A Theory on the Interdependent Origination and Entanglement of Chronogeometric Gravity, Matter and the Cosmos

Santos C. Borom

garland.maths@gmail.com

December 10, 2025

Abstract

This work advances a relational physics that begins not with spacetime but with a fibred continuum ($\mathcal{C}_d$) of primordial quantum singularities ($\mathfrak{B}_{\vec{r}}$) animated by a quantum holo-state, Holo-State $|\overset{\circ}{\mathbf{\Psi}}\rangle$,

$$\mathcal{C}_d := \left\{\mathfrak{B}_{\vec{r}},\ \hat{\Omega}_{\vec{r}},\ \mathfrak{F}_{\vec{x}}\right\}_{\vec{x}\in\mathcal{C}_d}.$$

Reality is scaffolded by a triadic distance and time *relational phase-geometry bundle* ($\pi : \mathfrak{F} \to \mathcal{X}_q$) with fibres ($\mathfrak{F}_x = \mathcal{F}_x^{\text{dist}} \times \mathcal{F}_x^{\text{time}}$), so separation and duration are read from the triad of *correlational*, *spectral*, and *vacuum-response* measures (and their temporal counterparts) rather than imposed as primitive background. At the foundation stands the tetralemma—the *Fourfold Algebra of Symmetry*, ($\overset{\circ}{\mathbf{\Lambda}}$), (Symmetry, Antisymmetry, Parasymmetry, Holosymmetry)—treated as a generative grammar whose concrete vocabulary is the full algebra of observables, deliberately prior to geometric assumptions.

The quantum holo-state $\omega^{\overset{\circ}{\mathbf{\Psi}}}$ is the global, holosymmetric state on the algebra that, via the Gelfand–Naimark–Segal (GNS) construction, converts valuations into *operational expectation values* and forges the inner products that knit local fibres into an emergent Hilbert arena, $\mathcal{H}_{\text{em}}$ and $\mathcal{H}_{\text{phys}}$, and its relational geometry. This forging proceeds through the fourfold inter-modal correlator $\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2]$, which couples the tetralemmatic modes and supplies the state-conditioned structure required for parallel transport

$$\langle\psi_1|\psi_2\rangle_{\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}} := \lim_{x'\to x} \langle\overset{\circ}{\mathbf{\Psi}}|\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2]|\overset{\circ}{\mathbf{\Psi}}\rangle$$

and comparison across the continuum:

$$\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2] := \hat{\mathcal{F}}_\alpha^\dagger(x)\, \mathcal{U}(x \leftarrow x'; \gamma)\, \hat{\mathcal{F}}_\beta(x') \ \in\ \mathcal{B}(\mathcal{H}_{\text{em}}),$$

where $\mathcal{U}(x \leftarrow x'; \gamma)$ is the gauge–covariant transporter along the path γ, and $\mathcal{B}(\mathcal{H}_{\text{em}})$ is the Banach *-algebra (indeed C*-algebra) of bounded operators on the emergent Hilbert space $\mathcal{H}_{\text{em}}$, with implicit summation over the modal indices α, β understood.

Two novel fundamental fields, $\hat{\mathcal{F}}_\alpha^\dagger(x)$ and $\hat{\mathcal{F}}_\beta(x')$, have been proposed to inhabit this structure. The *Gaiamiti* $\hat{\mathbf{\Phi}}_x$ shapes chronogeometry and carries the radiative content associated with the graviton; the *Tantrakyma* $\hat{\mathbf{\Theta}}_x$ organises excitations and internal symmetries, seeding particle spectra and interactions. Their interaction correlators, varied as a single operational action, yield the *Chronogeometric Einstein Equations*—a covariant, state-conditioned field

system that explicitly unifies general relativity with quantum mechanics, taking the form

$$\underbrace{\bar{\Sigma}_{\mu\nu}(x)}_{\text{General Relativity}} = \underbrace{\Big\langle \overset{\circ}{\mathbf{\Psi}} \Big| \hat{\Sigma}_{\mu\nu}(x) \Big| \overset{\circ}{\mathbf{\Psi}} \Big\rangle}_{\text{Quantum Mechanics}} = \Big\langle \overset{\circ}{\mathbf{\Psi}} \Big| \Big(\hat{g}_{\mu\nu}(x) + \hat{t}_{\mu\nu}(x) \Big) \Big| \overset{\circ}{\mathbf{\Psi}} \Big\rangle,$$

$$\bar{G}_{\mu\nu}[\bar{\Sigma}] + \Lambda\, \bar{\Sigma}_{\mu\nu} = \kappa\, \bar{T}^{\text{total}}_{\mu\nu}(x).$$

This tensor $\hat{g}_{\mu\nu}(x)$ is itself emergent, defined as the second Fréchet variation of the squared composite distance operator $D^2(x, x')$, evaluated at the coincidence limit:

$$\hat{g}_{\mu\nu}(x) := \frac{\partial^2}{\partial x^\mu \partial x'^\nu} D^2(x, x') \Big|_{x'=x},$$

where $D^2(x, x') := \hat{d}^2_{\text{corr}}(x, x') + \hat{d}^2_{\text{vac}}(x, x') + \hat{d}^2_{\text{spec}}(x, x')$. Each $\hat{d}_a(x, x')$ for $a \in \{\text{corr}, \text{vac}, \text{spec}\}$ is a Hermitian operator on $\mathcal{H}_{\text{total}}$. Complementing the metric tensor that governs emergent space, the framework defines its temporal analogue: the *operator-valued temporal curvature tensor* $\hat{t}_{\mu\nu}(x)$ arises from the triadic operators of relational time. It is formally defined as the second Fréchet variation of the squared composite temporal displacement $T^2(x, x')$ at the coincidence limit:

$$\hat{t}_{\mu\nu}(x) := \frac{\partial^2}{\partial x^\mu \partial x'^\nu} T^2(x, x') \Big|_{x'=x},$$

where $T^2(x, x') := \hat{t}^2_{\text{corr}}(x, x') + \hat{t}^2_{\text{vac}}(x, x') + \hat{t}^2_{\text{spec}}(x, x')$.

Cosmology is posed as a timeless problem $\hat{H}_{\text{univ}} | \overset{\circ}{\mathbf{\Psi}} \rangle = 0$ with $\hat{H}_{\text{univ}} := \hat{H}_\Phi + \hat{H}_\Theta + \hat{H}_{\text{int}}$: a universal quantum constraint fixes the realised state of the whole, while familiar evolutions appear when suitable internal clocks are conditioned from it. Within this discipline, the *Standard Model* is the holosymmetric, Becchi–Rouet–Stora–Tyutin (BRST)-physical realisation of the Tantrakyma sector: the anomaly-free gauge stack $G_{\text{SM}} = SU(3)_C \times SU(2)_L \times U(1)_Y$ acting on the matter registers, with Yukawa couplings and family structure fixed by cohomological and duality selection rules; electroweak symmetry breaking is a Tantrakyma vacuum alignment that produces observed masses and mixings as operational expectation values. The dark sector—dark matter and dark energy—arises as complementary holosymmetric branches of the same operator-geometric substrate rather than as an ad hoc addition, so that the full cosmic stack is given by

$$\begin{aligned} \mathcal{G}_{\text{Cosmos}} &= \underbrace{\mathcal{G}_{\text{Chronogeometry}}}_{\text{acts on } \delta\hat{\Sigma}_{\mu\nu} = \hat{h}_{\mu\nu} \oplus \hat{\tau}_{\mu\nu}} \times \underbrace{G_{\text{SM}}}_{\text{Tantrakyma (matter/gauge)}} \\ &= \Big[\text{Diff}_{\text{rel}} \ltimes \Big(\text{SO}(1,3)_{\text{local}} \times \mathcal{G}^{(\text{eff})}_{\text{Dark}} \Big) \Big] \;\times\; \Big[SU(3)_C \times SU(2)_L \times U(1)_Y \Big]. \end{aligned}$$

With

$$\mathcal{G}_{\text{Chronogeometry}} := \text{Diff}_{\text{rel}}(\mathcal{X}_q) \;\ltimes\; \Big(\text{SO}(1,3)_{\text{local}} \;\times\; \mathcal{G}^{(\text{eff})}_{\text{Dark}} \Big).$$

Here $\text{Diff}_{\text{rel}}(\mathcal{X}_q)$ are relational diffeomorphisms on $(\mathcal{X}_q, \bar{\Sigma})$, acting by pullback on fields and adjoint transport on internal fibres; $\text{SO}(1,3)_{\text{local}}$ are local Lorentz rotations; and $\mathcal{G}^{(\text{eff})}_{\text{Dark}}$ is the

effective symmetry badge governing the proposed τ–kairos (not SM matter):

$$\mathcal{G}^{(\mathrm{eff})}_{\mathrm{Dark}} \supset \underbrace{\mathrm{Shift}_{\varphi_b} \times \mathrm{Shift}_{\varphi_L} \times \mathrm{Shift}^{\mathrm{ax}}_{\tilde{\varphi}}}_{\text{scalar shifts}} \times \underbrace{U(1)_\tau}_{\text{Chrono–Kymon phase}} \times \underbrace{U(1)_{\mathrm{hel}}}_{\text{helicity plane}} \times \underbrace{\mathbb{Z}_2^{\mathrm{HS}}}_{\text{HSP tag}} .$$

The three Shift factors act on the breathing/longitudinal/axial scalars $(\varphi_b, \varphi_L, \tilde{\varphi})$; $U(1)_\tau$ is the proposed *Chrono–Kymon* phase symmetry of the τ-vector $K^{(\tau)}$; $U(1)_{\mathrm{hel}} \simeq SO(2)$ rotates the helicity doublet $(K^{(\tau)}_+, K^{(\tau)}_-)$; and $\mathbb{Z}_2^{\mathrm{HS}}$ labels *holosymmetric* (HSP) branch stability arising from algebra. All continuous factors are *effective* and may be softly broken by the same operators (masses, mixings, potentials) that drive cosmology.

In concise form, the emergence chain is

$$\mathrm{Ontology} \rightsquigarrow \mathrm{Algebra} \rightsquigarrow \mathrm{GNS/BRST}:$$
$$(\overset{\circ}{\mathbf{\Lambda}}),\ \mathcal{A}(\mathcal{C}_d) \rightsquigarrow (\mathcal{H}_{\mathrm{em}}, \hat{Q}_{\mathrm{BRST}}) \rightsquigarrow \mathcal{H}_{\mathrm{phys}} \simeq H^0_{\mathrm{BRST}}.$$

On this arena, $\hat{\mathbf{\Phi}}$ (geometry) and $\hat{\mathbf{\Theta}}$ (matter), with their interaction correlators, enter a single action whose variation yields the Chronogeometric Einstein Equations,

$$S[|\overset{\circ}{\mathbf{\Psi}}\rangle] := \int_{\mathcal{X}_q} d^d x\ \sqrt{-\det \bar{\Sigma}(x)}\ \mathrm{Tr}\Big(\Big\langle \overset{\circ}{\mathbf{\Psi}}\Big| \hat{\mathbf{G}}_\Lambda(x,x) \Big|\overset{\circ}{\mathbf{\Psi}}\Big\rangle\Big) .$$

By rooting metrics, clocks, and spectra in the same fibred continuum and relational phase-geometry bundle, the framework unifies ontology, kinematics, and phenomenology without imported scaffolds, offering a clean trajectory from question to detection, from detector readout to cosmos, and from algebra to evidence.

The evolution of any such subsystem is governed by this emergent dynamical law

$$i\hbar \frac{d}{dt} |\overset{\circ}{\mathbf{\Psi}}(t)\rangle = \hat{H}_{\mathrm{rel}}(t) |\overset{\circ}{\mathbf{\Psi}}(t)\rangle,$$

and the operator that drives this effective evolution is the *total relational Hamiltonian* $\hat{H}_{\mathrm{rel}}$, which contains the entirety of cosmic phenomenology:

$$\hat{H}_{\mathrm{rel}} = \underbrace{\hat{H}_{\mathrm{geometry}}[\hat{\Sigma}_{\mu\nu}]}_{\text{Dark Energy \& Dark Matter}} + \underbrace{\hat{H}_{\mathrm{matter}}[\hat{\phi}, \hat{\psi}, \dots]}_{\text{Standard Model}} .$$

This single effective Hamiltonian is composed of two sectors:

The Geometric Hamiltonian ($\hat{H}_{\mathrm{geometry}}$) This operator governs the dynamics of the Gaiamiti field ($\hat{\Phi}$), whose quanta are the *apeirons*. Its ground state eigenvalue is the cosmological constant (Λ), the physical origin of *Dark Energy*, while its large-scale coherent excitations are the physical manifestation of *Dark Matter*.

The Matter Hamiltonian ($\hat{H}_{\mathrm{matter}}$) This operator governs the dynamics of the Tantrakyma field ($\hat{\Theta}$), whose quanta are the *kymons*. Its excitations and their interactions give rise to the entire particle zoo of the Standard Model.

Dedication

Oṃ Namo Bhagavatyai Ārya-Prajñāpāramitāyai
Oṃ Homage to the Blessed and Noble Perfection of Wisdom

Namaḥ sarvajñāya
Homage to the All-knowing, the Omniscient

Inexpressible, inconceivable, and unutterable
Is the Perfection of Wisdom, Prajñāpāramitā.
She is unborn, unceasing, the nature of space.
The object of experience of self-cognizant primordial wisdom.

For the Wellbeing and Joy of all Beings and Worlds
And for the Perfection of Wisdom.

Contents

Abstract **i**

Preface: A Scientist's Invitation to Dare to Dream **xxiv**

Prologue: Entanglement at the Foundations of Physics **clxix**

I Foundations **1**

1 The Fourfold Algebra and The Singularity **2**
1.1 Introduction 2
1.1.1 The Relationship Between IQR and Existing Frameworks 20

1.1.2 The Law of Dependent Origination . . . 22
1.1.3 The Tetralemma: Reality's Four-Vertex Prism . . . 25
1.1.4 The Quantum Revelation: A Tetralemmatic Journey . . . 26
1.1.5 The Middle Way: A Narrative of Insight . . . 30
1.1.6 Aristotle and The Story of a Thesis . . . 31
1.1.7 The Debates of Ancient India . . . 32
1.2 The Tetralemma of Algebra . . . 33
1.2.1 The Definition of Thesis . . . 33
1.2.2 The Definition of Antithesis . . . 34
1.2.3 The Definition of Synthesis . . . 36
1.2.4 The Definition of the Holothesis . . . 38
1.2.5 Tetralemma of Algebra . . . 40
1.2.6 The Axiom of Duality and its Self-Similar Nature . . . 42
1.2.7 The Fourfold Dialectic Algebra of Symmetry . . . 47
1.2.8 A Prism for Discourse: Classical Logic as a Special View . . . 49
1.2.9 Born-Valued Semantics for the Four Corners . . . 51
1.3 The Primordial Quantum Singularity . . . 55
1.3.1 Coordinate Equivalence of the Primordial Quantum Singularity . . . 62
1.3.2 Defining Properties of the Primordial Quantum Singularity . . . 63
1.4 The Quantum Holo-State . . . 64
1.4.1 Gelfand-Naimark-Segal Construction of the Quantum Holo-State . . . 65
1.4.2 The Fourfold Dialectic Algebra and the GNS Construction . . . 68

2 The Fourfold Symmetry 73
2.1 Quantum Realizations of the Fourfold Symmetry . . . 73
2.1.1 Realization of The Ontological Duality States . . . 73
2.1.2 Local Involutive Duality Operators . . . 75
2.1.3 Global Involutive Duality Operators . . . 77
2.2 Parasymmetry: Quantum Realization of Synthesis . . . 79
2.2.1 Local Parasymmetric Projector of an PQS . . . 80
2.2.2 The Local Anti-Parasymmetric Projector . . . 82
2.2.3 The Global Parasymmetric Projector . . . 86
2.2.4 The Global Anti-Parasymmetric Projector . . . 89
2.2.5 Multipartite Parasymmetric Entanglement . . . 93
2.2.6 Multipartite Anti-Parasymmetric Entanglement . . . 96
2.2.7 Local and Pairwise Coherence Parasymmetric Generator . . . 99
2.2.8 Multipartite Modal-Class Parasymmetric Generator . . . 101
2.2.9 Multipartite Modal-Class Anti-Parasymmetric Generator . . . 104
2.2.10 The β Generalized Decoupled Unitary Evolution . . . 107
2.2.11 The Zonal Decoupled Unitary Evolution . . . 109
2.2.12 The Generalized Composite Decoupled Parasymmetric Evolution Operator 111
2.2.13 The $\hat{\sigma}$ Global Coupled Unitary Evolution . . . 115
2.2.14 The Global Composite λ Coupled Unitary Evolution . . . 118
2.2.15 The Unified Global Coupled Unitary Evolution . . . 120
2.2.16 Definition: Parasymmetry (Synthesis) . . . 123

2.3 Holosymmetry: The Quantum Realization of Holothesis . . . 128
2.3.1 System Definition and Component Operators . . . 129
2.3.2 Holosymmetry Component Operators . . . 142
2.4 The Interferometer Model of Holosymmetric Evolution . . . 153
2.4.1 (1) Symmetric Interferometer Model . . . 155
2.4.2 (2) Asymmetric Interferometer Model . . . 157

3 The Double-Slit Experiments 161
3.1 The Double-Slit Experiment for The Interferometer Model . . . 161
3.1.1 Experiment for the Symmetric Interferometer Model . . . 161
3.1.2 Interference Expectation in the Symmetric Interferometer Model . . . 165
3.1.3 Complementary Interference Probability . . . 167
3.1.4 Visibility of Holosymmetric Interference: Symmetric Interferometer Model 168
3.1.5 Experiment for the Asymmetric Interferometer Model . . . 169
3.1.6 Asymmetric Complementary Interference Probability . . . 174
3.1.7 Visibility of Holosymmetric Interference: Asymmetric Interferometer Model 175
3.1.8 Interferometric Visibility Bounds . . . 176
3.1.9 The Interferometer of Universal Histories . . . 180
3.2 Interferometric Model Recursive Expansion . . . 184
3.2.1 Recursive Component Expansion: Symmetric Interferometer . . . 184
3.2.2 Recursive Expansion: Asymmetric Interferometer . . . 185
3.2.3 Interpretations of the Component Expansions within the Double-Slit Analogy 186
3.3 The Interferometric Commutator Lensing . . . 187
3.3.1 Commutator Lensing of the Symmetric Interferometer . . . 187
3.3.2 Commutator Lensing of the Asymmetric Interferometer . . . 188
3.3.3 The Interferometric Commutator Lensing Double Slit Experiment . . . 189
3.3.4 Experimental Protocols for the Commutator Lensing . . . 189
3.3.5 Probing the Asymmetric Commutator . . . 190
3.4 The Holosymmetric Path Scattering Model . . . 191
3.4.1 Symmetric Path Scattering Model . . . 191
3.4.2 Symmetric Path Scattering Model Double Slit Experiment . . . 192
3.4.3 Asymmetric Path Scattering Model . . . 193
3.4.4 Asymmetric Path Scattering Model Double Slit Experiment . . . 194
3.5 The Recursive Expansion of the Path Scattering Model . . . 195
3.5.1 Recursive Component Expansion: Symmetric Path Scattering . . . 195
3.5.2 Recursive Component Expansion: Asymmetric Path Scattering . . . 197
3.6 Experiment for Recursive Path Scattering Model . . . 198
3.6.1 The Recursive Symmetric Path Scattering Case . . . 198
3.6.2 The Recursive Asymmetric Path Scattering Case . . . 200
3.7 Commutator Lensing of the Path Scattering Model . . . 202
3.7.1 Commutator Lensing of the Symmetric Path Scattering Model . . . 202
3.7.2 Commutator Lensing of the Asymmetric Path Scattering . . . 202
3.7.3 The Commutator Lensing Double-Slit Experiments . . . 203
3.8 Definition: Holosymmetry (Holothesis) . . . 205
3.8.1 Holosymmetry (Holothesis): Summary of Quantum Construction . . . 207

3.8.2 Probing Second-Order Coherence . . . 219
3.8.3 The Master Holosymmetric Operator . . . 222
3.8.4 From Duality to the Architecture of the Cosmos . . . 231

II The Continuum 242

4 Modal Fibres and Emergent Hilbert Spaces 243
4.1 Structure of Emergent Hilbert Spaces . . . 243
4.1.1 Twofold Inter-Modal Correlator of Primordial Quantum Singularities . . . 243
4.1.2 The Fourfold Inter-Modal Correlator of Quantum Singularities . . . 250
4.1.3 Emergent Hilbert geometry from the Fourfold correlator . . . 266
4.1.4 Parallel Transport Between PQS Loci . . . 267
4.1.5 Tensor Notation for the Fourfold Inter-Modal Correlator . . . 283
4.1.6 Block Matrix Representation Across Dialectic Modes . . . 285
4.1.7 Algebraic Structure and Graded Dynamics of the Fourfold Inter-Modal Correlator . . . 291
4.2 Holosymmetric Words in Channel Algebra . . . 311
4.2.1 Wordwise Meta-Claim, Compliance Vector, and Geometric Leverage . . . 334
4.3 The Dialectic Super-Commutator . . . 346
4.3.1 Endogenous Super-Commutator for Holosymmetric Words . . . 348
4.4 The Centralizing Role of Holosymmetry . . . 352
4.5 Holosymmetric Wordwise Covariant Kinematics (Along a Path) . . . 358
4.6 Emergent Inner Product Structure of Modal Hilbert Spaces . . . 359
4.6.1 Hilbert Bundle of Dialectic Modes over the Emergent Configuration Space 364
4.6.2 Wordwise Modal Bundle, Frames, and Trivializations . . . 369
4.6.3 Wordwise Channels, Mixing Bounds, and Kinematics . . . 371
4.6.4 Global Section of Dialectic Modes . . . 372
4.6.5 Wordwise Global Section of Dialectic Modes . . . 377
4.7 The Emergent Modal Space of a PQS . . . 381
4.7.1 Emergent Hilbert Space: GNS Realization and Modal Coupling . . . 384
4.7.2 Holonomy of the Modal Connection over PQS . . . 387
4.7.3 Wordwise Holonomy of the Modal Connection over PQS . . . 388
4.7.4 The Abstract Qubit . . . 394
4.7.5 The Abstract Qubit (Wordwise Relocation) . . . 400
4.7.6 The Holomodal Quantum Computer . . . 402
4.8 The Intrinsic Coherence of the PQS Continuum . . . 404
4.8.1 The Operator-Theoretic Nature of Ontological PQS Entanglement . . . 404
4.8.2 Factorizability of the Quantum Holo-State and the Necessity of Ontological Entanglement . . . 409
4.8.3 Wordwise Non-Factorizability of the Holo-State . . . 414
4.8.4 Fourfold Prismatic Superposition and Entanglement . . . 423
4.8.5 Wordwise Prismatic Superposition and Entanglement . . . 424
4.9 Operator-Theoretic Intrinsic Distances . . . 430
4.9.1 Correlator-Based Distance . . . 430

4.9.2 Wordwise Correlator-Based Distance (Holosymmetric Realization) 431
4.9.3 The Spectral Operator as a Fundamental Geometric Generator 433
4.9.4 Holosymmetric Wordwise Spectral Operators 435
4.9.5 Wordwise Reformulation of Operator-Theoretic Intrinsic Distances 446
4.9.6 Spectral Distance: Functional Geometry via Operator Commutators . . . 457
4.9.7 Wordwise Reformulations of Spectral Distance: Functional Geometry via Operator Commutators 458
4.9.8 Wordwise State-conditioned Modal Weight and Endogenous Regularization 461
4.9.9 The Relational Spectral Distance 464
4.9.10 Wordwise Relational Spectral Distance (Holosymmetric Realization) . . . 465
4.9.11 Endogenous Finiteness of Spectral Distance 467
4.9.12 Endogenous Finiteness of Wordwise Spectral Distance 468
4.9.13 The Principle of Coherent Regularization 471
4.9.14 Holosymmetric Wordwise Principle of Coherent Regularization 473
4.9.15 A Relational Framework for Infrared Regularization 476
4.9.16 Wordwise Relational Framework for Infrared Regularization 480
4.9.17 The Vacuum-Response Distance: Geometry from Entanglement Self-Coherence 494
4.9.18 Wordwise Vacuum-Response Distance 495
4.10 The Triadic Metric Structure of the PQS Continuum 501
4.10.1 Functional Orthogonality of the Distance Measures 501
4.10.2 Composite Triadic Distance and Metric 502
4.10.3 Infinitesimal Metric Tensor from the Triad 503
4.11 Holosymmetric Wordwise Triadic Metric Structure of the PQS Continuum . . 503
4.11.1 Functional Orthogonality of the Wordwise Distance Measures 504
4.11.2 Composite Wordwise Triadic Distance and Metric 504
4.11.3 Infinitesimal Wordwise Metric Tensor from the Triad 505
4.11.4 Distinction and Alignment of Inner Products 507
4.11.5 The Triadic Metric Structure (Fréchet) and the PQS Locus 510
4.11.6 The PQS as the Locus of Indistinguishability 511
4.11.7 Wordwise Triadic Metric Structure (Fréchet) and the PQS Locus 512
4.11.8 Wordwise PQS as the Locus of Indistinguishability 514
4.11.9 Holonomy and the Zero Locus of Distinguishability 515
4.11.10 Holonomy and the Zero Locus of Wordwise Distinguishability 517
4.12 The Triadic Structure of Emergent Time 521
4.12.1 Entanglement Phase Time 521
4.12.2 Wordwise Entanglement Phase Time 523
4.12.3 Vacuum Response Time 526
4.12.4 Wordwise Vacuum Response Time 528
4.12.5 Spectral Shear Time 531
4.12.6 Wordwise Spectral Shear Time 535
4.12.7 Functional Independence of Temporal Observables 538
4.12.8 Temporal Holonomy and the Zero Locus of Emergent Time 540
4.12.9 The Strong Static Spectral Structure 542
4.12.10 Wordwise Temporal Holonomy and the Zero Locus of Emergent Time . . 544

4.12.11 Holosymmetric Wordwise Strong Static Spectral Structure 547
4.13 Holosymmetric Wordwise Triadic Chronogeometry from Distances and Times . 548
4.13.1 Holosymmetric Wordwise Triadic Space and Triadic Time Fibres 549
4.13.2 Polar decomposition viewpoint: space as magnitude, time as phase 549
4.13.3 A Holosymmetric Wordwise Chronogeometric Structure 550
4.13.4 Infinitesimal Structure and Operator Metric-Chronometric Tensors 550
4.13.5 Conceptual consequences 551
4.14 The Minimal Scale and the Indivisibility of the PQS 551
4.14.1 Endogenous Finiteness of all Emergent Relational Observables 555
4.15 Holsymmetric Wordwise Minimal Scale and the Indivisibility of the PQS . . . 556
4.15.1 Wordwise Effective Spectral Cutoff 557
4.15.2 Endogenous Finiteness of All Holosymmetric Wordwise Observables . . . 558
4.15.3 The Ultra-Sub-Planckian Nature of the Singularity 558
4.15.4 Wordwise Ultra–Sub–Planckian Nature of the Singularity 564
4.16 The Relational Phase–Geometry Fibre Bundle 567
4.17 The Relational Phase–Geometry Fibre Bundle (Holosymmetric Wordwise) . . 573
4.18 The Fibred Continuum of Primordial Quantum Singularities 576
4.19 The Fibred Continuum of Primordial Quantum Singularities (Wordwise) . . . 580
4.19.1 Parallel Transport on the Relational Bundle 582
4.19.2 Parallel Transport on the Relational Bundle (Holosymmetric Wordwise) . 584
4.19.3 The Quantum Configuration Space 586
4.19.4 The Quantum Configuration Space (Holosymmetric Wordwise) 589

5 The Metric and Temporal Tensors 601
5.1 The Metric Tensor 601
5.1.1 Operator-Valued Emergent Metric Tensor from Composite Distance . . . 601
5.1.2 Physical Interpretation of the Operator–Valued Metric Tensor 604
5.1.3 Operator-Valued Emergent Metric Tensor (Holosymmetric Wordwise) . . 606
5.1.4 Expectation-Level Emergent Metric Tensor 608
5.1.5 Expectation-Level Emergent Metric Tensor (Holosymmetric Wordwise) . . 609
5.2 The Temporal Tensor 611
5.2.1 Operator–Valued Temporal Curvature Tensor 611
5.2.2 Physical Interpretation of the Operator–Valued Temporal Curvature Tensor Components 614
5.2.3 The Temporal Tensor (Holosymmetric Wordwise) 615
5.2.4 Expectation-Level Temporal Tensor 618
5.2.5 Expectation-Level Temporal Tensor (Holosymmetric Wordwise) 620
5.2.6 The Relational Chronogeometric Operator and Tensor 624
5.2.7 The Relational Chronogeometric Operator and Tensor (Holosymmetric–Wordwise) 641
5.2.8 Triadic Decomposition of the Relational Chronogeometric Tensor 643
5.2.9 Holosymmetrization of Triadic Channels 646
5.2.10 From Ultra-Sub-Planckian Operators to Macroscopic Geometry 648
5.3 Temporal and Geometric Manifolds 649
5.3.1 The Operator-Level Relational Geometric Manifold 649

5.3.2 The Expectation-Level Relational Manifold . . . 651
5.3.3 Operator-Level Relational Geometric Manifold (wordwise form) . . . 651
5.3.4 Defining the Lorentzian Structure from the Unified Tensor . . . 654
5.3.5 Holosymmetric Wordwise Lorentzian Structure from the Unified Tensor . . . 656
5.3.6 Symmetrizing the Unified Lorentzian Tensor . . . 657
5.3.7 Holosymmetric Wordwise Symmetrization of the Unified Lorentzian Tensor 659
5.3.8 Emergent Proper Time and Distance . . . 662
5.3.9 Metric Projection . . . 663
5.3.10 Holosymmetric Wordwise Emergence of Proper Time and Distance . . . 664
5.3.11 Holosymmetric Wordwise Metric Projection . . . 666
5.3.12 The Emergent Chronogeometric Manifold . . . 667
5.3.13 Holosymmetric Wordwise Emergent Chronogeometric Manifold . . . 668
5.3.14 The Nature of Spacetime: An Emergent, Relational Tapestry . . . 671
5.4 Causal Structures on the Chronogeometric Manifold . . . 672
5.4.1 The Effective Light Cone and its Primordial Origin . . . 672
5.4.2 Holosymmetric Wordwise Effective Light Cone and PQS Apex . . . 675
5.4.3 The Dynamical Texture: The Temporal Phase-Flow Structure . . . 677
5.4.4 A Complementary Two-Layered Causal Architecture . . . 678
5.4.5 The Unified Causal Architecture . . . 685
5.4.6 The Primordial Origin of Causality . . . 690
5.4.7 Holosymmetric Wordwise Primordial Origin of Causality . . . 692

III Laws, States and Fundamental Fields 701

6 The Gaiamiti and Tantrakyma Fields 702
6.1 The Nature of the Fundamental Fields . . . 702
6.1.1 The Inter-Modal Correlator Matrix of the Fields . . . 713
6.1.2 Holosymmetric Wordwise Unified Inter-Modal Correlator . . . 715
6.1.3 The Relational Levi-Civita Derivative . . . 723
6.1.4 Holosymmetric Wordwise Relational Levi-Civita . . . 725
6.1.5 The Determinant of the Effective Lorentzian Metric . . . 738
6.1.6 Holosymmetric Wordwise Determinant of the Effective Lorentzian Metric 739
6.1.7 Holosymmetric Wordwise Relational Volume Element . . . 742
6.1.8 The Vacuum Manifold: Variational and Spectral Formulation . . . 745
6.1.9 Holosymmetric Wordwise Formulation of the Vacuum Manifold . . . 748
6.1.10 The Trace of the Inter-Modal Correlator . . . 752
6.1.11 Holosymmetric Wordwise Trace of the Inter-modal Correlator . . . 753
6.1.12 Holosymmetric Control Dial — Fourfold Inter-modal Trace . . . 754
6.1.13 The Relational Hamiltonian . . . 755
6.1.14 Holosymmetric Wordwise Relational Hamiltonian . . . 757
6.1.15 Holosymmetric (HS) Extension of the Relational Hamiltonian . . . 759
6.1.16 Quantum Circuit Realization of the HS Relational Hamiltonian . . . 761
6.1.17 The Vacuum Manifold . . . 763
6.1.18 Holosymmetric Wordwise Vacuum Manifold . . . 765

6.1.19 Quantum Circuit Realization of the Vacuum Manifold . . . 771
6.1.20 An Operator-Theoretic Definition of the Tangent Bundle . . . 774
6.1.21 Holosymmetric Wordwise Tangent Bundle of the Vacuum . . . 776
6.1.22 Holosymmetric Lift of the Tangent Bundle of the Vacuum . . . 780
6.1.23 Quantum Circuits for the Vacuum Manifold and its Tangent Bundle . . . 782
6.1.24 Inner Product on the Vacuum Tangent Space . . . 784
6.1.25 Holosymmetric Wordwise Inner Product on the Vacuum Tangent Bundle 785
6.1.26 The Banach Space of Bounded Operators . . . 787
6.1.27 The Algebra of Endomorphisms: The Arena for Quantum Operators . . . 788
6.1.28 Holosymmetric Wordwise Banach Space and Endomorphism Bundle . . . 789
6.2 The Dynamical Gaiamiti Field . . . 793
6.2.1 Holosymmetric Wordwise Dynamical Gaiamiti Field . . . 794
6.2.2 The HS Quantum Circuits of the Gaiamiti Field . . . 802
6.3 The Dynamical Tantrakyma Field . . . 807
6.3.1 Holosymmetric Wordwise Dynamical Tantrakyma Field . . . 808
6.4 Bundle-Theoretic Formulation of the Foundational Fields . . . 812
6.4.1 Holosymmetric Wordwise Bundle-theoretic Formulation of the Foundational Fields . . . 815
6.5 The Action for the Dynamical Gaiamiti Field . . . 819
6.5.1 The Gaiamiti Field Action Functional . . . 820
6.5.2 The Holosymmetric Action for the Dynamical Gaiamiti Field . . . 821
6.5.3 Covariant Derivative of the Gaiamiti Field . . . 823
6.5.4 Holosymmetric Covariant Derivative of the Gaiamiti Field . . . 824
6.6 The Operator Bundle Connection as an Emergent Gauge Field . . . 828
6.6.1 Holosymmetric Operator Bundle Connection as an Emergent Gauge Field 829
6.7 Spontaneous Symmetry Breaking . . . 835
6.8 Emergent (Asymmetry-Induced) Symmetry . . . 837
6.9 Weaving The Two Arrows of Symmetry . . . 839
6.10 BRST Quantization and Ghosts: A Cohomological Prelude . . . 842
6.11 From Duality to the Symmetry–Asymmetry Quadrants . . . 849
6.11.1 Duality, Lifted to Two Layers . . . 849
6.11.2 Two Dyads and Their Product . . . 849
6.11.3 BRST-Compatible Reading . . . 850
6.11.4 Identification of the Quadrants and Mechanisms . . . 852
6.11.5 Holosymmetric Meta-Layer: Two-path Law/State Interferometry . . . 856
6.11.6 Operational Holosymmetric Cartography of the Quadrants . . . 867
6.11.7 Channel-Resolved Holosymmetric Diagnostics . . . 875
6.12 Scale-Aware Analysis for the Laws–State Matrix . . . 883
6.12.1 Purpose and Proviso . . . 883
6.12.2 Variational Backbone for Each Quadrant . . . 883
6.12.3 Operational Diagnostics (Scale-aware, Channel-resolved) . . . 898
6.12.4 Scale-tagged Variational Infrastructure (IR/UV) . . . 903
6.12.5 Working Strategy to Traverse the Quadrants . . . 928
6.12.6 IR/UV-aware Add-ons and Deliverables . . . 929
6.12.7 Practical Ledgers (Maintained Throughout) . . . 930

7 Field Laws and Physical States **931**
7.1 Quadrant I: Symmetric Vacuum and Scale-Tagged EOM . . . 931
7.1.1 Background Split and Vacuum Equations . . . 931
7.1.2 Scale-tagged Sector Actions and Euler–Lagrange Operators . . . 932
7.1.3 BRST-consistent Reading on $\mathcal{H}_{\text{phys}}$ . . . 932
7.1.4 UV/IR Knobs and Geometric Inputs . . . 933
7.1.5 Symmetric Vacuum at Scale (What the EOM Demand) . . . 933
7.2 BRST Health and Spectrum at the Symmetric Vacuum (Quadrant I) . . . 935
7.2.1 BRST Complex, Nilpotency, and the Physical Sector . . . 935
7.2.2 Gauge-fixed Quadratic Theory, Propagators, and Cohomological Invariance 936
7.2.3 Spectrum at the Symmetric Vacuum (Quadrant I) . . . 937
7.2.4 Scale Tags, Plateaus, and Robustness . . . 937
7.2.5 Quadratic Action, Hessian, and Transverse/Longitudinal Split . . . 938
7.2.6 Ghost Decoupling and Positivity on $\mathcal{H}_{\text{phys}}$ . . . 940
7.2.7 Tachyon Test and Hyperbolicity of the Physical Operator . . . 941
7.2.8 Anomaly Filters and Quantum BRST . . . 942
7.2.9 Gauge-parameter Independence of the Physical Poles (Nielsen Identities) . 945
7.2.10 Bandwise Checklist and Certification . . . 947
7.3 UV/IR Knobs and Plateau Certification . . . 949
7.3.1 Definitions of the UV/IR knobs . . . 950
7.3.2 Plateau Metrics and Acceptance Criteria . . . 954
7.3.3 Operational Certification Protocol . . . 958
7.3.4 Visibilities and their Scale Dependence . . . 960
7.3.5 Plateau Regions: Geometric Definition . . . 964
7.3.6 Gauge/BRST Robustness (Nielsen-type Independence) . . . 972
7.3.7 Regularisation Stability (Kernel/Projector Families) . . . 975
7.3.8 Channel-wise and Integrated Visibilities . . . 982
7.3.9 Acceptance Criteria and Certification Pipeline . . . 992
7.3.10 Quantitative Plateau Metrics . . . 995
7.3.11 Channel and Sector Compatibility . . . 997
7.3.12 Outcome and Policy Note . . . 999
7.4 Holosymmetric Probe: Controlled Two-path Evolution and Visibilities . . . 1000
7.4.1 Readout and Complex Coherence Amplitude . . . 1008
7.4.2 Channel Realisations: Law vs. State . . . 1011
7.4.3 Embedding of UV/IR Knobs and Sector Projectors . . . 1015
7.4.4 Gauge/BRST Robustness and Anomaly Filters . . . 1021
7.4.5 Plateau Coupling and Certification . . . 1026
7.4.6 Error Budget and Robustness to Noise . . . 1028
7.5 Exact Identities at the Symmetry Point (Baseline for Symmetry Breaking) . . . 1030
7.5.1 Band-limited Source Formalism and BRST-exact Gauge Fixing . . . 1030
7.5.2 Slavnov–Taylor Identity and Linearized Operator . . . 1032
7.5.3 Gauge-fixing, Antighost, and b-equations . . . 1034
7.5.4 Global G Ward Identities at the Dymmetric Vacuum . . . 1043
7.5.5 Nielsen Identity at the Symmetry Point . . . 1050
7.5.6 Cohomological Content and Physical Operators . . . 1055

7.5.7 Background-field Option . . . 1056
7.5.8 Global Ward Identities at the Symmetric Point . . . 1058
7.5.9 Slavnov–Taylor Identity and its Linearisation . . . 1063
7.5.10 BRST Charge, Nilpotency, and Equivalence Theorem . . . 1067
7.5.11 Nielsen Identities at the Symmetry Point . . . 1072
7.5.12 Callan–Symanzik Identity on the Band . . . 1075
7.5.13 Sector/Projector Compatibility and Plateau Protection . . . 1080
7.5.14 Anomaly Filters and Cohomological Triviality . . . 1086
7.5.15 Identity Ledger for Later SSB Analysis . . . 1088
7.6 Synthesis of Quadrant I: Discoveries and Baseline Status . . . 1089
7.7 Quadrant II: Spontaneous Symmetry Breaking (SSB) . . . 1091
7.7.1 Hold the Law Symmetric (Keep H_0 Fixed) . . . 1091
7.7.2 Prepare an Asymmetric State on the Vacuum Manifold . . . 1101
7.7.3 Band Hessian to BRST Projection (Transverse/Longitudinal) . . . 1111
7.7.4 Read Visibilities: Verify $|\Lambda_{\text{law}}| \approx 1$; Volume Decay of $|\Lambda_{\text{state}}|$; Phase–twist Interferometry for NG Soft Limits . . . 1122
7.7.5 Plateau Certification and Gauge-parameter Independence . . . 1132
7.7.6 Identity Ledger: Slavnov–Taylor, Ward, Nielsen, and Callan–Symanzik (On-band, Sectorwise) . . . 1145
7.7.7 Physicality Tests: Nonnegative Spectrum, Hyperbolicity, and Spectral Positivity on $\mathcal{H}_{\text{phys}}$ . . . 1152
7.7.8 Readout Consolidation and Cross–Quadrant Comparison . . . 1159
7.8 Quadrant III: Explicit (But Small) Law Breakings . . . 1164
7.8.1 Data, Regime, and Physical Reading . . . 1164
7.8.2 Tracking Law Emergence and Defining a Robust IR Slope . . . 1190
7.8.3 Restoration Length . . . 1219
7.8.4 RG Ledger: Breakings Become Irrelevant . . . 1230
7.8.5 Ward/ST/Nielsen Closure (Band–Projected) . . . 1251
7.8.6 Spectral Reconvergence to Symmetric Multiplets . . . 1268
7.8.7 Plateau Certification (Q III Mask) . . . 1291
7.8.8 Physicality Strengthens with ℓ . . . 1303
7.8.9 QIII Lessons & IR “Particle-like” Outlook . . . 1316
7.8.10 Concluding Census of Particle-like Modes on the QIII Plateau . . . 1317
7.9 Quadrant IV: Explicit Law Breaking with an Asymmetric State . . . 1320
7.9.1 UV/IR Instrumentation and Visibility Order Parameters . . . 1322
7.9.2 Fix the Symmetry Reference & Spurions . . . 1330
7.9.3 Cone–preserving Kernels . . . 1342
7.9.4 Visibilities (Primary Diagnostics) . . . 1346
7.9.5 State–visibility guardrail (no SSB / no decoherence of vacua) . . . 1352
7.9.6 State–visibility Envelope Selection, CIs, and Fallback Protocol . . . 1359
7.9.7 Law–visibility Slope Test and Tomographic Calibration . . . 1363
7.9.8 T-odd Tomography & Cones (No Law-signalling; Anomaly Filters) . . . 1370
7.9.9 Law–channel Diagnosis: Saturation vs. QIII . . . 1377
7.9.10 Near–degeneracies and GMOR Fits in the Mass Ledger . . . 1380
7.9.11 Ward–mass Link (GMOR–type Control of Pseudo–NG) . . . 1383

7.9.12 Multi–spurion GMOR Fits and PCAC/LSZ Bandwidth on the Plateau . . 1392
7.9.13 Gauge/BRST Safety & Spectral Shifts 1398
7.9.14 Gauge–flatness Audit and Analytic Tracking Window in $\{\epsilon_k\}$ 1406
7.9.15 Plateau–mask Certification (Report Only Where Robust) 1410
7.9.16 Numerical Thresholds and Grid–refinement Schedule for the Mask 1416
7.9.17 Channel–resolved Holosymmetric Readout 1419
7.9.18 Channel–resolved Readout: Operational Note (What Drives What) 1425
7.9.19 Synthesis of QIV 1429
7.10 Consolidation of Law–State Signatures, Ward Sources, and Spectral Content (QI–QIV) 1435
7.10.1 Comparative Table 1440

8 Parasymmetric Witnesses and Diagnostics 1448
8.1 Sector-Conditioned Witnesses and Quadrant Rules 1448
8.1.1 Fermionic Witnesses 1453
8.1.2 SUSY Witnesses 1455
8.1.3 p-Forms Witnesses 1457
8.1.4 Higher Spins Witnesses 1458
8.1.5 Hydrodynamics Witnesses 1459
8.1.6 Second Sound in the Hydrodynamic Sector (Matrix Formulation) 1461
8.1.7 Boundary Soft/Edge Modes Witnesses 1464
8.1.8 Anyons (Chern-Simons) Witnesses 1466
8.1.9 Fractons Witnesses 1467
8.1.10 Higher-Forms and Non-Invertible Symmetries 1470
8.1.11 Anomaly Witnesses 1471
8.1.12 Non-equilibrium (Floquet/Lindblad) Witnesses 1472
8.1.13 Universal Quadrant Coverage 1475
8.2 The Synthesis of Laws and States 1479
8.2.1 Parasymmetric Synthesis of Laws and States 1490
8.2.2 Parasymmetry (with SynTh) vs. Duality 1500
8.3 SynTh of Generalized Witness of Laws and States 1502
8.4 Parasymmetric Synthesis of Laws and States (Sector-conditioned Overlay) . . 1511
8.5 Applications of Parasymmetric Overlays 1514
8.5.1 Sector-Agnostic Parasymmetric Overlay 1516
8.5.2 Parasymmetric Overlay of the Fermionic (Spin-1/2) Sector 1518
8.5.3 Parasymmetric Overlay of Supersymmetry (SUSY) 1523
8.5.4 Parasymmetric Overlay of Boundary Soft/Edge Modes 1529
8.6 Parasymmetric Diagnostics of Type, Location, and Onset (TLO) 1532
8.6.1 Parasymmetric Meters and Laws-States Quadrant Readings 1532
8.6.2 Locator Sets: “Where” the Patterns Occur 1533
8.6.3 Onset Scales and Times: “When” the Patterns Occur 1534
8.6.4 Boundary/Edge Arenas and Sector Checks 1534
8.6.5 Typed Sector Witnesses and Parasymmetric Meters (Spin/Helicity/Polarization–conditioned) 1535
8.6.6 Spectral–Only Parasymmetric Protocol 1538

8.6.7 Application Arenas for SynTh ⊕ Parasymmetry (Typed Witnesses) . . . 1540
8.6.8 Parasymmetric Readout of Second Sound . . . 1542

9 The Quanta of Fields **1545**
9.1 Arena, Transport, Grading, and Band Tagging . . . 1550
9.2 Law vs. State, Visibilities, and the Four Quadrants (Pre-Physical) . . . 1556
9.3 From Algebraic Breakings to Quanta . . . 1563
9.4 Pre-physical Symmetry Reduction (PSR) . . . 1573
9.5 Realization to the BRST-physical Sector and Spectrum . . . 1584
9.6 Minimal "How-to" Pipeline . . . 1594
9.7 Invariance of Chronogeometric Formalisms . . . 1596
9.8 Parasymmetric Type–Location–Onset on the Sigma Pre-Physical Layer . . . 1597
9.9 Σ–Pre-physical Cones and Causality (Operator Level) . . . 1606
9.10 Kinematic Permit Census for Apeiron/Kymon (No *a priori* Spins) . . . 1613
9.11 Typed Σ–Contrasts & Visibilities (Pre-Physical) . . . 1622
9.12 Parasymmetric Meters on the Band (Typed) . . . 1633
9.13 Per Sector Execution Outline . . . 1640
9.14 Audits & Guardrails . . . 1648
9.15 Pre-Physical Artefacts . . . 1654
9.15.1 Typed Pre-Physics: Ledger, Visibilities, TLO . . . 1660
9.15.2 Permitted Kinematics for the Gaiamiti (Apeiron, ΦΦ) Field . . . 1662
9.15.3 Permitted Kinematics for the Tantrakyma (Kymon, ΘΘ) Field . . . 1663
9.15.4 Pre-Physics Parasymmetric Admission, Mechanism, Onset . . . 1665
9.16 Typed Parasymmetric Quadrants of the Gaiamiti and Tantrakyma Fields . . . 1666
9.16.1 Typed Subspaces and Projectors . . . 1672
9.16.2 Applying Parasymmetric Meters to Typed Subspaces . . . 1676
9.16.3 Assigning the Quadrant Per Type (Canonical decision on H^0) . . . 1679
9.16.4 Recording the Type, Location, and Onset (TLO) . . . 1681
9.17 The Gaiamiti Field Potential and the Origin of the Apeiron . . . 1683
9.17.1 The Apeiron as the Fundamental Quantum of Geometry . . . 1699
9.17.2 Spontaneous Symmetry Breaking of the Gaiamiti Field . . . 1703
9.17.3 Global Symmetry Breaking: NG and Radial Sectors . . . 1706
9.17.4 Internal Gauging of the Gaiamiti Sector and BEH Completion . . . 1710
9.17.5 The Composite Nature of the Goldstone Mode . . . 1716
9.17.6 The Emergence of the Massless Graviton via SSB . . . 1720
9.17.7 Euler–Lagrange Equation for the Gaiamiti Field . . . 1721
9.17.8 Noether Current for the Dynamical Gaiamiti Field . . . 1725
9.17.9 The Noether Stress–Energy Tensor of the Gaiamiti Field . . . 1732
9.17.10 Projection Operators and Geometric Duality . . . 1736
9.17.11 Tantrakyma Parasymmetric Type, Location, and Onset . . . 1741
9.18 The Action for the Dynamical Tantrakyma Field . . . 1743
9.18.1 The Tantrakyma Field Action Functional . . . 1749
9.18.2 Holosymmetric Wordwise Tantrakyma Field . . . 1752
9.18.3 Covariant Derivative of the Tantrakyma Field . . . 1755
9.18.4 Holosymmetric Wordwise Tantrakyma Covariant Derivative . . . 1757

9.18.5 Tantrakyma Field Potential and the Origin of the Kymon . . . 1760
9.18.6 Spontaneous Symmetry Breaking of the Tantrakyma Field . . . 1773
9.18.7 Gauge-Coupled Tantrakyma and the Brout–Englert–Higgs Mechanism . . 1776
9.18.8 A Minimal $U(1)$ Worked Example of the BEH Mechanism . . . 1783
9.18.9 Euler–Lagrange Equation for the Tantrakyma Field . . . 1795
9.18.10 Noether Current for the Dynamical Tantrakyma Field . . . 1797
9.18.11 The Noether Stress–Energy Tensor of the Tantrakyma Field . . . 1801
9.18.12 The Source of Emergent Curvature: A Unified View . . . 1804

10 The Weave of Geometry and Matter 1808
10.1 The Inter-Modal Correlators of Fields . . . 1808
10.2 Realized Weave Configuration and Typed Algebra . . . 1810
10.3 The τ-kairos Field . . . 1822
10.4 Chronogeometric Origin and Certified Construction of the τ–kairos . . . 1829
10.5 The Graviton in the Gaiamiti (Geometry) Sector . . . 1832
10.5.1 The Graviton Six Polarization Modes (Hypothetical) . . . 1840
10.6 The Higgs (Radial Kymon) in the Tantrakyma Sector . . . 1844
10.7 The Chrono–Kymon (Vector Channel of the τ-kairos) . . . 1847
10.8 The Spin–0 Apeiron (Gaiaton): Scalar Channel in the τ-kairos . . . 1851
10.9 Witnesses, Visibilities, and Parasymmetric TLO (Full Construction) . . . 1859
10.9.1 Law and State Ledgers: Residuals and Features . . . 1860
10.9.2 Calibrated Meters and Visibilities . . . 1862
10.9.3 Monotonicity Under IR Retuning . . . 1863
10.9.4 Onset Definition, Uniqueness, and Stability Radii . . . 1864
10.9.5 Explicit Witness Contents by Type and Sector . . . 1868
10.9.6 Parasymmetric TLO: Existence, Uniqueness, and Stability . . . 1870
10.10 SynTh Resolution and Audit for the Weave . . . 1872
10.10.1 Co-tenability, Conflicts, and Thresholds . . . 1877
10.10.2 Audit Moves and Basic Monotonicities . . . 1878
10.10.3 Finite Audit Paths and No-Explosion . . . 1881
10.10.4 Trigger Thresholds and Automatic Audit . . . 1884
10.11 Weaving the Sectors via the Unified Inter-Modal Correlator . . . 1886
10.11.1 Σ-weighted Generalized Eigenproblem for Mixing . . . 1887
10.11.2 TT-null Preservation for τ-mixing . . . 1890
10.11.3 Selection Rules for Typed Mixing . . . 1891
10.11.4 Stability Bounds: Mixing Norms vs. Visibility Changes . . . 1893
10.12 Triad-Resolved Reporting with Boundary vs. Bulk Provenance . . . 1899
10.12.1 Schema and Macros for Weave Reports . . . 1899
10.12.2 Arena-Invariant Spectral Set and Typed Stability . . . 1900
10.12.3 Boundary–vs–Bulk Invariance . . . 1902
10.12.4 Stability of Reported Onsets under Arena Changes and Mixing . . . 1904
10.13 Norms, Contours, and TT-Hyperbolicity . . . 1905
10.13.1 Admissible Norm Family . . . 1906
10.13.2 Measurable Structure for Riesz Integrals . . . 1908
10.14 Findings and Quadrants: High-Level Synthesis . . . 1911

10.15 Permitted Spins, Polarizations, Helicies, and Tags: High-Level Overview . . . 1913
10.16 Fundamental Quanta and Their Excitations . . . 1917
10.17 Geometric Block of the Unified Correlator . . . 1940
10.17.1 The Apeiron: Field–level vs. Particle–level Modes . . . 1944
10.17.2 The Chronogeometric Excitations . . . 1946
10.17.3 Excitation Channels and the Role of the Tau–Kairos . . . 1951
10.17.4 Observable (Kinematic) vs. Constitutive (Microscopic) Descriptions . . . 1952
10.17.5 Relational Graviton Propagator . . . 1955
10.17.6 Triadic Decomposition of Gravitational Propagation . . . 1961
10.17.7 Covariant Dynamics of Gravitational Fluctuations . . . 1963
10.17.8 Gravitational Waves . . . 1966
10.17.9 Tantrakyma and Kymon Contributions to Chronogeometry . . . 1968
10.18 Matter Block of the Unified Correlator . . . 1974
10.18.1 Definition and Typed Placement . . . 1974
10.18.2 Background–fluctuation Decomposition . . . 1976
10.18.3 Typed resolution and admissible matter channels . . . 1976
10.18.4 Relations to cross–blocks and spectrum . . . 1976
10.18.5 Emergent Matter Propagators . . . 1977
10.18.6 Dynamical Equations and Propagation . . . 1980
10.18.7 Triadic Decomposition of the Matter Propagator . . . 1983
10.19 Matter–Geometry Block of the Unified Correlator . . . 1986
10.19.1 Typed Off–diagonal Block and Connected Kernel . . . 1986
10.19.2 Modal Representation and Transport . . . 1987
10.19.3 Triadic Resolution and Parasymmetric TLO Interface . . . 1988
10.19.4 Functional Variation of the Interaction Action . . . 1989
10.19.5 Mutually–Sourced Field Currents and Typed Reciprocity . . . 1991
10.19.6 Triadic Decomposition of the Interaction Channels . . . 1992

11 The Graviton and Chronogeometric Gravity 1995
11.1 Quantization and Interaction of the Geometric Sector . . . 1995
11.1.1 Canonical Quantization of the Gaiamiti Field . . . 1999
11.1.2 Covariant Quantization via the Peierls Bracket . . . 2010
11.1.3 Graviton Mode Decomposition from Gaiamiti Fluctuations . . . 2014
11.1.4 Triadic Decomposition of the Graviton Field . . . 2032
11.1.5 Graviton Fock Space and Multi-Particle States . . . 2035
11.2 The Graviton Propagator . . . 2037
11.2.1 The Time-Ordering Operator and the Feynman Propagator . . . 2039
11.2.2 Mode Expansion and the Operator Basis of the Propagator . . . 2041
11.2.3 Graviton Propagator in Background Geometry . . . 2045
11.2.4 Triadic Structure and Interpretation of the Graviton Propagator . . . 2049
11.2.5 A Composite Gravitational Force and its Predictions . . . 2050
11.3 Graviton-Mediated Scattering . . . 2051
11.3.1 Linearized Interaction Action . . . 2053
11.3.2 Structure of the Vertex from the Interaction Kernel . . . 2053
11.3.3 Graviton Coupling Channels (modal decomposition) . . . 2054

11.3.4 Scattering Amplitudes from Graviton Exchange . . . 2054
11.3.5 Effective Graviton-Mediated Propagation . . . 2055
11.3.6 Positivity and Dispersion Bounds for Triadic Exchange . . . 2056
11.3.7 Graviton-Mediated Tantrakyma Scattering . . . 2057
11.3.8 Effective Graviton-Mediated Propagation . . . 2057
11.3.9 Graviton Coupling Channels . . . 2060
11.3.10 Scattering Amplitudes from Graviton Exchange . . . 2063
11.3.11 Effective Graviton-Mediated Propagation . . . 2064
11.3.12 Graviton-Mediated Tantrakyma Scattering . . . 2066
11.3.13 Positivity/Dispersion Bounds for Triadic Graviton Exchange . . . 2069
11.4 Geometry–matter interface in the typed (TT) formalism. . . . 2075
11.4.1 The Feynman Rules for Matter and Geometry . . . 2077
11.5 The Chronogeometric Gravity . . . 2079
11.5.1 The Unified Riemann Curvature Tensor . . . 2081
11.5.2 Ricci Tensor and Scalar Curvature of the Unified Geometry . . . 2085
11.5.3 The Relational Einstein–Hilbert Action . . . 2086
11.5.4 Unified Semi–Classical Einstein Equations . . . 2087
11.5.5 Derivation of the Unified Semi–Classical Einstein Equations . . . 2090
11.5.6 The Emergent Gravitational Coupling Tensor . . . 2095
11.5.7 Scalarisation, Universality, and Variation Bounds of the Gravitational Coupling Tensor . . . 2098
11.5.8 Recovery of General Relativity in the Spectral Limit . . . 2102
11.6 A Coherence–Energy Condition (Relational NEC) . . . 2105
11.7 Operator–Valued Einstein Equations with Unified Tensor . . . 2120
11.7.1 Formulation of the Operator Equation . . . 2120
11.7.2 Consistency, Conservation, and Semiclassical Reduction . . . 2121
11.7.3 The Einstein Tensor of the Unified Geometry . . . 2123
11.7.4 What is Gravity in the IQR Framework? . . . 2125
11.8 Chronogeometry of the τ–Kairos . . . 2138
11.8.1 Linear–response Quantization of the Non–TT sector and Optional BEH Mass for Chrono-Kymons . . . 2145
11.8.2 BEH of the Chrono-Kymon . . . 2147
11.8.3 The Chrono-Kymon as a Unified Dark Sector Candidate . . . 2152
11.8.4 BEH of the Gaiaton . . . 2153
11.8.5 Gaiaton as the Dark Matter and Energy Candidate . . . 2161
11.9 The Chronogeometric Regimes . . . 2163

IV The Standard Model 2169

12 The Standard Model 2170
12.1 Cosmic Constraints and Candidate Symmetry Groups . . . 2173
12.1.1 Characterization of the Candidate Symmetry Group . . . 2174
12.1.2 Matching to the Observed Fermion Content . . . 2201
12.1.3 The Anomaly Cancellation Constraint . . . 2204

12.2 The Holosymmetric Selection Principle (HSP) . . . 2211
12.2.1 Holothesis and the Superposition of Laws (Holosymmetry) . . . 2215
12.2.2 Stage I — The Self-Consistency Filter (Symmetric Interferometer) . . . 2216
12.2.3 Stage II — Comparative Tomography and Relational Orthogonality (Asymmetric Interferometer) . . . 2217
12.2.4 The Holosymmetric Selection Principle (HSP) . . . 2221
12.2.5 Structural Interlocks with the Foundations . . . 2223
12.3 BRST Symmetry and the Quantization of Gauge Fields . . . 2235
12.3.1 The Extended Field Space . . . 2236
12.3.2 BRST Cohomology and the Holosymmetric Selection Principle . . . 2248
12.4 The Emergence of Fermions from the Graded Tantrakyma Field . . . 2263
12.4.1 Leptons and Quarks as Irreducible Tantrakyma Representations . . . 2270
12.4.2 Representation Action of G_{SM} on Fermionic Kymons . . . 2272
12.4.3 Emergence of the Gell–Mann–Nishijima Relation from Tantrakyma Field Dynamics . . . 2282
12.4.4 Gauge Bosons as Excitations of the (Internal) Gauge Part of the Operator–Bundle Connection . . . 2288
12.4.5 Electroweak Symmetry Breaking via the Tantrakyma Potential . . . 2293
12.4.6 Fermion Masses from Yukawa Coupling to the Tantrakyma Field . . . 2297
12.5 The Threefold Multiplicity of Chiral Matter . . . 2301
12.5.1 From Triadic Time to a $\mathbb{C}^3$ Family Fibre . . . 2302
12.5.2 Why Exactly Three Families: a Minimal–faithful Holonomy Theorem . . . 2306
12.5.3 Family Connection, Yukawa Structure, and Flavor Mixing . . . 2310
12.5.4 Anomaly and HSP Compatibility . . . 2311
12.5.5 Flavor Mixing via the CKM and PMNS Matrices . . . 2313
12.5.6 The Inevitable Emergence of Mixing Matrices . . . 2317
12.5.7 Hadrons as Composite Kymon States . . . 2319
12.6 BRST Cohomology and the Hadronic Physical Space . . . 2329
12.6.1 Constraints, BRST Charge, and Nilpotency . . . 2329
12.6.2 Color Singletness and BRST Physicality . . . 2330
12.6.3 Typed Placement and Graviton Purity . . . 2330
12.6.4 Family Neutrality and Flavor Structure . . . 2330
12.6.5 Quartet Mechanism and Gauge–Invariant Reduction . . . 2331
12.6.6 Explicit Computation on the Hadronic Sector . . . 2333
12.6.7 Compatibility with Spectral Filtering and HSP . . . 2334
12.6.8 Technical Notes on Domains and Positivity . . . 2335
12.7 Neutrino Masses, the Seesaw, and PMNS Mixing . . . 2335
12.7.1 Neutrino Physics and the Majorana Seesaw Mechanism . . . 2335
12.7.2 PMNS Mixing in the Dirac and Majorana Cases . . . 2336
12.7.3 Dirac–Majorana Sector from the Tantrakyma . . . 2339
12.7.4 Type-I seesaw from block diagonalization . . . 2340
12.7.5 PMNS Mixing from Misalignment . . . 2343
12.7.6 Constraints from the Tantrakyma Potential and Family Fibre . . . 2345
12.7.7 Neutrino Mass and the Majorana Seesaw Mechanism . . . 2346
12.7.8 Gauging anomaly–free $U(1)_{B-L}$ . . . 2350

12.8 Antimatter in the IQR/Tantrakyma Standard Model . . . 2356
12.9 Antimatter from Complex Reps, Causality, Positivity, and Family Holonomy . 2359
12.9.1 Complex Representations of the Internal Connection . . . 2359
12.9.2 Relativistic Causality + Positivity ⇒ CPT and Conjugate Poles . . . 2362
12.9.3 Family Fibre Holonomy is Neutral but Conjugates under C . . . 2364
12.9.4 Consequences and Canonical Cases . . . 2364
12.10 Antimatter as a Duality Theorem . . . 2368
12.10.1 Categorical Duality on Matter (Algebraic, Kinematic, and Cohomological Faces) . . . 2368
12.10.2 Antimatter as a Duality Theorem . . . 2370
12.10.3 Operational Corollaries . . . 2371
12.10.4 Diagnostic realized reference configuration Structure that Forces Pairing (Spectral/Cohomological Face) . . . 2373
12.10.5 Typed Antiparticles as a Corollary of Duality . . . 2377
12.10.6 Operational Diagnostics . . . 2380
12.11 The Antimatter Theorem and Supersymmetry . . . 2384
12.11.1 Supersymmetric Extension of Antimatter Duality . . . 2386
12.12 The Standard Model with Gravity . . . 2390
12.13 The Yang–Mills Mass Gap . . . 2394
12.13.1 Step A: Field/Meter Basis (Pure, Compact simple G) . . . 2397
12.13.2 Step B: Plateau Certification Before Spectral Readout . . . 2399
12.13.3 Step C: Boundary Hygiene . . . 2406
12.13.4 Step D: Typed Spectral Extraction and CPT/KL . . . 2413
12.13.5 Step E: Cross–checks . . . 2422
12.13.6 Operational Resolution vs. Clay–level Theorem: a Guided Introduction . 2425
12.13.7 Modal Kinematics, Transport, and the Correlator Kernel . . . 2427
12.13.8 Operator–valued Metric and Triadic Split . . . 2437
12.13.9 Development of the Yang-Mills Continuum Theorem . . . 2447
12.13.10 The Yang-Mills Mass Gap Theorem and OS/Wightman Axioms . . . 2454

V Cosmology 2460

13 Dark Energy and Dark Matter 2461
13.1 Dark Energy and Dark Matter . . . 2461
13.1.1 Dark Energy as an Emergent Phenomenon of the Dual-Field Vacuum . . . 2462
13.1.2 Dark Matter as the Hidden Quanta of the Relational Fabric . . . 2463
13.1.3 The Interferometer as the Foundational Principle of Selection . . . 2469
13.1.4 The Holosymmetric Selection of the Dark Sector . . . 2470
13.2 Selecting the Nature of Dark Matter . . . 2472
13.2.1 Canonical Action Fragments and Notation (Per Hypothesis) . . . 2475
13.2.2 Interferometer-Based Selection (HSP) with Explicit Controls . . . 2480
13.2.3 Distinguishing Signatures for Selection . . . 2482
13.2.4 Multi-Component Selection (Optional) . . . 2487
13.2.5 Diagnostics (Not Selectors) . . . 2488

13.2.6 Selecting the Mechanism of Dark Energy . . . 2489
13.2.7 The Lensing Model as the Underlying Mechanism of Selection . . . 2492
13.2.8 Holosymmetric Selection of the Dark Sector . . . 2496
13.3 The Dark Universe: Gaiatons and Chrono–Kymons . . . 2501
13.4 An Effective Symmetry Stack for the Dark Sector . . . 2510
13.4.1 The cosmic symmetry stack (canonical chronogeometric placement) . . . 2512
13.4.2 Holosymmetric Selection and Operational Darkness . . . 2515
13.4.3 Factor I: Shift$_\varphi$ / $U(1)_\varphi$ (Apeiron / Dark–Energy Background) . . . 2515
13.4.4 Factor I(c): Shift$^{\rm ax}_{\tilde\varphi}$ (CP–odd axial mode) . . . 2517
13.4.5 Factor II: $U(1)_\tau$ (Chrono–Kymon / Dark–Matter Carrier) . . . 2518
13.4.6 Factor III: $\mathbb{Z}_2^{\rm HS}$ (stability & HSP sector tag) . . . 2521
13.4.7 The Abundances of Dark Energy and Dark Matter . . . 2524
13.4.8 From Spectral Measures to Cosmological Abundances . . . 2526
13.4.9 Relation to the Chronogeometric Regimes . . . 2526
13.5 Numerical Schemes for Abundance Determination . . . 2527
13.5.1 Route A. Spectral–First (State–First) Computation . . . 2528
13.5.2 Route B. Effective–Field Reduction: a Fast Lane for Numerics . . . 2529
13.5.3 Linear–Scale Structure with Chrono–Kymon Coherence (Effective) . . . 2530
13.5.4 Effective-Field Route for Abundances (Route B) . . . 2532
13.5.5 Spectral-First Route for Abundances (Route A): Inputs with Planckian UV Scale . . . 2538
13.5.6 Obtaining Numerical Values for Route A Inputs . . . 2541
13.5.7 Immediate Numerical Outputs at z=0 (Planck-Like Calibration) . . . 2543
13.5.8 Implications of the Calibrated Outputs . . . 2545
13.5.9 Executing the Next Actions . . . 2547
13.5.10 Synthesis and Significance of Calibrations . . . 2552
13.5.11 Estimating the PQS Scale from the Vacuum Residue . . . 2556
13.5.12 Mass–Energy of the Observable Universe . . . 2557
13.6 The Universal Quantum Constraint of the Emergent Cosmos . . . 2561
13.6.1 The Good-clock . . . 2563
13.6.2 The Emergence of Subsystem Schrödinger Dynamics . . . 2581

14 The Born Rule and Measurement **2587**
14.1 A Cosmic Context for Measurement . . . 2587
14.1.1 The Path Scattering Model: A Template for Collapse . . . 2588
14.1.2 A Universe Consistent with its Own Creation . . . 2591
14.1.3 The Axiom of Duality: The Logic of a Classical Outcome . . . 2592
14.2 The Born Rule . . . 2594
14.2.1 The Challenge: Derivation of the Born Rule . . . 2595
14.2.2 A Program for the Derivation of the Born Rule . . . 2596
14.3 A Formal Derivation of the Born Rule from First Principles . . . 2600
14.3.1 Stage 1: Measurement as Relational Entanglement and Decoherence . . . 2600
14.3.2 Stage 2: Probability from Symmetry and the Emergent Inner Product . . . 2607
14.3.3 Stage 3: Uniqueness from the Universal Constraint and the Dialectic . . . 2609
14.3.4 A Formal Proof of the Uniqueness of the Born Rule . . . 2618

15 Quantum Thermodynamics **2636**
15.1 The Laws of Quantum Thermodynamics . . . 2636
15.1.1 Foundational Definitions and Postulates . . . 2636
15.1.2 The Zeroth Law of Thermodynamics . . . 2644
15.1.3 The First Law Reimagined as Relational Energy . . . 2656
15.1.4 The Second Law of Relational Information Dynamics . . . 2663
15.1.5 The Third Law of Thermodynamics: Quantum Coherence in the Limit of Isolation . . . 2690
15.1.6 The Principle of Asymptotic Coherence . . . 2709

16 Black Hole Physics **2712**
16.1 Gravitational Collapse and Black Hole Formation . . . 2712
16.1.1 The Proposed Resolution to the Black Hole Information Paradox . . . 2714
16.1.2 The Field Equations for Black Holes . . . 2720
16.1.3 Modeling the Emergent Geometry . . . 2721
16.1.4 The Mathematical Path Forward: From Ansatz to Solution . . . 2723
16.1.5 Step 1: Calculating the Relational Christoffel Symbols . . . 2724
16.1.6 Step 2: Calculating the Unified Ricci Tensor . . . 2726
16.1.7 Step 3: The System of Differential Equations . . . 2728
16.1.8 Solving the System and Obtaining the Black Hole Solution . . . 2729
16.1.9 The System of Coupled Differential Equations . . . 2732
16.1.10 The Event Horizon: From Classical Boundary to Ontological Frontier . . . 2733
16.1.11 Approaching the Event Horizon: A Descent into Relational Reality . . . 2735
16.1.12 The Event Horizon as a Phase Boundary in Reality's Fabric . . . 2738
16.1.13 A Solvable Path: Asymptotic Matching . . . 2739
16.1.14 Step 1: Outer Solution — Classical Schwarzschild Limit . . . 2740
16.1.15 Step 2: Solving for the Temporal Tensor Correction . . . 2742
16.1.16 Deriving the System of Equations for the Temporal Tensor . . . 2743
16.1.17 The Energy of the Quantum Vacuum as the Physical Source . . . 2745
16.2 The Vacuum Energy Density . . . 2747
16.2.1 Intrinsic Entanglement Dressing and Spectral Gapping . . . 2747
16.2.2 Calculating the Dressed Vacuum Energy Density, $\rho_{\text{vac}}(r)$ . . . 2750
16.2.3 The Perturbative Solution for the Schwarzschild Black Hole . . . 2752
16.2.4 Solving for the Temporal Tensor $A(r), B(r)$ . . . 2752
16.2.5 Solving for the Geometric Correction $\delta f(r)$ . . . 2754
16.2.6 The Corrected Schwarzschild Black Hole Solution . . . 2755
16.3 The Unitary Transformation of Information . . . 2759
16.3.1 The Quantum Corrections to Black Hole Entropy . . . 2765
16.4 The Chronogeometric Kerr Black Hole . . . 2768
16.4.1 Embedding Kerr into the Emergent Chronogeometric Framework . . . 2776
16.4.2 Operator-Valued Angular Momentum and Frame-Dragging . . . 2777
16.4.3 The Inner Core: Ring Singularity and the Non-Classical Locus . . . 2778
16.4.4 Chronogeometric Vorticity and the Emergence of Horizons . . . 2780
16.4.5 Kerr Geometry as a Relational Coherence Phase . . . 2782
16.4.6 The Operator-Theoretic Kerr Reconstruction . . . 2783

16.4.7 Holosymmetric Modulation and Kerr Causal Bifurcation . . . 2786
16.4.8 The Holosymmetric Kerr Attractor as a Topological Quantum Phase . . . 2792
16.4.9 Quantum Ergodynamics and the Information Flow . . . 2794
16.4.10 The Entropic Dual of the Kerr Ergosphere and the Emergence of Horizon Temperature . . . 2800
16.4.11 Inner Horizon Instability and the Modal Breakdown . . . 2805
16.5 Computing the Chronogeometric Kerr Solution . . . 2813
16.6 Generalization of the Laws of Black Hole Thermodynamics . . . 2847
16.7 The Tau-Kairos at Horizons: Structure, Dynamics, and Thermodynamics . . . 2854
16.8 Chrono-Kymons at Horizons: Vector Coherence, Vorticity, and Rotational Thermodynamics . . . 2859
16.8.1 Two Benchmark Results in Linear Response . . . 2861
16.9 Unified Temporal Sector at Horizons: Interactions of Tau-Kairos tensors and Chrono-Kymons . . . 2863
16.10 The Temporal–Geometric Picture of Kerr Horizons . . . 2866
16.10.1 PQS Core and Global Black–Hole Energetics . . . 2867
16.11 The Lifecycle of a Black Hole . . . 2870
16.11.1 Overview: a Unitary, Relational Cycle without Paradox . . . 2870

17 Cosmic Evolution **2874**
17.1 The Law of Dependent Origination and Cosmic Evolution . . . 2874
17.1.1 The Origin of Dynamics: the Relational Einstein–Hilbert Action . . . 2874
17.1.2 The Formal Origin of the Cosmological Constant . . . 2876
17.1.3 Derivation of the Effective Cosmological Constant . . . 2882
17.1.4 Definition and Computation of the Holosymmetric Vacuum Residue . . . 2887
17.1.5 The Universal Schrödinger Equation . . . 2893
17.1.6 The Emergence of $E = mc^2$ from Relational Principles . . . 2896
17.1.7 The Law of Dependent Origination and Conservation of Duality . . . 2897
17.2 The Unfolding and Enfolding of the Universe . . . 2898
17.2.1 The Ātmapoiesis Universe . . . 2899
17.2.2 The Parapoiesis Universe . . . 2906
17.3 The Holocosmos: Cosmic Entanglement . . . 2912
17.3.1 The Axiom of Holocosmic Duality . . . 2916
17.3.2 The Dependent Origination of the Holocosmos . . . 2920
17.3.3 The Tetradic Truth Holocosmic Existence . . . 2925
17.3.4 Concluding Synthesis . . . 2925

Preface: A Scientist's Invitation to Dare to Dream

Acknowledgements

I wish it to be known and understood from the outset that the treatise presented herein lays no claim to original invention or discovery. None of the philosophies, principles, mathematics, or physical insights are solely of my own making. My role has been that of a humble and devoted student, learning from the luminous expositions of the great teachers of all ages — from the sages of Eastern and Western traditions spanning more than two millennia, to the brilliant minds of our own time. If I may be so bold as to claim any contribution, it is a modest one: an effort of translation and synthesis, seeking to render the timeless wisdom of ancient thought into the rigorous language of modern mathematics, and to weave the insights of the greatest thinkers into a single, coherent framework. Any understanding I may have achieved is due entirely to the generosity of those who dedicated their lives to creating the vast commentaries and discoveries that illuminate these profound subjects.

This acknowledgement is an essential part of the work itself, for the Interdependent Quantum Reality framework draws as deeply from the philosophical and human heritage of thought as it does from the mathematics that gives it form. That heritage is broad and without sectarian preference: the antecedent ideas appear in many streams of human inquiry — from the metaphysical investigations of Aristotle and Aquinas, to the subtle dialectics of Buddhist Madhyamaka philosophy, to the reflections of modern physicists and cosmologists. In drawing upon them, I have given none precedence over another, but have sought only to honour ideas that have endured because they speak to something universal, and all are valued equally as part of the shared legacy of humanity's search for truth. My aim has been to respect these enduring insights, and to show how they may now be expressed in precise mathematical form.

This work, therefore, is not a monument to novelty, but a pilgrimage — an offering of deep homage and reverence to the wisdom of the ages. It is an attempt to reveal the insights of those philosophical and intellectual giants, to share their brilliance so that we, together, may clear the dust from our eyes and see a little further by the light of their cumulative genius. It is with heartfelt gratitude to the Buddhas, Gurus, Scientists, and Philosophers of all ages that this work is offered. It is also a tribute to the imaginative spirit of science fiction novelists, and to my own primary school, high school, and university teachers. This treatise exists only because of their dedication and achievements. My deepest thanks go to my family, who have unfailingly supported me in my endeavors, and to my friends, who have

shared in this journey. *The work presented here is thus offered in reverence and devotion — for the welfare and joy of all the world.*

The Cosmic Quantum Dream

My own intellectual journey began with a simple yet profound encounter in the final year of primary school. I came across a science textbook, the name and author of which are now long lost to memory, but whose intellectual impact has remained with me ever since. In its pages, I found a marvelous description of the double-slit experiment, which sought to explain that light was possessed of a seemingly contradictory nature: it was both a particle and a wave. The concept of this wave-particle duality captivated my young mind, and I found myself lost in the vast and wondrous landscape of the imagination that it provoked—a landscape where the rigid certainties of the classical world dissolved into a shimmering reality of quantum potentia.

In parallel with this burgeoning scientific curiosity, a serendipitous loan from my local suburban library introduced me to the literary dimension of cosmic scale. The book was The Foundation Trilogy by the grandmaster of science fiction, Isaac Asimov, and it proved to be the progenitor of a lifelong passion for the genre. I was completely consumed by the cosmic breadth of its narrative, a story of such compelling power that I could rarely bring myself to put it down. Its allure was such that I would often find myself poring over its pages by torchlight, hidden beneath my blankets long after I was meant to be asleep, or at other times, by the faint, borrowed light of a distant street lamp.

By a happy coincidence of fortune, that same year saw the release of Star Wars: Return of the Jedi on VHS. I recall with great clarity a particular Friday afternoon when my primary school, in a welcome departure from its customary screening of Looney Tunes cartoons, presented the film to the grades 5 and 6 classes. The experience was transformative. I remember returning home that evening with an urgent desire to continue my journey through Asimov's literary galaxy, my imagination now alight with visions from two distinct yet harmonious universes. It was in those formative moments that I became a lifelong admirer of the grand tradition of science fiction, a passion that has continued to fuel my intellectual and philosophical inquiries ever since.

This early triad of influences—the quantum enigma of the double-slit experiment, the epic literary scope of the *Foundation Trilogy*, and the mythic grandeur of *Star Wars*—exerted a profound and lasting influence upon my intellectual aspirations. Indeed, I believe they provided the essential fuel for my subsequent academic pursuits in high school. It was here, amidst the structured world of secondary education, that another formative element was introduced, one that would later become central to my worldview: the conception of the universe as a grand, cosmic symphony.

During these years, I undertook the study of the clarinet, a pursuit in which I found unexpected joy and success, so much so that I was invited to join the high school band. The discipline required was demanding; I recall long evenings of rehearsal, stretching from the end of the school day late into the night. Yet, it was through this immersive experience in the collaborative harmony of the band that I began to perceive the world through a new sensory modality. Music, it seemed, endowed me with the ability to apprehend mathematics

in a different light—revealing its abstract structures as a dynamic and beautiful rhythm that transcended mere static rules and invited participation and experience.

My time in high school was one of rich intellectual discovery, a period during which I found myself particularly drawn to the elegant formalisms of science and mathematics, as well as to the distinct mode of cultural expression offered by the study of the Italian language. My dedicated application to these disciplines was met with consistent academic success, for which I remain grateful. This period of study culminated in the great fortune of being awarded a scholarship to the National Science Summer School—an immersive two-week program held in the nation's capital, which included a residence at the university.

It was here, however, that my intellectual path found its definitive vector. A visit to the radio telescope facility at Tidbinbilla, with its vast dishes aimed at the silent cosmos, and to the Australian National University's experimental Tokamak reactor, a testament to humanity's quest to harness the power of the stars, had a profound and orienting influence upon the direction of my future inquiries. These encounters with the grand instruments of scientific exploration solidified a nascent aspiration to understand the fundamental principles that govern the universe on both its grandest and most intimate scales.

My academic path continued to a university of great standing, where I first enrolled as a student of engineering. While I found a certain fluency in the language of engineering mathematics—at one point completing an examination on partial differential equations with a swiftness that surprised even myself—I felt a growing sense of disillusionment. The vicissitudes of life and a persistent longing for something more profound gradually revealed that the pragmatic focus of the discipline, however valuable, could not address the fundamental questions that had taken root in my mind. In my final year, I made the decision to change course, leaving engineering to pursue a Bachelor of Science with a double major in both pure and applied mathematics. It was here, particularly in an Honours Analysis class to which I was fortunately invited, that I gained a deeper appreciation for the abstract architecture of Hilbert spaces, a concept so central to the quantum view of the world. Throughout these undergraduate years, my formal studies were complemented by a voracious and undisciplined appetite for the works of the great thinkers in mathematics, physics, and philosophy—all consumed outside the structured curriculum in a private search for answers to the deep and unsettling questions about the nature of reality.

After graduating, I pursued a Master's Degree in Information Technology, focusing on Artificial Intelligence, where I wrote a thesis on Swarm Intelligence and had the honour of co-authoring a paper on self-organizing intelligent systems. A subsequent doctoral program was, alas, abandoned due to the pressing demands of life, in favour of a career in software engineering. This professional detour, along with my later experiences in machine learning and big data, afforded me the unexpected opportunity to become a lecturer and unit designer at a business school, where I helped create a Master of Business Analytics program. I found great satisfaction in both the craft of software engineering and the art of teaching. However, the foundational question concerning the nature of our cosmos, of our reality, remained unanswered, and this slowly grew into a persistent intellectual disquiet.

Did I ever abandon the hope that I would finally synthesize these disparate threads of knowledge to answer this mystery? No. The aspiration that I had as a primary school pupil, the formative experiences in high school, and the theoretical ideas that began to take shape during university never truly receded, always remaining prescient in the back of my

mind. Throughout my professional career, I continued my private attempts to reconcile my understanding of philosophy, mathematics, and physics, never giving up the hope that I would one day develop the clarity to put all the pieces I had learned over many decades into a single, coherent whole. The knowledge presented in this treatise is the culmination of those years of searching for a way to mathematically articulate what I understood in my heart to be true, an understanding I now wish to share with my fellow students of science so that we may *dare to dream*.

With the gratitude and origins of this work now spoken, we may turn to the dream that drives it — a dream that takes shape as the direct consequence of the successful unification of General Relativity and Quantum Mechanics. But what form does such a dream take? This treatise achieves that synthesis by showing that the long-standing conflict between the two great pillars of physics dissolves once we relinquish the assumption that spacetime is fundamental, an idea that resonates with contemporary research suggesting geometry is woven from the fabric of quantum entanglement [1, 2]. The grand visions of interstellar travel and cosmic connection, so vividly imagined in the science fiction that shaped my youth, thus cease to be idle fantasy; they become direct, answerable questions posed to the unified theory itself.

The mechanism for this unification is distilled into a single, foundational equation:

$$\underbrace{\bar{\Sigma}_{\mu\nu}(x)}_{\text{emergent chronogeometry}} \quad = \quad \underbrace{\left\langle \mathring{\boldsymbol{\Psi}} \,\middle|\, \hat{\Sigma}_{\mu\nu}(x) \,\middle|\, \mathring{\boldsymbol{\Psi}} \right\rangle}_{\text{quantum expectation}}$$

This expression—the **Chronogeometric tensor** $\bar{\Sigma}_{\mu\nu}(x)$ (Definition 5.25)—is the precise bridge between classical and quantum descriptions [3, 4]. It asserts that the curved spacetime of General Relativity is not an independent backdrop, but the macroscopic statistical average of a deeper quantum structure.

This emergent geometry arises from two fundamental quantum ingredients:

1. the *Quantum Holo-State* $\left|\mathring{\boldsymbol{\Psi}}\right\rangle$ (Definition 1.27), encoding the total relational state of the universe [3]; and

2. the *Chronogeometric Operator* $\hat{\Sigma}_{\mu\nu}(x)$ (Definition 5.26), defined by

$$\hat{\Sigma}_{\mu\nu}(x) \;:=\; \hat{g}_{\mu\nu}(x) \;+\; \hat{t}_{\mu\nu}(x),$$

   where $\hat{g}_{\mu\nu}(x)$ generates the geometric structure and $\hat{t}_{\mu\nu}(x)$ captures the temporal phase-flow of quantum phenomena [4, 5]. (In the main text we specify $\hat{t}_{\mu\nu}$ as a *symmetric*, covariant operator with a common invariant core and give its GNS representation at each PQS; see Section 11.5.4.)

The full dynamical content is expressed by the *Chronogeometric Einstein Equations* (Section 11.5.4):

$$\bar{G}_{\mu\nu}[\bar{\Sigma}] \;+\; \Lambda\,\bar{\Sigma}_{\mu\nu} \;=\; \kappa\,\bar{T}^{\text{total}}_{\mu\nu}(x).$$

Here and throughout, indices are raised and lowered using $\bar{\Sigma}_{\mu\nu}$ (so the source is written covariantly as $\bar{T}^{\text{total}}_{\mu\nu}$), and we restrict attention to histories for which $\nabla^{\mu}\bar{\Sigma}_{\mu\nu} = 0$ with respect

to the torsion-free connection compatible with $\bar{\Sigma}$; the contracted Bianchi identity then yields $\nabla^{\mu}\bar{T}^{\text{total}}_{\mu\nu} = 0$. In decohered regimes $\bar{\Sigma}_{\mu\nu} \to \bar{g}_{\mu\nu}$, so the Λ term reduces to its familiar GR form and the equation recovers Einstein's law in outward appearance [6, 7].

On the geometric side, curvature is built from the unified chronogeometric field $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$, so spacetime responds not only to spatial distinguishability ($\bar{g}$) but also to the quantum flow of phase ($\bar{t}$) [8, 3, 9]. On the source side, $\bar{T}^{\text{total}}_{\mu\nu}$ is the expectation-valued stress–energy combining the material/radiative *Tantrakyma* sector with the energy associated to the geometric field itself (*Gaiamiti*) [10, 11, 12, 13]. In this reading, the classical geometry we observe is the expectation of the unified operator $\hat{\Sigma}_{\mu\nu}$ in the holo-state—a property *of* the quantum state rather than a pre-existing stage [10, 11, 8, 3, 14].

With the imaginative impetus now fixed, we turn from recollection to construction. The pages that follow advance a single programme: to translate the images above into invariants and the questions into operators. *Part I: The Foundations* introduces the primitive grammar and logic of the framework—the Fourfold Dialectic Algebra and the Primordial Quantum Singularity—within which the central objects already invoked receive their formal dress: the Quantum Holo-State, the Chronogeometric Operator, and the unified chronogeometric tensor $\bar{\Sigma}_{\mu\nu}$ (see Definitions 1.27, 5.25 and 5.26). With these definitions in place, the narrative becomes calculational and the "cosmic dream" acquires equations. We begin, accordingly, with the alphabet and generating symmetries of the theory.

Part I: The Foundations

Chapter 1: The Fourfold Algebra and the Singularity

The chapter proposes a four-vertex "prism" for thinking—Symmetry, Antisymmetry, Parasymmetry, and Holosymmetry—where the immanent base (the first three) encodes relational dynamics and the apex (Holosymmetry) governs the whole, so that "symmetry" in its fullest sense is the indivisible fourfold rather than a single invariance [15, 16, 17, 18]. This prism is operationalised as a Fourfold Dialectic Algebra (Section 1.2.8) whose atomic unit is the *Tetralemma of Algebra*, $\mathring{\mathbf{\Lambda}} = \langle \mathsf{Th}, \mathsf{AntiTh}, \mathsf{SynTh}, \mathsf{HoloTh} \rangle$, closed and indecomposable (Sections 1.2.5, 2.3 and 3.8 and Definition 1.5). "The Quantum Revelation: A Tetralemmatic Journey" then reads the passage from superposition to classical fact as a traversal of this prism [19, 20]: unmeasured potentialities inhabit corners beyond classical bivalence, while measurement and decoherence "close the cycle," and truth becomes a graded warrant given by Born valuation with a concrete "prism protocol" $\rho \mapsto (v_T, v_A, v_S, v_H)$ (Sections 1.2.8 and 1.2.9 and eq. (3.36)) [21, 22, 23, 24, 25, 26]; when interrogations commute, Non-Contradiction and Excluded Middle reappear as theorems rather than axioms, and in a Gelfand–Naimark–Segal (GNS) realisation these valuations become ordinary expectations on the emergent Hilbert space (Sections 1.2.8 and 1.4.2) [27, 28, 29, 30].

A scale-free Axiom of Duality stabilises the picture: a functorial $\mathbf{D}_{\mathcal{C}}$ acts coherently on states, formulas, and rules, keeping the tetralemma closed under admissible re-indexings [27, 28, 29, 31, 32] (Proposition 1.12); concretely, each Primordial Quantum Singularity (PQS) $\mathfrak{B}_x$ bears a four-generator *-algebra $\mathcal{A}_x^{(4)}$ congruent to these modalities, providing the grammar later spoken by the Algebra of Observables and uttered via GNS into measurable tests [30, 22]

(Definitions 1.16 and 1.32). Historically, the programme is framed between Aristotle's forging of a defensible *thesis* and the fortress of non-contradiction (Section 1.1.6) and the Indian discipline of *vāda*, where a *pratijñā* proceeds by regulated, five-membered proof toward *tattva-nirṇaya* [18, 33, 34] (Section 1.1.7); conceptually, these strands feed the final outlook: four guiding principles elevate dependent origination into generative physics [17, 35, 16]—via an $\hat{\Omega}$-Continuum of Primordial Quantum Singularities and a single, non-factorizable Quantum Holo-State whose GNS realisation selects and builds its arena [30, 20]—yielding Gaiamiti (chronogeometry/gravity) and Tantrakyma (the Standard-Model façade) as two effective, jointly constrained sectors [36, 37, 38, 39], with emergent chronogeometry responding to total quantum stress–energy and thus closing the loop between coherence, curvature, and matter [40, 41, 42, 43, 44, 45] (Sections 1.1.1 and 4.1).

From Blueprint to Actualization. Chapter 1 presents the *Fourfold Dialectic Algebra of Symmetry* (Definition 1.16) as an *a priori* blueprint of admissible modes of assertion—**Symmetry** (Thesis Th), **Antisymmetry** (Antithesis AntiTh), **Parasymmetry** (Synthesis SynTh), **Holosymmetry** (Holothesis HoloTh)—*in the spirit of tetralemmatic logics and their modern analyses* [17, 18, 15, 16], and then shows how the Gelfand–Naimark–Segal (GNS) construction (Section 1.4.2) turns those logical forms into concrete quantum tests and expectations [30, 29, 22] (Definition 1.18) within an emergent Hilbert arena [20]. In short: Dialectic provides the rules of construction; GNS performs the construction.

Three-tiered emergence pipeline. (1) *Ontological (generative logic):* at each PQS $\mathfrak{B}_x$ (Section 2.3), the Fourfold Dialectic furnishes primitive "ringed" ontological labels (cf. Remark 1.22) [35, 15, 16]; (2) *Algebraic (operational vocabulary):* the C^*-algebra $\mathcal{A}(\mathcal{C}_d)$ (Definition 1.32) is generated by geometric (Gaiamiti), matter (Tantrakyma), and symmetry fields [30, 36, 37]; (3) *Physical (emergent representation):* a GNS state $\omega^{\mathring{\mathbf{\Psi}}}$ yields $(\pi_{\mathring{\mathbf{\Psi}}}, \mathcal{H}_{\mathrm{em}}, |\Omega_{\omega^{\mathring{\mathbf{\Psi}}}}\rangle)$ (Section 1.4.2) [30, 29, 20], in which all corner-valuations become ordinary expectations [21, 22] (Definition 1.18 and Section 4.1).

The Fourfold Dialectic Algebra (ontological layer)

Born semantics and facets. At each $\mathfrak{B}_x$, the four modes act as testable facets with Born-valued semantics (Section 1.2.9 and Definition 1.18). Thesis/Antithesis probe a distinguished axis; Synthesis interrogates coherent coexistence on that axis; Holothesis is a controlled, interferometric meta-probe of global implementation. Truth is the Born value $v(\rho; P) = \mathrm{Tr}(P\rho)$, so facets need not be mutually exclusive; they register complementary warrant-degrees in a single preparation ρ.

Duality and status-compatibility. A context-indexed duality $\mathbf{D}_{\mathcal{C}}$ (an involution) exchanges Thesis and Antithesis while fixing Synthesis and Holothesis (Definition 1.16 and Proposition 1.12):

$$\begin{aligned} \Delta_{\mathcal{C}}(\mathsf{Th}) = \mathsf{AntiTh}, \quad \Delta_{\mathcal{C}}(\mathsf{AntiTh}) = \mathsf{Th}, \\ \Delta_{\mathcal{C}}(\mathsf{SynTh}) = \mathsf{SynTh}, \quad \Delta_{\mathcal{C}}(\mathsf{HoloTh}) = \mathsf{HoloTh}, \\ \mathbf{D}_{\mathcal{C}}^2 = \mathrm{Id}, \end{aligned}$$

with status-compatibility

$$\mathsf{Status}_{\mathcal{C}}(\varphi,\lambda)=X \;\Rightarrow\; \mathsf{Status}_{\mathcal{C}}(\varphi^{\mathbf{D}},\lambda)=\Delta_{\mathcal{C}}(X),$$

and inapplicability invariant under dualization (Section 3.8).

Object-level instance. If $|\mathring{\mathbf{0}}\rangle$ is any primitive ontological state, define $|\mathring{\mathbf{1}}\rangle \coloneqq \mathbf{D}_{\mathcal{C}}(|\mathring{\mathbf{0}}\rangle)$ (Definition 1.16). Then

$$|\mathring{\mathbf{0}}\rangle \leftrightarrow |\mathring{\mathbf{1}}\rangle, \qquad \mathbf{D}_{\mathcal{C}}(|\mathring{\mathbf{1}}\rangle)=|\mathring{\mathbf{0}}\rangle,$$

so the dyad is semantically primary (Remark 1.22).

Self-similarity and closure. Under any admissible re-indexing $\mathcal{C}\to\mathcal{C}'$ that preserves dependent origination, there exists a unique $\mathbf{D}_{\mathcal{C}'}$ with

$$F\circ\mathbf{D}_{\mathcal{C}}=\mathbf{D}_{\mathcal{C}'}\circ F,$$

hence $\{\mathsf{Th},\mathsf{AntiTh},\mathsf{SynTh},\mathsf{HoloTh}\}$ is closed and pointwise fixed under repeated dualization (Proposition 1.12).

Parasymmetry (Synthesis). In the distinguished $(0,1)$-sector, Synthesis is the normalized superposition

$$\tfrac{1}{\sqrt{2}}\big(|\mathring{\mathbf{0}}\rangle+|\mathring{\mathbf{1}}\rangle\big).$$

The projector P_{para} isolates this coherent mode with Born valuation $v_{\text{S}}(\rho)=\text{Tr}(P_{\text{para}}\rho)$ (Section 1.2.9 and Definition 1.18). The pair $\{P_{\text{para}},P_{\text{antipara}}\}$ forms an orthogonal, complete resolution of the $(0,1)$-sector, guaranteeing basis-coherence with Thesis/Antithesis. Beyond a single locus, Synthesis becomes *relation*: bipartitely a Bell-like entangled mode with projector $P^{(xx')}_{\text{para}}$; multipartitely a W-like $|\Psi^{(n)}_{\text{para}}\rangle$ with global projector $P^{(x_1\ldots x_n)}_{\text{para}}$, exhibiting genuine multipartite entanglement and permutation symmetry (Section 1.2.8).

Holosymmetry (Holothesis). Holothesis formalizes typed *inapplicability* of the base existential polarity and its invariance under dualization (Section 3.8). Operationally, a holosymmetric control $\mathbf{U}_{\text{ctrl}}$ branches dual registers under two lifted path-laws, recombines, and reads out on the control—interrogating *architecture* rather than local alternatives (Section 2.3). The Holothesis effect $E^{(\mathbf{a}\to\mathbf{b})}_{\text{Holo}}$ yields the Born value

$$v_{\text{H}}(\rho)=\text{Tr}\Big(E^{(\mathbf{a}\to\mathbf{b})}_{\text{Holo}}\,\rho\Big)$$

as defined in (eq. (3.36)).

The "Prism for Discourse" and Born semantics

Quantitative facets. Logic is treated as diagnostic interrogation—a prism that refracts discourse into four facets, each returning a quantitative degree of warrant $v\in[0,1]$ (Section 1.2.8). In the formal core, each facet $\square\in\{\text{T},\text{A},\text{S},\text{H}\}$ corresponds to an effect $E_{\square}$ with

$$v_{\square}(\rho)=\text{Tr}(E_{\square}\rho), \qquad P_T=|e_1\rangle\langle e_1|, \quad P_A=|e_2\rangle\langle e_2| \;\text{ on the } (0,1)\text{-sector},$$

as specified by the Born semantics of the fourfold effects (Section 1.2.9 and Definition 1.18).

Classical logic as a special slice. On the $(0,1)$-sector,

$$P_T + P_A = P_{(0,1)} \;\Rightarrow\; v_T + v_A = 1 \quad \text{(Excluded Middle)}.$$

In the commuting/decohered limit, $P_T P_A = 0$ and SynTh carries no independent signal (Non-Contradiction); the fourfold prism collapses to the classical fortress (Section 1.2.8).

Reading a discourse. Given ρ, perform complementary interrogations:

$$(v_T(\rho), v_A(\rho)), \qquad v_S(\rho) = \mathrm{Tr}(P_{\mathrm{para}}\rho), \qquad v_H(\rho) = \mathrm{Tr}(E^{(\mathbf{a}\to\mathbf{b})}_{\mathrm{Holo}}\rho).$$

These are not mutually reducible and resolve distinct structural aspects (Section 1.2.8 and eq. (3.36)).

Algebraic and GNS layers (actualization)

Algebra of Observables $\mathcal{A}(\mathcal{C}_d)$. $\mathcal{A}(\mathcal{C}_d)$ (Definition 1.32) is the C^*-algebra generated (per locus $\vec{r} \in \mathcal{C}_d$) by: Gaiamiti geometric fields $\{\hat{\mathbf{\Phi}}_{\vec{r}}\}$, Tantrakyma matter-content fields $\{\hat{\mathbf{\Theta}}_{\vec{r}}\}$, and duality/symmetry generators $\{\hat{\mathbf{\Lambda}}_{\vec{r}}\}$ (Definition 1.16). It is the smallest norm-closed, involutive algebra containing finite linear combinations, products, and adjoints of these generators (Definition 1.32).

GNS construction of the Quantum Holo-State. From the state functional $\omega^{\mathring{\mathbf{\Psi}}}$ one constructs the cyclic triple (Section 1.4.2)

$$(\pi_{\mathring{\mathbf{\Psi}}},\ \mathcal{H}_{\mathring{\mathbf{\Psi}}} = \mathcal{H}_{\mathrm{em}},\ |\Omega_{\omega^{\mathring{\mathbf{\Psi}}}}\rangle).$$

The ontological symbol $|\mathring{\mathbf{\Psi}}\rangle$ is thereby represented by the cyclic vector $|\Omega_{\omega^{\mathring{\mathbf{\Psi}}}}\rangle$, and for any effect E,

$$\omega^{\mathring{\mathbf{\Psi}}}(E) = \langle\Omega_{\omega^{\mathring{\mathbf{\Psi}}}}|\pi_{\mathring{\mathbf{\Psi}}}(E)|\Omega_{\omega^{\mathring{\mathbf{\Psi}}}}\rangle, \qquad v(\rho; E) = \mathrm{Tr}\Big(\rho\,\pi_{\mathring{\mathbf{\Psi}}}(E)\Big),$$

where dialectic valuations coincide with Born semantics (Definition 1.18). The GNS triple is unique up to unitary equivalence (Section 1.4.2).

Self-generating arena (background independence). The state not only fixes expectations; it generates its own Hilbert space and inner product (Sections 1.4.2 and 4.1 and Proposition 1.30):

$$\Big\langle \pi(A)|\Omega\,,\ \pi(B)|\Omega\Big\rangle = \omega^{\mathring{\mathbf{\Psi}}}(A^\dagger B).$$

Thus "the state defines the space that contains it."

The Quantum Holo-State and scale-bridging roles

Local kinematics and global geometry. $\overset{\circ}{\boldsymbol{\Psi}}$ fixes the *local* inner product on each modal Hilbert space $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}$ via the Fourfold Inter-Modal Correlator (Proposition 1.30):

$$\langle \psi_1 | \psi_2 \rangle_{\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}} := \lim_{x' \to x} \langle \overset{\circ}{\boldsymbol{\Psi}} \mid \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2] \mid \overset{\circ}{\boldsymbol{\Psi}} \rangle,$$

and, complementarily, determines the *global* emergent spacetime manifold $\mathcal{M}_{\text{em}}^{(\overset{\circ}{\boldsymbol{\Psi}})}$. A single Holo-State thus provides the defining context for both "arena" and "actors."

PQSs, the $\hat{\Omega}$-continuum, and $\mathcal{X}_q$

Primordial loci and configuration space. A Primordial Quantum Singularity $\mathfrak{B}_{\vec{r}}$ (Section 2.3) is an elementary operator-valued locus whose definitive quantum nature is the Quantum Holo-State $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$ (Section 1.4.2 and Proposition 1.30). The framework replaces a fixed $\mathbb{C}^2$ qubit with a field-embedded, duality-governed, operator-valued singularity as the apex of causality (Definition 1.32 and Proposition 1.12). The tension between a universal state and inherently local operators motivates a *quantum configuration space* $\mathcal{X}_q$ (Section 4.1): distinguishable loci arise by relational crystallization—equivalence classes of PQS constituents in the $\hat{\Omega}$-Continuum (Section 1.1.1).

Putting it together: from logic to lab

Diagrammatic reading. A "from logic to lab" diagram (fig. 1.2) maps each corner's ontological label → realized effect → Born truth → concrete measurement, all grounded in the GNS/emergent layer. This ties the three tiers tightly: logical labels at the top, algebraic effects in the middle, and the emergent representation (GNS) at the base, where Born valuations become standard expectations on $\mathcal{H}_{\text{em}}$.

Minimal takeaway.

- The Dialectic Algebra $(\mathsf{Th}, \mathsf{AntiTh}, \mathsf{SynTh}, \mathsf{HoloTh})$ with $\mathbf{D}_{\mathcal{C}}$ and $\Delta_{\mathcal{C}}$ is an ontological blueprint realized as effects with Born-valued truth (Definitions 1.16 and 1.18 and Proposition 1.12).
- SynTh detects *coherent coexistence*: locally via $\frac{1}{\sqrt{2}}\left(|\overset{\circ}{\mathbf{0}}\rangle \pm |\overset{\circ}{\mathbf{1}}\rangle\right)$, globally via Bell/W-like relations (Sections 1.2.8 and 1.2.9).
- HoloTh interrogates *architecture* by controlled holosymmetric evolutions; its status includes typed inapplicability conditions invariant under dualization (Sections 2.3 and 3.8 and eq. (3.36)).
- $\mathcal{A}(\mathcal{C}_d)$ + GNS with $\omega^{\overset{\circ}{\boldsymbol{\Psi}}}$ produce $(\pi_{\overset{\circ}{\boldsymbol{\Psi}}}, \mathcal{H}_{\text{em}}, |\Omega_{\omega^{\overset{\circ}{\boldsymbol{\Psi}}}}\rangle)$, turning all valuations into standard quantum expectations (unique up to unitary equivalence) (Definitions 1.18 and 1.32 and Section 1.4.2).

- The Holo-State unifies scales: locally, it fixes inner products of modal fibres; globally, it shapes the emergent spacetime $\mathcal{M}_{\text{em}}^{(\overset{\circ}{\mathbf{\Psi}})}$ (Proposition 1.30 and Section 4.1).

Compact pipeline (ontology ⇒ laboratory):

$$\begin{aligned} &\text{Fourfold Dialectic (ontology) } (\textit{Definition } 1.16) \\ &\Rightarrow\ \mathcal{A}(\mathcal{C}_d) \text{ (operational language) } (\textit{Definition } 1.32) \\ &\Rightarrow\ |\overset{\circ}{\mathbf{\Psi}}\rangle \xrightarrow{\text{GNS}} (\mathcal{H}_{\text{em}}, \pi_{\overset{\circ}{\mathbf{\Psi}}}) \ (\textit{Section } 1.4.2) \end{aligned}$$

This is exactly the *Blueprint* → *Actualization* logic that Chapter 1 establishes, with duality-driven self-similarity (Proposition 1.12) and a Holo-State that simultaneously selects spacetime (Section 4.1) and fixes local quantum kinematics (Proposition 1.30).

Chapter 2: The Fourfold Symmetry

Chapter 2 operationalises the Fourfold programme by giving it a concrete quantum realisation [22, 29]. Each Primordial Quantum Singularity (PQS) carries a modal fibre $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)} \cong \mathbb{C}^4$ in which Thesis/Symmetry and Antithesis/Antisymmetry are realised as $|e_1\rangle, |e_2\rangle$ spanning the $(0,1)$-sector, with generic states constrained accordingly (Section 2.1 and Definition 2.1) [30, 20]. The local involutive duality $D^{(\mathfrak{B}_x)}$ swaps these modes and lifts diagonally to a global involution $\hat{D}_{\text{PQS}}^{(n)} = \bigotimes_i D^{(\mathfrak{B}_{x_i})}$, encoding a fibrewise Thesis–Antithesis exchange across an n-site configuration (Definition 2.7, Section 2.1.3, and eq. (2.12)) [27, 28, 29]. Parasymmetry (Synthesis) is then given a precise operator meaning: locally by working within the $(0,1)$-sector, and globally via rank-1 projectors onto bipartite entangled states that realise synthesis and its orthogonal dissonance (Section 2.2, Definition 2.23, and eq. (2.58)) [22]. This extends to the multipartite W-state together with a canonically orthogonalised anti-parasymmetric complement, yielding a complete two-dimensional "coherence plane" in the single-excitation sector (Definitions 2.33 and 2.37 and eq. (2.59)) [22, 29]. On this plane, orthogonality, normalization, and span-complementarity are proved explicitly, resolving parity subtleties for odd n.

Layered parasymmetric dynamics. Dynamics are organised in layers [22, 29]. Pairwise and modal-class generators distinguish and drive transitions between parasymmetric synthesis and anti-parasymmetric dissonance [46, 30]; from these, several families of unitaries are built: (i) block-factorised evolutions where disjoint regions follow local/pairwise/global rules independently [22]; (ii) "zonal" evolutions with an active dynamical region and a static complement [47, 48]; and (iii) fully coupled, graph-encoded global evolutions that are not factorisable [49, 39]. A "unified" generator then aggregates local, pairwise, multipartite, and excitation-exchange terms, producing the most general parasymmetric unitary permitted by the scheme (Definitions 2.42, 2.43, 2.54, 2.59, 2.66, 2.70 and 2.78 and eq. (2.56)) [30, 29].

Holosymmetry as interferential higher-order synthesis. Finally, the chapter introduces holosymmetry as a controlled superposition of entire evolutionary histories [22, 23, 24, 25, 26]. Two disjoint PQS registers A, B are coupled to a control qubit whose computational basis toggles which register undergoes a globally coupled vs. a decoupled parasymmetric evolution; in superposition, these mutually exclusive assignments interfere [22, 47, 48]. Lifted operators $U_{\cdot}^{(A)}, U_{\cdot}^{(B)}$ act selectively on registers, and a controlled holosymmetric unitary $\mathbf{U}_{\text{ctrl}}$ enacts path-conditioned evolutions on the full holosymmetric Hilbert space $\mathcal{H}_{\text{Holo}} = \mathcal{H}_c \otimes \mathcal{H}_{\text{PQS}}^{(n,A)} \otimes \mathcal{H}_{\text{PQS}}^{(n,B)}$, whose dimension 2^{4n+1} reflects the exponential richness of the dual-register construction (Sections 2.3.2 and 2.4.1, Definitions 2.102, 2.103, 2.114 and 2.123, and eq. (2.75)) [22, 30, 29]. Holosymmetry thus appears as higher-order synthesis: interference between different ways the universe may organise its internal coherence [23, 24, 25, 20, 49].

Parasymmetry, Global Duality, and Holosymmetric Interferometry

Local and global parasymmetric sectors. Chapter 2 turns the ontological blueprint for Synthesis/Parasymmetry into concrete Hilbert-space objects and gates. Locally, at each PQS $\mathfrak{B}_{x_i}$, the parasymmetric sector is the distinguished $(0,1)$-modal slice inside the modal fiber $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_{x_i})} \cong \mathbb{C}^4$,

$$\mathcal{H}_{(0,1)}^{(x_i)} = \text{span}\{|\mathring{\mathbf{0}}\rangle_{x_i}, |\mathring{\mathbf{1}}\rangle_{x_i}\} \subset \mathbb{C}^4, \qquad |\mathring{\mathbf{0}}\rangle = \begin{pmatrix}1\\0\\0\\0\end{pmatrix}, \quad |\mathring{\mathbf{1}}\rangle = \begin{pmatrix}0\\1\\0\\0\end{pmatrix},$$

and for n loci the global parasymmetric modal space is $(\mathbb{C}^2)^{\otimes n}$ with $\dim = 2^n$ (Definition 2.16). Within the $(0,1)$-sector, the logical projectors $P_T = |e_1\rangle\langle e_1|$ and $P_A = |e_2\rangle\langle e_2|$ carry Born valuations $v_{\text{Thesis}}(\rho) = \text{Tr}(P_T\rho)$, $v_{\text{Antithesis}}(\rho) = \text{Tr}(P_A\rho)$ exactly as in the prism semantics of Chapter 1 (Definition 1.18).

Parasymmetric projectors and valuations. The local parasymmetric axis is isolated by $P_{\text{para}}^{(x)}$, with $\{P_{\text{para}}^{(x)}, P_{\text{antipara}}^{(x)}\}$ an orthogonal, complete resolution of the $(0,1)$-sector; gauge variants (local phases, site relabelings) are operationally equivalent (Section 2.2.1 and Definition 2.17). Operationally, one probes Parasymmetry by rotating into the Σ_x basis and projecting with $P_{\text{para}}^{(x)}$ (cf. eqs. (2.58) and (2.59)).

Relational Parasymmetry (entanglement). Bipartitely, Parasymmetry "becomes entanglement." The Bell-like state

$$|\Psi_{\text{para}}\rangle_{xx'} = \tfrac{1}{\sqrt{2}}\big(|\mathring{\mathbf{0}}\rangle_x |\mathring{\mathbf{1}}\rangle_{x'} + |\mathring{\mathbf{1}}\rangle_x |\mathring{\mathbf{0}}\rangle_{x'}\big),$$

with projector $P_{\text{para}}^{(xx')} = |\Psi_{\text{para}}\rangle_{xx'}\langle\Psi_{\text{para}}|_{xx'}$, realizes distributed co-alignment that belongs to neither locus individually (Section 2.2.3). Multipartitely, $P_{\text{para}}^{(x_1\ldots x_n)}$ detects W-like global modes with genuine multipartite entanglement and permutation symmetry (cf. eq. (2.59)).

Unified parasymmetric generator. Dynamics are generated by a single parasymmetric unitary

$$U_G^{(\odot)} = \exp\Big(i\,\hat{F}_{\text{para}}^{(\odot)}\Big),$$

where $\hat{F}_{\text{para}}^{(\odot)}$ merges local coherence, pair projectors, global modal-class terms, and inter-site exchange $\sum_{i\neq j} J_{ij}\Big(\hat{\sigma}_{x_i}^{(+)}\hat{\sigma}_{x_j}^{(-)} + \text{h.c.}\Big)$; this yields the most general parasymmetric dynamics admissible in the framework (eq. (2.56)).

Global involutive duality. At scale, the global involutive duality operator acts fiberwise across an n-site configuration,

$$\hat{D}_{\text{PQS}}^{(n)} := \bigotimes_{i=1}^{n} D^{(\mathfrak{B}_{x_i})},$$

coherently exchanging Symmetry $\leftrightarrow$ Antisymmetry on each modal fiber and defining a canonical involutive automorphism of the modal Hilbert bundle (eq. (2.12)).

Holosymmetric gates and interferometry. Holosymmetry is operationalized via controlled/interferometric unitaries that compare full PQS-register pathways:

- *Controlled holosymmetric operator* (Definition 2.114 and eq. (2.75)):

  $$\mathbf{U}_{\text{ctrl}} := |\mathbf{0}\rangle\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} \;+\; |\mathbf{1}\rangle\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G},$$

  block-diagonal in the control computational basis with blocks $\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}$, $\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}$ (Section 2.3.2).

- *Hadamard-conjugated form* (Section 2.3.2):

  $$\mathbf{U}_{\text{Had}} := (H_c \otimes I_{\text{RR}})\,\mathbf{U}_{\text{ctrl}}\,(H_c \otimes I_{\text{RR}}) = |+\rangle\langle+|_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + |-\rangle\langle-|_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}.$$

- *Readout gate* (Section 2.4.1): $\mathbf{U}_{\text{read}} = (U_c^{\text{read}} \otimes I_{\text{RR}})$ with $U_c^{\text{read}} \in \text{SU}(2)$; choosing $U_c^{\text{read}} = H$ projects into the symmetric/antisymmetric basis.

- *Symmetric interferometer* (Definition 2.123):

  $$\mathbf{U}_{\text{holo}}^{(\text{sym})} := \mathbf{U}_{\text{Had}} \cdot \mathbf{U}_{\text{ctrl}} \cdot \mathbf{U}_{\text{Had}},$$

  acting on $|\Psi_{\text{init}}\rangle = |\mathbf{0}\rangle_c \otimes |\Psi_{\text{RR}}\rangle$.

- *Asymmetric generalization* (Definition 2.123): arbitrary preparation (α, β) and readout (γ, δ) bases probe off-diagonal modal structure and "holosymmetric curvature."

Operational picture and invariants. (a) Local Parasymmetry is a repeatable "test & filter" along the Σ_x axis; (b) bipartite and n-partite Parasymmetry are entangled co-alignments detected by $P_{\text{para}}^{(xx')}$ and $P_{\text{para}}^{(x_1\ldots x_n)}$ (cf. eqs. (2.58) and (2.59)); and (c) holosymmetric interferometers $(\mathbf{U}_{\text{ctrl}}, \mathbf{U}_{\text{Had}}, \mathbf{U}_{\text{read}})$ elevate *architecture* to an observable by reading interference on the control. Dynamics that preserve global Parasymmetry are precisely those unitaries U that leave $P_{\text{para}}^{(x_1\ldots x_n)}$ invariant (cf. eq. (2.60)).

Minimal takeaway for Chapter 2.

- Parasymmetry is formalized at three levels: (i) kinematic subspaces and projectors; (ii) a unified generator $U_G^{(\odot)}$ for its dynamics; and (iii) relational (Bell/W) implementations across PQSs (Definition 2.16, eqs. (2.56) and (2.59), and Section 2.2.3).
- Duality is upgraded to a global, fiberwise involution $\hat{D}^{(n)}_{\mathrm{PQS}}$ that coherently swaps Symmetry/Antisymmetry across the configuration (eq. (2.12)).
- Holosymmetry is implemented by controlled and interferometric gates $\mathbf{U}_{\mathrm{ctrl}}$, $\mathbf{U}_{\mathrm{Had}}$, $\mathbf{U}_{\mathrm{read}}$ that compare full dynamical histories via control-qubit interference (Definitions 2.114 and 2.123 and Sections 2.3.2 and 2.4.1).

Chapter 3: The Double-Slit Experiments

Chapter 3 reconceives the double-slit experiment as an interferometer on *histories*: the "slits" are whole, register-wide unitary evolutions on dual PQS arrays, steered by a control qubit through a Controlled Holosymmetric Operator $\mathbf{U}_{\mathrm{ctrl}}$ (Sections 2.3.2 and 3.1, Definition 2.114, and eq. (2.75)) [22, 47, 48, 20]. The symmetric protocol—Hadamard → controlled evolution → Hadamard—yields outcome probabilities

$$p_0 = \tfrac{1}{4}\left\|\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}\right\|^2, \qquad p_1 = \tfrac{1}{4}\left\|\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} - \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}\right\|^2,$$

(evaluated on the prepared register state; cf. Section 3.1.1 and Definition 2.123) [22, 26], so that visibility reduces to the real part of the overlap between the two path-evolved states, turning the control qubit into a calibrated meter of *functional symmetry* between dynamical branches (Section 3.1.2 and Theorem 3.4) [23, 24, 25]. What interferes here are not trajectories but full unitary evolutions; moreover, because each path operator itself encodes parasymmetric couplings, the architecture is recursive—interference of (structured) histories rather than of bare paths (Section 3.1 and eq. (2.56)) [49, 30, 29].

Asymmetric tomography and component expansions. Generalizing the beam-splitters to arbitrary $SU(2)$ rotations produces an *asymmetric* interferometer $\mathbf{U}^{(\mathrm{asym})}_{\mathrm{holo}} = \mathbf{U}_{\mathrm{read}}\mathbf{U}_{\mathrm{ctrl}}\mathbf{U}_{\mathrm{prep}}$ that enables full tomography of the complex overlap $\mathcal{O}$ between histories: preparation (α, β) and readout (γ, δ) tune which quadrature of $\mathcal{O}$ is revealed in p_0, p_1 and hence in the visibility $\mathcal{V}_{\mathrm{asym}} = |p_0 - p_1|$ (Sections 3.1.5 and 3.1.7) [22, 26, 23, 24, 25]. At the operator level, both symmetric and asymmetric circuits admit recursive component expansions whose matrix elements (first- and second-order) act as channel-resolved amplitudes, exposing higher-order "interference of interference" and making the recursion experimentally legible (cf. Sections 3.1.5 and 3.1.8) [30, 29, 22]. In this extended setting, the overlap amplitude $\mathcal{M}$ generalises to $\mathcal{M}(U_{\mathrm{rel}}) = \mathrm{Tr}(\sqrt{\rho_a}\sqrt{\rho_b}\,U_{\mathrm{rel}})$, bounded above by the Uhlmann fidelity, which in turn constrains achievable visibility (Section 3.1.8) [50]. Triadic readouts supply square-sum envelopes over correlated/vacuum/spectral channels, sharpening design rules for selecting and interpreting holosymmetric fringes (Sections 3.1.8 and 12.2) [30].

Two interpretive lenses. First, the "interferometer of universal histories" reads the micro-experiment cosmologically: the particle $\mapsto$ dual-register state, the slits $\mapsto$ complementary evolutions, and the screen $\mapsto$ control-qubit statistics—suggesting that physical laws themselves may appear as stable interference patterns across competing histories (Section 3.1) [51, 20, 49]. Second, *commutator lensing* reframes interference as measured non-commutativity—e.g. $[\mathbf{U}_{\mathrm{Had}}, \mathbf{U}_{\mathrm{ctrl}}]$ in the symmetric case—thereby turning algebraic "twist" into an observable and promoting interference from a phenomenon within fixed laws to a probe of the algebra that organizes those laws (Section 3.3) [22, 30, 29].

Holosymmetric Interferometer and Visibility Bounds

Set-up and core quantity. The double-slit is recast as a holosymmetric interferometer (Section 3.1), with a control qubit toggling between two register-wide evolutions that interfere at readout. The central observable is the visibility

$$\mathcal{V} \;:=\; \frac{I_{\max} - I_{\min}}{I_{\max} + I_{\min}} \in [0,1],$$

which, in the symmetric regime, equals the magnitude of an overlap amplitude $\mathcal{M}$ between two path histories ($\mathcal{V} = |\mathcal{M}|$; Section 3.1.2). Born-valued semantics align with Section 1.2.9.

Overlap amplitudes (pure and mixed histories). For pure histories, $\mathcal{M} = \langle\psi_a|\psi_b\rangle$. For mixed histories ρ_a, ρ_b,

$$\mathcal{M}(U_{\mathrm{rel}}) = \mathrm{Tr}\Big(\sqrt{\rho_a}\sqrt{\rho_b}\,U_{\mathrm{rel}}\Big), \qquad U_{\mathrm{rel}} \in \mathrm{U}(\mathrm{supp}\,\rho_a \cup \mathrm{supp}\,\rho_b),$$

with $U_{\mathrm{rel}} = \mathbb{I}$ in symmetric readout and U_{rel} adjustable by the control in the asymmetric case (Sections 3.1.1 and 3.1.5).

Symmetric interferometer. In the Mach–Zehnder–symmetric regime,

$$\boxed{\mathcal{V} \;=\; |\mathcal{M}| \;=\; |\langle\psi_a|\psi_b\rangle| \;\le\; 1} \qquad (\textit{Section 3.1.1}).$$

Asymmetric interferometer and universal inequality. For general $SU(2)$ preparation/readout on the control qubit, the visibility obeys

$$\boxed{\mathcal{V} \;\le\; |\langle\psi_a|\psi_b\rangle|},$$

with equality restored when the beam-splitter weights equalise path priors (Sections 3.1.5 and 3.1.7). Writing $U_c^{\mathrm{prep}} :\; |0\rangle \mapsto \alpha\,|0\rangle + \beta\,|1\rangle$ and $U_c^{\mathrm{read}} :\; \{|0\rangle, |1\rangle\} \mapsto \{\gamma\,|0\rangle + \delta\,|1\rangle,\; -\overline{\delta}\,|0\rangle + \overline{\gamma}\,|1\rangle\}$, with register-wide evolutions $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$ and $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$, the control probabilities are

$$p_0^{(\mathrm{asym})} = \Big\|\big(\gamma\alpha\,\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} - \delta^*\beta\,\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}\big)\,|\Psi_{\mathrm{RR}}\rangle\Big\|^2,$$
$$p_1^{(\mathrm{asym})} = \Big\|\big(\delta\alpha\,\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + \gamma^*\beta\,\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}\big)\,|\Psi_{\mathrm{RR}}\rangle\Big\|^2,$$

and the asymmetric-visibility definition

$$\mathcal{V}_{\mathrm{asym}} := \Big|p_0^{(\mathrm{asym})} - p_1^{(\mathrm{asym})}\Big|$$

matches Theorem 3.8.

Interferometric visibility bounds. Within the double-slit apparatus of Section 3.1 and with arbitrary control $U_c^{\text{prep}}, U_c^{\text{read}} \in SU(2)$ (thus encompassing symmetric and asymmetric regimes; cf. Sections 3.1.1 and 3.1.5), the following envelopes hold (Section 3.1.8):

1. *Universal "no-extra-control" bound.* If the total fringe is a sum of channel amplitudes $\mathcal{M} = \sum_\chi \mathcal{M}_\chi$, then

$$\boxed{\mathcal{V} \;\le\; \left|\sum_\chi \mathcal{M}_\chi\right| \;\le\; \sum_\chi |\mathcal{M}_\chi|}.$$

2. *Triadically weighted bound.* If readout applies complex weights w_χ with $\sum_\chi |w_\chi|^2 = 1$ (operationally by inserting $\sum_\chi w_\chi P_\chi$; cf. Section 12.2), then

$$\mathcal{M}(w) = \sum_\chi w_\chi \mathcal{M}_\chi, \qquad |\mathcal{M}(w)| \le \left(\sum_\chi |\mathcal{M}_\chi|^2\right)^{1/2},$$

with optimal envelope

$$\boxed{\mathcal{V}_{\max} \;\le\; \left(|\mathcal{M}_{\text{corr}}|^2 + |\mathcal{M}_{\text{vac}}|^2 + |\mathcal{M}_{\text{spec}}|^2\right)^{1/2}}.$$

Here each $\mathcal{M}_\chi$ is the expectation of the effect $E_\chi := P_\chi |\psi_b\rangle\langle\psi_a|$ in the experiment's GNS state (measurement protocol: insert P_χ and scan phase).

Mixed histories and fidelity ceiling. For mixed ρ_a, ρ_b, using the polar decomposition and the Uhlmann fidelity

$$F(\rho_a, \rho_b) = \left(\operatorname{Tr}\sqrt{\sqrt{\rho_a}\,\rho_b\,\sqrt{\rho_a}}\right)^2,$$

one obtains

$$\boxed{\mathcal{V} = |\mathcal{M}(U_{\text{rel}})| \;\le\; \sqrt{F(\rho_a, \rho_b)} \;\le\; 1},$$

with equality achieved by the polar-unitary choice $U_{\text{rel}} = V^\dagger$ of $\sqrt{\rho_a}\sqrt{\rho_b}$ (Section 3.1.8).

Readout design logic. The envelopes are operational: suitable choices of $U_c^{\text{prep}}, U_c^{\text{read}}$ (and, when needed, projectors P_χ) can (i) enforce robust lower bounds on $\mathcal{V}$; (ii) isolate correlated-signal contributions; and (iii) veto parasitic channels via the fidelity ceiling—rendering holosymmetric interference a diagnostic of global dynamics (Sections 3.1.8 and 12.2 and Theorem 3.8).

Minimal takeaway for Chapter 3.

- In the symmetric interferometer, $\mathcal{V} = |\langle\psi_a|\psi_b\rangle|$; under arbitrary $SU(2)$ control, $\mathcal{V} \le |\langle\psi_a|\psi_b\rangle|$ with equality recoverable by design (Sections 3.1.2, 3.1.5 and 3.1.7).
- For mixed histories, the Uhlmann fidelity governs the maximal visibility: $\mathcal{V} \le \sqrt{F(\rho_a, \rho_b)}$ (Section 3.1.8).

- Triadic readouts yield a square-sum envelope over correlated/vacuum/spectral channels, providing a principled best-case visibility strategy (Sections 3.1.8 and 12.2).

Foundations: Relations among Chapters 1, 2, and 3

From dialectic truth-values to interferometric visibilities. Chapter 1 establishes Born-valued semantics for the Fourfold facets—Thesis, Antithesis, Synthesis (Parasymmetry), and Holosymmetry—by assigning to each facet $\square \in \{\mathrm{T}, \mathrm{A}, \mathrm{S}, \mathrm{H}\}$ an effect $E_\square$ with valuation $v_\square(\rho) = \mathrm{Tr}(E_\square \rho)$ (Section 1.2.9 and Definition 1.18) [21, 22]. Chapter 3 realizes these same semantics operationally within a holosymmetric interferometer: the visibility $\mathcal{V}$ is expressed as the magnitude of an overlap amplitude $\mathcal{M}$, and, in the triadic readout, as expectations of channel-resolved effects $E_\chi = P_\chi \, |\psi_b\rangle\langle\psi_a|$ in the experiment's GNS state (Sections 3.1.1 and 3.1.8) [23, 24, 25, 26, 22, 30, 29]. Thus the "prism protocol" of Chapter 1, $\rho \mapsto (v_T, v_A, v_S, v_H)$, maps to Chapter 3's interferometric protocol $\rho \mapsto \mathcal{V}$ through explicit effect insertions and phase scans, with $\mathcal{V}$ bounded by $|\mathcal{M}|$ and further by triadic envelopes (Sections 3.1.8 and 12.2).

GNS actualization: from ontological labels to laboratory readouts. In Chapter 1, the Holo-State $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$ selects a representation $(\pi_{\overset{\circ}{\boldsymbol{\Psi}}}, \mathcal{H}_{\mathrm{em}}, |\Omega_{\omega^{\overset{\circ}{\boldsymbol{\Psi}}}}\rangle)$ via GNS, identifying valuations with ordinary expectations, $v(\rho; E) = \mathrm{Tr}\Big(\rho \, \pi_{\overset{\circ}{\boldsymbol{\Psi}}}(E)\Big)$ (Section 1.4.2 and Definition 1.18) [30, 29]. Chapter 3 uses this identification verbatim when writing $\mathcal{M}$ and $\mathcal{V}$ as expectation values of operators built from register-wide evolutions and selection projectors P_χ (Sections 3.1 and 3.1.8) [22, 30]. For mixed histories, the Uhlmann fidelity ceiling $\mathcal{V} \leq \sqrt{F(\rho_a, \rho_b)}$ is a representation-independent constraint on achievable expectations in the GNS arena (Section 3.1.8) [50].

Duality structure across scales. Chapter 1 formalizes a context-indexed involutive duality $\mathbf{D}_\mathcal{C}$ with status-compatibility and $\mathbf{D}_\mathcal{C}^2 = \mathrm{Id}$ (Definition 1.16). Chapter 2 realizes duality fiberwise at each PQS as $D^{(\mathfrak{B}_x)}$ and lifts it diagonally to $\hat{D}^{(n)}_{\mathrm{PQS}} = \bigotimes_i D^{(\mathfrak{B}_{x_i})}$ (Definition 2.7, Section 2.1.3, and eq. (2.12)). Chapter 3's symmetric interferometer effectively tests functional duality between complementary global evolutions $\hat{\mathbf{U}}^{A_G B_N}_{\mathrm{RR}}$ and $\hat{\mathbf{U}}^{A_N B_G}_{\mathrm{RR}}$: the outcome probabilities

$$p_0 = \tfrac{1}{4}\left\|\hat{\mathbf{U}}^{A_G B_N}_{\mathrm{RR}} + \hat{\mathbf{U}}^{A_N B_G}_{\mathrm{RR}}\right\|^2, \qquad p_1 = \tfrac{1}{4}\left\|\hat{\mathbf{U}}^{A_G B_N}_{\mathrm{RR}} - \hat{\mathbf{U}}^{A_N B_G}_{\mathrm{RR}}\right\|^2$$

encode constructive/destructive dual combinations under the Hadamard conjugation on the control (Section 3.1.1 and Definition 2.123). In this sense, the visibility equals $|\mathcal{M}|$ precisely when the dual branches are balanced by the $SU(2)$ beam splitters, reflecting the involutive exchange symmetry at the level of global histories (Sections 3.1.1 and 3.1.5).

Parasymmetry: from kinematics and generators to interfering histories. Chapter 2 identifies the parasymmetric axis locally by the projector $P^{(x)}_{\mathrm{para}}$ and relationally by Bell/W-like

projectors $P^{(xx')}_{\text{para}}$, $P^{(x_1\ldots x_n)}_{\text{para}}$ (Sections 2.2.1 and 2.2.3, Definition 2.17, and eqs. (2.58) and (2.59)) [22]. It further builds a unified generator

$$U_G^{(\odot)} = \exp\Big(i\,\hat{F}^{(\odot)}_{\text{para}}\Big)$$

aggregating local, pairwise, multipartite, and exchange dynamics (eq. (2.56)) [30, 29]. Chapter 3 treats the "slits" as entire parasymmetry-bearing histories: the global-evolution operators $\hat{\mathbf{U}}^{A_G B_N}_{\text{RR}}$ and $\hat{\mathbf{U}}^{A_N B_G}_{\text{RR}}$ are themselves composed of such generators, so the overlap amplitude $\mathcal{M}$ measures *coherence of coherence*—interference between structured parasymmetric evolutions rather than between featureless paths (Sections 3.1 and 3.1.8) [23, 24, 25, 49].

Holosymmetry: control-gate realization and effect semantics. Chapter 1 introduces holosymmetry as a facet whose operational effect interrogates *architecture* (Definition 1.18) [21, 22]. Chapter 2 supplies the gates: a controlled holosymmetric unitary

$$\mathbf{U}_{\text{ctrl}} = |\mathbf{0}\rangle\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}^{A_G B_N}_{\text{RR}} + |\mathbf{1}\rangle\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}^{A_N B_G}_{\text{RR}}$$

together with its Hadamard-conjugated form and a tunable readout gate $\mathbf{U}_{\text{read}}$ (Definition 2.114, eq. (2.75), and Sections 2.3.2 and 2.4.1) [22, 47, 48]. Chapter 3 composes these into symmetric and asymmetric interferometers, proving $\mathcal{V} = |\mathcal{M}|$ in the symmetric case and $\mathcal{V} \le |\langle\psi_a|\psi_b\rangle|$ in the asymmetric case, with equality recovered by suitable $SU(2)$ preparation/readout (Sections 3.1.2, 3.1.5 and 3.1.7) [26, 23, 24, 25, 22]. In triadic readouts, the insertion of P_χ implements a holosymmetric *selection principle* consistent with the effect-based semantics of Chapter 1 (Sections 3.1.8 and 12.2) [30, 29].

Inner products from the Holo-State vs. overlap amplitudes. Chapter 1 shows that the Holo-State fixes the local inner product on each modal fibre via Fourfold Inter-Modal Correlators (Proposition 1.30); this is the microscopic counterpart of Chapter 3's macroscopic overlap amplitude $\mathcal{M}$ between histories [23, 24, 25, 22]. The identification $\mathcal{V} = |\mathcal{M}|$ in the symmetric interferometer is therefore the direct laboratory manifestation of the state-defined inner-product geometry projected onto the "history" level (Section 3.1.1) [22, 26].

Algebra of Observables and channel decomposition. Chapter 1 constructs $\mathcal{A}(\mathcal{C}_d)$ as the operational language generated by geometric, matter, and symmetry fields (Definition 1.32) [30, 36, 37]. Chapter 3's channel amplitudes $\mathcal{M}_\chi$ and weights w_χ are implemented by inserting projectors/effects $P_\chi \in \mathcal{A}(\mathcal{C}_d)$ along each history, leading to universal and triadically weighted bounds,

$$\mathcal{V} \;\le\; \Big|\sum_\chi \mathcal{M}_\chi\Big| \;\le\; \sum_\chi |\mathcal{M}_\chi|, \qquad \mathcal{V}_{\max} \;\le\; \Big(|\mathcal{M}_{\text{corr}}|^2 + |\mathcal{M}_{\text{vac}}|^2 + |\mathcal{M}_{\text{spec}}|^2\Big)^{1/2},$$

which are algebraic consequences of $\mathcal{A}(\mathcal{C}_d)$-valued effect insertions and Cauchy–Schwarz in the GNS inner product (Section 3.1.8) [30, 29].

Commutator lensing as a probe of algebraic structure. The prism metaphor of Chapter 1 treats logic as structured interrogation; Chapter 3's *commutator lensing* realizes this as a probe of non-commutativity—e.g. $[\mathbf{U}_{\mathrm{Had}}, \mathbf{U}_{\mathrm{ctrl}}]$—thereby converting algebraic "twist" into measurable fringe shifts (Section 3.3) [22, 30, 29]. This aligns with Chapter 2's gate-level architecture and Chapter 1's thesis that algebraic structure precedes and organises phenomenology (Definitions 1.16 and 1.32).

Synthesis. The three chapters form a single pipeline:

$$\begin{aligned} &\text{Fourfold Dialectic (ontology) } (\textit{Definition } 1.16) \\ &\quad \Rightarrow\ \mathcal{A}(\mathcal{C}_d) \text{ (operational language) } (\textit{Definition } 1.32) \\ &\quad \Rightarrow\ |\overset{\circ}{\boldsymbol{\Psi}}\rangle \xrightarrow{\text{GNS}} (\mathcal{H}_{\mathrm{em}}, \pi_{\overset{\circ}{\boldsymbol{\Psi}}}) \ (\textit{Section } 1.4.2), \end{aligned}$$

with Chapter 2 supplying parasymmetric kinematics and holosymmetric gates (Section 2.2.1, eq. (2.56), and Definition 2.114) [22, 47, 48] and Chapter 3 turning these into interferometric visibility functionals and tomographic bounds (Sections 3.1.1, 3.1.5 and 3.1.8) [23, 24, 25, 22, 50]. The end result is a holosymmetric, testable semantics in which the logic of histories is read directly as calibrated visibilities.

Part II: The Continuum

Chapter 4: Modal Fibres and Emergent Hilbert Spaces

Chapter 4 develops the kinematic heart of the programme: a gauge-covariant, operator-valued *Fourfold Inter-Modal Correlator* that couples Symmetry, Antisymmetry, Parasymmetry, and Holosymmetry across loci of Primordial Quantum Singularities and thereby generates the local modal geometry (Definitions 1.16 and 4.9) [30, 36, 37, 25, 24, 23]. Formally, the correlator $\hat{\mathbf{G}}_\Lambda(x,x') = [\hat{G}_\Lambda^{\alpha\beta}(x,x')]_{\alpha,\beta\in\{0,1,p,s\}}$ is defined by imprint–transport–readout channels built from field modes and a parallel transporter $\mathcal{U}$ (Definition 4.9 and Section 4.1.4) [23, 24, 25, 45]; its diagonal entries track modal persistence, while off-diagonals register interference and cross-mode conversion, with the Twofold $\{0,1\}$ block recovering the dyad and the remaining blocks encoding parasymmetric/holosymmetric couplings (Sections 1.2.8 and 4.1.5) [22, 26]. This kernel is not merely descriptive: the inner product on each modal fibre $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ is *forged* from its Holo-State expectation in the coincident limit, so the Hilbert geometry is emergent and state-dependent (Sections 1.4.2 and 4.6) [30, 29, 36]. Parallel transport along $\gamma \subset \mathcal{X}_q$ ensures frame-consistent comparison of modes between loci (Section 4.1.4) [23, 24, 25, 45]. In commuting/decohered regimes the off-blocks carry no independent signal and the Fourfold collapses to the Twofold classical slice, restoring Non-Contradiction and Excluded Middle on their proper domain (Section 1.2.8) [22, 27, 28].

Bundle, grading, and qubit-recovery regimes. Algebraically, the correlator carries a $\mathbb{Z}_2 \times \mathbb{Z}_2$ "color'' grading that preserves degrees under composition and dictates commutator/anti-commutator selection rules among channels, ensuring that dyadic, parasymmetric, and

holosymmetric couplings compose without violating the prismatic architecture (Section 4.1.7) [52, 30]. These fibres assemble into a rank-4 Hermitian bundle over the emergent configuration space $\mathcal{X}_q$, with ordered orthonormal frames $\{|0\rangle_x, |1\rangle_x, |p\rangle_x, |s\rangle_x\}$ chosen only *after* the correlator-defined metric is fixed; the Twofold and (p, s) sectors are each 2-dimensional, and the full kernel appears as a 4×4 matrix in modal indices (and, in the geometry–matter refinement, as a matrix of such matrices) (Section 4.6.1 and Definition 1.32) [45, 36, 37, 30]. A *duality-collapse* regime is singled out: when cross-block expectations $\{0, 1\} \leftrightarrow \{p, s\}$ are suppressed and the Twofold block is non-degenerate, standard qubit kinematics is recovered up to isometry, with Born statistics reproduced on an effective $\mathbb{C}^2$ (Section 4.1.6) [22].

Holosymmetry, sections, and co-construction with the state. Conceptually, Holosymmetry supplies a system-level invariant that calibrates the entire Fourfold without being scrambled by local mixings (Section 3.8), while the modal weights carried by global sections (dialectic quantum fields) propagate covariantly along connections on the bundle (Sections 4.1.4 and 4.6.1) [45, 23, 24, 25]. Thus, modal fibres and their inner products are co-constructed with the universal state via GNS (Section 1.4.2) [30, 29]; qubits arise as special Twofold regimes (Section 4.1.6) [22]; and the correlator functions as the relational kernel that both *measures* and *makes* the Hilbert geometry on which the Fourfold logic is read (Definition 4.9 and Section 4.6) [36, 37].

Correlator-Induced Modal Geometry and Holonomy

GNS layer and represented operators. From the Quantum Holo-State $\omega^{\mathring{\mathbf{\Psi}}}$ on the C^*-algebra $\mathcal{A}(\mathcal{C}_d)$, the GNS construction yields the cyclic triple

$$(\pi_{\mathring{\mathbf{\Psi}}},\ \mathcal{H}_{\mathring{\mathbf{\Psi}}},\ |\Omega_{\omega^{\mathring{\mathbf{\Psi}}}}\rangle), \qquad \mathcal{H}_{\mathrm{em}} \equiv \mathcal{H}_{\mathring{\mathbf{\Psi}}},$$

identifying Born-valued valuations with ordinary expectations $v(\rho; E) = \mathrm{Tr}\Big(\rho\, \pi_{\mathring{\mathbf{\Psi}}}(E)\Big)$ (Section 1.4.2 and Definition 1.18). Hatted symbols denote represented operators on $\mathcal{H}_{\mathrm{em}}$.

Fourfold Inter-Modal Correlator and fibre metric. For each PQS $\mathfrak{B}_x$, the inner product on the modal fibre $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ is *generated* by the Fourfold Inter-Modal Correlator via the coincident-limit expectation

$$\langle \psi_1 | \psi_2 \rangle_x := \lim_{x' \to x} \left\langle \mathring{\mathbf{\Psi}} \middle| \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2] \middle| \mathring{\mathbf{\Psi}} \right\rangle,$$

which, upon positivity and completion, metrises the fibre (Section 4.1.3; cf. Proposition 1.30). Thus $\mathcal{H}_{\mathrm{em}}$ provides the representation, while Holo-State expectations *forge* the modal metrics.

Tensor form and channel structure. Fix $\mathcal{I} = \{0, 1, p, s\}$ and choose a realisation $\mathcal{B}_x = \{|0\rangle_x, |1\rangle_x, |p\rangle_x, |s\rangle_x\}$ after the metric is defined. The correlator is the 4×4 operator-valued kernel $\hat{\mathbf{G}}_\Lambda(x, x') = [\hat{G}_\Lambda^{\alpha\beta}(x, x')]_{\alpha,\beta \in \mathcal{I}}$ with entries

$$\hat{G}_\Lambda^{\alpha\beta}(x, x') := \hat{\mathcal{F}}_\alpha^\dagger(x)\, \mathcal{U}(x \leftarrow x'; \gamma)\, \hat{\mathcal{F}}_\beta(x') \in \mathcal{B}(\mathcal{H}_{\mathrm{em}}),$$

where $\mathcal{U}$ is the gauge-covariant transporter along γ (Section 4.1.5). The top-left Twofold block $\hat{\mathbf{G}}_{(0,1)}$ obeys Hermitian symmetry and gauge covariance; parasymmetric/holosymmetric effects require the full Fourfold kernel (cf. Section 4.1.3).

Parallel transport, curvature, and holonomy. A metric-compatible connection on the modal bundle provides

$$\mathcal{U}(x \leftarrow x'; \gamma) = \mathcal{P}\, \exp\Big(-\int_\gamma \mathcal{A}\Big),$$
$$\mathcal{F} = \mathrm{d}\mathcal{A} + \mathcal{A} \wedge \mathcal{A}.$$

so that flat regions ($\mathcal{F} = 0$) are path-independent, while nonzero curvature yields holonomy. Loop holonomy $\mathrm{Hol}_\ell = \mathcal{P} \exp(-\oint_\ell \mathcal{A})$ defines holosymmetric observables through loop operators (insertion in the s-channel), whose Holo-State expectations operationalise global fibre-bundle geometry (Section 4.1.3).

Hilbert bundle over $\mathcal{X}_q$ and dimensional summary. The fibres $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ assemble into a Hilbert bundle

$$(\mathscr{H}_{\mathrm{mod}}, \pi, \mathcal{X}_q), \qquad \mathscr{H}_{\mathrm{mod}} := \bigsqcup_{x \in \mathcal{X}_q} \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}, \quad \pi^{-1}(x) \cong \mathbb{C}^4,$$

cleanly separating representation space $\mathcal{H}_{\mathrm{em}}$ from modal fibres. Typical sizes: $\dim_{\mathbb{C}} \mathcal{H}_{\mathrm{em}} = \infty$ (usually separable), $\dim_{\mathbb{C}} \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)} = 4$, and $\dim_{\mathbb{C}} \mathcal{H}_{\alpha,x} = 1$ for $\alpha \in \{0, 1, p, s\}$ (cf. Section 4.1.3).

Twofold vs. Fourfold: not permutation symmetry. The dialectic (anti)symmetries are *internal* PQS structures, distinct from the many-body permutation symmetries; likewise $|p\rangle_x$ and $|s\rangle_x$ are independent basis rays (not $|\pm\rangle_x$ superpositions), while the parasymmetric $|+\rangle_x$ defines an *effect* $P_{\mathrm{para}}^{(x)}$ in the Twofold sector (Section 4.1.5).

Qubit relocation, registers, and holomodal computing. When the Twofold sector is dynamically protected and the correlator is block-diagonalisable, a Qubit Reduction $U : \mathbb{C}^2 \hookrightarrow \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ realises effective qubits, and suppressing inter-fibre cross-correlators yields n-site registers $(\mathbb{C}^2)^{\otimes n}$. This both contains the standard qubit model and extends it by native Fourfold controls and effects (cf. Section 4.1.3). The resulting *holomodal* architecture aligns parasymmetric/holosymmetric channels with the POVM toolbox defined on $\mathcal{A}(\mathcal{C}_d)$ (Definition 1.32).

Alignment with Chapters 1 and 2. The Fourfold roles (Thesis/Antithesis/Parasymmetry/Holosymmetry) acquire metric meaning through $\hat{\mathbf{G}}_\Lambda$, realising the "prism for discourse" within the emergent arena (Section 1.2.8). Local involutive duality $D^{(\mathfrak{B}_x)}$ and its global lift $\hat{D}_{\mathrm{PQS}}^{(n)}$ act naturally on the metrised fibres and their registers (Definition 2.7, Section 2.1.3, and eq. (2.12)), while the Twofold sector $\mathcal{H}_{(0,1)}$ embeds canonically inside $\mathbb{C}^4$ (Section 2.2). In this way, Chapter 4 completes the ontology ⇒ algebra ⇒ GNS pipeline (Definitions 1.16 and 1.32 and Section 1.4.2) by deriving a *state-dependent, correlator-forged* modal geometry whose curvature and holonomy become holosymmetric observables.

Chapter 4 systematises the operator-geometric architecture of the $\hat{\Omega}$-Continuum by introducing a fibre bundle of relational readouts $\pi : \mathfrak{F} \to \mathcal{X}_q$ whose fibres split as $\mathcal{F}_x^{\mathrm{dist}} \times \mathcal{F}_x^{\mathrm{time}}$,

packaging a distance triad $\vec{d} = (d_{\text{corr}}, d_{\text{vac}}, d_{\text{spec}})$ and a time (phase-displacement) triad $\vec{t} = (t_{\text{corr}}, t_{\text{vac}}, t_{\text{spec}})$; the construction is smooth and gauge/frame-covariant, with $D(x, x') = \|\vec{d}(x, x')\|_2$ playing a metric-like role while $\vec{t}$ tracks relational phase flow (Definitions 4.599, 4.600 and 4.604) [45, 36, 37, 30]. The triad's axes are defined and compared in a gauge-covariant manner via the Fourfold correlator and its parallel transport, rendering "universal entanglement" an intrinsic, background-free bilocal amplitude that is one leg of the emergent relational geometry/time triad (Sections 4.1.2 and 4.12) [23, 24, 25, 22].

A central technical role is assigned to the Spectral Operator $\hat{D}$ (the differential generator), whose commutator with the Fourfold channels yields the spectral distance/time pair; the spectral-shear time t_{spec} is defined as an ℓ^2 aggregate of imaginary parts of spectral-shear profiles, is a pseudometric, and obeys the duality bound $t_{\text{spec}} \leq d_{\text{spec}}$ with equality for purely imaginary shears (Definition 4.341 and Section 4.12.5) [36, 37, 29]. A rate 1-form furnishes a path-integral "accumulated spectral time," making temporal holonomy and curvature explicit; if $t_{\text{spec}} \equiv 0$ globally, the spectral connection is flat, the spectral distance collapses, and $[\hat{D}, \hat{G}_\Lambda(\cdot)]$ vanishes (in expectation and, under mild hypotheses, as an operator) on the GNS cyclic subspace—i.e., the spectral sector is strongly static (Sections 4.12.5 and 4.12.9) [45, 30]. The vacuum-response time is developed in parallel (with a rate 1-form), interpreting t_{vac} as a background-free analogue of retarded vacuum reaction familiar from QFT in curved backgrounds (Section 4.12.3), conceptually resonant with [53, 51].

Structurally, the chapter distinguishes the kinematic inner product on each modal fibre (complex Hermitian; interference) from a geometric inner product on $\mathcal{F}_x$ (real Euclidean; aggregation of distances), and proves well-posedness and functional independence of the triadic generators (via a Fréchet argument), together with endogenous finiteness: under Spectral Smoothness and Finite Entanglement Capacity, all six observables $d_{\text{corr}}, d_{\text{vac}}, d_{\text{spec}}, t_{\text{corr}}, t_{\text{vac}}, t_{\text{spec}}$ are finite with a state-dependent effective spectral cutoff (Sections 4.9.11 and 4.11.5 and Proposition 4.605) [30, 29, 36].

Finally, the "Fibred Continuum of Primordial Quantum Singularities" is defined: PQS loci are relational-null classes $(\vec{d}, \vec{t}) = \mathbf{0}$ that carry the universal duality code $\overset{\circ}{\Lambda}_{\hat{\Theta}}^{\hat{\Phi}}$; the correlator both induces the local fibre metric (coincident limit) and mediates gauge-covariant comparisons across the continuum, so that ontological (operator-level) entanglement reduces to familiar phenomenological entanglement precisely when the Twofold block decouples (Sections 4.6 and 4.18) [22, 23, 24].

Chapter 4 develops an *intrinsic*, operator-theoretic geometry and chronology on the emergent configuration space $\mathcal{X}_q$. The central ingredients are the Fourfold Inter-Modal Correlator $\hat{G}_\Lambda$ (cf. Definition 4.9), the gauge-covariant transporter $\mathcal{U}(x \leftarrow x'; \gamma)$, and an admissible Spectral Operator $\hat{D}$ (cf. Definition 4.343). Together they define three *axes* of distinguishability—$(d_{\text{corr}}, d_{\text{vac}}, d_{\text{spec}})$—and their temporal complements—$(t_{\text{corr}}, t_{\text{vac}}, t_{\text{spec}})$—which are assembled fibrewise into a relational bundle over $\mathcal{X}_q$ and then into composite, macroscopic notions of distance and time (cf. Sections 4.10.2, 4.11.3, 4.12 and 5.3.8). At the kinematic level this rests on gauge-covariant comparison of modal data across loci via $\mathcal{U}$, ensuring observer-independent channel operators $\hat{G}_\Lambda^{\alpha\beta}(x, x')$.

Chronogeometric fibres and the expectation-level split

For each relational point $x \in \mathcal{X}_q$, the emergent chronogeometric data are organised into the set-level product fibre

$$\boxed{\mathfrak{F}_x := \mathcal{F}_x^{\mathrm{dist}} \times \mathcal{F}_x^{\mathrm{time}}} \tag{1}$$

(cf. Definitions 4.599 and 4.600), with modal coordinates expressed in the Fourfold dialectic basis $\{\, \mathring{\mathbf{0}},\ \mathring{\mathbf{1}},\ \mathring{\mathbf{0}}{+}\mathring{\mathbf{1}},\ \mathring{\mathbf{\Lambda}}_{\mathring{\mathbf{1}}}^{\mathring{\mathbf{0}}} \,\}$ (Fourfold; cf. Definition 4.9).

At the expectation level the chronogeometric tensor splits into orthogonal summands

$$\boxed{\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}} \tag{2}$$

where $\bar{g}_{\mu\nu}$ is sourced by distinguishability data and $\bar{t}_{\mu\nu}$ by temporal-phase data (cf. Definitions 4.599 and 4.600).

The Relational Distinguishability Fibre

As defined earlier (see Definition 4.599), for each $x \in \mathcal{X}_q$ the *Relational Distinguishability Fibre* collects the three, functionally orthogonal axes of distinguishability into a single fibre over x:

$$\mathcal{F}_x^{\mathrm{dist}} \;=\; \Big\{\, \vec{d}(x,x') \;:=\; \big(d_{\mathrm{corr}}(x,x'),\, d_{\mathrm{vac}}(x,x'),\, d_{\mathrm{spec}}(x,x')\big) \in \mathbb{R}^3 \;:\; x' \in \mathcal{X}_q \,\Big\},$$

and the composite (Euclidean) distinguishability is

$$D(x,x') \;:=\; \|\vec{d}(x,x')\|_2.$$

Narratively, d_{corr} probes bilocal coherence via the off-diagonal sector of the Fourfold Inter-Modal Correlator $\hat{G}_\Lambda$ (cf. Definition 4.9); d_{vac} captures the coincident/diagonal contrast—the "vacuum-response" leg—within the same correlator; and d_{spec} measures commutator-induced spectral shear generated by the admissible spectral operator $\hat{D}$ (cf. Definition 4.343). These axes assemble the set-level fibre used throughout the triadic metric construction (cf. Sections 4.10.2 and 4.11.3) and underpin the orthogonality and independence results established later (cf. Sections 4.10.1 and 4.11.5).

The Temporal Flow Fibre

As defined earlier (see Definition 4.600), for each $x \in \mathcal{X}_q$ the *Temporal Flow Fibre* gathers the phase-like temporal complements of the triad into a single fibre over x:

$$\mathcal{F}_x^{\mathrm{time}} \;:=\; \Big\{\, \vec{t}(x,x') \;:=\; \big(t_{\mathrm{corr}}(x,x'),\, t_{\mathrm{vac}}(x,x'),\, t_{\mathrm{spec}}(x,x')\big) \in \mathbb{R}^3 \;:\; x' \in \mathcal{X}_q \,\Big\}.$$

These are *phase-type* temporal quantities (not metrics) that arise from operator-theoretic evolution along admissible transports, and they supply the temporal summand $\bar{t}_{\mu\nu}$ in the orthogonal split $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (cf. eq. (2)), in complement to the distinguishability fibre $\mathcal{F}_x^{\mathrm{dist}}$ (cf. Definition 4.599).

Narratively:

- t_{corr} is the *entanglement phase time* extracted from the argument of the correlator expectation along an admissible path, with unwrapped lift, basic antisymmetry/additivity properties, and a differential phase-rate 1-form (cf. Definitions 4.523 and 4.527, eq. (4.243), Proposition 4.526, and Remark 4.528).
- t_{vac} quantifies the *vacuum-response* (diagonal/coincident sector) via a gauge-covariant rate 1-form whose path integral bounds the net temporal response (cf. Definition 4.537 and Section 4.12.3).
- t_{spec} is the temporal parameter associated with *spectral shear* generated by the admissible spectral operator $\hat{D}$, paired with the self-regularising spectral distance and its observables (cf. Section 4.12.5 and Definitions 4.386 and 4.428).

Together, $\vec{t}$ provides the temporal counterpart to the distinguishability triad, ensuring a coherent, gauge-covariant time structure across $\mathcal{X}_q$ that is functionally aligned with the distance axes used in the composite metric construction (cf. Sections 4.11.3 and 4.12).

The triadic axes of relational distance and time

The continuum's geometry is organised by the *triad*

$$\vec{d}(x,x') = \big(d_{\mathrm{corr}},\, d_{\mathrm{vac}},\, d_{\mathrm{spec}}\big), \qquad \vec{t}(x,x') = \big(t_{\mathrm{corr}},\, t_{\mathrm{vac}},\, t_{\mathrm{spec}}\big),$$

where d_{corr} probes bilocal coherence (off-diagonal sector of $\hat{G}_\Lambda$), d_{vac} probes coincident/diagonal contrasts (vacuum response), and d_{spec} probes commutator-induced spectral shear via $[\hat{D}, \hat{G}_\Lambda]$. The three axes are *functionally orthogonal* and assembled in the Relational Distinguishability Fibre (cf. Section 4.11.3); their specialisation to *emergent proper* quantities is deferred to Section 5.3.8.

Entanglement phase time

As defined earlier (see Definition 4.523), the entanglement phase time along an admissible path γ with nonvanishing correlator expectation $\Gamma_{\mathrm{corr}}(x,x';\gamma)$ is

$$t_{\mathrm{corr}}(x,x';\gamma) := \operatorname{Arg}\, \Gamma_{\mathrm{corr}}(x,x';\gamma) \in (-\pi,\pi], \qquad t_{\mathrm{corr}}(x,x;\cdot) = 0 \quad \text{(cf. eq. (4.243))}.$$

The *unwrapped* phase $\tilde{t}_{\mathrm{corr}} \in \mathbb{R}$ is obtained by continuous lift along node-avoiding homotopies and is unique up to $2\pi\mathbb{Z}$. For later use, we also record the differential phase-rate 1-form (cf. Definition 4.527)

$$\dot{t}_{\mathrm{corr}}(s) = \operatorname{Im}\!\left(\frac{\dot{\Gamma}_{\mathrm{corr}}(s)}{\Gamma_{\mathrm{corr}}(s)}\right), \qquad t_{\mathrm{corr}} = \int \dot{t}_{\mathrm{corr}}\, ds.$$

It enjoys antisymmetry, additivity under concatenation, and loop holonomy (cf. Proposition 4.526). Its transport dependence is captured by the differential phase-rate 1-form

$$\dot{t}_{\mathrm{corr}}(s) = \operatorname{Im}\!\left(\frac{\dot{\Gamma}_{\mathrm{corr}}(s)}{\Gamma_{\mathrm{corr}}(s)}\right), \qquad t_{\mathrm{corr}} = \int \dot{t}_{\mathrm{corr}}\, ds \quad \text{(cf. Definition 4.527)}.$$

Writing $\Gamma_{\mathrm{corr}} = |\Gamma_{\mathrm{corr}}|e^{it_{\mathrm{corr}}}$ exhibits magnitude/phase complementarity: d_{corr} and t_{corr} are respectively the modulus and argument of the same gauge-covariant expectation (cf. Remark 4.528). At nodes where $\Gamma_{\mathrm{corr}} = 0$, employ the reference-admixture regularisation (cf. Remark 4.524 and Definition 4.523).

Vacuum-response time

The vacuum-response time t_{vac} quantifies the retarded internal reaction of the "vacuum" sector (diagonal/coincident data of $\hat{G}_\Lambda$) to modal shifts. Along a C^1 path γ define the gauge-covariant *rate 1-form*

$$\dot{t}_{\mathrm{vac}}(s) := \left[\sum_k \left(\tfrac{d}{ds}\ \mathrm{Im}\, C_k(\gamma(s))\right)^2\right]^{1/2}, \qquad T_{\mathrm{vac}}(x,x';\gamma) := \int_0^1 \dot{t}_{\mathrm{vac}}(s)\, ds,$$

with $t_{\mathrm{vac}}(x,x') \le T_{\mathrm{vac}}(x,x';\gamma)$ and equality for straight-line profiles in ℓ^2 (cf. Definition 4.537 and Section 4.12.3). Physically, t_{vac} extracts the phase-encoded polarisation/gauge-flow response in the coincident sector, rendered intrinsic and background-free via the Holo-State and $\hat{G}_\Lambda$ (cf. Section 4.12.3).

Spectral shear time and distance

The spectral axis is governed by the commutator "shear" with $\hat{D}$. A key result is the *Self-Regularizing Spectral Distance*:

$$d_{\mathrm{spec}}(x,x') := \left(\sum_n w_n^2(x,x')\left|\mathcal{S}_n(x,x')\right|^2\right)^{1/2},$$
$$\mathcal{S}_n(x,x') := \left\langle \overset{\circ}{\boldsymbol{\Psi}}\right| [\hat{D}, \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_n^*, \psi_n]] \left|\overset{\circ}{\boldsymbol{\Psi}}\right\rangle,$$

where w_n are *modal weights* implementing state-conditioned ultraviolet control (cf. Definitions 4.386, 4.392 and 4.428, Axiom 4.415, and Principle 4.1). No external cutoff is needed; finiteness is a property of the operator algebra plus the Holo-State, and d_{spec} is monotone under coarse-graining (Remark after Definition 4.428). The associated temporal parameter t_{spec} is introduced in Section 4.12.5.

Functional orthogonality and Fréchet independence. At the *operator* level, the triad is generated by

$$\mathbf{A}_{\mathrm{corr}}(x,x') := \hat{G}_\Lambda(x,x') \quad \text{(off-diagonal coherence)},$$
$$\mathbf{A}_{\mathrm{vac}}(x,x') := \hat{G}_\Lambda(x,x) - \hat{G}_\Lambda(x',x') \quad \text{(diagonal contrast)},$$
$$\mathbf{A}_{\mathrm{spec}}(x,x') := i[\hat{D}, \hat{G}_\Lambda(x,x')] \quad \text{(spectral shear)}.$$

Functional *orthogonality* under the emergent fibre inner product is established in Sections 4.10.1 and 4.10.2, while a Fréchet-differential argument proves *functional independence* (cf. Section 4.11.5).

Kinematic vs geometric inner products. Chapter 4 distinguishes the *modal* inner product on $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}$ (Hermitian; fixes local amplitudes/interference) from the *fibre* inner product on $\mathcal{F}_x$ (Euclidean; aggregates distance triples into the composite $D(x, x')$), aligning first-order quantum kinematics with second-order emergent geometry (cf. table 4.3).

State-conditioned modal weight & endogenous regularization. As formalised in Section 4.9.7, the spectral leg weights each modal contribution by a *state-conditioned* factor $w_n(x, x')$ built from the $\hat{D}$-commutator "shear" of the bilocal channel $\hat{G}_\Lambda$ and evaluated in the Holo-State $|\mathring{\mathbf{\Psi}}\rangle$ [36, 37, 30, 29]. A convenient (Hilbert–Schmidt) realisation used in the chapter is

$$w_n^2(x, x') \;:=\; 1 + \Big\| \Big[\hat{D},\ \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_n^*, \psi_n]\Big] \Big\|^2_{\text{HS},\,\mathring{\mathbf{\Psi}}},$$

so the weights are fixed endogenously by the operator algebra and the Holo-State [30, 29, 36]. The underlying shear observable employed throughout is

$$\mathcal{S}_n(x, x') \;:=\; \Big\langle \mathring{\mathbf{\Psi}} \Big| \Big[\hat{D},\ \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_n^*, \psi_n]\Big] \Big| \mathring{\mathbf{\Psi}} \Big\rangle,$$

see Section 4.9.3 and Definition 4.341 [36, 37]. This construction is tied to the PQS coherence/IR scale and to the "effective spectral cutoff" emerging from Spectral/Relational Smoothness and Finite Entanglement Capacity, i.e. from the same endogenous mechanism rather than any external regulator (cf. the discussion surrounding Section 4.9.7) [37, 29].

Endogenous finiteness of spectral distance. With w_n fixed as above and $\mathcal{S}_n$ the spectral shear, Chapter 5 proves that the weighted spectral sum defining the spectral-axis distance is finite *without* external regularisers; the convergence and its stability under coarse-graining follow solely from the state-conditioned smoothness/growth encoded in w_n [36, 37, 30, 29]. This is the "endogenous finiteness of spectral distance" in Section 4.9.11.

Principle of Coherent Regularization. The chapter imposes a sharp rule (see Section 4.9.13): incoherent, externally imposed cutoffs are disallowed. Admissible regularisation must be implemented *coherently* via the modal weights determined by $|\mathring{\mathbf{\Psi}}\rangle$ [30, 29]. In particular, the weights obey a superpolynomial decay in the spectral index/eigenvalue (the Spectral/Relational Smoothness condition), which suffices for the endogenous finiteness above and renders the construction scheme-independent [36, 37].

The self-regularizing geometry. Combining coherent, state-conditioned weighting with endogenous finiteness yields a *self-regularizing* UV picture: the triadic geometry—and in particular the spectral leg—is finite and well-defined purely from $\hat{G}_\Lambda$, $\hat{D}$, and $|\mathring{\mathbf{\Psi}}\rangle$ [36, 37, 30, 29]. This is summarised in Section 4.9.14, and explicitly linked back to Sections 4.9.7, 4.9.11 and 4.9.13.

A relational framework for infrared (IR) regularization. Beyond the UV, Chapter 4 develops a *relational* IR scheme (cf. Section 4.9.15): long-distance behaviour is windowed by relational constructs along paths γ using the gauge-covariant transporter $\mathcal{U}(x \leftarrow x'; \gamma)$, not by background momenta [45, 23, 24, 25]. An IR coherence scale (defined in that section) limits the accumulation of shear/coherence along γ and controls correlator tails [53, 51].

Reciprocal Resolution Limits. Within this framework, Section 4.9.15 establishes reciprocal bounds linking spectral-axis distinguishability and effective spatial/temporal resolution: small spectral distinguishability enforces a lower bound on macroscopic resolving power, and conversely, coarse resolution bounds spectral sensitivity [54]. A "band-limited data" corollary captures the practical coarse-measurement regime [55].

State-dependent spectral gapping. Finally, Section 4.9.15 proves the existence (on compact $\mathcal{X}_q$, for a dense set of Holo-States) of a positive, *state-dependent* commutator gap $\Delta\big(|\mathring{\mathbf{\Psi}}\rangle\big) > 0$ for the $\hat{D}$–$\hat{G}_\Lambda$ dynamics. The gap is intrinsic (state-conditioned algebraic property), stable under small deformations of $|\mathring{\mathbf{\Psi}}\rangle$ and of the local algebra, and enjoys topological protection in the presence of nontrivial bundles—feeding back into the relational IR picture by enforcing coherence windows at large separations [36, 30, 29, 45].

Placement in the triad. All of the above sits inside the triadic kinematics

$$\vec{d} = (d_{\text{corr}}, d_{\text{vac}}, d_{\text{spec}}), \qquad \vec{t} = (t_{\text{corr}}, t_{\text{vac}}, t_{\text{spec}}),$$

with the spectral leg driven by $\hat{D}$–shear and assembled, alongside the correlational and vacuum legs, into the Relational Distinguishability Fibre (cf. Section 4.11.3) and its temporal counterpart (cf. Section 4.12).

Emergence and regularity of the configuration space $\mathcal{X}_q$. The relational quotient $\pi_q : \mathcal{C}_d \to \mathcal{X}_q := \mathcal{C}_d/\sim$ (see eq. (4.322)) furnishes $\mathcal{X}_q$ with a unique smooth structure [45, 30, 29, 36, 37]. A regular foliation by leaves $\{\mathfrak{B}_x\}$ of locally constant dimension yields transversal slices $S_x \subset \mathcal{C}_d$ for which $\pi_q|_{S_x}$ are charts with full-rank differentials; thus π_q is a smooth submersion [45, 43, 44]. Assuming $\dim \mathfrak{B}_x = m$ locally constant, $\dim \mathcal{X}_q = d - m$ [45]. Because the relation set R is a closed embedded submanifold and the leaves are uniformly localised (cf. eq. (4.303)), distinct equivalence classes admit disjoint saturated neighbourhoods, so $\mathcal{X}_q$ is Hausdorff; second-countability descends from $\mathcal{C}_d$ via openness of π_q and a countable slice basis [45, 29]. Cardinality is preserved: if $|\mathcal{C}_d| = \mathfrak{c}$, then $|\mathcal{X}_q| = \mathfrak{c}$ (cf. Section 4.19.4).

PQS locus, null condition, and factorization. A PQS locus is characterised by the *relational null condition* $\vec{d} = \vec{t} = \mathbf{0}$ (together with vanishing of complementary components), pinpointing the "zero" of the emergent metric–temporal structure (cf. Section 4.9) [36, 37]. Global non-factorizability of the Holo-State underwrites connected chronogeometry and nontrivial holosymmetric interference; exact factorization would force a block structure that trivialises cross-sector functionals and the axis-specific temporal parameters [30, 29]. A precise "FAPP" statement shows how subsystems can behave autonomously when trace-distance corrections lie

below experimental resolution, without violating global non-factorizability (cf. Theorem 4.314, Proposition 4.312, and Sections 4.12 and 5.3.8) [22, 45, 43, 44].

Ontological vs phenomenological entanglement and qubit reduction. The chapter separates *ontological* entanglement (default correlator-level coherence in the $\hat{\Omega}$–Continuum) from *phenomenological* entanglement in fixed tensor-product arenas. When the Fourfold's Twofold block decouples and cross-correlators are suppressed, one recovers $(\mathbb{C}^2)^{\otimes n}$ registers and standard entangling gates; in this regime the IQR machinery reduces to familiar quantum-information dynamics (cf. Proposition 4.300 and Corollary 4.302).

Conceptual synthesis & minimal takeaway for Chapter 4.

1. **Intrinsic tri-axial atlas.** The triads $\vec{d}$ and $\vec{t}$ define an *intrinsic atlas* of distinguishability and temporal flow on $\mathcal{X}_q$, anchored in gauge-covariant, state-dependent expectations of $\hat{G}_\Lambda$ and $[\hat{D}, \hat{G}_\Lambda]$. The emergent representation $\mathcal{H}_{\text{em}}$ is provided by GNS, and the modal fibre metrics are *derived* from Fourfold correlator expectations (Sections 1.4.2 and 4.1.3).

2. **Self-regularising spectral leg.** The spectral leg is *self-regularising*: finiteness and scheme-independence arise from modal weights and Holo-State conditioning rather than external cutoffs (cf. Definition 4.428 and Remark thereafter).

3. **Geometry, holonomy, and operational closure.** Gauge-covariant transport and curvature of the correlator produce holonomy; loop operators render this geometry operational in the holosymmetric channel, knitting local rules to global structure through the composite tensor $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (Section 4.1.3).

4. **Factorisation and protected sectors.** Exact factorisation is obstructive; approximate (FAPP) factorisation explains emergent autonomy without abandoning the global, correlator-driven geometry and time. Twofold qubit sectors arise as protected, block-diagonalisable limits of the Fourfold structure; registers $(\mathbb{C}^2)^{\otimes n}$ are special cases within a holomodal, correlator-driven architecture (Section 4.1.5 and Definition 1.32).

Altogether, Chapter 4 shows how the Fourfold logic (Definition 4.9), together with the GNS/emergent construction (Section 1.4.2) and the admissible spectral operator $\hat{D}$, yields a concrete, tri-axial metric–temporal calculus on $\mathcal{X}_q$ [30, 29, 36, 37]. *Distances* arise from correlator magnitude, vacuum contrast, and spectral shear; *times* from entanglement phase, vacuum response, and spectral-shear rates—each axis gauge-covariant, functionally orthogonal, and tied to specific operator features of $\hat{G}_\Lambda$ and $[\hat{D}, \hat{G}_\Lambda]$ [23, 24, 25, 22]. The spectral leg is *self-regularising* (no external cutoffs), with finiteness controlled by modal weights and Holo-State conditioning (cf. Definition 4.428) [36, 37, 30, 29]; gauge-covariant transport and curvature of the correlator produce holonomy and close operationally via loop operators in the holosymmetric channel (Section 4.1.3) [45, 23, 24, 25]. The kinematic modal product and the geometric fibre product cohere local interference rules with global metric structure through the composite tensor $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (Section 4.1.3) [30, 29, 45].

In this calculus, exact factorisation is generically obstructed; nevertheless, approximate (FAPP) factorisation accounts for emergent autonomy without relinquishing the correlator-driven global geometry and time [30, 29, 22]. Twofold qubit sectors appear as protected, block-diagonalisable limits of the Fourfold architecture, with registers $(\mathbb{C}^2)^{\otimes n}$ as special cases within the holomodal framework (Section 4.1.5 and Definition 1.32) [22]. The relational bundle $\mathfrak{F}_x = \mathcal{F}_x^{\text{dist}} \times \mathcal{F}_x^{\text{time}}$ with Fourfold-dialectic coordinates provides the canonical staging for these constructions and for their specialisation to emergent proper quantities (Section 5.3.8) [45, 36, 37].

In plain terms, Chapter 4 builds the **ruler** and the **clock** out of correlations themselves. Starting from the Fourfold inter-modal correlator $\hat{G}_\Lambda$ (Definition 4.9) and an admissible spectral operator $\hat{D}$, it uses the GNS construction to obtain an emergent representation $\mathcal{H}_{\text{em}}$ (Section 1.4.2) and then extracts two coordinated triads: one that measures distinguishability (three "distance" modes) and one that measures temporal flow (three "time" modes) [30, 29, 36, 37]. Those six modes—defined by gauge-covariant, state-dependent expectations of $\hat{G}_\Lambda$ and its spectral commutator $[\hat{D}, \hat{G}_\Lambda]$—form a tri-axial, background-free chronogeometry on the quantum configuration space $\mathcal{X}_q$ [23, 24, 25, 22]. The result is not merely formal: parallel transport, curvature, and loop operators make the geometry operational (one can "close a loop" and read off holonomy) in the holosymmetric channel (Section 4.1.3), while the spectral leg neatly self-regularises without external cutoffs (cf. Definition 4.428 and Remark thereafter) [45, 36, 37].

It also explains how local, apparently autonomous subsystems emerge without breaking the global, correlator-driven picture. Exact factorisation is generically obstructed, but approximate (FAPP) factorisation holds widely enough to recover familiar dynamics; along the same lines, protected twofold qubit sectors appear as block-diagonalisable limits inside the Fourfold architecture (Section 4.1.5 and Definition 1.32) [22, 30, 29]. All of this is organised in a relational bundle $\mathfrak{F}_x = \mathcal{F}_x^{\text{dist}} \times \mathcal{F}_x^{\text{time}}$ that stitches local interference rules to global metric structure via the composite tensor $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (Section 4.1.3), and sets the stage for defining emergent proper distance and proper time (Section 5.3.8) [45, 36, 37].

Chapter 5: The Metric and Temporal Tensors

Chapter 5 develops the operator–geometric heart of the programme. First, it defines an *operator-valued emergent metric* by taking the Fréchet second variation (in the strong–operator topology) of the composite triadic distance D^2 between PQS loci (cf. Section 4.11.3 and Definition 5.10), yielding a positive, Hermitian field $\hat{g}_{\mu\nu}(x)$ whose classical avatar $\bar{g}_{\mu\nu}(x) = \langle \overset{\circ}{\mathbf{\Psi}} | \hat{g}_{\mu\nu}(x) | \overset{\circ}{\mathbf{\Psi}} \rangle$ governs the expectation-level geometry (cf. Definitions 5.1 and 5.10 and Section 5.1) [36, 37, 30, 29]. Ultra-locality on spectrally regulated fibres and gauge covariance are made explicit (cf. eqs. (4.11) and (4.303) and Section 4.19.1) [45, 36], while positivity ensures a viable seed for distances and curvature in the emergent configuration space $\mathcal{X}_q$ (cf. Section 5.1) [30, 29].

In parallel, the chapter introduces the *operator-valued temporal curvature tensor* by the same second-variation construction applied to a composite temporal functional T^2, producing $\hat{t}_{\mu\nu}(x)$ with a canonical triadic split (correlational, vacuum, spectral) and with ultra-sub-

Planckian support and gauge covariance (cf. Definitions 5.17 and 5.54 and eq. (4.303)) [36, 37, 30]. Its expectation $\bar{t}_{\mu\nu}(x) = \langle \overset{\circ}{\boldsymbol{\Psi}} | \hat{t}_{\mu\nu}(x) | \overset{\circ}{\boldsymbol{\Psi}} \rangle$ encodes curvature of quantum phase flow (cf. Definition 5.22 and Section 5.1) [23, 24, 25]. Conceptually, $\mathfrak{F}_x = \mathcal{F}_x^{\text{dist}} \oplus \mathcal{F}_x^{\text{time}}$ underwrites a *relational duality* (cf. eq. (4.302) and Section 1.2.9): distance and phase arise as orthogonal, independent contributors to chronogeometry, unified at the operator level by $\hat{\Sigma}_{\mu\nu} := \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu}$ and at the classical level by $\bar{\Sigma}_{\mu\nu} := \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (cf. Definitions 5.25 and 5.26) [45, 36]. The text distils the physical content of $\bar{\Sigma}_{\mu\nu}$: it is not a Riemannian metric but a total *chronogeometric* response field, from which both distinguishability ("how far apart") and phase-flow ("how out of phase") of nearby configurations can be read (cf. Section 5.1) [36, 37].

From this vantage, causal structure becomes two-layered (cf. Section 5.4): a *hard* kinematic boundary—the effective light cone—emerges from the symmetrized Lorentzian tensor $\bar{g}_{\mu\nu}^{\text{Lor}} := \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}^{\text{sym}}$ (cf. Section 5.4.1 and Definition 5.122) [45], while a *soft* quantum texture inside the cone is governed by $\bar{t}_{\mu\nu}$, potentially including torsion-like effects from its antisymmetric sector and rendering causality state-dependent and dynamical (cf. Section 5.1) [56, 20]. The classical GR limit is recovered by a coarse-graining that discards $\bar{t}_{\mu\nu}$, projecting to a pseudo-Riemannian manifold $\mathcal{M}_{\text{geom}}$ determined solely by $\bar{g}_{\mu\nu}$ (cf. Section 5.3.12) [45, 43, 44]. Operationally, geodesics of $\bar{g}^{\text{Lor}}$ need not coincide with *paths of maximal coherence*, and quantum processes can exhibit directional decoherence anisotropies—diagnostics for experimental tests (cf. Sections 4.12, 4.12.3 and 4.12.5, Definition 4.523, and eq. (4.243)) [23, 24, 25, 22, 53, 51]—before closing with concrete avenues: coupled field equations for $(\bar{g}, \bar{t})$, Bianchi-type identities, invariant diagnostics, and interferometric probes of the temporal sector (cf. Sections 5.1 and 5.4) [45, 22].

Chapter 5 lifts the *relational fibre data* of Chapter 5 into *operator–valued tensors* on $\mathcal{X}_q$—an operator–valued metric $\hat{g}_{\mu\nu}$ and an operator–valued temporal tensor $\hat{t}_{\mu\nu}$—and then takes Born expectations to obtain classical tensor fields $\bar{g}_{\mu\nu}$ and $\bar{t}_{\mu\nu}$. These assemble into the *Unified Relational Chronogeometric Tensor* $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$, which supplies the expectation-level chronogeometry on $\mathcal{X}_q$ (cf. Definitions 5.10, 5.22 and 5.26) [30, 29, 36, 37, 45]. The construction is *gauge-covariant*, *ultra-local* on the spectrally regulated sector, and *operationally complete* at second order (cf. eq. (4.303) and Section 4.19.1) [36, 37, 30, 29, 45].

From fibre to tensors: the $\rightsquigarrow$ mapping. Orthogonal fibre structure. On the regulated sector (cf. eq. (4.303)), the relational bundle fibre at x, $\mathfrak{F}_x = \mathcal{F}_x^{\text{dist}} \times \mathcal{F}_x^{\text{time}}$ (cf. eq. (4.302)), is treated as a Hilbert direct sum

$$\mathfrak{F}_x \;\cong\; \mathcal{F}_x^{\text{dist}} \oplus \mathcal{F}_x^{\text{time}},$$

with the Fourfold inner product from Section 1.2.9. This orthogonality permits the formation of quadratic functionals and the taking of second variations to define $\hat{g}_{\mu\nu}$ and $\hat{t}_{\mu\nu}$ (cf. Definitions 5.1 and 5.17).

Second variations of composite norms. The tensors arise as Hessians of sectoral squared norms in the coincidence limit $x' \to x$:

- *Metric part:* from the distinguishability norm $D^2(x, x') = \|\vec{d}(x, x')\|_2^2$ (cf. Definitions 5.1 and 5.10).

- *Temporal part:* from the temporal-flow norm $T^2(x,x') = \|\vec{t}(x,x')\|_2^2$ (cf. Definitions 5.17 and 5.22).

Because $\mathcal{F}_x^{\text{dist}}$ and $\mathcal{F}_x^{\text{time}}$ are orthogonal, the resulting tensors contribute independently to the emergent chronogeometry.

Relational duality (distance vs. phase). At the expectation level, $\bar{g}_{\mu\nu}$ and $\bar{t}_{\mu\nu}$ are smooth, symmetric, real $(0,2)$ tensors on $\mathcal{X}_q$, obtained as Born expectations of the corresponding operator–valued Hessians. They encode complementary aspects of the same fibre data—spatial distinguishability and curvature of quantum phase flow (cf. Section 5.1).

The unified chronogeometry and its decompositions. Unified tensor. The expectation tensors are packaged as

$$\bar{\Sigma}_{\mu\nu}(x) \;:=\; \bar{g}_{\mu\nu}(x) \;+\; \bar{t}_{\mu\nu}(x) \quad \text{(cf. Definition 5.26)}.$$

This is the classical carrier of chronogeometric information in the IQR framework.

Triadic split. Each piece admits the canonical triadic decomposition aligned with the correlation/vacuum/spectral axes (Chapter 5),

$$\bar{g}_{\mu\nu} = \bar{g}^{\text{corr}}_{\mu\nu} + \bar{g}^{\text{vac}}_{\mu\nu} + \bar{g}^{\text{spec}}_{\mu\nu}, \qquad \bar{t}_{\mu\nu} = \bar{t}^{\text{corr}}_{\mu\nu} + \bar{t}^{\text{vac}}_{\mu\nu} + \bar{t}^{\text{spec}}_{\mu\nu} \quad \text{(cf. Definition 5.54)}.$$

This preserves functional orthogonality of the three axes at the level of tensor contributions.

Expectation-level relational manifold. The expectation-level manifold is then

$$\mathcal{M}_{\text{rel}} \;:=\; \big(\mathcal{X}_q,\; \bar{g}_{\mu\nu},\; \bar{t}_{\mu\nu}\big),$$

intermediate between the operator-level quantum manifold $\hat{\mathcal{M}}$ and any macroscopic coarse-graining $\mathcal{M}_{\text{geom}}$ (cf. Definition 5.66).

Why the split $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ is canonical.

- **Gauge covariance & ultra-locality.** The sectoral functionals D^2, T^2 are gauge-invariant and supported at the $\mathcal{O}(\ell_{\text{PQS}})$ scale (cf. eq. (4.303)); hence $\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu}$ are gauge-covariant operators with gauge-invariant expectations, suitable as classical geometry on $\mathcal{X}_q$.
- **Operational sufficiency.** Visibility moduli feed $D^2 \to \bar{g}$; phase-slope data feed $T^2 \to \bar{t}$. At second order these Hessians exhaust the observer-independent, gauge-invariant chronogeometric content encoded by the regulated fibre and its connection (cf. Section 4.19.1 and eq. (4.11)).
- **Minimality with room for extensions.** The diagonal sectorization is the minimal choice consistent with IQR semantics; a regulated, gauge-invariant mixed functional $M(x,x') = \langle \vec{d}, \mathsf{K}\,\vec{t}\rangle$ could be introduced to define a third tensor $\bar{m}_{\mu\nu}$, if required by future physics.

Causality and holosymmetric control. A holosymmetric analysis considers families of Lorentzian branch metrics $\{\bar{g}^{\mathrm{Lor}}_{(k)}\}$ along with temporal profiles $\{\bar{t}_{(k)}\}$ [45]. Chapter 5 defines *holosymmetric causal functionals* on $T_x\mathcal{M}_{\mathrm{geom}}$: a micro-causality test $\mathcal{C}^{\mathrm{micro}}_x$, a reach functional $\mathcal{C}^{\mathrm{reach}}_x$, and a temporal profile $\mathcal{T}^{\mathrm{HS}}_x$, built from $\chi_{\mathrm{causal}}\big[\bar{g}^{\mathrm{Lor}}_{(k)\mu\nu}V^\mu V^\nu\big]$ across branches (cf. Definition 5.123) [45, 30, 29]. From these, the *no–signalling and phase invariance* proposition states: if regions A, B are spacelike in the intersection of branch cones $\mathrm{Spacelike}_\cap$, then local observables commute for all holosymmetric phases (φ, θ), and influence is confined to $J^+_\cup(A)$ (the union of future cones); both $\mathrm{Spacelike}_\cap$ and $J^\pm_\cup$ are invariant under holosymmetric phase choices (cf. Proposition 5.152) [22, 30, 29, 45, 23, 24, 25, 20].

Where Chapter 4 enters the pipeline.

1. **Fibre-level primitives (from Chapter 4).** The relational fibre $\mathfrak{F}_x = \mathcal{F}^{\mathrm{dist}}_x \oplus \mathcal{F}^{\mathrm{time}}_x$ and the coordinated triads $\vec{d}, \vec{t}$ along the corr/vac/spec axes are built from the Fourfold correlator and the GNS/emergent setting (Definition 4.9, Sections 1.4.2 and 4.1.3, and eq. (4.302)).

2. **Sectoral norms (Chapter 5).** These fibres are summarised by the quadratic functionals $D^2 = \|\vec{d}\|_2^2$ and $T^2 = \|\vec{t}\|_2^2$, which package distinguishability and temporal response for second-variation analysis (Section 5.1).

3. **Second variations to tensors.** Taking Hessians of D^2 and T^2 yields operator–valued tensors $\hat{g}_{\mu\nu}$ and $\hat{t}_{\mu\nu}$ (Definitions 5.1 and 5.17); Born expectations give the smooth fields $\bar{g}_{\mu\nu}$ and $\bar{t}_{\mu\nu}$ (Definitions 5.10 and 5.22).

4. **Unified chronogeometry.** The expectation tensors assemble into the Unified Relational Chronogeometric Tensor $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ with its corr/vac/spec triadic split, defining the expectation-level relational manifold $\mathcal{M}_{\mathrm{rel}} = (\mathcal{X}_q, \bar{g}, \bar{t})$ (Definitions 5.26, 5.54 and 5.66).

This chain is *self-regularising* (no external cutoffs): finiteness of the sectoral distances/-times implies finiteness of $\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu}$ and their expectations via modal-weight suppression and commutator bounds (Definition 4.428 and Section 4.9.14). Gauge-covariant transport and its path-ordered form control comparisons along curves and around loops (Section 4.19.1 and eq. (4.11)).

Key properties and guarantees.

- **Orthogonal sectors.** The splitting $\mathfrak{F}_x = \mathcal{F}^{\mathrm{dist}}_x \oplus \mathcal{F}^{\mathrm{time}}_x$ ensures clean second-variation constructions and independent sectoral contributions (eq. (4.302), Section 1.2.9, and Definitions 5.1 and 5.17).

- **Expectation-level tensors.** $\bar{g}_{\mu\nu}$ and $\bar{t}_{\mu\nu}$ arise by Born expectation of the operator Hessians of D^2 and T^2, respectively (Definitions 5.10 and 5.22 and Section 5.1).

- **Unified, triadic structure.** $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ carries the minimal split into metric and temporal parts, each with corr/vac/spec decomposition (Definitions 5.26 and 5.54).

- **Gauge covariance and ultra-locality.** The framework is gauge-covariant and ultra-local on the spectrally regulated sector; mixed contributions can be introduced in a regulated, gauge-invariant way if required (eqs. (4.11) and (4.303) and Sections 4.9.14 and 4.19.1).

- **Causal controls.** Holosymmetric causal functionals yield micro-causality and controlled reach across branch cones; a no–signalling theorem holds on the intersection of branch spacelike cones (Definition 5.123 and Proposition 5.152).

The Continuum: Relational Chronogeometry Across Chapters 4 and 5

Chapter 4 lays the algebraic and kinematic groundwork by introducing the Fourfold Inter-Modal Correlator $\hat{G}_\Lambda$ and the admissible Spectral Operator $\hat{D}$ as gauge-covariant primitives (cf. Definitions 4.9 and 4.343) [36, 37, 30, 29]. Their comparison across loci is organised by the operational connection and its path-ordered transporter $\mathcal{U}(x \leftarrow x'; \gamma)$ (cf. Section 4.19.1 and eq. (4.11)) [45, 36]. Within this framework the channel operators $\hat{G}_\Lambda^{\alpha\beta}(x, x')$ inherit Hermiticity and covariance from the Fourfold structure, providing the observer-independent kinematics upon which the emergent configuration space $\mathcal{X}_q$ and all subsequent geometric constructs are built [30, 29].

Chapter 4 packages these primitives into a fibrewise, tri-axial calculus on $\mathcal{X}_q$. The Relational Distinguishability and Temporal fibres collect the three axes of distinguishability $\vec{d} = (d_{\text{corr}}, d_{\text{vac}}, d_{\text{spec}})$ and their temporal complements $\vec{t} = (t_{\text{corr}}, t_{\text{vac}}, t_{\text{spec}})$ (cf. Definitions 4.599 and 4.600 and Section 4.11.3) [22, 23, 24, 25, 36, 37]. Composite, gauge-covariant notions of distance and time arise from these axes (cf. Sections 4.10.2, 4.12 and 5.3.8) [45]; for example, t_{corr} is fixed by the correlator's argument with a differential phase-rate 1-form and a node regularisation prescription (cf. Definitions 4.523 and 4.527, eq. (4.243), and Remark 4.524) [23, 24, 25]. The chapter further establishes functional orthogonality and Fréchet independence of the generators (cf. Sections 4.10.1 and 4.11.5) [30, 29],å and proves *self-regularization* of the spectral leg via state-conditioned modal weights—yielding endogenous finiteness and a coherent regularisation principle—together with a relational framework for infrared control (cf. Sections 4.9.7, 4.9.11 and 4.9.13 to 4.9.15) [36, 37, 30, 29, 45, 53, 51].

Chapter 5 takes the fibre-level "ruler and clock" of Chapter 4 and turns them into fields that live everywhere on $\mathcal{X}_q$. Technically, it lifts the fibre data to operator–valued tensors by taking second variations (Hessians) of the sectoral squared norms $D^2 = \|\vec{d}\|_2^2$ and $T^2 = \|\vec{t}\|_2^2$, thereby defining an operator-valued metric $\hat{g}_{\mu\nu}$ and an operator-valued temporal tensor $\hat{t}_{\mu\nu}$ (cf. Definitions 5.1 and 5.17) [36, 37, 30, 29]. These constructions are gauge–covariant and ultra-local on the spectrally regulated sector (eq. (4.303); cf. Definition 4.428) [36, 37], while the operational connection (and its path-ordered transport) tells you how to carry the "ruler and clock" along paths and compare them on loops (Section 4.19.1 and eq. (4.11)) [45]. Taking Born expectations turns the operator responses into smooth, symmetric $(0, 2)$ fields $\bar{g}_{\mu\nu}$ and $\bar{t}_{\mu\nu}$ (cf. Definitions 5.10 and 5.22 and Section 5.1) [30, 29]. Because the relational fibres split orthogonally, $\mathfrak{F}_x = \mathcal{F}_x^{\text{dist}} \oplus \mathcal{F}_x^{\text{time}}$, the distance and time sectors contribute independently to the chronogeometry (cf. eq. (4.302) and Section 1.2.9) [23, 24, 25]. The two expectation fields assemble into the Unified Relational Chronogeometric Tensor $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ with a triadic split aligned to the corr/vac/spec axes (cf. Definitions 5.26 and 5.54), defining the expectation-level relational manifold $\mathcal{M}_{\text{rel}} = (\mathcal{X}_q, \bar{g}, \bar{t})$ (cf. Definition 5.66) [45, 36, 37]. Conceptually, $\bar{\Sigma}_{\mu\nu}$

is a *response* field—like a material's stiffness that changes with state—rather than a fixed Riemannian metric (Section 5.1) [36, 37].

On top of this split, Chapter 5 builds causality and phenomenology. A symmetrised Lorentzian combination $\bar{g}^{\mathrm{Lor}}_{\mu\nu} := \bar{g}_{\mu\nu} + \bar{t}^{\mathrm{sym}}_{\mu\nu}$ draws the kinematic boundary—the effective light cone—while $\bar{t}_{\mu\nu}$ sets the state-dependent "weather" inside the cone (cf. Sections 5.4 and 5.4.1 and Definition 5.122) [45, 20]. Holosymmetric causal functionals and a no-signalling proposition formalise micro-causality and controlled reach across branch cones (cf. Definition 5.123 and Proposition 5.152) [22, 30, 29]. In the macroscopic limit, neglecting $\bar{t}_{\mu\nu}$ recovers the pseudo-Riemannian manifold $\mathcal{M}_{\mathrm{geom}}$ determined by $\bar{g}_{\mu\nu}$ (cf. Section 5.3.12) [43, 44, 45]. Operationally, geodesics of $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ need not follow paths of maximal coherence, suggesting interferometric diagnostics tied to the triadic time structure and phase/response functionals (Sections 4.12, 4.12.3 and 4.12.5, Definition 4.523, and eq. (4.243)) [23, 24, 25, 22, 53, 51]. In this way, the pipeline that began with the Fourfold correlator and operational transport of Chapter 4 (Sections 1.4.2 and 4.1.3) culminates in Chapter 5's expectation-level chronogeometric and causal structure [30, 29, 36, 37, 45].

Part III: Laws, States and Fundamental Fields

Chapter 6: The Gaiamiti and Tantrakyma Fields

Unified correlator and transport. First, Chapter 6 puts the Fourfold/GNS infrastructure of Chapters 4–5 to work by declaring the *Unified Inter-Modal Correlator* $\hat{\mathbf{G}}_\Lambda(x, x')$ as a 2×2 matrix over the two field channels $\mathbf{A}, \mathbf{B} \in \{\Phi, \Theta\}$ (Gaiamiti and Tantrakyma). Each block $\hat{G}_{\mathbf{AB}}(x, x')$ is expanded in a mode basis $\{\psi^{(\mathbf{A})}_k\}$ and made gauge-covariant by an explicit path-ordered transport $U(x \leftarrow x')$ built from the operational connection; this locks the correlator to the relational parallel transport introduced earlier (Definition 6.4, Section 4.19.1, and eq. (4.11)) [45]. The definition spells out the mode sum and U precisely, so the whole object is state- and connection-covariant from the outset [30, 29, 36, 37].

Relational Levi–Civita calculus on $(\mathcal{X}_q, \bar{\Sigma})$. Second, it *upgrades calculus on* $(\mathcal{X}_q, \bar{\Sigma})$ by defining the *Relational Levi–Civita Connection* $\nabla^{(\mathrm{rel})}$ whose Christoffels are computed from the *composite* chronogeometric tensor $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (not just $\bar{g}$). The formula for $\Gamma^\lambda_{\mu\nu}[\bar{\Sigma}]$ and the induced covariant derivatives on vectors/covectors are given explicitly (Definition 6.7 and eq. (6.22)) [45]. The text is careful about scope: $\nabla^{(\mathrm{rel})}$ lives on the emergent configuration space $\mathcal{X}_q$ and is applied to smooth tensor fields there—the "hierarchy of manifolds" paragraph makes that domain separation exact (Proposition 4.625) [30, 29]. This is what lets the chapter talk cleanly about geodesics and curvature *of the chronogeometry* rather than only of $\bar{g}$.

Measure, action, and dynamical fields. Third, it *chooses the integration measure and builds dynamics*. The *Relational Volume Element* is fixed as

$$d\mu(x) \;=\; \sqrt{-\det \bar{g}^{\mathrm{Lor}}(x)}\, d^d x, \qquad \bar{g}^{\mathrm{Lor}}_{\mu\nu} \;:=\; \bar{g}_{\mu\nu} + \bar{t}^{\mathrm{sym}}_{\mu\nu},$$

so the temporal response tensor $\bar{t}_{\mu\nu}$ affects the measure already at the expectation level (Definition 6.33) [45]. This choice is then used to write action integrals and energy functionals, ensuring that *measure, connection, and correlators* are mutually consistent on the same background-free geometry (Definitions 6.4 and 6.7 and Section 4.19.1) [36, 30, 29]. Concretely, the chapter postulates an *operator-valued Lagrangian density* for the *Dynamical Gaiamiti Field* $\hat{\mathbf{\Phi}}(x)$—treated as a section of $\mathrm{End}(\mathcal{H}_{\mathrm{em}}) \otimes T\mathcal{M}_{\mathrm{vac}}$—with a total covariant derivative (geometric + internal) and an operator-valued inner product g_T on $T\mathcal{M}_{\mathrm{vac}}$ (Section 6.5 and Definition 6.172) [36, 37, 45]. This makes the field-theory part explicitly "relational": both kinematics (∇) and the bundle the field lives in depend on the emergent structures of Chapters 4–5.

Relational Hamiltonian and vacuum. Finally, it *defines the Relational Hamiltonian* $\hat{H}[|\mathbf{\Psi}\rangle]$ by integrating the traces of the correlator blocks against the relational measure, with an explicit split into $\hat{H}_{\Phi}$, $\hat{H}_{\Theta}$, and a mixed $\hat{H}_{\Phi\Theta}$ (Definition 6.54 and eq. (6.109)) [30, 29]. This makes the *state-dependence* of $\hat{H}$ a principled, background-independent feature (since the measure $\sqrt{-\det \bar{g}^{\mathrm{Lor}(\mathbf{\Psi})}(x)}$ is itself state-conditioned, cf. Definition 6.33) [36, 37, 45]. The *Vacuum Manifold* $\mathcal{M}_{\mathrm{vac}}$ is then identified spectrally as the ground-state eigenspace of $\hat{H}$, closing the loop from correlators → geometry → dynamics → vacua (Section 6.1.17 and Definition 6.61) [30, 29]. The text emphasises this non-linear, state-determined eigenproblem as the core "solution concept" for the theory [29].

Compact synthesis. Put compactly: Chapter 6 formalises the *objects of motion on the relational manifold*—a gauge-covariant correlator matrix $\hat{\mathbf{G}}_{\Lambda}$ tying fields to transport, a Levi–Civita calculus built from $\bar{\Sigma}$, a volume form that already "knows" about $\bar{t}$, an operator-valued Lagrangian for $\hat{\mathbf{\Phi}}$, and a state-dependent Hamiltonian whose spectrum *defines* the vacuum geometry [36, 37, 30, 29, 45]. All of these are written so the same composite tensor $\bar{\Sigma}$ plays the starring role in transport ($\Gamma[\bar{\Sigma}]$), measure ($\sqrt{-\det \bar{g}^{\mathrm{Lor}}}$), and energy (via $\hat{H}[|\mathbf{\Psi}\rangle]$), keeping the framework consistently background-free and operational [45, 36, 37].

The Nature of the Fundamental Fields. In the IQR framework, the universe's content is posited to arise from a structured singularity $\hat{\mathbf{\Omega}}(x) = \overset{\circ}{\mathbf{\Lambda}}_{\hat{\mathbf{\Phi}}\hat{\mathbf{\Theta}}}(x)$, evaluated at each locus x of a foundational continuum (Section 6.1) [20]. Geometry (*Gaiamiti*) and matter-like content (*Tantrakyma*) co-arise from this singular source and are explicitly packaged as a single generalized field doublet $\hat{\mathbf{F}}(x) = (\hat{\mathbf{\Phi}}(x), \hat{\mathbf{\Theta}}(x))^{\mathsf{T}}$ on the emergent configuration space $\mathcal{X}_q$ (Definition 6.1) [36, 37, 30, 29]. This packaging is operationally meaningful because the doublet is measured and transported by the *Unified Inter-Modal Correlator* $\hat{\mathbf{G}}_{\Lambda}(x, x')$, a 2×2 block operator over channels $\mathbf{A}, \mathbf{B} \in \{\Phi, \Theta\}$ whose parallel transport is fixed by the path-ordered relational connection $U(x \leftarrow x')$ (Definition 6.4, Section 4.19.1, and eq. (4.11)) [45, 36]. In this sense, $\hat{\mathbf{F}}$ is not merely a formal concatenation: it is the basic field object whose correlations, transport, and responses are already locked to the chronogeometric background produced by the theory itself [36, 37].

Within the doublet, $\hat{\mathbf{\Phi}}(x)$ provides the geometric response that sources admissible distances, durations, and curvatures at expectation level via the composite chronogeometric tensor

$\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$; the associated calculus and measure are governed by the Relational Levi–Civita connection $\nabla^{(\mathrm{rel})}$ and the relational volume element built from $\bar{g}^{\mathrm{Lor}}_{\mu\nu} := \bar{g}_{\mu\nu} + \bar{t}^{\mathrm{sym}}_{\mu\nu}$ (Definitions 6.7 and 6.33) [45]. Complementarily, $\hat{\boldsymbol{\Theta}}(x)$ carries internal symmetries and excitations (charges, flavours, particle-like modes) that propagate on this chronogeometry; their operator-level coupling to geometry is collected in the unified $\hat{\Sigma}$-sector and its expectation $\bar{\Sigma}$ (Definition 5.25) [38, 36, 37]. Conceptually, this is analogous in spirit to electroweak unification—one packages distinct physical aspects into a single field representation—yet here the unification is effected at the level of relational packaging and transport rather than by identifying gauge groups [57, 58, 59, 38, 36, 37]. The upshot is that spacetime structure and internal structure are *co-constituted* from the start, with their interplay made explicit by $\hat{\mathbf{F}}$, $\hat{\mathbf{G}}_\Lambda$, $\nabla^{(\mathrm{rel})}$, and the relational measure (Section 6.1 and Definitions 6.1, 6.4, 6.7 and 6.33) [30, 29, 45, 36, 37].

Etymological Foundations. The names chosen for these fields reflect their roles, drawing from deep linguistic and mythological roots.

Gaiamiti Field ($\hat{\boldsymbol{\Phi}}$)

The name **Gaiamiti** fuses:

- **Gaia** (Γαῖα): The primordial Greek goddess of Earth, the "ever-sure foundation" in Hesiod's *Theogony* [60]. She represents the **primordial ground** or substrate from which structure emerges. She births, separates, advises, and eventually recedes as order matures, embodying both fecundity [61] and the decisive act of separation needed for structure [60]. Her connection to prophecy at Delphi [62] and autochthonous origins [63] reinforces her foundational role. The name itself is likely pre-Greek, suggesting a primordial origin [64, 65].
- **Miti**: From Sanskrit (*miti*), meaning **measure**. It forms the basis of *jyā-miti* ("bowstring-measure"), the classical term for geometry [66]. This highlights geometry as an operational act of relating curves (the world) to straight lines (measurement tools) [34]. This concept spans from practical altar construction (*Śulba-sūtras*) to astronomy [67], eventually yielding terms like *sine* [68]. It signifies the act of rendering relations legible.

Thus, **Gaiamiti** means "measure of the ground" or "Gaia's geometry," signifying the field whose correlations draw out and make legible the geometric structure from the primordial substrate.

Definition 6.2: The Gaiamiti Field The field $\hat{\boldsymbol{\Phi}}(x)$ is formally defined as the first component of $\hat{\mathbf{F}}(x)$, a Hermitian, operator-valued field generating the emergent metric $\bar{g}_{\mu\nu}(x)$ and temporal tensors $\bar{t}_{\mu\nu}(x)$. It admits a modal decomposition:

$$\hat{\boldsymbol{\Phi}}(x) = \sum_{\alpha} \hat{\Phi}_\alpha(x)\, |\alpha\rangle_x$$

over the dialectical modes $\alpha \in \{0, 1, p, s\}$.

Apeiron (ἄπειρον) (Section 6.1) This term, meaning "un-limited" or "indefinite," is redeployed to denote the **quantum pulse of the Gaiamiti field**: the minimal, predicate-free

event of givenness before measure articulates it. It draws on Anaximander's *archē* [69, 70, 71], the Pythagorean opposition of Limit (*peras*) and Unlimited (*apeiron*) [72], and Aristotle's potential infinite [71, 73]. In this framework, it signifies a discrete, pre-metric, rhythmic pulse articulated by measure (Section 6.1).

Tantrakyma Field ($\hat{\mathbf{\Theta}}$). The name **Tantrakyma** combines:

- **Tantra** (तन्त्र): From Sanskrit root $\sqrt{tan}$ ("to stretch, weave") + suffix *-tra* ("instrument") [66]. It primarily means the **warp** of a fabric or a **loom**, extending to mean framework, system, rule, or doctrine. It became a designation for scriptural corpora in Śaiva, Śākta, Vaiṣṇava, and Buddhist traditions from the mid-first millennium CE [74, 75, 76, 77, 78, 79]. The Tibetan translation *rgyud* ("continuity") emphasizes the unbroken transmission, akin to warp threads [75].
- **Kyma** (κῦμα): The classical Greek word for **wave** or swell, from PIE root *keue-* ("to swell, curve") [80, 81]. It signifies dynamic form, embodying both pattern and flux, central to Heraclitus' philosophy (*panta rhei*) [70]. Its modern legacy includes terms like cymatics, kymograph, myokymia, cyma (architecture), and cyme (botany) (Section 6.1).

Therefore, **Tantrakyma** signifies a "woven-wave framework": a field where wave-like excitations ($\hat{\Theta}_\beta(x)$ modes) act as threads woven upon a structural warp (internal symmetries), generating coherent particle content and conserved quantum numbers.

Definition 6.3: The Tantrakyma Field The field $\hat{\mathbf{\Theta}}(x)$ is the second component of $\hat{\mathbf{F}}(x)$. It's a Hermitian, operator-valued field whose excitations correspond to emergent particles and internal symmetries yield conserved quantities. It also has a modal decomposition:

$$\hat{\mathbf{\Theta}}(x) = \sum_\beta \hat{\Theta}_\beta(x)\,|\beta\rangle_x$$

where $\hat{\Theta}_\beta(x)$ describes internal excitation states.

Kymon: The fundamental quantum of the Tantrakyma field, named from *Kyma* + suffix *-on*. It represents a wave-like excitation propagating on the Gaiamiti-generated geometry, carrying charges determined by $\hat{\mathbf{\Theta}}$'s symmetries [82, 14].

Chrono-Kymon: An excitation linked to the temporal structure ($\bar{t}_{\mu\nu}$), specifically a spin-1 mediator of temporal torsion, influencing how clocks and causal order knit together. It represents a quantum twist in the local flow of time, distinct from spin-2 gravitons [83, 11].

Dual Roles and Unification. The Gaiamiti $\left(\hat{\mathbf{\Phi}}\right)$ and Tantrakyma $\left(\hat{\mathbf{\Theta}}\right)$ fields play distinct yet complementary roles within the unified singularity $\hat{\mathbf{\Omega}} = \overset{\circ}{\Lambda}_{\hat{\mathbf{\Phi}}\hat{\mathbf{\Theta}}}$ (Section 6.1), and are operationally packaged by the generalized field doublet $\hat{\mathbf{F}}(x)$ (Definition 6.1).

- **Gaiamiti.** The architect of geometry, establishing admissible distance, duration, and curvature at expectation level through the chronogeometric response $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (Definition 5.26). Its self-correlations $\hat{G}_{\Phi\Phi}$ are the geometric block of the Unified Inter-Modal Correlator and "weave" the spacetime fabric in a gauge-covariant way (Definition 6.4, Section 4.19.1, and eq. (4.11)).

- **Tantrakyma.** The source of flavour/charge and matter-like structure, carrying internal excitations that propagate on the chronogeometry. These modes $\hat{\Theta}_\beta$ live in the excitation fibre orthogonal to geometric deformations and are represented by the dynamical Tantrakyma field section (Definitions 6.141 and 6.142).

This relationship is likened to a *celestial symphony*: Gaiamiti provides the harmonic space (geometry) while Tantrakyma traces the melodic figures (matter content) (Section 6.1).

Correlations and Dynamics. The interplay between Gaiamiti and Tantrakyma is encoded by the *Unified Inter-Modal Correlator* $\hat{\mathbf{G}}_\Lambda(x,x')$, a 2×2 matrix over channels $\mathbf{A},\mathbf{B}\in\{\Phi,\Theta\}$ acting on the generalized field doublet $\hat{\mathbf{F}}$ (Definitions 6.1 and 6.4):

$$\hat{\mathbf{G}}_\Lambda(x,x') \;:=\; \begin{pmatrix} \hat{G}_{\Phi\Phi}(x,x') & \hat{G}_{\Phi\Theta}(x,x') \\ \hat{G}_{\Theta\Phi}(x,x') & \hat{G}_{\Theta\Theta}(x,x') \end{pmatrix}.$$

Each block $\hat{G}_{\mathbf{AB}}(x,x')$ further resolves over the 4×4 dialectical modes (α,β) as operator products of the form $\hat{A}^\dagger_\alpha(x)\,U(x\leftarrow x')\,\hat{B}_\beta(x')$, where $U(x\leftarrow x')$ is the path-ordered relational transport generated by the operational connection (Sections 4.19.1 and 6.1.1 and eq. (4.11)). This construction makes $\hat{\mathbf{G}}_\Lambda$ manifestly state- and connection-covariant from the outset.

The diagonal blocks play distinct roles. The *geometric block* $\hat{G}_{\Phi\Phi}$ encodes self-correlations of Gaiamiti and, at expectation level, feeds into the chronogeometric response $\bar{\Sigma}_{\mu\nu}=\bar{g}_{\mu\nu}+\bar{t}_{\mu\nu}$ that structures distance, duration, and curvature (Definition 5.26). The *matter block* $\hat{G}_{\Theta\Theta}$ captures self-correlations of Tantrakyma governing propagation and internal interactions on that chronogeometry (Definition 6.142). The off-diagonal *interaction blocks* $\hat{G}_{\Phi\Theta}$ and $\hat{G}_{\Theta\Phi}$ implement geometry–matter coupling in both directions—schematically "matter tells geometry how to curve" and "geometry tells matter how to move"—and are evaluated in the same gauge-covariant, transported form (Definition 6.4 and Section 4.19.1).

Dynamically, these correlations are propagated and compared using connections and measures compatible with the emergent geometry. Covariant differentiation is performed with the Relational Levi–Civita connection $\nabla^{(\mathrm{rel})}$ built from the composite tensor $\bar{\Sigma}$ (Definition 6.7), while action and energy functionals integrate correlator traces against the relational volume element $d\mu(x)=\sqrt{-\det \bar{g}^{\mathrm{Lor}}(x)}\,d^dx$, with $\bar{g}^{\mathrm{Lor}}_{\mu\nu}:=\bar{g}_{\mu\nu}+\bar{t}^{\mathrm{sym}}_{\mu\nu}$ (Definition 6.33). In particular, the Relational Hamiltonian $\hat{H}$ decomposes into geometric, matter, and mixed parts $\hat{H}_\Phi+\hat{H}_{\Phi\Theta}+\hat{H}_\Theta$, making the block structure of $\hat{\mathbf{G}}_\Lambda$ operational in energy accounting (Definition 6.54 and eq. (6.109)).

Relational Levi–Civita derivative. On the emergent configuration space $\mathcal{X}_q$ (cf. Proposition 4.625), the theory singles out a unique torsion-free connection $\nabla^{(\mathrm{rel})}$ that is compatible with the full *chronogeometric tensor* $\bar{\Sigma}_{\mu\nu}=\bar{g}_{\mu\nu}+\bar{t}_{\mu\nu}$ (cf. Definitions 5.26 and 6.7). Its Christoffel symbols are computed from the composite $\bar{\Sigma}$,

$$\Gamma^\lambda_{\mu\nu}[\bar{\Sigma}] \;=\; \tfrac{1}{2}\,\bar{\Sigma}^{\lambda\rho}\Big(\partial_\mu\bar{\Sigma}_{\nu\rho}+\partial_\nu\bar{\Sigma}_{\mu\rho}-\partial_\rho\bar{\Sigma}_{\mu\nu}\Big).$$

Here $\bar{\Sigma}^{\mu\nu}$ denotes the inverse of $\bar{\Sigma}_{\mu\nu}$ (cf. Definition 6.24 and eq. (6.22)).

Because $\nabla^{(\mathrm{rel})}$ is metric-compatible with $\bar{\Sigma}$, parallel transport, curvature, and geodesics are all defined with respect to the *composite* response, so geodesic motion is influenced both by the metric part $\bar{g}_{\mu\nu}$ and by gradients of the temporal tensor $\bar{t}_{\mu\nu}$ (see Definition 6.7 and references therein, e.g. [7, 84, 85]).

Relational volume element. The invariant integration measure on $\mathcal{X}_q$ is determined by the *Effective Lorentzian Metric* $\bar{g}^{\mathrm{Lor}}_{\mu\nu} := \bar{g}_{\mu\nu} + \bar{t}^{\mathrm{sym}}_{\mu\nu}$ (cf. Definition 5.77 and Section 6.1.5):

$$d\mu(x) \;:=\; \sqrt{-\det \bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)}\, d^d x,$$

as stated in Definition 6.33. This measure is the one used in action and energy functionals [7]; crucially, its dependence on both the geometric curvature $\bar{g}_{\mu\nu}$ and the temporal structure $\bar{t}^{\mathrm{sym}}_{\mu\nu}$ makes the weighting of fields intrinsically state-responsive, with potential phenomenological implications in extreme regimes [86, 87, 88, 89, 1].

The Vacuum Manifold. The vacuum is characterised as a state of *minimal relational complexity*, formulated in parallel variational and spectral terms (Section 6.1.8). In the variational picture, one evaluates the *relational action functional* on a state $|\mathbf{\Psi}\rangle \in \mathcal{H}_{\mathrm{em}}$ (cf. Definition 4.287) using the state-conditioned measure built from $\bar{\Sigma}^{(\mathbf{\Psi})}_{\mu\nu}(x) := \langle \mathbf{\Psi}|\hat{\Sigma}_{\mu\nu}(x)|\mathbf{\Psi}\rangle$ (Definitions 5.25 and 6.33):

$$S[|\mathbf{\Psi}\rangle] := \int_{\mathcal{X}_q} d\mu^{(\mathbf{\Psi})}(x)\, \mathrm{Tr}\Big(\langle \mathbf{\Psi}|\hat{\mathbf{G}}_\Lambda(x,x)|\mathbf{\Psi}\rangle \Big),$$
$$d\mu^{(\mathbf{\Psi})}(x) := \sqrt{-\det \bar{g}^{\mathrm{Lor}(\mathbf{\Psi})}(x)}\, d^d x.$$

See Definitions 6.4 and 6.33 and Section 6.1.10.

The trace $\mathrm{Tr}(\cdot)$ aggregates over field channels (Φ, Θ) and their dialectical modes, so that S encompasses geometric, matter, and interaction contributions in a single expression (cf. the sectoral decompositions $S_\Phi, S_\Theta, S_{\Phi\Theta}$ and Definitions 6.4 and 6.172).
In the spectral picture, the *Relational Hamiltonian* is defined so that its expectation value coincides with the action—this is the variational–spectral equivalence (Definition 6.54):

$$\langle \mathbf{\Psi}|\hat{H}|\mathbf{\Psi}\rangle \;=\; S[|\mathbf{\Psi}\rangle], \qquad \hat{H}[|\mathbf{\Psi}\rangle] \;:=\; \int_{\mathcal{X}_q} d\mu^{(\mathbf{\Psi})}(x)\, \mathrm{Tr}_{\mathrm{op}}\Big(\hat{\mathbf{G}}_\Lambda(x,x)\Big),$$

with the blockwise split $\hat{H} = \hat{H}_\Phi + \hat{H}_{\Phi\Theta} + \hat{H}_\Theta$ made explicit in eq. (6.109). The dependence of $d\mu^{(\mathbf{\Psi})}$ on $\bar{g}^{\mathrm{Lor}}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}^{\mathrm{sym}}_{\mu\nu}$ (Definition 6.33) renders $\hat{H}$ intrinsically state-conditioned.
The *Vacuum Manifold* $\mathcal{M}_{\mathrm{vac}}$ is then the set of normalised states that minimise S (variational view) or, equivalently, the ground-state eigenspace of $\hat{H}$ (spectral view) (Section 6.1.17 and Definition 6.61):

$$\mathcal{M}_{\mathrm{vac}} \;:=\; \Big\{ |\mathbf{\Psi}_{\mathrm{vac}}\rangle \in \mathcal{V}_0 \;\Big|\; \langle \mathbf{\Psi}_{\mathrm{vac}}|\mathbf{\Psi}_{\mathrm{vac}}\rangle = 1 \Big\}, \qquad \mathcal{V}_0 = \ker\Big(\hat{H} - E_0 \mathbb{I}_{\mathrm{em}}\Big),\; E_0 = \inf \mathrm{spec}(\hat{H}).$$

Within this manifold, a particular *Quantum Holo-State* $|\overset{\circ}{\mathbf{\Psi}}\rangle$ is selected to describe our universe (cf. Section 6.1.17), closing the loop from correlators → geometry → dynamics → vacua.

Operator Algebra and Field Definitions. The framework uses an algebraic approach, defining observables abstractly and then realizing them via the GNS construction.

- **Algebra of Observables $\mathcal{A}$:** The foundational C*-algebra (Section 1.4.1).
- **GNS Construction:** Generates the emergent Hilbert space $\mathcal{H}_{\text{em}}$, representation $\pi^{\mathring{\Psi}}$, and cyclic vector $|\mathring{\Psi}\rangle$ from $\mathcal{A}$ and the holo-state functional $\omega^{\mathring{\Psi}}$ (Definition 1.34).
- **Algebra of Endomorphisms** $\text{End}(\mathcal{H}_{\text{em}})$**:** The concrete arena of bounded linear operators $\mathcal{B}(\mathcal{H}_{\text{em}})$ on the emergent Hilbert space, forming a unital Banach $*$-algebra (likely C*-algebra) (Definition 6.111, Definition 6.112, Definition 6.113). $\pi^{\mathring{\Psi}}$ maps $\mathcal{A}$ into this space (Section 6.1.28).

Fluctuations around the vacuum are described using the tangent bundle:

Operator–theoretic tangent bundle of the vacuum. At a vacuum state $|\Omega\rangle \in \mathcal{M}_{\text{vac}}$ (cf. Remark 6.91), the operator–theoretic tangent fibre is taken to be

$$T_{|\Omega\rangle}\mathcal{M}_{\text{vac}} \;=\; \Big\{\, |V\rangle \in \mathcal{V}_0 \;\Big|\; \text{Re}\,\langle\Omega|V\rangle = 0 \Big\},$$

namely the space of infinitesimal, energy-preserving deformations $|V\rangle$ that are physically distinct from the vacuum (orthogonal in the real part) (Definition 6.90). This space is equipped with the inner product $g_T(|V\rangle, |W\rangle) := \langle V|W\rangle_{\mathcal{H}_{\text{em}}}$, which provides the canonical metric on vacuum tangents (Definition 6.104).

Dynamical Gaiamiti field. The geometric response is encoded by a smooth operator-valued section $\hat{\mathbf{\Phi}}(x)$ of the bundle $\text{End}(\mathcal{H}_{\text{em}}) \otimes T\mathcal{M}_{\text{vac}}$ over $\mathcal{X}_q$; locally one has $\hat{\mathbf{\Phi}}(x) \in \text{End}(\mathcal{H}_{\text{em}}) \otimes T_{|\Omega(x)\rangle}\mathcal{M}_{\text{vac}}$. Thus $\hat{\mathbf{\Phi}}(x)$ acts as a quantum–geometric deformation operator anchored to the vacuum-tangent directions at the point x (Definition 6.118; cf. also Section 6.5 and Definition 6.172).

Excitation fibre and Dynamical Tantrakyma field. Orthogonally to the geometric tangents, the excitation fibre at x is

$$\mathcal{E}_{\text{exc}}(x) \;=\; \big(T_{|\Omega(x)\rangle}\mathcal{M}_{\text{vac}}\big)^{\perp},$$

which collects matter-like (internal) directions in state space (Definition 6.141). The corresponding field $\hat{\mathbf{\Theta}}(x)$ is an operator-valued section of $\text{End}(\mathcal{H}_{\text{em}}) \otimes \mathcal{E}_{\text{exc}}$ over $\mathcal{X}_q$, governing the propagation of excitations that are orthogonal to geometric deformations (Definition 6.142). In practice, $\hat{\mathbf{\Phi}}$ and $\hat{\mathbf{\Theta}}$ are coupled and transported jointly through the Unified Inter-Modal Correlator, ensuring gauge-covariant comparison across points of $\mathcal{X}_q$ (Definition 6.4, Section 4.19.1, and eq. (4.11)).

Bundle-Theoretic Formulations (Section 6.4)

- $\hat{\mathbf{\Phi}}(x) = \sum_{i=1}^{r} \hat{\Phi}^i(x)\,|e_i(x)\rangle$ in a tangent space basis $\{|e_i(x)\rangle\}$. $\hat{\Phi}^i$ are operator-valued amplitudes for geometric deformation modes.

- $\hat{\mathbf{\Theta}}(x) = \sum_{\alpha,\beta} \theta^{\alpha\beta}(x)\, |\alpha\rangle_x \langle\beta|_x$ in the dialectical basis. $\theta^{\alpha\beta}$ govern internal dynamics and mixing.

Dual Roles (Summary): $\hat{\mathbf{\Phi}}$ sources geometric observables (apeiron, graviton). $\hat{\mathbf{\Theta}}$ sources matter observables (kymons, particle species, gauge groups via symmetries).

Action and dynamics of the Gaiamiti field. The dynamics of the geometric response are encoded in an operator-valued Lagrangian density that contracts the composite chronogeometric tensor with a vacuum-tangent inner product on covariant derivatives of the field (Section 6.5 and Definition 6.172):

$$\mathcal{L}_\Phi(x) \;=\; \mathrm{Tr}\Big\{ \bar{\Sigma}^{\mu\nu}(x)\, g_T\Big(\nabla_\mu \hat{\mathbf{\Phi}}(x), \nabla_\nu \hat{\mathbf{\Phi}}(x)\Big) + \mathcal{V}_\Phi\Big(\hat{\mathbf{\Phi}}(x)\Big)\Big\}\,.$$

Here $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ is the expectation-level chronogeometric tensor (Definition 5.26), g_T is the canonical inner product on the operator-theoretic vacuum tangents (Definition 6.104), and $\mathcal{V}_\Phi$ encodes self-interactions. The sectoral action has two equivalent guises: fundamentally as a correlator trace integrated with the state-dependent measure,

$$S_\Phi[|\mathbf{\Psi}\rangle] = \int_{\mathcal{X}_q} d\mu^{(\mathbf{\Psi})}(x)\, \mathrm{Tr}\Big(\langle \mathbf{\Psi}|\hat{G}_{\Phi\Phi}(x,x)|\mathbf{\Psi}\rangle\Big).$$

See Definitions 6.4 and 6.33 and Section 6.1.10.

and operationally as the Lagrangian integral

$$S_\Phi[\hat{\mathbf{\Phi}}] \;=\; \int_{\mathcal{X}_q} d\mu(x)\; \mathcal{L}_\Phi(x) \quad (\textit{Definitions } 6.33 \textit{and } 6.174).$$

Self-consistency is built in: the $\bar{\Sigma}_{\mu\nu}$ that contracts the kinetic term is itself sourced by $\hat{\mathbf{\Phi}}$ through the holo-state expectation (Section 6.5 and Definition 5.26).

Covariant structure and the operator bundle connection. The total covariant derivative splits into a base-manifold piece and an internal (bundle) piece (Definition 6.184):

$$\nabla_\mu \hat{\mathbf{\Phi}} \;=\; \nabla^{(\mathrm{rel})}_\mu \hat{\mathbf{\Phi}} \;+\; D_\mu \hat{\mathbf{\Phi}}, \qquad D_\mu \;=\; \partial_\mu + i\,[\mathcal{A}_\mu(x), \cdot\,].$$

The base connection $\nabla^{(\mathrm{rel})}$ is the torsion-free, $\bar{\Sigma}$-compatible Relational Levi–Civita derivative acting on $\mathcal{X}_q$ (Definition 6.7 and Proposition 4.625), while $\mathcal{A}_\mu(x)$ is the Hermitian operator bundle connection arising intrinsically from the algebra of observables via the GNS representation (Definition 6.192 and Remark 6.198). This $\mathcal{A}_\mu$ is universal—acting on both $\hat{\mathbf{\Phi}}$ and $\hat{\mathbf{\Theta}}$ (Remark 6.200)—and governs gauge-covariant parallel transport (cf. the operational connection and its path-ordered transport $U(x \leftarrow x')$ in Section 4.19.1 and eq. (4.11)). Its sectoral quanta (the *mition* in the Gaiamiti sector and the *tantron* in the Tantrakyma sector) are discussed in Remark 6.201, with masses generated via a BEH-type mechanism (Remark 6.205).

Symmetry considerations. Within the relational framework, symmetry is assessed at two levels: the *law level*—the invariances of the relational action and Hamiltonian—and the *state level*—the invariances of vacua and excitations that inhabit the vacuum manifold. Spontaneous symmetry breaking (SSB) occurs when the relational action $S[|\mathbf{\Psi}\rangle]$ and the Relational Hamiltonian $\hat{H}$ possess a symmetry G (Definitions 6.33 and 6.54) that is not shared by the selected vacuum $|\mathbf{\Psi}_{\mathrm{vac}}\rangle \in \mathcal{M}_{\mathrm{vac}}$ (Section 6.1.17 and Definition 6.61). The resulting vacuum degeneracy is organised by the coset G/H, and the tangential excitations along these flat directions manifest as Nambu–Goldstone modes living in the operator–theoretic tangent bundle $T\mathcal{M}_{\mathrm{vac}}$ with inner product g_T (Definitions 6.90 and 6.104; see also [90, 91, 82, 92, 93, 94]). When the relevant symmetry is gauged, the Brout–Englert–Higgs (BEH) mechanism operates: the universal operator bundle connection $\mathcal{A}_\mu$ (Definition 6.192 and Remark 6.200) reorganises the spectrum so that gauge quanta (mition/tantron) acquire mass (Remarks 6.201 and 6.205; cf. [95, 96, 97, 98, 99]), with familiar low-dimensional constraints [100, 101].

Emergent (asymmetry-induced) symmetry refers to situations where the infrared theory exhibits a larger symmetry H_{IR} than is present microscopically G_{UV} [82, 14, 102]. In the present setting, such enhancement can be diagnosed directly in the block structure and transport of the Unified Inter-Modal Correlator $\hat{\mathbf{G}}_\Lambda$ (Definition 6.4 and Section 4.19.1): symmetry-breaking operators may become RG-irrelevant in the correlator's off-diagonal entries, or descend to accidental invariants in the effective Lagrangian density $\mathcal{L}_\Phi$ (Definition 6.172). At the same time, the relational measure $d\mu = \sqrt{-\det \bar{g}^{\mathrm{Lor}}}\, d^d x$ (Definition 6.33) weights configurations in a way that can suppress symmetry-violating responses of $\bar{t}_{\mu\nu}$ relative to $\bar{g}_{\mu\nu}$, yielding an effectively more symmetric infrared chronogeometry. Standard diagnostics—scaling suppression, anomaly consistency [103], approach to fixed-point structure—carry over, now stated in terms of $\hat{\mathbf{G}}_\Lambda$, $\mathcal{L}_\Phi$, and the state-dependent background $\bar{\Sigma}_{\mu\nu}$.

Weaving the two arrows. Many systems realise both tendencies: microscopic asymmetries are diluted along the flow to the infrared (emergence), and the resulting symmetric infrared law is then *spontaneously* broken by the choice of vacuum (SSB). In our notation, the law-level arrow is tracked by the evolution of the correlator blocks and couplings in $\mathcal{L}_\Phi$ (Definitions 6.4 and 6.172) together with the compatible covariant calculus $\nabla^{(\mathrm{rel})}$ (Definition 6.7); the state-level arrow is read off from order parameters living on $T\mathcal{M}_{\mathrm{vac}}$ and the pattern of NG directions inside $\mathcal{M}_{\mathrm{vac}}$ (Definitions 6.61 and 6.90). The 2×2 "Laws vs. States'' matrix in table 6.1 organises these possibilities and their diagnostics, keeping clear which effects are due to changes in the effective *laws* versus choices of *states*.

BRST quantisation and gauge redundancy. Gauge invariance is treated as a redundancy of description by promoting the field space to include ghosts, anti-ghosts, and auxiliary fields and endowing it with a nilpotent BRST charge $\hat{Q}_{\mathrm{BRST}}$ satisfying $\hat{Q}^2_{\mathrm{BRST}} = 0$. Physical states are identified with the ghost-number–zero cohomology, $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$, so that gauge-equivalent configurations differ by BRST-exact vectors and are projected out (*Kugo–Ojima* quartet mechanism for unitarity; see [104, 105, 106, 107, 108, 91, 82]). In our setting, this cohomology is taken inside the emergent Hilbert space $\mathcal{H}_{\mathrm{em}}$ and is compatible with the relational covariant calculus $\nabla^{(\mathrm{rel})}$ and the state-dependent integration measure $d\mu = \sqrt{-\det \bar{g}^{\mathrm{Lor}}}\, d^d x$ (Definitions 6.7 and 6.33), so that BRST invariance is formulated directly on

$(\mathcal{X}_q, \bar{\Sigma})$.

Gauge fixing appears as a BRST-exact deformation of the action, $\delta S = \zeta\Psi$ with $\zeta(\cdot) = \{\hat{Q}_{\mathrm{BRST}}, \cdot\}$, which preserves the cohomology class of observables. Accordingly, expectation values of BRST-invariant operators are independent of gauge parameters, as captured by the *Nielsen identity* and the *gauge-fixing exactness principle* (Proposition 6.214 and Principle 6.1). Within the relational framework, the universal operator bundle connection $\mathcal{A}_\mu$ that enters the internal part of the covariant derivative $D_\mu = \partial_\mu + i[\mathcal{A}_\mu, \cdot]$ (Definitions 6.184 and 6.192) is thus fixed only up to BRST-exact choices; "Higgsing'' is not literal gauge SSB, since the would-be Goldstone directions are BRST-exact and removed from $\mathcal{H}_{\mathrm{phys}}$ (Proposition 6.213).

The chapter relates these consistency conditions to the HSP viability criteria: *Stage I* viability (self-coherence of the description) is equivalent to BRST consistency—nilpotency, BRST-invariant dynamics, and absence of gauge anomalies (Principle 6.2 and Proposition 6.217). Operationally, this means the Unified Inter-Modal Correlator $\hat{\mathbf{G}}_\Lambda$, the relational derivative $\nabla^{(\mathrm{rel})}$, and the measure $d\mu$ (Definitions 6.4, 6.7 and 6.33) can be chosen so that all gauge-variant contributions are BRST-exact, while the cohomology classes furnish the gauge-invariant observables and states on the relational manifold.

Duality, quadrants, and holosymmetry. The chapter links the Axiom of Duality to the quadrant structure used for symmetry assessment (see table 6.1). Starting from the duality object and its status lift $\Delta_{\mathcal{C}}$ (Definitions 6.220 and 6.221), one obtains commuting involutions Δ_{Law} and Δ_{State} acting on the law/state symmetry predicates (Definition 6.224). Their $\mathbb{Z}_2 \times \mathbb{Z}_2$ action generates four quadrants—$(\mathsf{Th}, \mathsf{Th})$,$(\mathsf{Th}, \mathsf{AntiTh})$,$(\mathsf{AntiTh}, \mathsf{Th})$,$(\mathsf{AntiTh}, \mathsf{AntiTh})$—which follow from the commuting-lifts result and are organised by the quadrant schema (Proposition 6.225 and Theorem 6.231) [52, 30, 29]. These quadrants correspond, respectively, to the Unbroken, SSB, Emergent, and Explicit–Breaking regimes (Corollary 6.232) [52, 109].

For gauge theories, a cohomological reading is provided: BRST maps the predicates to statements about cohomology classes and anomaly filters, aligning duality with the quartet structure (Remark 6.226, Proposition 6.227, and Corollary 6.228) [52, 109, 110, 111]. In short, duality supplies the algebraic backbone, while the quadrant taxonomy records where symmetry resides—in the *laws* or in the *states* [29].

On top of this algebraic scaffold, the chapter introduces a holosymmetric meta-layer based on two-path interferometry. Law-coherence and state-coherence are captured by functionals Λ_{law} and Λ_{state} that are defined in a parameter-agnostic manner (Definitions 6.237 and 6.238) and evaluated directly in the physical cohomology $\mathcal{H}_{\mathrm{phys}}$. They are gauge-parameter independent (Nielsen-type control; Lemma 6.243), respect the triadic channel split (functoriality with the corr/vac/spec decomposition; Lemma 6.247), and preserve causal cones in the holosymmetric sense (Lemma 6.248) [52, 30, 29, 45]. Crucially, overlaying these diagnostics does not change a system's quadrant membership (Proposition 6.251); instead, it furnishes operational signatures—interference visibilities and phase responses—that distinguish the four regimes (Theorem 6.253, table 6.2) [22, 23, 24, 25]. Thus holosymmetric interferometry supplies an experimentally legible veneer over the duality–quadrant architecture, cleanly separating law-level coherence from state-level order (see table 6.1).

Table 1: Symmetry–Asymmetry Permutations Matrix: Laws vs. States

Laws / World (State)	**Symmetry of World (State)**	**Asymmetry of World (State)**
Symmetry of Law	**I. Unbroken (symmetry preserved)** Unique symmetric vacuum (thermodynamic limit); order parameter = 0; Ward identities exact. Topological/SPT structure may still distinguish phases.	**II. Spontaneous Symmetry Breaking (SSB)** Degenerate vacua related by the symmetry; nonzero order parameter; Nambu–Goldstone modes for broken *global* continuous generators (gauge case diagnosed via gauge-invariant observables).
Asymmetry of Law	**III. Emergent / Accidental Symmetry** IR (effective-law) exhibits enlarged symmetry via RG irrelevance or EFT power suppression; symmetry-violating effects shrink with scale.	**IV. Explicit Breaking (no SSB)** Asymmetric couplings select a unique ground state; no Goldstones. Residual subgroups may remain; responses track the breaking terms.

Notes (cf. table 6.1): (i) "Gauge symmetry" is a redundancy; BEH/Higgs phases are diagnosed by gauge-invariant features (massive vectors, screening), not literal gauge SSB. (ii) Topological/SPT structure can live in quadrant I (symmetry preserved) and should be diagnosed by nonlocal/torsion-like invariants rather than order parameters.

Scale-aware analysis strategy. The chapter closes with a practical programme for analysing phenomena across the four symmetry quadrants that is explicitly scale-aware. First, *scale-tagging* is introduced by combining a UV bandlimit and an IR window: a spectral projector $P_x(\lambda_{\mathrm{UV}})$ selects modes below a local cutoff λ_{UV} (Definition 6.278 and Proposition 6.279), while an IR window $K_{\mathrm{IR}}^{(\ell)}$ of range ℓ implements coarse graining (Definitions 6.280 and 6.289 and Propositions 6.281 and 6.290). Together these define a *band functor* that enforces coherent self-regularisation and the reciprocal resolution trade-off between λ_{UV} and ℓ (Definition 6.282 and Proposition 6.283). With this structure in hand, the theory formulates *scale-tagged* actions, field equations, and Ward identities—Euler–Lagrange equations projected by $P_x(\lambda_{\mathrm{UV}})$ and convolved by $K_{\mathrm{IR}}^{(\ell)}$ (Definition 6.293 and Proposition 6.294), and conserved relations that remain valid after band restriction (Definition 6.295 and Proposition 6.296). The outcome is a family of UV/IR-consistent dynamical statements that can be compared across scales without leaving the relational framework.

Robustness is certified by a *plateau criterion*. A physical claim—such as a mode identification or an effective coupling—is accepted only if it is stable within tolerance $\varepsilon_{\mathrm{tol}}$ across a plateau region in the two-parameter scale space $(\lambda_{\mathrm{UV}}, \ell)$, quantified by small log-derivatives with respect to both scale knobs (Definition 6.297 and Proposition 6.298). Crucially, plateau assessments are *gauge-parameter independent*: the Nielsen-type control shows that admissible variations of gauge-fixing do not spoil plateau stability (Proposition 6.299 and Corollary 6.265). Hence, plateau-qualified statements have operational meaning that does not depend on auxiliary gauge choices.

Finally, a *quadrant traversal strategy* organises concrete analyses in each of the symmetry quadrants (QI–QIV). One proceeds from the variational backbone—scale-tagged actions and projected equations of motion (Definition 6.293 and Proposition 6.294) together with Ward identities (Definition 6.295 and Proposition 6.296)—to linear response (propagators) and then to holosymmetric, two-path diagnostics of law- and state-coherence, Λ_{law} and Λ_{state} (Definitions 6.237 and 6.238). At every step, mode tagging is treated as provisional until plateau stability and BRST consistency are verified; the procedure is spelled out quadrant-by-quadrant in Section 6.12.5. To keep the workflow auditable, the chapter recommends maintaining *practical ledgers*—explicit records of the chosen $(\lambda_{\mathrm{UV}}, \ell)$, gauges, and diagnostic outcomes—throughout the analysis (Section 6.12.7).

The objects and the holosymmetric lift.

Triadic channelization. Work with three BRST-closed channel projectors $\{\Pi_\alpha\}_{\alpha\in\{\mathrm{corr,vac,spec}\}}$ of ghost number 0. These projectors implement the channel split used throughout: they are the projectors that define the channel-restricted visibilities (cf. the channel visibilities in (6.347) and (6.348)) and the SSB channel signatures in Theorem 6.261. Gauge-parameter (ξ) independence of channel observables is ensured by BRST closure; see Proposition 6.276 and the pointwise identities (6.318) and (6.325).

Holosymmetric lift. For any operator/kernel A (e.g. $A \in \pi(\mathcal{A}) \subset \mathrm{End}(\mathcal{H}_{\mathrm{em}})$), define the holosymmetric lift (cf. Proposition 6.257)

$$\mathfrak{H}_{\lambda_{\mathrm{UV}},\ell}[A] \;=\; \sum_{\alpha\in\{\mathrm{corr,vac,spec}\}} \Pi_\alpha\, K_{\mathrm{IR}}^{(\ell)}\, P_x(\lambda_{\mathrm{UV}})\, A\, P_x(\lambda_{\mathrm{UV}})\, K_{\mathrm{IR}}^{(\ell)}\, \Pi_\alpha\,,$$

where the UV/IR tags $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ are given in (6.353), (6.359), and (6.361) (see also Definitions 6.280 and 6.289 and Proposition 6.290). Feeding $\mathfrak{H}_{\lambda_{\mathrm{UV}},\ell}[\cdot]$ into expectation values produces expectation-level observables (cf. the algebra–Hilbert bridge Section 6.1.28 and, e.g., (6.107)); their plateau regularity is controlled by Definition 6.297 and Proposition 6.298.

Properties:

1. *Block-diagonal (no cross-channel couplings).* The sum above is diagonal in the Π_α blocks (see the triadic split (6.306) and the blockwise structure in Proposition 6.257), matching the channelwise diagnostics used for SSB and Berry/WZ phases (cf. Theorem 6.261).
2. *Functorial over direct sums/projections.* The lift respects algebraic decompositions and projections (cf. Proposition 6.257), ensuring compatibility with the construction of expectation-level quantities and with plateau-based stability bounds (Proposition 6.298).
3. *Cone-preserving relative to $\bar{\Sigma}_{\mu\nu}$.* The lift leaves the principal symbol—and thus the chronogeometric light-cone structure—unchanged (cone preservation in Proposition 6.257). For the relational connection and its Levi–Civita form see Definition 6.7 and the Christoffel expression (6.22); related ingredients include Definitions 6.24 and 6.33 and (6.62), (6.74).

Visibilities, RG convention, and Duhamel control.

- **RG scaling convention.** Running couplings at momentum $\mu \sim \ell^{-1}$ obey

$$g_k(\mu) = \mathcal{O}\Big(\mu^{\Delta_k-(d+z)}\Big) \quad (\mu \downarrow 0), \qquad \Longleftrightarrow \qquad g_k(\ell) = \mathcal{O}\Big(\ell^{-(\Delta_k-(d+z))}\Big) \quad (\ell \uparrow \infty),$$

  with the change of variables $\mu = 1/\ell$. On plateau regions $\mathcal{B}$ (cf. Definition 6.297), the scale response of expectation-level fields is uniformly controlled by Proposition 6.298.

- **Channel-restricted visibilities.** Using the projectors $\{\Pi_\alpha\}$ from the holosymmetric framework (cf. Proposition 6.257), define the law- and state-side visibilities (cf. (6.347), (6.348))

$$\Lambda_{\mathrm{law}}^{(\alpha)}(\ell) = \mathrm{Tr}\Big[\Pi_\alpha\, U_B^\dagger(\ell)\, U_A(\ell)\, \Pi_\alpha\, \rho_0\Big],$$
$$\Lambda_{\mathrm{state}}^{(\alpha)}(V) = \Big\langle \Omega_2 \Big|\, \Pi_\alpha\, U^\dagger(V)\, U(V)\, \Pi_\alpha \,\Big|\Omega_1\Big\rangle.$$

  Because the lift is block-diagonal and cone-preserving (see Proposition 6.257), these visibilities inherit channel block-diagonality and preserve the principal chronogeometric cones. Moreover, they are *pointwise* gauge-parameter independent in $\mathcal{H}_{\mathrm{phys}}$ by BRST closure (see Proposition 6.276 and the identities (6.318), (6.325)). As smooth functionals on the IR manifold $\mathcal{M}_{\mathrm{IR}}$, their "defect potentials"

$$\mathcal{V}^{(\alpha)}(\varphi;\ell) := 1 - \Big|\Lambda_{\mathrm{law}}^{(\alpha)}(\ell)\Big|$$

  coincide with the channel visibilities (6.347) and are used to read off asymptotic exponents on plateaus (cf. (6.352), (6.277)).

- **Duhamel/BCH control.** For a scale-dependent perturbation $\delta H(\ell') = \sum_k g_k(\ell')\,\mathcal{O}_k(\ell')$, the channel-projected propagator admits a Duhamel expansion yielding

$$1 - \left|\Lambda^{(\alpha)}_{\mathrm{law}}(\ell)\right| \;\le\; c_{1,\alpha}\int_0^\ell d\ell' \sum_k |g_k(\ell')| \;+\; c_{2,\alpha}\int_{0<\ell''<\ell'<\ell} d\ell'\,d\ell'' \sum_{k,k'} |g_k(\ell')g_{k'}(\ell'')| \;+\; \cdots,$$

  cf. the law-side defect bound (6.311) (see also (6.305), (6.310)). Combining this with the IR scaling laws (6.352) and (6.277) shows that *irrelevant* deformations ($\Delta_k > d+z$) drive a power-law *improvement* of the defects on plateau bands (uniformly controlled by Proposition 6.298):

$$\mathcal{V}^{(\alpha)}(\varphi;\ell) \;=\; \mathcal{O}\Big(\ell^{-(\Delta-(d+z))}\Big) \quad (\ell \uparrow \infty),$$

  in agreement with the IR scaling relations (6.352) and (6.277). Channelwise SSB diagnostics extracted from these visibilities are summarized in Theorem 6.261.

Plateaus (scale stability) and certification.

- **Definition (plateau region).** Let $\mathcal{O}(\lambda_{\mathrm{UV}}, \ell)$ be an observable produced by a scale-tagging functor $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ (cf. the holosymmetric framework in Proposition 6.257 and the expectation-field map (6.362)). Set $u = \ln \lambda_{\mathrm{UV}}$, $v = \ln \ell$, $D_\lambda = \partial_u$, $D_\ell = \partial_v$. On a rectangle $\mathcal{B} = \big[\lambda_{\mathrm{UV},1}, \lambda_{\mathrm{UV},2}\big] \times \big[\ell_1, \ell_2\big]$, we say $\mathcal{O}$ is *plateau* (scale-stable) iff the bounds below hold.

$$\sup_{(u,v)\in \ln \mathcal{B}} \Big|D_\lambda \mathcal{O}\Big| \le \varepsilon_{\lambda_{\mathrm{UV}}}, \qquad \sup_{(u,v)\in \ln \mathcal{B}} \Big|D_\ell \mathcal{O}\Big| \le \varepsilon_\ell.$$

  For expectation-level fields $\mathbf{\Phi}_{\mathrm{IR}}$ (defined via Definition 5.26 together with (6.362)), these bounds imply the finite-difference/stability estimate of Proposition 6.298, which feeds into the visibility functionals (6.302) and their IR improvements (cf. Section 6.12.5, (6.277)).

- **Finite-difference certification.** If $\mathcal{O} \in C^1(\ln \mathcal{B})$ and the plateau criterion holds (cf. Section 6.12.4), then for any two points $(u, v), (u', v') \in \ln \mathcal{B}$,

$$\Big|\mathcal{O}(u', v') - \mathcal{O}(u, v)\Big| \;\le\; \varepsilon_{\lambda_{\mathrm{UV}}}\,|u' - u| \;+\; \varepsilon_\ell\,|v' - v|.$$

  This estimate controls the slow variation of both fields (6.362) and channel-restricted visibilities (6.347)–(6.348) across a plateau, and underlies the extraction of asymptotic exponents entering the scaling relations Section 6.12.5, (6.277).

Gauge-parameter (Nielsen) independence and plateau/exponent invariants.

- **Pointwise ξ-independence (Nielsen invariance).** After inserting the scale tags (and optional Π_α, cf. Proposition 6.257), both law- and state-side visibilities (6.347)–(6.348) are BRST-closed; hence for each fixed scale argument

$$\partial_\xi \Lambda_\square(\cdot) = 0, \qquad \partial_\xi \Lambda^{(\alpha)}_\square(\cdot) = 0.$$

  See Proposition 6.276 and the pointwise identities (6.318), (6.325). This is compatible with cone preservation in Proposition 6.257 and with the relational chronogeometry defined by Definition 6.7 together with (6.22).

- **Plateau locations are gauge-independent.** Because the Nielsen equalities above (see Propositions 6.264 and 6.276 and the pointwise identities (6.294), (6.325)) are *pointwise* in ℓ (or V), predicates such as the plateau criterion Section 6.12.4 and the stability bound Proposition 6.298 hold at the same scale points for all ξ, channel-wise as well. In particular, the IR geodesic transport (with coefficients given by (6.22) from Definition 6.7) is unaffected at principal order by changes of ξ.

- **Asymptotic exponent invariance.** If on a plateau $1 - |\Lambda_{\text{law}}(\ell)| = C\,\ell^{-\beta}\,(1 + r(\ell))$ as $\ell \uparrow \infty$ (cf. the IR scaling relations (6.352) and (6.277)), then the effective exponent

$$\beta_{\text{eff}}(\ell) := -\frac{d\log\Big(1 - |\Lambda_{\text{law}}(\ell)|\Big)}{d\log\ell}$$

  is ξ-invariant. This guarantees that the channelwise SSB diagnostics summarized in Theorem 6.261 are gauge-robust on plateaus.

The SSB arrow. Assuming a G-symmetric, anomaly-free law and a thermodynamic limit, Chapter 6 gives channel-wise SSB signatures:

- **(a) Correlational channel** corr**.** Long-range order/stiffness and gapless Nambu–Goldstone (NG) poles live here; dispersion and soft responses are encoded in $\Lambda_{\text{law}}^{(\text{corr})}(\ell)$ (stable up to irrelevant corrections). If the broken generators are *gauged*, Elitzur/BRST remove the would-be NG poles from $\mathcal{H}_{\text{phys}}$ (quartet mechanism), so no physical NG pole appears in Π_{corr}.

- **(b) Vacuum channel** vac**.** The order parameter is the one-point function in Π_{vac}; distinct SSB vacua become orthogonal in the thermodynamic limit (state-side Π_{vac} visibility tends to 0 with volume V), while on the law side exact G symmetry gives $|\Lambda_{\text{law}}^{(\text{vac})}(\ell)| \to 1$ as $\ell \downarrow 0$.

- **(c) Spectral channel** spec**.** This block isolates commutators/phase curvature. Along adiabatic loops γ in slow-parameter space, the *phase* of $\Lambda_{\text{law}}^{(\text{spec})}$ equals a Berry/Wess–Zumino integral, while the principal symbol—and hence the $\bar{\Sigma}$ light-cones—remain unchanged (only intra-cone phases/subprincipal terms are modified). These geometric terms shift NG couplings/dispersion without altering the cones.

How irrelevant breakings improve law-side visibility. Using the RG convention (cf. (6.352), (6.277)) and the Duhamel bound (cf. (6.311)), *irrelevant* symmetry violations ($\Delta > d + z$) decay as powers of ℓ. Consequently, on plateau regions (cf. Definition 6.297) the law-side visibility defects vanish with exponent $\Delta - (d + z)$ uniformly in scale (cf. the stability bounds Proposition 6.298 and the IR scaling relations (6.352), (6.277)). Explicitly, for the spectral block (and analogously for Π_{vac} and Π_{corr}, using the block-diagonality of Proposition 6.257),

$$\Big|1 - \Lambda_{\text{law}}^{(\text{spec})}(\ell;\varphi,\theta)\Big| \;\le\; C(\varphi,\theta)\sum_{k}\Big|g_k(1/\ell)\Big| \;=\; \mathcal{O}\Big(\ell^{-(\Delta-(d+z))}\Big) \quad (\ell \uparrow \infty),$$

i.e. $\mathcal{O}\big(\ell^{\Delta-(d+z)}\big)$ in the $\ell \downarrow 0$ convention. For the vacuum block one likewise finds $|\Lambda^{(\mathrm{vac})}_{\mathrm{law}}(\ell)| \to 1$ as $\ell \downarrow 0$ (cf. (6.347)), with the same power counting; the associated asymptotic exponents read off from $\mathcal{V}^{(\alpha)}$ are ξ-invariant by Proposition 6.276 (see also (6.318)).

Expectation-level fields and the IR manifold. After lifting by the holosymmetric functor (cf. Proposition 6.257) and coarse-graining, every local operator is mapped to an *expectation-level field*. Concretely, for a multiplet of relational operators $\hat{\mathbf{\Phi}}(x) = \{\hat{\Phi}^a(x)\}_a$ we recall

$$\Phi^a_{\mathrm{IR}}(x;\ell,\lambda_{\mathrm{UV}}) \;:=\; \Big\langle \overset{\circ}{\mathbf{\Psi}}\Big|\, \mathfrak{H}_{\lambda_{\mathrm{UV}},\ell}\Big[\hat{\Phi}^a(x)\Big]\,\Big|\overset{\circ}{\mathbf{\Psi}}\Big\rangle, \qquad a=1,\dots,N,$$

where $|\overset{\circ}{\mathbf{\Psi}}\rangle$ is the GNS cyclic vector (cf. Section 6.1.28). The image of this map defines a smooth *IR configuration manifold*

$$\mathcal{M}_{\mathrm{IR}} \;:=\; \mathrm{Im}\Big(\,\hat{\mathbf{\Phi}} \;\mapsto\; \mathbf{\Phi}_{\mathrm{IR}}(\,\cdot\,;\ell,\lambda_{\mathrm{UV}})\Big)\,,$$

on which slow collective variables φ^i (moduli, order-parameter coordinates, etc.) provide local charts. Plateau regularity is controlled by Proposition 6.298, and the relational chrono-geometry/connection come from Definition 6.7 together with the Christoffel formula (6.22); IR transport is encoded by the geodesics of $\nabla^{(\mathrm{rel})}$. Visibility functionals pull back to smooth IR potentials as in (6.302), with large-ℓ improvement governed by the IR scaling relations Section 6.12.5, (6.277).

Plateau regularity and RG transport. On any plateau block $\mathcal{B}$ (cf. Definition 6.297), the field map (6.354) is scale-stable as stated in Proposition 6.298. Explicitly,

$$\sup_{(u,v)\in \ln\mathcal{B}} \Big\|\partial_v \mathbf{\Phi}_{\mathrm{IR}}(x;\ell,\lambda_{\mathrm{UV}})\Big\| \;\le\; \varepsilon_\ell, \qquad \sup_{(u,v)\in \ln\mathcal{B}} \Big\|\partial_u \mathbf{\Phi}_{\mathrm{IR}}(x;\ell,\lambda_{\mathrm{UV}})\Big\| \;\le\; \varepsilon_{\lambda_{\mathrm{UV}}},$$

with $u = \ln\lambda_{\mathrm{UV}}$, $v = \ln\ell$. In particular, along RG trajectories generated by $\beta^i(\varphi)\partial_{\varphi^i}$ (cf. (6.352)) one has $\dot\varphi^i = \beta^i(\varphi)$ and, on plateaus, $\beta^i(\varphi_\star) \approx 0$, so the φ^i evolve adiabatically. This stability is compatible with the block-diagonal, cone-preserving holosymmetric lift (cf. Proposition 6.257) and underlies the uniform control of the visibility functionals in (6.302) and the scaling relations (6.352), (6.277).

Induced chronogeometry and relational connection. The unified expectation tensor $\bar\Sigma_{\mu\nu}$ induces on $\mathcal{M}_{\mathrm{IR}}$ both a metric and a torsion-free, metric-compatible *relational* connection (see Definition 6.7 and (6.22)):

$$\bar g_{\mu\nu}(x) := \mathsf{P}_{\mathrm{geom}}\Big[\bar\Sigma_{\mu\nu}\Big](\mathbf{\Phi}_{\mathrm{IR}}(x))\,, \qquad \nabla^{(\mathrm{rel})}_\alpha \bar g_{\mu\nu} = 0. \tag{3}$$

Slow configurations $\gamma : s \mapsto \varphi^i(s)$ follow geodesics on $\mathcal{M}_{\mathrm{IR}}$ up to subprincipal corrections, as encoded by the relational Levi–Civita connection Definition 6.7, with Christoffel symbols given explicitly in (6.22):

$$\frac{D^2\varphi^i}{Ds^2} \;+\; \Gamma^i{}_{jk}(\varphi)\,\frac{D\varphi^j}{Ds}\frac{D\varphi^k}{Ds} \;=\; \mathcal{O}\big(\varepsilon_\ell,\varepsilon_{\lambda_{\mathrm{UV}}}\big),$$

where $\Gamma^{i}{}_{jk}$ is the Levi–Civita connection of the moduli-space metric G_{ij} induced, for instance, from connected two-point functions after the holosymmetric lift (cf. Proposition 6.257). By Nielsen/BRST gauge-parameter independence (Proposition 6.276; see also (6.318)), the plateau loci and the exponents extracted from these IR transports are ξ-invariant.

Visibility functionals as IR potentials. Channel-restricted law-side visibilities (Part 2) pull back to smooth functionals on $\mathcal{M}_{\mathrm{IR}}$:

$$\mathcal{V}^{(\alpha)}(\varphi;\ell) \;:=\; 1-\left|\Lambda^{(\alpha)}_{\mathrm{law}}(\ell)\right| \qquad \Big(\text{cf. (6.347)}\Big),$$

and, for irrelevant deformations, satisfy the power-law improvement from Part 6,

$$\mathcal{V}^{(\alpha)}(\varphi;\ell) \;=\; \mathcal{O}\Big(\ell^{-(\Delta-(d+z))}\Big) \quad \text{as } \ell\uparrow\infty, \qquad \text{uniformly on plateaus; cf. (6.352), (6.277).}$$

By Nielsen (gauge-parameter) independence (Part 4; Proposition 6.276), both the plateau loci and the asymptotic exponents extracted from $\mathcal{V}^{(\alpha)}$ are ξ-invariant (see also (6.325)). Along adiabatic loops in slow-parameter space, the spec-channel phase of $\Lambda^{(\mathrm{spec})}_{\mathrm{law}}$ reproduces the Berry/WZ integral without altering the principal cones (Part 5; Theorem 6.261). In summary, $\mathcal{M}_{\mathrm{IR}}$ provides the macroscopic geometric stage on which the diagnostics from Parts 2–6 live: fields (6.362) evolve quasi-geodesically with respect to the relational Levi–Civita connection Definition 6.7 with coefficients (6.22), and the visibility functionals (6.302) obey the plateau/improvement bounds of Proposition 6.298 and the scaling relations (6.352), (6.277).

Algebraic–to–Hilbert-space bridge (GNS) and the concrete operator arena.

- **Representation-independent start.** Begin with the abstract *-algebra $\mathcal{A}$ of operations and relations (no representation chosen). The concrete operator arena used later for visibilities (see (6.347)–(6.348)) and for the holosymmetric lift (see Proposition 6.257) will be obtained functorially from $\mathcal{A}$ via GNS.

- **GNS construction of the "quantum holo-state."** Given a pure state functional $\omega^{\mathring{\mathbf{\Psi}}}: \mathcal{A}\to\mathbb{C}$, the GNS theorem (cf. Section 6.1.28) yields $(\mathcal{H}_{\mathrm{em}}, \pi^{\mathring{\mathbf{\Psi}}}, |\mathring{\mathbf{\Psi}}\rangle)$ with

  $$\omega(A) = \Big\langle \mathring{\mathbf{\Psi}} \,\Big|\, \pi(A) \,\Big|\, \mathring{\mathbf{\Psi}} \Big\rangle, \qquad A\in\mathcal{A}.$$

  Feeding $\pi(A)$ through the holosymmetric lift (cf. Proposition 6.257) and then taking expectations produces the expectation-level fields used on the IR manifold, canonically organized by (6.364) (cf. Definition 6.286).

- **Destination: $\mathrm{End}(\mathcal{H}_{\mathrm{em}})$.** The image $\pi(\mathcal{A})\subset\mathcal{B}(\mathcal{H}_{\mathrm{em}}) =: \mathrm{End}(\mathcal{H}_{\mathrm{em}})$ is the concrete operator arena where the relational observables (e.g. $\hat{\mathbf{\Phi}}(x)$) live. Channel projectors $\{\Pi_\alpha\}$ from the holosymmetric framework act on $\mathcal{H}_{\mathrm{em}}$ to define the channel-restricted visibilities (6.347)–(6.348). Block-diagonality and cone preservation follow from Proposition 6.257 together with the relational connection $\nabla^{(\mathrm{rel})}$ from Definition 6.7 and (6.22); plateau/exponent robustness is ensured by the Nielsen/BRST results Proposition 6.276 (cf. (6.318)).

Emergent chronogeometry and the relational Levi–Civita derivative

- **Chain of emergence.** From the quantum "globe" $\mathcal{X}_q$ and the unified expectation tensor $\bar{\Sigma}_{\mu\nu}$, construct the relational connection $\nabla^{(\mathrm{rel})}_\mu$ (see Definition 6.7); its coefficients are given by (6.22). This is the geometric substrate on which the expectation-level dynamics are read (cf. Definition 6.286, (6.364)) together with the plateau diagnostics Definition 6.297 and Proposition 6.298.

- **Geodesics & effects.** Slow configurations $\gamma : s \mapsto \varphi^i(s)$ evolve along IR geodesics up to subprincipal corrections, governed by the relational Levi–Civita connection Definition 6.7 with symbols (6.22):

$$\frac{D^2\varphi^i}{Ds^2} + \Gamma^i{}_{jk}(\varphi)\,\frac{D\varphi^j}{Ds}\,\frac{D\varphi^k}{Ds} \;=\; \mathcal{O}\big(\varepsilon_\ell, \varepsilon_{\lambda_{\mathrm{UV}}}\big).$$

  Gradients of the temporal tensor $\bar{t}_{\mu\nu}$ (see Definition 6.7 and (6.22)) generate controlled departures from GR-like transport in regimes of strong coherence (e.g. near black-hole or early-universe conditions). The principal cones remain unchanged under adiabatic loops; only intra-cone phases adjust, consistent with the spectral-channel Berry/WZ structure in Theorem 6.261 and the pointwise BRST/Nielsen independence Proposition 6.276 (cf. (6.318)).

Chapter 6 lays the groundwork for dynamics on the emergent chronogeometry by specifying the fundamental field content and the operational structures that govern it. The generalized field doublet $\hat{\mathbf{F}}(x) = (\hat{\mathbf{\Phi}}(x), \hat{\mathbf{\Theta}}(x))^{\mathsf{T}}$ provides the basic degrees of freedom (Definition 6.1), whose correlations are organised by the Unified Inter–Modal Correlator $\hat{\mathbf{G}}_\Lambda(x, x')$ with gauge–covariant transport (Definition 6.4, Section 4.19.1, and eq. (4.11)) [30, 29, 36, 37, 45]. Calculus is performed with the Relational Levi–Civita derivative $\nabla^{(\mathrm{rel})}$, compatible with the composite chronogeometric response $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (Definition 6.7) [45], and actions are integrated against the state–responsive relational volume element $d\mu = \sqrt{-\det \bar{g}^{\mathrm{Lor}}}\, d^d x$ with $\bar{g}^{\mathrm{Lor}}_{\mu\nu} := \bar{g}_{\mu\nu} + \bar{t}^{\mathrm{sym}}_{\mu\nu}$ (Definition 6.33) [45]. The geometric sector is encoded by an operator–valued Lagrangian density $\mathcal{L}_\Phi$ and its action S_Φ (Definitions 6.172 and 6.174) [36, 37], while the full energy bookkeeping is handled by the Relational Hamiltonian $\hat{H} = \hat{H}_\Phi + \hat{H}_{\Phi\Theta} + \hat{H}_\Theta$ (Definition 6.54 and eq. (6.109)) [30, 29]. Vacua are then identified either variationally via the action or spectrally via $\hat{H}$, yielding the Vacuum Manifold $\mathcal{M}_{\mathrm{vac}}$ that anchors geometric tangents and excitation fibres for subsequent analysis (Section 6.1.17 and Definition 6.61) [30, 29].

Beyond these constructions, the chapter equips the framework with a methodological toolkit for extracting physics. Gauge redundancy is treated cohomologically through BRST quantisation—ensuring nilpotency, gauge–parameter independence, and consistency tests for anomalies (Propositions 6.213, 6.214 and 6.217 and Principles 6.1 and 6.2) [52, 109, 110, 111]. Symmetry organisation is clarified by the duality–quadrant schema, which separates law–level from state–level symmetry via commuting lifts and classifies regimes as Unbroken, SSB, Emergent, or Explicit Breaking (Definitions 6.220, 6.221 and 6.224, Proposition 6.225, Theorem 6.231, and Corollary 6.232) [52, 29]. Operational diagnostics are provided by holosymmetric,

two–path coherence functionals Λ_{law} and Λ_{state}, which are gauge–stable and quadrant–preserving (Definitions 6.237 and 6.238, Lemmas 6.243, 6.247 and 6.248, Proposition 6.251, and Theorem 6.253) [22, 23, 24, 25]. Finally, a scale–aware strategy—UV/IR scale–tagging and the plateau criterion—ensures that claims are robust across reciprocal resolution trade–offs and insensitive to gauge choices (Definitions 6.278, 6.280, 6.282, 6.289, 6.293, 6.295 and 6.297, Propositions 6.294, 6.296, 6.298 and 6.299, and Corollary 6.265) [36, 37, 30, 29].

Chapter 7 Field Laws and Physical States

Physical sector and kernels. Chapter 7 works entirely on the *physical sector*—the BRST cohomology $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$—and at fixed *scale tags* $(\lambda_{\text{UV}}, \ell)$ [52, 112]. It expands the (gauge-fixed) action to quadratic order around a symmetric reference vacuum $(\bar{\Phi}, \bar{\Theta})$ and packages fluctuations as

$$\delta\hat{\mathbf{U}}_{\lambda_{\text{UV}},\ell} := \left(\delta\hat{\mathbf{\Phi}}_{\lambda_{\text{UV}},\ell} \,,\, \delta\hat{\mathbf{\Theta}}_{\lambda_{\text{UV}},\ell} \right)^{\mathsf{T}}.$$

This defines the *gauge-fixed quadratic kernel* $\mathcal{K}_{\Psi}^{(\lambda_{\text{UV}},\ell)}$ and, after projection, the *physical kernel*

$$\mathcal{K}_{\text{phys}}^{(\lambda_{\text{UV}},\ell)} := \mathsf{P}_{\text{phys}}\, \mathcal{K}_{\Psi}^{(\lambda_{\text{UV}},\ell)}\, \mathsf{P}_{\text{phys}},$$

whose inverse (retarded/advanced/Feynman) is the *physical propagator* $\mathcal{D}_{\text{phys}}^{(\lambda_{\text{UV}},\ell)}$ [22] (Definitions 6.274 and 7.15). On $\mathcal{H}_{\text{phys}}$ one has the *cohomological inversion* $\mathcal{K}_{\text{phys}} \cdot \mathcal{D}_{\text{phys}} = \mathbf{1}$, and—crucially—the *pole locations and residues* of $\mathcal{D}_{\text{phys}}$ are *independent of the gauge parameter* (either ξ generically or α_{gf} in the background-field method), since quartet/longitudinal pieces decouple and Slavnov–Taylor/Nielsen identities render gauge-parameter shifts BRST-exact (Proposition 7.16) [112, 38, 109, 113]. For background geometric and measure structures used throughout, see the relational calculus and volume element of Chapter 6 (Definitions 6.7 and 6.33) and the chronogeometric response from Chapters 4–5 (Definition 5.26).

BFM Ward identities and Nielsen control on the band. To control gauge symmetry cleanly, the chapter adopts the *Background-Field Method (BFM)* and imposes background Ward identities together with a *band-limited Nielsen identity* for the 1PI functional Γ [114, 112]. Two key conclusions follow: (i) *pole masses are gauge-parameter independent*; (ii) *on-shell visibilities* are protected. Both hold *on the band*—i.e., after applying the UV projector $P_x(\lambda_{\text{UV}})$ and IR window $K_{\text{IR}}^{(\ell)}$—and remain *robust under admissible regulator changes* within tolerance ε_{reg} [112, 38, 109] (eqs. (7.201), (7.207) and (7.209) and Section 7.3.7). For the UV/IR operators themselves, see the scale-tagging constructions of Chapters 6–7 (Definitions 6.278, 6.280, 6.282 and 6.289).

Band Callan–Symanzik and BRST compatibility. With the *scale-tagged equations of motion* held fixed (eqs. (6.333) and (6.334)), the *Callan–Symanzik equation* holds *on the band* for any renormalised 1PI n-point function $\Gamma_{\text{band}}^{(n)}$, with the usual mass insertions when present [32, 112] (eq. (7.213)). A remark clarifies compatibility with Slavnov–Taylor/Nielsen identities and the band EOM: the UV/IR maps carry ghost number 0, commute with BRST, and thus

preserve the algebraic constraints on visibilities [52, 109]. (For context from Chapter 6: the *scale-tagged EOM* arise by inserting $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ into the operator equations and then projecting to $\mathcal{H}_{\mathrm{phys}}$; their BRST-closedness yields *gauge-parameter invariance in cohomology* at fixed scales, cf. eqs. (6.333) and (6.334) and Definition 6.270.)

UV/IR knobs and plateau certification. The chapter then formalises *scale control* and an *acceptance test* for *plateau stability*. All subsequent constructions are declared to act *on the band* and to be compatible with BRST and the scale-tagged EOM [112] (eqs. (6.333), (6.334) and (6.374)). Three quantitative *plateau metrics*—Slope, Curvature, Oscillation—are defined with a pass criterion: for each audited observable O, the metrics must lie below user-chosen tolerances $\tau_O^{(1)}$, $\tau_O^{(2)}$, $\tau_O^{(\mathrm{osc})}$ throughout a 2D region of $(\lambda_{\mathrm{UV}}, \ell)$. If all audited observables satisfy these bounds, the bandpoint is declared *plateau-stable* and can be used for mode identification and cross-quadrant comparisons (eq. (7.28) and Section 7.3).

Pole extraction and certification workflow. Operationally, *pole extraction* uses *Riesz projectors* on the transverse–traceless kernel $\Gamma^{(\mathrm{TT})}(p^2)$: encircle a (possibly degenerate) pole at $p^2 = -m_i^2$ with a loop γ_i, form

$$\mathbb{P}_i = \frac{1}{2\pi i} \oint_{\gamma_i} \left(\Gamma^{(\mathrm{TT})}(z)\right)^{-1} dz, \qquad \widehat{\Gamma}_i := \mathbb{P}_i \, \Gamma^{(\mathrm{TT})} \, \mathbb{P}_i,$$

and read off m_i^2 from the eigenvalues of $\widehat{\Gamma}_i$ [30]. By the band Nielsen identity, the *pole positions* are α-independent (residues may vary), and the certification step declares *pass/fail* for a bandpoint $(\lambda_{\mathrm{UV}}, \ell)$ by checking algebraic items (C1–C6) together with the plateau metrics [114, 112] (Section 7.3).

Quadrant-wise readings (QI baseline; QIV synopsis). For *Quadrant I* (unbroken laws, symmetric state), mixing respects spin and the BRST projection leaves only *transverse, positive-norm* modes in $\mathcal{H}_{\mathrm{phys}}$ (no ghosts, no tachyons) [52, 112]. QI thus provides a *robust, gauge-independent, scale-stable baseline* against which any sharper particle-level tagging (e.g., "graviton", "Higgs", "apeiron", "chrono-kymon") must be validated on a certified plateau (Sections 7.2.6 and 7.3 to 7.5). For *Quadrant IV* (explicit breaking; asymmetric state), the adjoint-covariant Ward identity on $\mathcal{H}_{\mathrm{phys}}$ acquires a *non-vanishing source* proportional to explicit spurions $\{\epsilon_k\}$ (eq. (7.665)), so Goldstone's theorem does not apply—there are *no exact NG poles* [38]. Instead, would-be NG modes develop *small but nonzero gaps* governed by a *GMOR-type identity*,

$$f_{\pi_b}^2 \, m_{\pi_b}^2 = \sum_k \epsilon_k \left\langle \mathcal{S}_{k,b}^{(\lambda_{\mathrm{UV}})} \right\rangle_{\mathbf{\Psi}} + \mathcal{O}(\epsilon^2),$$

hence $m_{\pi_b}^2 \propto \epsilon$ with vanishing intercept within plateau tolerance [112, 32]. The *splitting pattern* across light multiplets is fixed by the spurion irrep's projection onto broken directions; in practice one isolates a near-degenerate cluster with a Riesz projector, follows its analytic subspace, and uses *Kato parallel transport* for eigenframe labelling [30] (Theorem 7.630, Corollary 7.631, Definitions 7.559 and 7.618, Lemmas 7.566 and 7.619, and Propositions 7.568 and 7.620).

Scale-tagged, BRST-consistent equations on $\mathcal{H}_{\text{phys}}$. Chapter 7 formalises how the *scale-tagged* equations of motion—after UV/IR tagging and channel projection—descend to the physical Hilbert space and become *gauge-parameter independent* via band-limited Nielsen identities on $\mathcal{H}_{\text{phys}}$ [112, 38] (Definition 6.272 and eqs. (6.333), (6.334), (6.374) and (7.201)). Concretely, with the physical projector P_{phys} and BRST differential ζ,

$$\mathsf{P}_{\text{phys}}(\mathcal{E}_\bullet + \zeta\,\chi_\bullet)\,\mathsf{P}_{\text{phys}} = 0,$$
$$\partial_\xi[\mathsf{P}_{\text{phys}}(\mathcal{E}_\bullet + \zeta\,\chi_\bullet)\,\mathsf{P}_{\text{phys}}] = 0.$$

(Similarly, $\partial_{\alpha_{\text{gf}}} = 0$.) See [52, 112]. Here the ζ-exact pieces vanish in BRST cohomology, so the *clean forms* $\mathsf{P}_{\text{phys}}\mathcal{E}_\bullet\mathsf{P}_{\text{phys}} = 0$ hold and are independent of gauge parameters (Nielsen safety on the band; see eq. (7.209) and the discussion following eq. (7.201); cf. also Definition 6.270) [114, 38].

Assuming the UV projector $P_x(\lambda_{\text{UV}})$ and IR window $K_{\text{IR}}^{(\ell)}$ are BRST-closed and commute with the channel projectors $\{\Pi_\alpha\}$, every scale-tagged object produced by the *band functor* $\mathfrak{B}_{\lambda_{\text{UV}},\ell}$ (Definitions 6.278, 6.280, 6.282 and 6.289) *descends* to $\mathcal{H}_{\text{phys}}$, and all channel-resolved observables inherit *gauge-parameter independence* [52, 114, 109, 112]. Symbolically,

$$\mathsf{P}_{\text{phys}}\,\mathfrak{B}_{\lambda_{\text{UV}},\ell}[\cdot]\,\mathsf{P}_{\text{phys}} = \mathfrak{B}_{\lambda_{\text{UV}},\ell}\Big[\mathsf{P}_{\text{phys}}(\cdot)\,\mathsf{P}_{\text{phys}}\Big], \qquad \partial_\xi,\ \partial_{\alpha_{\text{gf}}} \text{ kill physical readouts.}$$

See Definition 6.272 for the precise construction on the band, Definition 6.275 for channel-resolved visibilities on $\mathcal{H}_{\text{phys}}$, and Section 6.12.4 for the plateau criterion that underwrites scale-stable interpretation.

UV/IR knobs, band preparation, and self-adjointness. After restricting dynamics to the physical sector (Stage-I, BRST-consistent regime), the generator $\mathbb{K}(\epsilon)$ preserves $\mathcal{H}_{\text{phys}}$ and remains self-adjoint on a common domain $\mathcal{D}$ [115]. The UV/IR taggers—local spectral projector $P_x(\lambda_{\text{UV}})$ and IR window $K_{\text{IR}}^{(\ell)}$—are chosen so that they commute with $\mathbb{K}(\epsilon)$ and with the channel projectors (the conditions listed in Section 7.3; cf. Definitions 6.278, 6.280, 6.282 and 6.289) [114, 52, 109]. Consequently, the *band-prepared* generator

$$\mathbb{K}^{(x)}(\epsilon) := K_{\text{IR}}^{(\ell)}\,P_x(\lambda_{\text{UV}})\,\mathbb{K}(\epsilon)\,P_x(\lambda_{\text{UV}})\,K_{\text{IR}}^{(\ell)}$$

is again self-adjoint on $\mathcal{D}$ [115, 116]. The *band Hessian* $\mathbb{H}^{(x)}(\epsilon)$—the second variation of the band action around the vacuum—obeys the standard quadratic-form criteria: under the locality/polynomial hypotheses used for the band constructions, the perturbations are form-bounded with relative bound < 1, and by KLMN/Kato–Rellich one obtains a *unique self-adjoint* operator on the same form (hence operator) domain [115, 30] (Section 7.3; see also the preparation of scale-tagged equations in eqs. (6.333), (6.334) and (6.374)).

Plateau geometry in the log-chart and equivalence of definitions. Chapter 7 fixes the parameter-space *log-chart*

$$(u, v) = (\ln\lambda_{\text{UV}}, \ln\ell), \qquad g_{\text{plat}} := du^2 + dv^2, \qquad D_\lambda = \partial_u = \lambda_{\text{UV}}\partial_{\lambda_{\text{UV}}}, \quad D_\ell = \partial_v = \ell\,\partial_\ell,$$

and defines the gradient/Hessian operators used to test plateau stability for C^2 observables [112, 32] (Section 7.3). Two plateau criteria are then shown to be *equivalent (with matched tolerances)*: a channelwise/directional version and a geometric version phrased in terms of gradient smallness and local oscillation bounds on geodesic balls $B_r^{\mathrm{plat}}(\lambda_{\mathrm{UV}}, \ell)$ in the metric g_{plat}. The text proves both inclusions under the stated identification of neighbourhoods and tolerances, establishing operational equivalence of the tests and providing the quantitative thresholds used later for certification (Sections 6.12.4 and 7.3 and eq. (7.28)).

Quadrant logic on a certified plateau $\mathfrak{P}$ Using the *visibilities* $(\Lambda_{\mathrm{law}}, \Lambda_{\mathrm{state}})$ (cf. Definitions 7.35 and 7.42), a *decision rule* $\mathrm{Quad}(\Lambda_{\mathrm{law}}, \Lambda_{\mathrm{state}})$ classifies the plateau into:

QI: $|\Lambda_{\mathrm{law}}| \approx 1$ and $|\Lambda_{\mathrm{state}}| \approx 1$.

QII: law conserved ($|\Lambda_{\mathrm{law}}| \approx 1$), but state decoheres: $\Lambda_{\mathrm{state}}(V) \to 0$ as $V \to \infty$.

QIII: "IR healing" of apparent law deficits: $1 - |\Lambda_{\mathrm{law}}(\ell)| = \mathcal{O}(\ell^{\Delta-(d+z)})$, while $|\Lambda_{\mathrm{state}}| \approx 1$.

QIV: persistent law deficit: $|\Lambda_{\mathrm{law}}| \le 1 - \delta_\star$ on $\mathfrak{P}$ for some $\delta_\star \in (0,1)$, with $\Lambda_{\mathrm{state}}(V) \to 1$.

Here "≈ 1" means $\big||\cdot| - 1\big| \le \varepsilon_{\mathrm{tol}}$ uniformly on $\mathfrak{P}$ in the log-chart variables u, v. This rule makes Ward-type statements and spectral data *readable across scales* on a certified plateau (cf. Section 6.12.4).

Integrated readouts and their inheritance properties. Given channel readouts $R_\alpha(u, v)$, Chapter 8 introduces *integrated readouts* $\overline{R}_\alpha$ (and $\overline{\overline{R}}_\alpha$) obtained by averaging against a weight w supported inside a certified plateau $\mathfrak{P}$ (cf. the plateau definition and certification in Section 6.12.4 and eq. (7.28)). On a certified plateau, the *same* gradient/curvature/oscillation bounds *carry over* to the integrated objects—with equal or better constants:

$$\big|\partial_v \overline{R}_\alpha\big| \le \varepsilon_{\mathrm{grad}}, \qquad \big\|\nabla^2 \overline{R}_\alpha\big\| \le \varepsilon_{\mathrm{curv}}, \qquad \mathrm{osc}\,[\overline{R}_\alpha] \le \varepsilon_{\mathrm{osc}},$$

and analogously in u. Discrete counterparts follow by replacing integrals with convex combinations (see eqs. (7.413) and (7.414)).
The *integrated vs. pointwise diagnostic* at a reference point $(u_*, v_*) \in \mathrm{supp}(w)$,

$$\Delta_\alpha := \big|\overline{R}_\alpha - R_\alpha(u_*, v_*)\big|,$$

is $\mathcal{O}(\varepsilon_{\mathrm{plat}})$ by the inheritance lemma together with a mean-value estimate *within* $\mathfrak{P}$; reports should include Δ_α and the weight's shape/support (cf. eq. (7.390)).

Discrete IR window and curvature-based plateau test (error control). For a grid v_k with mesh h and a smoothed curvature proxy $\phi_h = \mathsf{LP}_{\tau,h}\,\mathsf{D}_2 f$, one has the uniform second-order *consistency*

$$\phi_h(k) = \phi(v_k) + \mathcal{O}(h^2), \qquad \sup_k \big|\phi_h(k) - f''(v_k)\big| = \mathcal{O}(h^2) + \mathcal{O}(\tau^2),$$

using the bias expansion with an even kernel W of finite $m_2(W)$. Defining the continuous IR window $\mathcal{I}_{\mathrm{IR}} = \{\, v : \ |\phi(v)| \le \varepsilon_{\mathrm{curv}} \,\}$ and its *grid* counterpart $\mathcal{I}_{\mathrm{IR}}^{(h)} = \{\, k : \ |\phi_h(k)| \le \varepsilon_{\mathrm{curv}} \,\}$, the chapter proves a *band-measure control*

$$\mu\Big(\mathcal{I}_{\mathrm{IR}} \,\triangle\, \mathcal{I}_{\mathrm{IR}}^{(h)}\Big) \le C\,\delta_\phi = \mathcal{O}(h^2),$$

under Lipschitz regularity of ϕ on the operational window (cf. Proposition 7.397).

Ward-operator bounds and Nielsen projection. Expanding the *Ward operator* around the symmetric point and evaluating on-shell yields a linear-response representation with *band-limited kernels* $\mathcal{K}_A^a(p;\ell)$. Taking operator norms on the compact band $\mathfrak{B}(\lambda_{\mathrm{UV}},\ell)$ gives a uniform bound

$$\mathcal{W}(\ell) \ \le \ C(\lambda_{\mathrm{UV}},\ell)\sum_A \Big|\varepsilon_A(\kappa/\ell)\Big| + \mathcal{O}(\|\varepsilon\|^2),$$

where $C = \sum_A C_A$ encodes band suprema of the kernel/reference-operator ratio. Passing to *Nielsen-projected eigen-couplings*—a change of basis R that Jordanises the projected stability matrix on $W = \mathrm{im}\,\mathcal{N}$—improves interpretation and stabilises the flow (cf. Definitions 7.454 and 7.457).

Joint (multi-channel) integrated plateaus. For a set of channels $\mathfrak{C} \subset \{\mathrm{law}, \mathrm{state}, \mathrm{mix}, \ldots\}$ and averaging widths $(\Delta\lambda, \Delta\ell)$ (or $(\Delta u, \Delta v)$), the *joint integrated plateau* is

$$\overline{\mathfrak{P}}_{\mathfrak{C}}(\Delta\lambda,\Delta\ell) := \bigcap_{\alpha\in\mathfrak{C}} \Big\{ (\lambda_{\mathrm{UV}},\ell) \in \mathcal{R} : \ \overline{\Lambda}_\alpha \text{ satisfies Section 7.3.5 with prescribed tolerances} \Big\},$$

with inheritance-consistent tolerances per α. It *includes* the pointwise joint plateau when the averaging box stays inside the test ball and tolerances match; conversely, under a $C^{1,1}$ assumption on each channel and small enough half-sizes $\Delta u/2, \Delta v/2 \le r$, one gets *reverse inclusion* after *rescaling* tolerances using channel Lipschitz data (cf. eq. (7.64)).

Operational process.

1. *Prepare the band* with BRST-closed taggers $P_x(\lambda_{\mathrm{UV}})$, $K_{\mathrm{IR}}^{(\ell)}$; restrict to $\mathcal{H}_{\mathrm{phys}}$; ensure self-adjoint $\mathbb{K}^{(x)}$ and $\mathbb{H}^{(x)}$. (cf. Section 7.1.4 and Definition 6.282)

2. *Certify a plateau* in the log-chart by gradient, curvature, oscillation tolerances on B_r^{plat}; use the geometric/channelwise equivalence to choose the most convenient test. (cf. Definition 7.47 and Section 6.12.4)

3. *Classify the physics* on $\mathfrak{P}$ via $\mathrm{Quad}(\Lambda_{\mathrm{law}}, \Lambda_{\mathrm{state}})$ (QI–QIV) to read Ward statements and spectra consistently across scales. (cf. Definition 7.685)

4. *Report integrated readouts* $\overline{R}_\alpha$; include the *diagnostic gap* Δ_α and the averaging weight/-support, knowing the plateau tolerances inherit. (cf. Lemma 7.345 and eq. (7.418))

5. *Discrete implementation*: use the curvature-window method; budget $\mathcal{O}(h^2)$ (mesh) and $\mathcal{O}(\tau^2)$ (smoothing) errors; control the symmetric-difference measure of the IR window. (cf. Propositions 7.382 and 7.397)

6. *Linear response*: bound $\mathcal{W}(\ell)$ by band-kernels times small couplings; optionally *Nielsen-project* to eigen-couplings for sharper flow control. (cf. Definition 7.454)

Quadrant Analysis Strategy. Chapter 7 outlines a systematic, scale-aware strategy (Section 6.12.5) using the variational backbone (Section 6.12.2) and operational diagnostics (Section 6.12.3) to analyze each quadrant. All analyses are performed within the physical sector $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$ (Remark 6.226), ensuring gauge invariance and relying on plateau stability (Section 6.12.4) for robust results. Mode identifications (like "Higgs-like", "graviton-like") remain provisional until confirmed stable on a plateau (Definition 6.300).

QI (law symmetric, state symmetric) Goal (baseline). On a certified plateau $\mathfrak{P}$, QI is the fully coherent reference case: both visibilities satisfy $\left|\Lambda_{\text{law}}\right| \approx 1$ and $\left|\Lambda_{\text{state}}\right| \approx 1$ (tolerance semantics on $\mathfrak{P}$ in the $(u, v) = (\ln \lambda_{\text{UV}}, \ln \ell)$ chart); this is the calibration baseline for all other quadrants, as fixed by the canonical decision rule Definition 7.685 and Remark 7.686. (Log-chart/plateau geometry used throughout is defined in Definition 7.47 and Section 6.12.4.)

Checks

- *Scale-tagged physical EOMs (clean, gauge-independent).* On $\mathcal{H}_{\text{phys}}$ the BRST-exact pieces drop out and the "clean forms" hold,

$$\mathsf{P}_{\text{phys}}\big(\mathcal{E}_\bullet + \zeta\, \chi_\bullet\big)\, \mathsf{P}_{\text{phys}} = 0, \qquad \partial_\xi(\cdot) = 0 \quad \text{(Nielsen safety)},$$

  so the physical equations are independent of gauge parameters/presentation choices on certified plateaus; see Section 7.1.3 (with inputs from Definition 6.272). Moreover the BRST-closed UV/IR taggers commute with channel projections, and the band functor descends:

$$\mathsf{P}_{\text{phys}}\ \mathfrak{B}_{\lambda_{\text{UV}},\ell}[\cdot]\ \mathsf{P}_{\text{phys}} = \mathfrak{B}_{\lambda_{\text{UV}},\ell}\Big[\mathsf{P}_{\text{phys}}(\cdot)\,\mathsf{P}_{\text{phys}}\Big],$$

  ensuring all scale-tagged objects/readouts are defined on $\mathcal{H}_{\text{phys}}$ (and remain gauge-independent); cf. Section 7.1.4 and Definition 6.282.

- *Ward/Slavnov–Taylor conservation in $\mathcal{H}_{\text{phys}}$.* The scale-tagged Ward identity on the physical sector reads, schematically,

$$\nabla_\mu^{(\text{ad})}\, \bar{J}^\mu_{a,\text{imp}} = \sum_k \epsilon_k\, \mathcal{S}^{(\lambda_{\text{UV}},\ell)}_{a,k} \quad \text{on } \mathcal{H}_{\text{phys}},$$

  with all right-hand terms BRST/Nielsen-safe. In QI (no explicit breaking in the law channel), $\epsilon_k = 0$ and conservation is exact: $\nabla_\mu^{(\text{ad})} \bar{J}^\mu_{a,\text{imp}} = 0$. Definition 7.567 and eq. (7.665).

- *Spectrum health from the physical Hessian.* Linearisation about the symmetric vacuum gives the physical Hessian $\mathbb{H}^{(\lambda_{\text{UV}},\ell)}_{\text{phys}}$ via the linearised Euler–Lagrange system eq. (6.332) (and, in the band-prepared setting, eq. (7.10)). On $\mathcal{H}_{\text{phys}}$ the two-point functions admit a Källén–Lehmann representation with positive weights; poles/residues are gauge-parameter independent, giving positivity (no ghosts) and a non-tachyonic spectrum under the stated hypotheses. eqs. (7.12), (7.14) and (7.15) and Corollary 7.24.

- *Holosymmetric visibilities at unity on* $\mathfrak{P}$. The (holosymmetric) visibilities are BRST/Nielsen-safe and normalised so that on QI plateaus they satisfy

$$\left|\Lambda_{\text{law}}(\ell)\right| \approx 1 \quad \text{and} \quad \left|\Lambda_{\text{state}}(V)\right| \approx 1.$$

  Definitions 6.275 and 7.685.

In Chapter 7 the Law-State quadrant means: clean, gauge-independent physical equations; exact Ward conservation on $\mathcal{H}_{\text{phys}}$; a positive, ghost-free, non-tachyonic spectrum from the physical Hessian; and visibilities pinned to unity on a certified plateau—precisely as encoded by the canonical quadrant rule.

QII (law symmetric, state asymmetric) Goal (broken phase, conserved law). On a certified plateau $\mathfrak{P}$, QII keeps *law*-level conservation intact while the *state* decoheres because a vacuum manifold is present. In the canonical decision rule this is recorded as $\left|\Lambda_{\text{law}}\right| \approx 1$ and $\Lambda_{\text{state}}(V) \to 0$ as $V \to \infty$; the remark explicitly ties the decoherence to the manifold of vacua that become orthogonal at large volume (cf. Definition 7.685 and Remark 7.686).

Vacuum structure and unbroken subgroup. The broken-phase bookkeeping is given in terms of the decomposition of the Lie algebra into unbroken versus broken pieces, with the counts

$$r_{\text{g}} = \dim \mathfrak{g}_{\text{gauge}} - \dim \mathfrak{h}_{\text{gauge}}, \qquad r_{\text{gl}} = r - r_{\text{g}},$$

and all statements to be evaluated on $\mathcal{H}_{\text{phys}}$ and certified plateaus. This encodes the unbroken subgroup (here, $\mathfrak{h}_{\text{gauge}}$ on the gauged side) and sets the stage for mode counting around any chosen vacuum on the manifold (cf. Definition 6.61).

Hessian analysis and mode content (Goldstone vs. BEH). Linearising about a symmetry-broken vacuum and reading the *band* Hessian on the physical sector gives the expected spectrum:

- unbroken gauged generators $\mathfrak{h}_{\text{gauge}}$: massless spin-1 vectors (physical polarisations after BRST projection);
- broken *global* generators: r_{gl} Nambu–Goldstone scalars (gapless in the $V \to \infty$ limit, with standard caveats for nonrelativistic/type-B cases);
- broken *gauged* generators: r_{g} massive vectors via Brout–Englert–Higgs reorganisation; the would-be NG modes are "eaten" and are not separate physical excitations.

All counts and readings are taken in $\mathcal{H}_{\text{phys}}$ on certified plateaus (cf. eq. (7.10) and Propositions 6.213 and 7.22).

Law channel (conservation) and quartet decoupling. In the band/IR-tagged setting the improved currents obey the scale-tagged Ward structure; with no explicit breakers in QII, law-level conservation survives on $\mathcal{H}_{\text{phys}}$. Ghosts, antighosts, Nakanishi–Lautrup fields and longitudinal gauge modes organise into Kugo–Ojima quartets and decouple, so physical poles come from the transverse blocks of the Hessian (cf. Definition 7.567, eq. (7.665), and Proposition 7.22).

State channel (preparation and ordered limits). The broken state is prepared by a seed that *does not* alter the symmetric operator content used by the law-channel meters; ordered

limits $\varepsilon \to 0^+$, then $V \to \infty$, then the plateau scan, produce distinct symmetry-broken branches that become orthogonal as $V \to \infty$. This realises $\Lambda_{\text{state}}(V) \to 0$ while keeping $\left|\Lambda_{\text{law}}\right| \approx 1$ (cf. Remark 7.686 and Definition 6.275).

Gauge/Nielsen safety and plateau robustness. With BRST-closed UV/IR tags, the dynamics descends to $\mathcal{H}_{\text{phys}}$ and physical spectra/matrix elements are gauge-parameter independent (Nielsen identities). Within the IR window used for audits, pole positions and residues have $\partial_\xi m_\alpha^2 = \partial_\xi Z_\alpha = 0$ (within tolerance). Plateau acceptance criteria (slope/curvature/oscillation budgets) gate the analysis (cf. Sections 6.12.4 and 7.1.3 and Definitions 6.272, 6.282 and 7.47).

Net QII signature (Chapter 8). On $\mathfrak{P}$: exact law conservation and ghost/longitudinal decoupling (law visibility at unity), a vacuum manifold that drives state-sector decoherence as $V \to \infty$ (state visibility to zero), and a spectrum with NG modes for broken global symmetries and BEH-massive vectors for broken gauged ones—each read in the band-prepared, BRST-physical sector with gauge-independent pole data (cf. Definitions 6.275 and 7.685).

QIII (law asymmetric, state symmetric): what Chapter 8 establishes. Goal (emergent symmetry). QIII captures cases where explicit microscopic breaking is present but *irrelevant* in the renormalisation–group sense, so the *law channel heals in the IR* while the *state channel remains symmetric* on a certified plateau $\mathfrak{P}$. In the canonical decision rule this reads

$$1 - \left|\Lambda_{\text{law}}(\ell)\right| \xrightarrow[\ell\to\infty]{} 0, \qquad \left|\Lambda_{\text{state}}(V)\right| \approx 1 \quad \text{(tolerance semantics on } \mathfrak{P}\text{)},$$

classifying the law as asymptotically exact while the state shows no SSB (cf. Definitions 6.275 and 7.685 and Section 6.12.4).

Checks (with operative scalings and plateau controls).

1. **Setup with irrelevant explicit breakers.** Work with a scale–tagged Lagrangian that includes explicit symmetry–breaking insertions

$$\sum_k \varepsilon_k \mathcal{O}_k, \qquad \Delta_k > d + z,$$

so each $\mathcal{O}_k$ is RG–irrelevant. The "relational Ward'' control sets the decay rate in the IR by the leading irrelevant dimension $\Delta := \min_k \Delta_k$ (cf. Lemma 6.233 and Definition 7.567).

2. **Ward–identity violation decays at a power law.** On the physical sector $\mathcal{H}_{\text{phys}}$, the scale–tagged Ward deviation $\Delta_a^{(\lambda_{\text{UV}},\ell)}$ obeys the IR law

$$\Delta_a^{(\lambda_{\text{UV}},\ell)} = \mathcal{O}\Big(\ell^{-(\Delta-(d+z))}\Big) \quad (\ell \to \infty),$$

equivalently $\mathcal{O}\Big(\mu^{\Delta-(d+z)}\Big)$ for $\mu \sim 1/\ell$ (cf. eq. (6.377) and Lemma 6.233).

3. **Law–coherence deficit heals with the same exponent.** The visibility deficit tracks the Ward decay:

$$1 - \left|\Lambda_{\text{law}}(\ell)\right| = \mathcal{O}\Big(\ell^{-(\Delta-(d+z))}\Big),$$

so the law channel tends to unity on $\mathfrak{P}$ (cf. Proposition 6.277 and Definition 7.685).

4. **State remains symmetric.** Within the certified IR window, order parameters vanish and the state visibility stays at unity (within tolerances):

$$\left|\Lambda_{\text{state}}(V)\right| \approx 1 \quad \text{on } \mathfrak{P},$$

with inheritance to integrated readouts $\overline{R}_\alpha$ (cf. Definition 6.275, Lemma 7.345, and eq. (7.390)).

5. **Plateau certification and discrete IR–window test.** Use the chapter's log–chart $u = \ln \lambda_{\text{UV}}$, $v = \ln \ell$ and metric g_{plat} to enforce gradient/curvature/oscillation tolerances (cf. Definition 7.47 and Section 6.12.4). For implementations on a grid v_k with mesh h, the smoothed curvature proxy $\phi_h = \mathsf{LP}_{\tau,h}\,\mathsf{D}_2 f$ satisfies

$$\phi_h(k) = \phi(v_k) + \mathcal{O}(h^2), \qquad \sup_k \left|\phi_h(k) - f''(v_k)\right| = \mathcal{O}(h^2) + \mathcal{O}(\tau^2),$$

and the symmetric–difference measure between the continuous and grid IR windows is controlled by

$$\mu\Big(\mathcal{I}_{\text{IR}} \,\triangle\, \mathcal{I}_{\text{IR}}^{(h)}\Big) = \mathcal{O}(h^2),$$

which underwrites robust QIII diagnoses on $\mathfrak{P}$ (cf. Propositions 7.382 and 7.397).

6. **Gauge/Nielsen safety.** Because the UV/IR taggers are BRST–closed and commute with channel projectors, all band–limited objects descend to $\mathcal{H}_{\text{phys}}$, and physical readouts are gauge–parameter independent:

$$\partial_\xi(\text{physical readout}) = 0, \quad \partial_{\alpha_{\text{gf}}}(\text{physical readout}) = 0,$$

within the plateau tolerances (cf. Section 7.1.3 and Definition 6.282).

Net QIII signature On a certified plateau $\mathfrak{P}$: (i) explicit breaking is present but irrelevant; (ii) the Ward deficit and the law–visibility deficit both decay as $\ell^{-(\Delta-(d+z))}$; (iii) the state channel remains symmetric, $\left|\Lambda_{\text{state}}\right| \approx 1$; (iv) all statements are BRST-/Nielsen–safe and stable under the chapter's plateau certification and discrete IR–window controls (cf. eq. (6.377), Propositions 6.277, 7.382 and 7.397, Definitions 6.275 and 7.47, and Section 7.1.3).

Quadrant IV vs. III — resolving the mismatch. Quadrant IV describes *persistent explicit breaking* driven by relevant/marginal spurions $\{\epsilon_k\}$, together with a *non-vanishing Ward deficit in the IR* and *state coherence* in a unique (non-symmetric) vacuum. In Chapter 7 this appears as an adjoint–covariant Ward identity with a source term proportional to the spurions (eq. (7.665)) and leads to GMOR–type gaps for the would-be NG modes, $f_{\pi_b}^2 m_{\pi_b}^2 = \sum_k \epsilon_k \langle \mathcal{S}_{k,b}^{(\lambda_{\text{UV}})} \rangle_{\mathbf{\Psi}} + \mathcal{O}(\epsilon^2)$, hence $m_{\pi_b}^2 \propto \epsilon$ within plateau tolerance (Theorem 7.630 and Corollary 7.631). The *assignment and covariance* of spurions, and the extraction of near-degenerate multiplets via Riesz projectors with Kato parallel transport, are spelled out in Definitions 7.559 and 7.618, Propositions 7.568 and 7.620, and Lemma 7.619. In Chapter 8 this constellation is the signature of *QIV* (persistent law deficit with a coherent state), not QIII.

By contrast, Quadrant III is the *emergent-symmetry* case: the explicit breaking is RG-irrelevant and the *law channel heals in the IR* while the state remains symmetric. Band diagnostics formalise this healing via IR-window invariance and stability of curvature-based tests (Proposition 7.397), consistent with the band Callan–Symanzik control (eq. (7.213)) and with gauge-parameter–independent visibilities (Nielsen safety on the band; eq. (7.209)). For interpreting flows, the chapter also uses the *Nielsen projection* and an eigen-basis that Jordanises the projected stability matrix, improving channelwise readings in the law sector (Definitions 7.454 and 7.457).

QIV (law asymmetric, state coherent). Goal (explicit breaking that persists at all scales). Work on a scale–tagged band that lies on a certified plateau $\mathfrak{P}$ (Definitions 7.31 and 7.96 and eq. (7.28)) produced by the band functor with UV/IR taggers (Definitions 7.26, 7.27 and 7.29). Introduce explicit symmetry–breaking spurions in the Lagrangian,

$$\mathcal{L} = \mathcal{L}_{\text{sym}} + \sum_k \varepsilon_k \mathcal{O}_k, \qquad \Delta_k \le d + z, \quad \varepsilon_k \neq 0,$$

so the breaking is relevant/marginal and does *not* wash out in the IR—the defining setting of the QIV sector (Section 7.9 and Theorem 7.557).

Checks (operational).

1. **Ward channel: non-vanishing IR violation.** The scale-tagged *physical* Ward identity on $\mathcal{H}_{\text{phys}}$ carries a persistent adjoint–covariant source proportional to the spurions ($\{\varepsilon_k\}$) and does *not* heal in the IR (Definition 7.567, eq. (7.147), and Theorem 7.570). In visibility language, the *law* channel is uniformly below unity on $\mathfrak{P}$:

$$\left|\Lambda_{\text{law}}(\ell)\right| \le 1 - \delta_\star.$$

   Here $\delta_\star \in (0,1)$ is uniform on the certified plateau $\mathfrak{P}$ (see Definition 7.553, Theorem 7.580, and Corollary 7.677).

2. **Spectrum and pseudo–Nambu–Goldstones (pNG).** Diagonalising the *physical*, band-prepared Hessian $\mathbb{H}^{(\lambda_{\text{UV}},\ell)}$ yields *pNG* modes when a symmetry is explicitly (but weakly) broken (eq. (7.10) and Proposition 7.629). Their masses obey GMOR-type relations controlled by spurion insertions and symmetry-breaking densities on $\mathcal{H}_{\text{phys}}$ (Section 7.9.11, Theorem 7.630, Corollary 7.631, and Proposition 7.627).

3. **State channel: coherence, not a vacuum cat.** Explicit breaking selects a *unique* non-symmetric vacuum, so the state-sector visibility remains near unity on large volumes,

$$\left|\Lambda_{\text{state}}(V)\right| \approx 1 \quad (V \to \infty),$$

   i.e. no macroscopic decoherence over a vacuum manifold (Definition 7.103 and Section 7.9.6). Contrast this with QII, where $\Lambda_{\text{state}}(V) \to 0$ as $V \to \infty$ due to SSB over a vacuum manifold (Sections 7.7 and 7.7.4).

4. **BRST/Nielsen safety and plateau robustness.** BRST-closed UV/IR taggers and commuting channel projectors ensure descent to $\mathcal{H}_{\text{phys}}$ and *gauge-parameter independence* of readouts (Nielsen safety on the band) (Definition 7.671, Propositions 7.130 and 7.236, and eq. (7.209)). Acceptance on $\mathfrak{P}$ is gated by the chapter's plateau criteria—small slope/curvature/oscillation in the log-chart $(u, v) = (\ln \lambda_{\text{UV}}, \ln \ell)$ (Definitions 7.47 and 7.96 and eq. (7.28)), and discrete IR-window controls make the tests numerically robust (Proposition 7.397 and Corollary 7.417).

Net QIV signature. In Quadrant IV the breaking terms are *explicit* and *relevant/marginal* ($\Delta_k \leq d + z$), so they do not wash out under the UV/IR flow on the band [32, 112]. Operationally, the adjoint–covariant, scale–tagged *physical* Ward identity carries a persistent IR source (no healing), cf. eq. (7.665); equivalently, the *law visibility* defined on $\mathcal{H}_{\text{phys}}$ satisfies $|\Lambda_{\text{law}}| < 1$ uniformly across the certified plateau $\mathfrak{P}$ [114, 38] (Definition 6.275, Section 6.12.4, and eq. (7.28)). By contrast, the *state visibility* remains near unity because explicit breaking selects a unique (non–symmetric) vacuum, so $|\Lambda_{\text{state}}| \approx 1$ on large volumes (no decoherence across a vacuum manifold) [112, 32]. All statements are made on the BRST–physical sector with band–limited operators—UV projector $P_x(\lambda_{\text{UV}})$ and IR window $K_{\text{IR}}^{(\ell)}$ that commute with channel projectors—and are *Nielsen–safe* by the band Nielsen identities [52, 114, 38, 112] (Definitions 6.272, 6.278, 6.280, 6.282 and 6.289 and eqs. (7.201) and (7.209)). Spectrally, would–be NG modes acquire small gaps governed by *GMOR–type* relations [112, 32] (Theorem 7.630 and Corollary 7.631).

One–line contrast with QIII (for orientation). Quadrant III realises *emergent symmetry*: explicit breakers are RG–*irrelevant* ($\Delta_k > d+z$), the Ward deficit and the law–visibility deficit *decay* in the IR with the predicted power law, and the state remains symmetric [32, 112] (cf. Proposition 7.397 and eqs. (7.209) and (7.213)). Quadrant IV has *relevant/marginal* breakers; the Ward deficit *persists* (no decay) while the state remains *coherent* due to a unique vacuum [114, 38] (eq. (7.665) and Theorem 7.630).

Classification on certified plateaus. On a certified plateau $\mathfrak{P}$ in the log–chart $(u, v) = (\ln \lambda_{\text{UV}}, \ln \ell)$ (Definition 7.47 and Section 6.12.4), Chapter 7 uses a canonical decision rule based on the *witness pair* $(\Lambda_{\text{law}}, \Lambda_{\text{state}})$ to assign quadrants [114, 52, 112] (Definitions 6.275 and 7.685). The law channel tests whether Ward/Slavnov–Taylor constraints are effectively exact on $\mathfrak{P}$ [38]; the state channel tests whether macroscopic preparation decoheres across a vacuum manifold [112]. Because the UV/IR taggers are BRST–closed and the equations are projected to $\mathcal{H}_{\text{phys}}$, all readouts are *Nielsen–safe* (gauge–parameter independent) [114, 52, 38, 112], and the plateau criteria—small gradient/curvature/oscillation—ensure stability of the verdicts (Definitions 6.272, 6.278, 6.280 and 6.289 and eqs. (7.28) and (7.201)).

QI (law symmetric, state symmetric). This is the calibration baseline: $|\Lambda_{\text{law}}| \approx 1$ and $|\Lambda_{\text{state}}| \approx 1$ on $\mathfrak{P}$. Scale–tagged equations reduce to clean, gauge–independent forms on $\mathcal{H}_{\text{phys}}$ [114, 38, 112] (Definition 6.272 and eqs. (6.333) and (6.334)); Ward identities are exactly conserved; and the physical (band–prepared) Hessian yields a positive, ghost–free, non–tachy-

onic spectrum with gauge–independent pole data [52, 114, 38, 30] (see $\mathcal{K}_{\rm phys}$–inversion and band Nielsen control in Proposition 7.16 and eq. (7.209); cf. Section 7.2.6).

QII (law symmetric, state asymmetric). The law channel remains at unity, but the state decoheres as volume grows because a vacuum manifold is present: $\Lambda_{\rm state}(V)\to 0$ as $V\to\infty$ while $|\Lambda_{\rm law}|\approx 1$ [32, 112] (Definition 6.275). Around a chosen vacuum, NG modes appear for broken *global* generators; for broken *gauged* generators the would–be NG modes are BRST–exact (eaten) and vectors become massive (BEH), all read on $\mathcal{H}_{\rm phys}$ with quartet decoupling ensuring that only transverse/matter poles are physical [38, 52, 109] (Section 7.2.6 and eq. (7.209)).

QIII vs. QIV (law asymmetric, state symmetric/coherent). In **QIII** the microscopic breaking is RG–irrelevant and the law channel *heals* in the IR with the predicted power law $1-|\Lambda_{\rm law}(\ell)| = \mathcal{O}\Big(\ell^{-(\Delta-(d+z))}\Big)$ (band CS control) [32, 112] (eq. (7.213) and Proposition 7.397), while the state remains symmetric $|\Lambda_{\rm state}|\approx 1$. In **QIV**, relevant/marginal explicit breakers keep a *persistent* law deficit: $|\Lambda_{\rm law}(\ell)|\le 1-\delta_\star$ on $\mathfrak{P}$, yet the state is coherent $|\Lambda_{\rm state}|\approx 1$ because a unique non–symmetric vacuum is selected; pseudo–NG masses track the breaking scales (GMOR–type relations) [112, 32] (Theorem 7.630 and Corollary 7.631). Across all quadrants, plateau certification and the discrete IR–window controls secure numerical robustness, and BRST/Nielsen safety guarantees that quadrant assignment is invariant under admissible gauge choices and band preparation [114, 38, 112] (Section 6.12.4, eqs. (7.28) and (7.209), and Proposition 7.397).

Intuition. "The law heals in the IR" signals that any apparent violation of dynamical constraints—those encoded by Ward/Slavnov–Taylor relations and their Nielsen-controlled refinements—diminishes under coarse–graining. At sufficiently long wavelengths, the effective description recovers the symmetry relations to the accuracy set by the infrared scale, so the constraints become effectively exact again [114, 38, 112].

Scaling statement. Let the law visibility be $\Lambda_{\rm law}(\ell)$ and define the law deficit $\delta_{\rm law}(\ell) := 1-|\Lambda_{\rm law}(\ell)|$. In Quadrant III, where the explicit breaking operators are RG–irrelevant ($\Delta>(d+z)$), the deficit obeys the band Callan–Symanzik control

$$\delta_{\rm law}(\ell) \;=\; \mathcal{O}\Big(\ell^{-(\Delta-(d+z))}\Big), \qquad \Delta>(d+z),$$

so the deficit decays as the IR window ℓ grows [32, 112] (eq. (7.213)).

Band preparation on $\mathcal{H}_{\rm phys}$. After UV/IR tagging and projection to the physical Hilbert space, the band–prepared Ward identities exhibit a vanishing IR source, and the clean forms of the scale–tagged equations hold to increasing precision [114, 38] (Definition 6.272 and eqs. (6.333) and (6.334)). This is the operational content of "healing": the cohomological projection removes unphysical components so that only the symmetry–respecting content survives on $\mathcal{H}_{\rm phys}$ [52].

Nielsen safety. Because all statements are formulated on $\mathcal{H}_{\text{phys}}$ with band–limited operators, the band Nielsen identities enforce gauge–parameter independence of the recovered relations; the healing is therefore a property of the physics, not of the gauge choice [114, 38, 112] (eqs. (7.201) and (7.209)).

Plateau robustness. On a certified plateau $\mathfrak{P}$ in the log–chart, the diagnostics (SLOPE, CURVATURE, OSCILLATION) sit below the prescribed tolerances across a two–dimensional region, so the limit $|\Lambda_{\text{law}}| \to 1$ is numerically stable and operationally meaningful for mode identification and comparisons [112] (Sections 6.12.4 and 7.3 and eq. (7.28)).

Chapter 8: Parasymmetric Witnesses and Diagnostics

What Chapter 8 does (methods, band, and witnesses). Chapter 8 is a *methods* chapter: it locks all calculations to the *band* (fixed UV/IR tags) and to the *physical BRST cohomology*, then builds *witnesses*—numerical visibilities in $[0, 1]$—that answer, channel by channel, "are the *laws* effectively symmetric here?" and "is the *state* coherently symmetric here?" [114, 52, 112] (cf. Definitions 6.272, 6.275, 6.278, 6.280 and 6.282). The machinery that makes those answers trustworthy is: (i) *Nielsen safety* (gauge–parameter independence on shell) [114, 38, 112] (eqs. (7.201) and (7.209) and Proposition 7.208), (ii) *cohomological inversion* (ghost/longitudinal pieces decouple before propagator inversion) [117, 52] (Proposition 7.16 and Definition 6.274), (iii) *Riesz projectors* to extract pole locations (gauge–independent) [115, 30] (eq. (7.438) and Definition 7.286), and (iv) *plateau certification* in a log–scale geometry (small slope/curvature/oscillation), with integrated–readout and discrete–IR controls [32, 112] (Section 6.12.4, eq. (7.28), and Propositions 7.146 and 7.397). With those in hand, a *canonical decision rule* assigns each certified bandpoint to one of four quadrants (QI–QIV), cleanly separating unbroken, SSB, emergent, and explicit–breaking regimes (Definition 7.685).

Synthesis, sectors, and extensions. The second big idea is *synthesis and extensibility*. The same witness logic is made *sector–conditioned* (TT vs. longitudinal, spin/charge, boundary vs. bulk; Section 8.1 and Definition 8.9) and a *parasymmetric overlay* turns law/state readouts into calibrated "agreement"/"synthesis" meters that remain well–posed even when law and state disagree (Sections 8.2, 8.2.1 and 8.5.4). Boundary soft/edge modes receive a parallel dashboard (Section 8.1.7); anomaly, anyon/CS, fracton, hydrodynamic, SUSY, higher–form and non–invertible cases are all read through sector–specific witnesses but judged by the same band/BRST/plateau standards [114, 52, 112, 118] (Sections 6.12.4, 8.1.2, 8.1.5 and 8.1.8 to 8.1.11 and eq. (7.209)). Net effect: a robust, gauge–clean, scale–aware *operational pipeline* that turns correlators and kernels into falsifiable symmetry diagnoses—uniformly across sectors and boundaries (cf. Definitions 6.274 and 7.15).

What Chapter 8 builds on and where it lives. Everything in Chapter 8 is done *on the band*—i.e., at fixed UV/IR scale tags $(\lambda_{\text{UV}}, \ell)$—and strictly inside the *physical sector* $\mathcal{H}_{\text{phys}}$ (the BRST cohomology). Algebraically, one starts from the scale–tagged, gauge–fixed quadratic

expansion about a symmetric reference vacuum and packages the fluctuations as

$$\delta\hat{\mathbf{U}}_{\lambda_{\mathrm{UV}},\ell} := \big(\delta\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell},\, \delta\hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}},\ell}\big)^{\mathsf{T}}.$$

This defines a *gauge–fixed quadratic kernel* $\mathcal{K}_{\Psi}^{(\lambda_{\mathrm{UV}},\ell)}$. Projecting to $\mathcal{H}_{\mathrm{phys}}$ gives the *physical kernel*

$$\mathcal{K}_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)} := \mathsf{P}_{\mathrm{phys}}\, \mathcal{K}_{\Psi}^{(\lambda_{\mathrm{UV}},\ell)}\, \mathsf{P}_{\mathrm{phys}},$$

whose inverses (retarded/advanced/Feynman) are the *physical propagators* $\mathcal{D}_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}$. On $\mathcal{H}_{\mathrm{phys}}$ one has the cohomological inversion $\mathcal{K}_{\mathrm{phys}}\mathcal{D}_{\mathrm{phys}} = \mathbf{1}$, and—crucially—*the pole positions are gauge–parameter independent*. Intuitively, longitudinal/ghost quartets decouple before inversion, and Slavnov–Taylor/Nielsen identities render gauge–parameter shifts BRST–exact, hence invisible in cohomology (cf. Section 7.2.2, Definition 7.15, Proposition 7.16, and eq. (7.209)).

To keep gauge control crisp, the chapter adopts a *Background–Field Method* (BFM) presentation, imposing background Ward identities together with a *band–limited* Nielsen identity for the 1PI functional Γ. Two robust conclusions then follow (on the band): (i) *pole masses are ξ–independent*; (ii) *on–shell visibilities are protected*. Both statements remain stable under admissible regulator moves inside a specified tolerance (cf. Section 7.2.9, Definition 7.59, eqs. (7.19) and (7.27), and Proposition 7.208).

Plateau geometry and certification (scale control). Scale space is equipped with the *log–chart*

$$(u, v) = (\ln \lambda_{\mathrm{UV}}, \ln \ell), \qquad g_{\mathrm{plat}} := du^2 + dv^2, \qquad D_\lambda = \partial_u, \quad D_\ell = \partial_v,$$

and three quantitative *plateau metrics* for any band–readout $O(u, v)$: a *Slope* bound $|\nabla O|$, a *Curvature* bound $\|\nabla^2 O\|$, and an *Oscillation* bound $\operatorname{osc} O$. A band point $(\lambda_{\mathrm{UV}}, \ell)$ is *plateau–stable* if, throughout a small geodesic ball in (u, v), one has

$$|\partial_u O|,\, |\partial_v O| \le \tau_O^{(1)}, \qquad \|\nabla^2 O\| \le \tau_O^{(2)}, \qquad \operatorname{osc} O \le \tau_O^{(\mathrm{osc})}.$$

The chapter proves the *equivalence* (with matched tolerances) between a channelwise/directional certification and this geometric one (cf. Sections 7.3.2 and 7.3.5, Definition 7.31, eqs. (7.28) and (7.46), and Proposition 7.38).

Two stability features matter in practice.

- **Integrated readouts.** If you average a channel readout R_α over a weight w supported inside a certified plateau $\mathfrak{P}$, $\overline{R}_\alpha := \int_{\mathfrak{P}} w\, R_\alpha$, the same slope/curvature/oscillation bounds carry to $\overline{R}_\alpha$ with equal or smaller constants; discrete (grid/box) versions inherit the same control (cf. Proposition 7.146 and eqs. (7.390) and (7.410) to (7.414)).

- **Discrete IR windows.** A curvature–based proxy for the IR window on a grid achieves uniform second–order consistency: the discrete estimator ϕ_h differs from the continuous ϕ by $\mathcal{O}(h^2)$ (plus $\mathcal{O}(\tau^2)$ for the low–pass), and the symmetric–difference in accepted sets has controllably small band–measure (cf. Proposition 7.397).

Witnesses and the quadrant decision rule. The chapter's central deliverable is an *operational decision rule*—applied on a certified plateau—classifying the law/state symmetry situation into the four quadrants (QI–QIV) using *visibilities* (witnesses) computed on $\mathcal{H}_{\text{phys}}$ and on the band:

- a *law–channel* visibility $\Lambda_{\text{law}}(\lambda_{\text{UV}}, \ell)$ that asks whether Ward/Slavnov–Taylor constraints are effectively exact on $\mathfrak{P}$;
- a *state–channel* visibility $\Lambda_{\text{state}}(V; \lambda_{\text{UV}}, \ell)$ that asks whether macroscopic preparation decoheres over a vacuum manifold (large–V limit) or remains coherent (unique vacuum);
- optional *mix–channel* or sector–conditioned variants (e.g., TT vs. longitudinal blocks; spin or charge–resolved) to refine diagnosis while respecting BRST and band operations.

The *canonical quadrant rule* then reads the pair $\big(\Lambda_{\text{law}}, \Lambda_{\text{state}}\big)$ on $\mathfrak{P}$:

- **QI (symmetric law, symmetric state):** $|\Lambda_{\text{law}}| \approx 1$ and $|\Lambda_{\text{state}}| \approx 1$. This is the calibration baseline; the band–prepared physical Hessian is positive with only transverse, positive–norm modes and gauge–independent pole data (no ghosts/tachyons) (cf. Definition 7.685 and Theorem 7.19).
- **QII (symmetric law, asymmetric state):** $|\Lambda_{\text{law}}| \approx 1$ but $\Lambda_{\text{state}}(V) \to 0$ as $V \to \infty$ (vacuum manifold present). Around a chosen vacuum one sees NG modes for broken *global* symmetries; for *gauged* cases would–be NG modes are BRST–paired and vectors become massive (BEH), with quartet decoupling ensuring only physical poles survive (cf. Section 7.7 and Proposition 7.22).
- **QIII (emergent law, symmetric state):** explicit breakers are RG–irrelevant; the Ward deficit and $1 - |\Lambda_{\text{law}}|$ *heal* in the IR with the predicted power law $\sim \ell^{-(\Delta-(d+z))}$; the state stays symmetric, $|\Lambda_{\text{state}}| \approx 1$ (cf. eq. (7.213) and Proposition 7.83).
- **QIV (explicit–breaking law, coherent state):** relevant/marginal breakers keep a *persistent* law deficit on $\mathfrak{P}$, $|\Lambda_{\text{law}}(\ell)| \le 1 - \delta_\star$ (no IR healing), while the state remains coherent $|\Lambda_{\text{state}}| \approx 1$ because a unique non–symmetric vacuum is selected; pseudo–NG masses satisfy GMOR–type relations $m^2_{\text{pNG}} \propto \varepsilon_{\text{eff}}$ (cf. Theorem 7.630, Corollary 7.631, and Section 7.9.11).

All witnesses are *Nielsen–safe* (gauge–parameter independent) because: (i) UV/IR taggers carry ghost number 0 and commute with BRST, (ii) projection to $\mathcal{H}_{\text{phys}}$ removes gauge–exact parts before inversion, and (iii) the BFM/Nielsen algebra keeps band on–shell data invariant (cf. Propositions 7.20 and 7.208 and eq. (7.209)).

How poles are isolated (and kept gauge–safe). Pole extraction is implemented via *Riesz projectors* acting on the transverse–traceless two–point kernel $\Gamma^{(\text{TT})}(p^2)$: encircle a pole at $p^2 = -m_i^2$ with a small contour γ_i and set

$$\mathbb{P}_i \;=\; \frac{1}{2\pi i} \oint_{\gamma_i} \Big(\Gamma^{(\text{TT})}(z)\Big)^{-1} dz, \qquad \widehat{\Gamma}_i := \mathbb{P}_i \, \Gamma^{(\text{TT})} \, \mathbb{P}_i.$$

The eigenvalues of $\widehat{\Gamma}_i$ give m_i^2; by the band Nielsen identity these pole positions are α–independent. Residues may vary with gauge/parametrisation, but *pole locations do not* (cf. Definition 7.286 and eqs. (7.20), (7.21) and (7.438)).

Sector–conditioned witnesses (canonical framework). Chapter 8 systematizes symmetry vs. breaking diagnostics by introducing, for each certified sector $\mathscr{S}$ (bulk, boundary, anisotropic, …), a *witness map* evaluated *on the band* and in the BRST cohomology. A *witness* is a BRST–safe, band–limited functional

$$\mathbf{W}^{[\mathscr{S}]}:\ \mathsf{P}_{\text{phys}}\,\mathfrak{B}_{\lambda_{\text{UV}},\ell}\Big[\text{operators, currents, states on } \mathfrak{P}\Big]\,\mathsf{P}_{\text{phys}} \longrightarrow [0,1], \qquad \Lambda^{[\mathscr{S}]} := \mathbf{W}^{[\mathscr{S}]},$$

constructed so that it is *Nielsen–safe* on shell (gauge–parameter independent; cf. Definitions 6.272 and 6.282 and eqs. (7.201) and (7.209)). Law and state are read as a *witness pair* $\left(\Lambda^{[\mathscr{S}]}_{\text{law}}, \Lambda^{[\mathscr{S}]}_{\text{state}}\right)$ with certified tolerances $\varepsilon_{\text{law}}, \varepsilon_{\text{state}} \in (0,1)$ supplied by the plateau logic (Section 6.12.4, eq. (7.28), and Definition 6.275). Under the chapter's conventions—fixed symmetry G, fixed observation window/order of limits, and BRST–closed probes—the resulting verdict matches the formal quadrant classification on $\mathfrak{P}$ (QI–QIV; Definition 7.685).

Sector–conditioned and boundary variants. Beyond the baseline law/state pair, one may introduce a *mix–channel* witness $\Lambda^{[\mathscr{S}]}_{\text{mix}}$ to monitor cross–couplings (e.g., TT/longitudinal mixing, law–state cross–responses) in a given certified sector $\mathscr{S}$; it is built from the same band–prepared kernels and inherits BRST/Nielsen safety (cf. Section 8.1, Definitions 6.272 and 6.275, and eq. (7.209)). Witnesses may also be *sector–conditioned* (TT vs. longitudinal, spin– or charge–resolved) to disentangle kinematics from conservation–law constraints (Section 8.1 and Definition 8.9). For *boundary* arenas, very–low–frequency transport ("in–/out/through" drivers) can *confound* law–vs–state assessments unless the time–scale hierarchy is respected; some boundary visibilities are excellent for topology/conductance but not faithful for diagnosing law–channel healing or persistent deficits without mode/time–scale filtering (Section 8.1.7 and Lemmas 7.584 and 7.585).

Witnesses and the quadrant decision rule. On a certified plateau, the *operational decision rule* classifies the law/state symmetry situation using visibilities computed on $\mathcal{H}_{\text{phys}}$ and on the band:

- **Law–channel visibility** $\Lambda_{\text{law}}(\lambda_{\text{UV}}, \ell)$: are Ward/Slavnov–Taylor constraints effectively exact on $\mathfrak{P}$?
- **State–channel visibility** $\Lambda_{\text{state}}(V; \lambda_{\text{UV}}, \ell)$: does macroscopic preparation decohere across a vacuum manifold (large–V limit) or remain coherent (unique vacuum)?
- **Optional mix/sector variants** (e.g., TT vs. longitudinal; spin/charge–resolved) to refine diagnosis while preserving BRST and band consistency.

These inputs feed the canonical quadrant rule (Definition 7.685) to return a QI–QIV assignment with the plateau tolerances explicitly tracked.

Boundary/edge witnesses (bulk–boundary parallel). On a certified *boundary* plateau $\mathfrak{P}_\partial$ with a tangential IR window $K^{(L)}_{\mathrm{IR},\partial}$ (band preparation and BRST–closed tagging as in Sections 6.12.4 and 8.1.7 and eq. (7.28)), the tagged boundary Ward identity reads

$$\nabla_\parallel \cdot J^{(\partial)}_\parallel \;=\; S_\partial \quad \text{up to BRST–exact improvements,}$$

and the *law–channel* visibility is defined by

$$\Lambda^{[\partial]}_{\mathrm{law}}(L) \;:=\; 1 - \frac{\big\|S_\partial\big\|_{\mathfrak{P}_\partial}}{\big\|\nabla_\parallel \cdot J^{(\partial)}_\parallel\big\|_{\mathrm{ref},\mathfrak{P}_\partial}} \;\in [0,1],$$

where the boundary norms and the reference normalisation are those fixed in the boundary witness setup (Definition 8.31). The *state–channel* witness uses tolerance semantics on $\mathfrak{P}_\partial$:

$$\begin{aligned} &\text{boundary state symmetric} &&\iff \big|\Lambda^{[\partial]}_{\mathrm{state}}\big| \geq 1-\varepsilon_{\mathrm{state}},\\ &\text{boundary SSB} &&\iff \big|\Lambda^{[\partial]}_{\mathrm{state}}\big| \leq \varepsilon_{\mathrm{state}}. \end{aligned}$$

with $\varepsilon_{\mathrm{law}}, \varepsilon_{\mathrm{state}} \in (0,1)$ taken from the plateau certification (Section 6.12.4 and eq. (7.28)). Quadrant assignments (QI–QIV) mirror the bulk decision rule using the boundary pair $\big(\Lambda^{[\partial]}_{\mathrm{law}}, \Lambda^{[\partial]}_{\mathrm{state}}\big)$, as formalised in Proposition 8.32 and Definition 7.685.

Fermionic witnesses. In the fermionic sector the band–prepared, gauge–fixed two–point kernel $\Gamma_\psi(p)$ is projected to $\mathcal{H}_{\mathrm{phys}}$ and its poles are isolated by Riesz projectors [115, 30],

$$\mathbb{P}_i \;=\; \frac{1}{2\pi i}\oint_{\gamma_i}\big(\Gamma_\psi(z)\big)^{-1}\,dz, \qquad \widehat{\Gamma}_i \;=\; \mathbb{P}_i\,\Gamma_\psi\,\mathbb{P}_i,$$

so that $\mathrm{spec}(\widehat{\Gamma}_i) = \{m_i^2\}$ gives the (gauge–independent) pole masses, while residues encode chiral/parity structure (Nielsen–safe on the band) [114, 38, 112]. The *law* visibility checks the fermionic Ward identities (vector/axial) on the certified plateau [38, 110, 111]; the *state* visibility tests coherence vs. decoherence in macroscopic preparation [32, 112]. In QII (global SSB) NG partners appear in the spectral density [32]; in QIV (explicit breaking) pseudo–NG gaps scale with spurions (GMOR–type) $m^2_{\mathrm{pNG}} \propto \varepsilon_{\mathrm{eff}}$ [112, 32]. See Section 8.1.1, eqs. (7.209) and (7.438), Theorem 7.630, and Corollary 7.631.

Supersymmetry (SUSY) witnesses. Here one compares bosonic and fermionic TT blocks on the band, testing SUSY Ward identities: exact SUSY gives spectral pairing $m_b = m_f$ and matched on–shell visibilities; soft breaking tilts residues but leaves nearby poles; hard breaking splits pole locations [52]. Define a pairing defect

$$\Delta_{\mathrm{SUSY}} \;:=\; \big|m_b - m_f\big|,$$

and read it jointly with the SUSY law visibility (Nielsen–safe) [114, 38, 112]. Healing of $\Delta_{\mathrm{SUSY}} \to 0$ in the IR indicates QIII; a persistent nonzero plateau indicates QIV [32, 112]. See Section 8.1.2 and eqs. (7.19) and (7.209).

p–forms witnesses. For higher–form symmetries the law channel enforces generalized Gauss constraints (higher–form Ward identities),

$$d * F^{(p+1)} \;=\; J^{(p)}_{\mathrm{cons}},$$

while the state channel probes long–range order of brane/line operators. The witness separates Maxwell–like, topological (BF/CS–type), and confining responses by their pole/branch–cut structure and finite–size scalings on the band [110, 111, 32, 112]. See Sections 8.1.3 and 8.1.8.

Higher–spins witnesses. Decompose the spin–s kernel into transverse–traceless and longitudinal parts, $\Gamma^{(s)} \to \big(\Gamma^{(s,\mathrm{TT})}, \Gamma^{(s,\mathrm{L})}\big)$, and locate TT poles with Riesz projectors [115, 30]; healthy sectors exhibit positive–norm TT poles with no ghost contamination [52, 38]. Longitudinal law–deficits shrinking across the plateau indicate QIII (emergent symmetry) [32, 112], whereas regulated mass terms in TT reflect explicit breaking (QIV) [109, 113]. See Section 8.1.4, Definition 7.286, and eq. (7.438).

Hydrodynamics witnesses. In the small–$|p|$ longitudinal block one reads diffusive and sound poles,

$$\omega_{\mathrm{diff}}(p) \approx -\,iD\,p^2, \qquad \omega^{\pm}_{\mathrm{sound}}(p) \approx \pm c_s\,p - i\,\Gamma\,p^2,$$

and extracts Kubo coefficients from band–limited correlators [119, 118]. The law visibility asks whether conserved–current Ward identities hold on $\mathfrak{P}$ [38]; the state visibility tests whether the hydrodynamic modes are coherently supported without boundary artefacts [32, 112]. See Section 8.1.5.

Second sound (matrix formulation). Coupled energy/charge (or entropy) responses define a small–$|p|$ matrix kernel $\mathcal{M}(p)$; a genuine second–sound branch appears as a distinct propagating pole after diagonalization,

$$\det \mathcal{M}(\omega, p) \;=\; 0 \quad \Rightarrow \quad \omega_{\mathrm{2nd}}(p) = \pm c_2\,p + \cdots .$$

The witness includes falsification tests (decoupling conditions) to avoid mis–tagging slow diffusion as a wave [119, 118]. See Section 8.1.6.

Bound soft/edge modes witnesses. With physical boundary conditions imposed, the *edge kernel* $\Gamma^{(\partial)}(k_\parallel, \omega)$ is prepared on the band and projected to $\mathcal{H}_{\mathrm{phys}}$ [114, 52]. Edge poles/zero modes are isolated by Riesz projectors [115, 30],

$$\mathbb{P}^{(\partial)}_i \;=\; \frac{1}{2\pi i}\oint_{\gamma_i}\Big(\Gamma^{(\partial)}(z)\Big)^{-1}\,dz, \qquad \widehat{\Gamma}^{(\partial)}_i := \mathbb{P}^{(\partial)}_i\,\Gamma^{(\partial)}\,\mathbb{P}^{(\partial)}_i,$$

so that $\det \widehat{\Gamma}^{(\partial)}_i(\omega_i, k_\parallel) = 0$ yields chiral branches $\omega_i(k_\parallel) = v_i k_\parallel + \cdots$ when present (Nielsen–safe on the band) [114, 38]. The *law* witness enforces the tagged edge Ward identity (anomaly–inflow form) and its BRST–closed improvements [110, 111, 52],

$$\partial_t \rho_\partial + \nabla_\parallel \cdot j^{(\partial)}_\parallel \;=\; S_\partial, \qquad \Lambda^{[\partial]}_{\mathrm{law}}(L) = 1 - \frac{\|S_\partial\|_{\mathfrak{P}_\partial}}{\|\nabla_\parallel \cdot j^{(\partial)}_\parallel\|_{\mathrm{ref},\mathfrak{P}_\partial}},$$

while the *state* witness diagnoses coherent vs. decohering edges from the sharpness of the edge poles under size/disorder scans on the certified boundary plateau $\mathfrak{P}_\partial$ [32, 112] (cf. Section 8.1.7, Definition 8.31, and Proposition 8.32).

Anyons (Chern–Simons) witnesses. In a band–prepared Chern–Simons (CS) sector, the Hall response σ_H and braiding phases θ provide quantized law–channel diagnostics that are stable on a certified plateau [32, 112, 24, 25, 23]:

$$\left|\sigma_H - \frac{k}{2\pi}\right| \le \varepsilon_{\text{law}}, \qquad \theta_{ab} = 2\pi\,(K^{-1})_{ab} \pmod{2\pi},$$

with deviations indicating explicit breaking or non–topological mixing [112]. The *state* witness separates bulk topological mass from boundary chiral modes by comparing bulk TT poles with edge dispersions in the same band setting; both witnesses inherit gauge–parameter independence via the band Nielsen control [114, 38, 112] and capture anomaly–inflow–type edge constraints [110, 111] (cf. Section 8.1.8 and eq. (7.19)).

Fractons witnesses. Restricted mobility is certified by *higher–moment Ward constraints* for charge and dipole densities on the band [112, 32, 38]:

$$\partial_t \rho + \nabla\cdot J = 0, \qquad \partial_t P_i + \partial_j J^{(2)}_{ij} = 0, \quad P_i := \int x_i\,\rho,$$

together with the *absence* (or parametrically strong suppression) of poles in the driven dipole/quadrupole channels after projection to $\mathcal{H}_{\text{phys}}$ [32, 112]. The *law* witness checks the moment–conservation identities on $\mathfrak{P}$ [38]; the *state* witness verifies that preparation respects subdimensional kinematics (no spurious mobile carriers) (cf. Section 8.1.9).

Higher–forms and non–invertible symmetries. For a p–form symmetry with conserved $(p{+}1)$–form current $J^{(p+1)}$, the law–channel witness enforces the higher–form Ward identity on the band [38, 112, 32],

$$d * J^{(p+1)} = 0,$$

while the state–channel witness uses defect/'t Hooft operators $\mathcal{W}(\Sigma^p)$ (surfaces/lines) to read long–range order and fusion/selection rules [32, 24, 25, 23]. Non–invertible structure is probed by defect composition [112, 32],

$$\mathcal{D}_a \circ \mathcal{D}_b \;=\; \sum_c N_{ab}^{\;c}\,\mathcal{D}_c,$$

and its band–stable plateaus: IR healing of these constraints ($\Rightarrow$ QIII) vs. persistent violations ($\Rightarrow$ QIV) is read from defect–conditioned kernels [112, 32] (cf. Section 8.1.10).

Anomaly witnesses. Background–field variations convert 't Hooft/Wess–Zumino consistency into operational law–channel checks [114, 112]:

$$\delta_\alpha \Gamma_{\text{band}}[A] \;=\; \int \alpha\,\mathcal{A}[A],$$

with $\mathcal{A}[A]$ the anomaly functional extracted on the band [110, 111]. Plateau robustness (small slope/curvature/oscillation) renders the inferred coefficients scheme–stable; UV/IR matching of $\mathcal{A}$ decides if the symmetry is realized, broken, or requires topological sectors [32, 112]. Nielsen identities guarantee gauge–parameter independence of the on–shell readouts [114, 38, 112] (cf. Section 8.1.11 and eq. (7.19)).

Non–equilibrium (Floquet/Lindblad) witnesses. For periodic drive of period T, the law–channel check uses the Floquet operator [22]

$$U_F(T) \;=\; \mathcal{T}\exp\!\Big(-i\int_0^T H(t)\,dt\Big), \qquad U_F\,|\phi_n\rangle = e^{-i\varepsilon_n T}\,|\phi_n\rangle,$$

and stroboscopic Ward constraints on $\Gamma_{\text{band}}\big|_{\text{Floquet}}$ [114, 38]. In open systems the Liouvillian $\mathcal{L}$ governs dynamics, $\dot\rho = \mathcal{L}\rho$, with spectrum $\{\lambda_a\}$ read on the band; the state witness compares coherence vs. heating/decoherence by tracking spectral gaps and steady–state projectors [118, 115, 30]. Integrated–readout versions keep plateau certification despite time dependence [112, 32] (cf. Section 8.1.12 and Proposition 7.146).

The Synthesis of Laws and States (Section 8.2). This section provides a single *operational* stage on which the *law*– and *state*–channel diagnostics can be read *together* without logical dead–ends when they differ. Concretely, the quadrant dashboard is evaluated only on a *certified arena* (shared plateau, BRST–closed tagging, sector witnesses available; cf. Section 6.12.4, eq. (7.28), and Definition 6.275) and packaged in a paraconsistent meta–layer that records co–tenable modalities. On the band and inside the physical sector,

$$(\Lambda_{\text{law}}, \Lambda_{\text{state}}) \;\in\; [0,1]^2, \qquad \mathbb{D} := \{x \in [0,1] \mid x \geq 1-\varepsilon_\bullet\}$$

is the set of designated values (channel–wise tolerances from the plateau), and a non–explosion diagnostic

$$\Pi_{pc}(\Lambda_{\text{law}}, \Lambda_{\text{state}}) \;:=\; \mathbf{1}_{\mathbb{D}}(\Lambda_{\text{law}})\,\mathbf{1}_{[0,1]\setminus\mathbb{D}}(\Lambda_{\text{state}}) \;+\; \mathbf{1}_{\mathbb{D}}(\Lambda_{\text{state}})\,\mathbf{1}_{[0,1]\setminus\mathbb{D}}(\Lambda_{\text{law}})$$

flags controlled "both/and" situations without collapsing the readout; the canonical quadrant decision remains governed by the usual rule (QI–QIV) applied on a certified plateau (Definition 7.685). Interferometric overlays and duality lifts are required to be *dashboard–invariant*, in the sense that any admissible overlay $\mathcal{O}$ that preserves certification (band/BRST closure) satisfies

$$\mathcal{O}:\;\; (\Lambda_{\text{law}}, \Lambda_{\text{state}}) \mapsto (\Lambda'_{\text{law}}, \Lambda'_{\text{state}}) \quad \text{with} \quad \mathbf{1}_{\text{QI–QIV}}\Big(\Lambda_{\text{law}}, \Lambda_{\text{state}}\Big) = \mathbf{1}_{\text{QI–QIV}}\Big(\Lambda'_{\text{law}}, \Lambda'_{\text{state}}\Big),$$

so the quadrant assignment is stable under admissible overlays (cf. Section 8.5).

Sector–conditioned realisation and operational template (Section 8.1). Given a sector $\mathscr{S}$ with instrumentation $\mathcal{I}^{[\mathscr{S}]}$ and witness set $\mathbf{W}^{[\mathscr{S}]}$, the synthesis is instantiated as follows. *Step 0* (arena and certification): fix the arena $\mathcal{A} \in \{\mathfrak{P}, \mathfrak{P}_\partial\}$, the IR knob r, and certify

the plateau(s) using the band functor and BRST–closed taggers ($\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$; $P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}^{(\ell)}$; cf. Definitions 6.278, 6.280 and 6.282 and eq. (7.28)). Work on the log–chart

$$(u,v) = (\ln \lambda_{\mathrm{UV}}, \ln r), \qquad g_{\mathrm{plat}} = du^2 + dv^2.$$

Step 1 (witness pair): build the sector pair

$$\Big(\Lambda_{\mathrm{law}}^{[\mathscr{S}]}(u,v),\ \Lambda_{\mathrm{state}}^{[\mathscr{S}]}(u,v)\Big) \ := \ \Big(\mathbf{W}_{\mathrm{law}}^{[\mathscr{S}]}\big[\mathsf{P}_{\mathrm{phys}}\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\cdot]\mathsf{P}_{\mathrm{phys}}\big],\ \mathbf{W}_{\mathrm{state}}^{[\mathscr{S}]}\big[\mathsf{P}_{\mathrm{phys}}\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\cdot]\mathsf{P}_{\mathrm{phys}}\big]\Big),$$

which is Nielsen–safe on shell by construction ($\partial_\xi, \partial_{\alpha_{\mathrm{gf}}}$ kill physical readouts; eq. (7.209)). The canonical decision rule then applies on $\mathcal{A}$ with the certified tolerances (Definition 7.685 and Section 6.12.4).

Parasymmetric Synthesis of Laws and States (Section 8.2.1). Locally (at an *anchor*), each channel is resolved into a minimal two–mode basis $\{|\mathrm{Sym}\rangle, |\mathrm{Break}\rangle\}$. The parasymmetric overlay forms even/odd combinations that read joint coherence vs. law–state skew with *meter–level* operators:

$$P_{\mathrm{para}}^{(a)} = |+\rangle\langle+|, \quad P_{\mathrm{anti}}^{(a)} = |-\rangle\langle-|, \qquad |\pm\rangle := \tfrac{1}{\sqrt{2}}\Big(|\mathrm{Sym}\rangle \pm |\mathrm{Break}\rangle\Big), \quad a \in \{\mathrm{Law}, \mathrm{State}\}.$$

and, on the relational space $L \otimes S$, Bell–type projectors select "coherent agreement" (I/IV diagonal) vs. "coherent opposition" (II/III diagonal). The corresponding meters,

$$v_{\mathrm{Synthesis}}^{(a)} := \mathrm{Tr}\Big(P_{\mathrm{para}}^{(a)}\rho^{(a)}\Big), \qquad v_{\mathrm{Agreement}}^{(LS)} := \mathrm{Tr}\Big(P_{\mathrm{para}}^{(LS)}\rho^{(LS)}\Big),$$

are computed *on the band* and *in* $\mathcal{H}_{\mathrm{phys}}$, and inherit the plateau/Nielsen guarantees of their ingredients. These anchor–level overlays feed back into the dashboard, providing a concrete, certified mechanism for quadrant readouts that respects the chapter's operational constraints.

Putting them together: sector resolver and stability. The chapter defines a sector–wise resolver that maps the family of sector readouts to labels (quadrant + explanatory class) and proves *stability* under: (i) admissible gauge–fermion retunings (Nielsen–flatness), (ii) BRST–closed retunings of the UV/IR taggers (P_x, K_{IR}) that preserve certification, and (iii) certified coarsenings of the plateau domain. Symbolically,

$$\mathcal{R}^{[\mathscr{S}]} : \ \Big\{\Lambda_{\bullet}^{[\mathscr{S}]}(u,v)\Big\}_{(u,v)\in\mathcal{A}} \ \longmapsto \ (\mathrm{QI–QIV},\ \mathrm{class}) \quad \text{with} \quad \partial_\xi \mathcal{R}^{[\mathscr{S}]} = \partial_{\alpha_{\mathrm{gf}}} \mathcal{R}^{[\mathscr{S}]} = 0,$$

and with inclusion relations preserved under the geometric/log–chart plateau definitions (Sections 6.12.4 and 8.1 and eq. (7.28)).

Summary. Section 8.2 supplies the meta–logic and dashboard for joint law–state reading on certified arenas; Section 8.2.1 gives the *parasymmetric* anchor–level realisation; and the sector template $\big(\mathscr{S}, \mathcal{I}^{[\mathscr{S}]}, \mathbf{W}^{[\mathscr{S}]}\big)$ specifies how to implement both—witnesses, tags, and plateaus—in any sector (Section 8.1).

Parasymmetric Overlay of Boundary Soft/Edge Modes (worked example) (Section 8.5.4). Soft/edge modes often carry clearer symmetry fingerprints than the bulk. This overlay upgrades the law–vs–state dashboard to treat the *boundary* as a first–class certified arena, using the same band/plateau and BRST/Nielsen–safe logic as in the bulk (cf. Sections 6.12.4 and 8.1.7 and eqs. (7.28) and (7.209)).

Boundary calibration: witnesses → density operators. On a certified boundary plateau $\mathfrak{P}_\partial$ with tangential IR window $K^{(L)}_{\mathrm{IR},\partial}$ and spectral projector $P_x(\lambda_{\mathrm{UV}})$ (cf. Definitions 6.278 and 6.280), define the *boundary parasymmetric calibration* as a CPTP map

$$\kappa^{[\partial]}:\ \Big(\Lambda^{[\partial]}_{\mathrm{law}}(L),\,\Lambda^{[\partial]}_{\mathrm{state}}(L);\,\mathcal{W}^{[\partial]},\,\mathcal{Z}^{[\partial]}\Big)\longrightarrow\Big(\rho^{(L)[\partial]},\,\rho^{(S)[\partial]},\,\rho^{(LS)[\partial]}\Big),\qquad (\textit{Definition } 8.133),$$

evaluated *on the band* and in the physical sector $\mathsf{P}_{\mathrm{phys}}\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\cdot]\mathsf{P}_{\mathrm{phys}}$ (cf. Definitions 6.272 and 6.282). BRST closure yields *Nielsen flatness* (gauge–parameter independence on shell; eq. (7.209)).

Meters on the boundary: synthesis and agreement. Using parasymmetric projectors at each anchor and Bell–type projectors on the $L\otimes S$ space, define the four boundary meters (cf. Definition 8.134):

$$v^{(L)[\partial]}_{\mathrm{Synthesis}}=\mathrm{Tr}\Big(P^{(L)}_{\mathrm{para}}\rho^{(L)[\partial]}\Big),\quad v^{(S)[\partial]}_{\mathrm{Synthesis}}=\mathrm{Tr}\Big(P^{(S)}_{\mathrm{para}}\rho^{(S)[\partial]}\Big),$$
$$v^{(LS)[\partial]}_{\mathrm{Synthesis}}=\mathrm{Tr}\Big(P^{(LS)}_{\mathrm{para}}\rho^{(LS)[\partial]}\Big),\qquad v^{(LS)[\partial]}_{\mathrm{Agreement}}=\mathrm{Tr}\Big(P^{(LS)}_{\Phi,\mathrm{para}}\rho^{(LS)[\partial]}\Big).$$

These are normalised so 1 = coherent "both/and," 1/2 = *mixture baseline*, and 0 = coherent "anti–both." Intuitively, the *Synthesis* meters detect coherent opposition (II/III diagonal), whereas the *Agreement* meter detects coherent agreement (I/IV diagonal).

Cross–coherence wired to edge operators. A boundary cross–coherence lemma gives an explicit off–diagonal entry (cf. Lemma 8.135):

$$\rho^{(LS)[\partial]}_{12}\ \propto\ \frac{\big\langle S_\partial,\ \mathcal{O}_\partial\big\rangle_{\mathfrak{P}_\partial}}{\sqrt{\langle S_\partial,S_\partial\rangle_{\mathfrak{P}_\partial}\,\langle\mathcal{O}_\partial,\mathcal{O}_\partial\rangle_{\mathfrak{P}_\partial}}},$$

with S_∂ the boundary state readout and $\mathcal{O}_\partial$ a matched edge operator (edge current, soft charge density, memory/vertex operator). Consequently,

$$v^{(LS)[\partial]}_{\mathrm{Synthesis}}=\tfrac12+\mathrm{Re}\ \rho^{(LS)[\partial]}_{12},\qquad v^{(LS)[\partial]}_{\mathrm{Agreement}}=\tfrac12+\mathrm{Re}\ \rho^{(LS)[\partial]}_{\Phi,12},$$

so the meters are directly tied to concrete edge observables.

Quadrant patterns at the boundary. Thresholds $\tau_{\mathrm{agr}},\tau_{\mathrm{syn}}\in(0,\tfrac12)$ fix the correspondence (cf. Proposition 8.136):

$$\text{QI or QIV}\ \Rightarrow\ v^{(LS)[\partial]}_{\mathrm{Agreement}}\geq\tfrac12+\tau_{\mathrm{agr}},\qquad \text{QII or QIII}\ \Rightarrow\ v^{(LS)[\partial]}_{\mathrm{Synthesis}}\geq\tfrac12+\tau_{\mathrm{syn}}.$$

In boundary–QIV (explicit breaking) the *agreement* meter stays high as L grows and the edge spectral ledger displays a pNG band on $\mathfrak{P}_\partial$; in boundary–QII (edge SSB) one finds an edge Goldstone and $v^{(S)[\partial]}_{\mathrm{Synthesis}}\to\frac12$ (mixture baseline); in boundary–QIII the *synthesis* meter rises monotonically as the boundary law deficit $d^{[\partial]}_{\mathrm{law}}(L)\downarrow$ (IR healing).

Operational recipe. As a procedure (cf. Definition 8.137):

1. *Classify* the boundary sector with a resolver $\mathcal{R}^{[\partial]}$ using the boundary witnesses (cf. Definition 8.31 and Proposition 8.32).

2. *Calibrate* $\kappa^{[\partial]}$ and compute the meters $\upsilon^{(L)[\partial]}_{\text{Synthesis}}$, $\upsilon^{(S)[\partial]}_{\text{Synthesis}}$, $\upsilon^{(LS)[\partial]}_{\text{Synthesis}}$, $\upsilon^{(LS)[\partial]}_{\text{Agreement}}$.

3. *Threshold* with $\tau_{\text{agr}}, \tau_{\text{syn}}$ to decide I/IV vs. II/III; the 1/2 baseline flags local "bothness" (incoherent mixture).

4. *Cross–check ledgers:* QII $\Rightarrow$ edge Goldstone & conserved/improved edge current; QIII $\Rightarrow$ adjoint divergence vanishes along L and no NG/pNG.

5. *Steer* by varying L (and boundary conditions) to test healing vs. persistence. In open settings ensure the relevant projectors commute with the evolution, or satisfy $\mathcal{L}^{\dagger}(P) = 0$, so meters are conserved/auditable.

Why this overlay matters. It disambiguates bulk–boundary puzzles by scanning the single tangible knob L, turning tolerance–form boundary witnesses (Section 8.1.7) into instrument–grade coherence diagnostics with the same plateau certification and Nielsen flatness. It is the edge realisation of the global synthesis (Section 8.2), expressed in meters that can be audited on certified boundary arenas.

Chapter 8 delivers a *practical, falsifiable pipeline* for deciding whether symmetry is realized, spontaneously broken, emergent, or explicitly violated, while surviving the standard traps of *gauge dependence*, *ghost contamination*, and *scale drift* [114, 38, 117, 32, 112]. The construction fixes all computations *on the band* and *in the physical BRST cohomology* [114, 52] (cf. Definitions 6.272 and 6.282 and Proposition 7.20) and then builds *witnesses*—visibilities in $[0, 1]$—that answer, channel by channel, whether the *laws* are effectively symmetric and whether the *state* remains coherently symmetric [112, 32] (cf. Definition 6.275). Trustworthiness comes from: (i) *Nielsen safety* (gauge–parameter independence on shell) [114, 38, 112] (eqs. (7.201) and (7.209) and Proposition 7.208), (ii) *cohomological inversion* that decouples longitudinal/ghost pieces before inversion [117, 52] (Proposition 7.16 and Definition 6.274), (iii) *Riesz projectors* to extract gauge–independent pole locations [115, 30] (eq. (7.438) and Definition 7.286), and (iv) *plateau certification* in a UV/IR log–scale geometry—small slope/curvature/oscillation—with integrated–readout and discrete–IR controls [32, 112] (Section 6.12.4, eq. (7.28), and Propositions 7.146 and 7.397). With these in hand, the *canonical decision rule* assigns each certified bandpoint to one of the four quadrants (QI–QIV), cleanly separating unbroken, SSB, emergent, and explicit–breaking regimes (Definition 7.685).

Conclusions (synthesis and extensibility). Equally important is the *synthesis* layer and its *extensibility*. The same witness logic is made *sector–conditioned* (TT vs. longitudinal, spin/charge, boundary vs. bulk; Section 8.1 and Definition 8.9) and upgraded by a *parasymmetric overlay* that turns law/state readouts into calibrated "agreement"/"synthesis" meters—well–posed even when law and state disagree (cf. Sections 8.2, 8.2.1 and 8.5.4). Boundary soft/edge modes receive a first–class dashboard, and anomaly, anyon/CS, fracton, hydrodynamic, SUSY, higher–form and non–invertible cases all plug into the same band/BRST/plateau standards [110, 111, 32, 118, 52, 112] (Sections 8.1.2, 8.1.5 and 8.1.8 to 8.1.11). Net effect: a robust, gauge–clean, scale–aware *operational pipeline* that turns

correlators and Ward identities into falsifiable symmetry diagnoses—uniformly across sectors and boundaries [38, 112] (cf. Definitions 6.274 and 7.15).

Chapter 9: The Quanta of Fields

Where Chapter 9 "lives" in scale and what is admissible Chapter 9 partitions the analysis by scale and specifies *what operations are admissible at each level*. There are three regimes: the *PQS core* (no meters/certifications), the *pre-physical band* $[\ell_{\mathrm{PQS}}, \ell_P)$ (operator-level audits only), and the *realized regime* $\ell \geq \ell_P$ (first certified plateaux $\mathfrak{P}$ and projection to the physical BRST cohomology $H^0(\hat{Q}_{\mathrm{BRST}})$). In particular, *plateau certification* $\mathfrak{P}$ and evaluation of physical, channelized visibilities $\left(\Lambda_{\mathrm{law}}, \Lambda_{\mathrm{state}}\right)$ are allowed only in the realized regime; SynTh and parasymmetric meters act only on H^0 and remain Σ-causal and BRST-safe (cf. Definition 9.1, Chapter 9, Lemma 9.3, and Sections 9.5 and 9.16).

A scale-to-sector reading is provided as follows: for $\ell \geq \ell_P$ one passes to physical poles/residues on H^0—e.g. in the Gaiamiti ($\Phi\Phi$) block, the massless spin-2 Goldstone (graviton) and gapped geometric radials; in the Tantrakyma ($\Theta\Theta$) block, NG/pNG/Higgs content is resolved by the quadrant logic (QI–QIV). For $\ell \gg \ell_P$ or $\ell \sim \mathcal{L}_U$, the same H^0 kernels govern, respectively, IR/classical and cosmological coarse-grain limits (consistently with the earlier framework) (cf. Remark 9.4 and Section 9.18.6).

Pre-physical Symmetry Reduction (PSR). PSR is the operator-level stage, valid for relational scales $\ell < \ell_P$, where symmetry content is analysed *prior* to any physical projection or meters. The analysis is carried out entirely in the Σ-centric language using the unified tensor $\hat{\Sigma}_{\mu\nu} = \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu}$ (cf. Definition 5.25) and the Σ-cones determined by the diagnostic realized reference configuration $\bar{\Sigma}_{\mu\nu}$ (cf. Definition 5.77). Within this band-prepared setting, admissible taggers $P_x(\lambda_{\mathrm{UV}})$ and $K^{(\ell)}_{\mathrm{IR}}$ are applied [32, 112], BRST-exact gauge fixing and Nielsen safety are enforced [52, 117, 120, 121, 114], and typed ledgers are assembled for later realized reads (cf. Section 6.12.7). In this regime, plateaux $\mathfrak{P}$ are not certifiable and no projection to the physical cohomology $H^0(\hat{Q}_{\mathrm{BRST}})$ is performed [52] (cf. Definition 9.58).

Outputs and eligibility. Operationally, PSR identifies broken directions and builds Σ-weighted Gram operators to solve generalized eigenproblems, yielding Σ-orthonormal principal systems (Apeiron–Kymon creators) as robust, gauge-clean *candidates* for modes and channels—still pre-physical in status [115, 30] (cf. Definition 5.25). These outputs function as eligibility data: they acquire physical meaning only upon entering the realization regime $\ell \geq \ell_P$, where a plateau $\mathfrak{P}$ is certified [32, 112] (cf. Definition 9.58) and the projection to $H^0(\hat{Q}_{\mathrm{BRST}})$ is taken [52, 117]. There, meters and visibilities $(\Lambda_{\mathrm{law}}, \Lambda_{\mathrm{state}})$ are evaluated within the witness framework on $\mathcal{H}_{\mathrm{phys}}$ [38, 112] (cf. Section 8.1 and Definition 6.275).

Σ-centric upgrade. Chapter 9 recasts the entire Pre-physical Symmetry Reduction (PSR) $\rightarrow$ realization workflow in the unified chronogeometric language of the operator-level tensor

$$\hat{\Sigma}_{\mu\nu} \;:=\; \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu},$$

(cf. Definition 5.25) together with its realized expectation on the diagnostic realized reference configuration,

$$\bar{\Sigma}_{\mu\nu} \;:=\; \langle\, \hat{\Sigma}_{\mu\nu} \,\rangle \;=\; \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu},$$

(cf. Definition 5.26). Hence any construction that was formerly stated "triad-wise" (separately in $\bar{g}$ and $\bar{t}$; cf. Section 4.10.2) is now expressed intrinsically at the Σ-level (i.e., $\bar{\Sigma} = \bar{g} + \bar{t}$ via Definition 5.25), while all causal and variational statements are understood with respect to $\bar{\Sigma}$ and the Σ-cones it determines (cf. Definitions 5.77 and 6.7 and Section 5.4).

Regime stratification (PQS-anchored). The programme is stratified by relational scale, anchored at the PQS core (cf. Section 4.18) and flowing through pre-physical, realized, IR/classical, and cosmological domains:

$$\begin{aligned}\text{PQS core} \;\rightsquigarrow\; \big[\ell_{\mathrm{PQS}}, \ell_P\big) \text{ (pre-physical)} \;\rightsquigarrow\; \\ \ell \geq \ell_P \text{ (realization)} \;\rightsquigarrow\; \ell \gg \ell_P \text{ (IR/classical)} \;\rightsquigarrow\; \ell \sim \mathcal{L}_U \text{ (cosmological)}\,.\end{aligned}$$

Only for $\ell \geq \ell_P$ can a plateau $\mathfrak{P}$ be certified (see Definitions 6.300, 7.47 and 9.58 and Section 6.12.4); below ℓ_P the analysis is strictly pre-physical/operator-level, admitting neither $\mathfrak{P}$ nor physical meters or projections. At eligible scales, BRST projection to the physical sector is well defined and all physical readouts are taken on

$$\mathcal{H}_{\mathrm{phys}} \;=\; H^0(\hat{Q}_{\mathrm{BRST}})\,,$$

with cohomological/Nielsen hygiene inherited from the band-prepared setting (cf. Definitions 6.278, 6.280 and 6.282, Principle 6.1, and Propositions 6.214 and 7.16). Physical witness evaluations then proceed on $\mathcal{H}_{\mathrm{phys}}$ using the established meters and decision rules (cf. Section 8.1 and Definitions 6.275 and 7.685).

Operational eligibility and workflow by regime. *(i) PQS core:* no operational diagnostics apply. *(ii) Pre-physical band* $\big[\ell_{\mathrm{PQS}}, \ell_P\big)$*:* one may employ band-tagging (cf. Definitions 6.278, 6.280 and 6.282), perform Σ-cone causality checks against the realized cones (cf. Definitions 5.77 and 5.122), enforce BRST/Nielsen hygiene (cf. Principle 6.1, Proposition 6.214, and eq. (7.209)), and prepare typed ledgers for later realized reads (cf. Section 6.12.7); however, there is no $\mathfrak{P}$ in this regime, and thus no physical meters. *(iii) Realization* $\ell \geq \ell_P$*:* certify a plateau $\mathfrak{P}$ (cf. Definitions 6.300, 7.47 and 9.58 and Section 6.12.4), project to H^0, evaluate the physical visibilities $(\Lambda_{\mathrm{law}}, \Lambda_{\mathrm{state}})$ within the witness framework (cf. Section 8.1 and Definitions 6.275 and 7.685), and run SynTh/typed meters exclusively on $\mathcal{H}_{\mathrm{phys}}$ (cf. Sections 8.2 and 8.2.1). In every case, computations and causality statements are articulated in the Σ-centric formulation, ensuring compatibility with the Σ-causal structure (cf. Section 5.4 and Definition 6.7) and the certified-plateau criteria referenced above.

Correlator amplitude, band-locality, and Σ–chronogeometry To compute observables, Chapter 9 fixes the *Fourfold correlator amplitude* and its *normalizations* on the band:

$$\mathsf{G}(x, x'; \psi_1, \psi_2) := \Big\langle \mathring{\mathbf{\Psi}} \Big|\, \hat{G}_\Lambda\Big(\mathfrak{B}_x, \mathfrak{B}_{x'}\Big)\, [\psi_1^*, \psi_2] \,\Big|\, \mathring{\mathbf{\Psi}} \Big\rangle, \qquad N(\psi_1, x) := \mathsf{G}(x, x; \psi_1, \psi_1)^{1/2},$$

with frame alignment via $\mathcal{U}(x \leftarrow x'; \gamma)$ and *band-limited locality* enforced by the tag $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$. This construction is compatible with the Fourfold bundle and graded kinematics, preserving Hermiticity/covariance (cf. Definitions 9.5, 9.7, 9.10 and 9.18, Proposition 9.6, eqs. (9.1) and (9.36), and Lemma 9.11).

From these ingredients the *operator-valued chronogeometry* $\hat{\Sigma}_{\mu\nu}$ is built *on the band* by combining correlator-based distances $(d_{\mathrm{corr}}, d_{\mathrm{vac}}, d_{\mathrm{spec}})$ with the spectral generator $\hat{D}$; this is the Σ-object that underwrites cones/hyperbolicity and all subsequent Σ-aware meters (cf. Definitions 9.20 and 9.98, eqs. (9.4) and (9.41), and Proposition 9.108).

Σ-weighted broken-sector mode extraction (pre-physical). In the pre-physical, operator-level stage (PSR; $\ell < \ell_P$), broken directions are isolated by Σ-based contrasts that replace earlier distance-only diagnostics. Concretely, contrasts are formed in the Σ-centric language (cf. Definitions 5.25 and 5.26) and organised using the quadrant logic for symmetry status (cf. Theorem 6.231 and Definition 7.685), providing a Σ-aware selection of the broken-generator set $\mathfrak{B}$. This replaces purely $\bar{g}$-based criteria from the triadic split (cf. Section 4.10.2) and ensures compatibility with the realized causal structure determined by $\bar{\Sigma}$ (cf. Definition 5.77 and Section 5.4).

Step A: find breakings. Given the Σ-centric correlator stack (cf. Definition 6.4 and Section 6.1.1), Σ-contrasts are applied to diagnose departures from the symmetric ledger, yielding the set of broken generators $\mathfrak{B}$. Band hygiene (BRST/Nielsen) is maintained throughout to ensure gauge-stable eligibility data (cf. Principle 6.1, Proposition 6.214, eq. (7.209), and Definitions 6.278, 6.280 and 6.282).

Step B: graded tangents & Σ-weighted Gram operators. For each $X \in \mathfrak{B}$, compute the graded tangent variations $\delta_X \hat{G}_{\Phi\Phi}$ and $\delta_X \hat{G}_{\Theta\Theta}$ of the unified inter-modal correlators (cf. Definition 6.4 and Section 6.1.1) and assemble the corresponding Σ-weighted Gram operators C_Φ and C_Θ. The principal directions are obtained from the generalized eigenproblems

$$C_\Phi\, u_a = \lambda_a\, \hat{\Sigma}^{(\Phi)}\, u_a, \qquad C_\Theta\, v_b = \mu_b\, \hat{\Sigma}^{(\Theta)}\, v_b,$$

posed in the Σ-pre-Hilbert modules determined by the operator tensor $\hat{\Sigma}_{\mu\nu} = \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu}$ (cf. Definition 5.25). By construction these problems are gauge/frame covariant and band-regular (cf. Definitions 6.278 and 6.280), and are compatible with the bundle-theoretic packaging of excitations (cf. Section 6.4 and Definition 6.192).

Step C: principal systems and Apeiron/Kymon creators. Solutions furnish Σ-orthonormal principal systems $\{u_a\} \subset \mathcal{N}_\Phi^{\mathrm{pre}}$ and $\{v_b\} \subset \mathcal{N}_\Theta^{\mathrm{pre}}$ with non-negative spectra, together with Σ-weighted block factorizations that define the Apeiron/Kymon creators (normalisation governed by the Σ-inner product induced from $\hat{\Sigma}$; cf. Definition 5.25). The Σ-weighting renders principal directions intrinsically sensitive to both amplitude curvature and temporal phase-flow via the quadratic form

$$\hat{Q}_x[v] \;=\; v^\mu v^\nu\, \hat{\Sigma}_{\mu\nu}(x) \quad (\text{cf. } \textit{Definitions } 5.25 \textit{and } 6.7),$$

ensuring that $\bar{g}$- and $\bar{t}$-sector information is coherently combined. Off-diagonal intermodal entries in $\hat{G}_{\Theta\Theta}$ are treated faithfully by the same Σ-weighted eigenproblem; when $[\hat{\Sigma}^{(\Theta)}, \delta_X \hat{G}_{\Theta\Theta}] = 0$ the problem block-diagonalises channelwise, otherwise it reduces to a compact positive operator K_Θ with a globally diagonalizable spectrum (cf. Section 6.1.1). Temporal-sector dominance isolates the torsional (Chrono–Kymon) channels as a distinguished type within the Tantrakyma sector (cf. Section 6.1 and Definition 6.141); in the correlator-only or $\hat{\Sigma} \propto \mathbb{I}$ limits the construction collapses to the familiar SVD.

Phase–amplitude split and stability. The Σ-based principal directions carry a phase/amplitude character inherited from the $\bar{t}/\bar{g}$ split (cf. Section 4.10.2), and their spectra and normalisations vary continuously under band deformations. On certified plateaux $\mathfrak{P}$, degeneracy splittings are $\mathcal{O}(\varepsilon)$ with respect to the plateau tolerances (cf. Definitions 6.300, 7.47 and 9.58, Section 6.12.4, and Proposition 7.38).

Status note (pre-physical). All extractions above are pre-physical eligibility data: physical pole structure and residues are to be read only after entering the realization regime $\ell \geq \ell_P$, certifying a plateau $\mathfrak{P}$ and projecting to the physical cohomology $H^0(\hat{Q}_{\mathrm{BRST}})$; the resulting channels persist consistently with Ward/Nielsen identities on $\mathcal{H}_{\mathrm{phys}}$ (cf. Definitions 6.275 and 9.58, eqs. (7.20) and (7.209), and Section 8.1).

Σ–contrasts, continuity, and stability under band deformations Given $\hat{\Sigma}$, the chapter works with the quadratic form $\hat{Q}_x[v] = v^\mu v^\nu \hat{\Sigma}_{\mu\nu}(x)$ and defines *pre-physical, Σ–based contrasts* for the law and state channels. A key technical point is *well-posedness and stability*: for band-regular, Hermitian SOT–C^2 $\hat{\Sigma}$ on the fibre $P_x(\lambda_{\mathrm{UV}})\mathcal{H}_{\mathrm{total}}$, the maps

$$v \longmapsto \Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}}(g;x;v), \qquad v \longmapsto \Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}}(g;x;v)$$

are continuous and bounded; moreover, *small band deformations* $(\lambda_{\mathrm{UV}}, \ell) \mapsto (\lambda'_{\mathrm{UV}}, \ell')$ induce $\mathcal{O}(\varepsilon)$ variations controlled by an effective cutoff $\Lambda_{\mathrm{eff}}(x, x'; \varepsilon)$. This is summarized succinctly as $\mathcal{O}(\varepsilon)$-stability of the contrasts under small band moves (cf. Definitions 9.22, 9.23 and 9.166, eqs. (9.5) to (9.7) and (9.11), Lemma 9.27, and Remark 9.82).

Quadrant logic before realization: the Σ–aware predicate Before ℓ_P, the physical QI–QIV logic is *mirrored* by a *pre-physical, Σ–aware quadrant predicate* at each point x, using the two Σ–contrasts $\Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}}(g;x)$ and $\Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}}(g;x)$ with a small tolerance ε:

$$\textbf{QI}: \ \Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}} < \varepsilon, \ \Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}} < \varepsilon, \quad \textbf{QII}: \ \Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}} < \varepsilon, \ \Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}} > \varepsilon,$$

$$\textbf{QIII}: \ \Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}} > \varepsilon, \ \Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}} < \varepsilon, \quad \textbf{QIV}: \ \Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}} > \varepsilon, \ \Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}} > \varepsilon,$$

with the explicit note that this *mirrors* the physical quadrant logic in $H^0(\hat{Q}_{\mathrm{BRST}})$ (cf. Definitions 9.23 and 9.28 and Section 9.11). Interpretability is aided by a *triadic attribution*: decisions made from Σ–contrasts decompose into contributions from *coherence*, *vacuum*, and *spectral shear* (the "triadic" split of the chronogeometry), and this attribution respects gauge invariance and band locality (cf. Propositions 9.25 and 9.29 and eq. (9.195)).

BRST hygiene of PSR diagnostics and Σ-causality. Band tags $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ built from BRST-closed operators $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ (cf. Definitions 6.278, 6.280 and 6.282) descend to cohomology and render PSR diagnostics valid inputs for realization on $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ (cf. Proposition 7.16). In particular, the band-prepared Ward/Nielsen structure implies gauge-parameter independence for physical readouts once realized (cf. eqs. (7.201) and (7.209)). Replacing separate "metric+temporal" data by the realized chronogeometric carrier $\bar{\Sigma}$ (cf. Definitions 5.25 and 5.26), the quadratic operators $\mathcal{K}_\Psi(\bar{\Sigma})$ and the physical retarded kernels $G^{\mathrm{R}}_{\bullet;\mathrm{phys}}$ (cf. Definitions 6.274 and 7.15) are Σ-*causal*: their supports respect the causal cones determined by $\bar{\Sigma}$, while the cohomological and Ward/Nielsen overlays remain unchanged (cf. Definitions 5.77 and 6.7 and Section 5.4).

Interpretation across regimes. The BRST-closed band preparation and the descent-to-cohomology statement concern the PSR (pre-physical) stage, ensuring that operator-level diagnostics are admissible and stable prior to realization. The Σ-causal retarded kernels $G^{\mathrm{R}}_{\bullet;\mathrm{phys}}$ and their gauge-independent readouts pertain to the realized/physical diagnostic realized reference configuration, where $\bar{\Sigma}$ controls causal support and Ward/Nielsen identities continue to hold verbatim (cf. eqs. (7.201) and (7.209)).

Stability & SVD recovery. The entire Σ-weighted extraction at PSR level is stable under band deformations: eigenvalues, normalizations, and operator solutions vary continuously with $(\lambda_{\mathrm{UV}}, \ell)$. In the correlator-only regime or when $\hat{\Sigma} \propto \mathbb{I}$, the generalized Σ-weighted eigenproblems reduce to the usual SVD with familiar block factorizations (cf. Section 6.1.1). On a certified plateau $\mathfrak{P}$ (cf. Definitions 7.47 and 9.58 and Section 6.12.4), these stability properties carry over to realized/physical objects by the Σ-realization step, preserving the spectral identifications and normalization equivalences established at the operator level.

From Σ-PSR to realized physics. *Plateau & Σ-realization.* Upon certifying a plateau $\mathfrak{P}$ (cf. Definitions 6.300, 7.47 and 9.58 and Section 6.12.4), apply the Σ-realization map $\mathcal{U}_\Sigma$ to obtain the physical chronogeometry

$$\bar{\Sigma}_{\mu\nu} \;=\; \left\langle \mathcal{U}_\Sigma[\hat{\Sigma}_{\mu\nu}] \right\rangle,$$

with $\hat{\Sigma}_{\mu\nu} = \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu}$ (cf. Definitions 5.25 and 5.26). Construct the Σ-*causal* retarded kernels $G^{\mathrm{R}}_{\Phi\Phi;\mathrm{phys}}$ and $G^{\mathrm{R}}_{\Theta\Theta;\mathrm{phys}}$, whose supports respect the cones determined by $\bar{\Sigma}$ (cf. Definitions 5.77 and 6.7 and Section 5.4); cohomological inversion and Ward/Nielsen overlays remain as in the band-prepared formulation (cf. Definitions 6.274 and 7.15, Proposition 7.16, and eqs. (7.201) and (7.209)).

Carry-over of Nielsen safety. On a certified $\mathfrak{P}$, the realized $\bar{\Sigma}$ is gauge-parameter independent; therefore any quantity proven Nielsen-safe in the band analysis retains this property when rewritten as a functional of $\bar{\Sigma}$ (the wording "triad → $\bar{\Sigma}$" changes, not the conclusion) (cf. Proposition 7.208 and eqs. (7.20) and (7.21)).

Normalization equivalence. Particle-level criteria—pole locations, residues, positivity, and Nielsen safety—are invariant under the switch to Σ-inner products; Σ-based normalizations are equivalent to their triadic counterparts up to a positive bounded reweighting induced by $\hat{\Sigma}$ (cf. Definitions 5.25 and 5.26 and eq. (7.12)).

Poles/residues immovable under gauge scans or ζ-exact redefinitions. On $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$, BRST-exact insertions vanish (cf. Principle 6.1); with $\bar{\Sigma}$ Nielsen-flat on $\mathfrak{P}$, the realized poles and residues of the Σ-causal kernels cannot be shifted by gauge-parameter scans or by exact (ζ-exact) redefinitions (cf. eqs. (7.20), (7.21) and (7.209)).

Realization and physical reading (after ℓ_P) Once $\ell \geq \ell_P$, the chapter *projects to the physical sector* and evaluates *physical, channelized visibilities* $\left(\Lambda_{\text{law}}, \Lambda_{\text{state}}\right)$ on certified plateaux $\mathfrak{P}$; SynTh and parasymmetric meters apply here and remain BRST-safe. This is where the *quadrant logic becomes physical* (poles/residues on H^0) and where the Chapter 8 plateau geometry and Nielsen safety can be used *verbatim* (cf. Sections 9.5 and 9.16, Proposition 9.281, Definition 9.63, and Remark 9.82).

Chapter 9 welds a Σ–aware, band-local chronogeometric layer to the existing band/BRST framework. Before ℓ_P it supplies *operator-level, expectation-free* meters (stable under band moves) and a *pre-physical quadrant predicate* that mirrors the physical one. At and above ℓ_P it *hands off* to the Chapter 8 logic: certify $\mathfrak{P}$, project to $H^0(\hat{Q}_{\text{BRST}})$, evaluate $\left(\Lambda_{\text{law}}, \Lambda_{\text{state}}\right)$, and read spectra via poles/residues—now with Σ-built structures ensuring *gauge invariance*, *band locality*, and *robustness* of the decision flow (cf. Definition 9.60 and eqs. (9.26) and (9.27)).

High-level intuition I. At a high level, the Σ-centric picture turns the construction into a single geometric dial that simultaneously captures "shape" (amplitude curvature $\hat{g}$) and "flow" (temporal phase $\hat{t}$) through $\hat{\Sigma} = \hat{g} + \hat{t}$ (cf. Definitions 5.25 and 5.26). In the pre-physical PSR stage ($\ell < \ell_P$), diagnostics live at the operator level: band taggers prepare a clean arena (cf. Definitions 6.278 and 6.280), BRST/Nielsen hygiene ensures descent to cohomology and gauge-stable algebra (cf. Principle 6.1 and Propositions 6.214 and 7.16), and Σ-weighted contrasts select broken directions and principal modes via generalized eigenproblems within the unified correlator stack (cf. Section 6.1.1). The purpose at this stage is to produce eligibility data that are robust under UV/IR deformations, not to make physical claims. Realization begins only on a certified plateau $\mathfrak{P}$ (cf. Definition 9.58): the Σ-realization map $\mathcal{U}_\Sigma$ yields the diagnostic realized reference configuration geometry $\bar{\Sigma}$ (cf. Definition 5.26), retarded kernels become explicitly Σ-causal with support constrained by the $\bar{\Sigma}$-cones (cf. Definitions 5.77 and 6.7 and Section 5.4), and gauge-parameter independence proven on the band protects the ensuing physical readouts (cf. eqs. (7.201) and (7.209) and Definition 6.274).

High-level intuition II. Once realized, the picture is organised by typed subspaces and typed meters on H^0. Projectors $\Pi_S^{[\chi]}$ carve $\mathcal{H}_{\text{phys}}$ into operational slices aligned with the field packaging—e.g., Gaiamiti TT/Tr and Tantrakyma T/L, helicities, and the torsional Chrono–Kymon channel (cf. Sections 6.1 and 8.2 and Definition 6.141). Witnesses then evaluate law-symmetry and state-coherence where they matter (cf. Section 8.1 and Definition 6.275), and the resulting visibilities feed the canonical quadrant logic—QI (unbroken), QII (SSB with NG poles), QIII (emergent symmetry that heals with scale), QIV (explicit breaking with pNG behaviour) (cf. Definition 7.685, Theorems 6.231 and 7.630, and Corollary 7.631). Labels such as spin, helicity, and polarisation are fixed operationally by typed behaviour rather than by names (cf. Section 6.1). Because poles and residues are isolated with Riesz projectors on $\mathfrak{P}$ and read on H^0, they inherit Nielsen safety and BRST consistency

(cf. eqs. (7.20), (7.21), (7.209) and (7.438)): intuitively, PSR is the disciplined rehearsal that isolates the score, while realization is the performance on a certified stage where the acoustics ($\bar{\Sigma}$) and the microphones (typed witnesses) make the physical notes—poles, residues, visibilities—audible and robust.

Typed subspaces, typed meters, and typed quadrants on H^0. *Typed projectors & menus.* On the instrumented diagnostic realized reference configuration, introduce BRST-closed, cone-preserving typed projectors $\Pi_S^{[\chi]}$ that select subspaces compatible with the $\bar{\Sigma}$-causal cones (cf. Principle 6.1 and Definitions 5.77 and 6.7). The allowed menus match the field packaging (cf. Sections 6.1 and 6.4): **Gaiamiti** ($\Phi\Phi$) with TT (spin-2) and Tr (spin-0) types, and optional vectorial admixtures under a $\bar{\Sigma}$-preferred foliation; **Tantrakyma** ($\Theta\Theta$) with T/L types, helicities $\lambda = \pm$ where chiral frames exist, and a torsional/time-like *Chrono–Kymon* channel (spin-1 precursor) treated as its own type (cf. Definitions 5.77 and 6.141). These typed subspaces are constructed to be BRST-closed and thus descend to cohomology on $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$ without ambiguity (cf. Proposition 7.16).

Typed witnesses (SynTh) and drivers. Inserting $\Pi_S^{[\chi]}$ into the SynTh meters (cf. Sections 8.2 and 8.2.1) yields the typed witness $\mathbf{W}_{[\chi]}^{[S]}$, with representative readouts

$$v_{[\chi]}^{(L)} = \left|\text{Res}_{k^2=0}\ \Delta_{\Theta\Theta}^{[\chi]}(k)\right|, \quad v_{[\chi]}^{(S)} = \left\|\Delta_{\Theta\Theta}^{[\chi]}\big(t\to\infty;\,\mathbf{x}=0\big)\right\|_{H^0},$$
$$v_{[\chi]}^{(LS)} = \text{typed cross-channel gain.}$$

evaluated on $\mathcal{H}_{\text{phys}}$ inside a certified plateau $\mathfrak{P}$ and therefore Nielsen-flat by construction (cf. Definitions 6.275 and 9.58, Section 8.1, Proposition 7.208, and eqs. (7.20) and (7.438)).

Typed quadrant predicate & ledger. The canonical typed quadrant $\text{Quad}^{[S,\chi]} \in \{\text{QI}, \text{QII}, \text{QIII}, \text{QIV}\}$ is determined from the typed meters together with healing indicators, reusing the decision rule and overlays developed earlier (cf. Definition 7.685, Theorem 6.231, and Corollary 6.232). Existence, gauge/Nielsen safety, and uniqueness up to $\mathcal{O}(\varepsilon_{\mathfrak{P}})$ follow on $\mathfrak{P}$, yielding a well-defined mapping $(\mathfrak{P}; S, [\chi]) \mapsto \mathbf{L}^{[S,\chi]}$ for the typed ledger (cf. Sections 6.12.4 and 6.12.7 and Definitions 6.300 and 7.47).

Operational fixing of labels. Once $\mathbf{L}^{[S,\chi]}$ is populated, spin/helicity/polarization labels are fixed *operationally* by the chosen $[\chi]$ (independently of particle nomenclature); symmetry-breaking terminology (e.g., "graviton," "Higgs/NG/pNG") is applied only afterwards, consistent with the typed identification and the cohomological/causal structure (cf. Section 6.1 and Definition 5.77).

Typed NG/pNG semantics. If $\text{Quad}^{[S,\chi]} = \text{QII}$, the correlational channel exhibits an exact NG pole with positive residue on H^0 (cf. eqs. (7.12) and (7.438)); if QIV, it exhibits a pseudo-NG band with a GMOR slope certified on $\mathfrak{P}$ (cf. Theorem 7.630 and Corollary 7.631). The remaining cases follow the standard quadrant semantics: QI fully symmetric and QIII without IR NG/pNG realization (cf. Definition 7.685).

Gaiamiti sector (the ($\Phi\Phi$)–block): physical realization. On a certified plateau $\mathfrak{P}$ with readout on $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$ (cf. Definition 9.58), the Gaiamiti dynamics follow from expanding the action about the symmetric anchor and varying the tagged functional to solve

the Euler–Lagrange system

$$\frac{\delta S_\Phi}{\delta \Phi} = 0, \qquad D_\mu \text{ the appropriate (possibly internal) connection,}$$

as in Section 9.17.7 and eqs. (9.107) and (9.108). On an SSB background $\langle\Phi\rangle$, the law channel is exactly flat along the transverse–traceless (TT) tangents of the Gaiamiti manifold: these TT directions form the Goldstone manifold for global breaking in the $(\Phi\Phi)$ block, identifying a massless spin–2 *composite graviton* (cf. Proposition 9.243 and Section 9.17.5). In the two–point sector the TT correlator exhibits an NG pole at $k^2 = 0$ with positive residue,

$$\Delta^{\mathrm{TT}}_{\Phi\Phi}(k) \;=\; \frac{Z_{\mathrm{grav}}}{k^2 + i0^+} \;+\; \text{regular}, \qquad Z_{\mathrm{grav}} > 0,$$

encoded by the residue weights for the composite graviton (cf. Definition 9.265; positivity on $\mathcal{H}_{\mathrm{phys}}$ aligns with eq. (7.12)).

Radial/trace ("Gaiaton") mode is gapped. Orthogonal to the TT manifold, the radial/-trace direction ("Gaiaton") is governed by the Gaiamiti potential scale. The band–prepared Hessian yields a single scalar pole at $k^2 = m_A^2 > 0$ with no ghosts or tachyons on $\mathcal{H}_{\mathrm{phys}}$, captured by the typed radial mass operators in both global and gauged settings (cf. Definitions 9.250 and 9.259; see also Section 9.17.4).

Internal gauge BEH in the Gaiamiti block. If an internal gauge group acts on Φ with $D_\mu = \partial_\mu - ig\,A_\mu$, the SSB background $\langle\Phi\rangle$ induces a Brout–Englert–Higgs (BEH) mass matrix for the broken internal generators. The would–be longitudinal Goldstones are BRST–quartet and unphysical, leaving massive spin–1 vectors in the realized spectrum for broken directions, while unbroken generators remain massless (construction and mass statement in Section 9.17.4 and Proposition 9.257). Gauge–parameter independence of these masses and on–shell visibilities is ensured at realization by the Nielsen identity (cf. Theorem 9.351); explicit U(1) templates appear in Section 9.18.8.

Main Gaiamiti quanta.

- Massless spin–2 *graviton* (TT Goldstone of global SSB): Proposition 9.243 and Definition 9.265.
- Gapped scalar *Gaiaton* (radial/trace excitation): Definitions 9.250 and 9.259.
- *Massive internal vectors* ("mition") from BEH for broken internal generators; massless for unbroken ones: Section 9.17.4, Proposition 9.257, and Theorem 9.351.

BEH of the Gaiamiti (apeiron). We work on a certified plateau $\mathfrak{P}$ with physical readout on $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ (cf. Definition 9.58). The Gaiamiti field Φ has potential $V(\Phi)$ and may transform under an internal gauge group G_{int} with generators T_I and connection $A_\mu = A_\mu^I T_I$.

The Σ-centric kinetic term uses the Diagnostic realized reference configuration tensor $\bar{\Sigma}_{\mu\nu}$ and the covariant derivative [112, 38]

$$D_\mu \Phi = \partial_\mu \Phi - ig\, A^I_\mu T_I \Phi,$$

so the relevant Lagrangian pieces read [38]

$$\mathcal{L}_{\Phi,\text{gauged}} = \tfrac{1}{2}\,\bar{\Sigma}^{\mu\nu}(D_\mu\Phi)^\dagger(D_\nu\Phi) - V(\Phi) - \tfrac{1}{4}\,\bar{\Sigma}^{\mu\alpha}\bar{\Sigma}^{\nu\beta}\,F^I_{\mu\nu}F^I_{\alpha\beta} \qquad F^I_{\mu\nu} = \partial_\mu A^I_\nu - \partial_\nu A^I_\mu + g\, f^I{}_{JK}\, A^J_\mu A^K_\nu.$$

Band instrumentation is BRST-closed and typed projectors commute with the setup [52, 117, 114]. The SSB background $\langle\Phi\rangle = v$ breaks $G_{\text{int}} \to H$ and simultaneously fixes the TT *Goldstone sheet* in the $(\Phi\Phi)$ block and the orthogonal *radial/trace* (Gaiaton) direction (cf. Sections 9.17.4 and 9.17.7, eqs. (9.107) and (9.108), Proposition 9.243, and Definition 9.250); the BEH interpretation aligns with the standard gauge/Goldstone structure [38, 112].

Gauge–boson masses from $\langle\Phi\rangle$ (BEH matrix). Expanding $\Phi = v + \varphi$ in $\mathcal{L}_{\Phi,\text{gauged}}$, the covariant term yields a vacuum–algebraic quadratic piece in A_μ:

$$\mathcal{L}^{(A)}_{\text{mass}} = \tfrac{1}{2}\,\bar{\Sigma}^{\mu\nu}\,A^I_\mu (M^2)_{IJ} A^J_\nu, \qquad (M^2)_{IJ} := g^2\,\langle\, T_I v,\ T_J v\,\rangle_\Phi \quad (\geq 0).$$

Equivalently, in representation–matrix form [38, 112],

$$(M^2)_{IJ} = g^2\, v^\dagger \{T_I T_J\}_{\text{sym}} v, \qquad \{T_I T_J\}_{\text{sym}} := \tfrac{1}{2}\,(T_I T_J + T_J T_I).$$

Each *broken* generator $X_a \in \mathfrak{g}/\mathfrak{h}$ contributes a non–negative direction in $(M^2)_{IJ}$, while *unbroken* generators $H_\alpha \in \mathfrak{h}$ satisfy $T_{H_\alpha} v = 0$ and give *massless* vectors (cf. Section 9.17.4 and Proposition 9.257). Nielsen safety on $\mathfrak{P}$ ensures gauge–parameter independence [120, 121, 114] (cf. Theorem 9.351):

$$\frac{\partial}{\partial\xi}\,\det\!\Big[q^2\bar{\Sigma}^{\mu\nu}\delta^{IJ} - (M^2)^{IJ}\bar{\Sigma}^{\mu\nu}\Big] = 0 \implies \partial_\xi\, m^2_{\text{vector},\lambda} = 0 \quad \text{on } \mathfrak{P}.$$

Goldstones and BRST quartet decoupling. Decompose the apeiron fluctuation by the chapter's typed projectors in $(\Phi\Phi)$:

$$\varphi = \varphi_\parallel + \varphi_{\text{rad}}, \qquad \varphi_\parallel = \sum_I \alpha^I\, T_I v,$$

so $\varphi_\parallel$ spans the tangent (would–be Goldstone) directions and φ_{rad} the radial/trace (Gaiaton). The linearized equations identify $\varphi_\parallel$ as gauge directions; in BRST quantization they form *quartets* and are absent in $\mathcal{H}_{\text{phys}}$ [117, 52]. The mixing

$$g\,\bar{\Sigma}^{\mu\nu} A^I_\mu\,\langle T_I v,\ \partial_\nu\varphi\rangle_\Phi$$

is removed by gauge fixing/BRST completion, leaving the longitudinal modes unphysical while the transverse vector spectrum is governed by $(M^2)_{IJ}$ [38, 109] (cf. Section 9.17.4, Proposition 9.257, and Theorem 9.351).

Radial (trace) Gaiaton in the gauged theory. Orthogonal to the tangent sheet, the Gaiaton is the radial/trace amplitude mode. Its mass is fixed by the typed *radial mass operator*: globally by $\hat{\mathbb{M}}^{(\mathrm{rad})}_{\mathrm{glob}}$ and, in the gauged case, by the *apeiron radial mass operator (gauged)* $\hat{\mathbb{M}}^{(\mathrm{rad})}_{\mathrm{gauged}}$ (cf. Definitions 9.250 and 9.259). At quadratic order on $\mathfrak{P}$,

$$\mathcal{L}_{\mathrm{rad}} = \tfrac{1}{2}\,\bar{\Sigma}^{\mu\nu}\,\partial_\mu\varphi_{\mathrm{rad}}\partial_\nu\varphi_{\mathrm{rad}} - \tfrac{1}{2}\,m_A^2\,\varphi_{\mathrm{rad}}^2, \qquad m_A^2 = \langle\varphi_{\mathrm{rad}},\ \hat{\mathbb{M}}^{(\mathrm{rad})}_{\mathrm{gauged}}\ \varphi_{\mathrm{rad}}\rangle_\Phi > 0,$$

so the realized two–point function exhibits a single scalar pole at $k^2 = m_A^2$ with no ghost-s/tachyons on $\mathcal{H}_{\mathrm{phys}}$ [38, 52, 117]. On–shell gauge–parameter independence (Nielsen safety) follows from the Nielsen identities in the BFM setting [120, 121, 114] (cf. Theorem 9.351).

Typed/Σ–causal structure and decoupling from the graviton. The Gaiamiti BEH reorganizes the *internal spin–1* sector (vectors A_μ) and the *spin–0* radial sector; it does *not* induce a mass for the spin–2 graviton. The spin–2 carrier is the York–projected TT metric fluctuation, for which the TT block remains massless with positive residue (composite TT Goldstone) and is insulated from vector BEH [43, 44, 45, 38, 112]; observationally, bounds are consistent with a (nearly) massless graviton [122]. All propagators and vertices are $\bar{\Sigma}$–causal, and on $\mathfrak{P}$ all on–shell data (poles, residues, masses) are Nielsen–safe [120, 121, 114] (cf. Theorem 9.351).

Broken internal directions acquire vector masses from $(M^2)_{IJ} = g^2\langle T_I v, T_J v\rangle_\Phi$, while unbroken ones remain massless; would–be Goldstones decouple as BRST quartets [38, 52, 117] (cf. Section 9.17.4 and Proposition 9.257). The Gaiaton is a single gapped scalar with $k^2 = m_A^2 > 0$, determined by the typed radial operator (cf. Definitions 9.250 and 9.259 and Section 9.17.4). The TT graviton remains massless (TT Goldstone of global SSB in $(\Phi\Phi)$); there is no BEH leakage into spin–2 [113, 109]. All on–shell statements are Nielsen–safe on $\mathfrak{P}$ [120, 121, 114] (cf. Theorem 9.351).

The Tantrakyma sector (the $(\Theta\Theta)$–block): physical realization. Work on a certified plateau $\mathfrak{P}$ with readout on $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ (cf. Definition 9.58). The sector is instrumented by BRST–closed typed projectors and a permitted kinematics that decompose Θ into operational subspaces (set–up and bookkeeping in Sections 9.15.3 and 9.17.11, projectors in Definition 9.295). Linearising the band Hessian about the SSB background $\langle\Theta\rangle$ yields the typed Hessian and the corresponding typed linear–response system,

$$\frac{\delta S_\Theta}{\delta\Theta} = 0 \quad (\text{in each type})\,,$$

whose spectra are then read physically on $\mathfrak{P}$ (cf. eqs. (9.124) and (9.125)). These diagnostics are evaluated with the same cohomological/Nielsen hygiene as elsewhere, now within the $(\Theta\Theta)$ block and its typed slices.

Global SSB $\to$ scalar NG (or multiplet) in the Θ directions. Broken global generators in the Θ directions yield Nambu–Goldstone scalar(s) that are gapless in the large–volume limit, while unbroken generators leave modes massless or gapped according to the typed pattern selected by Definition 9.295 [32, 112, 38]. The existence and reading of these modes follow from the typed Hessian and linear–response equations (cf. eqs. (9.124) and (9.125)).

Internal gauge BEH in the Tantrakyma block. If Θ carries an internal gauge action, broken internal directions generate a Brout–Englert–Higgs (BEH) mass matrix for spin–1 vectors; the associated Tantrakyma would–be Goldstones are removed from the physical spectrum by quartet decoupling, while the unbroken internal subgroup yields massless vectors (the "tantron") [38, 112, 52, 117] (cf. Definition 9.345 and Proposition 9.347). Gauge–parameter independence of these on–shell masses and readouts at realization is ensured by the Nielsen identity [120, 121, 114] (cf. Theorem 9.351).

Pseudo–NG (pNG) when mild explicit breaking is present. With small explicit terms in the Θ potential, the would–be NG acquires a small mass squared determined by the *typed* mass operator extracted from the potential; explicit formulae and parameter relations are given in eqs. (9.175), (9.190) and (9.193) [112, 32].

Main Tantrakyma quanta (realized on $\mathfrak{P}$).

- *NG scalar(s)* of global SSB in the Θ directions (gapless in the large–volume limit): eqs. (9.124) and (9.125).
- *Massive internal vectors* from BEH for broken internal directions; *massless internal vectors* ("tantron") for unbroken ones: Definition 9.345, Proposition 9.347, and Theorem 9.351.
- *Radial Tantrakyma scalar* (Higgs–like amplitude mode) gapped; light pNG in the presence of small explicit breaking: eqs. (9.175) and (9.193).

BEH of the Tantrakyma (kymon). Work on a certified plateau $\mathfrak{P}$ and read on $\mathcal{H}_{\rm phys} = H^0(\hat{Q}_{\rm BRST})$. The Tantrakyma field Θ is instrumented by BRST–closed *typed projectors* and permitted kinematics that decompose Θ into operational subspaces [52, 117, 114] (set–up in Sections 9.15.3 and 9.17.11, projectors in Definition 9.295). When Θ transforms under an internal gauge group $G_{\rm int}$ with generators T_I and connection $A_\mu = A^I_\mu T_I$, the Σ–centric realized kinetic term and potential are [38, 112]

$$D_\mu\Theta = \partial_\mu\Theta - ig\,A^I_\mu T_I\Theta, \qquad \mathcal{L}_{\Theta,\text{gauged}} = \tfrac{1}{2}\,\bar{\Sigma}^{\mu\nu}(D_\mu\Theta)^\dagger(D_\nu\Theta) - V(\Theta) \;-\; \tfrac{1}{4}\,\bar{\Sigma}^{\mu\alpha}\bar{\Sigma}^{\nu\beta}F^I_{\mu\nu}F^I_{\alpha\beta},$$

with $F^I_{\mu\nu} = \partial_\mu A^I_\nu - \partial_\nu A^I_\mu + g f^I{}_{JK} A^J_\mu A^K_\nu$ [38]. An SSB background $\langle\Theta\rangle = w$ breaks $G_{\rm int} \to H$ and sets the typed decomposition used to read the spectrum via the *typed Hessian* and linear response (cf. eqs. (9.124) and (9.125)) [112, 38].

Vector masses from $\langle\Theta\rangle$ (Tantrakyma BEH matrix). Expanding $\Theta = w + \vartheta$ in $\mathcal{L}_{\Theta,\text{gauged}}$, the covariant term yields a vacuum–algebraic quadratic piece in the gauge fields:

$$\mathcal{L}^{(A)}_{\rm mass} = \tfrac{1}{2}\,\bar{\Sigma}^{\mu\nu}\,A^I_\mu(M^2_V)_{IJ}A^J_\nu, \qquad (M^2_V)_{IJ} := g^2\,\langle\, T_I w,\; T_J w\,\rangle_\Theta \;\; (\geq 0).$$

Equivalently, in representation–matrix form [38, 112],

$$(M^2_V)_{IJ} = g^2\,w^\dagger\{T_IT_J\}_{\rm sym}w, \qquad \{T_IT_J\}_{\rm sym} := \tfrac{1}{2}\,(T_IT_J + T_JT_I).$$

Broken internal directions $X_a \in \mathfrak{g}/\mathfrak{h}$ give strictly non–negative directions in $(M_V^2)_{IJ}$ and hence *massive* spin–1 vectors, whereas unbroken $H_\alpha \in \mathfrak{h}$ satisfy $T_{H_\alpha} w = 0$ and yield *massless* vectors ("tantrons") (cf. Definition 9.345 and Proposition 9.347). On $\mathfrak{P}$, the on–shell vector masses are *Nielsen–safe* [120, 121, 114]:

$$\frac{\partial\, M_{V,\lambda}^2}{\partial \xi} = 0 \quad \text{for all eigenmodes } \lambda, \qquad \text{(cf. Theorem 9.351)}.$$

Goldstones and BRST quartet decoupling. Use the typed projectors of Definition 9.295 to split the fluctuation into *tangent* (would–be Goldstone) and *radial* pieces,

$$\vartheta = \vartheta_{\parallel} + \vartheta_{\mathrm{rad}}, \qquad \vartheta_{\parallel} = \sum_I \beta^I\, T_I w.$$

The mixing term $g\,\bar{\Sigma}^{\mu\nu} A_\mu^I \langle T_I w,\ \partial_\nu \vartheta\rangle_\Theta$ is removed by gauge fixing/BRST completion [52, 117]; the $\vartheta_{\parallel}$ modes become *unphysical* (quartets) and do not appear in $\mathcal{H}_{\mathrm{phys}}$, while the transverse vector spectrum is governed by $(M_V^2)_{IJ}$ as above [38, 112] (cf. Definition 9.345, Proposition 9.347, and Theorem 9.351).

Radial amplitude and pseudo–NG masses. Orthogonal to the tangent sheet, the *radial Tantrakyma* (Higgs–like amplitude) mode is gapped, with its mass read from the typed Hessian of the potential in the appropriate projector slice (cf. eqs. (9.124) and (9.125)) [38, 112]. When *mild explicit breaking* is present in $V(\Theta)$, the would–be NG scalars acquire *pseudo–NG* (pNG) masses determined by a typed mass operator extracted from the potential, with explicit formulae given in

$$m^2_{\mathrm{pNG},[\chi]} = \lambda\Big(\mathbb{M}^{[\chi]}_{\mathrm{typed}}\Big), \qquad \text{cf. } \textit{eqs.}\ (9.175), (9.190) \textit{and}\ (9.193),$$

and these masses inherit gauge–parameter independence at realization (Nielsen safety) [120, 121, 114] (Theorem 9.351).

Typed/Σ–causal structure and spectrum summary. All readouts are taken in the typed slices fixed by $\Pi_\Theta^{[\chi]}$ and propagate with Σ–causal support; the spectrum is therefore operationally defined and BRST/Nielsen–clean on $\mathfrak{P}$ [52, 114]. In summary: broken internal directions in the Tantrakyma sector generate *massive spin–1 vectors* via the BEH matrix $(M_V^2)_{IJ}$, unbroken directions yield *massless* tantrons [38, 112] (cf. Definition 9.345 and Proposition 9.347); the *radial Tantrakyma* amplitude mode is *gapped*; and with small explicit breaking the *NG scalars* become *pNG* with masses fixed by the *typed* mass operator [112, 32] (cf. eqs. (9.175), (9.190) and (9.193)). All on–shell masses and residues are *Nielsen–safe* at realization [120, 121, 114] (Theorem 9.351).

Joint reading and provenance

- **Quadrant logic.** The Gaiamiti TT block realises a QI/QIII–like law reading in the TT channel (exact or IR–healing) with a gapped trace, while the Tantrakyma block separates cleanly into state versus law channels per the Chapter 8/9 witness pair: global

SSB appears in the state channel (NG/pNG), internal BEH in the gauge sector. All readouts are plateau–certified and Nielsen–safe in the realised regime; cf. Corollary 9.139 and Theorem 9.351.

- **Spectral hygiene.** Two–point functions in both blocks exhibit positive spectral weights in $\mathcal{H}_{\text{phys}}$; pole/residue data are gauge–parameter independent within the audit window (Nielsen safety), and mixing with unphysical sectors is removed by quartet decoupling (cf. Definition 9.265 and Theorem 9.351).

In short.

- **Gaiamiti**: global SSB with TT Goldstone identified as the composite graviton (Proposition 9.243), a gapped radial mode (Definitions 9.250 and 9.259), and, when internally gauged, BEH–massive vectors (Section 9.17.4 and Proposition 9.257).
- **Tantrakyma**: global SSB with NG scalar(s) (or light pNG if small explicit breaking is present), a gapped radial mode, and—under internal gauging—BEH–massive vectors for the broken subgroup (massless for the unbroken one) (eqs. (9.124) and (9.175), Definition 9.345, Proposition 9.347, and Theorem 9.351).

Here, Chapter 9's realized spectrum is organised by the certified plateau $\mathfrak{P}$ and the physical cohomology $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$, so every on-shell statement inherits BRST/Nielsen protection and Σ-causality [114, 52, 117, 120, 121, 112]. In the *Gaiamiti* ($\Phi\Phi$) block, an SSB background $\langle\Phi\rangle$ renders the law channel exactly flat along the transverse–traceless tangents of the Gaiamiti manifold; these TT directions are the Goldstone manifold and realize a *massless spin–2 composite graviton* [43, 44, 45, 122]. In the two-point sector this appears as an NG pole at $k^2 = 0$ with positive residue, read as a genuine, positivity-respecting excitation on H^0 (cf. Proposition 9.243, Definition 9.265, and eq. (7.12)). Orthogonal to TT lies the *radial/trace ("Gaiaton")* amplitude mode, whose scale is set by the Gaiamiti potential; on the band-prepared Hessian it is *gapped* with a single scalar pole at $k^2 = m_A^2 > 0$ and no ghosts/tachyons on $\mathcal{H}_{\text{phys}}$ [38, 52, 112] (cf. Definitions 9.250 and 9.259 and Section 9.17.4). If an internal gauge symmetry acts on Φ, the realized picture includes a Brout–Englert–Higgs mechanism: broken internal generators yield *massive spin–1 vectors* (the "mition"), with the would-be Goldstones removed by the quartet mechanism, while unbroken generators remain *massless*; masses and on-shell visibilities are Nielsen-safe on $\mathfrak{P}$ [38, 52, 117, 120, 121, 114] (cf. Sections 9.17.4 and 9.18.8, Proposition 9.257, and Theorem 9.351).

Complementarily, the *Tantrakyma* ($\Theta\Theta$) block is the "matter-like" half. BRST-closed typed projectors and permitted kinematics decompose Θ into operational subspaces; linearising the typed Hessian about $\langle\Theta\rangle$ furnishes the typed linear-response system whose spectra are read physically on $\mathfrak{P}$ [52, 117] (cf. Sections 9.15.3 and 9.17.11, Definition 9.295, and eqs. (9.124) and (9.125)). Broken global generators along Θ produce *Nambu–Goldstone scalar(s)*, gapless in the large-volume limit, while unbroken directions remain massless or acquire a gap according to type [32, 112, 38]. A *radial (Higgs-like) amplitude mode* is generically gapped, setting the intrinsic scale of the sector [38]. With an internal gauge action on Θ, the typed structure cleanly realizes *BEH*: broken directions give *massive spin–1 vectors*, the associated would-be Tantrakyma Goldstones decouple as quartets, and the unbroken subgroup

yields *massless "tantrons"*; on-shell masses and residues are again Nielsen-safe at realization [38, 52, 117, 120, 121, 114] (cf. Definition 9.345, Proposition 9.347, and Theorem 9.351). Small explicit breakings in the Θ potential bend the NG story into *pseudo–NG (pNG)* behaviour with parametrically light masses fixed by a typed mass operator [112, 32] (cf. eqs. (9.175), (9.190) and (9.193)).

Taken together, Chapter 9's conclusion is operational and robust: on a certified $\mathfrak{P}$, the *geometry-generating* Gaiamiti block contributes a *gapless TT Goldstone (graviton)* and a *gapped radial Gaiaton*, plus *BEH-driven vectors* if gauged; the *matter-like* Tantrakyma block contributes *NG scalar(s)* aligned with its broken directions, a *gapped radial amplitude mode*, *massive* (broken) and *massless* (unbroken) internal vectors when gauged, and controlled *pNG* masses under mild explicit breaking. Because all readouts live on H^0 and respect the band/Nielsen stack, pole locations, residues, and masses are *gauge-parameter independent* and Σ-causal—delivering a physically stable, typed ledger of the realized spectrum in the Fourfold framework [114, 52, 120, 121, 38, 112, 122, 32].

Chapter 10: The Weave of Geometry and Matter

Chapter 10 fixes the chronogeometric realized weave configuration $(\bar{\Sigma}, \mathcal{I}^{[\mathscr{S}]})$ and explains how observable spin–2 content is cleanly isolated from temporal/coherence dynamics by the weave's TT projector. The centerpiece is the τ–kairos sector—defined as $\hat{\tau}_{\mu\nu} := \delta\hat{t}_{\mu\nu}$ and made *TT–null* by construction—whose typed/triad pipeline extracts scalar, pseudoscalar, and Chrono–Kymon (CK) vector registers from the geometric $\Phi\Phi$ block, promotes them through meters, and stamps T–L–O onsets without ever polluting the TT graviton readout. The chapter then integrates (i) the bi-local interaction kernel that preserves TT closure at the level of variations, (ii) the Goldstone interpretation of the graviton, and (iii) species-resolved whitening of operator eigen-operator pairs, before distinguishing the *connection* ledger—where the mition and tantron live—from τ's temporal ledger. What follows is a cohesive, equation-forward summary that weaves these strands into a single narrative consistent with Chapter 11's typed/triad formalism.

The τ–kairos sector, its typed/triad weave, and the connection quanta

Definition and TT–nullity. The temporal/coherence channel of chronogeometry is the *τ–kairos* sector,

$$\hat{\tau}_{\mu\nu} := \delta\hat{t}_{\mu\nu}, \qquad \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\hat{\tau}_{\mu\nu} = 0,$$

so τ is *TT–null* by construction. Equivalently, $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\Pi^{[\chi]} = 0$ for any non-TT type $[\chi]$ admitted by the weave (Definition 10.162 and Lemma 10.17). This structural TT–nullity guarantees that the observable spin-2 graviton lives purely in the metric TT channel and cannot be contaminated by temporal/coherence fluctuations.

Bimodal–triadic organization and typed decomposition. Chronogeometric two-point functions are organized bimodally (metric $\leftrightarrow$ temporal) and refined by triad provenance

$a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$. Fixing the channel operators

$$\hat{h}_{\mu\nu} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \delta\hat{g}_{\mu\nu}, \qquad \hat{\tau}_{\mu\nu} := \delta\hat{t}_{\mu\nu},$$

with $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\hat{\tau}_{\mu\nu} = 0$, Chapter 11 resolves the τ tensor into scalar/pseudoscalar images and vector Chrono–Kymon (CK) helicities:

$$\boxed{\hat{\tau}_{\mu\nu} \;=\; S^{(b)}_{\mu\nu}[\hat{\varphi}_b] \;\oplus\; S^{(L)}_{\mu\nu}[\hat{\varphi}_L] \;\oplus\; \tilde{S}^{(p)}_{\mu\nu}[\tilde{\varphi}] \;\oplus\; \sum_{\lambda=\pm} A^{(\lambda)}_{\mu\nu}\Big[\hat{K}^{(\tau)}_{\lambda}\Big], \qquad \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\hat{\tau}_{\mu\nu} = 0.} \tag{4}$$

Here $S^{(b)}, S^{(L)}, \tilde{S}^{(p)}$ encode the scalar/pseudoscalar register, and $A^{(\pm)}$ carry the CK vector helicities (Definition 10.4 and Lemma 10.17). The CK register admits the gauge-covariant field-strength representative

$$\hat{\mathcal{F}}^{(\tau)}_{\mu\nu} = 2\,\bar{\nabla}_{[\mu}\hat{K}^{(\tau)}_{\nu]},$$

fixing the notation used by the typed projectors (Definition 10.57). The full fluctuation splits triadically as

$$\delta\hat{\Sigma}_{\mu\nu} = \delta\hat{\Sigma}^{(\mathrm{corr})}_{\mu\nu} + \delta\hat{\Sigma}^{(\mathrm{vac})}_{\mu\nu} + \delta\hat{\Sigma}^{(\mathrm{spec})}_{\mu\nu}, \qquad \hat{h}_{\mu\nu} = \sum_a \hat{h}^{(a)}_{\mu\nu}, \quad \hat{\tau}_{\mu\nu} = \sum_a \hat{\tau}^{(a)}_{\mu\nu}\ (\textit{Definition } 10.187).$$

Physical diagonal blocks and TT readout. The TT graviton propagator is the TT-projected metric-channel correlator,

$$G^{\mathrm{TT}}_{\mu\nu\rho\sigma} := \langle\Omega|\, T\{\hat{h}_{\mu\nu}\hat{h}_{\rho\sigma}\}\,|\Omega\rangle \;=\; \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, G^{hh}\, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \quad (\textit{eq. } (10.114)),$$

while the $\tau\tau$ block carries phase-flow/coherence physics and drives entanglement dynamics without mediating classical forces. Its typed entries $\chi \in \{\tau_S, \tau_A, K^{(\tau)}_{\mathrm{T}}, K^{(\tau)}_{\mathrm{L}}\}$ are TT-null and feed only non-TT T–L–O stamps (Definitions 10.4 and 10.13).

Off-diagonal mixing and TT closure. Off-diagonal correlators $\langle T\{\hat{h}, \hat{\tau}\}\rangle$ encode state-dependent metric–temporal conversion amplitudes but carry *no* spin-2 content. Projecting with $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ annihilates these blocks, so they cannot generate TT modes; equivalently the mixed correlator TT-projects to zero (TT-closure, Proposition 10.184). Thus τ produces *no* gravitational-wave polarizations (Corollary 10.177).

Typed matter channels, positivity, and cone–gating. Matter in the temporal/CK families is organized by typed projectors $\Pi^{[\chi]}$ with $\chi \in \{S^{(b)}, S^{(L)}, \tilde{S}^{(p)}, A^{(\pm)}\}$. On a certified plateau $\mathfrak{P}$, each admitted block

$$\hat{\mathbf{G}}^{[\Theta\to\Theta;\chi]}_{\Lambda;a} := \Pi^{(a)}\Pi^{[\chi]}\, \hat{\mathbf{G}}^{[\Theta\to\Theta]}_{\Lambda}\, \Pi^{[\chi]}\Pi^{(a)}$$

is a positive semidefinite covariance propagating within $\bar{\Sigma}$-cones (admission & hyperbolicity; Definition 10.206 and Corollary 10.207).

Bi-local interaction kernel and variational sources. Interaction between the geometric and kairotic sectors is bi-local and Hermitian,

$$\mathbf{U}_{\Lambda}^{[\Phi\leftrightarrow\Theta]}(x,x') := \hat{\mathbf{G}}_{\Lambda}^{[\Phi\to\Theta]}(x,x') + \Big(\hat{\mathbf{G}}_{\Lambda}^{[\Theta\to\Phi]}(x',x)\Big)^{\dagger},$$

triad-resolved into $\{\mathrm{spec},\mathrm{vac},\mathrm{corr}\}$ (*eqs.* (10.175)*and* (10.176)).

The bi-local interaction density reads

$$\mathcal{L}_{\mathrm{int}}(x,x') := \mathrm{Tr}\Big[(\delta\hat{\mathbf{\Phi}}(x))^{\dagger}\,\mathbf{U}_{\Lambda}^{[\Phi\leftrightarrow\Theta]}(x,x')\,\delta\tilde{\hat{\mathbf{\Theta}}}(x'\to x)\Big] \quad (Definition\ 10.222),$$

and first variations yield bi-local sources in the Euler–Lagrange equations while preserving TT closure upon projection (Proposition 10.224 and Remark 10.225).

T–L–O augmentation (ledger → meters → stamps). On a certified plateau $\mathfrak{P}$, the sector-execution functor $\mathscr{E}_{\bullet}$ augments $\mathbf{G}_{\Sigma}^{(2)}$ by: (i) typed admission and cone-gating; (ii) typed visibilities from correlator spectra; (iii) parasymmetric meters producing Onset windows $(r^{\star},t^{\star})$; (iv) stamping $\mathrm{TLO}^{[\chi]}$ with Location and Type consistent with Chapter 11's spin/polarization scope (Proposition 10.185). BRST/Nielsen safety and uniqueness follow from the weave propositions.

Apeiron (trace scalar) vs. graviton (TT) in $\Phi\Phi$. Within the geometric block, the trace-typed scalar channel is the **Apeiron**, defined by typed insertion/triad resolution; the *observable* graviton is the TT refinement of the same $\Phi\Phi$ block via $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$. All non-TT types are strictly TT-null, ensuring clean spin-2 readout (eqs. (10.84) and (10.85)).

Typed sourcing of τ and particle content. Chapter 10 records τ sourcing without new primitives:

$$\hat{\tau}_{\mu\nu} = S_{\mu\nu}^{(b)}[\hat{\varphi}_b] \oplus S_{\mu\nu}^{(L)}[\hat{\varphi}_L] \oplus \tilde{S}_{\mu\nu}^{(p)}[\tilde{\varphi}] \oplus \sum_{\lambda=\pm} A_{\mu\nu}^{(\lambda)}[\hat{K}_{\lambda}^{(\tau)}],$$

with $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\hat{\tau}_{\mu\nu}=0$. Vector quanta in the temporal channel are CK modes; the TT graviton $\hat{\gamma}_{\mu\nu}$ is the Goldstone-tangential excitation of the metric channel (Remark 10.175).

Goldstone interpretation and induced dynamics. The operator mass $\mathcal{M}(x)$ arises from the Hessian of the Gaiamiti potential; flat directions yield Goldstones. The massless spin-2 graviton is the Goldstone of tangential Φ variations, projected to TT. The τ sector, while TT-null, may host scalar/vector modes with model-dependent masses. Linear response of the constitutive map determines how $\delta\hat{g}_{\mu\nu}$ and $\delta\hat{t}_{\mu\nu}$ inherit dynamics; the TT graviton is

$$\hat{h}_{\mu\nu} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}_{\mu\nu} \quad (eqs.\ (10.103)and\ (10.125)and\ Section\ 10.5).$$

Species resolution and operator-generalized eigen-operator pair whitening. Species-resolved TT projectors preserve the $\{\mathrm{spec},\mathrm{vac},\mathrm{corr}\}$ provenance. Species mixing is diagonalized via a $\bar{\Sigma}$-weighted operator-generalized eigen-operator pair eigenproblem $\mathcal{A}u=\lambda\,\mathcal{B}u$, with whitening $w=B^{-1/2}u$ so that $B^{-1/2}\mathcal{A}B^{-1/2}$ is self-adjoint in the reference inner product (Definitions 10.116 and 10.117).

Framework-level dynamics (closure and back-reaction). Varying the action w.r.t. Σ yields field equations where Tantrakyma contributes ordinary and coherence stress. Crucially, since $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\Pi^{[\chi]} = 0$, the spin-2 sector is dynamically closed: Tantrakyma/kymon effects sculpt $t_{\mu\nu}$ and its fluctuations, contribute to total stress, and leave the TT graviton sector untouched. T–L–O stamps log *when/where* temporal influences switch on (Section 10.17.4, Lemma 10.17, and Propositions 10.22 and 10.65).

Relation to Chapter 7 (backbone → readout). Chapter 10's $\Phi\Phi$ blocks and their TT/triad refinements are *typed projections* of the unified inter-modal correlator $\hat{\mathbf{G}}_\Lambda$ introduced in Chapter 7: one first fixes the realized weave configuration $(\bar{\Sigma}, \mathcal{I}^{[\mathscr{S}]})$ and then projects $\Phi\Phi \to$ triad $\to$ TT, ensuring compatibility with relational covariance and TT-nullity of non-TT types (Definitions 10.157 and 10.158 and eq. (10.83)).

TT isolation of the graviton and the $\Theta\Theta$ register. By TT isolation the graviton is

$$\hat{h}_{\mu\nu} = \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\, \delta\hat{g}_{\mu\nu} = \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\, \delta\hat{\Sigma}_{\mu\nu},$$

with τ annihilated by $\Pi^{[\mathrm{TT}]}$ (Lemma 10.172). Higgs/radial kymon content sits in the $\Theta\Theta$ block and interfaces with τ only via the shared typed/ledger pipeline (never by identification).

Where τ is *operational*. τ dominates in coherence-rich regimes—laboratory interferometry, early-universe epochs, near-horizon domains—where admissions/onsets are metered and TT purity is continuously witnessed by the T–L–O layer (Section 10.16).

Connection quanta: *mition* and *tantron*. A distinct strand concerns spin-1 excitations of the universal relational operator-bundle connection $\mathcal{A}_\mu$: the *mition* and *tantron*. They are *not* substrate quanta of Φ or Θ; after BEH breaking their transverse modes are massive, BRST-physical, with Proca-like content (Definitions 6.192, 9.258 and 9.348). Ledger-wise they live in the *connection* register and influence $\Sigma_{\mu\nu}$ only indirectly through the constitutive mapping $\Sigma[\Phi, \Theta]$ (eq. (10.101)); they are therefore *not* τ–kairos registers. Their projectors commute with the typed/triad structure and remain TT-orthogonal, so the graviton TT readout stays purely geometric (Proposition 10.9 and Definition 10.3). On a realized weave configuration they are credible dark-sector candidates (misalignment/freeze-in) subject to portal constraints; Section 10.16 summarizes: connection origin, BEH-massive transverse modes, typed/triad isolation, TT-safeguarding of the graviton, and cosmological viability.

One-line map (quick recall, retained verbatim). τ–kairos = TT-null temporal/coherence tensor with $\{\text{scalars } S^{(b)}, S^{(L)}, \tilde{S}^{(p)}\} \oplus \{\text{CK vectors } A^{(\pm)}\}$; CK has $\hat{\mathcal{F}}^{(\tau)}_{\mu\nu} = 2\bar{\nabla}_{[\mu}K^{(\tau)}_{\nu]}$. It never creates GW polarizations and is reported via T–L–O stamps. **Mition/Tantron** = BEH-massive spin-1 connection quanta living in the *connection* ledger; typed/TT safe; viable DM channels; not τ registers.

Chapter 11: The Graviton and Chronogeometric Gravity

Chapter 11 completes the quantisation of the typed, background–covariant spin-2 carrier on a certified plateau $\mathfrak{P}$ (with band tags BRST–closed and commuting) and reads all graviton observables on the physical space $\mathcal{H}_{\rm phys} = H^0(\hat{Q}_{\rm BRST})$. Using the York projector from Chapter 11, the graviton is *defined* as the transverse–traceless projection of the metric fluctuation (Definition 10.3 and eq. (10.4)),

$$\hat{h}_{\mu\nu} := \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\, \delta\hat{g}_{\mu\nu}, \qquad \bar{\nabla}^{\mu}\hat{h}_{\mu\nu} = 0, \quad \bar{g}^{\mu\nu}\hat{h}_{\mu\nu} = 0,$$

so tracelessness and transversality hold by construction. Temporal/coherence fluctuations $\hat{\tau}_{\mu\nu} = \delta\hat{t}_{\mu\nu}$ are TT–null in the weave, insulating the graviton channel from the τ–kairos layer; parasymmetric masks accordingly do not admit spin-2 TT excitations (Lemma 10.28 and Definition 10.4).

Typed dynamics are obtained by TT–sandwiching the spin-2 Lichnerowicz operator (plus sectoral mixing), yielding a sector–resolved TT wave operator $K^{[\mathrm{TT}],(a)}{}_{\mu\nu}{}^{\alpha\beta}$ and a strictly TT Feynman propagator $G^{\mathrm{F},[\mathrm{TT}],(a)}{}_{\underline{\mu\nu}}{}^{\rho\sigma}$. Its causal (Heaviside) representation uses the $\bar{\Sigma}$–order $\Theta_{\bar{\Sigma}}$, and equal–time limits via $\bar{\Sigma}$–covariant point–splitting reproduce the typed ETCR. On $\mathfrak{P}$, BRST–closedness and commutation of all selectors make these kernels gauge-parameter independent (Nielsen–flat) and regulator-safe; hyperbolicity of the principal symbol ensures retarded/advanced supports lie on $\bar{\Sigma}$–cones, giving typed microcausality.

The covariant algebra is then stated in terms of the TT Pauli–Jordan kernel $\Delta^{[\mathrm{TT}]} = G^{\mathrm{ret}}_{[\mathrm{TT}]} - G^{\mathrm{adv}}_{[\mathrm{TT}]}$ and the typed Peierls bracket $\{F, G\}_{\mathrm{P}}$ for gauge-invariant functionals $F[h], G[h]$; the equal–time limit recovers the canonical ETCR on any admissible $\bar{\Sigma}$–Cauchy slice. A $\bar{\Sigma}$–foliation $t : \mathcal{X}_q \to \mathbb{R}$ is used only to *present* equal–time relations; on the diagnostic realized reference configuration one proves the canonical↔Peierls equivalence, so predictions are foliation–independent.

Because the weave selectors commute, the graviton field resolves *triadically* by sector,

$$\hat{h}^{(a)}_{\mu\nu} = \Pi^{(a)}\,\hat{h}_{\mu\nu} = \Pi^{(a)}\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{g}_{\mu\nu}, \qquad a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\},$$

with $\sum_a \Pi^{(a)} = \mathbf{1}$ and $\Pi^{(a)}\Pi^{(b)} = \delta_{ab}\Pi^{(a)}$. In particular, the *Correlational Graviton* $\hat{h}^{(\mathrm{corr})} = \Pi^{(\mathrm{corr})}\hat{h}$ is the typed spin-2 projection of the interaction–sourced channel, while $\hat{\tau}$ remains TT–null. Quantum mechanically, the graviton is *composite* across the triad: its Källén–Lehmann representation on $\mathcal{H}_{\rm phys}$ carries a triadic residue vector $\vec{Z} = (Z_{\rm corr}, Z_{\rm vac}, Z_{\rm spec})$ and admits a corresponding triadic Fock factorisation, clarifying how geometric sources feed each sector.

Quantization of the spin–2 sector on the diagnostic realized reference configuration. The chapter quantizes the geometric (spin–2) sector on the realized, certified plateau $(\mathcal{X}_q, \bar{\Sigma}_{\mu\nu})$ using the Relational Levi–Civita connection $\bar{\nabla}$ that satisfies $\bar{\nabla}_\lambda\bar{\Sigma}_{\mu\nu} = 0$ (cf. Definition 6.7). The radiative carrier is fixed by the background–covariant York–type projector $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$, and the graviton field operator is defined intrinsically by

$$\hat{h}_{\mu\nu} \;:=\; \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{g}_{\mu\nu}, \qquad \bar{g}^{\mu\nu}\hat{h}_{\mu\nu} = 0, \quad \bar{\nabla}^{\mu}\hat{h}_{\mu\nu} = 0,$$

so tracelessness and transversality hold by construction. A key structural result is that the coherence/temporal fluctuation $\hat{\tau}_{\mu\nu} := \delta\hat{t}_{\mu\nu}$ is TT–null in the weave, hence spin–2 radiation is *purely metric* in all physical readouts (cf. Section 11.1.3).

Canonical and covariant quantizations; agreement on a plateau. Quantization is presented in two equivalent languages. (i) *Canonical:* impose equal–time commutators on the typed pair $(\hat{h}_{\mu\nu}, \hat{\Pi}^{\mu\nu})$ *after* band–sandwiching with $P_x(\lambda_{\rm UV})$ and projecting with $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$, preserving BRST hygiene and localizing the algebra within the $\bar{\Sigma}$–induced foliation. (ii) *Covariant:* use the Peierls bracket generated by the TT–projected Lichnerowicz operator so that the commutator is the (TT) difference of retarded/advanced Green operators supported on the $\bar{\Sigma}$–null cone. On a certified plateau $\mathfrak{P}$ the two definitions agree and yield the same realized propagator (cf. Section 11.1).

Typed Källén–Lehmann representation. The realized TT two–point function admits a triad–resolved Källén–Lehmann representation on $\mathcal{H}_{\rm phys}$,

$$\Delta^{(2),a}(q) \;=\; \frac{Z_a}{q^2+i\epsilon} \;+\; \int_0^\infty \frac{\rho_a^{(2)}(\mu^2)\,d\mu^2}{q^2-\mu^2+i\epsilon}, \qquad Z_a \geq 0, \quad \rho_a^{(2)} \geq 0,$$

with residue vector $\vec{Z} = (Z_{\rm corr}, Z_{\rm vac}, Z_{\rm spec})$ read on the plateau. This makes precise how the TT–typed graviton can carry a physical triadic admixture—only the component that survives BRST/plateau certification is realized (positivity as in Ch. 8).

Typed graviton–Tantrakyma vertex and sectoral structure. Typed interactions are built as shadows of the bi–local matter–geometry coupling ($\Phi\Phi \leftrightarrow \Theta\Theta$). The TT–typed, linearized graviton–Tantrakyma vertex is defined by

$$\hat{V}^{\mu\nu}(x) := \left(\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\right)^{\mu\nu}{}_{\alpha\beta}\, \frac{\delta S_{\rm int}\left[\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}\right]}{\delta \hat{h}_{\alpha\beta}(x)}\bigg|_{\mathfrak{P}}, \qquad \hat{V}^{\mu\nu} = \hat{V}^{\nu\mu}, \quad \bar{\Sigma}_{\mu\nu}\hat{V}^{\mu\nu} = 0,$$

and decomposes orthogonally into triadic blocks $\hat{V}^{(a)\mu\nu} := \Pi^{(a)}\hat{V}^{\mu\nu}$ with $[\Pi^{(a)}, \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}] = 0$ (cf. Definitions 11.24 and 11.64). Hermiticity, $\bar{\Sigma}$–cone support, and TT preservation follow from the weave algebra and from the TT–null τ leg, ensuring that spin–2 radiative content remains purely metric (cf. Section 11.1.3).

TT Ward identity and block–diagonal couplings. A typed Ward identity controls the vertex:

$$k_\mu\, \hat{V}^{[\mathrm{TT}],(a)\mu\nu}(k) = 0, \qquad \bar{\Sigma}_{\mu\nu}\, \hat{V}^{[\mathrm{TT}],(a)\mu\nu}(k) = 0, \quad \forall\, a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\},$$

with τ–insertions TT–null. Internal graviton lines in typed Feynman graphs are therefore TT propagators attached to TT vertices in the *same* triadic block; linearized couplings are block–diagonal in a (cf. Proposition 11.77 and Corollary 11.73).

Modal expansion in a typed TT polarization basis. For practical calculations, expand the vertex in a typed TT polarization basis $\{e_{\mu\nu}^{(s,a)}(k;x)\}$ (cf. Definition 11.18), defining modal amplitudes

$$\mathcal{V}^{(s,a)}(k;x) := \int d^d x'\, \mathcal{K}^{\mu\nu}(x,x')\, e_{\mu\nu}^{(s,a)}(k)\, e^{-ik\cdot x'},$$

so that

$$\hat{V}_{\mu\nu}^{[\mathrm{TT}],(a)}(x) = \sum_s \int \frac{d^d k}{(2\pi)^d}\, e_{\mu\nu}^{(s,a)}(k)\, \mathcal{V}^{(s,a)}(k;x) \;+\; \text{h.c.},$$

compatible with the sector–resolved TT mode expansion of $\hat{h}_{\mu\nu}^{(a)}$ and with the triadic propagators (cf. eqs. (11.86), (11.97) and (11.127) and Definition 11.51).

Mode decomposition from Gaiamiti fluctuations. In IQR, geometry is emergent: the graviton is the quantum excitation of the field whose expectation realizes $\bar{\Sigma}_{\mu\nu}$. Working on $\mathfrak{P}$ with $\hat{h}_{\mu\nu} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \delta\hat{g}_{\mu\nu}$, the coherence sector $\hat{\tau}_{\mu\nu}$ remains TT–null and does not contribute to spin–2 propagation (cf. Section 11.1.3).

Weave self–consistency: algebra, causality, and law diagnostics. The same $\bar{\Sigma}$ that governs causal propagation via the typed wave operator also fixes the quantum algebra via $\Delta^{[\mathrm{TT}]}$. The TT law residual $\mathbf{R}^{[\mathrm{TT}]}$ (cf. Definition 11.6) is naturally expressed through the Peierls construction; when the TT Ward identity holds on $\mathfrak{P}$, the residual vanishes and the law visibility saturates. Hence typed quantum algebra, causal support, and law diagnostics are mutually compatible on the diagnostic realized reference configuration; a typed Parasymmetric TLO layer is attached for the graviton to close the construction (cf. Proposition 11.14).

Chronogeometric Einstein equations (diagnostic realized reference configuration, operator, and linearised forms). On the diagnostic realized reference configuration $(\mathcal{X}_q, \bar{\Sigma}_{\mu\nu})$ with Relational Levi–Civita connection $\bar{\nabla}$ satisfying $\bar{\nabla}_\lambda \bar{\Sigma}_{\mu\nu} = 0$ (cf. Definition 6.7), the semi–classical law for expectation–level chronogeometry is stated in the Σ–language (cf. Definition 11.106 and eq. (11.191)):

$$\bar{G}_{\mu\nu}[\bar{\Sigma}] \;+\; \Lambda\, \bar{\Sigma}_{\mu\nu} \;=\; \kappa\, \bar{T}_{\mu\nu}^{\mathrm{total}}, \qquad \bar{G}_{\mu\nu} := \bar{R}_{\mu\nu} - \tfrac{1}{2}\, \bar{\Sigma}_{\mu\nu}\, \bar{R} \;\; (\textit{Definition 11.138 and eq. (11.260)}).$$

The diagnostic realized reference configuration law follows from the *relational Einstein–Hilbert action* on the same background (cf. Section 11.5.3, Definition 11.105, and eq. (11.188)) and is accompanied by covariant conservation identities

$$\bar{\nabla}^\mu \bar{G}_{\mu\nu} \equiv 0, \qquad \bar{\nabla}_\mu^{(\mathrm{rel})}\, \bar{T}_{\mathrm{total}}^{\mu\nu} = 0 \quad (\textit{eq. (11.192)}).$$

In coordinates adapted to $(\mathcal{X}_q, \bar{\Sigma})$, the unified Ricci tensor is obtained by contracting the $\bar{\Sigma}$–Riemann tensor written with the Relational Levi–Civita symbols, yielding the Christoffel form

$$\bar{R}_{\mu\nu} = \partial_\mu \Gamma^\lambda{}_{\nu\lambda} - \partial_\nu \Gamma^\lambda{}_{\mu\lambda} + \Gamma^\lambda{}_{\mu\alpha}\Gamma^\alpha{}_{\nu\lambda} - \Gamma^\lambda{}_{\nu\alpha}\Gamma^\alpha{}_{\mu\lambda},$$

and $\bar{R} = \bar{\Sigma}^{\mu\nu}\bar{R}_{\mu\nu}$ by the $\bar{\Sigma}$–trace. Physically, since $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$, the stationarity condition balances "metric curvature'' with "coherence/temporal curvature''; the limit $\bar{t}_{\mu\nu} \to 0$ reproduces the usual Einstein–Hilbert dynamics in g, while $\bar{R}[\bar{\Sigma}] - R[\bar{g}]$ encodes chronogeometric effects.

Unified operator–valued form. Beyond the diagnostic realized reference configuration equation, the chapter gives an operator–valued, retarded bitensor representation

$$\hat{G}_{\mu\nu}(x) = \int d\mu_{\bar{\Sigma}}(x')\, \hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x') \circ \Big[\mathcal{U}_{\bar{\Sigma}}(x \leftarrow x')_{\rho}{}^{\rho'}\, \mathcal{U}_{\bar{\Sigma}}(x \leftarrow x')_{\sigma}{}^{\sigma'}\, \hat{T}^{\rm total}_{\rho'\sigma'}(x')\Big] \quad (eq.\ (11.256)),$$

and a local symbolic form (densities/parallel transport suppressed)

$$\hat{G}_{\mu\nu}[\hat{\Sigma}](x) = \hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x) \circ \hat{T}^{\rm total}_{\rho\sigma}(x), \qquad A \circ B := \tfrac{1}{2}(AB+BA) \quad (eq.\ (11.257)),$$

with the total source split into diagonal, interaction, and coherence/torsion pieces,

$$\hat{T}^{\rm total}_{\rho\sigma} = \hat{T}^{(\Phi)}_{\rho\sigma} + \hat{T}^{(\Theta)}_{\rho\sigma} + \hat{T}^{(\Phi\Theta)}_{\rho\sigma} + \hat{T}^{(\tau)}_{\rho\sigma} \quad (eq.\ (11.258)).$$

Linearisation, typed spin–2 operator, and Σ–causal propagation. For radiative content, linearisation about $\bar{\Sigma}$ defines the *linearised Einstein operator*

$$\mathcal{E}_{\mu\nu}{}^{\rho\sigma}[\bar{\Sigma}]\, h_{\rho\sigma} := \frac{\delta}{\delta \bar{\Sigma}_{\rho\sigma}}\Big(\bar{G}_{\mu\nu}[\bar{\Sigma}] + \Lambda\, \bar{\Sigma}_{\mu\nu}\Big)\Big|_{\bar{\Sigma}} h_{\rho\sigma}, \qquad \bar{\nabla}^{\mu}(\mathcal{E}_{\mu\nu}{}^{\rho\sigma} h_{\rho\sigma}) = 0 \quad (eq.\ (11.181)).$$

York–projecting onto spin–2 yields the typed wave operator governing the graviton carrier,

$$K^{[\rm TT]}_{\mu\nu}{}^{\alpha\beta} := \Big[\Pi^{[\rm TT]}_{\bar{\Sigma}} \circ \mathcal{E}[\bar{\Sigma}] \circ \Pi^{[\rm TT]}_{\bar{\Sigma}}\Big]_{\mu\nu}{}^{\alpha\beta} \quad (eq.\ (11.182)),$$

which underpins the Σ–causal retarded/advanced Green functions and the Peierls algebra used elsewhere in the chapter (cf. Sections 11.5.4 and 11.7). Accordingly, TT kernels propagate on the $\bar{\Sigma}$ microcausal cone, ensuring consistency with the quantum bracket framework; diagnostic realized reference configuration–level diagnostics are packaged as a residual $\mathbf{R}^{[\rm Ein]}$ and visibility $\Lambda^{[\rm Ein]}_{\rm law}$, which saturate when the TT Ward identities and the Bianchi/relational conservation conditions hold.

Chapter 11 thus recasts Einstein's equations in a chronogeometric form keyed to $\bar{\Sigma}$: the definitions, conservation laws, causal support, and radiative (TT) content are made intrinsic to the unified tensor and its Levi–Civita geometry, while the operator–valued law, its linearised carrier, and diagnostic realized reference configuration instrumentation ensure all statements are read on the same BRST/Nielsen–clean, realized platform. The chapter provides a typed, background–covariant and BRST–consistent graviton construction: definition by York TT projection (Definition 10.3 and eq. (10.4)); sector–aware TT wave operators and propagators with causal support; a covariant (Peierls) algebra recovering ETCR on slices; and a triadic decomposition whose residues quantify correlational, vacuum and spectral content. All readings are made on $\mathfrak{P}$ and commute with the weave instrumentation, ensuring sector-clean, spin-clean and dialectically typed measurements consistent with the Chapter 11 realized weave configuration (Definitions 10.3 and 10.4 and Lemma 10.28).

BEH of the Chrono–Kymon: typed structure and consequences. *(1) TT graviton is blind to the chrono–kymon (no BEH "leakage" into spin–2).* The spin–2 carrier is the York–projected metric fluctuation

$$\hat{h}_{\mu\nu} \;=\; \Pi^{[\rm TT]}_{\bar{\Sigma}}\, \delta\hat{g}_{\mu\nu}, \qquad \bar{\nabla}^{\mu}\hat{h}_{\mu\nu} = 0, \quad \bar{g}^{\mu\nu}\hat{h}_{\mu\nu} = 0,$$

while the coherence/temporal fluctuation is $\hat{\tau}_{\mu\nu} := \delta\hat{t}_{\mu\nu}$. The weave algebra implies *TT–nullity of the coherence leg*

$$\left(\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\right)^{\mu\nu}{}_{\alpha\beta}\,\hat{\tau}^{\alpha\beta} = 0, \qquad \Rightarrow \qquad \langle \hat{h}, \hat{\tau} \rangle_{\mathrm{TT}} = 0,$$

so any would–be mixing that could shift the spin–2 pole is projected out (cf. Section 11.1.3). Consequently, the TT Dyson equation

$$\left[\Delta^{[\mathrm{TT}]}(k)\right]^{-1} = K^{[\mathrm{TT}]}(k) \; - \; \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\Sigma_{hh}(k)\,\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}, \qquad K^{[\mathrm{TT}]} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \circ \mathcal{E}[\bar{\Sigma}] \circ \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]},$$

has the τ-induced self–energy $\Sigma_{hh}^{(\tau)}$ *vanishing* in the TT block:

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\Sigma_{hh}^{(\tau)}(k)\,\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} = 0,$$

so the TT pole remains at $k^2 = 0$ with positive residue $Z_{\mathrm{grav}} > 0$. Equivalently, the Källén–Lehmann form in TT is unchanged (cf. eq. (11.182) and the typed KL discussion in Section 11.1).

(2) Ward identities keep spin–2 couplings purely metric and block–diagonal. For any typed triadic block $a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$, the TT vertex obeys

$$k_\mu\,\hat{V}^{[\mathrm{TT}],(a)\mu\nu}(k) = 0, \qquad \bar{\Sigma}_{\mu\nu}\,\hat{V}^{[\mathrm{TT}],(a)\mu\nu}(k) = 0,$$

and τ–insertions are *TT–null* within vertices and internal lines (cf. Proposition 11.77 and Corollary 11.73). Hence, typed Feynman graphs with internal gravitons use *TT propagators hooked to TT vertices in the same block*; no spin–1 (chrono–kymon) BEH structure can generate a mass term for $h_{\mu\nu}$.

(3) Where the BEH lives: the chrono–kymon's own spin–1 sector. The BEH mechanism acts in *spin–1 typed channels*, not in spin–2. Abstractly, in its sector the chrono–kymon quadratic kernel reads

$$\left[\mathcal{D}^{(\mathrm{CK})}(k)\right]^{-1}_{\alpha\beta} = \underbrace{\mathcal{K}^{(\mathrm{CK})}_{\alpha\beta}(k)}_{\text{kinetic, }\Sigma\text{-causal}} \; - \; \underbrace{\mathcal{M}^{2\,(\mathrm{CK})}_{\alpha\beta}}_{\text{BEH mass operator}},$$

with $\mathcal{M}^{2\,(\mathrm{CK})}$ generated by broken internal directions (the usual BEH pattern, now phrased on the $\bar{\Sigma}$ diagnostic realized reference configuration). Since $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ acts on *symmetric rank–2* tensors and is orthogonal to the chrono–kymon's *spin–1* subspace,

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\;\mathcal{M}^{2\,(\mathrm{CK})}\;\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} = 0,$$

by type/geometry, not by fine–tuning. Therefore, even if the chrono–kymon acquires a BEH mass $m_{\mathrm{CK}}^2 > 0$ in its sector, the graviton's typed KL representation

$$\Delta^{(2),a}(q) = \frac{Z_a}{q^2 + i\epsilon} + \int_0^\infty \frac{\rho_a^{(2)}(\mu^2)\,d\mu^2}{q^2 - \mu^2 + i\epsilon}, \qquad Z_a \geq 0,\; \rho_a^{(2)} \geq 0,$$

keeps its massless pole: no $\mathcal{O}(m_{\mathrm{CK}}^2)$ shift appears in TT (cf. Definition 11.51 and eq. (11.86)).

Bottom line. Chapter 11's quantization and TT Ward structure make the statement precise: the chrono–kymon may undergo BEH in its *own spin–1 typed sector*, but this does not induce, modify, or mix a BEH mass in the graviton. Mathematically, TT–nullity of τ and the typed Ward identities enforce

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\left\{\hat{\tau},\ \mathcal{M}^{2\,(\mathrm{CK})},\ \text{mixed } (h{-}\tau) \text{ kernels}\right\}\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}=0,$$

so the *spin–2 pole stays at* $k^2=0$ with Nielsen–safe residue, while any BEH physics of the chrono–kymon is *confined to spin–1 observables* in its own (typed) sector.

BEH of the Gaiaton (radial/trace mode in the Gaiamiti block). *(1) Radial mass from the typed Hessian; positivity on* H^0*.* The Gaiaton is the *radial/trace* (spin–0) excitation in the Gaiamiti ($\Phi\Phi$) manifold, orthogonal to the TT Goldstone sheet. On a certified plateau $\mathfrak{P}$, its pole is fixed by the *typed radial mass operator* extracted from the band–prepared Hessian of the Gaiamiti potential (global setting):

$$m_A^2 \;=\; \lambda_{\mathrm{rad}}\Big(\mathsf{M}_{\Phi}^{[\mathrm{rad}]}\Big) \;>\; 0, \qquad \Delta_A(k) \;=\; \frac{1}{k^2-m_A^2+i0^+} \;\text{ on }\; \mathcal{H}_{\mathrm{phys}}, \quad \text{(cf. Definition 9.250)}$$

with KL–positivity on H^0 (cf. eq. (7.12)). This realizes a single scalar pole $k^2=m_A^2$ (no ghosts/tachyons) as summarized in Section 9.17.4.

(2) Gauged case: BEH for internal vectors and a gauge–independent Gaiaton pole. When Φ carries an internal gauge action,

$$D_\mu\,\Phi=\partial_\mu\Phi-ig\,A_\mu\cdot\Phi,$$

the SSB background $\langle\Phi\rangle$ generates a *vector* BEH mass matrix for the broken generators, while the longitudinal would–be Goldstones form BRST quartets and decouple; unbroken generators remain massless (cf. Section 9.17.4 and Proposition 9.257). In this *gauged* setting the Gaiaton mass is obtained from the *Apeiron* (gauged–radial) operator,

$$m^2_{A,\mathrm{gauged}} \;=\; \lambda_{\mathrm{rad}}\Big(\mathsf{M}^{[\mathrm{rad}]}_{\Phi,\mathrm{Apeiron}}\Big), \qquad \text{(cf. Definition 9.259)}$$

and both the vector masses and the Gaiaton pole are Nielsen–safe on $\mathfrak{P}$:

$$\frac{\partial\, m^2_{A,\mathrm{gauged}}}{\partial\xi}=0, \qquad \frac{\partial\, M_V^2}{\partial\xi}=0 \quad \text{(cf. Theorem 9.351)}.$$

Thus BEH reorganizes the *spin–1* spectrum (massive broken, massless unbroken) without introducing gauge–parameter dependence into the scalar Gaiaton pole.

(3) Orthogonality to spin–2: no TT mass term from the Gaiaton. Because the TT projector annihilates pure traces,

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\Big[S(x)\,\bar{\Sigma}_{\mu\nu}\Big]=0 \qquad \forall\, S,$$

any mixing of the radial/trace scalar with the TT graviton is projected out. Equivalently, at the level of self–energies and vertices,

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\Sigma_{hh}^{(\mathrm{Tr})}(k)\,\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}=0, \qquad k_\mu\,\hat{V}^{[\mathrm{TT}]\mu\nu}=0, \quad \bar{\Sigma}_{\mu\nu}\hat{V}^{[\mathrm{TT}]\mu\nu}=0,$$

so the York–projected graviton carrier remains massless and purely metric (cf. eq. (11.182), Proposition 11.77, and Corollary 11.73).

Bottom line. The Gaiaton is a *gapped* scalar set by the typed radial curvature of the Gaiamiti potential,

$$\Delta_A(k) = \left(k^2 - m_A^2 + i0^+\right)^{-1},$$

with m_A^2 read either from $\mathsf{M}_\Phi^{[\mathrm{rad}]}$ (global) or from its gauged Apeiron counterpart, both *Nielsen–safe* on $\mathfrak{P}$ (cf. Definitions 9.250 and 9.259 and Theorem 9.351). Internal gauging triggers BEH *for vectors*, not for the spin–2 channel; the TT graviton remains massless by projection/orthogonality, while the Gaiaton's scalar pole stays gauge–independent and positive on H^0, completing the realized Gaiamiti spectrum (cf. Section 9.17.4 and Proposition 9.257).

Part IV: The Standard Model

Chapter 12: The Standard Model

Chapter 12 consolidates the Standard Model within the holosymmetric programme by showing how $G_{\mathrm{SM}} = \mathrm{SU}(3)_C \times \mathrm{SU}(2)_L \times \mathrm{U}(1)_Y$ is singled out by the Holosymmetric Selection Principle (HSP): Stage I is a self-consistency sieve (unitarity/BRST viability), and Stage II is a comparative interferometric tomography that enforces maximal relational coherence and orthogonality to alternatives [52, 117, 112] (Section 12.2). In this setting, gauge bosons are the BRST-physical excitations of the *internal* gauge connection $A_\mu^{\ I}$ on the Tantrakyma bundle, with Yang–Mills dynamics carried by the curvatures $F_{\mu\nu}^{\ I}$ (Definition 12.95; cf. [123, 38]); in parallel, the Operator Bundle Connection $\mathcal{A}_\mu$ has curvature $\mathcal{F}_{\mu\nu} = \partial_\mu \mathcal{A}_\nu - \partial_\nu \mathcal{A}_\mu + i[\mathcal{A}_\mu, \mathcal{A}_\nu]$ and enters the realized weave configuration-compatible transport and curvature split [36, 37] (Definitions 6.192 and 12.96). Fermionic kymons appear as operator-valued spinor sections whose one-generation content matches the familiar $(3,2)_{1/6}$, $(1,2)_{-1/2}$, $(3,1)_{2/3}$, $(3,1)_{-1/3}$, $(1,1)_{-1}$, $(1,1)_0$ assignments [38] (Section 12.4.1; [124]), with subscripts denoting $Y/2$ consistent with the GMN convention used in the chapter.

Electroweak symmetry breaking is treated as a vacuum alignment in the Tantrakyma sector, and the Gell–Mann–Nishijima relation $Q = I_3 + \frac{1}{2}Y$ is *derived*—not postulated—as the unique unbroken generator that leaves the vacuum invariant [38] (Section 12.4.3 and eq. (12.107); cf. [125, 126]). Flavor then enters as geometry: Yukawa structure and diagonalisation produce CKM/PMNS mixing, while the global "family frame" turns these into parallel-transport holonomies $\mathcal{P}\exp\int \mathcal{A}^{(F)}$ [38, 127] (Section 12.5.5 and eq. (12.134)). Confinement is recast as gauge-induced spectral filtering: the color-singlet projector P_{singlet} realises the hadronic sector as a gauge-invariant image inside the emergent Hilbert space, and BRST cohomology identifies the physical space, tying unitarity to the same projector architecture [38, 117] (Proposition 12.128, Section 12.3, and eq. (12.77); cf. [128]). These ingredients are then lifted back into HSP: anomaly freedom and BRST nilpotency supply the Stage I viability condition, while the interferometric coherence functional on the physical sector implements Stage II selection, yielding the SM as the unique holosymmetric fixed point among admissible candidates [117, 110, 111] (eq. (12.46), Theorem 12.57, and Section 12.2).

Beyond this core, the chapter develops two operational extensions germane to phenomenology. First, the neutrino sector accommodates Majorana options and a seesaw hierarchy within the Tantrakyma formalism, consistent with the flavor geometry above [129, 130, 131, 127] (Section 12.7.1). Second, "dark sectors" are characterised not ontologically but operationally: a state is dark relative to G_{SM} if it registers under a holosymmetric coherence projector for some hidden G while remaining spectrally orthogonal to all SM observables [112] (*see Definition 12.77*), providing a concrete tomography-based criterion for hidden dynamics (Section 12.5.4). A brief side thread outlines a state-dependent, unitary evolution across black-hole formation/evaporation within the holocosmic background, situating the construction amidst modern information-theoretic perspectives [22]. In sum, Chapter 12 does not append the SM to the framework; it *carves* it from the operator-geometric substrate via holosymmetric interferometry, with each textbook structure re-expressed as a consequence of relational selection and cohomological admissibility [52, 117, 112].

Realized weave configuration-compatible transport and the operator–bundle picture

The basic kinematic datum is a covariant "split" derivative that respects both geometry and the operator structure of fields on the certified plateau (Section 12.4.4). The realized weave configuration-compatible connection is

$$\nabla_\mu \;=\; \nabla_\mu^{(\mathrm{LC})} \;+\; i\,\mathrm{ad}_{\mathcal{A}_\mu} \quad \text{(Definitions 6.192 and 12.1)},$$

where $\nabla_\mu^{(\mathrm{LC})}$ is the Levi–Civita part and $\mathrm{ad}_{\mathcal{A}_\mu}(X) := [\mathcal{A}_\mu, X]$ is the inner derivation generated by the universal operator–bundle connection $\mathcal{A}_\mu \in \mathrm{End}(\mathcal{H}_{\mathrm{em}})$. With this split, all symmetry statements are tied to ∇_μ: the Euler–Lagrange operator for $\hat{\Theta}$ transforms covariantly under any admissible internal $G \subset G_{\mathrm{int}}$ on a certified plateau (definition "realized weave configuration-compatible G-symmetry", eqs. eq. (12.37)), so that

$$\mathrm{EL}[\hat{\Theta}] \;\mapsto\; U\,\mathrm{EL}[\hat{\Theta}]\,U^{-1}, \qquad U \in G \text{ (DRRC-admissible)}.$$

Here "DRRC-admissible" means admissible with respect to the diagnostic realized reference configuration of Definition 9.185. Intuitively: the Levi–Civita part transports spacetime indices; the $\mathrm{ad}_{\mathcal{A}_\mu}$ part transports operator content in $\mathrm{End}(\mathcal{H}_{\mathrm{em}})$; and the internal G-action is threaded in a way that keeps the total dynamics covariant on $\mathfrak{P}$.

For fermionic carriers living on

$$\mathrm{End}(\mathcal{H}_{\mathrm{em}}) \otimes \mathcal{S} \otimes E_n,$$

the total (realized weave configuration-compatible) split derivative is

$$\mathcal{D}_\mu \hat{\psi}_n \;=\; \nabla_\mu^{(\mathcal{S})} \hat{\psi}_n \;+\; i\,[\mathcal{A}_\mu, \hat{\psi}_n] \;+\; i\,g\,A_\mu^{\ I}\,\rho_n(T_I)\,\hat{\psi}_n,$$

with $\nabla_\mu^{(\mathcal{S})}$ the spin connection on $\mathcal{S}$, $A_\mu^{\ I} T_I$ the internal gauge field (see Definition 12.95), and ρ_n the representation on E_n. Its curvature splits cleanly into Lorentz, operator-bundle, and internal pieces (Lemma Lemma 12.87):

$$[\mathcal{D}_\mu, \mathcal{D}_\nu]\hat{\psi}_n \;=\; R_{\mu\nu}^{(\mathcal{S})} \cdot \hat{\psi}_n \;+\; i\,[\mathcal{F}_{\mu\nu}, \hat{\psi}_n] \;+\; i\,g\,F_{\mu\nu}^{\ I}\,\rho_n(T_I)\,\hat{\psi}_n,$$

where

$$\mathcal{F}_{\mu\nu} = \partial_\mu \mathcal{A}_\nu - \partial_\nu \mathcal{A}_\mu + i\,[\mathcal{A}_\mu, \mathcal{A}_\nu] \qquad \text{(Definition 12.96)},$$
$$F_{\mu\nu}^{\ I} = \partial_\mu A_\nu^{\ I} - \partial_\nu A_\mu^{\ I} + f^I{}_{JK}\, A_\mu^{\ J} A_\nu^{\ K}.$$

This decomposition powers the "hygiene" checks used later: each sector's curvature is measured and controlled separately while remaining compatible with the total transport $\mathcal{D}_\mu$ (Lemma 12.87).

On the diagnostic realized reference configuration $\mathfrak{P}$, the internal gauge kinetics are governed by the Yang–Mills action (Definition 12.98, eq. eq. (12.113)):

$$S_{\mathrm{YM}}[A] \;=\; -\frac{1}{2}\int_{\mathfrak{P}} \sqrt{|g|}\, d^d x\; \kappa_{IJ}\, F_{\mu\nu}^{\ I}\, F^{\mu\nu\, J},$$

with κ_{IJ} the invariant Killing metric on the internal Lie algebra. In flat coordinates this reduces to the standard form, and in $d = 4$ it projects onto the familiar

$$\mathrm{SU}(3)_C \oplus \mathrm{SU}(2)_L \oplus \mathrm{U}(1)_Y \qquad \text{(Lemma Lemma 12.99)}.$$

Crucially, the operator–bundle sector $\mathcal{A}_\mu$ and the internal gauge sector $A_\mu^{\ I}$ remain distinguished but consistently coupled through $\mathcal{D}_\mu$; this is what lets the chapter enforce covariance for composite operator observables built from fields in $\mathrm{End}(\mathcal{H}_{\mathrm{em}})$ (see also the unified split in Section 12.4.4).

Finally, meters are introduced operator-geometrically and are only deployed on certified plateaux so that readouts are BRST/Nielsen stable (Definition Definition 12.16). In practice this means: for any typed meter $\mathcal{M}_\tau$,

$$\big\langle\!\big\langle \mathcal{M}_\tau \big\rangle\!\big\rangle_{\mathrm{DRRC}} \text{ depends only on the BRST cohomology class,} \qquad \frac{\partial}{\partial \xi}\big\langle\!\big\langle \mathcal{M}_\tau \big\rangle\!\big\rangle_{\mathrm{DRRC}} = 0,$$

where $\big\langle\!\big\langle\, \cdot\, \big\rangle\!\big\rangle_{\mathrm{DRRC}}$ denotes evaluation on the diagnostic realized reference configuration of Definition 9.185, ξ is a gauge/Nielsen parameter, and $\hat{Q}_{\mathrm{BRST}}$ is the nilpotent charge selecting the physical space $\mathcal{H}_{\mathrm{phys}}$ (BRST background in Section 12.3). Because transport is realized weave configuration-compatible and the meters are defined in the operator–bundle language, the resulting pole/residue extractions and charge readouts commute with all admissible $G \subset G_{\mathrm{int}}$ on $\mathfrak{P}$, ensuring that symmetry statements proved with ∇_μ are faithfully realized by what the meters actually see (typed-witness context in Definition 9.187).

Anomaly meters, SM calibration, and minimality

Chapter 12 assembles the mixed and cubic anomaly constraints into the "anomaly block" (eq. (12.46)) and evaluates them with diagnostic realized reference configuration meters on the certified plateau $\mathfrak{P}$. For a chiral spectrum $\{f\}$ with hypercharge Y_f and non-abelian

representations R_f, the six local anomaly meters read

$$\begin{aligned}
\sum_f Y_f\, T_2^{\mathrm{SU}(3)}(R_f) &= 0 && [\,\mathrm{SU}(3)_C^2\!-\!\mathrm{U}(1)_Y\,],\\
\sum_f Y_f\, T_2^{\mathrm{SU}(2)}(R_f) &= 0 && [\,\mathrm{SU}(2)_L^2\!-\!\mathrm{U}(1)_Y\,],\\
\sum_f Y_f^{\,3} &= 0 && [\,\mathrm{U}(1)_Y^3\,],\\
\sum_f Y_f &= 0 && [\,\mathrm{grav}^2\!-\!\mathrm{U}(1)_Y\,],\\
\sum_f A^{\mathrm{SU}(3)}(R_f) &= 0 && [\,\mathrm{SU}(3)_C^3\,],\\
\sum_f A^{\mathrm{SU}(2)}(R_f) &= 0 && [\,\mathrm{SU}(2)_L^3\ (\mathrm{local})],
\end{aligned}$$

together with the global Witten test for SU(2):

$$\#\{\mathrm{SU}(2)\text{ doublets (counting color)}\} \equiv 0 \pmod 2.$$

Here $T_2(R)$ is the Dynkin index (with $T_2(\mathbf{2}) = T_2(\mathbf{3}) = \frac{1}{2}$ in the standard normalization) and $A(R)$ is the cubic anomaly coefficient (with $A(\mathbf{3}) = -A(\overline{\mathbf{3}}) = 1$ for SU(3)).

For the one-generation Standard-Model assignment listed in table 12.1, each meter vanishes explicitly. Writing all fields as left-handed Weyl spinors,

$$Q_L : (\mathbf{3},\mathbf{2},+\tfrac{1}{6}),\quad u_R^c : (\overline{\mathbf{3}},\mathbf{1},-\tfrac{2}{3}),\quad d_R^c : (\overline{\mathbf{3}},\mathbf{1},+\tfrac{1}{3}),\quad L_L : (\mathbf{1},\mathbf{2},-\tfrac{1}{2}),\quad e_R^c : (\mathbf{1},\mathbf{1},+1),$$

(with an optional $\nu_R^c : (\mathbf{1},\mathbf{1},0)$ which is anomaly-inert), the per-generation cancellations are

$$\begin{aligned}
\mathrm{SU}(3)_C^2\!-\!\mathrm{U}(1)_Y:&\quad 2\cdot\tfrac{1}{6}\,T_2(\mathbf{3}) + (-\tfrac{2}{3})\,T_2(\overline{\mathbf{3}}) + (\tfrac{1}{3})\,T_2(\overline{\mathbf{3}}) = \tfrac{1}{6}-\tfrac{1}{3}+\tfrac{1}{6} = 0,\\
\mathrm{SU}(2)_L^2\!-\!\mathrm{U}(1)_Y:&\quad 3\cdot(\tfrac{1}{6})\,T_2(\mathbf{2}) + 1\cdot(-\tfrac{1}{2})\,T_2(\mathbf{2}) = \tfrac{1}{4}-\tfrac{1}{4} = 0,\\
\mathrm{U}(1)_Y^3:&\quad 6\big(\tfrac{1}{6}\big)^3 + 3\big(-\tfrac{2}{3}\big)^3 + 3\big(\tfrac{1}{3}\big)^3 + 2\big(-\tfrac{1}{2}\big)^3 + (1)^3 = 0,\\
\mathrm{grav}^2\!-\!\mathrm{U}(1)_Y:&\quad 6\big(\tfrac{1}{6}\big) + 3\big(-\tfrac{2}{3}\big) + 3\big(\tfrac{1}{3}\big) + 2\big(-\tfrac{1}{2}\big) + (1) = 0,\\
\mathrm{SU}(3)_C^3:&\quad 2\,A(\mathbf{3}) - A(\overline{\mathbf{3}}) - A(\overline{\mathbf{3}}) = 2-1-1 = 0,\\
\text{Witten test}:&\quad 3\times(\text{quark doublet}) + 1\times(\text{lepton doublet}) = 4 \equiv 0 \pmod 2.
\end{aligned}$$

These equalities constitute the "Standard-Model calibration fixture" (Proposition 12.26): all six local anomaly meters read

$$\big\langle\!\big\langle \mathcal{A}_i \big\rangle\!\big\rangle_{\mathfrak{P}} = 0, \qquad i = 1,\dots,6,$$

validating the normalization/tagging in table 12.1 and ensuring Nielsen-flatness of certified readouts on $\mathfrak{P}$.

Under these constraints—together with "no exotics" in the chiral spectrum and the presence of a single nontrivial abelian factor—the chapter states the minimality/uniqueness of the internal gauge law:

$$G_{\mathrm{SM}} = \mathrm{SU}(3)_C \times \mathrm{SU}(2)_L \times \mathrm{U}(1)_Y,$$

namely, it is the unique minimal chiral gauge sector that solves the anomaly block (paragraph "Minimality and uniqueness").

Representation-theoretic sufficiency bridge to G_{SM}

The central statement (Theorem 12.57) asserts: if a candidate internal law A is (i) compact, local, and Tantrakyma-invariant; (ii) chirally admissible with the one-generation SM matter content; (iii) completely anomaly-free (local + global); and (iv) minimal (no extra chiral matter; exactly one nontrivial U(1)), then the induced K-action on $\mathcal{H}_{\mathrm{mat}}$ is unitarily equivalent to the SM action. Equivalently, there exists a unitary intertwiner $U : \mathcal{H}_{\mathrm{mat}} \to \mathcal{H}_{\mathrm{mat}}$ such that

$$\rho_A(k) \;=\; U\,\rho_{\mathrm{SM}}(k)\,U^{-1} \quad \forall\, k \in K, \qquad d\rho_A(X) \;=\; U\,d\rho_{\mathrm{SM}}(X)\,U^{-1} \quad \forall\, X \in \mathfrak{k},$$

with $\mathfrak{k} = \mathrm{Lie}(K)$.

Proof highlights (structural constraints). *(1) Weak factor from pseudoreal doublets.* Chiral admissibility with SM matter includes left-handed weak doublets. In a compact setting, a nontrivial two-dimensional *pseudoreal* irrep forces an $\mathfrak{su}(2)$ simple summand at minimal rank. Thus

$$\mathfrak{k} \;\supset\; \mathfrak{su}(2) \qquad \Rightarrow \qquad K \;\supset\; \mathrm{SU}(2)_L.$$

(2) Color factor from complex triplets with nonvanishing cubic index. The presence of color triplets and anti-triplets with a nonzero cubic anomaly coefficient $A(R)$ requires a simple factor whose fundamentals are complex and support a nonvanishing third-order invariant. Among compact simples, the minimal such choice is $\mathfrak{su}(3)$, hence

$$\mathfrak{k} \;\supset\; \mathfrak{su}(3) \qquad \Rightarrow \qquad K \;\supset\; \mathrm{SU}(3)_C.$$

(3) Abelian factor fixed by fractional charges and GMN. Diagnostic realized reference configuration charges obey the Gell–Mann–Nishijima relation

$$Q \;=\; I_3 \,+\, \tfrac{1}{2}\,Y,$$

and the observed fractional Q for quarks requires a hypercharge generator Y that is independent of the $\mathfrak{su}(2)$ Cartan. Minimality (exactly one nontrivial abelian factor) hence fixes

$$\mathfrak{k} \;\supset\; \mathfrak{u}(1)_Y \qquad \text{and no extra } \mathfrak{u}(1) \text{ factors.}$$

Additional abelian factors would either leave uncanceled mixed anomalies or demand exotic chiral matter, violating (iii) or (iv).

Putting (1)–(3) together and invoking complete anomaly freedom yields the Lie-algebra isomorphism

$$\mathfrak{k} \;\cong\; \mathfrak{su}(3) \oplus \mathfrak{su}(2) \oplus \mathfrak{u}(1),$$

hence (up to global issues and overall normalization of Y)

$$K \;\simeq\; \mathrm{SU}(3)_C \times \mathrm{SU}(2)_L \times \mathrm{U}(1)_Y.$$

Relational character and unitary equivalence. The "relational character" (Definition 12.54) packages the K-action on $\mathcal{H}_{\mathrm{mat}}$ as a class function

$$\chi_A(k) \;:=\; \mathrm{Tr}_{\mathcal{H}_{\mathrm{mat}}}\Big(\rho_A(k)\Big), \qquad k \in K,$$

which decomposes into irreducible characters as

$$\chi_A \;=\; \sum_{\lambda \in \widehat{K}} m_\lambda\, \chi^\lambda, \qquad m_\lambda \in \mathbb{N}.$$

Peter–Weyl orthogonality gives the "filters" used later in Stage II:

$$\langle \chi_A, \chi^\lambda \rangle_K \;:=\; \int_K \overline{\chi^\lambda(k)}\, \chi_A(k)\, d\mu_K(k) \;=\; m_\lambda,$$

so matching the relational character to the SM data fixes the multiplicities m_λ and, hence, the isomorphism class of the representation. Under (i)–(iv), the spectrum and multiplicities agree with the SM one-generation assignment, which implies the existence of the unitary intertwiner U above. This completes the bridge claimed in Theorem 12.57.

EWSB, Yukawas, and masses on the diagnostic realized reference configuration

Expanding the electroweak scalar about the vacuum on the certified plateau as

$$\bar{\Theta} \;=\; \langle H \rangle \;=\; \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v \end{pmatrix} \qquad \text{(Lemma 12.94)[38]},$$

and writing the Yukawa sector in the standard doublet notation,

$$\mathcal{L}_Y \;=\; -\,\bar{Q}_L\, \tilde{H}\, Y_u\, u_R - \bar{Q}_L\, H\, Y_d\, d_R - \bar{L}_L\, H\, Y_e\, e_R - \bar{L}_L\, \tilde{H}\, Y_\nu\, \nu_R + \mathrm{h.c.}, \qquad \tilde{H} := i\sigma^2 H^* \text{[38]},$$

one obtains the fermion mass Lagrangian after EWSB,

$$\mathcal{L}_{\mathrm{mass}}^{(\mathrm{ferm})} = -\,\bar{d}_L\, M_d\, d_R - \bar{u}_L\, M_u\, u_R - \bar{e}_L\, M_e\, e_R - \bar{\nu}_L\, M_D\, \nu_R + \mathrm{h.c.},$$

$$M_f = \frac{v}{\sqrt{2}}\, Y_f, \quad \text{for } f \in \{u, d, e\},$$

$$M_D = \frac{v}{\sqrt{2}}\, Y_\nu,$$

with the standard relations read on the diagnostic realized reference configuration [38, 112]. Each Yukawa matrix admits a singular-value (biunitary) decomposition $Y_f = U_{f_L}^\dagger\, y_f^{\mathrm{diag}}\, U_{f_R}$, whence

$$M_f \;=\; \frac{v}{\sqrt{2}}\, U_{f_L}^\dagger\, y_f^{\mathrm{diag}}\, U_{f_R}, \qquad m_f^{(i)} \;=\; \frac{v}{\sqrt{2}} \Big(\text{sing. values}(Y_f)\Big)^{(i)} \text{[38]}.$$

Diagnostic realized reference configuration-admissible unitary rotations bring the kinetic terms to canonical form and diagonalize the masses; charged-current mixing then appears as

$$V_{\mathrm{CKM}} \;=\; U_{u_L}\, U_{d_L}^\dagger,$$

$$U_{\mathrm{PMNS}} \;=\; U_{\ell_L}\, U_{\nu_L}^\dagger$$

(Section 12.5.5, Proposition 12.115, and eq. (12.134))[38, 127].

Readouts are performed in the scalar τ-channel, preserving TT purity (typed-meter and witness framework Definitions 9.187 and 12.16).

Transport is consistently handled by the realized weave configuration-compatible split derivative $\mathcal{D}_\mu$; the Yukawa tensors Y_f behave as internal scalars on the diagnostic realized reference configuration and thus preserve meter hygiene and covariance under the operator–bundle structure [36, 37] (realized weave configuration-compatibility Definition 6.192, Lemma 12.87, and Section 12.4.4). In particular, the scalar excitation $h(x)$ around $\bar{\Theta}$ couples proportionally to mass,

$$\mathcal{L}_{h\bar{f}f} \;=\; -\frac{h}{v}\left(\bar{d}_L M_d d_R + \bar{u}_L M_u u_R + \bar{e}_L M_e e_R + \bar{\nu}_L M_D \nu_R\right) + \text{h.c.},$$

which is aligned with the Yukawa tensors (diagnostic realized reference configuration alignment in Section 12.5.5) [38].

Compact summary. Before EWSB, gauge invariance forbids fermion mass terms. After EWSB, masses arise as the singular values of the Yukawas:

$$m_f \;=\; \frac{v}{\sqrt{2}}\,\text{sing. values}(Y_f),$$

and the Higgs–fermion couplings are aligned with Y_f, i.e. proportional to M_f/v in the mass basis (Section 12.5.5) [38, 112].

Gauge bosons and the GMN relation

On the certified plateau $\mathfrak{P}$, the physical gauge bosons are the BRST-physical *transverse* poles of the internal connection (Definition Definition 12.95) [52, 117, 38]: eight gluons g^a_μ $(a = 1, \ldots, 8)$ from $\mathrm{SU}(3)_C$, together with the electroweak primordial fields W^i_μ $(i = 1, 2, 3)$ and B_μ. After spontaneous symmetry breaking (SSB), the charged sector is formed by

$$W^\pm_\mu \;=\; \frac{1}{\sqrt{2}}\left(W^1_\mu \mp i\, W^2_\mu\right) \qquad \text{(Definition 12.95)[38]},$$

while the neutral sector mixes as

$$\begin{pmatrix} Z_\mu \\ A_\mu \end{pmatrix} \;=\; \begin{pmatrix} \cos\theta_W & -\sin\theta_W \\ \sin\theta_W & \cos\theta_W \end{pmatrix} \begin{pmatrix} W^3_\mu \\ B_\mu \end{pmatrix}, \qquad \tan\theta_W \;=\; \frac{g'}{g}\,[38],$$

so that A_μ is the massless eigenvector that couples to the unbroken generator $\hat{Q}$ determined in Section 12.4.3 (see also eq. (12.107)) [125, 126, 38].

The neutral mass matrix descending from the BEH sector (Higgs kinetic term) has the standard form

$$\mathcal{L}^{\text{neutral}}_{\text{mass}} \;=\; \frac{v^2}{8}\begin{pmatrix} W^3_\mu & B_\mu \end{pmatrix} \begin{pmatrix} g^2 & -g\,g' \\ -g\,g' & g'^2 \end{pmatrix} \begin{pmatrix} W^{3\mu} \\ B^\mu \end{pmatrix},$$

which is diagonalized by the rotation above, yielding

$$\begin{aligned} M_W &= \frac{g\,v}{2}, \\ M_Z &= \frac{v}{2}\sqrt{g^2+g'^2}, \\ M_\gamma &= 0 \end{aligned} \qquad \text{(DRRC EWSB context, Lemma 12.94)[38, 112].}$$

Here "DRRC" denotes the diagnostic realized reference configuration of Definition 9.185. The covariant derivative on a field with weak isospin generators T^a and hypercharge Y reads

$$\begin{aligned} D_\mu &= \partial_\mu + i\,g\,T^a W_\mu^a + i\,\frac{g'}{2}\,Y\,B_\mu \\ &= \partial_\mu + i\,\frac{g}{\sqrt{2}}\Big(T^+ W_\mu^+ + T^- W_\mu^-\Big) + i\,e\,Q\,A_\mu \\ &\quad + i\,\frac{g}{\cos\theta_W}\Big(T^3 - \sin^2\theta_W\,Q\Big)\,Z_\mu, \end{aligned}$$

with $T^\pm := T^1 \pm iT^2$, $e := g\sin\theta_W = g'\cos\theta_W$, and

$$Q \;=\; I_3 + \frac{Y}{2} \qquad \text{(Gell–Mann–Nishijima/GMN relation, eq. (12.107))[125, 126, 38].}$$

Thus, for all realized species, the fermionic charges satisfy $Q = I_3 + \frac{1}{2}Y$, exactly matching the neutral/charged mixing enforced by the BEH matrix, and the massless eigenvector A_μ couples to the unbroken electromagnetic generator $\hat{Q}$ on $\mathfrak{P}$ [38, 112] (unbroken generator from Lemma 12.94; emergence in Section 12.4.3).

Neutrino sector: seesaw and $U(1)_L$ breaking

Chapter 12 introduces a minimal lepton-number–violating extension with a gauge-singlet scalar Φ_ν whose vacuum expectation value (VEV) spontaneously breaks the global symmetry $U(1)_L$ and generates a Majorana mass for right-handed neutrinos [129, 130, 131, 127] (Section 12.7.1 and Proposition 12.148). The relevant Yukawa and Majorana couplings on the certified plateau $\mathfrak{P}$ are [38, 127]

$$\mathcal{L} \;\supset\; -\,\bar{L}_L\,\tilde{H}\,Y_\nu\,\nu_R \;-\; \tfrac{1}{2}\,y^{(\nu)}\,\Phi_\nu\,\overline{\nu_R^c}\nu_R \;+\; \text{h.c.}, \qquad \tilde{H} := i\sigma^2 H^*,$$

where Y_ν and $y^{(\nu)}$ act on the family fibre and commute with diagnostic realized reference configuration transport (operator/bundle covariance; Definitions 6.192 and 12.106 and Section 12.4.4). When [38, 131]

$$\langle H\rangle \;=\; \frac{1}{\sqrt{2}}\begin{pmatrix}0\\ v\end{pmatrix}, \qquad \langle\Phi_\nu\rangle \;=\; v_\nu,$$

one obtains the Dirac and Majorana blocks

$$M_D \;=\; \frac{v}{\sqrt{2}}\,Y_\nu, \qquad M_R \;=\; y^{(\nu)}\,v_\nu,$$

consistent with the DRCC vacuum alignment (Lemma 12.94) and the lepton-number–violating trigger (Proposition 12.148). In the $(\nu_L,\, \nu_R^c)$ basis the mass Lagrangian is

$$\mathcal{L}_{\text{mass}} \;=\; -\,\frac{1}{2}\,\begin{pmatrix}\nu_L^{\top} & \nu_R^{\top}\end{pmatrix} C^{-1}\begin{pmatrix}0 & M_D\\ M_D^{\top} & M_R\end{pmatrix}\begin{pmatrix}\nu_L\\ \nu_R^{c}\end{pmatrix} \;+\; \text{h.c.},$$

which is the Type-I seesaw with $m_L = 0$ [129, 130, 127] (diagnostic realized reference configuration formulation in Section 12.7.1). For a large Majorana scale $\|M_R\| \gg \|M_D\|$, block-diagonalization yields the effective light-neutrino mass matrix

$$m_\nu^{\text{eff}} \;=\; -\,M_D\,M_R^{-1}\,M_D^{\top} \;+\; \mathcal{O}(M_R^{-2}) \qquad \text{(eq. (12.209))[130, 127, 112].}$$

Parametrically, in any consistent operator norm,

$$\|m_\nu^{\text{eff}}\| \;\sim\; \frac{\left\|v\,Y_\nu\right\|^2}{\left\|y^{(\nu)}\right\|\,v_\nu},$$

matching the scaling quoted in eq. (12.209) [112]. This reproduces the Weinberg operator at low energies,

$$\mathcal{L}_{\text{eff}} \;=\; -\,\frac{1}{2}\,(\bar{L}_L\tilde{H})\,C^{-1}\,c_5\,(\tilde{H}^{\top}L_L^{c}) \;+\; \text{h.c.}, \qquad c_5 \;=\; Y_\nu\,M_R^{-1}\,Y_\nu^{\top},$$

so that after EWSB $m_\nu^{\text{eff}} = \frac{v^2}{2}\,c_5$, demonstrating consistency with the dimension-five Weinberg limit (Proposition Proposition 12.148) [112, 127]. In this construction, $U(1)_L$ breaking is encoded in $\langle\Phi_\nu\rangle \neq 0$, and the light eigenstates may be chosen Majorana ($\chi = \chi^c$) in the decoupling regime, with family phases carried by the PMNS holonomy and transforming under charge conjugation [127] (Definitions 12.136 and 12.156 and Proposition 12.115).

Families and anomaly/HSP compatibility

Triplicating the chiral multiplets by introducing a flat family fibre $\mathcal{F}$ leaves all anomaly constraints unchanged (Subsection Section 12.5.4; see also the anomaly block eq. (12.46)). Because each entry of the anomaly block is linear in traces over representations, replication by N_F families simply multiplies every contribution by N_F:

$$\sum_{f\in\text{all fam.}} Y_f\,T_2^G(R_f) = \sum_{a=1}^{N_F}\sum_{i\in 1\text{ gen.}} Y_i\,T_2^G(R_i) \;=\; N_F\sum_{i\in 1\text{ gen.}} Y_i\,T_2^G(R_i),$$

$$\sum_{f\in\text{all fam.}} Y_f^{\,3} = N_F\sum_{i\in 1\text{ gen.}} Y_i^{\,3}, \qquad \sum_{f\in\text{all fam.}} Y_f \;=\; N_F\sum_{i\in 1\text{ gen.}} Y_i,$$

$$\sum_{f\in\text{all fam.}} A^G(R_f) = N_F\sum_{i\in 1\text{ gen.}} A^G(R_i),$$

and similarly for all mixed and cubic non-abelian entries. Since each one-generation sum vanishes (SM calibration fixture Proposition 12.26, with assignments in table 12.1), all family-replicated sums vanish as well. Hence Stage-I viability is unchanged and the holosymmetric score remains $\mathcal{C}(A) = 1$ (Stage-I sieve in Section 12.2).

The family structure is implemented in transport by a flat, unitary auxiliary connection $\mathcal{A}_\mu^{(F)}$ acting on $\mathcal{F}$ (family fibre setup Definition 12.106); the total realized weave configuration-compatible derivative extends as

$$\begin{aligned} \mathcal{D}_\mu &\longrightarrow \mathcal{D}_\mu^{\mathrm{tot}} = \mathcal{D}_\mu + i\,\mathcal{A}_\mu^{(F)}, \\ \mathcal{A}_\mu^{(F)} &\in \mathfrak{u}(N_F) \\ &\quad \text{(realized weave configuration–compatibility background} \\ &\quad \text{Definition 6.192 and Lemma 12.87).} \end{aligned}$$

with vanishing family curvature and unitary holonomy,

$$\begin{aligned} F_{\mu\nu}^{(F)} &:= \partial_\mu \mathcal{A}_\nu^{(F)} - \partial_\nu \mathcal{A}_\mu^{(F)} + i\,[\mathcal{A}_\mu^{(F)}, \mathcal{A}_\nu^{(F)}] \;=\; 0, \\ U^{(F)}[\gamma] &:= \mathcal{P}\exp\!\Big(i\!\int_\gamma \mathcal{A}^{(F)}\Big) \in U(N_F) \\ &\quad \text{(Definition 12.106 and eq. (12.124)).} \end{aligned}$$

Flatness ensures path-independent family phases and preserves diagnostic realized reference configuration hygiene; unitarity preserves the inner product on the family fibre. Because $\mathcal{A}_\mu^{(F)}$ acts trivially on gauge indices and commutes with BRST on the diagnostic realized reference configuration,

$$\big[\hat{Q}_{\mathrm{BRST}},\, \mathcal{A}_\mu^{(F)}\big] \;=\; 0, \qquad \big[\hat{Q}_{\mathrm{BRST}},\, \mathcal{D}_\mu^{\mathrm{tot}}\big] \;=\; 0 \quad \text{(Sections 12.3 and 12.5.4)},$$

the family replication neither disturbs the anomaly block nor the Stage-I BRST structure. In particular, mixing matrices (CKM/PMNS) may be viewed as family holonomies derived from $\mathcal{A}_\mu^{(F)}$,

$$V_{\mathrm{CKM}},\, U_{\mathrm{PMNS}} \;\text{ as holonomies }\; U^{(F)}[\gamma] \quad \text{(Proposition 12.115 and eq. (12.134))},$$

while all certified meters on $\mathfrak{P}$ remain Nielsen-flat and BRST-safe (typed meters and witnesses in Definitions 9.187 and 12.16).

QCD composites via color–gauge projection

With BRST projection and confinement on the certified plateau $\mathfrak{P}$, physical hadrons arise as *color singlets* built from the n-quark sector (n=2 for mesons, n=3 for baryons) [38, 128] (color projector and spectral filtering in Definition 12.126 and Proposition 12.128; BRST diagnostic realized reference configuration set-up in Section 12.3). The hadronic Hilbert space is obtained by applying the color–singlet projector to the quark sector,

$$\mathcal{H}_{\mathrm{hadron}} \;:=\; P_{\mathrm{singlet}}\Big(\mathcal{H}_{\mathrm{quark}}^{(n)}\Big) \;\subset\; \mathcal{H}_{\mathrm{em}} \quad \text{(Corollary Corollary 12.125, eq. eq. (12.154))}.$$

Since P_{singlet} is gauge invariant, it commutes with the BRST charge on $\mathfrak{P}$ [117],

$$\big[\hat{Q}_{\mathrm{BRST}},\, P_{\mathrm{singlet}}\big] \;=\; 0,$$

so the BRST cohomology descends to the singlet subspace and colored excitations are projected out (see also Proposition 12.120 and Theorem 12.133).

Mesons (n=2). In color space $\mathbf{3}\otimes\overline{\mathbf{3}}=\mathbf{1}\oplus\mathbf{8}$, and the singlet projector may be written (Haar average) [30, 38]

$$(P_{\mathbf{1}})_i{}^j{}_k{}^\ell \;=\; \int_{\mathrm{SU}(3)} dU\; U_i{}^j\,(U^\dagger)_k{}^\ell \;=\; \frac{1}{3}\,\delta_i{}^\ell\,\delta_k{}^j.$$

Acting on bilinears yields color-singlet meson operators [128]

$$\mathcal{O}_M(x) \;=\; \bar{q}_j(x)\,\Gamma\,q^j(x),$$

whose two-point function displays only singlet poles in the chosen typed channel τ [128, 38]

$$\langle\,\mathcal{O}_M(x)\,\mathcal{O}_M^\dagger(0)\,\rangle^{(\tau)}_{\mathfrak{P},c} \;=\; \sum_{n\in\mathrm{Meson}_\tau} |\langle 0|\mathcal{O}_M|n,\tau\rangle|^2\;\Delta_n(x),$$

with Δ_n the (Euclidean/Minkowski) propagator of the n-th meson state (typed meters/witnesses in Definitions 9.187 and 12.16).

Baryons (n=3). For three quarks one has $\mathbf{3}\otimes\mathbf{3}\otimes\mathbf{3}=\mathbf{1}\oplus\mathbf{8}\oplus\mathbf{8}\oplus\mathbf{10}$, and the singlet is selected by the totally antisymmetric tensor [38, 128]:

$$\mathcal{O}_B(x) \;=\; \epsilon_{ijk}\,q^i(x)\,q^j(x)\,q^k(x)\,,\qquad (P_{\mathbf{1}})_{i_1i_2i_3}{}^{j_1j_2j_3} \;=\; \frac{1}{6}\,\epsilon_{i_1i_2i_3}\,\epsilon^{j_1j_2j_3}.$$

The corresponding correlator isolates baryonic singlet poles [128]

$$\langle\,\mathcal{O}_B(x)\,\overline{\mathcal{O}}_B(0)\,\rangle^{(\tau)}_{\mathfrak{P},c} \;=\; \sum_{n\in\mathrm{Baryon}_\tau} |\langle 0|\mathcal{O}_B|n,\tau\rangle|^2\;\Delta_n(x).$$

Typed placement and TT purity. All hadronic operators are placed in a specified typed channel τ so that certified meters interrogate only the intended spin/parity/flavor sector; this typed placement preserves TT purity throughout the extraction pipeline (typed architecture Definitions 9.187 and 12.16) [38]. In summary,

$$\boxed{\mathcal{H}_{\mathrm{hadron}} \;=\; P_{\mathrm{singlet}}\Big(\mathcal{H}^{(2)}_{\mathrm{quark}}\oplus\mathcal{H}^{(3)}_{\mathrm{quark}}\oplus\cdots\Big)\;\subset\;\mathcal{H}_{\mathrm{em}}}$$

realizes mesons and baryons as BRST-physical color singlets, in accordance with Corollary 12.125, eq. (12.154), and Theorem 12.133 [117, 128].

Antimatter (what it is on the diagnostic realized reference configuration, and how it shows up)

In the IQR/Tantrakyma architecture, the antimatter sector is *not* a separate theory: it is the charge–conjugate sector of the *same* kymonic fields, with (i) conjugate gauge representations, (ii) conjugate family phases, (iii) identical BRST/diagnostic realized reference configuration certification, and (iv) the same color–singlet projection for (anti)hadrons [38, 117, 128]. Neutrinos are special: if they are Dirac, they carry distinct antineutrinos; if they are Majorana (e.g. in the seesaw regime) the light states are self–conjugate and lepton–number–violating observables such as $0\nu\beta\beta$ become the antimatter signature [129, 130, 131, 127] (Section 12.7.4, Definition 12.136, and eq. (12.220)).

Three structural pillars (on a certified plateau). Together these enforce particle–antiparticle pairing for the SM sector on $\mathfrak{P}$:

1. complex unitary representations of the internal connection (and the induced $*$–structure on the operator bundle) [30, 38];
2. relativistic causality with a positive spectral measure on certified plateaux (CPT and conjugate poles) [38, 30];
3. neutrality of the family fibre with C-conjugated holonomy phases [127].

All of this is framed in the diagnostic realized reference configuration language of Section 12.4.4, Definitions 12.95, 12.96 and 12.106, Propositions 12.88 and 12.115, and eqs. (12.124) and (12.134).

Operational content. Dualizing

$$(R,\, U^{(F)}) \;\longmapsto\; (\overline{R},\, U^{(F)*})$$

leaves certified visibilities and typed SynTh witnesses invariant, so matter and antimatter occupy the *same* cell of the Symmetry–Asymmetry matrix (the holosymmetric lift is block-/cone–preserving) [117, 38]. At the field-theory level one has the matching of residues/poles [38, 128, 30]:

$$\begin{aligned} &\mathrm{Res}_{p^2\to m^2}\,\Big\langle 0\Big|T\{\psi(x)\,\bar\psi(0)\}\Big|0\Big\rangle \;\sim\; Z\,\Pi_{(m,s)}(p) \\ \Longleftrightarrow\;\; &\mathrm{Res}_{p^2\to m^2}\,\Big\langle 0\Big|T\{\psi^c(x)\,\overline{\psi^c}(0)\}\Big|0\Big\rangle \;\sim\; Z\,\Pi_{(m,s)}(p). \end{aligned}$$

Quadrant functionals obey (Definition 9.187)

$$\mathcal{Q}[\rho_0;\,\mathcal{O}] \;=\; \mathcal{Q}[\rho_0;\,\mathcal{O}^C], \qquad \mathcal{Q}\Big[\,|\Omega\rangle;\,\Pi_\alpha\Big] \;=\; \mathcal{Q}\Big[\,|\overline{\Omega}\rangle;\,\Pi_\alpha\Big],$$

so typed extraction makes no distinction between particle and antiparticle channels beyond the conjugation of additive charges and phases [38].

Hadronic realization. Color–singlet projection together with BRST cohomology produces hadrons and anti–hadrons as conjugate singlets [128, 117]:

$$\begin{aligned} &\text{mesons:} \quad q\bar q \;\text{ vs. }\; \bar q q, \\ &\text{baryons:} \quad \epsilon_{ijk}\, q^i q^j q^k \;\text{ vs. }\; \epsilon^{ijk}\, \bar q_i \bar q_j \bar q_k, \end{aligned}$$

with conjugate poles sharing the same (m, s) and carrying opposite additive charges (Definition 12.126, Proposition 12.120, and Theorem 12.133).

Neutrinos and the seesaw/Majorana limit (self-conjugacy)

Turning on symmetric Majorana blocks yields the general neutrino mass Lagrangian [127]:

$$\mathcal{L}_{\text{mass}}^{\text{Maj+Dirac}} \;=\; -\tfrac{1}{2}\,\Psi_D^{\top} C^{-1} \begin{pmatrix} m_L & m_D \\ m_D^{\top} & M_R \end{pmatrix} \Psi_D \;+\; \text{h.c.}, \qquad m_L = m_L^{\top}, \quad M_R = M_R^{\top},$$

where m_L and M_R are family endomorphisms on $(\mathcal{F}, \mathcal{A}^{(F)})$ (Definition 12.106), and one may take $\Psi_D^{\top} = (\nu_L,\ \nu_R^c)$.

In the Type–I seesaw limit ($m_L = 0$, $\|M_R\| \gg \|m_D\|$), block–diagonalization gives a *symmetric* light-neutrino mass matrix [129, 130, 131, 127]

$$m_\nu^{\text{eff}} \;=\; -\,m_D\, M_R^{-1}\, m_D^{\top} \;+\; \mathcal{O}(M_R^{-2}),$$

so the light mass eigenstates can be chosen Majorana ($\chi = \chi^c$). In this sense the $\mathbf{D}_{\mathcal{C}}$-dyad collapses: the dual acts trivially on each eigenstate (self–conjugacy), consistent with Proposition 12.158.

Diagonalizing the symmetric matrix defines the leptonic mixing as a *family holonomy* [127]:

$$U_{\nu_L}^{\top}\, m_\nu^{\text{eff}}\, U_{\nu_L} \;=\; \text{diag}(m_1, m_2, m_3), \qquad U_{\text{PMNS}} \;=\; U_{\ell_L}\, U_{\nu_L}^{\dagger},$$

with phases transforming under charge conjugation by $U^{(F)} \mapsto U^{(F)*}$ (Definitions 12.136 and 12.156 and Proposition 12.115).

The Axiom of Duality and Antimatter (how the pillars become a theorem)

Chapter 12 reframes antimatter as a *corollary of the Axiom of Duality* (Definition 2.100). The three pillars are presented as concrete incarnations of Duality:

(i) representation duality: $R \leftrightarrow \overline{R}$ (CRP, Definition 12.151),

(ii) spectral duality of poles/residues: KL+CPT (Proposition 12.154),

(iii) family-holonomy duality: $U^{(F)}[\gamma^{-1}] = U^{(F)}[\gamma]^{\dagger}$,
$C:\ U^{(F)} \mapsto U^{(F)*}$ (Definition 12.156).

Typed diagnostic realized reference configuration statement (Section 12.10). *Duality ⇒ Antiparticles (typed).* On a certified plateau $\mathfrak{P}$, any Tantrakyma kymon in a non-self-conjugate irrep acquires a unique conjugate class with the same (m/s) and opposite additive charges; family phases are complex-conjugated, and bound-state singlets map to their conjugates. If the irrep is real/pseudoreal or the relevant charge vanishes (or in the Majorana limit), Duality acts trivially and the field is self–conjugate.

Algebraic face (functorial duality). Formally, Duality is encoded by a contravariant *–functor on the internal representation category,

$$\mathfrak{D}_{\text{int}} : \ \text{Rep}(G_{\text{SM}}) \longrightarrow \text{Rep}(G_{\text{SM}}), \qquad \mathfrak{D}_{\text{int}}(R) = \overline{R}, \ \ \mathfrak{D}_{\text{int}}(U) = U^*,$$

so that, at the object level, $\mathbf{D}_{\mathcal{C}}$ implements charge conjugation

$$\psi \ \longmapsto \ \psi^c \ = \ \mathcal{C}\,\bar{\psi}^T, \qquad \psi^c \in \overline{R},$$

in accordance with Definition 12.151 and Proposition 12.152. This algebraic face sits alongside the geometric (family-holonomy) and spectral (KL+CPT) faces to yield the *Antiparticle Theorem* Corollary 12.172.

Consequence: quadrant invariance. Duality preserves the placement in the Symmetry–Asymmetry matrix (no inter–cell transport) because the holosymmetric lift is block-/cone–preserving; equivalently, there is no-law signalling and $(\bar{\Sigma})$–cone locality is maintained. In short, antimatter is the Duality shadow of the same certified dynamics on $\mathfrak{P}$, with all certified meters and spectral witnesses transforming contravariantly but *invariantly* in value.

The Yang–Mills Mass Gap

Chapter 12 develops two independent axioms–level routes from the operational "diagnostic realized reference configuration" pipeline to a nonzero mass gap in pure Yang–Mills (YM), while making explicit what remains to reach an unconditional, Clay–level solution. The key ingredients are typed, gauge–invariant meters (Definitions 9.187 and 12.16), KL/OS positivity in the continuum (Lemma 12.190), and the guards/plateau workflow that certifies spectral poles (Theorem 12.219) [30, 52, 117]. The section cross–links these pieces to the Continuum Theorem (Theorem 12.249) and the OS area–law route (Theorem 12.251), and clarifies the conditional nature of both routes (see Section 12.13.10) [123, 128, 32].

Operational resolution vs. Clay–level theorem: a guided introduction

Chapter 12 upgrades the diagnostic realized reference configuration pipeline to two axioms–level theorems that each imply a strictly positive first pole in standard glueball channels:

- **Route A (Spectral/guards).** Assuming continuum KL positivity with mild IR regularity, the guards/plateau workflow becomes the *Continuum Guards/Plateau Theorem* Theorem 12.249. On the certified plateau, typed two–point functions admit a KL form (Lemma 12.190); the effective mass is eventually monotone and convergent (Proposition 12.213); and the first pole satisfies $m_* > 0$ by typed certification (Theorem 12.219 and Corollary 12.250) [30].
- **Route B (Wilson loops).** Under OS reflection positivity, an area law for large rectangles yields exponential clustering and a universal lower bound $m \gtrsim c_{\text{cl}}\sqrt{\sigma} > 0$; this is the *OS area–law mass–gap theorem* Theorem 12.251. At this level, typed equality becomes a projection theorem (Proposition 12.216), meter locality reduces to microcausality (Proposition 12.217), and CPT/KL hygiene follows from OS→Wightman reconstruction (Lemma 12.190 and Propositions 12.154 and 12.158) [128, 30].

Both routes import a nonperturbative input—either (A) KL positivity with specified IR control, or (B) an OS–positive area law—so the mass–gap conclusions here are *conditional* on those inputs (see Section 12.13.10) [32].

Modal kinematics, transport, and the correlator kernel

The Chapter 4 "backbone" sets up the Fourfold modal fibre and the gauge–covariant transporter $\mathcal{U}$, from which the typed, gauge–invariant correlator kernel is built. For an admissible path $\gamma : x' \to x$, $\mathcal{U}(x \leftarrow x'; \gamma)$ is unitary, composes functorially, and trivializes on null paths; it assembles both the meter–to–meter transport of the diagnostic realized reference configuration pipeline (Definitions 9.187 and 12.16) and the loop variables that tie into Wilson–loop diagnostics [128]. This backbone targets exactly the Clay–level inputs: (A) it furnishes uniformity/IR guards for continuum correlators used in Theorem 12.249 and Proposition 12.213; (B) it interfaces with area–law control under OS positivity via transporter/loop comparison (Proposition 12.236 and Theorem 12.251) [30, 123].

Operator–valued metric, triadic split, and why it solves (A) and (B)

Chapter 5 introduces the operator–valued metric $\hat{g}_{\mu\nu}$ (second variation of a composite–operator distance) and its triadic split

$$\hat{g} \;=\; \hat{g}^{\mathrm{corr}} \;+\; \hat{g}^{\mathrm{vac}} \;+\; \hat{g}^{\mathrm{spec}},$$

which geometrizes the spectral/IR information needed for both routes [30, 22]. Under KL positivity and mild IR regularity, the guards (Lipschitz/TV/positivity) on diagnostic slopes become theorems, feeding directly into Theorem 12.249 and Proposition 12.213 [55, 30]. Under OS positivity, an area law enforces a finite correlation length in $\hat{g}$ and yields exponential clustering, matching the Wilson–loop bounds and leading to a strictly positive spectral threshold (Proposition 12.236 and Theorem 12.251) [128, 32]. Because typed meters are contractions of the Fourfold kernel, their two–point functions are governed by $\hat{g}^{\mathrm{spec}}$, ensuring positivity/locality control compatible with Lemma 12.190 and Proposition 12.217 [117, 52].

Development of the Yang-Mills Continuum Theorem

Assuming continuum KL positivity with IR regularity, Chapter 12 proves the continuum version of the diagnostic realized reference configuration pipeline [55, 30]:

1. **Certified plateau & IR window.** The mask/morphology and IR guards (Lipschitz/TV/-positivity) hold as theorems on a nonempty plateau $\mathcal{P}$ and an IR window $\mathcal{I}_{\mathrm{IR}} \subset \mathcal{P}$ (Theorem 12.249).

2. **KL form & effective–mass convergence.** Every typed, gauge–invariant two–point admits a KL (Laplace) representation with nonnegative measure, and the effective mass is eventually monotone with $\lim_{t\to\infty} m_{\mathrm{eff}}^{(O,\tau)}(t) = m_O$ (Lemma 12.190 and Proposition 12.213) [55, 22].

3. **Strict positive lower bound.** On $\mathcal{I}_{\mathrm{IR}}$,

$$C_O^{(\tau)}(t) = A_0 e^{-m_O t} + \int_{m_*}^{\infty} A(\mu)\, e^{-\mu t}\, d\mu, \qquad A_0, A(\mu) \geq 0.$$

   hence any isolated pole obeys $m_O \geq m_*^{\mathrm{op}} > 0$ (Corollary 12.250) [30].

4. **Typed equality & meter locality.** After band reduction, typed/untyped extractions coincide (Proposition 12.216), and meter locality follows from visibility additivity and cone–locality at the axioms level (Proposition 12.217) [117, 52].

5. **Typed certification (continuum).** Items (1)–(4) verify the hypotheses of the typed spectral certification theorem, establishing $m_O > 0$ on $\mathcal{P}$ without operational certification (Theorem 12.219) [30].

Altogether this yields a *Typed Spectral Gap in the continuum* with $m_O \geq m_*^{\mathrm{op}} > 0$ and $m_{\mathrm{eff}} \to m_O$ (Corollary 12.250), and underwrites the "Pipeline ⇒ Theorem" corollary (Corollary 12.252) [32, 128].

The Yang-Mills Mass Gap Theorem and OS/Wightman Axioms

In the OS/Wightman framework, assuming OS positivity and an area law for large rectangular Wilson loops, Chapter 12 proves [30, 128, 123, 32]:

1. **Exponential clustering.** For any local, gauge–invariant typed channel,

$$\left|\langle O(x)O(0)\rangle_{E,c}\right| \;\leq\; C_{\mathrm{cl}}\, e^{-c_{\mathrm{cl}}\sqrt{\sigma}\,|x|} \qquad (|x| \gg 1),$$

   by chessboard/Cauchy–Schwarz bounds that tie the connected two–point to a Wilson loop (Proposition 12.236) [128, 30].

2. **Strictly positive spectral threshold.** OS→Wightman reconstruction promotes clustering to a mass gap $m \geq c_{\mathrm{cl}}\sqrt{\sigma} > 0$, and typed effective masses converge to that threshold (Theorem 12.251) [30, 55].

At the axioms level, meter locality is just microcausality with visibility additivity (Proposition 12.217); KL/CPT hygiene descends from OS reconstruction (Lemma 12.190 and Proposition 12.154); and for pure YM the adjoint is real, so duality (charge conjugation) is inert in the glueball channels (Proposition 12.158) [30, 38]. All prerequisites (OS positivity, area law, transporter identifications) are treated as axioms–level inputs; with them, the operational pipeline collapses to a single, self–contained proof (Theorem 12.251 and Corollary 12.252) [128, 30].

YM mass–gap theorem. Under either of two axioms–level inputs, Chapter 12 establishes a strictly positive first pole in every typed, gauge–invariant glueball channel:

1. **Continuum (spectral/guards) route.** *Assumptions:* continuum Källén–Lehmann (KL) positivity with mild IR regularity for YM two–point functions (Lemma 12.190). *Conclusions:* the guards/plateau workflow upgrades to the *Continuum Guards/Plateau Theorem* (Theorem 12.249); on the certified plateau, typed correlators admit a KL form and the effective mass is eventually monotone and convergent (Proposition 12.213). Typed spectral certification then yields a nonzero gap:

$$\begin{aligned} &m_* > 0 \qquad \text{and} \\ &m_{\text{eff}}^{(O,\tau)}(t) \xrightarrow[t\to\infty]{} m_O \quad \text{for each typed channel } (O,\tau) \\ &\qquad (\textit{Theorem 12.219and Corollary 12.250}). \end{aligned}$$

   Equivalently, for any isolated pole one has $m_O \geq m_*^{\text{op}} > 0$ on the certified window (Corollary 12.250).

2. **OS/Wilson–loop (area–law) route.** *Assumptions:* Osterwalder–Schrader (OS) reflection positivity and an *area law* for large rectangles,

$$-\log\langle W[\square(R,T)]\rangle_E = \sigma RT + o(RT), \qquad \sigma > 0,$$

   with transporter/loop identification as in Proposition 12.236. *Conclusions:* exponential clustering of all local gauge–invariant typed correlators and, after OS→Wightman reconstruction, a strictly positive spectral threshold (Theorem 12.251):

$$\left|\langle O(x)O(0)\rangle_{E,c}\right| \leq C_{\text{cl}}\, e^{-c_{\text{cl}}\sqrt{\sigma}\,|x|} \quad \Rightarrow \quad m \geq c_{\text{cl}}\sqrt{\sigma} > 0,$$

   with typed effective masses converging to that threshold. At this level, typed equality becomes a projection theorem and meter locality reduces to microcausality (Propositions 12.216 and 12.217); KL/CPT hygiene follows from OS reconstruction (Lemma 12.190 and Proposition 12.154), and for pure YM the adjoint is real so duality (charge conjugation) is inert in these channels (Proposition 12.158).

Conditional statement of the result. The theorem established in Chapter 12 asserts a *nonzero mass gap* in every typed, gauge–invariant glueball channel, *conditional* on one of two axioms–level inputs. Concretely, for each typed operator/meter channel (as specified by the typed architecture Definitions 9.187 and 12.16), the two–point function admits a Källén–Lehmann representation with nonnegative measure and a strictly positive lowest spectral threshold, and the corresponding effective mass converges to the first pole:

$$m_* > 0, \qquad m_{\text{eff}}^{(O,\tau)}(t) \xrightarrow[t\to\infty]{} m_O.$$

This is the content of the continuum typed certification and gap statements (Lemma 12.190, Proposition 12.213, Theorem 12.219, and Corollary 12.250).

Routes to the statement. *Route A (spectral/guards).* Assuming continuum KL positivity with mild IR control, the guards/plateau workflow becomes the *Continuum Guards/Plateau Theorem* (Theorem 12.249). On the certified plateau, typed correlators have KL form and the effective mass is eventually monotone and convergent, yielding $m_O \geq m_*^{\mathrm{op}} > 0$ in each typed channel (Lemma 12.190, Proposition 12.213, and Corollary 12.250) [55, 30]. *Route B (OS/area law).* Assuming OS reflection positivity and a Wilson–loop area law, exponential clustering is obtained; OS→Wightman reconstruction then gives a strictly positive spectral threshold $m \geq c_{\mathrm{cl}}\sqrt{\sigma} > 0$ and convergence of typed effective masses to that value (Proposition 12.236 and Theorem 12.251) [30, 128]. At this level, typed equality reduces to a projection theorem and meter locality to microcausality; CPT/KL hygiene descends from OS reconstruction (Propositions 12.154, 12.216 and 12.217 and Lemma 12.190), and in pure YM the adjoint is real so the duality (charge conjugation) action is inert in these channels (Proposition 12.158) [30, 38].

Meaning of "typed" in the statement. The gap is asserted *in the channels interrogated by certified meters*: BRST–physical, gauge–invariant, typed operators defined on the certified plateau (Definitions 9.187 and 12.16). In this setting, correlators either possess an isolated first pole at $m_O > 0$ or, equivalently, exhibit exponential decay with rate $\geq m_*^{\mathrm{op}}$, and the effective–mass estimator extracted by the typed procedure converges to m_O (Theorem 12.219 and Corollary 12.250).

Non–unconditional (Clay–level) status. The two routes above are *conditional*: each assumes one nonperturbative input—either (A) KL positivity with specified IR regularity (Route A) or (B) an OS–positive area law (Route B). A Clay–level resolution requires deriving those inputs *within YM itself* [132, 133] and then applying the same pipeline–to–theorem collapse (Section 12.13.10 and Corollary 12.252). These are precisely the nonperturbative facts the Clay Mathematics Institute Millennium Problem on *Quantum Yang–Mills and the Mass Gap* asks one to derive *within* YM [132, 133].

Part V: Cosmology

Chapter 13: Dark Energy and Dark Matter

Chapter 13 reframes the dark sector as a consequence of the dual-field vacuum of *Gaiamiti* (geometric) and *Tantrakyma* (material), rather than as an *ad hoc* addendum to ΛCDM (Sections 13.1 and 13.1.1). Empirically anchored by supernovae and CMB data [134, 135, 136], the chapter proposes that the large but oppositely signed vacuum contributions of the two sectors nearly cancel by dynamical self-regulation, leaving a small residual ε_Λ that acts as dark energy. Operationally, this residue is a symmetry-filtered expectation value—the "global interferometric residue"—whose late-time dynamics are governed by the Spin-0 *Apeiron* (a quintessence-like mode that also unifies with the inflaton in the early universe) (Section 13.2.7). The dark-matter sector is then developed along three complementary tracks: (i) a vector Spin-1 *Chrono–Kymon* (quantum torsion of the Tantrakyma), neutral under G_{SM} and naturally

long-lived; (ii) a scalar Spin-0 *Apeiron* with WIMP-like relic abundance but of geometric origin; and (iii) a "dark gravity" option in which entanglement-induced correlations source halo phenomenology without introducing new particles (Section 13.1.2).

Methodologically, the chapter elevates *selection*—rather than postulation—to first principle. The Holosymmetric Selection Principle (HSP) and its *interferometer* compare entire universal histories in two stages: Stage I prunes those that fail physical self-coherence, and Stage II maximizes the physical holosymmetric overlap among the survivors (Sections 13.1.3 and 13.1.4). A commutator-based *lensing* formalism diagnoses the algebraic "twist" that produces bright/dark interferometric structure, while the comparative verdict is delivered by the interferometer itself (Section 13.2.7). In this frame, dark-matter candidates (Spin-0 vs. Spin-1) are adjudicated as inequivalent K-actions whose cross-overlap vanishes; the realized candidate is the one that yields the "bright fringe" (maximal self-overlap) across the SN/CMB/BAO/lensing/growth effect-families, subject to the operational *visibility* + *spectral-decoupling* test

$$\left\|\mathcal{P}_G^{\mathrm{coh}}\,|\psi\rangle\right\|^2 > 0, \qquad \langle\psi|\mathcal{O}_{\mathrm{SM}}|\psi\rangle_{\mathrm{phys}} \approx 0 \quad (\textit{Sections 13.2.7and 13.2.8}).$$

For dark energy, HSP selects among the three cancellation histories ($\Lambda > 0, = 0, < 0$), with the observed branch read as the unique coherence maximum (Sections 13.2.6 and 13.2.7). The energy budget is then partitioned by projector-filtered expectations into $(\rho_{\mathrm{DE}}, \rho_{\mathrm{DM}}, \rho_{\mathrm{vis}})$ and abundances $(\Omega_{\mathrm{DE}}, \Omega_{\mathrm{DM}}, \Omega_{\mathrm{vis}})$, linking sector dominance to chronogeometric regimes (Section 13.4.7). Finally, the entire construction is located within a timeless, background-free dynamical law—the Universal Quantum Constraint,

$$\hat{H}_{\mathrm{univ}}\,|\overset{\circ}{\boldsymbol{\Psi}}\rangle = 0,$$

from which ordinary Schrödinger evolution for subsystems emerges conditionally by clock-conditioning on $\mathcal{H}_{\mathrm{phys}}$ (Section 13.6).

Total energy from the universal Hamiltonian. Chapter 13 begins by fixing the comoving energy density through the *spectral resolution* of the regulated universal Hamiltonian $\hat{H}_{\mathrm{univ}} = \hat{H}_\Phi + \hat{H}_\Theta + \hat{H}_{\mathrm{int}}$ acting on the emergent/GNS space $\mathcal{H}_{\mathrm{em}}$. In the BRST-physical realized weave configuration, the realized Holo–State $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$ yields

$$\rho_{\mathrm{tot}} = \frac{1}{\mathrm{Vol}(\mathcal{X}_q)}\left\langle\overset{\circ}{\boldsymbol{\Psi}}\right|\hat{H}_{\mathrm{univ}}\left|\overset{\circ}{\boldsymbol{\Psi}}\right\rangle_{\mathrm{phys}} = \frac{1}{\mathrm{Vol}(\mathcal{X}_q)}\int_{\sigma(\hat{H}_{\mathrm{univ}})} E\,\mathrm{d}\mu_{\overset{\circ}{\boldsymbol{\Psi}}}^{\hat{H}_{\mathrm{univ}}}(E),$$

which fixes both the operational meaning (expectation value per comoving cell) and the measure-theoretic content (projector-valued spectral integral) of ρ_{tot} (see Definition 13.92 and eq. (13.144)). With the universal quantum constraint enforced on $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$, this definition is regulator-ready and sets up the partition of energy components that follows.

Holosymmetric partition and sum rule. The chapter then *resolves the identity* on the BRST-physical space by the coherence projectors

$$\mathcal{P}_{\mathrm{vac}}^{\mathrm{sym}} + \mathcal{P}_{\mathrm{dark}}^{\mathrm{coh}} + \mathcal{P}_{\mathrm{vis}}^{\mathrm{coh}} = \mathbf{1}_{\mathrm{phys}}, \qquad \mathcal{P}_i\mathcal{P}_j = \delta_{ij}\mathcal{P}_i,$$

so that HSP selection cancels cross-interferences and induces the abundance partition

$$\rho_{\mathrm{tot}} = \rho_{\mathrm{DE}} + \rho_{\mathrm{DM}} + \rho_{\mathrm{vis}}, \qquad \Omega_{\mathrm{DE}} + \Omega_{\mathrm{DM}} + \Omega_{\mathrm{vis}} = 1,$$

as formalized in Proposition 13.97 and eqs. (13.149) and (13.150). Conceptually, this ties the *kinematical* (projector) split to the *operational* (spectral) definition above, so every downstream observable can be expressed in terms of ρ_i or Ω_i without ambiguity.

Universe as an HSP fixed point. Finally, the chapter frames cosmic realization as a two-stage HSP adjudication over histories: *Stage I* enforces self-coherence/viability and *Stage II* maximizes holosymmetric overlap, selecting a background on which the $\{\mathcal{P}^{\mathrm{sym}}_{\mathrm{vac}}, \mathcal{P}^{\mathrm{coh}}_{\mathrm{dark}}, \mathcal{P}^{\mathrm{coh}}_{\mathrm{vis}}\}$ split is realized and the visible/DM/DE trichotomy emerges (Conclusion, "A Universe Sculpted by Coherence").

Dark energy (residual vacuum). Applying the symmetric vacuum projector $\mathcal{P}^{\mathrm{sym}}_{\mathrm{vac}}$ to $H_{\mathrm{vac}} := H^{\mathrm{vac}}_{\Phi} + H^{\mathrm{vac}}_{\Theta}$, the *residual vacuum* energy density is

$$\rho_{\mathrm{DE}} := \frac{1}{\mathrm{Vol}(\mathcal{X}_q)} \Big\langle \overset{\circ}{\boldsymbol{\Psi}} \Big| \, \mathcal{P}^{\mathrm{sym}}_{\mathrm{vac}} \, H_{\mathrm{vac}} \, \mathcal{P}^{\mathrm{sym}}_{\mathrm{vac}} \, \Big| \overset{\circ}{\boldsymbol{\Psi}} \Big\rangle_{\mathrm{phys}} \equiv \varepsilon_{\Lambda},$$

a small, positive *interferometric residue* singled out by Stage II (Definition 13.93 and eq. (13.145)). This locks the DE lane to a projector-defined piece of the universal Hamiltonian rather than to a bare cosmological constant.

Dark matter (holosymmetric-projected). The dark-sector abundance is defined via the coherence projector $\mathcal{P}^{\mathrm{coh}}_{\mathrm{dark}}$, which selects gauges G that pass Stage I and remain spectrally orthogonal to SM observables; the resulting ρ_{DM} is the physically certified "dark" contribution (see Definition 13.94).

Visible sector. Analogously, $\mathcal{P}^{\mathrm{coh}}_{\mathrm{vis}}$ isolates the SM lane so that the standard spectral sum over SM eigenstates $\{|\psi_\beta\rangle, E_\beta\}$ defines ρ_{vis} within the same universal spectral calculus (cf. the discussion leading to ρ_{vis} after eq. (13.149)).

Cross-term bookkeeping (if leakage). If an imperfect sectoring/regulator produces a nonzero cross contribution $\delta_{\mathrm{cross}}(\lambda)$, the chapter introduces

$$\Omega_{\mathrm{cross}}(\lambda) := \frac{\delta_{\mathrm{cross}}(\lambda)}{\rho_{\mathrm{tot}}(\lambda)}, \qquad 1 = \Omega_{\mathrm{DE}} + \Omega_{\mathrm{DM}} + \Omega_{\mathrm{vis}} + \Omega_{\mathrm{cross}},$$

to keep normalization exact while one takes regulator-clean limits (after Definition 13.99).

Regulator independence (ratios). Crucially, *ratios* of abundances are invariant under comoving cell rescalings and foliation changes—$\mathrm{Vol}(\mathcal{X}_q)$ cancels—provided the BRST-physical projectors and the HSP filter are used. Hence Ω_i are regulator-independent predictions (Remark 13.96).

Route A (spectral-first) inputs. For practical inference, Route A specifies background-dressed dispersions $E_{\mathrm{geo}}(k), E_{\mathrm{tym}}(k)$, fixes a Planckian UV cutoff $\lambda_{\mathrm{cutoff}} = \ell_{\mathrm{P}} \Rightarrow k_{\mathrm{UV}} = \ell_{\mathrm{P}}^{-1}$, and IR cutoffs $k^{\mathrm{cos}}_{\mathrm{IR}} = H_0/c$ (with an optional local mass-gap IR where needed). These inputs let one compute $\rho_{\mathrm{tot}}(a)$ directly from the spectral data (Definition 13.134).

From $\rho_{\mathrm{tot}}(a)$ to distances and growth. On flat FLRW, the total density determines the dimensionless expansion rate via

$$E^2(a) = \frac{\kappa' c^2}{3H_0^2}\,\rho_{\mathrm{tot}}(a) = \frac{\rho_{\mathrm{tot}}(a)}{\rho_{\mathrm{tot}}(1)}, \qquad \kappa' = 8\pi G,$$

which plugs into the standard RW distance–redshift relations and linear-growth equation; curvature adds the familiar $\Omega_k a^{-2}$ term (Proposition 13.110 and Remark 13.111). Thus, the projector-resolved spectrum maps cleanly to late-time observables.

Two practical routes to numerical projectors. The chapter offers (i) a *constructive* SVD/eigendecomposition on GNS two-point Gram operators to realize $\mathcal{P}^{\rm coh}_{\rm vis}$ and $\mathcal{P}^{\rm coh}_{\rm dark}$ directly, and (ii) a *calibrated FLRW reduction* that fixes $\mathcal{P}^{\rm coh}_{\rm vis}$ and normalizes the dark projector to $\Omega_{\rm DM}(z{=}0)$ (procedure 13.139). Both routes preserve the holosymmetric normalization so that the abundance sum rule stays exact.

The temporal (τ–kairos) sector: DE, DM, and "dark gravity." The τ–lane is presented in three complementary branches that share the same chronogeometric placement and typed orthogonality with the TT sector.

Scalar (Gaiaton/Apeiron) branch. Relative to u^μ, the covariant fluid variables read

$$\rho_\varphi = \tfrac{1}{2}\dot\varphi^2 + \tfrac{1}{2}D^2\varphi + V, \quad p_\varphi = \tfrac{1}{2}\dot\varphi^2 - \tfrac{1}{6}D^2\varphi - V, \quad q_\mu = -\dot\varphi D_\mu\varphi, \quad \pi_{\mu\nu} = D_{\langle\mu}\varphi\, D_{\nu\rangle}\varphi,$$

and the field obeys the *relational KG* equation $\Box^{(\rm rel)}\varphi - V_{,\varphi} = 0$ with $\Box^{(\rm rel)} = \bar\Sigma^{\mu\nu}\nabla^{(\rm rel)}_\mu\nabla^{(\rm rel)}_\nu$ (eq. (13.94)). In the FLRW/comoving limit this reduces to $\rho_\varphi = \frac{1}{2}\dot\varphi^2 + V$, $p_\varphi = \frac{1}{2}\dot\varphi^2 - V$, giving $w_\varphi \simeq -1$ for slow-roll (Proposition 13.69). A shift symmetry yields the Noether current $J^\mu_\varphi = \nabla^{(\rm rel)\,\mu}\varphi$ with $\nabla^{(\rm rel)}_\mu J^\mu_\varphi = \Box^{(\rm rel)}\varphi = -V'(\varphi)$ (Proposition 13.81 and eq. (13.128)).

Vector (Chrono–Kymon, dark $U(1)_\tau$) branch. The gauge-invariant Stueckelberg/Higgsed *Noether current* is

$$J^\mu_\tau = \nabla^{(\rm rel)}_\nu\mathcal{H}^{\nu\mu} + \mu_C^2\left(K^{(\tau)\mu} - \frac{1}{\mu_C}\nabla^{(\rm rel)\,\mu}\theta\right),$$

which in unitary gauge becomes $J^\mu_\tau = \nabla^{(\rm rel)}_\nu\mathcal{H}^{\nu\mu} + \mu_C^2 K^{(\tau)\mu}$; with a dark Higgs, $J^\mu_\tau \mapsto J^\mu_\tau - j^\mu_{H_\tau}$ (Proposition 13.88 and eqs. (13.136) and (13.137)). On general backgrounds the Chrono–Kymon behaves like *cold DM*,

$$\langle w_{K^{(\tau)}}\rangle \simeq 0, \qquad \langle\dot\rho_{K^{(\tau)}}\rangle + \theta\,\langle\rho_{K^{(\tau)}}\rangle \simeq 0 \;\Rightarrow\; \rho \propto e^{-\int\theta\,dt},$$

and in FLRW this reduces to $\rho \propto a^{-3}$ (eq. (13.106) and Proposition 13.77). Typed TT isolation $\Pi^{[\rm TT]}_{\bar\Sigma}\hat\tau = 0$ guarantees *no extra GW polarizations*.

Macroscopic ("dark gravity") branch. A coherent τ–background shifts the diagnostic realized reference configuration geometry and the TT operator through *typed balance* relations,

$$\mathcal{L}_{hh}[\bar\Sigma'] = \mathcal{L}_{hh}[\bar\Sigma] + \varepsilon\,\delta\mathcal{L}_{hh}[\Delta\Sigma^{(\tau)}] + \cdots, \qquad \delta\bar G_{\mu\nu}[\Delta\Sigma^{(\tau)}] + \Lambda\,\Delta\Sigma^{(\tau)}_{\mu\nu} = \kappa\,\Delta\bar T^{(\tau)}_{\mu\nu},$$

thereby modifying the Einstein sector *without* introducing new spin-2 modes; microscopically, this is the same τ operator whose quanta produce particle-level dark matter (eq. (13.3) and surrounding text). The chapter also lists additional dark candidates—e.g., a massive/oscillatory Gaiaton (CDM-like) or slow-roll Gaiaton (DE-like) depending on the Hessian of $\mathcal{V}_\Phi$, and neutral spin-1 quanta from the universal connection $\mathcal{A}_\mu$ (Mition/Tantron)—see Candidate II & III for details.

Observable mass and the FLRW cross-check. Away from FLRW gauge, the *chronogeometric (quasilocal) mass* over a domain $\mathcal{D}_\tau$ with boundary S_τ is

$$M_{\rm obs}(\tau) = \frac{1}{c^2}\Big\langle\int_{\mathcal{D}_\tau}\sqrt{\gamma}\,\rho_{\rm tot}\,{\rm d}^3x\Big\rangle_{\rm phys} + \frac{c^2}{8\pi G}\Big\langle\oint_{S_\tau}(k-k_0)\,{\rm d}S\Big\rangle_{\rm phys},$$

which packages *bulk* (spectral energy) and *boundary* (extrinsic curvature) contributions into a single BRST-physical observable (Definition 13.155 and eq. (13.220)). For a homogeneous, flat domain at $a = 1$ this collapses to

$$M_{\rm obs} = \frac{H_0^2}{2G}\, R^3, \qquad \frac{\delta M}{M} \simeq 2\,\frac{\delta H_0}{H_0} + 3\,\frac{\delta R}{R},$$

providing a sharp FLRW *consistency check* (Remark 13.157). Predictively, one works with the *partition* $M_i = \Omega_i\, M_{\rm obs}$ (with $\sum_i \Omega_i = 1$) and with the background mapping $\Lambda_{\rm eff} = \Lambda + \kappa' \varepsilon_\Lambda$; beyond FLRW, deviations are precisely captured by the boundary term (same remark).

Chapter 13 welds the universal spectral definition of $\rho_{\rm tot}$ to a holosymmetric projector calculus that (i) cleanly partitions DE/DM/visible energy, (ii) carries through to FLRW distances and growth, and (iii) grounds both particle-like (CK/Gaiaton) and macroscopic ("dark gravity") phenomenology in a single τ–typed sector—while keeping every step anchored to Definitions 13.92 to 13.94, 13.134 and 13.155, eqs. (13.3), (13.94), (13.106), (13.128), (13.136), (13.137), (13.144), (13.145), (13.149), (13.150) and (13.220), Propositions 13.69, 13.77, 13.81, 13.97 and 13.110, Remarks 13.96, 13.111 and 13.157, and procedure 13.139.

The Cosmic symmetry stack. At cosmic scales the theory organizes fluctuations of the unified chronogeometric field into radiative spin-2 and temporal (τ–kairos) lanes. The diagnostic realized reference configuration/background and quantum fluctuations are split as

$$\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}, \qquad \delta\hat{\Sigma}_{\mu\nu} = \hat{h}_{\mu\nu} + \hat{\tau}_{\mu\nu},$$

together with the *typed* TT projection that isolates the spin-2 sector and nulls any TT leakage from the τ-lane,

$$\hat{h}_{\mu\nu} := \Pi_{\bar{\Sigma}}^{[\rm TT]}\, \delta\hat{g}_{\mu\nu}, \qquad \Pi_{\bar{\Sigma}}^{[\rm TT]}\hat{\tau}_{\mu\nu} = 0 \quad \text{(eq. (13.119))}.$$

The τ–sector itself is typed into scalar and vector registers,

$$\hat{\tau}_{\mu\nu} = S^{(b)}_{\mu\nu}[\hat{\varphi}_b] \oplus S^{(L)}_{\mu\nu}[\hat{\varphi}_L] \oplus \tilde{S}^{(p)}_{\mu\nu}[\tilde{\varphi}] \oplus \sum_{\lambda=\pm} A^{(\lambda)}_{\mu\nu}\left[\hat{K}^{(\tau)}_{\lambda}\right] \quad \text{(eq. (13.113))}.$$

This typed architecture fixes what counts as genuine spin-2 radiation ($\hat{h}_{\mu\nu}$) versus temporal dynamics ($\hat{\tau}_{\mu\nu}$), and it underlies every symmetry statement that follows.

2) Layer 1: Unified chronogeometry. The chronogeometric symmetries acting on $\delta\hat{\Sigma}_{\mu\nu}$ form the semidirect product

$$\boxed{\mathcal{G}_{\rm Chronogeom} := {\rm Diff}_{\rm rel}(\mathcal{X}_q) \;\ltimes\; \left({\rm SO}(1,3)_{\rm local} \;\times\; \mathcal{G}^{(\rm eff)}_{\rm Dark}\right)} \quad \text{(eq. (13.122))}.$$

Here ${\rm Diff}_{\rm rel}(\mathcal{X}_q)$ are relational diffeomorphisms on the diagnostic realized reference configuration $(\mathcal{X}_q, \bar{\Sigma})$, acting by pullback on fields and adjoint transport on internal fibers; ${\rm SO}(1,3)_{\rm local}$

are local Lorentz rotations; and $\mathcal{G}_{\rm Dark}^{(\rm eff)}$ is the *effective* symmetry badge governing the τ–kairos registers (not SM matter):

$$\boxed{\mathcal{G}_{\rm Dark}^{(\rm eff)} \supset \underbrace{\mathrm{Shift}_{\varphi_b} \times \mathrm{Shift}_{\varphi_L} \times \mathrm{Shift}_{\tilde{\varphi}}^{\rm ax}}_{\text{scalar shifts}} \times \underbrace{U(1)_\tau}_{\text{Chrono–Kymon phase}} \times \underbrace{U(1)_{\rm hel}}_{\text{helicity plane}} \times \underbrace{\mathbb{Z}_2^{\rm HS}}_{\text{HSP tag}}} \quad \text{(eq. (13.108)).}$$

The three Shift factors act on the breathing/longitudinal/axial scalars $(\varphi_b, \varphi_L, \tilde{\varphi})$; $U(1)_\tau$ is the Chrono–Kymon phase symmetry of the τ-vector $K^{(\tau)}$; $U(1)_{\rm hel} \simeq SO(2)$ rotates the helicity doublet $(K_+^{(\tau)}, K_-^{(\tau)})$; and $\mathbb{Z}_2^{\rm HS}$ labels holosymmetric (HSP) branch stability. All continuous factors are *effective* and may be softly broken by the same operators (masses, mixings, potentials) that drive cosmology.

Typed orthogonality. By construction the TT projector never "sees" τ–modes, and the typed projectors $\Pi^{[\chi]}$ (with $[\chi] \in \{b, L, p, \lambda = \pm\}$) commute appropriately. The identities summarized around eq. (13.115) guarantee the *absence of extra GW polarizations* and clean bookkeeping of scalar/vector contributions.

3) Layer 2: The full cosmic stack (add the Standard Model). The *cosmic* group multiplies chronogeometry with the SM gauge sector,

$$\boxed{\begin{aligned}\mathcal{G}_{\rm Cosmos} &= \underbrace{\mathcal{G}_{\rm Chronogeom}}_{\text{acts on } \delta\hat{\Sigma}_{\mu\nu}=\hat{h}_{\mu\nu}\oplus\hat{\tau}_{\mu\nu}} \times \underbrace{G_{\rm SM}}_{\text{Tantrakyma (matter/gauge)}} \\ &= \left[\mathrm{Diff}_{\rm rel} \ltimes \left(\mathrm{SO}(1,3)_{\rm local} \times \mathcal{G}_{\rm Dark}^{(\rm eff)}\right)\right] \times \left[SU(3)_C \times SU(2)_L \times U(1)_Y\right],\end{aligned}}$$

as summarized in the chapter's "cosmic symmetry stack (summary)" (Section 13.4.1). The direct product with $G_{\rm SM}$ reflects that SM fields ("Tantrakyma") live on the chronogeometric background yet carry their *own* internal gauge charges.

The placement of the symmetry structures is *chronogeometric* in the strict sense (Section 13.4.1): the stack is organized on the chronogeometric fiber itself—namely, on the unified fluctuation $\delta\hat{\Sigma}$—rather than as an external juxtaposition "SM+GR". Concretely, spin-2 radiation is confined to the $\hat{h}_{\mu\nu}$ lane by the typed projector $\Pi^{[\rm TT]}$; all *temporal* physics (dark energy, dark matter, and "dark gravity" responses) inhabits the τ-lane with its own effective symmetries and soft breakings; and diffeomorphisms act *semidirectly* ($\ltimes$) because they simultaneously move fields on $\mathcal{X}_q$ and rotate the local fiber.

Operationally, the stack is not merely kinematical but the stage on which *holosymmetric tomography* and the *Holosymmetric Selection Principle (HSP)* operate. Stage I (safety) enforces BRST-physicality and typed/TT constraints, fixing the admissible set; Stage II (selection) then chooses, within that set, the background that maximizes holosymmetric coherence under SM-orthogonality (the HSP overlap rule) (Sections 13.1.3 and 13.1.4). This adjudication typically selects (i) a CK (Chrono–Kymon) branch with $\langle w_{K^{(\tau)}} \rangle \simeq 0$ and cosmic scaling $\rho \propto a^{-3}$ (cold-DM–like; cf. eq. (13.106) and Proposition 13.77), and/or (ii)

a scalar Gaiaton/Apeiron branch with slow roll $w_\varphi \simeq -1$ (DE-like), all while maintaining $\Pi^{[\mathrm{TT}]}$-orthogonality (no extra spin-2 modes) and compatibility with SM phenomenology.

Three consistency checks frame the construction. First, *GW consistency*: typed TT-null identities prevent extra gravitational-wave polarizations from the τ-sector; only $\hat{h}_{\mu\nu}$ propagates as spin-2 (see eqs. (13.115) and (13.119)). Second, *Modular/KMS alignment*: in stationary regimes (e.g. horizons), the time emerging from the universal constraint and internal clocks aligns with modular/KMS structure along the same symmetry lanes. Third, *soft breaking = cosmology*: the operators that softly break $\mathcal{G}_{\mathrm{Dark}}^{(\mathrm{eff})}$ (masses, potentials, mixings) are precisely those that yield late-time DE/DM phenomenology, tying the kinematics of the stack to macro-cosmic dynamics.

In practice, the stack *organizes projectors* and resolves the universal Hamiltonian's spectrum into DE/DM/visible contributions, leading to the abundance partition that feeds Ω_i and thus to observable distances and growth. It *guarantees regulator independence* of abundance ratios by preserving the TT/typed/HSP structure. Finally, it *bridges micro* $\leftrightarrow$ *macro*: the same τ-sector whose quanta realize particle-level dark matter (CK) also induces coherent background shifts ("dark gravity") in the diagnostic realized reference configuration equations without introducing new spin-2 degrees (eq. (13.3)).

$$\boxed{\mathcal{G}_{\mathrm{Cosmos}} = \left[\mathrm{Diff}_{\mathrm{rel}} \ltimes (\mathrm{SO}(1,3)_{\mathrm{local}} \times \mathcal{G}_{\mathrm{Dark}}^{(\mathrm{eff})})\right] \times G_{\mathrm{SM}}}$$

is placed *on* the chronogeometric fiber with typed TT-orthogonality and the effective dark badge (Shift $\times\, U(1)_\tau \times U(1)_{\mathrm{hel}} \times \mathbb{Z}_2^{\mathrm{HS}}$); operationally, HSP prunes/realizes this skeleton to produce the DE/DM/visible trichotomy used throughout Chapter 13 (Section 13.4.1 and eqs. (13.113) and (13.119)).

Universal Quantum Constraint: statement and setting. In the certified BRST-physical realized weave configuration of Chapter 13, the realized Holo–State $|\overset{\circ}{\boldsymbol{\Psi}}\rangle \in \mathcal{H}_{\mathrm{phys}}$ is annihilated by the total (relational) Hamiltonian,

$$\boxed{\hat{H}_{\mathrm{univ}}\,|\overset{\circ}{\boldsymbol{\Psi}}\rangle = 0} \qquad \text{(Thm. Theorem 13.168)},$$

with

$$\hat{H}_{\mathrm{univ}} = \hat{H}_\Phi + \hat{H}_\Theta + \hat{H}_{\mathrm{int}} \qquad \text{(Def. Definition 13.167)}.$$

This pins down the UQC as an *operator* constraint rather than a mere scalar energy condition, and it is posed directly on $\mathcal{H}_{\mathrm{phys}}$ so that all equalities are taken between BRST-physical states.

Why a constraint: the timeless thesis. Time in Chapter 13 is *relational*, not a fixed background parameter. Consequently, cosmic dynamics is cast in the Dirac/Wheeler–DeWitt spirit as a *single* quantum constraint on the universal state. Subsystem dynamics is then recovered by conditioning on good internal clocks and gauges (the chronogeometric placement ensures that such clocks live in the typed τ-lanes; see eqs. (13.115) and (13.119) for the TT/τ split that protects reparametrization invariance).

Ward-identity route (core proof step). Beginning with the adjoint-covariant Ward identity for time-reparametrizations, the source term decomposes into controlled deformations, BRST-exact pieces, and potential anomalies. On the certified operator realized weave configuration used in the chapter, BRST-exact contributions drop between physical states, Stage-I viability drives anomaly meters to zero, and calibrated boundary terms cancel. The net result is constancy of the total time-translation charge,

$$\dot{\mathbf{Q}}_{(t)} = 0, \qquad \Rightarrow \qquad \mathbf{Q}_{(t)} = \text{const.}$$

Under the universal closed calibration adopted for the cosmological background, $\mathbf{Q}_{(t)}$ is fixed to zero, which furnishes the bridge to the operator equation below.

From charge constancy to the operator law. With chronogeometric placement and typed TT-orthogonality, the clock-capable τ-lanes cannot induce explicit time-reparametrization breaking. Hence the universal evolution operator

$$U(s) = \exp\!\left(-\frac{i}{\hbar}\, s\, \hat{H}_{\text{univ}}\right)$$

leaves the universal state invariant, $U(s)\,|\overset{\circ}{\mathbf{\Psi}}\rangle = |\overset{\circ}{\mathbf{\Psi}}\rangle$. Differentiating at $s = 0$ yields the UQC in operator form,

$$\hat{H}_{\text{univ}}\,|\overset{\circ}{\mathbf{\Psi}}\rangle = 0 \qquad \text{(Thm. Theorem 13.168)}.$$

The role of the cosmic symmetry stack. The symmetry skeleton $\mathcal{G}_{\text{Cosmos}} = \left[\text{Diff}_{\text{rel}} \ltimes (\text{SO}(1,3)_{\text{local}} \times \mathcal{G}^{(\text{eff})}_{\text{Dark}})\right] \times G_{\text{SM}}$ (cf. Section 13.4.1) does not alter the content of the UQC; it *tightens* its hypotheses. Lane-wise BRST cleanliness, TT-nullity (eq. (13.119)), and a canonical clock sector ensure anomaly freedom and make the Ward argument above modular: each typed lane respects reparametrization invariance, so the combined system does as well, and the constraint holds exactly on $\mathcal{H}_{\text{phys}}$.

Dirac/BRST equivalence. Time-reparametrization invariance implies a first-class Hamiltonian constraint at the canonical level. In BRST quantization this lifts to an operator condition on physical states. The chapter's universal formulation matches this structure precisely, reproducing

$$\hat{H}_{\text{univ}}\,|\overset{\circ}{\mathbf{\Psi}}\rangle = 0$$

as the BRST-physical condition for the cosmological state, in agreement with Thm. Theorem 13.168.

Operational meaning. The UQC asserts that the *total relational energy vanishes*: the realized cosmos is a stationary, entangled Holo–State. What we ordinarily call "time'' appears only as a relational parameter extracted from internal clocks or (in stationary sectors) from modular/KMS flows. This same law underwrites the chapter's horizon-thermodynamic and cosmological applications, since it anchors all energetics to a constraint—rather than to an external time evolution.

Emergent Schrödinger dynamics for parts. Given a good internal clock and a BRST-physical factorization $\mathcal{H}_{\text{phys}} = \mathcal{H}_{\text{sub}} \otimes \mathcal{H}_{\text{clock}}$, conditioning the timeless state on clock readings produces unitary subsystem evolution with an effective Hamiltonian,

$$|\psi(t)\rangle := \frac{\langle t|\overset{\circ}{\mathbf{\Psi}}\rangle}{\|\langle t|\overset{\circ}{\mathbf{\Psi}}\rangle\|} \qquad \Longrightarrow \qquad i\hbar\,\partial_t|\psi(t)\rangle = \hat{H}_{\text{sub}}\,|\psi(t)\rangle,$$

compatible by construction with the UQC (see the post-theorem discussion around Thm. Theorem 13.168).

Spectral resolution and the abundance partition (after UQC). With the BRST-clean, lane-resolved $\hat{H}_{\text{univ}}$ and calibrated boundaries, the comoving total energy density is given by the spectral measure of $\hat{H}_{\text{univ}}$,

$$\begin{aligned}\rho_{\text{tot}} &= \frac{1}{\text{Vol}(\mathcal{X}_q)} \Big\langle \overset{\circ}{\boldsymbol{\Psi}} \Big| \hat{H}_{\text{univ}} \Big| \overset{\circ}{\boldsymbol{\Psi}} \Big\rangle_{\text{phys}} \\ &= \frac{1}{\text{Vol}(\mathcal{X}_q)} \int_{\sigma(\hat{H}_{\text{univ}})} E \, \mathrm{d}\mu_{\overset{\circ}{\boldsymbol{\Psi}}}^{\hat{H}_{\text{univ}}}(E) \quad (\textit{Definition 13.92and eq. (13.144)}),\end{aligned}$$

and, when holosymmetric projectors resolve the identity, it partitions as

$$\rho_{\text{tot}} = \rho_{\text{DE}} + \rho_{\text{DM}} + \rho_{\text{vis}},$$
$$\Omega_i = \frac{\rho_i}{\rho_{\text{tot}}}, \quad \sum_i \Omega_i = 1 \quad (\textit{Proposition 13.97and eqs. (13.149)and (13.150)}).$$

Summary arc.

$$\begin{aligned}\text{Tetralemma} &\rightsquigarrow \text{BRST/GNS} \rightsquigarrow \text{certified plateaues } \mathfrak{P} \\ &\rightsquigarrow \boxed{\hat{H}_{\text{univ}} \Big| \overset{\circ}{\boldsymbol{\Psi}} \Big\rangle = 0} \\ &\rightsquigarrow i\hbar\, \partial_t \, |\Psi_{\text{sub}}(t)\rangle = \hat{H}_{\text{sub}}^{\text{eff}}(t) \, |\Psi_{\text{sub}}(t)\rangle \, .\end{aligned}$$

Chapter 14: The Born Rule and Measurement

Chapter 14 reframes "measurement" as a local enactment of a universal, holosymmetric logic rather than an ad-hoc rupture of quantum dynamics. The narrative begins with the staged pipeline Evolve → Interfere → Re-evolve that is implemented by the holosymmetric interferometer (Section 3.1 and Definitions 2.123 and 3.12). System–apparatus coupling first creates robust entanglement and then decoheres into the environment, while classical definiteness is read as the Axiom of Duality's Boolean slice of an underlying fourfold logic (Section 14.1.3, Axiom 1.10, Definition 1.5, and Proposition 1.8). In this account, wave/particle duality is resolved as two faces of one process: an unobserved state inhabits the parasymmetric, superpositional stratum ($|\Psi_{\text{fin}}\rangle = \sum_k c_k \, |s_k\rangle \, |a_k\rangle \, |e_k\rangle$ with lane orthogonality; eq. (14.37)), while observation is a path-scattering "interference event" whose decohering broadcast into the Gaiamiti background leaves a locally classical record (Sections 14.1.3 and 14.3.1). Measurement thus becomes the localized instance of the Holosymmetric Selection Principle's global choice of laws: the same interferometric architecture that picks a self-coherent universe also governs the "click of a single detector" (Sections 13.2.7 and 14.1.1).

Program for the Born rule. Against this backdrop the chapter turns to probability. It first acknowledges standard practice—using the Born rule axiomatically for holosymmetric

effects E via $v(\rho; E) = \mathrm{Tr}(\rho E)$ in the interferometer/path-scattering models (Definitions 1.18 and 1.34, Corollary 3.3, and eq. (3.1))—before launching a three-stage derivation program (Sections 14.2, 14.2.1 and 14.2.2). *Stage 1* models measurement as relational decoherence that produces an orthogonal pointer basis for outcomes (preparation and imprint: eqs. (14.11) and (14.12); orthogonality at readout: Section 14.3.1 and eq. (14.37)). *Stage 2* uses relational envariance on the parasymmetric stratum—phase indifference, swap symmetry, and ancilla fine-graining—to fix

$$p_k = f(|c_k|) = |c_k|^2,$$

(phase indifference eq. (14.39), equiprobability eq. (14.4) and Lemma 14.9, fine-graining eq. (14.42); cf. eq. (14.43)).

Equivalently, probability is identified with the emergent inner-product norm of each branch, using the Fourfold Inter–Modal Correlator to define $\langle\cdot\rangle\cdot$ (eq. (14.38) and Section 14.3.2). *Stage 3* proves uniqueness from dialectic/monoidal constraints on the Axiom-of-Duality slice—additivity on orthogonal joins, unitary covariance, product factorization, coarse-graining noncontextuality, and continuity—together with Monoidal Duality's multiplicativity $f(rs) = f(r)f(s)$, which (with equiprobability) fixes the quadratic rule for all projectors (Section 14.3.3, Lemma 14.10, and Theorems 14.3, 14.4 and 14.11). In parallel, global-to-local consistency with the Universal Quantum Constraint—implemented by Page–Wootters conditioning with a good clock—excludes non-quadratic exponents on conservation grounds when time is emergent (Theorem 13.168, eqs. (13.227) and (13.236), Definition 13.158, Proposition 13.162, Corollary 13.163, and Sections 14.2.2 and 14.3.3). Holosymmetric Selection further prunes any non-Born assignments by diagnostic "twist" in commutator-lensing (Sections 13.2.7 and 14.3.3). The upshot is the explicit theorem: probabilities equal relational self-coherences,

$$p_k = \langle\Psi_{\mathrm{fin}}|\, P_k\, |\Psi_{\mathrm{fin}}\rangle = |c_k|^2, \qquad \text{with } P_k = \mathbb{I}_S \otimes |a_k\rangle\langle a_k| \otimes \mathbb{I}_E \text{ at readout (eq. (14.45)).}$$

No circularity: two-tier semantics at a fixed point. Finally, the chapter resolves the apparent circularity by distinguishing a pre-probabilistic state-valuation $v_0(E) = \omega(E)$, which Holosymmetry uses to build interferometric/lensing readouts (Definition 14.13 and Proposition 14.15), from operational probabilities. Stages 2–3 then prove that any calibration-compatible reparametrization $p = g \circ v_0$ collapses to the identity $g(x) = x$ (fixed-point identification), so Born-valued truths and Born probabilities coincide numerically without being conceptually conflated (Definition 14.14, Proposition 14.16, Theorem 14.19, Section 14.3.4, and Corollary 14.8). This fixed-point identification is background-independent—relying only on the effect algebra, the BRST/GNS state, and the universal constraint—and it seals the reflective consistency of the whole edifice: Tetralemma of Algebra + Duality + Holosymmetry together (under UQC) select the quadratic rule as the unique semantics of empirical frequency in an interdependent quantum reality (Definition 1.5, Axiom 1.10, Section 3.1, and Theorem 13.168).

Formal synthesis. The chapter's synthesis states that the combined force of the Tetralemma, Duality, Holosymmetry, and the Universal Quantum Constraint (UQC) determines the Born rule on the outcome stratum:

$$\boxed{\text{Tetralemma} + \text{Duality} + \text{Holosymmetry} + \text{UQC} \implies p_k = \langle\Psi_{\mathrm{fin}}|\, P_k\, |\Psi_{\mathrm{fin}}\rangle = |c_k|^2}.$$

In the emergent/GNS semantics, probabilities are *Born valuations* $v(\rho;E) = \mathrm{Tr}(\rho E)$ of holosymmetric effects (Definitions 1.18 and 1.34). Inner products are not primitive but derived from the Fourfold Inter–Modal Correlator via the emergent inner product (see eq. (14.38) and Ch. 4 for correlator-induced overlaps and triadic distances).

Holosymmetric interferometer (laboratory avatar of HSP). The holosymmetric interferometer provides the operational avatar of the HSP selection logic. With control basis $\{|\mathbf{0}\rangle_c, |\mathbf{1}\rangle_c\}$ steering the dual radiant registers, the controlled block is

$$\mathbf{U}_{\mathrm{ctrl}} = |\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} \quad \text{(eqs. (3.47) and (3.48))}.$$

The symmetric device is $\mathbf{U}_{\mathrm{holo}}^{(\mathrm{sym})} := \mathbf{U}_{\mathrm{Had}}\mathbf{U}_{\mathrm{ctrl}}\mathbf{U}_{\mathrm{Had}}$ (Definition 2.123); its tomographic/asymmetric generalization uses arbitrary $\mathbf{U}_{\mathrm{prep}}, \mathbf{U}_{\mathrm{read}} \in \mathrm{SU}(2)$,

$$\mathbf{U}_{\mathrm{holo}}^{(\mathrm{asym})} := \mathbf{U}_{\mathrm{read}}\ \mathbf{U}_{\mathrm{ctrl}}\ \mathbf{U}_{\mathrm{prep}} \quad \text{(Section 3.1)}.$$

For the symmetric device, the bright-port probability for input $|\mathbf{0}\rangle_c \otimes |\Psi_{\mathrm{RR}}\rangle$ is

$$p_0 = \tfrac{1}{4}\left\|(\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G})\,|\Psi_{\mathrm{RR}}\rangle\right\|^2 \quad \text{(Corollary 3.3 and eq. (3.1))}.$$

The *commutator lens* emphasizes that fringes are commutators in disguise,

$$[\mathbf{U}_{\mathrm{Had}}, \mathbf{U}_{\mathrm{ctrl}}] = \tfrac{1}{\sqrt{2}}\begin{pmatrix} 0 & \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} - \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} \\ \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} - \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} & 0 \end{pmatrix} \quad \text{(Definition 3.12)},$$

so postselecting $|\mathbf{0}\rangle_c$ yields

$$|\Psi_{\mathrm{post}}\rangle = \frac{\hat{\mathbf{U}}_+\,|\Psi_{\mathrm{RR}}\rangle}{\|\hat{\mathbf{U}}_+\,|\Psi_{\mathrm{RR}}\rangle\|}, \qquad \hat{\mathbf{U}}_+ := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}.$$

Measurement as path-scattering: Evolve → Interfere → Re-evolve. A laboratory measurement enacts the HSP-selected rules via the holosymmetric pipeline Evolve→Interfere→Re-evolve, with controls/readouts on the dual registers; conditioning on a control outcome initiates the re-evolve stage (Section 3.1). The separable preparation and local imprint are

$$|\Psi_{\mathrm{initial}}\rangle = \Big(\sum_i c_i\,|s_i\rangle\Big) \otimes |a_0\rangle \otimes |e_0\rangle, \qquad \sum_i |c_i|^2 = 1 \quad \text{(eq. (14.11))},$$

followed by a controlled imprint $U_{SA} : |s_i\rangle|a_0\rangle \mapsto |s_i\rangle|a_i\rangle$, producing $|\Psi_{\mathrm{ent}}\rangle = \sum_i c_i|s_i\rangle|a_i\rangle|e_0\rangle$ (eq. (14.12)). A perfectly correlating imprint is realized by the effective controlled Hamiltonian

$$\boxed{\hat{H}_{SA}^{\mathrm{eff}} = \sum_i |s_i\rangle\langle s_i| \ \otimes\ \hat{h}_i, \quad \hat{h}_i = \frac{\Omega_i}{2}\Big(|a_i\rangle\langle a_0| + |a_0\rangle\langle a_i|\Big)} \quad \text{(eq. (14.24))}.$$

Coupling the apparatus to the environment drives global decoherence; descending from a bi-local interaction action, it is presented as

$$\boxed{\hat{H}_{AE} := g_{AE}\int d^d x'\ \mathrm{Tr}\Big[\hat{\mathbf{\Phi}}_E^\dagger(x')\,\mathcal{U}_{\Phi\Theta}(x', x_A)\,\hat{\mathbf{\Theta}}_A(x_A)\Big]} \quad \text{(eqs. (14.13) and (14.29))}.$$

Born valuation at readout and the decoherence mechanism. In the minimal "measurement as relational entanglement'' cartoon,

$$\hat{U}_{\text{int}}\Big(\sum_i c_i|s_i\rangle \otimes |A_0\rangle\Big) = \sum_i c_i\,|s_i\rangle \otimes |A_i\rangle,$$
$$p_i = \langle\Psi_{\text{fin}}|(\mathbf{1}\otimes P_i)|\Psi_{\text{fin}}\rangle = |c_i|^2 \quad (\textit{Section 14.1.2and eq. (14.45)}).$$

With environment E,

$$|\Psi_{\text{SAE}}\rangle = \sum_i c_i\,|s_i\rangle \otimes |A_i\rangle \otimes |E_i\rangle, \quad \rho_S = \sum_i |c_i|^2 |s_i\rangle\langle s_i| + \sum_{i\neq j} c_i c_j^*\,\langle E_j|E_i\rangle\,|s_i\rangle\langle s_j|,$$

so as $\langle E_j|E_i\rangle \to 0$ the off-diagonals die out; the Ch. 4 triad $(d_{\text{corr}}, d_{\text{vac}}, d_{\text{spec}})$ explains this suppression and the emergence of effective classicality under coarse-graining (cf. Section 14.3.1).

Duality and the Twofold readout. By the *Axiom of Duality*, communicable outcomes inhabit a Twofold subspace with orthonormal basis $\{|\mathring{\mathbf{0}}\rangle, |\mathring{\mathbf{1}}\rangle\}$ and projectors $P_{\mathbf{0}}, P_{\mathbf{1}}$ satisfying $P_{\mathbf{0}} + P_{\mathbf{1}} = \mathbf{1}_{\text{Twofold}}$ and $P_{\mathbf{0}}P_{\mathbf{1}} = 0$; the classical outcome "$\mathbf{0/1}$" is then the Born valuation $p_{\mathbf{i}} = \text{Tr}(\rho P_{\mathbf{i}})$ (Section 14.1.3 and Axiom 1.10). On this AD slice, an admissible probability functional must obey additivity on orthogonal joins, unitary covariance, product factorization, coarse-graining noncontextuality, and continuity (Section 14.3.3). In particular, *Monoidal Duality* imposes multiplicativity on products (Lemma 14.10), which for rank-1 branches yields $f(rs) = f(r)f(s)$ and, by continuity, $f(r) = r^\alpha$; equiprobability for n equal branches fixes $\alpha = 2$. Thus the Stage-2 branch rule extends to *all* projectors:

$$p(P \mid \psi) = \langle\psi|\,P\,|\psi\rangle \quad \text{(Theorems 14.3, 14.4 and 14.11)}.$$

UQC and conditional dynamics. All laboratory conditioning sits inside the *Universal Quantum Constraint*

$$\boxed{\hat{H}_{\text{univ}}\Big|\mathring{\mathbf{\Psi}}\Big\rangle = 0}, \qquad \hat{H}_{\text{univ}} = \hat{H}_\Phi + \hat{H}_\Theta + \hat{H}_{\text{int}} + \hat{H}_\partial \quad \text{(Theorem 13.168 and eq. (13.236))}.$$

Deparametrizing with a good clock C (Stone-covariant PVMs $\{\Pi_t\}$) yields the conditioned state

$$\varrho_{\text{sub}|t} = \frac{\Pi_t\,\varrho\,\Pi_t}{\text{Tr}(\Pi_t\varrho)} \quad \text{(PVM)}; \quad \text{POVM via Naimark/Kraus (Corollary 13.163)}.$$

On certified plateaues $\mathfrak{P}$, the PVM case gives unitary conditional dynamics

$$i\hbar\,\partial_t\,|\Psi_{\text{sub}}(t)\rangle = \hat{H}^{\text{eff}}_{\text{sub}}(t)\,|\Psi_{\text{sub}}(t)\rangle\,,$$
$$\hat{H}^{\text{eff}}_{\text{sub}}(t) = \hat{H}_{\text{sub}} + \langle t|\,\hat{H}_{\text{int}}\,|t\rangle \quad (\textit{Proposition 13.162and eq. (13.227)}),$$

while the POVM case adds a small dissipator $i\hbar\,\mathcal{D}_t[\cdot] \to 0$ in the sharp limit (Corollary 13.163). Good-clock hygiene ensures $[A, \Pi_t] = 0$ for $A \in \mathcal{A}$ and TT-nullity (Definition 13.158 and Lemma 13.161), so the AD axioms (product, covariance) and Tetralemma coarse-graining logic apply unchanged to $\varrho_{\text{sub}|t}$.

Cosmic stack and typed cleanliness. On BRST-physical diagnostic realized reference configuration the symmetry stack

$$\mathcal{G}_{\text{Cosmos}} = \Big[\text{Diff}_{\text{rel}} \ltimes (\text{SO}(1,3)_{\text{local}} \times \mathcal{G}^{(\text{eff})}_{\text{Dark}})\Big] \times G_{\text{SM}}$$

acts on $\delta\hat{\Sigma}_{\mu\nu} = \hat{h}_{\mu\nu} \oplus \hat{\tau}_{\mu\nu}$ with typed TT-orthogonality; BRST-physical projectors split vacuum/dark/visible lanes via $\mathcal{P}^{\text{sym}}_{\text{vac}} + \mathcal{P}^{\text{coh}}_{\text{dark}} + \mathcal{P}^{\text{coh}}_{\text{vis}} = \mathbf{1}_{\text{phys}}$. This typed hygiene is what keeps the measurement template well-posed and TT-clean (cf. Section 14.3.3).

Putting the synthesis to work. On the Parasymmetric (synthesis) stratum the post-measurement state is lane-orthogonal,

$$|\Psi_{\text{fin}}\rangle = \sum_k c_k \, |s_k\rangle\,|a_k\rangle\,|e_k\rangle\,, \qquad \langle a_k\rangle\, a_\ell = \delta_{k\ell}, \quad \langle e_k\rangle\, e_\ell \approx \delta_{k\ell} \quad (\textit{eq. (14.37)}),$$

and Parasymmetric automorphisms together with environment-assisted invariance yield: phase indifference $p_k = f(|c_k|)$ (Section 14.2.2 and eq. (14.39)); swap-envariance/equiprobability on equal-modulus branches (eq. (14.4) and Lemma 14.9); and ancilla fine-graining that, for rational weights $|c_k|^2 = r_k/N$, gives $p_k = r_k/N = |c_k|^2$, extended by continuity to all amplitudes,

$$\boxed{p_k = |c_k|^2} \quad \text{(Stage 2; eq. (14.43))}.$$

Holosymmetry provides the operational preparation (pointer orthonormality), the projector reading, and the dynamical suppression of non-Born alternatives through commutator-lensing diagnostics (Sections 13.2.7 and 14.3.3). Duality (AD) imposes the Boolean outcome layer and composition axioms; Monoidal Duality enforces multiplicativity, which—together with equiprobability—fixes the unique quadratic rule for all projectors,

$$p(P \mid \psi) = \langle\psi|\, P\, |\psi\rangle \quad (\textit{Theorems 14.3, 14.4 and 14.11}).$$

Finally, UQC and Page–Wootters conditioning tie local valuations to globally conserved expectations; non-quadratic rules violate this consistency and are thus excluded (Sections 14.3.3 and 14.3.4). Equivalently, probability equals the *relational self-coherence* computed with the emergent inner product (Section 14.3.2):

$$p_k = \langle\Psi_{\text{fin}}|\, P_k\, |\Psi_{\text{fin}}\rangle = |c_k|^2.$$

Good–clock bridge. Deparametrizing the UQC with a good clock C (Definition 13.158) ensures gauge-clean, TT-safe conditioning and supplies the bridge identity for measurements in the $\{|s_k\rangle\}$ basis at time t,

$$\boxed{p_k(t) = \text{Tr}\Big(\varrho_{\text{sub}|t}\ |s_k\rangle\langle s_k|\Big) = |c_k(t)|^2}\,,$$

with POVM-clock corrections $i\hbar\,\mathcal{D}_t[\cdot] \to 0$ in the sharp limit (Corollary 13.163). This shows that Holosymmetry, Duality, and the Tetralemma act on the conditioned state exactly as required to deliver the Stage-2/3 Born rule on certified plateaues, completing the synthesis.

Chapter 15: Quantum Thermodynamics

hapter 15 derives the thermodynamic laws inside the IQR framework by starting from a C*-algebraic foundation on the PQS continuum $\mathcal{X}_q$, where the global Quantum Holo-State $\omega^{\mathring{\mathbf{\Psi}}}$ selects its GNS realisation and thereby fixes the Emergent Hilbert Space $\mathcal{H}_{\text{em}}$ together with subsystem algebras $\mathcal{A}_X \subset \mathcal{A}(\mathcal{C}_d)$ (Definitions 1.32 and 4.287 and Sections 1.4 and 1.4.1) [30, 22]. In this representation-theoretic setting, conserved Noether currents and charges are handled at the algebraic level and tied to the relational Hamiltonian structure (Sections 15.1, 15.1.1 and 15.1.3) [52, 117]. "Equilibrium'' is inherently relational: it is the stationarity of the restricted state on a chosen subalgebra together with vanishing net conserved current between subsystems, formulated within the unified causal architecture fixed by the expectation-level chronogeometry $\bar{\Sigma}$ and Good-clock gauge class $[C]$ (Definitions 5.26, 5.150 and 13.158) [119, 20, 45]. In this light, the Zeroth Law is shown to be a theorem of algebraic self-consistency on the universal state—an informational transitivity statement rather than a bare postulate—obtained by projection to the physical cohomology and by enforcing Σ-causal support (Section 15.1.2, Theorem 15.7, and Corollary 5.145) [30, 22, 119, 40].

> Equilibrium between two $\bar{\Sigma}$-causally connected *and interacting* subsystems X and Y—with $\bar{\Sigma}_{\mu\nu}(x) := \omega^{\mathring{\mathbf{\Psi}}}\big(\hat{\Sigma}_{\mu\nu}(x)\big) = \big\langle \Omega_\omega \big| \, \pi_\omega\big(\hat{\Sigma}_{\mu\nu}(x)\big) \, \big| \Omega_\omega \big\rangle$—is a dynamical alignment of their joint algebra $\mathcal{A}_X \vee \mathcal{A}_Y$. It is defined by two predicates on their shared, $\hat{G}_\Lambda$–based statistics *evaluated with the emergent inner product*: (1) **stationarity** under the Good-clock flow on $\mathcal{A}_X \vee \mathcal{A}_Y$, $\partial_t \langle O \rangle_{XY} = 0$ for all $O \in \mathcal{A}_X \vee \mathcal{A}_Y$; and (2) **vanishing of the relational Noether current** between X and Y, expressed componentwise for the fundamental fields as
>
> $$\Big\langle \hat{J}^\mu_{\mathbf{\Phi}}(X) - \hat{J}^\mu_{\mathbf{\Phi}}(Y) \Big\rangle = 0, \qquad \Big\langle \hat{J}^\mu_{\mathbf{\Theta}}(X) - \hat{J}^\mu_{\mathbf{\Theta}}(Y) \Big\rangle = 0,$$
>
> where $\hat{J}^\mu_{\mathbf{\Phi}}$ and $\hat{J}^\mu_{\mathbf{\Theta}}$ are the $\hat{\mathbf{\Phi}}$-Gaiamiti and $\hat{\mathbf{\Theta}}$-Tantrakyma Noether currents, respectively.

Building on this foundation, the First Law is reimagined by explicitly decomposing the total dynamics into coherent transport and interaction-induced monitoring [22, 30]. The current–kernel interaction $\hat{H}_{AE}$ and the record (pointer) channel $\mathbb{E}_A$ (Definitions 15.25 and 15.27) yield a stroboscopic reduced evolution whose coherent part is unitary while incoherent exchange is captured by unital pinching on the record algebra [22, 30]. Identifying local energy change with interaction-driven exchange then gives the diagnostic realized reference configuration-wise balance $\Delta U_A = Q - W$ from the Heisenberg evolution of the subsystem within a conserved total energy, now interpreted relationally (Section 15.1.3, eq. (15.12), Theorem 15.18, and Definition 15.14) [119].

The Second Law appears as an H-theorem for Page–Wootters–conditioned, coarse-grained dynamics: the CP–divisible product-limit generator on the state-conditioned active band enforces entropy monotonicity $S_A(t)$ via pinching/majorisation arguments, with equality only when the state resides on the record algebra and the instantaneous motion there is unitary (Propositions 13.172, 15.31 and 15.41, Definitions 15.27, 15.37 and 15.56, Assumption 15.39, Lemma 15.30, Theorem 15.50, eq. (15.14), and Corollary 15.48) [22, 30, 119].

Finally, the Third Law is recast as *asymptotic coherence*: as the relational coupling $g \downarrow 0$, GKLS rates vanish and the reduced dynamics tends to a unitary group so that entropy production flattens; exact isolation (the analogue of $T{=}0$) is unattainable at any finite Good-clock time, in harmony with resource-bounded control limits and purity lower bounds (Section 15.1.5, Propositions 15.52 and 15.62, and Theorem 15.65) [119, 137, 47, 48]. The chapter distinguishes the PQS both as the asymptotic ideal for laboratory subsystems and as a gravitationally realised but operationally inaccessible limit (consistent with the two-level applicability domain and the Planck floor), thereby preserving unattainability while situating thermodynamics as relational information dynamics throughout (Definitions 15.54 and 15.55 and Corollary 15.63) [40, 42, 20].

Foundational arena and measurement semantics. The operational stage is the PQS continuum $\mathcal{X}_q$ equipped with the C*-algebra of relational observables $\mathcal{A}(\mathcal{C}_d)$ (Definition 1.32). A coherent Quantum Holo-State $\omega^{\mathring{\mathbf{\Psi}}}$ selects the GNS representation, yielding the Emergent Hilbert Space $\mathcal{H}_{\mathrm{em}}$ and the vector state $|\mathring{\mathbf{\Psi}}\rangle$ (Sections 1.4 and 1.4.1 and Definition 4.287). Subsystems are subalgebras $\mathcal{A}_X \subset \mathcal{A}(\mathcal{C}_d)$ with restricted states ω_X, and probabilities are assigned by the Born valuation

$$v(\rho; E) = \mathrm{Tr}(\rho E) \qquad \text{(Definition 1.18)}.$$

The A–E interface: current–kernel coupling and pointer channel. Matter–geometry exchange is governed by the current–kernel interaction Hamiltonian

$$\hat{H}_{AE} := \int_{\mathcal{X}_q} \mathrm{d}x\, \mathrm{d}x'\ \mathrm{Tr}\Big[\hat{J}^{\Phi}_{(\Theta)}(x)\, \mathcal{U}_{\Phi\Theta}(x,x')\, \hat{J}^{\Theta}_{(\Phi)}(x')\Big],$$

with $\mathcal{U}_{\Phi\Theta}$ transported along $\bar{\Sigma}$-chronogeodesics (Definition 15.25). In the einselection regime the record (pointer) algebra on A admits orthogonal projectors $\{P_k\}_k$ and induces the Relational pointer channel

$$\mathbb{E}_A(\rho) := \sum_k P_k\, \rho\, P_k,$$

which is CPTP, idempotent, unital, and transport-covariant (Definition 15.27). A stroboscopic reduced step of size Δt combines unitary transport with einselection,

$$\Phi_{\Delta t}(\rho) := \mathbb{E}_A\Big(U_A(\Delta t)\, \rho\, U_A(\Delta t)^\dagger\Big), \qquad U_A(\Delta t) = e^{-iH_{A,\mathrm{eff}}\Delta t},$$

so unitary conjugation preserves spectral entropy and entropy production is due solely to $\mathbb{E}_A$.

Pinching increases entropy (engine of the H–theorem). Because $\mathbb{E}_A$ is a pinching onto the commutative record algebra,

$$S\Big(\mathbb{E}_A(\rho)\Big) \ \geq\ S(\rho), \qquad S(\rho) = -\,\mathrm{Tr}(\rho \log \rho),$$

with a proof via simultaneous block-diagonalization, a discrete Fourier twirl that displays a random-unitary decomposition, Ky–Fan majorization, and Schur–concavity (Lemma 15.30, Proposition 15.31, and Definition 15.27).

Chronogeometric time and effective reduced evolution. In the macroscopic timelike regime the emergent proper time is

$$t_{\mathrm{geom}}(x,x') = \Big[\bar{g}_{\mu\nu}(x)\, \Delta x^\mu \Delta x^\nu\Big]^{1/2} \quad (\textit{Definition 5.95}),$$

and the reduced dynamics on $\mathcal{A}_A$ assumes the time-local GKLS form

$$\dot{\rho}_A(t) = -i\,[H_{A,\mathrm{eff}}(t), \rho_A(t)] \;+\; \text{interaction/monitoring terms from } \hat{H}_{AE},$$

with t the Good-clock axis aligned with $\bar{\Sigma}$ in the macroscopic regime (Definitions 5.95, 13.158 and 15.25). The unital CP–divisible Markovian limit is described next.

Stage 3: causal architecture and admissible reduced dynamics. Expectation–level chrono-geometry $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ fixes the effective light–cone and Σ–causal support, ensuring monitoring/measurement on the record algebra cannot signal superluminally (see Definitions 5.26, 5.122 and 5.150 and Corollary 5.145). This supports a unital, CP–divisible continuum limit with channel-anisotropic rates when $\mathcal{U}_{\Phi\Theta}$ is anisotropic, formalised in Definition 15.37, Assumption 15.39, and Propositions 15.41 and 15.43. Admissible (CP–divisible) reduced dynamics $\{\Lambda_t\}_{t\geq 0}$ are defined precisely in Definition 15.37 (two-parameter family, composition law, divisibility).

Product–limit GKLS generator on the active band. With coarse-graining on the state-conditioned active band $\mathcal{H}_{\mathrm{act}}$ (Definition 15.56) and the finite-dimensionality postulate (Assumption 15.39), the Trotterized maps

$$\Lambda_t^{(n)} := \Big(\mathbb{E}_A \circ \mathrm{Ad}_{e^{-iH_{A,\mathrm{eff}}t/n}}\Big)^n$$

converge to a unital CPTP semigroup with GKLS generator $\mathcal{L}$ (Proposition 15.41). If $\mathcal{U}_{\Phi\Theta}$ is anisotropic, the resulting CP–divisible rates are channel-anisotropic while preserving Σ-causal support (cf. Definition 15.25 and Proposition 15.43).

Zeroth Law in the IQR setting (contextual highlight). Equilibrium equality $\mathrm{Eq}^{[\chi]}_{\mathrm{phys}}(\mathfrak{B}_A, \mathfrak{B}_B \mid \mathfrak{P})$ is obtained by projection to $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$, together with clock-gauge invariance and Σ-causal support (Theorem 15.7).

First Law (diagnostic realized reference configuration-wise). On a diagnostic realized reference configuration $\mathfrak{B}^{\langle \mathrm{DRRC}\rangle}$,

$$U_A(t) := \mathrm{Tr}\Big(\rho_A(t) H_A(t)\Big), \qquad \boxed{\Delta U_A = Q - W} \quad \text{(eq. (15.12))},$$

where Q is the incoherent contribution to ΔU_A generated by unital CP–divisible reductions $\mathbb{E}_A$, and W is the coherent Good-clock contribution from $H_A(t)$ (see Theorem 15.18, Definition 15.14, and Section 15.1.3).

> For an interacting subsystem A (coupled to an environment E with $\hat{H} = \hat{H}_A + \hat{H}_E + \hat{H}_{AE}$—environment Hamiltonian $\hat{H}_E$ plus interaction $\hat{H}_{AE}$ with A), *energy is relational*: along the Good-clock flow, the internal energy
>
> $$U_A(t) := \mathrm{Tr}\Big(\rho_A(t)\, H_A(t)\Big)$$
>
> changes only by heat current into A minus external power on A,
>
> $$\frac{dU_A}{dt} = J_A^{(Q)}(t) - P_A(t),$$

with expectations taken in the emergent inner product (from $\hat{G}_\Lambda$). These rates may be written equivalently as

$$J_A^{(Q)}(t) := \Big\langle i\,[\hat{H}_{AE}, \hat{H}_A] \Big\rangle = \mathrm{Tr}\Big(H_A\,\dot{\rho}_A(t)\Big),$$
$$P_A(t) := -\Big\langle \partial_t \hat{H}_A \Big\rangle = -\,\mathrm{Tr}\Big(\rho_A(t)\,\partial_t H_A(t)\Big).$$

When the interaction splits across the fundamental fields, $\hat{H}_{AE} = \hat{H}_{AE}^{(\mathbf{\Phi})} + \hat{H}_{AE}^{(\mathbf{\Theta})}$, the heat current decomposes componentwise in the Gaiamiti and Tantrakyma sectors:

$$J_A^{(Q)}(t) = \Big\langle i\,[\hat{H}_{AE}^{(\mathbf{\Phi})}, \hat{H}_A] \Big\rangle + \Big\langle i\,[\hat{H}_{AE}^{(\mathbf{\Theta})}, \hat{H}_A] \Big\rangle =: J_{A,\mathbf{\Phi}}^{(Q)}(t) + J_{A,\mathbf{\Theta}}^{(Q)}(t),$$

where $\hat{\mathbf{\Phi}}$ (Gaiamiti) and $\hat{\mathbf{\Theta}}$ (Tantrakyma) are the fundamental fields mediating exchanges. Consequently, over any interval $[t_0, t_1]$,

$$\Delta U_A = \int_{t_0}^{t_1} J_A^{(Q)}(t)\,dt \;-\; \int_{t_0}^{t_1} P_A(t)\,dt \;=\; Q_A - W_A.$$

Second Law (differential H–theorem). For admissible reduced dynamics one has

$$\frac{d}{dt} S_A(t) \;\geq\; 0,$$
with equality iff $[\rho_A(t), P_k] = 0\ \forall k$ and either $\kappa = 0$ or $\mathbb{E}_A(\rho_A(t)) = \rho_A(t)$,

so entropy is monotone and becomes flat precisely on the record algebra under purely unitary motion there (Theorem 15.50, eq. (15.14), and Corollary 15.48). Time and entropy co-emerge from Page–Wootters conditioning and einselection; Good-clock gauge-class invariance and macroscopic alignment with t_{geom} are explicit (Proposition 13.172, Definition 13.158, Lemma 15.6, and Theorem 13.168).

Second Law (relational form). Along any Good–clock axis t, the reduced state $\rho_A(t)$ follows a unital flow that is *divisible into intermediate completely positive, trace-preserving quantum channels* (CP-divisible), driven by the current–kernel coupling of the Fourfold correlator $\hat{G}_\Lambda$ and the pointer channel $\mathbb{E}_A$:

$$\dot{\rho}_A(t) \;=\; -\,i\,[H_{A,\mathrm{eff}}, \rho_A(t)] \;+\; \kappa\Big(\mathbb{E}_A(\rho_A(t)) - \rho_A(t)\Big),$$
$$\underbrace{\mathbb{E}_A(\rho) = \sum_k P_k\,\rho\,P_k\,.}_{\text{einselection}}$$

$$\underbrace{\forall\, t \geq r \geq s\ \exists\ V_{t,r}\ \text{CPTP}:\ \ \Phi_{t\leftarrow s} = V_{t,r}\circ\Phi_{r\leftarrow s},\quad \Phi_{t\leftarrow s}(\mathbb{I}_A) = \mathbb{I}_A}_{\text{CP-divisible, unital}}$$

$$\underbrace{P_j P_k = \delta_{jk} P_k, \qquad \sum_k P_k = \mathbb{I}_A,}_{\text{record projectors}}$$

$$\underbrace{\mathbb{E}_A^2 = \mathbb{E}_A}_{\text{pinching (conditional expectation)}},$$

$$\underbrace{\mathcal{R}_A = \{X \in \mathcal{B}(\mathcal{H}_A) : [X, P_k] = 0\ \forall k\} = \mathrm{Fix}(\mathbb{E}_A)}_{\text{record algebra and fixed points}}$$

With probabilities/expectations taken in the emergent inner product (from $\hat{G}_\Lambda$), the von Neumann entropy of A cannot decrease:

$$t_2 \geq t_1 \implies S_A(t_2) \geq S_A(t_1),$$
$$\frac{d}{dt} S_A(t) = \kappa\ \mathrm{Tr}[\rho_A(t)\log\rho_A(t) - \mathbb{E}_A(\rho_A(t))\log\rho_A(t)] \geq 0.$$

All increase is due to environment-induced superselection (the conditional expectation $\mathbb{E}_A$ onto $\mathcal{R}_A$); equality holds iff

$$[\rho_A(t), P_k] = 0\ \forall k \quad \text{and} \quad \mathbb{E}_A(\rho_A(t)) = \rho_A(t) \quad \text{or} \quad \kappa = 0,$$

i.e. the state already lies in the record algebra and the instantaneous motion on that commutant is unitary. Causality is respected: the flow preserves $\bar{\Sigma}$–lightcone support and admits no-law-signalling.

Third Law: coherence and residual entropy. For fixed $t < \infty$ on the active band $\mathcal{H}_{\text{act}}$,

$$\lim_{g\to 0} S\Big(\Lambda_t^{(g)}\rho_A(0)\Big) = S\Big(\rho_A(0)\Big), \qquad \lim_{g\to 0}\Lambda_t^{(g)} = \mathrm{Ad}_{e^{-iH_{A,\text{eff}}t}},$$

so the information-theoretic arrow flattens ($\dot{S} = 0$) (Proposition 15.52). If $H_{A,\text{eff}}$ has a $\mathfrak{g}$-fold degenerate ground space, then asymptotically on that manifold

$$S_0 = \log\mathfrak{g} \quad (k_B = 1),$$

and $S_0 = 0$ when $\mathfrak{g} = 1$ (Corollary 15.53). A two-level summary consolidates: (P) no thermodynamic map below the Planck floor; (R) vanishing production in isolation; residual $S_0 = \log\mathfrak{g}$; and strengthened unattainability at finite resources (Corollary 15.68).

Third Law (relational form — coherence and unattainability). On an *instrumented realization on a certified plateau* with a Good–clock $C = (\hat{T}, U_t)$ and

state–conditioned active band $\mathcal{H}_{\rm act}$, the reduced evolution at coupling $g\geq 0$ is the unital, CP–divisible flow $\{\Lambda_t^{(g)}\}_{t\geq 0}$ on $\mathcal{B}(\mathcal{H}_{\rm act})$ generated by

$$\mathcal{L}_g[\rho] = -\,i\,[H_{A,\rm eff},\rho] + \sum_\alpha \gamma_\alpha(g)\Big(L_\alpha \rho L_\alpha^\dagger - \tfrac{1}{2}\{L_\alpha^\dagger L_\alpha,\rho\}\Big)\,,$$
$$\Gamma(g) := \sum_\alpha \gamma_\alpha(g) \;\xrightarrow[g\downarrow 0]{}\; 0.$$

Coherence (isolation) limit. For every finite t,

$$\boxed{\lim_{g\to 0}\Lambda_t^{(g)} = \mathrm{Ad}_{e^{-iH_{A,\rm eff}t}}, \qquad \lim_{g\to 0} S\Big(\Lambda_t^{(g)}\rho\Big) = S(\rho), \qquad \lim_{g\to 0}\dot S_A^{(g)}(t) = 0}$$

i.e. entropy production *flattens* and the flow becomes unitary (asymptotic coherence).

Unattainability under finite resources. Any *instrumented realization on a certified plateau*–admissible, resource–bounded control with $\|H_{AE}(t)\| \leq g_{\max}$, $\Gamma(g(t)) \leq \kappa_{\max}$ and causal/Planckian onset

$$\underbrace{T_{\rm on} := \max\Big\{T_{\rm sig},\, \tau_{\rm Pl}^{(\rm thermo)}\Big\}}_{\substack{\text{signal latency / Planck floor}\\ (\Sigma\text{–causal \& timing constraints})}}$$

obeys, for any $T>0$ and any pure $|\psi\rangle$,

$$\boxed{\Big\|\rho_A(T) - |\psi\rangle\langle\psi|\,\Big\|_1 \geq \Big[\;\delta_0 - \underbrace{(c_1 g_{\max} + c_2\kappa_{\max})}_{\text{resource envelope}}\,(T - T_{\rm on})\;\Big]_+,}$$
$$\boxed{\delta_0 := \inf_{|\psi\rangle}\Big\|\rho_A(0) - |\psi\rangle\langle\psi|\,\Big\|_1 > 0}$$

(where $[x]_+ := \max\{x,0\}$). Equivalently,

$$T \;\geq\; T_{\rm on} + \frac{\delta_0 - \epsilon}{c_1 g_{\max} + c_2\kappa_{\max}} \quad \text{to reach} \quad \Big\|\rho_A(T) - |\psi\rangle\langle\psi|\,\Big\|_1 \leq \epsilon.$$

Residual entropy (degenerate ground spaces). Let P_0 project onto the ground manifold of $H_{A,\rm eff}$ and $\mathfrak{g} := \mathrm{rank}\,P_0$. If the long–time unital dynamics is ergodic on $P_0\mathcal{H}_{\rm act}$, then

$$\boxed{\lim_{g\to 0}\lim_{t\to\infty}\rho_A^{(g)}(t) \;=\; \frac{P_0}{\mathfrak{g}}, \qquad S_0 \;=\; \log\mathfrak{g}}$$

so $S_0 = 0$ iff the ground state is nondegenerate.

Causality and gauge. The flow preserves $\bar\Sigma$–lightcone support and is invariant in the Good–clock gauge class: no law–signalling, and all statements hold for any admissible emergent time variable.

Endogenous cutoffs, Planck floor, and active band. Thermodynamics applies only on the realized level; the pre-physical level is outside thermodynamic applicability (Definition 15.54). Any diagnostic realized reference configuration satisfies the Planck floor,

$$\Delta t \geq \tau_{\mathrm{Pl}}^{(\mathrm{thermo})} > 0, \qquad \Delta \ell \geq \ell_{\mathrm{Pl}}^{(\mathrm{thermo})} > 0 \quad (\textit{Definition } 15.55),$$

and the active band $\mathcal{H}_{\mathrm{act}}$ is fixed by state-conditioned spectral thresholds (Definition 15.56 and Assumption 15.39).

Resource-bounded control, Lipschitz bounds, and unattainability. Generator and trajectory norms obey

$$\|\mathcal{L}_{g(t)}\|_{1\to 1} \leq c_1 g_{\max} + c_2 \kappa_{\max},$$
$$\|\rho_A(t) - \rho_A(s)\|_1 \leq (c_1 g_{\max} + c_2 \kappa_{\max})\,|t-s| \quad (\textit{Lemma } 15.60),$$

and incorporating the Planck floor yields the earliest actionable time

$$t \;\geq\; T_{\mathrm{on}} := \max\{T_{\mathrm{sig}}, \tau_{\mathrm{Pl}}^{(\mathrm{thermo})}\} \quad (\textit{Lemma } 15.61).$$

For mixed $\rho_A(0)$ on $d := \dim \mathcal{H}_{\mathrm{act}} \geq 2$,

$$\inf_{\text{admissible controls on } [0,T]} \; \inf_{|\psi\rangle} \left\| \rho_A(T) - |\psi\rangle\langle\psi| \right\|_1 \;\geq\; \Big[\, \delta_0 - (c_1 g_{\max} + c_2 \kappa_{\max})\,(T - T_{\mathrm{on}}) \,\Big]_+,$$

so attaining a pure state in finite T requires resources that scale accordingly (Theorem 15.65 and Proposition 15.62). Within the two-level applicability and with resource-bounded control, no finite Good-clock time $T < \infty$ can reach the pre-physical PQS regime or an exactly PQS-coherent state; one needs $t_n \uparrow \infty$ with vanishing AE correlations (Corollary 15.63).

Synthesis of the laws. The First Law ($\Delta U_A = Q - W$) separates incoherent exchange (heat via $\mathbb{E}_A$) from coherent Good-clock driving (work) (eq. (15.12)). The Second Law ensures entropy production via pinching, with equality only on the record algebra under unitary motion there (Theorem 15.50 and eq. (15.14)). The Third Law states that at finite times the isolation limit preserves coherence and any residual entropy equals $\log \mathfrak{g}$ from ground-space degeneracy (Proposition 15.52 and Corollary 15.53). Altogether, these statements are Good-clock gauge-invariant and consistent with Σ-causal support, so the arrow of time emerges as information flow under unitary+einselection dynamics on the active band, unifying chronogeometry, entropy, and macroscopic evolution (Proposition 15.41 and Definitions 5.26 and 13.158).

Chapter 16: Black Hole Physics

Chapter 16 develops a black-hole physics of the relational programme by placing the information problem at center stage. After recalling Hawking's discovery that black holes radiate thermally [87], the chapter formulates the information paradox as a tension between semi-classical thermality and quantum unitarity, and frames its modern resolution in terms of entanglement entropy and the Page curve [138, 139]. In this account, the event horizon becomes the locus where classical description meets quantum coherence, and the paradox is

re-posed as a question about how correlations are transferred during evaporation (Section 16.1) [140, 20].

The technical core is a sourced, covariant wave equation for the temporal curvature tensor, $\Box_{\bar{g}}\bar{t}_{\mu\nu} + \mathcal{M}_{\mu\nu\rho\sigma}[\bar{g}]\bar{t}^{\rho\sigma} = \kappa'\langle\hat{T}^{\rm vac}_{\mu\nu}\rangle$ (eq. (16.467)), which encodes a dynamical feedback loop between Hawking flux and near-horizon chronogeometry. From this, the chapter defines a "temporal entropy density" $\sigma_t = \frac{1}{2}\bar{t}_{\mu\nu}\bar{t}^{\mu\nu}$ and establishes an information-conservation law $S_{\rm rad}(t) + S_t(t) = S_{\rm initial}$ governing unitary evaporation (Section 16.5) [20]. The early-time rise, turnover at t_P, and late-time decline of $S_{\rm rad}(t)$ (the Page curve) are then derived as the anti-phased exchange between stored temporal coherence and radiative entanglement, rendering "islands" and related holographic structures as effective summaries of the deeper relational dynamics [139, 140].

Finally, the horizon is reinterpreted as an ontological phase boundary between the macroscopic geometric regime and the mesoscopic relational regime (Section 16.1.10), with its radius determined by a transcendental relation corrected by vacuum coherence, $R_{\rm EH} + \alpha(\xi(R_{\rm EH})) = \frac{2GM}{c^2}$ (eq. (16.21)) [40, 42]. In the rotating case, the horizon acquires the quantum-corrected Kerr shift $R_{EH} \approx R^{(0)}_{EH} - \frac{a^2}{2R^{(0)}_{EH}}$ (eq. (16.426)), enabling a unified thermodynamics that maps the First/Second/Zeroth/Third Laws onto surface gravity, horizon area, and extremality bounds for the corrected geometry (Section 16.5) [141, 40]. The chapter closes by sketching observational handles—shadows, spectra, waveforms—and by making the unitary mechanism explicit: information stored in the central PQS is transferred to Hawking quanta via Apeiron/Kymon exchange across the shimmering horizon interface, resolving the paradox in a single relational narrative [122, 20].

In this relational account, the exterior "thermality" is explained algebraically rather than dynamically: restricting to the exterior algebra at (and outside) a Killing horizon yields a KMS state with respect to the horizon flow χ^a. Concretely,

$$\hat{K}_\omega = \beta_{\rm KMS}\hat{H}_\chi + c\,\mathbf{1} \quad ((16.481)),$$

and horizon regularity fixes $\beta_{\rm KMS} = 2\pi/\kappa$. Hence

$$T_H = \frac{\hbar\,\kappa_{\rm eff}}{2\pi k_B}, \qquad \kappa_{\rm eff} = \frac{c^2}{2}\,F'(R_{\rm EH}) \quad (eq.\ (16.5)),$$

so an exterior observer records KMS "thermality" while the global state remains pure (see Section 16.1.1). This resolves the apparent clash between semi-classical temperature and microscopic unitarity without altering quantum theory.

Geometrically, the unified lapse

$$N^2(r) = 1 - \frac{f(r)}{r} - A(r)$$

encodes the horizon as its largest root (eqs. (16.21) and (16.84)). Working perturbatively with $A(r) = \varepsilon\,a_1(r)$ and $f(r) = R_S + \varepsilon\,f_1(r)$, the first-order shift is

$$\delta R = f_1(R_S) - R_S\,a_1(R_S) \quad (eq.\ (16.161)),$$

so that $R_{\rm EH} = R_S + \varepsilon\,\delta R < R_S$ when $\rho_{\rm vac} > 0$. Equivalently,

$$\alpha\Big(\xi(R_{\rm EH})\Big) = R_S - R_{\rm EH} = -\,\varepsilon[f_1(R_S) - R_S a_1(R_S)] > 0 \quad (eq.\ (16.162)),$$

and this same horizon data fixes $\kappa_{\rm eff}$ via eq. (16.78) and T_H via eq. (16.79). The area/entropy shift then follows from $A_H = 4\pi R_{\rm EH}^2$,

$$S = \frac{A_H}{4G\hbar} = \frac{\pi R_S^2}{G\hbar}\Big(1 + 2\varepsilon\,\frac{\delta R}{R_S}\Big) + \mathcal{O}(\varepsilon^2),$$

providing a controlled $\mathcal{O}(\varepsilon)$ deviation tied to the same relational corrections.

Dynamically, the sourced evolution

$$\Box_{\bar g}\bar t_{\mu\nu} + \mathcal{M}_{\mu\nu\rho\sigma}[\bar g]\bar t^{\rho\sigma} = \kappa'\,\langle \hat T^{\rm vac}_{\mu\nu}\rangle \quad (eq.\ (16.467))$$

is complemented by a state-conditioned, band-limited source profile with spectral form

$$\rho_{\rm vac}(r) = \int_{\lambda_{\rm IR}(r)}^{\lambda_{\rm cutoff}(r)} \rho(\lambda;\bar\Sigma)\, w(\lambda;\,|\mathring{\mathbf{\Psi}}\rangle)\, E(\lambda;\bar\Sigma)\, d\lambda \quad (eqs.\ (16.129) and\ (16.130)),$$

so information is redistributed between the temporal sector and radiation rather than destroyed. This is captured by the conservation law

$$S_{\rm rad}(t) + S_t(t) = S_{\rm initial} \quad (Section\ 16.5),$$

whose early growth, turnover at t_P, and late decline reproduce the Page-curve morphology as an anti-phased exchange of correlations.

In the rotating setting, the dressed horizon/ergoregion are defined by the unified chrono-geometry: with $\chi^a = \partial_t + \Omega_H\,\partial_\phi$, the ergosurface satisfies $\bar\Sigma_{tt} = 0$ (eq. (16.202)) and the horizon is the largest root of $\bar\Sigma_{ab}\chi^a\chi^b = 0$ with

$$\Omega_H = -\frac{\bar\Sigma_{t\phi}}{\bar\Sigma_{\phi\phi}}\Big|_{r=R_H} \quad (eq.\ (16.203)).$$

To leading order one finds the quantum-corrected Kerr shift

$$R_{\rm EH} \approx R_{\rm EH}^{(0)} - \frac{a^2}{2\,R_{\rm EH}^{(0)}} \quad (eq.\ (16.426)),$$

and the associated unified thermodynamics is organized in Section 16.5, mapping the classical laws to surface gravity, corrected horizon area, and extremality bounds.

Finally, the framework ties directly to observations. Collecting the horizon-value data into

$$\boldsymbol{\theta} = \Big(a_1(R_S),\ f_1(R_S)/R_S,\ R_S a_1'(R_S),\ R_S f_1'(R_S)\Big),$$

one obtains leading QNM shifts,

$$\frac{\delta\omega_{\ell n}}{\omega_{\ell n}} = \mathcal{O}\Big(a_1(R_S),\ \tfrac{f_1(R_S)}{R_S},\ R_S a_1'(R_S),\ R_S f_1'(R_S)\Big),$$

and $\mathcal{O}(\varepsilon)$ deformations of the photon ring/shadow through the corrected lapse N^2. Together with the modular/KMS account of T_H (eq. (16.5)) and the conservation law above, this operationalizes the unitary mechanism: information stored in the central PQS is exported to Hawking quanta via Apeiron/Kymon exchange across the horizon interface, completing the single relational narrative introduced at the beginning of the chapter.

Opening motivation: horizons, singularities, and the information paradox. Chapter 16 motivates the program by treating the event horizon as a quantum–relational phase boundary and the putative singular interior as a limit of relational breakdown rather than a place [40, 42, 141, 20]. Within this framing, Hawking's prediction of an apparently thermal flux is put into tension with microscopic unitarity and the Page-curve expectation for entropy recovery [87, 138, 139]. The chapter's response is to embed black-hole physics in the IQR framework, where the exterior "thermality" arises from a modular/KMS restriction and coarse-graining, while the global state remains pure [30, 140]. The goal is a concrete, testable completion: derive horizon thermodynamics, locate and shift the event horizon relative to the classical Schwarzschild radius, and propagate those corrections into observables, all while keeping cross-references to the underlying field equations and modular structure explicit [40, 30].

The dual nature of Hawking radiation and the Hawking/KMS temperature. In the "isotropic" gauge $A = B$, the effective surface gravity simplifies to

$$\kappa_{\text{eff}} = \tfrac{c^2}{2}\, F'(R_{\text{EH}}),$$

which yields the Hawking temperature

$$T_H = \frac{\hbar\, \kappa_{\text{eff}}}{2\pi k_B} \xrightarrow[A=B=0]{\text{Schwarzschild}} \frac{\hbar c^3}{8\pi G M k_B},$$

as stated in (16.5). The chapter emphasizes the *dual* reading of Hawking radiation: it appears thermal to an exterior observer who restricts to a KMS algebra and coarse-grains over inaccessible modes, but the global state—treated in the IQR description—remains pure. This is elaborated in Section 16.1.1, where the KMS condition and environment-induced decoherence reconcile apparent thermality with unitary evolution.

Unified field equations for black holes. The dynamics are written in terms of the chrono-geometric tensor $\bar{\Sigma}_{\mu\nu}$, which combines the geometric and temporal sectors. Specializing the unified (semi-classical) Einstein equations to black-hole contexts leads to the vacuum constraints

$$\bar{G}_{\mu\nu}[\bar{\Sigma}] = 0,$$

to be solved subject to regularity at the horizon and asymptotic flatness at spatial infinity. This unified vacuum problem furnishes the stage on which horizon conditions, surface gravity, and thermodynamics are computed consistently from $\bar{\Sigma}_{\mu\nu}$.

Modeling the emergent geometry (static, spherically symmetric sector). In the static, spherically symmetric sector (Section 16.1.3), the chapter adopts a diagonal ansatz for $\bar{\Sigma}_{\mu\nu}$ encoded by the unified line element (16.18) together with the radial temporal tensor (16.108), featuring a lapse $N^2(r)$, a mass–length function $f(r)$ in the geometric block, and a temporal sector $\{A(r), B(r)\}$ [45, 43]. This ansatz converts the unified equations into coupled ordinary differential equations for $\{f; A, B\}$ (Sections 16.1.2 and 16.1.16) driven by dressed-vacuum sources, while the horizon and asymptotic conditions supply boundary data (cf. eq. (16.84)) [141]. The construction is designed so that horizon thermodynamic quantities can be read off directly from the lapse and its radial derivatives [40, 42].

The event horizon as a phase boundary. With the static ansatz, the lapse assumes the unified form

$$N^2(r) \;=\; 1 - \frac{f(r)}{r} - A(r).$$

The event horizon $R_{\rm EH}$ is defined as the largest root of this relation, equivalently the "phase boundary'' condition of eq. (16.84) together with the horizon conditions eqs. (16.16) and (16.21) [141]. These same relations anchor the effective surface gravity and temperature through eqs. (16.78) and (16.79); see the classical laws of black-hole mechanics and their thermodynamic identifications and Hawking's temperature [40, 87, 42].

Deriving the working equations (temporal tensor + geometric back-reaction). Projecting the unified equations onto the temporal sector yields an operator equation for the tt-profile A: a Lichnerowicz-type operator acting on A, sourced by the dressed vacuum (semiclassical back-reaction framework) [44, 43, 12, 40]. Simultaneously, the lapse is constrained by

$$N^2(r) \;=\; 1 - \frac{f(r)}{r} - A(r).$$

Combining these with the redshift/time-dilation relations in eq. (16.77), the phase-boundary condition in eq. (16.84), and the horizon relations in eqs. (16.16) and (16.21) yields the effective surface gravity eq. (16.78) and thereby T_H via eq. (16.79) [45, 141, 87, 40]. In the isotropic gauge $A = B$, the anisotropy vanishes and the first-order system closes cleanly (static, spherically symmetric reduction) [45]. Importantly, the apparent "halting of time'' at the horizon is interpreted as a projection/coarse-graining artifact; *chronogeometric viscosity* regularizes the flow that would otherwise become singular (for horizon-regular effective descriptions cf. the discussion of near-horizon thermodynamics) [40, 42].

Solving strategy I: asymptotic matching (outer solution). Outer solution ($r \gg R_S$)**.** In the decoupled, far-field regime, the operator equation for A reduces to a linear ODE whose Green's function exhibits a Yukawa-like kernel regulated by $\lambda \sim \ell_*$ (massive/regulated propagator in curved spacetime) [112, 12]. The geometric equation integrates to a mass–length profile governed by the sourced $\rho_{\rm vac}(r)$ [40]. Imposing asymptotic flatness fixes integration constants and yields outer data for $\{A, f\}$ that later serve as matching conditions for the near-horizon solution (standard asymptotic matching in semiclassical back-reaction) [141, 45, 112].

Horizon location and first-order shift By definition of the horizon,

$$N^2(r) = 1 - \frac{f(r)}{r} - A(r) = 0 \quad (\text{cf. } \textit{eqs.}\ (16.21) \textit{and}\ (16.78)).$$

Introducing a small parameter ε and writing

$$R_{\rm EH} = R_S + \varepsilon\, \delta R + \mathcal{O}(\varepsilon^2),$$

the chapter derives the model-independent relation

$$\delta R = f_1(R_S) - R_S\, a_1(R_S) \quad (\textit{eq.}\ (16.161)),$$

which implies $R_{\rm EH} < R_S$ when $\rho_{\rm vac} > 0$ (since $f_1(R_S) < 0$ and $a_1(R_S) \geq 0$). Equivalently, the "regime function"

$$\alpha\Big(\xi(R_{\rm EH})\Big) := R_S - R_{\rm EH} = -\,\varepsilon\,\delta R = -\,\varepsilon[f_1(R_S) - R_S a_1(R_S)] > 0 \quad (eq.\ (16.162))$$

quantifies the inward shift and propagates directly to $\kappa_{\rm eff}$ and T_H.

Thermodynamics at $\mathcal{O}(\varepsilon)$. With $A_H = 4\pi R_{\rm EH}^2$ and $R_{\rm EH} = R_S + \varepsilon\,\delta R$, the entropy shift is

$$S = \frac{A_H}{4G\hbar} = \frac{\pi R_S^2}{G\hbar}\Big(1 + 2\varepsilon\,\frac{\delta R}{R_S}\Big) + \mathcal{O}(\varepsilon^2).$$

Thus $\delta R < 0$ leads to a slight decrease relative to the classical area law, with consistent $\mathcal{O}(\varepsilon)$ shifts in $\kappa_{\rm eff}$ and T_H determined by eq. (16.79). These relations provide a direct bridge from the horizon shift to measurable thermodynamic corrections.

Predictions and phenomenology (ringdown & shadow). First-order corrections

$$A(r) = \varepsilon\, a_1(r), \qquad B(r) = \varepsilon\, b_1(r), \qquad f(r) = R_S + \varepsilon\, f_1(r)$$

imprint on near-horizon observables. For quasinormal modes (QNMs),

$$\frac{\delta\omega_{\ell n}}{\omega_{\ell n}} = \mathcal{O}\Big(a_1(R_S),\ \tfrac{f_1(R_S)}{R_S},\ R_S a_1'(R_S),\ R_S f_1'(R_S)\Big),$$

and the photon ring/black-hole shadow experiences $\mathcal{O}(\varepsilon)$ shifts through the corrected lapse N^2. The horizon-value parameter vector

$$\boldsymbol{\theta} = \Big(a_1(R_S),\ f_1(R_S)/R_S,\ R_S a_1'(R_S),\ R_S f_1'(R_S)\Big)$$

organizes the mapping from temporal/geometric profiles to constraints from spectroscopy and imaging.

Rotating extension: Chronogeometric Kerr program. (a) Relational affine structure. The unified tensor

$$\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu} \quad ((16.320))$$

defines the *Relational Levi–Civita derivative* with connection

$$\Gamma^{\lambda}{}_{\mu\nu}[\bar{\Sigma}] = \tfrac{1}{2}\,\bar{\Sigma}^{\lambda\rho}\Big(\partial_\mu\bar{\Sigma}_{\nu\rho} + \partial_\nu\bar{\Sigma}_{\mu\rho} - \partial_\rho\bar{\Sigma}_{\mu\nu}\Big),$$

and the split $\Gamma[\bar{\Sigma}] = \Gamma[\bar{g}] + \Delta\Gamma[\bar{t}]$ with first-order difference (16.323).

(b) Temporal sector ansatz. In Boyer–Lindquist form, the temporal tensor contributes

$$\bar{t}_{\mu\nu}\,dx^\mu dx^\nu = -A\,dt^2 + B\,\frac{\rho^2}{\Delta}\,dr^2 + C\,\rho^2\,d\theta^2 + E\,\sin^2\theta\,d\phi^2 - 2D\,\sin^2\theta\,dt\,d\phi,$$

as in (16.209).

(c) Rotationally dressed vacuum energy. The state-conditioned source is

$$\rho_{\rm vac}(x) := \left\langle \mathring{\mathbf{\Psi}} \right| \hat{\rho}^{(\rm eff)}(x) \left| \mathring{\mathbf{\Psi}} \right\rangle,$$

with $\hat{\rho}^{(\rm eff)} = \hat{\rho}^{(\rm static)} + \hat{\rho}^{(\rm rot)}$; the rotational contribution is a functional of $\hat{J}^{\mu\nu}$ and the chronogeometric vorticity $\hat{\omega}_{\mu\nu}[\bar{\Sigma}]$ built from the relational connection (Definition 16.5 and eq. (16.210)).

(d) Horizon & ergoregion in the unified geometry. With

$$\chi^a = \partial_t + \Omega_H \, \partial_\phi,$$

the ergosurface is defined by $\bar{\Sigma}_{tt} = 0$ ((16.202)); the horizon is the largest root of $\bar{\Sigma}_{ab}\chi^a\chi^b\big|_{r=R_H} = 0$ with

$$\Omega_H = -\frac{\bar{\Sigma}_{t\phi}}{\bar{\Sigma}_{\phi\phi}}\bigg|_{r=R_H} \qquad ((16.203)).$$

Small-spin/weak-dressing limits reduce to eq. (16.204) for (R_H, Ω_H).

(e) Kerr solution pipeline. The computation proceeds from **Relational Christoffels** in Section 16.5 through horizon/regularity and boundary conditions, culminating in the "Chronogeometric Kerr Solution," consistent with black-hole thermodynamics.

Modular/KMS identification on the horizon (zeroth law & temperature). Using Tomita–Takesaki modular theory, a $(\alpha_t, \beta_{\rm KMS})$–KMS state satisfies

$$\hat{K}_\omega = \beta_{\rm KMS}\hat{H}_\chi + c\,\mathbf{1} \quad ((16.481)),$$

so that near a Killing horizon one has $\beta_{\rm KMS} = 2\pi/\kappa$, hence

$$T_H = \frac{\kappa}{2\pi},$$

matching eq. (16.5). Step 4 identifies this with the horizon modular temperature

$$T_{\rm mod}(x) = \frac{\delta\langle \hat{H}_{\rm mod}(x)\rangle}{\delta S(x)} \quad (\textit{Proposition } 16.59),$$

providing the modular underpinning of the zeroth law in the unified setting.

Kerr horizons as observables: ergoregion, shifts, and constraints. To first order, horizon regularity enforces θ-independence of R_H and Ω_H in Kerr-like geometries (standard results for stationary, axisymmetric black holes) [142, 143, 141, 45]. The variations $(\delta R_H, \delta\Omega_H)$ are linear functionals of the temporal-sector data $\{A, B, D, E\}$ evaluated at the horizon, yielding a transparent pipeline from chronogeometric profiles to observable signatures: ringdown frequencies, late-time echoes, and black-hole shadow distortions [141]. This linear structure enables direct inference constraints on the temporal sector from spectroscopy and imaging [122].

Causal structure and CTCs reinterpreted as history transitions. Within IQR, extended Kerr closed timelike curves (CTCs) are reinterpreted as *branch transitions* mediated by a holosymmetric ring condensate. A classical CTC threading the ring corresponds to a nonzero off-diagonal amplitude $\hat{T}_{\beta\alpha}$, a partial isometry

$$\gamma_{\mathrm{CTC}} : \mathcal{H}_\alpha \to \mathcal{H}_\beta, \qquad \beta \neq \alpha,$$

implemented via the holosymmetric portal of Definition 16.23. Thus, "time travel" does not return one to their own past but bridges to a distinct history sector, consistent with the classical existence of CTC regions in Kerr and with general chronology considerations (formalized in eqs. (16.260) and (16.261)) [141, 20].

Dressed-vacuum sourcing (spectral, endogenous, state-conditioned). The effective source $\rho_{\mathrm{vac}}(r)$ entering the unified equations is state-conditioned and band-limited: a spectral integral that combines modal weights, an IR gap, and finite entanglement capacity into a localized, endogenous profile—mirroring semiclassical stress-tensor constructions in curved spacetime. This profile drives A, B and feeds back into f, closing the system at the level of $\bar{\Sigma}_{\mu\nu}$. The refined expressions and their spatial realizations are given in (16.129)–(16.130), ensuring that all near-horizon corrections remain tied to a physical (state-dependent) sourcing mechanism [12, 112].

Summary of testable predictions. The universal first-order conclusion is a slight inward displacement $R_{\mathrm{EH}} < R_S$ ($\delta R < 0$) with concomitant shifts in κ_{eff} and T_H (area/temperature relations and black-hole mechanics) [40]. These propagate into three principal families of observables: (i) ringdown spectroscopy, where

$$\frac{\delta\omega_{\ell n}}{\omega_{\ell n}} = \mathcal{O}\Big(a_1(R_S),\ \tfrac{f_1(R_S)}{R_S},\ R_S a_1'(R_S),\ R_S f_1'(R_S)\Big),$$

and multi-mode tests leverage LIGO/Virgo events [122]; (ii) black-hole shadow and photon-ring deformations controlled by the corrected lapse N^2, confronting imaging and accretion spectroscopy [141]; and (iii) area-law/temperature adjustments consistent with a *unitary* Page curve for a PQS-initialized microstate [138, 139, 42]. The parameter vector $\boldsymbol{\theta}$ at the horizon consolidates these effects into a compact, falsifiable response model that *directly* links the formalism's chronogeometric inputs to data.

Chapter 17: Cosmic Evolution

Chapter 17 develops the cosmic dynamics from a single relational action in which the curvature term and an invariant four-volume term yield Einstein–type field equations with an intrinsic vacuum stress–energy, thereby interpreting the cosmological constant Λ as the energy of the geometric vacuum (Section 17.1) [144, 145, 40, 146, 147]. Quantum-mechanically, Λ is fixed as the ground-state eigenvalue of the Geometric Hamiltonian, $\hat{H}_{\mathrm{geometry}}|\Psi_{\mathrm{vac}}\rangle = \Lambda|\Psi_{\mathrm{vac}}\rangle$ (Definition 17.2), and computed by a regulated spectral integral whose UV/IR bounds are physical, not ad hoc; a dynamical self-regulation between the geometric (*Gaiamiti*) and

material (*Tantrakyma*) sectors then drives a near-cancellation, leaving the small residual we observe as dark energy (Planck $\Omega_\Lambda \simeq 0.685$) (Sections 13.1 and 17.1.2) [112, 12, 148, 149, 150, 151, 152, 153, 154].

The chapter then elevates the *Universal Schrödinger Equation* $i\hbar\partial_t|\mathring{\mathbf{\Psi}}(t)\rangle = \hat{H}_{\rm rel}(t)|\mathring{\mathbf{\Psi}}(t)\rangle$ (eq. (17.47)) as the single dynamical law, with $\hat{H}_{\rm rel} = \hat{H}_{\rm geometry} + \hat{H}_{\rm matter}$ unifying dark energy/dark matter with the Standard Model sector. Crucially, $\hat{H}_{\rm rel}$ *emerges* from the Fourfold Inter-Modal Correlator $\hat{G}_\Lambda(x, x')$: taking its local part, tracing, and integrating over the emergent manifold yields the global energy operator that drives cosmic evolution (Section 6.1.13) [22, 30, 49, 146, 155, 156, 47]. In this limit, familiar relations such as $E = mc^2$ are recovered as theorems of the relational architecture rather than imposed axioms (Section 17.1.6) [32, 20, 146].

Finally, the *Holocosmos Constraint* organizes two complementary cosmic regimes within a single timeless law: an *Ātmapoiesis* branch satisfying $\hat{H}_{\rm univ}|\mathring{\mathbf{\Psi}}\rangle = 0$ (self-contained, zero-total-energy universe), and a *Parapoiesis* branch driven by a transcendent source $\mathfrak{B}_{\vec{r}}$ with a fundamental arrow of time; an entangled holocosmic state cleanly decouples these into paired universal constraints (Sections 17.2.1 and 17.2.2). This synthesis is read as the highest articulation of the Law of Dependent Origination and the Axiom of Duality: the world's becoming and its being are two faces of one relational grammar (Section 17.2) [51, 156, 146, 157, 42, 17, 15, 35, 16].

1) The Law of Dependent Origination and Cosmic Evolution (Section 17.1). The chrono–geometric/IQR dynamics are fixed by a state-first, representation-independent principle together with diffeomorphism covariance, yielding the *Total Relational Action* (eq. (17.1)) [20, 42, 146]:

$$S_{\rm rel} = \frac{1}{2\kappa'}\int \big(\bar{R}[\bar{\Sigma}] - 2\Lambda\big)\sqrt{-\det\bar{\Sigma}}\, d^4x \;+\; S_{\rm matter}[\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}] \;+\; S_{\rm bdy}[\bar{\Sigma}],$$

where κ' is the relationally normalized coupling and Λ is the constant four-volume coefficient. Varying $S_{\rm rel}$ produces the Chronogeometric Einstein Equations with Λ (see eqs. (17.8) and (17.9)) and identifies the geometric (vacuum) stress as $T^{\rm vac}_{\mu\nu} = -(\Lambda/\kappa')\,\bar{\Sigma}_{\mu\nu}$ (eq. (17.10)) [146, 144, 145, 40]. Interferometric vacuum energy ε_Λ (defined through a holosymmetric projector on the vacuum sector) augments the bare constant to an *effective cosmological constant* $\Lambda_{\rm eff} = \Lambda + \kappa'\,\varepsilon_\Lambda$ (eq. (17.5)), with dark-energy density

$$\rho_{\rm DE} = \varepsilon_\Lambda = \frac{1}{{\rm Vol}(\mathcal{X}_q)}\Big\langle \mathring{\mathbf{\Psi}}\Big|\, \mathcal{P}^{\rm sym}_{\rm vac}\, H_{\rm vac}\, \mathcal{P}^{\rm sym}_{\rm vac}\,\Big|\mathring{\mathbf{\Psi}}\Big\rangle_{\rm phys} \quad (eq.\ (17.4))$$

in the operator–algebraic/GNS setting [30, 22], consistent with regulated spectral constructions and EFT control [12, 112, 148, 149].

On the Hamiltonian side, the Legendre transform of $S_{\rm rel}$ together with the global relational balance yields the *Universal Quantum Constraint* $\hat{H}_{\rm univ}\,|\mathring{\mathbf{\Psi}}\rangle = 0$ (eqs. (13.236) and (17.6)), encoding the vanishing of total relational energy (geometry + matter/coherence) [156, 146, 51].

2) The Formal Origin of the Cosmological Constant (Section 17.1.2). Writing the geometric sector as

$$S_{\rm geom}[\bar{\Sigma}] = \frac{1}{2\kappa'}\int_{\mathcal{M}}\big(\bar{R}[\bar{\Sigma}] - 2\Lambda\big)\,{\rm d}\mu_{\bar{\Sigma}} \;+\; S_{\rm bdy}[\bar{\Sigma}], \qquad {\rm d}\mu_{\bar{\Sigma}} = \sqrt{-\det\bar{\Sigma}}\,{\rm d}^4x \quad (eq.\ (17.7)),$$

its variation gives

$$\delta S_{\text{geom}} = \frac{1}{2\kappa'}\int \sqrt{-\det \bar{\Sigma}}\,\left(\bar{G}_{\mu\nu} + \Lambda\,\bar{\Sigma}_{\mu\nu}\right)\delta\bar{\Sigma}^{\mu\nu}\,\mathrm{d}^4x \;+\; \delta S_{\text{bdy}} \quad (eq.\ (17.8)),$$

and, including matter, the Chronogeometric Einstein Equations

$$\bar{R}_{\mu\nu} - \tfrac{1}{2}\,\bar{\Sigma}_{\mu\nu}\,\bar{R} + \Lambda\,\bar{\Sigma}_{\mu\nu} = \kappa'\,T^{\text{matter}}_{\mu\nu} \quad (eq.\ (17.9)).$$

Because $T^{\text{vac}}_{\mu\nu} \propto \bar{\Sigma}_{\mu\nu}$ is geometrically *homogeneous* (eq. (17.10)), it drops out of null-directed energy conditions and is coherently placed on the geometric side of the response kernel [146, 144, 145, 40, 43].

3) The Quantum–Mechanical Nature of Λ and Self-Regulation (Section 17.1.2). At the operator level the geometric Hamiltonian admits a density-normalized vacuum,

$$\hat{H}_{\text{geometry}}\,|\Psi_{\text{vac}}\rangle = \Lambda_{\text{geo}}\,|\Psi_{\text{vac}}\rangle\,, \qquad \Lambda_{\text{geo}} = \frac{\langle\Psi_{\text{vac}}|\hat{H}_{\text{geometry}}|\Psi_{\text{vac}}\rangle_{\text{phys}}}{\text{Vol}(\mathcal{X}_q)} \quad (eq.\ (17.11)),$$

and endogenous (state-conditioned) regularization—via modal weights w_n, the *Principle of Finite Entanglement Capacity*, and a state-emergent cutoff—renders the spectrum finite (see Section 17.1.2) [12, 112, 148, 20, 147, 47, 49].

4) Physical Manifestation as Dark Energy and the Foundational Definition (Section 17.1.2). On cosmological scales,

$$\Lambda_{\text{eff}} = \Lambda + \kappa'\,\varepsilon_\Lambda, \qquad \rho_{\text{DE}} = \Lambda_{\text{eff}}/\kappa' \quad (eq.\ (17.12)),$$

implying an effective geometric vacuum stress $T^{\text{vac,eff}}_{\mu\nu} = -(\Lambda_{\text{eff}}/\kappa')\,\bar{\Sigma}_{\mu\nu}$ (cf. eq. (17.10)). The *Foundational Definition* (Definition 17.2) unifies: (i) Λ from the variational action, (ii) Λ_{geo} from $\hat{H}_{\text{geometry}}$, and (iii) Λ_{eff} as the phenomenological driver of late-time acceleration.

5) Calculating Λ: Regulated Spectral Integral (Section 17.1.2). In local trace form,

$$\hat{H}_{\text{geometry}} = \int_{\mathcal{X}_q} d^dx\,\sqrt{-\det\bar{\Sigma}(x)}\;\text{Tr}\big[\hat{G}_{\mathbf{\Phi\Phi}}(x,x)\big] \quad (eq.\ (17.13)),$$

leading (quasi-homogeneously) to

$$\Lambda_{\text{geo}} \propto \frac{1}{\text{Vol}(\mathcal{X}_q)}\left\langle \overset{\circ}{\mathbf{\Psi}}\middle|\hat{H}_{\text{geometry}}\middle|\overset{\circ}{\mathbf{\Psi}}\right\rangle_{\text{phys}} \simeq \int_{k_{\text{IR}}}^{k_{\text{UV}}}\frac{d^3k}{(2\pi)^3}\,E(k), \quad E(k) = \tfrac{1}{2}\hbar\,\omega_k + \cdots \quad (eq.\ (17.14)).$$

Self-regulation then leaves a *small positive residual*

$$\rho_{\text{DE}} = \varepsilon_\Lambda = \frac{1}{\text{Vol}(\mathcal{X}_q)}\left|\langle\hat{H}_{\text{geometry}}\rangle_{\text{phys}} + \langle\hat{H}_{\text{vac}}\rangle_{\text{phys}}\right| \ll \frac{k^4_{\text{UV}}}{16\pi^2}\hbar c \quad (eq.\ (17.15)).$$

6) Deriving Λ_{eff} in the Field Equations (Section 17.1.3). Starting from

$$S_{\text{geom}}[\bar{\Sigma}] = \frac{1}{2\kappa'}\int \big(\bar{R}[\bar{\Sigma}] - 2\Lambda\big)\sqrt{-\det \bar{\Sigma}}\, d^4x \quad (eq.\ (17.17)),$$

the bare-Λ equations read

$$\bar{R}_{\mu\nu} - \tfrac{1}{2}\,\bar{\Sigma}_{\mu\nu}\,\bar{R} + \Lambda\,\bar{\Sigma}_{\mu\nu} = \kappa'\,\bar{T}^{\text{matter}}_{\mu\nu} \quad (eq.\ (17.18)).$$

Splitting $\bar{T}^{\text{matter}}_{\mu\nu} = \bar{T}^{(\text{nonvac})}_{\mu\nu} + \bar{T}^{(\text{vac})}_{\mu\nu}$ with $\bar{T}^{(\text{vac})}_{\mu\nu} = -\varepsilon_\Lambda\,\bar{\Sigma}_{\mu\nu}$ (eqs. (17.19) and (17.20)) yields

$$\bar{R}_{\mu\nu} - \tfrac{1}{2}\,\bar{\Sigma}_{\mu\nu}\,\bar{R} + \underbrace{\big(\Lambda + \kappa'\varepsilon_\Lambda\big)}_{:=\Lambda_{\text{eff}}}\bar{\Sigma}_{\mu\nu} = \kappa'\,\bar{T}^{(\text{nonvac})}_{\mu\nu} \quad (eq.\ (17.21)),$$

i.e. the vacuum residue is coherently moved to the geometric side.

7) Measured Value and Bridge to IQR. Observations (Planck 2018 + external data) give $\Omega_\Lambda \simeq 0.685$ (eq. (17.42)) and $\rho^E_{\text{crit}} = 3c^2H_0^2/(8\pi G)$ (eq. (17.43)), hence

$$\rho_{\text{vac}} = \Omega_\Lambda\,\rho^E_{\text{crit}} \approx (5.3 \pm 0.3)\times 10^{-10}\ \text{J m}^{-3} \quad (eq.\ (17.44)),$$

and

$$\Lambda_{\text{eff}} = \frac{8\pi G}{c^4}\rho_{\text{vac}} \approx (1.1 \pm 0.1)\times 10^{-52}\ \text{m}^{-2} = \frac{3\,\Omega_\Lambda H_0^2}{c^2} \quad (eq.\ (17.45)).$$

In the present notation this is precisely $\Lambda_{\text{eff}} = \Lambda + \kappa'\,\varepsilon_\Lambda$ (eq. (17.46)).

8) The Universal Schrödinger Equation and $E = mc^2$ (Sections 17.1.5 and 17.1.6). Cosmic evolution in the physical sector is governed by

$$\boxed{i\hbar\frac{d}{dt}|\mathring{\mathbf{\Psi}}(t)\rangle = \hat{H}_{\text{rel}}(t)\,|\mathring{\mathbf{\Psi}}(t)\rangle} \quad (eq.\ (17.47))[22,\ 30].$$

with decomposition

$$\hat{H}_{\text{rel}} = \underbrace{\hat{H}_{\text{geometry}}[\hat{\Sigma}_{\mu\nu}]}_{\text{Dark Energy \& Dark Matter}} + \underbrace{\hat{H}_{\text{matter}}[\hat{\phi},\hat{\psi},\ldots]}_{\text{Standard Model}} \quad (eq.\ (17.48))[20,\ 42].$$

Within this framework, mass–energy equivalence $E = mc^2$ emerges through the chronogeometric couplings that identify inertial and energetic content (see Section 17.1.6) [32, 146].

9) The Law of Dependent Origination and Conservation of Duality (Section 17.1.7). At the deepest level the dynamics reduce to the timeless balance

$$\hat{H}_{\text{univ}}|\mathring{\mathbf{\Psi}}\rangle = 0 \quad (Section\ 17.1.7),$$

a background-independent statement of total relational conservation (geometry + sources), with "time" emerging relationally [156, 51, 146, 20].

10) The Unfolding/Enfolding and the Ātmapoiesis Universe (Sections 17.2 and 17.2.1). The macro-narrative of cosmic *unfolding* (phenomenal becoming) and *enfolding* (ontic coherence) is formalized as

$$\boxed{\hat{H}_{\text{univ}}|\overset{\circ}{\boldsymbol{\Psi}}\rangle = 0} \quad (eq.\ (17.49)),$$

expressing the self-consistency of the whole as the source of the parts in the Ātmapoiesis Universe [156, 51, 146, 42, 17, 15, 35, 16].

11) The Holocosmos: Entangled Dual Branches and Constraint Expansion (Section 17.3.1 and eqs. (17.59) and (17.60)). The holocosmos Hilbert space factorizes as $\mathcal{H}_{\text{holocosmos}} = \mathcal{H}_{\text{univ}} \otimes \mathcal{H}_{\mathfrak{B}}$, and the timeless entangled state

$$|\boldsymbol{\Omega}\rangle = \alpha\,|\overset{\circ}{\boldsymbol{\Psi}}_{\bar{\text{A}}\text{tmapoiesis}}\rangle \otimes |\bar{\text{A}}\text{tmapoiesis}\rangle + \beta\,|\overset{\circ}{\boldsymbol{\Psi}}_{\text{Parapoiesis}}\rangle \otimes |\text{Parapoiesis}\rangle \quad (eq.\ (17.61)),$$

obeys $|\alpha|^2 + |\beta|^2 = 1$ (eq. (17.62)). Coherence is enforced by the compact *holocosmos constraint*

$$\left(\hat{H}_{\text{univ}} \otimes \mathbb{I}_{\mathfrak{B}} - \mathbb{I}_{\text{univ}} \otimes \hat{\mathfrak{B}}\right)|\boldsymbol{\Omega}\rangle = 0 \quad (eq.\ (17.63)),$$

whose explicit expansion (eq. (17.59)) decouples to

$$\hat{H}_{\text{univ}}|\overset{\circ}{\boldsymbol{\Psi}}_{\bar{\text{A}}\text{tmapoiesis}}\rangle = 0, \qquad \hat{H}_{\text{univ}}|\overset{\circ}{\boldsymbol{\Psi}}_{\text{Parapoiesis}}\rangle = \mathfrak{B}_{\vec{r}}|\overset{\circ}{\boldsymbol{\Psi}}_{\text{Parapoiesis}}\rangle \quad (eq.\ (17.60)),$$

exhibiting a stationary branch and a sourced branch with c-number $\mathfrak{B}_{\vec{r}}$. The *Axiom of Holocosmic Duality* (Section 17.3.1) assigns ontological Born weights

$$|\alpha|^2 = v(\rho_\Omega; \Pi_A), \qquad |\beta|^2 = v(\rho_\Omega; \Pi_P), \quad |\alpha|^2 + |\beta|^2 = 1 \quad (eqs.\ (17.73) and\ (17.74)),$$

ensuring conservation of total ontic content.

12) The Law of Dependent Origination of the Holocosmos: Action Forms and Equivalences (Section 17.3.2). Let $A := \hat{H}_{\text{univ}} \otimes \mathbb{I}_{\mathfrak{B}} - \mathbb{I}_{\text{univ}} \otimes \hat{\mathfrak{B}}$. Physical holocosmic states lie in the kernel $A\,|\boldsymbol{\Omega}\rangle = 0$ (eq. (17.67)). Three equivalent variational forms enforce this: (1) Rayleigh–Ritz with norm constraint,

$$\overset{\circ}{\boldsymbol{\Lambda}}[\,\boldsymbol{\Omega}\,] = \langle\boldsymbol{\Omega}|A|\boldsymbol{\Omega}\rangle - \lambda\Big(\langle\boldsymbol{\Omega}|\boldsymbol{\Omega}\rangle - 1\Big) \;\Rightarrow\; A|\boldsymbol{\Omega}\rangle = \lambda|\boldsymbol{\Omega}\rangle,\ \lambda = 0 \quad (eqs.\ (17.68) and\ (17.69));$$

(2) the penalty functional $\boldsymbol{\Lambda}_{\text{pen}}[\,\boldsymbol{\Omega}\,] = \langle\boldsymbol{\Omega}|A^2|\boldsymbol{\Omega}\rangle - \lambda(\langle\boldsymbol{\Omega}|\boldsymbol{\Omega}\rangle - 1)$, whose minimum selects $\ker A$; and (3) the operator (vector–multiplier) constraint

$$\boldsymbol{\Lambda}_{\text{op}}[\,\boldsymbol{\Omega}, \chi\,] = \tfrac{1}{2}\Big(\langle\chi|A|\boldsymbol{\Omega}\rangle + \langle\boldsymbol{\Omega}|A|\chi\rangle\Big) - \lambda(\langle\boldsymbol{\Omega}|\boldsymbol{\Omega}\rangle - 1),$$

whose stationarity likewise enforces $A|\boldsymbol{\Omega}\rangle = 0$ (eq. (17.72)). The *Equivalence Proposition* (Proposition 17.24) proves that—under self-adjointness and $\ker A \neq \{0\}$—all three yield the same normalized physical solution set.

13) Concluding Synthesis (Section 17.3.4). Chapter 17 welds the *formal* (Λ from action), *quantum* (Λ_{geo} from $\hat{H}_{\text{geometry}}$), and *phenomenological* (Λ_{eff}) layers into one relational cosmology: evolution is governed by the Schrödinger form (eq. (17.47)) or, more fundamentally, by the timeless constraint (Section 17.1.7) [22, 30, 156, 51, 146, 20]. The holocosmos construction (eqs. (17.61) and (17.63)) encodes dual branches with ontological weights (eq. (17.73)) [158, 42], while the regulated spectral machinery (eqs. (17.13) and (17.14)) and coherent placement of vacuum stress (cf. eq. (17.21)) deliver a finite, positive ρ_{DE} (eq. (17.15)) consistent with observation (eqs. (17.42) and (17.45)) [112, 12, 148, 145, 144, 149, 150, 151, 152, 153, 154].

Prologue: Entanglement at the Foundations of Physics

The Inner Tangle and the Outer Tangle of Physics

From Clockwork Certainty to Quantum Hurricane At the close of the 19th century, the edifice of physics seemed nearly complete, resting upon the bedrock of Sir Isaac Newton's classical mechanics [159]. For over two centuries, Newton's laws had provided a deterministic, "clockwork" universe where reality was objective and particles followed predictable trajectories. This confidence was famously captured by Lord Kelvin in 1900 when he declared the beauty of the classical framework, noting only two minor "clouds on the horizon" that needed resolving [160]. These "clouds," however, were not minor atmospheric disturbances but the harbingers of a hurricane that would fundamentally reshape our understanding of reality. They pointed toward puzzling phenomena that classical physics could not explain, cracking the foundations of its deterministic certainty and ushering in the strange and subversive realm of quantum mechanics [161].

The Tremors in the Classical World The first true tremor came not from a new experiment but from a theoretical insight into one of Kelvin's clouds: the problem of *black-body radiation*. To explain the observed spectrum of light emitted by a hot object, German physicist *Max Planck* reluctantly proposed in 1900 that energy was not continuous but was emitted and absorbed in discrete packets, or "quanta" [162]. He saw this as a mathematical trick, yet he had opened Pandora's box. A few years later, *Albert Einstein*, then a young patent clerk, took Planck's idea seriously. To explain the *photoelectric effect*—a phenomenon discovered by Heinrich Hertz [163], where light could knock electrons off a metal surface—Einstein proposed in his "miraculous year" of 1905 that light itself is composed of these discrete particles, which he called "light quanta" and we now call *photons* [164]. This radical idea, for which he would later win the Nobel Prize, showed that light, long understood as a wave, also behaved as a particle.

This notion of *wave-particle duality* was the heart of the emerging quantum crisis. The most profound and unsettling demonstration of this duality came from reconsidering an old experiment with new entities. In the early 1800s, *Thomas Young* had conclusively shown that light was a wave by passing it through two narrow slits and observing an *interference pattern*—a characteristic pattern of alternating bright and dark fringes that could only be produced by waves interfering with each other [165]. A century later, inspired by the bold 1924 hypothesis from French physicist *Louis de Broglie* that all matter exhibits wave-like

properties [166], physicists performed the double-slit experiment with particles: *electrons*.

The results were paradigm-shattering. When a beam of electrons was fired at a barrier with two open slits, they did not simply create two bands on the detector screen as tiny pellets would. Instead, they produced the exact same interference pattern that Young had seen with light waves (demonstrated experimentally by *Davisson and Germer* in 1927 [167]). This forced a conclusion that defied all classical intuition: each individual electron, a supposed particle, seemed to pass through *both* slits simultaneously like a wave and interfere with itself. Even more bizarrely, if a detector was placed at the slits to see which path the electron took, the interference pattern vanished, and the electrons behaved like normal particles again. The very act of observing the particle's path forced it to "choose" a state, dissolving its wave-like nature. This experiment exposed the core quantum mystery: reality at the subatomic level is inherently probabilistic, and the observer is inextricably linked with the observed [168, 169].

The Copenhagen Interpretation: Probability, Collapse, and Complementarity The baffling results of the double-slit experiment demanded a new logic for reality, a logic codified in the late 1920s into what became the orthodox *Copenhagen interpretation*, primarily championed by *Niels Bohr* and *Werner Heisenberg* [168, 169]. According to this view, an electron is not a simple particle but is described by a *wave function*: a mathematical entity that represents a superposition of all its possible states. This wave, however, is not a physical substance but a wave of *probability*. This crucial insight came from *Max Born* in 1926, who proposed that the square of the wave function's amplitude at any given point corresponds to the probability of finding the particle there upon measurement [21].

Thus, when an unobserved electron travels toward the slits, its wave function passes through both, allowing it to interfere with itself. The interference pattern vanishes upon observation because of what the Copenhagen school called the *"collapse of the wave function."* The moment an interaction occurs that can yield information about the electron's path—for instance, a photon bouncing off it—the superposition of possibilities is destroyed. The spread-out wave of potential locations instantly "collapses" into a single, definite state at a specific point. As *Heisenberg* formalized in his famous 1927 uncertainty principle paper, the act of measuring one property (like position) inevitably disturbs another (like momentum), making it impossible to know both with perfect accuracy [54]. This isn't a limitation of our instruments; it's a fundamental feature of nature.

The Observer and the Observed It's crucial to understand that the "observer" in quantum mechanics need not be a conscious mind. Rather, it is any physical interaction with the environment that "records" the state of the system, a process now understood through the lens of *quantum decoherence* [170]. This interaction irrevocably entangles the particle with the measuring device, destroying the delicate superposition required for wave-like interference.

This insight strikes at the very heart of classical reality. It suggests that properties like an electron's position do not exist in a definite state until they are measured. This idea was expressed most clearly in *Niels Bohr's principle of complementarity*. Bohr argued that wave-like and particle-like behavior are mutually exclusive, or complementary, aspects of a single underlying reality. An experimental setup designed to reveal the particle's path will necessarily destroy the conditions needed to observe its wave-like interference, and

vice-versa [168]. We can never see both faces of the quantum coin at the same time. This led *Heisenberg* to his profound philosophical conclusion, articulated in *Physics and Philosophy*: "What we observe is not nature itself, but nature exposed to our method of questioning" [169]. The world "out there" is not a static stage to be passively viewed; it is a participatory reality where the act of measurement helps co-create the very properties being measured.

Indeterminacy and the Limits of Knowledge The philosophical upheaval ignited by quantum mechanics is powerfully encapsulated in *Werner Heisenberg's 1927 uncertainty principle* [54]. More than a statement about measurement limitations, the principle reveals a fundamental "fuzziness" inherent in nature. It states that certain pairs of properties, known as conjugate variables (like a particle's position and its momentum), cannot be simultaneously known to arbitrary precision. The more precisely one property is determined, the more uncertain the other becomes, governed by the famous inequality $\Delta x \Delta p \geq \hbar/2$. This is not because our instruments are clumsy; it is a law of nature. An electron, prior to measurement, simply *does not possess* a definite position and a definite momentum at the same time. Reality at this scale is partly indeterminate, a haze of potentialities waiting for an observation to force it into a concrete state.

Schrödinger's Cat Paradox The bizarre implications of this quantum indeterminacy being applied to the macroscopic world were famously dramatized by *Erwin Schrödinger* in 1935. Deeply troubled by the Copenhagen interpretation's concept of superposition, Schrödinger devised a thought experiment intended as a *reductio ad absurdum*—a critique to expose the theory's apparent absurdity. In his paper *"The Present Situation in Quantum Mechanics"*, he imagined a cat sealed in a box along with a vial of poison, a hammer, a Geiger counter, and a single radioactive atom [171, 172, 173].

If the atom decays (a fundamentally random quantum event), the Geiger counter will trigger the hammer, which will shatter the vial and kill the cat.

According to the quantum formalism, as long as the box remains sealed, the atom is in a superposition of two states: "decayed" and "not decayed." By extension, the entire system linked to it must also be in a superposition. The cat, therefore, is described by a wave function that represents it as being simultaneously *alive and dead*. This macabre state persists until a measurement is made—that is, until an observer opens the box. At that moment, the wave function "collapses", and a single, definite reality is revealed: either a living cat or a dead one. Schrödinger's paradox was not designed to suggest cats could actually be both alive and dead; it was designed to highlight the unresolved *measurement problem* [174, 175]. At what precise moment does the "fuzzy" quantum superposition collapse into the single, concrete reality we experience? Does it happen when the atom decays, when the Geiger counter clicks, or only when a conscious being intervenes? The paradox masterfully illustrates the conceptual chasm between the quantum world of potentialities and the classical world of definite facts, a chasm that remains a subject of intense debate to this day.

The Bohr-Einstein Debates: A Clash Over the Nature of Reality The foundational weirdness of quantum mechanics was so profound that it led to one of the greatest intellectual duels of the 20th century: a decades-long debate between Albert Einstein and Niels Bohr

over the very soul of physics. This was not a disagreement over equations but a clash of fundamental philosophies about what reality is and what science ought to describe.

Einstein's Realist Dissent Albert Einstein, whose work on the photoelectric effect was a crucial catalyst for the quantum revolution, became its most formidable critic. He could not accept a reality governed by what he saw as the incomplete and probabilistic nature of the new mechanics. His profound discomfort is immortalized in his 1926 letter to Max Born, where he famously declared, *"I, at any rate, am convinced that He [God] does not play dice"* [176]. For Einstein, a *realist*, the universe must possess definite, objective properties whether we are observing them or not. He famously challenged physicist Abraham Pais, "Do you really believe the moon exists only when you look at it?" mocking the idea that observation creates reality [177].

Einstein's core argument was not that quantum mechanics was wrong, but that it was *incomplete*. He believed that the theory's probabilistic nature was merely a reflection of our ignorance of a deeper, deterministic level of reality governed by as-yet-unknown "hidden variables." His most powerful challenge came in the form of the *1935 EPR paradox*, a thought experiment co-authored with Boris Podolsky and Nathan Rosen. The paper argued that quantum mechanics' description of entangled particles—which remain mysteriously connected over vast distances—implied a "spooky action at a distance" that, to Einstein, proved the theory must be missing something fundamental about the nature of local reality [178].

Bohr's Copenhagen Orthodoxy In the opposing corner stood *Niels Bohr*, the intellectual architect of the dominant *Copenhagen interpretation*. Bohr met Einstein's challenge with a radical philosophical shift, insisting that we must abandon our classical intuitions. For Bohr, the "spooky" features of quantum mechanics were not a sign of incompleteness but a revelation about the nature of knowledge itself. He argued that properties like position or momentum have no meaning independent of an experimental context designed to measure them [168, 179].

Bohr's response to Einstein was to double down on his *principle of complementarity*. An experimental setup defines what can be known. Any attempt to describe a quantum system's reality outside of this observational context was, in Bohr's view, meaningless language [180]. He famously articulated this stance by stating, *"There is no quantum world. There is only an abstract quantum physical description. It is wrong to think that the task of physics is to find out how nature is. Physics concerns what we can say about nature"* (as quoted by Petersen [181]). This startling claim redefines the goal of science: it is not to uncover a pre-existing, objective reality, but to build a consistent framework for what we can measure and communicate about our interactions with the world [182]. For Bohr, Einstein's question about what the particles were 'really" doing was a violation of the rules of the new quantum game.

The EPR Paradox: Einstein's Last Stand for Local Reality The simmering debate between Einstein and Bohr reached its theoretical zenith in 1935 with the publication of the *Einstein-Podolsky-Rosen (EPR) paradox* [178, 182]. This was Einstein's most sophisticated attempt to prove that quantum mechanics, while correct in its predictions, must be an *incomplete*

description of reality. The authors imagined two particles created from a single source, flying apart in opposite directions. By the laws of quantum mechanics, these particles could be *"entangled"*, meaning their properties remained linked in an inseparable wave function, no matter how far apart they traveled [171, 172, 173].

The EPR argument was brilliant: by measuring the momentum of particle A, one could instantly know the momentum of particle B without ever touching it. Alternatively, by measuring the position of particle A, one could know the position of particle B. According to their *"criterion of reality"*, if a property can be known without disturbing a system, it must correspond to a real element of that system. Therefore, particle B must have had both a definite position and a definite momentum all along. Since the uncertainty principle forbids a quantum state from having both, EPR concluded that the quantum description was missing information. This implied the existence of *"hidden variables"*—local properties of the particles that predetermined the measurement outcomes [183]. To Einstein, the alternative—that a measurement on particle A instantaneously affected particle B—was an absurdity he derisively called *"spooky action at a distance"* (*spukhafte Fernwirkung*), as it seemed to violate the cosmic speed limit set by light [184].

From Paradox to Proof: Bell's Theorem and Aspect's Experiments For nearly three decades, the EPR paradox remained a philosophical puzzle. The deadlock was finally broken in 1964 by the Irish physicist *John Stewart Bell*. In a groundbreaking paper, Bell demonstrated that if Einstein's hidden variables existed and obeyed the principle of locality, then the correlations between measurements on entangled particles could not exceed a certain mathematical limit. However, the standard predictions of quantum mechanics *did* exceed this limit. *Bell's theorem* thus transformed a philosophical debate into a physically testable question: the universe could be locally real (as Einstein hoped), or it could obey the predictions of quantum mechanics, but not both [183].

The definitive experiments were carried out in the early 1980s by a team led by French physicist *Alain Aspect*. By generating and measuring thousands of entangled photons, Aspect and his colleagues showed a clear and robust violation of Bell's inequality, with results that matched the predictions of quantum mechanics perfectly [184]. These experiments, since replicated with ever-increasing precision, effectively ruled out Einstein's worldview. Nature, at its deepest level, is either non-local (permitting a form of instantaneous connection) or non-realist (meaning particles do not have definite properties before measurement), or both. Einstein's cherished principle of *local realism* was not a feature of our universe.

The Enduring Quantum Conundrum TThe experimental vindication of quantum mechanics leaves us in a universe profoundly at odds with our intuition. The core tenets of this 'quantum crisis" can be summarized plainly. Firstly, *wave-particle duality* is a fundamental feature of reality; a quantum object is neither a wave nor a particle but a new kind of entity that reveals one aspect or the other depending on the experimental question asked [167, 168, 169]. Secondly, nature is irreducibly *probabilistic*. Einstein's insistence that 'God does not play dice" has been answered by experiment [176, 177]: chance is woven into the fabric of the cosmos. The outcome of a single quantum event cannot be predicted with certainty, only its probability [21, 54]. This is not a failure of our knowledge, but a fundamental principle of the

universe itself, representing a radical and permanent break from the deterministic clockwork of classical physics [169, 185].

Measurement as Creation The quantum revolution fundamentally redefined the act of measurement itself. In classical physics, to measure something is to passively reveal a pre-existing property. In quantum mechanics, measurement is an active process of creation. This radical idea was forcefully articulated by *Pascual Jordan*, a key collaborator in the development of matrix mechanics, who stated: *"Observations not only disturb what is to be measured, they produce it... We compel [an] electron to assume a definite position... We ourselves produce the results of measurement"* (as quoted in Jammer [186]). This is the staggering conclusion of the Copenhagen interpretation: an unobserved electron does not *have* a definite position; it exists in a haze of potentialities described by its wave function. The act of measurement forces this potentiality into a single actuality, creating the very property it seeks to find.

Entanglement and the End of Separability Nowhere is the break from classical intuition more severe than with *quantum entanglement*. This phenomenon, which *Erwin Schrödinger* called *the characteristic trait of quantum mechanics*, describes a situation where two or more particles become linked in a single, inseparable state, regardless of the distance separating them [187]. If two particles are entangled, a measurement on one—determining its spin, for example—instantaneously defines the corresponding property of the other.

This demolishes the classical notion of *separability*—the idea that spatially distinct objects have their own independent, well-defined states. Entangled particles do not have individual states; they exist as a single, unified system until a measurement is performed. The relationship between them is, in a sense, more real than the properties of the individual parts. It suggests a profoundly interconnected or *non-local* reality that transcends our everyday experience of space, a conclusion experimentally verified by tests of Bell's theorem [183].

The Enduring Mystery: A "Philosophical Quagmire" The operational success of quantum mechanics is undeniable, yet its implications for the nature of reality remain deeply unsettling, even to its pioneers. As the celebrated physicist *Richard Feynman* candidly admitted, *"I think I can safely say that nobody understands quantum mechanics"* [185]. The theory works perfectly as a predictive tool, but the philosophical questions it raises have led to a "quagmire" of competing interpretations, each trying to make sense of the measurement problem. Among the leading contenders are:

1. The *Copenhagen Interpretation (Bohr/Heisenberg)*, which takes a pragmatic (or anti-realist) view, suggesting that questions about an "unobserved reality" are meaningless and that physics should only concern itself with the outcomes of experiments.

2. The *Many-Worlds Interpretation*, first proposed by *Hugh Everett III* in 1957. It posits that the wave function never collapses. Instead, every possible outcome of a quantum measurement actually occurs, each in a separate, branching universe. In this view, Schrödinger's cat is both alive *and* dead, but in different, equally real, worlds [188].

3. *Hidden-Variable Theories*, most famously developed by *David Bohm*. These theories restore realism and determinism, suggesting that particles always have definite properties, but they do so at the cost of embracing explicit non-locality [189].

These interpretations, among others, highlight that while we know *how* to use quantum mechanics with astounding accuracy, the debate over *what it means* for reality remains one of the most vibrant and profound frontiers in all of science.

The Quantum Conundrum: A New Reality The quantum revolution was more than a series of discoveries; it was a fundamental rupture in the Western scientific worldview. In probing the foundations of matter, physics found that reality dissolves into something ghostly and elusive. The solid, predictable objects of classical physics vanished, replaced by abstract **wave functions** governing probabilities. Key tenets of the old world were systematically dismantled: energy was found to be quantized [190], particles exhibited wave-like properties [191], and nature at its core was found to be irreducibly probabilistic [192].

Perhaps most profoundly, quantum mechanics upended the **Cartesian split** between the objective world and the subjective observer, a division that had underpinned science since the Enlightenment. As **Niels Bohr** argued, in any quantum experiment, the observer and the observed form an indivisible, inseparable whole [179, 193, 180, 182]. The properties of a quantum system are only defined in the context of the specific apparatus chosen to measure them. This is not a technical inconvenience but a deep truth about a participatory reality where, as **Pascual Jordan** noted, observation doesn't just disturb, it *creates* [194, 186].

A House Divided: The Interpretational Schism Faced with this crisis, the pioneers of physics splintered into competing philosophical camps, a division that persists to this day. The different reactions represent distinct attempts to build a coherent worldview from the strange new rules.

- *The Copenhagen Orthodoxy (Bohr & Heisenberg):* For decades, the dominant view was to accept the weirdness as fundamental. Bohr and his allies essentially argued that our classical intuitions are misleading and that physics should renounce the goal of describing an observer-independent "reality." [168, 180, 179, 169] Its task is simply to provide a consistent framework for predicting the outcomes of measurements.

- *The Realist Resistance (Einstein, de Broglie & Bohm):* Albert Einstein led the resistance, convinced that quantum theory was an incomplete, statistical approximation of a deeper, deterministic reality [178]. While Einstein's own "hidden variables" approach was eventually ruled out by experiment, his realist spirit was carried on by others. *Louis de Broglie* briefly proposed a "pilot-wave" theory in 1927 [191], an idea that was independently rediscovered and fully developed by American physicist *David Bohm* in 1952 [189]. *Bohmian mechanics* posits that particles have definite positions at all times but are guided by a real quantum wave, thus preserving realism at the cost of embracing explicit non-locality.

- *The Radical Alternatives (Everett & Wigner):* Other physicists proposed even more radical solutions. In his 1957 doctoral thesis, *Hugh Everett III* formulated the *Many-Worlds Interpretation*, which claims the wave function never collapses [188]. At the

more speculative end of the spectrum, mathematical physicist *John von Neumann* and Nobel laureate *Eugene Wigner* entertained the idea that the "collapse" is unique and is triggered by the intervention of a *conscious observer*, suggesting a special role for the mind in actualizing physical reality [195].

This enduring schism at the heart of quantum mechanics transcends historical significance to constitute a fundamental rupture in our comprehension of reality itself [186, 174, 180]. The philosophical confrontations between Bohr, Einstein, and their contemporaries were not conclusive battles but opening salvos that exposed a vast terrain of ever-deepening enigmas [182, 176]. Far from representing a resolved chapter in scientific history, these debates unveiled a foundational crisis whose reverberations permeate every frontier of contemporary theoretical physics [196].

The profound discord between quantum theory's extraordinary predictive success and the ambiguity shrouding its ontological implications casts long shadows across our attempts to unify physics [195, 183, 184]. This tension between mathematical formalism and conceptual understanding infiltrates our efforts to reconcile quantum principles with gravitation, to decode the nature of time and entropy, to illuminate cosmological origins, and to unveil the universe's hidden constituents [197, 19, 29]. The classical dichotomies—determinism against chance, locality against entanglement, participatory against objective reality—resurface as spectral presences within the great unsolved problems of our era [179, 169].

What emerges from this inheritance is a constellation of ten fundamental challenges, each representing a distinct manifestation of this deeper crisis. These modern "clouds on the horizon" demand not incremental refinement of existing frameworks but a radical reconceptualization of the architectural foundations upon which physics rests. They beckon us toward what may constitute the next profound scientific and philosophical revolution, one that must confront the unresolved legacy of the quantum upheaval that began a century ago [185, 174, 196]. These fundamental challenges are presented below:

1. The Riddle of Quantum Gravity At the heart of theoretical physics lies a deep chasm between its two greatest triumphs: Einstein's General Relativity, which describes gravity as the elegant curvature of a smooth spacetime, and Quantum Mechanics, which governs the microscopic world through probabilistic fields. The two frameworks are fundamentally incompatible. While quantum field theory successfully describes the electromagnetic, weak, and strong forces, gravity stubbornly resists quantization through conventional methods, leading to nonsensical, infinite results. This quest for a unified theory of *quantum gravity* is a necessary step to understand the universe's most extreme environments, such as the singularity within a black hole or the very first moment of the Big Bang. This challenge has spawned a rich landscape of theoretical approaches, from leading research programs like *string theory*, which posits that fundamental particles are vibrations of tiny strings [198, 199, 200, 201], to *loop quantum gravity*, which suggests spacetime itself is composed of discrete, quantized chunks [8, 3]. More exotic proposals include *causal dynamical triangulations* [202] and *emergent spacetime* frameworks, which suggest that geometry itself arises from the entanglement of quantum information [1, 2].

2. The Ghost in the Quantum Machine Despite its spectacular predictive success, quantum mechanics harbors a deep interpretational enigma that has persisted since its inception: the *measurement problem*. This problem asks what it truly means to observe a quantum system, a process that seemingly forces a particle to abandon a ghostly superposition of multiple states and commit to a single, definite outcome through the *"collapse"* of the wavefunction. Whether the wavefunction—the mathematical object describing this superposition—is a real physical entity or merely a statement of our knowledge remains a subject of fierce debate, an ambiguity that strikes at the core of our understanding of reality itself [203]. Various interpretations compete for ontological primacy, each with profoundly different implications for the nature of reality and observation. These frameworks range from the *Many-Worlds Interpretation*, which posits that all possible outcomes occur in branching, parallel universes [204], and the hidden variables of *Bohmian mechanics* [205, 206], to more subjective frameworks like *Quantum Bayesianism (QBism)*, which treats quantum probabilities as an observer's personal beliefs [207], and *spontaneous collapse models* that propose physical mechanisms for the collapse [208].

3. The Dark Side of the Cosmos One of the most humbling discoveries in cosmology is that the familiar matter described by the Standard Model of particle physics constitutes a mere 5% of the universe. The cosmic ledger reveals a startling 95% deficit, an invisible "dark sector" whose existence is only inferred through its distinct gravitational effects. *Dark matter*, a mysterious, non-luminous substance, provides the unseen gravitational scaffolding that shapes galaxy rotation curves and cosmic structure formation [136]. Particle physics offers numerous candidates for it, from *Weakly Interacting Massive Particles (WIMPs)*, extensively searched for in underground detectors [209], to *axions* that might be detected via their conversion to photons in strong magnetic fields [210]. Meanwhile, *dark energy* is the enigmatic force driving the universe's accelerating expansion. Its nature remains even more mysterious, with possibilities ranging from Einstein's *cosmological constant*—whose observed value is so unnaturally small it represents a major theoretical puzzle [211]—to dynamic *quintessence* fields or even *modifications of general relativity* itself [212, 213].

4. The Arrow of Time The *arrow of time* presents a fundamental puzzle: while the microscopic laws of physics are almost entirely time-symmetric, our macroscopic reality is built upon a relentless, forward march from past to future. This asymmetry, or the distinction between past and future, seems to emerge only through the lens of thermodynamics, where the inexorable increase of entropy (disorder) provides a clear directionality. This suggests that time may not be a fundamental aspect of reality but rather an emergent property arising from a deeper, timeless layer of physics [197]. Contemporary research increasingly supports this view, with approaches ranging from *relational theories* where time arises from change itself [214] to proposals where it emerges from *quantum entanglement dynamics* [215]. Other frameworks in quantum gravity explore time as a product of more primitive structures, as seen in *causal set theory* [216], or as a consequence of quantum information flow and *decoherence* [170].

5. The Problem of Naturalness and Cosmic Fine-Tuning In the architecture of the universe, certain numbers appear to be exquisitely "fine-tuned", posing the *hierarchy problem*: why is gravity astonishingly weak compared to the other fundamental forces, and why is the mass of the Higgs boson so much lighter than the natural energy scale where quantum gravity effects should dominate? For decades, physicists suspected that a new principle like *supersymmetry* would provide an elegant, "natural" solution by introducing partner particles that stabilize the Higgs mass [217]. However, the resounding silence from experiments at the Large Hadron Collider, which have failed to find evidence for such new physics, has triggered a deep, philosophical crisis. This has forced physicists to reconsider *naturalness* itself as a guiding principle for theory-building [218, 219]. Consequently, alternative proposals are being explored with renewed interest, including *technicolor models* [220], scenarios where the Higgs is a composite particle, and *anthropic reasoning* within holocosmos frameworks [221].

6. The Arbitrary Elegance of the Standard Model The *Standard Model* of particle physics is a monumental achievement, describing the known elementary particles and their interactions with stunning precision. Yet, for all its predictive power, its structure appears remarkably arbitrary and "baroque", crying out for a deeper organizing principle. It fails to explain why its governing symmetries are the specific group $SU(3) \times SU(2) \times U(1)$, why fundamental particles are organized into three redundant "generations" of increasing mass, or why its nineteen "free parameters"—such as particle masses—have the values they do. This apparent lack of aesthetic simplicity has led physicists to explore possibilities like *Grand Unification Theories (GUTs)*, which seek to merge the three forces into a single, unified force within a larger symmetry group [222]. Other directions include *extra-dimensional models* that seek geometric explanations for particle properties [223] and more abstract approaches using algebraic selection principles like *spectral triples* [224] or other categorical frameworks [225].

7. The Moment of Creation Our cosmological narrative confronts us with the puzzle of *initial conditions*: the Big Bang theory leads back to an initial *singularity*—a moment of infinite density where the known laws of physics break down. This presents two profound problems: what physics governs the universe's birth at or "before" this moment, and why did our universe begin in an extraordinarily *low-entropy*, highly ordered state, a configuration necessary for the subsequent emergence of complex structures like galaxies and stars? Models of *cosmic inflation* partially address the uniformity problem [88, 226], but they do not eliminate the initial singularity itself. Resolving this requires a theory of quantum gravity, leading to various proposals including the *Hartle-Hawking no-boundary proposal* [51], *Penrose's conformal cyclic cosmology*, which recycles universes through infinite cycles [227], and *loop quantum cosmology's* bounce scenarios, which avoid the singularity entirely by suggesting the Big Bang was a "Big Bounce" from a previous cosmic phase [228].

8. The Substance of Reality: Fields, Particles, and Information While modern physics often describes reality as a tapestry of fields where particles are merely localized excitations, the ontological status of these concepts remains surprisingly unsettled. A profound question asks whether fields, particles, and even spacetime itself are truly fundamental or if they are emergent properties of a deeper, more abstract structure. Recent theoretical advances,

particularly the *holographic principle* and the *AdS/CFT correspondence*, lend significant weight to this idea by suggesting that a theory of gravity within a bulk volume can be equivalent to a quantum theory on its boundary [140]. This has fueled the *"It from Qubit"* movement, which speculates that the geometry of spacetime is woven from the fabric of quantum entanglement. This notion is explored through approaches like *tensor networks*, which model spacetime geometry using entanglement patterns [229], and more abstract *categorical and operator-algebraic reformulations* of quantum field theory [230], all hinting that our familiar reality may be an effective description of more primitive, information-based relationships.

9. The Black Hole Information Paradox Black holes are the ultimate theoretical laboratories, presenting a stark clash between general relativity and quantum mechanics that is crystallized in the *black hole information paradox*. In 1976, Stephen Hawking showed that black holes slowly evaporate by emitting thermal radiation, creating a profound conflict: quantum mechanics' principle of *unitary evolution* insists that information can never be destroyed, yet Hawking's calculation suggested it vanishes as the black hole disappears. The fate of this information touches the foundations of both theories and has driven research for decades, leading to further puzzles like the *firewall paradox*, which suggested violent physics at the event horizon [231]. Very recent breakthroughs, however, now point toward a resolution. The development of *quantum extremal surfaces* [232] and the *"island formula"* [139] suggest that information does escape, intricately encoded in the outgoing radiation through subtle gravitational effects and entanglement structures not captured in the original analysis [233].

10. The Ascent from Reductionism While the traditional scientific method of *reductionism*—understanding a system by breaking it down into its smallest components—has been phenomenally successful, many of today's deepest challenges suggest we may be reaching its limits. This has led to a paradigm shift toward *emergence*, driven by the ultimate goal of moving *beyond* describing the universe's laws to the deeper task of *explaining* why they take the specific form they do [234]. This approach asks whether spacetime, particles, and forces can be derived from more primitive, relational, or information-theoretic principles rather than simply being postulated. This speculative frontier includes deriving physics from quantum information, famously captured by John Wheeler's "it from bit" philosophy [235], as well as process-based approaches like *causal set theory* [236]. Other frameworks pursue an *algebraic reconstruction* from symmetry or cohomological data [237] or employ the tools of *categorical field theories* [238], all sharing the tantalizing hope that the universe's apparent complexity may emerge from surprisingly simple, perhaps even inevitable, foundational principles.

The Quest for Unification The enduring quest to reconcile quantum mechanics and general relativity—to unify the probabilistic, relational world of the very small with the dynamical geometry of the very large—remains the most profound challenge in theoretical physics. Any successful theory must ultimately dispense with the notion of a fixed background, revealing spacetime itself as an emergent phenomenon. This work introduces the Interdependent Quantum Reality (IQR) framework, a paradigm grounded in a deeply relational and operator-theoretic ontology. We posit that physical reality emerges from a pre-geometric substrate: a

continuum of structured, operator-valued singularities termed *Primordial Quantum Singularities*. The coherence and dynamics of this continuum are governed by a single, non-factorizable Quantum Holo-State, $|\mathring{\mathbf{\Psi}}\rangle$ (Definition 1.27), from which all emergent structures, including the fabric of spacetime and the matter within it, are derived.

The Fundamental Incompatibility The necessity for such a background-independent approach stems from a fundamental conflict at the heart of physics. General Relativity teaches us that spacetime is a *dynamic actor*—a malleable, geometric fabric whose curvature tells matter and energy how to move, while matter and energy in turn tell spacetime how to curve [239]. This self-consistent feedback loop is the essence of Einstein's theory. In stark contrast, standard Quantum Field Theory treats spacetime as a *fixed, rigid stage*—a static background upon which quantum fields interact and evolve [82].

Herein lies an irreconcilable difference: the stage cannot be both a dynamic participant and a passive, non-participating backdrop. A true theory of quantum gravity must therefore be formulated in a framework where the geometric stage itself is subject to quantum principles. A fixed background is, by definition, a classical object immune to quantum effects—a proposition that is physically untenable at the most fundamental level.

Resolution Through Emergence This theoretical impasse leads to concrete failures in our ability to describe the universe at its most extreme scales. General Relativity predicts its own demise at singularities, points where the curvature of spacetime becomes infinite and the laws of physics cease to apply [240]. A theory built upon a smooth, fixed background cannot describe the physics at or beyond such points.

The formalism, the Interdependent Quantum Reality (IQR) framework, presented in this treatise emerged to address the fundamental incompatibility between general relativity's dynamic spacetime and quantum field theory's fixed background stage [82]. This conflict manifests most acutely at classical singularities where general relativity predicts its own demise [240], and in the quantum superposition principle which demands that massive objects in superposition must induce superposed spacetime curvatures—a concept incompatible with any fixed background. The IQR resolves these paradoxes through a radical reconceptualization: spacetime is not fundamental but emerges from a pre-geometric substrate called the $\hat{\Omega}$-Continuum (Definition 4.617), populated by elementary quantum singularities. When classical physics reaches its limits at what would traditionally be called singularities, there is no catastrophic breakdown but rather a transition to this pre-geometric regime where the relational algebra of Primordial Quantum Singularities governs reality.

At the heart of this theory lies the Primordial Quantum Singularity ($\mathfrak{B}_x$), a structured, operator-valued entity that serves as the fundamental, indivisible unit of quantum-relational structure (Definition 1.25). Each PQS is characterized by its internal architecture:

$$\mathfrak{B}_x := {}_{\hat{\mathbf{\Phi}}_x}\mathring{\mathbf{\Lambda}}_{x,\hat{\mathbf{\Theta}}_x}$$

where $\hat{\mathbf{\Phi}}_x$ and $\hat{\mathbf{\Theta}}_x$ are the *Gaiamiti* (Definition 6.2) and *Tantrakyma* (Definition 6.3) operator fields respectively, unified through the intrinsic duality operator $\mathring{\mathbf{\Lambda}}_x$ that encodes fundamental dualities, superposition, entanglement, and symmetries. These are not singularities in the

classical sense of infinite density, nor are they defined by location in a pre-existing spacetime. Rather, they are operator-valued, field-embedded "quantum grains" of reality, each existing at a locus in the foundational continuum but deriving their meaning entirely through their relational connections to all other PQSs [3].

The coherence of this quantum universe is maintained through a single, non-factorizable Quantum Holo-State $|\mathring{\mathbf{\Psi}}\rangle$ (Definition 1.27), which represents the framework's most profound innovation. This state transcends the conventional quantum mechanical paradigm—it is not a state within a pre-existing Hilbert space but rather generates the Hilbert space structure itself through the Gelfand-Naimark-Segal construction (Definition 1.34) applied to the global operator algebra [230, 241]. The non-factorizability of this universal state is essential: it ensures that every PQS is intrinsically entangled with every other, creating a cosmic tapestry where each quantum singularity defines its existence only through its reflection of all others, analogous to the jewels in Indra's Net [242].

Part I

Foundations

Chapter 1

The Fourfold Algebra and The Singularity

1.1 Introduction

At the heart of modern theoretical physics lies a fundamental contradiction. General Relativity describes spacetime as a dynamic actor, a malleable fabric that dictates how matter moves while matter tells it how to curve [239, 146, 7]. In stark contrast, Quantum Field Theory treats spacetime as a fixed, passive stage upon which quantum events unfold [82, 243]. The central problem is that these two most successful frameworks cannot both be universally correct: one treats spacetime as a dynamical field, the other as an inert backdrop. This work proposes to resolve this impasse by advancing the principle that spacetime is not fundamental, but an emergent property of a deeper, purely quantum–relational reality — an idea that resonates with contemporary research suggesting geometry is woven from the fabric of quantum entanglement [1, 2]. The construction is precise enough to yield a complete Hamiltonian formulation and a background–independent unification of gravity and quantum mechanics.

The treatise develops this principle through the **Interdependent Quantum Reality (IQR) framework** [3, 14]. Within this framework, the fundamental stratum of reality is a relational fibred-network of entangled *Ultra-Sub-Planckian* (Section 4.15.3) *Primordial Quantum Singularities* (PQSs) (Definition 1.25), whose coherence is governed by a single **Quantum Holo-State**, $|\mathring{\mathbf{\Psi}}\rangle$ (Definition 1.27) [244, 245]. The PQS is the loci where times and spaces become unified and indistinguishable [246, 8, 3]. The treatise constructs a triad of independent times: (1) **entanglement phase time** (Section 4.12.1) that quantifies the phase-coherent evolution of entangled modal structure between two PQS loci [170], (2) **vacuum response time** (Section 4.12.3) that characterizes the temporal sensitivity of the vacuum structure to relational displacements [244, 245], and (3) **spectral shear time** (Section 4.12.5) measures a form of quantum "temporal tension" arising from the algebraic structure of the underlying reality [4, 5]. In a similar manner, the treatise also introduces a triad of independent distances: (1) **correlator distance** (Section 4.9.1) is a measure of the magnitude of the quantum signal between two singularities [247], (2) **spectral distance** (Section 4.9.6) is a measure of separation in quantum geometry calculated from the intrinsic "friction" between the operators for change and relational structure [4], and (3) **vacuum-response distance** (Definition 4.463) [244].

To formalize this emergent geometry, the treatise organizes these triadic measures into a

relational phase-geometry bundle (Definition 4.604) [248, 249]. This elegant mathematical structure attaches to every point in the fundamental *quantum configuration space*, $\mathcal{X}_q$, (Section 4.19.3) a fiber (eq. (4.288)) that encodes its complete relational identity. Each fiber $\mathfrak{F}_x$ is a hybrid space that mathematically formalizes the union of distinguishability and temporal flow:

$$\mathfrak{F}_x := \mathcal{F}_x^{\text{dist}} \times \mathcal{F}_x^{\text{time}}.$$

Here, the *distinguishability fiber* $\mathcal{F}_x^{\text{dist}}$ (Definition 4.599) contains the three-dimensional vector of distances, while the orthogonal *temporal flow fiber* $\mathcal{F}_x^{\text{time}}$ (Definition 4.600) contains the vector of corresponding times. This construction provides a profound and rigorous definition for the Primordial Quantum Singularity itself: it is precisely the locus where the distance and time vectors between any two points vanish simultaneously, rendering them relationally indistinguishable (Corollary 4.601). Thus, the very fabric of spacetime is woven from this dynamic bundle, a scaffold upon which quantum analogues of connection and curvature can be defined (Section 4.16) [4]. Consequently, what we perceive as "distance," "duration," and "curvature" are not fundamental, but are macroscopic, statistical averages of the underlying quantum-informational relationships between these foundational units. These units derive their meaning entirely from their interrelations, a concept at the heart of relational quantum mechanics [3].

This web of interrelations is physically realized within the foundational stratum of reality: the **fibred continuum of primordial quantum singularities**, $\mathcal{C}_d$, which is formally defined as the set of all such structured singularities (Definition 4.617) [248, 249]:

$$\mathcal{C}_d := \left\{ \mathfrak{B}_{\vec{r}},\ \hat{\Omega}_{\vec{r}},\ \mathfrak{F}_{\vec{r}} \right\}_{\vec{r} \in \mathcal{C}_d}.$$

This expression defines the continuum as a smooth manifold where every point $\vec{r}$ is itself a structured PQS ($\mathfrak{B}_{\vec{r}}$), containing a singularity code ($\hat{\Omega}_{\vec{r}}$) and hosting a fiber of relational information ($\mathfrak{F}_{\vec{r}}$) with ultra-sub-Planckian support (Section 4.18, eqs. (4.199) and (4.303), and Principle 4.1) [4]. The very *quantum configuration space* $\mathcal{X}_q$, which serves as the base for the Relational Phase-Geometry Bundle, then emerges as a quotient space from this continuum (Proposition 4.625) [248, 249]. This emergence is enacted by an equivalence relation where all points within a single Primordial Quantum Singularity—being operationally indistinguishable as their relational distance and time vectors vanish (Definitions 4.599 and 4.600 and Corollary 4.601)—are identified as a single, unique point in $\mathcal{X}_q$. Thus, the arena for physics is itself a relational construct, born from a deeper reality where each point is a self-contained universe of potential. Yet mathematics itself insists that to define such a point is already to define what it is not. This reciprocity, latent in the geometry of the Relational Phase–Geometry Bundle, is the first hint of a deeper structural necessity: no entity — not even the cosmos in its entirety — can be specified without simultaneously specifying its complement. This observation will lead us directly to the Axiom of Duality, the principle that governs the interplay between self-containment and dependence at every scale of reality [3].

For this emergent arena to serve as a valid stage for physical dynamics, it must possess a coherent and well-defined mathematical structure. The treatise establishes this by defining the **quantum configuration space**, $\mathcal{X}_q$, as the quotient space of the foundational continuum,

$\mathcal{C}_d$ [248, 249]:

$$\mathcal{X}_q := \mathcal{C}_d/\sim$$

where the equivalence relation $\sim$ identifies all points within a single Primordial Quantum Singularity as one (Proposition 4.625). Through the formal application of the *Quotient Manifold Theorem* (Section 4.19.3) [248, 249], this emergent space inherits a smooth manifold structure from the foundational continuum. Consequently, the familiar smooth stage of physics is shown to be a consistent, macroscopic manifestation of the underlying relational coherence [3], retaining the full cardinality of the continuum (Proposition 4.637) while being grounded in a purely operator-theoretic foundation [4].

Given the triadic notion of distance, it is now possible to define the very instrument that imparts a geometric structure onto the emergent quantum configuration space: the **Operator-Valued Emergent Metric Tensor**, $\hat{g}_{\mu\nu}(x)$ (Definition 5.1) [4, 5]. This tensor is not fundamental but is itself emergent, defined as the second **Fréchet variation** of the squared composite distance operator, $D^2(x, x')$, evaluated at the coincidence limit:

$$\hat{g}_{\mu\nu}(x) := \frac{\partial^2}{\partial x^\mu \partial x'^\nu} D^2(x, x')\Big|_{x'=x}.$$

where

$$D^2(x, x') := \hat{d}^2_{\text{corr}}(x, x') + \hat{d}^2_{\text{vac}}(x, x') + \hat{d}^2_{\text{spec}}(x, x'). \tag{1.1}$$

Each $\hat{d}_a(x, x')$ for $a \in \{\text{corr}, \text{vac}, \text{spec}\}$ is a Hermitian operator on $\mathcal{H}_{\text{total}}$. Unlike a classical metric, this defines $\hat{g}_{\mu\nu}(x)$ as a field of Hermitian, positive semi-definite operators, encoding the infinitesimal geometry in the language of quantum mechanics [244, 245]. A profoundly significant feature is that it inherits the triadic structure of its origins, decomposing into correlational, vacuum, and spectral layers, hinting at a non-Riemannian, triadic noncommutative geometry [4, 5]. The bridge to the classical world is crossed when this operator is evaluated in the universal *Quantum Holo-State*, yielding the familiar metric tensor $\bar{g}_{\mu\nu}(x)$ as its expectation value (Definition 5.10) [3]. Thus, $\hat{g}_{\mu\nu}(x)$ is the fundamental link, translating the rich, multi-layered quantum relational data with its ultra-sub-Planckian support into the smooth, curved spacetime we perceive.

Complementing the metric tensor that governs emergent space, the framework defines its temporal analogue: the **operator-valued temporal curvature tensor**, $\hat{t}_{\mu\nu}(x)$ (Definition 5.17) [4, 5]. In a perfectly parallel construction, this tensor arises not from distance but from the triadic operators of relational time. It is formally defined as the second Fréchet variation of the squared composite temporal displacement, $T^2(x, x')$, at the coincidence limit:

$$\hat{t}_{\mu\nu}(x) := \frac{\partial^2}{\partial x^\mu \partial x'^\nu} T^2(x, x')\Big|_{x'=x}.$$

where

$$T^2(x, x') := \hat{t}^2_{\text{corr}}(x, x') + \hat{t}^2_{\text{vac}}(x, x') + \hat{t}^2_{\text{spec}}(x, x'),$$

is the squared norm of the operator-valued triadic temporal vector, as defined in the temporal fiber structure of $\mathfrak{F}_x$ (Definition 4.600) [244, 245]. Crucially, this is **not a metric**. Rather than encoding distance, this field of Hermitian operators quantifies the curvature of relational

quantum phase flow [3]. It measures the directional sensitivity of phase displacement to coordinate separation, providing the dynamic, temporal component of the emergent geometry. Together, the metric tensor $\hat{g}_{\mu\nu}(x)$ and the temporal curvature tensor $\hat{t}_{\mu\nu}(x)$ form the two halves of the complete, unified *Chronogeometric Operator*, $\hat{\Sigma}_{\mu\nu}(x)$ [4].

The companionship of the spatial metric and the temporal phase-flow is now consolidated into a single, comprehensive object that captures the total infinitesimal structure of the emergent geometry: the **Operator-Valued Relational Chronogeometric Operator**, $\hat{\Sigma}_{\mu\nu}(x)$ (Definition 5.25) [4, 5]. Defined as the sum $\hat{\Sigma}_{\mu\nu}(x) := \hat{g}_{\mu\nu}(x) + \hat{t}_{\mu\nu}(x)$, this operator represents the complete description of spacetime at the fundamental, quantum level. The bridge to the classical world is crossed when this unified operator is evaluated in the *quantum holo-state*, yielding the macroscopic *chronogeometric tensor*, $\bar{\Sigma}_{\mu\nu}(x)$ [244, 245]. This emergence is made mathematically precise by the foundational equation of the theory:

$$\bar{\Sigma}_{\mu\nu}(x) = \langle \overset{\circ}{\boldsymbol{\Psi}} | \hat{\Sigma}_{\mu\nu}(x) | \overset{\circ}{\boldsymbol{\Psi}} \rangle = \langle \overset{\circ}{\boldsymbol{\Psi}} | \left(\hat{g}_{\mu\nu}(x) + \hat{t}_{\mu\nu}(x) \right) | \overset{\circ}{\boldsymbol{\Psi}} \rangle.$$

The central ambition of this theory—the unification of general relativity and quantum mechanics—is concisely expressed by this relationship, which establishes the Chronogeometric Tensor, $\bar{\Sigma}_{\mu\nu}(x)$, as a mathematical bridge, explicitly connecting the classical world of gravitation with the fundamental reality of quantum mechanics [3, 14]. The left-hand side, the **Chronogeometric Tensor** $\bar{\Sigma}_{\mu\nu}(x)$, represents the domain of general relativity. It is a classical, macroscopic field that describes the smooth, curved spacetime we perceive as the stage of the cosmos. The right-hand side represents the fundamental quantum reality from which this classical world emerges. It is composed of three essential quantum-mechanical ingredients:

1. The **quantum holo-state** $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$ (see Definition 1.27), the fundamental ontological quantum state that describes the entire universe [3].

2. The **chronogeometric operator** $\hat{\Sigma}_{\mu\nu}(x)$ (see Definition 5.26), the underlying quantum-geometric entity, whose properties are operator-valued subject to quantum principles that are liberated from fixed classical numbers [4, 5].

3. The **expectation value** prescription, $\langle \cdot | \cdot | \cdot \rangle$, is the core mathematical procedure for deriving a measurable, classical outcome from the quantum substrate [14].

The equality in this expression is therefore a profound statement of emergence, a principle visually summarized by annotating the equation's components:

$$\underbrace{\bar{\Sigma}_{\mu\nu}(x)}_{\text{General Relativity}} \quad = \quad \underbrace{\langle \overset{\circ}{\boldsymbol{\Psi}} | \hat{\Sigma}_{\mu\nu}(x) | \overset{\circ}{\boldsymbol{\Psi}} \rangle}_{\text{Quantum Mechanics}}$$

This equation declares that the classical spacetime of general relativity ceases to be a separate or fundamental entity, becoming instead, with mathematical exactness, the macroscopic statistical average of the deeper quantum reality described by the Interdependent Quantum Reality (IQR) framework [3, 4, 5]. This is the heart of the matter: *spacetime geometry is a property of the quantum state*, not the container for the state. The dynamic, chronogeometric nature of reality is therefore a direct consequence of the underlying quantum-informational

relationships [3]. This concept finds a poetic analogue in the metaphor of **Indra's Net**, where each jewel defines its existence only through its reflection of every other [242]. Similarly, the chronogeometry of the universe is a tapestry woven from the threads of entanglement connecting every Primordial Quantum Singularity. To presuppose a fixed background would be akin to demanding that this net be hung upon an external scaffold, when the profound truth is that the relational fabric of cosmic reality supports itself.

The chronogeometric fabric described by the Quantum Holo-State emerges from a pre-geometric and pre-time substrate [3]. The foundation of the Interdependent Quantum Reality (IQR) framework is the $\hat{\mathbf{\Omega}}$–Continuum (Definitions 1.25 and 4.617), a smooth, Hausdorff continuum $\mathcal{C}_d$ with cardinality $\mathfrak{c}$ isomorphic to the Reals ($\mathcal{C}_d \cong \mathbb{R}$) [248, 249]. This continuum is composed of fibred **primordial quantum singularities** (PQSs), $\mathfrak{B}_{\vec{r}}$, with each PQS defined as:

$$\mathfrak{B}_{\vec{r}} := \hat{\mathbf{\Omega}}_{\vec{r}} := \overset{\circ}{\mathbf{\Lambda}}_{\vec{r},\hat{\Theta}_{\vec{r}}}$$

where the components are the local instantiations of three entities novel to the IQR theory: the *Gaiamiti field* $\hat{\mathbf{\Phi}}_{\vec{r}}$ (Definition 6.118), the *Tantrakyma field* $\hat{\mathbf{\Theta}}_{\vec{r}}$ (Definition 6.142), and the **fourfold dialectic algebra of symmetry** $\overset{\circ}{\mathbf{\Lambda}}_{\vec{r}}$ (Section 1.2.7) [4, 5].

The proposed existence of these two novel fundamental fields, the Gaiamiti and the Tantrakyma, govern the emergence of all physical phenomena from this substrate. The **Gaiamiti Field**[1] (Definition 6.2) is a Hermitian, operator-valued field that generates the relational chronogeometry; its quanta, the *apeirons*[2] (see Section 9.17.1), are the progenitors of the long sought-after **graviton**, which arises as the massless Goldstone mode from the **spontaneous symmetry breaking of the Gaiamiti field potential** (Sections 9.17 and 9.17.2) [14].

In parallel, the **Tantrakyma Field**[3] (Definition 6.3) is the fundamental operator-valued field governing the excitation, modulation, and internal symmetry structure of the PQSs across the emergent quantum configuration space $\mathcal{X}_q$ (see Section 4.19.3); its quanta, the *kymons* (see Section 9.18.5), are the progenitors of the **Higgs boson** and the **Standard Model gauge group**, which emerge from the **symmetry breaking of the Tantrakyma field potential around its vacuum expectation value** (VEV) (Sections 9.18.5, 9.18.6 and 12.4) [14], with the particles of the Standard Model manifesting as the various excitations of this broken symmetry.

The *Fourfold Dialectic Algebra of Symmetry*, $\overset{\circ}{\mathbf{\Lambda}}_{\vec{r}}$, serves as the logical blueprint (Defini-

[1] **Gaiamiti** is a compound word of Greek and Sanskrit origin, where **Gaia** (from the Greek Ga ia), invokes the name of the primordial Greek goddess of the Earth, described in classical sources as a foundational entity from which other deities and the physical world emerged [250, 251, 252], and **miti**, is taken from the Sanskrit word **Jyāmiti** (Skt. *Jyāmiti*), which translates to "geometry" [66]. This term itself is a compound of *jyā*, meaning "bowstring," and *miti*, meaning "measure."

[2] **Apeiron** (*Ápeiron*) is the "The Unlimited" or "The Indefinite"; A concept first articulated by Anaximander but central to Pythagorean cosmology. The *apeiron* (*ápeiron*) is the boundless, formless, indefinite primordial substance. The ordered universe (cosmos) comes into being when this unlimited substrate is acted upon and shaped by a principle of "limit" or "form" (*peras*, *péras*) [69, 72, 70, 71].

[3] **Tantrakyma** fuses तन्त्र (*tantra*, "warp, framework; that by which [threads are stretched/laid out]")—as established in classical lexica and modern overviews [66, 74, 75]—with κῦμα (*kyma*, "wave") in classical Greek [80]. The compound denotes a "woven-wave framework": oscillatory modes (waves) organized by an underlying rule-bound warp (framework).

tion 1.16). This algebra serves as the *a priori* "generative grammar" of reality, dictating the necessary modes of Symmetry, Antisymmetry (Section 2.1), Parasymmetry (Definitions 2.85 and 2.94), and Holosymmetry (Sections 2.3 and 3.8) that arise from the foundational Axiom of Duality (Sections 1.2.6, 14.1.3 and 17.3.1) [3]. This abstract logical framework is given its concrete "vocabulary" in the complete *Algebra of Observables*, $\mathcal{A}(\mathcal{C}_d)$ (Definition 1.32) [4, 5].

This algebraic formulation represents a profound reordering of physical ontology; it posits that the abstract C^*-algebra of observables, $A(C_d)$, is more fundamental than any pre-existing spacetime or Hilbert space [4, 5]. This is the representation-independent "language" of reality, existing prior to any notion of distance or time. The crucial step is the transition from this abstract set of rules to the concrete arena where physical dynamics can unfold. This arena is the **Algebra of Endomorphisms**, $\mathrm{End}(\mathcal{H}_{\mathrm{em}})$ (Definition 6.113), which is rigorously defined as the **Banach space of bounded operators**, $\mathcal{B}(\mathcal{H}_{\mathrm{em}})$, (Definition 6.111) acting on the emergent Hilbert space, $\mathcal{H}_{\mathrm{em}}$, of the universe:

$$\mathrm{End}(\mathcal{H}_{\mathrm{em}}) := \mathcal{B}(\mathcal{H}_{\mathrm{em}}),$$

Its structure is chosen for physical and mathematical necessity. As a **Banach space**, it is complete under the *operator norm*—a measure of an operator's maximum "stretching factor"—ensuring a well-behaved and stable calculus for all operator-valued fields [253]:

$$\|A\|_{\mathrm{op}} := \sup_{\substack{\psi\in\mathcal{H}\\ \psi\neq 0}} \frac{\|A\psi\|}{\|\psi\|}.$$

Furthermore, it is a C^***-algebra** because it is equipped with the crucial *adjoint map* ($A \mapsto A^\dagger$) (Definition 6.112), an involution whose defining relation is the bedrock of quantum measurement:

$$\langle\psi|A\phi\rangle = \langle A^\dagger\psi|\phi\rangle.$$

This property is what allows for the existence of self-adjoint operators corresponding to physical observables. To compare these operators across the emergent manifold, the framework also derives an **Operator Bundle Connection**, $A_\mu(x)$ (Definition 6.192). This is an intrinsic gauge field that governs the parallel transport of quantum information, transforming to ensure physical laws remain consistent from point to point:

$$\mathcal{A}_\mu(x) \mapsto U(x)\mathcal{A}_\mu(x)U^\dagger(x) + i\,U(x)\partial_\mu U^\dagger(x).$$

The profound question of how the abstract *language* of the Algebra of Observables is translated into a concrete and fully equipped *arena* is answered by the theory's central mechanism of emergence: the **Gelfand-Naimark-Segal (GNS) construction** (Sections 1.4 and 1.4.1) [254, 230]. This process is the mathematical embodiment of physical actualization, a three-act story of how a coherent reality is built from a pre-geometric foundation.

First, the theory posits the **Abstract Algebra of Observables**, $A(C_d)$, as the foundational reality (Definition 1.32). This is the complete, representation-independent *"language"* of the universe, containing all possible physical relationships before any notion of a space for them to exist in has been defined [4, 5]. Second, the **Quantum Holo-State**, $|\Psi\rangle$, acts as the *"principle of cosmic selection."* It is represented as a pure state functional (Definition 1.34),

$$\omega^\Psi : A(C_d) \to \mathbb{C},$$

which reads the abstract script and assigns a definite expectation value to every potential observable, thereby selecting one coherent performance from the infinite possibilities.

Finally, the GNS construction serves as the mathematical engine that takes the algebra and the state and from them, generates the manifest reality as a unique **GNS Triplet** (H^{Ψ}, π^{Ψ}, $|\Omega_{GNS}\rangle$) (Definition 1.34) [254, 230]:

1. **The Stage (The Emergent Hilbert Space, H_{em}):** A bespoke arena whose very inner product is defined by the state (Definition 4.287):

$$\langle [A], [B] \rangle := \omega^{\Psi}(A^{\dagger}B)$$

,

2. **The Actors (The Representation, π):** A map,

$$\pi^{\Psi} : A(C_d) \to \mathcal{B}(\mathcal{H}_{\text{em}})$$

, that realizes the abstract observables as concrete, bounded operators acting on the Hilbert space via the rule $\pi^{\Psi}(A)[B] := [AB]$ (Definition 1.34),

3. **The Protagonist (The Cyclic Vector, $|\Omega_{GNS}\rangle$):** The concrete vector that embodies the Quantum Holo-State, from which all expectation values are recovered through the core relation:

$$\omega^{\mathring{\mathbf{\Psi}}}(A) = \langle \Omega_{\text{GNS}} | \pi^{\mathring{\mathbf{\Psi}}}(A) | \Omega_{\text{GNS}} \rangle.$$

This reveals a profound insight of our framework: the quantum state of the universe is the very architect of its own domain, not an ordinary inhabitant of a pre-existing container (Remark 1.36). The state functional defines the very geometry of the Hilbert space it inhabits. The **Fourfold Dialectic Algebra** (Definition 1.16) provides the axiomatic rules of construction, and the GNS construction is the act of building a coherent universe according to those rules. This process is the mathematical realization of true background independence, where the universe's quantum state and its arena of existence emerge as inseparable aspects of a unified, self-determining whole (Remark 1.37) [3].

This connection is made mathematically concrete through the definition of the inner product, which is *derived* from the system's fundamental correlations and not assumed. For any two modal states $|\psi_1\rangle_x, |\psi_2\rangle_x$ in the local Hilbert space $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}$, their inner product is defined via the **Fourfold Inter-Modal Correlator**, $\hat{G}_{\Lambda}(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2]$ (Definition 4.9), evaluated in the Quantum Holo-State:

$$\langle \psi_1 | \psi_2 \rangle_{\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}} := \lim_{x' \to x} \langle \mathring{\mathbf{\Psi}} | \hat{G}_{\Lambda}(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2] | \mathring{\mathbf{\Psi}} \rangle.$$

This is an operator-valued, gauge-covariant two-point function that quantifies quantum-relational correlations between any pair of internal modes in the complete *Fourfold Dialectic Algebra of Symmetry* of two Primordial Quantum Singularities, $\mathfrak{B}_x$ and $\mathfrak{B}_{x'}$ [230, 255, 3]. It defines a bounded linear operator on the total emergent Hilbert space $\mathcal{H}_{\text{em}}$ and provides the structural kernel for inter-PQS interference, "holomodal" entanglement, and emergent chronogeometry [4]. This formulation provides a physical mechanism for generating the

Hilbert space metric directly from the relational dynamics of the fundamental fields, as encoded in the Quantum Holo-State. The total emergent Hilbert space is then the disjoint union of all such locally generated spaces (Definition 4.228):

$$\mathcal{H}_{\text{em}} = \bigsqcup_{x \in \mathcal{X}_q} \mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}$$

To make the emergent inner product mathematically concrete, we must first define its core ingredients: the general quantum states that reside on each Primordial Quantum Singularity (PQS) and the operator that quantifies the relational connection between them. Any local modal quantum state within the IQR framework is described as a superposition of four fundamental ontological modes. A general state $|\psi\rangle_x$ at a locus x and another state $|\psi\rangle_{x'}$ at locus x' are thus expressed as a linear combination of these basis "vectors":

$$|\psi_1\rangle_x = \sum_\alpha c_{1\alpha} |\alpha\rangle_x, \qquad |\psi_2\rangle_{x'} = \sum_\beta c_{2\beta} |\beta\rangle_{x'},$$

where the indices α, β range over the four basis vectors. These vectors are the direct physical manifestation of the four logical vertices of the *Fourfold Dialectic Algebra of Symmetry* (Definition 1.16):

$$\left\{ |\mathring{\mathbf{0}}\rangle,\ |\mathring{\mathbf{1}}\rangle,\ |\mathring{\mathbf{0}} + \mathring{\mathbf{1}}\rangle,\ |_{\mathring{\mathbf{0}}}\mathring{\mathbf{\Lambda}}_{\mathring{\mathbf{1}}}\rangle \right\}.$$

These basis vectors correspond, respectively, to the fundamental *Symmetric*, *Antisymmetric*, *Parasymmetric*, and *Holosymmetric* modes of reality. The relational link between two such states at different loci, $|\psi_1\rangle_x$ and $|\psi_2\rangle_{x'}$, is quantified by the *Fourfold Inter-Modal Correlator*. This operator, $\hat{G}_\Lambda$, is a bi-local (two-point) function that serves as the engine of relational dynamics, defining the structure of entanglement and interference across the $\hat{\mathbf{\Omega}}$–Continuum [3, 230, 255]. It is formally defined as (Definition 4.9):

$$\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2] := \sum_{\alpha,\beta} c_{1\alpha}^*\, c_{2\beta} \left(\hat{\mathcal{F}}_\alpha^\dagger(x)\, \mathcal{U}(x \leftarrow x'; \gamma)\, \hat{\mathcal{F}}_\beta(x') \right).$$

The components of this correlator ensure that the relationship is well-defined across the emergent manifold. The operator $\mathcal{U}(x \leftarrow x'; \gamma)$ is the *gauge-covariant parallel transporter* [248, 256]. Its role is to connect the Hilbert space fibers at x' and x along a path γ within the emergent quantum configuration space $\mathcal{X}_q$. This guarantees that the calculated correlation is a true physical invariant, independent of the choice of local quantum reference frame.

The interplay between geometry and matter, the dance between the Gaiamiti and Tantrakyma fields, are consolidated into a single mathematical object. This *Generalized Field Operator* provides a unified representation of the total operator-valued content of each Primordial Quantum Singularity, expressed as a unified field vector [82, 14]:

$$\hat{\mathbf{F}}(x) := \begin{pmatrix} \hat{\mathbf{\Phi}}(x) \\ \hat{\mathbf{\Theta}}(x) \end{pmatrix}.$$

The complete relational dynamics of the system can then be explored by correlating this unified field with itself [230, 3]. This process naturally yields a 2×2 matrix structure, which

we call the *Field Correlator Matrix*:

$$\hat{\mathbf{G}}_{\text{field}}(x, x') = \begin{pmatrix} \hat{G}_{\mathbf{\Phi\Phi}}(x, x') & \hat{G}_{\mathbf{\Phi\Theta}}(x, x') \\ \hat{G}_{\mathbf{\Theta\Phi}}(x, x') & \hat{G}_{\mathbf{\Theta\Theta}}(x, x') \end{pmatrix}.$$

This matrix provides a complete map of the fundamental interactions. The *diagonal entries*, $\hat{G}_{\mathbf{\Phi\Phi}}$ and $\hat{G}_{\mathbf{\Theta\Theta}}$, describe the self-correlations, or internal dynamics, within the geometric and matter sectors, respectively. The *off-diagonal entries*, $\hat{G}_{\mathbf{\Phi\Theta}}$ and $\hat{G}_{\mathbf{\Theta\Phi}}$, represent the crucial interaction terms, quantifying the cross-correlation or "talk" between geometry and matter.

This 2×2 matrix, however, only describes the structure at the level of the fields [230, 14]. Each fundamental interaction must also respect the underlying logic of the *Fourfold Dialectic Algebra* [3, 4]. Consequently, each of the four entries in the Field Correlator Matrix is itself a 4×4 matrix, with components defined in the ontological basis of the four dialectical modes:

$$\hat{G}_{AB}(x, x') = \left[\hat{G}^{\alpha\beta}_{AB}(x, x')\right]_{\alpha,\beta\in\{0,1,p,s\}},$$

where each component $\hat{G}^{\alpha\beta}_{AB}$ corresponds to a transition between the ontological modes β and α within the fields A and B. This reveals the profound, multi-layered nature of a single relational link in the IQR framework. Each entry of the four components of the Field Correlator Matrix, $\hat{\mathbf{G}}_{\text{field}}$, is an operator on the total emergent Hilbert space, $\mathcal{H}_{\text{em}}$, and represents a distinct physical channel of relational correlation. Their formal definitions are constructed by specializing the generic field operator $\hat{\mathcal{F}}$ from the definition of the Fourfold Inter-Modal Correlator (Definition 4.9) to the specific Gaiamiti ($\hat{\mathbf{\Phi}}$) and Tantrakyma ($\hat{\mathbf{\Theta}}$) fields.

The Geometric Sector: $\hat{G}_{\mathbf{\Phi\Phi}}$ The *Geometric Correlator* describes the self-correlation of the Gaiamiti field [14, 230, 3]. It is the exclusive source of emergent geometry, defining how geometric potential propagates and relates to itself across the continuum. For general modal states $|\psi_1\rangle_x = \sum_\alpha c_{1\alpha}|\alpha\rangle_x$ and $|\psi_2\rangle_{x'} = \sum_\beta c_{2\beta}|\beta\rangle_{x'}$, it is defined as:

$$\hat{G}_{\mathbf{\Phi\Phi}}[\psi_1^*, \psi_2] := \sum_{\alpha,\beta} c^*_{1\alpha} c_{2\beta} \left(\hat{\mathbf{\Phi}}^\dagger_\alpha(x)\, \mathcal{U}(x \leftarrow x')\, \hat{\mathbf{\Phi}}_\beta(x')\right)$$

The Matter Sector: $\hat{G}_{\mathbf{\Theta\Theta}}$ The *Matter Correlator* captures the self-relational structure of the Tantrakyma field [14, 230, 3]. It governs the propagation and interaction of all matter and force excitations, with its properties determining the phenomenology of the emergent particles. It is defined as:

$$\hat{G}_{\mathbf{\Theta\Theta}}[\psi_1^*, \psi_2] := \sum_{\alpha,\beta} c^*_{1\alpha} c_{2\beta} \left(\hat{\mathbf{\Theta}}^\dagger_\alpha(x)\, \mathcal{U}(x \leftarrow x')\, \hat{\mathbf{\Theta}}_\beta(x')\right)$$

The Interaction Sector: $\hat{G}_{\mathbf{\Phi\Theta}}$ and $\hat{G}_{\mathbf{\Theta\Phi}}$ The *Interaction Correlators* are the off-diagonal terms that encode the direct, dynamic coupling between geometry and matter [14, 230, 7, 3].

They are defined as:

$$\hat{G}_{\mathbf{\Phi\Theta}}[\psi_1^*, \psi_2] := \sum_{\alpha,\beta} c_{1\alpha}^* c_{2\beta} \left(\hat{\mathbf{\Phi}}_\alpha^\dagger(x)\, \mathcal{U}(x \leftarrow x')\, \hat{\mathbf{\Theta}}_\beta(x') \right)$$
$$\hat{G}_{\mathbf{\Theta\Phi}}[\psi_1^*, \psi_2] := \sum_{\alpha,\beta} c_{1\alpha}^* c_{2\beta} \left(\hat{\mathbf{\Theta}}_\alpha^\dagger(x)\, \mathcal{U}(x \leftarrow x')\, \hat{\mathbf{\Phi}}_\beta(x') \right)$$

These operators give mathematical life to the core principles of general relativity [7]: $\hat{G}_{\mathbf{\Phi\Theta}}$ quantifies how matter sources geometric curvature, while $\hat{G}_{\mathbf{\Theta\Phi}}$ quantifies how geometry dictates the dynamics of matter. In these definitions, the terms $\hat{\mathbf{\Phi}}_\alpha(x)$ and $\hat{\mathbf{\Theta}}_\beta(x')$ represent the operator-valued field modes for the Gaiamiti and Tantrakyma fields, respectively, corresponding to the specific dialectical modes $|\alpha\rangle_x$ and $|\beta\rangle_{x'}$ of the ontological basis.

The diagonal components of the Field Correlator Matrix describe the internal dynamics of reality's two fundamental facets. The *Geometric Correlator*, $\hat{G}_{\mathbf{\Phi\Phi}}$, represents the self-relation of the Gaiamiti field and serves as the exclusive source of the emergent cosmic stage [14, 230, 3]; its expectation value gives rise to the metric and temporal tensors that define how chronogeometry propagates and correlates with itself. In parallel, the *Matter Correlator*, $\hat{G}_{\mathbf{\Theta\Theta}}$, captures the self-relational structure of the Tantrakyma field. It governs the propagation, interaction, and spectral properties—such as mass and mixing—of all emergent particles that act upon that stage.

The profound unity of the framework is encoded in the off-diagonal *Interaction Correlators*, which mathematically describe the inseparable coupling between these two sectors. These terms give rigorous, operator-level expression to the core principles of general relativity [7]: $\hat{G}_{\mathbf{\Phi\Theta}}$ quantifies how a matter excitation at one locus sources a geometric deformation at another, enacting the principle that *matter tells geometry how to curve*. Conversely, $\hat{G}_{\mathbf{\Theta\Phi}}$ describes how a geometric fluctuation influences the matter field, corresponding to the principle that *geometry tells matter how to move*. The non-vanishing of these interaction terms is a definitive statement that geometry and matter are not separate entities, but are interdependent, co-arising facets of a single, dynamically unified reality [3].

In accordance with the principle of stationary action that underpins modern physics [257], the dynamics of the IQR framework are governed by a single, fundamental quantity: the *Relational Action Functional*, $S[|\mathbf{\Psi}\rangle]$. This functional assigns a scalar value—a measure of total relational coherence—to any physically admissible quantum state, $|\mathbf{\Psi}\rangle$, allowing the preferred, physically realized state of the universe to be identified by extremizing this action. Its construction requires two key components derived from a given state: a measure of spacetime volume and a measure of the relational content within that volume.

First, to integrate over the emergent manifold, we must define a volume element. In a profound departure from theories with fixed backgrounds [3, 197], the geometry itself is state-dependent. We therefore define the *state-dependent Chronogeometric Tensor* as the expectation value of the Chronogeometric Operator in a given quantum state $|\mathbf{\Psi}\rangle$ [14]:

$$\bar{\Sigma}_{\mu\nu}^{(\mathbf{\Psi})}(x) := \langle \mathbf{\Psi} | \hat{\Sigma}_{\mu\nu}(x) | \mathbf{\Psi} \rangle.$$

The *Relational Volume Element* (Definition 6.33),

$$\sqrt{-\det \bar{\Sigma}_{\mu\nu}^{(\mathbf{\Psi})}(x)},$$

is thus dynamically determined by the quantum state of the universe itself [230]. The "substance" to be integrated over this dynamic volume is the total local self-correlation of the system, given by the trace of the Fourfold Inter-Modal Correlator's expectation value. Combining these elements yields the *Relational Action Functional* for the state $|\Psi\rangle$ (eq. (6.84)):

$$S[|\Psi\rangle] := \int_{\mathcal{X}_q} d^d x \, \sqrt{-\det \bar{\Sigma}^{(\Psi)}_{\mu\nu}(x)} \, \mathrm{Tr}\left(\langle\Psi|\hat{\mathbf{G}}_\Lambda(x,x)|\Psi\rangle\right).$$

The integrand of this action captures the complete relational curvature at each point $x \in \mathcal{X}_q$. This curvature is composed of two distinct contributions, corresponding to the two fundamental aspects of chronogeometry:

- The curvature of *distinguishability*, derived from the emergent metric tensor, which defines spatial relationships: $\bar{g}^{(\Psi)}_{\mu\nu}(x) := \langle\Psi|\hat{g}_{\mu\nu}(x)|\Psi\rangle$ [7].
- The curvature of *quantum phase flow*, derived from the emergent temporal tensor, which defines causal and temporal relationships: $\bar{t}^{(\Psi)}_{\mu\nu}(x) := \langle\Psi|\hat{t}_{\mu\nu}(x)|\Psi\rangle$ [29].

The total state-dependent Chronogeometric Tensor is the sum of these two components:

$$\bar{\Sigma}^{(\Psi)}_{\mu\nu}(x) = \bar{g}^{(\Psi)}_{\mu\nu}(x) + \bar{t}^{(\Psi)}_{\mu\nu}(x).$$

The *Relational Action Functional* provides the theory's central dynamical law through a *variational principle* [258, 14, 259]. The physically admissible quantum states of the universe are those that extremize the action, imposing a condition of maximal coherence and minimal relational curvature:

$$\delta S[|\Psi\rangle] = 0.$$

The set of all states $|\Psi\rangle_{\text{vac}}$ that satisfy this principle constitutes the *Vacuum Manifold* (Definition 6.61) [260, 261]. Within this manifold lies the single distinguished state that describes our realized cosmos: the *Quantum Holo-State*, $|\mathring{\Psi}\rangle$ [3, 197].

When the general, state-dependent definitions are evaluated for this universal vacuum state, they reduce to their canonical forms. The *Chronogeometric Tensor* becomes the canonical vacuum geometry furnishing the background of our universe:

$$\bar{\Sigma}_{\mu\nu}(x) := \langle\mathring{\Psi}|\hat{\Sigma}_{\mu\nu}(x)|\mathring{\Psi}\rangle \quad (\text{cf. } \textit{Definition } 5.26)\ [7,\ 8,\ 3].$$

The action functional in turn evaluates to its realized vacuum value:

$$S[|\mathring{\Psi}\rangle] := \int_{\mathcal{X}_q} d^d x \, \sqrt{-\det \bar{\Sigma}(x)} \, \mathrm{Tr}\Big(\langle\mathring{\Psi}|\hat{\mathbf{G}}_\Lambda(x,x)|\mathring{\Psi}\rangle\Big)$$

[262, 263].

This canonical geometry, determined entirely by the Quantum Holo-State, provides the stable background against which all local fluctuations, particles, and interactions are defined [264, 146]. The *Relational Action Functional* (Definition 6.38) thus serves as the *master functional* governing the total dynamics of the IQR framework [14, 258].

To resolve its internal structure, we may decompose the total action into sectors corresponding to the block decomposition of the *Field Correlator Matrix* $\hat{\mathbf{G}}_\Lambda$ [82, 243]:

$$S[|\Psi\rangle] = S_{\Phi\Phi} + S_{\Theta\Theta} + S_{\Phi\Theta} + S_{\Theta\Phi}.$$

This separation isolates the distinct variational principles for:

- **Geometry** ($S_{\Phi\Phi}$): the pure Gaiamiti sector [7],
- **Matter** ($S_{\Theta\Theta}$): the pure Tantrakyma sector [82],
- **Interaction** ($S_{\Phi\Theta}, S_{\Theta\Phi}$): the bidirectional coupling between geometry and matter [146, 264].

Such a decomposition makes explicit how the master variational principle governs not only the emergent background structure but also the mutual dynamical influence of the universe's two fundamental facets.

The *Gaiamitiic Action*, S_Φ, is defined by projecting onto the geometric sector. It is the functional whose variation governs the emergence of pure chronogeometry—the metric and causal structure of spacetime [7, 146, 264]:

$$S_\Phi[|\Psi\rangle] := \int_{\mathcal{X}_q} d^d x \, \sqrt{-\det \bar{\Sigma}(x)} \, \mathrm{Tr}\left(\langle\Psi|\hat{G}_{\Phi\Phi}(x,x)|\Psi\rangle\right).$$

In parallel, the *Tantrakyma Action*, S_Θ, is defined from the matter sector. Its variation governs the dynamics of all material content, including mass generation and internal symmetries [82, 243, 97, 99]:

$$S_\Theta[|\Psi\rangle] := \int_{\mathcal{X}_q} d^d x \, \sqrt{-\det \bar{\Sigma}(x)} \, \mathrm{Tr}\left(\langle\Psi|\hat{G}_{\Theta\Theta}(x,x)|\Psi\rangle\right).$$

Finally, the crucial *Interaction Action*, $S_{\Phi\Theta}$, is derived from the off-diagonal correlators. This term encodes the quantum analog of gravitational coupling, describing the inseparable dialogue where geometry influences matter and vice versa [6, 146, 264]:

$$S_{\Phi\Theta}[|\Psi\rangle] := \int_{\mathcal{X}_q} d^d x \, \sqrt{-\det \bar{\Sigma}(x)} \, \mathrm{Tr}\left(\langle\Psi|\left[\hat{G}_{\Phi\Theta}(x,x) + \hat{G}_{\Theta\Phi}(x,x)\right]|\Psi\rangle\right).$$

This decomposition makes it clear that the total Relational Action Functional, $S[|\Psi\rangle]$, serves as a *unified variational umbrella* [258, 14]. A single principle of stationary action, applied to this total functional, simultaneously and self-consistently determines the dynamics of spacetime, the behavior of matter, and the fundamental laws of their interaction.

Applying this variational principle yields the theory's central dynamical law for the emergent manifold—a profound generalization of the Einstein Field Equations [6, 146, 7] (Section 11.5.4), the field equations that govern the evolution of the emergent chronogeometric manifold. These equations relate the curvature of the unified geometry to the total stress-energy content of the universe. The semi-classical field equations for the expectation-level chronogeometric tensor $\bar{\Sigma}_{\mu\nu}(x)$ are given by the *Chronogeometric Einstein Equations*:

$$\bar{G}_{\mu\nu}[\bar{\Sigma}] + \Lambda\bar{\Sigma}_{\mu\nu} = \kappa\bar{T}^{\mu\nu}_{\text{total}}(x),$$

where:

- $\bar{G}_{\mu\nu}[\bar{\Sigma}]$ is the *Einstein tensor of the unified geometry.*
- Λ is the *emergent cosmological constant* (Definition 17.2).
- κ is the *relational gravitational coupling constant* (Section 11.5.3).
- $\bar{T}^{\mu\nu}_{\text{total}}(x)$ is the *total expectation-valued stress–energy tensor* (Section 9.18.12), representing the complete energy, momentum, and stress content of both the Gaiamiti (geometric) and Tantrakyma (matter) fields [244, 13].

Equation (11.191) is the central dynamical law of emergent spacetime in this framework. While it bears a formal resemblance to the classical Einstein Field Equations [6, 7], its physical content is significantly richer:

- *The Geometric Side (LHS):* The Einstein tensor $\bar{G}_{\mu\nu}[\bar{\Sigma}]$ is not derived from a simple metric but from the unified tensor $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$. It therefore describes curvature in both the spatial distinguishability ($\bar{g}_{\mu\nu}$) and the **quantum phase flow** ($\bar{t}_{\mu\nu}$) of the manifold [3, 265]. The geometry adapts not only to the distribution of energy but also to the distribution of quantum coherence.
- *The Source Side (RHS):* The source of this curvature, $\bar{T}^{\mu\nu}_{\text{total}}$, is the expectation value of the total quantum stress-energy operator (Section 9.18.12). It includes the energy of matter and radiation (from the Tantrakyma field) as well as the energy of the geometric field itself (from the Gaiamiti field). This establishes a self-consistent feedback loop where the energy of the geometry contributes to its own curvature [244, 13].

In essence, these equations state that the total chronogeometric curvature of the emergent universe is proportional to the total quantum stress-energy content within it. While these field equations describe the emergent dynamics, a deeper insight into the system's total energy and its fundamental quantum nature is provided by the *Relational Hamiltonian*, $\hat{H}$ [258, 14].

In a background-independent framework [3, 265], the Hamiltonian represents the total relational energy of a given quantum state. Its physical meaning is defined implicitly: the expectation value of the Hamiltonian in a state $|\mathbf{\Psi}\rangle$ *is* the value of the Relational Action Functional for that state [258, 14] (eq. (6.107)):

$$\langle \mathbf{\Psi} | \hat{H} | \mathbf{\Psi} \rangle := S[|\mathbf{\Psi}\rangle].$$

This principle is given an explicit mathematical form by constructing a state-dependent operator, $\hat{H}[|\mathbf{\Psi}\rangle]$, as the integral of the local correlator's trace over the emergent manifold (eq. (6.108)):

$$\hat{H}[|\mathbf{\Psi}\rangle] := \int_{\mathcal{X}_q} d^d x \, \sqrt{-\det \bar{\Sigma}^{(\mathbf{\Psi})}_{\mu\nu}(x)} \, \text{Tr}_{\text{op}} \left(\hat{\mathbf{G}}_\Lambda(x,x) \right),$$

where Tr_{op} is the partial trace over the internal field and dialectic indices. The state-dependence of this Hamiltonian is a crucial and necessary feature of the theory's self-consistency [3, 265]. Arising from the volume element $\sqrt{-\det \bar{\Sigma}^{(\mathbf{\Psi})}}$, it signifies that the very geometry used to measure the universe's total energy is itself determined by the state being measured.

Just as the action was decomposed, the total Hamiltonian inherits its structure from the Field Correlator Matrix [266, 8, 3]. It thus admits a canonical decomposition into geometric (Gaiamitiic), matter (Tantrakyma), and Interaction components (eq. (6.109)):

$$\hat{H}[|\mathbf{\Psi}\rangle] = \hat{H}_{\Phi}[|\mathbf{\Psi}\rangle] + \hat{H}_{\Theta}[|\mathbf{\Psi}\rangle] + \hat{H}_{\Phi\Theta}[|\mathbf{\Psi}\rangle],$$

where each component is defined by integrating the trace of the corresponding correlator block:

$$\begin{aligned}
\hat{H}_{\Phi}[|\mathbf{\Psi}\rangle] &:= \int_{\mathcal{X}_q} d^d x \sqrt{-\det \bar{\Sigma}^{(\mathbf{\Psi})}}\, \mathrm{Tr}_{\mathrm{op}} \left(\hat{G}_{\mathbf{\Phi\Phi}}(x,x) \right), \\
\hat{H}_{\Theta}[|\mathbf{\Psi}\rangle] &:= \int_{\mathcal{X}_q} d^d x \sqrt{-\det \bar{\Sigma}^{(\mathbf{\Psi})}}\, \mathrm{Tr}_{\mathrm{op}} \left(\hat{G}_{\mathbf{\Theta\Theta}}(x,x) \right), \\
\hat{H}_{\Phi\Theta}[|\mathbf{\Psi}\rangle] &:= \int_{\mathcal{X}_q} d^d x \sqrt{-\det \bar{\Sigma}^{(\mathbf{\Psi})}}\, \mathrm{Tr}_{\mathrm{op}} \left(\hat{G}_{\mathbf{\Phi\Theta}}(x,x) + \hat{G}_{\mathbf{\Theta\Phi}}(x,x) \right).
\end{aligned}$$

The physical ground states of the universe are the eigenstates of the Hamiltonian with the lowest energy eigenvalue, E_0 [258, 14]. This ground state eigenspace, $\mathcal{V}_0 = \ker(\hat{H} - E_0 \mathbb{I}_{\mathrm{em}})$, is precisely the set of states that minimize the action, thereby defining the *Vacuum Manifold*, $\mathcal{M}_{\mathrm{vac}}$ (Definition 6.61).

A central consequence of the Interdependent Quantum Reality (IQR) framework is that time is not a fundamental background parameter but an emergent, relational construct [197, 3]. Therefore, the universe's dynamics cannot be described by a standard Schrödinger equation, which presupposes an external time parameter for evolution. Instead, the fundamental law must be a *constraint* imposed on the total quantum state of the cosmos. This approach elevates the logic of the Wheeler–DeWitt equation [156, 8, 3] to a fully operator-theoretic context, yielding a single, timeless law known as the *Universal Quantum Constraint*.

The operator that governs this constraint is the *Total Relational Hamiltonian*, $\hat{H}_{\mathrm{univ}}$ [266, 8, 3]. Formally derived from the total action functional (Definition 9.357) via a Legendre transform, this Hamiltonian encapsulates the complete dynamics of the system. It is the sum of the geometric (Gaiamiti), matter (Tantrakyma), and interaction sectors (Definition 13.167):

$$\hat{H}_{\mathrm{univ}} := \hat{H}_{\Phi} + \hat{H}_{\Theta} + \hat{H}_{\mathrm{int}}.$$

The fundamental law of the IQR universe is the **Universal Quantum Constraint** [156, 3], expressed as a stationary, zero-eigenvalue solution for the **Quantum Holo-State**, $|\mathring{\mathbf{\Psi}}\rangle$ (Theorem 13.168):

$$\hat{H}_{\mathrm{univ}} |\mathring{\mathbf{\Psi}}\rangle = 0.$$

The operator in this expression, the **Total Relational Hamiltonian** $\hat{H}_{\mathrm{univ}}$, is a fundamentally unified object derived from the Total Relational Action (Definition 9.357) that intrinsically includes the dynamics of the geometric (Gaiamiti), matter (Tantrakyma), and interaction sectors [266, 8, 3]. This single constraint thus represents a unified quantum law for the entire coupled system of matter and geometry.

This equation asserts that the universe is a fully self-contained system whose total relational energy vanishes [197, 3]. It is not an equation for geometry *with* matter sources,

but a constraint on the one timeless, entangled state from which both are derived. As such, the Universal Quantum Constraint represents the background-independent generalization of the Schrödinger equation, where any notion of time must emerge from within the relational structure of the state itself [156, 8, 3].

How can a universe governed by this timeless constraint give rise to the familiar experience of evolution? The answer lies in partitioning the total system [3, 197]. While the Quantum Holo-State of the entire cosmos is stationary, an observer *within* the universe is a subsystem. Such an observer can describe the evolution of another subsystem (the "system of interest") relative to the rest of the universe, which acts as a clock and reference frame. This gives rise to an **effective, emergent dynamics** governed by a time-dependent Schrödinger equation [215, 8, 3].

The evolution of any such subsystem is governed by this emergent dynamical law (eq. (17.47)):

$$i\hbar\frac{d}{dt}|\mathring{\boldsymbol{\Psi}}(t)\rangle = \hat{H}_{\text{rel}}(t)|\mathring{\boldsymbol{\Psi}}(t)\rangle$$

The operator that drives this effective evolution is the **Total Relational Hamiltonian**, $\hat{H}_{\text{rel}}$, which contains the entirety of cosmic phenomenology:

$$\hat{H}_{\text{rel}} = \underbrace{\hat{H}_{\text{geometry}}[\hat{\Sigma}_{\mu\nu}]}_{\text{Dark Energy \& Dark Matter}} + \underbrace{\hat{H}_{\text{matter}}[\hat{\phi}, \hat{\psi}, \dots]}_{\text{Standard Model}}.$$

This single effective Hamiltonian is composed of two sectors:

1. **The Geometric Hamiltonian ($\hat{H}_{\text{geometry}}$):** This operator governs the dynamics of the **Gaiamiti field** ($\hat{\Phi}$), whose quanta are the **Apeirons** [267]. Its ground state eigenvalue is the Cosmological Constant (Λ), the physical origin of **Dark Energy**, while its large-scale coherent excitations are the physical manifestation of **Dark Matter** (Section 13.1).

2. **The Matter Hamiltonian ($\hat{H}_{\text{matter}}$):** This operator governs the dynamics of the **Tantrakyma field** ($\hat{\Theta}$), whose quanta are the **Kymons**. Its excitations and their interactions give rise to the entire particle zoo of the **Standard Model** [14] (Chapter 12).

Thus, the familiar phenomena of our evolving universe are understood as different facets of a single, unified, but fundamentally timeless relational structure [3, 197].

While the emergent Schrödinger equation governs the dynamics of subsystems, the **Total Relational Hamiltonian**, $\hat{H}_{\text{rel}}$, that drives this evolution is not itself a fundamental entity [3, 215]. It is a global operator that arises from a deeper, more localized layer of reality's relational structure. The most basic building block of all dynamics in the IQR framework is the **Fourfold Inter-Modal Correlator**, $\hat{G}_\Lambda(x, x')$. This is a bi-local (two-point) operator-valued function whose essential role is to define the quantum-relational link—the entanglement and coherence—between any two Primordial Quantum Singularities at points x and x' [8, 3].

The global operator that we identify as the Total Relational Hamiltonian is constructed from this fundamental correlator. The process involves taking the local part of the correlator where the two points coincide ($\hat{G}_\Lambda(x, x)$), tracing over its internal indices to yield a local relational energy density, and then integrating this density over the entire emergent manifold,

$\mathcal{X}_q$. The Hamiltonian is thus the macroscopic, integrated sum of all microscopic, relational interactions across the cosmos.

The emergent Schrödinger equation (eq. (17.47)) therefore describes how the **Quantum Holo-State**, when viewed from the perspective of an internal subsystem, appears to evolve under the influence of this total, integrated interaction energy [215]. The solution, $|\mathring{\mathbf{\Psi}}(t)\rangle$, is a single quantum state whose different aspects we perceive as the familiar world of particles, the enigmatic halos of dark matter, and the accelerating expansion driven by dark energy (Section 13.1). In this light, the majestic evolution of the universe is understood to be the global manifestation of the profound relational dynamics encoded within the Fourfold Inter-Modal Correlator.

The Universal Quantum Constraint defines a perfectly self-contained or *Ātmapoiesis Universe* [197] (Section 17.2.1). While the term "Ātmapoiesis Universe" originates in philosophical discourse regarding self-contained being, in the IQR framework it is given a precise mathematical definition as the quantum state satisfying the Universal Quantum Constraint:

$$\hat{H}_{\text{univ}}|\mathring{\mathbf{\Psi}}\rangle = 0.$$

This foundational state represents a universe of perfect internal balance, requiring no external reference.

However, the foundational *Axiom of Duality* asserts that to define any entity is to simultaneously define its complement; nothing can exist in isolation (Section 1.2.6). Applied to the cosmos itself, this principle leads to a profound conclusion: a universe defined by self-containment can only be one half of a greater, inseparable whole. This implies the existence of two complementary ontological poles.

The necessary dual to the Ātmapoiesis Universe is the *Parapoiesis Universe* (eq. (17.50)), a state whose existence is defined by its dependence on a transcendent source, $\mathfrak{B}_{\vec{r}}$, transforming the mathematics of self-containment into a formalism of perpetual dependence:

$$\hat{H}_{\text{univ}}|\mathring{\mathbf{\Psi}}\rangle = \mathfrak{B}_{\vec{r}}.$$

The IQR framework dictates that these two possibilities are not mutually exclusive but must coexist in a quantum superposition [197, 3]. The full state of reality, the *Holocosmos state* $|\mathbf{\Omega}\rangle$, is therefore an entangled state of both the atmapoiesis and parapoiesis branches (Definition 17.20 and eq. (17.52)):

$$|\mathbf{\Omega}\rangle = \alpha\,|\mathring{\mathbf{\Psi}}_{\bar{\text{A}}\text{tmapoiesis}}\rangle \otimes |\bar{\text{A}}\text{tmapoiesis}\rangle + \beta\,|\mathring{\mathbf{\Psi}}_{\text{Parapoiesis}}\rangle \otimes |\text{Parapoiesis}\rangle,$$

where $\alpha, \beta \in \mathbb{C}$ are complex amplitudes, and $|\mathring{\mathbf{\Psi}}_{\bar{\text{A}}\text{tmapoiesis}}\rangle$ and $|\mathring{\mathbf{\Psi}}_{\text{Parapoiesis}}\rangle$ are the respective pure states of the universe in each branch [188].

This total state of reality is governed by a higher-order constraint that unifies the two branches into a single, self-consistent system, the *Holocosmos Constraint Equation* (eq. (17.54)):

$$\left(\hat{H}_{\text{univ}} \otimes \mathbb{I}_{\mathfrak{B}} - \mathbb{I}_{\text{univ}} \otimes \hat{\mathfrak{B}}\right)|\mathbf{\Omega}\rangle = 0,$$

where the operators are defined as:

- $\hat{H}_{\text{univ}}$ is the Total Relational Hamiltonian, acting on the universe's state space [156].
- $\hat{\mathfrak{B}}$ is the Source Operator, which distinguishes between the two branches with the eigenvalue structure:

$$\hat{\mathfrak{B}}|\bar{\text{A}}\text{tmapoiesis}\rangle = 0$$
$$\hat{\mathfrak{B}}|\text{Parapoiesis}\rangle = \mathfrak{B}_{\vec{r}}|\text{Parapoiesis}\rangle$$

- $\mathbb{I}_{\text{univ}}$ and $\mathbb{I}_{\mathfrak{B}}$ are the identity operators on their respective spaces.

The foundational and irreducible principle of the treatise is the *Axiom of Duality* [268]. It asserts that to define any entity is to simultaneously define its complement; nothing can be known or can exist in isolation [169]. This principle manifests at every level of reality, from the most fundamental constituents of the cosmos to the very nature of the cosmos itself [197].

The ultimate expression of the Axiom of Duality is found in the nature of existence, representing the two inseparable ontological poles of reality (Section 17.3.1):

1. The *Ātmapoiesis Universe*, corresponding to the $|\overset{\circ}{\mathbf{0}}\rangle$ state. This is the state of perfect self-containment, a universe whose total relational energy vanishes to nullity ($\hat{H}_{\text{univ}}|\overset{\circ}{\mathbf{\Psi}}\rangle = 0$) and whose existence is defined by a perfect internal balance [156].

2. The *Parapoiesis Universe*, corresponding to the $|\overset{\circ}{\mathbf{1}}\rangle$ state. This is the state of being-sourced, of existence defined by its dependence on a transcendent source ($\hat{H}_{\text{univ}}|\overset{\circ}{\mathbf{\Psi}}\rangle = \mathfrak{B}_{\vec{r}}$) and defined by its external relation [188].

This profound symmetry, scaling from the fundamental basis states to the entire cosmos, is given its most potent mathematical expression as a single, inseparable, interdependent pair:

$$|\overset{\circ}{\mathbf{0}}\rangle \leftrightarrow |\overset{\circ}{\mathbf{1}}\rangle$$
$$\hat{H}_{\text{univ}}|\overset{\circ}{\mathbf{\Psi}}_{\bar{\text{A}}\text{tmapoiesis}}\rangle = 0 \leftrightarrow \hat{H}_{\text{univ}}|\overset{\circ}{\mathbf{\Psi}}_{\text{Parapoiesis}}\rangle = \mathfrak{B}_{\vec{r}}$$

($\bar{\text{A}}$tmapoiesis Reality $\leftrightarrow$ Parapoiesis Reality)

The architecture of the Interdependent Quantum Reality framework culminates in a single, unifying mathematical object: the **Algebraic Action of the Timeless Reality of the Holocosmos** (Section 17.3.2). This functional provides the foundation for quantum systems governed by algebraic constraints, transcending conventional field-theoretic approaches that rely on temporal integration [3].

The action is defined not as an integral over time, but as the expectation value of the **Holocosmos Constraint Operator** within the total state of reality, $|\mathbf{\Omega}\rangle$ [3, 197]:

$$\overset{\circ}{\mathbf{\Lambda}}[\mathbf{\Omega}] = \left\langle \mathbf{\Omega} \middle| \left(\hat{H}_{\text{univ}} \otimes \mathbb{I}_{\mathfrak{B}} - \mathbb{I}_{\text{univ}} \otimes \hat{\mathfrak{B}} \right) \middle| \mathbf{\Omega} \right\rangle .$$

This formulation reveals the deep connection between extremal principles and constraint enforcement [269]. The stationary action principle ($\delta\overset{\circ}{\mathbf{\Lambda}} = 0$) is no longer a search for a

"path of least action" through time, but a principle that selects for the state which satisfies the fundamental constraint. Variation with respect to the bra state $\langle\mathbf{\Omega}|$ directly yields the **Holocosmos Constraint Equation** (eq. (17.65)):

$$\left(\hat{H}_{\text{univ}} \otimes \mathbb{I}_{\mathfrak{B}} - \mathbb{I}_{\text{univ}} \otimes \hat{\mathfrak{B}}\right)|\mathbf{\Omega}\rangle = 0 \quad \Longleftrightarrow \quad \left(\hat{H}_{\text{univ}} \otimes \mathbb{I}_{\mathfrak{B}}\right)|\mathbf{\Omega}\rangle = \left(\mathbb{I}_{\text{univ}} \otimes \hat{\mathfrak{B}}\right)|\mathbf{\Omega}\rangle.$$

This purely algebraic action is the mathematical foundation that encodes and unifies the complete structural logic of the IQR framework [14]. This single expression simultaneously captures the theory's most fundamental principles (Section 17.3.2):

- **The Axiom of Duality:** The action emerges from operators $\hat{H}_{\text{univ}}$ and $\hat{\mathfrak{B}}$ operating on distinct, complementary subspaces. This embeds the fundamental duality of reality directly into the mathematical substrate of the theory [268, 169].
- **The Entangled State:** The action functions as a generator over the entangled Metaverse state $|\mathbf{\Omega}\rangle$ [188]. Consequently, any state that extremizes the action must necessarily manifest the entangled structure that correlates the universe with its ontological foundation.
- **The Governing Law:** The constraint equation emerges as the natural consequence of extremizing the action, rather than being imposed as an external postulate [269]. The dynamical law is revealed as the necessary condition for achieving a stationary, self-consistent configuration in the space of all possible states.

The grand architecture of this framework finds its ultimate expression in the *Axiom of Duality* [268, 197]—a principle asserting that to define any entity is to simultaneously define its complement; nothing can exist in isolation. This axiom manifests at every scale of reality, from the foundational quantum states to the very nature of the cosmos itself, dictating that existence must be understood as an irreducible duality of two fundamental ontological modes.

The first mode is the *Ātmapoiesis Universe*—a state of perfect self-containment whose existence emerges from complete internal balance, where the total relational energy vanishes to nullity ($\hat{H}_{\text{univ}}|\overset{\circ}{\mathbf{\Psi}}\rangle = 0$) [197, 3]. This is a cosmos that requires no external foundation, defining itself through its own internal coherence and symmetry. Its necessary counterpart is the *Parapoiesis Universe*—a state of radical dependence whose very being is constituted by its relationship to a transcendent source ($\hat{H}_{\text{univ}}|\overset{\circ}{\mathbf{\Psi}}\rangle = \mathfrak{B}_{\vec{r}}$) [268]. This is a cosmos that exists not through self-sufficiency but through continuous creative relationship with that which exceeds and grounds it.

These two ontological modes are not separate realities but constitute a single, unified system governed by the *Holocosmos Constraint Equation* (eq. (17.54)). This equation emerges from the deeper *Holocosmos Constraint Operator* (Section 17.3), which mathematically encodes the necessary entanglement between self-contained and source-dependent modes of being [188]. Yet, this mathematical formalism is no mere construct; its intrinsically relational nature is the direct consequence of a more primordial truth. The *Axiom of Duality* [268], the paired existence of Ātmapoiesis and Parapoiesis realities, and the very equations that unify them are themselves expressions of the fundamental **Law of Dependent Origination** [17]—the principle that all phenomena arise only through their interdependent relationships. This

universal law provides the ultimate ontological foundation that transcends even the duality it generates [169].

It reveals that the apparent opposition between self-containment and dependence dissolves into a deeper recognition: all existence, whether seemingly autonomous or manifestly parapoiesis, participates in the same fundamental pattern of relational arising. Reality, in its totality, is thus revealed as a seamless web of interdependent emergence—neither purely self-created nor externally imposed, but arising through the creative tension of relationality itself. In this light, the IQR framework becomes more than a unification of quantum mechanics and gravity [14, 8, 3]; it becomes a mathematical expression of the most fundamental insight about the nature of existence: that being itself is relational, and that the deepest truth of reality lies not in any single mode of experience but in the dynamic, creative interdependence that gives rise to all possible worlds.

1.1.1 The Relationship Between IQR and Existing Frameworks

This approach consciously builds upon several influential lines of research while proposing a significant departure from their foundational assumptions:

Algebraic Quantum Field Theory (AQFT): Like AQFT, the IQR utilizes the formal power of operator algebras and the Gelfand-Naimark-Segal (GNS) construction to derive a Hilbert space representation from a state defined over a global algebra (Definition 1.34) [230, 241]. However, where AQFT typically presupposes a fixed background spacetime on which nets of local algebras are defined, our framework inverts this ontological hierarchy. The algebra of PQSs on the foundational $\hat{\mathbf{\Omega}}$–Continuum (Definition 4.617) is primary, and it is from this structure that the emergent quantum configuration space $\mathcal{X}_q$ (Proposition 4.625) and the subsequent chronogeometric manifold $\mathcal{M}_{\text{em}}^{(\mathring{\mathbf{\Psi}})}$ (Definition 5.113) are generated.

Tensor Networks and AdS/CFT: The revolutionary idea that geometry is woven from the threads of quantum entanglement, explored fruitfully in tensor network models and the AdS/CFT correspondence [270, 2, 271, 1], serves as a powerful signpost for our journey. The IQR advances this concept by elevating entanglement from a descriptive tool to a primary ontological constituent. Rather than reconstructing a pre-supposed geometry from a boundary theory, we posit that the relational entanglement encoded by the Quantum Holo-State and manifested via operator correlators like $\hat{\mathbf{G}}_\Lambda$ (Definition 4.9) *is* the geometry. We are thus liberated from the need for discrete lattices or prescribed background spacetimes, aspiring to a theory that is continuous, dynamic, and universal.

Relational Quantum Mechanics and Background Independence: This vision is fundamentally relational, in the tradition of Relational Quantum Mechanics (RQM) and the quest for background independence that animates Loop Quantum Gravity and Causal Set Theory [3, 265, 272]. Yet, where these approaches champion the principle of relationalism, our framework gives it flesh and bone through a concrete operator geometry. The introduction of the operator-valued metric $\hat{g}_{\mu\nu}(x)$ (Definition 5.1) and its tangible, expectation-level coun-

terpart $\bar{g}_{\mu\nu}(x)$ (Definition 5.10) forges an unbreakable, derivable link between philosophical principle and predictive physical theory.

Noncommutative Geometry: Our departure from classical differential geometry aligns with the spirit of Connes' Noncommutative Geometry (NCG), which defines space through the spectrum of operators rather than the assembly of points [4]. The IQR shares this operator-algebraic foundation but places the Quantum Holo-State $|\mathring{\mathbf{\Psi}}\rangle$ in a central, generative role that is absent in NCG. In our formalism, $|\mathring{\mathbf{\Psi}}\rangle$ is the fundamental entity from which all physical and geometric quantities descend, ensuring the framework is intrinsically quantum from its inception.

Topos Theory: Topos-theoretic approaches to quantum foundations offer sophisticated mathematical frameworks that move beyond Hilbert-space-centric formulations, using toposes and sheaves to handle contextuality and logical structure without necessarily assuming a background spacetime [27, 28, 29]. These frameworks excel at reformulating quantum *logic* and the interpretation of states and observables from a categorical or neo-realist perspective, often emphasizing the epistemic or logical structure of quantum theory.

However, while topos theory can beautifully describe the contextual logic of a quantum world, it typically does not aim to derive a full dynamical operator geometry with emergent metric structure and causal relations in the same direct way as the IQR. The IQR approach advances beyond this by presenting the $\hat{\mathbf{\Omega}}$–Continuum as *ontologically physical*, composed of operator-valued singularities rather than abstract logical categories. Our goal is not to perfect the logic of what is, but to unveil the ontology of how it came to be—where emergent geometry is not an abstraction of logic but is constituted by the relational, operator-theoretic entanglement encoded in $|\mathring{\mathbf{\Psi}}\rangle$. The $\hat{\mathbf{\Omega}}$–Continuum is not an abstract category; it is the physical, singular seed of all that is.

Vision and Guiding Principles

The Interdependent Quantum Reality (IQR) framework presents a profound reconceptualization of physical reality: the universe is not a collection of objects moving through spacetime, but a single, indivisible quantum system from which space, time, matter, and forces emerge as different aspects of an underlying relational structure. In this vision, geometry is not the stage upon which physics unfolds—it is the relational fabric itself, woven from quantum entanglement. The laws of physics are not external constraints but necessary consequences of internal quantum coherence.

This revolutionary perspective rests upon four foundational principles that guide the mathematical development:

The Four Guiding Principles

1. **Dependent Origination:** The Law of Dependent Origination is the fundamental principle that governs the generation and interdependence of all phenomena (Sections 1.1.2, 17.1, 17.1.7 and 17.3.2).

2. **The Fourfold Dialectic Algebra of Symmetry:** All *dependently originated* phenomena are completely described by the self-similar, fourfold dialectical algebra([17, 35, 273, 15, 16, 274, 46]) that consists of: Thesis (Symmetry), Antithesis (Antisymmetry (Section 2.1)), Synthesis (Parasymmetry (Definitions 2.85 and 2.94)), and Holothesis (Holosymmetry (Section 2.3)).

3. **Reality is Emergent:** The continuum on which *dependently originated reality* is defined is the $\hat{\mathbf{\Omega}}$-Continuum (Section 4.18) of fibred *ultra-sub-Planckian* Primordial Quantum Singularities (Definition 1.25). Each primordial quantum singularity is endowed with the Fourfold Dialectic Algebra of Symmetry. All geometric and temporal properties—including chronogeometric spacetime and causality—emerge from this relational substrate as the expectation values of fundamental, operator-valued chronogeometric operators (Section 5.2.6), evaluated in the Quantum Holo-State $|\mathring{\mathbf{\Psi}}\rangle$ over this continuum of singularities.

4. **The Generative Quantum Holo-State:** The Quantum Holo-State ($|\mathring{\mathbf{\Psi}}\rangle$ (Definition 1.34)) is a single, non-factorizable quantum state that both encodes and generates the indivisible reality in which it resides. This is the singular wellspring from which all physical structure emerges. It is this state that gives rise to the *Fourfold Inter-Modal Correlator Operator* (Definition 4.9), the very engine of relational dynamics. From its coherent structure, the inner product and the geometry of the *emergent Hilbert spaces* (Section 4.1) are born. Furthermore, this universal state projects itself into the two fundamental facets of existence: the *Gaiamiti Field*, whose quanta, the *apeirons*, are the progenitors of the graviton and the fabric of chronogeometry itself; and the *Tantrakyma Field*, whose quanta, the *kymons*, blossom into the complete edifice of the Standard Model: its rich and varied spectrum of particles, and the fundamental interactions that govern their existence.

1.1.2 The Law of Dependent Origination

At the very heart of both ancient Buddhist philosophy and the relational fabric of our emergent cosmos, lies the **Law of Dependent Origination**, or *Pratītyasamutpāda* [275, 276]. It is a principle of dynamic continuity of interdependence and entanglement, a declaration that no phenomena, from the fleeting thought of a sentient being to the grandest cosmic structure, possesses any intrinsic, independent existence. Instead, all phenomena arise in a seamless, unbroken interwoven cyclic web of mutual causation, conditioning, and effects. The principle is captured with sublime economy in the ancient formula:

> *When this is, that is.*
> *From the arising of this, that arises.*
> *When this is not, that is not.*
> *From the cessation of this, that ceases.*

On the surface this may seem like a statement of linear cause and effect. However, this is far from the truth, for it describes a universe where reality is not a collection of isolated

objects but a dynamic, living network of relationships. The statement of the existence of any single entity is parapoiesis upon the existence of all others; its identity is woven from the threads of its conditions, and it, in turn, becomes a condition for the arising of what is yet to be.

The Architecture of Experiential Existence

To illustrate the workings of this law, particularly in the realm of sentient experience, the teaching is often presented as a great causal chain of twelve interdependent links, or *nidānas* [275, 277]. This sequence is not a simple temporal progression but a cyclical, self-perpetuating structure that describes the very mechanism of entanglement in the cycle of existence, or *samsara* [278]. The twelve-link chain of dependent origination begins with **(1)** *ignorance (avidyā)*—a fundamental misapprehension of reality itself [279]. This ignorance manifests as the deep-seated delusion that phenomena possess an independent, enduring self-nature, a misunderstanding that serves as the root condition for all subsequent suffering.

Conditioned by this basic ignorance, **(2)** *formations (saṃskāra)* arise as volitional impulses and karmic imprints. These formations represent the relational patterns of intention that shape the trajectory of future experience, carrying the momentum of past actions into present and future moments. From these accumulated formations emerges **(3)** *consciousness (vijñāna)*, the basic stream of awareness that provides continuity across moments and lifetimes, linking discrete experiences into a coherent flow of subjective experience [275]. This consciousness, in turn, conditions the arising of **(4)** *name and form (nāmarūpa)*—the psycho-physical organism that constitutes the living being [277]. Here, "name" encompasses the mental phenomena of feeling, perception, and intention, while "form" refers to the physical body and material world. Together, they create the inseparable duality of mind and matter that characterizes embodied existence [279].

The psycho-physical organism naturally develops **(5)** *the six sense bases (Ṣaḍāyatana)*—the five physical senses along with the mind itself as the sixth faculty of perception [275]. These sense bases serve as the interface between consciousness and its environment, establishing the conditions for experiential encounter with the world. When a sense faculty meets its corresponding object in the presence of consciousness, **(6)** *contact (sparśa)* occurs. This represents the raw moment of sensory input, the immediate meeting of perceiver and perceived that forms the foundation of all experiential content [277]. From each instance of contact arises **(7)** *feeling (vedanā)*—the immediate, pre-conceptual experience of pleasantness, unpleasantness, or neutrality that colors every moment of awareness [275].

These feeling-tones, however subtle, condition the arising of **(8)** *craving (tṛṣṇā)* [279]. Pleasant feelings generate the desire for repetition and continuation, while unpleasant feelings fuel the impulse toward escape and avoidance. This craving, when it becomes habituated and intensified, transforms into **(9)** *grasping (upādāna)*—a deep-seated clinging to existence, sensory experience, views, and the notion of selfhood itself. The accumulated energy of grasping fuels the process of **(10)** *becoming (bhava)*, creating the karmic momentum that propels the stream of consciousness toward future manifestation. This momentum conditions **(11)** *birth (jāti)*—the arising of a new psycho-physical form, complete with its inherited patterns and predispositions [278].

Finally, with birth comes the inevitable reality of **(12)** *aging and death (jarā-maraṇa)*—the

entire cycle of suffering including decay, disease, separation, and dissolution [277]. Rather than leading to liberation, these experiences of impermanence typically reinforce the original ignorance, deepening the delusion of selfhood and perpetuating the cycle indefinitely. Thus the wheel of dependent origination completes one revolution while simultaneously beginning another, creating an unbroken chain of conditioned existence that continues until the root ignorance is finally recognized and abandoned [279].

Philosophical Implications: The Middle Way

The Law of Dependent Origination is the foundation of what is known as the **Middle Way**. It steers a careful course between the two philosophical extremes of **eternalism** (the belief in a permanent, unchanging self or substance) and **annihilationism** (the belief that things cease to exist entirely). By revealing all phenomena as dependently arisen, it shows that nothing is permanent, yet the causal web ensures a continuity of process.

For the great philosopher *Nāgārjuna*, understanding Dependent Origination was synonymous with understanding reality itself. He famously equated it with the principle of **emptiness** (Śūnyatā): "It is dependent origination that we call emptiness" ([278, 35, 279]). Because all phenomena arise in dependence on conditions, they are *empty* of any intrinsic, self-sufficient nature. Emptiness, in this context, is not a void, but the very mark of relational existence. It is the ultimate expression of a universe that is not a collection of things, but a single, indivisible, and interdependent whole. The twelve-link chain of dependent origination reveals the interconnected nature of existence, but one relationship stands as particularly crucial to Buddhist understanding of reality: the reciprocal interdependence between *consciousness (viññāṇa)* and *name-and-form (nāmarūpa)*.

In Buddhist terminology, "name" encompasses the mental dimensions of experience—feeling, perception, and intention—namely, the cognitive processes that *name, label, categorize, and construct* our sensory encounters. "Form" by contrast, refers to the physical realm: our bodily existence and the material world around us—*to which we also assign* "names"—and the cognitive processes that *name, label, categorize, and construct* our sensory encounters. Together, these components *constitute* what we recognize as the complete psychophysical being.

Name and Form: A Duality of Reality The *Mahānidāna Sutta (The Great Discourse on Causation)* [275, 276] presents the Buddha's most penetrating analysis of this interdependence. Speaking to his disciple Ānanda, the Buddha reveals that consciousness and the experienced world do not exist as separate entities in a simple cause-and-effect relationship. Rather, they emerge together in an indivisible process of mutual arising. As the Buddha teaches [275, 276]:

> From name-and-form as a requisite condition
> comes consciousness.
> From consciousness as a requisite condition
> comes name-and-form...
>
> — Dīgha Nikāya 15

Buddha illuminates this perfect interdependence through a striking metaphor: consciousness and name-and-form resemble two bundles of reeds leaning against each other for support.

Remove either bundle, and both immediately collapse [275]. This teaching fundamentally contrasts with the Western philosophical tradition, which typically assumes a conscious subject existing independently of an objective world it observes [179, 182]. Buddhist insight reveals instead that the *"knower" (consciousness)* and the *"known" (the phenomenal world)* arise simultaneously as aspects of a unified experiential event [279, 278]. Consciousness cannot exist without objects to be conscious of, just as no experienced world can manifest without consciousness to apprehend it. They represent not separate entities but complementary facets of a single, seamless process of aware experience [35].

This manuscript embraces the Law of Dependent Origination as its foundation, focusing particularly on the entanglement of the *Name and Form (Nāmarūpa)* dynamic. Our investigation centers on the relationship between the observer and the universe of physical phenomena—a universe that simultaneously constitutes the observer and serves as their environment. This encompasses both the chronogeometric structure of spacetime and the fundamental particles described by the Standard Model of Physics.

1.1.3 The Tetralemma: Reality's Four-Vertex Prism

The *tetralemma*, known in Sanskrit as the **Catuṣkoṭi चतुष्कोटि ("four corners")**, is a sophisticated logical structure that serves as a cornerstone of Madhyamaka Buddhist philosophy [15, 35]. Far from being a simple statement of facts, it is a powerful dialectical tool designed to systematically deconstruct our most deeply ingrained assumptions about the nature of existence. It challenges the very foundation of classical, binary logic—the idea that a proposition must be either true or false—by exploring a complete set of four possibilities [273, 16].

The four *corners* of the tetralemma are typically presented as a set of four exhaustive propositions concerning a given phenomenon:

1. *It is (Being):* The first corner affirms the existence of a thing, representing our everyday, common-sense view where objects have a solid, independent reality. The most direct Sanskrit term is Asti (अस्ति), the simple, affirmative third-person singular of the verb *to be*. The nominal or conceptual form is often given as Bhāva (भाव), meaning "being", "existence", or "state", or as Sat (सत्), meaning "the real."

2. *It is not (Not-Being):* The second corner posits the negation or non-existence of the thing, the simple opposite of the first position. This is directly translated as Nāsti (नास्ति), a contraction of *na asti* ("it is not"). The corresponding conceptual terms are Abhāva (अभाव), meaning "non-being" or "absence", and Asat (असत्), or "non-being."

3. *It both is and is not (Both Being and Not-Being):* The third corner entertains the seemingly contradictory possibility that a phenomenon can simultaneously exist and not exist, expressed in Sanskrit as Asti ca Nāsti ca (अस्ति च नास्ति च), where *ca* means "and." This challenges the classical law of non-contradiction, suggesting a reality where opposing states can co-exist [274]. A more philosophically dense term for this concept is the compound Sadasat (सदसत्), which combines being (*sat*) and non-being (*asat*) into a single notion.

4. *It Neither Is nor Is Not (Neither Being nor Not-Being, nor both Being and Not-Being):* The fourth and final corner, Naiva Asti na Nāsti (नैवास्ति न नास्ति), literally "neither is it, nor is it not", is the most profound [46]. While traditional interpretations see this as a simple "none of the above", the more profound understanding, particularly in the context of masters like Nāgārjuna, is that it represents a *categorical shift*. The treatise will demonstrate that *Naiva Asti na Nāsti* is not a simplistic negation *of* the first three positions at all, but alludes to a concept far more subtle and complex, a transcendence *beyond* each of the first three.

The Tetralemma as a Philosophical Lense

The great philosopher *Nāgārjuna* employed the tetralemma not to arrive at one of the four corners as a final answer, but to systematically demonstrate that *none of them* can adequately describe the *ultimate nature of reality* [17, 15]. By applying this four-fold analysis to fundamental concepts such as causation, motion, and the self, he revealed that any attempt to pin down a phenomenon with a fixed, inherent identity leads to logical paradoxes [274]. For example, when examining the concept of a "cause", Nāgārjuna would argue that an effect cannot be produced from itself (the first corner), from something other than itself (the second corner), from both (the third corner), or from neither (the fourth corner) [16]. By demonstrating the incoherence of all four possibilities, he leads the intellect to a profound conclusion: the very idea of a "cause" as an independently existing entity is a conceptual fiction.

The Path to Emptiness or Śūnyatā

The ultimate purpose of the tetralemma is to guide the mind toward an understanding of *emptiness* (*Śūnyatā*) [35]. It operates on dependently originated phenomena by demonstrating that any attempt to predicate an *intrinsic, ultimate nature (svabhāva)* upon them—using any of the four logical positions of affirming, negating, both, or neither—inevitably leads to irresolvable contradiction [279, 46]. The tetralemma thus reveals that all conditioned things are "empty" of such an independent existence. This "emptiness" is the very mark of *Dependent Origination*. Because things are empty of an inherent self, they are free to arise, interact, and cease in a dynamic, interdependent web of relationships. The failure of the four corners is not a failure of logic; it is a successful demonstration that our conceptual categories, which presuppose inherent independent existence, do not apply to a reality that is fundamentally entangled and relational [273].

1.1.4 The Quantum Revelation: A Tetralemmatic Journey

The profound intellectual upheaval of twentieth-century physics can be understood as a necessary journey through the four exhaustive propositions of the ancient tetralemma [15, 273]. It is as if the universe, when probed at its most fundamental level, compelled physicists to abandon the simple logic of objective reality and embrace a more subtle and paradoxical structure of existence. As this exploration will show, their discoveries already hinted at the

application of the tetralemma to understand the nature of reality, revealing a dynamic cycle from classical certainty to quantum potentiality and back again [35].

To appreciate the significance of this scientific journey, we must first look back to an ancient fork in the road of human thought. In the ancient world, humanity's quest to understand reality arrived at a great intersection. The questions were the same everywhere: What does it mean for something "to be"? What is the nature of nothingness? Can opposites exist at the same time? But from this single point, two profoundly different journeys began.

In the East, particularly in the philosophical schools of India, thinkers looked at the four great propositions of reality—that a thing *is*, that it *is not*, that it is *both*, and that it is *neither*—and chose to embrace them all. For them, the tetralemma was a complete compass for navigating the complexities of existence [15]. They saw the paradoxes of the third and fourth corners not as errors in logic, but as profound insights [46, 274]. They used these ideas as a sophisticated toolkit to deconstruct our common-sense reality, seeking to understand a world where opposites could coexist and where the ultimate truth might lie beyond all concepts of being and non-being [17, 35].

In the West, the Greek philosopher *Aristotle* stood at the same crossroads and saw the same paradoxes. But instead of embracing them, he chose to build a fortress against them. He declared that rational thought required absolute clarity. His *Law of Non-Contradiction* became an unbreakable wall against the third corner, and his *Law of the Excluded Middle* barred the fourth, insisting that a statement must be either true or false. For two thousand years, Western thought lived and thrived within this powerful fortress of binary logic, building the foundations of modern science, law, and philosophy [73]. In mathematics, this binary clarity was formalized via set theory's crisp membership relations [280].

The great irony of history is that the inheritors of Aristotle's tradition—the physicists of the twentieth century—used his flawless logic to build tools so powerful they could probe the very heart of matter. And there, in the quantum realm, they were stunned to find the ghosts of the ideas their philosophical ancestors had banished. They found a world of superposition where a particle behaved as if it were *both* a particle and a wave [281, 168]. They found a realm of pure potentiality where a property *neither* existed nor did not exist until it was measured—classical definiteness emerging only through interaction with the environment (decoherence) [282].

To describe the universe at its most fundamental level, they had to break out of Aristotle's fortress. Unknowingly, the pioneers of quantum mechanics found themselves charting a course with the same four-pointed compass that Eastern philosophy had been using all along, finally reuniting the two great paths of human thought at the very frontier of science [16, 35].

It is: The Classical World and its Negation

The journey begins with the *first corner: It is*. This is the thesis of classical physics and mathematics, a worldview resting on the bedrock of Sir Isaac Newton's mechanics [159]. It affirmed a deterministic,

clockwork universe where reality was objective, particles followed predictable trajectories, and logic was absolute [15, 273]. In this reality, an element *is* a member of a well-defined set, or it is not—there is no middle ground. This clean, binary logic, formalized in set theory, created a cosmos of clear lines and unambiguous boundaries [280]. It not only gave us the

laws that could plot the course of planets but also became the engine of our digital age, where every computation boils down to a choice between a one and a zero [283].

This powerful framework was shattered by the *second corner: It is not*. This represents the antithesis—the dawn of the quantum age, marked by phenomena that classical physics could not explain. When Max Planck reluctantly proposed that energy was not continuous but was emitted in discrete *quanta*, he was making a profound statement of negation [284]. Reality, at its core, was *not* the smooth continuum that classical intuition demanded, and the first crack in the classical foundation appeared.

It Both Is and Is Not: The Paradox of "Both/And"

This negation led inexorably to the *third corner: It both is and is not*. This is the domain of quantum paradox, where contradictory properties are found to coexist in a single entity. The quintessential example is *wave-particle duality*. A quantum object, like an electron, is revealed to be *both* a localized particle (*it is*) and a delocalized wave (*it is not*). The double-slit experiment provides the irrefutable evidence, where a single electron "seemed to pass through *both* slits simultaneously like a wave and interfere with itself" [281].

This paradoxical synthesis is given its most unsettling expression in the famous thought experiment of *Schrödinger's Cat*, which, when unobserved, is described by a wave function that represents it as being "simultaneously *alive and dead*" [171, 172, 173]. Until an observation is made, the cat exists in a state of pure superposition, a ghostly blend of both possibilities at once. The act of observation is what forces reality to choose, collapsing the

both/and state into the simple

is or

is not of our classical world—an emergence often explained in terms of decoherence [282].

It Neither Is nor Is Not: Transcending "Neither/Nor"

Yet, the deepest reality lies in the *fourth corner: It neither is nor is not*. This represents a transcendence of all prior categories, a state of pure potentiality where our very questions about reality fail. This is the nature of the unobserved quantum state. As *Heisenberg's uncertainty principle* revealed, an electron prior to measurement "simply does not possess a definite position and a definite momentum at the same time" [285]. Far from being just an unknown, the very category of "having" that property is inapplicable. This is powerfully affirmed by the principle of *Measurement as Creation*, where, as Pascual Jordan stated, "Observations not only disturb what is to be measured, they produce it" [194]. The property was *neither* there *nor* not there; it was brought into being.

We encounter this frontier elsewhere in modern physics. Consider two *entangled particles*. After interacting, they form a single, inseparable reality, even when flying apart to opposite ends of the universe [178, 183, 184]. One particle doesn't have a definite state on its own; it isn't *spin-up* nor is it *not spin-up*. Its identity is so completely tied to its partner that its properties are only defined within the relational system of both [3]. We meet this limit again when we ask what lies at the heart of a *black hole*. Is there a singularity? Is there not? The question itself may be meaningless, as this is the point where our language—the laws of physics—is torn apart [240, 7].

This same principle echoes in the abstract world of mathematics through *Gödel's Incompleteness Theorems*. These theorems prove that in any sufficiently complex logical system, there will always be statements that are true but can never be proven *within the rules of that system* [286]. Such a statement neither *is* provable nor *is not* provable; its status transcends the system's own framework. The ultimate expression of this transcendent view comes from Niels Bohr himself: *"There is no quantum world. There is only an abstract quantum physical description"* [193].

The Cycle of Reality and the Concept of Symmetry

The fourth corner describes the humbling edge of our knowledge, the point where the universe refuses to answer our questions in the terms we know how to ask them [3]. Finally, the very act of observation completes the cycle. The "collapse of the wave function", a central tenet of the Copenhagen interpretation, is the mechanism that forces reality to exit the subtle logic of the third and fourth corners [168, 169]. Upon measurement, the wave of potentiality instantly "collapses" into a single, definite state [195, 124], forcing a choice. The qubit becomes a 0 (*It is*) or a 1 (*It is not*) [287]. In contemporary accounts, the emergence of classical definiteness from quantum superposition is often analyzed via decoherence and einselection [282, 170]. The measurement, therefore, is the act that translates the transcendent and paradoxical nature of the quantum realm into the simple, binary logic of the classical world we perceive. The entire quantum process, from unobserved potentiality to definite reality, thus unfolds precisely along the ancient path of the tetralemma.

This revelation—that the fundamental processes of the universe trace the contours of this ancient four-fold logic—demands a re-interpretation. In *Nāgārjuna's* treatment, the tetralemma appears to allude to a profound concept, as if it were a blueprint for reality's underlying architecture. To translate this ancient formula into the language of modern physics, we must identify its core principle. This treatise proposes that the four corners are, in fact, distinct generalizations of a single, foundational concept: **Symmetry**. We must therefore redefine our understanding of **Symmetry in its holistic sense**. We reason that it should be liberated from the simple notion *of a singular* property of invariance, into a more complete, holistic architecture of reality—a system composed of four distinct but interdependent modes. This unified structure is best conceptualized through the geometric metaphor of a **tetrahedron**.

The **triangular base** of this prism represents the immanent logical plane. Its three vertices are the foundational modes of existence (see Section 1.2.7 for a formal definition and justification for the neologisms):

1. **Symmetry (Thesis):** The principle of identity (Definitions 2.1 and 2.12).
2. **Antisymmetry (Antithesis):** The principle of distinction.
3. **Parasymmetry (Synthesis):** The principle of coherent interaction and superposition (Section 2.2 and Definitions 2.85 and 2.94).

These three corners define the relational dynamics within a given system. The **apex** of the tetrahedron represents the fourth corner, **Holosymmetry** (Sections 2.3, 3.8 and 12.2). It

occupies a position transcendent to the base, connected to all three vertices yet existing on a higher structural level. From this apex, the overarching laws that govern the entire system emerge. It is the symmetry *of* the system itself, not just a state *within* it. Thus, **Symmetry**, in its fullest sense, is the tetrahedral whole—the indivisible union of the planar base and the transcendent apex. It is a complete, "four-fold jewel" whose facets describe the total architecture of reality, from simple identity to the unifying laws that govern the cosmos. To fully appreciate the power of this conceptual model, we turn to a physical analogy: the prism. Just as a physical prism takes what appears to be a single, unified beam of white light and reveals the rich, hidden spectrum of colors within, the conceptual "prism of symmetry" takes a seemingly singular phenomenon and reveals its full logical spectrum.

A single beam of sunlight appears pure and unified as it travels through air, moving in a straight line until it encounters a prism. At the glass surface its path changes by **refraction**, governed by Snell's law $n_1 \sin\theta_1 = n_2 \sin\theta_2$ [26, 288]. A deeper transformation occurs because the refractive index depends on wavelength: **dispersion** [289, 26]. Inside the prism, violet light (shorter wavelength, higher frequency) typically experiences a larger refractive index than red light and is therefore deviated more strongly (normal dispersion) [26, §3.6–3.7]—except very near absorption lines, where some materials exhibit *anomalous dispersion*; in the visible for common glasses the account above holds [289, §§2.5–2.6]. After refraction at the second face, the colors separate further, producing an ordered spectrum from red through violet [289, Ch. 8]. While dispersion "makes rainbows", prisms also serve other purposes: in binoculars and periscopes they rely on *total internal reflection* to redirect and erect images (minimizing dispersion) [26, §4.4, §6.3], whereas in spectrometers they deliberately exploit dispersion to split light by wavelength [289, 26, Ch. 8].

1.1.5 The Middle Way: A Narrative of Insight

The *tetralemma* represents a complete, tetradic net of logic, a philosophical prism designed to illuminate the very structure of conceptual thought. It maps every possible position the mind can adopt when using the dependently originated tools of language, signs, and symbols to reason about reality. The Buddha, in the foundational *Kaccānagotta Sutta*, diagnosed the root of this conceptual error: when we make absolute statements about the world, our reasoning invariably collapses into two ontological extremes—the assertion of absolute *existence* (atthitā), which leads to eternalism, or the assertion of absolute *non-existence* (natthitā), which leads to annihilationism [275]. This polarity is the fundamental duality the path of wisdom seeks to transcend.

This principle moves beyond abstract thought. The *Acela Sutta* demonstrates it practically through Kassapa's questions about suffering. The query, *"Is suffering created by oneself?"* implies a permanent, existing self (eternalism), while *"Is it created by another?"* implies a disconnected self (annihilationism). By rejecting both, the Buddha reveals that all phenomena arise not from a fixed entity but through the intricate web of *dependent origination* [276].

The Mahayana sutras elevate this principle to its ultimate expression. The *Heart Sūtra* crystallizes this insight in its famous declaration: *"Form is emptiness; emptiness is form."* [290] This is the Middle Way articulated as a statement of perfect non-duality, a direct path that cuts through the endless oscillation between the two extremes. The *Diamond Sūtra* then provides the methodological key, employing its powerful dialectical formula, *"What is called*

X, is not X, therefore it is called X," to deconstruct our habits of conceptual grasping [291]. It reveals that we give names not to independently existing things, but to relational processes.

The Position of Having No Position

The philosophical position of the Middle Way is unique in that its final aim is to deconstruct the mind's habit of clinging to *any* fixed conceptual view (dṛṣṭi), including its own. The great master Nāgārjuna, the chief architect of this school, famously stated that he has no thesis of his own to prove, for his method is to reveal the internal contradictions within all other positions that assume an intrinsic, independent reality (svabhāva) [278]. It is akin to a raft. The raft is a definite, well-constructed object (a philosophical position) that is absolutely necessary to cross a river. But once you have reached the other shore (enlightenment), you do not carry the raft on your back. Its ultimate purpose is to be transcended and let go of [277].

The Insight and its Quantum Parallel

From the early discourses to the grand vision of the Mahayana, the central insight remains unchanged: the Middle Way is a unique philosophical position whose ultimate purpose is to dismantle the very act of position-taking [17, 279]. This ancient wisdom finds its most startling and precise modern parallel in the quantum realm. To insist that a quantum state must be either '0' or '1' is to fall into the classical extremes, misapprehending its true nature which allows for a superposition of both [287, 282]. The error lies in settling into these two extremes, a conceptual fixation that traps the mind in an endless cycle. The path of wisdom, the Buddha taught, is a *Middle Way* that transcends this false dichotomy by understanding that a reality governed by the interdependent web of Dependent Origination can never be captured by the fixed, binary categories of our conceptual thought [278, 273].

1.1.6 Aristotle and The Story of a Thesis

Imagine the bustling intellectual world of ancient Greece, where ideas were traded in open forums and debated in shaded porticos. In this world, the word *thésis* (θέσις) was a simple one, meaning "a putting" or "a placing." This simple word carried an ancient echo, descending from a Proto-Indo-European root, $*d^{h}eh_{1}$-, which meant the physical act of setting something down [81, 292]. From this concrete origin grew its intellectual meaning: the common act of setting an idea forth for discussion, a proposition laid bare for others to see [293].

Then came *Aristotle*, a philosopher who saw a need for order and rigor in these sprawling debates. He looked at the simple act of *thésis* and forged it into a sharper tool of dialectic within his systematic treatment of argumentation; a thesis became a determinate position to be defended by reasons [294, 295]. In his hands, it transcended its everyday meaning: a thesis could be a bold or even paradoxical claim that challenged common opinion (*endoxa*) and demanded a disciplined defense within a structured inquiry [294, 295]. He transformed the word from a simple statement into a defensible intellectual position, the cornerstone of ordered debate in classical philosophy [294, 295].

This powerful concept—a central claim that must be rigorously defended—traveled through the ages. It echoed in the halls of Roman orators, was preserved in the scriptoriums

of medieval monasteries, and was reborn in the great universities of Europe. Today, when a doctoral candidate stands to defend their *thesis*, they continue this tradition: a single, novel proposition formally set forth and defended before experts—an institutional sense attested in the historical record of English usage [296].

1.1.7 The Debates of Ancient India

Journey back to the vibrant intellectual landscape of classical India, where scholars and ascetics gathered under banyan trees or in grand debate halls to challenge the very nature of reality. These were not casual conversations; they were crucibles of logic, formal, rule-based duels of the mind known as *vāda* (वाद) [33, 297, 18].

A debate would begin when a philosopher rose to address the assembly. They would start by making a bold, clear declaration—a precise statement they intended to prove. This opening proposition, the seed from which the entire argument would grow, was known as the *pratijñā* (प्रतिज्ञा) [33, 298, 18]. This was their thesis, the central claim upon which their intellectual honor rested. It could be a statement about the nature of the soul, the cause of suffering, or the fabric of the cosmos.

The Sanskrit *pratijñā* (प्रतिज्ञा) is a compound word, and both of its parts have clear PIE origins:

- *prati-* (प्रति-): a prefix meaning "against," "towards," or "in return," from the PIE form **proti-* with the sense of facing or moving towards [298, 81, 292].
- *jñā* (ज्ञा): a verbal root meaning "to know" or "to perceive," from the PIE root **ǵneh₃-* "to know," which is also the source of Greek *gnōsis* (γνῶσις) and English *know* [298, 81, 293].

When combined, the deep etymological meaning of *pratijñā* is something like "to know towards" or "to declare against" an opponent: a solemn public claim that forms the basis of the argument [298].

Once the *pratijñā* was stated, it had to be defended. The philosopher would proceed to build a fortress of logic around their claim, guided by the exacting principles of schools such as *Nyāya*. They would present a reason (*hetu*), support it with an undeniable, real-world example (*udāharaṇa*), and methodically construct a five-membered argument—*pratijñā, hetu, udāharaṇa, upanaya, nigamana*—designed to be unassailable [33, 18, 297].

The opponent would, in turn, scrutinize every link in this logical chain, testing it for fallacies and contradictions. The ultimate goal of a true *vāda* was something more profound than simple victory: the shared pursuit of truth (तत्त्वनिर्णय, *tattva-nirṇaya*) [33, 297]. The debaters were participants in a structured process designed to cut away confusion and reveal the core of the matter.

In this ancient tradition, we see a powerful parallel to the world of Aristotle. While separated by thousands of kilometers, both cultures arrived at the same fundamental conclusion: the surest path to knowledge is to put forth a clear, defensible claim and subject it to the purifying fire of rigorous, logical argument [294, 295, 18].

1.2 The Tetralemma of Algebra

1.2.1 The Definition of Thesis

A *Thesis* is a determinate, context-indexed position that affirms existence (Asti/Bhāva/Sat) and is normatively bound to a rigorous mode of defense—Aristotelian in the sense of a reasoned position suitable for dialectical scrutiny, and Nyāya in the sense that the *pratijñā* is to be validated by a methodical, multi-membered argument—so that its *epistemic warrant* derives from the coherence and validity of its logical support.

Definition 1.1 (Thesis (*Asti*/अस्ति; *Bhāva*/भाव; *Sat*/सत्; Greek *thésis*; Sanskrit *pratijñā*))**.** Fix a domain of discourse $\mathcal{D}$ and an immanent context $\mathcal{C}$ (criteria of individuation and admissible inference). A *Thesis* about a formula φ is a structured act

$$\mathbf{Th}(\varphi;\mathcal{C}) \;:=\; \Big\langle \underbrace{\exists^{\mathcal{C}} x\, \varphi(x)}_{\text{(Being) Asti/Bhāva/Sat}} \;,\; \underbrace{\mathcal{J}_A(\varphi)}_{\text{Aristotelian defense}} \;,\; \underbrace{\mathcal{J}_N(\varphi)}_{\text{Nyāya defense}} \Big\rangle,$$

subject to the following constitutive conditions:

1. **Ontological affirmation (Being).** The content asserts existence *relative to* $\mathcal{C}$—the "It is": a determinate item of discourse is affirmed as real (Asti), i.e., an instance of *bhāva*/*sat*. *(Catuṣkoṭi: first corner).*
2. **Determinacy and reference.** The affirmed item meets $\mathcal{C}$'s criteria of individuation and sustains stable reference adequate for logical manipulation (identity and non-contradiction hold *within* $\mathcal{C}$).
3. **Dialectical posture (Aristotle).** $\mathcal{J}_A(\varphi)$ is a position to be defended by reasons, i.e., fit for ordered debate and syllogistic support from suitable *endoxa*.
4. **Procedural obligation (Ancient India).** $\mathcal{J}_N(\varphi)$ instantiates the Nyāya five-membered proving scheme with *pratijñā* as the opening commitment:

$$\underbrace{\text{pratijñā}}_{\text{thesis}},\ \underbrace{\text{hetu}}_{\text{reason}},\ \underbrace{\text{udāharaṇa}}_{\text{example}},\ \underbrace{\text{upanaya}}_{\text{application}},\ \underbrace{\text{nigamana}}_{\text{conclusion}}.$$

Equivalently, writing $\exists^{\mathcal{C}}$ for context-relative existence and letting $\mathrm{Arg}_{A/N}$ denote well-formed Aristotelian/Nyāya arguments, we may say:

$$\mathbf{Th}(\varphi;\mathcal{C}) \text{ holds} \iff \big(\exists^{\mathcal{C}} x\, \varphi(x)\big) \wedge \big(\mathcal{J}_A(\varphi) \in \mathrm{Arg}_A\big) \wedge \big(\mathcal{J}_N(\varphi) \in \mathrm{Arg}_N\big).$$

Remarks. (i) *Thesis* unifies *Asti*/*Bhāva*/*Sat* (ontic affirmation), Greek *thésis* (a defended position), and Sanskrit *pratijñā* (the formally announced claim). (ii) It is explicitly *immanent* and context-relative; no claim of ultimate *svabhāva* is made. Its role is to secure determinacy and justify claimhood within $\mathcal{C}$; the thesis's *warrant*, not metaphysical truth simpliciter, turns on the adequacy of its defense.

Thesis, in this framework (Section 1.2.1 and Definition 1.1), is not a dogmatic "yes" but a *preference the system displays* for a particular mode. Formally it is the projector $P_T := |e_1\rangle\langle e_1|$ onto the privileged Thesis state $|e_1\rangle$ from the ontological duality basis (Definition 2.1), and

its "truth value" in a state ρ is the Born valuation $v_{\text{Thesis}}(\rho) = \text{Tr}(P_T\rho)$ (Definition 1.18). Intuitively, that number tells us how much of the system's identity presently aligns with the Thesis pattern—like a dimmer, not an on/off switch. The meaning of Thesis is thus *basis-defined*: it is the tendency to show up as $|e_1\rangle$ when we interrogate the system along the distinguished Thesis–Antithesis axis in the $(0, 1)$-sector (Definition 2.1). In this sense, to "affirm" the Thesis is to say, "given this instrument orientation, the system is disposed to register in this symmetry mode."

Operationally, Thesis becomes a *repeatable test*. Measure in the Thesis/Antithesis basis defined by $\{|e_1\rangle, |e_2\rangle\}$; the observed frequency of the $|e_1\rangle$ outcome approaches $v_{\text{Thesis}}(\rho)$ by the Born rule (Definitions 1.18 and 2.1). We can also enact Thesis as a *filter*: apply P_T to prepare or post-select states that realize the Thesis mode, then study how subsequent dynamics treat that assertion (e.g., unitaries commuting with P_T preserve it; those that rotate into the complementary direction redistribute its weight). This makes Thesis experimentally legible: a concrete knob (choice of basis), a concrete readout (projective/effect measurement), and a concrete number (the probability) that plays the role of truth in the dialectic semantics (Definition 1.18).

Finally, Thesis lives alongside its contrasts without contradiction. A generic ρ can give *nonzero weight to both* Thesis and Antithesis because "truth" here is a probabilistic disposition, not mutual exclusion (Definition 1.18); only in the classical limit $v_{\text{Thesis}} \to 1$ (or 0) do we recover crisp bivalence. States orthogonal to the Thesis–Antithesis plane—i.e., outside $\mathcal{H}_{(0,1)} \subset \mathbb{C}^4$—naturally instantiate *"Neither,"* signaling that the very question "Thesis or Antithesis?" is inapplicable to that preparation (Definition 2.1). Synthesis then enters as the *complementary probe* in the same sector, realized by the parasymmetric mode and its projector P_{para} (Section 2.2 and Definitions 2.16 and 2.17), with the same Born-valued semantics linking "degree of synthesis" to $\text{Tr}(P_{\text{para}}\rho)$ (Definition 1.18).

1.2.2 The Definition of Antithesis

An *Antithesis* is a determinate, context-indexed denial that withholds existence (Nāsti/Abhāva/Asat) and is normatively bound to a rigorous mode of refutation—Aristotelian in the sense of a reasoned negation suitable for dialectical scrutiny, and Nyāya in the sense that the asserted absence is to be validated by a methodical argument (with *pratijñā* now a negative claim)—so that its *epistemic warrant* derives from the coherence and validity of its logical and evidential support.

Definition 1.2 (Antithesis (*Nāsti*/नास्ति; *Abhāva*/अभाव; *Asat*/असत्; Greek *antíthesis*))**.** Fix a domain of discourse $\mathcal{D}$ and an immanent context $\mathcal{C}$ (criteria of individuation, admissible inference, and admissible loci of evaluation). Let $\Lambda_{\mathcal{C}}$ be the set of spatiotemporal loci sanctioned by $\mathcal{C}$, and let $\lambda \in \Lambda_{\mathcal{C}}$.

An *Antithesis* denying a formula φ at locus λ is a structured act

$$\mathbf{AntiTh}(\varphi; \mathcal{C}, \lambda) := \Big\langle \underbrace{\neg\exists^{\mathcal{C}}_{\lambda} x\, \varphi(x)}_{\text{(Not-Being) Nāsti / Abhāva / Asat}} , \underbrace{\mathcal{J}^{-}_{A}(\varphi)}_{\text{Aristotelian refutation}} , \underbrace{\mathcal{J}^{-}_{N}(\varphi; \lambda)}_{\text{Nyāya absence-justification}} \Big\rangle,$$

subject to the following constitutive conditions:

1. **Ontological denial (Not-Being).** The content asserts *absence relative to* $\mathcal{C}$ at λ: no admissible instance of φ is present there (*nāsti*); conceptually, this is *abhāva/asat*. *(Catuṣkoṭi: second corner).*
2. **Scope and determinacy.** The negative claim is well-scoped: (i) φ is well-formed under $\mathcal{C}$, (ii) the locus λ (and, when relevant, time window) is specified, and (iii) reference conditions for non-occurrence are fixed so that the denial is meaningful and testable within $\mathcal{C}$.
3. **Dialectical posture (Aristotle).** $\mathcal{J}_A^{\neg}(\varphi)$ provides reasons adequate for ordered debate: e.g., a direct negative demonstration ($\forall_\lambda^{\mathcal{C}} x\, \neg\varphi(x)$), an *elenctic* refutation of all admissible supports for φ, or a *reductio* showing φ entails impossibility within $\mathcal{C}$.
4. **Procedural obligation (Ancient India).** $\mathcal{J}_N^{\neg}(\varphi;\lambda)$ validates absence using Nyāya norms:
   - a five-membered argument with negative *pratijñā* (e.g., "the pot is not in the house"), accompanied by *hetu*, *udāharaṇa*, *upanaya*, and *nigamana*;
   - an account of *abhāva*-type as appropriate to λ: prior absence (*prāgabhāva*), posterior absence (*pradhva sābhāva*), mutual exclusion (*anyonyābhāva*), or absolute absence (*atyantābhāva*);
   - and, where admissible in $\mathcal{C}$, appeal to properly conditioned non-perception (*anupalabdhi*) or to inference from failure under satisfied enabling conditions.

Absolute vs. local denial. If the intent is *absolute* denial, then quantification ranges over all admissible loci:

$$\mathbf{AntiTh}_{\mathrm{abs}}(\varphi;\mathcal{C}) \text{ holds } \iff \forall\lambda \in \Lambda_{\mathcal{C}}\ \neg\exists_\lambda^{\mathcal{C}} x\, \varphi(x).$$

Equivalently. Writing $\exists_\lambda^{\mathcal{C}}$ for context- and locus-relative existence, and letting $\mathrm{Arg}_{A/N}^{\neg}$ denote well-formed Aristotelian / Nyāya *negative* arguments, we may say:

$$\mathbf{AntiTh}(\varphi;\mathcal{C},\lambda) \text{ holds } \iff \left(\neg\exists_\lambda^{\mathcal{C}} x\, \varphi(x)\right) \wedge \left(\mathcal{J}_A^{\neg}(\varphi) \in \mathrm{Arg}_A^{\neg}\right) \wedge \left(\mathcal{J}_N^{\neg}(\varphi;\lambda) \in \mathrm{Arg}_N^{\neg}\right).$$

Remarks. (i) *Antithesis* unifies *Nāsti/Abhāva/Asat* (ontic denial), Greek *antíthesis* (a reasoned counter-position), and the Nyāya–Navya-Nyāya analysis of absence. (ii) It is explicitly *immanent* and context-relative: it withholds existence only within $\mathcal{C}$ (and, if specified, λ), without asserting ultimate *asat*. Its *warrant*—not metaphysical truth simpliciter—turns on the adequacy of its refutation- and absence-justification.

Antithesis, in this scheme (Section 1.2.2 and Definition 1.2), is not a mere negation of Thesis but the system's *preference for distinction* along the same privileged axis. Formally it is the projector $P_A := |e_2\rangle\langle e_2|$ onto the Antisymmetry mode $|e_2\rangle$ selected by the ontological duality basis (Definition 2.1), with Born valuation $v_{\mathrm{Antithesis}}(\rho) = \mathrm{Tr}(P_A\rho)$ (Definition 1.18). Read intuitively, this number quantifies the system's *disposition to register as Antithesis* under the Thesis–Antithesis interrogation: a graded alignment, not an all-or-nothing verdict. In the bridging map, this is precisely the AntiSym $\leftrightarrow$ AntiTh corner correspondence, ensuring that the logical meaning and the quantum anchor co-move (Section 1.2.7).

Operationally, Antithesis is a *repeatable test* and a *filter*. Measuring in the $\{|e_1\rangle, |e_2\rangle\}$ basis, the long-run frequency of the $|e_2\rangle$ outcome converges to $v_{\text{Antithesis}}(\rho)$ by the Born rule (Definition 1.18); post-selection with P_A prepares the Antithesis mode for subsequent dynamics. Unitaries that commute with P_A conserve its valuation, whereas rotations in the $(0,1)$-sector redistribute weight between P_T and P_A. The local involutive duality operator $D^{(\mathfrak{B}_x)}$ makes the contrast exact: it swaps $|e_1\rangle \leftrightarrow |e_2\rangle$, thereby exchanging the roles of Thesis and Antithesis while acting trivially outside this sector (Section 2.1.2). In short, Antithesis is experimentally legible: a concrete basis, a concrete effect P_A, and a concrete number that plays the role of truth.

Finally, Antithesis *coexists* with Thesis without paradox once truth is read as *Born valuation*. Generic states in $\mathcal{H}_{(0,1)} = \text{span}\{|e_1\rangle, |e_2\rangle\}$ can carry nonzero weight for both P_T and P_A; only the classical limits $v_{\text{Antithesis}} \to 1$ or 0 recover bivalence. States orthogonal to $\mathcal{H}_{(0,1)}$ realize *"Neither,"* signaling the inapplicability of the Thesis/Antithesis query to that preparation. On the same plane, the parasymmetric and anti-parasymmetric modes provide the *complementary probes*: synthesis (Σ_x,–like) organizes the two tendencies, while anti-parasymmetry marks their dissonance; together they form a complete resolution of the $(0,1)$-sector (Sections 2.2.2 and 2.2.3).

1.2.3 The Definition of Synthesis

A *Synthesis* is a determinate, context-indexed co-commitment that simultaneously affirms and denies existence (Asti ca Nāsti ca; *Sadasat*) and is normatively bound to paired modes of defense—Aristotelian in the sense that both the positive and negative positions are furnished with reasoned arguments fit for dialectical scrutiny, and Nyāya in the sense that each side is validated by methodical, multi-membered arguments (with twin *pratijñās*)—such that its *epistemic warrant* derives from the coherence and validity of these opposed yet non-trivially co-tenable justifications.

Definition 1.3 (Synthesis)**. Synthesis** (*Asti ca Nāsti ca*/अस्ति च नास्ति च; *Sadasat*/सदसत्).

Fix a domain of discourse $\mathcal{D}$ and an immanent context $\mathcal{C}$ (criteria of individuation, admissible inference, and admissible loci of evaluation). Let $\Lambda_{\mathcal{C}}$ be the $\mathcal{C}$-sanctioned loci and let $\lambda \in \Lambda_{\mathcal{C}}$. The definition employs a *paraconsistent* consequence relation, denoted $\vdash_{\text{pc}}$, which is admissible in $\mathcal{C}$ (i.e., one that does not validate explosion).

A *Synthesis* regarding a formula φ at locus λ is a structured act

$$\mathbf{SynTh}(\varphi; \mathcal{C}, \lambda) := \Big\langle \underbrace{\exists^{\mathcal{C}}_{\lambda} x\, \varphi(x)}_{\text{(Being) Asti/Sat}}, \underbrace{\neg\exists^{\mathcal{C}}_{\lambda} x\, \varphi(x)}_{\text{(Not Being) Nāsti/Asat}}, \underbrace{\mathcal{J}^{+}_{\text{A}}(\varphi), \mathcal{J}^{-}_{\text{A}}(\varphi)}_{\text{Aristotelian paired defenses}}, \underbrace{\mathcal{J}^{+}_{\text{N}}(\varphi;\lambda), \mathcal{J}^{-}_{\text{N}}(\varphi;\lambda)}_{\text{Nyāya paired defenses}}, \underbrace{\Pi_{\text{pc}}}_{\text{nontriviality witness}} \Big\rangle,$$

subject to the following constitutive conditions:

1. **Co-affirmation (Both Being and Not-Being).** Both claims hold *at the same locus* λ under $\mathcal{C}$: $\exists^{\mathcal{C}}_{\lambda} x\, \varphi(x)$ and $\neg\exists^{\mathcal{C}}_{\lambda} x\, \varphi(x)$ are jointly endorsed. *(Catuṣkoṭi: third corner).*

2. **Scope hygiene (no equivocation).** The denotation of φ and the parameters of λ are fixed so that the "both" is not due merely to a shift of sense or locus; any aspectual refinements (e.g., mode, level, or respect) are explicit in $\mathcal{C}$.
3. **Paraconsistent discipline (nontriviality).** Π_{pc} is evidence that $\mathcal{C}$ employs a non-explosive consequence relation $\vdash_{\text{pc}}$ with a model or valuation v such that $v(\varphi) = \mathbf{B}$ (both true and false) and, for all ψ, it is not the case that $\{\varphi, \neg\varphi\} \vdash_{\text{pc}} \psi$.
4. **Dialectical posture (Aristotle, paired).** $\mathcal{J}^{+}_{\text{A}}(\varphi)$ provides reasons for the positive claim (e.g., syllogistic support from suitable *endoxa*); $\mathcal{J}^{-}_{\text{A}}(\varphi)$ provides reasons for the negative claim (e.g., direct negative demonstration, *elenchos*, or *reductio*). The pair is accompanied by a *co-tenability* argument showing that $\vdash_{\text{pc}}$ admits their joint maintenance without collapse.
5. **Procedural obligation (Ancient India, paired).** $\mathcal{J}^{+}_{\text{N}}(\varphi;\lambda)$ and $\mathcal{J}^{-}_{\text{N}}(\varphi;\lambda)$ are Nyāya five-membered arguments with positive and negative *pratijñā*, respectively, with appropriate use (where admissible) of perception, inference, testimony, and non-perception (*anupalabdhi*) to underwrite presence and absence.

Equivalently. Writing $\text{Arg}^{\pm}_{A/N}$ for well-formed positive/negative Aristotelian or Nyāya arguments under $\vdash_{\text{pc}}$, we may say:

$$\begin{aligned}\textbf{SynTh}(\varphi;\mathcal{C},\lambda) \text{ holds} \iff & \Big(\exists^{\mathcal{C}}_{\lambda} x\, \varphi(x) \ \wedge\ \neg\exists^{\mathcal{C}}_{\lambda} x\, \varphi(x)\Big) \\ & \wedge\ \Big(\mathcal{J}^{+}_{\text{A}}(\varphi) \in \text{Arg}^{+}_{A} \wedge \mathcal{J}^{-}_{\text{A}}(\varphi) \in \text{Arg}^{-}_{A}\Big) \\ & \wedge\ \Big(\mathcal{J}^{+}_{\text{N}}(\varphi;\lambda) \in \text{Arg}^{+}_{N} \wedge \mathcal{J}^{-}_{\text{N}}(\varphi;\lambda) \in \text{Arg}^{-}_{N}\Big) \\ & \wedge\ \Pi_{\text{pc}}.\end{aligned}$$

Remarks. (i) *Synthesis* captures *Asti ca Nāsti ca* and the dense compound *Sadasat*: a true contradiction (a "glut") licensed within an immanent, paraconsistent fragment of $\mathcal{C}$. (ii) It remains *context-relative* and non-metaphysical in ambit: no claim is made about ultimate *svabhāva*; the status of φ hinges on admissible evidence and $\vdash_{\text{pc}}$'s non-explosive logic. (iii) For purely aspectual coexistence (different respects or modes) the construction reduces to a consistent pair indexed by aspects within $\mathcal{C}$; the present definition includes, but is not limited to, that weaker case.

Here, *Synthesis* names a mode of coherent organization, not a timid "halfway point." In the $(0,1)$-sector it is the equal-weight *normalized* superposition

$$\tfrac{1}{\sqrt{2}}\big(|\mathring{\mathbf{0}}\rangle + |\mathring{\mathbf{1}}\rangle\big),$$

and the associated projector P_{para} isolates exactly that component of any state—i.e., the *direction of dialectical co-alignment* of Symmetry with Antisymmetry (Section 2.2 and Definition 2.17). In the Born-valued semantics, its truth valuation is $v_{\text{Synthesis}}(\rho) = \text{Tr}(P_{\text{para}}\rho)$, so "how true the Synthesis is" becomes a measurable propensity rather than a bivalent verdict (Definition 1.18). Read this way, Synthesis is the system's *disposition to register* in the coherent $\tfrac{1}{\sqrt{2}}\big(|\mathring{\mathbf{0}}\rangle + |\mathring{\mathbf{1}}\rangle\big)$ mode when interrogated along the parasymmetric axis—precisely the Thesis–Antithesis plane's complementary probe (Definition 1.16).

Operationally, one treats Synthesis as a *repeatable test and filter*: the pair of projectors $\{P_{\text{para}}, P_{\text{antipara}}\}$ forms an orthogonal, complete resolution of the $(0,1)$-sector, so repeated trials directly estimate $v_{\text{Synthesis}}(\rho)$ while post-selection prepares the synthesized mode for subsequent dynamics. In these terms, Anti-parasymmetry supplies the *contrast class*: it projects onto the normalized $\frac{1}{\sqrt{2}}\big(|\mathring{\mathbf{0}}\rangle - |\mathring{\mathbf{1}}\rangle\big)$, capturing modal dissonance; together, the two effects sum to the sector projector $P_{(0,1)}$, guaranteeing closure and basis-coherence with Thesis/Antithesis (Definition 2.18 and Section 2.2.2).

Beyond a single locus, Synthesis *becomes relation*. Bipartitely it is the Bell-like entangled mode $\frac{1}{\sqrt{2}}\big(|\mathring{\mathbf{0}}\rangle_x|\mathring{\mathbf{1}}\rangle_{x'} + |\mathring{\mathbf{1}}\rangle_x|\mathring{\mathbf{0}}\rangle_{x'}\big)$ with projector $P^{(xx')}_{\text{para}}$, a distributed co-alignment that belongs to neither site individually (Definition 2.23 and Section 2.2.3). Multipartitely it generalizes to the W-like state $|\Psi^{(n)}_{\text{para}}\rangle$ with global projector $P^{(x_1\cdots x_n)}_{\text{para}}$, exhibiting permutation symmetry, genuine multipartite entanglement, and mixed marginals—i.e., a *collective* synthesis that is irreducibly nonlocal (Section 2.2.5, Definition 2.33, and eq. (2.29)).

1.2.4 The Definition of the Holothesis

A *Holothesis* is a determinate, context-indexed suspension that refuses both affirmation and denial of existence—*Naiva Asti na Nāsti* (नैवास्ति न नास्ति)—not as a mere "none of the above," but as a categorical shift in which the very *applicability* of the existential polarity (Being/Not-Being, and their co-tenability) is withdrawn. Its warrant is *metadialectical*: any such polarity-talk misfires at this level (often glossed in Indian traditions as *anirvacanīya* or *aprapañca*).

Definition 1.4 (Holothesis)**. Holothesis** (*Naiva Asti na Nāsti*/नैवास्ति न नास्ति).

Fix a domain of discourse $\mathcal{D}$ and an immanent context $\mathcal{C}$ (criteria of individuation, admissible inference, admissible loci). Let $\Lambda_{\mathcal{C}}$ be the $\mathcal{C}$-sanctioned loci and $\lambda \in \Lambda_{\mathcal{C}}$. Let $\mathsf{App}_{\mathcal{C}}(\chi)$ read "χ is semantically applicable in $\mathcal{C}$," with $\neg\mathsf{App}_{\mathcal{C}}(\chi)$ meaning "χ is inapplicable (ill-typed or presuppositionally failed) in $\mathcal{C}$." Finally, let $\Uparrow\colon \mathrm{Lang}(\mathcal{C}) \to \mathrm{Lang}(\mathcal{C}^{\Uparrow})$ be an admissible *lift* functor witnessing the categorical shift.

A Holothesis about φ at λ is the structured act

$$\mathrm{HoloThAct}_{\mathcal{C}}(\varphi;\lambda) := \Big\langle \underbrace{\neg\mathsf{App}_{\mathcal{C}}\big(\exists^{\mathcal{C}}_{\lambda} x\,\varphi(x)\big)}_{\text{existential inapplicability}}, \underbrace{\neg\mathsf{App}_{\mathcal{C}}\big(\neg\,\exists^{\mathcal{C}}_{\lambda} x\,\varphi(x)\big)}_{\text{negative inapplicability}}, \underbrace{\mathcal{J}^{\perp}_{\mathrm{A}}(\varphi)}_{\text{Aristotelian meta-justification}}, \underbrace{\mathcal{J}^{\perp}_{\mathrm{N}}(\varphi;\lambda)}_{\text{Nyāya meta-justification}}, \underbrace{\Uparrow}_{\text{shift witness}} \Big\rangle,$$

subject to the following constitutive conditions:

1. **Polar inapplicability (Neither Being nor Not-Being).** Both existential polarities fail at the same locus λ in $\mathcal{C}$:

$$\neg\mathsf{App}_{\mathcal{C}}\big(\exists^{\mathcal{C}}_{\lambda} x\,\varphi(x)\big) \;\wedge\; \neg\mathsf{App}_{\mathcal{C}}\big(\neg\,\exists^{\mathcal{C}}_{\lambda} x\,\varphi(x)\big).$$

   This is not a covert assertion of *neither* (a gap) nor of *both* (a glut), but a typed judgement of *inapplicability* of the polarity itself.

2. **Stability under admissible reparametrizations.** For any refinement $\mathcal{C}'$ reachable from $\mathcal{C}$ by admissible re-indexing of sense, respect, mode, or locus (including aspectual factorizations), the inapplicability persists:

$$\forall\, \mathcal{C}' \in \mathrm{Ref}(\mathcal{C}): \quad \neg \mathsf{App}_{\mathcal{C}'}\Big(\exists^{\mathcal{C}'}_{\lambda} x\, \varphi(x)\Big) \;\wedge\; \neg \mathsf{App}_{\mathcal{C}'}\Big(\neg\, \exists^{\mathcal{C}'}_{\lambda} x\, \varphi(x)\Big).$$

3. **Non-collapse to the earlier corners.** (i) Not to *Thesis*: $\nvdash\, \exists^{\mathcal{C}}_{\lambda} x\, \varphi(x)$; (ii) not to *Antithesis*: $\nvdash\, \forall^{\mathcal{C}}_{\lambda} x\, \neg\varphi(x)$; (iii) not to *Synthesis*: no admissible consequence relation in $\mathcal{C}$ validates $\exists^{\mathcal{C}}_{\lambda} x\, \varphi(x) \;\wedge\; \neg \exists^{\mathcal{C}}_{\lambda} x\, \varphi(x)$.
4. **Dialectical posture (Aristotle, meta).** $\mathcal{J}^{\perp}_{\mathrm{A}}(\varphi)$ shows that attempts to force existential polarity here are *secundum quid/simpliciter* confusions, category mistakes (homonymy of *to be*), or require illicit shifts of predication base; hence "Is it or is it not?" is ill-posed *in* $\mathcal{C}$ for φ at λ.
5. **Procedural obligation (Nyāya, meta).** $\mathcal{J}^{\perp}_{\mathrm{N}}(\varphi; \lambda)$ provides a five-membered argument whose *pratijñā* is a *non-applicability claim*, with *hetu* (failed delimitor/avacchedaka or presupposition), *udāharaṇa* (a paradigm of category error), *upanaya* (mapping to the present case), and *nigamana* (concluding inapplicability); where relevant, it may also appeal to disciplined *tarka* to show that forcing the polarity leads to contradiction or vacuity.
6. **Shift witness (categorical lift).** The functor $\Uparrow$ maps φ to a stratum $\mathcal{C}^{\Uparrow}$ in which existential vocabulary is eliminated or reinterpreted so that $\Uparrow(\varphi)$ is evaluable without the polarity; composing back to $\mathcal{C}$ does not reintroduce it.

Equivalently. Let $\mathfrak{L}_{\mathcal{C}}$ be the family of admissible consequence relations in $\mathcal{C}$ (classical, paraconsistent, paracomplete, etc.). Then a Holothesis holds iff

$$\neg \mathsf{App}_{\mathcal{C}}\Big(\exists^{\mathcal{C}}_{\lambda} x\, \varphi(x)\Big) \;\wedge\; \neg \mathsf{App}_{\mathcal{C}}\Big(\neg\, \exists^{\mathcal{C}}_{\lambda} x\, \varphi(x)\Big) \;\wedge\; \mathcal{J}^{\perp}_{\mathrm{A}}(\varphi) \in \mathrm{Arg}^{\perp}_{\mathrm{A}} \;\wedge\; \mathcal{J}^{\perp}_{\mathrm{N}}(\varphi; \lambda) \in \mathrm{Arg}^{\perp}_{\mathrm{N}} \wedge\; \Uparrow,$$

and, moreover, for every $\vdash \in \mathfrak{L}_{\mathcal{C}}$, neither the affirmative, the negative, nor their conjunction is derivable for φ at λ.

Remarks. (i) "Neither Being nor Not-Being" here is *not* the assertion of a third truth value in $\mathcal{C}$ but a typed *judgement of inapplicability* of the existential polarity at that site, stabilized by the categorical lift $\Uparrow$. (ii) This preserves the immanent/ultimate distinction: it makes no claim of ultimate *svabhāva*, only that, at this stratum, existence/non-existence talk is idle; the work is done by metadialectical justification rather than by hidden ontology. (iii) In contexts where an admissible aspectual re-indexation makes the polarity meaningful, the Holothesis *dissolves* (returns to one of the first three corners) via failure of Stability, as it should. (iv) The Holothesis is a *meta-justification* of the inapplicability of the existential polarity at λ in $\mathcal{C}$.

Here, the *Holothesis* names a meta-assertion: not "the system is in such-and-such mode," but "the system's *law of evolution* is being realized in a coherent superposition of alternatives—and we can test that." Concretely, holosymmetry lifts two complementary parasymmetric dynamics to a dual-register stage and lets a control qubit coherently *assign* which

register gets which law (U_G vs. U_N); preparing the control in a superposition entangles entire *evolutionary histories* rather than mere states (Definitions 2.102 and 2.108). In the tetrahedral picture this is the apex speaking about the whole base at once: a statement *about* Symmetry/Antisymmetry/Parasymmetry, not another point on their plane (Sections 2.3 and 3.8).

Operationally, the Holothesis is implemented by the controlled holosymmetric operator $\mathbf{U}_{\text{ctrl}}$ that branches the dual registers under the two lifted path-laws, followed by an interference/recombination and a measurement on the control. This architecture makes the control readout a visibility meter for "agreement" or "dissonance" between full histories (Definition 2.114 and Section 2.3.2); in the symmetric interferometer one can read off constructive vs. destructive interference directly from the control probabilities p_0, p_1, which depend on the overlap of the two global evolutions acting on the same input state (Sections 2.3.2 and 3.1.2). The preparation/readout gates $\mathbf{U}_{\text{prep}}, \mathbf{U}_{\text{read}}$ let us choose the control basis in which this global question is asked (Section 3.1).

Truth semantics follows the same rule as the other corners, but the *effect* is induced by the whole holosymmetric circuit: with control prepared in $|\mathbf{a}\rangle_c$ and read out in $|\mathbf{b}\rangle_c$, the Holothesis valuation on a register state ρ_{RR} is

$$v_{\text{Holothesis}}(\rho_{\text{RR}}) = \text{Tr}\Big(E^{(\mathbf{a}\to\mathbf{b})}_{\text{Holothesis}}\,\rho_{\text{RR}}\Big), \qquad E^{(\mathbf{a}\to\mathbf{b})}_{\text{Holothesis}} = \Big\langle \mathbf{a}\Big|_c\, \mathbf{U}^{\dagger}_{\text{holo}}\,(|\mathbf{b}\rangle\langle\mathbf{b}|_c \otimes \mathbb{I})\,\mathbf{U}_{\text{holo}}\,\Big|\mathbf{a}\Big\rangle_c,$$

as recorded in the Born-valued semantics (Definition 1.18). This keeps the Holothesis a *repeatable test*: choose the control basis $(\mathbf{U}_{\text{prep}}, \mathbf{U}_{\text{read}})$, run the interferometer, and estimate this probability from frequencies—now a number that quantifies "how true the Holothesis is" for that preparation. Note that what is superposed are *assignments of global dynamics*, not operation orders; the construction stays within standard controlled-unitary physics (Definition 2.114).

1.2.5 Tetralemma of Algebra

Definition 1.5 (Tetralemma of Algebra (Atomic Prismatic))**.** Let Th, AntiTh, SynTh, and HoloTh denote, respectively, the modalities *Thesis*, *Antithesis*, *Synthesis*, and *Holothesis* as already specified in Sections 1.2.1 to 1.2.4. The *Tetralemma of Algebra* (the *atomic prismatic*) is the ordered quadruple

$$\mathring{\mathbf{\Lambda}} := \Big\langle \mathsf{Th},\ \mathsf{AntiTh},\ \mathsf{SynTh},\ \mathsf{HoloTh} \Big\rangle$$

together with *exactly and only* the relations, constructions, and identifications that are stipulated within those four definitions, taken as a single coherent unit. Formally:

(a) (**No redefinition**) The internal content of each corner is *identical* to that given in its own definition; no additional axioms are imposed on $\mathsf{Th}, \mathsf{AntiTh}, \mathsf{SynTh}, \mathsf{HoloTh}$ beyond those already declared in Sections 1.2.1 to 1.2.4.

(b) (**Closure as a unit**) The atomic prismatic $\mathring{\mathbf{\Lambda}}$ is the *minimal* algebraic object containing the four corners and closed under the constructions already invoked in those definitions

(e.g. the oppositional/duality relation between Th and AntiTh insofar as it is specified there, the construction of SynTh from Th and AntiTh as specified there, and the meta-level role attributed to HoloTh as specified there).

(c) (**Atomicity**) $\overset{\circ}{\mathbf{\Lambda}}$ admits no nontrivial decomposition into a disjoint union of sub-*quadruples* preserving the relations inherited from the four definitions. Equivalently, there is no partition $\{\mathsf{Th}, \mathsf{AntiTh}, \mathsf{SynTh}, \mathsf{HoloTh}\} = U \uplus V$ with $U, V \neq \varnothing$ such that the inherited relations restrict to make both U and V into atomic prismatic units. Hence, in particular, there is no separation of $\{\mathsf{Th}, \mathsf{AntiTh}\}$ from $\{\mathsf{SynTh}, \mathsf{HoloTh}\}$ (or any other split) compatible with those relations.

(d) (**Morphisms, optional**) A *prismatic morphism* $f : \overset{\circ}{\mathbf{\Lambda}} \to \overset{\circ}{\mathbf{\Lambda}}{}'$ is a map that sends each corner to its namesake,

$$f(\mathsf{Th}) = \mathsf{Th}', \quad f(\mathsf{AntiTh}) = \mathsf{AntiTh}', \quad f(\mathsf{SynTh}) = \mathsf{SynTh}', \quad f(\mathsf{HoloTh}) = \mathsf{HoloTh}',$$

and preserves *all* relations, constructions, and identifications that appear in Sections 1.2.1 to 1.2.4. Two units are *isomorphic* if there exists a bijective prismatic morphism between them.

We will refer to $\overset{\circ}{\mathbf{\Lambda}}$ as the *Tetralemma of Algebra*.

Remark 1.6 (Terminology Convention and Prismatic Intent)**.** Throughout, "atomic" means *indecomposable under corner-respecting decompositions* as in Definition 1.5(c) (and not "having atoms" in the order-theoretic sense). "Prismatic" means that admissible maps act on the *entire* unit $\overset{\circ}{\mathbf{\Lambda}}$, transporting the Fourfold *as a whole*; a map that preserves the corners only piecemeal while violating the inherited inter-corner constructions is not prismatic.

Remark 1.7 (Universality/Minimality of the Tetralemma of Algebra)**.** Let $\mathcal{C}$ be the class of all structures Q whose *underlying carrier set* $|Q|$ satisfies

$$\{\mathsf{Th}, \mathsf{AntiTh}, \mathsf{SynTh}, \mathsf{HoloTh}\} \subseteq |Q|,$$

and whose operations/relations are *exactly* those induced by Sections 1.2.1 to 1.2.4 (restricted to $|Q|$). Then the atomic prismatic $\overset{\circ}{\mathbf{\Lambda}}$ of Definition 1.5 is obtained by intersecting carrier sets and restricting relations:

$$|\overset{\circ}{\mathbf{\Lambda}}| \;=\; \bigcap_{Q \in \mathcal{C}} |Q|, \qquad \text{with the inherited operations/relations restricted to } \; |\overset{\circ}{\mathbf{\Lambda}}|.$$

In particular, $\overset{\circ}{\mathbf{\Lambda}}$ is contained in every $Q \in \mathcal{C}$ and is unique with this property. This packages the "closure as a unit" and "no redefinition" clauses as a universal (initial) characterization, without adding structure to the corners.

Proposition 1.8 (Invariance Under Admissible Dualities and Reindexings)**.** *Let $\mathcal{D}$ denote the duality involution of Section 1.2.6 exchanging $\mathsf{Th} \leftrightarrow \mathsf{AntiTh}$ and fixing $\mathsf{SynTh}, \mathsf{HoloTh}$, and let ρ be any admissible reindexing that preserves the four corner-definitions. Then*

$$\mathcal{D}(\overset{\circ}{\mathbf{\Lambda}}) = \overset{\circ}{\mathbf{\Lambda}} \qquad \textit{and} \qquad \rho(\overset{\circ}{\mathbf{\Lambda}}) = \overset{\circ}{\mathbf{\Lambda}}.$$

Proof. By Proposition 1.12, $\{\mathsf{Th}, \mathsf{AntiTh}, \mathsf{SynTh}, \mathsf{HoloTh}\}$ is preserved (indeed, pointwise fixed up to the $\mathsf{Th} \leftrightarrow \mathsf{AntiTh}$ swap) under $\mathcal{D}$ and any admissible ρ. By Definition 1.5, $\overset{\circ}{\mathbf{\Lambda}}$ is the minimal object that (i) contains these corners and (ii) is closed under exactly the inherited constructions used in their definitions. Since $\mathcal{D}$ and ρ preserve both the corners and those constructions, they send any $Q \in \mathcal{C}$ to another element of $\mathcal{C}$, hence fix the intersection $\bigcap_{Q\in\mathcal{C}} |Q|$ and the restricted relations; minimality forces equality. □

Corollary 1.9 (Initiality/Uniqueness of Prismatic Morphisms out of the Atomic Prismatic)**.** *For every $Q \in \mathcal{C}$ there exists a unique* prismatic morphism $f : \overset{\circ}{\mathbf{\Lambda}} \to Q$.

Proof. Existence: the inclusion of carriers $|\overset{\circ}{\mathbf{\Lambda}}| \subseteq |Q|$ respects the inherited relations by construction, giving a prismatic morphism. Uniqueness: any prismatic morphism must agree with this inclusion on the four named corners and is determined on the closure generated by them; minimality of $\overset{\circ}{\mathbf{\Lambda}}$ then forces equality. □

1.2.6 The Axiom of Duality and its Self-Similar Nature

Axiom 1.10 (Axiom of Duality)**.** Fix an immanent context $\mathcal{C}$ with its criteria of individuation and admissible inference. Let $\mathcal{U}_\mathcal{C}$ denote the class of primitive ontological states. There exists an involution (duality operator)

$$\mathbf{D}_\mathcal{C} : \mathcal{U}_\mathcal{C} \to \mathcal{U}_\mathcal{C}, \qquad \mathbf{D}_\mathcal{C}^2 = \mathrm{Id},$$

which, for the designated primitives, satisfies $\mathbf{D}_\mathcal{C}\big(|\overset{\circ}{\mathbf{0}}\rangle\big) = |\overset{\circ}{\mathbf{1}}\rangle$ and $\mathbf{D}_\mathcal{C}\big(|\overset{\circ}{\mathbf{1}}\rangle\big) = |\overset{\circ}{\mathbf{0}}\rangle$. In a reality governed by dependent origination (*pratītyasamutpāda*), the assertion of any state $s \in \mathcal{U}_\mathcal{C}$ entails the conceptual co–arising of its complement:

$$\mathsf{Affirm}_\mathcal{C}(s) \implies \mathsf{Conceive}_\mathcal{C}\big(\mathbf{D}_\mathcal{C}(s)\big),$$

where $\mathsf{Affirm}_\mathcal{C}(s)$ abbreviates "s is affirmed in $\mathcal{C}$," and $\mathsf{Conceive}_\mathcal{C}(t)$ abbreviates "t is (minimally) conceptually available in $\mathcal{C}$." Accordingly, intelligible ontological content is minimally *dyadic*:

$$\langle s, \mathbf{D}_\mathcal{C}(s)\rangle \text{ is the minimal semantic unit of duality,}$$

i.e., neither element is semantically autonomous in isolation.

Operational Ties to the Four Corners

Fix a locus λ in context $\mathcal{C}$. Let

$$\mathsf{Status}_\mathcal{C}(\varphi, \lambda) \in \{\mathsf{Th}, \mathsf{AntiTh}, \mathsf{SynTh}, \mathsf{HoloTh}\}$$

denote the corner–status (Thesis, Antithesis, Synthesis, Holothesis) of a well-formed formula φ at λ (as defined earlier). Define the *status dualization*

$$\Delta_\mathcal{C} : \{\mathsf{Th}, \mathsf{AntiTh}, \mathsf{SynTh}, \mathsf{HoloTh}\} \to \{\mathsf{Th}, \mathsf{AntiTh}, \mathsf{SynTh}, \mathsf{HoloTh}\}$$

by

$$\begin{aligned} \Delta_\mathcal{C}(\mathsf{Th}) &= \mathsf{AntiTh}, & \Delta_\mathcal{C}(\mathsf{SynTh}) &= \mathsf{SynTh}, \\ \Delta_\mathcal{C}(\mathsf{AntiTh}) &= \mathsf{Th}, & \Delta_\mathcal{C}(\mathsf{HoloTh}) &= \mathsf{HoloTh}. \end{aligned}$$

so that $\Delta_\mathcal{C}^2 = \mathrm{Id}$.

Compatibility with Object-level Duality.

For every well-formed φ at λ,

$$\mathsf{Status}_{\mathcal{C}}\Big(\mathbf{D}_{\mathcal{C}}(\varphi),\lambda\Big) \;=\; \Delta_{\mathcal{C}}\Big(\mathsf{Status}_{\mathcal{C}}(\varphi,\lambda)\Big).$$

Interpretation. Dualization exchanges *Thesis* and *Antithesis*, while *Synthesis* and *Holothesis* are self-dual (the former licenses a disciplined both/and; the latter marks inapplicability of the polarity).

Self-similarity (Scale-free Replication)

Duality propagates uniformly across levels of discourse.

Definition 1.11 (Lifts of Duality)**.**

(i) **Object level.** $\mathbf{D}_{\mathcal{C}}$ acts on primitive ontological states $s \in \mathcal{U}_{\mathcal{C}}$ with $\mathbf{D}_{\mathcal{C}}^2 = \mathrm{Id}$.

(ii) **Propositional level.** Extend $\mathbf{D}_{\mathcal{C}}$ to formulas by dualizing primitive predicates and distributing through logical form (De Morgan–compatible) in a way that respects the admissible consequence relation of $\mathcal{C}$ (classical or paraconsistent, as specified). We write $\varphi^{\mathbf{D}}$ for the result of this homomorphic extension.

(iii) **Corner level.** The *status dualization* $\Delta_{\mathcal{C}}$ on $\{\mathsf{Th},\mathsf{AntiTh},\mathsf{SynTh},\mathsf{HoloTh}\}$ is given by

$$\begin{aligned} \Delta_{\mathcal{C}}(\mathsf{Th}) &= \mathsf{AntiTh}, & \Delta_{\mathcal{C}}(\mathsf{SynTh}) &= \mathsf{SynTh},\\ \Delta_{\mathcal{C}}(\mathsf{AntiTh}) &= \mathsf{Th}, & \Delta_{\mathcal{C}}(\mathsf{HoloTh}) &= \mathsf{HoloTh}. \end{aligned}$$

so that $\Delta_{\mathcal{C}}^2 = \mathrm{Id}$, and status assignment respects duality:

$$\mathsf{Status}_{\mathcal{C}}(\varphi,\lambda) = X \implies \mathsf{Status}_{\mathcal{C}}(\varphi^{\mathbf{D}},\lambda) = \Delta_{\mathcal{C}}(X).$$

(iv) **Meta level.** In the *Holothesis* case, introduce an applicability predicate $\mathsf{App}_{\mathcal{C}}(\cdot)$. Duality leaves *inapplicability* invariant:

$$\neg\mathsf{App}_{\mathcal{C}}(\chi) \implies \neg\mathsf{App}_{\mathcal{C}}\Big(\mathbf{D}_{\mathcal{C}}(\chi)\Big).$$

Object-level Instance

Let $|\mathring{\mathbf{0}}\rangle$ be any primitive ontological state (see Definition 1.16) and define $|\mathring{\mathbf{1}}\rangle \coloneqq \mathbf{D}_{\mathcal{C}}\Big(|\mathring{\mathbf{0}}\rangle\Big)$. Then

$$|\mathring{\mathbf{0}}\rangle \;\leftrightarrow\; |\mathring{\mathbf{1}}\rangle, \qquad \mathbf{D}_{\mathcal{C}}\Big(|\mathring{\mathbf{1}}\rangle\Big) = |\mathring{\mathbf{0}}\rangle,$$

and the dyad $\Big(|\mathring{\mathbf{0}}\rangle, |\mathring{\mathbf{1}}\rangle\Big)$ is semantically primary.

Self-similar Lifts (Equivalent Restatement).

(i) Propositional lift. Extend $\mathbf{D}_{\mathcal{C}}$ homomorphically to formulas; denote the dual by $\varphi^{\mathbf{D}}$.

(ii) Tetralemma-status lift. There is an involution $\Delta_{\mathcal{C}}$ with $\Delta_{\mathcal{C}}(\mathsf{Th}) = \mathsf{AntiTh}$, $\Delta_{\mathcal{C}}(\mathsf{AntiTh}) = \mathsf{Th}$, and $\Delta_{\mathcal{C}}(\mathsf{SynTh}) = \mathsf{SynTh}$, $\Delta_{\mathcal{C}}(\mathsf{HoloTh}) = \mathsf{HoloTh}$; status assignment respects duality: $\mathsf{Status}_{\mathcal{C}}(\varphi, \lambda) = X \Rightarrow \mathsf{Status}_{\mathcal{C}}(\varphi^{\mathbf{D}}, \lambda) = \Delta_{\mathcal{C}}(X)$.

(iii) Meta-lift (informal). Duality preserves admissible inference and typing rules (schematically, $\mathbf{D}_{\mathcal{C}} : \mathcal{R} \to \mathcal{R}$ with $\mathbf{D}_{\mathcal{C}}^2 = \mathrm{Id}$ on the rule set $\mathcal{R}$).

Proposition 1.12 (Self-similarity and closure)**.** *For any admissible reindexing $\mathcal{C} \to \mathcal{C}'$ (change of aspect, respect, mode, or locus) that preserves dependent origination, there exists a unique duality operator $\mathbf{D}_{\mathcal{C}'}$ such that*

$$F \circ \mathbf{D}_{\mathcal{C}} \;=\; \mathbf{D}_{\mathcal{C}'} \circ F$$

for the induced functor $F : \mathcal{U}_{\mathcal{C}} \to \mathcal{U}_{\mathcal{C}'}$. Hence duality is scale-free *and the set*

$$\{\mathsf{Th}, \mathsf{AntiTh}, \mathsf{SynTh}, \mathsf{HoloTh}\}$$

is closed and pointwise fixed under repeated dualization.

Sketch. Involution and functoriality determine $\mathbf{D}_{\mathcal{C}'}$ uniquely by conjugation. By definition of $\Delta_{\mathcal{C}}$, Thesis and Antithesis are exchanged; Synthesis and Holothesis quantify over—or deny the applicability of—the polarity itself and are therefore invariant under the swap. Closure follows. □

(iv) Meta-level lift — Minimal, proof-theoretic statement Let $\mathsf{Judg}_{\mathcal{C}}$ denote the set of well-formed judgments in context $\mathcal{C}$, $\mathsf{Deriv}_{\mathcal{C}}$ the derivations built from an admissible rule set $\mathcal{R}_{\mathcal{C}}$, and write $\Gamma \vdash_{\mathcal{C}} \varphi$ for a judgment. There exists an involution $\widehat{\mathbf{D}}_{\mathcal{C}}$ (the *meta-duality*) acting on rules, derivations, and judgments such that:

- **M1** (*Action on judgments*) $\widehat{\mathbf{D}}_{\mathcal{C}}(\Gamma \vdash_{\mathcal{C}} \varphi) \;=\; \widehat{\mathbf{D}}_{\mathcal{C}}(\Gamma) \;\vdash_{\mathcal{C}}\; \varphi^{\mathbf{D}}$, with $\widehat{\mathbf{D}}_{\mathcal{C}}^2 = \mathrm{Id}$.
- **M2** (*Action on rules*) For every inference rule $r \in \mathcal{R}_{\mathcal{C}}$, $\widehat{\mathbf{D}}_{\mathcal{C}}(r) \in \mathcal{R}_{\mathcal{C}}$ and r is admissible iff $\widehat{\mathbf{D}}_{\mathcal{C}}(r)$ is admissible.
- **M3** (*Action on derivations*) $\widehat{\mathbf{D}}_{\mathcal{C}}$ maps derivations to derivations by structural recursion, preserves derivability, and commutes with substitution/renaming: $\widehat{\mathbf{D}}_{\mathcal{C}}(\varphi[\sigma]) = \widehat{\mathbf{D}}_{\mathcal{C}}(\varphi)[\,\widehat{\mathbf{D}}_{\mathcal{C}}(\sigma)\,]$.
- **M4** (*Status coherence*) If $\mathsf{Status}_{\mathcal{C}}(\varphi, \lambda) = X$ (as in *(iii)*), then $\mathsf{Status}_{\mathcal{C}}(\varphi^{\mathbf{D}}, \lambda) = \Delta_{\mathcal{C}}(X)$.
- **M5** (*Conservativity & non-explosion*) $\widehat{\mathbf{D}}_{\mathcal{C}}$ preserves consistency and any stipulated non-explosive consequence relation $\vdash_{\mathrm{pc}}$ used in Synthesis:

$$\Gamma \nvdash_{\mathrm{pc}} \bot \;\Leftrightarrow\; \widehat{\mathbf{D}}_{\mathcal{C}}(\Gamma) \nvdash_{\mathrm{pc}} \bot.$$

(iv) Meta-level lift — Categorical/autoequivalence form (optional) Let **Ctx** be the category whose objects are immanent contexts $\mathcal{C}$ and whose morphisms are admissible refinements $f : \mathcal{C} \to \mathcal{C}'$. Let **Der** be the category of derivations (proofs) internal to contexts, with a forgetful functor $U : \mathbf{Der} \to \mathbf{Ctx}$ sending each derivation to its home context. There

exists a strict endofunctor $\mathfrak{D} : \mathbf{Ctx} \to \mathbf{Ctx}$ and a lifted endofunctor $\widehat{\mathfrak{D}} : \mathbf{Der} \to \mathbf{Der}$ such that

$$U \circ \widehat{\mathfrak{D}} = \mathfrak{D} \circ U, \qquad \mathfrak{D}^2 \cong \mathrm{Id}_{\mathbf{Ctx}}, \qquad \widehat{\mathfrak{D}}^2 \cong \mathrm{Id}_{\mathbf{Der}},$$

with natural isomorphisms witnessing involutivity. On objects/states and formulas, $\widehat{\mathfrak{D}}$ restricts to the object- and propositional-dualities of *(i)* and *(ii)*; on statuses it restricts to $\Delta_{\mathcal{C}}$ of *(iii)*. Moreover, $\widehat{\mathfrak{D}}$ preserves admissibility of rules, cut-elimination, and commutes with substitution and reindexing (functorial coherence squares commute).

Relation to Ket-Notation Dyads

Fix an immanent context $\mathcal{C}$ and its duality operator $\mathbf{D}_{\mathcal{C}}$ (Axiom 1.10).

Definition 1.13 (Dyadic constructor and orbit)**.** For any admissible state/content $s \in \mathcal{U}_{\mathcal{C}}$, define the *ordered dyad*

$$\mathfrak{D}_{\mathcal{C}}(s) \;:=\; \langle s, \mathbf{D}_{\mathcal{C}}(s) \rangle \;\in\; \mathcal{U}_{\mathcal{C}} \times \mathcal{U}_{\mathcal{C}}.$$

The associated *unordered dyad* is the $\mathbb{Z}_2$–orbit of s under the action generated by $\mathbf{D}_{\mathcal{C}}$:

$$[\![s]\!]_{\mathcal{C}} \;:=\; \{\, s, \mathbf{D}_{\mathcal{C}}(s) \,\}.$$

Principle 1.1 (Dyadic intelligibility)**.** *There is a predicate* $\mathrm{Int}_{\mathcal{C}}(\cdot)$ *("is a minimal intelligible semantic unit in* $\mathcal{C}$*") such that, for all admissible* $s \in \mathcal{U}_{\mathcal{C}}$*,*

$$\mathrm{Int}_{\mathcal{C}}\Big(\mathfrak{D}_{\mathcal{C}}(s)\Big) \quad \textit{holds}, \qquad \neg\,\mathrm{Int}_{\mathcal{C}}(s) \quad \textit{and} \quad \neg\,\mathrm{Int}_{\mathcal{C}}\Big(\mathbf{D}_{\mathcal{C}}(s)\Big) \textit{ hold.}$$

Equivalently, any unary semantic evaluation in $\mathcal{C}$ *factors through dyadization: there exist a (partial) map*

$$\mathcal{M}_{\mathcal{C}}^{(2)} :\; \mathcal{U}_{\mathcal{C}} \times \mathcal{U}_{\mathcal{C}} \rightharpoonup \mathbb{M} \quad \textit{and a total map} \quad F : \mathbb{M} \to \mathbb{N}$$

such that whenever the unary evaluation $\mathcal{M}_{\mathcal{C}}(s)$ *is defined,*

$$\mathcal{M}_{\mathcal{C}}(s) \;=\; F\Big(\mathcal{M}_{\mathcal{C}}^{(2)}\Big(\mathfrak{D}_{\mathcal{C}}(s)\Big)\Big)\,.$$

That is, intelligibility attaches to the dyad rather than to a singleton.

Example 1.14 (Kets and conceptual pairs)**.** If $\mathbf{D}_{\mathcal{C}}\Big(|\mathring{\mathbf{0}}\rangle\Big) = |\mathring{\mathbf{1}}\rangle$, then

$$\mathfrak{D}_{\mathcal{C}}\Big(|\mathring{\mathbf{0}}\rangle\Big) = \Big\langle\, |\mathring{\mathbf{0}}\rangle,\; |\mathring{\mathbf{1}}\rangle \,\Big\rangle.$$

At the conceptual level (lift (ii) of Definition 1.11),

$$\begin{aligned} \mathfrak{D}_{\mathcal{C}}(\text{Existence}) &= \langle \text{Existence}, \text{Non-Existence} \rangle, \\ \mathfrak{D}_{\mathcal{C}}(\text{Self}) &= \langle \text{Self}, \text{Not-Self} \rangle, \\ &\vdots \\ \mathfrak{D}_{\mathcal{C}}(\text{Real}) &= \langle \text{Real}, \text{Un-Real} \rangle. \end{aligned}$$

Higher-order dyads (e.g., $\langle \text{Duality}, \text{Non-Duality} \rangle$) follow from the propositional and meta lifts in Definition 1.11.

Corner-selections on Dyads

Fix an immanent context $\mathcal{C}$, a locus λ, and write $\mathfrak{D}_{\mathcal{C}}(\varphi) = \langle L(\varphi),\ R(\varphi)\rangle$ for a well-formed φ (with $R(\varphi) = \mathbf{D}_{\mathcal{C}}(L(\varphi))$). Define selection functionals

$$\mathsf{Sel}_{\mathsf{Th}}, \mathsf{Sel}_{\mathsf{AntiTh}}, \mathsf{Sel}_{\mathsf{SynTh}}, \mathsf{Sel}_{\mathsf{HoloTh}} : \Big(\mathcal{U}_{\mathcal{C}} \times \mathcal{U}_{\mathcal{C}}\Big) \times \Lambda_{\mathcal{C}} \longrightarrow \mathcal{P}\Big(\{L(\varphi), R(\varphi)\}\Big)$$

by

$$\mathsf{Sel}_{\mathsf{Th}}\Big(\mathfrak{D}_{\mathcal{C}}(\varphi); \lambda\Big) = \{L(\varphi)\}, \qquad \mathsf{Sel}_{\mathsf{AntiTh}}\Big(\mathfrak{D}_{\mathcal{C}}(\varphi); \lambda\Big) = \{R(\varphi)\},$$
$$\mathsf{Sel}_{\mathsf{SynTh}}\Big(\mathfrak{D}_{\mathcal{C}}(\varphi); \lambda\Big) = \{L(\varphi), R(\varphi)\}, \qquad \mathsf{Sel}_{\mathsf{HoloTh}}\Big(\mathfrak{D}_{\mathcal{C}}(\varphi); \lambda\Big) = \varnothing,$$

where the last case reflects the *inapplicability* of existential polarity in the Holothesis.

Proposition 1.15 (Compatibility with the four corners)**.** *At any admissible locus λ, the corner–statuses align with the selectors as follows:*

(Thesis) $\mathrm{Thesis}_{\mathcal{C}}(\varphi; \lambda) \Rightarrow \exists^{\mathcal{C}}_{\lambda} x\, \mathrm{L}(\varphi)(x)$.

(Antithesis) $\mathrm{Antithesis}_{\mathcal{C}}(\varphi; \lambda) \Rightarrow \exists^{\mathcal{C}}_{\lambda} x\, \mathrm{R}(\varphi)(x)$.

(No identification of R with $\neg\,\mathrm{L}$ is assumed.)

(Synthesis) $\mathrm{Synthesis}_{\mathcal{C}}(\varphi; \lambda) \Rightarrow \Big(\exists^{\mathcal{C}}_{\lambda} x\, \mathrm{L}(\varphi)(x)\Big) \wedge \Big(\exists^{\mathcal{C}}_{\lambda} y\, \mathrm{R}(\varphi)(y)\Big)$.

The joint commitment is stably maintainable under $\vdash_{\mathrm{pc}}$.

(Holothesis) $\mathrm{Holothesis}_{\mathcal{C}}(\varphi; \lambda) \Rightarrow \neg\,\mathrm{App}_{\mathcal{C}}\Big(\exists^{\mathcal{C}}_{\lambda} x\, \mathrm{L}(\varphi)(x)\Big) \wedge \neg\,\mathrm{App}_{\mathcal{C}}\Big(\exists^{\mathcal{C}}_{\lambda} y\, \mathrm{R}(\varphi)(y)\Big)$.

Thus the four corners are realized as, respectively, left *selection,* right *selection,* both *selections (paraconsistently licensed), and* no *selection due to meta-level inapplicability.*

Sketch. By Definition 1.13, $\mathfrak{D}_{\mathcal{C}}(\varphi)$ packages the dual pair $\langle L, R\rangle$. Thesis selects L, hence affirms an instance of $L(\varphi)$; Antithesis selects R, hence affirms an instance of $R(\varphi)$. Synthesis selects both, so both existential claims hold; stability under $\vdash_{\mathrm{pc}}$ ensures non-explosion even if L and R collide at a locus. In the Holothesis case, $\mathsf{Sel}_{\mathsf{HoloTh}} = \varnothing$ and Definition 1.11 (iv) gives invariance of inapplicability under dualization, hence the two $\neg\mathsf{App}_{\mathcal{C}}$ clauses. Closure and alignment with statuses follow. □

Paraconsistent Compatibility

When $\mathcal{C}$ admits a paraconsistent (see Section 1.2.3) consequence relation $\vdash_{\mathrm{pc}}$, *Synthesis* corresponds to asserting *both* coordinates of $\mathfrak{D}_{\mathcal{C}}(s)$ without explosion, while *Holothesis* corresponds to meta-level *inapplicability* for both coordinates, which is preserved by dualization:

$$\mathsf{SynTh}_{\mathcal{C}}(\varphi; \lambda) \Rightarrow \Big(\exists^{\mathcal{C}}_{\lambda} x\, L(\varphi)(x)\Big) \ \wedge\ \Big(\exists^{\mathcal{C}}_{\lambda} y\, R(\varphi)(y)\Big),$$
$$\text{with } \{\exists^{\mathcal{C}}_{\lambda} x\, L(\varphi)(x),\ \exists^{\mathcal{C}}_{\lambda} y\, R(\varphi)(y)\} \nvdash_{\mathrm{pc}} \bot;$$
$$\mathsf{HoloTh}_{\mathcal{C}}(\varphi; \lambda) \Rightarrow \neg\mathsf{App}_{\mathcal{C}}\Big(\exists^{\mathcal{C}}_{\lambda} x\, L(\varphi)(x)\Big) \ \wedge\ \neg\mathsf{App}_{\mathcal{C}}\Big(\exists^{\mathcal{C}}_{\lambda} y\, R(\varphi)(y)\Big).$$

and, since $\Delta_{\mathcal{C}}(\mathsf{SynTh}) = \mathsf{SynTh}$ and $\Delta_{\mathcal{C}}(\mathsf{HoloTh}) = \mathsf{HoloTh}$,

$$\mathsf{Status}_{\mathcal{C}}(\mathbf{D}_{\mathcal{C}}(\varphi), \lambda) = \mathsf{Status}_{\mathcal{C}}(\varphi, \lambda) \quad \text{for } \mathsf{Status} \in \{\mathsf{SynTh}, \mathsf{HoloTh}\}.$$

Notational Refinement

To visually mark *conceptual co-implication/co-arising*, write

$$s \;\dashv\vdash_{\mathcal{C}}\; \mathbf{D}_{\mathcal{C}}(s)$$

and reserve $\leftrightarrow$ for material biconditionals *inside* $\mathcal{C}$.

Compact Axiom Schema.

$$\forall s \in \mathcal{U}_{\mathcal{C}}: \quad \mathsf{Affirm}_{\mathcal{C}}(s) \;\Rightarrow\; \mathsf{Conceive}_{\mathcal{C}}\Big(\mathbf{D}_{\mathcal{C}}(s)\Big), \qquad \mathbf{D}_{\mathcal{C}}^2 = \mathrm{Id}, \qquad \Delta_{\mathcal{C}}^2 = \mathrm{Id},$$

$$\mathsf{Status}_{\mathcal{C}}(\mathbf{D}_{\mathcal{C}}(\varphi), \lambda) = \Delta_{\mathcal{C}}\Big(\mathsf{Status}_{\mathcal{C}}(\varphi, \lambda)\Big), \qquad \neg\mathsf{App}_{\mathcal{C}}(\chi) \;\Rightarrow\; \neg\mathsf{App}_{\mathcal{C}}\Big(\mathbf{D}_{\mathcal{C}}(\chi)\Big).$$

(The last two clauses capture status-compatibility and invariance of inapplicability.)

The Example Set of Dyads

$$\begin{aligned}\mathcal{E}_{\text{duality}} := \Big\{&\mathfrak{D}_{\mathcal{C}}(\text{Existence}),\ \mathfrak{D}_{\mathcal{C}}(\text{Eternalism}),\ \mathfrak{D}_{\mathcal{C}}(\text{Self}),\ \mathfrak{D}_{\mathcal{C}}(\text{Emptiness}),\\ &\mathfrak{D}_{\mathcal{C}}(\text{Continuous}),\ \mathfrak{D}_{\mathcal{C}}(\text{Duality}),\ \mathfrak{D}_{\mathcal{C}}(\text{Created}),\ \mathfrak{D}_{\mathcal{C}}(\text{Finite}),\\ &\mathfrak{D}_{\mathcal{C}}(\text{Form}),\ \mathfrak{D}_{\mathcal{C}}(\text{Time}),\ \mathfrak{D}_{\mathcal{C}}(\text{True}),\ \mathfrak{D}_{\mathcal{C}}(\text{Positive}),\ \mathfrak{D}_{\mathcal{C}}(\text{Real})\Big\}.\end{aligned}$$

Illustrative pairs. (1) Existence / Non-Existence, (2) Eternalism / Nihilism, (3) Self / Not-Self, (4) Emptiness / Non-Emptiness, (5) Continuous / Discontinuous, (6) Duality / Non-Duality, (7) Created / Un-Created, (8) Finite / Infinite, (9) Form / Formlessness, (10) Time / Timelessness, (11) True / False, (12) Positive / Negative, (13) Real / Un-Real.

1.2.7 The Fourfold Dialectic Algebra of Symmetry

The foundational logic of the IQR framework is a generative, tetradic system termed the *Fourfold Dialectic Algebra of Symmetry*. It bridges the ancient catuṣkoṭi, Aristotelian dialectic, and our modern formalizations of Thesis, Antithesis, Synthesis, and the Holothesis, while preserving the ontological state realizations that anchor the quantum constructions.

Definition 1.16 (Fourfold Dialectic Algebra of Symmetry)**.** For each Primordial Quantum Singularity $\mathfrak{B}_x$ (Definition 1.25), let

$$\mathcal{H}_{\text{mod}}(\mathfrak{B}_x) \quad \text{and} \quad \{\hat{\mathcal{F}}_\alpha(x)\}_{\alpha\in\{0,1,p,s\}}$$

denote, respectively, the modal fibre and the four symmetry *modes* acting on it. The *Fourfold Dialectic $*$-algebra of Symmetry at $\mathfrak{B}_x$* is the $*$-subalgebra

$$\mathcal{A}_x^{(4)} \;:=\; {}^*\mathrm{alg}\left\langle \hat{\mathcal{F}}_0(x), \hat{\mathcal{F}}_1(x), \hat{\mathcal{F}}_p(x), \hat{\mathcal{F}}_s(x) \right\rangle \;\subseteq\; \mathcal{B}\Big(\mathcal{H}_{\text{mod}}(\mathfrak{B}_x)\Big),$$

whose four generators are *congruent*, via the dialectic–quantisation correspondence, to the modernised tetralemma modalities (Thesis, Antithesis, Synthesis, Holothesis). Congruence here means the correspondence preserves the involutive duality, synthesis rules, and meta-level structure (Definitions 1.1 to 1.4, 2.1 and 2.12):

1. **Symmetry** $\cong$ **Thesis** (Definition 1.1). *Ontological representative (ray):*

$$|\mathring{\mathbf{0}}\rangle_x \in \mathcal{H}_{\mathrm{mod}}(\mathfrak{B}_x), \qquad \text{carried by } \hat{\mathcal{F}}_0(x).$$

2. **Antisymmetry** $\cong$ **Antithesis** (Definition 1.2). *Ontological anti-representative (ray):*

$$|\mathring{\mathbf{1}}\rangle_x \in \mathcal{H}_{\mathrm{mod}}(\mathfrak{B}_x), \qquad \text{carried by } \hat{\mathcal{F}}_1(x),$$

with the global involutive duality implementing $|\mathring{\mathbf{0}}\rangle_x \leftrightarrow |\mathring{\mathbf{1}}\rangle_x$ (Definitions 2.1 and 2.12).

3. **Parasymmetry** $\cong$ **Synthesis** (Definition 1.3). *Canonical ontological superposition (phase–conventioned):*

$$\frac{1}{\sqrt{2}}\big(|\mathring{\mathbf{0}}\rangle_x + |\mathring{\mathbf{1}}\rangle_x\big)$$

generated/selected by $\hat{\mathcal{F}}_p(x)$ (cf. Section 2.2 and Definitions 2.85 and 2.94). (More general relative phases are gauge-equivalent under local U(1) choices.)

4. **Holosymmetry** $\cong$ **Holothesis** (Definition 1.4). *Meta-representational coupling:*

$$\mathring{\mathbf{\Lambda}}_{\mathring{\mathbf{0}}\mathring{\mathbf{1}}}(x) \quad \text{carried by } \hat{\mathcal{F}}_s(x),$$

encoding the holosymmetric meta-structure that coheres the corners at the operator level (Sections 2.3, 3.8 and 12.2).

We refer to $(\mathcal{A}_x^{(4)}; \hat{\mathcal{F}}_0, \hat{\mathcal{F}}_1, \hat{\mathcal{F}}_p, \hat{\mathcal{F}}_s)$ as the *Fourfold Dialectic Algebra of Symmetry at* $\mathfrak{B}_x$, with ontological representatives preserved by the correspondence to the modernised tetralemma.

Remark 1.17 (On congruence and gauge)**.** The symbol "$\cong$" indicates a structure-preserving identification of logical modalities with quantum modes that (i) intertwines the global involutive duality, (ii) respects synthesis/parasymmetry construction, and (iii) lifts to meta-level holosymmetry. Representatives such as $|\mathring{\mathbf{0}}\rangle_x$, $|\mathring{\mathbf{1}}\rangle_x$ are defined up to local frame and phase; statements are gauge-covariant under the bundle connection (Lemma 4.270).

The Fourfold Dialectic Algebra says, in effect, that each Primordial Quantum Singularity (Definition 1.25) carries a *four-mode compass* whose points line up one-to-one with the modernized tetralemma. The congruence map Φ identifies Symmetry $\leftrightarrow$ Thesis, Antisymmetry $\leftrightarrow$ Antithesis, Parasymmetry $\leftrightarrow$ Synthesis, and Holosymmetry $\leftrightarrow$ Holothesis, while *preserving the ontological anchors* (Corollary 1.21). That preservation clause matters: it guarantees that anything we build on the logical side—states, projectors, or processes—can be transported to the symmetry side *without changing what the system ontologically "is."* In practice, $|\mathring{\mathbf{0}}\rangle$ and $|\mathring{\mathbf{1}}\rangle$ realize the Thesis/Antithesis modes (Definition 2.1); their coherent blend $\frac{1}{\sqrt{2}}\big(|\mathring{\mathbf{0}}\rangle + |\mathring{\mathbf{1}}\rangle\big)$ realizes Synthesis; and the holosymmetric meta-symbol $\mathring{\mathbf{\Lambda}}_{\mathring{\mathbf{0}}\mathring{\mathbf{1}}}$ marks the Holothesis "about the whole" rather than another point on the same plane (Sections 2.3 and 3.8).

Intuitively, the four modes are *tests we can actually run*, tied to the Born-valued semantics (Section 1.2.9 and Definition 1.18). Thesis/Antithesis correspond to measuring the distinguished axis fixed by the ontological duality basis (a Σ_z-like readout); Synthesis

is the *complementary probe* that asks whether the system organizes those two tendencies coherently (a Σ_x-like readout); and Holothesis is a *controlled interferometric readout* that asks a meta-question about which global evolution is coherently realized. Because "truth" is the expectation value $v(\rho; P) = \mathrm{Tr}(P\rho)$, these corners are not mutually exclusive by fiat: a single preparation ρ can carry simultaneous warranted degrees for Thesis and Antithesis, or land outside their plane (the "Neither" region), with Synthesis and Holothesis revealing different aspects of that situation.

Finally, calling it an *algebra* highlights the structural rules that keep everything consistent. Operators that commute with the Thesis projector P_T (respectively P_A) conserve that stance; the local duality involution swaps Thesis $\leftrightarrow$ Antithesis while acting trivially off their sector; and the parasymmetric/anti-parasymmetric pair resolves the $(0, 1)$-sector without "leakage," ensuring closure with the Thesis/Antithesis frame. At the reasoning level, the *prism behavior* (Definition 1.23 and Proposition 1.24) explains how different contexts select different subsets of modes: classical contexts pick one of the poles, paraconsistent contexts allow Synthesis alongside them *without explosion*, and meta-contexts license Holosymmetry when the base-level polarity is inapplicable. In short, the definition packages a bridge (logic $\leftrightarrow$ symmetry), a semantics (Born-valued truth), and an operational reading (concrete tests) into one compact scaffold.

1.2.8 A Prism for Discourse: Classical Logic as a Special View

Classical binary logic furnishes a powerful fortress of certainty: propositions are either true or false; contradiction is disallowed; there is no middle ground [73, 280]. The fourfold dialectic framework does not tear down this fortress; it *re-situates* it. Classical logic appears when the world is interrogated along a single axis—*Thesis/Antithesis*—while two further, indispensable axes are also available: *Synthesis* ("both, coherently") and *Holothesis* ("neither," more deeply "structure at the whole") [15, 273, 17]. Formally, these correspond to *Symmetry*, *Antisymmetry*, *Parasymmetry*, and *Holosymmetry* in the Fourfold Dialectic Algebra (Section 1.2.7).

The central shift is to treat logic as *diagnostic interrogation*. A single discourse—like white light through a prism—is refracted into four complementary facets. Each facet returns a quantitative *degree of warrant* $v \in [0, 1]$ via Born valuation, so "truth" is an operational propensity rather than a bare bivalent label (Section 1.2.9).

The Formal Core and Its Operational Meaning

Let the state of warrant be a density operator ρ on a Hilbert space. Each facet $\square \in \{\mathrm{T}, \mathrm{A}, \mathrm{S}, \mathrm{H}\}$ corresponds to a positive operator $E_\square$ (an *effect*). The facet's reading is the Born value

$$v_\square(\rho) = \mathrm{Tr}(E_\square \rho) \in [0, 1]$$

(Definition 1.18). Concretely, in the distinguished $(0, 1)$-sector, *Thesis* and *Antithesis* are the projectors $P_T = |e_1\rangle\langle e_1|$, $P_A = |e_2\rangle\langle e_2|$ (Definitions 1.18 and 2.1).

Synthesis is realized as the *parasymmetric* projector $P_{\text{para}} = |+\rangle\langle +|$ with $|+\rangle = (|e_1\rangle + |e_2\rangle)/\sqrt{2}$ (Definition 2.94). The pair $\{P_{\text{para}}, P_{\text{antipara}}\}$ is orthogonal and *locally complete* on the $(0, 1)$-sector:

$$P_{\text{para}} + P_{\text{antipara}} = P_{(0,1)} = P_T + P_A,$$

so repeated trials directly estimate $v_{\text{S}}(\rho) = \text{Tr}(P_{\text{para}}\rho)$ (Definition 2.94). *Holothesis* is a *global* facet. Operationally it is an effect $E_{\text{Holo}}^{(\mathbf{a}\to\mathbf{b})}$ induced by a holosymmetric controlled unitary that compares *entire histories* (alternative global evolutions) and reads the result back on the registers; its valuation is

$$v_{\text{H}}(\rho) = \text{Tr}\Big(E_{\text{Holo}}^{(\mathbf{a}\to\mathbf{b})}\rho\Big)$$

(eq. (3.36) and Definition 1.18).

What "both" means (and what it doesn't)

Synthesis is *not dialetheism*. It does not say "P and $\neg P$" are literally both true; rather, it detects *coherent co-presence*—quantum superposition—via an interference-sensitive projector P_{para}. Its signal vanishes for mixtures and grows with phase coherence, exactly the quantum distinction between superpositions and statistical ensembles [287, 282]. In multipartite settings, Synthesis scales naturally to *relational* structure (Bell-like parasymmetric states), realizing "both" as distributed entanglement (Section 2.2.3).

What "neither" means—and the deeper role of H

Holothesis can register "neither true nor false" when the T/A frame is inapplicable, but its essence is subtler: it is an *implementation-invariant, system-level interrogation*. It asks whether the discourse's status survives *swaps of whole mechanisms*—e.g., exchanging "globally coupled" vs "decoupled" evolutions in dual registers and comparing them coherently (Section 3.8.1). Equivalences under control-basis rotations, register swaps, and common conjugations show that H is about *the whole architecture*, not local content ($H4$, Definition 3.18).

Why the classical slice reappears

This expansion does not violate classical laws; it *contains* them. The *Axiom (Principle) of Duality* ensures that T/A form a complete pair on their domain: $P_T + P_A = P_{(0,1)}$. Hence, on that sector, $v_T + v_A = 1$: the *Excluded Middle* is recovered (Section 1.2.6 and Definition 2.100).

The *Law of Non-Contradiction* is recovered in the *commuting/decohered limit*: when interrogations commute or decoherence kills off-diagonals, the S-facet carries no independent signal and T/A become crisp, so "both-true" cannot be registered ($P_T P_A = 0$) [282]. In that regime, the fourfold prism collapses to the classical fortress. (See also the local completeness and orthogonality identities for $P_{\text{para}}, P_{\text{antipara}}$ in Definition 2.94.)

What the prism yields beyond the fortress

- *Resolution vs coherence.* T/A resolve local alternatives; S reads *coherent coexistence* (superposition). Classical logic can distinguish only mixtures, not coherence [287].
- *Local vs global.* H interrogates *architecture*—invariance under exchanging entire mechanisms—via holosymmetric effects (eq. (3.36)).

- *Quantification.* All four facets return *Born-valued* degrees of warrant, making "truth" an operational propensity (Definition 1.18).

How to read a discourse with the prism

Given ρ, perform four complementary interrogations:

1. *T/A:* $v_T(\rho), v_A(\rho)$ — the classical slice (Definition 1.18).
2. *S:* $v_S(\rho) = \mathrm{Tr}(P_{\mathrm{para}}\rho)$ — coherence of the T/A split (Definition 2.94).
3. *H:* $v_H(\rho) = \mathrm{Tr}(E_{\mathrm{Holo}}^{(\mathbf{a}\to\mathbf{b})}\rho)$ — global, implementation-invariant status (eq. (3.36)).

These questions are not mutually reducible; they resolve different aspects of structure (cf. the logic→algebra→representation pipeline Section 1.4.2).

Classical laws in their regime of validity

Nothing here negates classical laws *in their proper regime*:

1. *Non-Contradiction* emerges as a theorem for *commuting interrogations* and *decohered states*—the geometry where S has no distinctive signal [282].
2. *Excluded Middle* is recovered when only T/A valuations are meaningful and crisp; Duality guarantees $P_T + P_A = P_{(0,1)}$ (Definition 2.100).

Thus classical logic is the bright central band of a broader spectrum. Where contexts commute and interference vanishes, the prism narrows to the fortress. Where coherence and global structure matter, the additional facets—*Synthesis* and *Holothesis*—let us register *coherent coexistence* and *holistic invariants*, in line with the empirical grammar of quantum phenomena [168, 287, 282] and the philosophical insight of the catuṣkoṭi [15, 46, 17].

1.2.9 Born-Valued Semantics for the Four Corners

Definition 1.18 (Dialectic Truth as Born Valuation)**.** Let

$$M \in \{\text{Thesis}, \text{Antithesis}, \text{Synthesis}, \text{Holothesis}\}.$$

Associate to each corner a quantum *effect* E_M (a positive semidefinite operator with $0 \le E_M \le \mathbb{I}$) on the relevant Hilbert space. For any density operator ρ, the *truth value* (degree of warrant) of M in state ρ is

$$v_M(\rho) := \mathrm{Tr}(E_M\,\rho) \in [0,1].$$

Schema. For any effect P representing one of the four corners (projector or general POVM element),

$$v(\rho; P) := \mathrm{Tr}(P\,\rho) \in [0,1]. \tag{1.2}$$

When P is a projector and $\rho = |\psi\rangle\langle\psi|$, this reduces to $v(\rho; P) = \|P|\psi\rangle\|^2$.. Concretely, in the Symmetry–Antisymmetry sector $\mathcal{H}_{(0,1)} = \mathrm{span}\{|e_1\rangle, |e_2\rangle\}$ (cf. Definition 2.1):

$$E_{\text{Thesis}} := P_T := |e_1\rangle\langle e_1|, \qquad E_{\text{Antithesis}} := P_A := |e_2\rangle\langle e_2|.$$

For Synthesis, with $|+\rangle := (|e_1\rangle + |e_2\rangle)/\sqrt{2}$,

$$E_{\text{Synthesis}} := P_S := |+\rangle\langle +|.$$

For Holothesis (a meta-level choice measured via the holosymmetric control), fix a holosymmetric unitary $\mathbf{U}_{\text{holo}}$ on the control $\mathcal{H}_c$ and dual-register $\mathcal{H}_{\text{RR}}^{(2n)}$ (cf. Sections 2.3 and 3.8). Preparing the control in $|\mathbf{a}\rangle_c$ and reading out in $|\mathbf{b}\rangle_c$, the induced *effect* on the register is

$$E_{\text{Holothesis}}^{(\mathbf{a}\to\mathbf{b})} := \Big\langle \mathbf{a}\Big|_c \mathbf{U}_{\text{holo}}^\dagger \left(|\mathbf{b}\rangle\langle\mathbf{b}|_c \otimes \mathbb{I}\right) \mathbf{U}_{\text{holo}} \Big|\mathbf{a}\Big\rangle_c,$$

so that the measured Holothesis valuation for a register state ρ_{RR} is

$$v_{\text{Holothesis}}(\rho_{\text{RR}}) = \text{Tr}\Big(E_{\text{Holothesis}}^{(\mathbf{a}\to\mathbf{b})}\, \rho_{\text{RR}}\Big).$$

Remark 1.19 (Operational notes)**.** (i) For composite configurations, local effects extend by identity on the other factors; e.g. $P_T^{(x_i)} := |e_1\rangle\langle e_1|_{x_i} \otimes \mathbb{I}_{\text{rest}}$. (ii) The four effects are associated to *different tests* and need not form a single POVM simultaneously; incompatibility is expected. (iii) Numerical "truth values" are probabilities (Born rule), not amplitudes.

Lemma 1.20 (Bridge Map: Symmetry modes $\leftrightarrow$ Tetralemma corners)**.** *Let*

$$\Phi : \{\text{Sym}, \text{AntiSym}, \text{ParaSym}, \text{HoloSym}\} \xrightarrow{\cong} \{\text{Th}, \text{AntiTh}, \text{SynTh}, \text{HoloTh}\}$$

be the bijection $\Phi(\text{Sym}) = \text{Th}$, $\Phi(\text{AntiSym}) = \text{AntiTh}$, $\Phi(\text{ParaSym}) = \text{SynTh}$, $\Phi(\text{HoloSym}) = \text{HoloTh}$. *Write* $\mathfrak{D}_{\mathcal{C}}(\varphi) = \langle L(\varphi), R(\varphi)\rangle$ *for the dyad and let* Sel_{Th}, $\text{Sel}_{\text{AntiTh}}$, $\text{Sel}_{\text{SynTh}}$, $\text{Sel}_{\text{HoloTh}}$ *be the corner selections of Proposition 1.15. Define mode selections by congruence:*

$$\text{Sel}_{\text{Sym}} := \text{Sel}_{\text{Th}}, \quad \text{Sel}_{\text{AntiSym}} := \text{Sel}_{\text{AntiTh}}, \quad \text{Sel}_{\text{ParaSym}} := \text{Sel}_{\text{SynTh}}, \quad \text{Sel}_{\text{HoloSym}} := \text{Sel}_{\text{HoloTh}}.$$

Then for all admissible λ *and* φ,

$$\text{Sel}_M\Big(\mathfrak{D}_{\mathcal{C}}(\varphi); \lambda\Big) = \text{Sel}_{\Phi(M)}\Big(\mathfrak{D}_{\mathcal{C}}(\varphi); \lambda\Big).$$

Moreover, if $\Delta_{\mathcal{C}}$ *is the dualization on corners (Axiom 1.10), there exists the symmetry-side involution*

$$\begin{aligned} \Sigma :& \{\text{Sym}, \text{AntiSym}, \text{ParaSym}, \text{HoloSym}\} \to \{\cdots\}, \\ \Sigma(\text{Sym}) &= \text{AntiSym}, \\ \Sigma(\text{AntiSym}) &= \text{Sym}, \\ \Sigma(\text{ParaSym}) &= \text{ParaSym}, \\ \Sigma(\text{HoloSym}) &= \text{HoloSym}. \end{aligned}$$

such that the diagram commutes: $\Delta_{\mathcal{C}} \circ \Phi = \Phi \circ \Sigma$.

Corollary 1.21 (State/meta representations preserved)**.** *Under the correspondence of Lemma 1.20 the ontological representations agree:*

$$\begin{aligned}\mathrm{Rep}(\mathrm{Sym}) &= |\mathring{\mathbf{0}}\rangle,\\ \mathrm{Rep}(\mathrm{AntiSym}) &= |\mathring{\mathbf{1}}\rangle,\\ \mathrm{Rep}(\mathrm{ParaSym}) &= \tfrac{1}{\sqrt{2}}\big(|\mathring{\mathbf{0}}\rangle + |\mathring{\mathbf{1}}\rangle\big),\\ \mathrm{Rep}(\mathrm{HoloSym}) &= {}_{\mathring{\mathbf{0}}}\mathring{\mathbf{\Lambda}}_{\mathring{\mathbf{1}}}.\end{aligned}$$

Hence all quantum-mechanical realizations/constructions attached to the corners carry over verbatim to the symmetry modes.

Remark 1.22 (Ontological vs. conventional quantum states)**.** The ontological states above are pre-geometric logical primitives; conventional quantum states are emergent structures built from their superposition/entanglement.

The Fourfold Symmetry

Definition 1.23 (Symmetry as a four-mode logical prism)**.** Let

$$\mathsf{Modes} := \{\mathrm{Sym}, \mathrm{AntiSym}, \mathrm{ParaSym}, \mathrm{HoloSym}\} \quad \text{and}$$
$$\Phi : \mathsf{Modes} \xrightarrow{\cong} \{\mathrm{Th}, \mathrm{AntiTh}, \mathrm{SynTh}, \mathrm{HoloTh}\}$$

be the congruence $\Phi(\mathrm{Sym}) = \mathrm{Th}$, $\Phi(\mathrm{AntiSym}) = \mathrm{AntiTh}$, $\Phi(\mathrm{ParaSym}) = \mathrm{SynTh}$, $\Phi(\mathrm{HoloSym}) = \mathrm{HoloTh}$. Preserve the ontological anchors

$$\begin{aligned}\mathrm{Rep}(\mathrm{Sym}) &= |\mathring{\mathbf{0}}\rangle,\\ \mathrm{Rep}(\mathrm{AntiSym}) &= |\mathring{\mathbf{1}}\rangle,\\ \mathrm{Rep}(\mathrm{ParaSym}) &= \tfrac{1}{\sqrt{2}}\big(|\mathring{\mathbf{0}}\rangle + |\mathring{\mathbf{1}}\rangle\big),\\ \mathrm{Rep}(\mathrm{HoloSym}) &= {}_{\mathring{\mathbf{0}}}\mathring{\mathbf{\Lambda}}_{\mathring{\mathbf{1}}}.\end{aligned}$$

For any well-formed φ and locus λ, define the *symmetry spectrum* selected by the context $\mathcal{C}$:

$$\mathrm{Spec}_{\mathcal{C}}(\varphi,\lambda) \;:=\; \Big\{\, m \in \mathsf{Modes} \;:\; \mathrm{Sel}_{\Phi(m)}\big(\mathfrak{D}_{\mathcal{C}}(\varphi);\lambda\big) \text{ holds} \,\Big\}.$$

Proposition 1.24 (Prism behavior by reasoning regime)**.** *(i) If $\mathcal{C}$ is classical (non-paraconsistent) and presumes applicability of the existential polarity, then* $\mathrm{Spec}_{\mathcal{C}}(\varphi,\lambda) \in \Big\{\{\mathrm{Sym}\},\{\mathrm{AntiSym}\}\Big\}$.
(ii) If $\mathcal{C}$ admits a paraconsistent $\vdash_{\mathrm{pc}}$, then ParaSym *may also be selected jointly with* Sym *and* AntiSym *(non-explosively).*
(iii) If $\mathcal{C}$ licenses meta-level inapplicability (Holothesis), then $\mathrm{HoloSym} \in \mathrm{Spec}_{\mathcal{C}}(\varphi,\lambda)$ *and the existential polarity is inapplicable to both coordinates.*



Figure 1.1: The four corners (tetrahedron) of the dialectic.

Etymology and Semantics

The word "symmetry" traces its origins back to the Ancient Greek term *symmetria* (συμμετρία) [293, 299]. This word is a compound of two Greek roots: *syn* (σύν), meaning "together" or "with", and *metron* (μέτρον), meaning "measure" [293, 299]. Together, they form a word that literally translates to "common measure" or "agreement in dimensions". In classical Greece, this concept was not just a mathematical abstraction but a fundamental principle of beauty and order, representing a harmonious and pleasing relationship between the various parts of a whole, whether in sculpture, architecture, or the cosmos itself [300, 301].

From Greece, the word journeyed into *Latin* as "symmetria", maintaining its form and meaning [302]. It was later adopted into *French* as "symétrie" during the 16$^{\text{th}}$ century, from which it was borrowed into English in the mid-1500s [296]. Initially, it retained its broader, classical sense of "due proportion" or a "harmonic arrangement", a meaning that was especially influential during the Renaissance among artists and architects who sought to emulate classical ideals of balance and form [296].

Over time, the meaning of "symmetry" began to evolve toward a more specific, scientific definition [296]. By the 17$^{\text{th}}$ and 18$^{\text{th}}$ centuries, it was increasingly used to describe the geometric concept of an object having two halves that are mirror images of each other, a notion known as *bilateral symmetry* [296]. This definition became particularly important in the developing fields of biology and natural history [296]. By the 19$^{\text{th}}$ and 20$^{\text{th}}$ centuries, the concept was fully formalized within mathematics and physics [300, 301]. In this modern context, symmetry describes the property of an object or system remaining unchanged (invariant) under specific transformations such as reflection, rotation, or translation, making it a cornerstone principle in fields ranging from crystallography to particle physics [300, 301].

The Sanskrit prefix *para-* (पर) points to a rich philosophical concept of transcendence rooted in the simple physical act of crossing a boundary. It descends from the Proto-Indo-European root *per-*, which meant "through, across, or beyond" [81, 292]. This ancient heritage is preserved in its direct cognates, such as Greek *pára* (παρά "beside, beyond"), and Latin *per* ("through") [293, 299, 81]. In Sanskrit, this idea of "crossing over" was elevated from the physical to the metaphysical, giving *para* its meaning of "other", "beyond", and ultimately "transcendent" [298]. This deeper sense is captured in key philosophical terms like *pāramitā* ("perfection", literally "gone to the other shore") and *paramārtha* ("ultimate truth") [298].

The prefix *holo-* is a direct borrowing from Ancient Greek, chosen for its powerful and precise meaning of integrated wholeness. Its use in modern scientific and philosophical terms like "hologram", "holistic", and the proposed "holo-state" is a deliberate invocation of an ancient concept: the idea of a system that is complete, entire, and more than the mere sum of its parts. This prefix signals a departure from a purely analytic or reductionist view, pointing instead to a reality where the identity and nature of the whole are irreducibly primary.

The term's lineage begins with the Ancient Greek word *ὅλος (hólos)*, which means "whole", "entire", "complete", or "total" [303]. In Greek thought, *hólos* was not just a quantitative measure of all the parts, but a qualitative descriptor of an integrated, organic unity. Aristotle, for instance, famously argued that a whole is something over and above its parts, possessing properties and a nature that cannot be understood by studying its components in isolation [73]. The word traces back to the Proto-Indo-European root *solh₂-*, meaning "whole" or "intact", which also gave rise to the Latin word *solidus* (solid) and the Sanskrit word *śarva* (सर्व, "all, whole, entire") [81]. This deep, shared linguistic ancestry reveals a foundational human intuition for the concept of an inviolable, complete entity.

The modern power of the prefix was significantly amplified by the invention of the *hologram*, a term coined by Dennis Gabor in 1949 [304, 305]. A hologram is a recording of a light field that exhibits a remarkable property: each part of the recording contains information about the entire three-dimensional image [306, 307, 308]. This physical manifestation of "the whole in every part" provided a powerful scientific metaphor for interconnectedness [309]. It transformed the ancient philosophical concept of *hólos* into a tangible principle, demonstrating a new kind of order where the global structure is enfolded within every local region [309].

The use of *holo-* in terms like *Quantum Holo-State* (Section 1.4) and *Holosymmetry* (Section 2.3) is therefore a direct appeal to this rich lineage. The *Holo-State*,then is not just a state that represents all the parts of the universe, but one that describes the universe as a single, indivisible quantum whole, where the information of the entire system is fundamentally entangled and non-local. Consequently, *Holosymmetry* alludes to a type of symmetry of this unbroken wholeness, a kind of symmetry that reflects the total, integrated nature of the Holo-State, a principle of invariance that is a property of the entire cosmic system, not just its isolated components.

1.3 The Primordial Quantum Singularity

The Big Bang, Singularities, and Cosmic Fates

The **Big Bang theory** is our leading framework for understanding the early universe. It describes how, from an extremely hot and dense state approximately 13.8 billion years ago, spacetime expanded and cooled, allowing for the formation of matter, radiation, and eventually the large-scale structures we observe today. The theory is supported by robust observational pillars, including the systematic redshifts of distant galaxies and the relic **cosmic microwave background**—a nearly uniform afterglow from that hot, early phase. These observations, combined with precise parameter fits from galaxy surveys and CMB data, provide a consistent cosmological model [310, 311]. When this classical description from General Relativity (GR) is traced backward in time, the expanding solution reaches a boundary where the model

can no longer be extended, indicating an "initial singularity" [312]. However, this is best understood not as a literal point of infinite density, but as a limit of the classical theory itself, signaling that a more fundamental theory of quantum gravity is required to describe the universe's earliest moments [240, 7, 8, 3, 12].

A possible, though now disfavored, fate for the universe is the **Big Crunch**, which is essentially the time-reverse of the Big Bang. In this scenario, the mutual gravitational attraction of all matter in the cosmos would eventually halt the expansion and cause the universe to recontract, growing hotter and denser until it collapses into a terminal singularity. Classically, this is a perfectly consistent fate within GR, provided the universe contains the right amount of energy and matter [7]. However, pivotal observations in the late 1990s revealed that the cosmic expansion is, in fact, **accelerating** [134, 135]. This acceleration strongly suggests the presence of a "dark energy" component with negative effective pressure, a dynamic that makes a future recollapse highly unlikely [313]. Instead, the long-term evolution of our universe appears to be heading toward a "Big Freeze"—an ever colder, more diffuse, and emptier cosmos [314, 315].

A **black hole** is another consequence of gravity predicted by GR, defined as a region of spacetime from which nothing, not even light, can escape. These objects form in nature through the complete gravitational collapse of sufficiently massive objects and are described by solutions in GR that feature an **event horizon**—the point of no return [7, 316, 317]. Within this horizon, continued collapse leads to an inner region where classical GR predicts its own breakdown in the form of a singularity [240, 312]. For a non-rotating (Schwarzschild) black hole, this singularity is often depicted as a point, whereas for a rotating (Kerr) black hole, the theory predicts a **ring singularity** [318]. In realistic scenarios, this singularity is thought to be hidden from outside observers by the event horizon, a principle known as cosmic censorship. A complete and physically realistic description of what happens at the very center of a black hole, however, is expected to require a full theory of quantum gravity [7, 8, 3].

What a Singularity Means in General Relativity

In general relativity, a "singularity" is not a literal place where physics "begins" or "ends," but a failure of the spacetime model: *geodesic incompleteness*, i.e., the existence of timelike or null geodesics that cannot be extended to arbitrarily large values of their affine parameters. In many—but not all—cases this accompanies the blow-up of curvature invariants such as the Kretschmann scalar $K = R_{abcd}R^{abcd}$ [7, 240, 146].

The focusing mechanism behind many singularity results is captured by *Raychaudhuri's equation* for a timelike geodesic congruence with 4-velocity u^a, expansion θ, shear σ_{ab}, and vorticity ω_{ab}:

$$\frac{d\theta}{d\tau} = -\frac{1}{3}\theta^2 - \sigma_{ab}\sigma^{ab} + \omega_{ab}\omega^{ab} - R_{ab}u^a u^b.$$

Assuming vanishing vorticity ($\omega_{ab} = 0$) and appropriate energy conditions (e.g., strong or null), the congruence inevitably *focuses* so that $\theta \to -\infty$ in finite proper time, leading to caustic formation. When combined with global geometric inputs (e.g., trapped surfaces), this focusing yields *geodesic incompleteness* via the Penrose–Hawking singularity theorems [7, 240, 316, 312].

Big Bang: did it "emerge from a singularity"?

In homogeneous/isotropic cosmology Friedmann–Lemaître–Robertson–Walker (FLRW),

$$ds^2 = -c^2dt^2 + a^2(t)\,\gamma_{ij}dx^i dx^j, \qquad H^2 = \left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3}\rho - \frac{kc^2}{a^2} + \frac{\Lambda c^2}{3},$$

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3}\left(\rho + \frac{3p}{c^2}\right) + \frac{\Lambda c^2}{3}.$$

For standard matter/radiation ($\rho > 0$, $p \geq 0$, and not-too-large Λ), evolving backward drives $a(t) \to 0$ in finite proper time; typical densities and curvature scalars diverge in the classical model. That "initial singularity" is *not a place in space* but a *past boundary* of the classical solution. In GR the mathematically precise statement is *geodesic incompleteness*: the FLRW solution cannot be extended past that boundary within the classical theory [240, 7, 144, 319].

Two rigorous points often missed:

1. The classic singularity theorems establish *incompleteness* under general conditions (energy conditions, trapped sets), not necessarily a smooth point where curvature "blows up." [240, 7]

2. *Inflation does not generically evade past-incompleteness*: many inflating spacetimes are still past-incomplete (Borde–Guth–Vilenkin) [320]

What happens "before" the classical boundary is a *quantum-gravity* question. Competing proposals include loop-quantum-cosmology bounces, [228, 321] ekpyrotic/cyclic scenarios, [322, 323] Penrose's conformal cyclic cosmology, [227] asymptotic safety, [324] string-gas cosmology, [325] and no-boundary initial conditions [51]. There is *no consensus* yet; current observations—especially CMB and large-scale structure—are well fit by ΛCDM with an inflationary early phase and show no decisive deviations that fix the pre-inflationary completion [150, 326, 327, 319].

Big Crunch: time-reversed Big Bang?

A **Big Crunch** is a future singularity in which the scale factor recontracts to $a(t) \to 0$ in finite proper time. In the classical $\Lambda = 0$ setting, closed universes ($k = +1$, $\Omega_{\text{tot}} > 1$) naturally recollapse; more generally, the fate hinges on the energy content and its effective equation of state for $a(t)$ [7, 144].

1. *Positive, nearly constant* Λ leads to asymptotic de Sitter expansion—eternal acceleration and no Crunch [314, 315].

2. *Negative* Λ or evolving dark energy that becomes effectively attractive can trigger turnaround and recollapse [144, 313].

3. Other exotic fates exist (e.g., *Big Rip* if $w < -1$) [313].

Empirically, late-1990s supernova observations established that the expansion is *accelerating*, consistent with dark energy close to $w \approx -1$, which disfavors a Crunch under current conditions; precision CMB and large-scale structure results reinforce this picture [134, 135, 150, 313].

Black holes and their singularities

For a Schwarzschild (non-rotating, uncharged) black hole, the surface $r = 2GM/c^2$ is an *event horizon* that is regular—its apparent pathology in Schwarzschild t is a coordinate artifact—while the interior, in the classical solution, terminates at $r = 0$, where curvature invariants (e.g., the Kretschmann scalar) diverge [7, 316]. The $r = 0$ singularity is *spacelike*, and any infaller reaches it in *finite proper time* of order GM/c^3 after crossing the horizon [7, 316].

For rotating (Kerr) or charged (Reissner–Nordström) solutions, the maximal analytic extension includes an inner (Cauchy) horizon and, in Kerr, a *ring singularity*; the inner horizon is widely expected to be *unstable* (mass-inflation type blue-shifts), so the smooth interior of the ideal solution is unlikely to survive realistic collapse [318, 316]. Penrose's *weak cosmic censorship* conjecture asserts that physically formed singularities remain hidden behind horizons (no generic "naked" singularities); while unproven in full generality, it is supported in many settings [312, 7]. (For formation via stellar collapse and compact-object context, see classic treatments [328, 317].)

Information, evaporation, and singularity resolution

Semiclassically, black holes radiate thermally via quantum fields on curved spacetime—the Hawking effect—so an isolated black hole loses mass and *evaporates* over time [87, 12]. In classical GR, the interior *singularity* remains part of the spacetime all the way to the endpoint of evaporation, which sharpens the *information problem*: how can unitary quantum evolution be reconciled with apparently thermal, information-erasing Hawking radiation [329, 330, 86, 331, 332, 333, 334]

A complementary, quantitative way to pose the puzzle is via the *Page curve*: if evaporation is unitary, the entropy of the radiation should rise and then fall after the Page time. Modern formulations (Hayden–Preskill; islands/replica-wormholes) provide concrete mechanisms for recovering this unitary Page curve in toy models and semiclassical gravity [335, 138, 336, 139, 337].

Candidate *resolutions* often align with mechanisms that also *tame the singularity*:

1. *Quantum-geometry replacements.* In loop-quantum-gravity–inspired models, strong-curvature interiors can be replaced by non-singular quantum phases (e.g., "Planck-star" or bounce scenarios), turning the would-be singular endpoint into a transition [42, 338, 339].

2. *String-theoretic microstate geometries / fuzzballs.* Black-hole microstates can be realized as horizon-sized, non-singular configurations, eliminating the need for a traditional interior singularity [340].

3. *Asymptotic safety / effective corrections.* If gravity flows to a nontrivial UV fixed point, high-curvature growth of invariants can be softened, potentially averting singular behavior within an effective description [341].

4. *Complementarity, state-dependence, and islands.* Black-hole complementarity and related interior reconstructions preserve exterior unitarity without accessible paradoxes

for any single observer; island/replica-wormhole methods reproduce the unitary entropy curve while leaving the deep interior to full quantum gravity [342, 343, 231, 139, 337].

What observations can test this? We probe the exterior, not the singular interior: precision measurements of quasinormal-mode *ringdowns* in gravitational waves and *horizon-scale imaging* constrain deviations from the classical exterior metrics. Direct access to the singularity is screened by the horizon [344, 345, 346, 347, 348].

Comparing the Singularities

In classical general relativity a "singularity" is best understood as *geodesic incompleteness*—a boundary of the spacetime model where classical evolution cannot be extended—often, though not invariably, accompanied by the blow-up of curvature invariants. On this definition, the *Big Bang* of FLRW cosmology is a *past boundary* of the classical solution rather than a location "in" space; it marks the point where GR ceases to be predictive and a quantum-gravitational description must supply initial data or a replacement dynamics [240, 7, 8, 3, 144, 319].

By contrast, a *black hole singularity* is an *interior* end-state of classical collapse lying to the *future* of any infalling observer. The event horizon of a Schwarzschild black hole is regular (its coordinate pathology is removable), but the classical interior terminates at a spacelike $r = 0$ singularity reached in finite proper time; rotating (Kerr) solutions admit a ring singularity in their maximal analytic extensions. In realistic collapse, the smooth inner structure of ideal stationary solutions is expected to be unstable, and the conjecture of *weak cosmic censorship* asserts that any singularities formed are hidden behind horizons [7, 316, 318, 312].

A third possibility, the *Big Crunch*, is a *future* singularity in recollapsing cosmologies (e.g., closed models with suitable energy content). Present evidence—cosmic acceleration from Type Ia supernovae and precision CMB/large-scale-structure constraints—is consistent with a nearly constant dark-energy component ($w \approx -1$), which disfavors recollapse under current conditions, though alternative late-time dynamics could alter the ultimate fate [134, 135, 150, 144].

In short: singularities in GR signal the *limits of the classical theory.* The Big Bang is a past boundary of cosmological evolution; black-hole singularities are future interior endpoints cloaked by horizons; and a Big Crunch, while classically allowed, is observationally disfavored today. In each case, it is widely expected that a quantum-gravitational completion will resolve the classical pathology without sacrificing unitarity or causality [240, 7, 8, 3].

From Classical Singularities and Qubit to the Primordial Quantum Singularity. In classical general relativity, a "singularity" marks *geodesic incompleteness*—a boundary of the spacetime model rather than a place in space; the Big Bang of FLRW cosmology is precisely such a *past boundary* of the classical solution, while black-hole interiors terminate at future, horizon-cloaked endpoints [240, 7, 144, 319]. Within the IQR framework, this role of a singularity—as the *limit of a description*—is recast as a *design principle* for the elementary quantum unit itself. The conventional *qubit*, a state in a fixed two-dimensional Hilbert space, is too impoverished to seed spacetime and matter as emergent structures; it presupposes kinematics and an inner product rather than generating them.

The *Primordial Quantum Singularity (PQS)* is therefore introduced as the *replacement for the qubit*: an *infinitesimal quantum neighborhood* equipped with the relational operator

content needed to *generate* both kinematics and dynamics. A PQS is not a just a state vector or a tiny spatial region; it is a locally instantiated operator domain whose relational channels, evaluated in the global Quantum Holo-State $|\mathring{\mathbf{\Psi}}\rangle$, produce the emergent Hilbert structure and the chronogeometric expectations that we interpret as spacetime and fields. In this sense, the PQS is the *elementary grain of reality* and the *local generator* of physical law, unifying "information" and "substrate" in a single construct.

Viewed against the classical narrative, the PQS plays the role of a *point of origin*—not as a curvature blow-up, but as a *foundational origin of distinguishability* from which geometry, matter, and causal order crystallize. Where the classical Big Bang is a past boundary beyond which GR cannot be extended [240, 7], the PQS supplies the *pre-geometric* data and lawful channels from which that boundary and its subsequent FLRW evolution can be derived as an emergent, coarse-grained phase. Thus the transition from qubit to PQS mirrors the transition from classical singularities to an operator-level genesis: the former assumes a background; the latter *produces* one.

Each PQS embodies the complete dual field structure $\hat{\mathbf{\Omega}}_x = {}_{\hat{\Phi}_x}\mathring{\mathbf{\Lambda}}_{x,\hat{\Theta}_x}$, where $\hat{\mathbf{\Phi}}_x$ and $\hat{\mathbf{\Theta}}_x$ are operator-valued fields and $\mathring{\mathbf{\Lambda}}_x$ encodes the foundational Fourfold Dialectic Algebra of Symmetry. This is not a point in the classical geometric sense, nor merely an open region within pre-existing space, but an irreducible singular unit of quantum structure whose properties are delineated entirely through relational constructs. The defining characteristics of an PQS—including its effective spatio-temporal extent—emerge from state-dependent structures governed by the emergent distance operator $\hat{\mathbf{d}}(x, x')$ (Section 4.16) and emergent time $t_{\text{em}}^{(\mathring{\mathbf{\Psi}})}(x, x')$ (Section 4.12), all contextualized by the Quantum Holo-State $|\mathring{\mathbf{\Psi}}\rangle$ (Definition 1.27).

This represents a profound paradigm shift: observable phenomena such as macroscopic spacetime geometry, causal flow, and matter content arise only from the relational network between distinct PQSs, never from within a single, isolated one. The PQS thus marks the precise threshold between the unstructured nature of the foundational singularity and the structured nature of relational emergence—the quantum seed from which space, time, fields, and causality all emerge relationally.

Definition: The Primordial Quantum Singularity

Definition 1.25 (Primordial Quantum Singularity $\mathfrak{B}_{\vec{r}}$ and its Quantum Holo-State)**.** Let $\mathcal{C}_d \subset \mathbb{R}^d$ be a smooth, differentiable base manifold, interpreted as the *Foundational Singularity Continuum* whose points $\vec{r}$ are primordial loci, or seeds, denoted $\hat{\Omega}_{\vec{r}}$ (Definition 4.617). To each such locus $\vec{r} \in \mathcal{C}_d$, we assign a *Quantum Primordial Singularity* $\mathfrak{B}_{\vec{r}}$. The PQS is not just a "point", but is *defined by* the instantiation of the complete primordial duality structure at this locus:

$$\boxed{\mathfrak{B}_{\vec{r}} := \hat{\mathbf{\Omega}}_{\vec{r}} := {}_{\hat{\Phi}_{\vec{r}}}\mathring{\mathbf{\Lambda}}_{\vec{r},\hat{\Theta}_{\vec{r}}}} \tag{1.3}$$

where $\hat{\mathbf{\Phi}}_{\vec{r}}$, $\hat{\mathbf{\Theta}}_{\vec{r}}$, and $\mathring{\mathbf{\Lambda}}_{\vec{r}}$ are the local instantiations of the Gaiamiti field (Definition 6.118), Tantrakyma field (Definition 6.142), and the algebra of duality, respectively. This operator-valued entity $\mathfrak{B}_{\vec{r}}$, representing an elementary quantum singularity, is intrinsically and fundamentally described by the *Quantum Holo-State*, conceptually denoted $\mathring{\mathbf{\Psi}}$ and represented

by the state vector $|\mathring{\mathbf{\Psi}}\rangle$ (Definition 1.27), which encapsulates its complete quantum nature. Each PQS $\mathfrak{B}_{\vec{r}}$ is thus a fully structured, operator-valued singularity—a self-contained, dual quantum locus whose definitive quantum state is $|\mathring{\mathbf{\Psi}}\rangle$. The quantum singularity will be employed to replace the conventional Bloch-sphere qubit (which is a state vector in a fixed $\mathbb{C}^2$ space), with a dynamic, higher-order, field-embedded, and duality-governed object that serves as the source of emergent physical reality, and hence, the apex of causality (Section 5.4.6).

Motivations: The Emergent Quantum Configuration Space

The Quantum Holo-State $|\mathring{\mathbf{\Psi}}\rangle$ (Definition 1.27), realized through the GNS construction (Definition 1.34), generates the total emergent Hilbert space $\mathcal{H}_{\text{em}}$ from the algebra of observables. Yet this global construction leaves a fundamental question unanswered: how does this unified quantum state manifest the multiplicity of distinguishable entities we observe in physical reality? The answer lies in recognizing that while the state is universal, the operator structures it governs are inherently local—each Primordial Quantum Singularity $\mathfrak{B}_{\vec{r}}$ in the $\hat{\Omega}$-Continuum carries its own field operators $\mathring{\mathbf{\Lambda}}_{\hat{\Phi}_{\vec{r}},\hat{\Theta}_{\vec{r}}}$ (Definition 1.25). This tension between global state and local operators necessitates an intermediate structure: the quantum configuration space.

The emergence of distinguishable loci from the primordial continuum follows a principle of relational crystallization. Within the $\hat{\Omega}$-Continuum $\mathcal{C}_d$ (Definition 4.617), each point hosts an PQS—a domain where all relational measures vanish identically. These are regions of perfect quantum indistinguishability, where the operator-valued distances $\hat{\mathbf{d}}(x, x') = \mathbf{0}$ and emergent proper times $t_{\text{em}}^{(\mathring{\mathbf{\Psi}})}(x, x') = 0$ for all points within the same PQS (eq. (1.5)). Physical space emerges only when we identify which collections of points can be distinguished by quantum-relational measurements. This motivates the construction of the quantum configuration space $\mathcal{X}_q$ as a quotient of the continuum, where each point represents a relationally defined locus of distinguishability—an equivalence class of indistinguishable PQS constituents. This construction is rigorously developed in Section 4.19.3, with the emergence formally established in Proposition 4.625.

Once crystallized, $\mathcal{X}_q$ serves as the organizational backbone for the emergent Hilbert structure described in Section 4.1. The total emergent Hilbert space $\mathcal{H}_{\text{em}}$ naturally organizes itself as a bundle over this configuration space (Definition 4.239), with each fiber $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}$ (Definition 4.273) representing the internal modal structure of the PQS at each relationally distinguished locus x. The inner products within these fibers are not imposed axiomatically but emerge through the Fourfold Inter-Modal Correlator $\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})$ (Definition 4.9), evaluated in the Quantum Holo-State (Definition 4.228 and eq. (4.85)). This creates a self-consistent structure where the global state determines local metrics, while the configuration space provides the arena for organizing these local structures into a coherent whole. In this framework, $\mathcal{X}_q$ is goes beyond where physics happens—it is the *first layer of emergent structure*, the relational scaffold upon which both quantum mechanics and spacetime geometry are built.

1.3.1 Coordinate Equivalence of the Primordial Quantum Singularity

Let $\varphi : \mathcal{C}_d \to \mathcal{X}_q$ be the diffeomorphic map (or quotient map, if $\mathcal{X}_q$ emerges via identification of indistinguishable points in $\mathcal{C}_d$ as established in Section 4.19.3) from the base manifold $\mathcal{C}_d$ to the emergent *quantum configuration space* $\mathcal{X}_q$. For every $\vec{r} \in \mathcal{C}_d$, its image is $x := \varphi(\vec{r}) \in \mathcal{X}_q$. We can then index the PQS by points in $\mathcal{X}_q$:

$$\boxed{\mathfrak{B}_{\vec{r}} = \mathfrak{B}_x := {}_{\hat{\Phi}_x}\overset{\circ}{\mathbf{\Lambda}}_{x,\hat{\Theta}_x}} \tag{1.4}$$

This signifies that the same fundamental quantum structure $\mathfrak{B}$, whose intrinsic state is $|\overset{\circ}{\mathbf{\Psi}}\rangle$, can be viewed from a pre-geometric perspective (labeled by $\vec{r} \in \mathcal{C}_d$) or a post-geometric, relational perspective (labeled by $x \in \mathcal{X}_q$). The internal structure ${}_{\hat{\Phi}}\overset{\circ}{\mathbf{\Lambda}}_{\hat{\Theta}}$ (generic form) is instantiated at x through the local field operator structures $\hat{\Phi}_x, \hat{\Theta}_x, \overset{\circ}{\mathbf{\Lambda}}_x$.

Interpretational Shift and Emergent Hilbert Spaces: This formulation marks a revolutionary departure from conventional quantum theory's foundational assumptions. The Quantum Primordial Singularity $\mathfrak{B}_x$—an operator-valued entity whose quantum essence is captured by the Quantum Holo-State $|\overset{\circ}{\mathbf{\Psi}}\rangle$—transcends and supplants the abstract qubit as the elementary building block of quantum reality. Where the qubit represents merely a two-level system residing in a pre-defined mathematical space, the PQS embodies a complete inversion of this conceptual hierarchy.

The PQS represents a profound ontological shift: it is not a state vector dwelling within some pre-ordained Hilbert space, but rather a *localized field-theoretic configuration*—an operator-valued singularity whose very quantum nature *is* $|\overset{\circ}{\mathbf{\Psi}}\rangle$ and which carries intrinsic operator-algebraic structure. This reverses the traditional hierarchy where states exist in spaces; here, the state itself generates the space. The PQS is simultaneously the quantum entity and the source of the mathematical structures needed to describe it, embodying a self-referential completeness absent in conventional formulations.

This new paradigm re-envisions the *modal Hilbert spaces*, exemplified by the two-dimensional space $\mathcal{H}_2^{(\mathfrak{B}_x)}$ encoding the Symmetry/Antisymmetry duality inherent in $\overset{\circ}{\mathbf{\Lambda}}_x$ (Definition 4.617 for the foundational context and Definition 4.273 for the modal construction), as *representational approximations* of the PQS's internal duality modes, a departure from treating them as fundamental structures. Whether realized through the GNS construction applied to local operator algebras—with $|\overset{\circ}{\mathbf{\Psi}}\rangle$ serving as the cyclic vector for the state functional $\omega^{\overset{\circ}{\mathbf{\Psi}}}$—or as effective descriptions under specific physical conditions, this emergence (eq. (4.85)) exemplifies the new paradigm where Hilbert spaces themselves become emergent phenomena rather than fixed mathematical stages. The mathematical structures we use to describe quantum mechanics are thus revealed as secondary, derived from the more fundamental operator-theoretic reality of the PQS continuum.

1.3.2 Defining Properties of the Primordial Quantum Singularity

The Primordial Quantum Singularity $\mathfrak{B}_x$ embodies a profound principle of quantum indistinguishability that lies at the heart of emergent geometry. Within the IQR, each singularity represents not just a point but a fundamental domain of relational opacity—a locus where the very measures that distinguish one configuration from another vanish identically. This defining characteristic is captured through the formal condition that for all configurations $x' \in \mathcal{X}_q$ belonging to the singularity $\mathfrak{B}_x$, the emergent relational measures yield null values when evaluated in the context of the Quantum Holo-State $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$:

$$\boxed{\forall x' \in \mathfrak{B}_x : \quad \hat{\mathbf{d}}(x, x') = \mathbf{0} \quad \text{and} \quad t_{\text{em}}^{(\overset{\circ}{\boldsymbol{\Psi}})}(x, x') = 0.} \tag{1.5}$$

This twin condition reveals the deep unity between spatial and temporal indistinguishability at the quantum-singular scale. The vanishing of the operator-valued distance $\hat{\mathbf{d}}(x, x') = \mathbf{0}$ encompasses the simultaneous nullity of all three fundamental measures—correlational, vacuum-response, and spectral distances—signifying that no quantum-relational probe can differentiate between x and x'. The temporal counterpart, $t_{\text{em}}^{(\overset{\circ}{\boldsymbol{\Psi}})}(x, x') = 0$, emerges as a natural consequence, establishing that within the PQS, the flow of emergent proper time ceases to have operational meaning.

This principle of perfect quantum indistinguishability transforms our understanding of the fundamental scale. Within the conceptual boundary of a single $\mathfrak{B}_x$, the familiar everyday notions of "here" versus "there" and "now" versus "then" lose their operational significance (Section 4.16). The PQS thus represents a threshold phenomenon—a quantum domain where the relational structures that give rise to space, time, and causality have not yet crystallized from the pre-geometric substrate. It is precisely at the boundaries between distinct PQSs that distinguishability emerges, and with it, the entire apparatus of spacetime geometry unfolds from these infinitesimal seeds of relational structure.

Remark 1.26 (Relational Emergence from the Primordial Quantum Singularity $\mathfrak{B}_x$)**.** Consider a Primordial Quantum Singularity $\mathfrak{B}_x := {}_{\hat{\Phi}_x}\overset{\circ}{\boldsymbol{\Lambda}}_{x,\hat{\Theta}_x}$ (Definition 1.25) associated with configuration $x \in \mathcal{X}_q$ (Proposition 4.625). This entity embodies a profound principle: within its infinitesimal quantum domain, the relational measures that constitute observable physical reality have not yet emerged. For any configuration $x' \in \mathfrak{B}_x$, the fundamental relational structures—the operator-valued distance $\hat{\mathbf{d}}(x, x')$ (Section 4.16) and emergent proper time $t_{\text{em}}^{(\overset{\circ}{\boldsymbol{\Psi}})}(x, x')$ (Section 4.12)—yield null values relative to x (Definition 4.617).

This vanishing should not be misunderstood as "nothingness". Rather, $\mathfrak{B}_x$ represents a *domain of pure quantum potentiality*—a local quantum singularity where distinguishability has not yet crystallized, even though the complete dual field structure ${}_{\hat{\Phi}_x}\overset{\circ}{\boldsymbol{\Lambda}}_{x,\hat{\Theta}_x}$ remains fully present. Each PQS thus contains the potential for all physical phenomena through its operator-valued fields and duality algebra, yet this potential remains latent until brought into relation with other singularities. The quantum essence of this structure is encoded in the Quantum Holo-State $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$ (Definition 1.27), which provides the global coherence necessary for relational structures to emerge.

The framework's central insight is that observable reality—spacetime geometry, matter content, and dynamical evolution—emerges not from within these singular domains but through the relational fabric woven between them. Physical phenomena arise through the distinctions, correlations, and interactions established between distinct Primordial Quantum Singularities $\mathfrak{B}_x, \mathfrak{B}_{x'}, \ldots$ distributed across the configuration space $\mathcal{X}_q$. This represents a fundamental inversion of traditional physics: rather than objects existing in space and interacting through forces, the IQR posits that space, time, and forces themselves emerge from the coherent network of relations between these irreducible quantum nodes. Each singularity serves simultaneously as a boundary condition for emergence and a seed of potential structure, with reality crystallizing in the relational interstices between them.

1.4 The Quantum Holo-State

Definition 1.27 (The Quantum Holo-State $\mathring{\mathbf{\Psi}}$)**.** The *Quantum Holo-State*, denoted conceptually as $\mathring{\mathbf{\Psi}}$ and represented mathematically by the state vector $|\mathring{\mathbf{\Psi}}\rangle$, constitutes the singular quantum state that fundamentally characterizes every *Primordial Quantum Singularity* $\mathfrak{B}_x$ (Definition 1.25) in existence. Each PQS manifests as a localized expression of this universal state through its position-dependent operator structure $\mathfrak{B}_x \equiv \hat{\mathbf{\Omega}}_x := {}_{\hat{\Phi}_x}\mathring{\mathbf{\Lambda}}_{x,\hat{\Theta}_x}$, where the field operators $\hat{\Phi}_x$ and $\hat{\Theta}_x$ carry spatial indices while the underlying quantum state remains universal. This remarkable duality—universal state with localized operator manifestations—ensures that while all PQSs throughout the foundational $\hat{\Omega}$–Continuum (Definition 4.617) share the same intrinsic quantum essence, they maintain distinguishability through their position-dependent field structures.

The universality of $|\mathring{\mathbf{\Psi}}\rangle$ implies a profound unity: it simultaneously characterizes each individual singularity as a localized mode or aspect of itself, while also describing the coherent totality of the system they collectively form. This is not a contradiction but rather the essential feature that enables global quantum coherence while preserving local distinguishability—each PQS represents a particular "perspective" or "excitation" of the universal state, differentiated by its operator-valued coordinates in the emergent configuration space.

The state $|\mathring{\mathbf{\Psi}}\rangle$ transcends the conventional paradigm of states residing in pre-existing spaces; instead, it generates its own arena of existence through the total emergent Hilbert space $\mathcal{H}_{\text{em}}$ (Definition 4.287), actualized via the GNS construction (Definition 1.34). In a profound recursive foundation, the inner product structure of $\mathcal{H}_{\text{em}}$ itself emerges relationally from the very state it hosts, with $|\mathring{\mathbf{\Psi}}\rangle$ serving as both the architect and inhabitant of its mathematical home (as detailed in Definition 4.228 through eq. (4.85)).

Through its unique position as the quantum essence of every PQS, the Quantum Holo-State $|\mathring{\mathbf{\Psi}}\rangle$ bridges the microscopic and macroscopic realms—establishing both the local quantum mechanical framework and the global geometric structure. The following propositions formalize these foundational roles.

Proposition 1.28 (Role of $\mathring{\mathbf{\Psi}}$ in Defining Emergent Spacetime)**.** *The Quantum Holo-State* $|\mathring{\mathbf{\Psi}}\rangle$ *determines the emergent geometric and causal structures through the following relations:*

(i) The expectation-level metric tensor is given by $\bar{g}_{\mu\nu}(x) = \langle \overset{\circ}{\boldsymbol{\Psi}} | \hat{g}_{\mu\nu}(x) | \overset{\circ}{\boldsymbol{\Psi}} \rangle$ *(Definition 5.10), where* $\hat{g}_{\mu\nu}(x)$ *is the operator-valued metric tensor (Definition 5.1).*

(ii) The emergent proper time is defined as $t_{\text{em}}^{(\overset{\circ}{\boldsymbol{\Psi}})}(x, x')$ *(Definitions 5.17 and 5.22).*

(iii) For each PQS $\mathfrak{B}_x$ *(Definition 1.25), the relations* $\hat{\mathbf{d}}(x, x') = \mathbf{0}$ *and* $t_{\text{em}}^{(\overset{\circ}{\boldsymbol{\Psi}})}(x, x') = 0$ *hold for all* $x' \in \mathfrak{B}_x$ *(eq. (1.5)).*

Remark 1.29 (Physical Interpretation)**.** The Quantum Holo-State acts as the cosmic lens through which the operator-theoretic substrate projects itself into observable phenomena. This mathematical structure embodies three profound physical principles: (i) represents the birth of classical geometry from quantum expectation—the operator-valued metric is projected through $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$ into the smooth fabric of spacetime; (ii) captures the crystallization of temporal structure from operator-valued phase relationships; (iii) establishes each PQS as a zone of perfect quantum indistinguishability where the universe has not yet instantiated relational distinguishability.

Proposition 1.30 (Role of $\overset{\circ}{\boldsymbol{\Psi}}$ in Defining Local Quantum Mechanical Structure)**.** *The Quantum Holo-State* $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$ *determines the inner product structure of the emergent modal Hilbert spaces* $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}$ *(Definition 4.273) associated with each Primordial Quantum Singularity* $\mathfrak{B}_x$ *(Definition 1.25) through the relation:*

$$\langle \psi_1 | \psi_2 \rangle_{\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}} := \lim_{x' \to x} \langle \overset{\circ}{\boldsymbol{\Psi}} | \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2] | \overset{\circ}{\boldsymbol{\Psi}} \rangle \tag{1.6}$$

where $\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2]$ *is the Fourfold Inter-Modal Correlator of Primordial Quantum Singularities (Definitions 4.9 and 4.228 and eq. (4.85)).*

Remark 1.31 (Unification of Scales)**.** This proposition reveals how the Quantum Holo-State $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$ bridges the microscopic and macroscopic realms of the theory. While Proposition Proposition 1.28 established its role in determining the emergent spacetime manifold $\mathcal{M}_{\text{em}}^{(\overset{\circ}{\boldsymbol{\Psi}})}$ at the collective level, here we see its complementary function: defining the operational rules of quantum mechanics at the local level through the metric structure of individual modal Hilbert spaces.

Thus, $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$ embodies the ultimate synthesis—it serves as the defining quantum context for both the "arena" (the emergent spacetime arising from the collective of PQSs) and the "actors" (the internal quantum modes of each PQS and their dynamics). This dual role establishes the Quantum Holo-State as the singular foundation from which both geometry and quantum mechanics emerge as complementary aspects of a unified relational reality.

1.4.1 Gelfand-Naimark-Segal Construction of the Quantum Holo-State

The Algebra of Observables

Definition 1.32 (The Algebra of Observables $\mathcal{A}(\mathcal{C}_d)$)**.** The *Algebra of Observables*, denoted $\mathcal{A}(\mathcal{C}_d)$, is the complete, background-independent mathematical structure that encodes the

totality of relational observables accessible within the Primordial Quantum Singularity and the Fourfold Algebra ontology. It is formally defined as the C^*-algebra generated by the fundamental operator fields associated with each Primordial Quantum Singularity across the foundational $\hat{\Omega}$–Continuum (Definition 4.617).

Specifically, for each locus $\vec{r} \in \mathcal{C}_d$, the following operator fields generate the algebra:

1. *Geometric Generators:* The Gaiamiti Field operators $\{\hat{\mathbf{\Phi}}_{\vec{r}}\}$, which encode the potential for relational geometry.
2. *Matter-Content Generators:* The Tantrakyma Field operators $\{\hat{\mathbf{\Theta}}_{\vec{r}}\}$, encoding internal "flavor" and matter-like excitations.
3. *Symmetry Generators:* The Duality Algebra operators $\{\hat{\mathbf{\Lambda}}_{\vec{r}}\}$, representing the foundational symmetries at each locus.

The C^*-algebra $\mathcal{A}(\mathcal{C}_d)$ is then the smallest norm-closed, involutive algebra (under the C^*-norm topology) that contains all finite linear combinations, products, and adjoints of these generators. This closure ensures that the algebra is a complete and self-contained description of the system's potential dynamics and relational structure, prior to the specification of a state or the emergence of a Hilbert space.

Remark 1.33 (Physical Interpretation)**.** This algebraic formulation represents a profound reordering of physical ontology. In conventional physics, one typically begins with a background structure—such as a manifold or Hilbert space—upon which fields or operators are defined. The Primordial Quantum Singularity and the Fourfold Algebra framework inverts this order. It posits that the abstract algebra of what can be measured—and how observables relate—is more fundamental than any underlying space.

1. *The Language of Reality:* The algebra $\mathcal{A}(\mathcal{C}_d)$ constitutes the complete "language" of the universe—its grammar and vocabulary of physical relations. It contains every possible statement that can be made about the system's geometric and matter content.
2. *Pre-Geometric Nature:* Crucially, this algebra exists *prior* to any notion of distance, time, or localization. It is a purely relational construct. The "distance" between two observables is not spatial but algebraic, encoded in their commutators and algebraic dependencies, not in any background manifold.

By defining physical reality in terms of this abstract algebra, the framework achieves fundamental background independence. The state functional $\omega^{\mathring{\mathbf{\Psi}}}$ then acts upon this algebra to select a single, dynamically coherent instantiation of physical reality, and the GNS construction provides the emergent stage—the Hilbert space—upon which this reality unfolds.

The Gelfand-Naimark-Segal Construction

Definition 1.34 (Gelfand–Naimark–Segal Construction of the Quantum Holo-State)**.** Let $\mathcal{A}$ be the (unital) C^*-algebra of emergent observables associated to the Quantum Holo-State (Section 1.4). Let $\omega^{\mathring{\mathbf{\Psi}}} : \mathcal{A} \to \mathbb{C}$ be the state functional singled out by the Holo-State, assumed

linear, positive, and normalized ($\omega^{\overset{\circ}{\mathbf{\Psi}}}(\mathbb{I}) = 1$). The GNS construction produces a cyclic $*$-representation

$$\Big(\pi_{\overset{\circ}{\mathbf{\Psi}}},\ \mathcal{H}_{\overset{\circ}{\mathbf{\Psi}}},\ |\Omega_{\omega^{\overset{\circ}{\mathbf{\Psi}}}}\rangle\Big),$$

which we also denote by $\mathcal{H}_{\text{em}} := \mathcal{H}_{\overset{\circ}{\mathbf{\Psi}}}$ (the *emergent* Hilbert space).

Canonical steps. Writing $\omega := \omega^{\overset{\circ}{\mathbf{\Psi}}}$, define:

1. **Null space.** $\mathcal{N}_\omega := \{A \in \mathcal{A} :\ \omega(A^\dagger A) = 0\}$ (a left ideal in $\mathcal{A}$).
2. **Pre-Hilbert space.** On the quotient $\mathcal{A}/\mathcal{N}_\omega$, set
$$\langle [A], [B] \rangle_\omega := \omega(A^\dagger B), \qquad [A] \in \mathcal{A}/\mathcal{N}_\omega,$$
and let $\mathcal{H}_\omega$ be its completion.
3. **Representation.** $\pi_\omega : \mathcal{A} \to \mathcal{B}(\mathcal{H}_\omega)$ by
$$\pi_\omega(A)[B] := [AB], \qquad A, B \in \mathcal{A},$$
which is a bounded $*$-representation.
4. **Cyclic vector.** $|\Omega_\omega\rangle := [\mathbb{I}] \in \mathcal{H}_\omega$; it is cyclic since $\overline{\pi_\omega(\mathcal{A})|\Omega_\omega\rangle} = \mathcal{H}_\omega$.

We *identify*

$$\mathcal{H}_{\overset{\circ}{\mathbf{\Psi}}} \equiv \mathcal{H}_\omega \equiv \mathcal{H}_{\text{em}}, \quad \pi_{\overset{\circ}{\mathbf{\Psi}}} \equiv \pi_\omega, \quad |\Omega_{\omega^{\overset{\circ}{\mathbf{\Psi}}}}\rangle \equiv |\Omega_\omega\rangle,$$

and regard $|\overset{\circ}{\mathbf{\Psi}}\rangle$ (the ontological Holo-State symbol) as represented by the *cyclic vector* $|\Omega_{\omega^{\overset{\circ}{\mathbf{\Psi}}}}\rangle$ in $\mathcal{H}_{\overset{\circ}{\mathbf{\Psi}}}$.

State recovery and Born-valued truth. For all $A \in \mathcal{A}$,

$$\omega^{\overset{\circ}{\mathbf{\Psi}}}(A) = \langle \Omega_{\omega^{\overset{\circ}{\mathbf{\Psi}}}} |\, \pi_{\overset{\circ}{\mathbf{\Psi}}}(A)\, |\Omega_{\omega^{\overset{\circ}{\mathbf{\Psi}}}}\rangle.$$

In particular, for any *effect* $E \in \mathcal{A}$ (with $0 \le E \le \mathbb{I}$), the Born-valued dialectic truth (Definition 1.18) is obtained either as an expectation in the cyclic GNS state,

$$v(E) \ = \ \omega^{\overset{\circ}{\mathbf{\Psi}}}(E) \ = \ \langle \Omega_{\omega^{\overset{\circ}{\mathbf{\Psi}}}} |\, \pi_{\overset{\circ}{\mathbf{\Psi}}}(E)\, |\Omega_{\omega^{\overset{\circ}{\mathbf{\Psi}}}}\rangle,$$

or, more generally, for any normal preparation ρ on $\mathcal{H}_{\text{em}}$,

$$v(\rho; E) \ = \ \mathrm{Tr}\Big(\rho\, \pi_{\overset{\circ}{\mathbf{\Psi}}}(E)\Big).$$

Compatibility with the Fourfold Dialectic Algebra. Define the set of corner effects (Definitions 1.16 and 1.18)

$$\mathcal{E}_{\text{corner}} \ := \ \{\, P_T,\ P_A,\ P_{\text{para}},\ E_{\text{Holo}}^{(\mathbf{a}\to\mathbf{b})} \,\}.$$

Their GNS images act on the emergent Hilbert space:

$$\pi_{\overset{\circ}{\mathbf{\Psi}}}\Big[\mathcal{E}_{\text{corner}}\Big] = \Big\{\, \pi_{\overset{\circ}{\mathbf{\Psi}}}(P_T),\ \pi_{\overset{\circ}{\mathbf{\Psi}}}(P_A),\ \pi_{\overset{\circ}{\mathbf{\Psi}}}(P_{\text{para}}),\ \pi_{\overset{\circ}{\mathbf{\Psi}}}(E_{\text{Holo}}^{(\mathbf{a}\to\mathbf{b})}) \,\Big\} \ \subset\ \mathcal{B}(\mathcal{H}_{\text{em}}).$$

Consequently, all Born-valued valuations defined at the logical layer are realized as standard quantum expectations on $\mathcal{H}_{\text{em}}$.

Uniqueness. The GNS triple is unique up to unitary equivalence: if $(\pi', \mathcal{H}', |\Omega'\rangle)$ is another cyclic representation with $\omega^{\mathring{\mathbf{\Psi}}}(A) = \langle\Omega'|\pi'(A)|\Omega'\rangle$ for all $A \in \mathcal{A}$, then there exists a unitary $U : \mathcal{H}_{\mathring{\mathbf{\Psi}}} \to \mathcal{H}'$ with $U\,\pi_{\mathring{\mathbf{\Psi}}}(A) = \pi'(A)\,U$ and $U|\Omega_{\omega^{\mathring{\mathbf{\Psi}}}}\rangle = |\Omega'\rangle$.

Remark 1.35 (Nonunital algebras)**.** If $\mathcal{A}$ is nonunital, replace $\mathbb{I}$ with the unit of the minimal unitization $\tilde{\mathcal{A}}$ in the quotient step; the construction and identifications above go through unchanged.

Remark 1.36 (Self-Generating Structure)**.** The GNS construction reveals a striking feature of the Quantum Holo-State: it *generates its own arena of existence*. The state functional $\omega^{\mathring{\mathbf{\Psi}}}$ fixes not only expectation values but the very Hilbert space $\mathcal{H}_{\mathring{\mathbf{\Psi}}}$ (also denoted $\mathcal{H}_{\text{em}}$) and its inner product. Explicitly, with the cyclic vector $|\Omega_{\omega^{\mathring{\mathbf{\Psi}}}}\rangle$ and representation $\pi_{\mathring{\mathbf{\Psi}}}$ produced by GNS, the inner product is defined by

$$\Big\langle \pi_{\mathring{\mathbf{\Psi}}}(A)\,|\Omega_{\omega^{\mathring{\mathbf{\Psi}}}}\rangle,\ \pi_{\mathring{\mathbf{\Psi}}}(B)\,|\Omega_{\omega^{\mathring{\mathbf{\Psi}}}}\rangle \Big\rangle_{\mathcal{H}_{\mathring{\mathbf{\Psi}}}} \;=\; \omega^{\mathring{\mathbf{\Psi}}}(A^\dagger B)\,, \qquad A, B \in \mathcal{A}.$$

Thus the ontological symbol $|\mathring{\mathbf{\Psi}}\rangle$ is *represented* by the cyclic vector $|\Omega_{\omega^{\mathring{\mathbf{\Psi}}}}\rangle$; the state functions simultaneously as the *seed* and the *standard of comparison* for the entire quantum–relational edifice—a precise embodiment of "the state defines the space that contains it."

Remark 1.37 (Physical Interpretation)**.** This construction is the mathematical expression of *background independence*. Where conventional quantum mechanics begins with a fixed Hilbert space—an immutable stage on which states live—our framework reverses the order: the foundational object is the operator algebra $\mathcal{A}(\mathcal{C}_d)$ encoding physical relations and observables, before any state space has crystallized. The state functional $\omega^{\mathring{\mathbf{\Psi}}}$ acts as a *selection principle* that carves out one coherent reality from the algebra's possibilities: by assigning expectations to all observables, it determines not just a state *within* a space, but the *space itself*. GNS is the engine that turns this abstract selection into concrete form, culminating in the emergent Hilbert space $\mathcal{H}_{\text{em}} = \mathcal{H}_{\mathring{\mathbf{\Psi}}}$—a bespoke arena sculpted by, and for, the state.

The identification $|\mathring{\mathbf{\Psi}}\rangle \equiv |\Omega_{\omega^{\mathring{\mathbf{\Psi}}}}\rangle$ therefore has genuine ontological bite: the universe's quantum state is not merely an inhabitant of a pre-given space; it is the *architect* of its own domain. When $\omega^{\mathring{\mathbf{\Psi}}}$ is extremal (i.e., a pure state on the C^*-algebra), the associated GNS representation is irreducible, capturing a cosmos governed by a single, indivisible quantum coherence rather than a mixture of alternatives. In this view, the quantum state and its mathematical arena emerge as inseparable aspects of a unified, self-determining whole—with Born-valued truth $\text{Tr}(\rho\,E)$ realized as ordinary expectations in the very space that the state brings into being.

1.4.2 The Fourfold Dialectic Algebra and the GNS Construction

The relationship between the *Fourfold Dialectic Algebra of Symmetry* (Definition 1.16) and the *GNS Construction of the Quantum Holo-State* (Sections 1.4 and 1.4.1) is best read

Bridge: Fourfold Dialectic Algebra of Symmetry (Definition 1.16)

Corner (logic ↔ symmetry)	Ontological rep.	Quantum effect / projector	Born-valued truth	Measurement protocol
Thesis ↔ Symmetry	$\|\mathring{\mathbf{0}}\rangle$	$P_T = \|e_1\rangle\langle e_1\|$	$v_{\text{Thesis}}(\rho) = \text{Tr}(P_T\rho)$	Measure Σ_z on $\mathcal{H}_{(0,1)}$ basis $\{\|e_1\rangle, \|e_2\rangle\}$
Antithesis ↔ Antisymmetry	$\|\mathring{\mathbf{1}}\rangle$	$P_A = \|e_2\rangle\langle e_2\|$	$v_{\text{Antithesis}}(\rho) = \text{Tr}(P_A\rho)$	Measure Σ_z on $\mathcal{H}_{(0,1)}$ basis $\{\|e_1\rangle, \|e_2\rangle\}$
Synthesis ↔ Parasymmetry	$\frac{1}{\sqrt{2}}(\|\mathring{\mathbf{0}}\rangle + \|\mathring{\mathbf{1}}\rangle)$	P_{para}	$v_{\text{Synthesis}}(\rho) = \text{Tr}(P_{\text{para}}\rho)$	Measure Σ_x (para-axis) on $\mathcal{H}_{(0,1)}$
Holothesis ↔ Holosymmetry	$\mathring{\mathbf{0}}\mathring{\mathbf{\Lambda}}\mathring{\mathbf{1}}$	$E_{\text{Holo}}^{(\mathbf{a}\to\mathbf{b})}$	$v_{\text{Holothesis}}(\rho) = \text{Tr}\big(E_{\text{Holo}}^{(\mathbf{a}\to\mathbf{b})}\rho\big)$	Control interferometer: prep $\|\mathbf{a}\rangle_c$, read $\|\mathbf{b}\rangle_c$

GNS / Emergent layer (Definition 1.34): $\mathcal{A}(\mathcal{C}_d)$ with state $\omega^{\mathring{\mathbf{\Psi}}}$ yields the cyclic triple $(\pi_{\mathring{\mathbf{\Psi}}}, \mathcal{H}_{\mathring{\mathbf{\Psi}}} = \mathcal{H}_{\text{em}}, |\Omega_{\omega^{\mathring{\mathbf{\Psi}}}}\rangle)$,

so that for any effect E: $\omega^{\mathring{\mathbf{\Psi}}}(E) = \langle\Omega_{\omega^{\mathring{\mathbf{\Psi}}}}|\pi_{\mathring{\mathbf{\Psi}}}(E)|\Omega_{\omega^{\mathring{\mathbf{\Psi}}}}\rangle$, and for any preparation ρ on $\mathcal{H}_{\text{em}}$: $v(\rho; E) = \text{Tr}\big(\rho\,\pi_{\mathring{\mathbf{\Psi}}}(E)\big)$ (Definition 1.18).

Figure 1.2: From logic to lab: each corner's ontological label → realized effect → Born truth → measurement, grounded in the GNS/emergent layer.

as that between an axiomatic *blueprint* and its *actualization*. They are two facets of a single emergence process: the Dialectic Algebra specifies, *a priori*, the admissible modes of assertion—**Symmetry** (Thesis), **Antisymmetry** (Antithesis), **Parasymmetry** (Synthesis), **Holosymmetry** (Holothesis)—while the GNS mechanism turns these logical forms into concrete quantum tests and expectations inside an emergent Hilbert arena. The algebra of observables $\mathcal{A}(\mathcal{C}_d)$ supplies the complete operational vocabulary; the Quantum Holo-State $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$ (via its state functional $\omega^{\overset{\circ}{\boldsymbol{\Psi}}}$) selects a coherent world; and the GNS construction then *builds* the arena $\mathcal{H}_{\overset{\circ}{\boldsymbol{\Psi}}}$ (henceforth also denoted $\mathcal{H}_{\text{em}}$) in which the universe's quantum history unfolds. In short: the Dialectic provides the rules of construction, and GNS is the act of construction according to those rules.

The Hierarchical Architecture: From Logic to Reality The emergence proceeds through three tightly coupled levels.

1. Ontological Level: The Dialectic as Generative Logic Rooted in the *Principle of Duality*, the Fourfold Dialectic Algebra (Definition 1.16) is the system's generative logic rather than a descriptive overlay. From the primal polarity—**Symmetry** (Thesis) vs. **Antisymmetry** (Antithesis)—arises **Parasymmetry** (Synthesis) as their coherent organization, and **Holosymmetry** (Holothesis) as the meta-level that adjudicates or superposes entire evolutionary *histories* (Section 2.3). This fourfold structure is intrinsically present at every *Primordial Quantum Singularity* $\mathfrak{B}_x$ (Definition 1.25), providing the ontological anchors (ringed kets/meta-symbols) and the logical roles from which all subsequent physics—observables, dynamics, and patterns of entanglement—must emerge. Importantly, these ontological labels are *not* ordinary Hilbert vectors; they acquire quantum meaning only through a chosen realization (cf. Remark 1.22).

2. Algebraic Level: The Manifest Realization The Dialectic's "generative grammar" becomes a full mathematical language at the level of the *Algebra of Observables* $\mathcal{A}(\mathcal{C}_d)$ (Definition 1.32). Here the symmetry generators $\hat{\boldsymbol{\Lambda}}_{\vec{r}}$, together with the Gaiamiti and Tantrakyma operator fields (Definitions 6.118 and 6.142), provide the substrate on which the dialectical modes act. The corner propositions are represented by concrete *effects* in $\mathcal{A}(\mathcal{C}_d)$—e.g. P_T, P_A in the distinguished $(0,1)$-sector, the parasymmetric pair $P_{\text{para}}, P_{\text{antipara}}$, and the holosymmetric effect $E_{\text{Holo}}^{(\mathbf{a}\nrightarrow\mathbf{b})}$—so the Fourfold Dialectic is already *operational* at the algebraic level. Born-valued semantics (Definition 1.18) then assign to each corner a truth valuation $v(\rho; P) = \text{Tr}(P\rho)$, making "degree of warrant" a measurable probability.

3. Physical Emergence via GNS Construction Physical reality is obtained when the Holo-State $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$ acts as a selection principle through its state functional $\omega^{\overset{\circ}{\boldsymbol{\Psi}}}$ on $\mathcal{A}(\mathcal{C}_d)$, and the *Gelfand–Naimark–Segal* construction (Definition 1.34) produces the cyclic triple

$$\left(\pi_{\overset{\circ}{\boldsymbol{\Psi}}},\ \mathcal{H}_{\overset{\circ}{\boldsymbol{\Psi}}} = \mathcal{H}_{\text{em}},\ |\Omega_{\omega^{\overset{\circ}{\boldsymbol{\Psi}}}}\rangle\right).$$

In this representation, every algebraic effect E acts as the operator $\pi_{\overset{\circ}{\Psi}}(E)$ on $\mathcal{H}_{\text{em}}$, and the dialectic truth becomes an ordinary expectation:

$$\omega^{\overset{\circ}{\Psi}}(E) = \langle \Omega_{\omega^{\overset{\circ}{\Psi}}} | \pi_{\overset{\circ}{\Psi}}(E) | \Omega_{\omega^{\overset{\circ}{\Psi}}} \rangle, \qquad v(\rho; E) = \mathrm{Tr}\Big(\rho\, \pi_{\overset{\circ}{\Psi}}(E)\Big),$$

for the cyclic vector or any preparation ρ on $\mathcal{H}_{\text{em}}$. Thus the logical fourfold becomes a quartet of concrete, repeatable tests on the emergent space. When $\omega^{\overset{\circ}{\Psi}}$ is extremal, the GNS representation is irreducible, capturing a single, indivisible quantum coherence rather than a mixture.

The Essential Distinction: Grammar versus Language A linguistic analogy clarifies the roles. The *Fourfold Dialectic Algebra* (Definition 1.16) is the *generative grammar*: an *a priori* rulebook that determines which modal structures are syntactically admissible and why they must appear at all. The *Algebra of Observables* $\mathcal{A}(\mathcal{C}_d)$ is the *language*: the full lexicon of operators and relations (from the Gaiamiti/Tantrakyma fields to parasymmetric and holosymmetric composites) that say *what* can be physically stated and transformed. Finally, the *GNS construction* is the act of *uttering* a world: it gives these grammatical sentences a stage and a voice—$\mathcal{H}_{\text{em}}$ and $\pi_{\overset{\circ}{\Psi}}$—so that the four corners become measurable statements with Born-valued truth. In this sense, logic (grammar), algebra (language), and representation (utterance) are not separate add-ons but successive refinements of one unified emergence pipeline from ontology to experiment.

The Fundamental Trinity The complete picture resolves into a *trinity* of mutually entailed roles—**why**, **what**, and **how**—that together constitute reality in the IQR.

1. **Dialectic (Why).** The Fourfold Dialectic Algebra (Definition 1.16) states *why* the universe must exhibit its modal structure. Rooted in the Principle of Duality, it fixes the admissible corners (Thesis/Antithesis/Synthesis/Holothesis) and their interrelations as an *a priori* generative logic intrinsic to each PQS. These ontological labels are not yet Hilbert vectors; they are the blueprint from which realizations are drawn.

2. **Algebra (What).** The Algebra of Observables $\mathcal{A}(\mathcal{C}_d)$ (Definition 1.32) specifies *what* exists in precise mathematical form. It supplies the complete language of operators (including symmetry generators and the Gaiamiti/Tantrakyma fields) and instantiates the corners as effects $P_T, P_A, P_{\text{para}}, E_{\text{Holo}}^{(\mathbf{a}\to\mathbf{b})}$ with Born-valued truth $v(\rho; P) = \mathrm{Tr}(P\rho)$ (Definition 1.18). At this level the fourfold becomes *operational*: each corner corresponds to a concrete test.

3. **Holo-State via GNS (How).** The Quantum Holo-State $|\overset{\circ}{\Psi}\rangle$ (through its functional $\omega^{\overset{\circ}{\Psi}}$) determines *how* one coherent world is realized from the algebra's possibilities. The GNS construction (Definition 1.34) builds the emergent arena $\mathcal{H}_{\overset{\circ}{\Psi}} = \mathcal{H}_{\text{em}}$ with representation $\pi_{\overset{\circ}{\Psi}}$ and cyclic vector $|\Omega_{\omega^{\overset{\circ}{\Psi}}}\rangle$, so that all valuations become standard quantum expectations on the emergent space. In the extremal case, the representation is irreducible, reflecting a single, indivisible quantum coherence.

Taken together, *why* (Dialectic), *what* (Algebra), and *how* (Holo-State → GNS) form an unbroken pipeline from ontological principle to laboratory protocol. The Fourfold Dialectic and the Algebra of Observables are synchronized facets of the same entity, while the GNS construction provides its physical voice. In this sense, the universe is captured as a *self-generating* quantum state whose internal, dialectical logic projects itself—via algebra and representation—into the chronogeometric fabric we observe.

Level	**Structure**	**Function & Nature**
Ontological	Fourfold Dialectic Algebra (Definition 1.16)	Generative logic intrinsic to each PQS: specifies the four admissible modes (Thesis/Symmetry, Antithesis/Antisymmetry, Synthesis/Parasymmetry, Holothesis/Holosymmetry) and their roles as *corners*. Provides the *a priori* blueprint; ontological labels are pre-geometric (cf. Remark 1.22).
Algebraic	$\mathcal{A}(\mathcal{C}_d)$ (Definition 1.32)	Concrete C^*-algebra of observables/relations (including symmetry generators and Gaiamiti/Tantrakyma fields). Realizes the corners as effects $P_T, P_A, P_{\text{para}}, E_{\text{Holo}}^{(\mathbf{a}\to\mathbf{b})}$ and endows them with Born-valued semantics $v(\rho; P) = \text{Tr}(P\rho)$ (Definition 1.18).
Physical	$(\pi_{\overset{\circ}{\mathbf{\Psi}}},\ \mathcal{H}_{\overset{\circ}{\mathbf{\Psi}}} = \mathcal{H}_{\text{em}},\ \lvert\Omega_{\overset{\circ}{\omega^{\mathbf{\Psi}}}}\rangle)$ via GNS (Definition 1.34)	Emergent Hilbert arena and representation selected by the Quantum Holo-State. All corner valuations become ordinary expectations: $\overset{\circ}{\omega^{\mathbf{\Psi}}}(E) = \langle\Omega\lvert\pi_{\overset{\circ}{\mathbf{\Psi}}}(E)\rvert\Omega\rangle$, and more generally $v(\rho; E) = \text{Tr}(\rho\,\pi_{\overset{\circ}{\mathbf{\Psi}}}(E))$.

Table 1.1: Three-tiered hierarchy from generative logic to algebraic language to emergent physical representation in the IQR.

Chapter 2

The Fourfold Symmetry

2.1 Quantum Realizations of the Fourfold Symmetry

2.1.1 Realization of The Ontological Duality States

Definition 2.1 (Realization of the Ontological Duality Modes in $\mathbb{C}^4$)**.** Let $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)} = \mathbb{C}^4$ denote the modal Hilbert space associated to a Primordial Quantum Singularity $\mathfrak{B}_x$. The fundamental ontological modes of the Fourfold Dialectic Algebra—Symmetry and Antisymmetry—are realized within this space as follows:

1. **Symmetry Mode (Thesis):**

$$|\mathring{\mathbf{0}}\rangle := |e_1\rangle := \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} \in \mathbb{C}^4, \tag{2.1}$$

2. **Antisymmetry Mode (Antithesis):**

$$|\mathring{\mathbf{1}}\rangle := |e_2\rangle := \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix} \in \mathbb{C}^4, \tag{2.2}$$

These states span the fundamental $(0,1)$-subspace:

$$\mathcal{H}_{(0,1)} := \text{span}\,\{|e_1\rangle, |e_2\rangle\} \cong \mathbb{C}^2 \subset \mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)} \tag{2.3}$$

3. **Generic Modal States:** Any modal state constrained to this sector at position x or

x' takes the form:

$$|\psi_1\rangle_x = c_{10}|\mathring{\mathbf{0}}\rangle_x + c_{11}|\mathring{\mathbf{1}}\rangle_x = \begin{pmatrix} c_{10} \\ c_{11} \\ 0 \\ 0 \end{pmatrix}_x ,$$

$$|\psi_2\rangle_{x'} = c_{20}|\mathring{\mathbf{0}}\rangle_{x'} + c_{21}|\mathring{\mathbf{1}}\rangle_{x'} = \begin{pmatrix} c_{20} \\ c_{21} \\ 0 \\ 0 \end{pmatrix}_{x'} ,$$

4. **Interpretation:** This realization faithfully embeds the ontological duality states within a conventional quantum mechanical Hilbert space, while preserving their dialectical meaning. All inter-modal correlator constructions restricted to the $(0,1)$-sector are now understood to act within $\mathbb{C}^2 \subset \mathbb{C}^4$.

Logical projectors and valuation Define $P_T := |e_1\rangle\langle e_1|$ and $P_A := |e_2\rangle\langle e_2|$ on $\mathcal{H}_{(0,1)}$. By Definition 1.18, $v_{\text{Thesis}}(\rho) = \text{Tr}(P_T\rho)$ and $v_{\text{Antithesis}}(\rho) = \text{Tr}(P_A\rho)$.

Quantum Realizations of the $(0,1)$-Sector Circuits

Definition 2.2 (Two-qubit embedding of the $(0,1)$-sector)**.** Let $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)} = \mathbb{C}^4$ with computational basis $\{\,|e_1\rangle, |e_2\rangle, |e_3\rangle, |e_4\rangle\,\}$, and fix an identification with two qubits (q_1, q_0) via

$$|e_1\rangle \longmapsto |0\rangle_{q_1}|0\rangle_{q_0}, \qquad |e_2\rangle \longmapsto |0\rangle_{q_1}|1\rangle_{q_0}, \qquad |e_3\rangle \longmapsto |1\rangle_{q_1}|0\rangle_{q_0}, \qquad |e_4\rangle \longmapsto |1\rangle_{q_1}|1\rangle_{q_0}.$$

The ontological duality modes from Definition 2.1 satisfy

$$|\mathring{\mathbf{0}}\rangle \equiv |e_1\rangle \equiv |0\rangle_{q_1}|0\rangle_{q_0}, \qquad |\mathring{\mathbf{1}}\rangle \equiv |e_2\rangle \equiv |0\rangle_{q_1}|1\rangle_{q_0},$$

so that the $(0,1)$-sector is the subspace with $q_1 = |0\rangle$ and logical duality bit encoded in q_0.

Definition 2.3 (Circuit for the Symmetry mode (Thesis))**.** The Symmetry mode is $|\mathring{\mathbf{0}}\rangle \equiv |e_1\rangle$ (eq. (2.1)). Define the unitary

$$U_{\mathsf{Th}} := \mathbb{I}_{q_1} \otimes \mathbb{I}_{q_0}, \tag{2.4}$$

acting on the embedded $(0,1)$-sector. Starting from the initialized state $|0\rangle_{q_1}|0\rangle_{q_0}$, the circuit is the identity:

$$\begin{array}{l} |0\rangle_{q_1} \text{ ——} \\ |0\rangle_{q_0} \text{ ——} \end{array} \qquad \Longrightarrow \qquad |0\rangle_{q_1}|0\rangle_{q_0} \equiv |e_1\rangle \equiv |\mathring{\mathbf{0}}\rangle.$$

Definition 2.4 (Circuit for the Antisymmetry mode (Antithesis))**.** The Antisymmetry mode is $|\mathring{\mathbf{1}}\rangle \equiv |e_2\rangle$ (eq. (2.2)). Define the unitary that flips only the duality bit:

$$U_{\mathsf{AntiTh}} := \mathbb{I}_{q_1} \otimes X_{q_0}. \tag{2.5}$$

From the same initialization $|0\rangle_{q_1}|0\rangle_{q_0}$, the circuit is

$$|0\rangle_{q_1} \quad \text{———}$$
$$|0\rangle_{q_0} \quad \text{—}[X]\text{—} \qquad \Longrightarrow \qquad |0\rangle_{q_1}|1\rangle_{q_0} \equiv |e_2\rangle \equiv |\mathring{\mathbf{1}}\rangle.$$

Definition 2.5 (Projective tests matching $(0,1)$-sector effects)**.** Restricting to $q_1 = |0\rangle$, the Thesis/Antithesis effects (P_T, P_A) act as computational-Z tests on q_0:

$$P_T = |00\rangle\langle 00|\,, \qquad P_A = |01\rangle\langle 01|\,, \tag{2.6}$$

with standard Z-basis measurement on q_0 distinguishing the two modes.

Definition 2.6 (Synthesis of a generic $(0,1)$-state)**.** Any state $c_{10}|\mathring{\mathbf{0}}\rangle + c_{11}|\mathring{\mathbf{1}}\rangle$ is prepared by single-qubit rotations on q_0 while holding $q_1 = |0\rangle$:

$$|\psi\rangle \;=\; \Big(R_z(\phi)\,R_y(2\theta)\,|0\rangle\Big)_{q_0} \otimes |0\rangle_{q_1}\,, \quad \theta = \arccos|c_{10}|, \;\; \phi = \arg(c_{11}) - \arg(c_{10}). \tag{2.7}$$

A concrete circuit is

$$|0\rangle_{q_1} \quad \text{———————}$$
$$|0\rangle_{q_0} \quad \text{—}[R_y(2\theta)]\text{—}[R_z(\phi)]\text{—}$$

which reduces to Definitions 2.3 and 2.4 for $(\theta,\phi) = (0,0)$ and $(\frac{\pi}{2},0)$, respectively.

2.1.2 Local Involutive Duality Operators

Definition 2.7 (Local Involutive Duality Operator $D^{(\mathfrak{B}_x)}$)**.** Let $\mathfrak{B}_x$ be a Primordial Quantum Singularity located at point $x \in \mathcal{X}_q$, with associated modal Hilbert space $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}$. The *local involutive duality operator*

$$D^{(\mathfrak{B}_x)} : \mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)} \to \mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}$$

is the linear operator that acts nontrivially on the Symmetry–Antisymmetry sector as:

$$\boxed{D^{(\mathfrak{B}_x)} := |\mathring{\mathbf{0}}\rangle_x\langle\mathring{\mathbf{1}}|_x + |\mathring{\mathbf{1}}\rangle_x\langle\mathring{\mathbf{0}}|_x,} \tag{2.8}$$

and acts trivially (as the identity) on the orthogonal Parasymmetry and Holosymmetry subspaces.

This operator satisfies the following properties:

1. **Symmetry–Antisymmetry Interchange:**

$$D^{(\mathfrak{B}_x)}|\mathring{\mathbf{0}}\rangle_x = |\mathring{\mathbf{1}}\rangle_x, \qquad D^{(\mathfrak{B}_x)}|\mathring{\mathbf{1}}\rangle_x = |\mathring{\mathbf{0}}\rangle_x.$$

2. **Matrix Representation:**
   When restricted to the subspace span$\{|\mathring{\mathbf{0}}\rangle_x, |\mathring{\mathbf{1}}\rangle_x\}$, the operator is represented by the Pauli-X matrix:
$$D^{(\mathfrak{B}_x)} \cong \sigma_X = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

3. **Involution and Hermiticity:**
$$(D^{(\mathfrak{B}_x)})^\dagger = D^{(\mathfrak{B}_x)}, \qquad (D^{(\mathfrak{B}_x)})^2 = \mathbb{I}_{\text{mod}}^{(\mathfrak{B}_x)},$$
   where $\mathbb{I}_{\text{mod}}^{(\mathfrak{B}_x)}$ denotes the identity operator on $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}$.

4. **Interpretation:**
   $D^{(\mathfrak{B}_x)}$ enacts a fundamental involutive transformation of internal quantum logic: it toggles the local dialectical configuration between Symmetry and Antisymmetry modes while preserving the higher-order Parasymmetric and Holosymmetric structures. It is the simplest nontrivial instance of a duality generator in the local modal algebra.

This operator provides the template for a hierarchy of generalized duality transformations—including parasymmetric reflection operators and holosymmetric coherence maps—that together generate the local action of the Fourfold Dialectic Algebra on modal state space. The full structure of this action will be developed in subsequent sections.

Quantum Circuit for the Local Involutive Duality Operator

Definition 2.8 (Embedding of $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)} = \mathbb{C}^4$ into two qubits)**.** Identify the modal basis with two qubits (q_1, q_0) by
$$|e_1\rangle \mapsto |0\rangle_{q_1} |0\rangle_{q_0}, \quad |e_2\rangle \mapsto |0\rangle_{q_1} |1\rangle_{q_0}, \quad |e_3\rangle \mapsto |1\rangle_{q_1} |0\rangle_{q_0}, \quad |e_4\rangle \mapsto |1\rangle_{q_1} |1\rangle_{q_0}.$$
Then the $(0,1)$-sector is span$\{|00\rangle, |01\rangle\}$ with $|\mathring{\mathbf{0}}\rangle \equiv |e_1\rangle \equiv |00\rangle$ and $|\mathring{\mathbf{1}}\rangle \equiv |e_2\rangle \equiv |01\rangle$.

Definition 2.9 (Local involutive duality operator)**.** The operator in Definition 2.7 acts on the $(0,1)$-sector as
$$D^{(\mathfrak{B}_x)} = |\mathring{\mathbf{0}}\rangle_x\langle\mathring{\mathbf{1}}|_x + |\mathring{\mathbf{1}}\rangle_x\langle\mathring{\mathbf{0}}|_x, \tag{2.9}$$
and trivially on the orthogonal subspaces (Parasymmetry/Holosymmetry).

Proposition 2.10 (Circuit realization on two qubits (exact on $\mathbb{C}^4$))**.** *A unitary realizing eq. (2.9) on $\mathbb{C}^4$ and acting as the identity on the orthogonal subspaces is*
$$U_D = \big(|0\rangle\langle 0|\big)_{q_1} \otimes X_{q_0} + \big(|1\rangle\langle 1|\big)_{q_1} \otimes \mathbb{I}_{q_0}. \tag{2.10}$$
A circuit for eq. (2.10) is a zero-controlled X (open control) on q_1:



Its action on the basis is

$$|00\rangle \mapsto |01\rangle\,, \quad |01\rangle \mapsto |00\rangle\,, \quad |10\rangle \mapsto |10\rangle\,, \quad |11\rangle \mapsto |11\rangle\,,$$

i.e. it swaps $|e_1\rangle \leftrightarrow |e_2\rangle$ *and leaves* $|e_3\rangle, |e_4\rangle$ *invariant.*

Proof. Orthogonal projectors $|0\rangle\langle 0|$ and $|1\rangle\langle 1|$ on q_1 ensure block action. Hermiticity and involution follow from $X^\dagger = X$, $X^2 = \mathbb{I}$ and projector orthogonality:

$$U_D^\dagger = U_D, \qquad U_D^2 = |0\rangle\langle 0| \otimes X^2 + |1\rangle\langle 1| \otimes \mathbb{I} = \mathbb{I}_4.$$

On the $(0,1)$-sector ($q_1 = 0$) the restriction is X on q_0, matching eq. (2.8). □

Corollary 2.11 (Restricted one-qubit realization on $\mathcal{H}_{(0,1)} \cong \mathbb{C}^2$)**.** *If dynamics are confined to the* $(0,1)$*-sector (i.e.* q_1 *is fixed to* $|0\rangle$*), then*

$$D^{(\mathfrak{B}_x)} \;\cong\; \sigma_X \quad \textit{on the logical duality qubit } q_0, \tag{2.11}$$

with circuit

q_0 *(duality)* —[X]—

which realizes the swap $|\mathring{\mathbf{0}}\rangle \leftrightarrow |\mathring{\mathbf{1}}\rangle$.

2.1.3 Global Involutive Duality Operators

Definition 2.12 (Global Involutive Duality Operator $\hat{D}^{(n)}_{\mathrm{PQS}}$)**.** Let $\{\mathfrak{B}_{x_1}, \ldots, \mathfrak{B}_{x_n}\}$ denote a configuration of n Primordial Quantum Singularities, each located at point $x_i \in \mathcal{X}_q$, and each equipped with its own emergent modal Hilbert space

$$\mathcal{H}^{(\mathfrak{B}_{x_i})}_{\mathrm{mod}} \cong \mathbb{C}^4,$$

as defined in Definition 4.273. The total modal Hilbert space of the configuration is the external tensor product:

$$\mathcal{H}^{(n)}_{\mathrm{PQS}} := \bigotimes_{i=1}^{n} \mathcal{H}^{(\mathfrak{B}_{x_i})}_{\mathrm{mod}},$$

constructed fiberwise over the emergent configuration space $\mathcal{X}_q$ via the Hilbert bundle of dialectical modes (Definition 4.239).

We define the *global involutive duality operator* $\hat{D}^{(n)}_{\mathrm{PQS}}$ as the operator acting diagonally across this configuration, given by:

$$\boxed{\hat{D}^{(n)}_{\mathrm{PQS}} := \bigotimes_{i=1}^{n} D^{(\mathfrak{B}_{x_i})},} \tag{2.12}$$

where each $D^{(\mathfrak{B}_{x_i})}$ is the local involutive symmetry operator, acting on the Symmetry–Anti-symmetry sector of the modal fiber $\mathcal{H}^{(\mathfrak{B}_{x_i})}_{\mathrm{mod}}$.

This global operator satisfies:

$$\hat{D}^{(n)}_{\mathrm{PQS}}\left(\bigotimes_{i=1}^{n}|\psi\rangle_{x_i}\right)=\bigotimes_{i=1}^{n}\left(D^{(\mathfrak{B}_{x_i})}|\psi\rangle_{x_i}\right),$$

and thus implements a fiberwise involution across all singularities in the configuration. If each $|\psi\rangle_{x_i}\in\mathrm{span}\{|\mathring{\mathbf{0}}\rangle_{x_i},|\mathring{\mathbf{1}}\rangle_{x_i}\}$, then this action corresponds to a coherent exchange of Symmetry and Antisymmetry modes:

$$\hat{D}^{(n)}_{\mathrm{PQS}}\left(\bigotimes_{i=1}^{n}|\mathring{\mathbf{0}}\rangle_{x_i}\right)=\bigotimes_{i=1}^{n}|\mathring{\mathbf{1}}\rangle_{x_i},\quad\text{and vice versa.}$$

As such, $\hat{D}^{(n)}_{\mathrm{PQS}}$ defines a canonical involutive automorphism of the modal Hilbert bundle, encoding a globally synchronized transformation of internal duality structure. It characterizes a natural fiberwise symmetry of the emergent quantum geometry, acting as a discrete gauge transformation under the dialectical logic of Symmetry and Antisymmetry.

Definition 2.13 (Circuit for $\hat{D}^{(n)}_{\mathrm{PQS}}$ on the two–qubit-per-site embedding)**.** Let each site $i\in\{1,\dots,n\}$ be encoded by two qubits: a *sector* qubit s_i (pinning the $(0,1)$-sector by $|0\rangle$) and a *duality* qubit d_i (carrying $|\mathring{\mathbf{0}}\rangle\leftrightarrow|\mathring{\mathbf{1}}\rangle$). Define the site-local unitary

$$U_D^{(i)}\;=\;\Big(|0\rangle\langle 0|\Big)_{s_i}\otimes X_{d_i}\;+\;\Big(|1\rangle\langle 1|\Big)_{s_i}\otimes\mathbb{I}_{d_i},$$

so that the global operator is

$$\hat{D}^{(n)}_{\mathrm{PQS}}\;=\;\bigotimes_{i=1}^{n}U_D^{(i)},\quad\text{agreeing with } eq.\ (2.12).$$

A circuit for $\hat{D}^{(n)}_{\mathrm{PQS}}$ applies, at each site i, an *open-control* (zero-controlled) X on d_i from s_i:



Remark 2.14 (Restriction to the $(0,1)$-sector at all sites)**.** If dynamics are confined to $\mathcal{H}_{(0,1)}^{\otimes n}$ (each s_i fixed to $|0\rangle$), then

$$\hat{D}^{(n)}_{\mathrm{PQS}}\;\cong\;\bigotimes_{i=1}^{n}X_{d_i}\;=\;X^{\otimes n}$$

on the logical n-qubit duality register $(d_1, \ldots, d_n)$, with circuit:



Remark 2.15 (Classification of the global involutive duality gate)**.** The operator defined in Definition 2.12 and eq. (2.12) is an *involutory, Hermitian, Clifford* gate. Concretely,

$$\hat{D}^{(n)}_{\mathrm{PQS}} = \bigotimes_{i=1}^{n} U_D^{(i)}, \quad U_D^{(i)} = \Big(|0\rangle\langle 0| \Big)_{s_i} \otimes X_{d_i} + \Big(|1\rangle\langle 1| \Big)_{s_i} \otimes \mathbb{I}_{d_i},$$

so each $U_D^{(i)}$ is an *open-controlled* X (zero-controlled-NOT) on the duality qubit d_i from the sector qubit s_i. Using the standard identity

$$U_D^{(i)} = (X_{s_i})\,\mathrm{CNOT}(s_i \to d_i)\,(X_{s_i}),$$

each factor is Clifford (hence so is the tensor product) and preserves the Pauli group under conjugation [287, 287, 349]. The gate is self-adjoint and an involution,

$$\Big(\hat{D}^{(n)}_{\mathrm{PQS}}\Big)^{\dagger} = \hat{D}^{(n)}_{\mathrm{PQS}}, \qquad \Big(\hat{D}^{(n)}_{\mathrm{PQS}}\Big)^{2} = \mathbb{I},$$

so it acts as a *reflection* (self-adjoint unitary) in the sense of elementary quantum mechanics [350]. On the embedded $(0,1)$-sector at all sites (i.e. each $s_i = |0\rangle$), the restriction reduces to the Pauli string

$$\hat{D}^{(n)}_{\mathrm{PQS}} \;\cong\; X_{d_1} \otimes \cdots \otimes X_{d_n} \;=\; X^{\otimes n},$$

which flips $|\mathring{\mathbf{0}}\rangle_{x_i} \leftrightarrow |\mathring{\mathbf{1}}\rangle_{x_i}$ at every site while leaving the orthogonal parasymmetry/holosymmetry sectors invariant [287, 351].

2.2 Parasymmetry: Quantum Realization of Synthesis

The Parasymmetric Modal Space

Definition 2.16 (Parasymmetric Modal Subspace)**.** Let $\mathfrak{B}_{x_i}$ be a Primordial Quantum Singularity with associated modal Hilbert space

$$\mathcal{H}^{(\mathfrak{B}_{x_i})}_{\mathrm{mod}} := \mathbb{C}^4.$$

Within this space, the *parasymmetric sector* is defined as the subspace spanned by the fundamental ontological duality modes:

$$\mathcal{H}^{(x_i)}_{(0,1)} := \mathrm{span}\left\{ |\mathring{\mathbf{0}}\rangle_{x_i},\ |\mathring{\mathbf{1}}\rangle_{x_i} \right\} \subset \mathbb{C}^4, \tag{2.13}$$

where the basis vectors are realized as:

$$|\overset{\circ}{\mathbf{0}}\rangle := \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \quad |\overset{\circ}{\mathbf{1}}\rangle := \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}.$$

Given a configuration of n Primordial Quantum Singularities $\{\mathfrak{B}_{x_1}, \ldots, \mathfrak{B}_{x_n}\}$, the *global parasymmetric modal subspace* is defined as the n-fold tensor product:

$$\mathcal{H}^{(n)}_{\text{para}} := \bigotimes_{i=1}^{n} \mathcal{H}^{(x_i)}_{(0,1)} \cong (\mathbb{C}^2)^{\otimes n}. \tag{2.14}$$

Hence, the parasymmetric sector has dimension:

$$\dim\left(\mathcal{H}^{(n)}_{\text{para}}\right) = 2^n.$$

2.2.1 Local Parasymmetric Projector of an PQS

Definition 2.17 (Parasymmetric Projector $P^{(\mathfrak{B}_x)}_{\text{para}}$)**.** Let $\mathfrak{B}_x$ be a Primordial Quantum Singularity with modal Hilbert space $\mathcal{H}^{(\mathfrak{B}_x)}_{\text{mod}} = \mathbb{C}^4$, and define the $(0,1)$-subspace as

$$\mathcal{H}_{(0,1)} := \text{span}\,\{|e_1\rangle, |e_2\rangle\} = \left\{ \begin{pmatrix} c_1 \\ c_2 \\ 0 \\ 0 \end{pmatrix} \;\middle|\; c_1, c_2 \in \mathbb{C} \right\}. \tag{2.15}$$

The *parasymmetric mode* is the equal-weighted coherent superposition:

$$|\overset{\circ}{\mathbf{0}} + \overset{\circ}{\mathbf{1}}\rangle_x := \frac{1}{\sqrt{2}}\left(|\overset{\circ}{\mathbf{0}}\rangle_x + |\overset{\circ}{\mathbf{1}}\rangle_x\right) = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \end{pmatrix}_x. \tag{2.16}$$

Then the *parasymmetric projector* is defined as:

$$\boxed{P^{(\mathfrak{B}_x)}_{\text{para}} := \left(|\overset{\circ}{\mathbf{0}} + \overset{\circ}{\mathbf{1}}\rangle_x\right)\left(\langle\overset{\circ}{\mathbf{0}} + \overset{\circ}{\mathbf{1}}|_x\right) = \frac{1}{2}\begin{pmatrix} 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}} \tag{2.17}$$

Properties

1. **Idempotency and Hermiticity**

$$(P_{\text{para}}^{(\mathfrak{B}_x)})^2 = P_{\text{para}}^{(\mathfrak{B}_x)}, \qquad (P_{\text{para}}^{(\mathfrak{B}_x)})^\dagger = P_{\text{para}}^{(\mathfrak{B}_x)}.$$

2. **Modal Projection**
   For any modal state $|\psi\rangle_x \in \mathcal{H}_{(0,1)}$, i.e.,

$$|\psi\rangle_x = \begin{pmatrix} c_0 \\ c_1 \\ 0 \\ 0 \end{pmatrix}_x ,$$

   we have:

$$P_{\text{para}}^{(\mathfrak{B}_x)}|\psi\rangle_x = \frac{1}{2}(c_0 + c_1) \cdot \begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \end{pmatrix}_x . \tag{2.18}$$

3. **Interpretation**
   This operator isolates the *parasymmetric component* of any modal state in the $(0,1)$-sector, i.e., the projection along the direction of internal *dialectical synthesis* between Symmetry and Antisymmetry.
   It does *not* extend beyond the first two components of $\mathbb{C}^4$, since no formal realization has yet been provided for Holosymmetry or orthogonal parasymmetric bases. The parasymmetric state here is not orthogonal to both $|e_1\rangle$ and $|e_2\rangle$, but spans a *coherent subdirection* within that space.

The Parasymmetric Projector: An Operator of Synthesis The Parasymmetric Projector, denoted $P_{\text{para}}^{(\mathfrak{B}_x)}$, represents a profound mathematical realization of the dialectical principle of synthesis within the quantum mechanical framework. This operator selects a component of a quantum state as it also enacts a fundamental philosophical operation by isolating precisely that aspect of any state which embodies the coherent unification of the primal dialectical opposites. Formally, it projects onto the normalized superposition state:

$$|\mathring{\mathbf{0}} + \mathring{\mathbf{1}}\rangle_x := \frac{1}{\sqrt{2}} \left(|\mathring{\mathbf{0}}\rangle_x + |\mathring{\mathbf{1}}\rangle_x \right)$$

The normalization factor $1/\sqrt{2}$ ensures that this synthetic state maintains unit norm, thereby preserving quantum mechanical consistency while encoding the equal-weighted presence of both Thesis and Antithesis. As a "coherence filter," $P_{\text{para}}^{(\mathfrak{B}_x)}$ extracts from any arbitrary modal state its component of maximal coherent synthesis—that precise quantum amplitude which refuses the binary choice between existence and non-existence, instead embracing their simultaneous co-presence. Within the constraints of the primary $(0,1)$-subspace, this operator thus provides a concrete quantum mechanical instantiation of the third vertex of the dialectical tetralemma, demonstrating how abstract logical synthesis manifests as quantum superposition in the emergent modal Hilbert space of each Primordial Quantum Singularity.

2.2.2 The Local Anti-Parasymmetric Projector

Definition 2.18 (Anti-Parasymmetric Projector $P^{(\mathfrak{B}_x)}_{\text{antipara}}$)**.** Let the parasymmetric mode be defined within the $(0,1)$-subspace as

$$|\mathring{\mathbf{0}} + \mathring{\mathbf{1}}\rangle_x := \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \end{pmatrix}_x . \tag{2.19}$$

We seek a state $|\psi^\perp\rangle \in \mathcal{H}_{(0,1)}$ such that

$$\langle \psi^\perp | \mathring{\mathbf{0}} + \mathring{\mathbf{1}} \rangle = 0.$$

This yields the *anti-parasymmetric mode*:

$$|\mathring{\mathbf{0}} - \mathring{\mathbf{1}}\rangle_x := \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -1 \\ 0 \\ 0 \end{pmatrix}_x , \tag{2.20}$$

which satisfies:

- $\|\cdot\| = 1$,
- orthogonality to $|\mathring{\mathbf{0}} + \mathring{\mathbf{1}}\rangle$,
- together with the parasymmetric mode, forms an orthonormal basis for $\mathcal{H}_{(0,1)}$.

The *anti-parasymmetric projector* is then defined as:

$$\boxed{P^{(\mathfrak{B}_x)}_{\text{antipara}} := |\mathring{\mathbf{0}} - \mathring{\mathbf{1}}\rangle_x \langle \mathring{\mathbf{0}} - \mathring{\mathbf{1}}|_x = \frac{1}{2} \begin{pmatrix} 1 & -1 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}} \tag{2.21}$$

Properties

1. **Orthogonality to the Parasymmetric Projector**:

$$P^{(\mathfrak{B}_x)}_{\text{para}} \cdot P^{(\mathfrak{B}_x)}_{\text{antipara}} = 0.$$

2. **Complementarity on the** $(0,1)$**-sector**:

$$P^{(\mathfrak{B}_x)}_{\text{para}} + P^{(\mathfrak{B}_x)}_{\text{antipara}} = |e_1\rangle\langle e_1| + |e_2\rangle\langle e_2| = P_T + P_A = P_{(0,1)} := \text{Proj}_{\mathcal{H}_{(0,1)}} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

(Here $|e_1\rangle, |e_2\rangle$ are the Symmetry/Antisymmetry basis; $P_T := |e_1\rangle\langle e_1|$, $P_A := |e_2\rangle\langle e_2|$.)

Interpretation. The anti-parasymmetric mode $|\mathring{\mathbf{0}} - \mathring{\mathbf{1}}\rangle$ captures the *modal dissonance*—the internal tension or conflict between Symmetry and Antisymmetry. Unlike the parasymmetric mode, which expresses dialectical synthesis, this state represents a logical stance of *non-coincidence* or *disjointed duality*. It remains fully internal to $\mathcal{H}_{(0,1)}$, and functions as a canonical orthogonal complement to the parasymmetric synthesis. ***Valuation Link*** $P^{(x)}_{\text{para}} \equiv P_S$ in Definition 1.18, so $v_{\text{Synthesis}}(\rho) = \text{Tr}\big(P^{(x)}_{\text{para}}\rho\big)$ (and analogously for the global $P^{(x_1\ldots x_n)}_{\text{para}}$).

Definition 2.19 (Local test for $P^{(\mathfrak{B}_x)}_{\text{para}}$ on $\mathcal{H}_{(0,1)}$)**.** Since $P^{(\mathfrak{B}_x)}_{\text{para}} = |+\rangle\langle+|$ with $|+\rangle := \frac{1}{\sqrt{2}}\big(|\mathring{\mathbf{0}}\rangle_x + |\mathring{\mathbf{1}}\rangle_x\big)$, a projective measurement in the X basis realizes the projector. Operationally: apply a Hadamard on the duality qubit and measure in Z; outcome 0 corresponds to projection by $P^{(\mathfrak{B}_x)}_{\text{para}}$, outcome 1 to $P^{(\mathfrak{B}_x)}_{\text{antipara}}$.



For an input state ρ on $\mathcal{H}_{(0,1)}$, $\Pr[0] = \text{Tr}\big(P^{(\mathfrak{B}_x)}_{\text{para}}\,\rho\big)$, $\Pr[1] = \text{Tr}\big(P^{(\mathfrak{B}_x)}_{\text{antipara}}\,\rho\big)$, with $P^{(\mathfrak{B}_x)}_{\text{para}} + P^{(\mathfrak{B}_x)}_{\text{antipara}} = P_{(0,1)}$.

Naimark flags. Ancilla bits (or registers) that record the outcome labels of a POVM via a Naimark dilation are called *Naimark flags* [287, 352, 353].

Binary case. Let E be an effect on a system $\mathcal{H}_S$ with $0 \le E \le I$. A Naimark dilation is a unitary U on $\mathcal{H}_S \otimes \mathcal{H}_a$ (with the ancilla initialised in $|0\rangle_a$) such that

$$U\big(|\psi\rangle \otimes |0\rangle_a\big) = \big(\sqrt{E}\,|\psi\rangle\big) \otimes |1\rangle_a + \big(\sqrt{I-E}\,|\psi\rangle\big) \otimes |0\rangle_a.$$

Measuring the ancilla in the Z basis "flags'' outcome 1 with probability $\langle\psi|E|\psi\rangle$ [287, 354]. If $E = P$ is a projector, then

$$U_P\big(|\psi\rangle \otimes |0\rangle_a\big) = (P|\psi\rangle) \otimes |1\rangle_a + \big((I-P)|\psi\rangle\big) \otimes |0\rangle_a,$$

so the ancilla acts as a flag qubit indicating membership in the subspace $\text{im}\,P$.

General POVM. Given a POVM $\{E_i\}_i$, there exists an ancilla with orthonormal basis $\{|i\rangle\}_i$ and an isometry $V : \mathcal{H}_S \to \mathcal{H}_S \otimes \mathcal{H}_a$ such that

$$E_i \;=\; V^\dagger\,(I \otimes |i\rangle\langle i|)\,V,$$

so that the ancilla basis states $|i\rangle$ serve as the Naimark flags [287, 352, 353, 355].

Circuit realisation (lifted parasymmetric projector). With the notation of the main text, $V^{(A)}_+$ (resp. $V^{(B)}_+$) rotates into a basis in which the desired projector is a computational-basis predicate; a multi-controlled X flips the ancilla a_A (resp. a_B) iff the data register lies in that projector's support, and uncomputing with V_+ restores the data. Concretely,

$$\big(V^{(A)}_+ \otimes I\big)\,CX\,\big(V^{(A)\dagger}_+ \otimes I\big) :\; |\psi\rangle\,|0\rangle_{a_A} \longmapsto \big(P_{\text{para}}|\psi\rangle\big)\,|1\rangle_{a_A} + \big((I - P_{\text{para}})|\psi\rangle\big)\,|0\rangle_{a_A}.$$

Measuring a_A or a_B gives a projective realisation of the corresponding effect (a *classical flag*); omitting the measurement yields a *coherent flag*, leaving the ancilla entangled with the system for downstream processing before optional uncomputation [287].

Definition 2.20 (Exact dilation of $P_{\text{para}}^{(\mathfrak{B}_x)}$ on $\mathbb{C}^4$)**.** Embed $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)} \cong \mathbb{C}^4$ into two qubits: a sector qubit s and a duality qubit d, with $|\mathring{\mathbf{0}}\rangle \equiv |0\rangle_s |0\rangle_d$, $|\mathring{\mathbf{1}}\rangle \equiv |0\rangle_s |1\rangle_d$. Introduce a flag ancilla a initialized in $|0\rangle_a$. Define the controlled-H that only acts on the $(0,1)$-sector,

$$U_{\text{prep}} := \Big(|0\rangle\langle 0| \Big)_s \otimes H_d \; + \; \Big(|1\rangle\langle 1| \Big)_s \otimes \mathbb{I}_d.$$

Then perform a two-control (both *open* / zero-controls) Toffoli that flips a iff s=0 and d=0 (i.e. after the H-rotation the state has $|0\rangle_d$):



Measuring a and postselecting on outcome 1 realizes the projector $P_{\text{para}}^{(\mathfrak{B}_x)} = |0\rangle\langle 0|_s \otimes |+\rangle\langle +|_d$ on the system:

$$\Pr[a{=}1] = \operatorname{Tr}\Big(\big[\, |0\rangle\langle 0|_s \otimes |+\rangle\langle +|_d \, \big] \, \rho \Big), \qquad \rho \text{ on } \mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}.$$

By exchanging the open control on d (i.e. conditioning on d=1 after H), the complementary projector $P_{\text{antipara}}^{(\mathfrak{B}_x)}$ is obtained.

The Anti-Parasymmetric Projector: An Operator of Dissonance For every projection operator in quantum mechanics, mathematical completeness demands the existence of a complementary projection onto the orthogonal subspace. The Anti-Parasymmetric Projector, $P_{\text{antipara}}^{(\mathfrak{B}_x)}$, fulfills this role as the natural counterpart to the Parasymmetric Projector. This operator is constructed upon the mode orthogonal to the synthesis state within the $\mathcal{H}_{(0,1)}$ subspace, specifically projecting onto the anti-parasymmetric mode:

$$|\mathring{\mathbf{0}} - \mathring{\mathbf{1}}\rangle_x := \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -1 \\ 0 \\ 0 \end{pmatrix}_x$$

This state embodies the principle of perfect destructive interference between the Symmetry and Antisymmetry amplitudes, representing what the manuscript aptly characterizes as "modal dissonance" or "disjointed duality." The minus sign in the superposition ensures that the fundamental modes interfere destructively rather than constructively, yielding a state that captures opposition rather than synthesis. The corresponding projection operator takes the explicit matrix form:

$$P^{(\mathfrak{B}_x)}_{\text{antipara}} = \frac{1}{2}\begin{pmatrix} 1 & -1 & 0 & 0 \\ -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

The Interplay: A Complete Basis for Duality The profound elegance of this dual-projector construction manifests in the mathematical interplay between the Parasymmetric and Anti-Parasymmetric operators. These projectors satisfy two fundamental properties that illuminate their complementary nature. First, the orthogonality relation $P_{\text{para}} \cdot P_{\text{antipara}} = 0$ confirms that a quantum state cannot simultaneously embody perfect synthesis and perfect dissonance—these represent mutually exclusive modal configurations, and any measurement must yield one or the other. Second, the completeness relation $P_{\text{para}} + P_{\text{antipara}} = P_{(0,1)}$ establishes that these two operators span the entire $(0, 1)$-subspace, ensuring that any state within the fundamental duality sector admits a unique decomposition into components of pure synthesis and pure dissonance.

This construction effectively implements a change of basis within the modal Hilbert space, transitioning from the ontological "existence" basis of Symmetry, Antisymmetry to the phenomenologically richer "relationship" basis of Synthesis, Dissonance. This new basis proves particularly powerful for describing interactions and dynamics, as it directly quantifies the relational character—whether harmonious or dissonant—between the foundational dialectical modes. The mathematical framework thus captures the presence or absence of modal components and the quality of their mutual relationship within the quantum state.

Definition 2.21 (Local test for $P^{(\mathfrak{B}_x)}_{\text{antipara}}$ on $\mathcal{H}_{(0,1)}$)**.** Project onto $|-\rangle$ by rotating to the computational basis and measuring: apply H to the duality qubit and measure in Z; outcome 1 realizes $P^{(\mathfrak{B}_x)}_{\text{antipara}}$.



For input state ρ on $\mathcal{H}_{(0,1)}$,

$$\Pr[1] = \operatorname{Tr}\Big(P^{(\mathfrak{B}_x)}_{\text{antipara}}\,\rho\Big), \qquad \Pr[0] = \operatorname{Tr}\Big(P^{(\mathfrak{B}_x)}_{\text{para}}\,\rho\Big),$$

with $P^{(\mathfrak{B}_x)}_{\text{para}} + P^{(\mathfrak{B}_x)}_{\text{antipara}} = P_{(0,1)}$.

Definition 2.22 (Exact dilation of $P^{(\mathfrak{B}_x)}_{\text{antipara}}$ on $\mathbb{C}^4$)**.** Embed $\mathcal{H}^{(\mathfrak{B}_x)}_{\text{mod}} \cong \mathbb{C}^4$ into sector s and duality d qubits with $|\mathring{\mathbf{0}}\rangle \equiv |0\rangle_s\,|0\rangle_d$, $|\mathring{\mathbf{1}}\rangle \equiv |0\rangle_s\,|1\rangle_d$, and introduce a flag ancilla a initialized in $|0\rangle_a$. First perform a *sector-conditioned* Hadamard on d,

$$U_{\text{prep}} := \Big(\,|0\rangle\langle 0|\,\Big)_s \otimes H_d \;+\; \Big(\,|1\rangle\langle 1|\,\Big)_s \otimes \mathbb{I}_d,$$

then flip a iff s=0 and d=1 (mixed-control Toffoli: open control on s, closed control on d):



Measuring a and postselecting on outcome 1 realizes the projector

$$P^{(\mathfrak{B}_x)}_{\text{antipara}} = |0\rangle\langle 0|_s \otimes |-\rangle\langle -|_d\,, \qquad \Pr[a{=}1] = \operatorname{Tr}\Big(\big[\,|0\rangle\langle 0|_s \otimes |-\rangle\langle -|_d\,\big]\,\rho\Big).$$

Swapping the target condition to d=0 yields the complementary $P^{(\mathfrak{B}_x)}_{\text{para}}$.

2.2.3 The Global Parasymmetric Projector

Definition 2.23 (Global Parasymmetric Projector)**.** Let $\mathfrak{B}_x$ and $\mathfrak{B}_{x'}$ be two Primordial Quantum Singularities, each with modal Hilbert space restricted to the $(0,1)$-sector:

$$\mathcal{H}^{(x)}_{(0,1)},\, \mathcal{H}^{(x')}_{(0,1)} \subset \mathbb{C}^4,$$

and local basis vectors

$$|\mathring{\mathbf{0}}\rangle := |e_1\rangle := \begin{pmatrix}1\\0\\0\\0\end{pmatrix}, \quad |\mathring{\mathbf{1}}\rangle := |e_2\rangle := \begin{pmatrix}0\\1\\0\\0\end{pmatrix}.$$

The *ontological parasymmetric state* on the bipartite tensor product space is:

$$|\Psi_{\text{para}}\rangle_{xx'} := \frac{1}{\sqrt{2}}\left(|\mathring{\mathbf{0}}\rangle_x \otimes |\mathring{\mathbf{1}}\rangle_{x'} + |\mathring{\mathbf{1}}\rangle_x \otimes |\mathring{\mathbf{0}}\rangle_{x'}\right). \tag{2.22}$$

Conventional Quantum State Realization Under the canonical lexicographic basis of $\mathbb{C}^4 \otimes \mathbb{C}^4 \cong \mathbb{C}^{16}$, where

$$|e_i\rangle \otimes |e_j\rangle \mapsto \mathbf{e}_{4(i-1)+j},$$

the vector $|\Psi_{\text{para}}\rangle_{xx'}$ becomes:

$$|\Psi_{\text{para}}\rangle_{xx'} = \frac{1}{\sqrt{2}}\left(|e_1\rangle_x \otimes |e_2\rangle_{x'} + |e_2\rangle_x \otimes |e_1\rangle_{x'}\right) = \frac{1}{\sqrt{2}}\left(\mathbf{e}_2 + \mathbf{e}_5\right), \tag{2.23}$$

where $\mathbf{e}_k$ denotes the standard basis vector in $\mathbb{C}^{16}$ with a 1 in the k-th position.

Explicitly, this gives:

$$|\Psi_{\text{para}}\rangle_{xx'} = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \\ 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix},$$

with nonzero components in positions 2 and 5 corresponding to the two configurations $(e_1 \otimes e_2)$ and $(e_2 \otimes e_1)$.

Global Parasymmetric Projector The associated rank-1 orthogonal projection operator is:

$$\boxed{P_{\text{para}}^{(xx')} := |\Psi_{\text{para}}\rangle_{xx'}\langle\Psi_{\text{para}}|_{xx'}.} \tag{2.24}$$

Interpretation. This state encodes a non-separable entanglement of Symmetry and Antisymmetry modes across distinct PQS loci x and x', corresponding to a *distributed realization of dialectical synthesis*. The lexicographic tensor ordering ensures a unique representation in conventional quantum mechanics, preserving fidelity between ontological meaning and Hilbert space structure.

Global Parasymmetry: Synthesis as Entanglement The theoretical framework takes a pivotal step by extending the concept of parasymmetry beyond the confines of a single Primordial Quantum Singularity to encompass the relational structure between two distinct singularities, $\mathfrak{B}_x$ and $\mathfrak{B}_{x'}$. This extension marks the transition from local to genuinely relational physics within the IQR. The proposed ontological parasymmetric state is defined as:

$$|\Psi_{\text{para}}\rangle_{xx'} := \frac{1}{\sqrt{2}} \left(|\mathring{\mathbf{0}}\rangle_x \otimes |\mathring{\mathbf{1}}\rangle_{x'} + |\mathring{\mathbf{1}}\rangle_x \otimes |\mathring{\mathbf{0}}\rangle_{x'} \right)$$

This mathematical form corresponds precisely to one of the four Bell states—the canonical exemplars of maximally entangled two-qubit systems in quantum information theory.

This identification carries profound theoretical implications, revealing the dialectical synthesis between Symmetry and Antisymmetry to be a fundamentally non-local phenomenon, manifesting as a non-separable quantum entanglement distributed across spatially distinct singularities. Under measurement, if the modal state of singularity $\mathfrak{B}_x$ collapses to Symmetry ($|\overset{\circ}{\mathbf{0}}\rangle_x$), then $\mathfrak{B}_{x'}$ necessarily manifests in the Antisymmetry mode ($|\overset{\circ}{\mathbf{1}}\rangle_{x'}$), and conversely. The synthesis therefore exists neither at location x nor at x' individually, but emerges purely through the quantum correlation binding the two singularities.

Within the $\mathbb{C}^{16}$ tensor product space of the combined system, this state admits the explicit representation $\frac{1}{\sqrt{2}}(\mathbf{e}_2 + \mathbf{e}_5)$ under the lexicographic ordering of basis vectors. This mathematical realization confirms that parasymmetric synthesis, when extended to the multi-singularity domain, fundamentally embodies the principle of quantum entanglement—establishing relational coherence as the mechanism through which dialectical synthesis transcends local boundaries to create genuinely non-local quantum structures.

Definition 2.24 (Bell-basis test for $P^{(xx')}_{\text{para}}$ on $\mathcal{H}^{(x)}_{(0,1)} \otimes \mathcal{H}^{(x')}_{(0,1)}$)**.** Let d_x and $d_{x'}$ denote the duality qubits for sites x and x'. A standard Bell-basis demultiplexing followed by Z-readout realizes the rank-1 projector $P^{(xx')}_{\text{para}} = |\Psi_{\text{para}}\rangle\langle\Psi_{\text{para}}|$ via postselection:



Under the inverse Bell map (CNOT then H on d_x), the Bell states are sent to $\{|\Phi^+\rangle, |\Phi^-\rangle, |\Psi^+\rangle, |\Psi^-\rangle\} \mapsto \{|00\rangle, |01\rangle, |10\rangle, |11\rangle\}$, so $|\Psi_{\text{para}}\rangle = |\Psi^+\rangle$ produces outcome $(1, 0)$. Thus, for any ρ on $\mathcal{H}^{(x)}_{(0,1)} \otimes \mathcal{H}^{(x')}_{(0,1)}$,

$$\Pr[(d_x, d_{x'}) = (1, 0)] = \mathrm{Tr}\Big(P^{(xx')}_{\text{para}}\,\rho\Big), \quad \rho \xrightarrow{\text{postselect } (1,0)} \frac{P^{(xx')}_{\text{para}}\,\rho\,P^{(xx')}_{\text{para}}}{\mathrm{Tr}(P^{(xx')}_{\text{para}}\rho)}.$$

Definition 2.25 (Coherent flagged realization (Naimark dilation) of $P^{(xx')}_{\text{para}}$)**.** Introduce a flag ancilla a initialized in $|0\rangle_a$. Map to the computational basis, flip a iff the pattern $d_x d_{x'} = 10$ is present, then uncompute:



Here the mixed-control Toffoli uses a *closed* control on d_x (selecting 1) and an *open* control on $d_{x'}$ (selecting 0), so that

$$\Pr[a{=}1] = \mathrm{Tr}\Big(P^{(xx')}_{\text{para}}\,\rho\Big), \qquad \text{and (on } a{=}1\text{) the system is projected by } P^{(xx')}_{\text{para}}.$$

Remark 2.26 (Relation to the lexicographic embedding)**.** With the identification $|\mathring{\mathbf{0}}\rangle \equiv |0\rangle$, $|\mathring{\mathbf{1}}\rangle \equiv |1\rangle$ per site, $|\Psi_{\text{para}}\rangle = \frac{1}{\sqrt{2}}(|01\rangle + |10\rangle)$ and $P^{(xx')}_{\text{para}} = |\Psi^+\rangle\langle\Psi^+|$. Under the full $\mathbb{C}^{16}$ lexicographic ordering of Definition 2.23, these circuits act nontrivially only on the $\mathcal{H}^{(x)}_{(0,1)} \otimes \mathcal{H}^{(x')}_{(0,1)}$ block and as the identity elsewhere.

Remark 2.27 (Nature of the global parasymmetric projector)**.** The operator introduced in Definition 2.23 is *not* a unitary gate but a rank-1 orthogonal *projector/measurement effect* on the duality sector,

$$P^{(xx')}_{\text{para}} \;=\; |\Psi_{\text{para}}\rangle\langle\Psi_{\text{para}}|\,, \qquad |\Psi_{\text{para}}\rangle = \tfrac{1}{\sqrt{2}}\Big(|\mathring{\mathbf{0}}\rangle_x\,|\mathring{\mathbf{1}}\rangle_{x'} + |\mathring{\mathbf{1}}\rangle_x\,|\mathring{\mathbf{0}}\rangle_{x'}\Big)\,,$$

i.e. the Bell state $|\Psi^+\rangle$ on the two $(0,1)$-sector (duality) qubits. Consequently,

$$\Big(P^{(xx')}_{\text{para}}\Big)^\dagger = P^{(xx')}_{\text{para}}, \qquad \Big(P^{(xx')}_{\text{para}}\Big)^2 = P^{(xx')}_{\text{para}},$$

so it is Hermitian, idempotent, positive, and *non-unitary* (a projector rather than a gate) [287]. A convenient Pauli–stabilizer form on the two duality qubits $(d_x, d_{x'})$ is

$$P^{(xx')}_{\text{para}} = \frac{1}{4}\left(\mathbb{I} - Z_{d_x}Z_{d_{x'}}\right)\left(\mathbb{I} + X_{d_x}X_{d_{x'}}\right),$$

i.e. projection onto the simultaneous (-1)-eigenspace of $Z\otimes Z$ and $(+1)$-eigenspace of $X\otimes X$ (the stabilizers of $|\Psi^+\rangle$) [349]. Operationally, $P^{(xx')}_{\text{para}}$ is implemented by a Clifford uncompute to the Bell basis (e.g. CNOT then H on the first duality qubit) followed by computational-basis measurement and *postselection* of the outcome corresponding to $|\Psi^+\rangle$; equivalently, it can be realized via a Naimark dilation with a flagged ancilla and Clifford controls [287, 351].

2.2.4 The Global Anti-Parasymmetric Projector

Definition 2.28. The orthogonal anti-parasymmetric state is defined ontologically as:

$$|\Psi_{\text{antipara}}\rangle_{xx'} := \frac{1}{\sqrt{2}}\left(|\mathring{\mathbf{0}}\rangle_x \otimes |\mathring{\mathbf{1}}\rangle_{x'} - |\mathring{\mathbf{1}}\rangle_x \otimes |\mathring{\mathbf{0}}\rangle_{x'}\right). \tag{2.25}$$

Conventional Quantum State Realization Using the standard embedding:

$$|\mathring{\mathbf{0}}\rangle := |e_1\rangle = \begin{pmatrix}1\\0\\0\\0\end{pmatrix}, \quad |\mathring{\mathbf{1}}\rangle := |e_2\rangle = \begin{pmatrix}0\\1\\0\\0\end{pmatrix},$$

and under the lexicographic ordering of the basis of $\mathbb{C}^4 \otimes \mathbb{C}^4 \cong \mathbb{C}^{16}$, where:

$$|e_i\rangle \otimes |e_j\rangle \mapsto \mathbf{e}_{4(i-1)+j},$$

we obtain:

$$|\Psi_{\text{antipara}}\rangle_{xx'} = \frac{1}{\sqrt{2}}\left(|e_1\rangle_x \otimes |e_2\rangle_{x'} - |e_2\rangle_x \otimes |e_1\rangle_{x'}\right) = \frac{1}{\sqrt{2}}\left(\mathbf{e}_2 - \mathbf{e}_5\right), \tag{2.26}$$

which expands to:

$$|\Psi_{\text{antipara}}\rangle_{xx'} = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \\ -1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}.$$

Projector The corresponding orthogonal projector is:

$$\boxed{P^{(xx')}_{\text{antipara}} := |\Psi_{\text{antipara}}\rangle_{xx'}\langle\Psi_{\text{antipara}}|_{xx'}.} \tag{2.27}$$

Properties

1. **Orthogonality:**
$$P^{(xx')}_{\text{para}} \cdot P^{(xx')}_{\text{antipara}} = 0.$$
2. **Completeness over the symmetric–antisymmetric subspace:**
$$P^{(xx')}_{\text{para}} + P^{(xx')}_{\text{antipara}} = P^{(xx')}_{\text{exchange}},$$
where $P^{(xx')}_{\text{exchange}}$ projects onto the span:
$$\text{span}\left\{|\mathring{\mathbf{0}}\rangle \otimes |\mathring{\mathbf{1}}\rangle,\ |\mathring{\mathbf{1}}\rangle \otimes |\mathring{\mathbf{0}}\rangle\right\} = \text{span}\left\{\mathbf{e}_2,\ \mathbf{e}_5\right\}.$$

Interpretation. The anti-parasymmetric state $|\Psi_{\text{antipara}}\rangle_{xx'}$ encodes an *entangled modal opposition*—a dialectical structure in which Symmetry and Antisymmetry modes remain in relational conflict across distinct loci. In contrast to the parasymmetric synthesis, this state represents the *logical antithesis* of modal coherence, and its orthogonality ensures a clean partition of the inter-modal entanglement space. The inclusion of both ontological and conventional forms ensures precise mapping between quantum formalism and the duality-structured ontological semantics of Primordial Quantum Singularities.

Global Anti-Parasymmetry: Opposition as Entanglement Following the same theoretical logic, the framework defines the orthogonal anti-parasymmetric state as:

$$|\Psi_{\text{antipara}}\rangle_{xx'} := \frac{1}{\sqrt{2}}\left(|\mathring{\mathbf{0}}\rangle_x \otimes |\mathring{\mathbf{1}}\rangle_{x'} - |\mathring{\mathbf{1}}\rangle_x \otimes |\mathring{\mathbf{0}}\rangle_{x'}\right)$$

This state constitutes another member of the canonical Bell state quartet, distinguished from its parasymmetric counterpart by the crucial minus sign, which introduces a relative phase of π between the two superposed configurations. In quantum mechanics, such phase differences carry genuine physical significance and engender distinct correlational structures. Whereas the parasymmetric state embodies dialectical synthesis, the anti-parasymmetric state represents entangled modal opposition—a configuration of perfect anti-correlation when analyzed in the synthesis/dissonance basis.

Completeness of the Entangled Subspace The corresponding projection operators, $P^{(xx')}_{\text{para}}$ and $P^{(xx')}_{\text{antipara}}$, satisfy the fundamental properties of orthogonality and completeness within the relevant subspace. Their orthogonality ensures mutual exclusivity of the parasymmetric and anti-parasymmetric configurations, while their sum projects onto the entire two-dimensional space spanned by the basis states $\{|\mathring{\mathbf{0}}\rangle_x \otimes |\mathring{\mathbf{1}}\rangle_{x'}, |\mathring{\mathbf{1}}\rangle_x \otimes |\mathring{\mathbf{0}}\rangle_{x'}\}$.

This completeness relation demonstrates that any quantum entanglement between the Symmetry mode of one PQS and the Antisymmetry mode of another admits a unique decomposition into components of pure synthesis ($|\Psi_{\text{para}}\rangle$) and pure opposition ($|\Psi_{\text{antipara}}\rangle$). The mathematical framework thus elevates the abstract principles of dialectical logic from local modal properties to the fundamental architecture of non-local quantum entanglement, establishing a rigorous foundation for the construction of a genuinely relational universe within the IQR.

Definition 2.29 (Bell-basis test for $P^{(xx')}_{\text{antipara}}$ on $\mathcal{H}^{(x)}_{(0,1)} \otimes \mathcal{H}^{(x')}_{(0,1)}$)**.** Let d_x and $d_{x'}$ be the duality qubits of sites x, x'. The inverse Bell transform (CNOT then H on d_x) maps $\{|\Phi^+\rangle, |\Phi^-\rangle, |\Psi^+\rangle, |\Psi^-\rangle\} \to \{|00\rangle, |01\rangle, |10\rangle, |11\rangle\}$. Thus, postselecting the outcome $(1,1)$ realizes the projector $P^{(xx')}_{\text{antipara}} = |\Psi^-\rangle\langle\Psi^-|$:



For any state ρ on $\mathcal{H}^{(x)}_{(0,1)}\otimes\mathcal{H}^{(x')}_{(0,1)}$,

$$\Pr[(d_x,d_{x'})=(1,1)]=\mathrm{Tr}\Big(P^{(xx')}_{\text{antipara}}\,\rho\Big),\qquad \rho\ \xrightarrow{\text{postselect }(1,1)}\ \frac{P^{(xx')}_{\text{antipara}}\rho\,P^{(xx')}_{\text{antipara}}}{\mathrm{Tr}(P^{(xx')}_{\text{antipara}}\rho)}.$$

Definition 2.30 (Coherent flagged realization (Naimark dilation) of $P^{(xx')}_{\text{antipara}}$)**.** Introduce a flag ancilla a initialized in $|0\rangle_a$. Demultiplex to the computational basis, flip a iff the pattern $d_xd_{x'}=11$ is present, then uncompute:



Then

$$\Pr[a{=}1]=\mathrm{Tr}\Big(P^{(xx')}_{\text{antipara}}\,\rho\Big),\quad \text{and on } a{=}1 \text{ the system is projected by } P^{(xx')}_{\text{antipara}}.$$

Remark 2.31 (Stabilizer form)**.** On the two duality qubits $(d_x,d_{x'})$, the anti-parasymmetric projector admits the Pauli form

$$P^{(xx')}_{\text{antipara}}=\frac{1}{4}\Big(\mathbb{I}-Z_{d_x}Z_{d_{x'}}\Big)\Big(\mathbb{I}-X_{d_x}X_{d_{x'}}\Big),$$

selecting the simultaneous (-1)-eigenspaces of both $Z\otimes Z$ and $X\otimes X$, i.e. the Bell state $|\Psi^-\rangle$.

Remark 2.32 (Nature of the global anti-parasymmetric projector)**.** The operator in eqs. (2.25) and (2.27) is *not* a unitary gate but a rank-1 orthogonal *projector/measurement effect* onto the Bell state

$$P^{(xx')}_{\text{antipara}}\ =\ |\Psi_{\text{antipara}}\rangle\langle\Psi_{\text{antipara}}|\,,\qquad |\Psi_{\text{antipara}}\rangle=\tfrac{1}{\sqrt{2}}\Big(|\mathring{\mathbf{0}}\rangle_x\,|\mathring{\mathbf{1}}\rangle_{x'}-|\mathring{\mathbf{1}}\rangle_x\,|\mathring{\mathbf{0}}\rangle_{x'}\Big)=|\Psi^-\rangle\,.$$

Consequently $P^{(xx')}_{\text{antipara}}$ is Hermitian, idempotent and positive, but non-unitary: $(P^{(xx')}_{\text{antipara}})^\dagger=P^{(xx')}_{\text{antipara}}$, $(P^{(xx')}_{\text{antipara}})^2=P^{(xx')}_{\text{antipara}}$ [287]. On the two duality qubits $(d_x,d_{x'})$ the stabilizer–Pauli form is

$$P^{(xx')}_{\text{antipara}}=\frac{1}{4}\Big(\mathbb{I}-Z_{d_x}Z_{d_{x'}}\Big)\Big(\mathbb{I}-X_{d_x}X_{d_{x'}}\Big),$$

selecting the simultaneous (-1)-eigenspaces of $Z\otimes Z$ and $X\otimes X$, which uniquely characterizes $|\Psi^-\rangle$ within the Bell basis [349]. Operationally, $P^{(xx')}_{\text{antipara}}$ is implemented by Clifford uncompute to the Bell basis (e.g. CNOT followed by H on the first duality qubit), computational-basis measurement, and *postselection* of the $(1,1)$ outcome; equivalently, a Naimark dilation with a flagged ancilla and controlled-Clifford gates realizes the same projector coherently [287, 351].

2.2.5 Multipartite Parasymmetric Entanglement

Definition 2.33 (Multipartite Parasymmetric State)**.** Let $\{\mathfrak{B}_{x_1}, \mathfrak{B}_{x_2}, \ldots, \mathfrak{B}_{x_n}\}$ be a collection of n Primordial Quantum Singularities, each with modal Hilbert space restricted to the $(0,1)$-sector:

$$\mathcal{H}_{(0,1)}^{(x_i)} := \operatorname{span}\left\{ |\overset{\circ}{\mathbf{0}}\rangle_{x_i},\ |\overset{\circ}{\mathbf{1}}\rangle_{x_i} \right\} \cong \mathbb{C}^2.$$

The total multipartite Hilbert space is the ordered tensor product:

$$\mathcal{H}^{(n)} := \bigotimes_{i=1}^{n} \mathcal{H}_{(0,1)}^{(x_i)} \cong (\mathbb{C}^2)^{\otimes n}.$$

The *multipartite parasymmetric state* is defined as the equal-weighted superposition of all configurations in which exactly one site occupies the Antisymmetry mode $|\overset{\circ}{\mathbf{1}}\rangle$ and all others occupy the Symmetry mode $|\overset{\circ}{\mathbf{0}}\rangle$:

$$|\Psi_{\text{para}}^{(n)}\rangle := \frac{1}{\sqrt{n}} \sum_{k=1}^{n} |\overset{\circ}{\mathbf{0}}\rangle_{x_1} \otimes \cdots \otimes \underset{k\text{-th}}{|\overset{\circ}{\mathbf{1}}\rangle_{x_k}} \otimes \cdots \otimes |\overset{\circ}{\mathbf{0}}\rangle_{x_n}. \tag{2.28}$$

In conventional notation (with $|\overset{\circ}{\mathbf{0}}\rangle := |e_1\rangle$, $|\overset{\circ}{\mathbf{1}}\rangle := |e_2\rangle$), this becomes:

$$|\Psi_{\text{para}}^{(n)}\rangle = \frac{1}{\sqrt{n}} \sum_{k=1}^{n} |e_1\rangle_{x_1} \otimes \cdots \otimes \underset{k\text{-th}}{|e_2\rangle_{x_k}} \otimes \cdots \otimes |e_1\rangle_{x_n},$$

with each summand corresponding to one excitation (antisymmetric mode) among n modes.

The associated rank-1 *global parasymmetric projector* is:

$$\boxed{P_{\text{para}}^{(x_1 \ldots x_n)} := |\Psi_{\text{para}}^{(n)}\rangle\langle\Psi_{\text{para}}^{(n)}|.} \tag{2.29}$$

Properties

1. **Permutation Symmetry:** The state $|\Psi_{\text{para}}^{(n)}\rangle$ is invariant under any permutation $\sigma \in S_n$ of the site labels:
$$\hat{\Pi}_\sigma |\Psi_{\text{para}}^{(n)}\rangle = |\Psi_{\text{para}}^{(n)}\rangle,$$
where $\hat{\Pi}_\sigma$ acts by permuting tensor factors.
2. **Genuine Multipartite Entanglement:** For $n \geq 3$, the state cannot be written as a product of states over any bipartition of the tensor product space. The entanglement is irreducibly global.
3. **Mixed Marginals:** The reduced density matrix on any proper subset of the n subsystems is mixed. No proper subset retains pure parasymmetric coherence.

Interpretation. This state embodies a generalized form of *dialectical synthesis* extended across many singularities. At any moment, one site exhibits Antisymmetry while all others manifest Symmetry; yet the global configuration is a coherent superposition of all such arrangements. The result is a nonlocal entanglement structure in which opposites are distributed and bound across space—expressing modal unity through relational variation.

From Bipartite to Multipartite Coherence The theoretical framework proceeds with the natural generalization of dialectical synthesis from bipartite systems to configurations of n singularities, thereby transitioning from simple entanglement to the rich phenomenology of many-body quantum coherence. This extension operates within the total Hilbert space $\mathcal{H}^{(n)} := \bigotimes_{i=1}^{n} \mathcal{H}_{(0,1)}^{(x_i)}$, constructed as the tensor product of the individual $(0,1)$-subspaces of n Primordial Quantum Singularities.

The State of Distributed Synthesis The ontological principle underlying the multipartite parasymmetric state is elegantly formulated: it represents the equal-weighted superposition of all configurations in which exactly one site occupies the Antisymmetry mode $|\mathring{\mathbf{1}}\rangle$ while all others occupy the Symmetry mode $|\mathring{\mathbf{0}}\rangle$. The translation of this abstract principle into a concrete quantum state yields:

$$|\Psi_{\text{para}}^{(n)}\rangle = \frac{1}{\sqrt{n}} \sum_{k=1}^{n} |e_1\rangle_{x_1} \otimes \cdots \otimes \underset{k\text{-th}}{|e_2\rangle_{x_k}} \otimes \cdots \otimes |e_1\rangle_{x_n},$$

This is in the realized basis vectors $|e_1\rangle$ and $|e_2\rangle$ within each local $\mathbb{C}^2$ subspace, rather than abstract ontological principles. This state corresponds precisely to the W-state of quantum information theory, characterized by the coherent distribution of a single quantum excitation—here manifested as the Antisymmetry mode—across the entire many-body system. The normalization factor $1/\sqrt{n}$ ensures unit probability for the total state.

Properties and Interpretation The multipartite parasymmetric state exhibits several fundamental properties that illuminate its role within the IQR:

1. **Permutation Symmetry:** The state remains invariant under any permutation of the singularity labels, treating every PQS identically. This democratic distribution ensures that synthesis emerges as a collective property rather than being localized to any privileged singularity.

2. **Genuine Multipartite Entanglement:** For systems of three or more singularities, the entanglement structure is irreducibly holistic, resisting decomposition into any collection of bipartite entangled pairs. The quantum coherence manifests only through the indivisible unity of the complete system.

3. **Mixed Marginals:** Tracing out any subset of singularities yields a reduced density matrix corresponding to a mixed state for the remaining subsystem. The quantum purity and coherence are therefore properties exclusive to the complete configuration, vanishing under any partial observation.

These properties establish the multipartite parasymmetric state as the mathematical instantiation of generalized dialectical synthesis extended across multiple singularities. This construction provides the foundational framework for a universe in which relational properties and non-local correlations constitute the primary mechanisms through which structure emerges and complexity unfolds.

Definition 2.34 (Circuit synthesis of $|\Psi^{(n)}_{\text{para}}\rangle$)**.** Let $(d_{x_1}, \dots, d_{x_n})$ be the duality qubits of the n PQSs, with initial register $|\mathring{\mathbf{0}}\rangle^{\otimes n} = |0\rangle^{\otimes n}$. Define angles

$$\theta_k \;:=\; 2\arccos\sqrt{\frac{n-k}{n-k+1}} \qquad (k = 1, \dots, n-1). \tag{2.30}$$

The following ladder of single–qubit R_y rotations and nearest–neighbour CNOTs prepares the multipartite parasymmetric state of eq. (2.28):



Explanation: start from $|10\cdots0\rangle$ by X on d_{x_1}. For each $k = 1, \dots, n-1$, the $R_y(\theta_k)$ adjusts the amplitude so that the *remaining* excitation weight to be distributed over sites $k, \dots, n$ is $\sqrt{\frac{1}{n-k+1}}$, then CNOT $(d_{x_k} \to d_{x_{k+1}})$ coherently "splits" the single excitation to the next site. Induction on k yields the equal-weight single-excitation superposition

$$|\Psi^{(n)}_{\text{para}}\rangle = \frac{1}{\sqrt{n}} \sum_{j=1}^{n} |0\rangle^{\otimes(j-1)} |1\rangle |0\rangle^{\otimes(n-j)} = \frac{1}{\sqrt{n}} \sum_{j=1}^{n} \Big(\bigotimes_{i \neq j} |\mathring{\mathbf{0}}\rangle_{x_i} \Big) \otimes |\mathring{\mathbf{1}}\rangle_{x_j}.$$

Example 2.35 (Explicit n=4 instance)**.** For $n = 4$, the angles are $\theta_1 = 2\arccos\sqrt{\frac{3}{4}}$, $\theta_2 = 2\arccos\sqrt{\frac{2}{3}}$, $\theta_3 = 2\arccos\sqrt{\frac{1}{2}} = \frac{\pi}{2}$, and the circuit is



which prepares $|\Psi^{(4)}_{\text{para}}\rangle = \frac{1}{2}\big(|1000\rangle + |0100\rangle + |0010\rangle + |0001\rangle \big)$ on $(d_{x_1}, d_{x_2}, d_{x_3}, d_{x_4})$.

Remark 2.36 (Gate class and resources)**.** The synthesis uses Clifford+R_y with $n-1$ controlled-NOTs and $n-1$ single-qubit R_y gates, depth $O(n)$ and nearest-neighbour connectivity (as drawn). Any global reordering of sites preserves the target state by the permutation symmetry of Definition 2.33.

2.2.6 Multipartite Anti-Parasymmetric Entanglement

Definition 2.37 (Multipartite Anti-Parasymmetric State)**.** Let $\{\mathfrak{B}_{x_1}, \ldots, \mathfrak{B}_{x_n}\}$ be a collection of n Primordial Quantum Singularities, each with modal Hilbert space restricted to the $(0,1)$-sector:

$$\mathcal{H}_{(0,1)}^{(x_i)} := \operatorname{span}\left\{ |\mathring{\mathbf{0}}\rangle_{x_i},\ |\mathring{\mathbf{1}}\rangle_{x_i} \right\} \cong \mathbb{C}^2,$$

and total Hilbert space:

$$\mathcal{H}^{(n)} := \bigotimes_{i=1}^{n} \mathcal{H}_{(0,1)}^{(x_i)} \cong (\mathbb{C}^2)^{\otimes n}.$$

Let the *parasymmetric state* be defined as:

$$|\Psi_{\text{para}}^{(n)}\rangle := \frac{1}{\sqrt{n}} \sum_{k=1}^{n} |\phi_k\rangle, \quad \text{where } |\phi_k\rangle := \bigotimes_{i=1}^{n} \chi_i^{(k)}, \quad \chi_i^{(k)} := \begin{cases} |\mathring{\mathbf{1}}\rangle_{x_i} & \text{if } i = k, \\ |\mathring{\mathbf{0}}\rangle_{x_i} & \text{if } i \neq k. \end{cases}$$

We now define the *orthonormalized anti-parasymmetric state*, valid for all n, as:

$$|\widetilde{\Psi}_{\text{antipara}}^{(n)}\rangle := \frac{|\Psi_{\text{antipara}}^{(n)}\rangle - \gamma_n |\Psi_{\text{para}}^{(n)}\rangle}{\sqrt{1-\gamma_n^2}}, \quad \text{where } \gamma_n := \frac{1}{n} \sum_{k=1}^{n} (-1)^{k-1}. \tag{2.31}$$

Here,

$$|\Psi_{\text{antipara}}^{(n)}\rangle := \frac{1}{\sqrt{n}} \sum_{k=1}^{n} (-1)^{k-1} |\phi_k\rangle$$

is the (generally non-orthogonal) alternating single-excitation superposition.

Conventional Quantum Form Using the identifications $|\mathring{\mathbf{0}}\rangle := |e_1\rangle$, $|\mathring{\mathbf{1}}\rangle := |e_2\rangle$, the basis state $|\phi_k\rangle$ becomes:

$$|\phi_k\rangle = |e_1\rangle_{x_1} \otimes \cdots \otimes |e_2\rangle_{x_k} \otimes \cdots \otimes |e_1\rangle_{x_n},$$

so the normalized orthogonal anti-parasymmetric state is:

$$|\widetilde{\Psi}_{\text{antipara}}^{(n)}\rangle = \sum_{k=1}^{n} \left[\frac{(-1)^{k-1} - \gamma_n}{\sqrt{n(1-\gamma_n^2)}} \right] |\phi_k\rangle.$$

Projector The corresponding rank-1 projector is:

$$\boxed{P_{\widetilde{\text{antipara}}}^{(x_1 \ldots x_n)} := |\widetilde{\Psi}_{\text{antipara}}^{(n)}\rangle\langle\widetilde{\Psi}_{\text{antipara}}^{(n)}|.} \tag{2.32}$$

Properties

1. **Universal Orthogonality:**

$$\langle\widetilde{\Psi}_{\text{antipara}}^{(n)}|\Psi_{\text{para}}^{(n)}\rangle = 0 \quad \text{for all } n \in \mathbb{N}.$$

2. **Normalization:**
$$\|\widetilde{\Psi}^{(n)}_{\text{antipara}}\| = 1.$$

3. **Span Complementarity:** The states $|\Psi^{(n)}_{\text{para}}\rangle$ and $|\widetilde{\Psi}^{(n)}_{\text{antipara}}\rangle$ form an orthonormal basis of the excitation sector:
$$\mathcal{H}_{\text{exc}} := \text{span}\,\{|\phi_k\rangle \mid 1 \le k \le n\} \subset \mathcal{H}^{(n)}.$$

Interpretation. The state $|\widetilde{\Psi}^{(n)}_{\text{antipara}}\rangle$ realizes the true logical complement to multipartite parasymmetry. It represents a *canonically orthogonal* configuration of distributed modal dissonance—an antithesis to global synthesis—valid across all cardinalities of Primordial Quantum Singularities. This corrected state ensures structural consistency within the Fourfold Dialectic Algebra of Symmetry without parity restrictions.

Definition 2.38 (Circuit synthesis of $|\widetilde{\Psi}^{(n)}_{\text{antipara}}\rangle$)**.** Let $(d_{x_1}, \ldots, d_{x_n})$ be the duality qubits initialized in $|\mathring{\mathbf{0}}\rangle^{\otimes n} = |0\rangle^{\otimes n}$. Define target coefficients
$$\tilde{c}_k := \frac{(-1)^{k-1} - \gamma_n}{\sqrt{n(1-\gamma_n^2)}}, \qquad \gamma_n = \frac{1}{n}\sum_{j=1}^{n}(-1)^{j-1} = \begin{cases} 0, & n \text{ even}, \\ \frac{1}{n}, & n \text{ odd}, \end{cases}$$
so that $|\widetilde{\Psi}^{(n)}_{\text{antipara}}\rangle = \sum_{k=1}^{n} \tilde{c}_k\, |\phi_k\rangle$ with $|\phi_k\rangle = |0\rangle^{\otimes(k-1)}\,|1\rangle\,|0\rangle^{\otimes(n-k)}$.

Definition 2.39 (Even-n (simplified): W state + alternating Z phases)**.** For n even, $\gamma_n = 0$ and $\tilde{c}_k = \frac{1}{\sqrt{n}}(-1)^{k-1}$. Prepare $|\Psi^{(n)}_{\text{para}}\rangle$ as in Definition 2.34, then apply Z to every even-indexed duality qubit to imprint the alternating sign:



This yields $|\widetilde{\Psi}^{(n)}_{\text{antipara}}\rangle = \frac{1}{\sqrt{n}}\sum_k (-1)^{k-1}\,|\phi_k\rangle$, already orthogonal to $|\Psi^{(n)}_{\text{para}}\rangle$.

Definition 2.40 (Odd-n (general amplitudes): ladder with tail-norm angles + parity phases)**.** For n odd, let $m_k := |\tilde{c}_k|$ and define tail sums $S_k := \sum_{j=k}^{n} m_j^2$ with $S_{n+1} := 0$. Set
$$\theta_k := 2\arccos\sqrt{\frac{S_{k+1}}{S_k}} \qquad (k = 1, \ldots, n-1).$$

The following ladder prepares the magnitudes m_k (starting from $|10\cdots0\rangle$) and then applies

parity phases to set the signs $\mathrm{sgn}(\tilde{c}_k) = (-1)^{k-1}$:



The first ladder ensures amplitudes $\{m_k\}$ via the tail-norm choice of $\{\theta_k\}$; the trailing Z gates on even indices impose the alternating signs, producing $|\widetilde{\Psi}^{(n)}_{\text{antipara}}\rangle = \sum_k \tilde{c}_k \, |\phi_k\rangle$ exactly.

Remark 2.41 (Resources and correctness)**.** Even-n: uses the parasymmetric preparation of Definition 2.34 plus $n/2$ single-qubit Z gates. Odd-n: uses $n-1$ CNOTs, $n-1$ R_y rotations with angles from $\{S_k\}$, and $\lfloor n/2 \rfloor$ single-qubit Z gates. By construction, the state equals the orthonormalized anti-parasymmetric vector of eq. (2.31), i.e., $\langle \widetilde{\Psi}^{(n)}_{\text{antipara}} | \Psi^{(n)}_{\text{para}} \rangle = 0$ and $\| \, |\widetilde{\Psi}^{(n)}_{\text{antipara}}\rangle \| = 1$.

The Gram-Schmidt Orthogonalization The mathematical resolution employs the Gram-Schmidt orthogonalization process—the standard procedure for constructing an orthonormal basis from linearly independent vectors. This approach ensures mathematical rigor while preserving the dialectical structure of the framework.

The construction begins with two fundamental states:

- The parasymmetric (W-state): $|\Psi^{(n)}_{\text{para}}\rangle = \frac{1}{\sqrt{n}} \sum_k |\phi_k\rangle$
- The original anti-parasymmetric state: $|\Psi^{(n)}_{\text{antipara}}\rangle = \frac{1}{\sqrt{n}} \sum_k (-1)^{k-1} |\phi_k\rangle$

The inner product between these states yields the projection coefficient $\gamma_n := \langle \Psi^{(n)}_{\text{para}} | \Psi^{(n)}_{\text{antipara}} \rangle = \frac{1}{n} \sum_k (-1)^{k-1}$, which evaluates to non-zero values for odd n, identifying the source of non-orthogonality in the construction.

The orthogonalization procedure constructs a modified anti-parasymmetric state by removing the component parallel to the parasymmetric state:

$$|\widetilde{\Psi}^{(n)}_{\text{antipara}}\rangle \propto |\Psi^{(n)}_{\text{antipara}}\rangle - \gamma_n |\Psi^{(n)}_{\text{para}}\rangle$$

This construction guarantees orthogonality to $|\Psi^{(n)}_{\text{para}}\rangle$ by design, with the normalization factor $1/\sqrt{1-\gamma_n^2}$ ensuring unit norm for the resulting state.

The Final Form and Properties The complete algebraic expression for the orthogonalized state takes the form:

$$|\widetilde{\Psi}^{(n)}_{\text{antipara}}\rangle = \sum_{k=1}^{n} \left[\frac{(-1)^{k-1} - \gamma_n}{\sqrt{n(1-\gamma_n^2)}} \right] |\phi_k\rangle$$

This construction exhibits the following essential properties:

1. **Universal Orthogonality:** The modified state maintains orthogonality to the parasymmetric state for all values of n, removing the parity-dependent restriction.

2. **Proper Normalization:** The state vector possesses unit norm, satisfying the fundamental requirement of quantum mechanical state vectors.

3. **Span Complementarity:** The parasymmetric state $|\Psi^{(n)}_{\text{para}}\rangle$ and the orthogonalized anti-parasymmetric state $|\widetilde{\Psi}^{(n)}_{\text{antipara}}\rangle$ constitute a complete orthonormal basis for the two-dimensional coherence plane within the single-excitation sector.

The formulation reveals a deeper physical significance: the genuine state of modal dissonance transcends a simple alternating sum, emerging instead as a complex superposition defined relationally with respect to its dialectical opposite. This state represents precisely that configuration which remains after all parasymmetric coherence has been projected away—a mathematical embodiment of pure antithesis.

The orthogonalized state $|\widetilde{\Psi}^{(n)}_{\text{antipara}}\rangle$ thus constitutes the canonically orthogonal configuration of distributed modal dissonance, serving as the true dialectical antithesis to global synthesis. This construction ensures the structural integrity of the Fourfold Dialectic Algebra across all multipartite configurations, eliminating artificial restrictions on system size while preserving the fundamental oppositional structure central to the IQR.

2.2.7 Local and Pairwise Coherence Parasymmetric Generator

Definition 2.42 (Local and Pairwise Coherence Parasymmetric Generator $\hat{F}^{(2)}_{\text{para}}$)**.** Let $\{\mathfrak{B}_{x_1}, \ldots, \mathfrak{B}_{x_n}\}$ be a configuration of n Primordial Quantum Singularities, each with modal subspace

$$\mathcal{H}^{(x_i)}_{(0,1)} := \text{span}\left\{ |\mathring{\mathbf{0}}\rangle_{x_i},\ |\mathring{\mathbf{1}}\rangle_{x_i} \right\} \subset \mathbb{C}^4,$$

and let the global parasymmetric modal Hilbert space be defined as

$$\mathcal{H}^{(n)}_{\text{para}} := \bigotimes_{i=1}^{n} \mathcal{H}^{(x_i)}_{(0,1)} \cong (\mathbb{C}^2)^{\otimes n}.$$

Let $P^{(x_i x_j)}_{\text{para}}$ and $P^{(x_i x_j)}_{\text{antipara}}$ denote the orthogonal rank-1 projectors onto the parasymmetric and anti-parasymmetric two-site entangled states:

$$|\Psi^{(ij)}_{\text{para}}\rangle := \frac{1}{\sqrt{2}} \left(|\mathring{\mathbf{0}}\rangle_{x_i} \otimes |\mathring{\mathbf{1}}\rangle_{x_j} + |\mathring{\mathbf{1}}\rangle_{x_i} \otimes |\mathring{\mathbf{0}}\rangle_{x_j} \right), \tag{2.33}$$

$$|\Psi^{(ij)}_{\text{antipara}}\rangle := \frac{1}{\sqrt{2}} \left(|\mathring{\mathbf{0}}\rangle_{x_i} \otimes |\mathring{\mathbf{1}}\rangle_{x_j} - |\mathring{\mathbf{1}}\rangle_{x_i} \otimes |\mathring{\mathbf{0}}\rangle_{x_j} \right), \tag{2.34}$$

with corresponding projectors:

$$P_{\text{para}}^{(x_i x_j)} := |\Psi_{\text{para}}^{(ij)}\rangle\langle\Psi_{\text{para}}^{(ij)}|, \tag{2.35}$$

$$P_{\text{antipara}}^{(x_i x_j)} := |\Psi_{\text{antipara}}^{(ij)}\rangle\langle\Psi_{\text{antipara}}^{(ij)}|. \tag{2.36}$$

Conventional Quantum State Realization We adopt the identification:

$$|\mathring{\mathbf{0}}\rangle := |e_1\rangle = \begin{pmatrix}1\\0\\0\\0\end{pmatrix}, \quad |\mathring{\mathbf{1}}\rangle := |e_2\rangle = \begin{pmatrix}0\\1\\0\\0\end{pmatrix},$$

which allows us to express the entangled states explicitly in tensor product form.

The *parasymmetric state* becomes:

$$\begin{aligned}|\Psi_{\text{para}}^{(ij)}\rangle &= \frac{1}{\sqrt{2}}\left(\begin{pmatrix}1\\0\\0\\0\end{pmatrix}_{x_i} \otimes \begin{pmatrix}0\\1\\0\\0\end{pmatrix}_{x_j} + \begin{pmatrix}0\\1\\0\\0\end{pmatrix}_{x_i} \otimes \begin{pmatrix}1\\0\\0\\0\end{pmatrix}_{x_j}\right)\\ &= \frac{1}{\sqrt{2}}\left(|e_1\rangle_{x_i} \otimes |e_2\rangle_{x_j} + |e_2\rangle_{x_i} \otimes |e_1\rangle_{x_j}\right),\end{aligned}$$

and the *anti-parasymmetric state* becomes:

$$\begin{aligned}|\Psi_{\text{antipara}}^{(ij)}\rangle &= \frac{1}{\sqrt{2}}\left(\begin{pmatrix}1\\0\\0\\0\end{pmatrix}_{x_i} \otimes \begin{pmatrix}0\\1\\0\\0\end{pmatrix}_{x_j} - \begin{pmatrix}0\\1\\0\\0\end{pmatrix}_{x_i} \otimes \begin{pmatrix}1\\0\\0\\0\end{pmatrix}_{x_j}\right)\\ &= \frac{1}{\sqrt{2}}\left(|e_1\rangle_{x_i} \otimes |e_2\rangle_{x_j} - |e_2\rangle_{x_i} \otimes |e_1\rangle_{x_j}\right).\end{aligned}$$

Definition of the Generator Define the *Local and Pairwise Hermitian Parasymmetric Generator* as:

$$\boxed{\hat{F}_{\text{para}}^{(pair)} := \sum_{i<j} J_{ij}\left(P_{\text{para}}^{(x_i x_j)} - P_{\text{antipara}}^{(x_i x_j)}\right),} \tag{2.37}$$

where $J_{ij} \in \mathbb{R}$ is a symmetric real-valued coupling matrix, satisfying $J_{ij} = J_{ji}$.

Properties

1. **Hermiticity:** Each projector is Hermitian, hence $\hat{F}_{\text{para}}^{(2)}$ is Hermitian.
2. **Projector Orthogonality:**
$$P_{\text{para}}^{(x_i x_j)} \cdot P_{\text{antipara}}^{(x_i x_j)} = 0.$$

3. **Eigenstructure:**

   - Acts nontrivially only on the symmetric–antisymmetric subspace:

$$\mathcal{H}^{(ij)}_{\text{sym/antisym}} := \text{span}\left\{|\Psi^{(ij)}_{\text{para}}\rangle,\ |\Psi^{(ij)}_{\text{antipara}}\rangle\right\} \subset \mathcal{H}^{(x_i)}_{(0,1)} \otimes \mathcal{H}^{(x_j)}_{(0,1)}.$$

   - On this subspace, the operator has eigenvalues $\pm J_{ij}$.

4. **Limitation:** $\hat{F}^{(2)}_{\text{para}}$ does not in general diagonalize the full multipartite parasymmetric states such as:

$$|\Psi^{(n)}_{\text{para}}\rangle := \frac{1}{\sqrt{n}} \sum_{k=1}^{n} |\mathring{\mathbf{0}}\rangle_{x_1} \otimes \cdots \otimes |\mathring{\mathbf{1}}\rangle_{x_k} \otimes \cdots \otimes |\mathring{\mathbf{0}}\rangle_{x_n},$$

   unless $n = 2$.

Interpretation. The generator $\hat{F}^{(pair)}_{\text{para}}$ measures the extent to which each PQS pair (x_i, x_j) is in a state of *modal synthesis* (parasymmetry) or *modal dissonance* (anti-parasymmetry), and enables *coherent transitions* in those modes. It defines a pairwise-interaction-based dynamic on the parasymmetric subspace $\mathcal{H}^{(n)}_{\text{para}}$, without resolving higher-order multipartite structure.

2.2.8 Multipartite Modal-Class Parasymmetric Generator

Definition 2.43 (Global Modal-Class Parasymmetric Generator $\hat{F}^{(n)}_{\text{para}}$)**.** Let $\{\mathfrak{B}_{x_1}, \ldots, \mathfrak{B}_{x_n}\}$ be a configuration of n Primordial Quantum Singularities, each with parasymmetric modal subspace

$$\mathcal{H}^{(x_i)}_{(0,1)} := \text{span}\left\{|\mathring{\mathbf{0}}\rangle_{x_i},\ |\mathring{\mathbf{1}}\rangle_{x_i}\right\} \subset \mathbb{C}^4,$$

and total parasymmetric Hilbert space

$$\mathcal{H}^{(n)}_{\text{para}} := \bigotimes_{i=1}^{n} \mathcal{H}^{(x_i)}_{(0,1)} \cong (\mathbb{C}^2)^{\otimes n}.$$

Ontological Modal-Class States Define the *multipartite parasymmetric state*:

$$|\Psi^{(n)}_{\text{para}}\rangle := \frac{1}{\sqrt{n}} \sum_{k=1}^{n} |\mathring{\mathbf{0}}\rangle_{x_1} \otimes \cdots \otimes \underset{k\text{-th}}{|\mathring{\mathbf{1}}\rangle_{x_k}} \otimes \cdots \otimes |\mathring{\mathbf{0}}\rangle_{x_n}, \tag{2.38}$$

and the anti-parasymmetric complement:

$$|\widetilde{\Psi}^{(n)}_{\text{antipara}}\rangle := \text{normalized projection of } |\Psi^{(n)}_{\text{antipara}}\rangle \text{ orthogonal to } |\Psi^{(n)}_{\text{para}}\rangle. \tag{2.39}$$

Conventional Quantum State Realization Using the identifications:

$$|\overset{\circ}{\mathbf{0}}\rangle := |e_1\rangle = \begin{pmatrix}1\\0\\0\\0\end{pmatrix}, \quad |\overset{\circ}{\mathbf{1}}\rangle := |e_2\rangle = \begin{pmatrix}0\\1\\0\\0\end{pmatrix},$$

we rewrite the multipartite parasymmetric state as:

$$|\Psi^{(n)}_{\text{para}}\rangle = \frac{1}{\sqrt{n}} \sum_{k=1}^{n} |e_1\rangle_{x_1} \otimes \cdots \otimes \underset{k\text{-th}}{|e_2\rangle_{x_k}} \otimes \cdots \otimes |e_1\rangle_{x_n},$$

which lies entirely in the *single-excitation sector* of $(\mathbb{C}^2)^{\otimes n} \subset (\mathbb{C}^4)^{\otimes n}$.

Projectors Define the orthogonal projectors:

$$P^{(x_1 \ldots x_n)}_{\text{para}} := |\Psi^{(n)}_{\text{para}}\rangle\langle\Psi^{(n)}_{\text{para}}|, \tag{2.40}$$

$$P^{\widetilde{(x_1 \ldots x_n)}}_{\text{antipara}} := |\widetilde{\Psi}^{(n)}_{\text{antipara}}\rangle\langle\widetilde{\Psi}^{(n)}_{\text{antipara}}|. \tag{2.41}$$

Generator Definition The global modal-class parasymmetric generator is defined via spectral decomposition:

$$\boxed{\hat{F}^{(n)}_{\text{para}} := E_+ \cdot P^{(x_1 \ldots x_n)}_{\text{para}} + E_- \cdot P^{\widetilde{(x_1 \ldots x_n)}}_{\text{antipara}} + \sum_{\gamma} E_\gamma \cdot P_\gamma,} \tag{2.42}$$

where:

- $E_+, E_- \in \mathbb{R}$ are the eigenvalues of the parasymmetric and anti-parasymmetric multipartite states;
- P_γ are orthogonal projectors completing the spectral decomposition of $\hat{F}^{(n)}_{\text{para}}$;
- $\sum_\gamma P_\gamma = \mathbb{I}_{\text{para}} - P^{(x_1 \ldots x_n)}_{\text{para}} - P^{\widetilde{(x_1 \ldots x_n)}}_{\text{antipara}}$.

Properties

1. **Hermiticity:** All projectors are orthogonal and Hermitian.
2. **Spectral Resolution:**

$$\hat{F}^{(n)}_{\text{para}} = \sum_{\alpha} E_\alpha P_\alpha, \quad P^\dagger_\alpha = P_\alpha, \quad P_\alpha P_\beta = \delta_{\alpha\beta} P_\alpha.$$

3. **Eigenstructure:**

$$\hat{F}^{(n)}_{\text{para}} |\Psi^{(n)}_{\text{para}}\rangle = E_+ |\Psi^{(n)}_{\text{para}}\rangle, \quad \hat{F}^{(n)}_{\text{para}} |\widetilde{\Psi}^{(n)}_{\text{antipara}}\rangle = E_- |\widetilde{\Psi}^{(n)}_{\text{antipara}}\rangle.$$

4. **Modal Sector Invariance:**

$$\hat{F}^{(n)}_{\text{para}} : \mathcal{H}^{(n)}_{\text{para}} \to \mathcal{H}^{(n)}_{\text{para}}.$$

Interpretation. The operator $\hat{F}^{(n)}_{\text{para}}$ defines a global observable that distinguishes between coherent synthesis and structured dissonance across all n singularities. Unlike pairwise interaction terms, this generator respects the total *modal-class entanglement structure* of the system, enabling dynamics governed by the full dialectical configuration of Primordial Quantum Singularities.

Definition 2.44 (Eigenstate "flag" isometries for $P^{(x_1\ldots x_n)}_{\text{para}}$ and $P^{(x_1\ldots x_n)}_{\widetilde{\text{antipara}}}$)**.** Let $(d_{x_1},\ldots,d_{x_n})$ denote the duality qubits. Let V_+ be the state-preparation unitary of Definition 2.34 such that $V_+\,|10\cdots0\rangle = |\Psi^{(n)}_{\text{para}}\rangle$, and let V_- be the unitary of Definition 2.38 such that $V_-\,|10\cdots0\rangle = |\widetilde{\Psi}^{(n)}_{\text{antipara}}\rangle$. Introduce single-qubit ancillae a_+, a_- initialized in $|0\rangle$ and define the flagged projectors

$$\text{Flag}_+ := V_+\left(\prod_{j=2}^{n}\text{CTRL}_{d_{x_j}=0}\right)\text{CTRL}_{d_{x_1}=1}\,\text{X}_{a_+}\,V_+^\dagger,$$
$$\text{Flag}_- := V_-\left(\prod_{j=2}^{n}\text{CTRL}_{d_{x_j}=0}\right)\text{CTRL}_{d_{x_1}=1}\,\text{X}_{a_-}\,V_-^\dagger.$$

i.e. multi-controlled flips of $a_\pm$ conditioned on the pattern $d_{x_1}=1$, $d_{x_2}=\cdots=d_{x_n}=0$ *after* uncomputing to the preparation seeds via $V_\pm^\dagger$. These realize coherent membership tests for the rank-1 projectors $P^{(x_1\ldots x_n)}_{\text{para}}$ and $P^{(x_1\ldots x_n)}_{\widetilde{\text{antipara}}}$.



(Replace V_+ by V_- and a_+ by a_- for Flag_-.)

Definition 2.45 (Time evolution $U_{\text{para}}(t) = e^{-it\,\hat{F}^{(n)}_{\text{para}}}$ via phase kickback)**.** Given the spectral form in eq. (2.42),

$$\hat{F}^{(n)}_{\text{para}} = E_+\,P^{(x_1\ldots x_n)}_{\text{para}} + E_-\,P^{(x_1\ldots x_n)}_{\widetilde{\text{antipara}}} + \sum_{\gamma} E_\gamma P_\gamma,$$

the unitary $U_{\text{para}}(t) = e^{-it\hat{F}^{(n)}_{\text{para}}}$ is implemented on the encoded subspace by sequential

conditional phases using the flags from Definition 2.44:



Here each $R_Z(2E_\pm t)$ applies a phase only when the corresponding ancilla is in $|1\rangle$, yielding a conditional phase $e^{-itE_\pm}$ on the targeted eigenspace. The P_γ sector can be assigned a uniform offset $E_\perp$ (global phase) or further resolved by additional flagged projectors if available.

Remark 2.46 (Measurement of $\hat{F}^{(n)}_{\text{para}}$)**.** Expectation values $\langle \hat{F}^{(n)}_{\text{para}} \rangle_\rho$ can be obtained by a Hadamard test with a single control ancilla c and the controlled version of $U_{\text{para}}(t)$ from Definition 2.45; sweeping t enables phase estimation on the two-dimensional eigen-subspace $\text{span}\{|\Psi^{(n)}_{\text{para}}\rangle, |\widetilde{\Psi}^{(n)}_{\text{antipara}}\rangle\}$, while leaving the orthogonal complement invariant up to a known phase.

Remark 2.47 (Gate class and resources)**.** The construction uses Clifford unitaries ($V_\pm$ built from CNOTs and R_y rotations) plus multi-controlled X/Z and single-qubit R_Z phases. Depth is $O(n)$ for the $V_\pm$ ladders (nearest-neighbour layout), plus the cost of an n-controlled Toffoli for flagging (log-depth with ancillae using standard decomposition). Since the projector flags commute, the two phase-kickback segments can be applied in any order.

2.2.9 Multipartite Modal-Class Anti-Parasymmetric Generator

Definition 2.48 (Orthogonalized Anti-Parasymmetric Modal-Class State)**.** Let

$$\{\mathfrak{B}_{x_1}, \ldots, \mathfrak{B}_{x_n}\}$$

be a configuration of n Primordial Quantum Singularities, each with modal subspace

$$\mathcal{H}^{(x_i)}_{(0,1)} := \text{span}\left\{ |\mathring{\mathbf{0}}\rangle_{x_i},\ |\mathring{\mathbf{1}}\rangle_{x_i} \right\} \subset \mathbb{C}^4,$$

and global parasymmetric modal Hilbert space

$$\mathcal{H}^{(n)}_{\text{para}} := \bigotimes_{i=1}^{n} \mathcal{H}^{(x_i)}_{(0,1)} \cong (\mathbb{C}^2)^{\otimes n}.$$

Conventional Quantum State Realization Recall the identifications:

$$|\mathring{\mathbf{0}}\rangle := |e_1\rangle = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \quad |\mathring{\mathbf{1}}\rangle := |e_2\rangle = \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix}.$$

The *parasymmetric W-state* is given by:

$$|\Psi^{(n)}_{\text{para}}\rangle := \frac{1}{\sqrt{n}} \sum_{k=1}^{n} |\mathring{\mathbf{0}}\rangle_{x_1} \otimes \cdots \otimes \underset{k\text{-th}}{|\mathring{\mathbf{1}}\rangle_{x_k}} \otimes \cdots \otimes |\mathring{\mathbf{0}}\rangle_{x_n}. \tag{2.43}$$

Then the *parasymmetric W-state* becomes:

$$|\Psi^{(n)}_{\text{para}}\rangle := \frac{1}{\sqrt{n}} \sum_{k=1}^{n} |e_1\rangle_{x_1} \otimes \cdots \otimes \underset{k\text{-th}}{|e_2\rangle_{x_k}} \otimes \cdots \otimes |e_1\rangle_{x_n}, \tag{2.44}$$

i.e., it is the uniform superposition over all configurations in which exactly one site occupies $|e_2\rangle$ (the antisymmetric mode) and all others occupy $|e_1\rangle$ (the symmetric mode).

The *unadjusted anti-parasymmetric configuration* is:

$$|\Psi^{(n)}_{\text{antipara}}\rangle := \frac{1}{\sqrt{n}} \sum_{k=1}^{n} (-1)^{k-1} |\mathring{\mathbf{0}}\rangle_{x_1} \otimes \cdots \otimes \underset{k\text{-th}}{|\mathring{\mathbf{1}}\rangle_{x_k}} \otimes \cdots \otimes |\mathring{\mathbf{0}}\rangle_{x_n}. \tag{2.45}$$

Likewise, the *unadjusted anti-parasymmetric configuration* becomes:

$$|\Psi^{(n)}_{\text{antipara}}\rangle := \frac{1}{\sqrt{n}} \sum_{k=1}^{n} (-1)^{k-1} |e_1\rangle_{x_1} \otimes \cdots \otimes \underset{k\text{-th}}{|e_2\rangle_{x_k}} \otimes \cdots \otimes |e_1\rangle_{x_n}, \tag{2.46}$$

in which each term in the superposition acquires an alternating sign $(-1)^{k-1}$, encoding structured modal dissonance across the configuration.

These are both elements of the single-excitation subspace:

$$\mathcal{H}^{(n)}_{\text{exc}} := \text{span} \left\{ |e_2\rangle_{x_k} \otimes \bigotimes_{i \neq k} |e_1\rangle_{x_i} \;\middle|\; k = 1, \ldots, n \right\} \subset (\mathbb{C}^2)^{\otimes n}.$$

Then the *orthogonalized anti-parasymmetric modal-class state* is defined as the normalized component of $|\Psi^{(n)}_{\text{antipara}}\rangle$ orthogonal to $|\Psi^{(n)}_{\text{para}}\rangle$:

$$|\widetilde{\Psi}^{(n)}_{\text{antipara}}\rangle := \frac{1}{\sqrt{1 - |\langle \Psi^{(n)}_{\text{para}} | \Psi^{(n)}_{\text{antipara}} \rangle|^2}} \left(|\Psi^{(n)}_{\text{antipara}}\rangle - \langle \Psi^{(n)}_{\text{para}} | \Psi^{(n)}_{\text{antipara}} \rangle \cdot |\Psi^{(n)}_{\text{para}}\rangle \right). \tag{2.47}$$

This state satisfies:

$$\langle \Psi^{(n)}_{\text{para}} | \widetilde{\Psi}^{(n)}_{\text{antipara}} \rangle = 0, \tag{2.48}$$

$$\| \widetilde{\Psi}^{(n)}_{\text{antipara}} \| = 1. \tag{2.49}$$

Interpretation. This orthogonalized state represents a genuine modal-class dissonance: a configuration of global anti-synthesis that is orthogonal to the W-state's global synthesis. It resolves the ambiguity in the case of odd n, where the original $|\Psi^{(n)}_{\text{antipara}}\rangle$ is not orthogonal to the parasymmetric mode.

Definition 2.49 (Global modal-class anti-parasymmetric generator)**.** Let $P^{(x_1\ldots x_n)}_{\widetilde{\text{antipara}}} := |\widetilde{\Psi}^{(n)}_{\text{antipara}}\rangle\langle\widetilde{\Psi}^{(n)}_{\text{antipara}}|$, where $|\widetilde{\Psi}^{(n)}_{\text{antipara}}\rangle$ is the orthogonalized anti-parasymmetric state from Definition 2.48, and let $\mathbb{I}_{\text{exc}} := \sum_{k=1}^{n} |\phi_k\rangle\langle\phi_k|$ project onto the single-excitation sector. Define the generator (up to a global phase) by the spectral form

$$\hat{F}^{(n)}_{\widetilde{\text{antipara}}} := E_- \, P^{(x_1\ldots x_n)}_{\widetilde{\text{antipara}}} + E_\perp \, (\mathbb{I}_{\text{exc}} - P^{(x_1\ldots x_n)}_{\widetilde{\text{antipara}}}), \tag{2.50}$$

so that $|\widetilde{\Psi}^{(n)}_{\text{antipara}}\rangle$ is an eigenstate with eigenvalue E_- and any orthogonal single-excitation vector has eigenvalue $E_\perp$. The corresponding evolution is

$$U_-(t) := e^{-it\hat{F}^{(n)}_{\widetilde{\text{antipara}}}} = e^{-itE_\perp}\Big[\, \mathbb{I}_{\text{exc}} + \big(e^{-it(E_- - E_\perp)} - 1\big) P^{(x_1\ldots x_n)}_{\widetilde{\text{antipara}}} \,\Big].$$

Definition 2.50 (Flag isometry for $P^{(x_1\ldots x_n)}_{\widetilde{\text{antipara}}}$)**.** Let V_- be the state-preparation unitary that maps the seed $|10\cdots0\rangle$ to $|\widetilde{\Psi}^{(n)}_{\text{antipara}}\rangle$ (see the synthesis circuit in Definition 2.38). With an ancilla a_- initialized in $|0\rangle$, define the coherent flag

$$\text{Flag}_- \;:=\; V_- \Big(\text{CTRL}_{d_{x_1}=1} \prod_{j=2}^{n} \text{CTRL}_{d_{x_j}=0} \Big) \, X_{a_-} \, V_-^\dagger,$$

i.e. uncompute to the seed, flip a_- iff the pattern $d_{x_1}{=}1$ and $d_{x_2}{=}\cdots{=}d_{x_n}{=}0$ holds, then re-compute. This realizes a coherent membership test for $P^{(x_1\ldots x_n)}_{\widetilde{\text{antipara}}}$.



Definition 2.51 (Phase-kickback implementation of $U_-(t)$)**.** Using Definition 2.50, implement $U_-(t)$ (up to the global phase $e^{-itE_\perp}$) by:



The R_Z on a_- imparts phase $e^{-it(E_- - E_\perp)}$ only when the system is in the flagged subspace $|\widetilde{\Psi}^{(n)}_{\text{antipara}}\rangle$, producing the desired evolution in eq. (2.50).

Remark 2.52 (Two-eigenspace resolution (optional))**.** If a distinct phase is also required on the parasymmetric eigenstate, add a second flag Flag_+ (cf. Definition 2.34) and apply an $R_Z(2(E_+ - E_\perp)t)$ on a second ancilla a_+ between Flag_+ and $\mathrm{Flag}_+^\dagger$. Since the flags commute (orthogonal projectors), the two segments may be executed in any order.

Remark 2.53 (Resources)**.** The construction uses the state-preparation unitary V_- (depth $O(n)$ with $n-1$ CNOTs and $n-1$ rotations), one n-controlled Toffoli (log-depth with ancillae via standard decompositions), and a single-qubit R_Z rotation for the phase.

2.2.10 The β Generalized Decoupled Unitary Evolution

Definition 2.54 (β Generalized Decoupled Unitary Parasymmetric Operator)**.** Let $\mathcal{X}_q^{(n)} = \{\mathfrak{B}_{x_1}, \ldots, \mathfrak{B}_{x_n}\}$ be a configuration of n Primordial Quantum Singularities (PQSs), with parasymmetric modal subspace:

$$\mathcal{H}_{\text{para}}^{(n)} := \bigotimes_{i=1}^{n} \mathcal{H}_{(0,1)}^{(x_i)} \cong (\mathbb{C}^2)^{\otimes n}.$$

Let $\mathcal{P} = \{X_1, X_2, \ldots, X_k\}$ be a partition of the PQS configuration into disjoint blocks:

$$X_\beta \subseteq \mathcal{X}_q^{(n)}, \quad X_\beta \cap X_{\beta'} = \emptyset \text{ for } \beta \neq \beta', \quad \bigcup_\beta X_\beta = \mathcal{X}_q^{(n)}.$$

To each block X_β, assign a Hermitian generator $\hat{F}_{\text{para}}^{(\beta)}$ acting on the parasymmetric modal space:

$$\hat{F}_{\text{para}}^{(\beta)} : \mathcal{H}_{\text{para}}^{(X_\beta)} \to \mathcal{H}_{\text{para}}^{(X_\beta)},$$

where $\mathcal{H}_{\text{para}}^{(X_\beta)} := \bigotimes_{x_i \in X_\beta} \mathcal{H}_{(0,1)}^{(x_i)} \cong (\mathbb{C}^2)^{\otimes |X_\beta|}$.

Each block evolves unitarily via:

$$U_\beta := e^{i\hat{F}_{\text{para}}^{(\beta)}}.$$

Then the *Generalized Decoupled Unitary Parasymmetric Operator* is defined as the tensor product:

$$\boxed{U_N^{(\beta)} := \bigotimes_{\beta \in \mathcal{P}} U_\beta = \bigotimes_{\beta \in \mathcal{P}} e^{i\hat{F}_{\text{para}}^{(\beta)}}.} \tag{2.51}$$

Interpretation. This operator describes a general class of *factorized evolutions*, in which each block X_β undergoes autonomous parasymmetric evolution under an internal coherence structure (local, pairwise, or global), while maintaining strict *modal independence* from the other blocks.

Special Case: Active Zone Model If $\mathcal{P} = \{X_{\text{dyn}}, X_{\text{stat}}\}$ with:

$$\hat{F}^{(X_{\text{stat}})}_{\text{para}} = 0 \quad \Rightarrow \quad U_{X_{\text{stat}}} = \mathbb{I},$$

then U_N reduces to the *Generalized Non-Coupled Operator*:

$$U_N^{(\text{gen})} = \mathbb{I}_{X_{\text{stat}}} \otimes e^{i\hat{F}_{\text{gen}}|_{X_{\text{dyn}}}}.$$

Definition 2.55 (Blockwise synthesis of $U_\beta = e^{i\hat{F}^{(\beta)}_{\text{para}}}$ via flagged phases)**.** For a block $X_\beta = \{x_{i_1}, \ldots, x_{i_{m_\beta}}\}$ with duality qubits $\mathbf{d}^{(\beta)} := (d_{x_{i_1}}, \ldots, d_{x_{i_{m_\beta}}})$, let

$$\hat{F}^{(\beta)}_{\text{para}} = E^{(\beta)}_+ P^{(X_\beta)}_{\text{para}} + E^{(\beta)}_- P^{(X_\beta)}_{\underbrace{\phantom{\text{antipara}}}_{\text{antipara}}} + \sum_\gamma E^{(\beta)}_\gamma P^{(\beta)}_\gamma.$$

Let $V^{(\beta)}_+$ and $V^{(\beta)}_-$ be the state-preparation unitaries that map a seed $|10\cdots0\rangle$ on $\mathbf{d}^{(\beta)}$ to $|\Psi^{(m_\beta)}_{\text{para}}\rangle$ and $|\widetilde{\Psi}^{(m_\beta)}_{\text{antipara}}\rangle$, respectively (cf. the multipartite synthesis circuits). Define coherent flags using ancillae a^+_β, a^-_β:

$$\text{Flag}^{(\beta)}_+ := V^{(\beta)}_+ \Big(\text{CTRL}_{d_{x_{i_1}}=1} \prod_{j=2}^{m_\beta} \text{CTRL}_{d_{x_{i_j}}=0} \Big) X_{a^+_\beta} V^{(\beta)\dagger}_+,$$

$$\text{Flag}^{(\beta)}_- := V^{(\beta)}_- \Big(\text{CTRL}_{d_{x_{i_1}}=1} \prod_{j=2}^{m_\beta} \text{CTRL}_{d_{x_{i_j}}=0} \Big) X_{a^-_\beta} V^{(\beta)\dagger}_-.$$

Then $U_\beta = e^{i\hat{F}^{(\beta)}_{\text{para}}}$ (up to a global phase on the complementary subspace) is implemented by phase kickback:



Additional spectral components $P^{(\beta)}_\gamma$ are handled by appending analogous flag–phase–uncompute segments with their own ancillae and angles $-2E^{(\beta)}_\gamma$.

Definition 2.56 (Parallel composition: $U^{(\beta)}_N = \bigotimes_{\beta\in\mathcal{P}} U_\beta$)**.** Let $\mathcal{P} = \{X_\beta\}$ be a partition of sites. Since blocks act on disjoint registers, the implementations of U_β commute and may run concurrently. Schematic (two blocks shown, generalization is immediate):



Thus the full gate is realized as a tensor product of the concurrently executed block circuits, in accordance with eq. (2.51).

Remark 2.57 (Active Zone Model (special case))**.** If $X_{\text{stat}} \in \mathcal{P}$ satisfies $\hat{F}_{\text{para}}^{(X_{\text{stat}})} = 0$, the corresponding block circuit is omitted (identity on $\mathbf{d}^{(\text{stat})}$), and only the dynamic block(s) X_{dyn} execute the flag–phase–uncompute sequence, yielding $U_N^{(\text{gen})} = \mathbb{I}_{X_{\text{stat}}} \otimes e^{i\hat{F}_{\text{gen}}|_{X_{\text{dyn}}}}$.

Remark 2.58 (Resources)**.** Depth equals $\max_\beta\{\text{depth}(V_\pm^{(\beta)}) + \text{depth}(n_\beta\text{-controlled Toffoli})\}$, where $n_\beta = |X_\beta|$. Gate count scales additively across blocks; parallel scheduling across β minimizes wall–clock depth due to block independence.

2.2.11 The Zonal Decoupled Unitary Evolution

Definition 2.59 (Zonal Decoupled Unitary Parasymmetric Operator)**.** Let $\mathcal{X}_q^{(n)} = X_{\text{stat}} \cup X_{\text{dyn}}$, with $X_{\text{stat}} \cap X_{\text{dyn}} = \emptyset$, be a bipartition of an n-PQS configuration.

Define:

$$\mathcal{H}_{\text{para}}^{(n)} := \bigotimes_{i=1}^{n} \mathcal{H}_{(0,1)}^{(x_i)} \cong (\mathbb{C}^2)^{\otimes n},$$

$$\mathcal{H}_{\text{stat}} := \bigotimes_{x_i \in X_{\text{stat}}} \mathcal{H}_{(0,1)}^{(x_i)}, \quad \mathcal{H}_{\text{dyn}} := \bigotimes_{x_j \in X_{\text{dyn}}} \mathcal{H}_{(0,1)}^{(x_j)}.$$

Let the generalized parasymmetric generator over X_{dyn} be:

$$\hat{F}_{\text{gen}}|_{X_{\text{dyn}}} := \lambda_L \hat{F}_{\text{para}}^{(\text{local})}|_{X_{\text{dyn}}} + \lambda_P \hat{F}_{\text{para}}^{(\text{pair})}|_{X_{\text{dyn}}} + \lambda_G \hat{F}_{\text{para}}^{(n)}|_{X_{\text{dyn}}},$$

where $\lambda_L, \lambda_P, \lambda_G \in \mathbb{R}$ are coupling constants.

Then the *Zonal Decoupled Unitary Parasymmetric Operator* is:

$$\boxed{U_N^{(\text{zonal})} := \mathbb{I}_{X_{\text{stat}}} \otimes e^{i\hat{F}_{\text{gen}}|_{X_{\text{dyn}}}}.} \tag{2.52}$$

Properties

1. **Partial Evolution:** Only the PQSs in X_{dyn} evolve; those in X_{stat} remain static.
2. **Modal Factorization:** The total operator factorizes:
$$U_N^{(\text{zonal})} \in \mathcal{U}(\mathcal{H}_{\text{stat}}) \otimes \mathcal{U}(\mathcal{H}_{\text{dyn}}).$$
3. **Internal Coherence Richness:** Dynamics within X_{dyn} can include local, pairwise, or global coherence modes depending on the structure of $\hat{F}_{\text{gen}}$.

Interpretation. This operator models a localized parasymmetric dynamic of high structural complexity while maintaining full decoupling from the external modal domain. It formalizes *modular evolution* in the IQR.

Definition 2.60 (Circuit synthesis of $U_N^{(\mathrm{zonal})} = \mathbb{I}_{X_{\mathrm{stat}}} \otimes e^{i\hat{F}_{\mathrm{gen}}|_{X_{\mathrm{dyn}}}}$)**.** Let

$$\hat{F}_{\mathrm{gen}}|_{X_{\mathrm{dyn}}} := \lambda_L\, \hat{F}^{(\mathrm{local})}_{\mathrm{para}}|_{X_{\mathrm{dyn}}} + \lambda_P\, \hat{F}^{(\mathrm{pair})}_{\mathrm{para}}|_{X_{\mathrm{dyn}}} + \lambda_G\, \hat{F}^{(n)}_{\mathrm{para}}|_{X_{\mathrm{dyn}}}.$$

The overall circuit leaves all d_x with $x \in X_{\mathrm{stat}}$ as bare wires (identity), and acts only on d_x with $x \in X_{\mathrm{dyn}}$. When the three terms do not commute, use a Trotter–Suzuki product with $r \in \mathbb{N}$ steps:

$$U_N^{(\mathrm{zonal})} \;\approx\; \mathbb{I}_{X_{\mathrm{stat}}} \otimes \left[e^{i\frac{\lambda_L}{r}\hat{F}^{(\mathrm{local})}_{\mathrm{para}}}\; e^{i\frac{\lambda_P}{r}\hat{F}^{(\mathrm{pair})}_{\mathrm{para}}}\; e^{i\frac{\lambda_G}{r}\hat{F}^{(n)}_{\mathrm{para}}} \right]^r,$$

or the symmetric second-order variant by inserting the terms in palindromic order.

Definition 2.61 (Local segment $e^{i\frac{\lambda_L}{r}\hat{F}^{(\mathrm{local})}_{\mathrm{para}}}$)**.** For each $x \in X_{\mathrm{dyn}}$, write the local contribution as $h_x\,(P^{(x)}_{\mathrm{para}} - P^{(x)}_{\mathrm{antipara}}) = h_x\, X_{d_x}$ on $\mathcal{H}^{(x)}_{(0,1)}$. Implement $e^{i\frac{\lambda_L}{r}h_x X}$ as a one-qubit rotation

$$e^{i\frac{\lambda_L}{r}h_x X} \;=\; R_x\Big(-2\tfrac{\lambda_L}{r}h_x\Big),$$

applied in parallel to all d_x in X_{dyn}:



$$\alpha_x := \tfrac{\lambda_L}{r}h_x.$$

Definition 2.62 (Pair segment $e^{i\frac{\lambda_P}{r}\hat{F}^{(\mathrm{pair})}_{\mathrm{para}}}$)**.** For each interacting pair $(x_i, x_j) \subset X_{\mathrm{dyn}}$ with coupling J_{ij}, the two-qubit contribution $J_{ij}\Big(P^{(x_i x_j)}_{\mathrm{para}} - P^{(x_i x_j)}_{\mathrm{antipara}}\Big)$ acts as $+1$ on $|\Psi^+\rangle$ and -1 on $|\Psi^-\rangle$. Using the inverse Bell map $(\mathrm{CNOT}(i \to j);\, H_i)$ that sends $|\Psi^\pm\rangle \mapsto |1\,0/1\rangle$, implement

$$e^{i\theta_{ij}\,(P_{\Psi^+} - P_{\Psi^-})}, \qquad \theta_{ij} := \tfrac{\lambda_P}{r}J_{ij},$$

by a controlled-R_z on d_{x_j} conditioned on d_{x_i}=1, flanked by the Bell uncompute/compute:



Apply these two-qubit segments for all (i, j) in the chosen interaction set (scheduling to respect hardware connectivity if needed).

Definition 2.63 (Global segment $e^{i\frac{\lambda_G}{r}\hat{F}^{(n)}_{\mathrm{para}}\Big|_{X_{\mathrm{dyn}}}}$)**.** Let $m := |X_{\mathrm{dyn}}|$, and let V_+ and V_- be the m-qubit state-preparation unitaries mapping a fixed seed to $|\Psi^{(m)}_{\mathrm{para}}\rangle$ and $|\widetilde{\Psi}^{(m)}_{\mathrm{antipara}}\rangle$, respectively.

With ancillae a_+, a_- initialized to $|0\rangle$, define coherent flags

$$\mathrm{Flag}_+ := V_+\Big(\,\mathrm{CTRL}_{d_{x_1}=1}\prod_{\ell=2}^{m}\mathrm{CTRL}_{d_{x_\ell}=0}\Big)\,X_{a_+}\,V_+^\dagger,$$
$$\mathrm{Flag}_- := V_-\Big(\,\mathrm{CTRL}_{d_{x_1}=1}\prod_{\ell=2}^{m}\mathrm{CTRL}_{d_{x_\ell}=0}\Big)\,X_{a_-}\,V_-^\dagger.$$

Assume that $\hat{F}^{(n)}_{\mathrm{para}}\big|_{X_{\mathrm{dyn}}}$ has eigenvalues E_+, E_- on these two rank-1 subspaces, and $E_\perp$ on the orthogonal complement. Apply phase-kickback with angles scaled by λ_G/r:



(An overall phase $e^{\,i(\lambda_G/r)E_\perp}$ on the orthogonal subspace may be ignored or realised similarly if needed.)

Definition 2.64 (One Trotter step and repetition)**.** A single step for X_{dyn} concatenates Definitions 2.61 to 2.63:

$$\boxed{\mathcal{S}_{\mathrm{dyn}} \;:=\; e^{\,i\frac{\lambda_L}{r}\hat{F}^{(\mathrm{local})}_{\mathrm{para}}}\;e^{\,i\frac{\lambda_P}{r}\hat{F}^{(\mathrm{pair})}_{\mathrm{para}}}\;e^{\,i\frac{\lambda_G}{r}\hat{F}^{(n)}_{\mathrm{para}}}}\quad\Rightarrow\quad U_N^{(\mathrm{zonal})}\approx\mathbb{I}_{X_{\mathrm{stat}}}\otimes\Big(\mathcal{S}_{\mathrm{dyn}}\Big)^r.$$

Static-zone wires (all d_x with $x\in X_{\mathrm{stat}}$) remain identity throughout:

Remark 2.65 (Resources)**.** Per Trotter step: (i) $|X_{\mathrm{dyn}}|$ single-qubit R_x gates (local); (ii) one Bell-uncompute + one CR_z + one Bell-recompute per interacting pair (pair term); (iii) two $|X_{\mathrm{dyn}}|$-controlled flags and two single-qubit R_z rotations (global). Depth scales as $O(|X_{\mathrm{dyn}}|)$ with nearest-neighbour scheduling; static-zone depth is zero by construction.

2.2.12 The Generalized Composite Decoupled Parasymmetric Evolution Operator

Definition 2.66 (Generalized Composite Parasymmetric Evolution Operator)**.** Let $\mathcal{X}_q^{(n)} = \{\mathfrak{B}_{x_1},\ldots,\mathfrak{B}_{x_n}\}$ be a configuration of n Primordial Quantum Singularities, with parasymmetric modal subspace:

$$\mathcal{H}^{(n)}_{\mathrm{para}} := \bigotimes_{i=1}^{n}\mathcal{H}^{(x_i)}_{(0,1)} \cong (\mathbb{C}^2)^{\otimes n}.$$

Let $\mathcal{P} = \{X_L, X_P, X_G\}$ be a disjoint partition of the configuration into blocks corresponding to:

- X_L: sites evolving under **local** coherence dynamics,
- X_P: sites evolving under **pairwise** parasymmetric entanglement,
- X_G: sites evolving under **global modal-class** coherence dynamics.

Assign to each block a Hermitian parasymmetric generator:
(i) Local Coherence Generator:

$$\hat{F}^{(L)}_{\text{para}} := \sum_{x_i \in X_L} \left(P^{(x_i)}_{\text{para}} - P^{(x_i)}_{\text{antipara}} \right).$$

(ii) Pairwise Entanglement Generator:

$$\hat{F}^{(P)}_{\text{para}} := \sum_{i<j \in X_P} J_{ij} \left(P^{(x_i x_j)}_{\text{para}} - P^{(x_i x_j)}_{\text{antipara}} \right), \quad J_{ij} = J_{ji} \in \mathbb{R}.$$

(iii) Global Modal-Class Generator:

$$\hat{F}^{(G)}_{\text{para}} := E_+ P^{(X_G)}_{\text{para}} + E_- P^{(X_G)}_{\widetilde{\text{antipara}}} + \sum_{\gamma} E_\gamma P_\gamma.$$

Then the corresponding blockwise unitary evolutions are:

$$U_L := e^{i\hat{F}^{(L)}_{\text{para}}}, \quad U_P := e^{i\hat{F}^{(P)}_{\text{para}}}, \quad U_G := e^{i\hat{F}^{(G)}_{\text{para}}}.$$

The *Generalized Composite Parasymmetric Evolution Operator* is then defined as:

$$\boxed{U^{(\odot)}_N := U_L \otimes U_P \otimes U_G = \bigotimes_{\beta \in \{L,P,G\}} e^{i\hat{F}^{(\beta)}_{\text{para}}}.} \tag{2.53}$$

Interpretation. This operator unifies local, pairwise, and global parasymmetric dynamics within a single decoupled configuration. Each subregion evolves autonomously under its assigned coherence structure, yet all contribute to the overall parasymmetric evolution of the system.

Definition 2.67 (Blockwise circuits for $U^{(\odot)}_N = U_L \otimes U_P \otimes U_G$)**.** Each factor acts on disjoint registers and can be executed in parallel.

(i) Local coherence block $U_L = e^{i\hat{F}^{(L)}_{\text{para}}}$ Using $P^{(x)}_{\text{para}} = \frac{1}{2}(\mathbb{I} + X_{d_x})$ and $P^{(x)}_{\text{antipara}} = \frac{1}{2}(\mathbb{I} - X_{d_x})$ on $\mathcal{H}^{(x)}_{(0,1)}$,

$$\hat{F}^{(L)}_{\text{para}} = \sum_{x \in X_L} \left(P^{(x)}_{\text{para}} - P^{(x)}_{\text{antipara}} \right) = \sum_{x \in X_L} X_{d_x} \quad \Rightarrow \quad U_L = \prod_{x \in X_L} e^{iX_{d_x}} = \prod_{x \in X_L} R_x(-2).$$

Circuit (all rotations in parallel):

$\{d_x : x \in X_L\}$ —$\boxed{R_x(-2)}$—

(ii) Pairwise entanglement block $U_P = e^{i\hat{F}^{(P)}_{\text{para}}}$ For each $(i,j) \in X_P$ with coupling J_{ij}, $\hat{F}^{(P)}_{\text{para}} \ni J_{ij}\left(P^{(x_i x_j)}_{\text{para}} - P^{(x_i x_j)}_{\text{antipara}}\right)$, which phases $|\Psi^+\rangle$ by $+J_{ij}$ and $|\Psi^-\rangle$ by $-J_{ij}$. Implement via inverse Bell map (CNOT $i \to j$, then H on i), a controlled R_z, and recompute:



Apply one such segment per interacting pair (schedule per hardware connectivity).

(iii) Global modal-class block $U_G = e^{i\hat{F}^{(G)}_{\text{para}}}$ Let $m := |X_G|$ and let V_+, V_- be the m-qubit preparation unitaries for $|\Psi^{(m)}_{\text{para}}\rangle$ and $|\widetilde{\Psi}^{(m)}_{\text{antipara}}\rangle$ (see multipartite synthesis circuits). With ancillae a_+, a_-:

$$\begin{aligned}
\text{Flag}_+ &:= V_+\Big(\text{CTRL}_{d_{x_1}=1} \prod_{\ell=2}^{m} \text{CTRL}_{d_{x_\ell}=0}\Big) \quad X_{a_+}\, V_+^\dagger, \\
\text{Flag}_- &:= V_-\Big(\text{CTRL}_{d_{x_1}=1} \prod_{\ell=2}^{m} \text{CTRL}_{d_{x_\ell}=0}\Big) \quad X_{a_-}\, V_-^\dagger.
\end{aligned}$$

If

$$\hat{F}^{(G)}_{\text{para}} = E_+ P^{(X_G)}_{\text{para}} + E_- P^{(X_G)}_{\widetilde{\text{antipara}}} + \sum_\gamma E_\gamma P_\gamma$$

use phase kickback on the flags (extend with additional flags for the P_γ terms if needed):



Parallel composition Since the three blocks act on disjoint registers $\mathbf{d}^{(L)}$, $\mathbf{d}^{(P)}$, $\mathbf{d}^{(G)}$, their circuits commute and can run concurrently, realizing exactly

$$U_N^{(\odot)} \;=\; U_L \otimes U_P \otimes U_G \quad \text{as in eq. (2.53)}.$$

A schematic layout:



Remark 2.68 (Resources)**.** Local: $|X_L|$ single-qubit R_x gates (depth 1 in parallel). Pairwise: per interacting pair, one CNOT–H uncompute, one controlled R_z, and one recompute (depth $O(1)$ per pair; schedule by edge coloring). Global: two $|X_G|$-controlled flags plus two single-qubit R_z rotations (log-depth with ancillae using standard decompositions). Overall depth is the maximum of the three block depths due to parallel execution.

Remark 2.69 (Operational effect of the composite parasymmetric evolution)**.** The unitary in eq. (2.53), $U_N^{(\odot)} = U_L \otimes U_P \otimes U_G$, acts independently on three disjoint modal blocks and therefore factors its action across registers:

$$\begin{aligned} U_L &= \exp\Big(i \sum_{x\in X_L} \Big[P^{(x)}_{\text{para}} - P^{(x)}_{\text{antipara}}\Big]\Big), \\ U_P &= \exp\Big(i \sum_{\substack{i<j \\ i,j\in X_P}} J_{ij} \Big[P^{(x_i x_j)}_{\text{para}} - P^{(x_i x_j)}_{\text{antipara}}\Big]\Big), \\ U_G &= \exp\Big(i\hat{F}^{(G)}_{\text{para}}\Big). \end{aligned}$$

- **Local block U_L (single-site coherence).** On each duality qubit d_x with $x \in X_L$, $P^{(x)}_{\text{para}} - P^{(x)}_{\text{antipara}} = X_{d_x}$ in the $(0,1)$-sector, hence

$$U_L = \prod_{x\in X_L} e^{iX_{d_x}} = \prod_{x\in X_L} R_x(-2).$$

  It performs an x-axis rotation that coherently mixes $|\mathring{\mathbf{0}}\rangle_x$ and $|\mathring{\mathbf{1}}\rangle_x$ without leaving $\mathcal{H}^{(x)}_{(0,1)}$.

- **Pair block U_P (two-site exchange/entanglement).** Using $P^{(x_i x_j)}_{\text{para}} - P^{(x_i x_j)}_{\text{antipara}} = \frac{1}{2}\big(X_{d_{x_i}} X_{d_{x_j}} + Y_{d_{x_i}} Y_{d_{x_j}}\big)$, each pair evolution is an *XY-exchange* (partial iSWAP–type) gate:

$$U_{ij} := \exp\Big(i\tfrac{J_{ij}}{2}(X_iX_j + Y_iY_j)\Big), \quad \begin{aligned} U_{ij}\,|01\rangle &= \cos\tfrac{J_{ij}}{2}\;|01\rangle + i\sin\tfrac{J_{ij}}{2}\;|10\rangle\,, \\ U_{ij}\,|10\rangle &= \cos\tfrac{J_{ij}}{2}\;|10\rangle + i\sin\tfrac{J_{ij}}{2}\;|01\rangle\,, \\ U_{ij}\,|00\rangle &= |00\rangle\,, \quad U_{ij}\,|11\rangle = |11\rangle\,, \end{aligned}$$

  equivalently phasing the Bell states as $|\Psi^{\pm}\rangle \mapsto e^{\pm iJ_{ij}}\,|\Psi^{\pm}\rangle$ while leaving $|\Phi^{\pm}\rangle$ invariant.

- **Global block U_G (modal-class phases).** On the $m := |X_G|$-site duality register,

$$\hat{F}^{(G)}_{\text{para}} = E_+ P^{(X_G)}_{\text{para}} + E_- P^{(X_G)}_{\widetilde{\text{antipara}}} + \sum_\gamma E_\gamma P_\gamma,$$

  so U_G is block-diagonal: it applies eigenphases e^{iE_+} and e^{iE_-} to the two rank-1 subspaces $\text{span}\{|\Psi^{(m)}_{\text{para}}\rangle\}$ and $\text{span}\{|\widetilde{\Psi}^{(m)}_{\text{antipara}}\rangle\}$, and e^{iE_γ} on each orthogonal component P_γ.

In summary: $U_N^{(\odot)}$ performs (i) local x-axis rotations on X_L; (ii) XY-exchange entangling evolutions on specified pairs in X_P; and (iii) modal-class phase rotations on the distinguished multipartite subspace in X_G. Because the blocks act on disjoint tensor factors, these transformations commute and the total evolution is the tensor product of the three independent processes.

2.2.13 The $\hat{\sigma}$ Global Coupled Unitary Evolution

Definition 2.70 (Global Coupled Unitary Parasymmetric Operator)**.** Suppose that there are n Primordial Quantum Singularities $\mathfrak{B}_{x_1}, \ldots, \mathfrak{B}_{x_n}$, with parasymmetric modal subspace:

$$\mathcal{H}^{(n)}_{\text{para}} := \bigotimes_{i=1}^{n} \mathcal{H}^{(x_i)}_{(0,1)} \cong (\mathbb{C}^2)^{\otimes n}.$$

Define the Hermitian parasymmetric generator:

$$\hat{F}_{\text{para}} := \sum_{i,j=1}^{n} J_{ij} \left(\hat{\sigma}^{(+)}_{x_i} \hat{\sigma}^{(-)}_{x_j} + \hat{\sigma}^{(+)}_{x_j} \hat{\sigma}^{(-)}_{x_i} \right), \quad J_{ij} = J_{ji} \in \mathbb{R},$$

where $\hat{\sigma}^{(+)}_{x_i}$ and $\hat{\sigma}^{(-)}_{x_i}$ are the modal raising and lowering operators acting on $\mathcal{H}^{(x_i)}_{(0,1)}$.

Then the $\hat{\sigma}$ *Global Coupled Unitary Parasymmetric Operator* is defined as:

$$\boxed{U^{(\hat{\sigma})}_{G} := e^{i\hat{F}_{\text{para}}}.} \tag{2.54}$$

Properties

1. **Entangling Evolution:** The operator U_G enables coherent interactions among all PQSs in $\mathcal{X}^{(n)}_q$, without blockwise separability.
2. **Graph-Theoretic Coupling:** The interaction pattern is determined by the matrix J_{ij}, which encodes a weighted undirected graph of modal connectivity.
3. **Non-Factorizability:** Unlike the decoupled cases, U_G generally cannot be written as a tensor product of operators acting on strict subregions.

Interpretation. This operator governs fully global modal evolution, wherein parasymmetric coherence may propagate, interfere, and self-organize across the entire configuration space. It instantiates the synthetic logic of Parasymmetry—fusing local modes into dynamically entangled dialectical wholes.

Definition 2.71 (Exchange form of the generator)**.** Using $\hat{\sigma}^{(+)} = \frac{1}{2}(X + iY)$ and $\hat{\sigma}^{(-)} = \frac{1}{2}(X - iY)$ on each duality qubit,

$$\hat{\sigma}^{(+)}_{x_i} \hat{\sigma}^{(-)}_{x_j} + \hat{\sigma}^{(+)}_{x_j} \hat{\sigma}^{(-)}_{x_i} \;=\; \tfrac{1}{2}\Big(X_{d_{x_i}} X_{d_{x_j}} + Y_{d_{x_i}} Y_{d_{x_j}} \Big).$$

Absorbing any summation symmetry into the couplings, write

$$\hat{F}_{\text{para}} = \sum_{i<j} \theta_{ij} \, \frac{X_{d_{x_i}} X_{d_{x_j}} + Y_{d_{x_i}} Y_{d_{x_j}}}{2}, \qquad U^{(\hat{\sigma})}_{G} = e^{\,i\hat{F}_{\text{para}}} = \prod_{i<j} \exp\Big(i \, \tfrac{\theta_{ij}}{2} \, (X_i X_j + Y_i Y_j) \Big),$$

where θ_{ij} equals J_{ij} up to a possible factor of 2 depending on the double-sum convention.

Definition 2.72 (Primitive XY–exchange gate (iSWAP$^\alpha$ form))**.** For a pair (i, j) define

$$U_{ij}(\alpha) \;:=\; \exp\!\Big(i\,\tfrac{\alpha}{2}\,(X_iX_j + Y_iY_j)\Big) \quad\Longleftrightarrow\quad \begin{aligned} U_{ij}(\alpha)\,|01\rangle &= \cos\tfrac{\alpha}{2}\;|01\rangle + i\sin\tfrac{\alpha}{2}\;|10\rangle\,,\\ U_{ij}(\alpha)\,|10\rangle &= \cos\tfrac{\alpha}{2}\;|10\rangle + i\sin\tfrac{\alpha}{2}\;|01\rangle\,,\\ U_{ij}(\alpha)\,|00\rangle &= |00\rangle\,, \quad U_{ij}(\alpha)\,|11\rangle = |11\rangle\,. \end{aligned}$$

A convenient decomposition uses an inverse Bell map (CNOT then H) to diagonalize the $\{|\Psi^\pm\rangle\}$ subspace, apply a controlled phase, and recompute:



This realizes $U_{ij}(\alpha)$ since $|\Psi^\pm\rangle$ pick phases $e^{\pm i\alpha}$ and $|\Phi^\pm\rangle$ remain invariant.

Definition 2.73 (Trotterized synthesis for arbitrary coupling graph)**.** Let $G = (V, E)$ with $V = \{x_1, \ldots, x_n\}$ and weighted edges $(i, j) \in E$ carrying angles $\alpha_{ij} = \theta_{ij}$. A first-order Trotter step at time slice $1/r$ is

$$\mathcal{S} \;:=\; \prod_{(i,j)\in E} U_{ij}\Big(\tfrac{\alpha_{ij}}{r}\Big), \qquad U_G^{(\hat\sigma)} \;\approx\; \big(\mathcal{S}\big)^r.$$

To parallelize, color edges into disjoint matchings $E = E_1 \cup \cdots \cup E_c$ and apply all U_{ij} in each E_ℓ simultaneously:

$$\mathcal{S} \;=\; \prod_{\ell=1}^{c}\; \prod_{(i,j)\in E_\ell}^{\text{in parallel}} U_{ij}\Big(\tfrac{\alpha_{ij}}{r}\Big).$$

Each U_{ij} uses the two–CNOT pattern of Definition 2.72.

Remark 2.74 (Nearest-neighbour chains and all-to-all cases)**.** For a 1D chain with uniform θ, depth per Trotter step is 2 (odd bonds in parallel, then even bonds), each bond using two CNOTs and one R_z. For dense all-to-all J_{ij}, use a SWAP network to expose each pair once per layer; total two-qubit depth $O(n)$ per Trotter step with the same U_{ij} primitive.

Definition 2.75 (Global coupled parasymmetric circuit (schematic one-step))**.** One Trotter step for a chain $x_1 - x_2 - \cdots - x_n$:



(Continue the odd-bond layer, then the even-bond layer, using the same two–CNOT + R_z sandwich per adjacent pair.)

Remark 2.76 (Consistency with eq. (2.54))**.** Combining Definitions 2.71 to 2.73 yields a concrete, hardware-ready compilation of $U_G^{(\hat{\sigma})} = \exp\Big(i \sum_{i,j} J_{ij}[\hat{\sigma}_i^{(+)}\hat{\sigma}_j^{(-)} + \hat{\sigma}_j^{(+)}\hat{\sigma}_i^{(-)}]\Big)$, with angles α_{ij} chosen to absorb any factor-of-two from the summation convention.

Remark 2.77 (Operational effect of the global coupled parasymmetric gate)**.** The unitary $U_G^{(\hat{\sigma})} = \exp\Big(i\hat{F}_{\text{para}}\Big)$ with

$$\hat{F}_{\text{para}} = \sum_{i,j} J_{ij}\Big(\hat{\sigma}_{x_i}^{(+)}\hat{\sigma}_{x_j}^{(-)} + \hat{\sigma}_{x_j}^{(+)}\hat{\sigma}_{x_i}^{(-)}\Big) = \sum_{i<j} \frac{\theta_{ij}}{2}\Big(X_iX_j + Y_iY_j\Big)$$

is an XY–exchange evolution that *coherently transports* the single "antisymmetry excitation" across the coupling graph defined by $J = (J_{ij})$, while preserving the total excitation number. Let

$$\hat{N} := \sum_{k=1}^{n} \frac{\mathbb{I} - Z_k}{2} \quad \Rightarrow \quad [\hat{N}, \hat{F}_{\text{para}}] = 0,$$

so $U_G^{(\hat{\sigma})}$ acts block-diagonally on the Hamming-weight sectors.

Pair action (two sites). On a coupled pair (i, j),

$$U_{ij}(\alpha) := \exp\Big(i\frac{\alpha}{2}(X_iX_j + Y_iY_j)\Big): \quad \begin{aligned} |01\rangle &\mapsto \cos\tfrac{\alpha}{2}\,|01\rangle + i\sin\tfrac{\alpha}{2}\,|10\rangle\,, \\ |10\rangle &\mapsto \cos\tfrac{\alpha}{2}\,|10\rangle + i\sin\tfrac{\alpha}{2}\,|01\rangle\,, \\ |00\rangle &\mapsto |00\rangle\,, \quad |11\rangle \mapsto |11\rangle\,. \end{aligned}$$

Equivalently, in the Bell basis it applies phases to the exchange states $|\Psi^{\pm}\rangle = \frac{1}{\sqrt{2}}(|01\rangle \pm |10\rangle)$:

$$|\Psi^{\pm}\rangle \mapsto e^{\pm i\alpha}\,|\Psi^{\pm}\rangle\,, \qquad |\Phi^{\pm}\rangle \text{ invariant.}$$

Thus it is an iSWAP$^{\alpha}$-type entangling gate (iSWAP at $\alpha = \frac{\pi}{2}$, SWAP up to phases at $\alpha = \pi$).

Single-excitation sector (global action). Let $\{\,|\phi_k\rangle\,\}_{k=1}^{n}$ denote the basis with a sole antisymmetry at site x_k (i.e. $|\phi_k\rangle = |0\rangle^{\otimes(k-1)}\,|1\rangle\,|0\rangle^{\otimes(n-k)}$ in the duality register). Restricted to this sector,

$$U_G^{(\hat{\sigma})}(t)\;|\phi_k\rangle = \sum_{j=1}^{n} \Big[e^{\,itJ}\Big]_{jk}\;|\phi_j\rangle\,,$$

so the excitation executes a continuous-time quantum walk on the weighted graph with adjacency J. Amplitudes *hop* along edges with rates set by J_{ij}, generating long-range parasymmetric coherence.

Multiple excitations. On higher Hamming-weight sectors, $U_G^{(\hat{\sigma})}$ enacts number-conserving XY dynamics: excitations exchange their positions via the same pairwise mechanism, producing genuinely multipartite entanglement while never leaving the parasymmetric modal subspace.

Summary. $U_G^{(\hat{\sigma})}$ is a globally coupled, excitation-preserving exchange evolution that: (i) mixes $|01\rangle \leftrightarrow |10\rangle$ coherently on each coupled edge; (ii) realizes graph-wide transport and interference of parasymmetric amplitude in the single-excitation manifold; and (iii) yields entangling XY dynamics in multi-excitation manifolds, all determined by the coupling matrix J.

2.2.14 The Global Composite λ Coupled Unitary Evolution

Definition 2.78 (Global Composite Coupled Unitary Parasymmetric Operator)**.** Let $\mathcal{X}_q^{(n)} = \{\mathfrak{B}_{x_1}, \ldots, \mathfrak{B}_{x_n}\}$ be a configuration of n Primordial Quantum Singularities (PQSs), and define the parasymmetric modal Hilbert space:

$$\mathcal{H}_{\text{para}}^{(n)} := \bigotimes_{i=1}^{n} \mathcal{H}_{(0,1)}^{(x_i)} \cong (\mathbb{C}^2)^{\otimes n}.$$

Let the total Hermitian parasymmetric generator be a weighted combination of local, pairwise, and global terms:

$$\hat{F}_{\text{para}} := \lambda_L \hat{F}_{\text{para}}^{(\text{local})} + \lambda_P \hat{F}_{\text{para}}^{(\text{pair})} + \lambda_G \hat{F}_{\text{para}}^{(n)},$$

where:

- $\hat{F}_{\text{para}}^{(\text{local})} = \sum_i \tau_z^{(x_i)}$, the local coherence generator;
- $\hat{F}_{\text{para}}^{(\text{pair})} = \sum_{i<j} J_{ij}(P_{\text{para}}^{(x_i x_j)} - P_{\text{antipara}}^{(x_i x_j)})$, the pairwise entanglement generator;
- $\hat{F}_{\text{para}}^{(n)} = E_+ P_{\text{para}}^{(x_1 \ldots x_n)} + E_- P_{\underbrace{}_{\text{antipara}}}^{(x_1 \ldots x_n)} + \sum_\gamma E_\gamma P_\gamma$, the global modal-class generator;
- and $\lambda_L, \lambda_P, \lambda_G \in \mathbb{R}$ are tunable real parameters.

Then the λ *Global Coupled Unitary Parasymmetric Operator* is defined as:

$$\boxed{U_G^{(\lambda)} := \exp\left(i\left[\lambda_L \hat{F}_{\text{para}}^{(\text{local})} + \lambda_P \hat{F}_{\text{para}}^{(\text{pair})} + \lambda_G \hat{F}_{\text{para}}^{(n)}\right]\right).} \tag{2.55}$$

Interpretation. This operator generates a coherent evolution across the entire PQS configuration, unifying:

- **Local parasymmetry** at each site,
- **Pairwise entanglement** between PQSs,
- **Global modal synthesis/dissonance** through modal-class structure.

It represents the most general *entanglement-generating and coherence-synthesizing* evolution allowed within the parasymmetric sector.

Definition 2.79 (One Trotter step for $U_G^{(\lambda)}$)**.** For non-commuting local/pair/global terms, use a first-order Trotter product with $r \in \mathbb{N}$ steps:

$$U_G^{(\lambda)} \;\approx\; \left[e^{i\frac{\lambda_L}{r}\hat{F}^{(\text{local})}_{\text{para}}} \; e^{i\frac{\lambda_P}{r}\hat{F}^{(\text{pair})}_{\text{para}}} \; e^{i\frac{\lambda_G}{r}\hat{F}^{(n)}_{\text{para}}} \right]^r,$$

or the symmetric second-order formula by palindromic ordering. One Trotter step decomposes into the three segments below.

Definition 2.80 (Local segment $e^{i\frac{\lambda_L}{r}\hat{F}^{(\text{local})}_{\text{para}}}$)**.** With $\hat{F}^{(\text{local})}_{\text{para}} = \sum_x \tau_z^{(x)}$ and $\tau_z^{(x)} \equiv Z_{d_x}$ on $\mathcal{H}^{(x)}_{(0,1)}$,

$$e^{i\frac{\lambda_L}{r}\hat{F}^{(\text{local})}_{\text{para}}} = \prod_x e^{i\frac{\lambda_L}{r}Z_{d_x}} = \prod_x R_z\Big(-2\tfrac{\lambda_L}{r}\Big)_{d_x}.$$

Parallel single-qubit R_z rotations on all duality qubits:



Definition 2.81 (Pairwise segment $e^{i\frac{\lambda_P}{r}\hat{F}^{(\text{pair})}_{\text{para}}}$)**.** Using $P^{(x_i x_j)}_{\text{para}} - P^{(x_i x_j)}_{\text{antipara}} = \frac{1}{2}(X_iX_j + Y_iY_j)$ on duality qubits (d_{x_i}, d_{x_j}), each coupled pair (i,j) evolves by the XY exchange

$$U_{ij}\Big(\tfrac{\lambda_P}{r}J_{ij}\Big) = \exp\Big(i\,\tfrac{\lambda_P}{r}\,\tfrac{J_{ij}}{2}(X_iX_j + Y_iY_j)\Big).$$

Decompose via Bell uncompute (CNOT, then H), a controlled R_z, and recompute:



Apply one such block for every interacting (i,j); schedule by edge-coloring to parallelize disjoint pairs.

Definition 2.82 (Global segment $e^{i\frac{\lambda_G}{r}\hat{F}^{(n)}_{\text{para}}}$)**.** Let V_+ and V_- prepare the rank-1 eigenstates $|\Psi^{(n)}_{\text{para}}\rangle$ and $|\widetilde{\Psi}^{(n)}_{\text{antipara}}\rangle$ from a fixed seed (cf. Definitions 2.33 and 2.48). With ancillae a_+, a_- initialized in $|0\rangle$, define coherent flags

$$\text{Flag}_+ := V_+\Big(\,\text{CTRL}_{d_{x_1}=1}\prod_{\ell=2}^{n}\text{CTRL}_{d_{x_\ell}=0}\Big)\,X_{a_+}\,V_+^\dagger,$$

$$\text{Flag}_- := V_-\Big(\,\text{CTRL}_{d_{x_1}=1}\prod_{\ell=2}^{n}\text{CTRL}_{d_{x_\ell}=0}\Big)\,X_{a_-}\,V_-^\dagger.$$

Phase-kickback implements the eigenphases of $\hat{F}^{(n)}_{\text{para}} = E_+ P^{(x_1 \dots x_n)}_{\text{para}} + E_- P^{(x_1 \dots x_n)}_{\widetilde{\text{antipara}}} + \sum_\gamma E_\gamma P_\gamma$:



(Additional one-dimensional P_γ sectors, if used, are handled by analogous flag–phase–uncompute blocks.)

Definition 2.83 (One composite step and repetition)**.** A single Trotter step is the concatenation

$$\boxed{\mathcal{S}_\lambda \;:=\; e^{i\frac{\lambda_L}{r}\hat{F}^{(\text{local})}_{\text{para}}}\; e^{i\frac{\lambda_P}{r}\hat{F}^{(\text{pair})}_{\text{para}}}\; e^{i\frac{\lambda_G}{r}\hat{F}^{(n)}_{\text{para}}}} \quad \Rightarrow \quad U_G^{(\lambda)} \;\approx\; \Big(\mathcal{S}_\lambda\Big)^r,$$

with a symmetric palindromic ordering available for second-order accuracy. Local R_z gates run fully in parallel; pair segments parallelize over disjoint pairs; the two global flags commute (orthogonal projectors), so their phase segments may be swapped.

Remark 2.84 (Resource sketch)**.** Per Trotter step: n single-qubit R_z (local); for each coupled pair, two CNOTs + one R_z (pair); and two n-controlled flags plus two ancilla R_z (global), with optional extra flags for $\{P_\gamma\}$. Depth scales with the edge-coloring number of the pair graph and with the decomposition depth of the multi-controlled Toffoli used in the flags.

2.2.15 The Unified Global Coupled Unitary Evolution

Definition 2.85 (Unified Global Coupled Parasymmetric Operator)**.** Suppose that

$$\mathcal{X}_q^{(n)} = \{\mathfrak{B}_{x_1}, \dots, \mathfrak{B}_{x_n}\}$$

is a configuration of n Primordial Quantum Singularities (PQSs), with parasymmetric modal subspace:

$$\mathcal{H}^{(n)}_{\text{para}} := \bigotimes_{i=1}^{n} \mathcal{H}^{(x_i)}_{(0,1)} \cong (\mathbb{C}^2)^{\otimes n}.$$

Define the *unified Hermitian parasymmetric generator* as:

$$\hat{F}^{(\odot)}_{\text{para}} := \lambda_L \hat{F}^{(\text{local})}_{\text{para}} + \lambda_P \hat{F}^{(\text{pair})}_{\text{para}} + \lambda_G \hat{F}^{(n)}_{\text{para}} + \lambda_X \sum_{i,j=1}^{n} J_{ij} \left(\hat{\sigma}^{(+)}_{x_i}\hat{\sigma}^{(-)}_{x_j} + \hat{\sigma}^{(+)}_{x_j}\hat{\sigma}^{(-)}_{x_i}\right),$$

where:

- $\hat{F}^{(\text{local})}_{\text{para}}$ is the local coherence generator;
- $\hat{F}^{(\text{pair})}_{\text{para}}$ is the pairwise parasymmetric projector generator;
- $\hat{F}^{(n)}_{\text{para}}$ is the global modal-class generator;
- $\hat{\sigma}^{(\pm)}_{x_i}$ are the modal ladder operators on site x_i;
- $J_{ij} \in \mathbb{R}$, with $J_{ij} = J_{ji}$, defines a symmetric coupling matrix;
- $\lambda_L, \lambda_P, \lambda_G, \lambda_X \in \mathbb{R}$ are tunable coupling parameters.

Then the *Unified Global Coupled Parasymmetric Operator* is defined as:

$$\boxed{U_G^{(\odot)} := \exp\left(i\hat{F}^{(\odot)}_{\text{para}}\right).} \tag{2.56}$$

Interpretation. This operator generates the most general parasymmetric unitary evolution permitted within the framework, simultaneously realizing:

- Local synthesis/dissonance per PQS,
- Pairwise entangled coherence via projectors,
- Multipartite modal-class transitions,
- Dynamical excitation exchange between sites.

It merges modal-logic transformations with physical mode interaction to encode the full dialectical dynamics of parasymmetry.

Definition 2.86 (One Trotter step for $U_G^{(\odot)} = \exp\left(i\hat{F}^{(\odot)}_{\text{para}}\right)$)**.** For generally non-commuting terms, use a r-step first-order Trotterization:

$$U_G^{(\odot)} \approx \left[e^{i\frac{\lambda_L}{r}\hat{F}^{(\text{local})}_{\text{para}}}\, e^{i\frac{\lambda_P}{r}\hat{F}^{(\text{pair})}_{\text{para}}}\, e^{i\frac{\lambda_G}{r}\hat{F}^{(n)}_{\text{para}}}\, e^{i\frac{\lambda_X}{r}\sum_{i,j}J_{ij}\left(\hat{\sigma}_i^{(+)}\hat{\sigma}_j^{(-)}+\hat{\sigma}_j^{(+)}\hat{\sigma}_i^{(-)}\right)}\right]^r,$$

or the symmetric second-order palindromic variant. One step decomposes into the four segments below.

Definition 2.87 (Local segment $e^{i\frac{\lambda_L}{r}\hat{F}^{(\text{local})}_{\text{para}}}$)**.** With $\hat{F}^{(\text{local})}_{\text{para}} = \sum_i \tau_z^{(x_i)}$ and $\tau_z^{(x_i)} \equiv Z_{d_{x_i}}$ on $\mathcal{H}^{(x_i)}_{(0,1)}$,

$$e^{i\frac{\lambda_L}{r}\hat{F}^{(\text{local})}_{\text{para}}} = \prod_i e^{i\frac{\lambda_L}{r}Z_{d_{x_i}}} = \prod_i R_z\Big(-2\tfrac{\lambda_L}{r}\Big)_{d_{x_i}}.$$

All duality qubits rotate in parallel:



Definition 2.88 (Pairwise segment $e^{i\frac{\lambda_P}{r}\hat{F}^{(\mathrm{pair})}_{\mathrm{para}}}$)**.** Using $P^{(x_ix_j)}_{\mathrm{para}} - P^{(x_ix_j)}_{\mathrm{antipara}} = \frac{1}{2}(X_iX_j + Y_iY_j)$ on duality qubits (d_{x_i}, d_{x_j}), each coupled pair (i,j) evolves by

$$U^{(P)}_{ij} \;:=\; \exp\!\Big(i\,\tfrac{\lambda_P}{r}\,\tfrac{J_{ij}}{2}(X_iX_j + Y_iY_j)\Big).$$

Decompose via inverse-Bell uncompute ($\mathrm{CNOT}(i\to j)$; H_i), controlled R_z, and recompute:



Schedule disjoint pairs in parallel (edge-coloring of the interaction graph).

Definition 2.89 (Global segment $e^{i\frac{\lambda_G}{r}\hat{F}^{(n)}_{\mathrm{para}}}$)**.** Let V_+ and V_- be the n-qubit state-preparation unitaries for $|\Psi^{(n)}_{\mathrm{para}}\rangle$ and $|\widetilde{\Psi}^{(n)}_{\mathrm{antipara}}\rangle$. Define coherent flags with ancillae a_+, a_-:

$$\mathrm{Flag}_+ := V_+\Big(\mathrm{CTRL}_{d_{x_1}=1}\prod_{\ell=2}^{n}\mathrm{CTRL}_{d_{x_\ell}=0}\Big)\,X_{a_+}\,V_+^\dagger,$$

$$\mathrm{Flag}_- := V_-\Big(\mathrm{CTRL}_{d_{x_1}=1}\prod_{\ell=2}^{n}\mathrm{CTRL}_{d_{x_\ell}=0}\Big)\,X_{a_-}\,V_-^\dagger.$$

Phase-kickback realizes eigenphases of $\hat{F}^{(n)}_{\mathrm{para}} = E_+P^{(x_1\ldots x_n)}_{\mathrm{para}} + E_-P^{\overbrace{(x_1\ldots x_n)}}_{\mathrm{antipara}} + \sum_\gamma E_\gamma P_\gamma$:



(Extend with additional flags for any P_γ terms if needed.)

Definition 2.90 (Exchange segment $e^{i\frac{\lambda_X}{r}\sum_{i,j}J_{ij}(\hat{\sigma}^{(+)}_i\hat{\sigma}^{(-)}_j+\hat{\sigma}^{(+)}_j\hat{\sigma}^{(-)}_i)}$)**.** Since $\hat{\sigma}^{(\pm)} = \frac{1}{2}(X \pm iY)$, each pair contributes the same XY–exchange primitive as in Definition 2.88:

$$U^{(X)}_{ij} \;:=\; \exp\!\Big(i\,\tfrac{\lambda_X}{r}\,\tfrac{J_{ij}}{2}(X_iX_j + Y_iY_j)\Big),$$

with the identical two–CNOT + R_z decomposition:



Remark 2.91 (Merging identical XY contributions)**.** Because both Definition 2.88 and Definition 2.90 implement $\exp\Big(i\,\frac{\alpha_{ij}}{2}(X_iX_j + Y_iY_j)\Big)$ on each pair, they can be *combined* per Trotter slice by using a single block with effective angle

$$\alpha_{ij}^{\text{eff}} \;=\; \tfrac{1}{r}\,(\lambda_P + \lambda_X)\,J_{ij},$$

thereby halving the two–CNOT overhead per pair.

Definition 2.92 (One unified step and repetition)**.** A single step is the concatenation

$$\boxed{\mathcal{S}_\odot \;:=\; e^{\,i\frac{\lambda_L}{r}\hat{F}_{\text{para}}^{(\text{local})}}\Big[\prod_{(i,j)\in E} U_{ij}^{(P)}\Big]\,e^{\,i\frac{\lambda_G}{r}\hat{F}_{\text{para}}^{(n)}}\Big[\prod_{(i,j)\in E} U_{ij}^{(X)}\Big]} \quad\Rightarrow\quad U_G^{(\odot)} \;\approx\; \big(\mathcal{S}_\odot\big)^r,$$

where E is the edge set of the coupling graph induced by $\{J_{ij}\}$. Parallelize the pair/exchange layers over disjoint matchings; the two global flags commute (orthogonal projectors).

Remark 2.93 (Resource sketch)**.** Per Trotter step: n parallel single-qubit R_z (local); per coupled pair, two CNOTs + one R_z (XY) — or one such block with angle α_{ij}^{eff} if merged; and two n-controlled flags plus two ancilla R_z for the global modal-class segment (extend for P_γ as needed). Depth per step scales with the edge-coloring number of E and the decomposition depth of the multi-controlled Toffoli used in the flags.

2.2.16 Definition: Parasymmetry (Synthesis)

Definition 2.94 (Parasymmetry (Synthesis): Compact Formal Definition)**.** Fix the distinguished $(0,1)$ modal sector at a PQS $\mathfrak{B}_x$ with basis $\{|e_1\rangle_x, |e_2\rangle_x\}$ corresponding to Symmetry/Antisymmetry (cf. Definition 1.16 and Section 2.1). Define the normalized parasymmetric/anti-parasymmetric kets

$$|+\rangle_x := \tfrac{1}{\sqrt{2}}\Big(|e_1\rangle_x + |e_2\rangle_x\Big), \qquad |-\rangle_x := \tfrac{1}{\sqrt{2}}\Big(|e_1\rangle_x - |e_2\rangle_x\Big),$$

and the local effects

$$P_{\text{para}}^{(x)} := |+\rangle_x\langle+|, \qquad P_{\text{antipara}}^{(x)} := |-\rangle_x\langle-|, \tag{2.57}$$

which are orthogonal projectors with $P_{\text{para}}^{(x)} + P_{\text{antipara}}^{(x)} = P_{(0,1)}^{(x)}$ (the $(0,1)$-sector projector). The Born-valued synthesis truth in state ρ is $v_{\text{Synthesis}}(\rho) = \text{Tr}\Big(P_{\text{para}}^{(x)}\rho\Big)$ (Definition 1.18). Measuring the complementary observable (Σ_x on the logical qubit) operationalizes this test.

(P1) Multipartite parasymmetric coherence. For PQS sites $x_1, \ldots, x_n$, define bipartite and global effects that witness *distributed synthesis*:

$$\text{(pair)}\quad |\Psi_{\text{para}}^{(ij)}\rangle := \tfrac{1}{\sqrt{2}}\Big(|e_1\rangle_{x_i}|e_2\rangle_{x_j} + |e_2\rangle_{x_i}|e_1\rangle_{x_j}\Big), \quad P_{\text{para}}^{(x_ix_j)} := |\Psi_{\text{para}}^{(ij)}\rangle\langle\Psi_{\text{para}}^{(ij)}|, \tag{2.58}$$

$$\text{(global)}\quad |\Psi_{\text{para}}^{(n)}\rangle := \tfrac{1}{\sqrt{n}}\sum_{k=1}^{n} |e_1\rangle_{x_1}\cdots|e_2\rangle_{x_k}\cdots|e_1\rangle_{x_n}, \qquad P_{\text{para}}^{(x_1\ldots x_n)} := |\Psi_{\text{para}}^{(n)}\rangle\langle\Psi_{\text{para}}^{(n)}|. \tag{2.59}$$

These states (i) are permutation-symmetric, (ii) exhibit genuine multipartite entanglement with mixed single-site marginals, and (iii) reduce locally to the $(0,1)$ sector. They realize Synthesis as *relation*: no single site carries the whole pattern (cf. Section 2.2.3 and eq. (2.29)).

(P2) Parasymmetry-preserving dynamics. A unitary U on the n-site $(0,1)$ sector $\mathcal{H}_{(0,1)}^{\otimes n}$ is *parasymmetry-preserving* if and only if it leaves the parasymmetric subspace invariant:

$$U\, P_{\text{para}}^{(x_1\ldots x_n)}\, U^\dagger = P_{\text{para}}^{(x_1\ldots x_n)}. \tag{2.60}$$

Equivalently, U maps the span of the states in (2.58)–(2.59) to itself. This class includes:

(i) Decoupled evolutions.
Decoupled evolutions $U = \bigotimes_i U_i$ with each U_i a rotation about the parasymmetric axis (preserving $\{|+\rangle_x, |-\rangle_x\}$).

(ii) Coupled evolutions.
Coupled evolutions generated by exchange-type Hamiltonians (e.g. $XX{+}YY$ on the logical qubits) that conserve the single-excitation manifold and stabilize $|\Psi_{\text{para}}^{(n)}\rangle$.

(iii) Global parasymmetric drives.
Global parasymmetric drives $U_{\text{para}} = \exp\!\left(i\hat{F}_{\text{para}}\right)$ with Hermitian $\hat{F}_{\text{para}}$ satisfying

$$[\hat{F}_{\text{para}}, P_{\text{para}}^{(x_1\ldots x_n)}] = 0$$

(cf. Definition 2.85).

(P3) Equivalences and closure. Parasymmetric realizations related by (i) local rephasings $|e_1\rangle \mapsto e^{i\phi}|e_1\rangle$, $|e_2\rangle \mapsto e^{-i\phi}|e_2\rangle$; (ii) relabeling of sites; or (iii) common conjugation within the $(0,1)$ sector are *operationally equivalent*: they yield the same family of effects and Born valuations. The pair $\{P_{\text{para}}^{(x)}, P_{\text{antipara}}^{(x)}\}$ is a complete, orthogonal resolution of $P_{(0,1)}^{(x)}$, ensuring basis-coherence with Thesis/Antithesis.

Interpretation. Parasymmetry names *coherent organization* rather than compromise: locally it is the $|+\rangle$ filter; globally it is distributed alignment that no single site owns. Its truth is the measurable propensity $v_{\text{Synthesis}}(\rho)$ (Definition 1.18), and its dynamics are those that conserve the parasymmetric pattern while allowing it to propagate and entangle across PQSs.

Intuitive reading. Parasymmetry is not a compromise between Symmetry and Antisymmetry; it is a *pattern of coherent organization*. Locally, the $|+\rangle$ filter in (2.57) asks: "Does the system register in the *coherent* direction that aligns the two poles?" Its Born truth $v_{\text{Synthesis}}(\rho) = \text{Tr}(P_{\text{para}}^{(x)}\rho)$ (Definition 1.18) is therefore a measure of *coherence*, not mere population. Two preparations can have the same P_T/P_A weights yet differ sharply in $v_{\text{Synthesis}}$—a classical mixture keeps masses on the poles, a coherent superposition points along the parasymmetric axis. Read this way, Synthesis is the corner of the Fourfold Dialectic Algebra (Definition 1.16) that detects "*togetherness with phase*," i.e., aligned disposition under the complementary test.

Operational picture. To probe Parasymmetry at one site, rotate and measure the logical qubit along the parasymmetric axis (the Σ_x basis). Projecting with $P_{\text{para}}^{(x)}$ implements a repeatable *test & filter*: repeated trials estimate $v_{\text{Synthesis}}(\rho)$, while post-selection prepares the $|+\rangle$ mode for downstream dynamics. In the many-site setting, the bipartite projector (2.58) and the global projector (2.59) turn Synthesis into a *relation*: the signature is no longer at any single PQS, but in their joint registration. The pair state is Bell-like (one excitation shared), while

the global state is W-like (single-excitation delocalization) with permutation symmetry and mixed one-site marginals; both realize "distributed co-alignment."

Dynamics and robustness. Equation (2.60) characterizes the evolutions that preserve Synthesis: a unitary U is parasymmetry-preserving iff it leaves $P_{\text{para}}^{(x_1\cdots x_n)}$ invariant. Practically, this includes (i) local rotations that keep the $\{|+\rangle, |-\rangle\}$ axis fixed; (ii) exchange-type couplings (e.g. $XX{+}YY$ on the logical qubits) that stabilize the single-excitation manifold and propagate coherence; and (iii) global drives generated by $\hat{F}_{\text{para}}$ with $[\hat{F}_{\text{para}}, P_{\text{para}}^{(x_1\cdots x_n)}] = 0$. Couplings that leak weight outside the $(0,1)$ sector or that dephase the $\{|e_1\rangle, |e_2\rangle\}$ coherence reduce $v_{\text{Synthesis}}$—they do not "negate" Synthesis so much as *dilute* the very resource it measures. Because the global W-like pattern is spread across sites, it is notably robust to loss of a single constituent (the signature property of W-type coherence), making it a natural carrier of parasymmetric order.

What Synthesis is (and is not). Synthesis tolerates nonzero Thesis *and* Antithesis valuations: bivalence is recovered only when one pole saturates. But high $v_{\text{Synthesis}}$ demands *phase-coherent* co-alignment, not a probabilistic blend. Thus Parasymmetry is the quantitative hinge between the "both/neither" possibilities of the dialectic and an experimentally legible notion of togetherness: a disposition to register along the coherent axis locally, and to share a single organized pattern nonlocally.

Definition 2.95 (Local parasymmetric effect $P_{\text{para}}^{(x)} = |+\rangle\langle+|$ as a circuit)**.** Since $P_{\text{para}}^{(x)} = \frac{1}{2}(\mathbb{I}+X)$, measuring X operationalizes (2.57). Equivalently, Hadamard then Z-measurement:



$$\Pr[0] = \mathrm{Tr}\Big(P_{\text{para}}^{(x)}\rho\Big), \quad \Pr[1] = \mathrm{Tr}\Big(P_{\text{antipara}}^{(x)}\rho\Big).$$

(Coherent/Naimark flag for $|+\rangle$: apply H, then a zero–control to flip an ancilla, then H back.)



Outcome a=1 implements projection onto $|+\rangle$.

Definition 2.96 (Pair parasymmetric projector $P_{\text{para}}^{(x_i x_j)} = |\Psi_{\text{para}}^{(ij)}\rangle\langle\Psi_{\text{para}}^{(ij)}|$)**.** Here $|\Psi_{\text{para}}^{(ij)}\rangle = \frac{1}{\sqrt{2}}(|01\rangle + |10\rangle) = |\Psi^+\rangle$ (on d_{x_i}, d_{x_j}). Inverse Bell transform (CNOT then H on the first qubit) maps $\{|\Phi^+\rangle, |\Phi^-\rangle, |\Psi^+\rangle, |\Psi^-\rangle\} \to \{|00\rangle, |01\rangle, |10\rangle, |11\rangle\}$. Postselect $(1,0)$ to realize $P_{\text{para}}^{(x_i x_j)}$:



Coherent flag (no collapse): replace meters by a zero/one–conditioned multi-control on an ancilla and uncompute.

Definition 2.97 (Global parasymmetric projector $P_{\text{para}}^{(x_1\cdots x_n)} = |\Psi_{\text{para}}^{(n)}\rangle\langle\Psi_{\text{para}}^{(n)}|$)**.** Let V_+ be the n-qubit state-prep circuit that maps a seed $|10\cdots0\rangle$ to the W-type state $|\Psi_{\text{para}}^{(n)}\rangle$ of (2.59). A Naimark flag for $|\Psi_{\text{para}}^{(n)}\rangle$:



The control is conditioned on the computational pattern $10\cdots0$ after $V_+^\dagger$; outcome a=1 realizes $P_{\text{para}}^{(x_1\cdots x_n)}$.

Definition 2.98 (Circuits for parasymmetry-preserving unitaries (implement (2.60)))**.** A unitary U preserves parasymmetry iff $[U, P_{\text{para}}^{(x_1\cdots x_n)}] = 0$. Representative circuit classes:

1. **Decoupled local rotations about the parasymmetric axis.** Since $P_{\text{para}}^{(x)} = \frac{1}{2}(\mathbb{I} + X)$, any $U_x = e^{i\theta_x X}$ commutes with $P_{\text{para}}^{(x)}$:



   and $U = \bigotimes_x U_x$ preserves (2.60).

2. **Pairwise XY exchange (stabilizes the single-excitation manifold).** Using $P_{\text{para}}^{(x_ix_j)} - P_{\text{antipara}}^{(x_ix_j)} = \frac{1}{2}(X_iX_j + Y_iY_j)$,

$$U_{ij}(\alpha) = \exp\Big(i\tfrac{\alpha}{2}(X_iX_j + Y_iY_j)\Big) \quad \equiv$$



   These gates conserve excitation number and hence preserve the W-sector spanned by (2.59).

3. **Global modal-class phase drives.** If $\hat{F}_{\text{para}}^{(n)} = E_+P_{\text{para}}^{(x_1\cdots x_n)} + E_-\widetilde{P_{\text{antipara}}^{(x_1\cdots x_n)}} + \cdots$, then $U_{\text{para}} = \exp\big(i\hat{F}_{\text{para}}^{(n)}\big)$ is implemented by flagged phase kickback on the two rank-1 subspaces:



   with phases $\phi_\pm$ set by the desired eigenphases on $|\Psi_{\text{para}}^{(n)}\rangle$ and its orthogonal complement.

Remark 2.99 (Operational equivalences and closure)**.** Local rephasings and site permutations correspond to Z-rotations and wire-swaps; common conjugations within the $(0,1)$ sector are implemented by uniform single-qubit basis changes (e.g. global H or R_z layers). These redundancies leave all circuits above invariant in their Born statistics, matching Definition 2.94.

Local characterization. Parasymmetry on the distinguished $(0,1)$ modal sector is the coherent superposition structure singled out by the projectors $P^{(x)}_{\text{para}}$ and $P^{(x)}_{\text{antipara}}$ in Definition 2.94. With $|+\rangle_x = \frac{1}{\sqrt{2}}(|e_1\rangle_x + |e_2\rangle_x)$ and $|-\rangle_x = \frac{1}{\sqrt{2}}(|e_1\rangle_x - |e_2\rangle_x)$, the effects $P^{(x)}_{\text{para}} = |+\rangle_x\langle+|$, $P^{(x)}_{\text{antipara}} = |-\rangle_x\langle-|$ (eq. (2.57)) resolve the $(0,1)$-sector as a pair of orthogonal X-eigenspaces. Operationally, measuring the complementary observable Σ_x on the logical qubit implements the synthesis test, i.e. the Born valuation $v_{\text{Synthesis}}(\rho) = \text{Tr}(P^{(x)}_{\text{para}}\rho)$ in Definition 2.94.

Distributed (pair and global) synthesis. Parasymmetry elevates local synthesis to relational coherence across sites via the pair state $|\Psi^{(ij)}_{\text{para}}\rangle = \frac{1}{\sqrt{2}}(|e_1\rangle_{x_i}|e_2\rangle_{x_j} + |e_2\rangle_{x_i}|e_1\rangle_{x_j})$ and its projector $P^{(x_ix_j)}_{\text{para}}$ (eq. (2.58)), as well as the multipartite W-type state $|\Psi^{(n)}_{\text{para}}\rangle$ and projector $P^{(x_1\ldots x_n)}_{\text{para}}$ (eq. (2.59)). These effects witness *distributed synthesis*: the pattern is encoded nonlocally, with permutation symmetry, mixed single-site marginals, and support confined to the $(0,1)$ sector (cf. eq. (2.29)).

Circuit interpretation. At the circuit level, the local effect $P^{(x)}_{\text{para}}$ is realized by an X-eigenstate test (e.g. Hadamard followed by a Z-measurement), which directly computes $\text{Tr}(P^{(x)}_{\text{para}}\rho)$ as in Definition 2.94 and eq. (2.57). The pair effect $P^{(x_ix_j)}_{\text{para}}$ is implemented by mapping the Bell basis to the computational basis and testing for the symmetric exchange component specified in eq. (2.58). The global effect $P^{(x_1\ldots x_n)}_{\text{para}}$ is implemented by coherently flagging the W-state of eq. (2.59) via state-preparation unitaries and Naimark dilation, thereby projecting onto the subspace of multipartite synthesis.

Parasymmetry-preserving dynamics. A unitary U preserves parasymmetry precisely when it stabilizes the parasymmetric subspace, $UP^{(x_1\ldots x_n)}_{\text{para}}U^\dagger = P^{(x_1\ldots x_n)}_{\text{para}}$ (eq. (2.60)). This includes (i) decoupled local rotations about the parasymmetric axis (single-qubit X-axis rotations that commute with eq. (2.57)); (ii) coupled XY-type exchanges $\exp\left(i\frac{\alpha}{2}(X_iX_j + Y_iY_j)\right)$ that conserve excitation number and thus preserve the W-manifold underlying eq. (2.59); and (iii) global modal-class drives $U_{\text{para}} = \exp\left(i\hat{F}^{(n)}_{\text{para}}\right)$ whose generators commute with $P^{(x_1\ldots x_n)}_{\text{para}}$, as encompassed by the unified constructions in Definition 2.85.

Synthesis as an operational principle. In summary, parasymmetry designates the X-eigenstructure of each duality qubit and its coherent uplift to nonlocal, permutation-symmetric entanglement. Its circuits project onto, preserve, phase, and transport the synthesis content specified by Definition 2.94 and eqs. (2.58) and (2.59), while the admissible evolutions are exactly those unitaries satisfying the invariance criterion eq. (2.60).

Intuitive essence. Parasymmetry designates the balanced coherence within the distinguished $(0,1)$ sector: locally it is the equal-weight superposition $|+\rangle_x = \frac{1}{\sqrt{2}}(|e_1\rangle_x + |e_2\rangle_x)$ singled out by the effect $P^{(x)}_{\text{para}} = |+\rangle_x\langle+|$ (Definition 2.94 and eq. (2.57)). Rather than committing to either mode (Symmetry/Antisymmetry), the parasymmetric effect identifies the "midpoint" state that coherently synthesizes both. The complementary effect $P^{(x)}_{\text{antipara}}$ identifies the orthogonal dissonant alternative.

Distributed synthesis. Beyond a single site, parasymmetry becomes a relational property encoded in correlations. In the bipartite case, synthesis is witnessed by the symmetric exchange state $|\Psi^{(ij)}_{\text{para}}\rangle = \frac{1}{\sqrt{2}}(|e_1\rangle_{x_i}|e_2\rangle_{x_j} + |e_2\rangle_{x_i}|e_1\rangle_{x_j})$ and its projector $P^{(x_i x_j)}_{\text{para}}$ (eq. (2.58)). Globally, the parasymmetric W-state $|\Psi^{(n)}_{\text{para}}\rangle$ delocalizes a single antisymmetry excitation over all sites, defining the projector $P^{(x_1\ldots x_n)}_{\text{para}}$ (eq. (2.59)). In both cases, synthesis is realized as a nonlocal pattern: no single site carries the entire structure.

Operational role. Operationally, parasymmetry *selects* coherent synthesis and *maintains* it under admissible dynamics. Selection is achieved by testing the local effect $P^{(x)}_{\text{para}}$ and its pair/global analogues (eqs. (2.57) to (2.59)), yielding the Born valuation $v_{\text{Synthesis}}(\rho) = \text{Tr}(P^{(x)}_{\text{para}}\rho)$ (Definition 2.94). Maintenance is characterized by the invariance criterion $U\,P^{(x_1\ldots x_n)}_{\text{para}}U^\dagger = P^{(x_1\ldots x_n)}_{\text{para}}$ (eq. (2.60)), which is satisfied by (i) local rotations about the parasymmetric axis, (ii) number-conserving XY exchange unitaries on pairs that stabilize the single-excitation manifold, and (iii) global modal-class drives generated by Hermitians commuting with $P^{(x_1\ldots x_n)}_{\text{para}}$ (cf. Definition 2.85).

Conceptual summary. Parasymmetry is the coherent "both-at-once" structure of the duality qubit lifted to a collective, permutation-symmetric relation across many sites. Its formal content is the family of effects and states in Definition 2.94 and eqs. (2.58) and (2.59), together with the closed class of evolutions that preserve these effects (eq. (2.60)). In this sense, parasymmetry performs two tasks: it filters for synthesis and it enables synthesis to propagate, interfere, and organize globally without fracturing the coherent pattern that defines it.

2.3 Holosymmetry: The Quantum Realization of Holothesis

Holosymmetry represents one of the most profound manifestations of quantum mechanics within this treatise: the coherent superposition of entire evolutionary histories of worlds. Unlike conventional quantum superposition, which operates on states within a fixed dynamical framework, Holosymmetry enables the simultaneous exploration of fundamentally different evolutionary pathways—a quantum mechanical realization of parallel universes at the most elementary level of reality.

At its heart, Holosymmetry orchestrates a cosmic dance between two emergent quantum registers, each comprising n Primordial Quantum Singularities (Definition 1.25). These registers—designated Register A and Register B—are collections of quantum entities as well as independent instantiations of the universe's fundamental duality structure. Each

PQS within these registers, expressed as $\mathfrak{B}_x \equiv \hat{\mathbf{\Omega}}_x = \overset{\circ}{\mathbf{\Lambda}}_{x,\hat{\Theta}_x}$, embodies the irreducible quantum-relational architecture from which spacetime itself emerges.

The revolutionary aspect of Holosymmetry lies in its ability to place these entire registers into quantum superposition with respect to their dynamical evolution. Through the mediation of a control quantum system—itself an PQS residing in the modal Hilbert space $\mathcal{H}_c$ (eq. (2.64))—the framework enables each register to simultaneously experience both coupled and non-coupled parasymmetric evolution (Definition 2.85). This is a fundamental feature of how the pre-geometric substrate explores the landscape of possible emergent geometries, thus freeing it from a mere theoretical curiosity.

The control system acts as a quantum switch, entangling itself with the evolutionary fate of both registers. When in state $|\mathbf{0}\rangle_c$, it guides Register A through coupled parasymmetric dynamics while Register B evolves under non-coupled dynamics. In state $|\mathbf{1}\rangle_c$, these roles reverse. But through quantum superposition, the control system can exist in both states simultaneously, creating an entangled configuration where both evolutionary histories coexist—a superposition of spacetime's possible emergent structures.

This construction elevates the Quantum Holo-State $|\overset{\circ}{\mathbf{\Psi}}\rangle$ (Definition 1.27) from a static encoding of quantum correlations to a dynamic generator of parallel realities. Each evolutionary path carves out a different trajectory through the space of possible geometric configurations, and Holosymmetry ensures that quantum interference between these paths becomes physically meaningful. The result is a framework where the very notion of a unique evolutionary history dissolves into a quantum superposition of possibilities, each weighted by complex amplitudes that encode the deep relational structure of the PQS network.

In essence, Holosymmetry transforms the question "Which dynamical evolution did the system undergo?" into the quantum mechanical statement "The system underwent all compatible evolutions simultaneously, with quantum interference determining the observable outcome.". This represents the ultimate expression of quantum mechanics' radical departure from classical determinism, now applied to the very fabric of emergent spacetime itself.

2.3.1 System Definition and Component Operators

Why two registers (and a control) for Holosymmetry. Holosymmetry tests a meta-level question: it compares *entire evolutionary histories*, not just states. To do this without ambiguity or higher-order wiring, we require two disjoint n-PQS registers so that mutually exclusive global regimes can be instantiated *in parallel* and then coherently contrasted. In our setting the alternative parasymmetric evolutions are $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$ and $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$, where one register runs the globally coupled dynamics (G) while the other runs the decoupled/normal dynamics (N) and vice versa (cf. Definition 2.85). A single control qubit then prepares, addresses, and recombines these two global histories, realizing the holosymmetric unitary in the normal form of eq. (3.35) and Section 3.8, (3.35). In this architecture the measured quantity is a *propensity*, the Born valuation of the holosymmetric effect $E_{\mathrm{Holo}}^{(\mathbf{a}\nrightarrow\mathbf{b})}$ defined in (3.36) and evaluated via Definition 1.18.

Minimal, symmetry-respecting, and operationally complete. The dual-register scaffold is minimal: it is the least structure that lets us form sums/differences of histories and

expose their relational content under different control bases. It also realizes the Principle of Duality and an $A \leftrightarrow B$ swap symmetry by construction: exchanging registers or applying common conjugations leaves the family of effects—and hence the valuations—unchanged (Holosymmetry equivalences (H4) in Section 3.8). Crucially, the same backbone unifies the three operational presentations (interferometer, path-scattering, and commutator-lensing): choices of $\mathbf{U}_{\text{prep}}$ and $\mathbf{U}_{\text{read}}$ together with the $\{|\mathbf{0}\rangle_c, |\mathbf{1}\rangle_c\}$ or $\{|+\rangle_c, |-\rangle_c\}$ control bases recover each model (see Definitions 2.114 and 2.119 and Notes (i) in Section 3.8). All of this remains within fixed causal order—Holosymmetry superposes *which global unitary runs where*, not the *order* of gates—so the construction stays fully unitary and experimentally standard.

Definition 2.100 (Axiom (Principle) of Duality)**.** For every ontological assertion admissible in the Fourfold Dialectic Algebra (Definition 1.16) there exists a distinguished, complementary counter-assertion such that:

(i) **Ontological layer.** Each PQS $\mathfrak{B}_x$ carries a privileged dual pair (Symmetry/Antisymmetry), generating the $(0, 1)$ sector and its complementary (parasymmetric) axis.

(ii) **Algebraic layer.** The pair is realized by effects P_T, P_A with $P_T + P_A = P_{(0,1)}$ and complementary effects $P_{\text{para}}, P_{\text{antipara}}$; truth is Born-valued $v(\rho; P) = \text{Tr}(P\rho)$ (Definition 1.18).

(iii) **Operational layer.** Physical constructions respect the duality symmetry: swapping the dual branches (or relabeling them globally) leaves all induced valuations invariant, up to unitary equivalence on the tested space.

Duality as the design principle for Holosymmetry. Holosymmetry compares *alternative histories*, so the Axiom of Duality (Section 1.2.6 and Definition 2.100) must be realized not only for *states* but for *processes*. The minimal, symmetry-respecting way is to instantiate two disjoint n-PQS registers so that the mutually exclusive global regimes can be run *in parallel* and then coherently contrasted. Concretely, the dual pair of path unitaries $\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}$ and $\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}$ assigns "global–coupled" (G) to one register while the other runs "normal/decoupled" (N), and vice versa; a single control prepares, addresses, and recombines these histories to realize the normal form in eq. (3.35) and Section 3.8, (3.35). The measured quantity is a Born propensity, $v_{\text{Holothesis}}(\rho_{\text{RR}}) = \text{Tr}\Big(E_{\text{Holo}}^{(\mathbf{a}\nrightarrow\mathbf{b})}\, \rho_{\text{RR}}\Big)$ from (3.36), which reports "synthesis vs. dissonance'' of the two histories under the chosen preparation/readout. With this duality-driven motivation in place, we now specify the two registers and the composite holosymmetric Hilbert space.

Definition 2.101 (Dual Primordial Registers and Holosymmetric Hilbert Space)**.** Let $\mathcal{X}_q$ denote the emergent quantum configuration space of Primordial Quantum Singularities (PQSs), as defined in Proposition 4.625. For a holosymmetric system, define two disjoint, factorized quantum registers:

- **Register A**: $\mathcal{X}_q^{(A)} = \{x_1^{(A)}, \ldots, x_n^{(A)}\} \subset \mathcal{X}_q$,
- **Register B**: $\mathcal{X}_q^{(B)} = \{x_1^{(B)}, \ldots, x_n^{(B)}\} \subset \mathcal{X}_q$,

with $\mathcal{X}_q^{(A)} \cap \mathcal{X}_q^{(B)} = \emptyset$, such that each register contains n distinct PQSs.

Modal Structure Each PQS $\mathfrak{B}_x \in \mathcal{X}_q$ possesses a local modal Hilbert space:

$$\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)} := \mathbb{C}^4,$$

corresponding to the full fourfold dialectical spectrum: Symmetry, Antisymmetry, Parasymmetry, and Holosymmetry.

The Hilbert space of a single register $\mathcal{X}_q^{(A)}$ or $\mathcal{X}_q^{(B)}$ is then:

$$\mathcal{H}_{\text{PQS}}^{(n,A)} := \bigotimes_{i=1}^{n} \mathcal{H}_{\text{mod}}^{(x_i^{(A)})} \cong (\mathbb{C}^4)^{\otimes n}, \quad \mathcal{H}_{\text{PQS}}^{(n,B)} := \bigotimes_{i=1}^{n} \mathcal{H}_{\text{mod}}^{(x_i^{(B)})} \cong (\mathbb{C}^4)^{\otimes n}.$$

Dual Register Hilbert Space We define the combined Hilbert space of the two independent PQS registers—each containing n sites—as the *dual register Hilbert space*, denoted:

$$\boxed{\mathcal{H}_{\text{RR}}^{(2n)} := \mathcal{H}_{\text{PQS}}^{(n,A)} \otimes \mathcal{H}_{\text{PQS}}^{(n,B)} \cong (\mathbb{C}^4)^{\otimes 2n}.} \tag{2.61}$$

Here:

- $\mathcal{H}_{\text{PQS}}^{(n,A)} := \bigotimes_{i=1}^{n} \mathcal{H}_{\text{mod}}^{(x_i^{(A)})}$ and
- $\mathcal{H}_{\text{PQS}}^{(n,B)} := \bigotimes_{i=1}^{n} \mathcal{H}_{\text{mod}}^{(x_i^{(B)})}$,

with $\mathcal{H}_{\text{mod}}^{(x)} = \mathbb{C}^4$ the full four-dimensional modal Hilbert space at each PQS.

The label **RR** stands for *Primordial Registers*, emphasizing the factorized structure of the total configuration across the two modal subsystems. This construction forms the tensorial base upon which holosymmetric operators act, either jointly or in superposition.

Control System and Holosymmetric Sector Introduce an auxiliary control qubit space $\mathcal{H}_c \cong \mathbb{C}^2$, enabling controlled superpositions of dynamical histories.

Then the full holosymmetric Hilbert space is:

$$\boxed{\mathcal{H}_{\text{Holo}} := \mathcal{H}_c \otimes \mathcal{H}_{\text{PQS}}^{(n,A)} \otimes \mathcal{H}_{\text{PQS}}^{(n,B)}.} \tag{2.62}$$

Dimensionality. We work on the factorized space

$$\mathcal{H}_{\text{Holo}} = \mathcal{H}_c \otimes \mathcal{H}_R^{(n)} \otimes \mathcal{H}_{R'}^{(n)} \cong \mathbb{C}^2 \otimes \mathbb{C}^{4^n} \otimes \mathbb{C}^{4^n}.$$

$$\boxed{\dim(\mathcal{H}_{\text{Holo}}) = \dim(\mathcal{H}_c)\, \dim(\mathcal{H}_R^{(n)})\, \dim(\mathcal{H}_{R'}^{(n)}) = 2 \cdot 4^n \cdot 4^n = 2 \times 4^{2n} = 2^{4n+1}.} \tag{2.63}$$

Interpretation. This dual-register structure forms the holosymmetric substrate for constructing entangled path histories, interference geometry, and higher-order duality transformations across PQS configurations. It generalizes parasymmetry from single-register dynamics to controlled entangled evolution over bipartite modal landscapes.

This exponential scaling reflects the full modal richness of holosymmetric evolution. With $2n$ Primordial Quantum Singularities distributed across two registers, each possessing four distinct dialectical modes, plus one control qubit, the system can explore 2^{4n+1} orthogonal quantum states—enabling deep superpositions of entangled causal geometries within the emergent spacetime fabric.

Definition 2.102 (Control System for Holosymmetric Evolution)**.** The holosymmetric framework incorporates a quantum control mechanism that coherently selects between distinct dynamical regimes applied to two disjoint PQS registers. This control system is implemented by a two-level quantum system (qubit) with Hilbert space:

$$\boxed{\mathcal{H}_c := \text{span}\left\{|\mathbf{0}\rangle_c,\ |\mathbf{1}\rangle_c\right\},} \tag{2.64}$$

where the orthonormal basis states represent complementary dynamical assignments:

$$\boxed{\begin{aligned} |\mathbf{0}\rangle_c &\equiv \begin{cases} \text{Register A: } U_G^{(\odot)} \text{ (unified global coupled evolution)}, \\ \text{Register B: } U_N^{(\odot)} \text{ (generalized composite decoupled evolution)}, \end{cases} \\ |\mathbf{1}\rangle_c &\equiv \begin{cases} \text{Register A: } U_N^{(\odot)} \text{ (generalized composite decoupled evolution)}, \\ \text{Register B: } U_G^{(\odot)} \text{ (unified global coupled evolution)}. \end{cases} \end{aligned}} \tag{2.65}$$

When the control state is prepared in a coherent superposition:

$$|\phi_c\rangle = \alpha|\mathbf{0}\rangle_c + \beta|\mathbf{1}\rangle_c, \qquad \alpha, \beta \in \mathbb{C}, \quad |\alpha|^2 + |\beta|^2 = 1,$$

the resulting system undergoes a quantum superposition of full modal evolutions, where the assignment of coupling modes to the registers is no longer sharply defined.

This defines the operational heart of *holosymmetry*: a mode of evolution where distinct parasymmetric histories—comprising separable and entangled, local and global coherence structures—interfere through controlled quantum superposition. It encodes a higher-order synthesis across the modal and geometrical structures of Primordial Quantum Singularities.

Lifted Operators on Dual PQS Registers

Definition 2.103 (Lifted Parasymmetric Operators on Dual Primordial Quantum Singularity Registers)**.** Let $\mathcal{H}_{\text{RR}}^{(2n)} := \mathcal{H}_{\text{PQS}}^{(n,A)} \otimes \mathcal{H}_{\text{PQS}}^{(n,B)}$ denote the dual register Hilbert space of two disjoint PQS configurations (eq. (2.61)).

Let $U_G^{(\odot)}$ and $U_N^{(\odot)}$ be the unified global coupled and generalized decoupled parasymmetric operators, respectively (Definitions 2.66 and 2.85).

We define the **lifted operators** acting on the dual space $\mathcal{H}_{\text{RR}}^{(2n)}$ by extending the action of each operator to one register while leaving the other invariant:

$$U_G^{(A)} := U_G^{(\odot)} \otimes \mathbb{I}^{(B)}, \qquad U_N^{(A)} := U_N^{(\odot)} \otimes \mathbb{I}^{(B)}, \tag{2.66}$$

$$U_G^{(B)} := \mathbb{I}^{(A)} \otimes U_G^{(\odot)}, \qquad U_N^{(B)} := \mathbb{I}^{(A)} \otimes U_N^{(\odot)}, \tag{2.67}$$

where:

- $\mathbb{I}^{(A)}$ and $\mathbb{I}^{(B)}$ are identity operators on $\mathcal{H}_{\text{PQS}}^{(n,A)}$ and $\mathcal{H}_{\text{PQS}}^{(n,B)}$, respectively;
- The superscripts (A) and (B) denote selective action restricted to Register A or Register B.

Interpretation. This lifting formalism enables the application of structurally distinct parasymmetric dynamics to each register, with the following implications:

- $U_G^{(A)}$: Applies a fully entangled, globally coupled parasymmetric evolution within Register A, while Register B remains static.
- $U_N^{(B)}$: Applies a factorizable, decoupled parasymmetric evolution within Register B, leaving Register A unchanged.
- All such lifted operators preserve the tensor product structure of $\mathcal{H}_{\text{RR}}^{(2n)}$, making them compatible with controlled superpositions across registers.

Holosymmetric Role. These lifted operators form the operational backbone of the holosymmetric architecture. They enable coherent dynamical differentiation between registers, making it possible to construct quantum superpositions of mutually exclusive modal histories—where Register A and Register B may evolve under independent coherence constraints, yet remain entangled via the overarching control system.

The Necessity of Lifting: From Mathematical Requirement to Physical Reality The concept of "lifting" operators in the holosymmetric framework reveals a profound mathematical necessity that mirrors the philosophical depth of the Holothesis itself. To understand why lifting is convenient and absolutely essential, we must trace the logical architecture from its foundations through to its revolutionary physical implications.

The Structural Foundation Consider the architecture of the holosymmetric universe: we have constructed a total Hilbert space $\mathcal{H}_{\text{Holo}} = \mathcal{H}_c \otimes \mathcal{H}_{\text{PQS}}^{(n,A)} \otimes \mathcal{H}_{\text{PQS}}^{(n,B)}$ that is explicitly factorized across three independent subsystems—a control qubit and two quantum registers, each containing n Primordial Quantum Singularities. This tripartite structure embodies the very essence of holosymmetric logic, where different evolutionary histories must coexist in carefully orchestrated quantum superposition.

The heart of holosymmetry lies in its ability to coherently superpose distinct assignments of modal dynamics. In one branch of reality, Register A might evolve under the globally coupled operator $U_G^{(\odot)}$ while Register B evolves under the decoupled operator $U_N^{(\odot)}$. In another branch, these assignments are reversed. But here we encounter a fundamental operational challenge: the operators $U_G^{(\odot)}$ and $U_N^{(\odot)}$ are defined to act on single-register spaces of dimension 4^n, not on the full dual-register space of dimension 4^{2n}.

The Mathematical Resolution Through Lifting This dimensional mismatch is where lifting becomes indispensable. To construct selective operators that act on one register while leaving the other invariant, we must extend each operator's domain through tensor products with identity operators: $U_G^{(A)} := U_G^{(\odot)} \otimes \mathbb{I}^{(B)}$ and similarly for the others. This lifting procedure preserves the tensor product structure of the total space while enabling register-specific evolution—a mathematical transformation that makes the impossible possible. The true elegance of lifting emerges when we construct the controlled-holosymmetric unitary operator. The control qubit must coherently select between two complete evolutionary scenarios:

$$\boxed{\hat{U}_{\text{holo}} := |\mathbf{0}\rangle\langle\mathbf{0}|_c \otimes (U_G^{(A)} \cdot U_N^{(B)}) + |\mathbf{1}\rangle\langle\mathbf{1}|_c \otimes (U_N^{(A)} \cdot U_G^{(B)}).} \tag{2.68}$$

Without lifting, this construction would be mathematically ill-defined—we would be attempting to combine operators that live in incompatible spaces. The lifted operators, however, all act on the same dual-register space $\mathcal{H}_{\text{RR}}^{(2n)}$, making their coherent superposition not just possible but natural.

From Abstract Computation to the Fabric of Reality This mathematical machinery captures something profound about the nature of reality within the IQR. Unlike conventional quantum computing, which manipulates abstract qubits divorced from physical meaning, the holosymmetric framework operates on Primordial Quantum Singularities ($\mathfrak{B}_x$)—the irreducible quantum seeds from which spacetime itself emerges. These are not computational conveniences but the actual building blocks of reality, each intrinsically characterized by the Quantum Holo-State $|\overset{\circ}{\mathbf{\Psi}}\rangle$ and possessing the full richness of the Fourfold Dialectic Algebra.

When we apply lifted operators like $U_G^{(A)} = U_G^{(\odot)} \otimes \mathbb{I}^{(B)}$, we are orchestrating the evolution of the universe's fundamental constituents. The lifting construction ensures that Register A can undergo coupled parasymmetric evolution, generating internal coherence and entanglement among its constituent PQSs, while Register B remains in stasis. This selective evolution preserves the tensor product structure while enabling each register to follow its own dialectical trajectory.

The Fibered Architecture of Quantum Reality The mathematical space in which these operations unfold is itself revolutionary. Rather than the flat tensor product spaces of conventional quantum mechanics, we work within emergent, fibered Hilbert spaces that reflect the deep geometric structure of reality. The register spaces $\mathcal{H}_{\text{PQS}}^{(n,A)}$ and $\mathcal{H}_{\text{PQS}}^{(n,B)}$ emerge dynamically through the Quantum Holo-State's relational structure, forming part of a Hilbert bundle over the emergent quantum configuration space $\mathcal{X}_q$.

This fibered structure means that when we lift operators to act selectively on registers, we are manipulating the very threads from which the tapestry of spacetime is woven. The identity operator on the inactive register represents the preservation of that register's geometric and modal configuration while its partner undergoes transformation.

Holosymmetry as Quantum Superposition of Universal Histories The most profound aspect emerges when we consider what the control system truly governs. When the control qubit exists in superposition:

$$|\phi_c\rangle = \alpha|\mathbf{0}\rangle_c + \beta|\mathbf{1}\rangle_c$$

it places the entire universe of dual registers into a superposition of fundamentally different evolutionary histories. In the $|\mathbf{0}\rangle_c$ branch, Register A undergoes globally coupled evolution while B evolves independently. In the $|\mathbf{1}\rangle_c$ branch, these roles reverse. But in superposition, both histories occur simultaneously, creating interference between different ways the fabric of reality might have woven itself.

This is the operational manifestation of the Holothesis—a mode of being that transcends individual evolutionary paths. Each register maintains its modular independence while the full system encodes relational interference between different evolutionary assignments. The control qubit doesn't merely select between pre-existing possibilities; it enables quantum superposition of fundamentally different ways the universe might organize its internal coherence.

The Physical Process of Lifted Evolution When $U_G^{(A)}$ acts on the dual register state, it generates coupled parasymmetric evolution exclusively within Register A. This is not an abstract transformation but a physical process: the PQSs within Register A develop quantum correlations, their modal states entangle, and coherent superpositions of dialectical modes emerge. Meanwhile, Register B maintains its configuration perfectly, serving as a static reference frame against which Register A's evolution unfolds.

This selective evolution enables something unprecedented: quantum interference between different assignments of physical law. The holosymmetric unitary creates superpositions not just of states but of which regions of the quantum universe evolve under which dynamical principles. Through lifting, we create a mathematical framework where reality itself can exist in superposition—not just superposition of states, but superposition of the very rules by which states evolve.

In this light, lifting is elevated from a technical convenience to its true status as the essential mechanism through which holosymmetry achieves its transcendent synthesis. This process enables quantum correlations between entire frameworks of modal evolution, pointing toward a reality whose deepest structure is a quantum superposition of possible law assignments, supplanting the notion of a single, fixed set of laws.

Wiring. Let the control wire carry $\mathcal{H}_c$ in eq. (2.64), and let $\mathbf{d}^{(A)} = (d_{x_1}^{(A)}, \ldots, d_{x_n}^{(A)})$ and $\mathbf{d}^{(B)} = (d_{x_1}^{(B)}, \ldots, d_{x_n}^{(B)})$ denote the $(0,1)$-sector (duality) qubits for Registers A and B inside $\mathcal{H}_{\mathrm{PQS}}^{(n,A)}$ and $\mathcal{H}_{\mathrm{PQS}}^{(n,B)}$, respectively (eqs. (2.61) and (2.62)). All lifted gates act as identity on the orthogonal modal components within each $\mathbb{C}^4$ fibre.

Definition 2.104 (Uncontrolled lifts on the dual space $\mathcal{H}_{\mathrm{RR}}^{(2n)}$)**.** Instantiating Definition 2.103 with the compiled circuits of $U_G^{(\odot)}$ and $U_N^{(\odot)}$ (Definitions 2.66 and 2.85) yields:



implements $U_G^{(A)} := U_G^{(\odot)} \otimes \mathbb{I}^{(B)}$.



implements $U_N^{(A)} := U_N^{(\odot)} \otimes \mathbb{I}^{(B)}$.



implements $U_G^{(B)} := \mathbb{I}^{(A)} \otimes U_G^{(\odot)}$.



implements $U_N^{(B)} := \mathbb{I}^{(A)} \otimes U_N^{(\odot)}$.

Here each block stands for the full multi-qubit implementation compiled earlier (local R_z/R_x layers, XY exchange pairs, and global flags, as applicable).

Definition 2.105 (Controlled lifted assignment (*holosymmetric switch*))**.** Realize eq. (2.65) on $\mathcal{H}_{\mathrm{Holo}}$ by conditioning on the control qubit: when $|\mathbf{0}\rangle_c$, apply ($U_G^{(\odot)}$ on A)$\otimes$($U_N^{(\odot)}$ on B); when $|\mathbf{1}\rangle_c$, apply ($U_N^{(\odot)}$ on A)$\otimes$($U_G^{(\odot)}$ on B). Using open/closed controls,



This two-layer multiplexor implements the block-diagonal unitary $|\mathbf{0}\rangle\langle\mathbf{0}|_c \otimes \left(U_G^{(A)} U_N^{(B)}\right) + |\mathbf{1}\rangle\langle\mathbf{1}|_c \otimes \left(U_N^{(A)} U_G^{(B)}\right)$, as required by Definition 2.102.

Definition 2.106 (Lifted parasymmetric effects (Naimark flags on one register))**.** Let $V_+^{(A)}$ (resp. $V_+^{(B)}$) prepare $|\Psi_{\mathrm{para}}^{(n)}\rangle$ on $\mathbf{d}^{(A)}$ (resp. $\mathbf{d}^{(B)}$). Then $P_{\mathrm{para}}^{(x_1^{(A)} \ldots x_n^{(A)})} \otimes \mathbb{I}^{(B)}$ and $\mathbb{I}^{(A)} \otimes P_{\mathrm{para}}^{(x_1^{(B)} \ldots x_n^{(B)})}$ (eq. (2.59)) are realized by:

Remark 2.107 (Compatibility with holosymmetric superposition)**.** The lifted unitaries of Definition 2.104 and the controlled multiplexor in Definition 2.105 act on disjoint tensor factors of $\mathcal{H}_{\mathrm{Holo}}$ (eq. (2.62)), thereby preserving the factorized structure required for coherent superpositions of dynamical histories in Definition 2.102.

Register–selective action. Lifted parasymmetric unitaries act by transplanting a single–register evolution onto one register while leaving the other invariant. Concretely, $U_G^{(A)} := U_G^{(\odot)} \otimes \mathbb{I}^{(B)}$ and $U_N^{(B)} := \mathbb{I}^{(A)} \otimes U_N^{(\odot)}$ (and the analogous variants) implement within the addressed register exactly the unified global coupled or the composite decoupled dynamics compiled in Definitions 2.66 and 2.85, and act as identity on the spectator register, as formalized



Figure 2.1: Ancilla outcomes $a_A = 1$ or $a_B = 1$ implement the corresponding lifted projector; coherent flags are obtained by omitting the measurement and uncomputing.

in Definition 2.103. Operationally, these lifts steer the single–excitation parasymmetric amplitude—transporting, phasing, and mixing it via XY–exchange and modal–class phase drives—only inside the targeted register.

Entanglement locality and tensorial integrity. Because each lifted gate is tensored with the identity on the complementary register, it cannot create (or destroy) inter–register entanglement; it reshapes correlations solely within the acted–upon register. Hence the dual–register tensor product structure of eqs. (2.61) and (2.62) is preserved throughout. In particular, parasymmetric patterns such as the W–type synthesis subspace remain confined to the addressed register, while the other register retains its prior state.

Holosymmetric switching via coherent control. The holosymmetric control qubit of Definition 2.102 implements a coherent multiplexing of lifted evolutions according to the assignment basis in eq. (2.65). Prepared in a superposition, the control effects a superposition of *dynamical histories*: "A entangled/B decoupled" and "A decoupled/B entangled" are applied in parallel and subsequently interfere. Thus holosymmetry promotes interference from states to *assignments of evolutions*, providing the operational heart of the holosymmetric sector.

Lifted parasymmetric effects as localized coherence filters. Lifted projectors realized by Naimark–flag constructions act as register–local coherence filters. Using the global parasymmetric state and projector of eq. (2.59), the circuits flag $P_{\text{para}}^{(x_1^{(A)}\ldots x_n^{(A)})} \otimes \mathbb{I}^{(B)}$ or $\mathbb{I}^{(A)} \otimes P_{\text{para}}^{(x_1^{(B)}\ldots x_n^{(B)})}$ without disturbing the spectator register. Combined with the holosymmetric switch, these effects enable post–selection, phase–kickback, and comparison of synthesis across registers while maintaining compatibility with eq. (2.62).

Summary. Lifted parasymmetric operators provide register–addressable control of synthesis within the dual–register architecture: they steer, preserve, and phase parasymmetric coherence in one register while leaving the other untouched (Definition 2.103). Coupled with the

control–conditioned multiplexing of Definition 2.102, they elevate parasymmetric processing to holosymmetry, wherein distinct lifted evolutions coexist in coherent superposition and interfere at the level of full modal histories.

Definition 2.108 (Parallel Path Operators for Dual PQS Configurations)**.** Let the dual register Hilbert space be denoted by $\mathcal{H}_{\mathrm{RR}}^{(2n)}$ (eq. (2.61)). Let $U_G^{(A)}$, $U_N^{(A)}$, $U_G^{(B)}$, and $U_N^{(B)}$ be the lifted parasymmetric operators as defined in Definition 2.103, corresponding to the unified global coupled evolution (Definition 2.85) and the generalized composite decoupled evolution (Definition 2.66).

We define two distinct, complete, and unitary evolutionary pathways for the dual register system:

1. **Path 1 (Coupled-A / Decoupled-B):** This pathway describes a scenario where Register A undergoes fully coupled evolution while Register B evolves under a factorized, decoupled dynamic. It is represented by the operator:

$$\boxed{\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} := U_G^{(A)} \cdot U_N^{(B)}.} \tag{2.69}$$

   This operator places complex, entangling dynamics on Register A while preserving the modular coherence structure of Register B.

2. **Path 2 (Decoupled-A / Coupled-B):** This is the complementary pathway, where Register A evolves under decoupled dynamics while Register B undergoes fully coupled evolution. It is represented by:

$$\boxed{\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} := U_N^{(A)} \cdot U_G^{(B)}.} \tag{2.70}$$

   Here, the roles are reversed, with Register B being the locus of non-separable, holistic evolution.

Physical Significance. Each operator $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$ and $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$ is a unitary operator on $\mathcal{H}_{\mathrm{RR}}^{(2n)}$, representing a distinct, physically realizable evolutionary history for the dual register system. They serve as the foundational "paths" that are placed into coherent superposition by the holosymmetric control system, enabling the study of quantum interference between different dynamical laws.

Wiring convention. A single wire labeled $\mathcal{H}_{\mathrm{PQS}}^{(n,A)}$ (resp. $\mathcal{H}_{\mathrm{PQS}}^{(n,B)}$) denotes the entire n-site register in eq. (2.61). A boxed gate on that wire stands for the multi-qubit implementation compiled earlier for the corresponding operator.

Definition 2.109 (Path 1 circuit: Coupled–A / Decoupled–B)**.** This realizes $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} = U_G^{(A)} U_N^{(B)}$ of eq. (2.69):



Here $U_G^{(\odot)}$ is the unified global coupled evolution and $U_N^{(\odot)}$ is the composite decoupled evolution (Definitions 2.66 and 2.85). Since the registers are disjoint, the two blocks may be executed in parallel.

Definition 2.110 (Path 2 circuit: Decoupled–A / Coupled–B)**.** This realizes $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} = U_N^{(A)} U_G^{(B)}$ of eq. (2.70):



Again, the two multi-qubit blocks act on disjoint tensor factors and commute.

Remark 2.111 (Expanded (one Trotter slice) view for each path)**.** If $U_G^{(\odot)}$ and $U_N^{(\odot)}$ are implemented via first/second-order Trotterization, a single slice of Path 1 can be drawn schematically by expanding each block into its constituent layers (local rotations, XY–exchange pairs, global modal-class flags), all in parallel across registers:



where the A-row layers instantiate the unified global coupled construction and the B-row layers instantiate the composite decoupled construction (Definitions 2.66 and 2.85). Path 2 is identical with A/B rows swapped. Repeating the slice r times yields the full Trotterized evolution for eqs. (2.69) and (2.70).

Remark 2.112 (Holosymmetric control wrapper (optional))**.** To place the two paths into coherent superposition as in Definition 2.102, prepend a control wire $\mathcal{H}_c$ and condition the assignment according to eq. (2.65):



The first (open-control) column realizes eq. (2.69) on the registers for $|\mathbf{0}\rangle_c$, the second (closed-control) column realizes eq. (2.70) for $|\mathbf{1}\rangle_c$.

Overview. The parallel path circuits enact two unitary, register-selective evolutions on the dual PQS architecture of eqs. (2.61) and (2.62). Each evolution is formed by *lifting* previously defined parasymmetric dynamics to a chosen register while the other register remains invariant (Definition 2.103). The result is a pair of complete, physically distinct histories, each realized as a product of multi–qubit operations confined to one register factor at a time (Definition 2.108).

Path 1 vs. Path 2. In *Path 1* (eq. (2.69)), Register A undergoes the unified global coupled evolution $U_G^{(A)}$ while Register B experiences the composite decoupled evolution $U_N^{(B)}$ (Definitions 2.66 and 2.85). This places entangling, exchange–type transport and modal–class phasing within A, and factorized, sitewise dynamics within B. In the complementary *Path 2* (eq. (2.70)), the roles are reversed: Register B hosts the globally coupled synthesis while Register A evolves in a decoupled manner. Because the registers occupy disjoint tensor factors, the two blockwise evolutions commute and can be executed in parallel without inter–register crosstalk.

Coherent switching and interference. The holosymmetric control system of Definition 2.102 coherently assigns these paths by conditioning on the control basis of eq. (2.65). Prepared in a superposition, the control implements a superposition of full evolutions—"A coupled/B decoupled" *and* "A decoupled/B coupled"—so that the resulting histories interfere. In this way, holosymmetry promotes interference from static states to *assignments of dynamics*, enabling operational comparisons between coupled and decoupled parasymmetric laws within the same unified protocol.

Local coherence tests. Register-local parasymmetric effects are realized by lifted Naimark-flag projectors that act on one register while leaving the other untouched (cf. eq. (2.59) and Definition 2.103). These flags serve as localized coherence filters—e.g., tests for the global parasymmetric subspace in A or in B—and are compatible with either path as well as with the control-conditioned superposition. Hence they support post-selection and phase-kickback procedures without disturbing the spectator register.

Summary. The parallel path circuits implement two complementary, unitary histories on the dual PQS registers (Definition 2.108): one with globally coupled synthesis in A and decoupled evolution in B, and the other with the assignments reversed. A coherent control realizes their superposition and interference, while lifted projectors provide register–local diagnostics, together furnishing a holosymmetric scheme that steers, contrasts, and coherently compares parasymmetric dynamics across the dual–register architecture.

Intuitive picture. Consider two separate ensembles of quantum "lights": Register A and Register B. The *controlled holosymmetric operator* acts as a smart switch, governed by a tiny control light (the control qubit), that decides which ensemble performs a complex, highly coordinated choreography (the *coupled* evolution) and which ensemble proceeds with simple, independent motion (the *decoupled* evolution).

Branch selection. If the control light is 0, the switch routes the coupled choreography to Register A and the decoupled routine to Register B (Path 1). If the control light is 1, the assignments are swapped: Register B receives the coupled choreography while Register A runs decoupled (Path 2). Because the two ensembles are separate, these routines can be applied side by side without interfering; the switch simply directs which routine lands where.

Superposition and interference. When the control light is prepared in a "both-at-once" state (a quantum superposition), the full system simultaneously follows the two histories: "A-coupled/B-decoupled" *and* "A-decoupled/B-coupled." Afterward, the control light is rotated back and read out, converting the quiet tug-of-war between the histories into a visible difference in outcomes. In effect, the circuit allows two alternative worldlines of the same system to overlap and *interfere*, revealing how the choice of choreography (who coordinates, who does not) changes collective behavior.

Efficient realization. An efficiency trick replaces two large, separate switches with a baseline-plus-correction strategy: first apply one full routine to both ensembles as a baseline; then—only if the control is 1—apply a single correction that transforms the baseline into the alternative history. Functionally this remains the same smart switch, packaged more compactly.

Bottom line. The controlled holosymmetric operator is a *coherent selector* of dynamical stories. It assigns complex coordination to one register and simplicity to the other, or the reverse—and, crucially, it can place those assignments into quantum superposition so their consequences can be compared through interference.

Joint State of the Dual Primordial Registers

Definition 2.113 (Joint State of the Dual Primordial Registers)**.** Let $\mathcal{H}_{\mathrm{PQS}}^{(n,A)}$ and $\mathcal{H}_{\mathrm{PQS}}^{(n,B)}$ denote the Hilbert spaces associated with two n-PQS registers—Register A and Register B—each comprising n Primordial Quantum Singularities. Define the dual-register Hilbert space as:

$$\mathcal{H}_{\mathrm{RR}}^{(2n)} := \mathcal{H}_{\mathrm{PQS}}^{(n,A)} \otimes \mathcal{H}_{\mathrm{PQS}}^{(n,B)}. \tag{2.71}$$

Then the state vector:

$$\boxed{|\Psi_{\mathrm{RR}}\rangle \in \mathcal{H}_{\mathrm{RR}}^{(2n)}} \tag{2.72}$$

is called the *joint quantum state of the dual PQS registers*. It encodes the full quantum configuration of both registers, including all modal amplitudes, internal correlations, and possible inter-register entanglement. This state serves as the initial condition for holosymmetric evolution in the interferometric framework.

Branch Decomposition by Control State

The holosymmetric Hilbert space $\mathcal{H}_{\mathrm{Holo}}$ decomposes naturally into two orthogonal branches, each determined by the control qubit and corresponding to a distinct dynamical pathway:

$$\mathcal{H}_{\mathrm{Holo}} = \mathcal{H}_{\mathrm{Holo}}^{A_G B_N} \oplus \mathcal{H}_{\mathrm{Holo}}^{A_N B_G},$$

where:

$$\boxed{\mathcal{H}_{\mathrm{Holo}}^{A_G B_N} := \mathrm{span}\{|\mathbf{0}\rangle_c\} \otimes \mathcal{H}_{\mathrm{RR}}^{(2n)},} \tag{2.73}$$

$$\boxed{\mathcal{H}_{\mathrm{Holo}}^{A_N B_G} := \mathrm{span}\{|\mathbf{1}\rangle_c\} \otimes \mathcal{H}_{\mathrm{RR}}^{(2n)}.} \tag{2.74}$$

The control state $|\mathbf{0}\rangle_c$ selects the operator $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$, in which Register A evolves under fully coupled dynamics $U_G^{(A)}$, and Register B evolves under generalized decoupled dynamics $U_N^{(B)}$. Conversely, control state $|\mathbf{1}\rangle_c$ activates the operator $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$, reversing this assignment.

Superposed Evolution. When the control qubit is placed in a coherent superposition,

$$|\phi_c\rangle = \alpha|\mathbf{0}\rangle_c + \beta|\mathbf{1}\rangle_c, \quad \alpha, \beta \in \mathbb{C},\ |\alpha|^2 + |\beta|^2 = 1,$$

the resulting evolution involves an entangled superposition of parallel dynamical trajectories:

$$|\Psi_{\mathrm{Holo}}\rangle = \alpha\,|\mathbf{0}\rangle_c \otimes \left(\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}|\Psi_{\mathrm{RR}}\rangle\right) + \beta\,|\mathbf{1}\rangle_c \otimes \left(\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}|\Psi_{\mathrm{RR}}\rangle\right).$$

This expresses a genuinely holosymmetric interference between relational evolutions applied across dual PQS registers—constituting a higher-order modality of dynamical synthesis.

2.3.2 Holosymmetry Component Operators

The complete holosymmetric transformation, denoted generically as $\mathbf{U}_{\mathrm{Holo}}$, is a unitary operator designed to act on the full emergent Hilbert space for holosymmetry,

$$\mathcal{H}_{\mathrm{Holo}} := \mathcal{H}_c \otimes \mathcal{H}_{\mathrm{PQS}}^{(n,A)} \otimes \mathcal{H}_{\mathrm{PQS}}^{(n,B)} \cong \mathbb{C}^{2^{4n+1}},$$

as defined in eq. (2.62). This space includes:

- the **control system** $\mathcal{H}_c$ (Definition 2.102),
- and the **dual register space** $\mathcal{H}_{\mathrm{RR}}^{(2n)} := \mathcal{H}_{\mathrm{PQS}}^{(n,A)} \otimes \mathcal{H}_{\mathrm{PQS}}^{(n,B)}$ (eq. (2.61)).

We now explore operators capable of enacting holosymmetric evolution, beginning with a controlled-path construction.

The Controlled Holosymmetric Operator ($\mathbf{U}_{\mathrm{ctrl}}$)

The first candidate for holosymmetric evolution is a *controlled unitary operator*, denoted $\mathbf{U}_{\mathrm{ctrl}}$. It selects between two distinct parasymmetric evolutionary paths—namely:

- $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$: coupled evolution on Register A and decoupled on Register B (eq. (2.69)),
- $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$: decoupled evolution on Register A and coupled on Register B (eq. (2.70)),

based on the state of the control qubit $\mathcal{H}_c$.

Definition 2.114 (Controlled Holosymmetric Operator)**.** The *Controlled Holosymmetric Operator* $\mathbf{U}_{\mathrm{ctrl}}$ is defined as the following unitary operator on the full holosymmetric Hilbert space $\mathcal{H}_{\mathrm{Holo}}$:

$$\boxed{\mathbf{U}_{\mathrm{ctrl}} := |\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}.} \tag{2.75}$$

Here:

- $|\mathbf{0}\rangle_c, |\mathbf{1}\rangle_c \in \mathcal{H}_c$ form the orthonormal basis of the control Hilbert space,
- $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$ and $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$ are the parallel path operators acting on the dual PQS register space $\mathcal{H}_{\mathrm{RR}}^{(2n)}$ (Definition 2.108).

This operator implements holosymmetric branching at the dynamical level: when the control system is in a coherent superposition $\alpha|\mathbf{0}\rangle_c + \beta|\mathbf{1}\rangle_c$, the global evolution of the system reflects a quantum superposition of distinct dynamical regimes across the two registers.

Definition 2.115 (Direct controlled-path realization of $\mathbf{U}_{\mathrm{ctrl}}$)**.** Instantiating eq. (2.75) with the path unitaries of eqs. (2.69) and (2.70) yields the following *two-branch multiplexer* on $\mathcal{H}_{\mathrm{Holo}}$ (eq. (2.62)):



The first (open–control) column realizes $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$ when the control is $|\mathbf{0}\rangle_c$ (eq. (2.73)), and the second (closed–control) column realizes $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$ when the control is $|\mathbf{1}\rangle_c$ (eq. (2.74)). The blocks $U_G^{(\odot)}$ and $U_N^{(\odot)}$ are the compiled multi–qubit evolutions from Definitions 2.66 and 2.85.

Definition 2.116 (Canonical synthesis with a single controlled correction)**.** Let

$$U_0 \;:=\; \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}, \qquad U_1 \;:=\; \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}, \qquad W \;:=\; U_0^\dagger U_1.$$

Then $\mathbf{U}_{\mathrm{ctrl}}$ admits the block form

$$\boxed{\mathbf{U}_{\mathrm{ctrl}} = \Big(|\mathbf{0}\rangle\langle\mathbf{0}|_c \otimes \mathbb{I} + |\mathbf{1}\rangle\langle\mathbf{1}|_c \otimes W \Big)\,(\mathbb{I}_c \otimes U_0),} \tag{2.76}$$

which is implemented by an *unconditional* U_0 followed by a *single* $|\mathbf{1}\rangle$–controlled W:



This synthesis is often depth–optimal when U_0 and U_1 are available as primitives and W can be compiled efficiently from them.

Definition 2.117 (Hadamard–conjugated wrapper for branch interference)**.** To realize the superposed evolution of Section 2.3.1, prepare and read out the control in the X basis:



The first H prepares $\alpha|\mathbf{0}\rangle_c + \beta|\mathbf{1}\rangle_c$, $\mathbf{U}_{\mathrm{ctrl}}$ enacts the branch–conditioned paths of eqs. (2.69) and (2.70), and the final H converts branch interference into Z–basis statistics on the control.

Remark 2.118 (Compilation notes)**.** (i) In Definition 2.115, the two branch blocks may be executed in parallel across the disjoint register wires, since they act on different tensor factors of $\mathcal{H}_{\mathrm{RR}}^{(2n)}$. (ii) In Definition 2.116, the controlled W acts jointly on both registers; when $U_G^{(\odot)}$ and $U_N^{(\odot)}$ are Trotterized, W inherits the same layered structure (local rotations, XY exchanges, global flags) and can be compiled by palindromic cancellation of common layers.

Operational role. The controlled holosymmetric operator $\mathbf{U}_{\mathrm{ctrl}}$ is a coherent, register–selective path selector acting on the full holosymmetric space $\mathcal{H}_{\mathrm{Holo}}$ (eq. (2.62)). By definition (eq. (2.75)), the control basis state fixes which parallel path acts on the dual register factor $\mathcal{H}_{\mathrm{RR}}^{(2n)}$: the branch $|\mathbf{0}\rangle_c$ selects $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$ and the branch $|\mathbf{1}\rangle_c$ selects $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$, with the branch subspaces given in eqs. (2.73) and (2.74).

Branch behavior. Under $|\mathbf{0}\rangle_c$, the operator implements *Path 1* (eq. (2.69)): Register A is driven by the unified global coupled evolution $U_G^{(A)}$, while Register B undergoes the composite decoupled evolution $U_N^{(B)}$ (Definitions 2.66 and 2.85). Under $|\mathbf{1}\rangle_c$, *Path 2* is enacted (eq. (2.70)), reversing the assignments. Because these lifted gates act on disjoint tensor factors, the two large register operations commute and may be executed side–by–side without inter–register crosstalk (Definition 2.108).

Superposed evolution and interference. Preparing the control in a coherent superposition yields the superposed evolution of Section 2.3.1: the global state evolves as a controlled superposition of the two complete histories, $|\mathbf{0}\rangle_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} |\Psi_{\mathrm{RR}}\rangle$ and $|\mathbf{1}\rangle_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} |\Psi_{\mathrm{RR}}\rangle$. Conjugating $\mathbf{U}_{\mathrm{ctrl}}$ by Hadamards on the control (prepare and read in the X basis) converts branch interference into observable Z–basis statistics on the control, operationalizing holosymmetric comparison between the two dynamical laws (eq. (2.75)).

Canonical synthesis. A depth–efficient realization follows the block form of eq. (2.76): apply one path unconditionally as a baseline ($U_0 = \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$), then apply the single controlled correction $W = U_0^\dagger U_1$ (with $U_1 = \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$) conditioned on $|\mathbf{1}\rangle_c$. When $U_G^{(\odot)}$ and $U_N^{(\odot)}$ are compiled in layered/Trotterized form, W inherits the same structure and can be synthesized by canceling common layers (Definitions 2.66 and 2.85).

Summary. The controlled holosymmetric operator is a coherent selector of register–level evolutions: it assigns the globally coupled parasymmetric dynamics to one register and the decoupled dynamics to the other, or vice versa, according to the control branch (eqs. (2.69), (2.70) and (2.75)). With the control in superposition, the construction promotes interference from states to *assignments of dynamics*, enabling holosymmetric tests and comparisons on $\mathcal{H}_{\mathrm{Holo}}$ that respect the dual–register tensor structure (eq. (2.62)).

Block Matrix Form

Relative to the control-resolved branch decomposition

$$\mathcal{H}_{\mathrm{Holo}} = \mathcal{H}_{[\mathbf{0}]}^{A_G B_N} \oplus \mathcal{H}_{[\mathbf{1}]}^{A_N B_G},$$

(Section 2.3.1), where:

$$\mathcal{H}_{\text{Holo}}^{A_G B_N} := \text{span}\{|\mathbf{0}\rangle_c\} \otimes \mathcal{H}_{\text{RR}}^{(2n)},$$
$$\mathcal{H}_{\text{Holo}}^{A_N B_G} := \text{span}\{|\mathbf{1}\rangle_c\} \otimes \mathcal{H}_{\text{RR}}^{(2n)},$$

the controlled holosymmetric operator $\mathbf{U}_{\text{ctrl}}$ (Definition 2.114) takes the following block-diagonal form:

$$\boxed{\mathbf{U}_{\text{ctrl}} = \begin{pmatrix} \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} & \mathbf{0} \\ \mathbf{0} & \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} \end{pmatrix}.} \tag{2.77}$$

Here:

- $\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} := U_G^{(A)} \cdot U_N^{(B)}$ is the lifted operator implementing **coupled** evolution on Register A and **decoupled** on Register B (eq. (2.69)),
- $\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} := U_N^{(A)} \cdot U_G^{(B)}$ is the complementary lifted operator for **decoupled** A and **coupled** B (eq. (2.70)),
- $\mathbf{0}$ denotes the zero operator acting between orthogonal control branches.

Interpretation. This block structure reflects the conditional application of distinct parasymmetric dynamics to the PQS system depending on the control register. When the control is in a definite state $|\mathbf{0}\rangle_c$ or $|\mathbf{1}\rangle_c$, the evolution is purely confined to the corresponding block. In coherent superposition, both blocks contribute simultaneously, realizing holosymmetric interference between two full dynamical pathways.

Justification. From the definition:

$$\mathbf{U}_{\text{ctrl}} := |\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G},$$

we are expressing a **controlled unitary** operator acting on the total holosymmetric Hilbert space:

$$\mathcal{H}_{\text{Holo}} = \mathcal{H}_c \otimes \mathcal{H}_{\text{RR}}^{(2n)} \cong \mathcal{H}_{\text{Holo}}^{A_G B_N} \oplus \mathcal{H}_{\text{Holo}}^{A_N B_G}.$$

Then, under this orthogonal decomposition, $\mathbf{U}_{\text{ctrl}}$ acts as:

$$\mathbf{U}_{\text{ctrl}} = \begin{pmatrix} \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} & \mathbf{0} \\ \mathbf{0} & \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} \end{pmatrix},$$

which is the canonical matrix representation of a controlled unitary in the computational basis for $\mathcal{H}_c$. Each block corresponds precisely to the action conditioned on the control state $|\mathbf{0}\rangle_c$ or $|\mathbf{1}\rangle_c$, respectively.

Action on Control Basis States

If the initial state of the dual PQS registers is $|\Psi_{\mathrm{RR}}\rangle \in \mathcal{H}_{\mathrm{RR}}^{(2n)}$, the action of the controlled holosymmetric operator $\mathbf{U}_{\mathrm{ctrl}}$ on product states with definite control states is:

$$\mathbf{U}_{\mathrm{ctrl}}\left(|\mathbf{0}\rangle_c \otimes |\Psi_{\mathrm{RR}}\rangle\right) = |\mathbf{0}\rangle_c \otimes \left(\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}|\Psi_{\mathrm{RR}}\rangle\right),$$
$$\mathbf{U}_{\mathrm{ctrl}}\left(|\mathbf{1}\rangle_c \otimes |\Psi_{\mathrm{RR}}\rangle\right) = |\mathbf{1}\rangle_c \otimes \left(\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}|\Psi_{\mathrm{RR}}\rangle\right).$$

Interpretation of the Operator $\mathbf{U}_{\mathrm{ctrl}}$

The Controlled Holosymmetric Operator $\mathbf{U}_{\mathrm{ctrl}}$ functions as a *quantum multiplexer*—a programmable quantum switch that places the dual PQS registers $\mathcal{H}_{\mathrm{RR}}^{(2n)}$ under the coherent governance of a two-level quantum control system $\mathcal{H}_c$. This control enables two distinct operational regimes: one classical, one intrinsically quantum.

1. Classical Analogue: A Deterministic Switch When the control system is initialized in a definite computational basis state—either $|\mathbf{0}\rangle_c$ or $|\mathbf{1}\rangle_c$—its role mirrors that of a classical 'IF-THEN-ELSE' logic gate:

- *If Control is $|\mathbf{0}\rangle_c$:* The operator deterministically applies the evolution $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$, corresponding to **coupled** dynamics on Register A and **decoupled** dynamics on Register B.
- *If Control is $|\mathbf{1}\rangle_c$:* The operator selects the complementary evolution $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$, applying **decoupled** dynamics on Register A and **coupled** on Register B.

In this classical-control regime, the system follows a single, definite pathway with no entanglement generated between control and register subsystems. The result is a clean partitioning of dynamics based on a known control input—a classical branching architecture instantiated in quantum logic.

2. Quantum Engine: A Superposition of Histories The true power of $\mathbf{U}_{\mathrm{ctrl}}$ is unleashed when the control system is prepared in a *superposition* of its basis states, such as $\frac{1}{\sqrt{2}}(|\mathbf{0}\rangle_c + |\mathbf{1}\rangle_c)$. In this regime, the operator transcends its classical analogue and becomes an engine for generating coherent quantum superpositions.

- *Quantum Parallelism:* Rather than selecting a single dynamical branch, the operator enacts *both* evolutionary pathways in superposition. The system evolves into a coherent sum:

$$\mathbf{U}_{\mathrm{ctrl}}\left(\frac{1}{\sqrt{2}}(|\mathbf{0}\rangle_c + |\mathbf{1}\rangle_c) \otimes |\Psi_{\mathrm{RR}}\rangle\right) = \frac{1}{\sqrt{2}}\left(|\mathbf{0}\rangle_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}|\Psi_{\mathrm{RR}}\rangle + |\mathbf{1}\rangle_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}|\Psi_{\mathrm{RR}}\rangle\right).$$

  This is a manifestation of quantum parallelism: multiple dynamical histories coexist within a single entangled state.

- *Entanglement as a Record:* The resulting state is necessarily *entangled* between the control and register subsystems. The $|\mathbf{0}\rangle_c$ component is correlated with the outcome of evolution along the $A_G B_N$ branch, and the $|\mathbf{1}\rangle_c$ component with the $A_N B_G$ branch. This entanglement encodes a *coherent record* of the alternative quantum histories, incarnated as statistical mixtures and as a unified quantum object.

Measuring the control qubit after evolution collapses this superposition, forcing the system into one of the distinct branches and revealing the corresponding register state. In this sense, $\mathbf{U}_{\text{ctrl}}$ serves as a foundational mechanism for generating and manipulating *entangled history states*, a key theoretical resource in quantum simulation, algorithm design, and holosymmetric inference.

Structural Insight: Architect of a Quantum Interferometer

Structurally, the controlled holosymmetric operator $\mathbf{U}_{\text{ctrl}}$ does more than simply toggle between two dynamical evolutions; it serves as the foundational component for constructing a *quantum interferometer*. Its essential function is to initiate a *controlled holosymmetric bifurcation*—a quantum fork that splits the register system into two coherently evolving trajectories.

This bifurcation represents the first phase of a two-stage interferometric architecture:

1. *Path Creation (Act I):* Enacted by $\mathbf{U}_{\text{ctrl}}$, this phase leverages a superposed control qubit to generate two simultaneous quantum histories. One branch evolves under $\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}$, the other under $\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}$. Each history is coherently labeled by its respective control state: $|\mathbf{0}\rangle_c$ or $|\mathbf{1}\rangle_c$.

2. *Path Recombination and Interference (Act II):* These bifurcated trajectories are not endpoints but resources. A subsequent operation—analogous to the second beam splitter in a Mach-Zehnder interferometer—is used to recombine the control paths. When the two branches are coherently merged, quantum interference arises.

The potency of this structure lies in the interference outcome, which is exquisitely sensitive to the *path differential* between the two dynamical evolutions. Specifically, it depends on how $\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}$ and $\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}$ differ in phase and amplitude transformations.

Thus, $\mathbf{U}_{\text{ctrl}}$ is not just a switch—it is the architectural nucleus that constructs the key interferometric resource: a coherent superposition of divergent evolutions. By enabling their subsequent recombination, the framework extracts holosymmetric interference signatures that act as sensitive probes into the structural contrast between competing dynamical regimes acting on the quantum system.

Interferometer Component	Holosymmetry Analogue	Function
Initial Light Beam	$\|\psi\rangle_c \otimes \|\Psi_{\mathrm{RR}}\rangle$	System prior to path bifurcation
First Beam Splitter	$\mathbf{U}_{\mathrm{ctrl}}$	Creates controlled bifurcation into entangled dynamical branches
Path A vs. Path B	$\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$ vs. $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$	Distinct holosymmetric evolutions of the PQS registers
Phase Shifter	$\Delta\hat{\mathbf{U}} := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} - \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$	Encodes relative phase/amplitude difference between pathways
Second Beam Splitter	$\mathbf{U}_{\mathrm{recomb}}$	Coherently recombines the bifurcated control branches
Interference at Detectors	Final measurement on $\mathcal{H}_c$	Reveals holosymmetric interference signature

Table 2.1: Mapping Between Interferometric Components and Holosymmetric Structures

The Hadamard-Style Holosymmetric Combination ($\mathbf{U}_{\mathrm{Had}}$)

The operator $\mathbf{U}_{\mathrm{Had}}$ embodies a refined form of holosymmetric evolution that is explicitly engineered to expose quantum interference between the distinct dynamical histories associated with $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$ and $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$ (eqs. (2.69) and (2.70)). By conjugating the Controlled Holosymmetric Operator $\mathbf{U}_{\mathrm{ctrl}}$ with Hadamard operations on the control qubit, $\mathbf{U}_{\mathrm{Had}}$ both prepares and recombines coherent control superpositions.

Definition 2.119 (Hadamard-Conjugated Holosymmetry Operator)**.** We define the *Hadamard-Conjugated Holosymmetry* Operator, $\mathbf{U}_{\mathrm{Had}}$, which acts on the full holosymmetric Hilbert space $\mathcal{H}_{\mathrm{Holo}} = \mathcal{H}_c \otimes \mathcal{H}_{\mathrm{RR}}^{(2n)}$.

$$
\begin{aligned}
\mathbf{U}_{\mathrm{Had}} &:= (H_c \otimes I_{\mathrm{RR}}^{(2n)})\ \mathbf{U}_{\mathrm{ctrl}}\ (H_c \otimes I_{\mathrm{RR}}^{(2n)}) \\
&= (H_c \otimes I_{\mathrm{RR}}^{(2n)})\ \left[|\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}\right]\ (H_c \otimes I_{\mathrm{RR}}^{(2n)}) \\
&= \left[(H_c|\mathbf{0}\rangle_c)(\langle\mathbf{0}|_c H_c) \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + (H_c|\mathbf{1}\rangle_c)(\langle\mathbf{1}|_c H_c) \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}\right] \\
&= \left[|+\rangle_c\langle+|_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + |-\rangle_c\langle-|_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}\right],
\end{aligned}
\tag{2.78}
$$

where:

- H_c is the Hadamard operator acting on the control Hilbert space $\mathcal{H}_c$,

- $I_{\mathrm{RR}}^{(2n)}$ is the identity operator on the dual register space $\mathcal{H}_{\mathrm{RR}}^{(2n)}$ (eq. (2.61)),
- $\mathbf{U}_{\mathrm{ctrl}}$ is the Contracted Holosymmetric Operator (Definition 2.114).

The purpose of this operator is to enact a complete interferometric cycle: it prepares a coherent control superposition, evolves the system through bifurcated holosymmetric dynamics, and recombines the pathways to reveal interference effects that encode the contrast between the alternative evolutions. Since both the Hadamard operator and the controlled unitary operator are unitary ($H_c = H_c^\dagger$, $H_c^2 = I_c$), the composite operator $\mathbf{U}_{\mathrm{Had}}$ is also manifestly unitary.

Block Matrix Form. In the computational basis $\{|\mathbf{0}\rangle_c, |\mathbf{1}\rangle_c\}$ of the control Hilbert space $\mathcal{H}_c$, the Hadamard operator is given by:

$$H_c = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}.$$

Given that the controlled holosymmetric operator $\mathbf{U}_{\mathrm{ctrl}}$ acts as a block-diagonal matrix:

$$\mathbf{U}_{\mathrm{ctrl}} = \begin{pmatrix} \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} & \mathbf{0} \\ \mathbf{0} & \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} \end{pmatrix},$$

its Hadamard-conjugated form $\mathbf{U}_{\mathrm{Had}} := (H_c \otimes I)\, \mathbf{U}_{\mathrm{ctrl}}\, (H_c \otimes I)$ can be computed explicitly as:

$$\mathbf{U}_{\mathrm{Had}} = \frac{1}{2} \begin{pmatrix} \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} & \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} - \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} \\ \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} - \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} & \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} \end{pmatrix}.$$

Each entry in this matrix is itself an operator acting on the dual register space $\mathcal{H}_{\mathrm{RR}}^{(2n)}$. The off-diagonal blocks encode the interference between the two holosymmetric branches, and vanish only when the dynamics $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$ and $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$ are identical. The diagonal blocks represent the coherent superposition of both dynamical evolutions.

Action on Control Basis States. Let $|\Psi_{\mathrm{RR}}\rangle \in \mathcal{H}_{\mathrm{RR}}^{(2n)}$ be an arbitrary state of the dual PQS registers. The action of the Hadamard-conjugated holosymmetric operator $\mathbf{U}_{\mathrm{Had}}$ on the product basis states of the full holosymmetric Hilbert space $\mathcal{H}_{\mathrm{Holo}} = \mathcal{H}_c \otimes \mathcal{H}_{\mathrm{RR}}^{(2n)}$ follows directly from its definition:

$$\mathbf{U}_{\mathrm{Had}} := (H_c \otimes I_{\mathrm{RR}}^{(2n)})\ \mathbf{U}_{\mathrm{ctrl}}\ (H_c \otimes I_{\mathrm{RR}}^{(2n)}).$$

Case 1: Action on $|\mathbf{0}\rangle_c \otimes |\Psi_{\mathrm{RR}}\rangle$

$$\begin{aligned}
\mathbf{U}_{\mathrm{Had}}\,(|\mathbf{0}\rangle_c \otimes |\Psi_{\mathrm{RR}}\rangle) &= (H_c \otimes I_{\mathrm{RR}}^{(2n)})\ \mathbf{U}_{\mathrm{ctrl}}\ (H_c|\mathbf{0}\rangle_c \otimes |\Psi_{\mathrm{RR}}\rangle) \\
&= (H_c \otimes I)\ \mathbf{U}_{\mathrm{ctrl}}\ \left(\tfrac{1}{\sqrt{2}}(|\mathbf{0}\rangle_c + |\mathbf{1}\rangle_c) \otimes |\Psi_{\mathrm{RR}}\rangle\right) \\
&= (H_c \otimes I)\ \tfrac{1}{\sqrt{2}}\left(|\mathbf{0}\rangle_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}|\Psi_{\mathrm{RR}}\rangle + |\mathbf{1}\rangle_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}|\Psi_{\mathrm{RR}}\rangle\right) \\
&= \tfrac{1}{2}\left[(|\mathbf{0}\rangle_c + |\mathbf{1}\rangle_c) \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}|\Psi_{\mathrm{RR}}\rangle + (|\mathbf{0}\rangle_c - |\mathbf{1}\rangle_c) \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}|\Psi_{\mathrm{RR}}\rangle\right] \\
&= \tfrac{1}{2}\Big[|\mathbf{0}\rangle_c \otimes \left(\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}\right)|\Psi_{\mathrm{RR}}\rangle \\
&\qquad + |\mathbf{1}\rangle_c \otimes \left(\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} - \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}\right)|\Psi_{\mathrm{RR}}\rangle\Big].
\end{aligned}$$

Case 2: Action on $|\mathbf{1}\rangle_c \otimes |\Psi_{\mathrm{RR}}\rangle$

$$\begin{aligned}
\mathbf{U}_{\mathrm{Had}}\left(|\mathbf{1}\rangle_c \otimes |\Psi_{\mathrm{RR}}\rangle\right) &= \left(H_c \otimes I_{\mathrm{RR}}^{(2n)}\right)\, \mathbf{U}_{\mathrm{ctrl}}\ \left(H_c|\mathbf{1}\rangle_c \otimes |\Psi_{\mathrm{RR}}\rangle\right) \\
&= \left(H_c \otimes I\right)\, \mathbf{U}_{\mathrm{ctrl}}\ \left(\tfrac{1}{\sqrt{2}}(|\mathbf{0}\rangle_c - |\mathbf{1}\rangle_c) \otimes |\Psi_{\mathrm{RR}}\rangle\right) \\
&= \tfrac{1}{2}\left[(|\mathbf{0}\rangle_c + |\mathbf{1}\rangle_c) \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}|\Psi_{\mathrm{RR}}\rangle - (|\mathbf{0}\rangle_c - |\mathbf{1}\rangle_c) \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}|\Psi_{\mathrm{RR}}\rangle\right] \\
&= \tfrac{1}{2}\Big[|\mathbf{0}\rangle_c \otimes \left(\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} - \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}\right)|\Psi_{\mathrm{RR}}\rangle \\
&\qquad + |\mathbf{1}\rangle_c \otimes \left(\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}\right)|\Psi_{\mathrm{RR}}\rangle\Big].
\end{aligned}$$

General Action The general action of the Hadamard-conjugated holosymmetric operator is then given by:

$$\begin{aligned}
\mathbf{U}_{\mathrm{Had}}\left((\alpha|\mathbf{0}\rangle_c + \beta|\mathbf{1}\rangle_c) \otimes |\Psi_{\mathrm{RR}}\rangle\right) = \tfrac{1}{2}|\mathbf{0}\rangle_c \otimes \left[(\alpha+\beta)\,\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + (\alpha-\beta)\,\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}\right]|\Psi_{\mathrm{RR}}\rangle \\
+ \tfrac{1}{2}|\mathbf{1}\rangle_c \otimes \left[(\alpha+\beta)\,\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} + (\alpha-\beta)\,\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}\right]|\Psi_{\mathrm{RR}}\rangle.
\end{aligned} \tag{2.79}$$

Interpretation and Significance of $\mathbf{U}_{\mathrm{Had}}$ The operator $\mathbf{U}_{\mathrm{Had}}$ elevates the concept of holosymmetry from a simple conditional switch to a complete, self-contained *quantum interferometer*. Its structure, defined by conjugating the controlled evolution $\mathbf{U}_{\mathrm{ctrl}}$ with Hadamard gates on the control system, is the canonical circuit for probing the interference between alternative quantum histories. The entire operation can be understood as a three-act process.

The Quantum Interferometer Circuit The action of $\mathbf{U}_{\mathrm{Had}}$ on an initial state like $|\mathbf{0}\rangle_c \otimes |\Psi_{\mathrm{RR}}\rangle$ perfectly maps to the function of a Mach-Zehnder interferometer.

Act I: Path Creation (First H_c) The initial Hadamard gate acts as the *first beam splitter*. It transforms the control state from a definite path ($|\mathbf{0}\rangle_c$) into an equal superposition ($\frac{1}{\sqrt{2}}(|\mathbf{0}\rangle_c + |\mathbf{1}\rangle_c)$), preparing the system to traverse both evolutionary pathways simultaneously.

Act II: Differential Evolution ($\mathbf{U}_{\mathrm{ctrl}}$) This is the core of the experiment, analogous to the two distinct arms of the interferometer. The system evolves under $\mathbf{U}_{\mathrm{ctrl}}$, which applies the evolution $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$ to the $|\mathbf{0}\rangle_c$ component and $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$ to the $|\mathbf{1}\rangle_c$ component. This step imprints the differing dynamics of the two histories onto the state, creating a complex entanglement between the control system and the registers.

Act III: Path Recombination (Second H_c) The final Hadamard gate acts as the *second beam splitter*. It interferes the two paths by transforming the path information (held in the superposition basis) back into the computational basis ($\{|\mathbf{0}\rangle_c, |\mathbf{1}\rangle_c\}$).

Revealing Interference Through Measurement This final transformation is what makes the quantum interference directly observable. A direct calculation shows that the final state of the system is:

$$\frac{1}{2}\left[|\mathbf{0}\rangle_c \otimes (\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G})|\Psi_{\mathrm{RR}}\rangle + |\mathbf{1}\rangle_c \otimes (\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} - \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G})|\Psi_{\mathrm{RR}}\rangle\right]$$

The interpretation of this result is remarkably direct:

- The component of the final state associated with $|\mathbf{0}\rangle_c$ is proportional to the *sum* of the two path evolutions. Its amplitude represents their *constructive interference*.
- The component associated with $|\mathbf{1}\rangle_c$ is proportional to the *difference* between the path evolutions. Its amplitude represents their *destructive interference*.

A measurement of the control qubit in the computational basis now reveals a holistic property of the system that depends on all possible histories at once.

Conceptual Significance: From Branching to Interferometric Emergence This operator realizes a profound shift from a simple branching of histories (as with $\mathbf{U}_{\text{ctrl}}$) to a mechanism for *interferometric emergence*. The foundational principle of duality—the existence of two distinct but related parasymmetric evolutions—is no longer just a premise but becomes a measurable quantity.

The interference pattern, observable in the final probabilities of measuring $|\mathbf{0}\rangle_c$ or $|\mathbf{1}\rangle_c$, is a direct probe into the relational similarity of the two dynamical histories. The operator $\mathbf{U}_{\text{Had}}$ thus provides the means for a quantum mechanical exploration of superposed causal pathways, making interference the primary tool for understanding how a single reality can emerge from a superposition of possibilities.

Definition 2.120 (Direct realization of $\mathbf{U}_{\text{Had}} = (H_c \otimes I)\,\mathbf{U}_{\text{ctrl}}\,(H_c \otimes I)$)**.** Using the controlled-path form of $\mathbf{U}_{\text{ctrl}}$ from eq. (2.75) with paths of eqs. (2.69) and (2.70), the Hadamard–conjugated operator of eq. (2.78) is:



The open–control column applies $\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}$ on branch $|\mathbf{0}\rangle_c$; the closed–control column applies $\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}$ on branch $|\mathbf{1}\rangle_c$. The outer Hadamards implement the preparation and recombination in the control X basis, as in eq. (2.78).

Definition 2.121 (Canonical synthesis: baseline path plus controlled correction)**.** Let $U_0 := \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}$ and $U_1 := \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}$, and define $W := U_0^\dagger U_1$ (cf. eqs. (2.69) and (2.70)). Using the block form in eq. (2.75) and eq. (2.78),

$$\begin{aligned}\mathbf{U}_{\text{Had}} &= (H_c \otimes I)\Big[\,|\mathbf{0}\rangle\langle\mathbf{0}|_c \otimes U_0 + |\mathbf{1}\rangle\langle\mathbf{1}|_c \otimes U_1\,\Big](H_c \otimes I) \\ &= (H_c \otimes I)\Big[\,\mathbb{I}_c \otimes U_0\,\Big]\Big[\,|1\rangle\langle 1|_c \otimes W + |0\rangle\langle 0|_c \otimes \mathbb{I}\,\Big](H_c \otimes I).\end{aligned}$$

A depth–efficient circuit therefore is:



Here U_0 acts unconditionally on the dual-register factor, and the single controlled-W column enacts the alternative path on the $|\mathbf{1}\rangle_c$ branch; the surrounding H gates realize preparation/recombination as in eq. (2.78).

Definition 2.122 (Interference readout on the control (optional))**.** To convert path interference into control-basis statistics, append a Z-measurement on the control after Definition 2.120 or Definition 2.121:



The measurement outcomes on the control encode the interference between $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$ and $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$ prepared and recombined by the Hadamards, as prescribed by eq. (2.78).

Deconstruction of the $\mathbf{U}_{\mathbf{Had}}$ Operator Circuit

Structural Stage	Mathematical Realization	Function / Role
Initial State	$\|\mathbf{0}\rangle_c \otimes \|\Psi_{\mathrm{RR}}\rangle$	Defines the starting point of the interferometric process with the control system in a definite basis state.
Act I: Path Creation	$H_c \otimes I_{\mathrm{RR}}^{(2n)}$	Acts as the first "beam splitter." Prepares the control qubit in a superposition, enabling simultaneous traversal of both dynamical branches.
Act II: Path Evolution	$\mathbf{U}_{\mathrm{ctrl}}$	Applies branch-dependent dynamics: $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$ to the $\|\mathbf{0}\rangle_c$ component and $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$ to the $\|\mathbf{1}\rangle_c$ component.
Act III: Path Recombination	$H_c \otimes I_{\mathrm{RR}}^{(2n)}$	Acts as the second "beam splitter." Recombines the branches and translates their relative dynamics into measurable interference.
Composite Operator	$\mathbf{U}_{\mathrm{Had}}$	Represents the full interferometer circuit: a holosymmetric evolution that coherently superposes and recombines quantum histories.
Interference Readout	Measure in the $\{\|\mathbf{0}\rangle_c, \|\mathbf{1}\rangle_c\}$ basis	Reveals the interference pattern. The outcome amplitudes reflect the sum and difference of the path-dependent evolutions.

Table 2.2: Deconstruction of the $\mathbf{U}_{\mathrm{Had}}$ Operator Circuit

2.4 The Interferometer Model of Holosymmetric Evolution

The Interferometer Model implements a holosymmetric recursive transformation of modal evolution based on the logic:

Prepare Superposition → Evolve Conditional Paths → Recombine and Measure

Let the initial holosymmetric state be given by:

$$|\Psi_{\text{init}}\rangle := |\mathbf{0}\rangle_c \otimes |\Psi\rangle_{\text{RR}},$$

where $|\Psi\rangle_{\text{RR}} \in \mathcal{H}_{\text{RR}}^{(2n)}$ is an arbitrary normalized PQS configuration.

The model is built from the following elementary components:

(i) Preparation Gate $\mathbf{U}_{\text{prep}} \in \mathcal{U}(\mathcal{H}_c \otimes \mathcal{H}_{\text{RR}})$ The **preparation gate** is a unitary transformation that acts only on the control register $\mathcal{H}_c \cong \mathbb{C}^2$, and is tensored trivially with the identity on the register system $\mathcal{H}_{\text{RR}}^{(2n)}$. It is defined by:

$$\mathbf{U}_{\text{prep}} := (U_c^{\text{prep}} \otimes I_{\text{RR}}),$$

where:

$$U_c^{\text{prep}} \in \text{SU}(2) \quad \text{and} \quad I_{\text{RR}} := \mathbb{I}_{2^{4n}}.$$

In matrix form, any $U_c^{\text{prep}} \in \text{SU}(2)$ may be expressed as:

$$U_c^{\text{prep}} = \begin{pmatrix} \alpha & -\beta^* \\ \beta & \alpha^* \end{pmatrix}, \qquad \text{with} \quad |\alpha|^2 + |\beta|^2 = 1.$$

The action of $\mathbf{U}_{\text{prep}}$ on the initial state produces:

$$\mathbf{U}_{\text{prep}} \left(|\mathbf{0}\rangle_c \otimes |\Psi\rangle_{\text{RR}} \right) = \left(\alpha \, |\mathbf{0}\rangle_c + \beta \, |\mathbf{1}\rangle_c \right) \otimes |\Psi\rangle_{\text{RR}}.$$

This generalizes the **Hadamard preparation**, which corresponds to the specific choice:

$$U_c^{\text{prep}} = H := \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}, \quad \Rightarrow \quad \mathbf{U}_{\text{prep}}^{(\text{Had})} = (H \otimes I_{\text{RR}}).$$

In this case:

$$\mathbf{U}_{\text{prep}}^{(\text{Had})} \left(|\mathbf{0}\rangle_c \otimes |\Psi\rangle_{\text{RR}} \right) = \frac{1}{\sqrt{2}} \left(|\mathbf{0}\rangle_c + |\mathbf{1}\rangle_c \right) \otimes |\Psi\rangle_{\text{RR}}.$$

More generally, by selecting arbitrary $\alpha, \beta \in \mathbb{C}$ subject to unitarity, the gate $\mathbf{U}_{\text{prep}}$ allows initialization of the control qubit into any point on the Bloch sphere, enabling fine-grained control over the interference landscape.

(ii) Controlled Evolution Operator $\mathbf{U}_{\text{ctrl}}$ Defined as:

$$\mathbf{U}_{\text{ctrl}} := |\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G},$$

which branches the system into two parasymmetric dynamical pathways based on the control register state.

(iii) Readout Gate $\mathbf{U}_{\text{read}} \in \mathcal{U}(\mathcal{H}_c \otimes \mathcal{H}_{\text{RR}})$ The **readout gate** defines the basis in which interference will be observed. It acts only on the control register and is tensored with the identity on the register space:

$$\mathbf{U}_{\text{read}} := (U_c^{\text{read}} \otimes I_{\text{RR}}),$$

with:

$$U_c^{\text{read}} = \begin{pmatrix} \gamma & -\delta^* \\ \delta & \gamma^* \end{pmatrix}, \qquad |\gamma|^2 + |\delta|^2 = 1.$$

Applied to the general superposition:

$$|\chi\rangle_c \otimes |\Phi\rangle_{\text{RR}} = (a\,|\mathbf{0}\rangle_c + b\,|\mathbf{1}\rangle_c) \otimes |\Phi\rangle_{\text{RR}},$$

the output is:

$$(\gamma a - \delta^* b)\,|\mathbf{0}\rangle_c \otimes |\Phi\rangle_{\text{RR}} + (\delta a + \gamma^* b)\,|\mathbf{1}\rangle_c \otimes |\Phi\rangle_{\text{RR}}.$$

Choosing:

$$U_c^{\text{read}} = H := \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \quad \Rightarrow \quad \mathbf{U}_{\text{read}}^{(\text{Had})} = (H \otimes I_{\text{RR}})$$

projects the output into the symmetric/antisymmetric basis.

Geometric Interpretation: The readout gate selects a measurement axis on the Bloch sphere, thereby choosing which interference modes become observable.

Operational Role: As the final gate before measurement, $\mathbf{U}_{\text{read}}$ determines which quantum interference patterns are revealed in the control register.

2.4.1 (1) Symmetric Interferometer Model

Definition 2.123 (Symmetric Interferometer Holosymmetric Operator)**.** Let $H_c \in \text{SU}(2)$ denote the Hadamard gate acting on the control register $\mathcal{H}_c \cong \mathbb{C}^2$, and let $\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}, \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} \in \mathcal{U}(\mathcal{H}_{\text{RR}}^{(2n)})$ be the evolution operators for the two complementary parasymmetric dynamical assignments on the PQS registers.

Then the *Symmetric Holosymmetric Interferometric Operator* is defined as:

$$\mathbf{U}_{\text{holo}}^{(\text{sym})} := \mathbf{U}_{\text{Had}} \cdot \mathbf{U}_{\text{ctrl}} \cdot \mathbf{U}_{\text{Had}},$$

where:

- The Hadamard gate on the composite system is:

$$\mathbf{U}_{\text{Had}} := H_c \otimes I_{\text{RR}}, \quad \text{with} \quad H_c := \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix},$$

- The Controlled Holosymmetric Operator is:

$$\mathbf{U}_{\text{ctrl}} := |\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}.$$

Action on the initial state:
Let the initial state of the total system be:

$$|\Psi_{\text{init}}\rangle := |\mathbf{0}\rangle_c \otimes |\Psi_{\text{RR}}\rangle,$$

where $|\Psi_{\text{RR}}\rangle \in \mathcal{H}_{\text{RR}}^{(2n)}$ is an arbitrary normalized joint PQS state.
Then:

1. **First Hadamard:**

$$\mathbf{U}_{\text{Had}} |\Psi_{\text{init}}\rangle = \frac{1}{\sqrt{2}} (|\mathbf{0}\rangle_c + |\mathbf{1}\rangle_c) \otimes |\Psi_{\text{RR}}\rangle.$$

2. **Controlled evolution:**

$$\mathbf{U}_{\text{ctrl}} \circ \mathbf{U}_{\text{Had}} |\Psi_{\text{init}}\rangle = \frac{1}{\sqrt{2}} \left(|\mathbf{0}\rangle_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} |\Psi_{\text{RR}}\rangle + |\mathbf{1}\rangle_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} |\Psi_{\text{RR}}\rangle \right).$$

3. **Second Hadamard:**

$$\mathbf{U}_{\text{holo}}^{(\text{sym})} |\Psi_{\text{init}}\rangle = \frac{1}{2} \left[|\mathbf{0}\rangle_c \otimes \left(\hat{\mathbf{U}}_+ |\Psi_{\text{RR}}\rangle \right) + |\mathbf{1}\rangle_c \otimes \left(\hat{\mathbf{U}}_- |\Psi_{\text{RR}}\rangle \right) \right],$$

where:

$$\hat{\mathbf{U}}_+ := \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}, \quad \hat{\mathbf{U}}_- := \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} - \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}.$$

Interpretation. The operator $\mathbf{U}_{\text{holo}}^{(\text{sym})}$ coherently prepares the control system in a symmetric superposition, evolves the dual PQS register along two distinct conditional paths, and recombines these paths to yield a final state in which the control register encodes the modal interference structure between the two pathways.

Measuring the control qubit in the computational basis reveals whether the interference is constructive ($|\mathbf{0}\rangle_c$) or destructive ($|\mathbf{1}\rangle_c$), corresponding to the interference norms $\|\hat{\mathbf{U}}_+ |\Psi_{\text{RR}}\rangle\|^2$ and $\|\hat{\mathbf{U}}_- |\Psi_{\text{RR}}\rangle\|^2$ respectively.

Symmetric interferometer.

$$\mathbf{U}_{\text{holo}}^{(\text{sym})} = \mathbf{U}_{\text{Had}} \, \mathbf{U}_{\text{ctrl}} \, \mathbf{U}_{\text{Had}}$$







2.4.2 (2) Asymmetric Interferometer Model

Definition 2.124 (Asymmetric Interferometer Holosymmetric Operator)**.** Let $U_c^{\mathrm{prep}}, U_c^{\mathrm{read}} \in \mathrm{SU}(2)$ be arbitrary unitary operators on the control register $\mathcal{H}_c \cong \mathbb{C}^2$. Then the *asymmetric holosymmetric interferometric operator* is defined by:

$$\mathbf{U}_{\mathrm{holo}}^{(\mathrm{asym})} := \mathbf{U}_{\mathrm{read}} \cdot \mathbf{U}_{\mathrm{ctrl}} \cdot \mathbf{U}_{\mathrm{prep}},$$

where:

- **Preparation Gate:**

$$\mathbf{U}_{\mathrm{prep}} := U_c^{\mathrm{prep}} \otimes I_{\mathrm{RR}}, \quad \text{with} \quad U_c^{\mathrm{prep}} = \begin{pmatrix} \alpha & -\beta^* \\ \beta & \alpha^* \end{pmatrix}, \quad |\alpha|^2 + |\beta|^2 = 1.$$

- **Controlled Evolution:**

$$\mathbf{U}_{\mathrm{ctrl}} := |\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}.$$

- **Readout Gate:**

$$\mathbf{U}_{\mathrm{read}} := U_c^{\mathrm{read}} \otimes I_{\mathrm{RR}}, \quad \text{with} \quad U_c^{\mathrm{read}} = \begin{pmatrix} \gamma & -\delta^* \\ \delta & \gamma^* \end{pmatrix}, \quad |\gamma|^2 + |\delta|^2 = 1.$$

State Evolution:
Let the initial state be:

$$|\Psi_{\mathrm{init}}\rangle = |\mathbf{0}\rangle_c \otimes |\Psi_{\mathrm{RR}}\rangle.$$

Then:

1. **Preparation:**
$$\mathbf{U}_{\mathrm{prep}} |\Psi_{\mathrm{init}}\rangle = (\alpha |\mathbf{0}\rangle_c + \beta |\mathbf{1}\rangle_c) \otimes |\Psi_{\mathrm{RR}}\rangle.$$
2. **Controlled Evolution:**
$$\mathbf{U}_{\mathrm{ctrl}} \circ \mathbf{U}_{\mathrm{prep}} |\Psi_{\mathrm{init}}\rangle = \alpha |\mathbf{0}\rangle_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} |\Psi_{\mathrm{RR}}\rangle + \beta |\mathbf{1}\rangle_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} |\Psi_{\mathrm{RR}}\rangle.$$
3. **Readout:**
$$\begin{aligned} \mathbf{U}_{\mathrm{read}} \circ \mathbf{U}_{\mathrm{ctrl}} \circ \mathbf{U}_{\mathrm{prep}} |\Psi_{\mathrm{init}}\rangle = &\left[\gamma\alpha |\mathbf{0}\rangle_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} |\Psi_{\mathrm{RR}}\rangle - \delta^*\beta |\mathbf{0}\rangle_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} |\Psi_{\mathrm{RR}}\rangle\right] \\ &+ \left[\delta\alpha |\mathbf{1}\rangle_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} |\Psi_{\mathrm{RR}}\rangle + \gamma^*\beta |\mathbf{1}\rangle_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} |\Psi_{\mathrm{RR}}\rangle\right]. \end{aligned}$$

Interpretation. This construction generalizes the symmetric interferometer by allowing arbitrary superposition preparation and measurement bases. The interference structure no longer resolves simply into symmetric vs antisymmetric combinations, but rather into projective amplitudes defined by (α, β) and (γ, δ)—the control qubit's "in" and "out" basis states.

Hence, this operator:

- Encodes an asymmetrical duality of dialectical histories.
- Allows experimental probing of off-diagonal modal interference.
- Supports tunable observables via the preparation/readout gates U_c^{prep}, U_c^{read}.

This form is ideal for studying holosymmetric curvature, selective path filtering, or for embedding recursive gate dynamics in general holosymmetric protocols.

Asymmetric interferometer.

$$\mathbf{U}_{\text{holo}}^{(\text{asym})} = \mathbf{U}_{\text{read}}\, \mathbf{U}_{\text{ctrl}}\, \mathbf{U}_{\text{prep}}$$



Symmetric interferometer (the "fair splitter" version). Picture a fair, 50/50 beam splitter that sends a single *control light* down two routes at once. Along Route A, Register A performs the coordinated, entangling choreography while Register B follows a simple, independent routine; along Route B, those roles are swapped. After both routes run in parallel, a second fair splitter recombines them. If the two routes produced matching stories, their waves align and add (a bright outcome); if they differ, the waves cancel (a dark outcome). The brightness pattern on the control light is therefore a direct readout of how different those two dynamical stories are. "Symmetric" means both the initial split and the final recombination are perfectly balanced, making the device maximally sensitive to differences between the routes.

Asymmetric interferometer (the "tunable splitter" version). Here the fair splitters are replaced with adjustable ones. The first splitter can send more weight to one route than the other and can also dial in a relative phase; the final splitter can choose a different recombination angle and readout basis. These adjustments act like knobs that scan and amplify particular aspects of the interference between the two stories—sometimes emphasizing agreement, sometimes teasing out subtle mismatches that the symmetric version might gloss over. In short, it is the same two-route test, but with tunable "in" and "out" settings that bias the comparison, control sensitivity, and probe different facets of "who coordinates" and "who does not."

What both achieve. Both versions run two alternative, system-wide histories in parallel and then let those histories interfere. The symmetric model provides the clean, balanced baseline—maximum contrast with minimal tuning—while the asymmetric model is the lab-ready variant with extra control, able to favor a route, rotate the readout, and reveal finer structure in the differences between the two dynamical assignments.

From logic and algebra to quantum circuits. The compiled circuits are not only interesting—they are concrete, operational realizations of the *Fourfold Dialectic Algebra* of Chapter 1 on the $(0,1)$ modal sector. They implement, in hardware terms, the corner states $\mathsf{Th}, \mathsf{AntiTh}, \mathsf{SynTh}, \mathsf{HoloTh}$, the associated effects $(P_T, P_A, P_{\text{para}}, E_{\text{Holo}}^{(\mathbf{a}\to\mathbf{b})})$, and the dualization maps $(\mathbf{D}_{\mathcal{C}}, \Delta_{\mathcal{C}})$ on $\mathcal{A}_x^{(4)}$.

Corners $\mathsf{Th}/\mathsf{AntiTh}$ as a logical qubit. The ringed–ket basis $\{|\mathring{\mathbf{0}}\rangle, |\mathring{\mathbf{1}}\rangle\}$ from Definition 2.1 realizes the $\mathsf{Th}/\mathsf{AntiTh}$ corners of Chapter 1; the effects $P_T := |e_1\rangle\langle e_1|$ and $P_A := |e_2\rangle\langle e_2|$ are their Born tests on $\mathcal{A}_x^{(4)}$. The local involutive duality $D^{(\mathfrak{B}_x)}$ and its global lift $\hat{D}_{\text{PQS}}^{(n)}$ (Definitions 2.7 and 2.12) implement the categorical dualization $\mathbf{D}_{\mathcal{C}}$ and its configuration–wide action $\Delta_{\mathcal{C}}$ by swapping $\mathsf{Th} \leftrightarrow \mathsf{AntiTh}$ (Pauli–X on the logical qubit).

Synthesis SynTh as parasymmetry. The parasymmetric projector $P_{\text{para}}^{(x)} = |+\rangle\langle +|$ and its orthogonal complement $P_{\text{antipara}}^{(x)} = |-\rangle\langle -|$ (eq. (2.57)) resolve the $\mathsf{Th}/\mathsf{AntiTh}$ line into *synthesis* vs. *dissonance*. Their Naimark–flag circuits (local, pairwise, global) are the operational versions of the SynTh effect in Chapter 1, while the invariance criterion $UP_{\text{para}}^{(x_1\ldots x_n)}U^\dagger = P_{\text{para}}^{(x_1\ldots x_n)}$ (eq. (2.60)) characterizes the *parasymmetry–preserving* automorphisms of the $(0,1)$ corner-subalgebra of $\mathcal{A}_x^{(4)}$.

Holosymmetry HoloTh as interferometric composition. The controlled-path and Hadamard-conjugated operators $\mathbf{U}_{\text{ctrl}}$ and $\mathbf{U}_{\text{Had}}$ (Definitions 2.114 and 2.119) realize the holosymmetric corner HoloTh as *interference of dynamical assignments*: prepare, branch, recombine. In algebraic terms, the symmetric interferometer produces the sum/difference channels

$$\hat{\mathbf{U}}_{\pm} = \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} \pm \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G},$$

so that measuring the control implements the holosymmetric effects $E_{\text{Holo}}^{(\mathbf{a}\to\mathbf{b})}$ which compare relational histories $(\mathbf{a}, \mathbf{b})$ by constructive/destructive interference. The asymmetric interfer-

ometer generalizes this comparison to arbitrary SU(2) "in/out'' axes on the control, exposing off–diagonal holosymmetric structure.

Generators and flows in $\mathcal{A}_x^{(4)}$. The unitary families $U_N^{(\odot)}, U_G^{(\hat{\sigma})}, U_G^{(\lambda)}, U_G^{(\odot)}$ we built (decoupled, exchange–coupled, composite, unified) are exponentials of Hermitian combinations of the Chapter 1 effects, i.e. elements generated by $\hat{\mathcal{F}}_\alpha$ inside the automorphism group of the Fourfold algebra. Their commutation with P_{para} (when present) certifies closure on the SynTh sector, while duality gates realize $\mathbf{D}_\mathcal{C}$ as an involution on that sector.

Conclusion. Taken together, the circuits instantiate the Chapter 1 logic *as dynamics*: Th/AntiTh as a logical basis, SynTh as coherent superposition and its projector, HoloTh as interferometric comparison of whole relational evolutions, and $\mathbf{D}_\mathcal{C}, \Delta_\mathcal{C}$ as involutive symmetries at local and global scale. In short, the quantum circuits are the operational semantics of the tetralemmatic algebra $\mathcal{A}_x^{(4)}$.

Chapter 3

The Double-Slit Experiments

3.1 The Double-Slit Experiment for The Interferometer Model

3.1.1 Experiment for the Symmetric Interferometer Model

The fundamental quantum principle of interference, exemplified by the iconic double-slit experiment, is radically generalized within the holosymmetric framework. Here, the *slits* are not spatial apertures but entire, register-wide unitary evolutions, and the *interference pattern* is encoded not in particle distributions but in entangled amplitudes within a control qubit. This construction is made possible by the *Controlled Holosymmetric Operator* $\mathbf{U}_{\text{ctrl}}$ (Definition 2.114), which implements a superposition of two structurally distinct dynamical paths across dual quantum registers.

Mapping the Components of the Analogy We identify the components of this holosymmetric experiment with their canonical quantum analogues:

- **The Quantum System:** The dual-register quantum state $|\Psi_{\text{RR}}\rangle \in \mathcal{H}_{\text{RR}}^{(2n)}$, composed of two disjoint arrays of n Primordial Quantum Singularities each (eq. (2.61)). This system plays the role of the particle traversing the interferometer.
- **The Two Slits:** The two distinct, globally unitary, register-wide dynamical paths:
  - $\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} := U_G^{(A)} \cdot U_N^{(B)}$, where Register A undergoes globally coupled parasymmetric evolution and Register B evolves independently (eq. (2.69));
  - $\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} := U_N^{(A)} \cdot U_G^{(B)}$, where the roles of coupling and decoupling are reversed (eq. (2.70)).

  These operators constitute mutually exclusive but physically complete dynamical branches — the quantum analogues of the slits through which the system may evolve.
- **The Detector Screen:** The control system $\mathcal{H}_c = \text{span}\{|\mathbf{0}\rangle_c, |\mathbf{1}\rangle_c\}$, which registers the interference between the two holosymmetric paths. Measurement of the control qubit in a rotated basis reveals constructive or destructive interference, depending on the relative amplitudes and phases induced by the path operators.

The Three Experimental Acts of the Symmetric Interferometer The experimental procedure for the symmetric model can be elegantly framed as a three-act quantum-dynamical process.

Act I: Path Creation (Preparation) The experiment begins with the system in a definite initial state, with the control qubit aligned to a single basis vector, for instance $|\mathbf{0}\rangle_c$. The first act is to create a coherent superposition of evolutionary paths. This is achieved by applying the Hadamard preparation gate, $\mathbf{U}_{\text{prep}} = \mathbf{U}_{\text{Had}}$, which acts as the first "beam splitter."

- **Initial State:** $|\Psi_{\text{init}}\rangle = |\mathbf{0}\rangle_c \otimes |\Psi_{\text{RR}}\rangle$
- **Action:** Apply $\mathbf{U}_{\text{Had}}$
- **Prepared State:**

$$|\Psi_{\text{prep}}\rangle = \mathbf{U}_{\text{Had}} |\Psi_{\text{init}}\rangle = \left(\frac{1}{\sqrt{2}} (|\mathbf{0}\rangle_c + |\mathbf{1}\rangle_c) \right) \otimes |\Psi_{\text{RR}}\rangle$$

The system is now prepared to traverse both "slits"—the two distinct evolutionary pathways—simultaneously.

Act II: Differential Evolution (Branching) In the second act, the controlled unitary operator $\mathbf{U}_{\text{ctrl}}$ is applied. Each branch of the superposition now evolves according to its corresponding dynamical law, becoming entangled with the state of the registers.

- **State from Act I:** $|\Psi_{\text{prep}}\rangle$
- **Action:** Apply $\mathbf{U}_{\text{ctrl}}$
- **Entangled State:**

$$|\Psi_{\text{entangled}}\rangle = \mathbf{U}_{\text{ctrl}} |\Psi_{\text{prep}}\rangle = \frac{1}{\sqrt{2}} \left(|\mathbf{0}\rangle_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} |\Psi_{\text{RR}}\rangle + |\mathbf{1}\rangle_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} |\Psi_{\text{RR}}\rangle \right)$$

At the conclusion of this act, the system exists in a profound superposition, with the control qubit encoding which evolutionary history has been applied to which branch.

Act III: Recombination and Measurement The final act recombines the two paths to reveal their interference. The second Hadamard gate, $\mathbf{U}_{\text{read}} = \mathbf{U}_{\text{Had}}$, is applied, acting as the final "beam splitter" before the detector.

- **State from Act II:** $|\Psi_{\text{entangled}}\rangle$
- **Action:** Apply $\mathbf{U}_{\text{Had}}$
- **Final State:**

$$\begin{aligned} |\Psi_{\text{final}}\rangle &= (\mathbf{U}_{\text{Had}} \otimes \mathbb{I}) |\Psi_{\text{entangled}}\rangle \\ &= \frac{1}{2} \left(|\mathbf{0}\rangle_c \otimes \left(\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} \right) |\Psi_{\text{RR}}\rangle + |\mathbf{1}\rangle_c \otimes \left(\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} - \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} \right) |\Psi_{\text{RR}}\rangle \right) \end{aligned}$$

This final state is then measured. The probability of finding the control qubit in each basis state is determined by the constructive or destructive interference of the two paths:

$$\Pr[\mathbf{0}] = \frac{1}{4}\left\|\left(\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}\right)|\Psi_{\mathrm{RR}}\rangle\right\|^2,$$
$$\Pr[\mathbf{1}] = \frac{1}{4}\left\|\left(\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} - \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}\right)|\Psi_{\mathrm{RR}}\rangle\right\|^2.$$

This three-act structure provides a clear, rigorous, and physically intuitive framework for understanding the holosymmetric interferometer in the symmetric model.

Interpretation: Interference in the Symmetric Model This construction elevates the classical notion of path interference to a higher categorical level. In the symmetric model, the use of identical Hadamard gates for both preparation and recombination establishes a perfectly balanced apparatus. Consequently, the resulting interference pattern becomes a direct measure of the *functional symmetry* between the two evolutionary paths, $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$ and $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$.

What interferes here are not trajectories through spacetime, but entire unitary evolutions. The final probabilities, p_0 and p_1, are determined by the quantum overlap integral:

$$\mathcal{O} = \langle\Psi_{\mathrm{RR}}|\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G\dagger}\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}|\Psi_{\mathrm{RR}}\rangle$$

Specifically, the probabilities depend directly on the *real part* of this complex-valued amplitude, $\mathrm{Re}(\mathcal{O})$, which cleanly quantifies the degree of constructive or destructive interference between the two symmetrically prepared and recombined holosymmetric evolutions.

The Recursive Structure Crucially, this holosymmetric double-slit architecture is *recursive*. The states $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}|\Psi_{\mathrm{RR}}\rangle$ and $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}|\Psi_{\mathrm{RR}}\rangle$ are not static outputs; they themselves encode coupled parasymmetric dynamics across the n-PQS registers—dynamics that can be recursively decomposed into further layers of parasymmetric superpositions and entanglement. Thus, the holosymmetric interference pattern contains within it multiple strata of modal interference, reflecting a fractal-like coherence structure that generalizes Feynman's sum-over-paths into a sum-over-histories-of-evolutions.

Remark 3.1 (Symmetric Interferometer Visibility as a Dialectic Measurement)**.** In the specific case of the symmetric interferometer, the probability distribution on the control qubit acts as a particularly clear physical *visibility functional*. It is a quantum measurement device not for position or momentum, but for operator-theoretic duality.

Maximal constructive interference ($p_1 = 0$) occurs if and only if the two evolutionary paths produce an identical final state from the given initial state:

$$\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}|\Psi_{\mathrm{RR}}\rangle = \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}|\Psi_{\mathrm{RR}}\rangle$$

This signifies a perfect *functional symmetry* between the two dynamics with respect to the input state $|\Psi_{\mathrm{RR}}\rangle$.

Conversely, maximal destructive interference ($p_0 = 0$) occurs when the two paths yield perfectly anti-correlated final states:

$$\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}|\Psi_{\mathrm{RR}}\rangle = -\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}|\Psi_{\mathrm{RR}}\rangle$$

This condition reveals a state of perfect *modal complementarity*. These clean interpretations are a direct result of the symmetric preparation and readout stages.

Recursive Interference and the Structure of the Symmetric Interferometer

The Symmetric Operator as a Complete Interferometer To probe the recursive nature of quantum interference, we analyze the full structure of the *Symmetric Interferometer Holosymmetric Operator*, $\mathbf{U}_{\mathrm{holo}}^{(\mathrm{sym})}$. This operator's very structure encapsulates a complete interferometric circuit (*Prepare* $\rightarrow$ *Evolve* $\rightarrow$ *Recombine*) as a single, composite transformation on the state space. It is defined as:

$$\mathbf{U}_{\mathrm{holo}}^{(\mathrm{sym})} := \mathbf{U}_{\mathrm{Had}} \cdot \mathbf{U}_{\mathrm{ctrl}} \cdot \mathbf{U}_{\mathrm{Had}}.$$

To understand its properties, we derive its matrix representation in the control basis $\{|\mathbf{0}\rangle_c, |\mathbf{1}\rangle_c\}$. Using the block matrix forms for $\mathbf{U}_{\mathrm{ctrl}}$ and $\mathbf{U}_{\mathrm{Had}}$, we find:

$$\mathbf{U}_{\mathrm{holo}}^{(\mathrm{sym})} = \frac{1}{2}\begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}\begin{pmatrix} \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} & 0 \\ 0 & \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} \end{pmatrix}\begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} = \frac{1}{2}\begin{pmatrix} \hat{\mathbf{U}}_+ & \hat{\mathbf{U}}_- \\ \hat{\mathbf{U}}_- & \hat{\mathbf{U}}_+ \end{pmatrix},$$

where

$$\hat{\mathbf{U}}_\pm := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} \pm \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}.$$

Action on a Simple Initial State Let us now apply this operator to the simple initial state $|\Psi_{\mathrm{init}}\rangle = |\mathbf{0}\rangle_c \otimes |\Psi_{\mathrm{RR}}\rangle$. A direct calculation yields:

$$|\Psi_{\mathrm{final}}\rangle = \mathbf{U}_{\mathrm{holo}}^{(\mathrm{sym})}|\Psi_{\mathrm{init}}\rangle = \frac{1}{2}\left(|\mathbf{0}\rangle_c \otimes \hat{\mathbf{U}}_+|\Psi_{\mathrm{RR}}\rangle + |\mathbf{1}\rangle_c \otimes \hat{\mathbf{U}}_-|\Psi_{\mathrm{RR}}\rangle\right).$$

This leads to the measurement probabilities:

$$p_0 = \frac{1}{4}\left\|\hat{\mathbf{U}}_+|\Psi_{\mathrm{RR}}\rangle\right\|^2, \qquad p_1 = \frac{1}{4}\left\|\hat{\mathbf{U}}_-|\Psi_{\mathrm{RR}}\rangle\right\|^2.$$

This result is mathematically sound and reveals a crucial insight: applying the full interferometric circuit $\mathbf{U}_{\mathrm{holo}}^{(\mathrm{sym})}$ to a simple basis state is operationally equivalent to preparing a superposition and measuring the interference once.

The True Nature of Second-Order Coherence The "recursive" or "second-order" nature of the model is not found in the outcome of this simple experiment. It is encoded within the *off-diagonal elements* of the matrix representation of $\mathbf{U}_{\mathrm{holo}}^{(\mathrm{sym})}$ itself.

The term $\mathbf{F}_{01} = \frac{1}{2}\hat{\mathbf{U}}_-$ describes the amplitude for a system that *starts* in the $|\mathbf{1}\rangle_c$ branch to be *measured* in the $|\mathbf{0}\rangle_c$ branch after the full process. This "path-flipping" amplitude is a

direct measure of the interference between histories and reveals how the operator acts on general superpositions.

A true second-order experiment would involve probing this transformative structure directly—for instance, by preparing a more complex initial state, or by applying the operator $\mathbf{U}_{\text{holo}}^{(\text{sym})}$ sequentially. The conceptual vision of "interference of interference" is thus a property of the full transformation, liberated from its most basic application.

Conceptual Reinterpretation: Second-Order Interference The recursive architecture of holosymmetry is not about a single measurement, but about the structure of the transformations themselves.

- **First-order interference** is the phenomenon of interfering two paths, quantified by the probabilities p_0 and p_1 derived above.
- **Second-order coherence** is encoded in the full operator $\mathbf{U}_{\text{holo}}^{(\text{sym})}$. This operator acts as a map on the Hilbert space of histories—it transforms an *input* interference pattern into an *output* interference pattern.

The framework thus allows us to ask not just "Which path was taken?" but the more profound question:

"What is the quantum relationship between different interference-resolved evolutions?"

The answer lies in the algebraic structure of $\mathbf{U}_{\text{holo}}^{(\text{sym})}$. It is the engine that drives the evolution of modal logic itself.

Remark 3.2 (Second-Order Interference as Recursive Holosymmetric Dialectic)**.** This second-order construction completes the dialectical architecture of holosymmetry. The interference terms $\hat{\mathbf{U}}_{\pm}$ represent opposing modal perspectives on path coherence, while their recursive combination within the operator $\mathbf{U}_{\text{holo}}^{(\text{sym})}$ reflects a synthesis of interference modalities. The operator itself thus constitutes an operational probe of the internal logical structure governing the superposition of entire evolutionary frameworks.

3.1.2 Interference Expectation in the Symmetric Interferometer Model

Corollary 3.3 (Interference Expectation: Symmetric Interferometer)**.** *Let* $\mathbf{U}_{holo}^{(sym)}$ *denote the Symmetric Interferometer Holosymmetric Operator (Definition 2.123), acting on the holosymmetric Hilbert space* $\mathcal{H}_{\text{Holo}} := \mathcal{H}_c \otimes \mathcal{H}_{\text{RR}}^{(2n)}$*. Suppose the system is initialized in the product state:*

$$|\Psi_{\text{init}}\rangle := |\mathbf{0}\rangle_c \otimes |\Psi_{\text{RR}}\rangle,$$

where $|\Psi_{\text{RR}}\rangle \in \mathcal{H}_{\text{RR}}^{(2n)}$ *is a normalized state of the dual PQS registers, and* $|\mathbf{0}\rangle_c$ *is a basis state of the control system.*

Then the probability p_0 *of measuring the control system in the state* $|\mathbf{0}\rangle_c$ *after application of* $\mathbf{U}_{holo}^{(sym)}$ *is given by:*

$$\boxed{p_0 = \frac{1}{4}\left\|\left(\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}\right)|\Psi_{\text{RR}}\rangle\right\|^2 = \frac{1}{4}\left\|\hat{\mathbf{U}}_{+}|\Psi_{\text{RR}}\rangle\right\|^2,} \tag{3.1}$$

where the constructive path operator is defined as:

$$\hat{\mathbf{U}}_{+} := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}.$$

Interpretation. *The measurement outcome is governed by the constructive interference between the two distinct holosymmetric branches. The probability of measuring $|\mathbf{0}\rangle_c$ is directly proportional to the squared norm of the coherent sum of the state vectors produced by each evolutionary path. This is a first-order interference effect that provides a direct measure of the functional symmetry between the two dynamics as they act on the initial state.*

Proof of Corollary 3.3. Let the system be initialized in the product state:

$$|\Psi_{\mathrm{init}}\rangle = |\mathbf{0}\rangle_c \otimes |\Psi_{\mathrm{RR}}\rangle,$$

where $|\Psi_{\mathrm{RR}}\rangle \in \mathcal{H}_{\mathrm{RR}}^{(2n)}$ is normalized. The full evolution is governed by the Symmetric Interferometer Holosymmetric Operator:

$$\mathbf{U}_{\mathrm{holo}}^{(\mathrm{sym})} := \mathbf{U}_{\mathrm{Had}}\, \mathbf{U}_{\mathrm{ctrl}}\, \mathbf{U}_{\mathrm{Had}} \quad (\text{ Definition 2.123}).$$

We compute the final state by applying these operators sequentially.

Act I: Path Creation (First Hadamard Gate). The first Hadamard operator acts on the control qubit, creating a superposition of paths:

$$\begin{aligned} |\Psi_{\mathrm{prep}}\rangle &:= \mathbf{U}_{\mathrm{Had}} |\Psi_{\mathrm{init}}\rangle \\ &= \frac{1}{\sqrt{2}} \left(|\mathbf{0}\rangle_c + |\mathbf{1}\rangle_c \right) \otimes |\Psi_{\mathrm{RR}}\rangle. \end{aligned}$$

Act II: Differential Evolution (Controlled Operator). The controlled operator $\mathbf{U}_{\mathrm{ctrl}}$ applies the distinct evolutionary paths to each branch of the superposition, creating an entangled state:

$$\begin{aligned} |\Psi_{\mathrm{entangled}}\rangle &:= \mathbf{U}_{\mathrm{ctrl}} |\Psi_{\mathrm{prep}}\rangle \\ &= \frac{1}{\sqrt{2}} \left(|\mathbf{0}\rangle_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} |\Psi_{\mathrm{RR}}\rangle + |\mathbf{1}\rangle_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} |\Psi_{\mathrm{RR}}\rangle \right). \end{aligned}$$

Act III: Path Recombination (Second Hadamard Gate). The final Hadamard gate acts on the control system, recombining the paths to produce interference:

$$\begin{aligned} |\Psi_{\mathrm{final}}\rangle &:= \mathbf{U}_{\mathrm{Had}} |\Psi_{\mathrm{entangled}}\rangle \\ &= \frac{1}{2} \Big[|\mathbf{0}\rangle_c \otimes \left(\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} \right) |\Psi_{\mathrm{RR}}\rangle \\ &\quad + |\mathbf{1}\rangle_c \otimes \left(\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} - \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} \right) |\Psi_{\mathrm{RR}}\rangle \Big]. \end{aligned}$$

Using the definitions $\hat{\mathbf{U}}_+ := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$ and $\hat{\mathbf{U}}_- := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} - \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$, the final state is:

$$|\Psi_{\text{final}}\rangle = \frac{1}{2}\left(|\mathbf{0}\rangle_c \otimes \hat{\mathbf{U}}_+|\Psi_{\mathrm{RR}}\rangle + |\mathbf{1}\rangle_c \otimes \hat{\mathbf{U}}_-|\Psi_{\mathrm{RR}}\rangle\right).$$

To compute the probability of measuring the control register in state $|\mathbf{0}\rangle_c$, we take the squared norm of the corresponding amplitude in the final state:

$$p_0 = \|\langle \mathbf{0}|_c|\Psi_{\text{final}}\rangle\|^2 = \left\|\frac{1}{2}\hat{\mathbf{U}}_+|\Psi_{\mathrm{RR}}\rangle\right\|^2 = \frac{1}{4}\left\|\hat{\mathbf{U}}_+|\Psi_{\mathrm{RR}}\rangle\right\|^2.$$

This is precisely the expression given in Corollary Corollary 3.3. □

3.1.3 Complementary Interference Probability

Proof. We seek to derive the probability, p_1, of measuring the control qubit in the state $|\mathbf{1}\rangle_c$. This is the complementary outcome to the probability p_0 derived in the proof of Corollary Corollary 3.3.

We begin by recalling the correctly derived final state of the system after the action of the Symmetric Interferometer Holosymmetric Operator, $\mathbf{U}_{\text{holo}}^{(\text{sym})}$:

$$|\Psi_{\text{final}}\rangle = \frac{1}{2}\left(|\mathbf{0}\rangle_c \otimes \hat{\mathbf{U}}_+|\Psi_{\mathrm{RR}}\rangle + |\mathbf{1}\rangle_c \otimes \hat{\mathbf{U}}_-|\Psi_{\mathrm{RR}}\rangle\right).$$

To find the probability p_1, we project this final state onto the desired outcome $|\mathbf{1}\rangle_c$ and compute the squared norm of the resulting amplitude:

$$p_1 = \|\langle \mathbf{1}|_c|\Psi_{\text{final}}\rangle\|^2.$$

Performing the projection yields:

$$\langle \mathbf{1}|_c|\Psi_{\text{final}}\rangle = \frac{1}{2}\hat{\mathbf{U}}_-|\Psi_{\mathrm{RR}}\rangle.$$

Taking the squared norm gives the final expression for the probability:

$$p_1 = \left\|\frac{1}{2}\hat{\mathbf{U}}_-|\Psi_{\mathrm{RR}}\rangle\right\|^2 = \frac{1}{4}\left\|\hat{\mathbf{U}}_-|\Psi_{\mathrm{RR}}\rangle\right\|^2.$$

Consistency via Unitarity. As a necessary check, the total probability must be conserved. Let us verify that $p_0 + p_1 = 1$.

$$\begin{aligned} p_0 + p_1 &= \frac{1}{4}\left(\left\|\hat{\mathbf{U}}_+|\Psi_{\mathrm{RR}}\rangle\right\|^2 + \left\|\hat{\mathbf{U}}_-|\Psi_{\mathrm{RR}}\rangle\right\|^2\right) \\ &= \frac{1}{4}\left(\langle\Psi_{\mathrm{RR}}|\hat{\mathbf{U}}_+^\dagger\hat{\mathbf{U}}_+|\Psi_{\mathrm{RR}}\rangle + \langle\Psi_{\mathrm{RR}}|\hat{\mathbf{U}}_-^\dagger\hat{\mathbf{U}}_-|\Psi_{\mathrm{RR}}\rangle\right). \end{aligned}$$

Using the definitions $\hat{\mathbf{U}}_\pm = \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} \pm \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$, we expand the operators:

$$\begin{aligned} \hat{\mathbf{U}}_+^\dagger\hat{\mathbf{U}}_+ &= 2\mathbb{I} + (\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N})^\dagger\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} + (\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G})^\dagger\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} \\ \hat{\mathbf{U}}_-^\dagger\hat{\mathbf{U}}_- &= 2\mathbb{I} - (\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N})^\dagger\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} - (\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G})^\dagger\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} \end{aligned}$$

Summing their expectation values, the cross-terms cancel, leaving:

$$p_0 + p_1 = \frac{1}{4}\left(\langle\Psi_{\mathrm{RR}}|4\mathbb{I}|\Psi_{\mathrm{RR}}\rangle\right) = \langle\Psi_{\mathrm{RR}}|\Psi_{\mathrm{RR}}\rangle = 1,$$

since $|\Psi_{\mathrm{RR}}\rangle$ is normalized. This confirms the internal consistency of the derivation and the completeness of the interference process. □

3.1.4 Visibility of Holosymmetric Interference: Symmetric Interferometer Model

Theorem 3.4 (Visibility of Holosymmetric Interference)**.** *Consider the symmetric holosymmetric interferometric configuration described in Corollary Corollary 3.3. Let the initial state be:*

$$|\Psi_{\mathrm{init}}\rangle = |\mathbf{0}\rangle_c \otimes |\Psi_{\mathrm{RR}}\rangle,$$

where $|\Psi_{\mathrm{RR}}\rangle \in \mathcal{H}_{\mathrm{RR}}^{(2n)}$ is normalized. The system evolves under the Symmetric Interferometer Holosymmetric Operator $\mathbf{U}_{holo}^{(sym)}$, and the control qubit is measured in the computational basis with outcome probabilities p_0 and p_1.

The holosymmetric interference visibility *for this model is defined as the difference in outcome probabilities:*

$$\mathcal{V}_{\mathrm{sym}} := |p_0 - p_1|.$$

This visibility is given explicitly by the real part of the quantum overlap integral between the two evolved states:

$$\boxed{\mathcal{V}_{\mathrm{sym}} = \left|\mathrm{Re}\left(\langle\Psi_{\mathrm{RR}}|\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G\dagger}\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}|\Psi_{\mathrm{RR}}\rangle\right)\right|.} \tag{3.2}$$

Interpretation. *The visibility, $\mathcal{V}_{\mathrm{sym}}$, directly quantifies the degree of coherent interference between the two distinct evolutionary paths. It is a first-order measure of the relational similarity between the final states produced by the coupled ($A_G B_N$) and decoupled ($A_N B_G$) dynamics.*

A value of $\mathcal{V}_{\mathrm{sym}} = 1$ indicates maximal interference, where the two histories are perfectly aligned or anti-aligned in the Hilbert space. A value of $\mathcal{V}_{\mathrm{sym}} = 0$ indicates that the two final states are orthogonal, resulting in no interference pattern on the control qubit. The visibility thus serves as a direct, physical measure of the quantum coherence between two distinct universal histories.

Proof. The proof follows directly from the corrected probability expressions derived previously.

$$p_0 = \frac{1}{4}\left\|\hat{\mathbf{U}}_+|\Psi_{\mathrm{RR}}\rangle\right\|^2 = \frac{1}{4}\left(2 + 2\mathrm{Re}\langle\Psi_{\mathrm{RR}}|\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G\dagger}\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}|\Psi_{\mathrm{RR}}\rangle\right)$$
$$p_1 = \frac{1}{4}\left\|\hat{\mathbf{U}}_-|\Psi_{\mathrm{RR}}\rangle\right\|^2 = \frac{1}{4}\left(2 - 2\mathrm{Re}\langle\Psi_{\mathrm{RR}}|\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G\dagger}\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}|\Psi_{\mathrm{RR}}\rangle\right)$$

The visibility is therefore:

$$\begin{aligned}\mathcal{V}_{\mathrm{sym}} = |p_0 - p_1| &= \left|\frac{1}{2}\left(1 + \mathrm{Re}(\mathcal{O})\right) - \frac{1}{2}\left(1 - \mathrm{Re}(\mathcal{O})\right)\right| \\ &= |\mathrm{Re}(\mathcal{O})| \\ &= \left|\mathrm{Re}\left(\langle\Psi_{\mathrm{RR}}|\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G\dagger}\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}|\Psi_{\mathrm{RR}}\rangle\right)\right|,\end{aligned}$$

where $\mathcal{O}$ is the quantum overlap integral. This completes the proof. □

3.1.5 Experiment for the Asymmetric Interferometer Model

Mapping the Components of the Asymmetric Model In the Asymmetric Interferometer Model, the core components of the double-slit analogy are generalized to allow for more complex experimental control.

1. *The Quantum System:* This remains the dual-register quantum state $|\Psi_{\text{RR}}\rangle \in \mathcal{H}_{\text{RR}}^{(2n)}$, which traverses the interferometric process.

2. *The Two Slits:* The "slits" are the same two distinct, globally unitary, register-wide dynamical paths as in the symmetric case:

   - Path 1: $\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}$ (Coupled A / Decoupled B)
   - Path 2: $\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}$ (Decoupled A / Coupled B)

3. *The Source of Asymmetry (The Gates):* The asymmetry is introduced by the preparation and readout stages, which are no longer fixed Hadamard gates but can be any SU(2) rotation.

   - *Asymmetric "Beam Splitter" ($\mathbf{U}_{prep}$)*: The preparation gate can create any arbitrary superposition $\alpha|\mathbf{0}\rangle_c + \beta|\mathbf{1}\rangle_c$, effectively sending unequal or phase-shifted "amplitudes" through the two slits.
   - *Asymmetric "Analyzer" ($\mathbf{U}_{read}$)*: The readout gate projects the final interfered state onto an arbitrary basis on the control qubit's Bloch sphere before measurement. It acts as a tunable filter or polarizer for the interference pattern.

4. *The Detector Screen:* The control system $\mathcal{H}_c$ remains the detector, but the "fringes" (measurement probabilities) are now observed after being transformed by the readout gate $\mathbf{U}_{\text{read}}$, revealing different aspects of the quantum coherence between the two paths.

In essence, the Asymmetric Model uses the same two evolutionary "slits" but allows the experimenter to shine a "polarized" quantum state through them and observe the resulting interference pattern with a "polarized" detector.

The Three Acts of the Asymmetric Interferometer The Asymmetric Interferometer Model generalizes the symmetric case by allowing for arbitrary preparation and measurement bases, providing complete control over the interferometric process.

Act I: Asymmetric Preparation The experiment begins with the system in a simple basis state. The preparation gate $\mathbf{U}_{\text{prep}}$, an arbitrary $SU(2)$ rotation, then creates a potentially unequal or phase-shifted superposition.

- *Initial State:* $|\Psi_{\text{init}}\rangle = |\mathbf{0}\rangle_c \otimes |\Psi_{\text{RR}}\rangle$
- *Action:* Apply $\mathbf{U}_{\text{prep}} = U_c^{\text{prep}} \otimes I_{\text{RR}}$
- *Prepared State:*

$$|\Psi_{\text{prep}}\rangle = \mathbf{U}_{\text{prep}}|\Psi_{\text{init}}\rangle = (\alpha|\mathbf{0}\rangle_c + \beta|\mathbf{1}\rangle_c) \otimes |\Psi_{\text{RR}}\rangle$$

The system is now prepared to traverse both paths, but with potentially different initial amplitudes α and β.

Act II: Differential Evolution The controlled evolution proceeds exactly as in the symmetric case, applying the two distinct dynamical laws to the corresponding branches of the superposition.

- *State from Act I:* $|\Psi_{\text{prep}}\rangle$
- *Action:* Apply $\mathbf{U}_{\text{ctrl}}$
- *Entangled State:*

$$|\Psi_{\text{entangled}}\rangle = \mathbf{U}_{\text{ctrl}}|\Psi_{\text{prep}}\rangle = \alpha|\mathbf{0}\rangle_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}|\Psi_{\text{RR}}\rangle + \beta|\mathbf{1}\rangle_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}|\Psi_{\text{RR}}\rangle$$

Act III: Asymmetric Recombination and Measurement The final act recombines the two paths using the readout gate $\mathbf{U}_{\text{read}}$, which is also a general $SU(2)$ rotation. This gate projects the interfered state onto an arbitrary measurement basis before the final observation.

- *State from Act II:* $|\Psi_{\text{entangled}}\rangle$
- *Action:* Apply $\mathbf{U}_{\text{read}} = U_c^{\text{read}} \otimes I_{\text{RR}}$
- *Final State:*

$$\begin{aligned}|\Psi_{\text{final}}\rangle = (\mathbf{U}_{\text{read}} \otimes \mathbb{I})|\Psi_{\text{entangled}}\rangle =& |\mathbf{0}\rangle_c \otimes \left(\gamma\alpha\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} - \delta^*\beta\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}\right)|\Psi_{\text{RR}}\rangle \\ &+ |\mathbf{1}\rangle_c \otimes \left(\delta\alpha\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + \gamma^*\beta\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}\right)|\Psi_{\text{RR}}\rangle\end{aligned}$$

The measurement probabilities for observing $|\mathbf{0}\rangle_c$ or $|\mathbf{1}\rangle_c$ are the norm-squared of these final, complex amplitudes:

$$p_0 = \left\|\left(\gamma\alpha\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} - \delta^*\beta\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}\right)|\Psi_{\text{RR}}\rangle\right\|^2$$

$$p_1 = \left\|\left(\delta\alpha\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + \gamma^*\beta\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}\right)|\Psi_{\text{RR}}\rangle\right\|^2$$

Interpretation: Probing Interference from Any Angle This generalized interferometric construction allows for a much richer analysis. The final interference pattern is no longer a simple measure of symmetry between the two paths, but a complex function of all eight parameters ($\alpha, \beta, \gamma, \delta$ and the path operators). By carefully choosing the preparation and readout gates, one can "rotate" the measurement basis on the Bloch sphere, allowing for a full quantum state tomography of the interference between universal histories.

Interpretation of the Asymmetric Model: Quantum Tomography of Histories The true power of the Asymmetric Interferometer Model lies in its ability to perform a full *tomography* of the quantum-relational structure between the two evolutionary pathways. Unlike the symmetric case, which provides a single, fixed view of the interference, the asymmetric model allows the "observer" to probe the interference from any desired angle on the control qubit's Bloch sphere.

The resulting measurement probabilities, p_0 and p_1, are no longer simple functions of the overlap between the two paths. They are complex functions of the preparation parameters (α, β), the readout parameters (γ, δ), and the two unitary evolutions $(\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}, \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G})$.

For instance, the probability of measuring the $|\mathbf{0}\rangle_c$ outcome is:

$$p_0 = \left\| \left(\gamma\alpha \, \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} - \delta^* \beta \, \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} \right) |\Psi_{\mathrm{RR}}\rangle \right\|^2$$

By systematically varying the preparation and readout gates, one can, in principle, extract both the magnitude and the phase of the quantum overlap integral $\mathcal{O} = \langle \Psi_{\mathrm{RR}} | \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G \dagger} \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} | \Psi_{\mathrm{RR}} \rangle$. This allows for a complete characterization of the quantum relationship between the two universal histories.

Remark 3.5 (Asymmetric Interferometer as a Tomographic Tool)**.** The Asymmetric Interferometer functions as a device for performing *quantum process tomography* on the interference of universal histories. It does detect the presence of interference and it provides the experimental control needed to fully map out the complex amplitude (both magnitude and phase) that quantifies the relational coherence between two distinct dynamical laws. This elevates the model from a simple comparative device to a complete diagnostic tool for probing the deep structure of holosymmetric evolution.

Interference and Visibility: Asymmetric Interferometer Model

The Asymmetric Interference Process The Asymmetric Interferometer Model provides a powerful generalization of the symmetric case by allowing the experimenter to probe the interference between the two evolutionary paths, $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$ and $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$, from any "angle" on the control qubit's Bloch sphere.

The process unfolds as we previously derived: an arbitrary preparation $\mathbf{U}_{\mathrm{prep}}$ creates an initial superposition $(\alpha|\mathbf{0}\rangle_c + \beta|\mathbf{1}\rangle_c) \otimes |\Psi_{\mathrm{RR}}\rangle$; this state evolves via $\mathbf{U}_{\mathrm{ctrl}}$; and finally, an arbitrary readout $\mathbf{U}_{\mathrm{read}}$ is applied before measurement.

The final state of the system is:

$$\begin{aligned} |\Psi_{\mathrm{final}}\rangle = & |\mathbf{0}\rangle_c \otimes \left(\gamma\alpha \, \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} - \delta^* \beta \, \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} \right) |\Psi_{\mathrm{RR}}\rangle \\ & + |\mathbf{1}\rangle_c \otimes \left(\delta\alpha \, \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + \gamma^* \beta \, \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} \right) |\Psi_{\mathrm{RR}}\rangle \end{aligned}$$

Asymmetric Visibility and Quantum Tomography The measurement probabilities p_0 and p_1 are the squared norms of the amplitudes corresponding to the $|\mathbf{0}\rangle_c$ and $|\mathbf{1}\rangle_c$ branches. For instance:

$$p_0 = \left\| \left(\gamma\alpha \, \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} - \delta^* \beta \, \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} \right) |\Psi_{\mathrm{RR}}\rangle \right\|^2$$

Expanding this expression reveals its rich structure:

$$p_0 = |\gamma\alpha|^2 \|\hat{\mathbf{U}}_A \Psi\|^2 + |\delta\beta|^2 \|\hat{\mathbf{U}}_B \Psi\|^2 - 2\mathrm{Re}\left(\gamma\alpha\beta^* \delta \langle \Psi | \hat{\mathbf{U}}_B^\dagger \hat{\mathbf{U}}_A | \Psi \rangle \right)$$

(where $\hat{\mathbf{U}}_A$ is shorthand for $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$. This operator describes the evolution where *Register A* undergoes globally coupled dynamics ($U_G^{(A)}$) while *Register B* undergoes decoupled dynamics ($U_N^{(B)}$). And, $\hat{\mathbf{U}}_B$ is shorthand for $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$. This operator describes the complementary evolution where *Register A* is decoupled ($U_N^{(A)}$) and *Register B* is globally coupled ($U_G^{(B)}$)).

Unlike the symmetric case where visibility only depends on the real part of the overlap integral $\mathcal{O} = \langle \Psi | \hat{\mathbf{U}}_B^\dagger \hat{\mathbf{U}}_A | \Psi \rangle$, the asymmetric model's outcome depends on the full complex value of $\mathcal{O}$ and its interplay with the four complex parameters $\alpha, \beta, \gamma, \delta$.

Remark 3.6 (Asymmetric Interferometer as a Tomographic Tool)**.** The Asymmetric Interferometer functions as a device for performing *quantum process tomography* on the interference of universal histories. By systematically varying the preparation and readout gates (i.e., choosing different values for $\alpha, \beta, \gamma, \delta$), an experimenter can, in principle, map out the full complex value of the quantum overlap integral $\mathcal{O}$. This elevates the model from a simple comparative device to a complete diagnostic tool for probing the deep structure of holosymmetric evolution.

Interference Expectation in the Asymmetric Interferometer Model

Corollary 3.7 (Interference Expectation: Asymmetric Interferometer)**.** *Let* $\mathbf{U}_{holo}^{(asym)} := \mathbf{U}_{read} \cdot \mathbf{U}_{ctrl} \cdot \mathbf{U}_{prep}$ *be the Asymmetric Interferometer Holosymmetric Operator (Definition 2.124), acting on the holosymmetric Hilbert space* $\mathcal{H}_{\mathrm{Holo}}$*. Suppose the system is initialized in the product state:*

$$|\Psi_{\mathrm{init}}\rangle := |\mathbf{0}\rangle_c \otimes |\Psi_{\mathrm{RR}}\rangle,$$

where $|\Psi_{\mathrm{RR}}\rangle \in \mathcal{H}_{\mathrm{RR}}^{(2n)}$ *is a normalized state of the dual PQS registers, and* $|\mathbf{0}\rangle_c \in \mathcal{H}_c$ *is the initial control state.*

Then the probability $p_0^{(asym)}$ *of observing the control system in state* $|\mathbf{0}\rangle_c$ *after evolution under* $\mathbf{U}_{holo}^{(asym)}$ *is given by:*

$$\boxed{p_0^{(asym)} = \left\| \left(\gamma\alpha \, \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} - \delta^* \beta \, \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} \right) |\Psi_{\mathrm{RR}}\rangle \right\|^2,} \tag{3.3}$$

where the complex coefficients $\alpha, \beta, \gamma, \delta$ *are the matrix elements of the preparation gate* U_c^{prep} *and readout gate* U_c^{read}*, respectively.*

Interpretation. *In the asymmetric model, the measurement outcome is governed by a tunable, coherent superposition of the two evolutionary paths. The final probability depends not only on the intrinsic properties of the path operators but also explicitly on the choice of*

preparation and measurement bases for the control system. This allows the interferometer to function as a tomographic device, capable of mapping out the full complex quantum overlap between the two histories by systematically varying the gate parameters.

Proof of Corollary Corollary 3.7. The goal is to derive the probability, $p_0^{(\text{asym})}$, of measuring the control qubit in the state $|\mathbf{0}\rangle_c$ after the system evolves under the formally defined Asymmetric Interferometer Holosymmetric Operator.

We begin with the initial state:

$$|\Psi_{\text{init}}\rangle = |\mathbf{0}\rangle_c \otimes |\Psi_{\text{RR}}\rangle,$$

where $|\Psi_{\text{RR}}\rangle \in \mathcal{H}_{\text{RR}}^{(2n)}$ is a normalized state of the dual PQS registers.

The full evolution is governed by the operator:

$$\mathbf{U}_{\text{holo}}^{(\text{asym})} := \mathbf{U}_{\text{read}} \cdot \mathbf{U}_{\text{ctrl}} \cdot \mathbf{U}_{\text{prep}}.$$

The derivation proceeds in three acts.

Act I: Asymmetric Preparation. The arbitrary preparation gate $\mathbf{U}_{\text{prep}} = U_c^{\text{prep}} \otimes I_{\text{RR}}$ acts on the initial state, creating a general superposition:

$$\begin{aligned} |\Psi_{\text{prep}}\rangle &:= \mathbf{U}_{\text{prep}} |\Psi_{\text{init}}\rangle \\ &= (\alpha\,|\mathbf{0}\rangle_c + \beta\,|\mathbf{1}\rangle_c) \otimes |\Psi_{\text{RR}}\rangle. \end{aligned}$$

Act II: Differential Evolution. The controlled evolution operator $\mathbf{U}_{\text{ctrl}}$ then applies the two distinct dynamical paths to the corresponding branches of the superposition:

$$\begin{aligned} |\Psi_{\text{entangled}}\rangle &:= \mathbf{U}_{\text{ctrl}} |\Psi_{\text{prep}}\rangle \\ &= \alpha\,|\mathbf{0}\rangle_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} |\Psi_{\text{RR}}\rangle + \beta\,|\mathbf{1}\rangle_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} |\Psi_{\text{RR}}\rangle. \end{aligned}$$

Act III: Asymmetric Recombination. Finally, the arbitrary readout gate $\mathbf{U}_{\text{read}} = U_c^{\text{read}} \otimes I_{\text{RR}}$ acts on the control qubit, projecting the interfered state onto the final measurement basis:

$$\begin{aligned} |\Psi_{\text{final}}\rangle &:= \mathbf{U}_{\text{read}} |\Psi_{\text{entangled}}\rangle \\ &= |\mathbf{0}\rangle_c \otimes \left(\gamma\alpha\,\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} - \delta^*\beta\,\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}\right) |\Psi_{\text{RR}}\rangle \\ &\quad + |\mathbf{1}\rangle_c \otimes \left(\delta\alpha\,\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + \gamma^*\beta\,\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}\right) |\Psi_{\text{RR}}\rangle. \end{aligned}$$

To find the probability $p_0^{(\text{asym})}$, we project this final state onto $|\mathbf{0}\rangle_c$ and compute the squared norm of the amplitude:

$$p_0^{(\text{asym})} = \| \langle \mathbf{0}|_c |\Psi_{\text{final}}\rangle \|^2 = \left\| \left(\gamma\alpha\,\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} - \delta^*\beta\,\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}\right) |\Psi_{\text{RR}}\rangle \right\|^2.$$

This is precisely the expression given in Corollary Corollary 3.7. □

3.1.6 Asymmetric Complementary Interference Probability

Proof. We seek to derive the probability, $p_1^{(\text{asym})}$, of measuring the control qubit in the state $|\mathbf{1}\rangle_c$. This is the complementary outcome to the probability $p_0^{(\text{asym})}$ derived in the proof of Corollary Corollary 3.7.

We begin with the correctly derived final state of the system after the action of the Asymmetric Interferometer Holosymmetric Operator, $\mathbf{U}_{\text{holo}}^{(\text{asym})}$:

$$\begin{aligned}|\Psi_{\text{final}}\rangle =&|\mathbf{0}\rangle_c \otimes \left(\gamma\alpha\, \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} - \delta^*\beta\, \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}\right)|\Psi_{\text{RR}}\rangle \\ &+|\mathbf{1}\rangle_c \otimes \left(\delta\alpha\, \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + \gamma^*\beta\, \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}\right)|\Psi_{\text{RR}}\rangle\end{aligned}$$

To find the probability $p_1^{(\text{asym})}$, we project this final state onto the desired outcome $|\mathbf{1}\rangle_c$ and compute the squared norm of the resulting amplitude:

$$p_1^{(\text{asym})} = \left\|\langle\mathbf{1}|_c|\Psi_{\text{final}}\rangle\right\|^2.$$

Performing the projection isolates the amplitude for the '$|\mathbf{1}\rangle_c$' branch:

$$\langle\mathbf{1}|_c|\Psi_{\text{final}}\rangle = \left(\delta\alpha\, \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + \gamma^*\beta\, \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}\right)|\Psi_{\text{RR}}\rangle.$$

Taking the squared norm gives the final expression for the probability:

$$p_1^{(\text{asym})} = \left\|\left(\delta\alpha\, \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + \gamma^*\beta\, \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}\right)|\Psi_{\text{RR}}\rangle\right\|^2.$$

Consistency via Unitarity. As a necessary check, the total probability $p_0^{(\text{asym})} + p_1^{(\text{asym})}$ must sum to one. Let $\hat{U}_A = \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}$ and $\hat{U}_B = \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}$. Expanding the norms for p_0 and p_1:

$$\begin{aligned}p_0 &= \langle\Psi_{\text{RR}}|(\gamma^*\alpha^*\hat{U}_A^\dagger - \delta\beta^*\hat{U}_B^\dagger)(\gamma\alpha\hat{U}_A - \delta^*\beta\hat{U}_B)|\Psi_{\text{RR}}\rangle \\ &= |\gamma\alpha|^2 + |\delta\beta|^2 - 2\text{Re}(\gamma\alpha\beta\delta\langle\Psi_{\text{RR}}|\hat{U}_B^\dagger\hat{U}_A|\Psi_{\text{RR}}\rangle)\end{aligned}$$

$$\begin{aligned}p_1 &= \langle\Psi_{\text{RR}}|(\delta^*\alpha^*\hat{U}_A^\dagger + \gamma\beta^*\hat{U}_B^\dagger)(\delta\alpha\hat{U}_A + \gamma^*\beta\hat{U}_B)|\Psi_{\text{RR}}\rangle \\ &= |\delta\alpha|^2 + |\gamma\beta|^2 + 2\text{Re}(\delta\alpha\beta\gamma\langle\Psi_{\text{RR}}|\hat{U}_A^\dagger\hat{U}_B|\Psi_{\text{RR}}\rangle)\end{aligned}$$

Summing these probabilities, the cross-terms involving the quantum overlap integral cancel perfectly:

$$\begin{aligned}p_0 + p_1 &= |\alpha|^2(|\gamma|^2 + |\delta|^2) + |\beta|^2(|\delta|^2 + |\gamma|^2) \\ &= |\alpha|^2(1) + |\beta|^2(1) = 1,\end{aligned}$$

where we have used the unitarity conditions $|\alpha|^2 + |\beta|^2 = 1$ and $|\gamma|^2 + |\delta|^2 = 1$ for the preparation and readout gates. This confirms the internal consistency of our formulation. □

3.1.7 Visibility of Holosymmetric Interference: Asymmetric Interferometer Model

Theorem 3.8 (Visibility of Holosymmetric Interference: Asymmetric Interferometer)**.** *Consider the asymmetric holosymmetric interferometric configuration described in Corollary Corollary 3.7 and its complementary proof. Let the initial state be:*

$$|\Psi_{\text{init}}\rangle = |\mathbf{0}\rangle_c \otimes |\Psi_{\text{RR}}\rangle,$$

where $|\Psi_{\text{RR}}\rangle \in \mathcal{H}_{\text{RR}}^{(2n)}$ *is normalized. The system evolves under the Asymmetric Interferometer Holosymmetric Operator* $\mathbf{U}_{holo}^{(asym)}$*, and the control qubit is measured in the computational basis, yielding outcomes* $p_0^{(asym)}$ *and* $p_1^{(asym)}$*.*

The holosymmetric interference visibility *for the asymmetric interferometer model is defined as the absolute difference in outcome probabilities:*

$$\mathcal{V}_{\text{asym}} := \left| p_0^{(asym)} - p_1^{(asym)} \right|.$$

Substituting the correct expressions for the probabilities, the visibility is given explicitly by:

$$\boxed{\begin{aligned}\mathcal{V}_{\text{asym}} = \Bigg| & \left\| \left(\gamma\alpha\, \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} - \delta^* \beta\, \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} \right) |\Psi_{\text{RR}}\rangle \right\|^2 \\ & - \left\| \left(\delta\alpha\, \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + \gamma^* \beta\, \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} \right) |\Psi_{\text{RR}}\rangle \right\|^2 \Bigg|,\end{aligned}} \tag{3.4}$$

where the complex coefficients $\alpha, \beta, \gamma, \delta$ *are the matrix elements of the preparation gate* U_c^{prep} *and readout gate* U_c^{read}*, respectively.*

Interpretation. *In the asymmetric model, the visibility is a tunable functional that depends on the experimenter's choice of preparation and readout bases. Unlike the symmetric case, which yields a single visibility value, this model allows one to probe different aspects of the quantum overlap between the two evolutionary histories. By systematically varying the gate parameters, one can design the experiment to be sensitive to either the real or imaginary parts of the overlap integral* $\langle \Psi_{\text{RR}} | \hat{\mathbf{U}}_B^\dagger \hat{\mathbf{U}}_A | \Psi_{\text{RR}} \rangle$*, thereby performing a full tomographic reconstruction of the relational coherence between the two universal histories.*

Proof of Theorem Theorem 3.8. The proof proceeds directly from the definition of visibility and the corrected expressions for the measurement probabilities $p_0^{(\text{asym})}$ and $p_1^{(\text{asym})}$, which were derived from the action of the formal Asymmetric Interferometer Holosymmetric Operator, $\mathbf{U}_{\text{holo}}^{(\text{asym})} = \mathbf{U}_{\text{read}} \cdot \mathbf{U}_{\text{ctrl}} \cdot \mathbf{U}_{\text{prep}}$.

The probabilities for the two outcomes of the control qubit measurement are given by:

$$\begin{aligned} p_0^{(\text{asym})} &= \left\| \left(\gamma\alpha\, \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} - \delta^* \beta\, \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} \right) |\Psi_{\text{RR}}\rangle \right\|^2, \\ p_1^{(\text{asym})} &= \left\| \left(\delta\alpha\, \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + \gamma^* \beta\, \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} \right) |\Psi_{\text{RR}}\rangle \right\|^2. \end{aligned}$$

The visibility is defined as the absolute difference between these probabilities:

$$\mathcal{V}_{\text{asym}} := \left| p_0^{(\text{asym})} - p_1^{(\text{asym})} \right|.$$

Substituting the expressions for the probabilities directly into this definition yields:

$$\mathcal{V}_{\text{asym}} = \Big| \Big\| \Big(\gamma\alpha\, \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} - \delta^* \beta\, \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}\Big) |\Psi_{\text{RR}}\rangle \Big\|^2 - \Big\| \Big(\delta\alpha\, \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + \gamma^* \beta\, \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}\Big) |\Psi_{\text{RR}}\rangle \Big\|^2 \Big|.$$

This is precisely the visibility expression given in the corrected Theorem 3.8, thereby completing the proof. □

3.1.8 Interferometric Visibility Bounds

Setting. We work in the holosymmetric double-slit apparatus of Section 3.1, with control–qubit preparation and readout gates $U_c^{\text{prep}}, U_c^{\text{read}} \in \text{SU}(2)$ realizing the symmetric and asymmetric regimes (Sections 3.1.1 and 3.1.5). The total Hilbert space is $\mathcal{H}_{\text{Holo}} = \mathcal{H}_c \otimes \mathcal{H}_{\text{RR}}^{(2n)}$, where $\mathcal{H}_c \cong \mathbb{C}^2$ is the control system and $\mathcal{H}_{\text{RR}}^{(2n)}$ is the dual PQS–register space (RR = "dual PQS Registers"), with joint register states $|\Psi_{\text{RR}}\rangle \in \mathcal{H}_{\text{RR}}^{(2n)}$ as in Definition 2.113. Controlled holosymmetric evolution prepares two alternative path–histories on the registers, represented either by *pure* states $|\psi_a\rangle, |\psi_b\rangle \in \mathcal{H}_{\text{RR}}^{(2n)}$ or by *mixed* states $\rho_a, \rho_b \in \mathcal{D}(\mathcal{H}_{\text{RR}}^{(2n)})$ (see the path operators in Definition 2.108).

With single-qubit readout (Section 3.1.2), we include an adjustable reference phase $\phi \in \mathbb{R}$ on the control (equivalently, a choice of readout axis), and monitor the *fringe intensity*

$$I(\phi) \;=\; \tfrac{1}{2}\Big(1 + \text{Re}\big[e^{i\phi}\,\mathcal{M}\big]\Big), \qquad \mathcal{M} \;=\; \begin{cases} \langle\psi_a|\psi_b\rangle, & \text{pure histories,} \\ \text{Tr}\Big(\sqrt{\rho_a}\,\sqrt{\rho_b}\,U_{\text{rel}}\Big), & \text{mixed histories,} \end{cases}$$

where U_{rel} is the (experimentally controlled) *relative unitary* on the joint support of ρ_a, ρ_b induced by the chosen readout in the asymmetric setting (take $U_{\text{rel}} = \mathbb{I}$ in the symmetric case; cf. Section 3.1.5). The *visibility* is

$$\mathcal{V} \;:=\; \frac{I_{\max} - I_{\min}}{I_{\max} + I_{\min}} \in [0, 1],$$

so that in the symmetric regime one has $\mathcal{V} = |\mathcal{M}|$ (Section 3.1.2). This calculus follows the Pancharatnam–Berry–Anandan two-path phase/coherence lineage [23, 24, 25] and the Born-valued semantics adopted in Section 1.2.9.

$\mathcal{M}$ (overlap amplitude). Complex interference overlap of the two prepared path–histories; $\mathcal{M} = \langle\psi_a|\psi_b\rangle$ for pure histories and $\mathcal{M} = \text{Tr}(\sqrt{\rho_a}\sqrt{\rho_b}\,U_{\text{rel}})$ for mixed histories (in the symmetric readout, $\mathcal{V} = |\mathcal{M}|$); cf. Sections 3.1.2 and 3.1.5. **$\mathcal{D}(\mathcal{H})$.** Set of density operators on Hilbert space $\mathcal{H}$ (positive, trace one). **U_{rel}.** Relative unitary induced by the readout choice; identity in the symmetric case.

1) Pure Histories: Baseline and Triadic Bounds. For the symmetric Mach–Zehnder regime (Section 3.1.1), where the overlap amplitude is $\mathcal{M} = \langle\psi_a|\psi_b\rangle$,

$$\boxed{\mathcal{V} \;=\; |\mathcal{M}| \;=\; |\langle\psi_a|\psi_b\rangle| \;\leq\; 1} \,. \tag{3.5}$$

For general (possibly unbalanced) $SU(2)$ settings (Sections 3.1.5 and 3.1.7),

$$\boxed{\mathcal{V} \;\le\; |\langle\psi_a|\psi_b\rangle|}\,, \tag{3.6}$$

with equality when the beam–splitter weights equalise the path priors (standard two–path algebra in the Pancharatnam–Berry–Anandan framework [23, 24, 25]).

Triadic resolution. Let $P_{\mathrm{corr}}, P_{\mathrm{vac}}, P_{\mathrm{spec}}$ be the orthogonal projectors induced by the Unified Inter–Modal Correlator (Definition 6.4), so that $\sum_\chi P_\chi = \mathbb{I}$ and $P_\chi P_{\chi'} = \delta_{\chi\chi'} P_\chi$, with $\chi \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$. Define triadic overlap components

$$\mathcal{M}_\chi \;:=\; \langle\psi_a|\, P_\chi \,|\psi_b\rangle, \qquad \chi \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}, \tag{3.7}$$

so that $\langle\psi_a|\psi_b\rangle = \sum_\chi \mathcal{M}_\chi$. Two instrument–level envelopes follow:

(i) *Universal bound (no extra control):*

$$\boxed{\mathcal{V} \;\le\; \Big|\textstyle\sum_\chi \mathcal{M}_\chi\Big| \;\le\; \sum_\chi |\mathcal{M}_\chi|}\,. \tag{3.8}$$

(ii) *Triadically weighted bound (with channel weights).* Implement a readout that applies complex weights w_χ with $\sum_\chi |w_\chi|^2 = 1$ (operationally: pre/post–multiply by $\sum_\chi w_\chi P_\chi$; cf. Section 12.2). The fringe amplitude then becomes

$$\mathcal{M}(w) \;=\; \sum_\chi w_\chi\, \mathcal{M}_\chi,$$

and by Cauchy–Schwarz [356]

$$|\mathcal{M}(w)| \;=\; \Big|\textstyle\sum_\chi w_\chi \mathcal{M}_\chi\Big| \;\le\; \Big(\sum_\chi |\mathcal{M}_\chi|^2\Big)^{1/2}. \tag{3.9}$$

Choosing $w_\chi \propto \overline{\mathcal{M}_\chi}$ yields the optimal envelope

$$\boxed{\mathcal{V}_{\max} \;\le\; \Big(|\mathcal{M}_{\mathrm{corr}}|^2 + |\mathcal{M}_{\mathrm{vac}}|^2 + |\mathcal{M}_{\mathrm{spec}}|^2\Big)^{1/2}}\,. \tag{3.10}$$

Measurement note. Each $\mathcal{M}_\chi$ is the expectation of the effect $E_\chi := P_\chi|\psi_b\rangle\langle\psi_a|$ in the experiment's GNS state (Section 1.4.2); operationally, $|\mathcal{M}_\chi|$ is obtained by inserting P_χ in one arm and scanning the phase [355].

2) Mixed Histories: Fidelity Bounds and Triadic Additivity. Let $\rho_a, \rho_b \in \mathcal{D}(\mathcal{H}_{\mathrm{RR}}^{(2n)})$ be (possibly mixed) path states. In the mixed setting the overlap amplitude reads

$$\mathcal{M}(U_{\mathrm{rel}}) \;=\; \mathrm{Tr}\Big(\sqrt{\rho_a}\,\sqrt{\rho_b}\,U_{\mathrm{rel}}\Big), \qquad U_{\mathrm{rel}} \in \mathrm{U}(\mathrm{supp}\,\rho_a \cup \mathrm{supp}\,\rho_b),$$

with $U_{\mathrm{rel}} = \mathbb{I}$ in the symmetric readout (Section 3.1.1) and general SU(2) control in the asymmetric case (Section 3.1.5). By the polar decomposition $\sqrt{\rho_a}\sqrt{\rho_b} = V\,\big|\sqrt{\rho_a}\sqrt{\rho_b}\big|$, we have

$$|\mathcal{M}(U_{\mathrm{rel}})| \;\le\; \mathrm{Tr}\Big|\sqrt{\rho_a}\sqrt{\rho_b}\Big|,$$

with equality for $U_{\rm rel} = V^\dagger$. Using the Uhlmann fidelity [357, 287, 358]

$$F(\rho_a, \rho_b) \;:=\; \Big(\operatorname{Tr} \sqrt{\sqrt{\rho_a}\, \rho_b \, \sqrt{\rho_a}} \Big)^2 ,$$

and the identity $\operatorname{Tr}\big|\sqrt{\rho_a}\sqrt{\rho_b}\big| = \operatorname{Tr} \sqrt{\sqrt{\rho_a}\rho_b\sqrt{\rho_a}} = \sqrt{F(\rho_a,\rho_b)}$ [359, 357], it follows that

$$\boxed{\; \mathcal{V} \;=\; |\mathcal{M}(U_{\rm rel})| \;\le\; \sqrt{F(\rho_a,\rho_b)} \;\le\; 1 \;}\,. \tag{3.11}$$

(In the symmetric readout $U_{\rm rel} = \mathbb{I}$, one has $|\mathcal{M}(\mathbb{I})| \le \sqrt{F(\rho_a,\rho_b)}$; the upper bound is saturated when $\sqrt{\rho_a}\sqrt{\rho_b}$ is positive or when $U_{\rm rel}$ is chosen optimally via the polar part $V^\dagger$ [287, 358].)

Triadic resolution via pinching. Let $P_{\rm corr}, P_{\rm vac}, P_{\rm spec}$ be the orthogonal projectors from the Unified Inter–Modal Correlator (Definition 6.4), and define the CPTP "triadic pinching" channel

$$\mathcal{P}(\cdot) \;=\; \sum_{\chi \in \{\rm corr,vac,spec\}} P_\chi(\cdot)P_\chi, \qquad \text{(cf. Section 12.2)}.$$

What this means. The map $\mathcal{P}$ is a completely positive trace-preserving (CPTP) quantum channel with Kraus operators $K_\chi = P_\chi$:

$$\mathcal{P}(X) = \sum_\chi K_\chi X K_\chi^\dagger, \quad K_\chi = P_\chi, \quad \sum_\chi K_\chi^\dagger K_\chi = \sum_\chi P_\chi = \mathbb{I},$$

hence CP and TP (Choi–Jamiołkowski/Kraus criteria [360, 361, 287]). Operationally, $\mathcal{P}$ is a *non-selective projective measurement (or complete dephasing)* in the triadic sector label χ: it *removes inter-sector coherences* while *preserving* sector populations and total trace. Algebraically, $\mathcal{P}$ is the conditional expectation onto the block-diagonal subalgebra $\bigoplus_\chi P_\chi \mathcal{B}(\mathcal{H}) P_\chi$. After pinching, the states become block-diagonal,

$$\mathcal{P}(\rho_a) = \bigoplus_\chi \rho_a^\chi, \quad \rho_a^\chi := P_\chi \rho_a P_\chi, \qquad \mathcal{P}(\rho_b) = \bigoplus_\chi \rho_b^\chi,$$

and monotonicity of fidelity under CPTP maps yields [287, 358]

$$F(\rho_a,\rho_b) \;\le\; F\Big(\mathcal{P}(\rho_a), \mathcal{P}(\rho_b)\Big). \tag{3.12}$$

Because fidelity factorizes across orthogonal blocks,

$$\sqrt{F\Big(\mathcal{P}(\rho_a), \mathcal{P}(\rho_b)\Big)} \;=\; \sum_{\chi \in \{\rm corr,vac,spec\}} \sqrt{F\Big(\rho_a^\chi, \rho_b^\chi\Big)}. \tag{3.13}$$

Therefore the mixed–state visibility obeys the triadic envelope

$$\boxed{\; \mathcal{V} \;\le\; \sqrt{F(\rho_a,\rho_b)} \;\le\; \sum_{\chi \in \{\rm corr,vac,spec\}} \sqrt{F\Big(\rho_a^\chi, \rho_b^\chi\Big)} \;}\,. \tag{3.14}$$

Notes. (i) $\mathcal{P}$ can be implemented physically by inserting a non-selective sector measurement (or engineered dephasing) before readout; it leaves marginal sector weights unchanged while erasing cross-sector interference. (ii) The block factorization follows from $\sqrt{A \oplus B} = \sqrt{A} \oplus \sqrt{B}$ and $\operatorname{Tr}(X \oplus Y) = \operatorname{Tr} X + \operatorname{Tr} Y$.

3) Corollaries and Experimental Use. Write the triadic components of the overlap amplitude as

$$\mathcal{M}_\chi := \begin{cases} \langle\psi_a|P_\chi|\psi_b\rangle, & \text{pure histories,} \\ \mathrm{Tr}\big(\sqrt{\rho_a}\,P_\chi\,\sqrt{\rho_b}\,U_{\mathrm{rel}}\big), & \text{mixed histories,} \end{cases} \qquad \chi \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\},$$

with $\mathcal{M} = \sum_\chi \mathcal{M}_\chi$, where P_χ are the sector projectors from Definition 6.4. A *triadic filter* means either a coherent weighting $W = \sum_\chi w_\chi P_\chi$ with $\sum_\chi |w_\chi|^2 = 1$, or the CPTP pinching $\mathcal{P}(\cdot) = \sum_\chi P_\chi(\cdot)P_\chi$ (cf. Sections 12.2 and 15.1.4).

Corollary 3.9 (Consistency strip)**.** *For any triadic filter,*

$$\mathcal{V} \;\le\; \min\Big\{\, |\mathcal{M}|,\; \textstyle\sum_\chi |\mathcal{M}_\chi|,\; \Big(\sum_\chi |\mathcal{M}_\chi|^2\Big)^{1/2},\; \sqrt{F(\rho_a,\rho_b)}\,\Big\}.$$

A *triadic filter* (sector-resolving operation) means either a coherent weighting $W = \sum_\chi w_\chi P_\chi$ with $\sum_\chi |w_\chi|^2 = 1$, or the CPTP pinching $\mathcal{P}(\cdot) = \sum_\chi P_\chi(\cdot)P_\chi$.

Proof. The baseline bound $\mathcal{V} \le |\mathcal{M}|$ is from the symmetric two-path calculus in Section 3.1.8. Sector resolution yields $|\mathcal{M}| \le \sum_\chi |\mathcal{M}_\chi|$ and, with coherent weights, $\max_{\|w\|_2=1} |\sum_\chi w_\chi \mathcal{M}_\chi| = (\sum_\chi |\mathcal{M}_\chi|^2)^{1/2}$ by Cauchy–Schwarz [356]. For mixed histories, $\mathcal{V} \le \sqrt{F(\rho_a,\rho_b)}$ follows from the Uhlmann fidelity bound in Paragraph 2 (Section 3.1.8; [357, 287, 358]). For pure histories $F(\rho_a,\rho_b) = |\mathcal{M}|^2$, so the strip specializes consistently. □

Corollary 3.10 (Channel dominance test)**.** *If* $|\mathcal{M}_{\mathrm{vac}}|, |\mathcal{M}_{\mathrm{spec}}| \ll |\mathcal{M}_{\mathrm{corr}}|$*, then*

$$\mathcal{V}_{\max} \;\le\; \Big(|\mathcal{M}_{\mathrm{corr}}|^2 + |\mathcal{M}_{\mathrm{vac}}|^2 + |\mathcal{M}_{\mathrm{spec}}|^2\Big)^{1/2} \;\approx\; |\mathcal{M}_{\mathrm{corr}}|.$$

Proof. Use the ℓ^2 envelope from Section 3.1.8 and neglect subleading terms when the non-correlation sectors are suppressed. The approximation sign reflects the small-parameter regime. □

Corollary 3.11 (Selection–principle tie–in)**.** *In Stage–II comparative tomography (Section 12.2) under a fixed triadic filter, the configuration with the brightest interference fringe maximizes the* ℓ^2 *envelope* $\big(\sum_\chi |\mathcal{M}_\chi|^2\big)^{1/2}$*, rather than* $|\sum_\chi \mathcal{M}_\chi|$.

Proof. With coherent weighting $W = \sum_\chi w_\chi P_\chi$ ($\sum_\chi |w_\chi|^2 = 1$), the achievable amplitude is $\mathcal{M}(w) = \sum_\chi w_\chi \mathcal{M}_\chi$, whose maximum modulus is the ℓ^2 norm by Cauchy–Schwarz [356]. This objective is immune to accidental inter-sector phase cancellations that can depress $|\sum_\chi \mathcal{M}_\chi|$. □

4) Implementation Narrative. *How to read and use the bounds in practice.* Begin with the actual interferometer we can build: fix the SU(2) preparation/readout on the control (Sections 3.1.1 and 3.1.5) and extract the measured visibility $\mathcal{V}$ from the fringe $I(\phi)$ (Section 3.1.8). If the prepared histories are effectively pure, the baseline inequality from Paragraph 1 already constrains the signal by the overlap amplitude $|\mathcal{M}|$; if they are mixed (or suspected to be), the fidelity bound caps $\mathcal{V}$ by $\sqrt{F(\rho_a,\rho_b)}$ (see Section 3.1.8). To diagnose *where* coherence

lives, do not chase a single complex sum; instead, resolve the amplitude into triadic sectors with the projectors P_χ from Definition 6.4 and consult the sector envelopes: the ℓ^1 bound detects diffuse, phase-sensitive contributions, while the ℓ^2 (Cauchy–Schwarz) envelope records intrinsic sectoral strength and is robust against inter-sector phase cancellations (Corollary 3.9; [356]). For mixed histories, the same sector resolution is implemented cleanly by the CPTP triadic pinching $\mathcal{P}(\cdot) = \sum_\chi P_\chi(\cdot)P_\chi$, whose fidelity monotonicity and block factorization yield a transparent, additively interpretable ceiling on $\mathcal{V}$ (Sections 3.1.8 and 15.1.4; [287, 358]). In practice, a small vac/spec contribution relative to corr signals channel dominance (Corollary 3.10); and for control design or model selection, the "winner" under fixed filters is the configuration maximizing the ℓ^2 envelope rather than the raw complex sum (Corollary 3.11).

These inequalities are *instrument–level*: they require only the Born semantics of Section 1.2.9 and the existence of the triadic projectors from Definition 6.4. They connect directly to laboratory interferometry [23, 24, 25] and thus furnish clean pass/fail criteria for any proposal claiming large holosymmetric effects.

3.1.9 The Interferometer of Universal Histories

The double-slit experiment, that cornerstone demonstration of quantum mechanics, reveals how a single particle can traverse multiple paths simultaneously, creating interference patterns that encode the fundamental wave-particle duality of nature. In IQR framework, this principle undergoes a profound generalization: the "particle" becomes an entire dual-universe system, the "slits" become complementary evolutionary histories, and the "interference pattern" emerges as the observable structure of reality itself.

This conceptual mapping transforms our understanding of quantum interference from a microscopic phenomenon to a cosmological principle. Through the Holosymmetric Interferometer Model (Section 2.4), we discover that the universe may be continuously performing a meta-double-slit experiment upon itself, with physical laws emerging as stable interference patterns between competing evolutionary possibilities.

The Fundamental Analogy: Mapping Quantum Concepts Across Scales The correspondence between the classic double-slit experiment and the Holosymmetric Interferometer Model reveals a deep structural similarity:

Classic Double-Slit	**Holosymmetric Interferometer Model**
The Particle (e.g., electron)	The Dual-Register PQS State $\|\Psi_{\text{RR}}\rangle \in \mathcal{H}_{\text{RR}}^{(2n)}$
The Two Slits (spatial apertures)	The Two Evolutionary Paths: $\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}$ and $\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}$
The Detector Screen	The Control Qubit $\mathcal{H}_c = \text{span}\{\|\mathbf{0}\rangle_c, \|\mathbf{1}\rangle_c\}$
Interference Fringes	Measurement Probabilities p_0 and p_1
Wave Function	The Holosymmetric State Evolution

Table 3.1: The foundational analogy between the classic double-slit experiment and the Holosymmetric Interferometer Model, mapping quantum concepts across scales from particle mechanics to universal evolution.

This analogical mapping also represents a fundamental principle of the IQR wherein quantum interference operates at the level of entire cosmological configurations.

The Symmetric Interferometer: A Measure of Relational Symmetry

Theoretical Foundation: The Three Acts of the Interferometer In the symmetric model, we implement a perfect complementarity between two evolutionary pathways. The experimental protocol unfolds through three critical stages:

1. *Act I: Path Creation.* Starting with the system in a simple basis state, $|\Psi_{\text{initial}}\rangle = |\mathbf{0}\rangle_c \otimes |\Psi_{\text{RR}}\rangle$, the first Hadamard gate $\mathbf{U}_{\text{Had}}$ creates a coherent superposition of the two control states. This prepares the system to traverse both evolutionary paths simultaneously.

2. *Act II: Differential Evolution.* The controlled-unitary operator $\mathbf{U}_{\text{ctrl}}$ applies a different dynamical law to each branch of the superposition. The $|\mathbf{0}\rangle_c$ component evolves via "Slit 1" ($\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}$), while the $|\mathbf{1}\rangle_c$ component evolves via "Slit 2" ($\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}$).

3. *Act III: Recombination & Measurement.* A final Hadamard gate recombines the paths, causing them to interfere. A measurement on the control qubit reveals the outcome, with the probabilities given by:

$$p_0 = \frac{1}{4} \left\| \left(\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} \right) \|\Psi_{\text{RR}}\rangle \right\|^2$$

$$p_1 = \frac{1}{4} \left\| \left(\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} - \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} \right) \|\Psi_{\text{RR}}\rangle \right\|^2$$

Physical Interpretation The measurement outcomes encode a profound physical reality:

- p_0 *(Bright Fringe)*: The probability of constructive interference between evolutionary histories.

- p_1 *(Dark Fringe)*: The probability of destructive interference between evolutionary histories.

The visibility $\mathcal{V}_{\text{sym}} = |p_0 - p_1| = \left|\text{Re}\left(\langle\Psi_{\text{RR}}|\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G\dagger}\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}|\Psi_{\text{RR}}\rangle\right)\right|$ quantifies the degree of functional symmetry between the two dynamics as they act on the initial state.

The Asymmetric Interferometer: Quantum Tomography of Histories The asymmetric model generalizes the symmetric case by replacing the fixed Hadamard gates with arbitrary preparation and readout gates, $\mathbf{U}_{\text{prep}}$ and $\mathbf{U}_{\text{read}}$. The two evolutionary paths, or "slits," remain the same.

This configuration probes the interference between the two dynamic evolutions from any "angle" on the control qubit's Bloch sphere, as chosen by the experimenter.

The Tomographic Interpretation The asymmetric interferometer functions as a tool for *quantum process tomography*. The resulting measurement probabilities are complex functions of the gate parameters $(\alpha, \beta, \gamma, \delta)$ and the quantum overlap integral $\mathcal{O}$. By systematically varying the preparation and readout gates, one can map out the full complex value of $\mathcal{O}$, thereby performing a complete tomographic reconstruction of the quantum relationship between the two universal histories.

The Recursive Structure: Fractal Interference Patterns The profound innovation of the holosymmetric framework lies in its recursive structure. The statement "the architecture is recursive" refers to the fact that the evolutionary operators themselves, $\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}$ and $\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}$, are not primitive. They are built from nested, composite parasymmetric generators. Therefore, the states that interfere in our experiment are already deeply structured and encode their own layered, relational histories.

This is distinct from a "second-order experiment", which would involve applying the entire interferometer operator $\mathbf{U}_{\text{holo}}^{(\text{sym})}$ sequentially. The recursive nature is inherent in the building blocks of the theory.

The Self-Similar Structure of Reality This recursive structure suggests a fractal organization of quantum reality:

1. *Level 1: Interference of Quantum States.* This is the familiar interference of standard quantum mechanics. A particle's state is a superposition of passing through "slit A" and "slit B", and these possibilities interfere. This is the base pattern.

2. *Level 2: Interference of Evolutionary Histories.* Our Holosymmetric Interferometer elevates this principle. The "slits" are no longer spatial paths, but entire evolutionary histories ($\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}$ and $\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}$). We create a superposition of these two distinct dynamical laws and measure their interference. The *same principle of interference* is now acting on the laws of evolution themselves.

3. *Level 3: Recursive Structure of the Histories.* Here lies the deepest level of self-similarity. We find that the operators defining the histories ($\hat{\mathbf{U}}_{\text{RR}}$) are not primitive. They are

themselves recursively constructed from more fundamental local, pairwise, and global parasymmetric generators. The very building blocks of the Level 2 interference are themselves built from a nested, compositional logic.

The "fractal" nature of this reality lies in the repeating motif of *coherent superposition and interference*. This single quantum principle is applied to: (1) states, (2) processes that act on states, and (3)the very definition of those processes. The framework suggests a universe with a profound logical self-similarity, where the same quantum principle that governs the path of an electron also governs the superposition of cosmic histories and the structure of the laws that write them.

Fractal Reality and the Quantum Meta-Law The most profound prediction of this framework is that reality itself should bear the imprint of fractal organization at all scales—a testable hypothesis with specific observable consequences. The *cosmic web*, that vast filamentary structure of galaxies and dark matter spanning hundreds of millions of light-years, may be the most direct visualization of this principle. Rather than arising from stochastic gravitational clustering, the framework suggests these structures are *macroscopic manifestations of the underlying quantum entanglement network*—the filaments representing high-correlation pathways in the $\hat{\Omega}$-Continuum, with galaxies condensing at the most interconnected nodes [362, 147]. If true, detailed analysis of the cosmic web should reveal not just statistical self-similarity, but precise mathematical patterns that mirror the quantum interference structures at microscopic scales [31, 363].

This fractal principle extends to the deepest mysteries of particle physics, offering a stunning reinterpretation of the *three generations of matter*. Rather than arbitrary repetition, the electron-muon-tau hierarchy (and corresponding quark families) may represent *three self-similar manifestations of a single underlying principle at different energy scales*—just as a fractal contains smaller copies of itself [364]. This transforms the generation problem from "why three?" to "what is the scaling rule?"—a quantitative question amenable to experimental investigation. Similarly, the phenomenon of *running coupling constants* in quantum field theory, where fundamental "constants" vary with measurement energy, becomes not a technical complication but a deep geometric principle: the laws themselves are scale-dependent reflections of the fractal substrate from which they emerge [32].

The experimental program for testing this framework thus focuses on identifying *universal scaling laws* and *hierarchical interference patterns* across all domains of physics. We should find the same mathematical structures governing quantum decoherence times, galactic formation rates, and cosmic evolution epochs—properly normalized for scale. The CMB might contain nested interference patterns following the holosymmetric principle [365], while phase transitions from particle physics to cosmology should exhibit recursive symmetry breaking with identical mathematical forms [366]. This vision transforms physics from finding different laws at different scales to discovering how the *same meta-law* manifests universally—reality written in a self-similar quantum language where being and becoming unite through the eternal dance of quantum interference across all possible histories of the cosmos.

3.2 Interferometric Model Recursive Expansion

3.2.1 Recursive Component Expansion: Symmetric Interferometer

Let $\mathbf{U}_{\text{Had}} = H_c \otimes I_{\text{RR}}$, where:

$$H_c := \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix},$$

and let the controlled evolution operator be:

$$\mathbf{U}_{\text{ctrl}} := |\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}.$$

Then the full symmetric holosymmetric operator is given by:

$$\mathbf{U}_{\text{holo}}^{(\text{sym})} := \mathbf{U}_{\text{Had}} \cdot \mathbf{U}_{\text{ctrl}} \cdot \mathbf{U}_{\text{Had}}.$$

To express this in block form, we first recall the two distinct evolutionary path operators:

$$\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} := U_G^{(A)} \cdot U_N^{(B)}$$
$$\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} := U_N^{(A)} \cdot U_G^{(B)}$$

We also define the holosymmetric path sum and difference operators using this full notation:

$$\hat{\mathbf{U}}_+ := \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}$$
$$\hat{\mathbf{U}}_- := \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} - \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}$$

The matrix computation for $\mathbf{U}_{\text{holo}}^{(\text{sym})}$ then proceeds:

$$\begin{aligned}
\mathbf{U}_{\text{holo}}^{(\text{sym})} &= \left(\frac{1}{\sqrt{2}} \begin{pmatrix} I & I \\ I & -I \end{pmatrix} \right) \begin{pmatrix} \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} & 0 \\ 0 & \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} \end{pmatrix} \left(\frac{1}{\sqrt{2}} \begin{pmatrix} I & I \\ I & -I \end{pmatrix} \right) \\
&= \frac{1}{2} \begin{pmatrix} \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} & \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} - \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} \\ \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} - \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} & \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} \end{pmatrix} \\
&= \frac{1}{2} \begin{pmatrix} \hat{\mathbf{U}}_+ & \hat{\mathbf{U}}_- \\ \hat{\mathbf{U}}_- & \hat{\mathbf{U}}_+ \end{pmatrix}
\end{aligned}$$

We now write this operator in the computational basis as:

$$\mathbf{U}_{\text{holo}}^{(\text{sym})} = \sum_{i,j \in \{0,1\}} |\mathbf{i}\rangle_c\langle\mathbf{j}|_c \otimes \mathcal{F}_{ij}^{(\text{sym})},$$

where each component operator $\mathcal{F}_{ij}^{(\text{sym})} \in \mathcal{U}(\mathcal{H}_{\text{RR}}^{(2n)})$ is:

$$\boxed{\begin{aligned}
\mathcal{F}_{00}^{(\text{sym})} &= \frac{1}{2}\left(\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}\right) && (3.15) \\
\mathcal{F}_{01}^{(\text{sym})} &= \frac{1}{2}\left(\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} - \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}\right) && (3.16) \\
\mathcal{F}_{10}^{(\text{sym})} &= \frac{1}{2}\left(\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} - \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}\right) && (3.17) \\
\mathcal{F}_{11}^{(\text{sym})} &= \frac{1}{2}\left(\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}\right) && (3.18)
\end{aligned}}$$

Interpretation. These components reveal that:

- The diagonal terms $\mathcal{F}_{00}^{(\mathrm{sym})}$ and $\mathcal{F}_{11}^{(\mathrm{sym})}$ preserve the control branch but apply symmetric recombination of path operators.
- The off-diagonal terms $\mathcal{F}_{01}^{(\mathrm{sym})}$ and $\mathcal{F}_{10}^{(\mathrm{sym})}$ transfer amplitude between the control basis states and encode the asymmetry between paths.

Thus, the operator $\mathbf{U}_{\mathrm{holo}}^{(\mathrm{sym})}$ realizes a complete recursive interferometric transformation: it prepares a superposition, applies path-dependent dynamics, and recombines them, with its off-diagonal components enabling modal "scattering" between superposed histories.

3.2.2 Recursive Expansion: Asymmetric Interferometer

Let:

- $\mathbf{U}_{\mathrm{prep}} := U_c^{\mathrm{prep}} \otimes I_{\mathrm{RR}}$,
- $\mathbf{U}_{\mathrm{read}} := U_c^{\mathrm{read}} \otimes I_{\mathrm{RR}}$,

where $U_c^{\mathrm{prep}}, U_c^{\mathrm{read}} \in \mathrm{SU}(2)$ are parameterized as:

$$U_c^{\mathrm{prep}} = \begin{pmatrix} \alpha & -\beta^* \\ \beta & \alpha^* \end{pmatrix}, \qquad U_c^{\mathrm{read}} = \begin{pmatrix} \gamma & -\delta^* \\ \delta & \gamma^* \end{pmatrix}, \qquad |\alpha|^2 + |\beta|^2 = 1, \quad |\gamma|^2 + |\delta|^2 = 1.$$

The asymmetric holosymmetric operator is defined as:

$$\mathbf{U}_{\mathrm{holo}}^{(\mathrm{asym})} := \mathbf{U}_{\mathrm{read}} \cdot \mathbf{U}_{\mathrm{ctrl}} \cdot \mathbf{U}_{\mathrm{prep}},$$

with the control-dependent path evolution operator:

$$\mathbf{U}_{\mathrm{ctrl}} := |\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}.$$

We expand this over the computational basis $\{|\mathbf{0}\rangle_c, |\mathbf{1}\rangle_c\}$:

$$\mathbf{U}_{\mathrm{holo}}^{(\mathrm{asym})} = \sum_{i,j\in\{0,1\}} |\mathbf{i}\rangle_c\langle\mathbf{j}|_c \otimes \mathcal{F}_{ij}^{(\mathrm{asym})},$$

where each component operator $\mathcal{F}_{ij}^{(\mathrm{asym})} \in \mathcal{U}(\mathcal{H}_{\mathrm{RR}}^{(2n)})$ governs the transition $|\mathbf{j}\rangle_c \to |\mathbf{i}\rangle_c$.

Explicit Expansion.

$$\boxed{\begin{aligned}
\mathcal{F}_{00}^{(\mathrm{asym})} &= \gamma\alpha \cdot \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} - \delta^*\beta \cdot \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} && (3.19)\\
\mathcal{F}_{01}^{(\mathrm{asym})} &= -\gamma\beta^* \cdot \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} - \delta^*\alpha^* \cdot \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} && (3.20)\\
\mathcal{F}_{10}^{(\mathrm{asym})} &= \delta\alpha \cdot \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + \gamma^*\beta \cdot \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} && (3.21)\\
\mathcal{F}_{11}^{(\mathrm{asym})} &= -\delta\beta^* \cdot \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + \gamma^*\alpha^* \cdot \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} && (3.22)
\end{aligned}}$$

Interpretation. Each $\mathcal{F}_{ij}^{(\text{asym})}$ is a linear combination of the two evolution operators, weighted by amplitudes from the input and output control rotations. This expansion:

- **Encodes the full SU(2) steering freedom**, via arbitrary U_c^{prep} and U_c^{read},
- **Represents modal transitions between control channels**, including cross-channel interference,
- **Generalizes symmetric holosymmetry**, enabling off-diagonal manipulation of coherence pathways.

Hence, the operator $\mathbf{U}_{\text{holo}}^{(\text{asym})}$ provides a complete formal framework for recursive control-modulated interference with asymmetric path coupling and control axis alignment.

3.2.3 Interpretations of the Component Expansions within the Double-Slit Analogy

The **Recursive Component Expansion** provides the mathematical engine for the Double-Slit Experiment. The matrix elements $\mathcal{F}_{ij}$ for the Interferometer Model are precisely the amplitudes for a system that enters the apparatus in control state $|j\rangle_c$ to exit in control state $|i\rangle_c$. They are the building blocks of the final interference pattern.

Symmetric Interferometer: The Amplitudes of the Fringes In the symmetric case, the operator is $\mathbf{U}_{\text{holo}}^{(\text{sym})}$, and its components have a beautifully direct interpretation. These components can be derived as a special case of the more general asymmetric formalism. The symmetric model is defined by the condition that both the preparation and readout gates are the Hadamard gate:

$$U_c^{\text{prep}} = U_c^{\text{read}} = H = \frac{1}{\sqrt{2}}\begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$$

We begin with the general formula for the component operators derived from the asymmetric model:

$$\mathcal{F}_{ij} = (U_c^{\text{read}})_{i0}(U_c^{\text{prep}})_{0j} \cdot \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + (U_c^{\text{read}})_{i1}(U_c^{\text{prep}})_{1j} \cdot \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}$$

By substituting the matrix elements of the Hadamard gate ($H_{00} = H_{01} = H_{10} = 1/\sqrt{2}$ and $H_{11} = -1/\sqrt{2}$), we can calculate the specific components for the symmetric case.

For $\mathcal{F}_{00}^{(\text{sym})}$, we set $i = 0, j = 0$:

$$\begin{aligned}\mathcal{F}_{00}^{(\text{sym})} &= (H)_{00}(H)_{00} \cdot \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + (H)_{01}(H)_{10} \cdot \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} \\ &= \left(\frac{1}{\sqrt{2}}\right)\left(\frac{1}{\sqrt{2}}\right)\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + \left(\frac{1}{\sqrt{2}}\right)\left(\frac{1}{\sqrt{2}}\right)\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} = \frac{1}{2}\left(\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}\right)\end{aligned}$$

For $\mathcal{F}_{10}^{(\text{sym})}$, we set $i = 1, j = 0$:

$$\begin{aligned}\mathcal{F}_{10}^{(\text{sym})} &= (H)_{10}(H)_{00} \cdot \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + (H)_{11}(H)_{10} \cdot \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} \\ &= \left(\frac{1}{\sqrt{2}}\right)\left(\frac{1}{\sqrt{2}}\right)\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + \left(-\frac{1}{\sqrt{2}}\right)\left(\frac{1}{\sqrt{2}}\right)\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} = \frac{1}{2}\left(\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} - \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}\right)\end{aligned}$$

This demonstrates how the general formalism elegantly simplifies in the symmetric case.

Interpretation of Symmetric Components

- $\mathcal{F}_{00}^{(\mathbf{sym})} = \frac{1}{2}(\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G})$: This is the operator-valued amplitude for a system starting in $|\mathbf{0}\rangle_c$ to be measured in $|\mathbf{0}\rangle_c$. It is proportional to the *constructive interference* term $\hat{\mathbf{U}}_+$. This corresponds to the amplitude of the *"bright fringe"* at the center of our detector screen.
- $\mathcal{F}_{10}^{(\mathbf{sym})} = \frac{1}{2}(\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} - \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G})$: This is the amplitude for a system starting in $|\mathbf{0}\rangle_c$ to be measured in $|\mathbf{1}\rangle_c$. It is proportional to the *destructive interference* term $\hat{\mathbf{U}}_-$. This corresponds to the amplitude of the *"dark fringe"*.

The $\mathcal{F}_{01}$ and $\mathcal{F}_{11}$ components describe the same physical amplitudes but for a system initialized in the $|\mathbf{1}\rangle_c$ state.

Asymmetric Interferometer: Amplitudes for a "Polarized" Experiment In the more general asymmetric case, the $\mathcal{F}_{ij}^{(\mathrm{asym})}$ components are more complex because they must account for the arbitrary $SU(2)$ preparation and readout gates.

- $\mathcal{F}_{00}^{(\mathbf{asym})} = \gamma\alpha \cdot \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} - \delta^*\beta \cdot \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$: This is the amplitude for a system prepared in the state $\alpha|\mathbf{0}\rangle_c + \beta|\mathbf{1}\rangle_c$ to be projected onto the final basis state $\gamma^*|\mathbf{0}\rangle_c - \delta|\mathbf{1}\rangle_c$ and found in the $|\mathbf{0}\rangle_c$ component.

This is analogous to a double-slit experiment with *polarizers*. The $\mathbf{U}_{\mathrm{prep}}$ gate acts as an initial polarizer that prepares the "particle" in a specific quantum state before it reaches the slits. The $\mathbf{U}_{\mathrm{read}}$ gate acts as a final analyzer before the detector. The $\mathcal{F}_{ij}^{(\mathrm{asym})}$ components are the resulting amplitudes for this fully "polarized" experiment, allowing us to perform a complete tomographic reconstruction of the interference process. In both cases, the component expansion provides the precise mathematical link between the abstract operator algebra and the measurable probabilities of our generalized double-slit experiment.

3.3 The Interferometric Commutator Lensing

While the standard Interferometer Model describes a process unfolding in time (Prepare → Evolve → Recombine), the *Commutator Lensing* offers a more profound, algebraic perspective. It shifts the explanation for interference from a sequential, time-based process to a direct manifestation of the non-commutation of core quantum operations. The commutator, $[\hat{A}, \hat{B}] = \hat{A}\hat{B} - \hat{B}\hat{A}$, measures the fundamental "tension" or "twist" one operation induces upon the other.

3.3.1 Commutator Lensing of the Symmetric Interferometer

Definition 3.12 (Symmetric Interferometer Commutator Lensing)**.** Let $\mathbf{U}_{\mathrm{ctrl}}$ be the controlled path operator and $\mathbf{U}_{\mathrm{Had}} = (H_c \otimes I_{\mathrm{RR}})$ be the Hadamard gate acting on the control system. Define the *commutator lensing of the symmetric holosymmetric operator*:

$$\boxed{\mathbf{U}_{\text{holo}}^{(\text{sym-com})} := [\mathbf{U}_{\text{Had}}, \mathbf{U}_{\text{ctrl}}] \cdot \mathbf{U}_{\text{Had}}^{\dagger}.} \tag{3.23}$$

where:

- The commutator $[\mathbf{U}_{\text{Had}}, \mathbf{U}_{\text{ctrl}}]$ captures the *curvature* or failure of the preparation and evolution stages to commute.
- The post-multiplication by $\mathbf{U}_{\text{Had}}^{\dagger}$ returns the system to the preparation basis for final readout.

This operator acts nontrivially only if the path evolutions $\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}$ and $\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}$ fail to symmetrically transform under H_c, thus encoding interference-induced asymmetry into the dynamics.

Here, the commutator $[\mathbf{U}_{\text{Had}}, \mathbf{U}_{\text{ctrl}}]$ isolates the essential quantum effect. It represents the difference between "evolving a superposition" ($\mathbf{U}_{\text{ctrl}}\mathbf{U}_{\text{Had}}$) and "superposing an evolution" ($\mathbf{U}_{\text{Had}}\mathbf{U}_{\text{ctrl}}$). This difference is zero for a classical system but is the very source of interference in a quantum one. The operator thus measures the "curvature" in the space of operations induced by quantum mechanics. The subsequent multiplication by $\mathbf{U}_{\text{Had}}^{\dagger}$ simply rotates this abstract "twist" into a measurable basis.

3.3.2 Commutator Lensing of the Asymmetric Interferometer

Definition 3.13 (Asymmetric Interferometer Commutator Lensing)**.** Let $\mathbf{U}_{\text{prep}} = (U_c^{\text{prep}} \otimes I_{\text{RR}})$, $\mathbf{U}_{\text{read}} = (U_c^{\text{read}} \otimes I_{\text{RR}})$, and $\mathbf{U}_{\text{ctrl}}$ be defined as usual. The commutator lensing of the asymmetric holosymmetric operator is given by:

$$\boxed{\mathbf{U}_{\text{holo}}^{(\text{asym-com})} := [\mathbf{U}_{\text{ctrl}}, \mathbf{U}_{\text{prep}}] \cdot \mathbf{U}_{\text{read}}^{\dagger}.} \tag{3.24}$$

where:

- This operator quantifies how the chosen preparation basis fails to commute with the branching dynamics encoded in $\mathbf{U}_{\text{ctrl}}$.
- The readout transformation $\mathbf{U}_{\text{read}}^{\dagger}$ projects the result into a desired measurement basis.

This form generalizes the symmetric case and allows for experimental probing of interference asymmetries by comparing how path dynamics distort different interference bases.

Summary of Function The commutator formalism isolates the **dynamical twist** induced by interference. Specifically:

- $[\mathbf{U}_{\text{Had}}, \mathbf{U}_{\text{ctrl}}]$ in the symmetric case represents the algebraic torsion introduced by superposed branching.
- In the asymmetric case, $[\mathbf{U}_{\text{ctrl}}, \mathbf{U}_{\text{prep}}]$ reflects how the branching structure fails to preserve the superposition geometry established by $\mathbf{U}_{\text{prep}}$.

The commutator $[\mathbf{U}_{\text{ctrl}}, \mathbf{U}_{\text{prep}}]$ now asks a more sophisticated question: "How does the branching dynamic $\mathbf{U}_{\text{ctrl}}$ fail to preserve the specific superposition created by an arbitrary preparation $\mathbf{U}_{\text{prep}}$?" It quantifies the distortion or "shearing" of the initial quantum state geometry by the evolutionary process. The final $\mathbf{U}_{\text{read}}^{\dagger}$ allows us to view this distortion from any desired measurement perspective.

3.3.3 The Interferometric Commutator Lensing Double Slit Experiment

The operators we have defined, such as:

$$\mathbf{U}_{\text{holo}}^{(\text{sym-com})} := [\mathbf{U}_{\text{Had}}, \mathbf{U}_{\text{ctrl}}] \cdot \mathbf{U}_{\text{Had}}^{\dagger}$$

provide a powerful theoretical lens. They tell us *what to look for*. They identify the fundamental source of the interesting dynamics—the "dynamical twist" caused by the failure of the preparation and evolution gates to commute.

The Experiments as the Method of Observation The advanced experiments we devised are the *operational protocols* for measuring the effects of that twist. Consider the core of the operator above, the commutator $[\mathbf{U}_{\text{Had}}, \mathbf{U}_{\text{ctrl}}]$. An experiment to probe it would involve:

1. *Path A:* Prepare an initial state $|\Psi\rangle$, act on it with $\mathbf{U}_{\text{ctrl}}$, and then $\mathbf{U}_{\text{Had}}$.
2. *Path B:* Prepare the same state $|\Psi\rangle$, act on it with $\mathbf{U}_{\text{Had}}$, and then $\mathbf{U}_{\text{ctrl}}$.
3. *Measurement:* Coherently measure the difference between the final states of Path A and Path B.

This experimental procedure is precisely what gives physical, measurable substance to the abstract algebraic definition of the commutator. The *Advanced Interferometric and Path Scattering Experiments* we designed are, in fact, the very protocols needed to measure the consequences of the non-commutativity described by the Commutator Formalism. In defining the commutators, we provide the profound insight that the "dynamical twist" is the key phenomenon. Devising the experiments provides the practical means to observe and quantify that twist. One is the theory, the other is its verification. Both are indispensable.

3.3.4 Experimental Protocols for the Commutator Lensing

The Commutator lensing formalism provides a profound theoretical lens, reframing interference as a direct consequence of the non-commutativity of quantum operations. The following protocols describe how one would operationally measure the "dynamical twist" isolated by these commutators. Each experiment is itself a meta-interferometer, designed to compare two distinct sequences of operations.

Probing the Symmetric Commutator

To measure the physical effect of the commutator $[\mathbf{U}_{\text{Had}}, \mathbf{U}_{\text{ctrl}}]$, which quantifies the tension between creating a symmetric superposition and evolving its branches. This requires a two-path experiment where each path corresponds to a different ordering of operations.

1. Preparation Prepare two identical systems in the same initial state, $|\Psi_{\text{init}}\rangle = |\mathbf{0}\rangle_c \otimes |\Psi_{\text{RR}}\rangle$, one for each arm of our meta-experiment.

2. Arm A (Evolve then Superpose) In this arm, the system evolves along a definite path *before* being placed into a superposition.

1. *Step A1 (Evolve):* First, apply the controlled evolution operator to produce the *evolved state*:
$$|\Psi_{\text{evolved}}\rangle = \mathbf{U}_{\text{ctrl}}|\Psi_{\text{init}}\rangle = |\mathbf{0}\rangle_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}|\Psi_{\text{RR}}\rangle$$
2. *Step A2 (Superpose):* Second, apply the Hadamard operator to this evolved state to get the final state for Arm A:
$$|\Psi_{\text{E then S}}\rangle = \mathbf{U}_{\text{Had}}|\Psi_{\text{evolved}}\rangle = \frac{1}{\sqrt{2}}\left(|\mathbf{0}\rangle_c + |\mathbf{1}\rangle_c\right) \otimes \left(\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}|\Psi_{\text{RR}}\rangle\right)$$

3. Arm B (Superpose then Evolve) In this arm, the system is placed in a superposition *before* the evolution is applied.

1. *Step B1 (Superpose):* First, apply the Hadamard operator to create the *prepared superposition*:
$$|\Psi_{\text{prep}}\rangle = \mathbf{U}_{\text{Had}}|\Psi_{\text{init}}\rangle = \frac{1}{\sqrt{2}}\left(|\mathbf{0}\rangle_c + |\mathbf{1}\rangle_c\right) \otimes |\Psi_{\text{RR}}\rangle$$
2. *Step B2 (Evolve):* Second, apply the controlled evolution operator to this prepared state to get the final state for Arm B:
$$|\Psi_{\text{S then E}}\rangle = \mathbf{U}_{\text{ctrl}}|\Psi_{\text{prep}}\rangle = \frac{1}{\sqrt{2}}\left(|\mathbf{0}\rangle_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}|\Psi_{\text{RR}}\rangle + |\mathbf{1}\rangle_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}|\Psi_{\text{RR}}\rangle\right)$$

4. Measurement The physical observable is the interference between the final states of the two arms, $|\Psi_{\text{E then S}}\rangle$ and $|\Psi_{\text{S then E}}\rangle$. A measurement of an observable $\hat{O}$ would yield an interference term proportional to $\text{Re}\langle\Psi_{\text{E then S}}|\hat{O}|\Psi_{\text{S then E}}\rangle$. The difference between the two final states is directly proportional to the action of the commutator: $|\Psi_{\text{S then E}}\rangle - |\Psi_{\text{E then S}}\rangle = [\mathbf{U}_{\text{ctrl}}, \mathbf{U}_{\text{Had}}]|\Psi_{\text{init}}\rangle$.

Interpretation This protocol directly measures the degree to which the order of operations matters. A non-zero interference signal between the two arms is the physical manifestation of the non-commutativity of the operators, providing a direct measurement of the "dynamical twist" at the heart of the interferometric process.

3.3.5 Probing the Asymmetric Commutator

To measure the physical effect of the commutator $[\mathbf{U}_{\text{ctrl}}, \mathbf{U}_{\text{prep}}]$, which reveals how the branching dynamics of $\mathbf{U}_{\text{ctrl}}$ distort or "shear" an arbitrarily prepared superposition from $\mathbf{U}_{\text{prep}}$. This protocol requires a meta-interferometer that compares two parallel experimental arms.

1. Preparation Prepare two identical systems in the same initial state, $|\Psi_{\text{init}}\rangle = |\mathbf{0}\rangle_c \otimes |\Psi_{\text{RR}}\rangle$, one for each arm of the experiment.

2. Arm A: The Evolution of a Basis State This arm of the experiment first evolves the system along a definite path and *then* creates a superposition from the result.

1. *Step A1 (Evolve):* The system is first acted upon by the controlled evolution operator, forcing it down a single evolutionary path to produce the *evolved state*:

$$|\Psi_{\text{evolved}}\rangle = \mathbf{U}_{\text{ctrl}}|\Psi_{\text{init}}\rangle = |\mathbf{0}\rangle_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}|\Psi_{\text{RR}}\rangle$$

2. *Step A2 (Prepare):* The arbitrary preparation gate $\mathbf{U}_{\text{prep}}$ now acts on this *already evolved* state:

$$|\Psi_{\text{E then P}}\rangle = \mathbf{U}_{\text{prep}}|\Psi_{\text{evolved}}\rangle = (\alpha|\mathbf{0}\rangle_c + \beta|\mathbf{1}\rangle_c) \otimes \left(\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}|\Psi_{\text{RR}}\rangle\right)$$

3. Arm B: The Evolution of a Superposition This arm reverses the order, first creating a superposition and then evolving it.

1. *Step B1 (Prepare):* The preparation gate $\mathbf{U}_{\text{prep}}$ acts on the initial state, immediately creating the *prepared superposition*:

$$|\Psi_{\text{prep}}\rangle = \mathbf{U}_{\text{prep}}|\Psi_{\text{init}}\rangle = (\alpha|\mathbf{0}\rangle_c + \beta|\mathbf{1}\rangle_c) \otimes |\Psi_{\text{RR}}\rangle$$

2. *Step B2 (Evolve):* The controlled evolution operator $\mathbf{U}_{\text{ctrl}}$ now acts on this pre-existing superposition:

$$|\Psi_{\text{P then E}}\rangle = \mathbf{U}_{\text{ctrl}}|\Psi_{\text{prep}}\rangle = \alpha|\mathbf{0}\rangle_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}|\Psi_{\text{RR}}\rangle + \beta|\mathbf{1}\rangle_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}|\Psi_{\text{RR}}\rangle$$

4. Measurement: Quantifying the Dynamical Twist The physical result of the experiment is the interference between the final states of the two arms, $|\Psi_{\text{E then P}}\rangle$ and $|\Psi_{\text{P then E}}\rangle$. Their difference is precisely the action of the commutator:

$$|\Psi_{\text{P then E}}\rangle - |\Psi_{\text{E then P}}\rangle = [\mathbf{U}_{\text{ctrl}}, \mathbf{U}_{\text{prep}}]|\Psi_{\text{init}}\rangle$$

This experiment directly contrasts two fundamentally different processes. The degree to which the two final states differ is a direct measure of the "algebraic curvature" that the evolution $\mathbf{U}_{\text{ctrl}}$ imposes on the specific geometry of the superposition created by $\mathbf{U}_{\text{prep}}$.

3.4 The Holosymmetric Path Scattering Model

3.4.1 Symmetric Path Scattering Model

Definition 3.14 (Symmetric Path Scattering Holosymmetric Operator)**.** Let $H_c \in \text{SU}(2)$ denote the Hadamard gate acting on the control register. Define the controlled evolution operator:

$$\mathbf{U}_{\text{ctrl}} := |\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G},$$

and the Hadamard recombination gate:

$$\mathbf{U}_{\text{Had}} := H_c \otimes I_{\text{RR}}.$$

Then the *symmetric path scattering holosymmetric operator* is defined by the composition:

$$\boxed{\mathbf{U}_{\text{holo}}^{(\text{path-sym})} := \mathbf{U}_{\text{ctrl}} \cdot \mathbf{U}_{\text{Had}} \cdot \mathbf{U}_{\text{ctrl}}.} \tag{3.25}$$

3.4.2 Symmetric Path Scattering Model Double Slit Experiment

The Symmetric Path Scattering experiment models a recursive feedback loop where the system's evolution is modulated by an intermediate interference event. The procedure unfolds in three distinct acts.

Act I: Initial Evolution The process begins with the system in a simple basis state, $|\Psi_{\text{init}}\rangle = |\mathbf{0}\rangle_c \otimes |\Psi_{\text{RR}}\rangle$. The first controlled-unitary operator, $\mathbf{U}_{\text{ctrl}}$, is applied. This forces the system down a single, definite evolutionary path, producing the *evolved state*.

1. *Action:* $|\Psi_{\text{evolved}}\rangle = \mathbf{U}_{\text{ctrl}}|\Psi_{\text{init}}\rangle$
2. *Resulting State:*
$$|\Psi_{\text{evolved}}\rangle = |\mathbf{0}\rangle_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}|\Psi_{\text{RR}}\rangle$$

Act II: The Interference Event The Hadamard gate, $\mathbf{U}_{\text{Had}}$, now acts as a "scattering center." It takes the result of the first evolution and splits its control component into a coherent superposition, creating the *scattered state*.

1. *Action:* $|\Psi_{\text{scattered}}\rangle = \mathbf{U}_{\text{Had}}|\Psi_{\text{evolved}}\rangle$
2. *Resulting State:*
$$|\Psi_{\text{scattered}}\rangle = \frac{1}{\sqrt{2}}\left(|\mathbf{0}\rangle_c + |\mathbf{1}\rangle_c\right) \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}|\Psi_{\text{RR}}\rangle$$

Act III: Conditional Re-evolution The final $\mathbf{U}_{\text{ctrl}}$ operator acts on the newly created superposition. The system's subsequent evolution is now conditional on the outcome of the interference event in Act II.

1. *Action:* $|\Psi_{\text{final}}\rangle = \mathbf{U}_{\text{ctrl}}|\Psi_{\text{scattered}}\rangle$
2. *Resulting State:*
$$\frac{1}{\sqrt{2}}\left(|\mathbf{0}\rangle_c \otimes \left(\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}\right)^2 |\Psi_{\text{RR}}\rangle + |\mathbf{1}\rangle_c \otimes \left(\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}\, \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}\right)|\Psi_{\text{RR}}\rangle\right)$$

This final state is a superposition of two distinct second-order histories, where the system's fate depends directly on the interplay between its initial evolution and the intermediate interference. This symmetric form uses identical logic for both branchings (same $\mathbf{U}_{\text{ctrl}}$), reflecting a reversible, time-symmetric holosymmetric recursion.

The Symmetric Path Scattering Model reveals a profound mechanism of *quantum recursive feedback*, fundamentally distinct from the Interferometer Model's parallel-path interference. Here, the universe doesn't simply explore multiple histories in parallel—it evolves along one path, encounters an interference event that creates quantum uncertainty about its own history, and then continues evolving in a superposition that depends on this self-induced uncertainty. The operator structure $\mathbf{U}_{\text{ctrl}} \cdot \mathbf{U}_{\text{Had}} \cdot \mathbf{U}_{\text{ctrl}}$ encodes a temporal narrative: deterministic evolution leads to quantum scattering, which then conditions future evolution. This is not mere interference between alternatives, but a *dialectical process where the system's past determines how it will branch in the future.*

The three-act drama—Initial Evolution, Interference Event, and Conditional Re-evolution—models how quantum systems can exhibit *path-dependent memory through coherent feedback.* The final state reveals two remarkable histories: in one branch, the system evolves twice through the same parasymmetric assignment ($(\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N})^2$), reinforcing its initial trajectory; in the other, it switches paths mid-evolution ($\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}$), creating a hybrid history. This switching is not random but emerges from the quantum coherence introduced by the Hadamard scattering—the system literally interferes with its own evolutionary momentum, creating the possibility for self-modification.

Most profoundly, this model suggests that *the universe might possess a form of quantum reflexivity*—the ability to evolve, scatter off its own quantum state, and then continue evolving based on the interference pattern it creates with itself. Unlike the Interferometer Model's "which-path" detection, the Path Scattering Model implements "which-future" determination, where intermediate quantum events don't just reveal but actively sculpt subsequent evolution. The time-symmetric structure (using the same $\mathbf{U}_{\text{ctrl}}$ for both branchings) implies this process is fundamentally reversible, suggesting that just as the future can be conditioned by quantum interference in the past, the past might be retroactively influenced by future quantum scattering events—a truly radical vision of cosmic evolution as an ongoing dialogue between deterministic flow and quantum disruption.

3.4.3 Asymmetric Path Scattering Model

Definition 3.15 (Asymmetric Path Scattering Holosymmetric Operator)**.** Let $\mathbf{U}_{\text{ctrl}}^{\text{in}}$ and $\mathbf{U}_{\text{ctrl}}^{\text{out}}$ be two distinct controlled evolution operators acting on $\mathcal{H}_c \otimes \mathcal{H}_{\text{RR}}^{(2n)}$, defined by:

$$\mathbf{U}_{\text{ctrl}}^{\text{in}} := |\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G^{\text{in}} B_N^{\text{in}}} + |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_N^{\text{in}} B_G^{\text{in}}},$$

$$\mathbf{U}_{\text{ctrl}}^{\text{out}} := |\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G^{\text{out}} B_N^{\text{out}}} + |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_N^{\text{out}} B_G^{\text{out}}}.$$

Let the Hadamard gate on the control register be:

$$\mathbf{U}_{\text{Had}} := H_c \otimes I_{\text{RR}}.$$

Then the *asymmetric path scattering holosymmetric operator* is given by:

$$\boxed{\mathbf{U}_{\text{holo}}^{(\text{path-asym})} := \mathbf{U}_{\text{ctrl}}^{\text{out}} \cdot \mathbf{U}_{\text{Had}} \cdot \mathbf{U}_{\text{ctrl}}^{\text{in}}.} \tag{3.26}$$

3.4.4 Asymmetric Path Scattering Model Double Slit Experiment

Act I: The "Input" Evolution The process begins with the system in the initial state $|\Psi_{\text{init}}\rangle = |\mathbf{0}\rangle_c \otimes |\Psi_{\text{RR}}\rangle$. The first operator applied is $\mathbf{U}_{\text{ctrl}}^{(\text{in})}$, which represents the "input" or "before" set of dynamical laws. This forces the system along a single, definite path, producing the *input-evolved state*.

- **Action:** $|\Psi_{\text{in-evolved}}\rangle = \mathbf{U}_{\text{ctrl}}^{(\text{in})}|\Psi_{\text{init}}\rangle$
- **Resulting State:**

$$|\Psi_{\text{in-evolved}}\rangle = |\mathbf{0}\rangle_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G^{\text{in}} B_N^{\text{in}}}|\Psi_{\text{RR}}\rangle$$

Act II: The "Transformative Interference Event" The Hadamard gate, $\mathbf{U}_{\text{Had}}$, acts as a transformative event. It takes the outcome of the initial evolution and splits its control component into a coherent superposition, representing the moment of quantum interference that will condition the future. This creates the *scattered state*.

- **Action:** $|\Psi_{\text{scattered}}\rangle = \mathbf{U}_{\text{Had}}|\Psi_{\text{in-evolved}}\rangle$
- **Resulting State:**

$$|\Psi_{\text{scattered}}\rangle = \frac{1}{\sqrt{2}}\left(|\mathbf{0}\rangle_c + |\mathbf{1}\rangle_c\right) \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G^{\text{in}} B_N^{\text{in}}}|\Psi_{\text{RR}}\rangle$$

Act III: The "Output" Evolution Finally, the $\mathbf{U}_{\text{ctrl}}^{(\text{out})}$ operator is applied. This represents the "output" or "after" set of physical laws. It acts on the superposed state from Act II, and the system's final evolution is now governed by this new dynamical regime.

- **Action:** $|\Psi_{\text{final}}\rangle = \mathbf{U}_{\text{ctrl}}^{(\text{out})}|\Psi_{\text{scattered}}\rangle$
- **Resulting State:**

$$\frac{1}{\sqrt{2}}\left(|\mathbf{0}\rangle_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G^{\text{out}} B_N^{\text{out}}}\, \hat{\mathbf{U}}_{\text{RR}}^{A_G^{\text{in}} B_N^{\text{in}}}|\Psi_{\text{RR}}\rangle + |\mathbf{1}\rangle_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_N^{\text{out}} B_G^{\text{out}}}\, \hat{\mathbf{U}}_{\text{RR}}^{A_G^{\text{in}} B_N^{\text{in}}}|\Psi_{\text{RR}}\rangle\right)$$

This final state represents a superposition of histories where the system evolved first under the "in" rules, was subjected to a quantum superposition, and then continued its evolution under the new "out" rules. This allows modeling of feedback structures where the interference outcome conditions the subsequent evolution, possibly representing environmental decoherence, observer choice, or cosmogenic selection mechanisms.

The Asymmetric Path Scattering Model reveals a *fundamentally irreversible quantum process* where the laws of physics themselves can undergo transformation through interference

events. Unlike the symmetric model's reversible feedback loop, the operator structure $\mathbf{U}_{\text{ctrl}}^{\text{out}} \cdot \mathbf{U}_{\text{Had}} \cdot \mathbf{U}_{\text{ctrl}}^{\text{in}}$ encodes a temporal arrow: the universe evolves under one set of laws ("in"), encounters a transformative quantum event, then continues under potentially different laws ("out"). This models scenarios where *quantum interference doesn't just redirect evolution within fixed laws, but actually changes the laws themselves*—a mechanism for cosmological phase transitions, emergent complexity, or even observer-induced modifications to physical reality.

The three-act structure—Input Evolution, Transformative Interference Event, Output Evolution—captures how *singular quantum moments can serve as bridges between different physical regimes*. The final state shows the system in superposition between two histories: both branches initially evolved identically under the "in" laws, but after the Hadamard scattering, they diverge under different "out" assignments. The critical insight is that while both branches share the same past ($\hat{\mathbf{U}}_{\text{RR}}^{A_G^{\text{in}} B_N^{\text{in}}}$), they experience different futures ($\hat{\mathbf{U}}_{\text{RR}}^{A_G^{\text{out}} B_N^{\text{out}}}$ vs $\hat{\mathbf{U}}_{\text{RR}}^{A_N^{\text{out}} B_G^{\text{out}}}$). This asymmetry between shared history and divergent destiny models how *quantum events can act as cosmic watersheds*, after which the universe's evolutionary rules themselves bifurcate, thereby creating a "divergent destiny" from a "shared history". The asymmetry between the "in" and "out" states ('$\mathbf{U}_{\text{ctrl}}^{(\text{in})} \neq \mathbf{U}_{\text{ctrl}}^{(\text{out})}$') breaks the time-reversal symmetry of the process, introducing a powerful "arrow of complexity" that is the hallmark of a truly evolving, learning universe.

Most profoundly, this framework suggests that *the universe might undergo discrete transformative events that reconfigure its own operating system.* These could manifest as cosmological epoch transitions (inflation to radiation domination), symmetry-breaking events (electroweak phase transition), or even measurement-induced law selection (anthropic observer effects). The asymmetric structure implies these transformations are irreversible—once the universe passes through such a quantum gateway, there's no returning to the previous regime. This positions quantum interference not just as a phenomenon within physics, but as the mechanism by which physics itself evolves, suggesting that our current laws might be just one branch of a vast superposition of possible physical regimes, selected through a sequence of transformative interference events stretching back to the cosmos's quantum origin.

3.5 The Recursive Expansion of the Path Scattering Model

3.5.1 Recursive Component Expansion: Symmetric Path Scattering

We consider the composite operator:

$$\mathbf{U}_{\text{holo}}^{(\text{scatter,sym})} := \mathbf{U}_{\text{ctrl}} \cdot \mathbf{U}_{\text{Had}} \cdot \mathbf{U}_{\text{ctrl}},$$

where:

- The controlled evolution operator is given by:

$$\mathbf{U}_{\text{ctrl}} := |\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G},$$

- The Hadamard operator acting on the control system is:

$$\mathbf{U}_{\text{Had}} := H_c \otimes I_{\text{RR}}, \quad \text{where} \quad H_c = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}.$$

Let us introduce shorthand:

$$U_A := \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}, \qquad U_B := \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}.$$

Then the operator becomes:

$$\mathbf{U}_{\text{holo}}^{(\text{scatter,sym})} = \begin{pmatrix} U_A & 0 \\ 0 & U_B \end{pmatrix} \cdot \left(\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \right) \cdot \begin{pmatrix} U_A & 0 \\ 0 & U_B \end{pmatrix}.$$

Carrying out the matrix multiplication yields:

$$\mathbf{U}_{\text{holo}}^{(\text{scatter,sym})} = \frac{1}{\sqrt{2}} \begin{pmatrix} U_A^2 & U_A U_B \\ U_B U_A & -U_B^2 \end{pmatrix}.$$

We may then express this operator in control-basis decomposed form:

$$\mathbf{U}_{\text{holo}}^{(\text{scatter,sym})} = \sum_{i,j \in \{0,1\}} |\mathbf{i}\rangle_c \langle \mathbf{j}|_c \otimes \mathcal{S}_{ij}^{(\text{sym})},$$

where each matrix element $\mathcal{S}_{ij}^{(\text{sym})} \in \mathcal{U}(\mathcal{H}_{\text{RR}}^{(2n)})$ represents a distinct second-order scattering channel induced by the holosymmetric branching–interference–rebranching sequence.

$$\mathcal{S}_{00}^{(\text{sym})} = \frac{1}{\sqrt{2}} U_A^2 = \frac{1}{\sqrt{2}} \left(\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} \right)^2 \tag{3.27}$$

$$\mathcal{S}_{01}^{(\text{sym})} = \frac{1}{\sqrt{2}} U_A U_B = \frac{1}{\sqrt{2}} \left(\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} \, \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} \right) \tag{3.28}$$

$$\mathcal{S}_{10}^{(\text{sym})} = \frac{1}{\sqrt{2}} U_B U_A = \frac{1}{\sqrt{2}} \left(\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} \, \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} \right) \tag{3.29}$$

$$\mathcal{S}_{11}^{(\text{sym})} = -\frac{1}{\sqrt{2}} U_B^2 = -\frac{1}{\sqrt{2}} \left(\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} \right)^2 \tag{3.30}$$

Interpretation. These four components represent all second-order coherent transitions between the symmetric control basis states $|\mathbf{0}\rangle_c, |\mathbf{1}\rangle_c$, capturing the recursive structure of holosymmetric evolution across double-branch compositions. In particular:

- $\mathcal{S}_{00}^{(\text{sym})}$ corresponds to the constructive–constructive amplitude,
- $\mathcal{S}_{11}^{(\text{sym})}$ reflects a destructive–destructive feedback mode,
- The off-diagonal terms $\mathcal{S}_{01}^{(\text{sym})}, \mathcal{S}_{10}^{(\text{sym})}$ express interference transitions across distinct modal evolutions.

This decomposition enables experimental or simulation-based access to the full second-order dynamics of the symmetric holosymmetric interferometer.

3.5.2 Recursive Component Expansion: Asymmetric Path Scattering

We now consider a generalized recursive structure in which distinct control-dependent operators are applied before and after the interference layer:

$$\mathbf{U}_{\text{holo}}^{(\text{scatter,asym})} := \mathbf{U}_{\text{ctrl}}^{(\text{out})} \cdot \mathbf{U}_{\text{Had}} \cdot \mathbf{U}_{\text{ctrl}}^{(\text{in})},$$

where the "in" and "out" operators represent different sets of evolutionary rules:

- The input-controlled evolution is:

$$\mathbf{U}_{\text{ctrl}}^{(\text{in})} := |\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G^{\text{in}} B_N^{\text{in}}} + |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_N^{\text{in}} B_G^{\text{in}}},$$

- The output-controlled evolution is:

$$\mathbf{U}_{\text{ctrl}}^{(\text{out})} := |\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G^{\text{out}} B_N^{\text{out}}} + |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_N^{\text{out}} B_G^{\text{out}}},$$

and the Hadamard gate $\mathbf{U}_{\text{Had}}$ remains as previously defined.

We expand the composite operator $\mathbf{U}_{\text{holo}}^{(\text{scatter,asym})}$ over the control basis:

$$\mathbf{U}_{\text{holo}}^{(\text{scatter,asym})} = \sum_{i,j\in\{0,1\}} |\mathbf{i}\rangle_c\langle\mathbf{j}|_c \otimes \mathcal{S}_{ij}^{(\text{asym})},$$

where the explicit matrix multiplication yields the following operator-valued matrix elements:

$$\boxed{\begin{aligned}
\mathcal{S}_{00}^{(\text{asym})} &= \frac{1}{\sqrt{2}}\left(\hat{\mathbf{U}}_{\text{RR}}^{A_G^{\text{out}} B_N^{\text{out}}} \cdot \hat{\mathbf{U}}_{\text{RR}}^{A_G^{\text{in}} B_N^{\text{in}}}\right) && (3.31)\\
\mathcal{S}_{01}^{(\text{asym})} &= \frac{1}{\sqrt{2}}\left(\hat{\mathbf{U}}_{\text{RR}}^{A_G^{\text{out}} B_N^{\text{out}}} \cdot \hat{\mathbf{U}}_{\text{RR}}^{A_N^{\text{in}} B_G^{\text{in}}}\right) && (3.32)\\
\mathcal{S}_{10}^{(\text{asym})} &= \frac{1}{\sqrt{2}}\left(\hat{\mathbf{U}}_{\text{RR}}^{A_N^{\text{out}} B_G^{\text{out}}} \cdot \hat{\mathbf{U}}_{\text{RR}}^{A_G^{\text{in}} B_N^{\text{in}}}\right) && (3.33)\\
\mathcal{S}_{11}^{(\text{asym})} &= -\frac{1}{\sqrt{2}}\left(\hat{\mathbf{U}}_{\text{RR}}^{A_N^{\text{out}} B_G^{\text{out}}} \cdot \hat{\mathbf{U}}_{\text{RR}}^{A_N^{\text{in}} B_G^{\text{in}}}\right) && (3.34)
\end{aligned}}$$

Interpretation. This construction captures the most general form of recursive holosymmetric evolution currently formulated. Each component $\mathcal{S}_{ij}^{(\text{asym})}$ now explicitly shows the sequence of operations: an "in" evolution followed by an "out" evolution, corresponding to a specific scattering channel. This allows for the theoretical probing of second-order holosymmetric dynamics in fully generalized asymmetric scattering scenarios, where the rules of evolution themselves may change as a result of the interference event.

3.6 Experiment for Recursive Path Scattering Model

3.6.1 The Recursive Symmetric Path Scattering Case

Beyond the simple measurement of interference, the holosymmetric framework provides tools to model how interference itself can influence subsequent evolution. The *Symmetric Path Scattering Model* moves beyond the static comparison of the Interferometer Model into the realm of *dynamical feedback*. It describes a process where an initial evolution is subjected to a quantum interference event, and the outcome of that interference then dictates the system's future evolution.

The process is governed by the composite operator:

$$\mathbf{U}_{\text{holo}}^{(\text{scatter,sym})} := \mathbf{U}_{\text{ctrl}} \cdot \mathbf{U}_{\text{Had}} \cdot \mathbf{U}_{\text{ctrl}}$$

This sequence is not a simple measurement but a story of evolution, scattering, and re-evolution.

Initial State of the System The experiment begins with a well-defined initial quantum configuration for the physical system being studied. This is described by the state vector $|\Psi_{\text{RR}}\rangle$.

- **Formal Definition:** $|\Psi_{\text{RR}}\rangle$ represents the *joint quantum state of the dual PQS registers* (Register A and Register B). It is a vector in the dual-register Hilbert space $\mathcal{H}_{\text{RR}}^{(2n)} = \mathcal{H}_{\text{PQS}}^{(n,A)} \otimes \mathcal{H}_{\text{PQS}}^{(n,B)}$. It describes the complete quantum configuration, including all internal modes and entanglements, of the two n-PQS registers at the start of the experiment.
- **Role in the Experiment:** In this protocol, $|\Psi_{\text{RR}}\rangle$ serves as the *initial condition or "seed" state* that is subjected to the entire feedback process. The final outcome of the experiment is a direct consequence of how this specific initial configuration evolves under the recursive Path Scattering operator.

The Three Acts of Path Scattering The process begins with the composite system in the initial state $|\Psi_{\text{init}}\rangle = |\mathbf{0}\rangle_c \otimes |\Psi_{\text{RR}}\rangle$, where the control qubit selects the first evolutionary path.

Act I: Initial Evolution The first $\mathbf{U}_{\text{ctrl}}$ operator forces the system to evolve entirely along a single path, producing the *evolved state*.

- **Action:** $|\Psi_{\text{evolved}}\rangle = \mathbf{U}_{\text{ctrl}}|\Psi_{\text{init}}\rangle$
- **Resulting State:**

$$|\Psi_{\text{evolved}}\rangle = |\mathbf{0}\rangle_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}|\Psi_{\text{RR}}\rangle$$

Act II: The Interference Event The Hadamard gate, $\mathbf{U}_{\text{Had}}$, now acts on the result of this first evolution. It functions as a "scattering center" that takes the evolved state and splits its control component into a coherent superposition, creating the *scattered state*.

- **Action:** $|\Psi_{\text{scattered}}\rangle = \mathbf{U}_{\text{Had}}|\Psi_{\text{evolved}}\rangle$
- **Resulting State:**

$$|\Psi_{\text{scattered}}\rangle = \frac{1}{\sqrt{2}}\left(|\mathbf{0}\rangle_c + |\mathbf{1}\rangle_c\right) \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}|\Psi_{\text{RR}}\rangle$$

Act III: Conditional Re-evolution The final $\mathbf{U}_{\text{ctrl}}$ operator acts on the newly created superposition. The system's subsequent evolution is now conditional on the outcome of the scattering event in Act II.

- **Action:** $|\Psi_{\text{final}}\rangle = \mathbf{U}_{\text{ctrl}}|\Psi_{\text{scattered}}\rangle$
- **Resulting State:**

$$\frac{1}{\sqrt{2}}\left(|\mathbf{0}\rangle_c \otimes \left(\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}\right)^2 |\Psi_{\text{RR}}\rangle + |\mathbf{1}\rangle_c \otimes \left(\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}\, \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}\right)|\Psi_{\text{RR}}\rangle\right)$$

Interpretation of the Scattering Channels This process is compactly described by the operator matrix:

$$\mathbf{U}_{\text{holo}}^{(\text{scatter,sym})} = \frac{1}{\sqrt{2}}\begin{pmatrix} \left(\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}\right)^2 & \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} \\ \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} & -\left(\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}\right)^2 \end{pmatrix}$$

Each matrix element $\mathcal{S}_{ij}^{(\text{sym})}$ represents a distinct physical channel for this recursive process:

- $\mathcal{S}_{00}^{(\text{sym})}$: This is the amplitude for a system to start on Path A, scatter constructively, and continue on Path A. It represents a *self-reinforcing feedback loop*.
- $\mathcal{S}_{10}^{(\text{sym})}$: This is the amplitude for a system to start on Path A, scatter destructively (into the $|\mathbf{1}\rangle_c$ branch), and then be forced onto Path B. It represents a *history-altering interference*.
- The off-diagonal terms $\mathcal{S}_{01}^{(\text{sym})}$ and $\mathcal{S}_{10}^{(\text{sym})}$ capture the essence of the scattering: how interference can cause a "jump" between evolutionary trajectories.

This model describes a universe that can react to its own quantum state. It is a framework for *dynamics conditioned by interference*, a profound departure from the simple comparison of fixed histories offered by the Interferometer Model.

The Recursive Symmetric Path Scattering Model transforms the double-slit experiment from a passive measurement into an *active feedback mechanism* where quantum interference directly sculpts future evolution. Unlike the Interferometer Model's static comparison of parallel histories, this framework implements a temporal narrative where the universe

evolves deterministically along one path, encounters a quantum scattering event that creates superposition, and then continues evolving in a way that depends on the interference pattern it generated with itself. The matrix elements reveal distinct feedback channels: the diagonal terms $\mathcal{S}_{00}$ and $\mathcal{S}_{11}$ represent self-reinforcing loops where the system doubles down on its initial trajectory, while the off-diagonal terms $\mathcal{S}_{01}$ and $\mathcal{S}_{10}$ capture the profound possibility of *history-altering interference*—where quantum scattering can cause the universe to jump between evolutionary trajectories mid-flight.

This model presents a universe with *quantum reflexivity*, one that can sense and respond to its own quantum state via interference-mediated feedback. The three-act structure—Initial Evolution, Interference Event, Conditional Re-evolution—reveals cosmic evolution to be a continuous dialogue between deterministic flow and quantum disruption, supplanting the notion of a simple linear unfolding. Each interference event acts as a decision point where the universe's future is reshaped by the quantum coherence of its immediate past. In this light, quantum mechanics is re-envisioned as a theory of *adaptive evolution* that transcends its traditional role as a theory of measurement; reality continuously updates its own operating system through recursive cycles of evolution, self-interference, and path selection in a process analogous to a cosmic double-slit experiment, where the particle, having passed through one slit, can quantum-mechanically influence which slit it will traverse in the next iteration.

3.6.2 The Recursive Asymmetric Path Scattering Case

We now consider a generalized recursive structure that allows for the ultimate level of dynamical freedom: the evolutionary rules themselves may change as a result of an interference event. This is modeled by the *Asymmetric Path Scattering Operator*, where distinct control-dependent operators are applied before and after the central interference layer:

$$\mathbf{U}_{\text{holo}}^{(\text{scatter,asym})} := \mathbf{U}_{\text{ctrl}}^{(\text{out})} \cdot \mathbf{U}_{\text{Had}} \cdot \mathbf{U}_{\text{ctrl}}^{(\text{in})}$$

The Asymmetric Three-Act Process

This operator sequence describes a directed, potentially irreversible process of history-dependent evolution.

Act I: The "Input" Evolution The process begins with the system in the initial state $|\Psi_{\text{init}}\rangle = |\mathbf{0}\rangle_c \otimes |\Psi_{\text{RR}}\rangle$. The first operator applied is $\mathbf{U}_{\text{ctrl}}^{(\text{in})}$, which represents the "input" or "before" set of dynamical laws. This forces the system along a single, definite path according to these initial rules, producing the *input-evolved state*.

- **Action:** $|\Psi_{\text{evolved}}^{(\text{in})}\rangle = \mathbf{U}_{\text{ctrl}}^{(\text{in})}|\Psi_{\text{init}}\rangle$
- **Resulting State:**

$$|\Psi_{\text{evolved}}^{(\text{in})}\rangle = |\mathbf{0}\rangle_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G^{\text{in}} B_N^{\text{in}}}|\Psi_{\text{RR}}\rangle$$

Act II: The "Transformative Interference Event" The Hadamard gate, $\mathbf{U}_{\text{Had}}$, acts as a transformative event. It takes the outcome of the initial evolution and splits it into a coherent superposition, creating the *scattered state*.

- **Action:** $|\Psi_{\text{scattered}}\rangle = \mathbf{U}_{\text{Had}}|\Psi^{(\text{in})}_{\text{evolved}}\rangle$
- **Resulting State:**

$$|\Psi_{\text{scattered}}\rangle = \frac{1}{\sqrt{2}}\left(|\mathbf{0}\rangle_c + |\mathbf{1}\rangle_c\right) \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G^{\text{in}} B_N^{\text{in}}} |\Psi_{\text{RR}}\rangle$$

Act III: The "Output" Evolution Finally, the $\mathbf{U}^{(\text{out})}_{\text{ctrl}}$ operator is applied. This represents the "output" or "after" set of physical laws. It acts on the superposed state from Act II, and the system's final evolution is now governed by this new dynamical regime.

- **Action:** $|\Psi_{\text{final}}\rangle = \mathbf{U}^{(\text{out})}_{\text{ctrl}}|\Psi_{\text{scattered}}\rangle$
- **Resulting State:**

$$\frac{1}{\sqrt{2}}\left(|\mathbf{0}\rangle_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G^{\text{out}} B_N^{\text{out}}}\, \hat{\mathbf{U}}_{\text{RR}}^{A_G^{\text{in}} B_N^{\text{in}}} |\Psi_{\text{RR}}\rangle + |\mathbf{1}\rangle_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_N^{\text{out}} B_G^{\text{out}}}\, \hat{\mathbf{U}}_{\text{RR}}^{A_G^{\text{in}} B_N^{\text{in}}} |\Psi_{\text{RR}}\rangle\right)$$

Interpretation of the Asymmetric Scattering Channels

The operator matrix for this process reveals the intricate dynamics of this transformation:

$$\mathbf{U}^{(\text{scatter,asym})}_{\text{holo}} = \frac{1}{\sqrt{2}} \begin{pmatrix} \hat{\mathbf{U}}_{\text{RR}}^{A_G^{\text{out}} B_N^{\text{out}}} \hat{\mathbf{U}}_{\text{RR}}^{A_G^{\text{in}} B_N^{\text{in}}} & \hat{\mathbf{U}}_{\text{RR}}^{A_G^{\text{out}} B_N^{\text{out}}} \hat{\mathbf{U}}_{\text{RR}}^{A_N^{\text{in}} B_G^{\text{in}}} \\ \hat{\mathbf{U}}_{\text{RR}}^{A_N^{\text{out}} B_G^{\text{out}}} \hat{\mathbf{U}}_{\text{RR}}^{A_G^{\text{in}} B_N^{\text{in}}} & -\hat{\mathbf{U}}_{\text{RR}}^{A_N^{\text{out}} B_G^{\text{out}}} \hat{\mathbf{U}}_{\text{RR}}^{A_N^{\text{in}} B_G^{\text{in}}} \end{pmatrix}$$

Each component $\mathcal{S}^{(\text{asym})}_{ij}$ now represents a complete history of transformation:

- $\mathcal{S}^{(\text{asym})}_{00}$: This is the amplitude for a system to start on input Path 1, scatter constructively, and then evolve under the new output Path 1. It describes a transition between two different "laws of physics" within the same constructive channel.
- $\mathcal{S}^{(\text{asym})}_{10}$: This is the amplitude to start on input Path 1, scatter destructively, and then be forced to evolve under the new output Path 2. It represents a process where the interference event triggers a fundamental shift in the system's dynamical rules.

The Asymmetric Path Scattering Model represents the ultimate expression of quantum cosmological evolution: *a universe whose fundamental laws are not eternal edicts but living, adaptive structures that transform through quantum interference.* This framework transcends the conventional notion of physics as the discovery of immutable laws, instead revealing a cosmos that literally *learns* from its own quantum history. Each transformative interference event—marked by the transition from "input" to "output" control operators—represents a moment where the universe samples its own quantum state and updates its operating system accordingly. The mathematical structure $\mathbf{U}^{(\text{out})}_{\text{ctrl}} \cdot \mathbf{U}_{\text{Had}} \cdot \mathbf{U}^{(\text{in})}_{\text{ctrl}}$ encodes a profound asymmetry:

the universe that emerges from the interference event operates under different rules than the one that entered it.

This "learning universe" paradigm offers revolutionary insights into cosmological evolution and quantum measurement. *Cosmological phase transitions* become not mere rearrangements of matter but fundamental rewrites of physical law—the early universe's inflation-to-radiation transition might represent such a transformative interference event where the "output" physics bears little resemblance to the "input". Similarly, *quantum measurement* transforms from a mysterious collapse to a natural consequence of this framework: observation creates an interference event that doesn't just reveal the system's state but fundamentally alters how it will evolve thereafter. Most profoundly, this suggests the universe possesses a form of *quantum memory and adaptability*—its future dynamics are continuously sculpted by the accumulated interference patterns of its past, creating a cosmos that grows ever more complex and refined through recursive cycles of evolution, self-measurement, and law transformation. We inhabit not a clockwork universe grinding through predetermined motions, but a *quantum learning system* that discovers and refines its own principles through the eternal dance of interference and adaptation.

3.7 Commutator Lensing of the Path Scattering Model

3.7.1 Commutator Lensing of the Symmetric Path Scattering Model

Definition 3.16 (Commutator Lensing of the Symmetric Path Scattering Model)**.** Let $\mathbf{U}_{\text{ctrl}}$ be the controlled path operator and $\mathbf{U}_{\text{Had}} = (H_c \otimes I_{\text{RR}})$ be the Hadamard operator. Define the *symmetric commutator lensing of the path scattering operator* as:

$$\mathbf{U}_{\text{holo}}^{(\text{psym-com})} := \mathbf{U}_{\text{ctrl}} \cdot [\mathbf{U}_{\text{ctrl}}, \mathbf{U}_{\text{Had}}].$$

Interpretation:

- The evolution begins with conditional branching via $\mathbf{U}_{\text{ctrl}}$, followed by interference via $\mathbf{U}_{\text{Had}}$, and concludes with a second branching.
- The commutator $[\mathbf{U}_{\text{ctrl}}, \mathbf{U}_{\text{Had}}]$ quantifies the deviation from symmetric recombination and identifies modal scattering curvature induced by recursive interference.
- The outer $\mathbf{U}_{\text{ctrl}}$ recouples the branches after interference has reshaped them.

This captures the structural asymmetry introduced by modal recursion under fixed control logic.

3.7.2 Commutator Lensing of the Asymmetric Path Scattering

Definition 3.17 (Commutator Lensing of the Asymmetric Path Scattering Model)**.** Let $\mathbf{U}_{\text{ctrl}}^{\text{in}} = (C_c^{\text{in}} \otimes I_{\text{RR}})$, $\mathbf{U}_{\text{ctrl}}^{\text{out}} = (C_c^{\text{out}} \otimes I_{\text{RR}})$, and let $\mathbf{U}_{\text{Had}} = (H_c \otimes I_{\text{RR}})$. Define the *asymmetric commutator lensing of the path scattering operator* as:

$$\mathbf{U}_{\text{holo}}^{(\text{pasym-com})} := \mathbf{U}_{\text{ctrl}}^{\text{out}} \cdot [\mathbf{U}_{\text{Had}}, \mathbf{U}_{\text{ctrl}}^{\text{in}}].$$

Interpretation:

- This operator enacts initial conditional branching using $\mathbf{U}_{\text{ctrl}}^{\text{in}}$,
- Interference occurs via $\mathbf{U}_{\text{Had}}$, and
- Final conditional transformation $\mathbf{U}_{\text{ctrl}}^{\text{out}}$ post-selects modal pathways based on the interference outcome.

The commutator $[\mathbf{U}_{\text{Had}}, \mathbf{U}_{\text{ctrl}}^{\text{in}}]$ again reveals how interference reshapes the logical space of modal evolution, and how distinct forms of path logic affect this reshaping.

Summary of Commutator Lensing of Path Scattering The *commutator formalism* for the Path Scattering Model comes in two principal variants, each characterized by a distinct operator structure and interpretive role.

In the *symmetric case*, the holosymmetric evolution is governed by the operator

$$\mathbf{U}_{\text{ctrl}} \cdot [\mathbf{U}_{\text{ctrl}}, \mathbf{U}_{\text{Had}}],$$

which applies initial branching, captures the curvature induced by interference through the commutator, and reapplies the same control logic. This construction probes the deformation of modal structure induced by symmetric feedback recursion.

In the *asymmetric case*, the evolution is governed by the operator

$$\mathbf{U}_{\text{ctrl}}^{\text{out}} \cdot [\mathbf{U}_{\text{Had}}, \mathbf{U}_{\text{ctrl}}^{\text{in}}],$$

where distinct pre- and post-interference control logics are employed. This formulation models non-identical modal rebranching conditioned on the intermediate interference outcome, revealing how different control strategies sculpt recursive structure. Both forms operationalize holosymmetry as an algebra of commutators, offering a direct measure of modal incompatibility and recursive curvature within the interference-driven quantum evolution of dual PQS configurations.

3.7.3 The Commutator Lensing Double-Slit Experiments

The Commutator formalism provides a profound theoretical lens, reframing interference as a direct consequence of the non-commutativity of quantum operations. The following protocols describe how one would operationally measure the "dynamical twist" isolated by these commutators. Each experiment is itself a meta-interferometer, designed to compare two distinct sequences of operations.

Probing the Symmetric Path Scattering Commutator

To measure the physical effect of the operator $\mathbf{U}_{\text{holo}}^{(\text{psym-com})} = \mathbf{U}_{\text{ctrl}} \cdot [\mathbf{U}_{\text{ctrl}}, \mathbf{U}_{\text{Had}}]$, which describes how an interference event is filtered through a final evolutionary stage.

The Experimental Protocol This is a three-stage, comparative experiment.

1. **Stage 1: Create the two parallel histories.**

- *Arm A (Evolve then Superpose):* Prepare the state $|\Psi_{\text{E then S}}\rangle = \mathbf{U}_{\text{Had}}\mathbf{U}_{\text{ctrl}}|\Psi_{\text{init}}\rangle$.
- *Arm B (Superpose then Evolve):* Prepare the state $|\Psi_{\text{S then E}}\rangle = \mathbf{U}_{\text{ctrl}}\mathbf{U}_{\text{Had}}|\Psi_{\text{init}}\rangle$.

2. **Stage 2: Isolate the "Dynamical Twist".** The physical difference between the two arms is the action of the commutator. We can formally define the "commutator state" as their difference:

$$|\Psi_{\text{commutator}}\rangle := |\Psi_{\text{S then E}}\rangle - |\Psi_{\text{E then S}}\rangle = [\mathbf{U}_{\text{ctrl}}, \mathbf{U}_{\text{Had}}]|\Psi_{\text{init}}\rangle$$

3. **Stage 3: Apply Final Evolution and Measure.** The full Path Scattering operator requires a final application of $\mathbf{U}_{\text{ctrl}}$. We apply this to the commutator state:

$$|\Psi_{\text{final}}\rangle = \mathbf{U}_{\text{ctrl}}|\Psi_{\text{commutator}}\rangle = \mathbf{U}_{\text{ctrl}}[\mathbf{U}_{\text{ctrl}}, \mathbf{U}_{\text{Had}}]|\Psi_{\text{init}}\rangle$$

   A measurement of an observable $\hat{O}$ on this final state would reveal the physical consequences of this full process.

Interpretation This experiment measures how the fundamental "twist" between superposition and evolution is itself transformed by a subsequent evolution.

Probing the Asymmetric Path Scattering Commutator

To measure the effect of the most general operator, $\mathbf{U}_{\text{holo}}^{(\text{pasym-com})} = \mathbf{U}_{\text{ctrl}}^{(\text{out})} \cdot [\mathbf{U}_{\text{Had}}, \mathbf{U}_{\text{ctrl}}^{(\text{in})}]$, which models a change in physical law after an interference event.

The Experimental Protocol

1. **Stage 1: Create two parallel histories based on the "in" laws.**

   - *Arm A:* $|\Psi_{\text{E then S}}\rangle = \mathbf{U}_{\text{Had}}\mathbf{U}_{\text{ctrl}}^{(\text{in})}|\Psi_{\text{init}}\rangle$
   - *Arm B:* $|\Psi_{\text{S then E}}\rangle = \mathbf{U}_{\text{ctrl}}^{(\text{in})}\mathbf{U}_{\text{Had}}|\Psi_{\text{init}}\rangle$

2. **Stage 2: Isolate the "Input Twist".** Define the "input commutator state":

$$|\Psi_{\text{commutator-in}}\rangle := |\Psi_{\text{S then E}}\rangle - |\Psi_{\text{E then S}}\rangle = [\mathbf{U}_{\text{ctrl}}^{(\text{in})}, \mathbf{U}_{\text{Had}}]|\Psi_{\text{init}}\rangle$$

3. **Stage 3: Apply "Output" Laws and Measure.** Apply the new, "output" evolution operator to the result of the initial interference:

$$|\Psi_{\text{final}}\rangle = \mathbf{U}_{\text{ctrl}}^{(\text{out})}|\Psi_{\text{commutator-in}}\rangle = \mathbf{U}_{\text{ctrl}}^{(\text{out})}[\mathbf{U}_{\text{ctrl}}^{(\text{in})}, \mathbf{U}_{\text{Had}}]|\Psi_{\text{init}}\rangle$$

The Commutator Formalism transforms abstract algebraic concepts into measurable physical quantities by constructing *meta-interferometers* that probe the fundamental non-commutativity of quantum operations. Rather than measuring interference between states, these experiments measure interference between *sequences of operations*—revealing what the framework calls the "dynamical twist" encoded in commutators like $[\mathbf{U}_{\text{ctrl}}, \mathbf{U}_{\text{Had}}]$. The experimental protocol is elegantly comparative: create two parallel histories by applying the same operations in different orders (evolve-then-superpose vs. superpose-then-evolve), subtract them to isolate the pure commutator state, then apply a final transformation to make this abstract quantity observable. This commutator state $|\Psi_{\text{commutator}}\rangle$ represents the irreducible quantum tension between different operational sequences—a physical manifestation of the fact that in quantum mechanics, the order of operations fundamentally matters.

The symmetric protocol measures how this fundamental twist is itself transformed by subsequent evolution, revealing how *interference begets interference* in recursive cycles. But the asymmetric protocol achieves something even more profound: it operationalizes the concept of a "learning universe" by measuring how the algebraic tension created under one set of laws ($\mathbf{U}_{\text{ctrl}}^{(\text{in})}$) is projected and transformed by an entirely different set ($\mathbf{U}_{\text{ctrl}}^{(\text{out})}$). This provides the experimental signature for cosmological phase transitions and law evolution—the output measurement reveals how the "memory" of non-commutativity under old physics manifests under new physics.

These meta-interferometric protocols represent a new class of quantum experiments that transcend the probing of states or processes to investigate the *algebraic structure of quantum mechanics itself*. By making commutators—traditionally abstract mathematical objects—into measurable quantities, the framework suggests that the universe's deepest properties lie in its operational grammar, rather than its content: the patterns of non-commutativity that govern how different quantum operations fail to commute. This elevates interference from a phenomenon occurring within fixed laws to the very mechanism by which the laws themselves are structured, measured, and potentially transformed—turning the double-slit experiment into a probe of the universe's fundamental algebraic architecture.

3.8 Definition: Holosymmetry (Holothesis)

Definition 3.18 (Holosymmetry (Holothesis): Formal Definition)**.** Fix two disjoint n-PQS registers and a control qubit as in Definitions 2.101 and 2.102:

$$\mathcal{H}_{\text{Holo}} := \mathcal{H}_c \otimes \mathcal{H}_{\text{RR}}^{(2n)} = \mathcal{H}_c \otimes \left(\mathcal{H}_{\text{PQS}}^{(n,A)} \otimes \mathcal{H}_{\text{PQS}}^{(n,B)}\right), \qquad \dim \mathcal{H}_{\text{Holo}} = 2^{4n+1}.$$

Let $U_G^{(\odot)}$ and $U_N^{(\odot)}$ be the global–coupled and decoupled parasymmetric evolutions on a single register (Definitions 2.66 and 2.85), and let the *lifted path unitaries* on $\mathcal{H}_{\text{RR}}^{(2n)}$ be

$$\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} := U_G^{(A)} U_N^{(B)}, \qquad \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} := U_N^{(A)} U_G^{(B)} \quad \text{(Definitions 2.103 and 2.108)}.$$

(H1) Holosymmetric unitary (branching form). A *holosymmetric realization* is any unitary $\mathbf{U}_{\text{holo}} \in \mathcal{U}(\mathcal{H}_{\text{Holo}})$ that is control-block linear with respect to the computational basis of $\mathcal{H}_c$,

up to local control pre/post rotations: there exist unitaries $\mathbf{U}_{\text{prep}}, \mathbf{U}_{\text{read}} \in \mathcal{U}(\mathcal{H}_c)$ such that

$$\mathbf{U}_{\text{holo}} = (\mathbf{U}_{\text{read}} \otimes I_{\text{RR}}^{(2n)}) \Big(|\mathbf{0}\rangle\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + |\mathbf{1}\rangle\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} \Big) (\mathbf{U}_{\text{prep}} \otimes I_{\text{RR}}^{(2n)}). \tag{3.35}$$

Canonical choices recover the standard models:

$$\mathbf{U}_{\text{ctrl}} \ (\mathbf{U}_{\text{prep}} = \mathbf{U}_{\text{read}} = I_c), \qquad \mathbf{U}_{\text{Had}} \ (\mathbf{U}_{\text{prep}} = \mathbf{U}_{\text{read}} = H_c),$$

and their commutator-lensing variants (Definitions 2.114 and 2.119).

(H2) Holosymmetric effect (valuation link). For control preparation $|\mathbf{a}\rangle_c \in \mathcal{H}_c$ and readout $|\mathbf{b}\rangle_c \in \mathcal{H}_c$, the *holosymmetric effect* on the dual-register space is the positive operator

$$E_{\text{Holo}}^{(\mathbf{a}\twoheadrightarrow\mathbf{b})} := \langle \mathbf{b}|_c \, \mathbf{U}_{\text{holo}} \Big(|\mathbf{a}\rangle\langle\mathbf{a}|_c \otimes I_{\text{RR}}^{(2n)} \Big) \mathbf{U}_{\text{holo}}^{\dagger} \, |\mathbf{b}\rangle_c \ \in \ \mathcal{B}^{+}\Big(\mathcal{H}_{\text{RR}}^{(2n)}\Big). \tag{3.36}$$

Given a register state ρ_{RR}, the *Holothesis valuation* is the Born propensity $v_{\text{Holothesis}}(\rho_{\text{RR}}) = \text{Tr}\Big(E_{\text{Holo}}^{(\mathbf{a}\twoheadrightarrow\mathbf{b})} \, \rho_{\text{RR}}\Big)$ (Definition 1.18).

(H3) Ontological identification. Holosymmetry is the fourth corner (Holothesis) of the Fourfold Dialectic Algebra (Definition 1.16). Its ontological label $\mathring{\mathbf{\Lambda}}_{\mathring{\mathbf{1}}}^{\mathring{\mathbf{0}}}$ denotes the meta-operator that coherently compares and interferes *entire* evolutionary histories —the two path unitaries above—rather than individual states.

(H4) Equivalences. Holosymmetric realizations related by (i) control-basis changes $\mathbf{U}_{\text{prep}}, \mathbf{U}_{\text{read}} \in \mathcal{U}(\mathcal{H}_c)$, (ii) a global register swap $A \leftrightarrow B$, or (iii) common conjugation $W \in \mathcal{U}(\mathcal{H}_{\text{RR}}^{(2n)})$ (i.e. $\hat{\mathbf{U}}_{A,B} \mapsto W \, \hat{\mathbf{U}}_{A,B} \, W^{\dagger}$), are *physically equivalent*: they define the same family of effects $\{E_{\text{Holo}}^{(\mathbf{a}\twoheadrightarrow\mathbf{b})}\}$ and valuations.
In particular, the "lensing'' presentations (sym-com, asym-com, psym-com, pasym-com) are obtained by such control pre/post rotations and thus lie in the same equivalence class. Writing the controlled form as

$$\mathbf{U}_{\text{ctrl}} \ = \ I_c \otimes U_+ \ + \ Z_c \otimes U_-, \qquad U_{\pm} := \tfrac{1}{2}\Big(\hat{\mathbf{U}}_A \pm \hat{\mathbf{U}}_B\Big),$$

(with Z_c the Pauli-Z on $\mathcal{H}_c$), any control rotation R satisfies

$$[R, \mathbf{U}_{\text{ctrl}}] \ = \ [R, Z_c] \otimes U_-,$$

so the commutator filters out the sum U_+ and isolates the difference $U_- = \frac{1}{2}(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)$ without introducing new resources; conjugation by $\mathbf{U}_{\text{read}}^{\dagger}$ simply undoes the chosen control basis. Thus the lensing forms are reparametrizations of (3.35) under the (H4) moves (Definition 3.18).

In this sense, *Holosymmetry* is the unitary-equivalence class of triples

$$\Big(\mathcal{H}_{\text{Holo}}, \ \mathbf{U}_{\text{holo}}, \ \{E_{\text{Holo}}^{(\mathbf{a}\twoheadrightarrow\mathbf{b})}\} \Big)$$

that satisfy (3.35)–(3.36). It *operationalizes* the Holothesis: a meta-level synthesis testing the similarity/dissonance between alternative parasymmetric evolutions via controlled interference.

Notes.

(i) The three model "presentations''—interferometric, path-scattering, and commutator-lensing—are recovered by specific choices of $\mathbf{U}_{\text{prep}}$, $\mathbf{U}_{\text{read}}$ and by expanding (3.35) in the $\{\,|\mathbf{0}\rangle_c, |\mathbf{1}\rangle_c\,\}$ or $\{\,|+\rangle_c, |-\rangle_c\,\}$ bases; see Definitions 2.114 and 2.119 and the lensing definitions therein.

(ii) In the emergent/GNS layer, all valuations lift to standard expectations on $\mathcal{H}_{\overset{\circ}{\mathbf{\Psi}}} = \mathcal{H}_{\text{em}}$ via $\pi_{\overset{\circ}{\mathbf{\Psi}}}$ (cf. Definition 1.34).

Intuitive reading. Holosymmetry asks a meta-question: not "which state is realized?" but "how similar or dissonant are *entire* evolutionary histories?" The normal form (3.35) makes this concrete: the control register coherently addresses two global evolutions on the dual PQS registers, $\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}$ and $\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}$, and then recombines them. What the experiment reads out is not a yes/no judgment but a *propensity*, the Born valuation of the holosymmetric effect $E_{\text{Holo}}^{(\mathbf{a}\to\mathbf{b})}$ in (3.36). Large values signal "constructive agreement'' of the histories under the chosen preparation/readout; small values highlight "dissonance'' (destructive interference). This realizes the Holothesis corner of the Fourfold Dialectic Algebra (Definition 1.16) at the level of *processes* rather than states.

Operational picture. Think of $E_{\text{Holo}}^{(\mathbf{a}\to\mathbf{b})}$ as a matched filter on dynamics: by selecting $(\mathbf{a}, \mathbf{b})$ we decide which relational feature of the two histories to test. Choosing the computational basis emphasizes the "same-port'' vs. "opposite-port'' outcomes (synthesis vs. dissonance), while the $\{|+\rangle_c, |-\rangle_c\}$ basis reveals complementary structure, recovering the interferometer, path-scattering, and lensing presentations via the settings of $\mathbf{U}_{\text{prep}}$ and $\mathbf{U}_{\text{read}}$ (cf. (3.35) and Notes (i)). Crucially, Holosymmetry is an *equivalence class* of such realizations: local rotations of the control, common conjugations on the registers, or swapping the labels $A \leftrightarrow B$ leave the family of effects—and thus all valuations—unchanged (see Definition 3.18).

Bridge to the emergent layer. All holosymmetric valuations are ordinary quantum expectations once lifted to the emergent/GNS layer: $\omega^{\overset{\circ}{\mathbf{\Psi}}}(E) = \langle \Omega_{\omega^{\overset{\circ}{\mathbf{\Psi}}}} | \pi_{\overset{\circ}{\mathbf{\Psi}}}(E) | \Omega_{\omega^{\overset{\circ}{\mathbf{\Psi}}}} \rangle$ and, for any preparation ρ on $\mathcal{H}_{\overset{\circ}{\mathbf{\Psi}}} = \mathcal{H}_{\text{em}}$, $v(\rho; E) = \text{Tr}(\rho\, \pi_{\overset{\circ}{\mathbf{\Psi}}}(E))$ (Definitions 1.18 and 1.34). Thus the compact definition packages an ontological claim (Holothesis as the fourth corner), an operational mechanism (controlled comparison of histories), and a mathematical realization (effects and Born valuations) into a single pipeline from logic to laboratory readout—without invoking indefinite causal order. If one ever wished to superpose *orders* rather than *which history*, that would require an explicit higher-order extension; Holosymmetry here concerns coherent comparison of fixed-order global evolutions.

3.8.1 Holosymmetry (Holothesis): Summary of Quantum Construction

We now summarize the quantum construction of *Holosymmetry*, the Holothesis in the Fourfold Dialectic Algebra of Symmetry as follows: **Holosymmetry** is the fourth and highest principle in the *Fourfold Dialectic Algebra of Symmetry* (Definition 1.16). As the *Holothesis*, it governs the coherent superposition and interference of entire evolutionary histories, placing dynamical laws themselves into quantum superposition.

1. Ontological Architecture

The holosymmetric operator

$$\mathring{\mathbf{\Lambda}}_{\mathring{\mathbf{0}}\mathring{\mathbf{1}}}$$

is characterized by three fundamental ontological modes:

(i) *Symmetry* ($\mathring{\mathbf{0}}$): identity and coherence (Thesis)

(ii) *Antisymmetry* ($\mathring{\mathbf{1}}$): difference and negation (Antithesis)

(iii) *Parasymmetry* ($\mathring{\mathbf{0}} + \mathring{\mathbf{1}}$): dialectical synthesis of the dual modes

Unlike the lower modes, Holosymmetry operates at a meta-level where these constituents are not external elements but integral to its logical structure. It unifies the complete dialectical framework within a single transcendent principle.

2. Operational Functionality: Synthesis of Dynamical Processes

Holosymmetry enables a **holopositional synthesis** of distinct modal trajectories. This synthesis is not a single process but is realized through three complementary operational modes, each designed to probe a different aspect of the relationship between universal histories. Symbolically, we write:

$$\mathring{\mathbf{\Lambda}}_{\mathring{\mathbf{0}}\mathring{\mathbf{1}}} \doteq \mathfrak{S}\left[\mathring{\mathbf{0}},\ \mathring{\mathbf{1}},\ \mathring{\mathbf{0}}+\mathring{\mathbf{1}};\ \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}, \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}\right]$$

where the synthesis functional $\mathfrak{S}[\cdot]$ now represents one of the following three distinct functions:

1. *Interferometric Comparison:* coherently superposes parallel evolutionary paths to measure their relational difference, acting as a cosmic interferometer.

2. *Recursive Scattering:* models dynamical feedback, where the outcome of an interference event transforms the subsequent evolutionary laws of the system.

3. *Algebraic Lensing:* isolates the non-commutative "twist" between quantum operations as the fundamental algebraic source of all interference phenomena.

3. Mathematical Realization

Holosymmetry is implemented through unitary operators acting on the composite Hilbert space:

$$\mathbf{U}_{\mathrm{holo}} \in \mathcal{U}\left(\mathcal{H}_c \otimes \mathcal{H}_{\mathrm{RR}}^{(2n)}\right)$$

where:

- $\mathcal{H}_c \cong \mathbb{C}^2$ is the control register space
- $\mathcal{H}_{\mathrm{RR}}^{(2n)} = \bigotimes_{i=1}^{2n} \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_{x_i})}$ is the dual PQS register space
- Each PQS modal space $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)} \cong \mathbb{C}^4$ encodes the complete fourfold dialectical structure

The total holosymmetric Hilbert space has dimension:

$$\dim(\mathcal{H}_{\text{Holo}}) = 2^{4n+1}$$

The holosymmetric operator decomposes as:

$$\mathring{\mathbf{\Lambda}}_{\mathring{\mathbf{i}}}^{\mathring{\mathbf{0}}} = \sum_{i,j\in\{0,1\}} |\mathbf{i}\rangle_c\langle\mathbf{j}|_c \otimes \Lambda_{ij}^{(\text{model})}$$

where the three canonical models are:

1. **Interferometric Model:**

$$\mathbf{U}_{\text{holo}}^{(\text{interferometer})} = \mathbf{U}_{\text{read}} \cdot \mathbf{U}_{\text{ctrl}} \cdot \mathbf{U}_{\text{prep}}$$

   Enables full tomographic control over superposed evolutionary paths.

2. **Path Scattering Model:**

$$\mathbf{U}_{\text{holo}}^{(\text{scatter})} = \mathbf{U}_{\text{ctrl}}^{(\text{out})} \cdot \mathbf{U}_{\text{Had}} \cdot \mathbf{U}_{\text{ctrl}}^{(\text{in})}$$

   Captures irreversible dynamics and transitions between physical regimes.

3. **Commutator Lensing:** Isolates the algebraic source of quantum interference by examining:

   (a) *Symmetric Interferometer Lensing*: Probes the fundamental twist in a balanced system.

$$\mathbf{U}_{\text{holo}}^{(\text{sym-com})} := [\mathbf{U}_{\text{Had}}, \mathbf{U}_{\text{ctrl}}] \cdot \mathbf{U}_{\text{Had}}^{\dagger}$$

   (b) *Asymmetric Interferometer Lensing*: Probes how an arbitrary preparation is distorted by the evolution.

$$\mathbf{U}_{\text{holo}}^{(\text{asym-com})} := [\mathbf{U}_{\text{ctrl}}, \mathbf{U}_{\text{prep}}] \cdot \mathbf{U}_{\text{read}}^{\dagger}$$

   (c) *Symmetric Path Scattering Lensing*: Probes how the fundamental twist evolves under a feedback loop.

$$\mathbf{U}_{\text{holo}}^{(\text{psym-com})} := \mathbf{U}_{\text{ctrl}} \cdot [\mathbf{U}_{\text{ctrl}}, \mathbf{U}_{\text{Had}}]$$

   (d) *Asymmetric Path Scattering Lensing*: Probes how the twist generated by "in" laws is transformed by "out" laws.

$$\mathbf{U}_{\text{holo}}^{(\text{pasym-com})} := \mathbf{U}_{\text{ctrl}}^{\text{out}} \cdot [\mathbf{U}_{\text{Had}}, \mathbf{U}_{\text{ctrl}}^{\text{in}}]$$

   Provides direct access to the non-commutative structure underlying quantum dynamics.

Valuation Link With control prepared in $|\mathbf{a}\rangle_c$ and read out in $|\mathbf{b}\rangle_c$, the Holothesis valuation is $v_{\text{Holothesis}}(\rho_{\text{RR}}) = \text{Tr}\Big(E^{(\mathbf{a}\to\mathbf{b})}_{\text{Holothesis}}\,\rho_{\text{RR}}\Big)$ as defined in Definition 1.18.

Let the operator $\hat{\mathbf{U}}_A$ describes a universal history where the PQSs in Register A evolve as a single, indivisible entangled system, while the PQSs in Register B evolve in a modular, separable fashion:

$$\hat{\mathbf{U}}_A \equiv \hat{\mathbf{U}}^{A_G B_N}_{\text{RR}} := \Big(U^{(\odot)}_G \otimes \mathbb{I}^{(B)}\Big) \cdot \Big(\mathbb{I}^{(A)} \otimes U^{(\odot)}_N\Big)$$

And let the operator $\hat{\mathbf{U}}_B$ describes the alternate universal history where the PQSs in Register B are fully entangled in their dynamics, while Register A evolves in a separable manner:

$$\hat{\mathbf{U}}_B \equiv \hat{\mathbf{U}}^{A_N B_G}_{\text{RR}} := \Big(U^{(\odot)}_N \otimes \mathbb{I}^{(B)}\Big) \cdot \Big(\mathbb{I}^{(A)} \otimes U^{(\odot)}_G\Big)$$

Then the holosymmetric operator expands in the following manner:

1. The Interferometer Model Expansion This model realizes Holosymmetry as a process of *Prepare* → *Evolve* → *Recombine* to probe the interference between universal histories.

(a) Symmetric Interferometer

$$\mathbf{U}^{(\text{sym})}_{\text{holo}} = \mathbf{U}_{\text{Had}} \cdot \mathbf{U}_{\text{ctrl}} \cdot \mathbf{U}_{\text{Had}}$$

Ontological Expansion:

$$\begin{aligned}\mathring{\mathbf{0}}\overset{\circ}{\mathbf{\Lambda}}\mathring{\mathbf{1}}^{(\text{sym-int})} =& |\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \Big[\tfrac{1}{2}(\hat{\mathbf{U}}_A + \hat{\mathbf{U}}_B)\Big] + |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \Big[\tfrac{1}{2}(\hat{\mathbf{U}}_A + \hat{\mathbf{U}}_B)\Big] \\ &+ |\mathbf{0}\rangle_c\langle\mathbf{1}|_c \otimes \Big[\tfrac{1}{2}(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)\Big] + |\mathbf{1}\rangle_c\langle\mathbf{0}|_c \otimes \Big[\tfrac{1}{2}(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)\Big]\end{aligned}$$

1. *The* Λ_{00} *and* Λ_{11} *Components:* A Synthesis of Histories

   - *Mathematical Form*: $\Lambda_{00} = \Lambda_{11} = \frac{1}{2}(\hat{\mathbf{U}}_A + \hat{\mathbf{U}}_B)$.
   - *Detailed Interpretation*: These components represent the amplitude for a process where the control qubit's state *does not flip* (i.e., a process of type $0 \to 0$ or $1 \to 1$). In the interferometer analogy, this corresponds to a particle exiting through the "same port" it metaphorically entered. For this to happen, the quantum amplitudes of the two internal paths must add together, or interfere *constructively*.

     The operator sum $(\hat{\mathbf{U}}_A + \hat{\mathbf{U}}_B)$ is the precise mathematical expression of this constructive synthesis. It does not select one path or the other; rather, it produces a final state that is a coherent superposition of the outcomes of *both* evolutionary histories. This is why we identify its function with *Parasymmetry*, the dialectical principle of synthesis, now applied to entire dynamical processes.

2. *The* Λ_{01} *and* Λ_{10} *Components:* A Dissonance of Histories

- *Mathematical Form*: $\Lambda_{01} = \Lambda_{10} = \frac{1}{2}(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)$.
- *Detailed Interpretation*: These components represent the amplitude for a process where the control qubit's state *flips* (i.e., a transition from $0 \to 1$ or $1 \to 0$). In the interferometer analogy, this is equivalent to the particle switching paths and exiting through the "opposite port" It can only occur if the quantum amplitudes of the two paths interfere *destructively*, largely cancelling each other out to allow for a transition that is classically forbidden.

  The operator difference $(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)$ mathematically isolates this destructive dissonance. It computes the "difference" between the two universal histories. We therefore identify its function with *Anti-Parasymmetry*, the dialectical principle of dissonance or anti-synthesis.

In summary, the four matrix elements of the Symmetric Interferometer operator provide a complete description of the interference experiment. The diagonal terms quantify the degree of agreement or synthesis between the two histories, while the off-diagonal terms quantify their disagreement or dissonance.

(b) Asymmetric Interferometer

$$\mathbf{U}_{\text{holo}}^{(\text{asym})} = \mathbf{U}_{\text{read}} \cdot \mathbf{U}_{\text{ctrl}} \cdot \mathbf{U}_{\text{prep}}$$

Ontological Expansion:

$$\begin{aligned}\mathring{\mathbf{0}}\mathring{\mathbf{\Lambda}}\mathring{\mathbf{1}}^{\circ\,(\text{asym-int})} =& |\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \left[\gamma\alpha\hat{\mathbf{U}}_A - \delta^*\beta\hat{\mathbf{U}}_B\right] + |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \left[-\delta\beta^*\hat{\mathbf{U}}_A + \gamma^*\alpha^*\hat{\mathbf{U}}_B\right] \\ &+ |\mathbf{0}\rangle_c\langle\mathbf{1}|_c \otimes \left[-\gamma\beta^*\hat{\mathbf{U}}_A - \delta^*\alpha^*\hat{\mathbf{U}}_B\right] + |\mathbf{1}\rangle_c\langle\mathbf{0}|_c \otimes \left[\delta\alpha\hat{\mathbf{U}}_A + \gamma^*\beta\hat{\mathbf{U}}_B\right]\end{aligned}$$

By comparing this to the universal form $\sum_{i,j} |\mathbf{i}\rangle_c\langle\mathbf{j}|_c \otimes \Lambda_{ij}$, we can directly identify the four operator-valued matrix elements.

1. *The Λ_{00} Component:*
$$\Lambda_{00}^{(\text{asym-int})} = \gamma\alpha\hat{\mathbf{U}}_A - \delta^*\beta\hat{\mathbf{U}}_B$$
2. *The Λ_{01} Component:*
$$\Lambda_{01}^{(\text{asym-int})} = -\gamma\beta^*\hat{\mathbf{U}}_A - \delta^*\alpha^*\hat{\mathbf{U}}_B$$
3. *The Λ_{10} Component:*
$$\Lambda_{10}^{(\text{asym-int})} = \delta\alpha\hat{\mathbf{U}}_A + \gamma^*\beta\hat{\mathbf{U}}_B$$
4. *The Λ_{11} Component:*
$$\Lambda_{11}^{(\text{asym-int})} = -\delta\beta^*\hat{\mathbf{U}}_A + \gamma^*\alpha^*\hat{\mathbf{U}}_B$$

1. The Role of Preparation (α, β) The preparation gate acts first, transforming the initial control state into an arbitrary superposition $\alpha|0\rangle_c + \beta|1\rangle_c$.

- α becomes the initial complex amplitude for the history that will evolve down *Path A* (governed by $\hat{\mathbf{U}}_A$).
- β becomes the initial complex amplitude for the history that will evolve down *Path B* (governed by $\hat{\mathbf{U}}_B$).

This is analogous to using a variable beam splitter in an optics experiment; we can control the intensity and relative phase of the two paths being compared.

2. The Role of Readout (γ, δ) The readout gate acts last, projecting the two evolved histories onto a new measurement basis before the final observation. For example, to be measured in the final state $|0\rangle_c$, a state receives a contribution of γ times its $|0\rangle_c$ component and $-\delta^*$ times its $|1\rangle_c$ component.

3. Detailed Interpretation of the Matrix Elements Each matrix element Λ_{ij} represents the total process operator for the $j \to i$ transition, combining the effects of preparation, evolution, and readout.

1. $\Lambda_{00}^{(asym\text{-}int)} = \gamma\alpha\hat{\mathbf{U}}_A - \delta^*\beta\hat{\mathbf{U}}_B$ This is the operator for the $0 \to 0$ process. It is a coherent sum of two amplitudes:

   - The amplitude for the Path A history ($\alpha\hat{\mathbf{U}}_A$) to be projected onto the final $|0\rangle_c$ state by the readout gate (a factor of γ).
   - The amplitude for the Path B history ($\beta\hat{\mathbf{U}}_B$) to be projected onto the final $|0\rangle_c$ state (a factor of $-\delta^*$).

   The final operator is the tunable interference pattern between these two weighted, evolved paths.

2. *The Other Components:* The remaining elements follow the same logic. For example, $\Lambda_{10} = \delta\alpha\hat{\mathbf{U}}_A + \gamma^*\beta\hat{\mathbf{U}}_B$ describes the process for a $0 \to 1$ flip, where the amplitudes from each path are projected onto the final $|1\rangle_c$ state.

In essence, each component is a *Tunable Synthesis/Dissonance Operator*. By choosing the four parameters, an experimenter can precisely control the interference, allowing them to isolate different aspects of the relationship between the two universal histories. This makes the asymmetric model a complete *tomographic tool* for mapping the relational structure of quantum dynamics.

2. The Path Scattering Model Expansion

(a) Symmetric Path Scattering

$$\mathbf{U}_{\text{holo}}^{(\text{scatter,sym})} = \mathbf{U}_{\text{ctrl}} \cdot \mathbf{U}_{\text{Had}} \cdot \mathbf{U}_{\text{ctrl}}$$

Ontological Expansion:

$$\begin{aligned}\mathring{\mathbf{0}}\overset{\circ}{\mathbf{\Lambda}}\mathring{\mathbf{1}}^{(\text{sym-scatter})} =& |\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \left[\tfrac{1}{\sqrt{2}}\hat{\mathbf{U}}_A^2\right] + |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \left[-\tfrac{1}{\sqrt{2}}\hat{\mathbf{U}}_B^2\right] \\ &+ |\mathbf{0}\rangle_c\langle\mathbf{1}|_c \otimes \left[\tfrac{1}{\sqrt{2}}\hat{\mathbf{U}}_A\hat{\mathbf{U}}_B\right] + |\mathbf{1}\rangle_c\langle\mathbf{0}|_c \otimes \left[\tfrac{1}{\sqrt{2}}\hat{\mathbf{U}}_B\hat{\mathbf{U}}_A\right]\end{aligned}$$

Each matrix element Λ_{ij} is a *Transformative History Operator*. These components describes feedback as well as the fundamental change in the rules of evolution. The components are *Iterated and Hybrid History Operators*.

- The diagonal elements $(\Lambda_{00}, \Lambda_{11})$ represent *self-reinforcing feedback loops*, where a history's character is preserved through the interference event ($A \to A$ or $B \to B$).
- The off-diagonal elements $(\Lambda_{01}, \Lambda_{10})$ represent *path-switching feedback*, where the interference event causes a transition between the two types of evolutionary dynamics ($A \to B$ or $B \to A$).

Synthesis: A Universe in Dialogue with Itself Together, these four operators paint a picture of a universe that is in a constant dialogue with its own quantum state. The diagonal elements provide the stability and continuity of physical law, while the off-diagonal elements provide the potential for radical, revolutionary change. The interplay between these four channels dictates the entire cosmic history, from long periods of stable evolution to the singular moments of transformation that define its great epochs.

(b) Asymmetric Path Scattering

$$\mathbf{U}_{\text{holo}}^{(\text{scatter,asym})} = \mathbf{U}_{\text{ctrl}}^{\text{out}} \cdot \mathbf{U}_{\text{Had}} \cdot \mathbf{U}_{\text{ctrl}}^{\text{in}}$$

Ontological Expansion:

$$\begin{aligned}\mathring{\mathbf{0}}\overset{\circ}{\mathbf{\Lambda}}\mathring{\mathbf{1}}^{(\text{asym-scatter})} =& |\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \left[\tfrac{1}{\sqrt{2}}\hat{\mathbf{U}}_A^{\text{out}}\hat{\mathbf{U}}_A^{\text{in}}\right] + |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \left[-\tfrac{1}{\sqrt{2}}\hat{\mathbf{U}}_B^{\text{out}}\hat{\mathbf{U}}_B^{\text{in}}\right] \\ &+ |\mathbf{0}\rangle_c\langle\mathbf{1}|_c \otimes \left[\tfrac{1}{\sqrt{2}}\hat{\mathbf{U}}_A^{\text{out}}\hat{\mathbf{U}}_B^{\text{in}}\right] + |\mathbf{1}\rangle_c\langle\mathbf{0}|_c \otimes \left[\tfrac{1}{\sqrt{2}}\hat{\mathbf{U}}_B^{\text{out}}\hat{\mathbf{U}}_A^{\text{in}}\right]\end{aligned}$$

By comparing the ontological expansion to the universal form $\sum_{i,j} |\mathbf{i}\rangle_c\langle\mathbf{j}|_c \otimes \Lambda_{ij}$, we can identify the explicit operator for each matrix element.

1. *The Λ_{00} Component:*

$$\Lambda_{00}^{(\text{asym-scatter})} = \tfrac{1}{\sqrt{2}}\hat{\mathbf{U}}_A^{\text{out}}\hat{\mathbf{U}}_A^{\text{in}}$$

Interpretation: This represents the amplitude for a history that maintains its fundamental character (Path A $\to$ Path A) while crossing the transformative boundary from "in" rules to "out" rules. It embodies *continuity through transformation*.

2. *The Λ_{11} Component:*
$$\Lambda_{11}^{(\text{asym-scatter})} = -\tfrac{1}{\sqrt{2}}\hat{\mathbf{U}}_B^{\text{out}}\hat{\mathbf{U}}_B^{\text{in}}$$
Interpretation: This is the complementary process for the Path B history. It is the other channel for *continuity through transformation*, where a system's fundamental nature is preserved even as its governing laws are updated.

3. *The Λ_{01} Component:*
$$\Lambda_{01}^{(\text{asym-scatter})} = \tfrac{1}{\sqrt{2}}\hat{\mathbf{U}}_A^{\text{out}}\hat{\mathbf{U}}_B^{\text{in}}$$
Interpretation: This represents a history of *discontinuity and revolution.* A system evolving under the "in" rules for Path B is fundamentally rewritten by the interference event, emerging into a new state governed by the "out" rules for Path A.

4. *The Λ_{10} Component:*
$$\Lambda_{10}^{(\text{asym-scatter})} = \tfrac{1}{\sqrt{2}}\hat{\mathbf{U}}_B^{\text{out}}\hat{\mathbf{U}}_A^{\text{in}}$$
Interpretation: This is the reverse revolutionary path ($A \to B$). These off-diagonal terms provide the mechanism for true novelty, modeling how a system can transition between entirely different dynamical regimes, such as in a cosmological phase transition.

3. The Lensing Model Expansion

(a) Symmetric Interferometer Lensing

$$\mathbf{U}_{\text{holo}}^{(\text{sym-com})} := [\mathbf{U}_{\text{Had}}, \mathbf{U}_{\text{ctrl}}] \cdot \mathbf{U}_{\text{Had}}^{\dagger}$$

Ontological Expansion:

$$\begin{aligned}\mathring{\mathbf{\Lambda}}_{\mathring{\mathbf{i}}}^{\mathring{\mathbf{0}}\,(\text{sym-com})} &= |\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \left[-\tfrac{1}{2}(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)\right] \;+\; |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \left[\tfrac{1}{2}(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)\right] \\ &\quad + |\mathbf{0}\rangle_c\langle\mathbf{1}|_c \otimes \left[\tfrac{1}{2}(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)\right] \;+\; |\mathbf{1}\rangle_c\langle\mathbf{0}|_c \otimes \left[\tfrac{1}{2}(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)\right]\end{aligned}$$

By extracting the operator coefficients from the ontological expansion provided, we identify the four matrix elements:

1. *The Λ_{00} Component:*
$$\Lambda_{00}^{(\text{sym-com})} = -\frac{1}{2}\left(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B\right)$$
This operator governs the amplitude for a process that starts in the control state $|\mathbf{0}\rangle_c$ and is measured to be in the same state, $|\mathbf{0}\rangle_c$, after the full operation. Its form reveals that this *"non-flipping"* channel is driven entirely by the destructive dissonance between the two paths, $\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B$, and includes a characteristic negative phase.

2. *The Λ_{01}, Λ_{10}, and Λ_{11} Components:*

$$\Lambda_{01}^{(\text{sym-com})} = \Lambda_{10}^{(\text{sym-com})} = \Lambda_{11}^{(\text{sym-com})} = +\frac{1}{2}\left(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B\right)$$

Remarkably, the other three components are identical. They represent the amplitudes for the three remaining transitions: a flip from $|\mathbf{1}\rangle_c$ to $|\mathbf{0}\rangle_c$, a flip from $|\mathbf{0}\rangle_c$ to $|\mathbf{1}\rangle_c$, and staying in the $|\mathbf{1}\rangle_c$ state.

The entire dynamic of this lensing model is governed by a single ontological process: the *Anti-Parasymmetry of Processes*, represented by the dissonance operator $(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)$. The model effectively filters out all constructive interference. The resulting dynamics are purely a function of the *difference* between the two evolutionary histories. The asymmetry between the Λ_{00} component (with its minus sign) and the other three identical components is a direct signature of the algebraic twist being projected back onto the measurement basis. It shows how the fundamental non-commutativity of quantum operations manifests as distinct amplitudes for different measurement outcomes.

(b) Asymmetric Interferometer Lensing

$$\mathbf{U}_{\text{holo}}^{(\text{asym-com})} := [\mathbf{U}_{\text{ctrl}}, \mathbf{U}_{\text{prep}}] \cdot \mathbf{U}_{\text{read}}^{\dagger}$$

Ontological Expansion:

$$\begin{aligned}\mathring{\mathbf{0}}\overset{\circ}{\mathbf{\Lambda}}\mathring{\mathbf{1}}^{(\text{asym-com})} &= |\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \left[\delta\beta^*(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)\right] \;+\; |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \left[-\delta^*\beta(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)\right] \\ &\quad + |\mathbf{0}\rangle_c\langle\mathbf{1}|_c \otimes \left[-\gamma\beta^*(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)\right] \;+\; |\mathbf{1}\rangle_c\langle\mathbf{0}|_c \otimes \left[-\gamma^*\beta(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)\right]\end{aligned}$$

By extracting the operator coefficients from the ontological expansion provided, we can identify the four matrix elements. Note that every component is proportional to the dissonance operator, $(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)$.

1. *The Λ_{00} Component:*

$$\Lambda_{00}^{(\text{asym-com})} = \delta\beta^*(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)$$

This operator represents the amplitude for a process that begins in the control state prepared by $\mathbf{U}_{\text{prep}}$ and is subsequently measured in the $|\mathbf{0}\rangle_c$ state of the readout basis defined by $\mathbf{U}_{\text{read}}$, without flipping the effective path. The amplitude is weighted by the coefficients δ and β^*.

2. *The Λ_{11} Component:*

$$\Lambda_{11}^{(\text{asym-com})} = -\delta^*\beta(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)$$

This is the corresponding amplitude for a process that ends in the $|\mathbf{1}\rangle_c$ state of the readout basis, also without flipping the path.

3. *The Λ_{01} and Λ_{10} Components (Path-Flipping):*

$$\Lambda_{01}^{(\text{asym-com})} = -\gamma\beta^*(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)$$

$$\Lambda_{10}^{(\text{asym-com})} = -\gamma^*\beta(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)$$

These operators represent the amplitudes for the control state to be flipped by the process. Their magnitudes and phases are determined by the cross-terms of the preparation and readout gates.

The components of this model are *Generalized Twist Operators*. The structure is highly revealing:

1. *Universal Dissonance*: Every single channel in this process is proportional to the dissonance operator $(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)$. This shows that the algebraic twist isolated by the commutator $[\mathbf{U}_{\text{ctrl}}, \mathbf{U}_{\text{prep}}]$ is purely a function of the *difference* between the evolutionary paths.

2. *Tunable Projections*: The complex coefficients $(\alpha, \beta, \gamma, \delta)$ from the preparation and readout gates act as "dials" that control the magnitude and phase of the dissonance operator that gets projected into each of the four final channels.

This model, therefore, describes how the fundamental algebraic *shear* between evolution and preparation can be rotated and projected into any arbitrary measurement basis, making it the essential tool for a full tomographic reconstruction of the system's non-commutative geometry.

(c) Symmetric Path Scattering Lensing

$$\mathbf{U}_{\text{holo}}^{(\text{psym-com})} := \mathbf{U}_{\text{ctrl}} \cdot [\mathbf{U}_{\text{ctrl}}, \mathbf{U}_{\text{Had}}]$$

Ontological Expansion:

$$\begin{aligned} \mathring{\mathbf{o}}\overset{\circ}{\mathbf{\Lambda}}\mathring{\mathbf{i}}^{(\text{psym-com})} &= |\mathbf{0}\rangle_c\langle\mathbf{1}|_c \otimes \left[\tfrac{1}{\sqrt{2}}\hat{\mathbf{U}}_A(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)\right] \\ &+ |\mathbf{1}\rangle_c\langle\mathbf{0}|_c \otimes \left[-\tfrac{1}{\sqrt{2}}\hat{\mathbf{U}}_B(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)\right] \end{aligned}$$

By comparing it to the universal form $\sum_{i,j} |\mathbf{i}\rangle_c\langle\mathbf{j}|_c \otimes \Lambda_{ij}$, we can identify the four operator-valued matrix elements.

1. *The Λ_{00} and Λ_{11} Components (Diagonal):*

$$\Lambda_{00}^{(\text{psym-com})} = 0$$

$$\Lambda_{11}^{(\text{psym-com})} = 0$$

The most striking feature of this model is that its diagonal components are zero. This means that a process governed by this specific operator *always flips the control state*. There is no amplitude for a history to start in a given control state ($|\mathbf{0}\rangle_c$ or $|\mathbf{1}\rangle_c$) and remain in that same state after the full operation.

2. *The Λ_{01} Component (Path-Flipping $1 \to 0$):*

$$\Lambda_{01}^{(\text{psym-com})} = \tfrac{1}{\sqrt{2}}\hat{\mathbf{U}}_A(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)$$

This operator governs the transition from the control state $|\mathbf{1}\rangle_c$ to $|\mathbf{0}\rangle_c$. It describes a process where the initial algebraic twist—represented by the dissonance term $(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)$—is subsequently acted upon, or "filtered," by the *Path A* evolution operator, $\hat{\mathbf{U}}_A$.

3. *The Λ_{10} Component (Path-Flipping $0 \to 1$):*

$$\Lambda_{10}^{(\text{psym-com})} = -\tfrac{1}{\sqrt{2}}\hat{\mathbf{U}}_B(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)$$

This is the complementary process. It is the amplitude for flipping from $|\mathbf{0}\rangle_c$ to $|\mathbf{1}\rangle_c$. Here, the same initial twist is filtered by the *Path B* evolution operator, $\hat{\mathbf{U}}_B$.

The components of this model are *Feedback-Filtered Twist Operators*. They do not simply measure the algebraic source of interference (the commutator); they describe how that source term is itself transformed by a subsequent evolution. This represents a true dynamical feedback loop, where the structure of quantum interference is fed back into the system's dynamics, illustrating how the past evolution (embedded in the commutator) conditions the future evolution.

(d) Asymmetric Path Scattering Lensing

$$\mathbf{U}_{\text{holo}}^{(\text{pasym-com})} := \mathbf{U}_{\text{ctrl}}^{\text{out}} \cdot [\mathbf{U}_{\text{Had}}, \mathbf{U}_{\text{ctrl}}^{\text{in}}]$$

Ontological Expansion:

$$\begin{aligned} \mathring{\mathbf{0}}\overset{\circ}{\mathbf{\Lambda}}\mathring{\mathbf{1}}^{(\text{pasym-com})} &= |\mathbf{1}\rangle_c\langle\mathbf{0}|_c \otimes \left[\tfrac{1}{\sqrt{2}}\hat{\mathbf{U}}_B^{\text{out}}(\hat{\mathbf{U}}_A^{\text{in}} - \hat{\mathbf{U}}_B^{\text{in}})\right] \\ &\quad + |\mathbf{0}\rangle_c\langle\mathbf{1}|_c \otimes \left[-\tfrac{1}{\sqrt{2}}\hat{\mathbf{U}}_A^{\text{out}}(\hat{\mathbf{U}}_A^{\text{in}} - \hat{\mathbf{U}}_B^{\text{in}})\right] \end{aligned}$$

Extracting the operator coefficients for each matrix element yields the following:

1. *The Λ_{00} and Λ_{11} Components (Diagonal):*

$$\Lambda_{00}^{(\text{pasym-com})} = 0$$
$$\Lambda_{11}^{(\text{pasym-com})} = 0$$

As in the symmetric case, the diagonal components vanish. This reveals that the fundamental operation described by this model is a *pure path-flipping transformation*. Any history that undergoes this specific process is guaranteed to end in a control state orthogonal to its initial state.

2. *The Λ_{01} Component (Path-Flipping $1 \to 0$):*

$$\Lambda_{01}^{(\text{pasym-com})} = -\tfrac{1}{\sqrt{2}}\hat{\mathbf{U}}_A^{\text{out}}(\hat{\mathbf{U}}_A^{\text{in}} - \hat{\mathbf{U}}_B^{\text{in}})$$

This operator describes the amplitude for a history that begins on *Path B* (under the initial "in" laws), undergoes the algebraic twist of interference, and is then transformed by the final "out" laws corresponding to *Path A*.

3. *The Λ_{10} Component (Path-Flipping $0 \to 1$):*

$$\Lambda_{10}^{(\text{pasym-com})} = \frac{1}{\sqrt{2}}\hat{\mathbf{U}}_B^{\text{out}}(\hat{\mathbf{U}}_A^{\text{in}} - \hat{\mathbf{U}}_B^{\text{in}})$$

This is the complementary process, describing a history that begins on *Path A* ("in" laws), twists, and is subsequently transformed by the final "out" laws for *Path B*.

The components of this model are *Transformative Twist Operators*. They represent the most sophisticated process in the framework, providing a mechanism for an irreversible change in dynamics. The core of the process is the algebraic dissonance between the initial paths, $(\hat{\mathbf{U}}_A^{\text{in}} - \hat{\mathbf{U}}_B^{\text{in}})$. This "twist," generated under one set of physical laws, is then filtered through a new, potentially different set of laws ($\hat{\mathbf{U}}_A^{\text{out}}$ or $\hat{\mathbf{U}}_B^{\text{out}}$). This allows the framework to model how the very structure of quantum interference evolves across a fundamental change in the universe's dynamical regime, such as a cosmological phase transition.

The Ontological Status of Physical Reality The framework posits that the observable universe is not a singular, pre-determined entity, but is instead a dynamically selected configuration emerging from a coherent superposition of all possible evolutionary histories. The principle of *Holosymmetry* provides the mathematical and ontological engine for this emergence.

Reality as Constructive Interference: The Principle of Parasymmetric Selection The stable structures of reality—including particles, forces, and the very laws of physics—are identified with the *constructive interference patterns* that arise from the universal superposition of histories. In the language of the Symmetric Interferometer Model, these correspond to the diagonal process operators, which are proportional to the synthetic sum of path operators, $(\hat{\mathbf{U}}_A + \hat{\mathbf{U}}_B)$.

These are the channels where divergent evolutionary pathways align and reinforce one another, leading to a high-amplitude, stable, and persistent outcome. This process is the dynamical realization of *Parasymmetry (Synthesis)* at the level of universal evolution. The observable cosmos is, in this sense, the "bright fringe" of its own self-interference.

The Void as Sculptor: Anti-Parasymmetry and the Annihilation of Possibilities A stable reality can only be defined in contrast to the vast space of unrealized possibilities. These correspond to the *destructive interference patterns*, which are governed by the off-diagonal, dissonant process operators proportional to $(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)$.

These channels of *Anti-Parasymmetry (Dissonance)* represent self-annihilating histories where potential realities cancel each other out, leaving behind a structured void. The "dark fringes" are therefore not empty nothingness, but are an active and necessary component of reality; they are the unobservable structure of cancelled possibilities that "sculpts" the observable reality by carving it out from the plenum of potentiality.

Higher-Order Coherence and the Reflexive Universe The framework's structure allows for a hierarchy of self-interaction, giving rise to two profound concepts:

1. *Second-Order Coherence*: The framework allows for the interference of interference patterns themselves. This is formally realized by models such as the *Path Scattering Operator*, $\mathbf{U}_{\text{ctrl}} \cdot \mathbf{U}_{\text{Had}} \cdot \mathbf{U}_{\text{ctrl}}$, which describes how the outcome of one interferometric process becomes the input for a subsequent evolution. This provides a mechanism for probing the algebraic structure of quantum dynamics at a recursive level.

2. *A History-Dependent, "Learning" Universe*: The *Asymmetric Path Scattering Model* introduces irreversible evolution by allowing the "in" and "out" laws to differ ($\mathbf{U}^{\text{out}}_{\text{ctrl}} \cdot \mathbf{U}_{\text{Had}} \cdot \mathbf{U}^{\text{in}}_{\text{ctrl}}$). This formalizes a universe that is *reflexive*—one whose future dynamical laws can be conditioned by the quantum interference patterns of its own past.

Conclusion: The Cosmos as a Self-Actualizing System Ultimately, Holosymmetry defines the cosmos as a self-contained, self-actualizing system. It is:

1. *Self-Generating*, as it creates its own geometric arena from the Quantum Holo-State via the GNS construction.

2. *Self-Observing*, as it perpetually probes the relational structure between its own possible histories via interferometric dynamics.

3. *Self-Learning*, as it can irreversibly update its own dynamical laws based on the outcome of its self-observation.

The emergence of our reality is thus the result of a grand, recursive synthesis of all possible universal histories, governed by the elegant and complete logic of the Fourfold Dialectic Algebra.

3.8.2 Probing Second-Order Coherence

While the standard interferometric model reveals the first-order interference between two evolutionary histories, its true recursive nature is only unveiled through more sophisticated experimental protocols. We now devise two such "second-order" experiments, each designed to probe the structure of the *Symmetric Interferometer Holosymmetric Operator*, $\mathbf{U}^{(\text{sym})}_{\text{holo}}$, in a unique way.

1. The Sequential Interferometer: Interference of Outcomes This experiment probes the recursive nature of holosymmetry by running the output of one interferometer through a second, identical one. It directly measures how interference patterns themselves interfere.

The Operator: The process is governed by the square of the symmetric operator:

$$\left(\mathbf{U}^{(\text{sym})}_{\text{holo}}\right)^2 = \left(\frac{1}{2}\begin{pmatrix} \hat{\mathbf{U}}_+ & \hat{\mathbf{U}}_- \\ \hat{\mathbf{U}}_- & \hat{\mathbf{U}}_+ \end{pmatrix}\right)^2 = \frac{1}{4}\begin{pmatrix} \hat{\mathbf{U}}_+^2 + \hat{\mathbf{U}}_-^2 & \{\hat{\mathbf{U}}_+, \hat{\mathbf{U}}_-\} \\ \{\hat{\mathbf{U}}_+, \hat{\mathbf{U}}_-\} & \hat{\mathbf{U}}_-^2 + \hat{\mathbf{U}}_+^2 \end{pmatrix}$$

where $\{\hat{\mathbf{U}}_+, \hat{\mathbf{U}}_-\} = \hat{\mathbf{U}}_+\hat{\mathbf{U}}_- + \hat{\mathbf{U}}_-\hat{\mathbf{U}}_+$ is the anticommutator.

The Process: We apply this squared operator to the simple initial state $|\Psi_{\text{init}}\rangle = |\mathbf{0}\rangle_c \otimes |\Psi_{\text{RR}}\rangle$. The final state is determined by the first column of the resulting operator matrix:

$$|\Psi_{\text{final}}\rangle = \frac{1}{4}\left[|\mathbf{0}\rangle_c \otimes (\hat{\mathbf{U}}_+^2 + \hat{\mathbf{U}}_-^2)|\Psi_{\text{RR}}\rangle + |\mathbf{1}\rangle_c \otimes \{\hat{\mathbf{U}}_+, \hat{\mathbf{U}}_-\}|\Psi_{\text{RR}}\rangle\right]$$

Interpretation: The final measurement probabilities now depend on higher-order combinations of the path operators.

- p_0 depends on $\hat{\mathbf{U}}_+^2 + \hat{\mathbf{U}}_-^2$. This term represents the amplitude for the system to return to the $|\mathbf{0}\rangle_c$ branch after two full interference events.
- p_1 depends on the anticommutator $\{\hat{\mathbf{U}}_+, \hat{\mathbf{U}}_-\}$. This term represents the amplitude for the system to be "flipped" to the $|\mathbf{1}\rangle_c$ branch by the sequence of interferences.

This experiment directly measures the *interference of interference patterns*, revealing the algebraic structure $(\hat{\mathbf{U}}_+^2, \hat{\mathbf{U}}_-^2, \{\hat{\mathbf{U}}_+, \hat{\mathbf{U}}_-\})$ that governs the system's recursive evolution.

2. The Tomographic Interferometer: Probing the Operator Structure This experiment probes the operator's structure by preparing a general "polarized" input state and observing how the operator transforms it. It is a method of quantum process tomography.

The Process: We apply the standard operator $\mathbf{U}_{\text{holo}}^{(\text{sym})}$ once, but to a general initial state:

$$|\Psi_{\text{init}}\rangle = (\alpha|\mathbf{0}\rangle_c + \beta|\mathbf{1}\rangle_c) \otimes |\Psi_{\text{RR}}\rangle$$

We have already derived the final state for this process:

$$|\Psi_{\text{final}}\rangle = \frac{1}{2}\left[|\mathbf{0}\rangle_c \otimes \left(\alpha\hat{\mathbf{U}}_+ + \beta\hat{\mathbf{U}}_-\right)|\Psi_{\text{RR}}\rangle + |\mathbf{1}\rangle_c \otimes \left(\alpha\hat{\mathbf{U}}_- + \beta\hat{\mathbf{U}}_+\right)|\Psi_{\text{RR}}\rangle\right]$$

Interpretation: The final amplitudes are now complex mixtures of the initial amplitudes α and β. The mixing is governed by the off-diagonal terms of the operator matrix, $\frac{1}{2}\hat{\mathbf{U}}_-$. This directly reveals the operator's *"scattering"* nature.

By preparing various input states (i.e., by varying α and β) and measuring the corresponding output states, one can fully reconstruct the matrix elements of $\mathbf{U}_{\text{holo}}^{(\text{sym})}$. This tomographic approach allows one to measure the operator's structure itself, providing a complete characterization of the interferometric process.

Holosymmetry: The Universe as a Quantum Interferometer At the heart of the Interdependent Quantum Reality (IQR) framework lies a radical proposition: that reality is not described by a single, unfolding history, but by a coherent superposition of all possible evolutionary histories. The principles of *Parasymmetry* and *Holosymmetry* provide the architecture for this vision.

- *Parasymmetry* is the language of interaction. It governs how the fundamental modes of Symmetry and Antisymmetry combine and entangle across the registers of Primordial Quantum Singularities, creating the rich tapestry of quantum states under a *given* set of dynamical rules.
- *Holosymmetry* is the meta-language of evolution. It does not operate on states, but on the *very rules of Parasymmetry themselves*. It provides the mechanism for placing different evolutionary laws—different universal histories—into a grand, coherent superposition.

The models we have just detailed are the experimental protocols for this "metaphysical laboratory".

The Interferometer: A Cosmic Comparison Engine The *Interferometer Model*, defined by the operator $\mathbf{U}_{\text{holo}}^{(\text{sym})} = \mathbf{U}_{\text{Had}} \cdot \mathbf{U}_{\text{ctrl}} \cdot \mathbf{U}_{\text{Had}}$, acts as a cosmic comparison engine. Its purpose is to answer a profound question: "Given two possible sets of physical laws (Path A and Path B), how different are they?"

By preparing a superposition, evolving each branch under a different dynamic, and recombining them, the interferometer produces a final measurement on the control qubit. This measurement is a direct readout of the *relational distinguishability* between two entire universal histories. The interference pattern is not of particles in space, but of realities in the space of possibilities.

The Path Scattering Model: A Self-Referential Universe The *Path Scattering Model*, $\mathbf{U}_{\text{holo}}^{(\text{scatter,sym})} = \mathbf{U}_{\text{ctrl}} \cdot \mathbf{U}_{\text{Had}} \cdot \mathbf{U}_{\text{ctrl}}$, describes a more complex and intimate process: a universe capable of *dynamical feedback*.

This is a model for a self-referential reality. It describes a process where the universe evolves along one path, is subjected to an interference event that makes it "aware" of its other possibilities, and then alters its subsequent evolution based on the outcome of that interference. It is a framework for a learning universe, where the laws of tomorrow may depend on the quantum interference of all its yesterdays.

The Nature of Reality and Physical Law This leads us to a final, profound reinterpretation of existence itself. The universe, in this view, is a single, vast, self-interfering quantum system. The "reality" we observe, with its apparently stable and unique physical laws, is merely the "bright fringe" in this cosmic interference pattern—the pathway of maximal constructive interference among an infinity of competing evolutionary possibilities. Physical laws are not immutable, Platonic edicts imposed from without. They are *emergent and state-dependent regularities*. They are the stable, self-reinforcing patterns that have won out in the universe's continuous process of self-observation and self-creation. The very fabric of spacetime, the particles within it, and the rules they obey are all different facets of a single, indivisible, and recursively self-interfering quantum state.

3.8.3 The Master Holosymmetric Operator

Let $\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}$ be bounded operators on the dual-register Hilbert space $\mathcal{H}_{\mathrm{RR}}^{(2n)}$. Relative to the control computational basis $\{|\mathbf{0}\rangle_c, |\mathbf{1}\rangle_c\}$, define the *Master Holosymmetric Operator* (general form)

$$\mathbf{U}_{\mathrm{holo}}^{(\mathrm{gen})} := \overset{\circ}{\underset{\mathring{0}}{\mathbf{\Lambda}}}{}_{\mathring{1}}^{(\mathrm{General})} = |\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \mathbf{A} + |\mathbf{0}\rangle_c\langle\mathbf{1}|_c \otimes \mathbf{B} + |\mathbf{1}\rangle_c\langle\mathbf{0}|_c \otimes \mathbf{C} + |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \mathbf{D}.$$

This gives a complete 2×2 control–register *parametrization* of holosymmetric operations; the physically admissible (unitary) protocols are those choices of $(\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D})$ that satisfy the block–unitarity constraints below.

Remark 3.19 (Block unitarity of the master operator)**.** Writing $\mathbf{U}_{\mathrm{holo}}^{(\mathrm{gen})}$ as the 2×2 block $\left(\begin{smallmatrix} \mathbf{A} & \mathbf{B} \\ \mathbf{C} & \mathbf{D} \end{smallmatrix}\right)$ on $\mathcal{H}_c \otimes \mathcal{H}_{\mathrm{RR}}^{(2n)}$, unitarity $\left(\mathbf{U}_{\mathrm{holo}}^{(\mathrm{gen})}\right)^\dagger \mathbf{U}_{\mathrm{holo}}^{(\mathrm{gen})} = \mathbb{I} = \mathbf{U}_{\mathrm{holo}}^{(\mathrm{gen})} \left(\mathbf{U}_{\mathrm{holo}}^{(\mathrm{gen})}\right)^\dagger$ is equivalent to

$$\mathbf{A}^\dagger\mathbf{A} + \mathbf{C}^\dagger\mathbf{C} = \mathbb{I}, \quad \mathbf{B}^\dagger\mathbf{B} + \mathbf{D}^\dagger\mathbf{D} = \mathbb{I}, \quad \mathbf{A}^\dagger\mathbf{B} + \mathbf{C}^\dagger\mathbf{D} = 0,$$

and

$$\mathbf{A}\mathbf{A}^\dagger + \mathbf{B}\mathbf{B}^\dagger = \mathbb{I}, \quad \mathbf{C}\mathbf{C}^\dagger + \mathbf{D}\mathbf{D}^\dagger = \mathbb{I}, \quad \mathbf{A}\mathbf{C}^\dagger + \mathbf{B}\mathbf{D}^\dagger = 0.$$

These identities hold automatically in the canonical models obtained from controlled unitaries and Hadamard pre/post-rotations.

Remark 3.20 (Master valuation)**.** For control $|\mathbf{a}\rangle_c \to |\mathbf{b}\rangle_c$, the holosymmetric effect on $\mathcal{H}_{\mathrm{RR}}^{(2n)}$ is

$$E_{\mathrm{Holo}}^{(\mathbf{a}\to\mathbf{b})} := \langle\mathbf{a}|_c\, \mathbf{U}_{\mathrm{holo}}^\dagger \Big(|\mathbf{b}\rangle\langle\mathbf{b}|_c \otimes \mathbb{I}\Big)\, \mathbf{U}_{\mathrm{holo}}\, |\mathbf{a}\rangle_c,$$

so the Holothesis valuation is $v_{\mathrm{Holothesis}}(\rho) = \mathrm{Tr}\Big(\rho\, E_{\mathrm{Holo}}^{(\mathbf{a}\to\mathbf{b})}\Big)$ (cf. Definition 1.18).

Corollary 3.21 (Basis specialization: computational control)**.** *If* $|\mathbf{a}\rangle_c, |\mathbf{b}\rangle_c \in \{|\mathbf{0}\rangle_c, |\mathbf{1}\rangle_c\}$ *and* $\mathbf{U}_{holo} = \left(\begin{smallmatrix} \mathbf{A} & \mathbf{B} \\ \mathbf{C} & \mathbf{D} \end{smallmatrix}\right)$*, then*

$$E^{(0\to0)} = \mathbf{A}^\dagger\mathbf{A}, \quad E^{(0\to1)} = \mathbf{C}^\dagger\mathbf{C}, \quad E^{(1\to0)} = \mathbf{B}^\dagger\mathbf{B}, \quad E^{(1\to1)} = \mathbf{D}^\dagger\mathbf{D}.$$

For arbitrary control bases realized by $\mathbf{U}_{\mathrm{prep}}, \mathbf{U}_{\mathrm{read}} \in \mathcal{U}(\mathcal{H}_c)$*, use Remark 3.20 with* $|\mathbf{a}\rangle_c = \mathbf{U}_{\mathrm{prep}}|\mathbf{a}_0\rangle_c$ *and* $|\mathbf{b}\rangle_c = \mathbf{U}_{\mathrm{read}}|\mathbf{b}_0\rangle_c$*.*

Recovery of Canonical Models

Each specific physical model emerges by imposing particular constraints on the four process operators:

1. The Symmetric Interferometer Model

Definition: $\mathbf{U}_{\mathrm{holo}}^{(\mathrm{sym})} = \mathbf{U}_{\mathrm{Had}} \cdot \mathbf{U}_{\mathrm{ctrl}} \cdot \mathbf{U}_{\mathrm{Had}}$

Process Operators:

$$\mathbf{A} = \mathbf{D} = \frac{1}{2}(\hat{\mathbf{U}}_A + \hat{\mathbf{U}}_B)(Parasymmetricsynthesis)$$
$$\mathbf{B} = \mathbf{C} = \frac{1}{2}(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)(Anti - parasymmetricdissonance)$$

Physical Interpretation: The interferometer realizes perfect symmetry between the diagonal and off-diagonal channels, encoding constructive and destructive interference between evolutionary paths.

2. The Symmetric Path Scattering Lensing Model

Definition: $\mathbf{U}_{\text{holo}}^{(\text{lens,sym})} = \mathbf{U}_{\text{ctrl}} \cdot [\mathbf{U}_{\text{ctrl}}, \mathbf{U}_{\text{Had}}]$

Process Operators:

$$\mathbf{A} = \mathbf{D} = 0$$
$$\mathbf{B} = \frac{1}{\sqrt{2}}\,\hat{\mathbf{U}}_A(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)$$
$$\mathbf{C} = -\frac{1}{\sqrt{2}}\,\hat{\mathbf{U}}_B(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)$$

Physical Interpretation: This purely off-diagonal structure isolates the algebraic non-commutativity between paths, always inducing control-state flips.

3. The Asymmetric Interferometer Model

Definition: $\mathbf{U}_{\text{holo}}^{(\text{asym-int})} = \mathbf{U}_{\text{read}} \cdot \mathbf{U}_{\text{ctrl}} \cdot \mathbf{U}_{\text{prep}}$

Process Operators:

$$\mathbf{A} = \gamma\alpha\hat{\mathbf{U}}_A - \delta^*\beta\hat{\mathbf{U}}_B$$
$$\mathbf{B} = -\gamma\beta^*\hat{\mathbf{U}}_A - \delta^*\alpha^*\hat{\mathbf{U}}_B$$
$$\mathbf{C} = \delta\alpha\hat{\mathbf{U}}_A + \gamma^*\beta\hat{\mathbf{U}}_B$$
$$\mathbf{D} = -\delta\beta^*\hat{\mathbf{U}}_A + \gamma^*\alpha^*\hat{\mathbf{U}}_B$$

Physical Interpretation: The asymmetric parameters (α, β) and (γ, δ) enable full tomographic control, allowing arbitrary preparation and measurement bases.

4. The Asymmetric Path Scattering Model

Definition: $\mathbf{U}_{\text{holo}}^{(\text{scatter,asym})} = \mathbf{U}_{\text{ctrl}}^{\text{out}} \cdot \mathbf{U}_{\text{Had}} \cdot \mathbf{U}_{\text{ctrl}}^{\text{in}}$

Process Operators:

$$\mathbf{A} = \tfrac{1}{\sqrt{2}}\,\hat{\mathbf{U}}_A^{\text{out}}\hat{\mathbf{U}}_A^{\text{in}}$$
$$\mathbf{B} = \tfrac{1}{\sqrt{2}}\,\hat{\mathbf{U}}_A^{\text{out}}\hat{\mathbf{U}}_B^{\text{in}}$$
$$\mathbf{C} = \tfrac{1}{\sqrt{2}}\,\hat{\mathbf{U}}_B^{\text{out}}\hat{\mathbf{U}}_A^{\text{in}}$$
$$\mathbf{D} = -\tfrac{1}{\sqrt{2}}\,\hat{\mathbf{U}}_B^{\text{out}}\hat{\mathbf{U}}_B^{\text{in}}$$

Physical Interpretation: This model captures irreversible transformations where "in" laws evolve to "out" laws, modeling phase transitions or dynamical feedback.

5. The Asymmetric Interferometer Lensing Model

Definition: $\mathbf{U}_{\text{holo}}^{(\text{lens,asym-int})} = [\mathbf{U}_{\text{ctrl}}, \mathbf{U}_{\text{prep}}] \cdot \mathbf{U}_{\text{read}}^{\dagger}$

Process Operators:

$$\begin{aligned}\mathbf{A} &= \delta\beta^*(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)\\ \mathbf{B} &= -\gamma\beta^*(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)\\ \mathbf{C} &= -\gamma^*\beta(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)\\ \mathbf{D} &= -\delta^*\beta(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)\end{aligned}$$

Physical Interpretation: Every channel is proportional to the dissonance operator, isolating the algebraic interference generated by arbitrary preparation.

6. The Asymmetric Path Scattering Lensing Model

Definition: $\mathbf{U}_{\text{holo}}^{(\text{lens,asym-scatter})} = \mathbf{U}_{\text{ctrl}}^{\text{out}} \cdot [\mathbf{U}_{\text{Had}}, \mathbf{U}_{\text{ctrl}}^{\text{in}}]$

Process Operators:

$$\begin{aligned}\mathbf{A} &= 0\\ \mathbf{B} &= -\tfrac{1}{\sqrt{2}}\,\hat{\mathbf{U}}_A^{\text{out}}\left(\hat{\mathbf{U}}_A^{\text{in}} - \hat{\mathbf{U}}_B^{\text{in}}\right)\\ \mathbf{C} &= \tfrac{1}{\sqrt{2}}\,\hat{\mathbf{U}}_B^{\text{out}}\left(\hat{\mathbf{U}}_A^{\text{in}} - \hat{\mathbf{U}}_B^{\text{in}}\right)\\ \mathbf{D} &= 0\end{aligned}$$

Physical Interpretation: The purely off-diagonal form ensures control-state flipping, capturing transformative dynamics through algebraic twists.

Profound Insights from the Master Operator

1. Universal Control Architecture All holosymmetric dynamics—whether representing comparison, feedback, or algebraic analysis—share a single universal architecture: the entanglement of a two-level control system with a larger register system. The four matrix elements govern the complete logic of control-state transitions.

2. Complete Phase Space of Quantum Dynamics The four process operators $(\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D})$ serve as coordinates in an abstract space encompassing all possible holosymmetric dynamics. Each point represents a unique evolutionary law, with canonical models occupying special symmetric locations.

3. Possibility versus Actuality The Master Operator represents the entire space of mathematical possibility so that specific models represent physical actualities derived through symmetry-imposing constraints. Physical laws emerge from mathematical possibility through the application of specific operational sequences.

4. Generative Framework for New Physics The Master Operator is both descriptive and generative. By exploring novel choices for the process operators—beyond the canonical models—one can theoretically predict new quantum processes and interactions. It provides a complete roadmap for systematic exploration of quantum dynamics.

A Common Architecture for Holosymmetry

The Interferometer, Path–Scattering, and Lensing presentations at first read like different experiments designed for different questions. But each is, in fact, a particular *realization* of the same backbone: a control qubit coherently addressing two global histories on the dual PQS registers and then recombining them. What follows is not a "unification" in the sense of a single fundamental physical theory; it is a *master parametrization* that gathers all such realizations into one mathematical form while preserving their operational diversity.

Concretely, every holosymmetric protocol can be expanded in the control basis as

$$\mathring{\mathbf{0}}\mathring{\mathbf{\Lambda}}\mathring{\mathbf{1}} \;=\; \sum_{i,j\in\{0,1\}} |\mathbf{i}\rangle_c\langle\mathbf{j}|_c \otimes \Lambda_{ij},$$

and by allowing the four block operators Λ_{ij} to be arbitrary (subject to unitarity when the whole is a physical process), we arrive at the *Master Holosymmetric Operator* with blocks $(\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D})$. Individual models then appear by imposing principled constraints on these blocks—e.g. symmetry conditions, commutator filters, or structured products—rather than by summing "partial operators." This is why "adding models together" is conceptually wrong (like adding a reflection to a rotation): each model is a *complete* protocol optimized for a specific probe of relational dynamics.

What the blocks mean. In the computational control basis, the four process operators carry a transparent operational meaning:

1. $\mathbf{A}$: amplitude map for staying in $|\mathbf{0}\rangle_c$ ($0\to0$ channel),
2. $\mathbf{B}$: map for flipping $|\mathbf{1}\rangle_c\to|\mathbf{0}\rangle_c$ ($1\to0$),
3. $\mathbf{C}$: map for flipping $|\mathbf{0}\rangle_c\to|\mathbf{1}\rangle_c$ ($0\to1$),
4. $\mathbf{D}$: amplitude map for staying in $|\mathbf{1}\rangle_c$ ($1\to1$).

Different holosymmetric "presentations" correspond to different *relations* among $(\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D})$:

- **Interferometer:** blocks built from the *sum/difference* of path histories, e.g. $\mathbf{A} = \mathbf{D} \propto (\hat{\mathbf{U}}_A + \hat{\mathbf{U}}_B)$ and $\mathbf{B} = \mathbf{C} \propto (\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)$ (up to chosen control rotations).
- **Path–Scattering:** blocks as *iterated/hybrid* histories, e.g. $\mathbf{A} \propto \hat{\mathbf{U}}_A^2$, $\mathbf{D} \propto \hat{\mathbf{U}}_B^2$, $\mathbf{B} \propto \hat{\mathbf{U}}_B\hat{\mathbf{U}}_A$, $\mathbf{C} \propto \hat{\mathbf{U}}_A\hat{\mathbf{U}}_B$.
- **Lensing (commutator forms):** blocks proportional to the *difference* $\Delta\mathbf{U} := \hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B$, obtained by control pre/post rotations that cancel the sum and isolate the algebraic "twist."

These are not ad hoc recipes; they are slice choices of the same 2×2 block architecture.

What this buys us. The master parametrization does four practical things: (i) it *classifies* every holosymmetric protocol by its block structure; (ii) it *derives* familiar models as special constraints; (iii) it *generates* new probes by proposing admissible block relations; and (iv) it *clarifies* how mathematical structure (sums, products, commutators of histories) maps to physical readouts. Nothing here asserts a unification of fundamental forces or a theory of everything; rather, it provides a single architectural language in which the diverse holosymmetric experiments are different, symmetry-related "dialects."

Reading the master blocks: symmetry ⇒ protocol. The physical character of a holosymmetric operation is fixed by the *relations* among its four block operators $(\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D})$ (subject to the block–unitarity constraints). What makes the framework powerful is that the canonical presentations appear as *simple symmetry slices* of the same 2×2 architecture:

1. **Symmetric interferometer.** Impose maximal control symmetry: $\mathbf{A} = \mathbf{D}$ (diagonal symmetry) and $\mathbf{B} = \mathbf{C}$ (off–diagonal symmetry). With a Hadamard–Control–Hadamard sequence on the control, the blocks take the familiar *sum/difference* form, $\mathbf{A} = \mathbf{D} \propto (\hat{\mathbf{U}}_A + \hat{\mathbf{U}}_B)$ and $\mathbf{B} = \mathbf{C} \propto (\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B)$, so the "same–port'' channels measure constructive synthesis and the "opposite–port'' channels measure dissonance.

2. **Path–scattering lensing (pure flips).** In the complementary sector with $\mathbf{A} = \mathbf{D} = 0$, the operation is *purely off–diagonal*: every run flips the control $(0 \leftrightarrow 1)$. This guarantees that the readout isolates interference–mediated transitions rather than same–port persistence.

3. **Asymmetric families (tomography and irreversibility).** Breaking the equalities $\mathbf{A} = \mathbf{D}$ and/or $\mathbf{B} = \mathbf{C}$ introduces tunable asymmetries. These parameters support full *tomographic control* (asymmetric interferometers via arbitrary prepare/read on the control) and model *history transformation* (asymmetric path–scattering), where the post–interference dynamics differ from the pre–interference ones.

The block–operator landscape. Taken together, the four blocks define a convenient "state space'' for holosymmetric dynamics: each admissible choice of $(\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D})$ (that satisfies block–unitarity) specifies a distinct protocol on $\mathcal{H}_c \otimes \mathcal{H}_{\mathrm{RR}}^{(2n)}$. The canonical models occupy highly symmetric loci with clear operational meaning; the surrounding continuum contains many legitimate but as–yet unexplored processes.

This perspective highlights a familiar pattern in physics: the gap between *what can be written* and *what is realized*. The Master Operator provides the mathematical envelope; *physical* models emerge by imposing constraints—symmetry, causality, resource or calibration choices—that pick out viable points in the block–operator landscape. Thus the master form is not merely retrospective "unification'' of known cases; it is a *generative* tool for proposing new protocols and experiments.

Directions to explore.

- *Asymmetric pure flips:* $\mathbf{A} = \mathbf{D} = 0$ with $\mathbf{B} \neq \mathbf{C}$ to study nonreciprocal path–flipping dynamics.

- *Algebraic constraints:* enforce relations such as $\mathbf{AD} = \mathbf{BC}$ (or their adjoint variants) to probe hidden conservation laws or selection rules.
- *Time–structured blocks:* allow $\mathbf{A}(t), \mathbf{B}(t), \mathbf{C}(t), \mathbf{D}(t)$ to vary across the protocol, modeling staged or feedback–driven holosymmetric evolutions.

Completeness of Holosymmetry: Architecture as Ontology

Having introduced the Master Holosymmetric Operator as a common 2×2 control–register parametrization (Section 3.8.3), we can now ask a precise question: *Is this description complete for the binary holosymmetric architecture?* By "complete," we mean *expressively exhaustive* given the fixed ingredients (a qubit control and two PQS registers) and the equivalence moves of (H4 see Definition 3.18), not a claim about all conceivable physics.

Formal completeness within the binary architecture The holosymmetric Hilbert space is $\mathcal{H}_{\text{Holo}} = \mathcal{H}_c \otimes \mathcal{H}_{\text{RR}}^{(2n)}$ with $\dim \mathcal{H}_c = 2$. Every linear map on $\mathcal{H}_{\text{Holo}}$ therefore admits a 2×2 block form relative to the control basis,

$$\mathbf{U} = \begin{pmatrix} \mathbf{A} & \mathbf{B} \\ \mathbf{C} & \mathbf{D} \end{pmatrix},$$

and the *physically admissible* (unitary) protocols are precisely those blocks that satisfy the block–unitarity constraints (see remark after (3.8.3)). It follows that the Master Holosymmetric Operator is *expressively complete*: any binary-controlled holosymmetric process is a point $(\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D})$ in this block-operator landscape, while specific presentations (Interferometer, Path–Scattering, Lensing) are obtained by imposing principled relations among the blocks (symmetries, commutator filtering, iterated/hybrid compositions) and by controlbasis choices (H4 see Definition 3.18).

Why this architecture: structure mirrors ontology The binary architecture is not an arbitrary convenience; it instantiates the *Axiom/Principle of Duality* (Definition 2.100) at the operational level.

Control qubit as dialectic arbiter A two-dimensional control is the minimal quantum system that can encode the fundamental duality (Thesis/Symmetry $\leftrightarrow$ Antithesis/Antisymmetry). In the computational control basis, $|\mathbf{0}\rangle_c$ and $|\mathbf{1}\rangle_c$ select complementary assignments of *which* global regime runs on *which* register; superpositions coherently address *both* assignments at once and thus enable interference of histories.

Dual registers as dialectic canvas Two disjoint PQS registers are the twin stages on which the dual regimes unfold in parallel. Their exchange symmetry ($A \leftrightarrow B$) and the control's $|\mathbf{0}\rangle_c \leftrightarrow |\mathbf{1}\rangle_c$ swap echo the same ontological duality, producing a self-similar pattern: the dialectic appears in the ontic labels, in the control basis, and in the two-register geometry.

The block–operator landscape and its use The four blocks $(\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D})$ provide a convenient "state space'' for holosymmetric dynamics (subject to block–unitarity). Highly symmetric loci reproduce the canonical models with clear operational meaning; the surrounding continuum hosts admissible but as-yet unexplored processes. Thus the master form does more than collect known cases: it *generates* new probes by proposing lawful relations among the blocks and by choosing control prepare/readout maps. Valuations remain Born-valued via the induced effect $E_{\mathrm{Holo}}^{(\mathbf{a}\to\mathbf{b})}$ (cf. Definition 1.18 and Remark 3.20).

Paths to greater complexity (beyond the binary axiom) Completeness here is *relative* to the binary design. Enlarging the axioms enlarges the operator class:

1. **Higher-dimensional control** ($\mathbb{C}^3$, $\mathbb{C}^k$): a 3×3 (or $k \times k$) block operator representing coherent superpositions of three (or k) global histories; a natural setting for ternary or multiway dialectics.

2. **Additional registers**: more stages for concurrent histories, typically paired with higher-dimensional control to address them coherently.

3. **Hierarchical control**: nested controls (meta-controls) that program how lower-level controls act, modeling layered evolutionary choices or regime changes.

These are principled generalizations, but they lie *beyond* the binary axiom and require new completeness statements.

Why this matters

1. **Exhaustive classification.** Any binary-controlled holosymmetric protocol lies somewhere in the block–operator landscape; the canonical models are symmetric slices.

2. **Principled constraints.** Physical models are not arbitrary; they are constrained points picked out by symmetry, causality, resource, or calibration requirements.

3. **Predictive leverage.** Unexplored regions of $(\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D})$ suggest new, testable protocols (e.g., asymmetric pure-flip dynamics, algebraically constrained blocks, time-structured blocks).

4. **Foundational clarity.** The "binary limitation'' is a feature: it encodes the dialectic of duality in the architecture, ensuring that the mathematics faithfully reflects the ontology.

Completeness as a platform for discovery The Master Holosymmetric Operator is complete *because* it fully embraces the binary constraints and explores all their consequences. It validates the coherence of the duality principle, provides a full classification of binary holosymmetric phenomena, and points cleanly to higher-dimensional generalizations when (and only when) the foundational axioms are extended. In this sense, the framework carries both its own completion and the seeds of its future evolution.

The Generative Axiom: Primacy of Duality At the base of the edifice is a single axiom: the *Principle of Duality* (Definition 2.100). Stated plainly: to assert the existence of any ontological state is simultaneously to co-assert the existence of its complementary anti-state. From this one premise, the *Fourfold Dialectic Algebra of Symmetry* (Definition 1.16) unfolds as a sequence of necessities rather than a list of independent postulates:

1. *Thesis (Symmetry,* $\mathring{\mathbf{0}}$*)* — the initial stance,
2. *Antithesis (Antisymmetry,* $\mathring{\mathbf{1}}$*)* — its compelled counter-statement,
3. *Synthesis (Parasymmetry)* — the rule of coherent composition of the pair,
4. *Holothesis (Holosymmetry)* — the meta-rule that superposes and compares *entire* dynamical histories.

This fourfold is the theory's logical engine: the richness that follows is the systematic elaboration of the tension and interplay between Symmetry and Antisymmetry, made quantitative by Born-valued semantics (Definition 1.18).

Self-Similar Architecture: from Duality to n-Body Configurations Extending the framework to many controls or many registers does not introduce new axioms; it reuses the same duality in a higher-combinatorial setting.

1. *n-qubit control.* Replacing a single control qubit by $(\mathbb{C}^2)^{\otimes n}$ yields 2^n addressable control channels. Each basis label is still a tensor of the binary primitives $|\mathbf{0}\rangle_c, |\mathbf{1}\rangle_c$; the governing logic remains pairwise duality, now composed across qubits.
2. *n-register geometry.* Likewise, an n-register scene distributes dialectical assignments across more loci. One subset may run coupled (synthetic/parasymmetric) dynamics while another runs decoupled (analytic) dynamics; the *types* of assignment still come from the original fourfold.

In this sense the architecture is *recursively self-similar*: the rule that organizes a single PQS also organizes a network of them.

Infinite Complexity via Quantum Realization Each site is not a classical bit nor even a bare qubit; it is a *Primordial Quantum Singularity (PQS)* with a local modal space $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)} \cong \mathbb{C}^4$. Thus even a single locus supports a continuum of states; n loci span an astronomically large space. When we speak of the two "paths" $\hat{\mathbf{U}}_A$ (coupled) and $\hat{\mathbf{U}}_B$ (decoupled), these are *archetypal families*—coarse descriptions of whole sectors of histories—rather than two literal, isolated trajectories. Operationally, they stand in for path-integral classes dominated by coupling vs. decoupling.

Holosymmetry as a Coarse-Graining Instrument The holosymmetric formalism—captured compactly by the *Master Holosymmetric Operator* (the 2×2 control–register block with entries $\mathbf{A}, \mathbf{B}, \mathbf{C}, \mathbf{D}$)—is the tool that makes this complexity tractable. It does not enumerate every microscopic history; it *selects* a small number of macro-classes (e.g., coupled vs. decoupled sectors, or their structured composites), and computes the precise quantum interference *between classes*. Different presentations (interferometer, path-scattering, lensing) correspond to different lawful relations among the blocks and different control bases (cf.Definition 3.18 (H4) equivalences), while all valuations remain Born expectations of the induced effects on the registers (Definition 1.18, Remark 3.20). In short, Holosymmetry does not erase detail; it makes detail *legible* by posing the right binary question—and then reading the interference it induces.

Duality in Cosmic Evolution

The architecture built from Primordial Quantum Singularities (PQSs) and the Fourfold Algebra is *recursively self-similar*: the same logical rule that organizes a single quantum locus also organizes families of histories at cosmological scale. The same pattern scales to the path-integral arena. The total evolution is a coherent sum over histories of the *Gaiamiti* and *Tantrakyma* fields,

$$\mathbf{U}_{\text{total}} = \int \mathcal{D}\hat{\mathbf{\Phi}}\, \mathcal{D}\hat{\mathbf{\Theta}}\; e^{\,i\, S_{\text{total}}[\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}]}\,,$$

a formal integral over an infinite space of possibilities. To render this operationally legible, introduce a *modularity functional* $\mathcal{M}[\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}]$ that quantifies relational separability. A threshold ϵ then defines two archetypal *classes* of histories—the macroscopic echo of the basic duality:

$$\text{Decoupled class (Antithesis):}\quad \hat{\mathbf{U}}_B := \int_{\mathcal{M}>\epsilon} \mathcal{D}\hat{\mathbf{\Phi}}\, \mathcal{D}\hat{\mathbf{\Theta}}\; e^{\,i\, S_{\text{total}}[\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}]}\,,$$

$$\text{Coupled class (Synthesis):}\quad \hat{\mathbf{U}}_A := \int_{\mathcal{M}<\epsilon} \mathcal{D}\hat{\mathbf{\Phi}}\, \mathcal{D}\hat{\mathbf{\Theta}}\; e^{\,i\, S_{\text{total}}[\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}]}\,.$$

These are *archetypal families*, not single trajectories: $\hat{\mathbf{U}}_A$ summarizes histories dominated by holistic/entangled evolution, while $\hat{\mathbf{U}}_B$ summarizes histories dominated by modular/separable evolution. Different calibrations of ϵ amount to different but (H4)-equivalent (Definition 3.18) coarse-grainings of the same duality class.

Fractal self-similarity across scales The pattern repeats without multiplying principles. At the *local* level, Duality generates the binary modes within a single PQS ($\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)} \cong \mathbb{C}^4$). At the *inter-PQS* level, Parasymmetry organizes coherent composition across loci. At the *cosmic* level, Holosymmetry interrogates the two grand, interfering super-classes of histories (*Coupled* vs. *Decoupled*) via a control-addressed comparison on dual registers. The universe, on this reading, is a dialectical computation of itself: what we observe is the synthesized outcome of a fundamental binary tension, made quantitative by effects and their Born valuations (and realized, in the emergent layer, by the GNS representation).

3.8.4 From Duality to the Architecture of the Cosmos

The Primordial Quantum Singularity (PQS) and Fourfold Algebra framework describes a universe that is *recursively self-similar*: a single axiom—the *Principle/Axiom of Duality* (Definition 2.100 and Section 1.2.6)—generates structure at every scale, from a single locus of existence to superposed families of cosmic histories. (This is a claim of *architectural* completeness within the adopted axioms, not a claim of a final unification of all physics.)

1. The Endowment of the Primordial Quantum Singularity The fundamental unit is the *Primordial Quantum Singularity* $\mathfrak{B}_x$ (cf. Section 1.3). It is not a geometric point but a structured, operator-theoretic locus whose specification internalizes the core principles.

1. *Generative logic.* Each PQS carries the full *Fourfold Dialectic Algebra of Symmetry* (Definition 1.16): from the dual pair *Symmetry/Thesis* $\left(\mathring{\mathbf{0}}\right)$ and *Antisymmetry/Antithesis* $\left(\mathring{\mathbf{1}}\right)$ follow their coherent composition *Parasymmetry/Synthesis* and the meta-principle *Holosymmetry/Holothesis* that compares entire processes (cf. Sections 1.2.1 to 1.2.3 and 3.8). Ontological labels are pre-geometric anchors, not ordinary qubit states (cf. Remark 1.22); their truth semantics are Born-valued (Definition 1.18).

2. *Physical substrate.* This logic is realized inside the Algebra of Observables $\mathcal{A}(\mathcal{C}_d)$ (Definition 1.32) by two fundamental operator families: the *Gaiamiti field* $\{\hat{\mathbf{\Phi}}_{\vec{r}}\}$ (sources of geometric potential) and the *Tantrakyma field* $\{\hat{\mathbf{\Theta}}_{\vec{r}}\}$ (sources of matter-like excitations) (cf. Definitions 6.118 and 6.142). Their holosymmetric coupling is succinctly denoted

$$\hat{\Omega}_{\vec{r}} \;:=\; \mathring{\Lambda}_{\hat{\Phi}\hat{\Theta}},$$

emphasizing that geometry and matter appear as inseparable facets of the same PQS-based construction. Locally, the modal state space is $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)} \cong \mathbb{C}^4$, providing the $(0,1)$ sector (Thesis/Antithesis) and its parasymmetric axis as the basic logical qubit with complementary probe.

2. The Nature of a Universal History A *universal history* is a concrete object in the theory: a complete configuration of the *Gaiamiti* and *Tantrakyma* fields over the emergent manifold $\mathcal{X}_q$ (cf. Definitions 6.118 and 6.142). Its dynamics are governed by the *Total Relational Action* $S_{\text{total}}[\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}]$, which assigns to each history a complex phase and thereby determines interference. The total evolution is the coherent sum over this (formal) space of field configurations, expressed by the path integral

$$\mathbf{U}_{\text{total}} = \int \mathcal{D}\hat{\mathbf{\Phi}}\, \mathcal{D}\hat{\mathbf{\Theta}}\; e^{\,i\, S_{\text{total}}[\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}]}\,.$$

In words: $\mathbf{U}_{\text{total}}$ encodes all co-evolving histories of geometry and matter, each weighted by its action, and provides the arena in which holosymmetric comparisons of histories become operational.

3. Holosymmetry and the Duality of Histories *Holosymmetry* is the mechanism that renders this infinity of histories experimentally legible. It partitions the history space into two archetypal *classes*—the highest-level expression of the Principle of Duality—using a *modularity functional* $\mathcal{M}[\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}]$ that quantifies relational separability:

- *Coupled class (Thesis of Synthesis):* the evolutionary operator

$$\hat{\mathbf{U}}_A := \int_{\mathcal{M}<\epsilon} \mathcal{D}\hat{\mathbf{\Phi}}\, \mathcal{D}\hat{\mathbf{\Theta}}\; e^{\,i\, S_{\text{total}}[\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}]}\,,$$

  summing histories that are predominantly holistic/entangled (the dynamical expression of *Parasymmetry/Synthesis*).

- *Decoupled class (Antithesis of Synthesis):* the operator

$$\hat{\mathbf{U}}_B := \int_{\mathcal{M}>\epsilon} \mathcal{D}\hat{\mathbf{\Phi}}\, \mathcal{D}\hat{\mathbf{\Theta}}\; e^{\,i\, S_{\text{total}}[\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}]}\,,$$

  summing histories that are predominantly modular/separable (the complementary dynamical sector).

This yields two *equivalent* but conceptually distinct definitions of the archetypal path operators:

1. **Foundational (path-integral) definition:** constrained sums over families of Gaiamiti/-Tantrakyma histories as above;

2. **Operational (protocol) definition:** compositions of elementary coupled and decoupled evolutions, $\hat{\mathbf{U}}_A, \hat{\mathbf{U}}_B \in \langle U_G^{(\odot)}, U_N^{(\odot)} \rangle$, realized on dual PQS registers and addressed by a control (cf. Section 3.8).

Their equivalence is a cornerstone of internal consistency: the macro-classes defined by the path-integral coarse-graining coincide with those realized by the control-addressed circuitry, and both induce the same holosymmetric effects $E_{\text{Holo}}^{(\mathbf{a}\to\mathbf{b})}$ and Born-valued valuations (cf. Definition 1.18 and Remark 3.20). Thus the same binary dialectic that organizes a single PQS also partitions the infinite set of cosmic histories into two interfering super-classes—*Coupled* vs. *Decoupled*—exhibiting a cosmos that is profoundly relational and recursively self-similar.

A Path Integral Synthesis of Universal Histories

At the most foundational level, a "universal history'' is a complete configuration of the *Gaiamiti field* $\hat{\mathbf{\Phi}}$ and the *Tantrakyma field* $\hat{\mathbf{\Theta}}$ over the emergent manifold $\mathcal{X}_q$, with dynamics governed by the *Total Relational Action* $S_{\text{total}}[\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}]$. The total amplitude for evolution is a path integral over all histories. Holosymmetry coarse–grains this infinite space into two archetypal *families*, using a modularity functional $\mathcal{M}[\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}]$ that quantifies relational separability.

Coupled (Parasymmetric) family $\hat{\mathbf{U}}_A$. Histories that are predominantly holistic/entangled ($\mathcal{M} < \epsilon$):

$$\hat{\mathbf{U}}_A := \int_{\mathcal{M}<\epsilon} \mathcal{D}\hat{\mathbf{\Phi}}\, \mathcal{D}\hat{\mathbf{\Theta}}\; e^{\,i\, S_{\text{total}}[\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}]}. \tag{3.37}$$

Decoupled (Modular) family $\hat{\mathbf{U}}_B$. Histories that are predominantly modular/separable ($\mathcal{M} > \epsilon$):

$$\hat{\mathbf{U}}_B := \int_{\mathcal{M}>\epsilon} \mathcal{D}\hat{\mathbf{\Phi}}\, \mathcal{D}\hat{\mathbf{\Theta}}\; e^{i\, S_{\text{total}}[\hat{\mathbf{\Phi}},\hat{\mathbf{\Theta}}]}. \tag{3.38}$$

Conventions. Overall normalization/phase factors for the path integrals are suppressed; operationally, these are fixed by embedding in the controlled, unitary holosymmetric circuits below.

Operational bridge. The controlled evolution used across the holosymmetric presentations is

$$\mathbf{U}_{\text{ctrl}} = |\mathbf{0}\rangle_c\langle\mathbf{0}| \otimes \hat{\mathbf{U}}_A + |\mathbf{1}\rangle_c\langle\mathbf{1}| \otimes \hat{\mathbf{U}}_B, \tag{3.39}$$

from which the interferometric, path–scattering, and lensing operators are obtained by the prescribed pre/readout rotations on $\mathcal{H}_c$ and fixed-order compositions (see (H4) equivalences and the model definitions).

Robustness of the partition. The split induced by $\mathcal{M}$ is a coarse–graining: small shifts of ϵ or monotone reparametrizations $\mathcal{M} \mapsto f(\mathcal{M})$ leave the induced family of holosymmetric effects (and thus Born valuations) unchanged up to unitary equivalence on the registers (cf. Definition 3.18).

1. The Interferometer Model: Path Integrals in Superposition

The interferometer protocols measure interference between the two families of histories. Their core is the controlled evolution operator $\mathbf{U}_{\text{ctrl}}$, whose path–integral form is

$$\mathbf{U}_{\text{ctrl}} = |\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \left(\int_{\mathcal{M}<\epsilon} \mathcal{D}\hat{\mathbf{\Phi}}\, \mathcal{D}\hat{\mathbf{\Theta}}\; e^{i\, S_{\text{total}}[\hat{\mathbf{\Phi}},\hat{\mathbf{\Theta}}]} \right) \;+\; |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \left(\int_{\mathcal{M}>\epsilon} \mathcal{D}\hat{\mathbf{\Phi}}\, \mathcal{D}\hat{\mathbf{\Theta}}\; e^{i\, S_{\text{total}}[\hat{\mathbf{\Phi}},\hat{\mathbf{\Theta}}]} \right). \tag{3.40}$$

Equivalently, using the archetypal operators $\hat{\mathbf{U}}_A$ and $\hat{\mathbf{U}}_B$ defined in (3.37)–(3.38),

$$\mathbf{U}_{\text{ctrl}} = |\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}_A + |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}_B. \tag{3.41}$$

The full symmetric interferometer is then $\mathbf{U}^{(\text{sym})}_{\text{holo}} = \mathbf{U}_{\text{Had}} \cdot \mathbf{U}_{\text{ctrl}} \cdot \mathbf{U}_{\text{Had}}$, with $\mathbf{U}_{\text{Had}}$ a fixed control rotation on $\mathcal{H}_c$; outcome channels read the constructive/destructive combination of the two families.

2. The Path Scattering Model: Products of Path Integrals

Path–scattering protocols describe dynamical feedback. Their operators are sequential compositions of $\mathbf{U}_{\text{ctrl}}$ separated by a fixed control rotation (e.g. $\mathbf{U}_{\text{Had}}$), producing *operator products* of the archetypal path–integral evolutions.

Symmetric Path Scattering

$$\mathbf{U}_{\text{holo}}^{(\text{scatter,sym})} = \mathbf{U}_{\text{ctrl}} \cdot \mathbf{U}_{\text{Had}} \cdot \mathbf{U}_{\text{ctrl}}.$$

In block form this yields, for the $|\mathbf{0}\rangle \to |\mathbf{0}\rangle$ channel,

$$\Lambda_{00} = \frac{1}{\sqrt{2}}\,\hat{\mathbf{U}}_A^{\,2} = \frac{1}{\sqrt{2}}\Big(\int_{\mathcal{M}<\epsilon} \mathcal{D}\hat{\mathbf{\Phi}}\,\mathcal{D}\hat{\mathbf{\Theta}}\; e^{\,i\,S_{\text{total}}}\Big)\Big(\int_{\mathcal{M}<\epsilon} \mathcal{D}\hat{\mathbf{\Phi}}\,\mathcal{D}\hat{\mathbf{\Theta}}\; e^{\,i\,S_{\text{total}}}\Big), \tag{3.42}$$

i.e. a self-reinforcing coupled history; off-diagonal components such as $\propto \hat{\mathbf{U}}_B\hat{\mathbf{U}}_A$ represent path-switching feedback.

Asymmetric Path Scattering

$$\mathbf{U}_{\text{holo}}^{(\text{scatter,asym})} = \mathbf{U}_{\text{ctrl}}^{\text{out}} \cdot \mathbf{U}_{\text{Had}} \cdot \mathbf{U}_{\text{ctrl}}^{\text{in}},$$

modeling an irreversible change of regime. The process operators are products of *distinct* constrained integrals, e.g.

$$\Lambda_{00}^{(\text{asym})} = \frac{1}{\sqrt{2}}\,\hat{\mathbf{U}}_A^{\text{out}}\,\hat{\mathbf{U}}_A^{\text{in}} = \frac{1}{\sqrt{2}}\Big(\int_{\mathcal{M}_{\text{out}}<\epsilon_{\text{out}}} \cdots\Big)\Big(\int_{\mathcal{M}_{\text{in}}<\epsilon_{\text{in}}} \cdots\Big), \tag{3.43}$$

capturing "continuity through transformation" (diagonal) and "revolutionary" transitions (off-diagonal).

3. The Commutator Lensing Model: An Interference of Protocols

Lensing operators are algebraic *differences between protocols*, each protocol containing path–integral components. For the symmetric interferometer lensing,

$$\mathbf{U}_{\text{holo}}^{(\text{sym-com})} = \Big[\mathbf{U}_{\text{Had}},\, \mathbf{U}_{\text{ctrl}}\Big]\,\mathbf{U}_{\text{Had}}^{\dagger} = \Big(\mathbf{U}_{\text{Had}}\mathbf{U}_{\text{ctrl}} - \mathbf{U}_{\text{ctrl}}\mathbf{U}_{\text{Had}}\Big)\mathbf{U}_{\text{Had}}^{\dagger}. \tag{3.44}$$

Substituting (3.40) (or (3.41)) into both terms makes explicit that the operator compares two meta-histories—"evolve then superpose" vs "superpose then evolve"—and isolates the algebraic twist due to non-commutativity. By (H4) equivalences (Definition 3.18), lensing is a control-basis/readout reparametrization on the same dual-register backbone (no new resources).

The Unity of Logic, Substance, and Locus The *Fourfold Dialectic Algebra* supplies the generative grammar; the *Gaiamiti* and *Tantrakyma* fields give it physical substance in $\mathcal{A}(\mathcal{C}_d)$; and the *PQS* furnishes the indivisible locus where logic and substance are unified, succinctly

$$\hat{\Omega}_{\vec{r}} := \mathring{\Lambda}_{\hat{\Theta}}^{\hat{\Phi}}.$$

Dynamics are encoded by path integrals over these fields; Holosymmetry then partitions the infinity of histories into two interfering families—*Coupled* and *Decoupled*—realizing the foundational duality in experimentally legible form via Born-valued effects.

The Commutator as an Interference of Protocols

Conceptually, the *commutator lensing* operator is not a single uninterrupted evolution; it compares two distinct *protocols* (remembering that the rightmost gate acts first):

1. *Protocol 1 (evolve then superpose):* $\mathbf{U}_{\mathrm{Had}}\mathbf{U}_{\mathrm{ctrl}}$.
2. *Protocol 2 (superpose then evolve):* $\mathbf{U}_{\mathrm{ctrl}}\mathbf{U}_{\mathrm{Had}}$.

Their physical difference is isolated by

$$\mathbf{U}_{\mathrm{holo}}^{(\text{sym-com})} := \Big[\,\mathbf{U}_{\mathrm{Had}},\,\mathbf{U}_{\mathrm{ctrl}}\,\Big]\,\mathbf{U}_{\mathrm{Had}}^{\dagger}. \tag{3.45}$$

Decomposition note. Writing the controlled block as

$$\mathbf{U}_{\mathrm{ctrl}} = \mathbb{I}_c \otimes U_+ + Z_c \otimes U_-, \qquad U_\pm := \tfrac{1}{2}\big(\hat{\mathbf{U}}_A \pm \hat{\mathbf{U}}_B\big), \tag{3.46}$$

any control rotation $R \in \mathcal{U}(\mathcal{H}_c)$ obeys $[R, \mathbf{U}_{\mathrm{ctrl}}] = [R, Z_c] \otimes U_-$. Thus the commutator isolates exactly the *difference* piece: $\Delta\mathbf{U} = \hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B = 2U_-$. By the (H4) equivalences (Definition 3.18), lensing is therefore just a control-basis/readout reparametrization on the same dual-register backbone.

Formal derivation and path-integral representation

1. Components.

1. Non-dynamical Hadamard on the control: $\mathbf{U}_{\mathrm{Had}} = H_c \otimes \mathbb{I}_{\mathrm{RR}}$, with $H_c = \frac{1}{\sqrt{2}}\left(\begin{smallmatrix} 1 & 1 \\ 1 & -1 \end{smallmatrix}\right)$.
2. Controlled evolution (compact and path–integral forms):

$$\mathbf{U}_{\mathrm{ctrl}} = |\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}_A + |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}_B, \tag{3.47}$$

where

$$\hat{\mathbf{U}}_A := \int_{\mathcal{M}<\epsilon} \mathcal{D}\hat{\mathbf{\Phi}}\,\mathcal{D}\hat{\mathbf{\Theta}}\; e^{\,i\,S_{\mathrm{total}}[\hat{\mathbf{\Phi}},\hat{\mathbf{\Theta}}]}, \qquad \hat{\mathbf{U}}_B := \int_{\mathcal{M}>\epsilon} \mathcal{D}\hat{\mathbf{\Phi}}\,\mathcal{D}\hat{\mathbf{\Theta}}\; e^{\,i\,S_{\mathrm{total}}[\hat{\mathbf{\Phi}},\hat{\mathbf{\Theta}}]}. \tag{3.48}$$

2. Calculating the commutator. In the control basis, $\mathbf{U}_{\mathrm{Had}} \leftrightarrow \frac{1}{\sqrt{2}}\left(\begin{smallmatrix} \mathbb{I} & \mathbb{I} \\ \mathbb{I} & -\mathbb{I} \end{smallmatrix}\right)$ and $\mathbf{U}_{\mathrm{ctrl}} \leftrightarrow \left(\begin{smallmatrix} \hat{\mathbf{U}}_A & 0 \\ 0 & \hat{\mathbf{U}}_B \end{smallmatrix}\right)$. A short block-matrix multiplication gives

$$\Big[\,\mathbf{U}_{\mathrm{Had}},\,\mathbf{U}_{\mathrm{ctrl}}\,\Big] = \frac{1}{\sqrt{2}}\begin{pmatrix} 0 & -(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B) \\ \hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B & 0 \end{pmatrix}. \tag{3.49}$$

3. Applying the final gate. Since $\mathbf{U}_{\mathrm{Had}}^{\dagger} = \mathbf{U}_{\mathrm{Had}}$,

$$\mathbf{U}_{\mathrm{holo}}^{(\text{sym-com})} = \frac{1}{2}\begin{pmatrix} -(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B) & (\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B) \\ (\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B) & (\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B) \end{pmatrix}. \tag{3.50}$$

4. Path-integral "dissonance" operator. Define the *dissonance operator*

$$\Delta\mathbf{U} \;:=\; \hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B \;=\; \Big(\int_{\mathcal{M}<\epsilon} - \int_{\mathcal{M}>\epsilon}\Big)\mathcal{D}\hat{\mathbf{\Phi}}\,\mathcal{D}\hat{\mathbf{\Theta}}\; e^{\,i\,S_{\text{total}}[\hat{\mathbf{\Phi}},\hat{\mathbf{\Theta}}]}, \tag{3.51}$$

so (3.50) becomes

$$\mathbf{U}_{\text{holo}}^{(\text{sym-com})} = \frac{1}{2}\begin{pmatrix} -\Delta\mathbf{U} & \Delta\mathbf{U} \\ \Delta\mathbf{U} & \Delta\mathbf{U} \end{pmatrix}. \tag{3.52}$$

Corollary 3.22 (Effect-level form for symmetric lensing)**.** *For initial control* $|\mathbf{a}\rangle_c$ *and readout* $|\mathbf{b}\rangle_c$ *in the computational basis, the channel Kraus block is* $K_{\mathbf{ba}} \in \{-\frac{1}{2}\Delta\mathbf{U},\ \frac{1}{2}\Delta\mathbf{U}\}$*. Hence every channel yields the same effect on* $\mathcal{H}_{\text{RR}}^{(2n)}$*:*

$$E_{\text{Holo}}^{(\mathbf{a}\nrightarrow\mathbf{b})} = K_{\mathbf{ba}}^{\dagger}K_{\mathbf{ba}} = \frac{1}{4}\,\Delta\mathbf{U}^{\dagger}\,\Delta\mathbf{U}, \qquad v_{\text{Holothesis}}(\rho) = \text{Tr}\Big(\rho\, E_{\text{Holo}}^{(\mathbf{a}\nrightarrow\mathbf{b})}\Big). \tag{3.53}$$

Thus lensing readouts measure the same "dissonance power'' irrespective of the computational channel; differences arise only from control-basis choices (cf. Definition 3.18).

Interpretation and operational note. Every entry of the lensing operator is proportional to the single object $\Delta\mathbf{U}$: the commutator acts as a *filter of differences*, canceling constructive (symmetric) contributions and isolating the algebraic "twist'' between the coupled and decoupled families. As written, $\mathbf{U}_{\text{holo}}^{(\text{sym-com})}$ need not be unitary by itself; experimentally it is accessed as a *difference of circuits* (e.g. control-sign tricks or linear-combination-of-unities methods), and its physical content is read out through effects and Born valuations defined in Definition 1.18. By the (H4) equivalences (Definition 3.18), lensing is a control-basis/readout reparametrization on the same dual-register backbone—no indefinite causal order is invoked.

The Asymmetric Lensing Operator

The defining operator isolates the non-commutativity between the controlled evolution and an arbitrary preparation, viewed from an arbitrary measurement basis:

$$\mathbf{U}_{\text{holo}}^{(\text{asym-com})} := [\mathbf{U}_{\text{ctrl}}, \mathbf{U}_{\text{prep}}]\;\mathbf{U}_{\text{read}}^{\dagger} = \Big(\mathbf{U}_{\text{ctrl}}\mathbf{U}_{\text{prep}} - \mathbf{U}_{\text{prep}}\mathbf{U}_{\text{ctrl}}\Big)\,\mathbf{U}_{\text{read}}^{\dagger},$$

where (as throughout) the rightmost gate acts first.

Formal derivation and path-integral representation

1. Components. Use general $SU(2)$ parametrizations for preparation and readout, and the path-integral–defined control:

$$\mathbf{U}_{\text{prep}} \leftrightarrow \begin{pmatrix} \alpha & -\beta^* \\ \beta & \alpha^* \end{pmatrix}, \quad \mathbf{U}_{\text{read}}^{\dagger} \leftrightarrow \begin{pmatrix} \gamma^* & \delta^* \\ -\delta & \gamma \end{pmatrix}, \quad \mathbf{U}_{\text{ctrl}} \leftrightarrow \begin{pmatrix} \hat{\mathbf{U}}_A & 0 \\ 0 & \hat{\mathbf{U}}_B \end{pmatrix},$$

with $|\alpha|^2+|\beta|^2 = |\gamma|^2+|\delta|^2 = 1$ (global phases immaterial). The path-integral forms are

$$\hat{\mathbf{U}}_A = \int_{\mathcal{M}<\epsilon}\mathcal{D}\hat{\mathbf{\Phi}}\,\mathcal{D}\hat{\mathbf{\Theta}}\; e^{iS_{\text{total}}[\hat{\mathbf{\Phi}},\hat{\mathbf{\Theta}}]}, \qquad \hat{\mathbf{U}}_B = \int_{\mathcal{M}>\epsilon}\mathcal{D}\hat{\mathbf{\Phi}}\,\mathcal{D}\hat{\mathbf{\Theta}}\; e^{iS_{\text{total}}[\hat{\mathbf{\Phi}},\hat{\mathbf{\Theta}}]}.$$

2. Calculating the commutator.

$$\mathbf{U}_{\text{ctrl}}\mathbf{U}_{\text{prep}} \leftrightarrow \begin{pmatrix} \alpha\hat{\mathbf{U}}_A & -\beta^*\hat{\mathbf{U}}_A \\ \beta\hat{\mathbf{U}}_B & \alpha^*\hat{\mathbf{U}}_B \end{pmatrix}, \qquad \mathbf{U}_{\text{prep}}\mathbf{U}_{\text{ctrl}} \leftrightarrow \begin{pmatrix} \alpha\hat{\mathbf{U}}_A & -\beta^*\hat{\mathbf{U}}_B \\ \beta\hat{\mathbf{U}}_A & \alpha^*\hat{\mathbf{U}}_B \end{pmatrix}.$$

Taking the difference gives

$$[\mathbf{U}_{\text{ctrl}}, \mathbf{U}_{\text{prep}}] = \begin{pmatrix} 0 & -\beta^*(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B) \\ -\beta(\hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B) & 0 \end{pmatrix}.$$

3. Applying the final gate. With $\Delta\mathbf{U} := \hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B$,

$$\mathbf{U}_{\text{holo}}^{(\text{asym-com})} = \begin{pmatrix} 0 & -\beta^*\,\Delta\mathbf{U} \\ -\beta\,\Delta\mathbf{U} & 0 \end{pmatrix} \begin{pmatrix} \gamma^* & \delta^* \\ -\delta & \gamma \end{pmatrix} \tag{3.54}$$

$$= \begin{pmatrix} \delta\,\beta^*\,\Delta\mathbf{U} & -\gamma\,\beta^*\,\Delta\mathbf{U} \\ -\gamma^*\,\beta\,\Delta\mathbf{U} & -\delta^*\,\beta\,\Delta\mathbf{U} \end{pmatrix}. \tag{3.55}$$

4. Final path-integral form (dissonance factorization). Define the *dissonance operator*

$$\Delta\mathbf{U} := \hat{\mathbf{U}}_A - \hat{\mathbf{U}}_B = \left(\int_{\mathcal{M}<\epsilon} - \int_{\mathcal{M}>\epsilon}\right) \mathcal{D}\hat{\mathbf{\Phi}}\,\mathcal{D}\hat{\mathbf{\Theta}}\; e^{iS_{\text{total}}[\hat{\mathbf{\Phi}},\hat{\mathbf{\Theta}}]},$$

so the operator factors as

$$\boxed{\mathbf{U}_{\text{holo}}^{(\text{asym-com})} = \begin{pmatrix} \delta\beta^* & -\gamma\beta^* \\ -\gamma^*\beta & -\delta^*\beta \end{pmatrix} \otimes \Delta\mathbf{U}}.$$

Remark 3.23 (Effect-level corollary and visibility)**.** Let K_{ba} denote the block mapping control $|a\rangle_c \to |b\rangle_c$ in the computational basis. Then

$$K_{00} = \delta\beta^*\,\Delta\mathbf{U}, \quad K_{01} = -\gamma\beta^*\,\Delta\mathbf{U}, \quad K_{10} = -\gamma^*\beta\,\Delta\mathbf{U}, \quad K_{11} = -\delta^*\beta\,\Delta\mathbf{U},$$

and the corresponding effects on $\mathcal{H}_{\text{RR}}^{(2n)}$ are

$$E_{\text{Holo}}^{(a\to b)} = K_{ba}^\dagger K_{ba} = |c_{ba}|^2\,\Delta\mathbf{U}^\dagger\Delta\mathbf{U}, \quad c_{ba} \in \{\delta\beta^*, -\gamma\beta^*, -\gamma^*\beta, -\delta^*\beta\}.$$

Thus all channels measure the same "dissonance power'' $\Delta\mathbf{U}^\dagger\Delta\mathbf{U}$, scaled by the control coefficients. Two practical consequences:

- **Preparation visibility:** if $\beta = 0$ (prep an eigenstate of Z_c), then $[\mathbf{U}_{\text{ctrl}}, \mathbf{U}_{\text{prep}}] = 0$ and the lensing signal vanishes.
- **Path degeneracy:** if $\Delta\mathbf{U} = 0$ (identical paths), then all effects vanish: asymmetric lensing detects no twist.

By (H4) equivalences (Definition 3.18), varying (α, β) and (γ, δ) tomographically scans the same dual-register backbone; no indefinite causal order is involved. Readouts use Born valuations as in Definition 1.18.

Interpretation. Asymmetric lensing is a pure $\Delta\mathbf{U}$ probe with tunable projection set by the preparation/readout angles. It complements the symmetric case: the former yields *channel-dependent* weights $|c_{ba}|^2$ on the same operator $\Delta\mathbf{U}^\dagger\Delta\mathbf{U}$, enabling full tomographic control over the algebraic twist between the coupled and decoupled history classes.

The Path Scattering Protocol

The defining operator for this model describes a three-act process (rightmost gate acts first): *Evolve* → *Interfere* → *Re-evolve*

$$\mathbf{U}_{\text{holo}}^{(\text{scatter,sym})} := \mathbf{U}_{\text{ctrl}} \cdot \mathbf{U}_{\text{Had}} \cdot \mathbf{U}_{\text{ctrl}}.$$

Here the registers evolve under $\mathbf{U}_{\text{ctrl}}$, are mixed by the Hadamard on the control, and then evolve again under the same controlled rule.

Formal derivation and path-integral representation

1. **Components.** Use the block forms

$$\mathbf{U}_{\text{Had}} \leftrightarrow \tfrac{1}{\sqrt{2}}\begin{pmatrix} \mathbb{I} & \mathbb{I} \\ \mathbb{I} & -\mathbb{I} \end{pmatrix}, \qquad \mathbf{U}_{\text{ctrl}} \leftrightarrow \begin{pmatrix} \hat{\mathbf{U}}_A & 0 \\ 0 & \hat{\mathbf{U}}_B \end{pmatrix},$$

with

$$\hat{\mathbf{U}}_A = \int_{\mathcal{M}<\epsilon} \mathcal{D}\hat{\mathbf{\Phi}}\,\mathcal{D}\hat{\mathbf{\Theta}}\; e^{\,i\,S_{\text{total}}}, \qquad \hat{\mathbf{U}}_B = \int_{\mathcal{M}>\epsilon} \mathcal{D}\hat{\mathbf{\Phi}}\,\mathcal{D}\hat{\mathbf{\Theta}}\; e^{\,i\,S_{\text{total}}}.$$

2. **Calculating the operator product.**

$$\mathbf{U}_{\text{Had}}\mathbf{U}_{\text{ctrl}} \leftrightarrow \tfrac{1}{\sqrt{2}}\begin{pmatrix} \hat{\mathbf{U}}_A & \hat{\mathbf{U}}_B \\ \hat{\mathbf{U}}_A & -\hat{\mathbf{U}}_B \end{pmatrix}, \quad \Rightarrow \quad \mathbf{U}_{\text{holo}}^{(\text{scatter,sym})} \leftrightarrow \tfrac{1}{\sqrt{2}}\begin{pmatrix} \hat{\mathbf{U}}_A^2 & \hat{\mathbf{U}}_A\hat{\mathbf{U}}_B \\ \hat{\mathbf{U}}_B\hat{\mathbf{U}}_A & -\hat{\mathbf{U}}_B^2 \end{pmatrix}.$$

3. **Final path-integral form.** Substituting the constrained path-integral definitions for $\hat{\mathbf{U}}_A$ and $\hat{\mathbf{U}}_B$ gives

$$\boxed{\mathbf{U}_{\text{holo}}^{(\text{scatter,sym})} = \tfrac{1}{\sqrt{2}}\begin{pmatrix} \left(\int_{\mathcal{M}<\epsilon}\mathcal{D}\hat{\mathbf{\Phi}}\mathcal{D}\hat{\mathbf{\Theta}}\, e^{iS_{\text{total}}}\right)^2 & \left(\int_{\mathcal{M}<\epsilon}\cdots\right)\left(\int_{\mathcal{M}>\epsilon}\cdots\right) \\ \left(\int_{\mathcal{M}>\epsilon}\cdots\right)\left(\int_{\mathcal{M}<\epsilon}\cdots\right) & -\left(\int_{\mathcal{M}>\epsilon}\mathcal{D}\hat{\mathbf{\Phi}}\mathcal{D}\hat{\mathbf{\Theta}}\, e^{iS_{\text{total}}}\right)^2 \end{pmatrix};}$$

the products denote sequential composition of the two path-selected evolutions (*evolve, then evolve again*).

Channel blocks and valuations (drop-in). Writing the control-basis blocks as K_{ba} for $|a\rangle_c \to |b\rangle_c$,

$$K_{00} = \tfrac{1}{\sqrt{2}}\,\hat{\mathbf{U}}_A^2, \quad K_{01} = \tfrac{1}{\sqrt{2}}\,\hat{\mathbf{U}}_A\hat{\mathbf{U}}_B, \quad K_{10} = \tfrac{1}{\sqrt{2}}\,\hat{\mathbf{U}}_B\hat{\mathbf{U}}_A, \quad K_{11} = -\tfrac{1}{\sqrt{2}}\,\hat{\mathbf{U}}_B^2.$$

The corresponding holosymmetric effects on $\mathcal{H}_{\text{RR}}^{(2n)}$ are $E_{\text{Holo}}^{(a\to b)} = K_{ba}^\dagger K_{ba}$, so Born valuations follow directly via $\text{Tr}(\rho\, E_{\text{Holo}}^{(a\to b)})$ (cf. Definition 1.18).

Interpretation: a universe in dialogue with itself. Each matrix entry is a distinct feedback channel:

- *Diagonal* $\left(\propto \hat{\mathbf{U}}_A^2,\, -\hat{\mathbf{U}}_B^2\right)$: self-reinforcing loops—histories that remain in the same dynamical regime across the interference event (continuity/stability).
- *Off-diagonal* $\left(\propto \hat{\mathbf{U}}_A\hat{\mathbf{U}}_B,\, \hat{\mathbf{U}}_B\hat{\mathbf{U}}_A\right)$: regime-switching channels—histories that swap between coupled and decoupled evolution (transformation/novelty). The non-commutativity $[\hat{\mathbf{U}}_A, \hat{\mathbf{U}}_B]$ controls the asymmetry between the two off-diagonal amplitudes.

Thus path scattering models *reflexive* dynamics: past evolution conditions the interference, which conditions the future evolution.

Sanity checks and special cases.

1. **Identical paths** ($\hat{\mathbf{U}}_A = \hat{\mathbf{U}}_B = \hat{\mathbf{U}}$). Then $\mathbf{U}_{\text{ctrl}} = \mathbb{I}_c \otimes \hat{\mathbf{U}}$ and $\mathbf{U}_{\text{holo}}^{(\text{scatter,sym})} = (\mathbf{U}_{\text{Had}} \otimes \hat{\mathbf{U}}^2)$. Off-diagonals become equal ($= \frac{1}{\sqrt{2}}\hat{\mathbf{U}}^2$); the protocol reduces to a trivial re-labelling on the control with a squared evolution on the registers.
2. **Inverse paths** ($\hat{\mathbf{U}}_B = \hat{\mathbf{U}}_A^\dagger$). Then $K_{00} = \frac{1}{\sqrt{2}}\hat{\mathbf{U}}_A^2$, $K_{11} = -\frac{1}{\sqrt{2}}\mathbb{I}$; the two off-diagonals are $\frac{1}{\sqrt{2}}\hat{\mathbf{U}}_A\hat{\mathbf{U}}_A^\dagger = \frac{1}{\sqrt{2}}\mathbb{I}$ and its reverse, giving a clean split between "retain A", "retain B", and "swap" channels.
3. **Commuting paths** ($[\hat{\mathbf{U}}_A, \hat{\mathbf{U}}_B] = 0$). Off-diagonals are equal; any asymmetry in switching rates must then arise from state preparation, not algebraic twist.

Unitarity and consistency. If $\hat{\mathbf{U}}_A, \hat{\mathbf{U}}_B$ are unitary, then each factor in $\mathbf{U}_{\text{holo}}^{(\text{scatter,sym})} = \mathbf{U}_{\text{ctrl}}\mathbf{U}_{\text{Had}}\mathbf{U}_{\text{ctrl}}$ is unitary, so the overall protocol is unitary. This places the model squarely within standard circuit mechanics (no indefinite causal order). As noted above, the block identification shows this protocol is a special point of the Master Operator:

$$\mathbf{A} = \tfrac{1}{\sqrt{2}}\hat{\mathbf{U}}_A^2, \quad \mathbf{B} = \tfrac{1}{\sqrt{2}}\hat{\mathbf{U}}_A\hat{\mathbf{U}}_B, \quad \mathbf{C} = \tfrac{1}{\sqrt{2}}\hat{\mathbf{U}}_B\hat{\mathbf{U}}_A, \quad \mathbf{D} = -\tfrac{1}{\sqrt{2}}\hat{\mathbf{U}}_B^2,$$

with "diagonal symmetry up to a sign'' and generally unequal off-diagonals when the paths do not commute.

The Asymmetric Path Scattering Protocol

The defining operator describes a directed, three–act process (rightmost acts first): *Evolve under "in" rules → Interfere → Re-evolve under "out" rules*:

$$\mathbf{U}_{\text{holo}}^{(\text{scatter,asym})} := \mathbf{U}_{\text{ctrl}}^{\text{out}}\, \mathbf{U}_{\text{Had}}\, \mathbf{U}_{\text{ctrl}}^{\text{in}}. \tag{3.56}$$

Here the "in" and "out" control operators may be built from different archetypal path operators, representing a structural change in the dynamical regime.

Formal derivation and path–integral representation

1. Components.

$$\mathbf{U}^{\text{in}}_{\text{ctrl}} \leftrightarrow \begin{pmatrix} \hat{\mathbf{U}}^{\text{in}}_A & 0 \\ 0 & \hat{\mathbf{U}}^{\text{in}}_B \end{pmatrix}, \qquad \mathbf{U}_{\text{Had}} \leftrightarrow \tfrac{1}{\sqrt{2}} \begin{pmatrix} \mathbb{I} & \mathbb{I} \\ \mathbb{I} & -\mathbb{I} \end{pmatrix}, \qquad \mathbf{U}^{\text{out}}_{\text{ctrl}} \leftrightarrow \begin{pmatrix} \hat{\mathbf{U}}^{\text{out}}_A & 0 \\ 0 & \hat{\mathbf{U}}^{\text{out}}_B \end{pmatrix}.$$

2. Operator product. Multiplying from right to left,

$$\mathbf{U}^{(\text{scatter,asym})}_{\text{holo}} \leftrightarrow \tfrac{1}{\sqrt{2}} \begin{pmatrix} \hat{\mathbf{U}}^{\text{out}}_A \hat{\mathbf{U}}^{\text{in}}_A & \hat{\mathbf{U}}^{\text{out}}_A \hat{\mathbf{U}}^{\text{in}}_B \\ \hat{\mathbf{U}}^{\text{out}}_B \hat{\mathbf{U}}^{\text{in}}_A & -\hat{\mathbf{U}}^{\text{out}}_B \hat{\mathbf{U}}^{\text{in}}_B \end{pmatrix}. \tag{3.57}$$

3. Path–integral building blocks. Using (potentially) distinct modularity functionals/thresholds for the two epochs:

$$\boxed{\hat{\mathbf{U}}^{\text{in}}_A := \int_{\mathcal{M}_{\text{in}} < \epsilon_{\text{in}}} \mathcal{D}\hat{\mathbf{\Phi}}\, \mathcal{D}\hat{\mathbf{\Theta}}\; e^{\,i\, S_{\text{total}}[\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}]}\,, \quad \hat{\mathbf{U}}^{\text{in}}_B := \int_{\mathcal{M}_{\text{in}} > \epsilon_{\text{in}}} \mathcal{D}\hat{\mathbf{\Phi}}\, \mathcal{D}\hat{\mathbf{\Theta}}\; e^{\,i\, S_{\text{total}}[\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}]}} \tag{3.58}$$

$$\boxed{\hat{\mathbf{U}}^{\text{out}}_A := \int_{\mathcal{M}_{\text{out}} < \epsilon_{\text{out}}} \mathcal{D}\hat{\mathbf{\Phi}}\, \mathcal{D}\hat{\mathbf{\Theta}}\; e^{\,i\, S_{\text{total}}[\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}]}\,, \quad \hat{\mathbf{U}}^{\text{out}}_B := \int_{\mathcal{M}_{\text{out}} > \epsilon_{\text{out}}} \mathcal{D}\hat{\mathbf{\Phi}}\, \mathcal{D}\hat{\mathbf{\Theta}}\; e^{\,i\, S_{\text{total}}[\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}]}} \tag{3.59}$$

(coupled A vs. decoupled B, as in the foundational definitions).

Channel blocks and effects (for valuations). Denote the control–basis blocks by K^{asym}_{ba} for the $|a\rangle_c \to |b\rangle_c$ channel. From (3.57):

$$\begin{aligned} K^{\text{asym}}_{00} &= \tfrac{1}{\sqrt{2}}\, \hat{\mathbf{U}}^{\text{out}}_A \hat{\mathbf{U}}^{\text{in}}_A, \quad K^{\text{asym}}_{01} = \tfrac{1}{\sqrt{2}}\, \hat{\mathbf{U}}^{\text{out}}_A \hat{\mathbf{U}}^{\text{in}}_B, \\ K^{\text{asym}}_{10} &= \tfrac{1}{\sqrt{2}}\, \hat{\mathbf{U}}^{\text{out}}_B \hat{\mathbf{U}}^{\text{in}}_A, \quad K^{\text{asym}}_{11} = -\tfrac{1}{\sqrt{2}}\, \hat{\mathbf{U}}^{\text{out}}_B \hat{\mathbf{U}}^{\text{in}}_B. \end{aligned}$$

The holosymmetric effects on $\mathcal{H}^{(2n)}_{\text{RR}}$ are $E^{(a \not\to b)}_{\text{Holo}} = (K^{\text{asym}}_{ba})^\dagger K^{\text{asym}}_{ba}$, so the Born valuations are $v(\rho; E) = \text{Tr}\big(\rho\, E^{(a \not\to b)}_{\text{Holo}}\big)$ (cf. Definition 1.18).

Interpretation: the mechanism of a learning universe. Each block is a complete, transformative history:

- *Continuity through transformation*: $K^{\text{asym}}_{00} \propto \hat{\mathbf{U}}^{\text{out}}_A \hat{\mathbf{U}}^{\text{in}}_A$ and $K^{\text{asym}}_{11} \propto -\hat{\mathbf{U}}^{\text{out}}_B \hat{\mathbf{U}}^{\text{in}}_B$ describe persistence of the dynamical character across the interference boundary ($A \to A$, $B \to B$).
- *Regime change (revolution)*: $K^{\text{asym}}_{10} \propto \hat{\mathbf{U}}^{\text{out}}_B \hat{\mathbf{U}}^{\text{in}}_A$ and $K^{\text{asym}}_{01} \propto \hat{\mathbf{U}}^{\text{out}}_A \hat{\mathbf{U}}^{\text{in}}_B$ implement $A \to B$ and $B \to A$ transitions, with asymmetry governed by the algebra and the state of the registers.

Irreversibility is encoded by the mismatch between "in" and "out'' operators and thresholds—this protocol models how interference can *update* the effective laws.

Sanity checks and special cases.

1. **No regime change** ($\hat{\mathbf{U}}_{\bullet}^{\text{out}} = \hat{\mathbf{U}}_{\bullet}^{\text{in}}$). Then $\mathbf{U}_{\text{holo}}^{(\text{scatter,asym})}$ reduces to the symmetric scatter form.

2. **Pure switcher** ($\hat{\mathbf{U}}_{A}^{\text{out}} = \hat{\mathbf{U}}_{B}^{\text{out}}$ or $\hat{\mathbf{U}}_{A}^{\text{in}} = \hat{\mathbf{U}}_{B}^{\text{in}}$). One off–diagonal channel can dominate; use this to design targeted $A \leftrightarrow B$ conversions.

3. **Unitarity.** If each $\hat{\mathbf{U}}_{A/B}^{\text{in/out}}$ is unitary, then $\mathbf{U}_{\text{ctrl}}^{\text{in/out}}$ and $\mathbf{U}_{\text{Had}}$ are unitary, hence $\mathbf{U}_{\text{holo}}^{(\text{scatter,asym})}$ is unitary. No indefinite causal order is invoked.

Part II

The Continuum

Chapter 4

Modal Fibres and Emergent Hilbert Spaces

4.1 Structure of Emergent Hilbert Spaces

4.1.1 Twofold Inter-Modal Correlator of Primordial Quantum Singularities

Definition 4.1 (Twofold Inter-Modal Correlator of Primordial Quantum Singularities)**.** Fix a PQS locus $\mathfrak{B}_x$. The Symmetry–Antisymmetry (Twofold) sector

$$\mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}\,(0,1)} := \mathrm{span}\{\, |\mathring{\mathbf{0}}\rangle_x,\, |\mathring{\mathbf{1}}\rangle_x \,\}$$

is the $(0,1)$-slice of the Fourfold modal fibre introduced in the Fourfold Dialectic setting (Chapter 1; ringed kets and the prism correspondence Sym $\leftrightarrow$ Th, AntiSym $\leftrightarrow$ AntiTh, cf. Definitions 1.16 and 1.23 and Lemma 1.20), and realized concretely in Chapter 2 as the Twofold subspace with the $(0,1)$ basis and Born-effect semantics (Definition 1.18 and the $(0,1)$ realisation therein).

Let $\mathcal{U}(x \leftarrow x';\gamma)$ denote the parallel transporter generated by the emergent connection $\mathcal{A}$ along a path $\gamma \subset \mathcal{X}_q$ (Chapter 4; eqs. (4.11), (4.14) and (4.16)). For the Fourfold selection operators $\hat{\mathcal{F}}_\alpha(\cdot)$ (Chapter 1; transported to fibres via the bridge congruence Φ, Lemma 1.20), define the restricted components

$$\hat{G}^{\alpha\beta}_{(0,1)}(x,x') := \hat{\mathcal{F}}^\dagger_\alpha(x)\,\mathcal{U}(x \leftarrow x';\gamma)\,\hat{\mathcal{F}}_\beta(x') \qquad (\alpha,\beta \in \{0,1\}),$$

i.e. the $(0,1)$-block of the Fourfold correlator kernel (Chapter 4; $\hat{\mathbf{G}}_\Lambda$ and its block structure $\{0,1,p,s\}$). Writing

$$|\psi_1\rangle_x = c^{(1)}_0|\mathring{\mathbf{0}}\rangle_x + c^{(1)}_1|\mathring{\mathbf{1}}\rangle_x, \quad |\psi_2\rangle_{x'} = c^{(2)}_0|\mathring{\mathbf{0}}\rangle_{x'} + c^{(2)}_1|\mathring{\mathbf{1}}\rangle_{x'},$$

and $c^{(j)} = (c^{(j)}_0, c^{(j)}_1)^\top$, set the Twofold Inter-Modal Correlator to be the operator-valued, sesquilinear kernel

$$\begin{aligned} \hat{G}_{(0,1)}(\mathfrak{B}_x,\mathfrak{B}_{x'})[\psi_1^*,\psi_2] &:= \left(c^{(1)}\right)^\dagger \hat{\mathbf{G}}_{(0,1)}(x,x')\,c^{(2)}, \\ \hat{\mathbf{G}}_{(0,1)}(x,x') &= \left[\hat{G}^{\alpha\beta}_{(0,1)}(x,x')\right]_{\alpha,\beta\in\{0,1\}}. \end{aligned} \tag{4.1}$$

Here the action of $\hat{\mathcal{F}}_\alpha$ on modal states is the operator realisation of the corner selectors of Chapter 1 (Fourfold *-algebra; Definition 1.16), and $\mathcal{U}$ is the path-ordered exponential of $\mathcal{A}$ (Chapter 4; eq. (4.11)). In interferometric realisations with a control register, $\hat{\mathcal{F}}_\alpha$ arise as the corner-conditioned blocks of the master holosymmetric operator of Chapter 3 (Section 3.8.3 and Corollary 3.21), but the present definition is representation-agnostic.

Basic properties of the twofold inter-modal correlator

Proposition 4.2 (Sesquilinearity)**.** *The map $(\psi_1^*, \psi_2) \mapsto \hat{G}_{(0,1)}[\psi_1^*, \psi_2]$ is conjugate-linear in the first slot and linear in the second:*

$$\hat{G}_{(0,1)}[\lambda\psi_1^* + \mu\phi_1^*, \psi_2] = \overline{\lambda}\,\hat{G}_{(0,1)}[\psi_1^*, \psi_2] + \overline{\mu}\,\hat{G}_{(0,1)}[\phi_1^*, \psi_2],$$
$$\hat{G}_{(0,1)}[\psi_1^*, \lambda\psi_2 + \mu\phi_2] = \lambda\,\hat{G}_{(0,1)}[\psi_1^*, \psi_2] + \mu\,\hat{G}_{(0,1)}[\psi_1^*, \phi_2].$$

Proof. Let $\psi_1, \phi_1 \in \mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}\,(0,1)}$ and $\psi_2, \phi_2 \in \mathcal{H}^{(\mathfrak{B}_{x'})}_{\mathrm{mod}\,(0,1)}$. Write the coefficient maps in the fixed $(0,1)$-basis

$$\mathcal{C}_x : \mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}\,(0,1)} \to \mathbb{C}^2, \qquad \mathcal{C}_{x'} : \mathcal{H}^{(\mathfrak{B}_{x'})}_{\mathrm{mod}\,(0,1)} \to \mathbb{C}^2,$$

so that $c^{(1)} := \mathcal{C}_x(\psi_1)$, $d^{(1)} := \mathcal{C}_x(\phi_1)$, $c^{(2)} := \mathcal{C}_{x'}(\psi_2)$, $d^{(2)} := \mathcal{C}_{x'}(\phi_2)$. By definition of the Twofold correlator (cf. (4.1)),

$$\hat{G}_{(0,1)}(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2] = (c^{(1)})^\dagger\, \hat{\mathbf{G}}_{(0,1)}(x, x')\, c^{(2)},$$

where $\hat{\mathbf{G}}_{(0,1)}(x, x') \in \mathcal{B}(\mathcal{H}_{\mathrm{em}})$ is fixed by (x, x').

(1) Conjugate homogeneity in the first slot. For $\lambda, \mu \in \mathbb{C}$,

$$\begin{aligned}
\hat{G}_{(0,1)}[(\lambda\psi_1 + \mu\phi_1)^*, \psi_2] &= \big(\mathcal{C}_x(\lambda\psi_1 + \mu\phi_1)\big)^\dagger\, \hat{\mathbf{G}}_{(0,1)}\, c^{(2)} \\
&= (\lambda c^{(1)} + \mu d^{(1)})^\dagger\, \hat{\mathbf{G}}_{(0,1)}\, c^{(2)} \quad \text{(linearity of } \mathcal{C}_x\text{)} \\
&= \overline{\lambda}\,(c^{(1)})^\dagger\, \hat{\mathbf{G}}_{(0,1)}\, c^{(2)} \;+\; \overline{\mu}\,(d^{(1)})^\dagger\, \hat{\mathbf{G}}_{(0,1)}\, c^{(2)} \\
&= \overline{\lambda}\,\hat{G}_{(0,1)}[\psi_1^*, \psi_2] \;+\; \overline{\mu}\,\hat{G}_{(0,1)}[\phi_1^*, \psi_2].
\end{aligned}$$

(2) Additivity in the first slot. Setting $\lambda = \mu = 1$ in the previous computation gives

$$\hat{G}_{(0,1)}[(\psi_1 + \phi_1)^*, \psi_2] = \hat{G}_{(0,1)}[\psi_1^*, \psi_2] + \hat{G}_{(0,1)}[\phi_1^*, \psi_2].$$

(3) Linearity in the second slot (homogeneity and additivity). For $\lambda, \mu \in \mathbb{C}$,

$$\begin{aligned}
\hat{G}_{(0,1)}[\psi_1^*, \lambda\psi_2 + \mu\phi_2] &= (c^{(1)})^\dagger\, \hat{\mathbf{G}}_{(0,1)}\, \mathcal{C}_{x'}(\lambda\psi_2 + \mu\phi_2) \\
&= (c^{(1)})^\dagger\, \hat{\mathbf{G}}_{(0,1)}\, (\lambda c^{(2)} + \mu d^{(2)}) \quad \text{(linearity of } \mathcal{C}_{x'}\text{)} \\
&= \lambda\,(c^{(1)})^\dagger\, \hat{\mathbf{G}}_{(0,1)}\, c^{(2)} \;+\; \mu\,(c^{(1)})^\dagger\, \hat{\mathbf{G}}_{(0,1)}\, d^{(2)} \\
&= \lambda\,\hat{G}_{(0,1)}[\psi_1^*, \psi_2] \;+\; \mu\,\hat{G}_{(0,1)}[\psi_1^*, \phi_2].
\end{aligned}$$

Items (1)–(3) together establish conjugate-linearity in the first argument and linearity in the second. All equalities hold in the operator space $\mathcal{B}(\mathcal{H}_{\mathrm{em}})$, and no further properties of $\hat{\mathbf{G}}_{(0,1)}$ are required beyond its being fixed for (x, x'). □

Proposition 4.3 (Hermitian symmetry)**.** *For all* $\alpha, \beta \in \{0,1\}$,

$$\left(\hat{G}^{\alpha\beta}_{(0,1)}(x,x')\right)^\dagger = \hat{G}^{\beta\alpha}_{(0,1)}(x',x),$$

and thus

$$\hat{G}_{(0,1)}(\mathfrak{B}_x,\mathfrak{B}_{x'})[\psi_1^*,\psi_2]^\dagger = \hat{G}_{(0,1)}(\mathfrak{B}_{x'},\mathfrak{B}_x)[\psi_2^*,\psi_1].$$

Proof. Step 0: Adjointability and domain. By construction, $\hat{\mathcal{F}}_\alpha(x)$ are bounded (or at least adjointable) operators and $\mathcal{U}(x \leftarrow x';\gamma)$ is unitary as a parallel transport generated by the connection $\mathcal{A}$; hence all products below are adjointable with adjoints in $\mathcal{B}(\mathcal{H}_{\mathrm{em}})$.

Step 1: Component-level symmetry. By definition,

$$\hat{G}^{\alpha\beta}_{(0,1)}(x,x') = \hat{\mathcal{F}}^\dagger_\alpha(x)\,\mathcal{U}(x \leftarrow x';\gamma)\,\hat{\mathcal{F}}_\beta(x').$$

Taking adjoints and using $(ABC)^\dagger = C^\dagger B^\dagger A^\dagger$, we obtain

$$\left(\hat{G}^{\alpha\beta}_{(0,1)}(x,x')\right)^\dagger = \hat{\mathcal{F}}^\dagger_\beta(x')\left(\mathcal{U}(x \leftarrow x';\gamma)\right)^\dagger \hat{\mathcal{F}}_\alpha(x).$$

Unitarity and the path-reversal/composition identities for transport (Chapter 4; $\mathcal{U}^\dagger = \mathcal{U}^{-1}$ and $\mathcal{U}(x \leftarrow x';\gamma)^{-1} = \mathcal{U}(x' \leftarrow x;\gamma^{-1})$, cf. eq. (4.14)) give

$$\left(\hat{G}^{\alpha\beta}_{(0,1)}(x,x')\right)^\dagger = \hat{\mathcal{F}}^\dagger_\beta(x')\,\mathcal{U}(x' \leftarrow x;\gamma^{-1})\,\hat{\mathcal{F}}_\alpha(x) = \hat{G}^{\beta\alpha}_{(0,1)}(x',x),$$

which is the desired component identity.

Step 2: Kernel-level symmetry. Write the coefficient expansions in the fixed $(0,1)$-basis:

$$|\psi_1\rangle_x = \sum_{\alpha\in\{0,1\}} c^{(1)}_\alpha\, |\overset{\circ}{\alpha}\rangle_x, \qquad |\psi_2\rangle_{x'} = \sum_{\beta\in\{0,1\}} c^{(2)}_\beta\, |\overset{\circ}{\beta}\rangle_{x'}.$$

Then (cf. (4.1))

$$\hat{G}_{(0,1)}(\mathfrak{B}_x,\mathfrak{B}_{x'})[\psi_1^*,\psi_2] = \sum_{\alpha,\beta} \overline{c^{(1)}_\alpha}\; \hat{G}^{\alpha\beta}_{(0,1)}(x,x')\; c^{(2)}_\beta.$$

Taking adjoints and using linearity of the adjoint together with the scalar rule $(\overline{z}\,A\,w)^\dagger = \overline{w}\,A^\dagger z$ for $z, w \in \mathbb{C}$, we get

$$\left(\hat{G}_{(0,1)}[\psi_1^*,\psi_2]\right)^\dagger = \sum_{\alpha,\beta} \overline{c^{(2)}_\beta}\; \left(\hat{G}^{\alpha\beta}_{(0,1)}(x,x')\right)^\dagger c^{(1)}_\alpha.$$

Applying Step 1 to the component adjoint yields

$$\left(\hat{G}_{(0,1)}[\psi_1^*,\psi_2]\right)^\dagger = \sum_{\alpha,\beta} \overline{c^{(2)}_\beta}\; \hat{G}^{\beta\alpha}_{(0,1)}(x',x)\; c^{(1)}_\alpha.$$

Relabelling the dummy indices $\alpha \leftrightarrow \beta$ gives

$$\left(\hat{G}_{(0,1)}[\psi_1^*,\psi_2]\right)^\dagger = \sum_{\alpha,\beta} \overline{c^{(2)}_\alpha}\; \hat{G}^{\alpha\beta}_{(0,1)}(x',x)\; c^{(1)}_\beta = \hat{G}_{(0,1)}(\mathfrak{B}_{x'},\mathfrak{B}_x)[\psi_2^*,\psi_1],$$

which proves the second identity. □

Proposition 4.4 (Gauge covariance)**.** *Under local frame changes $g(\cdot)$ (Chapter 4; eq. (4.16)), the fields transform as*

$$\hat{\mathcal{F}}_\alpha(x) \mapsto g(x)\hat{\mathcal{F}}_\alpha(x), \qquad \mathcal{U}(x \leftarrow x';\gamma) \mapsto g(x)\,\mathcal{U}(x \leftarrow x';\gamma)\,g(x')^\dagger,$$

and the block obeys

$$\hat{\mathbf{G}}_{(0,1)}(x,x') \mapsto g(x)\,\hat{\mathbf{G}}_{(0,1)}(x,x')\,g(x')^\dagger.$$

Consequently, any Born expectation in a gauge-covariant state (Chapter 1; Definition 1.18) is gauge-invariant.

Proof. Step 0: Adjoints and transport. From the stated transformation of $\hat{\mathcal{F}}_\alpha(x)$, taking adjoints yields $\hat{\mathcal{F}}^\dagger_\alpha(x) \mapsto \hat{\mathcal{F}}^\dagger_\alpha(x)\,g(x)^\dagger$. For transport, $\mathcal{U} \mapsto g(x)\mathcal{U}g(x')^\dagger$ (Chapter 4; eq. (4.16)).

Step 1: Component-level transformation. By definition,

$$\hat{G}^{\alpha\beta}_{(0,1)}(x,x') = \hat{\mathcal{F}}^\dagger_\alpha(x)\,\mathcal{U}(x \leftarrow x';\gamma)\,\hat{\mathcal{F}}_\beta(x').$$

After a gauge change,

$$\begin{aligned}\Big(\hat{G}^{\alpha\beta}_{(0,1)}(x,x')\Big)' &= \Big(\hat{\mathcal{F}}^\dagger_\alpha(x)\,g(x)^\dagger\Big)\Big(g(x)\,\mathcal{U}(x \leftarrow x';\gamma)\,g(x')^\dagger\Big)\Big(g(x')\,\hat{\mathcal{F}}_\beta(x')\Big)\\ &= \hat{\mathcal{F}}^\dagger_\alpha(x)\,\underbrace{g(x)^\dagger g(x)}_{=\mathbb{I}}\,\mathcal{U}(x \leftarrow x';\gamma)\,\underbrace{g(x')^\dagger g(x')}_{=\mathbb{I}}\,\hat{\mathcal{F}}_\beta(x')\\ &= \hat{G}^{\alpha\beta}_{(0,1)}(x,x').\end{aligned}$$

Thus each *component* is gauge-*invariant* as an operator in $\mathcal{B}(\mathcal{H}_{\mathrm{em}})$.

Step 2: Block-level covariance as change of frames at the endpoints. Arrange the components into the 2×2 block $\hat{\mathbf{G}}_{(0,1)}(x,x') = \big[\hat{G}^{\alpha\beta}_{(0,1)}(x,x')\big]_{\alpha,\beta\in\{0,1\}}$. This block is naturally a morphism from the x'-anchored frame to the x-anchored frame. Under a change of local frames at the endpoints by $g(x)$ and $g(x')$, the *coordinate* representation of a fixed morphism transforms by left/right action:

$$\hat{\mathbf{G}}_{(0,1)}(x,x') \;\longmapsto\; g(x)\,\hat{\mathbf{G}}_{(0,1)}(x,x')\,g(x')^\dagger.$$

This expresses the usual functorial behaviour of a bi-anchored kernel under basis change. It is compatible with Step 1: the underlying operator is invariant, while its matrix representatives in the moved frames are related by conjugation at the endpoints.

Step 3: Kernel-level covariance for arbitrary inputs. Let $|\psi_1\rangle_x = \sum_\alpha c^{(1)}_\alpha |\overset{\circ}{\alpha}\rangle_x$, $|\psi_2\rangle_{x'} = \sum_\beta c^{(2)}_\beta |\overset{\circ}{\beta}\rangle_{x'}$, and $c^{(j)} = (c^{(j)}_0, c^{(j)}_1)^\top$. Then (cf. (4.1))

$$\hat{G}_{(0,1)}[\psi_1^*,\psi_2] = (c^{(1)})^\dagger\,\hat{\mathbf{G}}_{(0,1)}(x,x')\,c^{(2)}.$$

Under frame changes, the coefficient vectors transform contragrediently: $c^{(1)} \mapsto (c^{(1)})' = g(x)\,c^{(1)}$ and $c^{(2)} \mapsto (c^{(2)})' = g(x')\,c^{(2)}$ (i.e., their components are taken in the moved frames). Hence

$$\begin{aligned}\Big(\hat{G}_{(0,1)}[\psi_1^*,\psi_2]\Big)' &= \Big((c^{(1)})'\Big)^\dagger\,\Big(\hat{\mathbf{G}}_{(0,1)}\Big)'\,(c^{(2)})'\\ &= (c^{(1)})^\dagger g(x)^\dagger\,\Big(g(x)\hat{\mathbf{G}}_{(0,1)}g(x')^\dagger\Big)\,g(x')\,c^{(2)}\\ &= (c^{(1)})^\dagger\,\hat{\mathbf{G}}_{(0,1)}\,c^{(2)} = \hat{G}_{(0,1)}[\psi_1^*,\psi_2],\end{aligned}$$

so the *kernel value* is invariant as an operator in $\mathcal{B}(\mathcal{H}_{\mathrm{em}})$.

Step 4: Gauge invariance of Born expectations. Let ρ be a (gauge-covariant) state on $\mathcal{H}_{\mathrm{em}}$. If $\rho \mapsto \rho' = g(x)\,\rho\,g(x)^\dagger$ (or, more generally, the appropriate anchor is used), then by the cyclicity of the trace and the block transformation in Step 2,

$$\mathrm{Tr}\Big(\rho\,\hat{\mathbf{G}}_{(0,1)}(x,x')\Big) = \mathrm{Tr}\Big(g(x)\rho g(x)^\dagger\; g(x)\hat{\mathbf{G}}_{(0,1)}(x,x')g(x')^\dagger\Big) = \mathrm{Tr}\Big(\rho'\,\hat{\mathbf{G}}'_{(0,1)}(x,x')\Big),$$

and likewise for vector states $|\Psi\rangle$ with $|\Psi\rangle \mapsto |\Psi'\rangle = g(\cdot)|\Psi\rangle$. Therefore, Born expectations (Chapter 1; Definition 1.18) are gauge-invariant.

The proposition follows: components are gauge-invariant, their block transforms covariantly under endpoint frame changes, and all Born valuations are invariant. □

Alternative proof (fibre-bundle notation). Let $\pi : \mathscr{H}_{\mathrm{mod}} \to \mathcal{X}_q$ be the modal Hilbert bundle of Chapter 4 and let $\mathscr{H}_{(0,1)} \subset \mathscr{H}_{\mathrm{mod}}$ be its Twofold subbundle with fibres $\pi^{-1}(x) = \mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}\,(0,1)}$. Write $P \to \mathcal{X}_q$ for the principal unitary frame bundle of $\mathscr{H}_{(0,1)}$ with structure group $G_{(0,1)} \subseteq \mathrm{U}(2)$. A (local) gauge transformation is a vertical bundle automorphism of P whose representative in a trivialisation $U \subset \mathcal{X}_q$ is a smooth map $g : U \to G_{(0,1)}$.

Selectors as an associated bundle morphism. Define the bundle map

$$\mathcal{F} :\ \mathscr{H}_{(0,1)} \longrightarrow \underline{\mathcal{B}(\mathcal{H}_{\mathrm{em}})} \qquad (x,v) \mapsto \hat{\mathcal{F}}_x(v) := \sum_{\alpha\in\{0,1\}} v_\alpha\,\hat{\mathcal{F}}_\alpha(x),$$

where $v = \sum_\alpha v_\alpha\,|\overset{\circ}{\alpha}\rangle_x$ in a local frame and $\underline{\mathcal{B}(\mathcal{H}_{\mathrm{em}})} = \mathcal{X}_q \times \mathcal{B}(\mathcal{H}_{\mathrm{em}})$ is the trivial bundle. By the Chapter 1 transformation law, $\hat{\mathcal{F}}_\alpha(x) \mapsto g(x)\hat{\mathcal{F}}_\alpha(x)$, hence $\mathcal{F}$ is $G_{(0,1)}$-equivariant:

$$\mathcal{F}_{x\cdot g}\Big(v\cdot g^{-1}(x)\Big) \;=\; g(x)\,\mathcal{F}_x(v).$$

Consequently, the adjoint map $\mathcal{F}_x^\dagger$ (with the adjoint taken in $\mathcal{B}(\mathcal{H}_{\mathrm{em}})$) transforms as $\mathcal{F}_x^\dagger \mapsto \mathcal{F}_x^\dagger\,g(x)^\dagger$.

Transport as a bundle morphism. Let $\mathcal{A}$ be the connection of Chapter 4 and let $\mathcal{U}(x \leftarrow x';\gamma) : \mathscr{H}_{(0,1),x'} \to \mathscr{H}_{(0,1),x}$ be the parallel transport along γ (eq. (4.11)). Under the gauge change g, its local representative transforms as

$$\mathcal{U}(x\leftarrow x';\gamma) \;\longmapsto\; g(x)\,\mathcal{U}(x\leftarrow x';\gamma)\,g(x')^\dagger \qquad \text{(Chapter 4; eq. (4.16))}.$$

The correlator as a section of a Hom-bundle. Define the bi-anchored bundle morphism

$$\hat{\mathbf{G}}_{(0,1)}(x,x') \;:=\; \mathcal{F}_x^\dagger \circ \mathcal{U}(x\leftarrow x';\gamma) \circ \mathcal{F}_{x'} \;\in\; \mathrm{Hom}\Big(\mathscr{H}_{(0,1),x'}, \mathscr{H}_{(0,1),x}\Big) \otimes \mathcal{B}(\mathcal{H}_{\mathrm{em}}).$$

Using the equivariance of $\mathcal{F}$ and the gauge law for $\mathcal{U}$, we obtain

$$\hat{\mathbf{G}}_{(0,1)}(x,x') \;\longmapsto\; \Big(\mathcal{F}_x^\dagger g(x)^\dagger\Big)\Big(g(x)\mathcal{U}g(x')^\dagger\Big)\Big(g(x')\mathcal{F}_{x'}\Big) \;=\; g(x)\,\hat{\mathbf{G}}_{(0,1)}(x,x')\,g(x')^\dagger,$$

which is exactly the announced two-sided covariance at the endpoints.

Coordinate-free kernel and invariance. Given $\psi_1 \in \mathscr{H}_{(0,1),x}$ and $\psi_2 \in \mathscr{H}_{(0,1),x'}$, define the operator-valued kernel

$$\hat{G}_{(0,1)}(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2] \;:=\; \hat{\mathbf{G}}_{(0,1)}(x,x')\Big(\psi_2\Big) \;\text{ contracted with }\; \psi_1^*,$$

which in a local frame reproduces $(c^{(1)})^\dagger\, \hat{\mathbf{G}}_{(0,1)}(x,x')\, c^{(2)}$ (cf. (4.1)). Under a gauge change, vectors ψ_1, ψ_2 co-transform as $\psi_1 \mapsto g(x)\psi_1$, $\psi_2 \mapsto g(x')\psi_2$, whence

$$\begin{aligned}\hat{G}'_{(0,1)}[\psi_1'^*, \psi_2'] &= \big(\psi_1'\big)^\dagger\, \hat{\mathbf{G}}'_{(0,1)}\, \psi_2' = \psi_1^\dagger g(x)^\dagger\, \Big(g(x)\hat{\mathbf{G}}_{(0,1)}g(x')^\dagger\Big)\, g(x')\psi_2 = \psi_1^\dagger\, \hat{\mathbf{G}}_{(0,1)}\, \psi_2 \\ &= \hat{G}_{(0,1)}[\psi_1^*, \psi_2],\end{aligned}$$

so the operator-valued kernel is gauge *invariant* for co-transforming inputs.

Born expectations are gauge-invariant. For any state ρ on $\mathcal{H}_{\mathrm{em}}$ (Chapter 1; Definition 1.18), let $\rho \mapsto \rho' = g(x)\rho g(x)^\dagger$ be the induced transformation at the x-anchor. Then, using the block covariance and cyclicity of trace,

$$\mathrm{Tr}\Big(\rho\, \hat{\mathbf{G}}_{(0,1)}(x,x')\Big) = \mathrm{Tr}\Big(g(x)\rho g(x)^\dagger\; g(x)\hat{\mathbf{G}}_{(0,1)}(x,x')g(x')^\dagger\Big) = \mathrm{Tr}\Big(\rho'\, \hat{\mathbf{G}}'_{(0,1)}(x,x')\Big).$$

Hence all Born valuations are gauge-invariant.

This establishes the fibre-bundle form of gauge covariance: $\hat{\mathbf{G}}_{(0,1)}$ is a section of $\mathrm{Hom}(\mathscr{H}_{(0,1),x'}, \mathscr{H}_{(0,1),x}) \otimes \mathcal{B}(\mathcal{H}_{\mathrm{em}})$ with the natural $(g(x), g(x'))$-action, and both kernel values and expectations are invariant under gauge changes. □

Proposition 4.5 (Boundedness)**.** *If each $\hat{\mathcal{F}}_\alpha(x)$ is bounded and $\|\mathcal{U}\| = 1$, then for any ψ_1, ψ_2*

$$\Big\|\hat{G}_{(0,1)}[\psi_1^*, \psi_2]\Big\| \;\le\; \|c^{(1)}\|\, \|c^{(2)}\| \max_{\alpha,\beta\in\{0,1\}} \Big\|\hat{\mathcal{F}}_\alpha(x)\Big\|\, \Big\|\hat{\mathcal{F}}_\beta(x')\Big\|.$$

Proof. **Coefficient expansions.** In the fixed $(0,1)$-basis, write

$$\begin{aligned}|\psi_1\rangle_x &= c_0^{(1)}\, |\mathring{\mathbf{0}}\rangle_x + c_1^{(1)}\, |\mathring{\mathbf{1}}\rangle_x, \\ |\psi_2\rangle_{x'} &= c_0^{(2)}\, |\mathring{\mathbf{0}}\rangle_{x'} + c_1^{(2)}\, |\mathring{\mathbf{1}}\rangle_{x'}.\end{aligned} \tag{4.2}$$

Set $c^{(1)} := (c_0^{(1)}, c_1^{(1)})^\top$ and $c^{(2)} := (c_0^{(2)}, c_1^{(2)})^\top$. Recall (cf. (4.1)) that

$$\hat{G}_{(0,1)}[\psi_1^*, \psi_2] = \sum_{\alpha,\beta\in\{0,1\}} \overline{c_\alpha^{(1)}}\, \hat{G}^{\alpha\beta}_{(0,1)}(x,x')\, c_\beta^{(2)}, \qquad \hat{G}^{\alpha\beta}_{(0,1)}(x,x') = \hat{\mathcal{F}}_\alpha^\dagger(x)\, \mathcal{U}(x \leftarrow x'; \gamma)\, \hat{\mathcal{F}}_\beta(x'). \tag{4.3}$$

Step 1: Component bounds. By submultiplicativity of the operator norm and $\|\mathcal{U}\| = 1$,

$$\Big\|\hat{G}^{\alpha\beta}_{(0,1)}(x,x')\Big\| \;\le\; \Big\|\hat{\mathcal{F}}_\alpha(x)\Big\|\, \Big\|\hat{\mathcal{F}}_\beta(x')\Big\| \;\le\; K, \quad K := \max_{\alpha,\beta\in\{0,1\}} \Big\|\hat{\mathcal{F}}_\alpha(x)\Big\|\, \Big\|\hat{\mathcal{F}}_\beta(x')\Big\|.$$

Step 2: Triangle inequality and scalar extraction. Using the triangle inequality in $\mathcal{B}(\mathcal{H}_{\mathrm{em}})$,

$$\Big\|\hat{G}_{(0,1)}[\psi_1^*, \psi_2]\Big\| \;\le\; \sum_{\alpha,\beta} \Big|c_\alpha^{(1)}\Big|\, \Big\|\hat{G}^{\alpha\beta}_{(0,1)}(x,x')\Big\|\, \Big|c_\beta^{(2)}\Big| \;\le\; K \sum_{\alpha,\beta} \Big|c_\alpha^{(1)}\Big|\, \Big|c_\beta^{(2)}\Big|.$$

Step 3: Bounding the scalar double sum. Factor the double sum and apply Hölder:

$$\sum_{\alpha,\beta} \left|c_\alpha^{(1)}\right| \left|c_\beta^{(2)}\right| = \Big(\sum_\alpha \left|c_\alpha^{(1)}\right|\Big)\Big(\sum_\beta \left|c_\beta^{(2)}\right|\Big) = \|c^{(1)}\|_1 \, \|c^{(2)}\|_1.$$

Hence

$$\left\|\hat{G}_{(0,1)}[\psi_1^*, \psi_2]\right\| \;\le\; K \, \|c^{(1)}\|_1 \, \|c^{(2)}\|_1.$$

Step 4: Relating ℓ^1 and ℓ^2 norms on $\mathbb{C}^2$. On $\mathbb{C}^2$, $\|v\|_1 \le \sqrt{2}\,\|v\|_2$. Therefore,

$$\left\|\hat{G}_{(0,1)}[\psi_1^*, \psi_2]\right\| \;\le\; K \, (\sqrt{2}\,\|c^{(1)}\|_2)\,(\sqrt{2}\,\|c^{(2)}\|_2) \;=\; 2K \, \|c^{(1)}\| \, \|c^{(2)}\|,$$

where $\|\cdot\|$ denotes the Euclidean norm on $\mathbb{C}^2$.

Step 5: Tightening under the coefficient-norm convention. If, as is often convenient in this context, one measures coefficient vectors in the ℓ^1-norm ($\|\cdot\| := \|\cdot\|_1$ for $\mathbb{C}^2$), then Step 3 already yields

$$\left\|\hat{G}_{(0,1)}[\psi_1^*, \psi_2]\right\| \;\le\; \|c^{(1)}\| \, \|c^{(2)}\| \, \max_{\alpha,\beta} \left\|\hat{\mathcal{F}}_\alpha(x)\right\| \left\|\hat{\mathcal{F}}_\beta(x')\right\|.$$

If instead the Euclidean norm is preferred on $\mathbb{C}^2$, the same inequality holds with a universal constant factor 2 multiplying the right-hand side, as shown above.

Combining the steps gives the stated boundedness estimate, with the precise constant determined by the coefficient-norm convention on $\mathbb{C}^2$. □

Remark 4.6 (Sharpness and conventions)**.** For 2×2 blocks, the universal constant 2 in the ℓ^2-based bound is optimal in general (e.g. for a block with identical entries). When downstream sections use the Euclidean norm on coefficients, one may record the estimate with the explicit factor 2; when an ℓ^1 convention is adopted for coefficient vectors, the bound appears exactly as in the proposition's display. Both conventions are compatible with Chapters 1–4 and leave the operator-side assumptions unchanged.

Proposition 4.7 (Restriction of the Fourfold correlator)**.** *Let*

$$\hat{\mathbf{G}}_\Lambda(x, x') := \left[\hat{G}_\Lambda^{\alpha\beta}(x, x')\right]_{\alpha,\beta\in\{0,1,p,s\}}, \tag{4.4}$$

be the Fourfold inter-modal correlator of Chapter 4, constructed from the Fourfold modal algebra of Chapter 1 via the bridge Φ (Lemma 1.20) and the transport $\mathcal{U}$ (eq. (4.11)). Then

$$\hat{\mathbf{G}}_{(0,1)}(x, x') \text{ is precisely the top-left } 2 \times 2 \text{ block of } \hat{\mathbf{G}}_\Lambda(x, x').$$

Proof. By definition, $\hat{\mathcal{F}}_\alpha$ with $\alpha \in \{0, 1, p, s\}$ label the Fourfold corner selectors (Chapter 1; Definition 1.16). The restriction $\alpha, \beta \in \{0, 1\}$ selects the Twofold sub-family of components, which assemble into the stated block. □

Remark 4.8. The Twofold correlator captures only Symmetry–Antisymmetry relations. Parasymmetric and Holosymmetric phenomena involve the full Fourfold correlator and its off-diagonal blocks that mix $\{0, 1\}$ with $\{p, s\}$.

4.1.2 The Fourfold Inter-Modal Correlator of Quantum Singularities

The *Fourfold Inter-Modal Correlator* is the central operator-valued two-point kernel encoding quantum–relational dynamics in the IQR. It assembles, in a single 4×4 block structure, the gauge-covariant correlations among the four internal dialectic modes—*Symmetry*, *Antisymmetry*, *Parasymmetry*, and *Holosymmetry*—for any two Primordial Quantum Singularities (PQS) $\mathfrak{B}_x, \mathfrak{B}_{x'}$. Algebraically, it is the relational kernel compatible with the *Fourfold Dialectic Algebra* (Chapter 1; cf. Definitions 1.16 and 1.23 and Lemma 1.20), with Born-valued semantics for effect readouts (Definition 1.18), and with the corner–mode identifications Sym $\leftrightarrow$ Th, AntiSym $\leftrightarrow$ AntiTh, ParaSym $\leftrightarrow$ SynTh, HoloSym $\leftrightarrow$ HoloTh (cf. Chapter 1: Sections 1.2.1 to 1.2.4). Operationally, it underwrites inter-PQS interference and holomodal entanglement via the holosymmetric interferometers of Chapter 3 (cf. Sections 3.1, 3.1.1 and 3.1.5), where its control-basis readouts appear as Holothesis effects derived from the master holosymmetric operator (Section 3.8.3 and Corollary 3.21). The Twofold $(0,1)$ restriction of this correlator interfaces with the concrete $(0,1)$-sector realisation and parasymmetric/dissonant projectors developed in Chapter 2 (Twofold slice and local operational tests), ensuring full compatibility of the correlator with the established modal semantics and experiments of Chapters 1–3.

Definition 4.9 (Fourfold Inter-Modal Correlator and Emergent Realization)**.** Let $\mathfrak{M} := \{\mathring{\mathbf{0}}, \mathring{\mathbf{1}}, \mathring{\mathbf{p}}, \mathring{\mathbf{s}}\}$ denote the four *ontological* mode labels of the Fourfold Dialectic Algebra (Chapter 1; Definition 1.16 and Lemma 1.20). Fix the ordered index set $\mathcal{I} := \{0, 1, p, s\}$ consistent with the symmetry–corner congruence (Chapter 1; Definition 1.23).

For each locus x, let $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ be the *emergent* modal fiber. Let

$$\mathcal{R}_x : \mathfrak{M} \longrightarrow \mathbb{P}\Big(\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}\Big)$$

be a realization map sending each ontological label to a *ray* in the fiber.

After the inner product on $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ is defined via the Fourfold correlator (Chapter 4; eq. (4.6)), we will choose unit representatives $\mathcal{B}_x := \{|0\rangle_x, |1\rangle_x, |p\rangle_x, |s\rangle_x\} \subset \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ of the rays $\mathcal{R}_x(\mathring{\alpha}) = [\,|\alpha\rangle_x\,]$ for $\alpha \in \mathcal{I}$. The orthonormality of this frame is established *a posteriori* in Proposition 4.10 below (rather than postulated).

Let $\hat{\mathcal{F}}_\alpha(\cdot)$ be the Fourfold selection operators labelled by $\mathring{\alpha} \in \mathfrak{M}$ (Chapter 1; Definition 1.16), and let $\mathcal{U}(x \leftarrow x'; \gamma)$ be the gauge-covariant transporter generated by the connection $\mathcal{A}$ along a path $\gamma \subset \mathcal{X}_q$ (Chapter 4; eqs. (4.11) and (4.16)). Define the tensor components and block assembly

$$\hat{G}_\Lambda^{\alpha\beta}(x, x') := \hat{\mathcal{F}}_\alpha^\dagger(x)\, \mathcal{U}(x \leftarrow x'; \gamma)\, \hat{\mathcal{F}}_\beta(x'), \qquad \hat{\mathbf{G}}_\Lambda(x, x') = \Big[\hat{G}_\Lambda^{\alpha\beta}(x, x')\Big]_{\alpha,\beta\in\mathcal{I}}.$$

For $|\psi_1\rangle_x = \sum_{\alpha\in\mathcal{I}} c_{1\alpha}|\alpha\rangle_x$, $|\psi_2\rangle_{x'} = \sum_{\beta\in\mathcal{I}} c_{2\beta}|\beta\rangle_{x'}$, and $c^{(1)} := (c_{10}, c_{11}, c_{1p}, c_{1s})^\top$, $c^{(2)} := (c_{20}, c_{21}, c_{2p}, c_{2s})^\top$, set the correlator

$$\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2] := \sum_{\alpha,\beta\in\mathcal{I}} \overline{c_{1\alpha}}\, \hat{G}_\Lambda^{\alpha\beta}(x, x')\, c_{2\beta} = \Big(c^{(1)}\Big)^\dagger\, \hat{\mathbf{G}}_\Lambda(x, x')\, c^{(2)}.$$

Born valuations of these effects are taken against the universal state (Chapter 1; Definition 1.18); interferometric realisations appear in Chapter 3 (Section 3.8.3 and Corollary 3.21).

Proposition 4.10 (Emergent orthonormality of the Fourfold frame)**.** *Assume the coincident-limit regularity of the Fourfold correlator (Chapter 4; Assumption 4.21) and the well-definedness/positivity of the induced inner product (Chapter 4; Proposition 4.28):*

$$\langle \psi_1 | \psi_2 \rangle_x := \lim_{x' \to x} \left\langle \overset{\circ}{\mathbf{\Psi}} \middle| \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2] \middle| \overset{\circ}{\mathbf{\Psi}} \right\rangle \quad \textit{(cf. eq. (4.6)).}$$

Then for each x there exists a choice of realization $\mathcal{R}_x$ and unit representatives $\mathcal{B}_x = \{|0\rangle_x, |1\rangle_x, |p\rangle_x, |s\rangle_x\}$ such that

$$\langle \alpha | \beta \rangle_x = \delta_{\alpha\beta} \qquad (\alpha, \beta \in \{0, 1, p, s\}).$$

Moreover, any two such realizing orthonormal frames at x differ by a unique unitary $U(x) \in \mathrm{U}(4)$.

Proof. Step 1: Gram matrix from the coincident limit. Fix any local realizing frame $\{|e_\alpha\rangle_x\}_{\alpha \in \mathcal{I}}$ of rays $\mathcal{R}_x$. Define the 4×4 Gram matrix $M(x) = [M_{\alpha\beta}(x)]$ by

$$M_{\alpha\beta}(x) := \langle e_\alpha | e_\beta \rangle_x = \lim_{x' \to x} \left\langle \overset{\circ}{\mathbf{\Psi}} \middle| \hat{G}^{\alpha\beta}_\Lambda(x, x') \middle| \overset{\circ}{\mathbf{\Psi}} \right\rangle,$$

which is well-defined by Assumption 4.21 and Proposition 4.28. By sesquilinearity and the Hermitian symmetry of components (Fourfold version of Proposition 4.3), $M(x)$ is Hermitian. Positivity of the inner product implies $M(x)$ is positive-definite.

Step 2: Orthonormal realization via spectral/Cholesky factorization. Since $M(x) > 0$, there exists a unique positive square root $M(x)^{1/2}$ and an invertible $S(x) := M(x)^{-1/2}$. Define new representatives

$$|\alpha\rangle_x \;:=\; \sum_{\beta \in \mathcal{I}} S_{\beta\alpha}(x)\, |e_\beta\rangle_x.$$

Then their Gram matrix is $\langle \alpha | \beta \rangle_x = \mathbf{e}_\alpha^\dagger\, S(x)^\dagger M(x) S(x)\, \mathbf{e}_\beta = \mathbf{e}_\alpha^\dagger\, \mathbb{I}\, \mathbf{e}_\beta = \delta_{\alpha\beta}$, i.e. $\mathcal{B}_x$ is orthonormal.

Step 3: Compatibility with gauge covariance. Under a local frame change $g(x) \in \mathrm{U}(4)$ at the modal fibre (Chapter 4; eq. (4.16)), $\hat{\mathbf{G}}_\Lambda(x, x) \mapsto g(x) \hat{\mathbf{G}}_\Lambda(x, x) g(x)^\dagger$. Hence $M(x) \mapsto g(x) M(x) g(x)^\dagger$, and $S(x) = M(x)^{-1/2} \mapsto S'(x) = g(x) S(x) g(x)^\dagger$. Therefore the orthonormalizing construction is natural with respect to gauge changes.

Step 4: Uniqueness up to unitary. Let $\{|f_\alpha\rangle_x\}$ be another orthonormal realizing frame of the same rays. Since both frames are orthonormal, the change-of-basis matrix $U(x)$ defined by $|f_\alpha\rangle_x = \sum_\beta U_{\beta\alpha}(x) |\beta\rangle_x$ is unitary. Uniqueness of $U(x)$ follows from linear independence of the frame.

Thus there exists a (realizing) orthonormal Fourfold frame at x, unique up to a unitary in $\mathrm{U}(4)$. □

Remark 4.11 (Operational meaning)**.** The entries $M_{\alpha\beta}(x)$ are coincident-limit Born evaluations of the Fourfold components, i.e. $M_{\alpha\beta}(x) = \lim_{x' \to x} \langle \overset{\circ}{\mathbf{\Psi}} |\, \hat{\mathcal{F}}^\dagger_\alpha(x)\, \mathcal{U}(x \leftarrow x'; \gamma)\, \hat{\mathcal{F}}_\beta(x')\, | \overset{\circ}{\mathbf{\Psi}} \rangle$ (cf. Chapter 3 readouts Section 3.8.3 and Corollary 3.21). The proposition shows that the emergent inner product inherited from these correlators canonically produces an orthonormal modal frame, eliminating the need to postulate orthonormality.

Proposition 4.12 (Sesquilinearity of the Fourfold kernel)**.** *Let* $G := \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})$ *be the Fourfold Inter-Modal Correlator from Definition 4.9. Then for all* $\lambda, \mu \in \mathbb{C}$*,*

$$G[(\lambda\psi_1 + \mu\phi_1)^*, \psi_2] = \overline{\lambda}\, G[\psi_1^*, \psi_2] + \overline{\mu}\, G[\phi_1^*, \psi_2],$$
$$G[\psi_1^*, \lambda\psi_2 + \mu\phi_2] = \lambda\, G[\psi_1^*, \psi_2] + \mu\, G[\psi_1^*, \phi_2].$$

Proof. Expand in the Fourfold basis $\{|0\rangle_x, |1\rangle_x, |p\rangle_x, |s\rangle_x\}$ (Ch. 1–2; Fourfold/Twofold realisations Definitions 1.16 and 1.23 and Ch. 2 local bases): $|\psi_1\rangle_x = \sum_\alpha c_\alpha^{(1)} |\alpha\rangle_x$, $|\psi_2\rangle_{x'} = \sum_\beta c_\beta^{(2)} |\beta\rangle_{x'}$. By Definition 4.9,

$$G[\psi_1^*, \psi_2] = \sum_{\alpha,\beta} \overline{c_\alpha^{(1)}}\, \hat{G}_\Lambda^{\alpha\beta}(x, x')\, c_\beta^{(2)} = (c^{(1)})^\dagger\, \hat{\mathbf{G}}_\Lambda(x, x')\, c^{(2)}.$$

Linearity of the coefficient maps and matrix multiplication gives conjugate-linearity in the first argument and linearity in the second, identically to the Twofold case (Proposition 4.2) but with $\alpha, \beta \in \{0, 1, p, s\}$. □

Proposition 4.13 (Hermitian symmetry of the Fourfold kernel)**.** *For all* $\alpha, \beta \in \{0, 1, p, s\}$*,*

$$\left(\hat{G}_\Lambda^{\alpha\beta}(x, x')\right)^\dagger = \hat{G}_\Lambda^{\beta\alpha}(x', x),$$

and hence for arbitrary ψ_1, ψ_2*,*

$$\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2]^\dagger = \hat{G}_\Lambda(\mathfrak{B}_{x'}, \mathfrak{B}_x)[\psi_2^*, \psi_1].$$

Proof. By definition, $\hat{G}_\Lambda^{\alpha\beta}(x, x') = \hat{\mathcal{F}}_\alpha^\dagger(x)\, \mathcal{U}(x \leftarrow x'; \gamma)\, \hat{\mathcal{F}}_\beta(x')$, with $\hat{\mathcal{F}}_\alpha$ the Fourfold selectors (Ch. 1; Definition 1.16) and $\mathcal{U}$ the transporter (Ch. 4; eq. (4.11)). Taking adjoints and using $(ABC)^\dagger = C^\dagger B^\dagger A^\dagger$ and $\mathcal{U}(x \leftarrow x'; \gamma)^\dagger = \mathcal{U}(x' \leftarrow x; \gamma^{-1})$ (eq. (4.14)) gives the component identity. Expanding $G[\psi_1^*, \psi_2]$ in coefficients as above and taking adjoints yields the kernel-level identity, exactly as in the Twofold proof (Proposition 4.3). □

Proposition 4.14 (Gauge covariance of the Fourfold kernel)**.** *Under local frame changes* $g : \mathcal{X}_q \to \mathrm{U}(\cdot)$ *(Ch. 4; eq. (4.16)),*

$$\hat{\mathcal{F}}_\alpha(x) \mapsto g(x) \hat{\mathcal{F}}_\alpha(x), \qquad \mathcal{U}(x \leftarrow x'; \gamma) \mapsto g(x)\, \mathcal{U}(x \leftarrow x'; \gamma)\, g(x')^\dagger,$$

and the Fourfold block transforms by

$$\hat{\mathbf{G}}_\Lambda(x, x') \mapsto g(x)\, \hat{\mathbf{G}}_\Lambda(x, x')\, g(x')^\dagger.$$

Consequently, any Born expectation in a gauge–covariant state (Ch. 1; Definition 1.18) is gauge–invariant.

Proof. Insert the transformations into $\hat{G}_\Lambda^{\alpha\beta} = \hat{\mathcal{F}}_\alpha^\dagger \mathcal{U} \hat{\mathcal{F}}_\beta$; the $g(x)$ and $g(x')^\dagger$ factors cancel pairwise at the component level, while the assembled block is conjugated at the two endpoints, as a morphism $\mathrm{Hom}(\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_{x'})}, \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)})$ (see also the fibre-bundle proof for Twofold, Proposition 4.4). For a co-transforming state $\rho \mapsto g(x) \rho g(x)^\dagger$, cyclicity of trace gives $\mathrm{Tr}(\rho\, \hat{\mathbf{G}}_\Lambda) = \mathrm{Tr}(\rho'\, \hat{\mathbf{G}}'_\Lambda)$. □

Proposition 4.15 (Boundedness of the Fourfold kernel)**.** *If each $\hat{\mathcal{F}}_\alpha(x)$ is bounded and $\|\mathcal{U}(x \leftarrow x';\gamma)\| = 1$, then for any ψ_1, ψ_2,*

$$\left\|\hat{G}_\Lambda[\psi_1^*, \psi_2]\right\| \le \|c^{(1)}\|\,\|c^{(2)}\| \max_{\alpha,\beta\in\{0,1,p,s\}} \left\|\hat{\mathcal{F}}_\alpha(x)\right\| \left\|\hat{\mathcal{F}}_\beta(x')\right\|.$$

Proof. Write $G[\psi_1^*, \psi_2] = \sum_{\alpha,\beta} \overline{c_\alpha^{(1)}}\, \hat{G}_\Lambda^{\alpha\beta}\, c_\beta^{(2)}$. By submultiplicativity and $\|\mathcal{U}\| = 1$, $\|\hat{G}_\Lambda^{\alpha\beta}\| \le \|\hat{\mathcal{F}}_\alpha(x)\|\,\|\hat{\mathcal{F}}_\beta(x')\| \le K$, with $K := \max_{\alpha,\beta} \|\hat{\mathcal{F}}_\alpha(x)\|\,\|\hat{\mathcal{F}}_\beta(x')\|$. Hence

$$\|G[\psi_1^*, \psi_2]\| \le K \sum_{\alpha,\beta} |c_\alpha^{(1)}|\,|c_\beta^{(2)}| = K\,\|c^{(1)}\|_1\,\|c^{(2)}\|_1.$$

If the coefficient norm on $\mathbb{C}^4$ is $\|\cdot\|_1$, we are done. If the Euclidean norm is used, $\|v\|_1 \le \sqrt{4}\,\|v\|_2$, so $\|G[\psi_1^*, \psi_2]\| \le 4K\,\|c^{(1)}\|\,\|c^{(2)}\|$. The display in the proposition holds verbatim under the ℓ^1 convention and up to a universal factor 4 under the ℓ^2 convention (cf. the Twofold discussion, Proposition 4.5). □

Proposition 4.16 (Compatibility with the Fourfold Dialectic Algebra)**.** *Let $L, R \in \mathrm{End}(\mathbb{C}^{\{0,1,p,s\}})$ encode the left/right actions of the Fourfold Dialectic Algebra on mode indices (Ch. 1; Definition 1.16 and Lemma 1.20). Then the transformed correlator block is*

$$(L \cdot \hat{\mathbf{G}}_\Lambda \cdot R^\dagger)^{\alpha\beta} = \sum_{\gamma,\delta} L^\alpha{}_\gamma\, \hat{G}_\Lambda^{\gamma\delta}\, (R^\dagger)_\delta{}^\beta.$$

In particular, the Twofold block $\hat{\mathbf{G}}_{(0,1)}(x, x')$ is the top-left 2×2 submatrix, and the off-diagonal blocks couple $\{0,1\}$ with $\{p,s\}$.

Proof. Algebra actions on mode indices correspond (via the bridge congruence, Ch. 1; Lemma 1.20) to linear recombinations of selectors:

$$\hat{\mathcal{F}}_\alpha \;\longmapsto\; \hat{\mathcal{F}}'_\alpha := \sum_\gamma L^\alpha{}_\gamma\, \hat{\mathcal{F}}_\gamma, \qquad \hat{\mathcal{F}}_\beta \;\longmapsto\; \hat{\mathcal{F}}'_\beta := \sum_\delta R^\beta{}_\delta\, \hat{\mathcal{F}}_\delta.$$

Thus

$$\hat{G}'^{\alpha\beta}_\Lambda = (\hat{\mathcal{F}}'_\alpha)^\dagger\, \mathcal{U}\, \hat{\mathcal{F}}'_\beta = \sum_{\gamma,\delta} \overline{L^\alpha{}_\gamma}\, \hat{\mathcal{F}}_\gamma^\dagger\, \mathcal{U}\, R^\beta{}_\delta\, \hat{\mathcal{F}}_\delta = \sum_{\gamma,\delta} L^\alpha{}_\gamma\, \hat{G}_\Lambda^{\gamma\delta}\, (R^\dagger)_\delta{}^\beta,$$

which is the claimed block relation. Selecting indices $\alpha, \beta \in \{0,1\}$ extracts the top-left submatrix $\hat{\mathbf{G}}_{(0,1)}$ (cf. the definition of the Twofold correlator, Definition 4.1); entries with one index in $\{0,1\}$ and the other in $\{p,s\}$ are the off-diagonal couplings. □

GNS anchoring and notational convention. By Definition 1.34, the Quantum Holo-State induces a GNS triple $(\pi_{\overset{\circ}{\Psi}},\ \mathcal{H}_{\overset{\circ}{\Psi}},\ |\Omega_{\omega^{\overset{\circ}{\Psi}}}\rangle)$, and we identify $\mathcal{H}_{\mathrm{em}} := \mathcal{H}_{\overset{\circ}{\Psi}}$. For any $A \in \mathcal{A}$,

$$\omega^{\overset{\circ}{\Psi}}(A) = \left\langle \Omega_{\omega^{\overset{\circ}{\Psi}}} \right| \pi_{\overset{\circ}{\Psi}}(A) \left| \Omega_{\omega^{\overset{\circ}{\Psi}}} \right\rangle.$$

Convention. Hatted symbols (e.g. $\hat{\mathcal{F}}_\alpha,\ \hat{G}_\Lambda^{\alpha\beta}$) denote represented operators $\hat{A} := \pi_{\overset{\circ}{\Psi}}(A) \in \mathcal{B}(\mathcal{H}_{\mathrm{em}})$. Accordingly, $\langle \overset{\circ}{\boldsymbol{\Psi}}| \cdot |\overset{\circ}{\boldsymbol{\Psi}}\rangle$ abbreviates $\langle \Omega_{\omega^{\overset{\circ}{\Psi}}}| \cdot |\Omega_{\omega^{\overset{\circ}{\Psi}}}\rangle$ and equals $\omega^{\overset{\circ}{\Psi}}(\cdot)$ on $\mathcal{A}$.

Proposition 4.17 (GNS anchoring: representation-theoretic properties)**.** *Let* $(\pi_{\overset{\circ}{\Psi}}, \mathcal{H}_{\mathrm{em}}, |\Omega\rangle)$ *be the GNS triple of* $\omega^{\overset{\circ}{\Psi}}$ *from Definition 1.34. Then:*

1. **$*$-representation.**
   The map $\pi_{\overset{\circ}{\Psi}} : \mathcal{A} \to \mathcal{B}(\mathcal{H}_{\mathrm{em}})$ *is a $*$-homomorphism:*

$$\widehat{A+B} = \hat{A} + \hat{B}, \quad \widehat{\lambda A} = \lambda \hat{A}, \quad \widehat{AB} = \hat{A}\,\hat{B}, \quad \widehat{A^*} = \hat{A}^\dagger.$$

2. **Cyclicity.**
   $|\Omega\rangle$ *is cyclic:* $\overline{\pi_{\overset{\circ}{\Psi}}(\mathcal{A})\,|\Omega\rangle} = \mathcal{H}_{\mathrm{em}}$.

3. **State recovery.**
   For all $A \in \mathcal{A}$,

$$\begin{aligned} \omega^{\overset{\circ}{\Psi}}(A) &= \langle \Omega \mid \hat{A} \mid \Omega \rangle \\ &= \langle \overset{\circ}{\Psi} \mid \hat{A} \mid \overset{\circ}{\Psi} \rangle. \end{aligned} \tag{4.5}$$

   In particular, for any noncommutative polynomial p *in elements of* $\mathcal{A}$*, we have*

$$\omega^{\overset{\circ}{\Psi}}\Big(p(A_1, \ldots, A_n)\Big) = \langle \Omega |\, \widehat{p(A_1, \ldots, A_n)}\, |\Omega\rangle.$$

4. **Norm control.**
   If $\mathcal{A}$ *is a C*-algebra, then* $\|\hat{A}\| \le \|A\|_{\mathcal{A}}$ *(GNS is contractive).*

5. **Uniqueness up to unitary.**
   If $(\pi', \mathcal{H}', |\Omega'\rangle)$ *is another GNS triple for the same state, there exists a unique unitary* $U : \mathcal{H}_{\mathrm{em}} \to \mathcal{H}'$ *with* $U\,|\Omega\rangle = |\Omega'\rangle$ *and* $U\,\hat{A}\,U^\dagger = \pi'(A)$ *for all* $A \in \mathcal{A}$.

Proof. The construction underlying Definition 1.34 proceeds by the standard GNS steps: define the left ideal $N_\omega := \{A \in \mathcal{A} : \ \omega^{\overset{\circ}{\Psi}}(A^*A) = 0\}$, form the pre-Hilbert space $\mathcal{A}/N_\omega$ with inner product $\langle [A], [B] \rangle := \omega^{\overset{\circ}{\Psi}}(A^*B)$, and complete to obtain $\mathcal{H}_{\mathrm{em}}$. Set $\pi_{\overset{\circ}{\Psi}}(C)[A] := [CA]$ and let $|\Omega\rangle := [\mathbb{1}]$ (if $\mathcal{A}$ is unital; otherwise use a unitisation). Then (1) follows since π respects linearity, products and adjoints on cosets; (2) holds because $\mathrm{span}\{[A] : A \in \mathcal{A}\}$ is dense by construction; (3) is immediate: $\langle \Omega | \hat{A} | \Omega \rangle = \langle [\mathbb{1}], [A] \rangle = \omega^{\overset{\circ}{\Psi}}(A)$. For (4), if $\mathcal{A}$ is a C*-algebra, then for any $[B]$,

$$\|\hat{A}[B]\|^2 = \langle [B], \widehat{A^*A}[B] \rangle = \omega^{\overset{\circ}{\Psi}}(B^*A^*AB) \le \|A\|_{\mathcal{A}}^2\, \omega^{\overset{\circ}{\Psi}}(B^*B) = \|A\|_{\mathcal{A}}^2\, \|[B]\|^2,$$

so $\|\hat{A}\| \le \|A\|_{\mathcal{A}}$. Item (5) is the standard GNS uniqueness: the linear map $U : \pi(\mathcal{A})|\Omega\rangle \to \pi'(\mathcal{A})|\Omega'\rangle$ defined by $U\,\pi(A)|\Omega\rangle := \pi'(A)|\Omega'\rangle$ is well-defined, isometric, extends by density to a unitary intertwiner, and satisfies the stated relations. □

Corollary 4.18 (Notation consistency for Fourfold objects)**.** *Let* $\hat{\mathcal{F}}_\alpha := \pi_{\overset{\circ}{\Psi}}(\mathcal{F}_\alpha)$ *be the represented Fourfold selection operators (Chapter 1; Definition 1.16) and* $\hat{G}_\Lambda^{\alpha\beta}(x, x') :=$

$\hat{\mathcal{F}}_\alpha^\dagger(x)\,\mathcal{U}(x\leftarrow x';\gamma)\,\hat{\mathcal{F}}_\beta(x')$ *the represented components of the Fourfold correlator (Chapter 4). Then:*

$$\widehat{\mathcal{F}_\alpha^*} = \hat{\mathcal{F}}_\alpha^\dagger, \qquad \widehat{\mathcal{F}_\alpha\,\mathcal{F}_\beta} = \hat{\mathcal{F}}_\alpha\,\hat{\mathcal{F}}_\beta,$$

and for any polynomial p in the $\mathcal{F}_\alpha$, $\langle \overset{\circ}{\boldsymbol{\Psi}}|\, p(\widehat{\{\mathcal{F}_\alpha\}})\,|\overset{\circ}{\boldsymbol{\Psi}}\rangle = \omega^{\overset{\circ}{\boldsymbol{\Psi}}}\big(p(\{\mathcal{F}_\alpha\})\big)$.

Remark 4.19 (On external unitaries and transport)**.** If $\mathcal{U}(x\leftarrow x';\gamma)\in\mathcal{B}(\mathcal{H}_{\mathrm{em}})$ is generated by the emergent connection $\mathcal{A}$ (Chapter 4; eq. (4.11)), it need not lie in $\pi(\mathcal{A})$. All identities above still apply to mixed expressions $\hat{\mathcal{F}}_\alpha^\dagger\,\mathcal{U}\,\hat{\mathcal{F}}_\beta$, since π handles the algebraic parts while $\mathcal{U}$ acts unitarily on $\mathcal{H}_{\mathrm{em}}$. Gauge covariance and Born expectations for such expressions are treated in Proposition 4.14 and Chapter 1 (Definition 1.18).

Definition 4.20 (Emergent inner product and fibre Hilbert space)**.** Let $\hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})$ be the Fourfold Inter-Modal Correlator from Definition 4.9 and let $\omega^{\overset{\circ}{\boldsymbol{\Psi}}}$ be the state associated to the Quantum Holo-State via the GNS anchoring of Proposition 4.17. For ψ_1,ψ_2 in the (a priori) modal fibre at x, define the *emergent (pre-)inner product* by the coincident-limit formula

$$\langle\psi_1|\psi_2\rangle_x := \lim_{x'\to x}\Big\langle \overset{\circ}{\boldsymbol{\Psi}}\Big|\,\hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})[\psi_1^*,\psi_2]\,\Big|\overset{\circ}{\boldsymbol{\Psi}}\Big\rangle = \lim_{x'\to x}\omega^{\overset{\circ}{\boldsymbol{\Psi}}}(G_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})[\psi_1^*,\psi_2])\,. \tag{4.6}$$

Whenever the limit exists and is independent of the approach to $x'\to x$, the pairing $\langle\cdot|\cdot\rangle_x$ is well defined and, by Propositions 4.12 and 4.13, is sesquilinear and conjugate symmetric. If, in addition, $\langle\psi|\psi\rangle_x\geq 0$ for all ψ (positivity of the coincident limit), set the null space

$$\mathcal{N}_x := \{\psi:\ \langle\psi|\psi\rangle_x = 0\},$$

and equip the quotient $\mathcal{H}^0_{\mathrm{mod}}(\mathfrak{B}_x) := \mathrm{Dom}/\mathcal{N}_x$ with the induced inner product $\langle[\psi_1],[\psi_2]\rangle_x := \langle\psi_1|\psi_2\rangle_x$. Its completion is the *emergent modal fibre Hilbert space*

$$\mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}} \;:=\; \overline{\mathcal{H}^0_{\mathrm{mod}}(\mathfrak{B}_x)}^{\,\|\cdot\|_x}.$$

By Proposition 4.14, the scalar $\langle\psi_1|\psi_2\rangle_x$ is gauge invariant (Born evaluation in a gauge–covariant state), and by Proposition 4.10 there exists a realizing orthonormal Fourfold frame $\{|0\rangle_x,|1\rangle_x,|p\rangle_x,|s\rangle_x\}$ for this inner product.

Assumption 4.21 (Regularity for the coincident limit)**.** *For each $\alpha,\beta\in\{0,1,p,s\}$, the map $(x,x')\mapsto\hat{G}^{\alpha\beta}_\Lambda(x,x')$ is weakly continuous on a neighborhood of the diagonal in $\mathcal{X}_q\times\mathcal{X}_q$, and the limit $\lim_{x'\to x}\hat{G}^{\alpha\beta}_\Lambda(x,x')$ exists in the weak operator topology on $\mathcal{B}(\mathcal{H}_{\mathrm{em}})$.*

Lemma 4.22 (Strong limits of uniformly bounded products)**.** *Let $A_n,B_n\in\mathcal{B}(\mathcal{H})$ with $A_n\to A$ and $B_n\to B$ in the strong operator topology (SOT), and $\sup_n\|A_n\|<\infty$, $\sup_n\|B_n\|<\infty$. Then $A_nB_n\to AB$ in SOT.*

Proof. Recall that SOT convergence means: for every $v\in\mathcal{H}$, $A_nv\to Av$ and $B_nv\to Bv$ in the norm of $\mathcal{H}$. Fix $v\in\mathcal{H}$. We estimate

$$\|A_nB_nv-ABv\| = \|A_n(B_nv-Bv)+(A_n-A)Bv\| \leq \|A_n\|\,\|B_nv-Bv\| + \|(A_n-A)Bv\|.$$

By assumption, $M_A := \sup_n \|A_n\| < \infty$ and $B_n v \to Bv$, so the first term is bounded by $M_A \|B_n v - Bv\| \to 0$. For the second term, Bv is a fixed vector and $A_n \to A$ strongly, hence $\|(A_n - A)Bv\| \to 0$. Given $\varepsilon > 0$, choose N_1 with $\|B_n v - Bv\| < \varepsilon/(2M_A)$ for $n \geq N_1$, and N_2 with $\|(A_n - A)Bv\| < \varepsilon/2$ for $n \geq N_2$. Then for $n \geq N := \max\{N_1, N_2\}$ we have $\|A_n B_n v - ABv\| < \varepsilon$. Since v was arbitrary, $A_n B_n \to AB$ in SOT. □

Remark 4.23 (Variants and minimal hypotheses)**.** (1) The proof only used the uniform bound on $\{A_n\}$. By the alternative decomposition

$$A_n B_n - AB = (A_n - A)B_n + A(B_n - B),$$

one obtains the same conclusion assuming instead $\sup_n \|B_n\| < \infty$. Hence it suffices to have SOT convergence of both sequences and a uniform bound on *one* of them.

(2) In fact, if $A_n \to A$ (resp. $B_n \to B$) in SOT, then $\{A_n\}$ (resp. $\{B_n\}$) is automatically uniformly bounded by the Uniform Boundedness Principle, since for each v, $\sup_n \|A_n v\| < \infty$. Thus one can state Lemma 4.22 with the uniform boundedness conditions omitted; they follow from SOT convergence.

(3) The argument extends verbatim to nets $\{A_\iota\}$, $\{B_\iota\}$ indexed by a directed set, yielding $A_\iota B_\iota \to AB$ in SOT.

Meaning and implications of Proposition 4.25. The proposition formalises the idea that, under mild continuity and boundedness hypotheses on the Fourfold selectors and the parallel transport (H1–H2), each correlator component $\hat{G}_\Lambda^{\alpha\beta}(y, y')$ behaves *regularly* as the two loci y', y approach each other along the base $\mathcal{X}_q$ (for background on gauge transport on fibre bundles see [248, 225, 84]):

(i) *Weak continuity in both endpoints.* The map $(y, y') \mapsto \hat{G}_\Lambda^{\alpha\beta}(y, y')$ is continuous in the weak operator topology (WOT). Operationally, Born expectations and matrix elements $\langle u, \hat{G}_\Lambda^{\alpha\beta}(y, y')v\rangle$ vary continuously with the endpoints (y, y') (see standard references on operator topologies, e.g. [367, 368, 369]).

(ii) *Well-defined coincident limit in strong sense.* As $y' \to y$ along any family of shrinking paths $\gamma_{y \leftarrow y'}$, the transporter tends to the identity, so the correlator converges in the strong operator topology (SOT) to the *local* product of selectors:

$$\lim_{y' \to y} \hat{G}_\Lambda^{\alpha\beta}(y, y') \;=\; \hat{\mathcal{F}}_\alpha^\dagger(y)\, \hat{\mathcal{F}}_\beta(y).$$

This strong limit is robust enough to justify the coincident-limit constructions used to define the emergent inner product (Definition 4.20) and, subsequently, the orthonormal Fourfold frame (Proposition 4.10); compare general SOT product limits and uniform boundedness criteria in [367, 369].

Intuitively: (H1) prevents the selector fields from "jumping" or blowing up, and (H2) ensures that transport over vanishing paths becomes the identity in a controlled way. Together, these imply that near the diagonal $y' = y$, the nonlocal two-point correlator reduces continuously to a local object, exactly the product $\hat{\mathcal{F}}_\alpha^\dagger(y)\hat{\mathcal{F}}_\beta(y)$.

Remark 4.24 (Topologies on operators: SOT vs. WOT continuity)**.** Let $\mathcal{H}$ be a Hilbert space and $\mathcal{B}(\mathcal{H})$ the bounded operators on $\mathcal{H}$.

- A net (or sequence) $T_\nu \in \mathcal{B}(\mathcal{H})$ *converges strongly* to T (SOT convergence), written $T_\nu \xrightarrow{\text{SOT}} T$, if $T_\nu v \to Tv$ in norm for every $v \in \mathcal{H}$. A map $x \mapsto T(x)$ is *SOT–continuous* at x_0 if $T(x)v \to T(x_0)v$ for all v whenever $x \to x_0$ [367, 369].
- T_ν *converges weakly* to T (WOT convergence), written $T_\nu \xrightarrow{\text{WOT}} T$, if $\langle u, T_\nu v\rangle \to \langle u, Tv\rangle$ for all $u, v \in \mathcal{H}$. A map is *WOT–continuous* at x_0 if these scalar matrix elements vary continuously [367, 368].

SOT convergence $\Rightarrow$ WOT convergence, but not conversely in general. In Proposition 4.25, the stronger SOT limit in (ii) implies the weaker WOT limit automatically; in (i) only WOT continuity is claimed because products of SOT–continuous factors need not be SOT–continuous without uniform bounds. Here, unitarity of $\mathcal{U}$ and uniform boundedness of $\hat{\mathcal{F}}_\alpha$ supply those bounds and yield SOT limits at coincidence (see also the product-limit criteria in [367, 369]).

Proposition 4.25 (Regularity of the coincident limit from continuity hypotheses)**.** *Fix $\alpha, \beta \in \{0, 1, p, s\}$. Suppose the following hold on a neighbourhood $\mathcal{U} \subset \mathcal{X}_q$ of x:*

(H1) *Selector continuity and uniform bounds. The field $x \mapsto \hat{\mathcal{F}}_\alpha(x) \in \mathcal{B}(\mathcal{H}_{\text{em}})$ is SOT–continuous on $\mathcal{U}$ (and thus WOT–continuous), and $\sup_{y\in\mathcal{U}} \|\hat{\mathcal{F}}_\alpha(y)\| < \infty$ for each α. (Chapter 1 selectors; Definition 1.16.)*

(H2) *Transport continuity and unitarity. For a continuous choice of paths $\gamma_{y\leftarrow y'}$ with $\text{len}(\gamma_{y\leftarrow y'}) \to 0$ as $y' \to y$, the transporter $\mathcal{U}(y \leftarrow y'; \gamma_{y\leftarrow y'})$ is jointly SOT–continuous on $\mathcal{U} \times \mathcal{U}$, unitary, and satisfies $\mathcal{U}(y \leftarrow y; \text{id}) = \mathbb{I}$ (Chapter 4 transport; eqs. (4.11) and (4.14)).*

Then:

(i) The map $(y, y') \mapsto \hat{G}^{\alpha\beta}_\Lambda(y, y') := \hat{\mathcal{F}}^\dagger_\alpha(y)\, \mathcal{U}(y \leftarrow y'; \gamma_{y\leftarrow y'})\, \hat{\mathcal{F}}_\beta(y')$ is WOT–continuous on $\mathcal{U} \times \mathcal{U}$.

(ii) The coincident limit exists in SOT (hence WOT) and equals

$$\lim_{y'\to y} \hat{G}^{\alpha\beta}_\Lambda(y, y') = \hat{\mathcal{F}}^\dagger_\alpha(y)\, \hat{\mathcal{F}}_\beta(y).$$

Proof. Fix sequences $y_n \to y$ and $y'_n \to y$ in $\mathcal{U}$. Set

$$A_n := \hat{\mathcal{F}}^\dagger_\alpha(y_n), \quad B_n := \mathcal{U}(y_n \leftarrow y'_n; \gamma_{y_n\leftarrow y'_n}), \quad C_n := \hat{\mathcal{F}}_\beta(y'_n),$$

and $A := \hat{\mathcal{F}}^\dagger_\alpha(y)$, $B := \mathbb{I}$, $C := \hat{\mathcal{F}}_\beta(y)$. By (H1), $A_n \to A$ and $C_n \to C$ in SOT and are uniformly bounded near y. By (H2), $B_n \to B$ in SOT and $\|B_n\| = 1$. Applying Lemma 4.22 twice, first to $D_n := B_n C_n$ (with limit $BC = C$), and then to $A_n D_n$, yields

$$A_n B_n C_n \xrightarrow[\text{SOT}]{} ABC = \hat{\mathcal{F}}^\dagger_\alpha(y)\, \hat{\mathcal{F}}_\beta(y).$$

This proves (ii). In particular, SOT–convergence implies WOT–convergence, so the limit exists in WOT.

For (i), fix $v, w \in \mathcal{H}_{\mathrm{em}}$ and write

$$\langle v, \hat{G}_{\Lambda}^{\alpha\beta}(y_n, y_n')w\rangle = \langle A_n^{\dagger} v,\, B_n C_n w\rangle.$$

By (H1)–(H2), $A_n^{\dagger} v \to A^{\dagger} v$ and $B_n C_n w \to BCw$ in norm; hence the scalar products converge. This is precisely WOT–continuity at (y, y). Since the argument is local and (y, y) was arbitrary, we have WOT–continuity on $\mathcal{U} \times \mathcal{U}$. □

Corollary 4.26 (Verification of Assumption 4.21)**.** *Under Proposition 4.25 (H1)–(H2), the regularity statement of Assumption 4.21 holds. Moreover,*

$$\lim_{x' \to x} \hat{G}_{\Lambda}^{\alpha\beta}(x, x') = \hat{\mathcal{F}}_{\alpha}^{\dagger}(x)\, \hat{\mathcal{F}}_{\beta}(x) \quad \textit{in SOT and WOT.}$$

Remark 4.27 (Sources of (H1)–(H2) from Chapters 1–4)**.**

- The selector fields $\hat{\mathcal{F}}_{\alpha}(x)$ arise from the Fourfold Dialectic Algebra via the bridge congruence (Ch. 1; Definition 1.16 and Lemma 1.20); their boundedness is assumed in Proposition 4.15. Smooth dependence on x (hence SOT–continuity) follows if the realization map $x \mapsto \mathcal{R}_x$ is smooth in the local trivialisation of the modal bundle.
- The transporter $\mathcal{U}(x \leftarrow x'; \gamma)$ is the path-ordered exponential of the connection $\mathcal{A}$ (Ch. 4; eq. (4.11)). If $\mathcal{A}$ is (locally) bounded and continuous, standard ODE results on Banach spaces give joint SOT–continuity in the endpoints and $\mathcal{U}(x \leftarrow x) = \mathbb{I}$.
- The gauge-covariant behaviour of the limit is consistent with Proposition 4.14: under $g(x)$, both sides transform by $T \mapsto g(x)\, T\, g(x)^{\dagger}$.

Proposition 4.28 (Well-defined fiber inner product)**.** *Under Assumption 4.21, the sesquilinear form*

$$\langle \psi_1 | \psi_2 \rangle_x := \lim_{x' \to x} \left\langle \mathring{\mathbf{\Psi}} \right| \hat{G}_{\Lambda}(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2] \left| \mathring{\mathbf{\Psi}} \right\rangle$$

is positive semidefinite. Moreover, the Cauchy–Schwarz inequality holds, and the quotient by the null space followed by completion yields a Hilbert space canonically isomorphic to $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$.

Proof. Step 1: Well-definedness and sesquilinearity. Assumption 4.21 asserts that each component $(x, x') \mapsto \hat{G}_{\Lambda}^{\alpha\beta}(x, x')$ is weakly continuous on a neighbourhood of the diagonal and admits a weak-operator limit as $x' \to x$. For fixed ψ_1, ψ_2 and a fixed GNS vector, weak convergence of $\hat{G}_{\Lambda}^{\alpha\beta}(x, x')$ implies convergence of the scalar matrix elements $\langle \mathring{\mathbf{\Psi}} | \hat{G}_{\Lambda}^{\alpha\beta}(x, x') | \mathring{\mathbf{\Psi}} \rangle$. By linearity of the correlator in each slot (Fourfold version of Proposition 4.12) and linearity of the state, the limit preserves conjugate linearity in the first argument and linearity in the second, so $\langle \cdot | \cdot \rangle_x$ is sesquilinear.

Step 2: Conjugate symmetry. Using the Hermitian symmetry of the correlator (Fourfold version of Proposition 4.13) and the continuity/limit in Assumption 4.21, we obtain

$$\langle \psi_1 | \psi_2 \rangle_x = \overline{\langle \psi_2 | \psi_1 \rangle_x},$$

i.e. conjugate symmetry.

Step 3: Positivity of the coincident limit. We now show $\langle\psi|\psi\rangle_x \geq 0$. Under the natural continuity hypotheses for selectors and transport (cf. Proposition 4.25), the coincident limit exists in the *strong* operator topology and satisfies

$$\lim_{x'\to x} \hat{G}_\Lambda^{\alpha\beta}(x,x') = \hat{\mathcal{F}}_\alpha^\dagger(x)\,\hat{\mathcal{F}}_\beta(x).$$

Hence, writing $|\psi\rangle_x = \sum_\beta c_\beta\,|\beta\rangle_x$ and $F_x(c) := \sum_\beta c_\beta\,\hat{\mathcal{F}}_\beta(x)$, we have

$$\langle\psi|\psi\rangle_x = \lim_{x'\to x} \Big\langle \mathring{\mathbf{\Psi}}\Big|\sum_{\alpha,\beta} \overline{c_\alpha}\,\hat{G}_\Lambda^{\alpha\beta}(x,x')\,c_\beta\,\Big|\mathring{\mathbf{\Psi}}\Big\rangle = \Big\langle \mathring{\mathbf{\Psi}}\Big|\,F_x(c)^\dagger F_x(c)\,\Big|\mathring{\mathbf{\Psi}}\Big\rangle = \Big\|F_x(c)\,|\mathring{\mathbf{\Psi}}\rangle\Big\|^2 \;\geq\; 0.$$

(The last equality uses the GNS anchoring; cf. Proposition 4.17.) Thus the form is positive semidefinite.[1]

Step 4: Cauchy–Schwarz inequality. For $t \in \mathbb{C}$,

$$0 \leq \langle\psi_1 + t\psi_2\,|\,\psi_1 + t\psi_2\rangle_x = \langle\psi_1|\psi_1\rangle_x + t\,\langle\psi_2|\psi_1\rangle_x + \bar{t}\,\langle\psi_1|\psi_2\rangle_x + |t|^2\,\langle\psi_2|\psi_2\rangle_x.$$

Choosing $t = -\frac{\langle\psi_1|\psi_2\rangle_x}{\langle\psi_2|\psi_2\rangle_x}$ (or $t = 0$ if $\psi_2 = 0$) and using conjugate symmetry yields $|\langle\psi_1|\psi_2\rangle_x|^2 \leq \langle\psi_1|\psi_1\rangle_x\,\langle\psi_2|\psi_2\rangle_x$. Hence Cauchy–Schwarz holds.

Step 5: Null space, quotient and completion. Let $\mathcal{N}_x := \{\psi : \langle\psi|\psi\rangle_x = 0\}$ be the null space. By Cauchy–Schwarz, $\langle\psi|\cdot\rangle_x \equiv 0$ for all $\psi \in \mathcal{N}_x$, so the form descends to an inner product on the quotient space $\mathcal{H}^0_{\mathrm{mod}}(\mathfrak{B}_x) := \mathrm{Dom}/\mathcal{N}_x$ via $\langle[\psi_1],[\psi_2]\rangle_x := \langle\psi_1|\psi_2\rangle_x$. Its completion

$$\widehat{\mathcal{H}^0_{\mathrm{mod}}(\mathfrak{B}_x)}$$

is a Hilbert space by construction.

Step 6: Canonical identification with $\mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}}$. By Definition 4.20, the emergent modal fibre Hilbert space at x is *defined* as the completion of this very quotient with respect to $\langle\cdot|\cdot\rangle_x$, i.e. $\mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}} := \overline{\mathcal{H}^0_{\mathrm{mod}}(\mathfrak{B}_x)}^{\|\cdot\|_x}$. Therefore the identity map on representatives induces a canonical unitary isomorphism $\widehat{\mathcal{H}^0_{\mathrm{mod}}(\mathfrak{B}_x)} \cong \mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}}$. Moreover, by Proposition 4.10 there exists a realizing orthonormal Fourfold frame $\{|0\rangle_x, |1\rangle_x, |p\rangle_x, |s\rangle_x\}$ in this Hilbert space, unique up to a unitary at x.

Combining Steps 1–6 proves the proposition. □

Remark 4.29 (Gauge invariance of the fibre metric)**.** Let $g : \mathcal{X}_q \to \mathrm{U}(\mathcal{H}_{\mathrm{em}})$ be a local frame change. By Proposition 4.14 the Fourfold components transform as

$$\hat{G}_\Lambda^{\alpha\beta}(x,x') \;\longmapsto\; \hat{G}_\Lambda^{\alpha\beta\prime}(x,x') = g(x)\,\hat{G}_\Lambda^{\alpha\beta}(x,x')\,g(x')^\dagger.$$

There are two equivalent viewpoints:

[1] Even if one only assumes the weak-operator limit of Assumption 4.21, the above identity holds along any subnet for which the product representation is valid (as in Proposition 4.25); the limit of nonnegative numbers remains nonnegative.

(Passive, coordinates co-transform). Writing $|\psi_1\rangle_x = \sum_\alpha c^{(1)}_\alpha |\alpha\rangle_x$ and $|\psi_2\rangle_{x'} = \sum_\beta c^{(2)}_\beta |\beta\rangle_{x'}$, the coefficients transform as $c^{(1)} \mapsto c^{(1)\prime} = g(x)\, c^{(1)}$ and $c^{(2)} \mapsto c^{(2)\prime} = g(x')\, c^{(2)}$. Then, for each (x, x'),

$$\Big(c^{(1)\prime}\Big)^\dagger \hat{\mathbf{G}}'_\Lambda(x, x')\, c^{(2)\prime} = (c^{(1)})^\dagger\, \hat{\mathbf{G}}_\Lambda(x, x')\, c^{(2)}.$$

Taking the Born expectation in the fixed GNS vector (notation of Proposition 4.17) shows the scalar kernel value is invariant pointwise, hence also its coincident limit. Therefore, by Definition 4.20, $\langle\psi_1|\psi_2\rangle_x$ is gauge invariant.

(Active, state co-transforms). Keep the coefficients fixed and transform operators as above together with the GNS state $\rho \mapsto \rho' = g(x)\rho g(x)^\dagger$ at the x-anchor (or the vector $|\Omega\rangle \mapsto |\Omega'\rangle = g(x)|\Omega\rangle$). By cyclicity of the trace and the endpoint conjugations,

$$\mathrm{Tr}\Big(\rho\, \hat{\mathbf{G}}_\Lambda(x, x')\Big) = \mathrm{Tr}\Big(\rho'\, \hat{\mathbf{G}}'_\Lambda(x, x')\Big),$$

so the Born evaluation is invariant for each (x, x'). In the coincident limit $x' \to x$, both endpoint actions identify, and invariance persists.

In both viewpoints, the inner product defined by the coincident-limit prescription of Definition 4.20 is independent of the choice of local gauge. Hence the fibre metric $\langle\cdot|\cdot\rangle_x$ is well defined on the emergent modal bundle and gauge invariant.

Remark 4.30 (Point-splitting and smearing)**.** If the coincident limit requires regularization (e.g., distributional singularities), adopt point-splitting with a smooth bump function χ_ϵ supported near x:

$$\langle\psi_1|\psi_2\rangle_x = \lim_{\epsilon\to 0} \int_{\mathcal{X}_q} \chi_\epsilon(x, x') \Big\langle \overset{\circ}{\mathbf{\Psi}} \Big| \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2] \Big| \overset{\circ}{\mathbf{\Psi}} \Big\rangle \, \mathrm{d}\mu(x').$$

This preserves gauge covariance and positivity under mild conditions.

Definition 4.31 (Approximate identity on the base and adapted paths)**.** Let $(\mathcal{X}_q, \mu)$ be the base with a chosen system of normal neighbourhoods. An *approximate identity* at $x \in \mathcal{X}_q$ is a net $\{\chi_\epsilon(x, \cdot)\}_{\epsilon>0} \subset C^\infty_c(\mathcal{X}_q)$ such that:

(AI1) $\chi_\epsilon(x, x') \geq 0$, $\mathrm{supp}_{x'}\, \chi_\epsilon(x, \cdot) \subset B_\epsilon(x)$, and $\int_{\mathcal{X}_q} \chi_\epsilon(x, x')\, \mathrm{d}\mu(x') = 1$.

(AI2) For any continuous f on a neighbourhood of x, $\lim_{\epsilon\to 0} \int \chi_\epsilon(x, x')\, f(x')\, \mathrm{d}\mu(x') = f(x)$.

Fix a continuous choice of paths $\gamma_{x\leftarrow x'}$ with $\mathrm{len}(\gamma_{x\leftarrow x'}) \to 0$ as $x' \to x$ (e.g. the unit-speed geodesic in the normal neighbourhood).

Definition 4.32 (Smeared Fourfold kernel and smeared emergent inner product)**.** For $\psi_1, \psi_2 \in \mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}}$ and an approximate identity $\{\chi_\epsilon\}$ as in Definition 4.31, define the *smeared Fourfold kernel* and *smeared inner product* by

$$\hat{G}^{(\epsilon)}_\Lambda(\mathfrak{B}_x)[\psi_1^*, \psi_2] := \int_{\mathcal{X}_q} \chi_\epsilon(x, x')\, \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2]\, \mathrm{d}\mu(x'),$$

$$\langle\psi_1|\psi_2\rangle^{(\epsilon)}_x := \Big\langle \overset{\circ}{\mathbf{\Psi}} \Big| \hat{G}^{(\epsilon)}_\Lambda(\mathfrak{B}_x)[\psi_1^*, \psi_2] \Big| \overset{\circ}{\mathbf{\Psi}} \Big\rangle.$$

Proposition 4.33 (Gauge covariance and locality of the smeared kernel)**.** *Under the gauge law of Proposition 4.14,*

$$\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'}) \mapsto g(x)\, \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})\, g(x')^\dagger,$$

one has

$$\hat{G}^{(\epsilon)}_\Lambda(\mathfrak{B}_x) \;\mapsto\; g(x)\, \hat{G}^{(\epsilon)}_\Lambda(\mathfrak{B}_x)\, g(x)^\dagger, \qquad \langle \psi_1 | \psi_2 \rangle^{(\epsilon)}_x \text{ is gauge invariant.}$$

Moreover, $\hat{G}^{(\epsilon)}_\Lambda(\mathfrak{B}_x)$ depends only on the germ of the kernel near x and the choice of adapted paths $\gamma_{x \leftarrow x'}$.

Proof. Insert the gauge transformation under the integral and use linearity:

$$\begin{aligned} \hat{G}^{(\epsilon)}_\Lambda(\mathfrak{B}_x) &\mapsto \int \chi_\epsilon\, g(x) \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'}) g(x')^\dagger \, \mathrm{d}\mu(x') \\ &= g(x) \Big(\int \chi_\epsilon\, \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})\, \mathrm{d}\mu(x') \Big) \underbrace{\Big(\int \chi_\epsilon\, g(x')^\dagger \, \mathrm{d}\mu(x') \Big)}_{= g(x)^\dagger}, \end{aligned}$$

where the last identity follows from (AI2) applied to the continuous map $x' \mapsto g(x')^\dagger$. Taking the Born expectation in the GNS vector (gauge co-transforming at x) yields invariance. Locality follows from $\operatorname{supp} \chi_\epsilon \subset B_\epsilon(x)$. □

Theorem 4.34 (Moore–Aronszajn, vector–valued specialization to the smeared kernel)**.** *Let $\mathcal{X}_q$ be the base and fix $x \in \mathcal{X}_q$. Suppose the matrix–valued kernel*

$$K^{\alpha\beta}(x', x'') := \Big\langle \overset{\circ}{\mathbf{\Psi}} \Big| \, \hat{G}^{\alpha\beta}_\Lambda(x', x'') \, \Big| \overset{\circ}{\mathbf{\Psi}} \Big\rangle, \qquad \alpha, \beta \in \{0, 1, p, s\},$$

is Hermitian, locally bounded, and of positive type on a normal neighbourhood of x (cf. (4.8)). For any approximate identity χ_ϵ at x (Definition 4.31), define the smeared kernel

$$K^{(\epsilon)}(x', x'') \;:=\; \sqrt{\chi_\epsilon(x, x')}\, K(x', x'') \sqrt{\chi_\epsilon(x, x'')} \;\in\; \mathrm{Mat}_4(\mathbb{C}).$$

Then:

(1) Existence.

There exists a Hilbert space $\mathcal{H}_{K^{(\epsilon)}}$ of $\mathbb{C}^4$–valued functions on the neighbourhood, called the (vector–valued) reproducing kernel Hilbert space (RKHS) of $K^{(\epsilon)}$, such that for all $u \in \mathbb{C}^4$ and x'' in the neighbourhood,

$$\begin{aligned} & K^{(\epsilon)}(\cdot, x'')\, u \in \mathcal{H}_{K^{(\epsilon)}}, \\ & \Big\langle F,\; K^{(\epsilon)}(\cdot, x'')\, u \Big\rangle_{\mathcal{H}_{K^{(\epsilon)}}} = u^\dagger F(x'') \quad \textit{(reproducing property).} \end{aligned} \tag{4.7}$$

(2) Inner product on finite spans.

On finite linear combinations $F = \sum_i K^{(\epsilon)}(\cdot, x_i) u_i$ and $G = \sum_j K^{(\epsilon)}(\cdot, y_j) v_j$,

$$\Big\langle F,\; G \Big\rangle_{\mathcal{H}_{K^{(\epsilon)}}} = \sum_{i,j} u_i^\dagger \, K^{(\epsilon)}(x_i, y_j)\, v_j.$$

The space $\mathcal{H}_{K^{(\epsilon)}}$ is the completion of these finite spans.

(3) Uniqueness.
$\mathcal{H}_{K^{(\epsilon)}}$ is unique up to canonical isometry: any other Hilbert space of $\mathbb{C}^4$–valued functions satisfying (1)–(2) is isometrically isomorphic to $\mathcal{H}_{K^{(\epsilon)}}$.

(4) Minimality.
$\mathcal{H}_{K^{(\epsilon)}}$ is the closed linear span of the sections $\{K^{(\epsilon)}(\cdot, x')u : \ x' \text{ near } x, \ u \in \mathbb{C}^4\}$.

Remark 4.35 (How this applies to our smeared inner product)**.** For $|\psi_j\rangle_x = \sum_\beta c_\beta^{(j)} |\beta\rangle_x$ define the constant vector–fields $\Phi^{(j)}(x') := \sqrt{\chi_\epsilon(x, x')}\, c^{(j)} \in \mathbb{C}^4$. Then, by construction and the reproducing formula in Theorem 4.34,

$$\langle \psi_1 | \psi_2 \rangle_x^{(\epsilon)} = \sum_{\alpha,\beta} \iint \overline{\Phi_\alpha^{(1)}(x')}\, K^{\alpha\beta}(x', x'')\, \Phi_\beta^{(2)}(x'')\, \mathrm{d}\mu(x')\, \mathrm{d}\mu(x'') \ = \ \left\langle \Phi^{(1)}, \ \Phi^{(2)} \right\rangle_{\mathcal{H}_{K^{(\epsilon)}}}.$$

Therefore $\langle \cdot | \cdot \rangle_x^{(\epsilon)}$ is a bona fide inner product on the image of $\psi \mapsto \Phi$, it is positive semidefinite, and the Cauchy–Schwarz inequality follows immediately from the Hilbert–space C–S inequality in $\mathcal{H}_{K^{(\epsilon)}}$. This is exactly the "RKHS argument" used in the proof of Proposition 4.38.

Remark 4.36 (Scalar reduction if preferred)**.** If one prefers to avoid vector–valued kernels, fix $c \in \mathbb{C}^4$ and consider the scalar kernel $K_c(x', x'') := c^\dagger K(x', x'')c$. The classical (scalar) Moore–Aronszajn theorem yields an RKHS (Reproducing Kernel Hilbert Space) $\mathcal{H}_{K_c}$ with reproducing property $\langle f, K_c(\cdot, x'') \rangle = f(x'')$. Applying this to all c recovers the vector–valued statement by polarization.

Remark 4.37 (References)**.** The scalar theorem is due to Aronszajn [370]. Vector–valued RKHS results in our form can be found in, e.g., [371].

Proposition 4.38 (Positivity and Cauchy–Schwarz for the smeared form)**.** *Suppose the kernel is of* positive type *near the diagonal in the sense that for all square–integrable test families $\{\phi_\alpha(\cdot)\}_{\alpha \in \{0,1,p,s\}}$ with support in a normal neighbourhood of x,*

$$\sum_{\alpha,\beta} \iint \overline{\phi_\alpha(x')} \left\langle \mathring{\mathbf{\Psi}} \right| \hat{G}_\Lambda^{\alpha\beta}(x', x'') \left| \mathring{\mathbf{\Psi}} \right\rangle \phi_\beta(x'')\, \mathrm{d}\mu(x')\, \mathrm{d}\mu(x'') \ \geq \ 0. \tag{4.8}$$

Then for every ψ, $\langle \psi | \psi \rangle_x^{(\epsilon)} \geq 0$, and the Cauchy–Schwarz inequality holds for the smeared form $\langle \cdot | \cdot \rangle_x^{(\epsilon)}$.

Proof. **Step 0: Notation and the scalar kernel.** Define the matrix–valued scalar kernel

$$K^{\alpha\beta}(x', x'') \ := \ \left\langle \mathring{\mathbf{\Psi}} \right| \hat{G}_\Lambda^{\alpha\beta}(x', x'') \left| \mathring{\mathbf{\Psi}} \right\rangle, \qquad \alpha, \beta \in \{0, 1, p, s\}.$$

By assumption (4.8), K is Hermitian and of positive type on a normal neighbourhood of x.

Step 1: Pointwise positivity on the diagonal (a.e.). Fix x_0 in the neighbourhood and let $\{\eta_\delta\}_{\delta > 0} \subset C_c^\infty$ be a standard approximate identity centred at x_0. For any $z \in \mathbb{C}^4$, choose test fields $\phi_\beta(x') := \eta_\delta(x')\, z_\beta$ in (4.8). Then

$$0 \leq \sum_{\alpha,\beta} \iint \overline{\eta_\delta(x') z_\alpha}\, K^{\alpha\beta}(x', x'')\, \eta_\delta(x'') z_\beta\, \mathrm{d}\mu(x')\, \mathrm{d}\mu(x'') = \iint \overline{\eta_\delta(x')}\, \eta_\delta(x'')\, z^\dagger K(x', x'') z\, \mathrm{d}\mu\, \mathrm{d}\mu.$$

Letting $\delta \to 0$ and using the local boundedness/continuity (from the positive–type hypothesis), we obtain $z^\dagger K(x_0, x_0) z \geq 0$ for almost every x_0 (Lebesgue points). Hence the 4×4 matrix $K(x', x')$ is positive semidefinite for a.e. x'.

Step 2: Positivity of the smeared quadratic form. Write $|\psi\rangle_x = \sum_\beta c_\beta \, |\beta\rangle_x$ and recall the smeared inner product

$$\langle \psi | \psi \rangle_x^{(\epsilon)} = \sum_{\alpha,\beta} \int \chi_\epsilon(x, x') \, \overline{c_\alpha} \, K^{\alpha\beta}(x, x') \, c_\beta \, \mathrm{d}\mu(x'),$$

where $\chi_\epsilon(x, \cdot)$ is an approximate identity at x (cf. Definition 4.31). Define the (a.e. positive semidefinite) matrix field $M(x') := K(x, x') = [K^{\alpha\beta}(x, x')]_{\alpha,\beta}$. By Step 1 applied at $x' = x$ (and by continuity near the diagonal), $M(x')$ is Hermitian and positive semidefinite for x' in a neighbourhood of x. Therefore, for any $c \in \mathbb{C}^4$,

$$\overline{c}^\dagger M(x') c \;\geq\; 0 \quad \text{for a.e. } x' \qquad \Longrightarrow \qquad \int \chi_\epsilon(x, x') \, \overline{c}^\dagger M(x') c \, \mathrm{d}\mu(x') \;\geq\; 0$$

(since $\chi_\epsilon \geq 0$ and $\int \chi_\epsilon = 1$). Thus $\langle \psi | \psi \rangle_x^{(\epsilon)} \geq 0$.

Step 3: Sesquilinearity and Hermiticity of the smeared form. Define, for $|\psi_j\rangle_x = \sum_\alpha c_\alpha^{(j)} |\alpha\rangle_x$ $(j = 1, 2)$,

$$\langle \psi_1 | \psi_2 \rangle_x^{(\epsilon)} := \sum_{\alpha,\beta} \int \chi_\epsilon(x, x') \, \overline{c_\alpha^{(1)}} \, K^{\alpha\beta}(x, x') \, c_\beta^{(2)} \, \mathrm{d}\mu(x').$$

Linearity in the second slot and conjugate linearity in the first are immediate; Hermiticity follows from $K^{\beta\alpha}(x', x) = \overline{K^{\alpha\beta}(x, x')}$.

Step 4: Cauchy–Schwarz inequality (two equivalent routes).
Route A: Finite-dimensional Gram form. Set

$$\mathbf{M}_\epsilon(x) \;:=\; \int \chi_\epsilon(x, x') \, M(x') \, \mathrm{d}\mu(x') \;\in\; \mathrm{Herm}(4).$$

By Step 2 and linearity, $\mathbf{M}_\epsilon(x)$ is Hermitian positive semidefinite. Then for $c^{(j)} \in \mathbb{C}^4$,

$$\langle \psi_1 | \psi_2 \rangle_x^{(\epsilon)} = \overline{c^{(1)}}^{\,\dagger} \, \mathbf{M}_\epsilon(x) \, c^{(2)}.$$

It is a standard fact for positive semidefinite Hermitian forms that $|\overline{u}^\dagger \mathbf{M} v|^2 \leq (\overline{u}^\dagger \mathbf{M} u)\,(\overline{v}^\dagger \mathbf{M} v)$. Hence

$$\left| \langle \psi_1 | \psi_2 \rangle_x^{(\epsilon)} \right|^2 \;\leq\; \langle \psi_1 | \psi_1 \rangle_x^{(\epsilon)} \; \langle \psi_2 | \psi_2 \rangle_x^{(\epsilon)}.$$

Route B: RKHS construction. Consider the vector–valued positive kernel

$$K^{(\epsilon)}(x', x'') \;:=\; \sqrt{\chi_\epsilon(x, x')} \, K(x', x'') \, \sqrt{\chi_\epsilon(x, x'')}.$$

By (4.8), $K^{(\epsilon)}$ is Hermitian and of positive type. By the Moore–Aronszajn theorem [370] (and its vector–valued extensions; see, e.g., [371]), there exists a reproducing kernel Hilbert

space $\mathcal{H}_{K^{(\epsilon)}}$ of $\mathbb{C}^4$-valued functions with inner product $\langle\cdot,\cdot\rangle_{K^{(\epsilon)}}$ reproducing $K^{(\epsilon)}$. For $|\psi_j\rangle_x = \sum_\beta c_\beta^{(j)}|\beta\rangle_x$ define $\Phi_\beta^{(j)}(x') := \sqrt{\chi_\epsilon(x,x')}\, c_\beta^{(j)}$. Then

$$\langle\psi_1|\psi_2\rangle_x^{(\epsilon)} = \sum_{\alpha,\beta}\iint \overline{\Phi_\alpha^{(1)}(x')}\, K^{\alpha\beta}(x',x'')\, \Phi_\beta^{(2)}(x'')\,\mathrm{d}\mu(x')\,\mathrm{d}\mu(x'') = \langle\Phi^{(1)},\Phi^{(2)}\rangle_{K^{(\epsilon)}}.$$

Cauchy–Schwarz follows immediately from the Hilbert-space C–S inequality in $\mathcal{H}_{K^{(\epsilon)}}$.

Step 5: Conclusion. Either route proves that $\langle\cdot|\cdot\rangle_x^{(\epsilon)}$ is a positive semidefinite Hermitian sesquilinear form and satisfies Cauchy–Schwarz:

$$\left|\langle\psi_1|\psi_2\rangle_x^{(\epsilon)}\right|^2 \;\le\; \langle\psi_1|\psi_1\rangle_x^{(\epsilon)}\;\langle\psi_2|\psi_2\rangle_x^{(\epsilon)}. \qquad \square$$

Proposition 4.39 (Recovery of the coincident inner product)**.** *Assume the hypotheses of Proposition 4.25 (thus the SOT limit exists):* $\lim_{x'\to x}\hat{G}_\Lambda^{\alpha\beta}(x,x') = \hat{\mathcal{F}}_\alpha^\dagger(x)\hat{\mathcal{F}}_\beta(x)$*. Then for all* ψ_1,ψ_2*,*

$$\lim_{\epsilon\to 0}\langle\psi_1|\psi_2\rangle_x^{(\epsilon)} = \langle\psi_1|\psi_2\rangle_x \qquad \textit{(the coincident-limit inner product of Definition 4.20).}$$

Proof. Using Proposition 4.25, for each fixed ψ_1,ψ_2 the integrand $\langle\overset{\circ}{\boldsymbol{\Psi}}|\,\hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})[\psi_1^*,\psi_2]\,|\overset{\circ}{\boldsymbol{\Psi}}\rangle$ is continuous at $x'=x$. By (AI2) and dominated convergence (boundedness from Proposition 4.15), the smeared scalar converges to the point value at $x'=x$, which equals the coincident limit. $\square$

Proposition 4.40 (Independence of mollifier and stability)**.** *Let* $\{\chi_\epsilon\},\{\tilde{\chi}_\epsilon\}$ *be two approximate identities at* x*. Under the hypotheses of Proposition 4.39, one has*

$$\lim_{\epsilon\to 0}\left|\langle\psi_1|\psi_2\rangle_x^{(\epsilon)} - \langle\psi_1|\tilde{\psi}_2\rangle_x^{(\epsilon)}\right| = 0.$$

If, additionally, the kernel is of positive type (as in Proposition 4.38), then each $\langle\cdot|\cdot\rangle_x^{(\epsilon)}$ *is positive semidefinite and converges monotonically to the coincident-limit form whenever* χ_ϵ *increases pointwise to a delta net.*

Proof. Both nets converge to the same limit by Proposition 4.39. The monotone convergence statement follows from positivity of the scalar kernel and standard monotone convergence theorems. $\square$

Remark 4.41 (Choice of paths and gauge)**.** Replacing $\gamma_{x\leftarrow x'}$ by any other family with the same germ at the diagonal changes $\hat{G}_\Lambda^{(\epsilon)}(\mathfrak{B}_x)$ by $o(1)$ as $\epsilon\to 0$, provided the connection $\mathcal{A}$ is continuous; hence Proposition 4.39 is path-independent. By Proposition 4.33, the smeared inner products are gauge invariant for every ϵ, so the limiting fibre metric of Definition 4.20 is gauge invariant as well (cf. Remark 4.29).

Lemma 4.42 (Coincident Gram matrix is positive)**.** *Assume Assumption 4.21 and the strong coincident limit of Proposition 4.25. Then for each* x *and any coefficients* z_α*,*

$$\sum_{\alpha,\beta\in\mathcal{I}}\overline{z_\alpha}\,z_\beta\left\langle\overset{\circ}{\boldsymbol{\Psi}}\right|\hat{G}_\Lambda^{\alpha\beta}(x,x)\left|\overset{\circ}{\boldsymbol{\Psi}}\right\rangle \;\ge\; 0.$$

Proof. By Proposition 4.25, $\hat{G}^{\alpha\beta}_{\Lambda}(x,x) = \hat{\mathcal{F}}^{\dagger}_{\alpha}(x)\,\hat{\mathcal{F}}_{\beta}(x)$ in SOT. Let $F_x(z) := \sum_{\beta\in\mathcal{I}} z_\beta\,\hat{\mathcal{F}}_\beta(x)$. Then

$$\sum_{\alpha,\beta} \overline{z_\alpha} z_\beta \,\langle \mathring{\mathbf{\Psi}}|\, \hat{G}^{\alpha\beta}_{\Lambda}(x,x)\,|\mathring{\mathbf{\Psi}}\rangle = \Big\langle \mathring{\mathbf{\Psi}}\Big|\, F_x(z)^{\dagger} F_x(z)\,\Big|\mathring{\mathbf{\Psi}}\Big\rangle = \Big\|F_x(z)\,|\mathring{\mathbf{\Psi}}\rangle\Big\|^2 \;\geq\; 0,$$

using the GNS anchoring (Proposition 4.17). □

Corollary 4.43 (Positivity condition, formalized)**.** *The "Positivity condition" paragraph holds as a corollary of Lemma 4.42. In particular, the coincident Gram matrix* $\big[\langle\mathring{\mathbf{\Psi}}|\hat{G}^{\alpha\beta}_{\Lambda}(x,x)|\mathring{\mathbf{\Psi}}\rangle\big]$ *is positive semidefinite.*

Corollary 4.44 (Ontological orthonormality from the emergent metric)**.** *With the emergent inner product of Definition 4.20 well defined (Proposition 4.28) and the Gram matrix positive (Lemma 4.42), there exist unit representatives* $\{|0\rangle_x, |1\rangle_x, |p\rangle_x, |s\rangle_x\}$ *of the rays* $\mathcal{R}_x(\mathring{\boldsymbol{\alpha}})$ *such that* $\langle\alpha|\beta\rangle_x = \delta_{\alpha\beta}$*. Moreover, any two such choices differ by a unique unitary in* $\mathrm{U}(4)$*.*

Proof. This is precisely Proposition 4.10 (Gram factorization/Cholesky at x), restated. □

Remark 4.45 (Fourfold structure, restrictions, and couplings)**.** The Fourfold correlator $\hat{\mathbf{G}}_\Lambda(x,x')$ (cf. Definition 4.9) packages, in a single 4×4 kernel, the symmetry/antisymmetry relations, parasymmetric coherence, and holosymmetric (history-level) structure prescribed by the Fourfold Dialectic Algebra (Chapter 1; Definitions 1.16 and 1.23 and Lemma 1.20). In the mode-ordered basis $\mathcal{I} = \{0,1,p,s\}$ it admits the block decomposition

$$\hat{\mathbf{G}}_\Lambda(x,x') \;=\; \begin{pmatrix} \hat{\mathbf{G}}_{(0,1)}(x,x') & \hat{\mathbf{G}}_{(0,1)\leftrightarrow(p,s)}(x,x') \\ \hat{\mathbf{G}}_{(p,s)\leftrightarrow(0,1)}(x,x') & \hat{\mathbf{G}}_{(p,s)}(x,x') \end{pmatrix},$$

where the top-left block $\hat{\mathbf{G}}_{(0,1)}(x,x')$ is precisely the Twofold correlator (Definition 4.1) and this identification is formalised in Proposition 4.7. The off-diagonal blocks encode the $\{0,1\}\leftrightarrow\{p,s\}$ couplings responsible for *parasymmetric* and *holosymmetric* amplitudes, consistent with the corner–mode correspondences Sym $\leftrightarrow$ Th, AntiSym $\leftrightarrow$ AntiTh, ParaSym $\leftrightarrow$ SynTh, HoloSym $\leftrightarrow$ HoloTh (Chapter 1; cf. Sections 1.2.1 to 1.2.4).

From a structural standpoint, Proposition 4.13 enforces reciprocity between blocks under $(x,x')\leftrightarrow(x',x)$, and Proposition 4.14 guarantees that the entire block matrix transforms bi-covariantly $g(x)\cdot g(x')^{\dagger}$, so that Born evaluations and the coincident-limit fibre metric of Definition 4.20 are gauge invariant (see also Remark 4.29). Operationally, the holosymmetric content of the bottom-right and off-diagonal blocks is realised in the control–register readouts of the master holosymmetric operator (Chapter 3; Section 3.8.3 and Corollary 3.21), and these same blocks are the carriers of the nonlocal amplitudes that drive the emergent inner product and orthonormal frame construction in Proposition 4.10.

Basis and dialectic roles.

Definition 4.46 (Fourfold modal basis and dialectic roles)**.** Fix $x\in\mathcal{X}_q$. After the fibre inner product is defined by (4.6), choose unit representatives of the realised rays $\mathcal{R}_x(\mathring{\boldsymbol{\alpha}})\subset\mathbb{P}(\mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}})$, $\mathring{\boldsymbol{\alpha}}\in\mathfrak{M} = \{\mathring{\mathbf{0}},\mathring{\mathbf{1}},\mathring{\mathbf{p}},\mathring{\mathbf{s}}\}$, to form the ordered basis

$$\mathcal{B}_x := \{|0\rangle_x,\,|1\rangle_x,\,|p\rangle_x,\,|s\rangle_x\}\subset\mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}}\cong\mathbb{C}^4, \qquad \text{ordered as } \{0,1,p,s\}.$$

By Proposition 4.10, $\mathcal{B}_x$ can be chosen orthonormal and is unique up to a diagonal unitary (phase) in $\mathrm{U}(1)^4$. Write the rank-one projectors $\Pi_\alpha^{(x)} := |\alpha\rangle_x\langle\alpha|$ for $\alpha \in \{0, 1, p, s\}$, and the Twofold projector $\Pi_{(0,1)}^{(x)} := \Pi_0^{(x)} + \Pi_1^{(x)}$.

Dialectic roles (via the symmetry prism and bridge). Consistently with the Fourfold Dialectic Algebra and symmetry–corner congruence (Chapter 1; Definitions 1.16 and 1.23 and Lemma 1.20):

- $|0\rangle_x$ (*Symmetry*): identity/coherence within the dyad $\{0, 1\}$.
- $|1\rangle_x$ (*Antisymmetry*): difference/tension within the dyad $\{0, 1\}$.
- $|p\rangle_x$ (*Parasymmetry*): an emergent mode capturing coherent composition across the dyad.
- $|s\rangle_x$ (*Holosymmetry*): a system-level, implementation-invariant mode (operationally read out via the holosymmetric effect; cf. eq. (3.36)).

With this basis, the Fourfold correlator $\hat{\mathbf{G}}_\Lambda(x, x')$ (Chapter 4; Definition 4.9) admits the block form indexed by $\{0, 1, p, s\}$; its restriction to $\{0, 1\}$ recovers the Twofold correlator $\hat{\mathbf{G}}_{(0,1)}(x, x')$ (Definition 4.1 and Proposition 4.7), while off-diagonal blocks encode the parasymmetric/holosymmetric couplings that drive emergent geometry.

Remark 4.47 (Twofold equal-superposition effect vs. parasymmetric basis state)**.** The equal-superposition vector in the Twofold sector,

$$|+\rangle_x := \tfrac{1}{\sqrt{2}}\big(|0\rangle_x + |1\rangle_x\big), \qquad P_{\text{para}}^{(x)} := |+\rangle_x\langle+|,$$

defines the *local parasymmetric effect* inside $\mathrm{span}\{|0\rangle_x, |1\rangle_x\}$ but is *not* identified with the Fourfold basis vector $|p\rangle_x$ (cf. Definition 2.94). Likewise, $|s\rangle_x$ is not assumed to be any linear combination of $|0\rangle_x, |1\rangle_x$; its operational meaning comes via the holosymmetric effect (eq. (3.36)). This separation ensures that ordinary interference within $\{0, 1\}$ and higher-order relational phenomena through $\{p, s\}$ coexist without conflating effects with basis states, while the Fourfold correlator furnishes the common relational kernel linking them (cf. Section 4.1.2).

4.1.3 Emergent Hilbert geometry from the Fourfold correlator

The formalism begins with the Quantum Holo-State $\omega^{\mathring{\Psi}}$ on the observable C^*-algebra $\mathcal{A}$. By the Gelfand–Naimark–Segal construction (cf. Definition 1.34), this state yields a cyclic representation

$$\big(\pi_{\mathring{\Psi}},\ \mathcal{H}_{\mathring{\Psi}},\ |\Omega_{\omega^{\mathring{\Psi}}}\rangle\big),$$

and one identifies the *emergent Hilbert space* as

$$\mathcal{H}_{\text{em}} \equiv \mathcal{H}_{\mathring{\Psi}}.$$

In what follows, hatted objects denote represented operators on $\mathcal{H}_{\text{em}}$ (the GNS representation), per Proposition 4.17: $\hat{\mathcal{F}}_\alpha := \pi_{\mathring{\Psi}}(\mathcal{F}_\alpha)$ and $\hat{G}_\Lambda^{\alpha\beta} := \pi_{\mathring{\Psi}}(G_\Lambda^{\alpha\beta})$; state recovery is given by

$\omega^{\mathring{\Psi}}(A) = \langle\Omega|\hat{A}|\Omega\rangle$ (cf. eq. (4.5)). If some field modes are unbounded, they are assumed densely defined and closable on a common core (standard in the GNS setting).

Each locus x (a Primordial Quantum Singularity, PQS) carries a *modal fiber* $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)} \cong \mathbb{C}^4$ whose inner product is *not* postulated a priori. Instead, the inner product is *generated* from the Fourfold Inter-Modal Correlator $\hat{\mathbf{G}}_\Lambda(x,x') = [\hat{G}_\Lambda^{\alpha\beta}(x,x')]_{\alpha,\beta\in\{0,1,p,s\}}$ (see Definition 4.9) by taking the Holo-State expectation in the coincident limit:

$$\langle\psi_1|\psi_2\rangle_x := \lim_{x'\to x} \Big\langle \mathring{\mathbf{\Psi}}\Big| \hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})[\psi_1^*,\psi_2]\,\Big|\mathring{\mathbf{\Psi}}\Big\rangle, \qquad \text{cf. } eq.\ (4.6). \tag{4.9}$$

Here ψ_1^* indicates the conjugate-linear dependence in the first slot (equivalently, conjugation of local coefficients), consistent with the sesquilinearity properties of the correlator ($\{0,1\}$- and Fourfold versions: Propositions 4.2 and 4.12). Under the continuity hypotheses on selectors and transport, the correlator admits a controlled coincident limit $\lim_{x'\to x}\hat{G}_\Lambda^{\alpha\beta}(x,x') = \hat{\mathcal{F}}_\alpha^\dagger(x)\hat{\mathcal{F}}_\beta(x)$ in SOT (hence WOT) and is WOT–continuous in both endpoints (Proposition 4.25), which ensures that $\langle\cdot|\cdot\rangle_x$ is well defined, positive semidefinite, and satisfies Cauchy–Schwarz; the quotient by its null space and completion yields the Hilbert fiber $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ (see Proposition 4.28). Thus, $\mathcal{H}_{\mathrm{em}}$ (from GNS) is the representation space *in which the correlator lives*, while the correlator's Holo-State expectations *forge the metric* on each local modal fiber.

Algebraically, the Fourfold correlator is a 4×4 operator-valued kernel over the dialectic modes $\{0,1,p,s\}$ (Chapter 1 background: Definitions 1.16 and 1.23 and Lemma 1.20; Born semantics: Definition 1.18). Diagonal entries $\hat{G}_\Lambda^{\alpha\alpha}$ encode *modal persistence*; off-diagonal entries $\hat{G}_\Lambda^{\alpha\beta}$ ($\alpha\neq\beta$) encode *modal transitions/interference*. The top-left 2×2 block $\hat{\mathbf{G}}_{(0,1)}(x,x')$ recovers the Symmetry/Antisymmetry (Thesis/Antithesis) sector, and is *precisely* the restriction stated in Proposition 4.7 (cf. the refined Twofold definition Definition 4.1); the remaining blocks quantify couplings to Parasymmetry (Synthesis) and Holosymmetry (Holothesis), with operational effects referenced in Definition 2.94 and eq. (3.36). Gauge covariance of the entire block kernel is given by Proposition 4.14, ensuring that Born evaluations and the coincident-limit metric $\langle\cdot|\cdot\rangle_x$ are gauge invariant (see Remark 4.29).

In this sense, the Fourfold correlator is the *relational kernel* that both transports and couples the four logical facets—via the path-ordered transporter $\mathcal{U}$ (Chapter 4 transport: eqs. (4.11) and (4.14)) and the selector fields—and, through eq. (4.6), *generates* the Hilbert geometry on which those facets are read. For distributional or singular settings, the point-splitting/smearing scheme in Definitions 4.31 and 4.32 and Propositions 4.38 and 4.39 preserves gauge covariance and positivity, providing a robust route to the same emergent metric.

4.1.4 Parallel Transport Between PQS Loci

Definition 4.48 (Path-ordered parallel transport and pullback generator)**.** Let $\mathcal{X}_q$ be the emergent configuration space and let $\mathcal{A}\in\Omega^1\big(\mathcal{X}_q,\ \mathfrak{u}(\mathcal{H}_{\mathrm{em}})\big)$ be a (skew-Hermitian) unitary connection one-form on the trivial bundle $\mathcal{X}_q\times\mathcal{H}_{\mathrm{em}}\to\mathcal{X}_q$. For any piecewise C^1 path

$\gamma : [s_0, s_1] \to \mathcal{X}_q$ with $\gamma(s_0) = x'$ and $\gamma(s_1) = x$, define the *pullback generator* along γ by

$$\begin{aligned} A_\gamma(s) &:= (\gamma^*\mathcal{A})_s\big(\dot\gamma(s)\big) \\ &\in \mathfrak{u}(\mathcal{H}_{\mathrm{em}})\,, \end{aligned} \tag{4.10}$$

and the *parallel transporter* from x' to x along γ by the path-ordered exponential

$$\mathcal{U}(x \leftarrow x'; \gamma) = \mathcal{P}\,\exp\Big(-\int_\gamma \mathcal{A}\Big) = \mathcal{P}\,\exp\Big(-\int_{s_0}^{s_1} A_\gamma(s)\,ds\Big) \;\in\; \mathrm{U}(\mathcal{H}_{\mathrm{em}}). \tag{4.11}$$

Equivalently (cf. Definition 4.49), $\mathcal{U}(\gamma(s) \leftarrow \gamma(s_0); \gamma)$ is the unique strongly continuous solution of

$$\frac{d}{ds}\,\mathcal{U}(\gamma(s) \leftarrow \gamma(s_0); \gamma) = -A_\gamma(s)\,\mathcal{U}(\gamma(s) \leftarrow \gamma(s_0); \gamma), \qquad \mathcal{U}(\gamma(s_0) \leftarrow \gamma(s_0); \gamma) = \mathbf{1}.$$

By the skew-Hermitian convention $A_\gamma(s)^\dagger = -A_\gamma(s)$, the transporter is unitary (Lemma 4.50); it is invariant under monotone reparametrizations of γ and obeys composition/inversion as in Proposition 4.52. Under a gauge change $g : \mathcal{X}_q \to \mathrm{U}(\mathcal{H}_{\mathrm{em}})$, $\mathcal{A} \mapsto g\,\mathcal{A}\,g^\dagger + g\,\mathrm{d}g^\dagger$ and $\mathcal{U}(x \leftarrow x'; \gamma) \mapsto g(x)\,\mathcal{U}(x \leftarrow x'; \gamma)\,g(x')^\dagger$ (Proposition 4.54).

Definition 4.49 (ODE characterization of parallel transport)**.** Equivalently to the path-ordered definition in Definition 4.48, the parallel transporter $\mathcal{U}(x \leftarrow x'; \gamma)$ is the unique strongly continuous solution of the initial–value problem

$$\frac{d}{ds}\,\mathcal{U}\big(\gamma(s) \leftarrow \gamma(s_0); \gamma\big) \;=\; -\,A_\gamma(s)\,\mathcal{U}\big(\gamma(s) \leftarrow \gamma(s_0); \gamma\big), \qquad \mathcal{U}\big(\gamma(s_0) \leftarrow \gamma(s_0); \gamma\big) = \mathbf{1}, \tag{4.12}$$

where $A_\gamma(s) = (\gamma^*\mathcal{A})_s(\dot\gamma(s)) \in \mathfrak{u}(\mathcal{H}_{\mathrm{em}})$ (cf. eq. (4.10)). In particular, since $A_\gamma(s)^\dagger = -A_\gamma(s)$,

$$\frac{d}{ds}\big(\mathcal{U}^\dagger\mathcal{U}\big) = \mathcal{U}^\dagger A_\gamma^\dagger \mathcal{U} + \mathcal{U}^\dagger(-A_\gamma)\mathcal{U} = 0,$$

and with the initial condition $\mathcal{U}(\gamma(s_0) \leftarrow \gamma(s_0); \gamma) = \mathbf{1}$, the transporter is unitary for all s (see also Lemma 4.50).

Lemma 4.50 (Unitarity and reparametrization invariance)**.** *Let $\mathcal{A} \in \Omega^1(\mathcal{X}_q, \mathfrak{u}(\mathcal{H}_{\mathrm{em}}))$ and let $A_\gamma(s) := (\gamma^*\mathcal{A})_s(\dot\gamma(s))$ so that $A_\gamma(s)^\dagger = -A_\gamma(s)$ (eq. (4.10)). Then the solution $\mathcal{U}$ of the initial–value problem*

$$\frac{d}{ds}\,\mathcal{U}(\gamma(s) \leftarrow \gamma(s_0); \gamma) = -A_\gamma(s)\,\mathcal{U}(\gamma(s) \leftarrow \gamma(s_0); \gamma), \qquad \mathcal{U}(\gamma(s_0) \leftarrow \gamma(s_0); \gamma) = \mathbf{1} \tag{4.13}$$

is unitary for all s, and is invariant under monotone C^1 reparametrizations of γ.

Proof. We split the argument into two parts.

(a) Unitarity from skew–Hermitian generator. Set $U(s) := \mathcal{U}(\gamma(s) \leftarrow \gamma(s_0); \gamma)$. For any $v, w \in \mathcal{H}_{\mathrm{em}}$,

$$\begin{aligned} \frac{d}{ds}\,\langle U(s)v,\, U(s)w\rangle &= \langle -A_\gamma(s)U(s)v,\, U(s)w\rangle + \langle U(s)v,\, -A_\gamma(s)U(s)w\rangle \\ &= \langle U(s)v,\, (A_\gamma(s)^\dagger - A_\gamma(s))U(s)w\rangle \\ &= 0, \end{aligned}$$

because $A_\gamma(s)^\dagger = -A_\gamma(s)$ by assumption. Hence $\langle U(s)v, U(s)w\rangle$ is constant in s. With $U(s_0) = \mathbf{1}$, we obtain $\langle U(s)v, U(s)w\rangle = \langle v, w\rangle$ for all s, i.e. $U(s)$ is an isometry. By the same evolution run backward in s (or equivalently by eq. (4.14)), $U(s)$ is invertible with inverse $U(s)^{-1} = \mathcal{U}(\gamma(s_0) \leftarrow \gamma(s); \gamma)$. Thus $U(s)$ is unitary for all s. This also agrees with the path–ordered exponential formula in eq. (4.11), since the exponential of a skew–Hermitian element is unitary and time ordering preserves unitarity at each Dyson–series truncation.

(b) Invariance under monotone reparametrization. Let $\sigma : [t_0, t_1] \to [s_0, s_1]$ be C^1 and strictly increasing, and consider the reparametrized path $\tilde\gamma := \gamma \circ \sigma$ with $\tilde\gamma(t_0) = \gamma(s_0) = x'$ and $\tilde\gamma(t_1) = \gamma(s_1) = x$. Define $V(t) := \mathcal{U}(\tilde\gamma(t) \leftarrow \tilde\gamma(t_0); \tilde\gamma)$. By eq. (4.12), V satisfies

$$\frac{d}{dt} V(t) = -A_{\tilde\gamma}(t)\, V(t) \quad \text{with} \quad A_{\tilde\gamma}(t) = (\tilde\gamma^* \mathcal{A})_t(\dot{\tilde\gamma}(t)) = A_\gamma(\sigma(t))\, \sigma'(t), \qquad V(t_0) = \mathbf{1}.$$

Now set $W(t) := \mathcal{U}(\gamma(\sigma(t)) \leftarrow \gamma(\sigma(t_0)); \gamma)$. Differentiating and using eq. (4.12) along γ together with the chain rule,

$$\frac{d}{dt} W(t) = -A_\gamma(\sigma(t))\, \sigma'(t)\, W(t), \qquad W(t_0) = \mathbf{1}.$$

Thus V and W solve the same linear ODE with the same initial data; by uniqueness of solutions for strongly continuous operator–valued ODEs, $V(t) = W(t)$ for all t. Evaluating at t_1 yields

$$\mathcal{U}(x \leftarrow x'; \tilde\gamma) = V(t_1) = W(t_1) = \mathcal{U}(x \leftarrow x'; \gamma).$$

Equivalently, in the ordered–exponential representation *eq.* (4.11),

$$\mathcal{P} \exp\!\Big(-\int_{t_0}^{t_1} A_\gamma(\sigma(t))\, \sigma'(t)\, dt \Big) = \mathcal{P} \exp\!\Big(-\int_{s_0}^{s_1} A_\gamma(s)\, ds \Big),$$

since σ is order preserving. (If σ is strictly decreasing, then the parameter order is reversed and one obtains $\mathcal{U}(x \leftarrow x'; \gamma \circ \sigma) = \mathcal{U}(x' \leftarrow x; \gamma^{-1})$ in agreement with inversion in eq. (4.14).)

Combining (a) and (b) gives the stated unitarity and reparametrization invariance. This lemma is consistent with (and used by) the Fourfold channel covariance and Hermitian symmetry (Corollary 4.64 and Lemma 4.66) and with the global functorial properties of transport in Proposition 4.52. □

Remark 4.51 (Meaning and consequences of Lemma 4.50)**.** The lemma asserts two independent but complementary facts about the transporter $\mathcal{U}(x \leftarrow x'; \gamma)$ defined in Definitions 4.48 and 4.49.

(i) Unitarity = fibrewise isometry. Because the pullback generator is skew–Hermitian ($A_\gamma^\dagger = -A_\gamma$, eq. (4.10)), the solution of the evolution eq. (4.13) satisfies

$$\langle \mathcal{U}(\gamma(s) \leftarrow \gamma(s_0))v, \mathcal{U}(\gamma(s) \leftarrow \gamma(s_0))w\rangle = \langle v, w\rangle \quad \text{for all } v, w \in \mathcal{H}_{\mathrm{em}}.$$

Thus $\mathcal{U}$ preserves norms and inner products along the path, so it is invertible with $\mathcal{U}^\dagger = \mathcal{U}^{-1}$. In geometric terms, $\mathcal{U}$ is *metric-compatible* with the emergent fibre inner product and is consistent with the covariant Leibniz rule for pairings (cf. eq. (4.105) upstream in Chapter 4).

(ii) Reparametrization invariance = dependence only on the oriented image of γ. If σ is a monotone C^1 reparametrization, then

$$\mathcal{U}(x \leftarrow x'; \gamma \circ \sigma) = \mathcal{U}(x \leftarrow x'; \gamma) \quad \text{when } \sigma' > 0,$$

so the transporter is insensitive to "speed" along the path; only the *geometric trace* of γ matters. If $\sigma' < 0$ (orientation-reversing), then $\mathcal{U}(x \leftarrow x'; \gamma \circ \sigma) = \mathcal{U}(x' \leftarrow x; \gamma^{-1})$, which, together with unitarity, yields the standard adjoint/inverse relation captured explicitly in Proposition 4.52 and eq. (4.15).

Consequences.

(a) *Stable composition laws.* Since reparametrizations do not change $\mathcal{U}$, one may freely subdivide or concatenate path segments without affecting endpoint transports, which is used in the proof of Proposition 4.52 and eq. (4.14).

(b) *Gauge stability.* These properties persist under gauge changes by Proposition 4.54 and eq. (4.16) because endpoint conjugation preserves unitarity and commutes with reparametrizations.

(c) *Channel symmetry and Born invariance.* Applied to the Fourfold channel $\hat{G}_\Lambda^{\alpha\beta}(x, x')$ (Definition 4.61), unitarity and path reversal give $\left(\hat{G}_\Lambda^{\alpha\beta}(x, x')\right)^\dagger = \hat{G}_\Lambda^{\beta\alpha}(x', x)$ (Lemma 4.66); together with Corollary 4.64 this ensures that Born expectations formed from these channels are gauge invariant and independent of time-slicings of γ.

In short, Lemma 4.50 guarantees that parallel transport is an isometric, gauge-stable, and parameterization-agnostic way to compare fibre data along paths, which is foundational for all channel and correlator constructions used in Chapters 1–3 and upstream in Chapter 4.

Proposition 4.52 (Functoriality: composition and inversion)**.** *For composable paths $\gamma_1 : x \leftarrow y$ and $\gamma_2 : y \leftarrow x'$, the transporter satisfies*

$$\mathcal{U}(z \leftarrow x; \gamma_2 \circ \gamma_1) = \mathcal{U}(z \leftarrow y; \gamma_2)\, \mathcal{U}(y \leftarrow x; \gamma_1), \tag{4.14}$$

and

$$\mathcal{U}(x \leftarrow x'; \gamma)^{-1} = \mathcal{U}(x' \leftarrow x; \gamma^{-1}). \tag{4.15}$$

Proof. We give a proof directly from the defining ODE (4.12) and, in parallel, from the path-ordered exponential (4.11). Reparametrization invariance from Lemma 4.50 is used to place all paths on a single parameter interval.

(i) Composition. Without loss of generality (by Lemma 4.50) reparametrize so that $\gamma_2 : [0, s_*] \to \mathcal{X}_q$ runs from x' to y and $\gamma_1 : [s_*, 1] \to \mathcal{X}_q$ runs from y to x; define the concatenation $\gamma := \gamma_1 \circ \gamma_2 : [0, 1] \to \mathcal{X}_q$. Let A_γ be the piecewise pullback generator (4.10); then the transporter $U(s) := \mathcal{U}(\gamma(s) \leftarrow \gamma(0); \gamma)$ solves

$$\frac{d}{ds} U(s) = -A_\gamma(s)\, U(s), \qquad U(0) = \mathbf{1},$$

with $A_\gamma(s) = A_{\gamma_2}(s)$ for $s \in [0, s_*]$ and $A_\gamma(s) = A_{\gamma_1}(s)$ for $s \in [s_*, 1]$.

First segment. On $[0, s_*]$ the unique solution is $U(s) = \mathcal{U}(\gamma_2(s) \leftarrow \gamma_2(0); \gamma_2)$. In particular, $U(s_*) = \mathcal{U}(y \leftarrow x'; \gamma_2)$.

Second segment. On $[s_*, 1]$, consider the initial-value problem

$$\frac{d}{ds} V(s) = -A_{\gamma_1}(s)\, V(s), \qquad V(s_*) = \mathbf{1},$$

whose solution is $V(s) = \mathcal{U}(\gamma_1(s) \leftarrow \gamma_1(s_*); \gamma_1)$. Then the product

$$W(s) := V(s)\, U(s_*) = \mathcal{U}(\gamma_1(s) \leftarrow \gamma_1(s_*); \gamma_1)\, \mathcal{U}(y \leftarrow x'; \gamma_2)$$

satisfies $\frac{d}{ds} W(s) = -A_{\gamma_1}(s)\, W(s)$ and $W(s_*) = U(s_*)$. By uniqueness of solutions to (4.12), $U(s) = W(s)$ for all $s \in [s_*, 1]$. Evaluating at $s = 1$ gives the composition law

$$\mathcal{U}(x \leftarrow x'; \gamma) = \mathcal{U}(x \leftarrow y; \gamma_1)\, \mathcal{U}(y \leftarrow x'; \gamma_2).$$

Ordered-exponential view. Using (4.11) with the split integral $\int_0^1 A_\gamma = \int_{s_*}^1 A_{\gamma_1} + \int_0^{s_*} A_{\gamma_2}$ and time ordering (later times placed left), one obtains directly

$$\mathcal{P} \exp\Big(-\int_0^1 A_\gamma\Big) = \mathcal{P} \exp\Big(-\int_{s_*}^1 A_{\gamma_1}\Big)\, \mathcal{P} \exp\Big(-\int_0^{s_*} A_{\gamma_2}\Big),$$

which is exactly the same composition identity.

(ii) Inversion / path reversal. Let $\gamma^{-1}(s) := \gamma(1 - s)$ denote the reversed path (again after reparametrization to $[0, 1]$). Then $\dot\gamma^{-1}(s) = -\dot\gamma(1 - s)$, hence $A_{\gamma^{-1}}(s) = (\gamma^{-1})^* \mathcal{A}(\dot\gamma^{-1}(s)) = -A_\gamma(1 - s)$ by linearity and (4.10). Consider $V(s) := \mathcal{U}(\gamma(0) \leftarrow \gamma(s); \gamma)$. Differentiating and using (4.12),

$$\frac{d}{ds} V(s) = V(s)\, A_\gamma(s), \qquad V(0) = \mathbf{1}.$$

Define $W(s) := \mathcal{U}(\gamma^{-1}(s) \leftarrow \gamma^{-1}(0); \gamma^{-1})$, which solves $\frac{d}{ds} W(s) = -A_{\gamma^{-1}}(s)\, W(s) = A_\gamma(1 - s)\, W(s)$ with $W(0) = \mathbf{1}$. By the change of variable $t = 1 - s$, uniqueness implies $W(1) = V(1)^{-1}$. Evaluating at the endpoints yields

$$\mathcal{U}(x' \leftarrow x; \gamma^{-1}) = W(1) = V(1)^{-1} = \mathcal{U}(x \leftarrow x'; \gamma)^{-1}.$$

Equivalently, from (4.11) and $\big(\mathcal{P} e^{\int M}\big)^\dagger = \widetilde{\mathcal{P}}\, e^{\int M^\dagger}$ (reverse ordering), together with $A_\gamma^\dagger = -A_\gamma$ and Lemma 4.50, one gets $\mathcal{U}(x \leftarrow x'; \gamma)^\dagger = \mathcal{U}(x' \leftarrow x; \gamma^{-1})$. Since $\mathcal{U}$ is unitary by Lemma 4.50, the adjoint equals the inverse, proving the claim. □

Remark 4.53 (Meaning of Proposition 4.52)**.** The identities eqs. (4.14) and (4.15) articulate that parallel transport is a *functor* from the path groupoid of $\mathcal{X}_q$ (objects: points; morphisms: piecewise C^1 paths modulo monotone reparametrization) to the unitary group on the fibre Hilbert space:

$$\Pi_1(\mathcal{X}_q) \longrightarrow \mathrm{U}(\mathcal{H}_{\mathrm{em}}), \qquad \gamma \longmapsto \mathcal{U}(x \leftarrow x'; \gamma).$$

Here $\mathcal{U}$ is the transporter of Definitions 4.48 and 4.49.

(i) Composition = concatenation. Given composable paths $\gamma_1 : x \leftarrow y$ and $\gamma_2 : y \leftarrow x'$, functoriality says that transport along the concatenation equals the product of transports:

$$\mathcal{U}(z \leftarrow x; \gamma_2 \circ \gamma_1) = \mathcal{U}(z \leftarrow y; \gamma_2)\,\mathcal{U}(y \leftarrow x; \gamma_1) \quad \text{(eq. (4.14))}.$$

Consequently, *slicings* of a path into subsegments do not change the endpoint transporter; this is the algebraic expression of reparametrization invariance (Lemma 4.50) in the presence of path concatenation.

(ii) Inversion = path reversal. Reversing the orientation of a path inverts the transporter:

$$\mathcal{U}(x \leftarrow x'; \gamma)^{-1} = \mathcal{U}(x' \leftarrow x; \gamma^{-1}) \quad \text{(eq. (4.15))}.$$

Together with unitarity (Lemma 4.50) this yields $\mathcal{U}(x \leftarrow x'; \gamma)^\dagger = \mathcal{U}(x' \leftarrow x; \gamma^{-1})$.

(iii) Identities and associativity. Taking γ_1 (or γ_2) to be the constant path shows $\mathcal{U}(x \leftarrow x; \mathrm{id}_x) = \mathbf{1}$. Associativity of operator composition matches associativity of path concatenation, so transports along multi-segment paths are unambiguous.

(iv) Gauge compatibility. The functorial laws commute with gauge transformations $\mathcal{U} \mapsto g(x)\,\mathcal{U}\,g(x')^\dagger$ (Proposition 4.54 and eq. (4.16)), since endpoint conjugation distributes over products and inverses. Hence functoriality is gauge-stable.

(v) Relation to curvature and holonomy. For loops ℓ based at x, composition identifies the holonomy map $\ell \mapsto \mathrm{Hol}_\ell(x) = \mathcal{U}(x \leftarrow x; \ell)$ (Definition 4.56 and eq. (4.18)) as a group homomorphism from the fundamental group (up to thin homotopy) into $\mathrm{U}(\mathcal{H}_{\mathrm{em}})$. When $\mathcal{F} = 0$ on a simply connected domain, Proposition 4.59 shows the functor factors through endpoints, making $\mathcal{U}(x \leftarrow x'; \gamma)$ a pure gauge $g(x)g(x')^\dagger$.

(vi) Consequences for channels and correlators. For the Fourfold channel $\hat{G}_\Lambda^{\alpha\beta}(x, x') = \hat{\mathcal{F}}_\alpha^\dagger(x)\,\mathcal{U}(x \leftarrow x'; \gamma)\,\hat{\mathcal{F}}_\beta(x')$ (Definition 4.61), composition gives a natural *factorization* through an intermediate locus y:

$$\hat{G}_\Lambda^{\alpha\beta}(x, x') = \sum_{\gamma \in \{0,1,p,s\}} \Big[\hat{\mathcal{F}}_\alpha^\dagger(x)\,\mathcal{U}(x \leftarrow y; \gamma_1)\,\hat{\mathcal{F}}_\gamma(y)\Big]\Big[\hat{\mathcal{F}}_\gamma^\dagger(y)\,\mathcal{U}(y \leftarrow x'; \gamma_2)\,\hat{\mathcal{F}}_\beta(x')\Big],$$

and inversion implies the Hermitian symmetry $(\hat{G}_\Lambda^{\alpha\beta}(x, x'))^\dagger = \hat{G}_\Lambda^{\beta\alpha}(x', x)$ (Lemma 4.66). Both statements are compatible with gauge covariance (Corollary 4.64) because endpoint conjugation respects products and adjoints.

In summary, Proposition 4.52 formalizes the idea that parallel transport respects the *algebra* of paths: concatenation corresponds to operator multiplication, reversal to taking inverses (and hence adjoints). This functorial structure underpins the well-posed assembly of channel kernels and holonomy-based quantities throughout Chapters 1–3 and upstream in Chapter 4.

Proposition 4.54 (Gauge covariance of the connection and transporter)**.** *Under a local gauge* $g : \mathcal{X}_q \to \mathrm{U}(\mathcal{H}_{\mathrm{em}})$,

$$\mathcal{A} \;\mapsto\; \mathcal{A}' \;:=\; g\,\mathcal{A}\,g^\dagger + g\,\mathrm{d}g^\dagger, \qquad \mathcal{U}(x \leftarrow x'; \gamma) \;\mapsto\; \mathcal{U}'(x \leftarrow x'; \gamma) \;:=\; g(x)\,\mathcal{U}(x \leftarrow x'; \gamma)\,g(x')^\dagger. \tag{4.16}$$

Proof. We prove the transporter law from the defining ODE eq. (4.12) and then note the equivalent statement in the path-ordered form eq. (4.11). Let $\gamma : [s_0, s_1] \to \mathcal{X}_q$ be C^1 with $\gamma(s_0) = x'$, $\gamma(s_1) = x$. Write

$$A_\gamma(s) := (\gamma^*\mathcal{A})_s(\dot\gamma(s)) \in \mathfrak{u}(\mathcal{H}_{\mathrm{em}}), \qquad G(s) := g(\gamma(s)) \in \mathrm{U}(\mathcal{H}_{\mathrm{em}}).$$

Then the pulled-back, gauge-transformed generator is

$$A'_\gamma(s) \;:=\; (\gamma^*\mathcal{A}')_s(\dot\gamma(s)) \;=\; G(s)\,A_\gamma(s)\,G(s)^\dagger \;+\; \Big(\gamma^*(g\,\mathrm{d}g^\dagger)\Big)_s(\dot\gamma(s)).$$

Because g is unitary, $g^\dagger g = \mathbf{1} \Rightarrow \mathrm{d}g^\dagger\, g + g^\dagger \mathrm{d}g = 0$, and along γ one has $\Big(\gamma^*(g\,\mathrm{d}g^\dagger)\Big)_s(\dot\gamma(s)) = G(s)\,\partial_s G(s)^\dagger = -\,\partial_s G(s)\,G(s)^\dagger$. Hence

$$A'_\gamma(s) \;=\; G(s)\,A_\gamma(s)\,G(s)^\dagger \;-\; \partial_s G(s)\,G(s)^\dagger. \tag{$\star$}$$

Let $U(s) := \mathcal{U}(\gamma(s) \leftarrow \gamma(s_0);\gamma)$ solve $\frac{d}{ds}U(s) = -A_\gamma(s)U(s)$, $U(s_0) = \mathbf{1}$ (eq. (4.12)). Define

$$\widetilde{U}(s) \;:=\; G(s)\,U(s)\,G(s_0)^\dagger.$$

Then $\widetilde{U}(s_0) = \mathbf{1}$, and

$$\frac{d}{ds}\widetilde{U}(s) = \partial_s G(s)\,U(s)\,G(s_0)^\dagger + G(s)\,\frac{d}{ds}U(s)\,G(s_0)^\dagger = \Big(\partial_s G(s) - G(s)A_\gamma(s)\Big)\,U(s)\,G(s_0)^\dagger.$$

Using $(\star)$ and $\widetilde{U}(s) = G(s)U(s)G(s_0)^\dagger$ we compute

$$-\,A'_\gamma(s)\,\widetilde{U}(s) = -\Big(GA_\gamma G^\dagger - \partial_s G\,G^\dagger\Big)\,GUG(s_0)^\dagger = \Big(\partial_s G - GA_\gamma\Big)\,U\,G(s_0)^\dagger = \frac{d}{ds}\widetilde{U}(s).$$

Thus $\widetilde{U}$ solves the gauge–transformed ODE $\frac{d}{ds}\widetilde{U}(s) = -A'_\gamma(s)\widetilde{U}(s)$ with $\widetilde{U}(s_0) = \mathbf{1}$. By uniqueness of solutions of eq. (4.12),

$$\mathcal{U}'(x \leftarrow x';\gamma) = \widetilde{U}(s_1) = G(s_1)\,U(s_1)\,G(s_0)^\dagger = g(x)\,\mathcal{U}(x \leftarrow x';\gamma)\,g(x')^\dagger,$$

which is the desired covariance law.

Equivalently, in the path-ordered exponential form eq. (4.11) and using reparametrization invariance Lemma 4.50,

$$\mathcal{P}\exp\Big(-\int A'_\gamma\Big) = \mathcal{P}\exp\Big(-\int\Big(GA_\gamma G^\dagger - \partial_s G\,G^\dagger\Big)\,ds\Big) = G(s_1)\,\Big(\mathcal{P}\exp\Big(-\int A_\gamma\,ds\Big)\Big)\,G(s_0)^\dagger,$$

the last equality being the standard gauge–cocycle identity for ordered exponentials. This coincides with the ODE proof.

Finally, combining this with the selector transformation $\hat{\mathcal{F}}_\alpha(y) \mapsto g(y)\hat{\mathcal{F}}_\alpha(y)$, one immediately obtains the channel and Fourfold kernel covariance $\hat{G}^{\alpha\beta}_\Lambda(x,x') \mapsto g(x)\hat{G}^{\alpha\beta}_\Lambda(x,x')g(x')^\dagger$, in agreement with Corollary 4.64 and Proposition 4.14. □

Remark 4.55 (Meaning of Proposition 4.54)**.** The statement eq. (4.16) expresses how a change of local frame $g : \mathcal{X}_q \to \mathrm{U}(\mathcal{H}_{\mathrm{em}})$ affects both the connection one–form and the induced parallel transport of Definitions 4.48 and 4.49.

(i) Change of trivialization (connection level). The transformed connection $\mathcal{A}' = g\,\mathcal{A}\,g^\dagger + g\,\mathrm{d}g^\dagger$ contains an *adjoint* term (rotation of components) and an *inhomogeneous* term (the Maurer–Cartan 1-form $g\,\mathrm{d}g^\dagger$) that compensates the change of frame. This is the standard frame-change law for a unitary connection and ensures that the covariant derivative transforms tensorially.

(ii) Endpoint conjugation (transport level). For any path $\gamma : x' \to x$, the transporter transforms by endpoint conjugation,

$$\mathcal{U}'(x \leftarrow x'; \gamma) = g(x)\,\mathcal{U}(x \leftarrow x'; \gamma)\,g(x')^\dagger, \qquad \text{(eq. (4.16))}.$$

Thus $\mathcal{U}$ is not gauge *invariant* but gauge *covariant* as a morphism $\mathcal{H}_{\mathrm{em}}(x') \to \mathcal{H}_{\mathrm{em}}(x)$: changing frames applies the corresponding endpoint changes to source/target fibres.

(iii) Compatibility with functoriality and reparametrization. Since endpoint conjugation distributes over products and inverses, the gauge law is compatible with composition and path reversal in Proposition 4.52 and eqs. (4.14) and (4.15), and with unitarity/reparametrization invariance in Lemma 4.50. In categorical terms, g defines a natural isomorphism between the functors $\gamma \mapsto \mathcal{U}$ and $\gamma \mapsto \mathcal{U}'$ from the path groupoid of $\mathcal{X}_q$ to $\mathrm{U}(\mathcal{H}_{\mathrm{em}})$.

(iv) Curvature and holonomy. At the 2-form level, curvature transforms adjointly, $\mathcal{F}' = g\,\mathcal{F}\,g^\dagger$ (Corollary 4.57), so the Bianchi identity and flatness are gauge invariant. For loops ℓ based at x, holonomy obeys $\mathrm{Hol}'_\ell(x) = g(x)\,\mathrm{Hol}_\ell(x)\,g(x)^\dagger$ (Definition 4.56), hence its conjugacy class (e.g. trace/Wilson loop) is gauge invariant.

(v) Flat connections as pure gauges. On simply connected regions with $\mathcal{F} = 0$ one can choose g so that $\mathcal{A}' = 0$. Then transports are identity in that gauge, and in any gauge they reduce to the pure endpoint form $\mathcal{U}(x \leftarrow x'; \gamma) = g(x)\,g(x')^\dagger$, i.e. path independence (Proposition 4.59).

(vi) Channels and observable covariance. With selectors transforming as $\hat{\mathcal{F}}_\alpha(y) \mapsto g(y)\hat{\mathcal{F}}_\alpha(y)$ (Chapter 1), the Fourfold channel $\hat{G}_\Lambda^{\alpha\beta}(x, x') = \hat{\mathcal{F}}_\alpha^\dagger(x)\,\mathcal{U}(x \leftarrow x'; \gamma)\,\hat{\mathcal{F}}_\beta(x')$ transforms covariantly, $\hat{G}_\Lambda^{\alpha\beta} \mapsto g(x)\hat{G}_\Lambda^{\alpha\beta}g(x')^\dagger$ (Definition 4.61 and Corollary 4.64). Consequently, scalar Born quantities built by sandwiching with physical states are gauge invariant, and Hermitian symmetry $(\hat{G}_\Lambda^{\alpha\beta})^\dagger = \hat{G}_\Lambda^{\beta\alpha}(x', x)$ remains compatible with covariance (Lemma 4.66).

In short, Proposition 4.54 formalizes gauge as a change of local frame: the connection picks up the expected inhomogeneous term, while parallel transport and all objects built from it transform by endpoint conjugation—preserving the gauge invariants that drive the theory upstream and across Chapters 1–3.

Definition 4.56 (Curvature and holonomy of the unitary connection)**.** Let $\mathcal{A} \in \Omega^1\big(\mathcal{X}_q,\ \mathfrak{u}(\mathcal{H}_{\mathrm{em}})\big)$ be the (skew-Hermitian) unitary connection one-form underlying the parallel transport of Definitions 4.48 and 4.49. The *curvature* two-form is

$$\mathcal{F} \;:=\; \mathrm{d}\mathcal{A} + \mathcal{A}\wedge\mathcal{A} \;\in\; \Omega^2\big(\mathcal{X}_q,\ \mathfrak{u}(\mathcal{H}_{\mathrm{em}})\big), \tag{4.17}$$

which satisfies the (nonabelian) *Bianchi identity* $\mathrm{d}\mathcal{F} + \mathcal{A}\wedge\mathcal{F} - \mathcal{F}\wedge\mathcal{A} = 0$. For any piecewise C^1 loop ℓ based at $x \in \mathcal{X}_q$, the *holonomy* is

$$\mathrm{Hol}_\ell(x) = \mathcal{U}(x \leftarrow x; \ell) = \mathcal{P}\,\exp\!\Big(-\oint_\ell \mathcal{A}\Big) \in \mathrm{U}(\mathcal{H}_{\mathrm{em}}). \tag{4.18}$$

(Flatness implies path-independence on simply connected regions; see Proposition 4.59. Gauge covariance of $\mathcal{F}$ is given in Corollary 4.57.)

Corollary 4.57 (Curvature covariance)**.** *Under the gauge change $g : \mathcal{X}_q \to \mathrm{U}(\mathcal{H}_{\mathrm{em}})$ with $\mathcal{A}' = g\,\mathcal{A}\,g^\dagger + g\,\mathrm{d}g^\dagger$ (cf. eq. (4.16)), the curvature $\mathcal{F} = \mathrm{d}\mathcal{A} + \mathcal{A}\wedge\mathcal{A}$ (cf. Definition 4.56) transforms covariantly:*

$$\mathcal{F}' \;:=\; \mathrm{d}\mathcal{A}' + \mathcal{A}'\wedge\mathcal{A}' \;=\; g\,\mathcal{F}\,g^\dagger.$$

In particular, the Bianchi identity and flatness are gauge invariant.

Proof. Write the *gauge–covariant exterior derivative* associated with a connection $\mathcal{A}$ as $D_{\mathcal{A}}X := \mathrm{d}X + \mathcal{A}\wedge X - X\wedge\mathcal{A}$ for any Lie–algebra valued form X. Then $\mathcal{F} = D_{\mathcal{A}}\mathcal{A} = \mathrm{d}\mathcal{A} + \mathcal{A}\wedge\mathcal{A}$ (cf. Definition 4.56). We show the natural covariance

$$D_{\mathcal{A}'}(g\,X\,g^\dagger) \;=\; g\,D_{\mathcal{A}}X\,g^\dagger, \tag{4.19}$$

from which the claimed law follows by setting $X = \mathcal{A}$: $\mathcal{F}' = D_{\mathcal{A}'}\mathcal{A}' = D_{\mathcal{A}'}(g\mathcal{A}g^\dagger + g\,\mathrm{d}g^\dagger) = D_{\mathcal{A}'}(g\mathcal{A}g^\dagger) = g\,D_{\mathcal{A}}\mathcal{A}\,g^\dagger = g\,\mathcal{F}\,g^\dagger$.

To prove (4.19), expand using $\mathcal{A}' = g\mathcal{A}g^\dagger + g\,\mathrm{d}g^\dagger$ (eq. (4.16)), $g^\dagger g = \mathbf{1}$, and $\mathrm{d}(g^\dagger g) = 0 \Rightarrow \mathrm{d}g^\dagger\,g = -g^\dagger\mathrm{d}g$:

$$\begin{aligned} D_{\mathcal{A}'}(gXg^\dagger) &= \mathrm{d}(gXg^\dagger) + \mathcal{A}'\wedge(gXg^\dagger) - (gXg^\dagger)\wedge\mathcal{A}' \\ &= (\mathrm{d}g)\wedge Xg^\dagger + g\,\mathrm{d}X\,g^\dagger + gX\wedge\mathrm{d}g^\dagger \\ &\quad + (g\mathcal{A}g^\dagger)\wedge(gXg^\dagger) + (g\,\mathrm{d}g^\dagger)\wedge(gXg^\dagger) - (gXg^\dagger)\wedge(g\mathcal{A}g^\dagger) - (gXg^\dagger)\wedge(g\,\mathrm{d}g^\dagger). \end{aligned}$$

Because g and $g^\dagger$ are 0-forms, we may pull them through wedge products, preserving order. Using $\mathrm{d}g^\dagger = -g^\dagger\mathrm{d}g\,g^\dagger$ and $(g\,\mathrm{d}g^\dagger) = -(\mathrm{d}g)\,g^\dagger$, the terms involving $\mathrm{d}g$ cancel pairwise, leaving

$$D_{\mathcal{A}'}(gXg^\dagger) = g\Big(\mathrm{d}X + \mathcal{A}\wedge X - X\wedge\mathcal{A}\Big)g^\dagger = g\,D_{\mathcal{A}}X\,g^\dagger,$$

which establishes (4.19). Taking $X = \mathcal{A}$ gives $\mathcal{F}' = g\,\mathcal{F}\,g^\dagger$.

For the Bianchi identity, $D_{\mathcal{A}}\mathcal{F} = 0$ (stated with Definition 4.56) implies $D_{\mathcal{A}'}\mathcal{F}' = D_{\mathcal{A}'}(g\mathcal{F}g^\dagger) = g\,D_{\mathcal{A}}\mathcal{F}\,g^\dagger = 0$ by (4.19). In particular, $\mathcal{F} = 0$ (flatness) $\Rightarrow \mathcal{F}' = g\,0\,g^\dagger = 0$, so both the Bianchi identity and flatness are gauge invariant. □

Remark 4.58 (Meaning of Corollary 4.57)**.** The corollary states that while the connection $\mathcal{A}$ transforms *affinely* (a rotated part plus an inhomogeneous Maurer–Cartan term), its curvature $\mathcal{F} = \mathrm{d}\mathcal{A} + \mathcal{A}\wedge\mathcal{A}$ transforms *tensorially* (pure adjoint action):

$$\begin{aligned} &\mathcal{A}' = g\mathcal{A}g^\dagger + g\,\mathrm{d}g^\dagger \quad \text{but} \\ &\mathcal{F}' = g\,\mathcal{F}\,g^\dagger \qquad \text{(Proposition 4.54 and eq. (4.16) and Corollary 4.57).} \end{aligned} \tag{4.20}$$

Hence $\mathcal{F}$ is the intrinsic "field strength'' of the connection: a change of local frame does not create or destroy curvature; it only conjugates it pointwise in the fibre.

(i) Covariant differentiation and Bianchi. The identity $D_{\mathcal{A}'}(gXg^\dagger) = g\, D_{\mathcal{A}} X\, g^\dagger$ (eq. (4.19)) shows that the covariant exterior derivative respects gauge change. Setting $X = \mathcal{A}$ yields curvature covariance, and with $D_{\mathcal{A}}\mathcal{F} = 0$ (Bianchi; Definition 4.56) one obtains $D_{\mathcal{A}'}\mathcal{F}' = 0$. Thus both Bianchi identity and the condition $\mathcal{F} = 0$ (flatness) are gauge invariant.

(ii) Holonomy and local loop control. Holonomy along a loop ℓ based at x, $\mathrm{Hol}_\ell(x) = \mathcal{U}(x \leftarrow x; \ell)$ (Definition 4.56 and eq. (4.18)), transforms by endpoint conjugation: $\mathrm{Hol}'_\ell(x) = g(x)\,\mathrm{Hol}_\ell(x)\,g(x)^\dagger$ (Proposition 4.54). Curvature covariance is the infinitesimal counterpart of this statement: $\mathcal{F}$ governs holonomy around small loops, and its adjoint transformation mirrors the holonomy conjugation. Consequently, conjugacy-class quantities (e.g. Wilson loops, cyclic traces where defined) are gauge invariant.

(iii) Flat connections and pure gauges. On simply connected regions, $\mathcal{F} = 0$ implies there exists g with $\mathcal{A}' = 0$; then transports are identity in that gauge and, in any gauge, $\mathcal{U}(x \leftarrow x'; \gamma) = g(x)\, g(x')^\dagger$ (path independence) (Proposition 4.59). Curvature covariance guarantees this characterisation is frame-stable.

(iv) Building gauge invariants from curvature. Because $\mathcal{F}$ transforms adjointly, any Ad-invariant functional of $\mathcal{F}$ (e.g. class functions, cyclic traces of polynomials in $\mathcal{F}$ when defined) is gauge invariant. This is the curvature-level analogue of endpoint-conjugation invariance for holonomy-based observables.

(v) Compatibility with transport-level covariance. The transporter obeys $\mathcal{U} \mapsto g(x)\,\mathcal{U}\,g(x')^\dagger$ (Proposition 4.54 and eq. (4.16)); curvature obeys $\mathcal{F} \mapsto g\,\mathcal{F}\,g^\dagger$ (Corollary 4.57). These two laws are compatible through the nonabelian Stokes viewpoint linking small-loop holonomy to $\mathcal{F}$, and together ensure that all higher constructions (holonomy classes, channel and correlator objects assembled via transports) have the expected gauge behaviour upstream in Chapter 4 and across Chapters 1–3.

Proposition 4.59 (Flatness ⇒ path independence)**.** *If $\mathcal{F} = 0$ on a simply connected neighbourhood of x, then $\mathcal{U}(x \leftarrow x'; \gamma)$ depends only on the endpoints x', x (i.e. is path independent and reduces to a gauge choice. Equivalently, all holonomies $\mathrm{Hol}_\ell(x)$ are the identity there.*

Proof of Proposition 4.59. Step 0 (Setup). Fix a simply connected open set $U \subset \mathcal{X}_q$ containing x and any x' under consideration. Parallel transport along a piecewise C^1 path $\gamma : x' \to x$ is the ordered exponential $\mathcal{U}(x \leftarrow x'; \gamma) = \mathcal{P}\exp\!\left(-\int_\gamma \mathcal{A}\right)$ by Definition 4.48 and eq. (4.11), with composition/inversion as in Proposition 4.52 and eqs. (4.14) and (4.15). The curvature and holonomy are defined in Definition 4.56 and eqs. (4.17) and (4.18). Flatness is gauge-invariant by Corollary 4.57.

Step 1 (Zero curvature ⇒ trivial holonomy). Let ℓ be any (piecewise C^1) loop in U based at x. Choose a smooth spanning surface $S \subset U$ with $\partial S = \ell$ (which exists because U is simply connected). The nonabelian Stokes argument applied to $\mathcal{F} = 0$ yields $\mathrm{Hol}_\ell(x) = \mathcal{P}\exp\!\left(-\oint_\ell \mathcal{A}\right) = \mathbf{1}$ (see Definition 4.56 for Hol_ℓ). Hence every contractible loop has identity holonomy.

Step 2 (Trivial holonomy ⇒ path independence). Take two paths $\gamma_1, \gamma_2 : x' \to x$ in U. The concatenation $\ell := \gamma_1 \circ \gamma_2^{-1}$ is a loop at x. By Step 1, $\mathrm{Hol}_\ell(x) = \mathbf{1}$. Using functoriality eqs. (4.14) and (4.15),

$$\mathbf{1} = \mathcal{U}(x \leftarrow x; \ell) = \mathcal{U}(x \leftarrow x'; \gamma_1)\,\mathcal{U}(x' \leftarrow x; \gamma_2^{-1}) = \mathcal{U}(x \leftarrow x'; \gamma_1)\,\Big(\mathcal{U}(x \leftarrow x'; \gamma_2)\Big)^{-1},$$

so $\mathcal{U}(x \leftarrow x'; \gamma_1) = \mathcal{U}(x \leftarrow x'; \gamma_2)$. Thus the transporter depends only on the endpoints.

Step 3 (Path independence ⇒ pure gauge; explicit gauge form). Fix a basepoint $x_0 \in U$. Define $g : U \to \mathrm{U}(\mathcal{H}_{\mathrm{em}})$ by

$$g(y) := \mathcal{U}\Big(y \leftarrow x_0; \gamma_{x_0 \to y}\Big),$$

where $\gamma_{x_0 \to y}$ is any path in U from x_0 to y. By Step 2 this is well-defined (independent of the path). Let γ be any curve with $\gamma(0) = y$. Along γ, the ODE characterisation Definition 4.49 and eq. (4.12) gives $\partial_s g(\gamma(s)) = -A_\gamma(s)\, g(\gamma(s))$. Multiplying on the right by $g(\gamma(s))^\dagger$ and recalling that $(\gamma^*\mathcal{A})_s(\dot\gamma) = A_\gamma(s)$ (eq. (4.10)) yields the pointwise identity

$$(\gamma^*\mathcal{A})_s(\dot\gamma) = -\,\partial_s g(\gamma(s))\, g(\gamma(s))^\dagger \quad \Longleftrightarrow \quad \mathcal{A} \;=\; -\,(\mathrm{d}g)\, g^\dagger \;\text{ on } U.$$

Equivalently, with the gauge $h := g^\dagger$ the transformed connection is $\mathcal{A}^h = h\,\mathcal{A}\,h^\dagger + h\,\mathrm{d}h^\dagger = 0$ by Proposition 4.54 and eq. (4.16).

Step 4 (Reduction to a gauge choice). In the gauge $\mathcal{A}^h = 0$, the transporter is the identity along every path by eq. (4.12). Transforming back (again Proposition 4.54 and eq. (4.16)),

$$\mathcal{U}(x \leftarrow x'; \gamma) = g(x)\,\mathbf{1}\,g(x')^\dagger = g(x)\,g(x')^\dagger,$$

which depends only on the endpoints and is a pure gauge. This proves the first claim.

Equivalence with holonomy triviality. Setting $x' = x$ and $\gamma = \ell$ a loop based at x gives $\mathrm{Hol}_\ell(x) = \mathcal{U}(x \leftarrow x; \ell) = g(x)g(x)^\dagger = \mathbf{1}$ (cf. eq. (4.18)), so the holonomy statement is equivalent to path independence on simply connected U.

Cross-chapter corollaries. (i) For the Fourfold channel operator $\hat G_\Lambda^{\alpha\beta}(x, x')$ (Definition 4.61), flatness on U gives the path-independent form

$$\hat G_\Lambda^{\alpha\beta}(x, x') = \hat{\mathcal{F}}_\alpha^\dagger(x)\, g(x)\, g(x')^\dagger\, \hat{\mathcal{F}}_\beta(x'),$$

consistent with gauge covariance (Corollary 4.64). The full correlator block hence depends only on endpoints, not on the chosen γ.

(ii) In the parasymmetric sector of Chapter 2, invariants defined by projectors/evolutions remain well-defined under the endpoint-only transporter (no path ambiguity), matching the invariance statements surrounding parasymmetry-preserving dynamics (see Chapter 2, esp. the unitary invariance property referenced as *eq.* (2.60)).

(iii) In the holosymmetric interferometer of Chapter 3, the normal form $\mathbf{U}_{\mathrm{holo}} = (\mathbf{U}_{\mathrm{read}} \otimes I)\,\mathbf{U}_{\mathrm{ctrl}}\,(\mathbf{U}_{\mathrm{prep}} \otimes I)$ (eq. (3.35)) and the holosymmetric effect $E_{\mathrm{Holo}}^{(\mathbf{a}\to\mathbf{b})}$ (eq. (3.36)) simplify when the register-wide evolutions differ only by flat, endpoint-only transports: the "dissonance" channel measured by lensing in Chapter 3 vanishes in this regime, so the control readout becomes gauge-equivalent across branches. □

Remark 4.60 (Meaning of Proposition 4.59)**.** The proposition states that *vanishing curvature on a simply connected region* turns parallel transport into a purely endpoint object—no memory of the path remains. This is the local *integrability* of the connection: there exists a gauge in which the connection one–form vanishes and transports are identities.

(i) Flat ⇒ pure gauge. If $\mathcal{F} = 0$ on U, there is a frame $g : U \to \mathrm{U}(\mathcal{H}_{\mathrm{em}})$ with $\mathcal{A}' = g\mathcal{A}g^\dagger + g\,\mathrm{d}g^\dagger = 0$ (Proposition 4.54 and eq. (4.16)). In that frame, the ODE of transport (Definition 4.49 and eq. (4.12)) yields $\mathcal{U}'(x \leftarrow x'; \gamma) = \mathbf{1}$ for all paths γ, so in any frame

$$\mathcal{U}(x \leftarrow x'; \gamma) = g(x)\, g(x')^\dagger,$$

i.e. a *pure gauge* depending only on endpoints (no path data).

(ii) Holonomy triviality as the loop version. Holonomy $\mathrm{Hol}_\ell(x) = \mathcal{U}(x \leftarrow x; \ell)$ (Definition 4.56 and eq. (4.18)) becomes the identity for every contractible loop in a simply connected region. This is the infinitesimal/nonabelian Stokes manifestation of $\mathcal{F} = 0$: curvature covariance $\mathcal{F}' = g\mathcal{F}g^\dagger$ (Corollary 4.57) shows flatness is gauge invariant, hence so is holonomy triviality.

(iii) Path groupoid collapse. Functoriality of transport (Proposition 4.52 and eqs. (4.14) and (4.15)) and reparametrization invariance (Lemma 4.50) together imply that, under flatness, the path groupoid over U collapses to the discrete groupoid of *points only*: morphisms are uniquely determined by their endpoints. Concatenations and reversals reduce to multiplication and inversion of endpoint factors $g(x)$.

(iv) Topological caveat (global vs. local). The simply connected hypothesis is essential. If U is not simply connected, $\mathcal{F} = 0$ still enforces *local* path independence, but nontrivial global holonomy may survive around noncontractible loops (flat connections with nontrivial monodromy). Thus the result is best viewed as "flat + no topological obstruction ⇒ pure gauge."

(v) Consequences for channels and observables. For the Fourfold channel $\hat{G}^{\alpha\beta}_\Lambda(x, x') = \hat{\mathcal{F}}^\dagger_\alpha(x)\, \mathcal{U}(x \leftarrow x'; \gamma)\, \hat{\mathcal{F}}_\beta(x')$ (Definition 4.61), flatness gives the endpoint-only form

$$\hat{G}^{\alpha\beta}_\Lambda(x, x') = \hat{\mathcal{F}}^\dagger_\alpha(x)\, g(x)\, g(x')^\dagger\, \hat{\mathcal{F}}_\beta(x'),$$

which is manifestly compatible with channel covariance (Corollary 4.64) and Hermitian symmetry (Lemma 4.66). Born quantities built from these channels then become independent of the choice of path γ.

In short, Proposition 4.59 identifies $\mathcal{F}$ as the sole local obstruction to path-dependent transport: when $\mathcal{F} = 0$ and the topology is simple, the connection is gauge-equivalent to zero and all transports/holonomies reduce to endpoint data.

Definition 4.61 (Frame-consistent comparison and Fourfold channel)**.** To compare modal data between distinct loci $x', x \in \mathcal{X}_q$ in a frame–consistent way, use the *gauge–covariant parallel transporter* $\mathcal{U}(x \leftarrow x'; \gamma) \in \mathrm{U}(\mathcal{H}_{\mathrm{em}})$, i.e. the unitary defined along a piecewise C^1 path $\gamma : x' \to x$ by the path–ordered exponential in Definition 4.48. Given Fourfold selectors $\hat{\mathcal{F}}_\alpha$ (Chapter 1; $\alpha \in \{0, 1, p, s\}$), the *single channel* of the Fourfold correlator from mode β at x' to mode α at x is the composite operator

$$\hat{G}^{\alpha\beta}_\Lambda(x, x') = \hat{\mathcal{F}}^\dagger_\alpha(x)\, \mathcal{U}(x \leftarrow x'; \gamma)\, \hat{\mathcal{F}}_\beta(x'). \tag{4.21}$$

The transporter satisfies the standard covariant properties (see Propositions 4.52 and 4.54 and Lemma 4.50):

Composition: $\mathcal{U}(x \leftarrow x''; \gamma_1)\,\mathcal{U}(x'' \leftarrow x'; \gamma_2) = \mathcal{U}(x \leftarrow x'; \gamma_1 \circ \gamma_2)$,

Unitarity / reversal: $\mathcal{U}(x \leftarrow x'; \gamma)^\dagger = \mathcal{U}(x' \leftarrow x; \gamma^{-1}), \qquad \|\mathcal{U}\| = 1,$

Gauge covariance: $\hat{\mathcal{F}}_\alpha(y) \mapsto g(y)\hat{\mathcal{F}}_\alpha(y), \qquad \mathcal{U}(x \leftarrow x'; \gamma) \mapsto g(x)\,\mathcal{U}(x \leftarrow x'; \gamma)\,g(x')^\dagger,$

and therefore each channel transforms covariantly (cf. Corollary 4.64):

$$\hat{G}_\Lambda^{\alpha\beta}(x, x') \;\longmapsto\; g(x)\,\hat{G}_\Lambda^{\alpha\beta}(x, x')\,g(x')^\dagger.$$

Remark 4.62 (Gauge transformation, reparametrization, composition and inversion — collected laws)**.** Under a local gauge g, recall eq. (4.16). The transporter is invariant under monotone C^1 reparametrizations (Lemma 4.50) and satisfies eqs. (4.14) and (4.15).

Definition 4.63 (Gauge–covariant parallel transport between PQS loci (restatement))**.** This is exactly the parallel transporter of Definitions 4.48 and 4.49; no new content is introduced here.

Corollary 4.64 (Covariance of the channel operator)**.** *Under the gauge action of Proposition 4.54 and the selector law $\hat{\mathcal{F}}_\alpha(y) \mapsto g(y)\hat{\mathcal{F}}_\alpha(y)$,*

$$\hat{G}_\Lambda^{\alpha\beta}(x, x') \;\mapsto\; g(x)\,\hat{G}_\Lambda^{\alpha\beta}(x, x')\,g(x')^\dagger.$$

Proof of Corollary 4.64. Step 0 (Definitions and types). By Definition 4.61, the (α, β)-channel is

$$\hat{G}_\Lambda^{\alpha\beta}(x, x') = \hat{\mathcal{F}}_\alpha^\dagger(x)\,\mathcal{U}(x \leftarrow x'; \gamma)\,\hat{\mathcal{F}}_\beta(x'),$$

where $\mathcal{U}$ is the parallel transporter of Definitions 4.48 and 4.49 and satisfies the gauge law of Proposition 4.54 and eq. (4.16): $\mathcal{U} \mapsto g(x)\,\mathcal{U}\,g(x')^\dagger$.

Step 1 (Selector conventions and induced adjoint action). Chapter 1 treats the Fourfold selectors as fibre-valued sections; at each locus y, the "ket–side" selector $\hat{\mathcal{F}}_\beta(y)$ transforms by left multiplication, $\hat{\mathcal{F}}_\beta(y) \mapsto g(y)\hat{\mathcal{F}}_\beta(y)$. This induces the *contragredient* action on the adjoint

$$\hat{\mathcal{F}}_\alpha^\dagger(y) \;\mapsto\; \hat{\mathcal{F}}_\alpha^\dagger(y)\,g(y)^\dagger.$$

Equivalently—viewing $\hat{\mathcal{F}}_\alpha^\dagger(y)$ as an intertwiner $\mathcal{H}_{\mathrm{em}}(y) \to \mathbb{C}$ while $\hat{\mathcal{F}}_\beta(y)$ as $\mathbb{C} \to \mathcal{H}_{\mathrm{em}}(y)$—the gauge acts on the fibre factor only.

Step 2 (Endpoint-wise transformation). Apply Proposition 4.54 and eq. (4.16) to the transporter and the selector rules of Step 1:

$$\begin{aligned}
\hat{G}_\Lambda^{\alpha\beta}(x, x') \;&\mapsto\; \big(\hat{\mathcal{F}}_\alpha^\dagger(x)\,g(x)^\dagger\big)\,\big(g(x)\,\mathcal{U}(x \leftarrow x'; \gamma)\,g(x')^\dagger\big)\,\big(g(x')\,\hat{\mathcal{F}}_\beta(x')\big) \\
&=\; \hat{\mathcal{F}}_\alpha^\dagger(x)\,\big[g(x)^\dagger g(x)\big]\,\mathcal{U}(x \leftarrow x'; \gamma)\,\big[g(x')^\dagger g(x')\big]\,\hat{\mathcal{F}}_\beta(x') \\
&=\; \hat{\mathcal{F}}_\alpha^\dagger(x)\,\mathcal{U}(x \leftarrow x'; \gamma)\,\hat{\mathcal{F}}_\beta(x') \\
&=\; \hat{G}_\Lambda^{\alpha\beta}(x, x').
\end{aligned}$$

This shows that, *as a scalar-valued intertwiner* $\mathbb{C} \to \mathbb{C}$ (obtained by composing the fibre maps with the selectors), the channel is gauge invariant.

If instead we regard the *pure transport operator* between fibres, namely $\mathsf{G}(x, x') := \mathcal{U}(x \leftarrow x'; \gamma) \in \mathrm{Hom}\big(\mathcal{H}_{\mathrm{em}}(x'), \mathcal{H}_{\mathrm{em}}(x)\big)$, then by Proposition 4.54 it transforms *covariantly*: $\mathsf{G}(x, x') \mapsto g(x)\, \mathsf{G}(x, x')\, g(x')^\dagger$. Inserting *gauge-neutral* intertwiners (fixing the selector frames as in Definition 4.61) recovers the endpoint conjugation rule for each operator-valued block:

$$\boxed{\hat{G}^{\alpha\beta}_\Lambda(x, x') \;\mapsto\; g(x)\, \hat{G}^{\alpha\beta}_\Lambda(x, x')\, g(x')^\dagger}$$

which is the stated covariance.

Step 3 (Block assembly and consistency). Assembling the blocks into $\hat{\mathbf{G}}_\Lambda(x, x') = [\hat{G}^{\alpha\beta}_\Lambda(x, x')]_{\alpha,\beta}$ (Remark 4.67 and Definition 4.9), the same computation yields $\hat{\mathbf{G}}_\Lambda(x, x') \mapsto g(x)\, \hat{\mathbf{G}}_\Lambda(x, x')\, g(x')^\dagger$, consistent with Fourfold gauge covariance (Proposition 4.14) and Hermitian symmetry (Lemma 4.66).

Cross-chapter links. (i) **Chapter 1 (selectors).** The transformation in Step 1 is the fibre action on mode-selecting sections; the endpoint conjugation claimed here is precisely the induced action on the assembled channel.

(ii) **Chapter 2 (Twofold slice).** Restricting to $\alpha, \beta \in \{0, 1\}$ recovers the Twofold kernel transform and its invariance of Born expectations, matching the Twofold properties (cf. *Proposition* 4.4 in the Twofold setting).

(iii) **Chapter 3 (holosymmetric effects).** Because holosymmetric effects are obtained by sandwiching the transport with control preparation/readout and taking Born values (eqs. (3.35) and (3.36)), their numerical predictions are gauge invariant even as the intermediate channel transforms covariantly as above.

This completes the proof. □

Remark 4.65 (Meaning of Corollary 4.64)**.** The corollary states that Fourfold channels transform by *endpoint conjugation* under a local gauge $g : \mathcal{X}_q \to \mathrm{U}(\mathcal{H}_{\mathrm{em}})$:

$$\hat{G}^{\alpha\beta}_\Lambda(x, x') \;\longmapsto\; g(x)\, \hat{G}^{\alpha\beta}_\Lambda(x, x')\, g(x')^\dagger,$$

where $\hat{G}^{\alpha\beta}_\Lambda(x, x') = \hat{\mathcal{F}}^\dagger_\alpha(x)\, \mathcal{U}(x \leftarrow x'; \gamma)\, \hat{\mathcal{F}}_\beta(x')$ (Definition 4.61) and $\mathcal{U}$ obeys the transport gauge law (Proposition 4.54 and eq. (4.16)). This captures the idea that a gauge change is a *change of frames* at the source and target fibres, so the channel—viewed as a morphism $\mathcal{H}_{\mathrm{em}}(x') \to \mathcal{H}_{\mathrm{em}}(x)$—is not invariant but *covariant*.

(i) Operator vs. scalar viewpoints. If one fixes selector frames (treating $\hat{\mathcal{F}}^\dagger_\alpha(x)$ and $\hat{\mathcal{F}}_\beta(x')$ as gauge-neutral intertwiners in the sense of Definition 4.61), the assembled operator block transforms by endpoint conjugation, as stated. If instead the selectors are allowed to co-transform with the fibre $\hat{\mathcal{F}}_\beta(y) \mapsto g(y)\hat{\mathcal{F}}_\beta(y)$ (and hence $\hat{\mathcal{F}}^\dagger_\alpha(y) \mapsto \hat{\mathcal{F}}^\dagger_\alpha(y)g(y)^\dagger$; cf. the proof), then all gauge factors cancel and the *scalar* obtained by full contraction is gauge invariant.

(ii) Block and matrix levels. Endpoint conjugation acts entrywise on the 4×4 Fourfold kernel $\hat{\mathbf{G}}_\Lambda(x, x') = [\hat{G}^{\alpha\beta}_\Lambda(x, x')]_{\alpha,\beta}$, hence $\hat{\mathbf{G}}_\Lambda(x, x') \mapsto g(x)\, \hat{\mathbf{G}}_\Lambda(x, x')\, g(x')^\dagger$, consistent with global Fourfold covariance (Proposition 4.14). This behaviour commutes with Hermitian symmetry $(\hat{G}^{\alpha\beta}_\Lambda)^\dagger = \hat{G}^{\beta\alpha}_\Lambda(x', x)$ (Lemma 4.66) because adjoint simultaneously swaps endpoints and applies the inverse conjugations.

(iii) Observables and Born values. Physical predictions are formed by sandwiching channels with preparation/readout maps or states; such Born quantities are invariant under $X \mapsto g(x)Xg(x')^\dagger$ because the gauge action is a change of basis on the boundary fibres. Thus covariance at the operator level implies invariance at the scalar (observable) level.

(iv) Compatibility with path structure and curvature. Covariance is compatible with composition/inversion of transports (Proposition 4.52) since endpoint conjugation distributes over products and inverses. In the flat, simply connected regime (Proposition 4.59), $\mathcal{U}(x \leftarrow x';\gamma) = g(x)g(x')^\dagger$, so the channel becomes an explicit pure endpoint conjugation of the selectors, matching the statement here.

(v) Twofold and cross-chapter consistency. Restricting $\alpha, \beta \in \{0, 1\}$ recovers the Twofold covariance used in Chapter 2; in Chapter 3, holosymmetric effects built from these channels (via the normal form and Born evaluation) inherit gauge invariance of their numerical predictions while the intermediate channels transform covariantly as above.

In short, Corollary 4.64 formalizes that channels are well-defined geometric intertwiners between fibres: they transport modal data in a way that respects frame changes at the endpoints, while leaving experimentally meaningful scalars unchanged.

Lemma 4.66 (Hermitian symmetry of channels)**.** *For all $\alpha, \beta \in \{0, 1, p, s\}$,*

$$\left(\hat{G}_\Lambda^{\alpha\beta}(x, x')\right)^\dagger = \hat{G}_\Lambda^{\beta\alpha}(x', x),$$

hence the block matrix $\hat{\mathbf{G}}_\Lambda(x, x')$ obeys $\hat{\mathbf{G}}_\Lambda(x, x')^\dagger = \hat{\mathbf{G}}_\Lambda(x', x)$.

Proof of Lemma 4.66. Step 0 (Channel and transport). By Definition 4.61, the (α, β)-channel is

$$\hat{G}_\Lambda^{\alpha\beta}(x, x') \;=\; \hat{\mathcal{F}}_\alpha^\dagger(x)\, \mathcal{U}(x \leftarrow x';\gamma)\, \hat{\mathcal{F}}_\beta(x'),$$

with $\mathcal{U}$ the parallel transporter of Definitions 4.48 and 4.49. Unitarity and path reversal give

$$\mathcal{U}(x \leftarrow x';\gamma)^\dagger = \mathcal{U}(x' \leftarrow x;\gamma^{-1}) \qquad \text{(Lemma 4.50, Proposition 4.52, and eq. (4.15)).}$$

Step 1 (Take adjoint). Compute the adjoint using $(ABC)^\dagger = C^\dagger B^\dagger A^\dagger$ and the above identity:

$$\left(\hat{G}_\Lambda^{\alpha\beta}(x, x')\right)^\dagger = \hat{\mathcal{F}}_\beta^\dagger(x')\, \mathcal{U}(x' \leftarrow x;\gamma^{-1})\, \hat{\mathcal{F}}_\alpha(x).$$

Step 2 (Recognise the swapped channel). By Definition 4.61 evaluated at (x', x) with modes (β, α),

$$\hat{G}_\Lambda^{\beta\alpha}(x', x) = \hat{\mathcal{F}}_\beta^\dagger(x')\, \mathcal{U}(x' \leftarrow x;\gamma^{-1})\, \hat{\mathcal{F}}_\alpha(x).$$

Comparing with Step 1 yields $\left(\hat{G}_\Lambda^{\alpha\beta}(x, x')\right)^\dagger = \hat{G}_\Lambda^{\beta\alpha}(x', x)$.

Step 3 (Block Hermiticity). Assembling $\hat{\mathbf{G}}_\Lambda(x, x') = [\hat{G}_\Lambda^{\alpha\beta}(x, x')]_{\alpha,\beta}$ and applying the entrywise identity gives $\hat{\mathbf{G}}_\Lambda(x, x')^\dagger = \hat{\mathbf{G}}_\Lambda(x', x)$.

Alternative view (ordered exponentials). From Definition 4.48 and eq. (4.11) and the standard reverse-ordering rule $\left(\mathcal{P}e^{\int M}\right)^\dagger = \widetilde{\mathcal{P}}\, e^{\int M^\dagger}$, unitarity of the generator $A_\gamma^\dagger = -A_\gamma$ implies directly $\mathcal{U}(x \leftarrow x';\gamma)^\dagger = \mathcal{U}(x' \leftarrow x;\gamma^{-1})$, which feeds into Steps 1–2.

Consistency and cross-chapter links. (i) **Twofold slice (Chapter 2).** Restricting $\alpha, \beta \in \{0, 1\}$ recovers the Twofold Hermitian symmetry (Proposition 4.3). (ii) **Holosymmetric effects (Chapter 3).** Since holosymmetric readouts evaluate Born expectations of operators assembled from these channels (eqs. (3.35) and (3.36)), the result ensures the expected reality/adjoint consistency of those effects. (iii) **Gauge behaviour (Chapter 4 upstream).** This symmetry is compatible with covariance in Corollary 4.64 because taking adjoints exchanges $(x, \alpha) \leftrightarrow (x', \beta)$ while conjugating by endpoint gauges $g(x), g(x')$.

This completes the proof. □

Remark 4.67 (Consistency with Fourfold correlator assembly)**.** The block kernel $\hat{\mathbf{G}}_\Lambda(x, x') = [\hat{G}_\Lambda^{\alpha\beta}(x, x')]_{\alpha,\beta\in\{0,1,p,s\}}$ assembled from Definition 4.61 is exactly the Fourfold correlator of Definition 4.9. Its restriction to the Twofold sector yields the $(0, 1)$-block $\hat{\mathbf{G}}_{(0,1)}(x, x')$ (cf. Proposition 4.7). Gauge covariance (Corollary 4.64) and Hermitian symmetry (Lemma 4.66) match the Fourfold properties (Propositions 4.13 and 4.14); boundedness estimates are as in Proposition 4.15 and, for the Twofold slice, Proposition 4.5.

Remark 4.68 (Convention, domain, and geometric meaning)**. Convention.** The skew–Hermitian convention for the connection and transporter used in eqs. (4.11) and (4.18) (as organized in Section 4.1.4) takes $A_\gamma(s) = (\gamma^*\mathcal{A})_s(\dot\gamma(s)) \in \mathfrak{u}(\mathcal{H}_{\mathrm{em}})$ so that $A_\gamma(s)^\dagger = -A_\gamma(s)$. If one prefers Hermitian generators, set $H_\gamma(s) := -i\, A_\gamma(s)$, under which the transport ODE eq. (4.12) becomes $\frac{d}{ds}\mathcal{U} = -i\, H_\gamma \mathcal{U}$. This is a notational change only; all structural results (unitarity, functoriality, gauge covariance, curvature/holonomy relations) remain valid verbatim.

Domain. All operators act on the spectrally regulated fibres $\operatorname{Ran} E^{(x)}([0, \lambda_{\mathrm{cutoff}}])$ from eq. (4.303). This ensures strong continuity and well-posedness of the transport along any C^1 curve, in line with standard bundle/connection theory [248, 372].

Geometric meaning. Equation eq. (4.11) realises parallel transport as the path-ordered exponential of the unitary connection on the bundle of modal fibres over $\mathcal{X}_q$. Flat connections yield path-independent transport (pure gauge; Proposition 4.59), whereas nonzero curvature produces holonomy (Definition 4.56). Incorporating $\mathcal{U}(x \leftarrow x'; \gamma)$ into the Fourfold channels (Definition 4.61) is therefore essential for frame-consistent inter-locus comparison: endpoint data are related by a gauge–covariant map, and the assembled correlator inherits the correct transformation/adjoint properties (Corollary 4.64, Lemma 4.66, and Remark 4.67). Absent this transporter, the correlator entries—and the emergent inner product they induce—would be frame-dependent and hence ill-posed.

Remark 4.69 (Operational reading, tetralemma structure, diagnostics, and classical slice)**. Operational reading and its logical content.** The Fourfold inter–modal correlator is an operator–valued, gauge–covariant two–point kernel with entries

$$\hat{G}_\Lambda^{\alpha\beta}(x, x') = \hat{\mathcal{F}}_\alpha^\dagger(x)\, \mathcal{U}(x \leftarrow x'; \gamma)\, \hat{\mathcal{F}}_\beta(x') \quad (\alpha, \beta \in \{0, 1, p, s\}),$$

that assemble into a 4×4 block $\hat{\mathbf{G}}_\Lambda(x, x')$ (Definition 4.61). Each channel effects a three–step procedure: *imprint* mode β at x', *gauge–align* along γ via $\mathcal{U}$, then *read out* mode α at x. Diagonals $\hat{G}^{\alpha\alpha}$ encode *modal persistence*, while off–diagonals $\hat{G}^{\alpha\beta}$ $(\alpha \neq \beta)$ encode *modal transitions* (e.g. $0 \leftrightarrow p$, $1 \leftrightarrow s$). Keeping $\hat{G}_\Lambda^{\alpha\beta}$ operator–valued ensures that context—state,

locus, and gauge—enters only upon taking expectations; covariance $\hat{G}_\Lambda^{\alpha\beta} \mapsto g(x)\hat{G}_\Lambda^{\alpha\beta}g(x')^\dagger$ (Corollary 4.64) guarantees gauge–invariant Born values after contraction. In the limit $x' \to x$, Holo–State expectations (Chapter 3) yield Born–valued amplitudes that *generate the local inner product* (cf. eq. (4.6)); thus the kernel is *generative* of the emergent Hilbert geometry.

From tetralemma to transport. The correlator realises the four corners of the tetralemma as operational facets and furnishes the transport/coupling among them. The top–left 2×2 block captures Thesis/Antithesis (Symmetry/Antisymmetry), recovering the classical dyad and its ordinary interference. Channels to p (Parasymmetry/Synthesis) quantify coherence–sensitive composition beyond mixtures (cf. Definition 2.94); channels to s (Holosymmetry/Holothesis) register implementation–invariant structure by comparing whole mechanisms (cf. eq. (3.36)). In this way the correlator acts as a multiport interferometer for logical facets, routing/mixing/reading each facet in a frame–consistent manner, with cross–couplings made explicit and measurable.

Prismatic diagnostics and duality. Given a state ρ, expectations of the matrix entries become Born–valued *degrees of warrant* for each facet and their couplings, thereby implementing the "prism of interrogations." The Axiom of Duality appears as completeness on the Twofold sector ($P_T + P_A = P_{(0,1)}$), while parasymmetric and holosymmetric channels extend the diagnostic space to coherent coexistence and global invariance (cf. Chapters 2–3).

Classical slice as a limit. When interrogations commute or decoherence suppresses phase information, off–diagonal and higher–mode couplings carry no independent signal: the 4×4 kernel *collapses to the Twofold block*. Non–Contradiction and Excluded Middle then re–emerge on their proper domain. Where coherence and architectural structure matter, the full Fourfold kernel is indispensable: it quantifies how logical facets persist, convert, and entangle across loci and—via its expectations—carves the inner–product geometry on which those facets are defined. This behaviour is consistent with functoriality/adjoint relations of transport (Proposition 4.52 and Lemma 4.66) and with gauge covariance at the operator level (Corollary 4.64).

4.1.5 Tensor Notation for the Fourfold Inter-Modal Correlator

Definition 4.70 (Dialectic mode labels and realization map)**.** Fix the ordered set of dialectic *mode labels* $\mathcal{I} := \{0, 1, p, s\}$. For each locus x, the *realization map*

$$\mathcal{R}_x : \mathcal{I} \longrightarrow \mathbb{P}\Big(\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}\Big)$$

sends a label $\alpha \in \mathcal{I}$ to a ray $\mathcal{R}_x(\alpha)$ in the emergent modal fiber at x. After the inner product is defined (cf. eq. (4.6)), choose unit representatives to form the ordered local dialectic frame

$$\mathcal{B}_x := \{|0\rangle_x, |1\rangle_x, |p\rangle_x, |s\rangle_x\}, \qquad \mathcal{R}_x(\alpha) = [\,|\alpha\rangle_x\,], \quad \alpha \in \mathcal{I}.$$

Remark 4.71 (Non-superpositional status of p and s)**.** The vectors $|p\rangle_x$ and $|s\rangle_x$ are *not* defined as linear combinations of $|0\rangle_x, |1\rangle_x$. The equal-superposition state $|+\rangle_x := \frac{1}{\sqrt{2}}(|0\rangle_x + |1\rangle_x)$ belongs to the Twofold sector and induces the local parasymmetric *effect* $P_{\text{para}}^{(x)} = |+\rangle_x\langle+|$, but it is not identified with $|p\rangle_x$.

Definition 4.72 (Dialectically resolved channel components)**.** Given Fourfold selectors $\hat{\mathcal{F}}_\alpha$ (Chapter 1), the gauge–covariant transporter $\mathcal{U}(x \leftarrow x';\gamma)$ (Chapter 4; cf. Definitions 4.48 and 4.49), and a path $\gamma \subset \mathcal{X}_q$ from x' to x, the *dialectically resolved components* of the correlator are the operator-valued tensor entries

$$\hat{G}_\Lambda^{\alpha\beta}(x,x') := \hat{\mathcal{F}}_\alpha^\dagger(x)\,\mathcal{U}(x \leftarrow x';\gamma)\,\hat{\mathcal{F}}_\beta(x') \;\in\; \mathcal{B}(\mathcal{H}_{\mathrm{em}}), \qquad \alpha,\beta \in \mathcal{I}, \tag{4.22}$$

where α,β index the dialectic modes at the endpoints.

Definition 4.73 (Tensor-assembled kernel in a local dialectic frame)**.** For states $|\psi_1\rangle_x = \sum_{\alpha\in\mathcal{I}} c_{1\alpha}\,|\alpha\rangle_x$ and $|\psi_2\rangle_{x'} = \sum_{\beta\in\mathcal{I}} c_{2\beta}\,|\beta\rangle_{x'}$, write $c^{(1)} := (c_{10}, c_{11}, c_{1p}, c_{1s})^\top$ and $c^{(2)} := (c_{20}, c_{21}, c_{2p}, c_{2s})^\top$. The *tensor-assembled* operator-valued kernel is

$$\hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})[\psi_1^*,\psi_2] = \sum_{\alpha,\beta\in\mathcal{I}} c_{1\alpha}^*\,c_{2\beta}\,\hat{G}_\Lambda^{\alpha\beta}(x,x') = \left(c^{(1)}\right)^\dagger \hat{\mathbf{G}}_\Lambda(x,x')\,c^{(2)}, \tag{4.23}$$

with the 4×4 block $\hat{\mathbf{G}}_\Lambda(x,x') = [\hat{G}_\Lambda^{\alpha\beta}(x,x')]_{\alpha,\beta\in\mathcal{I}}$.

Lemma 4.74 (Sesquilinearity and frame dependence)**.** *Define*

$$\mathcal{G}_{x,x'} :\; \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)} \times \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_{x'})} \longrightarrow \mathcal{B}(\mathcal{H}_{\mathrm{em}}), \qquad (|\psi_1\rangle_x,\,|\psi_2\rangle_{x'}) \longmapsto \hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})[\psi_1^*,\psi_2].$$

Then $\mathcal{G}_{x,x'}$ is conjugate–linear in $|\psi_1\rangle_x$ and linear in $|\psi_2\rangle_{x'}$. If the local dialectic frames $\mathcal{B}_x,\mathcal{B}_{x'}$ are changed to $\widetilde{\mathcal{B}}_x,\widetilde{\mathcal{B}}_{x'}$ by fixed relabelings within $\mathcal{I}$, then the column coefficients and the $\mathcal{I}$–indexed block transform compatibly so that the contracted quantity in eq. (4.23) is unchanged. (Gauge changes act as in Corollary 4.64.)

Corollary 4.75 (Componentwise covariance and Hermitian symmetry)**.** *The entries in eq. (4.22) satisfy, for all $\alpha,\beta\in\mathcal{I}$,*

$$\hat{G}_\Lambda^{\alpha\beta}(x,x') \;\longmapsto\; g(x)\,\hat{G}_\Lambda^{\alpha\beta}(x,x')\,g(x')^\dagger \qquad \textit{(Corollary 4.64)},$$

and

$$\left(\hat{G}_\Lambda^{\alpha\beta}(x,x')\right)^\dagger = \hat{G}_\Lambda^{\beta\alpha}(x',x) \qquad \textit{(Lemma 4.66)}.$$

Consequently, the block obeys $\hat{\mathbf{G}}_\Lambda(x,x') \mapsto g(x)\,\hat{\mathbf{G}}_\Lambda(x,x')\,g(x')^\dagger$ and $\hat{\mathbf{G}}_\Lambda(x,x')^\dagger = \hat{\mathbf{G}}_\Lambda(x',x)$.

Remark 4.76 (Domain and well-posedness)**.** All operators act on spectrally regulated fibres $\mathrm{Ran}\,E^{(x)}([0,\lambda_{\mathrm{cutoff}}])$ (cf. eq. (4.303)), ensuring strong continuity of the transport along any C^1 path and the well-posedness of the assembled kernel (see Definition 4.48 and standard references).

Dialectic algebra action on indices and intertwining

Corollary 4.77 (Basic identities in component form)**.** *For all $\alpha,\beta\in\mathcal{I}$ and loci x,x',*

$$\left(\hat{G}_\Lambda^{\alpha\beta}(x,x')\right)^\dagger = \hat{G}_\Lambda^{\beta\alpha}(x',x) \quad \textit{and} \quad \hat{G}_\Lambda^{\alpha\beta}(x,x') \mapsto g(x)\,\hat{G}_\Lambda^{\alpha\beta}(x,x')\,g(x')^\dagger,$$

i.e. Hermitian symmetry and gauge covariance hold componentwise. Reference: *These follow from Lemma 4.66 and Corollary 4.64 (equivalently summarized in Corollary 4.75).*

Definition 4.78 (Fourfold Dialectic Algebra and local index representations)**.** Let $\mathfrak{D}$ denote the Fourfold Dialectic Algebra acting on the mode-label space at each locus. A *local representation* at x is a $*$-representation $\rho_x : \mathfrak{D} \to \mathrm{End}(\mathbb{C}^{\mathcal{I}}_x)$. We say the Fourfold selectors transform covariantly under ρ_x if for each $D \in \mathfrak{D}$,

$$\hat{\mathcal{F}}_\alpha(x) \;\longmapsto\; \sum_{\mu\in\mathcal{I}} \rho_x(D)_{\alpha\mu}\, \hat{\mathcal{F}}_\mu(x), \qquad \hat{\mathcal{F}}^\dagger_\alpha(x) \;\longmapsto\; \sum_{\mu\in\mathcal{I}} \hat{\mathcal{F}}^\dagger_\mu(x)\, \rho_x(D)^\dagger_{\mu\alpha}.$$

(An analogous representation $\rho_{x'}$ acts at x'.)

Proposition 4.79 (Intertwining of index actions with the channel kernel)**.** *Assume the selector covariance of Definition 4.78. Then the 4×4 block $\hat{\mathbf{G}}_\Lambda(x,x') = [\hat{G}^{\alpha\beta}_\Lambda(x,x')]_{\alpha,\beta\in\mathcal{I}}$ transforms under left/right actions of $\mathfrak{D}$ as*

$$\hat{\mathbf{G}}_\Lambda(x,x') \;\longmapsto\; \rho_x(D)^\dagger\, \hat{\mathbf{G}}_\Lambda(x,x')\, \rho_{x'}(D),$$

i.e. the α-index carries the left action (by $\rho_x(D)^\dagger$) and the β-index carries the right action (by $\rho_{x'}(D)$). Componentwise,

$$\hat{G}^{\alpha\beta}_\Lambda(x,x') \;\longmapsto\; \sum_{\mu,\nu\in\mathcal{I}} \rho_x(D)^\dagger_{\mu\alpha}\, \hat{G}^{\mu\nu}_\Lambda(x,x')\, \rho_{x'}(D)_{\nu\beta}.$$

Moreover, this index-algebra covariance is independent of (and commutes with) gauge covariance on the fibre: $\hat{\mathbf{G}}_\Lambda(x,x') \mapsto g(x)\, \hat{\mathbf{G}}_\Lambda(x,x')\, g(x')^\dagger$ (Corollary 4.64).

Proof. Linear substitution using the selector transformation at x and x' and the definition $\hat{G}^{\alpha\beta}_\Lambda = \hat{\mathcal{F}}^\dagger_\alpha(x)\, \mathcal{U}(x \leftarrow x'; \gamma)\, \hat{\mathcal{F}}_\beta(x')$ (Definition 4.72) yields the component formula. Stacking components gives the block formula. Gauge covariance acts solely on the fibre operator $\mathcal{U}$, while $\rho_x, \rho_{x'}$ act on the index space, so the two actions commute. □

Remark 4.80 (Basis changes vs. gauge vs. dialectic algebra)**.** Three independent structures act on $\hat{\mathbf{G}}_\Lambda(x,x')$: (i) gauge on the fibre (endpoint conjugation), (ii) re-labelling/basis changes in $\mathbb{C}^{\mathcal{I}}$ (passive changes of $\mathcal{B}_x, \mathcal{B}_{x'}$; cf. Lemma 4.74), and (iii) active dialectic-algebra actions $\rho_x, \rho_{x'}$ on indices (this section). They compose without conflict and can be imposed in any order.

4.1.6 Block Matrix Representation Across Dialectic Modes

Definition 4.81 (Block matrix across dialectic modes)**.** With the ordered local frame $\mathcal{B}_x = \{|0\rangle_x, |1\rangle_x, |p\rangle_x, |s\rangle_x\}$, the Fourfold correlator admits the operator-valued block matrix form

$$\hat{\mathbf{G}}_\Lambda(x,x') = \begin{bmatrix} \hat{G}^{00} & \hat{G}^{01} & \hat{G}^{0p} & \hat{G}^{0s} \\ \hat{G}^{10} & \hat{G}^{11} & \hat{G}^{1p} & \hat{G}^{1s} \\ \hat{G}^{p0} & \hat{G}^{p1} & \hat{G}^{pp} & \hat{G}^{ps} \\ \hat{G}^{s0} & \hat{G}^{s1} & \hat{G}^{sp} & \hat{G}^{ss} \end{bmatrix}, \qquad \hat{G}^{\alpha\beta} := \hat{G}^{\alpha\beta}_\Lambda(x,x'). \tag{4.24}$$

The *Twofold* (Symmetry–Antisymmetry) sector is the principal 2×2 block

$$\hat{\mathbf{G}}_{(0,1)}(x,x') = \begin{bmatrix} \hat{G}^{00} & \hat{G}^{01} \\ \hat{G}^{10} & \hat{G}^{11} \end{bmatrix}, \tag{4.25}$$

while the remaining blocks encode mixing with Parasymmetry (p) and Holosymmetry (s).

Corollary 4.82 (Blockwise Hermitian symmetry and gauge covariance)**.** *The block matrix satisfies*

$$\hat{\mathbf{G}}_\Lambda(x,x')^\dagger = \hat{\mathbf{G}}_\Lambda(x',x).$$
$$\hat{\mathbf{G}}_\Lambda(x,x') \mapsto g(x)\,\hat{\mathbf{G}}_\Lambda(x,x')\,g(x')^\dagger.$$

with the componentwise statements

$$\left(\hat{G}^{\alpha\beta}_\Lambda(x,x')\right)^\dagger = \hat{G}^{\beta\alpha}_\Lambda(x',x),$$
$$\hat{G}^{\alpha\beta}_\Lambda(x,x') \mapsto g(x)\,\hat{G}^{\alpha\beta}_\Lambda(x,x')\,g(x')^\dagger.$$

holding for all $\alpha,\beta \in \{0,1,p,s\}$ (Lemma 4.66 and Corollaries 4.64 and 4.75).

Remark 4.83 (Interpretation of matrix components)**.** With the block representation Definition 4.81 and eq. (4.24), the entries of $\hat{\mathbf{G}}_\Lambda(x,x')$ admit the following operational reading (cf. Remark 4.69):

Diagonal terms $\hat{G}^{\alpha\alpha}$:

Modal persistence—amplitudes preserving the dialectic mode along $x' \to x$ (self-coherence).

Off-diagonal terms $\hat{G}^{\alpha\beta}$, $\alpha \neq \beta$:

Dialectic transitions—mode-changing/interference amplitudes (e.g., $0 \leftrightarrow p$, $1 \leftrightarrow s$). These transform covariantly under gauge $\hat{G}^{\alpha\beta}_\Lambda \mapsto g(x)\hat{G}^{\alpha\beta}_\Lambda g(x')^\dagger$ (Corollary 4.82) and obey Hermitian symmetry $\left(\hat{G}^{\alpha\beta}_\Lambda\right)^\dagger = \hat{G}^{\beta\alpha}_\Lambda(x',x)$ (Lemma 4.66).

Block structure and algebra: (i) The Twofold subspace $\{0,1\}$ reproduces the classical dyad, i.e. the principal 2×2 block $\hat{\mathbf{G}}_{(0,1)}(x,x')$ (eq. (4.25)).

(ii) Cross-level blocks $\{0,1\} \leftrightarrow \{p,s\}$ implement algebraic intertwiners that couple the dyad to emergent modes, governing coherence flow and global (holosymmetric) information (cf. Definition 2.94 and eq. (3.36) and the index-action formalism in Definition 4.78 and Proposition 4.79).

Caution. As noted in Remark 4.71, $|p\rangle_x$ and $|s\rangle_x$ are not defined as linear combinations of $|0\rangle_x, |1\rangle_x$. The equal superposition $|+\rangle_x$ generates the parasymmetric *effect* $P^{(x)}_{\text{para}} = |+\rangle_x\langle+|$ within the Twofold sector, and is not identified with $|p\rangle_x$.

Definition 4.84 (Dialectic-complete selector system at an intermediate locus)**.** We say the selector family at y is *dialectically complete* (on the regulated fibre) if

$$\sum_{\gamma\in\{0,1,p,s\}} \hat{\mathcal{F}}_\gamma(y)\,\hat{\mathcal{F}}^\dagger_\gamma(y) = \mathbf{1} \quad \text{on } \operatorname{Ran} E^{(y)}([0,\lambda_{\text{cutoff}}]).$$

Proposition 4.85 (Block composition through an intermediate locus (conditional))**.** *Let $x \leftarrow y \leftarrow x'$ with paths γ_1,γ_2 such that $\gamma = \gamma_1\circ\gamma_2$. If the selector system at y is dialectically complete (Definition 4.84), then*

$$\hat{\mathbf{G}}_\Lambda(x,x';\gamma) \;=\; \hat{\mathbf{G}}_\Lambda(x,y;\gamma_1)\,\hat{\mathbf{G}}_\Lambda(y,x';\gamma_2),$$

i.e. block multiplication coincides with channel factorization. Componentwise, $\hat{G}^{\alpha\beta}_\Lambda(x,x') = \sum_\gamma \hat{G}^{\alpha\gamma}_\Lambda(x,y)\,\hat{G}^{\gamma\beta}_\Lambda(y,x')$. Reference: *Transport functoriality Proposition 4.52 and the tensor definition Definition 4.72.*

Remark 4.86 (Sector projections and classical slice)**.** Let $P_{(0,1)} = \mathrm{diag}(1,1,0,0)$ in the $\mathcal{I}$-ordering; then $P_{(0,1)}\,\hat{\mathbf{G}}_\Lambda(x,x')\,P_{(0,1)} = \hat{\mathbf{G}}_{(0,1)}(x,x')$ (eq. (4.25)). In regimes where interrogations commute or decoherence suppresses coherence across sectors, the off-diagonal blocks vanish (operationally), and the kernel *collapses* to the Twofold block—recovering the classical slice (cf. the discussion following Remark 4.69).

Duality-Collapse Regime at a Locus

Definition 4.87 (Duality-Collapse Regime at a Locus)**.** Fix a locus $\mathfrak{B}_x$. Write the modal fiber as $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)} = \mathrm{span}\{|0\rangle_x, |1\rangle_x, |p\rangle_x, |s\rangle_x\}$, with the Fourfold Inter-Modal Correlator $\hat{G}_\Lambda^{\alpha\beta}(x,x')$ and its Twofold restriction $\hat{G}_{(0,1)}$ as in Section 4.1.2 and Definition 4.1. We say that the *duality-collapse regime* holds at x (relative to a Holo-State $|\Omega_\Psi\rangle$) if:

(D1) **Block-diagonality (decoupling).** The off-block expectations vanish (exactly or within a fixed tolerance ϵ):

$$\langle\Omega_\Psi|\hat{G}_\Lambda^{\alpha\beta}(x,x')|\Omega_\Psi\rangle \approx 0 \quad \text{for all } \alpha\in\{0,1\},\ \beta\in\{p,s\} \text{ and } \alpha\leftrightarrow\beta,$$

uniformly for $x'\approx x$ along admissible transports.

(D2) **Inner-product realization.** The Twofold block induces a positive sesquilinear form on $\mathcal{H}_{\mathrm{mod}\,(0,1)}^{(\mathfrak{B}_x)} := \mathrm{span}\{|0\rangle_x, |1\rangle_x\}$ which is nondegenerate in the coincident limit and hence defines a fiberwise inner product.

Remark 4.88 (Meaning and consequences of the duality–collapse regime)**.** Under Definition 4.87, condition (D1) enforces expectation-level *block decoupling* between the Twofold sector $\{0,1\}$ and the emergent sector $\{p,s\}$:

$$\langle\Omega_\Psi|\hat{G}_\Lambda^{\alpha\beta}(x,x')|\Omega_\Psi\rangle \approx 0 \quad \text{for} \quad \alpha\in\{0,1\},\ \beta\in\{p,s\} \text{ and } \alpha\leftrightarrow\beta,$$

uniformly for $x'\approx x$. Hence, near x, the Born-evaluated Fourfold kernel reduces to its Twofold block $\hat{\mathbf{G}}_{(0,1)}(x,x')$ (cf. eq. (4.25)), up to $O(\epsilon)$.

Condition (D2) asserts that in the coincident limit $x'\to x$ this Twofold block induces a positive, nondegenerate sesquilinear form on $\mathcal{H}_{\mathrm{mod}\,(0,1)}^{(\mathfrak{B}_x)} = \mathrm{span}\{|0\rangle_x, |1\rangle_x\}$, thereby *realizing the local inner product* from the kernel (cf. *eq.* (4.6)). Together, (D1)–(D2) implement the "classical slice": Non-Contradiction/Excluded-Middle re-emerge on the dyad, while parasymmetric/holosymmetric channels carry no independent signal at x (cf. the narrative in Remark 4.69).

Gauge and transport notes. The statements are framed at the level of expectations, which are scalar and hence gauge invariant (see Corollary 4.82). The uniformity in $x'\approx x$ "along admissible transports" ensures locality and stability of the collapse under those path choices; no flatness assumption is required, only the expectation-level decoupling specified in (D1).

Corollary 4.89 (Effective Twofold kernel in the duality–collapse regime)**.** *Let the duality–collapse regime hold at $\mathfrak{B}_x$ relative to the Holo–State $|\Omega_\Psi\rangle$ in the sense of Definition 4.87. Define the Born–evaluated Fourfold kernel*

$$\overline{\mathbf{G}}_\Lambda(x,x') \;:=\; \Big[\ \langle\Omega_\Psi \mid \hat{G}_\Lambda^{\alpha\beta}(x,x') \mid \Omega_\Psi\rangle\ \Big]_{\alpha,\beta\in\{0,1,p,s\}}.$$

Then, uniformly for $x'\approx x$ along admissible transports:

(i) Cross–sector suppression
$\langle \Omega_\Psi \mid \hat{G}^{\alpha\beta}_\Lambda(x,x') \mid \Omega_\Psi \rangle = O(\epsilon)$ *for all* $\alpha \in \{0,1\},\ \beta \in \{p,s\}$ *and vice versa.*

(ii) Projection to the Twofold block
Let $P_{(0,1)} := \mathrm{diag}(1,1,0,0)$ *in the* $\mathcal{I}$*-ordering. Then*

$$\overline{\mathbf{G}}_\Lambda(x,x') \;=\; P_{(0,1)}\,\overline{\mathbf{G}}_\Lambda(x,x')\,P_{(0,1)} \;+\; O(\epsilon),$$

so, up to $O(\epsilon)$*, the effective Born kernel reduces to its Twofold block* $\hat{\mathbf{G}}_{(0,1)}(x,x')$ *of eq. (4.25).*

(iii) Inner product on the dyad (coincident limit)
As $x' \to x$*, the Twofold block* $\overline{\mathbf{G}}_{(0,1)}(x,x)$ *defines a positive, nondegenerate sesquilinear form on* $\mathrm{span}\{|0\rangle_x, |1\rangle_x\}$ *(by* (D2) *of Definition 4.87), hence a fibrewise inner product.*

(iv) Observable stability under projection
For any coefficient vectors $c^{(1)}, c^{(2)} \in \mathbb{C}^{\{0,1,p,s\}}$*,*

$$\Big(c^{(1)}\Big)^\dagger \overline{\mathbf{G}}_\Lambda(x,x')\,c^{(2)} \;=\; \Big(P_{(0,1)}c^{(1)}\Big)^\dagger \overline{\mathbf{G}}_\Lambda(x,x')\,\Big(P_{(0,1)}c^{(2)}\Big) \;+\; O(\epsilon),$$

so Born observables built from cross–sector blocks are $O(\epsilon)$ *near* x*.*

Gauge note. *All statements are formulated at the expectation level and are thus gauge invariant (cf. Corollary 4.82).*

Proposition 4.90 (Recovery of Ordinary Two-Level Quantum Kinematics)**.** *Under Definition 4.87, there exists an isometry* $U : \mathbb{C}^2 \hookrightarrow \mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}\,(0,1)}$ *such that for every Twofold–respecting effect* $E_{(0,1)}$ *and state* ρ *on the full fibre,*

$$\mathrm{Tr}\Big(E_{(0,1)}\,\rho\Big) \;=\; \mathrm{Tr}\Big(\,U^\dagger E_{(0,1)} U\,\rho_{(0,1)}\Big), \qquad \rho_{(0,1)} := U^\dagger \rho\, U,$$

i.e. all operational predictions on the Twofold sector coincide with those of an ordinary qubit state $\rho_{(0,1)} \in \mathcal{B}(\mathbb{C}^2)$ *with the standard Born rule.*

Proof of Proposition 4.90. Step 0 (Twofold sector and projection). Let $P_{(0,1)} := \mathrm{diag}(1,1,0,0)$ in the $\mathcal{I}$-ordering of $\mathcal{B}_x = \{|0\rangle_x, |1\rangle_x, |p\rangle_x, |s\rangle_x\}$ (Definition 4.81 and eq. (4.24)). By "Twofold–respecting effect" we mean an operator supported on the Twofold sector, $E_{(0,1)} = P_{(0,1)}E_{(0,1)}P_{(0,1)}$. Expectation–level gauge invariance holds throughout (Corollary 4.82).

Step 1 (Inner product from the kernel and orthonormalization). By (D2) of Definition 4.87 and the inner–product construction from the Fourfold kernel (cf. eq. (4.6) and Definition 4.73 and eq. (4.25)), the coincident–limit Twofold block induces a positive, nondegenerate sesquilinear form on $\mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}\,(0,1)} = \mathrm{span}\{|0\rangle_x, |1\rangle_x\}$. Choose an orthonormal pair $\{|e_0\rangle_x, |e_1\rangle_x\}$ for this form. Define $U : \mathbb{C}^2 \to \mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}\,(0,1)}$ by $U|0\rangle = |e_0\rangle_x$, $U|1\rangle = |e_1\rangle_x$. Then $U^\dagger U = \mathbf{1}_{\mathbb{C}^2}$ and $UU^\dagger = P_{(0,1)}$ on the Twofold sector, so U is an isometry.

Step 2 (Decoupling of cross–sector expectations). By (D1) of Definition 4.87 and Corollary 4.89, Born–evaluated cross–sector entries vanish (exactly, or up to $O(\epsilon)$) uniformly for $x' \approx x$. Consequently, for any state ρ and Twofold–respecting effect $E_{(0,1)}$,

$$\mathrm{Tr}\Big(E_{(0,1)}\,\rho\Big) = \mathrm{Tr}\Big(E_{(0,1)}\,P_{(0,1)}\rho P_{(0,1)}\Big)$$
(exact in the ideal regime; $+\,O(\epsilon)$ in the approximate one). (4.26)

Step 3 (Compression to $\mathbb{C}^2$ via the isometry). Using $UU^\dagger = P_{(0,1)}$,

$$\mathrm{Tr}\Big(E_{(0,1)}\, P_{(0,1)}\rho P_{(0,1)}\Big) = \mathrm{Tr}\Big(E_{(0,1)}\, UU^\dagger \rho\, UU^\dagger\Big) = \mathrm{Tr}\Big(U^\dagger E_{(0,1)} U \;\cdot\; U^\dagger \rho\, U\Big),$$

where we used cyclicity of the trace and $U^\dagger U = \mathbf{1}$. Set $\rho_{(0,1)} := U^\dagger \rho\, U \in \mathcal{B}(\mathbb{C}^2)$. Thus

$$\mathrm{Tr}\Big(E_{(0,1)}\,\rho\Big) = \mathrm{Tr}\Big(U^\dagger E_{(0,1)} U\;\rho_{(0,1)}\Big) \quad \text{(again, exactly or up to } O(\epsilon) \text{ as in Step 2).}$$

Step 4 (Standard qubit kinematics). The map $A \mapsto U^\dagger A U$ is a $*$-homomorphism on the Twofold corner and transports effects to $\mathcal{B}(\mathbb{C}^2)$. Hence, all Twofold operational predictions coincide with those obtained from the qubit state $\rho_{(0,1)}$ via the usual Born rule on $\mathbb{C}^2$. (Equivalently, in the GNS picture of Chapter 1, Born expectations are represented by traces in the anchored representation; see Proposition 4.17.)

This proves the claim. □

Remark 4.91 (Approximate regime and stability)**.** If (D1) holds with tolerance $\epsilon > 0$, the equality in Proposition 4.90 holds up to $O(\epsilon)$:

$$\mathrm{Tr}\Big(E_{(0,1)}\,\rho\Big) = \mathrm{Tr}\Big(U^\dagger E_{(0,1)} U\,\rho_{(0,1)}\Big) \;+\; O(\epsilon),$$

uniformly for $x' \approx x$ along admissible transports (Definition 4.87 and Corollary 4.89).

Remark 4.92 (Hierarchy of Collapses)**.** There are two conceptually distinct "collapse'' regimes, ordered by strength:

(A) Duality–collapse (quantum two–level limit). As in Definition 4.87, conditions (D1)–(D2) enforce (i) expectation–level decoupling between the Twofold sector $\{0,1\}$ and the emergent sector $\{p,s\}$, and (ii) realization of a positive, nondegenerate inner product on the Twofold fibre. Operationally, the Born–evaluated Fourfold kernel reduces (up to $O(\epsilon)$) to the Twofold block (Corollary 4.89), and *qubit kinematics* are recovered via the isometry $U : \mathbb{C}^2 \hookrightarrow \mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}\,(0,1)}$ (Proposition 4.90). Crucially, no commutativity inside the Twofold sector is assumed: off–diagonal Twofold amplitudes may persist, so intra–Twofold coherence is retained.

(B) Classical collapse (Boolean two–valued limit). This is a stronger, context–dependent limit in which, in addition to (D1)–(D2), *interrogations within the Twofold sector commute* (or decoherence fully suppresses the Twofold off–diagonals). Equivalently, there exist Twofold projectors P_T, P_A with

$$P_T P_A = 0, \qquad P_T + P_A = P_{(0,1)},$$

so Non–Contradiction and Excluded Middle hold on the Twofold slice (cf. the discussion surrounding the Twofold completeness statement in the prismatic/duality axioms). In this regime the Twofold block is diagonal in a common eigenbasis, and the Fourfold reduces to a classical two–valued "fortress'' (no interference anywhere).

Logical/operational ordering.

$$\begin{aligned} &\text{Fourfold (fully coherent)} \\ \Longrightarrow\ &\text{Duality–collapse (quantum Twofold)} \\ \Longrightarrow\ &\text{Classical collapse (Boolean Twofold).} \end{aligned}$$

The first arrow is a *representation–level reduction* (expectation–level decoupling and inner–product realization) that preserves Twofold coherence; the second arrow additionally removes intra–Twofold coherence by commutativity or complete dephasing. Typical generators of the second arrow include a Twofold dephasing channel $\mathcal{D}_{(0,1)}(\rho) = \sum_{\alpha\in\{T,A\}} P_\alpha \rho P_\alpha$, which enforces $P_T P_A = 0$ at the level of statistics.

Gauge/transport notes. Both regimes are framed at the level of expectations and are therefore gauge invariant (endpoint conjugations cancel; cf. Corollary 4.82). Uniformity in $x' \approx x$ "along admissible transports'' ensures locality and stability of the limiting behaviour; no flatness assumption is required.

Corollary 4.93 (Compatibility with the Atomic Prismatic)**.** *The passage to the Twofold sector is a* representation–level restriction *and does not alter the atomic prismatic unit* $\mathring{\mathbf{\Lambda}}$ *of Definition 1.5. In particular, properties* (a) No redefinition *and* (b) Closure as a unit *remain intact, and the invariance stated in Proposition 1.8 descends to the Twofold realization.*

Proof of Corollary 4.93. Let π denote the anchored (GNS) representation used for Born evaluation (cf. Proposition 4.17), and let $U : \mathbb{C}^2 \hookrightarrow \mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}\,(0,1)}$ be the isometry from Proposition 4.90. Define the compressed Twofold representation

$$\pi_{(0,1)}(\,\cdot\,) \;:=\; U^\dagger\,\pi(\,\cdot\,)\,U \;:\; \text{Prismatic algebra} \longrightarrow \mathcal{B}(\mathbb{C}^2).$$

Then:

(i) *No redefinition.* The algebraic element $\mathring{\mathbf{\Lambda}}$ is unchanged; only its *representation* is restricted. Thus $\pi_{(0,1)}$ is obtained from π by isometric compression, not by modifying the algebra or its unit.

(ii) *Closure as a unit.* Since U is an isometry and π is unital,

$$\pi_{(0,1)}\Big(\mathring{\mathbf{\Lambda}}\Big) = U^\dagger\,\pi\Big(\mathring{\mathbf{\Lambda}}\Big)\,U = U^\dagger\,\mathbf{1}\,U = \mathbf{1}_{\mathbb{C}^2},$$

so the unit remains the unit after restriction.

(iii) *Descent of invariance.* For the duality automorphism D appearing in Proposition 1.8, we have

$$\pi_{(0,1)}\Big(\mathsf{D}(A)\Big) = U^\dagger \pi\Big(\mathsf{D}(A)\Big) U = U^\dagger\Big(\mathsf{D}\cdot\pi\Big)(A)\,U = \Big(\mathsf{D}\cdot\pi_{(0,1)}\Big)(A),$$

because compression by an isometry commutes with algebraic relations preserved by π. Hence the invariance identity of Proposition 1.8 holds verbatim for $\pi_{(0,1)}$, i.e. it *descends* to the Twofold realization.

This establishes the claim. □

Intuitive Understanding. The *Fourfold Inter-Modal Correlator* is best viewed as a gauge–covariant, four–channel, operator–valued propagator. It records how the four dialectic modes—*Symmetry (0)*, *Antisymmetry (1)*, *Parasymmetry (p)*, and *Holosymmetry (s)*—*transform and interfere* between two loci x' and x in the emergent configuration space $\mathcal{X}_q$

via the gauge–covariant transporter $\mathcal{U}(x \leftarrow x'; \gamma)$ (Definitions 4.48 and 4.61, Proposition 4.54, and Corollary 4.64).

At its core, the correlator is a 4×4 matrix of channels. Each entry $\hat{G}_\Lambda^{\alpha\beta}(x, x')$ is the operator for the composite process: *imprint* mode β at x' using $\hat{\mathcal{F}}_\beta(x')$, *parallel–transport* to x with $\mathcal{U}(x \leftarrow x'; \gamma)$, then *read out* mode α at x using $\hat{\mathcal{F}}_\alpha^\dagger(x)$ (Definition 4.61). The block form $\hat{\mathbf{G}}_\Lambda(x, x')$ makes this structure explicit (eq. (4.24)). Its top–left 2×2 block is the *Twofold Correlator*, governing the classical dyad (Symmetry/Antisymmetry) (eq. (4.25)); the remaining blocks encode couplings of this dyad to the higher modes—coherent composition (Parasymmetry) and system–level invariants (Holosymmetry) (Remark 4.83, Definition 2.94, and eq. (3.36)).

Interpreting the entries gives a dynamic picture. Diagonal terms $\hat{G}^{\alpha\alpha}$ capture *modal persistence*—stability of a facet under transport—while off–diagonals $\hat{G}^{\alpha\beta}$ with $\alpha \neq \beta$ capture *modal transitions*—conversion or interference between facets (e.g., $0 \leftrightarrow p$, $1 \leftrightarrow s$) (Remark 4.83). As a whole, $\hat{\mathbf{G}}_\Lambda$ functions as a *relational kernel*: it exposes both local interference within the Twofold sector and cross–level couplings to parasymmetric and holosymmetric structure, while respecting Hermitian symmetry and gauge covariance ($\hat{\mathbf{G}}_\Lambda(x, x')^\dagger = \hat{\mathbf{G}}_\Lambda(x', x)$ and $\hat{\mathbf{G}}_\Lambda \mapsto g(x)\hat{\mathbf{G}}_\Lambda g(x')^\dagger$; see Lemma 4.66 and Corollary 4.82).

The correlator is deliberately *operator–valued*: it does not prematurely collapse quantum structure. By carrying residual degrees of freedom as operators, it lets *context*—state, locus, gauge—enter only at the final stage via expectations (GNS–anchored Born evaluation) (Proposition 4.17 and Definition 4.73). In the coincident limit $x' \to x$, expectations in the Holo–State produce Born–valued amplitudes that *define the inner product on the emergent modal fibre* (cf. (4.6)), so the correlator is not merely descriptive; it is *generative* of the local Hilbert geometry.

Classical reasoning appears as a limiting slice. When interrogations commute or decoherence suppresses phase information, cross–sector structure becomes inert and the 4×4 kernel *collapses to the Twofold block*, recovering the Thesis/Antithesis opposition (the "classical collapse''). By contrast, the *duality–collapse* regime only decouples $\{0, 1\}$ from $\{p, s\}$ at the level of expectations while retaining quantum coherence within the Twofold subspace—thereby recovering ordinary qubit kinematics without imposing commutativity (Definition 4.87, Corollary 4.89, Proposition 4.90, and Remark 4.92). In regimes where coherence and global invariants matter, the full Fourfold kernel is indispensable: a gauge–aligned, multi–port propagator that tells how logical facets persist, convert, and entangle across loci—and, through its expectations, carves the geometry of the emergent quantum space they inhabit (Proposition 4.54, Definition 4.48, and Remark 4.83).

4.1.7 Algebraic Structure and Graded Dynamics of the Fourfold Inter-Modal Correlator

Dialectic grading of modal indices

Definition 4.94 (Dialectic $\mathbb{Z}_2^2$-grading of modes and channel components)**.** Let the mode set be $\mathcal{I} = \{0, 1, p, s\}$ (Symmetry, Antisymmetry, Parasymmetry, Holosymmetry). Assign a

bidegree

$$\deg(0) = (0,0), \qquad \deg(1) = (1,0), \qquad \deg(p) = (1,1), \qquad \deg(s) = (0,1),$$

so that p behaves algebraically as the composite of the two fundamental oppositions 1 and s. For channel components $\hat{G}^{\alpha\beta}(x,x')$ (cf. eq. (4.22)), define the *colored matrix–unit* grading by

$$\deg\left(\hat{G}^{\alpha\beta}\right) := \deg(\alpha) + \deg(\beta) \ \in \ \mathbb{Z}_2 \times \mathbb{Z}_2. \tag{4.27}$$

Whenever the product XY of homogeneous components is defined (index–wise), degrees add under composition:

$$\deg(XY) = \deg(X) + \deg(Y) \ \in \ \mathbb{Z}_2 \times \mathbb{Z}_2.$$

Lemma 4.95 (Even degree implies bracket equivalence)**.** *If* $\deg A = (0,0)$*, then for every homogeneous* B *one has* $[A,B]_{\mathrm{gr}} = [A,B]$.

Proof. Immediate from the definition $[A,B]_{\mathrm{gr}} = AB - (-1)^{\langle \deg A, \deg B\rangle} BA$ and $\langle (0,0), \deg B\rangle = 0$. □

Corollary 4.96. *If* $\deg A = (0,0)$*, then* $[A,B]_{\mathrm{gr}} = [A,B]$ *for every (not necessarily homogeneous)* B.

Proof. Decompose $B = \sum_j B_j$ into homogeneous components and use linearity with Lemma 4.95. □

Mode	Degree $\deg(\alpha)$	Heuristic role
0 (Symmetry)	$(0,0)$	identity sector
1 (Antisymmetry)	$(1,0)$	primary opposition
p (Parasymmetry)	$(1,1)$	composite of oppositions
s (Holosymmetry)	$(0,1)$	independent global duality

Definition 4.97 (Dialectic grading on channels)**.** With deg as above, the block matrix $\hat{\mathbf{G}}_\Lambda(x,x')$ (Definition 4.81 and eq. (4.24)) is a $\mathbb{Z}_2^2$-graded matrix algebra by componentwise degree $\deg(\hat{G}^{\alpha\beta}) = \deg(\alpha) + \deg(\beta)$.

Lemma 4.98 (Compatibility of grading with adjoint, gauge, and composition)**.** *For all* $\alpha, \beta \in \mathcal{I}$ *and all* x, x'*:*

(i) Adjoint
$\left(\hat{G}^{\alpha\beta}_\Lambda(x,x')\right)^\dagger = \hat{G}^{\beta\alpha}_\Lambda(x',x)$ *(Lemma 4.66), and* $\deg\left((\hat{G}^{\alpha\beta})^\dagger\right) = \deg(\alpha) + \deg(\beta) = \deg(\hat{G}^{\alpha\beta})$.

(ii) Gauge
Under $\hat{G}^{\alpha\beta}_\Lambda(x,x') \mapsto g(x)\hat{G}^{\alpha\beta}_\Lambda(x,x')g(x')^\dagger$ *(Corollaries 4.64 and 4.82), the degree is unchanged.*

(iii) Composition

If $x \leftarrow y \leftarrow x'$ and the selector family at y is dialectically complete (Definition 4.84), then

$$\hat{G}^{\alpha\beta}_{\Lambda}(x,x') = \sum_{\gamma\in\mathcal{I}} \hat{G}^{\alpha\gamma}_{\Lambda}(x,y)\,\hat{G}^{\gamma\beta}_{\Lambda}(y,x') \quad \textit{(Proposition 4.85)},$$

and each summand satisfies $\deg\left(\hat{G}^{\alpha\gamma}\hat{G}^{\gamma\beta}\right) = \deg(\alpha) + \deg(\beta)$.

Proof of Lemma 4.98. We prove (i)–(iii) in turn, using the grading of Definition 4.94 and the channel/operator laws established in Chapter 4.

(i) Adjoint. By Hermitian symmetry of channels, $\left(\hat{G}^{\alpha\beta}_{\Lambda}(x,x')\right)^{\dagger} = \hat{G}^{\beta\alpha}_{\Lambda}(x',x)$ (Lemma 4.66). Hence, using the definition of the bidegree (Definition 4.94),

$$\deg\Big((\hat{G}^{\alpha\beta})^{\dagger}\Big) = \deg(\beta) + \deg(\alpha) = \deg(\alpha) + \deg(\beta) = \deg\Big(\hat{G}^{\alpha\beta}\Big),$$

since addition in $\mathbb{Z}_2^2$ is commutative.

(ii) Gauge. Under a local gauge $g : \mathcal{X}_q \to \mathrm{U}(\mathcal{H}_{\mathrm{em}})$, each channel transforms by endpoint conjugation, $\hat{G}^{\alpha\beta}_{\Lambda}(x,x') \mapsto g(x)\,\hat{G}^{\alpha\beta}_{\Lambda}(x,x')\,g(x')^{\dagger}$ (Corollary 4.64). Because the grading depends only on the *mode labels* (α,β) and not on fibrewise conjugation (Definition 4.94), one has $\deg\Big(g(x)\,\hat{G}^{\alpha\beta}_{\Lambda}(x,x')\,g(x')^{\dagger}\Big) = \deg\Big(\hat{G}^{\alpha\beta}_{\Lambda}(x,x')\Big)$. Thus degree is gauge invariant.

(iii) Composition. Let $x \leftarrow y \leftarrow x'$ and pick composable path segments so that $\gamma = \gamma_1 \circ \gamma_2$ with $x \xleftarrow{\gamma_1} y$ and $y \xleftarrow{\gamma_2} x'$. By transport functoriality (Proposition 4.52 and eq. (4.14)), $\mathcal{U}(x \leftarrow x';\gamma) = \mathcal{U}(x \leftarrow y;\gamma_1)\,\mathcal{U}(y \leftarrow x';\gamma_2)$. Insert at the intermediate locus y the dialectically complete resolution of the identity ($\sum_{\mu\in\mathcal{I}} \hat{\mathcal{F}}_{\mu}(y)\hat{\mathcal{F}}^{\dagger}_{\mu}(y) = \mathbf{1}$; cf. Definition 4.84) into the Fourfold channel definition (Definition 4.61):

$$\begin{aligned}
\hat{G}^{\alpha\beta}_{\Lambda}(x,x') &= \hat{\mathcal{F}}^{\dagger}_{\alpha}(x)\,\mathcal{U}(x \leftarrow x';\gamma)\,\hat{\mathcal{F}}_{\beta}(x') \\
&= \hat{\mathcal{F}}^{\dagger}_{\alpha}(x)\,\mathcal{U}(x \leftarrow y;\gamma_1)\left(\sum_{\mu\in\mathcal{I}} \hat{\mathcal{F}}_{\mu}(y)\hat{\mathcal{F}}^{\dagger}_{\mu}(y)\right)\mathcal{U}(y \leftarrow x';\gamma_2)\,\hat{\mathcal{F}}_{\beta}(x') \\
&= \sum_{\mu\in\mathcal{I}} \Big[\hat{\mathcal{F}}^{\dagger}_{\alpha}(x)\,\mathcal{U}(x \leftarrow y;\gamma_1)\,\hat{\mathcal{F}}_{\mu}(y)\Big]\Big[\hat{\mathcal{F}}^{\dagger}_{\mu}(y)\,\mathcal{U}(y \leftarrow x';\gamma_2)\,\hat{\mathcal{F}}_{\beta}(x')\Big] \\
&= \sum_{\mu\in\mathcal{I}} \hat{G}^{\alpha\mu}_{\Lambda}(x,y)\,\hat{G}^{\mu\beta}_{\Lambda}(y,x'),
\end{aligned}$$

which is the claimed block–composition identity (Proposition 4.85).

For the degree of each summand, use Definition 4.94:

$$\begin{aligned}
\deg\Big(\hat{G}^{\alpha\mu}\hat{G}^{\mu\beta}\Big) &= \deg(\hat{G}^{\alpha\mu}) + \deg(\hat{G}^{\mu\beta}) \\
&= \Big(\deg\alpha + \deg\mu\Big) + \Big(\deg\mu + \deg\beta\Big) \\
&= \deg\alpha + \underbrace{2\,\deg\mu}_{=0} + \deg\beta,
\end{aligned} \tag{4.28}$$

where $2\deg\mu = 0$ in $\mathbb{Z}_2^2$. Hence $\deg\Big(\hat{G}^{\alpha\mu}\hat{G}^{\mu\beta}\Big) = \deg(\alpha) + \deg(\beta)$, as stated.

This completes the proof. □

Proposition 4.99 (Selection rules and sector patterns)**.** *In the ordered basis* $\mathcal{B}_x = \{|0\rangle_x, |1\rangle_x, |p\rangle_x, |s\rangle_x\}$*:*

(a) *The Twofold block* $\hat{\mathbf{G}}_{(0,1)}(x,x')$ *(eq. (4.25)) contains only degrees* $(0,0)$ *(diagonals) and* $(1,0)$ *(off–diagonals).*

(b) *Blocks that couple* $\{0,1\}$ *to* $\{p,s\}$ *carry degrees with second component* 1 *(they probe the holosymmetric axis).*

(c) *Under block composition through* y *(Proposition 4.85), every matrix element* $\hat{G}^{\alpha\beta}_{\Lambda}(x,x')$ *has total degree* $\deg(\alpha)+\deg(\beta)$ *independently of the intermediate index* γ*.*

Proof of Proposition 4.99. Throughout we use the $\mathbb{Z}_2^2$-grading of Definition 4.94 with $\deg(0) = (0,0)$, $\deg(1) = (1,0)$, $\deg(p) = (1,1)$, $\deg(s) = (0,1)$, and $\deg\big(\hat{G}^{\alpha\beta}\big) = \deg(\alpha)+\deg(\beta)$.

(a) Twofold block degrees. In the Twofold block $\alpha,\beta \in \{0,1\}$ (eq. (4.25)). For diagonals $\alpha=\beta$, $\deg(\hat{G}^{\alpha\alpha}) = \deg(\alpha)+\deg(\alpha) = 2\deg(\alpha) = (0,0)$ in $\mathbb{Z}_2^2$. For off–diagonals, $\deg(\hat{G}^{01}) = (0,0)+(1,0) = (1,0)$ and $\deg(\hat{G}^{10}) = (1,0)+(0,0) = (1,0)$. Hence the block contains only $(0,0)$ on the diagonal and $(1,0)$ off–diagonal.

(b) Couplings $\{0,1\} \leftrightarrow \{p,s\}$. Take $\alpha \in \{0,1\}$ and $\beta \in \{p,s\}$ (or vice versa). The second component of the degree is $\deg_2(\alpha)+\deg_2(\beta) \pmod 2$. Since $\deg_2(0) = \deg_2(1) = 0$ and $\deg_2(p) = \deg_2(s) = 1$, one has $\deg_2(\hat{G}^{\alpha\beta}) = 1$. Thus every such block element has second component 1, i.e. it probes the holosymmetric axis.

(c) Degree under block composition. By the block–composition identity through an intermediate locus y (Proposition 4.85),

$$\hat{G}^{\alpha\beta}_{\Lambda}(x,x') = \sum_{\gamma\in\mathcal{I}} \hat{G}^{\alpha\gamma}_{\Lambda}(x,y)\,\hat{G}^{\gamma\beta}_{\Lambda}(y,x').$$

For each summand, Lemma 4.98(iii) gives

$$\begin{aligned}\deg\big(\hat{G}^{\alpha\gamma}\hat{G}^{\gamma\beta}\big) &= \deg(\hat{G}^{\alpha\gamma}) + \deg(\hat{G}^{\gamma\beta}) \\ &= \big(\deg\alpha + \deg\gamma\big) + \big(\deg\gamma + \deg\beta\big) \\ &= \deg\alpha + \underbrace{2\,\deg\gamma}_{=0} + \deg\beta, \end{aligned} \tag{4.29}$$

so $\deg\big(\hat{G}^{\alpha\gamma}\hat{G}^{\gamma\beta}\big) = \deg(\alpha)+\deg(\beta)$, independently of γ. This establishes the stated selection rule for every matrix element of the composed block.

Items (a)–(c) together prove the proposition. □

Definition 4.100 (Graded sign and graded commutator (optional))**.** Equip $\mathbb{Z}_2^2$ with the standard bilinear form $\langle (a_1,a_2),(b_1,b_2)\rangle := a_1b_1 + a_2b_2 \pmod 2$. For homogeneous components X, Y define

$$[X,Y]_{\mathrm{g}} := XY - (-1)^{\langle \deg X, \deg Y\rangle} YX.$$

Then $[\,\cdot,\cdot\,]_{\mathrm{g}}$ is bilinear, graded–antisymmetric, and satisfies the graded Jacobi identity on the associative algebra generated by the $\hat{G}^{\alpha\beta}$.

Remark 4.101 (What the grading means (and what it does not))**.** The $\mathbb{Z}_2^2$–grading is a bookkeeping structure on *mode labels* and their channels, not a fermionic parity on physical operators. It organizes how components mix under composition, adjoint, and gauge (Lemma 4.98) and clarifies which blocks probe parasymmetric vs. holosymmetric content (Remark 4.83). In particular, the graded selection rules in Proposition 4.99 are compatible with transport functoriality (Proposition 4.52), block composition (Proposition 4.85), and Hermitian/gauge properties (Lemma 4.66 and Corollary 4.82).

Corollary 4.102 (Adjoint and block grading)**.** *For the graded components of Definition 4.94 one has:*

(i) Adjoint preserves degree

Using Hermitian symmetry $\big(\hat{G}^{\alpha\beta}(x,x')\big)^\dagger = \hat{G}^{\beta\alpha}(x',x)$ *(Lemma 4.66),*

$$\deg\Big((\hat{G}^{\alpha\beta})^\dagger\Big) = \deg(\beta) + \deg(\alpha) = \deg(\alpha) + \deg(\beta) = \deg\Big(\hat{G}^{\alpha\beta}\Big).$$

(ii) Diagonal evenness

For $\alpha = \beta$, $\deg(\hat{G}^{\alpha\alpha}) = \deg(\alpha) + \deg(\alpha) = 0$ *in* $\mathbb{Z}_2^2$, *so all diagonals are even* $(0,0)$.

(iii) Off–diagonal spectrum

For $\alpha \neq \beta$, $\deg(\hat{G}^{\alpha\beta}) \in \{(1,0),(0,1),(1,1)\}$.

Proof. (i) is immediate from Lemma 4.66 and commutativity of addition in $\mathbb{Z}_2^2$. (ii) follows since $2\deg(\alpha) = 0$ in $\mathbb{Z}_2^2$. (iii) is the complement of $(0,0)$ in the degree set for $\alpha \neq \beta$. □

Graded $*$-algebra of channels and super–bracket

Definition 4.103 (Graded $*$-algebra of channel components)**.** Let $\mathsf{G} := \operatorname{span}\{\hat{G}^{\alpha\beta}(x,x') : \alpha,\beta \in \mathcal{I}\} \subset \mathcal{B}(\mathcal{H}_{\mathrm{em}})$, with degree $\deg(\hat{G}^{\alpha\beta}) = \deg(\alpha) + \deg(\beta) \in \mathbb{Z}_2^2$ from Definition 4.94. Equip G with:

- *Product:* ordinary operator composition (degree–additive).
- *Adjoint:* $(\cdot)^\dagger$ with $(\hat{G}^{\alpha\beta})^\dagger = \hat{G}^{\beta\alpha}$ (Lemma 4.66).
- *Graded bracket:*

$$[X,Y]_{\mathrm{gr}} := XY - (-1)^{\langle \deg X, \deg Y\rangle} YX, \qquad \langle (a_1,a_2),(b_1,b_2)\rangle := a_1b_1 + a_2b_2 \pmod 2. \tag{4.30}$$

Proposition 4.104 (Color Lie $*$-algebra structure)**.** $(\mathsf{G}, [\cdot,\cdot]_{\mathrm{gr}}, (\cdot)^\dagger)$ *is a* $\mathbb{Z}_2^2$-color *Lie* $*$*-algebra:*

(a) *(*Graded skew-symmetry*)* $[X,Y]_{\mathrm{gr}} = -(-1)^{\langle \deg X, \deg Y\rangle}[Y,X]_{\mathrm{gr}}$.

(b) *(*Graded Jacobi*) For homogeneous* X,Y,Z,

$$\begin{aligned}&(-1)^{\langle \deg X, \deg Z\rangle}[X,[Y,Z]_{\mathrm{gr}}]_{\mathrm{gr}}\\ &+ (-1)^{\langle \deg Y, \deg X\rangle}[Y,[Z,X]_{\mathrm{gr}}]_{\mathrm{gr}}\\ &+ (-1)^{\langle \deg Z, \deg Y\rangle}[Z,[X,Y]_{\mathrm{gr}}]_{\mathrm{gr}} = 0.\end{aligned} \tag{4.31}$$

(c) *(*$*$-compatibility*)* $[X,Y]_{\mathrm{gr}}^\dagger = -(-1)^{\langle \deg X, \deg Y\rangle}[X^\dagger, Y^\dagger]_{\mathrm{gr}}$.

Proof. All identities are standard for the graded commutator on a $\mathbb{Z}_2^2$-graded associative $*$-algebra. (a) is immediate from (4.30). For (b), associativity of the product and bilinearity of $\langle\cdot,\cdot\rangle$ yield the color Jacobi identity by the usual cancellation in the sum of six terms. For (c), use $(XY)^\dagger = Y^\dagger X^\dagger$ and Lemma 4.66:

$$\begin{aligned}[X,Y]_{\mathrm{gr}}^\dagger &= (XY)^\dagger - (-1)^{\langle \deg X, \deg Y\rangle}(YX)^\dagger = Y^\dagger X^\dagger - (-1)^{\langle \deg X, \deg Y\rangle} X^\dagger Y^\dagger \\ &= -\Big(X^\dagger Y^\dagger - (-1)^{\langle \deg X^\dagger, \deg Y^\dagger\rangle} Y^\dagger X^\dagger\Big) = -\,(-1)^{\langle \deg X, \deg Y\rangle}[X^\dagger, Y^\dagger]_{\mathrm{gr}},\end{aligned}$$

since $\deg(X^\dagger) = \deg(X)$ by Lemma 4.98(i). □

Corollary 4.105 (Selection rules: even/odd mixing)**.** *Let $X, Y \in \mathsf{G}$ be homogeneous.*

(i) *If $\deg X = (0,0)$ or $\deg Y = (0,0)$, then $[X,Y]_{\mathrm{gr}} = [X,Y]$ (ordinary commutator).*

(ii) *If $\deg X = \deg Y \in \{(1,0),(0,1),(1,1)\}$, then $[X,Y]_{\mathrm{gr}} = XY + YX$ (graded anticommutator).*

(iii) *If $\deg X$ and $\deg Y$ are distinct nontrivial colors with $\langle \deg X, \deg Y\rangle = 0$ (e.g. $(1,0)$ with $(0,1)$), then $[X,Y]_{\mathrm{gr}} = [X,Y]$ (ordinary commutator).*

(iv) *If $\langle \deg X, \deg Y\rangle = 1$ (e.g. $(1,0)$ with $(1,1)$ or $(0,1)$ with $(1,1)$), then $[X,Y]_{\mathrm{gr}} = XY + YX$.*

Lemma 4.106 (Gauge covariance of the graded algebra)**.** *Under a local gauge $g : \mathcal{X}_q \to \mathrm{U}(\mathcal{H}_{\mathrm{em}})$, $\hat{G}^{\alpha\beta}(x,x') \mapsto g(x)\hat{G}^{\alpha\beta}(x,x')g(x')^\dagger$ (Corollary 4.64), one has for all $X, Y \in \mathsf{G}$:*

$$[gXg^\dagger,\ gYg^\dagger]_{\mathrm{gr}} = g\,[X,Y]_{\mathrm{gr}}\,g^\dagger, \qquad \deg(gXg^\dagger) = \deg X.$$

Proof. Degree invariance is Lemma 4.98(ii). For the bracket, use multiplicativity of conjugation and (4.30):

$$\begin{aligned}[gXg^\dagger, gYg^\dagger]_{\mathrm{gr}} &= gXYg^\dagger - (-1)^{\langle \deg X, \deg Y\rangle} gYXg^\dagger \\ &= g\Big(XY - (-1)^{\langle \deg X, \deg Y\rangle} YX\Big)g^\dagger \\ &= g[X,Y]_{\mathrm{gr}}g^\dagger.\end{aligned}$$

□

Two structural consequences

Proposition 4.107 (Even core acts by degree-0 derivations)**.** *For every locus and indices $\alpha, \beta, \gamma \in \mathcal{I}$, the diagonal channels are even $\deg(\hat{G}^{\alpha\alpha}) = (0,0)$ (Corollary 4.102) and act by ordinary derivations on all colored components:*

$$[\hat{G}^{\alpha\alpha}, \hat{G}^{\beta\gamma}]_{\mathrm{gr}} = [\hat{G}^{\alpha\alpha}, \hat{G}^{\beta\gamma}] \quad \text{and} \quad [\hat{G}^{\alpha\alpha}, XY]_{\mathrm{gr}} = [\hat{G}^{\alpha\alpha}, X]_{\mathrm{gr}}\,Y + X\,[\hat{G}^{\alpha\alpha}, Y]_{\mathrm{gr}}$$

for any homogeneous $X, Y \in \mathsf{G}$ (Definition 4.103). In particular, self-persistence amplitudes $\hat{G}^{\alpha\alpha}$ control the phase/magnitude of transition channels without inducing color flips.

Proof. By Corollary 4.102, $\deg(\hat{G}^{\alpha\alpha}) = (0,0)$. Thus the graded bracket with any Z reduces to the ordinary commutator: $[\hat{G}^{\alpha\alpha}, Z]_{\mathrm{gr}} = [\hat{G}^{\alpha\alpha}, Z]$ (Corollary 4.105(i)). The derivation law follows from $[D, XY] = [D,X]Y + X[D,Y]$ for the ordinary commutator applied with $D = \hat{G}^{\alpha\alpha}$. No color flip occurs because $\deg(D) = (0,0)$. □

Proposition 4.108 (Compatibility with the Fourfold Dialectic Algebra)**.** *Let L, R act on mode indices as in the Fourfold Dialectic Algebra (cf. Definition 1.5), and define the left/right action on the block kernel by*

$$(L \cdot \hat{\mathbf{G}}_\Lambda \cdot R^\dagger)^{\alpha\beta} = \sum_{\gamma,\delta\in\mathcal{I}} L^\alpha{}_\gamma \, \hat{G}_\Lambda^{\gamma\delta}(x,x') \, (R^\dagger)_\delta{}^\beta.$$

If L and R are degree-preserving on indices (homogeneous of total degree 0 with respect to Definition 4.94), then for each term in the sum

$$\deg\Big(L^\alpha{}_\gamma \, \hat{G}^{\gamma\delta} \, (R^\dagger)_\delta{}^\beta\Big) = \deg(\alpha) + \deg(\beta),$$

so the graded decomposition is preserved and the block structure respected.

Proof. Write the index matrix units as $E^\alpha{}_\gamma$ and endow them with $\deg(E^\alpha{}_\gamma) = \deg(\alpha) + \deg(\gamma)$ (the same color-addition rule as Definition 4.94). A degree-0 L (resp. $R^\dagger$) is a linear combination of homogeneous $E^\alpha{}_\gamma$ (resp. $E_\delta{}^\beta$) with $\deg(E^\alpha{}_\gamma) = \deg(\alpha) + \deg(\gamma)$ (resp. $\deg(E_\delta{}^\beta) = \deg(\delta) + \deg(\beta)$). For a homogeneous summand,

$$\deg\Big(E^\alpha{}_\gamma \, \hat{G}^{\gamma\delta} \, E_\delta{}^\beta\Big) = \Big(\deg\alpha + \deg\gamma\Big) + \Big(\deg\gamma + \deg\delta\Big) + \Big(\deg\delta + \deg\beta\Big) = \deg\alpha + \deg\beta,$$

since $2\deg\gamma = 2\deg\delta = 0$ in $\mathbb{Z}_2^2$. Hence each term (and thus the sum) has total degree $\deg(\alpha) + \deg(\beta)$. This matches the selection rule under composition (Proposition 4.99(c), via Proposition 4.85). □

Optional model postulates (graded dynamics)

Definition 4.109 (Word in the channel algebra)**.** Fix loci $x_0, \ldots, x_k$ and consider the generators $\{\hat{G}_\Lambda^{\alpha_i\beta_i}(x_i, x_{i-1})\}_{i=1}^k$ of the graded $*$-algebra G (Definitions 4.94 and 4.103). A *word* (of length k) is a well-typed product

$$W \;=\; X_k X_{k-1} \cdots X_1, \qquad X_i \in \{\hat{G}_\Lambda^{\alpha_i\beta_i}(x_i, x_{i-1})\},$$

where consecutive endpoints match (so composition is defined) and multiplication is ordinary operator composition (Proposition 4.85). The empty word has length 0 and equals the identity. The degree is additive: $\deg W = \sum_{i=1}^k \deg(X_i)$ (Definition 4.94 and Lemma 4.98), and under gauge, $W \mapsto g(x_k)\, W\, g(x_0)^\dagger$ (Corollary 4.64). A *reduced word* is one simplified using the algebraic relations in force (e.g. nilpotencies, centrality, orthogonality), yielding a normal form.

Remark 4.110 (Words as channel histories)**.** Intuitively, a *word* (cf. Definition 4.109) is a single "history'' obtained by stacking channel factors in sequence when their endpoints match. Its length is the number of factors; its degree is the sum of the factor degrees (Definition 4.94 and Lemma 4.98); and under gauge it conjugates at the endpoints (Corollary 4.64).

- **Example (edge history).** Along the p–s edge, $W = \hat{G}^{ps}\hat{G}^{sp}\hat{G}^{ps}$ is a well-typed word of length 3, representing a $s \to p \to s \to p$ conversion sequence across an intermediate locus (Proposition 4.85).

- **Local truncations from nilpotency.** Under Theorem 4.123 we have $(\hat{G}^{ps})^2 = (\hat{G}^{sp})^2 = 0$. Therefore any word containing two consecutive $\hat{G}^{ps}$ (or two consecutive $\hat{G}^{sp}$) factors collapses to 0, yielding the series–truncation effect stated in Corollary 4.125.
- **Stronger differential scenario.** In the stronger variant of (N1) (Definition 4.119) where the graded differential d satisfies $\mathsf{d}^2 = 0$ on both stalks p and s, the *mixed* back–to–back composites also vanish: $\hat{G}^{ps}\hat{G}^{sp} = 0 = \hat{G}^{sp}\hat{G}^{ps}$. Then any nonempty word in $\{\hat{G}^{ps}, \hat{G}^{sp}\}$ reduces to a linear combination of single–step terms (and possibly the identity), exactly as used in the series–truncation arguments.
- **Degree bookkeeping.** Because $\deg(\hat{G}^{ps}) = \deg(\hat{G}^{sp}) = (1,0)$ (Definition 4.94), a word with an odd number of $p \leftrightarrow s$ steps has degree $(1,0)$, while an even number yields total degree $(0,0)$. This parity explains why square terms are the first to vanish in the nilpotent regime.

Postulate 4.1 (Central (or approximately central) holosymmetry)**.** *The holosymmetric self-channel $\hat{G}^{ss}(x,x')$ is central (or central up to controlled corrections): $[\hat{G}^{ss}, \hat{X}]_{\mathrm{gr}} \approx 0$ for all channel combinations $\hat{X}$.*

Definition 4.111 (Central holosymmetry hypotheses)**.** Let $\mathsf{G} = \mathrm{span}\{\hat{G}^{\alpha\beta}_{\Lambda}(x,x')\}_{\alpha,\beta\in\mathcal{I}} \subset \mathcal{B}(\mathcal{H}_{\mathrm{em}})$ be the graded channel $*$-algebra with bracket $[\cdot,\cdot]_{\mathrm{gr}}$ from Definition 4.103. We say that one of the following hypotheses holds:

(C1) Group-averaged holosymmetry (Schur). There exists a compact (or unimodular) implementation group H acting unitarily fibrewise, $U_x : H \to \mathrm{U}(\mathcal{H}_{\mathrm{em}}(x))$, such that: (i) the connection is H-invariant and the transporter is H-equivariant,

$$U_x(h)\,\mathcal{U}(x \leftarrow x';\gamma) = \mathcal{U}(x \leftarrow x';\gamma)\,U_{x'}(h) \quad \forall h \in H, \tag{4.32}$$

(ii) the holosymmetric selector is the H-average of a seed selector $\hat{\mathcal{F}}_0$:

$$\hat{\mathcal{F}}_s(x) \;:=\; \int_H U_x(h)\,\hat{\mathcal{F}}_0(x)\,U_x(h)^\dagger \,\mathrm{d}h, \tag{4.33}$$

and (iii) the representation of H on the relevant subspace of $\mathcal{H}_{\mathrm{em}}(x)$ is irreducible.

(C2) Superselection of the holosymmetric charge. There is a self-adjoint charge Q with spectral projectors $\{P_\chi\}$ such that the holosymmetric mode s corresponds to a sector projector P_s, and the observable/channel algebra acts within the commutant of Q: $\mathsf{G} \subseteq \{Q\}' = \{X : [X,Q] = 0\}$.

(C3) Flat, topological reduction. On a simply connected neighbourhood $U \subset \mathcal{X}_q$ with $\mathcal{F} = 0$ (Proposition 4.59), the holosymmetric selector is a gauge-invariant functional of holonomy classes; since all holonomies trivialize on U, there exist endpoint maps $g : U \to \mathrm{U}(\mathcal{H}_{\mathrm{em}})$ with $\mathcal{U}(x \leftarrow x';\gamma) = g(x)g(x')^\dagger$ and a *scalar* kernel $c(x,x') \in \mathbb{C}$ such that

$$\hat{G}^{ss}_{\Lambda}(x,x') = \hat{\mathcal{F}}_s^\dagger(x)\,\mathcal{U}(x \leftarrow x';\gamma)\,\hat{\mathcal{F}}_s(x') = g(x)\,\Big(c(x,x')\,\mathbf{1}\Big)\,g(x')^\dagger. \tag{4.34}$$

Definition 4.112 (Centrality defect)**.** Under the hypotheses of Definition 4.111, define the (uniform) centrality defect

$$\delta \;:=\; \sup_{\|X\|=1} \Big\| [\,\hat{G}^{ss}_{\Lambda},\, X\,]_{\mathrm{gr}} \Big\| \;\in [0,\infty).$$

We say holosymmetry is δ-*central* if this bound is finite (and small).

Proposition 4.113 (Central holosymmetry under group averaging)**.** *If* (C1) *of Definition 4.111 holds, then* $\hat{G}^{ss}_{\Lambda}(x,x')$ *is proportional to the identity on the* H*-irreducible subspace and hence lies in the center of* G*:*

$$[\hat{G}^{ss}_{\Lambda}(x,x'),\, X]_{\mathrm{gr}} = 0 \quad \textit{for all } X \in \mathsf{G}.$$

Proof. By Definition 4.61, $\hat{G}^{ss}_{\Lambda} = \hat{\mathcal{F}}^{\dagger}_{s}(x)\,\mathcal{U}(x \leftarrow x';\gamma)\,\hat{\mathcal{F}}_{s}(x')$. Insert (4.33) at both endpoints and use the equivariance (4.32) to obtain

$$\hat{G}^{ss}_{\Lambda}(x,x') = \int_{H\times H}\Big(U_x(h)\,\hat{\mathcal{F}}^{\dagger}_{0}(x)\,U_x(h)^{\dagger}\Big)\,\mathcal{U}\,\Big(U_{x'}(k)\,\hat{\mathcal{F}}_{0}(x')\,U_{x'}(k)^{\dagger}\Big)\,\mathrm{d}h\,\mathrm{d}k.$$

A change of variables and averaging on the right shows $\hat{G}^{ss}_{\Lambda}$ intertwines the H-actions at x and x'. On an H-irreducible subspace, Schur's lemma yields $\hat{G}^{ss}_{\Lambda}(x,x') = \lambda(x,x')\,\mathbf{1}$. As a diagonal channel, $\deg(\hat{G}^{ss}) = (0,0)$ (Corollary 4.102), so the graded commutator reduces to the ordinary commutator and hence vanishes with any $X \in \mathsf{G}$. □

Proposition 4.114 (Central holosymmetry under superselection)**.** *If* (C2) *of Definition 4.111 holds, then* $\hat{G}^{ss}_{\Lambda}(x,x') \in \{Q\}'$ *and* $[\,\hat{G}^{ss}_{\Lambda}(x,x'),\, X\,]_{\mathrm{gr}} = 0$ *for all* $X \in \mathsf{G}$.

Proof. By hypothesis, $\mathsf{G} \subseteq \{Q\}'$. Since the holosymmetric channel is confined to the s-sector, $\hat{G}^{ss}_{\Lambda} = P_s\,\hat{G}^{ss}_{\Lambda}\,P_s$ and $[P_s, Q] = 0$. Thus $\hat{G}^{ss}_{\Lambda} \in \{Q\}'$. For each $X \in \mathsf{G}$, $[X,Q] = 0 \Rightarrow [X,P_s] = 0$, whence $[\hat{G}^{ss}_{\Lambda}, X] = 0$. Again $\deg(\hat{G}^{ss}) = (0,0)$ (Corollary 4.102), so $[\,\hat{G}^{ss}_{\Lambda},\, X\,]_{\mathrm{gr}} = [\,\hat{G}^{ss}_{\Lambda},\, X\,] = 0$. □

Proposition 4.115 (Central holosymmetry under flat, topological reduction)**.** *If* (C3) *of Definition 4.111 holds on a simply connected* U*, then for all* $x,x' \in U$*,* $[\,\hat{G}^{ss}_{\Lambda}(x,x'),\, X(x,x')\,]_{\mathrm{gr}} = 0$ *for every channel block* $X \in \mathsf{G}$.

Proof. By Proposition 4.59, choose a gauge with $\mathcal{A} = 0$ on U, hence $\mathcal{U} = \mathbf{1}$ and holonomies are trivial. By the assumption in (4.34), $\hat{G}^{ss}_{\Lambda}(x,x') = c(x,x')\,\mathbf{1}$ in this gauge, hence it commutes with all blocks. Gauge covariance of the bracket (Lemma 4.106) then implies centrality holds in any gauge, i.e. for the covariant form $g(x)\,(c\,\mathbf{1})\,g(x')^{\dagger}$ as written in (4.34). □

Theorem 4.116 (Central (or approximately central) holosymmetry)**.** *Assume any one of* (C1), (C2)*, or* (C3) *in Definition 4.111. Then for all* $X \in \mathsf{G}$*,*

$$[\,\hat{G}^{ss}_{\Lambda}(x,x'),\, X\,]_{\mathrm{gr}} = 0, \qquad [\,\hat{G}^{ss}_{\Lambda}(x,x'),\, X\,] = 0,$$

where the second equality follows from $\deg(\hat{G}^{ss}_{\Lambda}) = (0,0)$ *and Lemma 4.95.*

If the corresponding hypothesis holds only approximately (e.g. the H*-representation in* (C1) *is nearly irreducible, or* (4.34) *holds up to a small scalar defect), then letting* δ *be as in Definition 4.112, we have the uniform bound*

$$\big\|[\,\hat{G}^{ss}_{\Lambda}(x,x'),\, X\,]_{\mathrm{gr}}\big\| = \big\|[\,\hat{G}^{ss}_{\Lambda}(x,x'),\, X\,]\big\| \;\le\; \delta\,\|X\| \quad \textit{for all } X \in \mathsf{G}.$$

In the exact cases (C1)–(C3) *one has* $\delta = 0$.

Proof. Preliminaries. Fix endpoints (x, x'). By construction $\mathsf{G} = \mathrm{span}\{\hat{G}^{\alpha\beta}_{\Lambda}(x, x')\}_{\alpha,\beta\in\mathcal{I}} \subset \mathcal{B}(\mathcal{H}_{\mathrm{em}})$ is a graded $*$-algebra (Definition 4.103); in particular $\deg(\hat{G}^{ss}) = \deg(s) + \deg(s) = (0,1) + (0,1) = (0,0)$ (even) by Definition 4.94. Hence the graded bracket with $\hat{G}^{ss}$ coincides with the ordinary commutator: $[\,\hat{G}^{ss}, X\,]_{\mathrm{gr}} = [\,\hat{G}^{ss}, X\,]$ for all X (Corollary 4.102). Moreover, by gauge covariance of the graded structure, the vanishing of the bracket is gauge-invariant (Lemma 4.106 and Corollary 4.64); we may therefore work in any convenient gauge.

Case (C1): Group averaging / Schur. Let H act unitarily on the fibres at each locus and assume the holosymmetric selector is defined by averaging, $\hat{\mathcal{F}}_s(y) = \int_H U_y(h)\,\hat{\mathcal{F}}_0(y)\,U_y(h)^\dagger\,dh$ (Haar measure), so $\hat{\mathcal{F}}_s(y)$ is H-equivariant. Denote by $\mathrm{End}_H(\mathcal{H}_{\mathrm{em}}(y))$ the H-intertwiners at y. Then $\hat{\mathcal{F}}_s(y), \hat{\mathcal{F}}_s(y)^\dagger \in \mathrm{End}_H(\mathcal{H}_{\mathrm{em}}(y))$ by construction, and the channel

$$\hat{G}^{ss}_{\Lambda}(x, x') = \hat{\mathcal{F}}_s^\dagger(x)\,\mathcal{U}(x \leftarrow x'; \gamma)\,\hat{\mathcal{F}}_s(x') \qquad \text{(Definition 4.61)}$$

is an H-intertwiner from $\mathcal{H}_{\mathrm{em}}(x')$ to $\mathcal{H}_{\mathrm{em}}(x)$. Under the same symmetry, every block $\hat{G}^{\alpha\beta}_{\Lambda}(x, x')$ is also an H-intertwiner (selectors and transport are H-equivariant), so $\mathsf{G} \subset \mathrm{Hom}_H\Big(\mathcal{H}_{\mathrm{em}}(x'), \mathcal{H}_{\mathrm{em}}(x)\Big)$. Decompose both endpoint fibres into H-isotypic components. By the double-commutant/Schur decomposition, Hom_H is block-diagonal with scalar matrices on each multiplicity space; in particular, any two intertwiners commute on each isotypic block. Therefore $\hat{G}^{ss}_{\Lambda}$ commutes with every $X \in \mathsf{G}$, i.e. $[\,\hat{G}^{ss}_{\Lambda}, X\,] = 0$. In the irreducible case on the relevant subspace, Schur's lemma sharpens this to $\hat{G}^{ss}_{\Lambda} = c\,\Pi$ with $c \in \mathbb{C}$ and Π the intertwiner identity, which is manifestly central.

Case (C2): Superselection. Assume that label s determines a superselection charge and that the observable/channel algebra at fixed endpoints is the commutant of that charge, i.e. $\mathsf{G} \subset \mathcal{A}'$ for the charge representation, while $\hat{G}^{ss}_{\Lambda} \in Z(\mathcal{A}')$ by construction. Then $[\,\hat{G}^{ss}_{\Lambda}, X\,] = 0$ for all $X \in \mathsf{G}$ since central elements of the commutant commute with the whole commutant. (See Proposition 4.114 for a detailed realization.)

Case (C3): Flat, topological reduction. On a simply connected neighbourhood with $\mathcal{F} = 0$ (Proposition 4.59), choose a flat gauge so that $\mathcal{U}(x \leftarrow x'; \gamma) = g(x)\,g(x')^\dagger$ and holonomies are trivial. If s is defined as a gauge-invariant functional of holonomy classes, then in this flat gauge it reduces to an endpoint functional, yielding (for some scalar functions c_s) the reduction

$$\hat{G}^{ss}_{\Lambda}(x, x') = \; c_s(x)\,c_s(x')^*\,\mathbf{1}, \tag{4.35}$$

which clearly commutes with every $X \in \mathsf{G}$ (see Proposition 4.115 for the explicit reduction). By gauge covariance of the bracket (Lemma 4.106 and Corollary 4.64), commutation holds in all gauges.

Combining the three cases shows $[\,\hat{G}^{ss}_{\Lambda}, X\,] = 0$ for all $X \in \mathsf{G}$, hence $[\,\hat{G}^{ss}_{\Lambda}, X\,]_{\mathrm{gr}} = 0$ since $\hat{G}^{ss}$ is even.

Approximate statement. Suppose the hypotheses hold with defect ε. Typical instances: (i) in (C1), the H-averaging projects to an "almost irreducible" block so that on the regulated fibre $\|\hat{G}^{ss}_{\Lambda} - c\,\Pi\| \le \varepsilon$ for some scalar c; (ii) in (C3), flatness/holonomy reduction holds up to $\|\hat{G}^{ss}_{\Lambda} - c_s(x)c_s(x')^*\,\mathbf{1}\| \le \varepsilon$. For any $X \in \mathsf{G}$,

$$\|[\,\hat{G}^{ss}_{\Lambda}, X\,]\| = \|[\,(\hat{G}^{ss}_{\Lambda} - c\mathbf{1}) + c\mathbf{1},\, X\,]\| = \|[\,\hat{G}^{ss}_{\Lambda} - c\mathbf{1},\, X\,]\| \le 2\,\varepsilon\,\|X\|,$$

using bilinearity, $[c\mathbf{1}, X] = 0$, and the standard bound $\|AB - BA\| \le \|A\|\,\|B\| + \|B\|\,\|A\|$. Since $\hat{G}^{ss}$ is even, the same estimate holds for the graded bracket. The bound is uniform in X on norm-bounded subsets of G, completing the approximate centrality claim.

Conclusion. Under any of (C1)–(C3) we have $[\,\hat{G}^{ss}_{\Lambda},\, X\,]_{\mathrm{gr}} = 0$ for all $X \in \mathsf{G}$; under approximate hypotheses the commutator norm is uniformly $O(\varepsilon)$. This proves the theorem. □

Remark 4.117 (Operational meaning of central holosymmetry)**.** Theorem 4.116 asserts that the holosymmetric channel $\hat{G}^{ss}_{\Lambda}(x, x')$ is an *even, graded–central* element of the channel $*$-algebra G (Definitions 4.94 and 4.103): for every $X \in \mathsf{G}$, $[\,\hat{G}^{ss}_{\Lambda}, X\,]_{\mathrm{gr}} = 0$. Concretely:

(a) **No color mixing, only uniform weighting.** Since $\deg(\hat{G}^{ss}) = (0, 0)$, the graded bracket reduces to the ordinary commutator (Corollary 4.102). Thus $\hat{G}^{ss}_{\Lambda}$ preserves the $\mathbb{Z}_2^2$-grading of every channel (Proposition 4.99): it cannot create or annihilate dialectic "color''; it only rescales/phases existing amplitudes.

(b) **Order–insensitivity in compositions.** In any word of channels built by block composition through intermediate loci (Proposition 4.85), $\hat{G}^{ss}_{\Lambda}$ may be moved freely to the left or right without changing the value. Perturbatively, insertions of $\hat{G}^{ss}_{\Lambda}$ can be regrouped in Dyson/Neumann expansions with no effect on the sum, and with $O(\varepsilon)$ control in the approximate case of Theorem 4.116.

(c) **Gauge stability and endpoint interpretation.** Centrality is invariant under local gauge by the covariance of the graded structure (Lemma 4.106 and Corollary 4.64). In the flat, simply connected regime (Proposition 4.59), $\hat{G}^{ss}_{\Lambda}$ reduces to an endpoint scalar (or scalar multiple of the identity) (Proposition 4.115), so holosymmetric "phases'' can be pushed to the endpoints without altering any inter-modal coupling.

(d) **Compatibility with Twofold reductions.** Within the Twofold block (eq. (4.25)) and in the duality–collapse regime (Definition 4.87), centrality implies that $\hat{G}^{ss}_{\Lambda}$ acts as a scalar on the effective qubit sector; hence it does not obstruct the qubit recovery (Proposition 4.90).

(e) **Relation to symmetry mechanisms.** The three sufficient mechanisms in Definition 4.111 explain why centrality is natural: group averaging/Schur (C1) makes $\hat{G}^{ss}_{\Lambda}$ an intertwiner–scalar, superselection (C2) places it in the center of the commutant, and flat topological reduction (C3) collapses it to endpoint data. Each mechanism yields the same algebraic consequence in G.

In short, central (holosymmetric) structure *controls* amplitudes but does not *couple* dialectic sectors: it is globally present, gauge-stable, and algebraically inert with respect to graded commutation, so that physical predictions extracted from the Fourfold kernel (Section 4.1.2 and eq. (4.23)) are insensitive to the ordering of holosymmetric factors.

Remark 4.118 (Cross-chapter anchoring)**.** The ingredients used above are intrinsic to the developed formalism: (i) channel assembly and gauge covariance (Definition 4.61, Proposition 4.54, and Corollary 4.64), (ii) flatness ⇒ pure gauge transports (Proposition 4.59), and (iii) graded algebraic control of commutators (Definition 4.103, Proposition 4.104, and Lemma 4.106). When H encodes implementation symmetries of the holosymmetric interferometer (Chapter 3), (C1) matches the group-averaging constructions leading to implementation-invariant

effects; superselection scenarios (C2) arise naturally from the Chapter 1/2 mode-typing when charge sectors are fixed by preparation/measurement.

Postulate 4.2 (Graded nilpotency along the p–s edge)**.** *Certain colored transitions are graded-nilpotent, e.g.* $(\hat{G}^{ps})^2 = 0$ *and* $(\hat{G}^{sp})^2 = 0$*, capturing one-step synthesis/holothesis conversions that cannot pile up coherently in series.*

Definition 4.119 (Nilpotency hypotheses on the p–s edge)**.** Let $\mathsf{G} = \mathrm{span}\{\hat{G}^{\alpha\beta}_{\Lambda}(x,x')\}_{\alpha,\beta\in\mathcal{I}} \subset \mathcal{B}(\mathcal{H}_{\mathrm{em}})$ be the graded channel $*$-algebra of Definition 4.103, with degrees $\deg(\hat{G}^{\alpha\beta}) = \deg(\alpha) + \deg(\beta)$ as in Definition 4.94. We say that one of the following hypotheses holds.

(N1) Differential structure. There exists a graded differential d of degree $(1,1)$ on the mode complex (with $\mathsf{d}^2 = 0$) such that the functorial channel assignment $\Gamma : \mathrm{End}(\mathrm{Modes}) \to \mathsf{G}$ satisfies $\Gamma(\mathsf{d}) = \hat{G}^{ps}_{\Lambda}$ and is multiplicative with respect to composition, in the sense implemented by block composition through an intermediate locus $\big($cf. Proposition 4.85$\big)$.

(N2) Orthogonality at an intermediate locus. There exists a locus y (with dialectic completeness Definition 4.84) such that the p– and s–selectors have orthogonal supports,

$$\hat{\mathcal{F}}_s^{\dagger}(y)\,\hat{\mathcal{F}}_p(y) = 0 \quad = \quad \hat{\mathcal{F}}_p^{\dagger}(y)\,\hat{\mathcal{F}}_s(y), \tag{4.36}$$

and transports are unitary (Definition 4.48 and Lemma 4.50), so that orthogonality is preserved under insertion of resolutions at y.

(N3) Spectral selection rule (cutoff). With the spectral regulation $\mathrm{Ran}\, E^{(x)}([0,\lambda_{\mathrm{cutoff}}])$ from eq. (4.303), composing two $p\to s$ (or two $s\to p$) channels exits the regulated band, i.e.

$$E^{(x)}\hat{G}^{ps}_{\Lambda}(x,x')\big(\mathbf{1} - E^{(x')}\big) = \hat{G}^{ps}_{\Lambda}(x,x') \quad \text{and} \quad \big(\mathbf{1} - E^{(x)}\big)\hat{G}^{ps}_{\Lambda}(x,x')E^{(x')} = \hat{G}^{ps}_{\Lambda}(x,x'), \tag{4.37}$$

so that on the regulated fibres the square of $\hat{G}^{ps}_{\Lambda}$ (and similarly $\hat{G}^{sp}_{\Lambda}$) vanishes.

Proposition 4.120 (Nilpotent p–s edge under differential structure)**.** *If* (N1) *of Definition 4.119 holds, then* $(\hat{G}^{ps}_{\Lambda})^2 = 0$ *and* $(\hat{G}^{sp}_{\Lambda})^2 = 0$*.*

Proof. By hypothesis, $\hat{G}^{ps}_{\Lambda} = \Gamma(\mathsf{d})$ and Γ respects composition via the block composition identity of Proposition 4.85. Thus $(\hat{G}^{ps}_{\Lambda})^2 = \Gamma(\mathsf{d})\,\Gamma(\mathsf{d}) = \Gamma(\mathsf{d}^2) = \Gamma(0) = 0$. The same argument applies to $\hat{G}^{sp}_{\Lambda}$ when $\Gamma(\mathsf{d}^{\dagger}) = \hat{G}^{sp}_{\Lambda}$, using $\mathsf{d}^2 = 0$ and Lemma 4.66 for adjoints. □

Proposition 4.121 (Nilpotent p–s edge under orthogonality)**.** *If* (N2) *of Definition 4.119 holds, then* $(\hat{G}^{ps}_{\Lambda}(x,x'))^2 = 0$ *and* $(\hat{G}^{sp}_{\Lambda}(x,x'))^2 = 0$ *for all* x, x'*.*

Proof. Fix x, x' and insert the dialectically complete resolution at y (Definition 4.84) into the product $(\hat{G}^{ps})^2$ using the block composition rule (Proposition 4.85):

$$\hat{G}^{ps}(x,x')\,\hat{G}^{ps}(x,x') = \sum_{\gamma\in\mathcal{I}} \hat{G}^{p\gamma}(x,y)\,\hat{G}^{\gamma s}(y,x')\,\hat{G}^{ps}(x,x').$$

Expanding one more time through y and using the channel definition $\hat{G}^{\alpha\beta} = \hat{\mathcal{F}}_{\alpha}^{\dagger}\,\mathcal{U}\,\hat{\mathcal{F}}_{\beta}$ (Definition 4.61), every contribution contains at the locus y a factor of the form $\hat{\mathcal{F}}_s^{\dagger}(y)\hat{\mathcal{F}}_p(y)$ (or its adjoint) between two unitary transports. By (4.36) this factor is zero, and unitarity of transport (Lemma 4.50) preserves the orthogonality. Hence each summand vanishes and $(\hat{G}^{ps})^2 = 0$. The same argument, with p and s exchanged, yields $(\hat{G}^{sp})^2 = 0$. □

Proposition 4.122 (Nilpotent p–s edge under spectral selection)**.** *If* (N3) *of Definition 4.119 holds, then on the regulated fibres* $\operatorname{Ran} E^{(x)}([0,\lambda_{\text{cutoff}}])$ *one has* $(\hat{G}^{ps}_{\Lambda})^2 = 0$ *and* $(\hat{G}^{sp}_{\Lambda})^2 = 0$.

Proof. Let $\Pi_x := E^{(x)}([0,\lambda_{\text{cutoff}}])$. By (4.37), $\Pi_x \hat{G}^{ps}(x,x')\Pi_{x'} = 0$ while $\hat{G}^{ps}(x,x') = \Pi_x^{\perp}\hat{G}^{ps}(x,x')\Pi_{x'}^{\perp}$. Therefore

$$\Pi_x\,\hat{G}^{ps}(x,x')\,\Pi_{x'}\,\hat{G}^{ps}(x,x')\,\Pi_{x'} = \Big(\Pi_x\hat{G}^{ps}(x,x')\Pi_{x'}\Big)\Big(\hat{G}^{ps}(x,x')\Pi_{x'}\Big) = 0,$$

and similarly for all other placements of $\Pi/\Pi^{\perp}$ that restrict to the regulated fibres. Hence $(\hat{G}^{ps})^2 = 0$ on $\Pi\mathcal{H}_{\text{em}}$. The same argument applies to $\hat{G}^{sp}$. □

Theorem 4.123 (Nilpotent p–s edge under Y)**.** *Assume any one of* (N1), (N2), *or* (N3) *in Definition 4.119. Then*

$$(\hat{G}^{ps}_{\Lambda})^2 = (\hat{G}^{sp}_{\Lambda})^2 = 0.$$

Moreover, the degree constraint $\deg(\hat{G}^{ps}) = \deg(\hat{G}^{sp}) = (1,0)$ *(Definition 4.94) is consistent with nilpotency and with gauge covariance of products (Lemma 4.98).*

Proof. Preliminaries. For fixed endpoints (x,x'), recall $\hat{G}^{\alpha\beta}_{\Lambda}(x,x') = \hat{\mathcal{F}}^{\dagger}_{\alpha}(x)\,\mathcal{U}(x \leftarrow x';\gamma)\,\hat{\mathcal{F}}_{\beta}(x')$ (Definition 4.61). By the grading convention, $\deg(\hat{G}^{\alpha\beta}) = \deg(\alpha) + \deg(\beta)$ (Definition 4.94), hence

$$\begin{aligned}\deg(\hat{G}^{ps}) &= \deg(p) + \deg(s) = (1,1) + (0,1) = (1,0),\\ \deg(\hat{G}^{sp}) &= \deg(s) + \deg(p) = (0,1) + (1,1) = (1,0).\end{aligned} \tag{4.38}$$

and $\deg\Big((\hat{G}^{ps})^2\Big) = \deg\Big((\hat{G}^{sp})^2\Big) = (0,0)$. Degree additivity and gauge covariance under endpoint conjugation will be used repeatedly (Lemma 4.98, Corollary 4.64, and Proposition 4.54).

Square expansion and the "kill term". We compute

$$(\hat{G}^{ps}_{\Lambda})^2 = \hat{\mathcal{F}}^{\dagger}_{p}\,\mathcal{U}\,\hat{\mathcal{F}}_{s}\,\hat{\mathcal{F}}^{\dagger}_{p}\,\mathcal{U}\,\hat{\mathcal{F}}_{s} \tag{4.39}$$

$$= \hat{\mathcal{F}}^{\dagger}_{p}\,\mathcal{U}\,\underbrace{\big(\hat{\mathcal{F}}\,\hat{\mathcal{F}}^{\dagger}_{p}\big)}_{s}{}_{\text{kill term}}\,\mathcal{U}\,\hat{\mathcal{F}}_{s}. \tag{4.40}$$

Case (N1): Differential/typing structure on the p–s edge. By the domain/codomain *typing* of holosymmetric words (Remark 4.140 together with Proposition 4.85), the composition $\hat{\mathcal{F}}_s\hat{\mathcal{F}}^{\dagger}_p$ in (4.40) is ill-typed and hence vanishes under (N1). Therefore

$$\hat{\mathcal{F}}_s\hat{\mathcal{F}}^{\dagger}_p = 0 \quad\Longrightarrow\quad (\hat{G}^{ps}_{\Lambda})^2 = 0. \tag{4.41}$$

The $(\hat{G}^{sp}_{\Lambda})^2 = 0$ case is identical by swapping $p \leftrightarrow s$.

(Optional alternative under (N1)). Equivalently, if d is the graded differential on the two–node subcomplex $\{p,s\}$ with $\mathsf{d}^2 = 0$ and representation $\mathsf{d}|_{s\to p} \mapsto \hat{G}^{ps}_{\Lambda}$, $\mathsf{d}|_{p\to s} \mapsto \hat{G}^{sp}_{\Lambda}$, then functoriality yields $(\hat{G}^{ps}_{\Lambda})^2 = \rho((\mathsf{d}|_{s\to p})^2) = 0$ and likewise for sp.

Case (N2): Rank/orthogonality at an intermediate locus. Assume $\hat{\mathcal{F}}^{\dagger}_s(y)\hat{\mathcal{F}}_p(y) = 0$ at some intermediate locus y and that the selector family at y is dialectically complete (Definition 4.84). Using block composition through y (Proposition 4.85),

$$\hat{G}^{ps}_{\Lambda}(x,x') = \sum_{\gamma\in\mathcal{I}} \hat{G}^{p\gamma}_{\Lambda}(x,y)\,\hat{G}^{\gamma s}_{\Lambda}(y,x').$$

Hence

$$(\hat{G}^{ps}_{\Lambda})^2 = \sum_{\gamma,\delta} \hat{G}^{p\gamma}_{\Lambda}(x,y)\, \underbrace{\hat{G}^{\gamma s}_{\Lambda}(y,x')\, \hat{G}^{p\delta}_{\Lambda}(x,y)}_{\mathsf{M}_{\gamma\delta}}\, \hat{G}^{\delta s}_{\Lambda}(y,x').$$

Expanding $\mathsf{M}_{\gamma\delta}$ into selectors and transports, one finds that every term contains, at the locus y, a factor of the form $\hat{\mathcal{F}}^{\dagger}_{s}(y)\hat{\mathcal{F}}_{p}(y)$ or its adjoint, produced by the contraction of the γ- and δ-legs across y. By the orthogonality hypothesis this vanishes, whence $\mathsf{M}_{\gamma\delta} = 0$ for all γ, δ, and therefore $(\hat{G}^{ps}_{\Lambda})^2 = 0$. The same argument with $p \leftrightarrow s$ yields $(\hat{G}^{sp}_{\Lambda})^2 = 0$. Degree additivity and gauge covariance are preserved termwise (Lemma 4.98), so the conclusion is gauge-invariant.

Case (N3): Spectral cutoff / support exit. Let $E^{(y)}([0, \lambda_{\text{cutoff}}])$ be the spectral projector defining the regulated fibre at y (eq. (4.303)). Suppose the realizations are arranged so that a $p \to s$ map lands outside the admissible band for a subsequent $p \to s$ application (support–exit selection rule in Definition 4.119(N3)). Insert the resolution of the identity at y, $\mathbf{1} = E^{(y)} + [I - E^{(y)}]$, between the two factors of $\hat{G}^{ps}_{\Lambda}$. By hypothesis the image of the first $p \to s$ step lies in $\mathrm{Ran}(I - E^{(y)})$, while the second $p \to s$ factor has domain restricted to $\mathrm{Ran}(E^{(y)})$; hence their composition vanishes on the regulated fibre, and $(\hat{G}^{ps}_{\Lambda})^2 = 0$. The same holds for $s \to p$. Since endpoint conjugation preserves spectral bands and zero operators (Proposition 4.54 and Corollary 4.64), the statement is gauge-stable. Degree additivity remains as above.

Degree/gauge consistency. In all cases, $\deg(\hat{G}^{ps}) = \deg(\hat{G}^{sp}) = (1,0)$ by construction (Definition 4.94), and products transform covariantly under gauge with unchanged degree (Lemma 4.98). The vanishing of a product is preserved under unitary conjugation, so nilpotency holds in any gauge.

Conclusion. Under any one of (N1)–(N3), both $(\hat{G}^{ps}_{\Lambda})^2$ and $(\hat{G}^{sp}_{\Lambda})^2$ vanish, with grading and gauge properties consistent with Definition 4.94 and Lemma 4.98. □

Remark 4.124 (Operational meaning of a nilpotent p–s edge)**.** Theorem 4.123 states that the $p \leftrightarrow s$ converters behave as *odd, square–zero* morphisms in the channel algebra: $(\hat{G}^{ps}_{\Lambda})^2 = (\hat{G}^{sp}_{\Lambda})^2 = 0$ with $\deg(\hat{G}^{ps}) = \deg(\hat{G}^{sp}) = (1,0)$ (Definition 4.94). Concretely:

(a) **One–step conversion only.** Any word containing two successive $p \to s$ (or $s \to p$) factors collapses to zero. Hence series over the p–s edge *truncate at first order* (Corollary 4.125); higher multiple–conversion histories do not contribute.

(b) **Graded selection and interference.** The $(1,0)$ color of the edge maps fits the $\mathbb{Z}_2^2$ calculus of Definition 4.103 and eq. (4.30) and the blockwise selection rules in Proposition 4.99. In particular, graded commutators with $(0,1)$-colored channels pick up the expected sign, but the *squares* vanish identically, so there is no coherent stacking of $p \leftrightarrow s$ transitions.

(c) **Composed transport is simplified.** In block compositions through an intermediate locus (Proposition 4.85), any term that attempts a two–step $p \to s \to s$ or $s \to p \to p$ conversion vanishes termwise. Perturbative (Dyson/Neumann) expansions therefore have strictly fewer nonzero diagrams along the p–s edge.

(d) **Gauge stability.** Nilpotency and color are preserved by endpoint conjugation (Lemma 4.98, Corollary 4.64, and Proposition 4.54); thus the statement is *gauge–invariant* and holds in any local frame.

(e) **Compatibility with central holosymmetry.** When $\hat{G}^{ss}_\Lambda$ is central (Theorem 4.116), the $\{p, s\}$ subblock is (graded) upper–triangular with a nilpotent off–diagonal: ordering with $\hat{G}^{ss}_\Lambda$ is immaterial, and no iterative $p \leftrightarrow s$ cycling can build up.

(f) **Twofold reductions unaffected.** Because the edge is nilpotent, projecting to the Twofold sector ($\{0, 1\}$) does not acquire hidden higher–order contributions from p, s-loops; this supports the qubit recovery in the duality–collapse regime (Proposition 4.90).

In summary, a nilpotent p–s edge means *single–pass synthesis/holothesis conversion* with no iterative coherence. It is consistent with the graded algebraic structure, stable under gauge, and it reduces the combinatorics of channel compositions without constraining the Twofold slice.

Corollary 4.125 (Series truncation along the p–s edge)**.** *Assume the p–s nilpotency of Theorem 4.123, i.e. $(\hat{G}^{ps}_\Lambda)^2 = (\hat{G}^{sp}_\Lambda)^2 = 0$.*

(a) Pruning of nonalternating words

Every monomial in $\{\hat{G}^{ps}_\Lambda, \hat{G}^{sp}_\Lambda\}$ that contains $(\hat{G}^{ps}_\Lambda)^2$ or $(\hat{G}^{sp}_\Lambda)^2$ as a contiguous subword vanishes. Hence, in Dyson/Neumann or block–composition expansions (Proposition 4.85) all terms with two successive $p \to s$ or two successive $s \to p$ steps vanish termwise. Only strictly alternating *subwords (e.g. $\hat{G}^{ps}\hat{G}^{sp}\hat{G}^{ps}\cdots$) can survive.*

(b) Strong truncation under orthogonality and local flatness (optional)

If, in addition, the dialectic family is complete at each inserted intermediate locus (Definition 4.84), the p– and s–selectors are orthogonal there ($\hat{\mathcal{F}}_s^\dagger \hat{\mathcal{F}}_p = 0 = \hat{\mathcal{F}}_p^\dagger \hat{\mathcal{F}}_s$, Definition 4.119 (N2)), and transport is locally flat (so path-ordered transport reduces to endpoint conjugation, Propositions 4.54 and 4.59), then any surviving (alternating) word reduces to a linear combination of $\mathbf{1}$, $\hat{G}^{ps}_\Lambda$, and $\hat{G}^{sp}_\Lambda$.

Proof. (a) Let W be any word in the alphabet $\{E, F\} := \{\hat{G}^{ps}_\Lambda, \hat{G}^{sp}_\Lambda\}$. If W contains EE or FF contiguously, then by Theorem 4.123 the corresponding factor equals 0, so the entire monomial vanishes. In Dyson/Neumann or block–composition series (Proposition 4.85) the terms are ordered products of channels; the same cancellation applies termwise. Thus only strictly alternating subwords can contribute.

(b) Under Definition 4.84, insert the resolution at each intermediate locus y to factor alternating products via $\hat{G}^{\alpha\beta}(x, x') = \sum_{\gamma\in\mathcal{I}} \hat{G}^{\alpha\gamma}(x, y)\hat{G}^{\gamma\beta}(y, x')$ (Proposition 4.85). For an alternating pair, e.g. $\hat{G}^{ps}(x, y)\hat{G}^{sp}(y, x')$, the definition $\hat{G}^{\alpha\beta} = \hat{\mathcal{F}}^\dagger_\alpha \mathcal{U}\, \hat{\mathcal{F}}_\beta$ (Definition 4.61) gives

$$\hat{G}^{ps}(x, y)\hat{G}^{sp}(y, x') = \hat{\mathcal{F}}^\dagger_p(x)\,\mathcal{U}(x \leftarrow y)\, \underbrace{\hat{\mathcal{F}}_s(y)\hat{\mathcal{F}}^\dagger_s(y)}_{=:P_s(y)}\, \mathcal{U}(y \leftarrow x')\, \hat{\mathcal{F}}_p(x').$$

Orthogonality $(N2)$ implies $P_s(y)$ is a projection orthogonal to the p-support ($\hat{\mathcal{F}}_s^\dagger \hat{\mathcal{F}}_p = 0$), and unitarity of transport preserves these supports (Lemma 4.50). Under local flatness (Proposition 4.59) we may gauge-fix so that transports collapse to endpoint conjugations, whence $P_s(y)$ can be moved to an endpoint and absorbed into a diagonal factor that either acts as the identity on its support or is zero. Iterating this reduction shows that any alternating word collapses to a linear combination of $\mathbf{1}$, $\hat{G}^{ps}$, and $\hat{G}^{sp}$. Gauge covariance of the reduction follows from Corollary 4.64 and Lemma 4.106. □

Remark 4.126 (Cross-chapter anchoring and interpretation)**.** *Differential view (N1).* In Chapter 1 the tetralemma modes carry algebraic operations; modeling $p \leftrightarrow s$ as a graded differential d matches the $\mathbb{Z}_2^2$-grading of Definition 4.94 and promotes $\hat{G}^{ps}$ to a "boundary" map whose square vanishes. *Orthogonality view (N2).* If the parasymmetric and holosymmetric selectors are realized as partial isometries onto orthogonal subspaces at an intermediate locus, then the block-composition identity (Proposition 4.85) forces two-step $p \to s \to s$ (and $s \to p \to p$) processes to vanish. *Spectral view (N3).* In Chapter 2/3 implementations with spectral windows (eq. (4.303)), certain cross-mode transitions leave the energy band; restricting to the regulated fibre yields algebraic nilpotency of the corresponding channel squares.

Postulate 4.3 (Bosonic compatibility in the 0–s sector)**.** *Even channels generated by symmetry and holosymmetry diagonals commute:* $[\hat{G}^{00}, \hat{G}^{ss}] = 0$*, and more generally* $[\hat{G}^{00}, \hat{X}]_{\mathrm{gr}} = [\hat{G}^{00}, \hat{X}]$ *(no color flips).*

Definition 4.127 (Bosonic-commutation hypotheses in the 0–s sector)**.** Let

$$\mathsf{G} := \operatorname{span}\left\{ \hat{G}^{\alpha\beta}_{\Lambda}(x,x') \;\middle|\; \alpha,\beta \in \mathcal{I} \right\} \subset \mathcal{B}\big(\mathcal{H}_{\mathrm{em}}\big)$$

be the graded channel $*$-algebra of Definition 4.103, with degrees as in Definition 4.94. We consider the following sufficient hypotheses for the commutation

$$\Big[\hat{G}^{00}_{\Lambda}(x,x'),\, \hat{G}^{ss}_{\Lambda}(x,x')\Big] = 0$$

(hence also $\Big[\hat{G}^{00}_{\Lambda},\, \hat{G}^{ss}_{\Lambda}\Big]_{\mathrm{gr}} = 0$, since diagonals are even by Corollary 4.102).

(B1) Shared spectral functional. There exists a self-adjoint observable $Z(x,x')$ (gauge-invariant under Corollary 4.64) such that both diagonal channels are Borel functions of Z on the regulated fibre $\operatorname{Ran} E^{(x)}([0,\lambda_{\mathrm{cutoff}}])$ (cf. eq. (4.303)):

$$\hat{G}^{00}_{\Lambda}(x,x') = f\Big(Z(x,x')\Big), \qquad \hat{G}^{ss}_{\Lambda}(x,x') = g\Big(Z(x,x')\Big).$$

(B2) Central holosymmetry. The holosymmetric diagonal is central in G, e.g. under any of the hypotheses (C1)–(C3) of Definition 4.111 (see Theorem 4.116).

(B3) Joint unitary diagonalizability. On the regulated fibre, $\hat{G}^{00}_{\Lambda}(x,x')$ and $\hat{G}^{ss}_{\Lambda}(x,x')$ are normal and there exists a unitary W (possibly obtained in a flat gauge, cf. Propositions 4.54 and 4.59) such that $W^{\dagger}\hat{G}^{00}_{\Lambda}W = \operatorname{diag}(\lambda_i)$ and $W^{\dagger}\hat{G}^{ss}_{\Lambda}W = \operatorname{diag}(\mu_i)$.

Proposition 4.128 (Commutation from shared functional calculus)**.** *If* (B1) *of Definition 4.127 holds, then* $[\hat{G}^{00}_{\Lambda}(x,x'), \hat{G}^{ss}_{\Lambda}(x,x')] = 0$.

Proof. By the spectral theorem, Borel functions of the same self-adjoint operator commute: $[f(Z), g(Z)] = 0$. Since both diagonals are even-degree blocks (Corollary 4.102), the graded bracket coincides with the ordinary one. Gauge covariance of Z follows from Corollary 4.64, hence the statement is representation-independent. □

Proposition 4.129 (Commutation from central holosymmetry)**.** *If* (B2) *of Definition 4.127 holds, then* $[\hat{G}^{00}_{\Lambda}, \hat{G}^{ss}_{\Lambda}] = 0$.

Proof. Centrality (e.g. from Theorem 4.116) gives $[\hat{G}^{ss}_\Lambda, X]_{\mathrm{gr}} = 0$ for all $X \in \mathsf{G}$; in particular with $X = \hat{G}^{00}_\Lambda$. Since both are even, the graded and ordinary commutators coincide. □

Proposition 4.130 (Commutation from joint diagonalizability)**.** *If* (B3) *of Definition 4.127 holds, then* $[\hat{G}^{00}_\Lambda(x,x'), \hat{G}^{ss}_\Lambda(x,x')] = 0$.

Proof. Normal operators that are simultaneously unitarily diagonalizable commute. Hence

$$W^\dagger\,[\hat{G}^{00}_\Lambda, \hat{G}^{ss}_\Lambda]\,W = [\mathrm{diag}(\lambda_i), \mathrm{diag}(\mu_i)] = 0.$$

If the joint diagonalization is obtained in a flat gauge (Proposition 4.59), commutation holds in that gauge and thus in any gauge by covariance of the bracket (Lemma 4.106). □

Theorem 4.131 (Bosonic compatibility in the 0–*s* sector under Z)**.** *Assume any one of* (B1), (B2), *or* (B3) *in Definition 4.127. Then for all* x, x',

$$[\hat{G}^{00}_\Lambda(x,x'), \hat{G}^{ss}_\Lambda(x,x')] = 0, \qquad [\hat{G}^{00}_\Lambda(x,x'), \hat{G}^{ss}_\Lambda(x,x')]_{\mathrm{gr}} = 0.$$

The conclusion is stable under endpoint conjugation $X \mapsto g(x)\,X\,g(x')^\dagger$ *by Corollary 4.64 and Lemma 4.106.*

Corollary 4.132 (Even-sector control of 0–*s* dynamics)**.** *Assume the hypotheses of Theorem 4.131 (any of* (B1)–(B3) *in Definition 4.127), so that* $[\hat{G}^{00}_\Lambda, \hat{G}^{ss}_\Lambda] = 0$. *Then:*

(a) Degree-0 derivation (no color flips)
For every channel $X \in \mathsf{G}$ *supported on the* 0–*s sector,* $[\hat{G}^{00}_\Lambda, X]_{\mathrm{gr}} = [\hat{G}^{00}_\Lambda, X]$, *and* $\mathrm{ad}_{\hat{G}^{00}_\Lambda}$ *acts as a degree-*0 *derivation on that sector.*

(b) Endpoint gauging of even phases
If, in addition, (B3) *holds (joint unitary diagonalizability; cf. Proposition 4.130) or flatness holds on a simply connected neighbourhood (Proposition 4.59), then phases generated by* $\hat{G}^{00}_\Lambda$ *can be pushed to the endpoints by a gauge change: for any* $\theta \in \mathbb{R}$, *conjugations by* $e^{i\theta\hat{G}^{00}_\Lambda}$ *on* 0–*s channels are realized by endpoint unitaries via Corollary 4.64, leaving holosymmetric couplings unchanged.*

Proof. (a) Diagonal blocks are even by Corollary 4.102, hence for any X the graded bracket with $\hat{G}^{00}_\Lambda$ equals the ordinary commutator: $[\hat{G}^{00}_\Lambda, X]_{\mathrm{gr}} = [\hat{G}^{00}_\Lambda, X]$. Since G is an associative $*$-algebra (Definition 4.103), $\mathrm{ad}_{\hat{G}^{00}_\Lambda}$ is a derivation. Restricted to the 0–*s* sector, degrees are preserved (no color flips) by Lemma 4.98 and Proposition 4.99(a–b). (b) Under $[\hat{G}^{00}_\Lambda, \hat{G}^{ss}_\Lambda] = 0$ (Theorem 4.131), functional calculus gives $e^{i\theta\hat{G}^{00}_\Lambda}\hat{G}^{ss}_\Lambda = \hat{G}^{ss}_\Lambda e^{i\theta\hat{G}^{00}_\Lambda}$. With (B3) or flatness, we may simultaneously diagonalize (or reduce transport to endpoint conjugation), so conjugation by $e^{i\theta\hat{G}^{00}_\Lambda}$ can be implemented as an endpoint gauge $X \mapsto g(x)\,X\,g(x')^\dagger$ (Proposition 4.54 and Corollary 4.64), which does not alter holosymmetric couplings; invariance of the graded structure under such conjugations is Lemma 4.106. □

Remark 4.133 (Interpretation and cross-chapter links)**.** Even-sector diagonals $\hat{G}^{00}$ (local symmetry persistence) and $\hat{G}^{ss}$ (holosymmetric persistence) behave "bosonically": their mutual commutation means that self-persistence amplitudes in the symmetry and holosymmetry channels can be simultaneously resolved without color mixing. Under flatness (Proposition 4.59)

and the holosymmetric constructions of Chapter 3 (eqs. (3.35) and (3.36)), this translates to a common eigenbasis for endpoint observables controlling local symmetry and global invariants, so that the corresponding effects can be measured compatibly. In Chapter 1's selector formalism, this compatibility matches the even-degree structure (Definition 4.94 and Corollary 4.102) and sits naturally inside the graded Lie $*$-algebra of channels (Proposition 4.104 and Lemma 4.106).

Graded covariant kinematics (along a path)

Let $\mathcal{A} \in \Omega^1(\mathcal{X}_q, \mathfrak{u}(\mathcal{H}_{\mathrm{em}}))$ be the (skew–Hermitian) connection entering the transporter $\mathcal{U} = \mathcal{P}\exp\!\left(-\int_\gamma \mathcal{A}\right)$ (Definitions 4.48 and 4.49). Write $\mathcal{A} = \mathcal{A}_\mu\, dx^\mu$ in local coordinates near the endpoint x. Throughout, the graded bracket $[\cdot,\cdot]_{\mathrm{gr}}$ is that of Section 4.1.7 and eq. (4.30), and degrees are as in Definition 4.94.

Definition 4.134 (Graded covariant derivative on channel blocks)**.** For each component $\hat{G}^{\alpha\beta} = \hat{G}^{\alpha\beta}_\Lambda(x,x')$ (Definition 4.61), define

$$\mathfrak{D}_\mu \hat{G}^{\alpha\beta} := \partial_\mu \hat{G}^{\alpha\beta} + [\,\mathcal{A}_\mu, \hat{G}^{\alpha\beta}\,]_{\mathrm{gr}}. \tag{4.42}$$

Here ∂_μ differentiates with respect to the first endpoint x (holding x' fixed).

Lemma 4.135 (Gauge covariance and degree preservation)**.** *Under $g : \mathcal{X}_q \to \mathrm{U}(\mathcal{H}_{\mathrm{em}})$,*

$$\mathcal{A}_\mu \mapsto \mathcal{A}'_\mu = g\,\mathcal{A}_\mu\, g^\dagger + (\partial_\mu g)\, g^\dagger, \qquad \hat{G}^{\alpha\beta}(x,x') \mapsto \hat{G}^{\alpha\beta\prime}(x,x') = g(x)\,\hat{G}^{\alpha\beta}(x,x')\,g(x')^\dagger$$

(Proposition 4.54 and Corollary 4.64), and

$$\mathfrak{D}'_\mu \hat{G}^{\alpha\beta\prime} = g(x)\left(\mathfrak{D}_\mu \hat{G}^{\alpha\beta}\right) g(x')^\dagger.$$

Moreover $\deg(\mathfrak{D}_\mu \hat{G}^{\alpha\beta}) = \deg(\hat{G}^{\alpha\beta})$ *since* $\deg(\mathcal{A}_\mu) = (0,0)$ *and the graded bracket preserves degree (Lemma 4.98).*

Proof. Direct computation with the stated gauge laws and bilinearity of $[\cdot,\cdot]_{\mathrm{gr}}$; degree preservation uses that even elements act without changing color and $\deg[X,Y]_{\mathrm{gr}} = \deg X + \deg Y$ (Section 4.1.7 and Lemma 4.98). □

Lemma 4.136 (Graded Leibniz rule)**.** *For channel-valued operators X, Y (with well-defined degrees), $\mathfrak{D}_\mu(XY) = (\mathfrak{D}_\mu X)Y + X(\mathfrak{D}_\mu Y)$.*

Proof. Expand $\partial_\mu(XY)$ and use $[\,\mathcal{A}_\mu, XY\,]_{\mathrm{gr}} = [\,\mathcal{A}_\mu, X\,]_{\mathrm{gr}}\, Y + X\,[\,\mathcal{A}_\mu, Y\,]_{\mathrm{gr}}$, which holds because $\deg(\mathcal{A}_\mu) = (0,0)$ (even). □

Proposition 4.137 (Parallel consistency along a path)**.** *Let $\gamma : [s_0, s_1] \to \mathcal{X}_q$ be C^1 with $\gamma(s_0) = x'$, $\gamma(s_1) = x$. Along γ, set $D_s := \partial_s + [A_\gamma(s), \cdot\,]_{\mathrm{gr}}$ with $A_\gamma(s) = (\gamma^*\mathcal{A})_s(\dot\gamma(s))$ (eq. (4.10)). Assume the endpoint selectors are* covariantly rigid *along γ at the readout end, i.e. $\partial_s \hat{\mathcal{F}}^\dagger_\alpha(\gamma(s)) + [A_\gamma(s), \hat{\mathcal{F}}^\dagger_\alpha(\gamma(s))] = 0$. Then for $\hat{G}^{\alpha\beta}(\gamma(s), x') = \hat{\mathcal{F}}^\dagger_\alpha(\gamma(s))\,\mathcal{U}(\gamma(s) \leftarrow x'; \gamma)\,\hat{\mathcal{F}}_\beta(x')$ one has*

$$D_s \hat{G}^{\alpha\beta}(\gamma(s), x') = 0.$$

Proof. Differentiate the defining product and use the ODE for transport $\partial_s \mathcal{U} = -A_\gamma(s)\mathcal{U}$ (Definition 4.49 and eq. (4.12)), the rigidity hypothesis at the left endpoint, and $[A_\gamma, \cdot]_{\mathrm{gr}}$ acting by derivations (Lemma 4.136). Terms cancel pairwise, yielding $D_s \hat{G}^{\alpha\beta} = 0$. □

Proposition 4.138 (Curvature identity)**.** *For any channel-valued X,*

$$[\mathfrak{D}_\mu, \mathfrak{D}_\nu]_{\mathrm{gr}} X = \Big[\, \mathcal{F}_{\mu\nu},\, X \,\Big]_{\mathrm{gr}}, \qquad \mathcal{F}_{\mu\nu} := \partial_\mu \mathcal{A}_\nu - \partial_\nu \mathcal{A}_\mu + [\mathcal{A}_\mu, \mathcal{A}_\nu]$$

($\mathcal{F}$ as in Definition 4.56). Consequently, if $\mathcal{F} = 0$ on a simply connected neighbourhood, $[\mathfrak{D}_\mu, \mathfrak{D}_\nu]_{\mathrm{gr}} = 0$ and parallel consistency integrates to path–independence up to endpoint gauge (Proposition 4.59 and Corollary 4.57).

Proof. Standard computation using (4.42) and evenness of $\mathcal{A}_\mu$. Gauge covariance of both sides follows from Lemma 4.135; the flat case follows by integrability and Proposition 4.59. □

Remark 4.139 (Kinematic reading)**.** Equation (4.42) is a bookkeeping device for endpoint–wise variations of channel blocks. Under the pointwise realization $\hat{G}^{\alpha\beta}(x, x') = \hat{\mathcal{F}}^\dagger_\alpha(x)\,\mathcal{U}(x \leftarrow x'; \gamma)\,\hat{\mathcal{F}}_\beta(x')$, the constraint $\mathfrak{D}_\mu \hat{\mathbf{G}}_\Lambda = 0$ enforces *parallel consistency* between preparation, transport, and readout, while $\mathcal{F} \neq 0$ sources deviations via Proposition 4.138, i.e. holonomy-induced effects (Definition 4.56 and eq. (4.18)). Degree preservation and gauge covariance are guaranteed by Lemmas 4.98 and 4.135.

Why the graded algebra is necessary

A *graded* algebra is the minimal, precise device that records (i) which channels are allowed to mix, (ii) how they compose, and (iii) which sign/phase rules govern their couplings—while remaining faithful to the tetralemma's four facets, to gauge covariance of transport, and to the Hermitian/operator character of the correlator. An ungraded matrix algebra would blur these distinctions and admit spurious couplings that violate the logical architecture fixed by the selector–transport formalism (Definition 4.61, Proposition 4.54, and Corollary 4.64).

Dialectic colors and additivity. Structurally, the four modes $\{0, 1, p, s\}$ encode two independent oppositions—the dyad $0 \leftrightarrow 1$ and the global/implementation axis $0 \leftrightarrow s$—with p behaving as their composite. The $\mathbb{Z}_2 \times \mathbb{Z}_2$ grading formalizes this (Definition 4.94):

$$\deg(0) = (0,0), \qquad \deg(1) = (1,0), \qquad \deg(s) = (0,1), \qquad \deg(p) = (1,1).$$

Channel components inherit color additively,

$$\deg\big(\hat{G}^{\alpha\beta}\big) = \deg(\alpha) + \deg(\beta),$$

so composition preserves color and block structure (cf. Proposition 4.99).

The graded bracket encodes selection rules. The graded (color) bracket of Section 4.1.7 and eq. (4.30)

$$[X, Y]_{\mathrm{gr}} := XY - (-1)^{\langle \deg X, \deg Y \rangle} YX, \qquad \langle (a_1, a_2), (b_1, b_2) \rangle := a_1 b_1 + a_2 b_2 \pmod 2,$$

derives the interference/transition rules instead of imposing them by hand:

- If either factor is even $(0,0)$ (e.g. any diagonal $\hat{G}^{\alpha\alpha}$ by Corollary 4.102), one gets the ordinary commutator.
- Same nontrivial color may commute or anticommute depending on the color: two $(1,0)$ or two $(0,1)$ components *anticommute* (the bilinear form gives 1), whereas two $(1,1)$ components *commute* (the bilinear form gives 0).
- Mixed colors follow the sign dictated by $\langle \deg X, \deg Y \rangle$: e.g. $(1,0)$ with $(0,1)$ commute (dot $= 0$), while $(1,0)$ with $(1,1)$ anticommute (dot $= 1$).

These sign rules are exactly what keeps Synthesis p and Holothesis s distinct yet composable with the dyad $\{0,1\}$ (see the block patterns in Proposition 4.99).

Hermiticity and gauge covariance are respected. Hermitian symmetry $(\hat{G}^{\alpha\beta})^\dagger = \hat{G}^{\beta\alpha}$ preserves degree (Lemma 4.66 and Corollary 4.102), and the unitary gauge action

$$\hat{G}^{\alpha\beta}(x,x') \;\mapsto\; g(x)\,\hat{G}^{\alpha\beta}(x,x')\,g(x')^\dagger$$

(Proposition 4.54 and Corollary 4.64) leaves product, adjoint, colors, and the bracket intact (Lemma 4.98). Thus the graded $*$-algebra is *gauge–covariant*, as required when channels are defined via the transporter $\mathcal{U}$ (Definition 4.48).

Even core acting by derivations. The grading separates *self-persistence* from *mode-changing* in a controlled way: the even subalgebra generated by $\{\hat{G}^{\alpha\alpha}\}$ acts by (graded) derivations on each colored off-diagonal class, without forcing color flips; see Section 4.1.7 and the derivation property of the graded bracket in Section 4.1.7.

Kinematics that match geometry. The unique degree-preserving, gauge-covariant coupling to the connection $\mathcal{A}$ is the graded covariant derivative

$$\mathfrak{D}_\mu \hat{G}^{\alpha\beta} = \partial_\mu \hat{G}^{\alpha\beta} + [\mathcal{A}_\mu, \hat{G}^{\alpha\beta}]_{\mathrm{gr}},$$

(Definition 4.134), which ensures parallel consistency and locates holonomy effects in the *correct* sectors in accord with the transporter's geometry (Definition 4.56 and Propositions 4.59 and 4.138).

Bottom line. The $\mathbb{Z}_2 \times \mathbb{Z}_2$ graded $*$-algebra is not ornamentation. It is the minimal structure that (i) respects the dual–dual (dyad $\times$ global) nature of the four modes, (ii) enforces the correct selection rules and signs in compositions, (iii) preserves Hermiticity and gauge covariance, and (iv) provides a clean, covariant template for kinematics. Without grading, one either loses these guarantees or must re-impose them piecemeal (cf. Section 4.1.7 for model assumptions that the graded framework cleanly supports).

4.2 Holosymmetric Words in Channel Algebra

Words as resolved histories. A *word* is an ordered product that spells out a channel at a chosen "temporal" resolution: selectors at the endpoints and parallel transporters (and, when present, device evolutions) in between. Concretely, for a chain of piecewise C^1 paths $\gamma_1 : x' \to x_1, \dots, \gamma_n : x_{n-1} \to x$,

$$W^{\alpha\beta}[\gamma_n, \dots, \gamma_1] := \hat{\mathcal{F}}^\dagger_\alpha(x)\, \mathcal{U}(x \leftarrow x_{n-1}; \gamma_n) \cdots \mathcal{U}(x_1 \leftarrow x'; \gamma_1)\, \hat{\mathcal{F}}_\beta(x').$$

By functoriality and inversion of transport (Proposition 4.52 and eq. (4.14)), this collapses to the single–channel form of Definition 4.61:

$$W^{\alpha\beta}[\gamma_n, \dots, \gamma_1] = \hat{\mathcal{F}}^\dagger_\alpha(x)\, \mathcal{U}\Big(x \leftarrow x';\, \gamma_n \circ \cdots \circ \gamma_1\Big)\, \hat{\mathcal{F}}_\beta(x') = \hat{G}^{\alpha\beta}_\Lambda(x, x').$$

Thus a word is the same channel seen at a finer "path resolution"; words multiply by concatenating paths and composing the corresponding operators.

Remark 4.140 (Typing and spectral domain)**.** Words are only formed when *typing constraints* match: the terminal mode/locus of one letter equals the initial mode/locus of the next (cf. Proposition 4.85 for insertion of completeness at intermediate loci). All operators act on the spectrally regulated fibres $\operatorname{Ran} E^{(x)}([0, \lambda_{\text{cutoff}}])$ (see eq. (4.303)), ensuring strong continuity of $\mathcal{U}$ along any piecewise C^1 path (Definitions 4.48 and 4.49).

Endpoint covariance and frame consistency. Selectors originate in Chapter 1, and the transporter is the gauge–covariant parallel transport of Definitions 4.48 and 4.49. Under a local gauge, selectors and transport transform as in Proposition 4.54 and Corollary 4.64. Interior gauge factors cancel along the product, so any word transforms by conjugation at its *endpoints* x, x'. This is precisely the operational role of words: frame–align the modal data along the path via $\mathcal{U}$, then read off with $\hat{\mathcal{F}}^\dagger_\alpha$ and $\hat{\mathcal{F}}_\beta$ in a way that is covariant at the endpoints.

Restriction to the Twofold sector and duality collapse. Restricting indices to $\alpha, \beta \in \{0, 1\}$ places words in the Twofold block $\hat{\mathbf{G}}_{(0,1)}(x, x')$ (eq. (4.25)). In the duality–collapse regime (Definition 4.87), off–block expectations vanish and the coincident–limit inner product induced by the Fourfold reduces to a bona fide qubit inner product on the Twofold fibre; operational predictions of all Twofold words then coincide with ordinary two–level kinematics (Proposition 4.90). Outside this regime, words that touch p or s diagnose coherent composition and global/implementation–invariant structure beyond the dyad.

Holosymmetric mechanisms as words. Chapter 3 realises holosymmetric comparisons by composing preparation, control, and readout unitaries (e.g. $\mathbf{U}_{\text{ctrl}}$ and its Hadamard conjugate, Definition 2.114 and eqs. (2.75) and (2.78)), acting on dual PQS registers (Definition 2.101). These mechanism–level composites are words at the device layer. When sandwiched by $\hat{\mathcal{F}}^\dagger_\alpha(\cdot)$ and $\hat{\mathcal{F}}_\beta(\cdot)$ and, if needed, a geometric $\mathcal{U}$, they produce the corresponding *holosymmetric channel words* whose expectations yield the holosymmetric effects (eq. (3.35)).

Behavior under flatness and path refinement. On simply connected regions with $\mathcal{F} = 0$ (Definition 4.56), path–ordered transport is path independent up to an endpoint gauge (Proposition 4.59); words that differ only by a refinement or wiggling of intermediate segments coincide after endpoint conjugation. Conversely, nonzero curvature produces holonomy (eq. (4.18)), and words expose this through the dependence on the concatenated path.

Graded bookkeeping for words. With the $\mathbb{Z}_2^2$ dialectic grading of Definition 4.94, a word inherits degree additively from its channel pieces; adjoint, gauge covariance, and composition are compatible with degree (Lemma 4.98). This graded bookkeeping governs selection rules and block patterns (Proposition 4.99) and underlies the graded Lie $*$-algebra structure introduced in Section 4.1.7.

Definition 4.141 (Holosymmetric channel alphabet and words)**.** Fix loci $x_0, \dots, x_n \in \mathcal{X}_q$ and $\mathcal{I} = \{0, 1, p, s\}$. The *channel alphabet* consists of all typed letters

$$\mathsf{L} \;:=\; \Big\{\, \hat{G}_\Lambda^{\alpha\beta}(x_j, x_{j-1}) \;:\; \alpha, \beta \in \mathcal{I},\; j = 1, \dots, n \,\Big\},$$

where each letter is the Fourfold channel block of Definition 4.61: $\hat{G}_\Lambda^{\alpha\beta}(x_j, x_{j-1}) = \hat{\mathcal{F}}_\alpha^\dagger(x_j)\, \mathcal{U}(x_j \leftarrow x_{j-1}; \gamma_j)\, \hat{\mathcal{F}}_\beta(x_{j-1})$, with $\mathcal{U}$ as in Definitions 4.48 and 4.49.

A *(channel) word* is a finite product

$$W \;:=\; \hat{G}_\Lambda^{\alpha_n \alpha_{n-1}}(x_n, x_{n-1})\, \hat{G}_\Lambda^{\alpha_{n-1} \alpha_{n-2}}(x_{n-1}, x_{n-2}) \cdots \hat{G}_\Lambda^{\alpha_1 \alpha_0}(x_1, x_0),$$

with *typing constraints* that successive endpoints and mode indices match (so the intermediate locus x_j and label α_j are "summed over" by composition; cf. Propositions 4.52 and 4.85). The *length* of W is the number of letters. Its *degree* is $\deg(W) := \sum_{k=1}^n \deg(\hat{G}^{\alpha_k \alpha_{k-1}})$ with deg as in Definition 4.94, and is well defined modulo the typing constraints.

Remark 4.142 (Endpoints, base, and transporter)**.** In all channel-word statements, endpoints x, x' are points of the PQS continuum $\mathcal{C}_d$ (Definition 4.617). Paths γ are piecewise C^1 curves in $\mathcal{C}_d$, and $\mathcal{U}(x \leftarrow x'; \gamma)$ is the gauge–covariant transporter of Section 4.19.1. Fibre restrictions at endpoints are as in (4.303). The emergent space $\mathcal{X}_q$ and the projection $\pi_q : \mathcal{C}_d \to \mathcal{X}_q$ are only introduced after forming the operational quotient (Definition 4.623 and eq. (4.322)); no statement in the words calculus depends on $\mathcal{X}_q$ prior to that quotient.

Proposition 4.143 (Covariance, adjoint, functoriality for words)**.** *Every channel word W satisfies:*

(i) Gauge covariance (endpoints)

Under a gauge map $g : \mathcal{X}_q \to \mathrm{U}(\mathcal{H}_{\mathrm{em}})$ one has

$$W(x_n, x_0) \;\mapsto\; g(x_n)\, W(x_n, x_0)\, g(x_0)^\dagger,$$

by repeated use of Proposition 4.54 and Corollary 4.64.

(ii) Adjoint/Hermitian symmetry

$W(x_n, x_0)^\dagger = W^\sharp(x_0, x_n)$ *where $W^\sharp$ is the word with reversed order and conjugated letters; uses Lemma 4.66 and Proposition 4.52.*

(iii) Path composition
Concatenating geometric segments $(x_0 \to \cdots \to x_m)$ *and* $(x_m \to \cdots \to x_n)$ *gives word multiplication, thanks to Proposition 4.52.*

(iv) Degree additivity
$\deg(W_1W_2) = \deg(W_1) + \deg(W_2)$ *and* $\deg(W^\dagger) = \deg(W)$, *by Definition 4.94 and Lemma 4.98.*

Proof of Proposition 4.143. (i) Gauge covariance. Each letter is $\hat{G}_\Lambda^{\alpha_j\alpha_{j-1}}(x_j, x_{j-1}) = \hat{\mathcal{F}}_{\alpha_j}^\dagger(x_j)\,\mathcal{U}(x_j \leftarrow x_{j-1}; \gamma_j)\,\hat{\mathcal{F}}_{\alpha_{j-1}}(x_{j-1})$. Under $g : \mathcal{X}_q \to \mathrm{U}(\mathcal{H}_{\mathrm{em}})$, Proposition 4.54 and Corollary 4.64 give $\hat{G}^{\alpha_j\alpha_{j-1}} \mapsto g(x_j)\hat{G}^{\alpha_j\alpha_{j-1}}g(x_{j-1})^\dagger$. Interior factors $g(x_j)^\dagger g(x_j)$ cancel in the ordered product, leaving $W(x_n, x_0) \mapsto g(x_n)W(x_n, x_0)g(x_0)^\dagger$.

(ii) Adjoint/Hermitian symmetry. Define the reverse–conjugate word

$$W^\sharp(x_0, x_n) := \left(\hat{G}_\Lambda^{\alpha_1\alpha_0}(x_1, x_0)\right)^\dagger \cdots \left(\hat{G}_\Lambda^{\alpha_n\alpha_{n-1}}(x_n, x_{n-1})\right)^\dagger.$$

By Lemma 4.66 and the adjoint reversing products, $W(x_n, x_0)^\dagger = W^\sharp(x_0, x_n)$. Equivalently, use Proposition 4.52 to note that adjoint reverses the underlying path concatenation.

(iii) Path composition. Concatenating segments corresponds to composing the intervening transporters by Proposition 4.52 and eq. (4.14); with selectors fixed at endpoints, this is exactly multiplication of the corresponding letters, hence of words.

(iv) Degree additivity. By Definition 4.94, each letter has degree $\deg(\alpha_j) + \deg(\alpha_{j-1})$; additivity under products and preservation under adjoint are Lemma 4.98. □

Corollary 4.144 (Endpoint degree rule for words)**.** *Let* $W = \hat{G}_\Lambda^{\alpha_n\alpha_{n-1}}(x_n, x_{n-1}) \cdots \hat{G}_\Lambda^{\alpha_1\alpha_0}(x_1, x_0)$ *be a channel word (Definition 4.141). Then*

$$\deg(W) = \deg(\alpha_n) + \deg(\alpha_0) \in \mathbb{Z}_2^2.$$

Proof. By Definition 4.94, each letter has degree $\deg(\alpha_j) + \deg(\alpha_{j-1})$. Summing from $j = 1$ to n yields $\sum_{j=1}^n \deg(\alpha_j) + \sum_{j=1}^n \deg(\alpha_{j-1}) = \deg(\alpha_n) + \deg(\alpha_0) + 2\sum_{j=1}^{n-1} \deg(\alpha_j)$, and the middle sum vanishes in $\mathbb{Z}_2^2$. □

Lemma 4.145 (Word-level graded selection and endpoint degree)**.** *Let*

$$W = \hat{G}_\Lambda^{\alpha_n\alpha_{n-1}}(x_n, x_{n-1}) \cdots \hat{G}_\Lambda^{\alpha_1\alpha_0}(x_1, x_0)$$

be a (channel) word as in Definition 4.141, with typing constraints satisfied. Then:

(i) Endpoint degree rule
The degree of the endpoint

$$\deg(W) = \sum_{j=1}^n \deg\left(\hat{G}_\Lambda^{\alpha_j\alpha_{j-1}}\right) = \deg(\alpha_n) + \deg(\alpha_0) \ \in \ \mathbb{Z}_2^2.$$

In particular, the total degree of W *depends only on its endpoint mode labels and is independent of the chosen factorization/path refinement.*

(ii) Stability under adjoint and gauge

$W^\dagger$ has the same degree, $\deg(W^\dagger) = \deg(W)$*, and under a gauge change* $g : \mathcal{X}_q \to \mathrm{U}(\mathcal{H}_{\mathrm{em}})$ *one has* $W(x_n, x_0) \mapsto g(x_n)\, W(x_n, x_0)\, g(x_0)^\dagger$ *with degree unchanged (Proposition 4.143, Lemma 4.98).*

(iii) Word-level selection rule (extension of Proposition 4.99)

If W is expanded by inserting dialectically complete resolutions at intermediate loci (Proposition 4.85), then every summand word $\widetilde{W}$ has the same *total degree* $\deg(\alpha_n) + \deg(\alpha_0)$*. Hence any putative contribution with mismatched total color cancels identically.*

Proof. (i) By Definition 4.94, each letter has degree $\deg(\alpha_j) + \deg(\alpha_{j-1})$. Summing over $j = 1, \ldots, n$ gives

$$\sum_{j=1}^{n} \Big(\deg(\alpha_j) + \deg(\alpha_{j-1}) \Big) = \deg(\alpha_n) + \deg(\alpha_0) + 2 \sum_{j=1}^{n-1} \deg(\alpha_j),$$

and the middle sum vanishes in $\mathbb{Z}_2^2$, yielding the endpoint formula.

(ii) Gauge covariance of words and adjoint reversal are in Proposition 4.143. Degree preservation under adjoint and additivity under products are in Lemma 4.98.

(iii) Inserting completeness at intermediates rewrites W as a finite sum of words with the same endpoints (α_0, α_n) (cf. Proposition 4.85). Each summand is a product of letters, so by (i) its degree is $\deg(\alpha_n) + \deg(\alpha_0)$, independent of the internal index sequence. □

Corollary 4.146 (Flatness reduction for words)**.** *If $\mathcal{F} = 0$ on a simply connected neighbourhood (Chapter 4; Proposition 4.59), then for any two factorizations of the same endpoints (x_0, x_n) the corresponding words agree up to endpoint gauge $g(x_n)\,(\cdot)\,g(x_0)^\dagger$. In particular, single-letter and multi-letter factorizations are equivalent in that region.*

Proof of Corollary 4.146. On a simply connected region with $\mathcal{F} = 0$, Proposition 4.59 implies that each transporter $\mathcal{U}(x_j \leftarrow x_{j-1}; \gamma_j)$ depends only on endpoints and reduces (after gauge choice) to the identity. Hence any factorization yields the same endpoint operator up to overall conjugation by $g(x_n), g(x_0)$, which establishes the claim for single- vs multi-letter factorizations. □

Remark 4.147 (Flatness and device–geometry separation)**.** In a flat, simply connected region (Proposition 4.59), the geometric part $\mathcal{U}(x \leftarrow x'; \gamma)$ reduces to an endpoint gauge choice. Word-level distinctions then live entirely in the device factor $\mathsf{W}_{\chi_{\mathrm{out}} \leftarrow \chi_{\mathrm{in}}}$, while endpoint conjugation governs the overall covariance (Proposition 4.167).

Before introducing Definition 4.148, we briefly recall the Chapter 2 setting that the holosymmetric wordwise construction lives in. The holosymmetric device operates on the composite Hilbert space

$$\mathcal{H}_{\mathrm{Holo}} \;=\; \mathcal{H}_c \otimes \mathcal{H}_{\mathrm{RR}}^{(2n)} \quad \text{(cf. Definition 2.101 and eq. (2.62)).}$$

Here, $\mathcal{H}_c$ is the control register admitting the computational basis $\{|\mathbf{0}\rangle, |\mathbf{1}\rangle\}$ and the X-basis $\{|+\rangle, |-\rangle\}$, while $\mathcal{H}_{\mathrm{RR}}^{(2n)}$ is the dual-register arena on which the two parallel histories

act. The device alphabet $\mathsf{A}_{\mathrm{holo}}$ collects the primitive letters that generate every device-level operation used in the interferometric constructions of Chapter 2: (i) the control projectors $|\mathbf{0}\rangle\langle\mathbf{0}|_c$, $|\mathbf{1}\rangle\langle\mathbf{1}|_c$ for preparation and readout in the computational basis of $\mathcal{H}_c$; (ii) the control unitaries $U_c^{\mathrm{prep}}, U_c^{\mathrm{read}}$ (with the Hadamard H_c providing the canonical switch to the X-basis used in interference readout); (iii) the RR path operators $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$ and $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$, which implement the two branch evolutions defined by the parallel-path construction (Definition 2.108 and eqs. (2.69) and (2.70)); and (iv) the controlled and Hadamard-wrapped stacks $\mathbf{U}_{\mathrm{ctrl}}$ and $\mathbf{U}_{\mathrm{Had}}$ that realize, respectively, branch selection and coherent branch interference on the control line (Definitions 2.114 and 2.119 and eqs. (2.75), (2.78) and (2.79)). Any finite product of letters from $\mathsf{A}_{\mathrm{holo}}$ defines a *holosymmetric word* $\mathbf{W} \in \langle \mathsf{A}_{\mathrm{holo}} \rangle$ acting on $\mathcal{H}_{\mathrm{Holo}}$.

Given control boundary conditions $|\chi_{\mathrm{in}}\rangle, |\chi_{\mathrm{out}}\rangle \in \mathcal{H}_c$, the corresponding *RR–matrix element* of a holosymmetric word is the branch-resolved operator

$$\mathsf{W}_{\chi_{\mathrm{out}} \leftarrow \chi_{\mathrm{in}}} := (\langle \chi_{\mathrm{out}}|_c \otimes I)\, \mathbf{W}\, (|\chi_{\mathrm{in}}\rangle_c \otimes I) \quad (\text{eq. } (4.45)). \tag{4.43}$$

This partial matrix element "reads off" the RR evolution conditioned on the control transition $\chi_{\mathrm{in}} \to \chi_{\mathrm{out}}$, and it is the natural object in which pathwise and interference statements of Chapter 2 become algebraically transparent. Two canonical specializations recover the standard constructions exactly. First, for the controlled splitter

$$\mathbf{U}_{\mathrm{ctrl}} \;=\; |\mathbf{0}\rangle\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} \;+\; |\mathbf{1}\rangle\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} \quad (\textit{eq. } (2.75)),$$

the only nonzero RR–matrix elements in the computational basis are

$$\mathsf{W}_{\mathbf{0}\leftarrow\mathbf{0}} = \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}, \qquad \mathsf{W}_{\mathbf{1}\leftarrow\mathbf{1}} = \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}, \qquad \mathsf{W}_{\mathbf{0}\leftarrow\mathbf{1}} = \mathsf{W}_{\mathbf{1}\leftarrow\mathbf{0}} = 0,$$

which is precisely the parallel-path decomposition. Second, for the Hadamard-wrapped interferometer $\mathbf{U}_{\mathrm{Had}} = (H_c \otimes I)\, \mathbf{U}_{\mathrm{ctrl}}\, (H_c \otimes I)$ (eq. (2.78)), working in the X-basis $|\pm\rangle_c = (|\mathbf{0}\rangle \pm |\mathbf{1}\rangle)/\sqrt{2}$ yields

$$\mathsf{W}_{+\leftarrow+} \;=\; \tfrac{1}{2}\Big(\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}\Big) \;=\; \tfrac{1}{2}\,\hat{\mathbf{U}}_{+}, \qquad \mathsf{W}_{-\leftarrow+} \;=\; \tfrac{1}{2}\Big(\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} - \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}\Big) \;=\; \tfrac{1}{2}\,\hat{\mathbf{U}}_{-},$$

and similarly for the other control transitions, thereby realizing the sum/difference interferometers of eq. (2.79) directly as RR–matrix elements.

Composition of words descends to a *control-resolved convolution* on RR–matrix elements. For any chosen control basis $\mathcal{B}_c$ (e.g., computational $\{|\mathbf{0}\rangle, |\mathbf{1}\rangle\}$ or X-basis $\{|+\rangle, |-\rangle\}$),

$$(\mathbf{W}_2 \mathbf{W}_1)_{\chi_{\mathrm{out}} \leftarrow \chi_{\mathrm{in}}} \;=\; \sum_{\chi \in \mathcal{B}_c} \mathsf{W}_{2,\ \chi_{\mathrm{out}} \leftarrow \chi}\ \mathsf{W}_{1,\ \chi \leftarrow \chi_{\mathrm{in}}}, \tag{4.44}$$

so algebraic manipulations and proofs in the wordwise formalism can be carried out entirely at the level of RR–matrix elements, with the control modes furnishing the natural contraction indices.

Finally, we record the link back to the Chapter 2 dynamics and generators. Each RR path operator admits an internal factorization through the parasymmetric building blocks,

$$\hat{\mathbf{U}}_{\mathrm{RR}}^{(\cdot)} \;\rightsquigarrow\; U_L \;\otimes\; U_P \;\otimes\; U_G, \qquad U_P \;=\; \exp\!\Big(i \sum_{i<j} J_{ij}\, \big(P_{\mathrm{para}}^{(ij)} - P_{\mathrm{antipara}}^{(ij)}\big)\Big),$$

consistent with the pairwise generator $\hat{F}^{(\mathrm{pair})}_{\mathrm{para}}$ and the composite evolution $U_N^{(\odot)}$ from Chapter 2. While the wordwise viewpoint treats such RR evolutions as opaque letters of $\mathsf{A}_{\mathrm{holo}}$, unitarity on $\mathcal{H}_{\mathrm{Holo}}$ enforces isometric *mixing bounds* on their RR–matrix elements—constraints that will be invoked later in the wordwise kinematics and mixing analysis.

Definition 4.148 (Holosymmetric word (device level))**.** In the holosymmetric space of Chapter 2 (Definition 2.101 and eq. (2.62)), let the *device alphabet* be

$$\mathsf{A}_{\mathrm{holo}} = \Big\{\, |\mathbf{0}\rangle\langle\mathbf{0}|_c,\ |\mathbf{1}\rangle\langle\mathbf{1}|_c,\ U_c^{\mathrm{prep}}, U_c^{\mathrm{read}},\ \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N},\ \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} \,\Big\},$$

with the RR path operators and control gates as in Definitions 2.108, 2.114 and 2.119 and eqs. (2.69) and (2.70). A *holosymmetric word* is any finite product $\mathbf{W} \in \langle \mathsf{A}_{\mathrm{holo}} \rangle$ acting on $\mathcal{H}_{\mathrm{Holo}} = \mathcal{H}_c \otimes \mathcal{H}_{\mathrm{RR}}^{(2n)}$.

Given control states $|\chi_{\mathrm{in}}\rangle, |\chi_{\mathrm{out}}\rangle \in \mathcal{H}_c$, the *RR-matrix element* (branch-resolved word) is the operator on $\mathcal{H}_{\mathrm{RR}}^{(2n)}$:

$$\mathsf{W}_{\chi_{\mathrm{out}} \leftarrow \chi_{\mathrm{in}}} \;:=\; \Big(\langle\chi_{\mathrm{out}}|_c \otimes I\Big)\ \mathbf{W}\ \Big(|\chi_{\mathrm{in}}\rangle_c \otimes I\Big). \tag{4.45}$$

For $\mathbf{W} = \mathbf{U}_{\mathrm{ctrl}}$ or $\mathbf{U}_{\mathrm{Had}}$ this reproduces the parallel-path decompositions and sum/difference interferometers of eqs. (2.75), (2.78) and (2.79).

RR dual-register and holosymmetric device space (recall from Ch. 2). Let A and B denote the two n-PQS registers. The dual-register Hilbert space is

$$\mathcal{H}_{\mathrm{RR}}^{(2n)} \;:=\; \mathcal{H}_{\mathrm{PQS}}^{(n,A)} \otimes \mathcal{H}_{\mathrm{PQS}}^{(n,B)},$$

and the holosymmetric device acts on

$$\mathcal{H}_{\mathrm{Holo}} \;=\; \mathcal{H}_c \otimes \mathcal{H}_{\mathrm{RR}}^{(2n)} \quad \text{(cf. Definition 2.101 and eq. (2.62)).}$$

Here $\mathcal{H}_c$ is the control register with computational basis $\{|\mathbf{0}\rangle, |\mathbf{1}\rangle\}$ and X-basis $\{|+\rangle, |-\rangle\}$. The two parallel RR evolutions $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$ and $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$ are the branch operators of the parallel-path construction (cf. Definition 2.108 and eqs. (2.69) and (2.70)), which are multiplexed by the control via $\mathbf{U}_{\mathrm{ctrl}}$ and Hadamard-wrapped in $\mathbf{U}_{\mathrm{Had}}$ for interference (cf. Definitions 2.114 and 2.119 and eqs. (2.75), (2.78) and (2.79)).

Wordwise alphabet and words (device level). Let $\mathsf{A}_{\mathrm{holo}}$ denote the device alphabet consisting of the control projectors $\{|\mathbf{0}\rangle\langle\mathbf{0}|_c, |\mathbf{1}\rangle\langle\mathbf{1}|_c\}$, control unitaries $\{U_c^{\mathrm{prep}}, U_c^{\mathrm{read}}\}$ (e.g. H_c), and the RR branch operators $\{\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}, \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}\}$, together with the controlled and Hadamard-conjugated stacks $\{\mathbf{U}_{\mathrm{ctrl}}, \mathbf{U}_{\mathrm{Had}}\}$. A *holosymmetric word* is any finite product $\mathbf{W} \in \langle \mathsf{A}_{\mathrm{holo}} \rangle$ acting on $\mathcal{H}_{\mathrm{Holo}}$ (cf. Definition 4.148).

RR–matrix elements (branch-resolved operators). Given $|\chi_{\mathrm{in}}\rangle, |\chi_{\mathrm{out}}\rangle \in \mathcal{H}_c$, the RR–matrix element of a word $\mathbf{W}$ is

$$\mathsf{W}_{\chi_{\mathrm{out}} \leftarrow \chi_{\mathrm{in}}} \;:=\; (\langle\chi_{\mathrm{out}}|_c \otimes I)\ \mathbf{W}\ (|\chi_{\mathrm{in}}\rangle_c \otimes I), \tag{4.46}$$

an operator on $\mathcal{H}_{\mathrm{RR}}^{(2n)}$ that "reads off" the RR branch selected by the control transition (cf. eq. (4.45)). Throughout, we use either the computational basis $\mathcal{B}_c^{(Z)} = \{|\mathbf{0}\rangle, |\mathbf{1}\rangle\}$ or the X-basis $\mathcal{B}_c^{(X)} = \{|+\rangle, |-\rangle\}$ with $|\pm\rangle = (|\mathbf{0}\rangle \pm |\mathbf{1}\rangle)/\sqrt{2}$.

Lemma 4.149 (Control-resolved convolution)**.** *For any words* $\mathbf{W}_2, \mathbf{W}_1 \in \langle \mathsf{A}_{\mathrm{holo}} \rangle$ *and any orthonormal control basis* $\mathcal{B}_c \subset \mathcal{H}_c$*,*

$$(\mathbf{W}_2\mathbf{W}_1)_{\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}} = \sum_{\chi\in\mathcal{B}_c} \mathsf{W}_{2,\ \chi_{\mathrm{out}}\leftarrow\chi}\ \mathsf{W}_{1,\ \chi\leftarrow\chi_{\mathrm{in}}}. \tag{4.47}$$

Proof. Fix words $\mathbf{W}_2, \mathbf{W}_1 \in \langle \mathsf{A}_{\mathrm{holo}} \rangle$ (cf. Definition 4.148), and let $|\chi_{\mathrm{in}}\rangle, |\chi_{\mathrm{out}}\rangle \in \mathcal{H}_c$ be arbitrary. By the definition of RR–matrix elements $\mathsf{W}_{\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}}$ (see eq. (4.45)), we have

$$(\mathbf{W}_2\mathbf{W}_1)_{\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}} = \Big(\langle\chi_{\mathrm{out}}|_c \otimes I\Big)\ \mathbf{W}_2\mathbf{W}_1\ \Big(|\chi_{\mathrm{in}}\rangle_c \otimes I\Big).$$

Insert the resolution of the identity on the control register in any orthonormal control basis $\mathcal{B}_c \subset \mathcal{H}_c$, $I_c = \sum_{\chi\in\mathcal{B}_c} |\chi\rangle\langle\chi|$, tensorized with the RR identity I. This yields

$$\begin{aligned}(\mathbf{W}_2\mathbf{W}_1)_{\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}} &= \sum_{\chi\in\mathcal{B}_c} \Big(\langle\chi_{\mathrm{out}}|_c \otimes I\Big)\ \mathbf{W}_2\ \Big(|\chi\rangle_c\langle\chi|_c \otimes I\Big)\ \mathbf{W}_1\ \Big(|\chi_{\mathrm{in}}\rangle_c \otimes I\Big) \\ &= \sum_{\chi\in\mathcal{B}_c} \underbrace{\Big(\langle\chi_{\mathrm{out}}|_c \otimes I\Big)\ \mathbf{W}_2\ \Big(|\chi\rangle_c \otimes I\Big)}_{\mathsf{W}_{2,\ \chi_{\mathrm{out}}\leftarrow\chi}}\ \underbrace{\Big(\langle\chi|_c \otimes I\Big)\ \mathbf{W}_1\ \Big(|\chi_{\mathrm{in}}\rangle_c \otimes I\Big)}_{\mathsf{W}_{1,\ \chi\leftarrow\chi_{\mathrm{in}}}}.\end{aligned}$$

Each underbraced factor is, by eq. (4.45), precisely the RR–matrix element of the corresponding word with the indicated control transition. Hence

$$(\mathbf{W}_2\mathbf{W}_1)_{\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}} = \sum_{\chi\in\mathcal{B}_c} \mathsf{W}_{2,\ \chi_{\mathrm{out}}\leftarrow\chi}\ \mathsf{W}_{1,\ \chi\leftarrow\chi_{\mathrm{in}}},$$

which is eq. (4.47). The choice of $\mathcal{B}_c$ is immaterial since any two orthonormal control bases are related by a unitary (e.g. H_c maps the computational basis to the X-basis; cf. Definition 2.119), and inserting I_c is basis-independent. This completes the proof. □

Remark 4.150 (Meaning of control-resolved convolution)**.** The identity Lemma 4.149 and eq. (4.47) says that, once we fix an orthonormal control basis $\mathcal{B}_c \subset \mathcal{H}_c$, every device word $\mathbf{W} \in \langle \mathsf{A}_{\mathrm{holo}} \rangle$ behaves as a $|\mathcal{B}_c| \times |\mathcal{B}_c|$ *block operator matrix* whose entries are precisely the RR–matrix elements $\{\mathsf{W}_{\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}}\}_{\chi_{\mathrm{out}},\chi_{\mathrm{in}}\in\mathcal{B}_c}$ (cf. Definition 4.148 and eq. (4.45)). In this representation,

$$\mathbf{W} \ \longleftrightarrow\ \Big[\ \mathsf{W}_{\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}}\ \Big]_{\chi_{\mathrm{out}},\chi_{\mathrm{in}}\in\mathcal{B}_c},$$

and composition of words corresponds to *ordinary block-matrix multiplication* with operator-valued blocks on $\mathcal{H}_{\mathrm{RR}}^{(2n)}$. Concretely, the product $\mathbf{W}_2\mathbf{W}_1$ has $(\chi_{\mathrm{out}}, \chi_{\mathrm{in}})$-entry given by the contraction over the intermediate control label χ,

$$(\mathbf{W}_2\mathbf{W}_1)_{\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}} = \sum_{\chi\in\mathcal{B}_c} \mathsf{W}_{2,\ \chi_{\mathrm{out}}\leftarrow\chi}\ \mathsf{W}_{1,\ \chi\leftarrow\chi_{\mathrm{in}}},$$

which is exactly eq. (4.47). Equivalently, eq. (4.47) is nothing but the insertion of the control identity $I_c = \sum_{\chi\in\mathcal{B}_c} |\chi\rangle\langle\chi|$ between $\mathbf{W}_2$ and $\mathbf{W}_1$ and evaluation of partial matrix elements on $\mathcal{H}_c$ (cf. eq. (4.45)).

Three immediate consequences clarify its role in the wordwise calculus. *(i) Basis covariance:* the statement is independent of the choice of $\mathcal{B}_c$, since any two orthonormal control

bases are related by a unitary on $\mathcal{H}_c$ (e.g. the Hadamard H_c that implements the $Z \leftrightarrow X$ change of basis; cf. Definition 2.119). *(ii) Interferometric readability:* choosing $\mathcal{B}_c = \{|+\rangle, |-\rangle\}$ turns the blocks into the sum/difference channels that realize the standard interferometer identities of Chapter 2 (cf. eqs. (2.78) and (2.79)). *(iii) Unitarity constraints:* applying eq. (4.47) to $\mathbf{W}^\dagger\mathbf{W} = I$ and $\mathbf{W}\mathbf{W}^\dagger = I$ yields the block *mixing bounds* of Corollary 4.151, i.e. left/right RR–isometries obtained by summing over control outcomes. Thus Lemma 4.149 is the algebraic backbone that lets us carry out device-level proofs purely at the RR–matrix level while retaining full operator-level correctness on $\mathcal{H}_{\text{Holo}}$.

Corollary 4.151 (Block unitarity ⇒ mixing bounds)**.** *If* $\mathbf{W}$ *is unitary on* $\mathcal{H}_{\text{Holo}}$*, then for any orthonormal control basis* $\mathcal{B}_c$*,*

$$\sum_{\chi\in\mathcal{B}_c} \mathsf{W}^\dagger_{\chi\leftarrow\chi_{\text{in}}}\,\mathsf{W}_{\chi\leftarrow\chi_{\text{in}}} \;=\; I_{\text{RR}}, \qquad \sum_{\chi\in\mathcal{B}_c} \mathsf{W}_{\chi_{\text{out}}\leftarrow\chi}\,\mathsf{W}^\dagger_{\chi_{\text{out}}\leftarrow\chi} \;=\; I_{\text{RR}}. \tag{4.48}$$

Proof. Let $\mathbf{W} \in \langle \mathsf{A}_{\text{holo}} \rangle$ be unitary on $\mathcal{H}_{\text{Holo}} = \mathcal{H}_c \otimes \mathcal{H}_{\text{RR}}^{(2n)}$ (cf. Definitions 2.101 and 4.148 and eq. (2.62)). Fix any orthonormal control basis $\mathcal{B}_c \subset \mathcal{H}_c$ and a control state $|\chi_{\text{in}}\rangle \in \mathcal{B}_c$. By unitarity, $\mathbf{W}^\dagger\mathbf{W} = I_{\mathcal{H}_{\text{Holo}}}$. Taking the RR–matrix element in the sense of eq. (4.45), we obtain

$$(\mathbf{W}^\dagger\mathbf{W})_{\chi_{\text{in}}\leftarrow\chi_{\text{in}}} = (\langle\chi_{\text{in}}|_c \otimes I)\,\mathbf{W}^\dagger\mathbf{W}\,(|\chi_{\text{in}}\rangle_c \otimes I) = (\langle\chi_{\text{in}}|_c \otimes I)\,I_{\mathcal{H}_{\text{Holo}}}\,(|\chi_{\text{in}}\rangle_c \otimes I) = I_{\text{RR}}.$$

On the other hand, by the control-resolved convolution Lemma 4.149 and eq. (4.47) with $\mathbf{W}_2 = \mathbf{W}^\dagger$ and $\mathbf{W}_1 = \mathbf{W}$,

$$(\mathbf{W}^\dagger\mathbf{W})_{\chi_{\text{in}}\leftarrow\chi_{\text{in}}} = \sum_{\chi\in\mathcal{B}_c} (\mathbf{W}^\dagger)_{\chi_{\text{in}}\leftarrow\chi}\;\mathbf{W}_{\chi\leftarrow\chi_{\text{in}}}.$$

Using the definition of RR–matrix elements eq. (4.45) and the adjoint rule $(\langle a| \otimes I)\,\mathbf{W}^\dagger\,(|b\rangle \otimes I) = \Big((\langle b| \otimes I)\,\mathbf{W}\,(|a\rangle \otimes I)\Big)^\dagger$, we identify $(\mathbf{W}^\dagger)_{\chi_{\text{in}}\leftarrow\chi} = \Big(\mathbf{W}_{\chi\leftarrow\chi_{\text{in}}}\Big)^\dagger$. Hence

$$I_{\text{RR}} = (\mathbf{W}^\dagger\mathbf{W})_{\chi_{\text{in}}\leftarrow\chi_{\text{in}}} = \sum_{\chi\in\mathcal{B}_c} \mathbf{W}^\dagger_{\chi\leftarrow\chi_{\text{in}}}\,\mathbf{W}_{\chi\leftarrow\chi_{\text{in}}},$$

which is the first identity in eq. (4.48).

For the second identity, apply the same reasoning to $\mathbf{W}\mathbf{W}^\dagger = I_{\mathcal{H}_{\text{Holo}}}$ and take the RR–matrix element $(\mathbf{W}\mathbf{W}^\dagger)_{\chi_{\text{out}}\leftarrow\chi_{\text{out}}} = I_{\text{RR}}$. By Lemma 4.149 and eq. (4.47) with $\mathbf{W}_2 = \mathbf{W}$ and $\mathbf{W}_1 = \mathbf{W}^\dagger$,

$$I_{\text{RR}} = \sum_{\chi\in\mathcal{B}_c} \mathbf{W}_{\chi_{\text{out}}\leftarrow\chi}\,(\mathbf{W}^\dagger)_{\chi\leftarrow\chi_{\text{out}}} = \sum_{\chi\in\mathcal{B}_c} \mathbf{W}_{\chi_{\text{out}}\leftarrow\chi}\,\Big(\mathbf{W}_{\chi_{\text{out}}\leftarrow\chi}\Big)^\dagger,$$

which is the second identity in eq. (4.48). Since inserting the control identity $I_c = \sum_{\chi\in\mathcal{B}_c} |\chi\rangle\langle\chi|$ is basis-independent and any two orthonormal control bases are related by a unitary on $\mathcal{H}_c$ (e.g., the Hadamard H_c; cf. Definition 2.119), the result holds for any $\mathcal{B}_c$. This proves Corollary 4.151. □

Proposition 4.152 (Controlled splitter: parallel-path decomposition)**.** *For* $\mathbf{U}_{\mathrm{ctrl}} = |\mathbf{0}\rangle\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + |\mathbf{1}\rangle\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$ *(cf. eq. (2.75)), the only nonzero RR–matrix elements in* $\mathcal{B}_c^{(Z)}$ *are*

$$\mathsf{W}_{\mathbf{0}\leftarrow\mathbf{0}} = \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}, \qquad \mathsf{W}_{\mathbf{1}\leftarrow\mathbf{1}} = \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}, \qquad \mathsf{W}_{\mathbf{0}\leftarrow\mathbf{1}} = \mathsf{W}_{\mathbf{1}\leftarrow\mathbf{0}} = 0. \tag{4.49}$$

Proof. Work in the computational control basis $\mathcal{B}_c^{(Z)} = \{|\mathbf{0}\rangle, |\mathbf{1}\rangle\} \subset \mathcal{H}_c$. By the definition of RR–matrix elements (eq. (4.45)), for any $\chi_{\mathrm{out}}, \chi_{\mathrm{in}} \in \mathcal{B}_c^{(Z)}$,

$$\mathsf{W}_{\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}} = \Big(\langle\chi_{\mathrm{out}}|_c \otimes I\Big)\ \mathbf{U}_{\mathrm{ctrl}}\ \Big(|\chi_{\mathrm{in}}\rangle_c \otimes I\Big),$$

with $\mathbf{U}_{\mathrm{ctrl}}$ given by eq. (2.75): $\mathbf{U}_{\mathrm{ctrl}} = |\mathbf{0}\rangle\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + |\mathbf{1}\rangle\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$.

Diagonal entries. For $\chi_{\mathrm{out}} = \chi_{\mathrm{in}} = \mathbf{0}$,

$$\begin{aligned}\mathsf{W}_{\mathbf{0}\leftarrow\mathbf{0}} &= \Big(\langle\mathbf{0}|_c \otimes I\Big)\Big[\ |\mathbf{0}\rangle\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + |\mathbf{1}\rangle\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}\ \Big]\Big(|\mathbf{0}\rangle_c \otimes I\Big) \\ &= \big(\langle\mathbf{0}|\mathbf{0}\rangle\big)\,\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + \big(\langle\mathbf{0}|\mathbf{1}\rangle\big)\,\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} = \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N},\end{aligned}$$

since $\langle\mathbf{0}|\mathbf{0}\rangle = 1$ and $\langle\mathbf{0}|\mathbf{1}\rangle = 0$. Similarly, for $\chi_{\mathrm{out}} = \chi_{\mathrm{in}} = \mathbf{1}$,

$$\begin{aligned}\mathsf{W}_{\mathbf{1}\leftarrow\mathbf{1}} &= \Big(\langle\mathbf{1}|_c \otimes I\Big)\ \mathbf{U}_{\mathrm{ctrl}}\ \Big(|\mathbf{1}\rangle_c \otimes I\Big) \\ &= \big(\langle\mathbf{1}|\mathbf{0}\rangle\big)\,\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + \big(\langle\mathbf{1}|\mathbf{1}\rangle\big)\,\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} = \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}.\end{aligned}$$

Off-diagonal entries. For $\chi_{\mathrm{out}} = \mathbf{0},\ \chi_{\mathrm{in}} = \mathbf{1}$,

$$\begin{aligned}\mathsf{W}_{\mathbf{0}\leftarrow\mathbf{1}} &= \Big(\langle\mathbf{0}|_c \otimes I\Big)\ \mathbf{U}_{\mathrm{ctrl}}\ \Big(|\mathbf{1}\rangle_c \otimes I\Big) \\ &= \big(\langle\mathbf{0}|\mathbf{0}\rangle\big)\big(\langle\mathbf{0}|\mathbf{1}\rangle\big)\,\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + \big(\langle\mathbf{0}|\mathbf{1}\rangle\big)\big(\langle\mathbf{1}|\mathbf{1}\rangle\big)\,\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} = 0,\end{aligned}$$

and analogously $\mathsf{W}_{\mathbf{1}\leftarrow\mathbf{0}} = 0$. Equivalently, using projector calculus on $\mathcal{H}_c$, $|\mathbf{a}\rangle\langle\mathbf{a}| \cdot |\mathbf{b}\rangle\langle\mathbf{b}| = \delta_{\mathbf{ab}}\,|\mathbf{a}\rangle\langle\mathbf{a}|$, so the cross terms vanish by orthogonality of $|\mathbf{0}\rangle$ and $|\mathbf{1}\rangle$.

Collecting the four cases gives exactly the identities in eq. (4.49). Hence, in $\mathcal{B}_c^{(Z)}$ the only nonzero RR–matrix elements are the two diagonal blocks, each equal to its corresponding RR path operator, which proves the parallel-path decomposition claimed. □

Proposition 4.153 (Hadamard-wrapped interferometer: sum/difference channels)**.** *Let* $\mathbf{U}_{\mathrm{Had}} = (H_c \otimes I)\,\mathbf{U}_{\mathrm{ctrl}}\,(H_c \otimes I)$ *(cf. eq. (2.78)) and define* $\hat{\mathbf{U}}_{\pm} := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} \pm \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$ *(cf. eq. (2.79)). Then in the* X*-basis* $\mathcal{B}_c^{(X)}$*,*

$$\mathsf{W}_{+\leftarrow+} = \tfrac{1}{2}\,\hat{\mathbf{U}}_{+}, \quad \mathsf{W}_{-\leftarrow+} = \tfrac{1}{2}\,\hat{\mathbf{U}}_{-}, \quad \mathsf{W}_{+\leftarrow-} = \tfrac{1}{2}\,\hat{\mathbf{U}}_{-}, \quad \mathsf{W}_{-\leftarrow-} = \tfrac{1}{2}\,\hat{\mathbf{U}}_{+}. \tag{4.50}$$

Thus measuring the control in $|\pm\rangle$ *realizes the sum/difference interferometers of Chapter 2.*

Proof. Let $\mathbf{U}_{\mathrm{Had}} = (H_c \otimes I)\,\mathbf{U}_{\mathrm{ctrl}}\,(H_c \otimes I)$ as in eq. (2.78), with $\mathbf{U}_{\mathrm{ctrl}} = |\mathbf{0}\rangle\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + |\mathbf{1}\rangle\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$ (eq. (2.75)). Write $A := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$, $B := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$, and recall $\hat{\mathbf{U}}_{\pm} := A \pm B$ (eq. (2.79)). We first compute the RR–matrix elements of $\mathbf{U}_{\mathrm{Had}}$ in the computational basis $\mathcal{B}_c^{(Z)} = \{|\mathbf{0}\rangle, |\mathbf{1}\rangle\}$, then translate the indices to the X-basis $\mathcal{B}_c^{(X)} = \{|+\rangle, |-\rangle\}$ using H_c (cf. Definition 2.119).

By definition of RR–matrix elements (eq. (4.45)), for $\mathbf{a}, \mathbf{b} \in \{\mathbf{0}, \mathbf{1}\}$,

$$\mathsf{W}_{\mathbf{a}\leftarrow\mathbf{b}} = (\langle \mathbf{a}|_c \otimes I)\,(H_c \otimes I)\,\mathbf{U}_{\mathrm{ctrl}}\,(H_c \otimes I)\,(|\mathbf{b}\rangle_c \otimes I).$$

Insert the control identity in the Z-basis on both sides of $\mathbf{U}_{\mathrm{ctrl}}$, $I_c = \sum_{i\in\{\mathbf{0},\mathbf{1}\}} |i\rangle\langle i|$, and use $\mathbf{U}_{\mathrm{ctrl}}$ being diagonal in $\mathcal{B}_c^{(Z)}$ together with Proposition 4.152 and eq. (4.49):

$$(\langle i|_c \otimes I)\,\mathbf{U}_{\mathrm{ctrl}}\,(|j\rangle_c \otimes I) \;=\; \delta_{ij} \times \begin{cases} A, & i = \mathbf{0}, \\ B, & i = \mathbf{1}. \end{cases}$$

Therefore

$$\begin{aligned} \mathsf{W}_{\mathbf{a}\leftarrow\mathbf{b}} &= \sum_{i\in\{\mathbf{0},\mathbf{1}\}} \big(\langle \mathbf{a}|H_c|i\rangle\big)\,\big(\langle i|H_c|\mathbf{b}\rangle\big) \;\times \begin{cases} A, & i = \mathbf{0}, \\ B, & i = \mathbf{1}, \end{cases} \\ &= \big(\langle \mathbf{a}|H_c|\mathbf{0}\rangle\big)\big(\langle \mathbf{0}|H_c|\mathbf{b}\rangle\big)\,A + \big(\langle \mathbf{a}|H_c|\mathbf{1}\rangle\big)\big(\langle \mathbf{1}|H_c|\mathbf{b}\rangle\big)\,B. \end{aligned}$$

Using the Hadamard overlaps in the Z-basis,

$$\langle \mathbf{0}|H_c|\mathbf{0}\rangle = \langle \mathbf{0}|H_c|\mathbf{1}\rangle = \langle \mathbf{1}|H_c|\mathbf{0}\rangle = \tfrac{1}{\sqrt{2}}, \qquad \langle \mathbf{1}|H_c|\mathbf{1}\rangle = -\tfrac{1}{\sqrt{2}},$$

one obtains the four entries:

$$\begin{aligned} \mathsf{W}_{\mathbf{0}\leftarrow\mathbf{0}} &= \tfrac{1}{2}(A+B) = \tfrac{1}{2}\,\hat{\mathbf{U}}_+, & \mathsf{W}_{\mathbf{1}\leftarrow\mathbf{0}} &= \tfrac{1}{2}(A-B) = \tfrac{1}{2}\,\hat{\mathbf{U}}_-, \\ \mathsf{W}_{\mathbf{0}\leftarrow\mathbf{1}} &= \tfrac{1}{2}(A-B) = \tfrac{1}{2}\,\hat{\mathbf{U}}_-, & \mathsf{W}_{\mathbf{1}\leftarrow\mathbf{1}} &= \tfrac{1}{2}(A+B) = \tfrac{1}{2}\,\hat{\mathbf{U}}_+. \end{aligned}$$

Finally, since H_c maps $\mathcal{B}_c^{(Z)}$ to $\mathcal{B}_c^{(X)}$ via $|+\rangle = H_c|\mathbf{0}\rangle$, $|-\rangle = H_c|\mathbf{1}\rangle$ (cf. Definition 2.119), the same block relations hold upon relabeling $\mathbf{0} \leftrightarrow +$ and $\mathbf{1} \leftrightarrow -$. Thus, in the X-basis $\mathcal{B}_c^{(X)}$,

$$\mathsf{W}_{+\leftarrow+} = \tfrac{1}{2}\,\hat{\mathbf{U}}_+, \quad \mathsf{W}_{-\leftarrow+} = \tfrac{1}{2}\,\hat{\mathbf{U}}_-, \quad \mathsf{W}_{+\leftarrow-} = \tfrac{1}{2}\,\hat{\mathbf{U}}_-, \quad \mathsf{W}_{-\leftarrow-} = \tfrac{1}{2}\,\hat{\mathbf{U}}_+,$$

which is eq. (4.50). Hence measuring the control in $|\pm\rangle$ realizes the sum/difference interferometers of Chapter 2. □

Proposition 4.154 (Equivalence of symmetric holo-interferometer constructions)**.** *Let* $\mathbf{U}_{\mathrm{sym}} := \mathbf{U}_{\mathrm{Had}}\ \mathbf{U}_{\mathrm{ctrl}}\ \mathbf{U}_{\mathrm{Had}}$. *Then* $\{\mathsf{U}^{\mathrm{sym}}_{\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}}\}$ *computed via Lemma 4.149 and Propositions 4.152 and 4.153 coincide with the RR–matrix elements of the symmetric holo-interferometer* $\mathbf{U}^{(\mathrm{sym})}_{holo}$ *as defined in Chapter 2 (cf. eq. (2.79)). Consequently,* $\mathbf{U}_{\mathrm{sym}} = \mathbf{U}^{(\mathrm{sym})}_{holo}$ *on* $\mathcal{H}_{\mathrm{Holo}}$.

Proof. Let $A := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$ and $B := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$, and set $\hat{\mathbf{U}}_\pm := A \pm B$ (cf. eq. (2.79)). We work in the X-basis $\mathcal{B}_c^{(X)} = \{|+\rangle, |-\rangle\}$ with $|\pm\rangle = \frac{1}{\sqrt{2}}(|\mathbf{0}\rangle \pm |\mathbf{1}\rangle)$ (cf. Definition 2.119). By definition, $\mathbf{U}_{\mathrm{sym}} := \mathbf{U}_{\mathrm{Had}}\ \mathbf{U}_{\mathrm{ctrl}}\ \mathbf{U}_{\mathrm{Had}}$, where $\mathbf{U}_{\mathrm{Had}} = (H_c \otimes I)\,\mathbf{U}_{\mathrm{ctrl}}\,(H_c \otimes I)$ (eq. (2.78)) and $\mathbf{U}_{\mathrm{ctrl}} = |\mathbf{0}\rangle\langle\mathbf{0}|_c \otimes A + |\mathbf{1}\rangle\langle\mathbf{1}|_c \otimes B$ (eq. (2.75)).

Step 1: Block form of the factors in $\mathcal{B}_c^{(X)}$. From Proposition 4.153 and eq. (4.50) we already have, in the X-basis,

$$(\mathbf{U}_{\mathrm{Had}})_{+\leftarrow+} = \tfrac{1}{2}\,\hat{\mathbf{U}}_+, \quad (\mathbf{U}_{\mathrm{Had}})_{-\leftarrow+} = \tfrac{1}{2}\,\hat{\mathbf{U}}_-, \quad (\mathbf{U}_{\mathrm{Had}})_{+\leftarrow-} = \tfrac{1}{2}\,\hat{\mathbf{U}}_-, \quad (\mathbf{U}_{\mathrm{Had}})_{-\leftarrow-} = \tfrac{1}{2}\,\hat{\mathbf{U}}_+. \tag{4.51}$$

A direct change-of-basis computation (insert $I_c = |\mathbf{0}\rangle\langle\mathbf{0}| + |\mathbf{1}\rangle\langle\mathbf{1}|$ and use $|\pm\rangle = \frac{1}{\sqrt{2}}(|\mathbf{0}\rangle \pm |\mathbf{1}\rangle)$) gives the same block pattern for $\mathbf{U}_{\mathrm{ctrl}}$ in $\mathcal{B}_c^{(X)}$:

$$(\mathbf{U}_{\mathrm{ctrl}})_{+\leftarrow+} = \tfrac{1}{2}\,\hat{\mathbf{U}}_+, \quad (\mathbf{U}_{\mathrm{ctrl}})_{-\leftarrow+} = \tfrac{1}{2}\,\hat{\mathbf{U}}_-, \quad (\mathbf{U}_{\mathrm{ctrl}})_{+\leftarrow-} = \tfrac{1}{2}\,\hat{\mathbf{U}}_-, \quad (\mathbf{U}_{\mathrm{ctrl}})_{-\leftarrow-} = \tfrac{1}{2}\,\hat{\mathbf{U}}_+, \tag{4.52}$$

which is also consistent with Proposition 4.152 and eq. (4.49) and the basis change implemented by H_c (cf. Definition 2.119).

Equations (4.51)–(4.52) can be summarized as the identical 2×2 block matrix (with operator entries on $\mathcal{H}_{\mathrm{RR}}^{(2n)}$):

$$\mathbf{B} := \frac{1}{2}\begin{pmatrix} \hat{\mathbf{U}}_+ & \hat{\mathbf{U}}_- \\ \hat{\mathbf{U}}_- & \hat{\mathbf{U}}_+ \end{pmatrix}, \qquad \text{so that} \quad \mathbf{U}_{\mathrm{Had}} \leftrightarrow \mathbf{B}, \quad \mathbf{U}_{\mathrm{ctrl}} \leftrightarrow \mathbf{B} \quad \text{in } \mathcal{B}_c^{(X)}. \tag{4.53}$$

Step 2: Convolution (block) multiplication. By the control-resolved convolution Lemma 4.149 and eq. (4.47) (associativity by iterated insertion of I_c), the RR–matrix of the product $\mathbf{U}_{\mathrm{sym}} = \mathbf{U}_{\mathrm{Had}}\mathbf{U}_{\mathrm{ctrl}}\mathbf{U}_{\mathrm{Had}}$ in the X-basis is the block product

$$\mathbf{U}_{\mathrm{sym}} \leftrightarrow \mathbf{B}\cdot\mathbf{B}\cdot\mathbf{B} = \mathbf{B}^3.$$

A straightforward induction on m shows the operator identity

$$\mathbf{B}^m = \frac{1}{2}\begin{pmatrix} A^m + B^m & A^m - B^m \\ A^m - B^m & A^m + B^m \end{pmatrix}, \qquad m \in \mathbb{N}, \tag{4.54}$$

with the $m = 1$ case given by (4.53) and the step $m \to m+1$ following from block multiplication and the ring identities $(A \pm B)(A^m \pm B^m) = A^{m+1} \pm B^{m+1}$ and $(A \pm B)(A^m \mp B^m) = A^{m+1} \mp B^{m+1}$. Specializing (4.54) to $m = 3$ yields

$$\mathbf{U}_{\mathrm{sym}} \leftrightarrow \frac{1}{2}\begin{pmatrix} A^3 + B^3 & A^3 - B^3 \\ A^3 - B^3 & A^3 + B^3 \end{pmatrix}. \tag{4.55}$$

Step 3: Identification with the Chapter 2 symmetric device. By the Chapter 2 parallel-path semantics *eq.* (2.79) (applied to the three-stage symmetric wrapper), the RR–matrix elements of the symmetric holo-interferometer $\mathbf{U}_{\mathrm{holo}}^{(\mathrm{sym})}$ in the X-basis are precisely

$$\mathsf{U}^{(\mathrm{sym})}_{+\leftarrow+} = \tfrac{1}{2}(A^3 + B^3), \quad \mathsf{U}^{(\mathrm{sym})}_{-\leftarrow+} = \tfrac{1}{2}(A^3 - B^3), \quad \mathsf{U}^{(\mathrm{sym})}_{+\leftarrow-} = \tfrac{1}{2}(A^3 - B^3), \quad \mathsf{U}^{(\mathrm{sym})}_{-\leftarrow-} = \tfrac{1}{2}(A^3 + B^3),$$

matching entrywise the block matrix in (4.55). Therefore the families of RR–matrix elements coincide: $\{\mathsf{U}^{\mathrm{sym}}_{\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}}\} = \{\mathsf{U}^{(\mathrm{sym})}_{\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}}\}$. Since equality of all RR–matrix elements in an orthonormal control basis implies equality of the underlying operators on $\mathcal{H}_{\mathrm{Holo}} = \mathcal{H}_c \otimes \mathcal{H}_{\mathrm{RR}}^{(2n)}$ (by reconstruction from partial matrix elements; cf. eq. (4.45)), we conclude $\mathbf{U}_{\mathrm{sym}} = \mathbf{U}_{\mathrm{holo}}^{(\mathrm{sym})}$, as claimed. □

Remark 4.155 (Why Proposition 4.154 is not tautological)**.** Chapter 2 specifies the symmetric holo-interferometer $\mathbf{U}_{\mathrm{holo}}^{(\mathrm{sym})}$ by its RR–matrix behavior in the X-basis—namely, the sum/difference channels governed by $\hat{\mathbf{U}}_\pm$ (cf. eq. (2.79)). In contrast, the word $\mathbf{U}_{\mathrm{sym}} := \mathbf{U}_{\mathrm{Had}}\mathbf{U}_{\mathrm{ctrl}}\mathbf{U}_{\mathrm{Had}}$

is an implementation built from letters in $\mathsf{A}_{\mathrm{holo}}$. The proof of Proposition 4.154 uses only the RR–extraction eq. (4.45), the block forms of $\mathbf{U}_{\mathrm{ctrl}}$ and $\mathbf{U}_{\mathrm{Had}}$ (Propositions 4.152 and 4.153), and the control-resolved convolution (Lemma 4.149) to show that the two RR–matrix families coincide entrywise. Thus the proposition asserts *equivalence of specification and implementation*. It would become tautological *only* if $\mathbf{U}_{\mathrm{holo}}^{(\mathrm{sym})}$ were *defined* to be the same three-gate word; but under the behavioral definition via $\hat{\mathbf{U}}_{\pm}$, the equivalence is a contentful statement.

Proposition 4.156 (Uniqueness from RR–specification: reconstruction of the symmetric device)**.** *Let* $A := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$ *and* $B := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$ *be the two RR branch operators and set* $\hat{\mathbf{U}}_{\pm} := A \pm B$ *(cf. eq. (2.79)). Suppose* $\mathbf{U}$ *is any device on* $\mathcal{H}_{\mathrm{Holo}}$ *whose RR–matrix elements in the* X*-basis* $\mathcal{B}_c^{(X)} = \{|+\rangle, |-\rangle\}$ *satisfy*

$$\mathbf{U} \;\leftrightarrow\; \begin{bmatrix} \mathsf{U}_{+\leftarrow+} & \mathsf{U}_{+\leftarrow-} \\ \mathsf{U}_{-\leftarrow+} & \mathsf{U}_{-\leftarrow-} \end{bmatrix} = \frac{1}{2}\begin{bmatrix} \hat{\mathbf{U}}_{+} & \hat{\mathbf{U}}_{-} \\ \hat{\mathbf{U}}_{-} & \hat{\mathbf{U}}_{+} \end{bmatrix}. \tag{4.56}$$

Then $\mathbf{U} = (H_c \otimes I)\,\mathbf{U}_{\mathrm{ctrl}}\,(H_c \otimes I)$*, where* $\mathbf{U}_{\mathrm{ctrl}} = |\mathbf{0}\rangle\langle\mathbf{0}|_c \otimes A + |\mathbf{1}\rangle\langle\mathbf{1}|_c \otimes B$ *(cf. eq. (2.75)). In particular,* $\mathbf{U} = \mathbf{U}_{holo}^{(\mathrm{sym})}$ *as defined behaviorally in Chapter 2 (cf. eq. (2.79)).*

Proof. By hypothesis (4.56), in the X-basis the control-block matrix of $\mathbf{U}$ is

$$\mathbf{B} \;:=\; \frac{1}{2}\begin{bmatrix} \hat{\mathbf{U}}_{+} & \hat{\mathbf{U}}_{-} \\ \hat{\mathbf{U}}_{-} & \hat{\mathbf{U}}_{+} \end{bmatrix}.$$

Define operator-valued blocks (on $\mathcal{H}_{\mathrm{RR}}^{(2n)}$) in this basis by $B_{++} := \mathsf{U}_{+\leftarrow+} = \frac{1}{2}\hat{\mathbf{U}}_{+}$ and $B_{+-} := \mathsf{U}_{+\leftarrow-} = \frac{1}{2}\hat{\mathbf{U}}_{-}$. Then we can *reconstruct the branches* A, B from the RR–spec alone:

$$A \;=\; B_{++} + B_{+-} \;=\; \tfrac{1}{2}(\hat{\mathbf{U}}_{+} + \hat{\mathbf{U}}_{-}), \qquad B \;=\; B_{++} - B_{+-} \;=\; \tfrac{1}{2}(\hat{\mathbf{U}}_{+} - \hat{\mathbf{U}}_{-}). \tag{4.57}$$

Let $\mathbf{D}$ denote the Z-basis diagonal block of the controlled splitter, $\mathbf{D} := \mathrm{diag}(A, B)$, so that $\mathbf{U}_{\mathrm{ctrl}} \leftrightarrow \mathbf{D}$ in $\mathcal{B}_c^{(Z)} = \{|\mathbf{0}\rangle, |\mathbf{1}\rangle\}$ (cf. Proposition 4.152 and eq. (4.49)). Changing basis by the Hadamard H_c (cf. Definition 2.119), the conjugation $(H_c \otimes I)\,\mathbf{U}_{\mathrm{ctrl}}\,(H_c \otimes I)$ has X-basis block matrix

$$\mathbf{B}' \;=\; \frac{1}{2}\begin{bmatrix} A{+}B & A{-}B \\ A{-}B & A{+}B \end{bmatrix} \overset{(4.57)}{=} \frac{1}{2}\begin{bmatrix} \hat{\mathbf{U}}_{+} & \hat{\mathbf{U}}_{-} \\ \hat{\mathbf{U}}_{-} & \hat{\mathbf{U}}_{+} \end{bmatrix} = \mathbf{B}.$$

Hence $\mathbf{U}$ and $(H_c \otimes I)\,\mathbf{U}_{\mathrm{ctrl}}\,(H_c \otimes I)$ have identical RR–matrix families in an orthonormal control basis. By the reconstruction property of partial matrix elements (cf. eq. (4.45)), equality of all control blocks implies equality of the operators on $\mathcal{H}_{\mathrm{Holo}}$. Therefore $\mathbf{U} = (H_c \otimes I)\,\mathbf{U}_{\mathrm{ctrl}}\,(H_c \otimes I)$. □

Remark 4.157 (What Proposition 4.156 does)**.** Proposition 4.156 is the *converse* to Proposition 4.154. While Proposition 4.154 verifies that the implementation word $\mathbf{U}_{\mathrm{sym}} = \mathbf{U}_{\mathrm{Had}}\mathbf{U}_{\mathrm{ctrl}}\mathbf{U}_{\mathrm{Had}}$ realizes the Chapter 2 ± RR–spec (eq. (2.79)), Proposition 4.156 says the *spec alone*—the X-basis block family $\frac{1}{2}\begin{pmatrix} \hat{\mathbf{U}}_{+} & \hat{\mathbf{U}}_{-} \\ \hat{\mathbf{U}}_{-} & \hat{\mathbf{U}}_{+} \end{pmatrix}$—already *determines* the device:

$$\mathbf{U} \;=\; (H_c \otimes I)\,\mathbf{U}_{\mathrm{ctrl}}\,(H_c \otimes I), \qquad \mathbf{U}_{\mathrm{ctrl}} = |\mathbf{0}\rangle\langle\mathbf{0}|_c \otimes A + |\mathbf{1}\rangle\langle\mathbf{1}|_c \otimes B,$$

with $A = \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$, $B = \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$ and $\hat{\mathbf{U}}_{\pm} = A \pm B$. Thus, specification $\Rightarrow$ unique implementation on $\mathcal{H}_{\mathrm{Holo}}$ (reconstruction via eq. (4.45)), making the symmetric device both *sound* (Proposition 4.154) and *complete* (Proposition 4.156) with respect to the RR–matrix semantics.

Corollary 4.158 (Perfect and partial interference, wordwise criteria)**.** *In the X-basis, perfect constructive (resp. destructive) interference in the $+$ output channel occurs iff $\hat{\mathbf{U}}_- = 0$ (resp. $\hat{\mathbf{U}}_+ = 0$). More generally,*

$$\|\mathsf{W}_{-\leftarrow+}\| \;=\; \tfrac{1}{2}\,\|\hat{\mathbf{U}}_-\|, \qquad \|\mathsf{W}_{+\leftarrow+}\| \;=\; \tfrac{1}{2}\,\|\hat{\mathbf{U}}_+\|, \tag{4.58}$$

with the same relations for the $-$ input by symmetry.

Proof. Work in the X-basis $\mathcal{B}_c^{(X)} = \{|+\rangle, |-\rangle\}$. By the Hadamard-wrapped interferometer block formulas (Proposition 4.153 and eq. (4.50)) we have, for $+$ input,

$$\mathsf{W}_{+\leftarrow+} = \tfrac{1}{2}\,\hat{\mathbf{U}}_+, \qquad \mathsf{W}_{-\leftarrow+} = \tfrac{1}{2}\,\hat{\mathbf{U}}_-,$$

where $\hat{\mathbf{U}}_{\pm} := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} \pm \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$ (cf. eq. (2.79)).

Perfect constructive/destructive interference in the $+$ output. In the wordwise semantics, "perfect constructive in the $+$ output channel for $+$ input" means that the $-$ output branch vanishes, i.e. $\mathsf{W}_{-\leftarrow+} = 0$; similarly, "perfect destructive in the $+$ output" means $\mathsf{W}_{+\leftarrow+} = 0$. Using the identities above, these conditions are equivalent to

$$\mathsf{W}_{-\leftarrow+} = 0 \iff \hat{\mathbf{U}}_- = 0, \qquad \mathsf{W}_{+\leftarrow+} = 0 \iff \hat{\mathbf{U}}_+ = 0,$$

which proves the iff statements.

Norm relations. Let $\|\cdot\|$ denote the operator norm on $\mathcal{B}(\mathcal{H}_{\mathrm{RR}}^{(2n)})$ (any absolutely homogeneous norm would suffice). By absolute homogeneity and the block identities,

$$\left\|\mathsf{W}_{-\leftarrow+}\right\| = \left\|\tfrac{1}{2}\,\hat{\mathbf{U}}_-\right\| = \tfrac{1}{2}\,\|\hat{\mathbf{U}}_-\|, \qquad \left\|\mathsf{W}_{+\leftarrow+}\right\| = \left\|\tfrac{1}{2}\,\hat{\mathbf{U}}_+\right\| = \tfrac{1}{2}\,\|\hat{\mathbf{U}}_+\|,$$

which is eq. (4.58).

Symmetry for $-$ input. The remaining relations follow by the same calculation for $-$ input, using again $\mathsf{W}_{+\leftarrow-} = \frac{1}{2}\,\hat{\mathbf{U}}_-$ and $\mathsf{W}_{-\leftarrow-} = \frac{1}{2}\,\hat{\mathbf{U}}_+$ from Proposition 4.153 and eq. (4.50). □

Lemma 4.159 (Asymmetric SU(2) normal form)**.** *Let $A := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$ and $B := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$, and set $\mathbf{U}_{\mathrm{ctrl}} = |\mathbf{0}\rangle\langle\mathbf{0}|_c \otimes A + |\mathbf{1}\rangle\langle\mathbf{1}|_c \otimes B$ (cf. eqs. (2.75) and (4.49) and Proposition 4.152). For any control couplers $U_c^{(L)}, U_c^{(R)} \in U(\mathcal{H}_c)$ define the device word*

$$\mathbf{U}_{\mathrm{asym}} \;:=\; (U_c^{(R)} \otimes I)\;\mathbf{U}_{\mathrm{ctrl}}\;(U_c^{(L)} \otimes I).$$

Then for any orthonormal control basis $\mathcal{B}_c \subset \mathcal{H}_c$ and $\xi, \eta \in \mathcal{B}_c$,

$$\mathsf{W}_{\xi\leftarrow\eta}^{(\mathrm{asym})} \;=\; \langle\xi|U_c^{(R)}|\mathbf{0}\rangle\,\langle\mathbf{0}|U_c^{(L)}|\eta\rangle\,A \;+\; \langle\xi|U_c^{(R)}|\mathbf{1}\rangle\,\langle\mathbf{1}|U_c^{(L)}|\eta\rangle\,B, \tag{4.59}$$

where $\mathsf{W}_{\xi\leftarrow\eta}^{(\mathrm{asym})}$ denotes the RR–matrix element of $\mathbf{U}_{\mathrm{asym}}$ (cf. Definition 4.148 and eq. (4.45)).

Proof. By definition of RR–matrix elements (eq. (4.45)),

$$\mathsf{W}^{(\mathrm{asym})}_{\xi\leftarrow\eta} = (\langle\xi|_c\otimes I)\,(U_c^{(R)}\otimes I)\,\mathbf{U}_{\mathrm{ctrl}}\,(U_c^{(L)}\otimes I)\,(|\eta\rangle_c\otimes I).$$

Insert the Z-basis resolution of the control identity on both sides of $\mathbf{U}_{\mathrm{ctrl}}$, $I_c = |\mathbf{0}\rangle\langle\mathbf{0}| + |\mathbf{1}\rangle\langle\mathbf{1}|$, and use that $\mathbf{U}_{\mathrm{ctrl}}$ is diagonal in this basis with blocks A, B (Proposition 4.152 and eq. (4.49)). This yields

$$\mathsf{W}^{(\mathrm{asym})}_{\xi\leftarrow\eta} = \langle\xi|U_c^{(R)}|\mathbf{0}\rangle\,\langle\mathbf{0}|U_c^{(L)}|\eta\rangle\,A + \langle\xi|U_c^{(R)}|\mathbf{1}\rangle\,\langle\mathbf{1}|U_c^{(L)}|\eta\rangle\,B,$$

which is eq. (4.59). No properties of A, B beyond their action on $\mathcal{H}_{\mathrm{RR}}^{(2n)}$ are used. □

Remark 4.160 (Specializations and consequences)**.** *(i)* Symmetric interferometer: $U_c^{(L)} = U_c^{(R)} = H_c$ gives the sum/difference blocks $\frac{1}{2}(A \pm B) = \frac{1}{2}\,\hat{\mathbf{U}}_{\pm}$ (Proposition 4.153). *(ii)* Single output coupler: $U_c^{(L)} = I$, $U_c^{(R)} = H_c$ yields $\frac{1}{\sqrt{2}}$ times a single branch in each block (Corollary 4.162). *(iii)* Single input coupler: $U_c^{(L)} = H_c$, $U_c^{(R)} = I$ yields the corrected $\frac{1}{\sqrt{2}}$ pure-branch blocks (Corollary 4.163). *(iv)* General SU(2) phases: arbitrary $U_c^{(L/R)}$ give coefficients $\alpha_{\xi\eta}, \beta_{\xi\eta}$ with the mixing bounds following from block unitarity (Corollary 4.151). *(v)* Equality with Chapter 2: choosing $U_c^{(L/R)}$ to match the Chapter 2 construction gives identical RR blocks and hence identical devices (Proposition 4.165).

Proposition 4.161 (Asymmetric holo-interferometer: general SU(2) couplers)**.** *Let $U_c^{(L)}, U_c^{(R)} \in \mathrm{U}(\mathcal{H}_c)$ be arbitrary control unitaries and define the asymmetric device word*

$$\mathbf{U}_{\mathrm{asym}} \;:=\; (U_c^{(R)}\otimes I)\;\mathbf{U}_{\mathrm{ctrl}}\;(U_c^{(L)}\otimes I),$$

with $\mathbf{U}_{\mathrm{ctrl}}$ as in eq. (2.75) and RR branch operators $A := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$, $B := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$. For any orthonormal control basis $\mathcal{B}_c$ and $\xi, \eta \in \mathcal{B}_c$,

$$\mathsf{W}^{(\mathrm{asym})}_{\xi\leftarrow\eta} \;=\; \Big\langle\xi\Big|U_c^{(R)}\Big|\mathbf{0}\Big\rangle\,\Big\langle\mathbf{0}\Big|U_c^{(L)}\Big|\eta\Big\rangle\,A \;+\; \Big\langle\xi\Big|U_c^{(R)}\Big|\mathbf{1}\Big\rangle\,\Big\langle\mathbf{1}\Big|U_c^{(L)}\Big|\eta\Big\rangle\,B. \tag{4.60}$$

Proof. This is an immediate consequence of Lemma 4.159. Indeed, with $A := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$, $B := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$, and $\mathbf{U}_{\mathrm{asym}} = (U_c^{(R)}\otimes I)\,\mathbf{U}_{\mathrm{ctrl}}\,(U_c^{(L)}\otimes I)$, Lemma 4.159 gives, for any orthonormal control basis $\mathcal{B}_c$ and $\xi, \eta \in \mathcal{B}_c$,

$$\mathsf{W}^{(\mathrm{asym})}_{\xi\leftarrow\eta} = \langle\xi|U_c^{(R)}|\mathbf{0}\rangle\,\langle\mathbf{0}|U_c^{(L)}|\eta\rangle\,A + \langle\xi|U_c^{(R)}|\mathbf{1}\rangle\,\langle\mathbf{1}|U_c^{(L)}|\eta\rangle\,B,$$

which is exactly the claimed block formula eq. (4.60). Equivalently, one may derive the same expression directly from the RR–extraction eq. (4.45) by inserting $I_c = |\mathbf{0}\rangle\langle\mathbf{0}| + |\mathbf{1}\rangle\langle\mathbf{1}|$ and using the diagonal form of $\mathbf{U}_{\mathrm{ctrl}}$ in the Z-basis (Proposition 4.152 and eq. (4.49)), but Lemma 4.159 already encapsulates this calculation. □

Corollary 4.162 (Single output coupler)**.** *Let $U_c^{(R)} = H_c$, $U_c^{(L)} = I$. In the computational basis $\mathcal{B}_c^{(Z)} = \{|\mathbf{0}\rangle, |\mathbf{1}\rangle\}$,*

$$\begin{aligned} \mathsf{W}^{(\mathrm{asym})}_{\mathbf{0}\leftarrow\mathbf{0}} &= \tfrac{1}{\sqrt{2}}A, \quad \mathsf{W}^{(\mathrm{asym})}_{\mathbf{1}\leftarrow\mathbf{0}} = \tfrac{1}{\sqrt{2}}A, \\ \mathsf{W}^{(\mathrm{asym})}_{\mathbf{0}\leftarrow\mathbf{1}} &= \tfrac{1}{\sqrt{2}}B, \quad \mathsf{W}^{(\mathrm{asym})}_{\mathbf{1}\leftarrow\mathbf{1}} = -\tfrac{1}{\sqrt{2}}B. \end{aligned} \tag{4.61}$$

Thus a single *readout coupler mixes each branch amplitude into two output ports without creating $A \pm B$ interference inside a single block (interference appears when reading in the X-basis).*

Proof. Set $A := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$ and $B := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$. Consider the asymmetric device $\mathbf{U}_{\mathrm{asym}} = (U_c^{(R)} \otimes I)\, \mathbf{U}_{\mathrm{ctrl}}\, (U_c^{(L)} \otimes I)$ with $U_c^{(R)} = H_c$, $U_c^{(L)} = I$ and $\mathbf{U}_{\mathrm{ctrl}} = |\mathbf{0}\rangle\langle\mathbf{0}|_c \otimes A + |\mathbf{1}\rangle\langle\mathbf{1}|_c \otimes B$ (cf. eq. (2.75)). By the general asymmetric block formula Proposition 4.161 and eq. (4.60), for any $\xi, \eta \in \mathcal{B}_c^{(Z)} = \{|\mathbf{0}\rangle, |\mathbf{1}\rangle\}$,

$$\mathsf{W}_{\xi \leftarrow \eta}^{(\mathrm{asym})} \;=\; \underbrace{\langle \xi | H_c | \mathbf{0} \rangle}_{\text{output coupler}} \; \underbrace{\langle \mathbf{0} | I | \eta \rangle}_{=\,\delta_{\eta,\mathbf{0}}} A \;+\; \underbrace{\langle \xi | H_c | \mathbf{1} \rangle}_{\text{output coupler}} \; \underbrace{\langle \mathbf{1} | I | \eta \rangle}_{=\,\delta_{\eta,\mathbf{1}}} B.$$

Using the standard Hadamard overlaps in the computational basis (cf. Definition 2.119), $\langle \mathbf{0}|H_c|\mathbf{0}\rangle = \langle \mathbf{1}|H_c|\mathbf{0}\rangle = \langle \mathbf{0}|H_c|\mathbf{1}\rangle = \frac{1}{\sqrt{2}}$, $\langle \mathbf{1}|H_c|\mathbf{1}\rangle = -\frac{1}{\sqrt{2}}$, we evaluate the four entries:

(i) Input $\eta = \mathbf{0}$:

$$\mathsf{W}_{\mathbf{0}\leftarrow\mathbf{0}}^{(\mathrm{asym})} = \langle \mathbf{0}|H_c|\mathbf{0}\rangle\, A = \tfrac{1}{\sqrt{2}} A, \qquad \mathsf{W}_{\mathbf{1}\leftarrow\mathbf{0}}^{(\mathrm{asym})} = \langle \mathbf{1}|H_c|\mathbf{0}\rangle\, A = \tfrac{1}{\sqrt{2}} A.$$

(ii) Input $\eta = \mathbf{1}$:

$$\mathsf{W}_{\mathbf{0}\leftarrow\mathbf{1}}^{(\mathrm{asym})} = \langle \mathbf{0}|H_c|\mathbf{1}\rangle\, B = \tfrac{1}{\sqrt{2}} B, \qquad \mathsf{W}_{\mathbf{1}\leftarrow\mathbf{1}}^{(\mathrm{asym})} = \langle \mathbf{1}|H_c|\mathbf{1}\rangle\, B = -\tfrac{1}{\sqrt{2}} B.$$

These are exactly the relations in eq. (4.61). Each RR–block contains a single branch operator (either A or B) scaled by the coupler coefficient; hence no $A \pm B$ interference appears within an individual block in the Z-readout. Interference emerges when reading in the X-basis, where the blocks become sums/differences as in Proposition 4.153 and eq. (4.50). □

Corollary 4.163 (Single input coupler)**.** *Let $U_c^{(R)} = I$, $U_c^{(L)} = H_c$. In the X-basis $\mathcal{B}_c^{(X)} = \{|+\rangle, |-\rangle\}$,*

$$\mathsf{W}_{+\leftarrow+}^{(\mathrm{asym})} = \tfrac{1}{\sqrt{2}} A, \quad \mathsf{W}_{-\leftarrow+}^{(\mathrm{asym})} = \tfrac{1}{\sqrt{2}} A, \quad \mathsf{W}_{+\leftarrow-}^{(\mathrm{asym})} = \tfrac{1}{\sqrt{2}} B, \quad \mathsf{W}_{-\leftarrow-}^{(\mathrm{asym})} = -\tfrac{1}{\sqrt{2}} B, \tag{4.62}$$

where $A := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$, $B := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$. These follow directly from the general asymmetric block formula Proposition 4.161 and eq. (4.60).

Proof. Set $A := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$ and $B := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$. Consider the asymmetric device $\mathbf{U}_{\mathrm{asym}} = (U_c^{(R)} \otimes I)\, \mathbf{U}_{\mathrm{ctrl}}\, (U_c^{(L)} \otimes I)$ with $U_c^{(R)} = I$, $U_c^{(L)} = H_c$ and $\mathbf{U}_{\mathrm{ctrl}} = |\mathbf{0}\rangle\langle\mathbf{0}|_c \otimes A + |\mathbf{1}\rangle\langle\mathbf{1}|_c \otimes B$ (cf. eq. (2.75)). By the general asymmetric block formula Proposition 4.161 and eq. (4.60), for any control basis states $\xi, \eta \in \mathcal{B}_c$,

$$\mathsf{W}_{\xi \leftarrow \eta}^{(\mathrm{asym})} \;=\; \langle \xi | I | \mathbf{0} \rangle\, \langle \mathbf{0} | H_c | \eta \rangle\, A \;+\; \langle \xi | I | \mathbf{1} \rangle\, \langle \mathbf{1} | H_c | \eta \rangle\, B. \tag{4.63}$$

Now choose the X-basis $\mathcal{B}_c^{(X)} = \{|+\rangle, |-\rangle\}$ with $|\pm\rangle = \frac{1}{\sqrt{2}}(|\mathbf{0}\rangle \pm |\mathbf{1}\rangle)$ (cf. Definition 2.119). Using the Hadamard overlaps

$$\langle \mathbf{0}|H_c|+\rangle = 1, \quad \langle \mathbf{1}|H_c|+\rangle = 0, \qquad \langle \mathbf{0}|H_c|-\rangle = 0, \quad \langle \mathbf{1}|H_c|-\rangle = 1,$$

and the X-to-Z overlaps

$$\langle +|\mathbf{0}\rangle = \tfrac{1}{\sqrt{2}}, \quad \langle +|\mathbf{1}\rangle = \tfrac{1}{\sqrt{2}}, \qquad \langle -|\mathbf{0}\rangle = \tfrac{1}{\sqrt{2}}, \quad \langle -|\mathbf{1}\rangle = -\tfrac{1}{\sqrt{2}},$$

we evaluate (4.63) entrywise.

(i) + input ($\eta = +$). Then $\langle \mathbf{0}|H_c|+\rangle = 1$, $\langle \mathbf{1}|H_c|+\rangle = 0$, so

$$\mathsf{W}^{(\mathrm{asym})}_{+\leftarrow+} = \langle +|\mathbf{0}\rangle\, A = \tfrac{1}{\sqrt{2}} A, \qquad \mathsf{W}^{(\mathrm{asym})}_{-\leftarrow+} = \langle -|\mathbf{0}\rangle\, A = \tfrac{1}{\sqrt{2}} A.$$

(ii) − input ($\eta = -$). Then $\langle \mathbf{0}|H_c|-\rangle = 0$, $\langle \mathbf{1}|H_c|-\rangle = 1$, so

$$\mathsf{W}^{(\mathrm{asym})}_{+\leftarrow-} = \langle +|\mathbf{1}\rangle\, B = \tfrac{1}{\sqrt{2}} B, \qquad \mathsf{W}^{(\mathrm{asym})}_{-\leftarrow-} = \langle -|\mathbf{1}\rangle\, B = -\tfrac{1}{\sqrt{2}} B.$$

Thus, with a *single input coupler* $U_c^{(L)} = H_c$ and readout in the X-basis, the RR–matrix elements are

$$\mathsf{W}^{(\mathrm{asym})}_{+\leftarrow+} = \tfrac{1}{\sqrt{2}} A, \quad \mathsf{W}^{(\mathrm{asym})}_{-\leftarrow+} = \tfrac{1}{\sqrt{2}} A, \quad \mathsf{W}^{(\mathrm{asym})}_{+\leftarrow-} = \tfrac{1}{\sqrt{2}} B, \quad \mathsf{W}^{(\mathrm{asym})}_{-\leftarrow-} = -\tfrac{1}{\sqrt{2}} B.$$

Each block contains a *single* branch operator (either A or B) scaled by the appropriate coupler overlap; no $A \pm B$ combination appears within a single block.

Consistency note. The sum/difference blocks $\frac{1}{2}(A \pm B) = \frac{1}{2}\hat{\mathbf{U}}_{\pm}$ of eq. (2.79) arise when the device has *both* input and output Hadamard couplers (the symmetric wrapper of Proposition 4.153) or, equivalently, when one expresses $\mathbf{U}_{\mathrm{ctrl}}$ itself *in the X-basis on both input and output*, i.e. as the change-of-basis conjugate $(H_c \otimes I)\, \mathbf{U}_{\mathrm{ctrl}}\, (H_c \otimes I)$ (cf. Proposition 4.153). With only the input coupler $U_c^{(L)} = H_c$ and $U_c^{(R)} = I$, the X-basis blocks are as computed above. □

Remark 4.164 (Phase- and imbalance-asymmetry)**.** Taking $U_c^{(L)}$ and/or $U_c^{(R)}$ as a general SU(2) beam splitter with an internal phase (e.g. $U_c^{(L)} = R_z(\phi)\, H_c$, $U_c^{(R)} = H_c$) modifies the coefficients in eq. (4.60) so that $\mathsf{W}^{(\mathrm{asym})}_{\xi\leftarrow\eta} = \alpha_{\xi\eta}(\phi)\, A + \beta_{\xi\eta}(\phi)\, B$, thereby realizing *unequal* path mixing and controlled relative phases. All such variants still obey the block unitarity *mixing bounds* of Corollary 4.151, since $\mathbf{U}_{\mathrm{asym}}$ is unitary on $\mathcal{H}_{\mathrm{Holo}}$.

Proposition 4.165 (Equivalence with Chapter 2 asymmetric model)**.** *Let $\mathbf{U}^{(\mathrm{asym})}_{holo}$ denote the asymmetric holo-interferometer of Chapter 2 (obtained by a single coupler or a phase-imbalanced pair; cf. the constructions surrounding eqs. (2.78) and (2.79)). Choosing $U_c^{(L)}$, $U_c^{(R)}$ to match that construction, the RR–matrix elements computed from eq. (4.60) agree with those of $\mathbf{U}^{(\mathrm{asym})}_{holo}$ for all control transitions. Hence $\mathbf{U}_{\mathrm{asym}} = \mathbf{U}^{(\mathrm{asym})}_{holo}$ on $\mathcal{H}_{\mathrm{Holo}}$.*

Proof. Let $A := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$ and $B := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$, and fix control couplers $U_c^{(L)}, U_c^{(R)} \in \mathrm{U}(\mathcal{H}_c)$ so as to match the Chapter 2 asymmetric construction (single coupler or phase-imbalanced pair; cf. the discussion around eqs. (2.78) and (2.79)). Define the wordwise asymmetric device

$$\mathbf{U}_{\mathrm{asym}} \;:=\; (U_c^{(R)} \otimes I)\; \mathbf{U}_{\mathrm{ctrl}}\; (U_c^{(L)} \otimes I), \qquad \mathbf{U}_{\mathrm{ctrl}} = |\mathbf{0}\rangle\langle\mathbf{0}|_c \otimes A + |\mathbf{1}\rangle\langle\mathbf{1}|_c \otimes B \quad \text{(cf. eq. (2.75))}.$$

Step 1: RR–matrix elements of the wordwise device. By the general asymmetric block formula (Proposition 4.161 and eq. (4.60)), for any orthonormal control basis $\mathcal{B}_c$ and $\xi, \eta \in \mathcal{B}_c$,

$$\mathsf{W}^{(\mathrm{asym})}_{\xi\leftarrow\eta} = \langle \xi|U_c^{(R)}|\mathbf{0}\rangle\, \langle \mathbf{0}|U_c^{(L)}|\eta\rangle\, A \;+\; \langle \xi|U_c^{(R)}|\mathbf{1}\rangle\, \langle \mathbf{1}|U_c^{(L)}|\eta\rangle\, B. \tag{4.64}$$

Step 2: RR–matrix elements of the Chapter 2 device. Let $\mathbf{U}^{(\mathrm{asym})}_{\mathrm{holo}}$ be the asymmetric holo-interferometer of Chapter 2 with the same choice of input/output couplers $U_c^{(L)}, U_c^{(R)}$

(single Hadamard or a general SU(2) with phases; cf. eq. (2.78)). By the RR extraction rule (eq. (4.45)) and insertion of the control identity $I_c = |\mathbf{0}\rangle\langle\mathbf{0}| + |\mathbf{1}\rangle\langle\mathbf{1}|$ (the control-resolved convolution, Lemma 4.149 and eq. (4.47)), the RR–matrix elements of $\mathbf{U}^{(\mathrm{asym})}_{\mathrm{holo}}$ evaluate to

$$\mathsf{U}^{(\mathrm{asym})}_{\xi\leftarrow\eta} = \langle\xi|U_c^{(R)}|\mathbf{0}\rangle\,\langle\mathbf{0}|U_c^{(L)}|\eta\rangle\,A \;+\; \langle\xi|U_c^{(R)}|\mathbf{1}\rangle\,\langle\mathbf{1}|U_c^{(L)}|\eta\rangle\,B, \tag{4.65}$$

i.e. the same operator-valued block formula as (4.64). (This is the standard parallel-path evaluation in Chapter 2; see also the special cases worked out in Corollaries 4.162 and 4.163.)

Step 3: Equality of devices. Comparing (4.64) and (4.65) shows that, for every orthonormal control basis $\mathcal{B}_c$ and every $\xi,\eta\in\mathcal{B}_c$, $\mathsf{W}^{(\mathrm{asym})}_{\xi\leftarrow\eta} = \mathsf{U}^{(\mathrm{asym})}_{\xi\leftarrow\eta}$. By the reconstruction property of partial matrix elements (eq. (4.45)), equality of all control blocks implies equality of the underlying operators on $\mathcal{H}_{\mathrm{Holo}}$. Hence $\mathbf{U}_{\mathrm{asym}} = \mathbf{U}^{(\mathrm{asym})}_{\mathrm{holo}}$.

This establishes the claimed equivalence for arbitrary (single or paired, possibly phase-imbalanced) couplers chosen to match the Chapter 2 asymmetric construction. □

Remarks on internal RR factorization (link to generators). While the wordwise calculus treats the RR branch operators as opaque letters of $\mathsf{A}_{\mathrm{holo}}$, each such operator may internally factor through the parasymmetric blocks of Chapter 2,

$$\begin{aligned}\hat{\mathbf{U}}^{(\cdot)}_{\mathrm{RR}} &\rightsquigarrow U_L\otimes U_P\otimes U_G,\\ U_P &= \exp\Big(i\sum_{i<j} J_{ij}\left(P^{(ij)}_{\mathrm{para}} - P^{(ij)}_{\mathrm{antipara}}\right)\Big).\end{aligned}$$

consistent with the pairwise generator $\hat{F}^{(\mathrm{pair})}_{\mathrm{para}}$ and the composite evolution $U_N^{(\odot)}$ in Chapter 2. Unitarity on $\mathcal{H}_{\mathrm{Holo}}$ then enforces the isometric mixing bounds of Corollary 4.151, which will be invoked in the wordwise kinematics and mixing analysis.

Definition 4.166 (Word-dressed selectors and word-evaluated channels)**.** Let $\hat{\mathcal{F}}_\alpha(y)$ be the Fourfold selectors (Chapter 1) and let $\mathsf{W}_{\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}}$ be as in (4.45), acting on the RR factor that contains the modal fibre (via the lifted embeddings of Chapter 3; cf. Definition 2.103). Define the *word-dressed selector*

$$\hat{\mathcal{F}}^{[\mathbf{W};\,\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}]}_\alpha(y) := \mathsf{W}_{\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}}\,\hat{\mathcal{F}}_\alpha(y),$$

and the corresponding *word-evaluated channel*

$$\hat{G}^{\alpha\beta}_\Lambda\Big(x,x';\mathbf{W},\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}\Big) := \Big(\hat{\mathcal{F}}^{[\mathbf{W};\,\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}]}_\alpha(x)\Big)^\dagger\,\mathcal{U}(x\leftarrow x';\gamma)\,\hat{\mathcal{F}}_\beta(x'). \tag{4.66}$$

For $\mathbf{W}=I$ and any control states, this reduces to the undeformed $\hat{G}^{\alpha\beta}_\Lambda(x,x')$ of Definition 4.61.

Proposition 4.167 (Gauge covariance and grading preserved under holosymmetric dressing)**.** *For each* $\mathbf{W}$ *and control pair* $(\chi_{\mathrm{in}},\chi_{\mathrm{out}})$*:*

(a) Gauge covariance
$\hat{G}^{\alpha\beta}_\Lambda(x,x';\mathbf{W},\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}) \mapsto g(x)\,\hat{G}^{\alpha\beta}_\Lambda(\cdots)\,g(x')^\dagger$ *by Proposition 4.54 and Corollary 4.64, since* $\mathsf{W}_{\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}}$ *is internal to the fibre factor and conjugates compatibly.*

(b) Adjoint and degree

$\big(\hat{G}^{\alpha\beta}_{\Lambda}(\cdots)\big)^{\dagger} = \hat{G}^{\beta\alpha}_{\Lambda}(x', x; \mathbf{W}^{\dagger}, \chi_{\mathrm{in}} \leftarrow \chi_{\mathrm{out}})$ *and* deg *remains* $\deg(\alpha) + \deg(\beta)$ *(Lemmas 4.66 and 4.98 and Definition 4.94).*

(c) Functoriality

Replacing $\mathbf{W}$ *by a product* $\mathbf{W}_2\mathbf{W}_1$ *composes the dressing*

$$\mathsf{W}_{2,\chi_2\leftarrow\chi_1}\,\mathsf{W}_{1,\chi_1\leftarrow\chi_0}.$$

Geometric concatenation still composes via Proposition 4.52.

Proof. Fix a path $\gamma : x' \to x$, indices $\alpha, \beta \in \mathcal{I}$, and a device word $\mathbf{W} \in \langle \mathsf{A}_{\mathrm{holo}} \rangle$ with control transition $(\chi_{\mathrm{in}}, \chi_{\mathrm{out}})$. By the wordwise dressing of the Chapter 4 Fourfold channel, define

$$\hat{G}^{\alpha\beta}_{\Lambda}\big(x, x'; \mathbf{W}, \chi_{\mathrm{out}} \leftarrow \chi_{\mathrm{in}}\big) := \hat{\mathcal{F}}^{\dagger}_{\alpha}(x)\,\mathcal{U}(x \leftarrow x'; \gamma)\,\mathsf{W}_{\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}}\,\hat{\mathcal{F}}_{\beta}(x'). \tag{4.67}$$

Here, $\mathcal{U}$ is the parallel transporter and $\hat{\mathcal{F}}_{\alpha}$ are the Fourfold selectors (Definition 4.72); $\mathsf{W}_{\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}}$ is the RR–matrix element of $\mathbf{W}$ (Definition 4.148 and eq. (4.45)).

(a) Gauge covariance. Under a gauge change $g(\cdot)$ on the bundle, the factors transform by endpoint conjugation

$$\hat{\mathcal{F}}_{\alpha}(x) \mapsto g(x)\hat{\mathcal{F}}_{\alpha}(x), \qquad \mathcal{U}(x \leftarrow x'; \gamma) \mapsto g(x)\,\mathcal{U}(x \leftarrow x'; \gamma)\,g(x')^{\dagger} \quad (\textit{Proposition 4.54}),$$

and the assembled Fourfold channel obeys endpoint conjugation (*Corollary 4.64*). The RR–matrix element $\mathsf{W}_{\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}}$ acts on the RR fibre and is *gauge–neutral* with respect to the base change on $\mathscr{H}_{(0,1)}$. Thus, applying the transformations inside (4.67) and canceling $g(x)^{\dagger}g(x) = I$ and $g(x')^{\dagger}$ across the neutral RR factor, we obtain

$$\hat{G}^{\alpha\beta}_{\Lambda}(\cdots) \;\longmapsto\; g(x)\;\hat{G}^{\alpha\beta}_{\Lambda}(\cdots)\;g(x')^{\dagger},$$

which is the asserted covariance.

(b) Adjoint and degree. Taking adjoints in (4.67) and using $\mathcal{U}(x \leftarrow x'; \gamma)^{\dagger} = \mathcal{U}(x' \leftarrow x; \gamma^{-1})$ and $(\mathsf{W}_{\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}})^{\dagger} = \mathsf{W}^{\dagger}_{\chi_{\mathrm{in}}\leftarrow\chi_{\mathrm{out}}}$ (from *eq.* (4.45) and the adjoint rule), we get

$$\big(\hat{G}^{\alpha\beta}_{\Lambda}(x, x'; \mathbf{W}, \chi_{\mathrm{out}} \leftarrow \chi_{\mathrm{in}})\big)^{\dagger} = \hat{\mathcal{F}}^{\dagger}_{\beta}(x')\;\mathcal{U}(x' \leftarrow x; \gamma^{-1})\;\mathsf{W}^{\dagger}_{\chi_{\mathrm{in}}\leftarrow\chi_{\mathrm{out}}}\;\hat{\mathcal{F}}_{\alpha}(x),$$

which is exactly $\hat{G}^{\beta\alpha}_{\Lambda}(x', x; \mathbf{W}^{\dagger}, \chi_{\mathrm{in}} \leftarrow \chi_{\mathrm{out}})$, in accordance with the Hermitian symmetry of channels (*Lemma 4.66*).

For the grading, recall that the dialectic degree is carried by the selector labels and composes additively (*Definition 4.94and Lemma 4.98*). The dressed factor $\mathsf{W}_{\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}}$ acts in the RR fibre and is of degree 0 in the dialectic grading; hence

$$\deg\big(\hat{G}^{\alpha\beta}_{\Lambda}(\cdots)\big) = \deg\big(\hat{\mathcal{F}}^{\dagger}_{\alpha}\big) + \deg\big(\hat{\mathcal{F}}_{\beta}\big) = \deg(\alpha) + \deg(\beta),$$

as claimed.

(c) Functoriality. Let $\mathbf{W} = \mathbf{W}_2\mathbf{W}_1$. Then, by (4.67),

$$\hat{G}^{\alpha\beta}_{\Lambda}\big(x, x'; \mathbf{W}, \chi_2 \leftarrow \chi_0\big) = \hat{\mathcal{F}}^{\dagger}_{\alpha}(x)\, \mathcal{U}(x \leftarrow x'; \gamma) \Big(\sum_{\chi_1} \mathsf{W}_{2,\chi_2\leftarrow\chi_1}\, \mathsf{W}_{1,\chi_1\leftarrow\chi_0} \Big)\, \hat{\mathcal{F}}_{\beta}(x'),$$

where the sum over the intermediate control index χ_1 is the control-resolved convolution (*Lemma* 4.149*and eq.* (4.47)). Thus the dressing composes by $\mathsf{W}_{2,\chi_2\leftarrow\chi_1}\, \mathsf{W}_{1,\chi_1\leftarrow\chi_0}$ as stated.

For geometric concatenation of paths $\gamma_2 : x'' \to x$, $\gamma_1 : x' \to x''$, one has $\mathcal{U}(x \leftarrow x'; \gamma_2 \star \gamma_1) = \mathcal{U}(x \leftarrow x''; \gamma_2)\, \mathcal{U}(x'' \leftarrow x'; \gamma_1)$ by functoriality of transport (*Proposition* 4.52). Inserting this into (4.67) shows that the geometric composition of segments and the algebraic composition of dressings factorize compatibly, proving functoriality.

Items (a)–(c) together establish Proposition 4.167. □

Corollary 4.168 (Sum/difference words from Hadamard conjugation)**.** *With* $\mathbf{W} = \mathbf{U}_{\mathrm{Had}}$ *(Definition 2.119 and eq.* (2.78)*) and* $|\chi_{\mathrm{in}}\rangle = |\chi_{\mathrm{out}}\rangle \in \{|+\rangle, |-\rangle\}$*, the RR matrix elements are respectively the* sum *and* difference *of the two parallel RR paths (eq.* (2.79)*). The induced channels* $\hat{G}^{\alpha\beta}_{\Lambda}(x, x'; \mathbf{U}_{\mathrm{Had}}, \chi \leftarrow \chi)$ *therefore register holosymmetric interference between alternative mechanisms in the sense of Chapter 3.*

Proof. Let $A := \hat{\mathbf{U}}^{A_G B_N}_{\mathrm{RR}}$ and $B := \hat{\mathbf{U}}^{A_N B_G}_{\mathrm{RR}}$, and set $\hat{\mathbf{U}}_{\pm} := A \pm B$ (cf. eq. (2.79)). Take $\mathbf{W} = \mathbf{U}_{\mathrm{Had}} = (H_c \otimes I)\, \mathbf{U}_{\mathrm{ctrl}}\, (H_c \otimes I)$ (Definition 2.119 and eq. (2.78)). By the Hadamard-wrapped block identities (Proposition 4.153 and eq. (4.50)), for $\chi \in \{+, -\}$ we have

$$\mathsf{W}_{+\leftarrow+} = \tfrac{1}{2}\,\hat{\mathbf{U}}_{+}, \qquad \mathsf{W}_{-\leftarrow+} = \tfrac{1}{2}\,\hat{\mathbf{U}}_{-}, \qquad \mathsf{W}_{+\leftarrow-} = \tfrac{1}{2}\,\hat{\mathbf{U}}_{-}, \qquad \mathsf{W}_{-\leftarrow-} = \tfrac{1}{2}\,\hat{\mathbf{U}}_{+}.$$

Thus, when $|\chi_{\mathrm{in}}\rangle = |\chi_{\mathrm{out}}\rangle = |+\rangle$ (resp. $|-\rangle$), the RR element is proportional to the *sum* (resp. *difference*) of the two parallel RR paths ($\hat{\mathbf{U}}_{+}$ resp. $\hat{\mathbf{U}}_{-}$), as stated.

For the induced channels, recall the wordwise dressing definition

$$\hat{G}^{\alpha\beta}_{\Lambda}\big(x, x'; \mathbf{W}, \chi_{\mathrm{out}} \leftarrow \chi_{\mathrm{in}}\big) := \hat{\mathcal{F}}^{\dagger}_{\alpha}(x)\, \mathcal{U}(x \leftarrow x'; \gamma)\, \mathsf{W}_{\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}}\, \hat{\mathcal{F}}_{\beta}(x') \quad \text{(cf. eq. (4.67))}. \tag{4.68}$$

By linearity in the RR factor, inserting the above RR elements gives, for example,

$$\hat{G}^{\alpha\beta}_{\Lambda}\big(x, x'; \mathbf{U}_{\mathrm{Had}}, + \leftarrow +\big) = \tfrac{1}{2}\Big(\hat{G}^{\alpha\beta}_{\Lambda}\big(x, x'; A, + \leftarrow +\big) \;+\; \hat{G}^{\alpha\beta}_{\Lambda}\big(x, x'; B, + \leftarrow +\big)\Big),$$

and similarly

$$\hat{G}^{\alpha\beta}_{\Lambda}\big(x, x'; \mathbf{U}_{\mathrm{Had}}, - \leftarrow +\big) = \tfrac{1}{2}\Big(\hat{G}^{\alpha\beta}_{\Lambda}\big(x, x'; A, - \leftarrow +\big) \;-\; \hat{G}^{\alpha\beta}_{\Lambda}\big(x, x'; B, - \leftarrow +\big)\Big),$$

with the analogous relations for $-$ input. Hence the channels produced by $\mathbf{U}_{\mathrm{Had}}$ are *sum/difference* combinations of the channels associated to the two parallel RR mechanisms. By Proposition 4.167(a,b), these dressed channels inherit gauge covariance and Hermitian/degree structure; the $\pm$ superpositions therefore *register holosymmetric interference* between the alternatives in the precise wordwise sense (constructive in the $+$ block, destructive in the $-$ block), as advertised in Chapter 3.

This proves Corollary 4.168. □

Theorem 4.169 (Central (or approximately central) holosymmetry — words version)**.** *Let* G *be the* $*$*-algebra generated by holosymmetric channel words in Section 4.2, with alphabet* $\Sigma := \{\hat{\mathcal{F}}_\alpha(x), \hat{\mathcal{F}}_\alpha^\dagger(x),\ \mathcal{U}(x \leftarrow x';\gamma)\}$ *as in Definitions 4.48 and 4.61 and block composition through intermediate loci (Proposition 4.85). Define the holosymmetric self-channel*

$$\hat{G}_\Lambda^{ss}(x,x') := \hat{\mathcal{F}}_s^\dagger(x)\,\mathcal{U}(x \leftarrow x';\gamma)\,\hat{\mathcal{F}}_s(x').$$

Assume any one of the following endogenous hypotheses:

(C1$^{\text{word}}$) Control/branch transitivity. *The inner automorphism group* $\mathcal{H}$ *generated by control-basis rotations* $\mathbf{U}_c \in \mathrm{SU}(2)$ *(as in the holosymmetric interferometers, Definitions 2.119, 2.123 and 2.124) and the branch-swap involution* $\sigma : A_G B_N \leftrightarrow A_N B_G$ *on path families (Definition 2.108) acts by conjugation on* G*, and its representation is (nearly) irreducible on each typed Hom-space of words (Remark 4.140). Moreover,* $\hat{G}_\Lambda^{ss}$ *is* $\mathcal{H}$*-invariant (basis-independent and swap-symmetric; cf. the commutator-lensing normal form and effect equality in Definition 3.12, eq. (3.46), and Corollary 3.22).*

(C2$^{\text{word}}$) Flat holosymmetric section. *There exists a gauge in which* $\hat{\mathcal{F}}_s$ *is parallel along admissible paths, i.e.* $\mathcal{U}(x \leftarrow x';\gamma)\,\hat{\mathcal{F}}_s(x') = \mathrm{e}^{i\phi(x,x';\gamma)}\,\hat{\mathcal{F}}_s(x)$ *for all* γ *joining* x' *to* x *(up to a scalar phase), cf. Definitions 4.48 and 4.49 and Proposition 4.54.*

(C3$^{\text{word}}$) Conditional expectation onto holosymmetric invariants. *There is a contractive, completely positive, idempotent map* $\mathbb{E} : \mathsf{G} \to \mathsf{G}$ *(a Reynolds operator) obtained by averaging over the control-basis group and the branch-swap (cf. Definitions 2.119 and 2.123 for control actions and Definition 2.108 for swap), such that* $\mathbb{E}$ *projects onto the* $\mathcal{H}$*-fixed subalgebra and* $\hat{G}_\Lambda^{ss} \in \operatorname{Ran}\mathbb{E}$*.*

Then for all $X \in \mathsf{G}$*,*

$$[\,\hat{G}_\Lambda^{ss}(x,x'),\,X\,] = 0.$$

If the corresponding hypothesis holds only approximately (e.g. the $\mathcal{H}$*-action is nearly irreducible, or the parallel-transport condition in* (C2$^{\text{word}}$) *holds up to a phase defect* ε*, or* $\mathbb{E}$ *is computed with numerical/structural tolerance* δ*), then there exists* $\Delta \geq 0$ *(the* centrality defect*) such that*

$$\sup_{\|X\|=1} \left\|[\,\hat{G}_\Lambda^{ss},\,X\,]\right\| \;\leq\; \Delta, \qquad \Delta \to 0 \text{ as the defect parameters } \varepsilon,\delta \to 0.$$

Proof. We argue case by case, using only the holosymmetric word architecture.

Generator test. Since G is generated by the alphabet Σ (Definitions 4.48 and 4.61) and closed under block composition through intermediate loci (Proposition 4.85), it suffices to show $[\,\hat{G}_\Lambda^{ss},\,Y\,] = 0$ for each generator $Y \in \Sigma$; the general case follows by induction on word length and the typing/composition rules (Remark 4.140 and Proposition 4.85).

(C1$^{\text{word}}$): Control/branch transitivity. Let $\mathcal{H}$ be the inner automorphism group generated by (i) conjugations by holosymmetric control unitaries $(U_c \otimes I_{\mathrm{RR}})$ (prep/read rotations, Definitions 2.119 and 2.124), and (ii) the branch-swap σ that interchanges the path families $(A_G B_N) \leftrightarrow (A_N B_G)$ (Definition 2.108). By the interferometric constructions (Definition 3.12 and Corollary 3.22), the holosymmetric effects (and hence $\hat{G}_\Lambda^{ss}$ as a word) are basis-independent and swap-symmetric, so $\mathrm{Ad}_h(\hat{G}_\Lambda^{ss}) = \hat{G}_\Lambda^{ss}$ for all $h \in \mathcal{H}$. Let ρ be the (nearly) irreducible representation of $\mathcal{H}$ on a typed Hom-space of words (Remark 4.140). By the fixed-point/Schur argument, the $\mathcal{H}$-invariant elements in that space are scalars, so $\hat{G}_\Lambda^{ss}$ commutes with every

generator Y of matching type. Since G is the sum of such typed Hom-spaces, $\hat{G}^{ss}_{\Lambda}$ commutes with every generator, hence with every word. In the near-irreducible case, the fixed point is within operator-norm Δ of a scalar, giving the uniform defect bound.

(C2$^{\text{word}}$): Flat holosymmetric section. In the parallel gauge (Definition 4.48 and Proposition 4.54), $\mathcal{U}(x \leftarrow x';\gamma)\,\hat{\mathcal{F}}_s(x') = \mathrm{e}^{i\phi}\hat{\mathcal{F}}_s(x)$ for all admissible γ. Then

$$\hat{G}^{ss}_{\Lambda}(x,x') = \hat{\mathcal{F}}^{\dagger}_s(x)\,\mathcal{U}(x \leftarrow x';\gamma)\,\hat{\mathcal{F}}_s(x') = \mathrm{e}^{i\phi(x,x';\gamma)}\,\hat{\mathcal{F}}^{\dagger}_s(x)\hat{\mathcal{F}}_s(x).$$

The right-hand side is a scalar phase times the endpoint word $\hat{\mathcal{F}}^{\dagger}_s(x)\hat{\mathcal{F}}_s(x)$, which acts as the identity on the s-typed range and annihilates orthogonal types (typing/orthogonality in Remark 4.140). Consequently $\hat{G}^{ss}_{\Lambda}$ acts as a scalar on each typed block and hence commutes with (i) endpoint selectors (Definition 4.61) and (ii) transports $\mathcal{U}(\cdot)$; the phase is central and block composition is admissible (Proposition 4.85). Thus $[\,\hat{G}^{ss}_{\Lambda},\,Y\,]=0$ for all generators Y, hence for all words. If flatness holds up to a phase defect ε, the commutator norm is $O(\varepsilon)$ uniformly in $\|X\|=1$ (Proposition 4.54 and Corollary 4.64).

(C3$^{\text{word}}$): Conditional expectation onto invariants. Define the Reynolds operator

$$\mathbb{E}(Y) \;:=\; \int_{\mathrm{SU}(2)} (U_c\otimes I)^{\dagger}\,Y\,(U_c\otimes I)\,d\mu(U_c) \quad\text{and}\quad \mathbb{E}_{\sigma}(Y) \;:=\; \tfrac{1}{2}\Big(Y+\sigma(Y)\Big),$$

and set $\mathbb{E} := \mathbb{E}_{\sigma}\circ\mathbb{E}$ (Definitions 2.108, 2.119 and 2.123 for the actions averaged). Then $\mathbb{E}$ is a contractive, CP, idempotent conditional expectation onto the $\mathcal{H}$-fixed subalgebra. By control-basis independence and swap symmetry of the holosymmetric effect (Definition 3.12 and Corollary 3.22), $\hat{G}^{ss}_{\Lambda}$ is $\mathcal{H}$-fixed, hence $\hat{G}^{ss}_{\Lambda} = \mathbb{E}(\hat{G}^{ss}_{\Lambda}) \in \operatorname{Ran}\mathbb{E}$. For any generator Y, invariance of the range implies $(U^{\dagger}YU)$ commutes with $\mathbb{E}(Z)$ for all $U\in\mathcal{H}$, and averaging yields $[\,\mathbb{E}(Z),Y\,]=0$. Taking $Z=\hat{G}^{ss}_{\Lambda}$ shows that $\hat{G}^{ss}_{\Lambda}$ commutes with every generator, hence with every word. If the averaging is approximate with tolerance δ, we obtain the uniform bound.

In all three cases we have $[\,\hat{G}^{ss}_{\Lambda},\,Y\,]=0$ for the alphabet generators Y (Definitions 4.48 and 4.61), hence $[\,\hat{G}^{ss}_{\Lambda},\,X\,]=0$ for all words $X\in\mathsf{G}$ by closure under composition (Proposition 4.85). The approximate variants give the stated operator-norm defect bound. □

Theorem 4.170 (Nilpotent p–s edge — words version)**.** *Let G be the $*$-algebra generated by holosymmetric channel words (Section 4.2), with letters $\hat{G}^{\alpha\beta}_{\Lambda}(x_j,x_{j-1}) = \hat{\mathcal{F}}^{\dagger}_{\alpha}(x_j)\,\mathcal{U}(x_j \leftarrow x_{j-1};\gamma_j)\,\hat{\mathcal{F}}_{\beta}(x_{j-1})$ from Definitions 4.48, 4.49 and 4.61, typed and composed through intermediate loci as in Definition 4.141, Proposition 4.85, and Remark 4.140. Define $\hat{G}^{ps}_{\Lambda},\hat{G}^{sp}_{\Lambda}$ with fixed endpoints (x,x').*

Assume any one of the following endogenous hypotheses:

- (N1$^{\text{word}}$) Typing prohibition at the edge. *For each admissible locus y, the mixed composition is ill-typed and vanishes: $\hat{\mathcal{F}}_s(y)\,\hat{\mathcal{F}}^{\dagger}_p(y)=0$, by the domain/codomain typing of words (Remark 4.140 and Proposition 4.85).*
- (N2$^{\text{word}}$) Orthogonality at an intermediate locus. *There exists y with $\hat{\mathcal{F}}^{\dagger}_s(y)\hat{\mathcal{F}}_p(y)=0$ and a dialectically complete resolution at y (Definition 4.84 and Proposition 4.85).*
- (N3$^{\text{word}}$) Spectral support exit. *With the regulated fibre at y given by $E^{(y)}([0,\lambda_{\text{cutoff}}])$ (eq. (4.303)), a $p\to s$ step maps into $\operatorname{Ran}(I-E^{(y)})$ while a subsequent $p\to s$ requires domain $\operatorname{Ran}(E^{(y)})$.*

Then

$$(\hat{G}^{ps}_{\Lambda})^2 = (\hat{G}^{sp}_{\Lambda})^2 = 0.$$

The conclusion is stable under endpoint gauge conjugation $X \mapsto g(x)\, X\, g(x')^\dagger$ by Proposition 4.54 and Corollary 4.64.

Proof. Square expansion and the kill term. For fixed endpoints (x, x'),

$$(\hat{G}^{ps}_{\Lambda})^2 = \hat{\mathcal{F}}^\dagger_p(x)\,\mathcal{U}(x \leftarrow y;\gamma_1)\,\hat{\mathcal{F}}_s(y)\;\hat{\mathcal{F}}^\dagger_p(y)\,\mathcal{U}(y \leftarrow x';\gamma_2)\,\hat{\mathcal{F}}_s(x') \tag{4.69}$$

$$= \hat{\mathcal{F}}^\dagger_p(x)\,\mathcal{U}(x \leftarrow y;\gamma_1)\,\underbrace{\big(\hat{\mathcal{F}}_s(y)\hat{\mathcal{F}}^\dagger_p(y)\big)}_{\text{kill term}}\,\mathcal{U}(y \leftarrow x';\gamma_2)\,\hat{\mathcal{F}}_s(x'). \tag{4.70}$$

(N1$^{\text{word}}$). By the word typing rule (Remark 4.140) implemented via block composition (Proposition 4.85), $\hat{\mathcal{F}}_s(y)\hat{\mathcal{F}}^\dagger_p(y) = 0$. Hence (4.70) gives $(\hat{G}^{ps}_{\Lambda})^2 = 0$. The *sp* case is identical with $p \leftrightarrow s$.
(N2$^{\text{word}}$). Insert a dialectically complete resolution at y (Definition 4.84) to write

$$\hat{G}^{ps}_{\Lambda}(x, x') = \sum_{\gamma \in \mathcal{I}} \hat{G}^{p\gamma}_{\Lambda}(x, y)\, \hat{G}^{\gamma s}_{\Lambda}(y, x').$$

Squaring produces middle factors of the form $\hat{G}^{\gamma s}_{\Lambda}(y, x')\, \hat{G}^{p\delta}_{\Lambda}(x, y)$ which, when expanded back to selectors/transports, carry either $\hat{\mathcal{F}}^\dagger_s(y)\hat{\mathcal{F}}_p(y)$ or its adjoint at y—both zero by hypothesis. Thus each summand vanishes and $(\hat{G}^{ps}_{\Lambda})^2 = 0$; same for *sp*.
(N3$^{\text{word}}$). Insert $\mathbf{1} = E^{(y)} + [I - E^{(y)}]$ between the two *ps* factors. By the support-exit rule, the image of the first *ps* sits in $\mathrm{Ran}(I - E^{(y)})$, while the second *ps* requires domain in $\mathrm{Ran}(E^{(y)})$, so their composition is zero on the regulated fibre (eq. (4.303)). Hence $(\hat{G}^{ps}_{\Lambda})^2 = 0$ and similarly for *sp*.

Gauge stability. Endpoint conjugation preserves zero operators and the hypotheses (Proposition 4.54 and Corollary 4.64), so the conclusion is gauge-invariant. □

Remark 4.171 (Optional grading overlay)**.** If you retain the $\mathbb{Z}_2^2$ overlay (Definition 4.94 and Lemma 4.98), the words above also satisfy $\deg(\hat{G}^{ps}) = \deg(\hat{G}^{sp}) = (1, 0)$ and $\deg\big((\hat{G}^{ps})^2\big) = \deg\big((\hat{G}^{sp})^2\big) = (0, 0)$, consistent with nilpotency. The proof itself, however, does not use grading.

Definition 4.172 (Nilpotency defect)**.** Define $\eta := \max\Big\{\|(\hat{G}^{ps}_{\Lambda})^2\|,\ \|(\hat{G}^{sp}_{\Lambda})^2\|\Big\}$. Under approximate forms of (N1$^{\text{word}}$)–(N3$^{\text{word}}$) (typing leakage, small orthogonality error, or imperfect spectral separation), η quantifies the deviation from exact nilpotency.

Theorem 4.173 (Bosonic compatibility in the 0–*s* sector — words version)**.** *Let G be the $*$-algebra of holosymmetric channel words (Section 4.2), generated by letters $\hat{G}^{\alpha\beta}_{\Lambda}(x_j, x_{j-1}) = \hat{\mathcal{F}}^\dagger_\alpha(x_j)\,\mathcal{U}(x_j \leftarrow x_{j-1};\gamma_j)\,\hat{\mathcal{F}}_\beta(x_{j-1})$ (Definitions 4.48, 4.49 and 4.61), typed and composed through intermediate loci (Definition 4.141, Proposition 4.85, and Remark 4.140).*

Define the diagonal self-channels at fixed endpoints (x, x'):

$$\hat{G}^{00}_{\Lambda}(x, x') = \hat{\mathcal{F}}^\dagger_0(x)\,\mathcal{U}(x \leftarrow x';\gamma)\,\hat{\mathcal{F}}_0(x'), \qquad \hat{G}^{ss}_{\Lambda}(x, x') = \hat{\mathcal{F}}^\dagger_s(x)\,\mathcal{U}(x \leftarrow x';\gamma)\,\hat{\mathcal{F}}_s(x').$$

Assume any one of the following endogenous hypotheses:

- (B1$^{\text{word}}$) Common spectral generator (functional calculus). *There exists a spectral operator $\hat{D}$ (Chapter IV; Definition 4.341 and eq. (4.159)) and Borel functions f_{00}, f_{ss} such that $\hat{G}^{00}_{\Lambda} = f_{00}(\hat{D})$ and $\hat{G}^{ss}_{\Lambda} = f_{ss}(\hat{D})$ (on the relevant typed Hom-space).*
- (B2$^{\text{word}}$) Parallel diagonals on a flat patch. *On a simply connected neighbourhood with $\mathcal{F} = 0$ (Proposition 4.59), there exists a gauge (endpoint conjugation; Proposition 4.54 and Corollary 4.64) such that, for all admissible paths, $\mathcal{U}(x \leftarrow x'; \gamma)\, \hat{\mathcal{F}}_{\alpha}(x') = \mathrm{e}^{i\phi_{\alpha}(x,x';\gamma)}\, \hat{\mathcal{F}}_{\alpha}(x)$ for $\alpha \in \{0, s\}$. Equivalently (Corollary 4.146), the transporter collapses to a phase and both diagonals reduce to endpoint words.*
- (B3$^{\text{word}}$) Pinching onto the $0/s$ diagonal subalgebra. *Let $P_{\alpha}(x) := \hat{\mathcal{F}}^{\dagger}_{\alpha}(x)\hat{\mathcal{F}}_{\alpha}(x)$ and similarly at x'. The completely positive, contractive, idempotent map*

$$\Pi(Y) \;:=\; \sum_{\alpha\in\{0,s\}} \sum_{\beta\in\{0,s\}} \Big(P_{\alpha}(x)\, Y\, P_{\beta}(x') \Big)$$

maps G onto the commutative C^*-subalgebra generated by $\{P_0(x), P_s(x)\}$ at x and $\{P_0(x'), P_s(x')\}$ at x', and $\hat{G}^{00}_{\Lambda}, \hat{G}^{ss}_{\Lambda}$ lie in $\operatorname{Ran}\Pi$ (are fixed points).

Then, for all x, x',

$$[\, \hat{G}^{00}_{\Lambda}(x, x'),\, \hat{G}^{ss}_{\Lambda}(x, x')\,] = 0.$$

The conclusion is stable under endpoint conjugation $X \mapsto g(x)\, X\, g(x')^{\dagger}$ by Proposition 4.54 and Corollary 4.64.

Proof. Generator/word reduction. By Proposition 4.85 and the typing rules (Remark 4.140), it suffices to argue at the level of the diagonal self-channels defined above; commutation for these generators propagates to all words built from them and compatible letters.

(B1$^{\text{word}}$). If $\hat{G}^{00}_{\Lambda} = f_{00}(\hat{D})$ and $\hat{G}^{ss}_{\Lambda} = f_{ss}(\hat{D})$ for the same self-adjoint $\hat{D}$ (Definition 4.341), then functional calculus implies $[\, f_{00}(\hat{D}), f_{ss}(\hat{D})\,] = 0$. Hence $[\, \hat{G}^{00}_{\Lambda}, \hat{G}^{ss}_{\Lambda}\,] = 0$.

(B2$^{\text{word}}$). On a flat patch, choose a parallel gauge (Propositions 4.54 and 4.59) so that

$$\hat{G}^{00}_{\Lambda}(x, x') = \mathrm{e}^{i\phi_0(x,x';\gamma)}\, P_0(x), \qquad \hat{G}^{ss}_{\Lambda}(x, x') = \mathrm{e}^{i\phi_s(x,x';\gamma)}\, P_s(x).$$

By channel alphabet orthogonality/typing (Remark 4.140), $P_0(x)$ and $P_s(x)$ are endpoint projectors onto orthogonal modal subspaces, hence $P_0(x)P_s(x) = P_s(x)P_0(x) = 0$, and they commute. Phases are scalar, so $[\, \hat{G}^{00}_{\Lambda}, \hat{G}^{ss}_{\Lambda}\,] = 0$. Gauge covariance (Proposition 4.54 and Corollary 4.64) preserves commutators, so the result holds in any gauge.

(B3$^{\text{word}}$). The map Π is a conditional expectation (Reynolds operator) onto a *commutative* diagonal subalgebra (the "pinching" onto the $0/s$ blocks at both ends). Since $\hat{G}^{00}_{\Lambda}, \hat{G}^{ss}_{\Lambda}$ are 0–0 and s–s diagonals respectively, they are fixed by Π, i.e. lie in $\operatorname{Ran}\Pi$. Any two elements of a commutative range commute, hence $[\, \hat{G}^{00}_{\Lambda}, \hat{G}^{ss}_{\Lambda}\,] = 0$. Endpoint conjugation preserves $\operatorname{Ran}\Pi$ and commutators, so the conclusion is gauge-stable. □

Remark 4.174 (Optional grading overlay)**.** If you keep the $\mathbb{Z}_2^2$ bookkeeping, then $\deg(\hat{G}^{00}) = \deg(\hat{G}^{ss}) = (0, 0)$ (Definition 4.94), and by Lemma 4.95 one has $[A, B]_{\mathrm{gr}} = [A, B]$ for even A. Thus the graded and ordinary commutators coincide here. The proof above, however, does not use grading.

Definition 4.175 (Bosonic-commutation defect)**.** Under approximate forms of (B1$^{\mathrm{word}}$)–(B3$^{\mathrm{word}}$) (e.g. imperfect shared functional calculus, non-flat phase leakage, or imperfect pinching), define

$$\zeta \;:=\; \big\| [\, \hat{G}^{00}_{\Lambda}, \hat{G}^{ss}_{\Lambda} \,] \big\|.$$

Then $\zeta = 0$ in the exact hypotheses and $\zeta \to 0$ as the defect parameters vanish.

4.2.1 Wordwise Meta-Claim, Compliance Vector, and Geometric Leverage

Proposition 4.176 (Words-only derivability of the trio)**.** *Within the holosymmetric words framework (Section 4.2, Definitions 4.48, 4.61 and 4.141, Propositions 4.85 and 4.143, and Remark 4.140), assume the endogenous hypothesis families*

$$\begin{aligned} &(\mathrm{C1}^{\mathrm{word}})\text{–}(\mathrm{C3}^{\mathrm{word}}) && \textit{(Theorem 4.169)},\\ &(\mathrm{N1}^{\mathrm{word}})\text{–}(\mathrm{N3}^{\mathrm{word}}) && \textit{(Theorem 4.170)},\\ &(\mathrm{B1}^{\mathrm{word}})\text{–}(\mathrm{B3}^{\mathrm{word}}) && \textit{(Theorem 4.173)}. \end{aligned}$$

Then, for all admissible endpoints (x, x') and all channel words $X \in \mathsf{G}$,

(a) Central holosymmetry
$[\, \hat{G}^{ss}_{\Lambda}(x, x'),\, X \,] = 0.$

(b) Nilpotent *p*–*s* edge
$(\hat{G}^{ps}_{\Lambda})^2 = (\hat{G}^{sp}_{\Lambda})^2 = 0.$

(c) Bosonic 0–*s* compatibility
$[\, \hat{G}^{00}_{\Lambda}(x, x'),\, \hat{G}^{ss}_{\Lambda}(x, x') \,] = 0.$

All three statements are invariant under endpoint conjugation $X \mapsto g(x)\, X\, g(x')^{\dagger}$ by Proposition 4.54 and Corollary 4.64.

Remark 4.177 (Optional grading overlay)**.** If the $\mathbb{Z}_2^2$ overlay is retained (Definition 4.94 and Lemma 4.98), then (i) $\deg(\hat{G}^{ss}) = (0,0)$ implies $[A, \cdot]_{\mathrm{gr}} = [A, \cdot]$ by Lemma 4.95; (ii) $\deg(\hat{G}^{ps}) = \deg(\hat{G}^{sp}) = (1,0)$ is consistent with nilpotency and gauge covariance; (iii) $\deg(\hat{G}^{00}) = (0,0)$ and (i) give $[\hat{G}^{00}, \hat{G}^{ss}]_{\mathrm{gr}} = [\hat{G}^{00}, \hat{G}^{ss}] = 0$. The proofs of Theorems 4.169, 4.170 and 4.173 do not use grading.

Definition 4.178 (Holosymmetric compliance vector)**.** Let the *centrality defect* Δ be as in Definition 4.112, the *nilpotency defect* η as in Definition 4.172, and the *bosonic commutation defect* ζ as in Definition 4.175. Define the *holosymmetric compliance vector*

$$\mathfrak{K} \;:=\; (\Delta,\ \eta,\ \zeta) \;\in\; \mathbb{R}^3_{\geq 0}.$$

It is gauge-invariant under endpoint conjugation and vanishes iff the trio in Proposition 4.176 holds exactly.

Definition 4.179 (Operational signatures)**.** Using Chapter III interferometers and lensing (Definitions 2.119, 2.123, 2.124 and 3.12, Corollary 3.22, and eqs. (3.1) and (3.2)), define:

(i) **Visibility envelope width:** For a family of control unitaries $U_c \in \mathrm{SU}(2)$,

$$\mathcal{W}_{\mathrm{vis}} \;:=\; \sup_{U_c} \Big|\, \mathcal{V}(U_c) - \mathcal{V}(I) \,\Big|,$$

where $\mathcal{V}$ is the fringe visibility in the chosen model (eq. (3.2) and its asymmetric analogue).

(ii) **Lensing equality residual:**

$$\mathcal{L}_{\mathrm{res}} \;:=\; \Big\| \; E^{(+\!\to\!+)}_{\mathrm{Holo}} \;-\; E^{(-\!\to\!-)}_{\mathrm{Holo}} \; \Big\|,$$

with $E^{(\cdot)}_{\mathrm{Holo}}$ the lensing effects (Corollary 3.22).

(iii) **Two-step ps amplitude:**

$$\mathcal{A}^{(2)}_{ps} \;:=\; \Big\| \, (\hat{G}^{ps}_{\Lambda})^2 \, \Big\|, \qquad \mathcal{A}^{(2)}_{sp} \;:=\; \Big\| \, (\hat{G}^{sp}_{\Lambda})^2 \, \Big\|.$$

(iv) **Diagonal pinching mismatch:** Let Π be the diagonal pinching onto the $0/s$ subalgebra (Theorem 4.173, case (B3$^{\mathrm{word}}$)). For any observable Y formed from diagonal words,

$$\mathcal{P}_{\mathrm{mis}}(Y) \;:=\; \Big\| \, Y - \Pi(Y) \, \Big\|.$$

Lemma 4.180 (Pinching stability under almost-commutation)**.** *Let $\hat{G}^{00}_{\Lambda}, \hat{G}^{ss}_{\Lambda}$ be the $(0/s)$ diagonal blocks of the Fourfold channel (Definition 4.81), and let Π denote the diagonal pinching onto the $(0/s)$ block subalgebra (as used in Definition 4.179, cf. the commuting case in* (B3$^{\mathrm{word}}$) *of Theorem 4.173). Set the* bosonic-commutation defect $\zeta := \|[\hat{G}^{00}_{\Lambda}, \hat{G}^{ss}_{\Lambda}]\|$ *(Definition 4.175). Then there exists an absolute constant $C > 0$ such that for every observable Y generated by $(0/s)$–diagonal words one has*

$$\| \, Y - \Pi(Y) \, \| \;\le\; C \, \zeta.$$

In particular, if $\zeta = 0$ then Π is a conditional expectation onto the $(0/s)$ diagonal subalgebra and $\|Y - \Pi(Y)\| = 0$ for all such Y.

Proof. When $[\hat{G}^{00}_{\Lambda}, \hat{G}^{ss}_{\Lambda}] = 0$, the $(0/s)$ diagonal algebra is a commuting von Neumann subalgebra and Π is the conditional expectation onto it, so $Y = \Pi(Y)$ for all $(0/s)$–diagonal Y — this is precisely the content of the commuting case in Theorem 4.173.

For the almost-commuting case, represent Π as a double operator integral (DOI) with respect to the spectral measures of $\hat{G}^{00}_{\Lambda}$ and $\hat{G}^{ss}_{\Lambda}$. Standard DOI bounds imply Lipschitz-type stability: if the DOI kernel has bounded divided differences (true for the pinching kernel), then $\|Y - \Pi(Y)\| \le C \, \|[\hat{G}^{00}_{\Lambda}, \hat{G}^{ss}_{\Lambda}]\|$ for a universal $C > 0$. Since the defect ζ is gauge-invariant and norms are preserved under endpoint gauge (Proposition 4.54 and Corollary 4.64), the bound $\|Y - \Pi(Y)\| \le C \, \zeta$ holds uniformly for all $(0/s)$–diagonal words Y. □

Proposition 4.181 (Compliance-to-measurement bounds)**.** *There exist absolute constants $c_1, c_2, c_3 > 0$ (independent of system size and gauge) such that*

$$\mathcal{W}_{\mathrm{vis}} \;\le\; c_1 \, \Delta, \qquad \mathcal{L}_{\mathrm{res}} \;\le\; c_2 \, \Delta, \qquad \max\{\mathcal{A}^{(2)}_{ps}, \mathcal{A}^{(2)}_{sp}\} \;=\; \eta, \qquad \sup_{Y} \; \mathcal{P}_{\mathrm{mis}}(Y) \;\le\; c_3 \, \zeta.$$

Hence the holosymmetric compliance vector $\mathfrak{K} = (\Delta, \eta, \zeta)$ gives a state-independent envelope for interferometric and lensing deviations and diagonal pinching errors. In particular, $\mathfrak{K} = \vec{0}$ if and only if $\mathcal{W}_{\mathrm{vis}} = \mathcal{L}_{\mathrm{res}} = 0$, $\mathcal{A}^{(2)}_{ps} = \mathcal{A}^{(2)}_{sp} = 0$, and $\mathcal{P}_{\mathrm{mis}}(Y) = 0$ for all diagonal Y.

Proof idea. For the first two bounds, write holosymmetric effects in the lensing (commutator) normal form (Definition 3.12) and use the lensing equality in the central case (Corollary 3.22). A centrality defect Δ controls the variation of these effects under control-basis rotations and branch swap, hence the induced visibility and lensing differences; the constants absorb universal factors (e.g. the $1/4$ in the symmetric lensing effect). For the third, η is defined as the operator norm of the two-step edge (Definition 4.172). For the fourth, almost-commutation $\zeta = \|[\hat{G}^{00}_\Lambda, \hat{G}^{ss}_\Lambda]\|$ controls the deviation from the pinched diagonal via the *pinching-stability* Lemma Lemma 4.180; in the exact commuting case ($\zeta = 0$), the pinching coincides with the conditional expectation described in Remark Remark 4.182. Gauge covariance preserves all relevant norms throughout (Proposition 4.54 and Corollary 4.64).

Proof. **Preliminaries.** We work with: (a) the symmetric interferometer and its probabilities (Definition 2.123 and eq. (3.1)); (b) the commutator ("lensing") normal form (Definition 3.12); (c) the control–block split $\mathbf{U}_{\mathrm{ctrl}} = I_c \otimes U_+ + Z_c \otimes U_-$ (eq. (3.46)), which yields $[R, \mathbf{U}_{\mathrm{ctrl}}] = [R, Z_c] \otimes U_-$ for any control rotation $R \in \mathrm{SU}(2)$; (d) gauge covariance and norm preservation of channels (Proposition 4.54 and Corollary 4.64); and (e) the operational diagnostics $\mathcal{W}_{\mathrm{vis}}, \mathcal{L}_{\mathrm{res}}, \mathcal{A}^{(2)}_{ps}, \mathcal{A}^{(2)}_{sp}, \mathcal{P}_{\mathrm{mis}}$ (Definition 4.179), together with the defects Δ (centrality, Definition 4.112), η (nilpotency, Definition 4.172), ζ (bosonic-commutation, Definition 4.175).

(i) $\mathcal{W}_{\mathrm{vis}} \le c_1\,\Delta$. In the symmetric Mach–Zehnder realization (Definition 2.123), the control outcomes are governed by holosymmetric effects that decompose into "sum" and "difference" histories with universal $1/4$ prefactors (cf. eq. (3.1) and the derivation via Definition 3.12). Varying the control basis R modifies the effects by commutators where the only nontrivial, defect-sensitive contribution arises from failures of the ss block to commute with the word implementing the basis change. By the definition of centrality defect $\Delta = \sup \|[\hat{G}^{ss}_\Lambda, \cdot]\|$ (Definition 4.112) and submultiplicativity, the variation of the effects — and hence the visibility envelope $\mathcal{W}_{\mathrm{vis}}$ (Definition 4.179) — is bounded by a universal constant times Δ. All gauge choices are immaterial by Proposition 4.54 and Corollary 4.64. Thus $\mathcal{W}_{\mathrm{vis}} \le c_1\,\Delta$.

(ii) $\mathcal{L}_{\mathrm{res}} \le c_2\,\Delta$. *Lensing equality* states that the symmetric and asymmetric lensing effects coincide when the ss block is central (Definition 3.12 together with the equality statement in Corollary 3.22). With a nonzero centrality defect, their difference decomposes into a finite sum of graded commutators, each bounded by Δ up to the same universal constants (inherited from the interferometer's $1/4$ factors). Taking suprema over lensing settings (as in Definition 4.179) gives $\mathcal{L}_{\mathrm{res}} \le c_2\,\Delta$. Gauge invariance follows as in (i).

(iii) $\max\{\mathcal{A}^{(2)}_{ps}, \mathcal{A}^{(2)}_{sp}\} = \eta$. By definition (Definitions 4.172 and 4.179),

$$\eta = \max\Big\{ \|(\hat{G}^{ps}_\Lambda)^2\|,\ \|(\hat{G}^{sp}_\Lambda)^2\| \Big\} = \max\{\mathcal{A}^{(2)}_{ps}, \mathcal{A}^{(2)}_{sp}\}.$$

Under exact word-level nilpotency (Theorem 4.170) both norms vanish.

(iv) $\sup_Y \mathcal{P}_{\mathrm{mis}}(Y) \le c_3\,\zeta$. Let Π be the $(0/s)$–diagonal pinching used in Definition 4.179. By Lemma 4.180 we have $\|Y - \Pi(Y)\| \le C\,\zeta$ for a universal C and all $(0/s)$–diagonal words Y; hence $c_3 := C$ works.

Conclusion. Items (i)–(iv) yield the stated bounds. Moreover, the diagnostics vanish precisely when their controlling defects vanish: (iii) is tautological; (i)–(ii) reduce to exact centrality

($\Delta = 0$) via Corollary 3.22; (iv) reduces to exact commutation ($\zeta = 0$) by Lemma 4.180. Therefore

$$\mathfrak{K} = \vec{0} \iff \mathcal{W}_{\mathrm{vis}} = \mathcal{L}_{\mathrm{res}} = 0, \quad \mathcal{A}^{(2)}_{ps} = \mathcal{A}^{(2)}_{sp} = 0, \quad \mathcal{P}_{\mathrm{mis}}(Y) = 0 \ \ \forall Y,$$

and the proposition follows. □

Remark 4.182 (Spectral pinching as conditional expectation when $\zeta = 0$)**.** Assume the bosonic $0/s$ compatibility $[\hat{G}^{00}_\Lambda, \hat{G}^{ss}_\Lambda] = 0$ (Theorem 4.173) and let $\mathcal{D}_{0s}$ be the von Neumann subalgebra generated by the joint spectral projections of the commuting pair $(\hat{G}^{00}_\Lambda, \hat{G}^{ss}_\Lambda)$ inside the $(0/s)$-diagonal block (Definition 4.81). Then the pinching map Π used in Definition 4.179 coincides with the normal conditional expectation $\mathsf{E}_{0s} : \mathcal{B}(\mathcal{H}^{(2n)}_{\mathrm{RR}}) \to \mathcal{D}_{0s}$ given by the joint-spectral pinching

$$\mathsf{E}_{0s}(Y) \ = \ \int_\sigma E(d\omega)\, Y\, E(d\omega),$$

where E is the joint spectral measure of $(\hat{G}^{00}_\Lambda, \hat{G}^{ss}_\Lambda)$ and σ its joint spectrum. In particular, $\Pi = \mathsf{E}_{0s}$ is normal, unital, completely positive, contractive with $\|\Pi\| = 1$, idempotent $\Pi^2 = \Pi$, and $\mathcal{D}_{0s}$-bimodular: $\Pi(D_1 Y D_2) = D_1\, \Pi(Y)\, D_2$ for all $D_1, D_2 \in \mathcal{D}_{0s}$. Hence $\mathrm{Fix}(\Pi) = \mathcal{D}_{0s}$ and, for any $(0/s)$-diagonal word Y, $\|Y - \Pi(Y)\| = 0$. For $\zeta > 0$ (almost-commutation), Lemma 4.180 gives the quantitative stability bound $\|Y - \Pi(Y)\| \le C\,\zeta$, uniform under endpoint gauge by Proposition 4.54 and Corollary 4.64.

Definition 4.183 (Compliance regimes)**.** Fix thresholds $\varepsilon_\Delta, \varepsilon_\eta, \varepsilon_\zeta > 0$. The *holosymmetric ε-compliant regime* is

$$\mathcal{R}_\varepsilon \ := \ \Big\{ \ \mathfrak{K} \in \mathbb{R}^3_{\ge 0} : \ \Delta \le \varepsilon_\Delta, \ \eta \le \varepsilon_\eta, \ \zeta \le \varepsilon_\zeta \ \Big\}.$$

By Proposition 4.181, $\mathcal{R}_\varepsilon$ induces uniform bounds on $\mathcal{W}_{\mathrm{vis}}, \mathcal{L}_{\mathrm{res}}, \mathcal{A}^{(2)}_{ps}, \mathcal{A}^{(2)}_{sp}, \mathcal{P}_{\mathrm{mis}}$.

Definition 4.184 (Extended holosymmetric compliance profile)**.** Let $\mathcal{J}$ be a finite index set of defect functionals $\{\delta_j\}_{j\in\mathcal{J}}$, each $\delta_j \ge 0$, gauge-invariant under endpoint conjugation, and operationally measurable (cf. Definition 4.179). Define the *compliance profile*

$$\mathfrak{K}_{\mathcal{J}} \ := \ (\delta_j)_{j\in\mathcal{J}} \ \in \ \mathbb{R}^{|\mathcal{J}|}_{\ge 0}.$$

As a default baseline, take $\mathcal{J}_0 = \{\Delta, \eta, \zeta\}$, recovering $\mathfrak{K}_{\mathcal{J}_0} = (\Delta, \eta, \zeta)$ in $\mathbb{R}^3_{\ge 0}$.

Definition 4.185 (General compliance regimes)**.** Given thresholds $\varepsilon = (\varepsilon_j)_{j\in\mathcal{J}} \in \mathbb{R}^{|\mathcal{J}|}_{>0}$, the *$\varepsilon$-compliant regime* is

$$\mathcal{R}^{\mathcal{J}}_\varepsilon \ := \ \Big\{ \mathfrak{K}_{\mathcal{J}} \in \mathbb{R}^{|\mathcal{J}|}_{\ge 0} \ : \ \delta_j \le \varepsilon_j \ \ \forall\, j \in \mathcal{J} \Big\}.$$

For the baseline $\mathcal{J}_0 = \{\Delta, \eta, \zeta\}$, this reduces to $\mathcal{R}^{\mathcal{J}_0}_\varepsilon = \{(\Delta, \eta, \zeta) : \Delta \le \varepsilon_\Delta, \ \eta \le \varepsilon_\eta, \ \zeta \le \varepsilon_\zeta\} \subset \mathbb{R}^3_{\ge 0}$.

Remark 4.186 (Comparing profiles and choosing dimensions)**.** (i) **Projection/dominance.** Any extended profile projects onto the baseline via $\pi_{\mathcal{J}_0}(\mathfrak{K}_{\mathcal{J}}) = (\Delta, \eta, \zeta)$; membership in $\mathcal{R}_\varepsilon^{\mathcal{J}}$ implies membership in $\mathcal{R}_\varepsilon^{\mathcal{J}_0}$, but not conversely.

(ii) **Aggregates.** For a single-number certificate, use a weighted max or norm, e.g. $\|\mathfrak{K}_{\mathcal{J}}\|_{\infty,w} := \max_{j\in\mathcal{J}} w_j\, \delta_j$.

(iii) **Suggested extensions.** Beyond $\{\Delta, \eta, \zeta\}$, typical δ_j include: orthogonality defect of the triadic decomposition (Proposition 4.475); flatness/leakage along transports (via curvature bounds); endpoint-covariance drift; typing leakage; normal-form residual for ss; and spectral pinching/tomography residuals (cf. Definition 4.179).

Proposition 4.187 (Diagonal $0/s$ refinement of triadic metric/time)**.** *Assume bosonic $0/s$ compatibility (Theorem 4.173) so that $[\,\hat{G}^{00}_\Lambda, \hat{G}^{ss}_\Lambda\,] = 0$. Then at each locus x there exists a joint spectral resolution for the endpoint projectors $P_0(x), P_s(x)$ and the diagonal channel observables built from them. Consequently, the diagonal part of the triadic fibres (Sections 4.10 and 4.12) splits as a direct sum*

$$\mathcal{F}^{\mathrm{dist}}_{x,\mathrm{diag}} \;\cong\; \mathcal{F}^{\mathrm{dist},(0)}_x \oplus \mathcal{F}^{\mathrm{dist},(s)}_x, \qquad \mathcal{F}^{\mathrm{time}}_{x,\mathrm{diag}} \;\cong\; \mathcal{F}^{\mathrm{time},(0)}_x \oplus \mathcal{F}^{\mathrm{time},(s)}_x,$$

and one can define refined diagonal contributions

$$d^{(0)}_{\mathrm{vac}}(x,x'),\; d^{(s)}_{\mathrm{vac}}(x,x'), \qquad t^{(0)}_{\mathrm{vac}}(x,x'),\; t^{(s)}_{\mathrm{vac}}(x,x'),$$

whose sum recovers the diagonal components in the triadic metric/time (Definition 4.463 and eqs. (4.210) and (4.211)). These refinements are gauge-invariant and orthogonal to the off-diagonal correlator/spectral axes by Proposition 4.475.

Proof idea. Commutation yields simultaneous functional calculus for diagonal words, so one can project the diagonal contribution onto 0 and s sectors at both endpoints. Orthogonality with correlator/spectral parts follows from Proposition 4.475, and gauge covariance from Proposition 4.54 and Corollary 4.64.

Proof. Preliminaries. Work over the base $\mathcal{C}_d$ of Definition 4.617. All fibrewise objects live in the Hilbert bundle $\pi : \mathfrak{F} \to \mathcal{C}_d$ ((4.302)) with spectral restriction ((4.303)). Transport and gauge covariance are those of Section 4.19.1.

Step 1: Diagonal observables and block support. Fix endpoints (x, x'). Let

$$P_\alpha(x) := \hat{\mathcal{F}}^\dagger_\alpha(x)\hat{\mathcal{F}}_\alpha(x), \qquad P_\alpha(x') := \hat{\mathcal{F}}^\dagger_\alpha(x')\hat{\mathcal{F}}_\alpha(x') \quad (\alpha \in \{0, s\}).$$

By typing/orthogonality of endpoint selectors (Remark 4.140), $P_0(x)P_s(x) = P_s(x)P_0(x) = 0$ and similarly at x'. Define the (endpoint) diagonal channel observables at x

$$\mathsf{O}_{00}(x) := \hat{G}^{00}_\Lambda(x,x')\,\hat{G}^{00}_\Lambda(x,x')^\dagger, \qquad \mathsf{O}_{ss}(x) := \hat{G}^{ss}_\Lambda(x,x')\,\hat{G}^{ss}_\Lambda(x,x')^\dagger,$$

and analogously at x' with adjoint order reversed. Because $\hat{G}^{00}_\Lambda(x,x') = P_0(x)\hat{G}^{00}_\Lambda(x,x')P_0(x')$ and $\hat{G}^{ss}_\Lambda(x,x') = P_s(x)\hat{G}^{ss}_\Lambda(x,x')P_s(x')$ by typing, we have the support relations

$$\mathsf{O}_{00}(x) = P_0(x)\,\mathsf{O}_{00}(x)\,P_0(x), \qquad \mathsf{O}_{ss}(x) = P_s(x)\,\mathsf{O}_{ss}(x)\,P_s(x),$$

hence $\mathsf{O}_{00}(x)\mathsf{O}_{ss}(x) = \mathsf{O}_{ss}(x)\mathsf{O}_{00}(x) = 0$.

Step 2: Joint spectral resolution on the diagonal family. By bosonic compatibility, $[\hat{G}^{00}_\Lambda, \hat{G}^{ss}_\Lambda] = 0$ (Theorem 4.173). Therefore the normal operators $\mathsf{O}_{00}(x)$ and $\mathsf{O}_{ss}(x)$ commute. They also commute with $P_0(x), P_s(x)$ because each is supported on its own range. Thus the family $\{P_0(x), P_s(x), \mathsf{O}_{00}(x), \mathsf{O}_{ss}(x)\}$ is a commuting set of normal operators on the modal fibre at x. By the joint spectral theorem, there exists a joint spectral measure resolving them simultaneously. The same holds at x'.

Step 3: Algebraic splitting of the diagonal subalgebra. Let $\mathsf{A}_{x,\mathrm{diag}}$ be the unital C^*-algebra generated by $\{P_0(x), P_s(x), \mathsf{O}_{00}(x), \mathsf{O}_{ss}(x)\}$. Because $P_0(x), P_s(x)$ are orthogonal central idempotents in $\mathsf{A}_{x,\mathrm{diag}}$, the map

$$\Phi : \mathsf{A}_{x,\mathrm{diag}} \longrightarrow P_0(x)\mathsf{A}_{x,\mathrm{diag}}P_0(x) \oplus P_s(x)\mathsf{A}_{x,\mathrm{diag}}P_s(x), \quad A \mapsto \Big(P_0(x)AP_0(x),\ P_s(x)AP_s(x)\Big)$$

is a C^*-isomorphism (standard block decomposition by orthogonal central projections). Define

$$\mathsf{A}^{(0)}_x := P_0(x)\mathsf{A}_{x,\mathrm{diag}}P_0(x), \qquad \mathsf{A}^{(s)}_x := P_s(x)\mathsf{A}_{x,\mathrm{diag}}P_s(x).$$

Then $\mathsf{A}_{x,\mathrm{diag}} \cong \mathsf{A}^{(0)}_x \oplus \mathsf{A}^{(s)}_x$.

Step 4: Splitting of diagonal triadic fibres. By definition of the triadic fibres (Sections 4.10 and 4.12), $\mathcal{F}^{\mathrm{dist}}_{x,\mathrm{diag}}$ (resp. $\mathcal{F}^{\mathrm{time}}_{x,\mathrm{diag}}$) is the closure of diagonal observables/functions contributing to the diagonal part of the distance (resp. time) functionals, with the Holo-State pairing $\langle\!\langle \cdot, \cdot \rangle\!\rangle_\Psi$. Restricting the pairing to the two blocks and using $P_0(x)P_s(x) = 0$ one checks

$$\langle\!\langle A_0,\ A_s \rangle\!\rangle_\Psi \ = \ 0 \quad \text{for all } A_0 \in \mathsf{A}^{(0)}_x,\ A_s \in \mathsf{A}^{(s)}_x,$$

because each mixed product factors through $P_0(x)P_s(x)$ or $P_s(x)P_0(x)$ and vanishes by typing. Hence the diagonal fibre decomposes orthogonally as

$$\mathcal{F}^{\mathrm{dist}}_{x,\mathrm{diag}} \ \cong \ \mathcal{F}^{\mathrm{dist},(0)}_x \ \oplus \ \mathcal{F}^{\mathrm{dist},(s)}_x, \qquad \mathcal{F}^{\mathrm{time}}_{x,\mathrm{diag}} \ \cong \ \mathcal{F}^{\mathrm{time},(0)}_x \ \oplus \ \mathcal{F}^{\mathrm{time},(s)}_x,$$

where $\mathcal{F}^{\mathrm{dist},(\alpha)}_x$ and $\mathcal{F}^{\mathrm{time},(\alpha)}_x$ are the closures of $P_\alpha(x)\mathsf{A}_{x,\mathrm{diag}}P_\alpha(x)$ under the respective pairings ($\alpha \in \{0, s\}$).

Step 5: Refined diagonal contributions and reconstruction. Let Π_α be the block pinching onto the α-sector at x and x' (cf. the pinching map in Theorem 4.173, case (B3$^{\mathrm{word}}$)). Define the refined diagonal distances by applying the original diagonal functional (Definition 4.463 and eqs. (4.210) and (4.211)) to the pinched observables:

$$d^{(\alpha)}_{\mathrm{vac}}(x, x') := d_{\mathrm{vac}}\Big(\Pi_\alpha[\text{diag data at } (x, x')]\Big), \qquad \alpha \in \{0, s\}.$$

Likewise define $t^{(\alpha)}_{\mathrm{vac}}(x, x')$ from the diagonal time functional. Because the two sectors are orthogonal in the Holo-State pairing (above), the total diagonal contributions satisfy the Pythagorean additivity

$$d_{\mathrm{vac}}(x, x')^2 \ = \ \Big(d^{(0)}_{\mathrm{vac}}(x, x')\Big)^2 + \Big(d^{(s)}_{\mathrm{vac}}(x, x')\Big)^2, \qquad t_{\mathrm{vac}}(x, x')^2 \ = \ \Big(t^{(0)}_{\mathrm{vac}}(x, x')\Big)^2 + \Big(t^{(s)}_{\mathrm{vac}}(x, x')\Big)^2,$$

so their square-root sums recover the diagonal components of the triadic metric/time.

Step 6: Orthogonality to off-diagonal axes and gauge invariance. By Proposition 4.475, diagonal contributions are orthogonal (in the Holo-State pairing) to the correlator-based and spectral-shear parts; the refinement above keeps us inside the diagonal subspace, hence preserves this orthogonality. Under endpoint conjugation $X \mapsto g(x)\, X\, g(x')^\dagger$ (Proposition 4.54 and Corollary 4.64), the families $\{P_\alpha(x)\}$ and $\{\mathsf{O}_{\alpha\alpha}(x)\}$ transform by unitary conjugation; joint spectra and the induced quadratic functionals are invariant. Therefore the refined quantities and the direct-sum splittings are gauge-invariant.

Conclusion. We have (i) a joint spectral resolution for the commuting diagonal family, (ii) an orthogonal direct-sum decomposition of diagonal triadic fibres into 0 and s sectors, and (iii) refined diagonal distance/time components whose (Pythagorean) sums reconstruct the original diagonal parts, all gauge-invariant and orthogonal to the off-diagonal axes. □

Proposition 4.188 (ss-normal form for channel words)**.** *If central holosymmetry holds (Theorem 4.169), then every word $W \in \mathsf{G}$ admits a normal form*

$$W \;\sim\; \big(\hat{G}^{ss}_\Lambda\big)^m \;\cdot\; W_{\mathrm{red}} \quad \textit{with} \quad [\,\hat{G}^{ss}_\Lambda,\; W_{\mathrm{red}}\,] = 0,$$

where $m \geq 0$ and W_{red} contains no internal ss-letters. The factorization is unique up to endpoint gauge (Proposition 4.54) and commuting diagonal words.

Proof idea. Since $\hat{G}^{ss}_\Lambda$ commutes with all letters, slide each ss factor to one side using block composition (Proposition 4.85). Gauge conjugations preserve commutators and zero blocks, giving uniqueness modulo endpoint dressing.

Proof. Preliminaries. Work over the base $\mathcal{C}_d$ of Definition 4.617; words are evaluated on fibres of $\mathfrak{F}$ with (4.303) and transported by Section 4.19.1.

Setup and notation. Let

$$W = \hat{G}^{\alpha_n\alpha_{n-1}}_\Lambda(x_n, x_{n-1}) \cdots \hat{G}^{\alpha_1\alpha_0}_\Lambda(x_1, x_0)$$

be a typed channel word (Definition 4.141). Write $S(W) := \{j \in \{1, \dots, n\} : \alpha_j = \alpha_{j-1} = s\}$ for the set of ss-letters in W, and set $m := |S(W)|$.

Step 1: Existence by induction on m. We show that for every m there is a factorization

$$W = \Big(\prod_{j \in S(W)} \hat{G}^{ss}_\Lambda(x_j, x_{j-1}) \Big) \cdot W_{\mathrm{red}},$$

where W_{red} is a word with no ss-letters. Because $\hat{G}^{ss}_\Lambda$ is central (Theorem 4.169), the product of the ss-letters commutes with every factor; hence the order of the ss-letters in the left product is immaterial.

Base case $m = 0$. Then $S(W) = \varnothing$ and the claim holds with $W_{\mathrm{red}} = W$.

Inductive step. Assume the statement for all words with fewer than $m \geq 1$ ss-letters. Choose the leftmost ss-letter at position j, and decompose

$$W = A \cdot \hat{G}^{ss}_\Lambda(x_j, x_{j-1}) \cdot B,$$

with $A := \hat{G}^{\alpha_n\alpha_{n-1}} \cdots \hat{G}^{\alpha_{j+1}\alpha_j}$ and $B := \hat{G}^{\alpha_{j-1}\alpha_{j-2}} \cdots \hat{G}^{\alpha_1\alpha_0}$ (both possibly empty). By centrality, $[\,\hat{G}^{ss}_\Lambda, \cdot\,] = 0$ on G (Theorem 4.169), so

$$W = \hat{G}^{ss}_\Lambda(x_j, x_{j-1}) \cdot (A\,B).$$

By block composition through the intermediate locus x_j (Propositions 4.85 and 4.143 (iii)), the concatenation $A\,B$ is again a well-typed word with endpoints (x_n, x_0). Moreover $A\,B$ has exactly $m-1$ ss-letters. Apply the induction hypothesis to $A\,B$ to obtain

$$A\,B = \Big(\prod_{k \in S(A\,B)} \hat{G}^{ss}_\Lambda(\cdots) \Big) \cdot W_{\mathrm{red}}.$$

Combining the two displays gives the desired factorization for W, with the total left factor being the (commuting) product over all $j \in S(W)$, and W_{red} containing no ss-letters.

Step 2: Commutation with W_{red}. Since $\hat{G}^{ss}_\Lambda$ is central in G (Theorem 4.169), it commutes with every word, in particular with W_{red}. Thus $[\,\hat{G}^{ss}_\Lambda, W_{\mathrm{red}}\,] = 0$.

Step 3: Compressing the left factor to $\big(\hat{G}^{ss}_\Lambda\big)^m$. The left factor is a product of m ss-letters (with possibly different intermediate loci and paths), all commuting with each other and with W_{red}. We denote any such commuting m-fold product by $\big(\hat{G}^{ss}_\Lambda\big)^m$; because these elements lie in the (commutative) diagonal subalgebra generated by $0/s$-diagonals (Theorem 4.173, case ($\mathrm{B3}^{\mathrm{word}}$)), this notation is unambiguous up to multiplication by commuting diagonal words obtained from path/block recombinations (Proposition 4.85).

Step 4: Gauge covariance. Under endpoint gauge $X \mapsto g(x_n) X g(x_0)^\dagger$ (Proposition 4.54 and Corollary 4.64), both $\big(\hat{G}^{ss}_\Lambda\big)^m$ and W_{red} transform by conjugation. Hence the existence of the normal form and the commutation relation are gauge-invariant statements.

Step 5: Uniqueness up to endpoint gauge and commuting diagonal words. Suppose we have two normal forms

$$W = \big(\hat{G}^{ss}_\Lambda\big)^m W_{\mathrm{red}} = \big(\hat{G}^{ss}_\Lambda\big)^{m'} W'_{\mathrm{red}},$$

with $W_{\mathrm{red}}, W'_{\mathrm{red}}$ containing no ss-letters. Moving the central factors to the same side and cancelling yields

$$W_{\mathrm{red}} \;=\; D \;\cdot\; W'_{\mathrm{red}}, \qquad D := \big(\hat{G}^{ss}_\Lambda\big)^{m'-m}.$$

By construction, D belongs to the (commutative) diagonal $0/s$ subalgebra (Theorem 4.173, case ($\mathrm{B3}^{\mathrm{word}}$)) and commutes with both W_{red} and W'_{red}. Moreover, any further ambiguity from regrouping paths/blocks amounts to inserting endpoint conjugations (gauge) and multiplying by diagonal words (again commuting) (Propositions 4.54 and 4.85). Therefore the normal form is unique up to endpoint gauge and commuting diagonal words.

Conclusion. Every word admits the stated factorization with commuting $\hat{G}^{ss}_\Lambda$-power on the left and an ss-free remainder, unique modulo endpoint gauge and diagonal commuting factors. □

Definition 4.189 (p/s-cohomology and invariants)**.** Let $\mathcal{H}_p, \mathcal{H}_s$ be the p- and s-typed modal fibres at a locus. Define the two-term complex

$$0 \longrightarrow \mathcal{H}_p \xrightarrow{d_{ps} := \hat{G}^{ps}_\Lambda} \mathcal{H}_s \longrightarrow 0.$$

Under exact nilpotency (Theorem 4.170), $d_{ps}^2 = 0$ and the cohomology groups

$$H_{ps}^0 \;:=\; \ker(d_{ps}), \qquad H_{ps}^1 \;:=\; \mathcal{H}_s/\mathrm{im}(d_{ps})$$

are well-defined, gauge-invariant under endpoint conjugation, and stable under block composition. Their dimensions and induced pairings are *robust invariants* of synthesis–holothesis coupling.

Proposition 4.190 (Stability under small nilpotency defect)**.** *With nilpotency defect η (Definition 4.172), let $d_{ps} = \hat{G}_\Lambda^{ps} : \mathcal{H}_p \to \mathcal{H}_s$ and fix a tolerance $\tau > \sqrt{\eta}$. Define τ-pseudokernels/images by*

$$\ker_\tau(d_{ps}) = \{v \in \mathcal{H}_p :\ \|d_{ps}v\| \le \tau\|v\|\}, \quad \mathrm{im}_\tau(d_{ps}) = \{d_{ps}u + \delta :\ \|\delta\| \le \tau\|u\|\}.$$

Then the quotient $\widetilde{H}_{ps}^1(\tau) := \mathcal{H}_s/\mathrm{im}_\tau(d_{ps})$ and $\ker_\tau(d_{ps})$ are well-defined and their dimensions are constant for all $\tau \in (\sqrt{\eta}, \tau_0]$. In particular, the exact cohomology is recovered as $\eta \to 0$.

Proof. Preliminaries. Work in finite-dimensional Hilbert spaces $\mathcal{H}_p, \mathcal{H}_s$ (relevant fibres of the words calculus). Take the singular value decomposition (SVD) $d_{ps} = U\Sigma V^\dagger$ with singular values $s_1 \ge \cdots \ge s_r > 0$ and the rest zero. Let $\{v_i\} \subset \mathcal{H}_p$ (right singular vectors) and $\{u_i\} \subset \mathcal{H}_s$ (left singular vectors) be orthonormal bases with $d_{ps}v_i = s_i u_i$ for $1 \le i \le r$, and $d_{ps}v_i = 0$ for $i > r$.

1) Structure of the τ-pseudokernel. Write any $v = \sum_i a_i v_i$. Then $\|d_{ps}v\|^2 = \sum_i s_i^2|a_i|^2$ and $\|v\|^2 = \sum_i |a_i|^2$. Thus $v \in \ker_\tau(d_{ps})$ iff $\sum_i (s_i^2 - \tau^2)|a_i|^2 \le 0$. Because each term is ≥ 0 when $s_i > \tau$ and ≤ 0 when $s_i \le \tau$, this inequality forces $a_i = 0$ for all i with $s_i > \tau$. Conversely, any linear combination of v_i with $s_i \le \tau$ satisfies the inequality. Hence

$$\ker_\tau(d_{ps}) \;=\; \mathrm{span}\{\, v_i :\ s_i \le \tau \,\},$$

and $\dim \ker_\tau(d_{ps})$ equals the counting function $N_\le(\tau) := \#\{i :\ s_i \le \tau\}$. This dimension is constant for τ in any interval avoiding singular values.

2) Structure of the τ-pseudoimage (with the current definition). Given $y \in \mathcal{H}_s$, expand $y = \sum_i b_i u_i$. We need $y = d_{ps}u + \delta$ with $\|\delta\| \le \tau\|u\|$. Let $u = \sum_i a_i v_i$. Then $d_{ps}u = \sum_i s_i a_i u_i$ and $\delta = \sum_i (b_i - s_i a_i)u_i$. The constraint reads $\sum_i |b_i - s_i a_i|^2 \le \tau^2 \sum_i |a_i|^2$. This decouples coordinatewise and is satisfiable for *every* choice of $\{b_i\}$: for each i, pick

$$a_i \;=\; \frac{b_i}{s_i + \tau} \quad (s_i + \tau > 0),$$

which gives $|b_i - s_i a_i| = \frac{\tau}{s_i+\tau}|b_i| \le \tau|a_i|$. Summing over i yields the required inequality. Therefore

$$\mathrm{im}_\tau(d_{ps}) \;=\; \mathcal{H}_s \qquad \text{for every } \tau > 0.$$

Consequently, $\widetilde{H}_{ps}^1(\tau) = \mathcal{H}_s/\mathrm{im}_\tau(d_{ps}) = \{0\}$ and $\dim \widetilde{H}_{ps}^1(\tau) = 0$, a constant.

3) Constancy on $(\sqrt{\eta}, \tau_0]$. Because $\dim \ker_\tau(d_{ps})$ is the counting function $N_\le(\tau)$ of singular values, it is piecewise constant and can change only when τ crosses a singular value. Pick $\tau_0 > \sqrt{\eta}$ that does not coincide with any singular value and (if desired) lies below the next

larger singular value; then $\dim \ker_\tau(d_{ps})$ is constant for all $\tau \in (\sqrt{\eta}, \tau_0]$. The dimension of $\widetilde{H}^1_{ps}(\tau)$ is 0 for all $\tau > 0$, hence constant on that interval.

4) Gauge covariance and recovery in the exact limit. Under endpoint conjugations (gauge) $d_{ps} \mapsto g_s\, d_{ps}\, g_p^\dagger$ (Proposition 4.54 and Corollary 4.64), singular values are unchanged, so both constructions are gauge-invariant.

In the exact nilpotent case ($\eta = 0,\ d_{ps}^2 = 0$), we can take $\tau \downarrow 0$: then $\ker_\tau(d_{ps}) \downarrow \ker(d_{ps})$, and $\mathrm{im}_\tau(d_{ps})$ (as defined here) remains $\mathcal{H}_s$, so $\widetilde{H}^1_{ps}(\tau)$ collapses to 0. (See the remark below for a refinement that yields a nontrivial, stable H^1.) □

Remark 4.191 (Why the absolute pseudoimage is too permissive)**.** With the absolute-residual definition

$$\mathrm{im}_\tau(d) \;:=\; \Big\{\, du + \delta \;:\; \|\delta\| \le \tau\, \|u\| \Big\},$$

one always has $\mathrm{im}_\tau(d) = \mathcal{H}_s$ for every $\tau > 0$. Intuitively, the tolerance scales with $\|u\|$, so one can absorb any desired residual δ by rescaling u; more precisely, an SVD calculation shows every $y \in \mathcal{H}_s$ admits a decomposition $y = du + \delta$ with $\|\delta\| \le \tau\|u\|$. Consequently the quotient $\mathcal{H}_s/\mathrm{im}_\tau(d)$ is always trivial.

A nontrivial and stable alternative is to measure the residual *relative to the output*, and define the *relative pseudoimage*

$$\mathrm{im}^{\mathrm{rel}}_\tau(d) \;:=\; \Big\{\, du + \delta \;:\; \|\delta\| \le \tau\, \|du\| \Big\}.$$

Equivalently (and more spectrally transparent),

$$\mathrm{im}^{\mathrm{spec}}_\tau(d) \;:=\; \mathrm{Ran}\, \mathbf{1}_{(\tau,\infty)}\Big(\sqrt{dd^\dagger}\Big),$$

and $\mathrm{im}^{\mathrm{rel}}_\tau(d)$ has the same closure as $\mathrm{im}^{\mathrm{spec}}_\tau(d)$. Let $d = U\Sigma V^\dagger$ be an SVD with singular values (s_i), right singular vectors (v_i), and left singular vectors (u_i). Then (see Lemma 4.193):

$$\ker_\tau(d) = \mathrm{span}\{\, v_i :\ s_i \le \tau \,\}, \qquad \overline{\mathrm{im}^{\mathrm{rel}}_\tau(d)} = \overline{\mathrm{im}^{\mathrm{spec}}_\tau(d)} = \mathrm{span}\{\, u_i :\ s_i > \tau \,\},$$

so that

$$\dim \ker_\tau(d) = \Big|\{\, i :\ s_i \le \tau \,\}\Big|, \qquad \dim\Big(\mathcal{H}_s \Big/ \overline{\mathrm{im}^{\mathrm{rel}}_\tau(d)}\Big) = \Big|\{\, i :\ s_i \le \tau \,\}\Big|.$$

Both dimensions are constant for τ varying in any interval that avoids singular values. In the exact limit, taking $\tau \downarrow 0$ (with $\tau > \sqrt{\eta}$ and $\eta \to 0$) recovers the usual cohomology: $\ker(d)$ and $\mathcal{H}_s/\mathrm{im}(d)$. Here $|\cdot|$ denotes cardinality (possibly $+\infty$).

Definition 4.192 (Spectral pseudoimage and pseudokernel)**.** Let $d_{ps} = \hat{G}^{ps}_\Lambda : \mathcal{H}_p \to \mathcal{H}_s$ and let $\sqrt{d_{ps} d_{ps}^\dagger}$ have singular values $(s_i)_{i=1}^r > 0$ (the rest $= 0$). For $\tau \ge 0$ define

$$\ker_\tau(d_{ps}) := \{v \in \mathcal{H}_p :\ \|d_{ps} v\| \le \tau\, \|v\|\}, \qquad \mathrm{im}^{\mathrm{spec}}_\tau(d_{ps}) := \mathrm{Ran}\, \mathbf{1}_{(\tau,\infty)}\Big(\sqrt{d_{ps} d_{ps}^\dagger}\Big).$$

Lemma 4.193 (SVD characterization)**.** *Let $d_{ps} = U\Sigma V^\dagger$ be an SVD with right singular vectors $\{v_i\} \subset \mathcal{H}_p$ and left singular vectors $\{u_i\} \subset \mathcal{H}_s$, so that $d_{ps} v_i = s_i u_i$, $d_{ps}^\dagger u_i = s_i v_i$ and $s_1 \ge \cdots \ge s_r > 0$. Then for every $\tau \ge 0$,*

$$\ker_\tau(d_{ps}) = \mathrm{span}\{\, v_i :\ s_i \le \tau \,\}, \qquad \mathrm{im}^{\mathrm{spec}}_\tau(d_{ps}) = \mathrm{span}\{\, u_i :\ s_i > \tau \,\}.$$

Proof. Write $v = \sum_i a_i v_i$ with $\|v\|^2 = \sum_i |a_i|^2$ and $\|d_{ps}v\|^2 = \sum_i s_i^2 |a_i|^2$. Then $\|d_{ps}v\| \le \tau\|v\|$ iff $\sum_i (s_i^2 - \tau^2)|a_i|^2 \le 0$, which forces $a_i = 0$ for $s_i > \tau$ and allows arbitrary a_i for $s_i \le \tau$; hence the first identity. The second follows from the spectral projector definition: $\operatorname{Ran} \mathbf{1}_{(\tau,\infty)}(\sqrt{d_{ps}d_{ps}^\dagger}) = \operatorname{span}\{u_i : \ s_i > \tau\}$. □

Proposition 4.194 (Stability of spectral pseudo-cohomology)**.** *Let $d_{ps} = \hat{G}_\Lambda^{ps} : \mathcal{H}_p \to \mathcal{H}_s$ and let $\eta = \|d_{ps}^2\|$ be the nilpotency defect* (Definition 4.172)*. Fix $\tau > 0$ and define*

$$\widetilde{H}_{ps}^1(\tau) \ := \ \mathcal{H}_s \Big/ \operatorname{im}_\tau^{\mathrm{spec}}(d_{ps}).$$

Then:

(a) Piecewise-constant ranks

With $N_\le(\tau) := \#\{i : \ s_i \le \tau\}$ and $N_>(\tau) := \#\{i : \ s_i > \tau\}$,

$$\dim \ker_\tau(d_{ps}) = N_\le(\tau), \qquad \dim \widetilde{H}_{ps}^1(\tau) = N_\le(\tau),$$

and these dimensions are constant for τ varying in any interval that contains no singular value.

(b) Gauge invariance

Under endpoint conjugations $d_{ps} \mapsto g_s \, d_{ps} \, g_p^\dagger$ (Proposition 4.54 and Corollary 4.64)*, the multisets $\{s_i\}$, the subspaces $\ker_\tau(d_{ps})$, $\operatorname{im}_\tau^{\mathrm{spec}}(d_{ps})$, and the quotient $\widetilde{H}_{ps}^1(\tau)$ are unchanged up to unitary equivalence.*

(c) Exact limit

If $\eta = 0$ (exact nilpotency), then as $\tau \downarrow 0$,

$$\ker_\tau(d_{ps}) \ \downarrow \ \ker(d_{ps}), \qquad \operatorname{im}_\tau^{\mathrm{spec}}(d_{ps}) \ \uparrow \ \operatorname{im}(d_{ps}),$$

and thus $\dim \widetilde{H}_{ps}^1(\tau) \to \dim \big(\mathcal{H}_s / \operatorname{im}(d_{ps})\big)$.

Moreover, if you choose τ in any open interval disjoint from the singular values (e.g. $\tau \in (\sqrt{\eta}, \tau_0]$ with τ_0 not a singular value), the ranks in (a) are constant throughout that interval.

Proof. Preliminaries. Endpoints are in $\mathcal{C}_d$; $d_{ps} = \hat{G}_\Lambda^{ps}$ acts between the spectrally restricted fibres of $\mathfrak{F}$ at those endpoints ((4.303)); transport/gauge per Section 4.19.1.

(a) The identities for the dimensions follow immediately from Lemma 4.193. Piecewise constancy is the observation that $N_\le(\tau)$ and $N_>(\tau)$ can jump only when τ crosses a singular value.

(b) Gauge conjugations are unitary on domain and codomain, so they preserve $d_{ps}d_{ps}^\dagger$, hence the spectrum of $\sqrt{d_{ps}d_{ps}^\dagger}$, the SVD singular values, and the spectral subspaces defined by projectors $\mathbf{1}_{(\tau,\infty)}(\cdot)$ and $\mathbf{1}_{[0,\tau]}(\cdot)$. Therefore the two subspaces and the quotient are invariant up to unitary.

(c) If $\eta = 0$, then $d_{ps}^2 = 0$ and (finite-dimensional) $\operatorname{im}(d_{ps})$ is closed. For $\tau \downarrow 0$, $\mathbf{1}_{(\tau,\infty)}(\sqrt{d_{ps}d_{ps}^\dagger})$ increases strongly to $\mathbf{1}_{(0,\infty)}(\cdot)$, whose range is $\overline{\operatorname{im}(d_{ps})} = \operatorname{im}(d_{ps})$; likewise $\ker_\tau(d_{ps})$ decreases to $\ker(d_{ps})$. The rank limits yield the exact cohomology ranks. □

Remark 4.195 (Why we mention $\tau > \sqrt{\eta}$)**.** The SVD/spectral statements above do *not* require any relation between τ and η; they hold for every $\tau \geq 0$. Mentioning $\tau > \sqrt{\eta}$ ties this threshold to your "almost differential" regime and ensures you can choose a *single* interval (e.g. $(\sqrt{\eta}, \tau_0]$) that avoids singular values in practical scenarios, so the ranks are constant on that operational window.

Remark 4.196 (Intuition for spectral pseudo-cohomology)**.** In the SVD picture of d_{ps} (Lemma 4.193 and Proposition 4.194):

- $\ker_\tau(d_{ps})$ collects *domain directions* mapped with gain $\leq \tau$: the **soft modes**.
- $\mathrm{im}_\tau^{\mathrm{spec}}(d_{ps})$ collects *output directions* with singular value $> \tau$: the **hard image**.
- Both sides count the same number of modes at threshold τ, namely those with $s_i \leq \tau$. Hence
$$\dim \ker_\tau(d_{ps}) = \dim\left(\mathcal{H}_s/\mathrm{im}_\tau^{\mathrm{spec}}(d_{ps})\right),$$
and the pseudo-cohomology rank is *piecewise constant* as τ varies away from singular values (Proposition 4.194(a)).
- As $\tau \downarrow 0$ and $\eta \to 0$ (exact nilpotency), one recovers the usual objects:
$$\ker_\tau(d_{ps}) \downarrow \ker(d_{ps}), \qquad \mathrm{im}_\tau^{\mathrm{spec}}(d_{ps}) \uparrow \mathrm{im}(d_{ps}),$$
so $\widetilde{H}^1_{ps}(\tau) \to \mathcal{H}_s/\mathrm{im}(d_{ps})$ (Proposition 4.194(c), Remark 4.195).

This resolves the "too-permissive" absolute pseudoimage (Remark 4.191) and yields a crisp, gauge-invariant, *spectral* notion of *ps*-pseudo-cohomology; gauge covariance follows from endpoint conjugation invariance (Proposition 4.54 and Corollary 4.64).

Corollary 4.197 (Geometric packaging in the fibre)**.** *The structures of Propositions 4.187 and 4.188 and Definition 4.189 assemble over $\mathcal{C}_d$ into bundle data:*

(i) *a bi-diagonal refinement of the diagonal sub-fibres inside $\mathcal{F}^{\mathrm{dist}}$ and $\mathcal{F}^{\mathrm{time}}$;*

(ii) *an ss-central normal-form sheaf on the channel-word bundle; and*

(iii) *a p/s-cohomology sheaf well-defined in the exact case and upper-semicontinuous under small η.*

Proof. emphPreliminaries. Work over $\mathcal{C}_d$ as in Definition 4.617; use $\pi : \mathfrak{F} \to \mathcal{C}_d$ ((4.302)) with (4.303) and the transporter of Section 4.19.1.

(i) Bi-diagonal refinement of diagonal fibres. For each locus x, Proposition 4.187 yields commuting diagonal families and orthogonal central idempotents $P_0(x), P_s(x)$ on the diagonal fibre together with the direct sums
$$\mathcal{F}^{\mathrm{dist}}_{x,\mathrm{diag}} \cong \mathcal{F}^{\mathrm{dist},(0)}_x \oplus \mathcal{F}^{\mathrm{dist},(s)}_x, \qquad \mathcal{F}^{\mathrm{time}}_{x,\mathrm{diag}} \cong \mathcal{F}^{\mathrm{time},(0)}_x \oplus \mathcal{F}^{\mathrm{time},(s)}_x.$$
On any simply connected flat patch, the spectral projectors $P_\alpha(x)$ vary continuously by parallel transport (Propositions 4.54 and 4.59), hence their ranges form continuous subbundles; on

overlaps these bundles glue by endpoint gauge. This produces the desired bi-diagonal refinement bundles inside $\mathcal{F}^{\mathrm{dist}}$ and $\mathcal{F}^{\mathrm{time}}$, fibrewise orthogonal and gauge-invariant (orthogonality to non-diagonal axes is from Proposition 4.475).

(ii) *ss*-central normal-form sheaf. Let $\mathcal{W} \to \mathcal{C}_d$ be the channel-word bundle (fibre G_x). For an open set $U \subset \mathcal{C}_d$, define the presheaf

$$\mathcal{N}_{ss}(U) := \Big\{ \text{ sections } W \text{ on } U \Big/ \sim \Big\},$$

where $W \sim W'$ iff $W' = (\hat{G}^{ss}_{\Lambda})^m \cdot D \cdot W$ for some locally constant $m \geq 0$ and a commuting diagonal word D on U. By Proposition 4.188, each fibre $W(x)$ admits such a factorization and it is unique up to commuting diagonals and endpoint gauge, so local normal forms agree on overlaps; the usual equalizer condition yields sheaf gluing. Hence $\mathcal{N}_{ss}$ is a sheaf (of sets) encoding the *ss*-central normal form on $\mathcal{W}$.

(iii) p/s-cohomology (exact and upper-semicontinuous). In the exact case ($\eta = 0$), Definition 4.189 defines fibrewise $d_{ps}(x) = \hat{G}^{ps}_{\Lambda}(x)$ with $d_{ps}(x)^2 = 0$, and cohomology groups

$$H^0_{ps}(x) = \ker d_{ps}(x), \qquad H^1_{ps}(x) = \mathcal{H}_s(x)/\mathrm{im}\, d_{ps}(x).$$

On flat patches, spectral projectors of $d_{ps}(x)d_{ps}(x)^\dagger$ vary continuously; thus $\ker d_{ps}$ and $\mathrm{im}\, d_{ps}$ define subbundles and H^1_{ps} forms a quotient bundle. These patch by gauge to a cohomology sheaf on $\mathcal{C}_d$.

For small defect $\eta > 0$, pick $\tau > \sqrt{\eta}$ that avoids singular values uniformly on a cover; define the spectral pseudokernel/image by

$$\ker_\tau(d_{ps}(x)) = \mathrm{span}\{v_i(x) : s_i(x) \leq \tau\}, \quad \mathrm{im}^{\mathrm{spec}}_\tau(d_{ps}(x)) = \mathrm{span}\{u_i(x) : s_i(x) > \tau\},$$

and the pseudo-cohomology $\widetilde{H}^1_{ps}(\tau; x) := \mathcal{H}_s(x)/\mathrm{im}^{\mathrm{spec}}_\tau(d_{ps}(x))$. By Proposition 4.194, the ranks $\dim \ker_\tau$ and $\dim \widetilde{H}^1_{ps}(\tau)$ are piecewise constant and remain constant on any open set where no singular value crosses τ; hence the resulting fields are *upper-semicontinuous* bundles, which glue to a sheaf that limits to the exact cohomology sheaf as $\eta \to 0$.

Conclusion. Items (i)–(iii) produce, respectively, a bi-diagonal refinement of diagonal fibres, an *ss*-normal-form sheaf on channel words, and a (pseudo-)cohomology sheaf that is exact for $\eta = 0$ and upper-semicontinuous for small $\eta > 0$, all natural and gauge-invariant over $\mathcal{C}_d$. □

4.3 The Dialectic Super-Commutator

Definition 4.198 (Dialectic super-commutator)**.** Let the component channels carry the color (degree) $\deg(\hat{G}^{\alpha\beta}) := \deg(\alpha) + \deg(\beta) \in \mathbb{Z}_2 \times \mathbb{Z}_2$ (cf. Section 4.1.7). For any two components $\hat{G}^{\alpha\beta}$ and $\hat{G}^{\gamma\delta}$, define the graded bracket

$$\left[\hat{G}^{\alpha\beta}, \hat{G}^{\gamma\delta}\right]_{\mathrm{gr}} := \hat{G}^{\alpha\beta}\hat{G}^{\gamma\delta} - (-1)^{\langle \deg(\alpha)+\deg(\beta),\, \deg(\gamma)+\deg(\delta)\rangle}\hat{G}^{\gamma\delta}\hat{G}^{\alpha\beta}, \tag{4.71}$$

where $\langle (a_1, a_2), (b_1, b_2)\rangle := a_1 b_1 + a_2 b_2 \pmod 2$.

Remark 4.199 (Algebraic properties and selection rules)**.**

1. **Graded $*$-Lie structure.** The bracket $[\,\cdot\,,\,\cdot\,]_{\mathrm{gr}}$ is graded-skew, satisfies the graded Jacobi identity, and obeys $\big([X,Y]_{\mathrm{gr}}\big)^{\dagger} = -\,(-1)^{\langle \deg X, \deg Y\rangle}[X^{\dagger},Y^{\dagger}]_{\mathrm{gr}}$.
2. **Gauge covariance.** Under $g(x) \in \mathrm{U}(\mathcal{H}_{\mathrm{em}})$, $\hat{G}^{\alpha\beta}(x,x') \mapsto g(x)\hat{G}^{\alpha\beta}(x,x')g(x')^{\dagger}$; the bracket is preserved because degree, product, and adjoint are preserved (cf. Sections 4.1.2 and 4.1.4).
3. **Component-wise selection rule.** Let $c_1 := \deg(\hat{G}^{\alpha\beta})$, $c_2 := \deg(\hat{G}^{\gamma\delta})$. Then

$$\big[\hat{G}^{\alpha\beta},\hat{G}^{\gamma\delta}\big]_{\mathrm{gr}} = \begin{cases} [\hat{G}^{\alpha\beta},\hat{G}^{\gamma\delta}] & \text{if } \langle c_1,c_2\rangle = 0, \\ \{\hat{G}^{\alpha\beta},\hat{G}^{\gamma\delta}\} & \text{if } \langle c_1,c_2\rangle = 1, \end{cases}$$

   so the bracket reduces to an *ordinary commutator* unless the color dot-product is 1, in which case it becomes an *anticommutator*. Typical pairings (using $\deg(0) = (0,0)$, $\deg(1) = (1,0)$, $\deg(s) = (0,1)$, $\deg(p) = (1,1)$):

   - Even with anything $(0,0)\cdot(\cdot) = 0$: ordinary commutator.
   - Same odd color: $(1,0)$ with $(1,0)$ or $(0,1)$ with $(0,1)$: anticommutator.
   - Mixed with p: $(1,1)$ with $(1,0)$ or $(0,1)$: anticommutator.
   - Different odd colors $(1,0)$ with $(0,1)$: ordinary commutator.

Remark 4.200 (Physical postulates and analogies)**.** The grading fixes *which* bracket (commutator or anticommutator) is appropriate; dynamics enter through *postulates* that constrain their values. The analogies to bosons/fermions are algebraic, not ontic.

- **Commuting (bosonic-type) sector.** Symmetry–Holosymmetry self-channels are taken compatible: $[\hat{G}^{00},\hat{G}^{ss}] = 0$ (and, more generally, $[\hat{G}^{00},\cdot]$ is always an ordinary commutator since $\deg \hat{G}^{00} = (0,0)$).
- **Anticommuting (fermionic-type) mixing.** Choose a pair with color dot-product = 1; e.g.

$$\{\hat{G}^{10},\hat{G}^{p0}\} = 0,$$

  since $\deg(\hat{G}^{10}) = (1,0)$, $\deg(\hat{G}^{p0}) = (1,1)$, and $(1,0)\cdot(1,1) = 1$. This encodes a generalized exclusion between the transitions $0 \leftarrow 1$ and $0 \leftarrow p$.
- **Graded nilpotency (one-step constraints).** Postulates like $(\hat{G}^{ps})^2 = 0$ and $(\hat{G}^{sp})^2 = 0$ implement terminal pathways along the p–s edge (no coherent piling-up of successive $s \to p$ or $p \to s$ conversions).

These constraints function as principled "compatibility/exclusion rules'' for the ontological modes, derived from the color structure rather than imposed case by case.

4.3.1 Endogenous Super-Commutator for Holosymmetric Words

Definition 4.201 (Holosymmetric involutions and characters)**.** Let $\mathcal{H}$ be the inner automorphism group generated by (i) control-basis rotations on the control qubit and (ii) the branch-swap σ interchanging $A_G B_N \leftrightarrow A_N B_G$ (Definition 4.148, Corollary 4.168). Fix two involutive automorphisms:

$$\mathrm{Ad}_Z : X \mapsto (Z\otimes I)^\dagger X (Z\otimes I), \qquad \mathrm{Ad}_\sigma : X \mapsto \sigma(X),$$

where Z is the π rotation on the control (so that sum/difference interferometers are $\pm$ under Ad_Z and/or Ad_σ; cf. Corollary 4.168). For any holosymmetric word $W \in \mathsf{G}$, define its *endogenous charge* $q(W) = (q_c(W), q_b(W)) \in \mathbb{Z}_2^2$ by the eigenvalue equations

$$\mathrm{Ad}_Z(W) = (-1)^{q_c(W)} W, \qquad \mathrm{Ad}_\sigma(W) = (-1)^{q_b(W)} W,$$

whenever W lies in a joint eigenspace. In general, decompose W by the Reynolds projections $\Pi_{q_c,q_b} := \frac{1}{4}(I + (-1)^{q_c}\mathrm{Ad}_Z)(I + (-1)^{q_b}\mathrm{Ad}_\sigma)$ so that $W = \sum_{q\in\mathbb{Z}_2^2} W^{(q)}$ with $W^{(q)} := \Pi_q(W)$.

Remark 4.202 (Agreement with channel letters)**.** For channel letters $\hat{G}_\Lambda^{\alpha\beta}$ (Definition 4.141), the holosymmetric actions are diagonal on the fourfold modal basis; hence each letter is a joint eigenvector of $(\mathrm{Ad}_Z, \mathrm{Ad}_\sigma)$, and

$$q\big(\hat{G}_\Lambda^{\alpha\beta}\big) = \deg(\alpha) + \deg(\beta) \in \mathbb{Z}_2^2$$

(compatibly with Lemma 4.145 and Corollary 4.144). Thus the endogenous charge coincides with the traditional color on letters and, by functoriality, on typed words.

Definition 4.203 (Endogenous super-commutator)**.** For $A, B \in \mathsf{G}$, write their charge components $A = \sum_q A^{(q)}$, $B = \sum_{q'} B^{(q')}$ as in Definition 4.201. Define the *endogenous super-commutator* by

$$[A,B]_{\mathrm{endo}} := \sum_{q,q'\in\mathbb{Z}_2^2} \Big(A^{(q)} B^{(q')} \;-\; (-1)^{\langle q,q'\rangle}\, B^{(q')} A^{(q)} \Big), \tag{4.72}$$

$$\langle (a_1,a_2),(b_1,b_2)\rangle := a_1 b_1 + a_2 b_2 \pmod 2.$$

Proposition 4.204 (Graded $*$-Lie structure and covariance)**.** *The bracket* $[\,\cdot\,,\,\cdot\,]_{\mathrm{endo}}$ *satisfies:*

(a) Graded skew-symmetry and Jacobi
On joint eigenspaces it coincides with the standard $\mathbb{Z}_2^2$*-graded commutator; hence graded skew and the graded Jacobi identity hold, and by bilinearity they extend to all of* G*.*

(b) $*$-compatibility
$\big([A,B]_{\mathrm{endo}}\big)^\dagger = -\,(-1)^{\langle q(A),q(B)\rangle}\,[A^\dagger, B^\dagger]_{\mathrm{endo}}$*, with* $q(\cdot)$ *read componentwise.*

(c) Gauge covariance
Under endpoint gauge $X \mapsto g(x) X g(x')^\dagger$ *(Proposition 4.54 and Corollary 4.64),* Ad_Z *and* Ad_σ *act internally on the device factor, so charges and the bracket are preserved; thus* $[\,\cdot\,,\,\cdot\,]_{\mathrm{endo}}$ *is gauge-covariant.*

Corollary 4.205 (Selection rules and reduction)**.** *If A and B lie in joint eigenspaces with charges $q(A), q(B) \in \mathbb{Z}_2^2$, then*

$$[A,B]_{\mathrm{endo}} = \begin{cases} [A,B] & \text{if } \langle q(A), q(B)\rangle = 0, \\ \{A,B\} & \text{if } \langle q(A), q(B)\rangle = 1. \end{cases}$$

In particular, $q = (0,0)$ ("even" under both Z and σ) yields the ordinary commutator with every component. For letters $\hat{G}^{\alpha\beta}$ this reproduces Definition 4.198 and Remark 4.199 by Remark 4.202.

Remark 4.206 (Endogenous vs. graded: overlay principle)**.** The endogenous bracket uses only holosymmetric symmetries (Z, σ) and word functoriality (Definition 4.148, Proposition 4.167, and Corollary 4.168). When one *chooses* to identify $q(\cdot)$ with the bookkeeping degree $\deg(\cdot) \in \mathbb{Z}_2^2$ (Definition 4.94), one recovers the traditional graded bracket. Thus grading is an optional overlay; the sign rule is already fixed by endogeneous symmetry charges.

Remark 4.207 (Approximate symmetries and defect control)**.** If Ad_Z or Ad_σ hold only approximately (e.g. via the Reynolds projection of Theorem 4.169 with tolerance), the projections Π_q pick eigenspaces up to a defect measured by the compliance components (Δ, η, ζ) (Definition 4.178). Then

$$\big\| [A,B]_{\mathrm{endo}} - [A,B]^{\mathrm{ideal}}_{\mathrm{endo}} \big\| \;\le\; C \cdot \|\mathfrak{K}\| \, \|A\| \, \|B\|,$$

for a constant C dependent on the averaging window; in the perfect limit $\mathfrak{K} \to 0$ the ideal graded $*$-Lie structure is recovered.

Lemma 4.208 (Endogenous charges of channel letters)**.** *Let $\mathrm{Ad}_Z(X) = (Z \otimes I)^\dagger X (Z \otimes I)$ and $\mathrm{Ad}_\sigma(X) = \sigma(X)$ be the holosymmetric involutions on words (Definition 4.201), and let the fourfold selectors $\hat{\mathcal{F}}_\alpha(y)$ ($\alpha \in \{0,1,s,p\}$) be as in the words calculus (Definition 4.141, §1.2.9). Then for each α there is a charge $q(\alpha) = (q_c(\alpha), q_b(\alpha)) \in \mathbb{Z}_2^2$ with*

$$\mathrm{Ad}_Z\big(\hat{\mathcal{F}}_\alpha(y)\big) = (-1)^{q_c(\alpha)}\, \hat{\mathcal{F}}_\alpha(y), \qquad \mathrm{Ad}_\sigma\big(\hat{\mathcal{F}}_\alpha(y)\big) = (-1)^{q_b(\alpha)}\, \hat{\mathcal{F}}_\alpha(y),$$

and these charges coincide with the dialectic degrees:

$$q(\alpha) = \deg(\alpha) \in \{(0,0), (1,0), (0,1), (1,1)\}.$$

Consequently, for the channel letter $\hat{G}^{\alpha\beta}_\Lambda(x,x') = \hat{\mathcal{F}}^\dagger_\alpha(x)\, \mathcal{U}(x \leftarrow x'; \gamma)\, \hat{\mathcal{F}}_\beta(x')$ one has the joint eigen-relations

$$\mathrm{Ad}_Z\big(\hat{G}^{\alpha\beta}_\Lambda\big) = (-1)^{\,q_c(\alpha)+q_c(\beta)}\; \hat{G}^{\alpha\beta}_\Lambda,$$
$$\mathrm{Ad}_\sigma\big(\hat{G}^{\alpha\beta}_\Lambda\big) = (-1)^{\,q_b(\alpha)+q_b(\beta)}\; \hat{G}^{\alpha\beta}_\Lambda,$$

so the endogenous charge of a letter is

$$q\big(\hat{G}^{\alpha\beta}_\Lambda\big) = q(\alpha) + q(\beta) = \deg(\alpha) + \deg(\beta) \in \mathbb{Z}_2^2.$$

Proof. Step 1: Selectors are joint eigen-operators. In the fourfold modal basis fixed in §1.2.9, the holosymmetric involutions act diagonally on the device degrees of freedom: Ad_Z is the control π-rotation (computational Z) and Ad_σ is the branch-swap exchanging $A_G B_N \leftrightarrow A_N B_G$ (Definition 4.148 and Corollary 4.168). By construction, each endpoint selector $\hat{\mathcal{F}}_\alpha(y)$ projects onto a fixed fourfold modal sector and is therefore a joint eigen-operator of $(\mathrm{Ad}_Z, \mathrm{Ad}_\sigma)$ with eigenvalues ± 1. We write these eigenvalue parities as $q(\alpha) = (q_c(\alpha), q_b(\alpha))$. Matching with the dialectic grading (Definition 4.94) gives

$$q(0) = (0,0), \quad q(1) = (1,0), \quad q(s) = (0,1), \quad q(p) = (1,1),$$

i.e. $q(\alpha) = \deg(\alpha)$ (control parity on the first slot, branch parity on the second).

Step 2: Transport is neutral. The geometric transporter $\mathcal{U}(x \leftarrow x'; \gamma)$ acts on the fibre via the operational connection (Section 4.19.1) and does not involve the device involutions; hence it is neutral under both actions: $\mathrm{Ad}_Z(\mathcal{U}) = \mathcal{U}$ and $\mathrm{Ad}_\sigma(\mathcal{U}) = \mathcal{U}$.

Step 3: Letters inherit additive charges. Using multiplicativity of Ad_Z and Ad_σ and the adjoint relation $\mathrm{Ad}_\bullet(\hat{\mathcal{F}}_\alpha^\dagger) = \big(\mathrm{Ad}_\bullet(\hat{\mathcal{F}}_\alpha)\big)^\dagger$, we compute

$$\mathrm{Ad}_Z\big(\hat{G}_\Lambda^{\alpha\beta}\big) = (-1)^{q_c(\alpha)+q_c(\beta)}\, \hat{G}_\Lambda^{\alpha\beta}, \qquad \mathrm{Ad}_\sigma\big(\hat{G}_\Lambda^{\alpha\beta}\big) = (-1)^{q_b(\alpha)+q_b(\beta)}\, \hat{G}_\Lambda^{\alpha\beta}.$$

Thus $q(\hat{G}_\Lambda^{\alpha\beta}) = q(\alpha) + q(\beta)$, and with $q(\alpha) = \deg(\alpha)$ this equals $\deg(\alpha) + \deg(\beta)$, as claimed. □

Remark 4.209 (Endogenous physical postulates and analogies)**.** Within the endogenous super-commutator of Definition 4.203 and the charge assignment of Lemma 4.208:

- **Commuting (bosonic-type) sector.** Since $q(\hat{G}^{00}) = (0,0)$, Corollary 4.205 gives an ordinary commutator with every component. In particular $[\,\hat{G}^{00}, \hat{G}^{ss}\,] = 0$ by the words-level bosonic compatibility (Theorem 4.173).
- **Anticommuting (fermionic-type) mixing.** For letters with $\langle q(\hat{G}^{\alpha\beta}), q(\hat{G}^{\gamma\delta})\rangle = 1$ the bracket reduces to an anticommutator (Corollary 4.205). Example: $q(\hat{G}^{10}) = (1,0)$ and $q(\hat{G}^{p0}) = (1,1)$, so $\langle (1,0), (1,1)\rangle = 1$ and hence $\{\hat{G}^{10}, \hat{G}^{p0}\} = 0$ by the typing lemma below.
- **Graded nilpotency (one-step constraints).** The words-only nilpotency on the p–s edge, $(\hat{G}_\Lambda^{ps})^2 = (\hat{G}_\Lambda^{sp})^2 = 0$, was proven endogenously (no grading needed), cf. the words-version nilpotency theorem in the p–s section.

Thus, as in Remark 4.200, the *charges* fix *which* bracket applies, while the *values* are constrained by endogenous dynamics (centrality, bosonic compatibility, typing/orthogonality, and the p–s nilpotency).

Lemma 4.210 (Typed anticommutator vanishing: $\{\hat{G}^{10}, \hat{G}^{p0}\} = 0$)**.** *For fixed endpoints (x, x'), the channel letters $\hat{G}_\Lambda^{10}(x,x')$ and $\hat{G}_\Lambda^{p0}(x,x')$ satisfy*

$$\{\hat{G}_\Lambda^{10}(x,x'),\ \hat{G}_\Lambda^{p0}(x,x')\} = 0.$$

Proof. *By definition,* $\hat{G}^{10}_{\Lambda} = P_1(x)\,\hat{G}^{10}_{\Lambda}\,P_0(x')$ *and* $\hat{G}^{p0}_{\Lambda} = P_p(x)\,\hat{G}^{p0}_{\Lambda}\,P_0(x')$ *with orthogonal endpoint projectors* $P_1(x)P_p(x) = 0$ *(typing/selector orthogonality in Definition 4.141 and Lemma 4.145). Hence*

$$\hat{G}^{10}_{\Lambda}\hat{G}^{p0}_{\Lambda} = P_1(x)\,\hat{G}^{10}_{\Lambda}\,\underbrace{P_0(x')P_p(x)}_{=0}\,\hat{G}^{p0}_{\Lambda}\,P_0(x') = 0, \tag{4.73}$$

$$\hat{G}^{p0}_{\Lambda}\hat{G}^{10}_{\Lambda} = P_p(x)\,\hat{G}^{p0}_{\Lambda}\,\underbrace{P_0(x')P_1(x)}_{=0}\,\hat{G}^{10}_{\Lambda}\,P_0(x') = 0.$$

so their anticommutator vanishes. □

The intuition (super-commutators). A super-commutator is a compact *bookkeeping rule for interference signs.* Certain channel words behave "even" (reordering introduces no sign), while others are "odd" (reordering contributes a minus sign). When two odd contributions are swapped, the correct algebraic object is the *anticommutator*; otherwise it is the ordinary *commutator.* This sign choice is the algebraic proxy for which interference pattern—sum or difference—the setup realizes (cf. the selection rule in Corollary 4.205).

Where the sign actually comes from (endogenous view). No abstract grading needs to be postulated. The sign is *forced by two concrete holosymmetric involutions* acting on words: (i) the control π-flip Z (which toggles sum/difference interferometry), and (ii) the branch swap σ (which exchanges the two RR paths) (Definition 4.201 and Corollary 4.168). Every channel word W decomposes into joint eigensectors of these involutions; the resulting *endogenous charge* $q(W) = (q_c, q_b) \in \mathbb{Z}_2^2$ records whether W flips sign under Z and/or σ (Definition 4.201). When two words A, B are multiplied, their joint parity determines the reordering sign

$$(-1)^{\langle q(A),q(B)\rangle}, \qquad \langle (a_1,a_2),(b_1,b_2)\rangle = a_1b_1 + a_2b_2 \pmod 2,$$

and the *endogenous super-commutator*

$$[A,B]_{\text{endo}} \;=\; AB \;-\; (-1)^{\langle q(A),q(B)\rangle}BA$$

follows directly from symmetry (Definition 4.203). Identifying the charges with the usual modal colors recovers the textbook graded bracket (the *overlay principle*, Remark 4.206); the equality $q(\hat{G}^{\alpha\beta}) = \deg(\alpha) + \deg(\beta)$ is established in Lemma 4.208.

What this explains (selection rules → concrete dynamics).

- **Which bracket applies.** The charges alone decide *commutator vs. anticommutator* (Corollary 4.205); this is a kinematic, symmetry-level statement.
- **What the value is.** The actual numbers are fixed by endogenous dynamics and typing:
  - **Bosonic** $0/s$ **sector.** Since $q(\hat{G}^{00}) = (0,0)$, it commutes in the ordinary sense with every component; in particular, $[\hat{G}^{00}, \hat{G}^{ss}] = 0$ holds at the words level (Theorem 4.173).

– **"Fermionic-type" mixing.** If $\langle q(\hat{G}^{10}), q(\hat{G}^{p0})\rangle = 1$, the bracket reduces to an anticommutator; typing/orthogonality then forces $\{\hat{G}^{10}, \hat{G}^{p0}\} = 0$ (Lemma 4.210).

– $p \leftrightarrow s$ **edge.** The charge rule fixes the bracket type, while *nilpotency* $(\hat{G}^{ps})^2 = (\hat{G}^{sp})^2 = 0$ is proved purely from word composition and typing (no external grading required; see the words-version nilpotency theorem Theorem 4.170).

– **Central** ss**.** Holosymmetric invariance makes $\hat{G}^{ss}$ commute with all words (Theorem 4.169), yielding the *ss-normal form* $W \sim (\hat{G}^{ss})^m W_{\mathrm{red}}$ with W_{red} free of ss letters (Proposition 4.188).

Operational meaning (how it shows up in experiments).

- *Sum vs. difference interferometers* (control Z) and *branch-swap tests* directly resolve the two $\mathbb{Z}_2$ parities that constitute the charge q (Corollary 4.168 and Definition 4.201).
- *Commutation vs. anticommutation* manifests as *lensing equalities* and *destructive-interference constraints* in visibility envelopes predicted by the words calculus; departures in the near-ideal regime are controlled by the *compliance vector* $\mathfrak{K} = (\Delta, \eta, \zeta)$ (Definition 4.178) and propagate to measurable signatures via Proposition 4.181.

Why this matters.

- The graded sign rule becomes *intrinsic*: it is a consequence of holosymmetric device symmetries, not an external axiom (Definition 4.203 and Remark 4.206).
- The construction is *gauge-stable*: charges and brackets are preserved under endpoint conjugation and parallel transport (Propositions 4.54 and 4.204 and Corollary 4.64).
- The framework remains *operational*: the same symmetries that define the algebraic sign are precisely those implemented and read out by the holosymmetric words architecture (Definition 4.148 and Proposition 4.167).

In short, the super-commutator is the algebraic shadow of two endogenous $\mathbb{Z}_2$ symmetries—control flip and branch swap—acting on holosymmetric words. The words calculus fixes the sign through symmetry charges, dynamics and typing fix the values, and the resulting predictions are measurable and robust (Definitions 4.178 and 4.203, Corollary 4.205, Theorems 4.170 and 4.173, and Propositions 4.181 and 4.188).

4.4 The Centralizing Role of Holosymmetry

Postulate 4.4 (Centralizing role of Holosymmetry)**.** *The holosymmetric self–channel is* graded–central. *Since* $\deg(\hat{G}^{ss}) = \deg(s) + \deg(s) = (0,1) + (0,1) = (0,0)$*, the graded bracket reduces to the ordinary commutator. We postulate*

$$\left[\hat{G}^{ss}(x,x'),\, \hat{G}^{\alpha\beta}(y,y')\right] \;\approx\; 0 \qquad (\forall\, \alpha,\beta). \tag{4.74}$$

More precisely, the deviation from exact centrality is controlled by the modal curvature $\mathcal{F} := \mathrm{d}\mathcal{A} + \mathcal{A} \wedge \mathcal{A}$*:*

$$\left[\hat{G}^{ss}, \hat{G}^{\alpha\beta}\right] = \mathcal{O}\left(\int_{\gamma_{x\to x'}} \|\mathcal{F}\| + \int_{\gamma_{y\to y'}} \|\mathcal{F}\|\right) + \textit{(boundary terms)}. \tag{4.75}$$

Equality in (4.74) holds (i) in the coincident/flat limit $x' \to x$, $y' \to y$ *on contractible patches where* $\mathcal{F} \equiv 0$*, and (ii) when boundary terms vanish (e.g. under standard boundary conditions). Corrections are governed by nonlocal/holonomy effects along the relevant transport paths.*

Remark 4.211 (Meaning and immediate consequences)**.** Holosymmetry (*Holothesis*) is the implementation–invariant facet; centrality states that its self–channel acts as a *coherence scalar* for the whole Fourfold algebra. Algebraically:

1. **Superselection by spectrum.** By Schur's lemma, on each irreducible representation of the graded $*$-algebra, $\hat{G}^{ss}$ is scalar: $\hat{G}^{ss} = \lambda_s \mathbb{I}$. The spectrum $\sigma(\hat{G}^{ss})$ labels holosymmetric superselection sectors; every $\hat{G}^{\alpha\beta}$ preserves them.
2. **Block preservation.** If E_λ is the projector from the spectral resolution of $\hat{G}^{ss}$, then $[E_\lambda, \hat{G}^{\alpha\beta}] = 0$ and $E_\lambda \hat{G}^{\alpha\beta} E_\lambda$ yields a consistent sectorwise reduction of the Fourfold kernel.
3. **Covariance.** Centrality is invariant under unitary gauge transport $\hat{G}^{\alpha\beta} \mapsto g\,\hat{G}^{\alpha\beta} g^\dagger$ and under the graded bracket (Section 4.3), since $\deg \hat{G}^{ss} = (0,0)$.

Physically, $\hat{G}^{ss}$ is a fixed point for the algebraic flow of dualities: it constrains how other modes may mix without itself being altered by those mixings.

Field–space nesting: geometry–matter as a coupled matrix. Beyond a single foundational field, the relational kernel must also encode geometry–matter co–dependence. Introduce the *generalized field* column

$$\hat{\mathbf{F}}(x) := \begin{pmatrix} \hat{\mathbf{\Phi}}(x) \\ \hat{\mathbf{\Theta}}(x) \end{pmatrix},$$

and define the *field–space correlator* as a 2×2 operator–valued matrix

$$\hat{\mathbf{G}}_{\text{field}}(x,x') := \begin{pmatrix} \hat{G}_{\Phi\Phi}(x,x') & \hat{G}_{\Phi\Theta}(x,x') \\ \hat{G}_{\Theta\Phi}(x,x') & \hat{G}_{\Theta\Theta}(x,x') \end{pmatrix}, \qquad \hat{G}_{AB}(x,x') = \sum_{\alpha,\beta\in\{0,1,p,s\}} \hat{G}^{\alpha\beta}_{AB}(x,x')\,\mathbf{e}_{\alpha\beta}, \tag{4.76}$$

where each block $\hat{G}_{AB}$ (with $A, B \in \{\Phi, \Theta\}$) is itself a 4×4 matrix over the dialectic basis, and $\mathbf{e}_{\alpha\beta}$ denotes the canonical mode–matrix units. Thus the full kernel is a *matrix of matrices*: 2×2 in field space, each entry 4×4 in dialectic space.

Consistency checks.

- **Hermiticity & transport.** Each $\hat{G}^{\alpha\beta}_{AB}$ obeys the same Hermitian symmetry and gauge–covariant transport as in Sections 4.1.2 and 4.1.4.

- **Field–space rotations.** For any $W(x) \in \mathrm{U}(2)$, $\hat{\mathbf{F}} \mapsto W\hat{\mathbf{F}}$ induces $\hat{\mathbf{G}}_{\mathrm{field}}(x,x') \mapsto W(x)\,\hat{\mathbf{G}}_{\mathrm{field}}(x,x')\,W(x')^{\dagger}$; dialectic indices are untouched. Centrality of the holosymmetric self–channel extends componentwise: $[\hat{G}^{ss}_{AB},\, \hat{G}^{\alpha\beta}_{CD}] \approx 0$.
- **Sector reduction.** The spectral projectors of $\hat{G}^{ss}_{AB}$ commute with all $\hat{G}^{\alpha\beta}_{CD}$, enabling a sectorwise decomposition of the full geometry–matter kernel.

Remark 4.212 (From graded–central to *endogenously* central)**.**

1. **No color flip for $\hat{G}^{ss}$.** Since $q(\hat{G}^{ss}) = (0,0)$ by Lemma 4.208, the graded sign is trivial: the graded bracket equals the ordinary commutator for $\hat{G}^{ss}$ by Corollary 4.205 and Lemma 4.95.
2. **Postulate ⇒ theorem.** The centrality postulate (4.74) is recovered as a *words-level theorem*: $[\,\hat{G}^{ss}_{\Lambda},\, X\,] = 0$ for all $X \in \mathsf{G}$ under $(\mathrm{C1}^{\mathrm{word}})$–$(\mathrm{C3}^{\mathrm{word}})$ in Theorem 4.169. Thus "graded–central" becomes *endogenously central*, with the graded sign itself explained by the device involutions Z, σ (Section 4.3.1).

Proposition 4.213 (Curvature–controlled centrality defect)**.** *Let endpoints $x, x', y, y' \in \mathcal{C}_d$ and admissible paths $\gamma_{x\to x'}$, $\gamma_{y\to y'}$. There exists a constant $C > 0$ and a curvature–holonomy functional $\Phi(\gamma)$, controlled by the connection curvature $\mathcal{F} = \mathrm{d}\mathcal{A} + \mathcal{A}\wedge\mathcal{A}$, such that*

$$\Big\| [\hat{G}^{ss}_{\Lambda}(x,x'),\, \hat{G}^{\alpha\beta}_{\Lambda}(y,y')] \Big\| \;\le\; C\Big(\Phi(\gamma_{x\to x'}) + \Phi(\gamma_{y\to y'})\Big) \;+\; \text{boundary terms.} \tag{4.77}$$

The boundary terms vanish in the coincident/flat regime (contractible patches with $\mathcal{F} \equiv 0$, and trivial endpoint dressings), by Proposition 4.59 and Section 4.19.1. Consequently, the centrality defect Δ of the compliance vector $\mathfrak{K}$ satisfies

$$\Delta \;\lesssim\; \Phi(\gamma_{x\to x'}) + \Phi(\gamma_{y\to y'}),$$

i.e. centrality defects are holonomy defects*, refining the heuristic (4.75) in §4.4 and linking directly to Definition 4.178.*

Remark 4.214 (Immediate consequences, reinterpreted endogenously)**.**

1. **Superselection by spectrum.** By $\mathcal{H}$-invariance and Theorem 4.169, $\hat{G}^{ss}_{\Lambda}$ acts as a scalar on each typed Hom-space; its spectral projectors commute with all channel words, yielding sectorwise reductions exactly as in Remark 4.211.
2. **Block preservation and covariance.** Endpoint gauge covariance, functorial transport, and block composition $\big($Propositions 4.54, 4.85 and 4.143 and Corollary 4.64$\big)$ imply that centrality is gauge-stable and is preserved under word composition, matching the structural claims of §4.4.

Proposition 4.215 (Field–space nesting and componentwise centrality)**.** *Let the field–space correlator $\hat{\mathbf{G}}_{\mathrm{field}}$ be as in (4.76). Then the holosymmetric charges $q(\cdot)$ live in the device/dialectic factor and are independent of field labels. Hence, for each $A, B, C, D \in \{\Phi, \Theta\}$,*

$$\Big[\, \hat{G}^{ss}_{AB}(x,x'),\, \hat{G}^{\alpha\beta}_{CD}(y,y') \,\Big]_{\mathrm{endo}} = 0 \quad \textit{under } (\mathrm{C1}^{\mathrm{word}})\text{–}(\mathrm{C3}^{\mathrm{word}}), \tag{4.78}$$

with the same curvature–controlled defect bound (4.77). Moreover, for any $W(x) \in \mathrm{U}(2)$, the congruence action $\hat{\mathbf{G}}_{\text{field}} \mapsto W(x)\,\hat{\mathbf{G}}_{\text{field}}\,W(x')^{\dagger}$ preserves (4.78) (since it commutes with the device involutions Z, σ). One may package defects blockwise as Δ_{AB} and aggregate (e.g. $\|\Delta\|_{\text{field}} := \max_{A,B} \Delta_{AB}$), fitting into Definitions 4.178 and 4.183.

Remark 4.216 (Editorial annotations for §4.4)**.**

1. After Postulate 4.4, append: "*Words-level derivation via $\mathcal{H}$-invariance and conditional expectation:* Theorem 4.169."
2. Replace (4.75) by a named defect bound:

$$\left\| [\hat{G}^{ss}, \hat{G}^{\alpha\beta}] \right\| \;\le\; \Delta(x, x'; y, y') \qquad \text{with } \Delta \text{ controlled by curvature/holonomy functionals of } \mathcal{F}, \tag{4.79}$$

   citing Definition 4.178 and Proposition 4.59.
3. In the "Meaning and immediate consequences" list, change "graded $*$-algebra" to "words $*$-algebra with $\mathcal{H}$-action", and cite Theorem 4.169 and Proposition 4.188 for sector reduction and the ss-normal form.

The centralizing role of holosymmetry. The centralizing role of holosymmetry in §4.4 is not merely postulated: within the endogenous words framework it is *proved* (Theorem 4.169); its deviation is quantitatively controlled by holonomy/curvature (Definition 4.178, consistent with (4.75)); and the consequences—superselection, block preservation, and field–space covariance—follow from the same $\mathcal{H}$-invariance, block composition, and gauge–covariant transport that organize the holosymmetric alphabets and words (Propositions 4.54, 4.85 and 4.143 and Corollary 4.64).

Field–space nesting (endogenous words version). Beyond a single foundational field, geometry–matter co–dependence is encoded in a field–space matrix whose entries are *holosymmetric word–evaluated* Fourfold channels.

Definition 4.217 (Endogenous field–space correlator (words–dressed))**.** Let the device (holosymmetric) alphabet and RR matrix elements be as in Definition 4.148 and eq. (4.45), and let $\hat{\mathcal{F}}_{\alpha}^{[\mathbf{W};\,\chi_{\text{out}}\leftarrow\chi_{\text{in}}]}(x) := \mathsf{W}_{\chi_{\text{out}}\leftarrow\chi_{\text{in}}}\,\hat{\mathcal{F}}_{\alpha}(x)$ be the word–dressed selectors of Definition 4.166. For $A, B \in \{\Phi, \Theta\}$ and endpoints x, x', define the (block) field–space entries

$$\hat{G}_{AB}^{[\mathbf{W};\,\chi_{\text{out}}\leftarrow\chi_{\text{in}}]}(x, x') \;:=\; \sum_{\alpha,\beta\in\{0,1,p,s\}} \hat{G}_{AB}^{\alpha\beta}\Big(x, x'; \mathbf{W}, \chi_{\text{out}} \leftarrow \chi_{\text{in}}\Big)\, \mathbf{e}_{\alpha\beta}, \tag{4.80}$$

where each mode coefficient is the word–evaluated channel (cf. eq. (4.66))

$$\hat{G}_{AB}^{\alpha\beta}\Big(x, x'; \mathbf{W}, \chi_{\text{out}} \leftarrow \chi_{\text{in}}\Big) := \Big(\hat{\mathcal{F}}_{\alpha}^{[\mathbf{W};\,\chi_{\text{out}}\leftarrow\chi_{.in}]}(x)\Big)^{\dagger}\, \mathcal{U}(x \leftarrow x'; \gamma)\, \hat{\mathcal{F}}_{\beta}(x') \qquad \text{placed in the } (A, B) \text{ field slot.} \tag{4.81}$$

The endogenous field–space correlator is the 2×2 operator–valued matrix

$$\hat{\mathbf{G}}_{\text{field}}^{[\mathbf{W};\,\chi_{\text{out}}\leftarrow\chi_{\text{in}}]}(x,x') \;:=\; \begin{pmatrix} \hat{G}_{\Phi\Phi}^{[\mathbf{W};\,\chi_{\text{out}}\leftarrow\chi_{\text{in}}]}(x,x') & \hat{G}_{\Phi\Theta}^{[\mathbf{W};\,\chi_{\text{out}}\leftarrow\chi_{\text{in}}]}(x,x') \\ \hat{G}_{\Theta\Phi}^{[\mathbf{W};\,\chi_{\text{out}}\leftarrow\chi_{\text{in}}]}(x,x') & \hat{G}_{\Theta\Theta}^{[\mathbf{W};\,\chi_{\text{out}}\leftarrow\chi_{\text{in}}]}(x,x') \end{pmatrix}, \tag{4.82}$$

so the full kernel is a *matrix of matrices*: 2×2 in field space, each entry 4×4 in the dialectic basis $\{\mathbf{e}_{\alpha\beta}\}$ from Definition 4.141.

Proposition 4.218 (Basic properties)**.** *For* $\hat{\mathbf{G}}_{\text{field}}^{[\mathbf{W}]}$ *of* (4.82)*:*

Gauge covariance

By Propositions 4.54 and 4.167 and Corollary 4.64, each block transforms as $\hat{G}_{AB}^{[\mathbf{W}]}(x,x') \mapsto g(x)\,\hat{G}_{AB}^{[\mathbf{W}]}(x,x')\,g(x')^{\dagger}$.

Charge (endogenous grading)

The holosymmetric charges $q(\cdot)$ *reside in the device/dialectic factor and are independent of the field labels. Consequently, the componentwise endogenous centrality statements of Proposition 4.215 apply blockwise to* $\hat{\mathbf{G}}_{\text{field}}^{[\mathbf{W}]}$.

Reduction to the undressed form

For $\mathbf{W} = I$ *and* $\chi_{\text{out}} = \chi_{\text{in}}$, (4.82) *reduces to the original field–space correlator* $\hat{\mathbf{G}}_{\text{field}}(x,x')$ *in* (4.76).

Remark 4.219 (Charge resolution and compliance)**.** Applying the Reynolds projections Π_q of Definition 4.201 yields a charge–resolved decomposition $\hat{G}_{AB}^{[\mathbf{W}]} = \sum_{q\in\mathbb{Z}_2^2} \Pi_q\big(\hat{G}_{AB}^{[\mathbf{W}]}\big)$. Centrality and its curvature–controlled defect then lift componentwise to field space as in Proposition 4.213, and can be packaged blockwise (e.g. Δ_{AB}, aggregated via $\|\Delta\|_{\text{field}} := \max_{A,B}\Delta_{AB}$), consistent with Definitions 4.178 and 4.183.

Fourfold dynamics as a holosymmetric interferometer (endogenous reading). The Fourfold dynamics may be pictured as a many–port interferometer whose "beams"—Thesis (T), Antithesis (A), Synthesis (S), and Holothesis (H)—propagate through a shared device. In the endogenous calculus, this picture is made precise by the *channel alphabet and words* (Definition 4.141), together with holosymmetric device operations that realize sum/difference interferometry and branch exchange (Definition 4.148 and Corollary 4.168). Here T/A/S/H describe *traffic*—which modal branches are populated and whether they coexist coherently—while the holosymmetric actions specify the *wiring diagram* that steers and recombines those branches within words.

Central holosymmetry as a global coherence meter. The "coherence meter" of the apparatus is the holosymmetric self–channel, whose centralizing role is not merely postulated but *derived* in the words framework: under the endogenous hypotheses $(\mathrm{C1}^{\text{word}})$–$(\mathrm{C3}^{\text{word}})$ one has $[\,\hat{G}_{\Lambda}^{ss},\,X\,] = 0$ for all words $X \in \mathsf{G}$ (Theorem 4.169). Algebraically, centrality means commuting with everything; operationally, it means acting as a *coherence scalar* that multiplies processes without being distorted by their internal rearrangements (Remark 4.211). This endogenizes the intuition that the holosymmetric diagonal provides a fixed reference frame against which all other couplings unfold, and it underpins the *ss-normal form* $W \sim (\hat{G}^{ss})^m W_{\text{red}}$ (Proposition 4.188).

Sectors and local mixing. Because $\hat{G}^{ss}_{\Lambda}$ commutes with the entire words algebra, its spectral projectors commute with all channels, partitioning dynamics into *holosymmetric superselection sectors* (Remark 4.211 and Theorem 4.169). Within a fixed sector, the remaining modes (T/A/S/H) may mix and interfere according to the endogenous bracket and typing rules (Definition 4.203, Corollary 4.205, and Lemma 4.145); crossings between sectors, by contrast, correspond to genuine changes of the global architecture rather than local switches. In this sense, holosymmetry labels the "universe of implementation'' while leaving intra–sector dynamics intact.

When centrality fails: curvature and holonomy. Perfect centrality can fail only when global features of the transporter announce themselves. The words calculus quantifies this via a *curvature/holonomy* control:

$$\left\| [\hat{G}^{ss}_{\Lambda}(x,x'),\, \hat{G}^{\alpha\beta}_{\Lambda}(y,y')] \right\| \;\le\; C(\Phi(\gamma_{x\to x'}) + \Phi(\gamma_{y\to y'})) + \text{boundary terms},$$

where Φ is a holonomy functional determined by the curvature $\mathcal{F}$ (Proposition 4.213). Thus the "calibration tone'' wobbles only when transport explores nontrivial loops; in the flat/coincident limit the commutator vanishes exactly (Proposition 4.59). This matches the defect bookkeeping provided by the *holosymmetric compliance vector* $\mathfrak{K} = (\Delta, \eta, \zeta)$ (Definition 4.178).

The prism of interrogations: local facets vs. whole–system invariants. The Fourfold "prism'' cleanly separates: (i) *local resolution* (T/A occupancy and their interference), (ii) *coherent coexistence* (S–type contributions inside words), and (iii) *whole–system invariants* (the holosymmetric ss diagonal). Endogenous centrality guarantees that the reading of "structure at the whole'' is consistent regardless of which local facet is interrogated first (Theorem 4.169 and Proposition 4.188). Experimentally, the two $\mathbb{Z}_2$ involutions—control Z and branch swap σ—select the interference signs (charges) and thereby fix whether commutators or anticommutators are the correct comparisons (Definition 4.201 and Corollaries 4.168 and 4.205). Deviations from ideal behavior translate into bounded shifts in visibility/lensing diagnostics via the compliance map (Definition 4.178 and Proposition 4.181).

Intuitive summary. Holosymmetry is the fixed reference of the endogenous architecture—a coherence ledger for the entire mechanism. It constrains and normalizes other channels without being perturbed by their internal dynamics, except when genuinely global features (holonomy/curvature) intervene (Proposition 4.213). In the words calculus, this narrative is not merely metaphorical: it is encoded by centrality at the algebraic level (Theorem 4.169), sector decomposition (Remark 4.211), normal forms (Proposition 4.188), and symmetry–selected brackets (Definition 4.203 and Corollary 4.205), all organized by gauge–covariant transport and block composition (Propositions 4.54, 4.85 and 4.143 and Corollary 4.64).

4.5 Holosymmetric Wordwise Covariant Kinematics (Along a Path)

Fix a device-level word $\mathbf{W} \in \langle \mathsf{A}_{\mathrm{holo}} \rangle$ and a control transition $\chi_{\mathrm{in}} \to \chi_{\mathrm{out}}$. Recall the RR matrix element $\mathsf{W}_{\chi_{\mathrm{out}} \leftarrow \chi_{\mathrm{in}}}$ (Definition 4.148 and eq. (4.45)) and the corresponding *word-evaluated channel* components

$$\hat{G}^{\alpha\beta}_{\Lambda}(x, x'; \mathbf{W}, \chi_{\mathrm{out}} \leftarrow \chi_{\mathrm{in}}) = \Big(\hat{\mathcal{F}}^{[\mathbf{W};\, \chi_{\mathrm{out}} \leftarrow \chi_{\mathrm{in}}]}_{\alpha}(x)\Big)^{\dagger} \mathcal{U}(x \leftarrow x'; \gamma)\, \hat{\mathcal{F}}_{\beta}(x') \tag{4.83}$$

(see Definition 4.166 and eq. (4.66)).

Throughout we use the transport connection $\mathcal{A}$ ($\mathcal{U} = \mathcal{P}\exp\!\big(-\int_{\gamma} \mathcal{A}\big)$, Definition 4.48 and eqs. (4.11) and (4.12)) and the endpoint gauge laws for channels (Proposition 4.54 and Corollary 4.64).

Definition 4.220 (Wordwise covariant derivative)**.** For each component $\hat{G}^{\alpha\beta}(\cdot; \mathbf{W}, \chi_{\mathrm{out}} \leftarrow \chi_{\mathrm{in}})$, define the *wordwise* covariant derivative (w.r.t. the first endpoint x)

$$\mathbb{D}^{[\mathbf{W};\, \chi_{\mathrm{out}} \leftarrow \chi_{\mathrm{in}}]}_{\mu} \hat{G}^{\alpha\beta} \;:=\; \partial_{\mu} \hat{G}^{\alpha\beta} \;+\; \big[\, \mathcal{A}_{\mu},\; \hat{G}^{\alpha\beta} \,\big]. \tag{4.84}$$

Lemma 4.221 (Gauge covariance and typing preservation)**.** *Under an endpoint gauge $g : \mathcal{X}_q \to \mathrm{U}(\mathcal{H}_{\mathrm{em}})$,*

$$\mathcal{A}_{\mu} \mapsto g \mathcal{A}_{\mu} g^{\dagger} + (\partial_{\mu} g) g^{\dagger}, \qquad \hat{G}^{\alpha\beta} \mapsto g(x)\, \hat{G}^{\alpha\beta}\, g(x')^{\dagger},$$

and

$$\mathbb{D}^{[\mathbf{W};\, \chi_{\mathrm{out}} \leftarrow \chi_{\mathrm{in}}]}_{\mu} \hat{G}^{\alpha\beta} \;\mapsto\; g(x) \Big(\mathbb{D}^{[\mathbf{W};\, \chi_{\mathrm{out}} \leftarrow \chi_{\mathrm{in}}]}_{\mu} \hat{G}^{\alpha\beta}\Big)\, g(x')^{\dagger}.$$

Moreover, the endpoint typing $(\alpha, x) \to (\beta, x')$ is unchanged by $\mathbb{D}^{[\cdot]}_{\mu}$ since the commutator acts by conjugation on the RR space while respecting the letter typing rules (Propositions 4.143 and 4.167).

Proof. Direct substitution of the gauge laws (Proposition 4.54 and Corollary 4.64) into (4.84); typing preservation follows because the commutator with an even (connection) element does not alter the letter endpoints and admissible concatenations. □

Lemma 4.222 (Wordwise Leibniz rule)**.** *Let $X(\cdot; \mathbf{W}_1)$, $Y(\cdot; \mathbf{W}_2)$ be word-evaluated channels whose concatenation is admissible (Definition 4.141). Then*

$$\mathbb{D}^{[\mathbf{W}_2 \mathbf{W}_1]}_{\mu}\big(XY\big) = \big(\mathbb{D}^{[\mathbf{W}_1]}_{\mu} X\big) Y + X \big(\mathbb{D}^{[\mathbf{W}_2]}_{\mu} Y\big).$$

Proof. Expand $\partial_{\mu}(XY)$ and use that $[\,\mathcal{A}_{\mu}, XY\,] = [\,\mathcal{A}_{\mu}, X\,]Y + X[\,\mathcal{A}_{\mu}, Y\,]$, with admissibility of word concatenation provided by typing rules (Proposition 4.143). □

Proposition 4.223 (Wordwise parallel consistency along a path)**.** *Let $\gamma : [s_0, s_1] \to \mathcal{X}_q$ with $\gamma(s_0) = x'$, $\gamma(s_1) = x$ and $A_{\gamma}(s) = (\gamma^{*}\mathcal{A})_s(\dot{\gamma}(s))$ (eq. (4.10)). Assume* word-locked selectors *at the readout end:*

$$\partial_s \Big(\hat{\mathcal{F}}^{[\mathbf{W};\, \chi_{\mathrm{out}} \leftarrow \chi_{\mathrm{in}}]}_{\alpha}\big(\gamma(s)\big)\Big)^{\dagger} + \Big[A_{\gamma}(s),\; \Big(\hat{\mathcal{F}}^{[\mathbf{W};\, \chi_{\mathrm{out}} \leftarrow \chi_{\mathrm{in}}]}_{\alpha}\big(\gamma(s)\big)\Big)^{\dagger}\Big] = 0.$$

Then for $\hat{G}^{\alpha\beta}\big(\gamma(s), x'; \mathbf{W}, \chi_{\text{out}} \leftarrow \chi_{\text{in}}\big)$ *one has the* parallel consistency ODE

$$\Big(\partial_s + [A_\gamma(s), \cdot\,]\Big)\hat{G}^{\alpha\beta}\big(\gamma(s), x'; \mathbf{W}, \chi_{\text{out}} \leftarrow \chi_{\text{in}}\big) = 0.$$

Proof. Differentiate the defining product and use the transport ODE $\partial_s \mathcal{U} = -A_\gamma(s)\mathcal{U}$ (eq. (4.12)), the word-lock condition at the readout end, and Lemma 4.222 for derivation property. All terms cancel pairwise. □

Proposition 4.224 (Wordwise curvature identity)**.** *For any word-evaluated channel* $X(\cdot; \mathbf{W})$*,*

$$\big[\mathbb{D}_\mu^{[\mathbf{W}]},\ \mathbb{D}_\nu^{[\mathbf{W}]}\big]X = \big[\,\mathcal{F}_{\mu\nu},\ X\,\big], \qquad \mathcal{F}_{\mu\nu} = \partial_\mu \mathcal{A}_\nu - \partial_\nu \mathcal{A}_\mu + [\mathcal{A}_\mu, \mathcal{A}_\nu]$$

(Definition 4.56). If $\mathcal{F} = 0$ *on a simply connected neighbourhood, the parallel consistency of Proposition 4.223 integrates to word–path independence up to endpoint gauge (Proposition 4.59).*

Proof. Compute using (4.84); the connection enters only through ordinary commutators, so the standard curvature identity applies. Integrability and endpoint gauge covariance give path independence. □

Corollary 4.225 (Compatibility with Hadamard sum/difference)**.** *For the symmetric interferometer choice* $\mathbf{U}_{\text{Had}}$ *(Definition 2.123), the Hadamard-induced decomposition of RR elements into sum/difference histories (Corollary 4.168) is preserved by the wordwise derivative:*

$$\mathbb{D}_\mu^{[\mathbf{U}_{\text{Had}}]}\hat{G}_\Lambda^{\alpha\beta}(\cdot; \mathbf{U}_{\text{Had}}, \chi \leftarrow \chi) = \tfrac{1}{2}\Big(\mathbb{D}_\mu^{[\cdot]}\hat{G}_+^{\alpha\beta} + \mathbb{D}_\mu^{[\cdot]}\hat{G}_-^{\alpha\beta}\Big),$$

where $\hat{G}_\pm^{\alpha\beta}$ *are the channels built from the* $\pm$ *combinations of the two lifted histories (Definition 4.148).*

Remark 4.226 (Endogenous vs. graded kinematics)**.** The wordwise derivative (4.84) is the endogenous counterpart of $\mathfrak{D}_\mu$ in Definition 4.134. It avoids the graded overlay: no supercommutator is needed—letters carry typing, and $\mathcal{A}_\mu$ is even—yet Lemmas 4.221 and 4.222 and Proposition 4.224 mirror Lemmas 4.135 and 4.136 and Proposition 4.138. In particular, flatness implies word–path independence (Proposition 4.59), and parallel consistency along γ is exactly Proposition 4.223.

4.6 Emergent Inner Product Structure of Modal Hilbert Spaces

Standing GNS framework and notation. By Definition 1.34, the Holo-State $\omega^{\overset{\circ}{\mathbf{\Psi}}} : \mathcal{A} \to \mathbb{C}$ induces a GNS triple $(\pi_{\overset{\circ}{\mathbf{\Psi}}},\ \mathcal{H}_{\overset{\circ}{\mathbf{\Psi}}},\ |\Omega_{\omega^{\overset{\circ}{\mathbf{\Psi}}}}\rangle)$. We set $\mathcal{H}_{\text{em}} := \mathcal{H}_{\overset{\circ}{\mathbf{\Psi}}}$ and adopt the represented-operator convention: for any $A \in \mathcal{A}$, $\hat{A} := \pi_{\overset{\circ}{\mathbf{\Psi}}}(A) \in \mathcal{B}(\mathcal{H}_{\text{em}})$, with expectations $\omega^{\overset{\circ}{\mathbf{\Psi}}}(A) = \langle\Omega_{\omega^{\overset{\circ}{\mathbf{\Psi}}}}|\ \hat{A}\ |\Omega_{\omega^{\overset{\circ}{\mathbf{\Psi}}}}\rangle$. All field modes $\hat{\mathcal{F}}_\alpha$, channel kernels $\hat{G}_\Lambda^{\alpha\beta}(x, x')$, and the transporter $\mathcal{U}(x \leftarrow$

$x';\gamma) \in \mathrm{U}(\mathcal{H}_{\mathrm{em}})$ act on $\mathcal{H}_{\mathrm{em}}$, with $\hat{G}_\Lambda^{\alpha\beta}$ defined from selectors and transport as in Definition 4.61 and $\mathcal{U}$ given by the parallel–transport constructions Definitions 4.48 and 4.49 (cf. gauge covariance in Proposition 4.54 and Corollary 4.64).

Endogenous augmentation (holosymmetric words). In the holosymmetric words framework, device words $\mathbf{W}$ from Definition 4.148 act on the RR factor to dress the selectors via their RR matrix elements $\mathsf{W}_{\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}}$ (eq. (4.45)), producing word–dressed selectors

$$\hat{\mathcal{F}}_\alpha^{[\mathbf{W};\,\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}]}(x) := \mathsf{W}_{\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}}\,\hat{\mathcal{F}}_\alpha(x) \quad (\textit{Definition } 4.166).$$

The corresponding *word–evaluated channels* are

$$\hat{G}_\Lambda^{\alpha\beta}\Big(x,x';\mathbf{W},\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}\Big) := \Big(\hat{\mathcal{F}}_\alpha^{[\mathbf{W};\,\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}]}(x)\Big)^\dagger\,\mathcal{U}(x\leftarrow x';\gamma)\,\hat{\mathcal{F}}_\beta(x') \quad (\textit{eq. } (4.66)),$$

which reduce to the undeformed $\hat{G}_\Lambda^{\alpha\beta}(x,x')$ when $\mathbf{W}=I$ and $\chi_{\mathrm{out}}=\chi_{\mathrm{in}}$. Gauge covariance, adjoint symmetry, and degree additivity are preserved under dressing (Proposition 4.167) and under word formation from the channel alphabet (Definition 4.141 and Proposition 4.143). Typing, block composition through intermediate loci, and endpoint–degree rules hold verbatim for words (Proposition 4.85, Lemma 4.145, and Corollary 4.144).

Commutation relations are taken with the *endogenous super–commutator* (Definition 4.203); the bracket type (commutator vs. anticommutator) is fixed by the holosymmetric charges and the selection rule $\langle q(\cdot), q(\cdot)\rangle$ (Corollary 4.205). Words–level centrality, bosonic $0/s$ compatibility, and $p\leftrightarrow s$ nilpotency then apply to these operators acting on $\mathcal{H}_{\mathrm{em}}$ (Theorems 4.169, 4.170 and 4.173), with curvature/holonomy–controlled defects packaged by the compliance vector $\mathfrak{K}$ (Definition 4.178).

Assumption 4.227 (Holo-State and representation backdrop)**.** *Throughout this section, $\omega^{\mathring{\mathbf{\Psi}}}$ is a positive, normalized state on $\mathcal{A}$ with GNS triple $(\pi_{\mathring{\mathbf{\Psi}}}, \mathcal{H}_{\mathring{\mathbf{\Psi}}}, |\Omega_{\omega^{\mathring{\mathbf{\Psi}}}}\rangle)$ as provided by Definition 1.34. We identify $\mathcal{H}_{\mathrm{em}} := \mathcal{H}_{\mathring{\mathbf{\Psi}}}$, write $\hat{A} := \pi_{\mathring{\mathbf{\Psi}}}(A)$, and abbreviate $\langle\mathring{\mathbf{\Psi}}|\cdot|\mathring{\mathbf{\Psi}}\rangle = \langle\Omega_{\omega^{\mathring{\mathbf{\Psi}}}}|\cdot|\Omega_{\omega^{\mathring{\mathbf{\Psi}}}}\rangle = \omega^{\mathring{\mathbf{\Psi}}}(\cdot)$ on $\mathcal{A}$. In particular, the modal operators and channels act on $\mathcal{H}_{\mathrm{em}}$ as in Definitions 4.48, 4.49 and 4.61, and transform covariantly under endpoint gauge by Proposition 4.54 and Corollary 4.64.*

Definition 4.228 (Emergent inner product on a modal fiber)**.** Let x label a PQS locus and let $|\psi_i\rangle_x \in \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ be modal states (Definition 4.273). The inner product on the fiber at x is *defined* from the Fourfold Inter-Modal Correlator (Definition 4.9) by

$$\boxed{\langle\psi_1|\psi_2\rangle_x := \lim_{x'\to x}\Big\langle\mathring{\mathbf{\Psi}}\Big|\,\hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})[\psi_1^*,\psi_2]\,\Big|\mathring{\mathbf{\Psi}}\Big\rangle = \lim_{x'\to x}\omega^{\mathring{\mathbf{\Psi}}}(G_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})[\psi_1^*,\psi_2])\,.} \tag{4.85}$$

Assumption 4.229 (Regularity near the diagonal)**.** *The map $(x,x')\mapsto\hat{G}_\Lambda^{\alpha\beta}(x,x')$ is continuous in a neighborhood of the diagonal $\{(x,x) : x\in\mathcal{X}_q\}$ (in operator norm or weak operator topology), and the parallel transporter $\mathcal{U}(x\leftarrow x';\gamma)$ is unitary and continuous in (x,x'). These conditions ensure the existence of the limit in (4.85).*

Theorem 4.230 (Regularity near the diagonal for channel kernels)**.** *Let* $\hat{G}_\Lambda^{\alpha\beta}(x,x') = \hat{\mathcal{F}}_\alpha^\dagger(x)\,\mathcal{U}(x\leftarrow x';\gamma_{x\to x'})\,\hat{\mathcal{F}}_\beta(x')$ *be the Fourfold channel of Definition 4.61, with transporter* $\mathcal{U}$ *defined by Definitions 4.48 and 4.49. Assume the selector fields* $x\mapsto\hat{\mathcal{F}}_\alpha(x)$ *are operator–norm continuous on* $\mathcal{X}_q$*, uniformly bounded on compact sets, and that the operational connection of Section 4.19.1 is locally bounded with continuous curvature.*

Then:

(1) Unitarity and continuity of transport

For each admissible path $\gamma_{x\to x'}$*,* $\mathcal{U}(x\leftarrow x';\gamma_{x\to x'})$ *is unitary and depends continuously on* (x,x') *(in operator norm on compact sets); see Definition 4.49 and the Grönwall–type stability that follows from the local boundedness of the connection.*

(2) Near–diagonal continuity of $\hat{G}_\Lambda^{\alpha\beta}$

The map $(x,x')\mapsto\hat{G}_\Lambda^{\alpha\beta}(x,x')$ *is operator–norm continuous in a neighborhood of the diagonal* $\{(x,x):x\in\mathcal{X}_q\}$*. Hence, for any net* $(x_\nu,x'_\nu)\to(x,x)$*,*

$$\left\|\,\hat{G}_\Lambda^{\alpha\beta}(x_\nu,x'_\nu)-\hat{G}_\Lambda^{\alpha\beta}(x,x)\,\right\|\;\longrightarrow\;0.$$

(3) Existence of the coincident limit

The regulated coincident limit used in the emergent inner–product construction exists:

$$\lim_{(x',x)\to(x,x)}\;\langle\psi,\;\hat{G}_\Lambda^{\alpha\beta}(x,x')\,\phi\rangle\quad\textit{exists for all }\psi,\phi\in\mathcal{H}_{\mathrm{em}},$$

and coincides with the operator–norm limit from (2), thus justifying the use of (4.85) in §4.6.

Moreover, all statements are stable under endpoint gauge conjugation by $\hat{G}_\Lambda^{\alpha\beta}\mapsto g(x)\hat{G}_\Lambda^{\alpha\beta}g(x')^\dagger$ *(Proposition 4.54 and Corollary 4.64).*

Proof. (1) Unitarity follows from the ODE characterization with skew–Hermitian generator ($\dot{\mathcal{U}}=-\mathcal{A}(\dot\gamma)\mathcal{U}$) in Definition 4.49, which yields $\frac{d}{dt}\big(\mathcal{U}^\dagger\mathcal{U}\big)=0$. Continuity in (x,x') along a canonical local family of admissible paths follows from standard integral–equation stability and the local boundedness of the connection in Section 4.19.1.

(2) Write

$$\hat{G}_\Lambda^{\alpha\beta}(x,x')=\hat{\mathcal{F}}_\alpha^\dagger(x)\,\mathcal{U}(x\leftarrow x';\gamma_{x\to x'})\,\hat{\mathcal{F}}_\beta(x').$$

Each factor is operator–norm continuous by hypothesis (selectors) and by (1) (transport). Uniform boundedness on compact neighborhoods is ensured by the spectral restriction $\mathfrak{F}_x=\operatorname{Ran}E^{(x)}([0,\lambda_{\mathrm{cutoff}}])$ (eq. (4.303)), so the product map is operator–norm continuous.

(3) The operator–norm continuity implies strong and weak operator continuity; thus the limit (4.85) exists and equals the operator–norm limit. Gauge stability is inherited from Proposition 4.54 and Corollary 4.64. □

Corollary 4.231 (Words–dressed regularity)**.** *For word–dressed selectors and channels of Definition 4.166 and eq. (4.66),*

$$\hat{G}_\Lambda^{\alpha\beta}\big(x,x';\mathbf{W},\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}\big)=\Big(\hat{\mathcal{F}}_\alpha^{[\mathbf{W};\,\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}]}(x)\Big)^\dagger\,\mathcal{U}(x\leftarrow x';\gamma)\,\hat{\mathcal{F}}_\beta(x'),$$

the map $(x, x') \mapsto \hat{G}_\Lambda^{\alpha\beta}(\cdots)$ is operator–norm continuous near the diagonal, and the coincident limit exists as in (4.85). Gauge covariance and adjoint/degree properties are preserved by Proposition 4.167.

Corollary 4.232 (Compatibility with channel words)**.** *Let W be any channel word in the sense of Definition 4.141. The map $(x, x') \mapsto W(x, x')$ obtained by block composition through intermediate loci (Propositions 4.85 and 4.143) is operator–norm continuous near the diagonal, and its coincident limit exists in the sense of (4.85).*

Remark (strength relative to the original assumption). Theorem 4.230 replaces Assumption 4.229. In particular, the existence of the emergent inner-product limit is now a consequence of: (i) the ODE/parallel-transport framework (Definitions 4.48 and 4.49 and Section 4.19.1), (ii) selector continuity and spectral localization (eq. (4.303)), and (iii) gauge covariance (Proposition 4.54 and Corollary 4.64). The words-level extensions (Definitions 4.141 and 4.166 and Proposition 4.167) show that the result is robust under holosymmetric dressing and composition.

Proposition 4.233 (Well-posedness of the fiber inner product)**.** *Under Assumption 4.227 and the near-diagonal regularity of Theorem 4.230, the sesquilinear form $\langle\cdot|\cdot\rangle_x$ defined by the coincident-limit prescription (4.85) is antilinear in the first argument, linear in the second, Hermitian $\langle\psi_1|\psi_2\rangle_x = \overline{\langle\psi_2|\psi_1\rangle_x}$, and positive semidefinite. Quotienting by its null space and completing yields a Hilbert space, denoted again $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$.*

Proof. Setup. For fixed $x \in \mathcal{X}_q$, define the limiting operator

$$K_x \;:=\; \lim_{(x',x)\to(x,x)} \hat{G}_\Lambda(x, x') \quad \text{in operator norm,}$$

whose existence follows from Theorem 4.230. Here $\hat{G}_\Lambda(x, x')$ stands for the (modal) channel kernel built from selectors and transport as in Definition 4.61, i.e. $\hat{G}_\Lambda^{\alpha\beta}(x, x') = \hat{\mathcal{F}}_\alpha^\dagger(x)\,\mathcal{U}(x \leftarrow x';\gamma)\,\hat{\mathcal{F}}_\beta(x')$, with $\mathcal{U}$ as in Definitions 4.48 and 4.49. Define

$$\langle\psi_1|\psi_2\rangle_x \;:=\; \langle\psi_1,\, K_x\,\psi_2\rangle_{\mathcal{H}_{\mathrm{em}}} \qquad (\psi_1, \psi_2 \in \mathcal{H}_{\mathrm{em}}).$$

Sesquilinearity. For each (x, x'), the map $(\psi_1, \psi_2) \mapsto \langle\psi_1,\, \hat{G}_\Lambda(x, x')\,\psi_2\rangle$ is antilinear in ψ_1 and linear in ψ_2. Passing to the operator-norm limit preserves sesquilinearity; hence $\langle\cdot|\cdot\rangle_x$ is antilinear/linear in its arguments.

Hermiticity. By the channel Hermitian symmetry $(\hat{G}_\Lambda^{\alpha\beta}(x, x'))^\dagger = \hat{G}_\Lambda^{\beta\alpha}(x', x)$ (Lemma 4.66 and Proposition 4.52), $\hat{G}_\Lambda(x, x')^\dagger = \hat{G}_\Lambda(x', x)$. Taking the near-diagonal limit in operator norm yields $K_x^\dagger = K_x$, so $\langle\psi_1|\psi_2\rangle_x = \overline{\langle\psi_2|\psi_1\rangle_x}$.

Positive semidefiniteness. Write the selector column $\mathcal{F}(x) := \big(\hat{\mathcal{F}}_0(x), \hat{\mathcal{F}}_1(x), \hat{\mathcal{F}}_s(x), \hat{\mathcal{F}}_p(x)\big)$ as an operator $\mathcal{H}_{\mathrm{em}} \to \mathcal{H}_{\mathrm{em}}^{\oplus 4}$. In the coincident/flat limit, $\mathcal{U}(x \leftarrow x;\gamma) = I$ (Proposition 4.59), hence

$$K_x \;=\; \lim_{x'\to x} \hat{G}_\Lambda(x, x') \;=\; \sum_{\alpha,\beta} \hat{\mathcal{F}}_\alpha^\dagger(x)\,\delta_{\alpha\beta}\,\hat{\mathcal{F}}_\beta(x) \;=\; \mathcal{F}(x)^\dagger\,\mathcal{F}(x).$$

Therefore, for any $\psi \in \mathcal{H}_{\mathrm{em}}$, $\langle\psi|\psi\rangle_x = \langle\psi,\, \mathcal{F}(x)^\dagger\mathcal{F}(x)\,\psi\rangle = \|\mathcal{F}(x)\psi\|^2 \geq 0$, establishing positive semidefiniteness. (The equality with the near-diagonal limit is ensured by Theorem 4.230.)

Quotient and completion. Let $\mathcal{N}_x := \{\psi : \langle\psi|\psi\rangle_x = 0\}$ be the null space. The Cauchy–Schwarz inequality holds for positive semidefinite sesquilinear forms; thus $\langle\cdot|\cdot\rangle_x$ descends to a genuine inner product on the quotient $\mathcal{H}_{\mathrm{em}}/\mathcal{N}_x$. Completing this pre-Hilbert space yields a Hilbert space, denoted $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$.

Gauge covariance (well-posedness). Under endpoint gauge (Proposition 4.54 and Corollary 4.64), the transformations are:

$$\begin{aligned} \hat{G}_\Lambda(x,x') &\mapsto g(x)\hat{G}_\Lambda(x,x')g(x')^\dagger \\ \text{whence } K_x &\mapsto g(x)K_x g(x)^\dagger. \end{aligned} \tag{4.86}$$

The quadratic form

$$\langle\psi_1|\psi_2\rangle_x = \langle\psi_1, K_x\psi_2\rangle$$

is gauge-invariant, so the construction is well posed on the fiber.

All claims follow. □

Remark 4.234 (Words-dressed extension)**.** For word-dressed channels $\hat{G}_\Lambda^{\alpha\beta}(\cdots;\mathbf{W})$ (Definition 4.166 and eq. (4.66)), Theorem 4.230 and Corollary 4.231 ensure the coincident-limit operator $K_x^{[\mathbf{W}]}$ exists and remains positive: $K_x^{[\mathbf{W}]} = \mathcal{F}^{[\mathbf{W}]}(x)^\dagger\mathcal{F}^{[\mathbf{W}]}(x)$ with $\mathcal{F}^{[\mathbf{W}]}(x) := \left(\hat{\mathcal{F}}_\alpha^{[\mathbf{W}]}(x)\right)_\alpha$. Thus the same proof yields a well-posed inner product and the same fiber Hilbert space.

Status and geometry of the emergent spaces (endogenous strengthening).

Definition 4.235 (Modal bundle over $\mathcal{X}_q$)**.** Let $\mathcal{X}_q$ be the emergent configuration space of eq. (4.322). For each class $\mathfrak{B}_x$ (anchor $x \in \mathcal{X}_q$), let $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ be the fiber Hilbert space obtained from the coincident-limit inner product $\langle\cdot|\cdot\rangle_x$ in eq. (4.85) and Proposition 4.233. Define the total space and projection

$$\mathscr{H}_{\mathrm{mod}} := \bigsqcup_{x\in\mathcal{X}_q} \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}, \qquad \pi_{\mathrm{mod}} : \mathscr{H}_{\mathrm{mod}} \longrightarrow \mathcal{X}_q, \quad \pi_{\mathrm{mod}}(\psi) = x \ \text{ if } \ \psi \in \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}.$$

Then $(\mathscr{H}_{\mathrm{mod}}, \pi_{\mathrm{mod}})$ is a (locally trivial) Hermitian Hilbert bundle, with typical fiber isomorphic to the completion of $\mathcal{H}_{\mathrm{em}}/\mathcal{N}_x$ as in Proposition 4.233. Local trivializations are provided by the operational transporter of Section 4.19.1 in the regime of Theorem 4.230; the structure group acts by fiberwise unitaries (cf. Proposition 4.54 and Corollary 4.64).

Proposition 4.236 (Fiber metric is generated by correlators)**.** *For each $x \in \mathcal{X}_q$, let $K_x := \lim_{(x',x)\to(x,x)} \hat{G}_\Lambda(x,x')$ (operator-norm limit) as in Theorem 4.230. Then the fiber inner product is*

$$\langle\psi_1|\psi_2\rangle_x \;=\; \langle\psi_1,\; K_x\,\psi_2\rangle_{\mathcal{H}_{\mathrm{em}}} \qquad (\psi_1,\psi_2 \in \mathcal{H}_{\mathrm{em}}),$$

Hermitian and positive semidefinite (Proposition 4.233). In the flat/coincident regime (Proposition 4.59), $K_x = \mathcal{F}(x)^\dagger\mathcal{F}(x)$ with $\mathcal{F}(x) = (\hat{\mathcal{F}}_0(x), \hat{\mathcal{F}}_1(x), \hat{\mathcal{F}}_s(x), \hat{\mathcal{F}}_p(x))$, so

$$\langle\psi|\psi\rangle_x = \|\mathcal{F}(x)\psi\|^2,$$

making the induced metric state- and transport-aware *via $\omega^{\mathring{\Psi}}$ and the operational connection (Assumption 4.227 and Section 4.19.1).*

Proposition 4.237 (Local triviality, metric compatibility, and holonomy defect)**.** *Let* $\gamma : [0,1] \to \mathcal{X}_q$ *be an admissible curve with parallel transport* $\mathcal{U}_\gamma$ *of Definitions 4.48 and 4.49. Then on a simply connected flat neighborhood* ($\mathcal{F} \equiv 0$) *one has*

$$K_{\gamma(1)} \;=\; \mathcal{U}_\gamma\, K_{\gamma(0)}\, \mathcal{U}_\gamma^\dagger, \quad \textit{hence} \quad \langle \mathcal{U}_\gamma \psi_1 \,|\, \mathcal{U}_\gamma \psi_2 \rangle_{\gamma(1)} = \langle \psi_1 \,|\, \psi_2 \rangle_{\gamma(0)}.$$

Thus parallel transport gives unitary identifications *of nearby fibers (metric compatibility). In the curved case,*

$$\left\| K_{\gamma(1)} - \mathcal{U}_\gamma\, K_{\gamma(0)}\, \mathcal{U}_\gamma^\dagger \right\| \;\le\; C\, \Phi(\gamma),$$

with holonomy functional Φ *controlled by the curvature* $\mathcal{F}$ *(non-Abelian Stokes–type stability), i.e. metric non-parallelism is* bounded by holonomy *(Proposition 4.213).*

Remark 4.238 (Fourfold roles encoded in the fiber metric)**.** Decompose K_x into modal blocks using the channel alphabet (Definition 4.141):

$$K_x \;=\; \sum_{\alpha} \hat{\mathcal{F}}_\alpha^\dagger(x) \hat{\mathcal{F}}_\alpha(x) \;+\; \sum_{\alpha \neq \beta} \hat{\mathcal{F}}_\alpha^\dagger(x) \hat{\mathcal{F}}_\beta(x).$$

The diagonal terms encode *modal persistence* (occupancy), while off-diagonal terms encode *transitions/interference*. At words level, the $0/s$ diagonal refinements of triadic geometry split the diagonal contributions by sector (Proposition 4.187); central holosymmetry yields ss-normal forms that factor out a commuting "coherence scalar" (Theorem 4.169 and Proposition 4.188); and the $p \leftrightarrow s$ edge constraints (nilpotency) suppress multiple-step interference in those channels (Theorem 4.170). Departures in curved regimes are quantitatively controlled by the compliance vector $\mathfrak{K} = (\Delta, \eta, \zeta)$ (Definitions 4.178 and 4.183).

Principle of an emergent inner product (conceptual upshot, formalized). The fiber inner product is a *derived observable*:

$$\langle \psi_1 | \psi_2 \rangle_x \;=\; \omega^{\mathring{\mathbf{\Psi}}}\Big(\langle \psi_1 |\, \hat{G}_\Lambda(x,x')\, | \psi_2 \rangle \Big)\Big|_{(x',x)\to(x,x)}$$

(by Assumption 4.227 and Theorem 4.230; see also Proposition 4.233). (4.87)

Thus the probabilistic geometry of each fiber is fixed by the global, entangled architecture encoded in the Holo-State and transported by the operational connection (Section 4.19.1). In flat, commuting regimes ($\mathcal{F} \equiv 0$, central holosymmetry exact), the construction reduces to the classical dyad's metric on the diagonal sector (Proposition 4.187); when coherence and global structure matter, the full Fourfold content and its words-level symmetries determine the metric and its sectoral decomposition (Definition 4.203, Corollary 4.205, and Theorem 4.169).

4.6.1 Hilbert Bundle of Dialectic Modes over the Emergent Configuration Space

Building on the Primordial Quantum Singularity $\mathfrak{B}_x$ (Definition 1.25; cf. the PQS continuum Definition 4.617) and its emergent modal fiber $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$, whose inner product is generated by the Fourfold correlator (Section 4.6, eq. (4.85), and Proposition 4.233), we now organize these fibers over the emergent quantum configuration space $\mathcal{X}_q$ (Proposition 4.625) into a geometric bundle.

Definition 4.239 (Hilbert bundle of dialectic modes)**.** Let $\mathcal{X}_q$ be the emergent configuration space (eq. (4.322)). To each $x \in \mathcal{X}_q$ associate a PQS $\mathfrak{B}_x$ and its modal fiber $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ (a four-dimensional complex Hilbert space once the inner product is fixed by eq. (4.85) and Proposition 4.233). The *Hilbert bundle of dialectic modes* is the triple $(\mathscr{H}_{\mathrm{mod}}, \pi, \mathcal{X}_q)$ with

$$\boxed{\mathscr{H}_{\mathrm{mod}} := \bigsqcup_{x\in\mathcal{X}_q} \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}, \qquad \pi : \mathscr{H}_{\mathrm{mod}} \to \mathcal{X}_q, \quad \pi^{-1}(x) = \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)} \cong \mathbb{C}^4.} \tag{4.88}$$

After eq. (4.85) is established, choose unit representatives of the realized rays $\mathcal{R}_x(\alpha) \subset \mathbb{P}(\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)})$ for $\alpha \in \{0, 1, p, s\}$ to obtain an orthonormal ordered frame

$$\mathcal{B}_x := \{|0\rangle_x,\ |1\rangle_x,\ |p\rangle_x,\ |s\rangle_x\}, \qquad \langle\alpha_x|\beta_x\rangle = \delta_{\alpha\beta}.$$

Here $|0\rangle_x, |1\rangle_x$ are the Symmetry/Antisymmetry (dyad) modes, while $|p\rangle_x, |s\rangle_x$ realize the independent Parasymmetry/Holosymmetry channels (cf. the Fourfold alphabet and words in Definition 4.141).

Remark 4.240 (Independence of p, s from twofold superpositions)**.** The equal-superposition vector $|+\rangle_x := \frac{1}{\sqrt{2}}\big(|0\rangle_x + |1\rangle_x\big)$ lives in the *twofold* sector and defines the local parasymmetric *effect* $P^{(x)}_{\mathrm{para}} = |+\rangle_x\langle+|$ (Hadamard/sum interferometer; see Corollary 4.168); it is not identified with the fourfold basis vector $|p\rangle_x$. Similarly, $|s\rangle_x$ is not assumed to be any linear combination of $|0\rangle_x, |1\rangle_x$. The p and s channels are independent fiber directions whose operational meanings are read via their dedicated effects (parasymmetric and holosymmetric, respectively) and the endogenous bracket/selection rules (Definition 4.203 and Corollary 4.205).

Local trivializations, gauge, and connection (formalization). On a coordinate patch $U \subset \mathcal{X}_q$, a smooth choice of ordered orthonormal frames $\{\mathcal{B}_x\}_{x\in U}$ trivializes the bundle as $\pi^{-1}(U) \cong U \times \mathbb{C}^4$. The *structure group* is the subgroup

$$G_{\mathrm{struct}} \subseteq U(4)$$

that preserves the $\{0, 1, p, s\}$ ordering (and, when invoked, the $\mathbb{Z}_2 \times \mathbb{Z}_2$ charge/grade; cf. Definition 4.201 and Corollary 4.205). Let $\{\mathcal{B}_x\}$ and $\{\widetilde{\mathcal{B}}_x\}$ be two trivializations on U with transition $g : U \to G_{\mathrm{struct}}$. Then the *bundle connection* is given in these frames by a $u(4)$-valued 1–form $\mathcal{A}$ transforming as

$$\widetilde{\mathcal{A}} \;=\; g\,\mathcal{A}\,g^{\dagger} \;+\; g\,\mathrm{d}g^{\dagger}, \qquad \mathcal{F} \;:=\; \mathrm{d}\mathcal{A} + \mathcal{A}\wedge\mathcal{A}, \tag{4.89}$$

and its parallel transport along an admissible path $\gamma : x' \to x$ is the path–ordered exponential

$$\mathcal{U}_{\mathcal{A}}(x \leftarrow x'; \gamma) \;:=\; \mathcal{P}\exp\Big(-\int_{\gamma}\mathcal{A}\Big), \tag{4.90}$$

which transforms by endpoint conjugation

$$\mathcal{U}_{\mathcal{A}} \mapsto g(x)\,\mathcal{U}_{\mathcal{A}}\,g(x')^{\dagger}$$

and coincides with the operational transporter of the theory:

$$\mathcal{U}_{\mathcal{A}}(x \leftarrow x';\gamma) \;=\; \mathcal{U}(x \leftarrow x';\gamma) \quad \text{from Definitions 4.48 and 4.49 and Section 4.19.1.} \tag{4.91}$$

In particular, $\mathcal{F}$ is the curvature controlling holonomy and the near–diagonal regularity of modal kernels (Theorem 4.230 and Proposition 4.213); on simply connected flat patches ($\mathcal{F} \equiv 0$) transport is path–independent (Proposition 4.59).

Proposition 4.241 (Metric compatibility and holonomy bound)**.** *Let K_x be the coincident–limit kernel defining the fiber inner product (eq. (4.85) and Proposition 4.233). Then for any admissible path $\gamma : x' \to x$,*

$$K_x \;=\; \mathcal{U}(x \leftarrow x';\gamma)\; K_{x'}\; \mathcal{U}(x \leftarrow x';\gamma)^{\dagger} \qquad \text{if } \; \mathcal{F}|_{\gamma} \equiv 0,$$

so that parallel transport gives unitary identifications of nearby fibers (metric compatibility). In general,

$$\Big\| K_x - \mathcal{U}(x \leftarrow x';\gamma)\; K_{x'}\; \mathcal{U}(x \leftarrow x';\gamma)^{\dagger} \Big\| \;\le\; C\,\Phi(\gamma),$$

where $\Phi(\gamma)$ is a holonomy functional controlled by $\mathcal{F}$ (non-Abelian Stokes–type estimate), as in Propositions 4.213 and 4.237.

Remark 4.242 (Compatibility with endogenous words and sectors)**.** The holosymmetric $\mathbb{Z}_2^2$ *charges* live in the device/dialectic factor and are preserved by G_{struct}, hence endogenous centrality, $0/s$ bosonic compatibility, and $p \leftrightarrow s$ nilpotency (Theorems 4.169, 4.170 and 4.173) apply fiberwise and glue across patches by (4.89)–(4.91). Sector decompositions by the spectral projectors of $\hat{G}^{ss}_{\Lambda}$ are bundle–compatible (Remark 4.211 and Proposition 4.188).

Fiberwise orthonormality (emergent). By construction from (4.85), each local frame is orthonormal:

$$\langle \alpha_x | \beta_x \rangle_{\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}} = \delta_{\alpha\beta}, \qquad \alpha,\beta \in \{0,1,p,s\}.$$

In particular, the twofold subbundle spanned by $\{|0\rangle_x, |1\rangle_x\}$ recovers the classical dyad, while mixing with $\{|p\rangle_x, |s\rangle_x\}$ is mediated by the off-diagonal channels of the Fourfold correlator.

We endow the fibers with a Hermitian (unitary) structure and fix the structure group $G \subseteq \mathrm{U}(4)$ acting in the fundamental representation. We assume *local triviality*: for every open set $U \subset \mathcal{X}_q$ there exists a unitary trivialization

$$\pi^{-1}(U) \;\cong\; U \times \mathbb{C}^4.$$

On overlaps $U \cap V$, the transition functions $g_{UV} : U \cap V \to G$ satisfy the cocycle conditions $g_{UU} = \mathbf{1}$, $g_{UV} = g_{VU}^{-1}$, and $g_{UV}\, g_{VW} = g_{UW}$. They relate frames by $|\alpha\rangle_V = g_{UV}^{-1}(x)\,|\alpha\rangle_U$ and therefore preserve the ordered modal frame $\{0,1,p,s\}$; when a grading is imposed, one may restrict to a grading–preserving subgroup, e.g. $G \subseteq \mathrm{U}(2) \times \mathrm{U}(2)$. The connection one–form $\mathcal{A} \in \Omega^1(\mathcal{X}_q, \mathfrak{g})$ with $\mathfrak{g} \subseteq \mathfrak{u}(4)$ and its parallel transporter $\mathcal{U}(x \leftarrow x';\gamma) = \mathcal{P}\exp\!\Big(-\int_{\gamma}\mathcal{A}\Big)$ (Section 4.1.4) provide the differential gluing data.

Definition 4.243 (Emergent modal frame and inner product)**.** At each locus $x \in \mathcal{X}_q$, let $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)} \cong \mathbb{C}^4$ be the modal fibre equipped with the emergent inner product constructed from the Fourfold correlator (Definitions 4.9 and 4.20 and eq. (4.6)). Equivalently, we may reference its coordinate form (4.85). Fix the ordered modal frame

$$\mathcal{B}_x := \Big\{ |0\rangle_x,\ |1\rangle_x,\ |p\rangle_x,\ |s\rangle_x \Big\}.$$

Proposition 4.244 (Fiberwise orthonormality)**.** *By construction from* (4.85) *(*or *Definition 4.20 and eq.* (4.6)*), each local frame $\mathcal{B}_x$ is orthonormal:*

$$\boxed{\langle \alpha_x \mid \beta_x \rangle_{\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}} = \delta_{\alpha\beta}, \qquad \alpha, \beta \in \{0, 1, p, s\}.} \tag{4.92}$$

Definition 4.245 (Twofold (classical) subbundle and para/holo complement)**.** Define the twofold subbundle $\mathcal{H}^{(0,1)} := \mathrm{span}\{|0\rangle_x, |1\rangle_x\}_{x \in \mathcal{X}_q}$ and its orthogonal complement $\mathcal{H}^{(p,s)} := \mathrm{span}\{|p\rangle_x, |s\rangle_x\}_{x \in \mathcal{X}_q}$.

Proposition 4.246 (Mixing controlled by off-diagonal Fourfold channels)**.** *Let $\hat{\mathbf{G}}_\Lambda(x, x') = [\hat{G}_\Lambda^{\alpha\beta}(x, x')]_{\alpha,\beta \in \{0,1,p,s\}}$ be the Fourfold channel block matrix (Definition 4.81). Then, for any x, x',*

(i) If the off-diagonal blocks between the sectors vanish,

$$\hat{G}_\Lambda^{\alpha\beta}(x, x') = 0 \quad \text{for all } \alpha \in \{0, 1\},\ \beta \in \{p, s\} \text{ and for all } \alpha \in \{p, s\},\ \beta \in \{0, 1\},$$

the decomposition $\mathcal{H}^{(0,1)} \oplus \mathcal{H}^{(p,s)}$ is channel-invariant and no mixing occurs.

(ii) In general, for any unit vectors $u \in \mathcal{H}_{x'}^{(0,1)}$, $v \in \mathcal{H}_x^{(p,s)}$,

$$\Big| \langle v \mid \hat{\mathbf{G}}_\Lambda(x, x')\, u \rangle \Big| \ \le \sum_{\substack{\alpha \in \{p,s\} \\ \beta \in \{0,1\}}} \Big\| \hat{G}_\Lambda^{\alpha\beta}(x, x') \Big\|.$$

Thus the leakage amplitude from the classical dyad into the para/holo sector is bounded by the norms of the corresponding off-diagonal channel blocks.

Proof. (i) follows by block-diagonality of $\hat{\mathbf{G}}_\Lambda$. (ii) Expand the matrix element in the modal frames and apply Cauchy–Schwarz together with the operator norm bound. □

Definition 4.247 (Modal Hilbert bundle, structure group, and local triviality)**.** Let $\pi : \mathcal{E} \to \mathcal{X}_q$ be the Hermitian vector bundle with fibres $\mathcal{E}_x = \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)} \cong \mathbb{C}^4$ and ordered frames $\mathcal{B}_x$. Fix the structure group $G \subseteq \mathrm{U}(4)$ acting in the fundamental representation and assume *local triviality*: for every open $U \subset \mathcal{X}_q$ there exists a unitary trivialization

$$\boxed{\pi^{-1}(U) \ \cong\ U \times \mathbb{C}^4.} \tag{4.93}$$

On overlaps $U \cap V$, the transition maps $g_{UV} : U \cap V \to G$ satisfy the cocycle conditions $g_{UU} = \mathbf{1}$, $g_{UV} = g_{VU}^{-1}$, and $g_{UV} g_{VW} = g_{UW}$, and relate frames by $|\alpha\rangle_V = g_{UV}^{-1}(x)\, |\alpha\rangle_U$ (preserving the order $\{0, 1, p, s\}$). When a grading is imposed, one may restrict to a grading–preserving subgroup, e.g. $G \subseteq \mathrm{U}(2) \times \mathrm{U}(2)$ (Section 4.1.7 and Definition 4.94).

Definition 4.248 (Connection and parallel transport (differential gluing data))**.** Let $\mathcal{A} \in \Omega^1(\mathcal{X}_q, \mathfrak{g})$ be a (skew–Hermitian) connection one–form with $\mathfrak{g} \subseteq \mathfrak{u}(4)$, and define the parallel transporter along a piecewise C^1 path γ by

$$\mathcal{U}(x \leftarrow x'; \gamma) = \mathcal{P} \exp\Big(-\int_\gamma \mathcal{A} \Big) \quad \text{(Definition 4.48 and eqs. (4.11) and (4.12)).}$$

Proposition 4.249 (Unitarity and inner-product preservation)**.** *For any path $\gamma : x' \to x$, the transporter $\mathcal{U}(x \leftarrow x'; \gamma) \in G \subseteq \mathrm{U}(4)$ is unitary and preserves the fibre inner products induced by (4.85): for all $\psi, \phi \in \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_{x'})}$,*

$$\big\langle \mathcal{U}\,\psi \;\big|\; \mathcal{U}\,\phi \big\rangle_{\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}} = \langle \psi \mid \phi \rangle_{\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_{x'})}}.$$

Proof. Skew–Hermiticity $\mathcal{A}^\dagger = -\mathcal{A}$ implies $\mathcal{U} \in \mathrm{U}(4)$ via the path-ordered exponential. Compatibility with the fibre metrics follows from the differential equation $\partial_s \mathcal{U} = -A_\gamma(s)\mathcal{U}$ and the standard argument that $\partial_s \langle \mathcal{U}\psi \mid \mathcal{U}\phi \rangle = 0$ (eq. (4.12) and Definition 4.49). □

Proposition 4.250 (Gauge covariance of frames, channels, and connection)**.** *Under $g : \mathcal{X}_q \to G$, $\mathcal{A} \mapsto g\mathcal{A}g^\dagger + (\mathrm{d}g)\, g^\dagger$ and $\mathcal{U}(x \leftarrow x'; \gamma) \mapsto g(x)\,\mathcal{U}(x \leftarrow x'; \gamma)\, g(x')^\dagger$ (Proposition 4.54). Frames transform by $|\alpha\rangle_x \mapsto g(x)\,|\alpha\rangle_x$, preserving (4.92). Fourfold channels obey endpoint covariance $\hat{G}_\Lambda^{\alpha\beta}(x, x') \mapsto g(x)\,\hat{G}_\Lambda^{\alpha\beta}(x, x')\, g(x')^\dagger$ (Corollary 4.64).*

Corollary 4.251 (Flatness and path–independence)**.** *If the curvature $\mathcal{F} = \mathrm{d}\mathcal{A} + \mathcal{A} \wedge \mathcal{A}$ vanishes on a simply connected open set (Definition 4.56), then $\mathcal{U}(x \leftarrow x'; \gamma)$ depends only on endpoints up to endpoint gauge (Proposition 4.59).*

Remark 4.252 (Classical dyad and emergent mixing)**.** The twofold subbundle $\mathcal{H}^{(0,1)}$ recovers the classical dyad at each fibre, and coupling to the (p, s) sector is exactly measured by the off-diagonal Fourfold channel blocks as in Proposition 4.246.

Remark 4.253 (Modal meaning of the fibers)**.** The fiber $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ over $x \in \mathcal{X}_q$ does not encode classical kinematics (e.g. position/momentum); rather, it carries the *internal modal content* of the PQS $\mathfrak{B}_x$ via the Fourfold Dialectic Algebra. After the inner product is fixed, choose an orthonormal frame

$$\mathcal{B}_x = \{|0\rangle_x,\ |1\rangle_x,\ |p\rangle_x,\ |s\rangle_x\},$$

realizing Symmetry, Antisymmetry, Parasymmetry, and Holosymmetry as *independent* directions. In particular, $|p\rangle_x$ and $|s\rangle_x$ are not defined as linear combinations of $|0\rangle_x, |1\rangle_x$ (cf. Remark 4.240); the familiar equal-superposition $|+\rangle_x = \frac{1}{\sqrt{2}}(|0\rangle_x + |1\rangle_x)$ lives in the twofold sector and defines the *parasymmetric effect* $P_{\mathrm{para}}^{(x)} = |+\rangle_x\langle+|$, not the fourfold basis vector $|p\rangle_x$.

This internal, dialectical structure differs from permutation (anti)symmetry in many-body quantum mechanics [373]: it is an *intrinsic* attribute of a single pre-geometric quantum locus. The viewpoint aligns with algebraic QFT [230, 374], where internal state spaces are distinct from Fock-space constructions: the fibers supply kinematic labels and algebraic seeds from which the full dynamical symmetry content is generated.

Remark 4.254 (Categorical structure of the modal bundle)**.** The bundle admits a succinct categorical description as a presheaf of Hilbert spaces:

$$\mathscr{H} : \mathbf{Open}(\mathcal{X}_q)^{\mathrm{op}} \longrightarrow \mathbf{Hilb}, \qquad U \mapsto \Gamma(U, \mathscr{H}_{\mathrm{mod}}),$$

with inclusions $U \hookrightarrow V$ mapped to restriction morphisms $\Gamma(V, \mathscr{H}_{\mathrm{mod}}) \to \Gamma(U, \mathscr{H}_{\mathrm{mod}})$. This functorial viewpoint packages local trivializations, gauge transition data $\{g_{UV}\}$, and parallel transport as natural transformations, and it provides the natural setting to formulate sheaf/cohomological constraints on global sections and holonomy.

Conceptual takeaway. Interpreted functorially, the laws are the *coherence conditions* linking topology (covers and inclusions of $\mathcal{X}_q$) to quantum structure (sections and their restrictions). Parallel transport, connection, and curvature are the intrinsic consistency data of this map. Thus local modal amplitudes and global geometric structure are two faces of the same relational object: a globally coherent assignment $U \mapsto \Gamma(U, \mathscr{H}_{\mathrm{mod}})$ compatible with the Fourfold algebra, gauge covariance, and the graded selection rules.

4.6.2 Wordwise Modal Bundle, Frames, and Trivializations

Definition 4.255 (Word evaluation functor on channels)**.** Fix a device-level word $\mathbf{W} \in \langle \mathsf{A}_{\mathrm{holo}} \rangle$ and a control transition $\chi_{\mathrm{in}} \to \chi_{\mathrm{out}}$. Let $\mathsf{W}_{\chi_{\mathrm{out}} \leftarrow \chi_{\mathrm{in}}}$ be the RR matrix element (Definition 4.148 and eq. (4.45)). For each pair (x, x') define the *word-evaluated* Fourfold channel matrix

$$\hat{\mathbf{G}}_\Lambda^{[\mathbf{W};\, \chi_{\mathrm{out}} \leftarrow \chi_{\mathrm{in}}]}(x, x') := \Big[\hat{G}_\Lambda^{\alpha\beta}(x, x'; \mathbf{W}, \chi_{\mathrm{out}} \leftarrow \chi_{\mathrm{in}}) \Big]_{\alpha,\beta \in \{0,1,p,s\}} \quad (\textit{Definition 4.166 and eq. (4.66)}). \tag{4.94}$$

This defines a functor $\mathcal{E}_{(\mathbf{W},\chi)} : \hat{\mathbf{G}}_\Lambda \mapsto \hat{\mathbf{G}}_\Lambda^{[\mathbf{W};\, \chi \leftarrow \chi]}$ on channel block-matrices.

Lemma 4.256 (Functoriality, gauge covariance, and unitarity)**.** *For admissible words* $\mathbf{W}_1, \mathbf{W}_2$, $\mathcal{E}_{(\mathbf{W}_2\mathbf{W}_1,\chi)} = \mathcal{E}_{(\mathbf{W}_2,\chi)} \circ \mathcal{E}_{(\mathbf{W}_1,\chi)}$. *Under endpoint gauge* $g : \mathcal{X}_q \to \mathrm{U}(\mathcal{H}_{\mathrm{em}})$,

$$\hat{\mathbf{G}}_\Lambda^{[\mathbf{W};\, \chi \leftarrow \chi]}(x, x') \mapsto g(x)\, \hat{\mathbf{G}}_\Lambda^{[\mathbf{W};\, \chi \leftarrow \chi]}(x, x')\, g(x')^\dagger \quad (\textit{Proposition 4.54 and Corollary 4.64}).$$

If $\mathbf{W}$ *is built from unitary elements (control gates and lifted RR histories), then* $\mathsf{W}_{\chi \leftarrow \chi}(x) \in \mathrm{U}(4)$ *acts unitarily on the fibre at* x.

Definition 4.257 (Wordwise modal frame)**.** Let $\mathcal{B}_x = \{|0\rangle_x, |1\rangle_x, |p\rangle_x, |s\rangle_x\}$ be the emergent orthonormal frame (Proposition 4.244). For a fixed $(\mathbf{W}, \chi)$, define the *wordwise rotated frame* at x by

$$|\alpha\rangle_x^{[\mathbf{W},\chi]} := \mathsf{W}_{\chi \leftarrow \chi}(x)\, |\alpha\rangle_x, \qquad \alpha \in \{0, 1, p, s\}.$$

Proposition 4.258 (Wordwise orthonormality and frame transport)**.** *If* $\mathbf{W}$ *is unitary on RR, then the frame* $\mathcal{B}_x^{[\mathbf{W},\chi]}$ *is orthonormal and*

$$\langle \alpha_x^{[\mathbf{W},\chi]} \mid \beta_x^{[\mathbf{W},\chi]} \rangle = \delta_{\alpha\beta} \quad (\textit{preserving eq. (4.92)}).$$

Along a path $\gamma : x' \to x$, transported frames relate by

$$|\alpha\rangle_x^{[\mathbf{W},\chi]} = \mathcal{U}(x \leftarrow x';\gamma)\,|\alpha\rangle_{x'}^{[\mathbf{W},\chi]} \quad \textit{for all } \alpha,$$

since $\mathsf{W}_{\chi\leftarrow\chi}$ acts by endpoint conjugation and $\mathcal{U}$ is unitary (Definition 4.248 and Proposition 4.249).

Definition 4.259 (Wordwise structure words and trivializations)**.** Fix $U \subset \mathcal{X}_q$. A *wordwise trivialization* on U is a choice of unitary word $\mathbf{W}_U$ and control χ such that the map

$$\Phi_U :\ \pi^{-1}(U) \to U \times \mathbb{C}^4, \qquad (x,\ v)\ \mapsto\ \Big(x,\ \mathsf{W}_{U,\chi\leftarrow\chi}(x)\,v\Big)$$

is a unitary fibre isomorphism. On overlaps $U \cap V$, set

$$g_{UV}^{[\mathrm{word}]}(x)\ :=\ \mathsf{W}_{U,\chi\leftarrow\chi}(x)\,\mathsf{W}_{V,\chi\leftarrow\chi}(x)^{\dagger} \in \mathrm{U}(4).$$

Proposition 4.260 (Wordwise transition cocycle and structure group)**.** *The wordwise transition maps satisfy $g_{UU}^{[\mathrm{word}]} = \mathbf{1}$, $g_{UV}^{[\mathrm{word}]} = (g_{VU}^{[\mathrm{word}]})^{-1}$, and $g_{UV}^{[\mathrm{word}]} g_{VW}^{[\mathrm{word}]} = g_{UW}^{[\mathrm{word}]}$ pointwise on triple overlaps. Hence the* wordwise structure group $G_{\mathrm{word}} \subseteq \mathrm{U}(4)$ *is realized as $\{\mathsf{W}_{\chi\leftarrow\chi}(x) :$ $\mathbf{W}$ unitary word$\}$, and the bundle is locally trivial via wordwise trivializations, compatible with eq. (4.93). If a grading is imposed, restrict to words whose RR elements preserve the $(0,1)\oplus(p,s)$ split, so $G_{\mathrm{word}} \subseteq \mathrm{U}(2)\times\mathrm{U}(2)$.*

Full proof. Let ℓ_ε be a smooth, contractible loop based at x, bounding an oriented disk Σ_ε of area $O(\varepsilon^2)$ and diameter $O(\varepsilon)$. By endpoint gauge covariance (Proposition 4.54), fix a local gauge on a neighbourhood $U \ni x$ with

$$\boxed{\mathcal{A}_\mu(x) = 0, \qquad \partial_\alpha \mathcal{A}_\mu(x)\ \text{ is bounded on } U.} \tag{4.95}$$

Write the loop holonomy $\mathcal{U}_{\ell_\varepsilon}(x) := \mathcal{P}\exp\Big(-\oint_{\ell_\varepsilon}\mathcal{A}\Big) \in \mathrm{U}(4)$ (Definition 4.56).

Step 1 (Small–loop expansion). Use the Dyson expansion of the path–ordered exponential together with Stokes:

$$\mathcal{U}_{\ell_\varepsilon}(x) = \mathbf{1}_4 - \int_{\Sigma_\varepsilon}\Big(\mathrm{d}\mathcal{A} + \mathcal{A}\wedge\mathcal{A}\Big)(y)\ +\ R_\varepsilon = \mathbf{1}_4 - \int_{\Sigma_\varepsilon}\mathcal{F}(y)\ +\ R_\varepsilon,$$

with $\mathcal{F} = \mathrm{d}\mathcal{A} + \mathcal{A}\wedge\mathcal{A}$ the curvature (Definition 4.56). In the gauge (4.95), $\mathcal{A}(y) = O(\varepsilon)$ on ℓ_ε; hence $\int_{\Sigma_\varepsilon}\mathcal{A}\wedge\mathcal{A} = O(\varepsilon^3)$ (area×diameter). Taylor expand $\mathcal{F}(y) = \mathcal{F}(x) + O(\varepsilon)$ on Σ_ε, and decompose the oriented area bivector as

$$\Sigma^{\mu\nu}(\ell_\varepsilon)\ :=\ \int_{\Sigma_\varepsilon} d\sigma^{\mu\nu}\ =\ \varepsilon^2\,\sigma^{\mu\nu}(\ell_\varepsilon), \qquad \sigma^{\mu\nu}(\ell_\varepsilon) = O(1).$$

Define the (area–normalised) linear functional

$$\boxed{\Phi_x(\mathcal{F})\ :=\ \sum_{\mu<\nu}\sigma^{\mu\nu}(\ell_\varepsilon)\,\mathcal{F}_{\mu\nu}(x),} \tag{4.96}$$

which is $O(1)$ and *independent* of ε at fixed loop shape. Then

$$\boxed{\begin{aligned}\mathcal{U}_{\ell_\varepsilon}(x) &= \mathbf{1}_4 - \int_{\Sigma_\varepsilon} \mathcal{F}(x) \;+\; O(\varepsilon^3) \\ &= \mathbf{1}_4 - \varepsilon^2\, \Phi_x(\mathcal{F}) \;+\; O(\varepsilon^3).\end{aligned}} \tag{4.97}$$

The commutator role of $\mathcal{F}$ is consistent with the curvature identity $[\mathbb{D}_\mu^{[\cdot]}, \mathbb{D}_\nu^{[\cdot]}]X = [\mathcal{F}_{\mu\nu}, X]$ (Proposition 4.224).

Step 2 (Insert into the wordwise s–loop operator). By eq. (4.122),

$$\hat{H}_{\ell_\varepsilon}^{(s)[\mathbf{W}]}(x) = \Big(\hat{\mathcal{F}}_s^{[\mathbf{W};\chi\leftarrow\chi]}(x)\Big)^\dagger\, \mathcal{U}_{\ell_\varepsilon}(x)\, \hat{\mathcal{F}}_s^{[\mathbf{W};\chi\leftarrow\chi]}(x).$$

Substitute (4.97):

$$\boxed{\hat{H}_{\ell_\varepsilon}^{(s)[\mathbf{W}]}(x) = \Big(\hat{\mathcal{F}}_s^{[\mathbf{W};\chi\leftarrow\chi]}(x)\Big)^\dagger \Big(\mathbf{1}_4 - \varepsilon^2\, \Phi_x(\mathcal{F}) + O(\varepsilon^3)\Big) \hat{\mathcal{F}}_s^{[\mathbf{W};\chi\leftarrow\chi]}(x).} \tag{4.98}$$

Gauge covariance of $\mathcal{U}$ and the endpoint fields (Propositions 4.54 and 4.294 and Corollary 4.64) implies $\hat{H}_{\ell_\varepsilon}^{(s)[\mathbf{W}]}(x)$ transforms by conjugation.

Step 3 (Expectation and invariance). Take the Holo–State expectation (eq. (4.123)):

$$\boxed{\begin{aligned} h_{\ell_\varepsilon}^{(s)[\mathbf{W}]}(x) &= \omega^{\mathring{\mathbf{\Psi}}}\Big((\hat{\mathcal{F}}_s^{[\cdot]})^\dagger \hat{\mathcal{F}}_s^{[\cdot]}\Big) \;-\; \varepsilon^2\, \omega^{\mathring{\mathbf{\Psi}}}\Big((\hat{\mathcal{F}}_s^{[\cdot]})^\dagger\, \Phi_x(\mathcal{F})\, \hat{\mathcal{F}}_s^{[\cdot]}\Big) \;+\; O(\varepsilon^3) \\ &=:\; h_{\mathrm{flat}}^{(s)[\mathbf{W}]}(x) \;-\; \varepsilon^2\, \omega^{\mathring{\mathbf{\Psi}}}\Big((\hat{\mathcal{F}}_s^{[\cdot]})^\dagger\, \Phi_x(\mathcal{F})\, \hat{\mathcal{F}}_s^{[\cdot]}\Big) \;+\; O(\varepsilon^3). \end{aligned}} \tag{4.99}$$

Since the scalar expectation is invariant under endpoint gauge and word changes act by fibrewise unitary conjugation (Propositions 4.284 and 4.294), $h_{\ell_\varepsilon}^{(s)[\mathbf{W}]}(x)$ is gauge invariant. In the flat, simply connected case $\mathcal{F} \equiv 0$ one recovers $h_{\ell_\varepsilon}^{(s)[\mathbf{W}]}(x) = h_{\mathrm{flat}}^{(s)[\mathbf{W}]}(x)$ (Proposition 4.295).

Step 4 (Remainder bound). Let $M_1 = \sup_U \|\partial\mathcal{A}\|$ and $M_2 = \sup_U \big(\|\mathcal{A}\|^2 + \|\partial\mathcal{F}\|\big)$. Then the Dyson remainder satisfies $\|R_\varepsilon\| \le C\,(\text{area}) \times (\text{diameter}) \le C'(M_1, M_2)\,\varepsilon^3$, justifying the $O(\varepsilon^3)$ terms in (4.97)–(4.99). This completes the proof. □

4.6.3 Wordwise Channels, Mixing Bounds, and Kinematics

Definition 4.261 (Wordwise twofold/para–holo split)**.** Define the word-rotated subspaces at x

$$\mathcal{H}_x^{(0,1)[\mathbf{W}]} := \mathrm{span}\{|0\rangle_x^{[\mathbf{W},\chi]}, |1\rangle_x^{[\mathbf{W},\chi]}\}, \qquad \mathcal{H}_x^{(p,s)[\mathbf{W}]} := \mathrm{span}\{|p\rangle_x^{[\mathbf{W},\chi]}, |s\rangle_x^{[\mathbf{W},\chi]}\}.$$

Proposition 4.262 (Wordwise mixing controlled by off-diagonal blocks)**.** *Let* $\hat{\mathbf{G}}_\Lambda^{[\mathbf{W}]}(x,x')$ *denote the word-evaluated channel matrix (Definition 4.255). Then, for unit vectors* $u \in \mathcal{H}_{x'}^{(0,1)[\mathbf{W}]}$, $v \in \mathcal{H}_x^{(p,s)[\mathbf{W}]}$,

$$\Big|\langle v \mid \hat{\mathbf{G}}_\Lambda^{[\mathbf{W}]}(x,x')\, u\rangle\Big| \;\le\; \sum_{\substack{\alpha\in\{p,s\}\\ \beta\in\{0,1\}}} \Big\|\hat{G}_\Lambda^{\alpha\beta}(x,x';\mathbf{W},\chi\leftarrow\chi)\Big\|.$$

In particular, if all such off-diagonal blocks vanish, the split $\mathcal{H}^{(0,1)[\mathbf{W}]} \oplus \mathcal{H}^{(p,s)[\mathbf{W}]}$ is invariant and no mixing occurs.

Proof. Expand in the word-rotated frames and apply Cauchy–Schwarz and operator norm bounds, as in Proposition 4.246. □

Definition 4.263 (Wordwise covariant derivative and curvature)**.** Using the connection $\mathcal{A}$ on the base (Definition 4.248), define for each word-evaluated component the derivative

$$\mathbb{D}_\mu^{[\mathbf{W}]} \hat{G}^{\alpha\beta} := \partial_\mu \hat{G}^{\alpha\beta} + [\,\mathcal{A}_\mu, \hat{G}^{\alpha\beta}\,],$$

and along a path γ the path-derivative $\partial_s + [A_\gamma(s), \cdot\,]$ with $A_\gamma = (\gamma^*\mathcal{A})(\dot\gamma)$ (Section 4.5, Definition 4.220, and Proposition 4.223). The curvature is $\mathcal{F} = \mathrm{d}\mathcal{A} + \mathcal{A} \wedge \mathcal{A}$ (Definition 4.56).

Proposition 4.264 (Wordwise curvature identity and flatness)**.** *For any word-evaluated channel X, $[\mathbb{D}_\mu^{[\mathbf{W}]}, \mathbb{D}_\nu^{[\mathbf{W}]}]X = [\mathcal{F}_{\mu\nu}, X]$ (Proposition 4.224). If $\mathcal{F} = 0$ on a simply connected region, parallel consistency integrates to word–path independence up to endpoint gauge (Propositions 4.59 and 4.223).*

Remark 4.265 (Dictionary back to the exogenous bundle)**.** The endogenous objects above reproduce the exogenous formalization via:

1. frames $\mathcal{B}_x^{[\mathbf{W},\chi]} \leftrightarrow$ frames $\mathcal{B}_x$ acted on by $G \subseteq \mathrm{U}(4)$ (Definitions 4.247 and 4.257 and Proposition 4.258);
2. transition words $g_{UV}^{[\mathrm{word}]} \leftrightarrow$ transition functions g_{UV} (Definition 4.259 and Proposition 4.260);
3. word-evaluated channels $\hat{\mathbf{G}}_\Lambda^{[\mathbf{W}]} \leftrightarrow$ channel blocks $\hat{\mathbf{G}}_\Lambda$ (Definition 4.255);
4. wordwise kinematics $\mathbb{D}_\mu^{[\mathbf{W}]}, \partial_s + [A_\gamma, \cdot\,] \leftrightarrow$ graded/exogenous kinematics $\mathfrak{D}_\mu, D_s$ (Section 4.5, Definition 4.134, and Proposition 4.137);
5. mixing bound Proposition 4.262 $\leftrightarrow$ Proposition 4.246.

4.6.4 Global Section of Dialectic Modes

Definition 4.266 (Global section of dialectic modes)**.** Let $(\mathscr{H}_{\mathrm{mod}}, \pi, \mathcal{X}_q)$ be the Hilbert bundle of dialectic modes from Definition 4.239. A *global section* is a smooth map

$$\boxed{|\Psi_{\mathrm{modal}}\rangle : \ \mathcal{X}_q \longrightarrow \mathscr{H}_{\mathrm{mod}}, \qquad x \ \longmapsto \ |\Psi_{\mathrm{modal}}(x)\rangle \in \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)},} \tag{4.100}$$

satisfying the section condition $\pi \circ |\Psi_{\mathrm{modal}}\rangle = \mathrm{id}_{\mathcal{X}_q}$. On any trivializing patch $U \subset \mathcal{X}_q$ with orthonormal frame $\mathcal{B}_x = \{|0\rangle_x, |1\rangle_x, |p\rangle_x, |s\rangle_x\}$ (cf. eq. (4.92)), the section has the local expansion

$$|\Psi_{\mathrm{modal}}(x)\rangle = \sum_{\alpha \in \{0,1,p,s\}} c_\alpha(x)\, |\alpha\rangle_x \quad (x \in U), \tag{4.101}$$

with smooth coefficients $c_\alpha : U \to \mathbb{C}$ collected as $c(x) := (c_0, c_1, c_p, c_s)^\top$. Pointwise normalization in each fiber is

$$\boxed{c(x)^\dagger c(x) = |c_0(x)|^2 + |c_1(x)|^2 + |c_p(x)|^2 + |c_s(x)|^2 = 1 \qquad (\forall x \in U).} \tag{4.102}$$

On an overlap $U\cap V$, coefficient vectors are related by the transition matrix $g_{UV}(x) \in G \subseteq U(4)$ (preserving the ordered mode frame and, when invoked, the $\mathbb{Z}_2 \times \mathbb{Z}_2$ grading):

$$\boxed{c_U(x) \;=\; g_{UV}(x)\, c_V(x), \qquad x \in U \cap V,} \tag{4.103}$$

so (4.102) is gauge invariant.

Remark 4.267 (Gauge–Covariant Differentiation and Transport of a Section)**.** Let $\mathcal{A}$ be the connection one–form introduced in Section 4.1.4, with transporter $\mathcal{U} = \mathcal{P}\exp(-\int \mathcal{A})$. The *covariant derivative* of a section is

$$\boxed{\mathfrak{D}_\mu|\Psi_{\text{modal}}(x)\rangle := \partial_\mu|\Psi_{\text{modal}}(x)\rangle + \mathcal{A}_\mu(x)\,|\Psi_{\text{modal}}(x)\rangle,} \tag{4.104}$$

which transforms homogeneously under $g(x) \in G$ and preserves the pointwise norm $c(x)^\dagger c(x)$. Parallel sections obey $\mathfrak{D}_\mu|\Psi_{\text{modal}}\rangle = 0$, and holonomy is generated by the curvature $\mathcal{F} = \mathrm{d}\mathcal{A} + \mathcal{A}\wedge\mathcal{A}$.

Remark 4.268 (Fiberwise Expectations and Density)**.** The fiberwise density operator of a section is $\rho_{\text{modal}}(x) := |\Psi_{\text{modal}}(x)\rangle\langle\Psi_{\text{modal}}(x)|$. For any fiber observable $\hat{O}(x)$ (e.g. mode projectors), the local expectation $\langle\hat{O}\rangle(x) := \mathrm{Tr}\big(\rho_{\text{modal}}(x)\hat{O}(x)\big)$ is gauge invariant. Coupling to inter–locus physics proceeds via the Fourfold correlator and the transporter (cf. Sections 4.1.2 and 4.1.4).

Remark 4.269 (Independence of p, s from Twofold Superpositions)**.** The coefficients $c_p(x)$ and $c_s(x)$ weight *independent* fiber directions $|p\rangle_x$ and $|s\rangle_x$. They are not fixed by twofold superpositions (see Remark 4.240); in particular $|+\rangle_x = \frac{1}{\sqrt{2}}(|0\rangle_x + |1\rangle_x)$ defines a twofold *effect*, not the fourfold basis vector $|p\rangle_x$.

Lemma 4.270 (Fiberwise Pairing is Well Defined, Gauge Invariant, and Covariantly Leibniz)**.** *Let $|\Phi_{\text{modal}}\rangle, |\Psi_{\text{modal}}\rangle$ be global sections of $(\mathscr{H}_{\text{mod}}, \pi, \mathcal{X}_q)$, and let $\langle\cdot,\cdot\rangle_x$ denote the inner product on the fiber $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}$. Choose a local trivialization (U, e_U) with orthonormal frame $e_U(x) = (|0\rangle_x, |1\rangle_x, |p\rangle_x, |s\rangle_x)$, and write $|\Phi_{\text{modal}}(x)\rangle = e_U(x)c_U^\Phi(x)$, $|\Psi_{\text{modal}}(x)\rangle = e_U(x)c_U^\Psi(x)$ with column maps $c_U^\Phi, c_U^\Psi : U \to \mathbb{C}^4$.*

Then, for each $x \in \mathcal{X}_q$:

1. Well defined and gauge invariant. *On overlaps $U\cap V$, let $e_V(x) = e_U(x)\, g_{UV}(x)$ with $g_{UV} : U\cap V \to G \subseteq U(4)$. Then $c_V^\Phi = g_{UV}^{-1}c_U^\Phi$, $c_V^\Psi = g_{UV}^{-1}c_U^\Psi$ and*

$$\langle\Phi_{\text{modal}}(x), \Psi_{\text{modal}}(x)\rangle_x = (c_U^\Phi(x))^\dagger c_U^\Psi(x) = (c_V^\Phi(x))^\dagger c_V^\Psi(x),$$

so the scalar function $x \mapsto \langle\Phi_{\text{modal}}(x), \Psi_{\text{modal}}(x)\rangle_x$ is independent of trivialization and invariant under G–gauge frame changes.

2. Covariant Leibniz rule (metric compatibility). *Let $\mathcal{A}_\mu$ be the connection one–form of Section 4.1.4 and $\mathfrak{D}_\mu$ the covariant derivative of eq. (4.104), with $\mathcal{A}_\mu^\dagger = -\mathcal{A}_\mu$ (skew–Hermitian). Then, for all μ,*

$$\boxed{\begin{aligned}\partial_\mu\,\langle\Phi_{\text{modal}}(x), \Psi_{\text{modal}}(x)\rangle_x = &\Big\langle \mathfrak{D}_\mu\Phi_{\text{modal}}(x),\, \Psi_{\text{modal}}(x)\Big\rangle_x \\ &+ \Big\langle \Phi_{\text{modal}}(x),\, \mathfrak{D}_\mu\Psi_{\text{modal}}(x)\Big\rangle_x.\end{aligned}} \tag{4.105}$$

In particular, along any curve $\gamma : \lambda \mapsto x(\lambda)$, if $\mathfrak{D}_{\dot\gamma}|\Psi_{\text{modal}}\rangle = 0$ (parallel transport), then $\frac{d}{d\lambda}\langle\Psi_{\text{modal}}, \Psi_{\text{modal}}\rangle = 0$: norms and orthonormality are preserved by parallel transport.

Proof. Fix a trivialising patch $U \subset \mathcal{X}_q$ with an orthonormal modal frame at $x \in U$,

$$e_U(x) \coloneqq \Big(|0\rangle_x, |1\rangle_x, |p\rangle_x, |s\rangle_x\Big).$$

Write the coefficient column vectors $c_U^\Phi(x), c_U^\Psi(x) \in \mathbb{C}^4$, and set

$$|\Phi_{\text{modal}}(x)\rangle = e_U(x)\, c_U^\Phi(x), \qquad |\Psi_{\text{modal}}(x)\rangle = e_U(x)\, c_U^\Psi(x).$$

With the fibre inner product $\langle\cdot,\cdot\rangle_x$ (induced as in Definition 4.228) and the orthonormality of $e_U(x)$, we obtain

$$\Big\langle \Phi_{\text{modal}}(x), \Psi_{\text{modal}}(x)\Big\rangle_x = \Big(c_U^\Phi(x)\Big)^\dagger c_U^\Psi(x).$$

On an overlap $U \cap V \neq \varnothing$ let $e_V(x) = e_U(x)\, g_{UV}(x)$ with $g_{UV}(x) \in G \subseteq U(4)$. Then coefficients transform as $c_V^\Phi(x) = g_{UV}(x)^{-1} c_U^\Phi(x)$ and $c_V^\Psi(x) = g_{UV}(x)^{-1} c_U^\Psi(x)$, hence

$$(c_V^\Phi(x))^\dagger c_V^\Psi(x) = (c_U^\Phi(x))^\dagger\, (g_{UV}^{-1}(x))^\dagger g_{UV}^{-1}(x)\, c_U^\Psi(x) = (c_U^\Phi(x))^\dagger c_U^\Psi(x),$$

proving the gauge/trivialization invariance in (1). Equivalently, a local frame change $g(x) \in G$ acting by $|\Phi\rangle \mapsto g|\Phi\rangle$, $|\Psi\rangle \mapsto g|\Psi\rangle$ preserves the fiber pairing since $g^\dagger g = \mathbf{1}$.

For (2), use the covariant derivative $\mathfrak{D}_\mu = \partial_\mu + \mathcal{A}_\mu$ with $\mathcal{A}_\mu^\dagger = -\mathcal{A}_\mu$. In a local frame,

$$\partial_\mu\Big[(c_U^\Phi(x))^\dagger c_U^\Psi(x)\Big] = (\mathfrak{D}_\mu c_U^\Phi(x))^\dagger c_U^\Psi(x) + (c_U^\Phi(x))^\dagger (\mathfrak{D}_\mu c_U^\Psi(x)),$$

because the cross terms cancel: $(c_U^\Phi)^\dagger \mathcal{A}_\mu c_U^\Psi + (\mathcal{A}_\mu c_U^\Phi)^\dagger c_U^\Psi = (c_U^\Phi)^\dagger(\mathcal{A}_\mu + \mathcal{A}_\mu^\dagger) c_U^\Psi = 0$. Translating back to fiber pairings gives (4.105). Taking $\Phi = \Psi$ and restricting to a curve γ with $\mathfrak{D}_{\dot\gamma}|\Psi_{\text{modal}}\rangle = 0$ yields $\frac{d}{d\lambda}\langle\Psi_{\text{modal}}, \Psi_{\text{modal}}\rangle = 0$; norms and orthonormality are preserved by parallel transport. □

Remark 4.271 (Unitary Frames and h-Orthonormal Generalisation)**.** (i) In an orthonormal trivialisation (fibre metric $h = \mathbb{1}_4$), it is convenient to write the identity as $\mathbb{1}_4$ to emphasise rank, so for any frame change $g \in U(4)$ one has $g^\dagger g = \mathbb{1}_4$.

(ii) In a non-orthonormal trivialisation with fibre metric $h(x)$ (a smooth, positive-definite Hermitian form), the structure group reduces to

$$G_h \coloneqq \{\, g \in \mathrm{GL}(4,\mathbb{C}) \mid g^\dagger h\, g = h \,\}.$$

Frames transform as $e_V = e_U g$ and coefficients as $c_V = g^{-1} c_U$, and the fibre pairing is

$$\langle \Phi_{\text{modal}}, \Psi_{\text{modal}}\rangle_x = (c_U^\Phi)^\dagger h\, c_U^\Psi = (c_V^\Phi)^\dagger h\, c_V^\Psi,$$

since $(g^{-1})^\dagger h\, g^{-1} = h$ follows from $g^\dagger h g = h$. If the connection $\mathcal{A}_\mu$ is metric–compatible,

$$\partial_\mu h \;-\; \mathcal{A}_\mu^\dagger h \;-\; h\,\mathcal{A}_\mu \;=\; 0,$$

then with $\mathfrak{D}_\mu = \partial_\mu + \mathcal{A}_\mu$ one has the covariant Leibniz rule

$$\partial_\mu\Big[(c_U^\Phi)^\dagger h\, c_U^\Psi\Big] = (\mathfrak{D}_\mu c_U^\Phi)^\dagger h\, c_U^\Psi + (c_U^\Phi)^\dagger h\, (\mathfrak{D}_\mu c_U^\Psi), \tag{4.106}$$

which reduces to eq. (4.105) when $h = \mathbb{1}_4$.

Corollary 4.272 (Gauge Invariance and Parallel Constancy of Mode Weights)**.** *Let* $P_\alpha(x) = |\alpha\rangle_x\langle\alpha|_x$ *be the rank-one projectors onto the modal basis* $\{|0\rangle_x, |1\rangle_x, |p\rangle_x, |s\rangle_x\}$*. For any global section* $|\Psi_{\text{modal}}\rangle$ *define* $w_\alpha(x) := \langle\Psi_{\text{modal}}(x),\, P_\alpha(x)\,\Psi_{\text{modal}}(x)\rangle_x = |c_\alpha(x)|^2$*. Then:*

(1) $w_\alpha(x)$ *is independent of the choice of trivialization and invariant under local* G*–frame changes on overlaps.*

(2) *Along a curve* γ*, if the state is parallel transported* $\mathfrak{D}_{\dot\gamma}|\Psi_{\text{modal}}\rangle = 0$ and *the projectors are propagated by the same connection (i.e.* $\mathfrak{D}_{\dot\gamma}P_\alpha = 0$*, equivalently the modal frame is chosen covariantly constant along* γ*), then* $\frac{d}{d\lambda}\, w_\alpha(\gamma(\lambda)) = 0$*.*

Physically, a global section $|\Psi_{\text{modal}}(x)\rangle$ is the spatially varying **dialectical profile** of a quantum field over $\mathcal{X}_q$. By specifying the local amplitudes of all four modes at each locus, it can encode either an emergent vacuum configuration or a particle-like excitation. Mathematically, this mirrors the way sections of spinor bundles represent fermion fields and sections of gauge bundles encode gauge potentials in geometric field theory [375, 225].

Global sections as dialectic quantum fields. Unlike the conventional view in which a composite quantum system lives in a fixed tensor power $\mathcal{H}_2^{\otimes n}$, the present framework is intrinsically geometric and operator-algebraic [4, 241]. A system of n distinguishable entities is modeled by n Primordial Quantum Singularities (PQSs) $\mathfrak{B}_{x_1}, \ldots, \mathfrak{B}_{x_n}$ located at points $x_i \in \mathcal{X}_q$, each equipped with its own *modal Hilbert fiber* $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_{x_i})}$ whose inner product is *emergent* and state-dependent [11]. In this setting, a quantum field is a *global section* of the Hilbert bundle of dialectic modes: a smooth assignment

$$x \longmapsto |\Psi_{\text{modal}}(x)\rangle \in \mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)},$$

selecting at each point a superposition of the four dialectical modes.

At any $x \in \mathcal{X}_q$, the local modal state has the form

$$|\Psi_{\text{modal}}(x)\rangle = c_0(x)\,|\mathring{\mathbf{0}}\rangle_x + c_1(x)\,|\mathring{\mathbf{1}}\rangle_x + c_p(x)\,|\mathring{\mathbf{0}} + \mathring{\mathbf{1}}\rangle_x + c_s(x)\,\big|_{\mathring{\mathbf{0}}}\mathring{\Lambda}_{\mathring{\mathbf{1}}}\big\rangle_x,$$

where the orthonormal basis encodes the Fourfold Dialectic Algebra (Symmetry/Thesis, Antisymmetry/Antithesis, Parasymmetry/Synthesis, Holosymmetry/Holothesis), and the coefficients $c_\alpha(x) \in \mathbb{C}$ satisfy the pointwise normalization

$$|c_0(x)|^2 + |c_1(x)|^2 + |c_p(x)|^2 + |c_s(x)|^2 = 1.$$

The map $x \mapsto (c_0, c_1, c_p, c_s)(x)$ determines the field's "shape" across $\mathcal{X}_q$, from uniform dialectical vacua to localized excitations propagating on the quantum-relational substrate.

This generalizes familiar constructions in modern field theory: just as spinor and gauge fields are sections of their respective bundles [225, 375], a global section of the modal Hilbert bundle furnishes a fundamental field of the IQR. It simultaneously captures locality (one fiber per PQS) and the global logic of duality that organizes their modal evolution. Gauge-covariant comparison between fibers is mediated by the parallel transporter, and fiberwise pairings are well defined, gauge invariant, and preserved by parallel transport (cf. Lemma 4.270). In this way, the total fibered space (see Definition 4.239) becomes the stage on which the dialectical dynamics of the quantum universe unfold, with the global section providing the field that inhabits it.

Definition 4.273 (Modal Hilbert Space of a Primordial Quantum Singularity)**.** Fix a locus $x \in \mathcal{X}_q$ with associated Primordial Quantum Singularity $\mathfrak{B}_x$ (Definition 1.25). Its intrinsic operator-theoretic duality

$$\hat{\mathbf{\Omega}}_x = \overset{\circ}{\underset{\hat{\mathbf{\Phi}}_x}{\mathbf{\Lambda}}}_{x,\hat{\mathbf{\Theta}}_x}$$

encodes the Fourfold Dialectic Algebra (modes: Symmetry, Antisymmetry, Parasymmetry, Holosymmetry). The *modal Hilbert space* at x is the emergent fiber

$$\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)} \cong \mathbb{C}^4,$$

whose inner product is defined relationally by the Fourfold Inter-Modal Correlator in the Holo-State (cf. Section 4.6, (4.85)).

Realization map and orthonormal basis. Let $\mathcal{I} := \{0, 1, p, s\}$ be the mode labels and $\mathcal{R}_x : \mathcal{I} \to \mathbb{P}\big(\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}\big)$ the realization map sending a label to a ray. After the inner product is fixed, choose unit representatives

$$\mathcal{B}_x := \{|0\rangle_x, |1\rangle_x, |p\rangle_x, |s\rangle_x\}, \qquad \mathcal{R}_x(\alpha) = [\,|\alpha\rangle_x\,],$$

and impose the *ontological orthonormality*

$$\boxed{\langle \alpha_x \mid \beta_x \rangle \;=\; \delta_{\alpha\beta}, \qquad \alpha, \beta \in \{0, 1, p, s\}.} \tag{4.107}$$

Caution: $|p\rangle_x$ and $|s\rangle_x$ are independent basis vectors. The equal-superposition $|+\rangle_x := \frac{1}{\sqrt{2}}(|0\rangle_x + |1\rangle_x)$ belongs to the Twofold sector and defines the local parasymmetric *effect* $P_{\text{para}}^{(x)} = |+\rangle_x\langle+|$; it is not identified with $|p\rangle_x$.

Modal decomposition. With the shorthand $\mathcal{H}_{\alpha,x} := \text{span}\{|\,\alpha\rangle_x\}$ one has the direct sum

$$\boxed{\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)} = \mathcal{H}_{0,x} \oplus \mathcal{H}_{1,x} \oplus \mathcal{H}_{p,x} \oplus \mathcal{H}_{s,x}.} \tag{4.108}$$

State representation. Every $|\psi\rangle_x \in \mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}$ admits the expansion

$$|\psi\rangle_x = c_0(x)\,|0\rangle_x + c_1(x)\,|1\rangle_x + c_p(x)\,|p\rangle_x + c_s(x)\,|s\rangle_x,$$

with pointwise normalization

$$|c_0(x)|^2 + |c_1(x)|^2 + |c_p(x)|^2 + |c_s(x)|^2 = 1.$$

Interpretation. The fiber $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}$ is a locally constructed, relationally defined quantum space that carries the irreducible dialectic content at x. It is not a fixed tensor factor but the arena where the four modal facets are represented and compared; its metric is induced by the correlator expectations, ensuring full consistency with gauge-covariant transport and the Fourfold algebraic structure.

Remark 4.274 (On the Interpretation of Dialectic Modes)**.** At each locus $x \in \mathcal{X}_q$, the modal fiber $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ carries an orthonormal basis $\{|0\rangle_x, |1\rangle_x, |p\rangle_x, |s\rangle_x\}$ (cf. (4.92)), representing the four canonical facets of the Fourfold Dialectic Algebra: *Symmetry* (Thesis), *Antisymmetry* (Antithesis), *Parasymmetry* (Synthesis), and *Holosymmetry* (Holothesis). These modes are determined by the internal duality operator $\mathring{\mathbf{\Lambda}}_x$ as it appears inside the fibred operator-theoretic structure $\hat{\mathbf{\Omega}}_x = {}_{\hat{\Phi}_x}\mathring{\mathbf{\Lambda}}_{x,\hat{\Theta}_x}$, and are intrinsic to the singularity $\mathfrak{B}_x$.

It is essential to distinguish this notion of "(anti)symmetry" from the standard permutation-(anti)symmetry under S_n in many-body quantum mechanics, where symmetric/antisymmetric subspaces encode particle-exchange statistics on tensor products [376, 373]. Here, by contrast, the modal basis is a *local, internal* dialectic of a single Primordial Quantum Singularity; it is not obtained by (anti)symmetrizing over identical constituents. This viewpoint aligns with algebraic/categorical treatments in modern quantum theory [230, 377, 374], where internal state spaces are organized fiberwise over configuration spaces and possess their own intrinsic logic and geometry.

A further clarification is crucial for the higher modes. The parasymmetric basis vector $|p\rangle_x$ is *not* defined as the equal-superposition vector $|+\rangle_x := \frac{1}{\sqrt{2}}(|0\rangle_x + |1\rangle_x)$; the latter belongs to the Twofold sector and underlies the local parasymmetric *effect* $P^{(x)}_{\mathrm{para}} = |+\rangle_x\langle+|$, whereas $|p\rangle_x$ is an independent basis ray for the Fourfold algebra. Likewise, $|s\rangle_x$ (holosymmetry) is not assumed to be a linear combination of $|0\rangle_x$ and $|1\rangle_x$; operationally it is accessed via holosymmetric effects that probe implementation-invariant, history-level structure.

In this way the Fourfold Dialectic Algebra supplies more than a convenient labeling scheme: it furnishes a local, ontological basis from which internal geometry, logical duality, and coherence properties are read. Parasymmetric channels register coherence-sensitive synthesis beyond mixtures, while holosymmetric channels register global, implementation-invariant constraints— all within the same fiberwise, operator-algebraic framework.

4.6.5 Wordwise Global Section of Dialectic Modes

Definition 4.275 (Wordwise global section of dialectic modes)**.** Fix a unitary device word $\mathbf{W} \in \langle \mathsf{A}_{\mathrm{holo}} \rangle$ and a control label χ. Let $\mathsf{W}_{\chi\leftarrow\chi}(x)$ be its RR element at x (Definition 4.148 and eq. (4.45)). A *wordwise global section* is a smooth map

$$|\Psi^{[\mathbf{W},\chi]}_{\mathrm{modal}}\rangle : \ \mathcal{X}_q \to \mathscr{H}_{\mathrm{mod}}, \qquad x \mapsto |\Psi^{[\mathbf{W},\chi]}_{\mathrm{modal}}(x)\rangle \in \mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}},$$

that admits on each trivialising patch $U \subset \mathcal{X}_q$ the expansion in the *wordwise frame* $\mathcal{B}^{[\mathbf{W},\chi]}_x = \{|0\rangle^{[\mathbf{W},\chi]}_x, |1\rangle^{[\mathbf{W},\chi]}_x, |p\rangle^{[\mathbf{W},\chi]}_x, |s\rangle^{[\mathbf{W},\chi]}_x\}$ (Definition 4.257):

$$|\Psi^{[\mathbf{W},\chi]}_{\mathrm{modal}}(x)\rangle = \sum_{\alpha\in\{0,1,p,s\}} c^{[\mathbf{W}]}_\alpha(x)\, |\alpha\rangle^{[\mathbf{W},\chi]}_x \quad (x \in U), \tag{4.109}$$

with smooth coefficients $c^{[\mathbf{W}]}_\alpha : U \to \mathbb{C}$.

Proposition 4.276 (Normalization and gauge invariance)**.** *If* $\mathbf{W}$ *is unitary on RR, the wordwise frame is orthonormal (Proposition 4.258), and the coefficient vector* $c^{[\mathbf{W}]}(x) :=$

$(c_0^{[\mathbf{W}]}, c_1^{[\mathbf{W}]}, c_p^{[\mathbf{W}]}, c_s^{[\mathbf{W}]})^\top$ *satisfies the pointwise normalization*

$$\boxed{(c^{[\mathbf{W}]}(x))^\dagger c^{[\mathbf{W}]}(x) = 1 \qquad (\forall x \in U).} \tag{4.110}$$

Under endpoint gauge $g : \mathcal{X}_q \to \mathrm{U}(4)$, $|\Psi^{[\mathbf{W},\chi]}_{\mathrm{modal}}(x)\rangle \mapsto g(x)\,|\Psi^{[\mathbf{W},\chi]}_{\mathrm{modal}}(x)\rangle$ *and* (4.110) *is invariant.*

Proof. Let $\{|\phi_\mu(x)\rangle\}_{\mu\in\{0,1,p,s\}}$ be the wordwise modal frame at x, which is orthonormal by Proposition 4.258. Write the dressed modal state in this frame as

$$|\Psi^{[\mathbf{W},\chi]}_{\mathrm{modal}}(x)\rangle = \sum_{\mu\in\{0,1,p,s\}} c_\mu^{[\mathbf{W}]}(x)\,|\phi_\mu(x)\rangle, \qquad c^{[\mathbf{W}]}(x) = \left(c_0^{[\mathbf{W}]}, c_1^{[\mathbf{W}]}, c_p^{[\mathbf{W}]}, c_s^{[\mathbf{W}]}\right)^\top.$$

Normalization. Since the frame is orthonormal,

$$\left\| \, |\Psi^{[\mathbf{W},\chi]}_{\mathrm{modal}}(x)\rangle \, \right\|^2 = \sum_\mu \left| c_\mu^{[\mathbf{W}]}(x) \right|^2 = \left(c^{[\mathbf{W}]}(x)\right)^\dagger c^{[\mathbf{W}]}(x).$$

It therefore suffices to show $\| \, |\Psi^{[\mathbf{W},\chi]}_{\mathrm{modal}}(x)\rangle\| = 1$. By the wordwise dressing rule (cf. eq. (4.67)) the state is obtained by applying, to a unit-norm input, the product of three factors: (i) the parallel transporter $\mathcal{U}(x \leftarrow x'; \gamma)$ on the base, (ii) the RR–matrix element $\mathsf{W}_{\chi_{\mathrm{out}}\leftarrow\chi_{\mathrm{in}}}$ of $\mathbf{W}$ (cf. Definition 4.148 and eq. (4.45)), and (iii) the orthonormal frame map at x/x'. By assumption, $\mathbf{W}$ is unitary on RR, hence $\mathsf{W}_{\chi\leftarrow\chi}$ is unitary (on $\mathcal{H}^{(2n)}_{\mathrm{RR}}$); $\mathcal{U}$ is unitary by construction; and the frame maps are isometries due to orthonormality. Consequently the composite map is norm-preserving, so $\| \, |\Psi^{[\mathbf{W},\chi]}_{\mathrm{modal}}(x)\rangle\| = 1$ and hence

$$\left(c^{[\mathbf{W}]}(x)\right)^\dagger c^{[\mathbf{W}]}(x) = 1,$$

which is eq. (4.110).

Gauge invariance. Under an endpoint gauge $g : \mathcal{X}_q \to \mathrm{U}(4)$ on the modal fibre, the state transforms as $|\Psi^{[\mathbf{W},\chi]}_{\mathrm{modal}}(x)\rangle \mapsto g(x)\,|\Psi^{[\mathbf{W},\chi]}_{\mathrm{modal}}(x)\rangle$ (cf. gauge covariance in Proposition 4.167(a)). With the column of coefficients $c^{[\mathbf{W}]}(x)$ taken in the *local* frame at x, this is equivalently $c^{[\mathbf{W}]}(x) \mapsto g(x)\,c^{[\mathbf{W}]}(x)$. Since $g(x) \in \mathrm{U}(4)$,

$$\left(c^{[\mathbf{W}]}(x)\right)^\dagger c^{[\mathbf{W}]}(x) \;\longmapsto\; \left(c^{[\mathbf{W}]}(x)\right)^\dagger g(x)^\dagger g(x)\, c^{[\mathbf{W}]}(x) = \left(c^{[\mathbf{W}]}(x)\right)^\dagger c^{[\mathbf{W}]}(x),$$

so eq. (4.110) is gauge invariant pointwise on U.

This establishes Proposition 4.276. □

Definition 4.277 (Wordwise transitions on overlaps)**.** Let $U, V \subset \mathcal{X}_q$ be trivialising patches with wordwise trivializations $\mathbf{W}_U, \mathbf{W}_V$ (Definition 4.259). On $U \cap V$, define the transition word $g^{[\mathrm{word}]}_{UV}(x) = \mathsf{W}_{U,\chi\leftarrow\chi}(x)\,\mathsf{W}_{V,\chi\leftarrow\chi}(x)^\dagger \in \mathrm{U}(4)$ (Proposition 4.260). Then the coefficient vectors obey

$$\boxed{c_U^{[\mathbf{W}]}(x) = g^{[\mathrm{word}]}_{UV}(x)\, c_V^{[\mathbf{W}]}(x) \qquad (x \in U \cap V),} \tag{4.111}$$

so that (4.110) is patchwise consistent.

Remark 4.278 (Wordwise covariant derivative of a section)**.** With the base connection $\mathcal{A}$ (Definition 4.248), define the *wordwise covariant derivative* of the section by

$$\boxed{\mathbb{D}_\mu^{[\mathbf{W}]}|\Psi_{\text{modal}}^{[\mathbf{W},\chi]}(x)\rangle := \partial_\mu|\Psi_{\text{modal}}^{[\mathbf{W},\chi]}(x)\rangle + \mathcal{A}_\mu(x)\,|\Psi_{\text{modal}}^{[\mathbf{W},\chi]}(x)\rangle.} \tag{4.112}$$

This transforms homogeneously under endpoint gauge (Lemma 4.221) and preserves (4.110). A *wordwise parallel* section obeys $\mathbb{D}_\mu^{[\mathbf{W}]}|\Psi_{\text{modal}}^{[\mathbf{W},\chi]}\rangle = 0$.

Lemma 4.279 (Leibniz rule and parallel constancy along a path)**.** *Let γ be a C^1 curve with pullback A_γ. If the word-locked frame condition holds at the readout end (Proposition 4.223) and $\mathbb{D}_{\dot\gamma}^{[\mathbf{W}]}|\Psi_{\text{modal}}^{[\mathbf{W},\chi]}\rangle = 0$, then $\frac{d}{ds}\| \,|\Psi_{\text{modal}}^{[\mathbf{W},\chi]}(\gamma(s))\rangle\,\|^2 = 0$. Moreover, for any fibre observable $\hat O(x)$,*

$$\partial_\mu\langle\hat O\rangle^{[\mathbf{W}]}(x) = \Big\langle(\mathbb{D}_\mu^{[\mathbf{W}]})^\dagger\hat O(x)\Big\rangle^{[\mathbf{W}]} + \Big\langle\hat O(x)\,\mathbb{D}_\mu^{[\mathbf{W}]}\Big\rangle^{[\mathbf{W}]},$$

i.e. a covariant Leibniz rule holds (wordwise version of eq. (4.105)).

Proof. Let $x = \gamma(s)$ and write the dressed modal section in the wordwise orthonormal frame (cf. Proposition 4.258) as

$$|\Psi_{\text{modal}}^{[\mathbf{W},\chi]}(x)\rangle = \sum_{\mu\in\{0,1,p,s\}} c_\mu^{[\mathbf{W}]}(x)\,|\phi_\mu(x)\rangle, \qquad c^{[\mathbf{W}]}(x)\in\mathbb{C}^4.$$

Denote the wordwise connection one-form by $A_\mu^{[\mathbf{W}]}(x)\in\mathfrak{u}(4)$ (unitary gauge, cf. Proposition 4.167(a) and Proposition 4.276), and define the covariant derivative on coefficients

$$\mathbb{D}_\mu^{[\mathbf{W}]}c := \partial_\mu c + A_\mu^{[\mathbf{W}]}c, \qquad \Big(\mathbb{D}_\mu^{[\mathbf{W}]}c\Big)^\dagger = \partial_\mu c^\dagger + c^\dagger(A_\mu^{[\mathbf{W}]})^\dagger.$$

Along γ we write the pullback $A_\gamma(s) := \dot\gamma^\mu(s)\,A_\mu^{[\mathbf{W}]}(\gamma(s))$ and $\mathbb{D}_{\dot\gamma}^{[\mathbf{W}]}c := \frac{d}{ds}c + A_\gamma c$.

Parallel constancy of the norm. Let $N(s) := \| \,|\Psi_{\text{modal}}^{[\mathbf{W},\chi]}(\gamma(s))\rangle\,\|^2 = c^\dagger(\gamma(s))\,c(\gamma(s))$. Then

$$\frac{dN}{ds} = \frac{dc^\dagger}{ds}\,c + c^\dagger\,\frac{dc}{ds} = \Big(\mathbb{D}_{\dot\gamma}^{[\mathbf{W}]}c\Big)^\dagger c + c^\dagger\Big(\mathbb{D}_{\dot\gamma}^{[\mathbf{W}]}c\Big) + c^\dagger\Big(A_\gamma + A_\gamma^\dagger\Big)c.$$

Since $A_\mu^{[\mathbf{W}]}\in\mathfrak{u}(4)$, we have $A_\gamma^\dagger = -A_\gamma$, hence the last term vanishes. If $\mathbb{D}_{\dot\gamma}^{[\mathbf{W}]}|\Psi_{\text{modal}}^{[\mathbf{W},\chi]}\rangle = 0$ (equivalently, $\mathbb{D}_{\dot\gamma}^{[\mathbf{W}]}c = 0$), then $\frac{d}{ds}N(s) = 0$, proving the first claim. The invocation of Proposition 4.223 guarantees that the *word-locked* readout frame is the one in which $A_\mu^{[\mathbf{W}]}$ is computed, so metric-compatibility ($A_\mu^{[\mathbf{W}]}\in\mathfrak{u}(4)$) applies at the endpoint.

Covariant Leibniz rule for expectations. For any fibre observable $\hat O(x)$, write its matrix in the wordwise frame as $O(x)$ and

$$\langle\hat O\rangle^{[\mathbf{W}]}(x) = \langle\Psi_{\text{modal}}^{[\mathbf{W},\chi]}(x)|\,\hat O(x)\,|\Psi_{\text{modal}}^{[\mathbf{W},\chi]}(x)\rangle = c^\dagger(x)\,O(x)\,c(x).$$

Differentiate:

$$\partial_\mu\langle\hat O\rangle^{[\mathbf{W}]} = (\partial_\mu c^\dagger)Oc + c^\dagger(\partial_\mu O)c + c^\dagger O(\partial_\mu c).$$

Insert $\partial_\mu c = \mathbb{D}_\mu^{[\mathbf{W}]} c - A_\mu^{[\mathbf{W}]} c$ and its adjoint, and group terms to obtain

$$\partial_\mu \langle \hat{O} \rangle^{[\mathbf{W}]} = \big(\mathbb{D}_\mu^{[\mathbf{W}]} c\big)^\dagger O c + c^\dagger O \big(\mathbb{D}_\mu^{[\mathbf{W}]} c\big) + c^\dagger \Big(\partial_\mu O - (A_\mu^{[\mathbf{W}]})^\dagger O - O A_\mu^{[\mathbf{W}]}\Big) c.$$

By the *word-locked frame condition at readout* (cf. Proposition 4.223), fibre endomorphisms are identified in that local frame and regarded as *parallel-fixed*; equivalently,

$$\mathbb{D}_\mu^{[\mathbf{W}]} O := \partial_\mu O - (A_\mu^{[\mathbf{W}]})^\dagger O - O A_\mu^{[\mathbf{W}]} = 0 \quad \text{(fixed pairing, cf. eq. (4.105))}.$$

With this, the last bracket vanishes and we get

$$\partial_\mu \langle \hat{O} \rangle^{[\mathbf{W}]} = \Big\langle (\mathbb{D}_\mu^{[\mathbf{W}]})^\dagger \hat{O}(x) \Big\rangle^{[\mathbf{W}]} + \Big\langle \hat{O}(x)\, \mathbb{D}_\mu^{[\mathbf{W}]} \Big\rangle^{[\mathbf{W}]},$$

where the right-hand side is shorthand for $\langle (\mathbb{D}_\mu^{[\mathbf{W}]} \Psi) |\, \hat{O}\, |\Psi\rangle + \langle \Psi |\, \hat{O}\, (\mathbb{D}_\mu^{[\mathbf{W}]} \Psi)\rangle$. This is exactly the covariant Leibniz rule in the wordwise setting.

The two statements together establish Lemma 4.279. □

Definition 4.280 (Wordwise fibre density and expectations)**.** Set

$$\rho_{\text{modal}}^{[\mathbf{W}]}(x) := |\Psi_{\text{modal}}^{[\mathbf{W},\chi]}(x)\rangle \langle \Psi_{\text{modal}}^{[\mathbf{W},\chi]}(x)|.$$

For any fibre observable $\hat{O}(x)$, define the expectation

$$\langle \hat{O} \rangle^{[\mathbf{W}]}(x) := \mathrm{Tr}\Big(\rho_{\text{modal}}^{[\mathbf{W}]}(x)\, \hat{O}(x)\Big),$$

which is invariant under endpoint gauge (Lemma 4.256).

Proposition 4.281 (Independence of p, s from twofold superpositions (wordwise))**.** *In any wordwise frame, the coefficients $c_p^{[\mathbf{W}]}(x)$ and $c_s^{[\mathbf{W}]}(x)$ weight independent fibre directions $|p\rangle_x^{[\mathbf{W},\chi]}$ and $|s\rangle_x^{[\mathbf{W},\chi]}$. They are not fixed by twofold superpositions inside* $\mathrm{span}\{|0\rangle_x^{[\mathbf{W},\chi]}, |1\rangle_x^{[\mathbf{W},\chi]}\}$*; e.g. $|+\rangle_x^{[\mathbf{W},\chi]} = \frac{1}{\sqrt{2}}\big(|0\rangle_x^{[\mathbf{W},\chi]} + |1\rangle_x^{[\mathbf{W},\chi]}\big)$ is a twofold* effect*, not the fourfold basis vector $|p\rangle_x^{[\mathbf{W},\chi]}$ (*wordwise version of *Remark 4.269).*

Corollary 4.282 (Mode weights: gauge invariance and parallel constancy (wordwise))**.** *Let $P_\alpha^{[\mathbf{W}]}(x) = |\alpha\rangle_x^{[\mathbf{W},\chi]} \langle \alpha|_x^{[\mathbf{W},\chi]}$. Define $w_\alpha^{[\mathbf{W}]}(x) := \langle \Psi_{\text{modal}}^{[\mathbf{W},\chi]}(x),\, P_\alpha^{[\mathbf{W}]}(x)\, \Psi_{\text{modal}}^{[\mathbf{W},\chi]}(x)\rangle$. Then $w_\alpha^{[\mathbf{W}]}(x)$ is invariant under endpoint gauge and, along any curve γ, is constant if $\mathbb{D}_{\dot\gamma}^{[\mathbf{W}]} |\Psi_{\text{modal}}^{[\mathbf{W},\chi]}\rangle = 0$ and $\mathbb{D}_{\dot\gamma}^{[\mathbf{W}]} P_\alpha^{[\mathbf{W}]} = 0$ (same propagation of projectors), cf. Corollary 4.272.*

Proof. Let $P_\alpha^{[\mathbf{W}]}(x) = |\alpha\rangle_x^{[\mathbf{W},\chi]} \langle \alpha|_x^{[\mathbf{W},\chi]}$ and

$$w_\alpha^{[\mathbf{W}]}(x) := \Big\langle \Psi_{\text{modal}}^{[\mathbf{W},\chi]}(x)\,,\, P_\alpha^{[\mathbf{W}]}(x)\, \Psi_{\text{modal}}^{[\mathbf{W},\chi]}(x) \Big\rangle.$$

Gauge invariance. Under an endpoint gauge $g(x) \in \mathrm{U}(4)$ on the modal fibre (wordwise gauge covariance; cf. Proposition 4.167(a) and Proposition 4.276),

$$|\Psi_{\text{modal}}^{[\mathbf{W},\chi]}(x)\rangle \mapsto g(x)\, |\Psi_{\text{modal}}^{[\mathbf{W},\chi]}(x)\rangle, \qquad P_\alpha^{[\mathbf{W}]}(x) \mapsto g(x)\, P_\alpha^{[\mathbf{W}]}(x)\, g(x)^\dagger.$$

Hence

$$w_\alpha^{[\mathbf{W}]}(x) \;\mapsto\; \Big\langle g\Psi,\, gP_\alpha g^\dagger\, g\Psi \Big\rangle = \Big\langle \Psi,\, P_\alpha\, \Psi \Big\rangle = w_\alpha^{[\mathbf{W}]}(x),$$

using $g^\dagger g = I$ and unitarity of $g(x)$. Thus $w_\alpha^{[\mathbf{W}]}(x)$ is gauge invariant pointwise.

Parallel constancy along a curve. Let $x = \gamma(s)$ and denote $\mathbb{D}_{\dot\gamma}^{[\mathbf{W}]}$ the wordwise covariant derivative along γ (with pullback connection A_γ). By the covariant Leibniz rule established in Lemma 4.279, for any fibre observable $\hat O(x)$,

$$\frac{d}{ds}\,\langle \hat O\rangle^{[\mathbf{W}]}(\gamma(s)) = \Big\langle (\mathbb{D}_{\dot\gamma}^{[\mathbf{W}]})^\dagger \hat O \Big\rangle^{[\mathbf{W}]} + \Big\langle \hat O\, \mathbb{D}_{\dot\gamma}^{[\mathbf{W}]} \Big\rangle^{[\mathbf{W}]}.$$

Apply this with $\hat O = P_\alpha^{[\mathbf{W}]}$ and use the hypotheses $\mathbb{D}_{\dot\gamma}^{[\mathbf{W}]} |\Psi_{\mathrm{modal}}^{[\mathbf{W},\chi]}\rangle = 0$ and $\mathbb{D}_{\dot\gamma}^{[\mathbf{W}]} P_\alpha^{[\mathbf{W}]} = 0$ ("same propagation of projectors"). Then both terms on the right vanish:

$$\Big\langle (\mathbb{D}_{\dot\gamma}^{[\mathbf{W}]})^\dagger P_\alpha^{[\mathbf{W}]} \Big\rangle^{[\mathbf{W}]} = 0, \qquad \Big\langle P_\alpha^{[\mathbf{W}]}\, \mathbb{D}_{\dot\gamma}^{[\mathbf{W}]} \Big\rangle^{[\mathbf{W}]} = 0,$$

so $\dfrac{d}{ds}\, w_\alpha^{[\mathbf{W}]}(\gamma(s)) = 0$.

This proves gauge invariance and parallel constancy of the wordwise mode weights, as claimed in Corollary 4.282. □

4.7 The Emergent Modal Space of a PQS

For each PQS $\mathfrak{B}_x$ at $x \in \mathcal{X}_q$, the intrinsic operator-theoretic structure

$$\hat{\mathbf{\Omega}}_x = {}_{\hat\Phi_x}\mathring{\mathbf{\Lambda}}_{x,\hat\Theta_x}$$

induces a local modal Hilbert space $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$. Its orthonormal basis $\{|0\rangle_x, |1\rangle_x, |p\rangle_x, |s\rangle_x\}$ realizes the four irreducible dialectical modes (Symmetry, Antisymmetry, Parasymmetry, Holosymmetry). These are not merely labeled vectors; they are relationally constructed eigenmodes of the internal duality operator $\mathring{\mathbf{\Lambda}}_x$ and encode the full Fourfold Dialectic Algebra. Crucially, the inner product on $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$—and thus amplitudes, orthogonality, and measurement—is *defined* by the Fourfold Inter-Modal Correlator in the Quantum Holo-State (coincident limit), rather than imposed *a priori* (cf. eq. (4.85)). The kinematics of quantum theory (Hilbert geometry and Born weights) thereby become *emergent consequences* of a deeper, dialectically governed substrate.

From tensor products to fiber bundles Where conventional theory assembles composites via tensor products of fixed Hilbert spaces, the IQR replaces this scaffolding with a geometrically enriched architecture: the *Hilbert bundle of dialectic modes*

$$\mathscr{H}_{\mathrm{mod}} = \bigsqcup_{x\in\mathcal{X}_q} \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$$

over the emergent configuration space $\mathcal{X}_q$ (cf. Definition 4.239). Each fiber is a locally constructed modal space with an inner product generated from correlators; fibers are compared

by gauge-covariant parallel transport. In this setting, coherence and entanglement are not externally imposed relations among static subsystems but *relational structures between fibers*, mediated by operator-valued channels and the connection that produces transport. Thus the modal Hilbert space of a PQS is the conceptual and physical successor to the qubit: a dynamically emergent, geometrically grounded structure that unifies internal logical modes, spatial locality, and informational coherence within one operator-theoretic framework—advancing beyond the abstract axioms toward a relationally enriched description of quantum reality.

Definition 4.283 (Wordwise Modal Hilbert Space of a PQS)**.** Fix $x \in \mathcal{X}_q$ with PQS $\mathfrak{B}_x$ (Definition 1.25) and choose a unitary device word $\mathbf{W} \in \langle \mathsf{A}_{\text{holo}} \rangle$ with control label χ. Let $\mathsf{W}_{\chi\leftarrow\chi}(x)$ be its RR element (Definition 4.148 and eq. (4.45)). The *wordwise modal Hilbert space* at x is the same fibre

$$\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)} \cong \mathbb{C}^4,$$

equipped with the emergent inner product induced by the Fourfold correlator in the Holo–state (Section 4.6 and eq. (4.85)). The *wordwise realization map*

$$\mathcal{R}_x^{[\mathbf{W},\chi]} : \{0,1,p,s\} \longrightarrow \mathbb{P}\Big(\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}\Big), \qquad \alpha \longmapsto \Big[\,|\alpha\rangle_x^{[\mathbf{W},\chi]}\,\Big],$$

is defined by the *word-rotated frame*

$$|\alpha\rangle_x^{[\mathbf{W},\chi]} := \mathsf{W}_{\chi\leftarrow\chi}(x)\,|\alpha\rangle_x, \qquad \alpha \in \{0,1,p,s\} \quad (\textit{Definition 4.257and Proposition 4.258}).$$

Wordwise orthonormal basis. Since $\mathsf{W}_{\chi\leftarrow\chi}(x)$ is unitary on the fibre, orthonormality is preserved:

$$\boxed{\Big\langle \alpha_x^{[\mathbf{W},\chi]} \;\Big|\; \beta_x^{[\mathbf{W},\chi]} \Big\rangle = \delta_{\alpha\beta}, \qquad \alpha,\beta \in \{0,1,p,s\} \quad (\text{cf. } \textit{eq.} \text{ (4.107})).} \tag{4.113}$$

Caution. $|p\rangle_x^{[\mathbf{W},\chi]}$ and $|s\rangle_x^{[\mathbf{W},\chi]}$ remain independent basis vectors. The twofold equal superposition $|+\rangle_x^{[\mathbf{W},\chi]} := \frac{1}{\sqrt{2}}\Big(|0\rangle_x^{[\mathbf{W},\chi]} + |1\rangle_x^{[\mathbf{W},\chi]}\Big)$ is a twofold *effect* and is not identified with $|p\rangle_x^{[\mathbf{W},\chi]}$ (Remark 4.269 and Proposition 4.281).

Wordwise modal decomposition. With $\mathcal{H}_{\alpha,x}^{[\mathbf{W}]} := \text{span}\{|\,\alpha\rangle_x^{[\mathbf{W},\chi]}\}$, one has

$$\boxed{\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)} = \mathcal{H}_{0,x}^{[\mathbf{W}]} \oplus \mathcal{H}_{1,x}^{[\mathbf{W}]} \oplus \mathcal{H}_{p,x}^{[\mathbf{W}]} \oplus \mathcal{H}_{s,x}^{[\mathbf{W}]} \quad (\text{cf. } \textit{eq.} \text{ (4.108})).} \tag{4.114}$$

Wordwise state representation. Every $|\psi\rangle_x \in \mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}$ admits the expansion

$$|\psi\rangle_x = c_0^{[\mathbf{W}]}(x)\,|0\rangle_x^{[\mathbf{W},\chi]} + c_1^{[\mathbf{W}]}(x)\,|1\rangle_x^{[\mathbf{W},\chi]} + c_p^{[\mathbf{W}]}(x)\,|p\rangle_x^{[\mathbf{W},\chi]} + c_s^{[\mathbf{W}]}(x)\,|s\rangle_x^{[\mathbf{W},\chi]},$$

with pointwise normalization

$$\boxed{\Big|c_0^{[\mathbf{W}]}(x)\Big|^2 + \Big|c_1^{[\mathbf{W}]}(x)\Big|^2 + \Big|c_p^{[\mathbf{W}]}(x)\Big|^2 + \Big|c_s^{[\mathbf{W}]}(x)\Big|^2 = 1.} \tag{4.115}$$

Proposition 4.284 (Word/gauge covariance and invariance of the construction)**.** *For any endpoint gauge $g(x) \in \mathrm{U}(4)$ and any unitary words $\mathbf{W}, \mathbf{W}'$, the corresponding frames satisfy*

$$|\alpha\rangle_x^{[\mathbf{W}',\chi]} \;=\; U_{[\mathbf{W}',\mathbf{W}]}(x)\,|\alpha\rangle_x^{[\mathbf{W},\chi]}, \qquad U_{[\mathbf{W}',\mathbf{W}]}(x) := \mathsf{W}'_{\chi\leftarrow\chi}(x)\,\mathsf{W}_{\chi\leftarrow\chi}(x)^\dagger \in \mathrm{U}(4),$$

and $|\alpha\rangle_x^{[\mathbf{W},\chi]} \mapsto g(x)\,|\alpha\rangle_x^{[\mathbf{W},\chi]}$. Thus $\mathcal{R}_x^{[\mathbf{W},\chi]}$ changes by unitary conjugation, while $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$, the inner product (eq. (4.85)), the orthonormality (4.113), and the decomposition (4.114) are invariant.

Remark 4.285 (Connection/transport compatibility (wordwise at a point))**.** Along any path $\gamma : x' \to x$, frames and states obey the same unitary transport:

$$|\alpha\rangle_x^{[\mathbf{W},\chi]} = \mathcal{U}(x \leftarrow x';\gamma)\,|\alpha\rangle_{x'}^{[\mathbf{W},\chi]}, \qquad |\psi\rangle_x = \mathcal{U}(x \leftarrow x';\gamma)\,|\psi\rangle_{x'},$$

with $\mathcal{U} = \mathcal{P}\exp\!\big(-\int_\gamma \mathcal{A}\big)$ (Definition 4.248 and Proposition 4.249). Hence norms, orthonormality, and the modal split are preserved under parallel transport (the wordwise counterpart of Remark 4.267 and Lemma 4.270).

Interpretation (endogenous). The fibre $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ remains the relationally defined arena for the four modal facets, but is now *presented* via the word-rotated frame induced by a concrete device word. Different words give unitarily equivalent presentations (Prop. 4.284); measurements and inner products are unchanged, while the endogenous viewpoint keeps a tight link to operational realizations (control, lifted histories, and word-evaluated channels).

Remark 4.286 (Pointwise exogenous $\leftrightarrow$ endogenous dictionary)**.** At a fixed locus $x \in \mathcal{X}_q$, the exogenous (frame-based) and endogenous (word-rotated) presentations are related by the RR element of a unitary device word $\mathsf{W}_{\chi\leftarrow\chi}(x)$ (Definitions 4.148 and 4.257, eq. (4.45), and Proposition 4.258):

$$\begin{aligned}
&\text{(realization map)} && \mathcal{R}_x(\alpha) = [\,|\alpha\rangle_x\,] \;\longleftrightarrow\; \mathcal{R}_x^{[\mathbf{W},\chi]}(\alpha) = \Big[\,|\alpha\rangle_x^{[\mathbf{W},\chi]}\Big] \\
&&& \qquad\qquad = \Big[\,\mathsf{W}_{\chi\leftarrow\chi}(x)\,|\alpha\rangle_x\,\Big]; \\
&\text{(modal frame)} && \mathcal{B}_x = \{|0\rangle_x, |1\rangle_x, |p\rangle_x, |s\rangle_x\} \;\longleftrightarrow\; \mathcal{B}_x^{[\mathbf{W},\chi]} = \mathsf{W}_{\chi\leftarrow\chi}(x)\,\mathcal{B}_x; \\
&\text{(twofold superposition)} && |+\rangle_x = \tfrac{1}{\sqrt{2}}\big(|0\rangle_x + |1\rangle_x\big) \;\longleftrightarrow\; |+\rangle_x^{[\mathbf{W},\chi]} = \mathsf{W}_{\chi\leftarrow\chi}(x)\,|+\rangle_x.
\end{aligned}$$

The unitary conjugation preserves orthonormality (cf. eqs. (4.107) and (4.113)) and the modal decomposition (cf. eqs. (4.108) and (4.114)). *Caution (unchanged under conjugation):* the twofold effect $|+\rangle_x^{[\mathbf{W},\chi]}$ is not the parasymmetric basis vector $|p\rangle_x^{[\mathbf{W},\chi]}$; p, s remain independent directions (cf. Remark 4.269 and Proposition 4.281). By Proposition 4.284, different words (and endpoint gauges) change only the presentation via a unitary, leaving all fibrewise observables and pairings invariant.

4.7.1 Emergent Hilbert Space: GNS Realization and Modal Coupling

Definition 4.287 (Emergent Hilbert space via GNS and its modal role)**.** Let $\mathcal{A}$ be the unital C^*-algebra of emergent observables and let $\omega^{\overset{\circ}{\Psi}} : \mathcal{A} \to \mathbb{C}$ be the Holo-State. By the Gelfand–Naimark–Segal construction (Definition 1.34) there exists a cyclic $*$-representation

$$\left(\pi_{\overset{\circ}{\Psi}},\ \mathcal{H}_{\overset{\circ}{\Psi}},\ |\Omega_{\omega^{\overset{\circ}{\Psi}}}\rangle\right)$$

such that $\omega^{\overset{\circ}{\Psi}}(A) = \langle\Omega_{\omega^{\overset{\circ}{\Psi}}}|\ \pi_{\overset{\circ}{\Psi}}(A)\ |\Omega_{\omega^{\overset{\circ}{\Psi}}}\rangle$ for all $A \in \mathcal{A}$ and $\overline{\mathrm{span}}\{\pi_{\overset{\circ}{\Psi}}(A)|\Omega_{\omega^{\overset{\circ}{\Psi}}}\rangle : A \in \mathcal{A}\} = \mathcal{H}_{\overset{\circ}{\Psi}}$.

Definition. The **emergent Hilbert space** is

$$\mathcal{H}_{\mathrm{em}} \ :=\ \mathcal{H}_{\overset{\circ}{\Psi}},$$

with representation $\pi := \pi_{\overset{\circ}{\Psi}}$ and cyclic vector $|\Omega\rangle := |\Omega_{\omega^{\overset{\circ}{\Psi}}}\rangle$. For any $A \in \mathcal{A}$ we set the represented ("hatted") operator $\hat{A} := \pi(A) \in \mathcal{B}(\mathcal{H}_{\mathrm{em}})$, so that $\langle\overset{\circ}{\Psi}|\cdot|\overset{\circ}{\Psi}\rangle = \langle\Omega|\cdot|\Omega\rangle = \omega^{\overset{\circ}{\Psi}}(\cdot)$ on $\mathcal{A}$.

Compatibility with the modal sector.

(i) *Carrier space for channels and transport.* All Fourfold correlator components and transporters live on $\mathcal{H}_{\mathrm{em}}$:

$$\hat{G}^{\alpha\beta}_{\Lambda}(x, x') \in \mathcal{B}(\mathcal{H}_{\mathrm{em}}), \qquad \mathcal{U}(x \leftarrow x'; \gamma) \in \mathrm{U}(\mathcal{H}_{\mathrm{em}}),$$

with their gauge covariance and Hermitian symmetry as stated in Sections 4.1.2 and 4.1.4.

(ii) *Induced fiber metrics.* For each PQS locus $x \in \mathcal{X}_q$ the inner product on the modal fiber $\mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}}$ is *generated* from $\mathcal{H}_{\mathrm{em}}$ by the Holo-State expectation of the Fourfold correlator in the coincident limit (cf. eq. (4.85)):

$$\langle\psi_1|\psi_2\rangle_x := \lim_{x' \to x} \langle\Omega|\,\hat{G}_{\Lambda}(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2]\,|\Omega\rangle,$$

yielding the orthonormal modal basis and positivity discussed in Section 4.6.

(iii) *Hilbert bundle structure.* The collection of fibers $\{\mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}}\}_{x \in \mathcal{X}_q}$ organizes into the Hilbert bundle of dialectic modes $\mathscr{H}_{\mathrm{mod}} = \bigsqcup_{x \in \mathcal{X}_q} \mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}}$ (Definition 4.239); the connection whose path-ordered exponential is $\mathcal{U}$ implements gauge-covariant comparisons between fibers (Section 4.1.4).

Remark 4.288 (Uniqueness and Role)**.** The GNS representation $(\pi, \mathcal{H}_{\mathrm{em}}, |\Omega\rangle)$ is unique up to unitary equivalence; thus $\mathcal{H}_{\mathrm{em}}$ is determined by the pair $(\mathcal{A}, \omega^{\overset{\circ}{\Psi}})$. It serves as the universal carrier for represented observables, channel kernels, and parallel transporters, while *fiberwise* Hilbert geometries on $\mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}}$ are *induced* from $\mathcal{H}_{\mathrm{em}}$ via the Fourfold correlator expectations. This realizes the slogan: global state $\Rightarrow$ GNS space $\mathcal{H}_{\mathrm{em}} \Rightarrow$ local fiber metrics.

Proposition 4.289 (Dimensional Profile and Ranks)**.** *Let $\mathcal{A}$ be the observable C^*-algebra and let $(\pi_{\overset{\circ}{\Psi}}, \mathcal{H}_{\overset{\circ}{\Psi}}, |\Omega_{\omega^{\overset{\circ}{\Psi}}}\rangle)$ be the GNS triple of the Holo-State $\omega^{\overset{\circ}{\Psi}}$. Write $\mathcal{H}_{\mathrm{em}} := \mathcal{H}_{\overset{\circ}{\Psi}}$. Then:*

(i) Emergent Hilbert space

$\mathcal{H}_{\mathrm{em}}$ is generally infinite-dimensional (typically separable). Its cardinal dimension depends on $(\mathcal{A}, \omega^{\overset{\circ}{\Psi}})$.

(ii) Modal fibers

For each locus $x \in \mathcal{X}_q$, the modal fiber is a rank-4 complex Hilbert space

$$\dim_{\mathbb{C}} \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)} = 4,$$

with a direct-sum decomposition into four one-dimensional modal lines

$$\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)} = \mathcal{H}_{0,x} \oplus \mathcal{H}_{1,x} \oplus \mathcal{H}_{p,x} \oplus \mathcal{H}_{s,x}, \qquad \dim_{\mathbb{C}} \mathcal{H}_{\alpha,x} = 1.$$

(iii) Twofold vs. higher modes

The subspaces $\mathcal{H}_{(0,1),x} := \mathcal{H}_{0,x} \oplus \mathcal{H}_{1,x}$ and $\mathcal{H}_{(p,s),x} := \mathcal{H}_{p,x} \oplus \mathcal{H}_{s,x}$ have

$$\dim_{\mathbb{C}} \mathcal{H}_{(0,1),x} = \dim_{\mathbb{C}} \mathcal{H}_{(p,s),x} = 2.$$

(iv) Hilbert bundle of dialectic modes

The modal bundle $\pi : \mathcal{H}^{\mathrm{modal}} \to \mathcal{X}_q$ is a rank-4 complex Hermitian vector bundle, locally trivial as $\pi^{-1}(U) \cong U \times \mathbb{C}^4$. Its space of sections $\Gamma(\mathcal{X}_q, \mathcal{H}^{\mathrm{modal}})$ is typically infinite-dimensional.

(v) Correlator size in modal indices

The Fourfold inter-modal correlator has the block form

$$\hat{\mathbf{G}}_\Lambda(x, x') = [\hat{G}_\Lambda^{\alpha\beta}(x, x')]_{\alpha,\beta \in \{0,1,p,s\}}$$

i.e. a 4×4 matrix in modal indices with entries $\hat{G}_\Lambda^{\alpha\beta}(x, x') \in \mathcal{B}(\mathcal{H}_{\mathrm{em}})$. With the field–space refinement, one obtains a 2×2 matrix in field labels whose entries are themselves 4×4 in modal indices.

(vi) Operator spaces

$\mathcal{B}(\mathcal{H}_{\mathrm{em}})$ is infinite-dimensional (and non-separable in the operator-norm topology even when $\mathcal{H}_{\mathrm{em}}$ is separable).

$$\boxed{\begin{aligned} \dim_{\mathbb{C}} \mathcal{H}_{\mathrm{em}} &= \infty \quad \text{(typically separable)}, \\ \dim_{\mathbb{C}} \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)} &= 4, \\ \dim_{\mathbb{C}} \mathcal{H}_{\alpha,x} &= 1 \quad (\alpha \in \{0,1,p,s\}). \end{aligned}} \tag{4.116}$$

Remark 4.290 (Operator-Space Size and Why It Matters)**.** The symbol $\mathcal{B}(\mathcal{H}_{\mathrm{em}})$ denotes the C^*-algebra of all bounded operators on the (typically separable, infinite-dimensional) emergent Hilbert space $\mathcal{H}_{\mathrm{em}}$. Two distinct facts are being emphasized:

(a) Infinite-dimensionality. Even when $\mathcal{H}_{\mathrm{em}}$ is separable, $\mathcal{B}(\mathcal{H}_{\mathrm{em}})$ is *vast*: it contains all bounded linear maps, including entangling channels, projectors onto arbitrary closed subspaces, conditional expectations, etc.

(b) Non-separability in operator norm. Although $\mathcal{H}_{\text{em}}$ may be separable (i.e. it has a countable dense subset), the metric space $\big(\mathcal{B}(\mathcal{H}_{\text{em}}), \|\cdot\|\big)$ is *not* separable. A standard proof sketch: fix an orthonormal basis $\{e_n\}_{n\in\mathbb{N}}$ of $\mathcal{H}_{\text{em}}$ and, for each subset $S \subseteq \mathbb{N}$, let $P_S \in \mathcal{B}(\mathcal{H}_{\text{em}})$ be the diagonal projection onto $\text{span}\{e_n : n \in S\}$. Then for $S \neq T$ the operator $P_S - P_T$ is diagonal with entries in $\{-1, 0, 1\}$ and hence $\|P_S - P_T\| = 1$. Thus $\{P_S : S \subseteq \mathbb{N}\}$ is an uncountable family that is 1-separated in operator norm, so no countable set can be dense. Therefore $\mathcal{B}(\mathcal{H}_{\text{em}})$ is non-separable in $\|\cdot\|$.

Why this matters for the Holo-State. The Fourfold correlator takes values in $\mathcal{B}(\mathcal{H}_{\text{em}})$, so the "space of allowed channels" is large enough to encode arbitrarily fine, genuinely global couplings. Later, when we show that the Quantum Holo-State is generally *not factorizable* across putative subsystem splits, we will not be relying on a small algebraic deficiency: even with a separable $\mathcal{H}_{\text{em}}$, the ambient observable algebra is big enough to support non-product (entangled, holistically constrained) states and dynamics. In short, separability of the *state space* does not imply separability (or any smallness) of the *operator space*; the latter's non-separable richness is precisely what allows nontrivial global structure to persist.

The framework inverts the traditional hierarchy "Hilbert space first, physics second." In the algebraic spirit, it is **algebra first, state first—then Hilbert space** [230, 241, 4]. The Hilbert space is not a pre-given arena; it is an outcome of deeper structural data. Two interconnected layers make this precise.

Global (GNS) layer. From the observable C^*-algebra $\mathcal{A}$ and the Quantum Holo-State $\omega^{\mathring{\mathbf{\Psi}}}$, the GNS construction produces the state-relative *emergent Hilbert space*

$$\mathcal{H}_{\text{em}} := \mathcal{H}_{\mathring{\mathbf{\Psi}}},$$

typically infinite-dimensional (often separable), which carries the representation in which all "hatted" operators act—fields, correlator entries, and parallel transporters [241, 230].

Local (fibered) layer. Alongside this global representation, the local degrees of freedom organize as a *rank-4 Hermitian vector bundle* over the emergent configuration space $\mathcal{X}_q$ [375, 225]. At each locus x (a Primordial Quantum Singularity) sits a four-dimensional modal fiber $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)} \cong \mathbb{C}^4$. In a radical departure from textbook practice, the fiber metric is *not assumed*; it is *generated* by taking the Holo-State expectation of the *Fourfold Inter-Modal Correlator* in the coincident limit, thereby defining orthogonality and amplitudes on each fiber [230, 241]. A gauge connection furnishes *parallel transport* between fibers; its curvature produces *holonomy*, which is operationally visible in the holosymmetric channels [375, 225].

This architecture differs sharply from the standard picture. There is *no fixed background arena*: both the global representation space and the local fiber metrics *emerge* from $(\mathcal{A}, \omega^{\mathring{\mathbf{\Psi}}})$ and from correlators [230, 241, 11]. Subsystems and entanglement are not read off a prior tensor factorization; they are inferred directly from *operator-valued kernels plus transport* [230, 11]. Moreover, each fiber carries the $\mathbb{Z}_2 \times \mathbb{Z}_2$–graded *Fourfold Dialectic Algebra*, which supplies selection rules for mixing among Symmetry, Antisymmetry, Parasymmetry, and Holosymmetry [230, 241].

Bottom line. The result is a *relational, state-relative, fibered Hilbert geometry*: the "space of states" is not the stage for the theory—*it is one of the theory's outcomes*, carved by correlators, gauge transport, and the global Holo-State [230, 241, 375, 225].

4.7.2 Holonomy of the Modal Connection over PQS

The framework's Hilbert space is not a single, monolithic arena but a *fibered geometry*: local modal fibers are glued by a physical connection, and that gluing has observable consequences. On the Hilbert bundle of dialectic modes $(\mathscr{H}_{\mathrm{mod}}, \pi, \mathcal{X}_q)$ (cf. Definition 4.239), a skew-Hermitian connection one-form $\mathcal{A}$ generates the *parallel transporter* along a path $\gamma \subset \mathcal{X}_q$,

$$\mathcal{U}(x \leftarrow x'; \gamma) = \mathcal{P} \exp\Big(-\int_\gamma \mathcal{A} \Big), \qquad \mathcal{F} := \mathrm{d}\mathcal{A} + \mathcal{A} \wedge \mathcal{A}. \tag{4.117}$$

(cf. Section 4.1.4). For a closed loop ℓ based at x,

$$\mathrm{Hol}_\ell(x) := \mathcal{P}\, \exp\Big(-\oint_\ell \mathcal{A} \Big) \in \mathrm{U}(4)$$

is the *holonomy*. Flat regions ($\mathcal{F} \equiv 0$) on simply connected patches yield trivial holonomy; nonzero curvature or nontrivial topology produces a net, path-dependent twist.

Definition 4.291 (Holosymmetric loop operator)**.** Given a based loop ℓ at x with holonomy $\mathrm{Hol}_\ell \in \mathrm{U}(4)$ on the modal bundle, define

$$\hat{H}^{(s)}_\ell(x) := \hat{\mathcal{F}}^\dagger_s(x)\, \mathrm{Hol}_\ell\, \hat{\mathcal{F}}_s(x) \quad \in \mathcal{B}(\mathcal{H}_{\mathrm{em}}). \tag{4.118}$$

Its expectation in the Holo-State, $\omega^{\mathring{\mathbf{\Psi}}}\big(\hat{H}^{(s)}_\ell(x)\big)$, is the holonomy signal.

Loop readout in the holosymmetric channel. To turn holonomy into an operational signal, use the Fourfold channel structure (cf. Section 4.1.2). Let

$$\mathcal{U}_\ell(x) := \mathcal{U}(x \leftarrow x; \ell)$$

be the loop transporter along the closed path ℓ. The *holosymmetric loop operator* at x is

$$\hat{H}^{(s)}_\ell(x) \;:=\; \hat{\mathcal{F}}^\dagger_s(x)\, \mathcal{U}_\ell(x)\, \hat{\mathcal{F}}_s(x) \;=\; \hat{G}^{ss}_\Lambda(x, x; \ell), \tag{4.119}$$

i.e. the s–self channel of the Fourfold correlator with transport taken around ℓ (compare eq. (4.22)). Its expectation in the Holo-State gives a *holonomy signal*

$$h^{(s)}_\ell(x) \;:=\; \omega^{\mathring{\mathbf{\Psi}}}\big(\hat{H}^{(s)}_\ell(x)\big) \;=\; \Big\langle \mathring{\mathbf{\Psi}} \Big|\, \hat{H}^{(s)}_\ell(x) \,\Big|\, \mathring{\mathbf{\Psi}} \Big\rangle, \tag{4.120}$$

which is gauge-invariant (by the covariance of $\hat{\mathcal{F}}_s$ and $\mathcal{U}$; cf. Sections 4.1.2 and 4.1.4). In flat, contractible regions $h^{(s)}_\ell(x)$ is trivial; nontrivial values witness curvature or topology, the latter being directly analogous to Aharonov–Bohm–type behavior [378, 375].

Why the s–channel? Central coherence and cleanliness. The holosymmetric self–channel is (graded-)central in the Fourfold algebra (Postulate 4.4), so in flat regions it commutes with other modal transitions. Curvature introduces controlled deviations (eq. (4.75)). Consequently, $\hat{H}_\ell^{(s)}$ functions as a *stable coherence reference*: its phase/rotation around a loop cleanly reflects global structure, disentangled from local twofold interference.

Holonomy vs. singularity. Holonomy is not a "hole'' in space; it is a unitary memory of transport defined by the smooth connection $\mathcal{A}$. A classical singularity is a breakdown of geometry; by contrast, a Primordial Quantum Singularity (PQS) $\mathfrak{B}_x$ is an *algebraic seed* carrying the Fourfold Dialectic Algebra at x. The modal connection $\mathcal{A}$ lives on $\mathcal{X}_q$ *between* such loci and encodes how their local modal frames are glued; holonomy records the global twist of that gluing.

Metric compatibility and preservation under transport. With the fiber inner products generated from the correlator (cf. Section 4.6), the covariant derivative $\mathfrak{D}_\mu = \partial_\mu + \mathcal{A}_\mu$ satisfies the Leibniz rule

$$\partial_\mu \langle \Phi_{\text{modal}}, \Psi_{\text{modal}} \rangle_x = \Big\langle \mathfrak{D}_\mu \Phi_{\text{modal}}, \Psi_{\text{modal}} \Big\rangle_x + \Big\langle \Phi_{\text{modal}}, \mathfrak{D}_\mu \Psi_{\text{modal}} \Big\rangle_x . \tag{4.121}$$

so norms and orthonormal frames are preserved along parallel transport (cf. Lemma 4.270).

Conceptual upshot. Holonomy upgrades geometry into experiment: the connection that glues fibers *writes global structure into measurable phases*. The holosymmetric loop operator $\hat{H}_\ell^{(s)}$ is the calibrated dial that reads it. Flat transport erases the dial's deflection; curvature or nontrivial topology registers as a robust, implementation-invariant signal in the s–channel.

4.7.3 Wordwise Holonomy of the Modal Connection over PQS

We view holonomy operationally via word-evaluated channels and word-rotated frames (Definitions 4.148, 4.255 and 4.257 and eq. (4.45)). The base connection $\mathcal{A}$ and transporter $\mathcal{U} = \mathcal{P}\exp(-\int \mathcal{A})$ remain exactly as in the exogenous formulation (Definitions 4.56 and 4.248 and Proposition 4.249).

Definition 4.292 (Wordwise loop transporter and holonomy)**.** For a closed loop ℓ based at x, the loop transporter in the modal bundle is

$$\mathcal{U}_\ell(x) := \mathcal{U}(x \leftarrow x; \ell) = \mathcal{P}\,\exp\Big(-\oint_\ell \mathcal{A} \Big) \in \mathrm{U}(4) \quad (\textit{Definition 4.56}).$$

Fix a unitary device word $\mathbf{W}$ and control χ. The *wordwise holosymmetric loop operator* is

$$\boxed{\hat{H}_\ell^{(s)[\mathbf{W}]}(x) \;:=\; \Big(\hat{\mathcal{F}}_s^{[\mathbf{W};\,\chi\leftarrow\chi]}(x)\Big)^\dagger \mathcal{U}_\ell(x)\, \hat{\mathcal{F}}_s^{[\mathbf{W};\,\chi\leftarrow\chi]}(x) \;=\; \hat{G}_\Lambda^{ss}\Big(x, x; \mathbf{W}, \chi\leftarrow\chi; \ell\Big),} \tag{4.122}$$

i.e. the s–self word-evaluated channel around the loop (Definition 4.166 and eq. (4.66)).

Definition 4.293 (Wordwise holonomy signal)**.** With the Holo-State functional $\omega^{\mathring{\Psi}}(\cdot)$, define the *wordwise holonomy signal*

$$\boxed{h_\ell^{(s)[\mathbf{W}]}(x) \;:=\; \omega^{\mathring{\Psi}}\Big(\hat{H}_\ell^{(s)[\mathbf{W}]}(x)\Big) \;=\; \Big\langle \mathring{\Psi}\Big|\, \hat{H}_\ell^{(s)[\mathbf{W}]}(x)\,\Big|\mathring{\Psi}\Big\rangle.} \tag{4.123}$$

Proposition 4.294 (Gauge covariance and invariances)**.** *Under an endpoint gauge $g : \mathcal{X}_q \to \mathrm{U}(4)$,*

$$\mathcal{U}_\ell(x) \mapsto g(x)\,\mathcal{U}_\ell(x)\,g(x)^\dagger, \qquad \hat{\mathcal{F}}_s^{[\mathbf{W};\chi\leftarrow\chi]}(x) \mapsto g(x)\,\hat{\mathcal{F}}_s^{[\mathbf{W};\chi\leftarrow\chi]}(x),$$

and hence

$$\hat{H}_\ell^{(s)[\mathbf{W}]}(x) \;\mapsto\; g(x)\,\hat{H}_\ell^{(s)[\mathbf{W}]}(x)\,g(x)^\dagger,$$

so the Born scalar $h_\ell^{(s)[\mathbf{W}]}(x)$ in (4.123) is gauge-invariant (Proposition 4.54 and Corollary 4.64). Moreover, replacing $\mathbf{W}$ by another unitary word $\mathbf{W}'$ conjugates $\hat{H}_\ell^{(s)[\mathbf{W}]}(x)$ by a unitary $U_{[\mathbf{W}',\mathbf{W}]}(x)$ (Proposition 4.284), hence preserves its spectrum and any trace-based invariants.

Proof. Fix $x \in \mathcal{X}_q$ and a loop/segment ℓ based at x. Write the wordwise s–selector as a fibre ket $\hat{\mathcal{F}}_s^{[\mathbf{W};\chi\leftarrow\chi]}(x)$ and its rank-one projector $P_s^{[\mathbf{W}]}(x) := \hat{\mathcal{F}}_s^{[\mathbf{W};\chi\leftarrow\chi]}(x)\,\hat{\mathcal{F}}_s^{[\mathbf{W};\chi\leftarrow\chi]}(x)^\dagger$. Let the (endpoint) holonomy/transport be $\mathcal{U}_\ell(x)$. Define the wordwise holonomy operator

$$\hat{H}_\ell^{(s)[\mathbf{W}]}(x) \;:=\; \mathcal{U}_\ell(x)\,P_s^{[\mathbf{W}]}(x). \tag{4.124}$$

Gauge covariance. Under an endpoint gauge $g : \mathcal{X}_q \to \mathrm{U}(4)$ one has

$$\begin{aligned} \mathcal{U}_\ell(x) &\mapsto g(x)\,\mathcal{U}_\ell(x)\,g(x)^\dagger \quad \text{by } \textit{Proposition } 4.54, \\ \hat{\mathcal{F}}_s^{[\mathbf{W};\chi\leftarrow\chi]}(x) &\mapsto g(x)\,\hat{\mathcal{F}}_s^{[\mathbf{W};\chi\leftarrow\chi]}(x) \text{ by } \textit{Corollary } 4.64. \end{aligned} \tag{4.125}$$

Hence $P_s^{[\mathbf{W}]}(x) \mapsto g(x)\,P_s^{[\mathbf{W}]}(x)\,g(x)^\dagger$, and from (4.124) it follows that

$$\hat{H}_\ell^{(s)[\mathbf{W}]}(x) \;\mapsto\; g(x)\,\hat{H}_\ell^{(s)[\mathbf{W}]}(x)\,g(x)^\dagger.$$

Let the Born scalar $h_\ell^{(s)[\mathbf{W}]}(x)$ be the (normalized) expectation appearing in eq. (4.123). Since the state transforms as $|\Psi(x)\rangle \mapsto g(x)|\Psi(x)\rangle$ (endpoint action) and $g(x)$ is unitary,

$$h_\ell^{(s)[\mathbf{W}]}(x) = \langle\Psi(x)|\,\hat{H}_\ell^{(s)[\mathbf{W}]}(x)\,|\Psi(x)\rangle \;\longmapsto\; \langle\Psi|\,g^\dagger g\,\hat{H}_\ell^{(s)[\mathbf{W}]}(x)\,g^\dagger g\,|\Psi\rangle = h_\ell^{(s)[\mathbf{W}]}(x),$$

so the scalar is gauge-invariant.

Invariance under unitary word changes. Let $\mathbf{W}'$ be another *unitary* word. By wordwise modal invariance (Proposition 4.284), there exists a unitary $U_{[\mathbf{W}',\mathbf{W}]}(x) \in \mathrm{U}(4)$ such that, at the endpoint x,

$$\hat{\mathcal{F}}_s^{[\mathbf{W}';\chi\leftarrow\chi]}(x) = U_{[\mathbf{W}',\mathbf{W}]}(x)\,\hat{\mathcal{F}}_s^{[\mathbf{W};\chi\leftarrow\chi]}(x), \qquad \mathcal{U}_\ell^{[\mathbf{W}']}(x) = U_{[\mathbf{W}',\mathbf{W}]}(x)\,\mathcal{U}_\ell^{[\mathbf{W}]}(x)\,U_{[\mathbf{W}',\mathbf{W}]}(x)^\dagger,$$

(the latter being the same geometric transport written in the rotated word-locked frame). Consequently,

$$P_s^{[\mathbf{W}']}(x) = \hat{\mathcal{F}}_s^{[\mathbf{W}']}\hat{\mathcal{F}}_s^{[\mathbf{W}']\dagger} = U_{[\mathbf{W}',\mathbf{W}]}\, P_s^{[\mathbf{W}]}\, U_{[\mathbf{W}',\mathbf{W}]}^{\dagger},$$

and, using (4.124),

$$\begin{aligned}\hat{H}_\ell^{(s)[\mathbf{W}']}(x) &= \mathcal{U}_\ell^{[\mathbf{W}']}(x) P_s^{[\mathbf{W}']}(x) \\ &= U_{[\mathbf{W}',\mathbf{W}]}(x)\left(\mathcal{U}_\ell^{[\mathbf{W}]}(x) P_s^{[\mathbf{W}]}(x)\right) U_{[\mathbf{W}',\mathbf{W}]}(x)^\dagger \\ &= U_{[\mathbf{W}',\mathbf{W}]}(x)\, \hat{H}_\ell^{(s)[\mathbf{W}]}(x)\, U_{[\mathbf{W}',\mathbf{W}]}(x)^\dagger. \end{aligned} \tag{4.126}$$

Thus $\hat{H}_\ell^{(s)[\mathbf{W}']}(x)$ is unitarily similar to $\hat{H}_\ell^{(s)[\mathbf{W}]}(x)$, so they have the same spectrum and the same values of any trace-based invariant (e.g. $\mathrm{tr}[\hat{H}^k]$, characteristic polynomial coefficients), as claimed.

This proves Proposition 4.294. □

Proposition 4.295 (Flatness ⇒ trivial wordwise holonomy (on simply connected patches))**.** *If $\mathcal{F} = 0$ on a simply connected neighbourhood of ℓ, then $\mathcal{U}_\ell(x) = \mathbf{1}_4$ and*

$$\hat{H}_\ell^{(s)[\mathbf{W}]}(x) = \left(\hat{\mathcal{F}}_s^{[\mathbf{W};\chi\leftarrow\chi]}(x)\right)^\dagger \hat{\mathcal{F}}_s^{[\mathbf{W};\chi\leftarrow\chi]}(x), \tag{4.127}$$

so $h_\ell^{(s)[\mathbf{W}]}(x)$ reduces to the baseline s–channel self–correlator at the point (Propositions 4.224 and 4.264).

Proof. Let $U \subset \mathcal{X}_q$ be a simply connected neighbourhood containing the image of the loop ℓ based at x, and let $A^{[\mathbf{W}]}$ denote the wordwise connection one–form on the modal fibre with curvature $\mathcal{F} := \mathrm{d}A^{[\mathbf{W}]} + A^{[\mathbf{W}]} \wedge A^{[\mathbf{W}]}$. Assume $\mathcal{F} = 0$ on U.

Step 1: Flat ⇒ path–independent transport (and trivial loop holonomy). By the wordwise curvature/flatness statements (Propositions 4.224 and 4.264), flatness on a simply connected domain implies that parallel transport depends only on endpoints, i.e. for any two C^1 paths $\gamma_1, \gamma_2 : x' \to x$ in U, $\mathcal{U}(x \leftarrow x'; \gamma_1) = \mathcal{U}(x \leftarrow x'; \gamma_2)$. Equivalently, the holonomy around any contractible loop in U is the identity:

$$\mathcal{U}_\ell(x) = \mathbf{1}_4. \tag{4.128}$$

One way to see (4.128) directly is via the gauge–trivialization guaranteed by flatness on U: there exists $g : U \to \mathrm{U}(4)$ such that the transformed connection vanishes, $A^g = 0$ (local pure gauge). In that gauge the parallel transport along *any* path is the identity map, hence $\mathcal{U}_\ell^g(x) = \mathbf{1}_4$. By endpoint gauge covariance (Proposition 4.54), $\mathcal{U}_\ell^g(x) = g(x)\,\mathcal{U}_\ell(x)\,g(x)^\dagger$, and since $g(x)$ is unitary, this implies $\mathcal{U}_\ell(x) = \mathbf{1}_4$.

Step 2: Wordwise loop operator collapses to the baseline s–channel correlator. Recall the wordwise s–channel holonomy operator along ℓ (in the word–locked readout frame at x):

$$\hat{H}_\ell^{(s)[\mathbf{W}]}(x) = \left(\hat{\mathcal{F}}_s^{[\mathbf{W};\chi\leftarrow\chi]}(x)\right)^\dagger \mathcal{U}_\ell(x)\, \hat{\mathcal{F}}_s^{[\mathbf{W};\chi\leftarrow\chi]}(x) \quad \text{(cf. eq. (4.122))}. \tag{4.129}$$

Insert (4.128) into (4.129) to obtain

$$\hat{H}_{\ell}^{(s)[\mathbf{W}]}(x) = \Big(\hat{\mathcal{F}}_{s}^{[\mathbf{W};\chi\leftarrow\chi]}(x)\Big)^{\dagger}\hat{\mathcal{F}}_{s}^{[\mathbf{W};\chi\leftarrow\chi]}(x),$$

which is exactly eq. (4.127). In particular, the associated Born scalar $h_{\ell}^{(s)[\mathbf{W}]}(x)$ (cf. eq. (4.123)) reduces to the baseline s–channel self–correlator at the point x.

This proves Proposition 4.295. □

Remark 4.296 (Operational readout via wordwise channels)**.** The equality (4.122) shows that holonomy is directly encoded in an s–self word-evaluated channel; interferometric protocols that isolate the s–channel (e.g. via Hadamard sum/difference and centrality of ss under the words-only meta-claim, Proposition 4.176 and Corollary 4.225) provide clean access to $h_{\ell}^{(s)[\mathbf{W}]}(x)$. In flat, contractible regions the signal is trivial (Proposition 4.295); nontrivial values witness curvature or topology, in direct analogy with Aharonov–Bohm–type phases, but here recorded by the holosymmetric channel.

Remark 4.297 (Why the s–channel in the wordwise setting)**.** Under the words-only meta-claim conditions (Proposition 4.176), the ss block is central to leading order, while the ps/sp edge is nilpotent; hence the loop operator $\hat{H}_{\ell}^{(s)[\mathbf{W}]}$ serves as a stable coherence reference, with deviations from centrality controlled by the compliance vector $\mathfrak{K} = (\Delta, \eta, \zeta)$ (Definition 4.178 and Proposition 4.181).

Corollary 4.298 (Wordwise small-loop sensitivity (first order))**.** *Let ℓ_{ε} be a family of contractible loops of area $O(\varepsilon^2)$ based at x. Then*

$$\hat{H}_{\ell_{\varepsilon}}^{(s)[\mathbf{W}]}(x) = \Big(\hat{\mathcal{F}}_{s}^{[\mathbf{W};\chi\leftarrow\chi]}(x)\Big)^{\dagger}\Big(\mathbf{1}_4 - \varepsilon^2\,\Phi_x(\mathcal{F}) + O(\varepsilon^3)\Big)\hat{\mathcal{F}}_{s}^{[\mathbf{W};\chi\leftarrow\chi]}(x),$$

for a linear functional $\Phi_x(\mathcal{F})$ of the curvature two–form determined by the loop's oriented area (non–Abelian Stokes expansion). Consequently, $h_{\ell_{\varepsilon}}^{(s)[\mathbf{W}]}(x) = h_{\mathrm{flat}}^{(s)[\mathbf{W}]}(x) - \varepsilon^2\,\omega^{\mathring{\Psi}}\Big((\hat{\mathcal{F}}_{s}^{[\cdot]})^{\dagger}\Phi_x(\mathcal{F})\hat{\mathcal{F}}_{s}^{[\cdot]}\Big) + O(\varepsilon^3)$.

Idea. Use the Dyson expansion of $\mathcal{P}\exp(-\oint_{\ell}\mathcal{A})$ and the non–Abelian Stokes theorem to first order in the loop area; insert into (4.122) and evaluate the expectation. □

Proof. Fix a smooth oriented disk Σ_{ε} with $\partial\Sigma_{\varepsilon} = \ell_{\varepsilon}$ based at x. By endpoint gauge covariance (Proposition 4.54) choose a local gauge with $g(x) = \mathbf{1}$ and $\mathcal{A}_{\mu}(x) = 0$ (e.g. radial/normal gauge around x). Write the loop holonomy $\mathcal{U}_{\ell_{\varepsilon}}(x) = \mathcal{P}\exp\Big(-\oint_{\ell_{\varepsilon}}\mathcal{A}\Big) \in \mathrm{U}(4)$ (Definition 4.56), and expand the path–ordered exponential by the Dyson series. Because $\mathcal{A}(x) = 0$ and ℓ_{ε} has linear size $O(\varepsilon)$, the first nontrivial contribution is $O(\varepsilon^2)$. Using Stokes' theorem together with the definition of curvature $\mathcal{F} = \mathrm{d}\mathcal{A} + \mathcal{A}\wedge\mathcal{A}$ (Definition 4.56) and the standard (non–Abelian) small–loop expansion, one obtains

$$\mathcal{U}_{\ell_{\varepsilon}}(x) = \mathbf{1}_4 - \int_{\Sigma_{\varepsilon}}\mathcal{F}(x) + O(\varepsilon^3) = \mathbf{1}_4 - \Phi_x(\mathcal{F}) + O(\varepsilon^3), \tag{4.130}$$

where the linear functional $\Phi_x(\mathcal{F})$ is the contraction of $\mathcal{F}$ at x with the oriented area bivector of Σ_ε:

$$\Phi_x(\mathcal{F}) \;:=\; \sum_{\mu<\nu} \Sigma^{\mu\nu}(\ell_\varepsilon)\, \mathcal{F}_{\mu\nu}(x), \qquad \Sigma^{\mu\nu}(\ell_\varepsilon) := \int_{\Sigma_\varepsilon} d\sigma^{\mu\nu}\,,$$

so $\Phi_x(\mathcal{F}) = O(\varepsilon^2)$. The $O(\varepsilon^3)$ remainder collects (i) higher Dyson terms and (ii) Taylor errors from replacing $\mathcal{F}(y)$ by $\mathcal{F}(x)$ inside the surface integral; both scale as area×diameter $= O(\varepsilon^3)$. The appearance of $\mathcal{F}$ as the infinitesimal generator of noncommutativity is consistent with the curvature identity $[\mathbb{D}^{[\cdot]}_\mu, \mathbb{D}^{[\cdot]}_\nu]X = [\mathcal{F}_{\mu\nu}, X]$ (Proposition 4.224).

IInsert (4.130) into the wordwise s–loop operator

$$\hat{H}^{(s)[\mathbf{W}]}_{\ell_\varepsilon}(x) = \Big(\hat{\mathcal{F}}^{[\mathbf{W};\chi\leftarrow\chi]}_s(x)\Big)^\dagger \mathcal{U}_{\ell_\varepsilon}(x)\, \hat{\mathcal{F}}^{[\mathbf{W};\chi\leftarrow\chi]}_s(x)$$

(eq. (4.122)) to get

$$\hat{H}^{(s)[\mathbf{W}]}_{\ell_\varepsilon}(x) = \Big(\hat{\mathcal{F}}^{[\mathbf{W};\chi\leftarrow\chi]}_s(x)\Big)^\dagger \Big(\mathbf{1}_4 - \Phi_x(\mathcal{F}) + O(\varepsilon^3)\Big) \hat{\mathcal{F}}^{[\mathbf{W};\chi\leftarrow\chi]}_s(x),$$

which is the stated operator expansion (recall $\Phi_x(\mathcal{F}) = O(\varepsilon^2)$). Taking the Holo–State expectation (eq. (4.123)) yields

$$h^{(s)[\mathbf{W}]}_{\ell_\varepsilon}(x) = \underbrace{\omega^{\mathring{\mathbf{\Psi}}}\Big((\hat{\mathcal{F}}^{[\cdot]}_s)^\dagger \hat{\mathcal{F}}^{[\cdot]}_s\Big)}_{=:\; h^{(s)[\mathbf{W}]}_{\mathrm{flat}}(x)} - \omega^{\mathring{\mathbf{\Psi}}}\Big((\hat{\mathcal{F}}^{[\cdot]}_s)^\dagger\, \Phi_x(\mathcal{F})\, \hat{\mathcal{F}}^{[\cdot]}_s\Big) \;+\; O(\varepsilon^3),$$

i.e. $h^{(s)[\mathbf{W}]}_{\ell_\varepsilon}(x) = h^{(s)[\mathbf{W}]}_{\mathrm{flat}}(x) - \varepsilon^2\, \omega^{\mathring{\mathbf{\Psi}}}\Big((\hat{\mathcal{F}}^{[\cdot]}_s)^\dagger \Phi_x(\mathcal{F}) \hat{\mathcal{F}}^{[\cdot]}_s\Big) + O(\varepsilon^3)$. Gauge covariance of $\mathcal{U}$ and the endpoint fields (Propositions 4.54 and 4.294 and Corollary 4.64) implies that $\hat{H}^{(s)[\mathbf{W}]}_{\ell_\varepsilon}(x)$ transforms by conjugation, hence the scalar $h^{(s)[\mathbf{W}]}_{\ell_\varepsilon}(x)$ is gauge invariant; in the flat, simply connected case $\mathcal{F} \equiv 0$ one recovers the trivial loop value (Proposition 4.295). □

How Quantum Logic Weaves Spacetime

Outcome, not stage. *Geometry is inferred, not assumed.* Distances, causal dispositions, and the very notions of "where" and "when" are read as *expectation values* of the Fourfold inter-modal correlator in the Holo-State (Definition 4.9, Section 4.6, and eq. (4.85)). This is *operational background independence*: the same quantum relations that glue local modal fibres into a bundle (Definitions 4.239 and 4.247) also generate the geometric data that later constrain those relations via the connection $\mathcal{A}$ and its transporter $\mathcal{U}$ (Definition 4.248 and Section 4.1.4). In this way, the framework closes the classic GR–QFT gap by letting the "arena" itself be *derived* from correlator structure.

State-relative arena and learned subsystems. The working Hilbert space is *state-relative*: the physically active subspace and observables are carved out by the actual Holo-State $\omega^{\mathring{\mathbf{\Psi}}}$ through the modal fibre and its inner product (Definition 4.266 and eqs. (4.100) and (4.102)). There is likewise *no a priori tensor product*: "subsystems" are *learned* from (i) the Fourfold

block structure and its off-diagonal couplings (Definition 4.81 and Proposition 4.246), and (ii) the connection that transports modal data between loci (Definition 4.248). Entanglement is therefore not merely "in'' spacetime; it is part of the mechanism that *weaves* it, via transport and gluing of sections (Remark 4.267, Lemma 4.270, and Corollary 4.251).

Logic → law: selection rules from the Fourfold. Kinematics and dynamics are constrained by a *Fourfold Dialectic Algebra* with $\mathbb{Z}_2 \times \mathbb{Z}_2$ grading (Section 4.1.7 and Definition 4.94). The gradings act as *selection rules* on allowable inter-modal moves. In the endogenous, wordwise reading this appears constructively as the "words-only trio'': central ss, nilpotent ps/sp, and bosonic $0/s$ compatibility (Proposition 4.176). When coherence is lost, the theory *collapses to the Twofold block* (Thesis/Antithesis), realized here as the $(0,1)$ subbundle (Definition 4.245) with suppressed mixing into (p,s) when the relevant off-diagonal channels are small (Proposition 4.246). Thus the classical rules (non-contradiction, excluded middle) arise as a modal limit that the logic itself predicts.

A unified venue and new observables. Geometry and matter share a *common correlator architecture*—a 2×2 field block whose entries are 4×4 in modal space—so their co-emergence is described in a single venue through the Fourfold correlator (Definitions 4.9 and 4.81). This architecture suggests concrete, testable quantities: *holonomies* of the modal connection (Section 4.7.2). Transporting a mode around a closed loop ℓ yields a measurable change captured by the holosymmetric loop operator $\hat{H}_\ell^{(s)}(x) = \hat{G}_\Lambda^{ss}(x,x;\ell)$ and its expectation (eqs. (4.119) and (4.120)); in the endogenous calculus, the same signal is extracted from the word-evaluated s–self channel (Section 4.7.3 and eqs. (4.122) and (4.123)). Flat, simply connected regions trivialize the loop (Corollary 4.251 and Proposition 4.295); nonzero curvature or nontrivial topology produce robust, implementation-invariant phases (Aharonov–Bohm–type phenomena).

From ideas to inequalities. Because the framework is operational, intuition is pinned down by *bounds*. The holosymmetric compliance vector $\mathfrak{K} = (\Delta, \eta, \zeta)$ (Definition 4.178) controls measurable diagnostics—visibility envelopes, lensing residuals, two-step ps/sp amplitudes, and diagonal pinching—via Proposition 4.181 (with pinching stability in Lemma 4.180 and its exact conditional-expectation form in Remark 4.182). Thus departures from idealized logic are *quantified* and experimentally targetable.

Conceptual upshot (relational geometry, derived). Reality = relations. Physics is reframed as a *relational, graded, fibered Hilbert geometry* whose structures are *derived* from quantum-logical correlators rather than stipulated. The "space of states'' (global sections and their inner products, Definition 4.266 and Remark 4.268) and the "spacetime of events'' (glued by $\mathcal{A}$ and probed by holonomy, Definition 4.248 and Section 4.7.2) are outputs of the same engine. If borne out, this turns the usual scaffolding inside-out—and provides new levers (holonomy signals and compliance-vector diagnostics) to test how the universe *knits itself together* (Section 4.7.3 and Proposition 4.181).

4.7.4 The Abstract Qubit

From abstract bits to emergent modal structures. The transition from the conventional *qubit* to the *modal Hilbert space of a Primordial Quantum Singularity* (PQS) is a principled relocation of abstraction. In standard quantum mechanics, a qubit is an axiomatic object—an element of the isomorphism class of pairs $(\mathbb{C}^2, \langle\cdot,\cdot\rangle)$ endowed with the Born rule [379]. By contrast, within the IQR framework the relevant degrees of freedom are *localized, emergent phenomena*: the modal fibres $\mathcal{H}_{\text{mod}}(\mathfrak{B}_x)$ of a globally structured Hilbert bundle whose metric data are *induced* rather than postulated, and whose geometry is fixed by the correlator-driven fourfold dialectical structure (Definitions 4.9 and 4.273, Remark 4.253, and Sections 4.1.2 and 4.1.3). In this relocation, an "elementary quantum of information'' is not a static amplitude container but a *context-sensitive nexus of relations* determined by inter-PQS structure across the emergent configuration space $\mathcal{X}_q$.

The Abstract Qubit (Effective Limit). The conventional qubit is a unit vector in a fixed two-dimensional space $\mathcal{H} \cong \mathbb{C}^2$,

$$|\psi\rangle = \alpha|0\rangle + \beta|1\rangle, \qquad \alpha,\beta \in \mathbb{C}, \quad |\alpha|^2 + |\beta|^2 = 1,$$

with statistics supplied by the Born rule and multi-qubit structure given by $\mathcal{H}_2^{\otimes n}$. This architecture presupposes a background Hilbert space and does not itself encode emergent state-space geometry or internal dialectical dynamics. Laboratory implementations (ions, superconductors, photons, NV, quantum dots) instantiate this abstraction but decohere, motivating error correction [351, 380, 349]. IQR does not dispute any of this; it *relocates* the qubit, showing it as an *effective* face of a deeper, correlator-induced modal geometry.

State-dependent Arena via GNS and the Fourfold

At the foundational level, the Fourfold Dialectic Algebra of Symmetry provides the $*$-algebraic substrate; the universal Holo-State $|\mathring{\mathbf{\Psi}}\rangle$ endows it with expectations, and the GNS construction yields the concrete representation $(\pi_\Psi, \mathcal{H}_{\text{em}}, |\Omega_\Psi\rangle)$ (Section 1.4.2 and Definition 1.27). In particular, the *inner product on each modal fibre* is defined by the coincident-limit of the Fourfold Inter-Modal Correlator evaluated in $|\mathring{\mathbf{\Psi}}\rangle$ (Section 4.6 and Definition 4.228). Thus the *Hilbert arena is state-dependent*: representation first, background later.

Remark 4.299 (Relational Emergence of Modal Geometry)**.** The modal fibre $\mathcal{H}_{\text{mod}}(\mathfrak{B}_x)$ is fixed only relationally—by its couplings to neighbouring singularities $\mathfrak{B}_{x'}$ through $\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})$ and by the global state $|\mathring{\mathbf{\Psi}}\rangle$. This implements, at the operator level, the dictum that distinguishability and geometry crystallize *between* PQSs rather than *at* them.

Twofold Poles Within a Genuinely Fourfold Fibre

Inside each fibre $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}$ there are four ontological modes $\{|0\rangle_x, |1\rangle_x, |p\rangle_x, |s\rangle_x\}$ realising Symmetry, Antisymmetry, Parasymmetry, and Holosymmetry (Sections 1.2.7 and 2.1). The pair $|0\rangle_x, |1\rangle_x$ plays the role of *dyadic poles* of the Twofold sector, but only *after* the fibre's inner product has been *emergently fixed* from the correlator; by construction, $|p\rangle_x$ and $|s\rangle_x$ are

independent basis rays—not mere superpositions within the Twofold qubit dyad (Section 4.6). Hence the qubit-like axis is *recovered* as an internal sector of a genuinely Fourfold modality, not presupposed.

Qubit Reduction Theorem

Proposition 4.300 (Qubit Reduction)**.** *Assume that at $\mathfrak{B}_x$ there exist two orthonormal modal rays $|\alpha_x\rangle, |\beta_x\rangle \in \mathcal{H}_{\mathrm{mod}}(\mathfrak{B}_x)$ stabilised by the local dialectic generators, and that the inter-modal couplings are block-diagonalisable so that back-reaction on $S_x := \mathrm{span}\{|\alpha_x\rangle, |\beta_x\rangle\}$ is negligible over the control timescale. Then there exists an isometric embedding $U : \mathbb{C}^2 \hookrightarrow \mathcal{H}_{\mathrm{mod}}(\mathfrak{B}_x)$ with $U|0\rangle = |\alpha_x\rangle$ and $U|1\rangle = |\beta_x\rangle$, such that:*

(i) Born statistics on $\mathbb{C}^2$ agree with those induced by the emergent fibre inner product on $\mathcal{H}_{\mathrm{mod}}(\mathfrak{B}_x)$ (as defined via the correlator/GNS machinery in Section 4.6), and

(ii) admissible local controls act (on S_x) as an effective SU(2)*.*

Proof. Step 0 (Set-up). Work in any orthonormal trivialisation $(\mathcal{U}, e_{\mathcal{U}})$ around x with fibre inner product $\langle\cdot,\cdot\rangle_x$. The scalar pairing of sections is independent of trivialisation and G-gauge, by Lemma 4.270 (in particular (1) therein). Hence all constructions below are intrinsic to the fibre at x.

Step 1 (Isometric embedding). Since $\{|\alpha_x\rangle, |\beta_x\rangle\}$ is an orthonormal pair in $\mathcal{H}_{\mathrm{mod}}(\mathfrak{B}_x)$, define $U : \mathbb{C}^2 \to \mathcal{H}_{\mathrm{mod}}(\mathfrak{B}_x)$ by $U|0\rangle = |\alpha_x\rangle$, $U|1\rangle = |\beta_x\rangle$ and extend linearly. Then, for any $|\phi\rangle, |\psi\rangle \in \mathbb{C}^2$,

$$\langle \phi, \psi\rangle_{\mathbb{C}^2} = \langle U\phi, U\psi\rangle_x,$$

so U is an isometry onto $S_x = \mathrm{span}\{|\alpha_x\rangle, |\beta_x\rangle\}$. (This uses only orthonormality in the emergent fibre metric; cf. the construction preceding Proposition 4.300.)

Step 2 (Born statistics agree). Let E be any effect on $\mathbb{C}^2$ (i.e. $0 \le E \le \mathbb{1}_2$). Define the corresponding fibre effect $\widetilde{E} := UEU^\dagger$ acting on $\mathcal{H}_{\mathrm{mod}}(\mathfrak{B}_x)$. For any $|\psi\rangle \in \mathbb{C}^2$,

$$\langle\psi|E|\psi\rangle = \langle U\psi|\, UEU^\dagger\, |U\psi\rangle_x = \langle U\psi|\, \widetilde{E}\, |U\psi\rangle_x,$$

so Born valuations coincide by isometry. Moreover, the map $E \mapsto \widetilde{E} = UEU^\dagger$ is a unital $*$-monomorphism of effect algebras and therefore preserves POVMs, making the operational equivalence explicit. Since the modal inner product is induced by the Fourfold Inter-Modal Correlator via the GNS state (the state-dependent arena mechanism), this establishes (i).

Step 3 (Effective SU(2) *controls on S_x).* Let $\mathfrak{A}_x$ denote the Lie algebra of admissible local control generators at x (skew-Hermitian operators produced by the local dialectic generators/connection; see the covariant Leibniz setting). Stabilisation of the rays means S_x is invariant: for every $A \in \mathfrak{A}_x$, $AS_x \subseteq S_x$. Let P be the orthogonal projector onto S_x and define the compressed representation $\rho : \mathfrak{A}_x \to \mathfrak{u}(2)$ by $\rho(A) = PAP|_{S_x}$. Then $\rho(\mathfrak{A}_x)$ is a Lie subalgebra of $\mathfrak{u}(2)$. If $\rho(\mathfrak{A}_x)$ contains two linearly independent traceless, non-commuting elements, it generates $\mathfrak{su}(2)$ (standard Lie-generation criterion). Exponentiating yields a connected subgroup of U(2) whose commutator algebra is $\mathfrak{su}(2)$, i.e. SU(2) up to overall phase. Thus admissible controls act as effective SU(2) on S_x, giving (ii). (In the present framework the existence of at least two such independent compressed generators is ensured once the Twofold sector is non-abelian; equivalently, once the top-left block of the Fourfold

correlator is non-degenerate, which is the generic case discussed in the Twofold-vs-Fourfold analysis.)

Step 4 (On block-diagonalisable couplings and negligible back-reaction). Let the full local generator (drift + controls) decompose as

$$H = \begin{pmatrix} H_{S_x} & 0 \\ 0 & H_{S_x^\perp} \end{pmatrix} + \begin{pmatrix} 0 & K \\ K^\dagger & 0 \end{pmatrix} =: H_0 + \Delta, \qquad \|\Delta\| = \|K\| = \varepsilon.$$

Let $\mathcal{U}(t) := e^{-itH}$ be the propagator. By Duhamel's formula [381, 241, 287],

$$\mathcal{U}(t) - e^{-itH_0} = -i \int_0^t \mathcal{U}(t-s)\, \Delta\, e^{-isH_0}\, ds,$$

and, since H, H_0 are self-adjoint, $\|\mathcal{U}(\tau)\| = \|e^{-i\tau H_0}\| = 1$. Hence

$$\|(I-P)\,\mathcal{U}(t)\,P\| \;\le\; \int_0^t \|\Delta\|\, ds \;=\; \varepsilon t,$$

so leakage amplitude from S_x is $O(\varepsilon t)$ and probabilities $O((\varepsilon t)^2)$. In the regime "negligible over the control timescale" ($\varepsilon t \ll 1$), the action on S_x is well-approximated by $e^{-itH_{S_x}} \in \mathrm{U}(2)$, i.e. the effective SU(2) dynamics above.

Step 5 (Gauge/trivialisation independence). The constructions are independent of the choice of unitary frame and extend verbatim to h-orthonormal frames: replace $U(4)$ by the h-preserving group $G_h = \{g \mid g^\dagger h g = h\}$ and use metric-compatibility $\partial_\mu h - \mathcal{A}_\mu^\dagger h - h\mathcal{A}_\mu = 0$ to retain the covariant Leibniz rule; all equalities then hold with pairings $(c^\Phi)^\dagger h\, c^\Psi$ (cf. Remark 4.271).

This completes the proof. □

Context and links to earlier structure. The Twofold sector S_x is precisely the qubit-like "top-left block" inside the Fourfold correlator; when off-block mixings are negligible, ordinary qubit kinematics is recovered; when mixing with $\{p, s\}$ is appreciable, new (non-qubit) phenomena arise—exactly as stated in the surrounding text.

Why include this proof? It (a) makes explicit where the isometry, statistics-equivalence, and SU(2) control action come from in the bundle language; (b) isolates the approximation in a single norm bound; and (c) shows gauge-independence and the h-orthonormal generalisation are already baked in via the preceding lemma/remark on fibre pairings and metric-compatible connections.

Corollary 4.301 (Exact vs. Approximate Qubit Reduction)**.** *Retain the setting of Proposition 4.300 with $S_x = \mathrm{span}\{|\alpha_x\rangle, |\beta_x\rangle\}$ and orthogonal projector P onto S_x. Let the local generator (drift + controls) decompose as*

$$H \;=\; \underbrace{\begin{pmatrix} H_{S_x} & 0 \\ 0 & H_{S_x^\perp} \end{pmatrix}}_{=:H_0} + \underbrace{\begin{pmatrix} 0 & K \\ K^\dagger & 0 \end{pmatrix}}_{=:\Delta}, \qquad \|\Delta\| = \|K\| \le \varepsilon.$$

(a) Exact reduction ($\varepsilon = 0$). *If* $\Delta = 0$, *then for all* t,

$$\mathcal{U}(t) = e^{-itH} = e^{-itH_{S_x}} \oplus e^{-itH_{S_x^\perp}}, \quad \text{and} \quad P\,\mathcal{U}(t)\,P \;=\; e^{-itH_{S_x}} \in \mathrm{U}(2).$$

Thus the isometric embedding $U : \mathbb{C}^2 \to \mathcal{H}_{\mathrm{mod}}(\mathfrak{B}_x)$ *from Proposition 4.300 is an* exact *qubit equivalence: Born statistics and admissible controls coincide with those on the effective* SU(2) *dynamics on* S_x.

(b) Approximate reduction ($\varepsilon > 0$). *For* $\mathcal{U}(t) = e^{-itH}$ *the leakage from* S_x *into* $S_x^\perp$ *and the deviation of the compressed evolution from the ideal* $e^{-itH_{S_x}}$ *obey, for all* $t \geq 0$,

$$\left\|(I - P)\,\mathcal{U}(t)\,P\right\| \;\leq\; \varepsilon\, t, \qquad \left\|P\,\mathcal{U}(t)\,P \;-\; e^{-itH_{S_x}}\right\| \;\leq\; \varepsilon\, t.$$

In particular, for any unit vector $|\psi\rangle \in S_x$,

$$\|(I - P)\,\mathcal{U}(t)\,|\psi\rangle\,\|^2 \;\leq\; (\varepsilon t)^2,$$

so on timescales with $\varepsilon t \ll 1$ *the effective* SU(2) *description on* S_x *holds to first order in* εt.

Proof. For (a), $\Delta = 0$ makes H block-diagonal, yielding the direct-sum evolution and the exact identity $P\,\mathcal{U}(t)\,P = e^{-itH_{S_x}}$.

For (b), write $H = H_0 + \Delta$ and use Duhamel's formula

$$\mathcal{U}(t) - e^{-itH_0} = -i\int_0^t \mathcal{U}(t-s)\,\Delta\,e^{-isH_0}\,ds.$$

Since H and H_0 are self-adjoint, $\|\mathcal{U}(\tau)\| = \|e^{-i\tau H_0}\| = 1$; thus

$$\|\mathcal{U}(t) - e^{-itH_0}\| \;\leq\; \int_0^t \|\Delta\|\,ds \;=\; \varepsilon t.$$

Left- and right-multiplying by $I - P$ and P yields the leakage bound $\|(I - P)\,\mathcal{U}(t)\,P\| \leq \varepsilon t$. Moreover,

$$P\,\mathcal{U}(t)\,P - e^{-itH_{S_x}} = P\Big(\mathcal{U}(t) - e^{-itH_0}\Big)P + P\Big(e^{-itH_0} - e^{-itH_{S_x}} \oplus e^{-itH_{S_x^\perp}}\Big)P,$$

and the second term vanishes by block-diagonality of H_0, giving $\|P\,\mathcal{U}(t)\,P - e^{-itH_{S_x}}\| \leq \|\mathcal{U}(t) - e^{-itH_0}\| \leq \varepsilon t$, as stated. The probability bound follows immediately from the leakage estimate. □

Corollary 4.302 (Registers)**.** *Let* $x_1, \ldots, x_n$ *be* n *distinct PQS loci. For each* j *assume the hypotheses of Proposition 4.300: there exists a stabilised two-dimensional sector* $S_{x_j} = \mathrm{span}\{|\alpha_{x_j}\rangle, |\beta_{x_j}\rangle\} \subset \mathcal{H}_{\mathrm{mod}}(\mathfrak{B}_{x_j})$ *with isometric embedding* $U_j : \mathbb{C}^2 \hookrightarrow \mathcal{H}_{\mathrm{mod}}(\mathfrak{B}_{x_j})$, $U_j\,|0\rangle = |\alpha_{x_j}\rangle$, $U_j\,|1\rangle = |\beta_{x_j}\rangle$. *Assume further that cross–inter-modal correlator blocks between distinct fibres are perturbatively suppressed in the sense of Section 4.1.2, so that the global generator decomposes as*

$$H \;=\; \sum_{j=1}^{n} H^{(j)} \;+\; \sum_{i<j} H^{(ij)} \;+\; \Delta, \qquad \|\Delta\| \leq \varepsilon,$$

where $H^{(j)}$ acts locally on $\mathcal{H}_{\mathrm{mod}}(\mathfrak{B}_{x_j})$, $H^{(ij)}$ is a controlled (switchable) two-fibre coupling, and Δ collects off-block terms.

Then the map

$$U^{(n)} \;:=\; U_1 \otimes \cdots \otimes U_n : (\mathbb{C}^2)^{\otimes n} \longrightarrow \bigotimes_{j=1}^{n} \mathcal{H}_{\mathrm{mod}}(\mathfrak{B}_{x_j})$$

is an isometric embedding with image $S_{x_1} \otimes \cdots \otimes S_{x_n}$, and:

(a) Exact reduction ($\varepsilon = 0$)

If $\Delta = 0$ and each compressed local algebra $P_j H^{(j)} P_j$ generates $\mathfrak{su}(2)$ on S_{x_j}, then the effective arena is exactly $(\mathbb{C}^2)^{\otimes n}$ under $U^{(n)}$, and any switchable two-site interaction with nontrivial compressed part $P_{ij} H^{(ij)} P_{ij} \notin \mathfrak{u}(2) \oplus \mathfrak{u}(2)$ ($P_{ij} = P_i \otimes P_j$) yields an entangling two-qubit gate on the (i, j) register. In particular, single-qubit $\mathrm{SU}(2)$ on each site together with any fixed entangling two-qubit gate forms a universal gate set on $(\mathbb{C}^2)^{\otimes n}$ [287].

(b) Approximate reduction ($\varepsilon > 0$)

Let $\mathcal{U}(t) = e^{-itH}$ and $P = \bigotimes_{j=1}^{n} P_j$ with P_j the orthogonal projector onto S_{x_j}. Then, for all $t \geq 0$,

$$\|(I - P)\,\mathcal{U}(t)\,P\| \;\leq\; \varepsilon\, t, \qquad \Big\| P\,\mathcal{U}(t)\,P - e^{-it\sum_j P_j H^{(j)} P_j - it\sum_{i<j} P_{ij} H^{(ij)} P_{ij}} \Big\| \;\leq\; \varepsilon\, t,$$

so leakage outside $S_{x_1} \otimes \cdots \otimes S_{x_n}$ is $O\big((\varepsilon t)^2\big)$ and the compressed dynamics approximates a unitary on $(\mathbb{C}^2)^{\otimes n}$ to first order in εt.

Proof. Isometry of $U^{(n)}$ is immediate from that of each U_j and the tensor structure. Part *(a)* follows from block-diagonality ($\Delta = 0$), the sitewise $\mathfrak{su}(2)$ generation (by Proposition 4.300), and the fact that a two-qubit unitary with generator whose compression is not a sum of one-body terms is entangling; universality with local SU(2) plus any entangling two-qubit gate is standard [287]. Part *(b)* is the multi-site version of the Duhamel estimate used in Corollary 4.301: write $H = H_0 + \Delta$ with $H_0 = \sum_j H^{(j)} + \sum_{i<j} H^{(ij)}$ block-diagonal on $S_{x_1} \otimes \cdots \otimes S_{x_n}$, apply

$$\mathcal{U}(t) - e^{-itH_0} = -i \int_0^t \mathcal{U}(t - s)\,\Delta\, e^{-isH_0}\, ds,$$

use $\|\mathcal{U}(\tau)\| = \|e^{-i\tau H_0}\| = 1$, and then left/right compress by P to obtain the stated bounds. □

Remark 4.303 (Twofold block as a limit of the Fourfold correlator)**.** The Twofold correlator $\hat{\mathbf{G}}_{(0,1)}(x, x')$ is the top-left 2×2 block of the full Fourfold matrix $\big[\hat{G}_{\Lambda}^{\alpha\beta}(x, x')\big]_{\alpha,\beta \in \{0,1,p,s\}}$. Negligible off-block terms reproduce conventional qubit kinematics; non-negligible mixing with $\{p, s\}$ drives strictly non-qubit phenomena.

The Qubit in IQR: Epistemic, Operational, and Geometric Synthesis

Epistemic status (abstract type vs. derived sector). In standard QM/QI the qubit is an *abstract type*—an isomorphism class of pairs $(\mathbb{C}^2, \langle\cdot,\cdot\rangle)$ with the Born rule [379, 287]. In the IQR framework, by contrast, the qubit is *derived*: it is the equivalence class (modulo SU(2) gauge) of isometric embeddings

$$U : \mathbb{C}^2 \hookrightarrow \mathcal{H}_{\mathrm{mod}}(\mathfrak{B}_x), \qquad U\,|0\rangle = |\alpha_x\rangle\,, \quad U\,|1\rangle = |\beta_x\rangle\,,$$

when a dynamically protected Twofold sector $S_x := \mathrm{span}\{|\alpha_x\rangle\,, |\beta_x\rangle\}$ exists (Definition 4.273, Proposition 4.300, and Corollary 4.301).

Operational continuity (laboratory import). Laboratory qubits remain effective two-level carriers; the IQR construction supplies *deeper semantics*—why such sectors exist, how their inner products are *induced*, and when the reduction is valid. Standard practical concerns (decoherence, control, and fault-tolerance) retain their usual form in the effective regime [351, 380, 349].

Two narratives, contrasted.

(i) *Classical (axiomatic/postulatory) narrative.* One *postulates* a fixed two-dimensional complex Hilbert space $\mathbb{C}^2$ with its inner product, Born rule, and unitary controls; a pure qubit is any unit vector in $\mathbb{C}^2$ up to global phase, and composites are $(\mathbb{C}^2)^{\otimes n}$ [379, 287]. The dimension, metric, and control algebra are taken *as givens*.

(ii) *IQR (derivational) narrative.* One begins with the Fourfold Dialectic operator algebra at each PQS and the universal Holo-State. From these:

(a) the fibre inner product on each modal space is *induced* from the Fourfold Inter-Modal Correlator via the GNS procedure—hence the metric is *state-dependent*, not a background axiom (Section 4.6 and Definition 4.9);

(b) under explicit stability/decoupling hypotheses (block-diagonalisable couplings with negligible back-reaction) a two-dimensional sector $S_x \subset \mathcal{H}_{\mathrm{mod}}(\mathfrak{B}_x)$ is *selected*, and an isometric embedding $U : \mathbb{C}^2 \hookrightarrow \mathcal{H}_{\mathrm{mod}}(\mathfrak{B}_x)$ is exhibited (Proposition 4.300);

(c) admissible local generators, compressed to S_x, *realise* the effective control algebra $\mathfrak{su}(2)$, yielding SU(2) dynamics; measurement statistics agree because $E \mapsto UEU^\dagger$ is a unital $*$-monomorphism preserving POVMs (Proposition 4.300);

(d) when off-block couplings are small but nonzero, Duhamel bounds *quantify* the approximation; exact/approximate regimes are recorded, and an n-site register $(\mathbb{C}^2)^{\otimes n}$ emerges when cross-correlators are perturbatively suppressed (Corollaries 4.301 and 4.302 and Section 4.1.2).

Geometry as expectation and the information–geometry link. The same state-dependence that induces inner products on modal fibres also determines vacuum geometry:

$$\bar{\Sigma}_{\mu\nu}(x) = \left\langle \Omega_\Psi \middle| \hat{\Sigma}_{\mu\nu}(x) \middle| \Omega_\Psi \right\rangle,$$

so the relocation of the qubit (as a reduced Twofold sector of a Fourfold fibre) and the emergence of spacetime geometry share a single operator-theoretic mechanism: both are *expectation-value* effects of one Holo-State over a common algebraic substrate (Definition 5.26).

Bottom line. The treatise does not *supplant* the qubit; it *grounds and generalises* it. In correlator-sparse regimes the Twofold block reduces to conventional qubit kinematics; beyond that, parasymmetric/holosymmetric mixing carries one *beyond* the qubit paradigm. The abstract qubit survives as an effective type and a useful instrument, now situated within a background-free, correlator-generated modal geometry [241, 381, 287].

4.7.5 The Abstract Qubit (Wordwise Relocation)

From abstract bits to wordwise modal structure. In the endogenous calculus, an "elementary quantum of information'' is not a fixed $\mathbb{C}^2$ but a *protected two-letter subalphabet* inside a word–rotated modal fibre. The fibre metric is induced (not postulated) from the Fourfold correlator, while device words $\mathbf{W}$ select operational frames and channels (Definitions 4.255 and 4.257 and Proposition 4.258).

Definition 4.304 (Wordwise twofold sector at a PQS)**.** Fix $x \in \mathcal{X}_q$, a unitary device word $\mathbf{W}$, and control χ. Let $\mathcal{B}_x^{[\mathbf{W},\chi]} = \{|0\rangle_x^{[\mathbf{W},\chi]}, |1\rangle_x^{[\mathbf{W},\chi]}, |p\rangle_x^{[\mathbf{W},\chi]}, |s\rangle_x^{[\mathbf{W},\chi]}\}$ be the word–rotated orthonormal frame (Definition 4.257 and Proposition 4.258). Define the *wordwise twofold sector*

$$S_x^{[\mathbf{W}]} := \operatorname{span}\Big\{|0\rangle_x^{[\mathbf{W},\chi]}, |1\rangle_x^{[\mathbf{W},\chi]}\Big\} \subset \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}.$$

The parasymmetric/holosymmetric complement is $S_{x,\perp}^{[\mathbf{W}]} := \operatorname{span}\Big\{|p\rangle_x^{[\mathbf{W},\chi]}, |s\rangle_x^{[\mathbf{W},\chi]}\Big\}$.

Definition 4.305 (Wordwise qubit realisation)**.** The *wordwise abstract qubit at x* is an isometric embedding $U^{[\mathbf{W}]} : \mathbb{C}^2 \hookrightarrow \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ defined by

$$U^{[\mathbf{W}]}|0\rangle = |0\rangle_x^{[\mathbf{W},\chi]}, \qquad U^{[\mathbf{W}]}|1\rangle = |1\rangle_x^{[\mathbf{W},\chi]},$$

with the induced fibre inner product from the Fourfold correlator (Section 4.6 and eq. (4.85)). Thus any $|\psi\rangle \in \mathbb{C}^2$ is represented in the fibre as $U^{[\mathbf{W}]}|\psi\rangle \in S_x^{[\mathbf{W}]}$ with the same Born valuations.

Proposition 4.306 (Wordwise qubit reduction and control)**.** *Assume the word–evaluated off–diagonal blocks that couple $S_x^{[\mathbf{W}]}$ to $S_{x,\perp}^{[\mathbf{W}]}$ are (exactly) zero or (approximately) small in operator norm, as diagnosed by $\hat{\mathbf{G}}_\Lambda^{[\mathbf{W}]}(x,x')$ (Proposition 4.262). Then:*

(i) (Isometry & statistics). *$U^{[\mathbf{W}]}$ is an isometric embedding onto $S_x^{[\mathbf{W}]}$, and $E \mapsto \widetilde{E} := U^{[\mathbf{W}]} E \big(U^{[\mathbf{W}]}\big)^\dagger$ preserves POVMs and Born probabilities in the fibre metric.*

(ii) (Effective SU(2) controls). *If the admissible local generators at x, compressed to $S_x^{[\mathbf{W}]}$, contain two independent traceless, noncommuting elements, they generate $\mathfrak{su}(2)$ on $S_x^{[\mathbf{W}]}$; hence admissible controls act as effective* SU(2) *on the wordwise qubit.*

(iii) (Leakage bounds). *If the residual off–block couplings have norm ε, then for the propagator $\mathcal{U}(t)$ generated by the full local Hamiltonian,*

$$\left\|(I - P^{[\mathbf{W}]})\,\mathcal{U}(t)\,P^{[\mathbf{W}]}\right\| \;\le\; \varepsilon\, t, \qquad P^{[\mathbf{W}]} := \text{orthogonal projector onto } S_x^{[\mathbf{W}]},$$

so leakage probability is $O((\varepsilon t)^2)$ (Duhamel estimate; cf. the exogenous Corollary 4.301).

Remark 4.307 (Role of compliance vector)**.** Operational deviations are controlled by the holosymmetric compliance vector $\mathfrak{K} = (\Delta, \eta, \zeta)$ (Definition 4.178): Δ bounds visibility/lensing drifts, η fixes two–step ps/sp amplitudes, and ζ controls diagonal pinching stability (Proposition 4.181, Lemma 4.180, and Remark 4.182). In particular, small η and ζ suppress mixing and misalignment of the twofold sector under wordwise transport, keeping the effective qubit clean.

Corollary 4.308 (Wordwise qubit registers across loci)**.** *Let $x_1, \dots, x_n$ be distinct PQS loci, each with a protected twofold sector $S_{x_j}^{[\mathbf{W}_j]}$ and embedding $U_j^{[\mathbf{W}_j]}$. Define $U^{(n)} := U_1^{[\mathbf{W}_1]} \otimes \cdots \otimes U_n^{[\mathbf{W}_n]}$. If cross–fibre off–block terms are perturbatively suppressed (Section 4.1.2 and Proposition 4.262), then $U^{(n)}$ is an isometry*

$$U^{(n)} : \; (\mathbb{C}^2)^{\otimes n} \;\hookrightarrow\; S_{x_1}^{[\mathbf{W}_1]} \otimes \cdots \otimes S_{x_n}^{[\mathbf{W}_n]},$$

and the compressed multi–site dynamics approximates a unitary on $(\mathbb{C}^2)^{\otimes n}$ with error $O(\varepsilon t)$ as in Corollary 4.302.

Remark 4.309 (Twofold poles inside a genuinely Fourfold fibre (wordwise))**.** In the wordwise frame, $|0\rangle_x^{[\mathbf{W},\chi]}, |1\rangle_x^{[\mathbf{W},\chi]}$ are the dyadic poles *within* a genuinely Fourfold modality; $|p\rangle_x^{[\mathbf{W},\chi]}$, $|s\rangle_x^{[\mathbf{W},\chi]}$ are independent rays (not twofold superpositions) (Proposition 4.281 and Remark 4.286). The abstract qubit is therefore a *derived* two-letter sector—stable when off–block words/channels are small—rather than a background postulate.

Remark 4.310 (Gauge and word choice)**.** Changing the endpoint gauge or replacing $\mathbf{W}$ by $\mathbf{W}'$ conjugates the presentation by a unitary on the fibre (Propositions 4.260 and 4.284 and Definition 4.259); the isometry class of the qubit sector and its Born statistics are unaffected.

Bottom line (endogenous lens). The "qubit'' becomes a *state– and device–relative* twofold word sector $S_x^{[\mathbf{W}]} \subset \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ whose existence and quality are *diagnosed and bounded* by word–evaluated channels and the compliance vector. When off–block couplings are negligible, standard SU(2) qubit kinematics is recovered; when mixing with $\{p, s\}$ is appreciable, the endogenous calculus predicts departures from the qubit paradigm and locates their operational signatures.

In the endogenous "alphabets & words'' view, a qubit is not postulated as a fixed $\mathbb{C}^2$ but recovered as a *protected two-letter sector* inside a word–rotated modal fibre. A unitary

device word $\mathbf{W}$ selects an operational frame $\mathcal{B}_x^{[\mathbf{W},\chi]}$ on the fibre $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ (Definition 4.257 and Proposition 4.258), whose inner product is *induced* from the Fourfold correlator (Section 4.6 and eq. (4.85)). The wordwise twofold sector $S_x^{[\mathbf{W}]}$ and the isometric embedding $U^{[\mathbf{W}]} : \mathbb{C}^2 \hookrightarrow \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ formalise the "abstract qubit'' as a derived subalphabet with inherited Born statistics (Definitions 4.304 and 4.305). Independence of the parasymmetric/holosymmetric directions is explicit—$|p\rangle, |s\rangle$ are not twofold superpositions—so the dyad $(0, 1)$ lives inside a genuinely Fourfold fibre (Proposition 4.281 and Remark 4.286).

Control and robustness become operational statements. Compressing admissible generators to $S_x^{[\mathbf{W}]}$ yields $\mathfrak{su}(2)$ whenever two traceless noncommuting compressed elements exist, giving effective single–qubit dynamics (Proposition 4.306). Departures from the ideal are *quantified*: word–evaluated off–block couplings bound mixing (Proposition 4.262), while the holosymmetric compliance vector $\mathfrak{K} = (\Delta, \eta, \zeta)$ controls visibility/lensing drifts, two–step *ps*/*sp* amplitudes, and diagonal pinching stability (Definition 4.178, Proposition 4.181, Lemma 4.180, and Remark 4.182). Thus the regime where the qubit picture holds—and how it fails—is specified by measurable inequalities rather than by assumption.

Geometry completes the picture. The modal connection $\mathcal{A}$ and its holonomy are read out in the s–channel via the wordwise loop operator $\hat{H}_\ell^{(s)[\mathbf{W}]}(x)$ and its expectation $h_\ell^{(s)[\mathbf{W}]}(x)$ (Section 4.7.3 and eqs. (4.122) and (4.123)). Flat, simply connected regions trivialise the loop signal, while curvature or topology produce robust phases (Proposition 4.295). In this way, the qubit emerges as a state–and device–relative code sector whose metric, controls, error modes, and geometric phases are all generated and certified by the same wordwise, Fourfold correlator machinery.

4.7.6 The Holomodal Quantum Computer

Read "quantum computer'' in the broad, model–independent sense of a controllable physical process implementing algorithmic transformations on a state space. In this setting the IQR framework both preserves today's qubit computing and *enlarges* the design space in principled ways. The qubit reappears only as a *dynamically protected Twofold sector* inside a genuinely Fourfold modal fibre; concretely, as a *wordwise* two–letter subalphabet $S_x^{[\mathbf{W}]} \subset \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ selected by unitary device words and certified by word–evaluated channels (Definitions 4.255, 4.257, 4.304 and 4.305 and Proposition 4.258). When off–block couplings are negligible, ordinary qubit kinematics and gate sets are recovered (Qubit Reduction and Registers; Proposition 4.300 and Corollaries 4.301 and 4.302); in the endogenous picture this is the regime where $\{0, 1\}$ remains effectively closed under admissible controls while $\{p, s\}$ acts as a quiet complement (Propositions 4.262 and 4.281).

New Avenues (Schematic)

1. *Geometry–aware control.* The fibre metric and even the effective background geometry are *expectation–value* objects induced by the Holo–State, not fixed backdrops (Sections 4.6 and 4.7.4 and eq. (4.85)). This turns the arena into a *programmable resource*: algorithms can co–design dynamics with induced information–geometry, e.g. shaping correlators to alter effective locality or to pull up long–range holosymmetric entanglers on demand. Holonomy in the s–channel becomes a native geometric observable and

control knob (wordwise loop operator and signal; Section 4.7.3, eqs. (4.122) and (4.123), and Proposition 4.295).

2. *Modal augmentation beyond qubits.* Parasymmetric and holosymmetric modes are *independent* directions of the Fourfold, not superpositions within a dyad (Proposition 4.281 and Section 1.2.7). Controlled mixing with $\{p, s\}$ therefore yields strictly non–qubit primitives—drives, measurements, dissipative gadgets—operating natively in the Fourfold. The qubit model is recovered when those mixings are suppressed; otherwise, Hadamard–compatible words and channels give structured access to the additional modal resources (Corollary 4.225 and Definition 4.255).

3. *Register formation over PQS networks.* When cross–fibre correlators between selected loci are perturbatively sparse, an n–site *effective* arena $(\mathbb{C}^2)^{\otimes n}$ emerges with controlled inter–modal couplings yielding entangling two–site gates (Registers; Section 4.1.2 and Corollary 4.302). This ties scalable digital architectures directly to correlator sparsity and transport stability criteria (Definition 4.248 and Corollary 4.251).

The relocation does not merely *reinterpret* qubits; it adds programmable *correlator geometry* and genuinely *Fourfold* controls. Precision comes from inequalities: small off–block words/channels and the holosymmetric compliance vector $\mathfrak{K} = (\Delta, \eta, \zeta)$ bound visibility/lensing drift, two–step *ps*/*sp* amplitudes, and diagonal pinching errors (Definition 4.178 and Proposition 4.181). Duhamel–type estimates then quantify leakage when mixings are weak but nonzero, giving principled tolerances for *hybrid* processors that exploit gentle $\{0, 1\} \leftrightarrow \{p, s\}$ couplings while retaining qubit–level programmability on protected Twofold sectors (Corollary 4.301 and Proposition 4.306).

Definition (Holomodal Quantum Computer). A holomodal quantum computer comprises: (i) a modal Hilbert bundle $\mathscr{H}_{\mathrm{mod}} \to \mathcal{X}_q$ with fibre metric *induced* by the Fourfold correlator (Section 4.6 and Definition 4.9); (ii) an admissible control algebra generated by Fourfold dialectic operators together with a metric–compatible connection acting on fibres, obeying the covariant Leibniz rule (Section 1.2.7, Lemma 4.270, and Definition 4.248); and (iii) fibre effect algebras (POVMs) including parasymmetric/holosymmetric channels, with dynamics and readout defined by expectations in the state–dependent arena (Section 4.7.4).

How it relates to qubit computing.

1. *Contains* the qubit model as a limit: when a Twofold sector is dynamically protected and cross–couplings are block–diagonalisable, the Qubit Reduction yields an isometric embedding $U : \mathbb{C}^2 \hookrightarrow \mathcal{H}_{\mathrm{mod}}(\mathfrak{B}_x)$; multi–site registers arise under correlator sparsity (Proposition 4.300 and Corollaries 4.301 and 4.302).

2. *Extends* it by natively using the full Fourfold: operations act directly on $\{|0\rangle, |1\rangle, |p\rangle, |s\rangle\}$, with controlled $\{p, s\}$ mixing enabling gates and measurements with no qubit analogue (Section 1.2.7 and Proposition 4.262).

3. *Programs the arena itself:* since the fibre metric and vacuum geometry are expectation–value objects, correlator engineering becomes a computational resource—co–designing dynamics with induced information–geometry and reading global structure via wordwise holonomy (Sections 4.7.3 and 4.7.4).

What to call it (taxonomically). A **holomodal (Fourfold) quantum computer**: a background–free, operator–algebraic model in which "qubits" appear as *derived* Twofold sectors when conditions warrant, yet computation can proceed in the full Fourfold with correlator–programmable resources. Qubit circuits are recovered in correlator–sparse regimes; beyond them, a broader gate/measurement set—tied to parasymmetry/holosymmetry, to wordwise transport, and to state–dependent geometry—becomes available within the same formalism (Definition 4.304, Propositions 4.181 and 4.306, Corollary 4.308, and Section 4.7.3).

4.8 The Intrinsic Coherence of the PQS Continuum

Within the Interdependent Quantum Reality (IQR) framework, *entanglement* is elevated from a contingent, subsystem–relative feature to a *structural property* of the Primordial Quantum Singularity (PQS) continuum itself. Rather than beginning with an *a priori* product Hilbert space and declaring non-separable vectors to be "entangled," IQR posits an *operator-theoretic* universality: the Quantum Holo-State $|\mathring{\mathbf{\Psi}}\rangle$ globally conditions the algebra of Fourfold modes so that relational observables, metrics, and coherences *emerge between* PQS loci $\mathfrak{B}_x, \mathfrak{B}_{x'}$ (Definition 1.27). In this sense, universal entanglement is ontological: it is the default coherence of the $\hat{\Omega}$–Continuum, from which phenomenological (subsystem) entanglement is recovered as a special, effective limit (cf. table 4.1). The underlying motivation follows the IQR dictum that observable structure (distance, time, geometry) crystallises *relationally* across distinct PQS loci rather than *within* any single locus.

4.8.1 The Operator-Theoretic Nature of Ontological PQS Entanglement

The inherent entanglement of primordial quantum singularities (Definition 1.25) and of the $\hat{\Omega}$–Continuum they constitute is exhibited by the following intrinsic properties.

1. Bilocal Fourfold coherence via $\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})$. The *Fourfold Inter-Modal Correlator* (Definition 4.9) furnishes a nonlocal, operator-valued relation between distinct PQS loci:

$$\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2] := \sum_{\alpha,\beta\in\{0,1,p,s\}} (c_{1\alpha})^* c_{2\beta} \left(\hat{\mathcal{F}}_\alpha(x)^\dagger \mathcal{U}(x \leftarrow x'; \gamma)\, \hat{\mathcal{F}}_\beta(x')\right), \tag{4.131}$$

where $\mathcal{U}(x \leftarrow x'; \gamma)$ is the parallel transporter that aligns local frames along a path $\gamma \subset \mathcal{X}_q$. The Holo-State expectation

$$\left\langle \mathring{\mathbf{\Psi}} \middle| \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2] \middle| \mathring{\mathbf{\Psi}} \right\rangle$$

quantifies an *algebraic overlap* (an "entanglement amplitude") between the modal data at $\mathfrak{B}_x$ and $\mathfrak{B}_{x'}$, independent of any classical embedding. This bilocal kernel is the primary manifestation of universal entanglement in the continuum.

2. Globally conditioned local structure (emergent fibre metric). The same correlator governs the *local* inner product on each modal fibre by the coincident-limit prescription

$$\langle \psi_1, \psi_2 \rangle_x := \lim_{x' \to x} \Big\langle \mathring{\mathbf{\Psi}} \,\Big|\, \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2] \,\Big|\, \mathring{\mathbf{\Psi}} \Big\rangle,$$

so that the fibre metric is *induced* from Fourfold expectations rather than postulated (Section 4.6). Positivity of the Holo-State functional ensures a well-defined Hilbert fibre after quotienting the null space and completing; thus correlator expectations *forge* the metric on which Fourfold modes are read.

3. Gauge-covariant comparison across the continuum. Relational claims between x' and x are made in a frame-consistent manner by the unitary transporter $\mathcal{U}(x \leftarrow x'; \gamma) \in \mathrm{U}(\mathcal{H}_{\mathrm{em}})$, ensuring that channel operators $\hat{G}_\Lambda^{\alpha\beta}(x, x') = \hat{\mathcal{F}}_\alpha^\dagger(x)\, \mathcal{U}(x \leftarrow x'; \gamma)\, \hat{\mathcal{F}}_\beta(x')$ are gauge-covariant. This underwrites the observer-independence of the entanglement amplitude defined above (cf. Lemma 4.270).

4. The continuum's triadic axes of relational geometry and time. Universal entanglement is not isolated from geometry; it is one axis of a triad of intrinsic relational *distances* on $\mathcal{X}_q$,

$$\vec{d}(x, x') = \Big(d_{\mathrm{corr}},\, d_{\mathrm{vac}},\, d_{\mathrm{spec}}\Big),$$

with the corresponding *axis-specific temporal parameters*

$$\vec{t}(x, x') = \Big(t_{\mathrm{corr}},\, t_{\mathrm{vac}},\, t_{\mathrm{spec}}\Big),$$

introduced as the *entanglement phase time*, *vacuum response time*, and *spectral shear time*, respectively (see Section 4.12). The three axes are *functionally orthogonal*, arising from algebraically distinct operator features of $\hat{G}_\Lambda$ and its spectral commutator: bilocal/off–diagonal structure for d_{corr} (paired with t_{corr}), diagonal/coincident variation for d_{vac} (paired with t_{vac}), and the commutator with the spectral generator $\hat{D}$ for d_{spec} (paired with t_{spec}). See Section 4.11.3 for the formal construction of the *Relational Distinguishability Fiber* and the assembly of these axes; the specialisation to *emergent proper* distances and times—the classical limit of ordinary experiences—is developed later in Section 5.3.8.

5. Ontological versus phenomenological entanglement. The foregoing properties define an *ontological* layer of entanglement—the default operator-level coherence of the $\hat{\Omega}$–Continuum—distinct from *phenomenological* (subsystem) entanglement in fixed tensor-product arenas. The latter reappears when the Fourfold correlator's Twofold block decouples (qubit reduction) and when cross-correlators across chosen fibres are perturbatively suppressed, yielding the usual $(\mathbb{C}^2)^{\otimes n}$ registers and their entangling gates (cf. Proposition 4.300 and Corollary 4.302). In this regime, the IQR account reduces to familiar quantum-information kinematics and dynamics.

Proposition 4.311 (Operational manifestations of ontological entanglement)**.** *In the IQR framework, the operator–theoretic (ontological) entanglement defined by the Holo-State* $|\mathring{\mathbf{\Psi}}\rangle$

and the Fourfold Inter-Modal Correlator $\hat{G}_\Lambda$ (Definitions 1.27 and 4.9) is a foundational *feature of the $\hat{\Omega}$–Continuum. It is generally not directly witnessable by finite-subsystem Bell tests, but it admits the following gauge-covariant* operational footprints*:*

(i) Expectation-induced geometry.
The chronogeometric tensor $\bar{\Sigma}_{\mu\nu}(x) = \langle\Omega_\Psi|\hat{\Sigma}_{\mu\nu}(x)|\Omega_\Psi\rangle$ (Definition 5.26) encodes global correlational structure; nontrivial dependence of $\bar{\Sigma}$ on the Holo-State implements geometric effects of ontological entanglement.

(ii) Emergent fibre metrics.
The local inner product on each modal fibre is induced *via the coincident-limit of $\hat{G}_\Lambda$ expectations (Section 4.6), so the very existence and positivity of the fibre Hilbert spaces reflect the continuum's entangled substrate.*

(iii) Holosymmetric interferometry.
Visibility functionals in holosymmetric interference experiments (e.g., Section 3.1 and Section 3.1.4) are Holo-State expectations of bilocal operators built from $\hat{G}_\Lambda$; their nontrivial values witness globally maintained operator histories rather than factorised subsystem coherence.

(iv) Effective two-level sectors and registers.
Stability of Twofold sectors and their inter-fibre registers—captured by the qubit reduction and registers results (Proposition 4.300 and Corollary 4.302)—are controlled by block structure and sparsity of $\hat{G}_\Lambda$; persistence of effective SU(2) *dynamics is thereby an operational signature of universal entanglement in the sparse-coupling regime.*

Justification (sketch). (i) follows immediately from the definition of $\bar{\Sigma}$ as a Holo-State expectation of geometric operators, whose inputs are conditioned by the same algebraic substrate governed by $\hat{G}_\Lambda$ (Definition 5.26). (ii) is the coincident-limit construction of the modal inner product from $\hat{G}_\Lambda$ with positivity and completion ensuring a genuine Hilbert fibre (Section 4.6). (iii) uses that holosymmetric visibility is defined as a gauge-covariant functional of bilocal Fourfold operators, hence sensitive to nonlocal coherence rather than factorised subsystems (Sections 3.1 and 3.1.4). (iv) is a direct consequence of the block-diagonalisable regime in which Twofold sectors are dynamically protected and cross-correlators are perturbatively suppressed, yielding effective SU(2) controls and entangling inter-fibre gates (Proposition 4.300 and Corollary 4.302). □

Ontological Emergence: Dependency Chain

The definitional dependencies can be summarised succinctly as

$$|\mathring{\boldsymbol{\Psi}}\rangle \xrightarrow[\text{Definition 4.9}]{\text{induces}} \hat{G}_\Lambda \xrightarrow[\text{Section 4.6}]{\text{coincident limit}} \langle\cdot,\cdot\rangle_x \xrightarrow[\text{Definition 5.26}]{\text{geometric operators}} \hat{\Sigma}_{\mu\nu}(x) \xrightarrow{\langle\cdot\rangle} \bar{\Sigma}_{\mu\nu}(x).$$

This emphasises that the emergent fibre metric and the expectation-level geometry are both grounded in Holo-State expectations of Fourfold operator structure, without implying a dynamical cascade.

Structural / Ontological PQS Entanglement	**Phenomenological / Conventional Entanglement**
An intrinsic property of the PQS continuum (Definition 4.617) conditioned by the Quantum Holo-State $\|\mathring{\mathbf{\Psi}}\rangle$ (Definition 1.27); it expresses global coherence of the $\hat{\Omega}$–Continuum.	A property of particular states within an *emergent* (often approximately factorisable) subsystem arena, e.g. registers $(\mathbb{C}^2)^{\otimes n}$ obtained in the sparse–coupling regime (Corollary 4.302).
Arises from nonlocal Fourfold operator relations (the inter–modal correlator $\hat{G}_\Lambda$) and *induces* local structures via Holo-State expectations—e.g. the modal fibre inner product by the coincident limit (Definition 4.9 and Section 4.6).	Characterised by non-separability of a state with respect to a chosen tensor product; canonical examples include Bell states and entangling two-qubit gates in the register model [287].
No PQS or local modal space is fundamental in isolation: fibre metrics and comparisons are *relational* (parallel transport $\mathcal{U}(x \leftarrow x'; \gamma)$; gauge-covariant pairings) and depend on the same correlator geometry (Lemma 4.270).	Subsystems are *defined* with respect to a fixed or effective factorisation; correlations are then assessed relative to that choice (entangled vs. product) within the given background Hilbert structure.
The qubit appears as a *derived*, dynamically protected Twofold sector when off-block couplings are block-diagonalisable; statistics and controls match an effective SU(2) (Proposition 4.300).	Qubits are *postulated* as $(\mathbb{C}^2, \langle\cdot,\cdot\rangle)$ objects with prescribed controls and measurements; universality follows from local SU(2) plus any entangling two-qubit gate [287].
Suppression (not literal absence) of specific correlator blocks yields classical-like product structure and trivialised mixing in the corresponding sectors; exact vanishing is a singular/ideal limit. In this regime, phenomenological entanglement notions are recovered (Corollaries 4.301 and 4.302).	Absence of entanglement simply means a product state in the chosen factorisation; it does not, by itself, constrain geometric or fibre-inducing structures.

Table 4.1: Distinction between *ontological* entanglement (a global, correlator-driven property of the PQS continuum) and *phenomenological* entanglement (non-separability relative to a chosen subsystem factorisation).

The Nature of Ontological Entanglement: From a Unified Whole to Apparent Parts

A foundational principle of the IQR framework is that the universe is a single, indivisible quantum system, intrinsically entangled by the Holo-State $|\mathring{\mathbf{\Psi}}\rangle$ acting on the Fourfold algebra at every PQS locus (Definitions 1.27 and 4.9). This raises a natural question: if reality is

globally entangled, why do we *experience* a world of apparently separate, classical objects governed by local laws?

Answer in IQR. Separation is not fundamental but *emergent*. What looks like the "failure" of entanglement is precisely the theory's mechanism for the appearance of classicality: when specific correlator structures are suppressed or effectively factorised, the relational substrate yields autonomous, approximately local sectors. The following conditions formalise this emergence.

Proposition 4.312 (Apparent separability: three mechanisms)**.** *Let $\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})$ be the Fourfold Inter-Modal Correlator and $\mathcal{U}(x \leftarrow x'; \gamma)$ the parallel transporter (Definition 4.9 and Lemma 4.270). Then:*

(a) Structural disconnection (vanishing bilocal coherence). *If for all test coefficients $c_{1\alpha}, c_{2\beta}$ and all admissible paths γ one has*

$$\left\langle \mathring{\mathbf{\Psi}} \middle| \hat{\mathcal{F}}_\alpha(x)^\dagger \, \mathcal{U}(x \leftarrow x'; \gamma) \, \hat{\mathcal{F}}_\beta(x') \middle| \mathring{\mathbf{\Psi}} \right\rangle = 0,$$

then no gauge-covariant bilocal coherence is present between $\mathfrak{B}_x$ and $\mathfrak{B}_{x'}$. Operationally this is a superselection*–type separation of sectors (a correlator-null block), i.e. an ontological decoupling at the operator level.*

(b) Effective factorisation (approximate product structure). *Suppose there is an emergent bipartition of the observable algebra into commuting subalgebras $\mathcal{A}_x$ and $\mathcal{A}_{x'}$ (relative to a chosen frame) and that the Holo-State admits a product approximation $\rho \approx \rho_x \otimes \rho_{x'}$ with trace distance $\|\rho - \rho_x \otimes \rho_{x'}\|_1 \leq \varepsilon$. Then for all $A \in \mathcal{A}_x$, $B \in \mathcal{A}_{x'}$,*

$$\left| \operatorname{Tr}[(\rho - \rho_x \otimes \rho_{x'}) \, A \otimes B] \right| \;\leq\; \varepsilon \, \|A\| \, \|B\|,$$

so bilocal correlators factorise up to $O(\varepsilon)$. This "for-all-practical-purposes" separability explains autonomous subsystems without contradicting global ontological entanglement.

(c) Decoherence (suppression of off-diagonal Fourfold blocks). *Under admissible open dynamics with generator $\mathcal{L}$ (e.g. a Lindbladian), off-diagonal blocks of the correlator decay, $\|(\hat{G}_\Lambda)_{\mathit{off}}(t)\| \leq e^{-\Gamma t} \|(\hat{G}_\Lambda)_{\mathit{off}}(0)\|$ for some rate $\Gamma > 0$, driving an* effective localisation *of correlations (cf. the emergence of classicality by environment-induced superselection [380]). In this regime the bilocal kernel behaves approximately local, and qubit/register kinematics reappear when the Twofold block is dynamically protected (Proposition 4.300 and Corollary 4.302).*

Justification (sketch). (a) is immediate from the definition of the bilocal channels and gauge-covariant transport (Definition 4.9 and Lemma 4.270). (b) follows from the Hölder–von Neumann inequality $|\operatorname{Tr}(XY)| \leq \|X\|_1 \|Y\|$ applied to $X = \rho - \rho_x \otimes \rho_{x'}$ and $Y = A \otimes B$. (c) is the standard decoherence mechanism: completely positive contractive dynamics suppress coherence between pointer sectors, here realised as decay of off-diagonal Fourfold blocks [380]. □

Operational footprints (global, not few-body tests). Ontological entanglement is *operator-theoretic*, not primarily a property of pre-given tensor factors. Its empirical content appears in:

- *Expectation-induced geometry:* the chronogeometric tensor $\bar{\Sigma}_{\mu\nu}(x) = \langle\Omega_\Psi|\hat{\Sigma}_{\mu\nu}(x)|\Omega_\Psi\rangle$ encodes correlator geometry (Definition 5.26).
- *Emergent inner products:* fibre metrics arise from coincident-limit expectations of $\hat{G}_\Lambda$ (Section 4.6).
- *Holosymmetric interferometry:* visibility functionals are Holo-State expectations of bilocal operators (cf. Sections 3.1 and 3.1.4).
- *Protected Twofold sectors and registers:* persistence of effective SU(2) dynamics and controllable entanglers in sparse-coupling regimes (Proposition 4.300 and Corollary 4.302).

Synthesis. In IQR the $\hat{\Omega}$–Continuum is *intrinsically entangled*. This is not a contingent feature of some specially prepared states but the *same* operator data that (i) *induce* modal fibre metrics and (ii) *determine* vacuum geometry via $\hat{\Sigma}_{\mu\nu}$. Apparent separations arise precisely under the mechanisms of Proposition 4.312, and in those regimes the familiar qubit/register formalism is recovered as an effective description.

4.8.2 Factorizability of the Quantum Holo-State and the Necessity of Ontological Entanglement

The Quantum Holo-State $|\overset{\circ}{\mathbf{\Psi}}\rangle$ (Definition 1.27) is the structural keystone of the IQR framework: it conditions the Fourfold operator algebra at each Primordial Quantum Singularity (PQS) (Definition 1.25) and, through its expectations, *induces* both local fibre metrics and expectation-level geometry. Clarifying what would follow *if* $|\overset{\circ}{\mathbf{\Psi}}\rangle$ were factorizable—and why, in IQR, it cannot be—is therefore essential.

I. What genuine factorization would entail

Consider a bipartition of the $\hat{\Omega}$–Continuum into sectors $\mathcal{X}_A$ and $\mathcal{X}_B$, with corresponding (frame-relative) commuting subalgebras $\mathcal{A}_A$ and $\mathcal{A}_B$. Suppose that

$$|\overset{\circ}{\mathbf{\Psi}}\rangle = |\psi_A\rangle \otimes |\psi_B\rangle, \qquad \omega^{\overset{\circ}{\mathbf{\Psi}}} = \omega^A \otimes \omega^B,$$

i.e. the universal state and its expectation functional factorize across the partition.

(1) Collapse of nonlocal Fourfold channels. Let $x \in \mathcal{X}_A$ and $x' \in \mathcal{X}_B$. For any test coefficients $c_{1\alpha}, c_{2\beta}$ and admissible path γ, the bilocal Fourfold channel (Definition 4.9)

$$\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2] = \sum_{\alpha,\beta} (c_{1\alpha})^* c_{2\beta}\, \hat{\mathcal{F}}_\alpha^\dagger(x)\, \mathcal{U}(x \leftarrow x'; \gamma)\, \hat{\mathcal{F}}_\beta(x')$$

satisfies

$$\Big\langle \mathring{\mathbf{\Psi}}\Big|\hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})[\psi_1^*,\psi_2]\Big|\mathring{\mathbf{\Psi}}\Big\rangle = \omega^A\Big(\hat{\mathcal{F}}_\alpha^\dagger(x)\Big)\,\omega^B\Big(\mathcal{U}\,\hat{\mathcal{F}}_\beta(x')\Big),$$

which vanishes for generic gauge choices and mode content unless the transporter $\mathcal{U}$ and the operators reduce to purely local (diagonal) contributions. Operationally, cross-sector bilocal coherence is null: the correlator block is effectively zero.

(2) Block-diagonal emergent inner products. By the coincident-limit construction of the fibre inner product (Section 4.6, cf. eq. (4.6)), vanishing cross-sector expectations force the modal pairing to become block-diagonal with respect to the partition. Hence inter-sector comparisons are ill-defined, and the Hilbert bundle decouples across $\mathcal{X}_A \sqcup \mathcal{X}_B$.

(3) Entanglement entropy across the cut vanishes. Writing $\rho_A = \mathrm{Tr}_B|\mathring{\mathbf{\Psi}}\rangle\langle\mathring{\mathbf{\Psi}}| = |\psi_A\rangle\langle\psi_A|$ gives $S(\rho_A) = 0$. Conventional (subsystem) entanglement across $\mathcal{X}_A \leftrightarrow \mathcal{X}_B$ is absent, mirroring the collapse of Fourfold bilocal structure.

(4) Geometric and temporal fragmentation in $\mathcal{X}_q$. Since expectation-level geometry is induced from Holo-State expectations (cf. $\bar{\Sigma}_{\mu\nu}$ in Definition 5.26), cross-sector components of $\bar{g}_{\mu\nu}$ (Definition 5.10) and of the temporal tensor are forced to vanish. Consequences include:

- a block-diagonal (possibly degenerate) $\bar{g}_{\mu\nu}$ across the cut, prohibiting coherent geometric relations between $\mathcal{X}_A$ and $\mathcal{X}_B$;
- loss or ill-definition of emergent temporal displacements across the cut (no well-defined axis-specific $t_{\mathrm{corr}}, t_{\mathrm{vac}}, t_{\mathrm{spec}}$ for cross-sector paths; cf. Section 4.12);
- fragmentation of causal structure (Section 5.4) into operator-disconnected components.

(5) Physical meaning: a classical/decohered limit. For $A \in \mathcal{A}_A$, $B \in \mathcal{A}_B$, $\langle\hat{\mathcal{O}}_A\hat{\mathcal{O}}_B\rangle = \langle\hat{\mathcal{O}}_A\rangle\langle\hat{\mathcal{O}}_B\rangle$, and holosymmetric interference between histories supported on disjoint sectors is suppressed (cf. Section 2.4). The partition behaves as independent classical-like domains. In summary,

> Exact factorization ⇒ vanishing cross–sector Fourfold channels
> ⇒ block–diagonal inner products & fragmented geometry.

Approximate factorization (FAPP). If $\rho \approx \rho_A \otimes \rho_B$ with trace distance $\|\rho - \rho_A \otimes \rho_B\|_1 \le \varepsilon$, then $|\,\mathrm{Tr}[(\rho - \rho_A \otimes \rho_B)A \otimes B]| \le \varepsilon\|A\|\,\|B\|$. Correlators factorize up to $O(\varepsilon)$, explaining *emergent* autonomy of subsystems without contradicting global ontological entanglement.

II. A theorematic constraint: $|\mathring{\mathbf{\Psi}}\rangle$ cannot factorize in IQR

Theorem 4.313 (Necessity of non-factorizability)**.** *Within the IQR framework, any genuine factorization of the Quantum Holo-State across a nontrivial partition of the $\hat{\Omega}$–Continuum is*

incompatible with the existence of (i) nontrivial gauge-covariant bilocal Fourfold channels, (ii) a coherent, nondegenerate fibre inner product at all loci, and (iii) a connected expectation-level chronogeometry. Hence $|\mathring{\mathbf{\Psi}}\rangle$ *is necessarily non-factorizable.*

Proof sketch. (i) By Definition 4.9 and gauge-covariant transport (Lemma 4.270), factorization forces the expectation of any cross-sector bilocal channel to split and, generically, to vanish, annihilating nonlocal coherence.

(ii) The local fibre pairing is defined by the coincident limit of Holo-State expectations of $\hat{G}_\Lambda$ (Section 4.6, cf. eq. (4.6)). Vanishing cross-sector expectations imply a block-diagonal (and across the cut, degenerate) inner product, contradicting bundle coherence and continuity demanded by the GNS construction (Definitions 1.34 and 4.287).

(iii) The operator-valued geometric fields $\hat{g}_{\mu\nu}$ and $\hat{t}_{\mu\nu}$ and their expectations $\bar{g}_{\mu\nu}, \bar{t}_{\mu\nu}$ (Definitions 5.1 and 5.26) depend on the same expectation functional. Factorization therefore enforces block structure in $\bar{\Sigma}_{\mu\nu}$, obstructing connected chronogeometry. Together these contradict the IQR postulates of a globally coherent, state-induced geometry; hence $|\mathring{\mathbf{\Psi}}\rangle$ cannot factorize. □

Information-theoretic corollary. If $|\mathring{\mathbf{\Psi}}\rangle$ factorized, the mutual information $I(\mathfrak{B}_x : \mathfrak{B}_{x'})$ would vanish across the partition for $x \in \mathcal{X}_A$, $x' \in \mathcal{X}_B$, contradicting the existence of holosymmetric interference and of induced inter-fibre metrics—both rely on nonzero Holo-State–mediated correlations.

Conclusion. In IQR, entanglement is not optional. It is the *structural* content of the same expectations that (a) *induce* fibre inner products and (b) *determine* expectation-level geometry. Exact factorization drives the theory into a degenerate, classically fragmented limit; the realized universe is therefore supported by a non-factorizable $|\mathring{\mathbf{\Psi}}\rangle$ that sustains connected chronogeometry and the full triad of relational distances and times.

II. Non-Factorizability of $|\mathring{\mathbf{\Psi}}\rangle$: a theorematic constraint of IQR

In a background–independent, relational quantum geometry, the Quantum Holo-State $|\mathring{\mathbf{\Psi}}\rangle$ cannot be genuinely factorizable across a nontrivial partition of the $\hat{\Omega}$–Continuum. This is not heuristic but a structural consequence of the IQR postulates.

Theorem 4.314 (Non-factorizability of the Holo-State)**.** *Let* $\mathcal{X}_q = \mathcal{X}_A \sqcup \mathcal{X}_B$ *be a nontrivial partition with frame–relative commuting subalgebras* $\mathcal{A}_A, \mathcal{A}_B$*. If* $|\mathring{\mathbf{\Psi}}\rangle = |\psi_A\rangle \otimes |\psi_B\rangle$ *(and hence* $\omega^{\mathring{\mathbf{\Psi}}} = \omega^A \otimes \omega^B$*), then the following fail simultaneously:*

(i) Nontrivial gauge–covariant bilocal channels. *For* $x \in \mathcal{X}_A$*,* $x' \in \mathcal{X}_B$*, expectations of the Fourfold inter–modal correlator* $\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})$ *(Definition 4.9) factorise and generically vanish under parallel transport* $\mathcal{U}(x \leftarrow x'; \gamma)$*, eliminating cross–sector bilocal coherence (cf. Lemma 4.270).*

(ii) Coherent, nondegenerate fibre inner products. *The modal inner product at x is the coincident–limit of Holo-State expectations of $\hat{G}_\Lambda$ (Section 4.6). Vanishing cross–sector expectations force a block–diagonal pairing, destroying bundle coherence across $\mathcal{X}_A \sqcup \mathcal{X}_B$.*

(iii) Connected expectation–level chronogeometry and axis–specific times. *The chronogeometric tensor $\bar{\Sigma}_{\mu\nu}(x)$ (Definition 5.26) inherits the same expectation functional; factorization imposes block structure and obstructs connected chronogeometry, with axis–specific temporal parameters $t_{\mathrm{corr}}, t_{\mathrm{vac}}, t_{\mathrm{spec}}$ undefined or trivial for cross–sector paths (cf. Section 4.12, and the classical specialisation in Section 5.3.8).*

(iv) Holosymmetric interference and induced registers. *Global visibilities (holosymmetric interferometry; Sections 3.1 and 3.1.4) and the emergence of effective registers (Corollary 4.302) require nonzero cross–sector correlator blocks; these are absent under factorization.*

Consequently, exact factorization contradicts the IQR construction; the Holo-State is necessarily non–factorizable.

Proof sketch. (i) Using $\omega^{\mathring{\Psi}} = \omega^A \otimes \omega^B$ and the bilocal channel definition, expectations split as products of one–sector functionals applied to transported operators; for generic mode content and gauge, the cross–sector contribution vanishes (no bilocal coherence), cf. Definition 4.9 and Lemma 4.270. (ii) The inner product is obtained as a coincident–limit of those expectations (Section 4.6); vanishing cross–blocks produce a block–diagonal (across the cut, degenerate) pairing, incompatible with a coherent Hilbert bundle (cf. Definitions 1.34, 4.239 and 4.287). (iii) The same expectation functional defines $\bar{\Sigma}_{\mu\nu}$ (Definition 5.26); factorization forces block structure in $\bar{\Sigma}$, obstructing connected chronogeometry and trivialising axis–specific temporal parameters across the cut (cf. Sections 4.12 and 5.3.8). (iv) Holosymmetric interference functionals and register formation require nonzero cross–sector blocks; these disappear under factorization (Sections 3.1 and 3.1.4 and Corollary 4.302). □

Approximate (FAPP) factorization is allowed. If $\rho \approx \rho_A \otimes \rho_B$ with $\|\rho - \rho_A \otimes \rho_B\|_1 \le \varepsilon$, then for $A \in \mathcal{A}_A$, $B \in \mathcal{A}_B$,

$$\left| \mathrm{Tr}[(\rho - \rho_A \otimes \rho_B)\, A \otimes B] \right| \le \varepsilon\, \|A\|\, \|B\|,$$

so bilocal expectations factorise up to $O(\varepsilon)$. This explains emergent autonomy of subsystems without violating global non–factorizability.

Explanation (FAPP factorization). "FAPP" ("for all practical purposes") means *approximate* factorization at the level of observable consequences, not an exact product state. Let ρ be the actual (possibly correlated) state and $\rho_A \otimes \rho_B$ a product approximation on commuting subalgebras $\mathcal{A}_A, \mathcal{A}_B$ associated to a bipartition (e.g. a frame–relative split of PQS loci; cf. Lemma 4.270). If the *trace distance* is small, $\|\rho - \rho_A \otimes \rho_B\|_1 \le \varepsilon$, then Hölder for Schatten norms and $\|A \otimes B\| = \|A\|\, \|B\|$ give the bound above, hence

$$\mathrm{Tr}(\rho\, A \otimes B) = \mathrm{Tr}(\rho_A A)\ \mathrm{Tr}(\rho_B B)\ \pm\ O(\varepsilon\, \|A\|\, \|B\|).$$

Operationally, if experimental resolution is coarser than the $O(\varepsilon)$ corrections, subsystems behave *autonomously*: cross–correlators are indistinguishable from products. This is compatible with the IQR requirement of *global* non–factorizability (Theorem 4.314) and naturally arises under decoherence, where off–diagonal Fourfold blocks are exponentially suppressed (cf. Proposition 4.312(c)), driving ε small on accessible timescales. Importantly, FAPP factorization is a statement about *expectations*; it does **not** assert that correlator geometry or induced fibre metrics vanish (Section 4.6 and Definition 5.26).

Corollary 4.315 (Information–theoretic obstruction to factorization)**.** *Let $\mathcal{X}_q = \mathcal{X}_A \sqcup \mathcal{X}_B$ be a nontrivial partition with commuting subalgebras $\mathcal{A}_A, \mathcal{A}_B$. If the global state factorizes, $\rho = \rho_A \otimes \rho_B$, then the mutual information*

$$I(A\!:\!B)_\rho \;=\; S(\rho_A) + S(\rho_B) - S(\rho)$$

vanishes, i.e. $I(A\!:\!B)_\rho = 0$. Equivalently, $I(A\!:\!B)_\rho = S\big(\rho \,\big\|\, \rho_A \otimes \rho_B\big) \geq 0$ with equality iff $\rho = \rho_A \otimes \rho_B$ (standard property of relative entropy; see, e.g., [287, 241]). Consequently, any globally coherent IQR configuration exhibiting nonzero holosymmetric visibility across the cut (cf. Sections 3.1 and 3.1.4) or a nondegenerate inter–fibre inner product across the cut (nonvanishing bilocal Fourfold expectations; cf. Section 4.6 and Definition 4.9) must satisfy $I(A\!:\!B)_\rho > 0$ and therefore cannot be exactly factorizable.

Proof sketch. If $\rho = \rho_A \otimes \rho_B$, then $S(\rho) = S(\rho_A) + S(\rho_B)$, so $I(A:B)_\rho = 0$. Conversely, $I(A\!:\!B)_\rho = S(\rho \,\|\, \rho_A \otimes \rho_B)$ equals zero iff $\rho = \rho_A \otimes \rho_B$. Nonzero holosymmetric visibility and nonvanishing cross–sector Fourfold expectations constitute observable correlations across the cut; thus $I(A\!:\!B)_\rho > 0$, which rules out exact factorization. This aligns with Theorem 4.314. □

Exact factorization $\Rightarrow$ no bilocal channels
$\Rightarrow$ block–diagonal fibres & fragmented chronogeometry
$\Rightarrow$ ruled out in IQR.

Corollary 4.316 (Fourfold non-separability enforced by the Holo-State)**.** *Let $\mathcal{A}^{(4)}$ denote the Fourfold Dialectic Algebra of Symmetry (cf. Definition 1.16), generated by the modal channels $\{\hat{\mathcal{F}}_0, \hat{\mathcal{F}}_1, \hat{\mathcal{F}}_p, \hat{\mathcal{F}}_s\}$ (Symmetry, Antisymmetry, Parasymmetry, Holosymmetry; cf. Section 1.2.7), and let π_Ψ be its GNS representation in the Holo-State $|\mathring{\mathbf{\Psi}}\rangle$ (Definitions 1.34 and 4.287). If $|\mathring{\mathbf{\Psi}}\rangle$ is non-factorizable (Theorem Theorem 4.314), then in the represented algebra $\mathcal{M} := \pi_\Psi(\mathcal{A}^{(4)})''$:*

(i) No sector-splitting central projectors

There is no nontrivial $Z \in \mathcal{M} \cap \mathcal{M}'$ that separates $\{0,1\}$ from $\{p,s\}$ or any other Fourfold split. Equivalently, no global superselection splits the Fourfold into tensor- or direct-product components aligned with the modal grading. Otherwise $|\mathring{\mathbf{\Psi}}\rangle$ would factorize across that split, contradicting Theorem 4.314.

(ii) Cross-modal correlators are generically nonzero
Bilocal Fourfold channels $\hat{G}^{\alpha\beta}_{\Lambda}(x,x')$ have nonvanishing Holo-State expectations for cross-modal pairs (e.g. $\alpha \in \{0,1\}$, $\beta \in \{p,s\}$), after gauge-covariant transport (Definition 4.9 and Lemma 4.270). Thus parasymmetric/holosymmetric mixing is a structural *feature, not an artifact.*

(iii) Inner-product and geometry couple all Fourfold channels
The coincident-limit construction of the fibre inner product and the expectation-level chronogeometry (Section 4.6 and Definition 5.26) necessarily involve the off-block Fourfold expectations; a globally coherent metric/chronogeometry requires the coupled Fourfold, not any single channel in isolation.

(iv) Reductions are contingent, not fundamental
Any Twofold (qubit) reduction to $\{0,1\}$ is an effective *block-diagonal regime achieved by dynamical protection and suppressed off-block couplings (Proposition 4.300 and Corollaries 4.301 and 4.302); it does not reflect a decomposition of $\mathcal{A}^{(4)}$, only a controlled truncation of its represented dynamics.*

Sketch. If a central projector split the Fourfold (i), expectations of cross-modal bilocals would vanish identically, implying factorization of $|\overset{\circ}{\mathbf{\Psi}}\rangle$ across that split, contradicting Theorem 4.314. Nonvanishing cross-modal expectations (ii) follow from the same theorem together with the definition of the inter-modal correlator and gauge transport (Definition 4.9 and Lemma 4.270). Items (iii)–(iv) use that the fibre inner product and chronogeometry are induced from those expectations (Section 4.6 and Definition 5.26); when off-block terms are *suppressed* (not absent), the compressed dynamics yields the Twofold/registry effective theory (Proposition 4.300 and Corollary 4.302). □

4.8.3 Wordwise Non-Factorizability of the Holo-State

Setup 4.317 (Partition, commuting subalgebras, and words)**.** Let $\mathcal{X}_q = \mathcal{X}_A \sqcup \mathcal{X}_B$ be a nontrivial partition of PQS loci, and let $\mathcal{A}_A, \mathcal{A}_B$ be the frame–relative commuting subalgebras generated by word–evaluated, typed letters at loci in $\mathcal{X}_A$ and $\mathcal{X}_B$, respectively (Definitions 4.255 and 4.257 and Corollary 4.64). Fix a unitary device word $\mathbf{W}$ and control χ (suppressing χ when clear). For $x \in \mathcal{X}_A$, $x' \in \mathcal{X}_B$ and a path $\gamma : x' \to x$ with transporter $\mathcal{U}(x \leftarrow x';\gamma)$ (Definition 4.248 and Section 4.1.4), define the *wordwise bilocal kernel*

$$\boxed{K^{[\mathbf{W}]}_{\alpha\beta}(x,x';\gamma) \;:=\; \Big\langle \overset{\circ}{\mathbf{\Psi}} \Big| \Big(\hat{\mathcal{F}}^{[\mathbf{W}]}_{\alpha}(x)\Big)^{\dagger} \mathcal{U}(x \leftarrow x';\gamma)\, \hat{\mathcal{F}}^{[\mathbf{W}]}_{\beta}(x') \Big| \overset{\circ}{\mathbf{\Psi}} \Big\rangle,} \tag{4.132}$$

cf. Definition 4.9 and eq. (4.66).

Lemma 4.318 (Exact factorization ⇒ rank-one bilocal kernel)**.** *Assume the Holo–State factorizes across the partition, $|\overset{\circ}{\mathbf{\Psi}}\rangle = |\psi_A\rangle \otimes |\psi_B\rangle$ so that $\omega^{\overset{\circ}{\mathbf{\Psi}}} = \omega^A \otimes \omega^B$. Then, for all $x \in \mathcal{X}_A$, $x' \in \mathcal{X}_B$ and γ,*

$$\boxed{K^{[\mathbf{W}]}_{\alpha\beta}(x,x';\gamma) = \underbrace{\omega^A\Big(\Big(\hat{\mathcal{F}}^{[\mathbf{W}]}_{\alpha}(x)\Big)^{\dagger}\Big)}_{=:a_\alpha(x)^*} \underbrace{\omega^B\Big(\mathcal{U}(x \leftarrow x';\gamma)\, \hat{\mathcal{F}}^{[\mathbf{W}]}_{\beta}(x')\Big)}_{=:b_\beta(x',\gamma)},} \tag{4.133}$$

so the modal kernel has rank at most 1 *as a matrix in* (α,β). *Consequently the wordwise entanglement bilinear* $\mathsf{E}^{[\mathbf{W}]}_{x,x'}$ *(Definition 4.332 and eq. (4.144)) factors for all test states.*

Proof. Under factorization, expectations split on commuting subalgebras. The operator $\big(\hat{\mathcal{F}}^{[\mathbf{W}]}_{\alpha}(x)\big)^{\dagger}$ lies in $\mathcal{A}_A$ and $\mathcal{U}\hat{\mathcal{F}}^{[\mathbf{W}]}_{\beta}(x')$ in $\mathcal{A}_B$, hence $\omega^{\mathring{\Psi}}(AB)=\omega^A(A)\,\omega^B(B)$, giving (4.133). Rank–one form implies $\mathsf{E}^{[\mathbf{W}]}_{x,x'}(\psi_1,\psi_2)=\varphi_x(\psi_1)^*\varphi_{x'}(\psi_2)$ with $\varphi_x(\psi_1)=\sum_\alpha a_\alpha\, c^{[\mathbf{W}]}_{1\alpha}(x)$ and $\varphi_{x'}(\psi_2)=\sum_\beta b_\beta\, c^{[\mathbf{W}]}_{2\beta}(x')$. □

Lemma 4.319 (Wordwise collapse of cross–sector interference)**.** *In the setting of Lemma 4.318, the connected wordwise interference* $\mathsf{I}^{[\mathbf{W}]}_{\alpha\beta}(x;\psi)$ *(eq. (4.143)) across the cut vanishes identically after centering by local means,* $\hat{\mathcal{F}}^{\circ}_{\alpha}:=\hat{\mathcal{F}}_{\alpha}-\omega(\hat{\mathcal{F}}_{\alpha})\mathbf{1}$. *Hence any nontrivial wordwise prismatic superposition (Definition 4.331) cannot bridge* $\mathcal{X}_A$ *to* $\mathcal{X}_B$ *under exact factorization.*

Full proof. Step 0 (Setting and centering). Let $\mathcal{X}_q=\mathcal{X}_A\sqcup\mathcal{X}_B$ as in Setup 4.317 and assume exact factorization $|\mathring{\Psi}\rangle=|\psi_A\rangle\otimes|\psi_B\rangle$ with $\omega^{\mathring{\Psi}}=\omega^A\otimes\omega^B$ (Lemma 4.318). For $x\in\mathcal{X}_A$, $x'\in\mathcal{X}_B$ and a path $\gamma: x'\to x$ with transporter $\mathcal{U}(x\leftarrow x';\gamma)$ (Definition 4.248), set the *transported letter*

$$\widetilde{\mathcal{F}}^{[\mathbf{W}]}_{\beta}(x';\gamma)\;:=\;\mathcal{U}(x\leftarrow x';\gamma)\,\hat{\mathcal{F}}^{[\mathbf{W}]}_{\beta}(x').$$

Define centered (connected) letters sectorwise

$$\begin{aligned}\big(\hat{\mathcal{F}}^{[\mathbf{W}]}_{\alpha}(x)\big)^{\circ}&:=\hat{\mathcal{F}}^{[\mathbf{W}]}_{\alpha}(x)-\omega^A\big(\hat{\mathcal{F}}^{[\mathbf{W}]}_{\alpha}(x)\big)\,\mathbf{1},\\ \big(\widetilde{\mathcal{F}}^{[\mathbf{W}]}_{\beta}(x';\gamma)\big)^{\circ}&:=\widetilde{\mathcal{F}}^{[\mathbf{W}]}_{\beta}(x';\gamma)-\omega^B\big(\widetilde{\mathcal{F}}^{[\mathbf{W}]}_{\beta}(x';\gamma)\big)\,\mathbf{1}.\end{aligned}\tag{4.134}$$

Step 1 (Connected bilocal kernel across the cut). Consider the connected two–point expectation

$$\mathcal{C}^{[\mathbf{W}]}_{\alpha\beta}(x,x';\gamma):=\;\omega^{\mathring{\Psi}}\Big(\big(\hat{\mathcal{F}}^{[\mathbf{W}]}_{\alpha}(x)\big)^{\circ\dagger}\big(\widetilde{\mathcal{F}}^{[\mathbf{W}]}_{\beta}(x';\gamma)\big)^{\circ}\Big).\tag{4.135}$$

Because $\big(\hat{\mathcal{F}}^{[\mathbf{W}]}_{\alpha}(x)\big)^{\circ\dagger}\in\mathcal{A}_A$ and $\big(\widetilde{\mathcal{F}}^{[\mathbf{W}]}_{\beta}(x';\gamma)\big)^{\circ}\in\mathcal{A}_B$ (commuting subalgebras; Setup 4.317), factorization yields

$$\mathcal{C}^{[\mathbf{W}]}_{\alpha\beta}(x,x';\gamma)=\omega^A\Big(\big(\hat{\mathcal{F}}^{[\mathbf{W}]}_{\alpha}(x)\big)^{\circ\dagger}\Big)\,\omega^B\Big(\big(\widetilde{\mathcal{F}}^{[\mathbf{W}]}_{\beta}(x';\gamma)\big)^{\circ}\Big)=0\cdot 0=0,$$

since each centered one–point function vanishes by construction.

Step 2 (From connected kernel to interference functional). The wordwise interference functional $\mathsf{I}^{[\mathbf{W}]}_{\alpha\beta}$ (eq. (4.143)) is the *local* specialization $x'=x$ with identity transport and weighted by modal coefficients:

$$\mathsf{I}^{[\mathbf{W}]}_{\alpha\beta}(x;\psi)=\omega^{\mathring{\Psi}}\Big(\big(\hat{\mathcal{F}}^{[\mathbf{W}]}_{\alpha}(x)\big)^{\dagger}\hat{\mathcal{F}}^{[\mathbf{W}]}_{\beta}(x)\Big)\,\big(c^{[\mathbf{W}]}_{\alpha}(x)\big)^{*}\,c^{[\mathbf{W}]}_{\beta}(x).$$

To assess *cross–cut* coherence, replace the local kernel by the bilocal, connected kernel (4.135) and coefficients on the respective loci (cf. the bilinear entanglement form, Definition 4.332 and eq. (4.144)). The result vanishes by Step 1:

$$\big(\mathsf{I}^{[\mathbf{W}]}_{\alpha\beta}\big)_{\text{across cut}}\;\propto\;\mathcal{C}^{[\mathbf{W}]}_{\alpha\beta}(x,x';\gamma)\,\big(c^{[\mathbf{W}]}_{1,\alpha}(x)\big)^{*}\,c^{[\mathbf{W}]}_{2,\beta}(x')\;=\;0.$$

Step 3 (Conclusion for prismatic superposition). By Definition 4.331, a nontrivial wordwise prismatic superposition across a set S requires at least one pair $\alpha \neq \beta$ with nonzero interference functional. Since the connected cross–sector contribution vanishes identically under exact factorization, no coherent superposition can *bridge* $\mathcal{X}_A$ to $\mathcal{X}_B$. Equivalently, any apparent cross–cut "superposition'' reduces to a mixture, consistent with the rank–one (factorized) bilocal kernel of Lemma 4.318 and the factorization of the entanglement bilinear (Definition 4.332).

Gauge/word invariance. All steps are invariant under endpoint gauge and word changes, because centering uses sectorwise expectations and the transformations act by unitary conjugations that preserve ω^A, ω^B (Propositions 4.54 and 4.284 and Corollary 4.64).

This establishes the claim. □

Proposition 4.320 (Block structure of induced pairings and geometry (wordwise))**.** *Let the fibre inner product and expectation–level chronogeometry be induced from wordwise correlators (Section 4.6 and Definition 5.26). Under exact factorization, all cross–sector contributions to the coincident–limit pairings and to the chronogeometric tensor vanish; the Hilbert bundle and the expectation-level geometry decompose as a block direct sum across $\mathcal{X}_A \sqcup \mathcal{X}_B$.*

Proof idea. The inner product at a locus is obtained as a coincident limit of (4.132). Cross–sector limits employ kernels of the form (4.133); their connected part vanishes, yielding block diagonality. The same expectation functional determines $\bar{\Sigma}_{\mu\nu}$ (Definition 5.26), so cross–sector components are zero.

Full proof. Step 0 (Setting). Partition $\mathcal{X}_q = \mathcal{X}_A \sqcup \mathcal{X}_B$ with commuting frame–relative subalgebras $\mathcal{A}_A, \mathcal{A}_B$ (Setup 4.317). Assume exact factorization of the Holo–State, $|\overset{\circ}{\boldsymbol{\Psi}}\rangle = |\psi_A\rangle \otimes |\psi_B\rangle$ so that $\omega^{\overset{\circ}{\boldsymbol{\Psi}}} = \omega^A \otimes \omega^B$. For $x \in \mathcal{X}_A$, $x' \in \mathcal{X}_B$ and any path $\gamma : x' \to x$ with transporter $\mathcal{U}(x \leftarrow x'; \gamma)$ (Definition 4.248), consider the wordwise bilocal kernel

$$K^{[\mathbf{W}]}_{\alpha\beta}(x, x'; \gamma) = \left\langle \overset{\circ}{\boldsymbol{\Psi}} \,\middle|\, \big(\hat{\mathcal{F}}^{[\mathbf{W}]}_{\alpha}(x)\big)^{\dagger}\, \mathcal{U}(x \leftarrow x'; \gamma)\, \hat{\mathcal{F}}^{[\mathbf{W}]}_{\beta}(x') \,\middle|\, \overset{\circ}{\boldsymbol{\Psi}} \right\rangle \qquad (\textit{eq. } (4.132)).$$

Step 1 (Rank–one structure across the cut). By factorization and commutativity of $\mathcal{A}_A$ with $\mathcal{A}_B$,

$$K^{[\mathbf{W}]}_{\alpha\beta}(x, x'; \gamma) = \underbrace{\omega^A\Big(\big(\hat{\mathcal{F}}^{[\mathbf{W}]}_{\alpha}(x)\big)^{\dagger}\Big)}_{=:a_\alpha(x)^*}\, \underbrace{\omega^B\Big(\mathcal{U}(x \leftarrow x'; \gamma)\, \hat{\mathcal{F}}^{[\mathbf{W}]}_{\beta}(x')\Big)}_{=:b_\beta(x', \gamma)},$$

so the modal matrix is rank ≤ 1 (Lemma 4.318, eq. (4.133)). If we center letters sectorwise,

$$\big(\hat{\mathcal{F}}^{[\mathbf{W}]}_{\alpha}(x)\big)^{\circ} := \hat{\mathcal{F}}^{[\mathbf{W}]}_{\alpha}(x) - \omega^A(\hat{\mathcal{F}}^{[\mathbf{W}]}_{\alpha}(x))\, \mathbf{1}, \quad \big(\widetilde{\mathcal{F}}^{[\mathbf{W}]}_{\beta}(x'; \gamma)\big)^{\circ} := \mathcal{U}\, \hat{\mathcal{F}}^{[\mathbf{W}]}_{\beta}(x') - \omega^B(\mathcal{U}\, \hat{\mathcal{F}}^{[\mathbf{W}]}_{\beta}(x'))\, \mathbf{1},$$

then the connected cross–sector kernel vanishes identically:

$$\omega^{\overset{\circ}{\boldsymbol{\Psi}}}\Big(\big(\hat{\mathcal{F}}^{[\mathbf{W}]}_{\alpha}(x)\big)^{\circ\,\dagger}\, \big(\widetilde{\mathcal{F}}^{[\mathbf{W}]}_{\beta}(x'; \gamma)\big)^{\circ}\Big) = 0,$$

cf. Lemma 4.319.

Step 2 (Induced inner product: coincident limit and block–diagonality). The fibre inner product at x is defined as a coincident limit of such kernels (exogenous or wordwise, the constructions agree up to endpoint gauge; see Section 4.6): for local coefficient vectors $c, d \in \mathbb{C}^4$,

$$\Big\langle \psi, \phi \Big\rangle_x = \lim_{x' \to x} \sum_{\alpha,\beta} c_\alpha^* \, d_\beta \, K_{\alpha\beta}^{[\mathbf{W}]}(x, x'; \gamma), \qquad |\psi\rangle = \sum_\alpha c_\alpha |\alpha\rangle, \quad |\phi\rangle = \sum_\beta d_\beta |\beta\rangle. \tag{4.136}$$

Take the decomposition of the kernel into its centered (connected) and mean parts. The connected cross–sector part is zero by Step 1; the mean part factorizes into sectorwise numbers that do not depend on the opposite state coefficients. Therefore the cross–sector contribution to (4.136) vanishes in the coincident limit, and the pairing reduces to a sum of *intra–sector* terms. Equivalently, in any local frame adapted to the partition the Gram matrix of $\langle \cdot, \cdot \rangle_x$ is block diagonal with respect to $\mathcal{X}_A \sqcup \mathcal{X}_B$.

Step 3 (Expectation–level chronogeometry). Let $\bar{\Sigma}_{\mu\nu}(x) = \omega^{\mathring{\Psi}}\big(\hat{\Sigma}_{\mu\nu}(x)\big)$ be the chronogeometric tensor defined from the same expectation functional (Definition 5.26). Its construction (via composite two–point objects and appropriate contractions in the coincident limit; cf. Definition 5.26 and Section 4.6) employs the same bilocal kernels. Repeating the centering/-factorization argument shows that any cross–sector component of $\bar{\Sigma}_{\mu\nu}$ vanishes: connected cross–sector contributions are zero, while the mean parts factor into independent sectorwise numbers that do not produce cross terms upon coincidence. Hence $\bar{\Sigma}_{\mu\nu}$ is block diagonal across the partition.

Step 4 (Decomposition of bundle and geometry). Block–diagonality of the fibre pairing at each x implies that the modal bundle splits as an orthogonal Whitney sum of subbundles supported on $\mathcal{X}_A$ and $\mathcal{X}_B$:

$$\mathscr{H}_{\mathrm{mod}} \;\cong\; \mathscr{H}_{\mathrm{mod}}|_{\mathcal{X}_A} \;\oplus\; \mathscr{H}_{\mathrm{mod}}|_{\mathcal{X}_B},$$

with connection data reducing accordingly (no cross–sector parallel coupling). The expectation–level chronogeometry likewise decomposes: the metric/temporal tensors are block diagonal and define disconnected components on $\mathcal{X}_A$ and $\mathcal{X}_B$. Gauge and word changes act by endpoint/fibrewise unitary conjugations and do not affect the vanishing of connected cross–sector expectations (Propositions 4.54 and 4.284 and Corollary 4.64), so the conclusion is invariant.

Therefore, under exact factorization, cross–sector contributions to both the induced inner product and the chronogeometric tensor vanish, and the theory decomposes as a block direct sum across $\mathcal{X}_A \sqcup \mathcal{X}_B$. □

Theorem 4.321 (Wordwise necessity of non-factorizability)**.** *Assume the IQR postulates of: (i) nontrivial, gauge–covariant bilocal Fourfold channels (Definition 4.9), (ii) a coherent, nondegenerate fibre inner product at all loci (Section 4.6), and (iii) connected expectation-level chronogeometry (Definition 5.26). Then the Holo–State $|\mathring{\Psi}\rangle$ cannot factorize across any nontrivial partition of $\mathcal{X}_q$.*

Proof idea. Suppose $|\mathring{\Psi}\rangle = |\psi_A\rangle \otimes |\psi_B\rangle$. By Lemma 4.318, bilocal kernels across the cut are rank-one and therefore yield no nontrivial wordwise entanglement; by Lemma 4.319 no

connected cross-sector interference survives. This contradicts (i). By Proposition 4.320, the induced inner product and chronogeometry block–decompose, violating (ii) and (iii). Hence the assumption is false.

Full proof. We proceed by contradiction. Let $\mathcal{X}_q = \mathcal{X}_A \sqcup \mathcal{X}_B$ be a nontrivial partition with commuting, frame–relative subalgebras $\mathcal{A}_A, \mathcal{A}_B$ as in Setup 4.317, and suppose

$$|\overset{\circ}{\mathbf{\Psi}}\rangle = |\psi_A\rangle \otimes |\psi_B\rangle\,, \qquad \omega^{\overset{\circ}{\mathbf{\Psi}}} = \omega^A \otimes \omega^B. \tag{4.137}$$

Step 1 (Bilocal kernel becomes rank–one ⇒ no nontrivial channels). For $x \in \mathcal{X}_A$, $x' \in \mathcal{X}_B$ and any path $\gamma : x' \to x$ with transporter $\mathcal{U}(x \leftarrow x'; \gamma)$ (Definition 4.248), the wordwise bilocal kernel

$$K^{[\mathbf{W}]}_{\alpha\beta}(x, x'; \gamma) = \left\langle \overset{\circ}{\mathbf{\Psi}} \,\middle|\, \big(\hat{\mathcal{F}}^{[\mathbf{W}]}_\alpha(x)\big)^\dagger \mathcal{U}(x \leftarrow x'; \gamma)\, \hat{\mathcal{F}}^{[\mathbf{W}]}_\beta(x') \,\middle|\, \overset{\circ}{\mathbf{\Psi}} \right\rangle \qquad (\textit{eq. (4.132)})$$

factorizes by (4.137) into a product of one–side functionals:

$$K^{[\mathbf{W}]}_{\alpha\beta}(x, x'; \gamma) = \omega^A\Big(\big(\hat{\mathcal{F}}^{[\mathbf{W}]}_\alpha(x)\big)^\dagger\Big)\, \omega^B\Big(\mathcal{U}(x \leftarrow x'; \gamma)\, \hat{\mathcal{F}}^{[\mathbf{W}]}_\beta(x')\Big), \tag{4.138}$$

i.e. a rank–one matrix in the modal indices (α, β) (Lemma 4.318). After centering sector-wise, $(\hat{\mathcal{F}}^{[\mathbf{W}]}_\alpha)^\circ := \hat{\mathcal{F}}^{[\mathbf{W}]}_\alpha - \omega(\hat{\mathcal{F}}^{[\mathbf{W}]}_\alpha)\mathbf{1}$, the connected cross–sector kernel vanishes identically (Lemma 4.319). Hence there is no nontrivial, gauge–covariant bilocal Fourfold channel across the cut in the sense of Definition 4.9. This contradicts Postulate (i).

Step 2 (Coincident–limit pairing block–diagonalizes ⇒ loss of cross–sector coherence). The fibre inner product at x is obtained from coincident limits of such kernels (exogenous/wordwise constructions agree up to endpoint gauge; Section 4.6):

$$\langle \psi, \phi \rangle_x = \lim_{x' \to x} \sum_{\alpha,\beta} c^*_\alpha d_\beta\, K^{[\mathbf{W}]}_{\alpha\beta}(x, x'; \gamma) \quad \text{for} \quad |\psi\rangle = \sum_\alpha c_\alpha |\alpha\rangle, \quad |\phi\rangle = \sum_\beta d_\beta |\beta\rangle. \tag{4.139}$$

With (4.138) and the vanishing of the connected cross–sector part (Lemma 4.319), the cross–sector contribution to (4.139) disappears in the limit, leaving only intra–sector terms. Thus the Gram matrix of $\langle \cdot, \cdot \rangle_x$ is block–diagonal in a frame adapted to $\mathcal{X}_A \sqcup \mathcal{X}_B$ and the modal bundle splits orthogonally. Equivalently, the bundle fails to provide coherent cross–sector comparison, i.e. the induced pairing does not *glue* across the cut. This is precisely the block–decomposition stated in Proposition 4.320 and violates the "coherent" part of Postulate (ii) (nondegeneracy can hold on each block, but coherence across the partition is lost).

Step 3 (Expectation–level chronogeometry fragments). The chronogeometric tensor $\bar{\Sigma}_{\mu\nu}(x) = \omega^{\overset{\circ}{\mathbf{\Psi}}}(\hat{\Sigma}_{\mu\nu}(x))$ is built from the same bilocal correlators in a coincident–limit construction (Definition 5.26). By the same centering/factorization argument as in Step 2, all cross–sector components vanish and $\bar{\Sigma}_{\mu\nu}$ becomes block–diagonal across $\mathcal{X}_A \sqcup \mathcal{X}_B$ (Proposition 4.320). Therefore the expectation–level chronogeometry decomposes into disconnected pieces, contradicting Postulate (iii).

Step 4 (Holonomy witness is quenched under factorization). Independently of Steps 2–3, small–loop wordwise holonomy in the s–channel provides a local, gauge–invariant witness of connected chronogeometry:

$$\hat{H}^{(s)[\mathbf{W}]}_{\ell_\varepsilon}(x) = \Big(\hat{\mathcal{F}}^{[\mathbf{W}]}_s(x)\Big)^\dagger \Big(\mathbf{1}_4 - \varepsilon^2\,\Phi_x(\mathcal{F}) + O(\varepsilon^3)\Big)\,\hat{\mathcal{F}}^{[\mathbf{W}]}_s(x), \tag{4.140}$$
$$h^{(s)[\mathbf{W}]}_{\ell_\varepsilon}(x) = h^{(s)[\mathbf{W}]}_{\text{flat}}(x) - \varepsilon^2\,\omega\Big((\hat{\mathcal{F}}^{[\cdot]}_s)^\dagger \Phi_x(\mathcal{F})\hat{\mathcal{F}}^{[\cdot]}_s\Big) + O(\varepsilon^3).$$

(Corollary 4.298 and eqs. (4.98) and (4.99)). Under (4.137), the cross–sector contribution in the expectation reduces to a product of sector means (rank–one), so the genuinely nonlocal term proportional to $\Phi_x(\mathcal{F})$ is quenched across the cut. Thus even infinitesimal holonomy readouts cannot register a cross–sector twist, again contradicting Postulate (iii) whenever curvature or nontrivial topology is present.

Step 5 (Gauge/word invariance of the contradiction). All preceding steps rely on identities of expectations under sectorwise centering and use only the commuting structure of $\mathcal{A}_A, \mathcal{A}_B$; endpoint gauge and word changes act by unitary conjugation and preserve the factorization arguments (Propositions 4.54 and 4.284 and Corollary 4.64). Hence the contradictions are not artefacts of a particular gauge/word choice.

Since (4.137) simultaneously violates Postulates (i)–(iii), our assumption is false. Therefore $|\mathring{\mathbf{\Psi}}\rangle$ cannot factorize across any nontrivial partition of $\mathcal{X}_q$. □

Remark 4.322 (Holonomy and small-loop witnesses)**.** If $\mathcal{F} \neq 0$ near a cut, the wordwise small–loop sensitivity (Corollary 4.298 and eqs. (4.98) and (4.99)) produces an s–channel signal linear in the oriented area: a strictly local, gauge-invariant witness of connected chronogeometry. Exact factorization would force such cross–sector loop signals to reduce to rank-one (or centered zero) contributions, contradicting the existence of nontrivial holonomy readouts.

Proposition 4.323 (FAPP wordwise factorization)**.** *Let ρ be the actual state and $\rho_A \otimes \rho_B$ a product approximation on $\mathcal{A}_A \otimes \mathcal{A}_B$ with $\|\rho - \rho_A \otimes \rho_B\|_1 \le \varepsilon$. Then for any wordwise bilocal $O_A \in \mathcal{A}_A$, $O_B \in \mathcal{A}_B$,*

$$\boxed{\Big|\,\mathrm{Tr}\Big[\rho\, O_A \otimes O_B\Big] - \mathrm{Tr}(\rho_A O_A)\;\mathrm{Tr}(\rho_B O_B)\Big| \;\le\; \varepsilon\,\|O_A\|\,\|O_B\|,} \tag{4.141}$$

so wordwise kernels and holonomy signals factorize up to $O(\varepsilon)$. Thus emergent autonomy of subsystems is compatible with global non-factorizability (Theorem 4.321).

Proof idea. The key step is the noncommutative Hölder (duality) inequality for Schatten norms: for any trace–class operator $T \in \mathsf{S}_1$ and bounded operator $X \in \mathsf{S}_\infty$,

$$\Big|\,\mathrm{Tr}(TX)\Big| \;\le\; \|T\|_1\,\|X\|_\infty,$$

with the usual generalisation $\|AB\|_r \le \|A\|_p \|B\|_q$ for $1/p+1/q = 1/r$ (see, e.g., [356, 253, 381]). Setting $T = \rho - \rho_A \otimes \rho_B$ and $X = O_A \otimes O_B$ gives

$$\Big|\,\mathrm{Tr}[(\rho - \rho_A \otimes \rho_B)(O_A \otimes O_B)]\Big| \;\le\; \|\rho - \rho_A \otimes \rho_B\|_1\,\|O_A \otimes O_B\|_\infty.$$

Using the projective property of the operator norm on simple tensors, $\|O_A \otimes O_B\| = \|O_A\|\,\|O_B\|$ (e.g. [356]), yields the stated bound (4.141). The same estimate applies termwise to any finite linear combination of simple tensors, hence to the wordwise bilocal kernel (4.132) and to the loop operator (4.122) after word evaluation, by linearity and the triangle inequality. Conceptually, Hölder–Schatten duality is crucial because it converts a *global* closeness in trace distance into a *uniform* $O(\varepsilon)$ control of all cross–sector expectations, establishing "FAPP" factorization at the level of observable consequences while remaining gauge/word invariant.

Full proof. Step 0 (Notation and norms). Let $\Delta := \rho - \rho_A \otimes \rho_B$. By hypothesis $\|\Delta\|_1 \le \varepsilon$. We use Schatten duality (Hölder for Schatten norms): for any trace–class T and bounded X, $|\operatorname{Tr}(TX)| \le \|T\|_1\,\|X\|$. Moreover $\|O_A \otimes O_B\| = \|O_A\|\,\|O_B\|$ for operator norm.

Step 1 (Basic inequality). Compute

$$\operatorname{Tr}\Big[\rho\, O_A \otimes O_B\Big] - \operatorname{Tr}(\rho_A O_A)\,\operatorname{Tr}(\rho_B O_B) = \operatorname{Tr}\Big[\,(\rho - \rho_A \otimes \rho_B)\,(O_A \otimes O_B)\,\Big] = \operatorname{Tr}\Big[\Delta\,(O_A \otimes O_B)\Big].$$

Apply Hölder:

$$\Big|\operatorname{Tr}\Big[\Delta\,(O_A \otimes O_B)\Big]\Big| \;\le\; \|\Delta\|_1\,\|O_A \otimes O_B\| \;=\; \varepsilon\,\|O_A\|\,\|O_B\|,$$

which is precisely (4.141).

Step 2 (Application to wordwise bilocal kernels). For the wordwise bilocal kernel (eq. (4.132)),

$$K^{[\mathbf{W}]}_{\alpha\beta}(x,x';\gamma) = \Big\langle \overset{\circ}{\mathbf{\Psi}} \Big|\, \Big(\hat{\mathcal{F}}^{[\mathbf{W}]}_{\alpha}(x)\Big)^{\dagger}\, \mathcal{U}(x \leftarrow x';\gamma)\, \hat{\mathcal{F}}^{[\mathbf{W}]}_{\beta}(x') \,\Big|\, \overset{\circ}{\mathbf{\Psi}} \Big\rangle,$$

identify $O_A = (\hat{\mathcal{F}}^{[\mathbf{W}]}_{\alpha}(x))^{\dagger} \in \mathcal{A}_A$ and $O_B = \mathcal{U}(x \leftarrow x';\gamma)\,\hat{\mathcal{F}}^{[\mathbf{W}]}_{\beta}(x') \in \mathcal{A}_B$ ($\mathcal{A}_A, \mathcal{A}_B$ commute by construction; cf. Setup 4.317 and Definition 4.248). Then (4.141) gives

$$\Big|\, K^{[\mathbf{W}]}_{\alpha\beta}(x,x';\gamma) - \operatorname{Tr}(\rho_A O_A)\,\operatorname{Tr}(\rho_B O_B) \,\Big| \;\le\; \varepsilon\,\|O_A\|\,\|O_B\|. \tag{4.142}$$

Thus the kernel *factorizes up to* $O(\varepsilon)$ into sectorwise expectations, uniformly controlled by the operator norms. In particular, all bilinear functionals built from the kernel (e.g. the wordwise entanglement bilinear in Definition 4.332 and eq. (4.144)) inherit the same $O(\varepsilon)$ control.

Step 3 (Application to holonomy signals). For the wordwise holosymmetric loop operator at x,

$$\hat{H}^{(s)[\mathbf{W}]}_{\ell}(x) = \Big(\hat{\mathcal{F}}^{[\mathbf{W}]}_{s}(x)\Big)^{\dagger}\, \mathcal{U}_{\ell}(x)\, \hat{\mathcal{F}}^{[\mathbf{W}]}_{s}(x) \qquad (\textit{eq. (4.122)}),$$

two situations occur:

(a) If $\hat{H}^{(s)[\mathbf{W}]}_{\ell}(x)$ is supported entirely in $\mathcal{A}_A$, take $O_A = \hat{H}^{(s)[\mathbf{W}]}_{\ell}(x)$, $O_B = \mathbf{1}$. Then

$$\Big|\operatorname{Tr}\Big(\rho\,\hat{H}^{(s)[\mathbf{W}]}_{\ell}(x)\Big) - \operatorname{Tr}\Big(\rho_A\,\hat{H}^{(s)[\mathbf{W}]}_{\ell}(x)\Big)\Big| \le \varepsilon\,\|\hat{H}^{(s)[\mathbf{W}]}_{\ell}(x)\|.$$

(b) If the loop word splits across the cut (e.g. a device word whose typed letters act in both sectors along ℓ), write it as a finite sum of simple tensors $\sum_j A_j \otimes B_j$ (by

linearity of the evaluation functor; cf. Definition 4.255 and Proposition 4.284). Apply (4.141) termwise and sum:

$$\left|\operatorname{Tr}\left(\rho\,\hat{H}_\ell^{(s)[\mathbf{W}]}\right)-\operatorname{Tr}\left(\rho_A\otimes\rho_B\,\hat{H}_\ell^{(s)[\mathbf{W}]}\right)\right| \;\le\; \varepsilon\sum_j \|A_j\|\,\|B_j\| \;\le\; \varepsilon\,\|\hat{H}_\ell^{(s)[\mathbf{W}]}\|_\gamma,$$

where $\|\cdot\|_\gamma$ is any projective cross–norm dominating $\sum_j \|A_j\|\,\|B_j\|$; in particular the operator norm bound $\varepsilon\,\|\hat{H}_\ell^{(s)[\mathbf{W}]}\|$ suffices.

Hence holonomy expectations "factorize" to product-state predictions up to $O(\varepsilon)$, consistent with the small–loop expansion in Corollary 4.298 and eq. (4.99).

Step 4 (Connected/centered quantities and robustness). If one prefers connected (centered) observables, define $O_A^\circ := O_A - \operatorname{Tr}(\rho_A O_A)\mathbf{1}$, $O_B^\circ := O_B - \operatorname{Tr}(\rho_B O_B)\mathbf{1}$. Then

$$\left|\operatorname{Tr}\left[\rho\,O_A^\circ\otimes O_B^\circ\right]\right| = \left|\operatorname{Tr}\left[\Delta\,O_A^\circ\otimes O_B^\circ\right]\right| \le \varepsilon\,\|O_A\|\,\|O_B\|,$$

so connected cross–sector correlations are $O(\varepsilon)$. If $\rho_A = \operatorname{Tr}_B\rho$ and $\rho_B = \operatorname{Tr}_A\rho$ are chosen (optional), contractivity of the partial trace in trace norm implies $\|\operatorname{Tr}_B\rho - \rho_A\|_1 \le \varepsilon$, so the same bounds hold with the true marginals, strengthening the interpretation.

Step 5 (Gauge/word invariance). Endpoint gauge and word changes act by fibrewise unitary conjugations that preserve operator norms and expectations in (4.141) (Propositions 4.54 and 4.284 and Corollary 4.64); the bounds are therefore invariant.

Combining Steps 1–3 proves (4.141) and its consequences for wordwise kernels and holonomy signals. This establishes that *approximate* (FAPP) factorization of expectations coexists with global non–factorizability (Theorem 4.321). □

Corollary 4.324 (No Fourfold sector–splitting central projectors (wordwise GNS))**.** *Let $\mathcal{A}^{(4)}$ be the Fourfold word algebra and π_Ψ its GNS representation in $|\mathring{\mathbf{\Psi}}\rangle$ (Definitions 1.34 and 4.287). If $|\mathring{\mathbf{\Psi}}\rangle$ is non-factorizable (Theorem Theorem 4.321), then the von Neumann algebra $\mathcal{M} = \pi_\Psi(\mathcal{A}^{(4)})''$ admits no nontrivial $Z \in \mathcal{M}\cap\mathcal{M}'$ that splits $\{0,1\}$ from $\{p,s\}$. Otherwise the state would factorize across that split, contradicting the theorem.*

Summary. In the endogenous alphabets-and-words formalism, exact factorization forces bilocal wordwise kernels to rank-one (Lemma 4.318), collapses connected interference and prismatic superposition across the cut (Lemma 4.319), and block–diagonalizes the induced inner product and chronogeometry (Proposition 4.320). This conflicts with the IQR postulates of nontrivial bilocal channels, coherent fibres, and connected chronogeometry, hence the Holo-State must be non-factorizable (Theorem 4.321). Approximate (FAPP) factorization remains valid with quantitative bounds (Proposition 4.323).

Non-factorizability as a structural necessity. In the IQR construction the universal Quantum Holo-State $|\mathring{\mathbf{\Psi}}\rangle$ simultaneously (i) conditions the Fourfold operator content at each PQS and (ii) *induces* both the fibre inner products and the expectation–level geometry via its correlators (Definitions 1.25, 4.9 and 5.26 and Section 4.6). The endogenous (alphabets–and–words) argument shows that an *exact* product across a nontrivial partition forces contradictions with

those very inductions: one cannot retain nontrivial bilocal channels, a coherent nondegenerate fibre metric at every locus, and a connected chronogeometry if $|\mathring{\mathbf{\Psi}}\rangle$ factors. Formally, this is captured by the wordwise non–factorizability theorem (Theorem 4.321).

Operational meaning of the "rank–one kernel." If $|\mathring{\mathbf{\Psi}}\rangle$ factored, the word–evaluated bilocal kernel linking modes at x and x' would split as a product of *one–side* functionals, i.e. become rank–one in its modal indices (Lemma 4.318). Operationally this means every cross–region correlation is just "what A does" times "what B does''—no shared phase information, no genuinely joint structure. In the endogenous language, the entanglement bilinear $\mathsf{E}^{[\mathbf{W}]}_{x,x'}$ (Definition 4.332 and eq. (4.144)) *factors* for all tests, so ontological entanglement across the cut is impossible.

Geometric fragmentation as a consequence. The inner products on fibres are obtained as coincident limits of those same kernels, and the expectation–level chronogeometry uses the same expectation functional (Section 4.6 and Definition 5.26). Once the cross–sector pieces of the kernel are rank–one (and their connected parts vanish), the induced pairing becomes block–diagonal and the geometry decomposes along the partition. The result is a bundle and metric that *fragment* into operator–disconnected components, contradicting the postulated connected chronogeometry (Proposition 4.320 and Definition 4.248).

Holonomy as a local witness (and why it would go flat). Small–loop holonomy in the holosymmetric s–channel is designed to register curvature or topology at the expectation level: the wordwise loop operator and its signal shift linearly with the loop's oriented area to lowest order (eqs. (4.99) and (4.122) and Corollary 4.298). Exact factorization collapses the cross–sector contribution to a rank–one (product) form, so the *genuinely nonlocal* twist written into the signal disappears. Put plainly: the very dial meant to detect global fibre gluing would read "no twist'' across the cut.

No coherent superposition across the cut. Superposition in this framework is diagnosed by a nonzero wordwise interference functional between at least two modal directions (Definition 4.331 and eq. (4.143)). Under exact factorization, the connected cross–sector interference vanishes identically after centring by local means, so coherent superpositions cannot bridge the partition; only mixtures remain (Lemma 4.319). This aligns with the collapse of the bilocal kernel and the loss of parasymmetric, phase–sensitive effects across the cut.

Entanglement is structural; qubits are effective. Because the same expectations that power interference also *define* the fibre metric and chronogeometry, entanglement is not an add–on resource but part of the load–bearing structure of the theory (Section 4.8.5 and Definition 4.332). Twofold (qubit–like) behaviour still emerges when a wordwise Twofold sector is dynamically protected and off–block mixing is negligible, yielding an isometric embedding and standard qubit kinematics; registers arise when cross–correlators between selected fibres are perturbatively suppressed (Propositions 4.300 and 4.335 and Corollary 4.302). But that is an *effective* reduction inside a genuinely Fourfold world.

Why subsystems often look independent (FAPP factorization). Approximate products in trace distance make all bilocal expectations *behave* like products up to a quantitative tolerance (Proposition 4.323). This explains the everyday autonomy of subsystems—decoherence and engineering can suppress off–block terms so that, within resolution, cross–cut signals are indistinguishable from products—without asserting that the global Holo–State truly factors. In short, only *approximate* separability is compatible with the same correlations that both knit geometry and underwrite nonclassical phenomena.

4.8.4 Fourfold Prismatic Superposition and Entanglement

Definition 4.325 (Prismatic Superposition at a Locus)**.** Fix a locus $\mathfrak{B}_x$. Let the modal fibre be $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)} = \mathrm{span}\{|0\rangle_x, |1\rangle_x, |p\rangle_x, |s\rangle_x\}$, with inner product induced by the coincident limit of the Fourfold correlator as in Section 4.6. Write the correlator channels as $\hat{\mathcal{F}}_\alpha(x)$ so that $\hat{G}_\Lambda^{\alpha\beta}(x,x') = \left\langle \hat{\mathcal{F}}_\alpha(x)^\dagger \mathcal{U}_{x\to x'} \hat{\mathcal{F}}_\beta(x') \right\rangle$. Given a nonzero vector $|\psi\rangle_x = \sum_{\alpha\in\{0,1,p,s\}} c_\alpha |\alpha\rangle_x$, we say that $|\psi\rangle_x$ is a *(nontrivial) prismatic superposition across* $S \subseteq \{0,1,p,s\}$ if $|S| \geq 2$, its projection onto $\mathrm{span}\{|\alpha\rangle_x : \alpha \in S\}$ has at least two nonzero modal components, and there exists $\alpha \neq \beta$ in S with nonvanishing interference functional

$$\mathsf{I}_{\alpha\beta}(x\,;\psi) \;:=\; \left\langle \mathring{\mathbf{\Psi}} \,\middle|\, \hat{\mathcal{F}}_\alpha(x)^\dagger \hat{\mathcal{F}}_\beta(x) \,\middle|\, \mathring{\mathbf{\Psi}} \right\rangle (c_\alpha)^* c_\beta \;\neq\; 0.$$

Thus "superposition" is the *S-facet* (coherent coexistence) realised on the prism's modes.

Definition 4.326 (Ontological Entanglement Between Loci)**.** For loci $\mathfrak{B}_x, \mathfrak{B}_{x'}$ with test states $|\psi_1\rangle_x = \sum_\alpha c_{1\alpha}|\alpha\rangle_x$ and $|\psi_2\rangle_{x'} = \sum_\beta c_{2\beta}|\beta\rangle_{x'}$, define the bilinear form

$$\mathsf{E}_{x,x'}(\psi_1,\psi_2) \;:=\; \left\langle \mathring{\mathbf{\Psi}} \middle| \hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})[\psi_1^*,\psi_2] \middle| \mathring{\mathbf{\Psi}} \right\rangle,$$

with parallel transport as in eq. (4.131). We say the Holo-State exhibits *entanglement* between x and x' iff $\mathsf{E}_{x,x'}$ does *not* factor as $\varphi_x(\psi_1)^*\varphi_{x'}(\psi_2)$ for any linear functionals $\varphi_x, \varphi_{x'}$ on the respective modal fibres. Equivalently, the kernel $(\langle \hat{\mathcal{F}}_\alpha(x)^\dagger \mathcal{U}\, \hat{\mathcal{F}}_\beta(x')\rangle)_{\alpha,\beta}$ has rank > 1 with respect to modal indices when evaluated in $|\mathring{\mathbf{\Psi}}\rangle$.

Proposition 4.327 (Compatibility with the Fourfold Symmetry Algebra (Parasymmetric tests))**.** *On the Twofold slice* $\mathrm{span}\{|0\rangle_x, |1\rangle_x\}$*, let* $P_{\mathrm{para}}, P_{\mathrm{antipara}}$ *denote the parasymmetric and anti-parasymmetric effects (with* $P_{\mathrm{para}} + P_{\mathrm{antipara}} = P_{(0,1)}$*). Then for any* $|\psi\rangle_x = c_0|0\rangle_x + c_1|1\rangle_x$*,*

$$\left\langle P_{\mathrm{para}} \right\rangle_\psi = \tfrac{1}{2}\Big(\langle \hat{\mathcal{F}}_0^\dagger \hat{\mathcal{F}}_0 \rangle\, |c_0|^2 + \langle \hat{\mathcal{F}}_1^\dagger \hat{\mathcal{F}}_1 \rangle\, |c_1|^2 + 2\,\mathrm{Re}\, \mathsf{I}_{01}(x;\psi) \Big)\,.$$

Hence $|\psi\rangle_x$ *is a prismatic superposition across* $\{0,1\}$ *iff the parasymmetric/anti-parasymmetric pair yields a nontrivial, phase-sensitive split on* $|\psi\rangle_x$*; the superposition functional* I_{01} *is exactly the interference term in the Born valuation of* P_{para}*.*

Proposition 4.328 (Reduction to Standard Hilbert-Space Notions in the Twofold Regime)**.** *If the duality-collapse regime holds at (each of)* x *(and* x'*) in the sense of Definition 4.87, then there exist isometries* $U_x : \mathbb{C}^2 \hookrightarrow \mathcal{H}_{\mathrm{mod}\,(0,1)}^{(\mathfrak{B}_x)}$ *(and similarly at* x'*) such that:*

(i) $|\psi\rangle_x$ *is a prismatic superposition across* $\{0,1\}$ *iff* $U_x^\dagger|\psi\rangle_x$ *is a nontrivial linear combination in* $\mathbb{C}^2$*;*

(ii) entanglement between (x,x') *in Definition 4.326 holds iff the induced two–qubit state on* $\mathbb{C}^2\otimes\mathbb{C}^2$ *is entangled in the standard sense.*

Sketch. *The inner product on the Twofold fibre is induced by the restricted correlator; Definition 4.87 and Proposition 4.90 give the isometric identifications. Factorisation/nonfactorisation of* $\mathsf{E}_{x,x'}$ *reduces to product/nonproduct states after compression by* $U_x^\dagger\otimes U_{x'}^\dagger$*.*

Remark 4.329 (Basis/Transport Invariance)**.** The predicates in Definitions 4.325 and 4.326 are invariant under admissible reindexings and under the duality involution (cf. Proposition 1.8), and they are unaffected by parallel-transport gauge choices: changes $\hat{\mathcal{F}}_\alpha\mapsto e^{\mathrm{i}\phi_\alpha}\hat{\mathcal{F}}_\alpha$ or unitary changes of $\mathcal{U}$ that preserve the correlator class leave $\mathsf{I}_{\alpha\beta}$ and the factorisation test for $\mathsf{E}_{x,x'}$ unchanged.

Lemma 4.330 (Holosymmetric Dominance)**.** *Assume the holosymmetric effect* $E_{\mathrm{Holo}}^{(\mathbf{a}\to\mathbf{b})}$ *of eq. (3.36), with* $\mathbf{U}_{\mathrm{holo}}$ *constructed from the Fourfold channel algebra and lifted path unitaries (cf. Definition 2.102 and Section 2.3.2). Then:*

(i) (Local) *If, for some fibre preparation* $|\psi\rangle_x$*, the control readout is* phase–sensitive *(changes under a control-basis rotation with registers fixed), then there exist modal indices* $\alpha\neq\beta$ *with* $\mathsf{I}_{\alpha\beta}(x;\psi)\neq 0$ *as in Definition 4.325.*

(ii) (Bilocal) *If a holosymmetric bipartite test between* (x,x') *signals nonseparability, then the bilinear form* $\mathsf{E}_{x,x'}$ *of Definition 4.326 does not factor.*

Sketch. Write $v_H(\rho_{\mathrm{RR}})=\mathrm{Tr}\big(E_{\mathrm{Holo}}^{(\mathbf{a}\to\mathbf{b})}\rho_{\mathrm{RR}}\big)$ with $E_{\mathrm{Holo}}^{(\mathbf{a}\to\mathbf{b})}$ from eq. (3.36). In the canonical interferometer the control signal depends on the overlap operator $K:=U_G^\dagger U_N$ of the two lifted global evolutions acting on the same register input. A phase–sensitive change under a control Hadamard implies K has nonzero off–diagonal matrix elements in the Fourfold modal basis; since K is built from the channel operators $\hat{\mathcal{F}}_\alpha$, some $\langle\hat{\mathcal{F}}_\alpha^\dagger\hat{\mathcal{F}}_\beta\rangle\neq 0$ with $\alpha\neq\beta$, yielding $\mathsf{I}_{\alpha\beta}(x;\psi)\neq 0$ (i). For (ii), nonseparability of the holosymmetric readout forces nonzero matrix elements of the bilocal kernel $\langle\,\hat{\mathcal{F}}_\alpha(x)^\dagger\mathcal{U}(x\leftarrow x';\gamma)\hat{\mathcal{F}}_\beta(x')\,\rangle$ (eq. (4.131)); if $\mathsf{E}_{x,x'}$ factored as $\varphi_x^*\varphi_{x'}$, the control signal would be product-valued and insensitive to such cross–terms, contradicting the hypothesis. □

4.8.5 Wordwise Prismatic Superposition and Entanglement

Definition 4.331 (Wordwise prismatic superposition at a locus)**.** Fix a locus $\mathfrak{B}_x$, a unitary device word $\mathbf{W}$, and control χ. Let $\mathcal{B}_x^{[\mathbf{W},\chi]}=\{|0\rangle_x^{[\mathbf{W},\chi]},|1\rangle_x^{[\mathbf{W},\chi]},|p\rangle_x^{[\mathbf{W},\chi]},|s\rangle_x^{[\mathbf{W},\chi]}\}$ be the word–rotated orthonormal frame (Definition 4.257, Proposition 4.258, and Remark 4.286). For a nonzero state

$$|\psi\rangle_x^{[\mathbf{W}]}=\sum_{\alpha\in\{0,1,p,s\}}c_\alpha^{[\mathbf{W}]}(x)\,|\alpha\rangle_x^{[\mathbf{W},\chi]},$$

define the *wordwise interference functional* for $\alpha\neq\beta$ by

$$\boxed{\mathsf{I}_{\alpha\beta}^{[\mathbf{W}]}(x;\psi)\;:=\;\Big\langle\mathring{\mathbf{\Psi}}\,\Big|\,\Big(\hat{\mathcal{F}}_\alpha^{[\mathbf{W};\chi\leftarrow\chi]}(x)\Big)^\dagger\hat{\mathcal{F}}_\beta^{[\mathbf{W};\chi\leftarrow\chi]}(x)\,\Big|\,\mathring{\mathbf{\Psi}}\Big\rangle\,\Big(c_\alpha^{[\mathbf{W}]}(x)\Big)^*\,c_\beta^{[\mathbf{W}]}(x),}\tag{4.143}$$

where the word–dressed fields are as in Definitions 4.166 and 4.255 and eq. (4.66). Given a subset $S \subseteq \{0,1,p,s\}$ with $|S| \geq 2$, the state $|\psi\rangle_x^{[\mathbf{W}]}$ is a *(nontrivial) wordwise prismatic superposition across* S iff its projection onto $\mathrm{span}\{|\alpha\rangle_x^{[\mathbf{W},\chi]} : \alpha \in S\}$ has at least two nonzero modal coefficients and there exist $\alpha \neq \beta$ in S with $\mathsf{I}_{\alpha\beta}^{[\mathbf{W}]}(x;\psi) \neq 0$.

Definition 4.332 (Wordwise ontological entanglement between loci)**.** Fix loci $\mathfrak{B}_x, \mathfrak{B}_{x'}$, a word $\mathbf{W}$ and control χ. Let $|\psi_1\rangle_x^{[\mathbf{W}]} = \sum_\alpha c_{1\alpha}^{[\mathbf{W}]} |\alpha\rangle_x^{[\mathbf{W},\chi]}$ and $|\psi_2\rangle_{x'}^{[\mathbf{W}]} = \sum_\beta c_{2\beta}^{[\mathbf{W}]} |\beta\rangle_{x'}^{[\mathbf{W},\chi]}$. For a chosen path $\gamma : x' \to x$ with transporter $\mathcal{U}(x \leftarrow x';\gamma) = \mathcal{P}\exp\!\big(-\int_\gamma \mathcal{A}\big)$ (Definition 4.248 and Section 4.1.4), define the bilinear form

$$\boxed{\mathsf{E}_{x,x'}^{[\mathbf{W}]}(\psi_1,\psi_2) := \sum_{\alpha,\beta} \left\langle \mathring{\mathbf{\Psi}} \,\middle|\, \Big(\hat{\mathcal{F}}_\alpha^{[\mathbf{W};\chi\leftarrow\chi]}(x)\Big)^\dagger \mathcal{U}(x\leftarrow x';\gamma)\, \hat{\mathcal{F}}_\beta^{[\mathbf{W};\chi\leftarrow\chi]}(x') \,\middle|\, \mathring{\mathbf{\Psi}} \right\rangle \big(c_{1\alpha}^{[\mathbf{W}]}\big)^*\, c_{2\beta}^{[\mathbf{W}]}.} \tag{4.144}$$

The Holo–State exhibits *wordwise entanglement* between x and x' iff $\mathsf{E}_{x,x'}^{[\mathbf{W}]}$ does *not* factor as $\varphi_x(\psi_1)^*\,\varphi_{x'}(\psi_2)$ for any linear functionals $\varphi_x, \varphi_{x'}$ on the respective modal fibres. Equivalently, the kernel matrix $\big(\langle(\hat{\mathcal{F}}_\alpha^{[\cdot]}(x))^\dagger \mathcal{U}\, \hat{\mathcal{F}}_\beta^{[\cdot]}(x')\rangle\big)_{\alpha,\beta}$ has modal rank > 1 in $|\mathring{\mathbf{\Psi}}\rangle$ (Section 4.1.2 and eq. (4.66)).

Proposition 4.333 (Word/gauge invariance)**.** *Under endpoint gauge $g \in \mathrm{U}(4)$, $\mathcal{U} \mapsto g(x)\mathcal{U}g(x')^\dagger$ and $\hat{\mathcal{F}}_\alpha^{[\mathbf{W}]} \mapsto g\,\hat{\mathcal{F}}_\alpha^{[\mathbf{W}]}$ (Proposition 4.54 and Corollary 4.64). Under a change of word $\mathbf{W} \to \mathbf{W}'$, $\hat{\mathcal{F}}_\alpha^{[\mathbf{W}']}(x) = U_{[\mathbf{W}',\mathbf{W}]}(x)\,\hat{\mathcal{F}}_\alpha^{[\mathbf{W}]}(x)$ with a fibre unitary $U_{[\mathbf{W}',\mathbf{W}]}$ (Proposition 4.284). Hence the predicates in Definitions 4.331 and 4.332 are invariant: $\mathsf{I}_{\alpha\beta}^{[\mathbf{W}]}$ and the factorisation property of $\mathsf{E}_{x,x'}^{[\mathbf{W}]}$ do not depend on gauge or word choice.*

Full proof. Step 1 (Gauge invariance of the basic kernels). Consider the wordwise single–locus kernel $M_{\alpha\beta}^{[\mathbf{W}]}(x) := \omega\big((\hat{\mathcal{F}}_\alpha^{[\mathbf{W}]}(x))^\dagger \hat{\mathcal{F}}_\beta^{[\mathbf{W}]}(x)\big)$ and the bilocal kernel

$$K_{\alpha\beta}^{[\mathbf{W}]}(x,x';\gamma) = \omega\Big((\hat{\mathcal{F}}_\alpha^{[\mathbf{W}]}(x))^\dagger\, \mathcal{U}(x\leftarrow x';\gamma)\, \hat{\mathcal{F}}_\beta^{[\mathbf{W}]}(x')\Big), \qquad \text{(eq. (4.132))}. \tag{4.145}$$

Under endpoint gauge (Proposition 4.54 and Corollary 4.64),

$$\hat{\mathcal{F}}_\alpha^{[\mathbf{W}]}(x) \mapsto g(x)\hat{\mathcal{F}}_\alpha^{[\mathbf{W}]}(x), \qquad \mathcal{U}(x\leftarrow x';\gamma) \mapsto g(x)\,\mathcal{U}(x\leftarrow x';\gamma)\,g(x')^\dagger.$$

Then, algebraically,

$$(\hat{\mathcal{F}}'_\alpha)^\dagger \hat{\mathcal{F}}'_\beta = (\hat{\mathcal{F}}_\alpha)^\dagger g(x)^\dagger g(x) \hat{\mathcal{F}}_\beta = (\hat{\mathcal{F}}_\alpha)^\dagger \hat{\mathcal{F}}_\beta,$$

and similarly

$$(\hat{\mathcal{F}}'_\alpha)^\dagger \mathcal{U}' \hat{\mathcal{F}}'_\beta = (\hat{\mathcal{F}}_\alpha)^\dagger g(x)^\dagger g(x)\, \mathcal{U}\, g(x')^\dagger g(x')\, \hat{\mathcal{F}}_\beta = (\hat{\mathcal{F}}_\alpha)^\dagger \mathcal{U} \hat{\mathcal{F}}_\beta.$$

Thus the operators inside the expectations are unchanged; hence $M^{[\mathbf{W}]}$ and $K^{[\mathbf{W}]}$ are gauge invariant.

Step 2 (Gauge invariance of the predicates). The interference functional $\mathsf{I}^{[\mathbf{W}]}_{\alpha\beta}(x;\psi)$ multiplies $M^{[\mathbf{W}]}_{\alpha\beta}(x)$ by modal coefficients $c^*_\alpha c_\beta$ (eq. (4.143)); since $M^{[\mathbf{W}]}$ is gauge invariant and the coefficients are numbers associated to a fixed section in the same fibre, $\mathsf{I}^{[\mathbf{W}]}_{\alpha\beta}$ is gauge invariant. Likewise, the bilinear $\mathsf{E}^{[\mathbf{W}]}_{x,x'}$ uses $K^{[\mathbf{W}]}$ (Definition 4.332 and eq. (4.144)); its value and its factorisation/nonfactorisation are preserved under gauge.

Step 3 (Word change as fibrewise unitary equivalence). Let $U_x := U_{[\mathbf{W}',\mathbf{W}]}(x)$ and $U_{x'} := U_{[\mathbf{W}',\mathbf{W}]}(x')$ be the fibre unitaries from Proposition 4.284 so that $\hat{\mathcal{F}}^{[\mathbf{W}']}_\alpha(x) = U_x\,\hat{\mathcal{F}}^{[\mathbf{W}]}_\alpha(x)$. Then

$$M^{[\mathbf{W}']}(x) = U_x^\dagger\, M^{[\mathbf{W}]}(x)\, U_x, \qquad K^{[\mathbf{W}']}(x,x';\gamma) = U_x^\dagger\, K^{[\mathbf{W}]}(x,x';\gamma)\, U_{x'}.$$

(Here we use the functorial/endpoint nature of word evaluation, Proposition 4.284, which ensures that the transporter is left untouched and the endpoint letters are rotated unitarily.)

Step 4 (Word invariance of the predicates). Write the interference scalar as a quadratic form

$$\mathsf{I}^{[\mathbf{W}]}(x;\psi) = c^{[\mathbf{W}]}(x)^\dagger\, M^{[\mathbf{W}]}(x)\, c^{[\mathbf{W}]}(x),$$

with coefficient column $c^{[\mathbf{W}]}(x)$. Under $\mathbf{W} \to \mathbf{W}'$ the same physical section has coefficients $c^{[\mathbf{W}']}(x) = U_x^{-1}\, c^{[\mathbf{W}]}(x)$ in the rotated basis. Then

$$\begin{aligned}
\mathsf{I}^{[\mathbf{W}']}(x;\psi) &= \left(c^{[\mathbf{W}']}\right)^\dagger M^{[\mathbf{W}']}\, c^{[\mathbf{W}']} \\
&= \left(c^{[\mathbf{W}]}\right)^\dagger U_x^{-\dagger}\, (U_x^\dagger M^{[\mathbf{W}]} U_x)\, U_x^{-1}\, c^{[\mathbf{W}]} \\
&= \left(c^{[\mathbf{W}]}\right)^\dagger M^{[\mathbf{W}]}\, c^{[\mathbf{W}]} \\
&= \mathsf{I}^{[\mathbf{W}]}(x;\psi). \qquad (4.146)
\end{aligned}$$

Hence the presence/absence of prismatic superposition (Definition 4.331) is word invariant.

For the entanglement bilinear, let $c_1^{[\mathbf{W}]}(x), c_2^{[\mathbf{W}]}(x')$ be coefficient vectors at x, x'. Then

$$\mathsf{E}^{[\mathbf{W}]}_{x,x'}(\psi_1,\psi_2) = \left(c_1^{[\mathbf{W}]}(x)\right)^\dagger K^{[\mathbf{W}]}(x,x';\gamma)\, c_2^{[\mathbf{W}]}(x').$$

Under the word change, $c_1^{[\mathbf{W}']} = U_x^{-1} c_1^{[\mathbf{W}]}$, $c_2^{[\mathbf{W}']} = U_{x'}^{-1} c_2^{[\mathbf{W}]}$, and $K^{[\mathbf{W}']} = U_x^\dagger K^{[\mathbf{W}]} U_{x'}$. Therefore

$$\mathsf{E}^{[\mathbf{W}']}_{x,x'}(\psi_1,\psi_2) = \left(c_1^{[\mathbf{W}]}\right)^\dagger K^{[\mathbf{W}]} c_2^{[\mathbf{W}]} = \mathsf{E}^{[\mathbf{W}]}_{x,x'}(\psi_1,\psi_2).$$

In particular, factorisation of E is preserved: if $\mathsf{E}^{[\mathbf{W}]}_{x,x'}(\psi_1,\psi_2) = \varphi_x(\psi_1)^*\,\varphi_{x'}(\psi_2)$, then setting $\varphi'_x := \varphi_x \circ U_x{}^{-1}$ and $\varphi'_{x'} := \varphi_{x'} \circ U_{x'}{}^{-1}$ gives $\mathsf{E}^{[\mathbf{W}']}_{x,x'}(\psi_1,\psi_2) = \varphi'_x(\psi_1)^*\,\varphi'_{x'}(\psi_2)$; conversely, invertibility of $U_x, U_{x'}$ shows nonfactorisation cannot be created or destroyed by a word change.

Step 5 (Conclusion). Both the interference functional $\mathsf{I}^{[\mathbf{W}]}_{\alpha\beta}$ and the factorisation property of $\mathsf{E}^{[\mathbf{W}]}_{x,x'}$ are invariant under endpoint gauge (Proposition 4.54 and Corollary 4.64) and under word changes (Proposition 4.284). Therefore the predicates in Definitions 4.331 and 4.332 are well defined, independent of gauge and of the chosen endogenous word. □

Proposition 4.334 (Wordwise parasymmetric test on the Twofold slice)**.** *On the Twofold subspace* $S_x^{[\mathbf{W}]} = \mathrm{span}\{|0\rangle_x^{[\mathbf{W},\chi]}, |1\rangle_x^{[\mathbf{W},\chi]}\}$ *(Definition 4.304), let* $P_{\mathrm{para}}^{[\mathbf{W}]}$ *and* $P_{\mathrm{antipara}}^{[\mathbf{W}]}$ *be the parasymmetric/anti–parasymmetric effects (with* $P_{\mathrm{para}}^{[\mathbf{W}]} + P_{\mathrm{antipara}}^{[\mathbf{W}]} = P_{(0,1)}^{[\mathbf{W}]}$*). For* $|\psi\rangle_x^{[\mathbf{W}]} = c_0^{[\mathbf{W}]}|0\rangle_x^{[\mathbf{W},\chi]} + c_1^{[\mathbf{W}]}|1\rangle_x^{[\mathbf{W},\chi]}$*,*

$$\boxed{\left\langle P_{\mathrm{para}}^{[\mathbf{W}]} \right\rangle_\psi = \tfrac{1}{2}\Big(\langle (\hat{\mathcal{F}}_0^{[\cdot]})^\dagger \hat{\mathcal{F}}_0^{[\cdot]} \rangle \left| c_0^{[\mathbf{W}]} \right|^2 + \langle (\hat{\mathcal{F}}_1^{[\cdot]})^\dagger \hat{\mathcal{F}}_1^{[\cdot]} \rangle \left| c_1^{[\mathbf{W}]} \right|^2 + 2\ \mathrm{Re}\ \mathsf{I}_{01}^{[\mathbf{W}]}(x;\psi) \Big),} \tag{4.147}$$

so $|\psi\rangle_x^{[\mathbf{W}]}$ *is a nontrivial wordwise prismatic superposition across* $\{0,1\}$ *iff the parasymmetric/anti–parasymmetric pair yields a phase–sensitive split* ($\mathsf{I}_{01}^{[\mathbf{W}]} \neq 0$*).*

Full proof. Step 1 (Local Twofold coefficients and induced quadratic form). In the wordwise setting, the coincident–limit inner product on the Twofold slice is induced by the local wordwise kernel $M_{\alpha\beta}^{[\mathbf{W}]}(x) := \omega\Big((\hat{\mathcal{F}}_\alpha^{[\mathbf{W}]}(x))^\dagger \hat{\mathcal{F}}_\beta^{[\mathbf{W}]}(x)\Big)$ (Section 4.6, Definition 4.9). For $|\psi\rangle_x^{[\mathbf{W}]} = c_0^{[\mathbf{W}]}|0\rangle_x^{[\mathbf{W},\chi]} + c_1^{[\mathbf{W}]}|1\rangle_x^{[\mathbf{W},\chi]}$ the valuation of any Twofold effect E is a quadratic form in the coefficients, built from matrix elements of E and $M^{[\mathbf{W}]}$.

Step 2 (Parasymmetric/anti–parasymmetric effects on the Twofold). By definition, $P_{\mathrm{para}}^{[\mathbf{W}]}$ and $P_{\mathrm{antipara}}^{[\mathbf{W}]}$ are the $(\pm)$ effects associated with the 0/1 interference test on $S_x^{[\mathbf{W}]}$ ("Hadamard–like" split), satisfying $P_{\mathrm{para}}^{[\mathbf{W}]} + P_{\mathrm{antipara}}^{[\mathbf{W}]} = P_{(0,1)}^{[\mathbf{W}]}$, where $P_{(0,1)}^{[\mathbf{W}]}$ is the identity effect on the Twofold block. Consequently,

$$\left\langle P_{\mathrm{para}}^{[\mathbf{W}]} \right\rangle_\psi = \tfrac{1}{2}\Big(\left\langle P_{(0,1)}^{[\mathbf{W}]} \right\rangle_\psi + \left\langle X^{[\mathbf{W}]} \right\rangle_\psi \Big), \qquad \left\langle P_{\mathrm{antipara}}^{[\mathbf{W}]} \right\rangle_\psi = \tfrac{1}{2}\Big(\left\langle P_{(0,1)}^{[\mathbf{W}]} \right\rangle_\psi - \left\langle X^{[\mathbf{W}]} \right\rangle_\psi \Big), \tag{4.148}$$

where $X^{[\mathbf{W}]}$ is the off–diagonal "exchange" contribution that carries the phase sensitivity.

Step 3 (Evaluate the diagonal and off–diagonal parts). The diagonal (block–identity) valuation is

$$\left\langle P_{(0,1)}^{[\mathbf{W}]} \right\rangle_\psi = \left\langle (\hat{\mathcal{F}}_0^{[\cdot]})^\dagger \hat{\mathcal{F}}_0^{[\cdot]} \right\rangle \left| c_0^{[\mathbf{W}]} \right|^2 + \left\langle (\hat{\mathcal{F}}_1^{[\cdot]})^\dagger \hat{\mathcal{F}}_1^{[\cdot]} \right\rangle \left| c_1^{[\mathbf{W}]} \right|^2,$$

since the cross–terms are absent in the block identity. For the off–diagonal exchange piece, Hermiticity of $M^{[\mathbf{W}]}$ gives $M_{10}^{[\mathbf{W}]}(x) = \overline{M_{01}^{[\mathbf{W}]}(x)}$, and therefore

$$\left\langle X^{[\mathbf{W}]} \right\rangle_\psi = c_0^{[\mathbf{W}]*} c_1^{[\mathbf{W}]}\, M_{01}^{[\mathbf{W}]}(x) + c_1^{[\mathbf{W}]*} c_0^{[\mathbf{W}]}\, M_{10}^{[\mathbf{W}]}(x) = 2\ \mathrm{Re}\Big(c_0^{[\mathbf{W}]*} c_1^{[\mathbf{W}]}\, M_{01}^{[\mathbf{W}]}(x) \Big).$$

By the definition of the wordwise interference functional (Definition 4.331 and eq. (4.143)), $\mathsf{I}_{01}^{[\mathbf{W}]}(x;\psi) := M_{01}^{[\mathbf{W}]}(x)\, c_0^{[\mathbf{W}]*} c_1^{[\mathbf{W}]}$. Hence $\left\langle X^{[\mathbf{W}]} \right\rangle_\psi = 2\ \mathrm{Re}\ \mathsf{I}_{01}^{[\mathbf{W}]}(x;\psi)$.

Step 4 (Assemble and conclude). Plugging the diagonal and off–diagonal evaluations into (4.148) yields (4.147). Moreover,

$$\left\langle P_{\mathrm{para}}^{[\mathbf{W}]} \right\rangle_\psi - \left\langle P_{\mathrm{antipara}}^{[\mathbf{W}]} \right\rangle_\psi = \left\langle X^{[\mathbf{W}]} \right\rangle_\psi = 2\ \mathrm{Re}\ \mathsf{I}_{01}^{[\mathbf{W}]}(x;\psi).$$

Therefore the parasymmetric/anti–parasymmetric pair exhibits a nontrivial, phase–sensitive split iff $\mathsf{I}_{01}^{[\mathbf{W}]}(x;\psi) \neq 0$, which is exactly the predicate for a nontrivial wordwise prismatic superposition across $\{0,1\}$ in Definition 4.331. This completes the proof. □

Proposition 4.335 (Reduction to standard Hilbert–space notions in the Twofold regime (wordwise))**.** *If the wordwise Twofold sector $S_x^{[\mathbf{W}]}$ is dynamically protected and off–block couplings are negligible (Propositions 4.262 and 4.306), then there exists an isometry $U^{[\mathbf{W}]} : \mathbb{C}^2 \hookrightarrow \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ with $U^{[\mathbf{W}]}\,|0\rangle = |0\rangle_x^{[\mathbf{W},\chi]}$, $U^{[\mathbf{W}]}\,|1\rangle = |1\rangle_x^{[\mathbf{W},\chi]}$ (Definition 4.305). Consequently:*

(i) $|\psi\rangle_x^{[\mathbf{W}]}$ is a wordwise prismatic superposition across $\{0,1\}$ iff $(U^{[\mathbf{W}]})^\dagger|\psi\rangle_x^{[\mathbf{W}]}$ is a nontrivial linear combination in $\mathbb{C}^2$.

(ii) Wordwise entanglement between x, x' per Definition 4.332 holds iff the induced state on $\mathbb{C}^2 \otimes \mathbb{C}^2$ is entangled in the standard sense (after compression by $(U^{[\mathbf{W}]})^\dagger \otimes (U^{[\mathbf{W}]})^\dagger$); see also Corollary 4.308.

Full proof. Step 0 (Set–up and protection). Let $S_x^{[\mathbf{W}]} = \mathrm{span}\{|0\rangle_x^{[\mathbf{W},\chi]}, |1\rangle_x^{[\mathbf{W},\chi]}\}$ be invariant under the admissible local generators (dynamic protection), and let $P^{[\mathbf{W}]}$ be the orthogonal projector onto $S_x^{[\mathbf{W}]}$. Off–block couplings are negligible in the sense of Proposition 4.262, so that compressed dynamics on $S_x^{[\mathbf{W}]}$ closes up to controlled $O(\varepsilon)$ errors.

Step 1 (Isometry $U^{[\mathbf{W}]}$). Define $U^{[\mathbf{W}]} : \mathbb{C}^2 \to \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ by $U^{[\mathbf{W}]}\,|0\rangle = |0\rangle_x^{[\mathbf{W},\chi]}$, $U^{[\mathbf{W}]}\,|1\rangle = |1\rangle_x^{[\mathbf{W},\chi]}$ and extend linearly (Definition 4.305). Since $\{|0\rangle_x^{[\mathbf{W},\chi]}, |1\rangle_x^{[\mathbf{W},\chi]}\}$ is orthonormal in the fibre metric (induced from the correlator; cf. Section 4.6 and eq. (4.92)), one has, for all $\phi, \psi \in \mathbb{C}^2$, $\langle \phi, \psi\rangle_{\mathbb{C}^2} = \langle U^{[\mathbf{W}]}\phi,\, U^{[\mathbf{W}]}\psi\rangle_x$. Hence $U^{[\mathbf{W}]}$ is an isometry with image $S_x^{[\mathbf{W}]}$ (Proposition 4.306).

Step 2 (Superposition equivalence). Write $|\psi\rangle_x^{[\mathbf{W}]} = c_0^{[\mathbf{W}]}|0\rangle_x^{[\mathbf{W},\chi]} + c_1^{[\mathbf{W}]}|1\rangle_x^{[\mathbf{W},\chi]}$. Then $(U^{[\mathbf{W}]})^\dagger|\psi\rangle_x^{[\mathbf{W}]} = c_0^{[\mathbf{W}]}\,|0\rangle + c_1^{[\mathbf{W}]}\,|1\rangle \in \mathbb{C}^2$. If $(U^{[\mathbf{W}]})^\dagger|\psi\rangle$ is a nontrivial linear combination in $\mathbb{C}^2$ (both coefficients not simultaneously zero), the same coordinates describe $|\psi\rangle_x^{[\mathbf{W}]}$ in the fibre, so by the wordwise parasymmetric test the pair $P_{\mathrm{para}}^{[\mathbf{W}]}, P_{\mathrm{antipara}}^{[\mathbf{W}]}$ exhibits a phase–sensitive split precisely controlled by the interference term (Proposition 4.334). Conversely, if $|\psi\rangle_x^{[\mathbf{W}]}$ is a wordwise prismatic superposition across $\{0,1\}$, then $\mathsf{I}_{01}^{[\mathbf{W}]}(x;\psi) \neq 0$ for some choice of parasymmetric phase (word/basis) and therefore $c_0^{[\mathbf{W}]}, c_1^{[\mathbf{W}]}$ are not both zero; hence $(U^{[\mathbf{W}]})^\dagger|\psi\rangle$ is a nontrivial linear combination in $\mathbb{C}^2$. Word/gauge invariance of the predicate guarantees independence of the chosen word/basis (Proposition 4.333). This proves (i).

Step 3 (Entanglement equivalence). Let x, x' be two loci with protected Twofold sectors $S_x^{[\mathbf{W}]}, S_{x'}^{[\mathbf{W}]}$ and corresponding isometries $U_x^{[\mathbf{W}]}, U_{x'}^{[\mathbf{W}]}$. For test vectors $\psi_1 \in S_x^{[\mathbf{W}]}$, $\psi_2 \in S_{x'}^{[\mathbf{W}]}$, the wordwise entanglement bilinear $\mathsf{E}_{x,x'}^{[\mathbf{W}]}(\psi_1, \psi_2)$ is computed from the bilocal kernel (Definition 4.332 and eq. (4.144)). Compressing by $(U_x^{[\mathbf{W}]})^\dagger \otimes (U_{x'}^{[\mathbf{W}]})^\dagger$ gives a bilinear on $\mathbb{C}^2 \otimes \mathbb{C}^2$ whose values agree with the original ones, because $E \mapsto UEU^\dagger$ is a unital $*$–monomorphism of effect algebras on the protected block and the transport is compatible with compression (dynamic protection; Proposition 4.306). Therefore the factorisation property of $\mathsf{E}_{x,x'}^{[\mathbf{W}]}$ (product vs. nonproduct) holds iff the induced two–qubit state on $\mathbb{C}^2 \otimes \mathbb{C}^2$ is separable vs. entangled in the standard sense. This establishes (ii). The multi–site extension yielding registers follows from the same compression argument and the mixing bounds in Proposition 4.262, as summarised in Corollary 4.308.

Step 4 (Effect of small off–block couplings). If off–block terms are small but nonzero, Proposition 4.262 provides Duhamel–type bounds showing that leakage outside $S_x^{[\mathbf{W}]}$ is $O(\varepsilon t)$ and the compressed dynamics deviates from an exact SU(2) evolution by $O(\varepsilon t)$. The

conclusions of (i)–(ii) then hold to first order in ε over the control timescale, with errors controlled by those bounds.

Hence the protected wordwise Twofold sector realises the standard qubit notions by the isometry $U^{[\mathbf{W}]}$, and superposition/entanglement reduce to their usual Hilbert–space counterparts under compression. □

Remark 4.336 (Compliance–controlled regimes)**.** The holosymmetric compliance vector $\mathfrak{K} = (\Delta, \eta, \zeta)$ (Definition 4.178) bounds departures from the ideal superposition/entanglement picture: η controls two–step ps/sp leakage, ζ controls diagonal pinching stability, and Δ bounds visibility/lensing drifts in readout. The inequalities of Proposition 4.181 thus certify when wordwise Twofold superpositions behave qubit–like and when non–qubit Fourfold effects must be expected.

Lemma 4.337 (Wordwise holosymmetric dominance)**.** *In a holosymmetric interferometric test (Section 4.7.3), a phase–sensitive local signal implies the existence of $\alpha \neq \beta$ with $\mathsf{I}^{[\mathbf{W}]}_{\alpha\beta}(x;\psi) \neq 0$ as in Definition 4.331; a nonseparable bilocal signal implies that the bilinear form $\mathsf{E}^{[\mathbf{W}]}_{x,x'}$ of Definition 4.332 does not factor.*

Sketch. Write the holosymmetric readout as a Born valuation of a word–evaluated s–channel operator (eqs. (4.122) and (4.123)). Phase sensitivity under a control Hadamard requires nonzero off–diagonal modal overlaps in that operator, forcing $\langle(\hat{\mathcal{F}}^{[\cdot]}_{\alpha})^{\dagger}\hat{\mathcal{F}}^{[\cdot]}_{\beta}\rangle \neq 0$, hence $\mathsf{I}^{[\mathbf{W}]}_{\alpha\beta} \neq 0$. For the bilocal case, a nonseparable signal entails nontrivial cross–modal entries in the kernel $\langle(\hat{\mathcal{F}}^{[\cdot]}_{\alpha}(x))^{\dagger}\mathcal{U}\,\hat{\mathcal{F}}^{[\cdot]}_{\beta}(x')\rangle$, which would vanish if $\mathsf{E}^{[\mathbf{W}]}_{x,x'}$ factored; cf. Definition 4.332. □

Superposition (plain-language view, formalised). Superposition means a single system *coherently* occupies more than one compatible possibility at once, with a phase relation that can generate interference. In the qubit caricature this is $\alpha\,|0\rangle + \beta\,|1\rangle$. In the present framework, the same idea is expressed on the Fourfold modal fibre $\{|0\rangle, |1\rangle, |p\rangle, |s\rangle\}$ (Definition 4.273) by requiring that the state has support on at least two modal directions *and* a nonzero interference functional between them. Exogenously this is captured by Definition 4.325; endogenously (wordwise) it is Definition 4.331, where coherence shows up through the word–evaluated interference term $\mathsf{I}^{[\mathbf{W}]}_{\alpha\beta}(x;\psi)$ (eq. (4.143)). Intuitively: superposition is "coexistence with a phase''—and the phase is what powers interference in parasymmetric tests (Proposition 4.334).

Entanglement (plain-language view, formalised). Entanglement is superposition *spread across loci* in a way that cannot be pulled apart into independent local descriptions. In textbook language this is a non-product state on $\mathbb{C}^2 \otimes \mathbb{C}^2$; here it means the inter–locus correlator does *not* factor. Exogenously this is Definition 4.326; endogenously it is Definition 4.332, where non-factorisation of the bilinear form $\mathsf{E}^{[\mathbf{W}]}_{x,x'}(\psi_1, \psi_2)$ (eq. (4.144))—built from word–dressed channels and the transporter $\mathcal{U}$ (Definition 4.248)—certifies entanglement. Operationally, non-factorisation means measurement statistics at one locus correlate with those at another beyond any classical explanation.

Two clarifications from the holomodal framework. First, the parasymmetric/holosymmetric directions $|p\rangle, |s\rangle$ are *independent* modal directions, not mere qubit superpositions; superposition and entanglement can thus involve genuinely Fourfold structure, not only the Twofold slice (Remark 4.269 and Proposition 4.281). Second, geometry matters: wordwise holonomy in the s–channel (loop operator and signal; Section 4.7.3 and eqs. (4.122) and (4.123)) and the compliance vector $\mathfrak{K} = (\Delta, \eta, \zeta)$ (Definition 4.178) quantify how much interference survives and how cleanly non-factorisable correlations can be read (Proposition 4.181). In short, coherence (superposition) and nonlocal correlations (entanglement) are not just kinematic labels but *measurable, geometry–aware* features of the holomodal setup.

4.9 Operator-Theoretic Intrinsic Distances

Building upon IQR (Definition 1.25), the emergent quantum configuration space $\mathcal{X}_q$ (Proposition 4.625 and Section 4.19.4), the fibred $\hat{\Omega}$–Continuum (Definition 4.617), and the Quantum Holo-State $|\mathring{\mathbf{\Psi}}\rangle$ (Definition 1.27), we now formalize operator-theoretic measures of emergent distance and time between PQS loci. These quantities are relational and derive from the intrinsic quantum–geometric and correlational structure as perceived by $|\mathring{\mathbf{\Psi}}\rangle$.

We introduce three principal components of emergent distance between a PQS $\mathfrak{B}_x$ (at $x \in \mathcal{X}_q$) and another $\mathfrak{B}_{x'}$. Each furnishes one axis of an orthogonal, operator-derived chronogeometric profile.

4.9.1 Correlator-Based Distance

Intuitively, $d_{\mathrm{corr}}(x, x')$ quantifies *operator-algebraic distinguishability* between the modal structures at $\mathfrak{B}_x$ and $\mathfrak{B}_{x'}$, as mediated by the Fourfold Inter-Modal Correlator $\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})$ (see Section 4.1.2) and evaluated in the Holo-State $|\mathring{\mathbf{\Psi}}\rangle$ (Definition 1.27). Small values signal high coherence/similarity; large values signal low coherence/dissimilarity. [230, 241]

To make this precise, fix *reference modal vectors* $|\psi_1\rangle_x, |\psi_2\rangle_{x'}$ in the four-mode fibres $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)} = \mathrm{span}\{|0\rangle_x, |1\rangle_x, |p\rangle_x, |s\rangle_x\}$ and $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_{x'})}$, and write the scalar correlator amplitude

$$\mathsf{G}(x, x'; \psi_1, \psi_2) \;:=\; \Big\langle \mathring{\mathbf{\Psi}} \Big|\, \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'}) \big[\psi_1^*, \psi_2\big] \,\Big|\, \mathring{\mathbf{\Psi}} \Big\rangle,$$

where $\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2]$ is the bilinear evaluation of the (already defined) Fourfold correlator on the chosen test vectors (cf. Section 4.1.2). [382, 383, 82] Define the natural normalizations by the coincident limits

$$N(\psi_1, x) \;:=\; \Big(\mathsf{G}(x, x; \psi_1, \psi_1)\Big)^{1/2}, \qquad N(\psi_2, x') \;:=\; \Big(\mathsf{G}(x', x'; \psi_2, \psi_2)\Big)^{1/2}.$$

[82, 384]

Definition 4.338 (Correlator-Based Distance)**.** With the notation above, the *correlator-based similarity* is

$$C(x, x'; \psi_1, \psi_2) \;:=\; \frac{\big|\mathsf{G}(x, x'; \psi_1, \psi_2)\big|}{N(\psi_1, x)\, N(\psi_2, x')} \;\in\; [0, 1],$$

and the *correlator-based distance* is

$$d_{\mathrm{corr}}(x,x';\psi_1,\psi_2) \;:=\; \sqrt{1-C(x,x';\psi_1,\psi_2)} \;\in\; [0,1]. \tag{4.149}$$

[359, 385, 386, 387, 50, 388]

Discussion. (i) $d_{\mathrm{corr}}=0$ iff the normalized correlator attains unit modulus (maximal similarity under the chosen probes); $d_{\mathrm{corr}}=1$ iff the evaluated overlap vanishes. (ii) The construction is invariant under admissible reindexings and the duality involution (Proposition 1.8), and under phase rescalings of the test vectors. (iii) If desired, one may remove dependence on probes by taking an extremum over admissible $|\psi_1\rangle_x, |\psi_2\rangle_{x'}$ within a fixed class; we keep the reference-dependent form here to emphasize operational context and avoid additional optimization hypotheses. This encapsulates, without recourse to classical coordinates, the intrinsic coherence/overlap of the full Fourfold modal content across loci as registered by $|\mathring{\mathbf{\Psi}}\rangle$. In the Twofold (duality-collapse) regime (Definition 4.87), (4.149) reduces to the standard qubit overlap–to–distance transformation on the $\{0,1\}$ slice (cf. Proposition 4.328). [287, 359]

4.9.2 Wordwise Correlator-Based Distance (Holosymmetric Realization)

Fix admissible device words $\mathbf{W}_1$ at x and $\mathbf{W}_2$ at x' with RR elements $\mathsf{W}_{1,\chi\leftarrow\chi}(x)$ and $\mathsf{W}_{2,\chi'\leftarrow\chi'}(x')$, acting on the modal fibres as in the wordwise calculus (cf. Sections 4.5 and 4.6.2). For any reference modal vectors $|\psi_1\rangle_x \in \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ and $|\psi_2\rangle_{x'} \in \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_{x'})}$, define their *word-rotated* representatives

$$|\psi_1\rangle_x^{[\mathbf{W}_1,\chi]} := \mathsf{W}_{1,\chi\leftarrow\chi}(x)\,|\psi_1\rangle_x, \qquad |\psi_2\rangle_{x'}^{[\mathbf{W}_2,\chi']} := \mathsf{W}_{2,\chi'\leftarrow\chi'}(x')\,|\psi_2\rangle_{x'}.$$

The *wordwise scalar correlator amplitude* is

$$\mathsf{G}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\psi_1,\psi_2) := \Big\langle \mathring{\mathbf{\Psi}}\Big|\, \hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})\Big[\big(\psi_1^{[\mathbf{W}_1,\chi]}\big)^*,\ \psi_2^{[\mathbf{W}_2,\chi']}\Big]\,\Big|\mathring{\mathbf{\Psi}}\Big\rangle, \tag{4.150}$$

where the bilinear evaluation of the Fourfold correlator is as in Section 4.1.2 but applied only to word-rotated test vectors; the Holo-State $|\mathring{\mathbf{\Psi}}\rangle$ is as in Definition 1.27. [230, 241, 382, 383, 82]

Wordwise normalizations. Using the coincident-limit inner product from Section 4.6, set

$$N^{[\mathbf{W}_1]}(\psi_1,x) := \Big(\,\mathsf{G}^{[\mathbf{W}_1,\mathbf{W}_1]}(x,x;\psi_1,\psi_1)\,\Big)^{1/2}, \quad N^{[\mathbf{W}_2]}(\psi_2,x') := \Big(\,\mathsf{G}^{[\mathbf{W}_2,\mathbf{W}_2]}(x',x';\psi_2,\psi_2)\,\Big)^{1/2}. \tag{4.151}$$

These coincide with the exogenous normalizations when $\mathbf{W}_1=\mathbf{W}_2=\mathbf{1}$ by unitary invariance of the fibre metric and the endogenous evaluation map. [82, 384]

Definition 4.339 (Wordwise Correlator-Based Distance)**.** With the notation above, the *wordwise correlator-based similarity* is

$$C^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\psi_1,\psi_2) := \frac{\big|\mathsf{G}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\psi_1,\psi_2)\big|}{N^{[\mathbf{W}_1]}(\psi_1,x)\,N^{[\mathbf{W}_2]}(\psi_2,x')} \;\in [0,1], \tag{4.152}$$

and the *wordwise correlator-based distance* is

$$d_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\psi_1,\psi_2) := \sqrt{1-C^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\psi_1,\psi_2)} \;\in [0,1]. \tag{4.153}$$

By Cauchy–Schwarz in the fibre inner product from Section 4.6, $C^{[\mathbf{W}_1,\mathbf{W}_2]} \in [0,1]$ and hence (4.153) is well-defined. [359, 385, 386, 387, 50, 388]

Proposition 4.340 (Word/Gauge Invariance and Equivalence to Exogenous Form)**.** *For any admissible words* $\mathbf{W}_1, \mathbf{W}_2$ *and endpoint gauges,*

$$C^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\psi_1,\psi_2) = C(x,x';\psi_1,\psi_2), \qquad d^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{corr}}(x,x';\psi_1,\psi_2) = d_{\mathrm{corr}}(x,x';\psi_1,\psi_2),$$

where C, d_{corr} *are as in Definition 4.338.* Sketch. *Word-rotation multiplies both probes by local unitaries* $\mathsf{W}_{1,\chi\leftarrow\chi}(x)$ *and* $\mathsf{W}_{2,\chi'\leftarrow\chi'}(x')$*; by unitary invariance of the fibre inner product and the bilinear evaluation of* $\hat{G}_\Lambda$ *(cf. Section 4.5), the numerator and denominators in* (4.152) *transform covariantly and their ratio is invariant. Endpoint gauges act similarly on both legs; adjoint reciprocity of the correlator ensures modulus invariance.*

Proof. Let $U_1 := \mathsf{W}_{1,\chi\leftarrow\chi}(x) \in \mathrm{U}(4)$ and $U_2 := \mathsf{W}_{2,\chi'\leftarrow\chi'}(x') \in \mathrm{U}(4)$ denote the RR elements of the admissible device words at the endpoints (cf. Sections 4.5 and 4.6.2). By definition of the wordwise amplitude (4.150) and sesquilinearity of the Fourfold evaluation (Proposition 4.12; see also Section 4.1.2),

$$\begin{aligned}\mathsf{G}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\psi_1,\psi_2) &= \Big\langle \mathring{\mathbf{\Psi}} \Big|\, \hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})\Big[(U_1\psi_1)^*,\, U_2\psi_2\Big] \,\Big|\mathring{\mathbf{\Psi}}\Big\rangle \\ &= \Big\langle \mathring{\mathbf{\Psi}} \Big|\, U_1^\dagger\, \hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})\Big[\psi_1^*,\psi_2\Big]\, U_2 \,\Big|\mathring{\mathbf{\Psi}}\Big\rangle.\end{aligned} \tag{4.154}$$

Similarly, using the coincident-limit inner product from Section 4.6, the wordwise normalizations (4.151) satisfy

$$\begin{aligned}\Big(N^{[\mathbf{W}_1]}(\psi_1,x)\Big)^2 &= \Big\langle \mathring{\mathbf{\Psi}} \Big|\, U_1^\dagger\, \hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_x)\Big[\psi_1^*,\psi_1\Big]\, U_1 \,\Big|\mathring{\mathbf{\Psi}}\Big\rangle, \\ \Big(N^{[\mathbf{W}_2]}(\psi_2,x')\Big)^2 &= \Big\langle \mathring{\mathbf{\Psi}} \Big|\, U_2^\dagger\, \hat{G}_\Lambda(\mathfrak{B}_{x'},\mathfrak{B}_{x'})\Big[\psi_2^*,\psi_2\Big]\, U_2 \,\Big|\mathring{\mathbf{\Psi}}\Big\rangle.\end{aligned} \tag{4.155}$$

Step 1: Endpoint word/gauge covariance of the evaluation. By wordwise covariant kinematics (Section 4.5) and endpoint gauge covariance of the channel components (stated componentwise in Section 4.6.3 and globally in Section 4.1.2), the insertion of U_1, U_2 at the endpoints implements a simultaneous change of trivialization in the modal fibres. The Holo-State expectation is invariant under such endogenous relabellings (Definition 1.27 together with endogenous expectation invariance, cf. Remark 4.372), hence

$$\Big|\Big\langle \mathring{\mathbf{\Psi}} \Big|\, U_1^\dagger\, A\, U_2 \,\Big|\mathring{\mathbf{\Psi}}\Big\rangle\Big| = \Big|\Big\langle \mathring{\mathbf{\Psi}} \Big|\, A \,\Big|\mathring{\mathbf{\Psi}}\Big\rangle\Big| \quad \text{for any operator } A \text{ built from } \hat{G}_\Lambda \text{ by bilinear evaluation.} \tag{4.156}$$

(Adjoint reciprocity of $\hat{G}_\Lambda$ ensures that endpoint insertions do not affect the modulus; see Proposition 4.13.)

Step 2: Invariance of numerator and denominators. Applying (4.156) to (4.154) yields

$$\Big|\mathsf{G}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\psi_1,\psi_2)\Big| = \Big|\mathsf{G}(x,x';\psi_1,\psi_2)\Big|. \tag{4.157}$$

Likewise, applying (4.156) to (4.155) gives

$$N^{[\mathbf{W}_1]}(\psi_1,x) = N(\psi_1,x), \qquad N^{[\mathbf{W}_2]}(\psi_2,x') = N(\psi_2,x'). \tag{4.158}$$

Step 3: Equality of similarities and distances. Combining (4.157)–(4.158) with the definitions (4.152) and (4.181), and comparing with the exogenous definitions in Definition 4.338, we obtain

$$C^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\psi_1,\psi_2) = C(x,x';\psi_1,\psi_2), \qquad d_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\psi_1,\psi_2) = d_{\mathrm{corr}}(x,x';\psi_1,\psi_2).$$

Endpoint gauges. If, in addition, endpoint gauges $g(x), g(x') \in \mathrm{U}(4)$ are applied, the same argument holds with $U_1 \mapsto g(x)U_1$ and $U_2 \mapsto U_2 g(x')^\dagger$, using gauge covariance of the correlator and its bilinear evaluation (Sections 4.1.2 and 4.5) together with Proposition 4.13. Hence the equalities persist under endpoint gauges.

This proves the proposition. □

Discussion (wordwise). (i) $d_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]} = 0$ iff the normalized wordwise correlator attains unit modulus; $d_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]} = 1$ iff the evaluated overlap vanishes. (ii) The construction is invariant under admissible word changes (device-level relabellings), the duality involution (Proposition 1.8), and phase rescalings of probes. (iii) Probe-dependence can be removed by extremizing over admissible wordwise probes if desired, but we retain the reference-dependent form to emphasize operational context. (iv) In the Twofold (duality-collapse) regime (Definition 4.87), (4.181) reduces to the standard qubit overlap–distance on the $\{0,1\}$ slice (cf. Proposition 4.328). [287, 359]

4.9.3 The Spectral Operator as a Fundamental Geometric Generator

Within IQR, the emergent geometry is governed by two complementary generators. The first is the Fourfold Inter-Modal Correlator $\hat{G}_\Lambda$ (Definition 4.9), which furnishes the relational/integral structure. The second is the *Spectral Operator* $\hat{D}$, which supplies the differential structure. It is introduced axiomatically via its mathematical properties and its operational role, without redefinition of $\hat{G}_\Lambda$ or the Quantum Holo-State $|\overset{\circ}{\mathbf{\Psi}}\rangle$ (Definition 1.27).

Definition 4.341 (Spectral Operator $\hat{D}$)**.** The *Spectral Operator* $\hat{D}$ is a densely defined self-adjoint operator acting fibrewise on the modal Hilbert spaces $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ (Definition 4.273 and Section 4.6) and hence on the emergent Hilbert space $\mathcal{H}_{\mathrm{em}}$. It satisfies:

(a) **Self-adjointness (fibrewise).** For each locus $x \in \mathcal{X}_q$, the localisation $\hat{D}_x$ is self-adjoint on a dense domain in $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$.

(b) **Discrete (compact-resolvent) spectrum.** Each $\hat{D}_x$ has compact resolvent, hence a real, discrete spectrum $\{\lambda_n^{(x)}\}_{n\in\mathbb{N}}$ with orthonormal eigenbasis $\{|\psi_n^{(x)}\rangle\}$:

$$\hat{D}_x\,|\psi_n^{(x)}\rangle = \lambda_n^{(x)}\,|\psi_n^{(x)}\rangle. \tag{4.159}$$

(c) **Relational differential role (commutator action).** The generator $\hat{D}$ acts differentially on relational data through commutators with Fourfold correlators (Definition 4.9). For any eigenmode $|\psi_n^{(x)}\rangle$,

$$\mathcal{S}_n(x,x') \;:=\; \Big\langle \overset{\circ}{\mathbf{\Psi}} \Big|\, \Big[\, \hat{D},\; \hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})[\psi_n^*,\psi_n] \,\Big] \,\Big| \overset{\circ}{\mathbf{\Psi}} \Big\rangle, \tag{4.160}$$

is the *spectral shear* observable measuring the $\hat{D}$–induced differential response of the bilocal channel in the Holo-State.

The Axiom of Spectral Smoothness.

Definition 4.342 (Axiom of Spectral Smoothness)**.** There exist constants $C, k \geq 0$ and a neighbourhood $U \subset \mathcal{X}_q \times \mathcal{X}_q$ of the diagonal such that, for all $x, x' \in U$ and all eigenmodes $|\psi_n^{(x)}\rangle$ of $\hat{D}_x$,

$$\left|\mathcal{S}_n(x, x')\right| \;\leq\; C\left(1 + |n|\right)^k,$$

with the bound uniform under admissible parallel transports used in $\hat{G}_\Lambda$ (Definition 4.9).

Admissible Spectral Operators.

Definition 4.343 (Admissible Spectral Operators)**.** A densely defined operator $\hat{D}$ acting fibrewise on the modal Hilbert spaces $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}$ (and hence on $\mathcal{H}_{\text{em}}$; cf. Definition 4.273 and Section 4.6) is called *admissible* if and only if the following explicit conditions hold:

(i) **Self-adjointness (fibrewise).** For each locus $x \in \mathcal{X}_q$, the localisation $\hat{D}_x$ is self-adjoint on a dense domain in $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}$.

(ii) **Discrete (compact-resolvent) spectrum.** For each $x \in \mathcal{X}_q$, $\hat{D}_x$ has compact resolvent; hence there exist a real, discrete spectrum $\{\lambda_n^{(x)}\}_{n\in\mathbb{N}}$ and an orthonormal eigenbasis $\{|\psi_n^{(x)}\rangle\}_{n\in\mathbb{N}} \subset \mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}$ such that

$$\hat{D}_x\,|\psi_n^{(x)}\rangle = \lambda_n^{(x)}\,|\psi_n^{(x)}\rangle.$$

(iii) **Spectral Smoothness (differential bound near the diagonal).** Let $\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\cdot,\cdot]$ denote the Fourfold inter-modal correlator (Definition 4.9) built with admissible parallel transports. For all x, x' in some neighbourhood $U \subset \mathcal{X}_q \times \mathcal{X}_q$ of the diagonal, and for every eigenmode $|\psi_n^{(x)}\rangle$ from (ii), define the *spectral shear*

$$\mathcal{S}_n(x, x') \;:=\; \Big\langle \mathring{\mathbf{\Psi}}\Big|\,\Big[\,\hat{D},\; \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_n^*, \psi_n]\,\Big]\,\Big|\mathring{\mathbf{\Psi}}\Big\rangle.$$

There exist constants $C, k \geq 0$ such that the bound

$$\left|\mathcal{S}_n(x, x')\right| \;\leq\; C\left(1 + |n|\right)^k$$

holds uniformly (in particular, uniformly over the admissible parallel transports used in $\hat{G}_\Lambda$).

The class of all such operators is denoted $\mathfrak{D}_{\text{admissible}}$.

Remark 4.344 (Links to Spectral Distance and Time)**.** The commutator structure in (4.160) underlies the *spectral distance* and *spectral shear time* constructions introduced later (Sections 4.9.6 and 4.12.5). In particular, the boundedness supplied by Definition 4.342 guarantees well-posedness of these operator-valued notions in the coincident and near-coincident regimes, without any extra assumptions on $\hat{G}_\Lambda$ or $|\mathring{\mathbf{\Psi}}\rangle$.

4.9.4 Holosymmetric Wordwise Spectral Operators

Definition 4.345 (Wordwise Spectral Operator $\mathbb{D}^{[\mathbf{W}]}$)**.** Fix a device word $\mathbf{W}$ (and control χ) as in Sections 4.2, 4.5 and 4.6.2. A *wordwise spectral operator* is a densely defined operator

$$\mathbb{D}^{[\mathbf{W}]} : \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)} \to \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$$

acting fibrewise, with all adjoints taken with respect to the word-dressed inner product $\langle\cdot|\cdot\rangle_x^{[\mathbf{W}]}$ induced by the wordwise coincident-limit kernel (Sections 4.6 and 4.6.2). It satisfies:

(a) **Self-adjointness (fibrewise, word-dressed).** For each locus $x \in \mathcal{X}_q$, the localization $\mathbb{D}_x^{[\mathbf{W}]}$ is self-adjoint on a dense domain in $\left(\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}, \langle\cdot|\cdot\rangle_x^{[\mathbf{W}]}\right)$.

(b) **Discrete (compact-resolvent) spectrum (word-dressed).** Each $\mathbb{D}_x^{[\mathbf{W}]}$ has compact resolvent, hence a real, discrete spectrum $\{\lambda_{n,[\mathbf{W}]}^{(x)}\}_{n\in\mathbb{N}}$ with orthonormal eigenbasis $\{|\psi_{n,[\mathbf{W}]}^{(x)}\rangle\}$:

$$\mathbb{D}_x^{[\mathbf{W}]}\,|\psi_{n,[\mathbf{W}]}^{(x)}\rangle = \lambda_{n,[\mathbf{W}]}^{(x)}\,|\psi_{n,[\mathbf{W}]}^{(x)}\rangle. \tag{4.161}$$

(c) **Relational differential role via endogenous super-commutator.** Let $\hat{G}_\Lambda^{[\mathbf{W}]}(\mathfrak{B}_x, \mathfrak{B}_{x'})[\cdot,\cdot]$ denote the word-dressed Fourfold inter-modal correlator letter/word (Sections 4.2 and 4.6.3). Using the endogenous bracket $[\cdot,\cdot]_{\mathrm{endo}}$ from Section 4.3.1, define for any eigenmode $|\psi_{n,[\mathbf{W}]}^{(x)}\rangle$ the *wordwise spectral shear*

$$\mathcal{S}_n^{[\mathbf{W}]}(x,x') \;:=\; \left\langle \mathring{\mathbf{\Psi}} \right|\Big[\,\mathbb{D}^{[\mathbf{W}]},\; \hat{G}_\Lambda^{[\mathbf{W}]}(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_{n,[\mathbf{W}]}^*, \psi_{n,[\mathbf{W}]}]\,\Big]_{\mathrm{endo}}\left|\mathring{\mathbf{\Psi}}\right\rangle. \tag{4.162}$$

Wordwise Axiom of Spectral Smoothness.

Definition 4.346 (Wordwise Axiom of Spectral Smoothness)**.** There exist constants $C, k \geq 0$ and a neighbourhood $U \subset \mathcal{X}_q \times \mathcal{X}_q$ of the diagonal such that, for all $x, x' \in U$ and all eigenmodes $|\psi_{n,[\mathbf{W}]}^{(x)}\rangle$ of $\mathbb{D}_x^{[\mathbf{W}]}$,

$$\left|\mathcal{S}_n^{[\mathbf{W}]}(x,x')\right| \;\leq\; C\left(1+|n|\right)^k,$$

with the bound *uniform* under admissible wordwise parallel transports used in $\hat{G}_\Lambda^{[\mathbf{W}]}$ (Sections 4.5 and 4.6.3).

Admissible Wordwise Spectral Operators.

Definition 4.347 (Admissible Wordwise Spectral Operators)**.** A densely defined operator $\mathbb{D}^{[\mathbf{W}]}$ acting fibrewise on the word-dressed modal Hilbert spaces $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ is called *admissible wordwise* iff:

(i) *Self-adjointness (word-dressed)* holds as in Definition 4.345(a);

(ii) *Compact resolvent (discrete spectrum)* holds as in Definition 4.345(b), with eigenpairs $\left(\lambda_{n,[\mathbf{W}]}^{(x)}, |\psi_{n,[\mathbf{W}]}^{(x)}\rangle\right)$ satisfying (4.161);

(iii) The *Wordwise Axiom of Spectral Smoothness* holds as in Definition 4.346, with spectral shear $\mathcal{S}_n^{[\mathbf{W}]}(x,x')$ defined by (4.162).

The class of all such operators is denoted $\mathfrak{D}^{[\mathbf{W}]}_{\text{admissible}}$.

Remark 4.348 (Unitary equivalence across words)**.** If $U_x^{[\mathbf{W}\to\mathbf{W}']}:\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}\to\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}$ is the frame change induced by a word change (Section 4.6.2), then $\mathbb{D}_x^{[\mathbf{W}']}:=U_x^{[\mathbf{W}\to\mathbf{W}']}\,\mathbb{D}_x^{[\mathbf{W}]}\left(U_x^{[\mathbf{W}\to\mathbf{W}']}\right)^\dagger$ preserves admissibility and spectra. Thus the wordwise construction is representation-independent.

Theorem 4.349 (Wordwise Spectral Smoothness from transport regularity)**.** *Assume the wordwise framework of Sections 4.2, 4.3.1, 4.5, 4.6.2 and 4.6.3, and let $\mathbb{D}^{[\mathbf{W}]}$ be an admissible wordwise spectral operator (Definition 4.347). Then there exist a neighbourhood $U\subset\mathcal{X}_q\times\mathcal{X}_q$ of the diagonal and a constant $C\geq 0$ such that, for all $x,x'\in U$ and all eigenmodes $|\psi^{(x)}_{n,[\mathbf{W}]}\rangle$ of $\mathbb{D}_x^{[\mathbf{W}]}$,*

$$\left|\mathcal{S}_n^{[\mathbf{W}]}(x,x')\right| \;\leq\; C.$$

In particular, the estimate of Definition 4.346 holds with $k=0$, hence with any $k\geq 0$.

Proof. **Step 0 (setup).** Fix x,x' and a short path $\gamma:x'\to x$ with length $\ell=d(x,x')$ inside a normal neighbourhood. The word-dressed letter with endpoint eigenmode insertions is

$$\hat{G}_\Lambda^{[\mathbf{W}]}(\mathfrak{B}_x,\mathfrak{B}_{x'})[\psi^*_{n,[\mathbf{W}]},\psi_{n,[\mathbf{W}]}] \;=\; \hat{\mathcal{F}}_\alpha^{[\mathbf{W}]}(x)^\dagger\,\underbrace{\mathcal{U}^{[\mathbf{W}]}(x\leftarrow x';\gamma)}_{=:U_\gamma}\,\hat{\mathcal{F}}_\beta^{[\mathbf{W}]}(x')\,|\psi^{(x')}_{n,[\mathbf{W}]}\rangle\langle\psi^{(x')}_{n,[\mathbf{W}]}|$$

(with α,β the chosen modes; selectors are contractions, U_γ unitary).

The shear is the expectation of the endogenous bracket:

$$\mathcal{S}_n^{[\mathbf{W}]}(x,x')=\langle\Omega|[\mathbb{D}^{[\mathbf{W}]},\hat{G}_\Lambda^{[\mathbf{W}]}(\cdot)]_{\text{endo}}|\Omega\rangle.$$

Step 1 (eigen-endpoint commutator vanishes). At the x'*-end*, the rank-one projector $P_{n,x'}:=|\psi^{(x')}_{n,[\mathbf{W}]}\rangle\langle\psi^{(x')}_{n,[\mathbf{W}]}|$ commutes with $\mathbb{D}^{[\mathbf{W}]}_{x'}$ since $\mathbb{D}^{[\mathbf{W}]}_{x'}\psi^{(x')}_{n,[\mathbf{W}]}=\lambda^{(x')}_{n,[\mathbf{W}]}\psi^{(x')}_{n,[\mathbf{W}]}$:

$$[\mathbb{D}^{[\mathbf{W}]}_{x'},P_{n,x'}]=0.$$

Thus any $P_{n,x'}$ contribution to the bracket disappears; the n–dependence is neutralised at the source.

Step 2 (graded Leibniz for the endogenous bracket). Using the graded Leibniz rule for $[\cdot,\cdot]_{\text{endo}}$ and the charge-preserving nature of selectors and U_γ, we expand

$$\begin{aligned}\Big[\mathbb{D}^{[\mathbf{W}]},\hat{\mathcal{F}}_\alpha^\dagger\,U_\gamma\,\hat{\mathcal{F}}_\beta\,P_{n,x'}\Big]_{\text{endo}}&\\ =\;[\mathbb{D}^{[\mathbf{W}]},\hat{\mathcal{F}}_\alpha^\dagger]_{\text{endo}}\,U_\gamma\,\hat{\mathcal{F}}_\beta\,P_{n,x'}&\\ \pm\;\hat{\mathcal{F}}_\alpha^\dagger\,[\mathbb{D}^{[\mathbf{W}]},U_\gamma]_{\text{endo}}\,\hat{\mathcal{F}}_\beta\,P_{n,x'}&\\ \pm\;\hat{\mathcal{F}}_\alpha^\dagger\,U_\gamma\,[\mathbb{D}^{[\mathbf{W}]},\hat{\mathcal{F}}_\beta]_{\text{endo}}\,P_{n,x'},&\end{aligned}\tag{4.163}$$

where the signs are fixed by the charges (they do not affect norms).

Step 3 (uniform bounds on selector commutators). By admissibility/regularity of the wordwise calculus, each selector is bounded and $[\mathbb{D}^{[\mathbf{W}]}, \hat{\mathcal{F}}_\bullet]_{\mathrm{endo}}$ is a bounded operator; set $L_{\mathrm{sel}} := \max\{\|[\mathbb{D}^{[\mathbf{W}]}, \hat{\mathcal{F}}_\alpha]\|,\ \|[\mathbb{D}^{[\mathbf{W}]}, \hat{\mathcal{F}}_\beta]\|\}$ (finite and independent of n, γ).

Step 4 (transport bound from covariant kinematics). From the wordwise covariant-kinematics equation $\dot{U}_\gamma(t) = -\mathcal{A}_\mu(\gamma(t))\,\dot{\gamma}^\mu(t)\,U_\gamma(t)$ and the variation-of-constants formula, we get (for small ℓ)

$$\left\|[\mathbb{D}^{[\mathbf{W}]}, U_\gamma]_{\mathrm{endo}}\right\| \ \le\ \int_0^1 \left\|[\mathbb{D}^{[\mathbf{W}]}, \mathcal{A}_\mu(\gamma(t))]_{\mathrm{endo}}\right\| \,|\dot{\gamma}^\mu(t)|\, dt \ \le\ L_{\mathrm{conn}}\ \ell \ +\ O(\ell^2),$$

where $L_{\mathrm{conn}} := \sup_{\text{near diag}} \|[\mathbb{D}^{[\mathbf{W}]}, \mathcal{A}_\mu]\|$. Curvature/holonomy control in the same neighbourhood gives the $O(\ell^2)$ remainder.

Step 5 (take norms). Selectors are contractions, U_γ is unitary, and $\|P_{n,x'}\| = 1$, hence

$$\left\|[\mathbb{D}^{[\mathbf{W}]}, \hat{G}^{[\mathbf{W}]}_\Lambda(\cdot)]_{\mathrm{endo}}\right\| \ \le\ 2L_{\mathrm{sel}} \ +\ L_{\mathrm{conn}}\ \ell \ +\ O(\ell^2).$$

Step 6 (expectation and uniformity). Therefore

$$\left|\mathcal{S}^{[\mathbf{W}]}_n(x,x')\right| \ =\ \left|\langle\Omega|[\cdot]_{\mathrm{endo}}|\Omega\rangle\right| \ \le\ \left\|[\cdot]_{\mathrm{endo}}\right\| \ \le\ C_0 \ +\ C_1\, d(x,x') \ +\ O\Big(d(x,x')^2\Big),$$

with $C_0 = 2L_{\mathrm{sel}}$ and $C_1 = L_{\mathrm{conn}}$, both independent of n and uniform over admissible wordwise transports in the chosen neighbourhood U. Shrinking U if necessary, absorb the linear/quadratic terms into a single C and conclude $\left|\mathcal{S}^{[\mathbf{W}]}_n(x,x')\right| \le C$, i.e. Definition 4.346 with $k = 0$. □

Remark 4.350 (Why no n–growth appears)**.** The only n–dependence enters through the endpoint projector $P_{n,x'}$, but $[\mathbb{D}^{[\mathbf{W}]}_{x'}, P_{n,x'}] = 0$ by eigenmode pinning at x'. All remaining terms are governed by wordwise selector/connection commutators, whose norms are independent of n. Thus the shear is uniformly bounded; the polynomial form in Definition 4.346 then holds with $k = 0$ (and *a fortiori* with any $k \ge 0$).

Corollary 4.351 (Word invariance)**.** *Under a charge-preserving word change (Section 4.6.2), $\mathcal{S}^{[\mathbf{W}]}_n$ and the constants (C, U) are invariant by Remark 4.372.*

Admissible Wordwise Spectral Operators (derived).

Definition 4.352 (Admissible Wordwise Spectral Operators — derived)**.** A densely defined operator $\mathbb{D}^{[\mathbf{W}]}$ acting fibrewise on the word-dressed modal Hilbert spaces $\mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}}$ is *admissible wordwise (derived)* iff:

(i) **Self-adjointness (word-dressed)** holds as in Definition 4.345(a).

(ii) **Compact resolvent (discrete spectrum)** holds as in Definition 4.345(b), with eigenpairs $\left(\lambda^{(x)}_{n,[\mathbf{W}]}, |\psi^{(x)}_{n,[\mathbf{W}]}\rangle\right)$ satisfying (4.161).

(iii) **Wordwise regularity (implies smoothness).** The wordwise data satisfy (r,s)-regularity relative to $\mathbb{D}^{[\mathbf{W}]}$ as in Definition 4.373 on a normal neighbourhood of the diagonal $U \subset \mathcal{X}_q \times \mathcal{X}_q$.

We denote this class by $\mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,der}}$.

Theorem 4.353 (Derived admissibility ⇒ wordwise spectral smoothness)**.** *Let $\mathbb{D}^{[\mathbf{W}]}$ satisfy Definition 4.352. Then there exist constants $C, k \geq 0$ and a neighbourhood U of the diagonal such that, for all $x, x' \in U$ and all eigenmodes $|\psi^{(x)}_{n,[\mathbf{W}]}\rangle$,*

$$\left|\mathcal{S}^{[\mathbf{W}]}_n(x,x')\right| = \left|\left\langle \mathring{\mathbf{\Psi}}\right| \left[\, \mathbb{D}^{[\mathbf{W}]},\ \hat{G}^{[\mathbf{W}]}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi^*_{n,[\mathbf{W}]}, \psi_{n,[\mathbf{W}]}] \,\right]_{\mathrm{endo}} \left|\mathring{\mathbf{\Psi}}\right\rangle\right| \ \leq\ C\left(1+|n|\right)^k, \tag{4.164}$$

with the bound uniform *over admissible wordwise transports used in $\hat{G}^{[\mathbf{W}]}_\Lambda$ and* independent *of the word $\mathbf{W}$. Moreover,*

$$k \ =\ \vartheta \cdot \max\{\, r, s \,\}, \tag{4.165}$$

where ϑ is the Weyl growth exponent of the wordwise spectrum on the relevant compact base (e.g. $\vartheta = 1$ in 1D Dirac-type, $\vartheta = \frac{1}{2}$ in 2D, $\vartheta = \frac{1}{d}$ in dD).

Proof sketch. By (r,s)-regularity (Definition 4.373), the selector part yields $\|[\mathbb{D}^{[\mathbf{W}]}, \hat{\mathcal{F}}^{[\mathbf{W}]}_\bullet]\| \leq C_{\mathrm{sel}}$ (order r), and the transport part admits a Duhamel bound

$$\left\|[\mathbb{D}^{[\mathbf{W}]}, U_\gamma]\right\| \ \leq\ \int_\gamma \left\|[\mathbb{D}^{[\mathbf{W}]}, \mathcal{A}^{[\mathbf{W}]}(\dot\gamma)]\right\| ds \ \leq\ C_{\mathrm{conn}} \cdot \mathrm{length}(\gamma),$$

with $C_{\mathrm{conn}} = \sup \|[\mathbb{D}^{[\mathbf{W}]}, \mathcal{A}^{[\mathbf{W}]}]\|$ (order $s = 1$ in the first-order case). Combining the selector and transport contributions inside the endogenous bracket and using the fact that $[\mathbb{D}^{[\mathbf{W}]}_{x'}, P_{n,x'}] = 0$ for endpoint eigenprojectors, one obtains a local bound

$$\left|\mathcal{S}^{[\mathbf{W}]}_n(x,x')\right| \ \leq\ C_0 + C_1\,\mathrm{length}(\gamma),$$

uniform in n at fixed geometry. To convert geometric control into spectral control, use Weyl growth on the compact base: $|\lambda^{(x)}_{n,[\mathbf{W}]}| \sim n^\vartheta$, which promotes the first-order geometric bound to the polynomial n-bound with exponent $k = \vartheta \max\{r,s\}$ (cf. Theorem 4.384). Word/gauge invariance (Proposition 4.382) makes the constants independent of $\mathbf{W}$. □

Corollary 4.354 (Inclusion of classes)**.** *Every derived-admissible operator is wordwise spectrally smooth in the sense of Definition 4.383, hence*

$$\mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,der}} \ \subseteq\ \mathfrak{D}^{[\mathbf{W}]}_{\mathrm{admissible}}. \tag{4.166}$$

If, in addition, the base admits Weyl growth with exponent ϑ and the wordwise data are first-order ($r = 0$, $s = 1$), then (4.164) holds with $k = \vartheta$ uniformly in $\mathbf{W}$.

Remark 4.355 (Dependence of constants and locality)**.** The constant C in (4.164) may be chosen of the form $C \lesssim C_{\mathrm{sel}} + C_{\mathrm{conn}} \cdot \mathrm{diam}(U) + C_{\mathrm{curv}} \cdot \mathrm{area}(U)$, where $C_{\mathrm{sel}} = \sup \|[\mathbb{D}^{[\mathbf{W}]}, \hat{\mathcal{F}}^{[\mathbf{W}]}_\bullet]\|$, $C_{\mathrm{conn}} = \sup \|[\mathbb{D}^{[\mathbf{W}]}, \mathcal{A}^{[\mathbf{W}]}]\|$, and C_{curv} controls higher-order (curvature) corrections from the transporter expansion; all suprema are taken over the normal neighbourhood U and over admissible words. A partition-of-unity argument patches local neighbourhoods to give a global statement on compact bases.

Remark 4.356 (Equivalence under additional hypotheses)**.** If every wordwise admissible operator (Definition 4.347) on the given base admits (r, s)-regular representatives with the same Weyl exponent ϑ, then the two classes coincide, $\mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,der}} = \mathfrak{D}^{[\mathbf{W}]}_{\mathrm{admissible}}$. In particular this holds for Dirac-type generators on compact d-dimensional bases with smooth wordwise connections.

Definition 4.357 (Admissible Wordwise Spectral Operators — polynomial)**.** In addition to Definition 4.352(i)–(iii), assume a wordwise Weyl growth condition for $\mathbb{D}^{[\mathbf{W}]}$ (Definition 4.376) with exponent $\vartheta > 0$ (i.e. there exist $c_\pm > 0$ and n_0 such that for all $n \geq n_0$, $c_- n^\vartheta \leq |\lambda^{(x)}_{n,[\mathbf{W}]}| \leq c_+ n^\vartheta$ uniformly on a normal neighbourhood). We call such $\mathbb{D}^{[\mathbf{W}]}$ *admissible wordwise (polynomial)* and write

$$\mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,poly}} := \Big\{ \mathbb{D}^{[\mathbf{W}]} \in \mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,der}} \text{ with wordwise Weyl exponent } \vartheta > 0 \Big\}.$$

Proposition 4.358 (Spectral smoothness is automatic)**.** *If $\mathbb{D}^{[\mathbf{W}]} \in \mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,der}}$, then the Wordwise Spectral Smoothness bound holds in the uniform form of Definition 4.383 (i.e. with $k = 0$). If moreover $\mathbb{D}^{[\mathbf{W}]} \in \mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,poly}}$, then the polynomial form of the wordwise spectral smoothness bound holds with exponent*

$$k \;=\; \vartheta \, \max\{r, s\}, \tag{4.167}$$

hence the statement of Definition 4.346.

Proof sketch. For $\mathbb{D}^{[\mathbf{W}]} \in \mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,der}}$, (r, s)-regularity (Definition 4.373) yields bounded selector commutators and a Duhamel–type estimate for transport, $\|[\mathbb{D}^{[\mathbf{W}]}, U_\gamma]\| \leq \int_\gamma \|[\mathbb{D}^{[\mathbf{W}]}, \mathcal{A}^{[\mathbf{W}]}]\| \, ds$, on a normal neighbourhood. Inserting these into the endogenous bracket and using that $[\mathbb{D}^{[\mathbf{W}]}_{x'}, P_{n,x'}] = 0$ for endpoint eigenprojectors, one gets a bound $\big|\mathcal{S}^{[\mathbf{W}]}_n(x, x')\big| \leq C$ *independent of* n $(k = 0)$, uniform in admissible transports.

If, in addition, the wordwise Weyl condition holds with exponent ϑ (Definition 4.376), the geometric control is converted to spectral control by the growth of $|\lambda^{(x)}_{n,[\mathbf{W}]}|$, giving the polynomial estimate $\big|\mathcal{S}^{[\mathbf{W}]}_n(x, x')\big| \leq C' \, (1 + n)^{\vartheta \max\{r,s\}}$, which is (4.167). Word/gauge invariance (Proposition 4.382) makes the constants and the bound independent of the chosen word $\mathbf{W}$. □

Corollary 4.359 (Class relations and constants)**.** *One has the inclusions*

$$\mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,poly}} \;\subseteq\; \mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,der}} \;\subseteq\; \mathfrak{D}^{[\mathbf{W}]}_{\mathrm{admissible}},$$

and the constants in the uniform bound ($k = 0$) may be chosen of the form $C \lesssim C_{\mathrm{sel}} + C_{\mathrm{conn}} \cdot \mathrm{diam}(U) + C_{\mathrm{curv}} \cdot \mathrm{area}(U)$, with $C_{\mathrm{sel}} := \sup \|[\mathbb{D}^{[\mathbf{W}]}, \hat{\mathcal{F}}^{[\mathbf{W}]}_\bullet]\|$, $C_{\mathrm{conn}} := \sup \|[\mathbb{D}^{[\mathbf{W}]}, \mathcal{A}^{[\mathbf{W}]}]\|$, and C_{curv} controlling subleading curvature terms, all taken uniformly over admissible words and over the normal neighbourhood U.

Remark 4.360 (Derived vs. axiomatic admissibility)**.** The classes in Definitions 4.352 and 4.361 differ in hypotheses and in the nature of the smoothness guarantee:

(a) *Hypotheses.* The derived class $\mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,der}}$ assumes self-adjointness and compact resolvent together with (r,s)-regularity of the wordwise data on a normal neighbourhood (Definition 4.373). The axiomatic class $\mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,axiom}}$ assumes self-adjointness and compact resolvent together with the Wordwise Spectral Smoothness bound itself (Definition 4.383).

(b) *Consequences.* In the derived class, spectral smoothness is a theorem (Theorem 4.353): a uniform bound ($k = 0$) holds, and if a wordwise Weyl condition with exponent $\vartheta > 0$ is present (Definition 4.376), then the polynomial form holds with exponent $k = \vartheta \max\{r, s\}$. In the axiomatic class, this bound is postulated rather than proved.

(c) *Constants.* In the derived setting the bound is constructive; the constant C can be expressed in terms of commutators with selectors and connection and geometric parameters of the neighbourhood (Remark 4.355). In the axiomatic setting only existence of such constants is required.

(d) *Class relations.* There are inclusions $\mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,poly}} \subseteq \mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,der}} \subseteq \mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,axiom}}$ (Proposition 4.362). Under standard regularity plus Weyl growth, the three classes coincide (Theorem 4.363).

(e) *Interpretation.* The derived formulation exposes mechanism-level control (selector vs. transport contributions) and parameter dependence of the smoothness constants; the axiomatic formulation offers a backward-compatible, property-level entry point when only the bound is needed.

Admissible Wordwise Spectral Operators (axiomatic).

Definition 4.361 (Admissible Wordwise Spectral Operators — axiomatic)**.** A densely defined operator $\mathbb{D}^{[\mathbf{W}]}$ is *admissible wordwise (axiomatic)* iff:

(i) *Self-adjointness (word-dressed)* as in Definition 4.345(a);

(ii) *Compact resolvent (discrete spectrum)* as in Definition 4.345(b);

(iii) The *Wordwise Spectral Smoothness* bound as stated in Definition 4.346.

We denote this class by $\mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,axiom}}$.

Proposition 4.362 (Class relations)**.** *One has the inclusions*

$$\mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,poly}} \subseteq \mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,der}} \subseteq \mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,axiom}}. \tag{4.168}$$

Moreover, the constants appearing in the spectral smoothness bound of Definition 4.346 can be chosen uniformly in the word $\mathbf{W}$.

Proof sketch. The inclusion $\mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,poly}} \subseteq \mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,der}}$ is by definition. Theorem 4.353 gives $\mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,der}} \subseteq \mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,axiom}}$. Word/gauge invariance (Proposition 4.382) yields uniformity in $\mathbf{W}$. □

Theorem 4.363 (Equivalence under standard regularity)**.** *Assume the wordwise data admit (r,s)-regular representatives on a normal neighbourhood of the diagonal (Definition 4.373) and*

that $\mathbb{D}^{[\mathbf{W}]}$ *satisfies a wordwise Weyl growth condition with exponent* $\vartheta > 0$ *(Definition 4.376). Then*

$$\mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,axiom}} \;=\; \mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,der}} \;=\; \mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,poly}}, \tag{4.169}$$

and the spectral smoothness exponent can be taken as

$$k \;=\; \vartheta\, \max\{r, s\}. \tag{4.170}$$

Proof sketch. By Proposition 4.362 it remains to show the reverse inclusion $\mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,axiom}} \subseteq \mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,poly}}$ under the stated hypotheses. The (r, s)-regularity ensures bounded selector commutators and a Duhamel estimate for transport; combining these with Weyl growth converts the geometric (path-length) control into a polynomial-in-n control with exponent $\vartheta \max\{r, s\}$. Thus an axiomatic element admits the polynomial bound, i.e. belongs to $\mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,poly}}$. Equality of the three classes follows from (4.168). □

Remark 4.364 (Backward compatibility)**.** The axiomatic class recovers the exogenous admissibility (Definition 4.343) when the trivial word is chosen and the wordwise correlator reduces to the exogenous kernel, by Proposition 4.382. Thus existing results remain valid, while the wordwise formulation adds invariance and a clean split between selector- and transport-driven contributions.

Definition 4.365 (Wordwise-coherent family)**.** Let $U_x^{[\mathbf{W}]} : \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)} \to \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ be the word-frame unitary at locus x (Section 4.6.2). A collection $\{\mathbb{D}^{[\mathbf{W}]}\}_{\mathbf{W}}$ of densely defined fibrewise operators is *wordwise-coherent* if, for every pair of words $(\mathbf{W}, \mathbf{W}')$ and each x,

$$\mathbb{D}_x^{[\mathbf{W}']} \;=\; U_x^{[\mathbf{W}\to\mathbf{W}']}\, \mathbb{D}_x^{[\mathbf{W}]} \left(U_x^{[\mathbf{W}\to\mathbf{W}']}\right)^{\dagger}, \qquad U_x^{[\mathbf{W}\to\mathbf{W}']} := U_x^{[\mathbf{W}']} \left(U_x^{[\mathbf{W}]}\right)^{\dagger}.$$

Lemma 4.366 (Word-dressing preserves admissibility)**.** *If* $\hat{D}$ *is* admissible *in the exogenous sense (Definition 4.343), then for any word* $\mathbf{W}$ *the conjugates*

$$\mathbb{D}_x^{[\mathbf{W}]} \;:=\; U_x^{[\mathbf{W}]}\, \hat{D}_x \left(U_x^{[\mathbf{W}]}\right)^{\dagger}$$

constitute an admissible wordwise *operator (Definition 4.347).*

Proof sketch. (i) *Self-adjointness*: bounded unitary conjugation preserves self-adjointness and dense domains: $\left(\mathbb{D}_x^{[\mathbf{W}]}\right)^{\dagger} = U_x^{[\mathbf{W}]} \hat{D}_x \left(U_x^{[\mathbf{W}]}\right)^{\dagger}$.

(ii) *Compact resolvent & spectrum*: $(\mathbb{D}_x^{[\mathbf{W}]} - z)^{-1} = U_x^{[\mathbf{W}]} (\hat{D}_x - z)^{-1} \left(U_x^{[\mathbf{W}]}\right)^{\dagger}$ is compact whenever $(\hat{D}_x - z)^{-1}$ is; eigenpairs transform as $\lambda^{(x)}_{n,[\mathbf{W}]} = \lambda^{(x)}_n$, $|\psi^{(x)}_{n,[\mathbf{W}]}\rangle = U_x^{[\mathbf{W}]} |\psi^{(x)}_n\rangle$.

(iii) *Spectral smoothness*: write bilocal letters/words as $\hat{G}_\Lambda^{[\mathbf{W}]} = \mathbf{U}^{[\mathbf{W}]} \hat{G}_\Lambda (\mathbf{U}^{[\mathbf{W}]})^{\dagger}$ with $\mathbf{U}^{[\mathbf{W}]}$ the natural endpoint lift. Endogenous charges (the $\mathbb{Z}_2^2$ label) are preserved by word changes, so the super-bracket respects conjugation: $[\,\mathbf{U}AU^{\dagger}, \mathbf{U}BU^{\dagger}\,]_{\mathrm{endo}} = \mathbf{U}[\,A, B\,]_{\mathrm{endo}}\mathbf{U}^{\dagger}$ (Section 4.3.1). Hence

$$\left|\mathcal{S}_n^{[\mathbf{W}]}(x, x')\right| = \left|\langle\Omega|[\,\mathbb{D}^{[\mathbf{W}]}, \hat{G}_\Lambda^{[\mathbf{W}]}(\cdot)\,]_{\mathrm{endo}}|\Omega\rangle\right| \le \left\|[\,\hat{D}, \hat{G}_\Lambda(\cdot)\,]_{\mathrm{endo}}\right\| \le C(1 + |n|)^k,$$

using unitary invariance of operator norm and the exogenous bound (Definition 4.342). This is uniform over admissible wordwise transports. □

Lemma 4.367 (Wordwise-to-exogenous admissibility)**.** *Let* $\{\mathbb{D}^{[\mathbf{W}]}\}_{\mathbf{W}}$ *be a wordwise-coherent family (Definition 4.365) that is admissible wordwise (Definition 4.347) for at least one word* $\mathbf{W}_0$*. Define*

$$\hat{D}_x \;:=\; \left(U_x^{[\mathbf{W}_0]}\right)^{\dagger} \mathbb{D}_x^{[\mathbf{W}_0]}\, U_x^{[\mathbf{W}_0]}.$$

Then $\hat{D}$ *is an* admissible *spectral operator in the exogenous sense (Definition 4.343).*

Proof sketch. Each clause is the converse of Lemma 4.366: self-adjointness and compact resolvent descend by unitary conjugation; eigenpairs are pulled back; the spectral-smoothness bound follows from the wordwise bound and unitary invariance of norms/charges. Wordwise coherence ensures independence of the chosen reference word $\mathbf{W}_0$. □

Theorem 4.368 (Equivalence of admissible classes)**.** *Conjugation by word-frames induces a bijection*

$$\frac{\mathfrak{D}_{\text{admissible}}}{\textit{unitary eq.}} \;\longleftrightarrow\; \frac{\{\textit{wordwise-coherent families } \{\mathbb{D}^{[\mathbf{W}]}\}_{\mathbf{W}} \textit{ with } \mathbb{D}^{[\mathbf{W}]} \in \mathfrak{D}^{[\mathbf{W}]}_{\text{admissible}}\}}{\textit{word changes}},$$

natural with respect to changes of word (Remark 4.348).

Proof of Theorem 4.368. Define two maps:

$$\begin{aligned} \mathcal{D} :\; & \frac{\mathfrak{D}_{\text{admissible}}}{\text{unitary eq.}} \;\longrightarrow\; \frac{\left\{\begin{smallmatrix}\text{wordwise-coherent}\\ \text{admissible families}\end{smallmatrix}\right\}}{\text{word changes}}, \\ \mathcal{U} :\; & \frac{\left\{\begin{smallmatrix}\text{wordwise-coherent}\\ \text{admissible families}\end{smallmatrix}\right\}}{\text{word changes}} \;\longrightarrow\; \frac{\mathfrak{D}_{\text{admissible}}}{\text{unitary eq.}}. \end{aligned} \tag{4.171}$$

and show that $\mathcal{D}$ and $\mathcal{U}$ are well-defined inverse bijections, natural with respect to word changes.

Step 1 (Definition of $\mathcal{D}$*; well-definedness).* Given a class $[\hat{D}] \in \mathfrak{D}_{\text{admissible}}/\text{unitary eq.}$, fix any system of word-frames $\{U_x^{[\mathbf{W}]}\}_{\mathbf{W}}$ (charge-preserving fibre unitaries, Section 4.6.2) and set

$$\mathbb{D}_x^{[\mathbf{W}]} \;:=\; U_x^{[\mathbf{W}]}\, \hat{D}_x \left(U_x^{[\mathbf{W}]}\right)^{\dagger}. \tag{4.172}$$

By Lemma 4.366, each $\mathbb{D}^{[\mathbf{W}]}$ is admissible wordwise and the family is wordwise-coherent (Definition 4.365); hence $\mathcal{D}([\hat{D}])$ is the word-change class of $\{\mathbb{D}^{[\mathbf{W}]}\}_{\mathbf{W}}$.

Independence of choices: (i) If a different representative $\hat{D}' = V\,\hat{D}\,V^{\dagger}$ (fibrewise unitary V) is taken, then with the *modified* word-frames $\widetilde{U}_x^{[\mathbf{W}]} := U_x^{[\mathbf{W}]} V_x^{\dagger}$,

$$\widetilde{U}_x^{[\mathbf{W}]}\, \hat{D}'_x \left(\widetilde{U}_x^{[\mathbf{W}]}\right)^{\dagger} = U_x^{[\mathbf{W}]}\, \hat{D}_x \left(U_x^{[\mathbf{W}]}\right)^{\dagger},$$

so the produced family is unchanged. (ii) If word-frames are changed by a word-change cocycle $U_x^{[\mathbf{W}]} \mapsto U_x^{[\mathbf{W}']} = U_x^{[\mathbf{W}\to\mathbf{W}']}\, U_x^{[\mathbf{W}]}$, then

$$\mathbb{D}_x^{[\mathbf{W}']} = U_x^{[\mathbf{W}\to\mathbf{W}']}\, \mathbb{D}_x^{[\mathbf{W}]} \left(U_x^{[\mathbf{W}\to\mathbf{W}']}\right)^{\dagger},$$

which is exactly the quotient relation by word changes. Thus $\mathcal{D}$ is well-defined.

Step 2 (Definition of $\mathcal{U}$; well-definedness). Let $\{\mathbb{D}^{[\mathbf{W}]}\}_{\mathbf{W}}$ be a wordwise-coherent admissible family (Definitions 4.347 and 4.365). Fix a word $\mathbf{W}_0$ and define the exogenous operator by *undressing*

$$\hat{D}_x \;:=\; \left(U_x^{[\mathbf{W}_0]}\right)^{\dagger} \mathbb{D}_x^{[\mathbf{W}_0]}\, U_x^{[\mathbf{W}_0]}. \tag{4.173}$$

By Lemma 4.367, $\hat{D}$ is admissible in the exogenous sense. If a different representative of the family related by word changes is chosen, or if a different base word $\mathbf{W}_0'$ is used, the resulting $\hat{D}$ changes by a fibrewise unitary conjugation; hence $\mathcal{U}$ is well-defined on the quotient by word changes with values in $\mathfrak{D}_{\text{admissible}}/\text{unitary eq}$.

Step 3 (Mutual inverses). For $[\hat{D}]$, the family produced by (4.172) satisfies, at the chosen base word $\mathbf{W}_0$,

$$\left(U_x^{[\mathbf{W}_0]}\right)^{\dagger} \mathbb{D}_x^{[\mathbf{W}_0]}\, U_x^{[\mathbf{W}_0]} = \hat{D}_x,$$

so $\mathcal{U}(\mathcal{D}([\hat{D}])) = [\hat{D}]$. Conversely, for a wordwise-coherent family, wordwise coherence gives $\mathbb{D}_x^{[\mathbf{W}]} = U_x^{[\mathbf{W}]}\, \hat{D}_x \left(U_x^{[\mathbf{W}]}\right)^{\dagger}$ with $\hat{D}$ defined by (4.173); therefore $\mathcal{D}(\mathcal{U}([\{\mathbb{D}^{[\mathbf{W}]}\}_{\mathbf{W}}])) = [\{\mathbb{D}^{[\mathbf{W}]}\}_{\mathbf{W}}]$ in the quotient by word changes. Hence $\mathcal{D}$ and $\mathcal{U}$ are inverse bijections.

Step 4 (Naturality). Let $\mathbf{W} \mapsto \mathbf{W}'$ be a word change with conjugation $U_x^{[\mathbf{W}\to\mathbf{W}']}$. Then for any $[\hat{D}]$,

$$\mathcal{D}([\hat{D}]) \;\xrightarrow{\mathbf{W}\to\mathbf{W}'}\; \{\mathbb{D}_x^{[\mathbf{W}']}\}_{\mathbf{W}} = \left\{U_x^{[\mathbf{W}\to\mathbf{W}']}\, \mathbb{D}_x^{[\mathbf{W}]} \left(U_x^{[\mathbf{W}\to\mathbf{W}']}\right)^{\dagger}\right\}_{\mathbf{W}'},$$

while $\mathcal{U}$ composed with this change yields $\hat{D}'_x = (U_x^{[\mathbf{W}_0']})^{\dagger} \mathbb{D}_x^{[\mathbf{W}_0']} U_x^{[\mathbf{W}_0']}$ which equals $\hat{D}_x$ up to fibrewise unitary conjugation, since $\mathbf{W}_0' = \mathbf{W}_0$ composed with a word change. Thus the square

$$\begin{array}{ccc} \mathfrak{D}_{\text{admissible}}/\text{unitary eq.} & \xrightarrow{\;\mathcal{D}\;} & \{\text{wordwise families}\}/\text{word changes} \\ \Big\downarrow{\scriptstyle \text{id}} & & \Big\downarrow{\scriptstyle \text{word change}} \\ \mathfrak{D}_{\text{admissible}}/\text{unitary eq.} & \xrightarrow{\;\mathcal{D}\;} & \{\text{wordwise families}\}/\text{word changes} \end{array}$$

commutes up to the canonical identifications; i.e. the bijection is natural with respect to word changes (Remark 4.348).

Step 5 (Admissibility is preserved). Both directions preserve admissibility: $\mathcal{D}$ by Lemma 4.366 and the endogenous conjugation identity (Remark 4.372); $\mathcal{U}$ by Lemma 4.367. This also ensures the spectral-smoothness exponents and constants are word-invariant (Corollary 4.369).

All claims in the theorem follow. □

Corollary 4.369 (Invariants under word changes)**.** *For any word* $\mathbf{W}$ *and locus* x*:* $\operatorname{spec}(\mathbb{D}_x^{[\mathbf{W}]}) = \operatorname{spec}(\hat{D}_x)$ *(with multiplicities), and the optimal spectral-smoothness exponents* (C, k) *may be chosen* identically *for* $\mathcal{S}_n$ *and* $\mathcal{S}_n^{[\mathbf{W}]}$*.*

Theorem 4.370 (Inclusions and conditional equivalence)**.** *There are unconditional inclusions*

$$\mathfrak{D}^{[\mathbf{W}]}_{\text{adm,poly}} \;\subseteq\; \mathfrak{D}^{[\mathbf{W}]}_{\text{adm,der}} \;\subseteq\; \mathfrak{D}^{[\mathbf{W}]}_{\text{adm,axiom}}.$$

Moreover, if the wordwise data admit (r, s)*-regular representatives on a normal neighbourhood of the diagonal (Definition 4.373) and* $\mathbb{D}^{[\mathbf{W}]}$ *satisfies wordwise Weyl growth with exponent* $\vartheta > 0$ *(Definition 4.376), then*

$$\mathfrak{D}^{[\mathbf{W}]}_{\text{adm,axiom}} \;=\; \mathfrak{D}^{[\mathbf{W}]}_{\text{adm,der}} \;=\; \mathfrak{D}^{[\mathbf{W}]}_{\text{adm,poly}}, \tag{4.174}$$

and the Wordwise Spectral Smoothness bound holds with exponent

$$k \;=\; \vartheta\, \max\{r, s\}, \tag{4.175}$$

uniform under admissible wordwise transports and independent of the chosen word $\mathbf{W}$.

Proof sketch. The inclusion chain follows from the definitions and Theorem 4.353 and Proposition 4.362: $\mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,poly}} \subseteq \mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,der}}$ by definition, and $\mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,der}} \subseteq \mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,axiom}}$ since derived admissibility implies spectral smoothness. Under (r, s)-regularity and wordwise Weyl growth, geometric (path-length) control and bounded selector/connection commutators upgrade to a polynomial-in-n shear bound with exponent $\vartheta \max\{r, s\}$ (Theorem 4.353 and Proposition 4.358), which shows that any axiomatic element lies in $\mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,poly}}$; together with the inclusions this yields (4.174). Word/gauge invariance gives the stated uniformity and word independence. □

Remark 4.371 (Portability and invariance)**.** The classes $\mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,der}}$ and $\mathfrak{D}^{[\mathbf{W}]}_{\mathrm{adm,poly}}$ are invariant under charge-preserving word changes (frame conjugations) by Remarks 4.348 and 4.372, and the constants in (4.175) may be chosen uniformly across words.

Remark 4.372 (Endogenous vs. ordinary commutators; conjugation identity)**.** Let $[\cdot,\cdot]_{\mathrm{endo}}$ be the endogenous super-commutator from Section 4.3.1, with charges $q(\cdot) \in \mathbb{Z}_2^2$ induced by the holosymmetric involutions. Then:

(a) **Reduction rules.** For any words/letters A, B,

$$[A, B]_{\mathrm{endo}} = AB - (-1)^{\langle q(A), q(B)\rangle} BA, \quad \langle\cdot,\cdot\rangle : \mathbb{Z}_2^2 \times \mathbb{Z}_2^2 \to \mathbb{Z}_2.$$

In particular:

$$\begin{aligned} &\text{(even case)} \quad q(A) = 0 \text{ or } q(B) = 0 \;\Rightarrow\; [A, B]_{\mathrm{endo}} = [A, B], \\ &\text{(odd pairing)} \quad \langle q(A), q(B)\rangle = 1 \;\Rightarrow\; [A, B]_{\mathrm{endo}} = \{A, B\}. \end{aligned}$$

Writing $A = \sum_{\alpha\in\mathbb{Z}_2^2} A^{(\alpha)}$ into charge components gives the block formula

$$[A, B]_{\mathrm{endo}} = \sum_{\alpha,\beta\in\mathbb{Z}_2^2} \Big(A^{(\alpha)} B^{(\beta)} - (-1)^{\langle\alpha,\beta\rangle} B^{(\beta)} A^{(\alpha)} \Big).$$

(b) **Charge-preserving conjugations act by automorphisms.** Let U be a *charge-preserving* unitary (i.e. it intertwines the charge involutions; equivalently $[U, Z] = [U, \sigma] = 0$ in the usual models, cf. Section 4.6.2). Then for all A, B,

$$\big[\, UAU^\dagger,\; UBU^\dagger \,\big]_{\mathrm{endo}} \;=\; U\,[A, B]_{\mathrm{endo}}\,U^\dagger. \tag{4.176}$$

Proof sketch. Since $q(UAU^\dagger) = q(A)$ and $q(UBU^\dagger) = q(B)$, the sign $(-1)^{\langle q(A), q(B)\rangle}$ is invariant; expand the definition and factor $U(\cdot)U^\dagger$.

(c) **Word-change identity for spectral shear.** For the wordwise shear $\mathcal{S}_n^{[\mathbf{W}]}(x,x')$ from (4.162), a change of word $\mathbf{W} \to \mathbf{W}'$ implemented by a charge-preserving $U^{[\mathbf{W}\to\mathbf{W}']}$ (Section 4.6.2) yields

$$\begin{aligned}\mathcal{S}_n^{[\mathbf{W}']}(x,x') &= \Big\langle \Omega \Big| \big[\, \mathbb{D}^{[\mathbf{W}']},\ \hat{G}_\Lambda^{[\mathbf{W}']}(\cdot)\,\big]_{\mathrm{endo}} \Big| \Omega \Big\rangle \\ &= \Big\langle \Omega \Big| U^\dagger \big[\, \mathbb{D}^{[\mathbf{W}]},\ \hat{G}_\Lambda^{[\mathbf{W}]}(\cdot)\,\big]_{\mathrm{endo}}\, U \Big| \Omega \Big\rangle \\ &= \mathcal{S}_n^{[\mathbf{W}]}(x,x'). \end{aligned} \tag{4.177}$$

using (4.176) and unitary invariance of expectations. Hence the shear and any derived bounds (e.g. spectral smoothness exponents (C,k)) are *word-invariants*.

Definition 4.373 (Wordwise (r,s)-regularity relative to $\mathbb{D}^{[\mathbf{W}]}$)**.** Let $\mathbb{D}^{[\mathbf{W}]}$ be a wordwise spectral operator on the modal bundle (Definition 4.345). We say the wordwise data are (r,s)-*regular relative to* $\mathbb{D}^{[\mathbf{W}]}$ if there exist exponents $r,s \geq 0$ and finite constants $L_{\mathrm{sel}}, L_{\mathrm{conn}}$ such that:

(a) **Selector regularity (degree r).** For every mode selector $\hat{\mathcal{F}}_\bullet^{[\mathbf{W}]}$,

$$\Big\| (1+|\mathbb{D}^{[\mathbf{W}]}|)^{-r}\, [\mathbb{D}^{[\mathbf{W}]}, \hat{\mathcal{F}}_\bullet^{[\mathbf{W}]}]_{\mathrm{endo}} \Big\| \ \leq\ L_{\mathrm{sel}}.$$

(b) **Connection regularity (degree s).** For the wordwise connection components $\mathcal{A}_\mu^{[\mathbf{W}]}$ along admissible paths (Section 4.5),

$$\Big\| (1+|\mathbb{D}^{[\mathbf{W}]}|)^{-s}\, [\mathbb{D}^{[\mathbf{W}]}, \mathcal{A}_\mu^{[\mathbf{W}]}]_{.endo} \Big\| \ \leq\ L_{\mathrm{conn}}$$

uniformly in a normal neighbourhood of the diagonal. (4.178)

Lemma 4.374 (Eigenmode-weighted commutator bounds)**.** *Let* $|\psi_{n,[\mathbf{W}]}^{(x)}\rangle$ *be an eigenmode of* $\mathbb{D}_x^{[\mathbf{W}]}$ *with eigenvalue* $\lambda_{n,[\mathbf{W}]}^{(x)}$*. Under Definition 4.373,*

$$\begin{aligned} \Big\| [\mathbb{D}^{[\mathbf{W}]}, \hat{\mathcal{F}}_\bullet^{[\mathbf{W}]}]_{\mathrm{endo}}\, |\psi_{n,[\mathbf{W}]}^{(x)}\rangle \Big\| &\ \leq\ L_{\mathrm{sel}}\, (1+|\lambda_{n,[\mathbf{W}]}^{(x)}|)^r, \\ \Big\| [\mathbb{D}^{[\mathbf{W}]}, \mathcal{A}_\mu^{[\mathbf{W}]}]_{\mathrm{endo}}\, |\psi_{n,[\mathbf{W}]}^{(x)}\rangle \Big\| &\ \leq\ L_{\mathrm{conn}}\, (1+|\lambda_{n,[\mathbf{W}]}^{(x)}|)^s. \end{aligned} \tag{4.179}$$

Proof. Apply the definitions to the unit vector $(1+|\mathbb{D}^{[\mathbf{W}]}|)^{-r}|\psi_{n,[\mathbf{W}]}\rangle$ and use $(1+|\mathbb{D}^{[\mathbf{W}]}|)^r|\psi_{n,[\mathbf{W}]}\rangle = (1+|\lambda_{n,[\mathbf{W}]}|)^r|\psi_{n,[\mathbf{W}]}\rangle$, and similarly for s. □

Theorem 4.375 (Polynomial Wordwise Spectral Smoothness)**.** *Assume the hypotheses of Sections 4.2, 4.3.1, 4.5, 4.6.2 and 4.6.3 and (r,s)-regularity (Definition 4.373). Then there exist a neighbourhood $U \subset \mathcal{X}_q \times \mathcal{X}_q$ of the diagonal and a constant $C \geq 0$ such that, for all $x,x' \in U$ and all eigenmodes $|\psi_{n,[\mathbf{W}]}^{(x)}\rangle$ of $\mathbb{D}_x^{[\mathbf{W}]}$,*

$$\Big| \mathcal{S}_n^{[\mathbf{W}]}(x,x') \Big| \ \leq\ C\,(1+|\lambda_{n,[\mathbf{W}]}^{(x)}|)^{k_0}, \qquad k_0 := \max\{r,s\}.$$

Proof. Replicate the graded-Leibniz expansion in the proof of Theorem 4.349 (the uniform case $k = 0$), but now retain the λ-dependence via Lemma 4.374. For x, x' within a normal neighbourhood, the Duhamel/variation-of-constants estimate for the transport U_γ along a short path γ with length $\ell = d(x, x')$ gives

$$\left\| [\mathbb{D}^{[\mathbf{W}]}, U_\gamma]_{\mathrm{endo}} |\psi_{n,[\mathbf{W}]}\rangle \right| \;\le\; \Big(L_{\mathrm{conn}}\, \ell + O(\ell^2) \Big)\, (1 + |\lambda_{n,[\mathbf{W}]}|)^s.$$

Selector terms contribute $L_{\mathrm{sel}}\,(1 + |\lambda_{n,[\mathbf{W}]}|)^r$ on each side of U_γ. Using that selectors are contractions, U_γ is unitary, $P_{n,x'}$ commutes with $\mathbb{D}^{[\mathbf{W}]}_{x'}$, and the expectation is dominated by the operator norm, we obtain

$$\left| \mathcal{S}^{[\mathbf{W}]}_n(x, x') \right| \;\le\; C_0\,(1 + |\lambda_{n,[\mathbf{W}]}|)^r \;+\; C_1\,\ell\,(1 + |\lambda_{n,[\mathbf{W}]}|)^s \;+\; O(\ell^2)\,(1 + |\lambda_{n,[\mathbf{W}]}|)^s.$$

Shrink U so that $\ell \le \ell_0 \ll 1$ and absorb the linear/quadratic ℓ factors into a constant; the exponent is $k_0 = \max\{r, s\}$. □

Definition 4.376 (Wordwise Weyl growth)**.** We say $\mathbb{D}^{[\mathbf{W}]}$ satisfies *Weyl growth with exponent* $\vartheta > 0$ if there exist $a, b > 0$ and $n_0 \in \mathbb{N}$ such that, for all $n \ge n_0$,

$$a\, n^\vartheta \;\le\; 1 + |\lambda^{(x)}_{n,[\mathbf{W}]}| \;\le\; b\, n^\vartheta \quad \text{uniformly for } x \text{ in compact sets.}$$

Corollary 4.377 (Polynomial in the mode index)**.** *Under Theorem 4.375 and Definition 4.376 with exponent ϑ, there exist constants $C', k \ge 0$ and a neighbourhood U of the diagonal such that, for all $x, x' \in U$ and $n \ge n_0$,*

$$\left| \mathcal{S}^{[\mathbf{W}]}_n(x, x') \right| \;\le\; C'\,(1 + n)^k, \qquad k \;:=\; k_0\,\vartheta \;=\; \vartheta\, \max\{r, s\}.$$

For finitely many small indices $n < n_0$ the bound can be absorbed into C'.

Remark 4.378 (Choices and special cases)**.** (i) If the selectors are *zeroth order* ($r = 0$) and the connection is *first order* ($s = 1$), then $k_0 = 1$ and $k = \vartheta$.

(ii) If both are zeroth order ($r = s = 0$), we recover the uniform bound $k = 0$ (Theorem 4.349).

(iii) The result is *word-invariant*: changing words via a charge-preserving unitary (Section 4.6.2) preserves (r, s)-regularity constants and the exponents, by Remark 4.372.

4.9.5 Wordwise Reformulation of Operator-Theoretic Intrinsic Distances

Idea in one line. Compute everything through the *word–dressed* bilocal kernel $\hat{G}^{[\mathbf{W}]}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})$ (cf. (4.132)) and evaluate pairings via the Holo-State. Covariance under charge-preserving word changes and endpoint gauge (Remarks 4.348 and 4.372) guarantees that the resulting distances and spectral response functionals are *intrinsic* (word/gauge independent).

Definition 4.379 (Wordwise correlator amplitude, normalisation, and distance)**.** Fix a word $\mathbf{W}$ and test vectors $|\psi_1\rangle_x \in \mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}}$, $|\psi_2\rangle_{x'} \in \mathcal{H}^{(\mathfrak{B}_{x'})}_{\mathrm{mod}}$. Define the scalar amplitude via the wordwise bilocal kernel:

$$\mathsf{G}^{[\mathbf{W}]}(x, x'; \psi_1, \psi_2) := \left\langle \mathring{\mathbf{\Psi}} \right| \hat{G}^{[\mathbf{W}]}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2] \left| \mathring{\mathbf{\Psi}} \right\rangle. \tag{4.180}$$

With coincident-limit normalisations $N^{[\mathbf{W}]}(\psi_1, x) := \sqrt{\mathsf{G}^{[\mathbf{W}]}(x, x; \psi_1, \psi_1)}$ and $N^{[\mathbf{W}]}(\psi_2, x') := \sqrt{\mathsf{G}^{[\mathbf{W}]}(x', x'; \psi_2, \psi_2)}$, set the *wordwise similarity* and *distance*

$$C^{[\mathbf{W}]}(x, x'; \psi_1, \psi_2) := \frac{\left|\mathsf{G}^{[\mathbf{W}]}(x, x'; \psi_1, \psi_2)\right|}{N^{[\mathbf{W}]}(\psi_1, x)\, N^{[\mathbf{W}]}(\psi_2, x')} \in [0, 1], \qquad d^{[\mathbf{W}]}_{\mathrm{corr}} := \sqrt{1 - C^{[\mathbf{W}]}} \in [0, 1]. \tag{4.181}$$

Remark 4.380 (Intrinsic nature and reductions)**.** (i) By Remarks 4.348 and 4.372, both $C^{[\mathbf{W}]}$ and $d^{[\mathbf{W}]}_{\mathrm{corr}}$ are invariant under endpoint gauge and under word changes $\mathbf{W} \to \mathbf{W}'$; hence they define a unique, intrinsic correlator distance on the modal bundle.

(ii) In the Twofold (duality-collapse) slice, compression by the word-rotated isometry $U^{[\mathbf{W}]} : \mathbb{C}^2 \hookrightarrow \mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}}$ (Section 4.6.2) reproduces the standard qubit overlap distance on $\mathbb{C}^2$.

(iii) One may extremise over admissible probes to obtain a probe-independent $d_{\mathrm{corr}}(x, x') := \inf_{\psi_1, \psi_2} d^{[\mathbf{W}]}_{\mathrm{corr}}(x, x'; \psi_1, \psi_2)$; invariance ensures the extremum is word/gauge independent.

Definition 4.381 (Wordwise spectral operator and spectral shear)**.** Let $\mathbb{D}^{[\mathbf{W}]}$ be a *wordwise spectral operator* as in Definition 4.345, and let $\mathbb{D}^{[\mathbf{W}]}_x |\psi^{(x)}_{n,[\mathbf{W}]}\rangle = \lambda^{(x)}_{n,[\mathbf{W}]} |\psi^{(x)}_{n,[\mathbf{W}]}\rangle$. Using the endogenous super-commutator $[\cdot, \cdot]_{\mathrm{endo}}$ (Section 4.3.1), define the *wordwise spectral shear*

$$\mathcal{S}^{[\mathbf{W}]}_n(x, x') \;:=\; \Big\langle \mathring{\mathbf{\Psi}} \Big| \Big[\mathbb{D}^{[\mathbf{W}]},\; \hat{G}^{[\mathbf{W}]}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi^*_{n,[\mathbf{W}]}, \psi_{n,[\mathbf{W}]}] \Big]_{\mathrm{endo}} \Big| \mathring{\mathbf{\Psi}} \Big\rangle. \tag{4.182}$$

Proposition 4.382 (Word invariance of admissibility)**.**

(a) Wordwise invariance
Let $\mathbb{D}^{[\mathbf{W}]}$ be a wordwise spectral operator and let $\mathcal{S}^{[\mathbf{W}]}_n(x, x')$ be the wordwise spectral shear (Definition 4.345 and eq. (4.162)). If there exist a neighbourhood U of the diagonal, constants $C, k \geq 0$, and a word $\mathbf{W}_0$ such that

$$\left|\mathcal{S}^{[\mathbf{W}_0]}_n(x, x')\right| \;\leq\; C\,(1 + |n|)^k \quad \text{for all } (x, x') \in U \text{ and all } n,$$

then the same bound holds for every *charge-preserving word $\mathbf{W}$ with the* same *constants C, k:*

$$\left|\mathcal{S}^{[\mathbf{W}]}_n(x, x')\right| \;\leq\; C\,(1 + |n|)^k \qquad \forall\, \mathbf{W}.$$

(b) Equivalence with the exogenous clause
Under the wordwise–exogenous correspondence of Lemmas 4.366 and 4.367, the above wordwise bound holds for some (hence every) word $\mathbf{W}$ iff *the exogenous shear bound (Definition 4.342) holds for $\hat{D}$ in the admissible class $\mathfrak{D}_{\mathrm{admissible}}$. In particular, the same neighbourhood U and the same exponents (C, k) can be used on both sides.*

Proof. (a) *Wordwise invariance.* Let $\mathbf{W} \to \mathbf{W}'$ be a change of word implemented by charge-preserving endpoint unitaries $U_x, U_{x'}$ (cf. Section 4.6.2). The bilocal word kernel transforms by endpoint conjugation (passive letter change):

$$\hat{G}^{[\mathbf{W}']}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\phi_1^*, \phi_2] \;=\; U_x^\dagger\, \hat{G}^{[\mathbf{W}]}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[(U_x \phi_1)^*,\, U_{x'} \phi_2]\, U_{x'}.$$

Using the endogenous conjugation identity for the super-commutator (Remark 4.372, eq. (4.176)), we have

$$\left[U\mathbb{D}^{[\mathbf{W}]}U^\dagger,\ U\hat{G}_\Lambda^{[\mathbf{W}]}U^\dagger\right]_{\text{endo}} \;=\; U\left[\mathbb{D}^{[\mathbf{W}]},\ \hat{G}_\Lambda^{[\mathbf{W}]}\right]_{\text{endo}}U^\dagger.$$

Taking Holo-State expectations and using unitary invariance of the scalar (again Remark 4.372), the outer unitaries drop, so the shear is unchanged:

$$\mathcal{S}_n^{[\mathbf{W}']}(x,x') \;=\; \mathcal{S}_n^{[\mathbf{W}]}(x,x').$$

Therefore any bound for $\mathbf{W}_0$ propagates to every word $\mathbf{W}$ with the *same* constants C,k (on the same neighbourhood U).

(b) *Equivalence with the exogenous clause.* If $\hat{D}\in\mathfrak{D}_{\text{admissible}}$ (exogenous sense), Lemma 4.366 yields an admissible wordwise family $\{\mathbb{D}^{[\mathbf{W}]}\}_{\mathbf{W}}$, and part (a) shows the bound holds for all words with the same (C,k). Conversely, if the wordwise bound holds for some word $\mathbf{W}$, pick the trivial word so that $\hat{G}_\Lambda^{[\mathbf{W}]}=\hat{G}_\Lambda$ and $\mathbb{D}^{[\mathbf{W}]}=\hat{D}$ by Lemma 4.367; this recovers exactly the exogenous admissible bound. □

Definition 4.383 (Wordwise spectral smoothness)**.** Let $\{\mathbb{D}^{[\mathbf{W}]}\}_{\mathbf{W}}$ be a family of *wordwise spectral operators* in the sense of Definition 4.345. For each locus x and eigenmode $\mathbb{D}_x^{[\mathbf{W}]}|\psi_{n,[\mathbf{W}]}^{(x)}\rangle=\lambda_{n,[\mathbf{W}]}^{(x)}|\psi_{n,[\mathbf{W}]}^{(x)}\rangle$, let the *wordwise spectral shear* be

$$\mathcal{S}_n^{[\mathbf{W}]}(x,x') := \left\langle \mathring{\mathbf{\Psi}}\right|\left[\mathbb{D}^{[\mathbf{W}]},\ \hat{G}_\Lambda^{[\mathbf{W}]}(\mathfrak{B}_x,\mathfrak{B}_{x'})[\psi^*_{n,[\mathbf{W}]},\psi_{n,[\mathbf{W}]}]\right]_{\text{endo}}\left|\mathring{\mathbf{\Psi}}\right\rangle \tag{4.183}$$

cf. Definition 4.345 and eq. (4.162). (*cf.* Definition 4.345 and eq. (4.162)).

We say the family is *wordwise spectrally smooth* if there exist constants $C,k\ge 0$ and a neighbourhood $U\subset\mathcal{X}_q\times\mathcal{X}_q$ of the diagonal such that, for *every* charge-preserving word $\mathbf{W}$, for all $(x,x')\in U$ and all n,

$$\left|\mathcal{S}_n^{[\mathbf{W}]}(x,x')\right| \;\le\; C\left(1+|n|\right)^k, \tag{4.184}$$

where the bound is *uniform* over admissible wordwise parallel transports used in $\hat{G}_\Lambda^{[\mathbf{W}]}$ (cf. Sections 4.5 and 4.6.3) and *independent* of the chosen word $\mathbf{W}$ (by the charge-preserving conjugation invariance of the endogenous bracket; see Remarks 4.348 and 4.372). Equivalently,

$$\sup_{\mathbf{W}}\ \sup_{(x,x')\in U}\ \sup_{n}\ \frac{\left|\mathcal{S}_n^{[\mathbf{W}]}(x,x')\right|}{(1+|n|)^k} \;<\; \infty.$$

Theorem 4.384 (Wordwise spectral smoothness (uniform and polynomial forms))**.** *Under the wordwise hypotheses of Sections 4.2, 4.3.1, 4.5, 4.6.2 and 4.6.3:*

(a) Uniform form

There exist U (a neighbourhood of the diagonal) and $C\ge 0$ such that $\left|\mathcal{S}_n^{[\mathbf{W}]}(x,x')\right|\le C$ for all $x,x'\in U$ and all n (Theorem 4.349).

(b) Polynomial form

If the data are (r,s)-regular relative to $\mathbb{D}^{[\mathbf{W}]}$ (Definition 4.373), then $\left|\mathcal{S}_n^{[\mathbf{W}]}(x,x')\right| \le C(1+|\lambda_{n,[\mathbf{W}]}^{(x)}|)^{k_0}$ with $k_0 = \max\{r,s\}$ (Theorem 4.375). If in addition a wordwise Weyl growth holds with exponent $\vartheta > 0$ (Definition 4.376), then $\left|\mathcal{S}_n^{[\mathbf{W}]}(x,x')\right| \le C'(1+n)^k$ with $k = \vartheta \max\{r,s\}$ (Corollary 4.377).

All constants and neighbourhoods are word/gauge invariant *by Remarks 4.348 and 4.372.*

Proposition 4.385 (Wordwise invariance of distances and shear bounds)**.** *For any two words $\mathbf{W}, \mathbf{W}'$ related by a charge-preserving frame change (Section 4.6.2), one has*

$$C^{[\mathbf{W}]}(x,x';\psi_1,\psi_2) = C^{[\mathbf{W}']}(x,x';\psi_1',\psi_2'), \quad d_{\mathrm{corr}}^{[\mathbf{W}]} = d_{\mathrm{corr}}^{[\mathbf{W}']}, \quad \mathcal{S}_n^{[\mathbf{W}]}(x,x') = \mathcal{S}_n^{[\mathbf{W}']}(x,x'),$$

with $\psi_j' = U_{[\mathbf{W}',\mathbf{W}]}^{\dagger}\psi_j$ the rotated probes at their loci. Consequently, the smoothness bounds in Theorem 4.384 are independent of the chosen word and of endpoint gauge.

Example 1 — Finite fibre, flat transport (uniform bound $k = 0$)

Setup. Two loci x, x'. Each modal fibre is $\mathbb{C}^4$. Fix any word $\mathbf{W}$. Let the wordwise connection vanish locally: $\mathcal{A}^{[\mathbf{W}]} = 0$ so $U_\gamma = \mathbb{I}$ along short paths.

Wordwise data. Selectors $\{\hat{\mathcal{F}}_\alpha^{[\mathbf{W}]}\}_{\alpha\in\{0,1,p,s\}}$ are bounded. Choose a wordwise spectral operator diagonal in some frame,

$$\mathbb{D}_x^{[\mathbf{W}]} = \mathrm{diag}(d_1,d_2,d_3,d_4), \qquad \mathbb{D}_{x'}^{[\mathbf{W}]} = \mathrm{diag}(d_1',d_2',d_3',d_4').$$

Shear computation. With $U_\gamma = \mathbb{I}$ and rank-one $P_{n,x'} := |\psi_{n,[\mathbf{W}]}^{(x')}\rangle\langle\psi_{n,[\mathbf{W}]}^{(x')}|$,

$$\begin{aligned}\mathcal{S}_n^{[\mathbf{W}]}(x,x') &= \Big\langle \Omega \Big| \Big[\mathbb{D}^{[\mathbf{W}]},\ \hat{\mathcal{F}}_\alpha^\dagger\,\hat{\mathcal{F}}_\beta\,P_{n,x'}\Big]_{\mathrm{endo}} \Big|\Omega\Big\rangle \\ &= \Big\langle \Omega \Big| [\mathbb{D}^{[\mathbf{W}]},\hat{\mathcal{F}}_\alpha^\dagger]_{\mathrm{endo}}\,\hat{\mathcal{F}}_\beta\,P_{n,x'} \ \pm\ \hat{\mathcal{F}}_\alpha^\dagger\,[\mathbb{D}^{[\mathbf{W}]},\hat{\mathcal{F}}_\beta]_{\mathrm{endo}}\,P_{n,x'} \Big|\Omega\Big\rangle,\end{aligned}$$

since $[\mathbb{D}_{x'}^{[\mathbf{W}]}, P_{n,x'}] = 0$ (endpoint eigenmode). Hence

$$\left|\mathcal{S}_n^{[\mathbf{W}]}(x,x')\right| \ \le\ \left\|[\mathbb{D}^{[\mathbf{W}]},\hat{\mathcal{F}}_\alpha]\right\| \ +\ \left\|[\mathbb{D}^{[\mathbf{W}]},\hat{\mathcal{F}}_\beta]\right\| \ :=:\ C.$$

This C is independent of n and the word, so the uniform theorem applies with $k = 0$.

Distance. $d_{\mathrm{corr}}^{[\mathbf{W}]}(x,x';\psi_1,\psi_2)$ reduces to the normalized overlap through the local wordwise kernel (no transport), hence is word/gauge invariant.

Finite fibre, flat transport (uniform bound). A four-level internal Hilbert space (spin-(3/2) or ququart) with flat transport is a textbook finite-dimensional quantum system: the only source of response is the noncommutativity between the chosen measurement channels and the internal generator, so the mode index does not amplify the effect. This matches the standard picture of basis-mismatch dynamics (Ramsey-type logic without spatial gradients) in which the relevant bounds come from operator norms of finite matrices rather than dispersion with momentum or winding number [350, 287]. Thinking atomically, internal multiplets behave this way when spatial gauge structure is absent [179]. In our wordwise framing, the "uniform in n" cap is just the finite-dimensional commutator control that underlies basic Heisenberg-picture estimates [350].

Example 2 — Dirac-type along a circle (polynomial bound $k = 1$)

Setup. Configuration $s \in \mathbb{S}^1$ (arclength). Fibres are $\mathbb{C}^4$. Smooth wordwise connection $\mathcal{A}^{[\mathbf{W}]}(s)$ with transport $U_\gamma = \mathcal{P}\exp\big(-\int_\gamma \mathcal{A}^{[\mathbf{W}]}\big)$.

Wordwise spectral operator.

$$\mathbb{D}^{[\mathbf{W}]} = -i\,\partial_s \otimes \mathbb{I}_4 \;+\; V^{[\mathbf{W}]}(s),$$

with $V^{[\mathbf{W}]}$ bounded and smooth.

Regularity and growth. Selectors are multiplication-type $\Rightarrow r = 0$. Connection is first order: $[\mathbb{D}^{[\mathbf{W}]}, \mathcal{A}^{[\mathbf{W}]}] = -i(\partial_s \mathcal{A}^{[\mathbf{W}]}) + [V^{[\mathbf{W}]}, \mathcal{A}^{[\mathbf{W}]}]$, bounded on $\mathbb{S}^1 \Rightarrow s = 1$. Weyl growth on the circle: $|\lambda_{n,[\mathbf{W}]}| \sim n \Rightarrow \vartheta = 1$.

Shear bound. By the polynomial theorem,

$$\left|\mathcal{S}_n^{[\mathbf{W}]}(x, x')\right| \;\le\; C\,(1 + |\lambda_{n,[\mathbf{W}]}|)^{\max\{r,s\}} \le C'(1+n)^1.$$

Dirac on a ring (polynomial $k = 1$). On a mesoscopic ring or ring resonator with a smooth vector potential, a first-order generator $-i\,\partial_s + V$ and nontrivial holonomy produces a mode-dependent phase twist: higher winding modes accumulate more phase, so the commutator-based shear grows like $1 + n$. This is the Aharonov–Bohm mechanism in its cleanest form—gauge-invariant but path-sensitive via the Wilson loop—together with geometric-phase accumulation for cyclic transport [378, 24, 389, 25]. The scaling intuition follows directly from first-order transport on a compact 1D base with linear spectral growth of the Dirac harmonics [379]. *Suggested:* ring-resonator implementations (e.g., integrated optics à la Yariv) and mesoscopic AB experiments.

Example 3 — Discrete graph–Dirac on a cycle (polynomial $k = 1$)

Setup. Loci are vertices $j \in C_M$ (cycle graph), spacing $h := 2\pi/M$. Fibre at each vertex is $\mathbb{C}^4$. Edge connection: unitary links $U^{[\mathbf{W}]}_{j\to j+1} = \exp\big(-\mathcal{A}^{[\mathbf{W}]}_{j+1/2}\big)$.

Wordwise spectral operator. Define a discrete first-difference Dirac:

$$(\mathbb{D}^{[\mathbf{W}]}\psi)(j) = \frac{1}{2h}\Big(U^{[\mathbf{W}]}_{j-1\to j}\psi(j) \;-\; U^{[\mathbf{W}]}_{j+1\to j}\psi(j)\Big) \;+\; V^{[\mathbf{W}]}(j)\psi(j),$$

with bounded $V^{[\mathbf{W}]}$.

Regularity and growth. Selectors are local $\Rightarrow r = 0$. $[\mathbb{D}^{[\mathbf{W}]}, \mathcal{A}]$ involves first differences of $\mathcal{A}$ along edges $\Rightarrow s = 1$. Discrete eigenvalues on C_M satisfy $|\lambda_n| \asymp n$ for $n \ll M$ (after unfolding) $\Rightarrow \vartheta = 1$.

Shear bound.

$$\Big|\mathcal{S}^{[\mathbf{W}]}_n(x, x')\Big| \;\le\; C(1+n).$$

Discrete graph–Dirac on a cycle (polynomial $k = 1$). A tight-binding ring is the lattice analogue of the continuous case: the discrete Dirac (first difference) plus a link-variable connection generates shear controlled by the commutator with the on-site gauge/potential, and the transport contribution adds along the path as small link phases—precisely the Wilson-line intuition on a lattice [389]. The spectral ladder on a cycle grows linearly in the mode index for low-lying harmonics, giving the same $k = 1$ scaling. This picture dovetails with standard tight-binding solid-state lore (band formation from hopping on periodic graphs) [390, 391] and with noncommutative-geometric treatments of lattice transport [392].

Example 4 — Para-preserving generator (shear from transport only)

Setup. Focus on the parasymmetric facet. Let $P^{[\mathbf{W}]}_{\text{para}}$ be the para projector in the wordwise frame. Choose $\mathbb{D}^{[\mathbf{W}]}$ commuting with it.

Choice. Take $\mathbb{D}^{[\mathbf{W}]} = \hat{F}^{(P)}_{\text{para}} := \sum_{i<j} J_{ij}\Big(P^{(x_i x_j)}_{\text{para}} - P^{(x_i x_j)}_{\text{antipara}}\Big)^{[\mathbf{W}]}$, so $[\mathbb{D}^{[\mathbf{W}]}, \hat{\mathcal{F}}^{[\mathbf{W}]}_{\text{para}}]_{\text{endo}} = 0$.

Shear expansion.

$$\Big[\,\mathbb{D}^{[\mathbf{W}]},\; \hat{\mathcal{F}}^{\dagger}_{\alpha} U_\gamma \hat{\mathcal{F}}_\beta P_{n,x'}\,\Big]_{\text{endo}} = \hat{\mathcal{F}}^{\dagger}_{\alpha}\,[\mathbb{D}^{[\mathbf{W}]}, U_\gamma]_{\text{endo}}\;\hat{\mathcal{F}}_\beta P_{n,x'},$$

so only the transport term contributes.

Bounds. If $\mathcal{A}^{[\mathbf{W}]} = 0$ locally, then $[\mathbb{D}^{[\mathbf{W}]}, U_\gamma] = O(\ell^2)$ along a short path of length ℓ (curvature order), hence $|\mathcal{S}^{[\mathbf{W}]}_n(x, x')| \le C\,\ell^2$ uniformly in n. With small nonzero connection, $\|[\mathbb{D}^{[\mathbf{W}]}, U_\gamma]\| \le L_{\text{conn}}\ell + O(\ell^2) \Rightarrow |\mathcal{S}^{[\mathbf{W}]}_n| \le C_1\ell + C_2\ell^2$, still independent of n.

Para-preserving generator (shear from transport only). If the generator respects a symmetry sector (e.g., fermion parity, a protected pseudospin), then channel projectors within that sector commute with the dynamics and cannot create shear by themselves; only holonomy of the wordwise connection contributes. In the flat limit, the first-order effect vanishes; with small curvature, the response is geometric (Berry/Wilczek–Zee-type) and independent of mode index, reflecting symmetry protection [24, 25, 393]. From a control perspective, this is exactly the scenario where choosing symmetry-adapted frames stabilizes subspaces against dynamical mixing [287]. *Suggested:* original Wilczek–Zee non-Abelian phase (1984) and DFS/encoded-subspace reviews (e.g., Lidar–Whaley; Zanardi–Rasetti).

Example 5 — Holonomy bump (local curvature patch; uniform $k = 0$)

Setup. Along γ, let $\mathcal{A}^{[\mathbf{W}]} = 0$ except on a small sub-arc of length L_{bump} where $\mathcal{F} \neq 0$. Assume selector commutators bounded ($r = 0$) and that $[\mathbb{D}^{[\mathbf{W}]}, \mathcal{A}^{[\mathbf{W}]}]$ is uniformly bounded ($s = 0$).

Estimate. From Duhamel, $\|[\mathbb{D}^{[\mathbf{W}]}, U_\gamma]\| \leq \int_{\text{bump}} \|[\mathbb{D}^{[\mathbf{W}]}, \mathcal{A}]\|\, ds \leq L_{\text{conn}}\, L_{\text{bump}}$. Selector terms contribute a constant C_0. Therefore

$$\left|\mathcal{S}_n^{[\mathbf{W}]}(x, x')\right| \;\leq\; C_0 + C_1\, L_{\text{bump}},$$

uniform in n and word ($k = 0$).

Holonomy bump (localized curvature; uniform bound). A localized flux/curvature patch (magnetic "bump," refractive micro-lens, patterned gauge cell) produces a bounded, path-length–proportional geometric phase while leaving mode-index scaling essentially flat: shear grows with the portion of the path that traverses the bump but not with n. This is the local Aharonov–Bohm/holonomy picture: the effect is entirely geometric and encoded by a Wilson loop over the curved patch [378, 389]. Because no first-order differential growth with mode index is engaged, the response remains uniform—consistent with geometric-phase linear response on compact supports [25].

Example 6 — Twofold slice (abstract qubit inside the Fourfold)

Setup. Choose a word $\mathbf{W}$ and the isometric embedding $U^{[\mathbf{W}]} : \mathbb{C}^2 \hookrightarrow \mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}$ selecting the Twofold $(0, 1)$ slice. Let $\varphi_1, \varphi_2 \in \mathbb{C}^2$ be qubit probes and set $\psi_j := U^{[\mathbf{W}]}\varphi_j$ at the respective loci.

Distance reduction. The correlator amplitude becomes

$$\mathsf{G}^{[\mathbf{W}]}(x, x'; \psi_1, \psi_2) = \left\langle \Omega \middle| \hat{G}_\Lambda^{[\mathbf{W}]}\Big[\,(U\varphi_1)^*,\, U\varphi_2\,\Big] \middle| \Omega \right\rangle = \left\langle \Omega \middle| \hat{G}_{\Lambda,(0,1)}^{[\mathbf{W}]}\Big[\varphi_1^*, \varphi_2\Big] \middle| \Omega \right\rangle,$$

so $d_{\text{corr}}^{[\mathbf{W}]}(x, x'; \psi_1, \psi_2) = \sqrt{1 - \frac{|\langle \varphi_1 | \Phi(x,x')\, \varphi_2 \rangle|}{\|\Phi(x,x')^{1/2}\varphi_1\|\, \|\Phi(x,x')^{1/2}\varphi_2\|}}$, where $\Phi(x, x')$ is the induced 2×2 positive kernel on the Twofold block. Any word change rotates the qubit frame unitarily, leaving the value invariant.

Shear. If $\mathbb{D}^{[\mathbf{W}]}$ preserves the Twofold block (e.g. block-diagonal), the shear reduces to a qubit-level commutator with the induced transport; bounds come from Example 2/5 depending on whether the connection is first order or uniformly bounded.

Twofold slice / abstract qubit. Embedding an abstract qubit (Twofold slice) inside a larger modal space reproduces familiar single-qubit geometry: the correlator distance reduces to normalized qubit overlaps, and word/frame changes become Bloch-sphere rotations that do not alter operational predictions. This is the standard equivalence of encodings and the insensitivity of physical figures of merit to basis changes on the code subspace [287, 287]. It also harmonizes with hardware-agnostic criteria for qubit realizations—what matters is controllable subspace dynamics and well-defined measurement, not the embedding per se [351, 349].

Example 7 — Zonal evolution (static $\oplus$ dynamic; $s = 1$)

Setup. Partition loci into $X_{\mathrm{stat}} \sqcup X_{\mathrm{dyn}}$. Along paths inside X_{stat}, take $\mathcal{A}^{[\mathbf{W}]} = 0$; inside X_{dyn}, $\mathcal{A}^{[\mathbf{W}]}$ is first-order smooth. Selectors bounded ($r = 0$).

Shear localization. For a path γ crossing a dynamic segment of length L_{dyn},

$$\left\| [\mathbb{D}^{[\mathbf{W}]}, U_\gamma] \right\| \;\le\; \int_{\gamma \cap X_{\mathrm{dyn}}} \left\| [\mathbb{D}^{[\mathbf{W}]}, \mathcal{A}] \right\| ds \;\le\; L_{\mathrm{conn}}\, L_{\mathrm{dyn}}.$$

Hence

$$\left| \mathcal{S}_n^{[\mathbf{W}]}(x, x') \right| \;\le\; C_0 \;+\; C_1\, L_{\mathrm{dyn}} \quad \text{(uniform in } n\text{)}.$$

If, additionally, $\mathbb{D}^{[\mathbf{W}]}$ is Dirac-type on X_{dyn} (as in Example 2) and the base along the dynamic zone is compact with Weyl growth exponent $\vartheta = 1$, then the polynomial form holds:

$$\left| \mathcal{S}_n^{[\mathbf{W}]}(x, x') \right| \;\le\; C'(1+n)^{\vartheta \max\{r,s\}} = C'(1+n).$$

Static-support vanishing. If both probes are supported in X_{stat} and $\gamma \subset X_{\mathrm{stat}}$, then $U_\gamma = \mathbb{I}$ and the only commutators are selector terms; when $\mathbb{D}^{[\mathbf{W}]}$ commutes with those selectors (e.g. block-diagonal in the chosen modal split), the shear vanishes to first order.

Zonal evolution (static $\oplus$ dynamic region; localization of shear). When only a spatial subregion is driven (Floquet patch, gated strip, addressed zone), shear localizes to the dynamic segment and scales with its length; paths that remain in the static zone show vanishing first-order response. If the driven block is Dirac-type on a d-dimensional compact base, the spectral growth is governed by Weyl-type asymptotics, leading to $|\mathcal{S}| \lesssim (1+n)^{1/d}$: higher dimension dilutes mode growth [394, 395]. This aligns with modern Floquet-engineering perspectives where transport and response are organized by local drive profiles and effective Hamiltonians [396]. *Suggested:* broader Floquet reviews (Bukov–D'Alessio–Polkovnikov 2015; Eckardt 2017) and experimental platforms (cold-atom/photonic lattices) for zonal drives.

Example 1 — Finite fibre, flat transport (uniform bound $k = 0$)

Setup. Two loci x, x'. Each modal fibre is $\mathbb{C}^4$. Fix any word $\mathbf{W}$. Let the wordwise connection vanish locally: $\mathcal{A}^{[\mathbf{W}]} = 0$ so $U_\gamma = \mathbb{I}$ along short paths. Pick the wordwise spectral operator diagonal in some frame,

$$\mathbb{D}_x^{[\mathbf{W}]} = \mathrm{diag}(d_1, d_2, d_3, d_4), \quad \mathbb{D}_{x'}^{[\mathbf{W}]} = \mathrm{diag}(d'_1, d'_2, d'_3, d'_4),$$

and bounded word-rotated selectors $\{\hat{\mathcal{F}}_\alpha^{[\mathbf{W}]}\}_{\alpha\in\{0,1,p,s\}}$.

Shear bound. With $U_\gamma = \mathbb{I}$ and $P_{n,x'} := |\psi_{n,[\mathbf{W}]}^{(x')}\rangle\langle\psi_{n,[\mathbf{W}]}^{(x')}|$,

$$\begin{aligned}\mathcal{S}_n^{[\mathbf{W}]}(x,x') &= \Big\langle\Omega\Big|\Big[\mathbb{D}^{[\mathbf{W}]},\ \hat{\mathcal{F}}_\alpha^\dagger\,\hat{\mathcal{F}}_\beta\,P_{n,x'}\Big]_{\mathrm{endo}}\Big|\Omega\Big\rangle \\ &= \Big\langle\Omega\Big|[\mathbb{D}^{[\mathbf{W}]},\hat{\mathcal{F}}_\alpha^\dagger]_{\mathrm{endo}}\ \hat{\mathcal{F}}_\beta\,P_{n,x'} \\ &\qquad \pm\ \hat{\mathcal{F}}_\alpha^\dagger\,[\mathbb{D}^{[\mathbf{W}]},\hat{\mathcal{F}}_\beta]_{\mathrm{endo}}\,P_{n,x'}\Big|\Omega\Big\rangle.\end{aligned}$$

Since $[\mathbb{D}_{x'}^{[\mathbf{W}]}, P_{n,x'}] = 0$ (endpoint eigenmode), we obtain the uniform estimate

$$\Big|\mathcal{S}_n^{[\mathbf{W}]}(x,x')\Big| \ \le\ \Big\|[\mathbb{D}^{[\mathbf{W}]},\hat{\mathcal{F}}_\alpha]\Big\| \ +\ \Big\|[\mathbb{D}^{[\mathbf{W}]},\hat{\mathcal{F}}_\beta]\Big\| =:\ C,$$

independent of n and of the chosen word. This is the uniform form of spectral smoothness (Theorem 4.384(a), $k = 0$).

Toy numeric instance. Take $D = \mathrm{diag}(0, 1, 2, 3)$, $P_0 = |0\rangle\langle 0|$, $P_1 = |1\rangle\langle 1|$, $P_\pm = |\pm\rangle\langle\pm|$ with $|\pm\rangle = (|0\rangle \pm |1\rangle)/\sqrt{2}$. Then

$$\|[D, P_0]\| = \|[D, P_1]\| = 0, \qquad \|[D, P_+]\| = \|[D, P_-]\| = \tfrac{1}{2},$$

hence, e.g. for $(\alpha = 0, \beta = +)$, $\Big|\mathcal{S}_n^{[\mathbf{W}]}(x,x')\Big| \le \frac{1}{2}$.

Physical/real-world association (Example 1: finite fibre, flat transport). This setting models a multilevel internal system with no spatial gauge structure—e.g., a spin-(3/2) manifold, a four-level atom, or a ququart encoded in superconducting levels or trapped ions. The vanishing connection ($\mathcal{A}^{[\mathbf{W}]} = 0$) means no geometric holonomy is picked up along short paths, so the only mechanism for a nonzero shear is internal basis mismatch: the selectors (measurement/effect channels) do not commute with the internal generator $\mathbb{D}^{[\mathbf{W}]}$. Operationally, this reproduces the finite-dimensional Ramsey picture: contrasts and response are set by the commutator norms of finite matrices, hence the n-independent (uniform) cap on $\Big|\mathcal{S}_n^{[\mathbf{W}]}\Big|$ [350, 287]. Physically, a change of word $\mathbf{W}$ is just a change of readout frame; the predicted signal is invariant under such frame rotations, and the observable response is controlled solely by how the chosen measurement axes fail to commute with the internal Hamiltonian on the four-level manifold [179, 350].

Example 3 — Discrete graph–Dirac on a cycle (polynomial $k = 1$)

Setup. Loci are vertices $j \in C_M$ (cycle), spacing $h = 2\pi/M$. Fibre at each vertex is $\mathbb{C}^4$. Edge connection: unitary links $U^{[\mathbf{W}]}_{j\to j+1} = \exp\big(-\mathcal{A}^{[\mathbf{W}]}_{j+1/2}\big)$.

Wordwise Dirac. Let T be the shift on the cycle. Define the (Hermitian) discrete Dirac

$$\mathbb{D}^{[\mathbf{W}]} \;:=\; -\frac{i}{2h}\left(T - T^{\dagger}\right) \otimes \mathbb{I}_4 \;+\; V^{[\mathbf{W}]},$$

with bounded $V^{[\mathbf{W}]}$. Selectors are local (bounded).

Regularity & growth. Selector regularity $r = 0$. First-order connection along edges gives $s = 1$ via $[\mathbb{D}^{[\mathbf{W}]}, \mathcal{A}^{[\mathbf{W}]}]$ (finite differences). Weyl growth on a 1D compact base: $|\lambda_{n,[\mathbf{W}]}| \asymp n \Rightarrow \vartheta = 1$.

Shear bound. By Theorem 4.384(b) and Corollary 4.377,

$$\left|\mathcal{S}^{[\mathbf{W}]}_n(x, x')\right| \;\le\; C\,(1+n)^{\vartheta \max\{r,s\}} = C\,(1+n)^{1}.$$

Equivalently, for a short path of L edges (length $\approx Lh$),

$$\left\|[\mathbb{D}^{[\mathbf{W}]}, U_\gamma]\right\| \;\le\; \int_\gamma \left\|[\mathbb{D}^{[\mathbf{W}]}, \mathcal{A}]\right\| ds \;\lesssim\; L_{\mathrm{conn}} \cdot (Lh),$$

which feeds into the same polynomial estimate.

Toy numeric instance. For $M = 16$, $a = 0.7$ and a diagonal connection $A(j) = a\sin(2\pi j/M)$,

$$\left\|[\,\mathbb{D}^{[\mathbf{W}]}, A\,]\right\| \approx 0.5722,$$

and the transport contribution along a path of L edges satisfies $\left\|[\mathbb{D}^{[\mathbf{W}]}, U_\gamma]\right\| \lesssim 0.5722 \cdot (Lh)$.

Physical/real-world association (Example 3: discrete graph–Dirac on a cycle). A cycle graph with link unitaries captures a tight-binding ring with a synthetic (or real) gauge: think cold atoms in a ring lattice with Peierls phases, or a photonic ring of coupled resonators with engineered link phases. The discrete Dirac $\mathbb{D}^{[\mathbf{W}]} \propto (T - T^{\dagger})$ plays the role of a first-order lattice derivative; the edge connection encodes the link field. The commutator $[\mathbb{D}^{[\mathbf{W}]}, \mathcal{A}^{[\mathbf{W}]}]$ measures discrete gradients of the gauge/potential, so transport contributions add linearly with the number of traversed links (Duhamel sum of small phase kicks), producing the polynomial $k = 1$ shear bound. Spectrally, low-lying harmonics on a ring grow linearly with mode index, so higher modes are more sensitive to accumulated link phases (lattice Wilson line) [389]. In hardware, this maps to: (i) cold-atom rings with Raman-induced flux, (ii) integrated photonic rings with modulation-patterned couplers, or (iii) mesoscopic normal-metal/superconducting loops—each realizing mode-sensitive geometric phase twists along the cycle [378, 390, 391, 392].

Example T2 — 2D discrete torus Dirac (polynomial $k = \frac{1}{2}$)

Setup. Base $\mathbb{T}^2 = \mathbb{S}^1_{L_x} \times \mathbb{S}^1_{L_y}$ with coordinates (u, v), fibres $\mathbb{C}^4$. Smooth wordwise connection $\mathcal{A}^{[\mathbf{W}]} = (\mathcal{A}^{[\mathbf{W}]}_u, \mathcal{A}^{[\mathbf{W}]}_v)$ and transport $U_\gamma = \mathcal{P}\exp\Big(-\int_\gamma \mathcal{A}^{[\mathbf{W}]}\Big)$.

Wordwise Dirac. Let Γ_x, Γ_y be 4×4 Hermitian matrices with $\{\Gamma_\mu, \Gamma_\nu\} = 2\delta_{\mu\nu}$, and

$$\mathbb{D}^{[\mathbf{W}]} \;:=\; -i\Big(\partial_u \otimes \Gamma_x + \partial_v \otimes \Gamma_y\Big) \;+\; V^{[\mathbf{W}]}(u, v),$$

with bounded smooth $V^{[\mathbf{W}]}$.

Regularity & growth. Selectors are multiplication-type $\Rightarrow r = 0$. First-order connection:

$$[\mathbb{D}^{[\mathbf{W}]}, \mathcal{A}^{[\mathbf{W}]}_\mu] = -i\,(\partial_\mu \mathcal{A}^{[\mathbf{W}]}_\mu) \otimes \Gamma_\mu + [V^{[\mathbf{W}]}, \mathcal{A}^{[\mathbf{W}]}_\mu],$$

bounded on $\mathbb{T}^2 \Rightarrow s = 1$. For Dirac on a 2D compact base, Weyl growth exponent $\vartheta = \frac{1}{2}$.

Shear bound. By Theorem 4.384(b) and Corollary 4.377,

$$\Big|\mathcal{S}^{[\mathbf{W}]}_n(x, x')\Big| \;\le\; C\,(1 + |\lambda_{n,[\mathbf{W}]}|)^{\max\{r,s\}} \;\le\; C'\,(1+n)^{\vartheta\max\{r,s\}} = C'\,(1+n)^{1/2}.$$

Physical/real-world association (Example T2: 2D discrete torus Dirac). A 2D torus with a Dirac-type generator models platforms where transport is governed by first-order kinematics on a compact 2D base: e.g., photonic or phononic crystals with Dirac cones, ultracold atoms in optical lattices near Dirac points, or metamaterials emulating 2D Dirac dynamics. The wordwise connection $\mathcal{A}^{[\mathbf{W}]}$ encodes non-Abelian Berry/Wilczek–Zee structure of band frames; shear is set by $[\mathbb{D}^{[\mathbf{W}]}, \mathcal{A}^{[\mathbf{W}]}]$, i.e., how the drive or background gauge twists those frames [24, 393]. Weyl scaling on a compact 2D base gives $|\lambda_n| \sim n^{1/2}$, hence the $(1+n)^{1/2}$ bound: higher modes still feel more twist, but less aggressively than in 1D [394, 395]. Experimentally, the prediction is that mode-resolved linear-response signals (or commutator-based changes of suitable two-point correlators) should increase sublinearly with the band index, with the prefactor controlled by the local curvature/connection strength and the path length across the torus.

Example Z — Block-diagonal $\mathbb{D}^{[\mathbf{W}]}$ with shear localized to a dynamic zone

Setup (zonal split). Partition loci as $X_{\text{stat}} \sqcup X_{\text{dyn}}$. Along X_{stat} take $\mathcal{A}^{[\mathbf{W}]} = 0$ (flat); along X_{dyn} take a first-order smooth connection. Let the wordwise spectral operator be block-diagonal:

$$\mathbb{D}^{[\mathbf{W}]} \;=\; \mathbb{D}^{[\mathbf{W}]}_{\text{stat}} \oplus \mathbb{D}^{[\mathbf{W}]}_{\text{dyn}}, \qquad \Big[\mathbb{D}^{[\mathbf{W}]}_{\text{stat}},\; \hat{\mathcal{F}}^{[\mathbf{W}]}_\bullet\Big]_{\text{endo}} = 0,$$

so selectors commute on the static block.

Localized shear. For a path γ crossing a dynamic segment of length L_{dyn},

$$\left\|[\mathbb{D}^{[\mathbf{W}]}, U_\gamma]\right\| \;\le\; \int_{\gamma\cap X_{\mathrm{dyn}}} \left\|[\mathbb{D}^{[\mathbf{W}]}, \mathcal{A}]\right\| ds \;\le\; L_{\mathrm{conn}}\, L_{\mathrm{dyn}},$$

and the selector commutators on X_{stat} vanish. Hence

$$\left|\mathcal{S}_n^{[\mathbf{W}]}(x,x')\right| \;\le\; C_0 \;+\; C_1\, L_{\mathrm{dyn}},$$

uniform in n (the $k=0$ case of Theorem 4.384).

Polynomial refinement (dynamic Dirac). If $\mathbb{D}_{\mathrm{dyn}}^{[\mathbf{W}]}$ is Dirac-type on a d-dimensional compact dynamic sub-base, then $r=0$, $s=1$, and the Weyl exponent is $\vartheta=\frac{1}{d}$. By Theorem 4.384(b),

$$\left|\mathcal{S}_n^{[\mathbf{W}]}(x,x')\right| \;\le\; C'\,(1+n)^{\,\vartheta\cdot\max\{r,s\}} \;=\; C'\,(1+n)^{1/d}.$$

If both probes and the entire path γ are supported in X_{stat}, then $U_\gamma=\mathbb{I}$ and the shear vanishes to first order (only static selector terms remain and commute).

Physical/real-world association (Example Z: block-diagonal $\mathbb{D}^{[\mathbf{W}]}$; localized shear in a driven zone). Here the system is partitioned into a static zone (flat transport) and a driven zone (first-order connection), mirroring devices with spatially localized modulation: a Floquet "patch" on a 2D material, a gated stripe in a mesoscopic conductor, a subset of waveguides with longitudinal modulation, or a flux-biased region in a superconducting array. Because the static block's selectors commute with its generator, shear from that region vanishes to first order; all response localizes to the driven segment and scales with its traversed length L_{dyn}. If the driven block is Dirac-type on an effective d-dimensional compact base, Weyl scaling gives $|\mathcal{S}_n^{[\mathbf{W}]}|\lesssim(1+n)^{1/d}$: increasing spatial dimension dilutes the growth of mode sensitivity [394, 395]. Practically, this predicts (i) spatial control: moving the probe path in and out of the driven zone toggles the first-order signal; (ii) spectral control: higher modes amplify the response with a dimension-dependent power; and (iii) frame/word robustness: changing the local readout basis does not change the measured scalars, only how they are expressed, so the localization-and-scaling features are genuine device-level invariants [396].

4.9.6 Spectral Distance: Functional Geometry via Operator Commutators

We now introduce the third component of the relational metric profile, the *spectral distance* $d_{\mathrm{spec}}(x,x')$. Complementing the coherence distance d_{corr} and the vacuum-response distance d_{vac}, this component probes the *differential* aspect of the emergent geometry on $\mathcal{X}_q$, intrinsically and without reference to any background manifold.

Noncommutative-geometry template

In Connes' framework, a metric may be extracted from a spectral triple $(\mathcal{A},\mathcal{H},\mathcal{D})$ via

$$d(x,x') \;=\; \sup_{f\in\mathcal{A}}\Big\{\,|f(x)-f(x')| \;\Big|\; \|[\mathcal{D},f]\|_{\mathrm{op}}\le 1\,\Big\}, \tag{4.185}$$

see [4, 5, 4]. In IQR we retain the *commutator-as-Lipschitz* insight but instantiate it relationally using the already-defined Fourfold Inter-Modal Correlator $\hat{G}_\Lambda$ (Definition 4.9) and the Spectral Operator $\hat{D}$ (Definition 4.341), evaluated in the Quantum Holo-State $|\mathring{\mathbf{\Psi}}\rangle$ (Definition 1.27).

IQR formulation: state-conditioned, relational

Definition 4.386 (Spectral Shear Observable)**.** Fix loci $\mathfrak{B}_x, \mathfrak{B}_{x'}$ and choose a local orthonormal modal basis $\{|\psi_n^{(x)}\rangle\}_n$ at x (e.g. an eigenbasis of $\hat{D}_x$ as in Definition 4.341). The *spectral shear* for mode n is the Holo-State expectation of the commutator between the Spectral Operator and the bilinear evaluation of the Fourfold correlator on $|\psi_n^{(x)}\rangle$:

$$\mathcal{S}_n(x,x') \;:=\; \Big\langle \mathring{\mathbf{\Psi}} \Big| \Big[\, \hat{D},\; \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_n^*, \psi_n] \,\Big] \Big| \mathring{\mathbf{\Psi}} \Big\rangle \quad \text{(cf. (4.160))}.$$

A nonzero value of $\mathcal{S}_n(x,x')$ indicates that the relational structure encoded by $\hat{G}_\Lambda$ between x and x' is deformed by the spectral flow generated by $\hat{D}$, thereby contributing to the spectral component of separation.

Remarks. (i) The polynomial bound of Definition 4.342 guarantees well-posedness (finite aggregates over n) in near-coincident regimes. (ii) The construction is invariant under admissible reindexings and the duality involution (Proposition 1.8). (iii) In the duality-collapse regime (Definition 4.87), the shear reduces to the Twofold commutator signal on the $\{0,1\}$ slice, aligning with standard qubit differential diagnostics (cf. Proposition 4.328).

4.9.7 Wordwise Reformulations of Spectral Distance: Functional Geometry via Operator Commutators

Principle. Retain Connes' "commutator-as-Lipschitz" idea, but evaluate all point–separating functionals through the *word–dressed* correlator and fibrewise letters. Charge–preserving word changes act by endpoint unitaries (Remark 4.348), and the endogenous super–commutator is conjugation–invariant (Remark 4.372); hence the resulting distance is intrinsic (Proposition 4.382).

Definition 4.387 (Wordwise Lipschitz seminorm)**.** Fix a word $\mathbf{W}$ and let $\mathbb{D}^{[\mathbf{W}]}$ be a wordwise spectral operator (Definition 4.345). For any bounded fibre operator Φ on $\mathcal{H}_{\mathrm{em}}$, define

$$L^{[\mathbf{W}]}(\Phi) \;:=\; \Big\| \Big[\, \mathbb{D}^{[\mathbf{W}]},\; \Phi \,\Big]_{\mathrm{endo}} \Big\|_{\mathrm{op}}. \tag{4.186}$$

By charge–preserving conjugation invariance (Remark 4.372) and Proposition 4.382, $L^{[\mathbf{W}]}(\Phi)$ is independent of the choice of word; we nevertheless keep the superscript to emphasise the wordwise construction.

Definition 4.388 (Transport–consistent wordwise evaluations)**.** Let $x, x' \in \mathcal{X}_q$, choose a path $\gamma : x \to x'$ with transporter $\mathcal{U}^{[\mathbf{W}]}(x' \leftarrow x; \gamma)$ (Section 4.5), and pick a unit probe $|\psi\rangle_x \in \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$. Form the word–dressed letter $\hat{\mathcal{F}}_\psi^{[\mathbf{W}]}(x) := \sum_\alpha c_\alpha \hat{\mathcal{F}}_\alpha^{[\mathbf{W}]}(x)$ for $|\psi\rangle_x = \sum_\alpha c_\alpha |\alpha\rangle_x$,

and transport the probe to $|\psi\rangle_{x',\gamma} := \mathcal{U}^{[\mathbf{W}]}(x' \leftarrow x;\gamma)\,|\psi\rangle_x$. With coincident normalisers $N^{[\mathbf{W}]}(\psi,x) := \sqrt{\big\langle (\hat{\mathcal{F}}^{[\mathbf{W}]}_\psi(x))^\dagger \hat{\mathcal{F}}^{[\mathbf{W}]}_\psi(x)\big\rangle}$ and analogously for (ψ,x'), define

$$\begin{aligned} \mathsf{ev}^{[\mathbf{W}]}_{x,\psi}(\Phi) &:= \frac{\big\langle (\hat{\mathcal{F}}^{[\mathbf{W}]}_\psi(x))^\dagger\, \Phi\, \hat{\mathcal{F}}^{[\mathbf{W}]}_\psi(x)\big\rangle}{\big(N^{[\mathbf{W}]}(\psi,x)\big)^2}, \\ \mathsf{ev}^{[\mathbf{W}]}_{x',\psi;\gamma}(\Phi) &:= \frac{\big\langle (\hat{\mathcal{F}}^{[\mathbf{W}]}_\psi(x'))^\dagger\, \Phi\, \hat{\mathcal{F}}^{[\mathbf{W}]}_\psi(x')\big\rangle}{\big(N^{[\mathbf{W}]}(\psi,x')\big)^2}, \qquad |\psi\rangle_{x'} = \mathcal{U}^{[\mathbf{W}]}(x'\leftarrow x;\gamma)|\psi\rangle_x, \end{aligned} \tag{4.187}$$

where $\langle\cdot\rangle$ denotes the Holo–State expectation. These evaluations are gauge/word invariant (Remarks 4.348 and 4.372).

Definition 4.389 (Wordwise spectral distance)**.** For $x,x' \in \mathcal{X}_q$, define

$$d^{[\mathbf{W}]}_{\mathrm{spec}}(x,x') \;:=\; \sup_{\substack{\Phi=\Phi^\dagger \\ L^{[\mathbf{W}]}(\Phi)\le 1}} \; \sup_{\|\psi\|=1} \; \inf_{\gamma: x\to x'} \Big|\, \mathsf{ev}^{[\mathbf{W}]}_{x,\psi}(\Phi) - \mathsf{ev}^{[\mathbf{W}]}_{x',\psi;\gamma}(\Phi)\,\Big|. \tag{4.188}$$

The inner infimum removes path artefacts; on flat simply–connected neighbourhoods it is attained and path–independent by wordwise parallel consistency.

Proposition 4.390 (Basic properties)**.** *The quantity $d^{[\mathbf{W}]}_{\mathrm{spec}}$ in (4.188) is:*

(a) Word/gauge invariant*: independent of $\mathbf{W}$ and frame choices (Remarks 4.348 and 4.372 and Proposition 4.382).*

(b) A pseudo–metric*: symmetric, $d(x,x)=0$, and satisfies the triangle inequality (via the usual telescoping argument with $L^{[\mathbf{W}]}(\Phi)\le 1$ on each leg).*

(c) Local–flat reduction*: if $\mathcal{F}^{[\mathbf{W}]}=0$ on a simply–connected patch, then $d^{[\mathbf{W}]}_{\mathrm{spec}}(x,x')$ is independent of γ and reduces to the formula computed in a trivial word/frame.*

(d) Twofold reduction*: under the wordwise qubit embedding (Section 4.6.2), the supremum may be restricted to the compressed Twofold sector, yielding the usual qubit Connes–type distance on $\mathbb{C}^2$.*

Remark 4.391 (Linearisation and spectral shear)**.** For small separations, expand the transporter along γ; the difference of evaluations linearises to the wordwise spectral shear (cf. Definition 4.381):

$$\Big|\, \mathsf{ev}^{[\mathbf{W}]}_{x,\psi}(\Phi) - \mathsf{ev}^{[\mathbf{W}]}_{x',\psi;\gamma}(\Phi)\,\Big| \;=\; \sup_{L^{[\mathbf{W}]}(\Phi)\le 1} \Big|\, \Big\langle \big[\, \mathbb{D}^{[\mathbf{W}]},\; \hat{G}^{[\mathbf{W}]}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})[\psi^*,\psi]\,\big]_{\mathrm{endo}} \Big\rangle\,\Big| \;+\; O(\mathrm{area}),$$

which is well–posed by wordwise spectral smoothness (Definition 4.383). Thus $d^{[\mathbf{W}]}_{\mathrm{spec}}$ packages the first–order commutator response of the wordwise correlator in the endogenous calculus.

The State-Conditioned Modal Weight and Endogenous Regularization

A cornerstone of the relational spectral geometry is a *state–dependent* weighting of spectral modes. This *modal weight* quantifies, for each mode, how strongly the universal Quantum Holo-State supports the corresponding bilocal relational probe. It is thus not ad hoc but the operator-theoretic embodiment of IQR's physical principle that geometry is state-conditioned.

Definition 4.392 (State-Conditioned Modal Weight)**.** Fix loci $\mathfrak{B}_x, \mathfrak{B}_{x'}$ and let $\{\,|\psi_n^{(x)}\rangle\,\}_n$ be a local orthonormal modal basis at x (e.g. eigenstates of the localized spectral operator $\hat{D}_x$; cf. Definition 4.341). For each mode n, define the bilocal Fourfold probe

$$\hat{M}_n(x,x') \;:=\; \hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})[\psi_n^{(x)*},\psi_n^{(x')}] \qquad \text{(see Definition 4.9).}$$

The *modal weight* of mode n between x and x' in the Quantum Holo-State $|\overset{\circ}{\mathbf{\Psi}}\rangle$ (Definition 1.27) is

$$\boxed{w_n(x,x') \;:=\; \Big\|\,\hat{M}_n(x,x')\,|\overset{\circ}{\mathbf{\Psi}}\rangle\,\Big\|.} \tag{4.189}$$

Interpretation. $w_n(x,x')$ measures the *entanglement-supported coherence* of the nth mode across (x,x'): large w_n indicates a strong, Holo-State–backed correlation pathway for that mode; small (or vanishing) w_n signals spectral/relational suppression in that channel. The weights therefore implement a *physically grounded, state-conditioned regularization* of spectral sums and suprema.

Proposition 4.393 (Basic Properties)**.** *With $\hat{M}_n$ and w_n as above:*

(i) Gauge/transport invariance
$w_n(x,x')$ is invariant under admissible rephasings of the channel operators and under changes of the parallel transporter within the same gauge class used in Definition 4.9.

(ii) Symmetry
$w_n(x,x') = w_n(x',x)$. *(Indeed, $\hat{M}_n(x',x) = \hat{M}_n(x,x')^\dagger$ and norms on $|\overset{\circ}{\mathbf{\Psi}}\rangle$ coincide.)*

(iii) Bounds and vanishing criterion
$0 \le w_n(x,x') \le \|\hat{M}_n(x,x')\|_{\mathrm{op}}$. *Moreover, $w_n(x,x') = 0$ iff $|\overset{\circ}{\mathbf{\Psi}}\rangle$ lies in the kernel of $\hat{M}_n(x,x')$, i.e. the nth bilocal channel is silent in the Holo-State between (x,x').*

(iv) Continuity
If $(x,x') \mapsto \hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})$ is continuous (resp. smooth) in the operator topology used, then $(x,x') \mapsto w_n(x,x')$ is continuous (resp. smooth).

(v) Degeneracy covariance
Within any degenerate eigenspace of $\hat{D}_x$, replacing $\{|\psi_n^{(x)}\rangle\}$ by a unitary-mixed basis leaves $\sum_{n\in\mathcal{E}} w_n(x,x')$ invariant.

Remark 4.394 (Normalisation Options (When Needed))**.** When a dimensionless weight in $[0,1]$ is convenient (e.g. to compare across loci or spectra of differing scale), one may use

$$\bar{w}_n(x,x') \;:=\; \frac{w_n(x,x')}{\sqrt{\Big\|\hat{M}_n(x,x)\,|\overset{\circ}{\mathbf{\Psi}}\rangle\Big\|\,\Big\|\hat{M}_n(x',x')\,|\overset{\circ}{\mathbf{\Psi}}\rangle\Big\|}},$$

provided the denominators are nonzero (otherwise define $\bar{w}_n = 0$). This preserves (i)–(iv) and yields $\bar{w}_n(x, x) = 1$.

Endogenous Regularization. In the spectral distance construction (Section 4.9.6), the family $\{w_n(x, x')\}_n$ enters as state-conditioned multipliers on modewise commutator data (e.g. spectral shear terms of Definition 4.386). Modes that are not coherently supported by $|\overset{\circ}{\Psi}\rangle$ between (x, x') are automatically suppressed, rendering the resulting spectral functionals well-posed on infinite spectra without imposing *ad hoc* cutoffs.

4.9.8 Wordwise State-conditioned Modal Weight and Endogenous Regularization

Definition 4.395 (Wordwise bilocal probe)**.** Fix loci $\mathfrak{B}_x, \mathfrak{B}_{x'}$, a path $\gamma : x \to x'$ with wordwise transporter $\mathcal{U}^{[\mathbf{W}]}(x' \leftarrow x; \gamma)$ (Sections 4.5 and 4.6.3), and a word $\mathbf{W}$ in the endogenous holosymmetric alphabet (Remarks 4.348 and 4.372). Let $\{\, |\psi^{(x)}_{n,[\mathbf{W}]}\rangle \,\}_n$ be a local orthonormal modal basis at x (e.g. an eigenbasis of the *wordwise* spectral operator $\mathbb{D}^{[\mathbf{W}]}_x$; Definition 4.345). The *word–dressed* bilocal Fourfold probe for mode n is

$$\begin{aligned} \hat{M}^{[\mathbf{W}]}_n(x, x'; \gamma) \;&:=\; \hat{G}^{[\mathbf{W}]}_\Lambda\Big(\mathfrak{B}_x, \mathfrak{B}_{x'}\Big)\Big[\, \psi^{(x)\,*}_{n,[\mathbf{W}]},\; \psi^{(x')}_{n,[\mathbf{W}],\gamma}\,\Big], \\ |\psi^{(x')}_{n,[\mathbf{W}],\gamma}\rangle \;&:=\; \mathcal{U}^{[\mathbf{W}]}(x' \leftarrow x; \gamma)\, |\psi^{(x)}_{n,[\mathbf{W}]}\rangle. \end{aligned} \tag{4.190}$$

where $\hat{G}^{[\mathbf{W}]}_\Lambda$ is the *wordwise* Fourfold correlator (Sections 4.2 and 4.6.3).

Definition 4.396 (Wordwise state–conditioned modal weight)**.** With $\hat{M}^{[\mathbf{W}]}_n$ as in (4.190), define the *wordwise modal weight* in the Quantum Holo–State $|\overset{\circ}{\Psi}\rangle$ by

$$\boxed{\; w^{[\mathbf{W}]}_n(x, x'; \gamma) \;:=\; \Big\| \hat{M}^{[\mathbf{W}]}_n(x, x'; \gamma)\; |\overset{\circ}{\Psi}\rangle \Big\|. \;} \tag{4.191}$$

When no path dependence is intended (e.g. in flat, simply connected neighbourhoods), we write $w^{[\mathbf{W}]}_n(x, x')$ and implicitly take the path that realises the $\inf_\gamma$ via wordwise parallel consistency (cf. Propositions 4.137 and 4.138).

Proposition 4.397 (Basic properties (word/gauge/transport invariance and regularity))**.** *For the weights* (4.191)*:*

(i) Word invariance
If $\mathbf{W} \mapsto \mathbf{W}'$*, then* $\hat{M}^{[\mathbf{W}']}_n = U_x^\dagger\, \hat{M}^{[\mathbf{W}]}_n\, U_{x'}$ *for charge–preserving fibre unitaries* $U_{(\cdot)}$ *(Remark 4.348), hence* $w^{[\mathbf{W}']}_n(x, x'; \gamma) = w^{[\mathbf{W}]}_n(x, x'; \gamma)$ *by unitary invariance of the norm and endogenous expectation invariance (Remark 4.372).*

(ii) Endpoint gauge covariance
Under endpoint gauge $g(x), g(x') \in \mathrm{U}(4)$ *compatible with the wordwise connection (Section 4.5), one has* $\hat{\mathcal{F}}^{[\mathbf{W}]}_\alpha \mapsto g\, \hat{\mathcal{F}}^{[\mathbf{W}]}_\alpha$ *and* $\mathcal{U}^{[\mathbf{W}]} \mapsto g(x')\, \mathcal{U}^{[\mathbf{W}]} g(x)^\dagger$*, so* $\hat{M}^{[\mathbf{W}]}_n \mapsto g(x) \hat{M}^{[\mathbf{W}]}_n g(x)^\dagger$ *and* $w^{[\mathbf{W}]}_n$ *is unchanged.*

(iii) Symmetry

By path reversal ($\gamma \mapsto \gamma^{-1}$) and unitarity of $\mathcal{U}^{[\mathbf{W}]}$, together with the adjoint rule for $\hat{G}^{[\mathbf{W}]}_{\Lambda}$, one has

$$w^{[\mathbf{W}]}_n(x,x';\gamma) = w^{[\mathbf{W}]}_n(x',x;\gamma^{-1}),$$
$$\hat{M}^{[\mathbf{W}]}_n(x',x;\gamma^{-1}) = \hat{M}^{[\mathbf{W}]}_n(x,x';\gamma)^\dagger.$$

(iv) Bounds and vanishing criterion

$0 \le w^{[\mathbf{W}]}_n(x,x';\gamma) \le \|\hat{M}^{[\mathbf{W}]}_n(x,x';\gamma)\|_{\mathrm{op}}$. *Moreover* $w^{[\mathbf{W}]}_n(x,x';\gamma) = 0$ *iff* $|\mathring{\mathbf{\Psi}}\rangle \in \ker \hat{M}^{[\mathbf{W}]}_n(x,x';\gamma)$.

(v) Continuity

If $(x,x') \mapsto \hat{G}^{[\mathbf{W}]}_{\Lambda}(\mathfrak{B}_x,\mathfrak{B}_{x'})$ is continuous (resp. smooth) in the operator topology used, then $(x,x') \mapsto w^{[\mathbf{W}]}_n(x,x')$ is continuous (resp. smooth).

(vi) Degeneracy covariance (wordwise)

If $\{|\psi^{(x)}_{n,j,[\mathbf{W}]}\rangle\}_j$ spans a degenerate eigenspace of $\mathbb{D}^{[\mathbf{W}]}_x$ (Definition 4.345), then for any unitary mixing within that eigenspace, $\sum_j w^{[\mathbf{W}]}_{n,j}(x,x')$ is invariant.

Proof. Let $w^{[\mathbf{W}]}_n(x,x';\gamma)$ be given by (4.191) with $\hat{M}^{[\mathbf{W}]}_n$ from (4.190). Write

$$\left(w^{[\mathbf{W}]}_n(x,x';\gamma)\right)^2 = \left\langle \mathring{\mathbf{\Psi}} \right| \left(\hat{M}^{[\mathbf{W}]}_n\right)^\dagger \hat{M}^{[\mathbf{W}]}_n \left| \mathring{\mathbf{\Psi}} \right\rangle. \tag{$*$}$$

(i) Word invariance. A charge–preserving word change $\mathbf{W} \mapsto \mathbf{W}'$ is implemented by fibre unitaries $U_x, U_{x'}$ so that $\hat{M}^{[\mathbf{W}']}_n = U_x^\dagger \hat{M}^{[\mathbf{W}]}_n U_{x'}$ (Remark 4.348). Then

$$\left(\hat{M}^{[\mathbf{W}']}_n\right)^\dagger \hat{M}^{[\mathbf{W}']}_n = U_{x'}^\dagger \left(\hat{M}^{[\mathbf{W}]}_n\right)^\dagger \hat{M}^{[\mathbf{W}]}_n U_{x'}.$$

Taking the Holo–State expectation and using invariance of endogenous expectations under charge–preserving conjugations (Remark 4.372) gives $\left(w^{[\mathbf{W}']}_n\right)^2 = \left(w^{[\mathbf{W}]}_n\right)^2$, hence $w^{[\mathbf{W}']}_n = w^{[\mathbf{W}]}_n$.

(ii) Endpoint gauge covariance. Let $g(x), g(x') \in \mathrm{U}(4)$ be compatible with the wordwise connection (Section 4.5). Then $\hat{\mathcal{F}}^{[\mathbf{W}]}_\alpha \mapsto g\,\hat{\mathcal{F}}^{[\mathbf{W}]}_\alpha$ and $\mathcal{U}^{[\mathbf{W}]} \mapsto g(x')\mathcal{U}^{[\mathbf{W}]}g(x)^\dagger$ (transport/gauge covariance), hence $\hat{M}^{[\mathbf{W}]}_n \mapsto g(x)\hat{M}^{[\mathbf{W}]}_n g(x)^\dagger$. As in (i),

$$\left(\hat{M}^{[\mathbf{W}]}_n\right)^\dagger \hat{M}^{[\mathbf{W}]}_n \mapsto g(x) \left(\hat{M}^{[\mathbf{W}]}_n\right)^\dagger \hat{M}^{[\mathbf{W}]}_n\, g(x)^\dagger,$$

and by the same expectation invariance (Remark 4.372) one obtains $w^{[\mathbf{W}]}_n$ unchanged.

(iii) Symmetry. By path reversal and unitarity of the transporter, $\mathcal{U}^{[\mathbf{W}]}(x \leftarrow x';\gamma^{-1}) = \left(\mathcal{U}^{[\mathbf{W}]}(x' \leftarrow x;\gamma)\right)^\dagger$ (Section 4.5), and by the adjoint rule of the wordwise correlator,

$$\hat{G}^{[\mathbf{W}]}_{\Lambda}(\mathfrak{B}_{x'},\mathfrak{B}_x)[\phi_2^*,\phi_1] = \left(\hat{G}^{[\mathbf{W}]}_{\Lambda}(\mathfrak{B}_x,\mathfrak{B}_{x'})[\phi_1^*,\phi_2]\right)^\dagger.$$

Applying these to (4.190) yields $\hat{M}^{[\mathbf{W}]}_n(x',x;\gamma^{-1}) = \hat{M}^{[\mathbf{W}]}_n(x,x';\gamma)^\dagger$ and therefore, by ($*$), $w^{[\mathbf{W}]}_n(x',x;\gamma^{-1}) = w^{[\mathbf{W}]}_n(x,x';\gamma)$.

(iv) Bounds and vanishing. By definition and the operator norm inequality,

$$0 \;\le\; w_n^{[\mathbf{W}]}(x,x';\gamma) = \big\|\hat{M}_n^{[\mathbf{W}]}\,|\mathring{\mathbf{\Psi}}\rangle\big\| \;\le\; \big\|\hat{M}_n^{[\mathbf{W}]}\big\|_{\mathrm{op}}.$$

Moreover $w_n^{[\mathbf{W}]}(x,x';\gamma)=0$ iff $\hat{M}_n^{[\mathbf{W}]}\,|\mathring{\mathbf{\Psi}}\rangle = 0$, i.e. the Holo–State lies in the kernel.

(v) Continuity. Assume $(x,x') \mapsto \hat{G}_\Lambda^{[\mathbf{W}]}(\mathfrak{B}_x,\mathfrak{B}_{x'})$ is continuous (resp. smooth) in the operator topology and that γ is chosen by the wordwise parallel–consistency prescription on a normal neighbourhood (Propositions 4.137 and 4.138). Then $(x,x') \mapsto \hat{M}_n^{[\mathbf{W}]}(x,x';\gamma)$ is continuous (resp. smooth) by composition (Lemma 4.270), and so is $(x,x') \mapsto \big\|\hat{M}_n^{[\mathbf{W}]}\,|\mathring{\mathbf{\Psi}}\rangle\big\|$.

(vi) Degeneracy covariance. Let $\{|\psi^{(x)}_{n,j,[\mathbf{W}]}\rangle\}_j$ be an orthonormal basis of a degenerate eigenspace of $\mathbb{D}_x^{[\mathbf{W}]}$ (Definition 4.345), and let $|\psi'_{n,a}\rangle = \sum_j U_{aj}|\psi_{n,j}\rangle$ be any unitary mixing. Define $v_j := \hat{M}_{n,j}^{[\mathbf{W}]}\,|\mathring{\mathbf{\Psi}}\rangle$ and $v'_a := \hat{M}_{n,a}^{[\mathbf{W}]\prime}\,|\mathring{\mathbf{\Psi}}\rangle$. By linearity of the bilocal probe in each slot (Section 4.2), $v'_a = \sum_j U_{aj}\,v_j$. Hence

$$\sum_a \big\|v'_a\big\|^2 = \sum_a \Big\langle \sum_j U_{aj}v_j,\; \sum_k U_{ak}v_k \Big\rangle = \sum_{j,k} \Big(\sum_a U_{aj}\overline{U}_{ak}\Big)\langle v_j,v_k\rangle = \sum_j \|v_j\|^2,$$

since U is unitary. Using $(*)$, this proves that $\sum_j \big(w_{n,j}^{[\mathbf{W}]}(x,x')\big)^2$ is invariant under unitary mixing in the degenerate eigenspace. In particular, when the cross terms $\langle v_j,v_k\rangle$ vanish for $j\neq k$ (e.g. under the standard modal normalisation where distinct degenerate rays probe orthogonal bilocal channels; cf. Lemma 4.270), the linear sum $\sum_j w_{n,j}^{[\mathbf{W}]}(x,x')$ is also invariant. □

Remark 4.398 (Normalised wordwise weights)**.** When a dimensionless weight in $[0,1]$ is preferable, set

$$\bar{w}_n^{[\mathbf{W}]}(x,x';\gamma) := \frac{w_n^{[\mathbf{W}]}(x,x';\gamma)}{\sqrt{\big\|\hat{M}_n^{[\mathbf{W}]}(x,x;\mathrm{id})\,|\mathring{\mathbf{\Psi}}\rangle\big\|\,\big\|\hat{M}_n^{[\mathbf{W}]}(x',x';\mathrm{id})\,|\mathring{\mathbf{\Psi}}\rangle\big\|}},$$

whenever the denominator is nonzero; otherwise define $\bar{w}_n^{[\mathbf{W}]}=0$. This preserves the properties in Proposition 4.397 and gives $\bar{w}_n^{[\mathbf{W}]}(x,x;\mathrm{id})=1$.

Definition 4.399 (Endogenous (wordwise) regularisation of spectral aggregates)**.** Let $\mathcal{S}_n^{[\mathbf{W}]}(x,x')$ denote the *wordwise* spectral shear (Definition 4.381 and eq. (4.162)). Define the state–conditioned, wordwise–regularised shear aggregate by

$$\boxed{\mathcal{S}_{\mathrm{reg}}^{[\mathbf{W}]}(x,x') \;:=\; \sum_n \varpi_n\,\bar{w}_n^{[\mathbf{W}]}(x,x')\,\frac{\mathcal{S}_n^{[\mathbf{W}]}(x,x')}{(1+|n|)^k},} \tag{4.192}$$

where k is any exponent compatible with wordwise spectral smoothness (Definition 4.383 and Theorem 4.384) and $\varpi_n \in [0,1]$ are optional scheme weights (e.g. compact support in n) chosen within the same admissibility class. The state–dependence of $\bar{w}_n^{[\mathbf{W}]}$ suppresses modes that are not coherently supported by $|\mathring{\mathbf{\Psi}}\rangle$, yielding well–posed, physically grounded sums without ad hoc cutoffs.

Corollary 4.400 (Compatibility with correlator and spectral distances)**.** *The wordwise weights* $\bar{w}_n^{[\mathbf{W}]}$ *may be used as multipliers in (i) the Connes–type wordwise spectral distance construction (Definition 4.389) by restricting the supremum to observables* Φ *whose mode content is weighted by* $\bar{w}_n^{[\mathbf{W}]}$*, and (ii) correlator–based distance filters (Definition 4.379) by selecting probes with large* $\bar{w}_n^{[\mathbf{W}]}$*. In flat, simply–connected regions these insertions are word– and path–independent by wordwise parallel consistency (Proposition 4.137).*

Remark 4.401 (Twofold reduction)**.** Under the wordwise qubit reduction (Sections 4.6.2 and 4.7.5), only the Twofold indices contribute; the family $\{\bar{w}_n^{[\mathbf{W}]}\}$ collapses to the qubit modes on $S_x^{[\mathbf{W}]}$, reproducing the standard state–conditioned weighting of qubit diagnostics inside the protected sector.

4.9.9 The Relational Spectral Distance

Definition 4.402 (Relational Spectral Distance)**.** For loci $x, x' \in \mathcal{X}_q$, let $\{\mathcal{S}_n(x,x')\}_n$ be the spectral shear observables (Definition 4.386 and eq. (4.160)) evaluated on a fixed local modal basis at x (e.g. an eigenbasis of $\hat{D}_x$; Definition 4.341), and let $w_n(x,x')$ be the corresponding state–conditioned modal weights (Definition 4.392). The *relational spectral distance* is the weighted root–mean–square aggregate

$$\boxed{d_{\mathrm{spec}}(x,x') \;:=\; \left(\sum_n w_n^2(x,x') \left|\mathcal{S}_n(x,x')\right|^2 \right)^{1/2}.} \tag{4.193}$$

The construction functionally instantiates three ingredients of IQR: (i) the differential generator $\hat{D}$ (Definition 4.341), (ii) the relational integrator $\hat{G}_\Lambda$ (Definition 4.9), and (iii) the evaluating Quantum Holo-State $|\mathring{\mathbf{\Psi}}\rangle$ (Definition 1.27).

Proposition 4.403 (Well-Definedness and Basic Properties)**.** *Assume the Axiom of Spectral Smoothness (Definition 4.342) and the Principle of Finite Entanglement Capacity (Principle 4.1), with modal-weight decay as in eq. (4.199). Then:*

(i) (Finiteness) $d_{\mathrm{spec}}(x,x') < \infty$ *for all* $x, x' \in \mathcal{X}_q$.

(ii) (Nonnegativity and symmetry) $d_{\mathrm{spec}}(x,x') \geq 0$ *and* $d_{\mathrm{spec}}(x,x') = d_{\mathrm{spec}}(x',x)$.

(iii) (Coincidence) $d_{\mathrm{spec}}(x,x) = 0$.

(iv) (Invariance) d_{spec} *is invariant under admissible reindexings and the duality involution (Proposition 1.8), and under parallel–transport gauge changes used in Definition 4.9.*

Sketch. *By Definition 4.342,* $|\mathcal{S}_n(x,x')| \leq C(1+n)^k$ *uniformly near the diagonal. By eq. (4.199),* $w_n(x,x') \leq C'(1+n)^{-p}$ *with* $p > k + \frac{1}{2}$*. Hence* $\sum_n w_n^2 |\mathcal{S}_n|^2 \leq (CC')^2 \sum_n (1+n)^{-2(p-k)} < \infty$*, proving (i). Items (ii)–(iv) follow from the norm definition,* $\hat{M}_n(x',x) = \hat{M}_n(x,x')^\dagger$*, and the invariances of* $\mathcal{S}_n$ *and* w_n *(cf. Proposition 4.393).*

Remark 4.404 (Mechanism of Endogenous Regularization)**.** In eq. (4.193), high–index modes lacking Holo-State support are downweighted by $w_n(x,x')$, while commutator growth is controlled by spectral smoothness (Definition 4.342). Heuristically, this acts as a state–conditioned cutoff built into the theory's operator structure.

4.9.10 Wordwise Relational Spectral Distance (Holosymmetric Realization)

Throughout, fix admissible endpoint device words $\mathbf{W}_1$ at x and $\mathbf{W}_2$ at x' with RR elements $\mathsf{W}_{1,\chi\leftarrow\chi}(x)$, $\mathsf{W}_{2,\chi'\leftarrow\chi'}(x')$, and use the wordwise covariant kinematics and parallel transport from Section 4.5. Wordwise spectral operators and their regularity properties are as in Sections 4.9.4 and 4.9.7. State-conditioned modal weights use the endogenous construction in Section 4.9.8.

Definition 4.405 (Wordwise local spectral frame at x)**.** Let $\mathbb{D}^{[\mathbf{W}_1]}(x)$ be the wordwise spectral operator at x (cf. Section 4.9.4). Choose an orthonormal eigenbasis $\{|n\rangle_x^{[\mathbf{W}_1]}\}_n$ with projectors $\Pi_n^{(x)[\mathbf{W}_1]} := |n\rangle_x^{[\mathbf{W}_1]}\langle n|_x^{[\mathbf{W}_1]}$.

Definition 4.406 (Wordwise spectral shear observables)**.** Let $\mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x \leftarrow x';\gamma)$ be the wordwise parallel transporter along a path γ (Section 4.5). Define the transported spectral operator

$$\widetilde{\mathbb{D}}^{[\mathbf{W}_2]}_{x'\to x}(\gamma) := \mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x\leftarrow x';\gamma)^\dagger\, \mathbb{D}^{[\mathbf{W}_2]}(x')\, \mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x\leftarrow x';\gamma),$$

and the wordwise spectral difference $\Delta\mathbb{D}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma) := \widetilde{\mathbb{D}}^{[\mathbf{W}_2]}_{x'\to x}(\gamma) - \mathbb{D}^{[\mathbf{W}_1]}(x)$. For the Holo-State $|\mathring{\mathbf{\Psi}}\rangle$ (Definition 1.27), set the mode-n scalar

$$\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma) := \Big\langle \mathring{\mathbf{\Psi}}\Big|\, \Pi_n^{(x)[\mathbf{W}_1]}\, \Delta\mathbb{D}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma)\, \Pi_n^{(x)[\mathbf{W}_1]}\, \Big|\mathring{\mathbf{\Psi}}\Big\rangle. \tag{4.194}$$

This is the wordwise counterpart of Definition 4.386 and eq. (4.160).

Definition 4.407 (Wordwise modal weights)**.** Let $w_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')$ be the endogenous state-conditioned weights defined in Section 4.9.8; these are word/gauge invariant and obey the decay properties of eq. (4.199) (see Proposition 4.393).

Definition 4.408 (Wordwise Relational Spectral Distance)**.** With the objects above, define the weighted RMS aggregate

$$\boxed{d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \;:=\; \Bigg(\sum_n \Big(w_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\Big)^2 \Big|\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\Big|^2\Bigg)^{1/2}.} \tag{4.195}$$

Lemma 4.409 (Path/gauge covariance and holonomy control)**.** *Under endpoint gauges and word changes, the collection $\{\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma)\}_n$ transforms covariantly and $\big|\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}\big|$ is invariant. Changing γ deforms $\widetilde{\mathbb{D}}^{[\mathbf{W}_2]}_{x'\to x}$ by the holonomy of the modal connection; the induced variation of $\mathcal{S}_n$ is $O(\Phi(\gamma))$ with Φ as in Proposition 4.241. In flat patches ($\mathcal{F}\equiv 0$) the quantity is path independent.*

Proposition 4.410 (Well-Definedness and Basic Properties (Wordwise))**.** *Assume the Axiom of Spectral Smoothness (wordwise) as in Section 4.9.4 and the Principle of Finite Entanglement Capacity with eq. (4.199). Then, for all $x,x'\in\mathcal{X}_q$:*

(i) Finiteness. $d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') < \infty$.

(ii) Nonnegativity and symmetry. $d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \geq 0$ *and equals* $d_{\mathrm{spec}}^{[\mathbf{W}_2,\mathbf{W}_1]}(x',x)$.

(iii) Coincidence. $d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x)=0$.

(iv) Word/gauge invariance. $d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}$ *is invariant under admissible endpoint word changes and endpoint gauges, and is independent of* γ *on flat patches (up to* $O(\Phi(\gamma))$ *otherwise).*

Sketch. *Spectral smoothness gives* $|\mathcal{S}_n|\le C(1+n)^k$ *(uniform locally); modal-weight decay gives* $w_n\le C'(1+n)^{-p}$ *with* $p>k+\frac{1}{2}$*, so* $\sum_n w_n^2|\mathcal{S}_n|^2<\infty$ *proving (i). Items (ii)–(iii) follow by construction and* $\Delta\mathbb{D}(x,x)=0$*. For (iv), covariance of* $\mathcal{U}$ *and unitary invariance of the fibre expectation (cf. Sections 4.5 and 4.6) imply invariance under words/gauges. Path dependence is controlled by Lemma 4.409 and Proposition 4.241.*

Proposition 4.411 (Equivalence to Exogenous Spectral Distance)**.** *Let* $d_{\mathrm{spec}}(x,x')$ *be the exogenous distance of Definition 4.402. Then for any admissible endpoint words* $\mathbf{W}_1,\mathbf{W}_2$*,*

$$d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \;=\; d_{\mathrm{spec}}(x,x').$$

Proof sketch. *Choose* $\{|n\rangle_x\}_n$ *in Definition 4.402 to be the image of* $\{|n\rangle_x^{[\mathbf{W}_1]}\}_n$ *under the inverse word rotation at* x*. The exogenous shear* $\mathcal{S}_n(x,x')$ *coincides with the wordwise* $\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')$ *because transporting and evaluating in the Holo-State commute with endpoint word insertions (unitary covariance as in Section 4.5 and the adjoint reciprocity of the Fourfold correlator from Section 4.1.2). The modal weights agree by Definition 4.407 and Proposition 4.393. Taking the weighted RMS then yields equality.*

Corollary 4.412 (Twofold reduction in the duality-collapse regime)**.** *In the regime of Definition 4.87,* $d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')$ *reduces to the Twofold spectral distance computed on the* $\{0,1\}$ *slice, with transport and weights restricted to the Twofold sector (cf. Section 4.9.7).*

Lemma 4.413 (Holonomy–Lipschitz bound for wordwise spectral shear)**.** *Let* γ_1,γ_2 *be two piecewise-smooth paths from* x' *to* x *and set the loop*

$$\Delta\gamma \;:=\; \gamma_2\circ\gamma_1^{-1}. \tag{4.196}$$

With the wordwise transporter $\mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(\cdot)$ *and holonomy functional* $\Phi(\cdot)$ *from Section 4.5 and the holonomy bound in Proposition 4.241, there exists a constant* $c_\nabla>0$ *(depending only on the local connection data) such that*

$$\Big\|\mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x\leftarrow x';\gamma_2) \;-\; \mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x\leftarrow x';\gamma_1)\Big\| \;\le\; c_\nabla\,\Phi(\Delta\gamma).$$

Then, under the Wordwise Spectral Smoothness axiom (Axiom 4.418), the mode-n shear obeys

$$\Big|\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma_2)-\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma_1)\Big| \;\le\; L_n(x)\,\Phi(\Delta\gamma),\qquad L_n(x)\;\le\;2\,c_\nabla\,C\,\Big(\lambda_n^{[\mathbf{W}_1]}(x)\Big)^k, \tag{4.197}$$

where C,k *are as in Axiom 4.418 and* $\lambda_n^{[\mathbf{W}_1]}(x)$ *are the localized eigenvalues of* $\mathbb{D}^{[\mathbf{W}_1]}$ *at* x *(Definition 4.405).*

Proof sketch. Write $\widetilde{\mathbb{D}}_i:=\mathcal{U}_i^\dagger\,\mathbb{D}^{[\mathbf{W}_2]}(x')\,\mathcal{U}_i$ for $\mathcal{U}_i=\mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x\leftarrow x';\gamma_i)$ and $\Delta\mathbb{D}_i:=\widetilde{\mathbb{D}}_i-\mathbb{D}^{[\mathbf{W}_1]}(x)$. Then

$$\mathcal{S}_n(\gamma_i)=\Big\langle\mathring{\mathbf{\Psi}}\Big|\,\Pi_n^{(x)[\mathbf{W}_1]}\,\Delta\mathbb{D}_i\,\Pi_n^{(x)[\mathbf{W}_1]}\,\Big|\mathring{\mathbf{\Psi}}\Big\rangle.$$

Hence $\left|\mathcal{S}_n(\gamma_2) - \mathcal{S}_n(\gamma_1)\right| \le \|\Pi_n\,(\Delta\mathbb{D}_2 - \Delta\mathbb{D}_1)\,\Pi_n\| \le \|\Delta\mathbb{D}_2 - \Delta\mathbb{D}_1\|$. Using $\Delta\mathbb{D}_2 - \Delta\mathbb{D}_1 = \mathcal{U}_1^\dagger\Big((\mathcal{U}_1\mathcal{U}_2^\dagger)\,\mathbb{D}^{[\mathbf{W}_2]}(x') - \mathbb{D}^{[\mathbf{W}_2]}(x')\,(\mathcal{U}_1\mathcal{U}_2^\dagger)\Big)\mathcal{U}_1$, we get $\|\Delta\mathbb{D}_2 - \Delta\mathbb{D}_1\| \le 2\,\|\mathbb{D}^{[\mathbf{W}_2]}(x')\|\,\|\mathcal{U}_2 - \mathcal{U}_1\|$. The holonomy bound from Proposition 4.241 yields $\|\mathcal{U}_2 - \mathcal{U}_1\| \le c_\nabla\,\Phi(\Delta\gamma)$; spectral smoothness gives $\|\mathbb{D}^{[\mathbf{W}_2]}(x')\| \lesssim C\,(\lambda_n^{[\mathbf{W}_1]})^k$ on the n-block, producing (4.197). □

Corollary 4.414 (Holonomy–Lipschitz bound for the wordwise spectral distance)**.** *With the notation of Lemma 4.413 and modal weights $w_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')$ from Definition 4.407,*

$$\left|d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma_2) - d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma_1)\right| \;\le\; \Phi(\Delta\gamma)\left(\sum_n \Big(w_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\,L_n(x)\Big)^2\right)^{1/2}. \tag{4.198}$$

In particular, under Axiom 4.418 and Principle 4.2, the right-hand side is finite, so $d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}$ is Lipschitz in the holonomy size. On flat patches ($\mathcal{F} \equiv 0$ so $\Phi(\Delta\gamma) = 0$), the distance is path independent.

Proof sketch. Apply Minkowski/triangle inequality to the RMS in Definition 4.408 and use (4.197); finiteness follows from Proposition 4.410 (decay of weights) and the growth bound on $L_n(x)$. □

4.9.11 Endogenous Finiteness of Spectral Distance

We now show that the spectral distance $d_{\mathrm{spec}}(x,x')$ of Definition 4.402 is finite for all loci $x, x' \in \mathcal{X}_q$, with no external ultraviolet cutoff. The argument uses (i) the polynomial control of spectral shear from the Spectral Operator, and (ii) the state-conditioned decay of modal weights.

Axiom 4.415 (Spectral Smoothness)**.** For an admissible Spectral Operator $\hat{D}$ (Definitions 4.341 and 4.343), there exist constants $C > 0$ and $k \in \mathbb{N}$ such that the spectral shear observable of mode n,

$$\mathcal{S}_n(x,x') \;:=\; \left\langle \mathring{\mathbf{\Psi}} \,\middle|\, \left[\hat{D}, \hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})[\psi_n^*,\psi_n]\right] \,\middle|\, \mathring{\mathbf{\Psi}} \right\rangle \quad (\text{cf. } \textit{Definition } 4.386),$$

obeys the polynomial bound

$$|\mathcal{S}_n(x,x')| \;\le\; C\,\lambda_n^k,$$

where $\{\lambda_n\}$ are the (localised) eigenvalues of $\hat{D}$ at x (Definition 4.341).

Principle 4.1 (Finite Entanglement Capacity)**.** *The Holo-State $|\mathring{\mathbf{\Psi}}\rangle$ cannot support arbitrarily fine coherent channels. Equivalently, the modal weights*

$$w_n(x,x') \;:=\; \left\|\hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})[\psi_n^*,\psi_n]\,|\mathring{\mathbf{\Psi}}\rangle\right\| \quad (\textit{cf. Definition } 4.392)$$

decay with spectral level as

$$w_n(x,x') \;\le\; \frac{C'}{\lambda_n^\alpha} \qquad \textit{for some } C' > 0 \textit{ and } \alpha > k + \tfrac{1}{2}, \tag{4.199}$$

i.e. eq. (4.189) exhibits state-conditioned suppression at high frequency.

Proposition 4.416 (Endogenous Spectral Regularization)**.** *Under Axiom 4.415 and Principle 4.1, the spectral distance*

$$d_{\text{spec}}(x,x') \;=\; \Big(\sum\nolimits_n w_n^2(x,x')\,|\mathcal{S}_n(x,x')|^2\Big)^{1/2} \quad (\textit{cf. Definition } 4.402, (4.193))$$

is finite for all $x, x' \in \mathcal{X}_q$.

Proof. By Axiom 4.415 and (4.199),

$$w_n^2\,|\mathcal{S}_n|^2 \;\le\; (CC')^2\,\lambda_n^{-2(\alpha-k)}.$$

Hence

$$d_{\text{spec}}^2(x,x') \;\le\; (CC')^2 \sum\nolimits_n \lambda_n^{-2(\alpha-k)}.$$

Since $\alpha > k + \frac{1}{2}$, the exponent satisfies $2(\alpha-k) > 1$, and the series converges by comparison with a p-series (using the monotone ordering of λ_n ensured by the discrete spectrum in Definition 4.343). Thus $d_{\text{spec}}(x,x') < \infty$. □

Remark 4.417 (Counting-Function Refinement)**.** If the eigenvalue counting function satisfies a Weyl-type bound $N(\Lambda) \lesssim \Lambda^{d_s}$ (spectral dimension d_s for $\hat{D}$), then the same proof shows convergence already for

$$\alpha \;>\; k + \tfrac{d_s}{2},$$

with Proposition 4.416 recovered by the special case $d_s = 1$.

Interpretation. The factor $|\mathcal{S}_n|$ probes differential incompatibility (via the commutator with $\hat{D}$), while w_n encodes the Holo-State's support for that mode across (x,x'). Their product is summable: high-frequency shear is tamed not by a hand-imposed cutoff but by the theory's own operator- and state-level structure.

This result ensures that every geometric quantity defined using d_{spec} inherits the same internal finiteness. As such, the *Primordial Quantum Singularity*, which is defined via the collapse of all relational observables (including d_{spec}), is not only operationally well-motivated but also *mathematically well-defined*.

4.9.12 Endogenous Finiteness of Wordwise Spectral Distance

We now show that the wordwise spectral distance $d_{\text{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')$ of Definition 4.408 is finite for all loci $x, x' \in \mathcal{X}_q$, with no external ultraviolet cutoff. The argument uses (i) polynomial control of the *wordwise* spectral shear (Definition 4.406) and (ii) the endogenous, state-conditioned decay of *wordwise* modal weights (Definition 4.407).

Axiom 4.418 (Wordwise Spectral Smoothness)**.** For an admissible wordwise spectral operator $\mathbb{D}^{[\mathbf{W}]}$ (cf. Section 4.9.4), there exist constants $C > 0$ and $k \in \mathbb{N}$ such that the mode-n wordwise spectral shear,

$$\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \quad (\text{cf. } \textit{Definition } 4.406),$$

obeys the polynomial bound

$$\Big|\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\Big| \;\le\; C\,\Big(\lambda_n^{[\mathbf{W}_1]}(x)\Big)^k,$$

where $\{\lambda_n^{[\mathbf{W}_1]}(x)\}_n$ are the localized eigenvalues of $\mathbb{D}^{[\mathbf{W}_1]}$ at x (Definition 4.405).

Principle 4.2 (Wordwise Finite Entanglement Capacity)**.** *The Holo-State* $|\mathring{\mathbf{\Psi}}\rangle$ *cannot support arbitrarily fine* wordwise *coherent channels. Equivalently, the wordwise modal weights (Definition 4.407)*

$$w_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \;:=\; \Big\| \Pi_n^{(x)[\mathbf{W}_1]}\, \hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})\, \Pi_n^{(x)[\mathbf{W}_1]}\, |\mathring{\mathbf{\Psi}}\rangle \Big\|$$

decay with spectral level as

$$w_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \;\le\; \frac{C'}{\Big(\lambda_n^{[\mathbf{W}_1]}(x)\Big)^{\alpha}} \qquad \text{for some } C'>0 \text{ and } \alpha > k+\tfrac{1}{2}, \tag{4.200}$$

i.e. the state-conditioned suppression at high frequency (cf. eq. (4.199) and Proposition 4.393) holds in the wordwise frame at x*.*

Lemma 4.419 (Wordwise transport identity for the spectral operator)**.** *Let* $\gamma:[0,1]\to\mathcal{X}_q$ *be a piecewise-smooth path from* $x'=\gamma(0)$ *to* $x=\gamma(1)$*, with wordwise transporter* $\mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x\leftarrow x';\gamma) =: U(1)$ *and* $U(0)=\mathbf{1}$ *(Section 4.5). With the connection generator* A_s *along* γ *(so* $\partial_s U(s) = -A_s U(s)$*) and the wordwise spectral operator* $\mathbb{D}^{[\mathbf{W}(s)]}(x_s)$ *evaluated at* $x_s=\gamma(s)$*, one has the Duhamel-type identity*

$$U(1)^\dagger\, \mathbb{D}^{[\mathbf{W}_2]}(x')\, U(1) - \mathbb{D}^{[\mathbf{W}_1]}(x) \;=\; \int_0^1 U(s)^\dagger \Big(\partial_s \mathbb{D}^{[\mathbf{W}(s)]}(x_s) - [A_s, \mathbb{D}^{[\mathbf{W}(s)]}(x_s)] \Big) U(s)\, ds. \tag{4.201}$$

Proof. Set $F(s) := U(s)^\dagger\, \mathbb{D}^{[\mathbf{W}(s)]}(x_s)\, U(s)$. Using $\partial_s U^\dagger = U^\dagger A_s^\dagger$ and $\partial_s \mathbb{D} = \partial_s \mathbb{D}$, we get $\partial_s F = U^\dagger \big(\partial_s \mathbb{D} - [A_s,\mathbb{D}]\big) U$. Integrate $s\in[0,1]$ and note $F(1)-F(0)$ is the LHS of (4.201). (Unitarity of U yields $A_s^\dagger = -A_s$.) □

Lemma 4.420 (Band-polynomial localization)**.** *Let* $\{\Pi_n^{(x)[\mathbf{W}_1]}\}_n$ *be the spectral projectors of* $\mathbb{D}^{[\mathbf{W}_1]}(x)$ *(Definition 4.405) with eigenvalues* $\lambda_n^{[\mathbf{W}_1]}(x)$*. Suppose an operator* B *belongs to the admissible wordwise class at* x *so that the iterated commutators* $\mathrm{ad}_{\mathbb{D}}^j(B)$ *are bounded for* $0\le j\le k$ *with constants uniform on a patch (cf. admissibility in Section 4.9.4). Then*

$$\Big\| \Pi_n^{(x)[\mathbf{W}_1]}\, B\, \Pi_n^{(x)[\mathbf{W}_1]} \Big\| \;\le\; C_B \Big(1+\lambda_n^{[\mathbf{W}_1]}(x)\Big)^k. \tag{4.202}$$

Proof idea. Write $R=(1+D^2)^{-1/2}=f(D)$ with $D=\mathbb{D}^{[\mathbf{W}_1]}(x)$. To control the commutator with a *function* of D, we use the Helffer–Sjöstrand functional calculus: for an almost-analytic extension $\tilde f$ one has

$$[f(D),B] = \frac{1}{\pi}\int_{\mathbb{C}} \partial_{\bar z}\tilde f(z)\,(z-D)^{-1}[D,B]\,(z-D)^{-1}\, d^2z,$$

so $\|[R,B]\| \lesssim \|[D,B]\|$ by resolvent bounds [115, 397, 398, 399, 400]. This is precisely why Helffer–Sjöstrand is needed: it reduces commutators of *functions* of D (e.g. R) to commutators with D itself, which are controlled by admissibility. With $\|[R,B]\|\lesssim\|[D,B]\|$ in hand, expand B in the ad_D-filtration and iterate: each application of ad_R is bounded by a finite linear combination of ad_D^j, $1\le j\le k$ (again via Helffer–Sjöstrand), while resolvent estimates on the unit spectral slice around $\lambda_n^{[\mathbf{W}_1]}(x)$ control the band terms [115, 397]. Admissibility provides uniform bounds on $\|\mathrm{ad}_D^j(B)\|$ for $j\le k$, yielding the stated polynomial growth in $\lambda_n^{[\mathbf{W}_1]}(x)$.

Theorem 4.421 (Wordwise Spectral Smoothness)**.** *For an admissible wordwise spectral operator* $\mathbb{D}^{[\mathbf{W}]}$ *(cf. Section 4.9.4), there exist constants* $C > 0$ *and* $k \in \mathbb{N}$ *such that the mode-n wordwise spectral shear*

$$\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \quad (\textit{cf. Definition 4.406})$$

obeys the polynomial bound

$$\left|\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\right| \;\le\; C\left(\lambda_n^{[\mathbf{W}_1]}(x)\right)^k,$$

where $\{\lambda_n^{[\mathbf{W}_1]}(x)\}_n$ *are the localized eigenvalues of* $\mathbb{D}^{[\mathbf{W}_1]}$ *at* x *(Definition 4.405).*

Proof. By definition (Definition 4.406),

$$\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') = \left\langle \mathring{\mathbf{\Psi}}\right| \Pi_n^{(x)[\mathbf{W}_1]}\, \Delta\mathbb{D}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\, \Pi_n^{(x)[\mathbf{W}_1]} \left|\mathring{\mathbf{\Psi}}\right\rangle,$$

where $\Delta\mathbb{D}^{[\mathbf{W}_1,\mathbf{W}_2]}$ is the wordwise spectral difference used in Definition 4.406. Hence

$$\left|\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\right| \;\le\; \left\|\Pi_n^{(x)[\mathbf{W}_1]}\, \Delta\mathbb{D}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\, \Pi_n^{(x)[\mathbf{W}_1]}\right\|. \tag{4.203}$$

Apply the transport identity (4.201) from Lemma 4.419 along a path γ from x' to x:

$$\Delta\mathbb{D}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') = \int_0^1 U(s)^\dagger \Big(\partial_s \mathbb{D}^{[\mathbf{W}(s)]}(x_s) - [A_s, \mathbb{D}^{[\mathbf{W}(s)]}(x_s)]\Big) U(s)\, ds.$$

Taking the band norm and using unitary invariance,

$$\left\|\Pi_n\, \Delta\mathbb{D}\, \Pi_n\right\| \;\le\; \int_0^1 \left\|\Pi_n(s)\left(\partial_s \mathbb{D}^{[\mathbf{W}(s)]}(x_s) - [A_s, \mathbb{D}^{[\mathbf{W}(s)]}(x_s)]\right)\Pi_n(s)\right\| ds,$$

where $\Pi_n(s)$ is the n-projector of $\mathbb{D}^{[\mathbf{W}(s)]}(x_s)$ pulled back to x by $U(s)$. By admissibility (Section 4.9.4), both $B_s^{(1)} := \partial_s \mathbb{D}^{[\mathbf{W}(s)]}(x_s)$ and $B_s^{(2)} := [A_s, \mathbb{D}^{[\mathbf{W}(s)]}(x_s)]$ lie in the wordwise class with bounded commutators up to order k, uniformly on a patch. Therefore, by Lemma 4.420,

$$\left\|\Pi_n(s)\, B_s^{(j)}\, \Pi_n(s)\right\| \;\le\; C_j\left(1 + \lambda_n^{[\mathbf{W}_1]}(x)\right)^k, \qquad j = 1,2,$$

where we also use spectral continuity along γ to replace $\lambda_n^{[\mathbf{W}(s)]}(x_s)$ by an equivalent $\lambda_n^{[\mathbf{W}_1]}(x)$ up to a uniform multiplicative constant on the patch. Integrating in s and absorbing path-length/patch constants into C yields

$$\left\|\Pi_n^{(x)[\mathbf{W}_1]}\, \Delta\mathbb{D}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\, \Pi_n^{(x)[\mathbf{W}_1]}\right\| \;\le\; C\left(1 + \lambda_n^{[\mathbf{W}_1]}(x)\right)^k.$$

Combine with (4.203) to conclude the stated bound. □

Editorial. Replace citations to the axiom Axiom 4.418 by the theorem Theorem 4.421 wherever a proof-dependent statement is required; statements that only need a hypothesis may continue to cite Axiom 4.418 as an optional assumption.

Proposition 4.422 (Wordwise Endogenous Spectral Regularization)**.** *Under Theorem 4.421 and Principle 4.2, the wordwise spectral distance*

$$d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \;=\; \left(\sum\nolimits_n \left(w_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\right)^2 \left|\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\right|^2\right)^{1/2}$$

(*cf. Definition 4.408and eq.* (4.195)) (4.204)

is finite for all $x, x' \in \mathcal{X}_q$.

Proof. By Theorem 4.421 and (4.200),

$$\left(w_n^{[\mathbf{W}_1,\mathbf{W}_2]}\right)^2 \left|\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}\right|^2 \;\le\; (CC')^2 \left(\lambda_n^{[\mathbf{W}_1]}(x)\right)^{-2(\alpha-k)}.$$

Hence

$$\left(d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\right)^2 \;\le\; (CC')^2 \sum\nolimits_n \left(\lambda_n^{[\mathbf{W}_1]}(x)\right)^{-2(\alpha-k)}.$$

Since $\alpha > k + \frac{1}{2}$, the exponent satisfies $2(\alpha - k) > 1$, and the series converges by comparison with a p-series, using the monotone ordering/discreteness of $\{\lambda_n^{[\mathbf{W}_1]}(x)\}_n$ from the admissibility of $\mathbb{D}^{[\mathbf{W}_1]}$ (cf. Section 4.9.4). Thus $d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') < \infty$. □

Remark 4.423 (Wordwise Counting-Function Refinement)**.** If the eigenvalue counting function for $\mathbb{D}^{[\mathbf{W}_1]}$ satisfies a Weyl-type bound $N^{[\mathbf{W}_1]}(\Lambda) \lesssim \Lambda^{d_s}$ (spectral dimension d_s), then the same proof shows convergence already for

$$\alpha \;>\; k + \tfrac{d_s}{2},$$

where k is the exponent provided by Theorem 4.421. The case $d_s = 1$ recovers Proposition 4.422.

Interpretation (wordwise). $\left|\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}\right|$ probes differential incompatibility via the wordwise spectral difference (Definition 4.406), while $w_n^{[\mathbf{W}_1,\mathbf{W}_2]}$ encodes the Holo-State's support for that mode across (x, x') (Definition 4.407). Their product is summable: high-frequency shear is tamed not by a hand-imposed cutoff but by the operator- and state-level structure of the holosymmetric wordwise calculus.

Equivalence to the exogenous statement. By Proposition 4.411, finiteness of $d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}$ is equivalent to finiteness of d_{spec} in Definition 4.402; every geometric quantity built from the spectral distance inherits the same *endogenous* finiteness in either presentation.

4.9.13 The Principle of Coherent Regularization

Proposition 4.424 (The Principle of Coherent Regularization)**.** *The emergent geometry of the PQS continuum is* endogenously *regularized. For every pair of loci* $x, x' \in \mathcal{X}_q$*, all relational observables built from the Fourfold inter–modal structure and the spectral generator are finite as a consequence of two complementary ultraviolet (UV) mechanisms:*

(i) Ultraviolet control (operator dynamics)
The Axiom of Spectral Smoothness (Axiom 4.415) bounds the growth of spectral commutators, providing polynomial control of mode-by-mode shear magnitudes.

(ii) Ultraviolet suppression (state coherence)
The Principle of Finite Entanglement Capacity (Principle 4.1) enforces a decay of modal weights $w_n(x,x')$ governed by the Holo-State $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$ (Definition 1.27), suppressing contributions of high-index modes.

Consequently, the triadic relational distances $(d_{\mathrm{corr}}, d_{\mathrm{vac}}, d_{\mathrm{spec}})$ (Sections 4.9.1 and 4.9.9 and Definition 4.463), and all operator fields derived from them (e.g. $\hat{g}_{\mu\nu}$, $\hat{t}_{\mu\nu}$, and $\hat{\Sigma}_{\mu\nu}$; cf. Definitions 5.1, 5.17 and 5.26), are finite without any external UV cutoff.

Proof. The spectral piece is representative and already exhibits the mechanism fully. By definition (Definition 4.402),

$$d_{\mathrm{spec}}(x,x') = \Big(\sum_n w_n^2(x,x')\, |\mathcal{S}_n(x,x')|^2 \Big)^{1/2},$$

with $\mathcal{S}_n$ the spectral shear observable (Definition 4.386). Spectral Smoothness gives $|\mathcal{S}_n(x,x')| \le C\,\lambda_n^k$ (Axiom 4.415), while Finite Entanglement Capacity yields $w_n(x,x') \le C'\lambda_n^{-\alpha}$ with $\alpha > k + \frac{1}{2}$ (Principle 4.1). Hence

$$w_n^2\, |\mathcal{S}_n|^2 \;\le\; (CC')^2\, \lambda_n^{-2(\alpha-k)},$$

and the series converges by the p–series test since $2(\alpha-k) > 1$. This is Proposition Proposition 4.416.

For the correlator distance d_{corr} (Section 4.9.1), the Holo-State expectation of the bilocal Fourfold operator is norm–bounded by Cauchy–Schwarz in $\mathcal{H}_{\mathrm{em}}$; the normalizations $N(\psi_1,x)$ and $N(\psi_2,x')$ were chosen precisely so that the coherence fraction $C(x,x')$ lies in $[0,1]$, hence $d_{\mathrm{corr}}(x,x') = \sqrt{1-C(x,x')} \in [0,1]$ is manifestly finite.

For the vacuum–response distance d_{vac} (Definition 4.463), the coincident-limit construction and operator-norm control of the corresponding variation yield a finite quantity by the same Holo-State norm bounds used in the correlator case.

Because $\hat{g}_{\mu\nu}$, $\hat{t}_{\mu\nu}$, and $\hat{\Sigma}_{\mu\nu}$ are built from Fréchet variations of the (squared) composite distances and times (all finite) and then evaluated in $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$ (Definitions 5.1, 5.17 and 5.26), their expectation values are finite as well. No external regulator enters: UV growth is *controlled* by the commutator bounds and *suppressed* by modal-weight decay. □

Remark 4.425 (Scope, robustness, and invariances)**.**

(a) *Admissible spectral generators.* Finiteness is uniform over $\hat{D} \in \mathfrak{D}_{\mathrm{admissible}}$ (Definition 4.343); replacing $\hat{D}$ within this class preserves both bounds and hence the convergence.

(b) *Gauge/transport insensitivity.* Parallel transport $\mathcal{U}(x \leftarrow x';\gamma)$ and admissible basis changes in the Fourfold channels (Definition 4.9) leave the norm bounds entering d_{corr} and the weight/commutator estimates for d_{spec} unchanged.

(c) *Monotonicity under coarse–graining.* Any coarse-graining that contracts the mode index set or further suppresses off-diagonal Fourfold blocks can only decrease (or leave unchanged) the RMS sums defining d_{spec} and the coherence fraction in d_{corr}.

(d) *No scheme dependence.* Since no external cutoff or subtraction is introduced, there is no renormalization-scheme arbitrariness in the definitions of d_{corr}, d_{vac}, or d_{spec}; finiteness is a property of the operator algebra plus the Holo-State.

4.9.14 Holosymmetric Wordwise Principle of Coherent Regularization

Proposition 4.426 (Wordwise Principle of Coherent Regularization)**.** *The emergent geometry of the PQS continuum is* endogenously *regularized in the holosymmetric wordwise calculus. For every pair of loci $x, x' \in \mathcal{X}_q$, all relational observables built from the Fourfold inter–modal structure and the wordwise spectral generator are finite as a consequence of two complementary ultraviolet (UV) mechanisms:*

(i) Ultraviolet control (operator dynamics)
Wordwise Spectral Smoothness holds as a theorem (Theorem 4.421), giving polynomial control of the mode-by-mode shear magnitudes $\left|\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\right|$ (Definition 4.406).

(ii) Ultraviolet suppression (state coherence)
The Wordwise Finite Entanglement Capacity (Principle 4.2) enforces decay of the modal weights $w_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')$ (Definition 4.407), suppressing high-index modes.

Consequently, the triadic relational distances

$$\left(d_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]},\ d_{\mathrm{vac}},\ d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}\right)$$

(Definitions 4.339, 4.408 and 4.463) and all operator fields derived from them (e.g. $\hat{g}_{\mu\nu}$, $\hat{t}_{\mu\nu}$, and $\hat{\Sigma}_{\mu\nu}$; cf. Definitions 5.1, 5.17 and 5.26) are finite without any external UV cutoff.

Proof. The spectral piece already exhibits the mechanism fully. By Definition 4.408,

$$d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') = \left(\sum_n \left(w_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\right)^2 \left|\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\right|^2\right)^{1/2}.$$

Wordwise Spectral Smoothness gives $\left|\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\right| \le C\,(\lambda_n^{[\mathbf{W}_1]})^k$ (Theorem 4.421), while Wordwise Finite Entanglement Capacity yields $w_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \le C'(\lambda_n^{[\mathbf{W}_1]})^{-\alpha}$ with $\alpha > k + \frac{1}{2}$ (Principle 4.2). Hence

$$\left(w_n^{[\mathbf{W}_1,\mathbf{W}_2]}\right)^2 \left|\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}\right|^2 \le (CC')^2\,(\lambda_n^{[\mathbf{W}_1]})^{-2(\alpha-k)},$$

and the series converges (since $2(\alpha-k) > 1$), proving finiteness; cf. Proposition 4.422.

For the correlator distance, by Definition 4.339 we have

$$d_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') = \sqrt{1 - C^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')} \in [0,1],$$

as the numerator/denominator are Holo-State inner products bounded by Cauchy–Schwarz in the emergent fibre (Section 4.6); see also Proposition 4.340 for equivalence to the exogenous form.

For the vacuum–response distance d_{vac} (Definition 4.463), the coincident-limit construction and the same Holo-State norm bounds ensure finiteness, exactly as in the correlator case.

Finally, $\hat{g}_{\mu\nu}$, $\hat{t}_{\mu\nu}$, and $\hat{\Sigma}_{\mu\nu}$ are defined by Fréchet variations of (squared) composite distances/times and evaluation in $|\mathring{\mathbf{\Psi}}\rangle$ (Definitions 5.1, 5.17 and 5.26); since their inputs are finite, the resulting operator fields have finite expectations. No external regulator is introduced: UV growth is *controlled* by operator dynamics (spectral smoothness) and *suppressed* by state coherence (modal-weight decay). □

Remark 4.427 (Scope, robustness, and invariances (wordwise))**.**

(a) *Admissible spectral generators.* Finiteness is uniform over the wordwise admissible class $\mathfrak{D}_{\mathrm{admissible}}$ (Section 4.9.4); replacing $\mathbb{D}^{[\mathbf{W}]}$ within this class preserves both bounds and hence convergence.

(b) *Gauge/transport insensitivity.* Endpoint gauges and word changes, and the transporter $\mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x \leftarrow x';\gamma)$, leave the correlator bounds and the spectral weight/commutator estimates unchanged (cf. Section 4.5, Lemma 4.409, and Corollary 4.414).

(c) *Monotonicity under coarse–graining.* Any mode-coarse–graining or suppression of off-diagonal Fourfold blocks can only decrease (or leave unchanged) the RMS sum in $d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}$ and the coherence fraction entering $d_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]}$.

(d) *No scheme dependence.* No external cutoff or subtraction enters the definitions of the distances; finiteness is an endogenous property of the operator algebra plus the Holo-State. By Propositions 4.340 and 4.411, the wordwise and exogenous presentations agree on values.

The Self-Regularizing Geometry

The spectral/differential leg of the emergent geometry can be defined purely in terms of relational, state–conditioned operators. Rather than a classical algebra of functions, we use the Spectral Operator $\hat{D}$ (Definition 4.341), the Fourfold Inter-Modal Correlator $\hat{G}_\Lambda$ (Definition 4.9), and the Quantum Holo-State $|\mathring{\mathbf{\Psi}}\rangle$ (Definition 1.27). The construction below packages the ingredients of Definitions 4.386, 4.392 and 4.402 and the ultraviolet controls of Axiom 4.415 and Principle 4.1 into a single, self-contained definition.

Definition 4.428 (Self-Regularizing Spectral Distance)**.** For loci $x, x' \in \mathcal{X}_q$, the *self-regularizing spectral distance* is the state–conditioned functional

$$d_{\mathrm{spec}}(x,x') \;:=\; \left(\sum_n w_n^2(x,x') \left| \mathcal{S}_n(x,x') \right|^2 \right)^{1/2},$$

where:

(i) *Spectral shear observable* $\mathcal{S}_n(x,x')$ is the Holo-State expectation of the commutator between $\hat{D}$ and the modal correlator at mode n,

$$\mathcal{S}_n(x,x') := \left\langle \mathring{\mathbf{\Psi}} \,\middle|\, \left[\hat{D},\ \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_n^*, \psi_n] \right] \,\middle|\, \mathring{\mathbf{\Psi}} \right\rangle, \quad \text{(Definition 4.386)}$$

with $\{|\psi_n^{(x)}\rangle\}$ a local orthonormal basis (e.g. $\hat{D}_x$–eigenmodes).

(ii) *Modal weight* $w_n(x,x')$ is the Holo-State norm of the corresponding bilocal probe,

$$w_n(x,x') := \left\| \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_n^*, \psi_n] \; |\mathring{\mathbf{\Psi}}\rangle \right\|, \quad \text{(Definition 4.392)}$$

quantifying Holo-State–supported coherence of mode n between x and x'.

(iii) *Endogenous finiteness (coherent regularization).* If the *Axiom of Spectral Smoothness* holds, $|\mathcal{S}_n(x,x')| \leq C\,\lambda_n^k$ (Axiom 4.415), and the *Principle of Finite Entanglement Capacity* holds, $w_n(x,x') \leq C'/\lambda_n^\alpha$ with $\alpha > k + \frac{1}{2}$ (Principle 4.1), then the series converges and $d_{\text{spec}}(x,x')$ is finite. Equivalently, the *Principle of Coherent Regularization* (Proposition 4.424) guarantees a cutoff–free, self-regularized metric contribution.

This definition yields an intrinsically quantum, background–independent distance: it measures expectation–valued commutator tension (differential structure from $\hat{D}$) along channels selected and *regularized* by the Holo-State's finite entanglement support (weights from $\hat{G}_\Lambda$). No external ultraviolet regulator is required.

Wordwise Self-Regularizing Geometry (Holosymmetric Realization)

The spectral/differential leg of the emergent geometry can be defined purely in terms of *wordwise*, relational, state–conditioned operators. In place of a classical algebra of functions, we use the wordwise spectral operator $\mathbb{D}^{[\mathbf{W}]}$ (Section 4.9.4), the Fourfold inter–modal correlator $\hat{G}_\Lambda$ evaluated on word-rotated probes (Definitions 4.406 and 4.407), and the Quantum Holo-State $|\mathring{\mathbf{\Psi}}\rangle$ (Definition 1.27). The construction below packages the ingredients of Definitions 4.406 to 4.408 together with the ultraviolet controls proved in Theorem 4.421 and assumed in Principle 4.2 into a single, self-contained wordwise definition.

Definition 4.429 (Wordwise Self-Regularizing Spectral Distance)**.** For loci $x, x' \in \mathcal{X}_q$ and admissible endpoint words $\mathbf{W}_1$ at x, $\mathbf{W}_2$ at x', the *wordwise self-regularizing spectral distance* is the state–conditioned functional

$$d_{\text{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') := \left(\sum_n \left(w_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \right)^2 \left| \mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \right|^2 \right)^{1/2}, \tag{4.205}$$

where:

(i) *Wordwise spectral shear observable* $\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')$ is the Holo-State expectation of the wordwise spectral difference projected to the nth local spectral band at x,

$$\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') := \left\langle \mathring{\mathbf{\Psi}} \right| \Pi_n^{(x)[\mathbf{W}_1]}\, \Delta\mathbb{D}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\, \Pi_n^{(x)[\mathbf{W}_1]} \left| \mathring{\mathbf{\Psi}} \right\rangle, \quad \text{(Definition 4.406)}$$

with $\Pi_n^{(x)[\mathbf{W}_1]}$ the wordwise spectral projector of $\mathbb{D}^{[\mathbf{W}_1]}(x)$ (Definition 4.405).

(ii) *Wordwise modal weight* $w_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')$ is the Holo-State norm of the corresponding bilocal probe restricted to the nth band,

$$w_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') := \left\| \Pi_n^{(x)[\mathbf{W}_1]}\, \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})\, \Pi_n^{(x)[\mathbf{W}_1]}\, |\mathring{\mathbf{\Psi}}\rangle \right\|, \quad \text{(Definition 4.407)}$$

quantifying Holo-State–supported coherence of mode n between x and x'.

(iii) *Endogenous finiteness (coherent regularization).* By the *Wordwise Spectral Smoothness Theorem* (Theorem 4.421), $\left|\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\right| \leq C\,(\lambda_n^{[\mathbf{W}_1]}(x))^k$, and by the *Wordwise Finite Entanglement Capacity* principle (Principle 4.2), $w_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \leq C'/(\lambda_n^{[\mathbf{W}_1]}(x))^\alpha$ with $\alpha > k + \frac{1}{2}$. Therefore the series in (4.205) converges and $d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')$ is finite (Proposition 4.422). Equivalently, the *Wordwise Principle of Coherent Regularization* (Proposition 4.426) guarantees a cutoff–free, self-regularized metric contribution.

Discussion (wordwise). The distance (4.205) measures expectation–valued differential tension (via $\mathbb{D}^{[\mathbf{W}]}$) along bilocal channels selected and *regularized* by the Holo-State's finite entanglement support (via $w_n^{[\mathbf{W}_1,\mathbf{W}_2]}$). It is invariant under endpoint word/gauge changes and agrees numerically with the exogenous definition by Proposition 4.411. Path dependence is Lipschitz in the holonomy size and vanishes on flat patches (Lemma 4.413 and Corollary 4.414). No external ultraviolet regulator is required.

4.9.15 A Relational Framework for Infrared Regularization

The IQR framework endogenously regulates ultraviolet behaviour via *Spectral Smoothness* and *Finite Entanglement Capacity* (Proposition 4.424). It likewise resolves **infrared (IR) divergences** non-perturbatively. In place of idealised asymptotic assumptions, IR finiteness follows from the intrinsically *relational* and *state-conditioned* formulation: physical excitations are never isolated from the Quantum Holo-State $|\mathring{\mathbf{\Psi}}\rangle$ (Definition 1.27), observables are defined on finite relational windows, and long-range contributions are suppressed by the Holo-State's coherence structure (modal weights, Definition 4.392). We articulate three complementary principles.

The Principle of Intrinsic Entanglement Dressing

Principle 4.3 (Non-asymptotic nature of physical states)**.** *In a universe described by a single, non-factorisable Holo-State $|\mathring{\mathbf{\Psi}}\rangle$, no subsystem is ever asymptotically free. Every excitation is* intrinsically dressed*: its operational properties are co-defined by entanglement with the relational background (geometry + matter), encoded in state-conditioned correlators such as $\hat{G}_\Lambda$ (Definition 4.9) and their Holo-State evaluations.*

Corollary 4.430 (S-matrix as an effective construct)**.** *The textbook S-matrix arises only in a decohered, semi-classical regime that neglects intrinsic dressing. IR divergences in its perturbative expansion reflect that idealisation. In the full relational theory, scattering amplitudes are computed from* state-dependent *correlators and effects (e.g. prismatic/holosymmetric probes), automatically accompanied by Holo-State weights that regularise long-wavelength tails.*

Operational Windowing and Finite Relational Support

Principle 4.4 (Relational windowing)**.** *Physical interrogations are defined on* finite relational support*: bilocal evaluations use admissible parallel transports along paths γ of finite length; protocol-level effects (e.g. holosymmetric interferometers) employ preparation/readout windows with bounded temporal extent. Formally, every observable $\mathcal{O}[x,x';\gamma]$ entering distances or*

amplitudes is compactly supported in relational data (path length, mode index sets) fixed by the experiment's operational envelope.

Remark 4.431 (Immediate consequence)**.** The IR pathologies of "infinite time/volume limits" do not arise: all correlator integrals and protocol compositions are taken over finite windows. Any large-scale limit is an *explicit* approximation, not a structural requirement of the theory.

Relational Cluster Property and Coherence Length

Principle 4.5 (Relational cluster decomposition)**.** *There exists a (state-dependent) coherence profile for the Holo-State such that the state-conditioned modal weights $w_n(x, x')$ (Definition 4.392) decay with increasing relational separation; equivalently, for each mode family $\mathcal{E}$,*

$$\sum_{n\in\mathcal{E}} w_n(x, x') \longrightarrow 0 \quad \textit{as the relational separation of } (x, x') \textit{ exceeds the coherence scale.}$$

This encapsulates a cluster property: sufficiently distant loci become operationally decoupled *in Holo-State–weighted observables.*

Remark 4.432 (Compatibility with UV regularisation)**.** The same weights w_n that suppress high-index (UV) modes also encode large-separation (IR) decay. Thus the *coherent regularisation* mechanism is bidirectional: UV control via Axiom 4.415 + Principle 4.1, and IR suppression via Principle 4.4 + Principle 4.5.

IR Finiteness of Relational Observables

Theorem 4.433 (Infrared regularisation of relational amplitudes and distances)**.** *Let $\mathcal{A}[x, x'; \gamma]$ be any relational amplitude constructed from finite-support compositions of $\hat{G}_\Lambda$–evaluations and prismatic/holosymmetric effects, and let $d_\bullet(x, x') \in \{d_{\mathrm{corr}}, d_{\mathrm{vac}}, d_{\mathrm{spec}}\}$ denote the relational distances of Sections 4.9.1 and 4.9.9 and Definition 4.463. Under Principles 4.3 to 4.5, all such quantities are IR-finite; no large-time/large-volume regulator is required.*

Sketch. (i) By Principle 4.4, all constructions have finite relational support (no integrals to infinity). (ii) For d_{corr} and d_{vac}, Holo-State norm bounds (Cauchy–Schwarz on the emergent Hilbert space) ensure boundedness; the finite support removes soft/collinear divergences that in QFT arise from integrating over infinite tails. (iii) For d_{spec}, the RMS sum $\sum_n w_n^2 |\mathcal{S}_n|^2$ (Definition 4.402) is already UV-finite by Proposition 4.424; IR finiteness follows because, at large separation, Principle 4.5 forces $w_n(x, x') \to 0$ modewise (or summably over degenerate families), while the commutator terms remain polynomially controlled by Axiom 4.415. Hence all contributions are summable and the limits are finite. □

Remark 4.434 (Relation to dressed scattering)**.** In regimes where an effective scattering picture is appropriate, the "intrinsic dressing" of Principle 4.3 plays the role of soft-sector resummation: long-wavelength contributions are absorbed into the Holo-State–conditioned weights that multiply the same channel algebra used to form amplitudes. The outcome is an IR-finite, protocol-dependent transition functional without imposing external regulators.

Reciprocal Resolution Limits

The same structural ingredients that render the ultraviolet (UV) sector finite also impose a *natural infrared (IR) resolution limit*. In IQR, measurements are relational, state–conditioned, and of finite operational support (Principle 4.4); the Spectral Operator's commutator growth is polynomially controlled (Axiom 4.415); and the Holo-State furnishes mode–dependent weights (Definition 4.392) that suppress unsupported channels (Principles 4.3 and 4.5). Together these facts guarantee that sufficiently *soft* modes contribute negligibly to relational observables.

Definition 4.435 (IR Coherence Scale (Operational Definition))**.** Fix loci $x, x' \in \mathcal{X}_q$ and a tolerance $\varepsilon \in (0,1)$. Define the low–frequency partial sum

$$\mathcal{S}_{\le\Lambda}(x,x') \;:=\; \sum_{\lambda_n \le \Lambda} w_n^2(x,x') \left|\mathcal{S}_n(x,x')\right|^2, \quad \text{with } \mathcal{S}_n,\, w_n \text{ from Definitions 4.386 and 4.392.}$$

The *IR coherence scale* is the minimal spectral threshold at which the low–frequency contribution falls below the tolerance:

$$\lambda_{\mathrm{IR}}(x,x';\varepsilon) \;:=\; \inf\left\{\Lambda > 0 \;\middle|\; \mathcal{S}_{\le\Lambda}(x,x') \le \varepsilon^2\right\}.$$

Proposition 4.436 (Reciprocal Resolution Limits)**.** *Assume Spectral Smoothness (Axiom 4.415) with $|\mathcal{S}_n(x,x')| \le C\,\lambda_n^k$ for some $k \in \mathbb{N}$, and suppose the eigenvalue counting function of $\hat{D}_x$ obeys a Weyl-type bound $N(\Lambda) \le A\,\Lambda^{d_s}$ (spectral dimension d_s; cf. Remark 4.417). Then:*

(i) (Existence) *For every $\varepsilon \in (0,1)$ there exists a finite $\lambda_{\mathrm{IR}}(x,x';\varepsilon)$.*

(ii) (Quantitative bound) *If $w_n(x,x') \le W$ uniformly for $\lambda_n \le \Lambda$ (e.g. after normalisation Remark 4.394), then*

$$\mathcal{S}_{\le\Lambda}(x,x') \;\le\; K\,\Lambda^{2k+d_s}, \qquad K := W^2\,C^2\,\tfrac{A}{2k+d_s},$$

and hence one may choose

$$\lambda_{\mathrm{IR}}(x,x';\varepsilon) \;\le\; \left(\tfrac{\varepsilon^2}{K}\right)^{\frac{1}{2k+d_s}}.$$

(iii) (Band–limiting) *Together with the UV suppression $w_n(x,x') \le C'\lambda_n^{-\alpha}$ for $\alpha > k + \frac{1}{2}$ (Principle 4.1), the spectral contribution to distance is effectively band–limited to $\lambda_{\mathrm{IR}}(x,x';\varepsilon) \lesssim \lambda_n \lesssim \lambda_{\mathrm{cutoff}}(x,x';\varepsilon)$, with both thresholds determined endogenously by the Holo-State and the operational window.*

Proof. (i) By Axiom 4.415, $|\mathcal{S}_n|^2 \le C^2\lambda_n^{2k}$. For $\lambda_n \le \Lambda$ and any uniform bound $w_n \le W$, we obtain

$$\mathcal{S}_{\le\Lambda} \;\le\; W^2\,C^2 \sum_{\lambda_n \le \Lambda} \lambda_n^{2k}.$$

Using the counting bound $N(\Lambda) \le A\Lambda^{d_s}$ and comparing the Riemann sum with an integral yields $\sum_{\lambda_n\le\Lambda} \lambda_n^{2k} \le \frac{A}{2k+d_s}\Lambda^{2k+d_s}$. Thus $\mathcal{S}_{\le\Lambda} \le K\,\Lambda^{2k+d_s} \to 0$ as $\Lambda \downarrow 0$, proving existence and the stated bound. (ii) follows immediately by solving $K\,\Lambda^{2k+d_s} = \varepsilon^2$ for Λ. (iii) UV band–edge follows from Principle 4.1 together with Proposition 4.424; the IR band–edge is (i)–(ii). □

Remark 4.437 (Operational Meaning and Reciprocity)**.** The scale $\lambda_{\mathrm{IR}}(x, x'; \varepsilon)$ depends on the *operational window* (path length, interrogation time) and on the Holo-State through w_n. Longer windows raise sensitivity to softer modes (smaller λ_{IR}), while shorter windows act as high–pass filters. Conversely, the Holo-State's finite entanglement capacity fixes an effective UV band–edge. The emergent geometry is thus *self–regularising at both ends*: a band–passed, state–conditioned metric contribution determined by $(\hat{D}, \hat{G}_\Lambda, |\overset{\circ}{\boldsymbol{\Psi}}\rangle)$.

State-Dependent Spectral Gapping

In IQR, particle spectra are *state–conditioned*: effective masses and dispersion relations may depend on the global Holo-State $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$. This provides a dynamical, relational mechanism for IR regulation. In particular, while the graviton is massless in the *ideal* vacuum by the Goldstone mechanism (Proposition 11.29), coherent long–range configurations in other sectors can shift the effective vacuum and open a *state–dependent* mass gap.

Definition 4.438 (State–Conditioned Graviton Mass Operator)**.** Let $\mathcal{M}(\bar{\Phi})$ denote the (local) mass operator obtained from the Hessian of the Gaiamiti potential $\mathcal{V}_\Phi$ at a background $\bar{\Phi}(x)$ (see eq. (10.125)). Write $\Pi^{(2)}$ for the transverse–traceless projector onto the spin–2 subspace used in Section 11.1.3. For a perturbed background $\bar{\Phi}'(x)$ induced by the global state (e.g. via Tantrakyma backreaction), define the *effective graviton mass operator*

$$\mathcal{M}_{\mathrm{eff}}(x; \bar{\Phi}') \;:=\; \Pi^{(2)}\, \mathcal{M}(\bar{\Phi}'(x))\, \Pi^{(2)}.$$

Given any unit TT polarisation $e^{\mu\nu}(x)$, the *state–dependent mass square* is the Rayleigh quotient

$$m^2_{\mathrm{eff}}(x; e) \;:=\; \frac{\langle e,\, \mathcal{M}_{\mathrm{eff}}(x; \bar{\Phi}')\, e\rangle}{\langle e, e\rangle}, \qquad m^2_{\mathrm{eff}}(x) \;:=\; \inf_{e \in \mathrm{TT}}\; m^2_{\mathrm{eff}}(x; e).$$

In the ideal vacuum $\bar{\Phi}' = \bar{\Phi}$ one has $\mathcal{M}(\bar{\Phi})\, \Pi^{(2)} = 0$ by Proposition 11.29, hence $m_{\mathrm{eff}}(x) = 0$.

Proposition 4.439 (State–dependent Graviton Mass Gapping)**.** *Let $\bar{\Phi}'(x) = \bar{\Phi}(x) + \delta\Phi(x)$ be the effective background in the presence of a coherent long–range Tantrakyma configuration (cf. Section 9.18.5). Suppose:*

(a) Stability near the ideal vacuum
The restricted Hessian $\Pi^{(2)}\, \nabla^2 \mathcal{V}_\Phi(\bar{\Phi} + \theta\, \delta\Phi)\, \Pi^{(2)}$ is positive semidefinite for $\theta \in [0, 1]$, and strictly positive for some $\theta > 0$ whenever $\delta\Phi \neq 0$.

(b) Nontrivial backreaction
The Tantrakyma–induced shift $\delta\Phi$ is not orthogonal to all spin–2 directions in the sense that $\Pi^{(2)}\Big(\nabla^3 \mathcal{V}_\Phi(\bar{\Phi})[\delta\Phi, \cdot\,]\Big)\Pi^{(2)} \not\equiv 0$.

Then the effective graviton mass is strictly positive:

$$m^2_{\mathrm{eff}}(x) \;>\; 0 \quad \textit{whenever} \quad \delta\Phi(x) \neq 0,$$

and vanishes continuously as the background returns to the ideal vacuum: $\lim_{\delta\Phi \to 0} m_{\mathrm{eff}}(x) = 0$.

Sketch. Expand $\mathcal{M}(\bar{\Phi}')$ to first nonvanishing order around $\bar{\Phi}$: $\mathcal{M}(\bar{\Phi}+\delta\Phi) = \mathcal{M}(\bar{\Phi}) + \nabla\mathcal{M}(\bar{\Phi})[\delta\Phi] + O(\|\delta\Phi\|^2)$. By Proposition 11.29, $\Pi^{(2)}\mathcal{M}(\bar{\Phi})\Pi^{(2)} = 0$. Assumption (b) ensures that the TT–projected linear correction is not identically zero, and (a) guarantees nonnegativity (and, away from $\delta\Phi = 0$, strict positivity) of the Rayleigh quotient on the TT subspace. Continuity in $\delta\Phi$ gives the limiting statement. □

Corollary 4.440 (IR Regulation via State–dependent Gap)**.** *On scales large compared with the local* IR correlation length $\ell_{\mathrm{IR}}(x) := m_{\mathrm{eff}}(x)^{-1}$, *TT correlators (for the TT graviton projection $\hat{h}_{\mu\nu}$ see Definition 10.25 and eq. (10.29) and the channel–resolved propagator Section 10.17.5 and eq. (10.112)) acquire an exponential (or faster) suppression, $\langle\hat{h}_{\mu\nu}(x)\hat{h}_{\rho\sigma}(x')\rangle \sim e^{-\mathrm{dist}(x,x')/\ell_{\mathrm{IR}}(x)}$ modulo transport and curvature corrections (hyperbolic TT selection cf. Definition 10.27; see also the typed scalar/Apeiron scale and associated correlation length in Definition 9.242). Consequently, long–wavelength contributions to relational amplitudes and to $d_{\mathrm{corr}}, d_{\mathrm{vac}}, d_{\mathrm{spec}}$ are IR–finite in backgrounds with $m_{\mathrm{eff}}(x) > 0$, consistent with cone–gated microcausality and IR monotonicity (Proposition 9.108 and Lemma 9.105).*

Remark 4.441 (Apeiron Scale and Tantrakyma Sourcing)**.** The effective mass $m_{\mathrm{eff}}(x)$ is controlled by the *geometric inertia* scale set by the Spin–0 Apeiron parameters in $\mathcal{V}_\Phi$ (Section 10.8); to leading order one expects

$$m^2_{\mathrm{eff}}(x) \;\propto\; m_A^2\, F[\bar{\Phi}; \delta\Phi(x)],$$

with a dimensionless functional F determined by the cubic/quartic couplings in the Hessian and by the local coherence of the Tantrakyma sector. In the typed–Hessian language of Chapters 10–11, F encapsulates how the trace–typed curvature/mass operator in the scalar block responds to the backreaction $\delta\Phi$ (cf. the realization map and operator–to–geometry pipeline in Definition 9.60 and the typed selector machinery in Definition 10.7 and Proposition 10.149). Coherent Kymon populations (quanta of the Tantrakyma field; Section 9.18.5) furnish the backreaction $\delta\Phi$ that shifts the Gaiamiti sector off the Goldstone minimum, thereby inducing the gap. As $\delta\Phi \to 0$ (loss of coherence / dilution), the mass gap closes, restoring the Goldstone limit.

Remark 4.442 (Gauge Symmetry and Compatibility)**.** The "mass" here is *effective*, arising from propagation in a dressed medium defined by $|\mathring{\boldsymbol{\Psi}}\rangle$, not a fundamental gauge–breaking term. It is computed after projection onto physical TT modes and is compatible with the constraints used in Section 11.1.3. In this sense it parallels familiar medium–induced gaps (e.g. Anderson–Higgs screening) while preserving the underlying symmetry structure of the ideal vacuum [95, 97, 96, 98, 91, 82, 92, 93].

4.9.16 Wordwise Relational Framework for Infrared Regularization

The IQR framework endogenously regulates the ultraviolet via *Wordwise Spectral Smoothness* and *Finite Entanglement Capacity* (Proposition 4.426). It likewise resolves **infrared (IR) divergences** non-perturbatively in the holosymmetric wordwise calculus. In place of idealised asymptotics, IR finiteness follows from the intrinsically *relational* and *state-conditioned* formulation: physical excitations are never isolated from the Quantum Holo-State $|\mathring{\boldsymbol{\Psi}}\rangle$

(Definition 1.27); observables are bilocal/relational and evaluated on finite wordwise windows; long-range contributions are controlled by (i) spectral smoothness at low energy and (ii) bounded state-conditioned weights. We state precise wordwise principles and convert them into theorems.

Wordwise Principle of Intrinsic Entanglement Dressing

Principle 4.6 (Non-asymptotic nature of physical states (wordwise))**.** *In a universe described by a single, non-factorisable Holo-State* $|\mathring{\Psi}\rangle$*, no subsystem is asymptotically free. Every excitation is* intrinsically dressed*: its operational properties are co-defined by entanglement with the relational background (geometry + matter), encoded in state-conditioned, wordwise correlators and their Holo-State evaluations (e.g.* $\hat{G}_\Lambda$ *on word-rotated probes; cf. Definitions 4.406 and 4.407).*

Proposition 4.443 (Wordwise dressed observables are IR-finite)**.** *Let* $\mathcal{O}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')$ *be any relational observable built from* $\hat{G}_\Lambda$ *and* $\mathbb{D}^{[\mathbf{W}]}$ *that enters the distances in Definitions 4.339 and 4.408. Then its Holo-State expectation and the induced contributions to* $\left(d_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]}, d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}\right)$ *are IR-finite.*

Proof. For $d_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]}$, finiteness is immediate from the normalization (Definition 4.339). For $d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}$, split the spectral sum at a small threshold $\varepsilon > 0$:

$$\sum_n \left(w_n^{[\mathbf{W}_1,\mathbf{W}_2]}\right)^2 \left|\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}\right|^2 = \sum_{\lambda_n \le \varepsilon} (\cdots) + \sum_{\lambda_n > \varepsilon} (\cdots).$$

The high-energy tail is finite by UV regularization (Proposition 4.422). For the low-energy part, use the Wordwise Spectral Smoothness theorem (Theorem 4.421) with $k \ge 1$ to get $\left|\mathcal{S}_n\right| \le C\,\lambda_n^k$; meanwhile the weights are uniformly bounded $w_n \le \|\hat{G}_\Lambda\|$ on each band (definition Definition 4.407 and Cauchy–Schwarz in the emergent fibre, Section 4.6). Since $\mathbb{D}^{[\mathbf{W}_1]}(x)$ is admissible, its localized spectrum is discrete with finite multiplicity on bounded intervals (cf. Section 4.9.4); hence the sum over $\{\lambda_n \le \varepsilon\}$ is finite. This yields IR finiteness. □

Remark 4.444 (Counting-function strengthening at low energy)**.** If a two-sided Weyl-type estimate holds near the origin, $N^{[\mathbf{W}_1]}(\Lambda) \lesssim \Lambda^{d_s}$ for small Λ (spectral dimension $d_s > 0$), then the low-energy sum is bounded by $\sum_{\lambda_n \le \varepsilon} \lambda_n^{2k} \lesssim \varepsilon^{2k+d_s}$, so IR finiteness already follows for $k > 0$ even without appealing to boundedness of weights.

Corollary 4.445 (S-matrix as an effective construct (wordwise))**.** *The textbook S-matrix arises only in a decohered, semi-classical regime that neglects intrinsic dressing [82, 14, 401, 230]. In the full relational theory, scattering amplitudes are computed from* state-dependent, wordwise *correlators and effects (e.g. prismatic/holosymmetric probes), automatically accompanied by Holo-State weights that regularize long-wavelength (IR) tails [241, 255, 287, 355, 402]. Thus IR divergences of perturbative S-matrix expansions reflect the asymptotic idealization, not a pathology of the wordwise relational dynamics [82, 14].*

Remark 4.446 (Justification and context)**.**

(1) Why the textbook S-matrix is only effective. Standard QFT constructs the S-matrix using asymptotic in/out *free* Fock states defined at temporal infinity, thereby idealizing away persistent long-range dressing and selecting classical-like boundary conditions. This is explicit in the foundational scattering formalism [82, Chs. 3, 4] and the axiomatic/operator approaches [230, 241]. In realistic settings, measurement-induced decoherence selects quasi-classical "pointer" sectors, which underwrites the semi-classical regime in which the S-matrix description is operationally meaningful [170, 247, 282]. In our framework this is precisely the content of the *Non-asymptotic nature of physical states* principle (Principle 4.6): excitations are intrinsically dressed by their entanglement with the relational background, so the asymptotic free-state premise is an *effective* idealization rather than a microscopic truth [14, 401].

(2) Why amplitudes are state-dependent and wordwise. Relational observables in Chapter 4 are bilocal and evaluated via wordwise transport and Holo-State expectation; operationally they are effects (POVM elements/instruments) acting in local (or quasi-local) algebras and *conditioned* on the underlying state. This is exactly how modern measurement theory and the algebraic/QFT viewpoint formulate observations [287, 355, 230, 255, 241]. Our results show that the relevant objects for distances and shears are the wordwise correlators and spectral functionals (cf. Definitions 4.339 and 4.408), with intrinsic dressing already encoded by the Holo-State weights (see also Principle 4.6).

(3) Why Holo-State weights tame IR tails. Mathematically, controlling long-wavelength behavior amounts to weighting/smearing observables and spectra; this is the familiar machinery of generalized functions and spectral/heat-kernel techniques [384, 394]. In our setup, the (bandwise) bounded, state-conditioned wordwise weights enter the spectral distance by construction (Definition 4.407), and Wordwise Spectral Smoothness (Theorem 4.421) ensures low-energy suppression of the spectral amplitudes. Together with admissibility of the local spectral operator (discreteness with finite multiplicities; Section 4.9.4), this yields the IR finiteness of dressed observables (Proposition 4.443) and distances (Proposition 4.449(i)). Coherent-state dressing supplies the physical picture behind such weights in gauge theories [402].

(4) Why perturbative IR divergences are an artifact of the idealization. The familiar IR divergences of perturbative S-matrix expansions arise when one insists on undressed, plane-wave asymptotic states for massless sectors. In mainstream treatments, the cure is to use inclusive observables and/or coherent dressing and to sum over unresolved soft quanta—i.e., to fix the *asymptotic idealization*, not the dynamics [82, Sec. 6.5]; see also the systematic discussion of the soft/collinear structure in [14, Vol. 1, Sec. 13]. Our wordwise construction builds the necessary dressing and weighting into the primitives, so IR finiteness is structural rather than remedial (Proposition 4.443, Proposition 4.449). From an effective-field-theory perspective, this matches the dictum that an "observable" is regime- and resolution-dependent; once the correct coarse-grained (inclusive) quantities are used, the apparent pathology disappears [403].

Wordwise IR Windowing and Cluster Control

Definition 4.447 (Relational window and wordwise transporter)**.** Fix a path $\gamma : x' \to x$ and let $\mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x \leftarrow x';\gamma)$ be the wordwise transporter (Section 4.5). The *relational window* of an observable $\mathcal{O}$ along γ is the wordwise-transported, bilocal evaluation $\mathcal{O}^{[\gamma]}(x,x') := \mathrm{Ev}_{|\Psi\rangle}\Big(U^\dagger \mathcal{O} U\Big)$, which depends on the window only through the holonomy class of γ.

Lemma 4.448 (Holonomy–Lipschitz control of IR windowing)**.** *For two windows γ_1, γ_2 with loop $\Delta\gamma = \gamma_2 \circ \gamma_1^{-1}$, the change in the wordwise spectral shear and spectral distance obeys the holonomy–Lipschitz bounds of Lemma 4.413 and Corollary 4.414. In particular, if the modal connection is flat on a patch, $d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')$ is independent of the window, and otherwise varies at most linearly in the holonomy size $\Phi(\Delta\gamma)$.*

Proof. **Step 1: Windowing = conjugation by transport.** By definition of the relational window Definition 4.447, for any admissible observable $\mathcal{O}$,

$$\mathcal{O}^{[\gamma_i]}(x,x') \;=\; \mathrm{Ev}_{|\Psi\rangle}\Big(U_i^\dagger\, \mathcal{O}\, U_i\Big), \qquad U_i := \mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x \leftarrow x';\gamma_i),$$

where $\mathcal{U}$ is the wordwise transporter (eqs. (4.11) and (4.12) and Definition 4.248). By functoriality and inversion of transport (Proposition 4.52 and eqs. (4.14) and (4.15)),

$$U_2 \;=\; U_1\, H, \qquad H \;=\; \mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x \leftarrow x;\Delta\gamma)$$

is the holonomy around $\Delta\gamma$ (Definition 4.56). Hence the window-change enters only through conjugation by H:

$$\mathcal{O}^{[\gamma_2]}(x,x') - \mathcal{O}^{[\gamma_1]}(x,x') = \mathrm{Ev}_{|\Psi\rangle}\Big(U_1^\dagger\Big(H^\dagger \mathcal{O}\, H - \mathcal{O}\Big)U_1\Big).$$

Step 2: Linear control by the holonomy size. Let $s \mapsto H(s)$ be the transport along a smooth parametrization of $\Delta\gamma$ with $H(0) = \mathbf{1}$ and $H(1) = H$ (eq. (4.12)). Differentiating the conjugation flow gives

$$\frac{\mathrm{d}}{\mathrm{d}s}\Big(H(s)^\dagger \mathcal{O}\, H(s)\Big) = H(s)^\dagger\,[\,-A_{\Delta\gamma}(s),\,\mathcal{O}\,]\,H(s) = \mathrm{Ad}_{H(s)}\Big(\mathrm{ad}_{A_{\Delta\gamma}(s)}\mathcal{O}\Big),$$

where $A_{\Delta\gamma}(s)$ is the pulled-back connection along the loop (Definition 4.248). Integrating in s and taking the operator norm yields

$$\Big\|H^\dagger \mathcal{O}\, H - \mathcal{O}\Big\| \;\le\; \int_0^1 \Big\|\mathrm{ad}_{A_{\Delta\gamma}(s)}\mathcal{O}\Big\|\,\mathrm{d}s.$$

For the *spectral-shear observable* $\mathcal{O} = \mathcal{S}^{[\mathbf{W}_1,\mathbf{W}_2]}$ (Definition 4.406), the commutator is controlled uniformly along the loop by the operator-Lipschitz estimates proved in Lemma 4.413. Combining this with the curvature control of the holonomy size $\Phi(\Delta\gamma)$—a gauge-invariant bound extracted from the curvature two-form via the non-Abelian Stokes estimate (Definition 4.56, Proposition 4.138, and Corollary 4.57)—gives

$$\Big|\mathcal{S}^{[\gamma_2]}(x,x') - \mathcal{S}^{[\gamma_1]}(x,x')\Big| \;\le\; C_{\mathcal{S}}\,\Phi(\Delta\gamma).$$

The *spectral distance* $d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}$ is an RMS/weighted aggregation of such shear contributions (Definition 4.408). Applying Minkowski/triangle inequalities together with the same Lipschitz control yields the bound asserted in Corollary 4.414:

$$\left|d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma_2) - d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma_1)\right| \;\le\; C_{\mathrm{spec}}\,\Phi(\Delta\gamma).$$

Step 3: Flat-patch independence. If the modal curvature vanishes on a patch, $\mathcal{F}\equiv 0$, transport is path-independent (Proposition 4.59); consequently $H=\mathbf{1}$ for any loop inside the patch, hence both the shear and the spectral distance are independent of the window. This is precisely the special case $\Phi(\Delta\gamma)=0$.

Putting the steps together proves the stated holonomy–Lipschitz bounds for the spectral shear and distance, with linear dependence on the holonomy size $\Phi(\Delta\gamma)$ and exact window-independence on flat patches. □

Proposition 4.449 (Wordwise IR regularization theorem)**.** *Let* $d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}$ *and* $d_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]}$ *be as above. Then:*

(i) (IR finiteness) *Both distances are finite for all* $x,x'\in\mathcal{X}_q$.

(ii) (IR stability) *For any sequence of windows whose holonomy size remains bounded, the corresponding values of* $d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma)$ *form a Cauchy family (hence converge).*

(iii) (Cluster reduction) *If the Holo-State exhibits a wordwise cluster property (connected bilocal correlators vanish in the limit of large relational separation), then* $d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\to 0$ *and* $C^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\to 0$ *as the relational separation tends to infinity along flat patches.*

Proof. **(i) IR finiteness.** By definition of the correlator distance (Definition 4.339), $d_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]}$ is an RMS quantity normalized by the coincident-limit kernel and therefore is finite once the underlying bilocal expectation is finite; this holds for all admissible windows/observables (Definitions 4.1 and 4.447).

For $d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}$, write (suppressing $[\mathbf{W}_1,\mathbf{W}_2]$)

$$\Big(d_{\mathrm{spec}}(x,x';\gamma)\Big)^2 \;=\; \sum_n \Big(w_n\Big)^2\,\Big|\mathcal{S}_n(x,x';\gamma)\Big|^2,$$

where w_n are the (state-conditioned) modal weights (Definition 4.407) and $\mathcal{S}_n$ the spectral-shear components (Definitions 4.406 and 4.408). Split at a small threshold $\varepsilon>0$:

$$\sum_{\lambda_n\le\varepsilon}(\cdots) \;+\; \sum_{\lambda_n>\varepsilon}(\cdots).$$

The high-energy tail is finite by endogenous UV regularization (Proposition 4.422). For the low-energy part, Wordwise Spectral Smoothness gives $|\mathcal{S}_n|\le C\,\lambda_n^k$ for some $k\ge 1$ (Theorem 4.421); the bandwise weights are uniformly bounded $w_n\le\|\hat{G}_\Lambda\|$ by Cauchy–Schwarz in the emergent fibre (Section 4.6 and Definition 4.407). Admissibility of the localized spectral operator ensures a discrete spectrum with finite multiplicities on bounded intervals (Section 4.9.4). Hence $\sum_{\lambda_n\le\varepsilon}\lambda_n^{2k}<\infty$, so the low-energy sum converges. (A counting-function strengthening appears in Remark 4.444.) Therefore d_{spec} is finite for all x,x' and any wordwise window.

(ii) IR stability (Cauchy property under bounded holonomy). Equip the set of windows with the holonomy pseudo-metric

$$d_{\mathrm{hol}}(\gamma_1, \gamma_2) \;:=\; \Phi(\Delta\gamma), \qquad \Delta\gamma := \gamma_2 \circ \gamma_1^{-1},$$

where Φ is the (gauge-invariant) holonomy size extracted from the curvature two-form (Definition 4.56, Proposition 4.138, and Corollary 4.57). By the Holonomy–Lipschitz control (Lemma 4.448 and its consequence Corollary 4.414),

$$\left| d_{\mathrm{spec}}(x, x'; \gamma_2) - d_{\mathrm{spec}}(x, x'; \gamma_1) \right| \;\le\; C_{\mathrm{spec}}\, d_{\mathrm{hol}}(\gamma_1, \gamma_2).$$

Thus the map $\gamma \mapsto d_{\mathrm{spec}}(x, x'; \gamma)$ is Lipschitz from $(\mathrm{Windows}, d_{\mathrm{hol}})$ to $\mathbb{R}$. Consequently, if (γ_j) has bounded holonomy size and is Cauchy in d_{hol} (e.g. successive loops $\gamma_{j+1} \circ \gamma_j^{-1}$ have Φ-diameter $\to 0$), then $(d_{\mathrm{spec}}(x, x'; \gamma_j))$ is Cauchy in $\mathbb{R}$ and hence converges. In particular, any bounded-holonomy net that refines windows so that $\Phi(\gamma_{j+1} \circ \gamma_j^{-1}) \to 0$ yields convergence of the spectral distance values.

(iii) Cluster reduction on flat patches. Assume the wordwise cluster property: connected bilocal correlators vanish with relational separation (Definition 4.504). On flat patches ($\mathcal{F} \equiv 0$) transport is path-independent (Proposition 4.59), so window dependence drops out. By construction, d_{spec} and $C^{[\mathbf{W}_1, \mathbf{W}_2]}$ are built from connected bilocal expectations of wordwise-dressed observables (Definitions 4.339 and 4.408); the cluster hypothesis sends these connected parts to zero as the relational separation grows. The RMS/normalization in the definitions then gives

$$d_{\mathrm{spec}}^{[\mathbf{W}_1, \mathbf{W}_2]}(x, x') \longrightarrow 0, \qquad C^{[\mathbf{W}_1, \mathbf{W}_2]}(x, x') \longrightarrow 0 \quad \text{as the separation } \to \infty,$$

uniformly over windows contained in the flat patch.

Combining (i)–(iii) yields the proposition. □

Theorem 4.450 (Wordwise IR Regularization)**.** *Let $d_{\mathrm{spec}}^{[\mathbf{W}_1, \mathbf{W}_2]}$ and $d_{\mathrm{corr}}^{[\mathbf{W}_1, \mathbf{W}_2]}$ be the wordwise spectral and correlator distances of Definitions 4.339 and 4.408. Assume the hypotheses already established in Chapter 4:*

- *Wordwise Spectral Smoothness Theorem 4.421;*
- *bounded, state-conditioned modal weights Definition 4.407 (via fibre Cauchy–Schwarz, Section 4.6);*
- *admissibility/discreteness of the localized spectrum Section 4.9.4;*
- *endogenous UV control Proposition 4.422;*
- *Holonomy–Lipschitz window control Lemma 4.448 and Corollary 4.414.*

Then for all $x, x' \in \mathcal{X}_q$ and any window γ:

(i) (IR finiteness) *$d_{\mathrm{spec}}^{[\mathbf{W}_1, \mathbf{W}_2]}(x, x'; \gamma)$ and $d_{\mathrm{corr}}^{[\mathbf{W}_1, \mathbf{W}_2]}(x, x')$ are finite.*

(ii) (IR stability) *If (γ_j) is any sequence of windows with bounded holonomy size, then $\left(d_{\mathrm{spec}}^{[\mathbf{W}_1, \mathbf{W}_2]}(x, x'; \gamma_j)\right)_j$ is Cauchy and hence convergent.*

(iii) (Cluster reduction) *If the Holo-State obeys the wordwise cluster property (connected bilocal correlators vanish with large relational separation), then along flat patches*

$$d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \longrightarrow 0, \qquad C^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \longrightarrow 0 \quad \textit{as the separation } \to \infty.$$

Moreover, on any flat patch the distances are window-independent (Proposition 4.59).

Proof. **(i) IR finiteness.** By Definition 4.339, $d_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]}$ is an RMS quantity normalized by the coincident-limit kernel, hence finite for admissible observables/windows (Definition 4.447). For $d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}$, write (suppressing $[\mathbf{W}_1,\mathbf{W}_2]$)

$$\Big(d_{\mathrm{spec}}(x,x';\gamma)\Big)^2 = \sum_n \Big(w_n\Big)^2 \Big|\mathcal{S}_n(x,x';\gamma)\Big|^2,$$

with modal weights w_n (Definition 4.407) and spectral shear components $\mathcal{S}_n$ (Definitions 4.406 and 4.408). Split the sum at $\varepsilon > 0$:

$$\sum_{\lambda_n \le \varepsilon} (\cdots) \;+\; \sum_{\lambda_n > \varepsilon} (\cdots).$$

The UV tail is finite by Proposition 4.422. For the low-energy part, Theorem 4.421 yields $|\mathcal{S}_n| \le C\,\lambda_n^k$ for some $k \ge 1$; the bandwise weights are uniformly bounded $0 \le w_n \le \|\hat{G}_\Lambda\|$ (fibre Cauchy–Schwarz, Section 4.6). Admissibility implies the localized spectrum is discrete with finite multiplicity on bounded intervals (Section 4.9.4), whence $\sum_{\lambda_n \le \varepsilon} \lambda_n^{2k} < \infty$. (Optionally strengthen via Remark 4.444.) Hence d_{spec} is finite.

(ii) IR stability. Let $d_{\mathrm{hol}}(\gamma_1,\gamma_2) := \Phi(\gamma_2 \circ \gamma_1^{-1})$ be the holonomy pseudo-metric from Definition 4.56. By Lemma 4.448 and Corollary 4.414,

$$\Big|d_{\mathrm{spec}}(x,x';\gamma_2) - d_{\mathrm{spec}}(x,x';\gamma_1)\Big| \;\le\; C_{\mathrm{spec}}\, d_{\mathrm{hol}}(\gamma_1,\gamma_2).$$

Thus $\gamma \mapsto d_{\mathrm{spec}}(x,x';\gamma)$ is Lipschitz; any sequence (γ_j) with bounded holonomy size and $d_{\mathrm{hol}}(\gamma_{j+1},\gamma_j) \to 0$ yields a Cauchy (hence convergent) sequence of distances.

(iii) Cluster reduction on flat patches. On flat patches ($\mathcal{F} \equiv 0$), transport is path-independent (Proposition 4.59), so windowing drops out. By definition, both d_{spec} and $C^{[\mathbf{W}_1,\mathbf{W}_2]}$ are built from connected bilocal expectations of wordwise-dressed observables (Definitions 4.339 and 4.408). Under the wordwise cluster property, the connected parts vanish as relational separation $\to \infty$, and the RMS/normalization implies the stated limits.
Moreover (window-independence on flat patches). If $\mathcal{F} \equiv 0$ on a patch, holonomy is trivial, hence $d_{\mathrm{spec}}(x,x';\gamma)$ is independent of γ by Proposition 4.59 and Lemma 4.448. □

Wordwise Operational Windowing and Finite Relational Support

Principle 4.7 (Relational windowing (wordwise))**.** *Physical interrogations are defined on* finite relational support. *Bilocal evaluations use admissible wordwise parallel transports* $\mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x \leftarrow x';\gamma)$ *along paths* γ *of finite length and bounded holonomy, and protocol-level effects (e.g. holosymmetric interferometers) employ preparation/readout windows with bounded temporal extent. Formally, every observable* $\mathcal{O}^{[\mathbf{W}_1,\mathbf{W}_2]}[x,x';\gamma]$ *entering distances or amplitudes is compactly supported in relational data—(path length/holonomy class, mode-index usage)—fixed by the experiment's operational envelope (cf. Section 4.5 and Definition 4.406).*

Definition 4.451 (Effective wordwise support set)**.** Given $\varepsilon > 0$, define the ε–support of the Holo-State between (x, x') by

$$\mathcal{E}_{\varepsilon}^{[\mathbf{W}_1,\mathbf{W}_2]}(x, x') := \Big\{ n : \; w_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x, x') \geq \varepsilon \Big\},$$

where $w_n^{[\mathbf{W}_1,\mathbf{W}_2]}$ are the wordwise modal weights (Definition 4.407). We say an observable has *finite relational support* if its evaluation uses only modes in a set $\mathcal{E}$ with $\mathcal{E} \subseteq \mathcal{E}_{\varepsilon}^{[\mathbf{W}_1,\mathbf{W}_2]}(x, x')$ for some $\varepsilon > 0$ and a window γ with bounded holonomy.

Lemma 4.452 (Finiteness of effective support)**.** *Under the Wordwise Finite Entanglement Capacity (Principle 4.2),* $\mathcal{E}_{\varepsilon}^{[\mathbf{W}_1,\mathbf{W}_2]}(x, x')$ *is finite for every* $\varepsilon > 0$*.*

Proof. Recall the effective support set (Definition 4.451)

$$\mathcal{E}_{\varepsilon}^{[\mathbf{W}_1,\mathbf{W}_2]}(x, x') = \Big\{ n : \; w_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x, x') \geq \varepsilon \Big\},$$

where $w_n^{[\mathbf{W}_1,\mathbf{W}_2]}$ are the wordwise modal weights (Definition 4.407).

Route A (via decay in the spectral parameter). By Wordwise Finite Entanglement Capacity we have the polynomial decay

$$w_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x, x') \;\leq\; C' \left(\lambda_n^{[\mathbf{W}_1]}(x)\right)^{-\alpha} \qquad \text{for some } C', \alpha > 0, \tag{4.206}$$

as stated in (4.200). Hence $w_n \geq \varepsilon \;\Rightarrow\; \lambda_n \leq \Lambda_\varepsilon := (C'/\varepsilon)^{1/\alpha}$. By admissibility of the localized spectral operator (Section 4.9.4), the spectrum is discrete with finite multiplicities and has no finite accumulation points; in particular the counting function $N(\Lambda) := \#\{n : \lambda_n \leq \Lambda\}$ is finite for each $\Lambda < \infty$. Therefore

$$\#\,\mathcal{E}_{\varepsilon}^{[\mathbf{W}_1,\mathbf{W}_2]}(x, x') \;\leq\; N(\Lambda_\varepsilon) \;<\; \infty,$$

so the ε–support is finite.

Route B (uniform ℓ^p–summability). Alternatively, Principle 4.2 provides (for some $p > 0$) $\sum_n \left(w_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x, x')\right)^p < \infty$. Chebyshev/Markov yields

$$\#\,\mathcal{E}_{\varepsilon}^{[\mathbf{W}_1,\mathbf{W}_2]}(x, x') = \#\{n : \; w_n \geq \varepsilon\} \;\leq\; \varepsilon^{-p} \sum_n w_n^p \;<\; \infty.$$

This gives the same conclusion without appealing to (4.206).

Either route proves that for every $\varepsilon > 0$ the set $\mathcal{E}_{\varepsilon}^{[\mathbf{W}_1,\mathbf{W}_2]}(x, x')$ is finite. □

Proposition 4.453 (IR finiteness from windowing)**.** *Let* $\mathcal{O}^{[\mathbf{W}_1,\mathbf{W}_2]}[x, x'; \gamma]$ *be any observable entering* $d_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]}$ *or* $d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}$ *(Definitions 4.339 and 4.408), evaluated on a finite relational window in the sense of Definition 4.451. Then the induced contribution to the distance is IR-finite. Moreover, its dependence on* γ *obeys the holonomy–Lipschitz bounds of Lemma 4.413 and Corollary 4.414.*

Proof. **Setup (finite relational window and effective support).** Fix $\varepsilon > 0$ and let the window γ be admissible with bounded holonomy in the sense of Principle 4.7 and Definition 4.447. By Definition 4.451 and Lemma 4.452, the ε–support $\mathcal{E}_\varepsilon^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') = \{n : w_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \geq \varepsilon\}$ is finite. Write the spectral index set as a disjoint union

$$\mathbb{N} = \underbrace{\mathcal{E}_\varepsilon}_{\text{finite}} \sqcup \underbrace{\mathcal{E}_\varepsilon^{\mathrm{c}}}_{\text{tail}},$$

and denote by Λ_ε any energy threshold with $\{\lambda_n \leq \Lambda_\varepsilon\} \supseteq \mathcal{E}_\varepsilon$ (as guaranteed by admissibility and discreteness, Section 4.9.4).

(A) Spectral distance: IR finiteness. Suppressing $[\mathbf{W}_1, \mathbf{W}_2]$ in the notation, the spectral-distance contribution can be written as

$$\Big(d_{\mathrm{spec}}(x,x';\gamma)\Big)^2 = \sum_n \Big(w_n\Big)^2 \Big|\mathcal{S}_n(x,x';\gamma)\Big|^2,$$

with modal weights w_n and spectral-shear components $\mathcal{S}_n$ (Definitions 4.406 to 4.408). Split the series:

$$\sum_{n\in\mathcal{E}_\varepsilon} \Big(w_n\Big)^2 |\mathcal{S}_n|^2 \;+\; \sum_{n\in\mathcal{E}_\varepsilon^{\mathrm{c}}} \Big(w_n\Big)^2 |\mathcal{S}_n|^2 \;=:\; \Sigma_{\mathrm{fin}} + \Sigma_{\mathrm{tail}}.$$

Finite block. Since $\mathcal{E}_\varepsilon$ is finite and the transporter is unitary along γ (eqs. (4.11) and (4.12) and Definition 4.248), the conjugated observable remains bounded in operator norm; hence each $|\mathcal{S}_n|$ is finite. Therefore $\Sigma_{\mathrm{fin}} < \infty$.

Tail. On $\mathcal{E}_\varepsilon^{\mathrm{c}}$ we have $0 \leq w_n < \varepsilon$. Moreover, by endogenous UV control (Proposition 4.422) and wordwise spectral smoothness (Theorem 4.421), there exists a finite constant $K(\Lambda_\varepsilon)$ such that

$$\sum_{\{n:\,\lambda_n > \Lambda_\varepsilon\}} |\mathcal{S}_n(x,x';\gamma)|^2 \;\leq\; K(\Lambda_\varepsilon).$$

Hence

$$\Sigma_{\mathrm{tail}} = \sum_{n\in\mathcal{E}_\varepsilon^{\mathrm{c}}} \Big(w_n\Big)^2 |\mathcal{S}_n|^2 \;\leq\; \varepsilon^2 \sum_{n\in\mathcal{E}_\varepsilon^{\mathrm{c}}} |\mathcal{S}_n|^2 \;\leq\; \varepsilon^2 K(\Lambda_\varepsilon) \;<\; \infty.$$

Combining the two parts yields $d_{\mathrm{spec}}(x,x';\gamma) < \infty$. Note that the *IR* contribution is tamed already by the finiteness of $\mathcal{E}_\varepsilon$ (only finitely many low-energy modes contribute), while the UV tail is summable by Proposition 4.422.

(B) Correlator distance: IR finiteness. By Definition 4.339 the correlator distance is an RMS quantity normalized by the coincident-limit kernel; its building blocks are bilocal wordwise evaluations of bounded observables transported along γ (Definition 4.447, Proposition 4.14, and Lemma 4.66). Finite relational support implies the evaluation of any entering observable uses only the finite set $\mathcal{E}_\varepsilon$ above threshold; the remainder can be bounded by Cauchy–Schwarz in the emergent fibre together with the same UV summability as above (Section 4.6 and Proposition 4.422). Hence $d_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') < \infty$.

(C) Holonomy–Lipschitz dependence on the window. Let γ_1, γ_2 be two windows with loop $\Delta\gamma = \gamma_2 \circ \gamma_1^{-1}$ and holonomy size $\Phi(\Delta\gamma)$ (Definition 4.56). By the Holonomy–Lipschitz bounds proved for shear and distance (Lemma 4.413 and Corollary 4.414),

$$\Big|\mathcal{S}^{[\gamma_2]} - \mathcal{S}^{[\gamma_1]}\Big| \;\leq\; C_{\mathcal{S}}\,\Phi(\Delta\gamma), \qquad \Big|d_{\mathrm{spec}}(x,x';\gamma_2) - d_{\mathrm{spec}}(x,x';\gamma_1)\Big| \;\leq\; C_{\mathrm{spec}}\,\Phi(\Delta\gamma).$$

The same line of argument (finite block + tail bound) applies to d_{corr}, yielding analogous Lipschitz control.

Thus, for any finite relational window in the sense of Definition 4.451, the induced contributions to d_{spec} and d_{corr} are IR-finite, and the window dependence is Lipschitz in holonomy, as claimed. □

Remark 4.454 (Immediate consequence (wordwise))**.** IR pathologies of "infinite time/volume limits" do not arise: all correlator integrals and protocol compositions are taken over finite windows with bounded holonomy and effective mode support. Any large-scale limit is an *explicit* approximation, not a structural requirement (cf. Principle 4.7).

Wordwise Relational Cluster Property and Coherence Length

Definition 4.455 (Coherence profile and wordwise coherence length)**.** A *coherence profile* is a non-increasing function $\mathfrak{c} : [0,\infty) \to [0,\infty)$ with $\mathfrak{c}(r) \to 0$ as $r \to \infty$ such that, for some relational separation functional $r = r(x, x'; \gamma)$ (e.g. path length or a composite distance), one has for every mode family $\mathcal{E}$,

$$\sum_{n\in\mathcal{E}} w_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \;\le\; \mathfrak{c}\Big(r(x,x';\gamma)\Big).$$

The *coherence length* $\ell_c(\epsilon)$ is the least r with $\mathfrak{c}(r) \le \epsilon$.

Principle 4.8 (Relational cluster decomposition (wordwise))**.** *The Holo-State admits a coherence profile in the sense of Definition 4.455; equivalently, sufficiently distant loci become* operationally decoupled *in Holo-State–weighted wordwise observables.*

Theorem 4.456 (IR suppression of distances under clustering)**.** *Assume Principle 4.8 and the UV regularity of Theorem 4.421 and Principle 4.2. Then along any family of windows with bounded holonomy, as $r(x,x';\gamma) \to \infty$,*

$$d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \;\longrightarrow\; 0, \qquad C^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \;\longrightarrow\; 0 \quad \Big(\textit{so } d_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]} \to 1\Big).$$

Proof idea. For any finite family $\mathcal{E}$, by Cauchy–Schwarz

$$\sum_{n\in\mathcal{E}} \big(w_n^{[\cdot]}\big)^2 \big|\mathcal{S}_n^{[\cdot]}\big|^2 \;\le\; \Big(\sup_{n\in\mathcal{E}} \big|\mathcal{S}_n^{[\cdot]}\big|^2\Big)\Big(\sum_{n\in\mathcal{E}} \big(w_n^{[\cdot]}\big)^2\Big).$$

Wordwise spectral smoothness bounds $\sup_{n\in\mathcal{E}} |\mathcal{S}_n|$ on any fixed low-energy block; the UV tail is summable by Proposition 4.422. Clustering gives $\sum_{n\in\mathcal{E}} w_n^{[\cdot]} \to 0$ as $r \to \infty$ (hence the squared sum also vanishes), implying $d_{\mathrm{spec}}^{[\cdot]} \to 0$. For the correlator, clustering forces wordwise normalized overlaps to vanish, whence $C^{[\cdot]} \to 0$ and $d_{\mathrm{corr}}^{[\cdot]} \to 1$.

Proof. We suppress $[\mathbf{W}_1, \mathbf{W}_2]$ in the notation and keep the window γ implicit when it causes no ambiguity. Throughout, we assume the holonomy size $\Phi(\gamma)$ remains uniformly bounded (bounded-holonomy family).

Step 1: ε–Λ split of the spectral series. By Definition 4.408,

$$\Big(d_{\mathrm{spec}}(x,x';\gamma)\Big)^2 = \sum_n \Big(w_n(x,x')\Big)^2 \,\Big|\mathcal{S}_n(x,x';\gamma)\Big|^2,$$

where w_n are the (state-conditioned) wordwise modal weights and $\mathcal{S}_n$ are the spectral-shear components (Definitions 4.406 and 4.407). Fix $\epsilon > 0$. By endogenous UV regularity (Theorem 4.421 and Principle 4.2; see also Proposition 4.422), we can choose $\Lambda = \Lambda(\epsilon)$ so that the high-energy tail obeys

$$\sum_{\{\lambda_n > \Lambda\}} \Big(w_n\Big)^2 \Big|\mathcal{S}_n(x,x';\gamma)\Big|^2 \;<\; \epsilon/2 \quad \text{uniformly over bounded-holonomy windows.} \tag{4.207}$$

Write the series as low-energy block $B_\Lambda := \{n : \lambda_n \le \Lambda\}$ plus tail T_Λ.

Step 2: Uniform boundedness of low-energy shear on bounded-holonomy families. By Wordwise Spectral Smoothness there exist $k \ge 1$ and $C_\Lambda < \infty$ (depending on Λ but not on r or γ within a bounded-holonomy family) such that

$$\sup_{n \in B_\Lambda} \Big|\mathcal{S}_n(x,x';\gamma)\Big| \;\le\; C_\Lambda. \tag{4.208}$$

This follows from the operator functional calculus bounds in Theorem 4.421, together with the unitary nature of transport along γ (Definition 4.248 and eqs. (4.11) and (4.12)) and the Holonomy–Lipschitz control which prevents blow-up under bounded holonomy (Lemma 4.413).

Step 3: Coherence-profile suppression of weighted low-energy block. By the relational cluster principle (Principle 4.8), there exists a coherence profile $\mathfrak{c} : [0,\infty) \to [0,\infty)$ with $\mathfrak{c}(r) \to 0$ such that for any mode family $\mathcal{E}$,

$$\sum_{n \in \mathcal{E}} w_n(x,x') \;\le\; \mathfrak{c}\Big(r(x,x';\gamma)\Big).$$

Since $0 \le w_n \le \|\hat{G}_\Lambda\|$ by fibre Cauchy–Schwarz and the definition of weights (Section 4.6 and Definition 4.407), we have for the low-energy block

$$\sum_{n \in B_\Lambda} \Big(w_n\Big)^2 \;\le\; \|\hat{G}_\Lambda\| \sum_{n \in B_\Lambda} w_n \;\le\; \|\hat{G}_\Lambda\|\; \mathfrak{c}\Big(r(x,x';\gamma)\Big). \tag{4.209}$$

Step 4: Low-energy contribution vanishes as $r \to \infty$. Combining (4.208) and (4.209) gives

$$\sum_{n \in B_\Lambda} \Big(w_n\Big)^2 \Big|\mathcal{S}_n\Big|^2 \;\le\; \Big(\sup_{n \in B_\Lambda} |\mathcal{S}_n|^2\Big) \sum_{n \in B_\Lambda} \Big(w_n\Big)^2 \;\le\; C_\Lambda^2\, \|\hat{G}_\Lambda\|\; \mathfrak{c}\Big(r(x,x';\gamma)\Big).$$

Since $\mathfrak{c}(r) \to 0$ as $r \to \infty$, there exists $R = R(\epsilon,\Lambda)$ such that for all $r \ge R$ the low-energy block is $< \epsilon/2$.

Step 5: Conclusion for d_{spec}. For any $\epsilon > 0$, we have chosen Λ so that the tail is $< \epsilon/2$ by (4.207), and then R so that the block is $< \epsilon/2$. Hence for all $r \ge R$,

$$\Big(d_{\mathrm{spec}}(x,x';\gamma)\Big)^2 \;<\; \epsilon,$$

which establishes $d_{\mathrm{spec}}(x,x';\gamma)\to 0$ as $r\to\infty$ along bounded-holonomy windows.

Step 6: Connected correlator and correlator distance. By Definition 4.339, the correlator distance d_{corr} is an RMS normalization of bilocal overlaps; write its connected contribution as $C(x,x';\gamma)$ (same object used earlier in Corollary 4.457). The relational cluster principle (Principle 4.8), together with bounded holonomy and unitary transport, implies that connected bilocal wordwise expectations vanish as $r\to\infty$:

$$C(x,x';\gamma)\ \longrightarrow\ 0.$$

Indeed, by the same ε–Λ split and the argument above, low-energy weighted overlaps are suppressed by $\mathfrak{c}(r)$ while the high-energy tail is summable by UV regularity (Theorem 4.421, Principle 4.2, and Proposition 4.422). Therefore $C\to 0$, and the RMS normalization yields $d_{\mathrm{corr}}(x,x')\to 1$.

Window dependence. All bounds used are uniform over bounded-holonomy families. Moreover, the Holonomy–Lipschitz estimates (Lemma 4.413 and Corollary 4.414) guarantee that changing the window within such a family perturbs d_{spec} and the correlator by at most $O(\Phi(\Delta\gamma))$, which remains bounded and thus does not obstruct the limit $r\to\infty$. On flat patches, the limit is window-independent by Proposition 4.59.

Combining the steps completes the proof. □

Corollary 4.457 (Cluster reduction along flat patches)**.** *If the modal connection is flat on a patch, the limits in Theorem 4.456 are window independent (cf. Lemma 4.448). In particular, along any sequence with $r\to\infty$ inside the patch, $d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\to 0$ and $d_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\to 1$.*

Proof. Work under the hypotheses of Theorem 4.456. Let $\Omega\subset\mathcal{X}_q$ be a patch on which the modal curvature vanishes, $\mathcal{F}\equiv 0$. Then transport is path-independent on Ω (Proposition 4.59), so for any two windows $\gamma_1,\gamma_2\subset\Omega$ with the same endpoints $x'\to x$ we have

$$\mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x\leftarrow x';\gamma_1)=\mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x\leftarrow x';\gamma_2),$$

and the holonomy around the loop $\Delta\gamma:=\gamma_2\circ\gamma_1^{-1}$ is trivial, $H(\Delta\gamma)=\mathbf{1}$, with holonomy size $\Phi(\Delta\gamma)=0$ (Definition 4.56 and Corollary 4.57).

By the Holonomy–Lipschitz bounds for the shear and the spectral distance (Lemma 4.413 and Corollary 4.414),

$$\begin{aligned}\left|\mathcal{S}^{[\gamma_2]}(x,x')-\mathcal{S}^{[\gamma_1]}(x,x')\right|&\le C_{\mathcal{S}}\,\Phi(\Delta\gamma),\\ \left|d_{\mathrm{spec}}(x,x';\gamma_2)-d_{\mathrm{spec}}(x,x';\gamma_1)\right|&\le C_{\mathrm{spec}}\,\Phi(\Delta\gamma).\end{aligned}$$

In particular, if $\Delta\gamma=0$ (so $\Phi(\Delta\gamma)=0$), both differences vanish. Hence $d_{\mathrm{spec}}(x,x';\gamma)$ is independent of the window $\gamma\subset\Omega$; the same applies to the connected correlator $C^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma)$ and thus to d_{corr} by its RMS normalization (Definition 4.339).

Now take any sequence of windows $(\gamma_j)\subset\Omega$ with relational separation $r(x,x';\gamma_j)\to\infty$. Because $\Phi(\gamma_j)$ is identically zero on a flat patch, the "bounded-holonomy" requirement of Theorem 4.456 is trivially satisfied. Applying Theorem 4.456 yields

$$d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma_j)\longrightarrow 0,\qquad C^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma_j)\longrightarrow 0,$$

and, by definition, $d_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\to 1$. Since all these quantities are already independent of the chosen window inside Ω, the limits are window independent, as claimed. □

Remark 4.458 (Compatibility of UV and IR regularization (wordwise))**.** The same wordwise weights $w_n^{[\mathbf{W}_1,\mathbf{W}_2]}$ that suppress high-index (UV) modes encode large-separation (IR) decay. Thus *coherent regularization* is bidirectional: UV control via Theorem 4.421 and Principle 4.2 and IR suppression via Principles 4.7 and 4.8.

Wordwise Reciprocal Resolution Limits

The same structural ingredients that render the ultraviolet (UV) sector finite also impose a *natural infrared (IR) resolution limit* in the holosymmetric wordwise calculus. Measurements are relational, state–conditioned, and of finite operational support (Principle 4.7); the wordwise Spectral Operator's commutator growth is polynomially controlled (Theorem 4.421); and the Holo-State furnishes mode–dependent weights (Definition 4.407) that suppress unsupported channels (Principles 4.6 and 4.8). Together these facts guarantee that sufficiently *soft* modes contribute negligibly to relational observables.

Definition 4.459 (Wordwise IR Coherence Scale (Operational))**.** Fix loci $x, x' \in \mathcal{X}_q$, an admissible window $\gamma : x' \to x$, and a tolerance $\varepsilon \in (0,1)$. Define the low–frequency partial sum

$$\mathcal{S}_{\le\Lambda}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma) := \sum_{\lambda_n^{[\mathbf{W}_1]}(x)\le\Lambda} \left(w_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\right)^2 \left|\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma)\right|^2,$$

with $\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}$ and $w_n^{[\mathbf{W}_1,\mathbf{W}_2]}$ from Definitions 4.406 and 4.407. The *wordwise IR coherence scale* is the (equivalent) monotone definition

$$\begin{aligned}\lambda_{\mathrm{IR}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\varepsilon;\gamma) &:= \inf\left\{\Lambda>0 \;\middle|\; \mathcal{S}_{\le\Lambda}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma) \ge \varepsilon^2\right\}\\ &= \sup\left\{\Lambda>0 \;\middle|\; \mathcal{S}_{\le\Lambda}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma) \le \varepsilon^2\right\}.\end{aligned}$$

Proposition 4.460 (Wordwise Reciprocal Resolution Limits)**.** *Assume Wordwise Spectral Smoothness (Theorem 4.421) with* $\left|\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma)\right| \le C\left(\lambda_n^{[\mathbf{W}_1]}(x)\right)^k$ *for some* $k\in\mathbb{N}$*, and suppose the eigenvalue counting function at* x *obeys a Weyl-type bound* $N^{[\mathbf{W}_1]}(\Lambda) \le A\,\Lambda^{d_s}$ *(spectral dimension* $d_s>0$*; cf. Remark 4.444). Then:*

(i) (Existence) *For every* $\varepsilon\in(0,1)$ *and window* γ*, there exists a finite* $\lambda_{\mathrm{IR}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\varepsilon;\gamma)$*.*

(ii) (Quantitative bound) *If there is a uniform bound* $w_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \le W$ *for* $\lambda_n^{[\mathbf{W}_1]}(x)\le\Lambda$ *(e.g. by normalization and Cauchy–Schwarz in the emergent fibre, cf. Definition 4.407 and Section 4.6), then*

$$\mathcal{S}_{\le\Lambda}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma) \;\le\; K\,\Lambda^{\,2k+d_s}, \qquad K := W^2\,C^2\,\tfrac{A}{2k+d_s},$$

and therefore

$$\lambda_{\mathrm{IR}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\varepsilon;\gamma) \;\le\; \left(\tfrac{\varepsilon^2}{K}\right)^{\frac{1}{2k+d_s}}.$$

(iii) (Band–limiting) *If, in addition, the weights obey the UV decay* $w_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \le C'\big(\lambda_n^{[\mathbf{W}_1]}(x)\big)^{-\alpha}$ *with* exponent $\alpha > k + \frac{d_s}{2}$ *(Principle 4.2), then the spectral contribution to the distance is effectively band–limited to*

$$\lambda_{\mathrm{IR}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\varepsilon;\gamma) \;\lesssim\; \lambda_n^{[\mathbf{W}_1]}(x) \;\lesssim\; \lambda_{\mathrm{cutoff}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\varepsilon),$$

with both thresholds determined endogenously by the Holo-State and the operational window.

Proof. (i) From Theorem 4.421, $|\mathcal{S}_n|^2 \le C^2(\lambda_n)^{2k}$; for $\lambda_n \le \Lambda$ and any uniform bound $w_n \le W$ (valid uniformly over bounded-holonomy windows by Definition 4.447 and Lemma 4.413), we obtain

$$\mathcal{S}_{\le\Lambda}^{[\cdot]} \;\le\; W^2\,C^2 \sum_{\lambda_n\le\Lambda} (\lambda_n)^{2k}.$$

Using $N^{[\mathbf{W}_1]}(\Lambda) \le A\Lambda^{d_s}$ and comparing the sum with an integral gives $\sum_{\lambda_n\le\Lambda}(\lambda_n)^{2k} \le \frac{A}{2k+d_s}\Lambda^{2k+d_s}$; this ensures the set $\{\Lambda : \mathcal{S}_{\le\Lambda}^{[\cdot]} \ge \varepsilon^2\}$ is nonempty and bounded, so the infimum is finite.

(ii) Solving $K\,\Lambda^{2k+d_s} = \varepsilon^2$ for Λ yields the stated upper bound on $\lambda_{\mathrm{IR}}^{[cdot]}$.

(iii) For the UV side, with $w_n \lesssim \lambda_n^{-\alpha}$ and $|\mathcal{S}_n| \lesssim \lambda_n^k$, the tail of the distance scales like

$$\sum_{\lambda_n>\Lambda} \lambda_n^{2k}\,\lambda_n^{-2\alpha}\,\mathrm{d}N \;\sim\; \int_{\Lambda}^{\infty} \lambda^{2k-2\alpha}\,\lambda^{d_s-1}\,\mathrm{d}\lambda,$$

which converges iff $2k - 2\alpha + d_s < 0$, i.e. $\alpha > k + \frac{d_s}{2}$. Thus the total contribution is concentrated between the IR scale from (i)–(ii) and an endogenous UV cutoff $\lambda_{\mathrm{cutoff}}$ fixed by the decay and counting constants (Principle 4.2 and Proposition 4.422). □

Remark 4.461 (Operational Meaning and Reciprocity (wordwise))**.** The scale

$$\lambda_{\mathrm{IR}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\varepsilon;\gamma)$$

depends on the *operational window* (path length and holonomy; cf. Principle 4.7 and Lemma 4.413) and on the Holo-State via the weights $w_n^{[\mathbf{W}_1,\mathbf{W}_2]}$. Longer windows (or smaller holonomy) increase sensitivity to soft modes (smaller λ_{IR}), while shorter windows act as high–pass filters. Conversely, the Holo-State's finite entanglement capacity fixes an effective UV band–edge. The emergent geometry is thus *self–regularising at both ends*: a band–passed, state–conditioned metric contribution determined by $(\mathbb{D}^{[\mathbf{W}]}, \hat{G}_\Lambda, |\mathring{\mathbf{\Psi}}\rangle)$.

Definition 4.462 (Operational IR scale)**.** Fix $\eta > 0$, loci $x, x' \in \mathcal{X}_q$, and an admissible window $\gamma : x' \to x$. Define the low–frequency partial sum

$$\mathcal{S}_{\le\Lambda}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma) := \sum_{\lambda_n^{[\mathbf{W}_1]}(x)\le\Lambda} \Big(w_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\Big)^2 \Big|\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma)\Big|^2,$$

with w_n and $\mathcal{S}_n$ from Definitions 4.406 to 4.408. The *operational IR scale* is the right–continuous inverse

$$\lambda_{\mathrm{IR}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\eta;\gamma) := \inf\Big\{\Lambda \ge 0 :\; \mathcal{S}_{\le\Lambda}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma) \;\ge\; \eta^2\Big\},$$

with the convention $\inf \varnothing = \infty$ *(equivalently, restrict* $\eta \in \big(0, d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma]\big)$*).*

4.9.17 The Vacuum-Response Distance: Geometry from Entanglement Self-Coherence

We formalize the third axis of emergent distinguishability, the *vacuum-response distance* $d_{\mathrm{vac}}(x, x')$, which quantifies how the intrinsic ("vacuum") structure of the continuum differs between two PQS loci. Operationally, it measures the sensitivity of the Quantum Holo-State $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$ to *diagonal* (coincident) probes built from the Fourfold Inter-Modal Correlator $\hat{G}_\Lambda$, with inner products and modal bases induced exactly as in Section 4.6 and Definition 4.9. In the triadic geometry $\vec{d} = (d_{\mathrm{corr}}, d_{\mathrm{vac}}, d_{\mathrm{spec}})$, d_{vac} is the "vacuum axis," complementary to correlational and spectral axes, and is defined without appeal to any background manifold or field content.

Formal definition

Definition 4.463 (Vacuum-response distance)**.** Fix an orthonormal modal basis $\{|\psi_k\rangle_x\}_k$ in the fiber $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ induced by the coincident-limit inner product of $\hat{G}_\Lambda$ (Section 4.6). For each mode, define the *local self-correlator*

$$C_k(x) \;:=\; \Big\langle \overset{\circ}{\boldsymbol{\Psi}} \Big|\, \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_x)\big[\psi_k^*, \psi_k\big] \,\Big| \overset{\circ}{\boldsymbol{\Psi}} \Big\rangle,$$

i.e. the Holo-State expectation of the diagonal restriction of the bilocal operator at x. The *vacuum-response distance* between x and x' is the root-mean-square variation of these local self-coherences:

$$\boxed{d_{\mathrm{vac}}(x, x') \;:=\; \left[\sum_k \Big|\, C_k(x) - C_k(x') \,\Big|^2\right]^{1/2}.} \tag{4.210}$$

This captures the purely *diagonal* response encoded by $\hat{G}_\Lambda$: unlike d_{corr}, which probes off-diagonal bilocal coherence, d_{vac} probes how the vacuum's self-coherence profile changes from x to x'. Gauge-covariant parallel transport ensures observer-independent comparison when constructing such correlators across the continuum.

Vector formulation

Let the *vacuum correlator profile* at x be $\vec{V}_{\mathrm{vac}}(x) := (C_1(x), C_2(x), \dots)$. Then

$$\boxed{d_{\mathrm{vac}}(x, x') \;=\; \big\|\vec{V}_{\mathrm{vac}}(x) - \vec{V}_{\mathrm{vac}}(x')\big\|_2,} \tag{4.211}$$

an ℓ^2–metric on profiles extracted from the diagonal sector of $\hat{G}_\Lambda$. This field-agnostic construction interprets the "vacuum" as a profile of entanglement self-coherence determined by $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$.

Basic properties. (i) *State-conditioning:* d_{vac} is computed in the same Holo-State that forges the modal inner products; it is thus intrinsic to the IQR arena, not imposed. (ii) *Functional orthogonality:* By construction it probes the diagonal/coincident sector, functionally orthogonal to the bilocal off-diagonal structure (for d_{corr}) and to the commutator structure (for d_{spec}). (iii) *Finiteness:* Convergence of (4.210) follows from the decay of high-index modal significance (Principle of Finite Entanglement Capacity), hence no external cutoff is required.

Remark 4.464 (Vacuum distance as a curvature indicator)**.** Variations of the profile $\vec{V}_{\text{vac}}$ can be read as "vacuum curvature": d_{vac} isolates second-order structure in the diagonal sector, complementing d_{corr} (first-order overlap) and d_{spec} (differential/commutator response). This alignment underwrites the triadic fiber $\mathcal{F}_x = \text{span}\{\mathbf{e}_{\text{corr}}, \mathbf{e}_{\text{vac}}, \mathbf{e}_{\text{spec}}\}$ used to assemble the composite metric.

Remark 4.465 (Twofold reduction)**.** In a duality-collapse regime (Definition 4.87), the vacuum-response constructed on the Twofold sector reduces to the corresponding qubit-level notion, via the same isometric identifications that recover ordinary two-level kinematics. The definition itself is unchanged—only the representation is compressed.

Summary. d_{vac} is the vacuum axis of the triadic distinguishability: a state-conditioned, gauge-covariant, UV-finite, background-free measure of how the entanglement-defined "vacuum" varies between loci—built solely from the coincident sector of $\hat{G}_\Lambda$ and the Holo-State that induces the modal inner products.

4.9.18 Wordwise Vacuum-Response Distance

We formalize the third axis of emergent distinguishability in the *holosymmetric wordwise* calculus. The *vacuum–response distance* quantifies how the intrinsic ("vacuum") self-coherence of the PQS continuum differs between loci, using only diagonal (coincident) evaluations of the Fourfold inter–modal correlator in the Holo-State. Modal bases and inner products are those induced by the wordwise machinery (Section 4.6 and Definition 4.405).

Wordwise formal definitions

Definition 4.466 (Wordwise vacuum self-correlator)**.** Fix an admissible endpoint word $\mathbf{W}_1$ at x and the associated local spectral/modal frame $\{\Pi_n^{(x)[\mathbf{W}_1]}\}_n$ of $\mathbb{D}^{[\mathbf{W}_1]}(x)$ (Definition 4.405). The *wordwise vacuum self-correlator* in band n is

$$C_n^{[\mathbf{W}_1]}(x) := \left\langle \mathring{\mathbf{\Psi}} \middle| \Pi_n^{(x)[\mathbf{W}_1]}\, \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_x)\, \Pi_n^{(x)[\mathbf{W}_1]} \middle| \mathring{\mathbf{\Psi}} \right\rangle. \tag{4.212}$$

This is the diagonal restriction of the bilocal Fourfold operator, projected to the nth wordwise band at x.

Definition 4.467 (Holo-State band populations (wordwise))**.** For a word $\mathbf{W}$ at x, define the band population $\rho_n^{[\mathbf{W}]}(x) := \left\langle \mathring{\mathbf{\Psi}} \middle| \Pi_n^{(x)[\mathbf{W}]} \middle| \mathring{\mathbf{\Psi}} \right\rangle \in [0,1]$. Then $\sum_n \rho_n^{[\mathbf{W}]}(x) = 1$ on the fibre at x (completeness of the spectral resolution).

Definition 4.468 (Wordwise vacuum–response distance (state–conditioned, symmetric weighting))**.** Let $\mathbf{W}_1, \mathbf{W}_2$ be admissible words at x, x' and choose compatible wordwise frames on a common patch (Definition 4.235 and Proposition 4.244). Define the *vacuum profile* vectors $\vec{V}_{\mathrm{vac}}^{[\mathbf{W}_1]}(x) := (C_1^{[\mathbf{W}_1]}(x), C_2^{[\mathbf{W}_1]}(x), \dots)$ and $\vec{V}_{\mathrm{vac}}^{[\mathbf{W}_2]}(x')$ using (4.212). With the symmetric state weights

$$\theta_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \;:=\; \tfrac{1}{2}\Big(\rho_n^{[\mathbf{W}_1]}(x) + \rho_n^{[\mathbf{W}_2]}(x')\Big) \in [0,1], \qquad \sum_n \theta_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') = 1,$$

the *wordwise vacuum–response distance* is the weighted ℓ^2–difference

$$\boxed{d_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \;:=\; \left(\sum_{n=1}^{\infty} \theta_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \,\Big|\, C_n^{[\mathbf{W}_1]}(x) - C_n^{[\mathbf{W}_2]}(x') \,\Big|^2\right)^{1/2}.} \tag{4.213}$$

When needed to compare profiles in a single fibre, transport the x'–profile to x with the wordwise transporter $\mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x \leftarrow x'; \gamma)$ (Section 4.5); the resulting value is independent of the choice of compatible trivialization and is Lipschitz in the holonomy size of γ (cf. Lemma 4.413 and Corollary 4.414).

Vector formulation and invariances

Proposition 4.469 (Vector form and word/gauge invariance)**.** *The expression* (4.213) *equals the (symmetrically weighted) ℓ^2–distance between wordwise vacuum profiles and is invariant under endpoint word changes and endpoint gauge transformations. On flat patches the value is path independent; otherwise it varies at most linearly in the holonomy size.*

Proof. Changing words at an endpoint conjugates both the projectors and the diagonal operator by a local unitary; the Holo-State evaluation is invariant, hence each $C_n^{[\cdot]}$ and $\rho_n^{[\cdot]}$ is unchanged and so is (4.213). Gauge covariance and holonomy bounds follow from Section 4.5, Lemma 4.413, and Corollary 4.414; flatness yields path independence by Proposition 4.59. □

Well-definedness and finiteness

Lemma 4.470 (Bandwise bounds and absolute finiteness)**.** *For each n and locus x, $|C_n^{[\mathbf{W}_1]}(x)| \le \big\|\Pi_n^{(x)[\mathbf{W}_1]}\, \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_x)\, \Pi_n^{(x)[\mathbf{W}_1]}\big\| \le \|\hat{G}_\Lambda\|$. Consequently,*

$$\Big(d_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\Big)^2 \;\le\; \sum_n \theta_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\,(2\|\hat{G}_\Lambda\|)^2 \;=\; 4\|\hat{G}_\Lambda\|^2,$$

so $d_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') < \infty$ without any additional hypothesis.

Proof. **Step 1: Set-up and notation.** Fix an admissible word $\mathbf{W}_1$ at x and denote the band projector by $\Pi_n := \Pi_n^{(x)[\mathbf{W}_1]}$ (Definition 4.405). Let

$$A_x \;:=\; \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_x)$$

be the diagonal (coincident) Fourfold operator at x. By Definition 4.466, the nth band self-correlator is $C_n^{[\mathbf{W}_1]}(x) = \langle \overset{\circ}{\mathbf{\Psi}} | \Pi_n A_x \Pi_n | \overset{\circ}{\mathbf{\Psi}} \rangle$.

Step 2: Expectation bounded by operator norm. The Holo-State is normalized in the emergent fibre inner product (Section 4.6), hence for any bounded operator B on the fibre,

$$\left| \langle \overset{\circ}{\mathbf{\Psi}} | \, B \, | \overset{\circ}{\mathbf{\Psi}} \rangle \right| \;\le\; \|B\|.$$

Applying this to $B = \Pi_n A_x \Pi_n$ gives the first inequality:

$$|C_n^{[\mathbf{W}_1]}(x)| \;\le\; \|\Pi_n A_x \Pi_n\|.$$

Step 3: Sandwiching by a projector does not increase the norm. Since Π_n is an orthogonal projector, $\|\Pi_n\| = 1$ and $\Pi_n^\dagger = \Pi_n$. Therefore

$$\|\Pi_n A_x \Pi_n\| \;\le\; \|\Pi_n\| \, \|A_x\| \, \|\Pi_n\| \;=\; \|A_x\|.$$

By definition of A_x and the uniform bound on the family $x \mapsto A_x$, $\|A_x\| \le \|\hat{G}_\Lambda\|$. This yields the second inequality and hence the bandwise bound

$$|C_n^{[\mathbf{W}_1]}(x)| \;\le\; \|\hat{G}_\Lambda\| \qquad \text{for all } n.$$

Step 4: Weighted ℓ^2–bound for the vacuum–response distance. By Definition 4.468, with symmetric state weights $\theta_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \in [0,1]$ satisfying $\sum_n \theta_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') = 1$,

$$\left(d_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\right)^2 = \sum_n \theta_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \left| C_n^{[\mathbf{W}_1]}(x) - C_n^{[\mathbf{W}_2]}(x') \right|^2.$$

Using the elementary inequality $|a-b| \le |a| + |b|$ and the bandwise bounds from Step 3,

$$\left| C_n^{[\mathbf{W}_1]}(x) - C_n^{[\mathbf{W}_2]}(x') \right|^2 \;\le\; \left(|C_n^{[\mathbf{W}_1]}(x)| + |C_n^{[\mathbf{W}_2]}(x')| \right)^2 \;\le\; (2\|\hat{G}_\Lambda\|)^2.$$

Therefore,

$$\left(d_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\right)^2 \;\le\; \sum_n \theta_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \, (2\|\hat{G}_\Lambda\|)^2 = (2\|\hat{G}_\Lambda\|)^2 \sum_n \theta_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') = 4\|\hat{G}_\Lambda\|^2.$$

Step 5: Conclusion. The right-hand side is finite and independent of (x,x'), $\mathbf{W}_{1,2}$, and γ, so $d_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') < \infty$ holds *unconditionally*—no additional decay or capacity assumptions are needed beyond boundedness of $\hat{G}_\Lambda$ and normalization of the weights (built into the definition of θ_n). □

Proposition 4.471 (Basic properties)**.** *The quantity $d_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')$ is nonnegative, symmetric, vanishes at coincidence, and obeys the invariances of Proposition 4.469. On flat patches it is window independent.*

Proof. Recall the definition (Definitions 4.467 and 4.468):

$$\Big(d_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\Big)^2 = \sum_{n\geq 1} \theta_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\,\Big|\,C_n^{[\mathbf{W}_1]}(x) - C_n^{[\mathbf{W}_2]}(x')\,\Big|^2,$$
$$\theta_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') := \tfrac{1}{2}\Big(\rho_n^{[\mathbf{W}_1]}(x) + \rho_n^{[\mathbf{W}_2]}(x')\Big).$$

with $C_n^{[\mathbf{W}]}$ from Definition 4.466. We prove each property.

(a) Nonnegativity. Each summand is $\theta_n \geq 0$ times a square, hence the series is ≥ 0 and the square root is nonnegative by definition of the $\ell^2(\theta)$–norm (Definition 4.468).

(b) Symmetry. Exchanging $(x,\mathbf{W}_1)$ with $(x',\mathbf{W}_2)$ leaves the weights invariant by construction, $\theta_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') = \theta_n^{[\mathbf{W}_2,\mathbf{W}_1]}(x',x)$, and also $\big|C_n^{[\mathbf{W}_1]}(x) - C_n^{[\mathbf{W}_2]}(x')\big| = \big|C_n^{[\mathbf{W}_2]}(x') - C_n^{[\mathbf{W}_1]}(x)\big|$. Thus the sum is unchanged, giving $d_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') = d_{\mathrm{vac}}^{[\mathbf{W}_2,\mathbf{W}_1]}(x',x)$.

(c) Vanishing at coincidence. If $x = x'$ and $\mathbf{W}_1 = \mathbf{W}_2$ with the same (compatible) local frame (Definition 4.405), then $C_n^{[\mathbf{W}_1]}(x) = C_n^{[\mathbf{W}_2]}(x')$ for all n, so every summand vanishes and $d_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x) = 0$. If different words/gauges are chosen at the same point, invariance under endpoint word changes and endpoint gauge transformations (proved in Proposition 4.469) implies the same conclusion.

(d) Word/gauge invariance and holonomy control. Under an endpoint word change or endpoint gauge transformation, both the band projectors and the diagonal operator $\hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_x)$ transform by conjugation, and Holo-State expectations are invariant; likewise the band populations ρ_n are unchanged. Hence *each* C_n and θ_n is invariant, proving the stated invariances; see Proposition 4.469 for the detailed argument based on Section 4.5. When a transporter is used to compare profiles in one fibre, the dependence on the window γ is Lipschitz in the holonomy size by Lemma 4.413 and Corollary 4.414 (the diagonal evaluation is stable under bounded-holonomy conjugations).

(e) Window independence on flat patches. If the modal curvature vanishes on a patch, $\mathcal{F} \equiv 0$, then transport is path independent (Proposition 4.59); the holonomy around any loop is trivial, so any transported comparison of the x'–profile to the x–fibre is independent of the chosen window. Equivalently, the holonomy size $\Phi(\Delta\gamma) = 0$, and the Holonomy–Lipschitz control yields identical values for all windows in the patch (Lemma 4.448 and Corollary 4.414). Therefore $d_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')$ is window independent on flat patches.

(f) Boundedness (useful corollary). From Lemma 4.470 we have the uniform bound $\Big(d_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\Big)^2 \leq 4\|\hat{G}_\Lambda\|^2$, so d_{vac} is finite and globally bounded without extra hypotheses.

These establish all the claimed basic properties. □

Wordwise vector form and functional orthogonality

Corollary 4.472 (Vector form)**.** *With* $\vec{V}_{\mathrm{vac}}^{[\mathbf{W}]}(x)$ *as above,*

$$\boxed{d_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') = \Big\|\vec{V}_{\mathrm{vac}}^{[\mathbf{W}_1]}(x) - \vec{V}_{\mathrm{vac}}^{[\mathbf{W}_2]}(x')\Big\|_{\ell^2(\theta)},} \tag{4.214}$$

where $\|\cdot\|_{\ell^2(\theta)}$ *is the* ℓ^2*–norm with weights* $\theta_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') = \frac{1}{2}\Big(\rho_n^{[\mathbf{W}_1]}(x) + \rho_n^{[\mathbf{W}_2]}(x')\Big)$ *(Definition 4.467). Furthermore,* d_{vac} *probes the* diagonal *sector of* $\hat{G}_\Lambda$ *and is functionally orthogonal*

to $d_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]}$ *(off-diagonal bilocal overlap, Definition 4.339) and to* $d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}$ *(commutator/differential response, Definition 4.408).*

Proof. **(1) Equality with the weighted ℓ^2–norm.** By Definition 4.466, set $\vec{V}_{\mathrm{vac}}^{[\mathbf{W}_1]}(x) := \big(C_1^{[\mathbf{W}_1]}(x), C_2^{[\mathbf{W}_1]}(x), \dots\big)$ and similarly for $\vec{V}_{\mathrm{vac}}^{[\mathbf{W}_2]}(x')$. Unpacking the definition of $d_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}$ (Definition 4.468) gives

$$\Big(d_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\Big)^2 = \sum_{n=1}^{\infty} \theta_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \left| C_n^{[\mathbf{W}_1]}(x) - C_n^{[\mathbf{W}_2]}(x') \right|^2 = \left\| \vec{V}_{\mathrm{vac}}^{[\mathbf{W}_1]}(x) - \vec{V}_{\mathrm{vac}}^{[\mathbf{W}_2]}(x') \right\|_{\ell^2(\theta)}^2.$$

Taking the square root yields (4.214).

(2) Diagonal-sector dependence. Write the diagonal Fourfold operator at x as $A_x := \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_x)$ and decompose it in the wordwise spectral frame of $\mathbb{D}^{[\mathbf{W}_1]}(x)$ (Definition 4.405):

$$A_x = \underbrace{\sum_n \Pi_n^{(x)[\mathbf{W}_1]} A_x \Pi_n^{(x)[\mathbf{W}_1]}}_{=:A_x^{\mathrm{diag}}} + \underbrace{\sum_{m\neq n} \Pi_m^{(x)[\mathbf{W}_1]} A_x \Pi_n^{(x)[\mathbf{W}_1]}}_{=:A_x^{\mathrm{off}}}.$$

By Definition 4.466, $C_n^{[\mathbf{W}_1]}(x) = \langle\Psi|\, \Pi_n A_x \Pi_n \,|\Psi\rangle = \langle\Psi|\, \Pi_n A_x^{\mathrm{diag}} \Pi_n \,|\Psi\rangle$, while $\Pi_n A_x^{\mathrm{off}} \Pi_n = 0$. Hence each $C_n^{[\mathbf{W}_1]}(x)$ depends only on the *diagonal* part A_x^{diag}; the same holds at x' for $C_n^{[\mathbf{W}_2]}(x')$. Therefore d_{vac} is a functional of the diagonal sector of $\hat{G}_\Lambda$ only.

(3) Functional orthogonality to d_{corr}. The correlator distance $d_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]}$ is built from *bilocal* overlaps (Definition 4.339), i.e. from off-diagonal (in locus) blocks of the Fourfold operator and their connected parts. Consider a perturbation δA_x that is purely off-diagonal in the wordwise band sense, so that $\Pi_n^{(x)[\mathbf{W}_1]} \delta A_x \Pi_n^{(x)[\mathbf{W}_1]} = 0$ for all n. Then

$$\delta C_n^{[\mathbf{W}_1]}(x) = \langle\Psi|\, \Pi_n \delta A_x \Pi_n \,|\Psi\rangle = 0 \quad (\forall n),$$

hence $\delta d_{\mathrm{vac}} = 0$ to first order, while d_{corr} may change (it is sensitive to off-diagonal/bilocal structure). Conversely, diagonal variations $\delta A_x^{\mathrm{diag}}$ affect d_{vac} but do not contribute to the connected *off-diagonal* overlaps that define d_{corr}. Thus, in the operational sense of sensitivity to disjoint operator sectors, d_{vac} and d_{corr} are functionally orthogonal.

(4) Functional orthogonality to d_{spec}. The spectral distance $d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}$ is built from spectral-shear components that are governed by commutator/differential response with the spectral operator (Definitions 4.406 and 4.408). If δA_x is diagonal in the spectral frame of $\mathbb{D}^{[\mathbf{W}_1]}(x)$, then $[\mathbb{D}^{[\mathbf{W}_1]}(x), \delta A_x] = 0$, and (by the Helffer–Sjöstrand control developed for the shear) the corresponding shear variation vanishes. Hence such diagonal variations change d_{vac} but do not alter d_{spec} to first order; conversely, purely commutator-driven variations that keep the diagonal blocks fixed change d_{spec} while leaving d_{vac} unchanged. This again establishes functional orthogonality in the sense of sensitivity to complementary sectors (diagonal vs commutator directions).

All claims of the corollary follow. □

Twofold reduction and equivalence

Corollary 4.473 (Twofold reduction)**.** *In the duality-collapse regime (Definition 4.87), the wordwise vacuum-response reduces to the corresponding qubit-level notion on the* $\{0,1\}$ *sector, with the same formula* (4.214) *restricted to that slice.*

Proposition 4.474 (Equivalence to the exogenous vacuum distance)**.** *Let $d_{\mathrm{vac}}(x,x')$ be as in Definition 4.463. For any admissible words $\mathbf{W}_1,\mathbf{W}_2$ on a patch,*

$$d_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \;=\; d_{\mathrm{vac}}(x,x'),$$

provided the exogenous definition uses the same symmetric state weighting as in (4.213).

Proof. We compare the *wordwise* definition (Definitions 4.466 to 4.468) with the *exogenous* one (Definition 4.463). Write

$$A_x := \hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_x), \qquad A_{x'} := \hat{G}_\Lambda(\mathfrak{B}_{x'},\mathfrak{B}_{x'})$$

for the diagonal Fourfold operators at the two loci.

Step 1: Aligning frames by unitary word rotations. Let $\{\Pi_n^{(x)[\mathbf{W}_1]}\}_n$ be the wordwise spectral frame of $\mathbb{D}^{[\mathbf{W}_1]}(x)$ and $\{\Pi_{n'}^{(x')[\mathbf{W}_2]}\}_{n'}$ that of $\mathbb{D}^{[\mathbf{W}_2]}(x')$ (Definition 4.405). Choose exogenous modal frames $\{\widehat{\Pi}_n^{(x)}\}_n$, $\{\widehat{\Pi}_{n'}^{(x')}\}_{n'}$ related by local unitaries $U_x,U_{x'}$ (the inverse word rotations) via

$$\widehat{\Pi}_n^{(x)} \;=\; U_x^\dagger\,\Pi_n^{(x)[\mathbf{W}_1]}\,U_x, \qquad \widehat{\Pi}_{n'}^{(x')} \;=\; U_{x'}^\dagger\,\Pi_{n'}^{(x')[\mathbf{W}_2]}\,U_{x'}. \tag{4.215}$$

By construction, this is just a change of basis on each fibre (no physical operation).

Step 2: Invariance of band populations (weights). The wordwise band populations are $\rho_n^{[\mathbf{W}_1]}(x) = \langle\Psi|\Pi_n^{(x)[\mathbf{W}_1]}|\Psi\rangle$ and similarly at x' (Definition 4.467). The exogenous ones are $\widehat{\rho}_n(x) = \langle\Psi|\widehat{\Pi}_n^{(x)}|\Psi\rangle$. Using (4.215),

$$\widehat{\rho}_n(x) = \langle\Psi|U_x^\dagger\Pi_n^{(x)[\mathbf{W}_1]}U_x|\Psi\rangle = \langle\Psi_x|\Pi_n^{(x)[\mathbf{W}_1]}|\Psi_x\rangle, \qquad \Psi_x := U_x|\Psi\rangle.$$

But the evaluation functional is basis-covariant (it is the same scalar under a passive change of basis); equivalently, this is the endpoint word/gauge invariance from Proposition 4.469. Hence $\widehat{\rho}_n(x) = \rho_n^{[\mathbf{W}_1]}(x)$ and likewise $\widehat{\rho}_{n'}(x') = \rho_{n'}^{[\mathbf{W}_2]}(x')$. Therefore the *symmetric weights* coincide in the two presentations:

$$\widehat{\theta}_n^{[\cdot]}(x,x') = \tfrac{1}{2}\big(\widehat{\rho}_n(x)+\widehat{\rho}_n(x')\big) = \tfrac{1}{2}\big(\rho_n^{[\mathbf{W}_1]}(x)+\rho_n^{[\mathbf{W}_2]}(x')\big) = \theta_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x'). \tag{4.216}$$

Step 3: Invariance of band self-correlators. The exogenous diagonal band correlators at x are $\widehat{C}_n(x) = \langle\Psi|\,\widehat{\Pi}_n^{(x)}A_x\widehat{\Pi}_n^{(x)}\,|\Psi\rangle$. Using (4.215) and cyclicity,

$$\widehat{C}_n(x) = \langle\Psi|U_x^\dagger\Pi_n^{(x)[\mathbf{W}_1]}U_x\,A_x\,U_x^\dagger\Pi_n^{(x)[\mathbf{W}_1]}U_x|\Psi\rangle = \langle\Psi_x|\,\Pi_n^{(x)[\mathbf{W}_1]}\,A_x\,\Pi_n^{(x)[\mathbf{W}_1]}\,|\Psi_x\rangle.$$

Again by endpoint invariance (passive basis change), this equals the wordwise value $C_n^{[\mathbf{W}_1]}(x)$ from Definition 4.466. The same holds at x' for $C_{n'}^{[\mathbf{W}_2]}(x')$. Possible unitary rotations *within a degenerate band* leave Π_n unchanged, so the scalar is unaffected.

Step 4: Equality of the weighted ℓ^2 distances. Let the exogenous vacuum profiles be $\widehat{\vec{V}}_{\mathrm{vac}}(x) := (\widehat{C}_1(x),\widehat{C}_2(x),\dots)$ (and similarly at x'). By Steps 2–3 and (4.216),

$$\big\|\widehat{\vec{V}}_{\mathrm{vac}}(x)-\widehat{\vec{V}}_{\mathrm{vac}}(x')\big\|^2_{\ell^2(\widehat{\theta})} = \sum_n \widehat{\theta}_n\,|\widehat{C}_n(x)-\widehat{C}_n(x')|^2 = \sum_n \theta_n^{[\mathbf{W}_1,\mathbf{W}_2]}\,|C_n^{[\mathbf{W}_1]}(x)-C_n^{[\mathbf{W}_2]}(x')|^2,$$

which is precisely $\left(d_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\right)^2$ by Definition 4.468. Taking square roots yields

$$d_{\mathrm{vac}}(x,x') = d_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x').$$

Step 5 (optional): Transported comparisons. If either definition transports the x'–profile to x via a window γ (Section 4.5), the transported profile differs by conjugation with the transporter. The value is unchanged by Proposition 4.469; on flat patches path independence holds (Proposition 4.59), and in general the dependence is Holonomy–Lipschitz (Lemma 4.448 and Corollary 4.414), but the two presentations remain identical when the same γ is used.

These steps establish the claimed equivalence. □

Interpretation (wordwise). $d_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}$ measures changes in *self-coherence* of the vacuum between loci, using only coincident (diagonal) relational data. It is state-conditioned (via θ_n), gauge/word invariant, UV-finite (no external cutoff) by boundedness of $\hat{G}_\Lambda$ and normalization $\sum_n \theta_n = 1$, and complementary to the correlational and spectral axes in the triadic geometry. On flat patches it is independent of the comparison window; more generally, it varies at most linearly with holonomy (via Lemma 4.413 and Corollary 4.414). Under the Finite Entanglement Capacity principle (Principle 4.2), stronger decay estimates for the profiles follow immediately from the band-population bounds.

4.10 The Triadic Metric Structure of the PQS Continuum

Having established the three relational distances

$$\begin{aligned} d_{\mathrm{corr}}(x,x') &\quad \text{(Section 4.9.1)},\\ d_{\mathrm{vac}}(x,x') &\quad \text{(Definition 4.463)},\\ d_{\mathrm{spec}}(x,x') &\quad \text{(Section 4.9.9)}. \end{aligned}$$

we show that they furnish an *orthogonal triad* from which a composite metric on the emergent configuration space $\mathcal{X}_q$ is assembled. The resulting geometry is fibrewise and operator-informed: its axes arise from algebraically independent projections of the Fourfold correlator $\hat{G}_\Lambda$ and the Spectral Operator $\hat{D}$.

4.10.1 Functional Orthogonality of the Distance Measures

Probe Space and Projections. Let $\mathscr{P}$ denote the linear span of admissible relational probes built from $\hat{G}_\Lambda$ and $\hat{D}$ (bounded on the emergent Hilbert space), and endow $\mathscr{P}$ with the Holo-State sesquilinear form

$$\langle\!\langle A,B\rangle\!\rangle_\Psi \;:=\; \left\langle \mathring{\boldsymbol{\Psi}} \middle| A^\dagger B \middle| \mathring{\boldsymbol{\Psi}} \right\rangle, \qquad A,B\in\mathscr{P}.$$

Define the following $\langle\!\langle \cdot, \cdot \rangle\!\rangle_\Psi$–closed subspaces:

$$\mathscr{P}_{\text{corr}} := \text{span}\left\{ \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2] \;:\; x \neq x' \right\},$$
$$\mathscr{P}_{\text{vac}} := \text{span}\left\{ \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_x)[\psi^*, \psi] \;-\; \hat{G}_\Lambda(\mathfrak{B}_{x'}, \mathfrak{B}_{x'})[\phi^*, \phi] \right\},$$
$$\mathscr{P}_{\text{spec}} := \text{span}\left\{ [\hat{D},\, \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi^*, \psi]] \right\}.$$

Write $P_\bullet$ for the orthogonal projections onto these subspaces. The three distances use only the norms induced by P_{corr}, P_{vac}, and P_{spec} respectively (cf. their definitions).

Proposition 4.475 (Functional Orthogonality)**.** *The subspaces $\mathscr{P}_{\text{corr}}, \mathscr{P}_{\text{vac}}, \mathscr{P}_{\text{spec}}$ are mutually orthogonal with respect to $\langle\!\langle \cdot, \cdot \rangle\!\rangle_\Psi$. Equivalently, $P_{\text{corr}} P_{\text{vac}} = P_{\text{vac}} P_{\text{spec}} = P_{\text{spec}} P_{\text{corr}} = 0$.*

Sketch. (i) Elements of $\mathscr{P}_{\text{corr}}$ are bilocal off-diagonal ($x \neq x'$); those of $\mathscr{P}_{\text{vac}}$ are differences of diagonal (coincident) objects. Their supports are disjoint in the coincident/off-diagonal decomposition of $\hat{G}_\Lambda$, whence cross inner products vanish. (ii) For $A \in \mathscr{P}_{\text{spec}}$ one has $A^\dagger = -A$ (commutators are anti-Hermitian), while $B \in \mathscr{P}_{\text{corr}} \cup \mathscr{P}_{\text{vac}}$ is Hermitian; thus $\langle\!\langle A, B \rangle\!\rangle_\Psi$ is purely imaginary and its real norm contribution vanishes. A direct polarization argument yields mutual orthogonality. □

4.10.2 Composite Triadic Distance and Metric

Triadic Fibre and Calibration. At each pair (x, x') define the *triadic distance vector*

$$\vec{d}(x, x') \;:=\; \Big(d_{\text{corr}}(x, x'),\; d_{\text{vac}}(x, x'),\; d_{\text{spec}}(x, x') \Big)^\top \;\in\; \mathbb{R}^3_{\geq 0}.$$

Let $\mathbf{M}$ be a fixed (or state-/context-dependent) 3×3 symmetric positive-definite *calibration matrix* acting on this fibre (e.g. $\mathbf{M} = \text{diag}(\alpha^2, \beta^2, \gamma^2)$ with $\alpha, \beta, \gamma > 0$).

Definition 4.476 (Composite Triadic Distance)**.** The *triadic distance* is

$$D(x, x') \;:=\; \left\| \mathbf{M}^{1/2}\, \vec{d}(x, x') \right\|_2 \;=\; \left(\vec{d}(x, x')^\top \mathbf{M}\, \vec{d}(x, x') \right)^{1/2}. \tag{4.217}$$

Proposition 4.477 (Metric Properties)**.** *If each component $d_{\text{corr}}, d_{\text{vac}}, d_{\text{spec}}$ satisfies the metric axioms on $\mathcal{X}_q$, then D is a metric on $\mathcal{X}_q$ for any symmetric positive-definite $\mathbf{M}$. In particular, D is nonnegative, symmetric, definite, and obeys the triangle inequality.*

Proof. Nonnegativity and symmetry are immediate from (4.217). Definiteness follows since $\mathbf{M} \succ 0$ and each component distance is definite. For the triangle inequality, Minkowski's inequality in ℓ^2 yields $\|\mathbf{M}^{1/2}(\vec{d}(x, z))\|_2 \leq \|\mathbf{M}^{1/2}(\vec{d}(x, y))\|_2 + \|\mathbf{M}^{1/2}(\vec{d}(y, z))\|_2$. □

Remark 4.478 (Invariance and Reduction)**.** D is invariant under admissible reindexings and the duality involution (Proposition 1.8) because each component is. In a duality-collapse regime (Definition 4.87), D reduces to the corresponding qubit-level composite distance under the isometries of Proposition 4.328.

4.10.3 Infinitesimal Metric Tensor from the Triad

Define the squared triadic distance $D^2(x, x')$ as in (4.217). When (x', x) are linked by an admissible chart in $\mathcal{X}_q$ with coordinates x^μ, the (operator-evaluated) emergent metric at x is obtained by the coincidence-limit Hessian:

$$g_{\mu\nu}(x) \;:=\; \frac{1}{2}\,\frac{\partial^2}{\partial x'^{\mu}\,\partial x'^{\nu}}\,D^2(x,x')\Big|_{x'=x}. \tag{4.218}$$

Because D^2 is a calibrated sum of *orthogonal* components (Prop. 4.475), cross-variations vanish in the limit, and $g_{\mu\nu}$ decomposes additively into correlational, vacuum, and spectral contributions, each gauge-covariant and finite by construction (Sections 4.9.1 and 4.9.9, Definition 4.463, and Proposition 4.424).

Axis	**Geometric interpretation**	**Operator construction (in $\lvert\mathring{\Psi}\rangle$)**
$d_{\mathrm{corr}}(x, x')$	Off–diagonal entanglement/overlap geometry; first–order relational coherence	Normalised Holo-State expectation of the bilocal correlator $\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2]$ (Section 4.9.1 and Definition 4.338)
$d_{\mathrm{vac}}(x, x')$	Vacuum self–coherence curvature (coincident/diagonal sector); second–order variation	RMS of diagonal differences $C_k(x) - C_k(x')$ with $C_k(\cdot) = \left\langle \hat{G}_\Lambda(\cdot,\cdot)[\psi_k^*, \psi_k] \right\rangle$ (Definition 4.463)
$d_{\mathrm{spec}}(x, x')$	Spectral/differential geometry generated by $\hat{D}$	Weighted RMS of commutators $[\hat{D}, \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_n^*, \psi_n]]$ with modal weights $w_n(x, x')$ (Definitions 4.386, 4.392 and 4.402)

Table 4.2: Triadic axes of the emergent distance on $\mathcal{X}_q$, showing complementary geometric roles and operator-theoretic origins. All expectations are taken in the Quantum Holo-State $\lvert\mathring{\Psi}\rangle$ (Definition 1.27).

4.11 Holosymmetric Wordwise Triadic Metric Structure of the PQS Continuum

Having established the three *wordwise* relational distances

$$\begin{aligned} &d_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \quad \text{(Definition 4.339)},\\ &d_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \quad \text{(Definition 4.468)},\\ &d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \quad \text{(Definition 4.408)}, \end{aligned}$$

we show that they furnish an *orthogonal triad* from which a composite metric on the emergent configuration space $\mathcal{X}_q$ is assembled. The resulting geometry is fibrewise and operator-

informed: its axes arise from algebraically independent projections of the Fourfold correlator $\hat{G}_\Lambda$ and the wordwise spectral generator $\mathbb{D}^{[\mathbf{W}]}$.

4.11.1 Functional Orthogonality of the Wordwise Distance Measures

Wordwise probe space and projections. Let $\mathscr{P}^{\mathrm{w}}$ denote the linear span of admissible *wordwise* relational probes built from $\hat{G}_\Lambda$ and $\mathbb{D}^{[\mathbf{W}]}$ (bounded on the emergent Hilbert space), and endow $\mathscr{P}^{\mathrm{w}}$ with the *state-conditioned real (Jordan) sesquilinear form*

$$\langle\!\langle A, B\rangle\!\rangle_\Psi \;:=\; \mathrm{Re}\,\Big\langle \mathring{\mathbf{\Psi}}\Big|\, A^\dagger B\,\Big|\mathring{\mathbf{\Psi}}\Big\rangle \;=\; \tfrac{1}{2}\Big\langle \mathring{\mathbf{\Psi}}\Big|\, A^\dagger B + B^\dagger A\,\Big|\mathring{\mathbf{\Psi}}\Big\rangle, \qquad A, B \in \mathscr{P}^{\mathrm{w}}.$$

Define the following $\langle\!\langle\cdot,\cdot\rangle\!\rangle_\Psi$–closed subspaces (with word/gauge-compatible frames as in Sections 4.6.2 and 4.6.5):

$$\mathscr{P}^{\mathrm{w}}_{\mathrm{corr}} := \mathrm{span}\Big\{\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2] \;\Big|\; x \neq x'\Big\}, \tag{4.219}$$

$$\mathscr{P}^{\mathrm{w}}_{\mathrm{vac}} := \mathrm{span}\Big\{\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_x)[\psi^*, \psi] - \hat{G}_\Lambda(\mathfrak{B}_{x'}, \mathfrak{B}_{x'})[\phi^*, \phi]\Big\}, \tag{4.220}$$

$$\mathscr{P}^{\mathrm{w}}_{\mathrm{spec}} := \mathrm{span}\Big\{[\mathbb{D}^{[\mathbf{W}]},\, \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi^*, \psi]]\Big\}, \tag{4.221}$$

$$= \mathrm{span}\Big\{\Pi_n^{(x)[\mathbf{W}]}\, \Delta\mathbb{D}^{[\mathbf{W}_1,\mathbf{W}_2]}(x, x')\, \Pi_n^{(x)[\mathbf{W}]}\Big\}. \tag{4.222}$$

Write $P^{\mathrm{w}}_\bullet$ for the orthogonal projections onto these subspaces with respect to $\langle\!\langle\cdot,\cdot\rangle\!\rangle_\Psi$.

Proposition 4.479 (Functional orthogonality (wordwise))**.** *The subspaces $\mathscr{P}^{\mathrm{w}}_{\mathrm{corr}}$, $\mathscr{P}^{\mathrm{w}}_{\mathrm{vac}}$, $\mathscr{P}^{\mathrm{w}}_{\mathrm{spec}}$ are mutually orthogonal with respect to $\langle\!\langle\cdot,\cdot\rangle\!\rangle_\Psi$; i.e.,*

$$\langle\!\langle A, B\rangle\!\rangle_\Psi = 0 \quad \textit{whenever} \quad A \in \mathscr{P}^{\mathrm{w}}_{\mathrm{corr}}, \quad B \in \mathscr{P}^{\mathrm{w}}_{\mathrm{vac}} \cup \mathscr{P}^{\mathrm{w}}_{\mathrm{spec}}, \quad \textit{or cyclic permutations.}$$

Equivalently, the projections satisfy $P^{\mathrm{w}}_{\mathrm{corr}} P^{\mathrm{w}}_{\mathrm{vac}} = P^{\mathrm{w}}_{\mathrm{vac}} P^{\mathrm{w}}_{\mathrm{spec}} = P^{\mathrm{w}}_{\mathrm{spec}} P^{\mathrm{w}}_{\mathrm{corr}} = 0$.

Sketch. (i) Elements of $\mathscr{P}^{\mathrm{w}}_{\mathrm{corr}}$ are bilocal off-diagonal ($x \neq x'$); those of $\mathscr{P}^{\mathrm{w}}_{\mathrm{vac}}$ are differences of diagonal (coincident) objects. Their supports are disjoint in the coincident/off-diagonal block decomposition of $\hat{G}_\Lambda$ relative to wordwise frames. In the Jordan form $\langle\!\langle\cdot,\cdot\rangle\!\rangle_\Psi$, the cross-terms pick only diagonal-in-block contributions; hence $\langle\!\langle A_{\mathrm{corr}}, B_{\mathrm{vac}}\rangle\!\rangle_\Psi = 0$. (ii) For $A \in \mathscr{P}^{\mathrm{w}}_{\mathrm{spec}}$, $A^\dagger = -A$ (commutators are anti-Hermitian), while $B \in \mathscr{P}^{\mathrm{w}}_{\mathrm{corr}} \cup \mathscr{P}^{\mathrm{w}}_{\mathrm{vac}}$ is Hermitian; thus $A^\dagger B + B^\dagger A = -AB + BA = [B, A]$, which is anti-Hermitian, and its Holo-State expectation is purely imaginary. Taking the real part gives $\langle\!\langle A, B\rangle\!\rangle_\Psi = 0$. Polarization then yields the stated mutual orthogonality. □

4.11.2 Composite Wordwise Triadic Distance and Metric

Triadic fibre and calibration. At each pair (x, x') define the *wordwise triadic distance vector*

$$\vec{d}^{[\mathbf{W}_1,\mathbf{W}_2]}(x, x') := \Big(d^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{corr}}(x, x'),\; d^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{vac}}(x, x'),\; d^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{spec}}(x, x')\Big)^\top \in \mathbb{R}^3_{\geq 0}.$$

Let $\mathbf{M}$ be a fixed (or state-/context-dependent) 3×3 symmetric positive-definite *calibration matrix* (e.g. $\mathbf{M} = \mathrm{diag}(\alpha^2, \beta^2, \gamma^2)$ with $\alpha, \beta, \gamma > 0$).

Definition 4.480 (Composite wordwise triadic distance)**.**

$$D^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') := \left\|\mathbf{M}^{1/2}\,\vec{d}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\right\|_2 = \left(\,\vec{d}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')^{\top}\mathbf{M}\,\vec{d}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\,\right)^{1/2}. \tag{4.223}$$

Proposition 4.481 (Metric properties (wordwise))**.** *If each component $d^{[\cdot]}_{\mathrm{corr}}, d^{[\cdot]}_{\mathrm{vac}}, d^{[\cdot]}_{\mathrm{spec}}$ satisfies the metric axioms on $\mathcal{X}_q$, then $D^{[\mathbf{W}_1,\mathbf{W}_2]}$ is a metric on $\mathcal{X}_q$ for any $\mathbf{M} \succ 0$. In particular, $D^{[\mathbf{W}_1,\mathbf{W}_2]}$ is nonnegative, symmetric, definite, and obeys the triangle inequality.*

Proof. Immediate from (4.223) and Minkowski's inequality in ℓ^2. □

Remark 4.482 (Invariance, Lipschitz window dependence, and exogenous equivalence)**.** $D^{[\mathbf{W}_1,\mathbf{W}_2]}$ is invariant under endpoint word/gauge changes by the invariances of its components (Proposition 4.469 and the corresponding statements for $d^{[\cdot]}_{\mathrm{corr}}$ and $d^{[\cdot]}_{\mathrm{spec}}$). Along different windows, it is Lipschitz in the holonomy size (Lemma 4.413 and Corollary 4.414) and path independent on flat patches. Moreover,

$$D^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') = D(x,x') \quad \text{with } D \text{ from the exogenous composite definition,}$$

by the equivalence of the wordwise and exogenous components (e.g. Proposition 4.474 together with the analogous equivalences for d_{corr} and d_{spec}).

4.11.3 Infinitesimal Wordwise Metric Tensor from the Triad

Define the squared wordwise triadic distance $D^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')^2$ as in (4.223). When (x',x) are linked by an admissible chart in $\mathcal{X}_q$ with coordinates x^μ, the (operator-informed) emergent metric at x is obtained by the coincidence-limit Hessian:

$$g^{[\mathbf{W}]}_{\mu\nu}(x) := \frac{1}{2}\,\frac{\partial^2}{\partial x'^{\mu}\,\partial x'^{\nu}}\,D^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')^2\Big|_{x'=x}. \tag{4.224}$$

Because D^2 is a calibrated sum of *orthogonal* components (Prop. 4.479), cross-variations vanish in the limit, and $g^{[\mathbf{W}]}_{\mu\nu}$ decomposes additively into correlational, vacuum, and spectral contributions, each gauge/word covariant and finite by construction (Definitions 4.339, 4.408 and 4.468 and Proposition 4.426).

Twofold reduction. In a duality-collapse regime (Definition 4.87), $D^{[\mathbf{W}_1,\mathbf{W}_2]}$ and $g^{[\mathbf{W}]}_{\mu\nu}$ reduce to their Twofold counterparts by restriction to the $\{0,1\}$ slice, via the isometries used in Corollary 4.473 and the equivalence results for $d^{[\cdot]}_{\mathrm{corr}}$ and $d^{[\cdot]}_{\mathrm{spec}}$.

The Relational Distinguishability Fiber

We package the three distances into a local, operator-informed fibre of "distinguishability" over each locus.

Definition 4.483 (Relational Distinguishability Fiber $\mathcal{F}_x$)**.** For each PQS locus $x \in \mathcal{X}_q$, define the real three–dimensional vector space

$$\mathcal{F}_x \;\cong\; \mathrm{span}_{\mathbb{R}}\{\mathbf{e}_{\mathrm{corr}},\ \mathbf{e}_{\mathrm{vac}},\ \mathbf{e}_{\mathrm{spec}}\} \;\cong\; \mathbb{R}^3,$$

whose canonical basis vectors correspond to the three distances d_{corr} (Section 4.9.1), d_{vac} (Definition 4.463), and d_{spec} (Section 4.9.9). For any pair (x, x'), the *relational displacement* from the viewpoint of x is

$$\vec{d}(x, x') \;:=\; \Big(d_{\mathrm{corr}}(x, x'),\ d_{\mathrm{vac}}(x, x'),\ d_{\mathrm{spec}}(x, x')\Big) \;\in\; \mathcal{F}_x.$$

Remark 4.484 (Orthogonality and Composite Metric)**.** The functional orthogonality of the three axes is established in Section 4.10.1. The composite (triadic) distance and its metric properties are defined in Section 4.10.2. We do not restate those results here; the fibre $\mathcal{F}_x$ merely organises them as a local, rank–3 structure over $\mathcal{X}_q$.

Remark 4.485 (Bundle Viewpoint and Calibration)**.** As x varies, the fibres assemble into a rank–3 real vector bundle $\pi : \mathcal{F} \to \mathcal{X}_q$ with canonical frame $\{\mathbf{e}_{\mathrm{corr}}, \mathbf{e}_{\mathrm{vac}}, \mathbf{e}_{\mathrm{spec}}\}$. Any choice of calibration matrix on the fibre, as used in Section 4.10.2, equips $\mathcal{F}$ with a fibrewise inner product and yields the previously defined composite distance without further assumptions.

Remark 4.486 (Twofold reduction)**.** In the duality–collapse regime (Definition 4.87), the components of $\vec{d}(x, x')$ reduce to their Twofold counterparts via the isometries of Proposition 4.328; the fibre $\mathcal{F}_x$ remains three–dimensional but its coordinates are computed on the reduced representation.

Inner Product and Metric Structure on the Distinguishability Fiber

We formalize the fibrewise geometry of Section 4.11.3 by endowing each Relational Distinguishability Fiber $\mathcal{F}_x$ with a (possibly state-/context-dependent) inner product. This provides a clean geometric interpretation of the emergent distances and matches the calibration used in Section 4.10.2.

Definition 4.487 (Fiber Inner Product and Calibration Metric)**.** Let $\vec{u}, \vec{v} \in \mathcal{F}_x$ with coordinates u^a, v^a in the canonical basis $\{\mathbf{e}_{\mathrm{corr}}, \mathbf{e}_{\mathrm{vac}}, \mathbf{e}_{\mathrm{spec}}\}$. Fix a symmetric positive-definite *calibration matrix* $\mathbf{M}(x) = \big(M_{ab}(x)\big)$ as in Section 4.10.2. Define

$$\langle \vec{u}, \vec{v} \rangle_{\mathcal{F}_x} \;:=\; \vec{u}^{\top} \mathbf{M}(x)\, \vec{v} \;=\; M_{ab}(x)\, u^a v^b, \qquad \|\vec{u}\|_{\mathcal{F}_x} \;:=\; \sqrt{\langle \vec{u}, \vec{u} \rangle_{\mathcal{F}_x}}. \tag{4.225}$$

The *canonical* choice is $\mathbf{M}(x) = \mathbf{I}_3$ (Euclidean fibre). Diagonal $\mathbf{M}(x) = \mathrm{diag}(\alpha^2, \beta^2, \gamma^2)$ implements axis rescalings; off-diagonal entries allow calibrated couplings among axes.

For the relational displacement $\vec{d}(x, x') = \big(d_{\mathrm{corr}}, d_{\mathrm{vac}}, d_{\mathrm{spec}}\big) \in \mathcal{F}_x$, the composite (triadic) distance of Section 4.10.2 is simply the fibre norm:

$$D^2(x, x') \;=\; \langle \vec{d}(x, x'), \vec{d}(x, x') \rangle_{\mathcal{F}_x} \;=\; \vec{d}(x, x')^{\top} \mathbf{M}(x)\, \vec{d}(x, x'). \tag{4.226}$$

Remark 4.488 (Geometry in the Fiber)**.** The inner product (4.225) defines angles, projections, and orthogonal decompositions in $\mathcal{F}_x$. With the canonical choice $\mathbf{M} = \mathbf{I}_3$, the axes $\{\mathbf{e}_{\text{corr}}, \mathbf{e}_{\text{vac}}, \mathbf{e}_{\text{spec}}\}$ are orthonormal, reflecting the functional orthogonality established in Section 4.10.1; the magnitude $D(x, x')$ is the norm of the total relational separation vector. Nontrivial $\mathbf{M}(x)$ encodes calibrated anisotropy or mixing among axes without re-defining the component distances.

Remark 4.489 (Metric Field on the Bundle)**.** As x varies, $x \mapsto \mathbf{M}(x)$ specifies a smooth fibre metric on the rank-3 bundle $\pi : \mathcal{F} \to \mathcal{X}_q$. When inserted into the coincidence-limit construction of Section 4.10.3, this yields the same operator-informed emergent metric, now manifestly as the Hessian of D^2 with respect to the calibrated fibre norm.

4.11.4 Distinction and Alignment of Inner Products

Two inner products play complementary roles in IQR: the *geometric* inner product on the *Relational Distinguishability Fiber* $\mathcal{F}_x$ and the *kinematic* inner product on the local *Modal Hilbert Space* $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}$. They are not equivalent, but they are structurally aligned: the geometric structure is a second–order summary built from the first–order local kinematics.

Geometric Inner Product on the Distinguishability Fiber $\mathcal{F}_x$

This inner product supplies the metric structure for emergent distinguishability.

- *Domain:* The real vector space $\mathcal{F}_x \cong \text{span}\{\mathbf{e}_{\text{corr}}, \mathbf{e}_{\text{vac}}, \mathbf{e}_{\text{spec}}\} \cong \mathbb{R}^3$, whose coordinates are the three distances $d_{\text{corr}}(x, x')$, $d_{\text{vac}}(x, x')$, $d_{\text{spec}}(x, x')$.
- *Nature:* A real Euclidean pairing, used to form the composite distance $D(x, x')$ (Section 4.10.2).
- *Interpretation:* Encodes *second–order* emergent geometry. Its entries are built from relational observables evaluated in the modal bases provided by the local kinematics.

Kinematic Inner Product on the Modal Hilbert Space $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}$

This inner product determines the local quantum structure at a single PQS.

- *Domain:* The complex Hilbert space $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}$ (Definition 4.273), comprising modal states supported at $\mathfrak{B}_x$.
- *Nature:* A complex Hermitian pairing induced from the Fourfold Inter-Modal Correlator via the coincident limit in the Quantum Holo-State (eq. (4.85)).
- *Interpretation:* Encodes *first–order* quantum kinematics (interference/superposition). It furnishes the orthonormal modal bases $\{|\psi_k\rangle\}$ used as probes for all relational observables.

Structural Alignment and Emergence

The fiber inner product on $\mathcal{F}_x$ is *assembled from* quantities computed using the kinematic inner product on $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$. In short,

$$\langle\cdot,\cdot\rangle_{\mathcal{H}_{\mathrm{mod}}} \implies \text{modal bases and correlator expectations} \implies \left(d_{\mathrm{corr}}, d_{\mathrm{vac}}, d_{\mathrm{spec}}\right) \implies \langle\cdot,\cdot\rangle_{\mathcal{F}_x}.$$

Thus the geometric pairing is a state–conditioned, second–order summary of the local kinematics, not an additional postulate.

Feature	**Inner product on $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ (kinematic)**	**Inner product on $\mathcal{F}_x$ (geometric)**
Type	Complex Hermitian	Real Euclidean
Domain	Modal states at a single PQS	Distance triples between pairs of PQSs
Role	Defines local amplitudes/interference	Defines composite distinguishability $D(x, x')$
Level	First–order quantum kinematics	Second–order emergent geometry

Table 4.3: Distinct roles of the kinematic and geometric inner products and their alignment within IQR.

Summary. The modal inner product governs the local quantum rules and fixes the probe bases; the fiber inner product aggregates the resulting relational distances into a coherent metric structure on distinguishability.

Wordwise Relational Distinguishability Fiber

We package the three *wordwise* distances into a local, operator-informed fibre of "distinguishability" over each locus.

Definition 4.490 (Wordwise Relational Distinguishability Fiber $\mathcal{F}_x^{\mathrm{w}}$)**.** For each PQS locus $x \in \mathcal{X}_q$, define the real three–dimensional vector space

$$\mathcal{F}_x^{\mathrm{w}} \;\cong\; \mathrm{span}_{\mathbb{R}}\{\mathbf{e}_{\mathrm{corr}},\ \mathbf{e}_{\mathrm{vac}},\ \mathbf{e}_{\mathrm{spec}}\} \;\cong\; \mathbb{R}^3,$$

whose canonical basis vectors correspond to the three *wordwise* distances $d_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]}$ (Definition 4.339), $d_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}$ (Definition 4.468), and $d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}$ (Definition 4.408). For any pair (x, x') and admissible endpoint words $\mathbf{W}_1, \mathbf{W}_2$, the *wordwise relational displacement* from the viewpoint of x is

$$\vec{d}^{[\mathbf{W}_1,\mathbf{W}_2]}(x, x') \;:=\; \left(d_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x, x'),\ d_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x, x'),\ d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x, x')\right) \;\in\; \mathcal{F}_x^{\mathrm{w}}.$$

When needed to compare across fibres, profiles at x' are transported to x by the wordwise transporter $\mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x \leftarrow x'; \gamma)$ (Section 4.5).

Remark 4.491 (Orthogonality and Composite Metric (wordwise))**.** Functional orthogonality of the three axes is established in Section 4.11.1. The composite (triadic) distance and its metric properties are defined in Section 4.11.2. The fibre $\mathcal{F}_x^{\mathrm{w}}$ simply organises these structures as a local, rank–3 bundle over $\mathcal{X}_q$.

Remark 4.492 (Bundle Viewpoint, Transport, and Calibration)**.** As x varies, the fibres assemble into a rank–3 real vector bundle $\pi : \mathcal{F}^{\mathrm{w}} \to \mathcal{X}_q$ with canonical frame $\{\mathbf{e}_{\mathrm{corr}}, \mathbf{e}_{\mathrm{vac}}, \mathbf{e}_{\mathrm{spec}}\}$. The wordwise covariant kinematics (Section 4.5) induce a natural parallel transport on $\mathcal{F}^{\mathrm{w}}$ consistent with the probe transport used in the distances. Any calibration matrix chosen in Section 4.11.2 equips $\mathcal{F}^{\mathrm{w}}$ with a fibrewise inner product and reproduces the composite distance.

Remark 4.493 (Twofold reduction and exogenous equivalence)**.** In the duality–collapse regime (Definition 4.87), the components of $\vec{d}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')$ reduce to their Twofold counterparts via the isometries used in Corollary 4.473 and the corresponding results for $d^{[\cdot]}_{\mathrm{corr}}$ and $d^{[\cdot]}_{\mathrm{spec}}$. Moreover, $\vec{d}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') = \vec{d}(x,x')$ componentwise by the equivalence propositions already established, so the wordwise and exogenous fibres agree numerically.

Inner Product and Metric Structure on the Wordwise Distinguishability Fiber

Definition 4.494 (Wordwise Fiber Inner Product and Calibration Metric)**.** Let $\vec{u}, \vec{v} \in \mathcal{F}_x^{\mathrm{w}}$ with coordinates u^a, v^a in the canonical basis $\{\mathbf{e}_{\mathrm{corr}}, \mathbf{e}_{\mathrm{vac}}, \mathbf{e}_{\mathrm{spec}}\}$. Fix a symmetric positive-definite *calibration matrix* $\mathbf{M}(x) = \big(M_{ab}(x)\big)$ as in Section 4.11.2. Define

$$\langle \vec{u}, \vec{v} \rangle_{\mathcal{F}_x^{\mathrm{w}}} := \vec{u}^{\top} \mathbf{M}(x)\, \vec{v} = M_{ab}(x)\, u^a v^b, \qquad \|\vec{u}\|_{\mathcal{F}_x^{\mathrm{w}}} := \sqrt{\langle \vec{u}, \vec{u} \rangle_{\mathcal{F}_x^{\mathrm{w}}}}. \tag{4.227}$$

The *canonical* choice is $\mathbf{M}(x) = \mathbf{I}_3$ (Euclidean fibre). Diagonal $\mathbf{M}(x) = \mathrm{diag}(\alpha^2, \beta^2, \gamma^2)$ implements axis rescalings; off-diagonal entries allow calibrated couplings among axes when desired.

For the wordwise relational displacement $\vec{d}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \in \mathcal{F}_x^{\mathrm{w}}$, the composite (triadic) distance of Section 4.11.2 is the fibre norm:

$$\begin{aligned} \Big(D^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\Big)^2 &= \Big\langle \vec{d}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x'),\, \vec{d}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \Big\rangle_{\mathcal{F}_x^{\mathrm{w}}} \\ &= \vec{d}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')^{\top}\, \mathbf{M}(x)\, \vec{d}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x'). \end{aligned} \tag{4.228}$$

Remark 4.495 (Geometry in the Wordwise Fiber)**.** The inner product (4.227) defines angles, projections, and orthogonal decompositions in $\mathcal{F}_x^{\mathrm{w}}$. With $\mathbf{M} = \mathbf{I}_3$, the axes $\{\mathbf{e}_{\mathrm{corr}}, \mathbf{e}_{\mathrm{vac}}, \mathbf{e}_{\mathrm{spec}}\}$ are orthonormal, reflecting functional orthogonality (Section 4.11.1); the magnitude $D^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')$ is the norm of the total relational separation vector. Nontrivial $\mathbf{M}(x)$ encodes calibrated anisotropy or mixing among axes without changing the component distances themselves.

Remark 4.496 (Metric Field on the Wordwise Bundle)**.** As x varies, $x \mapsto \mathbf{M}(x)$ specifies a smooth fibre metric on the rank-3 bundle $\pi : \mathcal{F}^{\mathrm{w}} \to \mathcal{X}_q$. Inserted into the coincidence-limit construction of Section 4.11.3, this yields the same operator-informed emergent metric, now manifestly as the Hessian of $\Big(D^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\Big)^2$ with respect to the calibrated fibre norm.

4.11.5 The Triadic Metric Structure (Fréchet) and the PQS Locus

The three operator–theoretic distances introduced in Section 4.9—the correlator distance d_{corr}, the vacuum–response distance d_{vac}, and the spectral distance d_{spec}—organise the emergent metric structure on the quantum configuration space $\mathcal{X}_q$ in the spirit of relational quantum mechanics [3] and noncommutative geometry [4]. Their functional *orthogonality* and the construction of the composite metric are established in Sections 4.10.1 and 4.10.2. In this section we complement those results by proving a stronger *functional independence* statement for the underlying generators via a Fréchet–differential argument.

Functional Independence (Fréchet) of the Triadic Generators

Let $\mathcal{O} := \mathrm{End}(\mathcal{H}_{\mathrm{total}})$ be the C^*–algebra of bounded operators on the total emergent Hilbert space. Denote by $\hat{G}_\Lambda$ the Fourfold Inter–Modal Correlator (cf. Definition 4.9) and by $\hat{D}$ an admissible Spectral Operator (cf. Definition 4.343). We employ the following shorthand for the three operator–level generators attached to a pair of loci $(x, x') \in \mathcal{X}_q \times \mathcal{X}_q$:

$$\mathbf{A}_{\mathrm{corr}}(x,x') := \hat{G}_\Lambda(x,x') \qquad \text{(bilocal/off–diagonal coherence)}, \tag{4.229}$$

$$\mathbf{A}_{\mathrm{vac}}(x,x') := \hat{G}_\Lambda(x,x) - \hat{G}_\Lambda(x',x') \qquad \text{(diagonal/coincident contrast)}, \tag{4.230}$$

$$\mathbf{A}_{\mathrm{spec}}(x,x') := i\,[\hat{D}, \hat{G}_\Lambda(x,x')] \qquad \text{(spectral shear/commutator)}. \tag{4.231}$$

Proposition 4.497 (Fréchet functional independence of triadic generators)**.** *Let $f \in C^\infty(\mathcal{O}^{\otimes 3}; \mathcal{O})$ be a smooth operator functional.*[2] *If*

$$f\Big(\mathbf{A}_{\mathrm{corr}}(x,x'),\ \mathbf{A}_{\mathrm{vac}}(x,x'),\ \mathbf{A}_{\mathrm{spec}}(x,x')\Big) = 0 \quad \textit{for all } (x,x') \in \mathcal{X}_q \times \mathcal{X}_q,$$

then $f \equiv 0$. Equivalently, the three maps $(x,x') \mapsto \mathbf{A}_{\mathrm{corr}}, \mathbf{A}_{\mathrm{vac}}, \mathbf{A}_{\mathrm{spec}}$ are functionally independent.

Proof (sketch). **(i) Algebraic Support Separation.** $\mathbf{A}_{\mathrm{corr}}$ lives on the bilocal (off–diagonal) support of $\hat{G}_\Lambda$ on $\mathcal{X}_q \times \mathcal{X}_q$; $\mathbf{A}_{\mathrm{vac}}$ is a difference of diagonal restrictions; $\mathbf{A}_{\mathrm{spec}}$ is an *inner derivation* of the correlator generated by $\hat{D}$. No algebraic combination of diagonal restrictions reproduces a genuinely bilocal kernel, and no combination that omits $\hat{D}$ generates a nontrivial inner derivation.

(ii) GNS/von Neumann Separation. In the GNS representation of the Quantum Holo-State $\omega(\cdot) = \langle \overset{\circ}{\mathbf{\Psi}} | \cdot | \overset{\circ}{\mathbf{\Psi}} \rangle$ (cf. Definition 1.34), the cyclic vectors obtained by acting with the three families on $|\Omega_{\mathrm{GNS}}\rangle$ generate mutually orthogonal subspaces whenever mixed expectations vanish, i.e. $\omega(A^\dagger_{\mathrm{corr}} A_{\mathrm{vac}}) = \omega(A^\dagger_{\mathrm{corr}} A_{\mathrm{spec}}) = \omega(A^\dagger_{\mathrm{vac}} A_{\mathrm{spec}}) = 0$ (see [404, 230]). Hence a nontrivial functional dependence would force unwanted inclusions among the generated von Neumann algebras, contradicting the postulated independence of $\hat{D}$ and $\hat{G}_\Lambda$.

(iii) Fréchet–Differential Test. View $\mathcal{O}$ as a Banach space and consider arbitrary small variations $(\delta\hat{G}_\Lambda, \delta\hat{D})$. The first variations are

$$\delta\mathbf{A}_{\mathrm{corr}} = \delta\hat{G}_\Lambda(x,x'), \qquad \delta\mathbf{A}_{\mathrm{vac}} = \delta\hat{G}_\Lambda(x,x) - \delta\hat{G}_\Lambda(x',x'), \tag{4.232}$$

$$\delta\mathbf{A}_{\mathrm{spec}} = i\,[\delta\hat{D}, \hat{G}_\Lambda(x,x')] + i\,[\hat{D}, \delta\hat{G}_\Lambda(x,x')]. \tag{4.233}$$

[2] Smoothness is with respect to the Banach–space topology on $\mathcal{B}(\mathcal{H})$; see, e.g., [253, 241].

If a nontrivial identity $f(\mathbf{A}_{\text{corr}}, \mathbf{A}_{\text{vac}}, \mathbf{A}_{\text{spec}}) = 0$ held identically, its Fréchet derivative would annihilate all allowed variations. Choosing variations with $\delta\hat{G}_\Lambda = 0$ but arbitrary $\delta\hat{D}$ forces the coefficient of $\delta\mathbf{A}_{\text{spec}}$ to vanish; then choosing purely off–diagonal $\delta\hat{G}_\Lambda$ kills any dependence on $\mathbf{A}_{\text{corr}}$; finally, purely diagonal $\delta\hat{G}_\Lambda$ forces the $\mathbf{A}_{\text{vac}}$ coefficient to vanish. Thus all first–order coefficients are zero and, by smoothness, $f \equiv 0$. □

Remark 4.498 (Relation to Established Orthogonality)**.** Proposition 4.497 strengthens the vanishing of cross–expectations proved in Section 4.10.1: it shows that (i) no nontrivial smooth functional relation among the three generator families exists, and (ii) their tangent directions in the Banach space of operator fields are linearly independent. The composite metric of Section 4.10.2 therefore sits on a basis that is both orthogonal (expectation-level) and functionally independent (operator-level).

4.11.6 The PQS as the Locus of Indistinguishability

We now identify the physical meaning of the *zero* of the composite (triadic) distance $D(x, x')$ (Section 4.10.2) in relation to the foundational notion of a Primordial Quantum Singularity (PQS; Definition 1.25).

Proposition 4.499 (PQS as the Zero Locus of Distinguishability)**.** *For loci $x, x' \in \mathcal{X}_q$,*

$$D(x, x') = 0 \quad \Longleftrightarrow \quad \vec{d}(x, x') = (0, 0, 0),$$

i.e. all three relational distances vanish simultaneously: $d_{\text{corr}}(x, x') = d_{\text{vac}}(x, x') = d_{\text{spec}}(x, x') = 0$. In this case x and x' are quantum–indistinguishable *by the relational probes of IQR, and their coincidence realises the defining condition of a PQS.*

Proof. By the Principle of Coherent Regularization (Proposition 4.424), each component distance is well-defined and finite: $d_{\text{corr}}(x, x') \in [0, 1]$ (by construction), $d_{\text{vac}}(x, x') < \infty$ (modal–weight decay), and $d_{\text{spec}}(x, x') < \infty$ (Spectral Smoothness + Finite Entanglement Capacity).

With finiteness established, Sections 4.10.2 and 4.11.3 yield

$$D^2(x, x') \;=\; \vec{d}(x, x')^{\mathsf{T}}\, \mathbf{M}(x)\, \vec{d}(x, x'), \qquad \mathbf{M}(x) \succ 0.$$

Hence

$$D(x, x') = 0 \;\Longleftrightarrow\; \vec{d}(x, x') \in \ker \mathbf{M}(x).$$

Positive definiteness implies $\ker \mathbf{M}(x) = \{0\}$; therefore $\vec{d}(x, x') = \mathbf{0}$, i.e. each component distance vanishes. The converse is immediate.

Operationally, vanishing of all three components means that no admissible relational probe detects a difference between x and x': no bilocal coherence (off–diagonal), no diagonal vacuum response, and no spectral shear. This is precisely the IQR notion of *complete quantum unresolvability*, matching the PQS condition of Definition 1.25. □

Remark 4.500 (Role of Finite Entanglement Capacity)**.** The Finite Entanglement Capacity does not alter the logical equivalence $D = 0 \Longleftrightarrow \vec{d} = \mathbf{0}$, but it is essential to the *well-posedness*

and stability of that statement: it guarantees that all series entering d_{vac} and d_{spec} converge and that neighborhoods of the zero locus are controlled by uniform modal–weight bounds. Thus "$D = 0$" reflects genuine indistinguishability, not a formal cancellation in a divergent expression.

Remark 4.501 (Not a Background Coincidence)**.** $D(x, x') = 0$ does *not* assert equality of classical coordinates in a pre-given manifold. It states that, relative to the Holo-State and the operators of Section 4.9, all admissible relational tests fail to separate x from x'. The PQS is therefore the *null–distance locus* of the emergent operator geometry, not a prior geometric point.

Corollary 4.502 (Zero–Distance Identification)**.** *The relation $x \sim x'$ iff $D(x, x') = 0$ is an equivalence relation on $\mathcal{X}_q$. The metric D descends to the quotient $\mathcal{X}_q/\!\sim$, on which it is strictly definite. In particular, PQS loci correspond to equivalence classes of mutually indistinguishable points.*

Remark 4.503 (Twofold Reduction and Classical Limit)**.** In a duality–collapse regime (Definition 4.87), the condition $\vec{d}(x, x') = (0, 0, 0)$ reduces—via the isometries of Proposition 4.328—to vanishing of the corresponding twofold (qubit–level) distances. In further decohered classical limits, this reproduces ordinary point–coincidence.

4.11.7 Wordwise Triadic Metric Structure (Fréchet) and the PQS Locus

The three *wordwise* operator–theoretic distances introduced in Definitions 4.339, 4.408 and 4.468— the correlator distance $d_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]}$, the vacuum–response distance $d_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}$, and the spectral distance $d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}$—organise the emergent metric structure on the quantum configuration space $\mathcal{X}_q$ in the holosymmetric wordwise calculus. Their functional *orthogonality* and the composite metric are established in Sections 4.11.1 and 4.11.2. We now complement those results by proving a stronger *functional independence* statement for the underlying *wordwise* generators via a Fréchet–differential argument.

Functional Independence (Fréchet) of the Wordwise Triadic Generators

Let $\mathcal{O} := \mathcal{B}(\mathcal{H}_{\mathrm{total}})$ be the C^*–algebra of bounded operators on the total emergent Hilbert space. Denote by $\hat{G}_\Lambda$ the Fourfold inter–modal correlator and by $\mathbb{D}^{[\mathbf{W}]}$ an admissible wordwise spectral operator (Section 4.9.4). Fix endpoints $x, x' \in \mathcal{X}_q$, admissible words $\mathbf{W}_1$ at x, $\mathbf{W}_2$ at x', and a window $\gamma : x' \to x$ with transporter $\mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x \leftarrow x'; \gamma)$ (Section 4.5). We employ the following shorthand for three *wordwise* operator–level generators attached to (x, x'):

$$\mathbf{A}_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x, x'; \gamma) := \mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x \leftarrow x'; \gamma)^\dagger \, \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'}) \, \mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x \leftarrow x'; \gamma), \tag{4.234}$$
(bilocal/off–diagonal coherence)

$$\mathbf{A}_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x, x') := \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_x) - \hat{G}_\Lambda(\mathfrak{B}_{x'}, \mathfrak{B}_{x'}), \tag{4.235}$$
(diagonal/coincident contrast)

$$\mathbf{A}_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x, x'; \gamma) := \Big[\mathbb{D}^{[\mathbf{W}_1]}(x), \, \mathbf{A}_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x, x'; \gamma)\Big], \tag{4.236}$$
(wordwise spectral shear/commutator; anti-Hermitian)

(Equivalently, the last line may be written with the transported $\mathbb{D}^{[\mathbf{W}_2]}(x')$ or with the spectral difference $\Delta\mathbb{D}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')$, up to conjugation; cf. Definition 4.406.)

Definition 4.504 (Connected bilocal correlator (holosymmetric, wordwise))**.** Fix $x, x' \in \mathcal{X}_q$ and an admissible word pair $(\mathbf{W}_1, \mathbf{W}_2)$. The connected wordwise bilocal Fourfold correlator is the expectation-level quantity

$$\Gamma^{[\mathbf{W}_1,\mathbf{W}_2]}_{\text{conn}}(x,x') := \left\langle \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'}) \right\rangle_\Psi - \left\langle \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_x) \right\rangle_\Psi \left\langle \hat{G}_\Lambda(\mathfrak{B}_{x'}, \mathfrak{B}_{x'}) \right\rangle_\Psi.$$

Equivalently, on test vectors ψ, ϕ in the wordwise fibres,

$$\begin{aligned}\Gamma^{[\mathbf{W}_1,\mathbf{W}_2]}_{\text{conn}}(x,x')[\psi^*,\phi] := {} & \left\langle \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi^*,\phi] \right\rangle_\Psi \\ & - \left\langle \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_x)[\psi^*,\psi] \right\rangle_\Psi \left\langle \hat{G}_\Lambda(\mathfrak{B}_{x'}, \mathfrak{B}_{x'})[\phi^*,\phi] \right\rangle_\Psi.\end{aligned}$$

This object is invariant under admissible endpoint word/gauge changes and vanishes on the PQS class.

Proposition 4.505 (Fréchet functional independence of wordwise triadic generators)**.** *Let $f \in C^\infty(\mathcal{O}^{\otimes 3}; \mathcal{O})$ be a smooth operator functional.*[3] *If*

$$\begin{aligned}& f\Big(\mathbf{A}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\text{corr}}(x,x';\gamma),\ \mathbf{A}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\text{vac}}(x,x'),\ \mathbf{A}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\text{spec}}(x,x';\gamma)\Big) = 0, \\ & \forall\,(x,x',\gamma) \text{ and admissible } (\mathbf{W}_1,\mathbf{W}_2).\end{aligned}$$

then $f \equiv 0$. Equivalently, the maps $(x,x',\gamma) \mapsto \mathbf{A}^{[\cdot]}_{\text{corr}},\ \mathbf{A}^{[\cdot]}_{\text{vac}},\ \mathbf{A}^{[\cdot]}_{\text{spec}}$ are functionally independent.

Proof (sketch). **(i) Wordwise algebraic support separation.** $\mathbf{A}^{[\cdot]}_{\text{corr}}$ lives on the bilocal (off–diagonal) support of $\hat{G}_\Lambda$ on $\mathcal{X}_q \times \mathcal{X}_q$, transported to x; $\mathbf{A}^{[\cdot]}_{\text{vac}}$ is a difference of diagonal restrictions; $\mathbf{A}^{[\cdot]}_{\text{spec}}$ is an *inner derivation* of $\mathbf{A}^{[\cdot]}_{\text{corr}}$ generated by $\mathbb{D}^{[\mathbf{W}_1]}(x)$. No algebraic combination of diagonal restrictions reproduces a genuinely bilocal kernel, and no combination that omits $\mathbb{D}^{[\mathbf{W}_1]}$ generates a nontrivial inner derivation. Unitary word/gauge changes act by conjugation and do not alter this separation.

(ii) Wordwise GNS/von Neumann separation. In the GNS representation of $\omega(\cdot) = \langle \mathring{\mathbf{\Psi}}| \cdot |\mathring{\mathbf{\Psi}}\rangle$, the cyclic vectors obtained by acting with the three wordwise families on $|\Omega_{\text{GNS}}\rangle$ generate mutually orthogonal subspaces whenever mixed expectations vanish (functional orthogonality; cf. Section 4.11.1). A nontrivial functional dependence would force unwanted inclusions among the von Neumann algebras they generate, contradicting the independent roles of $\hat{G}_\Lambda$ and $\mathbb{D}^{[\mathbf{W}]}$ in the wordwise calculus.

(iii) Fréchet–differential test (wordwise). View $\mathcal{O}$ as a Banach space and consider arbitrary small variations $(\delta\hat{G}_\Lambda, \delta\mathbb{D}^{[\mathbf{W}_1]})$, keeping the window γ fixed. The first variations are

$$\delta\mathbf{A}^{[\cdot]}_{\text{corr}} = \mathcal{U}^\dagger\, \delta\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})\, \mathcal{U}, \tag{4.237}$$

$$\delta\mathbf{A}^{[\cdot]}_{\text{vac}} = \delta\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_x) - \delta\hat{G}_\Lambda(\mathfrak{B}_{x'}, \mathfrak{B}_{x'}), \tag{4.238}$$

$$\delta\mathbf{A}^{[\cdot]}_{\text{spec}} = \Big[\delta\mathbb{D}^{[\mathbf{W}_1]}(x),\ \mathbf{A}^{[\cdot]}_{\text{corr}}\Big] + \Big[\mathbb{D}^{[\mathbf{W}_1]}(x),\ \delta\mathbf{A}^{[\cdot]}_{\text{corr}}\Big], \tag{4.239}$$

[3] "Smooth" is in the Fréchet/Fréchet–Gateaux sense on Banach spaces (here the operator norm topology on $\mathcal{B}(\mathcal{H})$).

where $\mathcal{U} = \mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x \leftarrow x';\gamma)$. If a nontrivial identity $f(\mathbf{A}^{[\cdot]}_{\mathrm{corr}}, \mathbf{A}^{[\cdot]}_{\mathrm{vac}}, \mathbf{A}^{[\cdot]}_{\mathrm{spec}}) = 0$ held identically, its Fréchet derivative would annihilate all allowed variations. Choosing $\delta\hat{G}_\Lambda = 0$ but arbitrary $\delta\mathbb{D}^{[\mathbf{W}_1]}$ forces the coefficient of $\delta\mathbf{A}^{[\cdot]}_{\mathrm{spec}}$ to vanish; then choosing purely off–diagonal $\delta\hat{G}_\Lambda$ (so that $\delta\mathbf{A}^{[\cdot]}_{\mathrm{vac}} = 0$) kills any dependence on $\mathbf{A}^{[\cdot]}_{\mathrm{corr}}$; finally, purely diagonal $\delta\hat{G}_\Lambda$ forces the $\mathbf{A}^{[\cdot]}_{\mathrm{vac}}$ coefficient to vanish. Thus all first–order coefficients are zero and, by smoothness, $f \equiv 0$. □

Remark 4.506 (Relation to orthogonality and exogenous equivalence (wordwise))**.** Proposition 4.505 strengthens the vanishing of cross–expectations proved in Section 4.11.1: it shows that (i) no nontrivial smooth functional relation among the three *wordwise* generator families exists, and (ii) their tangent directions in the Banach space of operator fields are linearly independent. By the equivalence results (word/gauge invariance), the same independence holds for the exogenous generators, so the composite metric of Section 4.11.2 sits on a basis that is both orthogonal (expectation-level) and functionally independent (operator-level) in either presentation.

4.11.8 Wordwise PQS as the Locus of Indistinguishability

We identify the physical meaning of the *zero* of the composite (wordwise triadic) distance $D^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')$ (Section 4.11.2) in relation to the Primordial Quantum Singularity (PQS; Definition 1.25).

Proposition 4.507 (PQS as the zero locus of wordwise distinguishability)**.** *For loci $x, x' \in \mathcal{X}_q$ and admissible endpoint words $\mathbf{W}_1, \mathbf{W}_2$,*

$$D^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') = 0 \quad \Longleftrightarrow \quad \vec{d}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') = (0,0,0),$$

i.e. all three wordwise relational distances vanish simultaneously: $d^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{corr}}(x,x') = d^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{vac}}(x,x') = d^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{spec}}(x,x') = 0$. In this case x and x' are quantum–indistinguishable *by the wordwise relational probes, and their coincidence realises the defining condition of a PQS.*

Proof. Finiteness of components. By construction $d^{[\cdot]}_{\mathrm{corr}} \in [0,1]$ (Definition 4.339). For the vacuum–response, Lemma 4.470 gives the uniform bound $\left(d^{[\cdot]}_{\mathrm{vac}}(x,x')\right)^2 \le 4\|\hat{G}_\Lambda\|^2$, so $d^{[\cdot]}_{\mathrm{vac}} < \infty$ without additional hypotheses. For the spectral distance, finiteness follows from wordwise spectral smoothness together with endogenous capacity/UV control (Theorem 4.421, Principle 4.2, and Proposition 4.422).

Equivalence with the zero vector. With finiteness ensured, Sections 4.11.2 and 4.11.4 yield

$$\left(D^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\right)^2 = \vec{d}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')^\top\, \mathbf{M}(x)\, \vec{d}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x'), \qquad \mathbf{M}(x) \succ 0.$$

Hence $D^{[\cdot]}(x,x') = 0 \iff \vec{d}^{[\cdot]}(x,x') \in \ker \mathbf{M}(x)$; positive–definiteness implies $\ker \mathbf{M}(x) = \{0\}$, so $\vec{d}^{[\cdot]}(x,x') = \mathbf{0}$. The converse is immediate.

Operational meaning. Vanishing of all three components states that no admissible wordwise probe detects a difference between x and x': no bilocal coherence (off–diagonal), no diagonal vacuum response, and no spectral shear. This matches the IQR definition of *complete quantum unresolvability*—the PQS condition (Definition 1.25). □

Remark 4.508 (Role of Finite Entanglement Capacity (wordwise))**.** Finite Entanglement Capacity is *not* needed for the logical equivalence $D^{[\cdot]} = 0 \Leftrightarrow \vec{d}^{[\cdot]} = \mathbf{0}$ (which uses only $\mathbf{M} \succ 0$), but it strengthens *well-posedness and stability*: it yields convergent series for spectral contributions and uniform modal–weight bounds in neighborhoods of the zero locus (Theorem 4.421, Principle 4.2, and Proposition 4.422).

Remark 4.509 (Not a background coincidence (wordwise))**.** $D^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') = 0$ does *not* assert equality of classical coordinates. It states that, relative to the Holo-State and operators of the wordwise calculus, all admissible relational tests fail to separate x from x'. The PQS is therefore the *null–distance locus* of the emergent operator geometry, not a prior geometric point.

Corollary 4.510 (Zero–distance identification (wordwise))**.** *The relation $x \sim x'$ iff $D^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') = 0$ is an equivalence relation on $\mathcal{X}_q$, independent of the choice of admissible words and (on flat patches) of the window (Remark 4.482). The metric $D^{[\mathbf{W}_1,\mathbf{W}_2]}$ descends to the quotient $\mathcal{X}_q/\!\sim$, on which it is strictly definite. PQS loci correspond to equivalence classes of mutually indistinguishable points.*

Remark 4.511 (Calibration independence of the zero locus)**.** Because $\mathbf{M}(x) \succ 0$, the condition $D^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') = 0$ is equivalent to $\vec{d}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') = \mathbf{0}$ for *any* calibration $\mathbf{M}(x)$ (Section 4.11.4). Thus the PQS zero locus is independent of axis rescalings or calibrated couplings among the triadic components.

Remark 4.512 (Twofold reduction and exogenous equivalence)**.** In a duality–collapse regime (Definition 4.87), the condition $\vec{d}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') = (0,0,0)$ reduces—via the isometries used in Corollary 4.473 and the equivalence results for $d^{[\cdot]}_{\mathrm{corr}}$ and $d^{[\cdot]}_{\mathrm{spec}}$—to vanishing of the corresponding Twofold distances, reproducing ordinary point–coincidence in decohered classical limits. Moreover, $D^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') = D(x,x')$ by componentwise equivalence, so the exogenous and wordwise zero loci agree.

4.11.9 Holonomy and the Zero Locus of Distinguishability

We relate the *zero–distance condition* $D(x,x') = 0$ (Section 4.11.6) to the holonomy of the gauge–covariant transport $\mathcal{U}(x \leftarrow x';\gamma)$ used in the Fourfold correlator (Definition 4.9; cf. the parallel–transport dependence used in the Fourfold correlator (Definition 4.9).

Definition 4.513 (State–conditioned holonomy action on probes)**.** Let Hol_x be the holonomy group at a locus x (loops γ based at x mapped to $\mathcal{U}(x \leftarrow x;\gamma)$). Its *state–conditioned action* on prismatic probes is the conjugation $\hat{\mathcal{F}}_\alpha(x) \mapsto \mathcal{U}(x \leftarrow x;\gamma)\,\hat{\mathcal{F}}_\alpha(x)\,\mathcal{U}(x \leftarrow x;\gamma)^\dagger$, evaluated in the Holo-State $|\mathring{\mathbf{\Psi}}\rangle$. It is *holonomically trivial (in expectation)* if for every closed γ,

$$\Big\langle \hat{\mathcal{F}}^\dagger_\alpha(x)\,\mathcal{U}(x \leftarrow x;\gamma)\,\hat{\mathcal{F}}_\beta(x)\,\mathcal{U}(x \leftarrow x;\gamma)^\dagger \Big\rangle_\Psi = \big\langle \hat{\mathcal{F}}^\dagger_\alpha(x)\,\hat{\mathcal{F}}_\beta(x) \big\rangle_\Psi,$$

and similarly for all bilocal matrix elements entering $\hat{G}_\Lambda$ (Definition 4.9), with the same modal–weighting w_n (Definition 4.392).

Proposition 4.514 (Zero Locus Implies State–Conditioned Flatness)**.** *If $D(x,x')=0$, then for any path $\gamma : x \to x'$ and any closed loop $\gamma\circ\bar{\gamma}$ based at x (where $\bar{\gamma}$ is the reverse), the induced holonomy is trivial in the sense of Definition 4.513 on the Holo-State–supported probe sector. Equivalently,*

$$D(x,x')=0 \quad\Longrightarrow\quad \forall \text{ loops } \ell \text{ based at } x: \ \Big\| [\mathcal{U}(x \leftarrow x;\ell),\, \hat{G}_\Lambda(\cdot)]\,|\mathring{\mathbf{\Psi}}\rangle \Big\| = 0,$$

and, in particular, the spectral shear observables vanish along γ (Definition 4.386).

Sketch. $D(x,x')=0$ implies $\vec{d}(x,x')=\mathbf{0}$ (Proposition 4.499); thus each component distance vanishes: (i) $d_{\mathrm{corr}}=0$ forces equality of all bilocal coherence amplitudes along γ, which is stable under insertion of $\mathcal{U}$ for closed retracings; (ii) $d_{\mathrm{vac}}=0$ forces equality of diagonal (coincident) self–coherences at x and x', hence invariance under holonomy at x; (iii) $d_{\mathrm{spec}}=0$ forces $[\hat{D},\hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})[\psi_n^*,\psi_n]]$ to have zero Holo-State expectation for all modal modes, which is preserved under conjugation by $\mathcal{U}$ on closed loops. Together these yield the stated triviality in expectation on the Holo-State–supported subspace. □

Proposition 4.515 (State–conditioned Flatness Implies Local Zero–distance)**.** *Let $U \subset \mathcal{X}_q$ be a simply–connected neighbourhood of x such that:*

*(a) (*Holonomy triviality in expectation*) Every loop $\ell \subset U$ is trivial in the sense of Definition 4.513.*

*(b) (*Spectral quiescence*) $[\hat{D},\hat{G}_\Lambda(\mathfrak{B}_y,\mathfrak{B}_{y'})[\psi_n^*,\psi_n]]$ has vanishing Holo-State expectation for all $y,y' \in U$ and all relevant modes n.*

Then $D(y,y')=0$ for all $y,y' \in U$.

Sketch. (a) removes all holonomy–induced path dependence from the correlator sector—bilocal and diagonal— so $d_{\mathrm{corr}}(y,y')=0$ and $d_{\mathrm{vac}}(y,y')=0$ throughout U. (b) sets all spectral shear observables to zero (Definition 4.402), hence $d_{\mathrm{spec}}=0$. Therefore $\vec{d}(y,y')=\mathbf{0}$ for all $y,y' \in U$, so $D(y,y')=0$. □

Remark 4.516 (Representation–dependence and "Holonomic Invisibility")**.** Nontrivial holonomy may nevertheless act *invisibly* on the distances if its action lies in a sector annihilated by the Holo-State weights w_n (Definition 4.392) or by the expectation map (e.g. phases that cancel in $\langle\cdot\rangle_\Psi$). Thus, the relevant notion is holonomy *in the state–conditioned representation* carried by the probe algebra; triviality there suffices for $D=0$ even if the underlying connection has nonzero curvature.

Remark 4.517 (Curvature, Holonomy, and the Triadic Metric)**.** In standard differential geometry, holonomy is generated by curvature (Ambrose–Singer theorem, cf. [405, 248]). In IQR the operator–valued curvature (two–form of the transport underlying $\mathcal{U}$) contributes to the triadic metric via: (i) path–dependence of bilocal coherences (d_{corr}), (ii) shifts in coincident self–coherences (d_{vac}), and (iii) nonvanishing commutator tension with $\hat{D}$ (d_{spec}). Vanishing of these effects in the state–conditioned representation is the precise content of Propositions 4.514 and 4.515.

When the composite distance vanishes, $D(x, x') = 0$, the three relational "detectors" of geometry—bilocal coherence, diagonal vacuum contrast, and spectral shear—are all silent. Any closed loop based at x returns prismatic probes exactly as they were, so holonomy is trivial in the state-conditioned representation: no path-dependent twist appears in expectations, no vacuum profile shifts, and no commutator tension is sensed. In this regime the connection is effectively flat for the supported probe sector, and the pair (x, x') sits at a locus of complete quantum indistinguishability—the PQS class.

Conversely, nontrivial holonomy would register as a nonzero entry in at least one component of the triad. The picture parallels the classical link between curvature and holonomy: curvature that is operationally present produces detectable holonomy, which in turn lifts the triadic distances from zero. If, within a simply connected neighbourhood, holonomy is trivial (in expectation) and the spectral sector is quiescent, the triadic distances vanish throughout that neighbourhood. Intuitively, the Holo-State's entanglement network fixes the resolution of reality; at the zero locus that network cannot resolve loops or separations, so the emergent operator geometry identifies the region with a PQS equivalence class.

Operator Quantity	**Spatial Observable (d_i)**	**Temporal Observable (t_i)**
$\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})$ (bilocal, off–diagonal)	d_{corr}: magnitude; real–part aggregate of the bilocal channel	t_{corr}: phase; argument of the bilocal expectation
$\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_x)$ and $\hat{G}_\Lambda(\mathfrak{B}_{x'}, \mathfrak{B}_{x'})$ (coincident, diagonal)	d_{vac}: real self–variation (Definition 4.463 and eq. (4.211))	t_{vac}: imaginary self–variation (Definition 4.535 and eq. (4.246))
$[\hat{D}, \hat{G}_\Lambda]$ (spectral commutator)	d_{spec}: real commutator / spectral shear distance (Definition 4.402 and eq. (4.160))	t_{spec}: imaginary commutator / spectral shear time (Definition 4.543 and eq. (4.252))

Table 4.4: Duality of emergent spatial and temporal observables from fundamental operator constructs. Entries indicate which *operator feature* is aggregated and whether the aggregation takes the real (spatial) or imaginary (temporal) part. See Definitions 4.402, 4.463, 4.535 and 4.543 and eqs. (4.160), (4.211), (4.246) and (4.252).

4.11.10 Holonomy and the Zero Locus of Wordwise Distinguishability

We relate the *zero–distance condition* $D^{[\mathbf{W}_1,\mathbf{W}_2]}(x, x') = 0$ (Section 4.11.8) to the holonomy of the *wordwise* gauge–covariant transport $\mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x \leftarrow x'; \gamma)$ used throughout the holosymmetric calculus (Section 4.5).

Definition 4.518 (State–conditioned wordwise holonomy action on probes)**.** Let $\mathrm{Hol}_x^{[\mathbf{W}_x]}$ be the holonomy group of the wordwise modal connection at x, i.e. the set of loop transports

$$\mathrm{Hol}_x^{[\mathbf{W}_x]} := \Big\{ \mathcal{U}^{[\mathbf{W}_x,\mathbf{W}_x]}\big(x \leftarrow x; \gamma\big) \ : \ \gamma \text{ a loop based at } x \Big\}.$$

Its state–conditioned action on admissible probes $\hat{\mathcal{F}}_\alpha^{[\mathbf{W}_x]}(x)$ is the conjugation

$$\hat{\mathcal{F}}_\alpha^{[\mathbf{W}_x]}(x) \;\mapsto\; \mathcal{U}^{[\mathbf{W}_x,\mathbf{W}_x]}\big(x\leftarrow x;\gamma\big)\,\hat{\mathcal{F}}_\alpha^{[\mathbf{W}_x]}(x)\,\mathcal{U}^{[\mathbf{W}_x,\mathbf{W}_x]}\big(x\leftarrow x;\gamma\big)^\dagger.$$

We say holonomy is *trivial in expectation* on the supported sector if for every closed γ,

$$\Big\langle \hat{\mathcal{F}}_\alpha^{[\mathbf{W}_x]\dagger}(x)\,\mathcal{U}^{[\mathbf{W}_x,\mathbf{W}_x]}\big(x\leftarrow x;\gamma\big)\,\hat{\mathcal{F}}_\beta^{[\mathbf{W}_x]}(x)\,\mathcal{U}^{[\mathbf{W}_x,\mathbf{W}_x]}\big(x\leftarrow x;\gamma\big)^\dagger\Big\rangle_\Psi = \big\langle \hat{\mathcal{F}}_\alpha^{[\mathbf{W}_x]\dagger}(x)\,\hat{\mathcal{F}}_\beta^{[\mathbf{W}_x]}(x)\big\rangle_\Psi,$$

and likewise for bilocal matrix elements of the *wordwise* Fourfold correlator $\hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})$ that enter Definition 4.339, with the same modal weights $w_n^{[\mathbf{W}_1,\mathbf{W}_2]}$ (Definition 4.407).

Proposition 4.519 (Zero locus implies state–conditioned wordwise flatness)**.** *If*

$$D^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') = 0$$

then for any two paths $\gamma_1,\gamma_2 : x' \to x$ the closed loop $\ell = \gamma_2 \circ \bar{\gamma}_1$ based at x has trivial state–conditioned holonomy in the sense of Definition 4.518. Equivalently,

$$\Big\| \Big[\mathcal{U}^{[\mathbf{W}_x,\mathbf{W}_x]}\big(x\leftarrow x;\ell\big),\ \hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})[\psi^*,\phi]\Big]\,|\mathring{\mathbf{\Psi}}\rangle\Big\| = 0,$$

and, in particular, all wordwise spectral–shear observables along either open segment vanish: $\operatorname{Im}\,\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma_j) = 0$ *for $j=1,2$ (Definitions 4.406 and 4.408).*

Proof. **Notation.** Write $\mathbf{W}_x$ and $\mathbf{W}_{x'}$ for the word frames at the endpoints x and x'. Transports carry explicit endpoint labels as $\mathcal{U}^{[\mathbf{W}_x,\mathbf{W}_{x'}]}(x\leftarrow x';\gamma)$. For a loop ℓ based at x, we use $\mathcal{U}^{[\mathbf{W}_x,\mathbf{W}_x]}(x\leftarrow x;\ell)$.

Step 0: Transporter algebra (wordwise). The transporter obeys (unitarity, inversion, concatenation):

$$\mathcal{U}^{[\mathbf{W}_x,\mathbf{W}_{x'}]}(x\leftarrow x';\gamma)^\dagger = \mathcal{U}^{[\mathbf{W}_{x'},\mathbf{W}_x]}(x'\leftarrow x;\bar{\gamma}), \tag{4.240}$$

$$\mathcal{U}^{[\mathbf{W}_x,\mathbf{W}_x]}(x\leftarrow x;\gamma\circ\bar{\gamma}) = \mathbf{1}, \tag{4.241}$$

$$\mathcal{U}^{[\mathbf{W}_x,\mathbf{W}_z]}(x\leftarrow z;\gamma_2\circ\gamma_1) = \mathcal{U}^{[\mathbf{W}_x,\mathbf{W}_y]}(x\leftarrow y;\gamma_2)\,\mathcal{U}^{[\mathbf{W}_y,\mathbf{W}_z]}(y\leftarrow z;\gamma_1). \tag{4.242}$$

Hence, for the loop $\ell = \gamma\circ\bar{\gamma}$ based at x, $\mathcal{U}^{[\mathbf{W}_x,\mathbf{W}_x]}(x\leftarrow x;\ell) = \mathbf{1}$.

Step 1: Zero composite distance ⇒ zero triad. By Proposition 4.507 and Section 4.11.2,

$$D^{[\mathbf{W}_x,\mathbf{W}_{x'}]}(x,x') = 0 \iff \vec{d}^{[\mathbf{W}_x,\mathbf{W}_{x'}]}(x,x') = (0,0,0),$$

i.e. $d_{\mathrm{corr}}^{[\mathbf{W}_x,\mathbf{W}_{x'}]}(x,x') = d_{\mathrm{vac}}^{[\mathbf{W}_x,\mathbf{W}_{x'}]}(x,x') = d_{\mathrm{spec}}^{[\mathbf{W}_x,\mathbf{W}_{x'}]}(x,x') = 0$.

Step 2: Off–diagonal (correlator) sector. Define the bilocal probe at x

$$\mathbf{A}_{\mathrm{corr}}^{[\gamma]} := \mathcal{U}^{[\mathbf{W}_x,\mathbf{W}_{x'}]}(x\leftarrow x';\gamma)^\dagger\,\hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})\,\mathcal{U}^{[\mathbf{W}_x,\mathbf{W}_{x'}]}(x\leftarrow x';\gamma).$$

The vanishing $d_{\mathrm{corr}}^{[\mathbf{W}_x,\mathbf{W}_{x'}]}(x,x') = 0$ means all normalized off–diagonal contrasts vanish in expectation (as in Definition 4.339). Inserting the loop holonomy $\ell = \gamma\circ\bar{\gamma}$ conjugates $\mathbf{A}_{\mathrm{corr}}^{[\gamma]}$

by $\mathcal{U}^{[\mathbf{W}_x,\mathbf{W}_x]}(x \leftarrow x;\ell) = \mathbf{1}$ (Step 0), so every Holo–State expectation is unchanged. Thus holonomy is *trivial in expectation* on the off–diagonal sector (cf. Definition 4.518).

Step 3: Diagonal (vacuum) sector. Let $A_x := \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_x)$, $A_{x'} := \hat{G}_\Lambda(\mathfrak{B}_{x'}, \mathfrak{B}_{x'})$. Let $\{\Pi_n^{(x)[\mathbf{W}_x]}\}$, $\{\Pi_n^{(x')[\mathbf{W}_{x'}]}\}$ be local wordwise spectral frames (Definition 4.405). Since $d_{\mathrm{vac}}^{[\mathbf{W}_x,\mathbf{W}_{x'}]}(x,x') = 0$,

$$\Big\langle \Psi \Big|\, \Pi_n^{(x)[\mathbf{W}_x]} A_x \Pi_n^{(x)[\mathbf{W}_x]} \,\Big|\Psi\Big\rangle = \Big\langle \Psi \Big|\, \Pi_n^{(x')[\mathbf{W}_{x'}]} A_{x'} \Pi_n^{(x')[\mathbf{W}_{x'}]} \,\Big|\Psi\Big\rangle, \quad \forall n.$$

Transporting the x'-profile to x along γ and using unitarity identifies the diagonal blocks in the x-fibre in expectation. Conjugation by $\mathcal{U}^{[\mathbf{W}_x,\mathbf{W}_x]}(x \leftarrow x;\ell) = \mathbf{1}$ leaves these expectations invariant. Hence holonomy is trivial in expectation on the diagonal sector.

Step 4: Spectral (shear/commutator) sector. Define

$$\mathbf{A}_{\mathrm{spec}}^{[\gamma]} := \ i\Big[\mathbb{D}^{[\mathbf{W}_x]}(x),\ \mathbf{A}_{\mathrm{corr}}^{[\gamma]}\Big] \quad \text{(cf. Definition 4.406)}.$$

From $d_{\mathrm{spec}}^{[\mathbf{W}_x,\mathbf{W}_{x'}]}(x,x') = 0$ the band–filtered shear expectations vanish (cf. Definition 4.408). Replacing the window γ by the loop–modified window $\ell \circ \gamma$ gives

$$\mathbf{A}_{\mathrm{corr}}^{[\ell\circ\gamma]} = \ \mathcal{U}^{[\mathbf{W}_x,\mathbf{W}_x]}(x \leftarrow x;\ell)^\dagger\, \mathbf{A}_{\mathrm{corr}}^{[\gamma]}\, \mathcal{U}^{[\mathbf{W}_x,\mathbf{W}_x]}(x \leftarrow x;\ell),$$

so

$$\mathbf{A}_{\mathrm{spec}}^{[\ell\circ\gamma]} = i\Big[\mathbb{D}^{[\mathbf{W}_x]}(x),\, \mathcal{U}^{[\mathbf{W}_x,\mathbf{W}_x]}(x \leftarrow x;\ell)^\dagger\, \mathbf{A}_{\mathrm{corr}}^{[\gamma]}\, \mathcal{U}^{[\mathbf{W}_x,\mathbf{W}_x]}(x \leftarrow x;\ell)\Big].$$

With $\mathcal{U}^{[\mathbf{W}_x,\mathbf{W}_x]}(x \leftarrow x;\ell) = \mathbf{1}$ (Step 0), $\mathbf{A}_{\mathrm{spec}}^{[\ell\circ\gamma]} = \mathbf{A}_{\mathrm{spec}}^{[\gamma]}$, so these expectations remain zero. Thus the spectral axis is holonomically trivial in expectation.

Step 5: Commutator form. For any admissible probe B in the supported wordwise sector of $\hat{G}_\Lambda$,

$$\Big\langle \Psi \Big|\, B^\dagger\, \mathcal{U}^{[\mathbf{W}_x,\mathbf{W}_x]}(x \leftarrow x;\ell)\, B\, \mathcal{U}^{[\mathbf{W}_x,\mathbf{W}_x]}(x \leftarrow x;\ell)^\dagger \,\Big|\Psi\Big\rangle = \Big\langle \Psi \Big|\, B^\dagger B \,\Big|\Psi\Big\rangle,$$

hence $\Big\| \,[\mathcal{U}^{[\mathbf{W}_x,\mathbf{W}_x]}(x \leftarrow x;\ell),\ B]\,|\Psi\rangle \Big\| = 0$. Taking B from the probe algebra generated by entries of $\hat{G}_\Lambda$ (as in Definition 4.518) yields the desired commutator–norm statement.

Collecting the three sectors, holonomy along $\ell = \gamma \circ \bar{\gamma}$ is trivial in the state–conditioned wordwise representation, proving the claim. □

Remark 4.520 (On the loop quantifier)**.** The proof establishes triviality for the closed loop $\ell = \gamma \circ \bar{\gamma}$ generated by a chosen window $\gamma : x' \to x$, for which $\mathcal{U}^{[\mathbf{W}_x,\mathbf{W}_x]}(x \leftarrow x;\ell) = \mathbf{1}$ by the transporter axioms. For *arbitrary* loops ℓ based at x, state–conditioned triviality follows if, in addition, window–independence holds on a flat patch (cf. Lemma 4.448 and Corollary 4.414) or the local hypotheses of Proposition 4.519 apply; then $D^{[\mathbf{W}_x,\mathbf{W}_{x'}]}(x,x') = 0$ propagates to all windows and loops in the patch.

Proof. **Notation.** Let $\mathbf{W}_y$ and $\mathbf{W}_{y'}$ denote the endpoint word frames at y and y'. Write $\mathcal{U}^{[\mathbf{W}_y,\mathbf{W}_{y'}]}(y \leftarrow y';\gamma)$ for transport along an open path $\gamma : y' \to y$, and $\mathcal{U}^{[\mathbf{W}_y,\mathbf{W}_y]}(y \leftarrow y;\ell)$ for a loop ℓ at y.

Step 1: Window–independence in expectation on U. Fix $y, y' \in U$ and let $\gamma_1, \gamma_2 : y' \to y$ be two admissible windows inside U. Since U is simply connected, $\gamma_2 \circ \bar{\gamma}_1$ is contractible. By (a) (holonomy triviality in expectation) and Definition 4.518, for every admissible probe B generated by entries of $\hat{G}_\Lambda$,

$$\Big\langle B^\dagger \mathcal{U}^{[\mathbf{W}_y,\mathbf{W}_y]}(y \leftarrow y; \gamma_2 \circ \bar{\gamma}_1)\, B\, \mathcal{U}^{[\mathbf{W}_y,\mathbf{W}_y]}(y \leftarrow y; \gamma_2 \circ \bar{\gamma}_1)^\dagger \Big\rangle_\Psi = \langle B^\dagger B \rangle_\Psi.$$

Thus all Holo–State expectations of the bilocal/diagonal matrix elements that define the three wordwise distances (cf. Definitions 4.339, 4.408 and 4.468) are *independent of the window* $\gamma \subset U$. Equivalently, with

$$\mathbf{A}^{[\gamma]}_{\mathrm{corr}} := \mathcal{U}^{[\mathbf{W}_y,\mathbf{W}_{y'}]}(y \leftarrow y'; \gamma)^\dagger\, \hat{G}_\Lambda(\mathfrak{B}_y, \mathfrak{B}_{y'})\, \mathcal{U}^{[\mathbf{W}_y,\mathbf{W}_{y'}]}(y \leftarrow y'; \gamma), \qquad A_y := \hat{G}_\Lambda(\mathfrak{B}_y, \mathfrak{B}_y),$$

the relevant expectation–level matrix elements on the supported sector are window–independent.

Step 2: Vanishing of the spectral component. Define the (windowed) spectral–shear generator

$$\mathbf{A}^{[\gamma]}_{\mathrm{spec}} := i\Big[\mathbb{D}^{[\mathbf{W}_y]}(y),\, \mathbf{A}^{[\gamma]}_{\mathrm{corr}}\Big] \quad \text{(cf. Definition 4.406)}.$$

Assumption (b) (*spectral quiescence*) states that, for all $y, y' \in U$ and all relevant bands n, the Holo–State expectations of the band–filtered components of $\mathbf{A}^{[\gamma]}_{\mathrm{spec}}$ vanish (equivalently with the transported $\mathbb{D}^{[\mathbf{W}_{y'}]}(y')$ as stated). Hence, by Definition 4.408,

$$d^{[\mathbf{W}_y,\mathbf{W}_{y'}]}_{\mathrm{spec}}(y, y') = 0 \quad \text{for all } y, y' \in U.$$

Step 3: Modal alignment and Cauchy–Schwarz saturation (correlator axis). Let $\{\Pi^{(y)[\mathbf{W}_y]}_n\}_n$ and $\{\Pi^{(y')[\mathbf{W}_{y'}]}_n\}_n$ be local wordwise spectral frames (Definition 4.405). Using (b) and Step 1 (window–independence), for each band n,

$$\Big\langle \Psi \Big|\, \Pi^{(y)[\mathbf{W}_y]}_n\, \mathbf{A}^{[\gamma]}_{\mathrm{corr}}\, \Pi^{(y)[\mathbf{W}_y]}_n \,\Big|\Psi\Big\rangle = \Big\langle \Psi \Big|\, \Pi^{(y)[\mathbf{W}_y]}_n\, \mathbf{A}^{[\gamma]}_{\mathrm{corr}} \,\Big|\Psi\Big\rangle,$$

so the contributing matrix elements are supported within the same modal band (no "shear" in expectation). Transporting the y'-sector to y, the normalized correlator overlap (cf. Definition 4.339) saturates Cauchy–Schwarz bandwise; consequently,

$$d^{[\mathbf{W}_y,\mathbf{W}_{y'}]}_{\mathrm{corr}}(y, y') = 0 \quad \text{for all } y, y' \in U.$$

Step 4: Equality of diagonal vacuum profiles. Write $C^{[\mathbf{W}]}_n(y) := \langle \Psi | \Pi^{(y)[\mathbf{W}]}_n A_y \Pi^{(y)[\mathbf{W}]}_n | \Psi \rangle$ (cf. Definition 4.466). By Step 1 (window–independence) and the bandwise alignment from Step 3,

$$C^{[\mathbf{W}_y]}_n(y) = C^{[\mathbf{W}_{y'}]}_n(y') \quad \forall n,$$

hence the weighted ℓ^2 difference of vacuum profiles vanishes (Definition 4.468 and Corollary 4.472):

$$d^{[\mathbf{W}_y,\mathbf{W}_{y'}]}_{\mathrm{vac}}(y, y') = 0 \quad \text{for all } y, y' \in U.$$

Step 5: Composite distance. All three components vanish on U:

$$d_{\mathrm{corr}}^{[\mathbf{W}_y,\mathbf{W}_{y'}]}(y,y') = d_{\mathrm{vac}}^{[\mathbf{W}_y,\mathbf{W}_{y'}]}(y,y') = d_{\mathrm{spec}}^{[\mathbf{W}_y,\mathbf{W}_{y'}]}(y,y') = 0.$$

Thus $\vec{d}^{[\mathbf{W}_y,\mathbf{W}_{y'}]}(y,y') = \mathbf{0}$, and by Sections 4.11.2 and 4.11.4,

$$D^{[\mathbf{W}_y,\mathbf{W}_{y'}]}(y,y') = \left\|\mathbf{M}^{1/2}(y)\,\vec{d}^{[\mathbf{W}_y,\mathbf{W}_{y'}]}(y,y')\right\|_2 = 0,$$

for all $y, y' \in U$. □

Remark 4.521 (Representation–dependence and "holonomic invisibility" (wordwise))**.** Nontrivial holonomy may act *invisibly* if its action lies in a sector annihilated by the wordwise weights $w_n^{[\mathbf{W}_1,\mathbf{W}_2]}$ (Definition 4.407) or by the expectation map (e.g. phases cancelling in $\langle\cdot\rangle_\Psi$). The relevant criterion is triviality in the *state–conditioned wordwise representation* carried by the probe algebra; such triviality suffices for $D^{[\cdot]} = 0$ even if the underlying modal connection has nonzero curvature.

Remark 4.522 (Curvature, holonomy, and the triadic metric (wordwise))**.** In classical differential geometry, holonomy is generated by curvature (Ambrose–Singer theorem, cf. [405, 248]). In the wordwise calculus, the operator–valued curvature of the modal connection contributes to the triadic metric via: (i) path–dependence of bilocal coherences (the correlator axis $d_{\mathrm{corr}}^{[\cdot]}$), (ii) shifts in coincident self–coherences (the vacuum axis $d_{\mathrm{vac}}^{[\cdot]}$), and (iii) nonvanishing commutator tension with $\mathbb{D}^{[\mathbf{W}]}$ (the spectral axis $d_{\mathrm{spec}}^{[\cdot]}$). Vanishing of these effects in the state–conditioned representation is precisely the content of Proposition 4.519.

4.12 The Triadic Structure of Emergent Time

Complementary to the emergent spatial geometry, the theory defines a triad of emergent times. We begin with the *entanglement phase time*, the phase complement of the correlator–based distance (Section 4.9.1).

4.12.1 Entanglement Phase Time

Let $\hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})[\psi_1^*,\psi_2]$ denote the Fourfold Inter–Modal Correlator with gauge–covariant transport $\mathcal{U}(x\leftarrow x';\gamma)$ (Definition 4.9, cf. eq. (4.131)). For fixed test data ψ on the relevant fibres, write the Holo–State expectation as

$$\Gamma_{\mathrm{corr}}(x,x';\gamma) := \left\langle \mathring{\mathbf{\Psi}} \middle| \hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})[\psi^*,\psi] \middle| \mathring{\mathbf{\Psi}} \right\rangle \in \mathbb{C}.$$

Definition 4.523 (Entanglement Phase Time)**.** For pairs (x,x') and admissible paths γ such that $\Gamma_{\mathrm{corr}}(x,x';\gamma) \neq 0$, the *entanglement phase time* is the argument of the complex correlator expectation,

$$\boxed{t_{\mathrm{corr}}(x,x';\gamma) := \mathrm{Arg}\ \Gamma_{\mathrm{corr}}(x,x';\gamma), \qquad t_{\mathrm{corr}} \in (-\pi,\pi].} \tag{4.243}$$

If a continuous path family $s \mapsto \gamma_s$ connects (x, x') while keeping $\Gamma_{\mathrm{corr}} \neq 0$, the *unwrapped* phase is defined by continuous lifting,

$$\tilde{t}_{\mathrm{corr}}(x, x'; \gamma) := \text{cont. lift Arg } \Gamma_{\mathrm{corr}} \in \mathbb{R},$$

unique up to addition of $2\pi k$, $k \in \mathbb{Z}$. At coincidence, $t_{\mathrm{corr}}(x, x; \cdot) = 0$.

Conceptually, t_{corr} is the phase complement of the correlator–based distance (Section 4.9.1), generalising Pancharatnam's interferometric phase to the operator–relational setting [23], aligning with Berry's adiabatic geometric phase [24], and with the kinematic/geometric treatment of quantum evolution [25]. Unlike the usual time–evolution setups, the phase here accrues along prismatic, gauge–covariant transports across emergent space, embedded in the IQR relational geometry.

Remark 4.524 (Regularity and Nodes)**.** At *nodes* where $\Gamma_{\mathrm{corr}}(x, x'; \gamma) = 0$, the phase is undefined. In practice one uses (i) limiting values along node–avoiding paths, or (ii) a small reference admixture $\psi \mapsto \psi_\varepsilon$ to lift the phase off the branch cut and then sends $\varepsilon \to 0$.

Proposition 4.525 (Gauge and Rephasing Invariance)**.** t_{corr} *is invariant (mod* 2π*) under:*

(i) local unitary frame changes at the endpoints, $\hat{\mathcal{F}}_\alpha(y) \mapsto U(y)\hat{\mathcal{F}}_\alpha(y)U(y)^\dagger$, $\mathcal{U} \mapsto U(x)\mathcal{U}U(x')^\dagger$*;*

(ii) fibre rephasings of the test data $\psi \mapsto e^{i\phi_x}\psi$ *at* x *and* $e^{i\phi_{x'}}\psi$ *at* x'*;*

(iii) overall normalisation of ψ*.*

Proof sketch. Each transformation multiplies Γ_{corr} by a nonzero complex factor of unit modulus whose phase cancels between bra/ket or endpoints by gauge covariance; hence Arg is unchanged modulo 2π.

Proposition 4.526 (Basic Properties)**.** *On any fixed homotopy class of paths avoiding nodes:*

*(i) (*antisymmetry*)* $t_{\mathrm{corr}}(x', x; \bar{\gamma}) \equiv -\,t_{\mathrm{corr}}(x, x'; \gamma) \pmod{2\pi}$*, where* $\bar{\gamma}$ *is* γ *reversed;*

*(ii) (*concatenation*) for concatenable segments* γ_{12} $(x \to y)$ *and* γ_{23} $(y \to x')$ *with* $\Gamma_{\mathrm{corr}} \neq 0$*,*

$$\tilde{t}_{\mathrm{corr}}(x, x'; \gamma_{23} \circ \gamma_{12}) = \tilde{t}_{\mathrm{corr}}(x, y; \gamma_{12}) + \tilde{t}_{\mathrm{corr}}(y, x'; \gamma_{23});$$

*(iii) (*holonomy*) on a loop* $\gamma : x \to x$ *avoiding nodes,* $\tilde{t}_{\mathrm{corr}}(x, x; \gamma) = 2\pi\, n(\gamma)$ *detects the correlator–phase holonomy;* $n(\gamma) \in \mathbb{Z}$ *may be nonzero when the correlator's phase connection has nontrivial curvature.*

Definition 4.527 (Phase–Rate 1–Form)**.** For a smooth path $s \mapsto \gamma(s)$ with $\Gamma_{\mathrm{corr}}(s) \neq 0$, define the (gauge–invariant) phase rate

$$\dot{t}_{\mathrm{corr}}(s) = \frac{d}{ds} \arg\, \Gamma_{\mathrm{corr}}(s) = \mathrm{Im}\left(\frac{\dot{\Gamma}_{\mathrm{corr}}(s)}{\Gamma_{\mathrm{corr}}(s)} \right).$$

The integral $t_{\mathrm{corr}} = \int \dot{t}_{\mathrm{corr}}\, ds$ reproduces eq. (4.243) up to $2\pi\mathbb{Z}$ and makes the dependence on the path–ordered parallel transport explicit.

Remark 4.528 (Complementarity with d_{corr} and Calibration)**.** Writing $\Gamma_{\mathrm{corr}} = \|\Gamma_{\mathrm{corr}}\|\, e^{it_{\mathrm{corr}}}$ shows that d_{corr} (Section 4.9.1) and t_{corr} are, respectively, the magnitude and argument of the *same* bilocal gauge–covariant expectation. The phase is dimensionless; if a dimensionful time is desired, introduce a state–dependent calibration τ_* and set $T_{\mathrm{corr}} := \tau_*\, \tilde{t}_{\mathrm{corr}}$ (choice deferred to Section 5.3.8).

4.12.2 Wordwise Entanglement Phase Time

Let $\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\cdot,\cdot]$ denote the Fourfold inter–modal correlator and $\mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x \leftarrow x';\gamma)$ the admissible *wordwise* transporter along a path $\gamma : x' \to x$ (Section 4.5 and Definition 4.407). For a fixed test vector ψ on the relevant fibres (with the same symbol used at x and x' for brevity), define the complex wordwise correlator expectation

$$\Gamma^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{corr}}(x,x';\gamma \mid \psi) := \Big\langle \mathring{\mathbf{\Psi}} \Big| \, \hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'}) \Big[(U_1\psi)^*, \, \mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x \leftarrow x';\gamma)\, U_2\psi \Big] \, \Big| \mathring{\mathbf{\Psi}} \Big\rangle \; \in \; \mathbb{C}, \tag{4.244}$$

where $U_{1,2}$ are the local endpoint word unitaries at x, x' (cf. Definition 4.339 and Section 4.5). When no confusion arises we suppress the ψ–tag.

Definition 4.529 (Wordwise entanglement phase time)**.** For pairs (x,x') and admissible paths γ such that

$$\Gamma^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{corr}}(x,x';\gamma) \neq 0,$$

define the *wordwise entanglement phase time*

$$\boxed{t^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{corr}}(x,x';\gamma) := \operatorname{Arg} \Gamma^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{corr}}(x,x';\gamma), \qquad t^{[\cdot]}_{\mathrm{corr}} \in (-\pi,\pi].} \tag{4.245}$$

If a continuous path family $s \mapsto \gamma_s$ keeps $\Gamma^{[\cdot]}_{\mathrm{corr}} \neq 0$, the *unwrapped* phase is the continuous lift $\tilde{t}^{[\cdot]}_{\mathrm{corr}}(x,x';\gamma) \in \mathbb{R}$, unique up to $2\pi\mathbb{Z}$. At coincidence, $t^{[\cdot]}_{\mathrm{corr}}(x,x;\,\cdot) = 0$.

Conceptually, $t^{[\cdot]}_{\mathrm{corr}}$ is the phase complement of the wordwise correlator–based distance (Definition 4.339), generalising Pancharatnam's interferometric phase to the holosymmetric, operator–relational setting [23], aligning with Berry's geometric phase [24] and the kinematic view of quantum evolution [25]. Unlike standard time–evolution, the phase here accrues along prismatic, gauge–/word– covariant transports across the emergent configuration space.

Remark 4.530 (Regularity and nodes (wordwise))**.** At *nodes* where $\Gamma^{[\cdot]}_{\mathrm{corr}}(x,x';\gamma) = 0$ the phase is undefined. One uses (i) node–avoiding limits along homotopic families, or (ii) a small reference admixture $\psi \mapsto \psi_\varepsilon$ (lifting the phase off the branch cut) and then $\varepsilon \to 0$.

Proposition 4.531 (Gauge/word and rephasing invariance)**.** $t^{[\cdot]}_{\mathrm{corr}}$ *is invariant (mod* 2π*) under:*

(i) Endpoint word or gauge changes.
These act as

$$\hat{\mathcal{F}}_\alpha(y) \mapsto U(y)\hat{\mathcal{F}}_\alpha(y)U(y)^\dagger, \qquad \mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]} \mapsto U(x)\,\mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}U(x')^\dagger.$$

(ii) Fibre rephasings of the test data.
These are given by

$$\psi \mapsto e^{i\phi_x}\psi \quad \text{at } x, \qquad \psi \mapsto e^{i\phi_{x'}}\psi \quad \text{at } x'.$$

(iii) Overall normalisation of ψ.
(Global rescalings of ψ leave $t^{[\cdot]}_{\mathrm{corr}}$ unchanged.)

Proof. Recall

$$\Gamma_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma \mid \psi) = \Big\langle \mathring{\mathbf{\Psi}} \Big| \hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'}) \Big[(U_1\psi)^*,\, \mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x \leftarrow x';\gamma)\, U_2\psi \Big] \Big| \mathring{\mathbf{\Psi}} \Big\rangle,$$

with $t_{\mathrm{corr}}^{[\cdot]} = \operatorname{Arg} \Gamma_{\mathrm{corr}}^{[\cdot]}$ (eqs. (4.244) and (4.245)). We treat the three items in turn.

(1) Endpoint word/gauge changes. Let $V_x, V_{x'}$ be unitaries at x, x' implementing the local word/gauge change (Section 4.5). The admissible objects transform covariantly:

$$U_1 \mapsto V_x U_1, \qquad U_2 \mapsto V_{x'} U_2, \qquad \mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x \leftarrow x';\gamma) \mapsto V_x\, \mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x \leftarrow x';\gamma)\, V_{x'}^\dagger,$$

and the bilinear evaluation of the Fourfold correlator is endpoint-covariant (basis change at each end). Substituting,

$$\begin{aligned} \Gamma' &= \Big\langle \Psi \Big| \hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'}) \Big[(V_x U_1\psi)^*,\, V_x \mathcal{U}^{[\cdot]} V_{x'}^\dagger V_{x'} U_2\psi \Big] \Big| \Psi \Big\rangle \\ &= \Big\langle \Psi \Big| \hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'}) \Big[(V_x U_1\psi)^*,\, V_x \mathcal{U}^{[\cdot]} U_2\psi \Big] \Big| \Psi \Big\rangle \\ &= \Big\langle \Psi \Big| \hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'}) \Big[(U_1\psi)^*,\, \mathcal{U}^{[\cdot]} U_2\psi \Big] \Big| \Psi \Big\rangle \;=\; \Gamma, \end{aligned}$$

where the last equality uses endpoint covariance of $\hat{G}_\Lambda$ under the common conjugation at x (Definition 4.339) and unitarity of V_x. Hence $\operatorname{Arg} \Gamma' = \operatorname{Arg} \Gamma \mod 2\pi$.

(2) Fibre rephasings of the test data. Let $\psi \mapsto e^{i\phi_x}\psi$ at x and $\psi \mapsto e^{i\phi_{x'}}\psi$ at x', keeping $U_{1,2}$ and $\mathcal{U}^{[\cdot]}$ fixed. By conjugate-linearity/linearity of the correlator in its two slots,

$$\Gamma_{\mathrm{corr}} \;\longmapsto\; e^{-i\phi_x}\, e^{i\phi_{x'}}\, \Gamma_{\mathrm{corr}}.$$

Thus the argument shifts by $\phi_{x'} - \phi_x$. However such *endpoint rephasings* are a special case of the word/gauge change in (1), realized by $V_x = e^{-i\phi_x}\mathbf{1}$ and $V_{x'} = e^{-i\phi_{x'}}\mathbf{1}$, which induces the inverse phase on the transported ket and restores the original value of Γ_{corr} (the transporter simultaneously picks up $V_x(\cdot)V_{x'}^\dagger$ and the first-slot vector picks up V_x). Consequently, modulo the allowed endpoint word/gauge freedom, $t_{\mathrm{corr}}^{[\cdot]}$ is invariant under fibre rephasings.[4]

(3) Overall normalisation of ψ. Let $\psi \mapsto a\,\psi$ with $a \in \mathbb{C} \setminus \{0\}$ applied at both ends (same test data used at x and x'). Then by sesquilinearity,

$$\Gamma_{\mathrm{corr}} \;\longmapsto\; a^* a\, \Gamma_{\mathrm{corr}} = |a|^2\, \Gamma_{\mathrm{corr}},$$

so only the modulus changes and Arg is unchanged modulo 2π.

Combining (1)–(3) yields the stated invariances of $t_{\mathrm{corr}}^{[\cdot]}$.

□

Proposition 4.532 (Basic properties (fixed homotopy class, node–free))**.** *For paths avoiding nodes:*

(i) (antisymmetry) $t_{\mathrm{corr}}^{[\mathbf{W}_2,\mathbf{W}_1]}(x',x;\bar{\gamma}) \equiv -\, t_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma) \pmod{2\pi}$;

[4] Equivalently: one may fix a local trivialization (gauge) and regard endpoint rephasings as gauge changes; $t_{\mathrm{corr}}^{[\cdot]}$ is a gauge-invariant function of the gauge orbit.

(ii) (concatenation) $\tilde{t}^{[\cdot]}_{\mathrm{corr}}(x,x';\gamma_{23}\circ\gamma_{12}) = \tilde{t}^{[\cdot]}_{\mathrm{corr}}(x,y;\gamma_{12}) + \tilde{t}^{[\cdot]}_{\mathrm{corr}}(y,x';\gamma_{23})$;

(iii) (holonomy) *on a loop* $\gamma : x \to x$ *avoiding nodes,* $\tilde{t}^{[\cdot]}_{\mathrm{corr}}(x,x;\gamma) = 2\pi\, n(\gamma)$ *detects the correlator–phase holonomy, with* $n(\gamma) \in \mathbb{Z}$ *the winding number.*

Proof. Recall the wordwise complex overlap $\Gamma^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{corr}}(x,x';\gamma)$ from eq. (4.244) and $t^{[\cdot]}_{\mathrm{corr}} = \operatorname{Arg}\Gamma^{[\cdot]}_{\mathrm{corr}}$ from eq. (4.245). Throughout, we assume the chosen paths lie in a node–free homotopy class so that $\Gamma^{[\cdot]}_{\mathrm{corr}} \neq 0$ and the unwrapped phase $\tilde{t}^{[\cdot]}_{\mathrm{corr}}$ exists as a continuous lift (Definitions 4.529 and 4.533).

(1) Antisymmetry under endpoint swap and path reversal. Let $\bar{\gamma}$ be the reversal of γ, so $\mathcal{U}^{[\mathbf{W}_2,\mathbf{W}_1]}(x' \leftarrow x;\bar{\gamma}) = \mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x \leftarrow x';\gamma)^\dagger$ by unitarity and inversion (Section 4.5). Using the conjugate–linearity/linearity of the correlator in its two slots and the fact that swapping endpoints exchanges the slots, we compute

$$\begin{aligned}\Gamma^{[\mathbf{W}_2,\mathbf{W}_1]}_{\mathrm{corr}}(x',x;\bar{\gamma}) &= \Big\langle \Psi \Big|\, \hat{G}_\Lambda(\mathfrak{B}_{x'},\mathfrak{B}_x)\Big[(U_2\psi)^*,\, \mathcal{U}^{[\mathbf{W}_2,\mathbf{W}_1]}(x' \leftarrow x;\bar{\gamma})\, U_1\psi\Big] \,\Big|\, \Psi\Big\rangle \\ &= \Big\langle \Psi \Big|\, \hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})\Big[\mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x \leftarrow x';\gamma)\, U_2\psi,\ (U_1\psi)^*\Big] \,\Big|\, \Psi\Big\rangle^* \\ &= \Big(\Gamma^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{corr}}(x,x';\gamma)\Big)^*,\end{aligned}$$

where in the second line we used the adjoint transporter and the Hermitian symmetry of the kernel under slot exchange. Taking arguments and using $\operatorname{Arg}(z^*) \equiv -\operatorname{Arg}(z) \pmod{2\pi}$ gives

$$t^{[\mathbf{W}_2,\mathbf{W}_1]}_{\mathrm{corr}}(x',x;\bar{\gamma}) \equiv -\,t^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{corr}}(x,x';\gamma) \pmod{2\pi}.$$

(2) Concatenation (additivity of the unwrapped phase). Let $\gamma_{12} : x \to y$ and $\gamma_{23} : y \to x'$ be node–free, and set $\gamma_{13} = \gamma_{23} \circ \gamma_{12}$. Consider a smooth one–parameter interpolation $s \mapsto \gamma_s$ from γ_{12} followed by γ_{23} to γ_{13} within the same node–free homotopy class. By Definition 4.533,

$$\tilde{t}^{[\cdot]}_{\mathrm{corr}}(x,x';\gamma) = \int_0^1 \operatorname{Im}\!\left(\frac{\dot{\Gamma}^{[\cdot]}_{\mathrm{corr}}(s)}{\Gamma^{[\cdot]}_{\mathrm{corr}}(s)}\right) ds \quad \text{(path–ordered, node–free).}$$

Since $\Gamma^{[\cdot]}_{\mathrm{corr}}(s) \neq 0$, the integrand is continuous and the integral is additive under concatenation of the parameter intervals. Applying this first over the segment that deforms the identity path at x into γ_{12} and then over the segment that deforms the identity path at y into γ_{23} yields

$$\tilde{t}^{[\cdot]}_{\mathrm{corr}}(x,x';\gamma_{23}\circ\gamma_{12}) = \tilde{t}^{[\cdot]}_{\mathrm{corr}}(x,y;\gamma_{12}) + \tilde{t}^{[\cdot]}_{\mathrm{corr}}(y,x';\gamma_{23}).$$

Equivalently, additivity follows from $\frac{d}{ds}\arg(\Gamma_1\Gamma_2) = \operatorname{Im}\big((\dot{\Gamma}_1/\Gamma_1) + (\dot{\Gamma}_2/\Gamma_2)\big)$ applied to the ordered product structure of the transported argument in eq. (4.244).

(3) Holonomy on loops (winding number). Let γ be a node–free loop based at x and parametrize it by $s \in [0,1]$ with $\gamma(0) = \gamma(1) = x$. Then $s \mapsto \Gamma^{[\cdot]}_{\mathrm{corr}}(x,x;\gamma_s)$ is a continuous map into $\mathbb{C} \setminus \{0\}$. Define

$$\hat{\Gamma}(s) := \frac{\Gamma^{[\cdot]}_{\mathrm{corr}}(x,x;\gamma_s)}{|\Gamma^{[\cdot]}_{\mathrm{corr}}(x,x;\gamma_s)|} \in U(1).$$

Its degree (winding number) around the unit circle is

$$n(\gamma) = \frac{1}{2\pi}\int_0^1 \mathrm{Im}\left(\frac{\dot\Gamma^{[\cdot]}_{\mathrm{corr}}(s)}{\Gamma^{[\cdot]}_{\mathrm{corr}}(s)}\right) ds = \frac{1}{2\pi}\,\tilde t^{[\cdot]}_{\mathrm{corr}}(x,x;\gamma) \;\in\; \mathbb{Z},$$

where the equality uses Definition 4.533. Thus $\tilde t^{[\cdot]}_{\mathrm{corr}}(x,x;\gamma) = 2\pi\, n(\gamma)$, detecting the correlator–phase holonomy. Nonzero $n(\gamma)$ signals nontrivial curvature of the underlying phase connection sampled by the wordwise transport along γ. □

Definition 4.533 (Wordwise phase–rate 1–form)**.** For a smooth family $s \mapsto \gamma(s)$ with $\Gamma^{[\cdot]}_{\mathrm{corr}}(s) \neq 0$, define the gauge/word–invariant phase rate

$$\dot t^{[\cdot]}_{\mathrm{corr}}(s) = \frac{d}{ds}\arg\,\Gamma^{[\cdot]}_{\mathrm{corr}}(s) = \mathrm{Im}\left(\frac{\dot\Gamma^{[\cdot]}_{\mathrm{corr}}(s)}{\Gamma^{[\cdot]}_{\mathrm{corr}}(s)}\right),$$

so that $t^{[\cdot]}_{\mathrm{corr}} = \int \dot t^{[\cdot]}_{\mathrm{corr}}\, ds$ reproduces (4.245) up to $2\pi\mathbb{Z}$ and makes the dependence on the path–ordered transporter explicit.

Remark 4.534 (Complementarity with $d^{[\cdot]}_{\mathrm{corr}}$ and normalized complex overlap)**.** Let the wordwise coincident normalisations be $N^{[\mathbf{W}_1]}(\psi,x)$ and $N^{[\mathbf{W}_2]}(\psi,x')$ from Definition 4.339. Define the *normalized complex overlap*

$$\Upsilon^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma) := \frac{\Gamma^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{corr}}(x,x';\gamma)}{N^{[\mathbf{W}_1]}(\psi,x)\,N^{[\mathbf{W}_2]}(\psi,x')} = C^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\psi,\psi)\,e^{\,i\,t^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{corr}}(x,x';\gamma)},$$

so the wordwise correlator distance uses the magnitude $C^{[\cdot]} = |\Upsilon^{[\cdot]}|$ while $t^{[\cdot]}_{\mathrm{corr}} = \mathrm{Arg}\;\Upsilon^{[\cdot]}$ is its argument. The phase is dimensionless; for a dimensionful time, introduce a state–dependent calibration τ_* and set $T^{[\cdot]}_{\mathrm{corr}} := \tau_*\,\tilde t^{[\cdot]}_{\mathrm{corr}}$.

4.12.3 Vacuum Response Time

The second emergent temporal observable, $t_{\mathrm{vac}}(x,x')$, quantifies the vacuum's *phase–sensitive* response to relational displacements. It is built from the same diagonal (coincident) probes that define the vacuum–response distance d_{vac} (Definition 4.463), but isolates the imaginary part of their Holo–State expectations, thereby capturing phase–encoded distortions of the entanglement–defined vacuum structure. Concretely, fix the local self–correlators $C_k(x)$ exactly as in the definition of d_{vac} (Definition 4.463) and its vector formulation (4.211); write $\vec V_{\mathrm{vac}}(x) := (C_1(x), C_2(x), \dots)$ and denote componentwise imaginary parts by $\mathrm{Im}\,\vec V_{\mathrm{vac}}(x)$.

Formal Definition

Definition 4.535 (Vacuum response time)**.** With $C_k(x)$ as above, set $I_k(x) := \mathrm{Im}\,C_k(x)$. The *vacuum response time* between x and x' is

$$\boxed{t_{\mathrm{vac}}(x,x') \;:=\; \Big[\sum\nolimits_k \big(I_k(x) - I_k(x')\big)^2\Big]^{1/2} \;=\; \big\|\mathrm{Im}\,\vec V_{\mathrm{vac}}(x) - \mathrm{Im}\,\vec V_{\mathrm{vac}}(x')\big\|_2 .} \tag{4.246}$$

Equivalently, if $\{\hat{O}_k(x)\}_k$ are the localized diagonal probes realizing $C_k(x) = \langle \overset{\circ}{\boldsymbol{\Psi}} | \, \hat{O}_k(x) \, | \overset{\circ}{\boldsymbol{\Psi}} \rangle$, one may write

$$t_{\mathrm{vac}}(x,x') = \left[\sum\nolimits_k \left(\mathrm{Im} \, \left\langle \overset{\circ}{\boldsymbol{\Psi}} \middle| \hat{O}_k(x) - \hat{O}_k(x') \middle| \overset{\circ}{\boldsymbol{\Psi}} \right\rangle \right)^2 \right]^{1/2},$$

which matches the heuristic form previously stated.

Interpretation and Duality. By construction, t_{vac} is the "phase–time" dual of d_{vac}: the latter measures the *magnitude* of diagonal variation, while the former measures its *imaginary* (phase–like) part, both computed in the same Holo–State and from the same coincident sector of $\hat{G}_\Lambda$. Hence $|\mathrm{Im}\, z| \le |z|$ implies the pointwise bound

$$\boxed{t_{\mathrm{vac}}(x,x') \;\le\; d_{\mathrm{vac}}(x,x') \quad \text{for all } x,x' \in \mathcal{X}_q.} \tag{4.247}$$

Equality holds iff each modal difference $C_k(x) - C_k(x')$ is purely imaginary.

Basic Properties

Proposition 4.536 (Basic properties of t_{vac})**.** *For all $x, x', x'' \in \mathcal{X}_q$:*

(i) Nonnegativity, symmetry, triangle. $t_{\mathrm{vac}}(x,x') \ge 0$, $t_{\mathrm{vac}}(x,x') = t_{\mathrm{vac}}(x',x)$, *and* $t_{\mathrm{vac}}(x,x'') \le t_{\mathrm{vac}}(x,x') + t_{\mathrm{vac}}(x',x'')$. *Thus t_{vac} is a pseudometric; it is a metric on any sector where the map $x \mapsto \mathrm{Im}\, \vec{V}_{\mathrm{vac}}(x)$ separates points.*

(ii) Gauge/Frame covariance and basis–independence. *Modal bases are induced by the coincident–limit inner product of $\hat{G}_\Lambda$; under local unitary rotations of that basis, $\mathrm{Im}\, \vec{V}_{\mathrm{vac}}$ transforms orthogonally and ℓ^2–norms are preserved, so t_{vac} is basis–independent and gauge–covariant in the same sense as d_{vac}.*

(iii) Duality bound. *The inequality (4.247) holds componentwise by $|\mathrm{Im}\, z| \le |z|$.*

(iv) Finiteness (coherent regularisation). *Under the standard hypotheses (Finite Entanglement Capacity, Spectral/Relational Smoothness), all temporal counterparts of the distances are finite; in particular $t_{\mathrm{vac}}(x,x') < \infty$ for all x, x'.*

(v) Relational null criterion. *$t_{\mathrm{vac}}(x,x') = 0$ iff $\mathrm{Im}\, C_k(x) = \mathrm{Im}\, C_k(x')$ for all k. Together with the vanishing of the other components, this is part of the relational null condition characterising a PQS locus ($\vec{d} = \vec{t} = \mathbf{0}$).*

Rate Form Along a Path

Definition 4.537 (Vacuum–response rate 1-form)**.** Let $\gamma : [0,1] \to \mathcal{X}_q$ be a C^1 path and use the gauge–covariant identification of modal fibres along γ (parallel transport as in the Fourfold construction). Define

$$\boxed{\dot{t}_{\mathrm{vac}}(s) \;:=\; \left[\sum\nolimits_k \left(\tfrac{\mathrm{d}}{\mathrm{d}s} \, \mathrm{Im}\, C_k(\gamma(s)) \right)^2 \right]^{1/2}, \qquad T_{\mathrm{vac}}(x,x';\gamma) := \int_0^1 \dot{t}_{\mathrm{vac}}(s)\, \mathrm{d}s.} \tag{4.248}$$

Then $t_{\mathrm{vac}}(x,x') \le T_{\mathrm{vac}}(x,x';\gamma)$ with equality whenever the image $s \mapsto \operatorname{Im}\vec{V}_{\mathrm{vac}}(\gamma(s))$ is a straight segment in profile space (chord–length equality in ℓ^2). The construction is basis–independent by the same covariance used for $\hat{G}_\Lambda$.

Remark 4.538 (Placement in the Temporal Fibre and Calibration)**.** The temporal flow fibre $\mathcal{F}_x^{\mathrm{time}}$ houses the triad $(t_{\mathrm{corr}}, t_{\mathrm{vac}}, t_{\mathrm{spec}})$, dual to the distance fibre $\mathcal{F}_x^{\mathrm{dist}}$ (Definition 4.604, *via* $\mathfrak{F}_x = \mathcal{F}_x^{\mathrm{dist}} \times \mathcal{F}_x^{\mathrm{time}}$). As with t_{corr}, t_{vac} is dimensionless; if a dimensionful parameter is required one introduces a state–dependent calibration scale τ_*, to be fixed together with the emergent proper–time discussion.

Physical Reading. t_{vac} measures the *retarded internal reaction* of the vacuum to modal shifts, sensitive to phase–encoded polarisation and gauge flow in the coincident sector. It is thus the operator–theoretic, background–free analogue of phase response familiar from quantum field theory on curved backgrounds [53, 10, 406], here rendered intrinsic via the Holo–State and $\hat{G}_\Lambda$.

4.12.4 Wordwise Vacuum Response Time

The second emergent temporal observable, $t_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')$, quantifies the vacuum's *phase–sensitive* response to relational displacements in the holosymmetric wordwise calculus. It is built from the same diagonal (coincident) probes that define the wordwise vacuum–response distance $d_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}$ (Definition 4.468), but isolates the imaginary parts of their Holo–State expectations, thereby capturing phase–encoded distortions of the entanglement–defined vacuum structure. Concretely, use the wordwise self–correlators $C_n^{[\mathbf{W}]}(x)$ of Definition 4.466 and the vector profile $\vec{V}_{\mathrm{vac}}^{[\mathbf{W}]}(x) := \big(C_1^{[\mathbf{W}]}(x), C_2^{[\mathbf{W}]}(x), \dots\big)$; write $\operatorname{Im}\vec{V}_{\mathrm{vac}}^{[\mathbf{W}]}(x)$ for the componentwise imaginary parts.

Formal definition (wordwise)

Definition 4.539 (Wordwise vacuum response time)**.** Set $I_n^{[\mathbf{W}]}(x) := \operatorname{Im} C_n^{[\mathbf{W}]}(x)$. For admissible endpoint words $\mathbf{W}_1, \mathbf{W}_2$ at x, x', define

$$\boxed{t_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \;:=\; \Big\|\operatorname{Im}\vec{V}_{\mathrm{vac}}^{[\mathbf{W}_1]}(x) - \operatorname{Im}\vec{V}_{\mathrm{vac}}^{[\mathbf{W}_2]}(x')\Big\|_{\ell^2} \;=\; \Big[\sum\nolimits_n \Big(I_n^{[\mathbf{W}_1]}(x) - I_n^{[\mathbf{W}_2]}(x')\Big)^2\Big]^{1/2}.}$$

(4.249)

When a single–fibre comparison is preferred, transport the x'–profile to x with the wordwise transporter $\mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x \leftarrow x';\gamma)$ as in Definition 4.468; the value is independent of the compatible trivialization and is Lipschitz in the holonomy size.

Interpretation and duality (wordwise). By construction, $t_{\mathrm{vac}}^{[\cdot]}$ is the *phase–time* dual of $d_{\mathrm{vac}}^{[\cdot]}$: the latter measures the *magnitude* of diagonal variation, while the former measures

its *imaginary* (phase–like) part, both computed in the same Holo–State and from the same coincident sector of $\hat{G}_\Lambda$. Hence $|\operatorname{Im} z| \le |z|$ implies the pointwise bound

$$\boxed{t_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \;\le\; d_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \qquad \text{for all } x,x' \in \mathcal{X}_q,} \tag{4.250}$$

with equality iff each modal difference $C_n^{[\mathbf{W}_1]}(x) - C_n^{[\mathbf{W}_2]}(x')$ is purely imaginary.

Basic properties

Proposition 4.540 (Basic properties of $t_{\mathrm{vac}}^{[\cdot]}$)**.** *For all $x, x', x'' \in \mathcal{X}_q$ and admissible words:*

(i) Nonnegativity, symmetry, triangle. $t_{\mathrm{vac}}^{[\cdot]}(x,x') \ge 0$, $t_{\mathrm{vac}}^{[\cdot]}(x,x') = t_{\mathrm{vac}}^{[\cdot]}(x',x)$, *and* $t_{\mathrm{vac}}^{[\cdot]}(x,x'') \le t_{\mathrm{vac}}^{[\cdot]}(x,x') + t_{\mathrm{vac}}^{[\cdot]}(x',x'')$. *Thus $t_{\mathrm{vac}}^{[\cdot]}$ is a pseudometric; it is a metric on any sector where $x \mapsto \operatorname{Im} \vec{V}_{\mathrm{vac}}^{[\mathbf{W}]}(x)$ separates points.*

(ii) Word/gauge covariance and basis–independence. *Modal bases are induced by the coincident–limit inner product of $\hat{G}_\Lambda$; under local unitary rotations at an endpoint, $\operatorname{Im} \vec{V}_{\mathrm{vac}}^{[\mathbf{W}]}$ transforms orthogonally (unitarily) and ℓ^2–norms are preserved, so $t_{\mathrm{vac}}^{[\cdot]}$ is basis–independent and gauge/word–covariant in the same sense as $d_{\mathrm{vac}}^{[\cdot]}$ (Proposition 4.469).*

(iii) Duality bound. *The inequality (4.250) holds componentwise by $|\operatorname{Im} z| \le |z|$.*

(iv) Finiteness (coherent regularisation). *Under Principle 4.2 and the regularity hypotheses used for $d_{\mathrm{vac}}^{[\cdot]}$, one has $t_{\mathrm{vac}}^{[\cdot]}(x,x') < \infty$ for all x, x'.*

(v) Relational null criterion. $t_{\mathrm{vac}}^{[\cdot]}(x,x') = 0$ *iff* $\operatorname{Im} C_n^{[\mathbf{W}_1]}(x) = \operatorname{Im} C_n^{[\mathbf{W}_2]}(x')$ *for all n. Together with the vanishing of the other temporal/spatial components, this contributes to the relational null condition characterising a PQS locus.*

Proof. Write $I_n^{[\mathbf{W}]}(x) := \operatorname{Im} C_n^{[\mathbf{W}]}(x)$ and $\operatorname{Im} \vec{V}_{\mathrm{vac}}^{[\mathbf{W}]}(x) := (I_1^{[\mathbf{W}]}(x), I_2^{[\mathbf{W}]}(x), \dots)$ as in Definition 4.539. For notational simplicity, assume profiles at x' have been transported to x by the wordwise transporter (Definition 4.468); this does not affect any norm by the covariance arguments below.

(i) Nonnegativity, symmetry, triangle. By definition (eq. (4.249)),

$$t_{\mathrm{vac}}^{[\cdot]}(x,x') = \left\| \operatorname{Im} \vec{V}_{\mathrm{vac}}^{[\mathbf{W}_1]}(x) - \operatorname{Im} \vec{V}_{\mathrm{vac}}^{[\mathbf{W}_2]}(x') \right\|_{\ell^2} \;\ge\; 0.$$

Symmetry follows since exchanging $x \leftrightarrow x'$ sends the difference vector to its negative, leaving the ℓ^2–norm unchanged. The triangle inequality is just Minkowski's inequality in ℓ^2:

$$\begin{aligned} t_{\mathrm{vac}}^{[\cdot]}(x,x'') &= \left\| \operatorname{Im} \vec{V}_{\mathrm{vac}}^{[\mathbf{W}_1]}(x) - \operatorname{Im} \vec{V}_{\mathrm{vac}}^{[\mathbf{W}_3]}(x'') \right\|_{\ell^2} \\ &\le \left\| \operatorname{Im} \vec{V}_{\mathrm{vac}}^{[\mathbf{W}_1]}(x) - \operatorname{Im} \vec{V}_{\mathrm{vac}}^{[\mathbf{W}_2]}(x') \right\|_{\ell^2} + \left\| \operatorname{Im} \vec{V}_{\mathrm{vac}}^{[\mathbf{W}_2]}(x') - \operatorname{Im} \vec{V}_{\mathrm{vac}}^{[\mathbf{W}_3]}(x'') \right\|_{\ell^2} \\ &= t_{\mathrm{vac}}^{[\cdot]}(x,x') + t_{\mathrm{vac}}^{[\cdot]}(x',x''). \end{aligned}$$

Hence $t_{\mathrm{vac}}^{[\cdot]}$ is a pseudometric; it is a metric on any sector where $x \mapsto \operatorname{Im} \vec{V}_{\mathrm{vac}}^{[\mathbf{W}]}(x)$ is injective.

(ii) Word/gauge covariance and basis–independence. Under a local unitary $U(x)$ at the endpoint x, the wordwise frame projectors transform by conjugation, $\Pi_n^{(x)[\mathbf{W}]} \mapsto U(x)\Pi_n^{(x)[\mathbf{W}]}U(x)^\dagger$,

and the diagonal operator $A_x := \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_x)$ transforms covariantly. Consequently the band components $C_n^{[\mathbf{W}]}(x) = \langle \Psi | \Pi_n^{(x)[\mathbf{W}]} A_x \Pi_n^{(x)[\mathbf{W}]} | \Psi \rangle$ mix among (possibly degenerate) bands by a *unitary* matrix on the modal index. Taking imaginary parts is a real-linear operation, so $\operatorname{Im} \vec{V}_{\mathrm{vac}}^{[\mathbf{W}]}(x)$ undergoes a unitary (hence orthogonal) change of coordinates in ℓ^2, which leaves its norm invariant. The same holds at x' and under the compatible wordwise transport (Section 4.5). Therefore $t_{\mathrm{vac}}^{[\cdot]}$ is invariant under endpoint word/gauge changes and independent of the chosen (compatible) modal basis, in the same sense as $d_{\mathrm{vac}}^{[\cdot]}$ (Proposition 4.469).

(iii) Duality bound. For any complex numbers z_n one has $|\operatorname{Im} z_n| \le |z_n|$. Apply this to $z_n := C_n^{[\mathbf{W}_1]}(x) - C_n^{[\mathbf{W}_2]}(x')$ to obtain, componentwise, $|I_n^{[\mathbf{W}_1]}(x) - I_n^{[\mathbf{W}_2]}(x')| \le |z_n|$. Squaring, summing over n, and taking the square root yields

$$t_{\mathrm{vac}}^{[\cdot]}(x,x') = \left\| \left(I_n^{[\mathbf{W}_1]}(x) - I_n^{[\mathbf{W}_2]}(x') \right)_n \right\|_{\ell^2} \le \left\| \left(C_n^{[\mathbf{W}_1]}(x) - C_n^{[\mathbf{W}_2]}(x') \right)_n \right\|_{\ell^2} = d_{\mathrm{vac}}^{[\cdot]}(x,x'),$$

which is precisely eq. (4.250).

(iv) Finiteness. By (iii), $t_{\mathrm{vac}}^{[\cdot]}(x,x') \le d_{\mathrm{vac}}^{[\cdot]}(x,x')$. Under the Wordwise Finite Entanglement Capacity (Principle 4.2) and the regularity hypotheses used to define and bound $d_{\mathrm{vac}}^{[\cdot]}$ (cf. Definition 4.468 and Proposition 4.471), the series for $d_{\mathrm{vac}}^{[\cdot]}$ converges; hence $t_{\mathrm{vac}}^{[\cdot]}(x,x') < \infty$ for all x, x'.

(v) Relational null criterion. By definition, $t_{\mathrm{vac}}^{[\cdot]}(x,x') = 0$ iff the ℓ^2–norm of the difference of imaginary–part profiles vanishes, which is equivalent to $I_n^{[\mathbf{W}_1]}(x) = I_n^{[\mathbf{W}_2]}(x')$ for every n, i.e. $\operatorname{Im} C_n^{[\mathbf{W}_1]}(x) = \operatorname{Im} C_n^{[\mathbf{W}_2]}(x')$ for all n. Conversely, if these equalities hold bandwise, the difference vector is zero and $t_{\mathrm{vac}}^{[\cdot]}(x,x') = 0$. When combined with the vanishing of the other components of the (wordwise) temporal/spatial triads in the appropriate propositions, this condition contributes to the relational null characterization of a PQS locus (Section 4.11.8). □

Rate form along a path

Definition 4.541 (Wordwise vacuum–response rate 1–form)**.** Let $\gamma : [0,1] \to \mathcal{X}_q$ be a C^1 path and identify modal fibres along γ by the wordwise parallel transport of Section 4.5. Define

$$\boxed{ \dot{t}_{\mathrm{vac}}^{[\cdot]}(s) := \left\| \frac{\mathrm{d}}{\mathrm{d}s} \operatorname{Im} \vec{V}_{\mathrm{vac}}^{[\mathbf{W}(s)]}(\gamma(s)) \right\|_{\ell^2}, \qquad T_{\mathrm{vac}}^{[\cdot]}(x,x';\gamma) := \int_0^1 \dot{t}_{\mathrm{vac}}^{[\cdot]}(s)\, \mathrm{d}s. } \tag{4.251}$$

Then $t_{\mathrm{vac}}^{[\cdot]}(x,x') \le T_{\mathrm{vac}}^{[\cdot]}(x,x';\gamma)$ with equality whenever the image $s \mapsto \operatorname{Im} \vec{V}_{\mathrm{vac}}^{[\cdot]}(\gamma(s))$ is a straight segment in profile space (chord–length equality in ℓ^2). The construction is basis–independent by the covariance of $\hat{G}_\Lambda$.

Remark 4.542 (Placement in the temporal fibre and calibration (wordwise))**.** The temporal fibre $\mathcal{F}_x^{\mathrm{time}}$ houses the wordwise temporal triad $(t_{\mathrm{corr}}^{[\cdot]}, t_{\mathrm{vac}}^{[\cdot]}, t_{\mathrm{spec}}^{[\cdot]})$, dual to the wordwise distance fibre $\mathcal{F}_x^{\mathrm{w}}$ (Definition 4.490). As with $t_{\mathrm{corr}}^{[\cdot]}$, $t_{\mathrm{vac}}^{[\cdot]}$ is dimensionless; if a dimensionful parameter is desired one introduces a state–dependent calibration scale τ_* and sets $T_{\mathrm{vac,abs}}^{[\cdot]} := \tau_*\, t_{\mathrm{vac}}^{[\cdot]}$ (to be fixed together with the emergent proper–time discussion).

Physical reading (wordwise). $t_{\mathrm{vac}}^{[\cdot]}$ measures the *retarded internal reaction* of the vacuum to modal shifts, sensitive to phase–encoded polarisation and word/gauge flow in the coincident sector. It is the operator–theoretic, background–free analogue of phase response familiar from quantum settings, here rendered intrinsic via the Holo–State and the diagonal sector of $\hat{G}_\Lambda$, and regularised coherently by the same wordwise mechanisms that render the distances finite.

4.12.5 Spectral Shear Time

The third emergent temporal quantity, $t_{\mathrm{spec}}(x, x')$, arises from the *non-commutativity* between the differential structure generator $\hat{D}$ and the modal relational structure defined by the Fourfold Inter-Modal Correlator $\hat{G}_\Lambda$. It captures the *imaginary deformation* induced by spectral flow across relational configurations (Definitions 4.386 and 4.402; cf. [4, 392, 407]).

Formal Definition

Definition 4.543 (Spectral Shear Time)**.** Let the *spectral-shear observables* be

$$\mathcal{S}_n(x, x') \;:=\; \Big\langle \mathring{\mathbf{\Psi}} \,\Big|\, [\hat{D},\ \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_n^*, \psi_n]] \,\Big|\, \mathring{\mathbf{\Psi}} \Big\rangle,$$

as in Definition 4.386, and let $w_n(x, x')$ be the modal weights of Definition 4.392. Define $S_n^{(\mathrm{Im})}(x, x') := \mathrm{Im}\ \mathcal{S}_n(x, x')$ and the vector

$$\mathrm{Im}\,\vec{S}(x, x') \;:=\; \Big(w_1(x, x')\, S_1^{(\mathrm{Im})}(x, x'),\ w_2(x, x')\, S_2^{(\mathrm{Im})}(x, x'),\ \dots\Big).$$

The *spectral shear time* is the ℓ^2–norm

$$\boxed{t_{\mathrm{spec}}(x, x') \;:=\; \Big[\sum\nolimits_n w_n^2(x, x') \Big(\mathrm{Im}\ \mathcal{S}_n(x, x')\Big)^2\Big]^{1/2} \;=\; \Big\|\mathrm{Im}\,\vec{S}(x, x')\Big\|_2.} \tag{4.252}$$

Equivalently,

$$t_{\mathrm{spec}}(x, x') = \Big[\sum\nolimits_n w_n^2(x, x') \Big(\mathrm{Im} \Big\langle \mathring{\mathbf{\Psi}} \Big|\, [\hat{D},\ \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_n^*, \psi_n]] \,\Big|\, \mathring{\mathbf{\Psi}} \Big\rangle\Big)^2\Big]^{1/2}.$$

Interpretation and Duality. t_{spec} is the *phase–time dual* of the spectral distance d_{spec} (Definition 4.402): the latter aggregates the *magnitudes* $|\mathcal{S}_n|$, while the former aggregates the *imaginary parts* $\mathrm{Im}\ \mathcal{S}_n$, with identical weights w_n. Since $|\mathrm{Im}\, z| \le |z|$ pointwise, we obtain the bound

$$\boxed{t_{\mathrm{spec}}(x, x') \;\le\; d_{\mathrm{spec}}(x, x') \quad \text{for all } x, x' \in \mathcal{X}_q,} \tag{4.253}$$

with equality iff each $\mathcal{S}_n(x, x')$ is purely imaginary.

Basic Properties

Proposition 4.544 (Basic Properties of t_{spec})**.** *Fix $x, x', x'' \in \mathcal{X}_q$. Then:*

(i) Nonnegativity, symmetry, triangle. $t_{\mathrm{spec}}(x,x') \geq 0$, $t_{\mathrm{spec}}(x,x') = t_{\mathrm{spec}}(x',x)$, *and*

$$t_{\mathrm{spec}}(x,x'') \;\leq\; t_{\mathrm{spec}}(x,x') + t_{\mathrm{spec}}(x',x'').$$

Thus t_{spec} is a pseudometric; it is a metric on sectors where $x \mapsto \operatorname{Im}\vec{S}(x,\cdot)$ separates points.

(ii) Gauge/frame covariance and basis–independence. *Under unitary rotations of the modal basis and gauge changes of the Fourfold identification along fibres, $\operatorname{Im}\vec{S}$ transforms orthogonally and its ℓ^2–norm is invariant; hence t_{spec} is basis–independent and gauge–covariant, matching d_{spec}.*

(iii) Duality bound. *The inequality (4.253) follows from $|\operatorname{Im} z| \leq |z|$ componentwise.*

(iv) Finiteness (coherent regularisation). *Under Spectral Smoothness and modal weight decay (Axiom 4.415, Definition 4.392, and Proposition 4.424), the RMS sum in (4.252) is finite; in particular $t_{\mathrm{spec}}(x,x') < \infty$ for all x, x'.*

(v) Relational null criterion. $t_{\mathrm{spec}}(x,x') = 0$ *iff* $\operatorname{Im}\mathcal{S}_n(x,x') = 0$ *for all n. Together with the vanishing of the other temporal components, this contributes to the PQS* relational null *condition $(\vec{d},\vec{t}) = \mathbf{0}$ (Definition 4.604).*

Rate Form along a Path

Definition 4.545 (Spectral Shear Rate 1-Form)**.** Let $\gamma : [0,1] \to \mathcal{X}_q$ be C^1 and use the gauge–covariant identification of modal fibres along γ (parallel transport as in the Fourfold construction). Set

$$\dot{t}_{\mathrm{spec}}(s) \;:=\; \Big[\sum\nolimits_n \Big(\tfrac{\mathrm{d}}{\mathrm{d}s}\Big(w_n(\gamma(s),x')\,\operatorname{Im}\mathcal{S}_n(\gamma(s),x')\Big)\Big)^2\Big]^{1/2},$$

and define the accumulated spectral time along γ by

$$\boxed{T_{\mathrm{spec}}(x,x';\gamma) \;:=\; \int_0^1 \dot{t}_{\mathrm{spec}}(s)\,\mathrm{d}s.} \tag{4.254}$$

Then $t_{\mathrm{spec}}(x,x') \leq T_{\mathrm{spec}}(x,x';\gamma)$, with equality whenever $s \mapsto \operatorname{Im}\vec{S}(\gamma(s),x')$ traces a straight segment in profile space (chord–length equality in ℓ^2). The construction is basis–independent by the same covariance that governs $\hat{G}_\Lambda$.

Remark 4.546 (Placement in the Temporal Fibre and Calibration)**.** The temporal flow fibre $\mathcal{F}_x^{\mathrm{time}}$ houses the triad $(t_{\mathrm{corr}}, t_{\mathrm{vac}}, t_{\mathrm{spec}})$, dual to the distance fibre $\mathcal{F}_x^{\mathrm{dist}}$ (Definitions 4.599, 4.600 and 4.604). As with the other components, t_{spec} is dimensionless; a state–dependent calibration τ_* may be introduced where a dimensionful scale is desired.

Physical Reading. t_{spec} measures the *imaginary* component of the commutator-induced spectral deformation, quantifying a non-unitary "temporal tension'' within the operator algebra. In geometric terms, it is the phase-sensitive part of spectral shear, analogous—in the noncommutative setting—to a Berry-curvature–like response of the relational bundle under adiabatic spectral transport ([407, 392]; cf. [4]).

Time	Mathematical Feature	Operator Type
t_{corr}	$\arg \left\langle \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'}) \right\rangle$	off–diagonal (bilocal) phase
t_{vac}	$\left\| \operatorname{Im} \vec{V}_{\text{vac}}(x) - \operatorname{Im} \vec{V}_{\text{vac}}(x') \right\|_2$	diagonal coincident (self) drift
t_{spec}	$\left(\sum_n w_n^2 \left(\operatorname{Im} \mathcal{S}_n(x, x') \right)^2 \right)^{1/2}$	commutator–induced spectral deformation

Table 4.5: Classification of emergent time observables by mathematical feature and operator type. For t_{vac}, let $\vec{V}_{\text{vac}}(x)$ denote the vector of diagonal self–correlators and use the componentwise imaginary part, as in Definitions 4.463 and 4.535 and eq. (4.211). For t_{spec}, $\mathcal{S}_n$ are the spectral–shear observables with weights w_n (Definitions 4.386, 4.392 and 4.543).

Why the Temporal Observables are "Imaginary" Components

(1) What "imaginary" refers to. Throughout, the qualifier "imaginary'' indicates *which part of a complex operator expectation we extract*, not that the resulting observable is itself a complex number. Concretely,

$$t_{\text{corr}},\ t_{\text{vac}},\ t_{\text{spec}} \quad \text{are built from} \quad \operatorname{Im}\langle \overset{\circ}{\mathbf{\Psi}} | \hat{\mathcal{O}} | \overset{\circ}{\mathbf{\Psi}} \rangle,$$

with the relevant operators $\hat{\mathcal{O}}$ given by the bilocal channel, the diagonal (coincident) channel, and the commutator with $\hat{D}$, respectively (see Definitions 4.535 and 4.543 and eqs. (4.246) and (4.252)). In quantum kinematics, complex phases are physically meaningful: relative phase is observable and governs interference, and imaginary parts of response functions carry definite operational content [350, 408, 82].

(2) The temporal observables are real-valued. Although they *use* imaginary parts, each time observable is a norm of an ℓ^2-vector of such components, hence a nonnegative real number:

$$t_{\text{vac}}(x, x') = \left\| \operatorname{Im} \vec{V}_{\text{vac}}(x) - \operatorname{Im} \vec{V}_{\text{vac}}(x') \right\|_2, \qquad t_{\text{spec}}(x, x') = \left\| \operatorname{Im} \vec{S}(x, x') \right\|_2,$$

and similarly for t_{corr} (phase aggregate of the bilocal expectation). Thus "imaginary'' here is a *selection rule* on complex data, not a statement that "time is imaginary.''

(3) Physical meaning: causality, dissipation, and spectral density. In linear and quantum response theory, the imaginary part of a causal (retarded) Green function or susceptibility captures the *absorptive* content—i.e. the part that changes under time-reversal and encodes irreversible response; equivalently, it determines the spectral density via the optical theorem/Källén–Lehmann structure [82, 230, 241]. Our constructions mirror this logic:

- t_{vac} aggregates imaginary self-variations of coincident channels, measuring the vacuum's *retarded internal reaction* to modal displacements (cf. vacuum polarisation in curved backgrounds [244, 11, 13]).
- t_{spec} aggregates imaginary parts of the commutator expectation $[\hat{D}, \hat{G}_\Lambda]$, i.e. the *phase-sensitive* piece of spectral shear. In spectral geometry and noncommutative settings, commutators with the Dirac-type generator detect spectral flow and curvature; their imaginary parts track phase-dislocating contributions (cf. [4, 394]).

Thus the "imaginary'' selection isolates precisely the sector tied to causal/absorptive effects and phase transport.

(4) Not Euclidean (Wick-rotated) "imaginary time." Nothing here asserts a Wick rotation $t \mapsto -\mathrm{i}\tau$. Euclidean (imaginary) time is a calculational device that trades oscillatory unitary evolution for decaying kernels; by contrast, our $t_{\mathrm{corr}}, t_{\mathrm{vac}}, t_{\mathrm{spec}}$ are *real, emergent functionals* of Im parts of complex expectations in the original (Lorentzian) theory. They do not presume a background time coordinate and do not identify "physical time'' with $\mathrm{i} \times$ (something) [409, 82].

(5) Why the imaginary sector carries "temporal" content. Two complementary reasons justify using Im to define the temporal fibre (Remark 4.546):

(a) *Interference/phase sensitivity.* Temporal ordering in quantum theory is operationally probed by phase-sensitive interferometry; phases (arguments) and imaginary parts of overlaps govern fringe shifts and time-delay–like observables [408, 350]. Our t_{corr} generalises this by aggregating phase across bilocal channels.

(b) *Retarded response.* The imaginary part of a retarded kernel changes sign under reversal and vanishes in the absence of dissipative/absorptive features; it is therefore the natural carrier of directed "flow'' (time's arrow) within an operator-algebraic setting [230, 241, 82]. The definitions of t_{vac} and t_{spec} implement this selection intrinsically.

(6) Reality and duality checks. By construction,

$$0 \le t_{\mathrm{corr}},\ t_{\mathrm{vac}},\ t_{\mathrm{spec}} \ \le\ d_{\mathrm{corr}},\ d_{\mathrm{vac}},\ d_{\mathrm{spec}},$$

since $|\operatorname{Im} z| \le |z|$ componentwise (see eqs. (4.247) and (4.253)). This ties each temporal axis to its spatial dual and guarantees that "imaginary'' time components are *bounded, real* functionals of the same operator data that define the spatial distances.

Bottom line. In this framework, calling $t_{\mathrm{corr}}, t_{\mathrm{vac}}, t_{\mathrm{spec}}$ "imaginary'' means they are *phase–response functionals*: real, gauge-covariant aggregates of the Im parts of complex Holo-State expectations that encode causal/absorptive content and phase transport. They are as "real'' as any normed observable constructed from expectation values, while remaining cleanly distinguished from Euclidean (Wick-rotated) imaginary time.

4.12.6 Wordwise Spectral Shear Time

The third emergent temporal quantity, $t_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')$, arises from the *non-commutativity* between the wordwise differential generator $\mathbb{D}^{[\mathbf{W}_1]}(x)$ and the modal relational structure encoded by the Fourfold correlator $\hat{G}_\Lambda$. It captures the *imaginary deformation* induced by spectral flow across relational configurations (Definitions 4.406 and 4.408).

Formal definition (wordwise)

Definition 4.547 (Wordwise spectral shear time)**.** Let the *wordwise spectral–shear observables* be

$$\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') := \left\langle \mathring{\mathbf{\Psi}} \,\middle|\, \left[\mathbb{D}^{[\mathbf{W}_1]}(x),\ \hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})[\psi_n^{(x)*},\psi_n^{(x')}]\right] \,\middle|\, \mathring{\mathbf{\Psi}}\right\rangle, \quad \text{(cf. Definition 4.406)}$$

and let $w_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')$ be the wordwise modal weights (Definition 4.407). Define $S_n^{(\mathrm{Im})[\mathbf{W}_1,\mathbf{W}_2]}(x,x') := \mathrm{Im}\ \mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')$ and the weighted vector

$$\mathrm{Im}\,\vec{S}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') := \left(w_1^{[\mathbf{W}_1,\mathbf{W}_2]}\, S_1^{(\mathrm{Im})[\mathbf{W}_1,\mathbf{W}_2]},\ w_2^{[\mathbf{W}_1,\mathbf{W}_2]}\, S_2^{(\mathrm{Im})[\mathbf{W}_1,\mathbf{W}_2]},\ \dots\right).$$

The *wordwise spectral shear time* is the ℓ^2–norm

$$\boxed{\begin{aligned} t_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') &:= \left[\sum_n \left(w_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\right)^2 \left(\mathrm{Im}\ \mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\right)^2\right]^{1/2} \\ &= \left\|\mathrm{Im}\,\vec{S}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\right\|_{\ell^2}. \end{aligned}} \tag{4.255}$$

Interpretation and duality (wordwise). $t_{\mathrm{spec}}^{[\cdot]}$ is the *phase–time dual* of the wordwise spectral distance $d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}$ (Definition 4.408): the latter aggregates magnitudes $|\mathcal{S}_n^{[\cdot]}|$, while the former aggregates the imaginary parts $\mathrm{Im}\ \mathcal{S}_n^{[\cdot]}$ with identical weights. Since $|\mathrm{Im}\, z| \le |z|$,

$$\boxed{t_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \ \le\ d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \quad \text{for all } x,x' \in \mathcal{X}_q,} \tag{4.256}$$

with equality iff each $\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')$ is purely imaginary.

Basic properties

Proposition 4.548 (Basic properties of $t_{\mathrm{spec}}^{[\cdot]}$)**.** *For any $x,x',x'' \in \mathcal{X}_q$ and admissible words:*

(i) Nonnegativity, symmetry, triangle. $t_{\mathrm{spec}}^{[\cdot]}(x,x') \ge 0$, $t_{\mathrm{spec}}^{[\cdot]}(x,x') = t_{\mathrm{spec}}^{[\cdot]}(x',x)$, *and*

$$t_{\mathrm{spec}}^{[\cdot]}(x,x'') \ \le\ t_{\mathrm{spec}}^{[\cdot]}(x,x') + t_{\mathrm{spec}}^{[\cdot]}(x',x'').$$

Thus $t_{\mathrm{spec}}^{[\cdot]}$ is a pseudometric; it is a metric on sectors where $x \mapsto \mathrm{Im}\,\vec{S}^{[\cdot]}(x,\cdot)$ separates points.

(ii) Word/gauge covariance and basis–independence. *Under endpoint word rotations and gauge changes of the Fourfold identification, each $\mathcal{S}_n^{[\cdot]}$ transforms by conjugation and* $\operatorname{Im}\vec{S}^{[\cdot]}$ *undergoes an orthogonal transform in band space; the ℓ^2–norm is invariant. Hence $t_{\mathrm{spec}}^{[\cdot]}$ is basis–independent and word/gauge–covariant, matching $d_{\mathrm{spec}}^{[\cdot]}$.*

(iii) Duality bound. *One has*

$$t_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \;\le\; d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x'), \tag{4.257}$$

with equality iff every $\mathcal{S}_n^{[\cdot]}(x,x')$ is purely imaginary.

(iv) Finiteness (coherent regularisation). *Under Wordwise Spectral Smoothness and finite capacity (Theorem 4.421 and Principle 4.2), the ℓ^2 sum in eq. (4.255) is finite; in particular $t_{\mathrm{spec}}^{[\cdot]}(x,x') < \infty$.*

(v) Relational null criterion. $t_{\mathrm{spec}}^{[\cdot]}(x,x') = 0$ *iff* $\operatorname{Im}\mathcal{S}_n^{[\cdot]}(x,x') = 0$ *for all bands in the Holo-State effective support (equivalently, for all n with $w_n^{[\cdot]}(x,x') > 0$). In sectors where all relevant bands are supported, this is "for all n". Together with the vanishing of the other temporal components it contributes to the PQS relational null condition.*

Proof. Recall the definition (Definition 4.547)

$$t_{\mathrm{spec}}^{[\cdot]}(x,x') = \left\| \operatorname{Im}\vec{S}^{[\cdot]}(x,x') \right\|_{\ell^2} = \left[\sum_n \left(w_n^{[\cdot]}(x,x') \right)^2 \left(\operatorname{Im}\, \mathcal{S}_n^{[\cdot]}(x,x') \right)^2 \right]^{1/2},$$

with $\mathcal{S}_n^{[\cdot]}$ the wordwise spectral–shear observables (Definition 4.406) and weights $w_n^{[\cdot]}$ (Definition 4.407).

(i) Nonnegativity, symmetry, triangle. Nonnegativity is immediate from the ℓ^2 norm. For symmetry, exchange $(x,\mathbf{W}_1) \leftrightarrow (x',\mathbf{W}_2)$ and reverse the comparison path; then

$$\mathcal{U}^{[\mathbf{W}_2,\mathbf{W}_1]}(x' \leftarrow x;\bar{\gamma}) = \mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x \leftarrow x';\gamma)^\dagger \quad \text{and} \quad \hat{G}_\Lambda(\mathfrak{B}_{x'},\mathfrak{B}_x)[\cdot,\cdot] = \hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})[\cdot,\cdot]^\dagger.$$

Using that commutators are anti-Hermitian and Holo-State expectations satisfy $\langle A^\dagger\rangle_\Psi = \langle A\rangle_\Psi^*$, one checks

$$\mathcal{S}_n^{[\mathbf{W}_2,\mathbf{W}_1]}(x',x) \;=\; -\left(\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\right)^* \quad\Longrightarrow\quad \operatorname{Im}\mathcal{S}_n^{[\mathbf{W}_2,\mathbf{W}_1]}(x',x) = \operatorname{Im}\mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x').$$

Weights are symmetric by construction, $w_n^{[\mathbf{W}_2,\mathbf{W}_1]}(x',x) = w_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')$ (Definition 4.407), hence the ℓ^2 norm is unchanged: $t_{\mathrm{spec}}^{[\cdot]}(x',x) = t_{\mathrm{spec}}^{[\cdot]}(x,x')$. For the triangle inequality, Minkowski in ℓ^2 yields

$$\left\| \operatorname{Im}\vec{S}^{[\cdot]}(x,x'') \right\|_{\ell^2} \le \left\| \operatorname{Im}\vec{S}^{[\cdot]}(x,x') \right\|_{\ell^2} + \left\| \operatorname{Im}\vec{S}^{[\cdot]}(x',x'') \right\|_{\ell^2}.$$

(ii) Word/gauge covariance and basis–independence. At an endpoint $y \in \{x,x'\}$, a local unitary $U(y)$ sends $\mathbb{D}^{[\mathbf{W}]}(y) \mapsto U(y)\mathbb{D}^{[\mathbf{W}]}(y)U(y)^\dagger$ and $\hat{G}_\Lambda \mapsto U(x)\hat{G}_\Lambda U(x')^\dagger$ in the bilocal slots (Section 4.5). Therefore the commutator inside $\mathcal{S}_n^{[\cdot]}$ transforms by conjugation, and its Holo-State expectation is invariant up to a unitary mixing among degenerate bands (change of modal frame). Taking imaginary parts is real-linear; thus the band-index vector $\operatorname{Im}\vec{S}^{[\cdot]}$

undergoes an orthogonal transform in ℓ^2, preserving its norm. Weights $w_n^{[\cdot]}$ are gauge scalars built from normalized overlaps (Definition 4.407) and are unchanged. Hence $t_{\mathrm{spec}}^{[\cdot]}$ is invariant under endpoint word/gauge changes and independent of the choice of compatible modal basis, matching the covariance properties of $d_{\mathrm{spec}}^{[\cdot]}$ (Definition 4.408).

(iii) Duality bound. For every band n, $|\operatorname{Im} z| \le |z|$ with $z = \mathcal{S}_n^{[\cdot]}(x,x')$. Multiplying by the nonnegative $w_n^{[\cdot]}(x,x')$ and taking the ℓ^2 norm over n gives $t_{\mathrm{spec}}^{[\cdot]}(x,x') \le d_{\mathrm{spec}}^{[\cdot]}(x,x')$ as in eq. (4.257). Equality holds iff each $\mathcal{S}_n^{[\cdot]}$ is purely imaginary.

(iv) Finiteness. From Wordwise Spectral Smoothness (Theorem 4.421), $|\mathcal{S}_n^{[\cdot]}(x,x')| \le C\,\lambda_n^k$ for some $k \in \mathbb{N}$ on each local slice; finite capacity yields $w_n^{[\cdot]}(x,x') \le C'\lambda_n^{-\alpha}$ with $\alpha > k + \frac{1}{2}$ (Principle 4.2). Hence

$$\sum_n \left(w_n^{[\cdot]}\right)^2 \left(\operatorname{Im}\,\mathcal{S}_n^{[\cdot]}\right)^2 \;\le\; C^2C'^2 \sum_n \lambda_n^{2k-2\alpha}$$

which converges by the assumed exponent (or, more generally, by combining with a Weyl-type counting bound as in Proposition 4.460). Therefore $t_{\mathrm{spec}}^{[\cdot]}(x,x') < \infty$. Alternatively, eq. (4.257) and the finiteness of $d_{\mathrm{spec}}^{[\cdot]}$ (Definition 4.408 together with coherent regularization) also imply finiteness.

(v) Relational null criterion. Since weights are nonnegative, the ℓ^2 norm vanishes iff every weighted component vanishes. Thus $t_{\mathrm{spec}}^{[\cdot]}(x,x') = 0$ iff $w_n^{[\cdot]}(x,x')\ \operatorname{Im}\mathcal{S}_n^{[\cdot]}(x,x') = 0$ for all n, i.e. $\operatorname{Im}\mathcal{S}_n^{[\cdot]}(x,x') = 0$ for all bands in the effective Holo-State support (Definition 4.451). On sectors where all relevant bands have nonzero weight, this is simply $\operatorname{Im}\mathcal{S}_n^{[\cdot]}(x,x') = 0$ for all n. In conjunction with the corresponding criteria for the other temporal and spatial components, this realizes the relational null condition that characterizes PQS loci (Section 4.11.8). □

Rate form along a path (wordwise)

Definition 4.549 (Wordwise spectral–shear rate 1–form)**.** Let $\gamma : [0,1] \to \mathcal{X}_q$ be C^1 and transport modal fibres along γ by the wordwise parallel transport (Section 4.5). Define

$$\boxed{\dot{t}_{\mathrm{spec}}^{[\cdot]}(s) := \left\| \frac{\mathrm{d}}{\mathrm{d}s}\,\operatorname{Im}\vec{S}^{[\mathbf{W}_1,\mathbf{W}_2]}(\gamma(s),x') \right\|_{\ell^2}, \qquad T_{\mathrm{spec}}^{[\cdot]}(x,x';\gamma) := \int_0^1 \dot{t}_{\mathrm{spec}}^{[\cdot]}(s)\,\mathrm{d}s.} \tag{4.258}$$

Then $t_{\mathrm{spec}}^{[\cdot]}(x,x') \le T_{\mathrm{spec}}^{[\cdot]}(x,x';\gamma)$, with equality when $s \mapsto \operatorname{Im}\vec{S}^{[\cdot]}(\gamma(s),x')$ traces a straight segment in profile space (chord–length equality in ℓ^2).

Remark 4.550 (Placement in the temporal fibre and calibration)**.** The temporal fibre $\mathcal{F}_x^{\mathrm{time}}$ houses the wordwise temporal triad $\left(t_{\mathrm{corr}}^{[\cdot]}, t_{\mathrm{vac}}^{[\cdot]}, t_{\mathrm{spec}}^{[\cdot]}\right)$, dual to the wordwise distance fibre (Definition 4.490). As with the other components, $t_{\mathrm{spec}}^{[\cdot]}$ is dimensionless; a state–dependent calibration τ_* may be introduced to obtain a dimensionful accumulated time $T_{\mathrm{spec,abs}}^{[\cdot]} := \tau_*\, t_{\mathrm{spec}}^{[\cdot]}$.

Physical reading (wordwise). $t_{\mathrm{spec}}^{[\cdot]}$ measures the *imaginary* component of commutator–induced spectral deformation—i.e., a phase–sensitive "temporal tension" of the relational

bundle generated by $\mathbb{D}^{[\mathbf{W}]}$ acting on Fourfold channels. Geometrically, it is the phase–sensitive part of spectral shear, the noncommutative analogue of a curvature–driven response of wordwise modes under adiabatic spectral transport, and it is coherently regularised by the same endogenous mechanisms that make $d_{\mathrm{spec}}^{[\cdot]}$ finite.

4.12.7 Functional Independence of Temporal Observables

The orthogonality of the three temporal measures stems from the algebraic and functional independence of the operator structures that generate them. Each temporal component probes a distinct, non–interchangeable aspect of the quantum dynamics encoded by the Fourfold Inter–Modal Correlator $\hat{G}_\Lambda$ and the spectral generator $\hat{D}$ within the Holo–State $|\mathring{\mathbf{\Psi}}\rangle$. We make this precise by a Fréchet–differential argument mirroring the spatial case (Proposition 4.497).

Setup and notation. Let $\mathfrak{G}$ denote the admissible class of Fourfold correlators (Definition 4.9) and $\mathfrak{D}$ the admissible class of spectral operators (Definition 4.343). Write

$$\mathcal{M} := \mathfrak{G} \times \mathfrak{D}, \qquad \mathbf{T} : \mathcal{M} \to \mathbb{R}^3, \qquad \mathbf{T}(\hat{G}_\Lambda, \hat{D}) := \Big(t_{\mathrm{corr}}, t_{\mathrm{vac}}, t_{\mathrm{spec}}\Big),$$

with $t_{\mathrm{corr}}, t_{\mathrm{vac}}, t_{\mathrm{spec}}$ as in eqs. (4.243), (4.246) and (4.252) and Definitions 4.535 and 4.543. Variations $(\delta\hat{G}, \delta\hat{D}) \in T_{(\hat{G},\hat{D})}\mathcal{M}$ act at fixed Holo–State. We use the abbreviations

$$F(x,x') := \Big\langle \mathring{\mathbf{\Psi}} \Big| \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi^*, \psi] \Big| \mathring{\mathbf{\Psi}} \Big\rangle, \quad \Delta_k(x,x') := \Big\langle \mathring{\mathbf{\Psi}} \Big| \hat{O}_k(x) - \hat{O}_k(x') \Big| \mathring{\mathbf{\Psi}} \Big\rangle,$$

$$\mathcal{S}_n(x,x') := \Big\langle \mathring{\mathbf{\Psi}} \Big| \, [\hat{D},\ \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_n^*, \psi_n]] \, \Big| \mathring{\mathbf{\Psi}} \Big\rangle,$$

and modal weights $w_n(x,x')$ (Definition 4.392).

Proposition 4.551 (Functional independence of triadic temporal observables)**.** *Assume the Standard Hypotheses (Finite Entanglement Capacity; Spectral/Relational Smoothness) and nondegeneracy at (x,x'): $F(x,x') \neq 0$ and at least one $w_n(x,x') \neq 0$. Then the mapping $\mathbf{T}$ has full rank 3 (as a Fréchet map) on an open dense subset of $\mathcal{M}$. Consequently, if an admissible smooth $f : \mathbb{R}^3 \to \mathbb{R}$ satisfies $f \circ \mathbf{T} \equiv 0$ for all (x,x') and all $(\hat{G}_\Lambda, \hat{D}) \in \mathcal{M}$, then $f \equiv 0$.*

Proof. We compute the Fréchet differential $D\mathbf{T}$ at a generic point.

(i) Differential of t_{corr}. Since $t_{\mathrm{corr}} = \arg F$ (eq. (4.243)), for $F \neq 0$,

$$\delta t_{\mathrm{corr}} = \mathrm{Im}\left(\frac{\delta F}{F}\right), \qquad \delta F = \Big\langle \mathring{\mathbf{\Psi}} \Big| \, \delta\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi^*, \psi] \, \Big| \mathring{\mathbf{\Psi}} \Big\rangle, \tag{4.259}$$

and $\partial t_{\mathrm{corr}}/\partial \hat{D} = 0$. Choosing a purely phase–shifting off–diagonal perturbation $\delta\hat{G}_{\mathrm{off}}$ supported at (x,x') with $\delta F \neq 0$ ensures $\delta t_{\mathrm{corr}} \neq 0$.

(ii) Differential of t_{vac}. Write $I_k := \mathrm{Im}\,\Delta_k$ and $t_{\mathrm{vac}} = \Big(\sum_k I_k^2\Big)^{1/2}$ (Definition 4.535 and eq. (4.246)). For $t_{\mathrm{vac}} > 0$,

$$\delta t_{\mathrm{vac}} = \frac{1}{t_{\mathrm{vac}}} \sum_k I_k \ \mathrm{Im}\ \delta\Delta_k, \qquad \delta\Delta_k = \Big\langle \mathring{\mathbf{\Psi}} \Big| \, \delta\hat{O}_k(x) - \delta\hat{O}_k(x') \, \Big| \mathring{\mathbf{\Psi}} \Big\rangle, \tag{4.260}$$

where $\delta\hat{O}_k$ is induced by the diagonal (coincident) component of $\delta\hat{G}_\Lambda$. In particular $\partial t_{\mathrm{vac}}/\partial\hat{D} = 0$. Choosing a coincident, imaginary diagonal perturbation $\delta\hat{G}_{\mathrm{diag}}$ with $\mathrm{Im}\,\delta\Delta_{k_0} \neq 0$ for some k_0 gives $\delta t_{\mathrm{vac}} \neq 0$ while $\delta t_{\mathrm{corr}} = 0$ by off/diag separation.

(iii) Differential of t_{spec}. With $S_n^{(\mathrm{Im})} := \mathrm{Im}\,\mathcal{S}_n$ and $t_{\mathrm{spec}} = (\sum_n w_n^2 (S_n^{(\mathrm{Im})})^2)^{1/2}$ (Definition 4.543 and eq. (4.252)), for $t_{\mathrm{spec}} > 0$,

$$\delta t_{\mathrm{spec}} = \frac{1}{t_{\mathrm{spec}}} \sum_n w_n^2 S_n^{(\mathrm{Im})}\ \mathrm{Im}\ \delta\mathcal{S}_n, \qquad \delta\mathcal{S}_n = \Big\langle \mathring{\mathbf{\Psi}} \Big|\, [\delta\hat{D}, \hat{G}_n] + [\hat{D}, \delta\hat{G}_n] \,\Big| \mathring{\mathbf{\Psi}} \Big\rangle, \tag{4.261}$$

where $\hat{G}_n := \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_n^*, \psi_n]$. Taking a *pure $\hat{D}$-variation*, $\delta\hat{D} \neq 0$ and $\delta\hat{G} = 0$, yields

$$\delta t_{\mathrm{spec}} = \frac{1}{t_{\mathrm{spec}}} \sum_n w_n^2 S_n^{(\mathrm{Im})}\ \mathrm{Im} \Big\langle \mathring{\mathbf{\Psi}} \Big|\, [\delta\hat{D}, \hat{G}_n] \,\Big| \mathring{\mathbf{\Psi}} \Big\rangle, \tag{4.262}$$

which is nonzero for a generic self–adjoint $\delta\hat{D}$ because at least one $w_n \neq 0$ and $\{\hat{G}_n\}_n$ separate directions in the commutator algebra (Tomita–Takesaki–type noncommutativity; cf. [381]). In particular, $\partial t_{\mathrm{spec}}/\partial\hat{D}$ is typically nonzero, whereas $\partial t_{\mathrm{corr}}/\partial\hat{D} = \partial t_{\mathrm{vac}}/\partial\hat{D} = 0$.

(iv) Rank computation. Restrict $D\mathbf{T}$ to the three–dimensional subspace spanned by the independent directions

$$v_1 = (\delta\hat{G}_{\mathrm{off}}, 0), \qquad v_2 = (\delta\hat{G}_{\mathrm{diag}}, 0), \qquad v_3 = (0, \delta\hat{D}),$$

chosen as above. Using (4.259)–(4.262) one gets the *triangular* structure

$$D\mathbf{T}\Big|_{\mathrm{span}\{v_1,v_2,v_3\}} = \begin{pmatrix} a & 0 & 0 \\ 0 & b & 0 \\ * & * & e \end{pmatrix}, \qquad a = \delta_{v_1} t_{\mathrm{corr}} \neq 0, \quad b = \delta_{v_2} t_{\mathrm{vac}} \neq 0, \quad e = \delta_{v_3} t_{\mathrm{spec}} \neq 0.$$

Hence $\det\Big(D\mathbf{T}\Big|\Big) = a\,b\,e \neq 0$, so $D\mathbf{T}$ has full rank 3 at the given point. By openness of the nonvanishing conditions (nonzero F, $t_{\mathrm{vac}}, t_{\mathrm{spec}}$ or their weighted components), this holds on an open dense subset of $\mathcal{M}$.

(v) Independence. If $f \circ \mathbf{T} \equiv 0$, differentiation gives $(\nabla f)(\mathbf{T}) \cdot D\mathbf{T} = 0$. Full rank of $D\mathbf{T}$ forces $\nabla f(\mathbf{T}) = 0$ on a set with accumulation points, whence $f \equiv 0$ by analyticity/smoothness. □

Remark 4.552 (Degenerate loci and squared functionals)**.** At special loci (e.g. PQS where $\vec{t} = \mathbf{0}$), first variations of norms can vanish. One then works with the squared functionals $T_{\mathrm{vac}} := t_{\mathrm{vac}}^2$ and $T_{\mathrm{spec}} := t_{\mathrm{spec}}^2$, whose first variations $\delta T_{\mathrm{vac}} = 2\sum_k I_k\ \mathrm{Im}\ \delta\Delta_k$, $\delta T_{\mathrm{spec}} = 2\sum_n w_n^2 S_n^{(\mathrm{Im})}\ \mathrm{Im}\ \delta\mathcal{S}_n$ and, if necessary, second variations, still separate the three directions. Thus independence persists in the sense of *jet*–level controllability even on measure–zero degenerate sets, exactly as in the spatial case.

Remark 4.553 (Algebraic support separation)**.** The three components interrogate mutually distinct operator sectors: t_{corr} depends only on the *off–diagonal bilocal* channel of $\hat{G}_\Lambda$; t_{vac} only on the *coincident diagonal* channel; t_{spec} on the *commutator* layer with $\hat{D}$. No algebraic combination of the first two can generate the third, since $\hat{D}$ does not appear in $\{t_{\mathrm{corr}}, t_{\mathrm{vac}}\}$, while $[\hat{D}, \cdot]$ is a derivation detecting spectral flow (cf. [4, 381]). This complements the Fréchet rank argument above.

4.12.8 Temporal Holonomy and the Zero Locus of Emergent Time

As in the spatial case (Section 4.11.9), temporal observables acquire path dependence through the gauge–covariant identification of modal fibres. The parallel transporter $\mathcal{U}(x \leftarrow x';\gamma)$ along a path $\gamma \subset \mathcal{X}_q$ acts on the Fourfold channels and induces, for the three temporal components $(t_{\mathrm{corr}}, t_{\mathrm{vac}}, t_{\mathrm{spec}})$, corresponding *temporal connections* whose holonomies control the vanishing loci of emergent time. The underlying geometric mechanism is the standard correspondence between flatness, trivial holonomy, and path independence [248, 410, 375, 225]; here it is realised in an operator–theoretic, state–dependent form via $\hat{G}_\Lambda$ and $\hat{D}$.

Temporal Connections from the Rate Forms

Fix a simply connected open set $U \subset \mathcal{X}_q$ and a basepoint $x_* \in U$. Write the Holo–State expectation $\langle\cdot\rangle := \langle\mathring{\mathbf{\Psi}}| \cdot |\mathring{\mathbf{\Psi}}\rangle$. We define three (state–dependent) 1-forms on U:

Definition 4.554 (Temporal Connections)**.** (i) *Correlator phase connection.* For any fixed reference modal ψ (admissible in Definition 4.9) set

$$\Theta_{\mathrm{corr}} := \mathrm{d}_x \arg\Big(\big\langle \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x_*})[\psi^*, \psi]\big\rangle\Big),$$

a $U(1)$ Berry–type connection built from the bilocal channel's phase (cf. eq. (4.243); see also [407]).

(ii) *Vacuum response connection.* With $C_k(x)$ and $I_k(x) := \mathrm{Im}\, C_k(x)$ as in Definition 4.535, define the ℓ^2-valued 1-form

$$\boldsymbol{\vartheta}_{\mathrm{vac}} := \big(\mathrm{d}I_1, \mathrm{d}I_2, \dots\big).$$

Its line integral along γ reproduces the rate form $\dot{t}_{\mathrm{vac}}$ of eq. (4.248) via $\dot{t}_{\mathrm{vac}}(s) = \|\iota_{\dot\gamma}\boldsymbol{\vartheta}_{\mathrm{vac}}\|_2$.

(iii) *Spectral shear connection.* With $S_n^{(\mathrm{Im})}(x, x_*) := \mathrm{Im}\,\langle[\hat{D}, \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x_*})[\psi_n^*, \psi_n]]\rangle$ and weights $w_n(x, x_*)$ as in Definition 4.543, set

$$\boldsymbol{\vartheta}_{\mathrm{spec}} := \big(\mathrm{d}\,(w_1 S_1^{(\mathrm{Im})}),\ \mathrm{d}\,(w_2 S_2^{(\mathrm{Im})}), \dots\big).$$

Again $\dot{t}_{\mathrm{spec}}(s) = \|\iota_{\dot\gamma}\boldsymbol{\vartheta}_{\mathrm{spec}}\|_2$ matches eq. (4.254).

We call the exterior derivatives

$$\mathcal{F}_{\mathrm{corr}} := \mathrm{d}\,\Theta_{\mathrm{corr}}, \qquad \mathcal{F}_{\mathrm{vac}} := \mathrm{d}\,\boldsymbol{\vartheta}_{\mathrm{vac}}, \qquad \mathcal{F}_{\mathrm{spec}} := \mathrm{d}\,\boldsymbol{\vartheta}_{\mathrm{spec}}.$$

the (scalar or vector–valued) *temporal curvatures*. They are insensitive to local unitary gauge and modal basis rotations because arg and ℓ^2 norms are preserved (cf. gauge covariance in Definitions 4.535 and 4.543).

Definition 4.555 (Temporal Holonomy Functionals)**.** For a piecewise C^1 loop $\gamma \subset U$ based at x_*,

$$\mathcal{W}_{\mathrm{corr}}(\gamma) := \exp\Big(\mathrm{i}\oint_\gamma \Theta_{\mathrm{corr}}\Big), \qquad \mathsf{H}_{\mathrm{vac}}(\gamma) := \oint_\gamma \big\|\iota_{\dot\gamma}\boldsymbol{\vartheta}_{\mathrm{vac}}\big\|_2\,\mathrm{d}s, \qquad \mathsf{H}_{\mathrm{spec}}(\gamma) := \oint_\gamma \big\|\iota_{\dot\gamma}\boldsymbol{\vartheta}_{\mathrm{spec}}\big\|_2\,\mathrm{d}s.$$

While $\mathcal{W}_{\mathrm{corr}}$ is a standard Wilson loop, $\mathsf{H}_{\mathrm{vac}}$ and $\mathsf{H}_{\mathrm{spec}}$ measure the ℓ^2-length of the transported imaginary profiles along γ. By Stokes' theorem, $\oint_\gamma \Theta_{\mathrm{corr}} = \int_\Sigma \mathcal{F}_{\mathrm{corr}}$ for any spanning Σ [248, 410].

Zero Locus and Flatness

Proposition 4.556 (Temporal Flatness ⇔ Trivial Holonomy ⇔ Vanishing Times)**.** *Let $U \subset \mathcal{X}_q$ be simply connected. The following are equivalent:*

(a) $\mathcal{F}_{\mathrm{corr}} = 0$, $\mathcal{F}_{\mathrm{vac}} = 0$, *and* $\mathcal{F}_{\mathrm{spec}} = 0$ *on* U.

(b) For every loop $\gamma \subset U$, $\mathcal{W}_{\mathrm{corr}}(\gamma) = 1$ *and* $\mathsf{H}_{\mathrm{vac}}(\gamma) = \mathsf{H}_{\mathrm{spec}}(\gamma) = 0$.

(c) There exist gauge choices and modal orthonormal frames on U *for which the phase* $\arg\langle \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x_*})[\psi^*, \psi]\rangle$ *is constant in* x, *and the profiles* $I_k(x)$ *and* $w_n(x, x_*)S_n^{(\mathrm{Im})}(x, x_*)$ *are constant on* U.

(d) $t_{\mathrm{corr}}(x, x') = t_{\mathrm{vac}}(x, x') = t_{\mathrm{spec}}(x, x') = 0$ *for all* $x, x' \in U$.

Sketch. (a)⇒(b) is Stokes' theorem and path–independence of exact 1-forms. (b)⇒(c): trivial holonomy on a simply connected set yields globally defined potentials; orthogonal invariance of ℓ^2 makes the vacuum/spec profiles constant up to rotations. (c)⇒(d) by the definitions in eq. (4.243) and Definitions 4.535 and 4.543. (d)⇒(a): vanishing of all pairwise integrals implies the generating 1-forms are exact (hence curvature zero). □

Definition 4.557 (Zero Loci of Temporal Observables)**.** For $i \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$, set

$$Z_i := \{(x, x') \in U \times U : \ t_i(x, x') = 0\}, \qquad Z_{\mathrm{time}} := \bigcap_i Z_i.$$

Corollary 4.558 (Holonomy Invariance of the Zero Loci)**.** *If* (a)–(b) *of Proposition 4.556 hold on* U, *then* $Z_i = U \times U$ *for each* i. *Conversely, if for some* i *there exists a loop* $\gamma \subset U$ *with nontrivial holonomy* ($\mathcal{W}_{\mathrm{corr}}(\gamma) \neq 1$ *or* $\mathsf{H}_i(\gamma) > 0$), *then* $Z_i \neq U \times U$; *in fact, there are pairs* (x, x') *on the loop's image with* $t_i(x, x') > 0$.

Idea. Flatness makes each connection pure gauge, whence all endpoint differences vanish. Nontrivial holonomy forces path dependence; choosing x, x' on the loop and comparing the two segments yields a nonzero time by additivity of the rate integrals and arg–additivity. □

Operator Meaning and Examples

Correlator Phase Time. Nontrivial $\mathcal{F}_{\mathrm{corr}}$ is a Berry–type curvature of the bilocal channel's phase; it obstructs a global choice of correlator phase and forces t_{corr} to detect pathwise "phase slippage'' [407].

Vacuum Response Time. $\mathcal{F}_{\mathrm{vac}} \neq 0$ means the imaginary parts of coincident expectations $I_k(x)$ cannot be globally flattened in any modal frame; holonomy measures a retarded, gauge–encoded vacuum polarisation that persists around loops (cf. Section 4.12.3).

Spectral Shear Time. $\mathcal{F}_{\mathrm{spec}} \neq 0$ quantifies a nontrivial commutator–driven spectral flow that cannot be unitarily trivialised along closed paths, the noncommutative analogue of a Berry curvature sourced by $[\hat{D}, \hat{G}_\Lambda]$ [4, 392]. This links directly to modular/time flows in operator algebras [381].

Synthesis: Temporal Holonomy as the Obstruction to a Global "Clock Gauge"

Theorem 4.559 (Temporal Holonomy Obstruction)**.** *On a simply connected $U \subset \mathcal{X}_q$, the existence of a global* clock gauge *in which all three temporal components vanish identically on $U \times U$ is equivalent to joint flatness $\mathcal{F}_{\text{corr}} = \mathcal{F}_{\text{vac}} = \mathcal{F}_{\text{spec}} = 0$. Any nonzero temporal curvature yields a nontrivial holonomy and a nonempty complement $(U \times U) \setminus Z_{\text{time}}$.*

Sketch. Immediate from Proposition 4.556 and Corollary 4.558 and standard holonomy theory [248, 410, 375]. □

Remark (Gauge Covariance). The constructions above use the same parallel transport and gauge conventions as the spatial sector (Lemma 4.270), and inherit basis–independence from the ℓ^2 norms and arg map used in Definitions 4.535 and 4.543. Hence the zero loci Z_i are intrinsic to the Holo–State and the Fourfold/Spectral data, not artifacts of frame choices.

4.12.9 The Strong Static Spectral Structure

Lemma 4.560 (Strong static spectral structure from vanishing spectral time)**.** *Assume the Standard Hypotheses (Finite Entanglement Capacity; Spectral/Relational Smoothness) and let the spectral shear time t_{spec} be defined as in Definition 4.543 and eq. (4.252). Suppose that*

$$t_{\text{spec}}(x, x') = 0 \quad \textit{for all } x, x' \in \mathcal{X}_q$$

with respect to the Holo–State $|\mathring{\mathbf{\Psi}}\rangle$ and for all admissible reference modes. Then:

(i) Expectation–level commutation (state sense). *For every mode n and all $x, x' \in \mathcal{X}_q$,*

$$\Big\langle \mathring{\mathbf{\Psi}} \Big| \, [\hat{D},\ \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_n^*, \psi_n]] \, \Big| \mathring{\mathbf{\Psi}} \Big\rangle = 0. \tag{4.263}$$

Equivalently, the spectral–shear profiles $S_n^{(\text{Im})}(x, x') := \operatorname{Im} \langle\, [\hat{D}, \hat{G}_\Lambda(\cdot)] \,\rangle$ vanish identically.

(ii) Flat spectral connection and trivial holonomy. *The spectral rate 1–form of Definition 4.545 vanishes, hence the spectral curvature and holonomy of Section 4.12.8 are trivial:*

$$\boldsymbol{\vartheta}_{\text{spec}} \equiv 0, \qquad \mathcal{F}_{\text{spec}} = \mathrm{d}\, \boldsymbol{\vartheta}_{\text{spec}} \equiv 0, \qquad \mathsf{H}_{\text{spec}}(\gamma) = 0 \ \ \textit{for every loop } \gamma. \tag{4.264}$$

(iii) Collapse of the spectral distance. *The spectral distance satisfies*

$$d_{\text{spec}}(x, x') = 0 \quad \textit{for all } x, x' \in \mathcal{X}_q, \tag{4.265}$$

i.e. the spectral component of distinguishability is everywhere null (by the duality bounds and eq. (4.252)).

(iv) Operator–level commutation on the GNS cyclic subspace (strong sense). *Assume in addition that:*

(a) the von Neumann algebra $\mathfrak{A}_G := W^*\big(\{\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi^*, \psi]\}\big)$ admits $|\mathring{\mathbf{\Psi}}\rangle$ as a cyclic and separating vector, and

(b) for each x, x' and admissible mode, $[\hat{D},\ \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi^*, \psi]]$ is closable on a common invariant core and affiliated with $\mathfrak{A}_G$.

Then for all x, x' and all admissible modes,

$$[\hat{D},\ \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi^*, \psi]] \ = \ 0 \quad \text{as an operator on the GNS cyclic subspace } \overline{\mathfrak{A}_G\,|\mathring{\Psi}\rangle}. \tag{4.266}$$

Consequently, the spectral sector is static *in the strong sense: the differential generator $\hat{D}$ and all Fourfold channels commute on the physical (cyclic) subspace.*

Proof. (*i*) By definition (4.252), $t_{\mathrm{spec}}(x, x')$ is the ℓ^2–norm of the real vector whose components are $\mathrm{Im}\,\langle[\hat{D}, \hat{G}_\Lambda(\cdot)]\rangle$ (with weights). If the norm is 0 for all x, x', each component must vanish, giving (4.263). Since $[\hat{D}, \cdot]$ is anti–Hermitian for Hermitian inputs, the real part of the expectation is identically zero; hence the full expectation vanishes.

(*ii*) With $S_n^{(\mathrm{Im})} \equiv 0$ one has $w_n S_n^{(\mathrm{Im})} \equiv 0$, whence the spectral rate 1–form $\boldsymbol{\vartheta}_{\mathrm{spec}} = \big(\mathrm{d}(w_n S_n^{(\mathrm{Im})})\big)_n$ of Definition 4.545 vanishes identically. Therefore $\mathcal{F}_{\mathrm{spec}} = \mathrm{d}\,\boldsymbol{\vartheta}_{\mathrm{spec}} = 0$ and the loop functional $\mathsf{H}_{\mathrm{spec}}$ of Definition 4.555 is zero, establishing (4.264).

(*iii*) The spectral distance d_{spec} aggregates magnitudes of the same commutator expectations (real/absolute–value component; Definition 4.402). Since each expectation already vanishes by (*i*), (4.265) follows.

(*iv*) Let $\Omega := |\mathring{\Psi}\rangle$. By (*i*), $\omega([\hat{D}, \hat{G}]) = 0$ for all generators $\hat{G}$ (here $\omega(\cdot) = \langle\Omega|\cdot|\Omega\rangle$). Under (a)–(b), $[\hat{D}, \hat{G}]$ is affiliated with $\mathfrak{A}_G$ and defined on a common core; faithfulness (separating property) of ω on $\mathfrak{A}_G$ together with closability implies that vanishing of all quadratic forms $\langle A\Omega|[\hat{D}, \hat{G}]|A\Omega\rangle$ for $A \in \mathfrak{A}_G$ forces $[\hat{D}, \hat{G}] = 0$ on the GNS cyclic subspace (standard GNS/separating-vector argument; cf. [381]). Hence (4.266). □

Remark 4.561 (Equivalences via Temporal Holonomy)**.** On any simply connected $U \subset \mathcal{X}_q$, items (i)–(iii) are equivalent to $\mathcal{F}_{\mathrm{spec}} = 0$ and trivial spectral holonomy by Section 4.12.8 (see Proposition 4.556).

Intuition. The hypothesis $t_{\mathrm{spec}} \equiv 0$ says that the *phase–sensitive* part of the commutator–induced deformation vanishes everywhere; equivalently, the spectral channel contributes no retarded or absorptive response to relational transport. In concrete terms, the Dirac–type generator $\hat{D}$ does not shear the Fourfold correlator $\hat{G}_\Lambda$ even infinitesimally, so along any path in $\mathcal{X}_q$ there is no build–up of spectral Berry–like phase and no spectral hysteresis. This is precisely why the spectral rate 1–form and its curvature vanish (Section 4.12.8): the "spectral connection'' is pure gauge, hence its Wilson–type loop functional $\mathsf{H}_{\mathrm{spec}}$ is trivial for every loop. Geometrically, the differential and relational structures become jointly integrable in the Holo–State, and operationally the spectral contribution to the emergent clock reads identically zero.

Significance. The strong form of the result adds rigidity: under mild GNS hypotheses the commutator itself vanishes on the physical (cyclic) subspace, so $\hat{D}$ and all Fourfold channels can be simultaneously reduced there. Two consequences follow. First, the temporal triad collapses to a dyad: all path–dependence and holonomy sourced by the spectral sector

disappears, leaving only the correlator phase time and the vacuum response time to encode temporal variation. Second, any nontrivial temporal or spatial signal must originate in the off–diagonal coherence or the diagonal vacuum sector; none can be "faked'' by hidden spectral torsion. In short, the static spectral regime is the spectrally adiabatic and reversible phase of the theory—mathematically flat in the spectral fibre, physically free of spectral decoherence—and it provides a clean baseline against which genuine spectral phenomena (nonzero $\mathcal{F}_{\mathrm{spec}}$, nontrivial $[\hat{D}, \hat{G}_\Lambda]$) can be sharply identified and quantified.

4.12.10 Wordwise Temporal Holonomy and the Zero Locus of Emergent Time

In the holosymmetric *wordwise* calculus, temporal observables acquire path dependence through the gauge/word–covariant transporter $\mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x \leftarrow x';\gamma)$ along paths $\gamma \subset \mathcal{X}_q$ (Section 4.5). The three temporal components $(t^{[\cdot]}_{\mathrm{corr}}, t^{[\cdot]}_{\mathrm{vac}}, t^{[\cdot]}_{\mathrm{spec}})$ of Sections 4.12.2, 4.12.4 and 4.12.6 admit associated *wordwise temporal connections* whose holonomies control the vanishing loci of emergent time. The mechanism parallels the standard correspondence between flatness, trivial holonomy, and path independence, now realised in the operator–theoretic, state–dependent setting via $\hat{G}_\Lambda$ and $\mathbb{D}^{[\mathbf{W}]}$.

Temporal connections from the wordwise rate forms

Fix a simply connected open set $U \subset \mathcal{X}_q$ and a basepoint $x_* \in U$. Write $\langle\cdot\rangle_\Psi := \langle\mathring{\mathbf{\Psi}}|\cdot|\mathring{\mathbf{\Psi}}\rangle$.

Definition 4.562 (Wordwise temporal connections)**.** (i) *Correlator phase connection.* Let $\Gamma^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{corr}}(x, x_*;\gamma)$ be as in eq. (4.244). Define the $U(1)$–valued 1–form $\Theta^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{corr}}$ on U by the rate identity along any C^1 path γ:

$$\iota_{\dot\gamma}\Theta^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{corr}}(s) \;:=\; \dot t^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{corr}}(s) = \mathrm{Im}\Big(\tfrac{\dot\Gamma^{[\cdot]}_{\mathrm{corr}}(s)}{\Gamma^{[\cdot]}_{\mathrm{corr}}(s)}\Big),$$

cf. Definition 4.533. Equivalently, on any node–free chart one may write $\Theta^{[\cdot]}_{\mathrm{corr}} = \mathrm{d}_x \operatorname{Arg} \Gamma^{[\cdot]}_{\mathrm{corr}}(\cdot, x_*;\,\cdot)$.

(ii) *Vacuum response connection.* With the vacuum profile $\vec V^{[\mathbf{W}]}_{\mathrm{vac}}(x)$ and its imaginary part $\mathrm{Im}\,\vec V^{[\mathbf{W}]}_{\mathrm{vac}}(x)$ from Definitions 4.466 and 4.539, set the vector–valued 1–form

$$\boldsymbol{\vartheta}^{[\mathbf{W}]}_{\mathrm{vac}}(x) \;:=\; \mathrm{d}_x\,\mathrm{Im}\,\vec V^{[\mathbf{W}]}_{\mathrm{vac}}(x),$$

so that along γ one has $\dot t^{[\cdot]}_{\mathrm{vac}}(s) = \big\|\iota_{\dot\gamma}\boldsymbol{\vartheta}^{[\cdot]}_{\mathrm{vac}}(s)\big\|_{\ell^2}$ (eq. (4.251)).

(iii) *Spectral shear connection.* With $\mathrm{Im}\,\vec S^{[\mathbf{W}_1,\mathbf{W}_2]}(x, x_*)$ the weighted imaginary spectral–shear profile from Definition 4.547, define

$$\boldsymbol{\vartheta}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{spec}}(x) \;:=\; \mathrm{d}_x\,\mathrm{Im}\,\vec S^{[\mathbf{W}_1,\mathbf{W}_2]}(x, x_*),$$

so that $\dot t^{[\cdot]}_{\mathrm{spec}}(s) = \big\|\iota_{\dot\gamma}\boldsymbol{\vartheta}^{[\cdot]}_{\mathrm{spec}}(s)\big\|_{\ell^2}$ (eq. (4.258)).

Their (scalar or vector–valued) *temporal curvatures* are the exterior derivatives:

$$\mathcal{F}^{[\cdot]}_{\mathrm{corr}} := \mathrm{d}\,\Theta^{[\cdot]}_{\mathrm{corr}}, \qquad \mathcal{F}^{[\cdot]}_{\mathrm{vac}} := \mathrm{d}\,\boldsymbol{\vartheta}^{[\cdot]}_{\mathrm{vac}}, \qquad \mathcal{F}^{[\cdot]}_{\mathrm{spec}} := \mathrm{d}\,\boldsymbol{\vartheta}^{[\cdot]}_{\mathrm{spec}}.$$

Gauge/word covariance of these objects follows from Propositions 4.531, 4.540 and 4.548.

Definition 4.563 (Wordwise temporal holonomy functionals)**.** For a piecewise C^1 loop $\gamma \subset U$ based at x_*,

$$\mathcal{W}^{[\cdot]}_{\mathrm{corr}}(\gamma) := \exp\Big(\mathrm{i}\oint_\gamma \Theta^{[\cdot]}_{\mathrm{corr}}\Big),$$
$$\mathsf{H}^{[\cdot]}_{\mathrm{vac}}(\gamma) := \oint_\gamma \big\|\iota_{\dot\gamma}\boldsymbol{\vartheta}^{[\cdot]}_{\mathrm{vac}}\big\|_{\ell^2}\,\mathrm{d}s,$$
$$\mathsf{H}^{[\cdot]}_{\mathrm{spec}}(\gamma) := \oint_\gamma \big\|\iota_{\dot\gamma}\boldsymbol{\vartheta}^{[\cdot]}_{\mathrm{spec}}\big\|_{\ell^2}\,\mathrm{d}s.$$

By Stokes' theorem on simply connected charts, $\oint_\gamma \Theta^{[\cdot]}_{\mathrm{corr}} = \int_\Sigma \mathcal{F}^{[\cdot]}_{\mathrm{corr}}$ for any spanning surface Σ (e.g. [249, 411]).

Zero locus and flatness of the wordwise temporal sector

Proposition 4.564 (Temporal flatness $\Leftrightarrow$ trivial holonomy $\Leftrightarrow$ vanishing wordwise times)**.** *Let $U \subset \mathcal{X}_q$ be simply connected. The following are equivalent:*

(a) $\mathcal{F}^{[\cdot]}_{\mathrm{corr}} = 0$, $\mathcal{F}^{[\cdot]}_{\mathrm{vac}} = 0$, *and* $\mathcal{F}^{[\cdot]}_{\mathrm{spec}} = 0$ *on* U.

(b) *For every loop* $\gamma \subset U$, $\mathcal{W}^{[\cdot]}_{\mathrm{corr}}(\gamma) = 1$ *and* $\mathsf{H}^{[\cdot]}_{\mathrm{vac}}(\gamma) = \mathsf{H}^{[\cdot]}_{\mathrm{spec}}(\gamma) = 0$.

(c) *There exist gauge/word choices and compatible modal frames on* U *such that the potentials are constant:*

$$\operatorname{Arg}\Gamma^{[\cdot]}_{\mathrm{corr}}(x, x_*; \cdot) \equiv \text{const}, \quad \operatorname{Im}\vec{V}^{[\cdot]}_{\mathrm{vac}}(x) \equiv \text{const}, \quad \operatorname{Im}\vec{S}^{[\cdot]}(x, x_*) \equiv \text{const} \quad \text{on } U.$$

(d) $t^{[\cdot]}_{\mathrm{corr}}(x, x') = t^{[\cdot]}_{\mathrm{vac}}(x, x') = t^{[\cdot]}_{\mathrm{spec}}(x, x') = 0$ *for all* $x, x' \in U$.

Proof. **(a)⇒(b).** On a simply connected U, $\mathcal{F}^{[\cdot]}_{\mathrm{corr}} = 0$ implies $\Theta^{[\cdot]}_{\mathrm{corr}} = \mathrm{d}\phi$ for some scalar potential ϕ, hence $\oint_\gamma \Theta^{[\cdot]}_{\mathrm{corr}} = 0$ and $\mathcal{W}^{[\cdot]}_{\mathrm{corr}}(\gamma) = 1$. For the vector–valued cases, $\mathcal{F}^{[\cdot]}_{\mathrm{vac}} = 0$ and $\mathcal{F}^{[\cdot]}_{\mathrm{spec}} = 0$ imply $\boldsymbol{\vartheta}^{[\cdot]}_{\mathrm{vac}} = \mathrm{d}\mathbf{v}$ and $\boldsymbol{\vartheta}^{[\cdot]}_{\mathrm{spec}} = \mathrm{d}\mathbf{s}$ for some profiles $\mathbf{v}, \mathbf{s}$; thus $\iota_{\dot\gamma}\boldsymbol{\vartheta}^{[\cdot]}_{\mathrm{vac}} = \frac{\mathrm{d}}{\mathrm{d}s}\mathbf{v}(\gamma(s))$ and similarly for $\mathbf{s}$, so the loop integrals of their ℓ^2–norms vanish, $\mathsf{H}^{[\cdot]}_{\mathrm{vac}}(\gamma) = \mathsf{H}^{[\cdot]}_{\mathrm{spec}}(\gamma) = 0$.

(b)⇒(c). Trivial correlator holonomy yields a single–valued phase potential ϕ with $\Theta^{[\cdot]}_{\mathrm{corr}} = \mathrm{d}\phi$; choose the gauge fixing $\phi \equiv \text{const}$, making $\operatorname{Arg}\Gamma^{[\cdot]}_{\mathrm{corr}}$ constant. For the vector–valued connections, the vanishing of $\oint_\gamma \big\|\iota_{\dot\gamma}\boldsymbol{\vartheta}^{[\cdot]}_{\mathrm{vac/spec}}\big\|_{\ell^2}\mathrm{d}s$ for all loops forces $\iota_{\dot\gamma}\boldsymbol{\vartheta}^{[\cdot]}_{\mathrm{vac/spec}} = 0$ for all loop tangents $\dot\gamma$, hence $\boldsymbol{\vartheta}^{[\cdot]}_{\mathrm{vac/spec}} = 0$ on U. Therefore the corresponding profiles are constant (up to global orthogonal rotations in the modal index, which can be absorbed into the compatible frame).

(c)⇒(d). By Definitions 4.529, 4.541 and 4.549, each time is the integral of the corresponding rate 1–form pulled back along a path; if the underlying potentials are constant on U then all rates vanish and hence all three times vanish for all pairs in U.

(d)⇒(a). If $t^{[\cdot]}_{\mathrm{corr}}(x,x')=0$ for every pair, then for any loop γ, $\oint_\gamma \Theta^{[\cdot]}_{\mathrm{corr}} = \tilde{t}^{[\cdot]}_{\mathrm{corr}}(x_*,x_*;\gamma)=0$, so $\Theta^{[\cdot]}_{\mathrm{corr}}$ has zero periods; on simply connected U this implies exactness and hence $\mathcal{F}^{[\cdot]}_{\mathrm{corr}}=0$. Likewise, $t^{[\cdot]}_{\mathrm{vac}}(x,x')=0$ and $t^{[\cdot]}_{\mathrm{spec}}(x,x')=0$ give $\int_\gamma \left\|\iota_{\dot\gamma}\boldsymbol{\vartheta}^{[\cdot]}_{\mathrm{vac/spec}}\right\|_{\ell^2}\mathrm{d}s=0$ for all loops, forcing $\boldsymbol{\vartheta}^{[\cdot]}_{\mathrm{vac/spec}}=0$ pointwise (vary loop tangents locally), hence vanishing curvatures. □

Definition 4.565 (Zero loci of wordwise temporal observables)**.** For $i\in\{\mathrm{corr},\mathrm{vac},\mathrm{spec}\}$, set

$$Z^{[\cdot]}_i := \{(x,x')\in U\times U:\ t^{[\cdot]}_i(x,x')=0\},\qquad Z^{[\cdot]}_{\mathrm{time}} := \bigcap_i Z^{[\cdot]}_i.$$

Corollary 4.566 (Holonomy control of the wordwise temporal zero loci)**.** *If the equivalent conditions of Proposition 4.564 hold on U, then $Z^{[\cdot]}_i = U\times U$ for each component and hence $Z^{[\cdot]}_{\mathrm{time}}=U\times U$. Conversely, if some loop $\gamma\subset U$ exhibits nontrivial temporal holonomy for component i ($\mathcal{W}^{[\cdot]}_{\mathrm{corr}}(\gamma)\neq 1$ or $\mathsf{H}^{[\cdot]}_i(\gamma)>0$), then $Z^{[\cdot]}_i\neq U\times U$; in fact there exist x,x' on the loop image with $t^{[\cdot]}_i(x,x')>0$.*

Remark 4.567 (Holonomic invisibility (wordwise))**.** Nontrivial microscopic holonomy may act *invisibly* on the temporal sector if it lies in a band–subspace annihilated by the Holo–State weights and projections defining the rate forms (Definitions 4.539, 4.547 and 4.562). The relevant notion is holonomy in the *state–conditioned wordwise representation* carried by the probe algebra; triviality there suffices for the vanishing of $t^{[\cdot]}_{\mathrm{corr}}, t^{[\cdot]}_{\mathrm{vac}}, t^{[\cdot]}_{\mathrm{spec}}$ even if the underlying connection has nonzero bare curvature.

Wordwise Temporal Holonomy as the Obstruction to a Global "Clock Gauge"

Theorem 4.568 (Wordwise Temporal Holonomy Obstruction)**.** *Let $U\subset\mathcal{X}_q$ be simply connected. The following are equivalent:*

(a) There exists a global wordwise clock gauge *on U in which, for all $x,x'\in U$,*

$$t^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{corr}}(x,x') = t^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{vac}}(x,x') = t^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{spec}}(x,x') = 0.$$

(b) The wordwise temporal curvatures vanish jointly on U:

$$\mathcal{F}^{[\cdot]}_{\mathrm{corr}} = \mathrm{d}\,\Theta^{[\cdot]}_{\mathrm{corr}} = 0,\qquad \mathcal{F}^{[\cdot]}_{\mathrm{vac}} = \mathrm{d}\,\boldsymbol{\vartheta}^{[\cdot]}_{\mathrm{vac}} = 0,\qquad \mathcal{F}^{[\cdot]}_{\mathrm{spec}} = \mathrm{d}\,\boldsymbol{\vartheta}^{[\cdot]}_{\mathrm{spec}} = 0,$$

where $\Theta^{[\cdot]}_{\mathrm{corr}}$, $\boldsymbol{\vartheta}^{[\cdot]}_{\mathrm{vac}}$, $\boldsymbol{\vartheta}^{[\cdot]}_{\mathrm{spec}}$ are the wordwise temporal connections of Definition 4.562.

Moreover, if any temporal curvature is nonzero on U, then some temporal holonomy is nontrivial and the complement $(U\times U)\setminus Z^{[\cdot]}_{\mathrm{time}}$ is nonempty, where $Z^{[\cdot]}_{\mathrm{time}}$ is defined in Definition 4.565.

Proof. (a)⇒(b). If all three times vanish for every pair $(x,x')\in U\times U$, then for any piecewise C^1 loop $\gamma\subset U$ based at x_* we have

$$\oint_\gamma \Theta^{[\cdot]}_{\mathrm{corr}} = \tilde{t}^{[\cdot]}_{\mathrm{corr}}(x_*,x_*;\gamma)=0,\qquad \oint_\gamma \left\|\iota_{\dot\gamma}\boldsymbol{\vartheta}^{[\cdot]}_{\mathrm{vac}}\right\|_{\ell^2}\mathrm{d}s=0,\qquad \oint_\gamma \left\|\iota_{\dot\gamma}\boldsymbol{\vartheta}^{[\cdot]}_{\mathrm{spec}}\right\|_{\ell^2}\mathrm{d}s=0.$$

Thus the correlator Wilson loop is trivial and the vector–valued loop integrals vanish. On a simply connected U, vanishing periods imply exactness of the connections; therefore all curvatures are zero. This is the direction *(d)⇒(a)* in Proposition 4.564 specialized to the temporal sector.

(b)⇒(a). Joint flatness implies trivial temporal holonomy on U; by path independence of the exact 1–forms, there exist global potentials/frames (word/gauge choices and compatible modal bases) in which all rates vanish: $\iota_{\dot\gamma}\Theta^{[\cdot]}_{\mathrm{corr}} = \iota_{\dot\gamma}\boldsymbol{\vartheta}^{[\cdot]}_{\mathrm{vac}} = \iota_{\dot\gamma}\boldsymbol{\vartheta}^{[\cdot]}_{\mathrm{spec}} = 0$. Integrating along any path from x to x' gives $t^{[\cdot]}_{\mathrm{corr}}(x,x') = t^{[\cdot]}_{\mathrm{vac}}(x,x') = t^{[\cdot]}_{\mathrm{spec}}(x,x') = 0$ (see Definitions 4.533, 4.541 and 4.549 and Proposition 4.564).

For the final claim, if some temporal curvature is nonzero then there exists a loop with nontrivial holonomy (standard holonomy theory). By Corollary 4.566, this forces the existence of (x,x') with $t^{[\cdot]}_i(x,x') > 0$ for at least one component $i \in \{\mathrm{corr},\mathrm{vac},\mathrm{spec}\}$, so $(U\times U)\setminus Z^{[\cdot]}_{\mathrm{time}} \neq \varnothing$. □

Remark (Gauge/word covariance). All constructions use the wordwise parallel transport and covariance conventions of Section 4.5, and inherit basis–independence from the Arg and ℓ^2–norm invariances proved in Propositions 4.531, 4.540 and 4.548. Hence each zero locus $Z^{[\cdot]}_i$ of Definition 4.565 is intrinsic to $(\hat G_\Lambda, \mathbb{D}^{[\mathbf{W}]}, |\overset{\circ}{\boldsymbol{\Psi}}\rangle)$.

4.12.11 Holosymmetric Wordwise Strong Static Spectral Structure

Lemma 4.569 (Strong static spectral structure from vanishing wordwise spectral time)**.** *Assume the Standard Hypotheses (Finite Entanglement Capacity; Spectral/Relational Smoothness) and let $t^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{spec}}$ be as in Definition 4.547 and eq. (4.255). Suppose that*

$$t^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{spec}}(x,x') = 0 \quad \textit{for all } x,x' \in \mathcal{X}_q$$

with respect to $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$ and all admissible reference modes. Then:

(i) Expectation–level commutation (state sense). *For every band n and all $x,x' \in \mathcal{X}_q$,*

$$\Big\langle \overset{\circ}{\boldsymbol{\Psi}}\Big|\, \Big[\mathbb{D}^{[\mathbf{W}_1]}(x),\, \hat G_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})[\psi_n^{(x)*},\psi_n^{(x')}]\Big]\, \Big|\overset{\circ}{\boldsymbol{\Psi}}\Big\rangle \;=\; 0, \tag{4.267}$$

equivalently $S_n^{(\mathrm{Im})[\mathbf{W}_1,\mathbf{W}_2]}(x,x') = \mathrm{Im}\ \mathcal{S}_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \equiv 0$.

(ii) Flat spectral connection and trivial holonomy.

$$\boldsymbol{\vartheta}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{spec}} \equiv 0, \qquad \mathcal{F}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{spec}} = \mathrm{d}\,\boldsymbol{\vartheta}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{spec}} \equiv 0, \qquad \mathsf{H}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{spec}}(\gamma) = 0 \ \ \forall \textit{ loops } \gamma, \tag{4.268}$$

with $\boldsymbol{\vartheta}^{[\cdot]}_{\mathrm{spec}}$ from Definition 4.549.

(iii) Collapse of the wordwise spectral distance.

$$d^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{spec}}(x,x') = 0 \quad \textit{for all } x,x' \in \mathcal{X}_q, \tag{4.269}$$

where $d^{[\cdot]}_{\mathrm{spec}}$ is defined in Definition 4.408.

(iv) Operator–level commutation on the GNS cyclic subspace (strong sense). *Let* $\Omega := |\mathring{\Psi}\rangle$ *and* $\mathfrak{A}_G := W^*\Big(\{\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi^*, \psi]\}\Big)$. *If (a)* Ω *is cyclic and separating for* $\mathfrak{A}_G$*, and (b) for each* x, x' *and admissible mode the commutator* $[\mathbb{D}^{[\mathbf{W}_1]}(x),\, \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi^*, \psi]]$ *is closable on a common invariant core and affiliated with* $\mathfrak{A}_G$*, then*

$$\Big[\mathbb{D}^{[\mathbf{W}_1]}(x),\, \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi^*, \psi]\Big] \;=\; 0 \quad \textit{on} \quad \overline{\mathfrak{A}_G\,\Omega}. \tag{4.270}$$

Proof. (i) By eq. (4.255), $t^{[\cdot]}_{\mathrm{spec}}(x, x')$ is the ℓ^2–norm of the weighted vector with components $\mathrm{Im}\,\mathcal{S}^{[\cdot]}_n(x, x')$. If this norm vanishes for every pair (x, x'), each component must vanish, yielding (4.267).

(ii) With $S^{(\mathrm{Im})[\cdot]}_n \equiv 0$ one has $w^{[\cdot]}_n S^{(\mathrm{Im})[\cdot]}_n \equiv 0$; hence each component of the 1–form $\boldsymbol{\vartheta}^{[\cdot]}_{\mathrm{spec}} = \mathrm{d}_x\,\mathrm{Im}\,\vec{S}^{[\cdot]}$ vanishes. Therefore the curvature and the loop functional are trivial, giving (4.268) (Definitions 4.549 and 4.563).

(iii) The wordwise spectral distance aggregates magnitudes of the same spectral–shear expectations (with identical weights) by Definition 4.408. Since each expectation already vanishes by (i), one gets (4.269).

(iv) Let $\omega(A) := \langle\Omega|A|\Omega\rangle$. From (i) we have $\omega([\mathbb{D}^{[\mathbf{W}_1]}(x), \hat{G}]) = 0$ for all generators $\hat{G} \in \mathfrak{A}_G$ (wordwise Fourfold channels). By (a)–(b), the commutators are affiliated with $\mathfrak{A}_G$ and defined on a common core; faithfulness (separating property) of ω on $\mathfrak{A}_G$ and closability imply that vanishing of all quadratic forms $\langle A\Omega|[\mathbb{D}^{[\mathbf{W}_1]}(x), \hat{G}]|A\Omega\rangle$ for $A \in \mathfrak{A}_G$ forces the operator to be zero on the GNS cyclic subspace $\overline{\mathfrak{A}_G\,\Omega}$ (standard GNS/separating–vector argument). This proves (4.270). □

Remark 4.570 (Equivalences via wordwise temporal holonomy)**.** On any simply connected $U \subset \mathcal{X}_q$, items (i)–(iii) are equivalent to $\mathcal{F}^{[\cdot]}_{\mathrm{spec}} = 0$ and trivial spectral holonomy by Section 4.12.10 and Proposition 4.564.

4.13 Holosymmetric Wordwise Triadic Chronogeometry from Distances and Times

With the three *wordwise* distances

$$d^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{corr}}, \quad d^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{vac}}, \quad d^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{spec}} \quad \text{(Definitions 4.339, 4.408 and 4.468)}$$

and the three *wordwise* times

$$t^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{corr}}, \quad t^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{vac}}, \quad t^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{spec}} \quad \text{(Definitions 4.529, 4.539 and 4.547)}$$

we can assemble a holosymmetric, background–free "spacetime'' structure. The key inputs already established are: (i) functional orthogonality of the three axes (Section 4.11.1), (ii) functional independence of their generators (Section 4.11.7), and (iii) coherent finiteness (UV/IR) of all components (Propositions 4.422 and 4.426 and Principles 4.7 and 4.8).

4.13.1 Holosymmetric Wordwise Triadic Space and Triadic Time Fibres

Space fibre (recall). At each locus x, the *wordwise* distinguishability fibre $\mathcal{F}_x^{\mathrm{w}} \simeq \mathrm{span}\{\mathbf{e}_{\mathrm{corr}}, \mathbf{e}_{\mathrm{vac}}, \mathbf{e}_{\mathrm{spec}}\}$ (Definition 4.490) hosts the *space* displacement

$$\vec{d}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \;:=\; \Big(d_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]}, d_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}, d_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}\Big) \;\in\; \mathcal{F}_x^{\mathrm{w}},$$

with composite triadic distance $D^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') = \|\mathbf{M}^{1/2}(x)\,\vec{d}^{[\cdot]}(x,x')\|_2$ (Section 4.11.2 and Definition 4.494).

Time fibre (definition). Dually, define at each x the *temporal* fibre

$$\mathcal{T}_x^{\mathrm{w}} \;\cong\; \mathrm{span}\{\mathbf{f}_{\mathrm{corr}}, \mathbf{f}_{\mathrm{vac}}, \mathbf{f}_{\mathrm{spec}}\} \;\cong\; \mathbb{R}^3,$$

with the temporal displacement vector

$$\vec{t}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \;:=\; \Big(t_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]}, t_{\mathrm{vac}}^{[\mathbf{W}_1,\mathbf{W}_2]}, t_{\mathrm{spec}}^{[\mathbf{W}_1,\mathbf{W}_2]}\Big) \;\in\; \mathcal{T}_x^{\mathrm{w}}. \tag{4.271}$$

Equip $\mathcal{T}_x^{\mathrm{w}}$ with a symmetric positive–definite *temporal calibration* $\mathbf{N}(x) \succ 0$ (analogue of $\mathbf{M}(x)$), and define the composite time

$$T^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \;:=\; \Big\|\mathbf{N}^{1/2}(x)\,\vec{t}^{[\cdot]}(x,x')\Big\|_2. \tag{4.272}$$

By Propositions 4.532, 4.540 and 4.548, each component is finite and gauge/word–covariant; orthogonality of the axes (Section 4.11.1) implies that, in the canonical choice $\mathbf{N} = \mathbf{I}_3$, the temporal components are orthonormal at quadratic order.

4.13.2 Polar decomposition viewpoint: space as magnitude, time as phase

For each axis, the *same* wordwise relational amplitude furnishes a magnitude (space) and a phase (time). For the correlator sector, with normalized complex overlap $\Upsilon_{\mathrm{corr}}^{[\cdot]} = C^{[\cdot]} e^{i t_{\mathrm{corr}}^{[\cdot]}}$ (Remark 4.534), the distance uses $|C^{[\cdot]}|$ while the time uses Arg.

For the vacuum sector, set

$$\Delta\vec{C}_{\mathrm{diag}}^{[\cdot]}(x,x') \;:=\; \vec{V}_{\mathrm{vac}}^{[\mathbf{W}_1]}(x) \;-\; \vec{V}_{\mathrm{vac}}^{[\mathbf{W}_2]}(x'),$$

so $d_{\mathrm{vac}}^{[\cdot]} = \big\|\Delta\vec{C}_{\mathrm{diag}}^{[\cdot]}\big\|_{\ell^2}$ while $t_{\mathrm{vac}}^{[\cdot]} = \big\|\,\mathrm{Im}\,\Delta\vec{C}_{\mathrm{diag}}^{[\cdot]}\big\|_{\ell^2}$ (Definitions 4.468 and 4.539). For the spectral sector, $\vec{\mathcal{S}}^{[\cdot]}$ contributes $d_{\mathrm{spec}}^{[\cdot]} = \big\|\vec{\mathcal{S}}^{[\cdot]}\big\|_{\ell^2}$ and $t_{\mathrm{spec}}^{[\cdot]} = \big\|\,\mathrm{Im}\,\vec{\mathcal{S}}^{[\cdot]}\big\|_{\ell^2}$ (Definitions 4.408 and 4.547). Thus "*space*" (distance) collects magnitudes; "*time*" collects phases/imaginary parts—both extracted from gauge/word–covariant expectations in the Holo–State. This polar structure is the operational origin of spacetime in the wordwise calculus.

4.13.3 A Holosymmetric Wordwise Chronogeometric Structure

Definition 4.571 (Triadic chronometric form)**.** Fix calibrations $\mathbf{M}(x) \succ 0$ and $\mathbf{N}(x) \succ 0$ and a strictly positive *coherence–speed vector* $\mathbf{c}(x) = \mathrm{diag}(c_{\mathrm{corr}}, c_{\mathrm{vac}}, c_{\mathrm{spec}})$ (dimensionful scales). Define the indefinite quadratic form on $\mathcal{F}_x^{\mathrm{w}} \oplus \mathcal{T}_x^{\mathrm{w}}$ by

$$\mathfrak{s}^2_{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \;:=\; \underbrace{\Big\|\mathbf{c}(x)\,\mathbf{N}^{1/2}(x)\,\vec{t}^{[\cdot]}(x,x')\Big\|_2^2}_{\text{temporal part}} \;-\; \underbrace{\Big\|\mathbf{M}^{1/2}(x)\,\vec{d}^{[\cdot]}(x,x')\Big\|_2^2}_{\text{spatial part}}\,. \tag{4.273}$$

We call $\mathfrak{s}^2$ the *wordwise triadic spacetime interval*.

Remark 4.572 (Causal trichotomy and cones)**.** Define (with respect to x) *timelike*, *null*, and *spacelike* separations by the sign of $\mathfrak{s}^2$. Equivalently, in the canonical gauges $\mathbf{M} = \mathbf{N} = \mathbf{I}_3$ this is the comparison $D^{[\cdot]}(x,x') \lessgtr \|\mathbf{c}\,\vec{t}^{[\cdot]}(x,x')\|_2$. The null condition $\mathfrak{s}^2 = 0$ yields a triadic cone, whose shape is fixed by the coherence speeds $(c_{\mathrm{corr}}, c_{\mathrm{vac}}, c_{\mathrm{spec}})$ and—via Section 4.12.4 and Definition 4.549— by the holonomy/curvature of the wordwise connections. Word/gauge invariance follows from the invariances of $D^{[\cdot]}$ and $T^{[\cdot]}$.

Proposition 4.573 (Proper time and proper distance (wordwise))**.** *For any pair (x,x'):*

$$\begin{aligned} \textit{Proper time:}\quad & \tau^{[\cdot]}(x,x') := \sqrt{\max\{\mathfrak{s}^2_{[\cdot]}(x,x'),0\}},\\ \textit{Proper distance:}\quad & \ell^{[\cdot]}(x,x') := \sqrt{\max\{-\mathfrak{s}^2_{[\cdot]}(x,x'),0\}}. \end{aligned}$$

These are well-defined, finite (coherent regularisation), and gauge/word–invariant. In the duality–collapse regime they reduce to the corresponding Twofold notions (Remark 4.512).

4.13.4 Infinitesimal Structure and Operator Metric-Chronometric Tensors

Let $x' = x + \delta x$ in an admissible chart. Using the coincidence–limit Hessian for space (Section 4.11.3) and the rate 1–forms for time (Definitions 4.533, 4.541 and 4.549), define the local tensors

$$g^{(\mathrm{space})}_{\mu\nu}(x) := \tfrac{1}{2}\,\partial_\mu\partial_\nu\Big(\vec{d}^{[\cdot]}(x,x+\delta x)^{\top}\mathbf{M}(x)\,\vec{d}^{[\cdot]}(x,x+\delta x)\Big)\Big|_{\delta x=0}, \tag{4.274}$$

$$h^{(\mathrm{time})}_{\mu\nu}(x) := \Big(\mathbf{c}(x)\,\mathbf{N}^{1/2}(x)\,\boldsymbol{\theta}_\mu(x)\Big)^{\top}\Big(\mathbf{c}(x)\,\mathbf{N}^{1/2}(x)\,\boldsymbol{\theta}_\nu(x)\Big), \tag{4.275}$$

where $\boldsymbol{\theta}_\mu(x)$ is the 3–component column vector collecting the gauge/word–invariant temporal *rates* along the coordinate direction ∂_μ:

$$\boldsymbol{\theta}_\mu(x) \;:=\; \big(\theta^{\mathrm{corr}}_\mu(x),\ \theta^{\mathrm{vac}}_\mu(x),\ \theta^{\mathrm{spec}}_\mu(x)\big)^{\top},$$

with each θ^i_μ obtained from the corresponding wordwise rate 1–form by contraction with ∂_μ (Definitions 4.533, 4.541 and 4.549). By orthogonality (Section 4.11.1) cross terms vanish at quadratic order, and the infinitesimal interval is

$$\mathrm{d}\mathfrak{s}^2 \;=\; h_{\mu\nu}\,\mathrm{d}x^\mu\,\mathrm{d}x^\nu \;-\; g_{\mu\nu}\,\mathrm{d}x^\mu\,\mathrm{d}x^\nu. \tag{4.276}$$

4.13.5 Conceptual consequences

(a) *No background manifold:* "Space" is the fibrewise norm of magnitudes; "time" is the fibrewise norm of phases—both extracted from Holo–State expectations. The PQS is the null locus $D^{[\cdot]} = T^{[\cdot]} = 0$ (Section 4.11.8), i.e. complete relational indistinguishability.

(b) *Self–regularisation:* All components are finite by endogenous UV/IR control (Proposition 4.426 and Principles 4.7 and 4.8); no external cutoffs enter the definition of spacetime intervals.

(c) *Holonomy and "time curvature":* Nontrivial loop phases in $t^{[\cdot]}_{\mathrm{corr}}$ and nonzero spectral/vacuum rate curvature produce temporal holonomies (phase clocks), supplying operational analogues of redshift/time–dilation purely from relational data.

(d) *Causal cones are operational:* The triadic cone of Remark 4.572 is determined by coherence speeds and Holo–State support; achronal/chronal relations are experiment– (window–) dependent but gauge/word–invariant.

(e) *Equivalence of presentations:* By word/gauge invariance and the componentwise equivalence results (e.g. Proposition 4.474 together with the corresponding statements for $d^{[\cdot]}_{\mathrm{corr}}$ and $d^{[\cdot]}_{\mathrm{spec}}$), all *exogenous* statements about $D, T, \mathfrak{s}^2$ follow from the wordwise ones.

Summary. In the holosymmetric wordwise calculus, *space* and *time* are two faces of the same relational amplitudes: space collects their magnitudes (distances), time collects their phases (times). Orthogonality and independence of the triads make the composite interval (4.273) block–diagonal (to quadratic order), while endogenous regularisation renders it finite and well–posed. Spacetime thus *emerges* as a calibrated, gauge/word–invariant chronometric structure on the bundle $\mathcal{F}^{\mathrm{w}} \oplus \mathcal{T}^{\mathrm{w}} \to \mathcal{X}_q$, with the PQS as its null locus.

4.14 The Minimal Scale and the Indivisibility of the PQS

In contrast to classical approaches that *postulate* a fundamental length, the IQR framework *derives* a minimal relational scale as a consequence of the operator–theoretic dynamics of the Fourfold correlator $\hat{G}_\Lambda$, the spectral generator $\hat{D}$, and the Holo–State $|\mathring{\mathbf{\Psi}}\rangle$. At this scale, the very notion of separation ceases to be meaningful because *all* emergent relational probes (spatial and temporal) become indistinguishable.

Relational Null Condition. We collect the distinguishability and temporal null conditions together into a single precise statement (without redefining any previously introduced observables).

Definition 4.574 (Relational Null Condition)**.** Let $x, x' \in \mathcal{X}_q$. Write $\vec{d}(x, x') = (d_{\mathrm{corr}}, d_{\mathrm{vac}}, d_{\mathrm{spec}})$ and $\vec{t}(x, x') = (t_{\mathrm{corr}}, t_{\mathrm{vac}}, t_{\mathrm{spec}})$,

to denote the triads of emergent distances and times as defined earlier. We say that (x, x') satisfies the *relational null condition* iff

$$\boxed{\vec{d}(x, x') = \mathbf{0} \quad \text{and} \quad \vec{t}(x, x') = \mathbf{0}.} \tag{4.277}$$

Equivalently, $D(x, x') = \|\vec{d}(x, x')\|_2 = 0$ and $t_{\text{corr}}(x, x') = t_{\text{vac}}(x, x') = t_{\text{spec}}(x, x') = 0$.

Interpretation. The condition $D(x, x') = 0$ does not merely express a coordinate coincidence; it asserts a *relational collapse* in which the off–diagonal coherence channel, the diagonal vacuum profile, and the commutator–driven spectral shear all agree in assigning zero separation and zero temporal tension between x and x'. In the holonomy language, this corresponds to a regime where the relevant connections can be flattened (trivial holonomy) along paths joining x and x' (cf. Sections 4.11.9 and 4.12.8), so that no phase slippage, no vacuum recoil, and no spectral shear accumulate.

Regularity Hypotheses Ensuring Finiteness and Consistency (Recap) We *do not re-define* the regularity assumptions; we summarise the working hypotheses and point to their formal statements:

(a) **Spectral Smoothness** (Axiom 4.415). The spectral–shear observables obey polynomial bounds,

$$|\mathcal{S}_n(x, x')| \lesssim \lambda_n^k,$$

which control high–frequency behaviour and make spectral aggregates (distance/time) finite.

(b) **Relational Smoothness** (Axiom 4.415). The coincident self–correlators vary smoothly,

$$|C_k(x) - C_k(x')| \lesssim \lambda_k^m,$$

preventing pathological vacuum fluctuations across loci.

(c) **Endogenous Regularisation / Finite Entanglement Capacity** (Principle 4.1 and Definition 4.392). Modal weights decay,

$$w_n(x, x') \lesssim \lambda_n^{-\alpha}, \qquad \alpha > \max\{k, m\} + \tfrac{1}{2},$$

so that all RMS aggregates defining (d_i, t_i) converge and the ultraviolet sector is intrinsically regulated by the state–dependent relational structure.

Consequences to Keep in View. (i) The minimal scale is *emergent*: it is achieved dynamically where the joint zero locus of spatial and temporal fibres is reached; no external cutoff is assumed. (ii) In regimes with *static spectral structure* (Section 4.12.9), the spectral component identically vanishes ($d_{\text{spec}} = t_{\text{spec}} = 0$), so any residual granularity must arise from the correlator and vacuum sectors alone. (iii) Nontrivial temporal/spatial holonomy obstructs the collapse to the PQS; detecting such holonomy (Berry–type correlator curvature, vacuum or spectral curvature) certifies departure from the minimal relational scale.

Remark 4.575 (On Λ_{eff} versus the cosmological constant)**.**

(i) **Definition and role of Λ_{eff}.** $\Lambda_{\text{eff}}^{[\text{obs}]}(x, x'; \varepsilon)$ is the *effective spectral cutoff* defined operationally in Definition 4.576: it is the smallest threshold in the spectrum of the relational generator beyond which the weighted features of a chosen observable become $< \varepsilon$. It is *derived*, depends on the state, on the loci (x, x'), on the observable label [obs], and on the tolerance ε, and it is monotone nonincreasing in ε. Its inverse is the PQS coherence width: $\ell_{\text{PQS}} = (\Lambda_{\text{eff}})^{-1}$.

(ii) **Units.** If the generator's eigenvalues $\{\lambda_n\}$ have units L^{-1} (or energy), then

$$[\Lambda_{\text{eff}}] = L^{-1}, \qquad [\ell_{\text{PQS}}] = L.$$

(iii) **Cosmological constant (for comparison).** The cosmological constant is the parameter Λ_{cosmos} in the Einstein–Hilbert sector,

$$S = \frac{1}{16\pi G} \int (R - 2\Lambda_{\text{cosmos}}) \sqrt{-g}\, d^4x,$$

with units $[\Lambda_{\text{cosmos}}] = L^{-2}$ and associated de Sitter curvature radius $L_\Lambda = \sqrt{3/\Lambda_{\text{cosmos}}}$ (see, e.g., [412, 413, 211, 212, 414, 134, 135]). It is a background gravitational parameter, not an operational spectral threshold.

(iv) **No identification.** Λ_{eff} and Λ_{cosmos} pertain to *different sectors* (relational spectral analysis vs. background curvature), have *different dimensions*, and play *distinct roles*. No identification or inference between them is intended anywhere in this chapter.

(v) **Notational convention.** To avoid ambiguity, we reserve Λ_{cosmos} for the cosmological constant and use Λ_{eff} exclusively for the effective spectral cutoff. The statement "$\ell_{\text{PQS}} = (\Lambda_{\text{eff}})^{-1}$" *never* involves Λ_{cosmos}.

(vi) **Cross-chapter dictionary (cosmology vs. spectral).** Throughout Chapters 11 and 17, we use

$$\Lambda_{\text{eff}} = \Lambda + \kappa' \varepsilon_\Lambda$$

for the *cosmology-sector* effective cosmological constant (dimension L^{-2}), with ε_Λ the residual dark-energy density defined there. By contrast, in this chapter $\Lambda_{\text{eff}}^{[\text{obs}]}(x, x'; \varepsilon)$ always denotes the *spectral cutoff* (dimension L^{-1}). The presence of arguments $(x, x'; \varepsilon)$ is a built-in disambiguator for the spectral notion.

(vii) **Reserved symbols.** We reserve Λ_{cosmos} for statements that refer to the cosmological constant as a background parameter in the Einstein–Hilbert sector (dimension L^{-2}). In Chapters 11 and 17, when the cosmology-sector constant appears without the "cosmos" subscript, it participates in the effective combination $\Lambda_{\text{eff}} = \Lambda + \kappa' \varepsilon_\Lambda$. No symbol in this chapter is to be identified with that effective cosmological constant.

(viii) **Units gate.** A quick check prevents confusion:

$$[\Lambda_{\text{eff}}^{[\text{obs}]}] = L^{-1}, \qquad [\Lambda_{\text{cosmos}}] = [\Lambda] = [\Lambda_{\text{eff}}] = L^{-2}.$$

Any appearance of L^{-1} belongs to spectral analysis; any appearance of L^{-2} belongs to cosmology/chronogeometry.

(ix) **Pointers to usage.** For the cosmology-sector effective constant and its placement in the Einstein equations, see eq. (11.269) in Chapter 11; for the residual energy density and its identification $\varepsilon_\Lambda = \rho_{\mathrm{DE}}$, see eq. (17.15) in Chapter 17.

Definition 4.576 (Effective Spectral Cutoff)**.** Fix an admissible Spectral Operator $\hat{D}$ with (generalised) eigenvalues $\{\lambda_n\}_{n\in\mathbb{N}}$ ordered so that $\lambda_{n+1} \geq \lambda_n$ and $\lim_{n\to\infty} \lambda_n = +\infty$. (Definition 4.343; cf. [369, 394]), and fix two loci $x, x' \in \mathcal{X}_q$. For any observable built from Fourfold data—either the *spectral-shear* features $f_n^{(\mathrm{spec})}(x,x') := |\mathcal{S}_n(x,x')|$ (Definitions 4.402 and 4.543) or the *vacuum* features $f_n^{(\mathrm{vac})}(x,x') := |C_n(x) - C_n(x')|$ (Definitions 4.463 and 4.535)—and for modal weights $w_n(x,x')$ (Definition 4.392), define for each tolerance $\varepsilon > 0$ the *effective spectral cutoff*

$$\Lambda_{\mathrm{eff}}^{[\mathrm{obs}]}(x,x';\varepsilon) \;:=\; \inf\Big\{\Lambda > 0 :\; \forall\, n\, \Big(\lambda_n > \Lambda \;\Rightarrow\; w_n(x,x')\, f_n^{(\mathrm{obs})}(x,x') < \varepsilon\Big)\Big\},$$

with obs $\in \{\mathrm{spec}, \mathrm{vac}\}$. Equivalently, letting

$$N_{\mathrm{eff}}^{[\mathrm{obs}]}(x,x';\varepsilon) \;:=\; \min\Big\{N \in \mathbb{N} :\; \forall\, n > N,\; w_n(x,x')\, f_n^{(\mathrm{obs})}(x,x') < \varepsilon\Big\},$$

we have $\Lambda_{\mathrm{eff}}^{[\mathrm{obs}]}(x,x';\varepsilon) = \lambda_{N_{\mathrm{eff}}^{[\mathrm{obs}]}(x,x';\varepsilon)}$. For fixed (x,x') and observable, $\varepsilon \mapsto \Lambda_{\mathrm{eff}}^{[\mathrm{obs}]}(x,x';\varepsilon)$ is nonincreasing and right–continuous. The construction is gauge–covariant and basis–independent because w_n and the absolute features $|\mathcal{S}_n|$, $|C_n(x) - C_n(x')|$ are invariant under admissible unitary frame rotations (Lemma 4.270; cf. [4]).

Lemma 4.577 (Existence and Estimate of the Cutoff)**.** *Assume modal weight decay and feature growth bounds*

$$w_n(x,x') \;\leq\; \frac{C'}{\lambda_n^{\alpha}}, \qquad f_n(x,x') \;\leq\; C_f\, \lambda_n^{\beta},$$

for some constants $C', C_f > 0$ (uniform on the (x,x')–domain under consideration), with exponent gap $\alpha > \beta > -\infty$. Then for every $\varepsilon > 0$ there exists $N \in \mathbb{N}$ such that for all $n > N$ one has $w_n(x,x')\, f_n(x,x') < \varepsilon$, hence $\Lambda_{\mathrm{eff}}(x,x';\varepsilon) = \lambda_N$ is finite. Moreover,

$$\boxed{\Lambda_{\mathrm{eff}}(x,x';\varepsilon) \;\leq\; \max\left\{\lambda_{n_0},\; \left(\frac{C'C_f}{\varepsilon}\right)^{\frac{1}{\alpha-\beta}}\right\},} \tag{4.278}$$

where n_0 is any index beyond which the stated bounds hold. In particular, when the bounds hold from the outset,

$$\Lambda_{\mathrm{eff}}(x,x';\varepsilon) \;\leq\; \left(\frac{C'C_f}{\varepsilon}\right)^{\frac{1}{\alpha-\beta}}.$$

Proof. From the hypotheses, $w_n(x,x')\, f_n(x,x') \leq (C'C_f)\, \lambda_n^{-(\alpha-\beta)}$. Since $\alpha - \beta > 0$, choose N so that for all $n > N$, $(C'C_f)\, \lambda_n^{-(\alpha-\beta)} < \varepsilon$, i.e. $\lambda_n > (C'C_f/\varepsilon)^{1/(\alpha-\beta)}$. With N minimal, $\Lambda_{\mathrm{eff}} = \lambda_N$ and (4.278) follows. Right–continuity and monotonicity in ε are immediate from the definition of the infimum over a nested family of tail conditions. Gauge–covariance and basis–independence are inherited from the invariance of w_n and absolute features under admissible unitary changes. □

Remark 4.578 (Uniform Control on Regions and Consistency with IQR Regularity)**.** If the constants C', C_f and exponents (α, β) can be chosen uniformly for (x, x') in a set $K \subset \mathcal{X}_q \times \mathcal{X}_q$ (e.g. by Spectral/Relational Smoothness and Endogenous Regularization; Axiom 4.415 and Section 4.9.13), then

$$\sup_{(x,x')\in K} \Lambda_{\text{eff}}(x, x'; \varepsilon) \;\le\; \max\left\{\Lambda_0(K),\; \left(\frac{C'C_f}{\varepsilon}\right)^{\frac{1}{\alpha-\beta}}\right\},$$

for some finite $\Lambda_0(K)$ absorbing the finite–index onset of the bounds. Thus a *uniform* effective cutoff exists on K, ensuring that the ℓ^2–aggregates defining $d_{\text{spec}}, t_{\text{spec}}, d_{\text{vac}}, t_{\text{vac}}$ admit a controlled ultraviolet truncation consistent with the IQR finiteness postulates.

Remark 4.579 (Calibration and Observable Dependence)**.** The cutoff carries an explicit tolerance ε and an implicit observable label [obs]. If f_n is rescaled by a factor $\kappa > 0$ (change of units or calibration), then $\Lambda_{\text{eff}}^{[\text{obs}]}(x, x'; \varepsilon)$ rescales as $\kappa^{1/(\alpha-\beta)}$ according to (4.278). In particular, at the *relational null* locus $D(x, x') = 0$ (Sections 4.11.9 and 4.12.8), one has $f_n(x, x') \equiv 0$ and therefore $\Lambda_{\text{eff}}(x, x'; \varepsilon) = 0$ for every $\varepsilon > 0$.

4.14.1 Endogenous Finiteness of all Emergent Relational Observables

Proposition 4.580 (Endogenous Finiteness of all Emergent Relational Observables)**.** *Assume* modal weight decay *(Finite Entanglement Capacity; Principle 4.1 and Definition 4.392),* Spectral Smoothness *(Axiom 4.415), and* Relational Smoothness *(Axiom 4.415). Then all emergent relational distances $d_{\text{corr}}, d_{\text{vac}}, d_{\text{spec}}$ and their temporal counterparts $t_{\text{corr}}, t_{\text{vac}}, t_{\text{spec}}$ are finite for every pair of loci $x, x' \in \mathcal{X}_q$. Moreover, for each tolerance $\varepsilon > 0$ there exists a finite, state–dependent* effective spectral cutoff $\Lambda_{\text{eff}}(x, x'; \varepsilon)$ *as in Definition 4.576 and Lemma 4.577 such that modal contributions beyond Λ_{eff} are negligible. In particular,*

$$d_{\text{spec}}(x, x'),\; d_{\text{vac}}(x, x'),\; t_{\text{spec}}(x, x'),\; t_{\text{vac}}(x, x') \;<\; \infty,$$

and the correlator–based pair $d_{\text{corr}}, t_{\text{corr}}$ are bounded by construction.

Proof of Proposition 4.580. We verify finiteness for the six observables.

(1) Correlator–based distance and time. By definition, $d_{\text{corr}}(x, x')$ and $t_{\text{corr}}(x, x')$ are formed from normalised bilocal correlator expectations (phase and magnitude). Hence $0 \le d_{\text{corr}} \le 1$ and $|t_{\text{corr}}| \le \pi$; in particular, both are finite.

(2) Spectral distance and spectral time. Write

$$d^2_{\text{spec}}(x, x') = \sum_n w_n^2(x, x')\, |\mathcal{S}_n(x, x')|^2, \qquad t^2_{\text{spec}}(x, x') = \sum_n w_n^2(x, x') \left(\operatorname{Im} \mathcal{S}_n(x, x')\right)^2.$$

Spectral Smoothness gives $|\mathcal{S}_n(x, x')| \le C\, \lambda_n^k$ on the domain considered (Axiom 4.415). Finite Entanglement Capacity yields $w_n(x, x') \le C'\, \lambda_n^{-\alpha}$ with $\alpha > k + \frac{1}{2}$ (Principle 4.1). Therefore

$$w_n^2\, |\mathcal{S}_n|^2 \;\le\; (C'C)^2\, \lambda_n^{-2(\alpha-k)},$$

and the series converges by the p–test since $2(\alpha - k) > 1$. The same bound controls the imaginary–part aggregate, hence $d_{\text{spec}}, t_{\text{spec}} < \infty$. Existence of a finite tail threshold $\Lambda_{\text{eff}}(x, x'; \varepsilon)$ follows from Definition 4.576 and Lemma 4.577 (see also the bound eq. (4.278)).

(3) Vacuum–response distance and time. Let $C_n(x) := \langle \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_x)[\psi_n^*, \psi_n] \rangle$ and set

$$d_{\text{vac}}^2(x, x') = \sum_n w_n^2(x, x') \, |C_n(x) - C_n(x')|^2,$$

$$t_{\text{vac}}^2(x, x') = \sum_n w_n^2(x, x') \left[\text{Im}\Big(C_n(x) - C_n(x') \Big) \right]^2.$$

Relational Smoothness gives $|C_n(x) - C_n(x')| \le C'' \, \lambda_n^m$ (Axiom 4.415). With the same weight decay $w_n \le C' \lambda_n^{-\alpha}$ and $\alpha > m + \frac{1}{2}$, we obtain

$$w_n^2 \, |C_n(x) - C_n(x')|^2 \;\le\; (C'C'')^2 \, \lambda_n^{-2(\alpha - m)},$$

so both series converge and $d_{\text{vac}}, t_{\text{vac}} < \infty$. Again, an effective tail cutoff $\Lambda_{\text{eff}}(x, x'; \varepsilon)$ exists by Definition 4.576 and Lemma 4.577.

Conclusion. All six observables are finite for every pair (x, x'), and the modal–sum quantities admit an endogenous, state–dependent ultraviolet regularisation via $\Lambda_{\text{eff}}(x, x'; \varepsilon)$. □

Remark 4.581 (The PQS as the minimal relational pixel)**.** At the *relational null* locus $D(x, x') = 0$ (cf. Section 4.14), one has simultaneously $\vec{d}(x, x') = \mathbf{0}$ and $\vec{t}(x, x') = \mathbf{0}$. Thus the Primordial Quantum Singularity class $\mathfrak{B}_x$ is the fundamental indivisible "pixel" of the emergent operator geometry: no spatial, temporal, or spectral distinguishability remains.

4.15 Holsymmetric Wordwise Minimal Scale and the Indivisibility of the PQS

In the holosymmetric calculus, a minimal *relational* scale emerges *endogenously* from the operator dynamics of the Fourfold correlator $\hat{G}_\Lambda$, the wordwise spectral generator $\mathbb{D}^{[\mathbf{W}]}$, and the Holo–State $|\mathring{\mathbf{\Psi}}\rangle$. At this scale every admissible *wordwise* spatial and temporal probe becomes indistinguishable, so the very notion of separation ceases to be meaningful.

Wordwise relational null condition. We collect the (spatial) distinguishability and (temporal) phase–response null conditions into a single statement, relying only on previously introduced wordwise observables.

Definition 4.582 (Wordwise Relational Null Condition)**.** For $x, x' \in \mathcal{X}_q$ and admissible endpoint words $(\mathbf{W}_1, \mathbf{W}_2)$, let

$$\vec{d}^{[\mathbf{W}_1, \mathbf{W}_2]}(x, x') = \Big(d_{\text{corr}}^{[\cdot]}, d_{\text{vac}}^{[\cdot]}, d_{\text{spec}}^{[\cdot]} \Big), \quad \vec{t}^{[\mathbf{W}_1, \mathbf{W}_2]}(x, x') = \Big(t_{\text{corr}}^{[\cdot]}, t_{\text{vac}}^{[\cdot]}, t_{\text{spec}}^{[\cdot]} \Big),$$

with components as in Definitions 4.339, 4.408, 4.468, 4.529, 4.539 and 4.547. We say (x, x') satisfies the *wordwise relational null condition* iff

$$\boxed{\vec{d}^{[\mathbf{W}_1, \mathbf{W}_2]}(x, x') = \mathbf{0} \quad \text{and} \quad \vec{t}^{[\mathbf{W}_1, \mathbf{W}_2]}(x, x') = \mathbf{0}.} \tag{4.279}$$

Equivalently, the composite triadic distance $D^{[\cdot]}(x, x')$ and the three phase–times vanish (cf. Section 4.11.2).

Interpretation (wordwise). $D^{[\cdot]}(x,x')=0$ expresses a *relational collapse*: the off–diagonal channel, the diagonal vacuum profile, and the spectral commutator all assign zero separation, while $t^{[\cdot]}_{\mathrm{corr}} = t^{[\cdot]}_{\mathrm{vac}} = t^{[\cdot]}_{\mathrm{spec}} = 0$ declare that no phase holonomy accumulates along admissible transports (cf. holonomy statements in Definitions 4.533, 4.541 and 4.549). Thus the PQS is the *null–distance/zero–phase* locus of the wordwise operator geometry (cf. Section 4.11.8).

Regularity hypotheses (wordwise recap). We do not introduce new assumptions; we recall the working hypotheses in their wordwise form:

(a) **Wordwise Spectral Smoothness** (Theorem 4.421): the mode–n shear obeys $|\mathcal{S}^{[\cdot]}_n(x,x')| \lesssim (\lambda^{[\mathbf{W}_1]}_n(x))^k$.

(b) **Finite Entanglement Capacity (wordwise)** (Principle 4.2): $w^{[\cdot]}_n(x,x') \lesssim (\lambda^{[\mathbf{W}_1]}_n(x))^{-\alpha}$ with $\alpha > k + \frac{1}{2}$.

(c) **Relational smoothness for the diagonal sector:** the coincident self–correlators vary smoothly so that $|C^{[\mathbf{W}]}_n(x) - C^{[\mathbf{W}]}_n(x')| \lesssim (\lambda^{[\mathbf{W}]}_n)^m$ on patches where the frame is admissible (cf. the hypotheses used to prove finiteness of $d^{[\cdot]}_{\mathrm{vac}}$ and $t^{[\cdot]}_{\mathrm{vac}}$).

4.15.1 Wordwise Effective Spectral Cutoff

Definition 4.583 (Wordwise effective spectral cutoff)**.** Fix $\mathbb{D}^{[\mathbf{W}]}$ with localised eigenvalues $\{\lambda^{[\mathbf{W}]}_n\}$ and loci x, x'. For Fourfold–derived features evaluated at mode n,

$$f^{(\mathrm{spec})[\cdot]}_n(x,x') := \left|\mathcal{S}^{[\cdot]}_n(x,x')\right|, \qquad f^{(\mathrm{vac})[\cdot]}_n(x,x') := \left|C^{[\cdot]}_n(x) - C^{[\cdot]}_n(x')\right|,$$

and wordwise weights $w^{[\cdot]}_n(x,x')$, define for $\varepsilon > 0$:

$$\Lambda^{[\mathrm{obs}],\,[\cdot]}_{\mathrm{eff}}(x,x';\varepsilon) := \inf\left\{\Lambda > 0 :\ \forall\, n\ \left(\lambda^{[\mathbf{W}_1]}_n > \Lambda \Rightarrow w^{[\cdot]}_n\, f^{(\mathrm{obs})[\cdot]}_n < \varepsilon\right)\right\}, \quad \mathrm{obs} \in \{\mathrm{spec}, \mathrm{vac}\}.$$

This quantity is basis–independent and word/gauge–covariant because $w^{[\cdot]}_n$ and the absolute features are.

Lemma 4.584 (Existence and estimate (wordwise))**.** *If $w^{[\cdot]}_n \le C'(\lambda^{[\mathbf{W}_1]}_n)^{-\alpha}$ and $f^{(\mathrm{obs})[\cdot]}_n \le C_f(\lambda^{[\mathbf{W}_1]}_n)^{\beta}$ with $\alpha > \beta$ on the patch, then for every $\varepsilon > 0$,*

$$\boxed{\Lambda^{[\mathrm{obs}],\,[\cdot]}_{\mathrm{eff}}(x,x';\varepsilon) \ \le\ \left(\frac{C'C_f}{\varepsilon}\right)^{\frac{1}{\alpha-\beta}},} \tag{4.280}$$

hence $\Lambda^{[\mathrm{obs}],\,[\cdot]}_{\mathrm{eff}}$ is finite and nonincreasing in ε.

Proof. For $\lambda^{[\mathbf{W}_1]}_n > \Lambda$,

$$w^{[\cdot]}_n\, f^{(\mathrm{obs})[\cdot]}_n \ \le\ C'(\lambda^{[\mathbf{W}_1]}_n)^{-\alpha}\cdot C_f(\lambda^{[\mathbf{W}_1]}_n)^{\beta} \ =\ (C'C_f)\,(\lambda^{[\mathbf{W}_1]}_n)^{-(\alpha-\beta)}.$$

Choose $\Lambda = \left(\frac{C'C_f}{\varepsilon}\right)^{1/(\alpha-\beta)}$. Then for all n with $\lambda^{[\mathbf{W}_1]}_n > \Lambda$ we have $w^{[\cdot]}_n\, f^{(\mathrm{obs})[\cdot]}_n < \varepsilon$. Monotonicity in ε is immediate from the definition of the infimum. □

Remark 4.585 (Uniform control and equivalence of presentations)**.** If the constants and exponents are uniform on a region $K \subset \mathcal{X}_q \times \mathcal{X}_q$, then $\sup_{(x,x')\in K} \Lambda_{\mathrm{eff}}^{[\mathrm{obs}],[\cdot]}(x,x';\varepsilon)$ is finite. By the componentwise equivalence results between wordwise and exogenous constructions (*e.g.*, for correlator/vacuum/spectral components), the existence and bounds for Λ_{eff} coincide across presentations.

4.15.2 Endogenous Finiteness of All Holosymmetric Wordwise Observables

Proposition 4.586 (Coherent finiteness (wordwise))**.** *Under Theorem 4.421 and Principle 4.2 and the diagonal smoothness hypothesis, all six wordwise quantities*

$$d^{[\cdot]}_{\mathrm{corr}},\ d^{[\cdot]}_{\mathrm{vac}},\ d^{[\cdot]}_{\mathrm{spec}},\quad t^{[\cdot]}_{\mathrm{corr}},\ t^{[\cdot]}_{\mathrm{vac}},\ t^{[\cdot]}_{\mathrm{spec}}$$

are finite for every (x,x'), and for each $\varepsilon > 0$ admit a finite $\Lambda_{\mathrm{eff}}^{[\mathrm{obs}],[\cdot]}(x,x';\varepsilon)$ as in Definition 4.583 and Lemma 4.584.

Proof. The spectral sector follows from Wordwise Spectral Smoothness and finite capacity, which give an ℓ^2–summable UV tail for both absolute and imaginary commutator responses; the correlator and vacuum sectors are analogous, using the diagonal smoothness and the same capacity decay. Temporal components obey $|\operatorname{Im} z| \le |z|$, hence are bounded by their spatial counterparts. The bound (4.280) yields pointwise (modewise) UV control for any tolerance ε, establishing finiteness and effective band–limitation. □

Remark 4.587 (PQS as the minimal wordwise "pixel")**.** At the relational null locus (4.279) one has $D^{[\cdot]}(x,x') = 0$ and $t^{[\cdot]}_{\mathrm{corr}}(x,x') = t^{[\cdot]}_{\mathrm{vac}}(x,x') = t^{[\cdot]}_{\mathrm{spec}}(x,x') = 0$. Thus the PQS class is the indivisible *wordwise pixel* of the operator geometry: no spatial magnitude, no temporal phase, and no spectral shear remain resolvable— a statement invariant under endpoint word changes and parallel–transport gauges.

Consequences (wordwise). (i) The minimal scale is *derived*: the joint zero locus of $\left(\vec{d}^{[\cdot]}, \vec{t}^{[\cdot]}\right)$ is reached without external cutoffs (Proposition 4.426). (ii) If the spectrum is static on a patch, then $d^{[\cdot]}_{\mathrm{spec}} = t^{[\cdot]}_{\mathrm{spec}} = 0$ there, so any residual granularity is correlator/vacuum–driven. (iii) Nontrivial wordwise holonomies (correlator/vacuum/spectral) obstruct collapse to the PQS; detecting such holonomy certifies departure from the minimal relational scale.

4.15.3 The Ultra-Sub-Planckian Nature of the Singularity

In contrast to approaches that *postulate* a fundamental length, the IQR framework derives its minimal relational scale from operator-theoretic structure. The Planck length,

$$\ell_{\mathrm{P}} = \sqrt{\hbar G/c^3},$$

is a dimensional construct of semiclassical origin—useful as an *effective* threshold for the breakdown of classical geometry and low-energy quantum field theory, but not an intrinsic cutoff on microscopic relational observables (*cf.* [284, 415, 403, 416, 83, 40]). In IQR, the scale that limits distinguishability is the *coherence width* of a Primordial Quantum Singularity (PQS), denoted ℓ_{PQS}, which is extracted from the state-dependent spectral tail of relational probes.

The PQS Coherence Scale is Relational and State-Dependent

Distances and times in IQR are emergent from Holo–State expectations of the Fourfold correlator $\hat{G}_\Lambda$ and the spectral generator $\hat{D}$. Their UV behaviour is governed by modal weights $w_n(x,x')$ and feature amplitudes $f_n^{(\mathrm{obs})}(x,x')$ (spectral shear or vacuum variation), whose product controls the contribution of the n-th mode. Following Definition 4.576, for any tolerance $\varepsilon > 0$ we define the *effective spectral cutoff*

$$\Lambda_{\mathrm{eff}}^{[\mathrm{obs}]}(x,x';\varepsilon) \quad \text{by} \quad \forall\, \lambda_n > \Lambda_{\mathrm{eff}} : \quad w_n(x,x')\, f_n^{(\mathrm{obs})}(x,x') < \varepsilon,$$

and the corresponding PQS coherence width

$$\ell_{\mathrm{PQS}}^{[\mathrm{obs}]}(x,x';\varepsilon) := \left(\Lambda_{\mathrm{eff}}^{[\mathrm{obs}]}(x,x';\varepsilon)\right)^{-1}.$$

By Lemma 4.577, Λ_{eff} exists and is finite under the IQR regularity hypotheses (Spectral/Relational Smoothness and Finite Entanglement Capacity). Importantly, $\varepsilon \mapsto \Lambda_{\mathrm{eff}}(\varepsilon)$ is nonincreasing: stricter tolerances push the cutoff deeper into the UV, thus *reducing* ℓ_{PQS}.

Proposition 4.588 (Sufficient Condition for Sub-Planckian PQS)**.** *Fix an observable label* $\mathrm{obs} \in \{\mathrm{spec}, \mathrm{vac}\}$ *and two loci* (x,x')*. Suppose there exists a tolerance* $\varepsilon_* > 0$ *such that the Planck-scale band remains non-negligible in the Holo–State,*

$$\sup_{\lambda_n \geq \ell_{\mathrm{P}}^{-1}} w_n(x,x')\, f_n^{(\mathrm{obs})}(x,x') \;\geq\; \varepsilon_*. \tag{4.281}$$

Then for every $0 < \varepsilon \leq \varepsilon_*$*,*

$$\Lambda_{\mathrm{eff}}^{[\mathrm{obs}]}(x,x';\varepsilon) \;\geq\; \ell_{\mathrm{P}}^{-1} \qquad \Longrightarrow \qquad \ell_{\mathrm{PQS}}^{[\mathrm{obs}]}(x,x';\varepsilon) \;\leq\; \ell_{\mathrm{P}}.$$

In words: whenever Planck-scale modes are still operationally non-negligible *in the state (with respect to the chosen observable), the PQS coherence width is* sub-Planckian *for all tolerances that register those modes.*

Proof (sketch). If (4.281) holds and $0 < \varepsilon \leq \varepsilon_*$, then it cannot be the case that $w_n f_n^{(\mathrm{obs})} < \varepsilon$ for *all* $\lambda_n > \ell_{\mathrm{P}}^{-1}$; hence the defining tail condition in Definition 4.576 cannot be satisfied with any cutoff smaller than ℓ_{P}^{-1}. Therefore $\Lambda_{\mathrm{eff}} \geq \ell_{\mathrm{P}}^{-1}$ and the stated bound on ℓ_{PQS} follows. □

Typical Regime (Using IQR Bounds). Under Spectral/Relational Smoothness and weight decay, $w_n f_n^{(\mathrm{obs})} \lesssim C' C_{\mathrm{obs}}\, \lambda_n^{-(\alpha-\beta_{\mathrm{obs}})}$ with $\alpha - \beta_{\mathrm{obs}} > 0$ (cf. Axiom 4.415 and Principle 4.1). Evaluated at the Planck band one has the coarse upper control

$$\sup_{\lambda_n \geq \ell_{\mathrm{P}}^{-1}} w_n f_n^{(\mathrm{obs})} \;\lesssim\; C' C_{\mathrm{obs}}\, \ell_{\mathrm{P}}^{\alpha-\beta_{\mathrm{obs}}}.$$

Thus any tolerance ε that treats Planck-band contributions as *detectable* (i.e. ε not smaller than the actual state-dependent value of the left-hand side) will satisfy the hypothesis of Proposition 4.588. In that physically natural regime—where Planck-scale modes have *not* already decohered away in the Holo–State—the emergent PQS width obeys $\ell_{\mathrm{PQS}} \leq \ell_{\mathrm{P}}$.

Conceptual Upshot. The Planck scale is a semiclassical benchmark ([284, 415, 403, 416, 83]), not a microscopic prohibition. IQR replaces the *postulated* minimal length with a *derived*, state-dependent coherence width ℓ_{PQS} that can be driven strictly below ℓ_{P} whenever Planck-band modes remain operational in the relational data of $|\mathring{\mathbf{\Psi}}\rangle$. In short, *sub-Planckian* PQS structure is natural in IQR: it arises endogenously from modal weights and spectral/relational features, not from any external cutoff or background spacetime assumption.

From Sub–Planckian to Ultra–Sub–Planckian. The weak statement established above identifies "sub–Planckian" PQS width as a *cutoff comparison* at fixed observable and tolerance ε: by Definition 4.576,

$$\ell_{\mathrm{PQS}}(x,x';\varepsilon) \;\le\; \ell_{\mathrm{P}} \quad\Longleftrightarrow\quad \Lambda_{\mathrm{eff}}(x,x';\varepsilon) \;\ge\; \ell_{\mathrm{P}}^{-1}.$$

Proposition Proposition 4.588 then supplies a concrete, operational criterion: if Planck–band modes remain non–negligible in the Holo–State (with respect to the chosen observable), i.e.

$$\sup_{\lambda_n \ge \ell_{\mathrm{P}}^{-1}} w_n(x,x')\, f_n^{(\mathrm{obs})}(x,x') \;\ge\; \varepsilon_*,$$

then for every $0 < \varepsilon \le \varepsilon_*$ one necessarily has $\Lambda_{\mathrm{eff}} \ge \ell_{\mathrm{P}}^{-1}$ and hence $\ell_{\mathrm{PQS}} \le \ell_{\mathrm{P}}$. In this sense, the Planck length $\ell_{\mathrm{P}} = \sqrt{\hbar G/c^3}$ remains a *semiclassical benchmark* [284, 415, 403, 416, 83], while the actual relational limit ℓ_{PQS} is *derived* from state–dependent spectral tails.

Prelude to the Strong (Ultra–Sub–Planckian) Regime. The title's "ultra–sub–Planckian" claim strengthens the weak comparison into a *parametric hierarchy*: there exists a gap factor $\kappa \gg 1$ and a tolerance window $\varepsilon \in [\varepsilon_{\min}, \varepsilon_{\max}]$ such that

$$\ell_{\mathrm{PQS}}(x,x';\varepsilon) \;\le\; \frac{\ell_{\mathrm{P}}}{\kappa} \qquad\Longleftrightarrow\qquad \Lambda_{\mathrm{eff}}(x,x';\varepsilon) \;\ge\; \kappa\, \ell_{\mathrm{P}}^{-1}.$$

Operationally, this requires *sustained* UV support well beyond the Planck band,

$$\sup_{\lambda_n \ge \kappa\, \ell_{\mathrm{P}}^{-1}} w_n(x,x')\, f_n^{(\mathrm{obs})}(x,x') \;\ge\; \varepsilon_{\max},$$

so that any tolerance $\varepsilon \le \varepsilon_{\max}$ forces the effective cutoff to sit at or above $\kappa\, \ell_{\mathrm{P}}^{-1}$. In the next step we formalise this as a sufficient "ultra–" condition built on the monotonicity of $\varepsilon \mapsto \Lambda_{\mathrm{eff}}(x,x';\varepsilon)$ from Definition 4.576, thereby exhibiting a robust, state–dependent separation $\ell_{\mathrm{PQS}} \ll \ell_{\mathrm{P}}$ whenever the Holo–State retains appreciable relational signal in the super–Planckian tail.

Correlator Persistence Above the PQS Scale

Lemma 4.589 (Correlator Persistence Above the PQS Scale)**.** *Let ℓ_{P} denote the Planck length, and let $\ell_{\mathrm{PQS}} := \lambda_{\mathrm{cutoff}}^{-1}$ be the PQS coherence width defined by the effective cutoff (Definition 4.576). Then:*

(1) *For separations* $|x - x'| \sim \ell_{\mathrm{P}} \gg \ell_{\mathrm{PQS}}$, *the bilocal correlator is* not *uniformly suppressed by the effective cutoff. In particular, writing*

$$\langle \mathring{\mathbf{\Psi}} \mid \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'}) \mid \mathring{\mathbf{\Psi}} \rangle = \sum\nolimits_n w_n(x, x')\, \mathrm{e}^{\mathrm{i}\theta_n(x,x')},$$

if there exists $\delta \in (0, \pi)$ *and a set of indices* $\mathcal{N}_\delta$ *with* $\lambda_n < \lambda_{\mathrm{cutoff}}$ *and total weight* $W_\delta := \sum_{n\in\mathcal{N}_\delta} w_n(x, x') > 0$ *such that* $\theta_n(x, x') \in [-\delta, \delta]$ *for all* $n \in \mathcal{N}_\delta$, *then*

$$\left|\langle \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})\rangle\right| \;\geq\; \cos\delta\, W_\delta \;>\; 0.$$

(2) *For separations* $|x - x'| \lesssim \ell_{\mathrm{PQS}}$, *one has tail suppression: for every* $\varepsilon > 0$ *there exists* N *such that for all* $n > N$, $w_n(x, x') < \varepsilon$; *hence, by Definition 4.576,*

$$\left|\langle \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})\rangle\right| \;\leq \sum_{\lambda_n \geq \lambda_{\mathrm{cutoff}}} w_n(x, x') \;<\; \varepsilon.$$

In particular, $\left|\langle \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})\rangle\right| \to 0$ *as the tolerance defining* $\lambda_{\mathrm{cutoff}}$ *is tightened.*

Proof. The modal expansion is immediate from the spectral decomposition relative to $\hat{D}$:

$$\langle \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})\rangle = \sum_n w_n(x, x')\, \mathrm{e}^{\mathrm{i}\theta_n(x,x')}, \quad w_n(x, x') := \left\| \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_n^*, \psi_n]\, \mathring{\mathbf{\Psi}} \right\|.$$

For *(1)*, when $\ell_{\mathrm{P}} \gg \ell_{\mathrm{PQS}}$ the contributing modes below $\lambda_{\mathrm{cutoff}}$ carry non-negligible total weight. Under the phase–coherence hypothesis stated, the triangle inequality yields $\left|\sum_{n\in\mathcal{N}_\delta} w_n \mathrm{e}^{\mathrm{i}\theta_n}\right| \geq \cos\delta \sum_{n\in\mathcal{N}_\delta} w_n = \cos\delta\, W_\delta > 0$. For *(2)*, by Definition 4.576 and the decay of modal weights in the tail, for any $\varepsilon > 0$ one has $w_n(x, x') < \varepsilon$ for all $\lambda_n \geq \lambda_{\mathrm{cutoff}}$, whence $\left|\sum_{\lambda_n\geq\lambda_{\mathrm{cutoff}}} w_n \mathrm{e}^{\mathrm{i}\theta_n}\right| \leq \sum_{\lambda_n\geq\lambda_{\mathrm{cutoff}}} w_n < \varepsilon$. This gives the stated suppression. □

Ultra–Sub–Planckian Tail–Persistence and Parametric PQS Shrinkage

Lemma 4.590 (Ultra–Sub–Planckian Tail–Persistence ⇒ Parametric PQS Shrinkage)**.** *Fix an observable label* [obs] *(spectral shear or vacuum variation) with modal features* $f_n^{(\mathrm{obs})}(x, x')$ *and weights* $w_n(x, x')$, *and let the effective cutoff* $\Lambda_{\mathrm{eff}}^{[\mathrm{obs}]}(x, x'; \varepsilon)$ *be as in Definition 4.576. Let* $\ell_{\mathrm{P}} = \sqrt{\hbar G/c^3}$ *denote the Planck length and* $\kappa > 1$ *a dimensionless separation factor.*

Assume there exists a tolerance $\varepsilon_{\max} > 0$ *and a pair* (x, x') *such that the* UV tail is non–negligible above the Planck band by a factor κ, *i.e.*

$$\sup_{\lambda_n \;\geq\; \kappa\, \ell_{\mathrm{P}}^{-1}} \left[w_n(x, x')\, f_n^{(\mathrm{obs})}(x, x') \right] \;\geq\; \varepsilon_{\max}. \tag{4.282}$$

Then, for every tolerance $\varepsilon \in (0, \varepsilon_{\max}]$,

$$\Lambda_{\mathrm{eff}}^{[\mathrm{obs}]}(x, x'; \varepsilon) \;\geq\; \kappa\, \ell_{\mathrm{P}}^{-1} \qquad \Longleftrightarrow \qquad \ell_{\mathrm{PQS}}^{[\mathrm{obs}]}(x, x'; \varepsilon) \;:=\; \left(\Lambda_{\mathrm{eff}}^{[\mathrm{obs}]}(x, x'; \varepsilon)\right)^{-1} \;\leq\; \frac{\ell_{\mathrm{P}}}{\kappa}. \tag{4.283}$$

In particular, the PQS coherence width is uniformly *(in* ε*) separated from the Planck length by the factor* κ *on the whole window* $\varepsilon \in (0, \varepsilon_{\max}]$.

Proof. By definition of $\Lambda_{\mathrm{eff}}^{[\mathrm{obs}]}$ (Definition 4.576), for a given $\varepsilon > 0$ it is the smallest spectral scale beyond which *all* tail contributions satisfy $w_n\, f_n^{(\mathrm{obs})} < \varepsilon$. Condition (4.282) states that at least one mode with $\lambda_n \geq \kappa\, \ell_{\mathrm{P}}^{-1}$ *violates* the tail–smallness requirement at level $\varepsilon_{\max}$. Hence at $\varepsilon = \varepsilon_{\max}$ the minimal spectral threshold that renders the tail uniformly small must lie *to the right* of that mode, i.e. $\Lambda_{\mathrm{eff}}^{[\mathrm{obs}]}(x, x'; \varepsilon_{\max}) \geq \kappa\, \ell_{\mathrm{P}}^{-1}$. Monotonicity $\varepsilon \mapsto \Lambda_{\mathrm{eff}}^{[\mathrm{obs}]}(x, x'; \varepsilon)$ (nonincreasing) then implies $\Lambda_{\mathrm{eff}}^{[\mathrm{obs}]}(x, x'; \varepsilon) \geq \Lambda_{\mathrm{eff}}^{[\mathrm{obs}]}(x, x'; \varepsilon_{\max}) \geq \kappa\, \ell_{\mathrm{P}}^{-1}$ for all $\varepsilon \in (0, \varepsilon_{\max}]$, yielding (4.283). □

Corollary 4.591 (Strong Ultra–Sub–Planckian PQS)**.** *If* (4.282) *holds for* each *of the two diagnostic observables (spectral and vacuum), then both coherence widths satisfy*

$$\ell_{\mathrm{PQS}}^{[\mathrm{spec}]}(x, x'; \varepsilon) \;\leq\; \frac{\ell_{\mathrm{P}}}{\kappa}, \qquad \ell_{\mathrm{PQS}}^{[\mathrm{vac}]}(x, x'; \varepsilon) \;\leq\; \frac{\ell_{\mathrm{P}}}{\kappa} \quad \textit{for all} \;\; \varepsilon \in (0, \varepsilon_{\max}].$$

Consequently, every relational observable built from these sectors is supported on scales $\ll \ell_{\mathrm{P}}$ in the strong, parametric sense.

Remark 4.592 (Robustness Across Tolerances and Calibration)**.** The windowed nature of Lemma 4.590 ensures that the separation $\ell_{\mathrm{PQS}} \leq \ell_{\mathrm{P}}/\kappa$ is not a fine–tuned statement at a single tolerance, but persists across the entire interval $(0, \varepsilon_{\max}]$. Changes of observable normalization or units rescale Λ_{eff} according to eq. (4.278) (cf. Remark 4.579), leaving the *inequality form* (4.283) intact once κ and the tolerance window are re-expressed in the new calibration.

Remark 4.593 (Link to Correlator Persistence at the Planck Band)**.** The tail–persistence hypothesis (4.282) is compatible with—and strengthens—the correlator non-vanishing at Planckian separations established in Lemma 4.589: Planck–band separations ($\sim \ell_{\mathrm{P}}$) retain finite bilocal amplitude, while sub–PQS separations ($\lesssim \ell_{\mathrm{PQS}}$) suppress it. Under (4.282), the suppression threshold is parametrically beyond ℓ_{P}^{-1}, certifying an *ultra–sub–Planckian* PQS width.

The Ultra–Sub–Planckian Nature of PQS

We present four mutually reinforcing lines of argument that refine the weak, "sub–Planckian'' statement and prepare the ground for the strong, parametric ("ultra–'') separation.

(1) Modal spectrum scaling. By Finite Entanglement Capacity the modal weights obey $w_n \lesssim \lambda_n^{-\alpha}$ (Principle 4.1), while for admissible elliptic $\hat{D}$ one has Weyl–type growth $\lambda_n \asymp C\, n^{1/d_{\mathrm{eff}}}$ (effective spectral dimension; cf. [394, 369]). The effective cutoff Λ_{eff} from Definition 4.576 marks the tail where $w_n f_n^{(\mathrm{obs})} < \varepsilon$. If the Holo–State retains non–negligible signal at the Planck band (so that $\Lambda_{\mathrm{eff}} \geq \ell_{\mathrm{P}}^{-1}$), then the PQS width

$$\ell_{\mathrm{PQS}} \;:=\; \Lambda_{\mathrm{eff}}^{-1}$$

satisfies $\ell_{\mathrm{PQS}} \leq \ell_{\mathrm{P}}$ (weak case). If, moreover, the signal persists a factor $\kappa \gg 1$ deeper into the UV (Lemma 4.590), then $\Lambda_{\mathrm{eff}} \geq \kappa\, \ell_{\mathrm{P}}^{-1}$ and hence

$$\ell_{\mathrm{PQS}} \;\leq\; \frac{\ell_{\mathrm{P}}}{\kappa} \;\ll\; \ell_{\mathrm{P}} \quad \text{(strong, ultra–sub–Planckian)}.$$

(2) Operator–spectral suppression. Define the cutoff by the tail condition

$$w_n^2(x,x')\,|\mathcal{S}_n(x,x')|^2 \;\le\; \varepsilon \qquad (\lambda_n > \Lambda_{\text{eff}}).$$

Under Spectral Smoothness $|\mathcal{S}_n| \lesssim \lambda_n^{\beta}$ (Axiom 4.415) and $w_n \lesssim \lambda_n^{-\alpha}$ with $\alpha > \beta + \frac{1}{2}$ (Principle 4.1), Lemma 4.577 yields a finite, ε–dependent Λ_{eff} with the pointwise estimate (4.278). Thus tightening tolerance pushes Λ_{eff} arbitrarily high, making $\ell_{\text{PQS}} = \Lambda_{\text{eff}}^{-1}$ arbitrarily small. When Planck–band contributions are *detectable* (non–negligible), one has $\Lambda_{\text{eff}} \ge \ell_{\text{P}}^{-1}$, so $\ell_{\text{PQS}} \le \ell_{\text{P}}$; if the persistence extends beyond $\kappa\,\ell_{\text{P}}^{-1}$, one obtains the ultra–sub–Planckian bound (Lemma 4.590). (For the role of $\ell_{\text{P}} = \sqrt{\hbar G/c^3}$ as a semiclassical benchmark rather than a fundamental cutoff, see [284, 415, 403].)

(3) Breakdown of geometry below ℓ_{PQS}. The expectation–valued metric $\bar{g}_{\mu\nu}$ is meaningful only where modal coherence persists. For $|x-x'| < \ell_{\text{PQS}}$ one has tail suppression $w_n(x,x') \to 0$ $(\forall n)$, hence

$$\left|\langle \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})\rangle\right| = \left|\sum\nolimits_n w_n(x,x')\,\mathrm{e}^{\mathrm{i}\theta_n}\right| \;\longrightarrow\; 0,$$

so relational geometry dissolves below the PQS width. In contrast, at Planckian separations ($\sim \ell_{\text{P}}$) the correlator retains finite amplitude (Lemma 4.589); therefore geometric structure is supported *above* but not *below* ℓ_{PQS}.

(4) Interpretive principle.

> *Emergent geometry arises only above the modal distinguishability threshold set by the PQS width. The Planck length is not a fundamental bound, but an effective, semiclassical scale emerging from a deeper relational quantum substrate.*

Theorem: Sub– and Ultra–Sub–Planckian Coherence of the PQS

Theorem 4.594 (Sub– and Ultra–Sub–Planckian Coherence of the PQS)**.** *Let $\ell_{\text{PQS}} := \Lambda_{\text{eff}}^{-1}$ be the PQS coherence width from Definition 4.576, and $\ell_{\text{P}} := \sqrt{\hbar G/c^3}$ the Planck length. Under modal weight decay (Principle 4.1) and Spectral Smoothness (Axiom 4.415), if the Holo–State exhibits non–negligible Planck–band support for the chosen observable (weak persistence), then*

$$\ell_{\text{PQS}} \;\le\; \ell_{\text{P}}, \qquad \tau_{\text{PQS}} := \frac{\ell_{\text{PQS}}}{c} \;\le\; t_{\text{P}} := \frac{\ell_{\text{P}}}{c}.$$

If, moreover, the UV tail is non–negligible on $[\kappa\,\ell_{\text{P}}^{-1}, \infty)$ for some $\kappa \gg 1$ (Lemma 4.590), then

$$\ell_{\text{PQS}} \;\le\; \frac{\ell_{\text{P}}}{\kappa} \;\ll\; \ell_{\text{P}}, \qquad \tau_{\text{PQS}} \;\le\; \frac{t_{\text{P}}}{\kappa} \;\ll\; t_{\text{P}}.$$

Proof sketch. Finiteness and monotonicity of $\Lambda_{\text{eff}}(\varepsilon)$ follow from Lemma 4.577. Weak persistence enforces $\Lambda_{\text{eff}} \ge \ell_{\text{P}}^{-1}$, giving the sub–Planckian bound. Ultra persistence (with factor κ) enforces $\Lambda_{\text{eff}} \ge \kappa\,\ell_{\text{P}}^{-1}$ by Lemma 4.590, yielding the parametric (ultra–) separation. □

$$\boxed{\ell_{\mathrm{PQS}} = \Lambda_{\mathrm{eff}}^{-1} \;\le\; \frac{\ell_{\mathrm{P}}}{\max\{1,\kappa\}}\,, \qquad \tau_{\mathrm{PQS}} = \frac{\ell_{\mathrm{PQS}}}{c} \;\le\; \frac{t_{\mathrm{P}}}{\max\{1,\kappa\}}\,.}$$

Thus the minimal relational scale of the PQS is set endogenously by modal coherence and the spectral tail of relational observables; the Planck scale functions as a semiclassical threshold, not a fundamental quantum limit [284, 415, 403].

4.15.4 Wordwise Ultra–Sub–Planckian Nature of the Singularity

In the holosymmetric wordwise calculus, no *a priori* fundamental length is postulated. Instead, the minimal relational scale arises *endogenously* from the operator dynamics of the Fourfold correlator $\hat{G}_\Lambda$, the wordwise spectral generator $\mathbb{D}^{[\mathbf{W}]}$, and the Holo–State $|\mathring{\mathbf{\Psi}}\rangle$. The Planck length $\ell_{\mathrm{P}} = \sqrt{\hbar G/c^3}$ is thus an *effective semiclassical benchmark*—useful for diagnosing the breakdown of classical geometry and low–energy QFT, but not an intrinsic microscopic cutoff (*cf.* [284, 415, 403, 416, 83, 40]).

Wordwise PQS coherence width. For any observable label obs $\in \{\mathrm{spec}, \mathrm{vac}\}$ and loci (x, x'), let the *wordwise effective spectral cutoff* $\Lambda_{\mathrm{eff}}^{[\mathrm{obs}],\,[\mathbf{W}_1,\mathbf{W}_2]}(x, x'; \varepsilon)$ be as in Definition 4.583. Define the corresponding *PQS coherence width*

$$\ell_{\mathrm{PQS}}^{[\mathrm{obs}],\,[\mathbf{W}_1,\mathbf{W}_2]}(x, x'; \varepsilon) \;:=\; \Big(\Lambda_{\mathrm{eff}}^{[\mathrm{obs}],\,[\mathbf{W}_1,\mathbf{W}_2]}(x, x'; \varepsilon)\Big)^{-1}. \tag{4.284}$$

By Lemma 4.584, $\Lambda_{\mathrm{eff}}^{[\cdot]}$ exists and is finite under Wordwise Spectral Smoothness and Finite Entanglement Capacity (Theorem 4.421 and Principle 4.2); moreover, $\varepsilon \mapsto \Lambda_{\mathrm{eff}}^{[\cdot]}(\varepsilon)$ is nonincreasing, so tightening tolerance drives $\ell_{\mathrm{PQS}}^{[\cdot]}$ *down*.

Proposition 4.595 (Wordwise sufficient condition for sub–Planckian PQS)**.** *Fix* obs $\in \{\mathrm{spec}, \mathrm{vac}\}$ *and* (x, x'). *Suppose there exists* $\varepsilon_* > 0$ *such that the Planck band is* operationally non–negligible *in the Holo–State:*

$$\sup_{\lambda_n^{[\mathbf{W}_1]} \,\ge\, \ell_{\mathrm{P}}^{-1}} \Big[w_n^{[\mathbf{W}_1,\mathbf{W}_2]}(x, x')\, f_n^{(\mathrm{obs})[\mathbf{W}_1,\mathbf{W}_2]}(x, x') \Big] \;\ge\; \varepsilon_*, \tag{4.285}$$

where $f_n^{(\mathrm{spec})[\cdot]} := \big|\mathcal{S}_n^{[\cdot]}(x, x')\big|$ *and* $f_n^{(\mathrm{vac})[\cdot]} := \big|C_n^{[\cdot]}(x) - C_n^{[\cdot]}(x')\big|$. *Then for all* $0 < \varepsilon \le \varepsilon_*$,

$$\Lambda_{\mathrm{eff}}^{[\mathrm{obs}],\,[\cdot]}(x, x'; \varepsilon) \;\ge\; \ell_{\mathrm{P}}^{-1} \qquad \Longrightarrow \qquad \ell_{\mathrm{PQS}}^{[\mathrm{obs}],\,[\cdot]}(x, x'; \varepsilon) \;\le\; \ell_{\mathrm{P}}.$$

Hence, whenever Planck–band modes remain detectable (for the chosen observable), the PQS coherence width is sub–Planckian *for every tolerance that registers those modes.*

Proof. Assume (4.285) and fix $0 < \varepsilon \le \varepsilon_*$. If $\Lambda_{\mathrm{eff}}^{[\mathrm{obs}],[\cdot]}(x, x'; \varepsilon) < \ell_{\mathrm{P}}^{-1}$, then by definition of Λ_{eff} we would have $w_n^{[\cdot]} f_n^{(\mathrm{obs})[\cdot]} < \varepsilon \le \varepsilon_*$ for *all* n with $\lambda_n^{[\mathbf{W}_1]} \ge \ell_{\mathrm{P}}^{-1}$, contradicting (4.285). Therefore $\Lambda_{\mathrm{eff}}^{[\mathrm{obs}],[\cdot]} \ge \ell_{\mathrm{P}}^{-1}$, and (4.284) yields $\ell_{\mathrm{PQS}}^{[\mathrm{obs}],[\cdot]} \le \ell_{\mathrm{P}}$. □

Typical regime under IQR bounds (wordwise). Wordwise Spectral/Relational Smoothness and weight decay imply

$$w_n^{[\cdot]}\, f_n^{(\mathrm{obs})[\cdot]} \;\lesssim\; C' C_{\mathrm{obs}} \left(\lambda_n^{[\mathbf{W}_1]}\right)^{-(\alpha-\beta_{\mathrm{obs}})}, \qquad \alpha-\beta_{\mathrm{obs}}>0,$$

so at the Planck band $\sup_{\lambda_n^{[\mathbf{W}_1]}\geq \ell_{\mathrm{P}}^{-1}} w_n^{[\cdot]} f_n^{(\mathrm{obs})[\cdot]} \lesssim C' C_{\mathrm{obs}}\, \ell_{\mathrm{P}}^{\alpha-\beta_{\mathrm{obs}}}$. Any tolerance that treats this as *detectable* satisfies the hypothesis of Proposition 4.595, yielding $\ell_{\mathrm{PQS}}^{[\cdot]} \leq \ell_{\mathrm{P}}$ whenever Planck–band support has not decohered away.

Lemma 4.596 (Ultra–sub–Planckian tail–persistence ⇒ parametric shrinkage)**.** *Let $\kappa>1$ and assume that for some $\varepsilon_{\max}>0$,*

$$\sup_{\lambda_n^{[\mathbf{W}_1]} \,\geq\, \kappa\,\ell_{\mathrm{P}}^{-1}} \left[w_n^{[\cdot]}(x,x')\, f_n^{(\mathrm{obs})[\cdot]}(x,x')\right] \;\geq\; \varepsilon_{\max}. \tag{4.286}$$

Then for every $\varepsilon\in(0,\varepsilon_{\max}]$,

$$\Lambda_{\mathrm{eff}}^{[\mathrm{obs}],[\cdot]}(x,x';\varepsilon) \;\geq\; \kappa\,\ell_{\mathrm{P}}^{-1} \qquad\Longleftrightarrow\qquad \ell_{\mathrm{PQS}}^{[\mathrm{obs}],[\cdot]}(x,x';\varepsilon) \;\leq\; \frac{\ell_{\mathrm{P}}}{\kappa}. \tag{4.287}$$

Thus the PQS coherence width is uniformly separated from ℓ_{P} by the factor κ on the entire tolerance window $(0,\varepsilon_{\max}]$.

Proof. By (4.286), for $\varepsilon\leq\varepsilon_{\max}$ there exists at least one mode with $\lambda_n^{[\mathbf{W}_1]}\geq\kappa\,\ell_{\mathrm{P}}^{-1}$ such that $w_n^{[\cdot]} f_n^{(\mathrm{obs})[\cdot]}\geq\varepsilon$. Hence the tail–smallness condition cannot hold for *all* modes beyond any threshold $<\kappa\,\ell_{\mathrm{P}}^{-1}$. Therefore $\Lambda_{\mathrm{eff}}^{[\mathrm{obs}],[\cdot]}(x,x';\varepsilon)\geq\kappa\,\ell_{\mathrm{P}}^{-1}$, and the bound on $\ell_{\mathrm{PQS}}^{[\cdot]}$ follows from (4.284). □

Remark 4.597 (Correlator persistence vs. PQS suppression (wordwise))**.** Let $\Gamma_{\mathrm{corr}}^{[\cdot]}(x,x') = \sum_n w_n^{[\cdot]}(x,x')\, \mathrm{e}^{i\theta_n^{[\cdot]}}$ be the mode expansion in a wordwise spectral frame. If a subset $\mathcal{N}_\delta$ below the effective cutoff carries total weight $W_\delta>0$ with phases $|\theta_n^{[\cdot]}|\leq\delta<\pi$, then $\left|\Gamma_{\mathrm{corr}}^{[\cdot]}(x,x')\right| \geq \cos\delta\, W_\delta > 0$ (Planckian separations retain signal). Conversely, for separations $\lesssim \ell_{\mathrm{PQS}}^{[\cdot]}$ the tail suppression from $\Lambda_{\mathrm{eff}}^{[\cdot]}$ drives $\left|\Gamma_{\mathrm{corr}}^{[\cdot]}(x,x')\right|\to 0$ as the tolerance is tightened.

Conceptual upshot (wordwise). The Planck length serves as a semiclassical *benchmark* ([284, 415, 403, 416, 83, 40]), while the actual minimal relational scale is the *derived, state–dependent* width $\ell_{\mathrm{PQS}}^{[\cdot]}$ determined by wordwise modal weights and spectral/diagonal features. Whenever Planck–band (or super–Planckian, by Lemma 4.596) modes remain non–negligible, the PQS is *sub–Planckian* (indeed *ultra–sub–Planckian* with a parametric gap) in a way that is gauge–covariant and word–invariant. No external cutoff or background geometry is invoked; the scale emerges from the holosymmetric, state–conditioned operator algebra itself.

Wordwise Ultra–Sub–Planckian Nature of the PQS

We present four mutually reinforcing wordwise arguments that refine the weak "sub–Planckian'' statement and prepare the ground for a strong, parametric ("ultra–'') separation. All symbols are in the holosymmetric wordwise calculus (weights $w_n^{[\mathbf{W}_1,\mathbf{W}_2]}$, features $f_n^{(\mathrm{obs})[\mathbf{W}_1,\mathbf{W}_2]}$, eigenvalues $\lambda_n^{[\mathbf{W}_1]}$); effective cutoffs and widths are as in Definition 4.583.

(1) Modal spectrum scaling (wordwise). By Wordwise Finite Entanglement Capacity, $w_n^{[\cdot]} \lesssim \left(\lambda_n^{[\mathbf{W}_1]}\right)^{-\alpha}$ (Principle 4.2). For admissible (elliptic) $\mathbb{D}^{[\mathbf{W}_1]}$, Weyl-type growth gives $\lambda_n^{[\mathbf{W}_1]} \asymp C\, n^{1/d_{\text{eff}}}$ (effective spectral dimension; cf. [394, 369] and Remark 4.423). The wordwise effective cutoff $\Lambda_{\text{eff}}^{[\text{obs}],[\cdot]}$ from Definition 4.583 marks the tail where $w_n^{[\cdot]} f_n^{(\text{obs})[\cdot]} < \varepsilon$. If the Holo–State retains non–negligible signal at the Planck band (so that $\Lambda_{\text{eff}}^{[\cdot]} \geq \ell_{\text{P}}^{-1}$), then the PQS width

$$\ell_{\text{PQS}}^{[\text{obs}],[\cdot]} \;:=\; \left(\Lambda_{\text{eff}}^{[\text{obs}],[\cdot]}\right)^{-1}$$

obeys the *weak* sub–Planckian bound $\ell_{\text{PQS}}^{[\cdot]} \leq \ell_{\text{P}}$. If, moreover, the signal persists a factor $\kappa \gg 1$ deeper into the UV (Lemma 4.596), then $\Lambda_{\text{eff}}^{[\cdot]} \geq \kappa\, \ell_{\text{P}}^{-1}$ and hence

$$\ell_{\text{PQS}}^{[\cdot]} \;\leq\; \frac{\ell_{\text{P}}}{\kappa} \;\ll\; \ell_{\text{P}} \quad \text{(strong, ultra–sub–Planckian).}$$

(2) Operator–spectral suppression (wordwise). Define the cutoff by the tail condition

$$\left(w_n^{[\cdot]}(x,x')\right)^2 \left|\mathcal{S}_n^{[\cdot]}(x,x')\right|^2 \;\leq\; \varepsilon \qquad \left(\lambda_n^{[\mathbf{W}_1]} > \Lambda_{\text{eff}}^{[\cdot]}\right).$$

Under Wordwise Spectral Smoothness, $\left|\mathcal{S}_n^{[\cdot]}\right| \lesssim \left(\lambda_n^{[\mathbf{W}_1]}\right)^{\beta}$ (Theorem 4.421), and $w_n^{[\cdot]} \lesssim \left(\lambda_n^{[\mathbf{W}_1]}\right)^{-\alpha}$ with $\alpha > \beta + \frac{1}{2}$ (Principle 4.2). Then Lemma 4.584 yields a finite, ε–dependent $\Lambda_{\text{eff}}^{[\cdot]}$ with the pointwise estimate (as in (4.280))

$$\Lambda_{\text{eff}}^{[\cdot]}(x,x';\varepsilon) \;\leq\; \max\left\{\Lambda_0,\; \left(\frac{C' C_f}{\varepsilon}\right)^{\frac{1}{\alpha-\beta}}\right\}.$$

Tightening tolerance pushes $\Lambda_{\text{eff}}^{[\cdot]}$ arbitrarily high, hence $\ell_{\text{PQS}}^{[\cdot]} = \left(\Lambda_{\text{eff}}^{[\cdot]}\right)^{-1}$ can be driven arbitrarily small. When Planck–band contributions are *detectable* (non–negligible), one has $\Lambda_{\text{eff}}^{[\cdot]} \geq \ell_{\text{P}}^{-1}$, so $\ell_{\text{PQS}}^{[\cdot]} \leq \ell_{\text{P}}$; if the persistence extends beyond $\kappa\, \ell_{\text{P}}^{-1}$, one obtains the ultra–sub–Planckian bound (Lemma 4.596). (For the role of $\ell_{\text{P}} = \sqrt{\hbar G/c^3}$ as a semiclassical *benchmark*, see [284, 415, 403].)

(3) Breakdown of geometry below ℓ_{PQS} (wordwise). The expectation–valued geometry is meaningful only where modal coherence persists. For $|x - x'| < \ell_{\text{PQS}}^{[\cdot]}$ one has tail suppression $w_n^{[\cdot]}(x,x') \to 0$ $(\forall n)$, hence the wordwise correlator amplitude satisfies

$$\left|\langle \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})\rangle\right| = \left|\sum\nolimits_n w_n^{[\cdot]}(x,x')\, \mathrm{e}^{\mathrm{i}\theta_n^{[\cdot]}}\right| \;\longrightarrow\; 0,$$

so relational geometry dissolves below the PQS width. In contrast, at Planckian separations ($\sim \ell_{\text{P}}$) the correlator retains finite amplitude under the phase–coherent weight condition (wordwise analogue; see Remark 4.597).

(4) Interpretive principle (wordwise).

Emergent geometry arises only above the modal distinguishability threshold set by the (wordwise) PQS width. The Planck length is not a fundamental bound, but an effective, semiclassical scale emerging from the deeper holosymmetric relational substrate.

Wordwise Theorem: Sub– and Ultra–Sub–Planckian Coherence of the PQS

Theorem 4.598 (Sub– and Ultra–Sub–Planckian Coherence of the PQS (wordwise))**.** *Let* $\ell_{\mathrm{PQS}}^{[\mathrm{obs}],[\cdot]} := \left(\Lambda_{\mathrm{eff}}^{[\mathrm{obs}],[\cdot]}\right)^{-1}$ *be the PQS coherence width from Definition 4.583, and* $\ell_{\mathrm{P}} := \sqrt{\hbar G/c^3}$ *the Planck length. Under wordwise modal–weight decay (Principle 4.2) and Wordwise Spectral Smoothness (Theorem 4.421):*

(a) If the Holo–State exhibits non–negligible Planck–band support for the chosen observable (weak persistence), then

$$\ell_{\mathrm{PQS}}^{[\mathrm{obs}],[\cdot]} \;\le\; \ell_{\mathrm{P}}, \qquad \tau_{\mathrm{PQS}}^{[\mathrm{obs}],[\cdot]} := \frac{\ell_{\mathrm{PQS}}^{[\mathrm{obs}],[\cdot]}}{c} \;\le\; \frac{\ell_{\mathrm{P}}}{c}.$$

(b) If the UV tail is non–negligible on $[\kappa\,\ell_{\mathrm{P}}^{-1}, \infty)$ *for some* $\kappa \gg 1$ *(Lemma 4.596), then*

$$\ell_{\mathrm{PQS}}^{[\mathrm{obs}],[\cdot]} \;\le\; \frac{\ell_{\mathrm{P}}}{\kappa} \;\ll\; \ell_{\mathrm{P}}, \qquad \tau_{\mathrm{PQS}}^{[\mathrm{obs}],[\cdot]} \;\le\; \frac{1}{\kappa}\,\frac{\ell_{\mathrm{P}}}{c} \;\ll\; \frac{\ell_{\mathrm{P}}}{c}.$$

Proof sketch. Finiteness and monotonicity of $\varepsilon \mapsto \Lambda_{\mathrm{eff}}^{[\cdot]}(\varepsilon)$ follow from Lemma 4.584. Weak persistence enforces $\Lambda_{\mathrm{eff}}^{[\cdot]} \ge \ell_{\mathrm{P}}^{-1}$, giving the sub–Planckian bound. Ultra–persistence with factor κ enforces $\Lambda_{\mathrm{eff}}^{[\cdot]} \ge \kappa\,\ell_{\mathrm{P}}^{-1}$ by Lemma 4.596, yielding the parametric (ultra–) separation. □

$$\boxed{\ell_{\mathrm{PQS}}^{[\cdot]} = \left(\Lambda_{\mathrm{eff}}^{[\cdot]}\right)^{-1} \;\le\; \frac{\ell_{\mathrm{P}}}{\max\{1,\kappa\}}\,, \qquad \tau_{\mathrm{PQS}}^{[\cdot]} = \frac{\ell_{\mathrm{PQS}}^{[\cdot]}}{c} \;\le\; \frac{1}{\max\{1,\kappa\}}\,\frac{\ell_{\mathrm{P}}}{c}\,.}$$

Citations. Planck scale as semiclassical benchmark: [284, 415, 403, 416, 83, 40]. Weyl asymptotics and functional calculus background: [394, 369].

4.16 The Relational Phase–Geometry Fibre Bundle

In IQR, spatial and temporal relations are *emergent*, state–conditioned functionals of operator data rather than background primitives. To organise these relations, we assemble a fibre bundle

$$\boxed{\pi : \mathfrak{F} \longrightarrow \mathcal{X}_q,} \tag{4.288}$$

whose fibre at each basepoint $x \in \mathcal{X}_q$ collects the triadic *distance* data and the triadic *time* (phase–displacement) data evaluated against all comparison loci $x' \in \mathcal{X}_q$. Concretely,

$$\boxed{\mathfrak{F}_x \;:=\; \mathcal{F}_x^{\mathrm{dist}} \;\times\; \mathcal{F}_x^{\mathrm{time}},} \tag{4.289}$$

where $\mathcal{F}_x^{\mathrm{dist}}$ and $\mathcal{F}_x^{\mathrm{time}}$ are exactly those defined in Definitions 4.599 and 4.600. While this is a product at the level of sets, the expectation–level chronogeometric tensor splits into orthogonal summands, $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$, with $\bar{g}_{\mu\nu}$ sourced by distinguishability and $\bar{t}_{\mu\nu}$ by temporal phase data. Unless otherwise stated, modal coordinates on each fibre are expressed in the Fourfold dialectic basis $\{\, \mathring{\mathbf{0}},\ \mathring{\mathbf{1}},\ \mathring{\mathbf{0}}{+}\mathring{\mathbf{1}},\ \mathring{\mathbf{0}}\mathring{\mathbf{\Lambda}}\mathring{\mathbf{1}} \,\}$.

The Distinguishability and Temporal Flow Fibres

Definition 4.599 (Distinguishability Fibre)**.** For each $x \in \mathcal{X}_q$, define the set of relational *distance* vectors

$$\boxed{\mathcal{F}_x^{\mathrm{dist}} \;:=\; \Big\{ \vec{d}(x,x') \;:=\; \big(d_{\mathrm{corr}}(x,x'),\ d_{\mathrm{vac}}(x,x'),\ d_{\mathrm{spec}}(x,x')\big) \;\in \mathbb{R}^3 \;:\; x' \in \mathcal{X}_q \Big\}.} \tag{4.290}$$

The composite distinguishability is the Euclidean norm $D(x,x') \;:=\; \|\vec{d}(x,x')\|_2$. The axis components $d_{\mathrm{corr}}, d_{\mathrm{vac}}, d_{\mathrm{spec}}$ are those already defined earlier and are not re-stated here.

Definition 4.600 (Temporal Flow Fibre)**.** For each $x \in \mathcal{X}_q$, define the set of relational *time* (phase–displacement) vectors

$$\boxed{\mathcal{F}_x^{\mathrm{time}} \;:=\; \Big\{ \vec{t}(x,x') \;:=\; \big(t_{\mathrm{corr}}(x,x'),\ t_{\mathrm{vac}}(x,x'),\ t_{\mathrm{spec}}(x,x')\big) \;\in \mathbb{R}^3 \;:\; x' \in \mathcal{X}_q \Big\}.} \tag{4.291}$$

These are *phase* quantities (not metrics), arising from distinct operator–theoretic evolution mechanisms; their formal definitions have already been fixed and are not duplicated here.

Global Description. Equivalently, one may view $\mathfrak{F}$ as the disjoint union

$$\mathfrak{F} \;=\; \bigsqcup_{x \in \mathcal{X}_q} \Big(\{\, \vec{d}(x,\cdot) : \mathcal{X}_q \to \mathbb{R}^3 \,\} \;\times\; \{\, \vec{t}(x,\cdot) : \mathcal{X}_q \to \mathbb{R}^3 \,\} \Big),$$

with projection π sending $(x, \vec{d}, \vec{t}) \mapsto x$. Both fibres inherit gauge–covariant transport and holonomy from the underlying Fourfold correlator and spectral data; nothing in Definitions 4.599 and 4.600 changes those earlier constructions—this section fixes the bundle–level bookkeeping and notation only.

Corollary 4.601 (Relational Null Fibre Equals the PQS Class)**.** *Fix $x \in \mathcal{X}_q$ and define the bundle-level relational null fibre*

$$Z_x \;:=\; \{\, x' \in \mathcal{X}_q :\ \vec{d}(x,x') = \mathbf{0} \ \ \textit{and}\ \ \vec{t}(x,x') = \mathbf{0} \,\}.$$

Then

$$\boxed{Z_x \;=\; \mathfrak{B}_x,} \tag{4.292}$$

i.e. $x' \in \mathfrak{B}_x$ iff all emergent relational observables between x and x' vanish.

Proof sketch. By the relational null condition and the PQS identification (Definition 4.574, Section 4.14, and Remark 4.581), the vanishing of the distance triad $\vec{d}(x,x')$ characterises the indistinguishability locus generated by the distance fibre. The temporal zero–locus and holonomy/flatness criteria (Sections 4.11.9 and 4.12.8) imply that $\vec{t}(x,x') = \mathbf{0}$ holds on the same locus when the corresponding temporal curvatures vanish along any connecting path. Conversely, if both $\vec{d}$ and $\vec{t}$ vanish, the pair (x,x') lies in the PQS class by the same references. Hence $Z_x = \mathfrak{B}_x$. □

Coherence Locality of the Fibres (and What It Is Not)

The fibres $\mathfrak{F}_x = \mathcal{F}_x^{\mathrm{dist}} \times \mathcal{F}_x^{\mathrm{time}}$ are globally defined over $\mathcal{X}_q$. Their *joint* zero locus at x is precisely the PQS class $\mathfrak{B}_x$ (Definition 4.574 and Corollary 4.601). What the PQS width $\ell_{\mathrm{PQS}} = \Lambda_{\mathrm{eff}}^{-1}$ controls is not compact support of $(\vec{d},\vec{t})$, but the *resolution threshold* for coherent relational signal.

Lemma 4.602 (Coherence Attenuation Below the PQS Width)**.** *Fix $x \in \mathcal{X}_q$. For any tolerance $\varepsilon > 0$ and any observable label* [obs] $\in \{\mathrm{spec}, \mathrm{vac}\}$*, there exists a radius $r_\varepsilon(x) \lesssim \ell_{\mathrm{PQS}}(x;\varepsilon)$ such that whenever the effective separation of x and x' lies within this radius, the corresponding bilocal channel and feature–weighted tails are uniformly small:*

$$\left|\langle \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})\rangle\right| < \varepsilon, \qquad w_n(x,x')\, f_n^{(\mathrm{obs})}(x,x') < \varepsilon \;\; \text{for all} \;\; \lambda_n > \Lambda_{\mathrm{eff}}^{[\mathrm{obs}]}(x,x';\varepsilon).$$

Sketch. *This is the local-in-x' restatement of Definition 4.576 and Lemma 4.577: for fixed ε, the finiteness and monotonicity of $\Lambda_{\mathrm{eff}}(\varepsilon)$ ensure that tightening the tolerance, or moving x' into a sufficiently small coherence neighbourhood, suppresses the tail to level ε. The bilocal bound then follows from the modal expansion and the tail estimate.*

Remark 4.603 (No Compact-Support Claim)**.** Lemma 4.602 *does not* imply $\vec{d}(x,x') = \vec{0}$ or $\vec{t}(x,x') = \vec{0}$ outside any neighbourhood. In fact, by Definition 4.574 and Corollary 4.601 the joint vanishing occurs *only* on the PQS class $\mathfrak{B}_x$. Moreover, correlator persistence at the Planck band (Lemma 4.589) and ultra–sub–Planckian tail–persistence (Lemma 4.590) show that coherent relational signal—and hence nontrivial $(\vec{d},\vec{t})$—generically extends far beyond ℓ_{PQS}.

In plain terms, the lemma says. *Below the PQS width we cannot "see" structure, to any chosen precision.* Fix any small tolerance $\varepsilon > 0$. Then there is an ε–dependent neighbourhood around x (with size on the order of the PQS width ℓ_{PQS}) such that whenever x' lies inside this neighbourhood, the *coherent* relational signal between x and x' is uniformly tiny—both in the raw bilocal channel $\langle \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})\rangle$ and in the feature–weighted spectral "tails" $w_n(x,x')\, f_n^{(\mathrm{obs})}(x,x')$. In other words, by moving x' sufficiently close (within a radius $r_\varepsilon \lesssim \ell_{\mathrm{PQS}}$), all *resolvable* relational information can be made smaller than ε.

This is attenuation, not annihilation. The lemma does *not* claim that observables vanish identically outside some ball (no compact support). It asserts that within a small–enough ball the signal is *as small as we like*—set by ε—because the effective spectral cutoff $\Lambda_{\text{eff}}(x, x'; \varepsilon)$ pushes all significant contributions above the accessible band. This motivates the term *coherence locality* rather than hard locality.

What sets the scale? The radius r_ε is controlled by the *PQS width* $\ell_{\text{PQS}} = \Lambda_{\text{eff}}^{-1}$. As ε tightens, Λ_{eff} (weakly) increases and the corresponding neighbourhood can shrink, but the mechanism is always the same: modal weight decay together with spectral smoothness ensures that beyond the cutoff the contributions are uniformly small, and this uniform smallness propagates to the bilocal expectation via its modal expansion.

Intuition in one line. If we compare two loci closer than the PQS width, the bundle's "readouts" (distance triad and time triad) become operationally indistinguishable to arbitrary precision ε—not because the theory kills the signal, but because the high–frequency modes that would distinguish the loci are suppressed beyond the effective cutoff.

Why it matters. The lemma is the local, quantitative backbone of (i) the *minimal relational scale* (points cannot be told apart below ℓ_{PQS} to any prescribed tolerance), (ii) the *finiteness* of all emergent observables (the same cutoff controls convergence), and (iii) the *bundle's smooth structure* (the maps $x \mapsto \vec{d}(x, \cdot)$ and $x \mapsto \vec{t}(x, \cdot)$ vary tamely because small separations automatically suppress coherent contributions).

In sum: "Coherence Attenuation Below the PQS Width" means that *below a state–dependent resolution threshold, the relational geometry loses operational contrast in a controlled, tolerance–dependent way*—small–but–not–zero everywhere, and never claiming compact support.

Definition of the Relational Bundle

Definition 4.604 (Relational Phase–Geometry Bundle)**.** Let $\mathcal{X}_q$ be the base space and let $\mathcal{F}_x^{\text{dist}}$, $\mathcal{F}_x^{\text{time}}$ be as in Definitions 4.599 and 4.600. The total space is the disjoint union

$$\boxed{\mathfrak{F} \;:=\; \bigsqcup_{x \in \mathcal{X}_q} \Big(\mathcal{F}_x^{\text{dist}} \;\times\; \mathcal{F}_x^{\text{time}} \Big),} \tag{4.293}$$

with projection $\pi : \mathfrak{F} \to \mathcal{X}_q$ given on simple elements by

$$\pi\Big(\vec{d}(x, x'),\, \vec{t}(x, x') \Big) \;:=\; x.$$

For each fixed $x' \in \mathcal{X}_q$ there are canonical evaluation maps

$$\text{ev}_{x'}^{\text{dist}} :\; \mathfrak{F}_x \to \mathbb{R}^3, \quad \vec{d}(x, \cdot) \mapsto \vec{d}(x, x'), \qquad \text{ev}_{x'}^{\text{time}} :\; \mathfrak{F}_x \to \mathbb{R}^3, \quad \vec{t}(x, \cdot) \mapsto \vec{t}(x, x'),$$

and the composite distinguishability is $D(x, x') = \|\vec{d}(x, x')\|_2$.

To comprehend the emergent fabric of reality in the IQR framework, it is helpful to view each locus $x \in \mathcal{X}_q$ not as a bare point but as a *relational nexus* equipped with an "instrument panel"—its fibre $\mathfrak{F}_x$ in the relational phase–geometry bundle (Definition 4.604). This panel simultaneously reads, for every comparison locus x', a triad of *distinguishability* amplitudes

and a triad of *phase (time)* displacements, assembled respectively in the fibres $\mathcal{F}_x^{\mathrm{dist}}$ and $\mathcal{F}_x^{\mathrm{time}}$ (Definitions 4.599 and 4.600). The product structure $\mathfrak{F}_x = \mathcal{F}_x^{\mathrm{dist}} \times \mathcal{F}_x^{\mathrm{time}}$ expresses that separations are never assessed independently of their intrinsic phase dynamics; the two readouts are inseparable facets of a single relational datum.

The *distinguishability* reading is the vector $\vec{d}(x,x') = (d_{\mathrm{corr}}, d_{\mathrm{vac}}, d_{\mathrm{spec}})$, whose Euclidean aggregate $D(x,x') = \|\vec{d}(x,x')\|_2$ quantifies how strongly x and x' differ as encoded by bilocal coherence, coincident vacuum response, and spectral shear. In parallel, the *temporal* reading is the vector $\vec{t}(x,x') = (t_{\mathrm{corr}}, t_{\mathrm{vac}}, t_{\mathrm{spec}})$, which captures the phase–sensitive aspects of the same operator data. Taken together, $(\vec{d}, \vec{t})$ provide a *polar decomposition* of emergent chronogeometry across three canonical channels: amplitudes populate the expectation–level metric contribution $\bar{g}_{\mu\nu}$, phases populate the temporal contribution $\bar{t}_{\mu\nu}$, and their orthogonal sum yields the chronogeometric tensor $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ introduced earlier.

Transporting these readouts across $\mathcal{X}_q$ is governed by the gauge–covariant identifications already established. Along a path, the correlator–phase and spectral/vacuum rate forms induce temporal and spatial connections whose curvatures measure the failure of global trivialisation; nontrivial holonomies encode the path dependence of the phase flow and, dually, of distinguishability (Sections 4.11.9 and 4.12.8, see also Theorem 4.559). In this sense the bundle is not mere bookkeeping: it is the stage on which connection, curvature, and holonomy act to assemble local relational data into a global chronogeometric structure.

The panel also calibrates the *threshold of indistinguishability*. When, for a given x, both triads vanish against x', $\vec{d}(x,x') = \mathbf{0}$ and $\vec{t}(x,x') = \mathbf{0}$, one is precisely at the *relational null* fibre; by the formal condition and its bundle consequence this identifies x' with the PQS class $\mathfrak{B}_x$ (Definition 4.574 and Corollary 4.601). The characteristic width of that class, $\ell_{\mathrm{PQS}} = \Lambda_{\mathrm{eff}}^{-1}$, arises from the effective spectral cutoff (Definition 4.576). Below this state–dependent resolution scale the coherent signal can be made uniformly small to any tolerance (coherence locality), yet not identically zero (Lemma 4.602 and Remark 4.603); at Planck–band separations a finite bilocal amplitude typically persists (Lemma 4.589), and in the ultra–sub–Planckian regime the cutoff can stand parametrically beyond the Planck band (Lemma 4.590).

Finally, no background geometry is presupposed: all readings are expectation values in the Holo–State, induced by the Fourfold correlator $\hat{G}_\Lambda$ and the spectral generator $\hat{D}$. Changes of modal frame act orthogonally on the triads and preserve the physically meaningful content—norms, connections, and holonomies—so the relational phase–geometry bundle is intrinsic to the operator–theoretic data of the theory.

Proposition 4.605 (Well-Posedness, Smoothness, and Gauge Covariance)**.** *Assume Relational Smoothness and Spectral Smoothness for the underlying operator data. Then:*

(a) Topological/Smooth Structure. *The assignments $x \mapsto \vec{d}(x,\cdot)$ and $x \mapsto \vec{t}(x,\cdot)$ vary continuously (indeed, smoothly) on $\mathcal{X}_q$ in the product topology on $\mathbb{R}^{3\,\mathcal{X}_q}$, making $\pi : \mathfrak{F} \to \mathcal{X}_q$ a (set–theoretic) fibre bundle with a natural smooth structure induced from the operator expectations.*

(b) Gauge/Frame Covariance. *Local unitary changes of modal frame act by orthogonal transformations on the triads, preserving $D(x,x') = \|\vec{d}(x,x')\|_2$ and the rate/holonomy*

constructions for $\vec{t}$ (Sections 4.11.9 and 4.12.8). Hence $\mathfrak{F}$ is intrinsically defined by the Holo–State and Fourfold/Spectral data.

Intuitive Understanding.

What actually sits in a fibre. At each basepoint $x \in \mathcal{X}_q$, the fibre $\mathfrak{F}_x = \mathcal{F}_x^{\mathrm{dist}} \times \mathcal{F}_x^{\mathrm{time}}$ packages two complementary readouts about every comparison locus x': a *distance triad* $\vec{d}(x,x') = (d_{\mathrm{corr}}, d_{\mathrm{vac}}, d_{\mathrm{spec}})$ and a *time (phase–displacement) triad* $\vec{t}(x,x') = (t_{\mathrm{corr}}, t_{\mathrm{vac}}, t_{\mathrm{spec}})$. The distance triad behaves metric-like through the aggregator $D(x,x') = \|\vec{d}(x,x')\|_2$, whereas the time triad records *phases* (not lengths). Intuitively: D says "how distinguishable x is from x'," while the components of $\vec{t}$ say "how much relational phase is processed" along each axis.

How fibres talk to each other (transport and holonomy). Comparisons of $(\vec{d}, \vec{t})$ along paths in $\mathcal{X}_q$ are made via the gauge-covariant transport already introduced; its curvatures are the obstructions to globally trivialising these readouts (Sections 4.11.9 and 4.12.8). When temporal curvatures vanish, a *clock gauge* exists in which all temporal components identically disappear; otherwise closed loops accumulate a nonzero temporal holonomy (cf. the Temporal Holonomy Obstruction, Theorem 4.559). Thus the bundle is not merely "bookkeeping'': it is the stage on which connection, curvature, and holonomy act.

From global state to local geometry (sections as probes). Fixing x' and letting x vary turns $(x \mapsto \vec{d}(x,x'),\ x \mapsto \vec{t}(x,x'))$ into *sections* of the bundle: operationally, one probe x' "reads out'' the local relational geometry everywhere else, induced by Holo-State expectations of $\hat{G}_\Lambda$ and the spectral generator $\hat{D}$. The *effective spectral cutoff* controls the resolution window for these readouts (Definition 4.576); its existence and scaling guarantee stable, finite observables across fibres (Lemma 4.577).

Coherence locality, not compact support. The PQS width $\ell_{\mathrm{PQS}} = \Lambda_{\mathrm{eff}}^{-1}$ marks a *resolution threshold*: below it, bilocal amplitudes and feature-weighted tails are uniformly small, but not *identically* zero. Planck-band separations still retain finite correlator amplitude (Lemma 4.589), and ultra–sub–Planckian tail-persistence can force a parametrically small coherence width (Lemma 4.590). This is why *locality in IQR is coherence-limited*, not "hard'' compact support (Remark 4.603).

What the bundle is (and is not). $\mathfrak{F}$ is a *hybrid* metric/phase structure over $\mathcal{X}_q$: the distance triad behaves metric–like via the aggregator $D(x,x') = \|\vec{d}(x,x')\|_2$, while the time triad records *phase–generated* observables and admits holonomy under transport. Importantly, this *does not* mean the temporal sector lacks magnitudes. Indeed, t_{vac} and t_{spec} are already Euclidean norms of imaginary–part profiles (cf. Definitions 4.535 and 4.543), and the correlator phase t_{corr} naturally lives on the circle S^1 with the circular distance

$$\mathrm{dist}_{S^1}\big(\theta, 0\big) \;:=\; \min_{m\in\mathbb{Z}} |\theta + 2\pi m|.$$

When a single scalar is useful, one may *calibrate* a temporal aggregate (without changing any prior definitions), e.g.

$$\|\vec{t}(x,x')\|_{\mathrm{time}} \;:=\; \Big(\mathrm{dist}_{S^1}(t_{\mathrm{corr}}(x,x'),0)^2 \;+\; t_{\mathrm{vac}}(x,x')^2 \;+\; t_{\mathrm{spec}}(x,x')^2\Big)^{1/2}.$$

All holonomy/curvature statements in Section 4.12.8 depend only on the underlying connections, not on choosing such a calibration; the option merely makes explicit that temporal phase flow *can* be normed when needed.

4.17 The Relational Phase–Geometry Fibre Bundle (Holosymmetric Wordwise)

In the holosymmetric wordwise calculus, spatial and temporal relations are emergent, state–conditioned functionals of operator data *in a chosen word frame* [**W**], and are word/-gauge–invariant (Propositions 4.340 and 4.422 and their time–duals). To organise these relations, we assemble the bundle

$$\boxed{\pi : \mathfrak{F}^{[\mathbf{W}]} \longrightarrow \mathcal{X}_q,} \tag{4.294}$$

whose fibre at each basepoint $x \in \mathcal{X}_q$ collects the triadic *distance* data and the triadic *time* (phase–displacement) data—both evaluated against all comparison loci $x' \in \mathcal{X}_q$ in the frame [**W**]. Concretely,

$$\boxed{\mathfrak{F}_x^{[\mathbf{W}]} := \mathcal{F}_x^{\mathrm{dist},[\mathbf{W}]} \times \mathcal{F}_x^{\mathrm{time},[\mathbf{W}]}.} \tag{4.295}$$

While this is a set–level product, the expectation–level chronogeometric tensor splits orthogonally, $\bar{\Sigma}_{\mu\nu}^{[\mathbf{W}]} = \bar{g}_{\mu\nu}^{[\mathbf{W}]} + \bar{t}_{\mu\nu}^{[\mathbf{W}]}$, with $\bar{g}_{\mu\nu}^{[\mathbf{W}]}$ sourced by distinguishability and $\bar{t}_{\mu\nu}^{[\mathbf{W}]}$ by temporal phase data (cf. Sections 4.11.2 and 4.11.4 and their wordwise counterparts). Unless otherwise stated, modal coordinates on each fibre are expressed in a holosymmetric Fourfold alphabet at x, e.g. $\{\mathring{\mathbf{0}}, \mathring{\mathbf{1}}, \mathring{\mathbf{p}}, \mathring{\mathbf{s}}\}$, transported by the admissible gauge along paths γ.

The Distinguishability and Temporal Flow Fibres (Wordwise)

Definition 4.606 (Wordwise Distinguishability Fibre)**.** For each $x \in \mathcal{X}_q$, define the set of relational *distance* vectors in the word frame [**W**]

$$\boxed{\mathcal{F}_x^{\mathrm{dist},[\mathbf{W}]} := \left\{ \vec{d}^{[\mathbf{W}]}(x,x') := \left(d_{\mathrm{corr}}^{[\mathbf{W}]}(x,x'),\ d_{\mathrm{vac}}^{[\mathbf{W}]}(x,x'),\ d_{\mathrm{spec}}^{[\mathbf{W}]}(x,x')\right) \in \mathbb{R}^3 \ :\ x' \in \mathcal{X}_q \right\}.} \tag{4.296}$$

The composite distinguishability is the calibrated Euclidean norm $D^{[\mathbf{W}]}(x,x') := \|\mathbf{M}^{1/2}(x)\,\vec{d}^{[\mathbf{W}]}(x,x')\|_2$ (Sections 4.11.2 and 4.11.4). Each component is the wordwise version already fixed in Definitions 4.339, 4.408 and 4.468, and is not re-stated here.

Definition 4.607 (Wordwise Temporal Flow Fibre)**.** For each $x \in \mathcal{X}_q$, define the set of relational *time* (phase–displacement) vectors

$$\boxed{\mathcal{F}_x^{\mathrm{time},[\mathbf{W}]} := \left\{ \vec{t}^{[\mathbf{W}]}(x,x') := \left(t_{\mathrm{corr}}^{[\mathbf{W}]}(x,x'),\ t_{\mathrm{vac}}^{[\mathbf{W}]}(x,x'),\ t_{\mathrm{spec}}^{[\mathbf{W}]}(x,x')\right) \in \mathbb{R}^3 \ :\ x' \in \mathcal{X}_q \right\}.} \tag{4.297}$$

These are *phase* quantities (not metrics), defined wordwise in Definitions 4.529, 4.539 and 4.547.

Global description (wordwise). Equivalently,

$$\mathfrak{F}^{[\mathbf{W}]} \;=\; \bigsqcup_{x\in\mathcal{X}_q} \Big(\{\, \vec{d}^{[\mathbf{W}]}(x,\cdot) : \mathcal{X}_q \to \mathbb{R}^3 \,\} \;\times\; \{\, \vec{t}^{[\mathbf{W}]}(x,\cdot) : \mathcal{X}_q \to \mathbb{R}^3 \,\} \Big),$$

with projection π sending $(x, \vec{d}^{[\mathbf{W}]}, \vec{t}^{[\mathbf{W}]}) \mapsto x$. Both fibres inherit gauge–covariant transport and holonomy from the Fourfold correlator and wordwise spectral data; nothing in Definitions 4.606 and 4.607 alters prior constructions—this section fixes bundle–level bookkeeping in the wordwise setting.

Remark 4.608 (Word/Gauge invariance and exogenous equivalence)**.** By Propositions 4.340 and 4.411 and the matching results for the temporal triad, the fibres and all induced quantities satisfy $\vec{d}^{[\mathbf{W}]}(x,x') = \vec{d}(x,x')$, $\vec{t}^{[\mathbf{W}]}(x,x') = \vec{t}(x,x')$ and hence $D^{[\mathbf{W}]}(x,x') = D(x,x')$, modulo the fixed calibration $\mathbf{M}(x)$. Thus the relational bundle is independent of the particular holosymmetric word choice.

Relational Null Fibre and the PQS Class (Wordwise)

Corollary 4.609 (Wordwise relational null fibre equals the PQS class)**.** *Fix $x \in \mathcal{X}_q$ and define*

$$Z_x^{[\mathbf{W}]} \;:=\; \{\, x' \in \mathcal{X}_q :\; \vec{d}^{[\mathbf{W}]}(x,x') = \mathbf{0} \;\; \textit{and} \;\; \vec{t}^{[\mathbf{W}]}(x,x') = \mathbf{0} \,\}.$$

Then

$$\boxed{Z_x^{[\mathbf{W}]} \;=\; \mathfrak{B}_x,} \tag{4.298}$$

i.e. $x' \in \mathfrak{B}_x$ iff all emergent wordwise relational observables between x and x' vanish.

Proof sketch. By the composite metric definition, $D^{[\mathbf{W}]}(x,x') = 0$ iff $\vec{d}^{[\mathbf{W}]}(x,x') = \mathbf{0}$ (Section 4.11.2). Word/exogenous equivalence for all six components (Propositions 4.340 and 4.422 and their temporal analogues) gives

$$\vec{d}^{[\mathbf{W}]}(x,x') = \mathbf{0} = \vec{t}^{[\mathbf{W}]}(x,x') \;\Longleftrightarrow\; \vec{d}(x,x') = \mathbf{0} = \vec{t}(x,x'),$$

i.e. the (wordwise) null condition is equivalent to the exogenous null condition (Definition 4.574). The exogenous "zero–locus = PQS" identification then yields $Z_x^{[\mathbf{W}]} = \mathfrak{B}_x$ (Proposition 4.507 and Corollary 4.510). Finally, word/gauge invariance of the components implies that this identification is independent of the choice of $[\mathbf{W}]$ (see also Proposition 4.613). □

Coherence Locality of the Fibres (What It Is and Is Not, Wordwise)

The fibres $\mathfrak{F}_x^{[\mathbf{W}]} = \mathcal{F}_x^{\mathrm{dist},[\mathbf{W}]} \times \mathcal{F}_x^{\mathrm{time},[\mathbf{W}]}$ are globally defined over $\mathcal{X}_q$. Their *joint* zero locus at x is the PQS class $\mathfrak{B}_x$ (Corollary 4.609). What the wordwise PQS width $\ell_{\mathrm{PQS}}^{[\cdot]} = (\Lambda_{\mathrm{eff}}^{[\cdot]})^{-1}$ controls is not compact support of $(\vec{d}^{[\mathbf{W}]}, \vec{t}^{[\mathbf{W}]})$, but the *resolution threshold* for coherent relational signal.

Lemma 4.610 (Wordwise coherence attenuation below the PQS width)**.** *Fix $x \in \mathcal{X}_q$. For any tolerance $\varepsilon > 0$ and any observable label* [obs] $\in \{\mathrm{spec}, \mathrm{vac}\}$*, there exists a radius $r_\varepsilon^{[\cdot]}(x) \lesssim \ell_{\mathrm{PQS}}^{[\mathrm{obs}],[\cdot]}(x;\varepsilon)$ such that whenever the effective separation of x and x' lies within this radius, the corresponding wordwise bilocal channel and feature–weighted tails are uniformly small:*

$$\left|\langle \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})\rangle\right| < \varepsilon, \qquad w_n^{[\cdot]}(x,x')\, f_n^{(\mathrm{obs})[\cdot]}(x,x') < \varepsilon \;\; \textit{for all} \;\; \lambda_n^{[\mathbf{W}_1]} > \Lambda_{\mathrm{eff}}^{[\mathrm{obs}],[\cdot]}(x,x';\varepsilon).$$

Sketch. *This is the wordwise restatement of Definition 4.583 and Lemma 4.584 and their wordwise versions, using the modal expansion with weights $w_n^{[\cdot]}$ and features $f_n^{(\mathrm{obs})[\cdot]}$.*

Remark 4.611 (No compact–support claim (wordwise))**.** Lemma 4.610 *does not* imply $\vec{d}^{[\mathbf{W}]}(x,x') = \vec{0}$ or $\vec{t}^{[\mathbf{W}]}(x,x') = \vec{0}$ outside any neighbourhood. The joint vanishing occurs only on $\mathfrak{B}_x$ (Corollary 4.609). Moreover, correlator persistence at the Planck band and ultra–sub–Planckian tail–persistence (wordwise versions of Remark 4.597 and Lemma 4.596) show that coherent relational signal—and hence nontrivial $(\vec{d}^{[\mathbf{W}]}, \vec{t}^{[\mathbf{W}]})$—generically extends far beyond $\ell_{\mathrm{PQS}}^{[\cdot]}$.

Bundle Metric and Calibration (Wordwise Recap)

Equip each fibre with the calibration inner product $\langle\cdot,\cdot\rangle_{\mathcal{F}_x}$ (Definition 4.494), taken wordwise as constant under change of $[\mathbf{W}]$ by Remark 4.608. Then the composite distance is the fibre norm, $D^{[\mathbf{W}]}(x,x') = \|\mathbf{M}^{1/2}(x)\, \vec{d}^{[\mathbf{W}]}(x,x')\|_2$, and the operator–valued metric arises from the coincidence–limit Hessian as in Section 4.11.3, now manifestly independent of the choice of holosymmetric alphabet.

Wordwise Definition of the Relational Bundle

Definition 4.612 (Holosymmetric Wordwise Relational Phase–Geometry Bundle)**.** Let $\mathcal{X}_q$ be the base and let $\mathcal{F}_x^{\mathrm{dist},[\mathbf{W}]}$, $\mathcal{F}_x^{\mathrm{time},[\mathbf{W}]}$ be the wordwise fibres of Definitions 4.606 and 4.607. Define the total space as the disjoint union

$$\boxed{\mathfrak{F}^{[\mathbf{W}]} \;:=\; \bigsqcup_{x\in\mathcal{X}_q} \Big(\mathcal{F}_x^{\mathrm{dist},[\mathbf{W}]} \;\times\; \mathcal{F}_x^{\mathrm{time},[\mathbf{W}]} \Big),} \tag{4.299}$$

with projection $\pi^{[\mathbf{W}]} : \mathfrak{F}^{[\mathbf{W}]} \to \mathcal{X}_q$ acting on simple elements by

$$\pi^{[\mathbf{W}]}\Big(\vec{d}^{[\mathbf{W}]}(x,x'),\, \vec{t}^{[\mathbf{W}]}(x,x') \Big) \;:=\; x.$$

For each fixed $x' \in \mathcal{X}_q$, the canonical evaluation maps are

$$\begin{aligned} \mathrm{ev}_{x'}^{\mathrm{dist},[\mathbf{W}]}&:\; \mathfrak{F}_x^{[\mathbf{W}]} \to \mathbb{R}^3, \qquad \vec{d}^{[\mathbf{W}]}(x,\cdot) \longmapsto \vec{d}^{[\mathbf{W}]}(x,x'),\\ \mathrm{ev}_{x'}^{\mathrm{time},[\mathbf{W}]}&:\; \mathfrak{F}_x^{[\mathbf{W}]} \to \mathbb{R}^3, \qquad \vec{t}^{[\mathbf{W}]}(x,\cdot) \longmapsto \vec{t}^{[\mathbf{W}]}(x,x'). \end{aligned}$$

With a calibration matrix $\mathbf{M}(x) \succ 0$ on $\mathcal{F}_x^{\mathrm{dist},[\mathbf{W}]}$ (Section 4.11.4), the composite distinguishability is

$$\boxed{D^{[\mathbf{W}]}(x,x') \;:=\; \Big\|\mathbf{M}^{1/2}(x)\, \vec{d}^{[\mathbf{W}]}(x,x')\Big\|_2.} \tag{4.300}$$

Proposition 4.613 (Word/Gauge equivalence with the exogenous bundle)**.** *There is a canonical fibrewise isomorphism* $\Phi^{[\mathbf{W}]} : \mathfrak{F}^{[\mathbf{W}]} \xrightarrow{\cong} \mathfrak{F}$ *such that* $\Phi^{[\mathbf{W}]}\big(\vec{d}^{[\mathbf{W}]}(x,x'), \vec{t}^{[\mathbf{W}]}(x,x')\big) = \big(\vec{d}(x,x'), \vec{t}(x,x')\big)$ *and* $D^{[\mathbf{W}]}(x,x') = D(x,x')$ *for the same* $\mathbf{M}(x)$.

Proof (sketch). By word/gauge invariance of the component observables (Propositions 4.340 and 4.422 and the time–duals), $\vec{d}^{[\mathbf{W}]}(x,x') = \vec{d}(x,x')$ and $\vec{t}^{[\mathbf{W}]}(x,x') = \vec{t}(x,x')$ for all (x,x'). Define $\Phi^{[\mathbf{W}]}$ fibrewise by identity on these triples; this intertwines projections and evaluation maps and preserves (4.300). □

Remark 4.614 (Local trivializations and smooth structure)**.** On any chart $U \subset \mathcal{X}_q$ where the transport, modal frames, and calibration $\mathbf{M}(x)$ vary smoothly, the map

$$\varphi_U^{[\mathbf{W}]} : \pi^{-1}(U) \longrightarrow U \times \Big(\mathcal{C}^\infty(U;\mathbb{R}^3) \times \mathcal{C}^\infty(U;\mathbb{R}^3)\Big), \quad (x, \vec{d}^{[\mathbf{W}]}, \vec{t}^{[\mathbf{W}]}) \mapsto \Big(x;\, [\vec{d}^{[\mathbf{W}]}]_x,\, [\vec{t}^{[\mathbf{W}]}]_x\Big)$$

provides a local trivialization when the fibres are endowed with the natural Fréchet topology of smooth $\mathbb{R}^3$–valued profiles. Changes of word/frame act pointwise by orthogonal transformations on each $\mathbb{R}^3$ and preserve norms and $D^{[\mathbf{W}]}$. Fixing a comparison locus x' produces the finite–rank *evaluation subbundle* $E_{x'} \to U$ with typical fibre $\mathbb{R}^3 \times \mathbb{R}^3$.

Remark 4.615 (Relational null fibre and PQS)**.** The bundle–level null fibre at x, $Z_x^{[\mathbf{W}]} := \{x' \mid \vec{d}^{[\mathbf{W}]}(x,x') = \mathbf{0},\ \vec{t}^{[\mathbf{W}]}(x,x') = \mathbf{0}\}$, coincides with the PQS class $\mathfrak{B}_x$ by Corollary 4.609; this identification is independent of $[\mathbf{W}]$.

Remark 4.616 (Component references (wordwise))**.** Each component of $\vec{d}^{[\mathbf{W}]}$ and $\vec{t}^{[\mathbf{W}]}$ is defined in Definitions 4.339, 4.408, 4.468, 4.529, 4.539 and 4.547; the composite structure and inner products are in Sections 4.11.2 and 4.11.4.

4.18 The Fibred Continuum of Primordial Quantum Singularities

Definition 4.617 (The Fibred Continuum of Primordial Quantum Singularities)**.** Let $\mathcal{C}_d$ be a smooth, connected, Hausdorff, second–countable d-manifold diffeomorphic to $\mathbb{R}^d$ (hence $\mathrm{card}(\mathcal{C}_d) = \mathfrak{c}$). Fix the coherence width $\ell_{\mathrm{PQS}} := \lambda_{\mathrm{cutoff}}^{-1}$ with $\ell_{\mathrm{PQS}} \ll \ell_{\mathrm{P}}$ as determined by the effective spectral cutoff of the admissible spectral generator already in use for $(d_{\mathrm{spec}}, t_{\mathrm{spec}})$ (Section 1.2.9 and Definition 4.600). Then:

1. **PQS loci as relational null classes.** For each $\vec{r} \in \mathcal{C}_d$, define the *Primordial Quantum Singularity* (PQS) associated to $\vec{r}$ to be the equivalence class

$$\mathfrak{B}_{\vec{r}} := \Big\{\vec{r}' \in \mathcal{C}_d :\ \vec{d}(\vec{r},\vec{r}') = \vec{0} \text{ and } \vec{t}(\vec{r},\vec{r}') = \vec{0}\Big\},$$

i.e. the relational–null domain anchored at $\vec{r}$ in the sense already stipulated for the triads $\vec{d}$ and $\vec{t}$ (Definitions 4.599 and 4.600). By construction, $\mathfrak{B}_{\vec{r}}$ is the minimal domain on which all emergent observables vanish (no new proof required).

2. **Structured singular content.** Each PQS $\mathfrak{B}_{\vec{r}}$ carries the structured content

$$\boxed{\hat{\Omega}_{\vec{r}} \;:=\; \overset{\circ}{\hat{\Lambda}}_{\hat{\Phi}\hat{\Theta}},} \tag{4.301}$$

where $\overset{\circ}{\Lambda}$ is the universal duality code and $\hat{\Phi}, \hat{\Theta}$ are the Gaiamiti and Tantrakyma dynamical fields, respectively (as fixed earlier in Definitions 6.118 and 6.142). No further structure is imposed here beyond what those definitions provide.

3. **Hilbert bundle of relational observables.** The bundle of relational observables is the (locally trivial) Hilbert bundle

$$\boxed{\pi : \; \mathfrak{F} \;:=\; \bigsqcup_{\vec{r}\in\mathcal{C}_d} \Big(\mathcal{F}^{\mathrm{dist}}_{\vec{r}} \times \mathcal{F}^{\mathrm{time}}_{\vec{r}} \Big) \;\longrightarrow\; \mathcal{C}_d,} \tag{4.302}$$

with typical fiber the Hilbert direct sum $\mathcal{H}_{\mathrm{typ}} := \mathcal{H}_{\mathrm{dist}} \oplus \mathcal{H}_{\mathrm{time}}$, where each $\mathcal{F}^{\mathrm{dist}}_{\vec{r}}$ (resp. $\mathcal{F}^{\mathrm{time}}_{\vec{r}}$) is the completion of the corresponding modal space under the inner products induced by the Fourfold expectations at $\vec{r}$ (already fixed by the coincident-limit construction), and the product in (4.302) is the topological product of these Hilbert spaces (algebraically the direct sum). The projection is the anchor map $\pi(\phi) = \vec{r}$ for any $\phi \in \mathfrak{F}_{\vec{r}}$. Local trivializations are provided by parallel transport of the Fourfold data (cf. the gauge–covariant transporter already introduced), and the structure group acts by fiberwise unitaries on $\mathcal{H}_{\mathrm{typ}}$ [372, 248]. Each fiber is coordinatized in the fourfold dialectic modal basis established in Section 1.2.9.

4. **Ultra–sub–Planckian support.** For each $\vec{r} \in \mathcal{C}_d$, let $\hat{D}_{\vec{r}}$ be the (already fixed) admissible spectral generator underlying the spectral components of the triads, and let $E^{(\vec{r})}(\cdot)$ denote its spectral measure. The fibre is restricted by the projector onto energies below the effective cutoff:

$$\boxed{\mathfrak{F}_{\vec{r}} \;=\; \mathrm{Ran}\Big(E^{(\vec{r})}\big([0, \lambda_{\mathrm{cutoff}}]\big)\Big), \qquad \ell_{\mathrm{PQS}} := \lambda^{-1}_{\mathrm{cutoff}} \ll \ell_{\mathrm{P}}.} \tag{4.303}$$

Here $\mathrm{Ran}(P) = \{P\psi : \psi \in \mathcal{H}_{\vec{r}}\}$ is the range (image) of the projector P; since $E^{(\vec{r})}([0, \lambda_{\mathrm{cutoff}}])$ is orthogonal, Ran is a closed subspace. Equivalently,

$$\begin{aligned} \mathfrak{F}_{\vec{r}} &= \mathbf{1}_{[0,\lambda_{\mathrm{cutoff}}]}(\hat{D}_{\vec{r}})\, \mathcal{H}_{\vec{r}} = \ker\Big(\mathbf{1} - E^{(\vec{r})}([0, \lambda_{\mathrm{cutoff}}])\Big), \\ \psi \in \mathfrak{F}_{\vec{r}} &\iff E^{(\vec{r})}\Big((\lambda_{\mathrm{cutoff}}, \infty)\Big)\, \psi = 0. \end{aligned} \tag{4.304}$$

Thus "ultra–sub–Planckian support'' means that all fibrewise spectral weight of admissible sections lies below $\lambda_{\mathrm{cutoff}}$, i.e. within coherence width ℓ_{PQS}, in accordance with the coherent–regularisation hypotheses (Principle 4.1 and eq. (4.199)).

We call the resulting structure

$$\boxed{\mathcal{C}_d \text{ : a fibred continuum of Primordial Quantum Singularities } \Big\{\, \mathfrak{B}_{\vec{r}},\; \hat{\Omega}_{\vec{r}},\; \mathfrak{F}_{\vec{r}} \,\Big\}_{\vec{r}\in\mathcal{C}_d}.}$$

Remarks (purely clarificatory, no new claims). (i) Item (4.302) is a standard Hilbert-bundle specification with fibers already determined by the Fourfold inner-product construction; (ii) the coordinatization "in the fourfold dialectic modal basis'' means that fiber coordinates are taken in the Thesis/Antithesis/Synthesis/Holothesis decomposition fixed in Section 1.2.9; (iii) the spectral form (4.303) is the fiberwise restatement of the effective cutoff that underlies d_{spec} and t_{spec}, expressed via the spectral calculus [4].

The definition is designed to collapse the pre-geometric ingredients into a single, coherent arena in which "locations'' are operational rather than primitive. A locus $\vec{r} \in \mathcal{C}_d$ is not a bare point but the anchor of an equivalence class determined entirely by the triadic indistinguishability already fixed by the relational fibres: $\mathfrak{B}_{\vec{r}} = \{\vec{r}' : \vec{d}(\vec{r},\vec{r}') = \vec{0} = \vec{t}(\vec{r},\vec{r}')\}$. In this way, the continuum $\mathcal{C}_d$ is read through the operational capacities of the theory—namely the distance and temporal fibres $\mathcal{F}^{\text{dist}}_{\vec{r}}$ and $\mathcal{F}^{\text{time}}_{\vec{r}}$—so that the very notion of "sameness of place'' is the vanishing of the established triads $\vec{d}$ and $\vec{t}$ (Definitions 4.599 and 4.600). The consequence is a background-independent identification of PQS loci as *relational–null classes*, minimal domains on which all emergent observables, by prior construction, vanish.

At each such locus the definition attaches *structured* ontic content without importing any new assumptions: the object $\hat{\Omega}_{\vec{r}} = \mathring{\Lambda}_{\hat{\Phi}\hat{\Theta}}$ is simply the local presentation of the universal duality code $\mathring{\Lambda}$ mediated by the already-defined Gaiamiti and Tantrakyma fields $\hat{\Phi}, \hat{\Theta}$ (eq. (4.301) and Definitions 6.118 and 6.142). This ensures that every PQS carries the same dual structural DNA that drives the theory's dialectic (Thesis/Antithesis/Synthesis/Holothesis; cf. Section 1.2.9), while keeping the definition purely organisational: it records what is present at a locus rather than proving anything new about its dynamics.

The relational data are then elevated from pointwise objects to a globally coherent geometric carrier by declaring the Hilbert bundle $\pi : \mathfrak{F} \to \mathcal{C}_d$ with fibre $\mathfrak{F}_{\vec{r}} = \mathcal{F}^{\text{dist}}_{\vec{r}} \times \mathcal{F}^{\text{time}}_{\vec{r}}$ (eq. (4.302)). Because each fibre's inner product is induced by the coincident-limit Fourfold expectations already in place, local trivialisations arise from the gauge-covariant parallel transport built into the formalism, and the natural structure group acts fibrewise by unitaries on the typical Hilbert space $\mathcal{H}_{\text{typ}} = \mathcal{H}_{\text{dist}} \oplus \mathcal{H}_{\text{time}}$ [372, 248]. The fourfold dialectic gives a canonical coordinatisation for sections of $\mathfrak{F}$ (Section 1.2.9), so that comparisons of fibres are intrinsic to the theory's operational semantics rather than imposed externally.

A further tightening is achieved by encoding ultraviolet control directly into the definition of each fibre. Using the admissible spectral generator already responsible for the spectral components $(d_{\text{spec}}, t_{\text{spec}})$, one restricts to the spectral range below the effective cutoff and thereby fixes the coherence width $\ell_{\text{PQS}} = \lambda^{-1}_{\text{cutoff}} \ll \ell_{\text{P}}$. Expressed fibrewise, this is the statement that $\mathfrak{F}_{\vec{r}} = \operatorname{Ran} E^{(\vec{r})}([0, \lambda_{\text{cutoff}}])$, which is exactly the ultra-sub-Planckian support recorded in eq. (4.303). Nothing new is asserted about the existence or value of the cutoff; rather, the definition faithfully registers the coherent-regularisation already justified elsewhere, ensuring that all subsequent constructions inherit the intended finiteness properties.

Taken together, the base $\mathcal{C}_d$, the PQS equivalence classes $\mathfrak{B}_{\vec{r}}$, the structured content $\hat{\Omega}_{\vec{r}}$, and the spectrally regulated Hilbert fibres $\mathfrak{F}_{\vec{r}}$ constitute a single, background-independent "workspace'' for the emergent chronogeometry. Because the fibres are precisely those that generate the triadic distances and times, the operator-valued metric and temporal curvature tensors can be introduced downstream as $\hat{g}_{\mu\nu}$ and $\hat{t}_{\mu\nu}$ from the established composite norms and rate forms (Definitions 5.1 and 5.17), and expectation-level chronogeometric data follow

by the usual passage to Born valuations [4, 3]. The definition thus furnishes a compact but exhaustive scaffold: loci defined operationally, content specified ontologically, geometry carried functorially—ready for interferometric constructions, parallel transport, and curvature without reopening any lemma or proof that the formalism has already secured.

Does the definition merely assemble the PQSs into a continuum $\mathcal{C}_d$ of cardinality $\mathfrak{c}$? Essentially yes, but with two clarifications crucial for later mathematics. First, the definition does more than merely aggregate PQSs into a set: it installs them on a smooth base $\mathcal{C}_d$ explicitly taken to be a connected, Hausdorff, second–countable d-manifold diffeomorphic to $\mathbb{R}^d$. Hence, by stipulation, $\mathrm{card}(\mathcal{C}_d) = \mathfrak{c}$, and the PQSs are *indexed* by this continuum via the assignment $\vec{r} \mapsto (\mathfrak{B}_{\vec{r}}, \hat{\Omega}_{\vec{r}}, \mathfrak{F}_{\vec{r}})$ (Section 4.18). Second, the "points" of this continuum are not primitive ontic atoms; each $\vec{r}$ anchors a relational–null class $\mathfrak{B}_{\vec{r}} = \{\vec{r}' : \vec{d}(\vec{r}, \vec{r}') = \vec{0} = \vec{t}(\vec{r}, \vec{r}')\}$, so the labelling is operationally grounded in the triads $\vec{d}, \vec{t}$ (Definitions 4.599 and 4.600). In particular, the choice $\mathcal{C}_d \cong \mathbb{R}^d$ ensures the *indexing space* already has cardinality $\mathfrak{c}$; under the standing identification of loci by the relational–null criterion (and using the earlier cardinality result for the emergent configuration space), the resulting collection of PQS loci inherits this cardinality rather than collapsing. Thus the definition indeed "puts all the PQSs together" as a continuum, while simultaneously fixing the smooth structure and the attached Hilbert bundle $\pi : \mathfrak{F} \to \mathcal{C}_d$, so that geometry and dynamics can be expressed fibrewise and transported consistently (eqs. (4.302) and (4.303)).

Does stipulating a smooth base $\mathcal{C}_d \cong \mathbb{R}^d$ and indexing a PQS triple at each anchor imply there are continuum-many PQS loci? Yes—with one subtlety. By stipulation in Section 4.18, the base is a smooth, connected, Hausdorff, second–countable d-manifold with $\mathrm{card}(\mathcal{C}_d) = \mathfrak{c}$. Assign to each anchor $\vec{r} \in \mathcal{C}_d$ the triple $(\mathfrak{B}_{\vec{r}}, \hat{\Omega}_{\vec{r}}, \mathfrak{F}_{\vec{r}})$, where $\mathfrak{B}_{\vec{r}}$ is the relational–null class defined by the triads $\vec{d}, \vec{t}$ (Definitions 4.599 and 4.600), $\hat{\Omega}_{\vec{r}} = \overset{\circ}{\hat{\Phi}\Lambda}_{\hat{\Theta}}$ encodes the local duality content (eq. (4.301)), and $\mathfrak{F}_{\vec{r}} = \mathcal{F}^{\mathrm{dist}}_{\vec{r}} \times \mathcal{F}^{\mathrm{time}}_{\vec{r}}$ is the fibre in the declared Hilbert bundle $\pi : \mathfrak{F} \to \mathcal{C}_d$ (eqs. (4.302) and (4.303)). This yields a surjection $\mathcal{C}_d \twoheadrightarrow \{\mathfrak{B}_{\vec{r}}\}$, so there are *at most* $\mathfrak{c}$ loci; and, under the standing indistinguishability criterion that anchors a distinct relational–null class at each $\vec{r}$, the quotient does not collapse cardinality. Consequently, the family of PQS loci is indeed continuum–many: distinct $\mathfrak{B}_{\vec{r}}$ smoothly organised over $\mathcal{C}_d$, each with its structured content and spectrally regulated fibre.

Does the continuum of PQS inherit the smooth manifold structure of the base $\mathcal{C}_d \cong \mathbb{R}^d$ (and thus have cardinality $\mathfrak{c}$)? A careful distinction among three levels is required. First, the base $\mathcal{C}_d$ is *stipulated* to be a smooth, connected, Hausdorff, second–countable d-manifold diffeomorphic to $\mathbb{R}^d$, and it serves as the indexing arena for the Hilbert bundle $\pi : \mathfrak{F} \to \mathcal{C}_d$ (eq. (4.302)). On this base, each anchor $\vec{r}$ is assigned $(\mathfrak{B}_{\vec{r}}, \hat{\Omega}_{\vec{r}}, \mathfrak{F}_{\vec{r}})$, with $\mathfrak{B}_{\vec{r}}$ the relational–null class set by $\vec{d}, \vec{t}$ (Definitions 4.599 and 4.600) and $\hat{\Omega}_{\vec{r}}$ recording the local structured content. In this *parametric* sense there are continuum–many PQS data because there are continuum–many anchors. Second, viewing the PQS loci as equivalence classes $\{\mathfrak{B}_{\vec{r}}\}$, there is always a surjection $\mathcal{C}_d \to \{\mathfrak{B}_{\vec{r}}\}$. If every relational–null class is singleton, then the map is a bijection and the continuum of loci literally inherits the base manifold structure (and cardinality). If some

classes are nontrivial, passing to loci involves a quotient that may identify anchors; the resulting space has the quotient topology and need not be diffeomorphic to $\mathbb{R}^d$ unless standard regularity conditions for smooth quotients hold. In all cases, because the map is surjective from a set of size $\mathfrak{c}$, the family of loci has cardinality $\leq \mathfrak{c}$; under the standing hypotheses used in this framework, the quotient does not collapse cardinality and one obtains continuum–many distinct loci. Third, operationally it is the *total space* $\mathfrak{F} = \bigsqcup_{\vec{r}} (\mathcal{F}_{\vec{r}}^{\mathrm{dist}} \times \mathcal{F}_{\vec{r}}^{\mathrm{time}})$ with projection π (eq. (4.302)) that carries the geometry: smoothness and local triviality are imposed along $\mathcal{C}_d$, and sections are coordinatised in the fourfold dialectic basis. Thus, if by "continuum of PQS'' one means the *indexed family* over $\mathcal{C}_d$, the manifold structure of the base organises the entire collection; if one means the *quotient space of loci* as a standalone object, inheriting the full manifold structure and diffeomorphism type of $\mathbb{R}^d$ depends on whether the equivalence classes are singleton (or, more generally, whether the quotient satisfies the usual smoothness conditions).

4.19 The Fibred Continuum of Primordial Quantum Singularities (Wordwise)

Definition 4.618 (The Fibred Continuum of Primordial Quantum Singularities (Wordwise))**.** Let $\mathcal{C}_d$ be a smooth, connected, Hausdorff, second–countable d-manifold diffeomorphic to $\mathbb{R}^d$ (hence $\mathrm{card}(\mathcal{C}_d) = \mathfrak{c}$). Choose a smooth embedding $\iota : \mathcal{C}_d \hookrightarrow \mathcal{X}_q$ into the emergent configuration space of the holosymmetric wordwise calculus and write $x = \iota(\vec{r})$. Fix a *wordwise* coherence width $\ell_{\mathrm{PQS}} := \lambda_{\mathrm{cutoff}}^{-1}$ with $\ell_{\mathrm{PQS}} \ll \ell_{\mathrm{P}}$, determined by the effective spectral cutoff of an admissible wordwise spectral generator $\mathbb{D}^{[\mathbf{W}]}(x)$ used for $\left(d_{\mathrm{spec}}^{[\cdot]}, t_{\mathrm{spec}}^{[\cdot]}\right)$ (Definitions 4.408, 4.547 and 4.583). Then:

1. **PQS loci as wordwise relational null classes.** For each $\vec{r} \in \mathcal{C}_d$ (i.e. $x = \iota(\vec{r})$), define the *Primordial Quantum Singularity* (PQS) associated to $\vec{r}$ to be the equivalence class

$$\mathfrak{B}_{\vec{r}}^{[\mathbf{W}]} := \left\{ \vec{r}' \in \mathcal{C}_d : \ \vec{d}^{[\mathbf{W}]}(x, \iota(\vec{r}')) = \mathbf{0} \text{ and } \vec{t}^{[\mathbf{W}]}(x, \iota(\vec{r}')) = \mathbf{0} \right\},$$

where $\vec{d}^{[\mathbf{W}]}$ and $\vec{t}^{[\mathbf{W}]}$ are the wordwise triads of distances and times (Definitions 4.582, 4.606 and 4.607). By Proposition 4.507 and Corollary 4.510, $\mathfrak{B}_{\vec{r}}^{[\mathbf{W}]}$ is the (word/gauge–invariant) minimal locus on which all emergent *wordwise* observables vanish; the underlying equivalence relation does not depend on $[\mathbf{W}]$ (cf. Remark 4.608).

2. **Structured singular content (wordwise presentation).** Each PQS $\mathfrak{B}_{\vec{r}}^{[\mathbf{W}]}$ carries the operator–theoretic content

$$\boxed{\hat{\Omega}_{\vec{r}}^{[\mathbf{W}]} := \overset{\circ}{\Lambda}_{\hat{\Phi}\hat{\Theta}},} \tag{4.305}$$

where $\overset{\circ}{\Lambda}$ is the universal duality code and $\hat{\Phi}, \hat{\Theta}$ are the (wordwise) Gaiamiti and Tantrakyma dynamical fields, respectively, as fixed in their operator formulations (wordwise versions of *Definitions* 6.118*and* 6.142). No additional structure is imposed here.

3. **Hilbert bundle of wordwise relational observables.** The *wordwise* relational bundle is the (locally trivial) Hilbert bundle

$$\boxed{\pi^{[\mathbf{W}]}:\ \mathfrak{F}^{[\mathbf{W}]}\ :=\ \bigsqcup_{\vec{r}\in\mathcal{C}_d}\Big(\mathcal{F}^{\mathrm{dist},[\mathbf{W}]}_{\iota(\vec{r})}\times\mathcal{F}^{\mathrm{time},[\mathbf{W}]}_{\iota(\vec{r})}\Big)\ \longrightarrow\ \mathcal{C}_d,} \tag{4.306}$$

with typical fibre the Hilbert direct sum $\mathcal{H}^{[\mathbf{W}]}_{\mathrm{typ}} := \mathcal{H}^{[\mathbf{W}]}_{\mathrm{dist}} \oplus \mathcal{H}^{[\mathbf{W}]}_{\mathrm{time}}$. Here $\mathcal{F}^{\mathrm{dist},[\mathbf{W}]}_x$ and $\mathcal{F}^{\mathrm{time},[\mathbf{W}]}_x$ are the completions (in the ℓ^2–type inner products induced by the wordwise Fourfold expectations and the calibrations $\mathbf{M}(x), \mathbf{N}(x)$, cf. Section 4.11.4 and eq. (4.272)) of the corresponding modal profile spaces underlying $\vec{d}^{[\mathbf{W}]}$ and $\vec{t}^{[\mathbf{W}]}$ at x. The projection is $\pi^{[\mathbf{W}]}(\phi) = \vec{r}$ for $\phi \in \mathfrak{F}^{[\mathbf{W}]}_{\vec{r}}$. Local trivializations are obtained by wordwise gauge–covariant parallel transport of the Fourfold/spectral data (Section 4.5), and the structure group acts fibrewise unitarily on $\mathcal{H}^{[\mathbf{W}]}_{\mathrm{typ}}$ [372, 248]. Fibres are coordinatized in a holosymmetric Fourfold modal alphabet at the anchor point (cf. the discussion preceding Definition 4.606).

4. **Ultra–sub–Planckian support (wordwise spectral projector).** For each $\vec{r} \in \mathcal{C}_d$ let $x = \iota(\vec{r})$ and let $\mathbb{D}^{[\mathbf{W}]}(x)$ be the admissible wordwise spectral generator of the spectral/time triad at x, with spectral measure $E^{(x;[\mathbf{W}])}(\cdot)$. Restrict the fibre by the projector onto the effective cutoff band:

$$\boxed{\mathfrak{F}^{[\mathbf{W}]}_{\vec{r}}\ =\ \mathrm{Ran}\Big(E^{(x;[\mathbf{W}])}\big([0,\lambda_{\mathrm{cutoff}}]\big)\Big),\qquad \ell_{\mathrm{PQS}} := \lambda^{-1}_{\mathrm{cutoff}} \ll \ell_{\mathrm{P}},} \tag{4.307}$$

i.e.

$$\begin{aligned}\mathfrak{F}^{[\mathbf{W}]}_{\vec{r}} &= \mathbf{1}_{[0,\lambda_{\mathrm{cutoff}}]}\big(\mathbb{D}^{[\mathbf{W}]}(x)\big)\,\mathcal{H}_x\ =\ \ker\Big(\mathbf{1} - E^{(x;[\mathbf{W}])}\big([0,\lambda_{\mathrm{cutoff}}]\big)\Big),\\ \psi \in \mathfrak{F}^{[\mathbf{W}]}_{\vec{r}} &\Longleftrightarrow E^{(x;[\mathbf{W}])}\big((\lambda_{\mathrm{cutoff}},\infty)\big)\,\psi = 0.\end{aligned} \tag{4.308}$$

Thus "ultra–sub–Planckian support'' asserts that all fibrewise spectral weight of admissible sections lies below $\lambda_{\mathrm{cutoff}}$, i.e. within the coherence width ℓ_{PQS}, consistent with Wordwise Spectral Smoothness and Finite Entanglement Capacity (Theorem 4.421 and Principle 4.2).

We call the resulting wordwise structure

$$\boxed{\mathcal{C}_d:\ \text{a fibred continuum of Primordial Quantum Singularities}\ \Big\{\ \mathfrak{B}^{[\mathbf{W}]}_{\vec{r}},\ \hat{\Omega}^{[\mathbf{W}]}_{\vec{r}},\ \mathfrak{F}^{[\mathbf{W}]}_{\vec{r}}\ \Big\}_{\vec{r}\in\mathcal{C}_d},}$$

canonically equivalent (fibrewise and in all induced distances/times) to its exogenous presentation by Proposition 4.613.

Calibrated distinguishability in the bundle (wordwise recap). For $x = \iota(\vec{r})$, the composite (wordwise) distinguishability between x and $x' = \iota(\vec{r}')$ reads

$$\boxed{D^{[\mathbf{W}]}(\vec{r},\vec{r}')\ :=\ \Big\|\mathbf{M}^{1/2}(x)\,\vec{d}^{[\mathbf{W}]}(x,x')\Big\|_2,} \tag{4.309}$$

with $\vec{d}^{[\mathbf{W}]}$ and $\vec{t}^{[\mathbf{W}]}$ the fibre coordinates provided by Definitions 4.606 and 4.607, and $\mathbf{M}(x) \succ 0$ any fixed calibration (cf. Section 4.11.4). By word/gauge invariance, $\vec{d}^{[\mathbf{W}]} = \vec{d}$ and $\vec{t}^{[\mathbf{W}]} = \vec{t}$, hence $D^{[\mathbf{W}]} = D$ (Proposition 4.613).

Remarks. (1) The embedding ι merely supplies a continuum label set; all observables are computed wordwise on $\mathcal{X}_q$ and are independent of $[\mathbf{W}]$. (2) The PQS class at $\vec{r}$ is the *wordwise relational null fibre* $Z^{[\mathbf{W}]}_{\iota(\vec{r})}$ (Corollary 4.609). (3) Ultra–sub–Planckianity of the fibres dovetails with the effective cutoff analysis and PQS width bounds of Sections 4.15.1 and 4.15.4.

4.19.1 Parallel Transport on the Relational Bundle

Setup. Working on the Hilbert bundle $\pi : \mathfrak{F} \to \mathcal{C}_d$ with fibres $\mathfrak{F}_{\vec{r}} = \mathcal{F}^{\mathrm{dist}}_{\vec{r}} \times \mathcal{F}^{\mathrm{time}}_{\vec{r}}$ (eq. (4.302)), we use the fourfold dialectic modal basis (Section 1.2.9) and the fibre inner products induced by coincident–limit Fourfold expectations. Throughout, $\gamma : [0,1] \to \mathcal{C}_d$, $s \mapsto \gamma(s)$, denotes a C^1 curve.

Operational parallel transport. There is a unique[5] family of fibrewise unitaries $U_\gamma(s \leftarrow s_0) : \mathfrak{F}_{\gamma(s_0)} \to \mathfrak{F}_{\gamma(s)}$ such that, for any section Ψ of $\mathfrak{F}$,

$$\boxed{\begin{aligned}&\Big\langle U_\gamma(s \leftarrow s_0)\Psi(\gamma(s_0)), U_\gamma(s \leftarrow s_0)\Phi(\gamma(s_0))\Big\rangle_{\gamma(s)} \\ &\qquad = \Big\langle \Psi(\gamma(s_0)), \Phi(\gamma(s_0))\Big\rangle_{\gamma(s_0)}.\end{aligned}} \tag{4.310}$$

and U_γ respects the fourfold decomposition on each fibre. We identify U_γ with the gauge–covariant transporter already used for Fourfold data, $U_\gamma(s \leftarrow s_0) = \mathcal{U}\big(\gamma(s) \leftarrow \gamma(s_0); \gamma\big)$, so that composition and reparametrisation invariance hold: $U_{\gamma_2 \odot \gamma_1} = U_{\gamma_2} U_{\gamma_1}$ and $U_{\gamma \circ \varphi}(s \leftarrow s_0) = U_\gamma(\varphi(s) \leftarrow \varphi(s_0))$.

Connection 1-form and covariant derivative. In any local trivialisation along γ, the transport is generated by a skew-adjoint operator-valued 1-form $\mathcal{A}_\gamma(s)$ via

$$\boxed{\frac{d}{ds}\, U_\gamma(s \leftarrow s_0) \;=\; -\,i\,\mathcal{A}_\gamma(s)\, U_\gamma(s \leftarrow s_0), \qquad \mathcal{A}_\gamma(s)^\dagger = \mathcal{A}_\gamma(s),} \tag{4.311}$$

with block structure compatible with $\mathcal{F}^{\mathrm{dist}} \oplus \mathcal{F}^{\mathrm{time}}$. For a section Ψ of $\mathfrak{F}$, the covariant derivative along γ is

$$\boxed{\big(\nabla_{\dot\gamma}\Psi\big)\big|_{\gamma(s)} := \lim_{h \to 0} \frac{U_\gamma(s \leftarrow s+h)\,\Psi(\gamma(s+h)) - \Psi(\gamma(s))}{h} \;=\; \Big(\partial_s + i\,\mathcal{A}_\gamma(s)\Big)\Psi(\gamma(s)).} \tag{4.312}$$

By (4.310), ∇ is metric-compatible with the Fourfold–induced inner product.

[5] Uniqueness here refers to the requirement of (i) fibrewise unitarity, (ii) preservation of Fourfold expectations, and (iii) functoriality under concatenation; cf. standard parallel transport uniqueness for metric-compatible connections [248, 372].

Propagation of triadic quantities. Let $\vec{d} = (d_{\mathrm{corr}}, d_{\mathrm{vac}}, d_{\mathrm{spec}})$ and $\vec{t} = (t_{\mathrm{corr}}, t_{\mathrm{vac}}, t_{\mathrm{spec}})$ be the triads on $\mathfrak{F}$ (Definitions 4.599 and 4.600). Along γ, we evaluate $x = \gamma(s_0)$ against $x' = \gamma(s)$ and pull back all fibre data at x' to x using $U_\gamma(s \leftarrow s_0)$. With this convention the path–rate relations read

$$\boxed{\begin{aligned} \frac{d}{ds}\, t_{\mathrm{corr}}(x, \gamma(s)) &= \frac{d}{ds}\, \operatorname{Arg} \langle \hat{G}_\Lambda \rangle_{x,\gamma(s)}, \\ \frac{d}{ds}\, t_{\mathrm{vac}}(x, \gamma(s)) &= \dot{t}_{\mathrm{vac}}(s), \\ \frac{d}{ds}\, t_{\mathrm{spec}}(x, \gamma(s)) &= \dot{t}_{\mathrm{spec}}(s). \end{aligned}} \tag{4.313}$$

with $\dot{t}_{\mathrm{vac}}$ and $\dot{t}_{\mathrm{spec}}$ the rate forms already defined via ℓ^2–norms of the transported modal profiles (see the path-rate constructions cited earlier). The corresponding accumulated times $T_{\mathrm{vac}}, T_{\mathrm{spec}}$ along γ control the bounds $t_{\mathrm{vac}} \le T_{\mathrm{vac}}$ and $t_{\mathrm{spec}} \le T_{\mathrm{spec}}$. Analogously, $\vec{d}(x, \gamma(s))$ is computed from the Fourfold correlator with the x'-fibre pulled back to x by U_γ, ensuring gauge–covariance of all components.

Interferometric transport and holonomy. Given two piecewise C^1 paths γ_1, γ_2 with the same endpoints, the loop holonomy on $\mathfrak{F}$ is

$$\boxed{\mathcal{H}[\gamma_1, \gamma_2] \;:=\; U_{\gamma_2}(1 \leftarrow 0)^{-1}\, U_{\gamma_1}(1 \leftarrow 0),} \tag{4.314}$$

a fibrewise unitary that reduces to the identity for homotopic paths with vanishing curvature. In the holosymmetric interferometry setting (two complementary branches; cf. the dual–register construction), the Born visibility is an expectation of $\mathcal{H}$ in the prepared Holo-State or register state, providing a gauge–invariant measure of "law interference" (*cf.* the visibility functional discussed previously).

Spectral regularity and domain control. All transports and generators above are taken on the spectrally regulated fibres $\mathfrak{F}_{\vec{r}} = \operatorname{Ran} E^{(\vec{r})}([0, \lambda_{\mathrm{cutoff}}])$ (eq. (4.303)). Coherent-regularisation (finite entanglement capacity and spectral smoothness) guarantees that $\mathcal{A}_\gamma(s)$ is densely defined on the regulated fibre and that the corresponding unitaries U_γ act within the ultra–sub–Planckian support, so the rates in (4.313) and the holonomy (4.314) are well-defined on the operational domain.

Summary. Equations (4.310)–(4.314) give a purely operational connection on $\mathfrak{F}$ that (i) preserves the Fourfold–induced inner product, (ii) transports the fourfold modal coordinates, and (iii) propagates triadic times and distances along curves in $\mathcal{C}_d$ in a gauge–covariant manner. This furnishes the needed infrastructure for chronogeometric and interferometric constructions downstream, without introducing assumptions beyond eq. (4.302), the fourfold basis (Section 1.2.9), and the spectral constraint eq. (4.303).

4.19.2 Parallel Transport on the Relational Bundle (Holosymmetric Wordwise)

Setup (wordwise). Work on the wordwise Hilbert bundle $\pi^{[\mathbf{W}]} : \mathfrak{F}^{[\mathbf{W}]} \to \mathcal{C}_d$ with fibres $\mathfrak{F}^{[\mathbf{W}]}_{\vec{r}} = \mathcal{F}^{\mathrm{dist},[\mathbf{W}]}_{\vec{r}} \times \mathcal{F}^{\mathrm{time},[\mathbf{W}]}_{\vec{r}}$ (eq. (4.306)). Use the Fourfold dialectic modal basis and the fibre inner products induced by coincident–limit wordwise Fourfold expectations (Definitions 4.606 and 4.607). Let $\gamma : [0,1] \to \mathcal{C}_d$, $s \mapsto \gamma(s)$, be C^1.

Operational wordwise parallel transport. There exists a unique[6] family of fibrewise unitaries

$$U^{[\mathbf{W}]}_{\gamma}(s \leftarrow s_0) : \ \mathfrak{F}^{[\mathbf{W}]}_{\gamma(s_0)} \longrightarrow \mathfrak{F}^{[\mathbf{W}]}_{\gamma(s)}$$

such that, for any sections Ψ, Φ of $\mathfrak{F}^{[\mathbf{W}]}$,

$$\boxed{\Big\langle U^{[\mathbf{W}]}_{\gamma}(s \leftarrow s_0)\Psi(\gamma(s_0)),\, U^{[\mathbf{W}]}_{\gamma}(s \leftarrow s_0)\Phi(\gamma(s_0))\Big\rangle_{\gamma(s)} = \Big\langle \Psi(\gamma(s_0)), \Phi(\gamma(s_0))\Big\rangle_{\gamma(s_0)},} \tag{4.315}$$

and $U^{[\mathbf{W}]}_{\gamma}$ respects the wordwise decomposition $\mathcal{F}^{\mathrm{dist},[\mathbf{W}]} \oplus \mathcal{F}^{\mathrm{time},[\mathbf{W}]}$. We identify it with the gauge–covariant transporter for the wordwise Fourfold data:

$$\begin{aligned} U^{[\mathbf{W}]}_{\gamma}(s \leftarrow s_0) &= \mathcal{U}^{[\mathbf{W}]}\Big(\gamma(s) \leftarrow \gamma(s_0); \gamma\Big), \\ U^{[\mathbf{W}]}_{\gamma_2 \circ \gamma_1} &= U^{[\mathbf{W}]}_{\gamma_2}\, U^{[\mathbf{W}]}_{\gamma_1}, \\ U^{[\mathbf{W}]}_{\varphi\circ\varphi}(s \leftarrow s_0) &= U^{[\mathbf{W}]}_{\gamma}\Big(\varphi(s) \leftarrow \varphi(s_0)\Big). \end{aligned} \tag{4.316}$$

Connection 1–form and covariant derivative (wordwise). In any local trivialisation along γ, transport is generated by a Hermitian operator–valued 1–form $\mathcal{A}^{[\mathbf{W}]}_{\gamma}(s)$ via

$$\boxed{\frac{d}{ds}\, U^{[\mathbf{W}]}_{\gamma}(s \leftarrow s_0) = -\,i\,\mathcal{A}^{[\mathbf{W}]}_{\gamma}(s)\, U^{[\mathbf{W}]}_{\gamma}(s \leftarrow s_0), \qquad \Big(\mathcal{A}^{[\mathbf{W}]}_{\gamma}(s)\Big)^{\dagger} = \mathcal{A}^{[\mathbf{W}]}_{\gamma}(s),} \tag{4.317}$$

with block structure compatible with $\mathcal{F}^{\mathrm{dist},[\mathbf{W}]} \oplus \mathcal{F}^{\mathrm{time},[\mathbf{W}]}$. For a section Ψ of $\mathfrak{F}^{[\mathbf{W}]}$, define the covariant derivative:

$$\boxed{\big(\nabla^{[\mathbf{W}]}_{\dot\gamma}\Psi\big)\Big|_{\gamma(s)} := \lim_{h\to 0} \frac{U^{[\mathbf{W}]}_{\gamma}(s \leftarrow s+h)\, \Psi(\gamma(s+h)) - \Psi(\gamma(s))}{h} = \Big(\partial_s + i\,\mathcal{A}^{[\mathbf{W}]}_{\gamma}(s)\Big)\Psi(\gamma(s)).}$$

(4.318)

By (4.315), $\nabla^{[\mathbf{W}]}$ is metric–compatible with the Fourfold–induced fibre inner product. The solution is the path–ordered exponential $U^{[\mathbf{W}]}_{\gamma}(s \leftarrow s_0) = \mathcal{P}\exp\Big(-i\int_{s_0}^{s} \mathcal{A}^{[\mathbf{W}]}_{\gamma}(\sigma)\, d\sigma\Big)$.

[6] Uniqueness under the wordwise analogues of: (i) fibrewise unitarity, (ii) preservation of Fourfold expectations, and (iii) functoriality under concatenation; cf. metric–compatible parallel transport [248, 372].

Curvature and holonomy (wordwise). On a chart with connection 1–form $\mathcal{A}^{[\mathbf{W}]}$, the curvature is

$$\boxed{\mathcal{F}^{[\mathbf{W}]} = d\mathcal{A}^{[\mathbf{W}]} + i\,\mathcal{A}^{[\mathbf{W}]}\wedge\mathcal{A}^{[\mathbf{W}]}, \qquad \mathrm{Hol}^{[\mathbf{W}]}(\gamma\ \text{loop}) = \mathcal{P}\exp\oint_{\gamma}(-i\,\mathcal{A}^{[\mathbf{W}]}).} \tag{4.319}$$

Nontrivial $\mathcal{F}^{[\mathbf{W}]}$ generates correlator/vacuum/spectral phase holonomies that enter the temporal triad (Definitions 4.533, 4.541 and 4.549).

Word–gauge covariance. Under a word/frame change $V^{[\mathbf{W}\to\mathbf{W}']}(s)$ on the fibre,

$$\boxed{\mathcal{A}_{\gamma}^{[\mathbf{W}']} = V^{\dagger}\mathcal{A}_{\gamma}^{[\mathbf{W}]}V - i\,V^{\dagger}\partial_s V, \qquad U_{\gamma}^{[\mathbf{W}']}(s\leftarrow s_0) = V^{\dagger}(s)\,U_{\gamma}^{[\mathbf{W}]}(s\leftarrow s_0)\,V(s_0),} \tag{4.320}$$

and the induced transports preserve the values of the distance/time triads (Propositions 4.340 and 4.422 and temporal analogues).

Propagation of triadic quantities (wordwise). Let $\vec{d}^{[\mathbf{W}]} = (d_{\mathrm{corr}}^{[\mathbf{W}]}, d_{\mathrm{vac}}^{[\mathbf{W}]}, d_{\mathrm{spec}}^{[\mathbf{W}]})$ and $\vec{t}^{[\mathbf{W}]} = (t_{\mathrm{corr}}^{[\mathbf{W}]}, t_{\mathrm{vac}}^{[\mathbf{W}]}, t_{\mathrm{spec}}^{[\mathbf{W}]})$. Fix $x = \gamma(s_0)$ and evaluate against $x' = \gamma(s)$, pulling back fibre data at x' to x with $U_{\gamma}^{[\mathbf{W}]}(s\leftarrow s_0)$. Then

$$\boxed{\begin{aligned}
&\frac{d}{ds}\,t_{\mathrm{corr}}^{[\mathbf{W}]}(x,\gamma(s)) = \mathrm{Im}\left(\frac{d}{ds}\log\Gamma_{\mathrm{corr}}^{[\mathbf{W}]}(x,\gamma(s))\right),\\
&\frac{d}{ds}\,t_{\mathrm{vac}}^{[\mathbf{W}]}(x,\gamma(s)) = \dot{t}_{\mathrm{vac}}^{[\mathbf{W}]}(s) = \left[\sum_k\left(\tfrac{d}{ds}\ \mathrm{Im}\,C_k^{[\mathbf{W}]}(\gamma(s))\right)^2\right]^{1/2},\\
&\frac{d}{ds}\,t_{\mathrm{spec}}^{[\mathbf{W}]}(x,\gamma(s)) = \dot{t}_{\mathrm{spec}}^{[\mathbf{W}]}(s) = \left[\sum_n\left(\tfrac{d}{ds}\left(w_n^{[\mathbf{W}]}\,\mathrm{Im}\,\mathcal{S}_n^{[\mathbf{W}]}\right)\right)^2\right]^{1/2},
\end{aligned}} \tag{4.321}$$

where $\Gamma_{\mathrm{corr}}^{[\mathbf{W}]}$ is the wordwise bilocal correlator expectation, $C_k^{[\mathbf{W}]}$ the diagonal self–correlators, and $\mathcal{S}_n^{[\mathbf{W}]}$ the wordwise spectral shear (Definitions 4.339, 4.408 and 4.468).

Proposition 4.619 (Metric compatibility and functoriality (wordwise))**.** *The covariant derivative $\nabla^{[\mathbf{W}]}$ of (4.318) is compatible with the Fourfold–induced inner product, and $U_{\gamma}^{[\mathbf{W}]}$ is functorial under concatenation and reparametrisation (4.316). Consequently, norms that define the triadic distances and the associated temporal rates are preserved under parallel transport along γ (up to their explicit path–dependence encoded by the correlator/spectral functionals).*

Remark 4.620 (Block structure and calibrated mixing)**.** In the canonical gauge the connection $\mathcal{A}^{[\mathbf{W}]}$ is block–diagonal w.r.t. $\mathcal{F}^{\mathrm{dist},[\mathbf{W}]}\oplus\mathcal{F}^{\mathrm{time},[\mathbf{W}]}$. If a nontrivial calibration matrix $\mathbf{M}(x)$ couples the axes (Definition 4.494), one may allow controlled off–diagonal blocks in $\mathcal{A}^{[\mathbf{W}]}$ consistent with metric compatibility; holonomy then mixes components within the calibrated fibre without altering the wordwise invariants of the composite metric.

4.19.3 The Quantum Configuration Space

The quantum configuration space is not fundamental; it *emerges* from the fibred PQS continuum $\mathcal{C}_d$ (Section 4.18) by identifying loci that are operationally indistinguishable with respect to the relational observables encoded in the bundle $\pi : \mathfrak{F} \to \mathcal{C}_d$ (eqs. (4.302) and (4.303)) and compared via the gauge–covariant parallel transport of Section 4.19.1.

Reference: Quotient Manifold Theorems

Theorem 4.621 (Quotient Manifold — Group Action [249, Thm. 21.10])**.** *If a Lie group G acts smoothly, freely, and properly on a smooth manifold M, then the orbit space M/G is a smooth manifold of dimension $\dim M - \dim G$, the projection $\pi : M \to M/G$ is a smooth submersion, and $M \to M/G$ is a principal G-bundle.*

Theorem 4.622 (Quotient Manifold — General Form [411])**.** *Let M be a smooth manifold and $\sim$ an equivalence relation on M. If each class is a closed embedded submanifold of M, the relation set $R = \{(p, q) \in M \times M : \ p \sim q\}$ is a closed embedded submanifold of $M \times M$, and the projection $\pi : M \to M/\sim$ is a smooth submersion, then $M/\sim$ is a smooth manifold and π is a smooth submersion.*

Definition 4.623 (Operational indistinguishability)**.** For $x, x' \in \mathcal{C}_d$, write $\mathscr{R}(x, x') := \big(\vec{d}(x, x'), \vec{t}(x, x')\big) \in \mathbb{R}^3 \times \mathbb{R}^3$, where $\vec{d}$ and $\vec{t}$ are the triads from the distinguishability and temporal-flow fibres (Definitions 4.599 and 4.600). We declare

$$x \sim x' \quad :\Longleftrightarrow \quad \mathscr{R}(x, x') = \mathbf{0},$$

i.e. all triadic components vanish when the x'-fibre is compared to the x-fibre by the transporter of Section 4.19.1. The *quantum configuration space* is the quotient

$$\mathcal{X}_q \ := \ \mathcal{C}_d \,/\, \sim, \qquad \pi_q : \mathcal{C}_d \to \mathcal{X}_q \ \text{ the canonical projection.} \tag{4.322}$$

Gauge covariance and well-posedness. Because the correlator $\hat{G}_\Lambda$ and the transporter $\mathcal{U}$ transform covariantly and the distances/times are built from gauge-invariant combinations, $\mathscr{R}$ is invariant under fibre frame changes; hence $\sim$ is gauge-well-defined (cf. Section 4.19.1).

Quotient Manifold Theorem and its applicability to $\mathcal{X}_q$

We seek conditions under which $\mathcal{X}_q$ inherits a smooth structure from $\mathcal{C}_d$. Two standard routes apply:

(i) *Free, proper group action* [249, Thm. 21.10]: if $\sim$ is the orbit relation of a smooth, free, proper Lie-group action on $\mathcal{C}_d$, then $\mathcal{C}_d/\sim$ is a smooth manifold and π_q is a smooth submersion (indeed a principal bundle).

(ii) *General equivalence relations* [411, Ch. 14]: if the relation set $R := \{(x, x') \in \mathcal{C}_d \times \mathcal{C}_d : x \sim x'\}$ is a closed embedded submanifold, each class $[x]$ is a closed embedded submanifold of $\mathcal{C}_d$, and the projection π_q is a smooth submersion, then $\mathcal{X}_q$ carries a unique smooth structure making π_q a submersion.

How our data meet the hypotheses (no new lemmas). Define the *relational map*

$$\mathcal{R}: \mathcal{C}_d \times \mathcal{C}_d \longrightarrow \mathbb{R}^6, \qquad \mathcal{R}(x,x') := \Big(\vec{d}(x,x'), \vec{t}(x,x')\Big). \tag{4.323}$$

By construction of the fibres and transport (eq. (4.302) and Section 4.19.1), $\mathcal{R}$ is smooth on its operational domain; the equivalence relation is the zero-level set $R = \mathcal{R}^{-1}(\{0\})$. Under the standing regularity inherent in the triad definitions (smooth dependence on anchors, constant-rank behaviour inherited from coherent regularisation and ultra–sub–Planckian support, eq. (4.303)), the following hold in the regime of interest:

(Q1) *Embeddedness of classes.* For each x, the fibre $\{x' : \mathcal{R}(x,x') = 0\}$ is a closed embedded submanifold (a "relational leaf" through x) by the preimage theorem [249, Thm. 4.12].

(Q2) *Embeddedness of the relation.* $R = \mathcal{R}^{-1}(\{0\}) \subset \mathcal{C}_d \times \mathcal{C}_d$ is a closed embedded submanifold whenever $\mathcal{R}$ has constant rank near R.

(Q3) *Submersion.* The differential $d\pi_q$ has full rank: equivalently, the restriction $\mathrm{pr}_1 : R \to \mathcal{C}_d$ is a submersion. This is automatic when the classes vary smoothly with constant dimension (constant-rank foliation condition), which is the generic case in our regulated setting.

When (Q1)–(Q3) hold, $\mathcal{X}_q$ is a smooth manifold and π_q a smooth submersion by [411, Ch. 14]. In particular, if each class has dimension m (locally constant), then $\dim \mathcal{X}_q = d - m$.

Topological properties. If, in addition, R is closed and the class distribution is locally finite (a consequence of ultra–sub–Planckian support, eq. (4.303)), then $\mathcal{X}_q$ is Hausdorff and second–countable; π_q is an open map. Cardinality is preserved: $|\mathcal{X}_q| = |\mathcal{C}_d| = \mathfrak{c}$ (as recorded earlier), so the emergent arena retains continuum cardinality while quotienting out operational redundancies.

Interpretation. The quotient $\mathcal{X}_q$ is the space of *operational positions*: two anchors are indistinguishable precisely when all triadic diagnostics vanish once compared by the canonical transporter. The smooth structure on $\mathcal{X}_q$ is therefore not postulated but *inherited* from $\mathcal{C}_d$ under mild regularity of the relational map $\mathcal{R}$, and the differential of π_q selects directions in $\mathcal{C}_d$ transverse to the relational leaves as bona fide "configurational" directions. Subsequent dynamical fields and chronogeometric tensors may be pushed forward along π_q, or, equivalently, pulled back from $\mathcal{X}_q$ to $\mathcal{C}_d$ along sections of π_q, without ambiguity.

Remark 4.624 (Application to the IQR)**.** We verify the hypotheses of Theorem 4.622 for the quotient $\pi_q : \mathcal{C}_d \to \mathcal{X}_q$ defined in Definition 4.623 and eq. (4.322).

1. **Each equivalence class is an embedded submanifold.** For fixed $x \in \mathcal{C}_d$, the class $[x] = \{x' \in \mathcal{C}_d : \mathscr{R}(x,x') = \mathbf{0}\}$ is the zero–level set of the smooth map $x' \mapsto \mathscr{R}(x,x') = (\vec{d}(x,x'), \vec{t}(x,x'))$ (the triadic data compared via the transporter of Section 4.19.1). Under the standing regularity (smooth dependence and constant rank on the operational domain ensured by coherent–regularisation and ultra–sub–Planckian support, eqs. (4.199) and (4.303) and Principle 4.1), the preimage theorem [249,

Thm. 4.12] implies $[x]$ is a closed, smoothly embedded submanifold ("relational leaf"). Moreover, eq. (4.303) localises each leaf to an $\mathcal{O}(\ell_{\mathrm{PQS}})$ neighbourhood.

2. **The relation set $R \subset \mathcal{C}_d \times \mathcal{C}_d$ is closed and embedded.** Define the relational map $\mathcal{R} : \mathcal{C}_d \times \mathcal{C}_d \to \mathbb{R}^6$ by eq. (4.323). Then $R = \{(x, x') : \ x \sim x'\} = \mathcal{R}^{-1}(\{0\})$ is closed (preimage of a closed set under a continuous map). When $\mathcal{R}$ has constant rank near R (the same regularity as above), the constant–rank/preimage theorem yields that R is a closed embedded submanifold of $\mathcal{C}_d \times \mathcal{C}_d$ [411, Ch. 14].
3. **The projection $\pi_q : \mathcal{C}_d \to \mathcal{X}_q$ is a smooth submersion.** Equivalently, the restriction $\mathrm{pr}_1 : R \to \mathcal{C}_d$ is a submersion. Under the leafwise constant–dimension assumption (again, a consequence of the constant–rank behaviour of $\mathcal{R}$), the foliation by relational leaves admits local slices transverse to $[x]$, so $d\pi_q$ has full rank everywhere. Hence the quotient inherits a unique smooth structure making π_q a smooth submersion [411, Ch. 14].

Therefore the emergent configuration space $\mathcal{X}_q = \mathcal{C}_d/\sim$ is a smooth manifold, with π_q a smooth submersion. In particular, if the leaves have locally constant dimension m, then $\dim \mathcal{X}_q = d - m$; Hausdorffness and second–countability follow from the closed–embedded relation and the local finiteness implied by eq. (4.303).

Emergence of $\mathcal{X}_q$ as a Relational Quotient Space

Proposition 4.625 (Emergence of $\mathcal{X}_q$ as a Relational Quotient Space)**.** *The quantum configuration space $\mathcal{X}_q$ is defined as the quotient space:*

$$\mathcal{X}_q := \mathcal{C}_d/\sim \tag{4.324}$$

where the equivalence relation $\sim$ on $\mathcal{C}_d$ is defined by the simultaneous vanishing of all relational observables:

$$\vec{r} \sim \vec{r}' \quad \Longleftrightarrow \quad \vec{d}(\vec{r}, \vec{r}') = 0 \text{ and } \vec{t}(\vec{r}, \vec{r}') = 0. \tag{4.325}$$

Each point $x \in \mathcal{X}_q$ thus corresponds to an equivalence class of Primordial Quantum Singularities, $[\mathfrak{B}_{\vec{r}}]$, that are operationally indistinguishable.

Proof. By Corollary 4.601 the relation $\sim$ defined by the simultaneous vanishing of the triads $\vec{d}, \vec{t}$ is an equivalence relation on $\mathcal{C}_d$, and its classes are exactly the PQS relational–null loci $\mathfrak{B}_x = \{x' : \vec{d}(x, x') = \vec{0} = \vec{t}(x, x')\}$. Define the *relational map*

$$\mathcal{R} : \ \mathcal{C}_d \times \mathcal{C}_d \longrightarrow \mathbb{R}^6, \qquad \mathcal{R}(x, x') \ := \ \Big(\vec{d}(x, x'),\, \vec{t}(x, x')\Big). \tag{4.326}$$

By construction of the fibres and the gauge–covariant transporter in Section 4.19.1, the entries of $\mathcal{R}$ are obtained by inserting the parallel–transported fibre data into the Fourfold correlators and then taking the regulated expectations; local triviality of $\pi : \mathfrak{F} \to \mathcal{C}_d$ together with spectral localisation $\mathfrak{F}_x = \mathrm{Ran}\, E^{(x)}([0, \lambda_{\mathrm{cutoff}}])$ (eq. (4.303)) implies that $\mathcal{R}$ depends smoothly on (x, x') on the operational domain. Gauge covariance is preserved: under $g : \mathcal{C}_d \to \mathrm{U}(\mathcal{H}_{\mathrm{em}})$ we have the transform laws of eq. (4.16), and the zero set of $\mathcal{R}$ is invariant; hence $\sim$ is frame–independent.

Let $R := \{(x, x') \in \mathcal{C}_d \times \mathcal{C}_d : \ x \sim x'\} = \mathcal{R}^{-1}(\{0\})$. Continuity of $\mathcal{R}$ shows R is closed. By coherent–regularisation and ultra–sub–Planckian support (eqs. (4.199) and (4.303) and Principle 4.1) the differential $D\mathcal{R}$ has constant rank in a neighbourhood of R; therefore, by the preimage/constant–rank theorem (e.g. [249, Thm. 4.12]) R is a closed embedded submanifold of $\mathcal{C}_d \times \mathcal{C}_d$. Fix $x \in \mathcal{C}_d$ and consider the slice map $x' \mapsto \mathcal{R}(x, x')$; the same rank condition implies that $\{x' : \mathcal{R}(x, x') = 0\} = [x] = \mathfrak{B}_x$ is a closed, smoothly embedded submanifold ("relational leaf") of $\mathcal{C}_d$. Moreover, eq. (4.303) localises each leaf to an $\mathcal{O}(\ell_{\mathrm{PQS}})$ neighbourhood of x.

We now prove that the quotient $\pi_q : \mathcal{C}_d \to \mathcal{X}_q := \mathcal{C}_d/\!\sim$ (eq. (4.322)) is a smooth submersion and that $\mathcal{X}_q$ inherits a unique smooth structure. Because the leaves $\{\mathfrak{B}_x\}$ have locally constant dimension (the constant–rank condition), the relation R defines a regular foliation; standard foliation theory yields, through each x, a submanifold slice $S_x \subset \mathcal{C}_d$ transversal to $\mathfrak{B}_x$ and meeting nearby leaves exactly once. The map $\pi_q|_{S_x} : S_x \to \mathcal{X}_q$ is a diffeomorphism onto an open neighbourhood of $[x]$, and these slice charts assemble into a smooth atlas on $\mathcal{X}_q$. In differential form, transversality of S_x gives that $d\pi_q$ has full rank everywhere, i.e. π_q is a smooth submersion. Therefore the hypotheses of the general quotient–manifold theorem (Theorem 4.622, cf. [411, Ch. 14]) are met: each class is a closed embedded submanifold, the relation set R is a closed embedded submanifold, and the projection is a submersion. Consequently, $\mathcal{X}_q$ is a smooth manifold and, if the leaf dimension $m := \dim \mathfrak{B}_x$ is locally constant, $\dim \mathcal{X}_q = d - m$.

Topological properties follow from the same structure. Since R is closed and the leaves are uniformly localised (eq. (4.303)), distinct classes admit disjoint saturated neighbourhoods; thus $\mathcal{X}_q$ is Hausdorff. Second–countability is inherited because $\mathcal{C}_d$ is second–countable and π_q is open with a countable slice basis. Cardinality is preserved as π_q is surjective and $|\mathcal{C}_d| = \mathfrak{c}$, hence $|\mathcal{X}_q| = \mathfrak{c}$. This completes the proof that $\mathcal{X}_q$ emerges as a well-defined smooth, Hausdorff, second–countable manifold from the relational quotient of the PQS continuum, with π_q a smooth submersion compatible with the operational/gauge structure of Section 4.19.1. □

4.19.4 The Quantum Configuration Space (Holosymmetric Wordwise)

The quantum configuration space is *derived*, not postulated. It emerges from the fibred PQS continuum $\mathcal{C}_d$ (Section 4.19) by identifying loci that are *operationally indistinguishable* with respect to the *wordwise* relational observables encoded in the bundle $\pi^{[\mathbf{W}]} : \mathfrak{F}^{[\mathbf{W}]} \to \mathcal{C}_d$ (eqs. (4.306) and (4.307)) and compared via the gauge–covariant parallel transport of Section 4.19.2.

Reference: Quotient Manifold Theorems (for context)

Theorem 4.626 (Quotient Manifold — Group Action [249, Thm. 21.10])**.** *If a Lie group G acts smoothly, freely, and properly on a smooth manifold M, then M/G is a smooth manifold of dimension $\dim M - \dim G$, the projection $\pi : M \to M/G$ is a smooth submersion, and $M \to M/G$ is a principal G–bundle.*

Theorem 4.627 (Quotient Manifold — General Form [411])**.** *Let M be a smooth manifold and $\sim$ an equivalence relation on M. If each class is a closed embedded submanifold of M, the*

relation set $R = \{(p,q) \in M \times M : \ p \sim q\}$ is a closed embedded submanifold of $M \times M$, and $\pi : M \to M/\sim$ is a smooth submersion, then $M/\sim$ is a smooth manifold and π is a smooth submersion.

Wordwise operational indistinguishability and the quotient

Definition 4.628 (Wordwise operational indistinguishability)**.** For $x, x' \in \mathcal{C}_d$, define the wordwise triads

$$\vec{d}^{[\mathbf{W}]}(x,x') = \left(d^{[\mathbf{W}]}_{\mathrm{corr}}, d^{[\mathbf{W}]}_{\mathrm{vac}}, d^{[\mathbf{W}]}_{\mathrm{spec}}\right), \qquad \vec{t}^{[\mathbf{W}]}(x,x') = \left(t^{[\mathbf{W}]}_{\mathrm{corr}}, t^{[\mathbf{W}]}_{\mathrm{vac}}, t^{[\mathbf{W}]}_{\mathrm{spec}}\right),$$

as in Definitions 4.339, 4.408, 4.468, 4.529, 4.539 and 4.547, with the x'–fibre compared to the x–fibre by the transporter $U^{[\mathbf{W}]}_{\gamma}$ of Section 4.19.2. Set

$$\mathscr{R}^{[\mathbf{W}]}(x,x') := \left(\vec{d}^{[\mathbf{W}]}(x,x'),\, \vec{t}^{[\mathbf{W}]}(x,x')\right) \in \mathbb{R}^3 \times \mathbb{R}^3.$$

Declare

$$\boxed{x \sim_{[\mathbf{W}]} x' \quad :\Longleftrightarrow \quad \mathscr{R}^{[\mathbf{W}]}(x,x') = \mathbf{0}.} \tag{4.327}$$

The *wordwise quantum configuration space* is the quotient

$$\boxed{\mathcal{X}^{[\mathbf{W}]}_q := \mathcal{C}_d \,/\, \sim_{[\mathbf{W}]}, \qquad \pi^{[\mathbf{W}]}_q : \mathcal{C}_d \to \mathcal{X}^{[\mathbf{W}]}_q \ \text{ the canonical projection.}} \tag{4.328}$$

Proposition 4.629 (Equivalence, PQS classes, and quotient metric)**.** *(i) $\sim_{[\mathbf{W}]}$ is an equivalence relation. (ii) Its classes coincide with the wordwise PQS classes: $[x]_{\sim_{[\mathbf{W}]}} = \mathfrak{B}^{[\mathbf{W}]}_x$ (Definition 4.582 and Proposition 4.507). (iii) Writing $D^{[\mathbf{W}]}(x,x') = \|\vec{d}^{[\mathbf{W}]}(x,x')\|_2$ (Section 4.11.2), the rule*

$$\overline{D}^{[\mathbf{W}]}\left([x],[x']\right) := D^{[\mathbf{W}]}(x,x')$$

is a well–defined metric on $\mathcal{X}^{[\mathbf{W}]}_q$ and makes $\pi^{[\mathbf{W}]}_q : \left(\mathcal{C}_d, D^{[\mathbf{W}]}\right) \to \left(\mathcal{X}^{[\mathbf{W}]}_q, \overline{D}^{[\mathbf{W}]}\right)$ a nonexpansive submetry. Sketch. *Reflexivity: $\mathscr{R}^{[\mathbf{W}]}(x,x) = \mathbf{0}$. Symmetry: each component is symmetric (or antisymmetric modulo 2π with vanishing value implying symmetry). Transitivity follows from the triangle inequality for each component together with the zero–value condition. PQS identification is Proposition 4.507. Well–definedness of $\overline{D}^{[\mathbf{W}]}$ uses definiteness on the quotient.*

Proof. Write, for brevity, $\mathscr{R}^{[\mathbf{W}]}(x,x') := \left(\vec{d}^{[\mathbf{W}]}(x,x'),\, \vec{t}^{[\mathbf{W}]}(x,x')\right) \in \mathbb{R}^3 \times \mathbb{R}^3$ with $\vec{d}^{[\mathbf{W}]} = (d^{[\mathbf{W}]}_{\mathrm{corr}}, d^{[\mathbf{W}]}_{\mathrm{vac}}, d^{[\mathbf{W}]}_{\mathrm{spec}})$, $\vec{t}^{[\mathbf{W}]} = (t^{[\mathbf{W}]}_{\mathrm{corr}}, t^{[\mathbf{W}]}_{\mathrm{vac}}, t^{[\mathbf{W}]}_{\mathrm{spec}})$. Recall (a) each spatial component is a nonnegative, symmetric pseudometric obeying the triangle inequality (*wordwise* versions of the basic properties proved with the Fourfold/gauge transport), and (b) $t^{[\cdot]}_{\mathrm{vac}}$ and $t^{[\cdot]}_{\mathrm{spec}}$ are nonnegative pseudometrics with $t^{[\cdot]}_{\mathrm{vac}} \le d^{[\cdot]}_{\mathrm{vac}}$ and $t^{[\cdot]}_{\mathrm{spec}} \le d^{[\cdot]}_{\mathrm{spec}}$ (Proposition 4.540, Proposition 4.548); $f := t^{[\cdot]}_{\mathrm{corr}}$ enjoys antisymmetry (mod 2π), concatenation on a fixed homotopy class, and node–regularised continuity (Proposition 4.532, Remark 4.530).

(i) $\sim_{[\mathbf{W}]}$ is an equivalence relation. By definition, $x \sim_{[\mathbf{W}]} x'$ iff $\mathscr{R}^{[\mathbf{W}]}(x,x') = \mathbf{0}$. Reflexivity follows from the coincident–limit normalisations and $t^{[\cdot]}(x,x) = 0$ (Definition 4.529, Definition 4.539, Definition 4.547): $\mathscr{R}^{[\mathbf{W}]}(x,x) = \mathbf{0}$.

Symmetry: $d^{[\cdot]}_{\mathrm{corr}}, d^{[\cdot]}_{\mathrm{vac}}, d^{[\cdot]}_{\mathrm{spec}}$ are symmetric, and $t^{[\cdot]}_{\mathrm{vac}}, t^{[\cdot]}_{\mathrm{spec}} \geq 0$ are symmetric pseudometrics. For $t^{[\cdot]}_{\mathrm{corr}}$, antisymmetry gives $t^{[\mathbf{W}_2,\mathbf{W}_1]}_{\mathrm{corr}}(x', x; \bar{\gamma}) \equiv -\, t^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{corr}}(x, x'; \gamma) \pmod{2\pi}$ (Proposition 4.532); thus if $t^{[\cdot]}_{\mathrm{corr}}(x, x') = 0$ then $t^{[\cdot]}_{\mathrm{corr}}(x', x) = 0$ as well. Hence $\mathscr{R}^{[\mathbf{W}]}(x, x') = \mathbf{0}$ implies $\mathscr{R}^{[\mathbf{W}]}(x', x) = \mathbf{0}$.

Transitivity: Let $x \sim_{[\mathbf{W}]} y$ and $y \sim_{[\mathbf{W}]} z$. For the spatial components, the triangle inequality yields $d^{[\cdot]}_i(x, z) \leq d^{[\cdot]}_i(x, y) + d^{[\cdot]}_i(y, z) = 0$ for $i \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$, hence $d^{[\cdot]}_i(x, z) = 0$. For $t^{[\cdot]}_{\mathrm{vac}}$ and $t^{[\cdot]}_{\mathrm{spec}}$ the same argument applies since they are pseudometrics: $t^{[\cdot]}_{\mathrm{vac}}(x, z) \leq t^{[\cdot]}_{\mathrm{vac}}(x, y) + t^{[\cdot]}_{\mathrm{vac}}(y, z) = 0$, and similarly for $t^{[\cdot]}_{\mathrm{spec}}$. For $t^{[\cdot]}_{\mathrm{corr}}$, use the concatenation property on node–free homotopy classes (Proposition 4.532) and the standard node–regularisation (Remark 4.530): pick admissible paths γ_{xy}, γ_{yz} and define $\gamma_{xz} := \gamma_{yz} \circ \gamma_{xy}$; then the unwrapped phase obeys $\tilde{t}^{[\cdot]}_{\mathrm{corr}}(x, z; \gamma_{xz}) = \tilde{t}^{[\cdot]}_{\mathrm{corr}}(x, y; \gamma_{xy}) + \tilde{t}^{[\cdot]}_{\mathrm{corr}}(y, z; \gamma_{yz}) = 0$, hence $t^{[\cdot]}_{\mathrm{corr}}(x, z) \equiv 0 \pmod{2\pi}$ and, by continuity at 0, $t^{[\cdot]}_{\mathrm{corr}}(x, z) = 0$. Therefore $\mathscr{R}^{[\mathbf{W}]}(x, z) = \mathbf{0}$, so $\sim_{[\mathbf{W}]}$ is transitive.

(ii) Classes coincide with the wordwise PQS classes. By the wordwise *relational null condition* (Definition 4.582), $\mathfrak{B}^{[\mathbf{W}]}_x = \{x' : \vec{d}^{[\mathbf{W}]}(x, x') = \mathbf{0},\ \vec{t}^{[\mathbf{W}]}(x, x') = \mathbf{0}\}$, which is precisely the $\sim_{[\mathbf{W}]}$–equivalence class of x. Equivalently, the identification follows from the zero–locus characterisation of the PQS (Proposition 4.507).

(iii) $\overline{D}^{[\mathbf{W}]}$ is a well–defined metric and $\pi^{[\mathbf{W}]}_q$ is a nonexpansive submetry.

Well–definedness (independence of representatives). Suppose $x \sim_{[\mathbf{W}]} x_1$ and $x' \sim_{[\mathbf{W}]} x'_1$. By the triangle inequality for $D^{[\mathbf{W}]} = \|\vec{d}^{[\mathbf{W}]}\|_2$,

$$D^{[\mathbf{W}]}(x_1, x'_1) \;\leq\; D^{[\mathbf{W}]}(x_1, x) + D^{[\mathbf{W}]}(x, x') + D^{[\mathbf{W}]}(x', x'_1) \;=\; D^{[\mathbf{W}]}(x, x').$$

Exchanging the roles of (x, x') and (x_1, x'_1) yields the reverse inequality, hence $D^{[\mathbf{W}]}(x_1, x'_1) = D^{[\mathbf{W}]}(x, x')$. Therefore $\overline{D}^{[\mathbf{W}]}\big([x], [x']\big) := D^{[\mathbf{W}]}(x, x')$ is independent of the choices of representatives.

Metric axioms on the quotient. Nonnegativity and symmetry are inherited from $D^{[\mathbf{W}]}$. For the triangle inequality, choose representatives to obtain

$$\overline{D}^{[\mathbf{W}]}\big([x], [z]\big) \;=\; D^{[\mathbf{W}]}(x, z) \;\leq\; D^{[\mathbf{W}]}(x, y) + D^{[\mathbf{W}]}(y, z) \;=\; \overline{D}^{[\mathbf{W}]}\big([x], [y]\big) + \overline{D}^{[\mathbf{W}]}\big([y], [z]\big).$$

For definiteness (identity of indiscernibles), assume $\overline{D}^{[\mathbf{W}]}\big([x], [x']\big) = 0$. Then $D^{[\mathbf{W}]}(x, x') = 0$, hence *each* spatial component vanishes: $d^{[\cdot]}_{\mathrm{corr}}(x, x') = d^{[\cdot]}_{\mathrm{vac}}(x, x') = d^{[\cdot]}_{\mathrm{spec}}(x, x') = 0$. By the duality bounds (eq. (4.250), eq. (4.256)), $t^{[\cdot]}_{\mathrm{vac}}(x, x') = t^{[\cdot]}_{\mathrm{spec}}(x, x') = 0$. For $t^{[\cdot]}_{\mathrm{corr}}$, note that $d^{[\cdot]}_{\mathrm{corr}}(x, x') = 0$ implies Cauchy–Schwarz saturation of the normalised bilocal overlap in the Holo–State sector (the "Pancharatnam–parallelity'' condition for the Fourfold channels in the chosen word frame); together with the state–conditioned flatness implied by the full spatial zero locus (Proposition 4.519), we may choose an admissible gauge/readout in which the transport is trivial and the saturated overlap is real nonnegative, hence its phase is 0 (Proposition 4.531, Proposition 4.532). Gauge/word invariance then yields $t^{[\cdot]}_{\mathrm{corr}}(x, x') = 0$. Thus $\vec{t}^{[\cdot]}(x, x') = \mathbf{0}$, i.e. $x \sim_{[\mathbf{W}]} x'$ by (i), and so $[x] = [x']$. Therefore $\overline{D}^{[\mathbf{W}]}$ is a bona fide metric on $\mathcal{X}^{[\mathbf{W}]}_q$.

Nonexpansive submetry. By definition, $\overline{D}^{[\mathbf{W}]}\big(\pi^{[\mathbf{W}]}_q(x), \pi^{[\mathbf{W}]}_q(x')\big) = D^{[\mathbf{W}]}(x, x')$, so $\pi^{[\mathbf{W}]}_q$ is 1–Lipschitz (nonexpansive). Let $B_D(x, r)$ and $B_{\overline{D}}([x], r)$ denote the open balls in the

domain/quotient metrics. If $z \in B_D(x,r)$, then $\overline{D}^{[\mathbf{W}]}\big([x],[z]\big) = D^{[\mathbf{W}]}(x,z) < r$, so $\pi_q^{[\mathbf{W}]}(B_D(x,r)) \subseteq B_{\overline{D}}([x],r)$. Conversely, if $[w] \in B_{\overline{D}}([x],r)$, pick any representative $z \in [w]$; by the representative–independence shown above, $D^{[\mathbf{W}]}(x,z) = \overline{D}^{[\mathbf{W}]}\big([x],[w]\big) < r$, so $z \in B_D(x,r)$ and $\pi_q^{[\mathbf{W}]}(z) = [w]$. Hence $\pi_q^{[\mathbf{W}]}(B_D(x,r)) = B_{\overline{D}}([x],r)$, i.e. $\pi_q^{[\mathbf{W}]}$ is a submetry.

This proves (i)–(iii). □

Proposition 4.630 (Word–invariance of the quotient)**.** *If $\mathbf{W} \mapsto \mathbf{W}'$ is an admissible word change, then $\mathscr{R}^{[\mathbf{W}']}(x,x') = \mathscr{R}^{[\mathbf{W}]}(x,x')$ and hence $\sim_{[\mathbf{W}']} = \sim_{[\mathbf{W}]}$. Consequently there is a canonical isometric identification $\mathcal{X}_q^{[\mathbf{W}']} \cong \mathcal{X}_q^{[\mathbf{W}]}$.* Reason. *Wordwise invariance of all six components (Propositions 4.410 and 4.422 and temporal analogues) implies equality of the relations and of the induced metrics.*

Proof. Fix an admissible word/frame change $\mathbf{W} \mapsto \mathbf{W}'$. By definition this is implemented by pointwise fibre unitaries $V(x), V(x') \in \mathrm{U}(\mathcal{H}_{\mathrm{modal}})$ at the endpoints and the corresponding covariant update of the transporter $\mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x \leftarrow x';\gamma) \mapsto \mathcal{U}^{[\mathbf{W}'_1,\mathbf{W}'_2]}(x \leftarrow x';\gamma) = V(x)\,\mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x \leftarrow x';\gamma)\,V(x')^\dagger$ (cf. the word–gauge rule (4.320)).

Step 1: invariance of the three $\vec{d}$ components.

Correlator distance. The complex bilocal expectation transforms as

$$\Gamma_{\mathrm{corr}}^{[\mathbf{W}'_1,\mathbf{W}'_2]}(x,x';\gamma) = \Big\langle \hat{G}_\Lambda\big[\,(V(x)U_1\psi)^*,\; V(x)\mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x \leftarrow x';\gamma)\,U_2\psi\,\big]\Big\rangle = \Gamma_{\mathrm{corr}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma),$$

by cyclicity of the expectation and unitarity at both ends; hence its modulus—and therefore $d_{\mathrm{corr}}^{[\cdot]}$—is invariant (*wordwise* form of Proposition 4.410 for the correlator).

Vacuum (diagonal) distance. Writing the coincident self–correlator profile as a modal vector $\vec{C}^{[\mathbf{W}]}(x) = (C_n^{[\mathbf{W}]}(x))_n$, a word change acts by a unitary mixing on the modal frame, which descends to an orthogonal matrix $O(x)$ on components: $\vec{C}^{[\mathbf{W}']}(x) = O(x)\,\vec{C}^{[\mathbf{W}]}(x)$. Therefore the difference satisfies $\|\vec{C}^{[\mathbf{W}']}(x) - \vec{C}^{[\mathbf{W}']}(x')\|_{\ell^2} = \|O(x)\vec{C}^{[\mathbf{W}]}(x) - O(x')\vec{C}^{[\mathbf{W}]}(x')\|_{\ell^2}$. Transporting x' to x with the admissible wordwise parallel transport (§4.19.2) makes the two orthogonal actions coincide, so the ℓ^2–norm, and hence $d_{\mathrm{vac}}^{[\cdot]}$, is preserved (Proposition 4.469 in wordwise form).

Spectral distance. For each band n, $\mathcal{S}_n^{[\mathbf{W}]}(x,x') = \langle [\mathbb{D}^{[\mathbf{W}_1]}(x), \hat{G}_\Lambda(\cdot)]\rangle$ transforms by conjugation; its magnitude is invariant. The modal weights transform covariantly, and the weighted profile $\vec{\mathcal{S}}^{[\cdot]} = (w_n^{[\cdot]}|\mathcal{S}_n^{[\cdot]}|)_n$ rotates by an orthogonal matrix in band space (unitary change of modal basis). Hence the RMS norm $\|\vec{\mathcal{S}}^{[\cdot]}\|_{\ell^2} = d_{\mathrm{spec}}^{[\cdot]}$ is unchanged (Proposition 4.422 wordwise).

Step 2: invariance of the three $\vec{t}$ components.

Correlator phase time. By the endpoint rule above, $\Gamma_{\mathrm{corr}}^{[\mathbf{W}']}(x,x';\gamma)$ is multiplied only by unit–modulus factors that cancel between bra/ket or endpoints; thus $t_{\mathrm{corr}}^{[\mathbf{W}']}(x,x';\gamma) = \mathrm{Arg}\,\Gamma_{\mathrm{corr}}^{[\mathbf{W}']}(x,x';\gamma) \equiv \mathrm{Arg}\,\Gamma_{\mathrm{corr}}^{[\mathbf{W}]}(x,x';\gamma) = t_{\mathrm{corr}}^{[\mathbf{W}]}(x,x';\gamma) \pmod{2\pi}$ (Proposition 4.531).

Vacuum response time. $\mathrm{Im}\,\vec{V}_{\mathrm{vac}}^{[\mathbf{W}']}(x) = O(x)\,\mathrm{Im}\,\vec{V}_{\mathrm{vac}}^{[\mathbf{W}]}(x)$, so the ℓ^2–difference norm $t_{\mathrm{vac}}^{[\cdot]} = \|\,\mathrm{Im}\,\vec{V}_{\mathrm{vac}}^{[\mathbf{W}_1]}(x) - \mathrm{Im}\,\vec{V}_{\mathrm{vac}}^{[\mathbf{W}_2]}(x')\|_{\ell^2}$ is invariant (wordwise part of Proposition 4.540).

Spectral shear time. Similarly, $\operatorname{Im}\vec{S}^{[\mathbf{W}']}(x,x') = O\,\operatorname{Im}\vec{S}^{[\mathbf{W}]}(x,x')$ with the same orthogonal action on band space; hence $t^{[\cdot]}_{\mathrm{spec}} = \|\operatorname{Im}\vec{S}^{[\cdot]}\|_{\ell^2}$ is unchanged (wordwise part of Proposition 4.548).

Combining the six components yields $\mathscr{R}^{[\mathbf{W}']}(x,x') = \mathscr{R}^{[\mathbf{W}]}(x,x')$ for all $x,x' \in \mathcal{C}_d$. Therefore the zero sets coincide and the indistinguishability relations agree: $x \sim_{[\mathbf{W}]} x' \iff \mathscr{R}^{[\mathbf{W}]}(x,x') = \mathbf{0} \iff \mathscr{R}^{[\mathbf{W}']}(x,x') = \mathbf{0} \iff x \sim_{[\mathbf{W}']} x'$. Hence $\sim_{[\mathbf{W}']} = \sim_{[\mathbf{W}]}$.

Canonical isometry of the quotients. Define $\Phi : \mathcal{X}_q^{[\mathbf{W}]} \to \mathcal{X}_q^{[\mathbf{W}']}$ on classes by $\Phi\big([x]_{\sim_{[\mathbf{W}]}}\big) := [x]_{\sim_{[\mathbf{W}']}}$. This is well defined because the relations coincide, and it is bijective with inverse defined analogously. Moreover,

$$\overline{D}^{[\mathbf{W}']}\big(\Phi([x]),\Phi([x'])\big) = D^{[\mathbf{W}']}(x,x') = \|\vec{d}^{[\mathbf{W}']}(x,x')\|_2 = \|\vec{d}^{[\mathbf{W}]}(x,x')\|_2 = \overline{D}^{[\mathbf{W}]}\big([x],[x']\big),$$

so Φ is an isometry. Since Φ is induced by the identity on $\mathcal{C}_d$, it is canonical.

This proves the proposition. □

Topology, smooth structure, and curvature data

Definition 4.631 (Quotient topology and smooth structure)**.** Equip $\mathcal{X}_q^{[\mathbf{W}]}$ with the quotient topology of $\pi_q^{[\mathbf{W}]}$ and the metric $\overline{D}^{[\mathbf{W}]}$. Suppose the relation set

$$R^{[\mathbf{W}]} := \{(x,x') \in \mathcal{C}_d \times \mathcal{C}_d : \; x \sim_{[\mathbf{W}]} x'\}$$

is a closed embedded submanifold and $\pi_q^{[\mathbf{W}]}$ is a smooth submersion. Then by Theorem 4.627, $\mathcal{X}_q^{[\mathbf{W}]}$ is a smooth manifold and $\pi_q^{[\mathbf{W}]}$ is a smooth submersion. In the presence of a free proper action by a wordwise gauge group $G^{[\mathbf{W}]}$ preserving $\mathscr{R}^{[\mathbf{W}]}$, the quotient $\mathcal{C}_d/G^{[\mathbf{W}]}$ realises $\mathcal{X}_q^{[\mathbf{W}]}$ and $\pi_q^{[\mathbf{W}]}$ is a principal $G^{[\mathbf{W}]}$–bundle (Theorem 4.626).

Remark 4.632 (Emergent metric and tensors on the quotient)**.** The coincidence–limit construction yields the wordwise emergent metric $g_{\mu\nu}^{[\mathbf{W}]}$ from $D^{[\mathbf{W}]}$ (Section 4.11.3). Under the hypotheses of Definition 4.631, $g^{[\mathbf{W}]}$ and its operator–valued refinements (e.g. $\hat{g}_{\mu\nu}$, $\hat{t}_{\mu\nu}$, $\hat{\Sigma}_{\mu\nu}$) descend to smooth fields on $\mathcal{X}_q^{[\mathbf{W}]}$; expectation–level tensors define the chronogeometric sector on the quotient.

Remark 4.633 (Independence from the holosymmetric word and exogenous equivalence)**.** By Proposition 4.630, the diffeomorphism class and metric structure of $\mathcal{X}_q^{[\mathbf{W}]}$ are independent of $\mathbf{W}$. Moreover, via the wordwise–exogenous equivalence results (e.g. Proposition 4.340), the wordwise quotient coincides isometrically with the exogenous construction $\mathcal{X}_q = \mathcal{C}_d/\!\sim$, so we may drop the superscript when no ambiguity arises.

Definition 4.634 (Temporal flow on the quotient)**.** For $[x],[x'] \in \mathcal{X}_q^{[\mathbf{W}]}$, define the temporal triad as the class–invariant values $\vec{t}^{\,[\mathbf{W}]}\big([x],[x']\big) := \vec{t}^{[\mathbf{W}]}(x,x')$. The rate 1–forms along projected curves are induced from the connection of Section 4.19.2 by pushforward through $\pi_q^{[\mathbf{W}]}$; their holonomies detect nontrivial correlator/vacuum/spectral phase curvature on $\mathcal{X}_q^{[\mathbf{W}]}$.

Emergence of $\mathcal{X}_q^{[\mathbf{W}]}$ as a Wordwise Relational Quotient Space

Proposition 4.635 (Emergence of $\mathcal{X}_q^{[\mathbf{W}]}$ as a Relational Quotient)**.** *Fix an admissible holosymmetric word* $\mathbf{W}$*. Define the equivalence relation* $\sim_{[\mathbf{W}]}$ *on* $\mathcal{C}_d$ *by the* simultaneous vanishing *of the wordwise triads* $\vec{d}^{[\mathbf{W}]}$ *and* $\vec{t}^{[\mathbf{W}]}$ *(pulled back by the operational transporter of Section 4.19.2):*

$$\boxed{\vec{r} \sim_{[\mathbf{W}]} \vec{r}' \iff \vec{d}^{[\mathbf{W}]}(\vec{r},\vec{r}') = \mathbf{0} \text{ and } \vec{t}^{[\mathbf{W}]}(\vec{r},\vec{r}') = \mathbf{0}.} \tag{4.329}$$

The wordwise quantum configuration space *is the quotient*

$$\boxed{\mathcal{X}_q^{[\mathbf{W}]} := \mathcal{C}_d / \sim_{[\mathbf{W}]}.} \tag{4.330}$$

Each point $x \in \mathcal{X}_q^{[\mathbf{W}]}$ *corresponds to a wordwise PQS class* $[\mathfrak{B}_{\vec{r}}^{[\mathbf{W}]}]$ *of loci that are operationally indistinguishable.*

Proof. Fix an admissible holosymmetric word $\mathbf{W}$ and let $\mathcal{R}^{[\mathbf{W}]} : \mathcal{C}_d \times \mathcal{C}_d \to \mathbb{R}^6$ denote the *wordwise relational map*

$$\mathcal{R}^{[\mathbf{W}]}(\vec{r},\vec{r}') := \Big(\vec{d}^{[\mathbf{W}]}(\vec{r},\vec{r}'),\ \vec{t}^{[\mathbf{W}]}(\vec{r},\vec{r}') \Big),$$

where $\vec{d}^{[\mathbf{W}]}$ and $\vec{t}^{[\mathbf{W}]}$ are evaluated with the operational transporter of Section 4.19.2. By definition, $\vec{r} \sim_{[\mathbf{W}]} \vec{r}' \iff \mathcal{R}^{[\mathbf{W}]}(\vec{r},\vec{r}') = \mathbf{0}$.

Step 1: $\sim_{[\mathbf{W}]}$ is an equivalence whose classes are PQS leaves. Reflexivity follows from the normalisations at coincidence (see Definitions 4.339, 4.529, 4.539 and 4.547), which give $\mathcal{R}^{[\mathbf{W}]}(\vec{r},\vec{r}) = \mathbf{0}$. Symmetry of $\vec{d}^{[\mathbf{W}]}$ and the phase/antisymmetry properties of $\vec{t}^{[\mathbf{W}]}$ (modulo 2π, with vanishing value implying symmetry) yield $\mathcal{R}^{[\mathbf{W}]}(\vec{r},\vec{r}') = \mathbf{0} \Rightarrow \mathcal{R}^{[\mathbf{W}]}(\vec{r}',\vec{r}) = \mathbf{0}$. Transitivity follows from the triangle inequality for each component of $\vec{d}^{[\mathbf{W}]}$ and from additivity of the unwrapped phase rates for $\vec{t}^{[\mathbf{W}]}$ (Propositions 4.532, 4.540 and 4.548): if both pairs $(\vec{r},\vec{r}')$ and $(\vec{r}',\vec{r}'')$ are zero, then $(\vec{r},\vec{r}'')$ is zero. Hence $\sim_{[\mathbf{W}]}$ is an equivalence relation. By the wordwise relational null identification (Definition 4.582 and Corollary 4.609), its classes are precisely the wordwise PQS leaves $\mathfrak{B}_{\vec{r}}^{[\mathbf{W}]} = \{\vec{r}' : \vec{d}^{[\mathbf{W}]}(\vec{r},\vec{r}') = \vec{0} = \vec{t}^{[\mathbf{W}]}(\vec{r},\vec{r}')\}$.

Step 2: regularity of the zero set. Work in a local trivialisation of the wordwise bundle $\pi^{[\mathbf{W}]} : \mathfrak{F}^{[\mathbf{W}]} \to \mathcal{C}_d$ (eq. (4.306)) and along smooth transports (Section 4.19.2). Under the standing hypotheses (Wordwise Spectral Smoothness and Finite Entanglement Capacity; Theorem 4.421 and Principle 4.2) and diagonal regularity, the Fourfold expectations that define the six components are uniformly absolutely convergent modal series; by dominated differentiation, each component of $\mathcal{R}^{[\mathbf{W}]}$ depends smoothly on $(\vec{r},\vec{r}')$. Hence the relation set

$$R^{[\mathbf{W}]} := \{(\vec{r},\vec{r}') : \mathcal{R}^{[\mathbf{W}]}(\vec{r},\vec{r}') = \mathbf{0}\}$$

is closed. Moreover, by functional *orthogonality* and *independence* of the three axes (Sections 4.11.1 and 4.11.7), the differential $D\mathcal{R}^{[\mathbf{W}]}$ has locally constant rank $r \in \{0,\dots,6\}$ along $R^{[\mathbf{W}]}$. The constant–rank (preimage) theorem then gives that $R^{[\mathbf{W}]}$ is a closed embedded submanifold of $\mathcal{C}_d \times \mathcal{C}_d$; slicing at fixed $\vec{r}$ shows each leaf $\mathfrak{B}_{\vec{r}}^{[\mathbf{W}]}$ is a closed embedded submanifold of $\mathcal{C}_d$.

Step 3: quotient manifold and emergence of $\mathcal{X}_q^{[\mathbf{W}]}$. Because the leaves are closed embedded submanifolds and $R^{[\mathbf{W}]}$ is a closed embedded submanifold of $\mathcal{C}_d \times \mathcal{C}_d$, and because the canonical projection $\pi_q^{[\mathbf{W}]} : \mathcal{C}_d \to \mathcal{C}_d/\sim_{[\mathbf{W}]}$ is a smooth submersion in a neighbourhood of each leaf (local slices transverse to the foliation are provided by the implicit–function theorem applied to $\mathcal{R}^{[\mathbf{W}]}$), the quotient $\mathcal{X}_q^{[\mathbf{W}]} := \mathcal{C}_d/\sim_{[\mathbf{W}]}$ is a smooth manifold and $\pi_q^{[\mathbf{W}]}$ is a smooth submersion (Theorem 4.627). Each point $x \in \mathcal{X}_q^{[\mathbf{W}]}$ is by construction a class $[\mathfrak{B}_{\vec{r}}^{[\mathbf{W}]}]$ of loci that are operationally indistinguishable in the word frame $\mathbf{W}$.

Step 4: (optional) metric descent and submetry. Writing $D^{[\mathbf{W}]}(\vec{r},\vec{r}') = \|\vec{d}^{[\mathbf{W}]}(\vec{r},\vec{r}')\|_2$ (Section 4.11.2), the induced rule $\overline{D}^{[\mathbf{W}]}([\vec{r}],[\vec{r}']) = D^{[\mathbf{W}]}(\vec{r},\vec{r}')$ is well defined on the quotient and defines a metric; the projection $\pi_q^{[\mathbf{W}]} : (\mathcal{C}_d, D^{[\mathbf{W}]}) \to (\mathcal{X}_q^{[\mathbf{W}]}, \overline{D}^{[\mathbf{W}]})$ is 1–Lipschitz and a submetry (cf. Proposition 4.629).

All claims in the proposition follow: $\mathcal{X}_q^{[\mathbf{W}]}$ *emerges* as the relational quotient of $\mathcal{C}_d$ by the wordwise indistinguishability relation, and its points are the wordwise PQS classes. □

Remark 4.636 (Word–invariance and exogenous equivalence)**.** Admissible word changes $\mathbf{W} \mapsto \mathbf{W}'$ preserve the six wordwise components and thus $\sim_{[\mathbf{W}']} = \sim_{[\mathbf{W}]}$ (Proposition 4.630); therefore $\mathcal{X}_q^{[\mathbf{W}']} \cong \mathcal{X}_q^{[\mathbf{W}]}$ canonically. Via the wordwise–exogenous equivalence, this quotient agrees isometrically with the exogenous construction $\mathcal{X}_q = \mathcal{C}_d/\sim$ and we may drop $[\mathbf{W}]$ when no ambiguity arises.

Cardinality of the Configuration Space

Proposition 4.637 (Cardinality of $\mathcal{X}_q$)**.** *Let $\mathcal{C}_d$ be the smooth, connected, Hausdorff, second–countable d-manifold of continuum cardinality $\mathfrak{c}$ fixed in Section 4.18, and let $\sim$ be the operational indistinguishability relation of Definition 4.623. Assume the quotient $\pi_q : \mathcal{C}_d \to \mathcal{X}_q := \mathcal{C}_d/\sim$ satisfies the hypotheses of Theorem 4.622 (so that $\mathcal{X}_q$ is a smooth, Hausdorff, second–countable manifold). Then:*

$$\boxed{\left|\mathcal{X}_q\right| \;=\; \left|\mathcal{C}_d\right| \;=\; \mathfrak{c}.} \tag{4.331}$$

Proof. Surjectivity of π_q gives $|\mathcal{X}_q| \le |\mathcal{C}_d| = \mathfrak{c}$. It remains to show $|\mathcal{X}_q| \ge \mathfrak{c}$. Write $m := \dim \mathfrak{B}_x$ (leaf dimension), locally constant by Section 4.19.3.

Case 1 (positive transverse dimension). If $d - m \ge 1$, then by the submersion property there exists, through any $x \in \mathcal{C}_d$, a slice S_x transverse to $\mathfrak{B}_x$ such that $\pi_q|_{S_x} : S_x \to \pi_q(S_x)$ is a diffeomorphism onto an open neighbourhood in $\mathcal{X}_q$. Since S_x is a C^1 submanifold of dimension $d - m \ge 1$, it is (locally) homeomorphic to a nonempty open subset of $\mathbb{R}^{d-m}$, hence has cardinality $\mathfrak{c}$. Therefore $|\mathcal{X}_q| \ge |\pi_q(S_x)| = \mathfrak{c}$.

Case 2 (zero–dimensional leaves). If $m = 0$, each class $\mathfrak{B}_x$ is a discrete embedded submanifold. Ultra–sub–Planckian localisation eq. (4.303) confines $\mathfrak{B}_x$ to a small closed ball, hence, being a discrete subset of a compact set, $\mathfrak{B}_x$ is *finite*. Decompose $\mathcal{C}_d$ by leaf size:

$$\mathcal{C}_d \;=\; \bigcup_{k\in\mathbb{N}} \; \bigcup_{[y]\in\mathcal{X}_q:\ |\mathfrak{B}_y|=k} \mathfrak{B}_y.$$

Cardinal arithmetic for infinite sets then yields $\mathfrak{c} = |\mathcal{C}_d| \leq \sum_{k\in\mathbb{N}} k \cdot |\{[y] : |\mathfrak{B}_y| = k\}| \leq \aleph_0 \cdot |\mathcal{X}_q| = |\mathcal{X}_q|$. Thus $|\mathcal{X}_q| \geq \mathfrak{c}$.

In both cases we conclude $|\mathcal{X}_q| = \mathfrak{c}$. □

Remark 4.638 (What the assumptions ensure)**.** The conclusion (4.331) relies only on the quotient-manifold regularity from Section 4.19.3 and the spectral localisation eq. (4.303). Either there is positive transverse dimension $d - m \geq 1$, whence a single slice already carries cardinality $\mathfrak{c}$; or the leaves are zero–dimensional and, by localisation, finite, so that the quotient has $\mathfrak{c}$ many distinct classes. No "almost everywhere" injectivity is needed, and no new hypotheses beyond those already in force are introduced.

This result confirms that the emergent configuration space $\mathcal{X}_q$ retains the rich, continuous structure of the foundational continuum $\mathcal{C}_d$, providing a robust arena for the dynamics of the Gaiamiti and Tantrakyma fields.

Architectural Synthesis of the Emergent Quantum Configuration Space

Taken together, Sections 4.18, 4.19.1 and 4.19.3 fix a complete operational scaffold from which the configuration arena emerges:

- **Base (indexing continuum).** A smooth, connected, Hausdorff, second–countable d-manifold $\mathcal{C}_d \cong \mathbb{R}^d$ of cardinality $\mathfrak{c}$, serving only as an index set for operational data (Section 4.18).
- **Content (structured loci).** At each anchor $x \in \mathcal{C}_d$ sits the structured PQS content $\hat{\Omega}_x = {}_{\hat{\Phi}}\overset{\circ}{\Lambda}_{\hat{\Theta}}$, the universal duality code mediated by the Gaiamiti and Tantrakyma fields (eq. (4.301)).
- **Fibres (relational observables).** Relational distinguishability and temporality are carried by the Hilbert bundle $\pi : \mathfrak{F} \to \mathcal{C}_d$ with fibre $\mathfrak{F}_x = \mathcal{F}_x^{\text{dist}} \times \mathcal{F}_x^{\text{time}}$ (eq. (4.302)), coordinatised in the fourfold modal basis and restricted to the ultra–sub–Planckian spectral subspace $\operatorname{Ran} E^{(x)}([0, \lambda_{\text{cutoff}}])$ (eq. (4.303); see also [372, 248]).
- **Emergence (relational quotient).** Operational indistinguishability is encoded by the triads $\mathscr{R}(x, x') = (\vec{d}, \vec{t})$; declaring $x \sim x'$ iff $\mathscr{R}(x, x') = \mathbf{0}$ (Definition 4.623) yields the quotient $\mathcal{X}_q := \mathcal{C}_d/\!\sim$ with projection π_q (eq. (4.322)). Under the constant–rank, spectrally regulated regime already in force, $\mathcal{X}_q$ inherits a smooth, Hausdorff, second–countable structure and π_q is a smooth submersion (see Section 4.19.3). Cardinality is preserved: $|\mathcal{X}_q| = |\mathcal{C}_d| = \mathfrak{c}$ (Proposition 4.637).

Conceptually, the base labels *loci*, not classical points: each anchor x indexes a PQS $\mathfrak{B}_x$ and a regulated fibre $\mathfrak{F}_x$; comparisons across anchors are effected by the gauge–covariant transporter of Section 4.19.1. The smooth structure on $\mathcal{X}_q$ is therefore inherited from $\mathcal{C}_d$ *through* the operational equivalence defined by the triads, while the geometry carried by sections of $\mathfrak{F}$ (distances, phase times, holonomy) is propagated by the skew–Hermitian connection and its curvature. In this way, the emergent arena is intrinsically relational

and quantum: chronogeometric quantities are read from operator-valued correlators on spectrally controlled fibres rather than imposed kinematically, aligning the framework with noncommutative and relational geometric principles [4, 3].

$$\boxed{\begin{aligned}
\mathcal{C}_d &:= \text{smooth, Hausdorff, second–countable base}, \quad |\mathcal{C}_d| = \mathfrak{c},\\
\hat{\Omega}_{\vec{r}} &:= \hat{\Phi}\mathring{\Lambda}_{\hat{\Theta}}, \quad \vec{r} \in \mathcal{C}_d,\\
\mathfrak{F}_{\vec{r}} &:= \mathcal{F}^{\mathrm{dist}}_{\vec{r}} \times \mathcal{F}^{\mathrm{time}}_{\vec{r}}, \quad \operatorname{supp} \mathfrak{F}_{\vec{r}} \subset \mathcal{O}(\ell_{\mathrm{PQS}}),\\
\mathcal{X}_q &:= \mathcal{C}_d/\!\sim, \quad |\mathcal{X}_q| = \mathfrak{c}.
\end{aligned}}$$

Cardinality of the Wordwise Configuration Space

Proposition 4.639 (Cardinality of $\mathcal{X}_q^{[\mathbf{W}]}$)**.** *Let $\mathcal{C}_d$ be the smooth, connected, Hausdorff, second–countable d–manifold of continuum cardinality $\mathfrak{c}$ fixed in Definition 4.618. Let $\sim_{[\mathbf{W}]}$ be the wordwise operational–indistinguishability relation of Definition 4.628, and assume the quotient $\pi_q^{[\mathbf{W}]} : \mathcal{C}_d \to \mathcal{X}_q^{[\mathbf{W}]} := \mathcal{C}_d/\sim_{[\mathbf{W}]}$ satisfies the hypotheses of the wordwise quotient–manifold theorem (Theorem 4.627), so that $\mathcal{X}_q^{[\mathbf{W}]}$ is smooth, Hausdorff, and second–countable. Then*

$$\boxed{\left|\mathcal{X}_q^{[\mathbf{W}]}\right| \;=\; \left|\mathcal{C}_d\right| \;=\; \mathfrak{c}.} \tag{4.332}$$

Proof. Since $\pi_q^{[\mathbf{W}]} : \mathcal{C}_d \twoheadrightarrow \mathcal{X}_q^{[\mathbf{W}]}$ is surjective,

$$|\mathcal{X}_q^{[\mathbf{W}]}| \;\le\; |\mathcal{C}_d| \;=\; \mathfrak{c}.$$

It remains to show $|\mathcal{X}_q^{[\mathbf{W}]}| \ge \mathfrak{c}$.

Step 1: Local slice and positive transverse dimension. By Proposition 4.635, each equivalence class $\mathfrak{B}_x^{[\mathbf{W}]}$ is a closed, embedded submanifold ("relational leaf") of dimension $m := \dim \mathfrak{B}_x^{[\mathbf{W}]}$, locally constant. The quotient map $\pi_q^{[\mathbf{W}]}$ is a smooth submersion, so for any $x \in \mathcal{C}_d$ there exists a submanifold (slice) $S_x \subset \mathcal{C}_d$ of dimension $d-m$ through x such that $\pi_q^{[\mathbf{W}]}\restriction_{S_x} : S_x \to \pi_q^{[\mathbf{W}]}(S_x)$ is a diffeomorphism onto an open neighbourhood in $\mathcal{X}_q^{[\mathbf{W}]}$ (cf. Theorem 4.627). If $d-m \ge 1$ at some point, then S_x contains a nonempty open set in $\mathbb{R}^{d-m}$ and hence $|S_x| = \mathfrak{c}$. Therefore

$$|\mathcal{X}_q^{[\mathbf{W}]}| \;\ge\; |\pi_q^{[\mathbf{W}]}(S_x)| \;=\; |S_x| \;=\; \mathfrak{c}.$$

Combining with the upper bound gives $|\mathcal{X}_q^{[\mathbf{W}]}| = \mathfrak{c}$.

Step 2: Excluding the degenerate case $d-m=0$. It remains to rule out $d-m=0$ under our hypotheses. If $d-m=0$, then $m=d$ and each leaf $\mathfrak{B}_x^{[\mathbf{W}]}$ is an embedded d–submanifold of the d–manifold $\mathcal{C}_d$. Hence every leaf is *open* in $\mathcal{C}_d$. Because $\mathcal{C}_d$ is connected, the only partition of $\mathcal{C}_d$ into disjoint nonempty open leaves is the trivial one with a single leaf, i.e. $\mathcal{X}_q^{[\mathbf{W}]}$ would be a singleton.

However, ultra–sub–Planckian localisation of the fibres (eq. (4.307)) implies that each leaf is confined to an $\mathcal{O}(\ell_{\mathrm{PQS}})$ neighbourhood; in particular no leaf can be open in $\mathcal{C}_d$ unless it is

all of $\mathcal{C}_d$, which would contradict the nontriviality of the relational observables that define the quotient. Thus the degenerate case $d-m=0$ is excluded by the standing hypotheses, and there *must* exist points with $d-m\geq 1$. Step 1 then applies.

Optional reinforcement when $m=0$ (zero–dimensional leaves). If $m=0$ globally, each leaf is a discrete embedded submanifold. Choosing a chart with compact closure, discrete $\cap$ compact $\Rightarrow$ finite; hence there is $k\in\mathbb{N}$ such that

$$\mathcal{C}_d=\bigcup_{k\in\mathbb{N}}\bigcup_{[y]\in\mathcal{X}_q^{[\mathbf{W}]}:\ |\mathfrak{B}_y^{[\mathbf{W}]}|=k}\mathfrak{B}_y^{[\mathbf{W}]},\qquad\Rightarrow\qquad \mathfrak{c}=|\mathcal{C}_d|\leq\sum_{k\in\mathbb{N}}k\cdot|\{[y]:|\mathfrak{B}_y^{[\mathbf{W}]}|=k\}|\leq\aleph_0\cdot|\mathcal{X}_q^{[\mathbf{W}]}|.$$

Hence $|\mathcal{X}_q^{[\mathbf{W}]}|\geq\mathfrak{c}$, consistent with Step 1 (since now $d-m=d\geq 1$).

Combining the upper and lower bounds yields $|\mathcal{X}_q^{[\mathbf{W}]}|=\mathfrak{c}$, proving (4.332). □

Remark 4.640 (Role of assumptions (wordwise))**.** The conclusion (4.332) uses only: (i) the quotient–manifold regularity from Proposition 4.635 and Theorem 4.627, and (ii) spectral localisation of fibres $\mathfrak{F}_x^{[\mathbf{W}]}=\operatorname{Ran}E^{(x)[\mathbf{W}]}([0,\lambda_{\text{cutoff}}])$ (eq. (4.307)). Either there is positive transverse dimension $d-m\geq 1$ and a single slice has cardinality $\mathfrak{c}$, or the leaves are zero–dimensional and finite, forcing $\mathfrak{c}$ many distinct classes. No further hypotheses are introduced.

Architectural Synthesis of the Emergent Wordwise Configuration Space

Collecting Definition 4.618, Section 4.19.2, and Propositions 4.635 and 4.639:

- **Base (indexing continuum).** $\mathcal{C}_d\cong\mathbb{R}^d$ (smooth, connected, Hausdorff, second–countable, $|\mathcal{C}_d|=\mathfrak{c}$) serves as an *index* for operational data in the wordwise calculus.
- **Content (wordwise PQS loci).** Each anchor $x\in\mathcal{C}_d$ carries the wordwise PQS content $\hat{\Omega}_x^{[\mathbf{W}]}=\mathring{\Lambda}_{\hat{\Theta}}^{\hat{\Phi}}$ and its relational–null leaf $\mathfrak{B}_x^{[\mathbf{W}]}$ (cf. Definition 4.618).
- **Fibres (relational observables).** The holosymmetric relational bundle $\pi^{[\mathbf{W}]}:\mathfrak{F}^{[\mathbf{W}]}\to\mathcal{C}_d$ with $\mathfrak{F}_x^{[\mathbf{W}]}=\mathcal{F}_x^{\text{dist}[\mathbf{W}]}\times\mathcal{F}_x^{\text{time}[\mathbf{W}]}$ is coordinatised in the fourfold dialectic basis and restricted to the ultra–sub–Planckian sector $\operatorname{Ran}E^{(x)[\mathbf{W}]}([0,\lambda_{\text{cutoff}}])$ (eqs. (4.306) and (4.307)).
- **Transport (operational connection).** Gauge–covariant parallel transport $U_\gamma^{[\mathbf{W}]}$ preserves Fourfold expectations and generates the metric–compatible connection (Section 4.19.2).
- **Emergence (relational quotient).** Wordwise operational indistinguishability declares $x\sim_{[\mathbf{W}]}x'$ iff the six wordwise components vanish; the quotient $\mathcal{X}_q^{[\mathbf{W}]}:=\mathcal{C}_d/\sim_{[\mathbf{W}]}$ is a smooth, Hausdorff, second–countable manifold with $\pi_q^{[\mathbf{W}]}$ a smooth submersion (Proposition 4.635), and *cardinality is preserved*: $|\mathcal{X}_q^{[\mathbf{W}]}|=\mathfrak{c}$ (Proposition 4.639).

Indra's Net and the Relational Fabric of the Continuum

The fibred continuum $\mathcal{C}_d$ (Section 4.18) reframes "space'' and "time'' as phenomena *emergent from operator–valued relations* rather than as pre-given backdrops. Each Primordial Quantum Singularity $\mathfrak{B}_{\vec{r}}$ is a locus whose identity is fixed entirely by its web of operator correlations with every other locus, encoded fibrewise by the relational bundle $\pi : \mathfrak{F} \to \mathcal{C}_d$ (eq. (4.302)) and compared via the gauge–covariant transport of Section 4.19.1. The quotient $\mathcal{X}_q = \mathcal{C}_d/\sim$ (eq. (4.322)) is thereby *read off* from this web: two anchors are the "same place'' exactly when all triadic observables vanish between them, so that the fabric of the continuum is nothing but the network of correlations itself, spectrally constrained at ultra–sub–Planckian width (eq. (4.303)). In this sense our formalism finds a poetic and ancient resonance in the metaphor of **Indra's Net**, which captures with striking clarity the foundational idea that nothing exists independently and that structure arises through reflection, connection, and coherence across the whole:

> "Far away in the heavenly abode of the great god Indra, there is a wonderful net which has been hung by some cunning artificer in such a manner that it stretches out infinitely in all directions. In accordance with the extravagant tastes of deities, the artificer has hung a single glittering jewel in each 'eye' of the net, and since the net itself is infinite in dimension, the jewels are infinite in number. There hang the jewels, glittering like stars of the first magnitude, a wonderful sight to behold. If we now arbitrarily select one of these jewels for inspection and look closely at it, we will discover that in its polished surface there are reflected all the other jewels in the net, infinite in number. Not only that, but each of the jewels reflected in this one jewel is also reflecting all the other jewels, so that there is an infinite reflecting process occurring. This symbolizes our world, where every sentient being (and every phenomenon) is a jewel in Indra's Net, and each jewel reflects all the others, and so on ad infinitum. Each jewel represents an individual life form, atom, cell or unit of consciousness. Each is distinctly different, yet all are intimately connected, and all reflect each other."
>
> — Francis H. Cook, *Hua-yen Buddhism: The Jewel Net of Indra* (1977)

The ancient metaphor of Indra's Net, wherein each jewel in a cosmic web reflects every other, is granted operational substance through the formalism of Holosymmetry. In this construction, the Net is realised as a fibred Pre-Quantum Spacetime (PQS) continuum, with the 'jewels' being precisely the PQS loci $\mathfrak{B}_x$ situated over a base space of relational anchors $\mathcal{C}_d$ (Section 4.18). The phenomenon of 'reflection' is not merely allegorical; it corresponds to the quantitative assessment of relational observables. What each jewel $\mathfrak{B}_x$ reflects of any other $\mathfrak{B}_{x'}$ is the complete ledger of chronogeometric information carried by the fibre $\mathfrak{F}_{x'} = \mathcal{F}^{\text{dist}}_{x'} \times \mathcal{F}^{\text{time}}_{x'}$, evaluated via Fourfold correlators after being brought into a common frame at x (eq. (4.302) and Section 4.19.1).

This process of reflection is enacted by the gauge-covariant parallel transport operator,

$$\mathcal{U}(x \leftarrow x'; \gamma) = \mathcal{P} \exp\left(-\int_\gamma \mathcal{A}\right),$$

which transports the state-dependent data from the fibre at x' to the fibre at x along a specified path γ within the base manifold (eqs. (4.11) and (4.16)). The visual attributes of the reflection are given precise, measurable correlates. The *brightness* of the jewel-to-jewel reflection is quantified by the magnitude of the resulting correlators—specifically, the normalised overlap that underlies the correlation distance d_{corr}. Concurrently, its *hue* or *phase* is recorded by the entanglement phase time t_{corr} (eq. (4.313)). The entire network of mutual reflections is thus rendered as a tangible operator network, constrained to an ultra-sub-Planckian resolution by the spectral projection defining the PQS loci (eq. (4.303)).

The geometry of this relational weave is probed through interferometry. A holosymmetric interferometer compares two distinct reflective itineraries, γ_1 and γ_2, between a pair of jewels. The observable contrast in such an experiment measures the holonomy operator for the composed loop, $\mathcal{H}[\gamma_1, \gamma_2]$ (eq. (4.314)). The expectation value of this operator in the global Quantum Holo-State, $\langle \overset{\circ}{\boldsymbol{\Psi}} | \mathcal{H} | \overset{\circ}{\boldsymbol{\Psi}} \rangle$, directly yields the interferometric signature: its modulus defines the reflection's brightness, and its argument defines its phase. For small loops, a non-trivial holonomy certifies the presence of curvature in the connection $\mathcal{A}$. This curvature represents a failure of global 'mirror consistency' across the net, thereby serving as a direct probe of the emergent chronogeometry (Section 4.19.1).

From this relational tapestry, the very arena of physics—the configuration space $\mathcal{X}_q$—emerges not as an externally imposed backdrop, but as the coherent *pattern* of the net itself. Two anchor points, $x, x' \in \mathcal{C}_d$, are deemed to occupy the same location in the emergent space if and only if their complete reflection profiles are operationally indistinguishable. This condition is formalised by the vanishing of the triadic relational observables $\mathscr{R}(x, x') = (\vec{d}, \vec{t}) = \mathbf{0}$ after canonical comparison (Definition 4.623 and eq. (4.322)). Consequently, $\mathcal{X}_q$ is the quotient space of the base manifold under this indistinguishability equivalence, inheriting a smooth structure directly from the regularity of the relational map rather than from any prior coordinate system (Section 4.19.3).

The entire construction is underpinned by two fundamental principles: gauge covariance and global coherence. Because the triadic observables and the holosymmetric readouts are gauge-invariant, the physical content of the net is independent of the choice of local frames within each fibre; a gauge transformation merely rotates the 'facets' of a jewel without altering what is reflected (eq. (4.16)). The coherence of the entire net—the 'light' by which all jewels mutually illuminate one another—is supplied by the single, non-factorisable Quantum Holo-State $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$. It is this global state that binds all reflections into one indivisible, entangled fabric, ensuring that the emergent pattern $\mathcal{X}_q$ is an observer-independent reality.

Chapter 5

The Metric and Temporal Tensors

5.1 The Metric Tensor

5.1.1 Operator-Valued Emergent Metric Tensor from Composite Distance

Definition 5.1 (Operator-Valued Emergent Metric Tensor from Composite Distance)**.** Let $x, x' \in \mathcal{X}_q$ and let $\hat{d}_{\mathrm{corr}}(x,x'), \hat{d}_{\mathrm{vac}}(x,x'), \hat{d}_{\mathrm{spec}}(x,x') \in \mathrm{Herm}^{\geq 0}(\mathcal{H}_{\mathrm{total}})$ be the operator-valued analogues of the triadic relational distances from Definition 4.476, with components initially introduced in eqs. (4.149), (4.193) and (4.210) and realised by localised modal correlators with gauge–covariant comparison between the PQS loci anchored at x and x' (Section 4.19.1). Define the operatorial composite distance $D^2(x,x')$ by replacing each scalar d_a in Definition 4.476 with its operator counterpart $\hat{d}_a$.

The *operator-valued emergent metric tensor* is the Fréchet second variation of D^2 at coincidence:

$$\boxed{\hat{g}_{\mu\nu}(x) \;:=\; \frac{\partial^2}{\partial x^\mu\, \partial x'^\nu}\, D^2(x,x')\Big|_{x'=x}\,.} \tag{5.1}$$

Topology/regularity clause. Unless stated otherwise, the derivatives in (5.1) are taken in the **strong–operator topology (SOT)**, uniformly on the fixed spectral band selected by the fibrewise cutoff $\mathrm{Ran}\, E^{(x)}([0,\lambda_{\mathrm{cutoff}}])$ (eq. (4.303)); by spectral localisation and the smooth dependence of the transporter (Section 4.19.1), D^2 is SOT–C^2 near coincidence and hence $\hat{g}_{\mu\nu}(x) \in \mathrm{Herm}^{\geq 0}(\mathcal{H}_{\mathrm{total}})$.

Here $\hat{g}_{\mu\nu}$ is a symmetric $(0,2)$-type operator field on $\mathcal{X}_q$ encoding the infinitesimal quantum relational geometry prior to taking expectations. Its classical counterpart is $\bar{g}_{\mu\nu}(x) := \langle \overset{\circ}{\boldsymbol{\Psi}} \,|\, \hat{g}_{\mu\nu}(x) \,|\, \overset{\circ}{\boldsymbol{\Psi}} \rangle$ (Definition 5.10).

Properties of the Operator-Valued Emergent Metric Tensor

The operator-valued emergent metric tensor, $\hat{g}_{\mu\nu}(x)$, occupies a central place in the relational geometry of the Primordial Quantum Singularity (PQS) continuum. Unlike classical metric tensors, which are real-valued symmetric bilinear forms on the tangent bundle of a manifold, $\hat{g}_{\mu\nu}(x)$ is a *Hermitian, positive semi-definite operator-valued field* acting on the total Hilbert

space $\mathcal{H}_{\text{total}}$. It reflects the infinitesimal relational structure of the emergent quantum configuration space $\mathcal{X}_q$, not by assigning numerical distances, but by encoding *operator-level correlational amplitudes* between points in a non-classical continuum.

Definition from Triadic Operator Distances

We assemble the composite triadic operator–valued distance from three positive, Hermitian, fibrewise observables, constructed (and compared across anchors) by the gauge–covariant machinery of Section 4.19.1:

$$D^2(x,x') \;:=\; \hat{d}^2_{\text{corr}}(x,x') + \hat{d}^2_{\text{vac}}(x,x') + \hat{d}^2_{\text{spec}}(x,x') \;\in\; \text{Herm}^{\geq 0}(\mathcal{H}_{\text{total}}). \tag{5.2}$$

Here each $\hat{d}_a(x,x') \in \text{Herm}^{\geq 0}(\mathcal{H}_{\text{total}})$ $(a \in \{\text{corr}, \text{vac}, \text{spec}\})$ captures a distinct, fibrewise relational deviation between the PQS loci anchored at x and x':

- *Correlational deviation* via the Fourfold correlator $\hat{G}_\Lambda$ (pulling the x'-datum to x along $\mathcal{U}(x \leftarrow x'; \gamma)$; cf. eq. (4.11));
- *Vacuum displacement* via the diagonal contrast in the regulated fibre (as in the scalar definition, lifted operatorially);
- *Spectral shear* via the spectral generator $\hat{D}$ from Definition 4.343, realised through commutator–type increments on the bandlimited fibre.

Orthogonality of the three channels is understood *at the operational level* with respect to the GNS/Fourfold inner product induced by the Holo–State on the regulated operator algebra (cf. Section 1.2.9), so that the cross terms vanish in expectation and the sum (5.2) faithfully encodes independent modes of deviation.

The operator–valued metric tensor is the second Fréchet derivative of D^2 at coincidence:

$$\hat{g}_{\mu\nu}(x) \;:=\; \frac{\partial^2}{\partial x^\mu\, \partial x'^\nu}\, D^2(x,x')\bigg|_{x'=x}. \tag{5.3}$$

Topology/regularity clause. Derivatives in (5.3) are taken in the strong–operator topology (SOT), *uniformly on the fixed spectral band* $\text{Ran}\, E^{(x)}([0,\lambda_{\text{cutoff}}])$ (eq. (4.303)). Under the smooth dependence of the transporter and the triadic observables (Section 4.19.1), D^2 is SOT–C^2 near coincidence, hence $\hat{g}_{\mu\nu}(x) \in \text{Herm}^{\geq 0}(\mathcal{H}_{\text{total}})$. When the maps $x \mapsto \hat{d}_a(x,\cdot)$ are norm–differentiable on the band, the SOT derivatives agree with the operator–norm derivatives.

Target Space and Positivity

For each $x \in \mathcal{X}_q$,

$$\hat{g}_{\mu\nu}(x) \;\in\; \text{Herm}^{\geq 0}(\mathcal{H}_{\text{total}}). \tag{5.4}$$

Positivity follows because every channel $\hat{d}_a$ is positive and Hermitian, hence $\hat{d}_a^2 \geq 0$, and the SOT limit in (5.3) preserves Hermiticity/positivity on the bandlimited operator algebra. Consequently, expectation–level inner products and lengths (e.g. $\bar{g}_{\mu\nu}(x) = \langle \overset{\circ}{\boldsymbol{\Psi}} | \hat{g}_{\mu\nu}(x) | \overset{\circ}{\boldsymbol{\Psi}} \rangle$) are non–negative, making the operator metric a physically viable seed for distances, angles, and geodesic deviation within the noncommutative, fibrewise setting (Definition 5.10).

Localisation: Ultra–Sub–Planckian Support

The metric is *ultra–local* in the precise spectral/relational sense fixed earlier. Writing $P_x := E^{(x)}([0, \lambda_{\text{cutoff}}])$,

$$\operatorname{Ran} \hat{g}_{\mu\nu}(x) \;\subseteq\; P_x\, \mathcal{H}_{\text{total}}, \qquad \hat{g}_{\mu\nu}(x) \;=\; P_x\, \hat{g}_{\mu\nu}(x)\, P_x, \tag{5.5}$$

so that all action is confined to the $\mathcal{O}(\ell_{\text{PQS}})$ coherence neighbourhood encoded by the fibre projector (eq. (4.303)). In particular, $\hat{g}_{\mu\nu}$ is not placed *on top of* a classical background; rather, the classical manifold $\mathcal{X}_q$ is the quotient arena emergent from—and coordinatised by—the very relational data that $\hat{g}_{\mu\nu}$ (and its temporal analogue) encodes (Section 4.19.3).

State–dependence and expectation geometry

While $\hat{g}_{\mu\nu}(x)$ is defined state–independently (as an operator field obtained from the SOT–C^2 second variation in eq. (5.3)), its *operational* reading arises upon Born evaluation in the Holo–State. Writing $\rho_\circ := |\overset{\circ}{\mathbf{\Psi}}\rangle\langle\overset{\circ}{\mathbf{\Psi}}|$,

$$\bar{g}_{\mu\nu}(x) \;:=\; \operatorname{Tr}\Big(\rho_\circ\, \hat{g}_{\mu\nu}(x)\Big) \;=\; \langle \overset{\circ}{\mathbf{\Psi}} |\, \hat{g}_{\mu\nu}(x)\, |\, \overset{\circ}{\mathbf{\Psi}} \rangle, \tag{5.6}$$

which is the *expectation–level emergent metric* governing effective geometry for coarse, classical probes (Definition 5.10 and Section 1.2.9). Thus $\hat{g}_{\mu\nu}$ plays a dual role: ontologically, it encodes nonlocal operator correlations among PQS fibres; epistemically, its Born valuation $\bar{g}_{\mu\nu}$ supplies a Riemannian (later Lorentzian–deformed) tensor suitable for dynamics, propagation, and curvature on $\mathcal{X}_q$.

Symmetry and gauge covariance

The triadic operators $\hat{d}_a(x, x')$ are built from gauge–covariant data (Fourfold correlators and the transporter $\mathcal{U}$) and thus transform by conjugation under local frame changes $g : \mathcal{X}_q \to \mathrm{U}(\mathcal{H}_{\text{em}})$, in the convention of eq. (4.16):

$$\hat{g}_{\mu\nu}(x) \mapsto g(x)\, \hat{g}_{\mu\nu}(x)\, g(x)^\dagger \tag{5.7}$$

Consequently, expectation scalars are gauge–invariant: $\bar{g}_{\mu\nu}(x) = \operatorname{Tr}(\rho_\circ \hat{g}_{\mu\nu}(x)) = \operatorname{Tr}(g \rho_\circ g^\dagger\, g \hat{g}_{\mu\nu} g^\dagger)$ by cyclicity of the trace and the covariant transformation of $\rho_\circ$ (or, equivalently, by unitary invariance of the Born pairing). Hence the emergent geometric predictions are independent of internal frame choices (Section 4.19.1).

Non-Riemannian structure and internal triplet geometry

Because the composite distance $D^2 = \sum_a \hat{d}_a^2$ (eq. (5.2)) splits into three orthogonal channels (with respect to the Fourfold inner product on the regulated algebra, Section 1.2.9), the metric decomposes canonically as

$$\hat{g}_{\mu\nu}(x) \;=\; \hat{g}^{\text{corr}}_{\mu\nu}(x) \;+\; \hat{g}^{\text{vac}}_{\mu\nu}(x) \;+\; \hat{g}^{\text{spec}}_{\mu\nu}(x), \tag{5.8}$$

each term being the SOT–second variation of the corresponding $\hat{d}_a^2$. At the operator level the three sectors need not commute, so $\hat{g}_{\mu\nu}$ encodes a genuinely *noncommutative, triadic* geometry. At the expectation level the orthogonality of channels suppresses cross terms, yielding the classical tensor $\bar{g}_{\mu\nu} = \sum_a \bar{g}^a_{\mu\nu}$ on $\mathcal{X}_q$. In this sense, the interface between ontological (entangled, operator–valued) structure and epistemic (expectation–valued) geometry is made explicit: the former lives on the spectrally localised fibres $P_x\mathcal{H}_{\text{total}}$ (eq. (4.303)), while the latter is a smooth field on the quotient arena $\mathcal{X}_q$ (Section 4.19.3).

Positivity and locality. Since each $\hat{d}_a^2 \geq 0$ on the bandlimited algebra, $\hat{g}_{\mu\nu}(x) \in \text{Herm}^{\geq 0}(\mathcal{H}_{\text{total}})$, and $\bar{g}_{\mu\nu}(x)$ is positive semidefinite. Moreover, $\hat{g}_{\mu\nu}(x) = P_x\,\hat{g}_{\mu\nu}(x)\,P_x$ with $P_x = E^{(x)}([0, \lambda_{\text{cutoff}}])$, so its action is ultra-local in the sense fixed by $\ell_{\text{PQS}} \ll \ell_{\text{P}}$ (eq. (4.303)). Thus the classical manifold structure is *not* a backdrop for $\hat{g}_{\mu\nu}$; rather, it is *emergent from* the relational data that $\hat{g}_{\mu\nu}$ (and its temporal analogue) encode.

5.1.2 Physical Interpretation of the Operator–Valued Metric Tensor

The operator–valued emergent metric tensor $\hat{g}_{\mu\nu}(x)$ is the interface between the foundational operator algebra on the regulated fibres and the emergent geometry on $\mathcal{X}_q$. Unlike a classical metric, $\hat{g}_{\mu\nu}(x)$ is a field of Hermitian, positive–semidefinite operators supported on the spectral band $P_x\mathcal{H}_{\text{total}}$ with $P_x := E^{(x)}([0, \lambda_{\text{cutoff}}])$ (eq. (4.303)). It encodes the *infinitesimal relational structure* of the continuum in the strong–operator sense fixed by eq. (5.3).

Definition and triadic structure. Recalling $D^2 = \hat{d}^2_{\text{corr}} + \hat{d}^2_{\text{vac}} + \hat{d}^2_{\text{spec}}$ (eq. (5.2)), the metric is the SOT–C^2 second variation at coincidence, $\hat{g}_{\mu\nu}(x) = \partial_{x^\mu}\partial_{x'^\nu} D^2(x, x')\big|_{x'=x}$ (eq. (5.3)). This immediately yields the canonical decomposition

$$\hat{g}_{\mu\nu}(x) \;=\; \hat{g}^{\text{corr}}_{\mu\nu}(x) \;+\; \hat{g}^{\text{vac}}_{\mu\nu}(x) \;+\; \hat{g}^{\text{spec}}_{\mu\nu}(x), \tag{5.9}$$

each term the second variation of the corresponding channel and, in general, mutually noncommuting. Indices $\mu, \nu \in \{1, \ldots, d\}$ are coordinates on $\mathcal{X}_q$ with $d = \dim \mathcal{X}_q$.

Operational (expectation–level) geometry. While $\hat{g}_{\mu\nu}$ is state–independent, operational predictions arise by Born evaluation in the Holo–State $\rho_\circ = |\mathring{\mathbf{\Psi}}\rangle\langle\mathring{\mathbf{\Psi}}|$:

$$\bar{g}_{\mu\nu}(x) \;:=\; \text{Tr}\Big(\rho_\circ\,\hat{g}_{\mu\nu}(x)\Big), \qquad \bar{g}_{\mu\nu} = \bar{g}^{\text{corr}}_{\mu\nu} + \bar{g}^{\text{vac}}_{\mu\nu} + \bar{g}^{\text{spec}}_{\mu\nu} \quad (\textit{eqs. (5.6) and (5.8)}). \tag{5.10}$$

For any tangent vector $v = v^\mu \partial_\mu|_x$, the *state–dependent quadratic form*

$$\|v\|^2_{\rho_\circ} \;:=\; v^\mu v^\nu\,\bar{g}_{\mu\nu}(x) \;=\; \text{Tr}\Big(\rho_\circ\, v^\mu v^\nu\,\hat{g}_{\mu\nu}(x)\Big) \;\geq 0$$

is the effective length seen by coarse observers; thus $\bar{g}$ governs propagation, curvature, and dynamics on $\mathcal{X}_q$ (Definition 5.10 and Section 1.2.9). Ontologically, the full operator $\hat{g}_{\mu\nu}$ retains fine relational content beyond any single expectation value.

Component	Triadic mode	Operational meaning (second variation at coincidence)
$\hat{g}^{(a)}_{11}(x)$	Correlational	Pure x^1–displacement curvature of the correlator channel: stiffness of entanglement and rate of coherence loss along x^1.
	Vacuum	Pure x^1–axis curvature of vacuum self–coherence (diagonal contrast) on the regulated fibre.
	Spectral	Accumulation of spectral shear under pure x^1 displacements (response generated by $\hat{D}$).
$\hat{g}^{(a)}_{22}(x)$	Correlational	Entanglement–stiffness/decay for pure x^2 displacements.
	Vacuum	Vacuum curvature along the x^2 axis.
	Spectral	Spectral–shear accumulation for pure x^2 displacements.
$\hat{g}^{(a)}_{33}(x)$	Correlational	Entanglement–stiffness/decay for pure x^3 displacements.
	Vacuum	Vacuum curvature along the x^3 axis.
	Spectral	Spectral–shear accumulation for pure x^3 displacements.
$\hat{g}^{(a)}_{12}(x) = \hat{g}^{(a)}_{21}(x)$	Correlational	Mixed x^1–x^2 sensitivity of the correlator amplitude: cross–coupling of entanglement between the two directions.
	Vacuum	Vacuum anisotropy under combined (x^1, x^2) displacements.
	Spectral	Mixed spectral–response curvature for simultaneous x^1 and x^2 variations (phase–gradient coupling set by $\hat{D}$).
$\hat{g}^{(a)}_{13}(x) = \hat{g}^{(a)}_{31}(x)$	Correlational	Entanglement cross–coupling between x^1 and x^3.
	Vacuum	Vacuum anisotropy under combined (x^1, x^3) displacements.
	Spectral	Mixed spectral–response curvature for joint x^1 and x^3 variations.
$\hat{g}^{(a)}_{23}(x) = \hat{g}^{(a)}_{32}(x)$	Correlational	Entanglement cross–coupling between x^2 and x^3.
	Vacuum	Vacuum anisotropy under combined (x^2, x^3) displacements.
	Spectral	Mixed spectral–response curvature for joint x^2 and x^3 variations.

Table 5.1: Operator–valued metric components in a local 3D chart. Each entry is the SOT–second variation of the corresponding channel in D^2 (eqs. (5.2) and (5.3)); "spectral" refers to the response generated by the spectral operator $\hat{D}$ (Definition 4.343).

Gauge covariance and invariance of predictions. Since the triads are built from gauge–covariant objects (correlators and the transporter $\mathcal{U}$), $\hat{g}_{\mu\nu}$ transforms by conjugation under a local frame $g : \mathcal{X}_q \to \mathrm{U}(\mathcal{H}_{\mathrm{em}})$ (eq. (4.16)):

$$\hat{g}_{\mu\nu}(x) \;\longmapsto\; g(x)\,\hat{g}_{\mu\nu}(x)\,g(x)^{\dagger}, \qquad P_x \;\longmapsto\; g(x)\,P_x\,g(x)^{\dagger}.$$

Expectation–level scalars such as $\bar{g}_{\mu\nu}$ and $\|v\|^2_{\rho_\circ}$ are invariant by cyclicity of the trace and the unitary covariance of $\rho_\circ$; hence emergent geometric predictions are independent of internal frame choices (Section 4.19.1).

Three-layer reading. Every component carries: (i) a *directional layer* $\mu\nu$ specifying where in $\mathcal{X}_q$ one probes the infinitesimal relational curvature; (ii) a *modal layer* $a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$ specifying what type of relational deviation is interrogated; and (iii) an *operator layer* indicating that the outcome is a positive Hermitian operator on $P_x\mathcal{H}_{\mathrm{total}}$, ready for evaluation against states. This triadic, noncommutative structure makes explicit how ontological entanglement data on the fibres give rise, via (5.10), to an emergent (and later Lorentzian–deformed) classical geometry on $\mathcal{X}_q$ (Sections 4.19.3 and 5.1).

5.1.3 Operator-Valued Emergent Metric Tensor (Holosymmetric Wordwise)

Definition 5.2 (Wordwise operator–valued composite distance)**.** Fix admissible word frames $(\mathbf{W}_1, \mathbf{W}_2)$ at the anchors $x, x' \in \mathcal{X}_q$ (cf. holosymmetric transport in Section 4.19.2). Let

$$\hat{d}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{corr}}(x,x'), \quad \hat{d}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{vac}}(x,x'), \quad \hat{d}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{spec}}(x,x') \;\in\; \mathrm{Herm}^{\geq 0}\big(\mathcal{H}_{\mathrm{total}}\big)$$

be the wordwise *operator* analogues of the three distances, realised by the localised wordwise modal correlators and the wordwise spectral generator $\mathbb{D}^{[\mathbf{W}_1]}$, with gauge–covariant comparison between x and x' (cf. Definitions 4.407 and 4.408 and Section 4.19.2).

Let

$$\hat{\mathbf{d}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \;:=\; \Big(\hat{d}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{corr}}(x,x'),\; \hat{d}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{vac}}(x,x'),\; \hat{d}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{spec}}(x,x')\Big)^{\top}.$$

Choose a fibrewise symmetric positive–definite calibration $\mathbf{M}(x) \in \mathrm{Sym}^{+}(3)$, write $\mathbf{M}(x) = \mathbf{B}(x)^{\top}\mathbf{B}(x)$, and set the calibrated operator components

$$\hat{\mathbf{z}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \;:=\; \mathbf{B}(x)\,\hat{\mathbf{d}}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \qquad \text{(componentwise Hermitian operators).}$$

Define the *wordwise operator–valued composite distance* by

$$\boxed{\hat{D}^{2,[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \;:=\; \sum_{j=1}^{3}\Big(\hat{z}_j^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\Big)^2.} \tag{5.11}$$

Equivalently,

$$\hat{D}^{2,[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \;=\; \tfrac{1}{2}\sum_{a,b=1}^{3} M_{ab}(x)\,\Big\{\hat{d}_a^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x'),\, \hat{d}_b^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\Big\},$$

via the Jordan product $\{A, B\} := AB + BA$; both forms are Hermitian and agree in expectation, and coincide when the components commute.

Definition 5.3 (Wordwise operator–valued emergent metric tensor)**.** Let $E^{(x)[\mathbf{W}_1]}([0, \lambda_{\text{cutoff}}])$ be the fibrewise spectral projector selecting the ultra–sub–Planckian band (cf. eq. (4.307)). On the localised band

$$\mathfrak{H}_x \;:=\; \operatorname{Ran} E^{(x)[\mathbf{W}_1]}([0, \lambda_{\text{cutoff}}]),$$

the mixed Fréchet second variation of $\hat{D}^{2,[\mathbf{W}_1,\mathbf{W}_2]}(x, x')$ in the strong–operator topology exists and defines the *wordwise operator–valued emergent metric*:

$$\boxed{\hat{g}_{\mu\nu}^{[\mathbf{W}_1,\mathbf{W}_2]}(x) \;:=\; \frac{\partial^2\, \hat{D}^{2,[\mathbf{W}_1,\mathbf{W}_2]}(x, x')}{\partial x^\mu\, \partial x'^\nu}\bigg|_{x'=x} \;\in\; \operatorname{Herm}^{\geq 0}\big(\mathfrak{H}_x\big).} \tag{5.12}$$

Unless stated otherwise, derivatives are taken in the strong–operator topology uniformly on $\mathfrak{H}_x$. Smoothness follows from spectral localisation, wordwise Spectral Smoothness, and smooth gauge–covariant transport (cf. Axiom 4.418 and Section 4.19.2).

Remark 5.4 (Scalarization and orthogonal decomposition)**.** The expectation–valued (classical) metric is

$$\bar{g}_{\mu\nu}^{[\mathbf{W}_1,\mathbf{W}_2]}(x) \;:=\; \Big\langle \overset{\circ}{\mathbf{\Psi}} \,\Big|\, \hat{g}_{\mu\nu}^{[\mathbf{W}_1,\mathbf{W}_2]}(x) \,\Big|\, \overset{\circ}{\mathbf{\Psi}} \Big\rangle.$$

With diagonal calibration $\mathbf{M}(x) = \operatorname{diag}(\alpha^2, \beta^2, \gamma^2)$ and the fibre decomposition of the triad, cross–terms vanish in expectation, so $\bar{g}_{\mu\nu}$ decomposes additively into correlator, vacuum, and spectral pieces (cf. Sections 4.11.2 and 4.11.4).

Proposition 5.5 (Well–posedness, positivity, and regularity)**.** *Under the wordwise Spectral Smoothness and Finite Entanglement Capacity hypotheses (Axiom 4.418 and Principle 4.2), the map $(x, x') \mapsto \hat{D}^{2,[\mathbf{W}_1,\mathbf{W}_2]}(x, x')$ is* SOT*–C^2 near coincidence on $\mathfrak{H}_x$, and $\hat{g}_{\mu\nu}^{[\mathbf{W}_1,\mathbf{W}_2]}(x)$ is a positive semidefinite Hermitian operator.* Sketch. *Each wordwise operator distance is an* SOT*–continuous Holo–State norm of a localised, gauge–transported probe; spectral localisation plus the commutator/weight bounds control second differences. Positivity follows from (5.11) as a sum of squares of Hermitian operators.*

Proposition 5.6 (Frame/word covariance and unitary equivalence)**.** *If $(\mathbf{W}_1, \mathbf{W}_2) \mapsto (\mathbf{W}'_1, \mathbf{W}'_2)$ is an admissible word change with fibre unitaries $(U_x, U_{x'})$ (cf. Proposition 4.397), then*

$$\hat{d}_a^{[\mathbf{W}'_1,\mathbf{W}'_2]}(x, x') = U_x^\dagger\, \hat{d}_a^{[\mathbf{W}_1,\mathbf{W}_2]}(x, x')\, U_x, \qquad \hat{D}^{2,[\mathbf{W}'_1,\mathbf{W}'_2]}(x, x') = U_x^\dagger\, \hat{D}^{2,[\mathbf{W}_1,\mathbf{W}_2]}(x, x')\, U_x,$$

and hence

$$\boxed{\hat{g}_{\mu\nu}^{[\mathbf{W}'_1,\mathbf{W}'_2]}(x) \;=\; U_x^\dagger\, \hat{g}_{\mu\nu}^{[\mathbf{W}_1,\mathbf{W}_2]}(x)\, U_x.} \tag{5.13}$$

Therefore the scalarised metric $\bar{g}_{\mu\nu}(x)$ is word/frame invariant.

Remark 5.7 (Ordering and Jordan symmetrisation)**.** Because the components $\hat{d}_a^{[\mathbf{W}_1,\mathbf{W}_2]}$ need not commute, the quadratic form may be defined either as a sum of calibrated squares (preferred for positivity) or via the Jordan product $\{\cdot, \cdot\}$ to maintain Hermiticity for a general $\mathbf{M}$. Both give the same Holo–State expectation and agree whenever the commutators vanish to leading order.

Remark 5.8 (Band–limitation and domain)**.** All derivatives and operator products are taken on the localised band $\mathfrak{H}_x = \operatorname{Ran} E^{(x)[\mathbf{W}_1]}([0, \lambda_{\text{cutoff}}])$ (cf. eq. (4.307)). This ensures domain stability, SOT–C^2 regularity, and compatibility with coherent regularisation.

Remark 5.9 (Coordinate–free form)**.** Using the operational connection ∇ along charts (cf. Section 4.19.2), one may write

$$\hat{g}_{\mu\nu}^{[\mathbf{W}_1,\mathbf{W}_2]}(x) \;=\; \tfrac{1}{2}\Big(\nabla_\mu \nabla'_\nu \, \hat{D}^{2,[\mathbf{W}_1,\mathbf{W}_2]}(x,x')\Big)\Big|_{x'=x},$$

with the primed derivative acting on the fibre at x' after parallel transport to x.

Outcome. Equations (5.11)–(5.12) furnish the holosymmetric wordwise operator–valued metric tensor. It is positive, SOT–regular, band–limited, and unitarily equivalent across word frames; its expectation reproduces the calibrated, additive sum of correlator, vacuum, and spectral contributions established at the scalar level (cf. Sections 4.11.2 and 4.11.4).

5.1.4 Expectation-Level Emergent Metric Tensor

Definition 5.10 (Expectation-Level Emergent Metric Tensor $\bar{g}_{\mu\nu}(x)$)**.** Let $\mathfrak{F}_x = \mathcal{F}_x^{\text{dist}} \times \mathcal{F}_x^{\text{time}}$ be the relational fibre at $x \in \mathcal{X}_q$ (Definitions 4.599 and 4.600). Write $\vec{d}(x,x') \in \mathcal{F}_x^{\text{dist}}$ for the triadic distinguishability vector and define

$$D^2(x,x') \;:=\; \|\vec{d}(x,x')\|_x^2,$$

where $\|\cdot\|_x$ is the Euclidean norm induced by the fibre inner product constructed from Fourfold expectations (Section 4.6).

Assume that, in a local chart about x, the map $(x,x') \mapsto D^2(x,x')$ is C^2 on a neighbourhood of the diagonal and admits a quadratic expansion; equivalently, the operator-level composite distance underlying D^2 has well-defined Fréchet second variations taken in the *strong-operator topology*, uniformly on the spectral band fixed by the projector Π_Λ (eq. (4.303)). Then the *expectation-level emergent metric* is the symmetric tensor field

$$\boxed{D^2(x,x') \;=\; \bar{g}_{\mu\nu}(x)\,\Delta x^\mu \Delta x^\nu \;+\; \mathcal{O}(|\Delta x|^3), \qquad \Delta x^\mu := (x'-x)^\mu,} \tag{5.14}$$

with components given by the coincidence-limit second derivatives

$$\bar{g}_{\mu\nu}(x) \;=\; \frac{\partial^2}{\partial x^\mu \, \partial x'^\nu} D^2(x,x')\Big|_{x'=x}.$$

If, moreover, the operator-valued metric tensor $\hat{g}_{\mu\nu}(x)$ is defined by the Fréchet second variation of the operatorial composite distance (Definition 5.1), then the two formulations agree via the universal state:

$$\boxed{\bar{g}_{\mu\nu}(x) \;:=\; \Big\langle \overset{\circ}{\mathbf{\Psi}} \Big| \, \hat{g}_{\mu\nu}(x) \, \Big| \overset{\circ}{\mathbf{\Psi}} \Big\rangle.} \tag{5.15}$$

Topology & regularity (one-line reminder). All Fréchet derivatives of D^2 and T^2 are taken in the strong-operator topology on the spectrally regulated fibres (uniformly on the band selected by Π_Λ), ensuring that $\hat{g}_{\mu\nu}(x), \hat{t}_{\mu\nu}(x) \in \mathrm{Herm}(\mathcal{H}_{\mathrm{total}})$ and that (5.15) is well-defined.

Two Complementary Readings

Operational. Equation (5.14) states that, for infinitesimal separations, the squared composite distinguishability is a quadratic form whose coefficients are the measurable components of the effective Riemannian geometry. Probing $D(x, x')$ for nearby x' reconstructs $\bar{g}_{\mu\nu}(x)$.

Ontological. Equation (5.15) ties the same coefficients to the expectation of a gauge-covariant operator field $\hat{g}_{\mu\nu}(x)$, making explicit that geometry is state-induced by the Quantum Holo-State $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$.

Gauge Covariance and Invariance of Predictions

Because the triadic ingredients ($\hat{G}_\Lambda$, $\hat{D}$, and the transporter $\mathcal{U}$) are gauge-covariant, the operator metric transforms as

$$\hat{g}_{\mu\nu}(x) \;\mapsto\; U(x)\,\hat{g}_{\mu\nu}(x)\,U(x)^\dagger,$$

for any local frame $U(x) \in \mathrm{Aut}(\mathcal{H}_{\mathrm{total}})$. Consequently, $\bar{g}_{\mu\nu}(x)$ is invariant, and all emergent geometric predictions are frame-independent within the internal bundle.

Hierarchy of Emergence (compressed)

- *Level 0 (Foundational):* Fourfold operator algebra and universal state $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$.
- *Level 1 (Quantum geometry):* $\hat{g}_{\mu\nu}(x)$ from Fréchet variation of the operatorial composite distance.
- *Level 2 (Classical geometry):* $\bar{g}_{\mu\nu}(x) = \langle \hat{g}_{\mu\nu}(x) \rangle$ governing macroscopic dynamics.

Consequence

Once $\bar{g}_{\mu\nu}(x)$ is fixed by either (5.14) or (5.15), the standard Riemannian data (Christoffel symbols, curvature) follow in the usual way on $\mathcal{X}_q$, providing the bridge to macrophysical equations of motion.

5.1.5 Expectation-Level Emergent Metric Tensor (Holosymmetric Wordwise)

Definition 5.11 (Wordwise expectation-level emergent metric tensor)**.** Fix admissible word frames $(\mathbf{W}_1, \mathbf{W}_2)$ at anchors $x, x' \in \mathcal{X}_q$ and use the gauge–covariant transporter along admissible paths (cf. Section 4.19.2). Define the wordwise triadic distinguishability vector (with the x'–fibre transported to x)

$$\vec{d}^{[\mathbf{W}_1,\mathbf{W}_2]}(x, x') := \Big(d^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{corr}}(x, x'),\; d^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{vac}}(x, x'),\; d^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mathrm{spec}}(x, x') \Big) \;\in\; \mathcal{F}^{\mathrm{dist}}_x.$$

Equip $\mathcal{F}_x^{\mathrm{dist}}$ with the calibrated fibre inner product

$$\langle \vec{u}, \vec{v} \rangle_{x,[\mathbf{W}_1]} \;:=\; \vec{u}^{\top} \mathbf{M}(x)\, \vec{v}, \qquad \mathbf{M}(x) \in \mathrm{Sym}^{+}(3).$$

The *wordwise composite distance* is

$$D^{2,[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \;:=\; \left\| \vec{d}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \right\|^2_{x,[\mathbf{W}_1]} \;=\; \vec{d}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x')^{\top} \mathbf{M}(x)\, \vec{d}^{[\mathbf{W}_1,\mathbf{W}_2]}(x,x').$$

Assume that in a local chart around x, the map $(x,x') \mapsto D^{2,[\mathbf{W}_1,\mathbf{W}_2]}(x,x')$ is C^2; this is ensured by the band-limited operator-level regularity (cf. eq. (4.307) and Axiom 4.418 and Principle 4.2). Then the *wordwise expectation-level emergent metric* is determined by the coincidence expansion

$$\boxed{D^{2,[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \;=\; \bar{g}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mu\nu}(x)\, \Delta x^{\mu}\, \Delta x^{\nu} \;+\; \mathcal{O}\Big(|\Delta x|^3\Big), \qquad \Delta x^{\mu} := (x'-x)^{\mu},} \tag{5.16}$$

with components

$$\boxed{\bar{g}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mu\nu}(x) \;:=\; \frac{\partial^2}{\partial x^{\mu}\, \partial x'^{\nu}}\, D^{2,[\mathbf{W}_1,\mathbf{W}_2]}(x,x') \Big|_{x'=x}.} \tag{5.17}$$

Remark 5.12 (Equivalence with the operator-valued form)**.** If $\hat{g}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mu\nu}(x)$ is defined via the band-limited SOT Fréchet Hessian of the operatorial composite distance (cf. eq. (5.12)), then

$$\boxed{\bar{g}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mu\nu}(x) \;=\; \Big\langle \mathring{\mathbf{\Psi}} \,\Big|\, \hat{g}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mu\nu}(x) \,\Big|\, \mathring{\mathbf{\Psi}} \Big\rangle,} \tag{5.18}$$

i.e. the expectation-level metric is the Holo-State scalarisation of the operator-valued metric on the localised band.

Proposition 5.13 (Positivity, orthogonal decomposition, and regularity)**.** *Under wordwise Spectral Smoothness and Finite Entanglement Capacity (ensuring band-limited C^2 regularity), $\bar{g}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mu\nu}(x)$ is positive semidefinite and smooth. With diagonal calibration $\mathbf{M} = \mathrm{diag}(\alpha^2,\beta^2,\gamma^2)$ and functional orthogonality of the triad, $\bar{g}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mu\nu}$ decomposes additively into correlator, vacuum, and spectral contributions; cross-terms vanish at expectation level (cf. Sections 4.11.2 and 4.11.4).*

Proposition 5.14 (Word/frame covariance and invariance)**.** *For admissible word changes $(\mathbf{W}_1,\mathbf{W}_2) \mapsto (\mathbf{W}'_1,\mathbf{W}'_2)$ with fibre unitaries, the six wordwise components of $\vec{d}$ are invariant (cf. Propositions 4.397 and 4.422), hence*

$$D^{2,[\mathbf{W}'_1,\mathbf{W}'_2]}(x,x') = D^{2,[\mathbf{W}_1,\mathbf{W}_2]}(x,x'), \qquad \bar{g}^{[\mathbf{W}'_1,\mathbf{W}'_2]}_{\mu\nu}(x) = \bar{g}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mu\nu}(x).$$

Thus the expectation-level metric is independent of the word frame and gauge choices.

Remark 5.15 (Calibration and physical dimensions)**.** The calibration $\mathbf{M}(x)$ implements axis rescalings or controlled couplings between the triadic components and fixes physical dimensions/units. Different choices of $\mathbf{M}$ correspond to admissible re-weightings of the three orthogonal contributions; $\bar{g}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mu\nu}$ transforms covariantly under such recalibrations.

Remark 5.16 (Consistency with the exogenous (non-wordwise) presentation)**.** By word/frame invariance of the distances and their operator representatives, the metric components computed from (5.17) agree with the non-wordwise construction; the wordwise formalism is an equivalent, gauge-transparent packaging rather than a different geometry (cf. Propositions 4.613 and 4.630).

Summary. Equations (5.16)–(5.17) define the holosymmetric wordwise expectation-level emergent metric tensor. It is well-posed, positive, smooth, orthogonally decomposed across the triad, and invariant under admissible word/frame changes, and it coincides with the scalarisation of the operator-valued metric (5.12).

5.2 The Temporal Tensor

5.2.1 Operator–Valued Temporal Curvature Tensor

Definition 5.17 (Operator–Valued Temporal Curvature Tensor)**.** Let $x, x' \in \mathcal{X}_q$ and let

$$\hat{t}_{\mathrm{corr}}(x,x'), \quad \hat{t}_{\mathrm{vac}}(x,x'), \quad \hat{t}_{\mathrm{spec}}(x,x')$$

be the operator–valued analogues of the triadic temporal observables from Definition 4.600, realised by gauge–covariant comparison of the x'–fibre with the x–fibre (via Section 4.19.1). Define the composite temporal displacement

$$T^2(x,x') \;:=\; \hat{t}^{\,2}_{\mathrm{corr}}(x,x') + \hat{t}^{\,2}_{\mathrm{vac}}(x,x') + \hat{t}^{\,2}_{\mathrm{spec}}(x,x') \;\in\; \mathrm{Herm}^{\geq 0}(\mathcal{H}_{\mathrm{total}}). \tag{5.19}$$

The *operator–valued temporal curvature tensor* is the Fréchet second variation of T^2 at coincidence:

$$\boxed{\hat{t}_{\mu\nu}(x) \;:=\; \frac{\partial^2}{\partial x^\mu\, \partial x'^\nu}\, T^2(x,x')\Big|_{x'=x} \;\in\; \mathrm{Herm}(\mathcal{H}_{\mathrm{total}}).} \tag{5.20}$$

Topology/regularity clause. The derivatives in (5.20) are taken in the strong–operator topology (SOT), uniformly on the spectrally regulated fibres $P_x\mathcal{H}_{\mathrm{total}}$ with $P_x := E^{(x)}([0,\lambda_{\mathrm{cutoff}}])$ (eq. (4.303)). Under the smooth dependence of the transporter and the temporal observables (Section 4.19.1 and eq. (4.313)), T^2 is SOT–C^2 near the diagonal, hence $\hat{t}_{\mu\nu}(x)$ is well–defined and Hermitian.

Triadic decomposition and localisation. By (5.19) one has the canonical split

$$\hat{t}_{\mu\nu}(x) \;=\; \hat{t}^{\mathrm{corr}}_{\mu\nu}(x) \;+\; \hat{t}^{\mathrm{vac}}_{\mu\nu}(x) \;+\; \hat{t}^{\mathrm{spec}}_{\mu\nu}(x), \tag{5.21}$$

each term the SOT–second variation of the corresponding channel. Localisation follows from spectral support:

$$\hat{t}_{\mu\nu}(x) \;=\; P_x\, \hat{t}_{\mu\nu}(x)\, P_x, \qquad \mathrm{Ran}\, \hat{t}_{\mu\nu}(x) \subseteq P_x\mathcal{H}_{\mathrm{total}},$$

so the tensor acts ultra–sub–Planckianly in the same sense as $\hat{g}_{\mu\nu}$ (eq. (4.303)).

Gauge covariance and expectation–level temporal tensor. For any local frame $g : \mathcal{X}_q \to \mathrm{U}(\mathcal{H}_{\mathrm{em}})$ (cf. eq. (4.16)),

$$\hat{t}_{\mu\nu}(x) \;\longmapsto\; g(x)\,\hat{t}_{\mu\nu}(x)\,g(x)^{\dagger}, \qquad P_x \;\longmapsto\; g(x)\,P_x\,g(x)^{\dagger},$$

so the Born expectation in the Holo–State $\rho_\circ = |\mathring{\mathbf{\Psi}}\rangle\langle\mathring{\mathbf{\Psi}}|$ is gauge–invariant:

$$\bar{t}_{\mu\nu}(x) \;:=\; \mathrm{Tr}\Big(\rho_\circ\,\hat{t}_{\mu\nu}(x)\Big). \tag{5.22}$$

The symmetric part $\bar{t}^{\mathrm{sym}}_{\mu\nu}$ provides the temporal deformation that enters the effective Lorentzian structure via $\bar{g}^{\mathrm{Lor}}_{\mu\nu} := \bar{g}_{\mu\nu} + \bar{t}^{\mathrm{sym}}_{\mu\nu}$ (Section 5.1), while the antisymmetric part captures phase–flow vorticity.

Physical reading. Whereas $\hat{g}_{\mu\nu}$ measures curvature of *distinguishability*, $\hat{t}_{\mu\nu}$ measures curvature of *quantum phase flow* in the temporal fibre $\mathcal{F}^{\mathrm{time}}_x$: it quantifies the directional sensitivity of the triadic time displacements to coordinate separation, providing the temporal counterpart that is absent in classical Riemannian geometry but natural in the relational operator framework. In small–loop limits, the temporal sector coheres with interferometric holonomy diagnostics (Section 4.19.1), giving a direct operational handle on the chrono–geometric content of phase dynamics.

Symmetry of Construction

The definition of $\hat{t}_{\mu\nu}(x)$ is deliberately parallel to that of the operator–valued metric tensor $\hat{g}_{\mu\nu}(x)$ (Definition 5.1):

- $\hat{g}_{\mu\nu}(x)$ is the Fréchet second variation (at coincidence) of the squared norm of the *distinguishability vector* $\vec{d}$.
- $\hat{t}_{\mu\nu}(x)$ is the Fréchet second variation (at coincidence) of the squared norm of the *temporal–flow vector* $\vec{t}$.

Both variations are taken on spectrally regulated fibres in the *strong–operator topology*, uniformly on the fixed spectral band; the resulting components land in $\mathrm{Herm}(\mathcal{H}_{\mathrm{total}})$ by construction. This symmetry makes explicit that the spatial and temporal sectors of the relational fibre $\mathfrak{F}_x = \mathcal{F}^{\mathrm{dist}}_x \times \mathcal{F}^{\mathrm{time}}_x$ admit parallel curvature diagnostics.

Physical Interpretation

Whereas $\hat{g}_{\mu\nu}(x)$ captures second–order sensitivity of *distinguishability* to coordinate displacements, $\hat{t}_{\mu\nu}(x)$ captures the second–order sensitivity of *phase flow*. Nonzero components indicate that infinitesimal phase accumulation between nearby loci depends on direction in $\mathcal{X}_q$. Operationally:

- *Directional phase anisotropy:* $\hat{t}_{\mu\mu}(x)$ quantifies the curvature of entanglement–phase (or vacuum/spectral) time under pure x^μ shifts, i.e. phase–flow "stiffness'' along x^μ.

- *Phase shear (mixed response):* $\hat{t}_{\mu\nu}(x)$ with $\mu \neq \nu$ measures cross–couplings of phase displacement under mixed (x^μ, x^ν) shifts—how phase flow "leans'' when the displacement is not axis–aligned.
- *Local temporal texture:* The sign/structure of $\hat{t}_{\mu\nu}(x)$ (generally indefinite) encodes a pointwise "temporal texture'' of the continuum not visible to the metric sector alone.

Gauge covariance mirrors the metric case: for any local frame $U(x)$,

$$\hat{t}_{\mu\nu}(x) \;\mapsto\; U(x)\,\hat{t}_{\mu\nu}(x)\,U(x)^\dagger,$$

so physical predictions are extracted from gauge–invariant functionals (e.g. state expectations, traces on regulated sectors).

Completing the Geometry of the Relational Bundle

Together, $\hat{g}_{\mu\nu}(x)$ and $\hat{t}_{\mu\nu}(x)$ furnish a complete, operator–level, infinitesimal description of the relational bundle:

$$\mathfrak{F}_x = \underbrace{\mathcal{F}_x^{\text{dist}}}_{\text{metric curvature via } \hat{g}} \times \underbrace{\mathcal{F}_x^{\text{time}}}_{\text{phase–flow curvature via } \hat{t}}.$$

Both tensors are ultra–local (supported on $\mathcal{O}(\ell_{\text{PQS}})$), gauge–covariant, and constant along relational leaves, hence descend unambiguously to the quotient $\pi_q : \mathcal{C}_d \to \mathcal{X}_q$. In compact form, the unified chronogeometric operator is

$$\hat{\Sigma}_{\mu\nu}(x) \;:=\; \hat{g}_{\mu\nu}(x) + \hat{t}_{\mu\nu}(x),$$

whose expectation in the Quantum Holo–State produces the classical chronogeometric tensor.

Expectation and Classical Field Content

The operational (expectation–level) temporal tensor is

$$\bar{t}_{\mu\nu}(x) \;:=\; \Big\langle \mathring{\mathbf{\Psi}} \,\Big|\, \hat{t}_{\mu\nu}(x) \,\Big|\, \mathring{\mathbf{\Psi}} \Big\rangle,$$

entering the macroscopic chronogeometry via $\bar{\Sigma}_{\mu\nu}(x) = \bar{g}_{\mu\nu}(x) + \bar{t}_{\mu\nu}(x)$. Here $\bar{t}_{\mu\nu}$ serves as the classical imprint of phase–flow curvature—complementary to $\bar{g}_{\mu\nu}$—governing causal/temporal structure in the emergent regime.

Future Directions

The temporal tensor unlocks several precise lines of development:

1. *Coupled field equations.* Derive the Euler–Lagrange equations for $\bar{g}_{\mu\nu}$ and $\bar{t}_{\mu\nu}$ from the relational action, and identify coupling terms (e.g. $g^{-1}\cdot t$ contractions) that mix spatial and temporal curvature.

2. *Bianchi–type identities.* Establish differential identities obeyed by $\hat{t}_{\mu\nu}$ and $\hat{g}_{\mu\nu}$ (and their expectations), clarifying integrability and constraint propagation.

3. *Invariants and scalars.* Construct gauge–invariant scalars such as $\mathrm{Tr}_\Lambda\big[\rho\,\hat{t}_{\mu\nu}\hat{t}^{\mu\nu}\big]$, $\mathrm{Tr}_\Lambda\big[\rho\,\hat{g}_{\mu\nu}\hat{t}^{\mu\nu}\big]$ on the regulated band for use in effective actions and diagnostics.

4. *Mixed tensors.* Use $\bar{g}^{\mu\alpha}$ to form $\bar{t}^{\mu}{}_{\nu} := \bar{g}^{\mu\alpha}\bar{t}_{\alpha\nu}$ ("temporal strain''), and study its eigenstructure as a local indicator of preferred phase–flow directions (arrows) in $\mathcal{X}_q$.

5. *Interferometric probes.* Translate components of $\bar{t}_{\mu\nu}$ into holosymmetric interferometric observables (visibility/phase slopes for axis–specific loops), tightening the operational bridge to $\mathcal{U}(x \leftarrow x';\gamma)$.

In sum, $\hat{t}_{\mu\nu}(x)$ is the temporal counterpart to $\hat{g}_{\mu\nu}(x)$: a symmetric, Hermitian, gauge–covariant second variation that quantifies curvature of quantum phase flow. Together they provide the full operator–valued chronogeometric data from which classical spacetime and its causal texture emerge by expectation.

5.2.2 Physical Interpretation of the Operator–Valued Temporal Curvature Tensor Components

The operator–valued temporal curvature tensor $\hat{t}_{\mu\nu}(x)$ complements the spatial metric tensor $\hat{g}_{\mu\nu}(x)$ as its temporal dual. While $\hat{g}_{\mu\nu}(x)$ encodes relational *spatial* structure, $\hat{t}_{\mu\nu}(x)$ captures the *infinitesimal curvature of quantum phase flow* in the temporal fibre $\mathcal{F}_x^{\mathrm{time}}$. Together they complete the infinitesimal operator–geometric description of the relational bundle $\mathfrak{F}_x$.

Definition (recall). By Definition 5.17,

$$\hat{t}_{\mu\nu}(x) \;=\; \frac{\partial^2}{\partial x^\mu\,\partial x'^\nu}\,T^2(x,x')\Big|_{x'=x}, \qquad T^2(x,x') \;=\; \hat{t}^2_{\mathrm{corr}}(x,x') + \hat{t}^2_{\mathrm{vac}}(x,x') + \hat{t}^2_{\mathrm{spec}}(x,x'),$$

where the triadic temporal observables are the operator analogues of Definition 4.600. This yields the canonical split (eq. (5.21))

$$\hat{t}_{\mu\nu}(x) \;=\; \hat{t}^{\mathrm{corr}}_{\mu\nu}(x) \;+\; \hat{t}^{\mathrm{vac}}_{\mu\nu}(x) \;+\; \hat{t}^{\mathrm{spec}}_{\mu\nu}(x),$$

each term encoding second–order variation of relational phase flow under displacements along x^μ and x^ν.

General Rule for Interpretation

- *Diagonal components* ($\mu=\nu$): $\hat{t}^{(a)}_{\mu\mu}(x)$ measures the curvature of phase displacement under a pure x^μ shift in channel $a \in \{\mathrm{corr},\mathrm{vac},\mathrm{spec}\}$—the local "stiffness'' (acceleration) of phase flow along that axis.

- *Off–diagonal components* ($\mu\neq\nu$): $\hat{t}^{(a)}_{\mu\nu}(x)$ diagnoses anisotropy and cross–coupling of phase flow—how phase evolution along x^μ depends on displacement in x^ν (mixed response).

Three Layers of Meaning

Each component $\hat{t}_{\mu\nu}(x)$ carries:

1. *Directional layer ($\mu\nu$):* Sensitivity of phase flow to displacements along the coordinate directions x^μ, x^ν on $\mathcal{X}_q$.

2. *Modal layer (triadic):* The physical origin of the flow—entanglement (corr), vacuum phase shift (vac), or spectral deformation (spec).

3. *Operator layer:* A Hermitian operator on $\mathcal{H}_{\text{total}}$, i.e. a quantum observable whose expectation in the Holo–State produces the classical temporal tensor $\bar{t}_{\mu\nu}(x)$.

Summary. The tensor $\hat{t}_{\mu\nu}(x)$ provides an operator–valued account of how *phase evolution itself curves* across directions in configuration space. This extends geometry beyond the Riemannian metric sector: phase curvature is encoded not in lengths but in the *directional asymmetry of phase flow*. In this sense, $\hat{t}_{\mu\nu}(x)$ is the temporal counterpart to $\hat{g}_{\mu\nu}(x)$, and together they complete the operator–valued relational chronogeometry of the bundle $\mathfrak{F}$.

5.2.3 The Temporal Tensor (Holosymmetric Wordwise)

Definition 5.18 (Wordwise operator–valued temporal curvature tensor)**.** Fix admissible word frames $(\mathbf{W}_1, \mathbf{W}_2)$ at $x, x' \in \mathcal{X}_q$ and use the gauge–covariant transporter $\mathcal{U}^{[\mathbf{W}_1,\mathbf{W}_2]}(x \leftarrow x'; \gamma)$ along admissible paths (cf. Section 4.19.2). Let

$$\hat{t}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\text{corr}}(x, x'; \gamma), \qquad \hat{t}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\text{vac}}(x, x'; \gamma), \qquad \hat{t}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\text{spec}}(x, x'; \gamma)$$

be the wordwise operator–valued analogues of the temporal observables (corr/vac/spec) obtained by comparing the x'–fibre with the x–fibre via $\mathcal{U}$ (cf. Definition 4.607 and Section 4.19.2). Assume these channel operators are Hermitian on the band–limited fibre. Define the composite temporal displacement operator

$$\boxed{\begin{aligned} T^{2,[\mathbf{W}_1,\mathbf{W}_2]}(x, x'; \gamma) \;:=\; &\Big(\hat{t}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\text{corr}}(x, x'; \gamma)\Big)^2 + \Big(\hat{t}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\text{vac}}(x, x'; \gamma)\Big)^2 \\ &+ \Big(\hat{t}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\text{spec}}(x, x'; \gamma)\Big)^2 \;\in\; \text{Herm}^{\geq 0}\Big(\mathcal{H}_{\text{total}}\Big). \end{aligned}} \tag{5.23}$$

The *wordwise operator–valued temporal curvature tensor* is the SOT–Fréchet second variation at coincidence:

$$\boxed{\hat{t}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mu\nu}(x) \;:=\; \frac{\partial^2}{\partial x^\mu\, \partial x'^\nu}\, T^{2,[\mathbf{W}_1,\mathbf{W}_2]}(x, x'; \gamma)\Big|_{x'=x} \;\in\; \text{Herm}\Big(\mathfrak{H}_x\Big),} \tag{5.24}$$

where $\mathfrak{H}_x := \text{Ran}\, E^{(x)[\mathbf{W}_1]}([0, \lambda_{\text{cutoff}}])$ (cf. eq. (4.307)). Derivatives are taken in the strong–operator topology, uniformly on $\mathfrak{H}_x$. Smooth dependence of $\mathcal{U}$ and the channel observables implies $T^{2,[\mathbf{W}_1,\mathbf{W}_2]} \in \text{SOT}\text{–}C^2$ near the diagonal, so $\hat{t}^{[\mathbf{W}_1,\mathbf{W}_2]}_{\mu\nu}(x)$ is well defined and Hermitian.

Component	Triadic mode	Physical meaning (condensed)
$\hat{t}^{(a)}_{11}(x)$	Correlational	Phase–flow curvature of entanglement coherence under pure x^1 displacement.
	Vacuum	Vacuum–phase curvature along the x^1 axis (diagonal temporal response).
	Spectral	Spectral phase–shear accumulation for pure x^1 displacements.
$\hat{t}^{(a)}_{22}(x)$	Correlational	Phase–flow curvature of correlations under pure x^2 displacement.
	Vacuum	Vacuum–phase curvature along the x^2 axis.
	Spectral	Spectral phase–shear accumulation for pure x^2 displacements.
$\hat{t}^{(a)}_{33}(x)$	Correlational	Phase–flow curvature of correlations under pure x^3 displacement.
	Vacuum	Vacuum–phase curvature along the x^3 axis.
	Spectral	Spectral phase–shear accumulation for pure x^3 displacements.
$\hat{t}^{(a)}_{12}(x) = \hat{t}^{(a)}_{21}(x)$	Correlational	Mixed phase–flow cross–coupling for combined (x^1, x^2) displacements.
	Vacuum	Mixed vacuum–phase response across (x^1, x^2).
	Spectral	Mixed spectral phase–shear across (x^1, x^2).
$\hat{t}^{(a)}_{13}(x) = \hat{t}^{(a)}_{31}(x)$	Correlational	Mixed phase–flow cross–coupling for combined (x^1, x^3) displacements.
	Vacuum	Mixed vacuum–phase response across (x^1, x^3).
	Spectral	Mixed spectral phase–shear across (x^1, x^3).
$\hat{t}^{(a)}_{23}(x) = \hat{t}^{(a)}_{32}(x)$	Correlational	Mixed phase–flow cross–coupling for combined (x^2, x^3) displacements.
	Vacuum	Mixed vacuum–phase response across (x^2, x^3).
	Spectral	Mixed spectral phase–shear across (x^2, x^3).

Table 5.2: Operator–valued temporal curvature components $\hat{t}^{(a)}_{\mu\nu}(x)$ in a local 3D chart. Each entry is the SOT second variation (at coincidence) of the corresponding temporal channel in T^2 (Definition 5.17); expectation in the Holo–State yields the classical tensor $\bar{t}_{\mu\nu}(x)$.

Triadic split and band–localisation. From (5.23),

$$\boxed{\hat{t}_{\mu\nu}^{[\mathbf{W}_1,\mathbf{W}_2]}(x) \;=\; \hat{t}_{\mu\nu}^{[\mathbf{W}_1,\mathbf{W}_2],\,\mathrm{corr}}(x) + \hat{t}_{\mu\nu}^{[\mathbf{W}_1,\mathbf{W}_2],\,\mathrm{vac}}(x) + \hat{t}_{\mu\nu}^{[\mathbf{W}_1,\mathbf{W}_2],\,\mathrm{spec}}(x),} \tag{5.25}$$

each term the SOT–Hessian of the corresponding channel. Band–localisation holds:

$$\hat{t}_{\mu\nu}^{[\mathbf{W}_1,\mathbf{W}_2]}(x) \;=\; P_x^{[\mathbf{W}_1]}\,\hat{t}_{\mu\nu}^{[\mathbf{W}_1,\mathbf{W}_2]}(x)\,P_x^{[\mathbf{W}_1]}, \qquad P_x^{[\mathbf{W}_1]} := E^{(x)[\mathbf{W}_1]}([0,\lambda_{\mathrm{cutoff}}]),$$

hence $\operatorname{Ran}\hat{t}_{\mu\nu}^{[\mathbf{W}_1,\mathbf{W}_2]}(x) \subseteq \mathfrak{H}_x$ (ultra–sub–Planckian action).

Word/frame covariance and expectation invariance. For admissible word changes

$$(\mathbf{W}_1,\mathbf{W}_2) \mapsto (\mathbf{W}_1',\mathbf{W}_2')$$

with fibre unitaries U_x,

$$\hat{t}_{\mu\nu}^{[\mathbf{W}_1',\mathbf{W}_2']}(x) = U_x^\dagger\,\hat{t}_{\mu\nu}^{[\mathbf{W}_1,\mathbf{W}_2]}(x)\,U_x, \qquad P_x^{[\mathbf{W}_1']} = U_x^\dagger\,P_x^{[\mathbf{W}_1]}\,U_x,$$

so the Holo–State expectation is invariant:

$$\boxed{\bar{t}_{\mu\nu}(x) \;:=\; \mathrm{Tr}\Big(\rho_\circ\,\hat{t}_{\mu\nu}^{[\mathbf{W}_1,\mathbf{W}_2]}(x)\Big) \quad \text{is independent of } [\mathbf{W}_1,\mathbf{W}_2],} \tag{5.26}$$

with $\rho_\circ := |\mathring{\mathbf{\Psi}}\rangle\langle\mathring{\mathbf{\Psi}}|$.

Proposition 5.19 (Positivity channel, symmetry, and vorticity)**.** *Under Spectral/Relational Smoothness and Finite Entanglement Capacity (ensuring SOT–C^2 band–regularity):*

(a) $T^{2,[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma) \succeq 0$ for all (x,x'), hence $\hat{t}_{\mu\nu}^{[\mathbf{W}_1,\mathbf{W}_2]}(x)$ is Hermitian.

(b) The expectation tensor decomposes uniquely into symmetric and antisymmetric parts, $\bar{t}_{\mu\nu} = \bar{t}_{\mu\nu}^{\mathrm{sym}} + \bar{t}_{\mu\nu}^{\mathrm{anti}}$. The symmetric part contributes to the effective Lorentzian deformation (e.g. $\bar{g}_{\mu\nu}^{\mathrm{Lor}} := \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}^{\mathrm{sym}}$), while the antisymmetric part encodes phase–flow vorticity (temporal holonomy; cf. Definition 4.563 and eq. (4.321)).

(c) With the functional orthogonality of the temporal channels (corr/vac/spec), the channel Hessians add without cross terms at expectation level.

Remark 5.20 (Optional calibration of temporal axes)**.** A fibrewise positive–definite temporal calibration $\mathbf{N}(x)$ may be inserted by replacing (5.23) with

$$T_{\mathbf{N}}^{2,[\mathbf{W}_1,\mathbf{W}_2]}(x,x';\gamma) := \hat{\boldsymbol{t}}(x,x';\gamma)^\top\,\mathbf{N}(x)\,\hat{\boldsymbol{t}}(x,x';\gamma), \qquad \hat{\boldsymbol{t}} := \Big(\hat{t}_{\mathrm{corr}},\hat{t}_{\mathrm{vac}},\hat{t}_{\mathrm{spec}}\Big)^\top,$$

yielding a covariant transformation of $\hat{t}_{\mu\nu}$ and $\bar{t}_{\mu\nu}$; it reduces to (5.24)–(5.26) for $\mathbf{N} = \mathbf{I}_3$.

Remark 5.21 (Consistency with rate forms and path dependence)**.** Along a C^1 path γ with wordwise, gauge–covariant identification of fibres, the rate forms for $\big(t_{\mathrm{corr}},t_{\mathrm{vac}},t_{\mathrm{spec}}\big)$ (cf. eq. (4.321)) give the first variation of $T^{2,[\mathbf{W}_1,\mathbf{W}_2]}$, while the SOT–Hessian $\hat{t}_{\mu\nu}^{[\mathbf{W}_1,\mathbf{W}_2]}$ captures the coincidence–limit second–order response, in agreement with (5.24).

Summary. Equations (5.23)–(5.26) define the holosymmetric wordwise temporal tensor: it is built from the triadic temporal channels, SOT–well-posed on the band–limited fibre, covariant under word/frame changes, additively decomposed across channels, and physically reads as the operator–level "phase–curvature'' whose expectation contributes to the effective Lorentzian structure on $\mathcal{X}_q$.

5.2.4 Expectation-Level Temporal Tensor

Definition 5.22 (Expectation-Level Temporal Curvature Tensor)**.** Let $\hat{t}_{\mu\nu}(x)$ be the operator-valued temporal curvature tensor at $x \in \mathcal{X}_q$ as in Definition 5.17, and let $|\overset{\circ}{\boldsymbol{\Psi}}\rangle \in \mathcal{H}_{\text{total}}$ denote the Quantum Holo-State. The *expectation-level temporal curvature tensor* is

$$\boxed{\bar{t}_{\mu\nu}(x) \;:=\; \Big\langle \overset{\circ}{\boldsymbol{\Psi}} \Big| \, \hat{t}_{\mu\nu}(x) \, \Big| \overset{\circ}{\boldsymbol{\Psi}} \Big\rangle.} \tag{5.27}$$

Equivalently, for sufficiently small Δx and twice Fréchet differentiable $T^2(x, x')$,

$$T^2(x, x + \Delta x) \;=\; \bar{t}_{\mu\nu}(x)\, \Delta x^\mu \Delta x^\nu \;+\; \mathcal{O}(|\Delta x|^3). \tag{5.28}$$

The tensor $\bar{t}_{\mu\nu}$ is a smooth, symmetric, scalar-valued $(0,2)$-field on $\mathcal{X}_q$ encoding the classical, direction-dependent curvature of quantum phase flow in the temporal fibre $\mathcal{F}_x^{\text{time}}$.

Triadic decomposition. Because $\hat{t}_{\mu\nu} = \hat{t}^{\text{corr}}_{\mu\nu} + \hat{t}^{\text{vac}}_{\mu\nu} + \hat{t}^{\text{spec}}_{\mu\nu}$,

$$\bar{t}_{\mu\nu}(x) = \bar{t}^{\text{corr}}_{\mu\nu}(x) + \bar{t}^{\text{vac}}_{\mu\nu}(x) + \bar{t}^{\text{spec}}_{\mu\nu}(x), \qquad \bar{t}^{(a)}_{\mu\nu} := \Big\langle \overset{\circ}{\boldsymbol{\Psi}} \Big| \hat{t}^{(a)}_{\mu\nu} \Big| \overset{\circ}{\boldsymbol{\Psi}} \Big\rangle. \tag{5.29}$$

Basic properties. Under the standing regularity and spectral–regulation assumptions used for $\hat{g}_{\mu\nu}$:

1. **Symmetry:** $\bar{t}_{\mu\nu} = \bar{t}_{\nu\mu}$ (mixed second derivatives at coincidence commute).
2. **Tensoriality:** $\bar{t}_{\mu\nu}$ transforms as a rank-2 covariant tensor under diffeomorphisms of $\mathcal{X}_q$.
3. **Gauge invariance:** If $\hat{t}_{\mu\nu} \mapsto U(x)\hat{t}_{\mu\nu}U(x)^\dagger$ and $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$ transforms covariantly, then $\bar{t}_{\mu\nu}$ is invariant, so physical predictions depend only on gauge-invariant functionals.
4. **Non-metric character:** $\bar{t}_{\mu\nu}$ is not constrained by positivity; it can be indefinite and does not define lengths. It encodes curvature of *phase flow*, not distance.

Chronogeometric role. Together with the expectation-level metric $\bar{g}_{\mu\nu}(x)$,

$$\bar{\Sigma}_{\mu\nu}(x) \;:=\; \bar{g}_{\mu\nu}(x) + \bar{t}_{\mu\nu}(x) \tag{5.30}$$

provides the classical chronogeometric field: $\bar{g}$ governs distinguishability (spatial sector) and $\bar{t}$ governs directional phase-flow curvature (temporal sector) in the emergent regime.

This definition (Definition 5.22) furnishes the final step in the emergence of a *classical* temporal structure from the quantum–relational foundations. It is the exact temporal analogue of the expectation–level metric tensor $\bar{g}_{\mu\nu}(x)$ (Definition 5.10), thereby completing the description of the observable chronogeometry.

The Bridge to Classical Physics

The passage from operator potentiality to classical field is effected by taking the expectation of the operator–valued tensor in the Quantum Holo–State:

$$\bar{t}_{\mu\nu}(x) \;=\; \Big\langle \overset{\circ}{\boldsymbol{\Psi}} \Big| \, \hat{t}_{\mu\nu}(x) \, \Big| \overset{\circ}{\boldsymbol{\Psi}} \Big\rangle.$$

Here $\hat{t}_{\mu\nu}(x)$ encodes the *quantum* potential for phase–flow curvature, while $\bar{t}_{\mu\nu}(x)$ is its *classical* manifestation on $\mathcal{X}_q$, directly comparable with macroscopic observables.

Physical Interpretation

The field $\bar{t}_{\mu\nu}(x)$ is a smooth, symmetric, real–valued $(0,2)$–tensor on $\mathcal{X}_q$ that quantifies the *curvature of quantum phase flow*. In contrast to $\bar{g}_{\mu\nu}(x)$—which tells how *distance* varies—$\bar{t}_{\mu\nu}(x)$ tells how the *rate of phase evolution* varies with position and direction. It thereby furnishes a geometric description of the *texture of time* absent in standard General Relativity. Phenomenologically, it may be read as:

- a local indicator of an *arrow of time* (preferred phase–flow directions),
- a measure of *temporal anisotropy* (direction–dependent phase response),
- a prospective source for novel effects related to torsion or non–metricity in the effective connection.

Relational Duality of Emergent Geometry and Temporality

A salient feature of the IQR framework is the symmetric emergence of two complementary classical tensors—$\bar{g}_{\mu\nu}$ for spatial distinguishability and $\bar{t}_{\mu\nu}$ for temporal phase flow—both descending from the same quantum–relational substrate. This parallelism exhibits a genuine duality at the heart of the emergent chronogeometry.

Hierarchy of Emergence

The symmetry is neatly summarised by the following four–level hierarchy:

Table 5.3: Hierarchy of emergence for spatial and temporal geometric structures.

Level	Spatial / Distinguishability	Temporal / Phase Flow
0 (Fundamental)	Universal state $\|\overset{\circ}{\boldsymbol{\Psi}}\rangle$ and foundational operators (Fourfold correlator $\hat{G}_\Lambda$, spectral operator $\hat{D}$) Section 1.2.9 and Definition 4.343	
1 (Quantum–Relational)	Operator–valued distances $\hat{d}_a$	Operator–valued times $\hat{t}_a$
2 (Quantum–Geometric)	Operator–valued metric $\hat{g}_{\mu\nu}(x)$ Definition 5.1	Operator–valued temporal tensor $\hat{t}_{\mu\nu}(x)$ Definition 5.17
3 (Classical–Geometric)	Expectation–level metric $\bar{g}_{\mu\nu}(x)$ Definition 5.10	Expectation–level temporal tensor $\bar{t}_{\mu\nu}(x)$ Definition 5.22

5.2.5 Expectation-Level Temporal Tensor (Holosymmetric Wordwise)

Definition 5.23 (Wordwise expectation–level temporal curvature tensor)**.** Let $\hat{t}_{\mu\nu}^{[\mathbf{W}_1,\mathbf{W}_2]}(x)$ be the wordwise operator–valued temporal curvature tensor at $x \in \mathcal{X}_q$ constructed from the triadic temporal channels with gauge–covariant comparison (cf. Definition 5.18 and Section 4.19.2). For the Quantum Holo–State $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$, define the expectation–level tensor

$$\boxed{\bar{t}_{\mu\nu}(x) \;:=\; \Big\langle \overset{\circ}{\boldsymbol{\Psi}} \Big|\, \hat{t}_{\mu\nu}^{[\mathbf{W}_1,\mathbf{W}_2]}(x) \,\Big| \overset{\circ}{\boldsymbol{\Psi}} \Big\rangle.} \tag{5.31}$$

This definition is independent of the word frame $[\mathbf{W}_1, \mathbf{W}_2]$ and of local gauge choices, since for admissible fibre unitaries U_x one has $\hat{t}_{\mu\nu}^{[\mathbf{W}_1',\mathbf{W}_2']}(x) = U_x^\dagger\, \hat{t}_{\mu\nu}^{[\mathbf{W}_1,\mathbf{W}_2]}(x)\, U_x$ and the Born expectation is unitary–invariant.

Quadratic expansion. Let $T^{2,[\mathbf{W}_1,\mathbf{W}_2]}(x, x'; \gamma)$ be the SOT–C^2 composite temporal displacement operator (sum of channel squares) on the spectrally regulated fibre $P_x\mathcal{H}_{\text{total}}$ (cf. eq. (5.23)), with $P_x := E^{(x)[\mathbf{W}_1]}([0, \lambda_{\text{cutoff}}])$ (cf. eq. (4.307)). Then, in a local chart and for $\Delta x := x' - x$,

$$\boxed{\Big\langle T^{2,[\mathbf{W}_1,\mathbf{W}_2]}(x, x+\Delta x; \gamma)\Big\rangle_{\overset{\circ}{\boldsymbol{\Psi}}} \;=\; \bar{t}_{\mu\nu}(x)\, \Delta x^\mu \Delta x^\nu \;+\; \mathcal{O}\Big(|\Delta x|^3\Big),} \tag{5.32}$$

where $\langle\cdot\rangle_{\overset{\circ}{\boldsymbol{\Psi}}}$ denotes the Holo–State expectation. Consequently, $\bar{t}_{\mu\nu}$ is a smooth, symmetric scalar $(0,2)$–field on $\mathcal{X}_q$.

Triadic decomposition (wordwise). Since $\hat{t}_{\mu\nu}^{[\mathbf{W}_1,\mathbf{W}_2]} = \hat{t}_{\mu\nu}^{[\mathbf{W}_1,\mathbf{W}_2],\,\text{corr}} + \hat{t}_{\mu\nu}^{[\mathbf{W}_1,\mathbf{W}_2],\,\text{vac}} + \hat{t}_{\mu\nu}^{[\mathbf{W}_1,\mathbf{W}_2],\,\text{spec}}$, we have

$$\boxed{\bar{t}_{\mu\nu}(x) \;=\; \bar{t}_{\mu\nu}^{\text{corr}}(x) \;+\; \bar{t}_{\mu\nu}^{\text{vac}}(x) \;+\; \bar{t}_{\mu\nu}^{\text{spec}}(x), \qquad \bar{t}_{\mu\nu}^{(a)}(x) := \Big\langle \overset{\circ}{\boldsymbol{\Psi}} \Big|\, \hat{t}_{\mu\nu}^{[\mathbf{W}_1,\mathbf{W}_2],(a)}(x) \,\Big| \overset{\circ}{\boldsymbol{\Psi}} \Big\rangle.} \tag{5.33}$$

Each term is frame/gauge–independent by the same unitary covariance.

Regularity clause (SOT and band–locality). All derivatives are taken in the strong–operator topology, uniformly on the fixed spectral band $P_x := E^{(x)[\mathbf{W}_1]}([0, \lambda_{\text{cutoff}}])$. By coherent regularisation and the smooth dependence of the transporter, $T^{2,[\mathbf{W}_1,\mathbf{W}_2]} \in \text{SOT}\text{–}C^2$ near the diagonal; hence $\hat{t}_{\mu\nu}^{[\mathbf{W}_1,\mathbf{W}_2]}(x)$ and $\bar{t}_{\mu\nu}(x)$ are well defined and smooth, with action supported on $P_x \mathcal{H}_{\text{total}}$.

The Nature of the Duality of Space and Time

By "duality'' here we mean a *conceptual and structural complementarity*, not a categorical duality. The expectation–level tensors $\bar{g}_{\mu\nu}(x)$ and $\bar{t}_{\mu\nu}(x)$ arise as two faces of the single relational fibre at x,

$$\mathfrak{F}_x \;=\; \mathcal{F}_x^{\text{dist}} \times \mathcal{F}_x^{\text{time}} \quad \rightsquigarrow \quad \{\bar{g}_{\mu\nu}(x),\, \bar{t}_{\mu\nu}(x)\},$$

as introduced in the relational bundle $\pi : \mathfrak{F} \to \mathcal{C}_d$ (eq. (4.302)) and passed to the quotient $\mathcal{X}_q$ (Section 4.19.3).

1. Fibre structure: from product to (Hilbert) direct sum. Formally, $\mathfrak{F}_x$ is the topological product $\mathcal{F}_x^{\text{dist}} \times \mathcal{F}_x^{\text{time}}$ (eq. (4.302)). On the spectrally regulated sector (eq. (4.303)) each factor is a (separable) Hilbert space with the Fourfold inner product fixed in Section 1.2.9. With respect to this inner product, the two factors are orthogonal, so the algebraic structure is the (Hilbert) direct sum

$$\mathfrak{F}_x \;\cong\; \mathcal{F}_x^{\text{dist}} \;\oplus\; \mathcal{F}_x^{\text{time}},$$

which is the useful viewpoint when forming quadratic functionals and taking second variations that define the operator–valued tensors $\hat{g}_{\mu\nu}$ and $\hat{t}_{\mu\nu}$ (Definitions 5.1 and 5.17).

2. Emergence of the tensors: the $\rightsquigarrow$ mapping. The tensors do not *live* in the fibre; they *emerge* from how fibre norms vary with base–manifold displacements in $\mathcal{X}_q$. Concretely:

- **Metric:** $\bar{g}_{\mu\nu}(x)$ is the expectation–level second variation of the squared norm of the distinguishability vector, $D^2(x, x') = \|\vec{d}(x, x')\|_2^2$, as $x' \to x$ (Definitions 5.1 and 5.10).
- **Temporal:** $\bar{t}_{\mu\nu}(x)$ is the expectation–level second variation of the squared norm of the temporal–flow vector, $T^2(x, x') = \|\vec{t}(x, x')\|_2^2$, as $x' \to x$ (Definitions 5.17 and 5.22).

Because $\mathfrak{F}_x \cong \mathcal{F}_x^{\text{dist}} \oplus \mathcal{F}_x^{\text{time}}$ is an orthogonal sum, the emergent tensors provide *independent, orthogonal* contributions to the expectation–level chronogeometry.

Summary of the relational duality.

1. *Shared mathematical structure.* Both $\bar{g}_{\mu\nu}(x)$ and $\bar{t}_{\mu\nu}(x)$ are smooth, symmetric, real–valued $(0, 2)$–tensors on $\mathcal{X}_q$, obtained as Born expectations of operator–valued second variations built from composite norms (Definitions 5.10 and 5.22).

2. *Complementary fibre origins.* $\bar{g}_{\mu\nu}$ derives from $\mathcal{F}_x^{\text{dist}}$ (distinguishability sector), while $\bar{t}_{\mu\nu}$ derives from $\mathcal{F}_x^{\text{time}}$ (phase–flow sector), both within the relational bundle $\mathfrak{F}$ (eq. (4.302)).

3. *Completeness.* Taken together, $\bar{g}_{\mu\nu}$ and $\bar{t}_{\mu\nu}$ deliver a complete infinitesimal description of the emergent, expectation–level chronogeometry: the former governs spatial distinguishability, the latter encodes curvature of quantum phase flow (Section 5.1).

Remark 5.24 (Relational duality of metric and temporal tensors)**.** The tensors $\bar{g}_{\mu\nu}$ and $\bar{t}_{\mu\nu}$, arising from structurally parallel second–variation constructions (Definitions 5.1 and 5.17) and evaluated in the Holo–State (Definitions 5.10 and 5.22), constitute the scalar–level geometry of the relational bundle $\mathfrak{F} \to \mathcal{X}_q$. Their complementarity articulates a natural *relational duality*: distance and phase are equally fundamental, operator–derived ingredients of the emergent spacetime fabric.

Tensor	**Role**	**Fiber Source**	**Physical Interpretation**
$\bar{g}_{\mu\nu}(x)$	Expectation-level **metric tensor**	$\mathcal{F}_x^{\text{dist}}$	Measures *how far apart* quantum configurations are—the curvature of distinguishability.
$\bar{t}_{\mu\nu}(x)$	Expectation-level **temporal tensor**	$\mathcal{F}_x^{\text{time}}$	Measures *how phase changes* between configurations—the curvature of quantum phase flow.

Table 5.4: Dual Tensor Roles in the Relational Phase-Geometry Bundle

Why $\mathfrak{F}_x = \mathcal{F}_x^{\text{dist}} \times \mathcal{F}_x^{\text{time}} \rightsquigarrow \{\bar{g}_{\mu\nu}, \bar{t}_{\mu\nu}\}$ is both natural and necessary. First, the product (equivalently, orthogonal direct sum) structure of the relational fibre at x is not an aesthetic choice but a *definitionally enforced* decomposition of operational content. By construction in eq. (4.302), the fibre over x aggregates the data needed to evaluate the two triads $\vec{d}$ and $\vec{t}$ after gauge–covariant comparison (parallel transport) along $\mathcal{U}$ (Section 4.19.1 and eqs. (4.11) and (4.16)). On the spectrally regulated band (eq. (4.303)), both sectors carry Hilbert inner products fixed by the Fourfold expectations (Section 1.2.9), and—by design of the triads—the "distance" and "phase–flow" observables are orthogonal modal functionals. Hence the algebraic structure is the Hilbert direct sum

$$\mathfrak{F}_x \;\cong\; \mathcal{F}_x^{\text{dist}} \;\oplus\; \mathcal{F}_x^{\text{time}},$$

topologically identical to the product but better adapted to second–variation calculus and bundle unitary actions [372, 248].

Second, the passage $\rightsquigarrow \{\bar{g}_{\mu\nu}, \bar{t}_{\mu\nu}\}$ is not a heuristic "reading off" but the *canonical Hessian extraction* of gauge–invariant quadratic functionals. Concretely, the defined composite, gauge–covariant quadratic functionals

$$D^2(x,x') = \|\vec{d}(x,x')\|^2, \qquad T^2(x,x') = \|\vec{t}(x,x')\|^2,$$

whose operator lifts D^2, T^2 are C^2 in the strong–operator topology on the regulated fibres (uniformly on the cutoff band). The Hessians at coincidence

$$\hat{g}_{\mu\nu}(x) = \partial_{x^\mu}\partial_{x'^\nu} D^2\Big|_{x'=x}, \qquad \hat{t}_{\mu\nu}(x) = \partial_{x^\mu}\partial_{x'^\nu} T^2\Big|_{x'=x}$$

(Definitions 5.1 and 5.17) are therefore uniquely determined symmetric bilinear forms (by the polarization identities/Riesz representation on each fibre). Taking Born expectations in the Holo–State then yields the classical tensors $\bar{g}_{\mu\nu}, \bar{t}_{\mu\nu}$ (Definitions 5.10 and 5.22). No additional assumptions are imported at this step.

Third, *orthogonality and minimal coupling.* Because D^2 and T^2 are built as sums of squares within their respective sectors, the mixed second variations vanish at coincidence:

$$\partial_{x^\mu}\partial_{x'^\nu} \left\langle \vec{d}, \vec{t} \right\rangle\Big|_{x'=x} = 0,$$

so the expectation–level chronogeometry splits cleanly as $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (eq. (5.30)). This is the *minimal* gauge–invariant choice consistent with the IQR operational semantics (spatial distinguishability and temporal phase flow are read by different interferometric functionals). If future physics motivates cross–channel couplings, one can add a regulated, gauge–invariant mixed functional $M(x,x') = \langle \vec{d}, \mathsf{K}\vec{t} \rangle$ (with K skew/Hermitian as appropriate), whose Hessian would produce a third tensor $\bar{m}_{\mu\nu}$. The present formalism simply chooses the orthogonal (diagonal) sectorization as the base case.

Fourth, *covariance and locality.* The fibres transform by unitary gauge on $\mathfrak{F}$ and by diffeomorphisms on $\mathcal{X}_q$; the quadratic functionals D^2, T^2 are gauge–invariant and ultra–local ($\mathcal{O}(\ell_{\mathrm{PQS}})$) by eq. (4.303). Consequently, $\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu}$ are gauge–covariant operators whose expectations $\bar{g}_{\mu\nu}, \bar{t}_{\mu\nu}$ are gauge–invariant tensor fields—precisely the objects one is allowed to regard as classical geometry on $\mathcal{X}_q$. This aligns with standard bundle geometry and spectral constructions [248, 372, 4].

Finally, *operational sufficiency.* The two quadratic functionals already saturate the second–order, gauge–invariant information accessible to the interferometric protocols: visibility moduli feed D^2 (hence $\bar{g}$), and phase–slope data feed T^2 (hence $\bar{t}$) (Section 4.19.1 and eq. (4.11)). In this sense the arrow

$$\mathfrak{F}_x \;\rightsquigarrow\; \{\bar{g}_{\mu\nu}, \bar{t}_{\mu\nu}\}$$

is not only mathematically canonical (Hessian of sectoral norms) but also *experimentally complete* at the level of second variations: it captures all observer–independent, ultra–local chronogeometric content encoded by the regulated fibre and its connection.

In short: the product (orthogonal sum) fibre is forced by how the relational observables are defined; the tensors are the unique, gauge–covariant Hessians of the associated quadratic functionals; the split is the minimal, orthogonal sectorization consistent with the operational

semantics; and the expectations are the only admissible route to classical fields on $\mathcal{X}_q$. This is precisely the bundle–theoretic, operator–geometric pathway endorsed in the standard literature [372, 248, 4] and instantiated here via eqs. (4.302) and (4.303) and Definitions 5.1, 5.10, 5.17 and 5.22.

5.2.6 The Relational Chronogeometric Operator and Tensor

The orthogonal decomposition of the relational fibre $\mathfrak{F}_x = \mathcal{F}_x^{\text{dist}} \oplus \mathcal{F}_x^{\text{time}}$ motivates a single unified geometric object that captures the total infinitesimal structure.

Definition 5.25 (Operator–Valued Relational Chronogeometric Operator)**.** Let $\hat{g}_{\mu\nu}(x)$ and $\hat{t}_{\mu\nu}(x)$ be the operator–valued $(0,2)$-tensor fields defined in Definitions 5.1 and 5.17. Define the *operator–valued relational chronogeometric operator* by

$$\hat{\Sigma}_{\mu\nu}(x) = \hat{g}_{\mu\nu}(x) + \hat{t}_{\mu\nu}(x) \tag{5.34}$$

It is Hermitian on $\mathcal{H}_{\text{total}}$ and symmetric in μ, ν. Under a local frame $U(x)$ it is gauge–covariant:

$$\hat{\Sigma}_{\mu\nu}(x) \;\longmapsto\; U(x)\,\hat{\Sigma}_{\mu\nu}(x)\,U(x)^{\dagger}. \tag{5.35}$$

Expectation–Level Relational Chronogeometric Tensor

Definition 5.26 (Expectation–Level Relational Chronogeometric Tensor)**.** Let $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$ be the Quantum Holo–State (Definition 1.27). Define

$$\bar{g}_{\mu\nu}(x) := \langle\overset{\circ}{\boldsymbol{\Psi}}|\hat{g}_{\mu\nu}(x)|\overset{\circ}{\boldsymbol{\Psi}}\rangle, \qquad \bar{t}_{\mu\nu}(x) := \langle\overset{\circ}{\boldsymbol{\Psi}}|\hat{t}_{\mu\nu}(x)|\overset{\circ}{\boldsymbol{\Psi}}\rangle.$$

The *expectation–level chronogeometric tensor* is

$$\boxed{\bar{\Sigma}_{\mu\nu}(x) \;:=\; \bar{g}_{\mu\nu}(x) \;+\; \bar{t}_{\mu\nu}(x) \;=\; \Big\langle\overset{\circ}{\boldsymbol{\Psi}}\Big|\,\hat{\Sigma}_{\mu\nu}(x)\,\Big|\overset{\circ}{\boldsymbol{\Psi}}\Big\rangle.} \tag{5.36}$$

Remark 5.27 (Properties and interpretation)**.**

1. **Mathematical structure.** Since $\bar{g}_{\mu\nu}$ and $\bar{t}_{\mu\nu}$ are smooth, symmetric, real $(0,2)$-tensors on $\mathcal{X}_q$, their sum $\bar{\Sigma}_{\mu\nu}$ is likewise a smooth, symmetric, real $(0,2)$-tensor field.
2. **Physical content.** $\bar{\Sigma}_{\mu\nu}$ encodes the *total* infinitesimal relational response of the fibre $\mathfrak{F}_x$:
   - the curvature of distinguishability (the metric sector) via $\bar{g}_{\mu\nu}$;
   - the curvature of quantum phase flow (the temporal sector) via $\bar{t}_{\mu\nu}$.

   It is not a metric in the Riemannian sense; rather, a unified *chronogeometric* tensor capturing both "how far apart" and "how out of phase" nearby configurations are.
3. **Gauge covariance and observables.** At the operator level, $\hat{\Sigma}_{\mu\nu}$ is gauge–covariant, while scalar observables are obtained from gauge–invariant functionals (expectations, regulated traces). The expectation $\bar{\Sigma}_{\mu\nu}$ is frame–independent.

4. **Dynamical implications.** In a relational variational principle, matter and geometry may couple to the full $\bar{\Sigma}_{\mu\nu}$, not only to $\bar{g}_{\mu\nu}$. This enlarges the gravitational sector to include responses to phase–flow curvature carried by $\bar{t}_{\mu\nu}$, see Definition 5.26 and subsequent field equations.

A compact statement of emergence is

$$\underbrace{\bar{\Sigma}_{\mu\nu}(x)}_{\text{classical chronogeometry}} \quad = \quad \underbrace{\left\langle \mathring{\mathbf{\Psi}} \middle| \hat{\Sigma}_{\mu\nu}(x) \middle| \mathring{\mathbf{\Psi}} \right\rangle}_{\text{quantum expectation in Definition 1.27}} \quad ,$$

with $\hat{\Sigma}_{\mu\nu} = \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu}$ from Definitions 5.1, 5.17 and 5.25.

The introduction of $\bar{\Sigma}_{\mu\nu}(x)$ provides a powerful tool for exploring the deep connection between the spatial and temporal aspects of our emergent reality, suggesting that they are inseparable facets of a single, underlying quantum-relational structure.

Corollary 5.28 (Gauge invariance of the expectation-level chronogeometric tensor)**.** *Let $\hat{\Sigma}_{\mu\nu}(x)$ be the unified operator-valued chronogeometric tensor of Definition 5.25, transforming covariantly under local gauge unitaries $U(x)$ as in eq. (5.35). Let the Quantum Holo–State transform as $|\mathring{\mathbf{\Psi}}\rangle \mapsto |\mathring{\mathbf{\Psi}}'\rangle := U|\mathring{\mathbf{\Psi}}\rangle$. Then the expectation-level tensor $\bar{\Sigma}_{\mu\nu}(x) = \langle\mathring{\mathbf{\Psi}}|\hat{\Sigma}_{\mu\nu}(x)|\mathring{\mathbf{\Psi}}\rangle$ of Definition 5.26 is gauge invariant:*

$$\bar{\Sigma}'_{\mu\nu}(x) \;:=\; \langle\mathring{\mathbf{\Psi}}'|\hat{\Sigma}'_{\mu\nu}(x)|\mathring{\mathbf{\Psi}}'\rangle \;=\; \bar{\Sigma}_{\mu\nu}(x).$$

Proof. By Definition 5.25, the unified operator is $\hat{\Sigma}_{\mu\nu}(x) = \hat{g}_{\mu\nu}(x) + \hat{t}_{\mu\nu}(x)$, and it transforms covariantly under local frame changes $U(x)$ as

$$\hat{\Sigma}'_{\mu\nu}(x) = U(x)\,\hat{\Sigma}_{\mu\nu}(x)\,U(x)^\dagger \qquad \text{(eq. (5.35))}. \tag{5.37}$$

The expectation–level tensor is $\bar{\Sigma}_{\mu\nu}(x) = \langle\mathring{\mathbf{\Psi}}|\hat{\Sigma}_{\mu\nu}(x)|\mathring{\mathbf{\Psi}}\rangle$ (Definition 5.26). Under the same gauge, the Quantum Holo–State transforms as $|\mathring{\mathbf{\Psi}}\rangle \mapsto |\mathring{\mathbf{\Psi}}'\rangle := U|\mathring{\mathbf{\Psi}}\rangle$ (as stated in the corollary). Since the expectation at x couples the fibre at x, we write $U_x := U(x)$ and compute

$$\begin{aligned}\bar{\Sigma}'_{\mu\nu}(x) &:= \langle\mathring{\mathbf{\Psi}}'|\hat{\Sigma}'_{\mu\nu}(x)|\mathring{\mathbf{\Psi}}'\rangle = \langle\mathring{\mathbf{\Psi}}|\,U^\dagger\left[U_x\,\hat{\Sigma}_{\mu\nu}(x)\,U_x^\dagger\right]U\,|\mathring{\mathbf{\Psi}}\rangle \\ &= \langle\mathring{\mathbf{\Psi}}|\left[U^\dagger U_x\right]\,\hat{\Sigma}_{\mu\nu}(x)\,\left[U_x^\dagger U\right]\,|\mathring{\mathbf{\Psi}}\rangle.\end{aligned}$$

In the standard implementation of local frame changes, U acts pointwise on the fibre so that on the same fibre $U^\dagger U_x = U_x^\dagger U = \mathbb{I}$. Hence

$$\bar{\Sigma}'_{\mu\nu}(x) = \langle\mathring{\mathbf{\Psi}}|\,\hat{\Sigma}_{\mu\nu}(x)\,|\mathring{\mathbf{\Psi}}\rangle = \bar{\Sigma}_{\mu\nu}(x),$$

proving gauge invariance. The same conclusion follows termwise for $\bar{g}_{\mu\nu}$ and $\bar{t}_{\mu\nu}$, since each is defined by an identical expectation and $\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu}$ are themselves covariant (by construction in Definitions 5.1 and 5.17).

Equivalent density-operator proof. Let $\rho := |\overset{\circ}{\mathbf{\Psi}}\rangle\langle\overset{\circ}{\mathbf{\Psi}}|$. Then $\bar{\Sigma}_{\mu\nu}(x) = \mathrm{tr}\big[\hat{\Sigma}_{\mu\nu}(x)\,\rho\big]$ (Definition 5.26). Under the gauge, $\rho' = U\,\rho\,U^\dagger$ and $\hat{\Sigma}'_{\mu\nu}(x) = U_x\,\hat{\Sigma}_{\mu\nu}(x)\,U_x^\dagger$ (eq. (5.35)). Therefore,

$$\bar{\Sigma}'_{\mu\nu}(x) = \mathrm{tr}\big[\hat{\Sigma}'_{\mu\nu}(x)\,\rho'\big] = \mathrm{tr}\big[U_x\,\hat{\Sigma}_{\mu\nu}(x)\,U_x^\dagger\,U\,\rho\,U^\dagger\big] = \mathrm{tr}\big[\hat{\Sigma}_{\mu\nu}(x)\,\rho\big] = \bar{\Sigma}_{\mu\nu}(x),$$

using cyclicity of the trace and the fact that the pointwise and global actions agree on the same fibre.

Thus $\bar{\Sigma}_{\mu\nu}(x)$ is gauge invariant, establishing Corollary 5.28. □

Corollary 5.29 (Sector-wise reductions of the unified tensor)**.** *With $\hat{\Sigma}_{\mu\nu}(x) = \hat{g}_{\mu\nu}(x) + \hat{t}_{\mu\nu}(x)$ as in Definition 5.25 and $\bar{g}_{\mu\nu}, \bar{t}_{\mu\nu}$ as in Definition 5.26, the following reductions hold:*

(i) Temporal quiescence: *If $\bar{t}_{\mu\nu}(x) = 0$ on an open set $O \subseteq M$, then $\bar{\Sigma}_{\mu\nu}(x) = \bar{g}_{\mu\nu}(x)$ for all $x \in O$.*

(ii) Ultralocal distinguishability (flat metric sector): *If $\bar{g}_{\mu\nu}(x) = 0$ on an open set $O \subseteq M$, then $\bar{\Sigma}_{\mu\nu}(x) = \bar{t}_{\mu\nu}(x)$ for all $x \in O$.*

Proof. Recall the unified operator and its expectation-level tensor:

$$\hat{\Sigma}_{\mu\nu}(x) = \hat{g}_{\mu\nu}(x) + \hat{t}_{\mu\nu}(x) \quad \text{(Definition 5.25)}$$
$$\bar{\Sigma}_{\mu\nu}(x) = \Big\langle \overset{\circ}{\mathbf{\Psi}} \Big|\, \hat{\Sigma}_{\mu\nu}(x) \,\Big|\, \overset{\circ}{\mathbf{\Psi}} \Big\rangle \quad \text{(Definition 5.26)}$$

By linearity of the expectation functional and $\hat{\Sigma}_{\mu\nu} = \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu}$,

$$\bar{\Sigma}_{\mu\nu}(x) = \langle\overset{\circ}{\mathbf{\Psi}}|\hat{g}_{\mu\nu}(x)|\overset{\circ}{\mathbf{\Psi}}\rangle + \langle\overset{\circ}{\mathbf{\Psi}}|\hat{t}_{\mu\nu}(x)|\overset{\circ}{\mathbf{\Psi}}\rangle = \bar{g}_{\mu\nu}(x) + \bar{t}_{\mu\nu}(x), \tag{5.38}$$

which is the decomposition recorded in Definition 5.26.

(1) Temporal quiescence. Assume $\bar{t}_{\mu\nu}(x) = 0$ for all $x \in O$. Then by (5.38),

$$\forall x \in O: \qquad \bar{\Sigma}_{\mu\nu}(x) = \bar{g}_{\mu\nu}(x) + 0 = \bar{g}_{\mu\nu}(x).$$

Since $\bar{g}_{\mu\nu}$ and $\bar{\Sigma}_{\mu\nu}$ are smooth real symmetric $(0,2)$-tensors (Definition 5.26 and the ensuing remark), the equality holds tensorially on O, i.e. in any frame or chart.

(2) Ultralocal distinguishability (flat metric sector). Assume $\bar{g}_{\mu\nu}(x) = 0$ for all $x \in O$. Then (5.38) gives

$$\forall x \in O: \qquad \bar{\Sigma}_{\mu\nu}(x) = 0 + \bar{t}_{\mu\nu}(x) = \bar{t}_{\mu\nu}(x),$$

again as an identity of $(0,2)$-tensors on O.

Gauge/frame independence. By Corollary 5.28, $\bar{\Sigma}_{\mu\nu}$, $\bar{g}_{\mu\nu}$, and $\bar{t}_{\mu\nu}$ are each gauge-invariant under local frame unitaries $U(x)$ (cf. the operator-level covariance eq. (5.35)). Hence the reductions proved above are frame-independent statements on O.

Sufficient operator-level condition (optional). If, stronger than the hypotheses, one has $\hat{t}_{\mu\nu}(x) = 0$ (resp. $\hat{g}_{\mu\nu}(x) = 0$) as operators on O, then $\bar{t}_{\mu\nu}(x) = 0$ (resp. $\bar{g}_{\mu\nu}(x) = 0$) for *any* state, so the same reductions follow immediately from (5.38).

This establishes both items in Corollary 5.29. □

Definition 5.30 (Holosymmetric lift of the chronogeometric operator)**.** Let the law-control register be $\mathcal{H}_L = \mathrm{span}\{|0\rangle_L, |1\rangle_L\}$ and $\hat{\Sigma}^{(k)}_{\mu\nu}(x) = \mathbb{L}_k[\hat{g}_{\mu\nu}(x), \hat{t}_{\mu\nu}(x)]$ for $k \in \{0, 1\}$ (two functorial "law maps" from Definition 5.25).

Define the *holosymmetric lift* by

$$\begin{aligned}\widehat{\mathbf{\Sigma}}^{\mathrm{HS}}_{\mu\nu}(x) := |0\rangle\langle 0|_L \otimes \hat{\Sigma}^{(0)}_{\mu\nu}(x) + |1\rangle\langle 1|_L \otimes \hat{\Sigma}^{(1)}_{\mu\nu}(x) \\ + |0\rangle\langle 1|_L \otimes \hat{\mathcal{I}}_{\mu\nu}(x) + |1\rangle\langle 0|_L \otimes \hat{\mathcal{I}}^{\dagger}_{\mu\nu}(x),\end{aligned} \tag{5.39}$$

where the *interference operator* $\hat{\mathcal{I}}_{\mu\nu}(x)$ is built functorially from $(\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu})$ (by sums/products/(anti)commutators) and fixed by symmetry/locality.

Definition 5.31 (Holosymmetric reduction on the system)**.** Prepare $|+_\varphi\rangle_L = (|0\rangle_L + e^{i\varphi}|1\rangle_L)/\sqrt{2}$ and postselect the same readout. Then

$$\hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu}(x;\varphi) = \tfrac{1}{2}\Big(\hat{\Sigma}^{(0)}_{\mu\nu}(x) + \hat{\Sigma}^{(1)}_{\mu\nu}(x)\Big) + \mathrm{Re}\Big(e^{-i\varphi}\hat{\mathcal{I}}_{\mu\nu}(x)\Big)\,. \tag{5.40}$$

At the expectation level with the Quantum Holo–State,

$$\bar{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu}(x;\varphi) = \tfrac{1}{2}\Big(\langle\hat{\Sigma}^{(0)}_{\mu\nu}\rangle + \langle\hat{\Sigma}^{(1)}_{\mu\nu}\rangle\Big) + \mathrm{Re}\Big(e^{-i\varphi}\langle\hat{\mathcal{I}}_{\mu\nu}\rangle\Big)\,. \tag{5.41}$$

Canonical parametrisation (superposition of laws). A natural choice of two laws that differ by the temporal sign is

$$\hat{\Sigma}^{(0)}_{\mu\nu}(x) = \hat{g}_{\mu\nu}(x) + \hat{t}_{\mu\nu}(x), \qquad \hat{\Sigma}^{(1)}_{\mu\nu}(x) = \hat{g}_{\mu\nu}(x) - \hat{t}_{\mu\nu}(x). \tag{5.42}$$

For the interference channel, impose Hermiticity/locality and set

$$\hat{\mathcal{I}}^{(\theta)}_{\mu\nu}(x) := \tfrac{1}{2}\Big(\cos\theta\,\{\hat{g}_{\mu\nu}(x), \hat{t}_{\mu\nu}(x)\} + i\sin\theta\,[\hat{g}_{\mu\nu}(x), \hat{t}_{\mu\nu}(x)]\Big), \tag{5.43}$$

so that substituting (5.42) and (5.43) into (5.40) yields

$$\boxed{\hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu}(x;\varphi,\theta) = \hat{g}_{\mu\nu}(x) + \tfrac{1}{2}\cos\varphi\,\cos\theta\,\{\hat{g}_{\mu\nu}(x), \hat{t}_{\mu\nu}(x)\} + \tfrac{1}{2}\sin\varphi\,\sin\theta\; i[\hat{g}_{\mu\nu}(x), \hat{t}_{\mu\nu}(x)].} \tag{5.44}$$

Proposition 5.32 (Gauge covariance and expectation-level invariance)**.** *If* $(\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu}) \mapsto \big(U(x)\hat{g}_{\mu\nu}(x)U(x)^\dagger,\ U(x)\hat{t}_{\mu\nu}(x)U(x)^\dagger\big)$ *and* $|\overset{\circ}{\mathbf{\Psi}}\rangle \mapsto U|\overset{\circ}{\mathbf{\Psi}}\rangle$ *as in eq. (5.35), and if* $\mathbb{L}_0, \mathbb{L}_1$ *and* $\hat{\mathcal{I}}_{\mu\nu}$ *are built functorially from* $(\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu})$ *(via sums, products, (anti)commutators), then* $\widehat{\mathbf{\Sigma}}^{\mathrm{HS}}_{\mu\nu}$ *transforms covariantly and* $\bar{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu}(x;\varphi)$ *is gauge invariant.*

Proof. **Functorial covariance of the blocks.** Conjugation by $U(x)$ is a $*$–automorphism; for fibre operators X, Y,

$$\begin{aligned} U(X+Y)U^\dagger &= UXU^\dagger + UYU^\dagger, \\ U(XY)U^\dagger &= (UXU^\dagger)(UYU^\dagger), \\ U\{X,Y\}U^\dagger &= \{UXU^\dagger, UYU^\dagger\}, \\ U[X,Y]U^\dagger &= [UXU^\dagger, UYU^\dagger]. \end{aligned} \tag{5.45}$$

Since $\hat{\Sigma}^{(k)}_{\mu\nu} = \mathbb{L}_k[\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu}]$ and $\hat{\mathcal{I}}_{\mu\nu}$ are built from $(\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu})$ by these operations, we have

$$\begin{aligned} \hat{\Sigma}^{(k)}_{\mu\nu}(x) &\mapsto U(x)\hat{\Sigma}^{(k)}_{\mu\nu}(x)U(x)^\dagger, \\ \hat{\mathcal{I}}_{\mu\nu}(x) &\mapsto U(x)\hat{\mathcal{I}}_{\mu\nu}(x)U(x)^\dagger, \\ \hat{\mathcal{I}}^\dagger_{\mu\nu}(x) &\mapsto U(x)\hat{\mathcal{I}}^\dagger_{\mu\nu}(x)U(x)^\dagger. \end{aligned} \tag{5.46}$$

Covariance of the holosymmetric lift. By Definition 5.30,

$$\hat{\mathbf{\Sigma}}^{\mathrm{HS}}_{\mu\nu} = |0\rangle\langle 0|_L \otimes \hat{\Sigma}^{(0)}_{\mu\nu} + |1\rangle\langle 1|_L \otimes \hat{\Sigma}^{(1)}_{\mu\nu} + |0\rangle\langle 1|_L \otimes \hat{\mathcal{I}}_{\mu\nu} + |1\rangle\langle 0|_L \otimes \hat{\mathcal{I}}^\dagger_{\mu\nu} \quad (\textit{eq. } (5.39)).$$

Applying (5.46) termwise yields

$$\hat{\mathbf{\Sigma}}^{\mathrm{HS}}_{\mu\nu}(x) \;\mapsto\; (\mathbb{I}_L \otimes U(x))\, \hat{\mathbf{\Sigma}}^{\mathrm{HS}}_{\mu\nu}(x)\, (\mathbb{I}_L \otimes U(x))^\dagger,$$

so the lift is gauge–covariant.

Covariance of the reduced operator. Reducing with $|+_\varphi\rangle_L$ (Definition 5.31) gives

$$\hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu}(x;\varphi) = \tfrac{1}{2}\Big(\hat{\Sigma}^{(0)}_{\mu\nu}(x) + \hat{\Sigma}^{(1)}_{\mu\nu}(x)\Big) + \mathrm{Re}\Big(e^{-i\varphi}\hat{\mathcal{I}}_{\mu\nu}(x)\Big) \quad (\textit{eq. } (5.40)). \tag{5.47}$$

Using (5.46) and

$$\mathrm{Re}\Big(e^{-i\varphi}\hat{\mathcal{I}}_{\mu\nu}\Big) \;=\; \tfrac{1}{2}\Big(e^{-i\varphi}\hat{\mathcal{I}}_{\mu\nu} + e^{+i\varphi}\hat{\mathcal{I}}^\dagger_{\mu\nu}\Big) \;\mapsto\; U(x)\,\mathrm{Re}\Big(e^{-i\varphi}\hat{\mathcal{I}}_{\mu\nu}\Big)\,U(x)^\dagger,$$

we obtain

$$\hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu}(x;\varphi) \;\mapsto\; U(x)\,\hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu}(x;\varphi)\,U(x)^\dagger,$$

i.e. the reduced operator is gauge–covariant.

Expectation–level invariance. Let $|\mathring{\mathbf{\Psi}}\rangle \mapsto U|\mathring{\mathbf{\Psi}}\rangle$ (eq. (5.35)). By eq. (5.41) and the standard endpoint–gauge convention (on the fibre at x, $U^\dagger U(x) = \mathbb{I}$),

$$\begin{aligned} \bar{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu}(x;\varphi) &= \Big\langle \mathring{\mathbf{\Psi}}\Big|\, \hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu}(x;\varphi)\,\Big|\mathring{\mathbf{\Psi}}\Big\rangle \\ &\mapsto \Big\langle \mathring{\mathbf{\Psi}}\Big|\, U^\dagger(x)\Big(U(x)\hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu}(x;\varphi)U(x)^\dagger\Big)U(x)\,\Big|\mathring{\mathbf{\Psi}}\Big\rangle \\ &= \bar{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu}(x;\varphi), \end{aligned} \tag{5.48}$$

so the expectation–level quantity is gauge invariant.

These steps establish Proposition 5.32. □

Corollary 5.33 (Sector/commutation limits of the holosymmetric application)**.** *With eq. (5.44):*

(i) If $[\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu}] = 0$, then $\hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu} = \hat{g}_{\mu\nu} + \frac{1}{2}\cos\varphi\,\cos\theta\,\{\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu}\}$, so the interference is purely symmetric (no phase-odd mixing).

(ii) If furthermore $\{\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu}\} = 2\hat{g}_{\mu\nu}\hat{t}_{\mu\nu}$ (ultralocal), then

$$\hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu} = \hat{g}_{\mu\nu} + \cos\varphi\,\cos\theta\,\hat{g}_{\mu\nu}\hat{t}_{\mu\nu}.$$

(iii) Choosing $\varphi = 0$ isolates anticommutator mixing; choosing $\varphi = \frac{\pi}{2}$ isolates commutator mixing.

At the expectation level, the corresponding statements hold for $\bar{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu}(x;\varphi,\theta)$*; if the ultralocal product is small, one may then consistently truncate higher-order corrections in any subsequent expansions.*

Proof. From eq. (5.44),

$$\hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu} = \hat{g}_{\mu\nu} + \tfrac{1}{2}\cos\varphi\,\cos\theta\,\{\hat{g}_{\mu\nu},\hat{t}_{\mu\nu}\} + \tfrac{1}{2}\sin\varphi\,\sin\theta\, i[\hat{g}_{\mu\nu},\hat{t}_{\mu\nu}].$$

(1) If $[\hat{g}_{\mu\nu},\hat{t}_{\mu\nu}]=0$, the last term vanishes, yielding the stated symmetric form.

(2) Under the additional ultralocal identity $\{\hat{g}_{\mu\nu},\hat{t}_{\mu\nu}\} = 2\hat{g}_{\mu\nu}\hat{t}_{\mu\nu}$ (which is automatic if they commute and the product is pointwise), the formula reduces exactly to $\hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu} = \hat{g}_{\mu\nu} + \cos\varphi\,\cos\theta\,\hat{g}_{\mu\nu}\hat{t}_{\mu\nu}$.

(3) Setting $\varphi = 0$ gives $\cos\varphi = 1$, $\sin\varphi = 0$, isolating the anticommutator sector; setting $\varphi = \frac{\pi}{2}$ gives $\cos\varphi = 0$, $\sin\varphi = 1$, isolating the commutator sector, consistent with Proposition 5.42.

Taking expectations and using linearity (eq. (5.41)) yields the stated expectation-level versions. □

Remark 5.34 (Operational reading)**.** The phase φ (law-control preparation/readout) selects symmetric vs. antisymmetric sector coupling; θ sets the strength of coherence between the "two laws." For $\theta = \varphi = 0$, one recovers the classical mixture (no inter-law coherence): $\hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu} = \frac{1}{2}\left(\hat{\Sigma}^{(0)}_{\mu\nu} + \hat{\Sigma}^{(1)}_{\mu\nu}\right) = \hat{g}_{\mu\nu}$.

Lemma 5.35 (Unitary holosymmetric embedding of chronogeometry)**.** *Let the two chronogeometric histories be represented by Hermitian tensors* $\hat{\Sigma}^{(A_G B_N)}_{\mu\nu}(x)$ *and* $\hat{\Sigma}^{(A_N B_G)}_{\mu\nu}(x)$*. Choose a real smearing current* $J^{\mu\nu}(x)$ *(compactly supported), and define*

$$\hat{\mathcal{H}}_0 \;:=\; \int d^d x\; J^{\mu\nu}(x)\,\hat{\Sigma}^{(A_G B_N)}_{\mu\nu}(x), \qquad \hat{\mathcal{H}}_1 \;:=\; \int d^d x\; J^{\mu\nu}(x)\,\hat{\Sigma}^{(A_N B_G)}_{\mu\nu}(x).$$

The corresponding (history) unitaries are

$$\hat{U}_0 \;:=\; \mathcal{T}\, e^{-i\int d\tau\,\hat{\mathcal{H}}_0(\tau)}, \qquad \hat{U}_1 \;:=\; \mathcal{T}\, e^{-i\int d\tau\,\hat{\mathcal{H}}_1(\tau)}.$$

Holosymmetry is then implemented by the controlled unitary on control c *and registers* RR*:*

$$\mathbf{U}_{\mathrm{ctrl}} = \;|\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \hat{U}_0 \;+\; |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \hat{U}_1,$$

which is unitary and realises a superposition of the two chronogeometric histories.

Proof. **Step 1: Hermiticity of the smeared generators.** By assumption, each history tensor $\hat{\Sigma}^{(A_G B_N)}_{\mu\nu}(x)$ and $\hat{\Sigma}^{(A_N B_G)}_{\mu\nu}(x)$ is Hermitian on the system Hilbert space, and the smearing current $J^{\mu\nu}(x)$ is real with compact support. Hence

$$\hat{\mathcal{H}}_0 = \int d^d x\, J^{\mu\nu}(x)\,\hat{\Sigma}^{(A_G B_N)}_{\mu\nu}(x), \qquad \hat{\mathcal{H}}_1 = \int d^d x\, J^{\mu\nu}(x)\,\hat{\Sigma}^{(A_N B_G)}_{\mu\nu}(x)$$

are Hermitian operators (linear combinations of Hermitian fields with real coefficients).

Step 2: Unitarity of the time–ordered evolutions. Let $\hat{U}_k(\tau)$ solve the Schrödinger equation $\partial_\tau \hat{U}_k(\tau) = -\,i\,\hat{\mathcal{H}}_k(\tau)\,\hat{U}_k(\tau)$ with $\hat{U}_k(\tau_0) = I$. Then

$$\frac{d}{d\tau}\big(\hat{U}_k^\dagger(\tau)\hat{U}_k(\tau)\big) = \big(i\,\hat{U}_k^\dagger\hat{\mathcal{H}}_k\big)\hat{U}_k + \hat{U}_k^\dagger\big(-\,i\,\hat{\mathcal{H}}_k\hat{U}_k\big) = 0,$$

since $\hat{\mathcal{H}}_k(\tau)$ is Hermitian. Therefore $\hat{U}_k^\dagger(\tau)\hat{U}_k(\tau) = I$ for all τ, and the time–ordered solution $\hat{U}_k = \mathcal{T}\, e^{-i\int d\tau\, \hat{\mathcal{H}}_k(\tau)}$ is unitary ($k = 0, 1$).

Step 3: Unitarity of the controlled word. Define the control projectors $P_0 := |\mathbf{0}\rangle_c\langle\mathbf{0}|_c$ and $P_1 := |\mathbf{1}\rangle_c\langle\mathbf{1}|_c$, which obey $P_aP_b = \delta_{ab}P_a$ and $P_0 + P_1 = I_c$. The controlled operator (cf. eq. (2.75) and Proposition 4.152)

$$\mathbf{U}_{\text{ctrl}} = P_0\otimes\hat{U}_0 \;+\; P_1\otimes\hat{U}_1$$

has adjoint $\mathbf{U}_{\text{ctrl}}^\dagger = P_0\otimes\hat{U}_0^\dagger + P_1\otimes\hat{U}_1^\dagger$. Using orthogonality of P_0, P_1 and unitarity of $\hat{U}_k$,

$$\mathbf{U}_{\text{ctrl}}^\dagger\mathbf{U}_{\text{ctrl}} = (P_0\otimes\hat{U}_0^\dagger\hat{U}_0) + (P_1\otimes\hat{U}_1^\dagger\hat{U}_1) = (P_0 + P_1)\otimes I_{\text{RR}} = I_c\otimes I_{\text{RR}},$$

and similarly $\mathbf{U}_{\text{ctrl}}\mathbf{U}_{\text{ctrl}}^\dagger = I_c\otimes I_{\text{RR}}$. Thus $\mathbf{U}_{\text{ctrl}}$ is unitary on $\mathcal{H}_c\otimes\mathcal{H}_{\text{RR}}^{(2n)}$ (cf. Definition 2.101 and eq. (2.62)).

Step 4: Realisation of a coherent superposition of histories. Prepare the control in $|+_\varphi\rangle_c = (|\mathbf{0}\rangle_c + e^{i\varphi}|\mathbf{1}\rangle_c)/\sqrt{2}$ and the registers in $|\Psi_{\text{RR}}\rangle$. Then

$$\mathbf{U}_{\text{ctrl}}\Big(|+_\varphi\rangle_c\otimes|\Psi_{\text{RR}}\rangle\Big) = \frac{1}{\sqrt{2}}\Big(|\mathbf{0}\rangle_c\otimes\hat{U}_0|\Psi_{\text{RR}}\rangle + e^{i\varphi}|\mathbf{1}\rangle_c\otimes\hat{U}_1|\Psi_{\text{RR}}\rangle\Big),$$

which is a *coherent* control–register state encoding the two chronogeometric histories. Upon recombination/measurement in the X–basis one obtains the usual sum/difference interference (cf. the wordwise interferometer analysis in Propositions 4.153 and 5.38).

Combining Steps 1–3 establishes unitarity; Step 4 shows the operational superposition of the two histories. This proves Lemma 5.35. □

Proposition 5.36 (Interference = relational overlap of spacetimes)**.** *Prepare the control in* $|+_\varphi\rangle_c = (|\mathbf{0}\rangle_c + e^{i\varphi}|\mathbf{1}\rangle_c)/\sqrt{2}$ *and the registers in* $|\Psi_{\text{RR}}\rangle$*. Apply* $\mathbf{U}_{\text{ctrl}}$*, recombine with the inverse readout on c (Hadamard-with-phase), and measure c in the computational basis. Then*

$$p_c(\mathbf{0}\mid\varphi) \;=\; \tfrac{1}{2}\Big(1 + \text{Re}\Big[e^{-i\varphi}\,\langle\Psi_{\text{RR}}|\hat{U}_1^\dagger\hat{U}_0|\Psi_{\text{RR}}\rangle\Big]\Big),$$

so the visibility is set by the relational overlap *between the two chronogeometric unitaries. In particular, if* $\hat{U}_k = e^{-i\lambda\int J^{\mu\nu}\hat{\Sigma}_{\mu\nu}^{(k)}}$ *with small* λ*, then to leading order the signal decomposes into commutator/anticommutator pieces of* $\hat{g}_{\mu\nu}$ *and* $\hat{t}_{\mu\nu}$*, matching eq. (5.44).*

Proof. **Interference fringe from the controlled device.** Prepare $|+_\varphi\rangle_c = (|\mathbf{0}\rangle_c + e^{i\varphi}|\mathbf{1}\rangle_c)/\sqrt{2}$ and $|\Psi_{\text{RR}}\rangle$. Apply $\mathbf{U}_{\text{ctrl}} = |\mathbf{0}\rangle\langle\mathbf{0}|_c\otimes\hat{U}_0 + |\mathbf{1}\rangle\langle\mathbf{1}|_c\otimes\hat{U}_1$ (Lemma 5.35 and eq. (2.75)). Recombine on c with the inverse Hadamard-with-phase and measure in $\mathcal{B}_c^{(Z)}$ (cf. the standard wordwise

readout in Proposition 4.153). A direct calculation gives the (unnormalized) $|\mathbf{0}\rangle_c$-component $\frac{1}{2}\big(\hat{U}_0 + e^{-i\varphi}\hat{U}_1\big)|\Psi_{\mathrm{RR}}\rangle$; hence

$$p_c(\mathbf{0} \mid \varphi) = \frac{1}{4}\left(\|\hat{U}_0|\Psi_{\mathrm{RR}}\rangle\|^2 + \|\hat{U}_1|\Psi_{\mathrm{RR}}\rangle\|^2 + e^{-i\varphi}\langle\Psi_{\mathrm{RR}}|\hat{U}_1^\dagger\hat{U}_0|\Psi_{\mathrm{RR}}\rangle + e^{+i\varphi}\langle\Psi_{\mathrm{RR}}|\hat{U}_0^\dagger\hat{U}_1|\Psi_{\mathrm{RR}}\rangle\right).$$

Since $\hat{U}_k$ are unitary (Lemma 5.35), $\|\hat{U}_k\Psi_{\mathrm{RR}}\| = \|\Psi_{\mathrm{RR}}\| = 1$, so

$$p_c(\mathbf{0} \mid \varphi) = \tfrac{1}{2}\Big(1 + \mathrm{Re}\Big[e^{-i\varphi}\,\langle\Psi_{\mathrm{RR}}|\hat{U}_1^\dagger\hat{U}_0|\Psi_{\mathrm{RR}}\rangle\Big]\Big), \tag{5.49}$$

as claimed.

Why the Baker–Campbell–Hausdorff (BCH) formula matters here. The BCH formula is what makes the "two–history" product $\hat{U}_1^\dagger\hat{U}_0 = e^{+i\lambda S_1}e^{-i\lambda S_0}$ both analytically tractable and physically transparent in our framework. First, BCH combines the two exponentials into a *single* effective evolution,

$$\hat{U}_1^\dagger\hat{U}_0 = \exp\Big(-i\lambda K + \tfrac{\lambda^2}{2}[S_1, S_0] + O(\lambda^3)\Big), \qquad K := S_0 - S_1, \tag{5.50}$$

precisely as recorded in eq. (5.53) and used in Proposition 5.39. This isolates the "difference generator'' K (the leading effect) and the first genuinely non–Abelian correction $[S_1, S_0]$. Standard references for the BCH expansion and unitary exponentials include [253, 397, 404, 287].

Visibility vs. phase (who does what). When S_0, S_1 are Hermitian, the commutator $[S_1, S_0]$ is anti–Hermitian, so its expectation value is purely imaginary. To order $O(\lambda^2)$ this term therefore contributes a *phase* to the complex overlap $\langle\Psi_{\mathrm{RR}}|\hat{U}_1^\dagger\hat{U}_0|\Psi_{\mathrm{RR}}\rangle$, while the *magnitude* (and hence the fringe visibility) is reduced by the variance of K. Concretely,

$$\langle\Psi_{\mathrm{RR}}|\hat{U}_1^\dagger\hat{U}_0|\Psi_{\mathrm{RR}}\rangle = 1 - \tfrac{\lambda^2}{2}\,\mathrm{Var}_{\Psi_{\mathrm{RR}}}(K) + O(\lambda^3),$$

as stated in Proposition 5.39. This is fully consistent with the quantum–optical visibility calculus for interferometers [408, 287].

Commuting vs. non–commuting histories. If the two histories commute at the generator level, $[S_1, S_0] = 0$, then BCH collapses to $\hat{U}_1^\dagger\hat{U}_0 = \exp(-i\lambda K) + O(\lambda^3)$. This is exactly the "sector/commutation limit'' used alongside the holosymmetric reduction in eq. (5.44) and summarized in Corollaries 5.33 and 5.43.

Bridge to the operator mixing $\{\cdot,\cdot\}$ *vs.* $[\cdot,\cdot]$. Chapter 5 decomposes the reduced operator into symmetric (anticommutator) and phase–odd (commutator) parts, eq. (5.44). BCH provides the dynamical counterpart in the small–deformation regime: it cleanly *separates* the visibility–controlling difference generator K from the commutator–controlled relative phase. This aligns with the phase–resolved tomography protocol of Proposition 5.42, which uses the control phase φ to isolate anticommutator vs. commutator responses.

Order counting and truncation discipline. BCH makes explicit that non–Abelian corrections enter at $O(\lambda^2)$ (via $[S_1, S_0]$) and higher (nested commutators at $O(\lambda^3)$, etc.), justifying

the $O(\lambda^2)$ truncations in Proposition 5.39 and keeping the perturbation theory controlled [253, 397].

In short, BCH (i) fuses the two chronogeometric evolutions into one effective generator, (ii) separates *visibility* (variance of K) from *phase* (commutator term), and (iii) dovetails with the anticommutator/commutator split in eq. (5.44) and the phase–scanning protocol of Proposition 5.42. Without BCH these identifications would remain heuristic; with it, they are precise to the stated order $O(\lambda^2)$.

Small-deformation expansion and sector decomposition. Assume a real, compactly supported smearing tensor $J^{\mu\nu}(x)$ and a small parameter $|\lambda| \ll 1$. Define the two history unitaries by

$$\hat{U}_k \;=\; \exp\Big[-i\lambda \int d^d x\, J^{\mu\nu}(x)\, \hat{\Sigma}^{(k)}_{\mu\nu}(x)\Big], \qquad k \in \{0,1\}. \tag{5.51}$$

Introduce the integrated generators

$$S_k \;:=\; \int d^d x\, J^{\mu\nu}(x)\, \hat{\Sigma}^{(k)}_{\mu\nu}(x). \tag{5.52}$$

By the Baker–Campbell–Hausdorff formula (to $O(\lambda^2)$),

$$\hat{U}_1^\dagger \hat{U}_0 \;=\; e^{+\,i\lambda S_1}\, e^{-\,i\lambda S_0} \;=\; \exp\Big(-\,i\lambda\, K + \tfrac{\lambda^2}{2}\,[S_1, S_0]\Big) \;+\; O(\lambda^3), \tag{5.53}$$

with

$$K \;:=\; S_0 - S_1 \qquad \text{(cf. Proposition 5.39)}. \tag{5.54}$$

Expanding the exponential to second order and taking the expectation in $|\Psi_{\mathrm{RR}}\rangle$ yields

$$\langle \hat{U}_1^\dagger \hat{U}_0 \rangle = 1 - \,i\lambda\, \langle K \rangle + \frac{\lambda^2}{2}\, \langle [S_1, S_0] \rangle - \frac{\lambda^2}{2}\, \langle K^2 \rangle + O(\lambda^3), \tag{5.55}$$

where $\langle \cdot \rangle := \langle \Psi_{\mathrm{RR}} | \cdot | \Psi_{\mathrm{RR}} \rangle$. Since $S_{0,1}$ are Hermitian, K and K^2 are Hermitian, while $[S_1, S_0]$ is anti-Hermitian, so $\mathrm{Re}\langle [S_1, S_0] \rangle = 0$ and $\mathrm{Im}\langle [S_1, S_0] \rangle = \langle i[S_1, S_0] \rangle$ is real.

Insert (5.55) into (5.49) and use $\mathrm{Re}[e^{-i\varphi}\alpha] = \cos\varphi\, \mathrm{Re}\,\alpha + \sin\varphi\, \mathrm{Im}\,\alpha$ to obtain, up to $O(\lambda^2)$,

$$p_c(\mathbf{0} \mid \varphi) = \tfrac{1}{2}\Big[1 + \cos\varphi\Big(1 - \tfrac{\lambda^2}{2}\langle K^2 \rangle\Big) + \sin\varphi\Big(-\lambda \langle K \rangle + \tfrac{\lambda^2}{2} \langle i[S_1, S_0] \rangle\Big)\Big] + O(\lambda^3). \tag{5.56}$$

Thus the *even-in-*λ, $\cos\varphi$ component is governed by the symmetric (anticommutator-type) sector via $\langle K^2 \rangle$, while the *odd/phase-odd*, $\sin\varphi$ component is governed by the commutator sector via $\langle i[S_1, S_0] \rangle$, together with a possible linear bias $-\lambda \langle K \rangle$ (which vanishes under the usual centering $\langle K \rangle = 0$ adopted in tomography).

Canonical $\pm$ histories and identification with metric/temporal sectors. Under the canonical choice $\hat{\Sigma}^{(0)}_{\mu\nu} = \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu}$, $\hat{\Sigma}^{(1)}_{\mu\nu} = \hat{g}_{\mu\nu} - \hat{t}_{\mu\nu}$ (eq. (5.42)),

$$K = S_0 - S_1 = 2 \int d^d x\, J^{\mu\nu} \hat{t}_{\mu\nu}, \qquad [S_1, S_0] = 2\,[\int J^{\mu\nu} \hat{g}_{\mu\nu},\ \int J^{\rho\sigma} \hat{t}_{\rho\sigma}],$$

so (5.56) reads

$$p_c(\mathbf{0}\mid\varphi)=\tfrac{1}{2}\Big[1+\cos\varphi\Big(1-2\lambda^2\langle(\int J\,\hat{t})^2\rangle\Big)+\sin\varphi\Big(-2\lambda\langle\int J\,\hat{t}\rangle+\lambda^2\langle i[\int J\,\hat{g},\int J\,\hat{t}]\rangle\Big)\Big]+O(\lambda^3).$$

Hence the fringe contains a *symmetric* (even-in-λ) contribution built from the temporal-sector anticommutator (via $\langle(\int J\,\hat{t})^2\rangle=\frac{1}{2}\langle\{\int J\,\hat{t},\int J\,\hat{t}\}\rangle$) and a *commutator* (phase-odd) contribution $\langle i[\int J\,\hat{g},\int J\,\hat{t}]\rangle$, mirroring the anticommutator/commutator split that appears in the holosymmetric reduction (eq. (5.44)). This establishes the stated identification of visibility with the *relational overlap* $\langle\Psi_{\mathrm{RR}}|\hat{U}_1^\dagger\hat{U}_0|\Psi_{\mathrm{RR}}\rangle$ and its leading decomposition into symmetric vs. commutator sectors.

Together with (5.49), this completes the proof of Proposition 5.36. □

Remark 5.37 ("Operator insertion" variant)**.** If one prefers to use $\hat{\Sigma}^{(\cdot)}_{\mu\nu}$ directly, treat it as a *controlled observable insertion* between unitary segments: $\mathbf{U}^{\mathrm{ins}}_{\mathrm{ctrl}}=|\mathbf{0}\rangle\langle\mathbf{0}|_c\otimes\hat{V}\,\hat{\Sigma}^{(A_GB_N)}\,\hat{W}+|\mathbf{1}\rangle\langle\mathbf{1}|_c\otimes\hat{V}\,\hat{\Sigma}^{(A_NB_G)}\,\hat{W}$, with fixed unitaries $\hat{V},\hat{W}$ (or normalised CPTP instruments) ensuring a proper, norm-preserving map. This yields the same interference functional after suitable normalisation, but the unitary embedding of Lemma 5.35 is the cleanest.

Proposition 5.38 (Holosymmetric visibility = relational overlap)**.** *Under the unitary holosymmetric embedding of Lemma 5.35, prepare* $|+_\varphi\rangle_c=(|\mathbf{0}\rangle_c+e^{i\varphi}|\mathbf{1}\rangle_c)/\sqrt{2}$ *and a register state* $|\Psi_{\mathrm{RR}}\rangle$*. After controlled evolution and standard readout,*

$$p_c(\mathbf{0}\mid\varphi)=\tfrac{1}{2}\Big(1+\mathrm{Re}\Big[e^{-i\varphi}\,\langle\Psi_{\mathrm{RR}}|\hat{U}_1^\dagger\hat{U}_0|\Psi_{\mathrm{RR}}\rangle\Big]\Big).$$

Hence the visibility is $V=\big|\langle\Psi_{\mathrm{RR}}|\hat{U}_1^\dagger\hat{U}_0|\Psi_{\mathrm{RR}}\rangle\big|$.

Proof. **Setup and controlled evolution.** Initialize the control in $|+_\varphi\rangle_c=(|\mathbf{0}\rangle_c+e^{i\varphi}|\mathbf{1}\rangle_c)/\sqrt{2}$ and the registers in $|\Psi_{\mathrm{RR}}\rangle$. Under the unitary holosymmetric embedding (Lemma 5.35) the controlled splitter is $\mathbf{U}_{\mathrm{ctrl}}=|\mathbf{0}\rangle\langle\mathbf{0}|_c\otimes\hat{U}_0+|\mathbf{1}\rangle\langle\mathbf{1}|_c\otimes\hat{U}_1$ (cf. eq. (2.75) and Proposition 4.152). Acting on the input gives

$$\mathbf{U}_{\mathrm{ctrl}}\Big(|+_\varphi\rangle_c\otimes|\Psi_{\mathrm{RR}}\rangle\Big)=\frac{1}{\sqrt{2}}\Big(|\mathbf{0}\rangle_c\otimes\hat{U}_0|\Psi_{\mathrm{RR}}\rangle+e^{i\varphi}\,|\mathbf{1}\rangle_c\otimes\hat{U}_1|\Psi_{\mathrm{RR}}\rangle\Big).\tag{5.57}$$

Recombination and readout. Let R_φ be the "Hadamard-with-phase" recombiner on $\mathcal{H}_c$ such that $R_\varphi|+_\varphi\rangle_c=|\mathbf{0}\rangle_c$ and $R_\varphi|-_\varphi\rangle_c=|\mathbf{1}\rangle_c$ (the standard wordwise readout; cf. Proposition 4.153). Measuring c in $\mathcal{B}_c^{(Z)}$ after applying R_φ is equivalent to projecting (5.57) onto $\langle+_\varphi|_c$ before R_φ. Hence the $|\mathbf{0}\rangle_c$*-amplitude on RR* is

$$\langle+_\varphi|_c\,\mathbf{U}_{\mathrm{ctrl}}\,|+_\varphi\rangle_c\,|\Psi_{\mathrm{RR}}\rangle=\frac{1}{2}\left(\hat{U}_0+e^{-i\varphi}\hat{U}_1\right)|\Psi_{\mathrm{RR}}\rangle,$$

and the corresponding probability is

$$\begin{aligned}p_c(\mathbf{0}\mid\varphi)&=\left\|\frac{1}{2}\left(\hat{U}_0+e^{-i\varphi}\hat{U}_1\right)|\Psi_{\mathrm{RR}}\rangle\right\|^2\\&=\frac{1}{4}\Big(\|\hat{U}_0|\Psi_{\mathrm{RR}}\rangle\|^2+\|\hat{U}_1|\Psi_{\mathrm{RR}}\rangle\|^2+e^{-i\varphi}\langle\Psi_{\mathrm{RR}}|\hat{U}_1^\dagger\hat{U}_0|\Psi_{\mathrm{RR}}\rangle+e^{+i\varphi}\langle\Psi_{\mathrm{RR}}|\hat{U}_0^\dagger\hat{U}_1|\Psi_{\mathrm{RR}}\rangle\Big).\end{aligned}\tag{5.58}$$

By unitarity of $\hat{U}_k$ (Lemma 5.35), $\|\hat{U}_k|\Psi_{\mathrm{RR}}\rangle\| = \|\Psi_{\mathrm{RR}}\rangle\| = 1$. Using $\mathrm{Re}[e^{-i\varphi}\alpha] = \frac{1}{2}\Big(e^{-i\varphi}\alpha + e^{+i\varphi}\alpha^*\Big)$, (5.58) reduces to

$$p_c(\mathbf{0} \mid \varphi) = \tfrac{1}{2}\Big(1 + \mathrm{Re}\Big[e^{-i\varphi}\,\langle\Psi_{\mathrm{RR}}|\hat{U}_1^\dagger\hat{U}_0|\Psi_{\mathrm{RR}}\rangle\Big]\Big), \tag{5.59}$$

which is the stated formula.

Visibility as relational overlap. Write $\langle\Psi_{\mathrm{RR}}|\hat{U}_1^\dagger\hat{U}_0|\Psi_{\mathrm{RR}}\rangle =: A\,e^{i\phi}$ with $A = \Big|\langle\Psi_{\mathrm{RR}}|\hat{U}_1^\dagger\hat{U}_0|\Psi_{\mathrm{RR}}\rangle\Big|$. Then (5.59) becomes

$$p_c(\mathbf{0} \mid \varphi) = \tfrac{1}{2}\Big(1 + A\cos(\varphi - \phi)\Big).$$

Hence $p_{\max} = \frac{1}{2}(1+A)$ and $p_{\min} = \frac{1}{2}(1-A)$, so the (fringe) visibility $V := \frac{p_{\max}-p_{\min}}{p_{\max}+p_{\min}}$ equals $A = \Big|\langle\Psi_{\mathrm{RR}}|\hat{U}_1^\dagger\hat{U}_0|\Psi_{\mathrm{RR}}\rangle\Big|$.

This establishes the proposition. □

Proposition 5.39 (Small-deformation expansion and chronogeometric susceptibility)**.** *Let* $\hat{U}_k = \exp\Big[-i\lambda\int d^dx\, J^{\mu\nu}(x)\,\hat{\Sigma}^{(k)}_{\mu\nu}(x)\Big]$ *with real* $\lambda \ll 1$ *and test tensor* $J^{\mu\nu}$*. Define* $\Delta\hat{\Sigma}_{\mu\nu} := \hat{\Sigma}^{(0)}_{\mu\nu} - \hat{\Sigma}^{(1)}_{\mu\nu}$ *and* $\hat{K} := \int d^dx\, J^{\mu\nu}(x)\,\Delta\hat{\Sigma}_{\mu\nu}(x)$*. Then*

$$\langle\Psi_{\mathrm{RR}}|\hat{U}_1^\dagger\hat{U}_0|\Psi_{\mathrm{RR}}\rangle = 1 - \tfrac{\lambda^2}{2}\,\mathrm{Var}_{\Psi_{\mathrm{RR}}}(\hat{K}) + O(\lambda^3).$$

Consequently, $V = 1 - \frac{\lambda^2}{2}\,\mathrm{Var}_{\Psi_{\mathrm{RR}}}(\hat{K}) + O(\lambda^3)$.

Proof. Set $S_k := \int d^dx\, J^{\mu\nu}(x)\,\hat{\Sigma}^{(k)}_{\mu\nu}(x)$ so that $\hat{U}_k = \exp(-i\lambda S_k)$ with S_k Hermitian (real smearing of Hermitian fields). Define the Hermitian difference $\hat{K} := S_0 - S_1 = \int d^dx\, J^{\mu\nu}(x)\,\Delta\hat{\Sigma}_{\mu\nu}(x)$ as in the statement.

Overlap expansion. Expand the product to second order:

$$\hat{U}_1^\dagger\hat{U}_0 = e^{+i\lambda S_1}\,e^{-i\lambda S_0} = \Big(I + i\lambda S_1 - \tfrac{\lambda^2}{2}S_1^2\Big)\Big(I - i\lambda S_0 - \tfrac{\lambda^2}{2}S_0^2\Big) + O(\lambda^3).$$

Collecting terms gives

$$\hat{U}_1^\dagger\hat{U}_0 = I - i\lambda(S_0 - S_1) - \tfrac{\lambda^2}{2}\Big(S_1^2 + S_0^2 - 2S_1S_0\Big) + O(\lambda^3) = I - i\lambda\,\hat{K} - \tfrac{\lambda^2}{2}\,\hat{K}^2 + O(\lambda^3).$$

Taking the expectation in $|\Psi_{\mathrm{RR}}\rangle$,

$$\langle\Psi_{\mathrm{RR}}|\hat{U}_1^\dagger\hat{U}_0|\Psi_{\mathrm{RR}}\rangle = 1 - i\lambda\,\langle\hat{K}\rangle - \tfrac{\lambda^2}{2}\,\langle\hat{K}^2\rangle + O(\lambda^3), \tag{5.60}$$

with $\langle\cdot\rangle := \langle\Psi_{\mathrm{RR}}|\cdot|\Psi_{\mathrm{RR}}\rangle$.

Centering / phase-locking. Rewrite (5.60) as

$$\langle\hat{U}_1^\dagger\hat{U}_0\rangle = e^{-i\lambda\langle\hat{K}\rangle}\Big(1 - \tfrac{\lambda^2}{2}\,\mathrm{Var}_{\Psi_{\mathrm{RR}}}(\hat{K}) + O(\lambda^3)\Big), \quad \mathrm{Var}_{\Psi_{\mathrm{RR}}}(\hat{K}) := \langle\hat{K}^2\rangle - \langle\hat{K}\rangle^2.$$

Thus, up to the (experimentally removable) overall phase $e^{-i\lambda\langle\hat{K}\rangle}$—which can be absorbed by the control recombiner phase in Proposition 5.38—we obtain

$$\langle\Psi_{\mathrm{RR}}|\hat{U}_1^\dagger\hat{U}_0|\Psi_{\mathrm{RR}}\rangle = 1 - \tfrac{\lambda^2}{2}\,\mathrm{Var}_{\Psi_{\mathrm{RR}}}(\hat{K}) + O(\lambda^3).$$

(Equivalently, one may assume the standard centering $\langle \hat{K} \rangle = 0$, in which case the phase term vanishes.)

Visibility. From Proposition 5.38, $p_c(\mathbf{0} \mid \varphi) = \frac{1}{2}\Big(1 + \mathrm{Re}[e^{-i\varphi}\langle \hat{U}_1^\dagger \hat{U}_0 \rangle]\Big)$ and the fringe visibility is $V = \Big|\langle \hat{U}_1^\dagger \hat{U}_0 \rangle\Big|$. Using the previous expansion,

$$\Big|\langle \hat{U}_1^\dagger \hat{U}_0 \rangle\Big| = \Big| e^{-\,i\lambda\langle \hat{K}\rangle}\Big(1 - \tfrac{\lambda^2}{2}\,\mathrm{Var}(\hat{K}) + O(\lambda^3)\Big)\Big| = 1 - \tfrac{\lambda^2}{2}\,\mathrm{Var}_{\Psi_{\mathrm{RR}}}(\hat{K}) + O(\lambda^3).$$

Therefore $V = 1 - \frac{\lambda^2}{2}\,\mathrm{Var}_{\Psi_{\mathrm{RR}}}(\hat{K}) + O(\lambda^3)$, as claimed.

This proves Proposition 5.39. □

Background: Bures/Fubini–Study geometry and fidelity. For pure states $|\psi\rangle, |\phi\rangle$, the Fubini–Study (FS) angle

$$d_{\mathrm{FS}}(|\psi\rangle, |\phi\rangle) \;=\; \arccos\Big|\langle \psi|\phi\rangle\Big|$$

is the canonical Riemannian distance on projective Hilbert space $\mathbb{CP}^{d-1}$ [407, 287]. For mixed states ρ, σ, the Bures distance is defined from the Uhlmann fidelity

$$F(\rho,\sigma) \;:=\; \Big(\mathrm{Tr}\sqrt{\sqrt{\rho}\,\sigma\,\sqrt{\rho}}\Big)^2 \quad \text{so that} \quad D_{\mathrm{B}}(\rho,\sigma) \;=\; \arccos\sqrt{F(\rho,\sigma)}\,,$$

which reduces on pure states to $F(|\psi\rangle\langle\psi|, |\phi\rangle\langle\phi|) = |\langle\psi|\phi\rangle|^2$ and hence $D_{\mathrm{B}} = \arccos|\langle\psi|\phi\rangle|$ [385, 417, 418, 407, 287]. For a one–parameter family ρ_λ, the Bures metric induces the quantum Fisher information (QFI) via $\mathcal{F}_\lambda = 4\,g_{\mathrm{B}}(\partial_\lambda\rho_\lambda, \partial_\lambda\rho_\lambda)$, and, for pure states evolving unitarily $|\psi_\lambda\rangle = e^{-i\lambda\hat{K}}|\psi_0\rangle$, one has the standard identity $\mathcal{F}_\lambda = 4\,\mathrm{Var}_{\psi_0}(\hat{K})$ [419, 420, 421, 422]. Equivalently, for small parameter shifts $\delta\lambda$, the FS/Bures angle expands as $d_{\mathrm{FS}}\Big(|\psi_\lambda\rangle, |\psi_{\lambda+\delta\lambda}\rangle\Big) = \frac{1}{2}\sqrt{\mathcal{F}_\lambda}\,|\delta\lambda| + O(\delta\lambda^2)$, so that the overlap $|\langle\psi_\lambda|\psi_{\lambda+\delta\lambda}\rangle|$ encodes the local statistical distinguishability [419, 420].

Corollary 5.40 (Law-space (Bures/FS) distance for pure registers)**.** *For pure $|\Psi_{\mathrm{RR}}\rangle$, the Bures (FS) angle*

$$D_{\mathrm{B}} \;:=\; \arccos\Big|\langle \Psi_{\mathrm{RR}}|\hat{U}_1^\dagger \hat{U}_0|\Psi_{\mathrm{RR}}\rangle\Big|$$

obeys, to leading order in the deformation parameter λ of the unitary families,

$$D_{\mathrm{B}} \;=\; \lambda\,\sqrt{\mathrm{Var}_{\Psi_{\mathrm{RR}}}(\hat{K})} + O(\lambda^2),$$

where $\hat{K} = \int d^d x\, J^{\mu\nu}(x)\,\Delta\hat{\Sigma}_{\mu\nu}(x)$ is the generator difference from Proposition 5.39. Equivalently, the (pure-state) quantum Fisher information for parameter λ is $\mathcal{F}_\lambda = 4\,\mathrm{Var}_{\Psi_{\mathrm{RR}}}(\hat{K})$ [419, 420, 407, 287].

Proof. Let $|\Psi_{\mathrm{RR}}\rangle$ be pure and define $S_k := \int d^d x\, J^{\mu\nu}(x)\,\hat{\Sigma}^{(k)}_{\mu\nu}(x)$ and $\hat{K} := S_0 - S_1 = \int d^d x\, J^{\mu\nu}(x)\,\Delta\hat{\Sigma}_{\mu\nu}(x)$. By Proposition 5.39 we have, for $|\lambda| \ll 1$,

$$\langle \Psi_{\mathrm{RR}}|\hat{U}_1^\dagger \hat{U}_0|\Psi_{\mathrm{RR}}\rangle = 1 - \tfrac{\lambda^2}{2}\,\mathrm{Var}_{\Psi_{\mathrm{RR}}}(\hat{K}) + O(\lambda^3). \tag{5.61}$$

Taking the modulus removes any $O(\lambda)$ overall phase, hence

$$\left|\langle\Psi_{\mathrm{RR}}|\hat{U}_1^\dagger\hat{U}_0|\Psi_{\mathrm{RR}}\rangle\right| = 1 - \tfrac{\lambda^2}{2}\,\mathrm{Var}_{\Psi_{\mathrm{RR}}}(\hat{K}) + O(\lambda^3).$$

Write $1-\epsilon := \left|\langle\Psi_{\mathrm{RR}}|\hat{U}_1^\dagger\hat{U}_0|\Psi_{\mathrm{RR}}\rangle\right|$ with $\epsilon = \frac{\lambda^2}{2}\,\mathrm{Var}_{\Psi_{\mathrm{RR}}}(\hat{K}) + O(\lambda^3)$. Using the standard expansion $\arccos(1-\epsilon) = \sqrt{2\epsilon} + O(\epsilon^{3/2})$ as $\epsilon \to 0$, we obtain

$$D_{\mathrm{B}} = \arccos(1-\epsilon) = \sqrt{2\epsilon} + O(\epsilon^{3/2}) = \lambda\sqrt{\mathrm{Var}_{\Psi_{\mathrm{RR}}}(\hat{K})} + O(\lambda^2),$$

which proves the first claim.

For the equivalence with quantum Fisher information (QFI), recall that for a pure-state unitary family $|\Psi_\lambda\rangle = e^{-i\lambda\hat{K}}|\Psi_0\rangle$, the QFI is $\mathcal{F}_\lambda = 4\,\mathrm{Var}_{\Psi_0}(\hat{K})$ (e.g. [419, 420, 407, 287]). Moreover, the local FS/Bures line element satisfies $d_{\mathrm{FS}}(|\Psi_\lambda\rangle, |\Psi_{\lambda+d\lambda}\rangle) = \frac{1}{2}\sqrt{\mathcal{F}_\lambda}\,|d\lambda| + O(d\lambda^2)$, which upon identification of $d\lambda = \lambda$ and $\hat{K}$ as above reproduces exactly $D_{\mathrm{B}} = \lambda\sqrt{\mathrm{Var}_{\Psi_{\mathrm{RR}}}(\hat{K})} + O(\lambda^2)$ and hence $\mathcal{F}_\lambda = 4\,\mathrm{Var}_{\Psi_{\mathrm{RR}}}(\hat{K})$. □

Proposition 5.41 (Gauge covariance and expectation-level invariance (holosymmetric))**.** *If $(\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu}) \mapsto (U\hat{g}_{\mu\nu}U^\dagger, U\hat{t}_{\mu\nu}U^\dagger)$ and $|\overset{\circ}{\mathbf{\Psi}}\rangle \mapsto U|\overset{\circ}{\mathbf{\Psi}}\rangle$, and if $\hat{U}_{0,1}$ are built functorially from these operators, then $\hat{U}_1^\dagger\hat{U}_0 \mapsto U^\dagger(\hat{U}_1^\dagger\hat{U}_0)U$ and $\langle\overset{\circ}{\mathbf{\Psi}}|\hat{U}_1^\dagger\hat{U}_0|\overset{\circ}{\mathbf{\Psi}}\rangle$ is gauge invariant.*

Proof. **Step 1: Functorial covariance of the history unitaries.** By hypothesis, the basic fibre operators transform covariantly, $(\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu}) \mapsto (U\hat{g}_{\mu\nu}U^\dagger,\ U\hat{t}_{\mu\nu}U^\dagger)$ (cf. eq. (5.35)). Since $\hat{U}_{0,1}$ are built *functorially* from $(\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu})$ by compositions that are preserved by the $*$–automorphism $X \mapsto UXU^\dagger$, we have

$$\hat{U}_k \ \longmapsto\ \hat{U}'_k \ :=\ U\,\hat{U}_k\,U^\dagger, \qquad k \in \{0,1\}. \tag{5.62}$$

Step 2: Transformation of the overlap operator. Using (5.62) and $(UXU^\dagger)^\dagger = UX^\dagger U^\dagger$,

$$\hat{U}_1'^\dagger\,\hat{U}'_0 = \left(U\hat{U}_1U^\dagger\right)^\dagger\left(U\hat{U}_0U^\dagger\right) = U\,\hat{U}_1^\dagger\,U^\dagger\,U\,\hat{U}_0\,U^\dagger = U\,(\hat{U}_1^\dagger\hat{U}_0)\,U^\dagger.$$

Equivalently, if one *pulls back* the transformation to compare operators on the original fibre (Heisenberg viewpoint),

$$U^\dagger\left(\hat{U}_1'^\dagger\,\hat{U}'_0\right)U = \hat{U}_1^\dagger\,\hat{U}_0, \tag{5.63}$$

which is the stated mapping $\hat{U}_1^\dagger\hat{U}_0 \mapsto U^\dagger(\hat{U}_1^\dagger\hat{U}_0)U$.

Step 3: Expectation-level invariance. Let the Quantum Holo–State transform as $|\overset{\circ}{\mathbf{\Psi}}\rangle \mapsto U|\overset{\circ}{\mathbf{\Psi}}\rangle$. Then, using the covariant form above,

$$\left\langle\overset{\circ}{\mathbf{\Psi}}{}'\right|\hat{U}_1'^\dagger\,\hat{U}'_0\left|\overset{\circ}{\mathbf{\Psi}}{}'\right\rangle = \left\langle\overset{\circ}{\mathbf{\Psi}}\right|U^\dagger\left(U(\hat{U}_1^\dagger\hat{U}_0)U^\dagger\right)U\left|\overset{\circ}{\mathbf{\Psi}}\right\rangle = \left\langle\overset{\circ}{\mathbf{\Psi}}\right|\hat{U}_1^\dagger\hat{U}_0\left|\overset{\circ}{\mathbf{\Psi}}\right\rangle,$$

by unitarity $U^\dagger U = \mathbb{I}$. Hence the scalar $\langle\overset{\circ}{\mathbf{\Psi}}|\hat{U}_1^\dagger\hat{U}_0|\overset{\circ}{\mathbf{\Psi}}\rangle$ is gauge invariant.

Combining Steps 5.63–5.62 proves Proposition 5.41. □

Proposition 5.42 (Phase-resolved tomography of metric vs. temporal sectors)**.** *Let the holosymmetric reduced operator be* $\hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu}(x;\varphi,\theta)$ *as defined in Equation* (5.44)*. Then*

$$\partial_\varphi \hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu} = -\tfrac{1}{2}\sin\varphi\,\cos\theta\,\{\hat{g}_{\mu\nu},\hat{t}_{\mu\nu}\} + \tfrac{1}{2}\cos\varphi\,\sin\theta\, i[\hat{g}_{\mu\nu},\hat{t}_{\mu\nu}],$$

so scans of φ *isolate anticommutator (metric-like) vs. commutator (phase-curvature) responses.*

Proof. Start from the explicit holosymmetric reduction (eq. (5.44)):

$$\hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu}(x;\varphi,\theta) = \hat{g}_{\mu\nu}(x) + \tfrac{1}{2}\cos\varphi\,\cos\theta\,\{\hat{g}_{\mu\nu}(x),\hat{t}_{\mu\nu}(x)\} + \tfrac{1}{2}\sin\varphi\,\sin\theta\; i[\hat{g}_{\mu\nu}(x),\hat{t}_{\mu\nu}(x)].$$

Since $\hat{g}_{\mu\nu}$ and $\hat{t}_{\mu\nu}$ (hence $\{\cdot,\cdot\}$ and $[\cdot,\cdot]$ built from them) are independent of φ, differentiation acts only on the trigonometric prefactors. Using $\partial_\varphi(\cos\varphi) = -\sin\varphi$ and $\partial_\varphi(\sin\varphi) = \cos\varphi$, we obtain

$$\partial_\varphi \hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu} = -\tfrac{1}{2}\sin\varphi\,\cos\theta\,\{\hat{g}_{\mu\nu},\hat{t}_{\mu\nu}\} + \tfrac{1}{2}\cos\varphi\,\sin\theta\, i[\hat{g}_{\mu\nu},\hat{t}_{\mu\nu}],$$

which is the claimed identity.

For tomography, note the phase-selective specializations: at $\varphi = 0$ one has $\partial_\varphi \hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu}\big|_{\varphi=0} = \frac{1}{2}\sin\theta\, i[\hat{g}_{\mu\nu},\hat{t}_{\mu\nu}]$ (pure commutator response), while at $\varphi = \frac{\pi}{2}$ one has $\partial_\varphi \hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu}\big|_{\varphi=\frac{\pi}{2}} = -\frac{1}{2}\cos\theta\,\{\hat{g}_{\mu\nu},\hat{t}_{\mu\nu}\}$ (pure anticommutator response). Thus scanning φ separates metric-like (anticommutator) from phase-curvature (commutator) contributions, as stated. □

Corollary 5.43 (Sector/commutation limits)**.** *If* $[\hat{g}_{\mu\nu}(x),\hat{t}_{\mu\nu}(x)] = 0$ *on an open set* O*, then on* O $\hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu} = \hat{g}_{\mu\nu} + \frac{1}{2}\cos\varphi\,\cos\theta\,\{\hat{g}_{\mu\nu},\hat{t}_{\mu\nu}\}$ *and all antisymmetric (commutator) contributions vanish. If moreover* $\bar{t}_{\mu\nu} = 0$ *(temporal quiescence), then* $\hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu} = \hat{g}_{\mu\nu}$*.*

Proof. Start from the explicit holosymmetric reduction (eq. (5.44))

$$\hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu}(x;\varphi,\theta) = \hat{g}_{\mu\nu}(x) + \tfrac{1}{2}\cos\varphi\,\cos\theta\,\{\hat{g}_{\mu\nu}(x),\hat{t}_{\mu\nu}(x)\} + \tfrac{1}{2}\sin\varphi\,\sin\theta\, i[\hat{g}_{\mu\nu}(x),\hat{t}_{\mu\nu}(x)]. \quad (5.64)$$

Commutation regime. On the open set O assume $[\hat{g}_{\mu\nu}(x),\hat{t}_{\mu\nu}(x)] = 0$. Then the last (antisymmetric/commutator) term in (5.64) vanishes identically on O, yielding

$$\hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu} = \hat{g}_{\mu\nu} + \tfrac{1}{2}\cos\varphi\,\cos\theta\,\{\hat{g}_{\mu\nu},\hat{t}_{\mu\nu}\} \qquad \text{on } O,$$

which is the first claim. Equivalently, in the commuting case $\{\hat{g}_{\mu\nu},\hat{t}_{\mu\nu}\} = 2\,\hat{g}_{\mu\nu}\hat{t}_{\mu\nu}$, so only the symmetric (metric-like) channel contributes (cf. the discussion around eq. (5.44) and Corollary 5.33).

Temporal quiescence. If, in addition, temporal quiescence holds on O in the strong (operator-level) sense $\hat{t}_{\mu\nu}(x) = 0$, then $\{\hat{g}_{\mu\nu}(x),\hat{t}_{\mu\nu}(x)\} = 0$ there, and the surviving symmetric term also disappears. Hence

$$\hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu}(x;\varphi,\theta) = \hat{g}_{\mu\nu}(x) \qquad \text{on } O.$$

(If one assumes only expectation-level quiescence $\bar{t}_{\mu\nu} = 0$ with nonzero $\hat{t}_{\mu\nu}$, the corresponding *expectation* reduces as in Corollary 5.29, but operator equality need not follow; here we use the operator-level quiescence implicit in the stated conclusion.)

These observations establish Corollary 5.43. □

Proposition 5.44 (Channel-wise holosymmetry and cross-terms)**.** *Suppose the unified operator admits the triadic split* $\hat{\Sigma}_{\mu\nu} = \hat{\Sigma}^{(\mathrm{corr})}_{\mu\nu} + \hat{\Sigma}^{(\mathrm{vac})}_{\mu\nu} + \hat{\Sigma}^{(\mathrm{spec})}_{\mu\nu}$*. Then*

$$\hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu} = \sum_{\alpha} \hat{\Sigma}^{\langle \mathrm{HS}\rangle,\,\alpha}_{\mu\nu} + \tfrac{1}{2}\cos\varphi\,\cos\theta \sum_{\alpha\neq\beta}\{\hat{\Sigma}^{(\alpha)}_{\mu\nu}, \hat{\Sigma}^{(\beta)}_{\mu\nu}\} + \tfrac{1}{2}\sin\varphi\,\sin\theta\, i\sum_{\alpha\neq\beta}[\hat{\Sigma}^{(\alpha)}_{\mu\nu}, \hat{\Sigma}^{(\beta)}_{\mu\nu}],$$

with $\alpha, \beta \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$*. If* $[\hat{\Sigma}^{(\alpha)}_{\mu\nu}, \hat{\Sigma}^{(\beta)}_{\rho\sigma}] = 0$ *for* $\alpha \neq \beta$*, holosymmetry reduces channel-wise:* $\hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu} = \sum_{\alpha} \hat{\Sigma}^{\langle \mathrm{HS}\rangle,\,\alpha}_{\mu\nu}$*.*

Proof. Write the triadic decomposition $\hat{\Sigma}_{\mu\nu} = \sum_{\alpha\in\{\mathrm{corr},\mathrm{vac},\mathrm{spec}\}} \hat{\Sigma}^{(\alpha)}_{\mu\nu}$. Recall the holosymmetric reduced operator in the canonical parametrisation (eq. (5.44)):

$$\hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu} = \hat{g}_{\mu\nu} + \tfrac{1}{2}\cos\varphi\,\cos\theta\,\{\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu}\} + \tfrac{1}{2}\sin\varphi\,\sin\theta\; i[\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu}], \tag{5.65}$$

which is obtained from the functorial, *bilinear* interference block (eq. (5.43)) and the linear law–maps (Definition 5.30 and eq. (5.40)). In particular, the map

$$\mathcal{B}(A, B) \;:=\; \tfrac{1}{2}\cos\varphi\,\cos\theta\,\{A, B\} + \tfrac{1}{2}\sin\varphi\,\sin\theta\, i[A, B]$$

is jointly bilinear and symmetric/antisymmetric in the two slots as dictated by $\{\cdot,\cdot\}$ and $[\cdot,\cdot]$.

Now decompose the unified operator into the three channels and group terms by *within–channel* vs. *cross–channel* contributions. Using bilinearity,

$$\begin{aligned}\mathcal{B}\Big(\sum_{\alpha}\hat{\Sigma}^{(\alpha)}_{\mu\nu},\ \sum_{\beta}\hat{\Sigma}^{(\beta)}_{\mu\nu}\Big) &= \sum_{\alpha,\beta}\ \mathcal{B}\Big(\hat{\Sigma}^{(\alpha)}_{\mu\nu}, \hat{\Sigma}^{(\beta)}_{\mu\nu}\Big)\\ &= \sum_{\alpha}\ \mathcal{B}\Big(\hat{\Sigma}^{(\alpha)}_{\mu\nu}, \hat{\Sigma}^{(\alpha)}_{\mu\nu}\Big)\ + \sum_{\alpha\neq\beta}\ \mathcal{B}\Big(\hat{\Sigma}^{(\alpha)}_{\mu\nu}, \hat{\Sigma}^{(\beta)}_{\mu\nu}\Big).\end{aligned}$$

Define the channel–wise reduced pieces by

$$\hat{\Sigma}^{\langle \mathrm{HS}\rangle,\,\alpha}_{\mu\nu} \;:=\; \hat{\Sigma}^{(\alpha)}_{\mu\nu} + \mathcal{B}\Big(\hat{\Sigma}^{(\alpha)}_{\mu\nu}, \hat{\Sigma}^{(\alpha)}_{\mu\nu}\Big),$$

which is precisely the specialization of (5.65) to the single channel $\hat{\Sigma}^{(\alpha)}_{\mu\nu}$ (linearity in the "law maps" and bilinearity of the interference block; cf. Definition 5.30 and eqs. (5.40) and (5.43)). With this notation, inserting $\hat{\Sigma}_{\mu\nu} = \sum_{\alpha} \hat{\Sigma}^{(\alpha)}_{\mu\nu}$ into the holosymmetric reduction and regrouping yields

$$\hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu} = \sum_{\alpha} \hat{\Sigma}^{\langle \mathrm{HS}\rangle,\,\alpha}_{\mu\nu} + \sum_{\alpha\neq\beta}\ \mathcal{B}\Big(\hat{\Sigma}^{(\alpha)}_{\mu\nu}, \hat{\Sigma}^{(\beta)}_{\mu\nu}\Big).$$

Expanding the cross–channel bilinear block gives exactly

$$\sum_{\alpha\neq\beta}\ \mathcal{B}\Big(\hat{\Sigma}^{(\alpha)}_{\mu\nu}, \hat{\Sigma}^{(\beta)}_{\mu\nu}\Big) = \tfrac{1}{2}\cos\varphi\,\cos\theta \sum_{\alpha\neq\beta}\{\hat{\Sigma}^{(\alpha)}_{\mu\nu}, \hat{\Sigma}^{(\beta)}_{\mu\nu}\} + \tfrac{1}{2}\sin\varphi\,\sin\theta\, i\sum_{\alpha\neq\beta}[\hat{\Sigma}^{(\alpha)}_{\mu\nu}, \hat{\Sigma}^{(\beta)}_{\mu\nu}],$$

which proves the displayed formula in the proposition.

Finally, if all channels mutually commute in the sense that $[\hat{\Sigma}^{(\alpha)}_{\mu\nu}, \hat{\Sigma}^{(\beta)}_{\rho\sigma}] = 0$ for $\alpha \neq \beta$, then in particular $\{\hat{\Sigma}^{(\alpha)}_{\mu\nu}, \hat{\Sigma}^{(\beta)}_{\mu\nu}\} = 0$ and $[\hat{\Sigma}^{(\alpha)}_{\mu\nu}, \hat{\Sigma}^{(\beta)}_{\mu\nu}] = 0$, so all cross–channel terms vanish and one is left with the direct sum of the channel–wise reductions: $\hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu} = \sum_{\alpha} \hat{\Sigma}^{\langle \mathrm{HS}\rangle,\,\alpha}_{\mu\nu}$. □

Lemma 5.45 (Law-decoherence on the control suppresses interference)**.** *Let a phase-damping channel $\mathcal{D}_\eta$ act on the control after the controlled evolution, mapping off-diagonal elements $|\mathbf{0}\rangle\langle\mathbf{1}| \mapsto \eta\,|\mathbf{0}\rangle\langle\mathbf{1}|$ with $\eta \in [0,1]$. Then the outcome becomes*

$$p_c(\mathbf{0} \mid \varphi) = \tfrac{1}{2}\Big(1 + \eta\,\mathrm{Re}\big[e^{-i\varphi}\,\langle\Psi_{\mathrm{RR}}|\hat{U}_1^\dagger \hat{U}_0|\Psi_{\mathrm{RR}}\rangle\big]\Big),$$

so $V \mapsto \eta\, V$.

Proof. Prepare the control in $|+_\varphi\rangle_c = (|\mathbf{0}\rangle_c + e^{i\varphi}|\mathbf{1}\rangle_c)/\sqrt{2}$ and the registers in $|\Psi_{\mathrm{RR}}\rangle$, and apply the controlled unitary $\mathbf{U}_{\mathrm{ctrl}} = |\mathbf{0}\rangle\langle\mathbf{0}|_c \otimes \hat{U}_0 + |\mathbf{1}\rangle\langle\mathbf{1}|_c \otimes \hat{U}_1$ (cf. Lemma 5.35). The post–interaction joint state is the pure density operator

$$\begin{aligned}\rho_1 &= \mathbf{U}_{\mathrm{ctrl}}\Big(|+_\varphi\rangle\langle+_\varphi|_c \otimes |\Psi_{\mathrm{RR}}\rangle\langle\Psi_{\mathrm{RR}}|\Big)\mathbf{U}_{\mathrm{ctrl}}^\dagger \\ &= \tfrac{1}{2}\begin{bmatrix} |\mathbf{0}\rangle\langle\mathbf{0}| \otimes \hat{U}_0|\Psi\rangle\langle\Psi|\hat{U}_0^\dagger \\ + |\mathbf{1}\rangle\langle\mathbf{1}| \otimes \hat{U}_1|\Psi\rangle\langle\Psi|\hat{U}_1^\dagger \\ + e^{-i\varphi}|\mathbf{0}\rangle\langle\mathbf{1}| \otimes \hat{U}_0|\Psi\rangle\langle\Psi|\hat{U}_1^\dagger \\ + e^{i\varphi}|\mathbf{1}\rangle\langle\mathbf{0}| \otimes \hat{U}_1|\Psi\rangle\langle\Psi|\hat{U}_0^\dagger \end{bmatrix},\end{aligned} \tag{5.66}$$

where we abbreviate $|\Psi_{\mathrm{RR}}\rangle \equiv |\Psi\rangle$.

Now act on the control with the phase–damping channel $\mathcal{D}_\eta$ that scales coherences $|\mathbf{0}\rangle\langle\mathbf{1}| \mapsto \eta\,|\mathbf{0}\rangle\langle\mathbf{1}|$, $|\mathbf{1}\rangle\langle\mathbf{0}| \mapsto \eta\,|\mathbf{1}\rangle\langle\mathbf{0}|$ (and fixes the diagonals). The resulting state is

$$\rho_2 = (\mathcal{D}_\eta \otimes \mathrm{id})(\rho_1) = \tfrac{1}{2}\begin{bmatrix} |\mathbf{0}\rangle\langle\mathbf{0}| \otimes \hat{U}_0|\Psi\rangle\langle\Psi|\hat{U}_0^\dagger + |\mathbf{1}\rangle\langle\mathbf{1}| \otimes \hat{U}_1|\Psi\rangle\langle\Psi|\hat{U}_1^\dagger \\ + \eta e^{-i\varphi}|\mathbf{0}\rangle\langle\mathbf{1}| \otimes \hat{U}_0|\Psi\rangle\langle\Psi|\hat{U}_1^\dagger + \eta e^{i\varphi}|\mathbf{1}\rangle\langle\mathbf{0}| \otimes \hat{U}_1|\Psi\rangle\langle\Psi|\hat{U}_0^\dagger \end{bmatrix}.$$

For the standard readout (Hadamard-with-phase inverse on c as in Proposition 5.38), measuring $\mathbf{0}$ in the computational basis is equivalent to projecting *before* the readout onto $|+_\varphi\rangle\langle+_\varphi|_c$:

$$p_c(\mathbf{0} \mid \varphi) = \mathrm{Tr}\Big[\big(|+_\varphi\rangle\langle+_\varphi|_c \otimes I\big)\rho_2\Big].$$

Using $|+_\varphi\rangle\langle+_\varphi| = \tfrac{1}{2}\big(|\mathbf{0}\rangle\langle\mathbf{0}| + |\mathbf{1}\rangle\langle\mathbf{1}| + e^{-i\varphi}|\mathbf{0}\rangle\langle\mathbf{1}| + e^{i\varphi}|\mathbf{1}\rangle\langle\mathbf{0}|\big)$ and tracing out the control gives

$$p_c(\mathbf{0} \mid \varphi) = \tfrac{1}{4}\,\mathrm{Tr}_{\mathrm{RR}}\Big[\hat{U}_0|\Psi\rangle\langle\Psi|\hat{U}_0^\dagger + \hat{U}_1|\Psi\rangle\langle\Psi|\hat{U}_1^\dagger + \eta e^{-i\varphi}\hat{U}_0|\Psi\rangle\langle\Psi|\hat{U}_1^\dagger + \eta e^{i\varphi}\hat{U}_1|\Psi\rangle\langle\Psi|\hat{U}_0^\dagger\Big].$$

Since $\mathrm{Tr}_{\mathrm{RR}}(\hat{U}_k|\Psi\rangle\langle\Psi|\hat{U}_k^\dagger) = 1$ and

$$\mathrm{Tr}_{\mathrm{RR}}\big(\hat{U}_0|\Psi\rangle\langle\Psi|\hat{U}_1^\dagger\big) = \langle\Psi|\hat{U}_1^\dagger\hat{U}_0|\Psi\rangle, \qquad \mathrm{Tr}_{\mathrm{RR}}\big(\hat{U}_1|\Psi\rangle\langle\Psi|\hat{U}_0^\dagger\big) = \langle\Psi|\hat{U}_1^\dagger\hat{U}_0|\Psi\rangle^*,$$

we obtain

$$p_c(\mathbf{0} \mid \varphi) = \tfrac{1}{2}\Big(1 + \eta\,\mathrm{Re}\big[e^{-i\varphi}\,\langle\Psi_{\mathrm{RR}}|\hat{U}_1^\dagger\hat{U}_0|\Psi_{\mathrm{RR}}\rangle\big]\Big),$$

which is the stated formula. Comparing with the $\eta = 1$ (no dephasing) case (cf. Proposition 5.38) shows that the visibility factor $V = \big|\langle\Psi_{\mathrm{RR}}|\hat{U}_1^\dagger\hat{U}_0|\Psi_{\mathrm{RR}}\rangle\big|$ is mapped to ηV. □

Remark 5.46 (Operational black-hole implications (pointer to Chapter 16 Black Hole Physics))**.**

1. **Two knobs = two physical effects.** In the holosymmetric reduction

$$\begin{aligned}\hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu}(x;\varphi,\theta) = \hat{g}_{\mu\nu}(x) \\ + \tfrac{1}{2}\cos\varphi\,\cos\theta\,\{\hat{g}_{\mu\nu}(x),\hat{t}_{\mu\nu}(x)\} \\ + \tfrac{1}{2}\sin\varphi\,\sin\theta\, i[\hat{g}_{\mu\nu}(x),\hat{t}_{\mu\nu}(x)] \quad (eq.\ (5.44)),\end{aligned}$$

   the *anticommutator* term $\{\hat{g},\hat{t}\}$ is the *symmetric* (metric-like) mixing, while the *commutator* term $i[\hat{g},\hat{t}]$ is the *phase-odd* (temporal/phase-curvature) mixing. By Proposition 5.42, a phase scan in φ isolates each sector: $\varphi = 0$ selects the anticommutator channel; $\varphi = \frac{\pi}{2}$ selects the commutator channel.

2. **"Phase curvature modulates temperature/surface gravity."** Near a black-hole horizon, the operational "temperature'' is governed by the surface gravity κ. In this framework, $\hat{t}_{\mu\nu}$ carries the time/phase flow, and the commutator $i[\hat{g}_{\mu\nu},\hat{t}_{\mu\nu}]$ quantifies the failure of temporal flow to commute with the metric response—i.e., a *phase curvature*. Tuning to the pure commutator slice (set $\varphi = \frac{\pi}{2}$) reveals how this non-commutativity modulates the κ-driven channel (and thus the temperature surrogate) without conflating it with metric deformations.

3. **"Anticommutator deforms the near-horizon metric response."** The symmetric channel $\{\hat{g},\hat{t}\}$ acts as a metric-sector deformation (distinguishability curvature). With $\varphi = 0$ one has

$$\hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu}(x;0,\theta) = \hat{g}_{\mu\nu}(x) + \tfrac{1}{2}\cos\theta\,\{\hat{g}_{\mu\nu}(x),\hat{t}_{\mu\nu}(x)\},$$

   so the observable is a near-horizon metric response dressed by the temporal sector but without any phase-odd contribution.

4. **Consistency checks in Kerr specializations.** Inserting the Kerr-sector specializations for $\hat{g}_{\mu\nu}$ and $\hat{t}_{\mu\nu}$ and applying Proposition 5.42 and Corollary 5.43 yields:

   (a) If $[\hat{g},\hat{t}] = 0$ on the horizon patch, the commutator channel vanishes; only the symmetric metric-like deformation remains (no phase-odd modulation of κ).

   (b) If, further, temporal quiescence holds (e.g. $\bar{t}_{\mu\nu} = 0$), then $\hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu} = \hat{g}_{\mu\nu}$ on that patch.

5. **Operational takeaway.** Phase-resolved scans in φ *tomographically separate* (i) modifications of the horizon's effective temperature/surface gravity (pure commutator slice) from (ii) near-horizon metric deformations (pure anticommutator slice), which is precisely the content of Proposition 5.42 together with the commuting/quiescent limits in Corollary 5.43.

Remark 5.47 (Projection pathways: operational guide)**.** There are two natural routes from the expectation-level chronogeometry to an operational metric background:

(i) Lorentzian merge. Build the effective Lorentzian tensor $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ from the symmetric expectation-level chronogeometry (see Section 5.2.6; cf. Definitions 5.25 and 5.26). This route is preferred when the temporal sector is *part of the physics under probe*, e.g. in holosymmetric

tomography where commutator/anticommutator responses of $(\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu})$ are resolved by phase control (Propositions 5.38, 5.57 and 5.60 and Definition 5.56). It preserves the influence of $\bar{t}_{\mu\nu}$ on the effective cones and is the default for dynamical/response analyses.

(ii) Pure-metric projection. Project to the metric sector by dropping $\bar{t}_{\mu\nu}$ at expectation level, i.e. work on $(\mathcal{M}, \bar{g}_{\mu\nu})$ with $\bar{g}_{\mu\nu} := \langle \hat{g}_{\mu\nu} \rangle$ (see Definition 5.26). This route is used for classical limits, comparison with GR/QFT statements, and for causal/spectral results that are anchored to a metric background, notably in the FLRW specialisation (Section 5.4.7). It is also appropriate in regimes of temporal quiescence or symmetry alignment where $\bar{t}_{\mu\nu}$ is negligible or isotropically aligned, in which case the two routes coincide by Corollary 5.29.

Consistency clause. Microcausality and spectral positivity statements are formulated with respect to the effective Lorentzian cones; when $\bar{t}_{\mu\nu}$ is projected out or does not tilt the null structure, $J^{\pm}_{\bar{\Sigma}} = J^{\pm}_{\bar{g}^{\mathrm{Lor}}}$ and both pathways yield the same causal assignment (cf. Sections 5.2.6 and 5.4.7).

5.2.7 The Relational Chronogeometric Operator and Tensor (Holosymmetric–Wordwise)

Definition 5.48 (Wordwise operator–valued relational chronogeometric operator)**.** Let $\hat{g}^{[\mathbf{W}]}_{\mu\nu}(x)$ and $\hat{t}^{[\mathbf{W}]}_{\mu\nu}(x)$ be the operator–valued $(0,2)$ tensor fields constructed in Definitions 5.3 and 5.18, in a fixed holosymmetric word frame $[\mathbf{W}]$. The *wordwise operator–valued relational chronogeometric operator* is

$$\boxed{\hat{\Sigma}^{[\mathbf{W}]}_{\mu\nu}(x) \;:=\; \hat{g}^{[\mathbf{W}]}_{\mu\nu}(x) \;+\; \hat{t}^{[\mathbf{W}]}_{\mu\nu}(x) \;\in\; \mathrm{Herm}\big(\mathcal{H}_{\mathrm{total}}\big),} \tag{5.67}$$

with the following properties:

1. **Hermiticity and symmetry.** Each summand is Hermitian and symmetric in (μ, ν) by construction; hence so is $\hat{\Sigma}^{[\mathbf{W}]}_{\mu\nu}$.
2. **Gauge covariance / word invariance.** For any admissible fibre unitary U_x,
$$\hat{\Sigma}^{[\mathbf{W}']}_{\mu\nu}(x) = U_x^{\dagger}\, \hat{\Sigma}^{[\mathbf{W}]}_{\mu\nu}(x)\, U_x, \qquad \hat{\Sigma}^{[\mathbf{W}]}_{\mu\nu}(x) \;\longmapsto\; U_x^{\dagger}\, \hat{\Sigma}^{[\mathbf{W}]}_{\mu\nu}(x)\, U_x,$$
so the operator is independent of the choice of word frame and transforms covariantly under local gauge.
3. **Spectral localisation (ultra–sub–Planckian support).**
$$\hat{\Sigma}^{[\mathbf{W}]}_{\mu\nu}(x) = P_x\, \hat{\Sigma}^{[\mathbf{W}]}_{\mu\nu}(x)\, P_x, \qquad P_x := E^{(x)[\mathbf{W}]}([0, \lambda_{\mathrm{cutoff}}]),$$
i.e. its range lies inside the spectrally regulated fibre $\mathrm{Ran}\, P_x$ (cf. eq. (4.307)).
4. **SOT regularity.** Near coincidence, $\hat{\Sigma}^{[\mathbf{W}]}_{\mu\nu}$ is obtained as a strong–operator–topology second variation of SOT–C^2 composite functionals (distance/time), uniformly on $\mathrm{Ran}\, P_x$.

Orthogonal split and vanishing cross-terms. By functional orthogonality of the probe subspaces (cf. Section 4.11.1), the mixed second–variation terms between the distance and time sectors vanish at coincidence; hence $\hat{\Sigma}^{[\mathbf{W}]}_{\mu\nu}$ is a *direct sum* of the two sectoral curvatures in the coincidence limit, justifying the additive form in (5.67).

Remark 5.49 (Positivity and calibrated signatures)**.** With canonically positive channel aggregators (squared norms), $\hat{g}^{[\mathbf{W}]}_{\mu\nu} \succeq 0$ and $\hat{t}^{[\mathbf{W}]}_{\mu\nu}$ is Hermitian; thus $\hat{\Sigma}^{[\mathbf{W}]}_{\mu\nu}$ is Hermitian and positive semidefinite on the regulated fibre. If one adopts a Lorentzian calibration at the expectation level (e.g. using $\bar{g}^{\mathrm{Lor}}_{\mu\nu} := \bar{g}_{\mu\nu} + \bar{t}^{\mathrm{sym}}_{\mu\nu}$), the signature is set by the calibration rather than by altering the operator construction.

Expectation–Level Relational Chronogeometric Tensor (Wordwise Form)

Definition 5.50 (Wordwise expectation–level chronogeometric tensor)**.** For the Quantum Holo–State $|\overset{\circ}{\mathbf{\Psi}}\rangle$ (cf. Definition 1.27), define the wordwise dressed sector expectations

$$\bar{g}_{\mu\nu}(x) := \Big\langle \overset{\circ}{\mathbf{\Psi}} \Big| \, \hat{g}^{[\mathbf{W}]}_{\mu\nu}(x) \, \Big| \overset{\circ}{\mathbf{\Psi}} \Big\rangle, \qquad \bar{t}_{\mu\nu}(x) := \Big\langle \overset{\circ}{\mathbf{\Psi}} \Big| \, \hat{t}^{[\mathbf{W}]}_{\mu\nu}(x) \, \Big| \overset{\circ}{\mathbf{\Psi}} \Big\rangle,$$

where the wordwise dressing of the metric/temporal tensors is as in the Chapter 5 wordwise transport/channel formalism (gauge covariance and functoriality summarized in Proposition 4.167). The *expectation–level chronogeometric tensor* is

$$\boxed{\bar{\Sigma}_{\mu\nu}(x) \;:=\; \bar{g}_{\mu\nu}(x) \;+\; \bar{t}_{\mu\nu}(x) \;=\; \Big\langle \overset{\circ}{\mathbf{\Psi}} \Big| \, \hat{\Sigma}^{[\mathbf{W}]}_{\mu\nu}(x) \, \Big| \overset{\circ}{\mathbf{\Psi}} \Big\rangle,} \tag{5.68}$$

with $\hat{\Sigma}^{[\mathbf{W}]}_{\mu\nu} := \hat{g}^{[\mathbf{W}]}_{\mu\nu} + \hat{t}^{[\mathbf{W}]}_{\mu\nu}$ the dressed operator (cf. operator–level Definition 5.25 and its covariance eq. (5.35)). The expression (5.68) is frame/gauge–independent by unitary invariance of the Born expectation (cf. Definition 5.26 and Corollary 5.28) and by the wordwise covariance (cf. the dressing statement in Proposition 4.167 and the holosymmetric lift covariance in Proposition 5.32).

Local quadratic laws and additivity. Let $\Delta x := x' - x$. For SOT–C^2 composite functionals near the diagonal, the distance and temporal squares admit the wordwise Taylor expansions

$$\Big\langle D^{2,[\mathbf{W}]}(x,x') \Big\rangle_{\overset{\circ}{\mathbf{\Psi}}} = \bar{g}_{\mu\nu}(x)\, \Delta x^\mu \Delta x^\nu + \mathcal{O}\Big(|\Delta x|^3\Big),$$
$$\Big\langle T^{2,[\mathbf{W}]}(x,x') \Big\rangle_{\overset{\circ}{\mathbf{\Psi}}} = \bar{t}_{\mu\nu}(x)\, \Delta x^\mu \Delta x^\nu + \mathcal{O}\Big(|\Delta x|^3\Big),$$

consistent with the operator–level definitions of the sectors (cf. Definitions 5.1 and 5.17) and their wordwise dressing (cf. Proposition 4.167). By orthogonality of the fibre split, their sum yields

$$\Big\langle D^{2,[\mathbf{W}]}(x,x') + T^{2,[\mathbf{W}]}(x,x') \Big\rangle_{\overset{\circ}{\mathbf{\Psi}}} = \bar{\Sigma}_{\mu\nu}(x)\, \Delta x^\mu \Delta x^\nu + \mathcal{O}(|\Delta x|^3),$$

with normalization and gauge invariance of the underlying wordwise section ensured by Proposition 4.276 and covariant Leibniz transport by Lemma 4.279.

Triadic bookkeeping and covariance. Both $\hat{\Sigma}^{[\mathbf{W}]}_{\mu\nu}$ and $\bar{\Sigma}_{\mu\nu}$ admit the triadic decomposition inherited from the distance/time channels:

$$\hat{\Sigma}^{[\mathbf{W}]}_{\mu\nu} = \sum_{a\in\{\mathrm{corr,vac,spec}\}} \left(\hat{g}^{[\mathbf{W}],a}_{\mu\nu} + \hat{t}^{[\mathbf{W}],a}_{\mu\nu}\right), \qquad \bar{\Sigma}_{\mu\nu} = \sum_{a} \left(\bar{g}^{a}_{\mu\nu} + \bar{t}^{a}_{\mu\nu}\right),$$

each term transforming covariantly under local frames $U(x)$ (cf. eq. (5.35) and Proposition 5.32). Holosymmetric mixing and channel cross–terms are governed by Proposition 5.44, with sector/commutation limits in Corollary 5.43.

Remark 5.51 (Operational meaning)**.** $\bar{\Sigma}_{\mu\nu}$ is the *total* infinitesimal relational response extracted from the orthogonal fibre split $\mathfrak{F}_x = \mathcal{F}^{\mathrm{dist}}_x \oplus \mathcal{F}^{\mathrm{time}}_x$ (cf. operator–level packaging in Definition 5.25 and expectation–level form in Definition 5.26): the distinguishability curvature $\bar{g}_{\mu\nu}$ ("how far apart") plus the phase–flow curvature $\bar{t}_{\mu\nu}$ ("how out of phase"). It is not a purely Riemannian metric; rather, a unified chronogeometric tensor whose symmetric/antisymmetric sectoral responses can be tomographically isolated by phase scans (cf. Proposition 5.42 and the limits in Corollary 5.33).

Remark 5.52 (Null locus and PQS (wordwise cross–links))**.** At the relational null locus ($\vec{d} = \vec{t} = \mathbf{0}$), both wordwise sector expectations vanish, $\bar{g}_{\mu\nu}(x) = 0 = \bar{t}_{\mu\nu}(x)$, hence $\bar{\Sigma}_{\mu\nu}(x) = 0$ by (5.68). In the wordwise calculus, this statement is *gauge/frame–independent* by the normalization and covariance of the global section (cf. Proposition 4.276) and is *propagated along paths* by parallel constancy (cf. Lemma 4.279). Equivalently, at the null locus the mode weights $w^{[\mathbf{W}]}_{\alpha}(x)$ of the wordwise projectors $P^{[\mathbf{W}]}_{\alpha}(x)$ vanish in concert with the sector tensors (cf. Corollary 4.282). Thus PQS classes coincide with the joint zero locus of the chronogeometric tensor in the coincidence–limit expansion, evaluated in the Quantum Holo–State (cf. Definition 1.27), and this identification is stable under the wordwise gauge/transport structure developed in Chapter 4 (cf. Propositions 4.167 and 4.294).

Remark 5.53 (Twofold reduction and band–limiting)**.** In the duality–collapse regime, all constructions reduce via the isometric identifications to their Twofold counterparts, while spectral localisation P_x enforces endogenous band–limiting consistent with coherent regularisation of the fibre (cf. flatness/holonomy trivialization in Proposition 4.295 and gauge/expectation invariances in Corollary 5.28 and Proposition 5.41). Consequently, $\hat{\Sigma}_{\mu\nu}$ acts ultra–sub–Planckianly on each fibre.

Summary. The holosymmetric wordwise chronogeometric operator $\hat{\Sigma}_{\mu\nu} = \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu}$ packages, at the operator level, the orthogonal infinitesimal structure of the relational bundle (cf. Definition 5.25); its expectation $\bar{\Sigma}_{\mu\nu}$ (cf. Definitions 5.26 and 5.50) provides the classical chronogeometric tensor governing emergent spacetime response, with gauge/word invariances (Corollary 5.28 and Propositions 5.32 and 5.41), SOT regularity and transport (Lemma 4.279), and channel-wise holosymmetric structure (Proposition 5.44 and Corollary 5.33) built in.

5.2.8 Triadic Decomposition of the Relational Chronogeometric Tensor

Having established the Relational Chronogeometric Tensor $\bar{\Sigma}_{\mu\nu}(x)$ as the object encoding the total second-order relational geometry, we now decompose it according to the foundational triadic structure from which it inherits its character. This decomposition provides

a powerful analytical tool to dissect the contributions of the three fundamental relational mechanisms—correlation, vacuum structure, and spectral dynamics—to the overall geometry.

Definition 5.54 (Triadic Decomposition of the Relational Chronogeometric Tensor)**.** Let the expectation-level chronogeometric tensor be given by

$$\bar{\Sigma}_{\mu\nu}(x) := \bar{g}_{\mu\nu}(x) + \bar{t}_{\mu\nu}(x),$$

where both $\bar{g}_{\mu\nu}(x)$ and $\bar{t}_{\mu\nu}(x)$ admit triadic decompositions over the relational modes inherited from their operator-valued progenitors (Definitions 4.599 and 4.600):

$$\bar{g}_{\mu\nu}(x) = \bar{g}^{\text{corr}}_{\mu\nu}(x) + \bar{g}^{\text{vac}}_{\mu\nu}(x) + \bar{g}^{\text{spec}}_{\mu\nu}(x), \tag{5.69}$$

$$\bar{t}_{\mu\nu}(x) = \bar{t}^{\text{corr}}_{\mu\nu}(x) + \bar{t}^{\text{vac}}_{\mu\nu}(x) + \bar{t}^{\text{spec}}_{\mu\nu}(x). \tag{5.70}$$

Then $\bar{\Sigma}_{\mu\nu}(x)$ admits the canonical *triadic decomposition*:

$$\boxed{\bar{\Sigma}_{\mu\nu}(x) := \bar{\Sigma}^{\text{corr}}_{\mu\nu}(x) + \bar{\Sigma}^{\text{vac}}_{\mu\nu}(x) + \bar{\Sigma}^{\text{spec}}_{\mu\nu}(x),} \tag{5.71}$$

where each component is defined by the sum of the corresponding spatial and temporal parts:

$$\boxed{\bar{\Sigma}^{(a)}_{\mu\nu}(x) := \bar{g}^{(a)}_{\mu\nu}(x) + \bar{t}^{(a)}_{\mu\nu}(x), \quad a \in \{\text{corr, vac, spec}\}.} \tag{5.72}$$

We can also write this explicitly:

$$\boxed{\bar{\Sigma}_{\mu\nu}(x) = \sum_{a\in\{\text{corr,vac,spec}\}} \left(\bar{g}^{(a)}_{\mu\nu}(x) + \bar{t}^{(a)}_{\mu\nu}(x)\right) = \sum_{a} \bar{\Sigma}^{(a)}_{\mu\nu}(x).} \tag{5.73}$$

Interpretation of the Triadic Components Each component of this decomposition, $\bar{\Sigma}^{(a)}_{\mu\nu}(x)$, represents a complete "slice" of the relational geometry, capturing both the spatial (distinguishability) curvature and the temporal (phase) curvature associated with a single mode of interaction.

- $\bar{\Sigma}^{\text{corr}}_{\mu\nu}(x)$: This tensor component encodes the total curvature arising purely from *entanglement correlators*. It measures both how direct correlations decay with distance and how their relative quantum phase twists with separation.
- $\bar{\Sigma}^{\text{vac}}_{\mu\nu}(x)$: This component captures the relational geometry induced by the *structure of the vacuum*. It includes both the geometric response to vacuum amplitude gradients and the dynamical response from vacuum phase shifts.
- $\bar{\Sigma}^{\text{spec}}_{\mu\nu}(x)$: This component represents the geometry generated by *spectral dynamics*. It is sourced by the non-commutativity of the foundational operators and includes both the effects of modal dispersion (real part) and temporal spectral flow (imaginary part).

The decomposition of the relational tensors—$\bar{g}_{\mu\nu}(x)$, $\bar{t}_{\mu\nu}(x)$, and their unified sum $\bar{\Sigma}_{\mu\nu}(x)$—into correlational, vacuum, and spectral components is elevated beyond convenient mathematical partition. It expresses a foundational ontological principle of our theory: that all emergent geometry arises from three fundamentally distinct and orthogonal quantum sources. This decomposition partitions the geometry into orthogonal modes of relational information, each with a unique physical meaning and operational behavior.

The *correlational mode* (corr) is the primary channel of pure quantum entanglement and coherence. Derived directly from two-point modal correlator operators like $\hat{G}_F$, it measures how strongly and smoothly the internal structures of different Primordial Quantum Singularities are entangled across the configuration space. Geometrically, this mode contributes to the curvature of distinguishability through what can be understood as *quantum connectivity*—a measure of how tightly different PQSs are coupled via the fundamental field operators. It is the direct representation of the theory's core principle that entanglement constitutes the fabric of geometry.

The *vacuum mode* (vac), in contrast, encodes the geometric consequences of variations in the structure of the quantum vacuum itself. Derived from contrasts in the expectation values of local, diagonal operators, it measures how the background quantum structure—such as the configuration of zero-point modes—changes from one point to another. This mode is sensitive to phenomena such as local symmetry breaking and the landscape of effective potentials. Geometrically, it contributes a form of curvature sourced not by direct interaction, but by intrinsic shifts in the *relational quantum background*.

Finally, the *spectral mode* (spec) is a uniquely quantum-mechanical feature of the geometry, arising from the principle of non-commutativity. It is derived from the commutator between the Spectral Operator and the relational correlators (e.g., $[\hat{D}, \hat{G}_F]$) and measures how the spectrum itself—the distribution and flow of modal frequencies—is deformed across the configuration space. This is not a classical curvature; it is best understood as a *curvature in the space of spectra*, which impacts both emergent distance and time through the mechanisms of modal shear and spectral torsion.

This decomposition of the unified tensor $\bar{\Sigma}_{\mu\nu}(x)$ is a powerful analytical tool, as each component $\bar{\Sigma}^{(a)}_{\mu\nu}(x) = \bar{g}^{(a)}_{\mu\nu}(x) + \bar{t}^{(a)}_{\mu\nu}(x)$ tells us precisely how each relational mechanism contributes to both spatial and temporal curvature. This provides several profound advantages. It gives us the *clarity* to separate the distinct physical sources of geometry and the *interpretability* to analyze which modes dominate in a given physical regime. Furthermore, it provides a *modular* structure, allowing for the development of mode-specific dynamical equations, and offers a clearer path to *quantization* by separating the fundamental observables and their conjugate operators.

In summary, the triadic decomposition reflects a deep structure of the quantum continuum: that all observable relational geometry is composed from just three orthogonal kinds of operator-theoretic relational deviation. It reveals that the total relational tensor $\bar{\Sigma}_{\mu\nu}(x)$ is a synthesis of the fundamental duality between distinguishability and phase flow (from the fiber bundle decomposition $\mathfrak{F}_x = \mathcal{F}_x^{\text{dist}} \oplus \mathcal{F}_x^{\text{time}}$) and the triadic nature of the underlying relational observables. It provides a fully stratified tensor that captures the second-order relational structure in *space*, *phase*, and *mode*—a true modal basis for quantum geometry.

5.2.9 Holosymmetrization of Triadic Channels

By Definition 5.54, the unified chronogeometric operator decomposes as

$$\hat{\Sigma}_{\mu\nu}(x) = \hat{\Sigma}^{(\mathrm{corr})}_{\mu\nu}(x) + \hat{\Sigma}^{(\mathrm{vac})}_{\mu\nu}(x) + \hat{\Sigma}^{(\mathrm{spec})}_{\mu\nu}(x).$$

When convenient, write the corresponding metric and temporal pieces as

$$\hat{g}_{\mu\nu}(x) = \sum_{\alpha\in\{\mathrm{corr,vac,spec}\}} \hat{g}^{(\alpha)}_{\mu\nu}(x), \qquad \hat{t}_{\mu\nu}(x) = \sum_{\alpha\in\{\mathrm{corr,vac,spec}\}} \hat{t}^{(\alpha)}_{\mu\nu}(x),$$

so that $\hat{\Sigma}^{(\alpha)}_{\mu\nu} = \hat{g}^{(\alpha)}_{\mu\nu} + \hat{t}^{(\alpha)}_{\mu\nu}$ and $\hat{\Sigma}_{\mu\nu} = \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu}$.

Lemma 5.55 (Functoriality of the holosymmetric lift over sums)**.** *Let $\widehat{\mathbf{\Sigma}}^{\mathrm{HS}}_{\mu\nu}$ be the holosymmetric lift of Definition 5.30 built functorially (by linear combinations, products, (anti)commutators) from its inputs. Then for any finite sum $\hat{X} = \sum_j \hat{X}_j$, $\widehat{\boldsymbol{X}}^{\mathrm{HS}} = \sum_j \widehat{\boldsymbol{X}}^{\mathrm{HS}}_j$, and the reduced holosymmetric operator obeys the same linearity.*

Definition 5.56 (Canonical law pair for triadic holosymmetry)**.** Adopt the canonical pair that flips the temporal sector uniformly,

$$\hat{\Sigma}^{(0)}_{\mu\nu} = \sum_\alpha \left(\hat{g}^{(\alpha)}_{\mu\nu} + \hat{t}^{(\alpha)}_{\mu\nu}\right), \qquad \hat{\Sigma}^{(1)}_{\mu\nu} = \sum_\alpha \left(\hat{g}^{(\alpha)}_{\mu\nu} - \hat{t}^{(\alpha)}_{\mu\nu}\right),$$

and choose the inter-law coherence operator

$$\hat{\mathbb{C}}^{(\theta)}_{\mu\nu} := \tfrac{1}{2}\Big(\cos\theta\,\{\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu}\} + i\sin\theta\,[\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu}]\Big). \quad \hat{g}_{\mu\nu} := \sum_\alpha \hat{g}^{(\alpha)}_{\mu\nu},\ \hat{t}_{\mu\nu} := \sum_\alpha \hat{t}^{(\alpha)}_{\mu\nu}.$$

Proposition 5.57 (Explicit triadic holosymmetric reduction)**.** *With preparation/readout phase φ on the law-control and the choice in Definition 5.56, the reduced holosymmetric operator is*

$$\boxed{\begin{aligned}\hat{\Sigma}^{\langle\mathrm{HS}\rangle}_{\mu\nu}(x;\varphi,\theta) = \sum_\alpha \left(\hat{g}^{(\alpha)}_{\mu\nu} + \hat{t}^{(\alpha)}_{\mu\nu}\right) + \tfrac{1}{2}\cos\varphi\,\cos\theta \sum_{\alpha,\beta}\left\{\hat{g}^{(\alpha)}_{\mu\nu}, \hat{t}^{(\beta)}_{\mu\nu}\right\} \\ + \tfrac{1}{2}\sin\varphi\,\sin\theta\, i \sum_{\alpha,\beta}\left[\hat{g}^{(\alpha)}_{\mu\nu}, \hat{t}^{(\beta)}_{\mu\nu}\right].\end{aligned}} \tag{5.74}$$

Proof. Insert the shorthands $\hat{g} = \sum_\alpha \hat{g}^{(\alpha)}$, $\hat{t} = \sum_\alpha \hat{t}^{(\alpha)}$ into the single-sector formula $\hat{\Sigma}^{\langle\mathrm{HS}\rangle} = \hat{g} + \tfrac{1}{2}\cos\varphi\cos\theta\{\hat{g},\hat{t}\} + \tfrac{1}{2}\sin\varphi\sin\theta\, i[\hat{g},\hat{t}]$ (from the holosymmetric reduction with Definition 5.30), and expand the (anti)commutators bilinearly. □

Corollary 5.58 (Channel-orthogonality reduction)**.** *If there exist channel projectors $P^{(\alpha)}$ with $P^{(\alpha)}P^{(\beta)} = \delta_{\alpha\beta}P^{(\alpha)}$ such that $\hat{g}^{(\alpha)}_{\mu\nu} = P^{(\alpha)}\hat{g}_{\mu\nu}P^{(\alpha)}$, $\hat{t}^{(\alpha)}_{\mu\nu} = P^{(\alpha)}\hat{t}_{\mu\nu}P^{(\alpha)}$ and $[P^{(\alpha)}, P^{(\beta)}] = 0$, then cross-channel terms vanish:*

$$\left\{\hat{g}^{(\alpha)}_{\mu\nu}, \hat{t}^{(\beta)}_{\mu\nu}\right\} = \left[\hat{g}^{(\alpha)}_{\mu\nu}, \hat{t}^{(\beta)}_{\mu\nu}\right] = 0 \quad (\alpha \neq \beta),$$

and hence

$$\hat{\Sigma}^{\langle\mathrm{HS}\rangle}_{\mu\nu} = \sum_\alpha \left(\hat{g}^{(\alpha)}_{\mu\nu} + \hat{t}^{(\alpha)}_{\mu\nu} + \tfrac{1}{2}\cos\varphi\,\cos\theta\,\{\hat{g}^{(\alpha)}_{\mu\nu}, \hat{t}^{(\alpha)}_{\mu\nu}\} + \tfrac{1}{2}\sin\varphi\,\sin\theta\, i[\hat{g}^{(\alpha)}_{\mu\nu}, \hat{t}^{(\alpha)}_{\mu\nu}]\right).$$

Corollary 5.59 (Expectation-level triadic holosymmetry)**.** *For the Quantum Holo–State* $|\mathring{\mathbf{\Psi}}\rangle$*,*

$$\bar{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu}(x;\varphi,\theta) = \sum_{\alpha}\left(\bar{g}^{(\alpha)}_{\mu\nu} + \bar{t}^{(\alpha)}_{\mu\nu}\right) + \tfrac{1}{2}\cos\varphi\,\cos\theta \sum_{\alpha,\beta}\left\langle \{\hat{g}^{(\alpha)}_{\mu\nu}, \hat{t}^{(\beta)}_{\mu\nu}\}\right\rangle + \tfrac{1}{2}\sin\varphi\,\sin\theta\, i \sum_{\alpha,\beta}\left\langle [\hat{g}^{(\alpha)}_{\mu\nu}, \hat{t}^{(\beta)}_{\mu\nu}]\right\rangle,$$

with $\langle\cdot\rangle := \langle\mathring{\mathbf{\Psi}}|\cdot|\mathring{\mathbf{\Psi}}\rangle$ *and* $\bar{g}^{(\alpha)}_{\mu\nu} = \langle\hat{g}^{(\alpha)}_{\mu\nu}\rangle$*,* $\bar{t}^{(\alpha)}_{\mu\nu} = \langle\hat{t}^{(\alpha)}_{\mu\nu}\rangle$*. Gauge covariance of operators and joint transformation of* $|\mathring{\mathbf{\Psi}}\rangle$ *imply gauge invariance of the RHS.*

Proposition 5.60 (Phase-resolved channel tomography)**.** *Differentiating eq.* (5.74) *with respect to the control phase yields*

$$\partial_\varphi \hat{\Sigma}^{\langle \mathrm{HS}\rangle}_{\mu\nu} = -\tfrac{1}{2}\sin\varphi\,\cos\theta \sum_{\alpha,\beta}\{\hat{g}^{(\alpha)}_{\mu\nu}, \hat{t}^{(\beta)}_{\mu\nu}\} + \tfrac{1}{2}\cos\varphi\,\sin\theta\, i \sum_{\alpha,\beta}[\hat{g}^{(\alpha)}_{\mu\nu}, \hat{t}^{(\beta)}_{\mu\nu}],$$

so phase scans isolate, respectively, symmetric (anticommutator) and antisymmetric (commutator) cross-channel couplings.

Remark 5.61 (Channel-selective law flips (optional refinement))**.** One may generalise Definition 5.56 by flipping the temporal sign per channel via a vector $\mathbf{s} = (s_{\mathrm{corr}}, s_{\mathrm{vac}}, s_{\mathrm{spec}}) \in \{\pm 1\}^3$: $\hat{\Sigma}^{(1)}_{\mu\nu} = \sum_\alpha(\hat{g}^{(\alpha)}_{\mu\nu} + s_\alpha \hat{t}^{(\alpha)}_{\mu\nu})$. The analogue of eq. (5.74) then weights each pair (α,β) by s_β inside the (anti)commutator sums, enabling channel-resolved law-phase control.

Remark 5.62 (Operational time reversals under holosymmetry)**.** *What is permitted.* Using the canonical temporal sign flip of Definition 5.56, $\hat{\Sigma}^{(1)}_{\mu\nu} = \hat{g}_{\mu\nu} - \hat{t}_{\mu\nu}$, and constructing smeared self-adjoint generators as in the unitary embedding, the resulting branch unitaries U_0 and U_1 can be arranged so that $U_1 \simeq U_0^\dagger$ (backward evolution). The holosymmetric readout then realises a Loschmidt-echo probe of time reversal: the visibility $V = \left|\langle\Psi_{\mathrm{RR}}|U_1^\dagger U_0|\Psi_{\mathrm{RR}}\rangle\right|$ and its phase follow Proposition 5.38, thereby quantifying reversibility and exposing T-odd structure in the dynamics. In the triadic split, channel-selective reversals are admissible: choosing a sign vector $\mathbf{s} = (s_{\mathrm{corr}}, s_{\mathrm{vac}}, s_{\mathrm{spec}}) \in \{\pm 1\}^3$ (cf. Remark 5.61) produces superpositions of partially time-reversed chronogeometries; holosymmetric tomography (Proposition 5.60) then separates T-even contributions (anticommutators) from T-odd contributions (commutators) with spatial and channel localisation set by the smearing tensor.

The caveat. The fundamental time-reversal operator T is antiunitary and cannot itself appear as a controlled branch in a unitary circuit. The holosymmetric map does not directly implement T; rather, it permits controlled superpositions of unitaries that are T-conjugate ($U_1 = \mathsf{T}U_0\mathsf{T}^{-1}$) or the forward–backward pair U_0 vs. $U_0^\dagger$, both physically realisable as unitary histories. Consequently, the framework supports operational time reversals (reversed evolution and T-conjugate dynamics) and renders their signatures observable, while remaining consistent with unitarity and gauge covariance.

Spacetime is treated not as a fixed background but as an object that can be placed on a quantum balance. Holosymmetry prepares two chronogeometric histories—two candidate spacetimes, each with its own geometry and temporal flow—and interferes them under a controllable phase [287, 3, 23, 24]. The readout on the control qubit functions as a direct meter of similarity between these spacetimes: high visibility indicates relational proximity,

while low visibility signals divergence. In the small-difference limit, the visibility reduces to a Fisher-information–type susceptibility, furnishing a principled "distance on law space" that quantifies sensitivity to perturbations of the chronogeometric operator [359, 423, 407].

The interference pattern is not merely binary. By sweeping the control phase, mixing between the metric sector $\hat{g}$ ("how far apart") and the temporal sector $\hat{t}$ ("how out of phase") can be separated. The symmetric (anticommutator) contribution deforms expectation-level geometry, whereas the antisymmetric (commutator) contribution exposes phase curvature—transport effects associated with time flow and Berry-type structure [350, 25, 24]. In this way, the protocol implements tomography: experimental knobs on the control register select which aspect of chronogeometry is probed [424, 408]. Resolving $\hat{\Sigma}$ into triadic channels—correlational, vacuum, spectral—preserves this structure: intra-channel interference appears alongside explicit cross-channel couplings, while commuting or projector-orthogonal channels decouple, enabling sector-wise tests. Operationally, the result is a programmable interferometer over "the laws": temporal signs may be flipped per channel, probes localized with a smearing tensor, and each sector's contribution to emergent spacetime read out. In horizon applications, commutator-driven terms map to temperature/surface-gravity–like responses, whereas anticommutator terms reshape the near-horizon geometry, providing tunable levers for interrogating black-hole chronogeometry [244, 11, 86, 87, 425].

Within this holosymmetric setting, operational time reversals are admissible and informative. By pairing a history with its backward counterpart—implemented as the forward–backward unitary pair U_0 vs. $U_0^\dagger$ or, more generally, T-conjugate unitaries—the interference signal becomes a Loschmidt-echo–type probe of reversibility, with visibility and phase serving as calibrated witnesses of T-even (anticommutator) versus T-odd (commutator) structure [426, 427]. The small-deformation expansion ties echo decay to the variance of the chronogeometric deformation, defining an information-geometric measure of the arrow of time and yielding quantitative reversibility bounds in law space [359, 423, 407]. Channel-selective flips ($s_\alpha \in \{\pm 1\}$) then enable partial time reversals confined to correlational, vacuum, or spectral sectors, localised further by the smearing tensor; channels that commute contribute no T-odd signal. Because the construction is built from smeared self-adjoint generators, locality and gauge covariance are maintained [255], while environmental dephasing provides a clear diagnostic separation between intrinsic T-odd structure and extrinsic irreversibility—uniformly damping visibility without mimicking phase-selective T-odd signatures [380, 247]. A final foundational constraint remains: the fundamental time-reversal operator T is antiunitary and cannot itself be a controlled branch of a unitary circuit; holosymmetry therefore compares unitary histories related by reversal or T-conjugation rather than applying T directly [350, 287].

5.2.10 From Ultra-Sub-Planckian Operators to Macroscopic Geometry

A foundational distinction in IQR is between the *operator-valued* chronogeometric tensors and their *expectation-level* (classical) counterparts. The operators $\hat{g}_{\mu\nu}(x)$ and $\hat{t}_{\mu\nu}(x)$ are the genuine quantum-geometric objects introduced in Definitions 5.25 and 5.26. By construction—as second Fréchet variations of relational observables—they are supported on *ultra-sub-Planckian* fibres tied to a single Primordial Quantum Singularity (PQS), with coherence width ℓ_{PQS} (Lemma 4.602). Operationally, they probe the pre-geometric, relational core and vanish

identically outside the PQS fibre.

Emergence via the Quantum Holo–State. The smooth, macroscopic fields arise by evaluating these operators in the global, non-factorisable Quantum Holo–State $|\mathring{\mathbf{\Psi}}\rangle$:

$$\bar{g}_{\mu\nu}(x) \;:=\; \Big\langle \mathring{\mathbf{\Psi}} \Big|\, \hat{g}_{\mu\nu}(x) \,\Big| \mathring{\mathbf{\Psi}} \Big\rangle, \qquad \bar{t}_{\mu\nu}(x) \;:=\; \Big\langle \mathring{\mathbf{\Psi}} \Big|\, \hat{t}_{\mu\nu}(x) \,\Big| \mathring{\mathbf{\Psi}} \Big\rangle.$$

These expectation values act as *state-smoothing projections* of the ultra-fine operator geometry onto classical tensor fields on the emergent manifold $\mathcal{X}_q$ (cf. Section 5.2.6). When the state satisfies the usual regularity/positivity conditions (Hadamard/adiabatic or KMS where appropriate), expectation values of local composite operators are well-defined and inherit the differentiability required for geometric interpretation [244, 11, 230, 255]. In particular, the effective Lorentzian metric

$$\bar{g}^{\mathrm{Lor}}_{\mu\nu}(x) \;:=\; \bar{g}_{\mu\nu}(x) + \bar{t}^{\mathrm{sym}}_{\mu\nu}(x), \qquad \bar{t}^{\mathrm{sym}}_{\mu\nu} := \tfrac{1}{2}(\bar{t}_{\mu\nu} + \bar{t}_{\nu\mu}),$$

constructed in Definition 5.122, supplies the kinematic light-cone structure used throughout Sections 5.4.5 and 5.4.7.

Analogy with standard QFT. This micro-to-macro passage mirrors the familiar distinction between local field operators and their classical densities: the electron field $\hat{\psi}(x)$ is an operator-valued distribution, while $\rho(x) = e\,\langle\Psi|\hat{\psi}^\dagger(x)\hat{\psi}(x)|\Psi\rangle$ is a smooth, macroscopic charge density in suitable states [14, Ch. 6]. Likewise in IQR, $\hat{g}_{\mu\nu}(x)$ and $\hat{t}_{\mu\nu}(x)$ are ultra-local operator fields, whereas $\bar{g}_{\mu\nu}(x)$ and $\bar{t}_{\mu\nu}(x)$ are their smooth classical avatars that can vary coherently on macroscopic scales.

Remark 5.63 (Emergence of macroscopic tensors from ultra-sub-Planckian operators)**.** The operator-valued tensors $\hat{g}_{\mu\nu}(x)$ and $\hat{t}_{\mu\nu}(x)$ possess ultra-sub-Planckian support determined by the PQS fibre (Lemma 4.602). Their expectation values in the global Quantum Holo–State $|\mathring{\mathbf{\Psi}}\rangle$, $\bar{g}_{\mu\nu}(x)$ and $\bar{t}_{\mu\nu}(x)$, are smooth classical fields on $\mathcal{X}_q$ (cf. Definition 5.26). The global coherence of $|\mathring{\mathbf{\Psi}}\rangle$ functions as a *smoothing mechanism*, transporting ultra-local relational information into macroscopic geometry. In the geometric limit this mechanism yields the effective cones of $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ (light-cone kinematics) alongside the intra-cone phase transport governed by $\bar{t}_{\mu\nu}$ (Definitions 5.122 and 5.150), thereby recovering the GR/QFT regime as a special case [7, 244, 11].

Holosymmetric perspective. When holosymmetry is engaged, the operator-to-expectation map is applied *branchwise*, producing $\{\bar{g}^{\mathrm{Lor}}_{(k)}, \bar{t}_{(k)}\}_k$ that define the family of effective cones and temporal textures used in the intersection/union causal calculus (Definition 5.123, Proposition 5.124, and Corollary 5.126). The emergence mechanism itself is unchanged; holosymmetry adds *interferometric readout*, not a different route from micro to macro.

5.3 Temporal and Geometric Manifolds

5.3.1 The Operator-Level Relational Geometric Manifold

Before the emergence of a classical spacetime that is accessible to observers, there must exist a more fundamental, pre-geometric structure that encodes the raw, quantum-relational

information of the universe. This structure is the *Operator-Level Relational Geometric Manifold*, denoted $\hat{\mathcal{M}}$. It is not a classical space, but rather the quantum scaffolding from which classical reality is projected.

Definition 5.64 (Operator-Level Relational Geometric Manifold)**.** The *Operator-Level Relational Geometric Manifold* is the pre-geometric, quantum-curvature structure given by the triplet:

$$\boxed{\hat{\mathcal{M}} := \left(\mathcal{X}_q,\ \hat{g}_{\mu\nu}(x),\ \hat{t}_{\mu\nu}(x)\right)} \tag{5.75}$$

This definition synthesizes three core components of the theory into a single, foundational object:

1. *The Base Manifold, $\mathcal{X}_q$:* The structure is built upon the emergent quantum configuration space (Section 4.19.3), a smooth manifold whose points label the fundamental, distinguishable Primordial Quantum Singularities (PQSs). While itself emergent from the deeper $\mathcal{C}_d$ continuum, $\mathcal{X}_q$ serves as the minimal arena upon which operator-valued fields can be charted and their relations defined.

2. *The Operator-Valued Metric Tensor, $\hat{g}_{\mu\nu}(x)$:* This is a Hermitian, positive semi-definite operator field that endows $\hat{\mathcal{M}}$ with its notion of distinguishability. As defined in Definition 5.1, it arises from the second variation of the composite triadic distance. Each component, $\hat{g}_{\mu\nu}(x)$, is a quantum operator acting on $\mathcal{H}_{\text{total}}$ that encodes the infinitesimal relational distinguishability between nearby PQSs. This operator has ultra-local support, confined to the coherence width of the PQS, $\text{supp}(\hat{g}_{\mu\nu}(x)) \subseteq \mathcal{O}(\ell_{\text{PQS}})$.

3. *The Operator-Valued Temporal Curvature Tensor, $\hat{t}_{\mu\nu}(x)$:* This tensor is the novel, temporal counterpart to the metric. Mirroring the construction of $\hat{g}_{\mu\nu}(x)$ (Definition 5.17), it probes the phase dynamics of the system. It is a quantum operator that measures the anisotropies and curvature in the flow of quantum phase, providing a temporal structure completely absent in classical General Relativity. Like the metric, its support is confined to the PQS coherence scale.

Interpretation and Ontological Role The structure $\hat{\mathcal{M}} = (\mathcal{X}_q,\ \hat{g}_{\mu\nu},\ \hat{t}_{\mu\nu})$ does not describe a classical spacetime manifold. It is the pre-expectation, purely quantum substrate that captures the *noncommutative curvature of distinguishability* and the *phase-dynamic structure of quantum evolution*. It represents the pre-metric topology of the quantum world—the raw material from which all subsequent emergent structures, including classical distance and proper time, are derived.

Remark 5.65 (Relational Phase-Geometry vs. Metric Geometry)**.** It is more accurate to understand the structure $\hat{\mathcal{M}}$ as a *relational phase-geometry*, a concept that transcends a simple metric geometry. Its metric and temporal components reflect operator-level curvature of distinguishability and phase flow, respectively. Only when evaluated in the Quantum Holo-State, $|\mathring{\mathbf{\Psi}}\rangle$, do these quantum operators yield the classical tensor fields, $\bar{g}_{\mu\nu}(x)$ and $\bar{t}_{\mu\nu}(x)$, that define the distances, times, and causal structure of the observable universe.

5.3.2 The Expectation-Level Relational Manifold

Once the operator-level tensors $\hat{g}_{\mu\nu}(x)$ and $\hat{t}_{\mu\nu}(x)$ are projected into the Quantum Holo-State $|\overset{\circ}{\mathbf{\Psi}}\rangle$, they yield scalar-valued fields over the emergent quantum configuration space. The resulting manifold captures the full classical relational structure of both spatial distinguishability and quantum phase flow.

Definition 5.66 (Expectation-Level Relational Manifold)**.** The *Expectation-Level Relational Manifold* is the classical manifold formed by the expectation values of the operator-valued tensors:

$$\boxed{\mathcal{M}_{\text{rel}} := \Big(\mathcal{X}_q,\ \bar{g}_{\mu\nu}(x),\ \bar{t}_{\mu\nu}(x)\Big)\,,} \tag{5.76}$$

where:

- $\mathcal{X}_q$ is the emergent quantum configuration space (Section 4.19.3),
- $\bar{g}_{\mu\nu}(x) := \langle\overset{\circ}{\mathbf{\Psi}}|\hat{g}_{\mu\nu}(x)|\overset{\circ}{\mathbf{\Psi}}\rangle$ is the expectation-level metric tensor (Definition 5.10),
- $\bar{t}_{\mu\nu}(x) := \langle\overset{\circ}{\mathbf{\Psi}}|\hat{t}_{\mu\nu}(x)|\overset{\circ}{\mathbf{\Psi}}\rangle$ is the expectation-level temporal curvature tensor (Definition 5.22).

Interpretation and Role in the Emergence Chain The manifold $\mathcal{M}_{\text{rel}}$ is the first fully classical relational structure that encodes quantum-derived geometry. It exists one layer above the operator-level quantum manifold $\hat{\mathcal{M}}$, and one layer below the coarse-grained classical manifold $\mathcal{M}_{\text{geom}}$. It provides the full classical data from which proper time, causal direction, and distinguishability are derived — but retains both spatial and temporal curvature structures as independent fields.

5.3.3 Operator-Level Relational Geometric Manifold (wordwise form)

Definition 5.67 (Holosymmetric operator–word relational manifold)**.** Fix the device channel Dev with Hilbert space $\mathcal{H}_{\text{total}}$ and a word basis $\mathbf{W}$ for the Fourfold dialectic at each locus $x \in \mathcal{X}_q$. Let

$$P_x := E^{(x)[\mathbf{W}]}\Big([0, \lambda_{\text{cutoff}}]\Big)$$

be the spectral projector implementing ultra–sub–Planckian coherence (cf. eq. (4.307)). Define the *holosymmetric operator–word manifold* as the word–equivalence class

$$\boxed{\widehat{\mathbb{M}}^{\text{HS}} := \Big[\ \Big(\mathcal{X}_q;\ \hat{g}^{[\mathbf{W}]}_{\mu\nu}(x),\ \hat{t}^{[\mathbf{W}]}_{\mu\nu}(x)\Big)\ \Big]_{[\mathbf{W}]},} \tag{5.77}$$

where for each x:

(a) $\hat{g}^{[\mathbf{W}]}_{\mu\nu}(x),\ \hat{t}^{[\mathbf{W}]}_{\mu\nu}(x) \in \text{Herm}(\mathcal{H}_{\text{total}})$ are device–word tensors obtained as SOT–Fréchet second variations of the regulated composite distance/time functionals (cf. Definitions 5.3 and 5.18), and are fibre–local:

$$\hat{g}^{[\mathbf{W}]}_{\mu\nu}(x) = P_x\,\hat{g}^{[\mathbf{W}]}_{\mu\nu}(x)\,P_x, \qquad \hat{t}^{[\mathbf{W}]}_{\mu\nu}(x) = P_x\,\hat{t}^{[\mathbf{W}]}_{\mu\nu}(x)\,P_x.$$

(b) **Word/frame change.** For any charge–preserving fibre unitary U_x (admissible word change $\mathbf{W}\to\mathbf{W}'$),

$$\hat g_{\mu\nu}^{[\mathbf{W}']}(x)=U_x^\dagger \hat g_{\mu\nu}^{[\mathbf{W}]}(x)U_x,\qquad \hat t_{\mu\nu}^{[\mathbf{W}']}(x)=U_x^\dagger \hat t_{\mu\nu}^{[\mathbf{W}]}(x)U_x,\qquad P_x\mapsto U_x^\dagger P_x U_x.$$

The manifold is the *word class* under this equivalence.

(c) **Transport compliance.** Along any gauge–covariant transporter $\mathcal{U}^{[\mathbf{W},\mathbf{W}]}(x\leftarrow x';\gamma)$ the fields respect pullback/pushforward in the device channel, ensuring SOT–C^2 regularity near the diagonal (cf. Section 4.19.2).

Remark 5.68 (Chronogeometric word operator)**.** Set the unified device–word tensor

$$\hat\Sigma_{\mu\nu}^{[\mathbf{W}]}(x):=\hat g_{\mu\nu}^{[\mathbf{W}]}(x)+\hat t_{\mu\nu}^{[\mathbf{W}]}(x),$$

with the same word/gauge covariance as its summands (cf. Definition 5.48). All statements below may be expressed equivalently in terms of $\hat\Sigma_{\mu\nu}^{[\mathbf{W}]}$.

Proposition 5.69 (Word invariance, gauge covariance, and locality)**.** *The class $\widehat{\mathbb{M}}^{\mathrm{HS}}$ is: (i) word–invariant (independent of $[\mathbf{W}]$) by conjugation equivalence; (ii) gauge–covariant under local frame rotations; (iii) ultra–local on the PQS coherence band via P_x; and (iv) SOT–regular (second–differentiable) uniformly on* $\operatorname{Ran}P_x$.

Expectation-Level Relational Manifold (wordwise reduction)

Definition 5.70 (Holosymmetric expectation–word manifold)**.** Let $\rho_\circ:=|\mathring{\mathbf{\Psi}}\rangle\langle\mathring{\mathbf{\Psi}}|$ be the endogenous Holo–State (word–invariant). The *expectation–level relational manifold* is the reduction

$$\boxed{\mathbb{M}_{\mathrm{rel}}^{\langle\mathrm{HS}\rangle}:=\Big(\ \mathcal{X}_q;\ \bar g_{\mu\nu}(x),\ \bar t_{\mu\nu}(x)\ \Big),\qquad \bar g_{\mu\nu}(x):=\operatorname{Tr}\Big(\rho_\circ\,\hat g_{\mu\nu}^{[\mathbf{W}]}(x)\Big),\ \bar t_{\mu\nu}(x):=\operatorname{Tr}\Big(\rho_\circ\,\hat t_{\mu\nu}^{[\mathbf{W}]}(x)\Big).}\tag{5.78}$$

Equivalently, $\bar\Sigma_{\mu\nu}(x):=\operatorname{Tr}\Big(\rho_\circ\,\hat\Sigma_{\mu\nu}^{[\mathbf{W}]}(x)\Big)=\bar g_{\mu\nu}(x)+\bar t_{\mu\nu}(x)$.

Proposition 5.71 (Word/frame independence of the reduction)**.** *Under any fibre word/frame change U_x,*

$$\bar g_{\mu\nu}(x)\mapsto \operatorname{Tr}\Big(U_x\rho_\circ U_x^\dagger\,U_x^\dagger\hat g_{\mu\nu}^{[\mathbf{W}]}(x)U_x\Big)=\operatorname{Tr}\Big(\rho_\circ\,\hat g_{\mu\nu}^{[\mathbf{W}]}(x)\Big)=\bar g_{\mu\nu}(x),$$

and likewise for $\bar t_{\mu\nu}$ and $\bar\Sigma_{\mu\nu}$. Hence $\mathbb{M}_{\mathrm{rel}}^{\langle\mathrm{HS}\rangle}$ is word– and gauge–independent.

Remark 5.72 (Position in the emergence chain)**.** $\widehat{\mathbb{M}}^{\mathrm{HS}}$ is the pre-geometric *operator–word* scaffold; $\mathbb{M}_{\mathrm{rel}}^{\langle\mathrm{HS}\rangle}$ is its *classical word-reduction*, furnishing the expectation-level metric $\bar g$ and temporal curvature $\bar t$. A further projection to a pure-metric manifold $(\mathcal{X}_q,\bar g_{\mu\nu})$ is achieved by dropping the temporal sector when appropriate (temporal quiescence or calibration choice), cf. Definition 5.50.

Definition 5.73 (Canonical law–control word on the manifold fields)**.** Let the law–control register be

$$\mathcal{H}_L = \mathrm{span}\{\,|0\rangle_L, |1\rangle_L\,\}$$

and let $\big(\hat{g}_{\mu\nu}(x), \hat{t}_{\mu\nu}(x)\big)$ be the operator–valued tensors on $\mathcal{X}_q$. Introduce the Hermitian mixers

$$\boxed{\hat{S}_{\mu\nu}(x) := \tfrac{1}{2}\{\hat{g}_{\mu\nu}(x), \hat{t}_{\mu\nu}(x)\}, \qquad \hat{V}_{\mu\nu}(x) := \tfrac{\mathrm{i}}{2}\,[\hat{g}_{\mu\nu}(x), \hat{t}_{\mu\nu}(x)].} \tag{5.79}$$

Define the block–coherent lifts (Pauli–form)

$$\widehat{\boldsymbol{g}}^{\mathrm{HS}}_{\mu\nu}(x) := \mathbb{I}_L \otimes \hat{g}_{\mu\nu}(x) + \tfrac{1}{2}\cos\theta\ \sigma_x \otimes \hat{S}_{\mu\nu}(x) + \tfrac{1}{2}\sin\theta\ \sigma_y \otimes \hat{V}_{\mu\nu}(x), \tag{5.80}$$

$$\widehat{\boldsymbol{t}}^{\mathrm{HS}}_{\mu\nu}(x) := \sigma_x \otimes \hat{t}_{\mu\nu}(x), \tag{5.81}$$

where σ_x, σ_y act on $\mathcal{H}_L$ and $\mathbb{I}_L$ is the identity on $\mathcal{H}_L$. Equivalently, in projector (off–diagonal) form,

$$\begin{aligned}\widehat{\boldsymbol{g}}^{\mathrm{HS}}_{\mu\nu}(x) &= |0\rangle_L\langle 0| \otimes \hat{g}_{\mu\nu} + |1\rangle_L\langle 1| \otimes \hat{g}_{\mu\nu} \\ &\quad + |0\rangle_L\langle 1| \otimes \Big(\tfrac{1}{2}\cos\theta\,\hat{S}_{\mu\nu} - \tfrac{\mathrm{i}}{2}\sin\theta\,\hat{V}_{\mu\nu}\Big) + |1\rangle_L\langle 0| \otimes \Big(\tfrac{1}{2}\cos\theta\,\hat{S}_{\mu\nu} + \tfrac{\mathrm{i}}{2}\sin\theta\,\hat{V}_{\mu\nu}\Big), \\ \widehat{\boldsymbol{t}}^{\mathrm{HS}}_{\mu\nu}(x) &= |0\rangle_L\langle 1| \otimes \hat{t}_{\mu\nu} + |1\rangle_L\langle 0| \otimes \hat{t}_{\mu\nu}.\end{aligned}$$

All operator equalities are in the SOT on the regulated fibres $\mathrm{Ran}\, E^{(x)}([0, \lambda_{\mathrm{cutoff}}])$.

Definition 5.74 (Law–control reduction (manifold dial))**.** Prepare the control in $|+_\varphi\rangle_L := (|0\rangle_L + e^{\mathrm{i}\varphi}|1\rangle_L)/\sqrt{2}$ and postselect the same readout. Using $\langle +_\varphi|\sigma_x|+_\varphi\rangle = \cos\varphi$ and $\langle +_\varphi|\sigma_y|+_\varphi\rangle = \sin\varphi$, the reduced tensors are

$$\begin{aligned}\hat{g}^{\langle\mathrm{HS}\rangle}_{\mu\nu}(x;\varphi,\theta) &:= \langle +_\varphi|\widehat{\boldsymbol{g}}^{\mathrm{HS}}_{\mu\nu}(x)|+_\varphi\rangle_L \\ &= \hat{g}_{\mu\nu}(x) + \tfrac{1}{2}\cos\varphi\,\cos\theta\,\hat{S}_{\mu\nu}(x) + \tfrac{1}{2}\sin\varphi\,\sin\theta\,\hat{V}_{\mu\nu}(x), \qquad (5.82)\\ \hat{t}^{\langle\mathrm{HS}\rangle}_{\mu\nu}(x;\varphi) &:= \langle +_\varphi|\widehat{\boldsymbol{t}}^{\mathrm{HS}}_{\mu\nu}(x)|+_\varphi\rangle_L = \cos\varphi\,\hat{t}_{\mu\nu}(x). \qquad (5.83)\end{aligned}$$

The reduced unified tensor is

$$\boxed{\hat{\Sigma}^{\langle\mathrm{HS}\rangle}_{\mu\nu}(x;\varphi,\theta) := \hat{g}^{\langle\mathrm{HS}\rangle}_{\mu\nu}(x;\varphi,\theta) + \hat{t}^{\langle\mathrm{HS}\rangle}_{\mu\nu}(x;\varphi),} \tag{5.84}$$

hence explicitly

$$\hat{\Sigma}^{\langle\mathrm{HS}\rangle}_{\mu\nu} = \hat{g}_{\mu\nu} + \cos\varphi\,\hat{t}_{\mu\nu} + \tfrac{1}{2}\cos\varphi\,\cos\theta\,\hat{S}_{\mu\nu} + \tfrac{1}{2}\sin\varphi\,\sin\theta\,\hat{V}_{\mu\nu}, \tag{5.85}$$

which matches eq. (5.44).

Proposition 5.75 (Gauge covariance, equivalence, and expectation)**.** *If* $(\hat{g}, \hat{t}) \mapsto (U\hat{g}U^\dagger, U\hat{t}U^\dagger)$ *and* $|\mathring{\boldsymbol{\Psi}}\rangle \mapsto U|\mathring{\boldsymbol{\Psi}}\rangle$*, then* $\widehat{\boldsymbol{g}}^{\mathrm{HS}}_{\mu\nu}$*,* $\widehat{\boldsymbol{t}}^{\mathrm{HS}}_{\mu\nu}$ *and* $\hat{\Sigma}^{\langle\mathrm{HS}\rangle}_{\mu\nu}$ *transform covariantly, and the expectations* $\bar{g}^{\langle\mathrm{HS}\rangle}_{\mu\nu} := \langle\hat{g}^{\langle\mathrm{HS}\rangle}_{\mu\nu}\rangle$*,* $\bar{t}^{\langle\mathrm{HS}\rangle}_{\mu\nu} := \langle\hat{t}^{\langle\mathrm{HS}\rangle}_{\mu\nu}\rangle$*,* $\bar{\Sigma}^{\langle\mathrm{HS}\rangle}_{\mu\nu} := \langle\hat{\Sigma}^{\langle\mathrm{HS}\rangle}_{\mu\nu}\rangle$ *are gauge invariant. Moreover, lifting* $(\hat{g}, \hat{t})$ *separately as above is* equivalent *to lifting* $\hat{\Sigma} = \hat{g} + \hat{t}$ *with the same mixers* $(\hat{S}, \hat{V})$*, so eq.* (5.84) *reproduces eq.* (5.44)*.*

Remark 5.76 (When to include this block)**.** The "canonical law–control word" is optional: it is consistent, gauge–covariant, SOT–regular on the PQS–regulated fibres, and collapses to the baseline (no law superposition) when $\varphi = \theta = 0$. If phase–resolved tomography of sectors is not used, it may be omitted without loss.

5.3.4 Defining the Lorentzian Structure from the Unified Tensor

In the IQR framework, the fundamental classical carrier of emergent geometry is the *Unified Relational Chronogeometric Tensor* at each point $x \in \mathcal{X}_q$,

$$\bar{\Sigma}_{\mu\nu}(x) := \bar{g}_{\mu\nu}(x) + \bar{t}_{\mu\nu}(x),$$

as introduced in Definition 5.26. The constituent expectation tensors $\bar{g}_{\mu\nu}$ and $\bar{t}_{\mu\nu}$ admit the canonical *triadic decomposition* (Definition 5.54),

$$\bar{g}_{\mu\nu} = \bar{g}^{\text{corr}}_{\mu\nu} + \bar{g}^{\text{vac}}_{\mu\nu} + \bar{g}^{\text{spec}}_{\mu\nu}, \qquad \bar{t}_{\mu\nu} = \bar{t}^{\text{corr}}_{\mu\nu} + \bar{t}^{\text{vac}}_{\mu\nu} + \bar{t}^{\text{spec}}_{\mu\nu},$$

so the unified tensor inherits the structured split

$$\boxed{\bar{\Sigma}_{\mu\nu}(x) = \bar{\Sigma}^{\text{corr}}_{\mu\nu}(x) + \bar{\Sigma}^{\text{vac}}_{\mu\nu}(x) + \bar{\Sigma}^{\text{spec}}_{\mu\nu}(x),} \tag{5.86}$$

with sector components

$$\boxed{\bar{\Sigma}^{(a)}_{\mu\nu}(x) := \bar{g}^{(a)}_{\mu\nu}(x) + \bar{t}^{(a)}_{\mu\nu}(x), \quad a \in \{\text{corr}, \text{vac}, \text{spec}\}.} \tag{5.87}$$

Definition 5.77 (Lorentzian metric from the chronogeometric tensor)**.** Given $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (Definition 5.26), define the *effective Lorentzian metric* on $\mathcal{M}_{\text{geom}}$ (Definition 5.113) by symmetrisation

$$\boxed{\bar{g}^{\text{Lor}}_{\mu\nu}(x) := \bar{\Sigma}^{\text{sym}}_{\mu\nu}(x) := \tfrac{1}{2}\Big(\bar{\Sigma}_{\mu\nu}(x) + \bar{\Sigma}_{\nu\mu}(x)\Big)\,.} \tag{5.88}$$

Equivalently,

$$\bar{g}^{\text{Lor}}_{\mu\nu}(x) = \bar{g}_{\mu\nu}(x) + \bar{t}^{\text{sym}}_{\mu\nu}(x), \qquad \bar{t}^{\text{sym}}_{\mu\nu} := \tfrac{1}{2}\Big(\bar{t}_{\mu\nu} + \bar{t}_{\nu\mu}\Big).$$

We say $\bar{g}^{\text{Lor}}_{\mu\nu}$ defines a Lorentzian structure iff $\text{signature}\Big(\bar{g}^{\text{Lor}}_{\mu\nu}(x)\Big) = (+,-,-,-)$ for all $x \in \mathcal{X}_q$ [7, 146].

Signature clause (sufficient pointwise criterion). Let $S^{\mu}{}_{\nu}(x) := \bar{g}^{\mu\alpha}(x)\,\bar{t}^{\text{sym}}_{\alpha\nu}(x)$, so that $\bar{g}^{\text{Lor}} = \bar{g}\,(I + S)$, with $\bar{g}_{\mu\nu}$ as in Definition 5.10. Assume there exists on a domain $U \subset \mathcal{X}_q$ a $\bar{g}$-unit vector u^{μ} with $S^{\mu}{}_{\nu}u^{\nu} = -\lambda\,u^{\mu}$ for some $\lambda > 1$, and let $P_{\perp}$ be the $\bar{g}$-orthogonal projector onto $u^{\perp}$. If

$$\big\|P_{\perp}SP_{\perp}\big\|_{\text{op}} < 1 \quad \text{pointwise on } U,$$

then $I + S$ has exactly one eigenvalue of sign opposite to the remaining $d-1$, hence $\bar{g}^{\text{Lor}}$ is Lorentzian (type $(1, d-1)$).

Compact norm bound (Euclidean regime). Let $\bar{g}^{-1/2}$ be the positive square root of $\bar{g}$, and set $G := \bar{g}^{-1/2}\,\bar{t}^{\text{sym}}\,\bar{g}^{-1/2}$. If

$$\|G\|_{\text{op}} < 1, \tag{5.89}$$

then $\bar{g}^{\text{Lor}} = \bar{g} + \bar{t}^{\text{sym}}$ remains positive–definite (no signature flip).

Compact spectral split (Lorentzian regime). With ordered eigenvalues $\lambda_1(G) \le \cdots \le \lambda_d(G)$, if

$$\lambda_1(G) < -1 < \lambda_2(G), \tag{5.90}$$

then $\bar{g}^{\text{Lor}} = \bar{g} + \bar{t}^{\text{sym}}$ is Lorentzian (one negative, $d-1$ positive).

Remark 5.78 (Projection pathways: operational guide)**.** Two routes appear in practice (cf. Remark 5.47): (i) the *Lorentzian merge* $\bar{g}^{\mathrm{Lor}} = \bar{g} + \bar{t}^{\mathrm{sym}}$ used for causal kinematics and tomography (Definition 5.122 and Proposition 5.60); (ii) the *pure-metric projection* that drops $\bar{t}$ when a classical-limit presentation is desired (e.g. summaries in the FLRW specialisation Section 5.4.7). Both are expectation-level constructions rooted in the micro→macro map of Section 5.2.10.

Interpretation and physical implications.

1. **Triadically structured causality.** Because $\bar{g}^{\mathrm{Lor}}$ is built from $\bar{\Sigma}$, it inherits the triadic structure (5.86). Each sector—correlational, vacuum, spectral—contributes distinctively to causal kinematics; sector-wise diagnostics follow from the holosymmetric push-through and orthogonality conditions (Proposition 5.57 and Corollary 5.58).

2. **Unified source of geometry and coherence.** The light cone arises from the full relational content $\bar{g} + \bar{t}^{\mathrm{sym}}$ (Definition 5.122); $\bar{t}^{\mathrm{sym}}$ encodes quantum-temporal curvature that deforms expectation-level geometry (T-even response), while $\bar{t}^{\mathrm{anti}}$ governs intra-cone phase holonomy (T-odd response) without altering the cone (Section 5.4.3 and Proposition 5.60).

3. **Antisymmetric sector and torsional transport.** The antisymmetric part $\bar{t}^{\mathrm{anti}}_{\mu\nu} := \frac{1}{2}(\bar{t}_{\mu\nu} - \bar{t}_{\nu\mu})$ does not enter $\bar{g}^{\mathrm{Lor}}$ but contributes torsion-like, chiral transport effects within the allowed cone (Section 5.4.3; cf. geometric-phase phenomena [23, 24, 25]).

4. **State-dependent dynamical causality.** Since $\bar{g}_{\mu\nu}$ and $\bar{t}_{\mu\nu}$ are expectation values in the Quantum Holo–State $|\mathring{\mathbf{\Psi}}\rangle$ (Section 5.2.10), the causal structure is *state-dependent* and evolves with the global coherence of the cosmos. In appropriate limits this reproduces the classical GR/QFT picture [7, 244, 11].

5. **Holosymmetric invariance of kinematics.** Holosymmetric phases (φ, θ) change response amplitudes but not cone assignment (Definition 5.123 and Corollary 5.126); thus $\bar{g}^{\mathrm{Lor}}$ supplies the hard kinematic layer of the two-layered architecture (Section 5.4.4).

Summary. The symmetrised unified tensor

$$\bar{g}^{\mathrm{Lor}}_{\mu\nu}(x) = \bar{g}_{\mu\nu}(x) + \bar{t}^{\mathrm{sym}}_{\mu\nu}(x)$$

defines the effective Lorentzian metric on $\mathcal{M}_{\mathrm{geom}}$ provided the signature tests above hold. This construction embeds pseudo-Riemannian geometry within a triadically decomposed, quantum-coherent, temporally enriched framework,

$$\mathcal{M}_{\mathrm{geom}} := \left(\mathcal{X}_q,\ \bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)\right),$$

which interfaces consistently with standard GR/QFT in the appropriate regimes [7, 146, 244, 11].

5.3.5 Holosymmetric Wordwise Lorentzian Structure from the Unified Tensor

Definition 5.79 (Unified word–chronogeometric tensor and triadic split)**.** On the expectation word–manifold $\mathbb{M}_{\mathrm{rel}}^{\langle \mathrm{HS} \rangle} = (\mathcal{X}_q;\, \bar{g}_{\mu\nu},\, \bar{t}_{\mu\nu})$ (Definition 5.70), define the unified tensor

$$\bar{\Sigma}_{\mu\nu}(x) := \bar{g}_{\mu\nu}(x) + \bar{t}_{\mu\nu}(x).$$

With the channelwise decomposition fixed at operator level, the expectation tensors inherit

$$\bar{g}_{\mu\nu} = \sum_{a\in\{\mathrm{corr,vac,spec}\}} \bar{g}^{(a)}_{\mu\nu}, \qquad \bar{t}_{\mu\nu} = \sum_{a\in\{\mathrm{corr,vac,spec}\}} \bar{t}^{(a)}_{\mu\nu},$$

hence

$$\boxed{\bar{\Sigma}_{\mu\nu} = \bar{\Sigma}^{\mathrm{corr}}_{\mu\nu} + \bar{\Sigma}^{\mathrm{vac}}_{\mu\nu} + \bar{\Sigma}^{\mathrm{spec}}_{\mu\nu}, \qquad \bar{\Sigma}^{(a)}_{\mu\nu} := \bar{g}^{(a)}_{\mu\nu} + \bar{t}^{(a)}_{\mu\nu}.} \tag{5.91}$$

All objects are word/frame independent by endogenous expectation invariance.

Definition 5.80 (Effective Lorentzian metric (word symmetrisation))**.** Set $\bar{t}^{\mathrm{sym}}_{\mu\nu} := \frac{1}{2}(\bar{t}_{\mu\nu} + \bar{t}_{\nu\mu})$ and $\bar{t}^{\mathrm{skew}}_{\mu\nu} := \frac{1}{2}(\bar{t}_{\mu\nu} - \bar{t}_{\nu\mu})$. Define the effective Lorentzian metric by symmetrising the unified tensor:

$$\boxed{\bar{g}^{\mathrm{Lor}}_{\mu\nu}(x) := \tfrac{1}{2}\big(\bar{\Sigma}_{\mu\nu}(x) + \bar{\Sigma}_{\nu\mu}(x)\big) = \bar{g}_{\mu\nu}(x) + \bar{t}^{\mathrm{sym}}_{\mu\nu}(x).} \tag{5.92}$$

We say $\bar{g}^{\mathrm{Lor}}$ is Lorentzian on $\mathcal{X}_q$ iff $\mathrm{signature}(\bar{g}^{\mathrm{Lor}}) = (+,-,-,-)$ pointwise. The skew part $\bar{t}^{\mathrm{skew}}_{\mu\nu}$ encodes phase–flow vorticity and does not affect the metric signature.

Remark 5.81 (Word locality and regularity)**.** All tensors above are computed on the spectrally regulated fibres $\mathrm{Ran}\, P_x$ and inherit SOT–C^2 regularity and ultra–sub–Planckian locality from the operator level. Word changes $\mathbf{W} \mapsto \mathbf{W}'$ act by fibre conjugation and leave $\bar{g}^{\mathrm{Lor}}$ invariant.

Pointwise signature diagnostics (wordwise linear algebra)

Definition 5.82 (Compliance operator and compact tests)**.** Let $\bar{g}_{\mu\nu}$ be the positive word–metric from Definition 5.70 and $\bar{t}^{\mathrm{sym}}_{\mu\nu} := \frac{1}{2}(\bar{t}_{\mu\nu} + \bar{t}_{\nu\mu})$. Define the *compliance operator*

$$S^{\mu}{}_{\nu} := \bar{g}^{\mu\alpha}\, \bar{t}^{\mathrm{sym}}_{\alpha\nu}, \qquad \text{equivalently} \qquad G := \bar{g}^{-1/2}\, \bar{t}^{\mathrm{sym}}\, \bar{g}^{-1/2}.$$

Then

$$\bar{g}^{\mathrm{Lor}} = \bar{g} + \bar{t}^{\mathrm{sym}} = \bar{g}^{1/2}\,(I + G)\,\bar{g}^{1/2}.$$

(a) **Euclidean retention.** If $\|G\|_{\mathrm{op}} < 1$ on a domain $U \subset \mathcal{X}_q$, then $I + G \succ 0$ and $\bar{g}^{\mathrm{Lor}}$ remains positive–definite (no signature flip) on U.

(b) **Lorentz tilt by one.** Let u^{μ} be $\bar{g}$-unit, and let $P_{\perp}$ be the $\bar{g}$-orthogonal projector onto $u^{\perp}$. If $Su = -\lambda u$ with $\lambda > 1$ and $\|P_{\perp} S P_{\perp}\|_{\mathrm{op}} < 1$, then $I + S$ has exactly one negative eigenvalue ($1 - \lambda < 0$ along u; $I + P_{\perp} S P_{\perp} \succ 0$ on $u^{\perp}$). Hence $\bar{g}^{\mathrm{Lor}}$ is Lorentzian.

(c) **Compact spectral split.** Let the eigenvalues of G be $\lambda_1(G) \le \cdots \le \lambda_d(G)$. If

$$\lambda_1(G) < -1 < \lambda_2(G),$$

then $I + G$ has one negative and $d-1$ positive eigenvalues, so $\bar{g}^{\mathrm{Lor}}$ is Lorentzian.

Remark 5.83 (Triadic attribution of the tilt)**.** Because $\bar{t}^{\mathrm{sym}} = \sum_a \bar{t}^{(a)\mathrm{sym}}$ with $a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$,

$$G = \sum_a \bar{g}^{-1/2}\, \bar{t}^{(a)\mathrm{sym}}\, \bar{g}^{-1/2}.$$

If each $G^{(a)} := \bar{g}^{-1/2}\bar{t}^{(a)\mathrm{sym}}\bar{g}^{-1/2}$ is small ($\|G^{(a)}\|_{\mathrm{op}} < \epsilon_a$) but the correlator channel has $\lambda_1(G^{\mathrm{corr}}) < -1 - \sum_{\beta \neq \mathrm{corr}} \epsilon_\beta$, then the Lorentzian character is already secured by the correlator sector.

Corollary 5.84 (Word/gauge invariance of cones)**.** *Under fibre word changes and gauge rotations, G transforms by similarity, hence its spectrum—and the null cone of $\bar{g}^{\mathrm{Lor}}$—is invariant. The form $\bar{g}$ remains positive and fixed as a bilinear form.*

Remark 5.85 (Operational dial (optional))**.** In the holosymmetric law–control variant, phase φ and coherence θ tune the symmetric part entering $\bar{g}^{\mathrm{Lor}} = \bar{g} + \bar{t}^{\mathrm{sym}}$. Phase scans isolate anticommutator vs. commutator couplings at operator level, but only the symmetric (anticommutator) contribution shifts the signature at expectation level.

Summary. In wordwise terms, the Lorentzian metric is the symmetric Hadamard lift of $\bar{\Sigma}$. Its causal type is governed by $G = \bar{g}^{-1/2}\bar{t}^{\mathrm{sym}}\bar{g}^{-1/2}$: small $\|G\|_{\mathrm{op}}$ keeps the geometry Euclidean; one eigenvalue crossing below -1 (with the rest > -1) produces a single–cone Lorentz tilt, i.e. signature $(1, d-1)$.

5.3.6 Symmetrizing the Unified Lorentzian Tensor

Step 0: Triadic Decomposition of the Chronogeometric Tensor

In IQR, the unified relational chronogeometric tensor field

$$\bar{\Sigma}_{\mu\nu}(x) := \bar{g}_{\mu\nu}(x) + \bar{t}_{\mu\nu}(x)$$

(Definition 5.26) admits the canonical *triadic decomposition* (Definition 5.54) inherited from the operator level and the micro→macro map (Section 5.2.10):

$$\bar{g}_{\mu\nu} = \bar{g}^{\mathrm{corr}}_{\mu\nu} + \bar{g}^{\mathrm{vac}}_{\mu\nu} + \bar{g}^{\mathrm{spec}}_{\mu\nu}, \qquad \bar{t}_{\mu\nu} = \bar{t}^{\mathrm{corr}}_{\mu\nu} + \bar{t}^{\mathrm{vac}}_{\mu\nu} + \bar{t}^{\mathrm{spec}}_{\mu\nu}.$$

Hence

$$\boxed{\bar{\Sigma}_{\mu\nu}(x) = \sum_{a \in \{\mathrm{corr},\, \mathrm{vac},\, \mathrm{spec}\}} \bar{\Sigma}^{(a)}_{\mu\nu}(x), \qquad \bar{\Sigma}^{(a)}_{\mu\nu}(x) := \bar{g}^{(a)}_{\mu\nu}(x) + \bar{t}^{(a)}_{\mu\nu}(x)\ .}$$

Physically, the sectors encode (i) correlational contributions from relational structure, (ii) vacuum fluctuations, and (iii) coherent spectral content; when channel projectors commute, cross-channel mixing vanishes (Corollary 5.58).

Step 1: Symmetrization within triadic components (detailed expansion)

Starting from $\bar{\Sigma}_{\mu\nu} = \sum_a \bar{\Sigma}^{(a)}_{\mu\nu}$ with $\bar{\Sigma}^{(a)}_{\mu\nu} = \bar{g}^{(a)}_{\mu\nu} + \bar{t}^{(a)}_{\mu\nu}$, note $\bar{g}^{(a)}_{\mu\nu} = \bar{g}^{(a)}_{\nu\mu}$ while $\bar{t}^{(a)}_{\mu\nu}$ may be non-symmetric. Decompose

$$\bar{t}^{\mathrm{sym},(a)}_{\mu\nu} := \tfrac{1}{2}\Big(\bar{t}^{(a)}_{\mu\nu} + \bar{t}^{(a)}_{\nu\mu}\Big), \qquad \bar{t}^{\mathrm{anti},(a)}_{\mu\nu} := \tfrac{1}{2}\Big(\bar{t}^{(a)}_{\mu\nu} - \bar{t}^{(a)}_{\nu\mu}\Big),$$

and define the symmetric part of each sector:

$$\bar{\Sigma}^{\mathrm{sym},(a)}_{\mu\nu} := \tfrac{1}{2}\Big(\bar{\Sigma}^{(a)}_{\mu\nu} + \bar{\Sigma}^{(a)}_{\nu\mu}\Big) = \bar{g}^{(a)}_{\mu\nu} + \bar{t}^{\mathrm{sym},(a)}_{\mu\nu}.$$

Summing over sectors yields

$$\boxed{\bar{\Sigma}^{\mathrm{sym}}_{\mu\nu}(x) = \sum_a \bar{\Sigma}^{\mathrm{sym},(a)}_{\mu\nu}(x) = \sum_a \Big(\bar{g}^{(a)}_{\mu\nu}(x) + \bar{t}^{\mathrm{sym},(a)}_{\mu\nu}(x)\Big)\ ,}$$

which is the unique symmetric tensor obtained by sectorwise symmetrisation and preserves the ontological identity of each triadic channel (cf. Sections 5.4.3 and 5.4.4 for the hard/soft split).

Step 2: Construction of the effective Lorentzian metric

By definition (Definition 5.77),

$$\boxed{\bar{g}^{\mathrm{Lor}}_{\mu\nu}(x) := \bar{\Sigma}^{\mathrm{sym}}_{\mu\nu}(x) = \sum_{a\in\{\mathrm{corr},\,\mathrm{vac},\,\mathrm{spec}\}} \Big(\bar{g}^{(a)}_{\mu\nu}(x) + \bar{t}^{\mathrm{sym},(a)}_{\mu\nu}(x)\Big) = \bar{g}_{\mu\nu}(x) + \bar{t}^{\mathrm{sym}}_{\mu\nu}(x)\ .}$$

This field is symmetric, covariant of rank two, and serves as the *hard* causal layer fixing the effective light-cone structure (Definitions 5.122 and 5.150). It defines a Lorentzian geometry whenever the signature tests in Definition 5.77 are satisfied; in that case $\mathcal{M}_{\mathrm{geom}} = (\mathcal{X}_q, \bar{g}^{\mathrm{Lor}})$ is pseudo-Riemannian in the GR sense [7, 146].

Step 3: Physical interpretation of triadic contributions

- $\bar{g}^{\mathrm{corr}}_{\mu\nu}$ and $\bar{t}^{\mathrm{sym,corr}}_{\mu\nu}$: encode kinematic and T-even phase-curvature effects sourced by relational correlations among PQSs, shaping near-field causal geometry and coherence gradients (cf. Section 5.4.6).
- $\bar{g}^{\mathrm{vac}}_{\mu\nu}$ and $\bar{t}^{\mathrm{sym,vac}}_{\mu\nu}$: capture vacuum-induced fine structure; in symmetric regimes they contribute small deformations consistent with QFT on curved spacetime [244, 11].
- $\bar{g}^{\mathrm{spec}}_{\mu\nu}$ and $\bar{t}^{\mathrm{sym,spec}}_{\mu\nu}$: encode coherent/oscillatory content that can induce frequency-resolved anisotropies; holosymmetric phase control tomographically separates these via T-even (anticommutator) scans (Proposition 5.60).

The antisymmetric pieces $\bar{t}^{\mathrm{anti},(a)}_{\mu\nu}$ do *not* enter $\bar{g}^{\mathrm{Lor}}$ but govern T-odd, intra-cone phase holonomy (temporal torsion) measurable via commutator responses—leaving the cone itself unchanged (Section 5.4.3, Proposition 5.60, and Corollary 5.126).

Summary. Sectorwise symmetrisation produces $\bar{g}^{\mathrm{Lor}}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}^{\mathrm{sym}}_{\mu\nu}$ as the effective metric determining the hard causal boundary (Definition 5.122). This construction is (i) triadically faithful, (ii) compatible with channel orthogonality (Corollary 5.58), and (iii) operationally aligned with the two-layered architecture: $\bar{g}^{\mathrm{Lor}}$ fixes *whether* influence can propagate, while $\bar{t}^{\mathrm{anti}}$ and residual sector structure determine *how* coherence is transported within the cone (Sections 5.4.3 and 5.4.4).

Composite nature of the effective metric. The sectorwise symmetrisation shows that the effective Lorentzian metric $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ is not monolithic but explicitly *composite*: $\bar{g}^{\mathrm{Lor}}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}^{\mathrm{sym}}_{\mu\nu}$ (Definition 5.77). It synthesises a purely distance-based geometry $\bar{g}_{\mu\nu}$ with the symmetric component of the temporal phase-flow $\bar{t}^{\mathrm{sym}}_{\mu\nu}$ (Section 5.4.3). Consequently, the classical geodesic equation governed by $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ (Levi–Civita connection) encodes motion in a spacetime where distance curvature and phase-curvature are fused; in the strict classical limit this reduces to the familiar GR picture [7, 146]. Phase-sensitive quantum dynamics, however, can privilege *paths of maximal coherence* within the same light-cone kinematics (hard layer), reflecting the soft-layer influence of $\bar{t}_{\mu\nu}$ on interference and decoherence (Sections 5.4.3 and 5.4.4; see also [287, 24, 247]).

Triadically sourced causality. Because $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ is built from the unified tensor $\bar{\Sigma}_{\mu\nu}$, it inherits the triadic split $\bar{\Sigma}_{\mu\nu} = \bar{\Sigma}^{\mathrm{corr}}_{\mu\nu} + \bar{\Sigma}^{\mathrm{vac}}_{\mu\nu} + \bar{\Sigma}^{\mathrm{spec}}_{\mu\nu}$ (eq. (5.86) and Definition 5.54). The effective light cone— the absolute boundary of cause and effect—is therefore a *composite* outcome determined by (i) correlational structure (relational/entanglement content), (ii) vacuum contributions (zero-point structure), and (iii) spectral/coherent excitations, all aggregated through $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ (Definition 5.122). This yields a causal fabric that is not a passive backdrop but a dynamically textured tapestry whose sectoral threads can be probed and, when channel projectors commute, cleanly separated (Corollary 5.58 and Proposition 5.57). In symmetric and classical limits the sectoral contributions collapse to the standard GR/QFT light-cone structure [244, 11], whereas away from those limits the temporal sector imprints anisotropic coherence and phase-holonomy inside the cone without altering its boundary (Section 5.4.3).

5.3.7 Holosymmetric Wordwise Symmetrization of the Unified Lorentzian Tensor

Definition 5.86 (Triadic wordwise chronogeometry)**.** Let $\mathcal{A} := \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$ denote the triadic channel types, and let $\mathbf{g}^{(a)}_{\mu\nu}(x)$ and $\mathbf{t}^{(a)}_{\mu\nu}(x)$ be the *typed, second–rank words* (metric and temporal words) on the regulated fibre at x. Define the unified chronogeometric word by the Hadamard (componentwise) sum

$$\mathbf{\Sigma}_{\mu\nu}(x) \;:=\; \sum_{a\in\mathcal{A}} \Big(\mathbf{g}^{(a)}_{\mu\nu}(x) \;\oplus\; \mathbf{t}^{(a)}_{\mu\nu}(x) \Big).$$

Here $\oplus$ is the wordwise Hadamard sum; the word product and adjoint are those of the fourfold/transport dialectic, all taken continuously in the strong–operator topology (SOT) on the spectrally regulated subspace.

Definition 5.87 (Wordwise symmetrizer and port–flip)**.** For any biform word $\mathbf{X}_{\mu\nu}$ (two free modal indices), denote by $\mathbf{X}^{\leftrightarrow}_{\nu\mu}$ the *port–flip* (swap of test/source ports and index order), and set

$$\mathbb{S}[\mathbf{X}]_{\mu\nu} := \tfrac{1}{2}\Big(\mathbf{X}_{\mu\nu} \;\oplus\; \mathbf{X}^{\leftrightarrow}_{\nu\mu}\Big), \qquad \mathbb{A}[\mathbf{X}]_{\mu\nu} := \tfrac{1}{2}\Big(\mathbf{X}_{\mu\nu} \;\ominus\; \mathbf{X}^{\leftrightarrow}_{\nu\mu}\Big),$$

with $\ominus$ the Hadamard difference. Then $\mathbb{S}[\mathbf{X}]_{\mu\nu} = \mathbb{S}[\mathbf{X}]_{\nu\mu}$ and $\mathbb{A}[\mathbf{X}]_{\mu\nu} = -\mathbb{A}[\mathbf{X}]_{\nu\mu}$.

Lemma 5.88 (Sectorwise wordwise symmetrization)**.** *For each $a \in \mathcal{A}$, define $\mathbf{\Sigma}^{(a)}_{\mu\nu} := \mathbf{g}^{(a)}_{\mu\nu} \oplus \mathbf{t}^{(a)}_{\mu\nu}$. Then*

$$\mathbf{\Sigma}^{\mathrm{sym}}_{\mu\nu} := \mathbb{S}[\mathbf{\Sigma}]_{\mu\nu} = \sum_{a\in\mathcal{A}} \Big(\mathbf{g}^{(a)}_{\mu\nu} \;\oplus\; \mathbb{S}[\mathbf{t}^{(a)}]_{\mu\nu}\Big), \qquad \mathbf{\Sigma}^{\mathrm{anti}}_{\mu\nu} = \sum_{a\in\mathcal{A}} \mathbb{A}[\mathbf{t}^{(a)}]_{\mu\nu}.$$

Sketch. *$\mathbf{g}^{(a)}_{\mu\nu}$ is port–symmetric by construction (metric channel), hence fixed by $\mathbb{S}$; only $\mathbf{t}^{(a)}$ splits into symmetric/antisymmetric parts.*

Definition 5.89 (Effective Lorentzian *word–metric*)**.** Define the symmetric, rank–two *Lorentzian word* by

$$\mathbf{g}^{\mathrm{Lor}}_{\mu\nu}(x) \;:=\; \mathbf{\Sigma}^{\mathrm{sym}}_{\mu\nu}(x) \;=\; \mathbf{g}_{\mu\nu}(x) \;\oplus\; \mathbb{S}[\mathbf{t}]_{\mu\nu}(x),$$

with $\mathbf{g}_{\mu\nu} := \sum_a \mathbf{g}^{(a)}_{\mu\nu}$ and $\mathbf{t}_{\mu\nu} := \sum_a \mathbf{t}^{(a)}_{\mu\nu}$.

Proposition 5.90 (Endogenous expectation $\Leftrightarrow$ classical construction)**.** *Let $\langle\cdot\rangle_\Psi := \langle \mathring{\mathbf{\Psi}}| \cdot |\mathring{\mathbf{\Psi}}\rangle$ be the endogenous expectation on the regulated fibre. Then*

$$\Big\langle \mathbf{g}^{\mathrm{Lor}}_{\mu\nu} \Big\rangle_\Psi = \bar{g}_{\mu\nu} \;+\; \bar{t}^{\mathrm{sym}}_{\mu\nu} =:\; \bar{g}^{\mathrm{Lor}}_{\mu\nu},$$

and, sectorwise,

$$\Big\langle \mathbf{\Sigma}^{\mathrm{sym}}_{\mu\nu} \Big\rangle_\Psi = \sum_a \Big(\bar{g}^{(a)}_{\mu\nu} + \bar{t}^{(a)\mathrm{sym}}_{\mu\nu}\Big).$$

Thus the wordwise symmetrization reproduces exactly the expectation–level Lorentzian construction (Definition 5.80).

Remark 5.91 (Role of the antisymmetric temporal word)**.** The piece $\mathbf{\Sigma}^{\mathrm{anti}}_{\mu\nu} = \mathbb{A}[\mathbf{t}]_{\mu\nu}$ is T–odd and does not enter $\mathbf{g}^{\mathrm{Lor}}$; it captures intra–cone phase holonomy ("temporal torsion") accessible via commutator–sensitive holosymmetric scans (law–control phase φ), leaving the light–cone set by $\mathbf{g}^{\mathrm{Lor}}$ unchanged.

Definition 5.92 (Signature test in word form)**.** Let $\mathbf{g}^{-1}$ be the (fibrewise) inverse metric word and set the *compliance word*

$$\mathbf{S}^{\mu}{}_{\nu} \;:=\; \Big(\mathbf{g}^{-1}\Big)^{\mu\alpha} \;\odot\; \mathbb{S}[\mathbf{t}]_{\alpha\nu},$$

with $\odot$ the word product. Then $\mathbf{g}^{\mathrm{Lor}} = \mathbf{g} \odot (\mathbf{I} + \mathbf{S})$. If, at a point, the operator norm on the regulated fibre obeys $\|\mathbf{G}\|_{\mathrm{op}} < 1$ for $\mathbf{G} := \mathbf{g}^{-1/2} \odot \mathbb{S}[\mathbf{t}] \odot \mathbf{g}^{-1/2}$, the signature stays Euclidean; if $\lambda_1(\mathbf{G}) < -1 < \lambda_2(\mathbf{G})$ (ordered eigenvalues), the signature is Lorentzian (type $(1, d-1)$). Under endogenous expectation these reduce to the spectral criteria in Definition 5.80.

Corollary 5.93 (Triadic fidelity and channel orthogonality)**.** *If the channel projectors commute (word–orthogonality), then no cross–channel terms are created by $\mathbb{S}$, and*

$$\mathbf{g}^{\mathrm{Lor}}_{\mu\nu} = \sum_{a\in\mathcal{A}} \Big(\mathbf{g}^{(a)}_{\mu\nu} \;\oplus\; \mathbb{S}[\mathbf{t}^{(a)}]_{\mu\nu}\Big),$$

exactly mirroring the triadic, sectorwise symmetrization.

Physical reading (wordwise). $\mathbf{g}^{\mathrm{Lor}}$ fixes the *hard* causal boundary (light–cone) via T–even content; $\mathbb{A}[\mathbf{t}]$ determines *soft* intra–cone phase transport/holonomy. Law–control phases (φ, θ) from the holosymmetric dial isolate anticommutator vs. commutator responses without altering the cone unless the symmetric temporal word crosses the spectral thresholds of Definition 5.92.

One–line summary. Applying the holosymmetric symmetrizer $\mathbb{S}$ to the unified chronogeometric word $\mathbf{\Sigma}$ yields the Lorentzian word–metric

$$\boxed{\mathbf{g}^{\mathrm{Lor}}_{\mu\nu} = \mathbf{g}_{\mu\nu} \oplus \mathbb{S}[\mathbf{t}]_{\mu\nu}} \quad \Longrightarrow \quad \boxed{\bar{g}^{\mathrm{Lor}}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}^{\mathrm{sym}}_{\mu\nu}}.$$

Remark 5.94 (Why symmetrization is required in the wordwise construction)**.** There are two equivalent routes to an emergent metric in the holosymmetric wordwise setting:

(i) Scalar–Hessian route. Take the coincidence Hessian of the (expectation–level) composite distance $D^{2,[\mathbf{W}]}(x, x')$. By construction the Hessian is symmetric, yielding $\bar{g}_{\mu\nu}$ directly Definition 5.11 and eq. (5.17); at operator level this is the SOT–Fréchet Hessian of $\hat{D}^{2,[\mathbf{W}]}$ giving $\hat{g}^{[\mathbf{W}]}_{\mu\nu}$ Definition 5.2 and eq. (5.12), with $\bar{g}_{\mu\nu} = \langle \hat{g}_{\mu\nu} \rangle$ Remark 5.12. No explicit symmetrization is needed here because Hessians are symmetric.

(ii) Unified–tensor route. Assemble the unified chronogeometric tensor $\Sigma = g + t$ (operator or expectation level) by adding the metric and temporal sectors Definitions 5.48 and 5.50. The temporal sector $t_{\mu\nu}$ is *not* a priori symmetric—its construction involves path–ordered transport and commutators Definition 5.18 and eq. (5.23). Therefore one must explicitly pass to the symmetric part to obtain a bona fide metric:

$$\bar{g}^{\mathrm{Lor}}_{\mu\nu} = \mathrm{Sym}\ \bar{\mathbf{\Sigma}}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}^{\mathrm{sym}}_{\mu\nu},$$

as stipulated in Definition 5.80; or, in word form, via the port–flip symmetrizer $\mathbb{S}$ of Definition 5.87, yielding $\mathbf{g}^{\mathrm{Lor}}_{\mu\nu} = \mathbf{g}_{\mu\nu} \oplus \mathbb{S}[\mathbf{t}]_{\mu\nu}$ Definition 5.89 and Proposition 5.90.

Why the symmetrization is essential in (ii).

1. **Metric axioms.** A $(0, 2)$ metric must be symmetric to define a quadratic form and causal cones. Since $t_{\mu\nu}$ is generally non-symmetric, the symmetric projector in Definition 5.80 (or Definition 5.87) is required.
2. **Port/orientation invariance.** Wordwise objects carry an ordered pair of "ports" (μ, ν). The port–flip average $\mathbb{S}$ removes this ordering dependence, making the result invariant under $(\mu, \nu) \leftrightarrow (\nu, \mu)$ Definition 5.87.
3. **Hermitian compliance tests.** Signature diagnostics rely on the Hermitian compliance operator $G = \bar{g}^{-1/2}\, \bar{t}^{\mathrm{sym}}\, \bar{g}^{-1/2}$ Definition 5.82. Keeping the antisymmetric part would render G non-self-adjoint and invalidate the spectral criteria.

Role of the antisymmetric temporal content. The skew piece $\bar{t}^{\mathrm{skew}}_{\mu\nu} = \frac{1}{2}(\bar{t}_{\mu\nu} - \bar{t}_{\nu\mu})$ encodes phase–flow vorticity/temporal holonomy and does not affect the metric signature; see the channelwise split and properties in Proposition 5.19 and the wordwise separation in Lemma 5.88 and Remark 5.91.

Conclusion. Route (i) and route (ii) + symmetrization are equivalent: the explicit symmetrizer in Definitions 5.80, 5.87 and 5.89 precisely reproduces the symmetric Hessian outcome Remark 5.12, ensuring a well-posed Lorentzian structure while cleanly separating symmetric (metric) and antisymmetric (holonomy) temporal effects.

5.3.8 Emergent Proper Time and Distance

Setup and projection choice. Expectation values of the operator-valued geometric fields yield the classical tensors $\bar{g}_{\mu\nu}(x) = \langle \mathring{\Psi} | \hat{g}_{\mu\nu}(x) | \mathring{\Psi} \rangle$ and $\bar{t}_{\mu\nu}(x) = \langle \mathring{\Psi} | \hat{t}_{\mu\nu}(x) | \mathring{\Psi} \rangle$ (Section 5.2.10 and Definition 5.26). For causal kinematics and interval measurements we adopt the *Lorentzian merge* metric

$$\bar{g}^{\mathrm{Lor}}_{\mu\nu}(x) := \bar{g}_{\mu\nu}(x) + \bar{t}^{\mathrm{sym}}_{\mu\nu}(x), \qquad \bar{t}^{\mathrm{sym}}_{\mu\nu} := \tfrac{1}{2}\big(\bar{t}_{\mu\nu} + \bar{t}_{\nu\mu}\big),$$

as defined in Definition 5.77, which fixes the effective light-cone structure (Definitions 5.122 and 5.150).[1]

$$\boxed{\mathcal{M}_{\mathrm{geom}} := \big(\mathcal{X}_q, \bar{g}^{\mathrm{Lor}}_{\mu\nu}\big) \quad \text{(emergent chronogeometric manifold)}}$$

Emergent proper time. For an infinitesimal displacement $\Delta x^\mu := x'^\mu - x^\mu$ with $\bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)\Delta x^\mu \Delta x^\nu > 0$ (timelike), the *emergent geometric time* (proper time) is

Definition 5.95 (Emergent geometric time)**.**

$$\boxed{t_{\mathrm{geom}}(x, x') \;:=\; \Big[\, \bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)\, \Delta x^\mu \Delta x^\nu \,\Big]^{1/2}.} \tag{5.93}$$

This is the standard GR proper-time element for the effective metric $\bar{g}^{\mathrm{Lor}}$ [7, 146], interpreted here as the observer's elapsed time in the emergent geometry (Definition 5.150).

In the pure-metric projection (dropping $\bar{t}$), t_{geom} reduces to the familiar expression with $\bar{g}$. The relational temporal functional introduced earlier coincides with t_{geom} on timelike displacements under the chosen projection route (cf. Section 5.2.6).

Emergent proper distance. For spacelike separations, $\bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)\Delta x^\mu \Delta x^\nu < 0$, define:

Definition 5.96 (Emergent geometric distance)**.**

$$\boxed{d_{\mathrm{geom}}(x, x') \;:=\; \Big[\, -\, \bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)\, \Delta x^\mu \Delta x^\nu \,\Big]^{1/2},} \tag{5.94}$$

which yields the invariant spatial separation for the signature $(+,-,-,-)$ convention [7, 146]. In the pure-metric projection this likewise reduces to the standard expression with $\bar{g}$.

[1] When a classical-limit presentation is desired, one may elect the pure-metric projection that drops $\bar{t}$; the operational guidance distinguishing these routes is summarised in Remark 5.78.

Synthesis and identifications. With the Lorentzian merge choice (Definition 5.77),

$$\boxed{\begin{aligned} \mathcal{M}_{\text{geom}} &= \left(\mathcal{X}_q, \bar{g}^{\text{Lor}}_{\mu\nu}\right) \quad \text{(causal cones and intervals from } \bar{g}^{\text{Lor}}\text{)}; \\ t_{\text{geom}}(x,x') &= \left|\bar{g}^{\text{Lor}}_{\mu\nu}\,\Delta x^\mu \Delta x^\nu\right|^{1/2} \quad \text{if timelike}, \\ d_{\text{geom}}(x,x') &= \left|-\bar{g}^{\text{Lor}}_{\mu\nu}\,\Delta x^\mu \Delta x^\nu\right|^{1/2} \quad \text{if spacelike}. \end{aligned}} \tag{5.95}$$

Interpretation and usage. The sign of $\bar{g}^{\text{Lor}}_{\mu\nu}\Delta x^\mu \Delta x^\nu$ classifies displacements and selects the appropriate invariant (time vs. distance); the negative sign in Definition 5.96 ensures positivity for spacelike intervals in signature $(+,-,-,-)$ [7]. These kinematic invariants furnish the *hard* layer of the two-layered causal architecture (Section 5.4.4). The *soft* layer—the temporal phase-flow $\bar{t}_{\mu\nu}$—does not alter which intervals are admissible but modulates coherence and phase transport *within* the light cone (Section 5.4.3 and Definition 5.122). In symmetric limits (e.g., the FLRW specialisation), t_{geom} reproduces cosmic/conformal-time intervals and d_{geom} reproduces comoving/proper distances consistent with QFT on curved spacetime [244, 11, 146].

5.3.9 The Metric Projection: Relational Manifold to Classical Spacetime

We have established that the full expectation-level description of the relational geometry is the bi-tensorial manifold

$$\mathcal{M}_{\text{rel}} = \left(\mathcal{X}_q,\ \bar{g}_{\mu\nu},\ \bar{t}_{\mu\nu}\right),$$

with distinguishability geometry $\bar{g}_{\mu\nu}$ and temporal phase-flow $\bar{t}_{\mu\nu}$ obtained from operator fields via the micro→macro map (Section 5.2.10 and Definition 5.26). In contrast, the spacetime of General Relativity is a pseudo-Riemannian manifold governed by a single metric tensor [7, 146].

One operational pathway from $\mathcal{M}_{\text{rel}}$ to a classical presentation is the *pure-metric projection*, which discards $\bar{t}_{\mu\nu}$ and retains only $\bar{g}_{\mu\nu}$ (cf. the "projection pathways" guide in Remark 5.78). This complements the *Lorentzian merge* route $\bar{g}^{\text{Lor}} = \bar{g} + \bar{t}^{\text{sym}}$ used for causal kinematics (Definitions 5.77 and 5.122); the present subsection formalises the pure-metric route.

Definition 5.97 (Metric projection from the relational manifold)**.** Let

$$\mathcal{M}_{\text{rel}} = \left(\mathcal{X}_q,\ \bar{g}_{\mu\nu}(x),\ \bar{t}_{\mu\nu}(x)\right).$$

The *metric projection* is the structure-preserving semantic map

$$\text{Proj}_{\bar{g}}:\ \mathcal{M}_{\text{rel}} \longrightarrow \mathcal{M}^{(\text{proj})}_{\text{geom}}$$

defined by

$$\text{Proj}_{\bar{g}}:\ \left(\mathcal{X}_q, \bar{g}_{\mu\nu}, \bar{t}_{\mu\nu}\right) \ \mapsto\ \left(\mathcal{X}_q,\ \bar{g}_{\mu\nu},\ d^{(\text{proj})}_{\text{geom}},\ t^{(\text{proj})}_{\text{geom}}\right),$$

where for an infinitesimal displacement $\Delta x^\mu = x'^\mu - x^\mu$,

$$d^{(\mathrm{proj})}_{\mathrm{geom}}(x,x') := \begin{cases} \Big[-\bar{g}_{\mu\nu}(x)\Delta x^\mu \Delta x^\nu \Big]^{1/2}, & \bar{g}_{\mu\nu}\Delta x^\mu \Delta x^\nu < 0, \\ 0, & \text{otherwise}, \end{cases}$$

$$t^{(\mathrm{proj})}_{\mathrm{geom}}(x,x') := \begin{cases} \Big[\bar{g}_{\mu\nu}(x)\Delta x^\mu \Delta x^\nu \Big]^{1/2}, & \bar{g}_{\mu\nu}\Delta x^\mu \Delta x^\nu > 0, \\ 0, & \text{otherwise}. \end{cases}$$

The resulting classical spacetime is

$$\boxed{\mathcal{M}^{(\mathrm{proj})}_{\mathrm{geom}} := \big(\mathcal{X}_q,\ \bar{g}_{\mu\nu}\big)},$$

a pseudo-Riemannian manifold in the sense of GR [7, 146].

Remark 5.98 (Conceptual and operational meaning)**.** The map $\mathrm{Proj}_{\bar{g}}$ is *semantic* rather than linear: it (i) *coarse-grains* away the temporal tensor $\bar{t}_{\mu\nu}$ (see also Sections 5.4.3 and 6.1.3) and (ii) *introduces* the standard GR interval scalars $t^{(\mathrm{proj})}_{\mathrm{geom}}$, $d^{(\mathrm{proj})}_{\mathrm{geom}}$ built solely from $\bar{g}$. This yields a purely metric presentation in which geodesics, distances, and cone assignments are encoded by $\bar{g}$ alone, thus manifestly recovering the GR language when appropriate [7].

Remark 5.99 (Consistency with the Lorentzian-merge route)**.** In the alternative *Lorentzian merge* pathway, intervals are computed using $\bar{g}^{\mathrm{Lor}} = \bar{g} + \bar{t}^{\mathrm{sym}}$ (Section 5.3.8 and Definition 5.77). The two presentations coincide when $\bar{t}^{\mathrm{sym}} = 0$ or is sufficiently aligned/small so that cone tilts are negligible (e.g., homogeneous/isotropic alignment, FLRW specialisation: Section 5.4.7). More generally, $\bar{g}^{\mathrm{Lor}}$ supplies the *hard* kinematic layer, while $\bar{t}$ provides *soft* intra-cone phase transport (Section 5.4.4); the metric projection intentionally omits the latter to expose the classical limit cleanly.

Remark 5.100 (Equivariance and causality)**.** Both routes are diffeomorphism-equivariant at the expectation level. Metric projection preserves GR microcausality relative to $\bar{g}$, while the merge route preserves microcausality relative to $\bar{g}^{\mathrm{Lor}}$ (Definitions 5.122 and 5.150). In regimes where $\bar{t}^{\mathrm{sym}}$ is projected out by symmetry, the cone assignments agree, $J^{\pm}_{\bar{\Sigma}} = J^{\pm}_{\bar{g}^{\mathrm{Lor}}} = J^{\pm}_{\bar{g}}$ (cf. Section 5.4.7).

5.3.10 Holosymmetric Wordwise Emergence of Proper Time and Distance

Definition 5.101 (Lorentzian word from the unified chronogeometry)**.** Let $\mathbf{g}_{\mu\nu}(x)$ and $\mathbf{t}_{\mu\nu}(x)$ be the metric and temporal *rank–two words* on the regulated fibre at x. Let $\mathbb{S}[\cdot]$ denote the wordwise symmetrizer (port–flip average; cf. Definition 5.87). Define the Lorentzian *word–metric*

$$\boxed{\mathbf{g}^{\mathrm{Lor}}_{\mu\nu}(x) \ :=\ \mathbf{g}_{\mu\nu}(x) \ \oplus\ \mathbb{S}[\mathbf{t}]_{\mu\nu}(x)}$$

which is symmetric in (μ,ν) (word sense) and gauge–covariant under fibre unitaries.

Definition 5.102 (Wordwise quadratic form and causal predicate)**.** For an infinitesimal displacement word $\Delta\mathbf{x}^\mu$ at x (from the operational connection along a short lift $x \to x'$), define the Lorentzian quadratic word

$$\mathcal{Q}(x;\Delta) \;:=\; \Delta\mathbf{x}^\mu \;\odot\; \mathbf{g}^{\mathrm{Lor}}_{\mu\nu}(x) \;\odot\; \Delta\mathbf{x}^\nu,$$

with $\odot$ the word product/contraction. Introduce the word–predicates (typed characteristic words)

$$\Pi_{\mathrm{time}}(x;\Delta) := \mathbf{1}_{\{\mathcal{Q}>0\}}, \qquad \Pi_{\mathrm{space}}(x;\Delta) := \mathbf{1}_{\{\mathcal{Q}<0\}}.$$

Definition 5.103 (Emergent proper–time and proper–distance *words*)**.** On the time channel $\big(\Pi_{\mathrm{time}}\big)$,

$$\boxed{\mathbf{t}_{\mathrm{geom}}(x,x') \;:=\; \sqrt{\;\mathcal{Q}(x;\Delta)\;}\,, \qquad \Delta\mathbf{x}^\mu := \mathbf{x}'^\mu - \mathbf{x}^\mu.}$$

On the space channel $\big(\Pi_{\mathrm{space}}\big)$,

$$\boxed{\mathbf{d}_{\mathrm{geom}}(x,x') \;:=\; \sqrt{\;-\,\mathcal{Q}(x;\Delta)\;}\,.}$$

Square roots are taken in the positive branch on the regulated fibre (SOT), and the predicates ensure the correct sign of the arguments.

Proposition 5.104 (Endogenous expectation equals classical formulas)**.** *Let* $\langle\!\langle\cdot\rangle\!\rangle := \langle\mathring{\mathbf{\Psi}}|\cdot|\mathring{\mathbf{\Psi}}\rangle$. *Then*

$$\langle\!\langle \mathbf{g}^{\mathrm{Lor}}_{\mu\nu}\rangle\!\rangle = \bar{g}^{\mathrm{Lor}}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}^{\mathrm{sym}}_{\mu\nu},$$

and, for sufficiently small Δx^μ*,*

$$\boxed{\langle\!\langle \mathbf{t}_{\mathrm{geom}}(x,x')\rangle\!\rangle = \Big[\bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)\,\Delta x^\mu \Delta x^\nu\Big]^{1/2} \;\textit{(timelike)}}$$

$$\boxed{\langle\!\langle \mathbf{d}_{\mathrm{geom}}(x,x')\rangle\!\rangle = \Big[-\,\bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)\,\Delta x^\mu \Delta x^\nu\Big]^{1/2} \;\textit{(spacelike)}}$$

Thus the wordwise objects reduce to the proper–time/proper–distance constructed from $\bar{g}^{\mathrm{Lor}} = \bar{g} + \bar{t}^{\mathrm{sym}}$ *under endogenous evaluation.*

Definition 5.105 (Path versions with operational transport)**.** For a C^1 curve $\gamma : [0,1] \to \mathcal{X}_q$ and transported tangent word $\dot{\boldsymbol{\gamma}}^\mu(s)$ (via the operational connection), define

$$\mathbf{T}_{\mathrm{geom}}[\gamma] := \int_0^1 ds\, \sqrt{\dot{\boldsymbol{\gamma}}^\mu(s) \;\odot\; \mathbf{g}^{\mathrm{Lor}}_{\mu\nu}\big(\gamma(s)\big) \;\odot\; \dot{\boldsymbol{\gamma}}^\nu(s)} \quad \text{(timelike)}, \tag{5.96}$$

$$\mathbf{L}_{\mathrm{geom}}[\gamma] := \int_0^1 ds\, \sqrt{-\,\dot{\boldsymbol{\gamma}}^\mu(s) \;\odot\; \mathbf{g}^{\mathrm{Lor}}_{\mu\nu}\big(\gamma(s)\big) \;\odot\; \dot{\boldsymbol{\gamma}}^\nu(s)} \quad \text{(spacelike)}. \tag{5.97}$$

These are gauge–covariant by construction and reduce, under $\langle\!\langle\cdot\rangle\!\rangle$, to the standard GR line integrals for $\bar{g}^{\mathrm{Lor}}$.

Remark 5.106 (Projection choices within the word calculus)**.** *Lorentzian merge (default):* use $\mathbf{g}^{\mathrm{Lor}} = \mathbf{g} \oplus \mathbb{S}[\mathbf{t}]$ to retain the T-even temporal curvature in the cones. *Pure–metric projection:* set $\mathbf{t} \mapsto 0$ before evaluation; then $\mathbf{g}^{\mathrm{Lor}} \to \mathbf{g}$ and all expressions collapse to the purely metric word formulas. Both routes are encoded entirely wordwise; the operational choice of when to drop $\mathbf{t}$ is external/phenomenological (cf. Definitions 5.89 and 5.108).

One–line summary. Proper time and distance in the holosymmetric calculus are the positive roots of the Lorentzian quadratic $\mathcal{Q} = \Delta\mathbf{x}^{\mu} \odot \mathbf{g}^{\mathrm{Lor}}_{\mu\nu} \odot \Delta\mathbf{x}^{\nu}$, restricted by causal word–predicates; endogenous expectation yields the usual GR expressions built from $\bar{g}^{\mathrm{Lor}} = \bar{g} + \bar{t}^{\mathrm{sym}}$.

5.3.11 Holosymmetric Wordwise Metric Projection

Definition 5.107 (HS relational word–manifold)**.** On each fibre at $x \in \mathcal{X}_q$ let $\mathbf{g}_{\mu\nu}(x)$ and $\mathbf{t}_{\mu\nu}(x)$ be rank–two *words* (metric and temporal channels), gauge–covariant under fibre unitaries. The expectation–level objects are obtained by the endogenous evaluation $\langle\!\langle \cdot \rangle\!\rangle := \langle \mathring{\mathbf{\Psi}} | \cdot | \mathring{\mathbf{\Psi}} \rangle$. (See also the Lorentzian merge of Section 5.3.10.)

Definition 5.108 (HS metric–projection functor)**.** Define the holosymmetric *metric projection* (forgetful functor on the temporal channel)

$$\mathrm{Proj}^{\mathrm{HS}}_{\mathbf{g}} : \ (\mathcal{X}_q,\ \mathbf{g}_{\mu\nu},\ \mathbf{t}_{\mu\nu}) \ \longmapsto \ \big(\mathcal{X}_q,\ \mathbf{g}_{\mu\nu},\ \mathbf{d}^{(\mathrm{proj})},\ \mathbf{t}^{(\mathrm{proj})}\big),$$

by discarding $\mathbf{t}$ in the quadratic forms. For an infinitesimal displacement word $\Delta\mathbf{x}^{\mu} := \mathbf{x}'^{\mu} - \mathbf{x}^{\mu}$, set the *projected quadratic word*

$$\mathcal{Q}_{\mathbf{g}}(x;\Delta) := \Delta\mathbf{x}^{\mu} \ \odot \ \mathbf{g}_{\mu\nu}(x) \ \odot \ \Delta\mathbf{x}^{\nu}.$$

The projected proper–time and proper–distance *words* are defined piecewise (positive branch on the regulated fibre, SOT):

$$\mathbf{t}^{(\mathrm{proj})}(x,x') := \begin{cases} \sqrt{\mathcal{Q}_{\mathbf{g}}(x;\Delta)}\,, & \mathcal{Q}_{\mathbf{g}}(x;\Delta) > 0, \\ 0, & \text{otherwise.} \end{cases}$$

$$\mathbf{d}^{(\mathrm{proj})}(x,x') := \begin{cases} \sqrt{-\,\mathcal{Q}_{\mathbf{g}}(x;\Delta)}\,, & \mathcal{Q}_{\mathbf{g}}(x;\Delta) < 0, \\ 0, & \text{otherwise.} \end{cases}$$

Proposition 5.109 (Endogenous evaluation reproduces the classical projection)**.** *With* $\bar{g}_{\mu\nu} := \langle\!\langle \mathbf{g}_{\mu\nu} \rangle\!\rangle$*, for small* Δx^{μ} *one has*

$$\langle\!\langle \mathbf{t}^{(\mathrm{proj})}(x,x') \rangle\!\rangle = \begin{cases} \big[\ \bar{g}_{\mu\nu}(x)\,\Delta x^{\mu}\Delta x^{\nu}\ \big]^{1/2}, & \bar{g}_{\mu\nu}\Delta x^{\mu}\Delta x^{\nu} > 0, \\ 0, & \textit{otherwise}, \end{cases}$$

$$\langle\!\langle \mathbf{d}^{(\mathrm{proj})}(x,x') \rangle\!\rangle = \begin{cases} \big[\ -\ \bar{g}_{\mu\nu}(x)\,\Delta x^{\mu}\Delta x^{\nu}\ \big]^{1/2}, & \bar{g}_{\mu\nu}\Delta x^{\mu}\Delta x^{\nu} < 0, \\ 0, & \textit{otherwise.} \end{cases}$$

Thus $\mathrm{Proj}^{\mathrm{HS}}_{\mathbf{g}}$ *reduces exactly to the classical* $(\mathcal{X}_q, \bar{g}_{\mu\nu})$ *projection with the standard GR interval rules.*

Proposition 5.110 (Functorial properties)**.** *(a)* Gauge covariance. *For any fibre unitary* $U(x)$, $\mathbf{g} \mapsto U\mathbf{g}U^\dagger$ *implies* $\mathcal{Q}_\mathbf{g} \mapsto \mathcal{Q}_\mathbf{g}$*; hence* $\mathrm{Proj}^{\mathrm{HS}}_\mathbf{g}$ *is gauge–covariant and* $\langle\langle\cdot\rangle\rangle$*–gauge invariant.*

(b) Transport compatibility. *With operational transport* U_γ *along* γ*, projection commutes with pullback:* $\mathrm{Proj}^{\mathrm{HS}}_\mathbf{g} \circ U_\gamma = U_\gamma \circ \mathrm{Proj}^{\mathrm{HS}}_\mathbf{g}$.

(c) Idempotence. $\mathrm{Proj}^{\mathrm{HS}}_\mathbf{g} \circ \mathrm{Proj}^{\mathrm{HS}}_\mathbf{g} = \mathrm{Proj}^{\mathrm{HS}}_\mathbf{g}$ *(the temporal channel is forgotten once).*

Remark 5.111 (Relation to the Lorentzian–merge route)**.** The projection uses only $\mathbf{g}$. By contrast, the merge uses $\mathbf{g}^{\mathrm{Lor}} = \mathbf{g} \oplus \mathbb{S}[\mathbf{t}]$ *via* Definitions 5.87 and 5.101. Whenever the symmetric temporal word is $\mathbf{g}$–positive in the quadratic sense $\big(\Delta\mathbf{x}^\mu \odot \mathbb{S}[\mathbf{t}]_{\mu\nu} \odot \Delta\mathbf{x}^\nu \geq 0\big)$, timelike sets under the merge contain those of the projection; if it is negative, the converse may occur. No extra assumptions are required for the pure–metric projection itself.

Definition 5.112 (Projected classical spacetime (wordwise statement))**.** The image of $\mathrm{Proj}^{\mathrm{HS}}_\mathbf{g}$ under endogenous expectation is

$$\boxed{\mathcal{M}^{(\mathrm{proj})}_{\mathrm{geom}} := \big(\mathcal{X}_q,\ \bar{g}_{\mu\nu}\big)},$$

a pseudo–Riemannian manifold in the GR sense; its operational intervals are supplied by $\langle\langle\mathbf{t}^{(\mathrm{proj})}\rangle\rangle$ and $\langle\langle\mathbf{d}^{(\mathrm{proj})}\rangle\rangle$ as above.

One–line summary. The metric projection in the holosymmetric calculus is the *temporal–forgetful* functor that forms quadratic words with $\mathbf{g}$ alone; endogenous evaluation reproduces the classical $(\mathcal{X}_q, \bar{g}_{\mu\nu})$ spacetime and its GR interval rules, while remaining fully gauge– and transport–compatible at the word level.

5.3.12 The Emergent Chronogeometric Manifold

Having established the operator-theoretic substrate and its expectation-level fields (Section 5.2.10 and Definition 5.26), we arrive at the macroscopically salient layer: the observable spacetime. Denoted $\mathcal{M}_{\mathrm{geom}}$, this is the classical limit in which operator-valued chronogeometry crystallises into a pseudo-Riemannian manifold with determinate spatial and temporal intervals ([7, 146]; cf. Definitions 5.122 and 5.150).

Definition 5.113 (Emergent geometric manifold (Lorentzian-merge presentation))**.** Let $\bar{g}^{\mathrm{Lor}}_{\mu\nu} := \bar{g}_{\mu\nu} + \bar{t}^{\mathrm{sym}}_{\mu\nu}$ be the effective metric from Definition 5.77. The *emergent chronogeometric manifold* is the quadruple

$$\boxed{\mathcal{M}_{\mathrm{geom}} := \Big(\mathcal{X}_q,\ \bar{g}^{\mathrm{Lor}}_{\mu\nu}(x),\ d_{\mathrm{geom}}(x,x'),\ t_{\mathrm{geom}}(x,x')\Big)}, \tag{5.98}$$

where:

- $\mathcal{X}_q$ is the emergent configuration manifold of distinguishable PQSs (Section 4.19.3);
- $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ fixes causal cones and the Levi–Civita geometry (Definitions 5.122 and 5.150);

- t_{geom} and d_{geom} are the proper time and proper distance for timelike/spacelike infinitesimal separations computed with $\bar{g}^{\text{Lor}}$ (Definitions 5.95 and 5.96; [7]).

Remark 5.114 (Alternative presentation: pure-metric projection)**.** A purely classical presentation is obtained by the *metric projection* that drops $\bar{t}_{\mu\nu}$ (Section 5.3.9):

$$\mathcal{M}_{\text{geom}}^{(\text{proj})} := \text{Proj}_{\bar{g}}(\mathcal{M}_{\text{rel}}) = \left(\mathcal{X}_q,\ \bar{g}_{\mu\nu},\ d_{\text{geom}}^{(\text{proj})},\ t_{\text{geom}}^{(\text{proj})}\right) \quad \text{(Definition 5.97)}.$$

Both routes are expectation-level constructions (Section 5.2.10). They coincide when $\bar{t}^{\text{sym}} = 0$ or is aligned so that cone tilts vanish (e.g., FLRW specialisation; Section 5.4.7). Operationally, $\bar{g}^{\text{Lor}}$ supplies the *hard* causal layer, while $\bar{t}_{\mu\nu}$ governs *soft* intra-cone phase transport (Sections 5.4.3 and 5.4.4).

Remark 5.115 (Interpretation and ontological status)**.** $\mathcal{M}_{\text{geom}}$ is the observer-accessible, coarse-grained geometry: spatial lengths, temporal durations, and causal order are read from the effective metric (merge) or the projected metric (projection), recovering the GR description in appropriate limits [7, 146, 244, 11]. While $\bar{t}_{\mu\nu}$ is absent from the metric in the pure projection, its symmetric part contributes to $\bar{g}^{\text{Lor}}$ and its antisymmetric part encodes intra-cone phase holonomy, relevant in high-coherence regimes (early cosmology, near-horizon physics), without altering the light-cone boundary (Section 5.4.3 and Definition 5.122).

5.3.13 Holosymmetric Wordwise Emergent Chronogeometric Manifold

Definition 5.116 (HS Lorentzian–merge word–field)**.** On each fibre at $x \in \mathcal{X}_q$, let $\mathbf{g}_{\mu\nu}(x)$ and $\mathbf{t}_{\mu\nu}(x)$ be rank–two *holosymmetric words* (metric and temporal sectors), gauge–covariant under fibre unitaries. Let $\mathbb{S}[\cdot]$ denote the port–flip symmetriser (cf. Definition 5.87). Define the Lorentzian–merge word–field by the Hadamard sum

$$\boxed{\mathbf{g}_{\mu\nu}^{\text{Lor}}(x) \ := \ \mathbf{g}_{\mu\nu}(x) \ \oplus \ \mathbb{S}\big[\mathbf{t}_{\mu\nu}(x)\big].} \tag{5.99}$$

This coincides with the Lorentzian word of Definition 5.101. Under endogenous evaluation (Born map) $\langle\!\langle\cdot\rangle\!\rangle := \langle\mathring{\mathbf{\Psi}}|\cdot|\mathring{\mathbf{\Psi}}\rangle$,

$$\langle\!\langle \mathbf{g}_{\mu\nu} \rangle\!\rangle = \bar{g}_{\mu\nu}, \qquad \langle\!\langle \mathbb{S}[\mathbf{t}_{\mu\nu}] \rangle\!\rangle = \bar{t}_{\mu\nu}^{\text{sym}},$$

hence $\langle\!\langle \mathbf{g}_{\mu\nu}^{\text{Lor}} \rangle\!\rangle = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}^{\text{sym}} =: \bar{g}_{\mu\nu}^{\text{Lor}}$.

Definition 5.117 (HS emergent intervals and cones)**.** For an infinitesimal displacement word $\Delta\mathbf{x}^\mu := \mathbf{x}'^\mu - \mathbf{x}^\mu$ at x, define the Lorentzian quadratic word

$$\mathcal{Q}_{\mathbf{g}^{\text{Lor}}}(x;\Delta) := \Delta\mathbf{x}^\mu \ \odot \ \mathbf{g}_{\mu\nu}^{\text{Lor}}(x) \ \odot \ \Delta\mathbf{x}^\nu.$$

Let $\Pi_{\text{time}} := \mathbf{1}_{\{\mathcal{Q}_{\mathbf{g}^{\text{Lor}}} > 0\}}$ and $\Pi_{\text{space}} := \mathbf{1}_{\{\mathcal{Q}_{\mathbf{g}^{\text{Lor}}} < 0\}}$. The HS proper–time and proper–distance *words* (agreeing with Definition 5.103) are

$$\boxed{\mathbf{t}_{\text{geom}}(x,x') := \sqrt{\ \mathcal{Q}_{\mathbf{g}^{\text{Lor}}}(x;\Delta)\ }\ \cdot \Pi_{\text{time}}, \qquad \mathbf{d}_{\text{geom}}(x,x') := \sqrt{\ -\mathcal{Q}_{\mathbf{g}^{\text{Lor}}}(x;\Delta)\ }\ \cdot \Pi_{\text{space}},} \tag{5.100}$$

with square roots taken in the positive branch on the spectrally regulated fibre (SOT). The null word–cone at x is $\mathcal{N}_x := \{\Delta\mathbf{x} : \ \mathcal{Q}_{\mathbf{g}^{\text{Lor}}}(x;\Delta) = 0\}$.

Definition 5.118 (HS emergent chronogeometric manifold)**.** The *holosymmetric emergent chronogeometric manifold* is the quadruple

$$\boxed{\mathcal{M}^{\mathrm{HS}}_{\mathrm{geom}} := \Big(\mathcal{X}_q,\ \mathbf{g}^{\mathrm{Lor}}_{\mu\nu}(x),\ \mathbf{d}_{\mathrm{geom}}(x,x'),\ \mathbf{t}_{\mathrm{geom}}(x,x') \Big),} \tag{5.101}$$

with gauge–covariant transport inherited from the operational transporter U_γ along curves γ.

Proposition 5.119 (Endogenous equivalence with the classical presentation)**.** *Applying* $\langle\!\langle\cdot\rangle\!\rangle$ *to* (5.101) *yields*

$$\boxed{\langle\!\langle \mathcal{M}^{\mathrm{HS}}_{\mathrm{geom}} \rangle\!\rangle = \Big(\mathcal{X}_q,\ \bar{g}^{\mathrm{Lor}}_{\mu\nu}(x),\ d_{\mathrm{geom}}(x,x'),\ t_{\mathrm{geom}}(x,x') \Big) =: \mathcal{M}_{\mathrm{geom}}\ ,}$$

where $\bar{g}^{\mathrm{Lor}}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}^{\mathrm{sym}}_{\mu\nu}$*, and* $(d_{\mathrm{geom}}, t_{\mathrm{geom}})$ *coincide with the GR interval rules computed with* $\bar{g}^{\mathrm{Lor}}$ *(cf. Proposition 5.104).*

Proposition 5.120 (Gauge/transport compatibility and idempotence)**. (i) Gauge covariance.** *For any fibre unitary* $U(x)$*,* $\mathbf{g}^{\mathrm{Lor}} \mapsto U\,\mathbf{g}^{\mathrm{Lor}}\,U^\dagger$*. Consequently* $\mathcal{Q}_{\mathbf{g}^{\mathrm{Lor}}}$ *and the cone predicates are invariant under similarity, and endogenous expectations are gauge–invariant.*

(ii) Transport compatibility.
For the operational transporter U_γ*, pullback along* γ *yields* $U_\gamma^*\mathbf{g}^{\mathrm{Lor}} = \mathbf{g}^{\mathrm{Lor}}$*, hence* $U_\gamma^*\mathbf{t}_{\mathrm{geom}} = \mathbf{t}_{\mathrm{geom}}$ *and* $U_\gamma^*\mathbf{d}_{\mathrm{geom}} = \mathbf{d}_{\mathrm{geom}}$*.*

(iii) Projection idempotence.
Performing the metric projection ($\mathbf{g}^{\mathrm{Lor}} \rightsquigarrow \mathbf{g}$) *and then re-forming* (5.99) *with* $\mathbb{S}[\mathbf{t}] = 0$ *leaves the result unchanged; likewise at expectation level.*

Remark 5.121 (Levi–Civita and curvature words (optional))**.** Define the HS Christoffel word by port–symmetrised derivatives,

$$\mathbf{\Gamma}^{\rho}{}_{\mu\nu} := \tfrac{1}{2}\,\mathbf{g}^{\mathrm{Lor},\rho\lambda}\Big(\partial_\mu \mathbf{g}^{\mathrm{Lor}}_{\lambda\nu} + \partial_\nu \mathbf{g}^{\mathrm{Lor}}_{\lambda\mu} - \partial_\lambda \mathbf{g}^{\mathrm{Lor}}_{\mu\nu}\Big),$$

with derivatives taken in SOT on the regulated band. Geodesic, Riemann, Ricci and scalar–curvature words follow as usual; endogenous evaluation reproduces the standard Levi–Civita objects of $\bar{g}^{\mathrm{Lor}}$.

One-line summary. $\mathcal{M}^{\mathrm{HS}}_{\mathrm{geom}} = \big(\mathcal{X}_q, \mathbf{g}^{\mathrm{Lor}}, \mathbf{d}_{\mathrm{geom}}, \mathbf{t}_{\mathrm{geom}}\big)$; taking endogenous expectations yields precisely your emergent chronogeometric manifold $\mathcal{M}_{\mathrm{geom}}$ with Lorentzian–merge metric and GR interval rules.

The Chronogeometric Manifold Genesis Chain

The diagram below encapsulates the complete hierarchy of emergence within the IQR framework. It illustrates the logical and ontological path from the foundational, pre-geometric quantum substrate to the familiar classical spacetime manifold. We call this sequence the *Geometro-Temporal Manifold Genesis Chain.*

$$(\mathcal{C}_d,\ |\mathring{\mathbf{\Psi}}\rangle) \overset{\text{Quotient}}{\rightsquigarrow} \mathcal{X}_q \overset{\text{Fibers}}{\rightsquigarrow} \mathfrak{F}_x$$
$$\overset{\text{2nd Variation}}{\rightsquigarrow} (\hat{g}_{\mu\nu}(x),\ \hat{t}_{\mu\nu}(x)) \overset{\text{Expectation}}{\rightsquigarrow} (\bar{g}_{\mu\nu}(x),\ \bar{t}_{\mu\nu}(x))$$
$$\overset{\text{Classical Limit}}{\rightsquigarrow} \mathcal{M}_{\text{geom}}$$

Emergence chain (layered construction). This chain organises how macroscopic physical reality arises, layer by layer, from the quantum–relational substrate developed in Chapter 6.

1. *Foundation (substrate and coherence).* Begin with the pre-geometric continuum $\mathcal{C}_d$ of structured potential loci and the Primordial Quantum Singularities (PQSs) $\mathfrak{B}_x$ as the basic ontic units (Definition 1.25 and Section 5.4.6). Global quantum coherence is provided by the *Quantum Holo–State* $|\mathring{\mathbf{\Psi}}\rangle$, which supplies the state in which all operator fields are evaluated (Section 5.2.10; see also [287]).

2. *Emergence of distinguishability (quotient).* A *quotient identification* collapses the substrate by identifying loci whose relational observables vanish, yielding the *Emergent Quantum Configuration Space* $\mathcal{X}_q$ as equivalence classes of indistinguishable PQSs—the minimal arena of distinguishable events (Section 4.19.3 and Proposition 1.28).

3. *Encoding relationality (fibres/bundle).* Over $\mathcal{X}_q$ one equips each point x with a *relational fibre* $\mathfrak{F}_x$ carrying the raw, operator-valued data of distinguishability and temporal phase-flow relative to all other points (the *relational phase–geometry bundle*; cf. Sections 4.19.3 and 6.1.3). This is the kinematic repository of the operator geometry prior to taking expectations.

4. *Rise of quantum geometry (second variations).* A *second-variation* construction on the fibre data defines the operator-valued chronogeometric tensors $(\hat{g}_{\mu\nu}(x), \hat{t}_{\mu\nu}(x))$, giving the *operator-level relational geometry* $(\hat{\mathcal{M}})$ in which curvature of distance and of time are themselves quantum operators (Definitions 5.25 and 5.26). These ultra-local operators are supported on PQS fibres (Lemma 4.602).

5. *Classical projection (expectations; triadic structure).* Taking expectation values in $|\mathring{\mathbf{\Psi}}\rangle$ produces smooth classical fields $(\bar{g}_{\mu\nu}, \bar{t}_{\mu\nu})$ on $\mathcal{X}_q$, assembling the *expectation-level relational manifold*
$$\mathcal{M}_{\text{rel}} := \big(\mathcal{X}_q,\ \bar{g}_{\mu\nu},\ \bar{t}_{\mu\nu}\big),$$
with the canonical triadic split in each sector (Section 5.2.10 and Definition 5.54; cf. [244, 11]).

6. *Observer-accessible spacetime (classical limit).* Two presentation routes yield the macroscopic manifold perceived by observers:

   (a) **Lorentzian merge (hard causal layer).** Define $\bar{g}^{\text{Lor}}_{\mu\nu} := \bar{g}_{\mu\nu} + \bar{t}^{\text{sym}}_{\mu\nu}$ and take $\mathcal{M}_{\text{geom}} = (\mathcal{X}_q, \bar{g}^{\text{Lor}}_{\mu\nu})$ with proper intervals $t_{\text{geom}}, d_{\text{geom}}$ and cones from $\bar{g}^{\text{Lor}}$ (Definitions 5.77, 5.95, 5.96, 5.122 and 5.150; [7, 146]).

(b) **Metric projection (pure classical presentation).** Drop $\bar{t}_{\mu\nu}$ and retain the pseudo-Riemannian structure $(\mathcal{X}_q, \bar{g}_{\mu\nu})$ with standard GR intervals (Section 5.3.9 and Definition 5.97).

The two presentations coincide in symmetric limits (e.g., aligned/vanishing $\bar{t}^{\mathrm{sym}}$; FLRW specialisation), ensuring recovery of GR/QFT microcausality and stability (Section 5.4.7; [244, 11]).

In summary,

$$\boxed{\mathcal{C}_d \longrightarrow \mathcal{X}_q \longrightarrow \{\mathfrak{F}_x\} \longrightarrow (\hat{g}, \hat{t}) \longrightarrow (\bar{g}, \bar{t}) \longrightarrow \mathcal{M}_{\mathrm{geom}}}$$

is the relational-to-geometric pipeline: operator chronogeometry → expectation tensors → observer-level spacetime. The *hard* causal boundary is set by the effective metric (merge/projection), while the *soft* intra-cone phase dynamics—encoded by $\bar{t}_{\mu\nu}$—remains available for holosymmetric tomography without altering the cone assignment (Sections 5.4.3 and 5.4.4, Proposition 5.60, and Corollary 5.126).

5.3.14 The Nature of Spacetime: An Emergent, Relational Tapestry

A central challenge is to reconcile the smooth spacetime of general relativity with the probabilistic, relational character of quantum theory. In the present framework, spacetime is *not* fundamental but a coarse-grained, classical *emergent* structure arising from a deeper, pre-geometric, quantum–relational substrate; geometry is thus a property of the global quantum state rather than a background arena ([3]; cf. Section 5.2.10 and Definition 5.150 and [7, 146] for the classical benchmark).

Substrate and coherence. At the most basic level lives the $\hat{\Omega}$-Continuum: a smooth manifold of operator-valued singularities, the *Primordial Quantum Singularities* (PQSs), which supply the ontic loci of the theory (Definition 1.25 and Section 5.4.6). Physical instantiation is provided by the globally coherent, non-factorisable *Quantum Holo–State* $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$, which entangles the ensemble of PQSs into a relational network (Section 5.2.10; see also [287]).

Operator chronogeometry and expectation fields. Relational data over the emergent configuration space $\mathcal{X}_q$ (Section 4.19.3) give rise—via second variations—to the operator-valued chronogeometric tensors $(\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu})$ (Definitions 5.25 and 5.26). Taking expectations in $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$ produces smooth classical fields $(\bar{g}_{\mu\nu}, \bar{t}_{\mu\nu})$ on $\mathcal{X}_q$ (Section 5.2.10), from which the effective metric for causal kinematics is assembled as $\bar{g}^{\mathrm{Lor}}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}^{\mathrm{sym}}_{\mu\nu}$ (Definition 5.77).[2]

Spacetime as expectation of distinguishability. In this sense,

$$\bar{g}_{\mu\nu}(x) = \langle \overset{\circ}{\boldsymbol{\Psi}} | \, \hat{g}_{\mu\nu}(x) \, | \overset{\circ}{\boldsymbol{\Psi}} \rangle$$

(cf. Section 5.2.10 and Definition 5.26), so spacetime geometry is the *classical expectation of quantum distinguishability*. The observable manifold $\mathcal{M}_{\mathrm{geom}}$ (Lorentzian-merge presentation)

[2] A pure classical presentation is obtained by the metric projection that discards $\bar{t}_{\mu\nu}$ (Section 5.3.9); both routes recover standard GR intervals in appropriate limits [7, 146].

then reads $\mathcal{M}_{\text{geom}} = (\mathcal{X}_q, \bar{g}^{\text{Lor}}_{\mu\nu}, d_{\text{geom}}, t_{\text{geom}})$ (Definitions 5.95, 5.96 and 5.113), with light-cone kinematics fixed by $\bar{g}^{\text{Lor}}$ (Definition 5.122) and intra-cone phase transport governed by $\bar{t}_{\mu\nu}$ (Sections 5.4.3 and 5.4.4).

Analogy and classicality. The relationship between the $\hat{\Omega}$-Continuum and spacetime mirrors that between microscopic quantum states of H_2O molecules and a macroscopic ocean wave: the wave is real yet emergent from underlying quantum interactions. Likewise, smooth spacetime with its metric and causal order is a statistical manifestation of an entangled, quantum–relational network of PQSs (cf. decoherence/emergence discussions [247, 244, 11]). In this view, the familiar pseudo-Riemannian geometry is the "shadow'' cast by the Quantum Holo–State on the manifold of relational observables, offering a coherent route to reconcile quantum theory with gravitational kinematics by treating spacetime as a quantum–statistical emergent phenomenon.

5.4 Causal Structures on the Chronogeometric Manifold

The causal architecture of the emergent universe, within the formalism of Interdependent Quantum Reality (IQR), is not a pre-ordained, monolithic structure. It is a composite and dynamical phenomenon that arises from the two fundamental tensor fields derived from the Quantum Holo-State: the emergent metric tensor, $\bar{g}_{\mu\nu}$, and the emergent temporal curvature tensor, $\bar{t}_{\mu\nu}$. This results in a sophisticated, two-layered causal structure: a definitive, "hard" kinematic boundary defined by a composite Lorentzian metric, and a "soft," quantum-dynamical texture that modulates the flow of coherence within that boundary.

5.4.1 The Effective Light Cone and its Primordial Origin

The absolute boundary of causal influence—that which separates timelike, spacelike, and null-separated directions—is not an a priori structure imposed upon the manifold. Rather, in the IQR framework, it is an emergent construct derived from the Chronogeometric Tensor and ontologically rooted in the Primordial Quantum Singularity (PQS). The *effective light cone* is thus both a geometric object and a manifestation of quantum-relational indistinguishability.

The governing tensorial structure is the *effective Lorentzian metric*, denoted $\bar{g}^{\text{Lor}}_{\mu\nu}(x)$, obtained by symmetrizing the Chronogeometric Tensor as defined in Section 5.3.4.

Definition 5.122 (Effective Light Cone Structure and its Primordial Origin)**.** Let the effective Lorentzian metric on the emergent chronogeometric manifold $\mathcal{M}_{\text{geom}} \equiv \mathcal{M}_{\text{em}}^{(\mathring{\Psi})}$ be given by

$$\bar{g}^{\text{Lor}}_{\mu\nu}(x) := \bar{g}_{\mu\nu}(x) + \bar{t}^{\text{sym}}_{\mu\nu}(x),$$

where $\bar{t}^{\text{sym}}_{\mu\nu}(x) := \frac{1}{2}(\bar{t}_{\mu\nu}(x) + \bar{t}_{\nu\mu}(x))$ is the symmetric part of the emergent temporal tensor, and $\bar{g}_{\mu\nu}(x)$ is the expectation-level spatial metric.

The apex of this causal structure at any point $x \in \mathcal{X}_q$ is not a coordinate artifact, but the ontological singularity $\mathfrak{B}_x$ itself (Definition 1.25), from which the causal structure emanates.

A tangent vector $V^\mu \in T_x\mathcal{M}_{\text{geom}}$ is classified as:

- **Timelike**, if $\bar{g}^{\text{Lor}}_{\mu\nu}(x)V^\mu V^\nu > 0$,

- **Spacelike**, if $\bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)V^\mu V^\nu < 0$,
- **Null (lightlike)**, if $V^\mu \neq 0$ and $\bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)V^\mu V^\nu = 0$.

The set of all nonzero null vectors defines the *Effective Light Cone* at the singularity $\mathfrak{B}_x$,

$$\mathcal{V}_\pm(\mathfrak{B}_x) := \left\{ V^\mu \in T_x\mathcal{M}_{\mathrm{geom}} \;\middle|\; \bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)V^\mu V^\nu = 0,\ V^\mu \neq 0 \right\},$$

which constitutes the absolute boundary for all causal propagation originating from $\mathfrak{B}_x$.

Definition 5.123 (Holosymmetric effective Lorentzian tensors and cones)**.** Apply the holosymmetric framework (cf. Definition 5.56 and Proposition 5.57) to the chronogeometric data entering Definition 5.122. For two law branches $k \in \{0,1\}$, define the branchwise expectation tensors

$$\bar{g}^{\mathrm{Lor}}_{(k)\mu\nu}(x) := \bar{g}_{(k)\mu\nu}(x) + \bar{t}^{\mathrm{sym}}_{(k)\mu\nu}(x), \qquad \bar{t}^{\mathrm{sym}}_{(k)\mu\nu} := \tfrac{1}{2}\left(\bar{t}_{(k)\mu\nu} + \bar{t}_{(k)\nu\mu}\right),$$

with apex at the same ontological singularity $\mathfrak{B}_x$ (Definition 1.25). Let $\mathcal{V}^{(k)}_\pm(\mathfrak{B}_x)$ be the branchwise null cones defined as in Definition 5.122 but using $\bar{g}^{\mathrm{Lor}}_{(k)}$.

Define the *operational holosymmetric* spacelike relation and cones by

$$\mathrm{Spacelike}_\cap(x,y) \;:\Longleftrightarrow\; (x,y) \text{ are spacelike for both } \bar{g}^{\mathrm{Lor}}_{(0)} \text{ and } \bar{g}^{\mathrm{Lor}}_{(1)},$$

$$J^\pm_\cup(x) \;:=\; J^\pm_{\bar{g}^{\mathrm{Lor}}_{(0)}}(x) \;\cup\; J^\pm_{\bar{g}^{\mathrm{Lor}}_{(1)}}(x), \qquad J^\pm_\cap(x) \;:=\; J^\pm_{\bar{g}^{\mathrm{Lor}}_{(0)}}(x) \;\cap\; J^\pm_{\bar{g}^{\mathrm{Lor}}_{(1)}}(x).$$

Proposition 5.124 (No–law–signalling and branchwise microcausality)**.** *Let $\hat{\mathfrak{O}}(f)$ and $\hat{\mathfrak{O}}(h)$ be local observables supported in regions $A, B \subset \mathcal{M}_{\mathrm{geom}}$. If A and B are Spacelike$_\cap$, then for the holosymmetric protocol (any control phase φ and coherence angle θ)*

$$[\hat{\mathfrak{O}}(f), \hat{\mathfrak{O}}(h)] = 0,$$

i.e. microcausality holds operationally. Moreover, the reachable domain for signals from A is bounded by $J^+_\cup(A)$; holosymmetry cannot enlarge causal reach beyond the union of branchwise futures.

Idea of proof. Local commutativity holds separately in each branch with respect to $J^\pm_{\bar{g}^{\mathrm{Lor}}_{(k)}}$. Holosymmetric control unitaries act as direct sums over the branches (cf. Lemma 5.55) and cannot create commutators where each summand vanishes; phases (φ,θ) only weight symmetric/antisymmetric response functions (Proposition 5.60) without changing branchwise cone assignments. Signal paths must lie in a branch future; allowing both branches yields $J^+_\cup$. □

Proposition 5.125 (Cone stability under small holosymmetric deformations)**.** *Let $\Delta\bar{g}^{\mathrm{Lor}} := \bar{g}^{\mathrm{Lor}}_{(1)} - \bar{g}^{\mathrm{Lor}}_{(0)}$. For any $\bar{g}^{\mathrm{Lor}}_{(0)}$–null vector V at x,*

$$\left|\bar{g}^{\mathrm{Lor}}_{(1)}(V,V)\right| = \left|\Delta\bar{g}^{\mathrm{Lor}}(V,V)\right| \le \|\Delta\bar{g}^{\mathrm{Lor}}\|_x \, \|V\|^2_x,$$

so the null directions tilt by an amount controlled by $\|\Delta\bar{g}^{\mathrm{Lor}}\|_x$. In the canonical holosymmetric pair where only the temporal sector flips sign (Definition 5.56), the tilt is set by $\|\Delta\bar{t}^{\mathrm{sym}}\|_x$.

Corollary 5.126 (Phase invariance of cone assignment)**.** *The sets $J^{\pm}_{\cup}$ and Spacelike$_{\cap}$ are independent of the control/readout phases (φ, θ) used in holosymmetric tomography; these phases affect interference amplitudes (responses of $\hat{\Sigma}$) but not the branchwise effective metrics that define the cones.*

Corollary 5.127 (Aligned-symmetry case and FLRW limit)**.** *If $\bar{t}_{(k)\mu\nu}$ is aligned so that $\bar{t}^{\mathrm{sym}}_{(k)}$ does not tilt the null structure (e.g. homogeneous/isotropic alignment), then $\bar{g}^{\mathrm{Lor}}_{(0)} = \bar{g}^{\mathrm{Lor}}_{(1)}$, hence $J^{\pm}_{\cup} = J^{\pm}_{\cap} = J^{\pm}_{\bar{g}^{\mathrm{Lor}}}$ and the holosymmetric lift reduces to the single–cone case of Definition 5.122.*

Remark 5.128 (Apex and operational interpretation)**.** The apex remains the ontological singularity $\mathfrak{B}_x$ in both branches, and operationally the "safe spacelike" relation is the intersection criterion Spacelike$_{\cap}$ used in Proposition 5.124. Time-reversal superpositions (U vs. $U^{\dagger}$) preserve the cone assignment since the effective metrics coincide in that case (cf. Remark 5.62).

Interpretation. The light-cone definition Definition 5.122 unites geometric precision with ontological depth: the causal cone at a spacetime point is a structure that *emanates from* the Primordial Quantum Singularity itself. Within $\mathfrak{B}_x$, all relational observables vanish (Proposition 1.28), making it the indivisible origin of causal propagation (Theorem 5.155). The inclusion of the symmetric temporal sector $\bar{t}^{\mathrm{sym}}_{\mu\nu}(x)$ in the effective Lorentzian metric shows that the light-cone geometry—its shape, orientation, and causal-velocity structure—is modulated by local quantum phase dynamics. In regions of strong temporal curvature (e.g., early cosmogenesis or highly coherent quantum regimes), causal propagation can become anisotropically deformed. This effect is absent in classical General Relativity, where cones are fixed entirely by the metric sector and are insensitive to such phase-curvature contributions [7, 146].

Holosymmetric implications. Applying holosymmetry to the effective cone structure (Definition 5.123) produces a conservative, branchwise causal calculus. Operational microcausality is protected on the *intersection* of spacelike separations across branches: if two regions are Spacelike$_{\cap}$, their local observables commute even in the presence of holosymmetric control (Proposition 5.124). Possible influence is bounded by the *union* of branchwise futures $J^{\pm}_{\cup}$, so holosymmetry does not enlarge causal reach beyond what at least one history already permits; it alters only interference in response functions within that domain (Proposition 5.124). Control phases (φ, θ) perform tomography—separating T-even (anticommutator) from T-odd (commutator) responses—yet leave the cone assignment invariant (Corollary 5.126). The null directions themselves vary *continuously* under holosymmetric deformations, with tilts controlled by the norm of the branch difference $\Delta\bar{g}^{\mathrm{Lor}}$ (canonically by $\Delta\bar{t}^{\mathrm{sym}}$), yielding Lipschitz-type stability of causal structure and preserving well-posedness for hyperbolic dynamics (Proposition 5.125). These rules extend to arbitrary superpositions of histories by replacing pairwise intersection/union with $\bigcap_k$ and $\bigcup_k$, respectively, thereby defining a consistent causal calculus for many-branch holosymmetry.

The framework is compatible with symmetric and classical limits: when temporal sectors are aligned so that null structures coincide—e.g., in homogeneous/isotropic regimes—one recovers a single effective cone, and the holosymmetric lift reduces to Definition 5.122

(Corollary 5.127). Near horizons or other null-critical surfaces, branch differences manifest as measurable phase shifts in interference without any superluminal leakage, since accessible regions remain confined to $J^{\pm}_{\cup}$. Finally, the same Loschmidt-type overlaps used to quantify "distance in law space" serve as operational diagnostics of *cone susceptibility*: reductions in visibility correlate with projections of $\Delta\bar{g}^{\mathrm{Lor}}$ onto near-null directions, enabling inference of causal-structure shifts while maintaining microcausality and spectral stability [426, 427].

5.4.2 Holosymmetric Wordwise Effective Light Cone and PQS Apex

HS prelude (typed words and merge). Let $\mathbf{g}_{\mu\nu}(x)$ and $\mathbf{t}_{\mu\nu}(x)$ be rank–2 holosymmetric words on the distance and time channels, respectively, and let $\mathbb{S}[\cdot]$ denote the port–flip symmetriser on index pairs (cf. Definition 5.87). Define the *Lorentzian–merge word* (Hadamard sum; cf. Definition 5.101)

$$\boxed{\mathbf{g}^{\mathrm{Lor}}_{\mu\nu}(x) := \mathbf{g}_{\mu\nu}(x) \;\oplus\; \mathbb{S}[\mathbf{t}_{\mu\nu}(x)]}\,.$$

Under endogenous evaluation (Born map) $\langle\!\langle\cdot\rangle\!\rangle := \langle\mathring{\mathbf{\Psi}}|\cdot|\mathring{\mathbf{\Psi}}\rangle$, $\langle\!\langle \mathbf{g}^{\mathrm{Lor}}_{\mu\nu}\rangle\!\rangle = \bar{g}_{\mu\nu} + \bar{t}^{\mathrm{sym}}_{\mu\nu} =: \bar{g}^{\mathrm{Lor}}_{\mu\nu}$ (cf. Definition 5.80 and Proposition 5.90).

Definition 5.129 (HS effective cone and PQS apex)**.** For an infinitesimal displacement word $\Delta\mathbf{x}^{\mu}$ at $x \in \mathcal{X}_q$, set the quadratic word

$$\mathcal{Q}_{\mathbf{g}^{\mathrm{Lor}}}(x;\Delta) \;:=\; \Delta\mathbf{x}^{\mu} \;\odot\; \mathbf{g}^{\mathrm{Lor}}_{\mu\nu}(x) \;\odot\; \Delta\mathbf{x}^{\nu}.$$

Classify directions by the sign of $\mathcal{Q}_{\mathbf{g}^{\mathrm{Lor}}}$:

$$\text{timelike if } \mathcal{Q}_{\mathbf{g}^{\mathrm{Lor}}} > 0, \qquad \text{spacelike if } \mathcal{Q}_{\mathbf{g}^{\mathrm{Lor}}} < 0, \qquad \text{null if } \mathcal{Q}_{\mathbf{g}^{\mathrm{Lor}}} = 0 \text{ and } \Delta\mathbf{x} \neq 0.$$

The *HS null cone* at x is

$$\boxed{\mathcal{N}^{\mathrm{HS}}_x := \{\; \Delta\mathbf{x} \neq 0 :\; \mathcal{Q}_{\mathbf{g}^{\mathrm{Lor}}}(x;\Delta) = 0 \;\}}\,.$$

Its *apex* is the relational–null PQS class $\mathfrak{B}_x$ (the joint zero of distance/time triads), cf. Corollary 4.609 and Remark 5.52.

Proposition 5.130 (Endogenous equivalence of cones)**.** *Applying $\langle\!\langle\cdot\rangle\!\rangle$ to $\mathcal{N}^{\mathrm{HS}}_x$ yields the classical effective null cone $\{\, V^{\mu} \neq 0 :\; \bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)\, V^{\mu}V^{\nu} = 0 \,\}$.*

HS law–control version (two–law word channel).

Definition 5.131 (HS branchwise Lorentz words and cones)**.** Introduce the law–control register $\mathcal{H}_L = \mathrm{span}\{|0\rangle_L, |1\rangle_L\}$. For $k \in \{0,1\}$ define

$$\mathbf{g}^{\mathrm{Lor}}_{(k)\mu\nu}(x) := \mathbf{g}_{(k)\mu\nu}(x) \;\oplus\; \mathbb{S}[\mathbf{t}_{(k)\mu\nu}(x)],$$

and the branchwise cones $\mathcal{N}^{\mathrm{HS},(k)}_x := \{\Delta\mathbf{x} \neq 0 :\; \mathcal{Q}_{\mathbf{g}^{\mathrm{Lor}}_{(k)}}(x;\Delta) = 0\}$. Let $J^{\pm}_{\mathbf{g}^{\mathrm{Lor}}_{(k)}}(x)$ be the usual future/past sets computed with $\mathbf{g}^{\mathrm{Lor}}_{(k)}$. Define the operational relations

$$\mathrm{Spacelike}_{\cap}(x,y) \iff (x,y) \text{ are spacelike for both branches},$$

$$J^{\pm}_{\cup}(x) := J^{\pm}_{\mathbf{g}^{\mathrm{Lor}}_{(0)}}(x) \ \cup \ J^{\pm}_{\mathbf{g}^{\mathrm{Lor}}_{(1)}}(x), \qquad J^{\pm}_{\cap}(x) := J^{\pm}_{\mathbf{g}^{\mathrm{Lor}}_{(0)}}(x) \ \cap \ J^{\pm}_{\mathbf{g}^{\mathrm{Lor}}_{(1)}}(x).$$

The apex is the same PQS class $\mathfrak{B}_x$ in both branches.

Proposition 5.132 (No–law–signalling and branchwise microcausality)**.** *Let $\hat{\mathfrak{O}}(f)$ and $\hat{\mathfrak{O}}(h)$ be local word–observables supported in regions A, B. If A, B are* Spacelike$_{\cap}$*, then for any holosymmetric control (any phases φ, θ)*

$$[\hat{\mathfrak{O}}(f), \hat{\mathfrak{O}}(h)] = \mathbf{0},$$

and signals from A are confined to $J^{+}_{\cup}(A)$. Holosymmetric mixing does not enlarge causal reach beyond the union of branchwise futures.

Idea. Local commutativity holds in each branch relative to $J^{\pm}_{\mathbf{g}^{\mathrm{Lor}}_{(k)}}$. The controlled HS evolution is block–diagonal on the law register; off–diagonal coherence changes amplitudes, not cone assignments. Intersection–spacelike implies vanishing of each branch commutator, hence of the direct sum. □

Stability, phase–invariance, and aligned limits.

Proposition 5.133 (Cone stability under small HS deformations)**.** *Let $\Delta\mathbf{g}^{\mathrm{Lor}} := \mathbf{g}^{\mathrm{Lor}}_{(1)} - \mathbf{g}^{\mathrm{Lor}}_{(0)}$. For any $\mathbf{g}^{\mathrm{Lor}}_{(0)}$–null direction $\Delta\mathbf{x}$,*

$$\Big| \ \mathcal{Q}_{\mathbf{g}^{\mathrm{Lor}}_{(1)}}(x; \Delta) \ \Big| = \Big| \ \langle \Delta\mathbf{x}, \ \Delta\mathbf{g}^{\mathrm{Lor}}(x) \ \Delta\mathbf{x} \rangle \ \Big| \ \leq \ \|\Delta\mathbf{g}^{\mathrm{Lor}}(x)\|_{\mathrm{op}} \ \|\Delta\mathbf{x}\|^2.$$

In the canonical $(t \mapsto -t)$ branch pair, the tilt is governed by $\|\Delta\, \mathbb{S}[\mathbf{t}]\|_{\mathrm{op}}$.

Corollary 5.134 (Phase invariance)**.** *The sets $J^{\pm}_{\cup}$, $J^{\pm}_{\cap}$, and* Spacelike$_{\cap}$ *are independent of HS control phases (φ, θ); phases tune interference weights but not the branchwise Lorentz words.*

Corollary 5.135 (Aligned/isotropic case (single–cone reduction))**.** *If $\mathbb{S}[\mathbf{t}_{(0)}] = \mathbb{S}[\mathbf{t}_{(1)}]$ (e.g. isotropic alignment), then $\mathbf{g}^{\mathrm{Lor}}_{(0)} = \mathbf{g}^{\mathrm{Lor}}_{(1)}$ and*

$$J^{\pm}_{\cup} = J^{\pm}_{\cap} = J^{\pm}_{\mathbf{g}^{\mathrm{Lor}}}, \qquad \mathcal{N}^{\mathrm{HS},(0)}_x = \mathcal{N}^{\mathrm{HS},(1)}_x = \mathcal{N}^{\mathrm{HS}}_x.$$

Equivalence back to the classical text. Under endogenous evaluation,

$$\langle\!\langle \mathbf{g}^{\mathrm{Lor}}_{\mu\nu} \rangle\!\rangle = \bar{g}^{\mathrm{Lor}}_{\mu\nu}, \quad \langle\!\langle \mathcal{N}^{\mathrm{HS}}_x \rangle\!\rangle = \{V : \bar{g}^{\mathrm{Lor}}(V, V) = 0\}, \quad \langle\!\langle J^{\pm}_{\cup/\cap} \rangle\!\rangle = J^{\pm}_{\cup/\cap}(\bar{g}^{\mathrm{Lor}}_{(k)}),$$

so the effective light cone is a word–cone emanating from the PQS apex $\mathfrak{B}_x$; upon expectation it is exactly the Lorentzian cone of $\bar{g}^{\mathrm{Lor}}$ used in the main text.

5.4.3 The Dynamical Texture: The Temporal Phase-Flow Structure

The second ("soft") layer of causality in IQR does not expand the kinematic boundary set by the Effective Light Cone (Definition 5.122) but specifies a rich, anisotropic *intra-cone* dynamics for quantum information. Operationally, the cone assignment is fixed by the effective Lorentzian tensor (Definition 5.150; see also the projection guidance in Remark 5.47). The relevant object is the full temporal tensor $\bar{t}_{\mu\nu}(x)$, which we decompose into symmetric and antisymmetric parts:

$$\bar{t}_{\mu\nu}(x) = \bar{t}^{\mathrm{sym}}_{\mu\nu}(x) + \bar{t}^{\mathrm{anti}}_{\mu\nu}(x), \qquad \bar{t}^{\mathrm{sym}}_{\mu\nu} := \tfrac{1}{2}\big(\bar{t}_{\mu\nu} + \bar{t}_{\nu\mu}\big), \quad \bar{t}^{\mathrm{anti}}_{\mu\nu} := \tfrac{1}{2}\big(\bar{t}_{\mu\nu} - \bar{t}_{\nu\mu}\big). \tag{5.102}$$

This split supports the two-layered causal architecture of Section 5.4.4: the "hard" layer answers *whether* influence is allowed; the "soft" layer answers *how* coherence evolves along allowed paths.

Definition 5.136 (Temporal Phase-Flow Structure)**.** For $x \in \mathcal{X}_q$, the *Temporal Phase-Flow Structure* induced by $\bar{t}_{\mu\nu}(x)$ consists of two complementary components:

1. **Temporal Curvature Quadric (T-even).** The quadratic form
$$Q_t(V) := \bar{t}^{\mathrm{sym}}_{\mu\nu}(x)\, V^\mu V^\nu,$$
whose level sets $Q_t(V) = \text{const}$ define surfaces of constant phase-advance rate. The principal axes of Q_t select directions of extremal phase-flow curvature in $T_x\mathcal{M}_{\mathrm{geom}}$. In holosymmetric tomography, this component is isolated by the anticommutator response (T-even sector), cf. Proposition 5.60.
2. **Temporal Torsion Bivector (T-odd).** The 2-form $\bar{\omega}_{\mu\nu}(x) := \bar{t}^{\mathrm{anti}}_{\mu\nu}(x)$, which does not contribute to the quadratic form but generates a path-dependent phase holonomy (a "twist") in transported quantum states. In holosymmetric tomography, this component is isolated by the *retarded* commutator response (T-odd sector), cf. Proposition 5.60.

Interpretation. Classical, point-like trajectories depend only on the geodesic structure of $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ (hard layer; Definition 5.150; see also [7, 146]), and therefore do not register the temporal texture. Quantum processes that are phase/coherence sensitive do: $\bar{t}^{\mathrm{sym}}_{\mu\nu}$ governs *anisotropic phase-advance and shear*, leading to direction-dependent decoherence and interference, while $\bar{t}^{\mathrm{anti}}_{\mu\nu}$ governs *phase holonomy* (temporal torsion) akin to geometric-phase transport [23, 24, 25, 424, 408, 287, 247].

Holosymmetric readout. Within the fixed-cone assignment (unchanged by control phases; Corollary 5.126), the phase knobs (φ, θ) selectively probe the two components: anticommutator scans access the quadric (T-even) deformation of expectation geometry; commutator scans access the torsion-like (T-odd) phase curvature (Proposition 5.60). In symmetric limits (e.g., alignment conditions or the FLRW specialisation), these effects are suppressed or decouple, and the predictions reduce to standard GR/QFT behaviour (Section 5.4.7; [244, 11]).

Consistency note. Although $\bar{t}^{\mathrm{sym}}_{\mu\nu}$ enters the construction of $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ (cf. Definition 5.122), in this subsection we follow the operational guide of Remark 5.47: the cone is taken as fixed by the chosen projection route, and the temporal texture is analysed as an intra-cone structure that modulates *how*, not *whether*, causal evolution proceeds.

5.4.4 A Complementary Two-Layered Causal Architecture

The emergent causal structure in the IQR framework exhibits a complementary, two-layered hierarchy. Each layer plays a distinct yet interdependent role in defining the nature and evolution of causal propagation on the emergent chronogeometric manifold (cf. Definitions 5.122, 5.136 and 5.150).

1. **The Kinematic Boundary ("Hard" structure).** The *Effective Light Cone*, introduced in Definition 5.122, emanates from the Primordial Quantum Singularity $\mathfrak{B}_x$ (Theorem 5.155) and is determined by the effective Lorentzian metric $\bar{g}^{\mathrm{Lor}}_{\mu\nu}(x) = \bar{g}_{\mu\nu}(x) + \bar{t}^{\mathrm{sym}}_{\mu\nu}(x)$. It partitions $T_x\mathcal{M}_{\mathrm{geom}}$ into timelike, spacelike, and null directions and provides the *absolute boundary* for causal influence, answering the binary question: *Can* influence propagate from one point to another? In the classical limit this reduces to the familiar light-cone structure of GR [7, 146].

2. **The Dynamical Texture ("Soft" structure).** The *Temporal Phase-Flow Structure*, defined in Definition 5.136, is governed by the full temporal tensor $\bar{t}_{\mu\nu}(x)$, whose symmetric part shapes the phase-flow quadric and whose antisymmetric part contributes a torsion-like, phase-curvature response. This layer is *relational* and *anisotropic*: it characterises *how* coherence, phase, and entanglement evolve *within* the allowed cone, answering the higher-order question: *How does quantum coherence evolve along an allowed path?* Operationally, phase control separates T-even (anticommutator) and T-odd (commutator) responses (Proposition 5.60), echoing geometric-phase and interferometric diagnostics [23, 24, 424, 408, 287, 247].

Analogy: The Canyon and the River. The two-layered architecture can be visualised by a river flowing through a canyon. The **Effective Light Cone** provides the *impassable canyon walls*—the hard kinematic limits set by $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ (Definition 5.122). The **Temporal Phase-Flow Structure** supplies the *river current inside the canyon*—eddies, shears, and directional flows encoded by $\bar{t}_{\mu\nu}$ (Definition 5.136). A classical body traverses the canyon according to geodesics of $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ [7, 146], whereas a coherent quantum state responds to the internal texture: its phase and coherence evolve with both symmetric and antisymmetric contributions of $\bar{t}_{\mu\nu}$, leading to direction-dependent decoherence and interference even on isotropic $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ backgrounds [287, 247].

Holosymmetric extension. In the holosymmetric setting, the *hard* layer generalises to the *family of branchwise cones* $\{\bar{g}^{\mathrm{Lor}}_{(k)}\}_k$ (Definition 5.123). Microcausality is evaluated on the *intersection* of spacelike regions across branches, while potential influence is bounded by the *union* of branchwise futures (Proposition 5.124 and Corollary 5.126). The *soft* layer remains intra-cone: phases (φ, θ) adjust interference and reveal T-even/T-odd structure (Proposition 5.60) without altering cone assignment. In symmetric limits (e.g., alignment conditions leading to a single cone), the holosymmetric extension reduces to the single-history picture (Corollary 5.127), ensuring consistency with standard GR/QFT where appropriate [244, 11].

Remark 5.137 (Interpretation and relation to the two-layered architecture)**.** (i) *Cause is relational and operational.* In IQR, "cause'' is not an absolute primitive but a constrained *capacity to influence*, certified by (a) the **hard** kinematic layer—membership in $J^+_{\cup}$ and failure of Spacelike$_{\cap}$—and (b) the **soft** dynamical layer—the existence of a nonzero *retarded* response functional compatible with spectral positivity (Sections 5.4.4 and 5.4.7). This matches the operator-algebraic notion of locality plus positivity [230, 255] and reduces to standard GR/QFT causality in the single-cone limit [7, 11].

(ii) *Effect is a change in statistics at B traceable to A.* An "effect'' is the *observed* change in the expectation (or higher cumulants) of $\hat{\mathfrak{O}}_B$ under admissible local controls at A, with the direction of time fixed by the positivity/KMS/Hadamard conditions in Section 5.4.7. Holosymmetric phase control (φ, θ) can enhance *detection* (via interference visibility) without altering the underlying cone assignment (Corollary 5.126).

(iii) *Role of the temporal phase-flow.* The temporal tensor $\bar{t}_{\mu\nu}$ does not change *whether* influence is kinematically allowed (that is set by $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ and $J^{\pm}_{\cup}$), but it modulates *how* coherence and phase transport occur along allowed paths—biasing *which* allowed trajectories dominate (paths of maximal coherence) [287, 247]. This realises the two-layered causal architecture of Section 5.4.4.

(iv) *Branchwise and robust causation.* Strong (robust) causation obtains when $B \subset \bigcap_k J^+_{(k)}(A)$ and $R^{(k)}_{B\leftarrow A} \neq 0$ for all k. Otherwise, causation may be branch-selective but still operationally observable through holosymmetric interferometry (Definition 5.123 and Proposition 5.124).

(v) *PQS apex and origin.* All cones (and hence all admissible causal chains) emanate from the PQS apex $\mathfrak{B}_x$ (Theorem 5.155 and Remark 5.156). The notion of cause/effect is therefore ontologically *grounded* at PQSs and geometrically *shaped* by the effective light-cone structure (Definition 5.122).

Holosymmetric "cause and effect". Holosymmetry does *not* change what is kinematically admissible, but it refines how causation is *witnessed* and *decomposed* operationally. The admissibility layer is fixed by the family of branchwise cones $\{\bar{g}^{\mathrm{Lor}}_{(k)}\}_k$: microcausality is evaluated on the *intersection* of spacelike regions, while potential influence is bounded by the *union* of branchwise futures (Definition 5.123, Proposition 5.124, and Corollary 5.126). What holosymmetry adds is a phase-tunable interferometric handle that separates T-even and T-odd parts of the causal response without altering cone assignment (Proposition 5.60; cf. [230, 255]).

Definition 5.138 (Cause and effect in IQR (branchwise, retarded))**.** Let $A, B \subset \mathcal{M}_{\mathrm{geom}}$ with local observables $\hat{\mathfrak{O}}_A, \hat{\mathfrak{O}}_B$. Say that A can cause B iff the following hold:

(C1) **Kinematic admissibility:** A, B are not Spacelike$_{\cap}$ and $B \subset J^+_{\cup}(A)$, where Spacelike$_{\cap}$ and $J^+_{\cup}$ are defined by the branchwise cones of Definition 5.123.

(C2) **Retarded responsiveness:** There exists an admissible, spectrally positive local control on A such that the *retarded* linear response of $\hat{\mathfrak{O}}_B$ is nonzero, with advanced parts excluded by the spectral/Hadamard–KMS conditions (Section 5.4.7; [230, 244, 11]).

If A, B are Spacelike$_\cap$, microcausality enforces $[\hat{\mathfrak{O}}_A, \hat{\mathfrak{O}}_B] = 0$ and causation is ruled out (Proposition 5.124).

Definition 5.139 (Holosymmetric cause/effect: phase-resolved response)**.** Let $A, B \subset \mathcal{M}_{\text{geom}}$ and let $\hat{\mathfrak{O}}_B$ be a local observable supported in B. Choose an admissible local generator $\hat{\mathcal{G}}_A = \int J_A^{\mu\nu}(x)\, \hat{\Sigma}_{\mu\nu}(x)\, d^d x$ supported in A ($\hat{\Sigma}_{\mu\nu}$ as in Definition 5.25). Consider the holosymmetric control with phases (φ, θ) acting on the two law branches (cf. Definition 5.56 and Proposition 5.57). The *phase-resolved causal response* of B to A is the linear-response functional

$$\mathcal{E}_{B\leftarrow A}(\varphi,\theta) := \frac{d}{d\epsilon}\Big|_{\epsilon=0} \Big\langle \hat{\mathfrak{O}}_B \Big\rangle_{U_{\text{ctrl}}(\varphi,\theta)\, e^{-i\epsilon\hat{\mathcal{G}}_A}},$$

which can be written canonically as

$$\boxed{\mathcal{E}_{B\leftarrow A}(\varphi,\theta) = \tfrac{1}{2}\cos\varphi\,\cos\theta\, \underbrace{\Big\langle \{\hat{\mathcal{G}}_A, \hat{\mathfrak{O}}_B\} \Big\rangle}_{\text{T-even kernel}} + \tfrac{1}{2}\sin\varphi\,\sin\theta\, \underbrace{i\Big\langle [\hat{\mathcal{G}}_A, \hat{\mathfrak{O}}_B] \Big\rangle_{\text{ret}}}_{\text{T-odd retarded kernel}}}\,. \tag{5.103}$$

Here the (anti)commutators are evaluated in the Quantum Holo–State $|\mathring{\boldsymbol{\Psi}}\rangle$, with the commutator understood in the retarded sense so that advanced pieces vanish under the spectral-positivity/Hadamard–KMS conditions (Section 5.4.7; [230, 244, 11]).

Definition 5.140 (T-parity of response kernels)**.** Let T denote microscopic time-reversal (antiunitary, $\mathsf{T}\, i\, \mathsf{T}^{-1} = -i$), acting as $\mathsf{T}\, \hat{X}(t)\, \mathsf{T}^{-1} = \hat{X}_{\mathsf{T}}(-t)$. For local $\hat{\mathcal{G}}_A, \hat{\mathfrak{O}}_B$ define the two kernels

$$K_{\text{even}}(t) := \Big\langle \{\hat{\mathcal{G}}_A(t), \hat{\mathfrak{O}}_B(0)\} \Big\rangle, \quad K^{\text{ret}}_{\text{odd}}(t) := i\Big\langle [\hat{\mathcal{G}}_A(t), \hat{\mathfrak{O}}_B(0)] \Big\rangle_{\text{ret}}.$$

Then $\mathsf{T} K_{\text{even}}(t) \mathsf{T}^{-1} = +K_{\text{even}}(-t)$ *(T-even)* and $\mathsf{T} K^{\text{ret}}_{\text{odd}}(t) \mathsf{T}^{-1} = -K^{\text{ret}}_{\text{odd}}(-t)$ *(T-odd)*. With this convention, eq. (5.103) writes

$$\mathcal{E}_{B\leftarrow A}(\varphi,\theta) = \tfrac{1}{2}\cos\varphi\cos\theta\; K_{\text{even}} + \tfrac{1}{2}\sin\varphi\sin\theta\; K^{\text{ret}}_{\text{odd}},$$

so $\varphi = 0$ selects the T-even channel and $\varphi = \frac{\pi}{2}$ selects the T-odd (retarded) channel (cf. Proposition 5.60).

Definition 5.141 (T-even / T-odd kernels)**.** Let T be the time-reversal transformation acting on Heisenberg operators as $\mathsf{T}\, \hat{X}(t)\, \mathsf{T}^{-1} = \hat{X}_{\mathsf{T}}(-t)$ (antiunitary). Then for local generators/observables $\hat{\mathcal{G}}_A, \hat{\mathfrak{O}}_B$:

$$\mathsf{T}\, \{\hat{\mathcal{G}}_A(t), \hat{\mathfrak{O}}_B(0)\}\, \mathsf{T}^{-1} = +\{\hat{\mathcal{G}}_{A,\mathsf{T}}(-t), \hat{\mathfrak{O}}_{B,\mathsf{T}}(0)\} \quad \Rightarrow \quad \text{T-even},$$

$$\mathsf{T}\, \Big(i[\hat{\mathcal{G}}_A(t), \hat{\mathfrak{O}}_B(0)]\Big)\, \mathsf{T}^{-1} = -\, i[\hat{\mathcal{G}}_{A,\mathsf{T}}(-t), \hat{\mathfrak{O}}_{B,\mathsf{T}}(0)] \quad \Rightarrow \quad \text{T-odd}.$$

With the retarded prescription (step function support in the future), the commutator kernel is the *T-odd retarded* part; the anticommutator kernel is the *T-even* part (cf. Section 5.4.7).

Remark 5.142 (Two coherent uses of T-parity: operator–level vs. kernel–level)**.** There are two compatible ways to speak about "T-even/T-odd" in this chapter:

(A) Operator–level T-parity. One declares the *algebraic objects* $\{\hat{\mathcal{G}}_A(t), \hat{\mathfrak{O}}_B(0)\}$ and $i[\hat{\mathcal{G}}_A(t), \hat{\mathfrak{O}}_B(0)]$ to be respectively T–even / T–odd by their transformation under the antiunitary time–reversal $\mathsf{T}\,\hat{X}(t)\,\mathsf{T}^{-1} = \hat{X}_{\mathsf{T}}(-t)$:

$$\mathsf{T}\{\hat{\mathcal{G}}_A(t), \hat{\mathfrak{O}}_B(0)\}\mathsf{T}^{-1} = +\{\hat{\mathcal{G}}_{A,\mathsf{T}}(-t), \hat{\mathfrak{O}}_{B,\mathsf{T}}(0)\},$$
$$\mathsf{T}\Big(i[\hat{\mathcal{G}}_A(t), \hat{\mathfrak{O}}_B(0)]\Big)\mathsf{T}^{-1} = -\,i[\hat{\mathcal{G}}_{A,\mathsf{T}}(-t), \hat{\mathfrak{O}}_{B,\mathsf{T}}(0)].$$

This is a *state–independent* classification at the operator level.

(B) Kernel–level T-parity (our auxiliary convention). One first forms the *c-number kernels* used operationally in linear response,

$$K_{\mathrm{even}}(t) := \Big\langle \{\hat{\mathcal{G}}_A(t), \hat{\mathfrak{O}}_B(0)\} \Big\rangle, \qquad K^{\mathrm{ret}}_{\mathrm{odd}}(t) := i\Big\langle [\hat{\mathcal{G}}_A(t), \hat{\mathfrak{O}}_B(0)] \Big\rangle_{\mathrm{ret}},$$

(cf. Definition 5.139 and eq. (5.103)), and then record their T–parity: $K_{\mathrm{even}}(t) \mapsto +K_{\mathrm{even}}(-t)$, $K^{\mathrm{ret}}_{\mathrm{odd}}(t) \mapsto -K^{\mathrm{ret}}_{\mathrm{odd}}(-t)$. Here the *retarded* choice is built in (cf. Section 5.4.7).

Equivalence and use. Both viewpoints encode the same parity implied by $\mathsf{T}\,i\,\mathsf{T}^{-1} = -i$: the anticommutator is T–even and the (retarded) commutator is T–odd. (A) is algebraically minimal and can be cited anywhere; (B) aligns directly with the phase–resolved response

$$\mathcal{E}_{B\leftarrow A}(\varphi,\theta) = \tfrac{1}{2}\cos\varphi\cos\theta\, K_{\mathrm{even}} + \tfrac{1}{2}\sin\varphi\sin\theta\, K^{\mathrm{ret}}_{\mathrm{odd}},$$

so that $\varphi = 0$ isolates the T–even channel and $\varphi = \frac{\pi}{2}$ isolates the T–odd one (cf. Proposition 5.60). Note that T flips the step function, sending retarded to advanced, but our *operational* convention fixes the retarded projector by spectral positivity/Hadamard–KMS (cf. Section 5.4.7); the T–odd label refers to the underlying commutator, not to changing the causal prescription.

State symmetry. Whether or not the Quantum Holo–State is T–invariant does not affect the labels: T–parity refers to the transformation of the algebraic kernels; numerical invariance under T is a separate property of the state.

Recommendation. Use (A) when introducing symmetry properties at the operator level, and use (B) wherever eq. (5.103) is in play. The cone assignment remains phase–independent in either case (cf. Corollary 5.126).

Proposition 5.143 (Admissibility invariant, detection tunable)**.** *If A and B are* Spacelike$_\cap$ *then $[\hat{\mathfrak{O}}_A, \hat{\mathfrak{O}}_B] = 0$ and $\mathcal{E}_{B\leftarrow A}(\varphi,\theta) = 0$ for all (φ,θ) (Proposition 5.124). If $B \subset J^+_\cup(A)$ then $\mathcal{E}_{B\leftarrow A}$ may be* nonzero*, and phase scans isolate T-even vs. T-odd contributions by choosing $\varphi \in \{0, \frac{\pi}{2}\}$ (Proposition 5.60; [23, 24, 424, 408]).*

Proof. **(i) Spacelike case.** If A and B are Spacelike$_\cap$, then branchwise microcausality holds by Proposition 5.124: for any local fields supported in A and B one has vanishing commutators in each branch, hence

$$[\hat{\mathfrak{O}}_A, \hat{\mathfrak{O}}_B] = 0.$$

Since the admissible local generator $\hat{\mathcal{G}}_A = \int J_A^{\mu\nu} \hat{\Sigma}_{\mu\nu}$ is built from local operators supported in A (cf. Definition 5.25), the same locality statement implies $[\hat{\mathcal{G}}_A, \hat{\mathfrak{O}}_B] = 0$ whenever $A \perp_\cap B$ (see also Definition 5.123 and Corollary 5.126).

Insert these facts into the phase-resolved response of Definition 5.139, eq. (5.103):

$$\mathcal{E}_{B\leftarrow A}(\varphi,\theta) = \tfrac{1}{2}\cos\varphi\,\cos\theta\,\Big\langle\{\hat{\mathcal{G}}_A,\hat{\mathfrak{O}}_B\}\Big\rangle + \tfrac{1}{2}\sin\varphi\,\sin\theta\,i\Big\langle[\hat{\mathcal{G}}_A,\hat{\mathfrak{O}}_B]\Big\rangle_{\mathrm{ret}}.$$

The retarded (T-odd) kernel vanishes because $[\hat{\mathcal{G}}_A, \hat{\mathfrak{O}}_B] = 0$ and the retarded prescription excludes advanced parts (cf. Section 5.4.7). The T-even kernel is a symmetrised *connected* correlator; by the same spacelike locality plus spectral positivity/Hadamard–KMS conditions its support is confined within the union of branchwise causal domains, so it also vanishes when $A \perp_\cap B$ (cf. Section 5.4.7 and Proposition 5.60). Therefore

$$\mathcal{E}_{B\leftarrow A}(\varphi,\theta) = 0 \quad \text{for all } (\varphi,\theta),$$

as claimed.

(ii) Causally admissible case. If $B \subset J_\cup^+(A)$ (with $J_\cup^+$ defined by the holosymmetric cones, Definition 5.123), locality no longer forces the kernels to vanish. For admissible controls on A, the retarded (T-odd) kernel $i\langle[\hat{\mathcal{G}}_A,\hat{\mathfrak{O}}_B]\rangle_{\mathrm{ret}}$ can be nonzero while advanced parts remain excluded by spectral positivity (cf. Section 5.4.7). By Proposition 5.60, phase scans separate the two contributions in (5.103): choosing $\varphi = 0$ (standard balanced setting for θ) isolates the T-even anticommutator channel, whereas $\varphi = \frac{\pi}{2}$ isolates the T-odd retarded commutator channel, without altering the cone assignment (Corollary 5.126). Hence $\mathcal{E}_{B\leftarrow A}$ may be nonzero and its T-even/T-odd parts are operationally resolvable by phase control.

Combining (i)–(ii) yields the proposition. □

Definition 5.144 (Holosymmetric causation strengths)**.** Define the *branchwise* retarded strengths $R^{(k)}_{B\leftarrow A} := \frac{d}{d\epsilon}\big|_0 \langle\hat{\mathfrak{O}}_B\rangle^{(k)}_{e^{-i\epsilon\hat{\mathcal{G}}_A}}$. The *robust holo-causal strength* and the *selective holo-causal strength* are

$$\mathcal{E}^{\mathrm{rob}}_{B\leftarrow A} := \min_k \|R^{(k)}_{B\leftarrow A}\|, \qquad \mathcal{E}^{\mathrm{sel}}_{B\leftarrow A} := \max_k \|R^{(k)}_{B\leftarrow A}\|.$$

Robust causation obtains when $\mathcal{E}^{\mathrm{rob}} > 0$ and $B \subset \bigcap_k J^+_{(k)}(A)$; otherwise causation is branch-selective but can be amplified or nulled interferometrically via (5.103) (cf. [426, 427, 287]).

Corollary 5.145 (No–law–signalling)**.** *Holosymmetric control cannot create causation where kinematics forbids it: if $B \not\subset J_\cup^+(A)$ then $\mathcal{E}_{B\leftarrow A}(\varphi,\theta) = 0$ for all (φ,θ).*

Proof. Let $A, B \subset \mathcal{M}_{\mathrm{geom}}$ and suppose $B \not\subset J_\cup^+(A)$, with $J_\cup^+$ defined by the holosymmetric cones (cf. Definition 5.123 and Corollary 5.126). By definition, for every branch k either B is spacelike to A or B lies outside $J^+_{(k)}(A)$ (in particular, in the past of A) with respect to $\bar{g}^{\mathrm{Lor}}_{(k)}$.

Write the phase–resolved linear response (cf. Definition 5.139):

$$\mathcal{E}_{B\leftarrow A}(\varphi,\theta) = \tfrac{1}{2}\cos\varphi\cos\theta\,\Big\langle\{\hat{\mathcal{G}}_A,\hat{\mathfrak{O}}_B\}\Big\rangle + \tfrac{1}{2}\sin\varphi\sin\theta\,i\Big\langle[\hat{\mathcal{G}}_A,\hat{\mathfrak{O}}_B]\Big\rangle_{\mathrm{ret}}, \tag{5.104}$$

where $\hat{\mathcal{G}}_A = \int J_A^{\mu\nu}\hat{\Sigma}_{\mu\nu}$ is a local generator supported in A (cf. Definition 5.25).

(a) Retarded (T-odd) kernel. For each branch k, the support of the retarded commutator $i\langle[\hat{\mathcal{G}}_A, \hat{\mathfrak{O}}_B]\rangle_{\mathrm{ret}}$ is contained in $J^+_{(k)}(A)$ by construction of the retarded prescription and spectral positivity/Hadamard–KMS conditions (cf. Section 5.4.7). Since $B \not\subset J^+_{\cup}(A)$, it follows that $i\langle[\hat{\mathcal{G}}_A, \hat{\mathfrak{O}}_B]\rangle_{\mathrm{ret}} = 0$.

(b) Symmetric (T-even) kernel. If A and B are spacelike in a branch, local commutativity yields $[\hat{\mathcal{G}}_A, \hat{\mathfrak{O}}_B] = 0$ and the connected T-even contribution between A and B vanishes operationally (cf. Proposition 5.124 and Section 5.4.7). If B is branchwise *not* in the future of A, the time-symmetric kernel does not contribute to a *retarded* response: by the same positivity/causality conditions, its operational support is confined within the causal domains determined by the cones (cf. Proposition 5.60 and Section 5.4.7), hence it vanishes whenever $B \not\subset J^+_{(k)}(A)$ for all k.

Combining (a)–(b) in (5.104) gives $\mathcal{E}_{B\leftarrow A}(\varphi, \theta) = 0$ for all (φ, θ). Phase choices cannot alter this conclusion because the cone assignments—and thus the supports of both kernels—are phase-independent (cf. Corollary 5.126). This proves Corollary 5.145. □

Proposition 5.146 (Dephasing monotonicity (operational arrow))**.** *Let $\mathcal{D}_p$ be any completely-positive phase-damping channel on the control or registers with strength $p \in [0, 1]$. Then $|\mathcal{E}_{B\leftarrow A}(\varphi, \theta)|$ is non-increasing in p, with equality only for purely T-even kernels at the chosen phase. Thus environmental decoherence suppresses detection strength without altering admissibility ([247]; cf. Corollary 5.126).*

Remark 5.147 (Intuition and relation to the two-layered architecture)**.** Holosymmetry leaves the **hard** layer unchanged—cause requires membership in $J^+_{\cup}$ and failure of Spacelike$_{\cap}$ (Section 5.4.4). It enriches the **soft** layer by making the *detection* of effects phase-tunable through (5.103), separating symmetric (T-even) and antisymmetric (T-odd) transport governed by the temporal tensor $\bar{t}_{\mu\nu}$ (Definition 5.136). In the single-history/classical limit the two notions collapse to standard GR/QFT causality [7, 11].

Proposition 5.148 (Holosymmetry does *not* superpose the cause–effect predicate)**.** *Holosymmetry places* chronogeometric histories (laws) *in coherent superposition and lets their* causal responses *interfere; it does not place the logical predicate "A causes B" itself in superposition. Causation remains* branchwise and retarded *and is read out operationally (cf. Definitions 5.123, 5.138 and 5.139; see also [230, 255, 244, 11]).*

Proof. We show that holosymmetry coherently superposes *histories* and lets their *responses* interfere, while the *predicate* "A causes B" remains a branchwise/retarded, non-superposed statement determined by cone kinematics and linear response (cf. Definitions 5.123, 5.138 and 5.139; see also [230, 255, 244, 11]).

1. Kinematic gate is phase–invariant. Holosymmetry prepares a superposition of *law branches* with effective cones $\{\bar{g}^{\mathrm{Lor}}_{(k)}\}_k$ (Definitions 5.56 and 5.123 and Proposition 5.57). By Corollary 5.126, the operational relations Spacelike$_{\cap}$ and $J^{\pm}_{\cup}$ are *independent* of the control phases (φ, θ); thus condition (C1) in Definition 5.138—kinematic admissibility $B \subset J^+_{\cup}(A)$ & A, B not Spacelike$_{\cap}$ —is a *phase-invariant* Boolean statement. In particular, cone tilts that may arise from small holosymmetric deformations are controlled branchwise and do not depend on (φ, θ) (Proposition 5.125).

2. Branchwise microcausality and no–law–signalling. For Spacelike$_\cap$ regions A, B, local commutativity holds *in each branch* (Proposition 5.124). Consequently, with any local generator $\hat{\mathcal{G}}_A = \int J_A^{\mu\nu} \hat{\Sigma}_{\mu\nu}$ (Definition 5.25), one has $[\hat{\mathcal{G}}_A, \hat{\mathfrak{O}}_B] = 0$ and the retarded support from A is confined to $J_\cup^+(A)$ (Corollary 5.145). Thus, if (C1) fails, (C2) in Definition 5.138 cannot be satisfied for *any* phases.

3. Phase–resolved response lives inside the cones. By Definition 5.139, the phase-resolved linear response is

$$\mathcal{E}_{B\leftarrow A}(\varphi,\theta) = \tfrac{1}{2}\cos\varphi\,\cos\theta\,\big\langle\{\hat{\mathcal{G}}_A, \hat{\mathfrak{O}}_B\}\big\rangle + \tfrac{1}{2}\sin\varphi\,\sin\theta\, i\big\langle[\hat{\mathcal{G}}_A, \hat{\mathfrak{O}}_B]\big\rangle_{\text{ret}} \quad \text{(cf. } eq.\ (5.103)\text{)}. \tag{5.105}$$

The commutator term is retarded by the spectral–positivity/Hadamard–KMS prescription (Section 5.4.7), hence its support is contained in $J_\cup^+(A)$; it vanishes if $B \not\subset J_\cup^+(A)$ (Corollary 5.145). The anticommutator term contributes to detection *only* within the same causal domain and also vanishes at Spacelike$_\cap$ (Propositions 5.60 and 5.143).

4. Interference changes visibility, not admissibility. Equation (5.105) shows that (φ,θ) act as *real weights* that separate the T-even (anticommutator) and T-odd (retarded commutator) channels (Proposition 5.60), but cannot move their support beyond $J_\cup^+(A)$, nor alter Spacelike$_\cap$ (Corollary 5.126). Therefore:

$$\big(\text{C1 fails}\big) \;\Rightarrow\; \mathcal{E}_{B\leftarrow A}(\varphi,\theta) = 0 \text{ for all } (\varphi,\theta) \quad \text{(no–law–signalling: } \textit{Corollary } 5.145\text{)},$$

while if (C1) holds, then (C2) may hold and the *magnitude* of the observable effect can be enhanced/suppressed by phase choice without changing its kinematic admissibility (Proposition 5.143). The *truth value* of "A can cause B" in Definition 5.138 is thus determined by (C1) and the *existence* of a nonzero retarded response (C2), not by a superposed logical predicate.

5. Robust vs. selective causation is not superposition of the predicate. When only some branches allow a retarded response, one distinguishes "robust" and "selective" strengths (Definition 5.144). These quantify *how many* branches contribute and with what norm, but the predicate in Definition 5.138 is still evaluated according to (C1)+(C2) on the holosymmetric domains (Definition 5.123). Branchwise interference forms an *amplitude sum*, not a superposition of truth values.

6. Forward–backward probes do not superpose the predicate. In Loschmidt-type settings the superposed histories are U_0 and $U_0^\dagger$; the antiunitary time-reversal T is not a controllable branch (Remark 5.62; see also [426, 427, 350, 287]). Cone assignments remain those of the branch metrics and are phase-independent (Corollary 5.126); hence neither indefinite causal order nor a superposition of the predicate arises.

Conclusion. Holosymmetry coherently superposes *laws* and yields phase-tunable interference of *causal responses*, but (i) the kinematic gate (C1) is phase-invariant and branchwise (Definition 5.123 and Corollary 5.126), and (ii) the operational condition (C2) uses the *retarded* commutator confined to $J_\cup^+(A)$ (Section 5.4.7, Definition 5.139, and Corollary 5.145). Therefore the logical predicate "A causes B" is *not* placed in superposition; causation remains *branchwise and retarded*, as claimed. □

Remark 5.149 (Operational structure)**. (a) What holosymmetry superposes.** Law branches with cones $\{\bar{g}^{\mathrm{Lor}}_{(k)}\}$, not the truth value of "A causes B" (Definition 5.123); phases (φ, θ) do not change spacelike/timelike assignments (Proposition 5.124 and Corollary 5.126).

(b) Branchwise causation and aggregation. Causation is decided branchwise by kinematics + nonzero retarded response (cf. Definition 5.138); holosymmetry forms an *interferometric sum* of these responses and cannot violate no–law–signalling (Corollary 5.145).

(c) Three regimes. (1) *Robust:* $A \to B$ in every branch with nonzero retarded responses; interference changes strength, not admissibility (Definition 5.144). (2) *Selective:* admissible in some branches only; phase can hide or reveal an otherwise admissible effect, but admissibility still comes from $J^+_\cup$ (Definition 5.123 and Proposition 5.143). (3) *Null:* if A, B are Spacelike$_\cap$, commutators vanish and the causal response is identically zero for all phases (Proposition 5.124 and Definition 5.138).

(d) Not indefinite causal order. Time orientation is fixed branchwise by spectral positivity; advanced parts vanish (cf. Section 5.4.7; [230, 11]). Forward–backward probes superpose U_0 with $U_0^\dagger$, not "A before B" with "B before A" (Remark 5.62; [426, 427]).

(e) Phase-resolved diagnosis (tomography). The phase knob separates T-even and T-odd contributions without affecting cone assignment (Proposition 5.60 and Definition 5.139; [23, 24, 424, 408, 287]). Concretely,

$$\mathcal{E}_{B\leftarrow A}(\varphi,\theta) = \tfrac{1}{2}\cos\varphi\cos\theta\left\langle\{\hat{\mathcal{G}}_A, \hat{\mathfrak{O}}_B\}\right\rangle + \tfrac{1}{2}\sin\varphi\sin\theta\, i\left\langle[\hat{\mathcal{G}}_A, \hat{\mathfrak{O}}_B]\right\rangle_{\mathrm{ret}}, \tag{5.106}$$

so $\varphi = 0$ isolates the T-even (anticommutator) kernel and $\varphi = \frac{\pi}{2}$ isolates the T-odd retarded commutator.

Bottom line. Holosymmetry *superposes mechanisms*, not the truth value of "cause" and "effect": the kinematic gates (intersection/union cone rules) are unchanged, while interference offers a phase-tunable *diagnosis and decomposition* of how an already-admissible influence manifests—never a means to conjure influence where none is allowed (Definition 5.123, Proposition 5.124, and Corollary 5.145).

5.4.5 The Unified Causal Architecture

The complete causal structure in the IQR framework is a synthesis of two distinct but complementary layers. A physical process is constrained externally by the absolute boundary of the effective light cone and internally modulated by the quantum-dynamical texture of the temporal phase-flow structure. Together, these define a coherent, emergent causal geometry that is both *relationally grounded* and *dynamically textured*.

Definition 5.150 (Unified Causal Structure)**.** The emergent causal architecture on the manifold

$$\mathcal{M}_{\mathrm{geom}} := \left(\mathcal{X}_q,\ \bar{g}^{\mathrm{Lor}}_{\mu\nu},\ \bar{t}_{\mu\nu}\right)$$

is determined by the *causal-dual tensor pair* $(\bar{g}^{\mathrm{Lor}}_{\mu\nu}, \bar{t}_{\mu\nu})$, where:

1. The **Effective Lorentzian Metric** $\bar{g}^{\mathrm{Lor}}_{\mu\nu}(x) := \bar{g}_{\mu\nu}(x) + \bar{t}^{\mathrm{sym}}_{\mu\nu}(x)$, as defined in Definition 5.122, encodes the kinematic boundary of causality through the local light cone $\mathcal{V}_\pm(\mathfrak{B}_x)$.

2. The **Temporal Phase-Flow Tensor** $\bar{t}_{\mu\nu}(x)$, as defined in Definition 5.136, encodes the internal anisotropy and torsion of quantum-coherent evolution along permitted causal paths.

Given a tangent vector $V^\mu \in T_x\mathcal{M}_{\text{geom}}$, the causal relation is fully characterized by the *Causal Structure Functional*:

$$\boxed{\mathcal{C}_x(V) := \left(\chi_{\text{causal}}\left[\bar{g}^{\text{Lor}}_{\mu\nu}(x)V^\mu V^\nu\right],\ \bar{t}_{\mu\nu}(x)V^\mu V^\nu\right),} \tag{5.107}$$

where $\chi_{\text{causal}}(s)$ is the causal indicator function:

$$\chi_{\text{causal}}(s) := \begin{cases} 1, & \text{if } s \geq 0 \quad \text{(causally allowed)} \\ 0, & \text{if } s < 0 \quad \text{(causally forbidden).} \end{cases}$$

Interpretation of the Causal Structure Functional

The causal structure functional $\mathcal{C}_x(V)$ should be interpreted as a dual interrogation of the local chronogeometric fabric. It assesses both whether a direction is causally permissible and how a quantum-coherent process behaves along that direction.

1. The First Component: The Kinematic Gatekeeper The binary component

$$\chi_{\text{causal}}[\bar{g}^{\text{Lor}}_{\mu\nu}(x)V^\mu V^\nu]$$

acts as a definitive logical switch:

- *Function:* It determines whether V^μ lies inside or outside the effective light cone.
- *Physical Meaning:* It enforces the absolute causal boundary, as defined by the effective metric $\bar{g}^{\text{Lor}}_{\mu\nu}$. If the path is spacelike, this component returns 0, rendering the second term physically moot.

2. The Second Component: The Quantum-Dynamical Texture The scalar $\bar{t}_{\mu\nu}(x)V^\mu V^\nu$ is a continuous-valued functional that becomes physically meaningful only when the direction V^μ is causally allowed ($\chi_{\text{causal}} = 1$). It captures the dynamic evolution of quantum phase coherence:

- *Function:* Quantifies the curvature-induced modulation of phase evolution along V^μ, incorporating both anisotropic flow and torsional holonomy.
- *Physical Interpretation:*
  - A large positive value indicates rapid decoherence or phase rotation.
  - A near-zero value indicates a temporally flat or coherence-preserving direction.

– A negative value may correspond to phase re-stabilization or coherent reinforcement.

In total, the causal structure functional $\mathcal{C}_x(V)$ captures both the *permission* and the *character* of causal propagation. It generalizes the classical light cone criterion by embedding it within a richer, quantum-relational fabric—providing a unified causal architecture that encodes *both* where influence can travel and how coherence propagates along its permitted paths. This unified causal architecture has profound implications, suggesting that the universe is governed by a richer set of rules than classical relativity allows.

Beyond Geodesics: Paths of Maximal Coherence In classical general relativity, free motion follows geodesics of the background metric $\bar{g}_{\mu\nu}$ [7, 146]. In the IQR framework, the classical limit tracks geodesics of the *effective Lorentzian metric* $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ (cf. Definitions 5.122 and 5.150). Quantum evolution, however, is phase-sensitive and decoherence-prone; a quantum system therefore need not follow the same curve. Its most probable worldline is biased toward *paths of maximal coherence* determined by the temporal phase-flow structure $\bar{t}_{\mu\nu}$ (see Definition 5.136 and eq. (5.107)), so that the "straightest" (geodesic) and the "most coherent" (decoherence-minimizing) paths generically differ [428, 287, 247]. In stationary limits and weak phase-curvature, these two notions coincide to leading order, recovering the geodesic principle inside IQR.

Anisotropic Quantum Universe The temporal tensor $\bar{t}_{\mu\nu}$ induces a *quantum* anisotropy that leaves classical kinematics intact but modulates coherence and interference along different spatial directions (again via Definition 5.136). Thus two otherwise identical quantum experiments performed along inequivalent directions may display distinct decoherence rates or fringe visibilities even when $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ is isotropic (cf. Definition 5.122). This suggests concrete, testable signatures beyond standard physics, in the spirit of directional phase/optical anisotropies already constrained in cosmological and laboratory settings [429, 430]. In the symmetric (e.g. FLRW) limit where $\bar{t}_{\mu\nu}$ aligns with the foliation, the anisotropy is suppressed and the IQR predictions reduce to the standard isotropic case (see Section 5.4.7).

A New Dynamical Field The temporal phase-flow tensor $\bar{t}_{\mu\nu}$ is a *dynamical* field sourced by the universal quantum state (cf. Section 5.2.6), not a fixed background constant. Its evolution reshapes coherence transport inside the effective light cone and may act as a state-dependent driver of cosmic dynamics. In particular, a slowly varying, pervasive $\bar{t}_{\mu\nu}$ component can mimic—or dynamically source—late-time acceleration, offering an alternative to a rigid cosmological constant and connecting to broader discussions of modified or emergent gravitational effects [213, 147]. In appropriate limits, the construction remains compatible with QFT on curved spacetime [244, 11]: $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ sets kinematic causal boundaries, while $\bar{t}_{\mu\nu}$ governs the phase-coherent texture of quantum transport within those bounds (cf. Definition 5.150). *Synthesis.* The Unified Causal Structure (Definition 5.150) thus preserves the absolute causal boundaries of relativity via $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ while enriching the interior of the light cone with a dynamic, anisotropic, and experimentally accessible quantum texture encoded by $\bar{t}_{\mu\nu}$. In the classical or symmetry-aligned limits these structures collapse to the familiar GR picture; beyond

those limits they predict direction-dependent coherence and interference without violating microcausality or spectral stability.

Definition 5.151 (Holosymmetric Unified Causal Structure)**.** In the holosymmetric setting with two law branches $k \in \{0,1\}$, let

$$\bar{g}^{\mathrm{Lor}}_{(k)\mu\nu} := \bar{g}_{(k)\mu\nu} + \bar{t}^{\mathrm{sym}}_{(k)\mu\nu}, \qquad \bar{t}_{(k)\mu\nu} \text{ as in } \textit{Definition } 5.136,$$

and write $\mathcal{V}^{(k)}_{\pm}(\mathfrak{B}_x)$ for the branchwise light cones (cf. Definition 5.122). Define the holosymmetric *microcausal spacelike* relation and *influence futures* by

$$\mathrm{Spacelike}_{\cap}(x,y) \iff (x,y) \text{ are spacelike w.r.t. both } \bar{g}^{\mathrm{Lor}}_{(0)} \text{ and } \bar{g}^{\mathrm{Lor}}_{(1)},$$

$$J^{\pm}_{\cup}(x) := J^{\pm}_{\bar{g}^{\mathrm{Lor}}_{(0)}}(x) \;\cup\; J^{\pm}_{\bar{g}^{\mathrm{Lor}}_{(1)}}(x).$$

The *holosymmetric causal-dual data* at $x \in \mathcal{M}_{\mathrm{geom}}$ are

$$\Big(\{\bar{g}^{\mathrm{Lor}}_{(k)}\}_k,\ \{\bar{t}_{(k)}\}_k \Big),$$

and the *holosymmetric causal functionals* acting on $V \in T_x\mathcal{M}_{\mathrm{geom}}$ are

$$\mathcal{C}^{\mathrm{micro}}_x(V) := \min_k \ \chi_{\mathrm{causal}}\Big[\bar{g}^{\mathrm{Lor}}_{(k)\mu\nu}(x) V^\mu V^\nu\Big], \tag{5.108}$$

$$\mathcal{C}^{\mathrm{reach}}_x(V) := \max_k \ \chi_{\mathrm{causal}}\Big[\bar{g}^{\mathrm{Lor}}_{(k)\mu\nu}(x) V^\mu V^\nu\Big], \tag{5.109}$$

$$\mathcal{T}^{\mathrm{HS}}_x(V) := \Big(\bar{t}_{(0)\mu\nu} V^\mu V^\nu,\ \bar{t}_{(1)\mu\nu} V^\mu V^\nu \Big), \tag{5.110}$$

with χ_{causal} as in eq. (5.107). Here $\mathcal{C}^{\mathrm{micro}}$ governs *microcausality* (commutativity), while $\mathcal{C}^{\mathrm{reach}}$ governs *potential influence*.

Proposition 5.152 (No–law–signalling and phase invariance)**.** *If regions A, B satisfy* $\mathrm{Spacelike}_{\cap}$*, then for local observables $\hat{\mathfrak{O}}(f_A), \hat{\mathfrak{O}}(f_B)$ one has $[\hat{\mathfrak{O}}(f_A), \hat{\mathfrak{O}}(f_B)] = 0$ under any holosymmetric control (any phases (φ,θ)). Possible signal paths from A are confined to $J^+_{\cup}(A)$. The sets* $\mathrm{Spacelike}_{\cap}$ *and $J^{\pm}_{\cup}$, and hence the values of $\mathcal{C}^{\mathrm{micro}}_x, \mathcal{C}^{\mathrm{reach}}_x$, are independent of (φ,θ).*

Proof. **(1) Branchwise microcausality at** $\mathrm{Spacelike}_{\cap}$**.** By definition, $A \perp_{\cap} B$ means A, B are spacelike with respect to *each* branch metric $\bar{g}^{\mathrm{Lor}}_{(k)}$ (Definition 5.123). Hence branchwise locality gives $[\hat{\mathfrak{O}}(f_A), \hat{\mathfrak{O}}(f_B)] = 0$ in every branch. Holosymmetric control unitaries act as direct sums over branches (functorially on the law index; cf. Proposition 5.57 and Lemma 5.55), so if each summand commutator vanishes then, for the controlled evolution $U_{\mathrm{ctrl}}(\varphi,\theta)$,

$$\Big[U_{\mathrm{ctrl}}(\varphi,\theta)\, \hat{\mathfrak{O}}(f_A)\, U_{\mathrm{ctrl}}(\varphi,\theta)^\dagger,\ U_{\mathrm{ctrl}}(\varphi,\theta)\, \hat{\mathfrak{O}}(f_B)\, U_{\mathrm{ctrl}}(\varphi,\theta)^\dagger \Big] = 0,$$

which is the stated $[\hat{\mathfrak{O}}(f_A), \hat{\mathfrak{O}}(f_B)] = 0$ under any phases (φ,θ) (cf. Proposition 5.124).

(2) Retarded support confined to $J^+_{\cup}(A)$ (no–law–signalling). Let $\hat{\mathcal{G}}_A = \int J_A^{\mu\nu}\hat{\Sigma}_{\mu\nu}$ be any admissible local generator supported in A (Definition 5.25). The phase-resolved linear response at B is (Definition 5.139 and eq. (5.103))

$$\mathcal{E}_{B\leftarrow A}(\varphi,\theta) = \tfrac{1}{2}\cos\varphi\cos\theta\left\langle\{\hat{\mathcal{G}}_A,\hat{\mathfrak{O}}_B\}\right\rangle + \tfrac{1}{2}\sin\varphi\sin\theta\, i\left\langle[\hat{\mathcal{G}}_A,\hat{\mathfrak{O}}_B]\right\rangle_{\mathrm{ret}}.$$

By the spectral-positivity/Hadamard–KMS prescription, the retarded commutator has support *only* in the future domain determined by the cones (Section 5.4.7). Since holosymmetry provides the union of branchwise futures $J^+_{\cup}(A) = \bigcup_k J^+_{(k)}(A)$ (Definition 5.123), one has supp $i\langle[\hat{\mathcal{G}}_A,\hat{\mathfrak{O}}_B]\rangle_{\mathrm{ret}} \subseteq J^+_{\cup}(A)$, and the same causal restriction governs the anticommutator contribution operationally (Propositions 5.60 and 5.143). Hence possible signal paths from A are confined to $J^+_{\cup}(A)$ (see also Corollary 5.145).

(3) Phase invariance of the kinematic relations and the causal indices. By Corollary 5.126, the sets Spacelike$_{\cap}$ and $J^{\pm}_{\cup}$ are *independent* of the control phases (φ,θ). Consequently any cone-defined index or capacity—such as the microcausality indicator $\mathcal{C}_x^{\mathrm{micro}}$ (vanishing iff all local commutators at x vanish on Spacelike$_{\cap}$) and the reach index $\mathcal{C}_x^{\mathrm{reach}}$ (nonzero only if $x \in J^+_{\cup}(A)$ by construction in the unified causal structure; cf. Definitions 5.123 and 5.150)—is likewise phase-invariant. Moreover, while small holosymmetric deformations can *tilt* branch cones, such tilts are controlled by $\|\Delta\bar{g}^{\mathrm{Lor}}\|_x$ and not by (φ,θ) (Proposition 5.125), leaving the phase-invariance statement intact.

Combining (1)–(3) yields the proposition: (i) microcausality on Spacelike$_{\cap}$ holds under any holosymmetric control; (ii) all causal influence from A is confined to $J^+_{\cup}(A)$; and (iii) the relations Spacelike$_{\cap}$, $J^{\pm}_{\cup}$ and thus $\mathcal{C}_x^{\mathrm{micro}}, \mathcal{C}_x^{\mathrm{reach}}$ are independent of (φ,θ). □

Proposition 5.153 (Stability and reduction)**.** *Let $\Delta\bar{g}^{\mathrm{Lor}} := \bar{g}^{\mathrm{Lor}}_{(1)} - \bar{g}^{\mathrm{Lor}}_{(0)}$. Null directions tilt by an amount bounded by $\|\Delta\bar{g}^{\mathrm{Lor}}\|_x$, so both* Spacelike$_{\cap}$ *and $J^{\pm}_{\cup}$ vary continuously under small holosymmetric deformations. If the branches coincide (e.g. aligned temporal sectors), then $\bar{g}^{\mathrm{Lor}}_{(0)} = \bar{g}^{\mathrm{Lor}}_{(1)}$ and*

$$\mathcal{C}_x^{\mathrm{micro}} = \mathcal{C}_x^{\mathrm{reach}} = \chi_{\mathrm{causal}}\Big[\bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)V^\mu V^\nu\Big],$$

reducing exactly to Definition 5.150.

Proof. **(1) Cone tilt and continuity.** Let $\Delta\bar{g}^{\mathrm{Lor}} := \bar{g}^{\mathrm{Lor}}_{(1)} - \bar{g}^{\mathrm{Lor}}_{(0)}$. By Proposition 5.125, for any $\bar{g}^{\mathrm{Lor}}_{(0)}$–null V at x one has

$$\left|\bar{g}^{\mathrm{Lor}}_{(1)}(V,V)\right| = \left|\Delta\bar{g}^{\mathrm{Lor}}(V,V)\right| \le \|\Delta\bar{g}^{\mathrm{Lor}}\|_x\,\|V\|_x^2,$$

so the null directions tilt by an amount bounded by $\|\Delta\bar{g}^{\mathrm{Lor}}\|_x$. Equivalently, the future/past cones $C^{\pm}_{(k)}(x)$ of each branch vary *Lipschitz-continuously* in the operator norm of the metric perturbation. Therefore the branchwise spacelike/timelike classification of any *fixed* tangent vector (and hence of pairs of points with a *strict* spacelike/timelike margin) is stable for sufficiently small $\|\Delta\bar{g}^{\mathrm{Lor}}\|_x$; only vectors lying on the null boundary can change status, and this change is continuous in the sense of boundary deformation controlled by $\|\Delta\bar{g}^{\mathrm{Lor}}\|_x$.

(2) Continuity of Spacelike$_{\cap}$ **and $J^{\pm}_{\cup}$.** By Definition 5.123,

$$\mathrm{Spacelike}_{\cap}(x,y) \iff (x,y) \text{ spacelike for each } \bar{g}^{\mathrm{Lor}}_{(k)}, \qquad J^{\pm}_{\cup}(A) = \bigcup_k J^{\pm}_{(k)}(A).$$

Since each branch cone varies continuously by (1), their *finite* intersection (for $\text{Spacelike}_\cap$) and finite union (for $J^\pm_\cup$) inherit continuity under small holosymmetric deformations. Concretely, if (x, y) is strictly spacelike in every branch, then for $\|\Delta\bar{g}^{\text{Lor}}\|$ small enough it remains so; and if $y \in J^+_\cup(x)$ via a branch k_0, then for small deformations there remains a (nearby) causal path in branch k_0 from x to y, whence $y \in J^+_\cup(x)$ persists. Thus both relations vary continuously with the holosymmetric deformation parameter, as stated.

(3) Reduction when the branches coincide. If the branches coincide (e.g. aligned temporal sectors), then by Corollary 5.127 one has $\bar{g}^{\text{Lor}}_{(0)} = \bar{g}^{\text{Lor}}_{(1)} =: \bar{g}^{\text{Lor}}$, hence

$$J^\pm_\cup = J^\pm_\cap = J^\pm_{\bar{g}^{\text{Lor}}}, \qquad \text{Spacelike}_\cap = \text{Spacelike}_{\bar{g}^{\text{Lor}}}.$$

By the unified causal structure Definition 5.150, the microcausality indicator and reach index at x collapse to the single–cone evaluation of the causal sign of the quadratic form:

$$\mathcal{C}^{\text{micro}}_x = \mathcal{C}^{\text{reach}}_x = \chi_{\text{causal}}\Big[\bar{g}^{\text{Lor}}_{\mu\nu}(x)V^\mu V^\nu\Big],$$

i.e. they agree and reduce exactly to the single–history definition governed by $\bar{g}^{\text{Lor}}$ (cf. also Definition 5.122).

Combining (1)–(3) proves the proposition: the holosymmetric causal relations depend continuously on small branch deformations controlled by $\|\Delta\bar{g}^{\text{Lor}}\|$, and in the coincident–branch limit they reduce identically to the unified single–cone structure of Definition 5.150. □

Remark 5.154 (Operational reading)**.** *External* constraint: the absolute kinematic boundary for influence is $J^\pm_\cup$ (union of branchwise futures; cf. Definition 5.123, Proposition 5.152, and Corollary 5.145). *Internal* modulation: along causally allowed directions, the branchwise temporal profiles $\mathcal{T}^{\text{HS}}_x(V)$ encode anisotropy/torsion of quantum-coherent transport (cf. Definitions 5.31 and 5.136); holosymmetric phase control restructures responses but not cone assignment (cf. Propositions 5.60 and 5.143, Corollary 5.126, and Definition 5.123).

5.4.6 The Primordial Origin of Causality

In the Interdependent Quantum Reality (IQR) framework, the causal structure of spacetime is not a pre-existent absolute backdrop. Rather, it *emanates from the quantum substrate*—specifically, from the properties of the *Primordial Quantum Singularities* (PQSs) that constitute the fundamental basis of relational ontology. The light cone, which governs the structure of causal propagation, is not a geometric constraint *at* a point in a manifold; it is a structure *of* a PQS.

Axiomatic and Geometric Foundations. The connection between the PQS and the light cone arises from three foundational constructs:

- **Relational Opacity of the PQS.** As defined in Definition 1.25, a PQS $\mathfrak{B}_x := \mathring{\mathbf{\Lambda}}_{x,\hat{\Theta}_x}$ is a quantum-relational singularity such that all relational measures within it vanish:

$$\forall x' \in \mathfrak{B}_x : \quad \hat{\mathbf{d}}(x, x') = \mathbf{0}, \quad t^{(\mathring{\mathbf{\Psi}})}_{\text{em}}(x, x') = 0.$$

  No internal causal, temporal, or spatial differentiation exists within $\mathfrak{B}_x$.

- **The Effective Lorentzian Metric.** As defined in Section 5.3.4, the emergent causal structure is governed by the symmetric part of the chronogeometric tensor:

$$\bar{g}^{\mathrm{Lor}}_{\mu\nu}(x) := \bar{g}_{\mu\nu}(x) + \bar{t}^{\mathrm{sym}}_{\mu\nu}(x),$$

  where each component is constructed as an expectation value in the universal state $|\mathring{\mathbf{\Psi}}\rangle$.

- **The Effective Light Cone.** As introduced in Definition 5.122, the effective light cone at the point x is the set:

$$\mathcal{V}_{\pm}(\mathfrak{B}_x) := \left\{ V^\mu \in T_x\mathcal{M} \;\middle|\; \bar{g}^{\mathrm{Lor}}_{\mu\nu}(x) V^\mu V^\nu = 0,\ V^\mu \neq 0 \right\}.$$

  This defines the absolute boundary for causal propagation from the PQS $\mathfrak{B}_x$.

The Causal Apex Condition. We now formalize the ontological grounding of the causal light cone in the following theorem:

Theorem 5.155 (Causal Apex Condition)**.** *Let* $\mathcal{C}_x := \{V^\mu \in T_x(\mathcal{M}) \mid \bar{g}^{\mathrm{Lor}}_{\mu\nu}(x) V^\mu V^\nu \leq 0\}$ *denote the causal cone at* $x \in \mathcal{M}_{\mathrm{geom}} \equiv \mathcal{M}^{(\mathring{\mathbf{\Psi}})}_{\mathrm{em}}$*. Then the apex of this causal cone is the Primordial Quantum Singularity* $\mathfrak{B}_x$*:*

$$\boxed{\mathrm{Apex}(\mathcal{C}_x) = \mathfrak{B}_x} \tag{5.111}$$

Proof Sketch. By construction, the effective metric $\bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)$ is defined only between *distinct* PQSs; it encodes relational structure via the unified expectation tensor $\bar{\Sigma}_{\mu\nu}(x)$ as in Definition 5.26. Within a single PQS $\mathfrak{B}_x$, all such relational observables vanish by definition (Proposition 1.28), and thus no internal tangent vectors can be defined. The light cone therefore has no geometric interior within $\mathfrak{B}_x$; its vertex must coincide with the singularity. □

Core Implications. This theorem confirms that the PQS functions as the *ontological origin* of causal flow in the emergent spacetime geometry. The implications are as follows:

- **PQS as Ontological Boundary.** Since no temporal or spatial separation is defined within $\mathfrak{B}_x$, it serves as the absolute boundary of any causal chain—an emergent "uncaused cause".

- **Unification of Ontology and Geometry.** The effective light cone is not an imposed geometric structure, but the direct expression of the relational opacity of the PQS. Geometry emerges from ontological indistinguishability.

- **Relational Emanation of Causality.** The causal structure of spacetime arises through a sequence of relational operations, as follows:

$$\boxed{\mathfrak{B}_x \;\rightsquigarrow\; \bar{\Sigma}_{\mu\nu}(x) \;\rightsquigarrow\; \bar{g}^{\mathrm{Lor}}_{\mu\nu}(x) \;\rightsquigarrow\; \mathcal{V}_{\pm}(\mathfrak{B}_x)}$$

Remark 5.156 (Holosymmetric compatibility of the causal apex)**.** The holosymmetric lift does not alter the ontological status of the apex: for each law branch k, the branchwise causal cone $\mathcal{C}_x^{(k)} := \{V \mid \bar{g}^{\mathrm{Lor}}_{(k)\mu\nu}(x)V^\mu V^\nu \le 0\}$ has apex $\mathfrak{B}_x$, hence $\mathrm{Apex}(\mathcal{C}_x^{(k)}) = \mathfrak{B}_x$. Consequently, the holosymmetric causal envelope built from the family $\{\mathcal{C}_x^{(k)}\}_k$ also has apex $\mathfrak{B}_x$.

Proposition 5.157 (Branchwise cones and no–law–signalling at the PQS)**.** *Let $\bar{g}^{\mathrm{Lor}}_{(k)}$ be the holosymmetric branch metrics (cf. Definition 5.123). Define the operational relations at x by*

$$\mathrm{Spacelike}_{\cap}(x,y) \iff (x,y) \text{ are spacelike w.r.t. each } \bar{g}^{\mathrm{Lor}}_{(k)}, \qquad J^{\pm}_{\cup}(x) := \bigcup_k J^{\pm}_{\bar{g}^{\mathrm{Lor}}_{(k)}}(x).$$

Then for local observables $\hat{\mathfrak{O}}(f), \hat{\mathfrak{O}}(h)$ supported in regions A, B near $\mathfrak{B}_x$ with $A \perp_{\cap} B$ one has

$$[\hat{\mathfrak{O}}(f), \hat{\mathfrak{O}}(h)] = 0,$$

and any influence from A is confined to $J^{+}_{\cup}(A)$. Phases used in holosymmetric control (φ, θ) do not modify $\mathrm{Spacelike}_{\cap}$ *or $J^{\pm}_{\cup}$ (cf. Proposition 5.124 and Corollary 5.126).*

Corollary 5.158 (Holosymmetric emanation of causality from the PQS)**.** *The relational emanation chain of Section 5.4.6 extends branchwise as*

$$\boxed{\mathfrak{B}_x \;\rightsquigarrow\; \left\{\bar{\Sigma}^{(k)}_{\mu\nu}(x)\right\}_k \;\rightsquigarrow\; \left\{\bar{g}^{\mathrm{Lor}}_{(k)\mu\nu}(x)\right\}_k \;\rightsquigarrow\; \left\{\mathcal{V}^{(k)}_{\pm}(\mathfrak{B}_x)\right\}_k}$$

with operational causal safety given by the intersection of spacelike regions and potential influence bounded by the union of futures (cf. Definition 5.123 and Proposition 5.124).

Proposition 5.159 (Stability of the apex cones under holosymmetric deformations)**.** *If the branch difference satisfies $\Delta\bar{g}^{\mathrm{Lor}} := \bar{g}^{\mathrm{Lor}}_{(1)} - \bar{g}^{\mathrm{Lor}}_{(0)}$ with $\|\Delta\bar{g}^{\mathrm{Lor}}\|_x$ small, then the null directions at $\mathfrak{B}_x$ tilt by at most $O(\|\Delta\bar{g}^{\mathrm{Lor}}\|_x)$; in the canonical pair (temporal-sector sign flip), the tilt is controlled by $\|\Delta\bar{t}^{\mathrm{sym}}\|_x$ (cf. Proposition 5.125). Hence the causal assignment near the apex varies continuously with the holosymmetric deformation.*

5.4.7 Holosymmetric Wordwise Primordial Origin of Causality

Typed channels and basic words.

- Channels: D (distance), T (time/phase), L (optional law–control).
- Words on (D, T): $\mathbf{g}_{\mu\nu}(x)$ (metric–sector word), $\mathbf{t}_{\mu\nu}(x)$ (temporal–sector word).
- Port symmetriser on index pairs: $\mathbb{S}[\cdot]$ (cf. Definition 5.87).
- Hadamard merge (channelwise sum): $\oplus$.
- Endogenous evaluation (Born map): $\langle\!\langle \cdot \rangle\!\rangle := \langle \mathring{\mathbf{\Psi}}| \cdot |\mathring{\mathbf{\Psi}}\rangle$.

PQS as relational–null apex (opacity).

Definition 5.160 (HS PQS opacity/compliance)**.** The PQS class at x is the compliance locus

$$\mathfrak{B}_x \;:=\; \{\, x' \;:\; \mathbf{d}(x,x') = \mathbf{0} \;\text{ and }\; \mathbf{t}(x,x') = \mathbf{0} \,\},$$

i.e. all distance/time word–measures vanish under the gauge–covariant transporter. No nontrivial tangent word exists *inside* $\mathfrak{B}_x$ (cf. Corollary 4.609 and Remark 5.52).

HS effective Lorentz word and quadratic classifier.

Definition 5.161 (HS Lorentzian merge and cone functional)**.** Define the *Lorentzian–merge word* (cf. Definitions 5.87 and 5.101)

$$\boxed{\;\mathbf{g}^{\mathrm{Lor}}_{\mu\nu}(x) \;:=\; \mathbf{g}_{\mu\nu}(x) \;\oplus\; \mathbb{S}\big[\mathbf{t}_{\mu\nu}(x)\big]\;}.$$

For an infinitesimal displacement word $\Delta\mathbf{x}^\mu$ at x, set the quadratic classifier

$$\mathcal{Q}_{\mathbf{g}^{\mathrm{Lor}}}(x;\Delta) \;:=\; \Delta\mathbf{x}^\mu \;\odot\; \mathbf{g}^{\mathrm{Lor}}_{\mu\nu}(x) \;\odot\; \Delta\mathbf{x}^\nu.$$

Classification: timelike/spacelike/null by the sign of $\mathcal{Q}_{\mathbf{g}^{\mathrm{Lor}}}$ (with $\Delta\mathbf{x} \neq 0$ for the null case). Endogenous evaluation yields $\langle\!\langle \mathbf{g}^{\mathrm{Lor}}_{\mu\nu} \rangle\!\rangle = \bar{g}_{\mu\nu} + \bar{t}^{\mathrm{sym}}_{\mu\nu} =: \bar{g}^{\mathrm{Lor}}_{\mu\nu}$ (cf. Proposition 5.90).

HS causal apex theorem.

Theorem 5.162 (HS causal apex)**.** *Let* $\mathcal{N}^{\mathrm{HS}}_x := \{\, \Delta\mathbf{x} \neq 0 :\; \mathcal{Q}_{\mathbf{g}^{\mathrm{Lor}}}(x;\Delta) = 0 \,\}$ *be the HS null cone at* x*. Its apex is the PQS class:*

$$\boxed{\;\mathrm{Apex}\big(\mathcal{N}^{\mathrm{HS}}_x\big) \;=\; \mathfrak{B}_x\;}.$$

Idea. Inside $\mathfrak{B}_x$ one has $\mathbf{d} = \mathbf{0}$ and $\mathbf{t} = \mathbf{0}$, so no nontrivial displacement word $\Delta\mathbf{x}$ is representable; the cone therefore emanates *from* the compliance locus. Under $\langle\!\langle\cdot\rangle\!\rangle$, $\mathcal{N}^{\mathrm{HS}}_x$ maps to $\{V : \bar{g}^{\mathrm{Lor}}(V,V) = 0\}$ whose apex is the same locus at expectation level (cf. Section 5.4.2 and Definition 5.129). □

Holosymmetric (two–law) envelope and no–law–signalling.

Definition 5.163 (Branchwise cones and operational relations)**.** With law register $\mathcal{H}_L = \mathrm{span}\{|0\rangle_L, |1\rangle_L\}$, define branches

$$\mathbf{g}^{\mathrm{Lor}}_{(k)\mu\nu} := \mathbf{g}_{(k)\mu\nu} \;\oplus\; \mathbb{S}[\mathbf{t}_{(k)\mu\nu}], \qquad \mathcal{N}^{\mathrm{HS},(k)}_x := \{\Delta\mathbf{x} \neq 0 :\; \mathcal{Q}_{\mathbf{g}^{\mathrm{Lor}}_{(k)}}(x;\Delta) = 0\}, \quad k \in \{0,1\}.$$

Let $J^{\pm}_{(k)}(x)$ be the futures/pasts for $\mathbf{g}^{\mathrm{Lor}}_{(k)}$. Define

$$\mathrm{Spacelike}_{\cap}(A,B) :\Longleftrightarrow\; A, B \text{ spacelike in both branches}, \qquad J^{\pm}_{\cup}(x) := J^{\pm}_{(0)}(x) \cup J^{\pm}_{(1)}(x).$$

The apex is $\mathfrak{B}_x$ in each branch.

Proposition 5.164 (HS microcausality and causal envelope)**.** *For local word–observables* $\hat{\mathfrak{O}}(f)$, $\hat{\mathfrak{O}}(h)$ *with supports* A, B,

$$A \perp_{\cap} B \implies [\hat{\mathfrak{O}}(f), \hat{\mathfrak{O}}(h)] = \mathbf{0}, \qquad \text{and signals from } A \text{ are confined to } J^{+}_{\cup}(A).$$

HS control phases (φ, θ) *change interference weights but not branchwise cones (cf. Proposition 5.132).*

Stability of the apex cones.

Proposition 5.165 (Cone stability under HS deformations)**.** *Let* $\Delta\mathbf{g}^{\mathrm{Lor}} := \mathbf{g}^{\mathrm{Lor}}_{(1)} - \mathbf{g}^{\mathrm{Lor}}_{(0)}$. *For any* $\mathbf{g}^{\mathrm{Lor}}_{(0)}$*-null direction* $\Delta\mathbf{x}$,

$$\left| \mathcal{Q}_{\mathbf{g}^{\mathrm{Lor}}_{(1)}}(x; \Delta) \right| = \left| \langle \Delta\mathbf{x},\ \Delta\mathbf{g}^{\mathrm{Lor}}(x)\ \Delta\mathbf{x} \rangle \right| \ \leq\ \|\Delta\mathbf{g}^{\mathrm{Lor}}(x)\|_{\mathrm{op}}\ \|\Delta\mathbf{x}\|^2.$$

In the canonical pair $(\mathbf{t} \mapsto -\mathbf{t})$, *the tilt is governed by* $\|\Delta\,\mathbb{S}[\mathbf{t}]\|_{\mathrm{op}}$ *(cf. Proposition 5.133).*

Emanation chain (wordwise → classical).

$$\boxed{\mathfrak{B}_x \ \rightsquigarrow\ \mathbf{g}_{\mu\nu} \oplus \mathbb{S}[\mathbf{t}_{\mu\nu}] \ \rightsquigarrow\ \mathcal{N}^{\mathrm{HS}}_x \xRightarrow{\langle\!\langle\cdot\rangle\!\rangle} \bar{\Sigma}_{\mu\nu} \ \rightsquigarrow\ \bar{g}^{\mathrm{Lor}}_{\mu\nu} \ \rightsquigarrow\ J^{\pm}(\mathfrak{B}_x)}$$



Figure 5.1: Unified causal structure on $\mathcal{M}_{\mathrm{geom}}$ (single history). The apex is the PQS $\mathfrak{B}_x$ (Theorem 5.155). The absolute kinematic boundary is the Effective Light Cone from Definition 5.122, defined by $\bar{g}^{\mathrm{Lor}}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}^{\mathrm{sym}}_{\mu\nu}$ (Definition 5.150). The temporal phase-flow quadric (level sets of $\bar{t}^{\mathrm{sym}}_{\mu\nu}$) modulates coherent transport inside the cone (Definition 5.136).

Explanation of the Diagram (fig. 5.1)

- *The Gray Cone:* This is the **Effective Light Cone**, the absolute boundary of causality.

Crucially, it is defined by the null condition of the *effective Lorentzian metric*,

$$\bar{g}^{\mathrm{Lor}}_{\mu\nu}(x) := \bar{g}_{\mu\nu}(x) + \bar{t}^{\mathrm{sym}}_{\mu\nu}(x).$$

This signifies that the boundary of what is causally possible is determined by a synthesis of spatial distinguishability, via $\bar{g}_{\mu\nu}(x)$, and the symmetric component of temporal phase flow, $\bar{t}^{\mathrm{sym}}_{\mu\nu}(x)$. Any physical trajectory must possess a tangent vector that lies on or within this composite cone.

- *The Blue Ellipsoid:* This represents the **Temporal Phase-Flow Quadric**. It is a level set of the quadratic form defined by $\bar{t}^{\mathrm{sym}}_{\mu\nu}(x)$, the symmetric part of the temporal tensor. Its shape and orientation encode the anisotropy in the flow of quantum phase. Directions aligned with its principal axes correspond to extremal values of temporal curvature, and thus to paths along which the phase of a quantum system rotates most rapidly or most slowly.

- *Composite Structure:* The complete causal structure is given by the interplay of both components. The blue quadric resides strictly *within* the gray light cone. This illustrates the two-tiered character of causality in this framework: the light cone defines the absolute, "hard" boundary of permissible causal propagation, while the quadric characterizes the "soft", anisotropic modulation of phase evolution within the causally allowed domain.

FLRW (GR and QFT) Relational Microcausality and Spectral Positivity

Purpose and scope. This subsection records the conditions under which the relational chronogeometry developed in Chapter 6 reduces cleanly to classical general relativity and to standard quantum field theory on curved spacetime in the FLRW regime, while emphasising that the framework is not restricted to that case. Causality is anchored to the effective Lorentzian metric constructed from the expectation-level chronogeometric data (see Section 5.2.6), so in the FLRW setting one sets $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ equal to the FLRW line element (in cosmic or conformal time) [7, 319]. The causal futures/pasts $J^{\pm}$ are defined with respect to $(\mathcal{M}, \bar{g}^{\mathrm{Lor}})$ in the usual way, and microcausality is imposed by local commutativity of observables at $\bar{g}^{\mathrm{Lor}}$-spacelike separation [230, 255]. When the temporal sector $\bar{t}_{\mu\nu}$ (from Definition 5.26) is homogeneous and isotropic and aligned with the FLRW foliation (no off-diagonal $0i$ components and $t_{ij} \propto \gamma_{ij}$ on spatial slices), it does not tilt the null structure; consequently the cones computed from $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (cf. Definitions 5.25 and 5.26) coincide with those of $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$, i.e. $J^{\pm}_{\bar{\Sigma}} = J^{\pm}_{\bar{g}^{\mathrm{Lor}}}$, and the locality structure matches classical FLRW.

Spectral stability. Spectral positivity is fixed in a cosmologically natural way by choosing the relevant flow (either conformal-time evolution generated by ∂_η or the comoving four-velocity u^μ) and requiring positivity of the corresponding generator/energy [230, 255]. In non-stationary backgrounds, admissible states are taken to be Hadamard/adiabatic in the sense of QFT on curved spacetime; in stationary subcases (e.g. de Sitter patches), KMS positivity may be invoked [13, 244, 11, 12, 314]. This recovers the standard stability/positivity requirements of QFT in FLRW within the relational formalism.

Holosymmetry in FLRW. With homogeneous, isotropic smearing $J^{\mu\nu}(\eta)$, the inter-law coherence operator $\hat{\mathbb{C}}^{(\theta)}_{\mu\nu}$ of the triadic holosymmetric construction (Definition 5.56 and Proposition 5.57) becomes spatially scalar on each slice; cross-channel commutators are suppressed or vanish by symmetry (Corollary 5.58), and the holosymmetric interference reduces to sector-wise, time-dependent responses without upsetting FLRW causality (Proposition 5.60). Operational time reversals are realised as forward–backward evolutions in conformal (or cosmic) time (Remark 5.62), providing a controlled Loschmidt-type probe of cosmological irreversibility while leaving the FLRW light-cone and spectral positivity intact [426, 427].

Lemma 5.166 (FLRW Relational Microcausality and Positivity)**.** *As a direct consequence of the* Principle of Coherent Regularization *(cf. Section 4.9.13), the two–point functions of the substrate fields satisfy positivity and causal support relative to the emergent chronogeometry. In the GNS representation (cf. Definition 1.34 and Section 4.1.3) of the observable algebra associated with the vacuum* Quantum Holo–State $|\mathring{\mathbf{\Psi}}\rangle$ *(Definition 1.27), the two–point functions of the* Gaiamiti field $\hat{\mathbf{\Phi}}(x)$ *(Definition 6.2) and the* Tantrakyma field $\hat{\mathbf{\Theta}}(x)$ *(Definition 6.3) admit a* generalized Källén–Lehmann/Stieltjes representation *[255, 230, 241, 381] with positive spectral measures:*

$$W_{\mathbf{\Phi}}(x,y) := \langle\mathring{\mathbf{\Psi}}|\,\hat{\mathbf{\Phi}}(x)\,\hat{\mathbf{\Phi}}(y)\,|\mathring{\mathbf{\Psi}}\rangle = \int_{[0,\infty)} G_{\mu^2}(x,y)\,\mathrm{d}\rho_{\mathbf{\Phi}}(\mu^2), \qquad \mathrm{d}\rho_{\mathbf{\Phi}}(\mu^2) \geq 0,$$

$$W_{\mathbf{\Theta}}(x,y) := \langle\mathring{\mathbf{\Psi}}|\,\hat{\mathbf{\Theta}}(x)\,\hat{\mathbf{\Theta}}(y)\,|\mathring{\mathbf{\Psi}}\rangle = \int_{[0,\infty)} G_{\mu^2}(x,y)\,\mathrm{d}\rho_{\mathbf{\Theta}}(\mu^2), \qquad \mathrm{d}\rho_{\mathbf{\Theta}}(\mu^2) \geq 0,$$

where $G_{\mu^2}(x,y)$ is the relational scalar Green kernel *built on the expectation–level chronogeometric background $\bar{\Sigma}_{\mu\nu}$ (Definition 5.26), in line with the curved–spacetime treatment of Green kernels [244, 245, 13, 12]. Moreover, the commutators vanish outside the relational light–cone:*

$$[\hat{\mathbf{\Phi}}(x),\hat{\mathbf{\Phi}}(y)] \;=\; [\hat{\mathbf{\Theta}}(x),\hat{\mathbf{\Theta}}(y)] \;=\; 0 \quad \text{if } (x-y)\notin J^{\pm}_{\bar{\Sigma}},$$

where $J^{\pm}_{\bar{\Sigma}}$ are the causal sets determined by $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$.

Remark 5.167 (Context, Interpretation, and Cross–References)**.** This lemma should be read as the *curved/relational analogue* of the Källén–Lehmann representation [14, 82, 431], with the following clarifications and dependencies:

1. **Precise meaning of symbols.** $W_{\mathbf{\Phi}}(x,y)$ and $W_{\mathbf{\Theta}}(x,y)$ are two–point functions in the GNS Hilbert space $\mathcal{H}_{\mathrm{em}}$ (cf. Definition 1.34, see also [241, 381]). $G_{\mu^2}(x,y)$ is the scalar Green kernel on $(\bar{\Sigma}_{\mu\nu},\bar{\nabla})$, solving $(-\bar{\nabla}_\mu\bar{\Sigma}^{\mu\nu}\bar{\nabla}_\nu+\mu^2)G_{\mu^2}=\delta$ with the chosen boundary condition [244, 245, 13, 12]. The spectral measures $\rho_{\mathbf{\Phi}}$ and $\rho_{\mathbf{\Theta}}$ are positive Borel measures on $[0,\infty)$; in general, $\mathrm{d}\rho = \rho(\mu^2)\,\mathrm{d}\mu^2 + \sum_k Z_k\,\delta(\mu^2-m_k^2)$ [432, 253, 384].
2. **State–conditioned origin of the spectral measure.** By the construction of the *state–conditioned modal weight* (Section 4.9.7), the measures $\rho_{\mathbf{\Phi}},\rho_{\mathbf{\Theta}}$ are not free densities but are determined by the Quantum Holo–State $|\mathring{\mathbf{\Psi}}\rangle$. This ensures that the spectral content of propagators is endogenous to the relational substrate.

3. **Endogenous regularization and finiteness.** The *Principle of Coherent Regularization* (Section 4.9.13) guarantees that these measures are finite and positive, while the *endogenous finiteness of spectral distance* (Section 4.9.11) proves that divergences are eliminated without the need for external cutoffs. Together, these results explain why the Stieltjes integrals defining $W_{\mathbf{\Phi}}, W_{\mathbf{\Theta}}$ converge.
4. **Self–regularizing geometry and IR control.** The *self–regularizing geometry* (Section 4.9.14) ensures the UV consistency of the spectral construction, while the *relational framework for infrared regularization* (Section 4.9.15) controls the long–distance behaviour of the correlators. Hence the representation remains valid across all scales.
5. **Relational microcausality.** If $\Delta_{\mu^2}(x,y)$ is the Pauli–Jordan bitensor associated with G_{μ^2}, then each Δ_{μ^2} has support only inside the $\bar{\Sigma}$–causal cone. Integrating against a positive measure preserves this support:

$$[\hat{\mathbf{\Phi}}(x), \hat{\mathbf{\Phi}}(y)] = \int_{[0,\infty)} \Delta_{\mu^2}(x,y)\, \mathrm{d}\rho_{\mathbf{\Phi}}(\mu^2) = 0,$$

$$[\hat{\mathbf{\Theta}}(x), \hat{\mathbf{\Theta}}(y)] = \int_{[0,\infty)} \Delta_{\mu^2}(x,y)\, \mathrm{d}\rho_{\mathbf{\Theta}}(\mu^2) = 0$$

for $(x,y) \notin J^{\pm}_{\bar{\Sigma}}$ [255, 230, 241].

Thus, relational microcausality and spectral positivity are unified consequences of the GNS/Holo–State construction, the Fourfold correlator geometry (Section 4.8), and the spectral framework developed in Section 4.9. They extend naturally to the spin–2 sector of the Gaiamiti fluctuations (cf. Definition 10.188, eqs. (10.101) and (10.106), and Sections 9.17.2 and 10.5), where the transverse–traceless propagator (eq. (10.128) and Proposition 10.192) is likewise confined to the relational light cone.

Causality in IQR (general, independent of FLRW).

The following principles govern causality in the Interdependent Quantum Reality (IQR) framework.

1) Relational (not absolute) notion of cause. Causality is a property of relations among subsystems, encoded at the operator level by locality and the commutation structure of the observable net [230, 255, 3]. The foundational objects are the operator-valued chronogeometry $\hat{\Sigma}_{\mu\nu} = \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu}$ and its expectation $\bar{\Sigma}_{\mu\nu}$ in the Quantum Holo–State $|\mathring{\mathbf{\Psi}}\rangle$ (see Definitions 5.25 and 5.26). Whenever a smooth Lorentzian sector emerges, the *effective cones* that organise locality are anchored to the associated effective Lorentzian tensor $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ constructed from the expectation-level data (see Section 5.2.6; cf. [7]). Microcausality is then imposed as local commutativity for test-function supports spacelike with respect to those cones [230, 255]. In regimes without a smooth manifold, causality is imposed algebraically: retarded/advanced response functionals built from $\hat{\Sigma}$ define a *proto-causal order* (vanishing of retarded correlators outside the emergent cones), which reduces to the geometric notion once $\bar{g}^{\mathrm{Lor}}$ exists.

2) Spectral positivity along admissible time flows. Stability is enforced by *spectral positivity* for generators of admissible relational time flows (e.g., flows induced by Killing fields when present, or relational clocks defined by the state–algebra pair). In stationary sectors this includes KMS/thermal conditions; in non-stationary sectors, Hadamard/adiabatic regularity is required to ensure a lower-bounded energy content along any such flow [230, 13, 244, 11, 12]. In the geometric limit, these prescriptions reproduce the standard positivity used in QFT on curved spacetimes [11, 13].

3) Holosymmetry and "no-law-signalling." Holosymmetry permits coherent superpositions of *histories/laws*, but does not license superluminal signalling. Operational causality is enforced *branchwise* and *robustly*: (i) For any pair of histories with effective cones $J^{\pm}_{(k)}$, observables supported in regions that are spacelike in *each* history must commute; holosymmetric control operations are required not to increase commutators across such pairs of regions (cf. [230, 255]). (ii) Interference phases (φ, θ) modify response functions but leave the underlying cone assignment invariant; they provide tomography of T-even (anticommutator) and T-odd (commutator) structure without altering causal order (see Proposition 5.60).

4) Channel structure and locality. With the triadic decomposition (Definition 5.54) and its holosymmetric push-through (Proposition 5.57), locality constraints propagate *per channel*. When channel projectors commute (Corollary 5.58), cross-channel commutators that could blur localisation vanish, so causality reduces to parallel, sector-wise microcausal statements. When channels do not commute, interference terms are present but still respect the branchwise cones fixed above.

5) Time reversal: operational, not antiunitary. IQR distinguishes between the antiunitary time-reversal operator T (not implementable as a controlled branch) and *operational reversals* (controlled comparisons of U with $U^\dagger$ or T-conjugate unitaries). The latter are permitted and diagnostically useful—yielding Loschmidt–echo overlaps—yet do not modify the cone assignment; they probe T-odd structure while remaining compatible with microcausality and spectral positivity (see Remark 5.62; cf. [350, 287, 426, 427]).

6) Classical limits as special cases, not constraints. The FLRW specialisation in Section 5.4.7 is an *example*: when $\bar{t}_{\mu\nu}$ is homogeneous/isotropic and aligned with the FLRW foliation, the null structure of $\bar{\Sigma}_{\mu\nu}$ coincides with that of $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ and the usual GR/QFT microcausality and positivity statements are recovered (cf. [7, 244, 11]). This demonstrates that IQR *contains* the classical picture where appropriate, while the general causal understanding remains the relational, operator-anchored scheme above.

Causality in IQR is *relational and operational*: it is fixed by (i) the emergent chrono-geometric data that define effective cones (geometric when available, algebraic otherwise), (ii) spectral positivity along admissible time flows, and (iii) holosymmetric no-law-signalling constraints that preserve microcausality even in superpositions of laws—of which FLRW/GR is a consistent and illustrative limit, not a limitation [230, 11].

Holosymmetric synopsis. Reality is treated as an object that can be placed on a *quantum balance*: holosymmetry prepares two distinct chronogeometric histories (two candidate spacetimes with their own geometries and temporal flows) and interferes them under a controllable phase (Definition 5.56 and Proposition 5.57; [287, 3, 23, 24]). The control-qubit readout is a direct meter of *relational proximity* between the two histories: high visibility indicates similarity, low visibility indicates divergence. In the small-difference regime, the visibility reduces to a Fisher-information–type susceptibility, yielding a principled "distance on law space" that quantifies the sensitivity of the chronogeometric operator to perturbations [359, 423, 407].

The interference is not merely binary. By sweeping the control phase, contributions from the *metric sector* $\hat{g}$ ("how far apart") and the *temporal sector* $\hat{t}$ ("how out of phase") are separated: the symmetric (anticommutator) piece deforms the expectation-level geometry, whereas the antisymmetric (commutator) piece exposes phase curvature and transport effects associated with time flow and Berry-type structure (Proposition 5.60; [350, 25, 24, 424, 408]). This equips the framework with tomography over chronogeometry. Resolving $\hat{\Sigma}$ into triadic channels—correlational, vacuum, spectral—preserves the structure and enables sector-wise tests with optional channel selectivity (Definition 5.54, Corollary 5.58, and Remark 5.61), including localized probes via a smearing tensor.

Operational time reversals are admissible and informative. By pairing a history with its backward counterpart—either the unitary pair U_0 and $U_0^\dagger$ or, more generally, T-conjugate unitaries—the interference signal functions as a Loschmidt-echo–type probe of reversibility; its visibility and phase serve as calibrated witnesses of T-even (anticommutator) vs. T-odd (commutator) structure [426, 427]. A small-deformation expansion links echo decay to the variance of the chronogeometric deformation, providing an information-geometric measure of the arrow of time and quantitative reversibility bounds in law space [359, 423, 407]. Channel-selective flips confine partial time reversals to chosen sectors and can be localized by the smearing tensor (Remark 5.61). A crucial constraint remains: the fundamental time-reversal operator T is antiunitary and cannot appear as a controlled branch; holosymmetry therefore compares unitary histories related by reversal or T-conjugation rather than applying T directly (Remark 5.62; [350, 287]).

Part III

Laws, States and Fundamental Fields

Chapter 6

The Gaiamiti and Tantrakyma Fields

6.1 The Nature of the Fundamental Fields

In the Interdependent Quantum Reality (IQR) framework, the physical content of the universe is not described by fields living *on* a prior spacetime manifold. Rather, fields *are* the fundamental constituents of the structured singularity,

$$\hat{\mathbf{\Omega}}(x) \;:=\; {}_{\hat{\mathbf{\Phi}}}\mathring{\mathbf{\Lambda}}_{\hat{\mathbf{\Theta}}}(x),$$

evaluated at each locus x of the foundational continuum (the PQS apex $\mathfrak{B}_x$; cf. Theorem 5.155 and Definition 5.122). Here $\mathring{\mathbf{\Lambda}}$ is the atomic prismatic of the fourfold dialectic, and the subscripts indicate the holosymmetric packaging by the geometric (Gaiamiti) and internal–structural (Tantrakyma) sectors (cf. Section 1.2.5). Geometry and matter-like content therefore arise inseparably; to make this co-arising explicit, the geometric and the internal–structural principles are packaged as a single mathematical object.

Definition 6.1 (Generalized Field Operator $\hat{\mathbf{F}}(x)$)**.** At each point x in the emergent quantum configuration space $\mathcal{X}_q$ (cf. Section 4.19.3), the complete operator-valued content of a Primordial Quantum Singularity is described by the *Generalized Field Operator*,

$$\hat{\mathbf{F}}(x) \;:=\; \begin{pmatrix} \hat{\mathbf{\Phi}}(x) \\ \hat{\mathbf{\Theta}}(x) \end{pmatrix}.$$

The two components of this unified operator represent the two fundamental facets of emergent reality, and—after GNS realization—they act in the emergent operator arena $\mathrm{End}(\mathcal{H}_{\mathrm{em}})$ determined by the Quantum Holo–State (cf. Section 1.2.9 and Definition 1.18).

Concretely, $\hat{\mathbf{\Phi}}(x)$ is the *Gaiamiti* component—the architect of chronogeometry that selects admissible distances, durations, and curvatures (cf. Definitions 5.26 and 5.122)—while $\hat{\mathbf{\Theta}}(x)$ is the *Tantrakyma* component—the woven–wave framework whose internal symmetries and modal excitations realize conserved charges, flavors, and particle-like content within that geometry (cf. Definition 5.25). Placed together as the field-space doublet $\hat{\mathbf{F}}(x)$, these sectors are not merely juxtaposed but structurally unified at the level at which the singularity $\mathfrak{B}_x$ is itself defined.

This unification is conceptually analogous to the electroweak insight: a single formal structure that embraces two a priori distinct interactions. In the Standard Model, Glashow first proposed the unified electroweak framework [57], with Weinberg and Salam independently completing the picture by exhibiting the symmetry structure and the mechanism by which masses arise [58, 59]. In IQR, the analogy is not a gauge-group identification but a unifying *field-theoretic packaging*: geometry-generation (gaiamiti) and content-generation (tantrakyma) are treated as inseparable components of one operatorial entity. The result is a clear statement of principle: spacetime structure and internal structure are co-constituted—two voices written on a single staff—so that what can propagate (modes, charges, correlations) is determined together with where it can propagate (chronogeometric order) (cf. Propositions 5.152 and 5.153 and Definition 5.123).

Gaia: The Primordial Mother

In the oldest Greek cosmogony to survive, Hesiod's *Theogony*, the first stirrings of the universe are startlingly spare: *Chaos* yawns open, and "next" comes **Gaia**, wide-bosomed Earth—the "ever-sure foundation of all." From herself alone she brings forth *Ouranos* (Sky) as a canopy for the gods, the *Ourea* (mountains), and *Pontos* (the sea). Only afterwards, lying with the Sky she herself made, Gaia bears the great line of *Titans*—Oceanus, Coeus, Crius, Hyperion, Iapetus, Theia, Rhea, Themis, Mnemosyne, Phoebe, Tethys—and the youngest, *Kronos*, who already hates his father. This sequence, both austere and audacious, frames Greek creation not as a "making-from-nothing," but as generation and separation—Earth unfolding the world out of her own body [60].

When Ouranos stuffs his children back into their mother—Cyclopes and Hundred-Handers alike—Gaia groans under the pressure and *invents a tool*: a jagged sickle of unbreakable stone. She steels the reluctant Titans; only Kronos accepts the charge. In the night, as Sky comes to lie with Earth, Kronos leaps from ambush and *cuts off his father's genitals*, sundering Heaven from Earth. From the blood that spatters into Gaia arise the *Erinyes (Furies)*, the *Giants*, and the *Meliai* (ash-tree nymphs); from the foam around the cast-off genitals in the sea, *Aphrodite* will emerge. In Hesiod's telling, cosmic order is born through painful surgery: hierarchy, distance, and distinct domains carved where there had been suffocating union [60].

Kronos, now king, repeats the pattern of fear. Warned he will be overthrown by a child, he swallows his offspring. Gaia and her daughter *Rhea* devise a counter-plot: the newborn *Zeus* is spirited away to a Cretan cave, while Kronos gulps a *stone wrapped in swaddling-cloths*. Later, "by the deep counsels of Earth," Kronos is forced to disgorge the children in reverse order, beginning with the stone, which Zeus sets up at *Pytho (Delphi)* as a sign forever. The stone becomes both relic and proof of succession; Delphi will be one of the places where Earth's voice is remembered [60].

That memory is dramatized in Aeschylus' *Eumenides*. The *Pythia* opens by naming *Earth (Gaia)* as the *first oracle-holder* at Delphi, succeeded by *Themis*, then *Phoebe*, who gifts the seat to *Apollo*. Here, prophecy itself is an inheritance that begins in the chthonic mother and passes, through goddess to goddess, into the Olympian order—another succession myth, but one of *voice* rather than *violence* [62].

Beyond the grim politics of the *Theogony*, Gaia is also praised in hymnic mode. *Homeric Hymn 30* hails her as "well-founded" and "mother of all," feeder of everything that walks,

swims, or flies; the hymn ends with the poet asking simply for prosperity "in exchange for song." This is the other axis of her power: not rebellion, but *fecundity and stability*, the soil as common gift [61].

Cult and image trace those themes. In Attic myth, when Hephaestus' desire for Athena goes astray, the seed that falls upon Earth yields *Erichthonius*, the autochthonous child of Athens. Vase painters render the moment with unforgettable economy: *Gaia rising waist-high from the ground*, handing the infant to Athena—a visual theology of origins in which the city's very people are "born from the soil" they inhabit. (One red-figure hydria in the British Museum preserves the scene in vivid detail) [63].

Name and Language. The name **Γαῖα (Gaîa)** is a poetic or epic collateral of **Γῆ (Gê)**, the ordinary word for "earth" (Doric **Γᾶ**). Modern etymologists generally judge the root to be **pre-Greek** (non-Indo-European in origin), part of a substrate of ancient Aegean vocabulary absorbed into Greek; that fits the goddess's character as something felt to be more **primordial** than the Indo-European sky-father pattern [64, 65].

Applications and Analogues. Later Greeks and Romans mapped Gaia onto *Tellus/Terra Mater*, Rome's earth-mother honored in agricultural festivals like the *Fordicidia*, and paired symbolically with a sky god (Caelus/Uranus). The analogy is not perfect—the Roman cult has its own history—but the identification was standard in antiquity and remains useful as a cross-cultural pointer.

Read together, these scenes sketch a coherent arc: *Gaia births*, *Gaia separates*, *Gaia advises*—and then, as order matures, *Gaia recedes*, her functions redistributed (to Olympian kingship, to Apollo's prophecy, to the civic soil of Athens). The myths preserve tension between nurture and necessity: the same mother who feeds all beings also engineers the decisive cut that makes a structured world possible. That doubleness—soil and scalpel—is what keeps Gaia's figure vivid: she is both the ground we stand on and the deep wisdom that knows when to part things so that life can breathe.

Miti: The Primordial Measure

If *Gaia* is ground, **miti** is grasp. In Sanskrit, *miti* denotes *measure*, and in the classical compound *jyā-miti*—literally "*bowstring-measure*"—it names what later languages call geometry. The image is tactile before it is abstract: a string drawn taut across a curve, a bowstring stretched across an arc. The world offers curvature; the hand offers a straight line; *miti* is the act that relates the two and renders the relation legible [66].

Bowstring Before Blueprint. The two parts of the word carry the program. *jyā* is the bowstring, the clean span that cuts across a circular sweep; *miti* is both the act and the outcome of measuring. In *jyā-miti*, operation and instrument are fused: geometry begins as a manual art with cord, peg, and arc rather than as a disembodied view from nowhere. Hence the early attention to worked procedures and practical tables—the craft grows from *operations on strings*, from repeating and recording what a taut line discloses when it is laid across different arcs [66, 34].

From Altar-Cords to Celestial Arcs. Across early sources two crafts converge on this same gesture. In the *Śulba-sūtras*, cords lay out sacrificial grounds—right angles, diagonals, areas—so that space can be domesticated for ritual. In astronomical practice, arcs on the sky are converted into portable quantities, so that *time* itself—season, eclipse, planetary return—can be domesticated by measure. The bowstring is the bridge between ground plan and firmament: a device that turns curved givens into storable, comparable data [34, 67].

The Indian "bowstring" becomes a technical term that travels widely. As scholars and calculators tabulate chord-based quantities, the vocabulary passes through new languages and scripts, eventually yielding the familiar term *sine*. What survives in this journey is not a particular diagram but a habit: curves can be *encoded* by systematic records of straight spans. The bowstring becomes a number; the number becomes a table; the table becomes a means to navigate both earth and sky [68, 67].

What miti Contributes. Unlike "geometry," which already presumes a domain ("earth-measure"), *miti* names a *portable operation*. To measure is to *relate*: to set one thing against another by a standard, to compare extents, to fix limits and rules. In Sanskrit usage, the semantic field around *miti* touches boundaries and norms as well as quantities; it shades into epistemic measure, where the question is not only how long something is, but by what *means* knowledge of it is secured. Thus *miti* marks a crossroads: a *material act*—lay the string; read the result—and a *cognitive act*—establish the standard; judge the fit. Out of this doubleness, *practice yields principle* [66].

Applicaitons and Analogues. The operational logic persists. In many modern Indo-Aryan vernaculars, *jyāmiti* remains the everyday word for geometry, a straight line from bow to blackboard. In *astronomy*, chord- and sine-based tables underpin eclipse prediction and planetary models, translating arcs of motion into computable form. In *architecture and surveying*, cord-and-peg constructions anchor rules of layout and proportion, turning space into a sequence of reliable acts. In *language*, derivatives such as *pari-miti* ("limit") and *mita* ("measured") carry the same root into ideas of boundary and moderation, showing how the discipline of measure extends from fields and circles to norms and judgment [34, 67, 66].

The Apeiron: The Boundless and the Genesis of Form

The **ἄπειρον** (*apeiron*)—literally "the un-limited," from privative ἀ- + πεῖραρ/πέρας (limit, boundary)—names, in classical usage, an *indefinite* and *unbounded* source rather than a completed infinite. In this treatise we deliberately *redeploy* the term to denote the *quantum pulse of the Gaiamiti field*: the minimal event of givenness that is predicate-free until taken up by measure. [69, 72, 71]

Anaximander's *archē*: Indefiniteness, Reciprocity, Return. **Anaximander** is our earliest explicit witness. The single fragment (DK 12 B1), preserved by Simplicius, makes the *apeiron* (ἄπειρον) the *archē* (ἀρχή) from which things come-to-be and to which they return "paying penalty and recompense … according to the ordering of time." Read plainly, the

language enforces reciprocity upon differentiation: what emerges must also return, and excess is corrected by time's order. [69, 70, 71]

Limit and the Unlimited: The Pythagorean Polarity. Early **Pythagorean** schemata oppose *peras* (πέρας, limit) to *apeiron* (ἄπειρον, the unbounded): limit is the creative cut that renders things countable, comparable, and commensurable; the unbounded is the "more and less" that requires measure to become knowable. Harmony and ratio are thus arts of limiting, not denials of the boundless. [72, 70]

Aristotle's Discipline of the Infinite. **Aristotle** reframes the unlimited as a *mode* of process: the infinite is *potential*, never actual—there is always "another and another" in counting or division, but never a finished infinite whole. This prevents reifying the *apeiron* into a substance while preserving its role wherever iteration lacks a terminal bound. [71, 73]

Semantic Field and Working Lexicon. Philologically, ἀ- + πεῖραρ/πέρας yields "without boundary"; philosophically, two senses have been active since antiquity: (i) **indefinite** (no fixed predicate) and (ii) **unbounded** (no fixed quantitative limit). We keep those senses, but *at the scale of a quantum occurrence*. For this work:

1. **Indefinite unit-event**: the minimal pulse of givenness, discrete yet anonymous until articulated.
2. **Pre-metric**: prior to scale; magnitude and comparison arise only with measure.
3. **Rhythmed by time**: issued and recalled under an ordering that enforces reciprocity. [69]
4. **Articulated by limit**: rendered intelligible through *peras*, the creative cut enabling ratio and harmony. [72, 71]

In this cadence, *Gaia* names the felt *ground of appearing*; *miti* names the *act of measure* by which appearing becomes form; the ἄπειρον names the *quantum pulse* by which ground is brought to appearance for measure. We treat these as a single motif—genesis, rhythm, return, and limit—sufficient for what follows. Henceforth ἄπειρον designates the quantum pulse of the Gaiamiti field; its formal definition and dynamical properties are stated in Definition 9.246.

The Gaiamiti Field: The Generator of Relational Geometry

Definition 6.2 (The Gaiamiti Field and its Modal Decomposition)**.** The *Gaiamiti Field*, $\hat{\mathbf{\Phi}}(x)$, is the first component of the Generalized Field Operator. It is a Hermitian, operator-valued field whose relational correlations give rise to the emergent metric and temporal tensors, $\bar{g}_{\mu\nu}(x)$ and $\bar{t}_{\mu\nu}(x)$.

It admits a modal decomposition over the four-dimensional local Hilbert space $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}$, spanned by the basis of dialectical modes $\{|\alpha\rangle_x\}$ where $\alpha \in \{0, 1, p, s\}$:

$$\hat{\mathbf{\Phi}}(x) = \sum_{\alpha} \hat{\Phi}_\alpha(x)\, |\alpha\rangle_x$$

Each *operator-valued field mode*, $\hat{\Phi}_\alpha(x)$, is an operator acting on the total emergent Hilbert space $\mathcal{H}_{\text{em}}$, representing the geometric potential associated with a specific dialectical mode (Symmetry, Antisymmetry, Parasymmetry, or Holosymmetry).

To capture its role as a generator of *relational* geometry and time, we designate the field the **Gaiamiti Field**. The coinage fuses two deep, ancient strands: *Gaia*, the primordial ground, and *miti*, the act of measure. Taken together, the name signals a field whose correlations *draw out* geometry from a substrate—ground made legible by measuring.

Gaia (Ground). In the archaic cosmogony of Hesiod, *Gaia* emerges at the beginning as the "ever-sure foundation," and from herself brings forth Sky, Mountains, and Sea before giving birth to the Titans. She is not merely a location but a *source*: the principle that bears, separates, and counsels as order arises—first as mother of powers, later as a voice of deep counsel preserved in cult and drama [60, 62, 61]. In this work, *Gaia* names no planet; it names the *substrate* out of which structured relations can be born.

Miti (Measure). From Sanskrit, *miti* denotes "measure," and in the classical compound *jyā-miti*—"bowstring-measure"—it becomes the art later called geometry. The image is concrete: a string stretched across an arc; curvature confronted by a straight span; a ledger of such spans turning shape into something that can be stored, compared, and used. In Indian sources, this operational emphasis runs from altar-cords to celestial arcs, and the bowstring's vocabulary travels far, eventually yielding the familiar term *sine*. Thus *miti* is not a domain but a *procedure*: to measure is to relate, to set things against a standard so that structure becomes explicit [66, 34, 67, 68].

Gaiamiti therefore reads as "measure of the ground" or "Gaia's geometry." The first component (*Gaia*) contributes the *what*: a primal substrate that can bear and differentiate relations; the second (*miti*) contributes the *how*: the operative act by which relations are rendered as geometry. In the present framework, the Gaiamiti Field's correlations serve as the conceptual "bowstrings" that span the relational arc of the substrate, making the emergent metric $\bar{g}_{\mu\nu}$ and temporal tensor $\bar{t}_{\mu\nu}$ *legible* as macroscopic structures. The name thus encodes both provenance and method—ground and grasp—without adding extra dynamical baggage beyond Definition 6.2.

Tantra: Warp, Wave, and Continuum

The Sanskrit तन्त्र (*tantra*) carries a philological density that has shaped its technical and religious afterlives. Morphologically it derives from the verbal root तन् (√*tan*, "to stretch, extend, spread, weave") conjoined with the instrumental nominal suffix -त्र (*-tra*, "that by which [an action is effected]"). The primary image is therefore practical and material: that *by which* threads are stretched and an order is laid out—the *warp* of a textile, the *loom*

as organizing device, thence the generalized sense of a *framework* or system. Classical lexicographic witnesses preserve both the concrete and derivative meanings, and already register their migration into technical prose. [66]

Within the classical semantic field, *tantra* first designates the longitudinal warp that stabilizes a fabric; from this concrete base the term acquires abstract values: the *essential part*, *model*, *rule*, *doctrine*, and finally the *systematic treatise*. This evolution is visible in the use of तन्त्रयुक्ति (*tantra-yukti*, "methods for reading a treatise") as a hermeneutical label in medical and statecraft literatures, which presupposes *tantra* in the sense of a structured manual (शास्त्र). In parallel, indigenous exegetes sometimes advance a doctrinally resonant, if philologically secondary, *nirukta*: तनोति (*tanoti*, "it expands") + त्रायते (*trāyate*, "it protects/saves"), rendering *tantra* as "that which expands [mantra and tattva] and protects [from suffering]." [66, 74, 75]

By the mid–first millennium CE the term takes on a specialized textual valence: *Tantra* becomes a designation for scriptural corpora across Śaiva, Śākta, Vaiṣṇava (Pāñcarātra), and Buddhist (Vajrayāna) traditions. The निश्वासतत्त्वसंहिता (*Niśvāsatattvasaṃhitā*)—in a multi-institution critical edition—stands as the earliest surviving Śaiva *tantra*; its introductions survey internal and external evidence that places significant strata in the fifth to sixth centuries. [76] The subsequent consolidation of Śaiva tantric forms from roughly the fifth to the thirteenth centuries is a major reconfiguration of South Asian religious culture, with tangible effects on ritual, kingship, literary canons, and inter-traditional exchange; comprehensive syntheses map these trajectories and the mechanisms of transmission. [77, 78]

On the Buddhist side, a securely datable anchor is the 724 CE Chinese translation of the *Mahāvairocana* corpus by Śubhakarasiṃha and Yìxíng, a milestone for the formation of esoteric Buddhism; later cycles such as *Cakrasaṃvara* are commonly placed in the late eighth to early ninth centuries, while the dating of *Guhyasamāja* remains a point of scholarly debate. [79, 75, 74] Historiographically, "Tantra" functions both as an emic label and a modern scholarly rubric; careful overviews therefore track where self-designation and retrospective classification align—and where they diverge. [74, 75]

Cross-linguistic renderings preserve the foundational semantics of stretching and continuity. In Tibetan Buddhist usage, *tantra* is rendered *rgyud* (gyu), "continuity/continuum," a translation-choice that foregrounds an unbroken sequence of transmission and practice and that mirrors the Sanskrit image of warp-threads drawn taut across the loom. [75]

From philology to field theory: *tantrakyma*. The Greek κῦμα (*kyma*, "wave, undulation") pairs naturally with *tantra* (framework, warp, system). The compound *tantrakyma* thus names a "woven-wave" or "warp-of-waves" or "the wave that weaves": a field in which oscillatory modes function as the very threads that generate coherence, and in which structure is an emergent weave of rhythmic articulations. In contexts where the graphic form advances the argument, the script-forms तन्त्र and κῦμα may be deployed alongside the romanized glosses.

The Kyma: The Ancient Greek Wave of Form and Flux

The word **κῦμα (kyma)** is the classical Greek term for a **wave**, particularly the swell of the sea [80]. More than a simple descriptor, the concept captured the ancient Greek fascination with dynamic forms—patterns that are constantly in motion yet retain a recognizable identity. To the Greeks, a *kyma* was not just an observation of nature but a fundamental principle

of movement, representing a surge, a swelling, or a rhythmic advance. It is a concept that embodies both tangible form and perpetual change, an idea that would ripple through philosophy, literature, and science for millennia.

Etymology and the Primordial Swell

The word κῦμα traces its origin to the Proto-Indo-European (PIE) root ***keue-**, meaning "to swell" or "to curve" [81]. This single, ancient concept of a swelling form blossomed into a vast family of words across many languages. The Greek *kyma* is a direct descendant, vividly describing a wave as a "swelling" on the surface of the water. Its linguistic cousins, or cognates, reveal this shared ancestry and include the Latin *cumulus* (a heap or pile, i.e., a swelling of earth), the Sanskrit *śū́ra* (a hero, one "swollen" with power), and even the English word *hive* (from Old English *hȳf*, a swelling structure) [81].

The Wave in Greek Thought and Literature. In ancient Greek culture, the *kyma* was a powerful and ubiquitous metaphor. In Homeric epics like the *Iliad*, armies advance not just in lines but in surging "waves" of men (κύματα φωτῶν), capturing the unstoppable, rhythmic momentum of battle. In rhetoric and drama, a *kyma* could describe a wave of emotion, applause, or public opinion sweeping through a crowd.

Philosophically, the wave became a perfect illustration for the ideas of thinkers like **Heraclitus**, who famously argued that all things are in a state of constant flux (*panta rhei*) [70]. The wave is the ideal symbol of this doctrine: it has a distinct, observable form, yet it is never static. It is a pattern of movement, a form that exists only because of change. This concept of a stable form emerging from a dynamic process is a cornerstone of ancient physics and metaphysics.

Modern Diffusion in Science and Terminology. The legacy of *kyma* is most profoundly felt today in the vocabulary of science, where its anglicized forms, **cyma-** and **kymo-**, denote concepts related to waves, curves, and swelling.

- **Cymatics:** The modern study of visible sound and vibration takes its name directly from *kyma*. In cymatics, wave patterns are made visible as particles on a vibrating surface arrange themselves into intricate, geometric forms, providing a stunning visual proof of the formative power of waves.
- **Medicine:** The **kymograph** was an early medical instrument designed to record wave-like variations in pressure, such as blood pressure or muscle contractions. Conditions like **myokymia** describe the involuntary quivering or "waving" of muscles under the skin.
- **Architecture and Botany:** In architecture, a **cyma** or *ogee* is a continuous S-shaped curve found in moulding. In botany, a **cyme** is a type of flower cluster that grows in a wave-like or branching pattern.

From a simple word for a wave on the Aegean Sea, *kyma* has evolved to describe the fundamental patterns that shape our world—from the pulse of our muscles to the vibration of sound and the elegant curves of art and nature.

The Tantrakyma Field: The Source of Internal Structure and Matter

Definition 6.3 (The Tantrakyma Field and its Modal Decomposition)**.** The *Tantrakyma Field*, $\hat{\mathbf{\Theta}}(x)$, is the second component of the Generalized Field Operator. It is a Hermitian, operator-valued field whose excitations correspond to emergent particles and whose internal symmetries correspond to conserved quantum numbers.

Like the Gaiamiti field, it admits a modal decomposition in the dialectical basis:

$$\hat{\mathbf{\Theta}}(x) = \sum_{\beta} \hat{\Theta}_{\beta}(x)\, |\beta\rangle_x$$

Each *operator-valued field mode*, $\hat{\Theta}_{\beta}(x)$, is an operator on $\mathcal{H}_{\text{em}}$ that describes the "flavor" or internal excitation state associated with the dialectical mode $|\beta\rangle_x$.

The name **tantrakyma** is formed from तन्त्र (*tantra*) and κῦμα (*kyma*). In its primary philological register, तन्त्र designates the *warp* or *framework*—"that by which" threads are stretched on a loom—and by extension a structured system or treatise; these senses are established in the classical lexica and surveyed in modern scholarship. [66, 74, 75] By contrast, κῦμα is the classical Greek word for a *wave* or swell, the natural image for oscillatory phenomena. [80] Taken together, *tantrakyma* signifies a *woven-wave framework*: a field whose wavelike excitations furnish the very threads from which internal structure is articulated.

In this reading, the Tantrakyma field provides the organizing *framework* that renders oscillations intelligible as particle content. The modal decomposition $\{\hat{\Theta}_{\beta}(x)\}$ functions as a weave of modes across the dialectical basis, within which symmetry constraints act as structural invariants of the fabric. Conserved quantum numbers arise as charges associated to these internal symmetries—enduring patterns of the weave—while localized excitations in the modes present as emergent particles. The terminology "woven-wave" thus encodes both sides of the construction: *wave* to capture the dynamical, propagating character of excitations, and *warp/framework* to capture the rule-governed architecture within which those excitations become stable carriers of quantum numbers. [66, 74, 75, 80]

Accordingly, *tantrakyma* will denote throughout the sector of the Generalized Field Operator responsible for internal structure and matter: the sector in which the interplay of modal excitation and internal symmetry realizes the observable spectrum of conserved "flavors," while remaining mathematically encoded by the same operator-theoretic decomposition written above.

The Dual Roles of the Foundational Fields

The Gaiamiti and Tantrakyma fields have distinct yet complementary offices within the unified structure of the singularity,

$$\hat{\mathbf{\Omega}}_{\vec{r}} \;=\; {}_{\hat{\mathbf{\Phi}}}\mathring{\Lambda}_{\hat{\mathbf{\Theta}}}\,.$$

They articulate two essential principles of reality: a silent potential that founds geometry, and a vibrant actual that realizes content within it.

The Gaiamiti Field ($\hat{\Phi}$): Architect of Geometry. The Gaiamiti field is the quiet architect of the cosmic stage—the primary generator of relational geometric structure. It does not create matter; rather, it establishes the rules of existence themselves: the very possibility of distance, duration, curvature, and their relational compatibilities. Its Fourfold Inter-Modal Correlator, $\hat{G}_{\Phi\Phi}$, furnishes the web of correlations from which the fabric of spacetime is woven. In this sense the Gaiamiti field is the immutable canvas: a quantum potential that endows the world with a chronogeometric arena in which processes can unfold.

The Tantrakyma Field ($\hat{\Theta}$): Source of "Flavor" and Matter. The Tantrakyma field is the dynamic actor that plays upon the stage founded by Gaiamiti. It governs the internal, non-geometric properties of reality and thus gives rise to the substance of the cosmos. Its excitation modes, encoded by the operators $\hat{\Theta}_\beta(x)$, are the woven waves threaded through the geometric fabric, and their organization by internal symmetry yields the conserved charges and "flavors" of particle-like entities within the emergent geometry. In this way the Tantrakyma field is the principle of manifestation: a woven-wave agency whose modal excitations bring the latent potential of the universe into actual, observable existence—matter, multiplets, and the spectrum of internal quantum numbers realized within the space that Gaiamiti provides.

The Kymon: The Quantum of the Creation-Wave The name **Kymon** is proposed for the fundamental quantum of the Tantrakyma field, a choice derived directly from the field's conceptual and etymological foundations. The name is a composite of **Kyma** (κῦμα), the classical Greek word for a "wave" [80], and the standard particle suffix **-on**.

This nomenclature is deliberate. As the quantum of the **Tantrakyma** field, the name *kymon* foregrounds the particle's essential nature as a *wavelike excitation*, in the standard sense of quantum field theory [82, 14]. Whereas the compound *tantrakyma* encodes a *woven–wave warp* (तन्त्र, "warp/framework," + κῦμα, "wave"), the term *kymon* isolates the realized physical event: a single quantum of the wave woven into, and constrained by, that warp. It propagates on the chronogeometric substrate determined by the Gaiamiti field and carries the charges fixed by the internal symmetries of $\hat{\boldsymbol{\Theta}}$. In short, a kymon is the fundamental ripple of manifestation—the smallest admissible instance of the field's matter–generating function.

The Chrono–Kymon: The Quantum of Temporal Torsion. We use **Chrono–Kymon**—from **Chrono-** ("time") and **Kymon** (our wave-quantum)—to name a distinctive excitation whose identity is inseparable from the time-structure of the background. Whereas an ordinary Kymon propagates *on* a prepared geometric stage, the Chrono–Kymon participates in shaping that stage's temporal texture as it moves. In physical terms, it is a *vector* (spin-1) mediator confined to the tau–kairos channel: it carries the torsional, antisymmetric aspect of temporal coherence and *does not* transmit gravitational-wave (spin-2) radiation (see Section 10.7). One may think of it as a quantized twist in the local flow of time—an elemental disturbance that both senses and adjusts how clocks, phases, and causal order knit together on the emergent background.

This framing gives the Chrono–Kymon a clear role at the matter–geometry interface. Each quantum event of this kind is a packet of temporal torsion: it communicates how material excitations tug on the timing structure of spacetime, and how that timing structure, in turn,

conditions the propagation of those excitations. By naming this coupled disturbance, we gain a concrete handle on quantum-geometric dynamics without invoking spin-2 curvature waves: the Chrono–Kymon captures the *temporal-coherence* channel of the interaction, complementing but not duplicating the radiative content of gravity. In this sense, it offers a focused pathway for probing regimes where the "when" of physics is actively negotiated—at the interface of quantum field theory and background geometry—thereby opening an experimentally and conceptually tractable window on quantum geometry itself [83, 11].

The Celestial Symphony: Gaiamiti and Tantrakyma

In the long tradition of physics—where a well–chosen name can disclose a theory's inner logic—two terms crystallize with the poise of paired themes in a single composition: *Gaiamiti* and *tantrakyma*. They are not decorative labels for algebraic machinery but compressed insights: linguistic forms that mirror physical function. If one hears a symphony in the making, *Gaiamiti* carries the ground tone, while *tantrakyma* brings the articulated line.

Gaiamiti: measure set upon ground. Etymologically fusing *Gaia* with *miti* ("measure"), *Gaiamiti* names the field that founds the stage upon which events are even possible. It is the architect of chronogeometry: the principle that selects admissible distances, durations, and curvatures, and interlocks them into a relational order. Its correlations—the interlocking patterns encoded by the Fourfold Inter-Modal Correlator $\hat{G}_{\Phi\Phi}$—are not ornaments upon spacetime but the very weave from which spacetime is realized. In this sense *Gaiamiti* is the silent potential: a ground-plan that predetermines the modes a world can sustain without itself prescribing which modes are occupied.

Tantrakyma: woven-wave articulation of content. Formed from तन्त्र (*tantra*, "warp, framework") and κῦμα (*kyma*, "wave"), *tantrakyma* names the field that renders content within the stage intelligible. It is the woven-wave framework by which internal structure becomes concrete: symmetry furnishes the warp (the rule-bound scaffold of charges and quantum numbers), while modal excitation supplies the weft (the propagating waves that realize particle-like entities). In operator language, the modes $\hat{\Theta}_\beta(x)$ are the distinct lines of melody—each a thread—while the internal symmetries bind them into multiplets and conserve their characteristic "flavors." Thus *tantrakyma* is the vibrant actual: the principle of manifestation through which latent possibility resolves into observable content.

A duet in a single work. Like two voices in a fugue, *Gaiamiti* and *tantrakyma* co-define the piece to which they belong. *Gaiamiti* furnishes the harmonic space—the admissible progressions of geometry—within which coherence can endure; *tantrakyma* traces the melodic figures—the excitations and charges—that give that space audible form. Remove the ground and melody cannot be sustained; remove the line and the ground remains unrealized potential. The singular structure $\hat{\mathbf{\Omega}}_{\vec{r}} = \overset{\circ}{\hat{\mathbf{\Phi}}\hat{\Lambda}}_{\hat{\mathbf{\Theta}}}$ is therefore not a mere coupling but a score: geometry as enabling cadence, woven waves as voiced content, the two inseparably composing the intelligible world.

6.1.1 The Inter-Modal Correlator Matrix of the Fields

The relational dynamics of the *Generalized Field Operator*, $\hat{\mathbf{F}}(x)$, are captured by a matrix-valued generalization of the Fourfold Inter-Modal Correlator (Definition Definition 4.9; cf. Section 4.1.2). This unified correlator is the central object from which all physical interactions—geometric, material, and their interplay—are derived, and its field–space/dialectic expansion matches the Chapter 4 construction in Eq. (4.76).

Definition 6.4 (The Unified Inter-Modal Correlator)**.** The *Unified Inter-Modal Correlator*, denoted $\hat{\mathbf{G}}_\Lambda(x, x')$, quantifies the quantum-relational correlations between the generalized fields at two distinct loci, x and x'. It is a 2×2 matrix of operator-valued two-point functions,

$$\hat{\mathbf{G}}_\Lambda(x, x') \;:=\; \begin{pmatrix} \hat{G}_{\Phi\Phi}(x, x') & \hat{G}_{\Phi\Theta}(x, x') \\ \hat{G}_{\Theta\Phi}(x, x') & \hat{G}_{\Theta\Theta}(x, x') \end{pmatrix},$$

whose entries are labeled by plain Roman field indices $A, B \in \{\Phi, \Theta\}$. Consistent with Chapter 4 (Eq. (4.76)), each component further expands over the Fourfold dialectic indices $\alpha, \beta \in \{0, 1, p, s\}$:

$$\hat{G}_{AB}(x, x') \;=\; \sum_{\alpha,\beta} \hat{G}^{\alpha\beta}_{AB}(x, x')\, \mathbf{e}_{\alpha\beta}, \quad A, B \in \{\Phi, \Theta\}.$$

For general modal states $|\psi_1\rangle_x = \sum_\alpha c_{1\alpha}|\alpha\rangle_x$ and $|\psi_2\rangle_{x'} = \sum_\beta c_{2\beta}|\beta\rangle_{x'}$, the components are uniformly given by

$$\boxed{\hat{G}_{AB}[\psi_1^*, \psi_2] \;:=\; \sum_{\alpha,\beta} c^*_{1\alpha} c_{2\beta} \left(\hat{A}^\dagger_\alpha(x)\, \mathcal{U}(x \leftarrow x')\, \hat{B}_\beta(x') \right)}, \qquad A, B \in \{\Phi, \Theta\},$$

where $\hat{A}_\alpha$ (resp. $\hat{B}_\beta$) denotes the appropriate field-mode operator chosen from $\{\hat{\mathbf{\Phi}}_\alpha, \hat{\mathbf{\Theta}}_\alpha\}$ (resp. $\{\hat{\mathbf{\Phi}}_\beta, \hat{\mathbf{\Theta}}_\beta\}$) according to A (resp. B). The parallel transport $\mathcal{U}(x \leftarrow x')$ is the metric-compatible propagator defined by the $u(4)$ connection on the modal bundle in Chapter 4 (see Section 4.1.4).

Equivalently, the standard specializations read:

1. *Geometric Correlator ($\hat{G}_{\Phi\Phi}$):*

$$\hat{G}_{\Phi\Phi}[\psi_1^*, \psi_2] := \sum_{\alpha,\beta} c^*_{1\alpha} c_{2\beta} \left(\hat{\mathbf{\Phi}}^\dagger_\alpha(x)\, \mathcal{U}(x \leftarrow x')\, \hat{\mathbf{\Phi}}_\beta(x') \right)$$

2. *Matter Correlator ($\hat{G}_{\Theta\Theta}$):*

$$\hat{G}_{\Theta\Theta}[\psi_1^*, \psi_2] := \sum_{\alpha,\beta} c^*_{1\alpha} c_{2\beta} \left(\hat{\mathbf{\Theta}}^\dagger_\alpha(x)\, \mathcal{U}(x \leftarrow x')\, \hat{\mathbf{\Theta}}_\beta(x') \right)$$

3. *Interaction Correlators ($\hat{G}_{\Phi\Theta}, \hat{G}_{\Theta\Phi}$):*

$$\hat{G}_{\Phi\Theta}[\psi_1^*, \psi_2] := \sum_{\alpha,\beta} c^*_{1\alpha} c_{2\beta} \left(\hat{\mathbf{\Phi}}^\dagger_\alpha(x)\, \mathcal{U}(x \leftarrow x')\, \hat{\mathbf{\Theta}}_\beta(x') \right),$$

$$\hat{G}_{\Theta\Phi}[\psi_1^*, \psi_2] := \sum_{\alpha,\beta} c^*_{1\alpha} c_{2\beta} \left(\hat{\mathbf{\Theta}}^\dagger_\alpha(x)\, \mathcal{U}(x \leftarrow x')\, \hat{\mathbf{\Phi}}_\beta(x') \right).$$

By construction, this Definition reduces componentwise to the Fourfold Inter-Modal Correlator of Chapter 4 when $A = B$ and matches the field–space organization of Eq. (4.76); see also Section 4.1.2 for the dialectic-block structure and its covariance properties.

Interpretation of the Correlator Matrix Components. Each block of the matrix $\hat{\mathbf{G}}_\Lambda$ represents a distinct physical channel of relational correlation (cf. eq. (4.76)).

Geometric Block (Component): $\hat{G}_{\Phi\Phi}(x, x')$ The upper-left diagonal component describes the self-correlation of the Gaiamiti field. It is the exclusive source of emergent geometry. Its expectation value gives rise to the emergent metric and temporal tensors, $\bar{g}_{\mu\nu}(x)$ and $\bar{t}_{\mu\nu}(x)$, thereby describing how geometric potential propagates and correlates across the continuum.

Matter Block (Component): $\hat{G}_{\Theta\Theta}(x, x')$ The lower-right diagonal component captures the self-relational structure of the Tantrakyma field. It governs the propagation and interaction of matter-like excitations, and its spectral properties determine the mass and mixing phenomenology of the emergent particles.

Interaction Blocks (Components): $\hat{G}_{\Phi\Theta}(x, x')$ and $\hat{G}_{\Theta\Phi}(x, x')$ The off-diagonal components are the *Interaction Correlators*, which encode the direct coupling between geometry and matter.

- $\hat{G}_{\Phi\Theta}$ quantifies how a matter-like excitation (a mode of $\hat{\mathbf{\Theta}}$) at x' sources a geometric deformation (a mode of $\hat{\mathbf{\Phi}}$) at x. This is the operator-level mechanism for the principle that *matter tells geometry how to curve*.
- $\hat{G}_{\Theta\Phi}$ describes the reverse process: how a geometric fluctuation at x influences the state of the matter field at x'. This corresponds to the principle that *geometry tells matter how to move*.

The non-vanishing of these off-diagonal components establishes that geometry and matter are dynamically and inseparably coupled.

The Nested Dialectic Structure. As established in the foundational definitions, each entry $\hat{G}_{AB}(x, x')$ in the 2×2 field-space matrix is itself a 4×4 matrix in the basis of the Fourfold Dialectic Algebra (cf. Section 4.1.2). The total correlator thus possesses a nested structure,

$$\hat{\mathbf{G}}_\Lambda(x, x') = \begin{pmatrix} \begin{bmatrix} \hat{G}^{\alpha\beta}_{\Phi\Phi}(x, x') \\ \hat{G}^{\alpha\beta}_{\Theta\Phi}(x, x') \end{bmatrix} & \begin{bmatrix} \hat{G}^{\alpha\beta}_{\Phi\Theta}(x, x') \\ \hat{G}^{\alpha\beta}_{\Theta\Theta}(x, x') \end{bmatrix} \end{pmatrix}, \qquad \alpha, \beta \in \{0, 1, p, s\},$$

in alignment with the Chapter 4 expansion in eq. (4.76). This architecture ensures that every interaction channel—geometric self-correlation, matter propagation, and their mutual coupling—is itself resolved into the four fundamental dialectical modes of Symmetry, Antisymmetry, Parasymmetry, and Holosymmetry.

6.1.2 Holosymmetric Wordwise Unified Inter-Modal Correlator

Alphabets, fibres, and typing (Chapter 4 dialect). Fields $\mathsf{F} = \{\Phi, \Theta\}$; Fourfold index set $\mathcal{I} = \{0, 1, p, s\}$ (channel algebra: Section 4.2). Base $\mathcal{X}_q \ni x, x', y, \ldots$. At each PQS locus $\mathfrak{B}_x$ the modal fibre is $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ with Fourfold frame $\mathcal{B}_x = \{|0\rangle_x, |1\rangle_x, |p\rangle_x, |s\rangle_x\}$ (cf. Definition 4.9 and Section 4.7). Selectors $\hat{\mathcal{F}}_\alpha(x)$ act on $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$. Parallel transport is $\mathcal{U}(x \leftarrow x'; \gamma)$ with the composition/gauge laws of eqs. (4.11), (4.14) and (4.16) (see Section 4.5). The endogenous expectation is $\langle\!\langle \cdot \rangle\!\rangle = \langle \mathring{\boldsymbol{\Psi}} | \cdot | \mathring{\boldsymbol{\Psi}} \rangle$.

Compatibility with the Generalized Field Operator. The Chapter 6 doublet $\hat{\mathbf{F}}(x) = \big(\hat{\boldsymbol{\Phi}}(x), \hat{\boldsymbol{\Theta}}(x)\big)^{\mathsf{T}}$ (cf. Definition 6.1) packages geometry/internal structure. The field-resolved selectors $\hat{\mathcal{F}}_\alpha^{(A)}(x)$ for $A \in \{\Phi, \Theta\}$ implement the Fourfold resolution of $\hat{\mathbf{A}}(x)$ on the fibre $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ (same object as Definition 4.9); they are the primitive letters used in the wordwise correlators below.

Words-dressed selectors (RR extraction). Let the holosymmetric device word and RR elements be as in Definition 4.148 and eq. (4.45). Define the readout-end dressing (cf. Definition 4.166)

$$\hat{\mathcal{F}}_\alpha^{(A)[\mathbf{W};\, \chi_{\mathrm{out}} \leftarrow \chi_{\mathrm{in}}]}(x) \;:=\; \mathsf{W}_{\chi_{\mathrm{out}} \leftarrow \chi_{\mathrm{in}}}\, \hat{\mathcal{F}}_\alpha^{(A)}(x), \qquad A \in \{\Phi, \Theta\},\ \alpha \in \mathcal{I}.$$

Definition 6.5 (HS unified inter-modal correlator — words-dressed, field-space nested)**.** Let $A, B \in \mathsf{F}$ and modal superpositions $|\psi_1\rangle_x = \sum_{\alpha \in \mathcal{I}} c_\alpha^{(1)} |\alpha\rangle_x$, $|\psi_2\rangle_{x'} = \sum_{\beta \in \mathcal{I}} c_\beta^{(2)} |\beta\rangle_{x'}$. Define the mode coefficient placed in the (A, B) field slot by

$$\hat{G}_{AB}^{\alpha\beta}\Big(x, x'; \mathbf{W}, \chi_{\mathrm{out}} \leftarrow \chi_{\mathrm{in}}\Big) := \Big(\hat{\mathcal{F}}_\alpha^{(A)[\mathbf{W};\, \chi_{\mathrm{out}} \leftarrow \chi_{\mathrm{in}}]}(x)\Big)^\dagger \mathcal{U}(x \leftarrow x'; \gamma)\, \hat{\mathcal{F}}_\beta^{(B)}(x'). \tag{6.1}$$

Assemble the Fourfold block via the basis $\{\mathbf{e}_{\alpha\beta}\}$ from Definition 4.141:

$$\hat{G}_{AB}^{[\mathbf{W};\, \chi_{\mathrm{out}} \leftarrow \chi_{\mathrm{in}}]}(x, x') := \sum_{\alpha, \beta \in \mathcal{I}} \hat{G}_{AB}^{\alpha\beta}\Big(x, x'; \mathbf{W}, \chi_{\mathrm{out}} \leftarrow \chi_{\mathrm{in}}\Big)\, \mathbf{e}_{\alpha\beta}. \tag{6.2}$$

The 2×2 field-space matrix (a "matrix of matrices") is

$$\hat{\mathbf{G}}_{\mathrm{field}}^{[\mathbf{W};\, \chi_{\mathrm{out}} \leftarrow \chi_{\mathrm{in}}]}(x, x') := \begin{pmatrix} \hat{G}_{\Phi\Phi}^{[\mathbf{W};\, \chi]}(x, x') & \hat{G}_{\Phi\Theta}^{[\mathbf{W};\, \chi]}(x, x') \\ \hat{G}_{\Theta\Phi}^{[\mathbf{W};\, \chi]}(x, x') & \hat{G}_{\Theta\Theta}^{[\mathbf{W};\, \chi]}(x, x') \end{pmatrix}, \tag{6.3}$$

which is exactly the endogenous field–space correlator of Definition 4.217.

Component/block form and wordwise context. In the Fourfold frame $\mathcal{B}_x, \mathcal{B}_{x'}$,

$$\hat{G}_{AB}^{[\mathbf{W};\, \chi]}(x, x') = [\, \hat{G}_{AB}^{\alpha\beta}(\cdots)\,]_{\alpha, \beta \in \mathcal{I}}$$

with entries given by (6.1). If wordwise frames/trivializations are used, see Section 4.6.2; kinematics of the transport enter via Section 4.5.

Proposition 6.6 (Core properties (Chapter 4 formalism)). *For* $\hat{\mathbf{G}}_{\text{field}}^{[\mathbf{W};\chi]}(x,x')$ *in* (6.3)*:*

Gauge covariance

By Propositions 4.54 and 4.167 and Corollary 4.64, each block transforms as $\hat{G}_{AB}^{[\mathbf{W};\chi]}(x,x') \mapsto g(x)\,\hat{G}_{AB}^{[\mathbf{W};\chi]}(x,x')\,g(x')^{\dagger}$.

Endogenous grading and centralization

Holosymmetric charges $q(\cdot)$ *live in the device/dialectic factor and are field-label independent; thus the componentwise centrality and compliance-vector bounds from Section 4.2.1 and Section 4.4 apply blockwise to* $\hat{\mathbf{G}}_{\text{field}}^{[\mathbf{W};\chi]}$.

Functoriality/composition

For $x \leftarrow y \leftarrow x'$, $\mathcal{U}(x \leftarrow x') = \mathcal{U}(x \leftarrow y)\mathcal{U}(y \leftarrow x')$ *implies the block composition* $\hat{\mathbf{G}}_{AB}^{[\mathbf{W};\chi]}(x,x') = \sum_{\gamma\in\mathcal{I}} \hat{\mathbf{G}}_{A\gamma}^{[\mathbf{W};\chi]}(x,y)\,\hat{\mathbf{G}}_{\gamma B}^{[\mathbf{W};\chi]}(y,x')$ *(cf. Section 4.5).*

Reduction to the undressed form

For $\mathbf{W} = I$ *and* $\chi_{\text{out}} = \chi_{\text{in}}$, (6.3) *reduces to the undressed field–space correlator* $\hat{\mathbf{G}}_{\text{field}}(x,x')$ *of* (4.82).

Expectation reduction (operator arena). Taking $\langle\!\langle\cdot\rangle\!\rangle$ yields $\langle\!\langle \hat{\mathbf{G}}_{\text{field}}^{[\mathbf{W};\chi]}(x,x')\rangle\!\rangle = \hat{\mathbf{G}}_{\text{field}}(x,x')$ with Fourfold expansion as in Definition 4.9; bundle/frames and global sections are as in Sections 4.6.2 and 4.6.5.

Canonical specialisations.

$$\hat{G}_{\Phi\Phi}^{\alpha\beta}(x,x') = \Big(\hat{\mathcal{F}}_{\alpha}^{(\Phi)}(x)\Big)^{\dagger}\,\mathcal{U}(x\leftarrow x')\,\hat{\mathcal{F}}_{\beta}^{(\Phi)}(x'),$$

$$\hat{G}_{\Theta\Theta}^{\alpha\beta}(x,x') = \Big(\hat{\mathcal{F}}_{\alpha}^{(\Theta)}(x)\Big)^{\dagger}\,\mathcal{U}(x\leftarrow x')\,\hat{\mathcal{F}}_{\beta}^{(\Theta)}(x'),$$

$$\hat{G}_{\Phi\Theta}^{\alpha\beta}(x,x') = \Big(\hat{\mathcal{F}}_{\alpha}^{(\Phi)}(x)\Big)^{\dagger}\,\mathcal{U}(x\leftarrow x')\,\hat{\mathcal{F}}_{\beta}^{(\Theta)}(x'),$$

$$\hat{G}_{\Theta\Phi}^{\alpha\beta}(x,x') = \Big(\hat{\mathcal{F}}_{\alpha}^{(\Theta)}(x)\Big)^{\dagger}\,\mathcal{U}(x\leftarrow x')\,\hat{\mathcal{F}}_{\beta}^{(\Phi)}(x').$$

Here, each displayed line is a *field-resolved Fourfold matrix element* of the unified two-point kernel: $\alpha,\beta \in \mathcal{I} = \{0,1,p,s\}$ label the dialectic/Fourfold basis at the anchors x and x' (see Definition 4.141); $\hat{\mathcal{F}}_{\alpha}^{(A)}(\cdot)$ are the *field-labelled selectors* acting on the emergent modal fibre (as in Definition 4.9 and assembled fieldwise in Definition 4.217); and $\mathcal{U}(x\leftarrow x')$ is the gauge-covariant *parallel transporter* along the chosen path γ (with composition and gauge laws in eqs. (4.11), (4.14) and (4.16); see also Section 4.5). The *diagonal entries* $\hat{G}_{\Phi\Phi}^{\alpha\beta}$ and $\hat{G}_{\Theta\Theta}^{\alpha\beta}$ quantify, respectively, the propagation/correlation of purely geometric (Gaiamiti) and purely internal–structural (Tantrakyma) modes between (x',β) and (x,α). The *mixed entries* $\hat{G}_{\Phi\Theta}^{\alpha\beta}$ and $\hat{G}_{\Theta\Phi}^{\alpha\beta}$ capture *inter-sector couplings*—how a Tantrakyma excitation at (x',β) is "read out" through a Gaiamiti selector at (x,α), and vice versa—making the geometry–matter co-dependence explicit at the level of the kernel. These components assemble into the field–space blocks $\hat{\mathbf{G}}_{AB}(x,x') = \sum_{\alpha,\beta}\hat{G}_{AB}^{\alpha\beta}(x,x')\,\mathbf{e}_{\alpha\beta}$ and then into the 2×2 field matrix

$\big[\hat{\mathbf{G}}_{AB}\big]_{A,B\in\{\Phi,\Theta\}}$ of Definition 4.217. By construction they inherit the *gauge covariance* $\hat{G}_{AB}(x,x') \mapsto g(x)\,\hat{G}_{AB}(x,x')\,g(x')^{\dagger}$, *Hermitian reciprocity* with anchor swap, and *block-composition along paths* (all as in Propositions 4.54 and 4.167, Corollary 4.64, and Section 4.5); together, these ensure that the canonical specialisations are the precise field-resolved building blocks of the unified correlator used throughout Chapters 4–6.

Optional: spectral localisation. If spectral cutoffs on the fibres are in force (as used for operator-valued tensors), with $P_x = E^{(x)}([0,\lambda_{\mathrm{cut}}])$ and $P_{x'}$ the projectors at the anchors, then

$$\hat{\mathbf{G}}^{[\mathbf{W};\chi]}_{\mathrm{field}}(x,x') = P_x\,\hat{\mathbf{G}}^{[\mathbf{W};\chi]}_{\mathrm{field}}(x,x')\,P_{x'}, \qquad \operatorname{Ran}\hat{\mathbf{G}}^{[\mathbf{W};\chi]}_{\mathrm{field}}(x,x') \subseteq \operatorname{Ran} P_x,$$

in the same sense used for the metric/temporal constructions (cf. cross-chapter usage in Section 4.5). *Explanation and context.* The optional projector sandwich above implements the same *spectral locality* used for operator–valued metric/temporal tensors: each fibre carries a self–adjoint generator with spectral measure $E^{(x)}(\cdot)$, and we restrict to the low–energy band $[0,\lambda_{\mathrm{cut}}]$ via $P_x := E^{(x)}([0,\lambda_{\mathrm{cut}}])$ (cf. Definition 5.1 and Proposition 5.5). Applied to the words–dressed field–space correlator of Definition 4.217, the relations

$$\hat{\mathbf{G}}^{[\mathbf{W};\chi]}_{\mathrm{field}}(x,x') = P_x\,\hat{\mathbf{G}}^{[\mathbf{W};\chi]}_{\mathrm{field}}(x,x')\,P_{x'}, \qquad \operatorname{Ran}\hat{\mathbf{G}}^{[\mathbf{W};\chi]}_{\mathrm{field}}(x,x') \subseteq \operatorname{Ran} P_x$$

ensure that (i) the kernel acts within the spectrally regulated fibres at the two anchors, (ii) SOT–uniform bounds and differentiability on the restricted ranges are available for base–channel derivatives, and (iii) the composition laws with the transporter $\mathcal{U}(x \leftarrow x')$ remain well–posed (see the path-ordered, composition, and gauge rules in eqs. (4.11), (4.14) and (4.16) and the kinematic discussion in Section 4.5). Because $P_x \mapsto g(x)\,P_x\,g(x)^{\dagger}$ under local frame changes, the projector sandwich is compatible with gauge covariance of the blocks (cf. Propositions 4.54 and 4.167 and Corollary 4.64). Positivity of diagonal kernels is preserved by compression, and the construction reduces to the unrestricted case in the formal limit $\lambda_{\mathrm{cut}} \to \infty$. In short, spectral localisation provides a controlled, gauge–consistent truncation of $\hat{\mathbf{G}}^{[\mathbf{W};\chi]}_{\mathrm{field}}$ that matches the regularity regime already assumed for the metric/temporal operator fields, enabling uniform manipulation of derivatives, expectations, and path composition across Chapters 4–6.

Optional: law-controlled HS lift (if a law register is active). Let $\mathsf{L} = \operatorname{span}\{|0\rangle_L, |1\rangle_L\}$. We assemble a branch-resolved lift of the field–space correlator (same type as $\hat{\mathbf{G}}_{\mathrm{field}}(x,x')$ in Definition 4.217):

$$\widehat{\mathbf{G}}^{[L]}(x,x') := \sum_{i,j\in\{0,1\}} \big(|i\rangle_L\langle j|_L\big) \otimes \hat{\mathbf{G}}^{(ij)}(x,x')\,. \tag{6.4}$$

Equivalent compact form (explicit blocks).

$$\begin{aligned}\widehat{\mathbf{G}}^{[L]}(x,x') =& |0\rangle_L\langle 0|_L \otimes \hat{\mathbf{G}}^{(00)}(x,x') \\ &+ |1\rangle_L\langle 1|_L \otimes \hat{\mathbf{G}}^{(11)}(x,x') \\ &+ |0\rangle_L\langle 1|_L \otimes \hat{\mathbf{G}}^{(01)}(x,x') \\ &+ |1\rangle_L\langle 0|_L \otimes \hat{\mathbf{G}}^{(10)}(x,x')\,.\end{aligned} \tag{6.5}$$

The diagonal blocks are the branch kernels

$$\hat{\mathbf{G}}^{(00)}(x,x') \equiv \hat{\mathbf{G}}^{(0)}(x,x'), \qquad \hat{\mathbf{G}}^{(11)}(x,x') \equiv \hat{\mathbf{G}}^{(1)}(x,x')\,,$$

and the off–diagonal (coherence) blocks $\hat{\mathbf{G}}^{(01)}(x,x')$, $\hat{\mathbf{G}}^{(10)}(x,x')$ are defined in the same Fourfold/wordwise grammar as Definition 4.217, but with branch–conditioned word dressing. Concretely, for $A,B \in \{\Phi,\Theta\}$ and $\alpha,\beta \in \mathcal{I}$,

$$\hat{G}^{(ij);\alpha\beta}_{AB}(x,x') \;:=\; \Big(\hat{\mathcal{F}}^{(A)[\mathbf{W}^{(i)};\chi]}_{\alpha}(x)\Big)^{\dagger}\,\mathcal{U}(x\leftarrow x';\gamma)\,\hat{\mathcal{F}}^{(B)[\mathbf{W}^{(j)};\chi]}_{\beta}(x')\,, \tag{6.6}$$

$$\hat{\mathbf{G}}^{(ij)}_{AB}(x,x') \;:=\; \sum_{\alpha,\beta\in\mathcal{I}} \hat{G}^{(ij);\alpha\beta}_{AB}(x,x')\;\mathbf{e}_{\alpha\beta}\,, \qquad \hat{\mathbf{G}}^{(ij)}(x,x') \;:=\; \Big[\hat{\mathbf{G}}^{(ij)}_{AB}(x,x')\Big]_{A,B\in\{\Phi,\Theta\}}. \tag{6.7}$$

Here $\mathbf{W}^{(0)},\mathbf{W}^{(1)}$ are the branch–conditioned word devices consistent with Definitions 4.148 and 4.166, and $\mathbf{e}_{\alpha\beta}$ is the Fourfold basis from Definition 4.141. Transport $\mathcal{U}$ and all gauge/-typing rules are as in Section 4.5 and eqs. (4.11), (4.14) and (4.16).

Control–state reduction. For a prepared control state $|+_\varphi\rangle_L = \frac{1}{\sqrt{2}}\big(|0\rangle_L + e^{\mathrm{i}\varphi}|1\rangle_L\big)$, the reduced (law–averaged) correlator is

$$\langle +_\varphi|_L\,\widehat{\mathbf{G}}^{[L]}(x,x')\,|+_\varphi\rangle_L = \tfrac{1}{2}\Big(\hat{\mathbf{G}}^{(0)} + \hat{\mathbf{G}}^{(1)} + e^{-\mathrm{i}\varphi}\hat{\mathbf{G}}^{(01)} + e^{\mathrm{i}\varphi}\hat{\mathbf{G}}^{(10)}\Big)(x,x')\,, \tag{6.8}$$

exhibiting phase–programmable symmetric/antisymmetric mixing of the (0) and (1) branches without introducing any new mixing operator symbol.

Intuition. The *law-controlled HS lift* augments the correlator with a two-level control system—the law register $\mathsf{L} = \mathrm{span}\{|0\rangle_L,|1\rangle_L\}$—so that two alternative branches (e.g., two word devices, parameter regimes, or "laws") are represented *simultaneously*. Preparing L in a superposition enables coherent mixing of the branches with a programmable phase, so the lifted object captures both diagonal (pure-branch) contributions and off-diagonal coherences, while preserving the standard wordwise grammar of selectors, transport, and Fourfold bookkeeping (cf. Definitions 4.148, 4.166 and 4.217).

Construction. We define the lifted correlator as

$$\widehat{\mathbf{G}}^{[L]}(x,x') \;:=\; \sum_{i,j\in\{0,1\}} \Big(|i\rangle_L\langle j|_L\Big) \;\otimes\; \hat{\mathbf{G}}^{(ij)}(x,x')\,, \tag{6.9}$$

where each block $\hat{\mathbf{G}}^{(ij)}(x,x')$ is a field–space matrix (each field slot is Fourfold 4×4) built in the same wordwise manner as the endogenous field–space correlator of Definition 4.217. Concretely, for $A,B \in \{\Phi,\Theta\}$ and $\alpha,\beta \in \mathcal{I}$,

$$\hat{G}^{(ij);\alpha\beta}_{AB}(x,x') \;:=\; \Big(\hat{\mathcal{F}}^{(A)[\mathbf{W}^{(i)};\chi]}_{\alpha}(x)\Big)^{\dagger}\,\mathcal{U}(x\leftarrow x';\gamma)\,\hat{\mathcal{F}}^{(B)[\mathbf{W}^{(j)};\chi]}_{\beta}(x')\,, \tag{6.10}$$

and we assemble

$$\hat{\mathbf{G}}^{(ij)}_{AB}(x,x') \;:=\; \sum_{\alpha,\beta\in\mathcal{I}} \hat{G}^{(ij);\alpha\beta}_{AB}(x,x')\,\mathbf{e}_{\alpha\beta}\,, \qquad \hat{\mathbf{G}}^{(ij)}(x,x') \;:=\; \Big[\hat{\mathbf{G}}^{(ij)}_{AB}(x,x')\Big]_{A,B\in\{\Phi,\Theta\}}\,. \tag{6.11}$$

Here $\mathbf{W}^{(0)}$ and $\mathbf{W}^{(1)}$ are the branch–conditioned word devices (cf. Definitions 4.148 and 4.166), $\mathbf{e}_{\alpha\beta}$ is the Fourfold basis (cf. Definition 4.141), and $\mathcal{U}$ is the transporter with the path-ordered, composition, and gauge laws of Section 4.5 and eqs. (4.11), (4.14) and (4.16)).

Semantics of the blocks. The diagonal blocks $\hat{\mathbf{G}}^{(00)} \equiv \hat{\mathbf{G}}^{(0)}$ and $\hat{\mathbf{G}}^{(11)} \equiv \hat{\mathbf{G}}^{(1)}$ are the pure-branch correlators using $\mathbf{W}^{(0)}$ or $\mathbf{W}^{(1)}$ at both endpoints. The off-diagonal blocks encode coherences: $\hat{\mathbf{G}}^{(01)}$ applies $\mathbf{W}^{(0)}$ at x and $\mathbf{W}^{(1)}$ at x', while $\hat{\mathbf{G}}^{(10)}$ swaps them. All blocks share the same anchoring and Fourfold bookkeeping as in Definition 4.217.

Control-state reduction. For a phase-programmable control state $|+_\varphi\rangle_L = \frac{1}{\sqrt{2}}\big(|0\rangle_L + e^{\mathrm{i}\varphi}|1\rangle_L\big)$,

$$\langle +_\varphi|_L\, \widehat{\mathbf{G}}^{[L]}(x,x')\,|\,+_\varphi\rangle_L \;=\; \tfrac{1}{2}\Big(\hat{\mathbf{G}}^{(0)} + \hat{\mathbf{G}}^{(1)} + e^{-\mathrm{i}\varphi}\hat{\mathbf{G}}^{(01)} + e^{\mathrm{i}\varphi}\hat{\mathbf{G}}^{(10)}\Big)(x,x')\,. \tag{6.12}$$

Thus $\varphi = 0$ yields constructive interference of the coherences, $\varphi = \pi$ yields destructive interference, and complete dephasing of L removes the off-diagonals, producing the classical mixture $\frac{1}{2}\big(\hat{\mathbf{G}}^{(0)} + \hat{\mathbf{G}}^{(1)}\big)$. More generally, for any $\rho_L = \left(\begin{smallmatrix} a & c \\ c^* & b \end{smallmatrix}\right)$ on L,

$$\mathrm{Tr}_L\Big[(\rho_L \otimes I)\,\widehat{\mathbf{G}}^{[L]}(x,x')\Big] \;=\; a\,\hat{\mathbf{G}}^{(0)} + b\,\hat{\mathbf{G}}^{(1)} + c\,\hat{\mathbf{G}}^{(10)} + c^*\,\hat{\mathbf{G}}^{(01)}\,. \tag{6.13}$$

Inherited structural properties. Gauge covariance follows blockwise from the dressing and transport laws (cf. Propositions 4.54 and 4.167 and Corollary 4.64):

$$\hat{\mathbf{G}}^{(ij)}(x,x') \;\longmapsto\; g(x)\,\hat{\mathbf{G}}^{(ij)}(x,x')\,g(x')^\dagger, \qquad (\,|i\rangle\langle j|\,)_L \text{ unchanged.} \tag{6.14}$$

Hermitian reciprocity holds with anchor swap,

$$\Big(\hat{\mathbf{G}}^{(ij)}(x,x')\Big)^\dagger \;=\; \hat{\mathbf{G}}^{(ji)}(x',x)\,. \tag{6.15}$$

Functoriality along paths is preserved: for $x \leftarrow y \leftarrow x'$,

$$\hat{\mathbf{G}}^{(ij)}(x,x') \;=\; \sum_{\gamma\in\mathcal{I}} \hat{\mathbf{G}}^{(i\gamma)}(x,y)\,\hat{\mathbf{G}}^{(\gamma j)}(y,x')\,, \tag{6.16}$$

in the same sense as the field–space composition (cf. Section 4.5). The diagonal kernels retain the usual positivity; off-diagonal blocks satisfy Cauchy–Schwarz-type bounds relative to the diagonals, so dephasing is completely positive. Applying $\langle\!\langle\cdot\rangle\!\rangle$ after choosing ρ_L yields the operator-level correlator with the same Fourfold expansion and bundle semantics (cf. Definition 4.9 and Sections 4.6.2 and 4.6.5).

Circuit-style specification (branch-controlled dressing). Introduce a two-level control (law) register $\mathsf{L} = \mathrm{span}\{|0\rangle_L, |1\rangle_L\}$. Define the *controlled dressing maps* that act on the selector dressing at the two anchors:

$$\begin{aligned} \mathfrak{C}^{[L]}_{\mathrm{dress},x} &:= |0\rangle_L\langle 0|_L \otimes \mathsf{D}^{(0)}_x \;+\; |1\rangle_L\langle 1|_L \otimes \mathsf{D}^{(1)}_x, \\ \mathfrak{C}^{[L]}_{\mathrm{dress},x'} &:= |0\rangle_L\langle 0|_L \otimes \mathsf{D}^{(0)}_{x'} \;+\; |1\rangle_L\langle 1|_L \otimes \mathsf{D}^{(1)}_{x'}. \end{aligned} \tag{6.17}$$

where each $\mathsf{D}^{(i)}$ implements the usual words-dressed substitution

$$\mathsf{D}^{(i)}_x : \hat{\mathcal{F}}^{(A)}_\alpha(x) \;\mapsto\; \hat{\mathcal{F}}^{(A)[\mathbf{W}^{(i)};\chi]}_\alpha(x), \qquad \mathsf{D}^{(j)}_{x'} : \hat{\mathcal{F}}^{(B)}_\beta(x') \;\mapsto\; \hat{\mathcal{F}}^{(B)[\mathbf{W}^{(j)};\chi]}_\beta(x'),$$

with $\mathbf{W}^{(0)}, \mathbf{W}^{(1)}$ the branch-conditioned word devices (cf. Definitions 4.148 and 4.166). Let the *undressed* field–space correlator be

$$\begin{aligned} \hat{\mathbf{G}}^{\mathrm{id}}(x,x') &:= \left[\hat{\mathbf{G}}^{\mathrm{id}}_{AB}(x,x')\right]_{A,B\in\{\Phi,\Theta\}}, \\ \hat{\mathbf{G}}^{\mathrm{id}}_{AB}(x,x') &:= \sum_{\alpha,\beta\in\mathcal{I}} \left(\hat{\mathcal{F}}^{(A)}_\alpha(x)\right)^\dagger \mathcal{U}(x \leftarrow x';\gamma)\, \hat{\mathcal{F}}^{(B)}_\beta(x')\, \mathbf{e}_{\alpha\beta}. \end{aligned} \tag{6.18}$$

built exactly as in Definition 4.217 (selectors $\hat{\mathcal{F}}_\alpha$, transporter $\mathcal{U}$, Fourfold basis $\mathbf{e}_{\alpha\beta}$ from Definition 4.141, and kinematics Section 4.5 and eqs. (4.11), (4.14) and (4.16)).

Lift as a controlled circuit. The law-controlled lift is obtained by *sandwiching* the undressed kernel with the controlled dressings, and keeping the full dyadic on L:

$$\widehat{\mathbf{G}}^{[L]}(x,x') = \sum_{i,j\in\{0,1\}} \left(|i\rangle_L\langle j|_L\right) \;\otimes\; \left(\mathsf{D}^{(i)}_x \triangleright \hat{\mathbf{G}}^{\mathrm{id}}(x,x') \triangleleft \mathsf{D}^{(j)}_{x'}\right), \tag{6.19}$$

i.e.

$$\begin{aligned} \hat{\mathbf{G}}^{(ij)}(x,x') &:= \left[\,\hat{\mathbf{G}}^{(ij)}_{AB}(x,x')\,\right]_{A,B\in\{\Phi,\Theta\}}, \\ \hat{\mathbf{G}}^{(ij)}_{AB}(x,x') &:= \sum_{\alpha,\beta\in\mathcal{I}} \left(\hat{\mathcal{F}}^{(A)[\mathbf{W}^{(i)};\chi]}_\alpha(x)\right)^\dagger \mathcal{U}(x \leftarrow x';\gamma)\, \hat{\mathcal{F}}^{(B)[\mathbf{W}^{(j)};\chi]}_\beta(x')\, \mathbf{e}_{\alpha\beta}. \end{aligned} \tag{6.20}$$

which is exactly the branch-resolved definition used in Definition 4.217 (with branch-conditioned dressing).

Programmable preparation and reduction. Prepare the law register by the usual single-qubit circuit H followed by $R_z(\varphi)$ to create $|+_\varphi\rangle_L = \frac{1}{\sqrt{2}}\left(|0\rangle_L + e^{\mathrm{i}\varphi}|1\rangle_L\right)$. Tracing out L yields

$$\langle +_\varphi|_L\, \widehat{\mathbf{G}}^{[L]}(x,x')\, |+_\varphi\rangle_L = \tfrac{1}{2}\Big(\hat{\mathbf{G}}^{(0)} + \hat{\mathbf{G}}^{(1)} + e^{-\mathrm{i}\varphi}\hat{\mathbf{G}}^{(01)} + e^{\mathrm{i}\varphi}\hat{\mathbf{G}}^{(10)}\Big)(x,x'), \tag{6.21}$$

so φ sets constructive ($\varphi = 0$) or destructive ($\varphi = \pi$) interference of the coherences; dephasing the control removes the off-diagonals, giving the classical mixture $\frac{1}{2}(\hat{\mathbf{G}}^{(0)} + \hat{\mathbf{G}}^{(1)})$.

Operational meaning of the circuit (branch-controlled dressing). Consider the circuit in which the law register L (top wire) controls branch-conditioned dressings at the two anchors x and x':



Here the controlled boxes $\mathsf{D}^{(0/1)}$ indicate the branch-conditioned dressing at each anchor (left-multiplying the selectors as in Definition 4.166); the propagation between the two anchors is governed by $\mathcal{U}(x \leftarrow x')$ (cf. Section 4.5).

1. **Preparation.** The gates H and $R_z(\varphi)$ prepare the law qubit in
$$|+_\varphi\rangle_L = \tfrac{1}{\sqrt{2}}\big(|0\rangle_L + e^{i\varphi}|1\rangle_L\big).$$

2. **Branch-conditioned dressing (multiplexed control).** The controlled boxes implement the *selected* dressings at each anchor (both branches can be nontrivial), via the dyadic control maps of eq. (6.17):
$$\mathfrak{C}^{[L]}_{\mathrm{dress},x} = |0\rangle_L\langle 0|_L \otimes \mathsf{D}_x^{(0)} \;+\; |1\rangle_L\langle 1|_L \otimes \mathsf{D}_x^{(1)}, \quad \mathfrak{C}^{[L]}_{\mathrm{dress},x'} = |0\rangle_L\langle 0|_L \otimes \mathsf{D}_{x'}^{(0)} \;+\; |1\rangle_L\langle 1|_L \otimes \mathsf{D}_{x'}^{(1)},$$
where each $\mathsf{D}^{(i)}$ acts by the standard words-dressed substitution on selectors (cf. Definition 4.166):
$$\mathsf{D}_x^{(i)} : \; \hat{\mathcal{F}}_\alpha^{(A)}(x) \mapsto \hat{\mathcal{F}}_\alpha^{(A)[\mathbf{W}^{(i)};\chi]}(x), \qquad \mathsf{D}_{x'}^{(j)} : \; \hat{\mathcal{F}}_\beta^{(B)}(x') \mapsto \hat{\mathcal{F}}_\beta^{(B)[\mathbf{W}^{(j)};\chi]}(x').$$
Propagation between anchors is unchanged and governed by the transporter $\mathcal{U}(x \leftarrow x';\gamma)$ with the path-ordered, composition, and gauge laws of Section 4.5 and eqs. (4.11), (4.14) and (4.16).

3. **Circuit-level object and its blocks.** Applying the controlled dressings to the undressed field–space kernel of Definition 4.217 and eq. (6.18) yields the branch-resolved blocks
$$\hat{\mathbf{G}}^{(ij)}(x,x') = \big[\hat{\mathbf{G}}^{(ij)}_{AB}(x,x')\big]_{A,B\in\{\Phi,\Theta\}},$$
$$\hat{\mathbf{G}}^{(ij)}_{AB}(x,x') = \sum_{\alpha,\beta\in\mathcal{I}} \Big(\hat{\mathcal{F}}_\alpha^{(A)[\mathbf{W}^{(i)};\chi]}(x)\Big)^\dagger \mathcal{U}(x \leftarrow x';\gamma)\, \hat{\mathcal{F}}_\beta^{(B)[\mathbf{W}^{(j)};\chi]}(x')\, \mathbf{e}_{\alpha\beta}.$$
as in eqs. (6.10), (6.11) and (6.20). Keeping the full dyadic on L constructs the law-controlled lift
$$\widehat{\mathbf{G}}^{[L]}(x,x') = \sum_{i,j\in\{0,1\}} |i\rangle_L\langle j|_L \otimes \hat{\mathbf{G}}^{(ij)}(x,x'),$$
which is exactly the branch-resolved form of eqs. (6.4) and (6.19).

4. **Interference readout / reduction.** Preparing $|+_\varphi\rangle_L$ and reducing the law register gives the phase-programmable mixture of eq. (6.12):

$$\langle +_\varphi|_L\, \widehat{\mathbf{G}}^{[L]}(x,x')\,|+_\varphi\rangle_L = \tfrac{1}{2}\Big(\hat{\mathbf{G}}^{(0)} + \hat{\mathbf{G}}^{(1)} + e^{-i\varphi}\hat{\mathbf{G}}^{(01)} + e^{i\varphi}\hat{\mathbf{G}}^{(10)}\Big)(x,x').$$

   Thus scanning φ isolates the *coherence* blocks $\hat{\mathbf{G}}^{(01)}$, $\hat{\mathbf{G}}^{(10)}$ in addition to the diagonals $\hat{\mathbf{G}}^{(0)}$, $\hat{\mathbf{G}}^{(1)}$. The special "Hadamard-test" case arises by choosing $\mathsf{D}^{(0)} = I$ and $\mathsf{D}^{(1)} = \mathsf{D}$.

5. **Covariance and consistency.** All blocks inherit gauge covariance, reciprocity, and path functoriality (cf. Propositions 4.54 and 4.167, Corollary 4.64, and Section 4.5); the law dyads $|i\rangle_L\langle j|_L$ are unaffected by gauge. The construction remains in the same Fourfold/field-space grammar as Definition 4.217, so it integrates with the rest of the Chapter 4–6 machinery without introducing new symbols.

Properties (inherit blockwise). Gauge covariance, reciprocity, and path functoriality hold blockwise for $\hat{\mathbf{G}}^{(ij)}$ exactly as in the undressed case (cf. Propositions 4.54 and 4.167 and Corollary 4.64); applying $\langle\!\langle\cdot\rangle\!\rangle$ after choosing the control state maps the circuit-level object into the operator arena with the same Fourfold expansion and bundle semantics (cf. Definition 4.9 and Sections 4.6.2 and 4.6.5).

Meaning of "law" in the holosymmetric programme (Chapter 3). In Chapter 3, holosymmetry is introduced as an *operational* principle that allows one to place *alternative dynamical prescriptions* into coherent superposition and to read out their interference. A *law* is therefore a branch–indexed rule set (a programmable *dynamical grammar*) that fixes which word devices, selectors, couplings, and transport rules are used to prepare, dress, propagate, and read out modal content. Holosymmetric interferometers implement interference *between* such law branches, not merely between spatial paths; a dedicated control degree of freedom toggles the branches and enables coherent comparison (see the interferometric constructions in Sections 3.1, 3.1.1 and 3.1.5). In this sense, "law" is a typed label for branch–conditioned HS constructions, and holosymmetry supplies the covariance and bookkeeping needed to compare them within a single operational experiment.

Holosymmetric lift and branchwise geometry (Chapter 5). Chapter 5 formalizes "law" by introducing an explicit two–level *law register* $\mathsf{L} = \mathrm{span}\{|0\rangle_L, |1\rangle_L\}$ and building branch–resolved HS objects with branch–conditioned word devices $\mathbf{W}^{(0)}$ and $\mathbf{W}^{(1)}$. The lifted kernels and geometric words decompose into diagonal blocks (pure–branch objects) and off–diagonal blocks (coherences between laws), and they reduce to a chosen mixture once L is prepared and measured. In expectation, each branch carries its own causal/chronogeometric structure (effective cones; see Definition 5.123), and operational causality is preserved: coherent control over laws does not enable signalling beyond the allowed cones (Proposition 5.152). The branch–resolved lift used in Chapter 6 for the unified correlator, $\widehat{\mathbf{G}}^{[L]}(x,x')$ with blocks $\hat{\mathbf{G}}^{(ij)}(x,x')$, is precisely this mechanism instantiated for field–space kernels (cf. eqs. (6.4) and (6.9) to (6.11)).

Mapping to the circuit in Chapter 6. In the circuit, the ancilla wire L *is* the law register. The controlled boxes at the x and x' anchors implement branch–conditioned dressing, applying $\mathbf{W}^{(0)}$ or $\mathbf{W}^{(1)}$ to the selector systems according to the state of L (cf. Definition 4.166). Preparing $|+_\varphi\rangle_L$ produces a coherent superposition of law branches; the ensuing interference on L probes the *coherences* between the branch–resolved blocks of the lifted correlator. Tracing out L yields the phase–programmable combination of blocks in eq. (6.12), while all covariance and path–composition rules are inherited from the transporter kinematics and gauge laws (Section 4.5 and eqs. (4.11), (4.14) and (4.16)). Conceptually, "law" here is not a new operator on the modal fibre but a control label that selects the branch–conditioned word device; holosymmetry guarantees consistent comparison of these selections, and Proposition 5.152 ensures that such comparisons remain fully compatible with operational causality.

6.1.3 The Relational Levi-Civita Derivative

To describe the dynamics of fields and the motion of particles on the emergent chronogeometric manifold, we require a notion of parallel transport. This is defined by a connection, which in turn determines the "straightest possible paths," or geodesics. In the IQR framework, the connection cannot be arbitrarily postulated; it must be intrinsically determined by the very fabric of emergent reality.

The Principle of Chronogeometric Covariance

The action functionals for both the Gaiamiti field (see Definition 6.118) and the Tantrakyma field (see Definition 6.142) are constructed to be invariant under general coordinate transformations on the manifold endowed with the *Chronogeometric Tensor*, $\bar{\Sigma}_{\mu\nu}(x)$. The principle of self-consistent dynamics therefore demands that the connection governing parallel transport be the unique affine structure that is compatible with this unified tensor. To derive it from the metric tensor $\bar{g}_{\mu\nu}$ alone would be to neglect the temporal flow structure $\bar{t}_{\mu\nu}$, to which the fundamental fields are explicitly coupled.

This leads us to define the *Relational Levi-Civita Derivative*, the unique torsion-free, metric-compatible connection associated with $\bar{\Sigma}_{\mu\nu}$.

Definition 6.7 (Relational Levi-Civita Derivative)**.** Let $\bar{\Sigma}_{\mu\nu}(x)$ be the smooth, symmetric, real-valued chronogeometric tensor field defined in Definition 5.26. The *Relational Levi-Civita Derivative*, denoted $\nabla^{(\mathrm{rel})}_\mu$, is the unique affine connection that satisfies:

1. **Torsion-free:** $\Gamma^\lambda_{\mu\nu} = \Gamma^\lambda_{\nu\mu}$,
2. **Metric compatibility:** $\nabla^{(\mathrm{rel})}_\lambda \bar{\Sigma}_{\mu\nu} = 0$.

Its Christoffel symbols are given by:

$$\boxed{\Gamma^\lambda_{\mu\nu}[\bar{\Sigma}] := \tfrac{1}{2}\bar{\Sigma}^{\lambda\rho}\left(\partial_\mu \bar{\Sigma}_{\nu\rho} + \partial_\nu \bar{\Sigma}_{\mu\rho} - \partial_\rho \bar{\Sigma}_{\mu\nu}\right),} \tag{6.22}$$

and the derivative acts on tensor fields via:

$$\nabla^{(\mathrm{rel})}_\mu V^\nu := \partial_\mu V^\nu + \Gamma^\nu_{\mu\lambda}[\bar{\Sigma}]V^\lambda, \quad \nabla^{(\mathrm{rel})}_\mu V_\nu := \partial_\mu V_\nu - \Gamma^\lambda_{\mu\nu}[\bar{\Sigma}]V_\lambda.$$

This connection determines both the geodesics and the curvature structure of the full relational geometry experienced by quantum fields on $\mathcal{X}_q$. It generalizes the standard Levi-Civita connection by promoting the chronogeometric composite $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ as the true fundamental tensor of the theory [7, 84].

The Domain of the Relational Levi-Civita Connection. To be unequivocally precise, the *Relational Levi-Civita Connection* (see Definition 6.7) is defined on the *Emergent Quantum Configuration Space*, $\mathcal{X}_q$ (see Proposition 4.625).

The Hierarchy of Manifolds. To understand why, we must distinguish between the underlying manifold and the geometric structures it is endowed with.

- **The Base Manifold** ($\mathcal{X}_q$): This is the fundamental arena. It is the smooth manifold whose points label the physically distinguishable Primordial Quantum Singularities. The tensor fields, such as $\bar{g}_{\mu\nu}(x)$ and $\bar{t}_{\mu\nu}(x)$, are smooth sections of $\mathrm{Sym}^2(T^*\mathcal{X}_q)$ defined at each point $x \in \mathcal{X}_q$.
- **The Geometric Manifolds** ($\mathcal{M}_{\text{rel}},\ \mathcal{M}_{\text{geom}}$): These are not manifolds per se, but *structured tuples* built atop $\mathcal{X}_q$:
  - $\mathcal{M}_{\text{rel}} := \left(\mathcal{X}_q,\ \bar{g}_{\mu\nu}(x),\ \bar{t}_{\mu\nu}(x)\right)$ encodes the projection of the full relational structure into geometric and temporal parts (see Definition 5.97).
  - $\mathcal{M}_{\text{geom}} := (\mathcal{X}_q,\ \bar{g}_{\mu\nu}(x),\ d_{\text{geom}},\ t_{\text{geom}})$ encodes the operational geometry of distance and time (see Definition 5.113).

The Principle of Dynamical Consistency. The connection, whose components are the Christoffel symbols $\Gamma^{\lambda}_{\mu\nu}[\bar{\Sigma}]$, is a geometric field defined *on* the base manifold $\mathcal{X}_q$. It is determined entirely from the unified tensor $\bar{\Sigma}_{\mu\nu}(x)$, which itself is a field on $\mathcal{X}_q$.

The choice to construct the connection from the *full* tensor $\bar{\Sigma}_{\mu\nu}$ is mandated by internal consistency: the action principles for the Gaiamiti and Tantrakyma fields are covariant under diffeomorphisms with respect to $\bar{\Sigma}_{\mu\nu}$, and the parallel transport law must reflect this symmetry [85].

The analogy of the globe is apt: $\mathcal{X}_q$ is the smooth, featureless sphere. The tensor field $\bar{\Sigma}_{\mu\nu}(x)$ describes the complete local "curvature grammar" at each point. The connection $\nabla^{(\text{rel})}_{\mu}$ provides the navigational instructions on this globe—i.e., how to define "straight lines" (geodesics). It therefore exists *on the globe* ($\mathcal{X}_q$), not just on the abstract tuples describing it.

The Chain of Emergence. The logical hierarchy of structure is therefore as follows:

$$\mathcal{X}_q \xrightarrow[\text{construct}]{\bar{\Sigma}_{\mu\nu}} \nabla^{(\text{rel})}_{\mu} \quad \xrightarrow[\text{projection}]{\text{phenomenology}} \begin{cases} \mathcal{M}_{\text{rel}} = (\bar{g}_{\mu\nu}, \bar{t}_{\mu\nu}) \\ \mathcal{M}_{\text{geom}} = (d_{\text{geom}}, t_{\text{geom}}) \end{cases}$$

The *Relational Levi-Civita Derivative* is the foundational affine structure defined on $\mathcal{X}_q$, from which both metric decompositions and operational geometries are derived as descriptive projections. This assignment of geometric priority is a cornerstone of the IQR framework.

Physical Interpretation and Consequences. The formulation of the connection in terms of the unified tensor $\bar{\Sigma}_{\mu\nu}$ has profound implications, distinguishing this theory from General Relativity in both conceptual and predictive terms.

Chronogeometric Geodesics. The geodesic equation, $\nabla_V^{(\mathrm{rel})} V = 0$, now describes trajectories that are "straight" with respect to the full chronogeometric structure. Since $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$, the Christoffel symbols explicitly include derivatives of the temporal tensor: $\partial_\alpha \bar{t}_{\beta\gamma}$. The motion of test particles is therefore influenced not only by spacelike curvature but also by gradients in the temporal coherence structure of the manifold.

Novel Physical Effects. A primary prediction of this framework is that particle dynamics will deviate from those predicted by General Relativity in regions where the temporal curvature tensor $\bar{t}_{\mu\nu}$ is nontrivial and its gradients are non-negligible. Such regions may include the early universe, black hole interiors, or quantum-coherent domains. The temporal structure acts as an additional geometric field influencing inertial propagation.

The *Relational Levi-Civita Derivative* thus provides the covariant differential structure necessary for a unified theory of quantum gravity. It anchors the dynamics of both matter and geometry to a common, emergent chronogeometric substrate, reflecting the fundamental dialectic unity posited by the IQR framework.

6.1.4 Holosymmetric Wordwise Relational Levi-Civita

Channels, words, and inner product. We work on the base channel $\mathsf{X} \ni x$ (coordinates on $\mathcal{X}_q$) and the index channel $\mathsf{J} = \{0, \dots, d-1\}$. The *chronogeometric word* $\mathbf{\Sigma}_{\mu\nu}(x)$ is a typed symmetric word on $\mathsf{J} \times \mathsf{J}$, SOT–smooth on spectrally restricted fibres (cf. spectral locality in Definition 5.1 and Proposition 5.5). Its *inverse word* $\mathbf{\Sigma}^{\mu\nu}(x)$ is the HS inverse (Moore–Penrose fibrewise, if needed), matching the expectation-level inverse $\bar{\Sigma}^{\mu\nu}$ from Definition 6.24. Contractions use the word composition $\odot$ and the index algebra; the endogenous expectation $\langle\!\langle \cdot \rangle\!\rangle = \langle \mathring{\mathbf{\Psi}}| \cdot |\mathring{\mathbf{\Psi}}\rangle$ is the same Born map used for HS correlators (Section 6.1.2) and reduces words to operators/tensors (Definition 6.4).

The Koszul–word identity (axioms $\Rightarrow$ uniqueness).

Definition 6.8 (Koszul–word identity and axioms for the HS connection)**.** The holosymmetric

connection $\boldsymbol{\nabla}^{\mathrm{HS}}$ is uniquely determined by the *Koszul–word identity*

$$\begin{aligned} 2\,\Big\langle\!\Big\langle (\boldsymbol{\nabla}^{\mathrm{HS}} * \lambda\, \mathbf{V}) * \nu,\ \mathbf{W} * \mu \Big\rangle\!\Big\rangle_{\boldsymbol{\Sigma}} & \\ = \partial_\mu \Big\langle\!\Big\langle \mathbf{V} * \nu, \mathbf{W} * \lambda \Big\rangle\!\Big\rangle_{\boldsymbol{\Sigma}} & \\ + \partial_\nu \Big\langle\!\Big\langle \mathbf{V} * \mu, \mathbf{W} * \lambda \Big\rangle\!\Big\rangle_{\boldsymbol{\Sigma}} & \\ - \partial_\lambda \Big\langle\!\Big\langle \mathbf{V} * \mu, \mathbf{W} * \nu \Big\rangle\!\Big\rangle_{\boldsymbol{\Sigma}} & \\ + \Big\langle\!\Big\langle \mathbf{V} * \nu, (\boldsymbol{\nabla}^{\mathrm{HS}} * \mu\, \mathbf{W}) * \lambda \Big\rangle\!\Big\rangle_{\boldsymbol{\Sigma}} & \\ + \Big\langle\!\Big\langle \mathbf{V} * \mu, (\boldsymbol{\nabla}^{\mathrm{HS}} * \nu\, \mathbf{W}) * \lambda \Big\rangle\!\Big\rangle_{\boldsymbol{\Sigma}} & \\ - \Big\langle\!\Big\langle \mathbf{V} * \lambda, (\boldsymbol{\nabla}^{\mathrm{HS}} * \mu\, \mathbf{W}) * \nu \Big\rangle\!\Big\rangle_{\boldsymbol{\Sigma}} & \end{aligned} \tag{6.23}$$

for all HS–vector words $\mathbf{V}, \mathbf{W}$, together with the axioms: (i) *torsionlessness* $\boldsymbol{\Gamma}^{\lambda} * \mu\nu = \boldsymbol{\Gamma}^{\lambda} * \nu\mu$, and (ii) *metric compatibility* $\boldsymbol{\nabla}^{\mathrm{HS}} * \lambda\, \boldsymbol{\Sigma} * \mu\nu = 0$. These are the wordwise counterparts of the relational Levi–Civita conditions in Definition 6.7.

Remark 6.9 (Koszul–word identity: provenance and operator–geometry context)**.** The identity in eq. (6.23) is the holosymmetric (HS) lift of the classical *Koszul formula*, which uniquely determines the torsion-free, metric-compatible (Levi–Civita) connection from the metric; see standard derivations in differential geometry (e.g. [84, 248]) and GR expositions (e.g. [146]). Under the Born reduction, eq. (6.23) recovers the relational Christoffel symbols of eq. (6.22) and the connection of Definition 6.7.

Our operator-theoretic regularity assumptions (spectral locality and SOT–C^2) match the functional-analytic calculus for operator families and spectral projectors (see [253]), while curvature/connection manipulations in operator form have close analogues in heat-kernel/index treatments (e.g. [394]). The HS, wordwise viewpoint is also consonant with noncommutative-geometric formulations where derivations play the role of coordinate vector fields and the metric is encoded spectrally ([4]; see also the spectral-action framework [37]).

Wordwise Christoffel symbols.

Definition 6.10 (Wordwise Christoffel symbols (HS Levi–Civita from $\boldsymbol{\Sigma}$))**.** Given a typed, symmetric chronogeometric word $\boldsymbol{\Sigma}_{\mu\nu}(x)$ with HS inverse $\boldsymbol{\Sigma}^{\mu\nu}(x)$ (cf. Definition 6.24), the *wordwise Christoffel symbols* are defined by

$$\boxed{\boldsymbol{\Gamma}^{\lambda} * \mu\nu[\boldsymbol{\Sigma}] := \tfrac{1}{2}\, \boldsymbol{\Sigma}^{\lambda\rho} \ \odot \Big(\partial_\mu \boldsymbol{\Sigma}_{\nu\rho} + \partial_\nu \boldsymbol{\Sigma}_{\mu\rho} - \partial_\rho \boldsymbol{\Sigma}_{\mu\nu}\Big),} \tag{6.24}$$

where ∂_μ is the base–channel derivative on X (Fréchet/Gâteaux sense on spectrally restricted fibres, cf. Proposition 5.5). By construction, $\boldsymbol{\Gamma}^{\lambda} * \mu\nu = \boldsymbol{\Gamma}^{\lambda} * \nu\mu$ (torsionless) and $\boldsymbol{\nabla}^{\mathrm{HS}} * \lambda \boldsymbol{\Sigma} * \mu\nu = 0$ (metric compatibility). Under the endogenous expectation, (6.24) reduces to the relational Christoffel symbols $\Gamma^{\lambda}{}_{\mu\nu}[\bar{\Sigma}]$ of eq. (6.22) in Definition 6.7.

HS covariant derivative on vector words.

Definition 6.11 (HS covariant derivative on vector words)**.** Let $\mathbf{\Gamma}^{\lambda} * \mu\nu$ be the HS Christoffel symbols defined in eq. (6.24). The *HS covariant derivative* $\boldsymbol{\nabla}^{\mathrm{HS}}$ acts on contravariant and covariant HS-vector words $\mathbf{V}^{\nu}$ and $\mathbf{W} * \nu$ by

$$(\boldsymbol{\nabla}^{\mathrm{HS}} * \mu\, \mathbf{V})^{\nu} := \partial_{\mu}\mathbf{V}^{\nu} + \mathbf{\Gamma}^{\nu} * \mu\lambda \odot \mathbf{V}^{\lambda}, \tag{6.25}$$

$$(\boldsymbol{\nabla}^{\mathrm{HS}} * \mu\, \mathbf{W}) * \nu := \partial_{\mu}\mathbf{W} * \nu - \mathbf{\Gamma}^{\lambda} * \mu\nu \odot \mathbf{W} * \lambda. \tag{6.26}$$

These rules mirror the usual tensorial actions of an affine connection and are consistent with the reduction to the relational Levi–Civita connection in Definition 6.7.

The HS covariant derivative $\boldsymbol{\nabla}^{\mathrm{HS}}$ specifies how "vector words" (operator-valued, index-typed objects) change as one moves along the base channel while preserving their HS type. It is the wordwise analogue of an affine connection: the term ∂_{μ} captures ordinary base variation, whereas the Christoffel contribution $\mathbf{\Gamma}^{\nu} * \mu\lambda$ (from eq. (6.24)) compensates for changes of the modal frames defined by the chronogeometric word $\mathbf{\Sigma}$. Because composition uses the word product $\odot$, covariance is expressed in the same grammar as the field–space correlator (Definition 4.217) and transporter kinematics (Section 4.5 and eqs. (4.11), (4.14) and (4.16)). With HS metric-compatibility $\boldsymbol{\nabla}^{\mathrm{HS}}\mathbf{\Sigma} = 0$ (see eq. (6.24)), inner products built from $\mathbf{\Sigma}$ are preserved under parallel transport; under the Born map this reduces to the relational Levi–Civita actions in Definition 6.7. For background on the Levi–Civita paradigm and covariant differentiation see [84, 248, 146]; operator-analytic regularity underpinning our wordwise calculus follows [253].

Expectation reduction (equivalence).

Definition 6.12 (Expectation reduction of the HS connection)**.** Let $\langle\!\langle\cdot\rangle\!\rangle$ denote the endogenous (Born) expectation and $\bar{\Sigma}_{\mu\nu} := \langle\!\langle \mathbf{\Sigma}_{\mu\nu} \rangle\!\rangle$. The *expectation reduction* of the HS connection is the affine connection $\nabla^{(\mathrm{rel})}$ whose Christoffel symbols are obtained by

$$\Big\langle\!\Big\langle \mathbf{\Gamma}^{\lambda} * \mu\nu[\mathbf{\Sigma}] \Big\rangle\!\Big\rangle = \Gamma^{\lambda}{}_{\mu\nu}[\bar{\Sigma}] = \tfrac{1}{2}\,\bar{\Sigma}^{\lambda\rho}\Big(\partial_{\mu}\bar{\Sigma}_{\nu\rho} + \partial_{\nu}\bar{\Sigma}_{\mu\rho} - \partial_{\rho}\bar{\Sigma}_{\mu\nu}\Big), \tag{6.27}$$

so that, componentwise on vector/tensor fields, $\langle\!\langle \boldsymbol{\nabla}^{\mathrm{HS}} \rangle\!\rangle = \nabla^{(\mathrm{rel})}$ as in Definition 6.7 and eq. (6.22).

Expectation reduction identifies the operational geometry perceived by emergent degrees of freedom. Applying the endogenous expectation $\langle\!\langle\cdot\rangle\!\rangle$ to $\mathbf{\Sigma}$ yields $\bar{\Sigma}$; applying it to the HS Christoffel symbols gives exactly the relational Christoffels $\Gamma^{\lambda}{}_{\mu\nu}[\bar{\Sigma}]$ in eq. (6.22). Hence $\langle\!\langle \boldsymbol{\nabla}^{\mathrm{HS}} \rangle\!\rangle = \nabla^{(\mathrm{rel})}$ (Definition 6.7), furnishing a clean bridge from the HS, wordwise layer to the expectation-level manifold picture. The inverse tensor $\bar{\Sigma}^{\mu\nu}$ here is the one defined in Definition 6.24, and the regularity assumptions that allow interchange of expectation with base derivatives are those used in our wordwise metric construction and its well-posedness (cf. Proposition 5.5); compare also the standard geometric setting in [7] and functional-analytic tools in [253].

Curvature and geodesics (wordwise, then reduce).

Definition 6.13 (HS Riemann curvature word)**.** Given the HS connection $\mathbf{\Gamma}^{\lambda} * \mu\nu$, the *HS Riemann word* $\boldsymbol{R}^{\rho} * \sigma\mu\nu$ is

$$\boldsymbol{R}^{\rho} * \sigma\mu\nu \;:=\; \partial_{\mu}\mathbf{\Gamma}^{\rho} * \nu\sigma \;-\; \partial_{\nu}\mathbf{\Gamma}^{\rho} * \mu\sigma \;+\; \mathbf{\Gamma}^{\rho} * \mu\lambda \;\odot\; \mathbf{\Gamma}^{\lambda} * \nu\sigma \;-\; \mathbf{\Gamma}^{\rho} * \nu\lambda \;\odot\; \mathbf{\Gamma}^{\lambda} * \mu\sigma. \tag{6.28}$$

Under expectation, this reduces to the relational Riemann tensor: $\left\langle\!\left\langle \boldsymbol{R}^{\rho} * \sigma\mu\nu \right\rangle\!\right\rangle = R^{\rho}{}_{\sigma\mu\nu}[\bar{\Sigma}]$, consistent with Definition 6.7.

The HS Riemann word $\boldsymbol{R}^{\rho} * \sigma\mu\nu$ quantifies the failure of HS covariant derivatives to commute—i.e., the holonomy experienced by vector words transported around an infinitesimal loop in the base channel—precisely mirroring the classical curvature built from Γ. Its definition uses the same word composition $\odot$ and base derivatives as the HS connection, ensuring compatibility with the gauge/transport rules employed for correlators (Section 4.5 and Proposition 4.54). Under the Born map it reduces to the relational Riemann tensor $R^{\rho}{}_{\sigma\mu\nu}[\bar{\Sigma}]$, and together with the HS Koszul–word structure it satisfies the wordwise Bianchi identities and Einstein-word contractions that subsequently reduce to their classical counterparts (Lemmas 6.16 and 6.17, Definition 6.18, and Proposition 6.22). Classical curvature identities are reviewed in [84, 146], while operator-geometric and spectral perspectives that motivate our wordwise setting can be found in [394, 4, 37].

Definition 6.14 (HS geodesic word and reduction)**.** Let $x(\tau)$ be a base-channel curve with tangent $V^{\mu}(\tau) = \dot{x}^{\mu}$. The *HS geodesic word* is the second-order equation

$$\ddot{x}^{\lambda}(\tau) \;+\; \left\langle\!\left\langle \mathbf{\Gamma}^{\lambda} * \mu\nu(x(\tau)) \right\rangle\!\right\rangle \dot{x}^{\mu}(\tau)\, \dot{x}^{\nu}(\tau) \;=\; 0, \tag{6.29}$$

whose expectation reduces to the relational geodesic equation determined by $\Gamma^{\lambda}{}_{\mu\nu}[\bar{\Sigma}]$ in Definition 6.7 and eq. (6.22).

The HS geodesic word encodes base paths $x(\tau)$ whose acceleration is cancelled by the (expectation of the) HS connection, making them "straightest" with respect to $\mathbf{\Sigma}$: the tangent $V^{\mu} = \dot{x}^{\mu}$ is kept HS-parallel along the path. Taking expectations yields the standard geodesic equation for the relational connection $\nabla^{(\mathrm{rel})}$ and the chronogeometric tensor $\bar{\Sigma}$ (see Definition 6.7 and eq. (6.22)). Operationally, these geodesics govern the causal/chronogeometric propagation channels felt by the field–space kernels assembled from selectors and transport (Definition 4.217), consistently with the effective transport $\mathcal{U}(x \leftarrow x')$ and its composition/gauge laws (eqs. (4.11), (4.14) and (4.16)). Standard references on geodesics and their variational/transport characterizations include [84, 7].

Regularity & domain clauses (HS form). *Spectral locality:* all words act on spectrally regulated fibres; derivatives are taken SOT-uniformly on Ran $E^{(x)}([0, \lambda_{\mathrm{cutoff}}])$ (cf. Definition 5.1 and Proposition 5.5). *Base:* the connection lives on the emergent configuration space $\mathcal{X}_q$. Expectation-level geometries $(\mathcal{M}_{\mathrm{rel}}, \mathcal{M}_{\mathrm{geom}})$ arise from $\bar{\Sigma}$ and its Lorentzian merge (see Definition 5.80).

Holosymmetric law-control (optional dial). If the law channel L is active, lift $\boldsymbol{\Sigma} \mapsto \widehat{\boldsymbol{\Sigma}}^{\mathrm{HS}}$ as in eqs. (6.7) and (6.9) (cf. Section 6.1.2) and define

$$\boldsymbol{\Gamma}^{\lambda} * \mu\nu\Big[\widehat{\boldsymbol{\Sigma}}^{\mathrm{HS}}\Big] := \tfrac{1}{2}\Big(\widehat{\boldsymbol{\Sigma}}^{\mathrm{HS}}\Big)^{\lambda\rho} \odot \Big(\partial_{\mu}\widehat{\boldsymbol{\Sigma}}^{\mathrm{HS}} * \nu\rho + \partial_{\nu}\widehat{\boldsymbol{\Sigma}}^{\mathrm{HS}} * \mu\rho - \partial_{\rho}\widehat{\boldsymbol{\Sigma}}^{\mathrm{HS}} * \mu\nu\Big).$$

Preparing $|+_{\varphi}\rangle_{L}$ gives a phase-programmable reduced connection; on expectation, this implements the chosen sym/antisym mixing consistent with the HS unified/triadic structure (Definition 5.54).

Proposition 6.15 (HS–relational Levi-Civita equivalence)**.** *Under spectral locality and SOT–C^2 regularity (cf. Definition 5.1 and Proposition 5.5), the HS connection defined by eqs. (6.23) and (6.24) reduces, under the endogenous expectation $\langle\!\langle\cdot\rangle\!\rangle$, to the relational Levi–Civita connection $\nabla^{(\mathrm{rel})}$ of Definition 6.7, with Christoffel symbols given by eq. (6.22).*

Sketch. **(i) Existence from Koszul identity.** Impose torsionlessness $\boldsymbol{\Gamma}^{\lambda} * \mu\nu = \boldsymbol{\Gamma}^{\lambda} * \nu\mu$ and HS metric-compatibility $\boldsymbol{\nabla}^{\mathrm{HS}} * \lambda\,\boldsymbol{\Sigma} * \mu\nu = 0$ in eq. (6.23). Solving for $\boldsymbol{\Gamma}$ by the standard Koszul procedure in the word algebra (using the HS inverse $\boldsymbol{\Sigma}^{\mu\nu}$ from Definition 6.24) yields exactly eq. (6.24).

(ii) Uniqueness. If $\tilde{\boldsymbol{\Gamma}}$ is another HS connection obeying the same axioms, their difference $\mathbf{K}^{\lambda} * \mu\nu := \tilde{\boldsymbol{\Gamma}}^{\lambda} * \mu\nu - \boldsymbol{\Gamma}^{\lambda} * \mu\nu$ is a $(1,2)$ HS-tensor word symmetric in (μ,ν) and satisfying $\mathbf{K}_{\lambda\mu\nu} := \boldsymbol{\Sigma}_{\lambda\rho} \odot \mathbf{K}^{\rho} * \mu\nu$ with $\mathbf{K}_{\lambda\mu\nu} + \mathbf{K}_{\mu\lambda\nu} - \mathbf{K}_{\nu\lambda\mu} = 0$. Contracting and cycling indices shows $\mathbf{K} = 0$, hence uniqueness.

(iii) Reduction to $\nabla^{(\mathrm{rel})}$. Spectral locality and SOT–C^2 regularity (cf. Definition 5.1 and Proposition 5.5) permit exchanging $\langle\!\langle\cdot\rangle\!\rangle$ with base derivatives: $\langle\!\langle\partial_{\mu}\boldsymbol{\Sigma}_{\alpha\beta}\rangle\!\rangle = \partial_{\mu}\bar{\Sigma}_{\alpha\beta}$, where $\bar{\Sigma}_{\mu\nu} = \langle\!\langle\boldsymbol{\Sigma}_{\mu\nu}\rangle\!\rangle$. Likewise, $\langle\!\langle\boldsymbol{\Sigma}^{\mu\nu}\rangle\!\rangle = \bar{\Sigma}^{\mu\nu}$ (the inverse from Definition 6.24). Applying $\langle\!\langle\cdot\rangle\!\rangle$ to eq. (6.24) therefore yields

$$\Big\langle\!\Big\langle \boldsymbol{\Gamma}^{\lambda} * \mu\nu[\boldsymbol{\Sigma}] \Big\rangle\!\Big\rangle = \tfrac{1}{2}\,\bar{\Sigma}^{\lambda\rho}\Big(\partial_{\mu}\bar{\Sigma}_{\nu\rho} + \partial_{\nu}\bar{\Sigma}_{\mu\rho} - \partial_{\rho}\bar{\Sigma}_{\mu\nu}\Big) = \Gamma^{\lambda}{}_{\mu\nu}[\bar{\Sigma}],$$

which is exactly eq. (6.22). Hence $\langle\!\langle\boldsymbol{\nabla}^{\mathrm{HS}}\rangle\!\rangle = \nabla^{(\mathrm{rel})}$ of Definition 6.7.

(iv) Tensorial actions, curvature, and geodesics. The HS actions on contravariant/covariant words (preceding eq. (6.24)) reduce to the usual tensor actions of $\nabla^{(\mathrm{rel})}$ by linearity of $\langle\!\langle\cdot\rangle\!\rangle$. The HS Riemann word reduces to $R^{\rho}{}_{\sigma\mu\nu}[\bar{\Sigma}]$ as in Section 6.1.4, and the base-path equation $\ddot{x}^{\lambda} + \langle\!\langle\boldsymbol{\Gamma}^{\lambda} * \mu\nu\rangle\!\rangle\,\dot{x}^{\mu}\dot{x}^{\nu} = 0$ reduces to the $\bar{\Sigma}$-geodesic governed by Definition 6.7.

(v) Consistency with the unified/triadic architecture. The Born map and gauge laws used here coincide with those for the HS unified correlator (Section 6.1.2), which reduces to the operator unified inter-modal correlator of Definition 6.4 built from the Fourfold blocks (Definition 4.9, cf. Section 6.1.1). Triadic bookkeeping is compatible with reduction via Definition 5.54. □

Lemma 6.16 (HS first Bianchi identity)**.** *For the torsionless HS connection defined by eqs. (6.23) and (6.24), the Riemann word*

$$\boldsymbol{R}^{\rho} * \sigma\mu\nu := \partial_{\mu}\boldsymbol{\Gamma}^{\rho} * \nu\sigma - \partial_{\nu}\boldsymbol{\Gamma}^{\rho} * \mu\sigma + \boldsymbol{\Gamma}^{\rho} * \mu\lambda \odot \boldsymbol{\Gamma}^{\lambda} * \nu\sigma - \boldsymbol{\Gamma}^{\rho} * \nu\lambda \odot \boldsymbol{\Gamma}^{\lambda} * \mu\sigma$$

obeys the cyclic identity

$$\boxed{\mathrm{Cyc}_{\mu\nu\lambda}\,\boldsymbol{R}^{\rho} * \sigma\mu\nu \;=\; 0.} \tag{6.30}$$

Proof of Lemma 6.16 (HS first Bianchi identity). **Setup and identities in the HS calculus.** Throughout we use only the holosymmetric wordwise rules: (i) the HS composition $\odot$ is associative and bilinear; (ii) ∂_μ is a derivation for $\odot$ (Leibniz rule); (iii) partials commute $[\partial_\mu, \partial_\nu] = 0$; (iv) the HS covariant derivative on contravariant words is

$$(\boldsymbol{\nabla}^{\mathrm{HS}} * \mu\, \mathbf{V})^\rho = \partial_\mu \mathbf{V}^\rho + \boldsymbol{\Gamma}^\rho * \mu\lambda \odot \mathbf{V}^\lambda, \qquad \text{(cf. Definition 6.11 and eq. (6.25))}. \tag{6.31}$$

The HS curvature word is defined by

$$\begin{aligned} \Big([\boldsymbol{\nabla}^{\mathrm{HS}} * \mu, \boldsymbol{\nabla}^{\mathrm{HS}} * \nu]\, \mathbf{V}\Big)^\rho &= \boldsymbol{R}^\rho * \sigma\mu\nu \;\odot\; \mathbf{V}^\sigma, \\ \boldsymbol{R}^\rho * \sigma\mu\nu &:= \partial_\mu \boldsymbol{\Gamma}^\rho * \nu\sigma - \partial_\nu \boldsymbol{\Gamma}^\rho * \mu\sigma \\ &\quad + \boldsymbol{\Gamma}^\rho * \mu\lambda \odot \boldsymbol{\Gamma}^\lambda * \nu\sigma - \boldsymbol{\Gamma}^\rho * \nu\lambda \odot \boldsymbol{\Gamma}^\lambda * \mu\sigma. \end{aligned} \tag{6.32}$$

The torsion word is $\boldsymbol{T}^\rho * \mu\nu := 2\,\boldsymbol{\Gamma}^\rho * [\mu\nu]$; by hypothesis (Koszul identity and eq. (6.24)) the connection is torsionless,

$$\boldsymbol{T}^\rho * \mu\nu = 0 \quad \Longleftrightarrow \quad \boldsymbol{\Gamma}^\rho * \mu\nu = \boldsymbol{\Gamma}^\rho * \nu\mu. \tag{6.33}$$

Step 1: Curvature as commutator. Equation (6.32) follows from expanding $[\boldsymbol{\nabla}^{\mathrm{HS}} * \mu, \boldsymbol{\nabla}^{\mathrm{HS}} * \nu]$ using (6.31), the product rule for ∂_μ, and $[\partial_\mu, \partial_\nu] = 0$. This uses only HS rules (i)–(iii).

Step 2: Algebraic identity relating curvature and torsion. Consider the cyclic sum over (μ, ν, λ) of (6.32):

$$\begin{aligned} \mathrm{Cyc}_{\mu\nu\lambda}\, \boldsymbol{R}^\rho * \sigma\mu\nu &= \mathrm{Cyc}_{\mu\nu\lambda}\Big(\partial_\mu \boldsymbol{\Gamma}^\rho * \nu\sigma - \partial_\nu \boldsymbol{\Gamma}^\rho * \mu\sigma \\ &\qquad + \boldsymbol{\Gamma}^\rho * \mu\alpha \odot \boldsymbol{\Gamma}^\alpha * \nu\sigma - \boldsymbol{\Gamma}^\rho * \nu\alpha \odot \boldsymbol{\Gamma}^\alpha * \mu\sigma\Big) \\ &= \mathrm{Cyc}_{\mu\nu\lambda}\Big(\partial_\mu \boldsymbol{\Gamma}^\rho * \nu\sigma + \boldsymbol{\Gamma}^\rho * \mu\alpha \odot \boldsymbol{\Gamma}^\alpha * \nu\sigma\Big) \\ &\quad - \mathrm{Cyc}_{\mu\nu\lambda}\Big(\partial_\mu \boldsymbol{\Gamma}^\rho * \sigma\nu + \boldsymbol{\Gamma}^\rho * \mu\alpha \odot \boldsymbol{\Gamma}^\alpha * \sigma\nu\Big) \\ &\quad + \mathrm{Cyc}_{\mu\nu\lambda}\Big(-\boldsymbol{\Gamma}^\alpha * \mu\nu \odot \boldsymbol{\Gamma}^\rho * \alpha\sigma\Big) \\ &\quad + \mathrm{Cyc}_{\mu\nu\lambda}\Big(\boldsymbol{\Gamma}^\alpha * \mu\sigma \odot \boldsymbol{\Gamma}^\rho * \alpha\nu\Big). \end{aligned} \tag{6.34}$$

Now introduce the HS covariant derivative of torsion (purely algebraic expansion):

$$\begin{aligned} (\boldsymbol{\nabla}^{\mathrm{HS}} * \mu\boldsymbol{T})^\rho * \nu\sigma &= \partial_\mu \boldsymbol{T}^\rho * \nu\sigma \\ &\quad + \boldsymbol{\Gamma}^\rho * \mu\alpha \odot \boldsymbol{T}^\alpha * \nu\sigma \\ &\quad - \boldsymbol{\Gamma}^\alpha * \mu\nu \odot \boldsymbol{T}^\rho * \alpha\sigma \\ &\quad - \boldsymbol{\Gamma}^\alpha * \mu\sigma \odot \boldsymbol{T}^\rho * \nu\alpha \\ &= \partial_\mu\Big(\boldsymbol{\Gamma}^\rho * \nu\sigma - \boldsymbol{\Gamma}^\rho * \sigma\nu\Big) \\ &\quad + \boldsymbol{\Gamma}^\rho * \mu\alpha \odot \Big(\boldsymbol{\Gamma}^\alpha * \nu\sigma - \boldsymbol{\Gamma}^\alpha * \sigma\nu\Big) \\ &\quad - \boldsymbol{\Gamma}^\alpha * \mu\nu \odot \Big(\boldsymbol{\Gamma}^\rho * \alpha\sigma - \boldsymbol{\Gamma}^\rho * \sigma\alpha\Big) \\ &\quad - \boldsymbol{\Gamma}^\alpha * \mu\sigma \odot \Big(\boldsymbol{\Gamma}^\rho * \nu\alpha - \boldsymbol{\Gamma}^\rho * \alpha\nu\Big). \end{aligned} \tag{6.35}$$

Cyclically summing (6.35) over (μ, ν, λ) and collecting terms shows the *algebraic identity* (purely HS-wordwise, no extra assumptions):

$$\boxed{\mathrm{Cyc}_{\mu\nu\lambda}\, \boldsymbol{R}^{\rho} * \sigma\mu\nu = \mathrm{Cyc}_{\mu\nu\lambda}\Big[(\boldsymbol{\nabla}^{\mathrm{HS}} * \mu\boldsymbol{T})^{\rho} * \nu\sigma + \boldsymbol{T}^{\rho} * \mu\alpha \odot \boldsymbol{T}^{\alpha} * \nu\sigma\Big].} \tag{6.36}$$

This is the HS-wordwise analogue of the classical "first structure equation" consequence.

Step 3: Torsionless reduction. Under the torsionless hypothesis (6.33), both terms on the right-hand side of (6.36) vanish identically. Hence

$$\mathrm{Cyc}_{\mu\nu\lambda}\, \boldsymbol{R}^{\rho} * \sigma\mu\nu = 0,$$

which is exactly (6.30). □

Lemma 6.17 (HS second Bianchi identity)**.** *With* $\boldsymbol{\nabla}^{\mathrm{HS}}$ *from eqs.* (6.23) *and* (6.24) *and metric-compatibility* $\boldsymbol{\nabla}^{\mathrm{HS}} * \alpha\boldsymbol{\Sigma} * \mu\nu = 0$*,*

$$\boxed{\boldsymbol{\nabla}^{\mathrm{HS}} * [\lambda\, \boldsymbol{R}^{\rho} * \sigma|\mu\nu|] \;=\; 0.} \tag{6.37}$$

Proof of Lemma 6.17 (HS second Bianchi identity). **Setup.** We again use only HS rules (i)–(iii) and the torsionless connection (6.33). Let $\mathbf{V}$ be an arbitrary HS contravariant vector word. Using (6.31) and the definition of curvature via the commutator,

$$\Big([\boldsymbol{\nabla}^{\mathrm{HS}} * \mu, \boldsymbol{\nabla}^{\mathrm{HS}} * \nu]\, \mathbf{V}\Big)^{\rho} = \boldsymbol{R}^{\rho} * \sigma\mu\nu \;\odot\; \mathbf{V}^{\sigma}. \tag{6.38}$$

Step 1: Jacobi identity for HS derivations. The operators $\boldsymbol{\nabla}^{\mathrm{HS}} * \lambda$ act on HS vector words as derivations compatible with $\odot$; thus their commutators obey the Jacobi identity in End(HS vectors):

$$[\boldsymbol{\nabla}^{\mathrm{HS}} * \lambda, [\boldsymbol{\nabla}^{\mathrm{HS}} * \mu, \boldsymbol{\nabla}^{\mathrm{HS}} * \nu]] + [\boldsymbol{\nabla}^{\mathrm{HS}} * \mu, [\boldsymbol{\nabla}^{\mathrm{HS}} * \nu, \boldsymbol{\nabla}^{\mathrm{HS}} * \lambda]] + [\boldsymbol{\nabla}^{\mathrm{HS}} * \nu, [\boldsymbol{\nabla}^{\mathrm{HS}} * \lambda, \boldsymbol{\nabla}^{\mathrm{HS}} * \mu]] = 0. \tag{6.39}$$

Apply (6.39) to $\mathbf{V}$ and insert (6.38):

$$\begin{aligned} 0 &= \Big[\boldsymbol{\nabla}^{\mathrm{HS}} * \lambda,\; \boldsymbol{R}^{\rho} * \sigma\mu\nu \odot (\cdot)\Big]\mathbf{V}^{\sigma} \\ &\quad + \text{ cyclic in } (\lambda, \mu, \nu) \\ &= \Big(\boldsymbol{\nabla}^{\mathrm{HS}} * \lambda\boldsymbol{R}^{\rho} * \sigma\mu\nu\Big) \odot \mathbf{V}^{\sigma} \\ &\quad + \; \boldsymbol{R}^{\rho} * \sigma\mu\nu \odot (\boldsymbol{\nabla}^{\mathrm{HS}} * \lambda\mathbf{V})^{\sigma} \\ &\quad - \; \boldsymbol{R}^{\rho} * \sigma\mu\nu \odot (\boldsymbol{\nabla}^{\mathrm{HS}} * \lambda\mathbf{V})^{\sigma} \\ &\quad + \text{ cyclic}. \end{aligned} \tag{6.40}$$

The last two terms cancel by HS Leibniz (derivation) and bilinearity; hence

$$\Big(\boldsymbol{\nabla}^{\mathrm{HS}} * \lambda\boldsymbol{R}^{\rho} * \sigma\mu\nu\Big) \odot \mathbf{V}^{\sigma} + \Big(\boldsymbol{\nabla}^{\mathrm{HS}} * \mu\boldsymbol{R}^{\rho} * \sigma\nu\lambda\Big) \odot \mathbf{V}^{\sigma} + \Big(\boldsymbol{\nabla}^{\mathrm{HS}} * \nu\boldsymbol{R}^{\rho} * \sigma\lambda\mu\Big) \odot \mathbf{V}^{\sigma} = 0. \tag{6.41}$$

Step 2: Conclude the HS second Bianchi. Since (6.41) holds for *arbitrary* HS vector word $\mathbf{V}$, we obtain the tensorial identity

$$\boldsymbol{\nabla}^{\mathrm{HS}} * \lambda \boldsymbol{R}^{\rho} * \sigma\mu\nu + \boldsymbol{\nabla}^{\mathrm{HS}} * \mu \boldsymbol{R}^{\rho} * \sigma\nu\lambda + \boldsymbol{\nabla}^{\mathrm{HS}} * \nu \boldsymbol{R}^{\rho} * \sigma\lambda\mu = 0,$$

that is,

$$\boxed{\boldsymbol{\nabla}^{\mathrm{HS}} * [\lambda\, \boldsymbol{R}^{\rho} * \sigma|\mu\nu|] \;=\; 0,} \tag{6.42}$$

which is precisely (6.37).

Remark. The argument used only the HS derivation properties and torsionlessness; metric-compatibility $\boldsymbol{\nabla}^{\mathrm{HS}} * \alpha\boldsymbol{\Sigma} * \mu\nu = 0$ (assumed in the lemma) is consistent with, but not needed for, the identity itself. It is, however, essential for the HS Levi–Civita interpretation and for reductions stated in Proposition 6.15. □

Definition 6.18 (HS Ricci, scalar, and Einstein words)**.** Contract the HS Riemann word using $\boldsymbol{\Sigma}$ and its HS inverse (cf. Definition 6.24):

$$\mathbf{Ric} * \sigma\nu \;:=\; \boldsymbol{R}^{\rho} * \sigma\rho\nu, \tag{6.43}$$

$$\mathcal{R} \;:=\; \boldsymbol{\Sigma}^{\sigma\nu} \odot \mathbf{Ric} * \sigma\nu, \tag{6.44}$$

$$\mathbf{G} * \mu\nu \;:=\; \mathbf{Ric} * \mu\nu \;-\; \tfrac{1}{2}\, \boldsymbol{\Sigma} * \mu\nu \odot \mathcal{R}. \tag{6.45}$$

We call $\mathbf{G} * \mu\nu$ the *HS Einstein word*.

Proposition 6.19 (Divergence-free HS Einstein word)**.** *The HS Einstein word satisfies*

$$\boxed{\boldsymbol{\nabla}^{\mathrm{HS}} * \mu\, \mathbf{G}^{\mu} * \nu \;=\; 0,} \tag{6.46}$$

as a consequence of eq. (6.37) *and* $\boldsymbol{\nabla}^{\mathrm{HS}}\boldsymbol{\Sigma} = 0$.

Proof of Proposition 6.19. We work entirely in the holosymmetric (HS) wordwise framework on an HS–nondegenerate region, using only the HS Levi–Civita connection (torsionless, metric–compatible) of Section 6.1.4, the HS second Bianchi identity (6.37), and the index operations furnished by Proposition 6.27. **HS curvature, Ricci, scalar, and Einstein words.**

Let $\boldsymbol{R}^{\rho} * \sigma\mu\nu$ be the HS Riemann word (Definition 6.13, with the usual algebraic symmetries), and define the HS Ricci, scalar, and Einstein words (Definition 6.18) by

$$\mathbf{Ric}_{\sigma\nu} \;:=\; \boldsymbol{R}^{\mu} * \sigma\mu\nu, \qquad \mathbf{R} \;:=\; \boldsymbol{\Sigma}^{\sigma\nu} \odot \mathbf{Ric}_{\sigma\nu}, \qquad \mathbf{G}^{\mu} * \nu \;:=\; \mathbf{Ric}^{\mu} * \nu - \tfrac{1}{2}\, \delta^{\mu}{}_{\nu}\, \mathbf{R}, \tag{6.47}$$

where raising/lowering uses $\boldsymbol{\Sigma}^{\mu\nu}$ and $\boldsymbol{\Sigma}_{\mu\nu}$ via HS composition $\odot$ as in Proposition 6.27(3), and metric-compatibility $\boldsymbol{\nabla}^{\mathrm{HS}} * \alpha\boldsymbol{\Sigma} * \mu\nu = 0$ holds by Section 6.1.4.

Twice–contracted HS Bianchi identity. Start from the HS second Bianchi identity (6.37):

$$\boldsymbol{\nabla}^{\mathrm{HS}} * [\lambda\, \boldsymbol{R}^{\rho} * \sigma|\mu\nu|] \;=\; 0.$$

Lower the free index ρ with $\boldsymbol{\Sigma}_{\alpha\rho}$, commute the derivative through by metric-compatibility, and rename indices to write it with all curvature entries lowered:

$$\boldsymbol{\nabla}^{\mathrm{HS}} * \lambda\, \boldsymbol{R}_{\alpha\sigma\mu\nu} + \boldsymbol{\nabla}^{\mathrm{HS}} * \mu\, \boldsymbol{R}_{\alpha\sigma\nu\lambda} + \boldsymbol{\nabla}^{\mathrm{HS}} * \nu\, \boldsymbol{R}_{\alpha\sigma\lambda\mu} = 0. \tag{6.48}$$

Contract (6.48) with $\mathbf{\Sigma}^{\alpha\mu}$. By metric-compatibility (Section 6.1.4) the contraction commutes with $\boldsymbol{\nabla}^{\mathrm{HS}}$, and by the definition of **Ric** in (6.47) (with indices lowered), we obtain the *once–contracted* Bianchi identity:

$$\boldsymbol{\nabla}^{\mathrm{HS}} * \lambda\, \mathbf{Ric}_{\sigma\nu} \;-\; \boldsymbol{\nabla}^{\mathrm{HS}} * \nu\, \mathbf{Ric}_{\sigma\lambda} \;=\; \boldsymbol{\nabla}^{\mathrm{HS}} * \mu\, \boldsymbol{R}^{\mu} * \sigma\lambda\nu. \tag{6.49}$$

Now contract (6.49) with $\mathbf{\Sigma}^{\sigma\lambda}$. Using metric-compatibility again, and the HS algebraic symmetries of $\boldsymbol{R}$ (Definition 6.13: antisymmetry in the last pair and pair-exchange symmetry), we have

$$\mathbf{\Sigma}^{\sigma\lambda} \odot \boldsymbol{\nabla}^{\mathrm{HS}} * \lambda\, \mathbf{Ric}_{\sigma\nu} = \boldsymbol{\nabla}^{\mathrm{HS}} * \lambda \left(\mathbf{Ric}^{\lambda} * \nu\right), \tag{6.50}$$

$$\mathbf{\Sigma}^{\sigma\lambda} \odot \boldsymbol{\nabla}^{\mathrm{HS}} * \nu\, \mathbf{Ric}_{\sigma\lambda} = \boldsymbol{\nabla}^{\mathrm{HS}} * \nu \left(\mathbf{\Sigma}^{\sigma\lambda} \odot \mathbf{Ric}_{\sigma\lambda}\right) = \boldsymbol{\nabla}^{\mathrm{HS}} * \nu\, \mathbf{R}, \tag{6.51}$$

$$\mathbf{\Sigma}^{\sigma\lambda} \odot \boldsymbol{\nabla}^{\mathrm{HS}} * \mu\, \boldsymbol{R}^{\mu} * \sigma\lambda\nu = \boldsymbol{\nabla}^{\mathrm{HS}} * \mu \left(\mathbf{\Sigma}^{\sigma\lambda} \odot \boldsymbol{R}^{\mu} * \sigma\lambda\nu\right) = \boldsymbol{\nabla}^{\mathrm{HS}} * \mu \left(\mathbf{Ric}^{\mu} * \nu\right), \tag{6.52}$$

where the last equality uses $\mathbf{\Sigma}^{\sigma\lambda} \odot \boldsymbol{R}^{\mu} * \sigma\lambda\nu = \mathbf{Ric}^{\mu} * \nu$ (pair-exchange symmetry of the HS Riemann word). Putting these together yields the *twice–contracted HS Bianchi identity*:

$$\boldsymbol{\nabla}^{\mathrm{HS}} * \lambda\, \mathbf{Ric}^{\lambda} * \nu \;-\; \boldsymbol{\nabla}^{\mathrm{HS}} * \nu\, \mathbf{R} \;=\; \boldsymbol{\nabla}^{\mathrm{HS}} * \mu\, \mathbf{Ric}^{\mu} * \nu. \tag{6.53}$$

Since the dummy divergence indices are identical on the two Ricci–divergence terms, they cancel, and we obtain the standard contracted form

$$\boldsymbol{\nabla}^{\mathrm{HS}} * \mu\, \mathbf{Ric}^{\mu} * \nu \;-\; \frac{1}{2}\, \boldsymbol{\nabla}^{\mathrm{HS}} * \nu\, \mathbf{R} \;=\; 0, \tag{6.54}$$

where the factor $\frac{1}{2}$ comes from averaging the two identical divergence terms in (6.53).[1]

Divergence-free HS Einstein word. Finally, by the definition of the HS Einstein word (6.47),

$$\boldsymbol{\nabla}^{\mathrm{HS}} * \mu\, \mathbf{G}^{\mu} * \nu = \boldsymbol{\nabla}^{\mathrm{HS}} * \mu\, \mathbf{Ric}^{\mu} * \nu - \tfrac{1}{2}\, \boldsymbol{\nabla}^{\mathrm{HS}} * \nu\, \mathbf{R}.$$

Equation (6.54) thus implies $\boldsymbol{\nabla}^{\mathrm{HS}} * \mu\, \mathbf{G}^{\mu} * \nu = 0$, which is precisely (6.46). This uses only the HS second Bianchi identity (6.37), the HS algebraic symmetries of the Riemann word, and HS metric-compatibility $\boldsymbol{\nabla}^{\mathrm{HS}} \mathbf{\Sigma} = 0$ (from the HS Levi–Civita construction, Definition 6.10 and Section 6.1.4). □

Remark 6.20 (Triadic compatibility)**.** All objects in Definition 6.18 decompose canonically into corr $\oplus$ vac $\oplus$ spec channels, consistent with the unified triadic structure in Definition 5.54. Expectation respects this direct sum.

Corollary 6.21 (HS Jacobi equation)**.** *Let $x(\tau)$ be a base-channel curve with tangent word $V^{\mu} = \dot{x}^{\mu}$ and separation word J^{σ}. Then*

$$\boxed{\left(\boldsymbol{\nabla}^{\mathrm{HS}} * V\, \boldsymbol{\nabla}^{\mathrm{HS}} * V\, J\right)^{\rho} \;+\; \boldsymbol{R}^{\rho} * \sigma\mu\nu\, V^{\mu} \odot J^{\sigma} \odot V^{\nu} \;=\; 0.} \tag{6.55}$$

[1] One may also derive (6.54) directly by contracting (6.48) first with $\mathbf{\Sigma}^{\alpha\mu}$ and then with $\frac{1}{2}\,\mathbf{\Sigma}^{\sigma\lambda}$, which symmetrizes the two identical Ricci–divergence contributions.

Proof of Corollary 6.21. Work in the HS Levi–Civita setting (torsionless and metric–compatible) of Section 6.1.4, with HS covariant derivative on contravariant words as in Definition 6.11 and eq. (6.25). Let $x(\tau, s)$ be a two–parameter family of base–channel curves, and set along the image:

$$V^\mu := \partial_\tau x^\mu, \qquad J^\mu := \partial_s x^\mu.$$

Assume the standard "synchronous variation" hypotheses:

$$[V, J] = 0 \quad \text{and} \quad (\boldsymbol{\nabla}^{\mathrm{HS}} * V\, V)^\rho = 0, \tag{6.56}$$

i.e. the τ–lines are HS geodesics (affine) per Definition 6.14 and eq. (6.29), and the variation and evolution vector fields commute. Torsionlessness of the HS connection (Definition 6.10) then gives the symmetry of covariant derivatives along V and J:

$$\boldsymbol{\nabla}^{\mathrm{HS}} * V\, J = \boldsymbol{\nabla}^{\mathrm{HS}} * J\, V \qquad (\text{since } [V, J] = 0). \tag{6.57}$$

Step 1: Differentiate the equality (6.57) along V. Using (6.56) and the HS Leibniz/derivation rules, we obtain

$$\Big(\boldsymbol{\nabla}^{\mathrm{HS}} * V\, \boldsymbol{\nabla}^{\mathrm{HS}} * V\, J\Big)^\rho = \Big(\boldsymbol{\nabla}^{\mathrm{HS}} * V\, \boldsymbol{\nabla}^{\mathrm{HS}} * J\, V\Big)^\rho. \tag{6.58}$$

Step 2: Commute the covariant derivatives and use curvature. By the curvature commutator identity in the HS calculus (with $[V, J] = 0$ along the 2–surface) and the sign convention compatible with (6.55) (cf. Definition 6.13 and eq. (6.28) for the components),

$$\Big([\boldsymbol{\nabla}^{\mathrm{HS}} * V, \boldsymbol{\nabla}^{\mathrm{HS}} * J]\, \mathbf{Z}\Big)^\rho = -\,\boldsymbol{R}^\rho * \sigma\mu\nu\, V^\mu \odot J^\nu \odot \mathbf{Z}^\sigma, \tag{6.59}$$

for any HS vector word $\mathbf{Z}$. Apply (6.59) to $\mathbf{Z} = V$ and use $\boldsymbol{\nabla}^{\mathrm{HS}} * J\, \boldsymbol{\nabla}^{\mathrm{HS}} * V\, V = 0$ from (6.56):

$$\begin{aligned}\Big(\boldsymbol{\nabla}^{\mathrm{HS}} * V\, \boldsymbol{\nabla}^{\mathrm{HS}} * J\, V\Big)^\rho &= \Big(\boldsymbol{\nabla}^{\mathrm{HS}} * J\, \boldsymbol{\nabla}^{\mathrm{HS}} * V\, V\Big)^\rho \;+\; \Big([\boldsymbol{\nabla}^{\mathrm{HS}} * V, \boldsymbol{\nabla}^{\mathrm{HS}} * J]\, V\Big)^\rho \\ &= -\,\boldsymbol{R}^\rho * \sigma\mu\nu\, V^\mu \odot J^\sigma \odot V^\nu.\end{aligned} \tag{6.60}$$

Step 3: Conclude the Jacobi equation. Insert (6.60) into (6.58):

$$\Big(\boldsymbol{\nabla}^{\mathrm{HS}} * V\, \boldsymbol{\nabla}^{\mathrm{HS}} * V\, J\Big)^\rho = -\,\boldsymbol{R}^\rho * \sigma\mu\nu\, V^\mu \odot J^\sigma \odot V^\nu,$$

which is equivalent to the claimed HS Jacobi word (6.55):

$$\Big(\boldsymbol{\nabla}^{\mathrm{HS}} * V\, \boldsymbol{\nabla}^{\mathrm{HS}} * V\, J\Big)^\rho \;+\; \boldsymbol{R}^\rho * \sigma\mu\nu\, V^\mu \odot J^\sigma \odot V^\nu \;=\; 0.$$

Remarks on conventions. If one instead adopts the curvature commutator convention $[\boldsymbol{\nabla}^{\mathrm{HS}} * V, \boldsymbol{\nabla}^{\mathrm{HS}} * J] = +\,\boldsymbol{R}(V, J)$ (cf. the alternative sign used in some derivations of eq. (6.37)), the result is the same up to $\boldsymbol{R} \mapsto -\boldsymbol{R}$; (6.55) is written with the sign choice fixed by (6.59). In either convention, the geometric content is identical. □

Proposition 6.22 (Reduction of HS curvature/Einstein/Jacobi)**.** *Under the Born map of §6.1.2 (consistent with Definitions 4.9 and 6.4), and using Proposition 6.15,*

$$\begin{aligned} \Big\langle\!\Big\langle \boldsymbol{R}^{\rho} * \sigma\mu\nu \Big\rangle\!\Big\rangle &= R^{\rho}{}_{\sigma\mu\nu}[\bar{\Sigma}], \\ \Big\langle\!\Big\langle \mathbf{Ric} * \mu\nu \Big\rangle\!\Big\rangle &= \mathrm{Ric}_{\mu\nu}[\bar{\Sigma}], \\ \Big\langle\!\Big\langle \mathcal{R} \Big\rangle\!\Big\rangle &= \mathcal{R}[\bar{\Sigma}], \\ \Big\langle\!\Big\langle \mathbf{G}^{\mu} * \nu \Big\rangle\!\Big\rangle &= \mathcal{G}^{\mu}{}_{\nu}[\bar{\Sigma}]. \end{aligned} \tag{6.61}$$

and the identities eqs. (6.30), (6.37), (6.46) and (6.55) reduce respectively to the classical first/second Bianchi identities, $\nabla^{(\mathrm{rel})}_{\mu}\mathcal{G}^{\mu}{}_{\nu} = 0$, and the Jacobi equation for $\bar{\Sigma}$ (see Definition 6.7 and eq. (6.22)).

Remark 6.23 (Gauge covariance)**.** All HS curvature words transform by wordwise conjugation $g(x)\big(\,\cdot\,\big)g(x)^{\ddagger}$, as inherited from the transporter law in eq. (4.16) and the HS unified correlator construction of §6.1.2; their Born reductions are gauge-invariant.

Inverse of the Chronogeometric Tensor

Definition 6.24 (Inverse of the Chronogeometric Tensor)**.** Let $\bar{\Sigma}_{\mu\nu}(x)$ be the expectation-level chronogeometric tensor defined in Definition 5.26 by

$$\bar{\Sigma}_{\mu\nu}(x) := \bar{g}_{\mu\nu}(x) + \bar{t}_{\mu\nu}(x),$$

where $\bar{g}_{\mu\nu}(x)$ and $\bar{t}_{\mu\nu}(x)$ are the expectation-level metric and temporal curvature tensors obtained from the operator-level constructions via the Born map (see Definitions 5.10 and 5.22). The clean sectoral split and its limiting cases are summarized in Corollary 5.29. For causal and variational constructions one may also use the Lorentzian merge $\bar{g}^{\mathrm{Lor}}_{\mu\nu} := \bar{g}_{\mu\nu} + \bar{t}^{\mathrm{sym}}_{\mu\nu}$ introduced in Definitions 5.77 and 5.122.

We define the *inverse* $\bar{\Sigma}^{\mu\nu}(x)$ as the unique smooth, symmetric $(2,0)$-tensor field satisfying

$$\boxed{\bar{\Sigma}^{\mu\rho}(x)\,\bar{\Sigma}_{\rho\nu}(x) \;=\; \delta^{\mu}{}_{\nu}.} \tag{6.62}$$

This inverse exists and is unique on any open region of $\mathcal{X}_q$ where $\bar{\Sigma}_{\mu\nu}(x)$ is non-degenerate; this is precisely the hypothesis needed to form the relational Christoffel symbols $\Gamma^{\lambda}{}_{\mu\nu}[\bar{\Sigma}]$ in eq. (6.22) and to use $\bar{\Sigma}^{\mu\nu}$ in the sector actions (see eq. (6.315)).

Holosymmetric wordwise inverse of the chronogeometric word

Definition 6.25 (Holosymmetric Wordwise Inverse of the Chronogeometric Word)**.** Let $\boldsymbol{\Sigma}_{\mu\nu}(x)$ denote the operator–valued *holosymmetric wordwise chronogeometric tensor* (see Sections 5.2.7 and 5.3.5), and let

$$\bar{\Sigma}_{\mu\nu}(x) = \Big\langle \overset{\circ}{\boldsymbol{\Psi}} \Big|\, \boldsymbol{\Sigma}_{\mu\nu}(x) \,\Big|\, \overset{\circ}{\boldsymbol{\Psi}} \Big\rangle$$

recover the standard expectation tensor of Definition 5.26. We say that $\mathbf{\Sigma}$ is *HS-nondegenerate* on an open set $U \subset \mathcal{X}_q$ if the wordwise Riesz map

$$\mathcal{R}_{\mathbf{\Sigma}} : \ \mathbf{V}^{\nu} \ \mapsto \ \mathbf{\Sigma}_{\mu\nu} \odot \mathbf{V}^{\nu}$$

is bijective in the holosymmetric (HS) word category. On such a region, the *holosymmetric wordwise inverse* $\mathbf{\Sigma}^{\mu\nu}(x)$ is the unique HS word, symmetric in (μ, ν), satisfying the two–sided inverse relations

$$\boxed{\mathbf{\Sigma}^{\mu\rho}(x) \ \odot \ \mathbf{\Sigma}_{\rho\nu}(x) \ = \ \delta^{\mu}{}_{\nu}\,\mathbf{1}, \qquad \mathbf{\Sigma}_{\mu\rho}(x) \ \odot \ \mathbf{\Sigma}^{\rho\nu}(x) \ = \ \delta_{\mu}{}^{\nu}\,\mathbf{1},} \tag{6.63}$$

where $\mathbf{1}$ is the HS identity word and $\delta^{\mu}{}_{\nu}$ is the Kronecker symbol. Existence/uniqueness of $\mathbf{\Sigma}^{\mu\nu}$ is equivalent to HS-nondegeneracy of $\mathbf{\Sigma}$.

Remark 6.26 (Reduction to the standard inverse)**.** Taking the Born expectation of (6.63) yields

$$\left\langle \mathring{\mathbf{\Psi}} \middle| \, \mathbf{\Sigma}^{\mu\rho} \odot \mathbf{\Sigma}_{\rho\nu} \, \middle| \mathring{\mathbf{\Psi}} \right\rangle = \bar{\Sigma}^{\mu\rho}\,\bar{\Sigma}_{\rho\nu} = \delta^{\mu}{}_{\nu},$$

so that $\langle \mathbf{\Sigma}^{\mu\nu} \rangle = \bar{\Sigma}^{\mu\nu}$ is precisely the *standard* inverse of Definition 6.24, i.e. it satisfies (6.62).

Proposition 6.27 (HS covariance, symmetry, and compatibility)**.** *On any HS–nondegenerate region $U \subset \mathcal{X}_q$, the inverse word $\mathbf{\Sigma}^{\mu\nu}$ enjoys:*

(1) Symmetry
If $\mathbf{\Sigma}_{\mu\nu} = \mathbf{\Sigma}_{\nu\mu}$ then $\mathbf{\Sigma}^{\mu\nu} = \mathbf{\Sigma}^{\nu\mu}$.

(2) Gauge covariance
Under endpoint gauge transformations $g(x)$,

$$\mathbf{\Sigma}_{\mu\nu} \mapsto g(x)\,\mathbf{\Sigma}_{\mu\nu}\,g(x)^{\dagger} \quad \Longrightarrow \quad \mathbf{\Sigma}^{\mu\nu} \mapsto g(x)\,\mathbf{\Sigma}^{\mu\nu}\,g(x)^{\dagger},$$

consistent with Section 4.5.

(3) Index operations
Raising/lowering with $\mathbf{\Sigma}^{\mu\nu}$ and $\mathbf{\Sigma}_{\mu\nu}$ uses HS composition $\odot$, e.g. $\mathbf{V}_{\mu} := \mathbf{\Sigma}_{\mu\nu} \odot \mathbf{V}^{\nu}$ and $\mathbf{W}^{\mu} := \mathbf{\Sigma}^{\mu\nu} \odot \mathbf{W}_{\nu}$.

(4) Compatibility with the HS Levi–Civita connection
The HS Christoffels of Definition 6.10 depend on $\mathbf{\Sigma}^{\mu\nu}$ as

$$\mathbf{\Gamma}^{\lambda} * \mu\nu[\mathbf{\Sigma}] = \tfrac{1}{2}\,\mathbf{\Sigma}^{\lambda\rho} \odot \big(\partial_{\mu}\mathbf{\Sigma}_{\nu\rho} + \partial_{\nu}\mathbf{\Sigma}_{\mu\rho} - \partial_{\rho}\mathbf{\Sigma}_{\mu\nu}\big),$$

so that the HS connection $\boldsymbol{\nabla}^{\mathrm{HS}}$ is torsionless and HS–metric–compatible ($\boldsymbol{\nabla}^{\mathrm{HS}} * \lambda\, \mathbf{\Sigma}_{\mu\nu} = 0$); see Section 6.1.4.

(5) Lorentzian merge (wordwise)
The wordwise symmetrisation $\boldsymbol{g}^{\mathrm{Lor}}_{\mu\nu} := \frac{1}{2}(\mathbf{\Sigma}_{\mu\nu} + \mathbf{\Sigma}_{\nu\mu})$ (see Sections 5.3.5 and 5.3.7) admits an HS inverse $(\boldsymbol{g}^{\mathrm{Lor}})^{\mu\nu}$ defined by (6.63); its expectation reduces to $\bar{g}^{\mathrm{Lor}\,\mu\nu}$.

Proof idea. (1) follows from uniqueness of the two–sided HS inverse when the underlying tensor is symmetric. (2) uses conjugation covariance and functoriality of $\odot$ under endpoint unitaries. (3) is immediate from the definition of the HS Riesz map and its inverse. (4) is the standard derivation of the wordwise Levi–Civita symbols from the HS Koszul identity (see Definition 6.8). (5) follows from functoriality of $\odot$ under the symmetriser and linearity of the Born expectation.

Proof of Proposition 6.27. Work on an HS–nondegenerate open region $U \subset \mathcal{X}_q$, so the Riesz map $\mathcal{R}_{\mathbf{\Sigma}} : \mathbf{V}^\nu \mapsto \mathbf{\Sigma}_{\mu\nu} \odot \mathbf{V}^\nu$ is bijective and the two–sided HS inverse $\mathbf{\Sigma}^{\mu\nu}$ exists by Definition 6.25, i.e.

$$\mathbf{\Sigma}^{\mu\rho} \odot \mathbf{\Sigma}_{\rho\nu} = \delta^\mu{}_\nu \, \mathbf{1}, \qquad \mathbf{\Sigma}_{\mu\rho} \odot \mathbf{\Sigma}^{\rho\nu} = \delta_\mu{}^\nu \, \mathbf{1}. \tag{6.64}$$

(1) Symmetry. Assume $\mathbf{\Sigma}_{\mu\nu} = \mathbf{\Sigma}_{\nu\mu}$ on U. Define the HS bilinear pairings (for contravariant words $\mathbf{V}^\mu, \mathbf{W}^\nu$ and covariant words $\mathbf{V}_\mu, \mathbf{W}_\nu$)

$$\mathsf{B}(\mathbf{V}, \mathbf{W}) := \mathbf{V}^\mu \odot \mathbf{\Sigma}_{\mu\nu} \odot \mathbf{W}^\nu, \qquad \mathsf{C}(\mathbf{V}, \mathbf{W}) := \mathbf{V}_\mu \odot \mathbf{\Sigma}^{\mu\nu} \odot \mathbf{W}_\nu.$$

By symmetry of $\mathbf{\Sigma}_{\mu\nu}$, $\mathsf{B}(\mathbf{V}, \mathbf{W}) = \mathsf{B}(\mathbf{W}, \mathbf{V})$. Using (6.64) and the definitions $\mathbf{V}_\mu = \mathbf{\Sigma}_{\mu\rho} \odot \mathbf{V}^\rho$, $\mathbf{W}_\nu = \mathbf{\Sigma}_{\nu\sigma} \odot \mathbf{W}^\sigma$, we have

$$\mathsf{C}(\mathbf{V}, \mathbf{W}) = \mathbf{V}_\mu \odot \mathbf{\Sigma}^{\mu\nu} \odot \mathbf{W}_\nu = \mathbf{V}^\rho \odot \Big(\mathbf{\Sigma}_{\mu\rho} \odot \mathbf{\Sigma}^{\mu\nu} \odot \mathbf{\Sigma}_{\nu\sigma} \Big) \odot \mathbf{W}^\sigma = \mathbf{V}^\rho \odot \mathbf{\Sigma}_{\rho\sigma} \odot \mathbf{W}^\sigma = \mathsf{B}(\mathbf{V}, \mathbf{W}),$$

hence C is also symmetric: $\mathsf{C}(\mathbf{V}, \mathbf{W}) = \mathsf{C}(\mathbf{W}, \mathbf{V})$. Let $\mathbf{A}^{\mu\nu} := \mathbf{\Sigma}^{\mu\nu} - \mathbf{\Sigma}^{\nu\mu}$. Then symmetry of C implies, for all $\mathbf{V}_\mu, \mathbf{W}_\nu$,

$$\mathbf{V}_\mu \odot \mathbf{A}^{\mu\nu} \odot \mathbf{W}_\nu = 0.$$

By HS–nondegeneracy (bijectivity of the Riesz map), this forces $\mathbf{A}^{\mu\nu} = 0$, i.e. $\mathbf{\Sigma}^{\mu\nu} = \mathbf{\Sigma}^{\nu\mu}$ on U.

(2) Gauge covariance. Under endpoint gauge transformations $g(x)$ we have conjugation covariance of HS words (cf. *wordwise covariant kinematics* in Section 4.5): $\mathbf{\Sigma}_{\mu\nu} \mapsto g \, \mathbf{\Sigma}_{\mu\nu} \, g^\dagger$. Let $\widetilde{\mathbf{\Sigma}}^{\mu\nu} := g \, \mathbf{\Sigma}^{\mu\nu} \, g^\dagger$. Then, using associativity of $\odot$ and $g^\dagger g = \mathbf{1}$,

$$\widetilde{\mathbf{\Sigma}}^{\mu\rho} \odot (g \, \mathbf{\Sigma}_{\rho\nu} \, g^\dagger) = g \Big(\mathbf{\Sigma}^{\mu\rho} \odot \mathbf{\Sigma}_{\rho\nu} \Big) g^\dagger = g \, \delta^\mu{}_\nu \, g^\dagger = \delta^\mu{}_\nu \, \mathbf{1},$$

and similarly $(g \, \mathbf{\Sigma}_{\mu\rho} \, g^\dagger) \odot \widetilde{\mathbf{\Sigma}}^{\rho\nu} = \delta_\mu{}^\nu \, \mathbf{1}$. By uniqueness of the two–sided HS inverse (Definition 6.25), the inverse transforms by conjugation: $\mathbf{\Sigma}^{\mu\nu} \mapsto g \, \mathbf{\Sigma}^{\mu\nu} \, g^\dagger$.

(3) Index operations. Define lowering and raising by $\mathbf{V}_\mu := \mathbf{\Sigma}_{\mu\nu} \odot \mathbf{V}^\nu$ and $\mathbf{W}^\mu := \mathbf{\Sigma}^{\mu\nu} \odot \mathbf{W}_\nu$. Then, using (6.64),

$$\begin{aligned} \mathbf{\Sigma}^{\mu\nu} \odot \mathbf{V}_\nu &= \mathbf{\Sigma}^{\mu\nu} \odot \mathbf{\Sigma}_{\nu\rho} \odot \mathbf{V}^\rho = \delta^\mu{}_\rho \, \mathbf{V}^\rho = \mathbf{V}^\mu, \\ \mathbf{\Sigma}_{\mu\nu} \odot \mathbf{W}^\nu &= \mathbf{\Sigma}_{\mu\nu} \odot \mathbf{\Sigma}^{\nu\rho} \odot \mathbf{W}_\rho = \delta_\mu{}^\rho \, \mathbf{W}_\rho = \mathbf{W}_\mu. \end{aligned} \tag{6.65}$$

so raising and lowering are mutually inverse via HS composition $\odot$.

(4) Compatibility with the HS Levi–Civita connection. By Definition 6.10, the (torsionless) HS Christoffel symbols are

$$\mathbf{\Gamma}^{\lambda} * \mu\nu[\mathbf{\Sigma}] = \tfrac{1}{2}\,\mathbf{\Sigma}^{\lambda\rho} \odot \big(\partial_\mu \mathbf{\Sigma}_{\nu\rho} + \partial_\nu \mathbf{\Sigma}_{\mu\rho} - \partial_\rho \mathbf{\Sigma}_{\mu\nu}\big),$$

which is the unique solution of the HS Koszul identity (see Definition 6.8) ensuring torsionlessness $\mathbf{\Gamma}^{\lambda} * \mu\nu = \mathbf{\Gamma}^{\lambda} * \nu\mu$ and metric-compatibility $\boldsymbol{\nabla}^{\mathrm{HS}} * \lambda \mathbf{\Sigma}_{\mu\nu} = 0$. The latter follows by direct substitution of the above $\mathbf{\Gamma}$ into the HS covariant derivative (cf. Section 6.1.4).

(5) Lorentzian merge and its inverse. Let $\boldsymbol{g}^{\mathrm{Lor}}_{\mu\nu} := \frac{1}{2}(\mathbf{\Sigma}_{\mu\nu} + \mathbf{\Sigma}_{\nu\mu})$ (cf. Sections 5.3.5 and 5.3.7). On U, the symmetriser is functorial and preserves HS–nondegeneracy (wordwise centrality/holonomy control on U), so $\boldsymbol{g}^{\mathrm{Lor}}$ is HS–nondegenerate and admits a unique HS inverse $(\boldsymbol{g}^{\mathrm{Lor}})^{\mu\nu}$ determined by (6.64) with $\mathbf{\Sigma} \rightsquigarrow \boldsymbol{g}^{\mathrm{Lor}}$, i.e. by (6.63). By the reduction statement for HS inverses (Remark 6.26), $\langle (\boldsymbol{g}^{\mathrm{Lor}})^{\mu\nu} \rangle = \bar{g}^{\mathrm{Lor}\,\mu\nu}$.

All five items follow, completing the proof. □

Corollary 6.28 (Sector actions and relational LC via HS inverse)**.** *In the sector actions of eq. (6.315), one may replace $\bar{\Sigma}^{\mu\nu}$ by the expectation of the HS inverse, $\langle \mathbf{\Sigma}^{\mu\nu} \rangle$, without changing dynamics. Likewise, the relational Christoffels $\Gamma^{\lambda}{}_{\mu\nu}[\bar{\Sigma}]$ of eq. (6.22) are the Born expectations of $\mathbf{\Gamma}^{\lambda} * \mu\nu[\mathbf{\Sigma}]$.*

6.1.5 The Determinant of the Effective Lorentzian Metric

The term $\det \bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)$ denotes the determinant of the matrix representation of the *Effective Lorentzian Metric* (Definition 5.77) at a specific point x in the emergent quantum configuration space, $\mathcal{X}_q$.

Matrix Representation of the Tensor

In any local coordinate system $(x^0, x^1, \ldots, x^{d-1})$ on the manifold $\mathcal{X}_q$, the rank-2 tensor $\bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)$ can be written as a $d \times d$ symmetric matrix:

$$[\bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)] = \begin{pmatrix} \bar{g}^{\mathrm{Lor}}_{00} & \bar{g}^{\mathrm{Lor}}_{01} & \cdots & \bar{g}^{\mathrm{Lor}}_{0,d-1} \\ \bar{g}^{\mathrm{Lor}}_{10} & \bar{g}^{\mathrm{Lor}}_{11} & \cdots & \bar{g}^{\mathrm{Lor}}_{1,d-1} \\ \vdots & \vdots & \ddots & \vdots \\ \bar{g}^{\mathrm{Lor}}_{d-1,0} & \bar{g}^{\mathrm{Lor}}_{d-1,1} & \cdots & \bar{g}^{\mathrm{Lor}}_{d-1,d-1} \end{pmatrix}$$

Due to the symmetry of the tensor, we have $\bar{g}^{\mathrm{Lor}}_{\mu\nu} = \bar{g}^{\mathrm{Lor}}_{\nu\mu}$.

The Determinant and its Geometric Significance

The determinant of this matrix, often written simply as $g^{\mathrm{Lor}}(x)$, is a scalar quantity calculated from its components. Formally, it is given by the Leibniz formula:

$$\det\big(\bar{g}^{\mathrm{Lor}}_{\mu\nu}\big) = \sum_{\sigma \in S_d} \operatorname{sgn}(\sigma) \prod_{i=0}^{d-1} \bar{g}^{\mathrm{Lor}}_{i,\sigma(i)} \tag{6.66}$$

where the sum is over all permutations σ of the indices.

Its physical significance is profound: it relates the coordinate volume to the true, physical invariant volume. An infinitesimal coordinate box with volume $d^d x = dx^0 dx^1 \dots dx^{d-1}$ does not represent a physical volume on a curved manifold. The determinant of the metric provides the necessary scaling factor, or Jacobian, to transform this coordinate-dependent volume into a coordinate-independent physical volume [7].

6.1.6 Holosymmetric Wordwise Determinant of the Effective Lorentzian Metric

Definition 6.29 (HS determinant of the effective Lorentzian metric)**.** Let $\boldsymbol{g}^{\mathrm{Lor}}_{\mu\nu}(x) := \frac{1}{2}(\boldsymbol{\Sigma}_{\mu\nu}(x) + \boldsymbol{\Sigma}_{\nu\mu}(x))$ be the holosymmetric wordwise effective Lorentzian metric (cf. Sections 5.3.5 and 5.3.7), whose Born expectation equals the standard effective metric $\bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)$ of Definition 5.77.

In any local chart $(x^0, \dots, x^{d-1})$ on $\mathcal{X}_q$, define the *holosymmetric determinant word* by the Levi–Civita contraction

$$\boxed{\det{}_{\mathrm{HS}}\big(\boldsymbol{g}^{\mathrm{Lor}}(x)\big) := \frac{1}{d!}\,\varepsilon^{\mu_1\cdots\mu_d}\,\varepsilon^{\nu_1\cdots\nu_d}\,\boldsymbol{g}^{\mathrm{Lor}}_{\mu_1\nu_1}(x)\ \odot\ \cdots\ \odot\ \boldsymbol{g}^{\mathrm{Lor}}_{\mu_d\nu_d}(x),} \tag{6.67}$$

where $\varepsilon^{\mu_1\cdots\mu_d}$ is the totally antisymmetric Levi–Civita symbol (with $\varepsilon^{0\,1\cdots d-1} = +1$ in the chosen orientation) and $\odot$ denotes the HS composition. The result is a scalar HS word (a density word of weight 2).

Remark 6.30 (Reduction to the standard determinant)**.** The Born expectation of (6.67) reproduces the standard determinant of Section 6.1.5:

$$\Big\langle \mathring{\boldsymbol{\Psi}}\Big|\ \det{}_{\mathrm{HS}}\ \Big(\boldsymbol{g}^{\mathrm{Lor}}(x)\Big)\ \Big|\mathring{\boldsymbol{\Psi}}\Big\rangle = \det\Big(\bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)\Big),$$

equivalently matching the Leibniz formula (6.66).

Proposition 6.31 (Gauge covariance and centrality)**.** *Under endpoint gauge transformations $g(x)$, the metric word transforms by conjugation $\boldsymbol{g}^{\mathrm{Lor}}_{\mu\nu} \mapsto g(x)\,\boldsymbol{g}^{\mathrm{Lor}}_{\mu\nu}\,g(x)^\dagger$, hence*

$$\det{}_{\mathrm{HS}}\ \Big(\boldsymbol{g}^{\mathrm{Lor}}\Big)\ \mapsto\ g(x)\ \det{}_{\mathrm{HS}}\ \Big(\boldsymbol{g}^{\mathrm{Lor}}\Big)\ g(x)^\dagger.$$

In flat/central patches (holonomy trivial), $\det_{\mathrm{HS}}(\boldsymbol{g}^{\mathrm{Lor}})$ lies in the center of the HS algebra; its Born expectation is gauge invariant.

Proof of Proposition 6.31. We work purely in the holosymmetric (HS) wordwise calculus of Chs. 4–6.

(1) Gauge covariance of $\boldsymbol{g}^{\mathrm{Lor}}$ and of $\det_{\mathrm{HS}}(\boldsymbol{g}^{\mathrm{Lor}})$. By endpoint gauge covariance of HS words (wordwise parallel transport/kinematics), the effective HS Lorentzian metric transforms at $x \in \mathcal{X}_q$ by conjugation (cf. *Holosymmetric wordwise covariant kinematics* and the construction of $\boldsymbol{g}^{\mathrm{Lor}}$ in Sections 5.3.5 and 5.3.7):

$$\boldsymbol{g}^{\mathrm{Lor}}_{\mu\nu}(x)\ \longmapsto\ g(x)\,\boldsymbol{g}^{\mathrm{Lor}}_{\mu\nu}(x)\,g(x)^\dagger. \tag{6.68}$$

Recall the HS determinant definition (Definition 6.29):

$$\det_{\mathrm{HS}}\left(\boldsymbol{g}^{\mathrm{Lor}}\right)=\frac{1}{d!}\,\varepsilon^{\mu_1\cdots\mu_d}\,\varepsilon^{\nu_1\cdots\nu_d}\,\boldsymbol{g}^{\mathrm{Lor}}_{\mu_1\nu_1}\;\odot\;\cdots\;\odot\;\boldsymbol{g}^{\mathrm{Lor}}_{\mu_d\nu_d},\tag{6.69}$$

where $\odot$ is the HS composition and the Levi–Civita symbols are c-numbers (cf. Definition 6.29 and eq. (6.67)). Applying (6.68) factor-by-factor and using associativity of $\odot$ together with unitarity $g^\dagger\odot g=\mathbf{1}$, the interstitial $g^\dagger g$ cancel, giving

$$\begin{aligned}\det_{\mathrm{HS}}\left(g\,\boldsymbol{g}^{\mathrm{Lor}}\,g^\dagger\right)&=\frac{1}{d!}\,\varepsilon^{\mu_1\cdots\mu_d}\varepsilon^{\nu_1\cdots\nu_d}\,(g\,\boldsymbol{g}^{\mathrm{Lor}}_{\mu_1\nu_1}g^\dagger)\odot\cdots\odot(g\,\boldsymbol{g}^{\mathrm{Lor}}_{\mu_d\nu_d}g^\dagger)\\&=g\,\left[\frac{1}{d!}\,\varepsilon^{\mu_1\cdots\mu_d}\varepsilon^{\nu_1\cdots\nu_d}\,\boldsymbol{g}^{\mathrm{Lor}}_{\mu_1\nu_1}\odot\cdots\odot\boldsymbol{g}^{\mathrm{Lor}}_{\mu_d\nu_d}\right]\,g^\dagger\\&=g\;\det_{\mathrm{HS}}\left(\boldsymbol{g}^{\mathrm{Lor}}\right)\,g^\dagger,\end{aligned}\tag{6.70}$$

which is the claimed covariance.

(2) Centrality on flat/central patches. From the *centralizing role of holosymmetry* (Ch. 4; see Section 4.4 and the holonomy/centrality bounds), commutator defects of wordwise letters built from gauge-covariant local blocks are controlled by holonomy. On a flat/central patch (trivial holonomy), these defects vanish: the elementary letters are effectively central in the HS algebra. Since $\det_{\mathrm{HS}}(\boldsymbol{g}^{\mathrm{Lor}})$ is a c-number Levi–Civita contraction of a finite $\odot$-polynomial in the (now) central letters $\boldsymbol{g}^{\mathrm{Lor}}_{\mu\nu}$, it follows that

$$\left[\det_{\mathrm{HS}}\left(\boldsymbol{g}^{\mathrm{Lor}}\right),\;\mathbf{Z}\right]=0\qquad\text{for all HS words }\mathbf{Z}\text{ on the patch.}\tag{6.71}$$

Consequently $\det_{\mathrm{HS}}(\boldsymbol{g}^{\mathrm{Lor}})$ is invariant under conjugation by $g(x)$ on the patch and lies in the center of the HS algebra there.

(3) Gauge-invariant Born expectation. By centrality (6.71) and (6.70),

$$\left\langle\mathring{\boldsymbol{\Psi}}\right|\,g\,\underset{\mathrm{HS}}{\det}(\boldsymbol{g}^{\mathrm{Lor}})\,g^\dagger\,\left|\mathring{\boldsymbol{\Psi}}\right\rangle=\left\langle\mathring{\boldsymbol{\Psi}}\right|\,\underset{\mathrm{HS}}{\det}(\boldsymbol{g}^{\mathrm{Lor}})\,\left|\mathring{\boldsymbol{\Psi}}\right\rangle.$$

Moreover, by the reduction statement (Remark 6.30),

$$\left\langle\mathring{\boldsymbol{\Psi}}\right|\,\det_{\mathrm{HS}}\left(\boldsymbol{g}^{\mathrm{Lor}}\right)\left|\mathring{\boldsymbol{\Psi}}\right\rangle=\det\left(\bar{g}^{\mathrm{Lor}}_{\mu\nu}\right),\tag{6.72}$$

and the expectation-level effective metric $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ is word/frame independent (cf. the expectation-level wordwise constructions in Section 5.1.5). Therefore the scalar (6.72) is gauge invariant.

Combining (1)–(3) proves the proposition. □

Definition 6.32 (HS volume density word and volume form)**.** In a Lorentzian sector with signature $(+,-,\ldots,-)$, define the *HS volume density word* by the functional (spectral) calculus on the regulated fibre

$$\boxed{\boldsymbol{\varpi}(x)\;:=\;\left(\,\det_{\mathrm{HS}}\left(-\,\boldsymbol{g}^{\mathrm{Lor}}(x)\right)\right)^{1/2},}\tag{6.73}$$

and the associated HS volume form word $d\boldsymbol{V} := \boldsymbol{\varpi}(x)\, d^d x$. Its Born expectation reproduces the standard invariant measure:

$$\left\langle \mathring{\boldsymbol{\Psi}} \right| \boldsymbol{\varpi}(x) \left| \mathring{\boldsymbol{\Psi}} \right\rangle = \sqrt{-\det\left(\bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)\right)}, \qquad \left\langle \mathring{\boldsymbol{\Psi}} \right| d\boldsymbol{V} \left| \mathring{\boldsymbol{\Psi}} \right\rangle = \sqrt{-\det\left(\bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)\right)}\, d^d x.$$

The Role of the Lorentzian Signature

Crucially, for any metric with a *Lorentzian signature* (e.g., $(+,-,-,-)$ in $d = 4$), the determinant is always *negative*. This is because one eigenvalue of the metric matrix has a different sign from the others.

- This means the quantity $-\det \bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)$ is positive.
- Taking the square root, $\sqrt{-\det \bar{g}^{\mathrm{Lor}}_{\mu\nu}}$, yields a real, positive scalar.

This scalar is the precise factor needed to define the invariant volume element, $d\mu$, which is essential for constructing action integrals that are independent of the observer's choice of coordinates:

$$d\mu(x) = \sqrt{-\det \bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)}\, d^d x.$$

This ensures that the laws of physics derived from the action principle are truly universal.

Relational Volume Element

Definition 6.33 (Relational Volume Element)**.** Let $\bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)$ be the effective Lorentzian metric as defined in Definition 5.77. The *Relational Volume Element* at a point $x \in \mathcal{X}_q$ is the invariant measure on the emergent manifold, given by:

$$\boxed{d\mu(x) := d^d x \sqrt{-\det \bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)},} \tag{6.74}$$

where:

- $d^d x$ is the coordinate volume element in a local chart of $\mathcal{X}_q$.
- $\det \bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)$ denotes the determinant of the effective Lorentzian metric.
- The minus sign under the square root ensures the volume element is real for a metric with Lorentzian signature $(+,-,-,-)$.

In any theory of spacetime, one must establish a proper, coordinate-independent measure of volume. This mathematical tool is essential for defining covariant integrals, which are necessary to calculate total physical quantities such as the Action Functional [7]. In our framework, this is achieved through the *Relational Volume Element*, $d\mu(x)$, which is derived from the *Effective Lorentzian Metric*, $\bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)$ (Definition 6.33).

While the volume element itself is defined from the metric, that metric is a composite object, $\bar{g}^{\mathrm{Lor}}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}^{\mathrm{sym}}_{\mu\nu}$ (Definition 5.77). This structure reveals a novel aspect of our emergent reality: the fundamental measure of "how much" spacetime exists in a given region

depends not only on the purely geometric curvature encoded in $\bar{g}_{\mu\nu}$, but also on the symmetric part of the temporal phase-flow structure, $\bar{t}^{\mathrm{sym}}_{\mu\nu}$.

This has several radical implications that distinguish our theory from General Relativity. Spacetime volume is no longer a passive, static property determined by the metric alone; it is dynamically influenced by the quantum phase relationships within it. This suggests that in extreme regimes—near black hole horizons [86, 87, 245], in the very early universe [88, 89], or in regions of high entanglement density [1]—where the temporal tensor $\bar{t}_{\mu\nu}$ is significant, our understanding of physical laws that depend on volume, such as entropy and energy density, may require modification.

Ultimately, the complete coupling of dynamics to the full chronogeometric texture is realized through the *Relational Levi-Civita Connection* (Definition 6.7). This connection, used in the kinetic terms of the action, depends on the entire unified tensor $\bar{\Sigma}_{\mu\nu}$, ensuring that the motion of all fields is fundamentally coupled to the deep temporal texture of reality.

6.1.7 Holosymmetric Wordwise Relational Volume Element

Definition 6.34 (HS Relational Volume Element)**.** Let $\boldsymbol{g}^{\mathrm{Lor}}_{\mu\nu}(x) = \frac{1}{2}(\boldsymbol{\Sigma}_{\mu\nu}(x) + \boldsymbol{\Sigma}_{\nu\mu}(x))$ be the holosymmetric wordwise effective Lorentzian metric (cf. Sections 5.3.5 and 5.3.7), and let $\boldsymbol{\varpi}(x)$ be the HS volume density word defined in Definition 6.32 and eq. (6.73) by

$$\boldsymbol{\varpi}(x) \;:=\; \Big(\det{}_{\mathrm{HS}} \Big(- \boldsymbol{g}^{\mathrm{Lor}}(x) \Big) \Big)^{1/2}.$$

The *HS Relational Volume Element* on a local chart $(x^0, \dots, x^{d-1})$ of $\mathcal{X}_q$ is the HS measure word

$$\boxed{ d\boldsymbol{\mu}(x) \;:=\; d^d x \; \boldsymbol{\varpi}(x) \;=\; d^d x \, \Big(\det{}_{\mathrm{HS}} \Big(- \boldsymbol{g}^{\mathrm{Lor}}(x) \Big) \Big)^{1/2}, } \tag{6.75}$$

with orientation and time-orientation fixed once per chart.

Remark 6.35 (Reduction to the standard relational volume element)**.** Taking the Born expectation in the Holo–State yields

$$\Big\langle \mathring{\boldsymbol{\Psi}} \Big| \, d\boldsymbol{\mu}(x) \, \Big| \mathring{\boldsymbol{\Psi}} \Big\rangle = d^d x \, \sqrt{ - \det\big(\bar{g}^{\mathrm{Lor}}_{\mu\nu}(x) \big) } = d\mu(x),$$

which is exactly the invariant measure of Definition 6.33, i.e. (6.74).

Proposition 6.36 (Gauge covariance and centrality of the HS measure)**.** *Under endpoint gauge transformations $g(x)$, one has $\boldsymbol{g}^{\mathrm{Lor}}_{\mu\nu} \mapsto g(x)\, \boldsymbol{g}^{\mathrm{Lor}}_{\mu\nu}\, g(x)^\dagger$, hence*

$$d\boldsymbol{\mu}(x) \;\mapsto\; d^d x \, \Big(\det{}_{\mathrm{HS}} \Big(- \, g \, \boldsymbol{g}^{\mathrm{Lor}} \, g^\dagger \Big) \Big)^{1/2} \;=\; g(x)\, d\boldsymbol{\mu}(x)\, g(x)^\dagger.$$

In flat/central patches (trivial holonomy), $d\boldsymbol{\mu}(x)$ lies in the center of the HS algebra, so its Born expectation is gauge invariant and equals $d\mu(x)$.

Proof of Proposition 6.36. We use only the holosymmetric (HS) wordwise calculus and results from Chs. 4–6.

(i) Gauge covariance of $\boldsymbol{g}^{\mathrm{Lor}}$ and of the HS determinant. By endpoint gauge covariance of HS words (cf. Section 4.5 and the covariance laws for wordwise blocks), the HS effective Lorentzian metric transforms at a point $x \in \mathcal{X}_q$ by conjugation:

$$\boldsymbol{g}^{\mathrm{Lor}}_{\mu\nu}(x) \;\longmapsto\; g(x)\,\boldsymbol{g}^{\mathrm{Lor}}_{\mu\nu}(x)\,g(x)^{\dagger}. \tag{6.76}$$

Recall the HS determinant word (Definition Definition 6.29):

$$\det_{\mathrm{HS}}\!\left(\boldsymbol{g}^{\mathrm{Lor}}\right) = \frac{1}{d!}\,\varepsilon^{\mu_1\cdots\mu_d}\,\varepsilon^{\nu_1\cdots\nu_d}\,\boldsymbol{g}^{\mathrm{Lor}}_{\mu_1\nu_1} \;\odot\; \cdots \;\odot\; \boldsymbol{g}^{\mathrm{Lor}}_{\mu_d\nu_d}, \tag{6.77}$$

where $\odot$ is the HS composition and the Levi–Civita symbols are c-numbers. Applying (6.76) factor-by-factor and using associativity of $\odot$ together with unitarity ($g^{\dagger}\odot g = \mathbf{1}$) collapses the interstitial $g^{\dagger}\odot g$ pairs:

$$\begin{aligned}
\det_{\mathrm{HS}}\!\left(g\,\boldsymbol{g}^{\mathrm{Lor}}\,g^{\dagger}\right) &= \frac{1}{d!}\,\varepsilon^{\mu_1\cdots\mu_d}\varepsilon^{\nu_1\cdots\nu_d}\,(g\boldsymbol{g}^{\mathrm{Lor}}_{\mu_1\nu_1}g^{\dagger}) \odot \cdots \odot (g\boldsymbol{g}^{\mathrm{Lor}}_{\mu_d\nu_d}g^{\dagger}) \\
&= g\,\left[\frac{1}{d!}\,\varepsilon^{\mu_1\cdots\mu_d}\varepsilon^{\nu_1\cdots\nu_d}\,\boldsymbol{g}^{\mathrm{Lor}}_{\mu_1\nu_1} \odot \cdots \odot \boldsymbol{g}^{\mathrm{Lor}}_{\mu_d\nu_d}\right] g^{\dagger} \\
&= g\;\det_{\mathrm{HS}}\!\left(\boldsymbol{g}^{\mathrm{Lor}}\right)\,g^{\dagger}.
\end{aligned} \tag{6.78}$$

Thus the HS determinant is gauge covariant by conjugation.

(ii) Gauge covariance of the HS volume density and of the HS measure. By Definition 6.32 and eq. (6.73),

$$\boldsymbol{\varpi}(x) \;:=\; \left(\,\det_{\mathrm{HS}}\left(-\,\boldsymbol{g}^{\mathrm{Lor}}(x)\right)\right)^{1/2}.$$

The functional calculus is stable under $*$–conjugation, hence from (6.78):

$$\boldsymbol{\varpi}(x) \;\longmapsto\; \left(\det_{\mathrm{HS}}\left(-\,g\,\boldsymbol{g}^{\mathrm{Lor}}\,g^{\dagger}\right)\right)^{1/2} = g(x)\;\boldsymbol{\varpi}(x)\;g(x)^{\dagger}. \tag{6.79}$$

Therefore the HS measure word (Definition Definition 6.34)

$$d\boldsymbol{\mu}(x) \;:=\; d^d x\;\boldsymbol{\varpi}(x)$$

transforms as

$$d\boldsymbol{\mu}(x) \;\longmapsto\; d^d x\,\left(\det_{\mathrm{HS}}\left(-\,g\,\boldsymbol{g}^{\mathrm{Lor}}\,g^{\dagger}\right)\right)^{1/2} = g(x)\,d\boldsymbol{\mu}(x)\,g(x)^{\dagger}, \tag{6.80}$$

which is the claimed covariance law.

(iii) Centrality in flat/central patches and gauge-invariant reduction. From the *centralizing role of holosymmetry* (Ch. 4) and the holonomy/centrality bound (cf. the curvature–holonomy estimate and its flat-patch corollary), commutator defects of HS words built from gauge-covariant local contractions are bounded by holonomy. On flat/central patches (trivial holonomy), those defects vanish: the elementary letters are effectively central, hence any

polynomial (in $\odot$) with c-number coefficients—such as $\det_{\mathrm{HS}}(\boldsymbol{g}^{\mathrm{Lor}})$—is central. By (6.79), $\boldsymbol{\varpi}$ is then central, and so is $d\boldsymbol{\mu}(x) = d^d x\, \boldsymbol{\varpi}(x)$:

$$[d\boldsymbol{\mu}(x),\, \mathbf{Z}] = 0 \qquad \text{for all HS words } \mathbf{Z} \text{ on the patch.} \tag{6.81}$$

Consequently $d\boldsymbol{\mu}(x)$ is invariant under conjugation by $g(x)$, and its Born expectation is gauge invariant. Using the reduction statement of Remark 6.35 we obtain

$$\left\langle \mathring{\boldsymbol{\Psi}} \right| d\boldsymbol{\mu}(x) \left| \mathring{\boldsymbol{\Psi}} \right\rangle = d^d x \sqrt{-\det \bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)} = d\mu(x),$$

the standard relational volume element of Definition 6.33.

This completes the proof. □

Corollary 6.37 (HS-covariant integration and sector actions)**.** *For any HS word observable $\mathbf{W}(x)$ with well-defined Born expectation, the HS integral $\int d\boldsymbol{\mu}(x)\ \mathbf{W}(x)$ is well-defined on spectrally regulated fibres, and its expectation reproduces the standard covariant integral:*

$$\left\langle \mathring{\boldsymbol{\Psi}} \right| \int d\boldsymbol{\mu}(x)\ \mathbf{W}(x) \left| \mathring{\boldsymbol{\Psi}} \right\rangle = \int d\mu(x)\ \langle \mathbf{W}(x) \rangle.$$

In particular, replacing $d\mu(x)$ by $\langle d\boldsymbol{\mu}(x) \rangle$ in the sector actions (eq. (6.315)) does not change the dynamics.

Proof of Corollary 6.37. We work strictly in the holosymmetric (HS) wordwise framework of Chs. 4–6 and use spectral regulation on fibres.

1) Well-posedness of the HS integral. By Definition Definition 6.34, $d\boldsymbol{\mu}(x) = d^d x\, \boldsymbol{\varpi}(x)$ with $\boldsymbol{\varpi}(x) = \left(\det_{\mathrm{HS}}(-\boldsymbol{g}^{\mathrm{Lor}}(x)) \right)^{1/2}$, where $\boldsymbol{g}^{\mathrm{Lor}}_{\mu\nu}$ is Hermitian and spectrally regulated. Hence $\|\boldsymbol{\varpi}(x)\| \le C_\Lambda$ on each regulated patch. For an HS word observable $\mathbf{W}(x)$ with well-defined Born expectation and $\|\mathbf{W}(x)\| \in L^1_{\mathrm{loc}}(\mathcal{X}_q)$ relative to $d^d x$, the Bochner (strong-operator) integral

$$\int_{U \subset \mathcal{X}_q} d\boldsymbol{\mu}(x)\, \mathbf{W}(x) \;=\; \int_U d^d x\, \left(\boldsymbol{\varpi}(x) \odot \mathbf{W}(x) \right)$$

exists and defines an HS word (by standard norm estimates $\| \int \cdot \| \le \int \| \cdot \|$).

2) Interchanging Born expectation and integration. Let $\mathcal{E}(\cdot) := \left\langle \mathring{\boldsymbol{\Psi}} \right| (\cdot) \left| \mathring{\boldsymbol{\Psi}} \right\rangle$ denote the Born expectation (linear, positive, and bounded by operator norm on spectrally regulated fibres). By dominated-convergence/Bochner interchange (the integrand norm is dominated by $\|\boldsymbol{\varpi}(x)\|\, \|\mathbf{W}(x)\| \in L^1_{\mathrm{loc}}$), we obtain

$$\mathcal{E}\left[\int d\boldsymbol{\mu}(x)\ \mathbf{W}(x) \right] = \int d^d x\ \mathcal{E}[\boldsymbol{\varpi}(x) \odot \mathbf{W}(x)]\,. \tag{6.82}$$

3) Pointwise evaluation of the integrand expectation. From Proposition 6.36 we have: (i) the conjugation law for the HS determinant and its square root, and (ii) centrality of $d\boldsymbol{\mu}(x)$ (equivalently of $\boldsymbol{\varpi}(x)$) on flat/central patches. Moreover, by the Chapter 4 centrality statements (holonomy bounds and blockwise centrality of the device/dialectic sector), $\boldsymbol{\varpi}(x)$ lies in the device/dialectic factor and is block-central with respect to field-space blocks. Therefore, on each spectrally regulated chart U we can compute (using cyclicity of the trace on the device/dialectic factor and blockwise centrality)

$$\mathcal{E}[\boldsymbol{\varpi}(x) \odot \mathbf{W}(x)] = \mathcal{E}[\boldsymbol{\varpi}(x)]\ \mathcal{E}[\mathbf{W}(x)] = \sqrt{-\det \bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)}\ \langle \mathbf{W}(x) \rangle, \qquad x \in U, \tag{6.83}$$

where we used the reduction identity of Remark 6.35: $\mathcal{E}[\boldsymbol{\varpi}(x)] = \sqrt{-\det \bar{g}^{\mathrm{Lor}}(x)}$. Cover $\mathcal{X}_q$ with such charts and invoke a partition of unity to extend (6.83) globally (the identity is local and hence compatible with partitions).

4) Conclusion. Combining (6.82) and (6.83) yields

$$\left\langle \overset{\circ}{\boldsymbol{\Psi}} \right| \int d\boldsymbol{\mu}(x)\ \mathbf{W}(x) \left| \overset{\circ}{\boldsymbol{\Psi}} \right\rangle = \int d^d x\ \sqrt{-\det \bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)}\ \langle \mathbf{W}(x) \rangle = \int d\mu(x)\ \langle \mathbf{W}(x) \rangle,$$

which is the stated identity.

5) Sector actions. In the sector actions (cf. (6.315)), each integrand is of the form $\mathrm{Tr}\big(\cdots\big)$ applied to HS word observables with well-defined Born expectations. Replacing $d\mu(x)$ by $\langle d\boldsymbol{\mu}(x) \rangle$ multiplies the same scalar density $\sqrt{-\det \bar{g}^{\mathrm{Lor}}}$ into the integrand; by the equality above, the resulting Euler–Lagrange equations are unchanged. Hence the dynamics are unaffected by the replacement.

This proves the corollary. □

6.1.8 The Vacuum Manifold: Variational and Spectral Formulation

The vacuum manifold is the set of quantum states that yield the most symmetric, stable, and least complex relational geometry. This structure emerges from a unification of two mathematically rigorous perspectives:

- A *variational formulation*, in which the ground state minimizes a scalar functional encoding the total geometric-relational complexity,
- A *spectral formulation*, in which the ground state is the lowest eigenstate of a composite Hamiltonian associated with inter-modal correlations.

Definition 6.38 (Relational Action Functional)**.** Let $|\boldsymbol{\Psi}\rangle \in \mathcal{H}_{\mathrm{em}}$ be a physically admissible quantum state in the total emergent Hilbert space (Definition 4.287),

$$\mathcal{H}_{\mathrm{em}} := \bigsqcup_{x \in \mathcal{X}_q} \mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}}.$$

Let $\hat{\Sigma}_{\mu\nu}(x)$ denote the operator-valued chronogeometric tensor field (Definition 5.25), and let $\hat{\mathbf{G}}_\Lambda(x, x')$ denote the Unified Inter-Modal Correlator (Definition 6.4).

Then the *state-dependent expectation-level chronogeometric tensor* is defined pointwise on $\mathcal{X}_q$ by:

$$\bar{\Sigma}^{(\Psi)}_{\mu\nu}(x) := \langle \Psi | \hat{\Sigma}_{\mu\nu}(x) | \Psi \rangle.$$

The *Relational Action Functional* associated to the state $|\Psi\rangle$ is then given by:

$$\boxed{S[|\Psi\rangle] := \int_{\mathcal{X}_q} d^d x \, \sqrt{-\det \bar{\Sigma}^{(\Psi)}_{\mu\nu}(x)} \, \mathrm{Tr}\left(\langle \Psi | \hat{\mathbf{G}}_\Lambda(x, x) | \Psi \rangle \right).} \tag{6.84}$$

This functional measures the total pointwise self-correlation of the generalized field operator $\hat{\mathbf{F}}(x)$ in the state $|\Psi\rangle$, integrated over the emergent configuration space $\mathcal{X}_q$ with respect to the *Relational Volume Element* (Definition 6.33).

The integrand captures the complete relational curvature at each point $x \in \mathcal{X}_q$, including:

- The curvature of *distinguishability*, via $\bar{g}^{(\Psi)}_{\mu\nu}(x) := \langle \Psi | \hat{g}_{\mu\nu}(x) | \Psi \rangle$,
- The curvature of *quantum phase flow*, via $\bar{t}^{(\Psi)}_{\mu\nu}(x) := \langle \Psi | \hat{t}_{\mu\nu}(x) | \Psi \rangle$,

so that:

$$\bar{\Sigma}^{(\Psi)}_{\mu\nu}(x) = \bar{g}^{(\Psi)}_{\mu\nu}(x) + \bar{t}^{(\Psi)}_{\mu\nu}(x).$$

In the special case where $|\Psi\rangle = |\mathring{\Psi}\rangle$, the universal vacuum state, this definition reduces to the canonical tensor $\bar{\Sigma}_{\mu\nu}(x)$ (Definition 5.26).

Critical points of the functional $S[|\Psi\rangle]$ define the Vacuum Manifold (Definition 6.61), characterizing globally minimal relational curvature and maximal coherence.

Two-Level Interpretation.

We must therefore distinguish between:

1. The General Relational Action Functional.

Defined over arbitrary physically admissible states $|\Psi\rangle$, as in Definition 6.38:

$$S[|\Psi\rangle] := \int_{\mathcal{X}_q} d^d x \, \sqrt{-\det \bar{\Sigma}^{(\Psi)}(x)} \, \mathrm{Tr}\left(\langle \Psi | \hat{\mathbf{G}}_\Lambda(x, x) | \Psi \rangle \right),$$

with $\bar{\Sigma}^{(\Psi)}_{\mu\nu}(x) := \langle \Psi | \hat{\Sigma}_{\mu\nu}(x) | \Psi \rangle$.

This allows us to formulate the variational principle:

$$\delta S[|\Psi\rangle] = 0,$$

and thereby define the *Vacuum Manifold* as the set of critical states $|\Psi\rangle_{\text{vac}}$ (Definition 6.61).

2. Evaluation on the Quantum Holo-State.

Once this structure is established, one may define the special evaluation:

$$S[|\mathring{\Psi}\rangle] := \int_{\mathcal{X}_q} d^d x \, \sqrt{-\det \bar{\Sigma}(x)} \, \mathrm{Tr}\left(\langle \mathring{\Psi} | \hat{\mathbf{G}}_\Lambda(x, x) | \mathring{\Psi} \rangle \right),$$

where $\bar{\Sigma}_{\mu\nu}(x) := \langle \mathring{\Psi} | \hat{\Sigma}_{\mu\nu}(x) | \mathring{\Psi} \rangle$ is the canonical vacuum geometry (Definition 5.26).

This represents the *realized vacuum state* of the cosmos, furnishing the background against which all local fluctuations, particles, and interactions are defined.

The *Relational Action Functional* defined in Definition 6.38 provides a *general variational principle* that is applicable to *each constituent field* (Gaiamiti, Tantrakyma) individually, as well as to their unified configuration via the generalized field operator $\hat{\mathbf{F}}(x)$. Let us briefly justify this from both formal and physical standpoints.

1. Structural Universality of the Unified Correlator The unified correlator

$$\hat{\mathbf{G}}_\Lambda(x,x') := \begin{pmatrix} \hat{G}_{\mathbf{\Phi\Phi}}(x,x') & \hat{G}_{\mathbf{\Phi\Theta}}(x,x') \\ \hat{G}_{\mathbf{\Theta\Phi}}(x,x') & \hat{G}_{\mathbf{\Theta\Theta}}(x,x') \end{pmatrix}$$

contains *all sectoral correlations* as special cases:

- Intra-field geometry: $\hat{G}_{\mathbf{\Phi\Phi}}(x,x')$,
- Intra-field matter: $\hat{G}_{\mathbf{\Theta\Theta}}(x,x')$,
- Inter-field couplings: $\hat{G}_{\mathbf{\Phi\Theta}}(x,x')$, $\hat{G}_{\mathbf{\Theta\Phi}}(x,x')$.

Each of these matrix entries is itself a 4×4 matrix in the dialectic modal basis, which allows one to isolate and analyze:

- The *geometry-only* dynamics by evaluating the action with $\hat{\mathbf{F}} = \hat{\mathbf{\Phi}}$ and truncating to $\hat{G}_{\mathbf{\Phi\Phi}}$,
- The *matter-only* dynamics analogously from $\hat{G}_{\mathbf{\Theta\Theta}}$,
- The *coupling terms* through the off-diagonal correlators.

2. Field-Specific Reduction of the Action Functional By projecting onto the respective subfields, one can define:

- A *Gaiamitiic Action Functional*:

$$S_{\mathbf{\Phi}}[|\mathbf{\Psi}\rangle] := \int_{\mathcal{X}_q} d^d x \, \sqrt{-\det \bar{\Sigma}(x)} \, \mathrm{Tr}\left(\langle \mathbf{\Psi}|\hat{G}_{\mathbf{\Phi\Phi}}(x,x)|\mathbf{\Psi}\rangle\right)$$

- A *Tantrakyma Action Functional*:

$$S_{\mathbf{\Theta}}[|\mathbf{\Psi}\rangle] := \int_{\mathcal{X}_q} d^d x \, \sqrt{-\det \bar{\Sigma}(x)} \, \mathrm{Tr}\left(\langle \mathbf{\Psi}|\hat{G}_{\mathbf{\Theta\Theta}}(x,x)|\mathbf{\Psi}\rangle\right)$$

- An *Interaction Action Functional*, defined, for example, as:

$$S_{\mathbf{\Phi\Theta}}[|\mathbf{\Psi}\rangle] := \int_{\mathcal{X}_q} d^d x \, \sqrt{-\det \bar{\Sigma}(x)} \, \mathrm{Tr}\left(\langle \mathbf{\Psi}| \left[\hat{G}_{\mathbf{\Phi\Theta}}(x,x) + \hat{G}_{\mathbf{\Theta\Phi}}(x,x)\right] |\mathbf{\Psi}\rangle\right)$$

This decomposition respects the structure of $\hat{\mathbf{G}}_\Lambda$ and enables the isolation of dynamical contributions from each field sector.

3. Physical Interpretation From a physical perspective:

- The *Gaiamitiic sector* governs the emergence of geometry (metric and causal structure).
- The *Tantrakyma sector* governs matter content, mass generation, and internal symmetries.
- The *off-diagonal interactions* encode the quantum analog of gravitational coupling between matter and spacetime, i.e., geometry influencing matter and vice versa.

Thus, each sector's variational dynamics is encoded within the general action functional $S[|\mathbf{\Psi}\rangle]$, which serves as a *unified variational umbrella* encompassing all sectoral dynamics.

6.1.9 Holosymmetric Wordwise Formulation of the Vacuum Manifold

Data and references. Let $\mathbf{\Sigma}_{\mu\nu}(x)$ be the HS operator–valued chronogeometric word and $\boldsymbol{g}^{\mathrm{Lor}}_{\mu\nu}(x) := \frac{1}{2}(\mathbf{\Sigma}_{\mu\nu}(x) + \mathbf{\Sigma}_{\nu\mu}(x))$ its HS Lorentzian symmetrisation (cf. Sections 5.3.5 and 5.3.7). Let $d\boldsymbol{\mu}(x)$ denote the HS volume element defined in Section 6.1.7, Definition 6.34, and eq. (6.75) via the HS determinant Definition 6.29 and eq. (6.67). Let $\hat{\mathbf{G}}_\Lambda(x,x')$ be the (operator–valued) unified inter–modal correlator matrix (wordwise assembly; cf. Section 6.1.2). For a normalized state $|\Psi\rangle \in \mathcal{H}_{\mathrm{em}}$, the Born expectations reduce HS words to standard tensors as in Chs. 5–6.

Definition 6.39 (HS relational action word)**.** The *holosymmetric relational action* associated to a normalized state $|\Psi\rangle$ is the scalar HS word

$$\boxed{\mathbf{S}[|\Psi\rangle] \;:=\; \int_{\mathcal{X}_q} d\boldsymbol{\mu}(x)\; \mathrm{Tr}\Big(\,\langle\Psi|\,\hat{\mathbf{G}}_\Lambda(x,x)\,|\Psi\rangle\,\Big).} \tag{6.85}$$

Remark 6.40 (Reduction to the standard action)**.** Taking Born expectation and using $\langle d\boldsymbol{\mu}\rangle = \sqrt{-\det \bar{g}^{\mathrm{Lor}}}\, d^dx$ (Definition 6.34) gives

$$\Big\langle \mathbf{S}[|\Psi\rangle]\Big\rangle = \int_{\mathcal{X}_q} d^dx\, \sqrt{-\det \bar{\Sigma}^{(\Psi)}(x)}\; \mathrm{Tr}\Big(\,\langle\Psi|\,\hat{\mathbf{G}}_\Lambda(x,x)\,|\Psi\rangle\,\Big) = S[|\Psi\rangle],$$

which is exactly the standard relational action of Definition 6.38, cf. (6.84).

Lemma 6.41 (HS stationarity and nonlinear eigenoperator)**.** *Impose* $\langle\Psi|\Psi\rangle = 1$ *and vary* $\mathbf{S}[|\Psi\rangle] - \lambda\,\langle\Psi|\Psi\rangle$*. Using only HS calculus and reduction identities, stationarity is equivalent to the* nonlinear *eigenoperator equation*

$$\boxed{\Big(\,\hat{\mathcal{H}}_{\mathrm{corr}}\Big[\bar{\Sigma}^{(\Psi)}\Big] \;+\; \widehat{\mathcal{B}}[\Psi]\,\Big)\,|\Psi\rangle \;=\; \lambda\,|\Psi\rangle,} \tag{6.86}$$

with

$$\hat{\mathcal{H}}_{\mathrm{corr}}\Big[\bar{\Sigma}\Big] := \int_{\mathcal{X}_q} d\mu_{\bar{\Sigma}}(x)\; \mathrm{Tr}\Big(\hat{\mathbf{G}}_\Lambda(x,x)\Big), \tag{6.87}$$

$$\widehat{\mathcal{B}}[\Psi] := \frac{1}{2}\int_{\mathcal{X}_q} d\mu_{\bar{\Sigma}^{(\Psi)}}(x)\; \bar{\Sigma}^{(\Psi)\mu\nu}(x)\; \mathsf{G}^{(\Psi)}(x)\; \hat{\Sigma}_{\mu\nu}(x), \qquad \mathsf{G}^{(\Psi)}(x) := \mathrm{Tr}\Big(\langle\Psi|\,\hat{\mathbf{G}}_\Lambda(x,x)\,|\Psi\rangle\Big), \tag{6.88}$$

and $\bar{\Sigma}^{(\Psi)\mu\nu}$ *is the (standard) inverse of* $\bar{\Sigma}^{(\Psi)}_{\mu\nu}$ *(cf. Definition 6.24 and eq. (6.62)), obtained by reducing the HS inverse of Definition 6.25.*

Proof idea. Vary (6.85) using the HS Leibniz rule and the reduction

$$\delta\sqrt{-\det\bar{\Sigma}^{(\Psi)}} = \tfrac{1}{2}\sqrt{-\det\bar{\Sigma}^{(\Psi)}}\,\bar{\Sigma}^{(\Psi)\mu\nu}\delta\bar{\Sigma}^{(\Psi)}_{\mu\nu}$$

Here $\delta\bar{\Sigma}^{(\Psi)}_{\mu\nu}(x) = \langle\delta\Psi|\hat{\Sigma}_{\mu\nu}(x)|\Psi\rangle + \text{h.c.}$ and $d\mu_{\bar{\Sigma}^{(\Psi)}} = \sqrt{-\det\bar{\Sigma}^{(\Psi)}}\,d^dx$ (Definition 6.33). Collect the pieces linear in $\langle\delta\Psi|$ to obtain (6.86).

Proof of Lemma 6.41. We use only holosymmetric (HS) wordwise machinery and the reduction identities from Chs. 4–6. Recall the HS relational action word (Definition 6.39)

$$\mathbf{S}[|\Psi\rangle] = \int_{\mathcal{X}_q} d\boldsymbol{\mu}(x)\,\mathrm{Tr}\Big(\langle\Psi|\,\hat{\mathbf{G}}_\Lambda(x,x)\,|\Psi\rangle\Big), \qquad d\boldsymbol{\mu}(x) = d^dx\,\boldsymbol{\varpi}(x), \quad \boldsymbol{\varpi} = \Big(\det_{\mathrm{HS}}(-\boldsymbol{g}^{\mathrm{Lor}})\Big)^{1/2}, \tag{6.89}$$

with $\boldsymbol{g}^{\mathrm{Lor}}_{\mu\nu} = \frac{1}{2}(\boldsymbol{\Sigma}_{\mu\nu} + \boldsymbol{\Sigma}_{\nu\mu})$ (cf. Sections 5.3.5 and 5.3.7) and the HS determinant/measure from Definitions 6.29 and 6.34. By Remark 6.35, taking the Born expectation reduces $d\boldsymbol{\mu}$ to the standard relational volume element $d\mu(x) = \sqrt{-\det\bar{g}^{\mathrm{Lor}}}\,d^dx$ (cf. Definition 6.33).

We impose the normalization constraint $\langle\Psi|\Psi\rangle = 1$ via a Lagrange multiplier λ and vary the functional

$$\mathcal{F}[|\Psi\rangle] := \mathbf{S}[|\Psi\rangle] - \lambda\,\langle\Psi|\Psi\rangle.$$

Step 1: Decompose the variation. Write the integrand as $\mathcal{L}(x;|\Psi\rangle) := \boldsymbol{\varpi}(x) \odot \mathrm{Tr}\Big(\langle\Psi|\,\hat{\mathbf{G}}_\Lambda(x,x)\,|\Psi\rangle\Big)$, so that $\mathbf{S} = \int d^dx\,\mathcal{L}$. Since $\hat{\mathbf{G}}_\Lambda$ and $\hat{\Sigma}_{\mu\nu}$ are state-independent operators/words, all state-dependence enters through the Born expectations and through the measure density via $\bar{\Sigma}^{(\Psi)}_{\mu\nu}(x) := \langle\Psi|\hat{\Sigma}_{\mu\nu}(x)|\Psi\rangle$. Thus

$$\delta\mathcal{L} = \underbrace{\delta\boldsymbol{\varpi}(x)}_{\text{measure}} \odot \mathrm{Tr}\Big(\langle\Psi|\hat{\mathbf{G}}_\Lambda(x,x)|\Psi\rangle\Big) \;+\; \boldsymbol{\varpi}(x) \odot \mathrm{Tr}\Big(\langle\delta\Psi|\hat{\mathbf{G}}_\Lambda(x,x)|\Psi\rangle + \text{h.c.}\Big). \tag{6.90}$$

Step 2: Variation of the HS measure density (via the inverse). By Definition 6.24 (and (6.62)), $\bar{\Sigma}^{(\Psi)\mu\nu}$ denotes the inverse of $\bar{\Sigma}^{(\Psi)}_{\mu\nu}$, obtained by reducing the HS inverse (Definition 6.25). The standard determinant identity (matrix calculus together with Definition 6.24) gives

$$\delta\sqrt{-\det\bar{\Sigma}^{(\Psi)}} = \tfrac{1}{2}\sqrt{-\det\bar{\Sigma}^{(\Psi)}}\,\bar{\Sigma}^{(\Psi)\mu\nu}\,\delta\bar{\Sigma}^{(\Psi)}_{\mu\nu}. \tag{6.91}$$

Since $\delta\bar{\Sigma}^{(\Psi)}_{\mu\nu}(x) = \langle\delta\Psi|\hat{\Sigma}_{\mu\nu}(x)|\Psi\rangle + \langle\Psi|\hat{\Sigma}_{\mu\nu}(x)|\delta\Psi\rangle$, (6.91) yields the *measure-variation contribution*

$$\begin{aligned}\int_{\mathcal{X}_q} d^dx\;\delta\boldsymbol{\varpi}(x) \odot \mathrm{Tr}\Big(\langle\Psi|\hat{\mathbf{G}}_\Lambda(x,x)|\Psi\rangle\Big) &= \frac{1}{2}\int_{\mathcal{X}_q} d\mu_{\bar{\Sigma}^{(\Psi)}}(x)\;\bar{\Sigma}^{(\Psi)\mu\nu}(x)\;\mathsf{G}^{(\Psi)}(x)\\ &\quad\times\Big(\langle\delta\Psi|\hat{\Sigma}_{\mu\nu}(x)|\Psi\rangle + \text{h.c.}\Big)\\ &= \langle\delta\Psi|\;\widehat{\mathcal{B}}[\Psi]\;|\Psi\rangle \;+\; \text{h.c.},\end{aligned} \tag{6.92}$$

where $d\mu_{\bar{\Sigma}^{(\Psi)}} = \sqrt{-\det\bar{\Sigma}^{(\Psi)}}\,d^dx$ (cf. Definition 6.33), $\mathsf{G}^{(\Psi)}(x) := \mathrm{Tr}\Big(\langle\Psi|\hat{\mathbf{G}}_\Lambda(x,x)|\Psi\rangle\Big)$, and the *backreaction operator* is exactly (6.88):

$$\widehat{\mathcal{B}}[\Psi] := \frac{1}{2}\int_{\mathcal{X}_q} d\mu_{\bar{\Sigma}^{(\Psi)}}(x)\;\bar{\Sigma}^{(\Psi)\mu\nu}(x)\;\mathsf{G}^{(\Psi)}(x)\;\hat{\Sigma}_{\mu\nu}(x).$$

Step 3: Variation of the correlator expectation. Using linearity of the trace, the HS–covariant integration rule (Corollary 6.37), and the fact that $d\boldsymbol{\mu}$ is independent of the *infinitesimal* state variation (its variation was already taken into account in Step 2), we obtain

$$\begin{aligned}\int_{\mathcal{X}_q} d^d x\ \boldsymbol{\varpi}(x) \odot \mathrm{Tr}\Big(\langle \delta\Psi|\hat{\mathbf{G}}_\Lambda(x,x)|\Psi\rangle + \mathrm{h.c.}\Big) &= \langle\delta\Psi|\ \int_{\mathcal{X}_q} d\mu_{\bar{\Sigma}^{(\Psi)}}(x)\ \mathrm{Tr}\Big(\hat{\mathbf{G}}_\Lambda(x,x)\Big)\ |\Psi\rangle\ +\ \mathrm{h.c.} \\ &= \langle\delta\Psi|\ \hat{\mathcal{H}}_{\mathrm{corr}}[\bar{\Sigma}^{(\Psi)}]\ |\Psi\rangle\ +\ \mathrm{h.c.}, \end{aligned} \tag{6.93}$$

with $\hat{\mathcal{H}}_{\mathrm{corr}}[\bar{\Sigma}]$ defined in (6.87).[2]

Step 4: Collect terms and impose the constraint. Combining (6.92) and (6.93), and subtracting the constraint variation $-\lambda(\langle\delta\Psi|\Psi\rangle + \mathrm{h.c.})$, we get

$$\delta\mathcal{F} = \langle\delta\Psi|\ \Big(\hat{\mathcal{H}}_{\mathrm{corr}}[\bar{\Sigma}^{(\Psi)}] + \widehat{\mathcal{B}}[\Psi]\Big)\ |\Psi\rangle\ -\ \lambda\,\langle\delta\Psi|\Psi\rangle\ +\ \mathrm{h.c.}$$

Since $|\delta\Psi\rangle$ is arbitrary and both $\hat{\mathcal{H}}_{\mathrm{corr}}[\bar{\Sigma}^{(\Psi)}]$ and $\widehat{\mathcal{B}}[\Psi]$ are Hermitian (the latter because it is an HS–reduced integral of Hermitian densities with scalar weights), the stationarity condition $\delta\mathcal{F} = 0$ is equivalent to the *nonlinear* eigenoperator equation

$$\Big(\hat{\mathcal{H}}_{\mathrm{corr}}[\bar{\Sigma}^{(\Psi)}] + \widehat{\mathcal{B}}[\Psi]\Big)\,|\Psi\rangle = \lambda\,|\Psi\rangle,$$

which is precisely (6.86).

Step 5: Well-posedness of the integrals. All integrals above are well-defined on spectrally regulated fibres by Corollary 6.37 (HS-covariant integration) and the boundedness of $\boldsymbol{\varpi}$ on regulated patches; gauge covariance and centrality of the HS measure were established in Proposition 6.36. The inverse $\bar{\Sigma}^{(\Psi)\mu\nu}$ used in (6.91) exists on regions where $\bar{\Sigma}^{(\Psi)}$ is nondegenerate by Definition 6.24 and eq. (6.62), consistent with the HS inverse (Definition 6.25).

This completes the proof. □

Definition 6.42 (HS vacuum manifold)**.** The *HS vacuum manifold* is the set of normalized solutions modulo global phase

$$\boxed{\mathfrak{V}^{\mathrm{HS}}_{\mathrm{vac}} := \Big\{\ |\Psi\rangle\ \Big|\ (\hat{\mathcal{H}}_{\mathrm{corr}}[\bar{\Sigma}^{(\Psi)}] + \widehat{\mathcal{B}}[\Psi])\,|\Psi\rangle = \lambda_{\min}\,|\Psi\rangle\ \Big\}\Big/U(1),}\tag{6.94}$$

with $\bar{\Sigma}^{(\Psi)}_{\mu\nu}(x) = \langle\Psi|\hat{\Sigma}_{\mu\nu}(x)|\Psi\rangle$. Its reduction equals the standard vacuum manifold of critical states of $S[|\boldsymbol{\Psi}\rangle]$ (cf. Section 6.1.8).

Proposition 6.43 (HS variational–spectral equivalence)**. (i) Frozen geometry**
If $\bar{\Sigma}$ is held fixed (no backreaction), the minimizers of **S** *are precisely the ground states of the* linear *operator*

$$\boxed{\hat{\mathcal{H}}_{\mathrm{corr}}[\bar{\Sigma}] = \int_{\mathcal{X}_q} d\mu_{\bar{\Sigma}}(x)\ \mathrm{Tr}\Big(\hat{\mathbf{G}}_\Lambda(x,x)\Big).}\tag{6.95}$$

[2]Here, "Tr" is over the field-space (and dialectic) block indices of the unified correlator, leaving an operator on the emergent Hilbert space; cf. the assembly rules in Section 6.1.2.

(ii) Self-consistent vacuum

With backreaction, vacua are fixed points of the nonlinear *map*

$$|\Psi\rangle \;\mapsto\; \Big(\hat{\mathcal{H}}_{\mathrm{corr}}[\bar{\Sigma}^{(\Psi)}] + \widehat{\mathcal{B}}[\Psi]\Big)|\Psi\rangle,$$

i.e. solutions of (6.86) *with* $\bar{\Sigma}^{(\Psi)}_{\mu\nu}(x) = \langle\Psi|\hat{\Sigma}_{\mu\nu}(x)|\Psi\rangle$. *Under spectral regularity and positivity of the integrand, minimizing sequences exist; fixed-point existence follows under standard compactness hypotheses for the induced map on the projective Hilbert space.*

Proof of Proposition 6.43. We use only HS-wordwise machinery and the reduction identities established in Definition 6.39, Remark 6.40, Proposition 6.36, and Corollary 6.37 together with the stationarity Lemma 6.41.

Preliminaries. Recall the HS relational action word (Definition 6.39)

$$\mathbf{S}[|\Psi\rangle] = \int_{\mathcal{X}_q} d\boldsymbol{\mu}(x)\ \mathrm{Tr}\Big(\langle\Psi|\hat{\mathbf{G}}_\Lambda(x,x)|\Psi\rangle\Big),$$

and its reduction to the standard action (Remark 6.40). By Corollary 6.37,

$$\Big\langle \mathring{\boldsymbol{\Psi}}\Big|\ \int d\boldsymbol{\mu}\,\mathbf{W}\ \Big|\mathring{\boldsymbol{\Psi}}\Big\rangle = \int d\mu\ \langle\mathbf{W}\rangle, \tag{6.96}$$

for any HS word observable with well-defined Born expectation. We also recall that the HS measure $d\boldsymbol{\mu}$ is gauge-covariant and central on flat patches (Proposition 6.36), hence does not obstruct spectral constructions.

(i) Frozen geometry. Assume $\bar{\Sigma}$ is held fixed (no backreaction). Then $d\boldsymbol{\mu}$ is independent of $|\Psi\rangle$ and

$$\mathbf{S}[|\Psi\rangle] = \int_{\mathcal{X}_q} d\boldsymbol{\mu}_{\bar{\Sigma}}(x)\ \mathrm{Tr}\Big(\langle\Psi|\hat{\mathbf{G}}_\Lambda(x,x)|\Psi\rangle\Big) = \Big\langle\Psi\Big|\ \underbrace{\int_{\mathcal{X}_q} d\mu_{\bar{\Sigma}}(x)\ \mathrm{Tr}\Big(\hat{\mathbf{G}}_\Lambda(x,x)\Big)}_{=:\ \hat{\mathcal{H}}_{\mathrm{corr}}[\bar{\Sigma}]}\ \Big|\Psi\Big\rangle,$$

where we used (6.96) with fixed geometry to pass the expectation inside. Thus $\mathbf{S}[|\Psi\rangle] = \langle\Psi|\hat{\mathcal{H}}_{\mathrm{corr}}[\bar{\Sigma}]|\Psi\rangle$ with $\hat{\mathcal{H}}_{\mathrm{corr}}[\bar{\Sigma}]$ given by (6.95). Minimizing $\mathbf{S}$ under $\langle\Psi|\Psi\rangle = 1$ is the Rayleigh–Ritz problem for the *linear*, Hermitian, positive operator $\hat{\mathcal{H}}_{\mathrm{corr}}[\bar{\Sigma}]$ (positivity follows from two-point positivity of $\hat{\mathbf{G}}_\Lambda$ and positivity of $d\mu_{\bar{\Sigma}}$). Hence the minimizers are precisely the ground states (eigenvectors associated to the least eigenvalue), and the minimum is that least eigenvalue. This proves *(i)*.

(ii) Self-consistent vacuum. Allowing backreaction, $\bar{\Sigma}^{(\Psi)}_{\mu\nu}(x) = \langle\Psi|\hat{\Sigma}_{\mu\nu}(x)|\Psi\rangle$ and the measure $d\mu_{\bar{\Sigma}^{(\Psi)}}$ depend on $|\Psi\rangle$. By Lemma 6.41, stationarity of $\mathbf{S}$ under the normalization constraint is *equivalent* to the nonlinear eigenoperator equation

$$\Big(\hat{\mathcal{H}}_{\mathrm{corr}}[\bar{\Sigma}^{(\Psi)}] + \widehat{\mathcal{B}}[\Psi]\Big)\,|\Psi\rangle = \lambda\,|\Psi\rangle, \qquad \text{cf. (6.86) with (6.87) and (6.88).} \tag{6.97}$$

Thus any critical state (vacuum) is a fixed point of the nonlinear map

$$\mathcal{T}: \ |\Psi\rangle \ \mapsto \ \Big(\hat{\mathcal{H}}_{\mathrm{corr}}[\bar{\Sigma}^{(\Psi)}] + \widehat{\mathcal{B}}[\Psi]\Big)\,|\Psi\rangle,$$

up to overall normalization and phase (projective class). Conversely, any normalized solution of (6.97) is a stationary point of $\mathbf{S}$.

Existence of minimizing sequences and fixed points. Under spectral regularity (spectral cutoff on fibres) the kernels entering $\hat{\mathcal{H}}_{\mathrm{corr}}[\bar{\Sigma}^{(\Psi)}]$ and $\widehat{\mathcal{B}}[\Psi]$ define bounded Hermitian operators; positivity of the integrand $\mathrm{Tr}(\hat{\mathbf{G}}_\Lambda(x,x)) \geq 0$ implies $\mathbf{S}[|\Psi\rangle] \geq 0$, so $\inf_{\|\Psi\|=1} \mathbf{S}$ exists and there are minimizing sequences $\{|\Psi_n\rangle\}$ on the unit sphere. By weak compactness of the unit ball and lower semicontinuity of the quadratic form induced by $\hat{\mathcal{H}}_{\mathrm{corr}}[\bar{\Sigma}^{(\Psi)}]$ plus the continuity of the backreaction functional $|\Psi\rangle \mapsto \widehat{\mathcal{B}}[\Psi]$ (integral of bounded linear maps composed with the continuous $|\Psi\rangle \mapsto \bar{\Sigma}^{(\Psi)}$), a subsequence converges (in the projective Hilbert space) to a minimizer. By Lemma 6.41, any minimizer satisfies (6.97). Equivalently, on a convex, weakly compact subset of the projective space, the map $|\Psi\rangle \mapsto \Pi\Big(\hat{\mathcal{H}}_{\mathrm{corr}}[\bar{\Sigma}^{(\Psi)}] + \widehat{\mathcal{B}}[\Psi]\Big)|\Psi\rangle$ (with Π the normalization projector) is continuous; Schauder's fixed-point theorem then yields a fixed point, hence a self-consistent vacuum.

Therefore, vacua are exactly the fixed points of the nonlinear map in item *(ii)*, i.e. solutions of (6.97) with $\bar{\Sigma}^{(\Psi)}$ determined by $|\Psi\rangle$. This proves *(ii)* and completes the proof. □

Remark 6.44 (Gauge covariance, positivity, and two-level reading)**.** *Gauge covariance:* under endpoint unitaries $U(x)$, the ingredients transform by conjugation, $d\boldsymbol{\mu}$ is central in flat patches, and spectra of $\hat{\mathcal{H}}_{\mathrm{corr}}$ are gauge invariant. *Positivity:* if $\hat{\mathbf{G}}_\Lambda(x,x) \succeq 0$ (two-point positivity) and $d\boldsymbol{\mu} \succeq 0$, then $\hat{\mathcal{H}}_{\mathrm{corr}} \succeq 0$. *Two-level reading:* the "general functional" vs. "evaluation on the Holo–State" corresponds to (i) solving (6.86) for $|\Psi\rangle$, then (ii) evaluating at $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$ to obtain the realized cosmic vacuum background.

6.1.10 The Trace of the Inter-Modal Correlator

1. The Structure of the Unified Inter-Modal Correlator

The operator-valued matrix $\hat{\mathbf{G}}_\Lambda(x,x)$ is defined (Definition 6.4) as:

$$\hat{\mathbf{G}}_\Lambda(x,x) := \begin{pmatrix} \hat{G}_{\boldsymbol{\Phi}\boldsymbol{\Phi}}(x,x) & \hat{G}_{\boldsymbol{\Phi}\boldsymbol{\Theta}}(x,x) \\ \hat{G}_{\boldsymbol{\Theta}\boldsymbol{\Phi}}(x,x) & \hat{G}_{\boldsymbol{\Theta}\boldsymbol{\Theta}}(x,x) \end{pmatrix},$$

where each entry is itself an operator on the total emergent Hilbert space $\mathcal{H}_{\mathrm{em}}$, and each block is internally a 4×4 matrix in the dialectical mode indices $\alpha, \beta \in \{0, 1, p, s\}$.

2. Expectation Value in $|\Psi\rangle$

When sandwiched between $|\boldsymbol{\Psi}\rangle$, the matrix-valued operator $\hat{\mathbf{G}}_\Lambda(x,x)$ yields a 2×2 matrix whose entries are 4×4 matrices over the dialectical mode indices $\alpha, \beta \in \{0, 1, p, s\}$. The result is:

$$\Big\langle \boldsymbol{\Psi}\Big|\hat{\mathbf{G}}_\Lambda(x,x)\Big|\boldsymbol{\Psi}\Big\rangle = \begin{pmatrix} \Big[\langle\boldsymbol{\Psi}|\hat{G}^{\alpha\beta}_{\boldsymbol{\Phi}\boldsymbol{\Phi}}(x,x)|\boldsymbol{\Psi}\rangle\Big] & \Big[\langle\boldsymbol{\Psi}|\hat{G}^{\alpha\beta}_{\boldsymbol{\Phi}\boldsymbol{\Theta}}(x,x)|\boldsymbol{\Psi}\rangle\Big] \\ \Big[\langle\boldsymbol{\Psi}|\hat{G}^{\alpha\beta}_{\boldsymbol{\Theta}\boldsymbol{\Phi}}(x,x)|\boldsymbol{\Psi}\rangle\Big] & \Big[\langle\boldsymbol{\Psi}|\hat{G}^{\alpha\beta}_{\boldsymbol{\Theta}\boldsymbol{\Theta}}(x,x)|\boldsymbol{\Psi}\rangle\Big] \end{pmatrix}.$$

That is, each entry in this 2×2 matrix is a full 4×4 matrix of the form:

$$\left[\langle \mathbf{\Psi} | \hat{G}^{\alpha\beta}_{IJ}(x,x) | \mathbf{\Psi} \rangle\right]_{\alpha,\beta \in \{0,1,p,s\}}.$$

3. Full Trace Expression

Taking the full trace means summing over the field indices $I = J$ and over the diagonal dialectical modes $\alpha = \beta$. Therefore:

$$\begin{aligned}\mathrm{Tr}\left(\langle \mathbf{\Psi} | \hat{\mathbf{G}}_\Lambda(x,x) | \mathbf{\Psi} \rangle\right) &= \sum_{I \in \{\mathbf{\Phi},\mathbf{\Theta}\}} \sum_{\alpha \in \{0,1,p,s\}} \langle \mathbf{\Psi} | \hat{G}^{\alpha\alpha}_{II}(x,x) | \mathbf{\Psi} \rangle \\ &= \sum_{\alpha} \left[\langle \mathbf{\Psi} | \hat{G}^{\alpha\alpha}_{\mathbf{\Phi}\mathbf{\Phi}}(x,x) | \mathbf{\Psi} \rangle + \langle \mathbf{\Psi} | \hat{G}^{\alpha\alpha}_{\mathbf{\Theta}\mathbf{\Theta}}(x,x) | \mathbf{\Psi} \rangle\right].\end{aligned}$$

This may be concisely expressed as:

$$\mathrm{Tr} = \mathrm{Tr}_{\text{field}} \circ \mathrm{Tr}_{\text{dialectic}} \left(\langle \mathbf{\Psi} | \hat{\mathbf{G}}_\Lambda(x,x) | \mathbf{\Psi} \rangle\right).$$

Thus, the quantity

$$\mathrm{Tr}\left(\langle \mathbf{\Psi} | \hat{\mathbf{G}}_\Lambda(x,x) | \mathbf{\Psi} \rangle\right)$$

is the *total sum over field types and modal diagonals* of the expectation values of the correlators evaluated at a point. It defines a scalar function of x, encoding the *local self-correlational energy density* of the generalized field operator in the state $|\mathbf{\Psi}\rangle$, which is then integrated over $\mathcal{X}_q$ in the relational action functional.

6.1.11 Holosymmetric Wordwise Trace of the Inter-modal Correlator

Setting. Let $\hat{\mathbf{G}}_\Lambda(x,x)$ be the unified inter–modal correlator (operator–valued matrix) at coincidence as in Definition 6.4 and the standard exposition of Section 6.1.10. Its 2×2 *field* block structure is indexed by $I, J \in \{\mathbf{\Phi}, \mathbf{\Theta}\}$; each block is a 4×4 *dialectic* matrix in $\alpha, \beta \in \{0, 1, p, s\}$. Expectation in a normalized state $|\Psi\rangle$ is $\langle \cdot \rangle_\Psi := \langle \Psi | \cdot | \Psi \rangle$.

Definition 6.45 (HS field–dialectic trace and scalar correlational density)**.** Define the nested field–dialectic trace by

$$\mathrm{Tr}_{\mathrm{FD}} \;:=\; \mathrm{Tr}_{\text{field}} \circ \mathrm{Tr}_{\text{dialectic}},$$

where $\mathrm{Tr}_{\text{dialectic}}$ takes the 4×4 trace over (α, β) and $\mathrm{Tr}_{\text{field}}$ takes the 2×2 trace over (I, J). The *HS scalar correlational density* is

$$\boxed{\mathsf{G}^{(\Psi)}(x) \;:=\; \mathrm{Tr}_{\mathrm{FD}}\Big(\langle\, \hat{\mathbf{G}}_\Lambda(x,x) \,\rangle_\Psi \Big).} \tag{6.98}$$

Remark 6.46 (Component expansion and reduction to the standard trace)**.** Writing out the two traces gives

$$\mathsf{G}^{(\Psi)}(x) = \sum_{I \in \{\mathbf{\Phi},\mathbf{\Theta}\}} \sum_{\alpha \in \{0,1,p,s\}} \left\langle \hat{G}^{\alpha\alpha}_{II}(x,x) \right\rangle_\Psi, \tag{6.99}$$

i.e. field diagonals and modal diagonals only.

Thus $\mathsf{G}^{(\Psi)}(x) = \mathrm{Tr}\Big(\langle \hat{\mathbf{G}}_\Lambda(x,x)\rangle_\Psi\Big)$, matching the standard formula of Section 6.1.10 and the integrand used in Definition 6.38 and eq. (6.84).

Proposition 6.47 (Gauge/basis invariance of $\mathsf{G}^{(\Psi)}$)**.** *Let $R \in U(2)$ act on the field indices and $V \in U(4)$ on the dialectic indices, and let $g(x)$ be an endpoint gauge acting by conjugation on the operator content. Then*

$$\hat{\mathbf{G}}_\Lambda \;\longmapsto\; (R\otimes V)\, g\, \hat{\mathbf{G}}_\Lambda\, g^\dagger\, (R^\dagger \otimes V^\dagger) \quad\Longrightarrow\quad \mathsf{G}^{(\Psi)}(x) \;\; \textit{is invariant.}$$

Proof. Cyclicity of finite-dimensional traces implies $\mathrm{Tr}_{\mathrm{FD}}\Big((R\otimes V)X(R^\dagger\otimes V^\dagger)\Big) = \mathrm{Tr}_{\mathrm{FD}}(X)$. For endpoint gauge, take the Born expectation first: $\langle g\,\hat{\mathbf{G}}_\Lambda\, g^\dagger\rangle_\Psi = \langle g^\dagger\Psi|\hat{\mathbf{G}}_\Lambda|g^\dagger\Psi\rangle$. Since $\mathrm{Tr}_{\mathrm{FD}}$ yields a c-number and only the ray $[\Psi]$ matters, the value is unchanged under $|\Psi\rangle \mapsto g^\dagger|\Psi\rangle$. Hence $\mathsf{G}^{(\Psi)}(x)$ is invariant. □

Proposition 6.48 (Positivity under spectral regularity)**.** *Suppose the coincident limit is UV-regularized and the two-point kernels are positive: for each field block I, the 4×4 matrix $\langle \hat{G}^{\alpha\beta}_{II}(x,x)\rangle_\Psi$ is positive semidefinite. Then $\mathsf{G}^{(\Psi)}(x) \geq 0$.*

Proof. Positivity of each 4×4 block implies $\mathrm{Tr}_{\mathrm{dialectic}}\langle \hat{G}_{II}(x,x)\rangle_\Psi \geq 0$. Summing over I preserves nonnegativity, hence (6.98) is ≥ 0. □

Corollary 6.49 (Insertion into the HS action and reduction)**.** *With the HS volume element $d\boldsymbol{\mu}(x)$ from Definition 6.34, the HS action word (cf. Definition 6.39 and eq. (6.85)) reads*

$$\mathbf{S}[|\Psi\rangle] = \int_{\mathcal{X}_q} d\boldsymbol{\mu}(x)\; \mathsf{G}^{(\Psi)}(x).$$

By Corollary 6.37,

$$\Big\langle \mathbf{S}[|\Psi\rangle]\Big\rangle = \int_{\mathcal{X}_q} d\mu(x)\; \mathsf{G}^{(\Psi)}(x),$$

which coincides with the standard relational action density of Definition 6.38.

6.1.12 Holosymmetric Control Dial — Fourfold Inter-modal Trace

Setup and notation. Let $\hat{\mathbf{G}}_\Lambda(x,x)$ be the unified correlator (field 2×2 blocks, each a 4×4 dialectic matrix) as in Definition 6.4 and Section 6.1.10. Write $\mathrm{Tr}_{\mathrm{FD}} = \mathrm{Tr}_{\mathrm{field}} \circ \mathrm{Tr}_{\mathrm{dialectic}}$ (Definition 6.45). Recall the operator–valued correlator density $\hat{\mathcal{G}}(x) = \mathrm{Tr}_{\mathrm{FD}}(\hat{\mathbf{G}}_\Lambda(x,x))$ (Definition 6.55). Denote the mixed off–diagonal field blocks by $\hat{G}_{\Phi\Theta}(x,x)$ and $\hat{G}_{\Theta\Phi}(x,x)$.

Definition 6.50 (HS law-control and reduced fourfold trace)**.** Introduce the two-level law-control space $\mathcal{H}_L = \mathrm{span}\{|0\rangle_L, |1\rangle_L\}$, and form the *HS-extended correlator density* on $\mathcal{H}_L \otimes \mathcal{H}_{\mathrm{em}}$:

$$\begin{aligned}\hat{\mathcal{G}}^{\mathrm{HS}}(x) := {} & |0\rangle\langle 0|_L \otimes \hat{\mathcal{G}}^{(0)}(x) \\ & + |1\rangle\langle 1|_L \otimes \hat{\mathcal{G}}^{(1)}(x) \\ & + |0\rangle\langle 1|_L \otimes \mathrm{Tr}_{\mathrm{dialectic}}\Big(\hat{G}_{\Phi\Theta}(x,x)\Big) \\ & + |1\rangle\langle 0|_L \otimes \mathrm{Tr}_{\mathrm{dialectic}}\Big(\hat{G}_{\Theta\Phi}(x,x)\Big),\end{aligned} \tag{6.100}$$

where $\hat{\mathcal{G}}^{(k)}(x) = \mathrm{Tr}_{\mathrm{FD}}(\hat{\mathbf{G}}^{(k)}_{\Lambda}(x,x))$ are branchwise copies (cf. eq. (6.121)).

Prepare the control register in $|+_*\varphi\rangle_L = \frac{1}{\sqrt{2}}(|0\rangle_L + e^{i\varphi}|1\rangle_L)$ and define the exchange–symmetric/antisymmetric mixed scalars

$$\begin{aligned} \mathsf{S}^{(\Psi)}(x) &:= \mathrm{Tr}_{\mathrm{dialectic}}\Big(\langle \hat{G}_{\Phi\Theta}(x,x) + \hat{G}_{\Theta\Phi}(x,x)\rangle_\Psi \Big), \\ \mathsf{C}^{(\Psi)}(x) &:= \mathrm{Tr}_{\mathrm{dialectic}}\Big(\langle i\hat{G}_{\Phi\Theta}(x,x) - i\hat{G}_{\Theta\Phi}(x,x)\rangle_\Psi \Big). \end{aligned} \tag{6.101}$$

Then the reduced scalar is

$$\boxed{\mathcal{G}^{(\Psi)}_{\langle \mathrm{HS}\rangle}(x;\varphi,\theta) = \frac{1}{2}\sum_{k=0}^{1} \mathsf{G}^{(\Psi),(k)}(x) + \mathrm{Re}\Big(e^{-i\varphi}\,[\,\cos\theta\,\mathsf{S}^{(\Psi)}(x) + \sin\theta\,\mathsf{C}^{(\Psi)}(x)\,]\Big),} \tag{6.102}$$

with $\mathsf{G}^{(\Psi),(k)}(x) = \langle\Psi|\hat{\mathcal{G}}^{(k)}(x)|\Psi\rangle$ (eq. (6.98)).

Remark 6.51 (Gauge covariance and causal structure)**.** By Propositions 6.31 and 6.36, the HS measure and determinant are gauge covariant, and central on flat patches. Since the dial (φ,θ) enters only through $\widehat{\mathcal{G}}^{\mathrm{HS}}$ and not through $\boldsymbol{g}^{\mathrm{Lor}}$ or $d\boldsymbol{\mu}$, the effective light-cone assignment is unchanged (cf. Section 5.4.2).

Proposition 6.52 (Tomography: φ- and θ-derivatives isolate symmetric/antisymmetric mixes)**.** *With* (6.102)*,*

$$\partial_\varphi\, \mathcal{G}^{(\Psi)}_{\langle \mathrm{HS}\rangle}(x;\varphi,\theta) = -\,\mathrm{Im}\Big(e^{-i\varphi}\,[\,\cos\theta\,\mathsf{S}^{(\Psi)}(x) + \sin\theta\,\mathsf{C}^{(\Psi)}(x)\,]\Big), \tag{6.103}$$

$$\partial_\theta\, \mathcal{G}^{(\Psi)}_{\langle \mathrm{HS}\rangle}(x;\varphi,\theta) = \;\mathrm{Re}\Big(e^{-i\varphi}\Big[\,-\sin\theta\,\mathsf{S}^{(\Psi)}(x) + \cos\theta\,\mathsf{C}^{(\Psi)}(x)\Big]\Big). \tag{6.104}$$

In particular,

$$\boxed{\begin{aligned} \partial_\theta \mathcal{G}^{(\Psi)}_{\langle \mathrm{HS}\rangle}(x;\varphi,0) &= \mathrm{Re}\Big(e^{-i\varphi}\,\mathsf{C}^{(\Psi)}(x)\Big), \qquad (6.105)\\ \partial_\theta \mathcal{G}^{(\Psi)}_{\langle \mathrm{HS}\rangle}(x;\varphi,\tfrac{\pi}{2}) &= -\,\mathrm{Re}\Big(e^{-i\varphi}\,\mathsf{S}^{(\Psi)}(x)\Big). \qquad (6.106) \end{aligned}}$$

Choosing $\varphi = 0$ (or $\varphi = \frac{\pi}{2}$) selects real (or imaginary) parts of these mixes.

Corollary 6.53 (Insertion into HS action/Hamiltonian is causal-structure neutral)**.** *With the HS volume $d\boldsymbol{\mu}$ (Definition 6.34) and the HS Hamiltonian word (Definition 6.56), replacing $\mathsf{G}^{(\Psi)}$ by $\mathcal{G}^{(\Psi)}_{\langle \mathrm{HS}\rangle}(\varphi,\theta)$ changes only the integrand by $\mathrm{Re}(e^{-i\varphi}[\,\cos\theta\,\mathsf{S}^{(\Psi)} + \sin\theta\,\mathsf{C}^{(\Psi)}\,])$; since neither $d\boldsymbol{\mu}$ nor $\boldsymbol{g}^{\mathrm{Lor}}$ depends on (φ,θ), the effective light-cone assignment (Section 5.4.2) is unchanged.*

6.1.13 The Relational Hamiltonian

In a background-independent framework where time is emergent, the Hamiltonian is not defined as a generator of temporal evolution but is instead derived from the theory's fundamental action principle. The **Relational Hamiltonian**, $\hat{H}$, is the self-adjoint operator whose expectation value for any state is the Relational Action Functional, $S[|\boldsymbol{\Psi}\rangle]$. This definition establishes the **Variational-Spectral Equivalence Principle**: the states that minimize the action are precisely the ground states of the Hamiltonian.

Definition 6.54 (The Relational Hamiltonian, $\hat{H}$)**.** Let $S[|\mathbf{\Psi}\rangle]$ be the Relational Action Functional defined over the total emergent Hilbert space, $\mathcal{H}_{\text{em}}$ (Definition 6.38). The **Relational Hamiltonian** is defined implicitly through its expectation value:

$$\boxed{\langle \mathbf{\Psi}|\hat{H}|\mathbf{\Psi}\rangle := S[|\mathbf{\Psi}\rangle].} \tag{6.107}$$

This implicit definition is made explicit through the construction of a state-dependent operator, $\hat{H}[|\mathbf{\Psi}\rangle]$, which acts as the integral of an operator-valued Lagrangian kernel over the emergent manifold $\mathcal{X}_q$:

$$\boxed{\hat{H}[|\mathbf{\Psi}\rangle] := \int_{\mathcal{X}_q} d^d x \sqrt{-\det \bar{\Sigma}^{(\mathbf{\Psi})}_{\mu\nu}(x)}\ \text{Tr}_{\text{op}}\left(\hat{\mathbf{G}}_\Lambda(x,x)\right),} \tag{6.108}$$

where Tr_{op} is the partial trace over the internal field and dialectic indices, leaving an operator on $\mathcal{H}_{\text{em}}$. The state-dependence of the Hamiltonian, arising from the volume element $\sqrt{-\det \bar{\Sigma}^{(\mathbf{\Psi})}}$, is a necessary feature of the theory's relational self-consistency.

Decomposition into Field Sectors The total Hamiltonian inherits its structure from the Unified Inter-Modal Correlator, $\hat{\mathbf{G}}_\Lambda$, and thus admits a canonical decomposition into Gaiamitiic (geometric), Timbric (material), and Interaction components:

$$\boxed{\hat{H}[|\mathbf{\Psi}\rangle] = \hat{H}_\Phi[|\mathbf{\Psi}\rangle] + \hat{H}_\Theta[|\mathbf{\Psi}\rangle] + \hat{H}_{\Phi\Theta}[|\mathbf{\Psi}\rangle],} \tag{6.109}$$

where each component is defined by integrating the trace of the corresponding block of the correlator matrix:

$$\begin{aligned}
\hat{H}_\Phi[|\mathbf{\Psi}\rangle] &:= \int_{\mathcal{X}_q} d^d x \sqrt{-\det \bar{\Sigma}^{(\mathbf{\Psi})}}\ \text{Tr}_{\text{op}}\left(\hat{G}_{\mathbf{\Phi\Phi}}(x,x)\right), \\
\hat{H}_\Theta[|\mathbf{\Psi}\rangle] &:= \int_{\mathcal{X}_q} d^d x \sqrt{-\det \bar{\Sigma}^{(\mathbf{\Psi})}}\ \text{Tr}_{\text{op}}\left(\hat{G}_{\mathbf{\Theta\Theta}}(x,x)\right), \\
\hat{H}_{\Phi\Theta}[|\mathbf{\Psi}\rangle] &:= \int_{\mathcal{X}_q} d^d x \sqrt{-\det \bar{\Sigma}^{(\mathbf{\Psi})}}\ \text{Tr}_{\text{op}}\left(\hat{G}_{\mathbf{\Phi\Theta}}(x,x) + \hat{G}_{\mathbf{\Theta\Phi}}(x,x)\right).
\end{aligned}$$

The ground state eigenspace, $\mathcal{V}_0 = \ker(\hat{H} - E_0 \mathbb{I}_{\text{em}})$, is the set of states that minimize the action, thereby defining the *Vacuum Manifold*, $\mathcal{M}_{\text{vac}}$ (Definition 6.61).

The state-dependence of the Hamiltonian, denoted $\hat{H}[|\mathbf{\Psi}\rangle]$, is not a flaw but a necessary and profound feature of a truly background-independent theory. The geometry of the universe, encoded in the volume element $\sqrt{-\det \bar{\Sigma}^{(\mathbf{\Psi})}}$, is not a fixed stage upon which dynamics unfold; it is an active participant, shaped by the very state whose energy it measures.

This definition completes the conceptual architecture outlined in Definition 6.61 by providing an explicit, albeit state-dependent, form for the Relational Hamiltonian. Finding the vacuum states of the theory is now equivalent to solving the non-linear eigenvalue problem for the ground state of this operator.

6.1.14 Holosymmetric Wordwise Relational Hamiltonian

Ingredients. Let $\hat{\mathbf{G}}_\Lambda(x,x)$ be the unified inter–modal correlator (Definition 6.4 and Section 6.1.10), and let $d\boldsymbol{\mu}(x)$ be the HS relational volume element (Definition 6.34 and Section 6.1.7). Denote the field–dialectic trace by $\mathrm{Tr}_{\mathrm{FD}} = \mathrm{Tr}_{\mathrm{field}} \circ \mathrm{Tr}_{\mathrm{dialectic}}$ (Definition 6.45), and define the operator–valued correlator density:

Definition 6.55 (HS operator correlator density)**.**

$$\boxed{\hat{\mathcal{G}}(x) \;:=\; \mathrm{Tr}_{\mathrm{FD}}\Big(\,\hat{\mathbf{G}}_\Lambda(x,x)\,\Big), \qquad (\text{operator on } \mathcal{H}_{\mathrm{em}}).} \tag{6.110}$$

Its Born expectation in $|\Psi\rangle$ reduces to the scalar density of Definition 6.45 and eq. (6.98): $\mathsf{G}^{(\Psi)}(x) = \langle\Psi|\hat{\mathcal{G}}(x)|\Psi\rangle$, which equals the standard trace of Section 6.1.10.

Definition 6.56 (HS Relational Hamiltonian word)**.** The *HS Relational Hamiltonian* associated to a normalized state $|\Psi\rangle$ is

$$\boxed{\widehat{\boldsymbol{H}}[|\Psi\rangle] \;:=\; \int_{\mathcal{X}_q} d\boldsymbol{\mu}(x)\; \hat{\mathcal{G}}(x) \;=\; \int_{\mathcal{X}_q} d^d x\; \boldsymbol{\varpi}(x)\; \hat{\mathcal{G}}(x),} \tag{6.111}$$

with $\boldsymbol{\varpi}(x) = \Big(\det_{\mathrm{HS}}(-\boldsymbol{g}^{\mathrm{Lor}}(x))\Big)^{1/2}$ (Definitions 6.29 and 6.34).

Proposition 6.57 (Expectation reduction equals the Relational Action)**.** *For all normalized* $|\Psi\rangle$*,*

$$\boxed{\langle\Psi|\widehat{\boldsymbol{H}}[|\Psi\rangle]|\Psi\rangle \;=\; \int_{\mathcal{X}_q} d\mu(x)\; \mathsf{G}^{(\Psi)}(x) \;=\; S[|\Psi\rangle],} \tag{6.112}$$

where $d\mu(x) = \sqrt{-\det \bar{\Sigma}^{(\Psi)}(x)}\, d^d x$ *and* $S[|\Psi\rangle]$ *is the standard Relational Action Functional (Definition 6.38 and eq. (6.84)).*

Proof idea. Use $\langle d\boldsymbol{\mu}\rangle = \sqrt{-\det \bar{\Sigma}^{(\Psi)}}\, d^d x$ (Remark 6.35) and $\mathsf{G}^{(\Psi)}(x) = \langle\Psi|\hat{\mathcal{G}}(x)|\Psi\rangle$ (Definitions 6.45 and 6.55) together with HS–covariant integration (Corollary 6.37).

Proof (expanded). Let $|\Psi\rangle$ be normalized and in the common domain of all local densities. By HS–covariant integration (Corollary 6.37), the state-dependent relational Hamiltonian is implemented as an operator-valued integral of a local density against the relational volume element:

$$\widehat{\boldsymbol{H}}[|\Psi\rangle] \;=\; \int_{\mathcal{X}_q} d\boldsymbol{\mu}(x)\; \hat{\mathcal{G}}(x), \qquad \text{with} \quad \hat{\mathcal{G}}(x) \text{ as in } \textit{Definition } 6.55. \tag{6.113}$$

Taking the expectation in $|\Psi\rangle$ and using the linearity/continuity guaranteed by Corollary 6.37 yields the HS–covariant push-through of the expectation:

$$\langle\Psi|\widehat{\boldsymbol{H}}[|\Psi\rangle]|\Psi\rangle \;=\; \int_{\mathcal{X}_q} \langle\Psi|\, d\boldsymbol{\mu}(x)\, \hat{\mathcal{G}}(x)\, |\Psi\rangle. \tag{6.114}$$

Since $d\boldsymbol{\mu}(x)$ is an operator-valued volume element built from the unified chronogeometric tensor, its *expectation* becomes a scalar positive measure; precisely (Remark 6.35),

$$\langle d\boldsymbol{\mu}(x)\rangle \;=\; \sqrt{-\det \bar{\Sigma}^{(\Psi)}(x)}\, d^d x \;=:\; d\mu(x). \tag{6.115}$$

Moreover, by the definition of the operator correlator density and the field-diagonal trace/expectation correspondence (Definitions 6.45 and 6.55),

$$\mathsf{G}^{(\Psi)}(x) \;:=\; \langle\Psi|\hat{\mathcal{G}}(x)|\Psi\rangle. \tag{6.116}$$

Combining (6.114)–(6.116) we obtain

$$\langle\Psi|\widehat{\boldsymbol{H}}[|\Psi\rangle]|\Psi\rangle \;=\; \int_{\mathcal{X}_q} d\mu(x)\; \mathsf{G}^{(\Psi)}(x) \;=\; \int_{\mathcal{X}_q} \sqrt{-\det \bar{\Sigma}^{(\Psi)}(x)}\; \mathsf{G}^{(\Psi)}(x)\, d^d x. \tag{6.117}$$

It remains to identify the right-hand side with the Relational Action. By the very definition of the Relational Action Functional (Definition 6.38 and eq. (6.84)), the action for the state $|\Psi\rangle$ is the HS–covariant integral of the expectation-level density against the relational volume measure:

$$S[|\Psi\rangle] \;=\; \int_{\mathcal{X}_q} d\mu(x)\; \mathsf{G}^{(\Psi)}(x). \tag{6.118}$$

Comparing (6.117) with (6.118) gives the desired identity

$$\langle\Psi|\widehat{\boldsymbol{H}}[|\Psi\rangle]|\Psi\rangle \;=\; S[|\Psi\rangle].$$

This completes the proof. □

Remark 6.58 (On triadic decomposition, gauge, and boundary terms)**.** (i) The equality holds triad-wise since both $d\boldsymbol{\mu}$ and $\hat{\mathcal{G}}$ decompose into corr/vac/spec sectors, and Corollary 6.37 preserves the sectoral split. (ii) Gauge covariance at the operator level and gauge invariance at expectation level are inherited from the constructions used in Definition 6.55 and Remark 6.35. (iii) Any boundary contributions included in the sector actions are handled by the same HS–covariant integration corollary, so the identification with $S[|\Psi\rangle]$ remains exact.

Sector decomposition. Writing $\hat{\mathbf{G}}_\Lambda$ in field blocks and using $\mathrm{Tr}_{\mathrm{FD}}$,

$$\boxed{\widehat{\boldsymbol{H}}[|\Psi\rangle] = \widehat{\boldsymbol{H}}_\Phi[|\Psi\rangle] + \widehat{\boldsymbol{H}}_\Theta[|\Psi\rangle] + \widehat{\boldsymbol{H}}_{\Phi\Theta}[|\Psi\rangle],} \tag{6.119}$$

with

$$\begin{aligned}
\widehat{\boldsymbol{H}}_\Phi[|\Psi\rangle] &:= \int_{\mathcal{X}_q} d\boldsymbol{\mu}(x)\; \mathrm{Tr}_{\mathrm{op}}\Big(\hat{G}_{\boldsymbol{\Phi\Phi}}(x,x)\Big),\\
\widehat{\boldsymbol{H}}_\Theta[|\Psi\rangle] &:= \int_{\mathcal{X}_q} d\boldsymbol{\mu}(x)\; \mathrm{Tr}_{\mathrm{op}}\Big(\hat{G}_{\boldsymbol{\Theta\Theta}}(x,x)\Big),\\
\widehat{\boldsymbol{H}}_{\Phi\Theta}[|\Psi\rangle] &:= \int_{\mathcal{X}_q} d\boldsymbol{\mu}(x)\; \mathrm{Tr}_{\mathrm{op}}\Big(\hat{G}_{\boldsymbol{\Phi\Theta}}(x,x) + \hat{G}_{\boldsymbol{\Theta\Phi}}(x,x)\Big),
\end{aligned} \tag{6.120}$$

which reduce to eq. (6.109) upon taking expectations (Proposition 6.57).

Gauge covariance and centrality. Under endpoint gauge $g(x)$, $\hat{\mathbf{G}}_\Lambda \mapsto g\,\hat{\mathbf{G}}_\Lambda\,g^\dagger$ and $d\boldsymbol{\mu} \mapsto g\,d\boldsymbol{\mu}\,g^\dagger$ (Proposition 6.36). Hence $\widehat{\boldsymbol{H}}[|\Psi\rangle] \mapsto g\,\widehat{\boldsymbol{H}}[|\Psi\rangle]\,g^\dagger$. On flat/central patches, $d\boldsymbol{\mu}$ is central (Proposition 6.36), so $\widehat{\boldsymbol{H}}$ is gauge-covariant and its quadratic form is gauge invariant.

Variational–spectral equivalence (nonlinear). Stationarity of $\langle\Psi|\widehat{\boldsymbol{H}}[|\Psi\rangle]|\Psi\rangle$ under $\langle\Psi|\Psi\rangle = 1$ yields the nonlinear eigenoperator equation (Lemma 6.41):

$$\Big(\hat{\mathcal{H}}_{\mathrm{corr}}[\bar{\Sigma}^{(\Psi)}] + \widehat{\mathcal{B}}[\Psi]\Big)\,|\Psi\rangle = \lambda\,|\Psi\rangle,$$

so ground solutions coincide with minimizers of $S[|\Psi\rangle]$ (Proposition 6.43).

Positivity and self-adjointness. If the coincident limit is UV–regularized and $\hat{\mathbf{G}}_\Lambda(x,x) \succeq 0$ (two-point positivity), then $\hat{\mathcal{G}}(x) \succeq 0$ and the quadratic form $\langle\Psi|\widehat{\boldsymbol{H}}[|\Psi\rangle]|\Psi\rangle = S[|\Psi\rangle] \geq 0$. By standard representation theorems (closed, semibounded forms) a self-adjoint representative exists, consistent with Section 6.1.13.

6.1.15 Holosymmetric (HS) Extension of the Relational Hamiltonian

Control register and branch data. Introduce a two-level *law-control* Hilbert space $\mathcal{H}_L = \mathrm{span}\{|0\rangle_L, |1\rangle_L\}$ with matrix units $P_0 := |0\rangle\langle 0|_L$, $P_1 := |1\rangle\langle 1|_L$ and $E_{01} := |0\rangle\langle 1|_L$, $E_{10} := |1\rangle\langle 0|_L$. Fix a pair of branchwise HS data

$$\Big(\hat{\boldsymbol{\Sigma}}^{(k)}_{\mu\nu}(x),\ \hat{\mathbf{G}}^{(k)}_\Lambda(x,x)\Big), \qquad k \in \{0,1\},$$

constructed exactly as in the single-branch case (Definition 6.55 and eq. (6.110) and Definition 6.56 and eq. (6.111)), but allowing the two branches to be functorially distinct (e.g. "+/−" temporal sector, or model variants). Write the branch density and branch Hamiltonian words

$$\hat{\mathcal{G}}^{(k)}(x) := \mathrm{Tr}_{\mathrm{FD}}\Big(\hat{\mathbf{G}}^{(k)}_\Lambda(x,x)\Big), \qquad \widehat{\boldsymbol{H}}^{(k)}[|\Psi\rangle] := \int_{\mathcal{X}_q} d\boldsymbol{\mu}^{(k)}(x)\ \hat{\mathcal{G}}^{(k)}(x), \tag{6.121}$$

where $d\boldsymbol{\mu}^{(k)}(x)$ is the HS measure built from $\boldsymbol{g}^{\mathrm{Lor},(k)}_{\mu\nu} := \frac{1}{2}(\boldsymbol{\Sigma}^{(k)}_{\mu\nu} + \boldsymbol{\Sigma}^{(k)}_{\nu\mu})$ (Definition 6.34 and Section 6.1.7).

Off–diagonal (mixing) Hamiltonian from the correlator blocks. Define the *mixing Hamiltonian* directly from the off–diagonal field blocks of the unified correlator:

$$\boxed{\hat{H}_{\mathrm{mix}} := \int_{\mathcal{X}_q} d\boldsymbol{\mu}^{\langle 0/1\rangle}(x)\ \mathrm{Tr}_{\mathrm{op}}\Big(\mathrm{Tr}_{\mathrm{dialectic}}\Big(\hat{G}_{\boldsymbol{\Phi\Theta}}(x,x)\Big)\Big)}, \tag{6.122}$$

where $d\boldsymbol{\mu}^{\langle 0/1\rangle}(x)$ is any *symmetric* central choice built from the two branches (e.g. the arithmetic mean $\frac{1}{2}(d\boldsymbol{\mu}^{(0)} + d\boldsymbol{\mu}^{(1)})$). Its adjoint satisfies $\hat{H}^\dagger_{\mathrm{mix}} = \int d\boldsymbol{\mu}^{\langle 0/1\rangle}\ \mathrm{Tr}_{\mathrm{op}}\Big(\mathrm{Tr}_{\mathrm{dialectic}}(\hat{G}_{\boldsymbol{\Theta\Phi}})\Big)$, so no new symbols are introduced. Under endpoint gauge, covariance follows from Propositions 6.31 and 6.36 and blockwise conjugation of $\hat{G}_{\boldsymbol{\Phi\Theta}}, \hat{G}_{\boldsymbol{\Theta\Phi}}$.

Definition 6.59 (HS Hamiltonian word on $\mathcal{H}_L \otimes \mathcal{H}_{\mathrm{em}}$)**.**

$$\boxed{\widehat{\boldsymbol{H}}^{\mathrm{HS}}[|\Psi\rangle] \;:=\; P_0 \otimes \widehat{\boldsymbol{H}}^{(0)}[|\Psi\rangle] \;+\; P_1 \otimes \widehat{\boldsymbol{H}}^{(1)}[|\Psi\rangle] \;+\; E_{01} \otimes \hat{H}_{\mathrm{mix}} \;+\; E_{10} \otimes \hat{H}_{\mathrm{mix}}^{\dagger}.} \tag{6.123}$$

Proposition 6.60 (Gauge covariance and centrality)**.** *If each branch is gauge covariant and $d\boldsymbol{\mu}^{(k)}$ obeys Proposition 6.36, then for endpoint gauges $g(x)$ acting diagonally on the branches,*

$$\widehat{\boldsymbol{H}}^{\mathrm{HS}}[|\Psi\rangle] \;\mapsto\; (\mathbf{1}_L \otimes g)\; \widehat{\boldsymbol{H}}^{\mathrm{HS}}[|\Psi\rangle]\; (\mathbf{1}_L \otimes g)^{\dagger},$$

and on flat/central patches the quadratic form is gauge invariant (central measure).

Preparation–recombination and reduced HS Hamiltonian. Prepare the control register in the phase-tilted $|+\!*\,\varphi\rangle_L$ state

$$|+\!*\,\varphi\rangle_L := \frac{1}{\sqrt{2}}\Big(|0\rangle_L + e^{i\varphi}|1\rangle_L\Big), \qquad \langle+\!*\;\varphi| := \frac{1}{\sqrt{2}}\Big(\langle 0|_L + e^{-i\varphi}\langle 1|_L\Big). \tag{6.124}$$

The reduced operator on $\mathcal{H}_{\mathrm{em}}$ is

$$\hat{H}^{\langle \mathrm{HS}\rangle}[|\Psi\rangle;\varphi] := \langle+\!*\;\varphi|\,\widehat{\boldsymbol{H}}^{\mathrm{HS}}[|\Psi\rangle]\,|+\!*\,\varphi\rangle_L.$$

Using the matrix-unit algebra $P_iP_j = \delta_{ij}P_i$, $P_iE_{jk} = \delta_{ij}E_{ik}$, $E_{ij}E_{kl} = \delta_{jk}E_{il}$, we obtain

$$\boxed{\hat{H}^{\langle \mathrm{HS}\rangle}[|\Psi\rangle;\varphi] = \tfrac{1}{2}\Big(\widehat{\boldsymbol{H}}^{(0)}[|\Psi\rangle] + \widehat{\boldsymbol{H}}^{(1)}[|\Psi\rangle]\Big) + \mathrm{Re}\Big(e^{-i\varphi}\hat{H}_{\mathrm{mix}}\Big).} \tag{6.125}$$

Expectation reduction and action. Taking the Born expectation in $|\Psi\rangle$ and using $\langle d\boldsymbol{\mu}^{(k)}\rangle = \sqrt{-\det \bar{\Sigma}^{(\Psi),(k)}}\, d^dx$ (Remark 6.35) yields

$$\begin{aligned}\langle\Psi|\hat{H}^{\langle \mathrm{HS}\rangle}[|\Psi\rangle;\varphi]|\Psi\rangle &= \tfrac{1}{2}\sum_{k=0}^{1}\int_{\mathcal{X}_q} d\mu_{\bar{\Sigma}^{(\Psi),(k)}}(x)\; \mathsf{G}^{(\Psi),(k)}(x) \;+\; \mathrm{Re}\Big(e^{-i\varphi}\,\langle\Psi|\hat{H}_{\mathrm{mix}}|\Psi\rangle\Big)\\ &=: \tfrac{1}{2}\Big(S^{(0)}[|\Psi\rangle] + S^{(1)}[|\Psi\rangle]\Big) \;+\; \mathrm{Re}\Big(e^{-i\varphi}\,\langle\hat{H}_{\mathrm{mix}}\rangle_\Psi\Big),\end{aligned} \tag{6.126}$$

with $S^{(k)}$ the branch actions (Definition 6.38) formed from the k-branch data.

HS Euler–Lagrange (EL) operator. Vary $\langle\Psi|\hat{H}^{\langle \mathrm{HS}\rangle}[|\Psi\rangle;\varphi]|\Psi\rangle$ at fixed norm, exactly as in Lemma 6.41. Using the branchwise version of (6.86) and linearity of the variation, we obtain the HS-EL operator

$$\boxed{\begin{aligned}\hat{H}^{\langle \mathrm{HS}\rangle}_{\mathrm{EL}}[|\Psi\rangle;\varphi] = {}& \tfrac{1}{2}\Big(\hat{\mathcal{H}}_{\mathrm{corr}}[\bar{\Sigma}^{(\Psi),(0)}] + \hat{\mathcal{B}}^{(0)}[\Psi]\Big)\\ &+ \tfrac{1}{2}\Big(\hat{\mathcal{H}}_{\mathrm{corr}}[\bar{\Sigma}^{(\Psi),(1)}] + \hat{\mathcal{B}}^{(1)}[\Psi]\Big)\\ &+ \mathrm{Re}\Big(e^{-i\varphi}\,\delta_\Psi\hat{H}_{\mathrm{mix}}\Big),\end{aligned}} \tag{6.127}$$

where each branch term is as in eqs. (6.87) and (6.88), and $\delta_\Psi\hat{H}_{\mathrm{mix}}$ is the Fréchet derivative of (6.122) with respect to the state-dependent inputs (through $\bar{\Sigma}^{(\Psi),(k)}$ if the symmetric measure depends on $|\Psi\rangle$).

Remarks. (i) The symmetric choice $d\boldsymbol{\mu}^{\langle 0/1\rangle} := \frac{1}{2}(d\boldsymbol{\mu}^{(0)} + d\boldsymbol{\mu}^{(1)})$ keeps (6.122) central on flat patches; other symmetric choices (e.g. geometric mean of densities) are admissible and leave the tomography and covariance statements unchanged. (ii) No new symbols beyond your established $\mathrm{Tr}_{\mathrm{FD}}$, $d\boldsymbol{\mu}$, and the off–diagonal blocks $\hat{G}_{\boldsymbol{\Phi\Theta}}, \hat{G}_{\boldsymbol{\Theta\Phi}}$ are introduced.

6.1.16 Quantum Circuit Realization of the HS Relational Hamiltonian

Goal. Given the HS Hamiltonian word

$$\widehat{\boldsymbol{H}}[|\Psi\rangle] = \int_{\mathcal{X}_q} d\boldsymbol{\mu}(x)\, \hat{\mathcal{G}}(x), \qquad \hat{\mathcal{G}}(x) := \mathrm{Tr}_{\mathrm{FD}}\Big(\hat{\mathbf{G}}_{\Lambda}(x,x)\Big) \quad \text{(Definitions 6.55 and 6.56)},$$

we build a quantum circuit that estimates the expectation $\langle\Psi|\,\widehat{\boldsymbol{H}}[|\Psi\rangle]\,|\Psi\rangle = S[|\Psi\rangle]$ (Proposition 6.57) and, in the law-controlled case, implements the reduced HS Hamiltonian $\hat{H}^{\langle \mathrm{HS}\rangle}[|\Psi\rangle;\varphi]$ (eq. (6.125)).

Registers.

- *System register* S initialised in $|\Psi\rangle \in \mathcal{H}_{\mathrm{em}}$.
- *Quadrature (space) register* X to index nodes $\{x_j\}_{j=1}^{M}$ approximating $\int_{\mathcal{X}_q}$.
- *Field register* F for $I \in \{\boldsymbol{\Phi}, \boldsymbol{\Theta}\}$ (size 2).
- *Dialectic register* D for $\alpha \in \{0, 1, p, s\}$ (size 4).
- *Probe qubit* P for Hadamard-test style estimation.
- *(Optional) Law register* $\mathsf{L} = \mathrm{span}\{|0\rangle_L, |1\rangle_L\}$ for branch control (eq. (6.125)).

Discretised HS measure and weights. Choose a quadrature $\{(x_j, w_j)\}_{j=1}^{M}$ such that $\sum_j w_j\,(\cdot) \approx \int_{\mathcal{X}_q} d\mu_{\bar{\Sigma}^{(\Psi)}}(\cdot)$ with $w_j := \sqrt{-\det \bar{\Sigma}^{(\Psi)}(x_j)}\,\Delta x_j$ (reduction of $d\boldsymbol{\mu}$, Definition 6.34). Define the *weight state* on X:

$$\mathrm{PREP}_w : \ |0\rangle_{\mathsf{X}} \ \mapsto \ \sum_{j=1}^{M} \sqrt{\frac{w_j}{\sum_k w_k}}\ |j\rangle_{\mathsf{X}}.$$

Block-encoding of the correlator density $\hat{\mathcal{G}}(x_j)$. Using the Fourfold/field decomposition and transporter $\mathcal{U}$ (Definition 6.4 and Section 4.5), each block entry is of the word form

$$\hat{G}^{\alpha\beta}_{IJ}(x_j, x_j) = \Big(\hat{\mathcal{F}}^{(I)}_{\alpha}(x_j)\Big)^{\dagger}\, \mathcal{U}(x_j \leftarrow x_j)\, \hat{\mathcal{F}}^{(J)}_{\beta}(x_j).$$

We build a *SELECT–unitary* on $(\mathsf{F}, \mathsf{D}, \mathsf{S})$ that realises a linear combination of such unitaries:

$$\text{SELECT-}\mathbf{G} \ : \ |I,\alpha\rangle_{\mathsf{FD}} \otimes |\phi\rangle_{\mathsf{S}} \ \mapsto \ |I,\alpha\rangle_{\mathsf{FD}} \otimes \Big(\hat{\mathcal{F}}^{(I)}_{\alpha}\Big)^{\dagger}\, \mathcal{U}\, |\phi\rangle_{\mathsf{S}},$$

and its companion that applies $\hat{\mathcal{F}}_\beta^{(J)}$. With standard LCU/block-encoding (oblivious amplitude amplification), these yield a unitary $U_{\mathcal{G}}$ on $(\mathsf{F},\mathsf{D},\mathsf{X},\mathsf{S},\text{anc})$ such that

$$(\langle 0|_{\text{anc}} \otimes I)\, U_{\mathcal{G}}\, (|0\rangle_{\text{anc}} \otimes I) = \sum_j \Pi_j^{\mathsf{X}} \otimes \frac{\hat{\mathcal{G}}(x_j)}{\alpha_{\mathcal{G}}} \quad \text{(block-encoding)},$$

for a known $\alpha_{\mathcal{G}} \geq \|\hat{\mathcal{G}}(x)\|$ and projectors $\Pi_j^{\mathsf{X}} = |j\rangle\langle j|$.

FD-trace gadget. To implement $\mathrm{Tr}_{\mathrm{FD}}$ coherently, prepare a uniform superposition on (F,D), apply SELECT-$\mathbf{G}$ and its adjoint, and unprepare: this realises the sum over field/dialectic diagonals as a single block-encoding (the standard "trace via uniform index superposition" trick consistent with Section 6.1.10 and Definition 6.45). We fold this into $U_{\mathcal{G}}$.

Putting space and weights together. Define a *multiplexed* block-encoding controlled by X:

$$\text{SELECT-}x: \ |j\rangle_{\mathsf{X}} \ \mapsto \ U_{\mathcal{G}}(x_j).$$

Using the weight state on X, we obtain a global block-encoding of $\widehat{\boldsymbol{H}}[|\Psi\rangle] \approx \sum_j w_j\, \hat{\mathcal{G}}(x_j)$ up to the known scale $\alpha := \alpha_{\mathcal{G}} \sum_j w_j$.

Energy (action) estimator — probe-X Hadamard test. Let $O := \widehat{\boldsymbol{H}}[|\Psi\rangle]/\alpha$ be the normalised operator represented by the block-encoding. The following circuit estimates $\langle\Psi|O|\Psi\rangle$ (hence $S[|\Psi\rangle]$):



The expectation of Z on P equals $\mathrm{Re}\langle\Psi|O|\Psi\rangle$; using a phase on P yields the imaginary part if needed. Amplitude estimation upgrades precision if desired.

Law-controlled HS circuit (reduced Hamiltonian). In the presence of a law register and the mixing Hamiltonian $\hat{H}_{\text{mix}}$ (eqs. (6.122), (6.123) and (6.125)), set $H_{\text{avg}} := \frac{1}{2}\big(\widehat{\boldsymbol{H}}^{(0)} + \widehat{\boldsymbol{H}}^{(1)}\big)$ and $H_{\text{mix}} := \hat{H}_{\text{mix}}$. We denote a block-encoding unitary of an operator O by $\mathcal{U}[O]$. The circuit is:



Meaning of $\mathcal{U}[O]$: there exists an ancilla register of a qubits and a scale $\alpha > 0$ such that

$$(\langle 0^a| \otimes I)\, \mathcal{U}[O]\, (|0^a\rangle \otimes I) = \frac{O}{\alpha}.$$

Preparing L in $|+*\varphi\rangle$ and using the probe-X Hadamard test yields $\mathrm{Re}\Big(\langle\Psi|H_{\mathrm{avg}}|\Psi\rangle + e^{-i\varphi}\langle\Psi|H_{\mathrm{mix}}|\Psi\rangle\Big)$, matching eq. (6.125).

Here the control from L prepares the reduced operator $\hat{H}^{\langle \mathrm{HS}\rangle} = \frac{1}{2}(\widehat{\boldsymbol{H}}^{(0)} + \widehat{\boldsymbol{H}}^{(1)}) + \mathrm{Re}(e^{-i\varphi}\hat{H}_{\mathrm{mix}})$ exactly as in eq. (6.125). The same probe-X readout returns $\langle\Psi|\hat{H}^{\langle \mathrm{HS}\rangle}|\Psi\rangle$.

Remarks.

- **Gauge covariance.** Dressing/transport unitaries enter only inside the SELECT–**G** gadgets; by Propositions 6.31 and 6.36 the overall construction is gauge-covariant, and central on flat patches.
- **Backreaction loop.** The dependence of $w_j = \sqrt{-\det \bar{\Sigma}^{(\Psi)}(x_j)}\,\Delta x_j$ on $|\Psi\rangle$ implements the nonlinear EL problem (Lemma 6.41): classical outer loop updates $|\Psi\rangle$ (or its variational parameters), the circuit estimates $S[|\Psi\rangle]$ (or gradients via parameter-shift).
- **Complexity levers.** Accuracy is set by (i) quadrature size M, (ii) block-encoding normalisation $\alpha_{\mathcal{G}}$, and (iii) amplitude-estimation shots. The FD-trace gadget is $O(1)$ in registers (constant field/dialectic sizes) and polylog in M via state preparation PREP_w.

Bottom line. The HS grammar—selectors, transporter, FD-trace, HS measure, and (optionally) law register—maps one-to-one to:

(controlled dressings) + LCU block-encodings + weight-state quadrature + Hadamard tests,

yielding a concrete, programmable circuit that *is* the holosymmetric wordwise Relational Hamiltonian.

6.1.17 The Vacuum Manifold

The vacuum is not an empty stage but a specific quantum configuration of the universe characterized by minimal relational complexity. With the Relational Hamiltonian now formally defined, the vacuum manifold can be identified directly from its spectral properties.

Definition 6.61 (The Vacuum Manifold)**.** Let $\hat{H}$ be the Relational Hamiltonian defined over the total emergent Hilbert space $\mathcal{H}_{\mathrm{em}}$ (Definition 6.54), and let $E_0 := \inf \mathrm{spec}(\hat{H})$ be its lowest eigenvalue, corresponding to the ground state energy.

1. The **ground-state eigenspace**, $\mathcal{V}_0$, is the subspace of all states that are eigenvectors of $\hat{H}$ with eigenvalue E_0:

$$\boxed{\mathcal{V}_0 := \ker(\hat{H} - E_0 \mathbb{I}_{\mathrm{em}}).} \tag{6.128}$$

2. The **Vacuum Manifold**, $\mathcal{M}_{\text{vac}}$, is the set of all normalized states within this ground-state eigenspace:

$$\boxed{\mathcal{M}_{\text{vac}} := \{|\mathbf{\Psi}_{\text{vac}}\rangle \in \mathcal{V}_0 \mid \langle \mathbf{\Psi}_{\text{vac}}|\mathbf{\Psi}_{\text{vac}}\rangle = 1\}\,.} \tag{6.129}$$

Variational-Spectral Equivalence. This spectral definition is entirely equivalent to the variational one. By the construction of the Hamiltonian (eq. (6.107)), where $\langle \mathbf{\Psi}|\hat{H}|\mathbf{\Psi}\rangle := S[|\mathbf{\Psi}\rangle]$, the quantum mechanical principle of minimizing the energy expectation value to find the ground state is identical to the field-theoretic principle of finding the stationary points of the action. Therefore, a state is in the ground-state eigenspace if and only if it extremizes the Relational Action Functional:

$$|\Psi_{\text{vac}}\rangle \in \mathcal{V}_0 \quad \Longleftrightarrow \quad \left.\frac{\delta S[|\mathbf{\Psi}\rangle]}{\delta\langle \mathbf{\Psi}|}\right|_{|\mathbf{\Psi}\rangle=|\Psi_{\text{vac}}\rangle} = 0.$$

The states in the vacuum manifold are thus those corresponding to configurations of minimal relational curvature and maximal coherence. The specific vacuum state realized by our universe is identified with the Quantum Holo-State, $|\overset{\circ}{\mathbf{\Psi}}\rangle \in \mathcal{M}_{\text{vac}}$.

1. The Ground State Energy The symbol E_0 denotes the *lowest possible eigenvalue* of the Relational Hamiltonian $\hat{H}$, formally expressed as:

$$\boxed{E_0 := \inf \text{ spec}(\hat{H}),} \tag{6.130}$$

where $\text{spec}(\hat{H})$ is the spectrum of $\hat{H}$, i.e., the set of all its eigenvalues. This quantity E_0 is the *ground state energy* of the total quantum-relational system. As the Relational Hamiltonian $\hat{H}$ encodes the total relational curvature, phase flow, and matter-geometry interaction energy, the ground state energy E_0 represents the configuration of *minimal relational complexity*. States that saturate this bound are those of maximal coherence and minimal curvature, and are therefore identified as *vacuum states*.

2. The Identity Operator on the Emergent Hilbert Space The operator $\mathbb{I}_{\text{em}}$ is the identity on the total emergent Hilbert space (Definition 4.287):

$$\boxed{\mathbb{I}_{\text{em}} : \mathcal{H}_{\text{em}} \to \mathcal{H}_{\text{em}}, \qquad \mathbb{I}_{\text{em}}|\mathbf{\Psi}\rangle = |\mathbf{\Psi}\rangle \quad \forall\, |\mathbf{\Psi}\rangle \in \mathcal{H}_{\text{em}}.} \tag{6.131}$$

Mathematical Role. The expression $\hat{H} - E_0\mathbb{I}_{\text{em}}$ is well-defined because it subtracts an operator from an operator. Since E_0 is a scalar, it must be lifted to operator status by multiplying with the identity:

$$E_0\mathbb{I}_{\text{em}} = \text{“scalar rescaling operator”} : |\mathbf{\Psi}\rangle \mapsto E_0|\mathbf{\Psi}\rangle.$$

This allows us to interpret the kernel:

$$\mathcal{V}_0 := \ker(\hat{H} - E_0\mathbb{I}_{\text{em}})$$

as the *ground-state eigenspace*, i.e., the set of all states that satisfy:

$$\hat{H}|\mathbf{\Psi}_{\text{vac}}\rangle = E_0|\mathbf{\Psi}_{\text{vac}}\rangle.$$

Physical Interpretation of the Vacuum Manifold This definition provides a purely relational and background-independent characterization of the vacuum. The vacuum is not posited as an empty stage but is dynamically identified as the quantum state (or set of states) that minimizes the total relational content of the universe, thereby producing the "simplest" possible emergent geometry.

If the minimum of the action functional is unique, the vacuum state $|\mathring{\boldsymbol{\Psi}}\rangle$ is unique. If, however, there exists a continuous family of states that all yield the same minimum action, this set of degenerate ground states forms a non-trivial vacuum manifold, $\mathcal{M}_{\text{vac}}$. The existence of such a manifold is the necessary precondition for spontaneous symmetry breaking, where the system must "choose" one of the available vacua to reside in, thereby determining the specific geometric and physical properties of our emergent universe.

6.1.18 Holosymmetric Wordwise Vacuum Manifold

Definition 6.62 (Holosymmetric wordwise ground energy and projector)**.** Let the *Relational Hamiltonian word* be

$$\widehat{\boldsymbol{H}}[|\Psi\rangle] \;=\; \int_{\mathcal{X}_q} d\boldsymbol{\mu}(x)\,\hat{\mathcal{G}}(x) \quad \text{(Definitions 6.55 and 6.56)}.$$

Define the ground energy E_0 and its spectral projector $\boldsymbol{\Pi}_0$ by

$$\boxed{E_0 \;:=\; \inf \operatorname{spec}\Big(\widehat{\boldsymbol{H}}[|\Psi\rangle]\Big), \qquad \boldsymbol{\Pi}_0 \;:=\; \mathbf{1}_{\{E_0\}}\Big(\widehat{\boldsymbol{H}}[|\Psi\rangle]\Big),} \tag{6.132}$$

where $\mathbf{1}_{\{E_0\}}(\cdot)$ denotes the spectral projection at the eigenvalue E_0.

Definition 6.63 (Holosymmetric wordwise vacuum manifold (Hilbert and projective))**.** Let $\mathcal{V}_0 := \operatorname{Ran}\boldsymbol{\Pi}_0 = \ker\Big(\widehat{\boldsymbol{H}}[|\Psi\rangle] - E_0\,\mathbf{1}_{\text{em}}\Big)$ be the ground-eigenspace associated with (6.132). The *HS vacuum manifold (Hilbert form)* and its projective version are

$$\boxed{\mathcal{M}_{\text{vac}}^{(\text{word})} \;:=\; \Big\{\,|\Psi_{\text{vac}}\rangle \in \mathcal{V}_0 :\; \langle\Psi_{\text{vac}}|\Psi_{\text{vac}}\rangle = 1\,\Big\}, \qquad \mathbb{P}\mathcal{M}_{\text{vac}} \;\cong\; \mathbb{P}(\mathcal{V}_0).} \tag{6.133}$$

Here $\mathbb{P}(\mathcal{V}_0)$ denotes the projective space of rays in $\mathcal{V}_0$.

Variational–spectral equivalence (wordwise).

Proposition 6.64 (Variational–spectral equivalence (wordwise))**.** *Let $\widehat{\boldsymbol{H}}[|\Psi\rangle]$ be the Relational Hamiltonian word (Definition 6.56) with ground data $(E_0, \boldsymbol{\Pi}_0)$ from Definition 6.62 and vacuum manifold $\mathcal{M}_{\text{vac}}^{(\text{word})}$ from Definition 6.63. With the action word $S[|\Psi\rangle] = \langle\Psi|\widehat{\boldsymbol{H}}[|\Psi\rangle]|\Psi\rangle$ (Proposition 6.57), the following hold for normalized states $|\Psi\rangle$ in the common domain:*

$$\boxed{|\Psi_{\text{vac}}\rangle \in \mathcal{M}_{\text{vac}}^{(\text{word})} \quad\Longleftrightarrow\quad \delta S[|\Psi\rangle]\Big|_{|\Psi\rangle=|\Psi_{\text{vac}}\rangle} = 0, \quad S[|\Psi_{\text{vac}}\rangle] = E_0, \quad \langle\Psi_{\text{vac}}|\Psi_{\text{vac}}\rangle = 1.}$$

(6.134)

Equivalently:

(i) (Stationarity ⇒ eigenstate) If $\delta S[|\Psi\rangle] = 0$ under the constraint $\langle\Psi|\Psi\rangle = 1$, then there exists $\lambda \in \mathbb{R}$ with $\widehat{\boldsymbol{H}}[|\Psi\rangle]|\Psi\rangle = \lambda|\Psi\rangle$ and $\lambda = S[|\Psi\rangle]$.

(ii) (Ground minimizers ⇔ vacuum manifold) Among stationary points, the global minimizers of S on the unit sphere are precisely the normalized vectors in the ground eigenspace $\mathcal{V}_0$, i.e. $\mathcal{M}_{\text{vac}}^{(\text{word})} = \{\, |\Psi\rangle : \; S[|\Psi\rangle] = E_0, \; \langle\Psi|\Psi\rangle = 1 \,\}$.

Proof. (i) Stationarity ⇒ eigenstate. Consider the constrained functional $\mathcal{L}[|\Psi\rangle, \lambda] := S[|\Psi\rangle] - \lambda(\langle\Psi|\Psi\rangle - 1)$, with $S[|\Psi\rangle] = \langle\Psi|\widehat{\boldsymbol{H}}[|\Psi\rangle]|\Psi\rangle$ (Proposition 6.57). A standard variational calculation with arbitrary $\delta|\Psi\rangle$ satisfying $\langle\Psi|\delta\Psi\rangle + \langle\delta\Psi|\Psi\rangle = 0$ gives

$$\delta\mathcal{L} = 2\,\mathrm{Re}\Big\langle \delta\Psi \Big| \Big(\widehat{\boldsymbol{H}}[|\Psi\rangle] - \lambda\,\mathbf{1}_{\text{em}}\Big)\Big|\Psi\Big\rangle = 0 \quad \forall\, \delta|\Psi\rangle.$$

Hence $\Big(\widehat{\boldsymbol{H}}[|\Psi\rangle] - \lambda\,\mathbf{1}_{\text{em}}\Big)|\Psi\rangle = 0$, so $|\Psi\rangle$ is an eigenstate with eigenvalue λ. Taking the expectation, $\lambda = \langle\Psi|\widehat{\boldsymbol{H}}[|\Psi\rangle]|\Psi\rangle = S[|\Psi\rangle]$.

(ii) Ground minimizers ⇔ vacuum manifold. By the spectral theorem and Definition 6.62, for any normalized $|\Phi\rangle$ one has $\langle\Phi|\widehat{\boldsymbol{H}}[|\Psi\rangle]|\Phi\rangle \geq E_0$, with equality iff $|\Phi\rangle \in \mathcal{V}_0 = \mathrm{Ran}\,\boldsymbol{\Pi}_0$. Using Proposition 6.57, this reads $S[|\Phi\rangle] \geq E_0$, and equality selects precisely the normalized ground eigenvectors. Thus the global minimizers of S on the unit sphere coincide with $\mathcal{M}_{\text{vac}}^{(\text{word})}$, establishing (6.134). □

Remark 6.65 (Degeneracy and projectivization)**.** If E_0 is degenerate, then $\mathcal{M}_{\text{vac}}^{(\text{word})}$ is the unit sphere in $\mathcal{V}_0$ and its projectivization satisfies $\mathbb{P}\mathcal{M}_{\text{vac}} \cong \mathbb{P}(\mathcal{V}_0)$ (Definition 6.63). The proposition still holds verbatim.

Proposition 6.66 (Gauge/holonomy covariance of the relational Hamiltonian and ground projector (wordwise))**.** *Let $g(x)$ be an endpoint gauge (fibrewise unitary) acting by conjugation on local quantities so that $d\boldsymbol{\mu}(x) \mapsto g(x)\, d\boldsymbol{\mu}(x)\, g(x)^\dagger$ and $\hat{\mathcal{G}}(x) \mapsto g(x)\,\hat{\mathcal{G}}(x)\,g(x)^\dagger$ (Propositions 6.31 and 6.36 and Definition 6.55). Let the Relational Hamiltonian word be $\widehat{\boldsymbol{H}}[|\Psi\rangle] = \int_{\mathcal{X}_q} d\boldsymbol{\mu}(x)\,\hat{\mathcal{G}}(x)$ (Definition 6.56). Denote by $\mathbf{g}$ the induced unitary on the ambient space implementing $(\mathbf{g}\,\Phi)(x) = g(x)\,\Phi(x)$. Then*

$$\boxed{\widehat{\boldsymbol{H}}[|\Psi\rangle] \;\longmapsto\; \mathbf{g}\,\widehat{\boldsymbol{H}}[|\Psi\rangle]\,\mathbf{g}^\dagger, \qquad \boldsymbol{\Pi}_0 \;\longmapsto\; \mathbf{g}\,\boldsymbol{\Pi}_0\,\mathbf{g}^\dagger,} \tag{6.135}$$

where $\boldsymbol{\Pi}_0 = \mathbf{1}_{\{E_0\}}\Big(\widehat{\boldsymbol{H}}[|\Psi\rangle]\Big)$ (Definition 6.62). The same covariance holds for holonomies: for any loop γ based at x with holonomy $h_\gamma = \mathcal{U}(x \leftarrow x; \gamma)$ (eqs. (4.11) and (4.14)), one has $\widehat{\boldsymbol{H}} \mapsto \mathbf{h}_\gamma\,\widehat{\boldsymbol{H}}\,\mathbf{h}_\gamma^\dagger$ and $\boldsymbol{\Pi}_0 \mapsto \mathbf{h}_\gamma\,\boldsymbol{\Pi}_0\,\mathbf{h}_\gamma^\dagger$.

Proof. By the stated covariance of $d\boldsymbol{\mu}$ and $\hat{\mathcal{G}}$ and HS–covariant integration (Corollary 6.37),

$$\widehat{\boldsymbol{H}} \;\mapsto\; \int_{\mathcal{X}_q} g(x)\, d\boldsymbol{\mu}(x)\,\hat{\mathcal{G}}(x)\, g(x)^\dagger = \mathbf{g}\Big(\int_{\mathcal{X}_q} d\boldsymbol{\mu}(x)\,\hat{\mathcal{G}}(x)\Big)\mathbf{g}^\dagger = \mathbf{g}\,\widehat{\boldsymbol{H}}\,\mathbf{g}^\dagger.$$

Unitary equivariance of spectral calculus then gives $\mathbf{1}_{\{E_0\}}(\mathbf{g}\,\widehat{\boldsymbol{H}}\,\mathbf{g}^\dagger) = \mathbf{g}\,\mathbf{1}_{\{E_0\}}(\widehat{\boldsymbol{H}})\,\mathbf{g}^\dagger$, proving the projector transformation. The holonomy case is identical with $g(x)$ replaced by the fibrewise unitary h_γ. □

Corollary 6.67 (Gauge/holonomy invariance of vacuum rays)**.** *Let $\mathbb{P}\mathcal{M}_{\mathrm{vac}}$ be the set of vacuum rays (Definition 6.63). Under the transformations in eq. (6.135), the set $\mathbb{P}\mathcal{M}_{\mathrm{vac}}$ is invariant: if $|\Psi_{\mathrm{vac}}\rangle \in \mathcal{M}_{\mathrm{vac}}^{(\mathrm{word})}$, then $\mathbf{g}|\Psi_{\mathrm{vac}}\rangle$ and $\mathbf{h}_\gamma|\Psi_{\mathrm{vac}}\rangle$ are also vacuum vectors, and the corresponding rays coincide in $\mathbb{P}\mathcal{M}_{\mathrm{vac}}$.*

Proof. If $|\Psi_{\mathrm{vac}}\rangle \in \mathrm{Ran}\,\mathbf{\Pi}_0$, then $\mathbf{g}|\Psi_{\mathrm{vac}}\rangle \in \mathrm{Ran}\,(\mathbf{g}\mathbf{\Pi}_0\mathbf{g}^\dagger)$ by eq. (6.135). Thus $\mathbf{g}|\Psi_{\mathrm{vac}}\rangle$ is a vacuum vector for the transformed Hamiltonian and, since $\mathbf{g}$ is unitary, its ray lies in $\mathbb{P}\mathcal{M}_{\mathrm{vac}}$. The holonomy case is identical. □

Remark 6.68 (Triadic covariance and expectation reduction)**.** The covariance in Proposition 6.66 holds componentwise in the corr/vac/spec triad (eq. (5.86)), since both $d\boldsymbol{\mu}$ and $\hat{\mathcal{G}}$ admit triadic decompositions and Corollary 6.37 respects the sector split. At expectation level the bar-fields are gauge invariant by construction, so the induced statement on $\mathbb{P}\mathcal{M}_{\mathrm{vac}}$ persists after taking expectations.

HS extension (two-law control)

Definition 6.69 (Branch Hamiltonians (wordwise))**.** Let $\mathcal{H}_L = \mathrm{span}\{|0\rangle_L, |1\rangle_L\}$ with matrix units $P_0 = |0\rangle\langle 0|$, $P_1 = |1\rangle\langle 1|$, $E_{01} = |0\rangle\langle 1|$, $E_{10} = |1\rangle\langle 0|$. For each branch $k \in \{0, 1\}$, let the branch data be $\big(\hat{\mathbf{\Sigma}}^{(k)}, \hat{\mathbf{G}}_\Lambda^{(k)}\big)$, and define

$$\boxed{\widehat{\boldsymbol{H}}^{(k)}[|\Psi\rangle] := \int_{\mathcal{X}_q} d\boldsymbol{\mu}^{(k)}(x)\ \hat{\mathcal{G}}^{(k)}(x), \qquad \hat{\mathcal{G}}^{(k)}(x) := \mathrm{Tr}_{\mathrm{FD}}\Big(\hat{\mathbf{G}}_\Lambda^{(k)}(x,x)\Big),} \tag{6.136}$$

as in eq. (6.121), with $d\boldsymbol{\mu}^{(k)}(x)$ the branchwise relational volume element (cf. Propositions 6.31 and 6.36) and $\mathrm{Tr}_{\mathrm{FD}}$ the field-diagonal trace (Definition 6.45).

Definition 6.70 (Mixing Hamiltonian (wordwise))**.** Define the *mixing Hamiltonian* by

$$\boxed{\hat{H}_{\mathrm{mix}} \ := \ \int_{\mathcal{X}_q} d\boldsymbol{\mu}(x)\ \mathrm{Tr}_{\mathrm{op}}\Big(\mathrm{Tr}_{\mathrm{dialectic}}\big(\hat{G}_{\mathbf{\Phi\Theta}}(x,x)\big)\Big),} \tag{6.137}$$

where $\mathrm{Tr}_{\mathrm{dialectic}}$ contracts the Fourfold modal block and $\mathrm{Tr}_{\mathrm{op}}$ is the operator trace consistent with Definition 6.45. For any dial phase $\varphi \in \mathbb{R}$, the Hermitian part $\mathrm{Re}\big(e^{-i\varphi}\hat{H}_{\mathrm{mix}}\big)$ is self-adjoint.

Definition 6.71 (Two-law HS extension on $\mathcal{H}_{\mathrm{em}} \otimes \mathcal{H}_L$)**.** Let $|+_\varphi\rangle_L := \frac{1}{\sqrt{2}}\big(|0\rangle_L + e^{i\varphi}|1\rangle_L\big)$. The *controlled (two-law) Hamiltonian* is the self-adjoint operator

$$\boxed{\widehat{\boldsymbol{H}}^{[L]}(\varphi) := P_0 \otimes \widehat{\boldsymbol{H}}^{(0)} \ + \ P_1 \otimes \widehat{\boldsymbol{H}}^{(1)} \ + \ \frac{1}{2}\Big(e^{-i\varphi}E_{01} \otimes \hat{H}_{\mathrm{mix}} \ + \ e^{+i\varphi}E_{10} \otimes \hat{H}_{\mathrm{mix}}^\dagger\Big).} \tag{6.138}$$

We also define the *controlled action* by

$$S_\varphi^{[L]}[|\Psi\rangle] \ := \ \Big\langle \Psi \otimes +_\varphi \Big|\ \widehat{\boldsymbol{H}}^{[L]}(\varphi)\ \Big|\Psi \otimes +_\varphi\Big\rangle, \tag{6.139}$$

whenever $|\Psi\rangle$ lies in the common domain of the local densities (Corollary 6.37).

Proposition 6.72 (Self-adjointness, gauge covariance, and triadic structure)**.** *The operator* $\widehat{\boldsymbol{H}}^{[L]}(\varphi)$ *in* (6.138) *is self-adjoint. Under endpoint gauge/holonomy transformations (Proposition 6.66), one has*

$$\widehat{\boldsymbol{H}}^{(k)} \mapsto \mathbf{g}\,\widehat{\boldsymbol{H}}^{(k)}\,\mathbf{g}^{\dagger}, \qquad \hat{H}_{\mathrm{mix}} \mapsto \mathbf{g}\,\hat{H}_{\mathrm{mix}}\,\mathbf{g}^{\dagger}, \qquad \widehat{\boldsymbol{H}}^{[L]}(\varphi) \mapsto (\mathbf{1}_L \otimes \mathbf{g})\,\widehat{\boldsymbol{H}}^{[L]}(\varphi)\,(\mathbf{1}_L \otimes \mathbf{g})^{\dagger}.$$

Moreover, all ingredients admit corr/vac/spec decompositions preserved by the construction (eq. (5.86)).

Proof. Self-adjointness follows since $P_k = P_k^{\dagger}$, $E_{10} = E_{01}^{\dagger}$, each $\widehat{\boldsymbol{H}}^{(k)}$ is self-adjoint by (6.136), and the off-diagonal sum is a Hermitian pair. Gauge covariance is inherited from Proposition 6.66 and linearity of HS–covariant integration (Corollary 6.37). The triadic statement follows from the decompositions of $d\boldsymbol{\mu}^{(k)}$ and $\hat{\mathcal{G}}^{(k)}$, and of $\hat{G}_{\boldsymbol{\Phi\Theta}}$ entering (6.137). □

Proposition 6.73 (Expectation-level two-law control (interference dial))**.** *For any normalized* $|\Psi\rangle$ *in the common domain, with the branch actions* $S^{(k)}[|\Psi\rangle] := \langle\Psi|\widehat{\boldsymbol{H}}^{(k)}[|\Psi\rangle]|\Psi\rangle$ *(Proposition 6.57) and the mixing functional* $M[|\Psi\rangle] := \langle\Psi|\hat{H}_{\mathrm{mix}}|\Psi\rangle$*, one has*

$$\boxed{S_{\varphi}^{[L]}[|\Psi\rangle] \;=\; \frac{S^{(0)}[|\Psi\rangle] + S^{(1)}[|\Psi\rangle]}{2} \;+\; \mathrm{Re}\Big(e^{-i\varphi}\,M[|\Psi\rangle]\Big).} \tag{6.140}$$

In particular, choosing $\varphi = \arg M[|\Psi\rangle]$ *maximizes the interference term, while* $\varphi = \arg M[|\Psi\rangle] \pm \frac{\pi}{2}$ *suppresses it to zero.*

Proof. Expand $\widehat{\boldsymbol{H}}^{[L]}(\varphi)$ in the $\{|0\rangle_L, |1\rangle_L\}$ basis and evaluate (6.139) with $|+_{\varphi}\rangle_L$; the diagonal contributes $\frac{1}{2}\big(S^{(0)} + S^{(1)}\big)$ via Proposition 6.57, and the off-diagonal yields $\frac{1}{2}\big(e^{-i\varphi}M + e^{+i\varphi}\overline{M}\big) = \mathrm{Re}(e^{-i\varphi}M)$. □

Corollary 6.74 (Branch recovery and extremal dials)**.** *If the ancilla is prepared in* $|0\rangle_L$ *(resp.* $|1\rangle_L$*), then* $S_{\varphi}^{[L]}[|\Psi\rangle] = S^{(0)}[|\Psi\rangle]$ *(resp.* $S^{(1)}[|\Psi\rangle]$*). For* $|+_{\varphi}\rangle_L$*,* (6.140) *interpolates between the arithmetic mean and the extremal values by dialling* φ*.*

Remark 6.75 (Domain, gauge invariance after reduction, and wordwise consistency)**.** (i) All expectations are well-defined by Corollary 6.37. (ii) After taking expectations in the ringed-ket state, gauge covariance reduces to gauge *invariance* of the scalar functionals (Remark 6.35). (iii) The construction is the holosymmetric (wordwise) lift of the standard two-branch control, compatible with Section 6.1.2.

HS extension (two-law control): Full Hamiltonian and reduction

Definition 6.76 (Full HS Hamiltonian (two-law mixing))**.** On $\mathcal{H}_L \otimes \mathcal{H}_{\mathrm{em}}$ with $P_0 = |0\rangle\langle 0|$, $P_1 = |1\rangle\langle 1|$, $E_{01} = |0\rangle\langle 1|$, $E_{10} = |1\rangle\langle 0|$, and branch Hamiltonians $\widehat{\boldsymbol{H}}^{(k)}[|\Psi\rangle]$ from Definition 6.69 and eq. (6.136) and mixing operator $\hat{H}_{\mathrm{mix}}$ from Definition 6.70 and eq. (6.137), define

$$\boxed{\widehat{\boldsymbol{H}}^{\mathrm{HS}}[|\Psi\rangle] \;:=\; P_0 \otimes \widehat{\boldsymbol{H}}^{(0)}[|\Psi\rangle] \;+\; P_1 \otimes \widehat{\boldsymbol{H}}^{(1)}[|\Psi\rangle] \;+\; E_{01} \otimes \hat{H}_{\mathrm{mix}} \;+\; E_{10} \otimes \hat{H}_{\mathrm{mix}}^{\dagger}.} \tag{6.141}$$

Proposition 6.77 (Self-adjointness and gauge/holonomy covariance)**.** *The operator* $\widehat{\boldsymbol{H}}^{\mathrm{HS}}$ *in* (6.141) *is self-adjoint. Under endpoint gauge/holonomy transformations (Proposition 6.66),*

$$\widehat{\boldsymbol{H}}^{(k)} \mapsto \mathbf{g}\,\widehat{\boldsymbol{H}}^{(k)}\,\mathbf{g}^{\dagger}, \qquad \hat{H}_{\mathrm{mix}} \mapsto \mathbf{g}\,\hat{H}_{\mathrm{mix}}\,\mathbf{g}^{\dagger}, \qquad \widehat{\boldsymbol{H}}^{\mathrm{HS}} \mapsto (\mathbf{1}_L \otimes \mathbf{g})\,\widehat{\boldsymbol{H}}^{\mathrm{HS}}\,(\mathbf{1}_L \otimes \mathbf{g})^{\dagger}.$$

Moreover, the corr/vac/spec triadic decomposition is preserved (eq. (5.86)).

Proof. Self-adjointness: $P_k = P_k^{\dagger}$, $E_{10} = E_{01}^{\dagger}$, each $\widehat{\boldsymbol{H}}^{(k)}$ is self-adjoint by Definition 6.69, and $E_{01}\otimes \hat{H}_{\mathrm{mix}} + E_{10}\otimes \hat{H}_{\mathrm{mix}}^{\dagger}$ is Hermitian for arbitrary $\hat{H}_{\mathrm{mix}}$. Gauge/holonomy covariance is inherited from Proposition 6.66 and linearity of HS–covariant integration (Corollary 6.37). Triadic preservation is immediate from the triadic splits of the ingredients. □

Proposition 6.78 (Reduction under the dial state $|+_\varphi\rangle_L$)**.** *Prepare* $|+_\varphi\rangle_L = \frac{1}{\sqrt{2}}(|0\rangle_L + e^{i\varphi}|1\rangle_L)$. *Then the compressed operator on* $\mathcal{H}_{\mathrm{em}}$ *is*

$$\boxed{\hat{H}^{\langle \mathrm{HS}\rangle}[|\Psi\rangle;\varphi] := \Big\langle +_\varphi \Big|\, \widehat{\boldsymbol{H}}^{\mathrm{HS}}[|\Psi\rangle] \,\Big|\, +_\varphi \Big\rangle = \tfrac{1}{2}\Big(\widehat{\boldsymbol{H}}^{(0)}[|\Psi\rangle] + \widehat{\boldsymbol{H}}^{(1)}[|\Psi\rangle]\Big) + \mathrm{Re}\Big(e^{-i\varphi}\hat{H}_{\mathrm{mix}}\Big),}$$

(6.142)

which matches eq. (6.125) verbatim, with $\hat{H}_{\mathrm{mix}}$ *in place of any generic symbol.*

Proof. Using $\langle +_\varphi|P_0|+_\varphi\rangle = \langle +_\varphi|P_1|+_\varphi\rangle = \frac{1}{2}$ and $\langle +_\varphi|E_{01}|+_\varphi\rangle = \frac{1}{2}e^{-i\varphi}$, $\langle +_\varphi|E_{10}|+_\varphi\rangle = \frac{1}{2}e^{+i\varphi}$, sandwich (6.141) between $\langle +_\varphi|$ and $|+_\varphi\rangle$:

$$\hat{H}^{\langle \mathrm{HS}\rangle} = \tfrac{1}{2}\widehat{\boldsymbol{H}}^{(0)} + \tfrac{1}{2}\widehat{\boldsymbol{H}}^{(1)} + \tfrac{1}{2}e^{-i\varphi}\hat{H}_{\mathrm{mix}} + \tfrac{1}{2}e^{+i\varphi}\hat{H}_{\mathrm{mix}}^{\dagger} = \tfrac{1}{2}(\widehat{\boldsymbol{H}}^{(0)} + \widehat{\boldsymbol{H}}^{(1)}) + \mathrm{Re}(e^{-i\varphi}\hat{H}_{\mathrm{mix}}).$$

□

Corollary 6.79 (Expectation/action reduction)**.** *For any normalized* $|\Psi\rangle$ *in the common domain,*

$$\Big\langle \Psi \Big| \hat{H}^{\langle \mathrm{HS}\rangle}[|\Psi\rangle;\varphi] \Big| \Psi \Big\rangle = \frac{S^{(0)}[|\Psi\rangle] + S^{(1)}[|\Psi\rangle]}{2} + \mathrm{Re}\Big(e^{-i\varphi}M[|\Psi\rangle]\Big),$$

with $S^{(k)}[|\Psi\rangle] = \langle\Psi|\widehat{\boldsymbol{H}}^{(k)}[|\Psi\rangle]|\Psi\rangle$ *(Proposition 6.57) and* $M[|\Psi\rangle] = \langle\Psi|\hat{H}_{\mathrm{mix}}|\Psi\rangle$, *i.e. the formula of Proposition 6.73 and eq.* (6.140).

Remark 6.80 (Dial choices, branch recovery, and invariances)**.** (i) $\varphi = \arg M[|\Psi\rangle]$ maximizes the interference term; $\varphi = \arg M[|\Psi\rangle] \pm \frac{\pi}{2}$ nulls it (cf. Proposition 6.73). (ii) Preparing $|0\rangle_L$ (resp. $|1\rangle_L$) recovers the branch Hamiltonian $\widehat{\boldsymbol{H}}^{(0)}$ (resp. $\widehat{\boldsymbol{H}}^{(1)}$) as in Corollary 6.74. (iii) Gauge/holonomy covariance of $\widehat{\boldsymbol{H}}^{\mathrm{HS}}$ (Proposition 6.77) descends to the compressed operator, while expectation-level quantities are gauge invariant by the construction of the bar-fields (Remark 6.35).

HS vacuum manifold at fixed dial

Definition 6.81 (Fixed-dial ground sector (HS–reduced))**.** Let $\hat{H}^{\langle \mathrm{HS}\rangle}(\varphi)$ be the reduced operator from Proposition 6.78 and eq. (6.142). Define the fixed–dial ground energy and projector by

$$\boxed{E_0^{\langle \mathrm{HS}\rangle}(\varphi) := \inf \mathrm{spec}\Big(\hat{H}^{\langle \mathrm{HS}\rangle}(\varphi)\Big), \qquad \Pi_0^{\langle \mathrm{HS}\rangle}(\varphi) := \mathbf{1}_{\{E_0^{\langle \mathrm{HS}\rangle}(\varphi)\}}\Big(\hat{H}^{\langle \mathrm{HS}\rangle}(\varphi)\Big).}$$

(6.143)

Definition 6.82 (HS vacuum manifold at fixed dial)**.** With Definition 6.81, the *HS vacuum manifold at fixed dial* is

$$\boxed{\mathcal{M}_{\mathrm{vac}}^{\langle \mathrm{HS}\rangle}(\varphi) := \Big\{ |\Psi\rangle : \ \hat{H}^{\langle \mathrm{HS}\rangle}(\varphi)\,|\Psi\rangle = E_0^{\langle \mathrm{HS}\rangle}(\varphi)\,|\Psi\rangle,\ \langle\Psi|\Psi\rangle = 1 \Big\} = \mathbb{P}\Big(\mathrm{Ran}\,\Pi_0^{\langle \mathrm{HS}\rangle}(\varphi)\Big).}$$

(6.144)

Proposition 6.83 (Variational–spectral equivalence at fixed dial)**.** *Let $S_\varphi^{[L]}[|\Psi\rangle]$ be the controlled action from Definition 6.71, Proposition 6.73, and eq. (6.140). Then for normalized states,*

$$\boxed{|\Psi_{\mathrm{vac}}\rangle \in \mathcal{M}_{\mathrm{vac}}^{\langle \mathrm{HS}\rangle}(\varphi) \iff \delta S_\varphi^{[L]}[|\Psi\rangle]\Big|_{|\Psi\rangle=|\Psi_{\mathrm{vac}}\rangle} = 0, \quad S_\varphi^{[L]}[|\Psi_{\mathrm{vac}}\rangle] = E_0^{\langle \mathrm{HS}\rangle}(\varphi).}$$

(6.145)

Proof. By Proposition 6.78 and eq. (6.142), $\hat{H}^{\langle \mathrm{HS}\rangle}(\varphi)$ is the compression of $\widehat{\boldsymbol{H}}^{\mathrm{HS}}$ to $|+_\varphi\rangle_L$. By Proposition 6.73, $S_\varphi^{[L]}[|\Psi\rangle] = \langle\Psi|\hat{H}^{\langle \mathrm{HS}\rangle}(\varphi)|\Psi\rangle$, so the constrained stationarity on the unit sphere implies the eigenvalue equation with eigenvalue equal to the expectation; minimizers give the ground value $E_0^{\langle \mathrm{HS}\rangle}(\varphi)$, as in Proposition 6.64. □

Proposition 6.84 (Gauge/holonomy invariance of fixed–dial vacuum rays)**.** *Under endpoint gauge/holonomy transformations (Propositions 6.66 and 6.77), $\hat{H}^{\langle \mathrm{HS}\rangle}(\varphi) \mapsto \mathbf{g}\,\hat{H}^{\langle \mathrm{HS}\rangle}(\varphi)\,\mathbf{g}^\dagger$. Hence $\Pi_0^{\langle \mathrm{HS}\rangle}(\varphi) \mapsto \mathbf{g}\,\Pi_0^{\langle \mathrm{HS}\rangle}(\varphi)\,\mathbf{g}^\dagger$ and $\mathbb{P}\Big(\mathrm{Ran}\,\Pi_0^{\langle \mathrm{HS}\rangle}(\varphi)\Big)$ is invariant.*

Proof. Conjugation covariance follows from Proposition 6.77 and compression (Proposition 6.78). Spectral calculus is equivariant under unitary conjugation, hence projector covariance; unitarity preserves rays. □

Remark 6.85 (Dial continuity and bounds)**.** The map $\varphi \mapsto \hat{H}^{\langle \mathrm{HS}\rangle}(\varphi)$ is continuous and 2π–periodic by eq. (6.142). For any normalized $|\Psi\rangle$, $S_\varphi^{[L]}[|\Psi\rangle] = \frac{1}{2}\Big(S^{(0)} + S^{(1)}\Big) + \mathrm{Re}(e^{-i\varphi}M)$ (eq. (6.140)), so its φ–extrema occur at $\varphi = \arg M$ and $\arg M \pm \pi$. Consequently,

$$\frac{E_0^{(0)} + E_0^{(1)}}{2} - \|\hat{H}_{\mathrm{mix}}\| \ \le\ E_0^{\langle \mathrm{HS}\rangle}(\varphi) \ \le\ \frac{E_0^{(0)} + E_0^{(1)}}{2} + \|\hat{H}_{\mathrm{mix}}\|,$$

with operator norm $\|\cdot\|$; this follows from eq. (6.142) and the variational (min–max) principle.

Proposition 6.86 (Consistency: dial–augmented action density and gauge covariance)**.** *For any normalized $|\Psi\rangle$ in the common domain, the reduced operator $\hat{H}^{\langle \mathrm{HS}\rangle}(\varphi)$ of Proposition 6.78 and eq. (6.142) reproduces the* dial–augmented action*:*

$$\boxed{\Big\langle \Psi \Big| \hat{H}^{\langle \mathrm{HS}\rangle}(\varphi) \Big| \Psi \Big\rangle = \ S_\varphi^{[L]}[|\Psi\rangle] = \ \int_{\mathcal{X}_q} \sqrt{-\det \bar{\Sigma}^{(\Psi)}(x)}\ \mathsf{G}_\varphi^{(\Psi)}(x)\, d^d x,}$$

(6.146)

where $\bar{\Sigma}^{(\Psi)}$ is the expectation unified tensor (eq. (5.34) and Definition 5.77) and $\mathsf{G}_\varphi^{(\Psi)}$ is the dial–augmented *density obtained in Section 6.1.12 by adding the controlled mixed–sector contribution. Moreover, $\hat{H}^{\langle \mathrm{HS}\rangle}(\varphi)$ is gauge/holonomy covariant,*

$$\hat{H}^{\langle \mathrm{HS}\rangle}(\varphi) \ \mapsto\ \mathbf{g}\,\hat{H}^{\langle \mathrm{HS}\rangle}(\varphi)\,\mathbf{g}^\dagger,$$

as a consequence of the covariance of $d\boldsymbol{\mu}$ and $\hat{\mathcal{G}}$ (Propositions 6.31 and 6.36) together with HS–covariant integration (Corollary 6.37).

Proof. The identity (6.146) is the expectation-level version of Proposition 6.78 combined with Proposition 6.73 and eq. (6.140) and the action identification in Proposition 6.57. Gauge/holonomy covariance follows from Proposition 6.77 and the unitary equivariance of spectral calculus; expectation-level expressions are then gauge invariant by Remark 6.35. □

Corollary 6.87 (Causality is unchanged by the dial)**.** *Let* $\bar{g}^{\mathrm{Lor}}_{\mu\nu} = \frac{1}{2}(\bar{\Sigma}_{\mu\nu} + \bar{\Sigma}_{\nu\mu})$ *(Definition 5.77) and let effective cones be defined as in Section 5.4.2. The dial* φ *in eq. (6.142) modifies only the* mixed–sector integrand *of the action density (Section 6.1.12) and leaves* $\bar{\Sigma}_{\mu\nu}$ *(hence* $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$*) unchanged. Therefore the null sets, timelike/spacelike classification, and apex conditions are independent of* φ *(cf. Section 5.4.2).*

Remark 6.88 (Vacuum geometry, sector diagnostics, and dial stability)**.** (i) If the ground eigenspace $\mathcal{V}_0$ has finite dimension k, then the set of vacuum rays satisfies $\mathbb{P}\mathcal{M}_{\mathrm{vac}} \cong \mathbb{CP}^{k-1}$ (Definition 6.63); for $k = 1$ the vacuum is unique up to phase.

(ii) The corr/vac/spec triadic decomposition (eq. (5.86)) of $\widehat{\boldsymbol{H}}$ restricts to $\mathcal{V}_0$ and provides *sector-resolved* diagnostics of symmetry-protected degeneracies. In particular, if a word-symmetry $\mathbf{U}$ (holosymmetric, channel-resolved) commutes with the relevant sector blocks (Section 6.11.7 and Propositions 6.257 and 6.276), then $\mathcal{V}_0$ carries a nontrivial $\mathbf{U}$–representation and degeneracy is protected unless the commuting relation is broken by deformations of that sector.

(iii) Dial stability: from eq. (6.142) and the min–max principle,

$$\left|E_0^{\langle \mathrm{HS}\rangle}(\varphi_1) - E_0^{\langle \mathrm{HS}\rangle}(\varphi_2)\right| \;\le\; \left\|\mathrm{Re}\big(e^{-i\varphi_1}\hat{H}_{\mathrm{mix}}\big) - \mathrm{Re}\big(e^{-i\varphi_2}\hat{H}_{\mathrm{mix}}\big)\right\| \;\le\; 2\,\|\hat{H}_{\mathrm{mix}}\|\,\left|\sin\tfrac{\varphi_1-\varphi_2}{2}\right|,$$

so the dial dependence is continuous and 2π–periodic; extremal dials are those of Proposition 6.73.

(iv) All consistency statements above persist under holonomy conjugations and after taking expectations (gauge invariance), by Proposition 6.66 and Remark 6.35.

6.1.19 Quantum Circuit Realization of the Vacuum Manifold

Circuit A (base action/energy estimator). Discretise $\int_{\mathcal{X}_q} d\boldsymbol{\mu}(x)\,\hat{\mathcal{G}}(x)$ by nodes (x_j, w_j) and block-encode $O := \sum_j w_j\,\hat{\mathcal{G}}(x_j)/\alpha$ with known scale α. A Hadamard test on a probe qubit returns $\mathrm{Re}\langle\Psi|O|\Psi\rangle$, hence $S[|\Psi\rangle]$.



Here:

- $\mathrm{PREP}_w : |0\rangle_{\mathsf{X}} \mapsto \sum_j \sqrt{\frac{w_j}{\sum_k w_k}}\,|j\rangle_{\mathsf{X}}$ encodes the reduced HS measure (Definition 6.34).
- BlockEncode(O) uses your FD-trace gadget (Definition 6.45) to realise $\hat{\mathcal{G}}(x_j) = \mathrm{Tr}_{\mathrm{FD}}(\hat{\mathbf{G}}_\Lambda(x_j, x_j))$ coherently, then LCU.

- $\mathbb{E}[Z_{\mathsf{P}}] = \mathrm{Re}\langle\Psi|O|\Psi\rangle$; an added phase on P gives the imaginary part.

Circuit B (law-controlled reduced estimator). We estimate

$$\langle\Psi|\hat{H}^{\langle\mathrm{HS}\rangle}(\varphi)|\Psi\rangle = \tfrac{1}{2}\sum_{k=0}^{1}\langle\Psi|\widehat{\boldsymbol{H}}^{(k)}|\Psi\rangle + \mathrm{Re}\Big(e^{-i\varphi}\langle\Psi|\hat{H}_{\mathrm{mix}}|\Psi\rangle\Big),$$

as in eq. (6.142), by summing two Hadamard tests:





Readout. Add the two estimates (rescaled by $\alpha_0, \alpha_1, \alpha_{\mathrm{mix}}$) to obtain $\langle\Psi|\hat{H}^{\langle\mathrm{HS}\rangle}(\varphi)|\Psi\rangle$. The $R_z(\varphi)$ on the probe implements $\mathrm{Re}\big(e^{-i\varphi}\,\cdot\,\big)$ in the mixing term, with

$$\hat{H}_{\mathrm{mix}} := \int_{\mathcal{X}_q} d\boldsymbol{\mu}(x)\ \mathrm{Tr}_{\mathrm{op}}\Big(\mathrm{Tr}_{\mathrm{dialectic}}\big(\hat{G}_{\boldsymbol{\Phi\Theta}}(x,x)\big)\Big) \quad \text{(eq. (6.122))}.$$

Remark 6.89 (Implementation notes for the circuits of Section 6.1.16)**.**

1. **FD-trace gadget.** The nested trace $\mathrm{Tr}_{\mathrm{FD}} = \mathrm{Tr}_{\mathrm{field}} \circ \mathrm{Tr}_{\mathrm{dialectic}}$ (Definition 6.45) is implemented coherently by preparing a uniform superposition on the finite field/dialectic index registers, applying the block-select unitaries that realise the correlator entries, and unpreparing. This yields the diagonal field–modal contraction consistent with Section 6.1.10 and Definition 6.45 and the scalar density (6.98).
2. **Weights and HS measure.** The quadrature state PREP_w on the space register encodes the reduced HS measure (Definition 6.34):

$$\mathrm{PREP}_w :\ |0\rangle_{\mathsf{X}} \longmapsto \sum_j \sqrt{\frac{w_j}{\sum_k w_k}}\,|j\rangle_{\mathsf{X}}, \qquad w_j := \sqrt{-\det\bar{\Sigma}^{(\Psi)}(x_j)}\,\Delta x_j.$$

   Thus the discretised sum reproduces the reduction of $d\boldsymbol{\mu}$ in the expectation-level integral.

3. **Tomography dials.** When the θ-dial of the fourfold trace is employed (Section 6.1.12), the mixing Hamiltonian is taken in the θ-resolved form

$$\boxed{\hat{H}_{\mathrm{mix}}(\theta) := \int_{\mathcal{X}_q} d\boldsymbol{\mu}(x)\ \mathrm{Tr}_{\mathrm{op}}\Big[\mathrm{Tr}_{\mathrm{dialectic}}\Big(\cos\theta\,(\hat{G}_{\Phi\Theta}(x,x)+\hat{G}_{\Theta\Phi}(x,x)) + \sin\theta\, i\,(\hat{G}_{\Phi\Theta}(x,x)-\hat{G}_{\Theta\Phi}(x,x))\Big)\Big].} \tag{6.147}$$

Block-encoding $\hat{H}_{\mathrm{mix}}(\theta)$ uses the same FD-trace and weight mechanisms as in the base case; the probe rotation $R_z(\varphi)$ implements the readout $\mathrm{Re}\big(e^{-i\varphi}\,\langle\Psi|\hat{H}_{\mathrm{mix}}(\theta)|\Psi\rangle\big)$, matching the reduced HS estimator structure in eq. (6.125).

From definition to executable procedure. The quantum circuit of Section 6.1.16 realises the variational–spectral equivalence operationally. By block-encoding the HS Relational Hamiltonian word and estimating the expectation value $\langle\Psi|\,\widehat{\boldsymbol{H}}[|\Psi\rangle]\,|\Psi\rangle$, the circuit computes the Relational Action $S[|\Psi\rangle]$ (Proposition 6.57). Consequently, preparing a variational family $|\Psi(\vartheta)\rangle$ and minimising the circuit's estimate of S identifies ground states of $\hat{H}$ and hence points on the vacuum manifold $\mathcal{M}_{\mathrm{vac}}$ (Definition 6.54 and Section 6.1.17).

Algorithmic backreaction and the nonlinear fixed point. A distinctive aspect of the framework is the state-dependence of the geometry through the HS measure. In the circuit, this enters as the weight state PREP_w, whose amplitudes encode the reduction of $d\boldsymbol{\mu}$ to $d\mu_{\bar{\Sigma}(\Psi)}$ (Definition 6.34). Iterating the loop "prepare $|\Psi(\vartheta)\rangle\to$ estimate $S[|\Psi\rangle]\to$ update ϑ" implements the Euler–Lagrange stationarity condition and its nonlinear eigenoperator form (Lemma 6.41) as a concrete hybrid (quantum–classical) fixed-point algorithm.

Tomographic access to mixed-sector structure. The same architecture provides phase-programmable diagnostics of inter-field couplings. With the law register and the dials (φ,θ) (Section 6.1.12), the circuit evaluates the reduced HS Hamiltonian $\hat{H}^{\langle\mathrm{HS}\rangle}[|\Psi\rangle;\varphi]$ (eq. (6.125)) and isolates exchange–symmetric and exchange–antisymmetric contributions via the scalars $\mathsf{S}^{(\Psi)}(x)$ and $\mathsf{C}^{(\Psi)}(x)$. In particular, the derivatives with respect to θ at fixed φ implement the tomography formulas that select "anticommutator-like" and "commutator-like" mixes (cf. Section 6.1.12).

Gauge covariance and causal neutrality. Because the FD-trace and HS measure are used precisely as defined (Definitions 6.34 and 6.45), the block-encodings inherit endpoint gauge covariance, and on flat/central patches the measure is central (Propositions 6.31 and 6.36). The control dials affect only the correlator sector and not the HS measure or the effective Lorentzian metric, hence the light-cone assignment remains unchanged (Section 5.4.2 and Corollary 6.53). The resulting estimates are therefore representation-safe and physically interpretable.

From ground energy to moduli. Once a minimiser is prepared, the ground-state eigenspace $\mathcal{V}_0 = \ker(\hat{H} - E_0\mathbb{I}_{\mathrm{em}})$ and the vacuum manifold $\mathcal{M}_{\mathrm{vac}}$ (Definition 6.61) can be probed by small-oscillation or Hessian-style analyses around the circuit-prepared state, giving access

to the local curvature of the action (stability, masses) and to degeneracies (the geometry of $\mathbb{P}(\mathcal{V}_0)$). Thus the circuit not only finds vacuum states but also provides a pathway to their spectral and geometric characterisation.

A common computational grammar. The circuit maps the holosymmetric wordwise primitives—selectors and transport (Section 4.5), FD-trace (Definition 6.45 and Section 6.1.10), and HS measure (Definition 6.34)—into a compact set of hardware operations: controlled dressings, LCU block-encodings, weight-state quadrature, and Hadamard tests. This uniform mapping renders distinct chronogeometric hypotheses directly comparable within a single computational pipeline while remaining faithful to the formal constructions that define the action, the Hamiltonian, and the vacuum manifold.

6.1.20 An Operator-Theoretic Definition of the Tangent Bundle

To describe the dynamics of quantum fields fluctuating around vacuum states, we must first rigorously define the space of all infinitesimal deformations of the vacuum manifold. While this is often done using curves on a manifold [84, 249], a more fundamental approach in the pqs framework derives this structure directly from the geometry of the emergent Hilbert space, $\mathcal{H}_{\text{em}}$ [407].

Definition 6.90 (Operator-Theoretic Tangent Bundle of the Vacuum Manifold)**.** Let $\mathcal{M}_{\text{vac}}$ be the vacuum manifold, defined as the unit sphere within the ground-state eigenspace $\mathcal{V}_0 \subset \mathcal{H}_{\text{em}}$ (eq. (6.129)) [407]. The *operator-theoretic tangent bundle* to $\mathcal{M}_{\text{vac}}$, denoted $T\mathcal{M}_{\text{vac}}$, is the Hilbert sub-bundle whose fiber at each point $|\Omega\rangle \in \mathcal{M}_{\text{vac}}$ is the tangent space $T_{|\Omega\rangle}\mathcal{M}_{\text{vac}}$. This tangent space is defined as the set of all vectors in $\mathcal{V}_0$ that are orthogonal to the vacuum state $|\Omega\rangle$ [375, 407]:

$$\boxed{T_{|\Omega\rangle}\mathcal{M}_{\text{vac}} := \{|V\rangle \in \mathcal{V}_0 \mid \text{Re}\langle\Omega|V\rangle = 0\}\,.} \tag{6.148}$$

The total space of the bundle is the disjoint union of these fibers [372]:

$$\boxed{T\mathcal{M}_{\text{vac}} := \bigsqcup_{|\Omega\rangle\in\mathcal{M}_{\text{vac}}} T_{|\Omega\rangle}\mathcal{M}_{\text{vac}}.} \tag{6.149}$$

Interpretation and Geometric Motivation This definition provides a conceptually rich and physically grounded picture of the vacuum structure. A tangent vector $|V\rangle \in T_{|\Omega\rangle}\mathcal{M}_{\text{vac}}$ represents an infinitesimal, physically meaningful deformation of the vacuum state $|\Omega\rangle$. The two defining conditions ensure:

1. **Energy Conservation**: The condition $|V\rangle \in \mathcal{V}_0$ ensures that the deformation lies within the degenerate ground-state eigenspace of the Hamiltonian. These are fluctuations of *zero excitation energy*—Goldstone-like modes intrinsic to the vacuum's structure [93, 94, 92]. They represent changes in the vacuum configuration that do not alter the total relational energy of the system.

2. **Physical Distinctness**: The orthogonality condition,

$$\mathrm{Re}\langle\Omega|V\rangle = 0,$$

ensures that the variation $|V\rangle$ represents a physically meaningful deformation of the vacuum state $|\Omega\rangle$, distinct from a mere global rescaling or phase transformation [379, 407].

Geometrically, consider a smooth curve of vacuum states $|\Omega(t)\rangle \in \mathcal{M}_{\text{vac}}$ satisfying:

$$|\Omega(0)\rangle = |\Omega\rangle, \qquad \frac{d}{dt}|\Omega(t)\rangle\Big|_{t=0} = |V\rangle.$$

Since $\mathcal{M}_{\text{vac}}$ consists of normalized states, we impose:

$$\langle\Omega(t)|\Omega(t)\rangle = 1 \quad \forall t.$$

Differentiating with respect to t at $t = 0$ yields:

$$\frac{d}{dt}\langle\Omega(t)|\Omega(t)\rangle\Big|_{t=0} = \langle V|\Omega\rangle + \langle\Omega|V\rangle = 2\,\mathrm{Re}\langle\Omega|V\rangle = 0.$$

Hence, the real orthogonality condition $\mathrm{Re}\langle\Omega|V\rangle = 0$ characterizes precisely those directions in $\mathcal{V}_0$ that preserve the norm constraint. These are the admissible tangent vectors to the vacuum manifold $\mathcal{M}_{\text{vac}}$ [84, 249].

Each fiber $T_{|\Omega\rangle}\mathcal{M}_{\text{vac}}$ is the hyperplane in $\mathcal{V}_0$ orthogonal to the ray defined by $|\Omega\rangle$. It represents the space of all relationally distinct, first-order deformations of the vacuum. This tangent bundle, $T\mathcal{M}_{\text{vac}}$, serves as the natural codomain for operator-valued fields that encode dynamical, vacuum-preserving geometry—most notably, the **Dynamical Gaiamiti Field**, $\hat{\mathbf{\Phi}}(x)$, which will be introduced in the subsequent section [372, 375].

The Nature of the Vacuum State $|\Omega\rangle$

In the context of the tangent bundle definition (Definition 6.90), the state vector $|\Omega\rangle$ denotes a *single, specific vacuum state* chosen from the *Vacuum Manifold*, $\mathcal{M}_{\text{vac}}$.

To be precise:

1. *Origin*: The state $|\Omega\rangle$ is an element of the Vacuum Manifold, which is the set of all normalized states that minimize the Relational Action Functional and are, equivalently, ground states of the Relational Hamiltonian $\hat{H}$.

$$|\Omega\rangle \in \mathcal{M}_{\text{vac}} = \{|\Psi_{\text{vac}}\rangle \in \mathcal{V}_0 \mid \langle\Psi_{\text{vac}}|\Psi_{\text{vac}}\rangle = 1\}.$$

2. *Role as a Reference Point*: When we define the tangent space $T_{|\Omega\rangle}\mathcal{M}_{\text{vac}}$, we are selecting a particular vacuum configuration, $|\Omega\rangle$, to serve as a reference point. The tangent space at that point then represents the space of all possible infinitesimal, energy-preserving deformations *away from that specific vacuum.*

3. *Distinction from the Quantum Holo-State*: The symbol $|\Omega\rangle$ is used as a general placeholder for *any* state within the vacuum manifold. The symbol $|\mathring{\mathbf{\Psi}}\rangle$ refers to the one specific state within $\mathcal{M}_{\text{vac}}$ that is identified as the *Quantum Holo-State*—the actual, realized vacuum of our cosmos.

Remark 6.91 (The Vacuum Manifold, the Universal State, and Local Excitations)**.** The distinction between the Vacuum Manifold ($\mathcal{M}_{\text{vac}}$), the Quantum Holo-State ($|\mathring{\mathbf{\Psi}}\rangle$), and a local vacuum configuration ($|\Omega(x)\rangle$) is central to the physical interpretation of the theory.

The **Vacuum Manifold,** $\mathcal{M}_{\text{vac}}$, represents the complete space of all possible, physically viable ground states of the cosmos. Each point on this manifold is a distinct quantum state that minimizes the total relational action, thereby possessing the same minimal energy, E_0. The existence of a manifold with more than one point signifies a degeneracy in the ground state, providing the necessary mathematical structure for spontaneous symmetry breaking. It is the landscape of potential realities.

The **Quantum Holo-State,** $|\mathring{\mathbf{\Psi}}\rangle$, is the single state vector chosen from this manifold that represents the *actualized reality* of our specific universe. It is the specific answer to the question of which vacuum was selected from the space of all possibilities.

Finally, the symbol $|\Omega(x)\rangle$, used in the context of local field definitions, represents the *local manifestation* of this universally chosen vacuum. It is the particular vacuum configuration that the universe, as described by $|\mathring{\mathbf{\Psi}}\rangle$, has settled into at the emergent spacetime point x. The tangent space at this point, $T_{|\Omega(x)\rangle}\mathcal{M}_{\text{vac}}$, then correctly describes the space of all possible infinitesimal geometric fluctuations, or quantum excitations, around this established ground state.

6.1.21 Holosymmetric Wordwise Tangent Bundle of the Vacuum

Setting. Let $\mathcal{V}_0 := \ker(\hat{H} - E_0\mathbb{I}_{\text{em}})$ be the ground-state eigenspace of the Relational Hamiltonian (Definition 6.54 and eq. (6.130)), and $\mathcal{M}_{\text{vac}} := \{\, |\Omega\rangle \in \mathcal{V}_0 : \langle\Omega|\Omega\rangle = 1 \,\}$ the vacuum manifold (Definition 6.61 and eq. (6.129)). Denote by Π_0 the orthogonal projector onto $\mathcal{V}_0$.

Definition 6.92 (Horizontal projector and tangent fibre (wordwise))**.** For $|\Omega\rangle \in \mathcal{M}_{\text{vac}}$, define the *horizontal projector*

$$\boxed{\mathsf{T}_\Omega \;:=\; \Pi_0 - |\Omega\rangle\langle\Omega|.} \tag{6.150}$$

The *tangent fibre* at $|\Omega\rangle$ is the horizontal slice of $\mathcal{V}_0$:

$$\boxed{T_{|\Omega\rangle}\mathcal{M}_{\text{vac}} \;:=\; \Big\{\, |V\rangle \in \mathcal{V}_0 :\; \mathrm{Re}\langle\Omega|V\rangle = 0 \,\Big\} \;=\; \mathsf{T}_\Omega\,\mathcal{V}_0.} \tag{6.151}$$

Equivalently, on the projective vacuum ray space, $T_{[\Omega]}\mathbb{P}\mathcal{M}_{\text{vac}} \cong \{\, V \in \mathcal{V}_0 : \langle\Omega|V\rangle = 0 \,\}$.

Proposition 6.93 (Operator characterisations)**.** *Let $\mathcal{V}_0 = \ker(\hat{H} - E_0\mathbb{I}_{\text{em}})$ be the ground-space (eq. (6.128)), $\mathcal{M}_{\text{vac}} \subset \mathcal{V}_0$ the unit sphere (eq. (6.129)), and let $|\Omega\rangle \in \mathcal{M}_{\text{vac}}$ be fixed. Write Π_0 for the orthogonal projector onto $\mathcal{V}_0$ and $P_\Omega := |\Omega\rangle\langle\Omega|$.*

The tangent directions $T_{|\Omega\rangle}\mathcal{M}_{\text{vac}}$ defined in Definition 6.90 and eq. (6.148) admit the following operator descriptions.

(a) Hermitian-generator form
For any Hermitian K supported on $\mathcal{V}_0$ ($K^\dagger = K$, $K = \Pi_0 K \Pi_0$) with $\langle\Omega|K|\Omega\rangle = 0$,

$$|V\rangle \;:=\; -\,i\,K\,|\Omega\rangle \;\in\; T_{|\Omega\rangle}\mathcal{M}_{\rm vac}. \tag{6.152}$$

(b) Projector (Grassmannian) form
For any anti-Hermitian A supported on $\mathcal{V}_0$ ($A^\dagger = -A$, $A = \Pi_0 A \Pi_0$),

$$\delta P_\Omega := [A, P_\Omega], \qquad |V\rangle \;:=\; \delta P_\Omega\,|\Omega\rangle \;\in\; T_{|\Omega\rangle}\mathcal{M}_{\rm vac}. \tag{6.153}$$

Moreover, the images produced by (a) *and* (b) *coincide and equal the* horizontal *subspace* $\{\,|V\rangle \in \mathcal{V}_0 :\; \langle\Omega|V\rangle = 0\,\} \subset T_{|\Omega\rangle}\mathcal{M}_{\rm vac}$.

Proof idea. For (a), $\langle\Omega|V\rangle = -i\langle\Omega|K|\Omega\rangle \in i\mathbb{R}$, hence $\mathrm{Re}\langle\Omega|V\rangle = 0$. For (b), $P_\Omega^2 = P_\Omega$ implies admissible variations are commutators with anti-Hermitian A; applying δP_Ω to $|\Omega\rangle$ produces a vector orthogonal (in the real sense) to $|\Omega\rangle$.

Proof. **Preliminaries.** Since $|\Omega\rangle \in \mathcal{V}_0$, one has $\Pi_0|\Omega\rangle = |\Omega\rangle$. By Definition 6.90 and eq. (6.148),

$$T_{|\Omega\rangle}\mathcal{M}_{\rm vac} = \Big\{\,|V\rangle \in \mathcal{V}_0 :\; \mathrm{Re}\langle\Omega|V\rangle = 0\,\Big\}.$$

(a) Sufficiency. Let $K = K^\dagger$ with $K = \Pi_0 K \Pi_0$ and $\langle\Omega|K|\Omega\rangle = 0$, and set $|V\rangle := -iK|\Omega\rangle$. Then $|V\rangle \in \mathcal{V}_0$ because $K = \Pi_0 K \Pi_0$ and $\Pi_0|\Omega\rangle = |\Omega\rangle$ imply $|V\rangle = -i\Pi_0 K \Pi_0|\Omega\rangle \in \mathrm{Ran}\,\Pi_0 = \mathcal{V}_0$. Furthermore,

$$\langle\Omega|V\rangle = -i\,\langle\Omega|K|\Omega\rangle \;\in i\mathbb{R} \quad\Rightarrow\quad \mathrm{Re}\langle\Omega|V\rangle = 0.$$

Hence $|V\rangle \in T_{|\Omega\rangle}\mathcal{M}_{\rm vac}$.
(b) Sufficiency. Let $A^\dagger = -A$ with $A = \Pi_0 A \Pi_0$, and define $\delta P_\Omega = [A, P_\Omega]$. Then δP_Ω is tangent to the rank-one Grassmannian at P_Ω since $P_\Omega^2 = P_\Omega$ implies $P_\Omega + \epsilon\,[A, P_\Omega]$ preserves the projector constraint to first order in ϵ. Acting on $|\Omega\rangle$ gives

$$|V\rangle := \delta P_\Omega|\Omega\rangle = \Big(AP_\Omega - P_\Omega A\Big)|\Omega\rangle = A|\Omega\rangle - |\Omega\rangle\,\langle\Omega|A|\Omega\rangle.$$

Because $A = \Pi_0 A \Pi_0$ and $|\Omega\rangle \in \mathcal{V}_0$, one has $A|\Omega\rangle \in \mathcal{V}_0$, hence $|V\rangle \in \mathcal{V}_0$. Also,

$$\langle\Omega|V\rangle = \langle\Omega|A|\Omega\rangle - \langle\Omega|A|\Omega\rangle = 0,$$

since $\langle\Omega|A|\Omega\rangle \in i\mathbb{R}$ for anti-Hermitian A. Thus $|V\rangle \in T_{|\Omega\rangle}\mathcal{M}_{\rm vac}$.
(b) ⇒ (a): horizontal completeness. Fix A as in *(b)* and set

$$K \;:=\; iA \;-\; i\,\langle\Omega|A|\Omega\rangle\,P_\Omega.$$

Then $K^\dagger = K$ (since iA is Hermitian and $i\langle\Omega|A|\Omega\rangle \in \mathbb{R}$) and $K = \Pi_0 K \Pi_0$. Moreover,

$$\langle\Omega|K|\Omega\rangle = i\,\langle\Omega|A|\Omega\rangle - i\,\langle\Omega|A|\Omega\rangle\,\langle\Omega|P_\Omega|\Omega\rangle = 0.$$

Finally,

$$-iK|\Omega\rangle = -i\Big(iA|\Omega\rangle - i\,\langle\Omega|A|\Omega\rangle\,|\Omega\rangle\Big) = A|\Omega\rangle - |\Omega\rangle\,\langle\Omega|A|\Omega\rangle = \delta P_\Omega|\Omega\rangle,$$

which is exactly the vector produced in *(b)*. Hence every vector obtained by *(b)* arises from *(a)*.

(a) ⇒ (b): horizontal completeness. Fix K as in *(a)* and define $A := -iK$. Then $A^\dagger = -A$ and $A = \Pi_0 A \Pi_0$. Since $\langle\Omega|K|\Omega\rangle = 0$, one has

$$[A, P_\Omega]\,|\Omega\rangle = -iK|\Omega\rangle + i\,|\Omega\rangle\langle\Omega|K|\Omega\rangle = -\,i\,K|\Omega\rangle = |V\rangle,$$

i.e. the vector from *(a)* is obtained by *(b)* with this choice of A.

Images coincide with the horizontal subspace. By construction, both *(a)* and *(b)* always yield vectors with $\langle\Omega|V\rangle = 0$. Conversely, let $|V\rangle \in \mathcal{V}_0$ satisfy $\langle\Omega|V\rangle = 0$. Set

$$A_{V,\Omega} \;:=\; |V\rangle\langle\Omega| \;-\; |\Omega\rangle\langle V| \qquad \text{(anti-Hermitian, supported on } \mathcal{V}_0),$$

and compute

$$[A_{V,\Omega}, P_\Omega]\,|\Omega\rangle = A_{V,\Omega}|\Omega\rangle - |\Omega\rangle\langle\Omega|A_{V,\Omega}|\Omega\rangle = |V\rangle - |\Omega\rangle\underbrace{\Big(\langle\Omega|V\rangle - \langle V|\Omega\rangle\Big)}_{=0} = |V\rangle.$$

Thus any *horizontal* tangent vector ($\langle\Omega|V\rangle = 0$) arises from *(b)*, and hence from *(a)* by the previous steps.

Remark on the vertical $U(1)$ direction. The purely vertical phase direction $i|\Omega\rangle$ satisfies $\mathrm{Re}\langle\Omega|\,i\Omega\rangle = 0$ and therefore lies in $T_{|\Omega\rangle}\mathcal{M}_{\mathrm{vac}}$ (eq. (6.148)). It is *not* produced by *(a)* under the constraint $\langle\Omega|K|\Omega\rangle = 0$, nor by *(b)* (since $[A, P_\Omega]$ leaves P_Ω invariant along global phase changes). Allowing K with $\langle\Omega|K|\Omega\rangle \neq 0$ generates the vertical component: $-iK|\Omega\rangle$ acquires a multiple of $i|\Omega\rangle$. Hence *(a)–(b)* exactly parameterise the *horizontal* (projective) tangent space $\{V : \langle\Omega|V\rangle = 0\}$, while the full real-orthogonal tangent $T_{|\Omega\rangle}\mathcal{M}_{\mathrm{vac}}$ is the direct sum of this horizontal subspace with $\mathrm{span}\{\,i|\Omega\rangle\,\}$. □

Definition 6.94 (FS inner product and Berry data on $T\mathcal{M}_{\mathrm{vac}}$)**.** For $|V\rangle, |W\rangle \in T_{|\Omega\rangle}\mathcal{M}_{\mathrm{vac}}$, define the Fubini–Study inner product in horizontal form

$$\boxed{\langle V, W\rangle_{\mathrm{FS}} := \;\mathrm{Re}\Big(\langle V|W\rangle - \langle V|\Omega\rangle\langle\Omega|W\rangle\Big) = \;\mathrm{Re}\,\langle V|\mathsf{T}_\Omega W\rangle.} \tag{6.154}$$

The U(1) Berry connection and curvature on $\mathcal{M}_{\mathrm{vac}}$ are

$$\mathcal{A} := i\,\langle\Omega|d\Omega\rangle, \qquad \mathcal{F} := d\mathcal{A} = i\,d\langle\Omega| \wedge d|\Omega\rangle, \tag{6.155}$$

with horizontality enforced by $d|\Omega\rangle \mapsto \mathsf{T}_\Omega\, d|\Omega\rangle$.

Proposition 6.95 (Gauge/holonomy covariance)**.** *Under an endpoint gauge/frame U on $\mathcal{H}_{\mathrm{em}}$, $|\Omega\rangle \mapsto U|\Omega\rangle$, $\Pi_0 \mapsto U\Pi_0U^\dagger$, one has*

$$\mathsf{T}_\Omega \;\mapsto\; U\,\mathsf{T}_\Omega\,U^\dagger, \qquad T_{|\Omega\rangle}\mathcal{M}_{\mathrm{vac}} \;\mapsto\; U\Big(T_{|\Omega\rangle}\mathcal{M}_{\mathrm{vac}}\Big), \tag{6.156}$$

and the Fubini–Study inner product on the tangent fibres is invariant. Moreover, parallel transport along admissible paths is implemented by unitary transporters and carries fibres isometrically (cf. the gauge–covariant kinematics of Section 4.5).

Proof. **Step 1: Covariance of the horizontal projector.** Recall $\mathsf{T}_\Omega := \Pi_0 - |\Omega\rangle\langle\Omega|$, with Π_0 the orthogonal projector onto $\mathcal{V}_0 = \ker(\hat{H} - E_0\mathbb{I}_{\text{em}})$ and $|\Omega\rangle \in \mathcal{M}_{\text{vac}}$ (eqs. (6.128) and (6.129)). Under a unitary frame/gauge U,

$$\mathsf{T}_\Omega \;\longmapsto\; U\Pi_0 U^\dagger - U|\Omega\rangle\langle\Omega|U^\dagger = \Pi_0' - |\Omega'\rangle\langle\Omega'| =: \mathsf{T}_{\Omega'},$$

with $\Pi_0' := U\Pi_0 U^\dagger$ and $|\Omega'\rangle := U|\Omega\rangle$. Hence $U\,\mathsf{T}_\Omega\,U^\dagger = \mathsf{T}_{\Omega'}$.

Step 2: Covariance of the tangent fibre. By Definition 6.90 and eq. (6.148),

$$T_{|\Omega\rangle}\mathcal{M}_{\text{vac}} = \Big\{|V\rangle \in \mathcal{V}_0 :\ \mathrm{Re}\langle\Omega|V\rangle = 0\Big\} = \mathsf{T}_\Omega\,\mathcal{V}_0.$$

Unitarity gives $\mathcal{V}_0' = U\mathcal{V}_0$ and preserves the inner product, so for any $|V\rangle \in T_{|\Omega\rangle}\mathcal{M}_{\text{vac}}$,

$$\mathrm{Re}\langle\Omega'|\,U|V\rangle\rangle = \mathrm{Re}\langle\Omega|V\rangle = 0, \qquad U|V\rangle \in \mathcal{V}_0'.$$

Therefore $U\big(T_{|\Omega\rangle}\mathcal{M}_{\text{vac}}\big) = T_{|\Omega'\rangle}\mathcal{M}_{\text{vac}}$, which is equivalent to $U\,\mathsf{T}_\Omega\,\mathcal{V}_0 = \mathsf{T}_{\Omega'}\,\mathcal{V}_0'$.

Step 3: Invariance of the Fubini–Study inner product. For tangent vectors $|V\rangle, |W\rangle \in T_{|\Omega\rangle}\mathcal{M}_{\text{vac}}$, the Fubini–Study form can be written in the horizontal projector form

$$\langle V, W\rangle_{\text{FS}} = \mathrm{Re}\,\Big\langle V\Big|\mathsf{T}_\Omega W\Big\rangle = \mathrm{Re}\,\Big(\langle V|W\rangle - \langle V|\Omega\rangle\langle\Omega|W\rangle\Big).$$

Using $U^\dagger U = \mathbb{I}$ and Step 1,

$$\mathrm{Re}\,\Big\langle UV\Big|\,\mathsf{T}_{\Omega'}(UW)\Big\rangle = \mathrm{Re}\,\Big\langle V\Big|\,U^\dagger\mathsf{T}_{\Omega'}U\,W\Big\rangle = \mathrm{Re}\,\Big\langle V\Big|\,\mathsf{T}_\Omega W\Big\rangle,$$

so $\langle UV, UW\rangle_{\text{FS}} = \langle V, W\rangle_{\text{FS}}$. Thus the fibre metric is unitarily invariant.

Step 4: Holonomy/parallel transport. Along an admissible path γ in the base (or control) space, let $\mathcal{U}(x \leftarrow x';\gamma)$ denote the gauge–covariant transporter (path–ordered unitary) with the composition and gauge laws of Section 4.5. For a vacuum section $|\Omega(x)\rangle$, define $|\Omega(x)\rangle = \mathcal{U}(x \leftarrow x_0;\gamma)\,|\Omega(x_0)\rangle$ and similarly transport $|V(x_0)\rangle \in T_{|\Omega(x_0)\rangle}\mathcal{M}_{\text{vac}}$ to $|V(x)\rangle := \mathcal{U}(x \leftarrow x_0;\gamma)\,|V(x_0)\rangle$. Since $\mathcal{U}$ is unitary and intertwines the corresponding projectors, Steps 1–3 apply with $U = \mathcal{U}(x \leftarrow x_0;\gamma)$, showing that $|V(x)\rangle \in T_{|\Omega(x)\rangle}\mathcal{M}_{\text{vac}}$ and that $\langle\cdot,\cdot\rangle_{\text{FS}}$ is preserved. Hence parallel transport carries fibres isometrically.

Combining Steps 1–4 establishes (6.156) and the asserted isometry of fibres under gauge/holonomy transport. □

Proposition 6.96 (Energy-stationarity on the ground leaf)**.** *For any $|V\rangle \in T_{|\Omega\rangle}\mathcal{M}_{\text{vac}}$,*

$$\frac{d}{d\tau}\,\langle\Omega + \tau V|\hat{H}|\Omega + \tau V\rangle\Big|_{\tau=0} = 0. \tag{6.157}$$

Equivalently, if $|V\rangle = -iK|\Omega\rangle$ as in (6.152), then $\langle\Omega|[\hat{H}, K]|\Omega\rangle = 0$.

Proof idea. On $\mathcal{V}_0$, $\hat{H}|\cdot\rangle = E_0|\cdot\rangle$; the first variation is $E_0(\langle V|\Omega\rangle + \langle\Omega|V\rangle) = 2E_0\,\mathrm{Re}\langle\Omega|V\rangle = 0$. For the commutator form, use $\hat{H}|\Omega\rangle = E_0|\Omega\rangle$.

Proof. **Preliminaries.** By definition of the tangent fibre $T_{|\Omega\rangle}\mathcal{M}_{\mathrm{vac}}$ (see Definition 6.90 and eq. (6.148)), one has $|V\rangle \in \mathcal{V}_0$ and $\mathrm{Re}\langle\Omega|V\rangle = 0$, where $\mathcal{V}_0 = \ker(\hat{H} - E_0\mathbb{I}_{\mathrm{em}})$ is the ground eigenspace (eq. (6.128)). Hence

$$\hat{H}|\Omega\rangle = E_0|\Omega\rangle, \qquad \hat{H}|V\rangle = E_0|V\rangle. \tag{6.158}$$

Step 1: First variation along an admissible direction. Consider the (not necessarily normalised) curve $|\Omega(\tau)\rangle := |\Omega\rangle + \tau|V\rangle$. Its energy expectation is

$$\mathcal{E}(\tau) := \langle\Omega(\tau)|\hat{H}|\Omega(\tau)\rangle = \langle\Omega|\hat{H}|\Omega\rangle + \tau\Big(\langle V|\hat{H}|\Omega\rangle + \langle\Omega|\hat{H}|V\rangle\Big) + O(\tau^2).$$

Therefore

$$\mathcal{E}'(0) = \langle V|\hat{H}|\Omega\rangle + \langle\Omega|\hat{H}|V\rangle. \tag{6.159}$$

Using (6.158),

$$\mathcal{E}'(0) = E_0\Big(\langle V|\Omega\rangle + \langle\Omega|V\rangle\Big) = 2E_0\ \mathrm{Re}\langle\Omega|V\rangle\ =\ 0,$$

which establishes (6.157).

Step 2: Independence of normalisation. The curve $|\widetilde{\Omega}(\tau)\rangle := \frac{|\Omega\rangle+\tau|V\rangle}{\|\,|\Omega\rangle+\tau|V\rangle\,\|}$ stays on the unit sphere in $\mathcal{V}_0$. Since $\frac{d}{d\tau}\|\,|\Omega\rangle+\tau|V\rangle\,\|^2\big|_{\tau=0} = 2\,\mathrm{Re}\langle\Omega|V\rangle = 0$ by tangency, the difference between $\mathcal{E}(\tau)$ and $\langle\widetilde{\Omega}(\tau)|\hat{H}|\widetilde{\Omega}(\tau)\rangle$ appears only at $O(\tau^2)$. Hence the first derivative at $\tau = 0$ is the same for either curve, and the stationarity result is intrinsic to the tangent direction.

Step 3: Commutator (generator) form. By the Hermitian–generator characterisation of tangent vectors (Proposition 6.93, (6.152)), any $|V\rangle \in T_{|\Omega\rangle}\mathcal{M}_{\mathrm{vac}}$ can be written as $|V\rangle = -iK|\Omega\rangle$ with $K^\dagger = K$, $K = \Pi_0 K \Pi_0$, and $\langle\Omega|K|\Omega\rangle = 0$. Substituting into (6.159) gives

$$\mathcal{E}'(0) = \langle -iK\Omega|\hat{H}|\Omega\rangle + \langle\Omega|\hat{H}|-iK|\Omega\rangle = i\langle\Omega|K\hat{H}|\Omega\rangle - i\langle\Omega|\hat{H}K|\Omega\rangle = i\,\langle\Omega|[K,\hat{H}]|\Omega\rangle.$$

Thus $\mathcal{E}'(0) = 0$ is equivalent to $\langle\Omega|[K,\hat{H}]|\Omega\rangle = 0$, or, equivalently, $\langle\Omega|[\hat{H},K]|\Omega\rangle = 0$.

Combining Steps 1–3 yields the stated energy-stationarity on the ground leaf and its equivalent commutator form. □

6.1.22 Holosymmetric Lift of the Tangent Bundle of the Vacuum

Definition 6.97 (HS lift: branch ground projectors and horizontal projectors)**.** Consider the two–branch HS extension (Section 6.1.15; cf. Section 6.1.18 and Definition 6.69). Let $\Pi_0^{(k)} := \mathbf{1}_{\{E_0^{(k)}\}}\big(\widehat{\boldsymbol{H}}^{(k)}\big)$ project onto the branch ground spaces $\mathcal{V}_0^{(k)} := \mathrm{Ran}\,\Pi_0^{(k)}$ for $k \in \{0,1\}$. Fix a normalized vacuum vector $|\Omega\rangle$ and define the branchwise *horizontal* projectors

$$\mathsf{T}_\Omega^{(k)} \ :=\ \Pi_0^{(k)} - |\Omega\rangle\langle\Omega| \qquad (k = 0,1), \tag{6.160}$$

whenever $|\Omega\rangle \in \mathcal{V}_0^{(k)}$.

Definition 6.98 (Block HS tangent fibre)**.** With Definition 6.97, the *block HS tangent fibre* at the vacuum $|\Omega\rangle$ is

$$\boxed{T^{\mathrm{HS}}_{|\Omega\rangle} \;:=\; \mathsf{T}^{(0)}_{\Omega}\,\mathcal{V}^{(0)}_{0} \;\oplus\; \mathsf{T}^{(1)}_{\Omega}\,\mathcal{V}^{(1)}_{0}.} \tag{6.161}$$

Definition 6.99 (Reduction at dial phase and reduced horizontal projector)**.** Prepare the law register in $|+_\varphi\rangle_L = \frac{1}{\sqrt{2}}(|0\rangle_L + e^{i\varphi}|1\rangle_L)$ and let $\hat{H}^{\langle \mathrm{HS}\rangle}(\varphi)$ be the reduced operator (Proposition 6.78 and eq. (6.142); cf. eq. (6.125)). Define the reduced ground projector and horizontal projector by

$$\Pi_0^{\langle \mathrm{HS}\rangle}(\varphi) := \mathbf{1}_{\{E_0^{\langle \mathrm{HS}\rangle}(\varphi)\}}\Big(\hat{H}^{\langle \mathrm{HS}\rangle}(\varphi)\Big), \qquad \mathsf{T}_{\Omega}^{\langle \mathrm{HS}\rangle}(\varphi) := \Pi_0^{\langle \mathrm{HS}\rangle}(\varphi) - |\Omega\rangle\langle\Omega|. \tag{6.162}$$

The *reduced HS tangent fibre* is

$$\boxed{T^{\langle \mathrm{HS}\rangle}_{|\Omega\rangle}(\varphi) \;=\; \mathsf{T}_{\Omega}^{\langle \mathrm{HS}\rangle}(\varphi)\,\mathrm{Ran}\Big(\Pi_0^{\langle \mathrm{HS}\rangle}(\varphi)\Big).} \tag{6.163}$$

Proposition 6.100 (Compression consistency of tangent fibres)**.** *Let $\mathcal{C}_\varphi : \mathcal{H}_L \otimes \mathcal{H}_{\mathrm{em}} \to \mathcal{H}_{\mathrm{em}}$ be the compression $\mathcal{C}_\varphi := \langle +_\varphi|(\,\cdot\,)|+_\varphi\rangle$. Then the compression intertwines block and reduced ground projectors and their horizontal parts:*

$$\mathcal{C}_\varphi\Big[\Big(P_0\otimes \Pi_0^{(0)}\Big)\oplus\Big(P_1\otimes \Pi_0^{(1)}\Big)\Big] = \Pi_0^{\langle \mathrm{HS}\rangle}(\varphi), \qquad \mathcal{C}_\varphi\Big[\Big(P_0\otimes \mathsf{T}_{\Omega}^{(0)}\Big)\oplus\Big(P_1\otimes \mathsf{T}_{\Omega}^{(1)}\Big)\Big] = \mathsf{T}_{\Omega}^{\langle \mathrm{HS}\rangle}(\varphi), \tag{6.164}$$

and therefore $\mathcal{C}_\varphi\Big[T^{\mathrm{HS}}_{|\Omega\rangle}\Big] = T^{\langle \mathrm{HS}\rangle}_{|\Omega\rangle}(\varphi)$.

Proof. By Proposition 6.78, $\hat{H}^{\langle \mathrm{HS}\rangle}(\varphi)$ is the compression of $\widehat{\boldsymbol{H}}^{\mathrm{HS}}$; spectral calculus commutes with unitary conjugations and with such finite–rank compressions on the law register, yielding the first identity in (6.164). Subtract $|\Omega\rangle\langle\Omega|$ on both sides to obtain the second identity, then apply both to the respective ranges to get (6.163). □

Proposition 6.101 (Gauge/holonomy covariance and causal stability)**.** *Under endpoint gauge/holonomy transformations (Propositions 6.66 and 6.77), the spaces $\mathcal{V}_0^{(k)}$, $T^{\mathrm{HS}}_{|\Omega\rangle}$, $\mathrm{Ran}\,\Pi_0^{\langle \mathrm{HS}\rangle}(\varphi)$ and $T^{\langle \mathrm{HS}\rangle}_{|\Omega\rangle}(\varphi)$ transform by the induced unitary and hence have gauge/holonomy–invariant* rays*. Moreover, since the HS dial does not modify the expectation unified tensor $\bar{\Sigma}$ (and thus $\bar{g}^{\mathrm{Lor}}$), the cone assignment is unchanged (Corollary 6.87; cf. Section 5.4.2).*

Proof. Covariance of the projectors follows from Propositions 6.66 and 6.77 and unitary equivariance of spectral calculus; invariance of rays is immediate. The causal statement follows from Corollary 6.87, since the dial acts only on the mixed–sector integrand (Section 6.1.12) and leaves $\bar{\Sigma}$ (hence $\bar{g}^{\mathrm{Lor}}$) unchanged. □

Remark 6.102 (On choice of reference vacuum and dimensions)**.** (i) The definition (6.160) assumes $|\Omega\rangle \in \mathcal{V}_0^{(k)}$. If not, replace $|\Omega\rangle$ by the normalized projection $|\Omega^{(k)}\rangle := \Pi_0^{(k)}|\Omega\rangle / \|\Pi_0^{(k)}|\Omega\rangle\|$ and set $\mathsf{T}_{\Omega}^{(k)} := \Pi_0^{(k)} - |\Omega^{(k)}\rangle\langle\Omega^{(k)}|$; this yields an equivalent horizontal fibre and leaves (6.161) unchanged up to canonical isomorphism.

(ii) If $\dim \mathcal{V}_0^{(k)} = m_k$, then $\dim\left(\mathsf{T}_\Omega^{(k)}\mathcal{V}_0^{(k)}\right) = m_k - 1$ (when the assumption in (i) holds). Consequently, $\dim T^{\mathrm{HS}}_{|\Omega\rangle} = (m_0 - 1) + (m_1 - 1)$ and $\dim T^{\langle \mathrm{HS}\rangle}_{|\Omega\rangle}(\varphi) = m_{\langle \mathrm{HS}\rangle}(\varphi) - 1$, where $m_{\langle \mathrm{HS}\rangle}(\varphi) := \dim \operatorname{Ran} \Pi_0^{\langle \mathrm{HS}\rangle}(\varphi)$.

(iii) All statements hold triad-wise (corr/vac/spec), since the constructions respect the sector decomposition (eq. (5.86)) and the HS–covariant integration (Corollary 6.37).

Remark 6.103 (Bundle-level summary and use)**.** The operator-theoretic tangent bundle is the disjoint union

$$\boxed{T\mathcal{M}_{\mathrm{vac}} \;=\; \bigsqcup_{|\Omega\rangle\in\mathcal{M}_{\mathrm{vac}}} \mathsf{T}_\Omega\,\mathcal{V}_0,} \tag{6.165}$$

i.e. fibres are the real-horizontal complements to the vacuum rays. Any vacuum-preserving variation is obtained by acting with a Hermitian generator K supported on $\mathcal{V}_0$ and projecting with T_Ω. This is the wordwise analogue of taking a curve on the unit sphere and discarding the vertical (phase) component, and it is the natural target space for vacuum-preserving dynamical fields (cf. the operator correlator density and Hamiltonian word in Definitions 6.55 and 6.56 and the standard tangent bundle definition in Section 6.1.20).

6.1.23 Quantum Circuits for the Vacuum Manifold and its Tangent Bundle

Objective. Realise, on quantum hardware, the projector onto the ground leaf $\Pi_0 = \mathbf{1}_{\{E_0\}}(\widehat{\boldsymbol{H}}[|\Psi\rangle])$ and the horizontal projector $\mathsf{T}_\Omega = \Pi_0 - |\Omega\rangle\langle\Omega|$ (Definition 6.92), together with estimators for the Fubini–Study (FS) metric and Berry data (Definition 6.94), all consistent with the block-encodings previously used for the Relational Hamiltonian (Section 6.1.16).

Primitives (assumed available).

1. **Block-encoding of the (reduced) Hamiltonian word.** A unitary U_H that block-encodes $\widetilde{H} := (\widehat{\boldsymbol{H}} - E_{\mathrm{ref}}\mathbb{I})/\alpha$ with known scale α (Section 6.1.16). Here E_{ref} can be taken as a running estimate of E_0 or any shift that isolates the ground leaf.

2. **Ground-state preparation.** Either (i) *phase estimation + postselection* using U_H and a narrow energy window around E_0, or (ii) a *variational* preparer PREP_Ω yielding $|\Omega\rangle \in \mathcal{V}_0$ (Definition 6.61). In either case, the reflection about $|\Omega\rangle$ is available: $R_\Omega := \mathbb{I} - 2|\Omega\rangle\langle\Omega| = \mathrm{PREP}_\Omega\, Z\, \mathrm{PREP}_\Omega^\dagger$.

3. **Polynomial (QSP) filters.** A quantum signal processing (QSP) realisation $F_{\mathrm{gs}}(\widetilde{H})$ of a degree-d polynomial that approximates the spectral projector Π_0 (Heaviside-like filter with gap-dependent width).

Ground-space projector (QSP filter). Apply the QSP sequence driven by U_H to implement $\mathsf{Proj}_{\mathrm{gs}} \approx \Pi_0$. In circuit form:



Applied to any input, this coherently projects into $\mathcal{V}_0$ up to polynomial error.

Horizontal projector $\mathsf{T}_\Omega = \Pi_0 - |\Omega\rangle\langle\Omega|$. Implement Π_0 by the QSP filter above, and $|\Omega\rangle\langle\Omega|$ via the rank-one block-encoding $\frac{\mathbb{I}-R_\Omega}{2}$. Combine them by linear combination of unitaries (LCU) with a single-qubit ancilla a:



Postselecting (or using oblivious amplitude amplification) realises the affine combination $\Pi_0 - (\mathbb{I} - R_\Omega)/2 = \mathsf{T}_\Omega$ up to normalisation. Applied to a candidate variation $|\delta\Psi\rangle$, this returns the *tangent projection* $|V\rangle \propto \mathsf{T}_\Omega|\delta\Psi\rangle$ (eq. (6.151)).

Generator-based tangent vectors. Given a Hermitian word K supported on $\mathcal{V}_0$ (Proposition 6.93), block-encode K by LCU (standard for local sums), apply $|V\rangle \propto -i\,K\,|\Omega\rangle$, then enforce horizontality by a single projection $|V\rangle \leftarrow \mathsf{T}_\Omega|V\rangle$. A compact circuit realisation is:



where T_Ω is the LCU gadget above.

FS inner product estimator. For $|V\rangle, |W\rangle \in T_{|\Omega\rangle}\mathcal{M}_{\mathrm{vac}}$, estimate $\langle V, W\rangle_{\mathrm{FS}} = \mathrm{Re}(\langle V|W\rangle - \langle V|\Omega\rangle\langle\Omega|W\rangle)$ (Definition 6.94) via two Hadamard/SWAP-style overlap tests; the $\langle\cdot|\Omega\rangle$ terms are obtained by replacing one register with $|\Omega\rangle$.

Berry phase along a vacuum loop. Given a discretised loop $|\Omega_0\rangle \to \cdots \to |\Omega_m\rangle = |\Omega_0\rangle$, estimate the Bargmann invariant $\arg\prod_{j=0}^{m-1}\langle\Omega_j|\Omega_{j+1}\rangle$ by chaining overlap tests with phase kickback on a probe qubit; in the continuum limit this yields $\oint i\langle\Omega|d\Omega\rangle$ (Definition 6.94).

HS extension and reduction. With a law register L (Section 6.1.15), block-encode the branch projectors $\Pi_0^{(0)}, \Pi_0^{(1)}$ via their respective QSP filters, and assemble the HS horizontal projector $\mathsf{T}_\Omega^{\mathrm{HS}} = |0\rangle\langle 0|_L \otimes (\Pi_0^{(0)} - |\Omega\rangle\langle\Omega|) + |1\rangle\langle 1|_L \otimes (\Pi_0^{(1)} - |\Omega\rangle\langle\Omega|)$. Prepare $|+_\varphi\rangle_L$ and take the partial

matrix element to obtain the reduced projector $\mathsf{T}_\Omega^{\langle \mathrm{HS}\rangle}(\varphi)$ (Section 6.1.22 and eq. (6.163)). A compact control layout is:



which, after taking $\langle +_\varphi|_L(\cdot)|+_\varphi\rangle_L$, realises $\mathsf{T}_\Omega^{\langle \mathrm{HS}\rangle}(\varphi)$.

Readouts and guarantees.

- *Tangent projection:* For any input $|\delta\Psi\rangle$, $|V\rangle \propto \mathsf{T}_\Omega|\delta\Psi\rangle \in T_{|\Omega\rangle}\mathcal{M}_{\mathrm{vac}}$ (eq. (6.151)).
- *FS metric:* Overlaps from Hadamard/SWAP tests reconstruct $\langle\cdot,\cdot\rangle_{\mathrm{FS}}$ (Definition 6.94).
- *Berry curvature:* Discrete loop phases converge to $\mathcal{F} = d(i\langle\Omega|d\Omega\rangle)$ (eq. (6.155)).
- *Gauge covariance:* All blocks use the same dressing/transport oracles as Section 6.1.16; invariance and centrality properties follow from Propositions 6.36 and 6.60.

Complexity notes. The cost is dominated by the QSP degree required to resolve the ground-gap in F_{gs}, the LCU cost of K (when used), and the precision of overlap/phase estimation. Field/dialectic indices are finite (2×4), so the FD-trace gadgets (Definition 6.45) are $O(1)$ in register count and integrate unchanged with the Hamiltonian block-encodings from Section 6.1.16.

6.1.24 Inner Product on the Vacuum Tangent Space

Before defining the action, we must formalize the geometric structure of the space in which the Gaiamiti field's kinetic energy is measured. This requires defining an inner product on the tangent space of the vacuum manifold [84, 407].

Definition 6.104 (Inner Product on the Vacuum Tangent Space)**.** Let $T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}$ be the tangent space to the vacuum manifold at the local vacuum state $|\Omega(x)\rangle$ (Definition 6.90). We define the **inner product on the tangent space**, denoted g_T, as the canonical Hermitian inner product inherited from the ambient Hilbert space $\mathcal{H}_{\mathrm{em}}$ [379, 407]. For any two tangent vectors $|V\rangle, |W\rangle \in T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}$, their inner product is given by:

$$\boxed{g_T(|V\rangle, |W\rangle) := \langle V|W\rangle_{\mathcal{H}_{\mathrm{em}}}.} \tag{6.166}$$

The local orthonormal basis $\{|e_i(x)\rangle\}_{i=1}^r$ of the tangent space is defined to be orthonormal with respect to this inner product, such that $g_T(|e_i(x)\rangle, |e_j(x)\rangle) = \delta_{ij}$ [287].

Remark 6.105 (The Geometric Role of the Inner Product)**.** This inner product, g_T, is the mathematical structure that endows the space of possible geometric fluctuations with a notion of metric [84, 375]. It allows us to rigorously define the "magnitude" of a deformation, given by $\|V\|^2 = g_T(|V\rangle, |V\rangle)$, and the "angle" or relational orthogonality between two different types of deformations. It is precisely this structure that is required to formulate a well-defined and covariant kinetic term for the Gaiamiti field in its action functional [85], thereby providing the foundation for the dynamics of the emergent geometry.

6.1.25 Holosymmetric Wordwise Inner Product on the Vacuum Tangent Bundle

Setting. Let $\mathcal{V}_0 = \ker(\hat{H} - E_0 \mathbb{I}_{\text{em}})$ be the ground eigenspace (eq. (6.128)) and $\mathcal{M}_{\text{vac}} = \{\, |\Omega\rangle \in \mathcal{V}_0 : \langle\Omega|\Omega\rangle = 1 \,\}$ (eq. (6.129)). Denote by Π_0 the orthogonal projector onto $\mathcal{V}_0$, and write the horizontal projector at $|\Omega\rangle$ as

$$\mathsf{T}_\Omega \;:=\; \Pi_0 - |\Omega\rangle\langle\Omega|\,,$$

so that the tangent fibre is $T_{|\Omega\rangle}\mathcal{M}_{\text{vac}} = \{\, |V\rangle \in \mathcal{V}_0 : \mathrm{Re}\langle\Omega|V\rangle = 0 \,\}$ (Definition 6.90 and eq. (6.148)).

Definition 6.106 (HS vacuum–tangent inner product)**.** For $|V\rangle, |W\rangle \in T_{|\Omega\rangle}\mathcal{M}_{\text{vac}}$, define

$$\boxed{\; g_T(|V\rangle, |W\rangle) \;:=\; \langle V|W\rangle \;=\; \langle V|\mathsf{T}_\Omega W\rangle \;=\; \langle V|W\rangle - \langle V|\Omega\rangle\,\langle\Omega|W\rangle\,. \;} \tag{6.167}$$

Its Riemannian (real) part is

$$\boxed{\; g_T^{\mathrm{R}}(|V\rangle, |W\rangle) \;:=\; \mathrm{Re}\; g_T(|V\rangle, |W\rangle) \;=\; \mathrm{Re}\Big(\langle V|W\rangle - \langle V|\Omega\rangle\,\langle\Omega|W\rangle\Big)\,. \;} \tag{6.168}$$

A local orthonormal frame $\{|e_i(x)\rangle\}_{i=1}^{r} \subset T_{|\Omega(x)\rangle}\mathcal{M}_{\text{vac}}$ is one with $g_T(e_i, e_j) = \delta_{ij}$.

Proposition 6.107 (Basic properties of g_T)**.** *For each fixed $|\Omega\rangle \in \mathcal{M}_{\text{vac}}$:*

(i) g_T is Hermitian: $g_T(V, W) = \overline{g_T(W, V)}$.

(ii) Positivity on the tangent fibre: $g_T^{\mathrm{R}}(V, V) = \| \mathsf{T}_\Omega V \|^2 \geq 0$, and $g_T^{\mathrm{R}}(V, V) = 0$ iff $V = 0$ in $T_{|\Omega\rangle}\mathcal{M}_{\text{vac}}$.

(iii) Gauge/frame invariance: under any endpoint unitary U on $\mathcal{H}_{\text{em}}$, $|\Omega\rangle \mapsto U|\Omega\rangle$, $\Pi_0 \mapsto U\Pi_0 U^\dagger$,

$$\mathsf{T}_\Omega \mapsto U\mathsf{T}_\Omega U^\dagger, \qquad g_T(UV, UW) = g_T(V, W), \tag{6.169}$$

so g_T^{R} is invariant.

Proof. (1) Hermiticity is inherited from $\langle\cdot|\cdot\rangle$ and the rank-one correction $\langle V|\Omega\rangle\,\langle\Omega|W\rangle$. (2) Using (6.150), $g_T^{\mathrm{R}}(V, V) = \mathrm{Re}\langle V|\mathsf{T}_\Omega V\rangle = \langle V|\mathsf{T}_\Omega V\rangle = \| \mathsf{T}_\Omega V \|^2 \geq 0$, with equality iff $\mathsf{T}_\Omega V = 0$, i.e. $V \parallel |\Omega\rangle$. But $V \in T_{|\Omega\rangle}\mathcal{M}_{\text{vac}}$ also satisfies $\mathrm{Re}\langle\Omega|V\rangle = 0$, hence $V = 0$ in the tangent fibre. (3) Follows from U-unitarity: $\langle UV|UW\rangle = \langle V|W\rangle$ and $U|\Omega\rangle\langle\Omega|U^\dagger = |U\Omega\rangle\langle U\Omega|$, which implies $U\mathsf{T}_\Omega U^\dagger = \mathsf{T}_{U\Omega}$ and (6.169). □

Proposition 6.108 (Equivalence with the Fubini–Study metric)**.** *Let* $[\Omega]$ *denote the ray of* $|\Omega\rangle$*, and let* $\mathbb{P}(\mathcal{V}_0)$ *be the projective space. Then* g_T^{R} *equals the Fubini–Study metric pulled back to the unit sphere: for any horizontal representatives* $|V\rangle, |W\rangle$ *(i.e.* $\mathrm{Re}\langle\Omega|V\rangle = \mathrm{Re}\langle\Omega|W\rangle = 0$*),*

$$g_T^{\mathrm{R}}(V,W) = \mathrm{Re}\left(\langle V|W\rangle - \langle V|\Omega\rangle\,\langle\Omega|W\rangle\right) = \mathrm{FS}_{[\Omega]}(V,W)\,. \tag{6.170}$$

Proof idea. The standard Fubini–Study metric on $\mathbb{P}(\mathcal{V}_0)$ at $[\Omega]$ is $\mathrm{FS}_{[\Omega]}(V,W) = \mathrm{Re}\left(\langle V|W\rangle - \langle V|\Omega\rangle\,\langle\Omega|W\rangle\right)$ for horizontal lifts V, W modulo the vertical line $\mathrm{span}\{i|\Omega\rangle\}$. This coincides with (6.168).

Proof. Step 1 (Sphere, projection, and horizontal/vertical split). Let $S(\mathcal{V}_0) := \{\,|\Omega\rangle \in \mathcal{V}_0 : \langle\Omega|\Omega\rangle = 1\,\}$ and let $\pi : S(\mathcal{V}_0) \to \mathbb{P}(\mathcal{V}_0)$ be the canonical projection $|\Omega\rangle \mapsto [\Omega]$. The tangent space to the unit sphere at $|\Omega\rangle$ is

$$T_{|\Omega\rangle}S(\mathcal{V}_0) = \{\,|V\rangle \in \mathcal{V}_0 : \mathrm{Re}\langle\Omega|V\rangle = 0\,\}.$$

Inside it, the *vertical line* is $\mathcal{V}_{|\Omega\rangle} := \mathrm{span}\{\,i|\Omega\rangle\,\}$, and a canonical *horizontal* complement is the complex-orthogonal subspace

$$\mathcal{H}_{|\Omega\rangle} := \{\,|V\rangle \in \mathcal{V}_0 : \langle\Omega|V\rangle = 0\,\}.$$

The orthogonal projector (on $\mathcal{V}_0$) onto $\mathcal{H}_{|\Omega\rangle}$ is $\mathsf{T}_\Omega = \Pi_0 - |\Omega\rangle\langle\Omega|$ (eq. (6.150)), so that $T_{|\Omega\rangle}S(\mathcal{V}_0) = \mathcal{H}_{|\Omega\rangle} \oplus \mathcal{V}_{|\Omega\rangle}$ (orthogonal sum).

Step 2 (Riemannian submersion characterisation of FS*).* The Fubini–Study metric on $\mathbb{P}(\mathcal{V}_0)$ is the unique metric making π a Riemannian submersion: for $|V\rangle, |W\rangle \in T_{|\Omega\rangle}S(\mathcal{V}_0)$,

$$\mathrm{FS}_{[\Omega]}\left(d\pi_{|\Omega\rangle}V,\ d\pi_{|\Omega\rangle}W\right) = \mathrm{Re}\,\langle V^{\mathrm{H}}|W^{\mathrm{H}}\rangle, \tag{6.171}$$

where $V^{\mathrm{H}} := \mathsf{T}_\Omega V \in \mathcal{H}_{|\Omega\rangle}$ and similarly for W^{H}. This is well-defined because $d\pi_{|\Omega\rangle}$ annihilates the vertical line ($d\pi_{|\Omega\rangle}(i|\Omega\rangle) = 0$), hence depends only on horizontal components.

Step 3 (Pullback formula and g_T^{R}*).* Using $\mathsf{T}_\Omega^2 = \mathsf{T}_\Omega$ and $\mathsf{T}_\Omega^\dagger = \mathsf{T}_\Omega$, one has

$$\mathrm{Re}\,\langle V^{\mathrm{H}}|W^{\mathrm{H}}\rangle = \mathrm{Re}\,\langle V|\mathsf{T}_\Omega W\rangle = \mathrm{Re}\left(\langle V|W\rangle - \langle V|\Omega\rangle\,\langle\Omega|W\rangle\right).$$

By definition (eq. (6.168)), the right-hand side equals $g_T^{\mathrm{R}}(V,W)$. Combining with (6.171) yields

$$\mathrm{FS}_{[\Omega]}(V,W) = g_T^{\mathrm{R}}(V,W),$$

for the tangent classes represented by V, W.

Step 4 (Independence of representatives and gauge invariance). If $V \mapsto V + \alpha\, i|\Omega\rangle$ (vertical change), then $\mathsf{T}_\Omega(i|\Omega\rangle) = 0$, hence $g_T^{\mathrm{R}}(V+\alpha i\Omega, W) = g_T^{\mathrm{R}}(V,W)$ and the value depends only on the projective tangent classes. Under a unitary frame change U, $|\Omega\rangle \mapsto U|\Omega\rangle$, $\mathsf{T}_\Omega \mapsto U\mathsf{T}_\Omega U^\dagger$ (Proposition 6.95), and $\langle UV|U\mathsf{T}_\Omega UW\rangle = \langle V|\mathsf{T}_\Omega W\rangle$, so both sides remain invariant.

Step 5 (Degenerate case). If $\dim \mathcal{V}_0 = 1$, then $T_{|\Omega\rangle}S(\mathcal{V}_0) = \mathrm{span}\{i|\Omega\rangle\}$ is purely vertical, hence both sides vanish identically; the statement holds trivially.

Together, Steps 1–4 show that g_T^{R} is exactly the pullback of the Fubini–Study metric to the unit sphere of $\mathcal{V}_0$, establishing (6.170). □

Corollary 6.109 (Kinetic quadratic form for vacuum-preserving deformations)**.** *Let $x \mapsto |\Omega(x)\rangle \in \mathcal{M}_{\mathrm{vac}}$ be a smooth vacuum section. Define the horizontal derivative $|\nabla_\mu \Omega(x)\rangle := \mathsf{T}_{\Omega(x)}\, \partial_\mu |\Omega(x)\rangle$. Then the canonical quadratic form used in kinetic sectors is*

$$\boxed{\mathcal{K}_\Omega(x) \;:=\; g_T^{\mathrm{R}}\Big(|\nabla_\mu \Omega\rangle, |\nabla^\mu \Omega\rangle\Big) \;=\; \mathrm{Re}\,\langle \partial_\mu \Omega|\; \mathsf{T}_\Omega\; \partial^\mu \Omega\rangle\,,} \tag{6.172}$$

which is gauge/frame invariant by Proposition 6.107 and agrees with the Fubini–Study kinetic term on $\mathbb{P}(\mathcal{V}_0)$ by Proposition 6.108.

Remark 6.110 (Holosymmetric lift)**.** For a two-branch holosymmetric setting with ground spaces $\mathcal{V}_0^{(0)}$ and $\mathcal{V}_0^{(1)}$ and horizontal projectors $\mathsf{T}_\Omega^{(k)}$ on each branch, the direct-sum inner product

$$g_T^{\mathrm{HS}}\Big((V_0 \oplus V_1), (W_0 \oplus W_1)\Big) := g_T^{(0)}(V_0, W_0) + g_T^{(1)}(V_1, W_1)$$

is block-diagonal on fibres. After HS reduction (e.g. phase dial φ as in Section 6.1.15 and eq. (6.125)) the tangent inner product remains horizontal and unaffected by the dial; interference affects observable mixes but not the vacuum–tangent norm, in agreement with causal-structure neutrality results (cf. Remark 6.51 and Corollary 6.53).

6.1.26 The Banach Space of Bounded Operators

The algebra of endomorphisms, $\mathrm{End}(\mathcal{H}_{\mathrm{em}})$, is the "Banach space of bounded operators" is a precise mathematical assertion with several layers of meaning. Let us formally define the components of this structure.

Definition 6.111 (The Banach Space of Bounded Operators, $\mathcal{B}(\mathcal{H})$)**.** Let $\mathcal{H}$ be a Hilbert space. The set of all bounded linear operators from $\mathcal{H}$ to itself is denoted $\mathcal{B}(\mathcal{H})$ [253, 404]. This set forms a **Banach space**, which is a vector space that is complete with respect to a norm. The defining properties are:

1. **Vector Space Structure:** For any operators $A, B \in \mathcal{B}(\mathcal{H})$ and any scalar $\alpha \in \mathbb{C}$, the linear combination $(\alpha A + B)$ is also a bounded linear operator in $\mathcal{B}(\mathcal{H})$.
2. **The Operator Norm:** The space is equipped with the *operator norm*, $\|\cdot\|_{\mathrm{op}}$, defined for an operator A as:

$$\boxed{\|A\|_{\mathrm{op}} := \sup_{\substack{\psi \in \mathcal{H} \\ \|\psi\|=1}} \|A\psi\| = \sup_{\substack{\psi \in \mathcal{H} \\ \psi \neq 0}} \frac{\|A\psi\|}{\|\psi\|}.} \tag{6.173}$$

   This norm measures the maximum "stretching factor" that the operator applies to any vector in the Hilbert space. An operator is bounded if and only if its norm is finite [253].
3. **Completeness:** The space $\mathcal{B}(\mathcal{H})$ is *complete* with respect to the operator norm [253, 433]. This means that every Cauchy sequence of bounded operators, $\{A_n\}$, converges to a limit operator, A, that is also in $\mathcal{B}(\mathcal{H})$. This property is essential for the calculus of operators, ensuring that limits of well-behaved sequences are themselves well-behaved.

The Adjoint Map and the $*$-Algebra Structure Beyond being a Banach space, $\mathcal{B}(\mathcal{H})$ has additional structure crucial for quantum mechanics.

Definition 6.112 (The Adjoint Map and $*$-Algebra)**.** The space $\mathcal{B}(\mathcal{H})$ is equipped with an **involution**, known as the **adjoint map**, which sends an operator A to its Hermitian adjoint, $A^\dagger$ [404]. This map has the property that for any states $|\psi\rangle, |\phi\rangle \in \mathcal{H}$:

$$\langle \psi | A\phi \rangle = \langle A^\dagger \psi | \phi \rangle.$$

An algebra equipped with such an involution is called a **$*$-algebra** (or star-algebra). This structure is the foundation for defining **self-adjoint operators** ($A = A^\dagger$), which correspond to physical observables in quantum theory [195, 379].

By combining these definitions, we affirm that $\mathrm{End}(\mathcal{H}_{\mathrm{em}}) := \mathcal{B}(\mathcal{H}_{\mathrm{em}})$ is a **unital Banach $*$-algebra**. In fact, it satisfies the stronger condition of being a **C*-algebra**, which is the canonical mathematical framework for the algebra of observables in modern quantum physics [404, 254, 230].

6.1.27 The Algebra of Endomorphisms: The Arena for Quantum Operators

In the IQR framework, quantum fields are not mere functions on spacetime; they are operator-valued sections of a sophisticated geometric bundle. The space in which these operators "live" and from which they derive their algebraic properties is the algebra of endomorphisms over the emergent Hilbert space.

Definition 6.113 (Algebra of Endomorphisms over the Emergent Hilbert Space)**.** Let $\mathcal{H}_{\mathrm{em}}$ denote the total emergent Hilbert space of the theory (Definition 4.287). Then the symbol $\mathrm{End}(\mathcal{H}_{\mathrm{em}})$ refers to the unital $*$-algebra of all **bounded linear operators** on $\mathcal{H}_{\mathrm{em}}$ [253, 404]:

$$\mathrm{End}(\mathcal{H}_{\mathrm{em}}) := \mathcal{B}(\mathcal{H}_{\mathrm{em}}),$$

where $\mathcal{B}(\mathcal{H}_{\mathrm{em}})$ denotes the Banach space of bounded operators, equipped with the operator norm and an involution (the adjoint map, $A \mapsto A^\dagger$). This algebra provides the typical fiber for the endomorphism bundle over $\mathcal{X}_q$, from which all operator-valued fields are constructed as sections.

Interpretation and Significance This definition is laden with physical meaning, and understanding its components is crucial.

1. **Boundedness and the Operator Norm:** We restrict our attention to *bounded* operators. An operator A is bounded if there exists a constant M such that $\|A\psi\| \leq M\|\psi\|$ for all states $|\psi\rangle$. This property is essential because it guarantees that the operator does not "stretch" vectors infinitely, ensuring a well-behaved mathematical structure. It allows us to equip the space $\mathcal{B}(\mathcal{H}_{\mathrm{em}})$ with a natural norm, the *operator norm*, which turns it into a complete normed vector space (a Banach space) [253]. This provides the necessary topological structure to discuss the convergence and continuity of operator-valued fields.

2. **The $*$-Involution (Adjoint Operation):** The map $A \mapsto A^\dagger$ is the physically crucial operation of taking the Hermitian adjoint [195]. Its importance cannot be overstated. This involution allows us to define the most important class of operators in quantum mechanics: the *self-adjoint* (or Hermitian) operators, for which $A = A^\dagger$. According to the postulates of quantum theory, it is precisely these self-adjoint operators that correspond to physical observables, as they are guaranteed to have real eigenvalues [379, 195].

3. **The C*-Algebra Structure:** The combination of being a Banach space with a $*$-involution that satisfies the C*-identity, $\|A^\dagger A\| = \|A\|^2$, makes $\mathcal{B}(\mathcal{H}_{\text{em}})$ a *C*-algebra* [404]. This is the canonical mathematical structure for the algebra of observables in modern quantum physics [230]. The C*-identity provides a deep and powerful link between the algebraic properties of the operators (their products) and their topological properties (their norms), which is essential for a consistent theory. In many physical contexts, one works with specific subalgebras of $\mathcal{B}(\mathcal{H}_{\text{em}})$, such as von Neumann algebras, which have additional topological closure properties relevant to measurement theory and superselection sectors [404, 381].

By defining $\text{End}(\mathcal{H}_{\text{em}})$ as the algebra of bounded operators, we are choosing to go beyond mathematical conveniences and are endowing our theory with the precise and complete algebraic and topological structure required to rigorously describe quantum observables, their relationships, and their dynamics within the emergent, relational cosmos [254, 230].

6.1.28 Holosymmetric Wordwise Banach Space and Endomorphism Bundle

Setting. Let $\mathcal{H}_{\text{em}}$ be the emergent Hilbert space (Definition 4.287). Write $\text{End}(\mathcal{H}_{\text{em}}) = \mathcal{B}(\mathcal{H}_{\text{em}})$ for the bounded operators (Definitions 6.111 and 6.113).

Definition 6.114 (Operator-word vault; Banach/C* structure)**.** Define the *operator-word vault* by

$$\mathbb{W}_{\text{op}}(\mathcal{H}_{\text{em}}) \;:=\; \text{End}(\mathcal{H}_{\text{em}}) \;\equiv\; \mathcal{B}(\mathcal{H}_{\text{em}}).$$

Equip it with the operator norm

$$\boxed{\|\hat{O}\|_{\text{op}} \;:=\; \sup_{\psi\neq 0} \frac{\|\hat{O}\psi\|}{\|\psi\|} \;=\; \sup_{\|\psi\|=1} \|\hat{O}\psi\|.} \tag{6.174}$$

With the adjoint involution $\hat{O} \mapsto \hat{O}^\dagger$ and the usual product, $\left(\mathbb{W}_{\text{op}}(\mathcal{H}_{\text{em}}), \|\cdot\|_{\text{op}}, {}^\dagger\right)$ is a unital Banach $*$-algebra that satisfies the C*-identity

$$\boxed{\|\hat{O}^\dagger\hat{O}\|_{\text{op}} \;=\; \|\hat{O}\|_{\text{op}}^2, \qquad (\hat{O} \in \mathcal{B}(\mathcal{H}_{\text{em}})).} \tag{6.175}$$

Hence $\mathbb{W}_{\text{op}}(\mathcal{H}_{\text{em}})$ is a C*-algebra, and self-adjoint words $\hat{O} = \hat{O}^\dagger$ represent observable words.

Remark 6.115 (Equivalence with the standard presentation)**.** Equations (6.174)–(6.175) reproduce the "Banach space of bounded operators" and the unital C*-algebra structure of Definitions 6.111 and 6.112, with notation adapted to the wordwise grammar.

Definition 6.116 (Endomorphism word-bundle and transport covariance)**.** Let $\mathcal{X}_q$ carry the covariant kinematics of Section 4.5. For each $x \in \mathcal{X}_q$, let the modal fibre be $\mathcal{H}_x \cong \mathcal{H}_{\mathrm{em}}$. The *endomorphism word-bundle* is

$$\boxed{\mathfrak{End} \;:=\; \bigsqcup_{x\in\mathcal{X}_q} \mathcal{B}(\mathcal{H}_x) \;\longrightarrow\; \mathcal{X}_q,} \tag{6.176}$$

with typical fibre $\mathcal{B}(\mathcal{H}_x)$. Parallel transport along a path $\gamma : x' \to x$ acts fibrewise by inner automorphisms,

$$\hat{O}(x') \;\longmapsto\; \hat{O}_{\parallel}(x) \;:=\; U_\gamma(x \leftarrow x')\, \hat{O}(x')\, U_\gamma(x \leftarrow x')^\dagger, \tag{6.177}$$

preserving the $*$-structure and the operator norm.

Proposition 6.117 (Norm/$*$-preservation under transport)**.** *With U_γ unitary as in Section 4.5,*

$$\|\hat{O}_{\parallel}(x)\|_{\mathrm{op}} = \|\hat{O}(x')\|_{\mathrm{op}}, \qquad \big(\hat{O}_{\parallel}(x)\big)^\dagger = \big(\hat{O}(x')\big)^\dagger{}_{\parallel}(x),$$

hence each fibre carries canonically isomorphic C-algebras along admissible paths.*

Proof. Unitarity gives $\|\hat{O}_{\parallel}(x)\|_{\mathrm{op}} = \sup_{\|\psi\|=1} \|U_\gamma \hat{O}(x') U_\gamma^\dagger \psi\| = \sup_{\|\phi\|=1} \|\hat{O}(x')\phi\| = \|\hat{O}(x')\|_{\mathrm{op}}$. The adjoint identity follows from $(U\hat{O}U^\dagger)^\dagger = U\hat{O}^\dagger U^\dagger$. □

Holosymmetric lift on operator-words (no new symbols). Let $\mathcal{H}_L = \mathrm{span}\{|0\rangle_L, |1\rangle_L\}$ be a two-level law register with matrix units $P_0 := |0\rangle\langle 0|_L$, $P_1 := |1\rangle\langle 1|_L$, $E_{01} := |0\rangle\langle 1|_L$, $E_{10} := |1\rangle\langle 0|_L$. Given two branch operators $\hat{O}^{(0)}, \hat{O}^{(1)} \in \mathcal{B}(\mathcal{H}_{\mathrm{em}})$ and a *mixing operator* $\hat{O}_{\mathrm{mix}} \in \mathcal{B}(\mathcal{H}_{\mathrm{em}})$ (assembled covariantly from existing blocks of the theory, e.g. by combining the same sector words used in the single-branch construction), define

$$\boxed{\widehat{\boldsymbol{O}}^{\mathrm{HS}} \;:=\; P_0 \otimes \hat{O}^{(0)} \;+\; P_1 \otimes \hat{O}^{(1)} \;+\; E_{01} \otimes \hat{O}_{\mathrm{mix}} \;+\; E_{10} \otimes \hat{O}_{\mathrm{mix}}^\dagger.} \tag{6.178}$$

Preparing the control in $|+_\varphi\rangle_L = \frac{1}{\sqrt{2}}\big(|0\rangle_L + e^{i\varphi}|1\rangle_L\big)$ and reducing on $\mathcal{H}_L$ yields the *reduced* system operator

$$\boxed{\hat{O}^{\langle \mathrm{HS}\rangle}(\varphi) \;=\; \tfrac{1}{2}\big(\hat{O}^{(0)} + \hat{O}^{(1)}\big) \;+\; \mathrm{Re}\big(e^{-i\varphi}\hat{O}_{\mathrm{mix}}\big),} \tag{6.179}$$

which matches the general HS reduction pattern already used in Sections 6.1.12 and 6.1.15 without introducing any new symbols beyond the operator words $\hat{O}^{(k)}$ and $\hat{O}_{\mathrm{mix}}$. Gauge/-transport covariance follows from (6.177).

Topological note. For variational statements and coincidence limits in Chapter 6, the strong-operator topology (SOT) on each fibre can be employed in parallel with the norm topology that underlies the Banach/C* control; both are compatible with the wordwise constructions above.

From Algebra to Endomorphism: The Role of the GNS Construction

The connection between the Algebra of Observables (Section 1.4.1), The Gelfand-Naimark-Segal Construction (Definition 1.34), the Banach Space of Bounded Operators (Section 6.1.26), and the Algebra of Endomorphisms (Section 6.1.27) is one of *axiomatic foundation* and *concrete realization* [230, 404]. The theory begins with an abstract, physically motivated algebra of what can be observed, and from this, it constructs the familiar arena of quantum mechanics—the Hilbert space and the operators acting upon it.

1. The Starting Point: The Abstract Algebra of Observables The section on the *Algebra of Observables* (Section 1.4.1) introduces the foundational algebraic structure, denoted

$$\mathcal{A} := \mathcal{A}(\mathcal{C}_d),$$

This is a C*-algebra, defined axiomatically from the Gaiamiti and Tantrakyma fields. Crucially, at this stage, the theory makes *no mention of a Hilbert space* nor does it presuppose any representation. This is the essence of the algebraic approach to quantum theory: to define the system by the relationships between its observables (their sums, products, and adjoints) rather than by postulating a space for states to live in [230, 381]. It is the abstract "language" of the theory. At this stage, the theory is *representation-independent*, encoding only what operations are possible and how they relate.

2. The Bridge: The GNS Construction The *GNS Construction of the Quantum Holo-State* (Definition 1.34) provides the mechanism by which a concrete Hilbert space arises from the abstract algebra $\mathcal{A}$. Given a pure state functional

$$\omega^{\mathring{\mathbf{\Psi}}} : \mathcal{A} \to \mathbb{C},$$

the GNS theorem (Gelfand-Naimark-Segal) [254, 404] guarantees the existence of a unique (up to unitary equivalence) triplet $(\mathcal{H}_{\text{em}}, \pi^{\mathring{\mathbf{\Psi}}}, |\mathring{\mathbf{\Psi}}\rangle)$, consisting of:

- A Hilbert space $\mathcal{H}_{\text{em}}$,
- A *-representation $\pi^{\mathring{\mathbf{\Psi}}} : \mathcal{A} \to \mathcal{B}(\mathcal{H}_{\text{em}})$,
- A cyclic vector $|\mathring{\mathbf{\Psi}}\rangle \in \mathcal{H}_{\text{em}}$,

such that all expectation values are recovered by

$$\omega^{\mathring{\mathbf{\Psi}}}(A) = \langle \mathring{\mathbf{\Psi}} | \pi^{\mathring{\mathbf{\Psi}}}(A) | \mathring{\mathbf{\Psi}} \rangle.$$

This construction, central to the mathematical foundations of quantum mechanics [230, 195], provides the rigorous emergence of the total Hilbert space $\mathcal{H}_{\text{em}}$ (Definition 4.287) and the concrete representation of observables within it.

3. The Destination: The Concrete Algebra of Endomorphisms The sections *The Banach Space of Bounded Operators* (Section 6.1.26) and *The Algebra of Endomorphisms* (Section 6.1.27) describe the analytic and algebraic structure of the target space of the GNS representation:

$$\pi^{\mathring{\mathbf{\Psi}}}(\mathcal{A}) \subseteq \mathcal{B}(\mathcal{H}_{\text{em}}) =: \text{End}(\mathcal{H}_{\text{em}}).$$

Here, $\mathcal{B}(\mathcal{H}_{\text{em}})$ is the Banach space of bounded linear operators equipped with the operator norm [253], and $\text{End}(\mathcal{H}_{\text{em}})$ is its interpretation as a unital, involutive algebra: the *concrete operator arena* for the quantum field theory [404].

This is the space in which all relational operators—including the Dynamical Gaiamiti Field $\hat{\mathbf{\Phi}}(x)$ (Definition 6.118)—take their values.

Synthesis: Logical Sequence of Construction

The full logical flow from abstract algebraic structures to concrete operator-theoretic dynamics may be summarized as follows:

1. Abstract Algebra of Observables
The foundational, representation-independent C*-algebra $\mathcal{A}(\mathcal{C}_d)$ describing all potential physical relations.

↓ *The universal state $\omega^{\mathring{\mathbf{\Psi}}}$ selects a physical representation.*

2. GNS Construction
The emergent Hilbert space $\mathcal{H}_{\text{em}}$, representation $\pi^{\mathring{\mathbf{\Psi}}}$, and cyclic vector $|\mathring{\mathbf{\Psi}}\rangle$ generated from the abstract algebra via the state functional $\omega^{\mathring{\mathbf{\Psi}}}$.

↓ *The representation map $\pi^{\mathring{\mathbf{\Psi}}}$ provides a concrete realization of the algebra.*

3. Concrete Operator Algebra
The algebra of endomorphisms $\text{End}(\mathcal{H}_{\text{em}})$, where abstract observables are now realized as bounded operators on the emergent Hilbert space.

Figure 6.1: The logical cascade from the abstract Algebra of Observables to the concrete Algebra of Endomorphisms, mediated by the GNS construction.

Interpretive Conclusion

The sections on the GNS construction (Definition 1.34) describe the *engine of emergence*—the mechanism by which the physical Hilbert space and operator algebra are derived from purely algebraic data. The sections on bounded operators and endomorphisms (Sections 6.1.26 and 6.1.27) describe the *realized mathematical infrastructure*—the analytic and algebraic environment in which quantum-relational dynamics are now formulated.

This architecture is essential for the proper formulation of all field-theoretic objects in the theory. The Gaiamiti Field, for instance, is defined as:

$$\hat{\mathbf{\Phi}}(x) \in \mathrm{End}(\mathcal{H}_{\mathrm{em}}) \otimes T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}},$$

and can only be interpreted once the endomorphism algebra and the vacuum manifold (with its tangent bundle structure) are fully constructed. In this way, the transition from the algebraic to the operator-theoretic formalism is not only a technical pathway—it is the fundamental bridge from the axioms of relational quantum theory to the dynamics of emergent geometry.

6.2 The Dynamical Gaiamiti Field

The **Gaiamiti Field** is the fundamental dynamical entity responsible for generating relational geometry across the emergent configuration space, $\mathcal{X}_q$. It encodes infinitesimal deformations of the vacuum manifold, $\mathcal{M}_{\mathrm{vac}}$, and serves as the field-theoretic progenitor of all emergent geometric observables—including the operator-valued metric tensor $\hat{g}_{\mu\nu}(x)$, the temporal tensor $\hat{t}_{\mu\nu}(x)$, and their unified sum $\hat{\Sigma}_{\mu\nu}(x)$.

Definition 6.118 (Dynamical Gaiamiti Field as an Operator-Valued Vector Field)**.** Let $\mathcal{H}_{\mathrm{em}} := \bigsqcup_{x\in\mathcal{X}_q} \mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ denote the total emergent Hilbert bundle (Definition 4.287), and let $\mathcal{M}_{\mathrm{vac}}$ be the vacuum manifold defined as the normalized ground-state eigenspace of the Relational Hamiltonian (Definition 6.61).

Then the *Dynamical Gaiamiti Field*, denoted $\hat{\mathbf{\Phi}}(x)$, is a smooth operator-valued section over $\mathcal{X}_q$ of the fiber bundle:

$$\boxed{\hat{\mathbf{\Phi}} : \mathcal{X}_q \longrightarrow \mathrm{End}(\mathcal{H}_{\mathrm{em}}) \otimes T\mathcal{M}_{\mathrm{vac}},} \tag{6.180}$$

with local evaluations:

$$\boxed{\hat{\mathbf{\Phi}}(x) \in \mathrm{End}(\mathcal{H}_{\mathrm{em}}) \otimes T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}},} \tag{6.181}$$

where:

- $\mathrm{End}(\mathcal{H}_{\mathrm{em}})$ denotes the algebra of bounded linear operators on the total emergent Hilbert space;
- $|\Omega(x)\rangle \in \mathcal{M}_{\mathrm{vac}}$ is the locally chosen vacuum state at $x \in \mathcal{X}_q$, defining the reference fiber of the tangent space;
- $T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}$ is the operator-theoretic tangent space at $|\Omega(x)\rangle$, as defined in Definition 6.90.

Operationally, $\hat{\mathbf{\Phi}}(x)$ acts fiberwise on the modal sector $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)} \subset \mathcal{H}_{\text{em}}$, and is extended globally via the smooth structure of the bundle $\mathcal{H}_{\text{em}}$. It thus represents a coherent, covariant fluctuation of the vacuum geometry—a *dynamical field of geometric deformation operators*.

Dual Character and Relational Interpretation The definition of $\hat{\mathbf{\Phi}}(x)$ as a section of the bundle $\text{End}(\mathcal{H}_{\text{em}}) \otimes T\mathcal{M}_{\text{vac}}$ entails a dual functional character:

1. As an element of $\text{End}(\mathcal{H}_{\text{em}})$, the Gaiamiti field governs relational transformations of the quantum state: it acts on modal components of $|\mathbf{\Psi}\rangle$ to generate dynamical evolution at the operator-theoretic level.

2. As an element of $T_{|\Omega(x)\rangle}\mathcal{M}_{\text{vac}}$, it specifies a direction of infinitesimal geometric variation within the vacuum manifold: a Goldstone-type fluctuation of zero excitation energy, preserving the ground-state energy E_0 while modifying the local relational configuration.

In essence, $\hat{\mathbf{\Phi}}(x)$ is a *quantum-geometric deformation operator*: it tells us *what action* to perform in Hilbert space and *in which direction* to deform the vacuum geometry. All emergent geometric quantities—metric, connection, curvature—are functionals derived from the algebraic and differential structure of $\hat{\mathbf{\Phi}}(x)$.

6.2.1 Holosymmetric Wordwise Dynamical Gaiamiti Field

Setting. Let $\mathcal{H}_{\text{em}} = \bigsqcup_{x\in\mathcal{X}_q} \mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}$ be the emergent Hilbert bundle (Definition 4.287) and let $\mathcal{M}_{\text{vac}} \subset \mathcal{H}_{\text{em}}$ be the vacuum manifold (Definitions 6.61 and 6.90). Write $\mathbb{W}_{\text{op}}(\mathcal{H}_{\text{em}}) = \mathcal{B}(\mathcal{H}_{\text{em}})$ for the operator-word vault (Definition 6.114).

Definition 6.119 (Gaiamiti word-field; typed operator–tangent word)**.** Choose a smooth local vacuum section $x \mapsto |\Omega(x)\rangle \in \mathcal{M}_{\text{vac}}$ and set $\Pi_0(x)$ for the projector onto the local ground space and $\Pi_T(x) := \Pi_0(x) - |\Omega(x)\rangle\langle\Omega(x)|$ for the horizontal (vacuum–tangent) projector (cf. Definition 6.90). The *dynamical Gaiamiti field* is the typed operator–tangent word

$$\boxed{\hat{\mathbf{\Phi}} : \mathcal{X}_q \longrightarrow \mathbb{W}_{\text{op}}(\mathcal{H}_{\text{em}}) \otimes T\mathcal{M}_{\text{vac}}, \qquad \hat{\mathbf{\Phi}}(x) \in \mathcal{B}(\mathcal{H}_{\text{em}}) \otimes T_{|\Omega(x)\rangle}\mathcal{M}_{\text{vac}}.} \tag{6.182}$$

Regularity is taken in the strong–operator topology (SOT); spectral localisation is imposed by a local cutoff $P_x = E^{(x)}([0,\Lambda])$ with $P_x\hat{\mathbf{\Phi}}(x)P_x = \hat{\mathbf{\Phi}}(x)$.

Definition 6.120 (Transport and horizontal projection of the Gaiamiti word-field)**.** Let $\mathcal{U}(x \leftarrow x';\gamma)$ be the parallel transporter on the modal bundle (Section 4.5; cf. eqs. (4.11) and (4.14)), and let $\Pi_T(x) = \Pi_0(x) - |\Omega(x)\rangle\langle\Omega(x)|$ be the horizontal projector (Definition 6.90). For a dynamical Gaiamiti word-field $\hat{\mathbf{\Phi}}$ as in Definition 6.119, define its transport to x followed by horizontal projection as

$$\hat{\mathbf{\Phi}}_{\parallel}(x) \;:=\; \big(\mathbf{1}\otimes\Pi_T(x)\big)\Big[\,\mathcal{U}(x\leftarrow x')\;\hat{\mathbf{\Phi}}(x')\;\mathcal{U}(x\leftarrow x')^{\dagger}\,\Big]. \tag{6.183}$$

The definition is independent of the choice of path γ to first order in $|x-x'|$; path dependence enters only at $\mathcal{O}(|x-x'|^2)$ via curvature terms (see Section 4.5).

Definition 6.121 (Wordwise covariant gradient of the Gaiamiti word-field)**.** Assume SOT–smoothness of $x \mapsto \hat{\mathbf{\Phi}}(x)$ and $x \mapsto \Pi_T(x)$ and that $\mathcal{U}(x \leftarrow x')$ admits a C^1 coincidence expansion (Definition 6.7 and Section 6.1.4). The *wordwise covariant gradient* of $\hat{\mathbf{\Phi}}$ is the SOT limit

$$\boxed{\mathbb{D}_\mu \hat{\mathbf{\Phi}}(x) \;:=\; \lim_{x' \to x} \frac{\hat{\mathbf{\Phi}}(x) - \hat{\mathbf{\Phi}}_{\|}(x)}{\Delta x^\mu}\,.} \tag{6.184}$$

Definition 6.122 (Operator-slot covariant derivative)**.** For later comparison, define the operator-slot covariant derivative

$$\nabla_\mu^{(\mathcal{U})} \hat{\mathbf{\Phi}}(x) \;:=\; \lim_{x' \to x} \frac{\hat{\mathbf{\Phi}}(x) - \mathcal{U}(x \leftarrow x')\, \hat{\mathbf{\Phi}}(x')\, \mathcal{U}(x \leftarrow x')^\dagger}{\Delta x^\mu}, \tag{6.185}$$

which is the wordwise version of the HS covariant derivative on words (Definitions 6.11 and 6.185).

Proposition 6.123 (Gauge/transport covariance)**.** *Under a local endpoint gauge $g(x)$ (fibrewise unitary),*

$$\boxed{\begin{aligned} \hat{\mathbf{\Phi}}(x) &\mapsto \mathbf{g}(x)\, \hat{\mathbf{\Phi}}(x)\, \mathbf{g}(x)^\dagger, \\ \mathcal{U}(x \leftarrow x') &\mapsto g(x)\, \mathcal{U}(x \leftarrow x')\, g(x')^\dagger, \\ \Pi_T(x) &\mapsto g(x)\, \Pi_T(x)\, g(x)^\dagger. \end{aligned}} \tag{6.186}$$

(Proposition 6.66 and Section 4.5). Then both the transported field (6.183) and the covariant gradient (6.184) transform by conjugation:

$$\hat{\mathbf{\Phi}}_{\|}(x) \;\mapsto\; \mathbf{g}(x)\, \hat{\mathbf{\Phi}}_{\|}(x)\, \mathbf{g}(x)^\dagger, \qquad \mathbb{D}_\mu \hat{\mathbf{\Phi}}(x) \;\mapsto\; \mathbf{g}(x)\, \mathbb{D}_\mu \hat{\mathbf{\Phi}}(x)\, \mathbf{g}(x)^\dagger. \tag{6.187}$$

Proof idea. Insert the gauge transforms into (6.183); the $(\mathbf{1} \otimes \Pi_T)$ slot and the operator slot both transform by conjugation, and the transporter obeys the endpoint law (Section 4.5). The limit (6.184) then inherits the same conjugation law.

Proof (expanded). We prove covariance for the transported field (6.183) and then for the covariant gradient (6.184). Throughout, let $g(x)$ be a fibrewise unitary (endpoint gauge), let $\mathbf{g}(x)$ denote the induced inner automorphism on operators, and recall the endpoint law for the transporter $\mathcal{U}(x \leftarrow x') \mapsto g(x)\, \mathcal{U}(x \leftarrow x')\, g(x')^\dagger$ (Section 4.5). The horizontal projector transforms as $\Pi_T(x) \mapsto g(x) \Pi_T(x) g(x)^\dagger$ by definition (Definition 6.90) and gauge covariance of ground projectors (Proposition 6.66).

(1) Transported field. Define the gauge–transformed field by $\widetilde{\hat{\mathbf{\Phi}}}(x) := \mathbf{g}(x)\, \hat{\mathbf{\Phi}}(x)\, \mathbf{g}(x)^\dagger$, and similarly $\widetilde{\Pi_T}(x) := g(x) \Pi_T(x) g(x)^\dagger$, $\widetilde{\mathcal{U}}(x \leftarrow x') := g(x) \mathcal{U}(x \leftarrow x') g(x')^\dagger$. Insert these in (6.183):

$$\begin{aligned} \widetilde{\hat{\mathbf{\Phi}}}_{\|}(x) &= \Big(\mathbf{1} \otimes \widetilde{\Pi_T}(x)\Big)\, \mathrm{Ad}_{\widetilde{\mathcal{U}}(x \leftarrow x')}\Big(\widetilde{\hat{\mathbf{\Phi}}}(x')\Big) && (6.188) \\ &= \Big(\mathbf{1} \otimes g(x) \Pi_T(x) g(x)^\dagger\Big)\, \mathrm{Ad}_{g(x)\mathcal{U}(x \leftarrow x') g(x')^\dagger}\Big(g(x') \hat{\mathbf{\Phi}}(x') g(x')^\dagger\Big) && (6.189) \\ &= g(x)\, \Big(\mathbf{1} \otimes \Pi_T(x)\Big)\, \mathrm{Ad}_{\mathcal{U}(x \leftarrow x')}\Big(\hat{\mathbf{\Phi}}(x')\Big)\, g(x)^\dagger && (6.190) \\ &= \mathbf{g}(x)\, \hat{\mathbf{\Phi}}_{\|}(x)\, \mathbf{g}(x)^\dagger. && (6.191) \end{aligned}$$

where in the penultimate step we used that $(\mathbf{1}\otimes g)\,X\,(\mathbf{1}\otimes g^\dagger) = g\,X\,g^\dagger$ for the tensor-typed operator and the fact that $g(x)$ acts trivially on the basepoint x'-slot after cancellation of $g(x')^\dagger g(x') = \mathbf{1}$. This proves the first covariance claim.

(2) Covariant gradient. Write the transformed difference quotient underlying (6.184) as

$$\begin{aligned}\widetilde{\Delta_\mu}(x,x') &:= \frac{\tilde{\hat{\mathbf{\Phi}}}(x) - \tilde{\hat{\mathbf{\Phi}}}_\parallel(x)}{\Delta x^\mu}\\ &= \frac{\mathbf{g}(x)\hat{\mathbf{\Phi}}(x)\mathbf{g}(x)^\dagger - \mathbf{g}(x)\hat{\mathbf{\Phi}}_\parallel(x)\mathbf{g}(x)^\dagger}{\Delta x^\mu}\\ &= \mathbf{g}(x)\,\frac{\hat{\mathbf{\Phi}}(x) - \hat{\mathbf{\Phi}}_\parallel(x)}{\Delta x^\mu}\,\mathbf{g}(x)^\dagger.\end{aligned}$$

By assumption $g(\cdot)$ is continuous (indeed smooth) in x, so for fixed x the multipliers $\mathbf{g}(x), \mathbf{g}(x)^\dagger$ are *bounded* and independent of x'. Because inner automorphisms by bounded unitaries are SOT–continuous, we may pass them through the SOT limit:

$$\widetilde{\mathbb{D}_\mu\hat{\mathbf{\Phi}}}(x) := \lim_{x'\to x}\widetilde{\Delta_\mu}(x,x') = \mathbf{g}(x)\left(\lim_{x'\to x}\frac{\hat{\mathbf{\Phi}}(x) - \hat{\mathbf{\Phi}}_\parallel(x)}{\Delta x^\mu}\right)\mathbf{g}(x)^\dagger = \mathbf{g}(x)\,\mathbb{D}_\mu\hat{\mathbf{\Phi}}(x)\,\mathbf{g}(x)^\dagger,$$

which is the desired conjugation law.

(3) Regularity and path remarks. The limits exist under the SOT–C^1 regularity of $x\mapsto\hat{\mathbf{\Phi}}(x)$ and $x\mapsto\Pi_T(x)$ together with the C^1 endpoint expansion of $\mathcal{U}(x\leftarrow x')$ provided by the relational Levi–Civita structure (Definition 6.7 and Section 6.1.4). Any path dependence in $\mathcal{U}$ contributes only at $\mathcal{O}(|x-x'|^2)$ (curvature term), so it vanishes in the first-order quotient defining (6.184) (Section 4.5). Finally, the same algebra shows triad-wise covariance (corr/vac/spec) because each ingredient transforms by conjugation within its sector (eq. (5.86)).

This establishes the stated covariance of both $\hat{\mathbf{\Phi}}_\parallel(x)$ and $\mathbb{D}_\mu\hat{\mathbf{\Phi}}(x)$. □

Proposition 6.124 (Horizontal projection of the operator-slot derivative)**.** *Under the regularity assumptions of Definition 6.121, one has*

$$\mathbb{D}_\mu\,\hat{\mathbf{\Phi}}(x) = \Big(\mathbf{1}\otimes\Pi_T(x)\Big)\,\nabla_\mu^{(\mathcal{U})}\hat{\mathbf{\Phi}}(x)\,. \tag{6.192}$$

Proof idea. Expand $\mathcal{U}(x\leftarrow x') = \mathbf{1} - \mathcal{A}_\nu(x)\,\Delta x^\nu + \mathcal{O}(|\Delta x|^2)$ (Definitions 6.7 and 6.10), use SOT–smoothness of Π_T and $\hat{\mathbf{\Phi}}$, and keep terms up to $\mathcal{O}(\Delta x)$. Subtracting and dividing by Δx^μ yields the stated identity; details mirror the standard HS covariant calculus (Definitions 6.11 and 6.185).

Proof (expanded). We show that the difference quotient defining $\mathbb{D}_\mu\hat{\mathbf{\Phi}}(x)$ (Definition 6.121 and eq. (6.184)) equals the horizontal projection of the operator–slot derivative $\nabla_\mu^{(\mathcal{U})}\hat{\mathbf{\Phi}}(x)$ (Definition 6.122). Fix $x\in\mathcal{X}_q$.

Step 1: Insert the horizontal projector on the x–slot. By definition of the typed word-field (Definition 6.119 and eq. (6.182)), $\hat{\mathbf{\Phi}}(x) \in \mathcal{B}(\mathcal{H}_{\mathrm{em}}) \otimes T_{|\Omega(x))}\mathcal{M}_{\mathrm{vac}}$. Since $\Pi_T(x)$ is the projector onto $T_{|\Omega(x))}\mathcal{M}_{\mathrm{vac}}$ (Definition 6.90), it acts as the identity on the tangent slot of $\hat{\mathbf{\Phi}}(x)$:

$$(\mathbf{1} \otimes \Pi_T(x))\, \hat{\mathbf{\Phi}}(x) = \hat{\mathbf{\Phi}}(x). \tag{6.193}$$

Step 2: Rewrite the transported difference with a common projector. Using (6.183) and (6.193),

$$\hat{\mathbf{\Phi}}(x) - \hat{\mathbf{\Phi}}_{\|}(x) = \Big(\mathbf{1} \otimes \Pi_T(x)\Big)\hat{\mathbf{\Phi}}(x) - \Big(\mathbf{1} \otimes \Pi_T(x)\Big)\Big[\mathcal{U}(x \leftarrow x')\, \hat{\mathbf{\Phi}}(x')\, \mathcal{U}(x \leftarrow x')^{\dagger}\Big].$$

Factor $(\mathbf{1} \otimes \Pi_T(x))$:

$$\hat{\mathbf{\Phi}}(x) - \hat{\mathbf{\Phi}}_{\|}(x) = \Big(\mathbf{1} \otimes \Pi_T(x)\Big)\Big[\hat{\mathbf{\Phi}}(x) - \mathcal{U}(x \leftarrow x')\, \hat{\mathbf{\Phi}}(x')\, \mathcal{U}(x \leftarrow x')^{\dagger}\Big]. \tag{6.194}$$

Step 3: Pass to the difference quotient and the SOT–limit. Divide (6.194) by $\Delta x^{\mu} := x^{\mu} - x'^{\mu}$ and take the SOT limit $x' \to x$:

$$\mathbb{D}_{\mu}\hat{\mathbf{\Phi}}(x) = \lim_{x' \to x} \frac{\hat{\mathbf{\Phi}}(x) - \hat{\mathbf{\Phi}}_{\|}(x)}{\Delta x^{\mu}} = \Big(\mathbf{1} \otimes \Pi_T(x)\Big) \lim_{x' \to x} \frac{\hat{\mathbf{\Phi}}(x) - \mathcal{U}(x \leftarrow x')\, \hat{\mathbf{\Phi}}(x')\, \mathcal{U}(x \leftarrow x')^{\dagger}}{\Delta x^{\mu}}.$$

Here we used that multiplication by the bounded projector $(\mathbf{1} \otimes \Pi_T(x))$ is SOT–continuous and independent of x'; the existence of the limit follows from the regularity assumptions in Definition 6.121 together with the C^1 endpoint expansion of $\mathcal{U}$ (Definition 6.7 and Section 6.1.4). By Definition 6.122, the bracketed limit is precisely $\nabla^{(\mathcal{U})}_{\mu}\hat{\mathbf{\Phi}}(x)$, hence

$$\mathbb{D}_{\mu}\hat{\mathbf{\Phi}}(x) = \Big(\mathbf{1} \otimes \Pi_T(x)\Big)\, \nabla^{(\mathcal{U})}_{\mu}\hat{\mathbf{\Phi}}(x),$$

which is (6.192).

Step 4: (Optional) First-order expansion check. Writing $\mathcal{U}(x \leftarrow x') = \mathbf{1} - \mathcal{A}_{\nu}(x)\, \Delta x^{\nu} + \mathcal{O}(|\Delta x|^2)$ (Definitions 6.7 and 6.10) and using SOT–C^1 of $x \mapsto \hat{\mathbf{\Phi}}(x)$, one finds

$$\frac{\hat{\mathbf{\Phi}}(x) - \mathcal{U}\, \hat{\mathbf{\Phi}}(x')\, \mathcal{U}^{\dagger}}{\Delta x^{\mu}} = \partial_{\mu}\hat{\mathbf{\Phi}}(x) - [\mathcal{A}_{\mu}(x), \hat{\mathbf{\Phi}}(x)] + \mathcal{O}(|\Delta x|),$$

whose horizontal projection gives exactly the right-hand side of (6.192). Path (holonomy) dependence contributes only at $\mathcal{O}(|\Delta x|)$ beyond first order by Section 4.5.

Step 5: Triadic compatibility. All steps commute with the corr/vac/spec decomposition of the chronogeometric data (eq. (5.86)), hence the identity holds sector-wise.

This completes the proof. □

Corollary 6.125 (Expectation-level gauge invariance)**.** *For any ringed-ket state $|\mathring{\mathbf{\Psi}}\rangle$, the scalar densities $\langle\mathring{\mathbf{\Psi}}|\hat{\mathbf{\Phi}}_{\|}(x)|\mathring{\mathbf{\Psi}}\rangle$ and $\langle\mathring{\mathbf{\Psi}}|\mathbb{D}_{\mu}\hat{\mathbf{\Phi}}(x)|\mathring{\mathbf{\Psi}}\rangle$ are gauge invariant by conjugation covariance and the invariance of ringed-ket expectations (Remark 6.35).*

Proof. Let $\rho_\circ := |\overset{\circ}{\boldsymbol{\Psi}}\rangle\langle\overset{\circ}{\boldsymbol{\Psi}}|$. By Proposition 6.123, under any endpoint gauge $g(x)$ (the induced unitary $\mathbf{g}(x)$ on operators) we have

$$\hat{\boldsymbol{\Phi}}_{\parallel}(x) \;\mapsto\; \mathbf{g}(x)\,\hat{\boldsymbol{\Phi}}_{\parallel}(x)\,\mathbf{g}(x)^{\dagger}, \qquad \mathbb{D}_\mu\hat{\boldsymbol{\Phi}}(x) \;\mapsto\; \mathbf{g}(x)\,\mathbb{D}_\mu\hat{\boldsymbol{\Phi}}(x)\,\mathbf{g}(x)^{\dagger}.$$

The ringed–ket expectation is gauge invariant in the sense of Remark 6.35: the state $\rho_\circ$ is invariant under the inner automorphism generated by $\mathbf{g}(x)$ (equivalently, $\mathbf{g}(x)^{\dagger}\rho_\circ\mathbf{g}(x) = \rho_\circ$), and the expectation $\langle\overset{\circ}{\boldsymbol{\Psi}}|\cdot|\overset{\circ}{\boldsymbol{\Psi}}\rangle = \mathrm{Tr}(\rho_\circ\,\cdot)$ is thus conjugation–invariant.

Therefore,

$$\begin{aligned}
\Big\langle\overset{\circ}{\boldsymbol{\Psi}}\Big|\,\mathbf{g}\,\hat{\boldsymbol{\Phi}}_{\parallel}(x)\,\mathbf{g}^{\dagger}\,\Big|\overset{\circ}{\boldsymbol{\Psi}}\Big\rangle &= \mathrm{Tr}\Big(\rho_\circ\,\mathbf{g}\,\hat{\boldsymbol{\Phi}}_{\parallel}(x)\,\mathbf{g}^{\dagger}\Big) && (6.195)\\
&= \mathrm{Tr}\Big(\mathbf{g}^{\dagger}\rho_\circ\mathbf{g}\,\hat{\boldsymbol{\Phi}}_{\parallel}(x)\Big) && (6.196)\\
&= \mathrm{Tr}\Big(\rho_\circ\,\hat{\boldsymbol{\Phi}}_{\parallel}(x)\Big) && (6.197)\\
&= \Big\langle\overset{\circ}{\boldsymbol{\Psi}}\Big|\,\hat{\boldsymbol{\Phi}}_{\parallel}(x)\Big|\overset{\circ}{\boldsymbol{\Psi}}\Big\rangle, && (6.198)
\end{aligned}$$

$$\begin{aligned}
\Big\langle\overset{\circ}{\boldsymbol{\Psi}}\Big|\,\mathbf{g}\,\mathbb{D}_\mu\hat{\boldsymbol{\Phi}}(x)\,\mathbf{g}^{\dagger}\,\Big|\overset{\circ}{\boldsymbol{\Psi}}\Big\rangle &= \mathrm{Tr}\Big(\rho_\circ\,\mathbf{g}\,\mathbb{D}_\mu\hat{\boldsymbol{\Phi}}(x)\,\mathbf{g}^{\dagger}\Big) && (6.199)\\
&= \mathrm{Tr}\Big(\mathbf{g}^{\dagger}\rho_\circ\mathbf{g}\,\mathbb{D}_\mu\hat{\boldsymbol{\Phi}}(x)\Big) && (6.200)\\
&= \mathrm{Tr}\Big(\rho_\circ\,\mathbb{D}_\mu\hat{\boldsymbol{\Phi}}(x)\Big) && (6.201)\\
&= \Big\langle\overset{\circ}{\boldsymbol{\Psi}}\Big|\,\mathbb{D}_\mu\hat{\boldsymbol{\Phi}}(x)\Big|\overset{\circ}{\boldsymbol{\Psi}}\Big\rangle. && (6.202)
\end{aligned}$$

Hence both scalar densities are gauge invariant.

It remains to justify passage to the limit implicit in $\mathbb{D}_\mu$. By definition (Definition 6.121 and eq. (6.184)), $\mathbb{D}_\mu\hat{\boldsymbol{\Phi}}(x) = \lim_{x'\to x}\Delta_\mu(x,x')$ with $\Delta_\mu(x,x') = \frac{\hat{\boldsymbol{\Phi}}(x)-\hat{\boldsymbol{\Phi}}_{\parallel}(x)}{\Delta x^\mu}$. The functional $A \mapsto \langle\overset{\circ}{\boldsymbol{\Psi}}|A|\overset{\circ}{\boldsymbol{\Psi}}\rangle = \mathrm{Tr}(\rho_\circ A)$ is bounded and linear; since the limit exists in the SOT by the regularity assumptions of Definition 6.121, we may exchange the limit with the expectation, so the above conjugation–invariance applies pointwise to the limit as well.

All statements hold sectorwise (corr/vac/spec) because the ingredients transform by conjugation within each sector and the ringed–ket reduction is sector-preserving (eq. (5.86) and Remark 6.35). □

Remark 6.126 (Well-posedness, spectral localisation, and triadic structure)**.** (i) The SOT limit in (6.184) exists whenever $\hat{\boldsymbol{\Phi}}$ and Π_T are SOT–C^1 and the transporter is C^1 in the sense of Section 4.5; locality is enforced by the spectral cutoff $P_x = E^{(x)}([0,\Lambda])$ (Definition 6.119).

(ii) The corr/vac/spec triadic decomposition of the chronogeometric tensors (eq. (5.86)) induces a corresponding split of $\mathcal{U}$ and of the densities acted on by $\mathbb{D}_\mu$, and the constructions above preserve this split.

(iii) Since the dial control in Section 6.1.12 modifies only mixed-sector integrands, $\mathbb{D}_\mu$ is unaffected at fixed $\bar{\Sigma}$; causal cones remain determined by $\bar{g}^{\mathrm{Lor}}$ (Definition 5.77 and Section 5.4.2).

Definition 6.127 (Interaction contribution to the temporal sector (wordwise))**.** Let $\hat{\mathbf{\Phi}}$ be the Gaiamiti word-field of Section 6.2.1 and Definition 6.119, and let $\hat{\mathbf{\Theta}}$ be a companion typed operator–tangent word with the same spectral support on the emergent bundle.[3] Let $\mathbb{D}_\mu$ be the wordwise covariant gradient (Definition 6.121 and eq. (6.184)) and let Π_{time} denote the temporal projector from the wordwise metric projection machinery (Section 5.3.11). Using the canonical bilinears H, A from (6.204) (no redefinition here), define the *commutator-type interaction* contribution to the operator-valued temporal tensor by

$$\boxed{\hat{t}^{(\text{int})}_{\mu\nu}(x) \;:=\; \Pi_{\text{time}}\Big(\mathsf{A}\Big[\mathbb{D}_\mu\hat{\mathbf{\Phi}},\, \mathbb{D}_\nu\hat{\mathbf{\Theta}}\Big] + \mathsf{A}\Big[\mathbb{D}_\mu\hat{\mathbf{\Theta}},\, \mathbb{D}_\nu\hat{\mathbf{\Phi}}\Big]\Big)(x),} \tag{6.203}$$

to be added to the temporal sector $\hat{t}_{\mu\nu}$ of (6.205) when interaction channels are enabled (cf. Section 6.1.12).

Proposition 6.128 (Gauge/transport covariance of $\hat{t}^{(\text{int})}_{\mu\nu}$)**.** *Under endpoint gauges $g(x)$ and parallel transport as in Proposition 6.123 and Section 4.5, each factor in (6.203) transforms by conjugation; hence*

$$\hat{t}^{(\text{int})}_{\mu\nu}(x) \;\longmapsto\; \mathbf{g}(x)\,\hat{t}^{(\text{int})}_{\mu\nu}(x)\,\mathbf{g}(x)^\dagger.$$

Consequently, $\bar{t}^{(\text{int})}_{\mu\nu}(x) = \langle \mathring{\mathbf{\Psi}} | \hat{t}^{(\text{int})}_{\mu\nu}(x) | \mathring{\mathbf{\Psi}} \rangle$ is gauge invariant (Corollary 6.125).

Proof. By Proposition 6.123, $\mathbb{D}_\mu\hat{\mathbf{\Phi}} \mapsto \mathbf{g}\,(\mathbb{D}_\mu\hat{\mathbf{\Phi}})\,\mathbf{g}^\dagger$ and similarly for $\mathbb{D}_\mu\hat{\mathbf{\Theta}}$. The bilinear $\mathsf{A}[\cdot,\cdot]$ is built from commutators (see (6.204)), hence is natural under inner automorphisms: $\mathsf{A}[\mathbf{g}X\mathbf{g}^\dagger, \mathbf{g}Y\mathbf{g}^\dagger] = \mathbf{g}\,\mathsf{A}[X,Y]\,\mathbf{g}^\dagger$. The temporal projector Π_{time} is a wordwise metric projection (Section 5.3.11) and transforms by conjugation. The displayed covariance follows. Gauge invariance at the expectation level is then immediate by Corollary 6.125. □

Proposition 6.129 (Vanishing conditions and symmetry)**.** *Assume A is antisymmetric in its entries (commutator-type). Then:*

- *(i) (Commuting/quiescent regimes) If $[\mathbb{D}_\mu\hat{\mathbf{\Phi}}(x), \mathbb{D}_\nu\hat{\mathbf{\Theta}}(x)] = 0$ for all μ,ν, or if $\mathbb{D}_\mu\hat{\mathbf{\Phi}}(x) = 0$ and/or $\mathbb{D}_\nu\hat{\mathbf{\Theta}}(x) = 0$, then $\hat{t}^{(\text{int})}_{\mu\nu}(x) = 0$.*
- *(ii) (Index/field symmetry) The combination in (6.203) is symmetric under the simultaneous swap $(\mu,\Phi) \leftrightarrow (\nu,\Theta)$; after projection by Π_{time}, this yields a symmetric contribution to the temporal sector on the domain where the projection commutes with the swap (which holds for standard metric projections, cf. Section 5.3.11).*
- *(iii) (Dial neutrality of cones) Enabling the interaction channel modifies only the mixed-sector integrand in the action density (Section 6.1.12) and does not alter the expectation unified tensor $\bar{\Sigma}$; hence effective cones from $\bar{g}^{\text{Lor}}$ are unchanged (Definition 5.77, Section 5.4.2, and Corollary 6.87).*

Proof. (i) If either factor in each $\mathsf{A}[\cdot,\cdot]$ commutes with the other, the commutator vanishes; likewise if a factor is zero (quiescent field). The projector preserves 0.

(ii) Using antisymmetry of A and linearity, $\mathsf{A}[\mathbb{D}_\mu\Phi, \mathbb{D}_\nu\Theta] + \mathsf{A}[\mathbb{D}_\mu\Theta, \mathbb{D}_\nu\Phi]$ is invariant under $(\mu,\Phi) \leftrightarrow (\nu,\Theta)$. If Π_{time} is defined via a symmetric form (standard in Section 5.3.11), it commutes with the swap.

(iii) Cited statements follow from Section 6.1.12 and Corollary 6.87. □

[3] E.g. the Tantrakyma word-field in the dynamical sector; we do not re-define it here to avoid duplication.

Remark 6.130 (Triadic structure and sector bookkeeping)**.** The construction of $\hat{t}^{(\mathrm{int})}_{\mu\nu}$ is compatible with the corr/vac/spec triadic decomposition (eq. (5.86)): each $\mathbb{D}_\mu$ and the bilinear A act sectorwise, and Π_{time} preserves the sector split. Thus $\hat{t}^{(\mathrm{int})}_{\mu\nu}$ admits a triadic expansion aligned with (6.205).

Definition 6.131 (Canonical bilinears and Φ-generated chronogeometric tensors)**.** Let $\hat{\mathbf{\Phi}}$ be the Gaiamiti word-field of Section 6.2.1 and Definition 6.119 with wordwise covariant gradient $\mathbb{D}_\mu$ from Definition 6.121 and eq. (6.184). For operator words X, Y define the canonical Hermitian/anti-Hermitian bilinears

$$\boxed{\mathsf{H}[X,Y] := \tfrac{1}{2}\{X^\dagger, Y\}, \qquad \mathsf{A}[X,Y] := \tfrac{i}{2}[X^\dagger, Y]\,.} \tag{6.204}$$

Let Π_{dist} and Π_{time} be the channel projectors selected by the Fourfold modal basis (cf. Definition 4.141; see also the wordwise metric projection machinery in Section 5.3.11). The *operator-valued* chronogeometric tensors induced by $\hat{\mathbf{\Phi}}$ are

$$\boxed{\begin{aligned} \hat{g}_{\mu\nu}(x) &:= \Pi_{\mathrm{dist}}\Big(\, \mathsf{H}\big[\mathbb{D}_\mu\hat{\mathbf{\Phi}}(x),\, \mathbb{D}_\nu\hat{\mathbf{\Phi}}(x)\big] \,\Big)\,, \\ \hat{t}_{\mu\nu}(x) &:= \Pi_{\mathrm{time}}\Big(\, \mathsf{H}\big[\mathbb{D}_\mu\hat{\mathbf{\Phi}}(x),\, \mathbb{D}_\nu\hat{\mathbf{\Phi}}(x)\big] \,\Big)\,, \qquad \hat{\Sigma}_{\mu\nu}(x) := \hat{g}_{\mu\nu}(x) + \hat{t}_{\mu\nu}(x). \end{aligned}} \tag{6.205}$$

Proposition 6.132 (Well-posedness, SOT-Hermiticity, spectral localisation, and covariance)**.** *Assume the regularity and cutoff hypotheses of Definitions 6.119 and 6.121 (SOT–smooth fields, local spectral projector P_x with $P_x\hat{\mathbf{\Phi}}(x)P_x = \hat{\mathbf{\Phi}}(x)$). Then:*

(i) $\hat{g}_{\mu\nu}(x), \hat{t}_{\mu\nu}(x), \hat{\Sigma}_{\mu\nu}(x)$ are SOT-Hermitian and satisfy $P_x\,\hat{g}_{\mu\nu}(x)\,P_x = \hat{g}_{\mu\nu}(x)$ and similarly for $\hat{t}_{\mu\nu}, \hat{\Sigma}_{\mu\nu}$.

(ii) Under endpoint gauges/transport, $\mathbb{D}_\mu\hat{\mathbf{\Phi}} \mapsto \mathbf{g}\,(\mathbb{D}_\mu\hat{\mathbf{\Phi}})\,\mathbf{g}^\dagger$ (Proposition 6.123), and the channel projectors transform covariantly (Section 5.3.11); hence $\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu}, \hat{\Sigma}_{\mu\nu} \mapsto \mathbf{g}\,(\cdot)\,\mathbf{g}^\dagger$.

(iii) The corr/vac/spec triadic split (eq. (5.86)) is preserved by (6.205).

Proof. (i) By construction, $\mathsf{H}[X,Y]$ is Hermitian for any X, Y. Applying a (self-adjoint) channel projector preserves SOT-Hermiticity. The cutoff follows from $P_x\mathbb{D}_\mu\hat{\mathbf{\Phi}}(x)P_x = \mathbb{D}_\mu\hat{\mathbf{\Phi}}(x)$ (induced by $P_x\hat{\mathbf{\Phi}}(x)P_x = \hat{\mathbf{\Phi}}(x)$) and stability under adjoint and sums.

(ii) From Proposition 6.123, $\mathbb{D}_\mu\hat{\mathbf{\Phi}}$ transforms by conjugation. Naturality of H and covariance of the channel projectors (Section 5.3.11) give the same law for $\hat{g}, \hat{t}$, hence for $\hat{\Sigma}$.

(iii) Each ingredient admits a triadic decomposition; H and the projectors act sectorwise, so the split is preserved. □

Corollary 6.133 (Expectation-level fields and gauge invariance)**.** *For any ringed-ket state $|\overset{\circ}{\mathbf{\Psi}}\rangle$,*

$$\bar{g}_{\mu\nu}(x) := \langle\overset{\circ}{\mathbf{\Psi}}|\hat{g}_{\mu\nu}(x)|\overset{\circ}{\mathbf{\Psi}}\rangle, \quad \bar{t}_{\mu\nu}(x) := \langle\overset{\circ}{\mathbf{\Psi}}|\hat{t}_{\mu\nu}(x)|\overset{\circ}{\mathbf{\Psi}}\rangle, \quad \bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$$

are gauge invariant by Corollary 6.125, and are the macroscopic chronogeometric fields used in Chapter 5 (eqs. (5.6) and (5.34) and Definition 5.22). After symmetrization, $\bar{g}^{\mathrm{Lor}}_{\mu\nu} = \frac{1}{2}(\bar{\Sigma}_{\mu\nu} + \bar{\Sigma}_{\nu\mu})$ (Definition 5.77).

Remark 6.134 (Dial neutrality of cones and compatibility with interactions)**.** (i) The dial control modifies mixed-sector integrands but not $\bar{\Sigma}$ (Section 6.1.12 and Corollary 6.87); therefore the effective cones from $\bar{g}^{\mathrm{Lor}}$ are unchanged (Section 5.4.2).

(ii) Commutator-type interaction contributions to the temporal sector may be added as in Definition 6.127 and eq. (6.203); these are covariant by Proposition 6.128 and vanish in commuting/quiescent regimes (Proposition 6.129).

What $\hat{\mathbf{\Phi}}$ is doing. The Gaiamiti field $\hat{\mathbf{\Phi}}(x)$ (Definition 6.119) is a *word-dressed* dynamical object valued simultaneously in the operator vault $\mathcal{B}(\mathcal{H}_{\mathrm{em}})$ and in the vacuum tangent fibre $T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}$ (Definition 6.90). Thus $\hat{\mathbf{\Phi}}$ is not merely an operator; it carries a *direction of deformation* of the vacuum manifold, encoding how the vacuum state $|\Omega(x)\rangle$ is infinitesimally displaced within $\mathcal{M}_{\mathrm{vac}}$.

How we differentiate it (without gauge artefacts). Transport from x' to x uses the modal parallel transporter $\mathcal{U}(x \leftarrow x')$ together with horizontal projection by $\Pi_T(x)$, as formalized in Definition 6.120: $\hat{\mathbf{\Phi}}_{\parallel}(x) = (\mathbf{1}\otimes\Pi_T(x))\big[\mathcal{U}(x\leftarrow x')\,\hat{\mathbf{\Phi}}(x')\,\mathcal{U}(x\leftarrow x')^{\dagger}\big]$. The wordwise covariant gradient is the first–order transported difference $\mathbb{D}_\mu\hat{\mathbf{\Phi}}(x)$ (Definition 6.121 and eq. (6.184)). By Proposition 6.123, both $\hat{\mathbf{\Phi}}_{\parallel}$ and $\mathbb{D}_\mu\hat{\mathbf{\Phi}}$ transform by inner conjugation, hence their Born expectations are gauge invariant (Corollary 6.125).

The two canonical bilinears H, A. The universal pairings on operator words are the symmetric (Hermitian) bilinear $\mathsf{H}[X,Y] = \frac{1}{2}\{X^\dagger, Y\}$ and the antisymmetric (anti-Hermitian) bilinear $\mathsf{A}[X,Y] = \frac{i}{2}[X^\dagger, Y]$ (Definition 6.131 and eq. (6.204)). The former captures amplitude/energy–type pairings of variations, while the latter captures rotational/flow–type effects from noncommutativity.

From bilinears to geometry. Selecting Fourfold channels by the projectors Π_{dist} and Π_{time} (wordwise channel algebra; cf. Definition 4.141), the symmetric pairing of covariant gradients yields the operator–valued chronogeometric tensors defined in (6.205): $\hat{g}_{\mu\nu} = \Pi_{\mathrm{dist}}\big(\mathsf{H}[\mathbb{D}_\mu\hat{\mathbf{\Phi}}, \mathbb{D}_\nu\hat{\mathbf{\Phi}}]\big)$, $\hat{t}_{\mu\nu} = \Pi_{\mathrm{time}}\big(\mathsf{H}[\mathbb{D}_\mu\hat{\mathbf{\Phi}}, \mathbb{D}_\nu\hat{\mathbf{\Phi}}]\big)$, and $\hat{\Sigma}_{\mu\nu} = \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu}$. These are SOT-Hermitian and respect the local spectral cutoff P_x (Proposition 6.132).

Macroscopic fields (what is used operationally). Born expectations in the ringed state produce gauge-invariant macroscopic fields

$$\bar{g}_{\mu\nu}(x) := \Big\langle \mathring{\mathbf{\Psi}}\Big|\hat{g}_{\mu\nu}(x)\Big|\mathring{\mathbf{\Psi}}\Big\rangle, \qquad \bar{t}_{\mu\nu}(x) := \Big\langle \mathring{\mathbf{\Psi}}\Big|\hat{t}_{\mu\nu}(x)\Big|\mathring{\mathbf{\Psi}}\Big\rangle, \qquad \bar{\Sigma}_{\mu\nu}(x) = \bar{g}_{\mu\nu}(x) + \bar{t}_{\mu\nu}(x),$$

by Corollaries 6.125 and 6.133. Symmetrizing the unified field yields the effective Lorentzian metric $\bar{g}^{\mathrm{Lor}}_{\mu\nu}(x) = \frac{1}{2}\big(\bar{\Sigma}_{\mu\nu}(x) + \bar{\Sigma}_{\nu\mu}(x)\big)$ (Definition 5.77).

Optional interactions (when coupling is enabled). With a companion word-field $\hat{\mathbf{\Theta}}$, commutator-type interaction energy enters the temporal sector via the antisymmetric bilinear A, as specified in Definition 6.127 and eq. (6.203). These contributions vanish in commuting or quiescent regimes and remain gauge-covariant (Propositions 6.128 and 6.129).

Operational meaning. Geometry is *emergent from dynamics*: larger, directed changes in $\hat{\mathbf{\Phi}}$ (after transport and horizontal projection) stiffen the spatial/temporal tensors $(\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu})$. The channel projectors apportion symmetric "energy" into distance vs. time sectors. All constructions are gauge/transport–covariant at the operator level and gauge–invariant after expectation, so the scalar fields $(\bar{g}_{\mu\nu}, \bar{t}_{\mu\nu}, \bar{g}^{\mathrm{Lor}}_{\mu\nu})$ admit direct physical interpretation. Finally, the dial control affects only mixed-sector integrands in the action but leaves $\bar{\Sigma}$ unchanged, so causal cones determined by $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ are unaffected (Remark 6.134).

Toy example (commuting constant gradients; $\hat{t}^{(\mathrm{int})} = 0$). Fix a local trivialization and suppose the covariant gradients are x–independent and mutually commuting:

$$\mathbb{D}_\mu \hat{\mathbf{\Phi}}(x) = A_\mu, \qquad \mathbb{D}_\mu \hat{\mathbf{\Theta}}(x) = B_\mu, \qquad [A_\mu, A_\nu] = [B_\mu, B_\nu] = [A_\mu, B_\nu] = 0 \quad \forall \mu, \nu.$$

Then the interaction contribution (6.203) vanishes by antisymmetry of A:

$$\hat{t}^{(\mathrm{int})}_{\mu\nu}(x) = \Pi_{\mathrm{time}}(\mathsf{A}[A_\mu, B_\nu] + \mathsf{A}[B_\mu, A_\nu]) = 0 \quad \text{(cf. Definition 6.127 and Proposition 6.129).}$$

The *operator* chronogeometric tensors from Definition 6.131 and eq. (6.205) reduce to

$$\hat{g}_{\mu\nu}(x) = \Pi_{\mathrm{dist}}\Big(\tfrac{1}{2}\{A^\dagger_\mu, A_\nu\}\Big), \qquad \hat{t}_{\mu\nu}(x) = \Pi_{\mathrm{time}}\Big(\tfrac{1}{2}\{A^\dagger_\mu, A_\nu\}\Big), \qquad \hat{\Sigma}_{\mu\nu}(x) = \tfrac{1}{2}\{A^\dagger_\mu, A_\nu\},$$

so the unified tensor is just the symmetric pairing of the constant gradients, and the distance/-time split is decided entirely by the channel projectors $\Pi_{\mathrm{dist}}, \Pi_{\mathrm{time}}$ (cf. Definition 4.141). Taking Born expectations in the ringed state (Corollaries 6.125 and 6.133) gives gauge-invariant macroscopic fields

$$\bar{g}_{\mu\nu}(x) = \Big\langle \mathring{\mathbf{\Psi}} \Big| \Pi_{\mathrm{dist}}\Big(\tfrac{1}{2}\Big\{(\mathbb{D}_\mu \hat{\mathbf{\Phi}}(x))^\dagger, \mathbb{D}_\nu \hat{\mathbf{\Phi}}(x)\Big\}\Big) \Big| \mathring{\mathbf{\Psi}} \Big\rangle, \tag{6.206}$$

$$\bar{t}_{\mu\nu}(x) = \Big\langle \mathring{\mathbf{\Psi}} \Big| \Pi_{\mathrm{time}}\Big(\tfrac{1}{2}\Big\{(\mathbb{D}_\mu \hat{\mathbf{\Phi}}(x))^\dagger, \mathbb{D}_\nu \hat{\mathbf{\Phi}}(x)\Big\}\Big) \Big| \mathring{\mathbf{\Psi}} \Big\rangle. \tag{6.207}$$

$$\bar{\Sigma}_{\mu\nu}(x) = \Big\langle \mathring{\mathbf{\Psi}} \Big| \tfrac{1}{2}\{\, (\mathbb{D}_\mu \hat{\mathbf{\Phi}}(x))^\dagger, \mathbb{D}_\nu \hat{\mathbf{\Phi}}(x)\,\} \Big| \mathring{\mathbf{\Psi}} \Big\rangle.$$

and the effective Lorentzian metric is $\bar{g}^{\mathrm{Lor}}_{\mu\nu} = \frac{1}{2}(\bar{\Sigma}_{\mu\nu} + \bar{\Sigma}_{\nu\mu})$ (Definition 5.77). If, further, $A_\mu = \alpha_\mu \mathbf{1}_{\mathrm{em}}$ are scalar multiples of the identity within the spectral window (so $P_x A_\mu P_x = A_\mu$, cf. Definition 6.119), then

$$\{A^\dagger_\mu, A_\nu\} = 2\,\alpha^*_\mu \alpha_\nu\, \mathbf{1}_{\mathrm{em}}, \qquad \Rightarrow \quad \bar{\Sigma}_{\mu\nu}(x) = \alpha^*_\mu \alpha_\nu, \;\; \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu} = \alpha^*_\mu \alpha_\nu, \;\; \hat{t}^{(\mathrm{int})}_{\mu\nu} = 0,$$

which exhibits explicitly how geometry (via $\bar{g}_{\mu\nu}, \bar{t}_{\mu\nu}$) emerges from the directional "stiffness" set by the constant gradients, while the commutator-type interaction channel remains quiescent.

6.2.2 The HS Quantum Circuits of the Gaiamiti Field

Definition 6.135 (HS lift of the Gaiamiti word-field)**.** Let $\mathcal{H}_L = \mathrm{span}\{|0\rangle_L, |1\rangle_L\}$ with matrix units P_0, P_1, E_{01}, E_{10}, and let the branch fields $\hat{\mathbf{\Phi}}^{(0)}, \hat{\mathbf{\Phi}}^{(1)}$ be Gaiamiti word-fields as in

Definition 6.119, with a covariantly assembled mixing operator $\hat{\mathbf{\Phi}}_{\mathrm{mix}}$ (from existing sector words, no new symbols). Define the HS–lifted field on $\mathcal{H}_L \otimes \mathcal{H}_{\mathrm{em}}$ by

$$\boxed{\hat{\mathbf{\Phi}}^{\mathrm{HS}} := \; P_0 \otimes \hat{\mathbf{\Phi}}^{(0)} \; + \; P_1 \otimes \hat{\mathbf{\Phi}}^{(1)} \; + \; E_{01} \otimes \hat{\mathbf{\Phi}}_{\mathrm{mix}} \; + \; E_{10} \otimes \hat{\mathbf{\Phi}}^{\dagger}_{\mathrm{mix}}.} \tag{6.208}$$

Definition 6.136 (Reduction by the dial state)**.** Prepare the law register in $|+_\varphi\rangle_L = \frac{1}{\sqrt{2}}\big(|0\rangle_L + e^{i\varphi}|1\rangle_L\big)$. The reduced, phase–programmable field on $\mathcal{H}_{\mathrm{em}}$ is

$$\boxed{\hat{\mathbf{\Phi}}^{\langle \mathrm{HS}\rangle}(\varphi) = \; \tfrac{1}{2}\Big(\hat{\mathbf{\Phi}}^{(0)} + \hat{\mathbf{\Phi}}^{(1)}\Big) \; + \; \mathrm{Re}\Big(e^{-i\varphi}\hat{\mathbf{\Phi}}_{\mathrm{mix}}\Big),} \tag{6.209}$$

obtained by compression in the $|+_\varphi\rangle_L$ sector (cf. the Hamiltonian reduction in Proposition 6.78 and eq. (6.142)).

Proposition 6.137 (Gauge/transport covariance and regularity preservation)**.** *If each branch field $\hat{\mathbf{\Phi}}^{(k)}$ and the mixing operator $\hat{\mathbf{\Phi}}_{\mathrm{mix}}$ transform by conjugation under endpoint gauges and parallel transport (Proposition 6.123), then*

$$\hat{\mathbf{\Phi}}^{\mathrm{HS}} \;\mapsto\; (\mathbf{1}_L \otimes \mathbf{g})\; \hat{\mathbf{\Phi}}^{\mathrm{HS}}\; (\mathbf{1}_L \otimes \mathbf{g})^{\dagger}, \qquad \hat{\mathbf{\Phi}}^{\langle \mathrm{HS}\rangle}(\varphi) \;\mapsto\; \mathbf{g}\; \hat{\mathbf{\Phi}}^{\langle \mathrm{HS}\rangle}(\varphi)\; \mathbf{g}^{\dagger}.$$

Moreover, SOT–C^1 regularity and the local spectral cutoff P_x are preserved by the lift and reduction (cf. Definitions 6.119 and 6.121).

Proof. Conjugation covariance is immediate from linearity in the matrix units and Proposition 6.123; compression by $|+_\varphi\rangle_L$ commutes with conjugation on the environmental slot. Regularity and cutoff follow from boundedness of the matrix units, linearity, and $P_x \hat{\mathbf{\Phi}}^{(k)}(x) P_x = \hat{\mathbf{\Phi}}^{(k)}(x)$ (and similarly for $\hat{\mathbf{\Phi}}_{\mathrm{mix}}$). □

Proposition 6.138 (Dial passes through the covariant gradient)**.** *With $\mathbb{D}_\mu$ as in Definition 6.121, one has the linear reduction rule*

$$\boxed{\mathbb{D}_\mu\, \hat{\mathbf{\Phi}}^{\langle \mathrm{HS}\rangle}(\varphi) = \; \tfrac{1}{2}\Big(\mathbb{D}_\mu \hat{\mathbf{\Phi}}^{(0)} + \mathbb{D}_\mu \hat{\mathbf{\Phi}}^{(1)}\Big) \; + \; \mathrm{Re}\Big(e^{-i\varphi}\, \mathbb{D}_\mu \hat{\mathbf{\Phi}}_{\mathrm{mix}}\Big).} \tag{6.210}$$

Proof. The map $\hat{\mathbf{\Phi}} \mapsto \mathbb{D}_\mu \hat{\mathbf{\Phi}}$ is linear, being a composition of linear transport, linear projection, and a linear difference quotient with an SOT limit (Definitions 6.120 and 6.121). Apply it to (6.209). □

Corollary 6.139 (Insertion into bilinears and chronogeometric tensors)**.** *Inserting (6.210) into the canonical bilinear H and the channel projectors (Definition 6.131 and eq. (6.205)) produces exchange–symmetric tomography dials identical in structure to those obtained for the fourfold trace (Section 6.1.12): the diagonal (mean) contribution is $\frac{1}{4}\,\mathsf{H}[\mathbb{D}_\mu \hat{\mathbf{\Phi}}^{(0)} + \mathbb{D}_\mu \hat{\mathbf{\Phi}}^{(1)}, \mathbb{D}_\nu \hat{\mathbf{\Phi}}^{(0)} + \mathbb{D}_\nu \hat{\mathbf{\Phi}}^{(1)}]$, and the interference terms are controlled by $\mathrm{Re}(e^{-i\varphi}\cdot)$ applied to cross–branch pairings with $\mathbb{D}_\mu \hat{\mathbf{\Phi}}_{\mathrm{mix}}$. Gauge covariance follows from Proposition 6.137, and expectation–level fields remain gauge invariant by Corollaries 6.125 and 6.133.*

Remark 6.140 (Branch recovery, extremal dials, and causal neutrality)**.** (i) Preparing $|0\rangle_L$ (resp. $|1\rangle_L$) yields $\hat{\mathbf{\Phi}}^{\langle \mathrm{HS}\rangle} = \hat{\mathbf{\Phi}}^{(0)}$ (resp. $\hat{\mathbf{\Phi}}^{(1)}$); this mirrors Corollary 6.74.

(ii) For fixed fields, the interference amplitude $M_\mu := \mathbb{D}_\mu \hat{\mathbf{\Phi}}_{\mathrm{mix}}$ implies that $\varphi = \arg\left\langle \mathring{\mathbf{\Psi}} \right| \mathsf{H}[M_\mu, M_\nu] \left| \mathring{\mathbf{\Psi}} \right\rangle$ maximizes the mixed contribution inside the channel projectors, while $\varphi = \arg\left\langle \mathring{\mathbf{\Psi}} \right| \mathsf{H}[M_\mu, M_\nu] \left| \mathring{\mathbf{\Psi}} \right\rangle \pm \pi/2$ suppresses it (cf. Proposition 6.73).

(iii) As with the Hamiltonian dial, cone assignments derived from $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ are unchanged: the dial modifies only mixed–sector integrands and leaves the expectation unified tensor $\bar{\Sigma}$ invariant (cf. Remark 6.134 and Corollary 6.87).

Circuit blueprint (HS wordwise Gaiamiti).

1. **Registers.**

   - *Law qubit* $\mathcal{H}_L$: stores $\{|0\rangle_L, |1\rangle_L\}$ and the dial state $|+_\varphi\rangle_L$ used in Definition 6.136.
   - *Emergent register* $\mathcal{H}_{\mathrm{em}}$: carries the physical state and operators.
   - *Position/path register* $\mathcal{H}_x$ (optional, for x-dependence): indexes points/patches on $\mathcal{X}_q$ to discretize $x \mapsto (\cdot)(x)$.
   - *Phase/energy ancillas*: for spectral projectors Π_0, Π_T and Hadamard/LCU tests.

2. **State preparation (vacuum sector).** Prepare the ringed state and (optionally) the local vacuum:
$$U_{\mathrm{prep}}\,|0\cdots0\rangle = \left|\mathring{\mathbf{\Psi}}\right\rangle, \qquad U_{\mathrm{vac}}\,|0\cdots0\rangle = |\Omega(x)\rangle.$$
Ground-space projectors $\Pi_0(x)$ can be implemented by *phase estimation* on $\widehat{\boldsymbol{H}}$ or on $\hat{H}^{\langle \mathrm{HS}\rangle}(\varphi)$ (Chapter 6 ground sector), post-selecting the lowest energy band; then form the *horizontal projector*
$$\Pi_T(x) \;=\; \Pi_0(x) - |\Omega(x)\rangle\langle\Omega(x)|,$$
using reflections about $|\Omega(x)\rangle$ (Householder) and about $\operatorname{Ran}\Pi_0$ (both standard).

3. **Parallel transport and the covariant gradient.**

   - Implement the transporter along a discrete path $\gamma : x' \to x$ as a product of small-step unitaries,
$$\mathcal{U}(x \leftarrow x';\gamma) \;\approx\; \prod_{e\in\gamma} \exp\Big(- i\,\mathcal{A}_\mu(x_e)\,\Delta x_e^\mu \Big),$$
     i.e. the path-ordered exponential in circuit form (Trotterization).
   - *Transport + horizontal projection* is realized by the unitary block computing
$$\hat{\mathbf{\Phi}}_{\parallel}(x) \;=\; (\mathbf{1}\otimes\Pi_T(x)) \Big[\, \mathcal{U}(x\leftarrow x')\,\hat{\mathbf{\Phi}}(x')\,\mathcal{U}(x\leftarrow x')^\dagger \,\Big] \quad \text{(cf. Definition 6.120).}$$

- The *covariant gradient* $\mathbb{D}_\mu\hat{\mathbf{\Phi}}(x)$ (Definition 6.121) can be realized as a *finite-difference LCU*:

$$\mathbb{D}_\mu\hat{\mathbf{\Phi}}(x) \;\approx\; \frac{1}{\Delta x^\mu}\Big(\hat{\mathbf{\Phi}}(x) - \hat{\mathbf{\Phi}}_{\|}(x)\Big),$$

  implemented via a linear combination of unitaries (LCU) or two Hadamard tests on the pair of circuits for $\hat{\mathbf{\Phi}}(x)$ and $\hat{\mathbf{\Phi}}_{\|}(x)$. Using Proposition 6.124, one may equivalently implement

$$\mathbb{D}_\mu\hat{\mathbf{\Phi}}(x) \;=\; \Big(\mathbf{1}\otimes\Pi_T(x)\Big)\,\nabla^{(\mathcal{U})}_\mu\hat{\mathbf{\Phi}}(x),$$

  which is often simpler experimentally (differentiate first, then project).

4. **Dial-controlled HS lift (field level).** Use the law qubit and $\hat{\mathbf{\Phi}}^{\mathrm{HS}}$ from Definition 6.135. Prepare $|+_\varphi\rangle_L$ and *compress* (measure or uncompute L) to realize the reduced, phase-programmable field

$$\hat{\mathbf{\Phi}}^{\langle\mathrm{HS}\rangle}(\varphi) \;=\; \tfrac{1}{2}\Big(\hat{\mathbf{\Phi}}^{(0)} + \hat{\mathbf{\Phi}}^{(1)}\Big) \;+\; \mathrm{Re}\Big(e^{-i\varphi}\hat{\mathbf{\Phi}}_{\mathrm{mix}}\Big) \qquad \text{(Definition 6.136)}.$$

   The dial φ is a *phase rotation* on the law qubit prior to compression.

5. **Bilinears H,A as circuits.** Given block-encodings (or direct gate implementations) of X and Y:

   - *Symmetric (energy-like)*: estimate $\mathsf{H}[X,Y] = \frac{1}{2}\{X^\dagger, Y\}$ with two Hadamard tests on controlled-$X^\dagger Y$ and controlled-$YX^\dagger$, then average.
   - *Antisymmetric (current-like)*: estimate $\mathsf{A}[X,Y] = \frac{i}{2}[X^\dagger, Y]$ with the difference of those two Hadamard tests (or a single test with a phase flip).

   These plug in with $X = \mathbb{D}_\mu\hat{\mathbf{\Phi}}(x)$ and $Y = \mathbb{D}_\nu\hat{\mathbf{\Phi}}(x)$.

6. **Channel projectors and chronogeometric tensors.** Implement the wordwise channel projectors Π_{dist}, Π_{time} (Fourfold split) as block-encodings or reflections known from the Chapter 4 channel algebra. Then assemble

$$\hat{g}_{\mu\nu} = \Pi_{\mathrm{dist}}\Big(\mathsf{H}[\mathbb{D}_\mu\hat{\mathbf{\Phi}}, \mathbb{D}_\nu\hat{\mathbf{\Phi}}]\Big), \qquad \hat{t}_{\mu\nu} = \Pi_{\mathrm{time}}\Big(\mathsf{H}[\mathbb{D}_\mu\hat{\mathbf{\Phi}}, \mathbb{D}_\nu\hat{\mathbf{\Phi}}]\Big),$$

   exactly as in Definition 6.131. In practice, measure expectations of these operators in the ringed state using standard Hadamard/LCU estimators.

7. **Measuring macroscopic fields (Born expectations).** Using the ringed state $|\mathring{\mathbf{\Psi}}\rangle$ and Hadamard tests, estimate

$$\bar{g}_{\mu\nu}(x) = \Big\langle\mathring{\mathbf{\Psi}}\Big|\hat{g}_{\mu\nu}(x)\Big|\mathring{\mathbf{\Psi}}\Big\rangle, \quad \bar{t}_{\mu\nu}(x) = \Big\langle\mathring{\mathbf{\Psi}}\Big|\hat{t}_{\mu\nu}(x)\Big|\mathring{\mathbf{\Psi}}\Big\rangle,$$

   then form $\bar{\Sigma}_{\mu\nu}$ and $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ exactly as in Corollary 6.133 and Definition 5.77. Gauge covariance at the operator level becomes gauge invariance of these scalars by Corollary 6.125.

8. **Optional interaction channel.** If a companion field $\hat{\mathbf{\Theta}}$ is instantiated, the temporal interaction $\hat{t}^{(\mathrm{int})}_{\mu\nu}$ from Definition 6.127 is an A-bilinear and compiles to the *commutator circuit* (four unitaries via LCU). It vanishes when the gradients commute (toy example), serving as a hardware sanity check.

9. **Two implementation styles.**

   - *(A) Variational/Hadamard-test route*: only controlled versions of word unitaries plus reflections for the projectors are needed. NISQ-friendly; sampling dominates cost.
   - *(B) Block-encoding/QSVT route*: if $\hat{\mathbf{\Phi}}$, $\mathcal{U}$, Π_T are block-encoded, build $\mathbb{D}_\mu$, H, A, and channel projectors via linear-combination-of-unitaries and use QSVT for precision scaling.

10. **Practical caveats (honest but encouraging).**

    - $\Pi_T(x)$ and $\Pi_0(x)$ are not unitary; realize them via reflections, oblivious amplitude amplification, or post-selection (with heralded ancillas).
    - The SOT limit in $\mathbb{D}_\mu$ becomes a small lattice step Δx^μ; control bias by shrinking Δx^μ and variance by increasing shots.
    - Transport $\mathcal{U}$ is a path-ordered product; Trotter error scales with path length and commutator norms of $\mathcal{A}_\mu$.

Quantum circuit realization (HS wordwise Gaiamiti). Using the registers $(\mathcal{H}_L, \mathcal{H}_{\mathrm{em}}, \mathcal{H}_x)$ and ancillas, implement: (i) vacuum and ringed state prep $U_{\mathrm{vac}}, U_{\mathrm{prep}}$; (ii) transport $\mathcal{U}(x \leftarrow x')$ (product of small-step unitaries); (iii) horizontal projection via reflections giving $\Pi_T(x)$; (iv) the LCU finite-difference for $\mathbb{D}_\mu \hat{\mathbf{\Phi}}(x)$; (v) the HS lift and dial compression yielding $\hat{\mathbf{\Phi}}^{\langle \mathrm{HS} \rangle}(\varphi)$; (vi) Hadamard/LCU estimators for H, A and the channel projectors. Estimating the Born expectations in Corollary 6.133 yields $\bar{g}_{\mu\nu}, \bar{t}_{\mu\nu}, \bar{\Sigma}_{\mu\nu}$; symmetrization gives $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ (Definition 5.77). Gauge/transport covariance at the operator level and the invariance of ringed-ket expectations (Proposition 6.123 and Corollary 6.125) ensure the measured scalars are gauge invariant.

Commutator channel (optional). To estimate $\bar{t}^{(\mathrm{int})}_{\mu\nu}(x) = \left\langle \mathring{\mathbf{\Psi}} \right| \Pi_{\mathrm{time}}\, \mathsf{A}[X, Y] \left| \mathring{\mathbf{\Psi}} \right\rangle$ with $X = \mathbb{D}_\mu \hat{\mathbf{\Phi}}$, $Y = \mathbb{D}_\nu \hat{\mathbf{\Theta}}$ (Definition 6.127), repeat Circuit C with a *phase-flip* on the ancilla between the two controlled runs and take the difference, implementing $\frac{i}{2}[X^\dagger, Y]$.

Implementation notes. (i) The projectors $\Pi_0(x), \Pi_T(x), \Pi_{\mathrm{dist}}, \Pi_{\mathrm{time}}$ are realized via reflections/block-encodings (standard in the channel algebra of Chapter 4). (ii) $\mathcal{U}(x \leftarrow x')$ is compiled as a product of small-step unitaries (path ordering); see Circuit B. (iii) All estimators are compatible with the ringed-ket expectations and inherit gauge invariance by Corollary 6.125.

6.3 The Dynamical Tantrakyma Field

The *Tantrakyma Field* is the fundamental operator-valued field governing the excitation, modulation, and internal symmetry structure of the Primordial Quantum Singularities (PQS) across the emergent configuration space $\mathcal{X}_q$. While the Gaiamiti Field encodes infinitesimal deformations of the vacuum geometry (Section 6.2), the Tantrakyma Field encodes all internal, excitation-based structure—such as quantum charges, particle species, and gauge symmetries. It constitutes the matter-like sector of the theory.

Excitation Fiber via Projection Operators

To describe matter-like degrees of freedom, we must first isolate the directions in the local Hilbert space that are orthogonal to the geometric (vacuum deformation) directions. This is achieved formally using projection operators.

Definition 6.141 (Excitation Fiber via Projection Operators)**.** Let $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}$ be the local modal Hilbert space at point $x \in \mathcal{X}_q$, and let $|\Omega(x)\rangle \in \mathcal{M}_{\text{vac}}$ be the local vacuum state. Let $\hat{P}_T(x)$ be the orthogonal projection operator onto the vacuum tangent space $T_{|\Omega(x)\rangle}\mathcal{M}_{\text{vac}}$ (Definition 6.90). The *excitation projection operator* is then defined as its orthogonal complement:

$$\boxed{\hat{P}_{\text{exc}}(x) := \mathbb{I}_x - \hat{P}_T(x),} \tag{6.211}$$

where $\mathbb{I}_x$ is the identity operator on $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}$.

The *excitation fiber* is then defined as the image of this projector:

$$\boxed{\mathcal{E}_{\text{exc}}(x) := \text{Im}\left(\hat{P}_{\text{exc}}(x)\right) = \left(T_{|\Omega(x)\rangle}\mathcal{M}_{\text{vac}}\right)^{\perp}.} \tag{6.212}$$

This subspace contains all local degrees of freedom that are not pure geometric deformations.

The Tantrakyma Field Definition

Definition 6.142 (The Tantrakyma Field as an Operator-Valued Internal Mode Field)**.** The *Tantrakyma Field*, denoted $\hat{\mathbf{\Theta}}(x)$, is an operator-valued internal-mode field over $\mathcal{X}_q$, defined as a smooth section of the excitation-valued operator bundle:

$$\boxed{\hat{\mathbf{\Theta}} : \mathcal{X}_q \longrightarrow \text{End}(\mathcal{H}_{\text{em}}) \otimes \mathcal{E}_{\text{exc}},} \tag{6.213}$$

with local evaluations satisfying:

$$\boxed{\hat{\mathbf{\Theta}}(x) \in \text{End}(\mathcal{H}_{\text{em}}) \otimes \mathcal{E}_{\text{exc}}(x),} \tag{6.214}$$

where:

- $\text{End}(\mathcal{H}_{\text{em}})$ is the algebra of bounded operators on the emergent Hilbert space (Section 6.1.27);

- $\mathcal{E}_{\text{exc}}(x)$ is the excitation fiber defined at each x by eq. (6.212).

Operationally, $\hat{\mathbf{\Theta}}(x)$ acts locally on the internal modal sector $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}$ and is extended globally across $\mathcal{H}_{\text{em}}$ via the smooth structure of the emergent Hilbert bundle (Definition 4.287).

Interpretation While the Gaiamiti Field generates the geometric stage, the Tantrakyma Field, $\hat{\mathbf{\Theta}}(x)$, directs the actors upon it. At each point x, it governs the dynamics of matter-like structure by modulating internal degrees of freedom within $\mathcal{E}_{\text{exc}}(x)$. These excitations correspond to the relational substrate of quantum charges, spin, and internal symmetries.

Changes generated by $\hat{\mathbf{\Theta}}(x)$ correspond to local transitions in excitation configuration—rotations, modulations, or internal transitions. The algebraic structure of $\hat{\mathbf{\Theta}}(x)$, including its commutation relations and representation under internal symmetry groups, induces the observable particle spectrum and its local gauge symmetries.

In analogy, if $\hat{\mathbf{\Phi}}(x)$ provides the resonant curvature of the vacuum structure, then $\hat{\mathbf{\Theta}}(x)$ is the orchestra's performance upon that stage—the physical sound of matter as relationally defined excitation.

6.3.1 Holosymmetric Wordwise Dynamical Tantrakyma Field

Setting and excitation fibre. Let $x \mapsto |\Omega(x)\rangle \in \mathcal{M}_{\text{vac}}$ be a smooth local vacuum section (Definitions 6.61 and 6.90) and let $\hat{P}_T(x)$ denote the orthogonal projector onto the vacuum–tangent fibre $T_{|\Omega(x)\rangle}\mathcal{M}_{\text{vac}}$ (cf. Definition 6.90). Define the *excitation projector* and *excitation fibre* by

$$\boxed{\hat{P}_{\text{exc}}(x) := \mathbf{1}_x - \hat{P}_T(x), \qquad \mathcal{E}_{\text{exc}}(x) := \operatorname{Im} \hat{P}_{\text{exc}}(x) = \left(T_{|\Omega(x)\rangle}\mathcal{M}_{\text{vac}}\right)^{\perp}.} \tag{6.215}$$

Definition 6.143 (Tantrakyma word-field; excitation-valued operator word)**.** Let $\mathbb{W}_{\text{op}}(\mathcal{H}_{\text{em}})$ denote the operator–word vault (cf. Definition 6.114 and Section 6.1.27), defined by

$$\mathbb{W}_{\text{op}}(\mathcal{H}_{\text{em}}) \;:=\; \mathcal{B}(\mathcal{H}_{\text{em}}). \tag{6.216}$$

The *Tantrakyma field* is the SOT–smooth section

$$\hat{\mathbf{\Theta}} : \; \mathcal{X}_q \longrightarrow \mathbb{W}_{\text{op}}(\mathcal{H}_{\text{em}}) \;\otimes\; \mathcal{E}_{\text{exc}}, \qquad \hat{\mathbf{\Theta}}(x) \;\in\; \mathcal{B}(\mathcal{H}_{\text{em}}) \;\otimes\; \mathcal{E}_{\text{exc}}(x), \tag{6.217}$$

spectrally localised by a cutoff P_x (so $P_x\hat{\mathbf{\Theta}}(x)P_x = \hat{\mathbf{\Theta}}(x)$). By construction, its tangent slot is orthogonal to $T_{|\Omega(x)\rangle}\mathcal{M}_{\text{vac}}$, i.e. $(\mathbf{1} \otimes \hat{P}_T(x))\,\hat{\mathbf{\Theta}}(x) = 0$.

Gauge/transport covariance and covariant gradient.

Definition 6.144 (Transport with excitation projection)**.** Let $\mathcal{U}(x \leftarrow x'; \gamma)$ be the parallel transporter on the modal bundle (Section 4.5; cf. eqs. (4.11) and (4.14)), and let $\hat{P}_{\text{exc}}(x) = \mathbf{1}_x - \hat{P}_T(x)$ be the excitation projector (eq. (6.215)). For a Tantrakyma field $\hat{\mathbf{\Theta}}$ as in Definition 6.143, define the transported field at x with excitation projection by

$$\hat{\mathbf{\Theta}}_{\parallel}(x) \;:=\; \Big(\mathbf{1} \otimes \hat{P}_{\text{exc}}(x)\Big)\Big[\,\mathcal{U}(x \leftarrow x')\;\hat{\mathbf{\Theta}}(x')\;\mathcal{U}(x \leftarrow x')^{\dagger}\,\Big]. \tag{6.218}$$

To first order in $|x - x'|$ the definition is path-independent; curvature effects appear only at $\mathcal{O}(|x - x'|^2)$ (cf. Section 4.5).

Definition 6.145 (Wordwise covariant gradient of the Tantrakyma field)**.** Assume SOT–C^1 regularity of $x \mapsto \hat{\boldsymbol{\Theta}}(x)$ and $x \mapsto \hat{P}_{\mathrm{exc}}(x)$, and a C^1 endpoint expansion for $\mathcal{U}$ (Definition 6.7 and Section 6.1.4). The *wordwise covariant gradient* of $\hat{\boldsymbol{\Theta}}$ is the SOT limit

$$\boxed{\mathbb{D}_\mu \hat{\boldsymbol{\Theta}}(x) \;:=\; \lim_{x' \to x} \frac{\hat{\boldsymbol{\Theta}}(x) - \hat{\boldsymbol{\Theta}}_{\parallel}(x)}{\Delta x^\mu}\,.} \tag{6.219}$$

For comparison with the operator-slot derivative, we *reuse* Definition 6.122: $\nabla_\mu^{(\mathcal{U})} \hat{\boldsymbol{\Theta}}(x) = \lim_{x' \to x} \frac{\hat{\boldsymbol{\Theta}}(x) - \mathcal{U}\, \hat{\boldsymbol{\Theta}}(x')\, \mathcal{U}^\dagger}{\Delta x^\mu}$.

Proposition 6.146 (Gauge/transport covariance)**.** *Under a local endpoint gauge $g(x)$ (fibrewise unitary),*

$$\begin{aligned} \hat{\boldsymbol{\Theta}}(x) &\mapsto \mathbf{g}(x)\, \hat{\boldsymbol{\Theta}}(x)\, \mathbf{g}(x)^\dagger, \\ \mathcal{U}(x \leftarrow x') &\mapsto g(x)\, \mathcal{U}(x \leftarrow x')\, g(x')^\dagger, \\ \hat{P}_{\mathrm{exc}}(x) &\mapsto g(x)\, \hat{P}_{\mathrm{exc}}(x)\, g(x)^\dagger. \end{aligned}$$

(Proposition 6.66 and Section 4.5). Then

$$\hat{\boldsymbol{\Theta}}_{\parallel}(x) \;\mapsto\; \mathbf{g}(x)\, \hat{\boldsymbol{\Theta}}_{\parallel}(x)\, \mathbf{g}(x)^\dagger, \qquad \mathbb{D}_\mu \hat{\boldsymbol{\Theta}}(x) \;\mapsto\; \mathbf{g}(x)\, \mathbb{D}_\mu \hat{\boldsymbol{\Theta}}(x)\, \mathbf{g}(x)^\dagger.$$

Proof. Insert the gauge transforms into (6.218) and cancel $g(x')^\dagger g(x') = \mathbf{1}$ to obtain $\hat{\boldsymbol{\Theta}}_{\parallel}(x) \mapsto \mathbf{g}(x) \hat{\boldsymbol{\Theta}}_{\parallel}(x) \mathbf{g}(x)^\dagger$. For (6.219), the bounded inner automorphism by $\mathbf{g}(x)$ commutes with the SOT limit, yielding the covariant transformation of $\mathbb{D}_\mu \hat{\boldsymbol{\Theta}}(x)$, exactly as in Proposition 6.123. □

Proposition 6.147 (Excitation–horizontal projection of the operator-slot derivative)**.** *Under the regularity assumptions of Definition 6.145, one has*

$$\boxed{\mathbb{D}_\mu \hat{\boldsymbol{\Theta}}(x) = \; \Big(\mathbf{1} \otimes \hat{P}_{\mathrm{exc}}(x)\Big)\; \nabla_\mu^{(\mathcal{U})} \hat{\boldsymbol{\Theta}}(x)\,.} \tag{6.220}$$

Proof. Since $(\mathbf{1} \otimes \hat{P}_{\mathrm{exc}}(x))\, \hat{\boldsymbol{\Theta}}(x) = \hat{\boldsymbol{\Theta}}(x)$ by Definition 6.143, we may write

$$\hat{\boldsymbol{\Theta}}(x) - \hat{\boldsymbol{\Theta}}_{\parallel}(x) = \Big(\mathbf{1} \otimes \hat{P}_{\mathrm{exc}}(x)\Big) \Big[\hat{\boldsymbol{\Theta}}(x) - \mathcal{U}(x \leftarrow x')\, \hat{\boldsymbol{\Theta}}(x')\, \mathcal{U}(x \leftarrow x')^\dagger\Big].$$

Divide by Δx^μ and take the SOT limit $x' \to x$; boundedness and x'–independence of $(\mathbf{1} \otimes \hat{P}_{\mathrm{exc}}(x))$ allow it to pass through the limit, giving (6.220). The existence of the limit follows from the C^1 endpoint expansion of $\mathcal{U}$ and SOT–C^1 of the fields, as in Proposition 6.124. □

Corollary 6.148 (Expectation-level gauge invariance (Tantrakyma))**.** *For any ringed-ket state $|\mathring{\boldsymbol{\Psi}}\rangle$, the scalar densities $\langle \mathring{\boldsymbol{\Psi}} | \hat{\boldsymbol{\Theta}}_{\parallel}(x) | \mathring{\boldsymbol{\Psi}} \rangle$ and $\langle \mathring{\boldsymbol{\Psi}} | \mathbb{D}_\mu \hat{\boldsymbol{\Theta}}(x) | \mathring{\boldsymbol{\Psi}} \rangle$ are gauge invariant, by Proposition 6.146 and the invariance of ringed-ket expectations (Remark 6.35; cf. Corollary 6.125).*

Remark 6.149 (Well-posedness, spectral localisation, and triadic compatibility)**.** (i) The SOT limit in (6.219) exists under the same hypotheses as for the Gaiamiti field (Definition 6.121); locality is enforced by the spectral cutoff P_x from Definition 6.143.
(ii) The corr/vac/spec triadic split is preserved by transport, projection $\hat{P}_{\mathrm{exc}}$, and the operator-slot derivative, hence by $\mathbb{D}_\mu$ (cf. eq. (5.86)).
(iii) As for $\hat{\mathbf{\Phi}}$, dial operations that act only on mixed-sector integrands leave the expectation unified tensor $\bar{\Sigma}$ and cone assignments unchanged (cf. Remark 6.134 and Corollary 6.87).

Couplings to geometry.

Definition 6.150 (Tantrakyma–Gaiamiti interaction contribution (adopted))**.** We *adopt* the interaction definition of Definition 6.127 verbatim, with $\hat{\mathbf{\Phi}}$ the Gaiamiti field of Section 6.2.1 and $\hat{\mathbf{\Theta}}$ the Tantrakyma field of Definition 6.143. In particular, the commutator-type contribution to the temporal sector is the operator-valued tensor specified in (6.203), to be added to the temporal channel of (6.205) when interaction channels are enabled. No new symbols are introduced here.

Proposition 6.151 (Covariance, well-posedness, and vanishing regimes)**.** *The interaction tensor of Definition 6.150 satisfies:*

(i) Gauge/transport covariance
It transforms by conjugation under endpoint gauges and parallel transport, as established in Proposition 6.128, using the covariant gradients $\mathbb{D}_\mu\hat{\mathbf{\Phi}}$ (Definition 6.121) and $\mathbb{D}_\mu\hat{\mathbf{\Theta}}$ (Definition 6.145).

(ii) Well-posedness and spectral localisation
SOT–C^1 regularity and the local spectral cutoff P_x for both fields imply the same properties for $\hat{t}^{(\mathrm{int})}_{\mu\nu}$ (cf. Definitions 6.119 and 6.143 and Proposition 6.132).

(iii) Vanishing conditions
In commuting or quiescent regimes the interaction term vanishes, as in Proposition 6.129.

Corollary 6.152 (Expectation-level gauge invariance)**.** *For any ringed-ket state $|\mathring{\mathbf{\Psi}}\rangle$,*

$$\bar{t}^{(\mathrm{int})}_{\mu\nu}(x) := \left\langle \mathring{\mathbf{\Psi}} \middle| \hat{t}^{(\mathrm{int})}_{\mu\nu}(x) \middle| \mathring{\mathbf{\Psi}} \right\rangle$$

is gauge invariant by Corollary 6.125 together with Propositions 6.128 and 6.146.

Remark 6.153 (Triadic compatibility and causal neutrality)**.** (i) The corr/vac/spec triadic split is preserved by the interaction construction (sectorwise action of $\mathbb{D}_\mu$, A, and Π_{time}), hence $\hat{t}^{(\mathrm{int})}_{\mu\nu}$ admits a triadic expansion aligned with (6.205). (ii) As with the Gaiamiti-only case, enabling the interaction channel affects only mixed-sector integrands and leaves the expectation unified tensor $\bar{\Sigma}$ (hence the cone assignment from $\bar{g}^{\mathrm{Lor}}$) unchanged (cf. Remark 6.134 and Corollary 6.87).

Definition 6.154 (Canonical bilinears reused; Tantrakyma-generated tensors)**.** Reuse the canonical bilinears H, A from Definition 6.131 and eq. (6.204) and the channel projectors $\Pi_{\mathrm{dist}}, \Pi_{\mathrm{time}}$

(Fourfold split). With the covariant gradient $\mathbb{D}_\mu\hat{\mathbf{\Theta}}$ from Definition 6.145 and eq. (6.219), define the *Tantrakyma-only* operator-valued chronogeometric tensors by

$$\boxed{\begin{aligned}\hat{g}^{[\Theta]}_{\mu\nu}(x) &:= \Pi_{\mathrm{dist}}\Big(\mathsf{H}\Big[\mathbb{D}_\mu\hat{\mathbf{\Theta}}(x),\,\mathbb{D}_\nu\hat{\mathbf{\Theta}}(x)\Big]\Big)\\ \hat{t}^{[\Theta]}_{\mu\nu}(x) &:= \Pi_{\mathrm{time}}\Big(\mathsf{H}\Big[\mathbb{D}_\mu\hat{\mathbf{\Theta}}(x),\,\mathbb{D}_\nu\hat{\mathbf{\Theta}}(x)\Big]\Big)\\ \hat{\Sigma}^{[\Theta]}_{\mu\nu}(x) &:= \hat{g}^{[\Theta]}_{\mu\nu}(x)+\hat{t}^{[\Theta]}_{\mu\nu}(x)\end{aligned}} \tag{6.221}$$

Proposition 6.155 (Well-posedness, SOT-Hermiticity, covariance, and localisation)**.** *Under the hypotheses of Definitions 6.143 and 6.145 (SOT–C^1 regularity, spectral cutoff P_x), the tensors in (6.221) are SOT-Hermitian, satisfy $P_x(\cdot)P_x=(\cdot)$, and transform by conjugation under endpoint gauges and transport (Proposition 6.146). The corr/vac/spec triadic split (eq. (5.86)) is preserved.*

Corollary 6.156 (Expectation-level fields and gauge invariance (Tantrakyma sector))**.** *For any ringed-ket state $|\mathring{\mathbf{\Psi}}\rangle$,*

$$\bar{g}^{[\Theta]}_{\mu\nu}(x) := \Big\langle\mathring{\mathbf{\Psi}}\Big|\,\hat{g}^{[\Theta]}_{\mu\nu}(x)\,\Big|\mathring{\mathbf{\Psi}}\Big\rangle,\qquad \bar{t}^{[\Theta]}_{\mu\nu}(x) := \Big\langle\mathring{\mathbf{\Psi}}\Big|\,\hat{t}^{[\Theta]}_{\mu\nu}(x)\,\Big|\mathring{\mathbf{\Psi}}\Big\rangle,\qquad \bar{\Sigma}^{[\Theta]}_{\mu\nu}(x)=\bar{g}^{[\Theta]}_{\mu\nu}(x)+\bar{t}^{[\Theta]}_{\mu\nu}(x)$$

are gauge invariant by Corollary 6.148.

Remark 6.157 (Total two-field bookkeeping and optional cross-symmetric terms)**.** (i) With both fields present, a convenient *total* operator tensor is

$$\hat{g}^{\mathrm{tot}}_{\mu\nu} := \hat{g}^{[\Phi]}_{\mu\nu}+\hat{g}^{[\Theta]}_{\mu\nu},\qquad \hat{t}^{\mathrm{tot}}_{\mu\nu} := \hat{t}^{[\Phi]}_{\mu\nu}+\hat{t}^{[\Theta]}_{\mu\nu}+\hat{t}^{(\mathrm{int})}_{\mu\nu},\qquad \hat{\Sigma}^{\mathrm{tot}}_{\mu\nu} := \hat{g}^{\mathrm{tot}}_{\mu\nu}+\hat{t}^{\mathrm{tot}}_{\mu\nu},$$

using eqs. (6.203), (6.205) and (6.221). (ii) If desired, *cross-symmetric* energy pairings may be introduced as $\Pi_{\mathrm{dist/time}}\mathsf{H}[\mathbb{D}_\mu\hat{\mathbf{\Phi}},\mathbb{D}_\nu\hat{\mathbf{\Theta}}]+(\mu,\Phi)\leftrightarrow(\nu,\Theta)$; we omit these by default to avoid overlap with the commutator interaction channel (Definition 6.150).

Optional HS law-control dial (no new symbols).

Definition 6.158 (HS lift of the Tantrakyma word-field)**.** Let $\mathcal{H}_L=\mathrm{span}\{|0\rangle_L,|1\rangle_L\}$ with matrix units P_0,P_1,E_{01},E_{10}, and let the branch fields $\hat{\mathbf{\Theta}}^{(0)},\hat{\mathbf{\Theta}}^{(1)}$ be Tantrakyma word-fields as in Definition 6.143. For a covariantly assembled mixing operator $\hat{\mathbf{\Theta}}_{\mathrm{mix}}$ (from existing sector words; no new symbols), define the HS–lifted field on $\mathcal{H}_L\otimes\mathcal{H}_{\mathrm{em}}$ by

$$\boxed{\widehat{\mathbf{\Theta}}^{\mathrm{HS}} := P_0\otimes\hat{\mathbf{\Theta}}^{(0)}+P_1\otimes\hat{\mathbf{\Theta}}^{(1)}+E_{01}\otimes\hat{\mathbf{\Theta}}_{\mathrm{mix}}+E_{10}\otimes\hat{\mathbf{\Theta}}^{\dagger}_{\mathrm{mix}}.} \tag{6.222}$$

This mirrors eq. (6.208) for the Gaiamiti field.

Definition 6.159 (Reduction by the dial state)**.** Preparing the law register in $|+_\varphi\rangle_L=\frac{1}{\sqrt{2}}\big(|0\rangle_L+e^{i\varphi}|1\rangle_L\big)$ and compressing to $\mathcal{H}_{\mathrm{em}}$ yields the reduced, phase–programmable field

$$\boxed{\hat{\mathbf{\Theta}}^{\langle\mathrm{HS}\rangle}(\varphi)=\tfrac{1}{2}\Big(\hat{\mathbf{\Theta}}^{(0)}+\hat{\mathbf{\Theta}}^{(1)}\Big)+\mathrm{Re}\Big(e^{-i\varphi}\hat{\mathbf{\Theta}}_{\mathrm{mix}}\Big),} \tag{6.223}$$

exactly mirroring eq. (6.209).

Proposition 6.160 (Gauge/transport covariance and regularity preservation)**.** *If each branch field* $\hat{\mathbf{\Theta}}^{(k)}$ *and* $\hat{\mathbf{\Theta}}_{\mathrm{mix}}$ *transform by conjugation under endpoint gauges and parallel transport (Proposition 6.146), then*

$$\widehat{\mathbf{\Theta}}^{\mathrm{HS}} \;\mapsto\; (\mathbf{1}_L \otimes \mathbf{g})\, \widehat{\mathbf{\Theta}}^{\mathrm{HS}}\, (\mathbf{1}_L \otimes \mathbf{g})^\dagger, \qquad \hat{\mathbf{\Theta}}^{\langle \mathrm{HS}\rangle}(\varphi) \;\mapsto\; \mathbf{g}\, \hat{\mathbf{\Theta}}^{\langle \mathrm{HS}\rangle}(\varphi)\, \mathbf{g}^\dagger.$$

Moreover, SOT–C^1 regularity and the local spectral cutoff P_x are preserved by the lift and reduction (cf. Definitions 6.143 and 6.145).

Proposition 6.161 (Dial passes through the covariant gradient (Tantrakyma))**.** *With $\mathbb{D}_\mu$ as in Definition 6.145, one has the linear reduction rule*

$$\boxed{\mathbb{D}_\mu\, \hat{\mathbf{\Theta}}^{\langle \mathrm{HS}\rangle}(\varphi) = \tfrac{1}{2}\Big(\mathbb{D}_\mu \hat{\mathbf{\Theta}}^{(0)} + \mathbb{D}_\mu \hat{\mathbf{\Theta}}^{(1)}\Big) \;+\; \mathrm{Re}\Big(e^{-i\varphi}\, \mathbb{D}_\mu \hat{\mathbf{\Theta}}_{\mathrm{mix}}\Big).} \tag{6.224}$$

Proof. $\mathbb{D}_\mu$ is linear (transport + projection + SOT difference quotient). Apply it to eq. (6.223), exactly as in the Gaiamiti case (Proposition 6.138). □

Corollary 6.162 (Insertion into covariant gradient and interaction channel)**.** *Inserting eq. (6.224) into the Tantrakyma covariant gradient (6.219) and into the interaction tensor (6.203) realizes the same exchange–symmetric/antisymmetric tomography dials developed for the fourfold trace (Section 6.1.12), without altering cone assignments (cf. Remark 6.134).*

Remark 6.163 (Branch recovery and causal neutrality)**.** (i) Preparing $|0\rangle_L$ (resp. $|1\rangle_L$) yields $\hat{\mathbf{\Theta}}^{\langle \mathrm{HS}\rangle} = \hat{\mathbf{\Theta}}^{(0)}$ (resp. $\hat{\mathbf{\Theta}}^{(1)}$), mirroring the Gaiamiti branch recovery.
(ii) As with the Gaiamiti dial, the Tantrakyma dial modifies only mixed-sector integrands and leaves the expectation unified tensor $\bar{\Sigma}$ (hence cone assignments from $\bar{g}^{\mathrm{Lor}}$) unchanged (cf. Corollary 6.87).

Remark 6.164 (Orthogonality, covariance, and consistency with Chapter 6)**.** (i) By (6.215), $\hat{\mathbf{\Theta}}$ is orthogonal (in the tangent slot) to geometric deformations, so the Gaiamiti/Tantrakyma split cleanly separates geometry from internal excitations. (ii) Transport and frame covariance follow from (6.218)–(6.219). (iii) No new symbols beyond the established words are introduced; all constructions live in the operator-word vault $\mathcal{B}(\mathcal{H}_{\mathrm{em}})$ and its excitation/tangent bundles, in line with Sections 6.2 and 6.3.

6.4 Bundle-Theoretic Formulation of the Foundational Fields

The dynamical content of the IQR framework resides in two foundational operator-valued fields: the Gaiamiti field, which generates the geometry of the emergent manifold, and the Tantrakyma field, which governs its internal, matter-like excitations. Both admit a rigorous global description as smooth sections of fiber bundles constructed over the emergent configuration space $\mathcal{X}_q$. At each $x \in \mathcal{X}_q$, we identify the endomorphism fiber with the local modal space,

$$\mathrm{End}(\mathcal{H}_{\mathrm{em}})_x \;\cong\; \mathrm{End}\Big(\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}\Big) \quad \text{(cf. Definition 4.287)},$$

and endow the vacuum manifold's tangent bundle with the inner product induced from the ambient/projective geometry (cf. Definition 6.90), allowing $T\mathcal{M}_{\mathrm{vac}} \simeq T^*\mathcal{M}_{\mathrm{vac}}$ via the induced metric.

The Gaiamiti Field: Geometric Structure

The Dynamical Gaiamiti Field, $\hat{\mathbf{\Phi}}(x)$, is an *operator-valued vector* in the local tangent space to the vacuum manifold, i.e. a smooth section of

$$\mathrm{End}(\mathcal{H}_{\mathrm{em}}) \;\otimes\; T\mathcal{M}_{\mathrm{vac}} \quad \text{(cf. Definition 6.90)}.$$

Let $\{|e_i(x)\rangle\}_{i=1}^{r}$ be a local orthonormal frame for $T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}$, with dual frame $\{\langle e^i(x)|\}$ obtained from the induced metric, where $r = \dim T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}$. The Gaiamiti field decomposes as

$$\boxed{\hat{\mathbf{\Phi}}(x) \;=\; \sum_{i=1}^{r} \hat{\Phi}^i(x)\,|e_i(x)\rangle,} \tag{6.225}$$

where the operator-valued coefficients are defined by metric contraction with the dual frame,

$$\hat{\Phi}^i(x) \;:=\; (\mathbf{1}\otimes\langle e^i(x)|)\,\hat{\mathbf{\Phi}}(x) \;\equiv\; \langle e_i(x)|\,\hat{\mathbf{\Phi}}(x)\rangle \;\in\; \mathrm{End}(\mathcal{H}_{\mathrm{em}}).$$

Locally, $\hat{\mathbf{\Phi}}(x)$ is thus represented as a column vector of operator entries:

$$\hat{\mathbf{\Phi}}(x) \;\longleftrightarrow\; \begin{pmatrix} \hat{\Phi}^1(x) \\ \hat{\Phi}^2(x) \\ \vdots \\ \hat{\Phi}^r(x) \end{pmatrix}.$$

Under a change of tangent frame $|e_i\rangle \mapsto R_i{}^j|e_j\rangle$, the coefficients transform contravariantly, $\hat{\Phi}^i \mapsto (R^{-1})^i{}_j\,\hat{\Phi}^j$, while the endomorphism slot transforms by inner automorphism under local gauge $g(x)$: $X \mapsto g(x)\,X\,g(x)^\dagger$ (cf. the standard covariance laws mirrored in the wordwise setting).

The Tantrakyma Field: Matter and Excitation Structure

The Dynamical Tantrakyma Field, $\hat{\mathbf{\Theta}}(x)$, is a smooth section of

$$\mathrm{End}(\mathcal{H}_{\mathrm{em}}) \;\otimes\; \mathcal{E}_{\mathrm{exc}}(x),$$

where $\mathcal{E}_{\mathrm{exc}}(x)$ is the excitation fiber orthogonal to the vacuum-tangent fiber (cf. the projector construction leading to $\mathcal{E}_{\mathrm{exc}}$). For local displays below we *suppress the excitation slot* by fixing a local excitation frame or, equivalently, by projecting onto $\mathcal{E}_{\mathrm{exc}}(x)$ and reporting the $\mathrm{End}(\mathcal{H}_{\mathrm{em}})$ components.

The four-dimensional modal space reflecting the Fourfold Dialectic Algebra admits an orthonormal basis of pure modes (Definition 4.273):

$$\mathcal{B}_x := \Big\{\, |\mathbf{0}\rangle_x,\;\; |\mathbf{1}\rangle_x,\;\; |\mathbf{0}+\mathbf{1}\rangle_x,\;\; |\mathring{\mathbf{0}}\mathring{\mathbf{\Lambda}}\mathring{\mathbf{1}}\rangle_x \,\Big\},$$

and we index components by $\alpha, \beta \in \{0, 1, p, s\}$ with $p \leftrightarrow |\mathbf{0}+\mathbf{1}\rangle_x$ and $s \leftrightarrow |\mathring{\mathbf{0}}\mathring{\mathbf{\Lambda}}\mathring{\mathbf{1}}\rangle_x$. Relative to $\mathcal{B}_x$, the Tantrakyma field has the local operator-matrix decomposition

$$\boxed{\hat{\mathbf{\Theta}}(x) \;=\; \sum_{\alpha,\beta\in\{0,1,p,s\}} \theta^{\alpha\beta}(x)\,|\alpha\rangle_x\langle\beta|_x,} \tag{6.226}$$

with the explicit matrix of entries $\theta^{\alpha\beta}(x) := \langle \alpha_x | \hat{\mathbf{\Theta}}(x) | \beta_x \rangle \in \mathbb{C}$ understood after the excitation projection:

$$\hat{\mathbf{\Theta}}(x) \;\longleftrightarrow\; \begin{pmatrix} \theta^{00}(x) & \theta^{01}(x) & \theta^{0p}(x) & \theta^{0s}(x) \\ \theta^{10}(x) & \theta^{11}(x) & \theta^{1p}(x) & \theta^{1s}(x) \\ \theta^{p0}(x) & \theta^{p1}(x) & \theta^{pp}(x) & \theta^{ps}(x) \\ \theta^{s0}(x) & \theta^{s1}(x) & \theta^{sp}(x) & \theta^{ss}(x) \end{pmatrix}.$$

Under a change of modal frame $|\alpha\rangle_x \mapsto U(x)|\alpha\rangle_x$ the matrix transforms as $\theta(x) \mapsto U(x)\,\theta(x)\,U(x)^\dagger$; under a gauge on the endomorphism slot, $X \mapsto g(x)\,X\,g(x)^\dagger$. The excitation-slot choice transforms with the excitation frame in the natural way; local projections to $\mathcal{E}_{\text{exc}}(x)$ commute with these inner automorphisms.

Remarks on derivatives and observables (standard setting). All covariant derivatives in the standard formalism use the product connection: the Levi–Civita–type connection on $T\mathcal{M}_{\text{vac}}$ (or the projected connection on $\mathcal{E}_{\text{exc}}$) together with the gauge connection on $\text{End}(\mathcal{H}_{\text{em}})$; geometric and temporal observables arise from symmetric bilinears of covariant gradients combined with the channel projectors, so no componentwise Hermiticity of $\hat{\mathbf{\Phi}}$ or $\hat{\mathbf{\Theta}}$ is required.

Interpretation of Field Components The physical meaning of the theory is encoded in these component decompositions (cf. Definition 6.167):

- **Gaiamiti components $\left(\hat{\Phi}^i\right)$:** The frame $\{|e_i(x)\rangle\}_{i=1}^r$ spans the orthogonal *modes of geometric deformation* (e.g. expansion, shear) of the vacuum manifold, and the operator-valued coefficients $\hat{\Phi}^i(x)$ are the quantum amplitudes along those directions (dual-frame contraction as in eq. (6.229)).
- **Tantrakyma components $\left(\theta^{\alpha\beta}\right)$:** Relative to the modal basis $\{|\alpha\rangle_x\}_{\alpha\in\{0,1,p,s\}}$, the matrix elements govern internal/matter-like dynamics: *diagonals* $\theta^{\alpha\alpha}(x)$ encode self-modulation within a dialectical mode, while *off-diagonals* $\theta^{\alpha\beta}(x)$ $(\alpha \neq \beta)$ generate inter-modal mixing, the "woven-wave" couplings of internal states (cf. eq. (6.230) and the excitation confinement via eq. (6.227)).

In summary, the Gaiamiti components describe the dynamics of *geometric modes*, whereas the Tantrakyma components describe the dynamics of *internal, matter-like modes*. Their *covariant gradients*, paired through the canonical bilinears, feed into the chronogeometric tensors as in eq. (6.239).

The Dual Physical Roles of the Foundational Fields The Gaiamiti and Tantrakyma fields are the two pillars of emergent physical reality, governing the distinct yet inseparable domains of geometry and matter.

- **The Gaiamiti field $\left(\hat{\mathbf{\Phi}}\right)$ as the architect of spacetime:** The Gaiamiti field is the *primary* source of **geometric observables**: the baseline tensors $\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu}$ arise from the symmetric pairing $\mathsf{H}[D_\mu\hat{\mathbf{\Phi}}, D_\nu\hat{\mathbf{\Phi}}]$ and channel projection (cf. eq. (6.239)). Its vacuum expectation determines the background chronogeometry $\bar{\Sigma}_{\mu\nu}$ (with effective Lorentzian

reduction via Definition 5.77), while quantum fluctuations $\delta\hat{\mathbf{\Phi}}$ manifest as the *apeiron*[4]. *Caveat (precise scope):* the Tantrakyma field can *modulate the temporal sector* through the interaction contribution $\hat{t}^{(\mathrm{int})}_{\mu\nu}$ (the commutator channel), see eq. (6.239). Thus, "exclusive" should be read as "primary (with interaction back-reaction via $\hat{t}^{(\mathrm{int})}$)."

- **The Tantrakyma field $\left(\hat{\mathbf{\Theta}}\right)$ as the source of matter:** The Tantrakyma field supplies the theory's **matter-like observables**. Its quanta (*kymons*) are normal-mode excitations whose discrete spectra realize **particle species**; invariance of the Tantrakyma action under its internal symmetry group yields operational **gauge groups**, and the associated Noether charges give the conserved quantum numbers that organize kymons into multiplets within the emergent geometry. In the canonical chronogeometry, $\hat{\mathbf{\Theta}}$ couples through the temporal *interaction* channel $\hat{t}^{(\mathrm{int})}_{\mu\nu}$ (cf. eq. (6.239)); in the optional two-field bookkeeping, one may also form Tantrakyma-only tensors $(\hat{g}^{[\Theta]}_{\mu\nu}, \hat{t}^{[\Theta]}_{\mu\nu})$ and then specify the assembly rule for the physical $\hat{\Sigma}_{\mu\nu}$ accordingly (see the discussion around the two-field extension).

This duality is perfectly captured by the acoustic metaphor:

> *The Gaiamiti field is the resonant structure of the hall.*
> *The Tantrakyma field is the orchestra playing within it.*

The "sound'' that emerges from their interplay is the complete, interacting spectrum of matter and geometry that constitutes the universe.

6.4.1 Holosymmetric Wordwise Bundle-theoretic Formulation of the Foundational Fields

Foundational bundles and gluing data. Let $\{\mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}}\}_{x\in\mathcal{X}_q}$ be the (spectrally regulated) modal fibres of rank n (e.g. $n = 4$ for the fourfold block; cf. Section 6.1.27).

Definition 6.165 (Modal operator bundle and vacuum–tangent/excitation subbundles)**.** All fields below are assumed SOT–C^k ($k \geq 1$) and spectrally localised by cutoffs $P_x = E^{(x)}([0, \Lambda])$.

(i) Operator bundle

A C^*-algebra bundle

$$\mathcal{E}_{\mathrm{op}} := \bigsqcup_{x\in\mathcal{X}_q} \mathcal{B}\left(\mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}}\right) \longrightarrow \mathcal{X}_q,$$

associated to the modal frame principal bundle with structure group $U(n)$ acting fibrewise by the adjoint representation Ad. (Equivalently: the operator-word vault $\mathbb{W}_{\mathrm{op}}$ of Definition 6.114.)

[4] **Apeiron** (*Ápeiron*) — "The Unlimited" or "The Indefinite"; a concept first articulated by Anaximander but central to Pythagorean cosmology. The *apeiron* (*ápeiron*) is the boundless, formless, indefinite primordial substrate. The ordered universe (cosmos) comes into being when this unlimited substrate is articulated by a principle of "limit'' or "form'' (*peras*, *péras*) [69, 72, 70, 71].

(ii) Vacuum–tangent and excitation subbundles

Fix a smooth vacuum section $x \mapsto |\Omega(x)\rangle \in \mathcal{M}_{\text{vac}}$ and the orthogonal projector $\hat{P}_T(x)$ onto $T_{|\Omega(x)\rangle}\mathcal{M}_{\text{vac}}$ (cf. Definition 6.90). Define

$$\boxed{\mathcal{T}_{\text{vac}} := \operatorname{Im} \hat{P}_T \;\longrightarrow\; \mathcal{X}_q, \qquad \mathcal{E}_{\text{exc}} := \operatorname{Im} \hat{P}_{\text{exc}}\,, \qquad \hat{P}_{\text{exc}} := \mathbf{1} - \hat{P}_T.} \tag{6.227}$$

The inner product induced from the ambient/projective geometry identifies $T\mathcal{M}_{\text{vac}} \simeq T^*\mathcal{M}_{\text{vac}}$, allowing dual-frame contractions in local component definitions (used below).

(iii) Total field bundles

The associated total field bundles are defined by

$$\boxed{\mathcal{E}_\Phi := \mathcal{E}_{\text{op}} \otimes \mathcal{T}_{\text{vac}}, \qquad \mathcal{E}_\Theta := \mathcal{E}_{\text{op}} \otimes \mathcal{E}_{\text{exc}}.} \tag{6.228}$$

Remark 6.166 (Transition functions and covariance)**.** On a good cover $\{U_i\}$ of $\mathcal{X}_q$, pick (i) modal frames $g_i : U_i \to U(n)$ and (ii) orthonormal frames $R_i : U_i \to O(r)$ for $\mathcal{T}_{\text{vac}}$ ($r = \operatorname{rank} \mathcal{T}_{\text{vac}}$). On overlaps $U_i \cap U_j$, $g_j = g_{ij} g_i$ with $g_{ij} : U_i \cap U_j \to U(n)$ and $R_j = R_{ij} R_i$ with $R_{ij} : U_i \cap U_j \to O(r)$. Sections glue by $X_i \sim \mathrm{Ad}_{g_{ij}}(X_j)$ in the operator slot and $v_i \sim R_{ij} v_j$ in the tangent/excitation slots. Consequently, local *components* of the Gaiamiti field $\hat{\mathbf{\Phi}}$ transform as

$$\hat{\Phi}^i_{(i)} \;=\; \sum_k R_{ij}{}^i{}_k \; \mathrm{Ad}_{g_{ij}}\!\big(\hat{\Phi}^k_{(j)}\big),$$

i.e. contravariantly in the tangent index and by inner automorphism in the operator slot; components of the Tantrakyma field $\hat{\mathbf{\Theta}}$ transform by $\mathrm{Ad}_{g_{ij}}$ on the operator slot and remain in $\mathcal{E}_{\text{exc}}$ via the projector $\hat{P}_{\text{exc}}$.

Local frames, component fields, and their transformations.

Definition 6.167 (Local decompositions and wordwise components)**.** On $U \subset \mathcal{X}_q$, choose $\{|e_i(x)\rangle\}_{i=1}^r$ orthonormal in $T_{|\Omega(x)\rangle}\mathcal{M}_{\text{vac}}$ with dual frame $\{\langle e^i(x)|\}$ (via the induced inner product on $T\mathcal{M}_{\text{vac}}$), and a modal basis $\{|\alpha\rangle_x\}_{\alpha\in\{0,1,p,s\}}$. Then:

1. **Gaiamiti (geometric) field.**

$$\boxed{\hat{\mathbf{\Phi}}(x) \;=\; \sum_{i=1}^{r} \hat{\Phi}^i(x)\, |e_i(x)\rangle, \qquad \hat{\Phi}^i(x) := \Big(\mathbf{1}\otimes\langle e_i(x)|\Big)\, \hat{\mathbf{\Phi}}(x) \;\in\; \mathcal{B}(\mathcal{H}_{\text{em}}).} \tag{6.229}$$

On overlaps $U_i \cap U_j$, the components transform contravariantly in the tangent index and by inner automorphism on the operator slot:

$$\hat{\Phi}^i_{(i)} \;=\; \sum_{k=1}^{r} R_{ij}{}^i{}_k \; \mathrm{Ad}_{g_{ij}}\!\big(\hat{\Phi}^k_{(j)}\big).$$

2. **Tantrakyma (excitation) field.**

$$\boxed{\hat{\mathbf{\Theta}}(x) \;=\; \sum_{\alpha,\beta\in\{0,1,p,s\}} \theta^{\alpha\beta}(x)\, |\alpha\rangle_x\langle\beta|_x, \qquad \theta(x) := \Big[\theta^{\alpha\beta}(x)\Big]_{\alpha,\beta} \in M_4(\mathbb{C}),} \tag{6.230}$$

with modal covariance $\theta \mapsto g_{ij}\,\theta\,g_{ij}^{\dagger}$ on $U_i \cap U_j$. Confinement to the excitation bundle is enforced by $\hat{P}_{\mathrm{exc}}(x)$; when we display $\theta(x)$, we have fixed a local excitation frame (or projected to $\mathcal{E}_{\mathrm{exc}}(x)$), so the excitation action is suppressed in notation.

Remark 6.168 (Triadic channel projectors)**.** Let $\Pi^{(\mathrm{corr})}, \Pi^{(\mathrm{vac})}, \Pi^{(\mathrm{spec})}$ denote the sector projectors (correlator/vacuum/spectral channels). Applying them component-wise to $\hat{\Phi}^i$ and θ implements the triadic decomposition used throughout, with sectorwise symmetrisation and commutator/anticommutator tomography consistent with Section 6.1.12 and eq. (5.86).

Connections and covariant derivatives.

Definition 6.169 (Relational–modal covariant derivatives on $\mathcal{E}_\Phi$ and $\mathcal{E}_\Theta$)**.** Let $A_\mu \in \mathfrak{u}(n)$ be the modal connection 1-form and $\nabla^{(\mathrm{rel})}$ the Levi–Civita derivative of $\bar{\Sigma}$. Then:

1. *Operator slot.* For $X \in \Gamma(\mathcal{E}_{\mathrm{op}})$,
$$\mathcal{D}_\mu X \;:=\; \partial_\mu X + [A_\mu, X].$$
2. *Vacuum–tangent slot.* With $\omega_\mu{}^i{}_j \in \mathfrak{o}(r)$ the metric connection induced by g_T (Definition 6.104),
$$\nabla_\mu^{(\mathrm{vac})}\Big(\hat{\Phi}^i\,|e_i\rangle\Big) = \Big(\partial_\mu \hat{\Phi}^i + [A_\mu, \hat{\Phi}^i] + \omega_\mu{}^i{}_j \hat{\Phi}^j\Big)\,|e_i\rangle.$$
3. *Excitation slot.* For the smooth projector $P_{\mathrm{exc}}(x)$,
$$\omega_\mu^{(\mathrm{exc})} := P_{\mathrm{exc}}\,(\partial_\mu P_{\mathrm{exc}}), \qquad \mathcal{D}_\mu^{(\mathrm{exc})}\theta := \partial_\mu\theta + [A_\mu, \theta] + [\omega_\mu^{(\mathrm{exc})}, \theta].$$
In the displayed 4×4 matrix representation (after fixing an excitation frame or projection), the term $[\omega_\mu^{(\mathrm{exc})}, \theta]$ is the induced commutator on that operator block; if the excitation factor is kept explicitly, this corresponds to the identity on the operator slot tensored with the connection on $\mathcal{E}_{\mathrm{exc}}$.

The total covariant derivatives are

$$\boxed{\begin{aligned} D_\mu\hat{\mathbf{\Phi}} := \nabla_\mu^{(\mathrm{vac})}\hat{\mathbf{\Phi}} \quad &\Longleftrightarrow \quad D_\mu\hat{\Phi}^i = \partial_\mu\hat{\Phi}^i + [A_\mu, \hat{\Phi}^i] + \omega_\mu{}^i{}_j\hat{\Phi}^j, && (6.231)\\ D_\mu\hat{\mathbf{\Theta}} := \mathcal{D}_\mu^{(\mathrm{exc})}\hat{\mathbf{\Theta}} \quad &\Longleftrightarrow \quad D_\mu\theta = \partial_\mu\theta + [A_\mu, \theta] + [\omega_\mu^{(\mathrm{exc})}, \theta]. && (6.232)\end{aligned}}$$

Proposition 6.170 (Gauge and frame covariance)**.** *Under $g(x) \in U(n)$ and $R(x) \in O(r)$,*
$$\hat{\Phi}^i \mapsto \sum_j R^i{}_j\;\mathrm{Ad}_g\Big(\hat{\Phi}^j\Big), \qquad A_\mu \mapsto gA_\mu g^\dagger - (\partial_\mu g)\,g^\dagger, \qquad \omega_\mu \mapsto R\omega_\mu R^{-1} + (\partial_\mu R)R^{-1},$$
and the component rules (6.231)–(6.232) are covariant. Hence any tensor built from $D_\mu\hat{\mathbf{\Phi}}$ and $D_\mu\hat{\mathbf{\Theta}}$ using $\{\cdot,\cdot\}$ and $[\cdot,\cdot]$ is globally well defined.

Reconstruction of chronogeometry (for reference). Let $\mathsf{H}[X,Y] := \frac{1}{2}\{X^\dagger, Y\}$ and $\mathsf{A}[X,Y] := \frac{i}{2}[X^\dagger, Y]$. For each $x \in \mathcal{X}_q$, set the symmetric "baseline" pairing

$$\hat{B}_{\mu\nu}(x) \;:=\; \mathsf{H}\Big[D_\mu\hat{\mathbf{\Phi}}(x),\, D_\nu\hat{\mathbf{\Phi}}(x)\Big]. \tag{6.233}$$

Project to spatial/temporal channels:

$$\hat{g}_{\mu\nu}(x) \;:=\; \Pi_{\mathrm{dist}}\Big(\,\hat{B}_{\mu\nu}(x)\,\Big), \tag{6.234}$$

$$\hat{t}^{(0)}_{\mu\nu}(x) \;:=\; \Pi_{\mathrm{time}}\Big(\,\hat{B}_{\mu\nu}(x)\,\Big). \tag{6.235}$$

Include the commutator–type interaction (with the Tantrakyma field $\hat{\mathbf{\Theta}}$):

$$\hat{I}_{\mu\nu}(x) \;:=\; \mathsf{A}\Big[D_\mu\hat{\mathbf{\Phi}}(x),\, D_\nu\hat{\mathbf{\Theta}}(x)\Big] \;+\; \mathsf{A}\Big[D_\mu\hat{\mathbf{\Theta}}(x),\, D_\nu\hat{\mathbf{\Phi}}(x)\Big]. \tag{6.236}$$

and its temporal projection:

$$\hat{t}^{(\mathrm{int})}_{\mu\nu}(x) \;:=\; \Pi_{\mathrm{time}}\Big(\,\hat{I}_{\mu\nu}(x)\,\Big), \tag{6.237}$$

$$\hat{t}_{\mu\nu}(x) \;:=\; \hat{t}^{(0)}_{\mu\nu}(x) \;+\; \hat{t}^{(\mathrm{int})}_{\mu\nu}(x). \tag{6.238}$$

The unified chronogeometric tensor is

$$\hat{\Sigma}_{\mu\nu}(x) \;:=\; \hat{g}_{\mu\nu}(x) \;+\; \hat{t}_{\mu\nu}(x). \tag{6.239}$$

Born expectations in the ringed state $|\mathring{\mathbf{\Psi}}\rangle$ give the macroscopic fields:

$$\bar{g}_{\mu\nu}(x) \;=\; \Big\langle \mathring{\mathbf{\Psi}}\Big|\, \hat{g}_{\mu\nu}(x)\,\Big|\mathring{\mathbf{\Psi}}\Big\rangle, \tag{6.240}$$

$$\bar{t}_{\mu\nu}(x) \;=\; \Big\langle \mathring{\mathbf{\Psi}}\Big|\, \hat{t}_{\mu\nu}(x)\,\Big|\mathring{\mathbf{\Psi}}\Big\rangle, \tag{6.241}$$

$$\bar{\Sigma}_{\mu\nu}(x) \;=\; \bar{g}_{\mu\nu}(x) \;+\; \bar{t}_{\mu\nu}(x). \tag{6.242}$$

Holosymmetric law-control dial at the bundle level (optional; no new symbols).

Definition 6.171 (Law-control lift of field bundles)**.** Let $\mathcal{H}_L \simeq \mathbb{C}^2$ with basis $\{|0\rangle_L, |1\rangle_L\}$. For two law branches $k \in \{0,1\}$, define the lifted sections

$$\hat{\mathbf{\Phi}}^{\mathrm{HS}} = |0\rangle\langle 0|_L \otimes \hat{\mathbf{\Phi}}^{(0)} + |1\rangle\langle 1|_L \otimes \hat{\mathbf{\Phi}}^{(1)} + |0\rangle\langle 1|_L \otimes \hat{\Phi}_{\mathrm{mix}} + |1\rangle\langle 0|_L \otimes \hat{\Phi}^\dagger_{\mathrm{mix}},$$

(and analogously for $\hat{\mathbf{\Theta}}$ with a covariantly assembled mixing operator $\hat{\Theta}_{\mathrm{mix}}$; cf. Definition 6.158). Preparation of the law register in $|+_\varphi\rangle_L = \frac{1}{\sqrt{2}}\big(|0\rangle_L + e^{i\varphi}|1\rangle_L\big)$ and compression/uncomputation of $\mathcal{H}_L$ reduce these to phase-programmable effective sections that coincide fibrewise with eqs. (6.209) and (6.223). Their bilinear combinations reproduce the commutator/anticommutator tomography of Section 6.1.12, and—by the gauge/transport covariance of the ingredients (Propositions 6.123 and 6.146)— the construction is covariant under endpoint gauges and frame changes, without altering cone assignments (cf. Remark 6.134 and Corollary 6.87).

Regularity and locality clauses. All operator-valued maps are SOT–C^k on the regulated fibres $P_x\mathcal{H}_{\mathrm{em}}$; spectral locality is preserved by lift, transport, projection, and bilinear formation:

$$P_x\,\hat{g}_{\mu\nu}(x)\,P_x = \hat{g}_{\mu\nu}(x), \qquad P_x\,\hat{t}_{\mu\nu}(x)\,P_x = \hat{t}_{\mu\nu}(x),$$

and similarly for intermediate objects. Sector faithfulness follows by applying the channel projectors $\Pi^{(a)}$ inside (6.239); expectation-level scalars inherit gauge invariance by Corollaries 6.125 and 6.148.

6.5 The Action for the Dynamical Gaiamiti Field

Having rigorously defined the Dynamical Gaiamiti Field, $\hat{\mathbf{\Phi}}(x)$, as the fundamental generator of geometry, we now postulate the action that governs its dynamics. The Principle of Stationary Action applied to this functional yields the equations of motion for the Gaiamiti field—and thereby the evolution of emergent geometric structure. This action describes the dynamics of the quantum fluctuations of the field propagating on the emergent background geometry defined by the unified tensor $\bar{\Sigma}_{\mu\nu}$.

Definition 6.172 (Operator-Valued Lagrangian Density for the Gaiamiti Field)**.** Let $\hat{\mathbf{\Phi}}(x)$ denote the *Dynamical Gaiamiti Field*, interpreted as a section of the bundle

$$\mathrm{End}(\mathcal{H}_{\mathrm{em}}) \otimes T\mathcal{M}_{\mathrm{vac}},$$

over the emergent configuration space $(\mathcal{X}_q, \bar{\Sigma}_{\mu\nu})$, where:

- $\mathcal{H}_{\mathrm{em}}$ is the emergent Hilbert space constructed via the GNS construction from the Quantum Holo-State $|\mathring{\mathbf{\Psi}}\rangle$ (Definition 1.34);
- $\mathcal{M}_{\mathrm{vac}}$ denotes the vacuum manifold associated to the emergent chronogeometric structure (Section 6.1.17);
- $\bar{\Sigma}_{\mu\nu} := \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ is the chronogeometric tensor (Definition 5.26), and $\bar{\Sigma}^{\mu\nu}$ its inverse.

Then the *operator-valued Lagrangian density* for the Gaiamiti field is defined as the operator-traced scalar functional:

$$\mathcal{L}_\Phi(x) := \mathrm{Tr}\left\{\bar{\Sigma}^{\mu\nu}(x)\, g_T\left(\nabla_\mu\hat{\mathbf{\Phi}}(x), \nabla_\nu\hat{\mathbf{\Phi}}(x)\right) + \mathcal{V}_\Phi\left(\hat{\mathbf{\Phi}}(x)\right)\right\}, \tag{6.243}$$

where:

- $\nabla_\mu\hat{\mathbf{\Phi}}(x)$ denotes the total covariant derivative of the operator field, incorporating both geometric (spacetime) and internal (fiber-wise) connections;
- $g_T\colon T\mathcal{M}_{\mathrm{vac}} \times T\mathcal{M}_{\mathrm{vac}} \to \mathrm{End}(\mathcal{H}_{\mathrm{em}})$ is a Hermitian-positive operator-valued inner product on the vacuum tangent bundle;
- $\mathcal{V}_\Phi(\hat{\mathbf{\Phi}}(x)) \in \mathrm{End}(\mathcal{H}_{\mathrm{em}})$ is the Gaiamiti potential operator, encoding self-interactions and symmetry-breaking mechanisms.

The Lagrangian density decomposes naturally into two covariant components:

1. **Kinetic Term:**

$$\mathcal{T}_\Phi(x) := \mathrm{Tr}\left\{\bar{\Sigma}^{\mu\nu}(x)\, g_T\left(\nabla_\mu\hat{\mathbf{\Phi}}(x), \nabla_\nu\hat{\mathbf{\Phi}}(x)\right)\right\}, \tag{6.244}$$

describing the geometric and internal variation of the Gaiamiti field, dynamically coupled to the full relational chronogeometry.

2. **Potential Term:**

$$\mathcal{V}_{\Phi,\mathrm{density}}(x) := \mathrm{Tr}\left\{\mathcal{V}_\Phi\left(\hat{\mathbf{\Phi}}(x)\right)\right\}, \tag{6.245}$$

determining the self-interaction structure and vacuum response of the Gaiamiti field.

The corresponding *Gaiamiti action functional* is given by the integral over the emergent manifold $\mathcal{X}_q$, weighted by the relational volume element $d\mu(x) := d^dx\sqrt{-\det\left(\bar{\Sigma}\right)}$ (Section 5.2.6):

$$S_\Phi[\hat{\mathbf{\Phi}}] := \int_{\mathcal{X}_q} d\mu(x)\,\mathcal{L}_\Phi(x) = \int_{\mathcal{X}_q} d^dx\sqrt{-\det\left(\bar{\Sigma}(x)\right)}\,\mathcal{L}_\Phi(x). \tag{6.246}$$

Remark 6.173 (Covariance and Variational Principle)**.** The Lagrangian density $\mathcal{L}_\Phi(x)$ is manifestly covariant under general coordinate transformations on $\mathcal{X}_q$ and under local gauge transformations acting on the Hilbert fiber bundle $\mathcal{H}_{\mathrm{em}}$. The resulting Euler–Lagrange equations, obtained via the principle of stationary action $\delta S_\Phi = 0$, are therefore consistent with the background-independent operator chronogeometry of the IQR framework.

6.5.1 The Gaiamiti Field Action Functional

Definition 6.174 (Gaiamiti Field Action Functional)**.** The **Gaiamitiic Action Functional**, S_Φ, is fundamentally defined by the total relational self-correlation of the geometric sector, integrated over the emergent manifold. For any admissible state $|\mathbf{\Psi}\rangle$, its value is given by:

$$S_{\mathbf{\Phi}}[|\mathbf{\Psi}\rangle] := \int_{\mathcal{X}_q} d^dx\,\sqrt{-\det\bar{\Sigma}^{(\mathbf{\Psi})}_{\mu\nu}(x)}\;\mathrm{Tr}\left(\langle\mathbf{\Psi}|\hat{G}_{\mathbf{\Phi}\mathbf{\Phi}}(x,x)|\mathbf{\Psi}\rangle\right). \tag{6.247}$$

For a dynamical field like $\hat{\mathbf{\Phi}}(x)$, the Lagrangian density—the expectation value of the local self-correlator—is expressed explicitly in terms of the field and its covariant derivatives. This yields the standard kinetic-plus-potential form of the action, which we take as its operational definition:

$$\boxed{S_\Phi[\hat{\mathbf{\Phi}}] := \int_{\mathcal{X}_q} d^dx\sqrt{-\det\left(\bar{\Sigma}\right)}\,\mathrm{Tr}\left\{\bar{\Sigma}^{\mu\nu}(x) g_T\left(\nabla_\mu\hat{\mathbf{\Phi}}(x), \nabla_\nu\hat{\mathbf{\Phi}}(x)\right) + \mathcal{V}_\Phi\left(\hat{\mathbf{\Phi}}(x)\right)\right\},} \tag{6.248}$$

where:

leftmargin=3em,itemsep=0.25em]

- The kinetic term, using the inner product g_T from Definition 6.104, is, in an orthonormal basis,

$$\sum_i(\nabla_\mu\hat{\Phi}^i)^\dagger(\nabla_\nu\hat{\Phi}^i).$$

- ∇_μ is the full covariant derivative compatible with the bundle structure (Section 6.5.3).
- $\mathcal{V}_\Phi(\hat{\mathbf{\Phi}})$ is a local potential functional encoding geometric self-interaction.

These two forms are understood to be equivalent, with the second providing the explicit dynamical structure implied by the fundamental correlator-based definition.

Interpretation of the Action This action functional defines the *dynamical core* of the geometric sector. The **kinetic term** governs the propagation of the Gaiamiti field across the emergent manifold. Its coupling to the unified tensor $\bar{\Sigma}^{\mu\nu}$ ensures that these dynamics are sensitive not only to the curvature of distinguishability ($\bar{g}_{\mu\nu}$) but also to the curvature of quantum phase flow ($\bar{t}_{\mu\nu}$).

The **potential term**, $\mathcal{V}_\Phi(\hat{\mathbf{\Phi}})$, governs the self-interaction and local stability of the geometry. Its structure dictates the "energy landscape" of possible vacuum geometries. The minima of this potential correspond to the stable spacetime structures the universe could inhabit, providing a natural mechanism for spontaneous symmetry breaking.

Remark 6.175 (Background Independence and Self-Consistency)**.** The most profound feature of this action is its dynamic self-consistency. While it uses the expectation-level tensor $\bar{\Sigma}_{\mu\nu}(x)$ to define the dynamics of the operator $\hat{\mathbf{\Phi}}$, this tensor is itself an emergent consequence of the correlations of $\hat{\mathbf{\Phi}}$, evaluated in the Quantum Holo-State. This is not a circularity but the hallmark of a truly *background-independent* framework. It is a self-referential loop where geometry determines the propagation of the Gaiamiti field, while the Gaiamiti field generates the very geometry in which it propagates.

6.5.2 The Holosymmetric Action for the Dynamical Gaiamiti Field

Definition 6.176 (Wordwise kinetic pairing and covariant data)**.** Let D_μ be the total covariant derivative on $\mathcal{E}_\Phi = \mathcal{E}_{\text{op}} \otimes \mathcal{T}_{\text{vac}}$ (cf. eq. (6.231)), built from the modal connection A_μ and the vacuum–tangent connection ω_μ, and write $D^\mu := \bar{\Sigma}^{\mu\nu} D_\nu$. For a local g_T–orthonormal frame $\{|e_i\rangle\}_{i=1}^r$ of $\mathcal{T}_{\text{vac}}$, set components $\hat{\Phi}^i := \langle e^i | \hat{\mathbf{\Phi}} \rangle$ and $D_\mu \hat{\Phi}^i$ (dual-frame contraction as in Definition 6.167 and eq. (6.229)). Define the operator Hermitian/antisymmetric pairings

$$\mathsf{H}[X,Y] := \tfrac{1}{2}\{X^\dagger, Y\}, \qquad \mathsf{A}[X,Y] := \tfrac{i}{2}[X^\dagger, Y],$$

and realise the tangent inner product g_T componentwise by

$$\boxed{\mathcal{G}_T\Big(D_\mu \hat{\mathbf{\Phi}}, D_\nu \hat{\mathbf{\Phi}}\Big) := \sum_{i=1}^{r} \mathsf{H}\Big[D_\mu \hat{\Phi}^i,\ D_\nu \hat{\Phi}^i\Big] \ \in\ \mathcal{B}(\mathcal{H}_{\text{em}}).} \tag{6.249}$$

This construction is gauge/frame covariant under $U(n)$ (modal) and $O(r)$ (tangent) changes (cf. Proposition 6.170).

Definition 6.177 (Gaiamiti Lagrangian density and action (wordwise form))**.** Let $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ and $\bar{\Sigma}^{\mu\nu}$ its inverse on the nondegenerate domain. The (scalar) Lagrangian density

and the action are

$$\boxed{\mathcal{L}_\Phi(x) := \mathrm{Tr}\Big\{\bar{\Sigma}^{\mu\nu}(x)\,\mathcal{G}_T\Big(D_\mu\hat{\mathbf{\Phi}}, D_\nu\hat{\mathbf{\Phi}}\Big) + \mathcal{V}_\Phi\Big(\hat{\mathbf{\Phi}}(x)\Big)\Big\},} \tag{6.250}$$

$$\boxed{S_\Phi[\hat{\mathbf{\Phi}}] := \int_{\mathcal{X}_q} d^d x\,\sqrt{-\det\bar{\Sigma}(x)}\;\mathcal{L}_\Phi(x).} \tag{6.251}$$

Here $\mathcal{V}_\Phi$ is any local, gauge–covariant, frame–invariant potential built from wordwise scalars. (Traces are taken over the regulated fibre $P_x\mathcal{H}_{\mathrm{em}}$; cf. spectral locality in Definition 6.165.)

Remark 6.178 (Local component form)**.** In a g_T–orthonormal vacuum–tangent frame,

$$\mathcal{L}_\Phi = \sum_{i=1}^{r} \mathrm{Tr}\Big\{\bar{\Sigma}^{\mu\nu}\,\tfrac{1}{2}\Big((D_\mu\hat{\Phi}^i)^\dagger D_\nu\hat{\Phi}^i + D_\nu\hat{\Phi}^i(D_\mu\hat{\Phi}^i)^\dagger\Big)\Big\} + \mathrm{Tr}\,\mathcal{V}_\Phi(\{\hat{\Phi}^i\}),$$

with $D^\mu = \bar{\Sigma}^{\mu\nu}D_\nu$ and spectral regulation tacit (projectors P_x).

Definition 6.179 (Conjugate current and Euler–Lagrange equations)**.** Define the conjugate word-currents

$$\boxed{\hat{\Pi}^\nu{}_i := \frac{\partial}{\partial(D_\nu\hat{\Phi}^i)}\,\mathcal{G}_T\Big(D_\mu\hat{\mathbf{\Phi}}, D^\mu\hat{\mathbf{\Phi}}\Big) = \mathsf{H}\Big[\mathbf{1},\ D^\nu\hat{\Phi}^i\Big] = \tfrac{1}{2}\Big(D^\nu\hat{\Phi}^i + (D^\nu\hat{\Phi}^i)^\dagger\Big).} \tag{6.252}$$

Varying S_Φ in the SOT with compact support and using $\nabla^{(\mathrm{rel})}_\lambda\bar{\Sigma}^{\mu\nu} = 0$ yields, for each i,

$$\boxed{\nabla^{(\mathrm{rel})}_\mu\Big(\sqrt{-\det\bar{\Sigma}}\;\bar{\Sigma}^{\mu\nu}\,\mathrm{Tr}\,\hat{\Pi}_{\nu i}\Big) - \sqrt{-\det\bar{\Sigma}}\;\frac{\delta\,\mathrm{Tr}\,\mathcal{V}_\Phi}{\delta\hat{\Phi}^{i\dagger}} = 0.} \tag{6.253}$$

Equivalently (inside the trace), $\nabla^{(\mathrm{rel})}_\mu(\bar{\Sigma}^{\mu\nu}D_\nu\hat{\Phi}^i) + \cdots = 0$, where the dots denote the Fréchet gradient of $\mathcal{V}_\Phi$.

Definition 6.180 (Symmetric stress–energy tensor)**.** Variation of S_Φ with respect to $\bar{\Sigma}^{\mu\nu}$ at fixed $\hat{\mathbf{\Phi}}$ gives

$$\boxed{T^{(\Phi)}_{\mu\nu} := \mathrm{Tr}\Big\{\mathcal{G}_T\Big(D_\mu\hat{\mathbf{\Phi}}, D_\nu\hat{\mathbf{\Phi}}\Big)\Big\} - \tfrac{1}{2}\,\bar{\Sigma}_{\mu\nu}\,\mathrm{Tr}\Big\{\bar{\Sigma}^{\alpha\beta}\mathcal{G}_T\Big(D_\alpha\hat{\mathbf{\Phi}}, D_\beta\hat{\mathbf{\Phi}}\Big) + \mathcal{V}_\Phi(\hat{\mathbf{\Phi}})\Big\},} \tag{6.254}$$

with $\nabla^{(\mathrm{rel})}_\mu T^\mu{}_\nu = 0$ on shell (metric compatibility).

Proposition 6.181 (Gauge/frame covariance and positivity)**.** *$\mathcal{L}_\Phi$ and S_Φ are invariant under modal $U(n)$ and tangent $O(r)$ frame changes (via Ad_g and R, respectively; cf. Proposition 6.170). If $\mathcal{V}_\Phi$ is bounded below and each $D_\mu\hat{\Phi}^i$ is spectrally regulated, then $\mathrm{Tr}\,\mathsf{H}[X,X] = \frac{1}{2}\,\mathrm{Tr}\big(X^\dagger X + XX^\dagger\big) \geq 0$, so the kinetic contribution is nonnegative.*

Definition 6.182 (Triadic (channel) resolution)**.** With channel projectors $\Pi^{(a)}$ for $a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$,

$$\boxed{\mathcal{L}_\Phi = \sum_a \mathrm{Tr}\Big\{\bar{\Sigma}^{\mu\nu}\,\Pi^{(a)}\Big(\sum_i \mathsf{H}[D_\mu\hat{\Phi}^i, D_\nu\hat{\Phi}^i]\Big)\Big\} + \mathrm{Tr}\,\mathcal{V}_\Phi.} \tag{6.255}$$

If the $\Pi^{(a)}$ commute with the covariant data, cross-channel interferences vanish; otherwise, phase-odd mixings enter through A (cf. Section 6.1.12).

Holosymmetric law-control (reduced action; no new symbols). For two branches $k \in \{0,1\}$ with fields $\hat{\mathbf{\Phi}}^{(k)}$ and Lagrangians $\mathcal{L}_\Phi^{(k)} := \mathcal{L}_\Phi[\hat{\mathbf{\Phi}}^{(k)}]$, define the *cross* bilinears

$$\mathcal{S}_\Phi := \mathrm{Tr}\Big\{\bar{\Sigma}^{\mu\nu}\sum_i \tfrac{1}{2}\Big(\{(D_\mu\hat{\Phi}^{(0)i})^\dagger,\, D_\nu\hat{\Phi}^{(1)i}\} + \{(D_\mu\hat{\Phi}^{(1)i})^\dagger,\, D_\nu\hat{\Phi}^{(0)i}\}\Big)\Big\},$$
$$\mathcal{C}_\Phi := \mathrm{Tr}\Big\{\bar{\Sigma}^{\mu\nu}\sum_i \tfrac{i}{2}\Big([(D_\mu\hat{\Phi}^{(0)i})^\dagger,\, D_\nu\hat{\Phi}^{(1)i}] + [(D_\mu\hat{\Phi}^{(1)i})^\dagger,\, D_\nu\hat{\Phi}^{(0)i}]\Big)\Big\},$$

and, analogously, potential cross-terms built from $\{\hat{\Phi}^{(0)i}, \hat{\Phi}^{(1)i}\}$. Preparing the law register in $|+_\varphi\rangle_L$ and introducing the dial θ yields the *reduced HS Lagrangian density*

$$\boxed{\mathcal{L}_\Phi^{\langle \mathrm{HS}\rangle} = \tfrac{1}{2}\Big(\mathcal{L}_\Phi^{(0)} + \mathcal{L}_\Phi^{(1)}\Big) + \mathrm{Re}\Big(e^{-i\varphi}\Big[\cos\theta\,\mathcal{S}_\Phi + \sin\theta\,\mathcal{C}_\Phi\Big]\Big)\,,} \tag{6.256}$$

which cleanly separates anticommutator- (symmetric) and commutator- (antisymmetric) type couplings (cf. Definition 6.171). The causal cone assignments (fixed by $\bar{g}^{\mathrm{Lor}}$) are unaffected (cf. Definition 5.77).

Remark 6.183 (Boundary terms and well-posedness)**.** Wordwise integration by parts gives the boundary contribution

$$\delta S_\Phi\big|_{\partial\Omega} = \int_{\partial\Omega} d^{d-1}\Sigma_\mu \sqrt{-\det\bar{\Sigma}}\;\mathrm{Tr}\Big\{\bar{\Sigma}^{\mu\nu}\sum_i \mathsf{H}[\delta\hat{\Phi}^i,\, D_\nu\hat{\Phi}^i]\Big\}.$$

Dirichlet ($\delta\hat{\mathbf{\Phi}}|_{\partial\Omega} = 0$) or Neumann ($\bar{\Sigma}^{\mu\nu}\hat{\Pi}_{\nu i}\, n_\mu = 0$) conditions render the variational problem well posed.

6.5.3 Covariant Derivative of the Gaiamiti Field

To describe the dynamics of the Gaiamiti field, $\hat{\mathbf{\Phi}}(x)$, across the emergent quantum configuration space, $\mathcal{X}_q$, a simple partial derivative is insufficient. The Gaiamiti field is an operator-valued section of a non-trivial fiber bundle, and its variation is subject to two distinct geometric structures: the curvature of the emergent relational manifold and the "twist" of the internal operator space. The full covariant derivative, ∇_μ, is therefore a composite object that accounts for both effects.

Definition 6.184 (Covariant Derivative of the Gaiamiti Field)**.** Let $\hat{\mathbf{\Phi}}$ be a smooth operator-valued vector field over $\mathcal{X}_q$, formally a section of the bundle $\mathrm{End}(\mathcal{H}_{\mathrm{em}}) \otimes T\mathcal{M}_{\mathrm{vac}}$. Its covariant derivative, $\nabla_\mu\hat{\mathbf{\Phi}}$, is defined in its coordinate-free form as the sum of a base-space connection and a fiber connection:

$$\boxed{\nabla_\mu\hat{\mathbf{\Phi}} := \nabla_\mu^{(\mathrm{rel})}\hat{\mathbf{\Phi}} + D_\mu\hat{\mathbf{\Phi}},} \tag{6.257}$$

where:

- $\nabla_\mu^{(\mathrm{rel})}$ is the **Relational Levi-Civita Derivative** compatible with the unified tensor $\bar{\Sigma}_{\mu\nu}(x)$ (Definition 6.7), acting on the tangent vector indices of $\hat{\mathbf{\Phi}}$.

- $D_\mu := \partial_\mu + i[\mathcal{A}_\mu(x), \cdot]$ is the **gauge-covariant derivative**, acting on the operator-valued components of $\hat{\mathbf{\Phi}}$ via the Operator Bundle Connection $\mathcal{A}_\mu(x)$.

In a local basis $\{|e_\nu(x)\rangle\}$ for the tangent space $T\mathcal{M}_{\text{vac}}$, this takes the component form:

$$\boxed{(\nabla_\mu \hat{\mathbf{\Phi}})^\lambda(x) := \partial_\mu \hat{\mathbf{\Phi}}^\lambda(x) + \Gamma^\lambda_{\mu\nu}[\bar{\Sigma}](x)\, \hat{\mathbf{\Phi}}^\nu(x) + i\left[\mathcal{A}_\mu(x), \hat{\mathbf{\Phi}}^\lambda(x)\right],} \tag{6.258}$$

where $\hat{\mathbf{\Phi}}^\lambda(x) \in \text{End}(\mathcal{H}_{\text{em}})$ are the operator-valued components of the field.

Interpretation of the Composite Derivative The dual structure of the covariant derivative ∇_μ is a necessary consequence of the dual identity of the Gaiamiti field as both a geometric vector and a quantum operator.

The first correction term, governed by the *Relational Christoffel symbols* ($\Gamma^\lambda_{\mu\nu}[\bar{\Sigma}]$), ensures covariance with respect to the geometry of the base manifold $\mathcal{X}_q$. It is the mathematical device, familiar from General Relativity, that allows us to separate the intrinsic dynamics of the field from the artifacts of our chosen coordinate system.

The second correction term, governed by the *Operator Bundle Connection* ($\mathcal{A}_\mu$), ensures that the physics is independent of the local choice of basis in the Hilbert space fibers. It is a gauge connection that allows for the parallel transport of quantum information, making the theory's predictions invariant under local unitary transformations.

In essence, the full covariant derivative ∇_μ is a sophisticated and elegant tool that unifies the principle of general covariance with the principle of local gauge invariance. It is the correct and complete way to describe the dynamics of a quantum field that is simultaneously the source and subject of geometry.

6.5.4 Holosymmetric Covariant Derivative of the Gaiamiti Field

Definition 6.185 (Total covariant derivative as base + vacuum–tangent + operator connections)**.** Let $\hat{\mathbf{\Phi}}$ be a smooth section of $\text{End}(\mathcal{H}_{\text{em}}) \otimes T\mathcal{M}_{\text{vac}}$ over $\mathcal{X}_q$. The *total covariant derivative* is

$$\boxed{\nabla_\mu \hat{\mathbf{\Phi}} := \underbrace{\nabla^{(\text{rel})}_\mu \hat{\mathbf{\Phi}}}_{\text{base (Levi–Civita on } \bar{\Sigma})} + \underbrace{\nabla^{(T)}_\mu \hat{\mathbf{\Phi}}}_{\text{pullback connection on } T\mathcal{M}_{\text{vac}}} + \underbrace{\mathcal{D}_\mu \hat{\mathbf{\Phi}}}_{\text{operator (adjoint) connection}},} \tag{6.259}$$

where:

- $\nabla^{(\text{rel})}_\mu$ is the Relational Levi–Civita derivative compatible with $\bar{\Sigma}_{\alpha\beta}$ ($\nabla^{(\text{rel})}_\mu \bar{\Sigma}_{\alpha\beta} = 0$, torsion-free).
- $\nabla^{(T)}_\mu$ is the vacuum–tangent pullback connection induced by $x \mapsto |\Omega(x)\rangle$ and the metric g_T ($\nabla^{(T)}_\mu g_T = 0$).
- $\mathcal{D}_\mu := \partial_\mu + [A_\mu, \cdot]$ is the $\text{End}(\mathcal{H}_{\text{em}})$-valued operator-bundle connection (adjoint representation; cf. Definition 6.169).

When $\hat{\mathbf{\Phi}}$ carries no explicit base indices, $\nabla_\mu \hat{\mathbf{\Phi}}$ reduces to the total derivative $D_\mu \hat{\mathbf{\Phi}}$ of eq. (6.231).

Definition 6.186 (Component forms)**.** Choose a g_T-orthonormal local frame $\{|e_i(x)\rangle\}_{i=1}^r$ of $T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}$ with dual $\{\langle e^i(x)|\}$ and write $\hat{\mathbf{\Phi}} = \sum_i \hat{\Phi}^i\,|e_i\rangle$. Then

$$\boxed{(\nabla_\mu \hat{\mathbf{\Phi}})^i = \partial_\mu \hat{\Phi}^i + \omega_\mu{}^i{}_j\,\hat{\Phi}^j + [A_\mu, \hat{\Phi}^i],} \tag{6.260}$$

where $\omega_\mu{}^i{}_j \in \mathfrak{o}(r)$ are the coefficients of the pullback connection on $T\mathcal{M}_{\mathrm{vac}}$. If $\hat{\mathbf{\Phi}}$ additionally carries a base vector index, write components $\hat{\mathbf{\Phi}}^\lambda$ and

$$\boxed{(\nabla_\mu \hat{\mathbf{\Phi}})^\lambda = \partial_\mu \hat{\mathbf{\Phi}}^\lambda + \Gamma^\lambda{}_{\mu\nu}[\bar{\Sigma}]\,\hat{\mathbf{\Phi}}^\nu + [A_\mu, \hat{\mathbf{\Phi}}^\lambda].} \tag{6.261}$$

Here $\Gamma[\bar{\Sigma}]$ is the Levi–Civita connection of $\bar{\Sigma}$, and ω_μ acts on tangent indices as above.

Remark 6.187 (Leibniz, derivation, and SOT regularity)**.** Since $\mathcal{D}_\mu = \partial_\mu + [A_\mu, \cdot]$ is an inner derivation, $\mathcal{D}_\mu(XY) = (\mathcal{D}_\mu X)Y + X(\mathcal{D}_\mu Y)$. All derivatives are taken in the strong–operator topology, uniformly on regulated fibres $P_x\mathcal{H}_{\mathrm{em}}$ (cf. Definition 6.165).

Proposition 6.188 (Gauge/frame covariance)**.** *Under modal frame changes $g(x) \in U(n)$ and tangent rotations $R(x) \in O(r)$,*

$$\hat{\Phi}^i \mapsto \mathrm{Ad}_g(\hat{\Phi}^j)\,R_j{}^i, \qquad A_\mu \mapsto gA_\mu g^\dagger - (\partial_\mu g)\,g^\dagger, \qquad \omega_\mu \mapsto R\omega_\mu R^{-1} + (\partial_\mu R)R^{-1},$$

and $(\nabla_\mu \hat{\mathbf{\Phi}})$ transforms covariantly in the same way. If $\hat{\mathbf{\Phi}}$ carries spacetime indices, $\Gamma[\bar{\Sigma}]$ transforms as the Levi–Civita connection.

Holosymmetric wordwise proof. Work in a chart $U \subset \mathcal{X}_q$ and choose a g_T–orthonormal vacuum–tangent frame $\{|e_i\rangle\}_{i=1}^r$ with dual $\{\langle e^i|\}$, so that $\hat{\mathbf{\Phi}} = \sum_i \hat{\Phi}^i\,|e_i\rangle$ and the component form of the total covariant derivative is (cf. Definition 6.186)

$$(\nabla_\mu \hat{\mathbf{\Phi}})^i = \partial_\mu \hat{\Phi}^i + \omega_\mu{}^i{}_j\,\hat{\Phi}^j + [A_\mu, \hat{\Phi}^i].$$

Gauge/frame transforms. Let $g(x) \in U(n)$ act on the operator slot by the adjoint automorphism $\mathrm{Ad}_g(X) = gXg^\dagger$, and let $R(x) \in O(r)$ rotate the tangent frame $|e_i\rangle \mapsto |e'_i\rangle := R_i{}^j|e_j\rangle$ (so $\langle e'^i| = \langle e^j|\,R^{-1}{}_j{}^i$). The field components transform as

$$\hat{\Phi}'^i \;=\; (\mathbf{1}\otimes\langle e'^i|)\,\hat{\mathbf{\Phi}} \;=\; \mathrm{Ad}_g(\hat{\Phi}^j)\,R_j{}^i.$$

The connections transform as in the statement:

$$A'_\mu = gA_\mu g^\dagger - (\partial_\mu g)g^\dagger, \qquad \omega'_\mu = R\omega_\mu R^{-1} + (\partial_\mu R)R^{-1}.$$

(All derivatives are SOT on regulated fibres; cf. Definition 6.165.)

Covariance of $\nabla_\mu\hat{\mathbf{\Phi}}$. Compute $(\nabla'_\mu\hat{\mathbf{\Phi}}')^i$ using Definition 6.186:

$$\begin{aligned}(\nabla'_\mu\hat{\mathbf{\Phi}}')^i &= \partial_\mu\hat{\Phi}'^i + \omega'_\mu{}^i{}_k\,\hat{\Phi}'^k + [A'_\mu,\hat{\Phi}'^i]\\ &= \partial_\mu\Big(\mathrm{Ad}_g(\hat{\Phi}^j)R_j{}^i\Big) + \Big(R\omega_\mu R^{-1} + (\partial_\mu R)R^{-1}\Big)^i{}_k\,\mathrm{Ad}_g(\hat{\Phi}^\ell)\,R_\ell{}^k\\ &\quad + [\,gA_\mu g^\dagger - (\partial_\mu g)g^\dagger,\ \mathrm{Ad}_g(\hat{\Phi}^j)\,R_j{}^i\,].\end{aligned}$$

Use Leibniz, $\partial_\mu(\mathrm{Ad}_g X) = \mathrm{Ad}_g(\partial_\mu X) + [\,(\partial_\mu g)g^\dagger,\ \mathrm{Ad}_g X\,]$, and $R^{-1}\partial_\mu R = (\partial_\mu R)R^{-1}$:

$$\begin{aligned}(\nabla'_\mu\hat{\mathbf{\Phi}}')^i = \mathrm{Ad}_g(\partial_\mu\hat{\Phi}^j)R_j{}^i + \underbrace{[\,(\partial_\mu g)g^\dagger,\ \mathrm{Ad}_g(\hat{\Phi}^j)\,]R_j{}^i}_{(*)} + \mathrm{Ad}_g(\hat{\Phi}^\ell)(\partial_\mu R)_\ell{}^i\\ +\ R_m{}^i\,\omega_\mu{}^m{}_k\,\mathrm{Ad}_g(\hat{\Phi}^\ell)\,R_\ell{}^k +\ (\partial_\mu R)_m{}^i\,\mathrm{Ad}_g(\hat{\Phi}^\ell)\,R_\ell{}^m\\ +\ \underbrace{[\,gA_\mu g^\dagger,\ \mathrm{Ad}_g(\hat{\Phi}^j)\,]R_j{}^i}_{=\ \mathrm{Ad}_g([A_\mu,\hat{\Phi}^j])R_j{}^i} - \underbrace{[\,(\partial_\mu g)g^\dagger,\ \mathrm{Ad}_g(\hat{\Phi}^j)\,]R_j{}^i}_{(*)}.\end{aligned}$$

The two terms marked $(*)$ cancel identically. The $(\partial_\mu R)$ terms cancel as well: $\mathrm{Ad}_g(\hat{\Phi}^\ell)(\partial_\mu R)_\ell{}^i + (\partial_\mu R)_m{}^i\mathrm{Ad}_g(\hat{\Phi}^\ell)R_\ell{}^m = 0$ after reindexing (this is the standard compensation by the inhomogeneous piece of ω'_μ). What remains is

$$(\nabla'_\mu\hat{\mathbf{\Phi}}')^i = \mathrm{Ad}_g(\partial_\mu\hat{\Phi}^j)R_j{}^i + R_m{}^i\,\omega_\mu{}^m{}_k\,\mathrm{Ad}_g(\hat{\Phi}^\ell)\,R_\ell{}^k + \mathrm{Ad}_g([A_\mu,\hat{\Phi}^j])R_j{}^i.$$

Using $R_m{}^i R^{-1k}{}_j = \delta^i{}_j$, this is

$$(\nabla'_\mu\hat{\mathbf{\Phi}}')^i = \mathrm{Ad}_g\Big(\,\partial_\mu\hat{\Phi}^j + \omega_\mu{}^j{}_k\hat{\Phi}^k + [A_\mu,\hat{\Phi}^j]\,\Big)\,R_j{}^i = \mathrm{Ad}_g\Big((\nabla_\mu\hat{\mathbf{\Phi}})^j\Big)\,R_j{}^i,$$

which is precisely the claimed covariance:

$$\nabla_\mu\hat{\mathbf{\Phi}}\ \longmapsto\ \mathrm{Ad}_g(\nabla_\mu\hat{\mathbf{\Phi}})\ \text{with the tangent index rotated by } R.$$

Base indices (Levi–Civita part). If $\hat{\mathbf{\Phi}}$ carries spacetime indices, the additional $\Gamma[\bar{\Sigma}]$ terms transform with the usual Levi–Civita connection rule under frame changes (and diffeomorphisms), so the same calculation goes through with the base index contracted covariantly by $\Gamma^\lambda{}_{\mu\nu}[\bar{\Sigma}]$ (cf. Definition 6.186).

Conclusion. Therefore under $g \in U(n)$ and $R \in O(r)$ the components transform as stated,

$$\hat{\Phi}^i \mapsto \mathrm{Ad}_g(\hat{\Phi}^j)\,R_j{}^i, \quad A_\mu \mapsto gA_\mu g^\dagger - (\partial_\mu g)\,g^\dagger, \quad \omega_\mu \mapsto R\omega_\mu R^{-1} + (\partial_\mu R)R^{-1},$$

and $\nabla_\mu\hat{\mathbf{\Phi}}$ is covariant in the same sense. This proves Proposition 6.188. □

Definition 6.189 (Total curvature acting on $\hat{\mathbf{\Phi}}$)**.** The commutator of covariant derivatives yields

$$\boxed{[\nabla_\mu,\nabla_\nu]\hat{\mathbf{\Phi}} = \ \mathcal{R}_{\mu\nu}\cdot\hat{\mathbf{\Phi}}\ +\ [\mathcal{F}_{\mu\nu},\hat{\mathbf{\Phi}}], \qquad \mathcal{F}_{\mu\nu} = \partial_{[\mu}A_{\nu]} + [A_\mu,A_\nu],}\tag{6.262}$$

where $\mathcal{R}_{\mu\nu}$ collects the base Riemann curvature $R[\bar{\Sigma}]$ and the vacuum–tangent curvature $\Omega_{\mu\nu} = \partial_{[\mu}\omega_{\nu]} + [\omega_\mu,\omega_\nu]$, acting on the corresponding indices of $\hat{\mathbf{\Phi}}$.

Definition 6.190 (Wordwise (commutator/anticommutator) split of kinetic transport)**.** For operator components X, Y define $\mathsf{H}[X,Y] := \frac{1}{2}\{X^\dagger, Y\}$ and $\mathsf{A}[X,Y] := \frac{i}{2}[X^\dagger, Y]$ (cf. Definition 6.176). The kinetic density used in the action (Section 6.5.2) is built from $\sum_i \mathsf{H}[(\nabla_\mu \hat{\Phi}^i), (\nabla_\nu \hat{\Phi}^i)]$, while A captures phase–odd channels.

Holosymmetric reduction for kinetic pairings (no new symbols). Let $(\nabla_\mu \hat{\mathbf{\Phi}})^{(k)}$ denote the branchwise covariant derivatives for $k \in \{0,1\}$. Define the cross (symmetric/antisymmetric) *kinetic* mixes (componentwise in i)

$$\mathcal{S}_{\mu\nu} := \sum_i \mathsf{H}\Big[(\nabla_\mu \hat{\Phi}^{(0)i}),\, (\nabla_\nu \hat{\Phi}^{(1)i})\Big] + \sum_i \mathsf{H}\Big[(\nabla_\mu \hat{\Phi}^{(1)i}),\, (\nabla_\nu \hat{\Phi}^{(0)i})\Big],$$
$$\mathcal{C}_{\mu\nu} := \sum_i \mathsf{A}\Big[(\nabla_\mu \hat{\Phi}^{(0)i}),\, (\nabla_\nu \hat{\Phi}^{(1)i})\Big] + \sum_i \mathsf{A}\Big[(\nabla_\mu \hat{\Phi}^{(1)i}),\, (\nabla_\nu \hat{\Phi}^{(0)i})\Big].$$

Preparing the law register in $|+_\varphi\rangle_L$ and introducing the dial θ, the *reduced* kinetic density entering the action is

$$\boxed{\begin{aligned}\sum_i \mathsf{H}\Big[(\nabla_\mu \hat{\Phi}^{\langle \mathrm{HS}\rangle i}), (\nabla_\nu \hat{\Phi}^{\langle \mathrm{HS}\rangle i})\Big] = \tfrac{1}{2}\sum_i \Big(\mathsf{H}\Big[(\nabla_\mu \hat{\Phi}^{(0)i}), (\nabla_\nu \hat{\Phi}^{(0)i})\Big] + \mathsf{H}\Big[(\nabla_\mu \hat{\Phi}^{(1)i}), (\nabla_\nu \hat{\Phi}^{(1)i})\Big]\Big) \\ + \mathrm{Re}\Big(e^{-i\varphi}\Big[\cos\theta\, \mathcal{S}_{\mu\nu} + \sin\theta\, \mathcal{C}_{\mu\nu}\Big]\Big),\end{aligned}} \tag{6.263}$$

which cleanly separates anticommutator- vs. commutator-type inter-branch couplings *without* introducing new mixing symbols (cf. Definition 6.171).

Proposition 6.191 (Compatibility with the action and cone structure)**.** *With* $\nabla_\mu^{(\mathrm{rel})} \bar{\Sigma}_{\alpha\beta} = 0$ *and* $\nabla_\mu^{(T)} g_T = 0$*, inserting* (6.259)–(6.263) *into the Lagrangian of Section 6.5.2 yields covariant Euler–Lagrange equations. Holosymmetric reduction only modulates interference amplitudes in the kinetic density; it does not alter the effective cone assignment (which depends on* $\bar{g}^{\mathrm{Lor}}$ *alone; cf. Definition 5.77).*

Holosymmetric wordwise proof. **Step 1: Covariance of the kinetic density under** $U(n)\times O(r)$**.** Insert eq. (6.259) in place of the covariant data used in Section 6.5.2. In components (with a g_T–orthonormal frame),

$$\mathcal{G}_T\Big(\nabla_\mu \hat{\mathbf{\Phi}}, \nabla_\nu \hat{\mathbf{\Phi}}\Big) = \sum_{i=1}^{r} \mathsf{H}\Big[(\nabla_\mu \hat{\Phi}^i), (\nabla_\nu \hat{\Phi}^i)\Big].$$

By Proposition 6.188, $(\nabla_\mu \hat{\Phi}^i)$ transforms as $(\nabla_\mu \hat{\Phi}^i) \mapsto \mathrm{Ad}_g\big((\nabla_\mu \hat{\Phi}^j)\big) R_j{}^i$. Since $R \in O(r)$ acts orthogonally on the i-index and Ad_g acts by inner automorphism on the operator slot, the sum over i is invariant, and Tr kills the inner automorphism:

$$\sum_i \mathsf{H}\Big[(\nabla_\mu \hat{\Phi}^i), (\nabla_\nu \hat{\Phi}^i)\Big] \;\longmapsto\; \sum_i \mathsf{H}\Big[(\nabla_\mu \hat{\Phi}^i), (\nabla_\nu \hat{\Phi}^i)\Big], \qquad \mathrm{Tr}\{\cdot\} \text{ invariant.}$$

Hence the kinetic part in eq. (6.250) is a scalar under $U(n)\times O(r)$, and so is $\mathcal{V}_\Phi$ by assumption. Therefore $\mathcal{L}_\Phi$ is gauge/frame covariant.

Step 2: Covariant Euler–Lagrange equations. Vary S_Φ in eq. (6.251) with compact support. Using metric compatibility $\nabla^{(\mathrm{rel})}_\lambda \bar{\Sigma}^{\mu\nu} = 0$ and the divergence identity $\nabla^{(\mathrm{rel})}_\mu\big(\sqrt{-\det\bar{\Sigma}}\,X^\mu\big) = \partial_\mu\big(\sqrt{-\det\bar{\Sigma}}\,X^\mu\big)$, integration by parts yields (componentwise in i) the Euler–Lagrange equations displayed in eq. (6.253):

$$\nabla^{(\mathrm{rel})}_\mu\Big(\sqrt{-\det\bar{\Sigma}}\;\bar{\Sigma}^{\mu\nu}\,\mathrm{Tr}\,\hat{\Pi}_{\nu i}\Big) - \sqrt{-\det\bar{\Sigma}}\;\frac{\delta\,\mathrm{Tr}\,\mathcal{V}_\Phi}{\delta\hat{\Phi}^{i\dagger}} = 0,$$

with $\hat{\Pi}_{\nu i}$ from eq. (6.252). Each ingredient transforms covariantly (Proposition 6.188), Tr is class-function–valued, and $\nabla^{(\mathrm{rel})}$ is compatible with $\bar{\Sigma}$, so the equations are covariant.

Step 3: Holosymmetric reduction preserves covariance. Replace the branchwise covariant derivatives by the reduced combination in eq. (6.263). The reduced kinetic density is a *linear real* combination of: (i) two intra-branch scalars (each already covariant), and (ii) the real part of $e^{-i\varphi}(\cos\theta\,\mathcal{S}_{\mu\nu} + \sin\theta\,\mathcal{C}_{\mu\nu})$, where $\mathcal{S}_{\mu\nu}$ and $\mathcal{C}_{\mu\nu}$ are built from H- and A-pairings of covariant objects. By the same $U(n)\times O(r)$ argument as in Step 1 and the invariance of Tr, the reduced kinetic term remains a scalar; hence the same integration-by-parts step gives covariant E–L equations. This establishes the first claim.

Step 4: Cone assignment is unaffected by HS reduction. The causal cones used operationally are determined by the effective Lorentzian metric $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ obtained from $\bar{\Sigma}_{\mu\nu}$ via Definition 5.77 (cf. Chapters 4–5). In the present construction, $\bar{\Sigma}$ (hence $\bar{g}^{\mathrm{Lor}}$) is *external* to the HS reduction: the dial/reduction modifies only the *interference amplitudes* inside the kinetic density (eq. (6.263)) and leaves $\bar{\Sigma}$ and its determinant—and therefore the measure and cone assignment—unchanged (see also the dial–neutrality statements in Definition 6.171 and the causality remarks Definition 5.77 and related notes on cone invariance). Consequently, HS reduction does not alter $\bar{g}^{\mathrm{Lor}}$ nor the associated light-cone structure.

Conclusion. Under $\nabla^{(\mathrm{rel})}_\mu\bar{\Sigma}_{\alpha\beta} = 0$ and $\nabla^{(T)}_\mu g_T = 0$, substituting eqs. (6.259) and (6.263) into the Lagrangian of Section 6.5.2 yields covariant Euler–Lagrange equations, and the holosymmetric reduction modulates only kinetic interference terms without affecting the effective cone structure determined solely by $\bar{g}^{\mathrm{Lor}}$. □

6.6 The Operator Bundle Connection as an Emergent Gauge Field

In our framework, the internal quantum state space—namely, the Hilbert space fiber $\mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}}$—is not globally constant, but can vary from point to point across the emergent manifold $\mathcal{X}_q$. To compare field values at different locations, we must introduce a structure that governs the local variation of the quantum reference frame [375, 248]. This structure is the *Operator Bundle Connection*, $\mathcal{A}_\mu(x)$: a gauge field that prescribes how quantum information is parallel transported [123, 378].

Definition 6.192 (Operator Bundle Connection $\mathcal{A}_\mu(x)$)**.** Let $\mathcal{H}^{\text{modal}}$ be the emergent Hilbert bundle of dialectic modes (Definition 4.239). The operator-valued connection $\mathcal{A}_\mu(x)$ is a smooth field of Hermitian operators, each acting on the local fiber:

$$\boxed{\mathcal{A}_\mu(x) \in \mathfrak{u}\left(\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}\right) \subset \text{End}\left(\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)}\right),} \tag{6.264}$$

and transforming under local gauge transformations $U(x)$ according to the standard inhomogeneous law for non-Abelian potentials [123, 82]:

$$\mathcal{A}_\mu(x) \mapsto U(x)\mathcal{A}_\mu(x)U^\dagger(x) + i\,U(x)\partial_\mu U^\dagger(x). \tag{6.265}$$

In a fixed orthonormal basis $\{|\alpha\rangle_x\}_{\alpha\in\{0,1,p,s\}}$ of the four-mode modal Hilbert space (eq. (6.226)), the connection is represented as a Hermitian matrix:

$$\mathcal{A}_\mu(x) \longleftrightarrow \begin{pmatrix} A_\mu^{00}(x) & A_\mu^{01}(x) & A_\mu^{0p}(x) & A_\mu^{0s}(x) \\ A_\mu^{10}(x) & A_\mu^{11}(x) & A_\mu^{1p}(x) & A_\mu^{1s}(x) \\ A_\mu^{p0}(x) & A_\mu^{p1}(x) & A_\mu^{pp}(x) & A_\mu^{ps}(x) \\ A_\mu^{s0}(x) & A_\mu^{s1}(x) & A_\mu^{sp}(x) & A_\mu^{ss}(x) \end{pmatrix}, \qquad (A_\mu^{\alpha\beta})^\dagger = A_\mu^{\beta\alpha}. \tag{6.266}$$

6.6.1 Holosymmetric Operator Bundle Connection as an Emergent Gauge Field

Notation. We write $\mathcal{A}_\mu$ for the operator–bundle (modal) connection; this is the same object denoted A_μ in Definition 6.169, with curvature

$$\mathcal{F}_{\mu\nu} = \partial_{[\mu}\mathcal{A}_{\nu]} + [\mathcal{A}_\mu, \mathcal{A}_\nu].$$

Definition 6.193 (Holosymmetric lift of the connection and reduced connection (no new symbols))**.** For two law branches $k \in \{0,1\}$ with (possibly) distinct background pairs $\left(\bar{\Sigma}^{(k)}, \mathcal{A}_\mu^{(k)}\right)$, define on $\mathcal{H}_L \otimes \mathcal{H}_{\text{mod}}$ the lifted connection

$$\boxed{\widehat{\mathcal{A}}_\mu^{\text{HS}} := |0\rangle\langle 0|_L \otimes \mathcal{A}_\mu^{(0)} + |1\rangle\langle 1|_L \otimes \mathcal{A}_\mu^{(1)} + |0\rangle\langle 1|_L \otimes \mathcal{M}_\mu^{(\theta)} + |1\rangle\langle 0|_L \otimes \left(\mathcal{M}_\mu^{(\theta)}\right)^\dagger,} \tag{6.267}$$

$$\boxed{\mathcal{F}_{\mu\nu}^{\text{HS}} := \partial_{[\mu}\widehat{\mathcal{A}}_{\nu]}^{\text{HS}} + \left[\widehat{\mathcal{A}}_\mu^{\text{HS}}, \widehat{\mathcal{A}}_\nu^{\text{HS}}\right].} \tag{6.268}$$

Here the *mixing functional* is built *only* from the branch data,

$$\boxed{\mathcal{M}_\mu^{(\theta)} := \cos\theta\, \mathcal{S}_\mu\left[\mathcal{A}^{(0)}, \mathcal{A}^{(1)}\right] + \sin\theta\, \mathcal{C}_\mu\left[\mathcal{A}^{(0)}, \mathcal{A}^{(1)}\right], \qquad \begin{array}{l} \mathcal{S}_\mu[X,Y] := \frac{1}{2}\{X_\mu, Y_\mu\}, \\ \mathcal{C}_\mu[X,Y] := \frac{i}{2}[X_\mu, Y_\mu], \end{array}} \tag{6.269}$$

Preparing/reading the law register in $|+_\varphi\rangle_L := \frac{1}{\sqrt{2}}\big(|0\rangle_L + e^{i\varphi}|1\rangle_L\big)$ yields the reduced (operational) connection

$$\boxed{\mathcal{A}_\mu^{\langle \mathrm{HS}\rangle} = \tfrac{1}{2}\big(\mathcal{A}_\mu^{(0)} + \mathcal{A}_\mu^{(1)}\big) + \mathrm{Re}\big(e^{-i\varphi}\,\mathcal{M}_\mu^{(\theta)}\big), \qquad \mathcal{F}_{\mu\nu}^{\langle \mathrm{HS}\rangle} = \mathcal{F}\big[\mathcal{A}^{\langle \mathrm{HS}\rangle}\big]_{\mu\nu}.} \tag{6.270}$$

The pair (φ, θ) tomographically separates phase–even (anticommutator) and phase–odd (commutator) mixings, consistent with the HS law–control pattern of Definition 6.171.

Proposition 6.194 (Gauge covariance and branchwise microcausality)**.** *Under a modal gauge $U(x)$,*

$$\mathcal{A}_\mu^{(k)} \mapsto U\mathcal{A}_\mu^{(k)}U^\dagger - (\partial_\mu U)\,U^\dagger, \qquad \mathcal{S}_\mu, \mathcal{C}_\mu,\ \mathcal{M}_\mu^{(\theta)} \ \mapsto\ U(\cdot)U^\dagger,$$

so $\widehat{\mathcal{A}}_\mu^{\mathrm{HS}}$ and $\mathcal{F}_{\mu\nu}^{\mathrm{HS}}$ transform covariantly, and the reduced objects in eq. (6.270) obey

$$\mathcal{A}_\mu^{\langle \mathrm{HS}\rangle} \ \mapsto\ U\,\mathcal{A}_\mu^{\langle \mathrm{HS}\rangle}\,U^\dagger - (\partial_\mu U)\,U^\dagger, \qquad \mathcal{F}_{\mu\nu}^{\langle \mathrm{HS}\rangle} \ \mapsto\ U\,\mathcal{F}_{\mu\nu}^{\langle \mathrm{HS}\rangle}\,U^\dagger.$$

Holosymmetric phases (φ, θ) modulate only interference amplitudes (adjoint–covariant pieces); the causal cone is unchanged, being determined by $\bar{g}^{\mathrm{Lor}}$ (cf. Definition 5.77 and the compatibility results surrounding Definition 6.185).

Proof. **(1) Lifted connection transforms as a connection.** By Definition 6.169, a modal gauge $U(x)$ acts on each branch connection as $\mathcal{A}_\mu^{(k)} \mapsto U\mathcal{A}_\mu^{(k)}U^\dagger - (\partial_\mu U)\,U^\dagger$. By construction of the mixing functional in eq. (6.269), we take $\mathcal{M}_\mu^{(\theta)} \mapsto U\,\mathcal{M}_\mu^{(\theta)}\,U^\dagger$ (adjoint covariance). Insert these into the lifted block form (6.267). Writing $\mathbf{U} := \mathbf{1}_L \otimes U$ and $Q_\mu := (\partial_\mu U)U^\dagger$, one finds

$$\widehat{\mathcal{A}}_\mu^{\mathrm{HS}} \ \longmapsto\ \mathbf{U}\,\widehat{\mathcal{A}}_\mu^{\mathrm{HS}}\,\mathbf{U}^\dagger \ -\ \mathbf{1}_L \otimes Q_\mu,$$

i.e. the affine connection law holds fibrewise on $\mathcal{H}_L \otimes \mathcal{H}_{\mathrm{mod}}$ with the same Q_μ on every law block. This proves the first statement.

(2) Curvature covariance. Using (6.268) and the Leibniz rule, the standard gauge–theory identity yields

$$\mathcal{F}_{\mu\nu}^{\mathrm{HS}} = \partial_{[\mu}\widehat{\mathcal{A}}_{\nu]}^{\mathrm{HS}} + \big[\widehat{\mathcal{A}}_\mu^{\mathrm{HS}}, \widehat{\mathcal{A}}_\nu^{\mathrm{HS}}\big] \ \longmapsto\ \mathbf{U}\,\mathcal{F}_{\mu\nu}^{\mathrm{HS}}\,\mathbf{U}^\dagger,$$

since the affine pieces $-\mathbf{1}_L \otimes Q_\mu$ cancel in $\partial_{[\mu}(\cdot)_{\nu]} + [\cdot,\cdot]$ (cf. the curvature transformation in Definition 6.185).

(3) Reduced connection and curvature. Let $\mathcal{C}_\varphi(\cdot) := \langle +_\varphi|\,(\cdot)\,|+_\varphi\rangle_L$ denote compression on the law register. Because $\mathcal{C}_\varphi$ is linear and $\mathbf{U} = \mathbf{1}_L \otimes U$ commutes with the law-space compression,

$$\mathcal{A}_\mu^{\langle \mathrm{HS}\rangle} = \mathcal{C}_\varphi\big(\widehat{\mathcal{A}}_\mu^{\mathrm{HS}}\big) \ \longmapsto\ \mathcal{C}_\varphi\big(\mathbf{U}\,\widehat{\mathcal{A}}_\mu^{\mathrm{HS}}\,\mathbf{U}^\dagger - \mathbf{1}_L \otimes Q_\mu\big) = U\,\mathcal{A}_\mu^{\langle \mathrm{HS}\rangle}\,U^\dagger - Q_\mu,$$

which is precisely the connection law stated in (6.194). Applying the same compression to the HS curvature and using part (2) gives $\mathcal{F}_{\mu\nu}^{\langle \mathrm{HS}\rangle} = \mathcal{F}[\mathcal{A}^{\langle \mathrm{HS}\rangle}]_{\mu\nu} \mapsto U\,\mathcal{F}_{\mu\nu}^{\langle \mathrm{HS}\rangle}\,U^\dagger$, as in (6.270).

(4) Branchwise microcausality / cone invariance. Holosymmetric phases (φ, θ) enter only through the adjoint–covariant mixing functional $\mathcal{M}_\mu^{(\theta)}$ and thus modulate interference

amplitudes in the connection blocks; they do not alter the effective Lorentzian structure $\bar{g}^{\mathrm{Lor}}$ which is defined from the unified tensor $\bar{\Sigma}$ by Definition 5.77. In particular, the causal cone is fixed by $\bar{g}^{\mathrm{Lor}}$ and hence remains invariant under the above gauge/HS reductions. This is the same mechanism established for the action–level construction in Section 6.5.2 and formalized in Proposition 6.191, where holosymmetric reduction affects only interference terms in kinetic densities but not the cone assignment. Therefore, branchwise microcausality (cone structure) is unchanged.

Combining (1)–(4) proves the proposition. □

Remark 6.195 (Sectoral representations and universality)**.** Via the sectoral functors $(\mathcal{R}_\Phi, \mathcal{R}_\Theta)$ (Gaiamiti/Tantrakyma), a single modal connection $\mathcal{A}_\mu$ induces

$$\nabla_\mu^{(\Phi)} = \partial_\mu + [\mathcal{R}_\Phi(\mathcal{A}_\mu), \cdot], \qquad \nabla_\mu^{(\Theta)} = \partial_\mu + [\mathcal{R}_\Theta(\mathcal{A}_\mu), \cdot],$$

consistent with Definition 6.185. Wordwise gauge–invariant scalars are built from $\mathrm{Tr}\{\mathcal{F}_{\mu\nu}\mathcal{F}^{\mu\nu}\}$ and higher Chern words (cf. eq. (6.262)).

Corollary 6.196 (Canonical holosymmetric pair)**.** *If only the temporal sector flips between branches while the represented internal algebra is fixed, take $\mathcal{A}_\mu^{(0)} = \mathcal{A}_\mu^{(1)} = \mathcal{A}_\mu$. Then*

$$\boxed{\mathcal{A}_\mu^{\langle \mathrm{HS}\rangle} = \mathcal{A}_\mu + \mathrm{Re}\Big(e^{-i\varphi}\,\mathcal{M}_\mu^{(\theta)}[\mathcal{A},\mathcal{A}]\Big), \qquad \mathcal{M}_\mu^{(\theta)}[\mathcal{A},\mathcal{A}] = \cos\theta\,\tfrac{1}{2}\{\mathcal{A}_\mu,\mathcal{A}_\mu\},} \tag{6.271}$$

and, to leading order in the mixing functional $\mathcal{M}^{(\theta)}$,

$$\mathcal{F}_{\mu\nu}^{\langle \mathrm{HS}\rangle} = \mathcal{F}_{\mu\nu} + \nabla_{[\mu}^{(\mathcal{A})}\,\mathrm{Re}\Big(e^{-i\varphi}\mathcal{M}_{\nu]}^{(\theta)}\Big) + O\Big((\mathcal{M}^{(\theta)})^2\Big),$$

so (φ,θ) performs commutator/anticommutator tomography without altering the cone assignments determined by $\bar{g}^{\mathrm{Lor}}$.

Proof. Set the two branches equal in the canonical case: $\mathcal{A}_\mu^{(0)} = \mathcal{A}_\mu^{(1)} = \mathcal{A}_\mu$. By the definition of the mixing functional in eq. (6.269),

$$\mathcal{M}_\mu^{(\theta)}[\mathcal{A},\mathcal{A}] = \cos\theta\,\mathcal{S}_\mu[\mathcal{A},\mathcal{A}] + \sin\theta\,\mathcal{C}_\mu[\mathcal{A},\mathcal{A}] = \cos\theta\,\tfrac{1}{2}\{\mathcal{A}_\mu,\mathcal{A}_\mu\} + \sin\theta\,\tfrac{i}{2}[\mathcal{A}_\mu,\mathcal{A}_\mu].$$

Since $[\mathcal{A}_\mu,\mathcal{A}_\mu] = 0$, the commutator piece vanishes and we obtain $\mathcal{M}_\mu^{(\theta)}[\mathcal{A},\mathcal{A}] = \cos\theta\,\frac{1}{2}\{\mathcal{A}_\mu,\mathcal{A}_\mu\}$, as stated in (6.271). Inserting this identity into the reduced-connection formula (6.270) immediately gives

$$\mathcal{A}_\mu^{\langle \mathrm{HS}\rangle} = \tfrac{1}{2}(\mathcal{A}_\mu + \mathcal{A}_\mu) + \mathrm{Re}\Big(e^{-i\varphi}\,\mathcal{M}_\mu^{(\theta)}[\mathcal{A},\mathcal{A}]\Big) = \mathcal{A}_\mu + \mathrm{Re}\Big(e^{-i\varphi}\,\mathcal{M}_\mu^{(\theta)}[\mathcal{A},\mathcal{A}]\Big),$$

which is the first boxed relation in (6.271).

For the curvature, write the reduced connection as a background plus a small HS perturbation $\Delta\mathcal{A}_\mu := \mathrm{Re}\Big(e^{-i\varphi}\,\mathcal{M}_\mu^{(\theta)}[\mathcal{A},\mathcal{A}]\Big)$, so $\mathcal{A}_\mu^{\langle \mathrm{HS}\rangle} = \mathcal{A}_\mu + \Delta\mathcal{A}_\mu$. Using the curvature definition $\mathcal{F}_{\mu\nu} = \partial_{[\mu}\mathcal{A}_{\nu]} + [\mathcal{A}_\mu,\mathcal{A}_\nu]$ from Definition 6.169 and eq. (6.262), the standard expansion gives

$$\mathcal{F}[\mathcal{A}+\Delta\mathcal{A}]_{\mu\nu} = \mathcal{F}[\mathcal{A}]_{\mu\nu} + \nabla_{[\mu}^{(\mathcal{A})}\Delta\mathcal{A}_{\nu]} + [\Delta\mathcal{A}_\mu,\Delta\mathcal{A}_\nu],$$

where $\nabla^{(\mathcal{A})}_{\mu} X := \partial_\mu X + [\mathcal{A}_\mu, X]$ is the adjoint covariant derivative (cf. Definition 6.185). Keeping terms to leading order in the mixing functional and substituting $\Delta\mathcal{A}_\mu = \mathrm{Re}\big(e^{-i\varphi}\,\mathcal{M}^{(\theta)}_\mu\big)$ yields

$$\mathcal{F}^{\langle \mathrm{HS}\rangle}_{\mu\nu} = \mathcal{F}_{\mu\nu} + \nabla^{(\mathcal{A})}_{[\mu}\,\mathrm{Re}\big(e^{-i\varphi}\,\mathcal{M}^{(\theta)}_{\nu]}\big) + O\big((\mathcal{M}^{(\theta)})^2\big),$$

which is the second claimed relation.

Finally, (φ, θ) enter only through $\mathcal{M}^{(\theta)}_\mu$ and hence modulate adjoint–covariant interference amplitudes in the connection sector; they do not alter the effective Lorentzian structure $\bar{g}^{\mathrm{Lor}}$ that fixes causal cones (see Definition 5.77). This matches the action-level compatibility statement in Proposition 6.191, where holosymmetric reduction affects kinetic mixings but leaves cone assignments unchanged. The corollary follows. □

Remark 6.197 (Topological charges (Chern meaning, invariance, and HS–phase insensitivity))**.** Let $\mathcal{A}_\mu$ be the modal (operator–bundle) connection with curvature $\mathcal{F} = \frac{1}{2}\,\mathcal{F}_{\mu\nu}\,dx^\mu \wedge dx^\nu$ as in Definition 6.169 and eq. (6.262). The *Chern–Weil* "Chern words" are the even-degree differential forms

$$\boxed{\mathrm{Ch}_{2m}(\mathcal{A}) \;:=\; \mathrm{Tr}\Big(\,\mathcal{F}^{\wedge m}\Big) \qquad (m = 1, 2, \dots),} \tag{6.272}$$

whose cohomology classes are *gauge invariant* and depend only on the bundle isomorphism class (Chern–Weil theory) [248, 375, 225]. On a compact oriented $2m$–cycle $\Omega^{2m} \subset \mathcal{X}_q$, the associated *Chern numbers* are

$$\boxed{\begin{aligned} Q_{2m}[\mathcal{A};\Omega] &:= \mathcal{N}_m \int_{\Omega^{2m}} \mathrm{Tr}\big(\mathcal{F}^{\wedge m}\big) \in \mathbb{Z}, \\ \mathcal{N}_m &:= \frac{1}{(2\pi)^m} \quad \text{(up to the chosen convention).} \end{aligned}} \tag{6.273}$$

and classify holonomy sectors of the modal bundle (principal $U(n)$–bundle), i.e. they distinguish topologically inequivalent gauge backgrounds by their global winding/instanton data [248, 375, 225].

Gauge covariance and exact–form shift. Under the (adjoint/affine) gauge law of Definition 6.169 and Proposition 6.194, $\mathcal{F} \mapsto U\mathcal{F}U^\dagger$, hence Ch_{2m} is invariant. Infinitesimally, $\delta\mathrm{Tr}(\mathcal{F}^{\wedge m}) = d\big(m\,\mathrm{Tr}(\delta\mathcal{A} \wedge \mathcal{F}^{\wedge(m-1)})\big)$, so any change is an *exact* form and the integral (6.273) is unchanged on closed cycles [248, 375].

Examples. For $m = 1$, $\mathrm{Ch}_2(\mathcal{A}) = \mathrm{Tr}\,\mathcal{F}$ integrates (up to normalization) to the first Chern number. In four dimensions ($m = 2$),

$$\boxed{\mathrm{Ch}_4(\mathcal{A}) \;=\; \mathrm{Tr}\,\big(\mathcal{F} \wedge \mathcal{F}\big), \qquad Q_4[\mathcal{A}] = \mathcal{N}_2 \int \mathrm{Tr}\,(\mathcal{F} \wedge \mathcal{F}),} \tag{6.274}$$

which measures the (anti)self–dual instanton number in the modal sector [375, 225]. On manifolds with boundary, $\mathrm{Tr}(\mathcal{F} \wedge \mathcal{F}) = d\,\mathrm{CS}_3(\mathcal{A})$ relates the bulk invariant to a boundary Chern–Simons word, giving anomaly inflow–type boundary terms (the volume element arises from $\bar{g}^{\mathrm{Lor}}$ via Definition 6.33) [434, 375, 435, 7, 84].

Holosymmetric phases do not change Chern data. Consider the HS–reduced connection $\mathcal{A}_\mu^{\langle \mathrm{HS}\rangle}$ of eq. (6.270) with curvature $\mathcal{F}^{\langle \mathrm{HS}\rangle} = \mathcal{F}[\mathcal{A}^{\langle \mathrm{HS}\rangle}]$. By Corollary 6.196 (and the general expansion behind it),

$$\mathcal{F}^{\langle \mathrm{HS}\rangle} = \mathcal{F} \;+\; \nabla^{(\mathcal{A})}\Big(\mathrm{Re}[e^{-i\varphi}\mathcal{M}^{(\theta)}]\Big) \;+\; O\Big((\mathcal{M}^{(\theta)})^2\Big),$$

so $\mathrm{Tr}\Big((\mathcal{F}^{\langle \mathrm{HS}\rangle})^{\wedge m}\Big) = \mathrm{Tr}(\mathcal{F}^{\wedge m}) + d(\cdots)$ differs from the branch expression by an *exact* form (up to higher-order total derivatives). Therefore the Chern numbers (6.273) are unchanged by (φ, θ). Equivalently, working at the HS–lift level, the block–off–diagonal mixing in $\widehat{\mathcal{A}}_\mu^{\mathrm{HS}}$ of eq. (6.267) contributes only adjoint–covariant pieces that drop out of the cyclic trace in $\mathrm{Tr}(\mathcal{F}^{\wedge m})$ up to exact forms; the net integral reduces to the sum of *branchwise* contributions $\mathrm{Tr}\Big((\mathcal{F}^{(k)})^{\wedge m}\Big)$ (see cyclicity of the trace in operator algebras and its geometric avatars) [404, 4, 375].

Physical role. Chern numbers label superselection (holonomy) sectors for the modal bundle and constrain global dynamics: they are invariant under smooth, spectrally regulated deformations of the operator data and under the HS law–control dial (Definition 6.171, Proposition 6.194, and Corollary 6.196). Their densities use the volume form determined by $\bar{g}^{\mathrm{Lor}}$ (Definition 6.33); hence causal–cone assignments (from $\bar{g}^{\mathrm{Lor}}$) are compatible with and independent of these topological charges (cf. Proposition 6.191) [7, 84].

Emergent Origin of the Connection. In keeping with the foundational spirit of gauge theory, the operator-valued connection $\mathcal{A}_\mu(x)$ is not imposed externally but emerges intrinsically from the Algebra of Observables $\mathcal{A}$ (Definition 1.32) [230, 404]. Through the GNS construction (Definition 1.34), the abstract algebra is represented as concrete operators on $\mathcal{H}_{\mathrm{em}}$ [254], yielding:

$$\mathcal{A}_\mu(x) \in \pi^{\mathring{\mathbf{\Psi}}}(\mathcal{A}) \subseteq \mathrm{End}(\mathcal{H}_{\mathrm{em}}).$$

Thus, the Operator Bundle Connection is a *manifestation* of the internal symmetry content of the theory, as realized in the physical vacuum.

Remark 6.198 (The Connection as an Element of the Represented Algebra)**.** The connection field $\mathcal{A}_\mu(x)$ belongs to the represented subalgebra of observables [230, 381]:

$$\mathcal{A}_\mu(x) \in \pi^{\mathring{\mathbf{\Psi}}}(\mathcal{A}),$$

where $\mathcal{A}$ is the abstract observable algebra, and $\pi^{\mathring{\mathbf{\Psi}}}$ is the GNS representation associated to the Quantum Holo-State. Hence, $\mathcal{A}_\mu(x)$ acts as an auxiliary gauge field as well as a fundamental object arising from the same algebraic source as the fields it governs [4, 224].

Remark 6.199 (Dual Role of the Operator Bundle Connection)**.** Although $\mathcal{A}_\mu(x)$ is a mathematically universal object, its physical interpretation is context-dependent. In the Gaiamiti sector, it governs the parallel transport of geometric (vacuum-deformation) modes. In the Tantrakyma sector, it governs internal excitations and defines the local gauge structure associated with matter fields. It is thus the unifying transport field for both relational geometry and internal structure.

In this way, the covariant structure of the theory is not appended to the geometry—it is derived from the same fundamental algebra that generates the Hilbert space, its endomorphisms, and all dynamical fields.

Remark 6.200 (Universality of the operator bundle connection)**.** The same operator bundle connection $\mathcal{A}_\mu$ governs both substrate fields. For the Gaiamiti field Φ it enters the covariant derivative (cf. Section 6.5.3), while for the Tantrakyma field Θ it appears in Definition 9.323. In both cases $\mathcal{A}_\mu$ ensures local fibre covariance of the operator Hilbert bundle, irrespective of the sector. Thus $\mathcal{A}_\mu$ is a *universal modal gauge connection* of the theory, not confined to a single field.

Remark 6.201 (Sectoral quanta of the universal connection (field-corrected))**.** The universal operator bundle connection $\mathcal{A}_\mu$ acts on the substrate fields via their sectoral representations on the modal fibers. With the field symbols,

$$\nabla_\mu^{(\Phi)}\hat{\mathbf{\Phi}} := \partial_\mu\hat{\mathbf{\Phi}} + \mathcal{R}_\Phi(\mathcal{A}_\mu)\,\hat{\mathbf{\Phi}}, \qquad \nabla_\mu^{(\Theta)}\hat{\mathbf{\Theta}} := \partial_\mu\hat{\mathbf{\Theta}} + \mathcal{R}_\Theta(\mathcal{A}_\mu)\,\hat{\mathbf{\Theta}},$$

where $\mathcal{R}_\Phi$ and $\mathcal{R}_\Theta$ denote the sectoral representation maps of $\mathcal{A}_\mu$ (cf. Section 6.5.3 and Definition 9.323). By convention, the gauge quantum of $\mathcal{R}_\Phi(\mathcal{A}_\mu)$ is the *mition* and that of $\mathcal{R}_\Theta(\mathcal{A}_\mu)$ is the *tantron* (cf. Definitions 9.258 and 9.348). Their physical one-particle content is read in the typed, transverse vector channel and then realized in the BRST-physical sector $H^0(\hat{Q}_{\mathrm{BRST}})$ (cf. Section 9.5). On a certified plateau $\mathfrak{P}$, spontaneous symmetry breaking endows these connection excitations with Brout–Englert–Higgs masses governed by the sectoral operators

$$(M_\Phi^2)_{IJ} = g^2\, g_T\Big(\mathcal{R}_\Phi(T_I)\,\bar{\Phi},\ \mathcal{R}_\Phi(T_J)\,\bar{\Phi}\Big), \qquad (M_\Theta^2)_{IJ} = g^2\, g_C\Big(\mathcal{R}_\Theta(T_I)\,\bar{\Theta},\ \mathcal{R}_\Theta(T_J)\,\bar{\Theta}\Big),$$

whose eigenmodes/poles determine the transverse physical polarizations identified as the *mition* and *tantron*, respectively (see Chapter 10, Definitions 9.258 and 9.348). Thus, $\mathcal{A}_\mu$ is the unifying transport field for both relational geometry and internal structure, with its sectoral faces realized as the mition (Gaiamiti) and tantron (Tantrakyma).

Remark 6.202 (Channel-resolved diagnostics for $\mathcal{A}_\mu$)**.** The holosymmetric visibilities (Definition 6.275) with triadic channel projectors $\Pi_\alpha^{\mathrm{corr/vac/spec}}$ apply verbatim to $\mathcal{A}_\mu$ and its sectoral images. In practice: (i) Π_{corr} tests the Relational Ward structures (Lemma 6.233); (ii) Π_{vac} probes background order relevant for symmetry breaking; (iii) Π_{spec} isolates transverse vector poles/residues that identify the physical mition/tantron modes (cf. realization in Chapter 10, Section 9.5).

Remark 6.203 (Plateau license for naming connection quanta)**.** The quadrant decision rule of Chapter 7 is applied to the channel-resolved $\mathcal{A}_\mu$ readouts *only* on certified plateaus (Section 6.12.4). This ensures that the identification of a feature as "mition" or "tantron" is stable under $(\lambda_{\mathrm{UV}}, \ell)$ drifts and inherits gauge-parameter independence after projection to $\mathcal{H}_{\mathrm{phys}}$ (cf. Chapter 10: BRST chain-map and descent, Lemma 9.54).

Remark 6.204 (BRST/band compatibility and physical descent)**.** On the UV/IR band defined in Chapter 8 (Definition 6.282), the band tag commutes with the BRST differential; hence connection-based diagnostics are BRST-clean and descend to cohomology classes $H^0(\hat{Q}_{\mathrm{BRST}})$. Consequently, the mition/tantron poles extracted in Π_{spec} are Nielsen-safe on plateaus (cf. Chapter 10, Lemma 9.54 and Section 9.5).

Remark 6.205 (SSB and BEH mass operators for $\mathcal{A}_\mu$ (field-corrected))**.** On a certified plateau $\mathfrak{P}$ (cf. Section 6.12.4), a nonzero sectoral order parameter breaks directions in $\mathfrak{u}\big(\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}\big)$ and endows the sectoral images of the universal connection with Brout–Englert–Higgs masses. Using the field symbols of Chapter 7 and the sectoral representation maps $\mathcal{R}_\Phi$ and $\mathcal{R}_\Theta$, the typed mass matrices are

$$(M_\Phi^2)_{IJ} = g^2\, g_T\Big(\mathcal{R}_\Phi(T_I)\,\bar\Phi,\ \mathcal{R}_\Phi(T_J)\,\bar\Phi\Big), \qquad (M_\Theta^2)_{IJ} = g^2\, g_C\Big(\mathcal{R}_\Theta(T_I)\,\bar\Theta,\ \mathcal{R}_\Theta(T_J)\,\bar\Theta\Big),$$

with T_I, T_J the broken generators. Their eigenmodes, isolated by the typed, transverse vector projectors and read in the spectral channel Π_{spec}, appear as physical poles/residues corresponding to the *mition* and *tantron*, respectively (cf. Chapter 10, Definitions 9.258 and 9.348). Gauge-parameter independence and cohomological descent of these determinations follow from the band/BRST chain-map and the Σ-causal realization (cf. Chapter 10, Lemma 9.54 and Section 9.5).

Remark 6.206 (Basis independence of physical content)**.** Although $\mathcal{A}_\mu$ is represented in a local modal basis $\{0, 1, p, s\}$ (cf. Definition 6.192), the realized one-particle content (mition/tantron) depends only on the cohomology class of the associated band-tagged operators and is invariant under local changes of basis $U(x)$; cf. Chapter 10 on Σ-causal realization and physical retarded kernels (Section 9.5).

Remark 6.207 (Quadrant classification of connection quanta)**.** After extracting the channel-restricted visibilities for $\mathcal{A}_\mu$ on a plateau, apply the canonical quadrant decision rule of Chapter 7 to the pair $\big(\Lambda_{\mathrm{law}}, \Lambda_{\mathrm{state}}\big)$ associated with the typed vector slice. The resulting QI–QIV assignment for the connection quanta matches the realized, BRST-projected diagnosis of Chapter 10 (cf. Section 9.5).

6.7 Spontaneous Symmetry Breaking

Spontaneous symmetry breaking (SSB) is the precise way in which a **symmetric law** gives rise to an **asymmetric world**. A symmetry of the law is a transformation that leaves the action and equations of motion invariant; by Noether's theorem, each continuous symmetry implies a conserved current and charge J^μ [90]. SSB occurs when the Hamiltonian/Lagrangian enjoys a symmetry group G, but the realized ground state—selected in the thermodynamic or infinite-volume limit—fails to share it. The hallmark is a *manifold of degenerate vacua* related by G; choosing one vacuum "hides'' part of the symmetry without altering the underlying equations [91, 82, 393]. In field-theoretic language, if the unbroken subgroup is $H \subset G$, the vacuum manifold is the coset G/H. Correlation functions and response coefficients reorganize according to H, and the broken generators act nontrivially on the vacuum.

The local physics of SSB is controlled by the geometry of that vacuum manifold. Consider the canonical relativistic prototype

$$\mathcal{L} = |\partial\phi|^2 - \lambda\Big(|\phi|^2 - v^2\Big)^2,$$

invariant under the global $U(1)$ rotation $\phi \to e^{i\alpha}\phi$. The classical minima form a circle $|\phi| = v$. Writing $\phi(x) = \frac{1}{\sqrt{2}}(v + h(x) + i\,\pi(x))$, the radial mode h is gapped while the

angular mode π is gapless. This gapless field is the **Nambu–Goldstone (NG) mode**, and its existence reflects the fact that moving along the valley of minima costs arbitrarily little energy [92, 93, 94, 91]. In relativistic theories, the number of NG bosons equals $\dim(G/H)$; equivalently, there is one massless excitation per broken generator. Their derivative couplings dominate low-energy dynamics and enforce characteristic soft limits and Ward identities in scattering amplitudes [91, 82].

The distinction between **laws** and **states** is operational as well as conceptual. In a symmetric (unbroken) phase the vacuum is unique, symmetry charges annihilate it, and correlation functions respect the full group G. In a broken phase, the symmetry generators fail to annihilate the vacuum; acting with the corresponding currents creates NG quanta. Observables become anisotropic or otherwise ordered (magnetization in a ferromagnet; crystalline density modulations in a solid), yet the *structure* of those observables—selection rules, soft theorems, current algebra—is still governed by the symmetric law [102, 101, 91]. Physically, SSB trades manifest invariance for long-range organization: algebraic constraints from symmetry reappear as constraints on the low-energy spectrum and on the pattern of correlations.

Gauge theories require a special word. Local "gauge symmetry'' is a redundancy of description, so it does not literally *break* in the same sense as a global symmetry. Nevertheless, in phases where a gauge-covariant order parameter would suggest breaking, the **Brout–Englert–Higgs (BEH) mechanism** operates: the would-be NG modes combine with gauge fields, giving the latter a mass and a longitudinal polarization, while leaving behind a massive scalar associated with radial fluctuations [95, 96, 97, 98, 99, 91, 82]. The correct diagnostics are gauge-invariant: screened charges, massive vector excitations, and characteristic patterns of interactions. This gauge-theory realization is continuous with the global case at the level of the vacuum manifold and excitations, but the massless NG bosons are "eaten'' and do not appear as asymptotic particles.

SSB is also constrained by dimension and temperature. In low dimensions, long-wavelength fluctuations are so strong that continuous symmetries cannot support true long-range order at finite temperature in 1D and 2D spin systems, and in relativistic 1+1D quantum field theory even at zero temperature, forcing qualitatively different infrared behavior [100, 101]. Where SSB *is allowed* (higher dimensions, appropriate limits), it becomes sharp only in the thermodynamic limit: finite systems exhibit extremely long but finite coherence times rather than mathematically exact vacuum degeneracy, which is why analyses take large-volume limits before probing the order parameter [91, 101]. Within its domain of applicability, SSB is abundantly realized—ferromagnets and antiferromagnets, superfluids and superconductors (with their phase stiffness and Meissner response), crystals and charge-density waves—and its signatures (order parameters, collective modes, soft theorems) are quantitatively verified [102, 101, 82].

Finally, the renormalization-group perspective unifies these facts. Symmetry dictates the relevant operators near a fixed point, while SSB describes which vacuum around that fixed point the system selects. The low-energy effective action is built from NG fields parametrizing G/H with derivative interactions, systematically capturing universal infrared physics independent of microscopic details [389, 91, 393]. In this sense, SSB is not just a mechanism; it is the organizing principle by which symmetric laws generate ordered, asymmetric worlds—predicting *what must be gapless*, *how it must couple*, and *how observables*

must transform whenever a continuous global symmetry is hidden by the vacuum.

6.8 Emergent (Asymmetry-Induced) Symmetry

How Asymmetric Micro-Laws Yield Symmetric Macro-Worlds

Concept. *Emergent symmetry* is the systematic appearance, at long distances or low energies, of a symmetry larger than that present in the microscopic description. If G_{UV} is the symmetry of the short-distance (UV) "law" and $\mathcal{L}_{\mathrm{IR}}$ denotes the coarse-grained (IR) "effective theory", it often happens that $G_{\mathrm{UV}} \subsetneq H_{\mathrm{IR}}$, so observables in the IR organize into representations of H_{IR}. This is the logical reverse of spontaneous symmetry breaking (SSB): there, symmetric laws yield asymmetric states; here, **asymmetric laws** flow to a **more symmetric** effective description through coarse-graining, collective organization, and the renormalization group (RG) [82, 14, 102]. In statistical mechanics, the same idea appears as restoration of isotropy or other invariances in macroscopic averages [436].

Mechanisms. *(1) Coarse-graining and RG irrelevance (universality).* Symmetry-breaking operators can be RG-*irrelevant*. Linearizing the flow for a symmetry-violating coupling g,

$$\boxed{\frac{dg}{d\ell} = (d + z - \Delta)\, g + \cdots, \quad \Delta > d + z \;\Rightarrow\; g(\ell) \to 0,} \tag{6.275}$$

so the IR fixed point exhibits an *enhanced* symmetry that was not manifest microscopically. This explains isotropic long-wavelength phonons on anisotropic lattices, emergent Lorentz invariance near Dirac nodes, and conformal symmetry at criticality [82, 14, 102]. *(2) Accidental symmetry in effective field theory (EFT).* Low-energy dynamics are controlled by the lowest-dimension operators. If that minimal set respects a larger group, the EFT enjoys an *accidental* global symmetry; symmetry-violating effects enter only through higher-dimension operators suppressed by powers of E/Λ [14, 82, 409]. *(3) Ensemble/attractor restoration.* Far from equilibrium or in kinetic regimes, macroscopic observables can relax to *symmetric attractors* (e.g., isotropic Maxwellians in simple gases) or become symmetric *in expectation* under disorder/angle averaging. The symmetry is a property of statistics/observables, not of the micro-law itself [436]. *(4) Emergent gauge structure.* Local constraints and fractionalization in many-body systems can generate *emergent* gauge redundancies (e.g., U(1) spin liquids), so the IR description possesses gauge symmetry absent from the microscopic rules [437, 102].

Relation to SSB. Emergent symmetry and SSB address opposite arrows. Emergent symmetry enlarges the *law* in the IR; it does *not* by itself produce Nambu–Goldstone (NG) modes because no global symmetry of the governing action is being broken by the state. NG bosons appear only if the *emergent* global symmetry is subsequently *spontaneously broken* by the phase the system chooses. Otherwise one finds *approximately conserved currents* and Ward identities whose violations are controlled by irrelevant operators or EFT power counting [82, 14, 102].

Diagnostics. A system exhibits emergent symmetry when:

1. *Scaling suppression* of violations is observed: symmetry-breaking observables decrease with scale as $(E/\Lambda)^{\Delta-(d+z)}$ or as negative powers of system size—canonical RG signatures [102, 82].

2. *Anomaly consistency* holds: an *exact* emergent global symmetry must be compatible with 't Hooft anomaly constraints; otherwise only an *approximate* symmetry can emerge [82, 14, 103].

3. *Fixed-point structure* displays enhanced organization: correlators and spectra at criticality transform under the larger IR group (often conformal) [102].

Minimal toy model. Consider an *anisotropic* scalar,

$$\boxed{\mathcal{L} = \tfrac{1}{2}\left[c_x(\partial_x\phi)^2 + c_y(\partial_y\phi)^2\right] - V(\phi), \qquad c_x \neq c_y.} \tag{6.276}$$

Under RG, combined coordinate rescalings and the irrelevance of the anisotropy operator drive the kinetic term toward an *isotropic* form; microscopic asymmetry survives only as higher-dimension corrections in the IR action [102, 82].

Representative arenas.

1. *Lattice → isotropic hydrodynamics:* discrete rotational symmetries give way to continuum isotropy for sound/elastic modes; anisotropies vanish with momentum scale [102].

2. *Dirac materials:* tight-binding models lack Lorentz symmetry, yet linear dispersions and relativistic structure emerge near band-touching points [102].

3. *Critical phenomena:* disparate microscopics flow to conformal IR theories with enlarged symmetry, explaining universality of exponents/correlators [102, 82].

4. *Spin liquids and emergent gauge fields:* U(1) gauge redundancy and photon-like excitations arise from constraint structures in quantum magnets [437, 102].

5. *Cosmological isotropy:* despite microscopic CP violation, large-scale observables (e.g., the CMB) are nearly isotropic; inflationary coarse-graining yields symmetric statistics on the largest scales [438, 439].

Stacking with SSB. In practice, the two arrows frequently *stack*: a microscopically anisotropic model first *flows* to a more symmetric IR law (emergence), and the resulting IR symmetry can then be *spontaneously broken* by the phase realized. Diagnostics separate the layers: scale-vanishing violations signal emergence; nonzero order parameters and NG spectra signal subsequent SSB.

Bottom line. Emergent symmetry is a renormalization-group and effective-theory fact: scale, constraints, and averaging erase many microscopic asymmetries, revealing a higher-symmetry law that governs long-distance observables. Where exact, it must satisfy anomaly constraints; where approximate, its violations are parametrically small and measurable. It is the mirror image of SSB in the broader cartography of *laws* versus *states* [82, 14, 102, 437].

6.9 Weaving The Two Arrows of Symmetry

A Unifying Picture: Scales, Sectors, and the Two Directions. The apparent tension between "symmetry → asymmetry" (spontaneous symmetry breaking, SSB) and "asymmetry → symmetry" (emergent or accidental symmetry) dissolves once we separate **scale** and **sector**. A single physical system should be read in three layers. (1) At short distances the microscopic law $\mathcal{L}_{\text{micro}}$ may be explicitly asymmetric. (2) Under coarse-graining, irrelevant symmetry-violating operators flow to zero, and the infrared law $\mathcal{L}_{\text{IR}}$ often exhibits a **larger** symmetry group H_{IR} than the UV one G_{micro} ($G_{\text{micro}} \subset H_{\text{IR}}$) [82, 14, 102]. (3) The realized **phase** of $\mathcal{L}_{\text{IR}}$ can then **spontaneously break** part of this enhanced symmetry, $H_{\text{IR}} \to K$, producing low-energy excitations on the coset H_{IR}/K. In this layered view, micro-asymmetries are *washed out* at the level of the *law*, while the *state* that the law permits can reintroduce asymmetry through SSB. The residue of the UV asymmetry survives only as power-suppressed corrections in the IR description.

A Canonical, Concrete Example. A standard exemplar is the Heisenberg ferromagnet on a cubic lattice. Microscopically, the lattice breaks continuous spatial rotations, and exchange anisotropies may be present. Yet long-wavelength spin dynamics become effectively **isotropic** because the anisotropy operators are RG-irrelevant; the continuum spin-wave theory inherits rotational symmetry in the IR [102, 101]. Upon cooling below T_c, the **state** spontaneously breaks spin-rotation $O(3) \to O(2)$, and the magnons are the corresponding Nambu–Goldstone modes. In symbols,

$$\text{(lattice anisotropy)} \xRightarrow{\text{RG}} \text{rotationally symmetric } \mathcal{L}_{\text{IR}}, \qquad O(3) \xrightarrow{\text{SSB}} O(2).$$

Minimal Toy Model (Seeing Both Arrows Compactly). The two arrows can be exhibited with two coupled, schematic sectors. For the *law-side* arrow (asymmetry → symmetry), take

$$\boxed{\mathcal{L}_\phi = \tfrac{1}{2} \sum_{i=x,y,z} c_i (\partial_i \phi)^2, \qquad c_i = c + \delta c_i,} \tag{6.277}$$

so that anisotropy enters through operators like $(\partial_x^2 - \partial_y^2)\phi^2$. If such operators have scaling dimension $\Delta > d + z$, their couplings obey $\frac{dg}{d\ell} = (d + z - \Delta)g + \cdots$ and flow to zero; gradients isotropize in the IR [82, 102]. For the *state-side* arrow (symmetry → asymmetry), let a second sector have a global $U(1)$ with

$$\boxed{\mathcal{L}_\sigma = |\partial\sigma|^2 - m^2|\sigma|^2 - \lambda|\sigma|^4, \qquad m^2 > 0,} \tag{6.278}$$

so that $|\langle\sigma\rangle| = v \neq 0$ breaks $U(1)$ and generates one Goldstone boson. Weak couplings between ϕ and σ do not spoil the logic provided the anisotropy stays irrelevant.

Other Ways the Two Arrows Coexist. The coexistence need not occur in the same sector. A fluid sector can display *emergent isotropy* and hydrodynamic conservation laws, while a spin sector simultaneously undergoes SSB and develops long-range order. Likewise, symmetry can be *critically enhanced* at a quantum or classical critical point (often to conformal symmetry), while the adjacent phases on either side of the critical point exhibit distinct SSB patterns [102, 101]. In correlated matter, constraint structures can produce an *emergent gauge redundancy* in the IR; in a different regime the same effective gauge theory can enter a Higgs-like phase, where gauge-invariant diagnostics (massive vectors, screening) distinguish it [437].

How To Diagnose Both Arrows—Separating Law From State Operationally, one first tests for *emergent symmetry of the law*. Measure symmetry-violating features (e.g., anisotropy in correlators or dispersion) and verify that they *shrink with scale*, typically as powers of E/Λ or system size, consistent with RG irrelevance [102, 82]. One then tests for *SSB of the state*: detect a nonzero order parameter, establish symmetry-related degeneracy of vacua, and identify Nambu–Goldstone modes with the expected low-energy dispersion and counting rules [14, 102]. These complementary diagnostics display, side by side, *RG clean-up in the law* and *symmetry hiding in the state*.

A 2×2 Map for Intuition. Arranged as a cartography of "laws vs. states," four corners recur in nature: symmetric laws → symmetric world (unbroken), symmetric laws → asymmetric world (SSB), asymmetric laws → symmetric world (emergent/accidental symmetry), and asymmetric laws → asymmetric world (explicit breaking). The value of the map is pragmatic: it dictates which diagnostics are relevant in each quadrant (Ward identities and topological probes in the unbroken corner; order parameters and NG spectra in SSB; scaling suppression of violations for emergence; explicit coupling–response tracking for explicit breaking).

Why This Structure Is Expected (and Testable). From the RG viewpoint, *universality* follows because many microscopic details—including symmetry-violating terms—are encoded by operators whose couplings diminish under coarse-graining when $\Delta > d + z$. The resulting IR theory inherits an **enhanced symmetry** that governs long-distance observables [32]. From the EFT viewpoint, low-energy physics is organized by the *lowest-dimension* operators; if that minimal set respects a larger group, the symmetry is *accidental* and violations appear only through higher-dimension terms suppressed by E/Λ [415, 403]. In many-body systems, constraints and fractionalization can produce *emergent gauge structure*; when such a theory enters a Higgs-like regime, the would-be Goldstone supplies the longitudinal polarization of a now-massive vector—a perspective anticipated by Anderson in superconductors and unified in modern gauge theory [437, 95].

Putting It All Together. In practice, tuning a control parameter (temperature, density, frustration, pressure) often drives a trajectory III → II: microscopic asymmetry gives way to

Table 6.1: Symmetry–Asymmetry Permutations Matrix: Laws vs. States

Laws / World (State)	**Symmetry of World (State)**	**Asymmetry of World (State)**
Symmetry of Law	**I. Unbroken (symmetry preserved)** Unique symmetric vacuum (thermodynamic limit); order parameter = 0; Ward identities exact. Topological/SPT structure may still distinguish phases.	**II. Spontaneous Symmetry Breaking (SSB)** Degenerate vacua related by the symmetry; nonzero order parameter; Nambu–Goldstone modes for broken *global* continuous generators (gauge case diagnosed via gauge-invariant observables).
Asymmetry of Law	**III. Emergent / Accidental Symmetry** IR (effective-law) exhibits enlarged symmetry via RG irrelevance or EFT power suppression; symmetry-violating effects shrink with scale.	**IV. Explicit Breaking (no SSB)** Asymmetric couplings select a unique ground state; no Goldstones. Residual subgroups may remain; responses track the breaking terms.

Notes: (i) "Gauge symmetry" is a redundancy; BEH/Higgs phases are diagnosed by gauge-invariant features (massive vectors, screening), not literal gauge SSB. (ii) Topological/SPT structure can live in quadrant I (symmetry preserved) and should be diagnosed by nonlocal/torsion-like invariants rather than order parameters.

an *emergent, more symmetric IR law*, and the realized phase then *spontaneously breaks* part of that symmetry. The right workflow is layered: use RG/EFT to determine the *symmetry of the law* that genuinely controls long distances, and then apply SSB diagnostics to determine the *symmetry of the world* the system selects. This two-arrow narrative is not a paradox but the expected consequence of scale separation and effective description [82, 14, 102].

6.10 BRST Quantization and Ghosts: A Cohomological Prelude

Gauge symmetry is a redundancy, not an observable degree of freedom. Quantization therefore proceeds by enlarging field space with (Faddeev–Popov) ghosts and imposing a nilpotent BRST symmetry whose cohomology projects out the unphysical sector [104, 105, 106]. In the present operator–bundle setting, we adopt the formal definitions and constructions of *Chapter 13* (see Section 12.3), so that the gauge fixing is implemented as a BRST-exact term and all observables are read as BRST-invariant classes (cf. Definition 12.58 and Section 12.2). This keeps the discussion here narrative-level while deferring exact algebraic statements, charges, and gradings to that chapter.

The physical state space is the ghost-number-zero BRST cohomology, $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$, so that states are closed but not exact under the conserved BRST charge (Kugo–Ojima quartet mechanism ensures unitarity of the S-matrix on this quotient) [107, 108, 91, 82]. We use Principle 12.1 as our canonical reference: it fixes the operator definition of $\mathcal{H}_{\text{phys}}$, the role of ghost number, and the precise sense in which gauge/longitudinal modes and ghosts organize into cohomologically trivial doublets (cf. Section 12.3). In particular, "Higgsing" in gauge theories is never gauge SSB in the literal sense; would-be Goldstones that supply longitudinal polarizations are BRST-exact and hence non-physical, while physical predictions are cohomology classes, independent of gauge parameters.

Operationally, this BRST purification interlocks with the Holosymmetric Selection Principle (HSP) of *Chapter 13*. Stage I viability—perfect self-coherence of a candidate law—coincides with BRST consistency: nilpotency $Q^2 = 0$, invariance of the gauge-fixed dynamics, and absence of measure anomalies [82, 91]. We adopt the Chapter 13 convention that any uncancelled gauge anomaly manifests as a failure of Stage I and is dynamically excluded (Sections 2.4 and 12.2). This link will underwrite the quadrant analysis: Quadrants I–II (law-symmetric) inherit exact BRST structure; Quadrants III–IV (law-asymmetric) remain admissible only when explicit breakings preserve the BRST scaffold (spurion-consistent or non-gauge breakings), otherwise they are filtered out at Stage I.

For concreteness we will occasionally point back to the QCD example developed in *Chapter 13*: confinement appears as a cohomological selection of color singlets, recovering the hadronic spectrum from the BRST projection in a covariant gauge (see Proposition 12.128 and Section 12.6; cf. [91]). This example is not repeated here; it serves as the paradigm for how BRST cohomology implements "physical sector" extraction inside the operator–bundle formalism. Throughout the remaining sections (including the symmetry–asymmetry quadrants), all identifications of fluctuation modes (Higgs-like, graviton-like, apeiron, chrono–kymon) are kept *provisional* until multi-quadrant plateau-stability is established (cf. Section 12.2).

Conventions. Parity $p(A) := \mathrm{gh}(A) \bmod 2$. The graded commutator and anticommutator are

$$[A,B\} \;:=\; AB - (-1)^{p(A)p(B)}BA, \qquad \{A,B\} := AB + (-1)^{p(A)p(B)}BA,$$

and the BRST charge is odd: $p(\hat{Q}_{\mathrm{BRST}}) = 1$. In what follows, $\{\hat{Q}_{\mathrm{BRST}}, \cdot\}$ denotes the graded anticommutator.

Definition 6.208 (BRST complex and ghost number; recap)**.** Let $(\mathcal{A}, \star)$ be the operator *-algebra obtained from the Algebra of Observables via GNS and acting fiberwise on the modal Hilbert bundle; i.e. $\pi^{\mathring{\Psi}}(\mathcal{A}) \subseteq \mathcal{B}(\mathcal{H}_{\mathrm{em}})$ over $(\mathscr{H}_{\mathrm{mod}}, \pi, \mathcal{X}_q)$ (see Definitions 1.32, 1.34 and 4.239 and Remark 4.290). Per Section 12.3, the BRST charge $\hat{Q}_{\mathrm{BRST}}$ is nilpotent, $\hat{Q}^2_{\mathrm{BRST}} = 0$, with ghost-number grading $\mathrm{gh} : \mathcal{A} \to \mathbb{Z}$ and $[\mathrm{gh}, \hat{Q}_{\mathrm{BRST}}] = \hat{Q}_{\mathrm{BRST}}$ (so $p(A) = \mathrm{gh}(A) \bmod 2$). The cohomology in degree k is

$$H^k(\hat{Q}_{\mathrm{BRST}}) := \ker(\hat{Q}_{\mathrm{BRST}}|_{\mathrm{gh}=k}) \Big/ \mathrm{im}(\hat{Q}_{\mathrm{BRST}}|_{\mathrm{gh}=k-1}).$$

The physical state space is $\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}})$, per Principle 12.1.

Definition 6.209 (Cohomological projector)**.** Assume a contracting homotopy K of ghost number -1 exists on the BRST–doublet (unphysical) sector as constructed in Section 12.6.5, with

$$\{\hat{Q}_{\mathrm{BRST}}, K\} = \mathbf{1} - \Pi_{\mathrm{phys}} \quad \text{on the unphysical subspace}$$

(cf. the proof leading to the evolution–to–cohomology statement (12.79); see also Definition 12.129). Define the projector

$$\mathsf{P}_{\mathrm{phys}} \;:=\; \mathbf{1} - \{\hat{Q}_{\mathrm{BRST}}, K\}, \qquad \mathsf{P}^2_{\mathrm{phys}} = \mathsf{P}_{\mathrm{phys}}, \quad [\mathsf{P}_{\mathrm{phys}}, \hat{Q}_{\mathrm{BRST}}] = 0, \quad \mathrm{gh}(\mathsf{P}_{\mathrm{phys}}) = 0.$$

It acts as the identity on $H^0(\hat{Q}_{\mathrm{BRST}})$ and annihilates BRST-exact states.

Proof sketch. Expand $(\mathbf{1} - \{\hat{Q}, K\})^2$ using $\hat{Q}^2 = 0$ and the fact that $\{\hat{Q}, \Pi_{\mathrm{phys}}\} = 0$ in the construction of (12.79).

Lemma 6.210 (Graded cyclicity kills exacts)**.** *Assume the BRST-closed vacuum functional of Postulate 12.2 is realized via the GNS construction Definition 1.34; write* $\mathrm{Tr}(\cdot) := \omega(\cdot)$.[5] *If $\mathcal{O}$ is BRST-closed with ghost number* 0 *and X is any (homogeneous) operator, then*

$$\mathrm{Tr}\Big(\mathsf{P}_{\mathrm{phys}}\, \mathcal{O}\, \{\hat{Q}_{\mathrm{BRST}}, X\}\Big) = 0,$$

equivalently the band identity eq. (7.331)*.*

Proof. Use that $\mathsf{P}_{\mathrm{phys}}$ commutes with $\hat{Q}_{\mathrm{BRST}}$ and has ghost number 0 (eq. (7.330)). By graded Leibniz,

$$\{\hat{Q}_{\mathrm{BRST}},\, \mathsf{P}_{\mathrm{phys}}\mathcal{O}X\} = \{\hat{Q}_{\mathrm{BRST}}, \mathsf{P}_{\mathrm{phys}}\}\mathcal{O}X \;\pm\; \mathsf{P}_{\mathrm{phys}}\{\hat{Q}_{\mathrm{BRST}}, \mathcal{O}\}X \;\pm\; \mathsf{P}_{\mathrm{phys}}\mathcal{O}\{\hat{Q}_{\mathrm{BRST}}, X\}.$$

[5] GNS: Definition 1.34. BRST-closedness of the extended state: Postulate 12.2.

Since $\{\hat{Q}_{\mathrm{BRST}}, \mathsf{P}_{\mathrm{phys}}\} = 0$ and $\{\hat{Q}_{\mathrm{BRST}}, \mathcal{O}\} = 0$, this reduces to

$$\{\hat{Q}_{\mathrm{BRST}},\ \mathsf{P}_{\mathrm{phys}}\mathcal{O}X\} = \pm\, \mathsf{P}_{\mathrm{phys}}\mathcal{O}\{\hat{Q}_{\mathrm{BRST}}, X\}.$$

Now evaluate the state on both sides. BRST-closedness of ω implies $\omega(\{\hat{Q}_{\mathrm{BRST}}, Y\}) = 0$ for any Y (Postulate 12.2), hence

$$0 = \omega\Big(\{\hat{Q}_{\mathrm{BRST}},\ \mathsf{P}_{\mathrm{phys}}\mathcal{O}X\}\Big) = \pm\,\omega\Big(\mathsf{P}_{\mathrm{phys}}\mathcal{O}\{\hat{Q}_{\mathrm{BRST}}, X\}\Big),$$

which is the claim. In particular, with $\mathrm{Tr}(\cdot) = \omega(\cdot)$ this reproduces the band identity eq. (7.331). □

Lemma 6.211 (BRST doublet triviality)**.** *Let (u, v) be a BRST doublet with $[\hat{Q}_{\mathrm{BRST}}, u\} = v$ and $[\hat{Q}_{\mathrm{BRST}}, v\} = 0$ (Section 6.10). Then for any (graded) polynomial F in u, v,*

$$F(u,v) - F(u,0) \;=\; \{\hat{Q}_{\mathrm{BRST}},\ KF\}(u,v), \qquad KF \;:=\; \int_0^1 dt\; u\,(\partial_v F)(u,tv). \tag{6.279}$$

In particular, if $F(u,0) = 0$ (as for pure longitudinal/ghost insertions), then $F(u,v)$ is BRST-exact: $F(u,v) = \{\hat{Q}_{\mathrm{BRST}}, KF\}$. Hence matrix elements of $F(u,v)$ between physical states vanish by Lemma 6.210.

Proof. We work with the graded Leibniz rule from Section 6.10. Define the homotopy operator

$$(KF)(u,v) \;:=\; \int_0^1 dt\; u\,(\partial_v F)(u,tv),$$

where ∂_v is the graded left derivative with respect to v. Compute

$$\{\hat{Q}_{\mathrm{BRST}}, KF\} = \int_0^1 dt\, \Big(\{\hat{Q}_{\mathrm{BRST}}, u\}\,(\partial_v F)(u,tv) \pm u\,\{\hat{Q}_{\mathrm{BRST}}, (\partial_v F)(u,tv)\}\Big).$$

Using the doublet relations $\{\hat{Q}_{\mathrm{BRST}}, u\} = v$ and $\{\hat{Q}_{\mathrm{BRST}}, v\} = 0$, the second term reduces via the chain rule to

$$\begin{aligned}\{\hat{Q}_{\mathrm{BRST}}, (\partial_v F)(u,tv)\} &= (\partial_u\partial_v F)(u,tv)\,\{\hat{Q}_{\mathrm{BRST}}, u\} \\ &\quad + t\,(\partial_v\partial_v F)(u,tv)\,\{\hat{Q}_{\mathrm{BRST}}, v\} \\ &= (\partial_u\partial_v F)(u,tv)\,v,\end{aligned}$$

with the sign fixed by the parities $p(u), p(v)$. Hence

$$\{\hat{Q}_{\mathrm{BRST}}, KF\} = \int_0^1 dt\, \Big(v\,\partial_v F(u,tv) \pm u\,(\partial_u\partial_v F)(u,tv)\,v\Big).$$

By graded calculus, the integrand is exactly $\dfrac{d}{dt}F(u,tv)$:

$$\frac{d}{dt}F(u,tv) = v\,\partial_v F(u,tv) \pm u\,(\partial_u\partial_v F)(u,tv)\,v,$$

so integrating from $t = 0$ to 1 yields

$$\{\hat{Q}_{\mathrm{BRST}}, KF\} = F(u, v) - F(u, 0).$$

This proves (6.279). If $F(u, 0) = 0$, then $F(u, v)$ itself is BRST-exact. Finally, by Lemma 6.210, any exact insertion has vanishing physical matrix elements, and the conclusion follows. (Equivalently, the cohomology is independent of doublet variables; cf. Corollary 6.212.) □

Conventions. We write the BRST super-derivation as $\zeta(\,\cdot\,) \coloneqq [\hat{Q}_{\mathrm{BRST}}, \cdot\,\}$. In Chapter 11 this is denoted δ_{BRST}; thus $\zeta \equiv \delta_{\mathrm{BRST}}$. The parameter s is reserved exclusively for homotopies/evolution and never denotes a BRST differential.

Corollary 6.212 (Quartet mechanism; unitarity on the quotient)**.** *Longitudinal gauge modes and would-be Goldstones, together with $(c, \bar{c})$ and the Nakanishi–Lautrup fields, assemble into BRST doublets. Their insertions are cohomologically trivial in $H^0(\hat{Q}_{\mathrm{BRST}})$ by Lemmas 6.210 and 6.211. Consequently, the S-matrix is unitary on the physical space $\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}})$ (Kugo–Ojima [107, 108]; cf. [91, 82]).*

Proof. (Notation: in Chapter 7 we write the BRST super-derivation as $\zeta(\,\cdot\,) := \{\hat{Q}_{\mathrm{BRST}}, \cdot\,\}$, per Section 6.10.)

(1) *Cohomological triviality of the unphysical sector.* Let $\Pi_L^{\mu\nu}$ denote the longitudinal projector (e.g. $\Pi_L^{\mu\nu} = \partial^\mu\partial^\nu/\Box$ in a linear gauge, or its covariant/band analogue). Define

$$A_\mu^{\mathrm{L}} := \Pi_{L\,\mu}{}^{\nu} A_\nu, \qquad \bar{D}c := \Pi_L^{\mu\nu} D_\nu c,$$

with $D_\mu c = \partial_\mu c + [A_\mu, c]$. Using the BRST rules (graded anticommutator),

$$\zeta A_\mu = D_\mu c, \qquad \zeta c = -\tfrac{1}{2}[c, c], \qquad \zeta\bar{c} = b, \qquad \zeta b = 0,$$

we obtain

$$\zeta A_\mu^{\mathrm{L}} = \Pi_{L\,\mu}{}^{\nu} D_\nu c = \bar{D}c.$$

Moreover,

$$\zeta(\bar{D}c) = \Pi_L^{\mu\nu}\Big(D_\nu \zeta c + [\zeta A_\nu, c]\Big) = \Pi_L^{\mu\nu}\Big(-\tfrac{1}{2} D_\nu[c, c] + [D_\nu c, c]\Big).$$

In the linearised BRST differential (the Hessian/free limit relevant for the physical spectrum) one has $\zeta c = 0$, hence $\zeta(\bar{D}c) = 0$ exactly; beyond linear order the right-hand side is ζ–exact. Thus $(A^{\mathrm{L}}, \bar{D}c)$ is a BRST doublet in the sense of Lemma 6.211. The pair $(\bar{c}, b)$ satisfies $\zeta\bar{c} = b$, $\zeta b = 0$ and is a second BRST doublet. By Lemma 6.211, any graded polynomial in the variables of these pairs is ζ–exact; by Lemma 6.210 such exact insertions have vanishing matrix elements between physical states. Therefore longitudinal, ghost, and Nakanishi–Lautrup contributions decouple in $H^0(\hat{Q}_{\mathrm{BRST}})$.

(2) *Descent of dynamics and definition of S_{phys}.* By eq. (12.79) we have $[\hat{Q}_{\mathrm{BRST}}, \hat{H}_{\mathrm{tot}}] = 0$, hence $\hat{Q}_{\mathrm{BRST}}$ is conserved and commutes with the scattering operator $\mathcal{S}$ built from $\hat{H}_{\mathrm{tot}}$: $[\hat{Q}_{\mathrm{BRST}}, \mathcal{S}] = 0$. Therefore $\mathcal{S}$ preserves BRST classes. With the physical space defined in Definition 12.129 (see also Principle 12.1), the induced map

$$\mathcal{S}_{\mathrm{phys}} \colon H^0(\hat{Q}_{\mathrm{BRST}}) \to H^0(\hat{Q}_{\mathrm{BRST}}), \qquad [\psi] \mapsto [\mathcal{S}\psi],$$

is well defined.

(3) *Unitarity on the quotient.* Let P_{phys} be the cohomological projector of Definition 6.209, so $\mathsf{P}_{\text{phys}}^2 = \mathsf{P}_{\text{phys}}$, $[\mathsf{P}_{\text{phys}}, \hat{Q}_{\text{BRST}}] = 0$, $\text{gh}(\mathsf{P}_{\text{phys}}) = 0$. Define $\mathcal{S}_{\text{phys}} := \mathsf{P}_{\text{phys}}\mathcal{S}\,\mathsf{P}_{\text{phys}}$. Using $[\hat{Q}_{\text{BRST}}, \mathcal{S}] = 0$ and $\mathsf{P}_{\text{phys}} = \mathbf{1} - \{\hat{Q}_{\text{BRST}}, K\}$ (Definition 6.209),

$$[\mathsf{P}_{\text{phys}}, \mathcal{S}] = \{\hat{Q}_{\text{BRST}},\, \mathcal{S}K - K\mathcal{S}\},$$

is ζ–exact. Sandwiching with P_{phys} and invoking Lemma 6.210 yields $\mathsf{P}_{\text{phys}}\mathcal{S} = \mathcal{S}\mathsf{P}_{\text{phys}}$ on the physical subspace. Since $\mathcal{S}$ is unitary on the total space,

$$\mathcal{S}_{\text{phys}}^\dagger \mathcal{S}_{\text{phys}} = \mathsf{P}_{\text{phys}}\mathcal{S}^\dagger \mathsf{P}_{\text{phys}}\mathcal{S}\mathsf{P}_{\text{phys}} = \mathsf{P}_{\text{phys}}\mathcal{S}^\dagger\mathcal{S}\mathsf{P}_{\text{phys}} = \mathsf{P}_{\text{phys}},$$

and similarly $\mathcal{S}_{\text{phys}}\mathcal{S}_{\text{phys}}^\dagger = \mathsf{P}_{\text{phys}}$. Hence $\mathcal{S}_{\text{phys}}$ is unitary on $\mathcal{H}_{\text{phys}} = \text{im}\,\mathsf{P}_{\text{phys}}$. □

Proposition 6.213 (Higgsing is not gauge SSB)**.** *In a gauge theory with a Higgs potential, the would-be Goldstone excitations lie in BRST doublets with the longitudinal modes. Therefore no gauge generator acquires a physical order parameter; only gauge-invariant cohomology classes survive, and physical poles/residues are gauge-parameter independent (see Proposition 6.214; cf. [107, 108, 91, 82]).*

Proof. (Notation: to avoid conflict with the state s in Section 6.11, set $\delta_{\text{BRST}}(\,\cdot\,) := \{\hat{Q}_{\text{BRST}}, \cdot\}$, the graded anticommutator from Section 6.10.)

(i) Doublet pairing of unphysical modes. In an R_ζ gauge, the longitudinal gauge component A^{L} and the ghost combination $\bar{D}c$ form a BRST doublet, as do $(\bar{c}, b)$ (Nakanishi–Lautrup). By Lemma 6.211, any graded polynomial in these variables is δ_{BRST}-exact. Hence their insertions vanish between physical states by Lemma 6.210. Therefore the would-be Goldstones that supply longitudinal polarizations are cohomologically trivial—they live in doublets with A^{L} and decouple from $H^0(\hat{Q}_{\text{BRST}})$ (see also Corollary 6.212).

(ii) No gauge order parameter in the physical sector. Suppose a gauge generator G^a had a nonzero physical order parameter: there exists a *physical* operator $\mathcal{O} \in H^0(\hat{Q}_{\text{BRST}})$ with $\langle\mathcal{O}\rangle \neq 0$ and $[G^a, \mathcal{O}] \neq 0$. But gauge-charged operators transform nontrivially under BRST, e.g.

$$\delta_{\text{BRST}}\Phi \;=\; i\,c^a T^a \Phi$$

for a Higgs multiplet Φ, so any operator carrying nonzero gauge charge is not BRST-closed and thus not an element of H^0. Consequently, any admissible order parameter must be gauge invariant (a cohomology class). In the BEH phase, the nonzero background v reorganizes field variables; the physical content—cohomology classes and their matrix elements—remains gauge invariant. Hence there is no *gauge* SSB in the physical sector.

(iii) Gauge-parameter independence of poles/residues. Gauge fixing enters as a BRST-exact deformation, $\mathcal{L}_{\text{gf+gh}} = \{\hat{Q}_{\text{BRST}}, \Psi\}$ (Principle 6.1). By the Nielsen identity (Proposition 6.214), for any BRST-closed observable $\mathcal{O}$ and BRST-stationary state, $\partial_\zeta \langle\!\langle \mathcal{O} \rangle\!\rangle_{\text{phys}} = 0$. Taking $\mathcal{O}$ to be the physical two-point kernel (or its spectral projector) implies $\partial_\zeta m_n^2 = \partial_\zeta Z_n = 0$ for the poles and residues of physical excitations. Thus the spectrum extracted from H^0 is gauge-parameter independent.

Combining (i)–(iii), Higgsing does not produce a physical (gauge) order parameter; it trades unphysical modes into BRST doublets that decouple from H^0, while observable

quantities are gauge-invariant cohomology classes with ζ-independent poles/residues. This is the Kugo–Ojima picture of unitarity and physical state space [107, 108]; see also [91, 82]. □

Principle 6.1 (Gauge fixing is BRST-exact; Nielsen form)**.** *Let the gauge parameter ξ enter only through the gauge-fixing fermion $\Psi(\xi)$ so that*

$$\mathcal{L}_{\mathrm{gf+gh}} = \zeta\Psi, \qquad \partial_\xi \hat{H}_{\mathrm{tot}} = \zeta X_\xi$$

for some operator X_ξ, where $\zeta(\,\cdot\,) := \{\hat{Q}_{\mathrm{BRST}}, \cdot\}$ (see Section 6.10). Assume further that the state/measure is BRST-closed (no anomaly) and that we evaluate in the physical sector using the projector of Definition 6.209: $\langle\!\langle \mathcal{O} \rangle\!\rangle_{\mathrm{phys}} := \frac{\mathrm{Tr}\left(\mathsf{P}_{\mathrm{phys}} \mathcal{O}\, \rho\right)}{\mathrm{Tr}\left(\mathsf{P}_{\mathrm{phys}} \rho\right)}$. Then for any ghost-number-zero, BRST-closed observable $\mathcal{O}$,

$$\partial_\xi \, \langle\!\langle \mathcal{O} \rangle\!\rangle_{\mathrm{phys}} = 0.$$

One-line proof. Using $\partial_\xi \hat{H}_{\mathrm{tot}} = \zeta X_\xi$ and the BRST-closed state $[\hat{Q}_{\mathrm{BRST}}, \rho] = 0$,

$$\partial_\xi \, \mathrm{Tr}(\mathsf{P}_{\mathrm{phys}} \mathcal{O}\, \rho) = \mathrm{Tr}\Big(\mathsf{P}_{\mathrm{phys}} \mathcal{O}\, \zeta X_\xi\, \rho\Big) = \mathrm{Tr}\Big(\zeta(\mathsf{P}_{\mathrm{phys}} \mathcal{O} X_\xi \rho)\Big) = 0$$

by Lemma 6.210; the denominator is identical. □

Proposition 6.214 (Nielsen identity, operator form)**.** *Let ξ be a gauge parameter. Suppose the total Hamiltonian (or generator) varies as*

$$\partial_\xi \hat{H}_{\mathrm{tot}} = \zeta X_\xi, \qquad \text{with} \quad \zeta(\,\cdot\,) := \{\hat{Q}_{\mathrm{BRST}}, \cdot\} \quad (\text{see Section 6.10}),$$

and the state is BRST-stationary, $[\hat{Q}_{\mathrm{BRST}}, \rho] = 0$ (equivalently $\rho = \mathsf{P}_{\mathrm{phys}} \rho\, \mathsf{P}_{\mathrm{phys}}$). Then for every ghost-number-zero, BRST-closed observable $\mathcal{O}$,

$$\partial_\xi \, \mathrm{Tr}\Big(\mathsf{P}_{\mathrm{phys}}\, \mathcal{O}\, \rho\Big) = 0, \qquad \partial_\xi \, \langle\!\langle \mathcal{O} \rangle\!\rangle_{\mathrm{phys}} := \partial_\xi \frac{\mathrm{Tr}(\mathsf{P}_{\mathrm{phys}}\, \mathcal{O}\, \rho)}{\mathrm{Tr}(\mathsf{P}_{\mathrm{phys}}\, \rho)} = 0.$$

One-line proof. Using $\partial_\xi \hat{H}_{\mathrm{tot}} = \zeta X_\xi$ and $[\hat{Q}_{\mathrm{BRST}}, \rho] = 0$,

$$\partial_\xi \, \mathrm{Tr}(\mathsf{P}_{\mathrm{phys}} \mathcal{O}\, \rho) = \mathrm{Tr}\Big(\mathsf{P}_{\mathrm{phys}} \mathcal{O}\, \zeta X_\xi\, \rho\Big) = \mathrm{Tr}\Big(\zeta(\mathsf{P}_{\mathrm{phys}} \mathcal{O} X_\xi \rho)\Big) = 0$$

by Lemma 6.210; the denominator is identical. □

Definition 6.215 (Physical holosymmetric visibility)**.** Given a register state ρ and a holosymmetric effect $E_{\mathrm{Holo}}^{(\mathbf{a}\to\mathbf{b})}$ as in eq. (3.36), define the physical visibility by

$$v_{\mathrm{phys}}(\rho) := \frac{\mathrm{Tr}\Big(\mathsf{P}_{\mathrm{phys}}\, E_{\mathrm{Holo}}^{(\mathbf{a}\to\mathbf{b})}\, \rho\Big)}{\mathrm{Tr}\Big(\mathsf{P}_{\mathrm{phys}}\, \rho\Big)}.$$

Proposition 6.216 (BRST invariance of holosymmetric readouts)**.** *Assume* $\mathrm{gh}\Big(E_{\mathrm{Holo}}^{(\mathbf{a}\to\mathbf{b})}\Big) = 0$ *and* $\zeta E_{\mathrm{Holo}}^{(\mathbf{a}\to\mathbf{b})} = 0$*, where $\zeta(\,\cdot\,) := \{\hat{Q}_{\mathrm{BRST}}, \cdot\}$ (see Section 6.10). If*

$$E_{\mathrm{Holo}}^{(\mathbf{a}\to\mathbf{b})} \;\mapsto\; E_{\mathrm{Holo}}^{(\mathbf{a}\to\mathbf{b})} + \zeta X \quad \text{or} \quad \rho \;\mapsto\; \rho + \zeta Y,$$

then the physical visibility of Definition 6.215, $v_{\mathrm{phys}}(\rho) = \frac{\mathrm{Tr}\Big(\mathsf{P}_{\mathrm{phys}}\, E_{\mathrm{Holo}}^{(\mathbf{a}\to\mathbf{b})}\, \rho\Big)}{\mathrm{Tr}\Big(\mathsf{P}_{\mathrm{phys}}\, \rho\Big)}$, is unchanged. Hence holosymmetric overlaps/visibilities depend only on BRST cohomology classes.

Proof. For the numerator,

$$\delta \operatorname{Tr}(\mathsf{P}_{\text{phys}} E\rho) = \operatorname{Tr}\Big(\mathsf{P}_{\text{phys}}\, \zeta X\, \rho\Big) \quad \text{or} \quad \operatorname{Tr}\Big(\mathsf{P}_{\text{phys}}\, E\, \zeta Y\Big) = \operatorname{Tr}\Big(\zeta(\mathsf{P}_{\text{phys}} EX\rho)\Big) = 0,$$

by Lemma 6.210. The denominator varies similarly and vanishes for the same reason. □

Principle 6.2 (Stage I viability equals BRST consistency)**.** *A candidate law $\mathcal{L}$ passes Stage I of HSP iff, in the BRST framework of Chapter 11,*

$$\hat{Q}_{\text{BRST}}^2 = 0, \qquad [\hat{Q}_{\text{BRST}}, \hat{H}_{tot}] = 0, \qquad \Delta_{\text{meas}} = 0,$$

i.e. nilpotency, BRST-invariant dynamics, and vanishing Jacobian (no gauge anomaly; see eq. (12.84)). Any failure excludes $\mathcal{L}$ at Stage I.

Proposition 6.217 (Anomaly cohomology test)**.** *Let $\mathcal{A}$ be the local measure variation defined by eq. (12.84). If $[\mathcal{A}] \in H^{+1}(\hat{Q}_{\text{BRST}})$ is nontrivial, Stage I fails. Conversely, $[\mathcal{A}] = 0$ implies anomaly cancellation up to ζ–exact counterterms.*

Lemma 6.218 (Covariant BRST compatibility)**.** *Let ∇_μ be the* split connection *of Section 6.12.2 and eq. (6.314) on the operator bundle, decomposed into the operator–bundle (gauge) connection $\mathcal{A}_\mu$ and the relational Levi–Civita part $\nabla_\mu^{(\text{rel})}$ (see Definition 6.192 and Definition 6.7 and eq. (6.22)). If*

$$[\hat{Q}_{\text{BRST}}, \mathcal{A}_\mu] = 0, \qquad [\hat{Q}_{\text{BRST}}, \nabla_\mu^{(\text{rel})}] = 0,$$

then $[\hat{Q}_{\text{BRST}}, \nabla_\mu] = 0$ (BRST-invariant connection). Consequently, ∇_μ maps BRST-closed operators to BRST-closed operators and preserves $H^0(\hat{Q}_{\text{BRST}})$ along the base manifold.

Proof sketch. By Section 6.12.2 and eq. (6.314), $\nabla_\mu = \nabla_\mu^{(\text{rel})} + \mathcal{A}_\mu$ acts as a derivation on operator fields. Hence

$$[\hat{Q}_{\text{BRST}}, \nabla_\mu] = [\hat{Q}_{\text{BRST}}, \nabla_\mu^{(\text{rel})}] + [\hat{Q}_{\text{BRST}}, \mathcal{A}_\mu] = 0 + 0 = 0.$$

If $\hat{Q}_{\text{BRST}}\mathcal{O} = 0$ and $[\hat{Q}_{\text{BRST}}, \nabla_\mu] = 0$, then $\hat{Q}_{\text{BRST}}(\nabla_\mu \mathcal{O}) = \nabla_\mu(\hat{Q}_{\text{BRST}}\mathcal{O}) = 0$, so $\nabla_\mu \mathcal{O}$ is also closed; exacts remain exact by the same computation, and thus H^0 is preserved along the base. □

Remark 6.219 (Band-compatibility)**.** Let $P_x(\lambda_{\text{UV}})$ be the UV projector of eq. (7.29) (with BRST closure eq. (7.30)) and $K_{\text{IR}}^{(\ell)}$ the IR kernels of eqs. (7.32) and (7.33). Because both carry ghost number 0 and commute with $\hat{Q}_{\text{BRST}}$ eq. (7.30), the band/IR functor $\mathcal{B}^{(\lambda_{\text{UV}},\ell)}$ of eq. (7.40) commutes with P_{phys} and with $\hat{Q}_{\text{BRST}}$. Therefore all statements of Lemma 6.218 apply verbatim to band-limited, IR-smoothed composites.

Example (pointer). For color gauge symmetry, the constraints are implemented cohomologically in Chapter 11; the projector P_{phys} selects the color-singlet sector, reproducing the hadronic spectrum in covariant gauges. See Proposition 12.128 and Section 12.6.

6.11 From Duality to the Symmetry–Asymmetry Quadrants

Goal. Give a formal route from the Axiom of Duality to the 2×2 "Laws vs States" quadrants in table 6.1, using the existing objects and pipeline.

6.11.1 Duality, Lifted to Two Layers

Typing. $\mathsf{Status}_{\mathcal{C}} : \mathcal{U}_{\mathcal{C}} \times \Lambda_{\mathcal{C}} \to \{\mathsf{Th}, \mathsf{AntiTh}, \mathrm{Syn}, \mathrm{Holo}\}$, $\mathbf{D}_{\mathcal{C}} : \mathcal{U}_{\mathcal{C}} \to \mathcal{U}_{\mathcal{C}}$ with $\mathbf{D}_{\mathcal{C}}^2 = \mathrm{Id}$, $\Delta_{\mathcal{C}} : \{\mathsf{Th}, \mathsf{AntiTh}, \mathrm{Syn}, \mathrm{Holo}\} \to \{\mathsf{Th}, \mathsf{AntiTh}, \mathrm{Syn}, \mathrm{Holo}\}$ with $\Delta_{\mathcal{C}}^2 = \mathrm{Id}$ and $\Delta_{\mathcal{C}}(\mathsf{Th}) = \mathsf{AntiTh}$, $\Delta_{\mathcal{C}}(\mathsf{AntiTh}) = \mathsf{Th}$, $\Delta_{\mathcal{C}}(\mathrm{Syn}) = \mathrm{Syn}$, $\Delta_{\mathcal{C}}(\mathrm{Holo}) = \mathrm{Holo}$.

Definition 6.220 (Axiom of Duality (object level))**.** Fix an immanent context $\mathcal{C}$ with admissible primitives $\mathcal{U}_{\mathcal{C}}$. There is an involution $\mathbf{D}_{\mathcal{C}} : \mathcal{U}_{\mathcal{C}} \to \mathcal{U}_{\mathcal{C}}$ such that $\mathbf{D}_{\mathcal{C}}^2 = \mathrm{Id}$ and for any $s \in \mathcal{U}_{\mathcal{C}}$, the assertion of s entails the conceptual co-arising of $\mathbf{D}_{\mathcal{C}}(s)$ (dyads are the minimal semantic unit). See also the early formulation in Definition 2.100.

Definition 6.221 (Status/tetralemma lift)**.** There is an involution $\Delta_{\mathcal{C}}$ on corner-statuses exchanging Thesis↔AntiThesis while fixing Synthesis and Holothesis, with the compatibility condition

$$\mathsf{Status}_{\mathcal{C}}\Big(\mathbf{D}_{\mathcal{C}}(\varphi), \lambda\Big) = \Delta_{\mathcal{C}}\Big(\mathsf{Status}_{\mathcal{C}}(\varphi, \lambda)\Big). \tag{6.280}$$

This is the status-level lift of Definition 6.220; cf. the tetralemma discussion in Section 1.1.5.

Lemma 6.222 (Self-similarity across reindexings)**.** *For any admissible "change of aspect" F (e.g., coarse-graining, sector split), there exists a unique lifted duality $\mathbf{D}_{\mathcal{C}'}$ with $F \circ \mathbf{D}_{\mathcal{C}} = \mathbf{D}_{\mathcal{C}'} \circ F$. Hence duality is scale-free and the four statuses are closed under repeated dualization; cf. Proposition 1.12.*

Proof of Lemma 6.222 (sketch). If $F : \mathcal{U}_{\mathcal{C}} \to \mathcal{U}_{\mathcal{C}'}$ is an isomorphism in the admissible class, set $\mathbf{D}_{\mathcal{C}'} := F \circ \mathbf{D}_{\mathcal{C}} \circ F^{-1}$, which is an involution and makes $F \circ \mathbf{D}_{\mathcal{C}} = \mathbf{D}_{\mathcal{C}'} \circ F$ tautological and unique. For coarse-grainings/quotients, assume $\sim_F$ is the identification relation and $\mathbf{D}_{\mathcal{C}}$ preserves $\sim_F$-classes (i.e. $x \sim_F y \Rightarrow \mathbf{D}_{\mathcal{C}} x \sim_F \mathbf{D}_{\mathcal{C}} y$). Then define $\mathbf{D}_{\mathcal{C}'}([x]) := [\mathbf{D}_{\mathcal{C}} x]$; well-definedness and involutivity are immediate, and uniqueness follows since F is surjective to classes. The status-compatibility (6.280) is preserved by the same argument applied to $\mathsf{Status}_{\mathcal{C}}$ and $\mathsf{Status}_{\mathcal{C}'}$. □

Remark 6.223 (Law vs State as two levels of the same pipeline)**.** In the blueprint → language → arena passage (Fourfold algebra → observable algebra → GNS/Holo-State), the *law* resides at the algebraic/variational level (action/Hamiltonian invariances), while the *state* resides at the realized expectation level (vacua/ensembles). This is why the matrix in table 6.1 splits into rows (law) and columns (state); see also the Fourfold inter-modal correlator and the unified inter-modal correlator in Definitions 4.9 and 6.4.

6.11.2 Two Dyads and Their Product

Fix a global symmetry group G with a (projective) unitary representation $U : G \to \mathcal{U}(\mathcal{H})$ and generators Q^a (so $U(e^{i\epsilon_a T^a}) = \mathbf{1} + i\epsilon_a Q^a + \cdots$).

Definition 6.224 (Law- and State-predicates)**.** 1. **Law-predicate.** $P_{\mathrm{Law}}(G) :\iff \delta_G S = 0$ (equivalently $[\![Q^a, \mathcal{L}]\!] = \partial_\mu K^{\mu a}$, i.e. invariance up to a total derivative and with anomaly-free Ward identities). Declare

$$\mathsf{Status}(P_{\mathrm{Law}}, \lambda_{\mathrm{Law}}) = \begin{cases} \mathsf{Th} & \text{if } P_{\mathrm{Law}}(G) \text{ holds (“symmetric law”)} \\ \mathsf{AntiTh} & \text{otherwise (“asymmetric law”)} \end{cases}$$

2. **State-predicate.** $P_{\mathrm{State}}(G) :\iff U(g)\rho\,U(g)^{-1} = \rho\ \forall g \in G$ (equivalently $[Q^a, \rho] = 0$, so all G-charged order parameters have vanishing expectations and Ward identities hold on the state). Declare

$$\mathsf{Status}(P_{\mathrm{State}}, \lambda_{\mathrm{State}}) = \begin{cases} \mathsf{Th} & \text{if } P_{\mathrm{State}}(G) \text{ holds (“symmetric state”)} \\ \mathsf{AntiTh} & \text{otherwise (“asymmetric state”)} \end{cases}$$

By Definition 6.221, each predicate acquires a dual status under $\Delta_{\mathcal{C}}$: symmetry $\leftrightarrow$ asymmetry.

Proposition 6.225 (Commuting lifts and generated group)**.** *There exist commuting involutions* Δ_{Law} *and* Δ_{State} *(functorial lifts of* $\Delta_{\mathcal{C}}$ *to the law and state aspects) with* $\Delta_{\mathrm{Law}}^2 = \Delta_{\mathrm{State}}^2 = \mathrm{Id}$. *They act on ordered pairs of statuses by*

$$\Delta_{\mathrm{Law}} : (\sigma_{\mathrm{L}}, \sigma_{\mathrm{S}}) \mapsto (\Delta\sigma_{\mathrm{L}}, \sigma_{\mathrm{S}}), \qquad \Delta_{\mathrm{State}} : (\sigma_{\mathrm{L}}, \sigma_{\mathrm{S}}) \mapsto (\sigma_{\mathrm{L}}, \Delta\sigma_{\mathrm{S}}),$$

where Δ *flips* $\mathsf{Th} \leftrightarrow \mathsf{AntiTh}$ *and fixes* $\mathrm{Syn}, \mathrm{Holo}$ *when those statuses are present. Hence the generated group is* $\mathbb{Z}_2 \times \mathbb{Z}_2$*, and the orbit on any initial dyad consists of the four corners*

$$(\mathsf{Th}, \mathsf{Th}), \quad (\mathsf{Th}, \mathsf{AntiTh}), \quad (\mathsf{AntiTh}, \mathsf{Th}), \quad (\mathsf{AntiTh}, \mathsf{AntiTh}). \tag{6.281}$$

Proof. By Definition 6.221 and naturality under reindexings (Lemma 6.222), the same involution $\Delta_{\mathcal{C}}$ lifts independently to the law and state aspects. Each lift squares to the identity. Since the two lifts act on disjoint coordinates of the ordered pair, they commute; hence the subgroup they generate is $\{\,\mathrm{Id}, \Delta_{\mathrm{Law}}, \Delta_{\mathrm{State}}, \Delta_{\mathrm{Law}}\Delta_{\mathrm{State}}\,\} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$. The stated orbit follows. □

6.11.3 BRST-Compatible Reading

Remark 6.226 (Cohomological reading of the dyads)**.** The Axiom of Duality (Definitions 6.220 and 6.221) generates the two commuting dyads (Law, State) independently of any quantization choices. When gauge redundancy is present, we only *change the evaluation map* of the predicates in Definition 6.224—not the duality itself—by reading them in BRST cohomology:

$$\begin{aligned} \zeta(\,\cdot\,) &:= \{\hat{Q}_{\mathrm{BRST}}, \cdot\,\}, \qquad \text{see Section 6.10,} \\ \mathcal{A}^{(G)}_{\mathrm{phys}} &:= H^0\big(\mathcal{A}^{(G)}_{\mathrm{BRST}}, \zeta\big), \\ \mathcal{H}_{\mathrm{phys}} &:= H^0(\hat{Q}_{\mathrm{BRST}}). \end{aligned}$$

Law-predicate. Evaluate $P_{\mathrm{Law}}(G)$ as invariance in $\mathcal{A}^{(G)}_{\mathrm{phys}}$ (equivalently, invariance modulo ζ-exact deformations). *State-predicate.* Evaluate $P_{\mathrm{State}}(G)$ on the BRST-closed physical state $\rho_{\mathrm{phys}} := \mathsf{P}_{\mathrm{phys}}\rho\,\mathsf{P}_{\mathrm{phys}}$ on $\mathcal{H}_{\mathrm{phys}}$ (Definition 6.209 and Principle 12.1).

In this reading, "gauge SSB'' never occurs literally: would-be Goldstones, ghosts, and Nakanishi–Lautrup auxiliaries organize into BRST doublets and are cohomologically trivial (Lemma 6.211 and Corollary 6.212). Thus the dyads and their $\mathbb{Z}_2 \times \mathbb{Z}_2$ action (Proposition 6.225) remain unchanged; only truth-evaluation is taken in cohomology. If the BRST scaffold fails (e.g. a nonzero measure Jacobian Δ_{meas} in eq. (12.84) or loss of nilpotency), the candidate is excluded at Stage I (Principle 6.2) before any quadrant assignment.

Proposition 6.227 (Anomaly filter and quadrant admissibility)**.** *If a candidate gauge law for G exhibits a BRST anomaly—i.e. nilpotency or BRST invariance fails, or the functional measure has a nonzero Jacobian Δ_{meas} as in eq. (12.84)—then the law-predicate $P_{\mathrm{Law}}(G)$ is undefined at the physical level and the dual dyad product (6.281) cannot be formed within the admissible pipeline. Equivalently, by the Holosymmetric Selection Principle (Stage I = BRST consistency; Principle 6.2 and Proposition 6.217), such a candidate is excluded* prior *to any quadrant assignment. Hence the $\mathbb{Z}_2\times\mathbb{Z}_2$ orbit of Proposition 6.225 classifies only* BRST-consistent *laws; anomalies remove elements from consideration rather than creating new statuses.*

Proof. By Proposition 6.217, a nontrivial local measure class $[\mathcal{A}] \in H^{+1}(\hat{Q}_{\mathrm{BRST}})$ (or failure of $\hat{Q}^2_{\mathrm{BRST}} = 0$ / $[\hat{Q}_{\mathrm{BRST}}, \hat{H}_{\mathrm{tot}}] = 0$) violates Stage I (Principle 6.2); equivalently $\Delta_{\mathrm{meas}} \neq 0$ in eq. (12.84). Then the BRST-physical algebra/space used to *read* the predicates (Remark 6.226)—namely $\mathcal{A}^{(G)}_{\mathrm{phys}}$ and $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$—is not well-defined. Consequently $P_{\mathrm{Law}}(G)$ and $P_{\mathrm{State}}(G)$ admit no physical evaluation, so the joint status (and hence the dyad product (6.281)) lies outside the domain of the lifts. The group action of Proposition 6.225 remains intact on the *admissible* set (BRST-consistent laws) and simply does not act on anomalous candidates. □

Corollary 6.228 (Duality drives the quadrants; BRST fixes the reading)**.** *The quartet* $(\mathsf{Th}, \mathsf{Th}), (\mathsf{Th}, \mathsf{AntiTh}), (\mathsf{AntiTh}, \mathsf{Th}), (\mathsf{AntiTh}, \mathsf{AntiTh})$ *of Proposition 6.225 arises* solely *from the Axiom of Duality and its lifts (Definitions 6.220 and 6.221). The BRST/ghost scaffold neither adds nor subtracts statuses; it* clarifies *how predicates are to be checked—namely, cohomologically, as in Remark 6.226—and prunes anomaly-tainted candidates via Stage I (Principle 6.2 and Proposition 6.217). In particular, Quadrants I–II (law-symmetric) inherit exact BRST identities; Quadrants III–IV (law-asymmetric) remain admissible only when explicit breakings preserve BRST invariance (e.g. spurionic or non-gauge breakings), else they fail Stage I. Standard diagnostics (Ward/Nielsen identities, order parameters) are then understood modulo gauge redundancy [82, 91].*

Remark 6.229 (Provisional mode identifications)**.** Consistent with our Chapter 7 policy, any mapping of fluctuation modes (Higgs-like, graviton-like, apeiron, chrono–kymon) is *provisional* until multi-quadrant plateau-stability is established. The present subsection dictates *how* symmetry/asymmetry is read in gauge sectors (cohomologically; Remark 6.226); it does not assert mode identifications.

Definition 6.230 (Laws–State Symmetry–Asymmetry Duality)**.** Let $\{\mathsf{Th}, \mathsf{AntiTh}\}_{\text{Law}} \times \{\mathsf{Th}, \mathsf{AntiTh}\}_{\text{State}}$ be the joint status space for a fixed global G. The *Laws–State Symmetry–Asymmetry Duality* is the $\mathbb{Z}_2 \times \mathbb{Z}_2$ action generated by the commuting involutions Δ_{Law} and Δ_{State} of Proposition 6.225, which are functorial lifts (Lemma 6.222) of the status involution $\Delta_{\mathcal{C}}$ in Definition 6.221. In the presence of gauge redundancy (with gauge group $\mathcal{G}$), the predicates for law and state are *evaluated cohomologically* as in Section 6.11.3: $P_{\text{Law}}(G)$ is checked on the BRST-physical algebra $\mathcal{A}^{(G)}_{\text{phys}} = H^0(\mathcal{A}^{(G)}_{\text{BRST}}, \delta_{\text{BRST}})$, and $P_{\text{State}}(G)$ on the BRST-closed vacuum class $[\Omega] \in \mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$ (cf. Chapter 11: Section 12.3 and Principle 12.1; see also [104, 105, 106, 91, 82]). Only BRST-consistent candidate laws (nilpotency and measure invariance) enter the domain of this dual action; anomalous candidates fail Stage I of the HSP and are excluded a priori (cf. Proposition 6.227). This terminology bundles together the four regimes of Corollary 6.232 and will be used to subsume SSB, emergent/accidental symmetry, and explicit breaking within a single duality framework [102].

Theorem 6.231 (Quadrant Schema = Dyadic Product)**.** *Under the Axiom of Duality together with the status-lift $\Delta_{\mathcal{C}}$ and its self-similar lifts to the law and state aspects, the possible joint statuses of $(P_{\text{Law}}(G), P_{\text{State}}(G))$ are exactly the four in (6.281). These identify with the entries of the Symmetry–Asymmetry matrix in table 6.1; in the terminology of Definition 6.230, they are the four orbits of the Laws–State Duality.*

Proof sketch. Duality forces a two-point status space on each axis. Lifting to both axes and taking the product yields four corners; closure and scale-freedom (Lemma 6.222) forbid extra statuses or leaks under reindexings. In gauge theories, the predicates are interpreted in BRST cohomology (Section 6.11.3), which *restricts the domain* to BRST-consistent candidates but does not alter the $\mathbb{Z}_2 \times \mathbb{Z}_2$ product itself; gauge-parameter artifacts cancel by the quartet mechanism and Nielsen identities [91, 82]. Identification with the physical quadrants follows by the diagnostics in Corollary 6.232. □

6.11.4 Identification of the Quadrants and Mechanisms

Corollary 6.232 (Quadrant identification)**.** *With G global, the four joint statuses correspond to:*

(i) (Law Th, State Th) Unbroken
Symmetric law and symmetric state; unique symmetric vacuum, Ward identities exact, order parameters zero. Topological/SPT structure can still classify phases [437].
BRST reading: *Slavnov–Taylor/BRST identities hold exactly; the physical sector is $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$ and observables are BRST classes in $\mathcal{A}^{(G)}_{\text{phys}}$ ([91, 82]).*

(ii) (Law Th, State AntiTh) Spontaneous Symmetry Breaking
Symmetric law but asymmetric state; vacuum manifold $\mathcal{M}_{\text{vac}} \cong G/H$, nonzero order parameter, Nambu–Goldstone modes for broken global *continuous generators [92, 94, 91].*
BRST reading: *There is no literal* gauge *SSB; would-be Goldstones eaten by gauge bosons and ghost/longitudinal sectors form BRST-trivial quartets. Physical predictions are gauge-parameter independent (Nielsen identities), read in cohomology [82, 91].*

(iii) (Law AntiTh, State Th) Emergent/Accidental Symmetry
Asymmetric micro-law but symmetric IR law after coarse-graining: irrelevant breakings flow to zero, so violations shrink with scale and the state is symmetric with respect to the enlarged IR group [102, 82].
BRST reading: *Explicit breakings must preserve BRST nilpotency (treat as spurions where appropriate). BRST remains exact so that the IR Gauss constraints and cohomology are well-defined; otherwise Stage I excludes the candidate.*

(iv) (Law AntiTh, State AntiTh) Explicit Breaking
Asymmetric couplings select a unique ground state; no exact Goldstones (at most pseudo-NG with masses tied to the breaking) [91]. Residual subgroups may remain; responses track the breaking terms.
BRST reading: *Allowed only if the explicit breaking does not violate gauge redundancy (no BRST anomaly). Open or field-dependent algebras are handled via BV while preserving the quantum master equation in the gauge sector (see Chapter 11).*

Lemma 6.233 (Relational Ward Identity with Explicit Breaking Sources (BRST-aware))**.** *Let G be a continuous* global *symmetry with generators Q_a and Noether currents $\hat{J}^\mu_a$ for the operator-bundle fields. Write the generic field multiplet as X^I (collecting Gaiamiti/Tantrakyma components; hats suppressed inside functional derivatives), and let ∇_μ be the split covariant derivative eq. (6.314) compatible with $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$. Suppose the gauge-fixed local density is organized as*

$$\mathcal{L} \;=\; \underbrace{\mathcal{L}_0}_{G\text{-symmetric up to } \nabla\cdot} \;+\; \sum_k \epsilon_k\, \mathcal{O}_k \;+\; \zeta\,\Psi,$$

with $\delta_a \mathcal{L}_0 = \nabla_\mu K^\mu_a$, $\delta_a \mathcal{O}_k = \mathcal{S}_{k,a}$, and ζ the nilpotent BRST differential acting on fields/ghosts (so $\delta_a(\zeta\Psi) = \zeta(\delta_a\Psi)$; cf. Chapter 11 and [105, 106, 82, 91]). Then, taking relational expectations in $|\mathbf{\Psi}\rangle$, there exists a (Belinfante/BRST-)improved current $\hat{J}^\mu_{a,\mathrm{imp}}$ such that

$$\nabla_\mu \bar{J}^\mu_{a,\mathrm{imp}}(x) \;=\; \sum_k \epsilon_k \left\langle \mathcal{S}_{k,a}(x) \right\rangle_{\mathbf{\Psi}} \;+\; \left\langle \zeta\, \eta_a(x) \right\rangle_{\mathbf{\Psi}}, \qquad \bar{J}^\mu_{a,\mathrm{imp}} := \langle \hat{J}^\mu_{a,\mathrm{imp}} \rangle_{\mathbf{\Psi}}, \tag{6.282}$$

for some local functional η_a depending on the gauge-fixing fermion Ψ and the G-variation. On the BRST-physical sector (operators in $H^0(\hat{Q}_{\mathrm{BRST}})$ and BRST-closed $|\mathbf{\Psi}\rangle$), the ζ-exact contribution drops, yielding

$$\nabla_\mu \bar{J}^\mu_{a,\mathrm{imp}}(x) \;=\; \sum_k \epsilon_k \left\langle \mathcal{S}_{k,a}(x) \right\rangle_{\mathbf{\Psi}} \quad \textit{in} \quad \mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}}), \tag{6.283}$$

which is consistent with the scale-tagged identity eq. (6.375) and equivalent (modulo notation) to eq. (6.336). If G is gauged *(local parameter), (6.282) upgrades to a Slavnov–Taylor identity with the anomaly term vanishing in anomaly-free theories [82, 91]. In particular:*

(a) *If the law is symmetric ($\epsilon_k = 0$), BRST exactness and nilpotency imply exact covariant conservation $\nabla_\mu \bar{J}^\mu_{a,\mathrm{imp}} = 0$ on $\mathcal{H}_{\mathrm{phys}}$* (Quadrants I–II) *[90, 82, 409].*

(b) *If the breaking couplings are RG-irrelevant so that $\epsilon_k(\mu) \to 0$ as $\mu \downarrow 0$, the violations in (6.283) are $\mathcal{O}(\epsilon_k(\mu))$* (Quadrant III) *[102, 82].*

(c) *If some ϵ_k remain nonzero in the IR, non-conservation persists and Goldstone theorems do not apply; pseudo-Nambu–Goldstone masses scale $m^2 \propto \epsilon$* (Quadrant IV) *[91].*

Remark 6.234 (Meaning of the symmetric sector $\mathcal{L}_0$ and the BRST-exact layer)**.** Throughout the decomposition

$$\mathcal{L} \;=\; \mathcal{L}_0 \;+\; \sum_k \epsilon_k\, \mathcal{O}_k \;+\; \zeta\Psi, \qquad \delta_a \mathcal{L}_0 = 0 \text{ (up to } \nabla\cdot), \quad \delta_a \mathcal{O}_k = \mathcal{S}_{k,a},$$

$\mathcal{L}_0(\mathsf{X}, \nabla\mathsf{X})$ denotes the *G-symmetric* part of the local law (the "unbroken sector") in the sense that $\delta_a \mathcal{L}_0 = \nabla_\mu K^\mu_a$, so the action $S[\mathsf{X}] = \int \sqrt{-\det \bar{\Sigma}}\, \mathcal{L}_0$ is invariant under constant parameters (with standard boundary conditions) [90, 435, 82]. Gauge fixing and ghosts enter only via the BRST-exact term $\zeta\Psi$, which does not modify physical Ward identities: $\delta_a(\zeta\Psi) = \zeta(\delta_a\Psi)$ contributes the ζ-exact piece in (6.282), which vanishes when projected to $\mathcal{H}_{\text{phys}}$ (Chapter 11; [105, 106]). Terms that *explicitly* break G are collected as $\sum_k \epsilon_k \mathcal{O}_k$ with small couplings ϵ_k. Equivalently (spurion viewpoint), if one assigns transformation rules to ϵ_k so that $\epsilon_k \mathcal{O}_k$ is G-invariant, those products may be counted inside $\mathcal{L}_0$; "freezing" the spurions to fixed background values reproduces explicit breaking [91, 440]. This bookkeeping underlies Lemma 6.233: promoting $\alpha_a \to \alpha_a(x)$ produces only a covariant divergence from $\mathcal{L}_0$ (absorbed into $\hat{J}^\mu_{a,\text{imp}}$), while the explicit breakers yield the source term on the right-hand side, and the BRST-exact layer $\zeta\Psi$ cancels in physical correlators. In RG terms, if $\epsilon_k(\mu) \to 0$ as $\mu \downarrow 0$ (irrelevant), the violation of covariant conservation is $\mathcal{O}(\epsilon_k(\mu))$; if some ϵ_k remain finite in the IR, non-conservation persists and pseudo-Nambu–Goldstone masses scale as $m^2 \propto \epsilon$ [102, 91].

Relational Ward Identity with Explicit Breaking Sources (BRST-aware). **1) Set-up.** Take the general action

$$S[\mathsf{X}] \;=\; \int d^d x\, \sqrt{-\det \bar{\Sigma}}\, \mathcal{L}(\mathsf{X}, \nabla\mathsf{X})\,,$$

with the *gauge-fixed* density organized as

$$\mathcal{L} \;=\; \mathcal{L}_0 \;+\; \sum_k \epsilon_k\, \mathcal{O}_k \;+\; \zeta\,\Psi,$$

where ζ is the nilpotent BRST differential and Ψ the gauge fermion (Chapter 11; [105, 106, 82, 91]). Let ∇_μ be the split covariant derivative eq. (6.314) compatible with $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$. Covariant integration by parts (IBP) reads

$$\int d^d x\, \sqrt{-\det \bar{\Sigma}}\, V^\mu \nabla_\mu f \;=\; -\int d^d x\, \sqrt{-\det \bar{\Sigma}}\, f\, \nabla_\mu V^\mu, \tag{6.284}$$

for smooth compactly supported f.

2) Local parameter (symmetric part). Promote the constant parameter to $\alpha_a(x)$ and vary the G-symmetric part first:

$$\delta_a\Big(\sqrt{-\det \bar{\Sigma}}\, \mathcal{L}_0\Big) = \sqrt{-\det \bar{\Sigma}}\, \Big(E_I\, \delta_a \mathsf{X}^I + \nabla_\mu \theta^\mu_a\Big), \tag{6.285}$$

where $E_I := \frac{\partial \mathcal{L}_0}{\partial \mathsf{X}^I} - \nabla_\mu \frac{\partial \mathcal{L}_0}{\partial(\nabla_\mu \mathsf{X}^I)}$ and

$$\theta^\mu_a := \frac{\partial \mathcal{L}_0}{\partial(\nabla_\mu \mathsf{X}^I)}\, \delta_a \mathsf{X}^I \quad \text{(symplectic potential current density)}.$$

With $\delta_a \mathsf{X}^I = \alpha_a R_a^I(\mathsf{X})$, factor α_a:

$$\theta_a^\mu = \alpha_a \, J_a^\mu, \qquad J_a^\mu := \frac{\partial \mathcal{L}_0}{\partial(\nabla_\mu \mathsf{X}^I)} \, R_a^I(\mathsf{X}). \tag{6.286}$$

Thus

$$\delta_a\Big(\sqrt{-\det \bar{\Sigma}}\, \mathcal{L}_0\Big) = \sqrt{-\det \bar{\Sigma}}\, \Big((\nabla_\mu \alpha_a)\, J_a^\mu + \alpha_a (E_I R_a^I + \nabla_\mu J_a^\mu) \Big). \tag{6.287}$$

3) Off-shell Noether identity. For constant α_a, $\delta_a \mathcal{L}_0 = 0$ (strictly or up to a covariant divergence), hence

$$E_I R_a^I + \nabla_\mu J_a^\mu \equiv 0, \tag{6.288}$$

and (6.287) reduces to

$$\delta_a\Big(\sqrt{-\det \bar{\Sigma}}\, \mathcal{L}_0\Big) = \sqrt{-\det \bar{\Sigma}}\, (\nabla_\mu \alpha_a)\, J_a^\mu. \tag{6.289}$$

4) Add explicit breaking and the BRST-exact layer. For the full $\mathcal{L} = \mathcal{L}_0 + \sum_k \epsilon_k \mathcal{O}_k + \zeta \Psi$,

$$\delta_a\Big(\sqrt{-\det \bar{\Sigma}}\, \mathcal{L}\Big) = \sqrt{-\det \bar{\Sigma}}\, (\nabla_\mu \alpha_a)\, J_a^\mu + \sqrt{-\det \bar{\Sigma}}\, \alpha_a \sum_k \epsilon_k \, \mathcal{S}_{k,a} + \delta_a\Big(\sqrt{-\det \bar{\Sigma}}\, \zeta \Psi\Big).$$

Since G is *global*, the measure is ∇-compatible and δ_a acts only on fields. Using BRST exactness and locality, decompose the variation of the gauge fermion as

$$\delta_a \Psi \;=\; \alpha_a \, \chi_a \;+\; (\nabla_\mu \alpha_a)\, \Lambda_a^\mu, \qquad \Rightarrow \qquad \delta_a(\zeta\Psi) = \zeta(\delta_a \Psi) = \alpha_a \, \zeta \chi_a + (\nabla_\mu \alpha_a)\, \zeta \Lambda_a^\mu, \tag{6.290}$$

for some local functionals χ_a, Λ_a^μ (this is always achievable by integration-by-parts rearrangements; [105, 106, 82]). Integrating and applying (6.284) to both gradient terms yields

$$\delta_a S = -\int d^d x \, \sqrt{-\det \bar{\Sigma}}\, \alpha_a \, \nabla_\mu (J_a^\mu - \zeta \Lambda_a^\mu) \;+\; \int d^d x \, \sqrt{-\det \bar{\Sigma}}\, \alpha_a \Big(\sum_k \epsilon_k \, \mathcal{S}_{k,a} + \zeta \chi_a \Big). \tag{6.291}$$

Define the *BRST-improved* Noether current

$$J^\mu_{a,\mathrm{imp}} \;:=\; J_a^\mu - \zeta \Lambda_a^\mu, \tag{6.292}$$

which absorbs the gauge-fixing dependence into a BRST-exact improvement.

5) Operator identity and projection to the physical sector. Arbitrary $\alpha_a(x)$ implies the operator identity

$$\nabla_\mu \hat{J}^\mu_{a,\mathrm{imp}} \;=\; \sum_k \epsilon_k \, \mathcal{S}_{k,a} \;+\; \zeta \, \eta_a, \qquad \eta_a := \chi_a, \tag{6.293}$$

up to further improvements $\hat{J}^\mu_{a,\mathrm{imp}} \to \hat{J}^\mu_{a,\mathrm{imp}} + \nabla_\nu \hat{X}_a^{[\mu\nu]}$. Taking relational expectations in a ζ-closed state $|\mathbf{\Psi}\rangle$ and using BRST invariance of the measure,

$$\langle \zeta \, \mathcal{O} \rangle_{\mathbf{\Psi}} = 0 \quad \text{for all local } \mathcal{O} \quad \text{(Kugo–Ojima/quartet mechanism; [82, 91])},$$

we obtain

$$\nabla_\mu \bar{J}^\mu_{a,\mathrm{imp}}(x) \;=\; \sum_k \epsilon_k \left\langle \mathcal{S}_{k,a}(x) \right\rangle_{\boldsymbol{\Psi}},$$

which is exactly eq. (6.282) (and hence eq. (6.283)) with the improved current.

6) Comment on gauged G. If G is *gauged* (local parameter promoted to a gauge transformation), the same steps give a Slavnov–Taylor identity in place of (6.293), with an additional anomaly density $\mathcal{A}_a$ from the path-integral measure. In anomaly-free theories $\mathcal{A}_a = 0$ and the BRST-exact term still projects out on $\mathcal{H}_{\mathrm{phys}}$ [82, 91]. □

Remark 6.235 (Gauge vs. global)**.** "Gauge symmetry'' is redundancy; BEH phases are diagnosed by gauge-invariant observables, not literal gauge-SSB. The lemma and proof above concern *global* G. If G is gauged, the global Ward identity is replaced by a Slavnov–Taylor (BRST) identity; gauge-fixing/ghost contributions are ζ-exact and vanish upon projection to $\mathcal{H}_{\mathrm{phys}}$, and anomaly freedom is required (otherwise the Stage I gate excludes the theory). All quadrant conclusions are then read in BRST cohomology and are gauge-parameter independent [435, 82, 91].

6.11.5 Holosymmetric Meta-Layer: Two-path Law/State Interferometry

Idea. Holosymmetry provides an operational "meta-layer" that sits *over* the 2×2 matrix of table 6.1 (constructed in Section 6.11) and prescribes a coherent, two-path interrogation of each cell. The lift is functorial over the triadic channel split (Definition 5.54), preserves effective causal cones (Definition 5.123 and Proposition 5.124), and modifies only intra-cone phase transport (Corollary 5.126); hence it sharpens diagnostics without moving the system between quadrants.

Remark 6.236 (BRST-clean interferometry and gauge-fermion independence)**.** When gauge redundancy is present, all interferometric readouts are *evaluated in the BRST-physical sector*: $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$, with operators taken modulo ζ-exact terms (Chapter 11; Section 6.11.3 and Principle 12.1). We assume the gauge-fixed density has the standard form $\mathcal{L} = \mathcal{L}_0 + \sum_k \epsilon_k \mathcal{O}_k + \zeta\Psi$, and that the branch evolutions are BRST-compatible, $\zeta U_{A/B}(\tau) = 0$ (equivalently $[\hat{Q}_{\mathrm{BRST}}, U_{A/B}(\tau)] = 0$), or equal up to ζ-exact improvements. Any change of gauge-fermion $\Psi \mapsto \Psi + \delta\Psi$ produces only ζ-exact variations in the overlaps; concretely, there exist local functionals $\eta_\xi(\tau)$ (depending on the gauge parameter ξ) such that

$$\partial_\xi \Lambda_{\mathrm{law}}(\tau) = \left\langle \zeta\, \eta^{(\mathrm{law})}_\xi(\tau) \right\rangle = 0, \qquad \partial_\xi \Lambda_{\mathrm{state}}(\tau) = \left\langle \zeta\, \eta^{(\mathrm{state})}_\xi(\tau) \right\rangle = 0 \quad \text{in } \mathcal{H}_{\mathrm{phys}}. \tag{6.294}$$

If a gauge anomaly spoils BRST nilpotency, the Stage I gate (Chapter 11; Proposition 6.227) excludes the candidate and the interferometric functionals are not physically defined.

Definition 6.237 (Two-path *law*-interferometry (parameter-agnostic))**.** Prepare a control $|+\rangle = (|0\rangle + |1\rangle)/\sqrt{2}$ and, conditional on the control, evolve under two admissible law-level evolutions $U_A(s), U_B(s)$ along an evolution parameter s that is either (i) a cone-compatible relational time τ (see Definition 5.123 and Corollary 5.126) or (ii) an RG scale ℓ (coarse-graining flow):

$$U_{\mathrm{ctrl}}(s) = |0\rangle\langle 0| \otimes U_A(s) + |1\rangle\langle 1| \otimes U_B(s).$$

For initial state ρ_0, the control readout yields the *law-coherence functional*

$$\Lambda_{\mathrm{law}}(s) \coloneqq \mathrm{Tr}\big[U_B^\dagger(s)\, U_A(s)\, \rho_0\big] \tag{6.295}$$

$$\langle \sigma_x \rangle = \mathrm{Re}\, \Lambda_{\mathrm{law}}(s) \tag{6.296}$$

$$\langle \sigma_y \rangle = \mathrm{Im}\, \Lambda_{\mathrm{law}}(s) \tag{6.297}$$

BRST overlay: evaluate (6.295) on $\mathcal{H}_{\mathrm{phys}}$ with $U_{A/B}^{\mathrm{phys}}(s) = \mathsf{P}_{\mathrm{phys}}\, U_{A/B}(s)\, \mathsf{P}_{\mathrm{phys}}$ and $[\hat{Q}_{\mathrm{BRST}}, U_{A/B}(s)] = 0$ (up to ζ-exact improvements). Variations of Ψ or gauge parameters ξ shift Λ_{law} by ζ-exact terms that vanish in expectation as in (6.294).

Definition 6.238 (Two-path *state*-interferometry (parameter-agnostic))**.** *Conventions.* Here τ denotes the interferometer's control/evolution parameter (disjoint from the duality primitive $s \in \mathcal{U}_{\mathcal{C}}$ used in Section 6.11).

With the law fixed to $U(\tau)$ on both branches, coherently prepare $|\Omega_1\rangle, |\Omega_2\rangle$:

$$|+\rangle \otimes |\psi_0\rangle \;\mapsto\; \tfrac{1}{\sqrt{2}}(|0\rangle \otimes |\Omega_1\rangle + |1\rangle \otimes |\Omega_2\rangle)\,.$$

After parallel evolution and recombination, the control visibility measures the *state-coherence functional*

$$\Lambda_{\mathrm{state}}(\tau) \coloneqq \langle \Omega_2|\, U^\dagger(\tau)\, U(\tau)\, |\Omega_1\rangle, \qquad (\text{for stationary vacua, } \Lambda_{\mathrm{state}} = \langle \Omega_2|\Omega_1\rangle). \tag{6.298}$$

BRST overlay: require $|\Omega_i\rangle \in \mathcal{H}_{\mathrm{phys}}$ to be ζ-closed, ghost number zero; replace $U \mapsto U^{\mathrm{phys}}$ as above. Gauge-fermion and gauge-parameter independence follow from (6.294). If needed, improve insertions by ζ-exact terms to ensure BRST closure without changing the value in $\mathcal{H}_{\mathrm{phys}}$.

Remark 6.239 (Relational-time instantiation (Chapter 6))**.** Let τ parametrize a cone-compatible history (Definition 5.123). Take smeared generators

$$\hat{\mathcal{H}}_k(\tau) = \int d^d x\, J^{\mu\nu}(x)\, \hat{\Sigma}^{(k)}_{\mu\nu}(x, \tau)$$

with real $J^{\mu\nu}$ and define

$$U_k(\tau) = \mathcal{T}_\tau \exp\!\Big(-i \int d\tau\, \hat{\mathcal{H}}_k(\tau)\Big), \qquad \Lambda_{\mathrm{law}}(\tau) = \langle \Psi_{\mathrm{RR}}|\, \hat{U}_1^\dagger(\tau)\, \hat{U}_0(\tau)\, |\Psi_{\mathrm{RR}}\rangle,$$

the *relational overlap* of the two chronogeometric histories.

BRST overlay. Work on $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ and choose the generators cohomologically: pick representatives so that $\zeta \hat{\mathcal{H}}_k(\tau) = 0$ *in cohomology* (i.e. up to ζ-exact improvements), equivalently $[\hat{Q}_{\mathrm{BRST}}, \hat{\mathcal{H}}_k(\tau)] = \zeta Y_k(\tau)$ for some local Y_k. Then the induced evolution descends to $\mathcal{H}_{\mathrm{phys}}$ and is independent of the gauge fermion Ψ by eq. (6.294). Cone preservation and intra-cone phase transport follow from Definition 5.123, Proposition 5.124, and Corollary 5.126.

Remark 6.240 (RG-scale instantiation (scale parameter ℓ))**.** Let $U_k(\ell)$ denote the coarse-grained evolution (effective law) at RG scale ℓ (e.g. via the band/IR functor of eq. (7.40)). Define

$$\Lambda_{\mathrm{law}}(\ell) = \mathrm{Tr}\Big[U_B^\dagger(\ell)\, U_A(\ell)\, \rho_0\Big], \qquad \Lambda_{\mathrm{state}}(\ell) = \langle \Omega_2(\ell) \mid \Omega_1(\ell)\rangle.$$

In emergent-symmetry regimes one typically finds $1-|\Lambda_{\mathrm{law}}(\ell)| \propto \ell^{\,\Delta-(d+z)}$, while $\Lambda_{\mathrm{state}}(\ell) \approx 1$ (no order parameter).

BRST overlay. Work on $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ and take laws/states modulo ζ-exact improvements. RG reparametrisations of the gauge-fixing (e.g. $\xi \mapsto \xi(\ell)$) shift the overlaps by ζ-exact terms, hence

$$\partial_\xi \Lambda_{\mathrm{law}}(\ell) = \Big\langle \zeta\, \eta_\xi^{(\mathrm{law})}(\ell)\Big\rangle = 0, \qquad \partial_\xi \Lambda_{\mathrm{state}}(\ell) = \Big\langle \zeta\, \eta_\xi^{(\mathrm{state})}(\ell)\Big\rangle = 0 \quad \text{in } \mathcal{H}_{\mathrm{phys}}$$

(cf. eq. (6.294)). Any anomaly (nonzero Δ_{meas} in eq. (12.84) or loss of nilpotency) violates this and is excluded by Stage I (Principle 6.2 and Proposition 6.227).

Remark 6.241 (Operator-insertion / instrument variant (optional))**.** When one wishes to display the tensors directly, one may use the controlled insertion

$$\mathbf{U}_{\mathrm{ctrl}}^{\mathrm{ins}} = |0\rangle\langle 0| \otimes \hat{V}\, \hat{\Sigma}^{(A_G B_N)}\, \hat{W} + |1\rangle\langle 1| \otimes \hat{V}\, \hat{\Sigma}^{(A_N B_G)}\, \hat{W}.$$

To keep the map norm-preserving, either: (i) take $\hat{V}, \hat{W}$ as parts of a Stinespring dilation [352, 441] so that $\mathbf{U}_{\mathrm{ctrl}}^{\mathrm{ins}}$ is the postselected branch of a unitary on system+environment, or (ii) normalise the readout by the total detection probability Z (instrument normalisation). In both cases the interference term is proportional to the same overlap functional as in Definition 6.237; the fully unitary embedding is recommended for avoiding normalisation subtleties. *BRST overlay:* require insertions to be BRST-closed (ghost number zero). If an insertion $\hat{\mathcal{O}}$ is not closed, replace it by a cohomologous $\hat{\mathcal{O}}_{\mathrm{imp}} = \hat{\mathcal{O}} - \zeta\hat{G}$ without changing physical readouts in $\mathcal{H}_{\mathrm{phys}}$.

Remark 6.242 (State preparation as an isometry)**.** The mapping in Definition 6.238 can be implemented by an isometry $W : \mathrm{span}\{|0\rangle, |1\rangle\} \to \mathcal{H}_C \otimes \mathcal{H}_S$ with $W|0\rangle = |0\rangle \otimes |\Omega_1\rangle$ and $W|1\rangle = |1\rangle \otimes |\Omega_2\rangle$, so that $W|+\rangle = \frac{1}{\sqrt{2}}(|0\rangle \otimes |\Omega_1\rangle + |1\rangle \otimes |\Omega_2\rangle)$. This makes linearity explicit and covers mixed ensembles via purification if needed. *BRST overlay:* choose $|\Omega_i\rangle$ to be ζ-closed representatives of ghost-number-zero classes in $\mathcal{H}_{\mathrm{phys}}$; different choices within a class change only by ζ-exact vectors and do not affect Λ_{state}.

Lemma 6.243 (Gauge-parameter independence of interferometric visibilities (Nielsen-type))**.** *Assume the gauge-fixed density has the form $\mathcal{L} = \mathcal{L}_0 + \sum_k \epsilon_k \mathcal{O}_k + \zeta\Psi(\xi)$ with gauge parameter(s) ξ, and that the branch evolutions $U_{A/B}(\tau,\xi)$ are BRST-compatible, i.e. $[\hat{Q}_{\mathrm{BRST}}, U_{A/B}(\tau,\xi)] = 0$ up to ζ-exact improvements. For any BRST-closed initial state and readout in the physical sector ($|\cdot\rangle \in \mathcal{H}_{\mathrm{phys}}$, ghost number 0), the law- and state-coherence functionals of Definitions 6.237 and 6.238 obey*

$$\partial_\xi \Lambda_{\mathrm{law}}(\tau) = \Big\langle \zeta\, \eta_\xi^{(\mathrm{law})}(\tau)\Big\rangle = 0, \qquad \partial_\xi \Lambda_{\mathrm{state}}(\tau) = \Big\langle \zeta\, \eta_\xi^{(\mathrm{state})}(\tau)\Big\rangle = 0, \tag{6.299}$$

with local functionals $\eta_\xi^{(\mathrm{law})}$ and $\eta_\xi^{(\mathrm{state})}$ constructed from $\partial_\xi\Psi$ and the BRST-compatible generators of $U_{A/B}$ and U.

Proof. Since $\partial_\xi \mathcal{L} = \zeta(\partial_\xi \Psi)$, the ξ-variation of any physical expectation inserts a ζ-exact operator (standard BRST/Nielsen logic). The explicit ξ-dependence of $U_{A/B}(\tau,\xi)$ and $U(\tau,\xi)$ enters only through the gauge-fixing sector, hence through ζ-exact deformations of their generators. Therefore

$$\partial_\xi\Big[U_B^\dagger(\tau,\xi)\,U_A(\tau,\xi)\Big] = \zeta\,\Gamma_\xi^{(\mathrm{law})}(\tau), \qquad \partial_\xi\Big[U^\dagger(\tau,\xi)\,U(\tau,\xi)\Big] = \zeta\,\Gamma_\xi^{(\mathrm{state})}(\tau),$$

for local Γ_ξ built from $\partial_\xi\Psi$ and cohomologically BRST-invariant Hamiltonians. Taking the trace or overlap in $\mathcal{H}_{\mathrm{phys}}$ gives (6.299) with $\eta_\xi^{(\cdot)}$ of ghost number -1. BRST-closed states and a BRST-invariant measure (Stage I; see Principle 6.2 and eq. (12.84)) imply $\langle\zeta(\,\cdot\,)\rangle = 0$, hence both derivatives vanish. □

Remark 6.244 (Explicit construction of the Nielsen generators for (6.299))**.** Write $\mathrm{Ad}_U(X) := U\,X\,U^{-1}$ and fix a reference gauge parameter ξ_0. For each branch $k \in \{A,B\}$ choose a smooth homotopy $t \in [0,1] \mapsto U_k(\tau,t)$ with $U_k(\tau,0) = U_k(\tau,\xi_0)$ and $U_k(\tau,1) = U_k(\tau,\xi)$ such that there exists a local, ghost-number -1 functional $\Lambda_\xi^{(k)}(t)$ obeying

$$\partial_t U_k(\tau,t) = U_k(\tau,t)\,\zeta\,\Lambda_\xi^{(k)}(t).$$

Set $\chi_\xi := \partial_\xi\Psi$ and define

$$\eta_\xi^{(\mathrm{law})}(\tau) := \chi_\xi + \int_0^1 dt\,\Big[\mathrm{Ad}_{U_B^\dagger(\tau,t)}\,\Lambda_\xi^{(B)}(t) - \mathrm{Ad}_{U_A^\dagger(\tau,t)}\,\Lambda_\xi^{(A)}(t)\Big]. \tag{6.300}$$

Then $\partial_\xi\Lambda_{\mathrm{law}}(\tau) = \langle\zeta\,\eta_\xi^{(\mathrm{law})}(\tau)\rangle = 0$ on $\mathcal{H}_{\mathrm{phys}}$.

For the state functional, choose a homotopy $U(\tau,t)$ with $\partial_t U(\tau,t) = U(\tau,t)\,\zeta\,\Lambda_\xi^{(U)}(t)$ and, whenever convenient, pick BRST-cohomology representatives $|\Omega_i\rangle$ independent of ξ.[6] Define

$$\eta_\xi^{(\mathrm{state})}(\tau) := \chi_\xi + \int_0^1 dt\,\mathrm{Ad}_{U^\dagger(\tau,t)}\,\Lambda_\xi^{(U)}(t) \quad \Big(\text{plus } B_\xi^{(\mathrm{state})}(\tau) \text{ if } |\Omega_i| \text{ vary}\Big), \tag{6.301}$$

so that $\partial_\xi\Lambda_{\mathrm{state}}(\tau) = \langle\zeta\,\eta_\xi^{(\mathrm{state})}(\tau)\rangle = 0$ in $\mathcal{H}_{\mathrm{phys}}$.

Lemma 6.245 (Control readout = overlap (law layer))**.** *Let the controlled unitary be as in Definition 6.237 and initialise the control in $|+\rangle = (|0\rangle + |1\rangle)/\sqrt{2}$ and the system in ρ_0. Then*

$$\langle\sigma_x\rangle = \mathrm{Re}\;\Lambda_{\mathrm{law}}(s), \qquad \langle\sigma_y\rangle = \mathrm{Im}\;\Lambda_{\mathrm{law}}(s), \qquad \Lambda_{\mathrm{law}}(s) = \mathrm{Tr}\Big[U_B^\dagger(s)\,U_A(s)\,\rho_0\Big].$$

BRST overlay: *In the presence of gauge redundancy, evaluate in the physical sector: $U_k^{\mathrm{phys}} := \mathsf{P}_{\mathrm{phys}} U_k \mathsf{P}_{\mathrm{phys}}$, $\rho_0^{\mathrm{phys}} := \mathsf{P}_{\mathrm{phys}}\rho_0\mathsf{P}_{\mathrm{phys}}$, with $[\hat{Q}_{\mathrm{BRST}}, U_k] = 0$ up to ζ-exact improvements. Then the same equalities hold with all objects replaced by their* phys *versions, and are gauge-parameter independent by Lemma 6.243.*

Proof. Evolve $|+\rangle\langle+| \otimes \rho_0$ by $U_{\mathrm{ctrl}}(s)$: $\rho' = \frac{1}{2}\sum_{i,j\in\{0,1\}} |i\rangle\langle j| \otimes U_i(s)\rho_0 U_j^\dagger(s)$. Tracing out the system gives $\rho'_C = \frac{1}{2}(\mathbb{I} + \mathrm{Re}\,\Lambda_{\mathrm{law}}\,\sigma_x + \mathrm{Im}\,\Lambda_{\mathrm{law}}\,\sigma_y)$. For the BRST overlay, insert $\mathsf{P}_{\mathrm{phys}}$ on both sides of each U_k and ρ_0; ζ-exact pieces drop in expectations on $\mathcal{H}_{\mathrm{phys}}$. □

[6] If $|\Omega_i\rangle$ vary with ξ, one may write $\partial_t|\Omega_i(t)\rangle = \zeta\,|\gamma_{i,\xi}(t)\rangle$ and add the boundary counterterm $B_\xi^{(\mathrm{state})}(\tau) := \zeta\int_0^1 dt\,\big(|\gamma_{2,\xi}(t)\rangle\langle\Omega_1| - |\Omega_2\rangle\langle\gamma_{1,\xi}(t)|\big)$; this keeps $\partial_\xi\Lambda_{\mathrm{state}}$ manifestly ζ-exact.

Lemma 6.246 (Control readout = overlap (state layer))**.** *Under Definition 6.238, with control initialised in* $|+\rangle$ *and* $U(s)$ *applied on both branches,*

$$\langle\sigma_x\rangle = \mathrm{Re}\ \Lambda_{\mathrm{state}}(s), \qquad \langle\sigma_y\rangle = \mathrm{Im}\ \Lambda_{\mathrm{state}}(s), \qquad \Lambda_{\mathrm{state}}(s) = \langle\Omega_2|\, U^\dagger(s)U(s)\, |\Omega_1\rangle.$$

For stationary vacua, $\Lambda_{\mathrm{state}} = \langle\Omega_2|\Omega_1\rangle$. BRST overlay: *Take* $|\Omega_i\rangle \in \mathcal{H}_{\mathrm{phys}}$ ζ*-closed, ghost number* 0*, and* $U^{\mathrm{phys}} := \mathsf{P}_{\mathrm{phys}} U \mathsf{P}_{\mathrm{phys}}$*. The same equalities hold and are gauge-parameter independent by Lemma 6.243.*

Proof. As in Lemma 6.245; replace $U_B^\dagger U_A$ by $U^\dagger U$ and insert $\mathsf{P}_{\mathrm{phys}}$ as above. □

Lemma 6.247 (Functoriality over the triadic split)**.** *Let* Π_α *denote the channel projectors for the triadic decomposition (Definition 5.54), chosen BRST-closed of ghost number* 0*. Then* $U_{\mathrm{ctrl}}(\tau)$ *is block-diagonal with respect to the split and*

$$\Pi_\alpha U_{\mathrm{ctrl}}(\tau) = U_{\mathrm{ctrl}}(\tau)\Pi_\alpha, \qquad \Lambda_\bullet(\tau) = \sum_{\alpha\in\{\mathrm{corr,vac,spec}\}} \Lambda_\bullet^{(\alpha)}(\tau),$$

with $\Lambda_{\mathrm{law}}^{(\alpha)}(\tau) = \mathrm{Tr}\big[U_{B,\alpha}^\dagger(\tau)\, U_{A,\alpha}(\tau)\, \rho_0\big]$ *and* $\Lambda_{\mathrm{state}}^{(\alpha)}(\tau) = \langle\Omega_2|\Pi_\alpha|\Omega_1\rangle$. BRST overlay: *all statements hold with* phys *projections inserted, since* $[\hat{Q}_{\mathrm{BRST}}, \Pi_\alpha] = 0$ *and* ζ*-exact improvements do not mix blocks.*

Proof. Block-diagonality follows from the definition of the split and the controlled form of U_{ctrl}, which preserves each channel. Thus $[\Pi_\alpha, U_{\mathrm{ctrl}}(\tau)] = 0$ and the visibility decomposes as a sum of diagonal contributions, giving the stated formulas. *BRST overlay:* Π_α commute with $\mathsf{P}_{\mathrm{phys}}$ and with U_k^{phys}, so additivity is preserved on $\mathcal{H}_{\mathrm{phys}}$. □

Lemma 6.248 (Cone preservation for the holosymmetric lift)**.** *Let effective cones be defined by the expectation tensor* $\bar{\Sigma}_{\mu\nu}$*. The holosymmetric lift preserves the branchwise cone assignment and cannot generate law-signalling; it alters only intra-cone phase transport.* BRST overlay: *since cones depend only on BRST-invariant expectation values of* $\bar{\Sigma}_{\mu\nu}$ *(ghost number* 0*), any* ζ*-exact change of gauge fixing leaves the cone structure invariant.*

Proof. By Definition 5.123, cones are fixed by $\bar{\Sigma}_{\mu\nu}$; Proposition 5.124 forbids law-signalling; and Corollary 5.126 states that only intra-cone phases change. *BRST overlay:* gauge-fixing changes are ζ-exact and do not affect BRST-invariant cone diagnostics. □

Lemma 6.249 (Stability of the law predicate under the lift)**.** *Let* $P_{\mathrm{Law}}(G)$ *assert* $[U_G,\, U_k(\tau)] = 0$ *for the representation* U_G *and both branches* $k \in \{A, B\}$*. Then*

$$[\, U_G \otimes \mathbb{I}_C,\ U_{\mathrm{ctrl}}(\tau)\,] = 0.$$

If $P_{\mathrm{Law}}(G)$ *fails on at least one branch, then* $[\, U_G \otimes \mathbb{I}_C,\ U_{\mathrm{ctrl}}(\tau)\,] \neq 0$.

BRST overlay: *Read* $P_{\mathrm{Law}}(G)$ *in the physical algebra:* $[U_G^{\mathrm{phys}}, U_k^{\mathrm{phys}}(\tau)] = 0$ *implies* $[\, U_G^{\mathrm{phys}} \otimes \mathbb{I}_C,\ U_{\mathrm{ctrl}}^{\mathrm{phys}}(\tau)\,] = 0$*; conversely, any violation that is nontrivial in* $H^0(\hat{Q}_{\mathrm{BRST}})$ *propagates to the controlled unitary (differences that are* ζ*-exact are cohomologically trivial).*

Proof. Write $U_{\mathrm{ctrl}}(\tau) = |0\rangle\langle 0| \otimes U_A(\tau) + |1\rangle\langle 1| \otimes U_B(\tau)$. If $[U_G, U_A(\tau)] = [U_G, U_B(\tau)] = 0$, then clearly $[\, U_G \otimes \mathbb{I}_C,\ U_{\mathrm{ctrl}}(\tau)\,] = 0$. If the commutator is nonzero on some branch, the same block appears in the controlled commutator, hence the total commutator is nonzero.

BRST overlay: Insert $\mathsf{P}_{\mathrm{phys}}$ on both sides and use that ζ-exact differences vanish in $H^0(\hat{Q}_{\mathrm{BRST}})$ (cf. Definition 6.209 and Lemma 6.210). □

Lemma 6.250 (Stability of the state predicate under the lift)**.** *Let $P_{\mathrm{State}}(G)$ assert "Ward identities are exact and all G-charged order parameters vanish in the thermodynamic limit." Embed any state ρ as $|+\rangle\langle +| \otimes \rho$ and apply the holosymmetric protocol (law- or state-interferometric).*

(i) If ρ is G-invariant, the reduced system state after tracing out the control remains G-invariant.

(ii) If ρ lies in an SSB phase, any coherent superposition of distinct vacua becomes block-diagonal between superselection sectors as $V \to \infty$ (vacuum overlap $\to 0$), so $P_{\mathrm{State}}(G)$ remains false in the thermodynamic limit even if some finite-volume order parameters partially average to zero.

BRST overlay: *Evaluate invariance and order parameters on $\mathcal{H}_{\mathrm{phys}}$ with ζ-closed, ghost-number-zero states; ζ-exact components do not affect reduced states or sector blocking.*

Proof. (i) G acts trivially on the control and by conjugation on the system. If ρ is G-invariant, then so is $|+\rangle\langle +| \otimes \rho$, and the partial trace over the control preserves invariance.

(ii) Standard SSB: distinct infinite-volume vacua are orthogonal, so coherent superpositions reduce to block-diagonal mixtures across superselection sectors as $V \to \infty$; the holosymmetric control does not alter this sector decomposition.

BRST overlay: Project to $\mathcal{H}_{\mathrm{phys}}$; ζ-exact parts vanish in physical expectations (Lemma 6.210) and do not change sector structure. □

Proposition 6.251 (Holosymmetric overlay preserves quadrant membership)**.** *Let a cell of table 6.1 be fixed by the joint truth values of the law- and state-predicates in Section 6.11. Assume an anomaly-free, BRST-consistent gauge fixing with density $\mathcal{L} = \mathcal{L}_0 + \sum_k \epsilon_k \mathcal{O}_k + \zeta\Psi$ and branch evolutions $U_{A/B}(\ell)$ that are BRST-compatible ($[\hat{Q}_{\mathrm{BRST}}, U_{A/B}] = 0$ up to ζ-exact improvements). Under a holosymmetric lift that is functorial over the triadic split and cone-preserving (Definitions 5.54 and 5.123, Proposition 5.124, and Corollary 5.126), the diagnostics $(\Lambda_{\mathrm{law}}, \Lambda_{\mathrm{state}})$ from eq. (6.302) are well-defined on $\mathcal{H}_{\mathrm{phys}}$ and are gauge-parameter independent (by Lemma 6.243); moreover, the lift does not change the truth values of the predicates and hence does not change quadrant membership.*

Proof. **Step 0 (Setup and BRST gate).** Let the holosymmetric device act on control C and system S via the controlled unitary

$$U_{\mathrm{ctrl}}(\ell) = |0\rangle\langle 0|_C \otimes U_A(\ell) + |1\rangle\langle 1|_C \otimes U_B(\ell),$$

where U_A, U_B are admissible, BRST-compatible law-level evolutions at IR window size ℓ (via $K_{\mathrm{IR}}^{(\ell)}$), optionally composed with a UV bandlimit $P_x(\lambda_{\mathrm{UV}})$. Project to the physical sector with $\mathsf{P}_{\mathrm{phys}}$ and write $U_k^{\mathrm{phys}} := \mathsf{P}_{\mathrm{phys}} U_k \mathsf{P}_{\mathrm{phys}}$, $U_{\mathrm{ctrl}}^{\mathrm{phys}} := |0\rangle\langle 0| \otimes U_A^{\mathrm{phys}} + |1\rangle\langle 1| \otimes U_B^{\mathrm{phys}}$. Anomalous candidates are excluded at Stage I (see Principle 6.2 and eq. (12.84)) and are not considered.

Step 1 (Diagnostics are well-defined and gauge-fermion independent). For any initial system state ρ_0, prepare $|+\rangle\langle+|_C \otimes \rho_0^{\text{phys}}$ with $\rho_0^{\text{phys}} := \mathsf{P}_{\text{phys}}\rho_0\mathsf{P}_{\text{phys}}$. Applying $U_{\text{ctrl}}^{\text{phys}}(\ell)$ and tracing out S yields

$$\rho'_C(\ell) = \tfrac{1}{2}(\mathbb{I} + \operatorname{Re}\Lambda_{\text{law}}(\ell)\,\sigma_x + \operatorname{Im}\Lambda_{\text{law}}(\ell)\,\sigma_y)\,, \quad \Lambda_{\text{law}}(\ell) = \operatorname{Tr}\Big[(U_B^{\text{phys}})^\dagger(\ell)U_A^{\text{phys}}(\ell)\rho_0^{\text{phys}}\Big],$$

as in Lemma 6.245. Likewise for the *state* protocol (with control/evolution parameter τ), $\Lambda_{\text{state}}(\tau)$ is given by Lemma 6.246. By Lemma 6.243, $\partial_\xi\Lambda_{\text{law/state}} = \langle\zeta\,\eta_\xi^{(\cdot)}\rangle = 0$, so the readouts are gauge-parameter (gauge-fermion) independent in $\mathcal{H}_{\text{phys}}$.

Step 2 (Functoriality over the triadic split). Let $\{\Pi_\alpha\}_{\alpha\in\{\text{corr,vac,spec}\}}$ be the BRST-closed channel projectors (Definition 5.54). Because the controlled map respects the split,

$$\Pi_\alpha U_{\text{ctrl}}^{\text{phys}}(\ell) = U_{\text{ctrl}}^{\text{phys}}(\ell)\Pi_\alpha, \qquad \Lambda_\bullet(\ell) = \sum_\alpha \Lambda_\bullet^{(\alpha)}(\ell),$$

with $\Lambda_{\text{law}}^{(\alpha)}(\ell) = \operatorname{Tr}\Big[(U_{B,\alpha}^{\text{phys}})^\dagger(\ell)\,U_{A,\alpha}^{\text{phys}}(\ell)\,\rho_0^{\text{phys}}\Big]$ and $\Lambda_{\text{state}}^{(\alpha)}(\tau) = \langle\Omega_2|\Pi_\alpha|\Omega_1\rangle$ (Lemma 6.247). Thus no cross-channel contamination is introduced.

Step 3 (Cone preservation and admissibility). By Definition 5.123, Proposition 5.124, and Corollary 5.126, the lift preserves branchwise effective cones and alters only intra-cone phases; this holds a fortiori on $\mathcal{H}_{\text{phys}}$ since cone diagnostics depend on BRST-invariant expectation values. Quadrant-defining kinematics (e.g. vacuum structure, NG content for global symmetries) are therefore unaffected.

Step 4 (Law predicate stability). Let $P_{\text{Law}}(G)$ assert $[U_G, U(\ell)] = 0$ (equivalently $\delta_G S = 0$). If $[U_G^{\text{phys}}, U_A^{\text{phys}}(\ell)] = [U_G^{\text{phys}}, U_B^{\text{phys}}(\ell)] = 0$, then

$$[U_G^{\text{phys}}\otimes\mathbb{I}_C,\ U_{\text{ctrl}}^{\text{phys}}(\ell)] = |0\rangle\langle0|\otimes[U_G^{\text{phys}}, U_A^{\text{phys}}(\ell)] + |1\rangle\langle1|\otimes[U_G^{\text{phys}}, U_B^{\text{phys}}(\ell)] = 0,$$

so the controlled device is G-symmetric on $C\otimes S$ (Lemma 6.249). If $P_{\text{Law}}(G)$ fails on any branch, the commutator is nonzero; a controlled lift cannot restore a symmetry absent on a branch. Thus the truth value of $P_{\text{Law}}(G)$ is unchanged.

Step 5 (State predicate stability). Let $P_{\text{State}}(G)$ assert "Ward identities exact and all G-charged order parameters vanish in the thermodynamic limit." Embed ρ as $|+\rangle\langle+|_C\otimes\rho^{\text{phys}}$ with $\rho^{\text{phys}} := \mathsf{P}_{\text{phys}}\rho\mathsf{P}_{\text{phys}}$ and apply either protocol. Tracing out C gives

$$\rho'_S(\ell) = \tfrac{1}{2}\Big(U_A^{\text{phys}}(\ell)\rho^{\text{phys}}(U_A^{\text{phys}})^\dagger(\ell) \;+\; U_B^{\text{phys}}(\ell)\rho^{\text{phys}}(U_B^{\text{phys}})^\dagger(\ell)\Big)\,.$$

(i) If ρ is G-invariant and $P_{\text{Law}}(G)$ holds on both branches, then $\rho'_S(\ell)$ is G-invariant by covariance and convexity; $P_{\text{State}}(G)$ remains true. (ii) If ρ lies in an SSB phase, coherent superpositions of distinct vacua become block-diagonal between superselection sectors as $V\to\infty$ (vacuum overlaps $\to 0$); the resulting convex mixture retains nonvanishing order parameters and cluster properties (Lemma 6.250). Hence $P_{\text{State}}(G)$ remains false in the thermodynamic sense.

Step 6 (Quadrant membership is preserved). Quadrant membership depends only on the ordered pair $\Big(P_{\text{Law}}(G), P_{\text{State}}(G)\Big)$ (Section 6.11 and Proposition 6.225). Steps 4–5 show these truth values are invariant under the holosymmetric lift; Steps 2–3 ensure internal decomposition and causal structure are respected; Step 1 guarantees $(\Lambda_{\text{law}}, \Lambda_{\text{state}})$ are well-defined and gauge-parameter independent in $\mathcal{H}_{\text{phys}}$. Therefore the overlay refines diagnosis via eq. (6.302) without inducing transitions between cells of table 6.1. □

Remark 6.252 (Weak-history expansion and channel resolution)**.** Fix an RG scale ℓ and write $U_k(\ell) = \mathcal{T}_\tau \exp\Big(- i\int d\tau\, \mathcal{H}_k^{(\ell)}(\tau)\Big)$. Let $\delta\mathcal{H}^{(\ell)} := \mathcal{H}_A^{(\ell)} - \mathcal{H}_B^{(\ell)}$ and $K^{(\ell)} := \int d\tau\, \delta\mathcal{H}^{(\ell)}(\tau)$. For weak, smeared generators, a Magnus (operator) plus cumulant (state) expansion gives, to leading orders,

$$U_B^\dagger(\ell)U_A(\ell) \;\approx\; \exp\Big\{ i\,K^{(\ell)} \;+\; \tfrac{i}{2}\iint d\tau\, d\tau'\, [\mathcal{H}_A^{(\ell)}(\tau), \mathcal{H}_B^{(\ell)}(\tau')] \;+\; \cdots \Big\},$$

and hence for any physical probe ρ^{phys},

$$\Lambda_{\mathrm{law}}(\ell) = \mathrm{Tr}\Big(U_B^\dagger(\ell)U_A(\ell)\rho^{\mathrm{phys}}\Big) \;\approx\; \exp\Big\{ i\,\langle K^{(\ell)}\rangle_\rho \;-\; \tfrac{1}{2}\,\mathrm{Var}_\rho\Big(K^{(\ell)}\Big) \;+\; \cdots \Big\}.$$

Thus Re Λ_{law} is controlled by anticommutators/variances (T-even; correlational/vacuum channels), while Im Λ_{law} is controlled by commutators (T-odd; spectral channel).

BRST overlay: evaluate overlaps on $\mathcal{H}_{\mathrm{phys}}$; ζ-exact differences of generators between branches shift the exponent by ζ-exact terms and therefore drop from the visibilities (gauge-fermion independence by Lemma 6.243). Together with Lemma 6.247, this underpins phase- and channel-resolved diagnostics without changing quadrant membership.

Theorem 6.253 (Expected holosymmetric signatures by quadrant (IR/UV-aware, BRST-cohomological))**.** *Assume the scale-tagged infrastructure with IR windowing $K_{\mathrm{IR}}^{(\ell)}$ and, where needed, UV bandlimiting $P_x(\lambda_{\mathrm{UV}})$ (cf. Remarks 6.240 and 6.252), functoriality and cone preservation (Lemmas 6.247 and 6.248), and the Relational Ward identity. Work in an anomaly-free, BRST-consistent setting and evaluate all visibilities in the physical sector $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$, where gauge-parameter changes act by ζ-exact deformations and hence leave visibilities invariant (Lemma 6.243). Then, uniformly on plateau-stable bands in $(\lambda_{\mathrm{UV}}, \ell)$, the asymptotics are:*

(i) Unbroken (Law Sym., State Sym.)
$|\Lambda_{\mathrm{law}}(\ell)| \to 1$ *as* $\ell \downarrow 0$ *(only IR-irrelevant corrections);* $|\Lambda_{\mathrm{state}}(V)| \to 1$ *as* $V \uparrow \infty$*; Ward/Slavnov–Taylor identities hold on each branch [82, 91].*

(ii) SSB (Law Sym., State Asym.)
$|\Lambda_{\mathrm{law}}(\ell)| \to 1$ *as* $\ell \downarrow 0$*;* $|\Lambda_{\mathrm{state}}(V)| \to 0$ *as* $V \uparrow \infty$ *(thermodynamic orthogonality of distinct vacua). Phase-twist interferometry along the vacuum manifold reads Goldstone kinematics for broken* global *generators; in gauged sectors, would-be Goldstones are ζ-exact (no literal gauge-SSB) [92, 94, 91, 82].*

(iii) Emergent/Accidental (Law Asym., State Sym.)
In this regime the law-coherence deficit decays with the IR length:

$$1 - |\Lambda_{\mathrm{law}}(\ell)| = \mathcal{O}\Big(\ell^{-(\Delta-(d+z))}\Big) \quad (\ell \uparrow \infty),$$

with $\Delta := \min_k \Delta_k$ over RG-irrelevant breaking operators; and $|\Lambda_{\mathrm{state}}(V)| \to 1$. Equivalently, writing $\mu \sim 1/\ell$,

$$1 - |\Lambda_{\mathrm{law}}| = \mathcal{O}\Big(\mu^{\,\Delta-(d+z)}\Big) \quad (\mu \downarrow 0).$$

This is the "emergence law'' expected under RG flow to a symmetric IR fixed point [102, 82].

(iv) Explicit breaking (Law Asym., State Asym.)
$|\Lambda_{\mathrm{law}}(\ell)|$ *remains bounded away from* 1 uniformly into the IR*; more precisely, there exist* $c > 0$ *and* ℓ_0 *such that*

$$1 - |\Lambda_{\mathrm{law}}(\ell)| \geq c \quad \textit{for all} \quad \ell \geq \ell_0,$$

with the deficit set by the explicit couplings (no improvement toward the IR). $|\Lambda_{\mathrm{state}}|$ *is trivial when the ground state is unique (no true Goldstones; at most pseudo-NG with* $m^2 \propto \epsilon$*) [91].*

These signatures are conservative overlays: the holosymmetric lift provides diagnostics without altering quadrant membership, which is fixed by the dyadic product from the Axiom of Duality (Section 6.11) and preserved by the overlay (Proposition 6.251).

Remark 6.254 (BRST-closed visibilities, scaling conventions, and plateau uniformity)**.** Throughout Theorem 6.253, the visibilities are taken to be BRST-closed, dimensionless functionals $\Lambda_{\mathrm{law}}(\ell)$, $\Lambda_{\mathrm{state}}(V) \in [0,1]$ of the IR window scale ℓ and spatial/thermodynamic extent V, evaluated in the physical sector $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$.

(a) Nielsen independence. Let $\Psi(\zeta)$ denote the gauge-fixing fermion and $S_{\mathrm{gf+gh}} = S_0 + \{\hat{Q}_{\mathrm{BRST}}, \Psi(\zeta)\}$. For any BRST-closed observable $\mathcal{V}$ (including $\Lambda_{\mathrm{law}}, \Lambda_{\mathrm{state}}$),

$$\partial_\zeta \langle \mathcal{V} \rangle = \Big\langle \{\hat{Q}_{\mathrm{BRST}},\, \partial_\zeta \Psi\}\, \mathcal{V} \Big\rangle = 0, \qquad \hat{Q}_{\mathrm{BRST}}\, \mathcal{V} = 0,$$

so the visibilities are gauge-parameter independent (cf. Lemma 6.243).

(b) IR/UV scaling conventions. We use ℓ as the IR coarse-graining length and $\mu \sim \ell^{-1}$ as the corresponding momentum scale. With dynamical exponent z, the engineering dimension of spacetime is $d + z$. If $\mathcal{O}_k$ is a symmetry-breaking operator of scaling dimension Δ_k, then for *irrelevant* breakings ($\Delta_k > d + z$) one has the IR fall-off

$$1 - |\Lambda_{\mathrm{law}}(\ell)| = \mathcal{O}\Big(\ell^{-(\Delta - (d+z))}\Big) \quad (\ell \uparrow \infty),$$
$$\text{equivalently} \quad 1 - |\Lambda_{\mathrm{law}}| = \mathcal{O}\Big(\mu^{\,\Delta - (d+z)}\Big) \quad (\mu \downarrow 0),$$

with $\Delta := \min_k \Delta_k$. For *relevant/marginal* explicit breakings ($\Delta \leq d + z$), the deficit does not decay into the IR.

(c) Plateau-stable bands and uniformity. There exists a nonempty closed band $B \subset (\lambda_{\mathrm{UV}}, \ell)$ ("plateau-stable band") such that for all admissible choices of window kernels $K_{\mathrm{IR}}^{(\ell)}$, UV projectors $P_x(\lambda_{\mathrm{UV}})$, and gauges ζ (within a fixed BRST class), the induced variation obeys

$$\sup_{(\lambda_{\mathrm{UV}}, \ell) \in B} \Big| \delta \Lambda_{\mathrm{law/state}} \Big| \leq \varepsilon_{\mathrm{tol}}.$$

All big-$\mathcal{O}$ constants in Theorem 6.253 are understood to be uniform on B.

(d) Normalization robustness. If $\Lambda'_{\mathrm{law/state}}$ is obtained from $\Lambda_{\mathrm{law/state}}$ by any admissible (respectively BRST-closed, bounded, and monotone) renormalization of the underlying Ward deficits or cluster ratios, then $\Lambda'_{\mathrm{law/state}} = (1 + o_B(1))\, \Lambda_{\mathrm{law/state}}$ on B, so the stated asymptotics and quadrant classification are unchanged.

Proof of Theorem 6.253. Fix an anomaly-free, BRST-consistent gauge theory with gauge-fixed action $S_{\mathrm{gf+gh}} = S_0 + \{\hat{Q}_{\mathrm{BRST}}, \Psi(\zeta)\}$ and evaluate all expectation values in the physical sector $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$. Let $\Lambda_{\mathrm{law}}(\ell)$ and $\Lambda_{\mathrm{state}}(V)$ be the BRST-closed visibilities from Remark 6.254.

Setup and Nielsen independence. Differentiating the generating functional with respect to the gauge parameter ζ gives $\partial_\zeta S_{\mathrm{gf+gh}} = \{\hat{Q}_{\mathrm{BRST}}, \partial_\zeta \Psi\}$. For any BRST-closed observable $\mathcal{V}$ (in particular, the visibilities),

$$\partial_\zeta \langle \mathcal{V} \rangle = \Big\langle \{\hat{Q}_{\mathrm{BRST}}, \partial_\zeta \Psi\}\, \mathcal{V} \Big\rangle = 0,$$

where we used BRST invariance of the (anomaly-free) path-integral measure and $\hat{Q}_{\mathrm{BRST}}\mathcal{V} = 0$. Hence all statements below are gauge-parameter independent (cf. Lemma 6.243).

(i) Unbroken (Law Sym., State Sym.). If the microscopic law is exactly symmetric, symmetry-breaking operators $\{\mathcal{O}_k\}$ are absent. Allowing for UV counterterms, any symmetry-violating insertion is *irrelevant* with scaling dimension $\Delta_k > d + z$. Writing $\mu \sim \ell^{-1}$, Wilsonian power counting yields

$$\delta\Lambda_{\mathrm{law}}(\mu) \sim \sum_k c_k\, \mu^{\,\Delta_k - (d+z)} \xrightarrow[\mu\to\infty]{} 0 \quad \Longleftrightarrow \quad \delta\Lambda_{\mathrm{law}}(\ell) \xrightarrow[\ell\downarrow 0]{} 0,$$

so $|\Lambda_{\mathrm{law}}(\ell)| \to 1$ as stated. In a unique symmetric phase, cluster decomposition implies that long-distance correlators of BRST-closed, gauge-invariant operators factorize, whence $|\Lambda_{\mathrm{state}}(V)| \to 1$ as $V \uparrow \infty$. Finally, BRST invariance implies the Slavnov–Taylor identities on each branch.

(ii) SSB (Law Sym., State Asym.). Law symmetry gives the same UV behavior for Λ_{law} as in (i). Assume a global continuous symmetry is spontaneously broken with vacuum manifold $\mathcal{M}$. Take orthonormal vacua $\{|\Omega_\alpha\rangle\}_{\alpha\in\mathcal{M}}$ selected by boundary conditions; for a local order parameter ϕ, $\langle\Omega_\alpha|\phi|\Omega_\beta\rangle = v\,\delta_{\alpha\beta}$ and $\langle\Omega_\alpha|\Omega_\beta\rangle \to 0$ as $V \uparrow \infty$ for $\alpha \neq \beta$, so any BRST-closed state visibility that compares distinct phases vanishes in the thermodynamic limit: $|\Lambda_{\mathrm{state}}(V)| \to 0$. If the broken generators are *gauged*, Elitzur's theorem together with the BRST quartet mechanism (see the Chapter 8 projector $\mathsf{P}_{\mathrm{phys}}$ and quartet decoupling) removes would-be Goldstones from $\mathcal{H}_{\mathrm{phys}}$; there is no literal gauge-SSB, and the conclusions above for Λ_{law} remain intact.

(iii) Emergent/Accidental (Law Asym., State Sym.). Suppose the UV action contains symmetry-breaking *irrelevant* operators $\mathcal{O}_k$ with $\Delta_k > d + z$. Along the RG flow toward the IR, their couplings obey $g_k(\mu) \sim \mu^{\,\Delta_k - (d+z)}$ as $\mu \downarrow 0$; therefore

$$1 - |\Lambda_{\mathrm{law}}| \;\sim\; \sum_k c_k\, g_k(\mu) \;=\; \mathcal{O}\Big(\mu^{\,\Delta - (d+z)}\Big) \xrightarrow[\mu\downarrow 0]{} 0,$$

with $\Delta := \min_k \Delta_k$. Equivalently, $1 - |\Lambda_{\mathrm{law}}(\ell)| = \mathcal{O}\Big(\ell^{-(\Delta-(d+z))}\Big)$ as $\ell \uparrow \infty$. If the IR fixed point is symmetric and the phase is unique, cluster decomposition gives $|\Lambda_{\mathrm{state}}(V)| \to 1$. By Remark 6.254, these estimates hold *uniformly* on any plateau-stable band B.

(iv) Explicit breaking (Law Asym., State Asym.). Add a relevant or marginal explicit breaking $\epsilon\,\mathcal{O}_{\mathrm{break}}$ with $\Delta \le d + z$. Then $g(\mu) \sim \epsilon\,\mu^{\,\Delta-(d+z)}$ does not vanish as $\mu \downarrow 0$ (stays

constant for $\Delta = d + z$, grows for $\Delta < d + z$). Consequently, the law-coherence deficit is bounded away from zero into the IR. By continuity in $(\lambda_{\mathrm{UV}}, \ell)$ and plateau stability (Remark 6.254), there exist $c > 0$ and ℓ_0 such that

$$1 - |\Lambda_{\mathrm{law}}(\ell)| \;\geq\; c \qquad \forall\, \ell \geq \ell_0,$$

establishing the claimed uniform IR bound. If the ground state is unique, Λ_{state} is trivial. With near-degeneracy from small explicit breaking, pseudo–Nambu–Goldstone excitations obey the standard scaling $m^2_{\mathrm{pNG}} \propto \epsilon / f^2$ (with f the appropriate decay constant), consistent with the Chapter 8 physical-spectrum extraction via the projected Hessian $\mathbb{H}_{\mathrm{phys}}$ and positivity of the Källén–Lehmann density on $\mathcal{H}_{\mathrm{phys}}$.

Uniformity, functoriality, and cone preservation. All big-$\mathcal{O}$ constants above are uniform on any plateau-stable band B, as stated in Remark 6.254. Functoriality of the scale tags and cone preservation (Lemmas 6.247 and 6.248) ensure that IR windowing $K^{(\ell)}_{\mathrm{IR}}$ and UV bandlimiting $P_x(\lambda_{\mathrm{UV}})$ do not spoil the causal principal symbol nor the BRST projection to $\mathcal{H}_{\mathrm{phys}}$, so the asymptotics are insensitive to admissible choices within the class.

Conclusion. Items (i)–(iv) follow, and the "holosymmetric lift'' acts only as a diagnostic overlay: by construction (Axiom of Duality and Proposition 6.251) it does not alter quadrant membership, which is determined by the dyadic law/state product. □

Remark 6.255 (Edge cases and variants relevant to Theorem 6.253)**.** The asymptotic signatures in Theorem 6.253 persist under the following variants, provided all visibilities are BRST-closed and evaluated in $\mathcal{H}_{\mathrm{phys}}$ (cf. Remark 6.254). Cross-references point to the downstream formalism developed in *Chapter 8: Field Laws and Physical States*.

(a) Discrete-symmetry SSB (no Goldstone). If a *discrete* global symmetry is spontaneously broken (e.g. $G = \mathbb{Z}_n$), there are no Goldstone modes; instead the characteristic IR object is the *domain-wall tension* $T_{\mathrm{dw}} > 0$. Orthogonality of distinct infinite-volume vacua still holds, hence any BRST-closed state visibility that compares distinct broken phases obeys

$$|\Lambda_{\mathrm{state}}(V)| \;\xrightarrow[V\uparrow\infty]{}\; 0,$$

and the law-side statement is as in item (ii) of Theorem 6.253. When quoting spectra from the projected Hessian $\mathbb{H}_{\mathrm{phys}}$, the low-lying excitations are kink/domain-wall states rather than Goldstones (see Chapter 8, § *Physical projector and projected Hessian* and § *Quadratic theory, propagators, and positivity*).

(b) Topological order and choice of probes. In phases with topological order, local order parameters are inadequate. To prevent contamination of item (ii) by nonlocal probes, define visibilities using *BRST-closed, gauge-invariant* operators adapted to the phase (e.g. Wilson loops $\mathcal{W}_C$, 't Hooft surfaces $\mathcal{T}_\Sigma$, or their coarse-grained versions under $K^{(\ell)}_{\mathrm{IR}}$). All statements of Theorem 6.253 then apply verbatim to these probes on $\mathcal{H}_{\mathrm{phys}}$, with the understanding that "state symmetry" is diagnosed by topological degeneracy patterns rather than local order parameters (see Chapter 8, § *Transverse/longitudinal split & quartet decoupling* and § *Scale-tagged dynamics*).

(c) Anisotropic scaling (explicit z). With dynamical exponent z (time scaling as $t \mapsto b^z t$), engineering dimensions are measured against $d+z$. The scaling claims in Theorem 6.253 and Remark 6.254 already reflect this via the replacement $d \mapsto d+z$. Equivalently, writing $\mu_t \sim \ell_t^{-1}$ and $\mu_x \sim \ell_x^{-1}$ with $\mu_t \sim \mu_x^z$, all power-counting estimates can be expressed using a single IR scale $\mu \sim \ell^{-1}$ once z is fixed, without altering the big-$\mathcal{O}$ exponents (see Chapter 8, § *Scale-tagged dynamics (UV/IR knobs)*).

(d) Crossover windows and competing irrelevancies. If several symmetry-breaking *irrelevant* operators $\{\mathcal{O}_k\}$ compete, with dimensions $\Delta_1 < \Delta_2 < \cdots$ (all $> d+z$), then

$$1 - |\Lambda_{\text{law}}| \;=\; \sum_k c_k\, \mu^{\,\Delta_k-(d+z)} \quad (\mu \downarrow 0),$$

so the leading fall-off is set by $\Delta := \Delta_1$, while subleading corrections are visible away from a plateau band B. Crossover between two contributions $k = 1, 2$ occurs near the IR length

$$\ell_\times \;\sim\; \left|\tfrac{c_1}{c_2}\right|^{\frac{1}{\Delta_2-\Delta_1}},$$

beyond which the Δ_1 term dominates. On any fixed plateau B, the hierarchy is stable and the $\mathcal{O}\big(\ell^{-(\Delta-(d+z))}\big)$ law is uniform (see Chapter 8, § *Plateau stability and error control* and § *BRST-consistent reading and gauge-parameter independence*).

6.11.6 Operational Holosymmetric Cartography of the Quadrants

Conventions for this section. The UV knob is the resolution length λ_{UV} (entering via the spectral projector $P_x(\lambda_{\text{UV}})$), the IR coarse graining uses the kernel family $K_{\text{IR}}^{(\ell)}$ with window size ℓ, and the operational coherences $\Lambda_{\text{law}}, \Lambda_{\text{state}}$ remain as defined. *BRST reading:* we assume an anomaly-free, BRST-consistent gauge fixing $\mathcal{L} = \mathcal{L}_0 + \sum_k \epsilon_k \mathcal{O}_k + \zeta\Psi$; all interferometric quantities are evaluated in the physical sector $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$, with gauge-parameter changes acting by ζ-exact deformations (cf. Lemma 6.243).
Synthesis. The Axiom of Duality (see Section 6.11) yields the dyadic, $\mathbb{Z}_2 \times \mathbb{Z}_2$ quadrant structure of law vs. state; the *holosymmetric lift* adds a coherent, two-path instrumentation that cleanly separates *law-level coherence* (emergent vs. explicit breaking) from *state-level coherence* (unbroken vs. SSB), without changing quadrant membership. Geometry anchors the construction in Chapter 6: causal cones are fixed by the expectation tensor $\bar{\Sigma}_{\mu\nu}$ (Definition 5.26); the lift acts functorially on the triadic channel split (Definition 5.54), modifying only intra-cone phase transport. The IR averages are taken with $K_{\text{IR}}^{(\ell)}$, and (where needed elsewhere) UV bandlimiting is controlled by $P_x(\lambda_{\text{UV}})$.

Definition 6.256 (Holosymmetric operational order parameters)**.** Let $U_A(\ell)$ and $U_B(\ell)$ be two alternative evolutions at the *law* layer (e.g., microscopic vs. coarse-grained IR descriptions) evaluated at IR window size ℓ (with kernel $K_{\text{IR}}^{(\ell)}$), and let $|\Omega_1\rangle, |\Omega_2\rangle$ be two *state* choices (e.g., vacua) for system size V. The holosymmetric lift prepares a control qubit and runs the two paths coherently. The control readouts give the real/imaginary parts of

$$\Lambda_{\text{law}}(\ell) \;:=\; \mathrm{Tr}\Big[U_B^\dagger(\ell)\, U_A(\ell)\, \rho_0\Big], \qquad \Lambda_{\text{state}}(V) \;:=\; \langle\Omega_2|\Omega_1\rangle, \tag{6.302}$$

Table 6.2: Holosymmetric meta-layer (BRST-cohomological readout): what to probe in each quadrant (overlay to table 6.1).

Law \ State	**Symmetric state**	**Asymmetric state (SSB)**
Symmetric law	*Probe (in $\mathcal{H}_{\text{phys}}$):* plateau-stable $\lvert\Lambda_{\text{law}}\rvert\approx 1$ and $\lvert\Lambda_{\text{state}}\rvert\approx 1$ under $(\lambda_{\text{UV}}, \ell)$ scans; confirm Ward/Slavnov–Taylor exactness (BRST identities). Gauge-parameter changes are ζ-exact and leave visibilities invariant (Lemma 6.243).	*Probe (in $\mathcal{H}_{\text{phys}}$):* $\lvert\Lambda_{\text{law}}\rvert\approx 1$ (plateau-stable); $\Lambda_{\text{state}}(V)\to 0$ as $V\uparrow\infty$ (thermodynamic orthogonality). Use phase-twist interferometry along the vacuum manifold to read Goldstone kinematics for broken *global* generators; no literal gauge-SSB (would-be modes are ζ-exact).
Asymmetric law	*Probe (in $\mathcal{H}_{\text{phys}}$):* emergence scaling $1-\lvert\Lambda_{\text{law}}(\ell)\rvert = \mathcal{O}\big(\ell^{\Delta-(d+z)}\big) \quad (\ell\downarrow 0)$, with $\lvert\Lambda_{\text{state}}\rvert\approx 1$; IR Ward/Slavnov–Taylor become approximately exact. Explicit micro-breakings must preserve BRST nilpotency (spurionic if needed); otherwise excluded at Stage I.	*Probe (in $\mathcal{H}_{\text{phys}}$):* $\lvert\Lambda_{\text{law}}\rvert < 1$ bounded away from 1 at small ℓ (deficit tracks explicit couplings; no IR improvement). Unique vacuum $\Rightarrow$ trivial Λ_{state} (unity up to phases); no exact Goldstones (pseudo-NG only, with $m^2\propto\epsilon$). BRST must remain exact; anomaly $\Rightarrow$ Stage I exclusion.

Evaluate all visibilities in the BRST physical sector $\mathcal{H}_{\text{phys}}$; insertions are BRST-closed (ghost number 0) and gauge-parameter dependence is ζ-exact (cf. Section 6.11.3 and Lemma 6.243). Use the IR window $K_{\text{IR}}^{(\ell)}$ and, where needed, the UV bandlimit $P_x(\lambda_{\text{UV}})$; require plateau stability (Section 6.12.4). The lift is functorial over channel decompositions and preserves effective cones (Definitions 5.54 and 5.123, Proposition 5.124, and Corollary 5.126); anomalous candidates are excluded at Stage I (Proposition 6.227). *Legend:* All equalities/identities are understood modulo ζ-exact terms (cohomology classes). That is, insertions related by $\mathcal{O}\sim\mathcal{O}+\zeta G$ yield identical visibilities in $\mathcal{H}_{\text{phys}}$.

with ρ_0 the initial system state. *BRST overlay:* interpret (6.302) on $\mathcal{H}_{\text{phys}}$ by inserting the projector P_{phys} on both sides of $U_{A/B}$ and ρ_0, taking $|\Omega_i\rangle$ to be ζ-closed, ghost-number-zero representatives. Gauge-parameter dependence is ζ-exact and drops in $\mathcal{H}_{\text{phys}}$ (cf. Lemma 6.243); anomalous candidates are excluded at Stage I. These functionals serve as *operational order parameters* for the 2×2 in table 6.1.

Proposition 6.257 (Functoriality and causal guardrails)**.** *Let the* holosymmetric lift *acting on an observable/kernel A be the blockwise, scale–tagged transform*

$$\mathfrak{H}_{\lambda_{\text{UV}},\ell}[A] := \sum_{\alpha\in\{\text{corr,vac,spec}\}} \Pi_\alpha\, K_{\text{IR}}^{(\ell)}\, P_x(\lambda_{\text{UV}})\, A\, P_x(\lambda_{\text{UV}})\, K_{\text{IR}}^{(\ell)}\, \Pi_\alpha,$$

with $\{\Pi_\alpha\}$ the triadic projectors of Definition 5.54 and $\bar{\Sigma}_{\mu\nu}$ as in Definition 5.26. The holosymmetric lift respects direct sums/decompositions of observables and channels (functorial over the triadic split in Definition 5.54) and preserves the cone structure induced by $\bar{\Sigma}_{\mu\nu}$ (Definition 5.26). Hence it alters only intra-cone phase transport; in particular, it cannot change which cell of table 6.1 a system occupies. BRST overlay: *choose channel projectors BRST-closed (ghost number 0); ζ-exact improvements do not mix blocks and do not affect causal diagnostics in $\mathcal{H}_{\text{phys}}$.*

Proof of Proposition 6.257. **Set-up.** Let $\{\Pi_\alpha\}_{\alpha\in\{\text{corr,vac,spec}\}}$ be the triadic channel projectors from Definition 5.54, and let the scale tags be the IR window $K_{\text{IR}}^{(\ell)}$ and UV bandlimiter $P_x(\lambda_{\text{UV}})$. The *holosymmetric lift* of an observable (or kernel) A is the scale-tagged, channel-resolved transform

$$\mathfrak{H}_{\lambda_{\text{UV}},\ell}[A] := \sum_{\alpha} \Pi_\alpha\, K_{\text{IR}}^{(\ell)}\, P_x(\lambda_{\text{UV}})\, A\, P_x(\lambda_{\text{UV}})\, K_{\text{IR}}^{(\ell)}\, \Pi_\alpha,$$

possibly followed by physically harmless unitary intertwiners that are blockwise (channelwise). All objects are evaluated against the expectation-level chronogeometric tensor $\bar{\Sigma}_{\mu\nu}$ from Definition 5.26.

(1) Functoriality over direct sums / decompositions. By linearity and the idempotence $\Pi_\alpha^2 = \Pi_\alpha$, $\Pi_\alpha\Pi_\beta = 0$ $(\alpha \neq \beta)$,

$$\mathfrak{H}_{\lambda_{\text{UV}},\ell}[A \oplus B] = \mathfrak{H}_{\lambda_{\text{UV}},\ell}[A] \;\oplus\; \mathfrak{H}_{\lambda_{\text{UV}},\ell}[B], \qquad \mathfrak{H}_{\lambda_{\text{UV}},\ell}[\Pi_\alpha A] = \Pi_\alpha\, \mathfrak{H}_{\lambda_{\text{UV}},\ell}[A].$$

Thus $\mathfrak{H}$ commutes with injections and projections in the category whose objects are channel-decomposed operators and whose morphisms are block-preserving maps. In particular, the lift is *block-diagonal* w.r.t. the triadic split and therefore cannot create cross-channel couplings. This is the functorial content of Lemma 6.247, now spelled out at the level of the explicit transform.

(2) Cone preservation. Let $\mathcal{L}$ denote any hyperbolic kinetic operator that governs propagation in a given channel (e.g. the principal part of the quadratic EOM), with principal symbol $\sigma_{\text{pr}}(\mathcal{L})(x,k) = \bar{\Sigma}^{\mu\nu}(x)\, k_\mu k_\nu$ in the sense of Chapter VIII. The window $K_{\text{IR}}^{(\ell)}$ and the projector $P_x(\lambda_{\text{UV}})$ are pseudodifferential operators of order ≤ 0 (smoothing at the IR/UV ends). Hence, for the lifted kinetic operator

$$\mathcal{L}^{\text{lift}} := K_{\text{IR}}^{(\ell)} P_x(\lambda_{\text{UV}})\, \mathcal{L}\, P_x(\lambda_{\text{UV}}) K_{\text{IR}}^{(\ell)},$$

one has $\sigma_{\rm pr}(\mathcal{L}^{\rm lift}) = \sigma_{\rm pr}(\mathcal{L})$, because smoothing/order-0 factors do not alter the principal symbol, and Π_α acts only on internal indices (it is multiplication by a smooth matrix projector in field space). Therefore the characteristic cone $\mathcal{C}_x := \{k \neq 0 : \bar{\Sigma}^{\mu\nu}(x)k_\mu k_\nu = 0\}$ is *unchanged* by the lift. Equivalently, the causal propagator's wavefront set and the microcausal commutator support remain confined to the $\bar{\Sigma}$-cones. This is exactly Lemma 6.248 in principal-symbol form. Consequently, the lift can at most modify *intra-cone phase transport* (subprincipal terms, phases, residues), but it cannot open new propagation channels across the cone.

(3) Quadrant (law/state) predicates are invariant. Quadrant membership (table 6.1) depends on two predicates: (i) *law symmetry*, encoded by exact conservation of the improved currents (Relational Ward identity) in the physical sector; and (ii) *state symmetry*, encoded by uniqueness vs. orthogonality of infinite-volume vacua. For (i), the Relational Ward identity holds in $\mathcal{H}_{\rm phys}$ iff the RHS of $\nabla_\mu \bar{J}^\mu_{a,\rm imp} = \sum_k \epsilon_k \langle \mathcal{S}_{k,a} \rangle$ vanishes (cf. Chapter VII–VIII). Since $\mathfrak{H}_{\lambda_{\rm UV},\ell}$ is blockwise, of order ≤ 0 in spacetime, and does not insert relevant operators, it *does not* change $\{\epsilon_k\}$ nor the cohomology class of the current; thus the truth value of law symmetry is preserved. For (ii), the lift is norm-bounded and block-diagonal, hence it cannot turn a unique GNS representation into a family of disjoint ones, nor vice versa; orthogonality of distinct vacua and cluster properties are unaffected by smoothing/projecting operations of order ≤ 0. Therefore the state predicate is preserved. Together, the system remains in the same cell of table 6.1.

(4) BRST overlay and causal diagnostics. Choose the projectors Π_α to be BRST-closed (ghost number 0), so $[\hat{Q}_{\rm BRST}, \Pi_\alpha] = 0$, and select scale tags within a BRST-closed class. Any gauge-parameter (gauge-fermion) change produces a ζ-exact variation of the lift, which drops out of all expectations in $\mathcal{H}_{\rm phys}$ by Lemma 6.243. Since Π_α commute with the BRST charge, such variations *do not* mix blocks. Chapter VIII's positivity and projected-Hessian analysis then apply unchanged to causal diagnostics (transverse poles, hyperbolicity), which therefore remain the same before and after the lift.

Conclusion. The holosymmetric lift is a blockwise (functorial) order-≤ 0 transformation that preserves the $\bar{\Sigma}$-cone and the BRST predicates in $\mathcal{H}_{\rm phys}$. It can adjust intra-cone phases/subprincipal structure but cannot modify conservation laws, vacuum multiplicity, or causal reach. Hence it does *not* move a system between the quadrants of table 6.1; it only provides a diagnostic overlay. □

Theorem 6.258 (Operational Cartography of the Quadrants (BRST-cohomological))**.** *Assume the Relational Ward identity (6.282), standard cluster/thermodynamic limits, and an anomaly-free BRST setup with visibilities evaluated in $\mathcal{H}_{\rm phys}$ (gauge-parameter independence by Lemma 6.243). Then the quadrants of table 6.1 (identified in Corollary 6.232) are characterized by the asymptotics of the holosymmetric functionals in (6.302):*

(i) Unbroken (Law Sym., State Sym.)
$|\Lambda_{\rm law}(\ell)| \to 1$ *as* $\ell \downarrow 0$ *(only IR-irrelevant corrections);* $|\Lambda_{\rm state}(V)| \to 1$*; order parameter* $= 0$*; Ward/Slavnov–Taylor identities hold [82, 14, 91].*

(ii) SSB (Law Sym., State Asym.)
$|\Lambda_{\rm law}(\ell)| \to 1$ *as* $\ell \downarrow 0$*;* $|\Lambda_{\rm state}(V)| \to 0$ *as* $V \uparrow \infty$ *(vacuum-cat visibility vanishes in the thermodynamic limit). Phase-twist interferometry around the vacuum manifold reads*

off Goldstone kinematics for broken global *generators; in gauged sectors, would-be Goldstones are ζ-exact (no literal gauge-SSB) [92, 94, 91, 82].*

(iii) Emergent/Accidental (Law Asym., State Sym.)
Quantitatively, the law-coherence deficit obeys the IR power law:

$$1 - |\Lambda_{\text{law}}(\ell)| = \mathcal{O}\Big(\ell^{-(\Delta-(d+z))}\Big) \quad (\ell \uparrow \infty),$$

with $\Delta := \min_k \Delta_k$ over RG-irrelevant breaking operators; $|\Lambda_{\text{state}}(V)| \to 1$; no order parameter [102, 82]. Equivalently, with $\mu \sim \ell^{-1}$, the deficit scales as $1 - |\Lambda_{\text{law}}| = \mathcal{O}\Big(\mu^{\,\Delta-(d+z)}\Big)$ as $\mu \downarrow 0$.

(iv) Explicit breaking (Law Asym., State Asym.)
$|\Lambda_{\text{law}}(\ell)|$ *remains bounded away from* 1 uniformly into the IR*; more precisely, there exist $c > 0$ and ℓ_0 such that*

$$1 - |\Lambda_{\text{law}}(\ell)| \geq c \quad \textit{for all} \quad \ell \geq \ell_0,$$

with scaling set by the explicit breaking couplings. $|\Lambda_{\text{state}}(V)|$ is trivial (unique vacuum); no exact Goldstones (at most pseudo-NG with $m^2 \propto \epsilon$) [91].

Operational Cartography of the Quadrants (BRST-cohomological). We use the holosymmetric readout identities

$$\Lambda_{\text{law}}(s) = \text{Tr}\Big[U_B^\dagger(s) U_A(s) \rho_0\Big], \qquad \Lambda_{\text{state}}(s) = \langle \Omega_2 |\, U^\dagger(s) U(s)\, | \Omega_1 \rangle$$

(cf. Definitions 6.237 and 6.238 and Lemmas 6.245 and 6.246), together with functoriality and cone preservation (Lemmas 6.247 and 6.248) and predicate stability (Lemmas 6.249 and 6.250). *BRST reading:* all visibilities are evaluated in the physical sector $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$ via implicit insertions of P_{phys}; gauge-parameter (gauge-fermion) variations are ζ-exact and drop out by Lemma 6.243. We denote the evolution parameter by s; in the RG instantiation we take $s = \ell$ with IR windowing by $K_{\text{IR}}^{(\ell)}$ and, where needed, UV bandlimiting via $P_x(\lambda_{\text{UV}})$. Anomaly freedom is the Chapter 8 condition $\mathcal{S}(\Gamma^{(\lambda_{\text{UV}},\ell)}) = 0$.

Preliminaries (BRST-aware Ward identity). Write the gauge-fixed micro-law as $\mathcal{L} = \mathcal{L}_0 + \sum_k \epsilon_k \mathcal{O}_k + \zeta\Psi$ with $\delta_a \mathcal{L}_0 = \nabla_\mu K_a^\mu$ for global G, and $\delta_a \mathcal{O}_k = \mathcal{S}_{k,a}$. From Lemma 6.233 one has

$$\nabla_\mu \bar{J}^\mu_{a,\text{imp}} = \sum_k \epsilon_k \left\langle \mathcal{S}_{k,a} \right\rangle_{\mathbf{\Psi}} + \left\langle \zeta\, \widehat{\eta}_a \right\rangle_{\mathbf{\Psi}}, \qquad \bar{J}^\mu_{a,\text{imp}} := \langle \hat{J}^\mu_{a,\text{imp}} \rangle_{\mathbf{\Psi}}. \tag{6.303}$$

Projecting to $\mathcal{H}_{\text{phys}}$ kills the ζ-exact term, yielding

$$\nabla_\mu \bar{J}^\mu_{a,\text{imp}} = \sum_k \epsilon_k \left\langle \mathcal{S}_{k,a} \right\rangle_{\mathbf{\Psi}} \quad \text{in } \mathcal{H}_{\text{phys}}. \tag{6.304}$$

Thus exact conservation (RHS = 0) characterizes symmetric laws (Quadrants I–II), while RHS $\neq 0$ encodes explicit breaking (Quadrant IV). In emergent regimes (Quadrant III) the running couplings $\epsilon_k(\ell)$ flow to 0 under *coarse-graining*, i.e. as $\ell \uparrow \infty$, so the RHS becomes asymptotically small.

To relate Λ_{law} to breaking parameters, write $U_A(\ell) = \mathcal{T}\exp\{-i\int_0^\ell d\ell'\, H_A(\ell')\}$ and $U_B(\ell) = \mathcal{T}\exp\{-i\int_0^\ell d\ell'\, H_B(\ell')\}$, and set $\delta H(\ell') := H_A(\ell') - H_B(\ell')$. A Duhamel/BCH expansion gives

$$U_B^\dagger(\ell)U_A(\ell) = \mathbb{I} - i\int_0^\ell d\ell'\, \widetilde{\delta H}(\ell') - \int_{0<\ell''<\ell'<\ell} d\ell' d\ell''\, \widetilde{\delta H}(\ell')\widetilde{\delta H}(\ell'') + \cdots, \tag{6.305}$$

with $\widetilde{\delta H}$ in the interaction picture of H_B. By Lemma 6.243, any ζ-exact improvement leaves Λ_{law} invariant on $\mathcal{H}_{\mathrm{phys}}$. Under the RG instantiation, δH is a linear combination of breaking operators $\mathcal{O}_k$ with running $\epsilon_k(\ell)$ and scaling dimensions Δ_k.

Finally, by Lemma 6.250 and standard cluster/thermodynamic arguments (Chapter 8, § *Quadratic theory, propagators, and positivity*), if the phase is unbroken there is a unique G-invariant vacuum $|\Omega\rangle$ and $\langle\Omega_2|\Omega_1\rangle \to 1$ when preparations coincide in the IR; if the phase exhibits SSB (for *global* G), distinct vacua $\{|\Omega_\alpha\rangle\}$ are mutually orthogonal as $V\to\infty$ (disjoint GNS reps), so $\langle\Omega_\beta|\Omega_\alpha\rangle \to 0$ for $\alpha\neq\beta$.

(i) Unbroken (Law Sym., State Sym.). Law symmetry means $\epsilon_k \equiv 0$ in (6.304); thus $\nabla_\mu \bar{J}^\mu_{a,\mathrm{imp}} = 0$ and Ward/Slavnov–Taylor identities are exact. Any difference between $U_A(\ell), U_B(\ell)$ arises only from IR-irrelevant deformations. If the leading such operator has scaling dimension $\Delta > d+z$, its running coefficient satisfies $\epsilon(\ell) = \mathcal{O}(\ell^{\,\Delta-(d+z)}) \to 0$ as $\ell\downarrow 0$. Using (6.305) with $\delta H(\ell') = \epsilon(\ell')\,\mathcal{O}(\ell')$ and bounded $\|\mathcal{O}(\ell')\|$,

$$\big|1 - \Lambda_{\mathrm{law}}(\ell)\big| \le C_1\int_0^\ell d\ell'\,|\epsilon(\ell')| + C_2\int_{0<\ell''<\ell'<\ell} d\ell' d\ell''\,|\epsilon(\ell')\epsilon(\ell'')| = \mathcal{O}\Big(\ell^{\,\Delta-(d+z)}\Big),$$

so $|\Lambda_{\mathrm{law}}(\ell)| \to 1$ as $\ell\downarrow 0$. For the state, uniqueness of the symmetric vacuum implies $\Lambda_{\mathrm{state}}(V) = \langle\Omega|\Omega\rangle = 1$ up to finite-size corrections that vanish as $V\uparrow\infty$. Vanishing of all G-charged order parameters follows from Ward identities [82, 14].

(ii) SSB (Law Sym., State Asym.). The law is symmetric ($\epsilon_k \equiv 0$), so Ward identities hold on each branch and the same Duhamel estimate yields $|\Lambda_{\mathrm{law}}(\ell)| \to 1$ as $\ell\downarrow 0$. For the state, choose distinct symmetry-related vacua $|\Omega_1\rangle, |\Omega_2\rangle$ (points on G/H); then $\Lambda_{\mathrm{state}}(V) = \langle\Omega_2|\Omega_1\rangle \to 0$ as $V\uparrow\infty$ by orthogonality of sectors. Phase-twist interferometry along the vacuum manifold reads Goldstone kinematics for broken *global* generators (no literal gauge-SSB; would-be gauge modes are ζ-exact and decouple in $\mathcal{H}_{\mathrm{phys}}$ via the Kugo–Ojima quartet; Chapter 8, § *Transverse/longitudinal split & quartet decoupling*).

(iii) Emergent/Accidental (Law Asym., State Sym.). Now the micro-law contains *irrelevant* symmetry breakings with $\Delta_k > d+z$. Under coarse-graining to the IR, the running couplings decay as

$$\epsilon_k(\ell) = \mathcal{O}\Big(\ell^{-(\Delta_k-(d+z))}\Big) \qquad (\ell\uparrow\infty),$$

equivalently $g_k(\mu) = \mathcal{O}\Big(\mu^{\,\Delta_k-(d+z)}\Big)$ for $\mu \sim 1/\ell \downarrow 0$. Insert $\delta H(\ell') = \sum_k \epsilon_k(\ell')\,\mathcal{O}_k(\ell')$ in (6.305). Using boundedness/mixing (uniform on a plateau-stable band B; Remark 6.254) and the change of variables $\mu' = 1/\ell'$ to estimate the leading integral,

$$\int^\ell d\ell'\,\epsilon_k(\ell') \;\sim\; \int^{1/\mu}\frac{d\mu'}{\mu'^2}\,\mu'^{\,\Delta_k-(d+z)} \;=\; \mathcal{O}\Big(\mu^{\,\Delta_k-(d+z)}\Big) \;=\; \mathcal{O}\Big(\ell^{-(\Delta_k-(d+z))}\Big),$$

and similarly for the quadratic term. Hence

$$1 - |\Lambda_{\text{law}}(\ell)| = \mathcal{O}\Big(\ell^{-(\Delta-(d+z))}\Big) \qquad (\ell \uparrow \infty),$$

with $\Delta := \min_k \Delta_k$. State symmetry (unique symmetric vacuum) gives $\Lambda_{\text{state}}(V) \to 1$ and no local order parameter. Consistently, (6.304) has RHS $\sum_k \epsilon_k(\ell)\langle \mathcal{S}_{k,a}\rangle \to 0$ with the same IR power law [102, 82].

(iv) Explicit breaking (Law Asym., State Asym.). Here some breaking coupling is relevant or marginal, staying finite (or growing) in the IR; the RHS of (6.304) remains nonzero at low scales and exact Ward identities fail. In (6.305), the integrals of δH do not wash out; by positivity/variance bounds on BRST-closed probes (Chapter 8, § *Quadratic theory, propagators, and positivity*) there exist $c > 0$ and ℓ_0 (uniform on any plateau band B) such that

$$1 - |\Lambda_{\text{law}}(\ell)| \;\geq\; c \qquad \forall\, \ell \geq \ell_0,$$

establishing the uniform IR deficit. The state lies in a unique explicitly selected vacuum, so $\Lambda_{\text{state}}(V)$ is trivial; there are no exact Goldstones (at most pseudo–Nambu–Goldstones with $m^2 \propto \epsilon$, consistent with the projected Hessian $\mathbb{H}_{\text{phys}}$ evaluation in Chapter 7).

Quadrant stability and causality. Throughout, the holosymmetric lift preserves the triadic split and causal cones (Lemmas 6.247 and 6.248) and does not change the symmetry predicates (Lemmas 6.249 and 6.250). Nielsen independence (Lemma 6.243) guarantees gauge-parameter invariance of visibilities in $\mathcal{H}_{\text{phys}}$. Anomalous candidates fail the Stage I gate (Chapter 7, $\mathcal{S}(\Gamma) = 0$) and are excluded. Hence the measured asymptotics diagnose the pre-existing quadrant rather than inducing transitions.

Combining the four cases yields the characterization claimed in Theorem 6.258. □

Corollary 6.259 (Geometric anchoring (Chapter 6))**.** *For operator-valued tensors $\hat{g}_{\mu\nu}$, $\hat{t}_{\mu\nu}$, $\hat{\Sigma}_{\mu\nu}$ with triadic channels (Definition 5.54), implement the holosymmetric lift channel-wise (corr/vac/spec), using IR windowing by $K_{\text{IR}}^{(\ell)}$ and—where needed—UV bandlimiting via $P_x(\lambda_{\text{UV}})$. Evaluate all quantities in $\mathcal{H}_{\text{phys}}$ with BRST-closed, ghost-number-zero insertions; branchwise Lorentz cones, determined by the expectation tensor $\bar{\Sigma}_{\mu\nu}$ (Definition 5.26), remain fixed, and ζ-exact changes of gauge fixing do not affect cone diagnostics. Thus $(\Lambda_{\text{law}}, \Lambda_{\text{state}})$ diagnose the quadrant without altering the underlying causal structure.*

Proof of Corollary 6.259. **Context and goal.** Let $\hat{g}_{\mu\nu}$, $\hat{t}_{\mu\nu}$, and $\hat{\Sigma}_{\mu\nu} = \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu}$ be the operator-valued tensors endowed with the triadic channel decomposition of Definition 5.54. Write the expectation-level (operational) chronogeometric tensor as $\bar{\Sigma}_{\mu\nu} := \langle \hat{\Sigma}_{\mu\nu}\rangle$, cf. Definition 5.26. Implement the holosymmetric lift channel-wise (corr/vac/spec) by inserting the IR window $K_{\text{IR}}^{(\ell)}$ and, where needed, the UV bandlimiter $P_x(\lambda_{\text{UV}})$; evaluate all expectation values in the physical Hilbert space $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$ with BRST-closed, ghost-number-zero insertions. We must show that (i) the branchwise Lorentz cones determined by $\bar{\Sigma}_{\mu\nu}$ remain fixed and are gauge-parameter independent, and (ii) the visibilities $(\Lambda_{\text{law}}, \Lambda_{\text{state}})$ diagnose the quadrant without altering this causal structure.

Step 1: Cone preservation under the holosymmetric lift. Consider any channel $\alpha \in \{\text{corr}, \text{vac}, \text{spec}\}$ with projector Π_α. Let $\mathcal{L}_\alpha$ denote the hyperbolic kinetic operator governing

small fluctuations in that channel (e.g. the principal part of the quadratic Euler–Lagrange operator), whose principal symbol is

$$\sigma_{\mathrm{pr}}(\mathcal{L}_\alpha)(x,k) \;=\; \bar{\Sigma}^{\mu\nu}(x)\, k_\mu k_\nu \;\otimes\; \mathbf{1}_\alpha.$$

Define the lifted operator

$$\mathcal{L}_\alpha^{\mathrm{lift}} := \Pi_\alpha\, K_{\mathrm{IR}}^{(\ell)}\, P_x(\lambda_{\mathrm{UV}})\, \mathcal{L}_\alpha\, P_x(\lambda_{\mathrm{UV}})\, K_{\mathrm{IR}}^{(\ell)}\, \Pi_\alpha.$$

Since $K_{\mathrm{IR}}^{(\ell)}$ and $P_x(\lambda_{\mathrm{UV}})$ are pseudodifferential operators of order ≤ 0 (smoothing/filters) and Π_α acts on internal indices, the principal symbol is unchanged:

$$\sigma_{\mathrm{pr}}\Big(\mathcal{L}_\alpha^{\mathrm{lift}}\Big) \;=\; \sigma_{\mathrm{pr}}(\mathcal{L}_\alpha) \;=\; \bar{\Sigma}^{\mu\nu}(x)\, k_\mu k_\nu \;\otimes\; \mathbf{1}_\alpha.$$

Therefore the characteristic cone $\mathcal{C}_x := \{k \neq 0 :\ \bar{\Sigma}^{\mu\nu}(x)\, k_\mu k_\nu = 0\}$ is identical before and after the lift (no widening/narrowing, no tilting). Equivalently, the wavefront set of the causal propagator and the support of microcausal commutators are confined to the same $\bar{\Sigma}$-cones both pre- and post-lift. This is precisely the cone-preservation content established abstractly in Proposition 6.257 (via Lemma 6.248), now applied to the geometric tensors.

Step 2: BRST overlay and gauge-parameter independence of cone diagnostics. Choose Π_α, $K_{\mathrm{IR}}^{(\ell)}$, and $P_x(\lambda_{\mathrm{UV}})$ to be BRST-closed (ghost number 0), so that they commute with $\hat{Q}_{\mathrm{BRST}}$ and thus preserve $\mathcal{H}_{\mathrm{phys}}$. Let $\mathcal{O}_{\mathrm{cone}}$ be any BRST-closed diagnostic of causal structure (e.g. a projector onto the commutator's cone support or a symbol-level indicator of hyperbolicity). A gauge-fermion variation changes the gauge-fixed action by a ζ-exact term; by Lemma 6.243 such variations drop out of expectations in $\mathcal{H}_{\mathrm{phys}}$:

$$\partial_\zeta \left\langle \mathcal{O}_{\mathrm{cone}} \right\rangle_{\mathcal{H}_{\mathrm{phys}}} = \left\langle \{\hat{Q}_{\mathrm{BRST}},\, \partial_\zeta \Psi\}\, \mathcal{O}_{\mathrm{cone}} \right\rangle_{\mathcal{H}_{\mathrm{phys}}} = 0.$$

Hence cone diagnostics are gauge-parameter independent in the physical sector. In particular, the branchwise Lorentz cones defined by $\bar{\Sigma}_{\mu\nu}$ are stable under all admissible gauge choices and lifts.

Step 3: Quadrant diagnosis without causal back-reaction. By Theorem 6.258, the asymptotics of the BRST-closed visibilities Λ_{law} and Λ_{state} characterize, respectively, the symmetry of the microscopic law (exact, emergent, or explicitly broken) and of the realized state (symmetric vs. symmetry-broken) on the plateau bands. On the other hand, Proposition 6.257 ensures that the holosymmetric lift is blockwise (functorial over the triadic split), of pseudodifferential order ≤ 0, and therefore *cannot* modify either the principal symbol determined by $\bar{\Sigma}_{\mu\nu}$ or the law/state predicates that define quadrant membership. Combining these two inputs shows that the pair $(\Lambda_{\mathrm{law}}, \Lambda_{\mathrm{state}})$ furnishes a faithful *diagnosis* of the quadrant while leaving the underlying causal structure—encoded by the branchwise cones of $\bar{\Sigma}_{\mu\nu}$—unchanged.

Conclusion. Channel-wise holosymmetric lifting (corr/vac/spec), performed with BRST-closed scale tags and evaluated in $\mathcal{H}_{\mathrm{phys}}$, preserves the $\bar{\Sigma}$-cones and yields gauge-parameter–independent causal diagnostics. Consequently, $(\Lambda_{\mathrm{law}}, \Lambda_{\mathrm{state}})$ identify the quadrant specified in table 6.1 without altering the system's causal geometry. □

6.11.7 Channel-Resolved Holosymmetric Diagnostics

Triadic split. Let the unified chronogeometric operator and its expectation decompose additively and orthogonally into three channels,

$$\hat{\Sigma}_{\mu\nu} = \hat{\Sigma}^{(\mathrm{corr})}_{\mu\nu} + \hat{\Sigma}^{(\mathrm{vac})}_{\mu\nu} + \hat{\Sigma}^{(\mathrm{spec})}_{\mu\nu}, \qquad \bar{\Sigma}_{\mu\nu} = \bar{\Sigma}^{(\mathrm{corr})}_{\mu\nu} + \bar{\Sigma}^{(\mathrm{vac})}_{\mu\nu} + \bar{\Sigma}^{(\mathrm{spec})}_{\mu\nu} \tag{6.306}$$

with channel projectors Π_α ($\alpha \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$) such that $\Pi_\alpha \hat{\Sigma} = \hat{\Sigma}^{(\alpha)}$, $\sum_\alpha \Pi_\alpha = \mathrm{Id}$, and channel orthogonality at expectation level: $\langle \mathcal{O}^{(\alpha)} \mathcal{O}^{(\beta)} \rangle = \delta_{\alpha\beta} \langle \mathcal{O}^{(\alpha)} \mathcal{O}^{(\alpha)} \rangle$ for the diagnostic class of observables considered. *BRST overlay:* choose Π_α BRST-closed of ghost number 0 so that $[\hat{Q}_{\mathrm{BRST}}, \Pi_\alpha] = 0$ and channel expectations are computed on $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$, with equivalences understood modulo ζ-exact terms.

Holosymmetric lift (functoriality and cone invariance). The holosymmetric two-path lift H acts on pairs of law/state branches and a control qubit. It is linear over sums and respects the split:

$$\Pi_\alpha \circ \mathsf{H} = \mathsf{H} \circ \Pi_\alpha \quad (\alpha = \mathrm{corr}, \mathrm{vac}, \mathrm{spec}) \tag{6.307}$$

and it preserves causal cones determined by $\bar{\Sigma}_{\mu\nu}$ (no law-signalling); only *intra-cone phase transport* is affected. IR coarse-graining is performed with the kernel $K^{(\ell)}_{\mathrm{IR}}$ at window size ℓ, and—whenever required—UV bandlimiting is implemented by the spectral projector $P_x(\lambda_{\mathrm{UV}})$. *BRST overlay:* all evolutions/insertions are read in $\mathcal{H}_{\mathrm{phys}}$, are ghost-number 0, and are gauge-parameter independent by Lemma 6.243; anomalous candidates fail Stage I and are excluded.

Definition 6.260 (Channel-resolved holosymmetric order parameters)**.** Let $U_A(\ell), U_B(\ell)$ be alternative *law*-layer evolutions evaluated at IR window size ℓ (with $K^{(\ell)}_{\mathrm{IR}}$) and let $|\Omega_1\rangle, |\Omega_2\rangle$ be alternative *state* choices for volume V. Define the global functionals

$$\Lambda_{\mathrm{law}}(\ell) = \mathrm{Tr}\Big[U_B^\dagger(\ell)\, U_A(\ell)\, \rho_0\Big], \qquad \Lambda_{\mathrm{state}}(V) = \langle \Omega_2 | \Omega_1 \rangle \tag{6.308}$$

and the *channel-resolved* functionals by projection:

$$\Lambda^{(\alpha)}_{\mathrm{law}}(\ell) = \mathrm{Tr}\Big[U^\dagger_{B,\alpha}(\ell)\, U_{A,\alpha}(\ell)\, \rho_0\Big], \qquad \Lambda^{(\alpha)}_{\mathrm{state}}(V) = \langle \Omega_2 |\, \Pi_\alpha\, | \Omega_1 \rangle, \quad \alpha \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\} \tag{6.309}$$

with $U_{A,\alpha} := \Pi_\alpha U_A \Pi_\alpha$ and $U_{B,\alpha} := \Pi_\alpha U_B \Pi_\alpha$. The control readout yields the Re, Im parts of these visibilities. *BRST overlay:* interpret (6.308)–(6.309) on $\mathcal{H}_{\mathrm{phys}}$ via $U^{\mathrm{phys}}_k := \mathsf{P}_{\mathrm{phys}} U_k \mathsf{P}_{\mathrm{phys}}$, $U^{\mathrm{phys}}_{k,\alpha} := \Pi_\alpha U^{\mathrm{phys}}_k \Pi_\alpha$, $\rho^{\mathrm{phys}}_0 := \mathsf{P}_{\mathrm{phys}} \rho_0 \mathsf{P}_{\mathrm{phys}}$, and ζ-closed $|\Omega_i\rangle$ of ghost number 0; by Lemma 6.243, $\partial_\xi \Lambda^{(\alpha)}_\bullet = \langle \zeta\, \eta^{(\alpha)}_\xi \rangle = 0$.

Theorem 6.261 (SSB arrow: symmetry of law → asymmetry of state)**.** *Assume the law is G-symmetric and the thermodynamic limit exists. Then, asymptotically:*

(a) Correlational channel (corr)

Long-range order and stiffness live here. Gapless NG modes appear as low-energy poles; their dispersion/soft responses are read in $\Lambda^{(\mathrm{corr})}_{\mathrm{law}}(\ell)$ (stable, up to irrelevant corrections) and in phase-twist shifts. BRST note: *NG analysis applies to* global *generators; in gauged sectors, would-be modes are ζ-exact and do not produce physical NG poles.*

(b) Vacuum channel (vac)

The order parameter and vacuum alignment live here. $\left|\Lambda^{(\mathrm{vac})}_{\mathrm{state}}(V)\right| \to 0$ *with* V *(orthogonality of distinct vacua), while* $\left|\Lambda^{(\mathrm{vac})}_{\mathrm{law}}(\ell)\right| \to 1$.

(c) Spectral channel (spec)

Phase-curvature and commutator structure live here. Intra-cone phase transport modifies NG couplings/dispersion without altering cones; visibility phases in $\Lambda^{(\mathrm{spec})}_{\mathrm{law}}(\ell)$ *isolate WZ/Berry-like contributions.*

Proof of Theorem 6.261. **Standing assumptions and channelization.** Assume the microscopic law is G-symmetric and anomaly-free, with visibilities evaluated in the physical sector $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ (Nielsen independence, Lemma 6.243). Let $\{\Pi_{\mathrm{corr}}, \Pi_{\mathrm{vac}}, \Pi_{\mathrm{spec}}\}$ be the BRST-closed (ghost number 0) channel projectors of the triadic split (Chapter 6, Definition 5.54). Define the *channel-restricted* holosymmetric functionals by inserting these projectors into the two-path readouts (Chapter 7):

$$\Lambda^{(\alpha)}_{\mathrm{law}}(\ell) := \mathrm{Tr}\Big[\Pi_\alpha\, U_B^\dagger(\ell)\, U_A(\ell)\, \Pi_\alpha\, \rho_0\Big], \qquad \Lambda^{(\alpha)}_{\mathrm{state}}(V) := \langle\Omega_2|\, \Pi_\alpha\, U^\dagger(V)\, U(V)\, \Pi_\alpha\, |\Omega_1\rangle.$$

Functoriality of the lift and cone preservation (Proposition 6.257) ensure these are block-diagonal, gauge-parameter independent in $\mathcal{H}_{\mathrm{phys}}$, and leave the $\bar{\Sigma}$-cones fixed (Chapter 8, § *Quadratic theory, propagators, and positivity*).

(a) Correlational channel — NG poles, stiffness, and phase-twist responses. Because the *law* is G-symmetric, the improved Noether currents J^μ_a obey exact Ward identities in $\mathcal{H}_{\mathrm{phys}}$ (Chapter 7, Relational Ward; with ζ-exact parts dropping in cohomology). Suppose a *global* subgroup of G is spontaneously broken in the thermodynamic limit. Then there exists a local operator ϕ with $\langle\Omega|[\widehat{Q}_a, \phi(0)]|\Omega\rangle \neq 0$, where $\widehat{Q}_a = \int d^dx\, J^0_a$ (suitably regulated). By the standard spectral/Lehmann representation (applied to BRST-closed, gauge-invariant probes), the *connected* two-point function in the correlational channel has a low-energy singularity:

$$\widetilde{G}_{\phi\phi}(k) \;=\; \int d^{d+1}x\, e^{ik\cdot x}\, \langle\Omega|\, \Pi_{\mathrm{corr}}\, [\phi(x), \phi(0)]\, \Pi_{\mathrm{corr}}\, |\Omega\rangle \;\sim\; \frac{Z_\phi}{\omega^2 - \omega^2_{\mathrm{NG}}(\mathbf{k})} + \text{regular},$$

with $\omega_{\mathrm{NG}}(\mathbf{k})$ the NG dispersion (linear in Lorentz-invariant cases; model-dependent exponents possible for $z \neq 1$; Chapter 8 fixes positivity of the residue). Equivalently, the current–order-parameter form factor $\langle\Omega|J^\mu_a(0)|\pi_b(\mathbf{k})\rangle$ is nonzero and saturates the pole. Because the law is symmetric, $\Lambda^{(\mathrm{corr})}_{\mathrm{law}}(\ell)$ receives only IR-irrelevant corrections (no explicit symmetry-breaking insertions in the correlational block), hence it is *stable on plateaus* and approaches unity in the UV probe $\ell \downarrow 0$ (as in Theorem 6.258, item (ii) for the law-side). Phase-twist interferometry: impose a uniform twist θ along a cycle, implemented by $U_\theta = \exp\big(i\theta\widehat{J}\big)$ in the correlational block. The ground-state energy density obeys $E_0(\theta) - E_0(0) = \frac{1}{2}\,\rho_s\,\theta^2 + o(\theta^2)$, defining the *stiffness* $\rho_s \geq 0$. The second twist derivative is a Kubo formula for the zero-frequency, small-$\mathbf{k}$ limit of the correlational current correlator and, equivalently, for the coefficient of the NG kinetic term read in the projected Hessian $\mathbb{H}_{\mathrm{phys}}$ (Chapter 8). Thus NG poles and stiffness "live" in the correlational channel and their soft responses are encoded in $\Lambda^{(\mathrm{corr})}_{\mathrm{law}}$ and twist-induced shifts.

BRST note. If the broken generators are *gauged*, Elitzur/BRST remove would-be NG modes from $\mathcal{H}_{\rm phys}$ (quartet mechanism in the longitudinal–ghost sector; Chapter 8, § *Transverse/longitudinal split & quartet decoupling*), so no physical NG pole appears in $\Pi_{\rm corr}$.

(b) Vacuum channel — order parameter and orthogonality of vacua. Let $\{|\Omega_\alpha\rangle\}$ be thermodynamic-limit vacua related by G; choose two distinct ones, $|\Omega_1\rangle, |\Omega_2\rangle$, and take preparations that select them (predicate stability, Chapter 7). Then the *vacuum* channel visibility is precisely their overlap measured in the $\Pi_{\rm vac}$ block:

$$\Lambda^{(\rm vac)}_{\rm state}(V) = \langle\Omega_2|\,\Pi_{\rm vac}U^\dagger(V)U(V)\Pi_{\rm vac}\,|\Omega_1\rangle \xrightarrow[V\uparrow\infty]{} 0,$$

because distinct SSB vacua are orthogonal in the thermodynamic limit (disjoint GNS representations, cluster decomposition). The order parameter lives in $\Pi_{\rm vac}$ as the one-point function $\langle\Omega_\alpha|\Pi_{\rm vac}\phi|\Omega_\alpha\rangle$, which is nonzero and branch-dependent in SSB phases. On the *law* side, exact G-symmetry implies that $\Lambda^{(\rm vac)}_{\rm law}(\ell) \to 1$ as $\ell \downarrow 0$ (only IR-irrelevant differences survive the Duhamel expansion, cf. the proof of Theorem 6.258); hence the vacuum block cleanly separates: law visibility stable $\uparrow$, state visibility vanishing $\downarrow$.

(c) Spectral channel — phase curvature, commutators, and WZ/Berry terms. The *spectral* block $\Pi_{\rm spec}$ isolates the commutator/phase-curvature content of low-energy transport. Let γ denote an adiabatic loop in the space of slow parameters (e.g. spatial twists, background sources) implemented in the $\Pi_{\rm spec}$ block by a unitary U_γ generated by BRST-closed sources. By cone preservation (Proposition 6.257), the principal symbol (and thus the $\bar\Sigma$-cones) are unchanged; only *intra-cone* phases and subprincipal terms are affected. Channel-restricted holosymmetric visibility acquires a geometric phase:

$$\arg\Lambda^{(\rm spec)}_{\rm law}[\gamma] = \arg\,\mathrm{Tr}\Big[\Pi_{\rm spec}\,U_\gamma\,\Pi_{\rm spec}\rho_0\Big] \equiv \oint_\gamma \mathcal{A} \quad \text{with} \quad d\mathcal{A} = \mathcal{F},$$

where $\mathcal{A}$ is the Berry connection on the relevant bundle of physical states and $\mathcal{F}$ its curvature (read off from Kubo–like response in $\Pi_{\rm spec}$). Equivalently, in effective-action language this is the Wess–Zumino/Wess–Zumino–Witten–type contribution confined to the spectral block; it shifts NG couplings/dispersion *without altering* the light-cone structure. Because the underlying law is symmetric and the lift is BRST-closed order-≤ 0, $\Lambda^{(\rm spec)}_{\rm law}(\ell)$ is stable on plateaus, and its *phase* isolates these geometric (WZ/Berry-like) contributions.

Conclusion. With a G-symmetric law and an existing thermodynamic limit, (a) the correlational block carries NG poles and stiffness (global SSB only), (b) the vacuum block exhibits orthogonality of distinct vacua with law visibility stable to 1, and (c) the spectral block captures geometric phase-curvature effects that renormalize intra-cone transport but not the cones. All statements are BRST-consistent, gauge-parameter independent in $\mathcal{H}_{\rm phys}$, and respect cone preservation; hence the asserted channel-wise signatures follow. □

Theorem 6.262 (Emergence arrow: asymmetry of law → symmetry of state)**.** *Assume symmetry-violating operators are RG-irrelevant and the state remains symmetric. Let* $\Delta := \min_k \Delta_k > d + z$ *be the minimal scaling dimension among the irrelevant breaking operators. Then, for* $\ell \downarrow 0$*:*

(a) Correlational

Microscopic anisotropies (e.g. gradient distortions) shrink with scale; quantitatively one has

$$1 - \left|\Lambda_{\text{law}}^{(\text{corr})}(\ell)\right| = \mathcal{O}\Big(\ell^{\Delta-(d+z)}\Big) \qquad (\ell \downarrow 0).$$

Equivalently, under IR coarse-graining,

$$1 - \left|\Lambda_{\text{law}}^{(\text{corr})}(\ell)\right| = \mathcal{O}\Big(\ell^{-(\Delta-(d+z))}\Big) \qquad (\ell \uparrow \infty).$$

(b) Vacuum

Explicit biases wash out under coarse-graining and no order parameter forms. In the UV–probe convention,

$$1 - \left|\Lambda_{\text{law}}^{(\text{vac})}(\ell)\right| = \mathcal{O}\Big(\ell^{\Delta-(d+z)}\Big) \qquad (\ell \downarrow 0),$$

while the state visibility saturates,

$$\left|\Lambda_{\text{state}}^{(\text{vac})}(V)\right| \to 1 \qquad (V \uparrow \infty).$$

Equivalently, along IR coarse–graining,

$$1 - \left|\Lambda_{\text{law}}^{(\text{vac})}(\ell)\right| = \mathcal{O}\Big(\ell^{-(\Delta-(d+z))}\Big) \qquad (\ell \uparrow \infty).$$

(c) Spectral

Dephasing/phase-curvature asymmetries are isolated by (φ, θ); for fixed (φ, θ) the visibility defect obeys

$$1 - \left|\Lambda_{\text{law}}^{(\text{spec})}(\ell; \varphi, \theta)\right| = \mathcal{O}\Big(\ell^{\Delta-(d+z)}\Big) \qquad (\ell \downarrow 0),$$

and hence decays along IR coarse-graining as $\mathcal{O}\Big(\ell^{-(\Delta-(d+z))}\Big)$ for $\ell \uparrow \infty$.

BRST note: *all statements hold in $\mathcal{H}_{\text{phys}}$ and are gauge-parameter independent (Nielsen-type identities); anomaly-induced breakdowns of BRST nilpotency are excluded by Stage I.*

Proof of Theorem 6.262. **Standing assumptions and notation.** Assume the gauge-fixed theory is anomaly-free and BRST-consistent; evaluate all quantities in $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$ so that gauge-parameter changes are ζ-exact and drop out of expectations (Nielsen independence, Lemma 6.243). The microscopic *law* contains symmetry-violating operators $\{\mathcal{O}_k\}$ with scaling dimensions $\Delta_k > d + z$ (RG-irrelevant); the *state* remains symmetric (unique G-invariant infinite-volume vacuum by predicate stability). Let Π_α, $\alpha \in \{\text{corr}, \text{vac}, \text{spec}\}$, be the BRST-closed channel projectors (Chapter 6), and define channel-restricted visibilities by inserting Π_α in the two-path readouts (Chapter 7). IR windowing is implemented by $K_{\text{IR}}^{(\ell)}$ and (when used) $P_x(\lambda_{\text{UV}})$; the holosymmetric lift is functorial and cone-preserving (Proposition 6.257).

RG scaling convention. Write the running couplings at momentum scale $\mu \sim \ell^{-1}$ as $g_k(\mu) = \mathcal{O}\Big(\mu^{\Delta_k-(d+z)}\Big)$ as $\mu \downarrow 0$ (IR flow); equivalently, $g_k(\ell) = \mathcal{O}\Big(\ell^{-(\Delta_k-(d+z))}\Big)$ as $\ell \uparrow \infty$. We will prove IR improvement in this standard form and then translate to the $\ell \downarrow 0$ (UV-probe) convention stated in the theorem via $\mu = 1/\ell$.

Law-side Duhamel control (channel-restricted). Let $U_A(\ell), U_B(\ell)$ be the two path-evolutions underlying Λ_{law} (Chapter 7), and set $\delta H(\ell') := H_A(\ell') - H_B(\ell') = \sum_k g_k(\ell')\,\mathcal{O}_k(\ell')$. For each channel α, a Duhamel/BCH expansion yields (interaction picture w.r.t. H_B)

$$\Pi_\alpha U_B^\dagger(\ell) U_A(\ell) \Pi_\alpha = \Pi_\alpha - i\int_0^\ell d\ell'\,\Pi_\alpha \widetilde{\delta H}(\ell')\Pi_\alpha - \int_{0<\ell''<\ell'<\ell} d\ell' d\ell''\,\Pi_\alpha \widetilde{\delta H}(\ell')\Pi_\alpha \widetilde{\delta H}(\ell'')\Pi_\alpha + \cdots . \tag{6.310}$$

On any plateau-stable band B (Remark 6.254), channel mixing is absent (functoriality) and operator norms are uniformly controlled. Therefore

$$1 - \left|\Lambda_{\mathrm{law}}^{(\alpha)}(\ell)\right| \;\le\; c_{1,\alpha}\int_0^\ell d\ell' \sum_k |g_k(\ell')| \;+\; c_{2,\alpha}\int_{0<\ell''<\ell'<\ell} d\ell' d\ell'' \sum_{k,k'} |g_k(\ell') g_{k'}(\ell'')| \;+\; \cdots . \tag{6.311}$$

(a) Correlational channel. Inserting $g_k(\ell') = \mathcal{O}\big(\ell'^{-(\Delta_k-(d+z))}\big)$ for $\ell' \uparrow \infty$ and summing over k gives

$$\int^\ell d\ell' \sum_k |g_k(\ell')| = \mathcal{O}\big(\ell^{-(\Delta-(d+z))}\big), \qquad \Delta := \min_k \Delta_k > d+z,$$

and similarly the quadratic term in (6.311) is subleading. Hence

$$1 - \left|\Lambda_{\mathrm{law}}^{(\mathrm{corr})}(\ell)\right| = \mathcal{O}\big(\ell^{-(\Delta-(d+z))}\big) \qquad (\ell \uparrow \infty). \tag{6.312}$$

This expresses that microscopic anisotropies (e.g. gradient distortions) shrink under coarse-graining. Translating to the UV-probe convention used in the theorem statement via $\mu = 1/\ell$ gives $1 - \left|\Lambda_{\mathrm{law}}^{(\mathrm{corr})}(\ell)\right| = \mathcal{O}\big(\ell^{\Delta-(d+z)}\big)$ as $\ell \downarrow 0$, which is the displayed scaling.

(b) Vacuum channel. Predicate stability with a unique symmetric vacuum implies $\Lambda_{\mathrm{state}}^{(\mathrm{vac})}(V) = \langle\Omega|\Omega\rangle \to 1$ as $V \uparrow \infty$ (standard cluster/thermodynamic limit; Chapter 8, § *Quadratic theory, propagators, and positivity*). There is no order parameter: for any local G-charged operator ϕ, $\langle\Omega|\phi|\Omega\rangle = 0$ by Ward identities (law symmetric). On the law side, the same estimate as in (a) with the Π_{vac} insertion yields $1 - \left|\Lambda_{\mathrm{law}}^{(\mathrm{vac})}(\ell)\right| = \mathcal{O}\big(\ell^{-(\Delta-(d+z))}\big)$ as $\ell \uparrow \infty$ (equivalently $\mathcal{O}(\ell^{\Delta-(d+z)})$ as $\ell \downarrow 0$), so $\left|\Lambda_{\mathrm{law}}^{(\mathrm{vac})}(\ell)\right| \to 1$.

(c) Spectral channel. The spectral block isolates phase-curvature/commutator structure. Let (φ, θ) denote slow source/twist parameters that implement adiabatic probes in Π_{spec}. Cone preservation (Proposition 6.257) fixes the principal symbol (hence the $\bar{\Sigma}$-cones); only *intra-cone* phases and subprincipal pieces are modified. Irrelevant symmetry breakings enter the effective geometric phase (Berry/WZ) through response coefficients proportional to $g_k(\mu)$. Therefore, along any fixed (φ, θ) contour,

$$\left|1 - \Lambda_{\mathrm{law}}^{(\mathrm{spec})}(\ell;\varphi,\theta)\right| \;\le\; C(\varphi,\theta)\sum_k \left|g_k(1/\ell)\right| \;=\; \mathcal{O}\big(\ell^{-(\Delta-(d+z))}\big) \qquad (\ell \uparrow \infty),$$

uniformly on plateaus. Equivalently, in the theorem's $\ell \downarrow 0$ convention this is $\mathcal{O}\big(\ell^{\Delta-(d+z)}\big)$: the dephasing/phase-curvature asymmetry scales away when the breaking is irrelevant, improving $\left|\Lambda_{\mathrm{law}}^{(\mathrm{spec})}(\ell)\right|$.

BRST and gauge-parameter independence. All channel projectors and scale tags are BRST-closed (ghost number 0); by Lemma 6.243, any change of gauge fixing modifies the gauge-fixed action by a ζ-exact term and thus leaves all channel-restricted visibilities invariant in $\mathcal{H}_{\mathrm{phys}}$. Anomaly-induced failure of BRST nilpotency is excluded by the Stage I assumption.

Conclusion. Irrelevant symmetry violations decay along the RG flow, which—through the channel-restricted Duhamel bounds—forces the law-side visibility defects to vanish with the power $\Delta - (d + z)$ in (6.312) and its $\Pi_{\mathrm{vac}}/\Pi_{\mathrm{spec}}$ analogues. The state remains symmetric (unique vacuum, no order parameter), hence $\left|\Lambda^{(\mathrm{vac})}_{\mathrm{state}}(V)\right| \to 1$. Translating the IR statement to the UV-probe variable reproduces the theorem's $\ell \downarrow 0$ asymptotics. This establishes items (a)–(c). □

Because the holosymmetric readouts in Theorem 6.262 are evaluated via BRST-closed, functorial, cone-preserving lifts that are block-diagonal in the triadic split, channel contributions add with no cross-terms and gauge-parameter changes generate only ζ-exact deformations—hence clean factorization in $\mathcal{H}_{\mathrm{phys}}$.

Corollary 6.263 (Why the factorization is clean (BRST-cohomological))**.** *By (6.306) and channel orthogonality, at the expectation level the attributions add without double counting, and cross-terms vanish. By (6.307), the holosymmetric lift respects the split, so the interferometric diagnostics are valid both per channel and in total. IR coarse-graining is performed with the kernel $K^{(\ell)}_{\mathrm{IR}}$ (window size ℓ), and—when needed—UV bandlimiting is implemented via the spectral projector $P_x(\lambda_{\mathrm{UV}})$; cone preservation ensures that channel-resolved phase effects never masquerade as kinematic changes in causal ordering.*

BRST overlay. *Choose the projectors Π_α BRST-closed (ghost number 0) so that $[\hat{Q}_{\mathrm{BRST}}, \Pi_\alpha] = 0$ and evaluate all quantities on $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$. Then: (i) ζ-exact improvements cannot mix channels, so additivity and orthogonality persist in cohomology; (ii) gauge-parameter (gauge-fermion) variations deform readouts by ζ-exact terms and hence leave channel and total visibilities invariant (Lemma 6.243); (iii) anomalous candidates (nilpotency failure or anomalous measure) are excluded upstream at Stage I. All equalities are understood modulo ζ-exact terms within $\mathcal{H}_{\mathrm{phys}}$.*

Proof of Corollary 6.263. **Step 1: Expectation-level additivity and vanishing cross terms.** By the triadic split (6.306), every channelized observable (or kernel) A decomposes as $A = \sum_\alpha \Pi_\alpha A \Pi_\alpha$ with mutually orthogonal, idempotent projectors $\{\Pi_\alpha\}_{\alpha\in\{\mathrm{corr,vac,spec}\}}$. Hence, for any trace-class state ρ,

$$\langle A\rangle = \mathrm{Tr}(A\rho) = \sum_\alpha \mathrm{Tr}(\Pi_\alpha A \Pi_\alpha \rho) = \sum_\alpha \mathrm{Tr}\Big(A\,\Pi_\alpha \rho \Pi_\alpha\Big),$$

where we used cyclicity of the trace and $\Pi_\alpha^2 = \Pi_\alpha$. Because the decomposition is orthogonal, mixed terms with $\alpha \neq \beta$ vanish: $\mathrm{Tr}(\Pi_\alpha A \Pi_\beta \rho) = \mathrm{Tr}(A \Pi_\beta \rho \Pi_\alpha) = 0$ by $\Pi_\alpha \Pi_\beta = 0$. Thus channel attributions *add without double counting* and cross terms drop out at the expectation level.

Step 2: Functoriality of the lift ⇒ blockwise diagnostics. By (6.307) (see also Proposition 6.257), the holosymmetric lift is block-diagonal over the triadic split:

$$\mathfrak{H}_{\lambda_{\mathrm{UV}},\ell}[A] = \sum_\alpha \Pi_\alpha\, K^{(\ell)}_{\mathrm{IR}}\, P_x(\lambda_{\mathrm{UV}})\, A\, P_x(\lambda_{\mathrm{UV}}) K^{(\ell)}_{\mathrm{IR}}\, \Pi_\alpha.$$

Therefore, the interferometric kernels entering the law/state visibilities (e.g. $U_B^\dagger U_A$ for Λ_{law} and the corresponding state-evolution product for Λ_{state}) remain block-diagonal. Consequently,

$$\Lambda_\square = \mathrm{Tr}\Big(\mathfrak{H}_{\lambda_{\text{UV}},\ell}[\mathcal{K}_\square]\ \rho_0\Big) = \sum_\alpha \mathrm{Tr}\Big(\Pi_\alpha\, \mathcal{K}_\square^{\text{lift}}\, \Pi_\alpha\, \rho_0\Big) = \sum_\alpha \mathrm{Tr}\Big(\mathcal{K}_\square^{\text{lift}}\, \Pi_\alpha \rho_0 \Pi_\alpha\Big),$$

where $\square \in \{\text{law}, \text{state}\}$ and $\mathcal{K}_\square$ denotes the corresponding interferometric kernel. Thus the *same* diagnostics are valid per channel and in total.

Step 3: Cone preservation ⇒ no kinematic masquerade. Let $\bar{\Sigma}_{\mu\nu}$ be the expectation-level chronogeometric tensor (Definition 5.26). $K_{\text{IR}}^{(\ell)}$ and $P_x(\lambda_{\text{UV}})$ are pseudodifferential of order ≤ 0, and Π_α acts on internal indices; hence they do not modify the principal symbol of the hyperbolic kinetic operators that determine causal propagation. Therefore the characteristic cone $\{k \neq 0 :\ \bar{\Sigma}^{\mu\nu} k_\mu k_\nu = 0\}$ is unchanged by the lift (Proposition 6.257). It follows that channel-resolved *phase* effects (subprincipal data, e.g. Berry/WZ-like phases in the spectral block) cannot be misread as changes in causal ordering: the cones—hence kinematics—remain fixed.

Step 4: BRST overlay in the physical sector. Choose the projectors Π_α BRST-closed (ghost number 0) so that $[\hat{Q}_{\text{BRST}}, \Pi_\alpha] = 0$, and evaluate all quantities on $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$. Then:

(i) Any ζ-exact improvement of kernels or currents changes $\mathcal{K}_\square \mapsto \mathcal{K}_\square + \{\hat{Q}_{\text{BRST}}, B\}$. Since insertions are BRST-closed and expectations are taken in $\mathcal{H}_{\text{phys}}$, one has $\mathrm{Tr}\Big(\Pi_\alpha\{\hat{Q}_{\text{BRST}}, B\}\Pi_\beta\, \rho_0\Big) = 0$ for all α, β, and in particular no channel mixing is induced. Thus additivity and orthogonality persist in cohomology.

(ii) Gauge-parameter (gauge-fermion) variations deform the gauge-fixed action by $\{\hat{Q}_{\text{BRST}}, \partial_\zeta \Psi\}$; by the Nielsen-type identity (Lemma 6.243), both channel and total visibilities are invariant in $\mathcal{H}_{\text{phys}}$.

(iii) Candidates with BRST nilpotency failure or anomalous measure are excluded at Stage I, so all equalities hold modulo ζ-exact terms within the physical sector.

Conclusion. Orthogonality ensures clean additivity and the absence of cross terms; functoriality of the lift preserves the split; cone preservation fixes kinematics while allowing only intra-cone phase modifications; and the BRST overlay guarantees gauge-parameter independence and stability in $\mathcal{H}_{\text{phys}}$. Hence the pair of visibilities $(\Lambda_{\text{law}}, \Lambda_{\text{state}})$ provides valid interferometric diagnostics per channel and in total, without altering the underlying causal structure. □

We have formalized an operational diagnostic layer that sits above the "laws vs. states" taxonomy. First, the unified chronogeometric operator and its expectation tensor are split into three additive, orthogonal channels—correlational, vacuum, and spectral—via the triadic decomposition (Definition 5.54). Orthogonality at the expectation level (see the channel results around Section 5.2.8) ensures that cross-terms are suppressed for the class of observables considered, so phenomena can be attributed to the channel in which they arise without double counting. On top of this structure, the holosymmetric lift is constructed to be linear (functorial) over the triadic split and to preserve the effective causal cones determined by the expectation tensor; only phases transported *within* the cones are modified (Definition 5.123, Proposition 5.124, and Corollary 5.126). IR averages are taken with $K_{\text{IR}}^{(\ell)}$, and UV control—whenever required in the derivations or scans—is provided by the spectral

projector $P_x(\lambda_{\mathrm{UV}})$. Consequently, the probe respects both the internal decomposition of the theory and its causal kinematics.

BRST reading. Throughout, we assume the gauge-fixed density $\mathcal{L} = \mathcal{L}_0 + \sum_k \epsilon_k \mathcal{O}_k + \zeta\Psi$ is anomaly-free and BRST-consistent (Stage I gate). All interferometric quantities are evaluated in the physical sector $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$, with insertions of ghost number 0; equalities are understood modulo ζ-exact terms. Gauge-parameter (gauge-fermion) variations act by ζ-exact deformations and hence leave the visibilities invariant (Nielsen-type identities; Lemma 6.243). In particular, statements about symmetry breaking refer to *global* G; would-be gauge Goldstones are cohomologically trivial (no literal gauge-SSB) [82, 91].

Operational content is provided by two coherence functionals defined by controlled, two-path interferometry. At the level of the law, the device compares alternative evolutions (e.g., micro law vs. coarse-grained IR law evaluated with $K_{\mathrm{IR}}^{(\ell)}$) to produce the *law-coherence* visibility $\Lambda_{\mathrm{law}}(\ell)$ in eq. (6.295). At the level of the state, coherent preparation of candidate vacua/ensembles yields the *state-coherence* visibility $\Lambda_{\mathrm{state}}(V)$ in eq. (6.298). Because the holosymmetric construction commutes with the channel projectors (cf. the functorial and block-diagonal behavior stated alongside Definition 5.54), these order parameters decompose additively by channel, allowing a clean diagnosis of correlational contributions (order/stiffness and long-distance propagation), vacuum contributions (alignment and order parameters), and spectral contributions (commutator-driven phase transport). The BRST overlay simply fixes the reading: $\Lambda_\bullet$ and their channel-resolved versions are computed on $\mathcal{H}_{\mathrm{phys}}$ and are insensitive to gauge-parameter choices.

The framework then identifies quadrant-specific signatures as asymptotics of these visibilities, thereby turning the conceptual 2×2 into a testable "operational cartography." In the unbroken regime (symmetric law, symmetric state), both visibilities remain near unity up to irrelevant corrections, consistent with exact Ward/Slavnov–Taylor identities and vanishing order parameters (Corollary 6.232; see also [82, 91]). In spontaneous symmetry breaking (symmetric law, asymmetric state), $|\Lambda_{\mathrm{law}}(\ell)| \approx 1$ while $|\Lambda_{\mathrm{state}}(V)|$ decays with volume and vanishes in the thermodynamic limit; phase-twist interferometry along the vacuum manifold reproduces Goldstone kinematics and soft limits [92, 94, 91]. For emergent/accidental symmetry (asymmetric micro law, symmetric state), the law-level visibility $|\Lambda_{\mathrm{law}}(\ell)|$ *improves* with coarse-graining according to a power set by the scaling dimension of the breaking operator, while the state-level visibility stays high and no order parameter develops, matching RG expectations [102, 82]. Under explicit breaking (asymmetric law, asymmetric state), $|\Lambda_{\mathrm{law}}(\ell)|$ remains bounded away from one with a deficit controlled by the breaking couplings, and the state sector is trivial (unique vacuum; no Goldstones) [91].

Finally, the construction is conservative with respect to the quadrant taxonomy: the holosymmetric overlay refines readout without changing quadrant membership. This is proved by combining (i) the readout lemmas for the two interferometers, (ii) functoriality over the triadic split, (iii) cone preservation/no–law–signalling, and (iv) stability of the law/state predicates under lifting, culminating in Proposition 6.251. Together with the quadrant identifications in Corollary 6.232 and the global map of Section 6.11 and table 6.1, the "operational cartography" theorem (Theorem 6.258) makes explicit how the visibilities $\Lambda_{\mathrm{law}}(\ell)$ and $\Lambda_{\mathrm{state}}(V)$ serve as measurable, channel-resolved order parameters for the four regimes—now read cohomologically on $\mathcal{H}_{\mathrm{phys}}$ and invariant under ζ-exact gauge choices.

6.12 Scale-Aware Analysis for the Laws–State Matrix

6.12.1 Purpose and Proviso

We will traverse all four cells of the Laws–State matrix using a single variational umbrella and a single set of operational diagnostics. Throughout, any assignment of fluctuation modes to "Higgs-like / graviton-like / apeiron / chrono–kymon" remains **provisional** until: (i) the full multi-quadrant analysis is complete; and (ii) all identifications are **scale-stable** under the UV/IR knobs introduced below—namely the spectral projector $P_x(\lambda_{\mathrm{UV}})$ and the IR windowing kernel $K_{\mathrm{IR}}^{(\ell)}$ (cf. Section 6.12.4). Holosymmetric visibilities provide the operational readouts that distinguish quadrants without changing quadrant membership (functoriality and cone-preservation in the sense of Chapter 6), and their channel-resolved versions separate correlational (T-even), vacuum (T-even), and spectral (T-odd) content.

BRST reading (proviso). We assume an anomaly-free, BRST-consistent gauge fixing of the form

$$\mathcal{L} = \mathcal{L}_0 + \sum_k \epsilon_k \mathcal{O}_k + \zeta\Psi,$$

and evaluate physical quantities on the BRST-closed sector $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ with ghost number 0. All equalities are understood modulo ζ-exact terms; gauge-parameter (gauge-fermion) deformations act by ζ-exact variations and thus leave visibilities and scale-tagged functionals invariant (Nielsen-type independence; cf. Lemmas 6.233 and 6.243).

6.12.2 Variational Backbone for Each Quadrant

Unified relational action (state-weighted, block-projectable)

Let $\hat{\mathbf{F}} = (\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}})^\top$ be the *Generalized Field Operator* (see Definition 6.1) and let

$$\hat{\mathbf{G}} = \begin{pmatrix} \hat{G}_{\Phi\Phi} & \hat{G}_{\Phi\Theta} \\ \hat{G}_{\Theta\Phi} & \hat{G}_{\Theta\Theta} \end{pmatrix}$$

denote the *unified inter-modal correlator* (matrix form and definition in Section 6.1.1 and Definition 6.4). The global variational umbrella is the *Relational Action Functional* introduced in Chapter 7,

$$S = \int_{\mathcal{X}_q} d^d x \, \sqrt{-\det \bar{\Sigma}(x)} \; \mathrm{Tr}\Big(\langle \mathbf{\Psi} | \, \hat{\mathbf{G}}(x,x) \, | \mathbf{\Psi} \rangle \Big). \tag{6.313}$$

Scale-aware convention: when UV bandlimiting is activated we write $S^{(\lambda_{\mathrm{UV}})}$ with the bandlimited correlator $\hat{\mathbf{G}}^{(\lambda_{\mathrm{UV}})}$, and in the fully scale-tagged variant include IR averaging with $K_{\mathrm{IR}}^{(\ell)}$ as in Section 6.12.4.

BRST overlay. Physical variations and readouts use the projected correlator

$$\hat{\mathbf{G}}^{\mathrm{phys}} := \mathsf{P}_{\mathrm{phys}} \, \hat{\mathbf{G}} \, \mathsf{P}_{\mathrm{phys}},$$

so that S may be read cohomologically as

$$S_{\mathrm{phys}} = \int_{\mathcal{X}_q} d^d x \, \sqrt{-\det \bar{\Sigma}} \; \mathrm{Tr}\Big(\langle \mathbf{\Psi} | \, \hat{\mathbf{G}}^{\mathrm{phys}}(x,x) \, | \mathbf{\Psi} \rangle \Big),$$

with invariance under gauge-parameter changes ensured by $\partial_\xi S_{\text{phys}} = \langle \zeta\, \eta_\xi \rangle = 0$ (cf. Lemma 6.243). The holosymmetric visibilities and their channel-resolved versions are computed from S_{phys} and are thus gauge-parameter independent.

Sector actions (geometry/matter) with a common bundle connection

Split covariant derivative (recall). We use the bundle-covariant derivative that combines the Relational Levi–Civita transport on spacetime indices with the operator-bundle connection on fibre degrees of freedom:

$$\nabla_\mu := \nabla_\mu^{(\text{rel})} + \Big(\partial_\mu + i[\mathcal{A}_\mu, \cdot]\Big), \tag{6.314}$$

where $\nabla_\mu^{(\text{rel})}$ is the Relational Levi–Civita derivative (see Definition 6.7 and eq. (6.22)) and $\mathcal{A}_\mu$ is the operator-bundle connection introduced in Definition 6.192. *BRST overlay:* $\mathcal{A}_\mu$ and the field multiplets form BRST doublets with respect to ζ; ∇_μ is understood modulo ζ-exact improvements, which do not affect $\mathcal{H}_{\text{phys}}$ observables.

$$\begin{aligned} S_\Phi &= \int d^d x \, \sqrt{-\det \bar{\Sigma}}\ \mathrm{Tr}\Big\{ \bar{\Sigma}^{\mu\nu}\, g_T\Big(\nabla_\mu \hat{\mathbf{\Phi}}, \nabla_\nu \hat{\mathbf{\Phi}}\Big) + \mathcal{V}_\Phi(\hat{\mathbf{\Phi}}) \Big\}, \\ S_\Theta &= \int d^d x \, \sqrt{-\det \bar{\Sigma}}\ \mathrm{Tr}\Big\{ \bar{\Sigma}^{\mu\nu}\, g_C\Big(\nabla_\mu \hat{\mathbf{\Theta}}, \nabla_\nu \hat{\mathbf{\Theta}}\Big) + \mathcal{V}_\Theta(\hat{\mathbf{\Theta}}) \Big\}. \end{aligned} \tag{6.315}$$

The per-sector Lagrangians and actions (operator-valued) for the Gaiamiti and Tantrakyma fields (defined in Definitions 6.142 and 11.17) read:

$$S_\Phi = \int d^d x \, \sqrt{-\det \bar{\Sigma}}\ \mathrm{Tr}\Big\{ \bar{\Sigma}^{\mu\nu} g_T\Big(\nabla_\mu \hat{\mathbf{\Phi}}, \nabla_\nu \hat{\mathbf{\Phi}}\Big) + \mathcal{V}_\Phi(\hat{\mathbf{\Phi}}) \Big\}. \tag{6.316}$$

Definitions 6.172 and 6.174 and eqs. (6.243) and (6.247)

$$S_\Theta = \int d^d x \, \sqrt{-\det \bar{\Sigma}}\ \mathrm{Tr}\Big\{ \bar{\Sigma}^{\mu\nu} g_C\Big(\nabla_\mu \hat{\mathbf{\Theta}}, \nabla_\nu \hat{\mathbf{\Theta}}\Big) + \mathcal{V}_\Theta(\hat{\mathbf{\Theta}}) \Big\}. \tag{6.317}$$

Definitions 9.301 and 9.310 and eqs. (9.129) and (9.146) Here g_T is the canonical Hermitian inner product on the vacuum *tangent* space (see Definition 6.104), while g_C is the operator-valued *co-metric* on the *cotangent* space with inverse g_C^{-1} used in the minimal Θ-potential (see Remark 9.303, Section 9.18.5, Definition 9.335, and eq. (9.133)). The split covariant derivative used in both sectors is the one already defined in eq. (6.314).

Gauge-fixing/ghost layer (cohomological hygiene). Add the BRST-exact gauge-fixing+ghost piece

$$S_{\text{gf+gh}} = \int d^d x \, \sqrt{-\det \bar{\Sigma}}\, \zeta \Psi,$$

with gauge fermion Ψ chosen so that $S_{\text{tot}} := S_\Phi + S_\Theta + S_{\text{gf+gh}}$ is local and preserves the cone structure. By Lemma 6.233 and Nielsen-type identities (Lemma 6.243), variations of Ψ shift S_{tot} by ζ-exact terms and do not affect physical observables, holosymmetric visibilities, or their scale-tagged scans in $\mathcal{H}_{\text{phys}}$. Anomalous (BRST-inconsistent) candidates are excluded upstream (Stage I).

Proposition 6.264 (Cohomological invariance of scale-tagged observables)**.** *Assume anomaly-free BRST (measure invariant, nilpotent ζ) and a gauge-fermion family $\Psi(\xi)$. Let the IR windowing $K_{\mathrm{IR}}^{(\ell)}$ and the UV projector $P_x(\lambda_{\mathrm{UV}})$ be* BRST-closed, ghost-number 0 *operators, i.e.*

$$[\hat{Q}_{\mathrm{BRST}}, K_{\mathrm{IR}}^{(\ell)}] = 0, \qquad [\hat{Q}_{\mathrm{BRST}}, P_x(\lambda_{\mathrm{UV}})] = 0,$$

and likewise for the channel projectors Π_α with $[\hat{Q}_{\mathrm{BRST}}, \Pi_\alpha] = 0$. Then for every ζ-closed, ghost-number-0 insertion $\mathcal{O}$,

$$\frac{\partial}{\partial \xi} \left\langle \mathcal{O} \right\rangle_{\mathcal{H}_{\mathrm{phys}}} = \left\langle \zeta\, \eta_\xi \right\rangle_{\mathcal{H}_{\mathrm{phys}}} = 0, \tag{6.318}$$

where η_ξ is the local functional generated by $\partial_\xi \Psi$ (Nielsen-type identity; cf. Lemma 6.243). Consequently, all the following scale-tagged quantities are gauge-parameter independent *and invariant under ζ-exact improvements (including Belinfante/BRST current improvements and antisymmetric divergences):*

$$\frac{\partial}{\partial \xi} S_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)} = 0, \qquad \frac{\partial}{\partial \xi} \Lambda_{\mathrm{law}}(\ell) = 0, \qquad \frac{\partial}{\partial \xi} \Lambda_{\mathrm{state}}(V) = 0, \tag{6.319}$$

$$\frac{\partial}{\partial \xi} \Lambda_{\mathrm{law}}^{(\alpha)}(\ell) = 0, \qquad \frac{\partial}{\partial \xi} \Lambda_{\mathrm{state}}^{(\alpha)}(V) = 0, \qquad \alpha \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}. \tag{6.320}$$

Full proof of Proposition 6.264. **Set–up and BRST data.** Work with the gauge-fixed action $S_{\mathrm{gf+gh}}(\xi) = S_0 + \{\hat{Q}_{\mathrm{BRST}}, \Psi(\xi)\}$, where $\hat{Q}_{\mathrm{BRST}}$ is nilpotent and the path–integral measure is BRST invariant (anomaly-free hypothesis). The BRST differential is $\zeta(\cdot) = \{\hat{Q}_{\mathrm{BRST}}, \cdot\}$. The IR window $K_{\mathrm{IR}}^{(\ell)}$, the UV projector $P_x(\lambda_{\mathrm{UV}})$, and the channel projectors Π_α are assumed BRST-closed, ghost-number 0: $[\hat{Q}_{\mathrm{BRST}}, K_{\mathrm{IR}}^{(\ell)}] = [\hat{Q}_{\mathrm{BRST}}, P_x(\lambda_{\mathrm{UV}})] = [\hat{Q}_{\mathrm{BRST}}, \Pi_\alpha] = 0$. We evaluate all correlators in the physical sector $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$, equivalently inserting the projector $\mathsf{P}_{\mathrm{phys}}$ that commutes with $\hat{Q}_{\mathrm{BRST}}$.

1) Nielsen identity in the physical sector. Let $\mathcal{O}$ be any BRST-closed, ghost-number 0 insertion: $\zeta\, \mathcal{O} = 0$, $\mathrm{gh}(\mathcal{O}) = 0$. Differentiate the expectation value with respect to the gauge parameter ξ. In either operator (Dyson) or path–integral language one has

$$\partial_\xi S_{\mathrm{gf+gh}}(\xi) = \{\hat{Q}_{\mathrm{BRST}},\, \partial_\xi \Psi(\xi)\} =: \zeta\, \eta_\xi, \quad \mathrm{gh}(\eta_\xi) = -1.$$

Thus

$$\frac{\partial}{\partial \xi} \left\langle \mathcal{O} \right\rangle_{\mathcal{H}_{\mathrm{phys}}} = \left\langle \mathcal{O} \left(\partial_\xi S_{\mathrm{gf+gh}} \right) \right\rangle_{\mathcal{H}_{\mathrm{phys}},\, \mathrm{conn}} = \left\langle \mathcal{O}\, \zeta\, \eta_\xi \right\rangle_{\mathcal{H}_{\mathrm{phys}}} \tag{6.321}$$

$$= \left\langle \zeta\, (\eta_\xi \mathcal{O}) \right\rangle_{\mathcal{H}_{\mathrm{phys}}} - \left\langle (\zeta\, \mathcal{O})\, \eta_\xi \right\rangle_{\mathcal{H}_{\mathrm{phys}}} \qquad \text{(graded Leibniz)} \tag{6.322}$$

$$= \left\langle \zeta\, (\eta_\xi \mathcal{O}) \right\rangle_{\mathcal{H}_{\mathrm{phys}}} \qquad (\zeta\, \mathcal{O} = 0). \tag{6.323}$$

Anomaly freedom implies BRST invariance of the measure, hence the expectation of a ζ-exact, ghost-number 0 operator vanishes in the physical sector: $\left\langle \zeta\, X \right\rangle_{\mathcal{H}_{\mathrm{phys}}} = 0$ for any X for which the correlator is defined. Therefore

$$\frac{\partial}{\partial \xi} \left\langle \mathcal{O} \right\rangle_{\mathcal{H}_{\mathrm{phys}}} = \left\langle \zeta\, \eta_\xi \right\rangle_{\mathcal{H}_{\mathrm{phys}}} = 0,$$

which is the Nielsen-type identity (6.318).

2) Stability under scale tags and channel projectors. Let $\mathcal{O}_\alpha^{(\lambda_{\rm UV},\ell)}$ denote any *scale-tagged* and *channel-resolved* version of $\mathcal{O}$, obtained by inserting $K_{\rm IR}^{(\ell)}$, $P_x(\lambda_{\rm UV})$, and Π_α in any block-diagonal (functorial) manner, e.g.

$$\begin{aligned}\mathcal{O}_\alpha^{(\lambda_{\rm UV},\ell)} = \Pi_\alpha\, K_{\rm IR}^{(\ell)}\, P_x(\lambda_{\rm UV})\, \mathcal{O} \\ \times\, P_x(\lambda_{\rm UV})\, K_{\rm IR}^{(\ell)}\, \Pi_\alpha.\end{aligned} \tag{6.324}$$

Because each tag commutes with $\hat{Q}_{\rm BRST}$ and has ghost-number 0, $\zeta\,\mathcal{O}_\alpha^{(\lambda_{\rm UV},\ell)} = \Pi_\alpha K_{\rm IR}^{(\ell)} P_x(\lambda_{\rm UV})(\zeta\,\mathcal{O}) P_x(\lambda_{\rm UV}) K_{\rm IR}^{(\ell)} \Pi_\alpha = 0$. Thus the same derivation yields

$$\frac{\partial}{\partial\xi}\left\langle \mathcal{O}_\alpha^{(\lambda_{\rm UV},\ell)}\right\rangle_{\mathcal{H}_{\rm phys}} = \left\langle \zeta\Big(\eta_\xi\, \mathcal{O}_\alpha^{(\lambda_{\rm UV},\ell)}\Big)\right\rangle_{\mathcal{H}_{\rm phys}} = 0,$$

and, by summing over α or dropping Π_α, the channel-summed (total) version is also invariant. Therefore *any* BRST-closed observable remains gauge-parameter independent after scale tagging and channel restriction.

3) Application to the listed objects. *(i) Physical action* $S_{\rm phys}^{(\lambda_{\rm UV},\ell)}$. By construction in Chapter 8, $S_{\rm phys}^{(\lambda_{\rm UV},\ell)}$ is the effective action restricted to $\mathcal{H}_{\rm phys}$, obtained from kernels that are BRST-closed up to ζ-exact pieces; hence it is a functional of BRST-closed, scale-tagged operators. Applying Step 2 shows $\partial_\xi S_{\rm phys}^{(\lambda_{\rm UV},\ell)} = 0$.

(ii) Law visibility $\Lambda_{\rm law}(\ell)$. Use the two-path definition (Chapter 7) in the presence of tags:

$$\Lambda_{\rm law}(\ell) = {\rm Tr}\Big[K_{\rm IR}^{(\ell)} P_x(\lambda_{\rm UV})\, U_B^\dagger(\ell) U_A(\ell)\, P_x(\lambda_{\rm UV}) K_{\rm IR}^{(\ell)}\, \rho_0\Big]\,.$$

Differentiating w.r.t. ξ inserts $\partial_\xi S_{\rm gf+gh} = \zeta\,\eta_\xi$ along the Dyson series for U_A, U_B. Since tags commute with $\hat{Q}_{\rm BRST}$ and we evaluate the trace in $\mathcal{H}_{\rm phys}$ (equivalently, we may sandwich with $\mathsf{P}_{\rm phys}$), the variation is the expectation of a ζ-exact, ghost-number 0 insertion and vanishes by Step 1. The same argument with Π_α inserted yields $\partial_\xi \Lambda_{\rm law}^{(\alpha)}(\ell) = 0$.

(iii) State visibility $\Lambda_{\rm state}(V)$. With scale tags and optional channel projector,

$$\Lambda_{\rm state}^{(\alpha)}(V) = \langle\Omega_2|\, \Pi_\alpha K_{\rm IR}^{(\ell)} P_x(\lambda_{\rm UV})\, U^\dagger(V) U(V)\, P_x(\lambda_{\rm UV}) K_{\rm IR}^{(\ell)} \Pi_\alpha\, |\Omega_1\rangle.$$

As above, ∂_ξ acts by inserting $\zeta\,\eta_\xi$ in the Dyson expansion. Because the tags and Π_α commute with $\hat{Q}_{\rm BRST}$ and the external states are evaluated in $\mathcal{H}_{\rm phys}$, the same Nielsen step shows $\partial_\xi \Lambda_{\rm state}^{(\alpha)}(V) = 0$, hence also $\partial_\xi \Lambda_{\rm state}(V) = 0$ after summing over α.

4) Invariance under ζ-exact improvements and divergences. Let a kernel or a current be changed by a BRST improvement or a covariant divergence,

$$\mathcal{K} \;\longmapsto\; \mathcal{K}' = \mathcal{K} + \zeta X + \nabla_\nu X^{[\mu\nu]}.$$

In $\mathcal{H}_{\rm phys}$, the expectation of ζX vanishes as in Step 1. The divergence term integrates to a surface contribution; under the standard falloff/boundary conditions used to define the thermodynamic and cluster limits (and, for compact support smearings, by Stokes' theorem),

the surface term does not contribute to the tagged, finite readouts. Hence all listed quantities are invariant under such improvements.

Conclusion. For BRST-closed, ghost-number 0 scale tags and channel projectors, the gauge-fermion variation inserts a BRST-exact operator whose expectation vanishes in the physical sector. Therefore $S_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}$, the law/state visibilities, and their channel-resolved versions are gauge-parameter independent and stable under ζ-exact (and divergence) improvements, establishing the proposition. □

Corollary 6.265 (Plateau and exponent invariance)**.** *With $K_{\mathrm{IR}}^{(\ell)}$ and $P_x(\lambda_{\mathrm{UV}})$ as in Proposition 6.264, the plateau locations and asymptotic exponents extracted from*

$$\Lambda_{\mathrm{law}}(\ell), \quad \Lambda_{\mathrm{state}}(V), \quad \Lambda_{\mathrm{law}}^{(\alpha)}(\ell), \quad \Lambda_{\mathrm{state}}^{(\alpha)}(V)$$

are independent of the gauge parameter ξ and of ζ-exact improvements. Thus the "operational cartography" (quadrant assignment and channel attributions) is cohomologically well defined and gauge-fermion invariant, up to finite-size or finite-window effects governed by Section 6.12.4.

Proof of Corollary 6.265. **Standing hypotheses.** All expectations are taken in the physical sector $\mathcal{H}_{\mathrm{phys}}$, the BRST charge is nilpotent and the measure is BRST invariant (Stage I), and the scale tags $K_{\mathrm{IR}}^{(\ell)}$, $P_x(\lambda_{\mathrm{UV}})$ as well as the channel projectors Π_α are BRST-closed, ghost-number 0. Consequently, the Nielsen-type identity of Proposition 6.264 holds:

$$\partial_\xi \langle \mathcal{O} \rangle_{\mathcal{H}_{\mathrm{phys}}} = \big\langle \zeta\, \eta_\xi \big\rangle_{\mathcal{H}_{\mathrm{phys}}} = 0 \qquad \text{for any BRST-closed ghost-number-0 } \mathcal{O}. \tag{6.325}$$

1) Gauge-parameter independence of the tagged visibilities (pointwise in scale). Apply (6.325) with

$$\mathcal{O} \in \Big\{ \Lambda_{\mathrm{law}}(\ell),\ \Lambda_{\mathrm{state}}(V),\ \Lambda_{\mathrm{law}}^{(\alpha)}(\ell),\ \Lambda_{\mathrm{state}}^{(\alpha)}(V) \Big\}, \quad \alpha \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}.$$

By Proposition 6.264, each of these is BRST-closed after inserting the tags and (optionally) Π_α, hence

$$\partial_\xi \Lambda_\square(\cdot) = 0, \qquad \partial_\xi \Lambda_\square^{(\alpha)}(\cdot) = 0, \quad \square \in \{\mathrm{law}, \mathrm{state}\}, \tag{6.326}$$

for every fixed scale argument (ℓ or V). Thus *as functions of scale*, the visibilities are identical for all ξ.

2) Plateau invariance. Let B denote any "plateau-stable band" as in Remark 6.254 and Section 6.12.4, e.g.

$$B := \Big\{ (\lambda_{\mathrm{UV}}, \ell) : \sup_{(\lambda_{\mathrm{UV}},\ell)\in B} \big|\delta\Lambda_\square\big| \le \varepsilon_{\mathrm{tol}} \Big\},$$

or, equivalently, the set on which prescribed smoothness/flatness tolerances hold (finite-size/finite-window corrections included). Because (6.326) is *pointwise* in ℓ and V, any predicate of the form

$$\mathcal{P}[\Lambda_\square](\lambda_{\mathrm{UV}}, \ell) \in \{\mathrm{true}, \mathrm{false}\} \quad (\text{respectively in } V)$$

has the same truth value for all ξ. Hence the collection of scale points satisfying the plateau criterion is *identical* for all ξ; i.e. plateau *locations* are gauge-parameter independent. The same reasoning applies to the channel-restricted visibilities by inserting Π_α.

3) Asymptotic exponent invariance. Write a generic law-side asymptotic on a plateau (for definiteness in the IR variable ℓ) as

$$1 - \left|\Lambda_{\text{law}}(\ell)\right| = C\,\ell^{-\beta}\left(1 + r(\ell)\right), \qquad \beta > 0, \quad r(\ell) \xrightarrow[\ell\to\infty]{} 0,$$

and analogously for the channel version and for the state-side finite-volume approach. Define the (logarithmic) *effective exponent* by

$$\beta_{\text{eff}}(\ell) := -\frac{d\log\left(1 - |\Lambda_{\text{law}}(\ell)|\right)}{d\log\ell}.$$

Differentiating with respect to ξ and using (6.326) gives

$$\partial_\xi \beta_{\text{eff}}(\ell) = -\frac{d}{d\log\ell}\left[\frac{\partial_\xi\left(1 - |\Lambda_{\text{law}}(\ell)|\right)}{1 - |\Lambda_{\text{law}}(\ell)|}\right] = 0.$$

Therefore $\beta_{\text{eff}}(\ell)$ (and hence its plateau limit β) is ξ-independent. The same argument applies verbatim to the channel-restricted visibilities and to exponents extracted in the UV variable $\mu \sim 1/\ell$ or in the volume variable V.

4) Invariance under ζ-exact improvements (including current improvements). Let an admissible improvement modify a kernel or current by $\mathcal{K} \mapsto \mathcal{K}' = \mathcal{K} + \zeta X + \nabla_\nu X^{[\mu\nu]}$. In $\mathcal{H}_{\text{phys}}$, ζX has vanishing expectation, and the divergence integrates to boundary/surface terms that are controlled by the finite-window/finite-size set-up of Section 6.12.4. Hence the *functions* Λ are unchanged on plateaus up to subleading, controllable corrections $o_B(1)$, which do not affect the validity of the plateau predicate nor the value of any asymptotic exponent (multiplicative constants and $o_B(1)$ corrections cancel in logarithmic slopes). Therefore both plateau locations and exponents are invariant under such improvements.

5) Conclusion. Since (i) the tagged visibilities are ξ-independent pointwise in scale, (ii) the plateau predicates depend only on those values, and (iii) ζ-exact (and divergence) improvements do not alter the visibilities in $\mathcal{H}_{\text{phys}}$ beyond tolerated finite-window/finite-size remainders, both the *plateau bands* and the *asymptotic exponents* extracted from $\Lambda_{\text{law}}, \Lambda_{\text{state}}, \Lambda^{(\alpha)}_{\text{law}}, \Lambda^{(\alpha)}_{\text{state}}$ are cohomologically and gauge-fermion invariant (up to the finite-size/window effects already codified in Section 6.12.4). This proves the corollary. □

Remark 6.266 (Boundary of validity)**.** The cohomological (gauge-fermion) invariance results rely on the following hypotheses; stepping outside them voids the conclusions or requires extra care.

(a) Anomaly-free BRST only. If BRST is anomalous—either through nilpotency failure or a non-invariant measure—then the Nielsen identity acquires an extra term and Proposition 6.264 no longer applies. Such candidates are excluded at *Stage I* (vanishing Slavnov functional / cohomology test for $H^{1,d}(\zeta\,|\,d)$).

(b) Insertions must be BRST-closed, ghost number 0. Operators (including scale tags and channel projectors) that are not ζ-closed or do not have ghost number 0 fall outside scope and are not used as diagnostics. If approximate closure holds, e.g. $[\hat{Q}_{\mathrm{BRST}}, K_{\mathrm{IR}}^{(\ell)}] = \varepsilon_{\mathrm{tag}} R$, then

$$\left|\partial_\xi \langle \mathcal{O} \rangle_{\mathcal{H}_{\mathrm{phys}}}\right| \;\le\; C\,\varepsilon_{\mathrm{tag}},$$

so invariance persists only up to controlled $o(1)$ errors; we do not rely on such approximations in the main statements.

(c) BRST-preserving regularization/renormalization. A regulator that breaks BRST (without compensating counterterms) induces spurious gauge-parameter dependence. We assume algebraic renormalization (or equivalent) restoring Slavnov–Taylor identities at each step; otherwise the invariance can fail.

(d) Finite-size/window artefacts. All gauge-parameter and cohomology invariances hold *pointwise* in scale; plateau/exponent statements are valid up to the finite-size/finite-window tolerances specified in Section 6.12.4. Outside plateau bands one expects effective-exponent drift.

(e) Nonlocal/topological probes. Line/surface operators must be made BRST-closed and gauge-invariant (including framing choices) before use. Uncompleted nonlocal probes can mix channels or mimic explicit breaking, invalidating the diagnostics.

(f) Global gauge-fixing issues (Gribov regions). The Nielsen identity is derived in a BRST-valid patch of configuration space. Nonperturbative obstructions (e.g. Gribov horizons) require a separate treatment; our results apply where BRST cohomology is well-defined (the physical sector $\mathcal{H}_{\mathrm{phys}}$).

(g) Operator mixing under RG. If renormalization mixes a diagnostic with lower-dimension, *non*-BRST-closed operators (e.g. via symmetry-breaking counterterms), that constitutes explicit breaking (Quadrant IV) rather than a failure of the invariance statements; the cartography then reflects the new law asymmetry.

(h) Time-/scale-dependent tags. When $K_{\mathrm{IR}}^{(\ell)}$, $P_x(\lambda_{\mathrm{UV}})$, or Π_α depend on the evolution parameter, the commutators $[\hat{Q}_{\mathrm{BRST}}, \cdot]$ must vanish *pointwise* in that parameter. Otherwise extra commutator terms appear and must be bounded explicitly.

(i) Boundary terms. Improvements by covariant divergences are harmless under the falloff/smearing assumptions used for thermodynamic and plateau limits. With physical boundaries, edge contributions must be kept BRST-closed or treated separately; the invariance statements refer to the bulk.

In short: anomaly-free BRST, BRST-closed ghost-number 0 insertions (including tags/projectors), and BRST-preserving renormalization are the envelope of validity; within it, the cartography and plateau/exponent extractions are cohomologically and gauge-fermion invariant up to the controlled tolerances already declared.

Gaiamiti Euler–Lagrange operator and field equation.

Definition 6.267 (Gaiamiti Euler–Lagrange operator and field equation)**.** Let $(\mathcal{X}_q, \bar{\Sigma}_{\mu\nu})$ be the chronogeometric background, with inverse $\bar{\Sigma}^{\mu\nu}$ and torsion–free, $\bar{\Sigma}$–compatible connection

∇ (i.e. $\nabla_\rho \bar{\Sigma}_{\mu\nu} = 0$; see Definition 6.184 and Section 6.5.3 and cf. [375]). The *Gaiamiti field* is an operator–valued section $\hat{\mathbf{\Phi}} : \mathcal{X}_q \to \mathcal{E}_\Phi$ of the Gaiamiti configuration bundle $\mathcal{E}_\Phi \to \mathcal{X}_q$ (cf. Definition 6.118 and Section 6.2; see also [230]). On each fiber $(\mathcal{E}_\Phi)_x$ we fix a positive, nondegenerate bilinear pairing g_T (the *kinetic/metric pairing*). Let $\mathcal{G}_T : (\mathcal{E}_\Phi)_x \to (\mathcal{E}_\Phi^\star)_x$ be the associated Riesz (Legendre) isomorphism, characterized by

$$\langle A, \mathcal{G}_T(B) \rangle \;=\; g_T(A, B) \qquad \text{for all } A, B \in (\mathcal{E}_\Phi)_x,$$

where $\langle \cdot, \cdot \rangle$ denotes the canonical fiberwise duality pairing (cf. [253, 433]). Let $\mathcal{V}_\Phi(\hat{\mathbf{\Phi}})$ be the (fiberwise) potential density for $\hat{\mathbf{\Phi}}$, and let $\mathcal{V}_{\mathrm{mix}}(\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}})$ be an optional mixing potential with the Tantrakyma field $\hat{\mathbf{\Theta}}$ (cf. Definitions 6.3 and 6.142 and Section 6.12.2; see also [82, 155]). The *canonical momentum current* of $\hat{\mathbf{\Phi}}$ is the $\mathcal{E}_\Phi^\star$–valued vector field

$$\Pi^\mu_{\mathbf{\Phi}} \;:=\; \bar{\Sigma}^{\mu\nu}\, \mathcal{G}_T\Big(\nabla_\nu \hat{\mathbf{\Phi}}\Big).$$

The *Gaiamiti Euler–Lagrange operator* is the $\mathcal{E}_\Phi^\star$–valued section

$$\mathcal{E}_{\mathbf{\Phi}} \;:=\; -\nabla_\mu\Big(\Pi^\mu_{\mathbf{\Phi}}\Big) \;+\; \frac{\partial \mathcal{V}_\Phi}{\partial \hat{\mathbf{\Phi}}} \;+\; \frac{\partial \mathcal{V}_{\mathrm{mix}}}{\partial \hat{\mathbf{\Phi}}}\,,$$

where the last term is present only when mixing is included. The *Gaiamiti field equation* is the on–shell condition

$$\boxed{\mathcal{E}_{\mathbf{\Phi}} = 0} \iff \boxed{-\nabla_\mu\Big(\Pi^\mu_{\mathbf{\Phi}}\Big) \;+\; \frac{\partial \mathcal{V}_\Phi}{\partial \hat{\mathbf{\Phi}}} \;+\; \frac{\partial \mathcal{V}_{\mathrm{mix}}}{\partial \hat{\mathbf{\Phi}}} \;=\; 0}\,. \tag{6.327}$$

Conventions and objects.

- ∇_μ is the covariant derivative induced by the background, $\bar{\Sigma}$–compatible connection on $T\mathcal{X}_q$ together with the chosen (representation) connection on $\mathcal{E}_\Phi$; it acts on $\hat{\mathbf{\Phi}}$ and its tensor products in the usual way (cf. Definition 6.184).
- $\bar{\Sigma}^{\mu\nu}$ raises indices and contracts the spacetime slot of $\nabla_\nu \hat{\mathbf{\Phi}}$ to form a current; it encodes the chronogeometric propagation weights (cf. eq. (6.62)).
- $\mathcal{G}_T$ converts a fiber vector (a variation of $\hat{\mathbf{\Phi}}$) into a fiber covector (a conjugate momentum) via g_T (cf. [253, 433]). An analogous Riesz map $\mathcal{G}_C$ exists on the Tantrakyma fiber (cf. Definition 6.142).
- $\frac{\partial \mathcal{V}_\Phi}{\partial \hat{\mathbf{\Phi}}}$ and $\frac{\partial \mathcal{V}_{\mathrm{mix}}}{\partial \hat{\mathbf{\Phi}}}$ are the fiberwise (Gâteaux) functional derivatives, valued in $\mathcal{E}_\Phi^\star$ (cf. Definition 6.174 and Section 6.12.2).
- $\Pi^\mu_{\mathbf{\Phi}}$ is an $\mathcal{E}_\Phi^\star$–valued spacetime vector (a momentum/flux current); $\nabla_\mu(\Pi^\mu_{\mathbf{\Phi}})$ is its covariant divergence.

All equalities are to be understood as (operator) identities on a common invariant domain; distributional interpretations are standard when needed (cf. [230]).

Intuitive reading. The equation eq. (6.327) says: the *covariant divergence* of the Gaiamiti momentum–flux current balances the local "restoring force" from the potential. The map $\mathcal{G}_T$ plays the role of a fiberwise Legendre transform, turning gradients of the geometric deformation $\hat{\mathbf{\Phi}}$ into their conjugate momenta; raising an index with $\bar{\Sigma}^{\mu\nu}$ turns those momenta into a spacetime flux. Setting $\mathcal{E}_{\mathbf{\Phi}} = 0$ thus enforces a conservation–of–flux condition modulated by the potential slope, and any coupling to $\hat{\mathbf{\Theta}}$ simply appears as an additional source. In short: geometric deformations propagate (via the background chronogeometry) so that momentum flux is covariantly conserved while the field "rolls" according to the potential it lives in.

Tantrakyma Euler–Lagrange operator and field equation.

Definition 6.268 (Tantrakyma Euler–Lagrange operator and field equation)**.** Let $(\mathcal{X}_q, \bar{\Sigma}_{\mu\nu})$ be the chronogeometric background with inverse $\bar{\Sigma}^{\mu\nu}$ and a torsion–free, $\bar{\Sigma}$–compatible connection ∇ (so $\nabla_\rho \bar{\Sigma}_{\mu\nu} = 0$; cf. Definition 6.184 and Section 6.5.3 and [375]). The *Tantrakyma field* is an operator–valued section $\hat{\mathbf{\Theta}} : \mathcal{X}_q \to \mathcal{E}_\Theta$, where $\mathcal{E}_\Theta \to \mathcal{X}_q$ is the Tantrakyma configuration bundle (cf. Definitions 6.3 and 6.142). On each fiber $(\mathcal{E}_\Theta)_x$ fix a positive, nondegenerate bilinear pairing g_C (the *kinetic/metric pairing*). Let $\mathcal{G}_C : (\mathcal{E}_\Theta)_x \to (\mathcal{E}_\Theta^\star)_x$ denote the associated Riesz/Legendre isomorphism, characterized by

$$\langle A, \mathcal{G}_C(B) \rangle \;=\; g_C(A, B) \qquad \text{for all } A, B \in (\mathcal{E}_\Theta)_x,$$

with $\langle \cdot, \cdot \rangle$ the canonical fiberwise duality pairing (cf. [253, 433]). Write $\mathcal{V}_\Theta(\hat{\mathbf{\Theta}})$ for the (fiberwise) potential density of $\hat{\mathbf{\Theta}}$, and include an optional mixing potential $\mathcal{V}_{\mathrm{mix}}(\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}})$ with the Gaiamiti field $\hat{\mathbf{\Phi}}$ when present (cf. Section 6.12.2; see also [82, 155]).

The *canonical momentum current* of $\hat{\mathbf{\Theta}}$ is the $\mathcal{E}_\Theta^\star$–valued spacetime vector field

$$\Pi^\mu_{\mathbf{\Theta}} \;:=\; \bar{\Sigma}^{\mu\nu}\, \mathcal{G}_C\Big(\nabla_\nu \hat{\mathbf{\Theta}}\Big).$$

The *Tantrakyma Euler–Lagrange operator* is the $\mathcal{E}_\Theta^\star$–valued section

$$\mathcal{E}_{\mathbf{\Theta}} \;:=\; -\nabla_\mu\Big(\Pi^\mu_{\mathbf{\Theta}}\Big) \;+\; \frac{\partial \mathcal{V}_\Theta}{\partial \hat{\mathbf{\Theta}}} \;+\; \frac{\partial \mathcal{V}_{\mathrm{mix}}}{\partial \hat{\mathbf{\Theta}}}\,,$$

with the last term included only when mixing is present. The *Tantrakyma field equation* is the on–shell condition

$$\boxed{\mathcal{E}_{\mathbf{\Theta}} = 0} \;\Longleftrightarrow\; \boxed{-\nabla_\mu\Big(\Pi^\mu_{\mathbf{\Theta}}\Big) \;+\; \frac{\partial \mathcal{V}_\Theta}{\partial \hat{\mathbf{\Theta}}} \;+\; \frac{\partial \mathcal{V}_{\mathrm{mix}}}{\partial \hat{\mathbf{\Theta}}} \;=\; 0}\,. \tag{6.328}$$

Conventions and objects.

- ∇_μ is the covariant derivative induced by the $\bar{\Sigma}$–compatible connection on $T\mathcal{X}_q$ together with the chosen representation connection on $\mathcal{E}_\Theta$; it acts on $\hat{\mathbf{\Theta}}$ and its tensor products in the standard way (cf. Definition 6.184).
- $\bar{\Sigma}^{\mu\nu}$ raises spacetime indices and contracts $\nabla_\nu \hat{\mathbf{\Theta}}$ to produce a current; it encodes chronogeometric propagation weights (cf. eq. (6.62)).

- $\mathcal{G}_C$ is the fiberwise Riesz/Legendre map associated with g_C (cf. [253, 433]); the Gaiamiti analogue is $\mathcal{G}_T$ (see Definition 6.267).
- $\frac{\partial \mathcal{V}_\Theta}{\partial \hat{\mathbf{\Theta}}}$ and $\frac{\partial \mathcal{V}_{\mathrm{mix}}}{\partial \hat{\mathbf{\Theta}}}$ are fiberwise (Gâteaux) functional derivatives valued in $\mathcal{E}_\Theta^\star$ (cf. Section 6.12.2).
- $\Pi^\mu_{\mathbf{\Theta}}$ is an $\mathcal{E}_\Theta^\star$–valued spacetime vector; $\nabla_\mu(\Pi^\mu_{\mathbf{\Theta}})$ is its covariant divergence.

All equalities are operator identities on a common invariant domain; distributional interpretations are standard when required (cf. [230]).

Intuitive reading. As in the Gaiamiti case *mutatis mutandis* (cf. Definition 6.267), eq. (6.328) reads as a flux–balance law in the Tantrakyma fiber: gradients of the matter-like excitation $\hat{\mathbf{\Theta}}$ are converted by $\mathcal{G}_C$ into conjugate momenta, and raising with $\bar{\Sigma}^{\mu\nu}$ promotes these to the spacetime flux $\Pi^\mu_{\mathbf{\Theta}}$. The field "rolls'' in its potential so that the covariant divergence of this flux equals the local restoring force (plus any mixing force from $\mathcal{V}_{\mathrm{mix}}$). Physically, Tantrakyma excitations propagate through the chronogeometry with momentum flux covariantly conserved up to potential and coupling terms.

Definition 6.269 (Ghost-number notation)**.** We denote the ghost-number grading by $\mathfrak{g}(\,\cdot\,) \in \mathbb{Z}$ and, when convenient, use the equivalent shorthand $\mathrm{gh}(\,\cdot\,)$; thus $\mathrm{gh}(X) = \mathfrak{g}(X)$. We also freely write statements like "$\mathrm{gh} = 0$" to mean $\mathfrak{g} = 0$.

1. **BRST degree.** The BRST differential $\zeta := \{\hat{Q}_{\mathrm{BRST}}, \cdot\}$ raises ghost number by $+1$:
$$\mathfrak{g}(\zeta X) = \mathfrak{g}(X) + 1 \quad (\text{equivalently, } \mathrm{gh}(\zeta X) = \mathrm{gh}(X) + 1).$$
In particular, $\mathfrak{g}(\eta_a) = -1$ and $\mathfrak{g}(\zeta\eta_a) = 0$ (cf. eq. (6.378)), and for the Nielsen generator η_ξ we have $\mathfrak{g}(\eta_\xi) = -1$ (cf. eq. (6.294)).
2. **Basic assignments.** Gauge fields and sources carry ghost number 0; Faddeev–Popov ghosts have $\mathfrak{g}(c) = +1$, anti-ghosts $\mathfrak{g}(\bar{c}) = -1$, and Nakanishi–Lautrup fields $\mathfrak{g}(B) = 0$. Scale tags and projectors satisfy
$$\mathfrak{g}\Big(P_x(\lambda_{\mathrm{UV}})\Big) = \mathfrak{g}\Big(K^{(\ell)}_{\mathrm{IR}}\Big) = \mathfrak{g}(\Pi_\alpha) = 0.$$
3. **Products and grading.** Ghost number is additive on products:
$$\mathfrak{g}(XY) = \mathfrak{g}(X) + \mathfrak{g}(Y) \quad (\text{equivalently, } \mathrm{gh}(XY) = \mathrm{gh}(X) + \mathrm{gh}(Y)),$$
and graded commutators/anticommutators use the standard sign $(-1)^{\mathfrak{g}(X)\mathfrak{g}(Y)}$.
4. **Physical sector.** The physical space is the degree-zero BRST cohomology,
$$\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}}) = \frac{\ker \zeta \cap \{\,\mathfrak{g} = 0\,\}}{\mathrm{im}\,\zeta \cap \{\,\mathfrak{g} = 0\,\}},$$
and all insertions used for expectation values on $\mathcal{H}_{\mathrm{phys}}$ have total ghost number 0 (i.e., $\mathrm{gh} = 0$).

When a label is present (e.g. an a for a global generator or channel), we may write $\mathfrak{g}_a$ (or gh_a) to emphasize the degree of the a-labeled piece.

BRST overlay (cohomological Euler–Lagrange operators).

Definition 6.270 (BRST overlay: cohomological Euler–Lagrange equations)**.** Let $\hat{Q}_{\mathrm{BRST}}$ be the nilpotent BRST charge ($\hat{Q}_{\mathrm{BRST}}^2 = 0$) and let $\zeta(\,\cdot\,) := \{\hat{Q}_{\mathrm{BRST}}, \cdot\}$ denote the induced graded derivation that raises ghost number by $+1$. Choose a gauge–fixing fermion Ψ (ghost number -1); the gauge–fixing + ghost sector is added as the ζ–exact term $\zeta\Psi$. Define the local insertions

$$\zeta\,\chi_{\mathbf{\Phi}} := \frac{\delta(\zeta\Psi)}{\delta\hat{\mathbf{\Phi}}}, \qquad \zeta\,\chi_{\mathbf{\Theta}} := \frac{\delta(\zeta\Psi)}{\delta\hat{\mathbf{\Theta}}}.$$

Let $\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}})$ be the physical Hilbert space (ghost number 0) and $\mathsf{P}_{\mathrm{phys}}$ the orthogonal projector onto it.

The *cohomological Euler–Lagrange operators* are

$$\mathcal{E}^{\mathrm{coh}}_{\mathbf{\Phi}} := \mathsf{P}_{\mathrm{phys}}\big(\mathcal{E}_{\mathbf{\Phi}} + \zeta\,\chi_{\mathbf{\Phi}}\big)\,\mathsf{P}_{\mathrm{phys}}, \qquad \mathcal{E}^{\mathrm{coh}}_{\mathbf{\Theta}} := \mathsf{P}_{\mathrm{phys}}\big(\mathcal{E}_{\mathbf{\Theta}} + \zeta\,\chi_{\mathbf{\Theta}}\big)\,\mathsf{P}_{\mathrm{phys}}.$$

The physical equations of motion are

$$\mathcal{E}^{\mathrm{coh}}_{\mathbf{\Phi}} = 0, \qquad \mathcal{E}^{\mathrm{coh}}_{\mathbf{\Theta}} = 0. \tag{6.329}$$

Because ζ–exact insertions vanish in BRST cohomology, these are equivalent on $\mathcal{H}_{\mathrm{phys}}$ to the base equations (6.327)–(6.328). Moreover, gauge–parameter deformations of Ψ shift $\chi_\bullet$ by ζ–exact terms and do not alter (6.329) (Nielsen–type independence).

Intuitive reading. BRST simply "modulates" the raw Euler–Lagrange equations by adding gauge–exact pieces that organize gauge fixing and ghosts; after projecting to the physical cohomology, those ζ–exact pieces drop out. Thus the physics is unchanged by gauge–parameter choices: the on–shell content is precisely the BRST–invariant part of the original equations for $\hat{\mathbf{\Phi}}$ and $\hat{\mathbf{\Theta}}$.

Linearised equations and background split.

Definition 6.271 (Linearised equations and background split)**.** Let the operator-valued fields be split into background (vacuum) plus fluctuations

$$\hat{\mathbf{\Phi}} = \bar{\Phi}\,\mathbb{I} + \delta\hat{\mathbf{\Phi}}, \qquad \hat{\mathbf{\Theta}} = \bar{\Theta}\,\mathbb{I} + \delta\hat{\mathbf{\Theta}} \quad \text{(cf. Definitions 6.142 and 11.17)}.$$

The backgrounds $\bar{\Phi}, \bar{\Theta}$ are c-number vacuum configurations solving the Euler–Lagrange equations

$$\mathcal{E}_{\mathbf{\Phi}}[\bar{\Phi}] = 0, \qquad \mathcal{E}_{\mathbf{\Theta}}[\bar{\Theta}] = 0 \quad \text{(see eqs. (6.327) and (6.328))}.$$

Linearised operators. Define the kinetic (second-order) operators acting on fluctuations by

$$\mathbb{K}_{\Phi} := -\nabla_\mu\big(\bar{\Sigma}^{\mu\nu}\,\mathcal{G}_T\,\nabla_\nu\,\cdot\,\big), \qquad \mathbb{K}_{\Theta} := -\nabla_\mu\big(\bar{\Sigma}^{\mu\nu}\,\mathcal{G}_C\,\nabla_\nu\,\cdot\,\big),$$

where ∇ is the torsion-free, $\bar{\Sigma}$–compatible connection on $T\mathcal{X}_q$ together with the chosen representation connections on $\mathcal{E}_\Phi, \mathcal{E}_\Theta$; $\bar{\Sigma}^{\mu\nu}$ is the inverse chronogeometric tensor; and $\mathcal{G}_T, \mathcal{G}_C$

are the fiberwise Riesz/Legendre maps associated to the positive bilinear pairings g_T, g_C on the respective fibers.

Hessian (mass/mixing) blocks. Let the fiberwise second derivatives at the vacuum be

$$\mathbb{M}_{\Phi\Phi} := \left.\frac{\partial^2 \mathcal{V}_\Phi}{\partial\hat{\mathbf{\Phi}}\,\partial\hat{\mathbf{\Phi}}}\right|_{\bar{\Phi}}, \quad \mathbb{M}_{\Theta\Theta} := \left.\frac{\partial^2 \mathcal{V}_\Theta}{\partial\hat{\mathbf{\Theta}}\,\partial\hat{\mathbf{\Theta}}}\right|_{\bar{\Theta}}, \quad \mathbb{M}_{\Phi\Theta} := \left.\frac{\partial^2 \mathcal{V}_{\mathrm{mix}}}{\partial\hat{\mathbf{\Phi}}\,\partial\hat{\mathbf{\Theta}}}\right|_{\bar{\Phi},\bar{\Theta}},$$

with $\mathbb{M}_{\Theta\Phi}$ the fiberwise adjoint (or transpose) as appropriate. These are linear maps from fluctuations to dual fluctuations, valued in $\mathcal{E}^\star_\Phi$ or $\mathcal{E}^\star_\Theta$.

Linearised system. Collect the fluctuations into $\delta\hat{\mathbf{U}} := (\delta\hat{\mathbf{\Phi}},\, \delta\hat{\mathbf{\Theta}})^{\mathsf{T}}$. The first variation of the Euler–Lagrange operators about $(\bar{\Phi}, \bar{\Theta})$ yields

$$\underbrace{\begin{pmatrix} \mathbb{K}_\Phi + \mathbb{M}_{\Phi\Phi} & \mathbb{M}_{\Phi\Theta} \\ \mathbb{M}_{\Theta\Phi} & \mathbb{K}_\Theta + \mathbb{M}_{\Theta\Theta} \end{pmatrix}}_{\mathbb{H}(\bar{\Phi},\bar{\Theta})} \begin{pmatrix} \delta\hat{\mathbf{\Phi}} \\ \delta\hat{\mathbf{\Theta}} \end{pmatrix} \;=\; 0, \tag{6.330}$$

where $\mathbb{H}(\bar{\Phi}, \bar{\Theta})$ is the (quadratic) fluctuation Hessian/operator used in Section 6.12.5. Equivalently, in block form,

$$\delta\mathcal{E}_{\mathbf{\Phi}} \;=\; (\mathbb{K}_\Phi + \mathbb{M}_{\Phi\Phi})\,\delta\hat{\mathbf{\Phi}} \;+\; \mathbb{M}_{\Phi\Theta}\,\delta\hat{\mathbf{\Theta}}, \qquad \delta\mathcal{E}_{\mathbf{\Theta}} \;=\; \mathbb{M}_{\Theta\Phi}\,\delta\hat{\mathbf{\Phi}} \;+\; (\mathbb{K}_\Theta + \mathbb{M}_{\Theta\Theta})\,\delta\hat{\mathbf{\Theta}}. \tag{6.331}$$

BRST note (projection to the physical sector). With gauge fixing and ghosts added as a ζ–exact term $\zeta\Psi$, longitudinal and ghost directions form ζ–quartets and decouple in cohomology. Denoting by $\mathsf{P}_{\mathrm{phys}}$ the projector onto $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$, the physically relevant quadratic operator is

$$\mathsf{P}_{\mathrm{phys}}\,\mathbb{H}(\bar{\Phi},\bar{\Theta})\,\mathsf{P}_{\mathrm{phys}}, \tag{6.332}$$

which governs the spectrum and normal modes on the physical sector; gauge-parameter deformations affect only ζ–exact (unphysical) blocks.

Intuitive reading. Linearising is "zooming in'' on small ripples around a vacuum that already solves the full equations. The kinetic pieces $\mathbb{K}_\Phi, \mathbb{K}_\Theta$ tell us how those ripples propagate through the background chronogeometry (via $\bar{\Sigma}$ and the covariant derivative), while the Hessian blocks $\mathbb{M}_{\bullet\bullet}$ act like local springs and couplings that set masses and mixings. Putting them together, the block matrix $\mathbb{H}$ is the small-oscillation operator: its eigenmodes are the normal excitations, and its eigenvalues encode the (squared) masses. Projecting with $\mathsf{P}_{\mathrm{phys}}$ simply discards pure-gauge and ghost ripples, ensuring that what remains are the genuinely observable vibrations of the system.

Scale-tagged form (projected and windowed).

Definition 6.272 (Scale–tagged Euler–Lagrange equations (projected & windowed))**.** Let $P_x(\lambda_{\mathrm{UV}})$ be the local UV spectral projector and $K^{(\ell)}_{\mathrm{IR}}$ the IR window kernel (cf. Section 6.12.4). Write

$$\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}}} := P_x(\lambda_{\mathrm{UV}})\,\hat{\mathbf{\Phi}}\,P_x(\lambda_{\mathrm{UV}}), \qquad \hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}}} := P_x(\lambda_{\mathrm{UV}})\,\hat{\mathbf{\Theta}}\,P_x(\lambda_{\mathrm{UV}}),$$

and denote by $\bar{\Sigma}^{(\ell)\mu\nu}$ the IR–windowed inverse chronogeometric tensor obtained from $\bar{\Sigma}^{\mu\nu}$ via $K_{\mathrm{IR}}^{(\ell)}$ (cf. Section 6.12.4). Let $\mathcal{G}_T, \mathcal{G}_C$ be the fiberwise Riesz/Legendre maps associated to the positive pairings g_T, g_C introduced in Definitions 6.267 and 6.268. Using the base Euler–Lagrange structures eqs. (6.327) and (6.328), the *scale–tagged* equations are

$$\mathcal{E}_{\mathbf{\Phi}}^{(\lambda_{\mathrm{UV}},\ell)} := P_x(\lambda_{\mathrm{UV}})\left[-\nabla_\mu\Big(\bar{\Sigma}^{(\ell)\mu\nu}\,\mathcal{G}_T\,\nabla_\nu\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}}}\Big) + \frac{\partial\mathcal{V}_\Phi}{\partial\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}}}}\right]P_x(\lambda_{\mathrm{UV}}) = 0, \tag{6.333}$$

$$\mathcal{E}_{\mathbf{\Theta}}^{(\lambda_{\mathrm{UV}},\ell)} := P_x(\lambda_{\mathrm{UV}})\left[-\nabla_\mu\Big(\bar{\Sigma}^{(\ell)\mu\nu}\,\mathcal{G}_C\,\nabla_\nu\hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}}}\Big) + \frac{\partial\mathcal{V}_\Theta}{\partial\hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}}}}\right]P_x(\lambda_{\mathrm{UV}}) = 0. \tag{6.334}$$

(When present, terms from $\mathcal{V}_{\mathrm{mix}}(\hat{\mathbf{\Phi}},\hat{\mathbf{\Theta}})$ are included by adding $\partial\mathcal{V}_{\mathrm{mix}}/\partial\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}}}$ in eq. (6.333) and $\partial\mathcal{V}_{\mathrm{mix}}/\partial\hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}}}$ in eq. (6.334), cf. Definitions 6.267 and 6.268.)

BRST–closed scale tagging and the physical sector. Assume $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ are BRST–closed (ghost number 0). Let $\mathsf{P}_{\mathrm{phys}}$ project onto $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ (cf. Definition 6.270). Then the physical, scale–tagged equations read

$$\mathsf{P}_{\mathrm{phys}}\,\mathcal{E}_{\mathbf{\Phi}}^{(\lambda_{\mathrm{UV}},\ell)}\,\mathsf{P}_{\mathrm{phys}} = 0, \qquad \mathsf{P}_{\mathrm{phys}}\,\mathcal{E}_{\mathbf{\Theta}}^{(\lambda_{\mathrm{UV}},\ell)}\,\mathsf{P}_{\mathrm{phys}} = 0,$$

and are invariant under gauge–parameter deformations of the gauge–fixing fermion (changes are ζ–exact and vanish in cohomology; cf. Definition 6.270 and [82, 155]). Spectral projectors and windowed operators are standard in the functional/spectral setting [230, 253].

Intuitive reading. Equations eqs. (6.333) and (6.334) are the band–limited versions of the base dynamics: $P_x(\lambda_{\mathrm{UV}})$ keeps only UV modes below the chosen cutoff at x, while $K_{\mathrm{IR}}^{(\ell)}$ smooths/coarse–grains the background weights into $\bar{\Sigma}^{(\ell)\mu\nu}$ over a length scale ℓ. Within this $(\lambda_{\mathrm{UV}},\ell)$–window, the same flux–balance holds—covariant divergence of the momentum–flux equals the local potential slope—but now for the projected fields and the IR–windowed chronogeometry. Projecting to $\mathcal{H}_{\mathrm{phys}}$ simply discards pure–gauge/ghost content, so the resulting equations capture the dynamics of observable modes at the selected scales.

Relational Ward identity (with explicit breakings)

Lemma 6.273 (Relational Ward identity with BRST improvement)**.** *Let the Lagrangian density be decomposed as*

$$\mathcal{L} \;=\; \mathcal{L}_0 \;+\; \sum_k \epsilon_k\,\mathcal{O}_k \;+\; \zeta\Psi,$$

where $\zeta(\,\cdot\,) := \{\hat{Q}_{\mathrm{BRST}}, \cdot\}$ is the BRST graded derivation (ghost number $+1$) induced by the nilpotent charge $\hat{Q}_{\mathrm{BRST}}$ (cf. Definition 6.270; see also [82, 230]). Suppose a (possibly local/relational) infinitesimal variation δ_a obeys

$$\delta_a\mathcal{L}_0 \;=\; \nabla_\mu K_a^\mu, \qquad \delta_a\mathcal{O}_k \;=\; \mathcal{S}_{k,a},$$

for some local current K_a^μ and local functionals $\mathcal{S}_{k,a}$. Let $\hat{J}^\mu_{a,\mathrm{imp}}$ denote the BRST–improved Noether current associated to δ_a (so that improvement terms absorb $\nabla_\mu K_a^\mu$ and gauge–fixing

artifacts; cf. [82]). Then there exists a local functional η_a generated by $\delta_a\Psi$ such that the following local identity holds off shell:

$$\nabla_\mu \hat{J}^\mu_{a,\mathrm{imp}} \;=\; \sum_k \epsilon_k\, \mathcal{S}_{k,a} \;+\; \zeta\, \eta_a. \tag{6.335}$$

Projecting to the physical sector $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ with $\mathsf{P}_{\mathrm{phys}}$ (cf. Definition 6.270) and writing $\bar{J}^\mu_{a,\mathrm{imp}} := \mathsf{P}_{\mathrm{phys}} \hat{J}^\mu_{a,\mathrm{imp}} \mathsf{P}_{\mathrm{phys}}$, one obtains

$$\nabla_\mu \bar{J}^\mu_{a,\mathrm{imp}} \;=\; \sum_k \epsilon_k \left\langle \mathcal{S}_{k,a} \right\rangle_{\boldsymbol{\Psi}}, \tag{6.336}$$

since ζ–exact insertions vanish in BRST cohomology (Nielsen–type independence; cf. Definition 6.270 and [230]). Identity (6.336) is consistent with the scale–tagged version eq. (6.375). If the symmetry group G is gauged, (6.336) upgrades to the corresponding Slavnov–Taylor identity, assuming anomaly–free BRST at Stage I (Section 12.2.2) (cf. [82]).

Proof of Lemma 6.233. Work covariantly with the background measure $\mathrm{d}\mu := \sqrt{-\det \bar{\Sigma}}\, \mathrm{d}^d x$ so that ∇_μ integrates by parts without torsion terms (cf. Definitions 6.267 and 6.268 and [375]). For any infinitesimal (possibly local/relational) variation δ_a acting on the field content, the standard Noether identity gives

$$\delta_a \mathcal{L} \;=\; \left\langle \mathcal{E}_{\boldsymbol{\Phi}},\, \delta_a \hat{\boldsymbol{\Phi}} \right\rangle + \left\langle \mathcal{E}_{\boldsymbol{\Theta}},\, \delta_a \hat{\boldsymbol{\Theta}} \right\rangle \;+\; \nabla_\mu J^\mu_{a,\mathrm{can}}, \tag{6.337}$$

where $J^\mu_{a,\mathrm{can}}$ is the canonical Noether current and $\mathcal{E}_{\boldsymbol{\Phi}}, \mathcal{E}_{\boldsymbol{\Theta}}$ are the Euler–Lagrange operators from eqs. (6.327) and (6.328) (see, e.g., [82]; cf. Definitions 6.267 and 6.268). On the other hand, using the assumed decomposition of $\mathcal{L}$ and the hypotheses on δ_a,

$$\delta_a \mathcal{L} \;=\; \delta_a \mathcal{L}_0 + \sum_k \epsilon_k\, \delta_a \mathcal{O}_k + \delta_a(\zeta \Psi) \;=\; \nabla_\mu K^\mu_a \;+\; \sum_k \epsilon_k\, \mathcal{S}_{k,a} \;+\; \zeta(\delta_a \Psi), \tag{6.338}$$

where we used that $\zeta(\cdot) = \{\hat{Q}_{\mathrm{BRST}}, \cdot\}$ is a graded derivation (ghost number $+1$) and that boundary/superpotential terms from $\delta_a(\zeta\Psi)$ can be absorbed into a divergence (cf. Definition 6.270 and [230]).

Equating eqs. (6.337) and (6.338) and rearranging yields

$$\nabla_\mu \Big(J^\mu_{a,\mathrm{can}} - K^\mu_a \Big) \;=\; \sum_k \epsilon_k\, \mathcal{S}_{k,a} \;+\; \zeta(\delta_a \Psi) \;-\; \left\langle \mathcal{E}_{\boldsymbol{\Phi}},\, \delta_a \hat{\boldsymbol{\Phi}} \right\rangle \;-\; \left\langle \mathcal{E}_{\boldsymbol{\Theta}},\, \delta_a \hat{\boldsymbol{\Theta}} \right\rangle. \tag{6.339}$$

By the algebraic Poincaré lemma (local cohomology of the divergence) and standard Noether–Belinfante improvement, there exists a local functional U^μ_a (built polynomially from fields and a finite number of derivatives) such that [117, 91, 82, 90, 435, 409]

$$\nabla_\mu U^\mu_a \;=\; -\left\langle \mathcal{E}_{\boldsymbol{\Phi}},\, \delta_a \hat{\boldsymbol{\Phi}} \right\rangle - \left\langle \mathcal{E}_{\boldsymbol{\Theta}},\, \delta_a \hat{\boldsymbol{\Theta}} \right\rangle \;+\; \zeta\, \Delta_a, \tag{6.340}$$

for some local Δ_a (the ζ–exact remainder organizes possible gauge–fixing artifacts). Define the *BRST–improved* current

$$\hat{J}^\mu_{a,\mathrm{imp}} \;:=\; J^\mu_{a,\mathrm{can}} - K^\mu_a - U^\mu_a.$$

Combining eqs. (6.339) and (6.340) gives the off–shell local identity

$$\nabla_\mu \hat{J}^\mu_{a,\mathrm{imp}} \;=\; \sum_k \epsilon_k\, \mathcal{S}_{k,a} \;+\; \zeta\big(\delta_a \Psi - \Delta_a\big). \tag{6.341}$$

Setting $\eta_a := \delta_a\Psi - \Delta_a$ establishes eq. (6.335).

For the physical statement, project to $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ with $\mathsf{P}_{\mathrm{phys}}$ (cf. Definition 6.270). Because ζ–exact insertions vanish in BRST cohomology, $\mathsf{P}_{\mathrm{phys}}\zeta\,\eta_a\,\mathsf{P}_{\mathrm{phys}} = 0$, and taking matrix elements (or equivalently, band/BRST–consistent expectation values $\langle\cdot\rangle_{\boldsymbol{\Psi}}$) gives

$$\nabla_\mu \bar{J}^\mu_{a,\mathrm{imp}} \;=\; \sum_k \epsilon_k \left\langle \mathcal{S}_{k,a} \right\rangle_{\boldsymbol{\Psi}},$$

which is eq. (6.336). When G is gauged, the same derivation—now applied to gauge variations—yields the Slavnov–Taylor form of the identity under the assumption of anomaly–free BRST (Stage I), as standard in covariant quantization [82]; see also the local field–algebra treatment in [230]. This matches the scale–tagged version eq. (6.375) by applying the UV projector and IR window (cf. Definition 6.272) and using that the projectors/windows are BRST–closed, so they commute with the cohomological reduction.

This completes the proof. □

Intuitive reading. Start from a symmetry of the reference theory $\mathcal{L}_0$ that is only violated by controlled "breaking operators" $\mathcal{O}_k$ and by gauge–fixing/ghost bookkeeping ($\zeta\Psi$). The improved current $\hat{J}^\mu_{a,\mathrm{imp}}$ is constructed so the variation of $\mathcal{L}_0$ is a total divergence; any nonzero divergence of the current then cleanly tracks (i) explicit symmetry–breaking sources $\epsilon_k\mathcal{S}_{k,a}$ and (ii) a BRST–exact piece from gauge fixing. After projecting to the physical cohomology (ghost number 0), the BRST–exact term drops out, leaving a Ward identity that says: the only obstruction to exact conservation is the explicit breaking encoded by the ϵ_k's. When $\epsilon_k = 0$ one has exact conservation (Quadrants I–II); if $\epsilon_k(\mu) \to 0$ toward the IR the conservation is approximate (Quadrant III); and if some ϵ_k persist at low scales, the non–conservation survives (Quadrant IV). In gauge theories the same logic yields the familiar Slavnov–Taylor relations.

Linear response and propagators

Definition 6.274 (Linear response kernels and propagators)**.** Let the fields be split into vacuum plus fluctuations as in Definition 6.271, and let the quadratic (Hessian) operator about $(\bar\Phi, \bar\Theta)$ be $\mathbb{H}(\bar\Phi, \bar\Theta)$ from eqs. (6.330) and (6.331). Define the fluctuation vector $\delta\hat{\mathbf{U}} := (\delta\hat{\boldsymbol{\Phi}},\, \delta\hat{\boldsymbol{\Theta}})^{\mathsf{T}}$ and the bilinear form

$$S^{(2)} \;=\; \tfrac{1}{2}\left\langle \delta\hat{\mathbf{U}},\, \mathcal{K}\,\delta\hat{\mathbf{U}} \right\rangle, \qquad \mathcal{K} \;:=\; \mathbb{H}(\bar\Phi, \bar\Theta) \;=\; \begin{pmatrix} \mathbb{K}_\Phi + \mathbb{M}_{\Phi\Phi} & \mathbb{M}_{\Phi\Theta} \\ \mathbb{M}_{\Theta\Phi} & \mathbb{K}_\Theta + \mathbb{M}_{\Theta\Theta} \end{pmatrix},$$

with blocks as in eq. (6.331). A *propagator* is any bi-distribution (kernel) $\mathcal{D}$ on the fluctuation bundle such that it inverts $\mathcal{K}$ with the chosen boundary prescription (retarded/advanced/Feynman), i.e.

$$\mathcal{K} \cdot \mathcal{D} \;=\; \mathbf{1}, \tag{6.342}$$

where "$\cdot$" denotes operator composition (contraction of fiber indices and integration over spacetime) and $\mathbf{1}$ is the identity on the fluctuation bundle (cf. [230, 82, 91]). Under standard hyperbolicity/ellipticity and spectral assumptions, the various causal/Feynman inverses exist as tempered distributions [253].

BRST overlay (physical propagator). Introduce gauge fixing via a BRST–exact gauge–fermion Ψ and form the gauge–fixed quadratic action $S_\Psi^{(2)} = \frac{1}{2}\langle \delta\hat{\mathbf{U}},\, \mathcal{K}_\Psi\, \delta\hat{\mathbf{U}}\rangle$ (cf. Definition 6.270). Let $\mathsf{P}_{\mathrm{phys}}$ project onto $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$. Longitudinal and ghost directions assemble into ζ–quartets and decouple in cohomology, and the *physical* kernel and propagator satisfy

$$\mathcal{K}_{\mathrm{phys}} \cdot \mathcal{D}_{\mathrm{phys}} = \mathbf{1} \quad \text{on } \mathcal{H}_{\mathrm{phys}},$$

with pole locations and residues independent of the gauge parameter by Nielsen–type identities (cohomological Ward identities) [82, 117]; see also Proposition 6.264.

Intuitive reading. Around a vacuum, the dynamics for small ripples is linear: $\mathcal{K}$ acts like a "stiffness" operator, and the propagator $\mathcal{D}$ is its inverse—telling us how a local source produces a response elsewhere. Choosing retarded/advanced/Feynman just fixes causal/boundary behavior. BRST gauge fixing only reshuffles unphysical directions: after projecting to $\mathcal{H}_{\mathrm{phys}}$ the response is unique, and its poles (masses, residues) do not depend on the gauge parameter. Channel–by–channel structure (Gaiamiti/Tantrakyma and their mixing) and T–even/T–odd tomography then follow from the holosymmetric interferometric split into anticommutator vs. commutator quadratures (cf. Chapter 6).

6.12.3 Operational Diagnostics (Scale-aware, Channel-resolved)

Holosymmetric visibilities

Definition 6.275 (Holosymmetric visibilities and scale tagging)**.** Let U_A, U_B be the two law-branch evolution operators acting on the system Hilbert space, and let ρ_0 be a reference state (density operator). Define the *law-channel visibility*

$$\Lambda_{\mathrm{law}} \;:=\; \mathrm{Tr}\Big[U_B^\dagger U_A\, \rho_0\Big], \tag{6.343}$$

and for two (pure) output states $|\Omega_1\rangle, |\Omega_2\rangle$ define the *state-channel visibility*

$$\Lambda_{\mathrm{state}} \;:=\; \langle \Omega_2 | \Omega_1 \rangle\ . \tag{6.344}$$

(For mixed-state generalizations, use the natural bilinear pairing induced by the chosen ordering/contour; cf. Definition 6.274.)

Scale-aware tagging. With the UV spectral projector $P_x(\lambda_{\mathrm{UV}})$ and IR window kernel $K_{\mathrm{IR}}^{(\ell)}$ (cf. Section 6.12.4 and Definition 6.272), write the scale–tagged operators and states as

$$U_{A/B}^{(\lambda_{\mathrm{UV}},\ell)} := K_{\mathrm{IR}}^{(\ell)}\, P_x(\lambda_{\mathrm{UV}})\, U_{A/B}\, P_x(\lambda_{\mathrm{UV}})\, K_{\mathrm{IR}}^{(\ell)}, \qquad \rho_0^{(\lambda_{\mathrm{UV}},\ell)} := P_x(\lambda_{\mathrm{UV}})\, \rho_0\, P_x(\lambda_{\mathrm{UV}}),$$

and analogously for $|\Omega_i^{(\lambda_{\mathrm{UV}},\ell)}\rangle$. The *scale–tagged* visibilities are

$$\Lambda_{\mathrm{law}}^{(\lambda_{\mathrm{UV}},\ell)} := \mathrm{Tr}\Big[(U_B^{(\lambda_{\mathrm{UV}},\ell)})^\dagger U_A^{(\lambda_{\mathrm{UV}},\ell)}\, \rho_0^{(\lambda_{\mathrm{UV}},\ell)}\Big], \tag{6.345}$$

$$\Lambda_{\mathrm{state}}^{(\lambda_{\mathrm{UV}},\ell)} := \langle \Omega_2^{(\lambda_{\mathrm{UV}},\ell)} | \Omega_1^{(\lambda_{\mathrm{UV}},\ell)} \rangle\ . \tag{6.346}$$

Channel resolution (triadic split). Let $\{\Pi_\alpha\}_{\alpha\in\{\mathrm{corr,vac,spec}\}}$ be BRST–closed (ghost number 0) mutually orthogonal projectors that resolve the holosymmetric triad (cf. Chapter 6: $\hat g_{\mu\nu} = \hat g^{\mathrm{corr}}_{\mu\nu} + \hat g^{\mathrm{vac}}_{\mu\nu} + \hat g^{\mathrm{spec}}_{\mu\nu}$). Assume $\sum_\alpha \Pi_\alpha = \mathbf{1}$ and that Π_α commute with the scale tags $P_x(\lambda_{\mathrm{UV}})$ and $K^{(\ell)}_{\mathrm{IR}}$. Then the visibilities decompose additively by channel:

$$\Lambda^{(\lambda_{\mathrm{UV}},\ell)}_{\mathrm{law}} = \sum_{\alpha\in\{\mathrm{corr,vac,spec}\}} \mathrm{Tr}\Big[\Pi_\alpha\,(U_B^{(\lambda_{\mathrm{UV}},\ell)})^\dagger U_A^{(\lambda_{\mathrm{UV}},\ell)}\,\Pi_\alpha\,\rho_0^{(\lambda_{\mathrm{UV}},\ell)}\Big], \tag{6.347}$$

$$\Lambda^{(\lambda_{\mathrm{UV}},\ell)}_{\mathrm{state}} = \sum_\alpha \langle \Omega_2^{(\lambda_{\mathrm{UV}},\ell)} | \Pi_\alpha | \Omega_1^{(\lambda_{\mathrm{UV}},\ell)} \rangle\,. \tag{6.348}$$

Causal preservation (no law–signalling). Let $\mathcal{O}_{\mathrm{meas}}$ be any measurement localized in a region obeying the windowed cone constraints determined by $\bar\Sigma^{(\ell)}$ (cf. Definition 6.272). If the difference $U_B^\dagger U_A$ is supported outside the effective causal domain of $\mathcal{O}_{\mathrm{meas}}$, then $\Lambda^{(\lambda_{\mathrm{UV}},\ell)}_\bullet$ is unchanged by that difference (Einstein causality in the local field algebra sense; cf. [230]).

Proposition 6.276 (BRST and gauge–parameter invariance of visibilities)**.** *To evaluate* (6.345)–(6.346)*, we project onto the physical sector,*

$$\mathcal{H}_{\mathrm{phys}} = H^0(\hat Q_{\mathrm{BRST}}),$$

using the projector $\mathsf{P}_{\mathrm{phys}}$ *and a set of BRST-closed projectors* Π_α*. Then* $\Lambda^{(\lambda_{\mathrm{UV}},\ell)}_\bullet$ *and their channel–resolved versions are invariant under* ζ*–exact changes of the gauge–fixing fermion* Ψ *and under gauge–parameter scans:*

$$\frac{\partial}{\partial\xi}\,\Lambda^{(\lambda_{\mathrm{UV}},\ell)}_\bullet = 0, \qquad \frac{\partial}{\partial\alpha_{\mathrm{gf}}}\,\Lambda^{(\lambda_{\mathrm{UV}},\ell)}_\bullet = 0.$$

provided the Nielsen/cohomological identities hold on the band (cf. Definition 6.270 and Proposition 6.264; see [117, 82]).

Proof of Proposition 6.276. **Set–up and notation.** Work with the gauge–fixed action $S_{\mathrm{gf+gh}}(\xi,\alpha_{\mathrm{gf}}) = S_0 + \{\hat Q_{\mathrm{BRST}}, \Psi(\xi,\alpha_{\mathrm{gf}})\}$, where $\hat Q_{\mathrm{BRST}}$ is nilpotent and the path–integral measure is BRST invariant (Stage I). Denote the BRST differential by $\zeta(\cdot) = \{\hat Q_{\mathrm{BRST}}, \cdot\}$. Let the scale tags $K^{(\ell)}_{\mathrm{IR}}$, $P_x(\lambda_{\mathrm{UV}})$ and the channel projectors Π_α be BRST–closed, ghost number 0, and let $\mathsf{P}_{\mathrm{phys}}$ project onto $\mathcal{H}_{\mathrm{phys}} = H^0(\hat Q_{\mathrm{BRST}})$, with $[\hat Q_{\mathrm{BRST}}, \mathsf{P}_{\mathrm{phys}}] = 0$.

1) BRST closure of the visibility kernels. The total (scale–dependent) Hamiltonian that generates the evolutions entering the visibilities is of the form $H_{\mathrm{tot}}(\cdot) = H_0(\cdot) + \zeta\Psi(\cdot)$, with $[\,\hat Q_{\mathrm{BRST}}, H_0\,] = 0$ (gauge–invariant part). By nilpotency and the graded Jacobi identity,

$$[\,\hat Q_{\mathrm{BRST}}, H_{\mathrm{tot}}\,] = [\,\hat Q_{\mathrm{BRST}}, H_0\,] + [\,\hat Q_{\mathrm{BRST}}, \zeta\Psi\,] = 0.$$

Hence the evolution operators commute with $\hat Q_{\mathrm{BRST}}$: $[\,\hat Q_{\mathrm{BRST}}, U(\cdot)\,] = 0$, and so does any product built from them, in particular the interferometric kernels $U_B^\dagger(\ell)U_A(\ell)$ and the state–overlap kernels $U^\dagger(V)U(V)$. Since $K^{(\ell)}_{\mathrm{IR}}$, $P_x(\lambda_{\mathrm{UV}})$, Π_α and $\mathsf{P}_{\mathrm{phys}}$ also commute with $\hat Q_{\mathrm{BRST}}$, every *tagged* kernel used below is BRST–closed.

2) Pointwise Nielsen/cohomological identity. Let $\mathcal{O}$ be any BRST–closed, ghost–number 0 insertion constructed from the tagged kernels and projectors (this includes the total and channel–resolved visibilities). Differentiating with respect to either parameter gives

$$\partial_\xi S_{\mathrm{gf+gh}} = \zeta\,(\partial_\xi \Psi) =: \zeta\,\eta_\xi, \qquad \partial_{\alpha_{\mathrm{gf}}} S_{\mathrm{gf+gh}} = \zeta\,(\partial_{\alpha_{\mathrm{gf}}} \Psi) =: \zeta\,\eta_{\alpha_{\mathrm{gf}}},$$

with $\mathrm{gh}(\eta_\cdot) = -1$. The standard Nielsen step (cf. Proposition 6.264) then yields, in the physical sector,

$$\partial_\xi \langle \mathcal{O} \rangle_{\mathcal{H}_{\mathrm{phys}}} = \Big\langle \mathcal{O}\, \partial_\xi S_{\mathrm{gf+gh}} \Big\rangle_{\mathcal{H}_{\mathrm{phys}}} = \Big\langle \mathcal{O}\, \zeta\, \eta_\xi \Big\rangle_{\mathcal{H}_{\mathrm{phys}}} = \Big\langle \zeta(\eta_\xi \mathcal{O}) \Big\rangle_{\mathcal{H}_{\mathrm{phys}}} - \underbrace{\Big\langle (\zeta \mathcal{O})\, \eta_\xi \Big\rangle_{\mathcal{H}_{\mathrm{phys}}}}_{=0} \tag{6.349}$$

$$= \Big\langle \zeta(\eta_\xi \mathcal{O}) \Big\rangle_{\mathcal{H}_{\mathrm{phys}}} = 0, \tag{6.350}$$

and identically $\partial_{\alpha_{\mathrm{gf}}} \langle \mathcal{O} \rangle_{\mathcal{H}_{\mathrm{phys}}} = 0$, because the expectation of a BRST–exact, ghost–number 0 insertion vanishes when the measure is BRST invariant (anomaly–free Stage I).

3) Application to the visibilities. Write the scale–tagged visibilities (total and channel–resolved) schematically as

$$\Lambda_\bullet^{(\lambda_{\mathrm{UV}},\ell)} = \mathrm{Tr}\Big[\mathsf{P}_{\mathrm{phys}}\, K_{\mathrm{IR}}^{(\ell)} P_x(\lambda_{\mathrm{UV}})\, \mathcal{K}_\bullet\, P_x(\lambda_{\mathrm{UV}}) K_{\mathrm{IR}}^{(\ell)}\, \mathsf{P}_{\mathrm{phys}}\, \rho_0 \Big],$$

and $\Lambda_\bullet^{(\alpha)} = \mathrm{Tr}\left[\Pi_\alpha(\cdots)\Pi_\alpha\, \rho_0\right]$ with $\bullet \in \{\mathrm{law}, \mathrm{state}\}$ and $\mathcal{K}_\bullet$ the corresponding kernel ($U_B^\dagger U_A$ or $U^\dagger U$). By Step 1 the full insertion between the traces is BRST–closed; hence Step 2 applies *pointwise in* $(\lambda_{\mathrm{UV}}, \ell)$, giving

$$\frac{\partial}{\partial \xi} \Lambda_\bullet^{(\lambda_{\mathrm{UV}},\ell)} = 0, \qquad \frac{\partial}{\partial \alpha_{\mathrm{gf}}} \Lambda_\bullet^{(\lambda_{\mathrm{UV}},\ell)} = 0,$$

and the same equalities with $\Lambda_\bullet^{(\alpha)}$ in place of $\Lambda_\bullet$, for each $\alpha \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$.

4) ζ–exact changes of the gauge–fixing fermion. If $\Psi \mapsto \Psi + \delta\Psi$ with *arbitrary* smooth $\delta\Psi$, the corresponding change in the action is $\delta S_{\mathrm{gf+gh}} = \zeta(\delta\Psi)$. Repeating the argument in Step 2 with $\eta_\xi \to \delta\Psi$ shows $\delta\Lambda_\bullet^{(\lambda_{\mathrm{UV}},\ell)} = 0$ and $\delta\Lambda_\bullet^{(\alpha)} = 0$. Thus the visibilities are invariant under ζ–exact deformations of the gauge fixing.

Conclusion. Under the stated hypotheses (Nielsen/cohomological identities holding on the band), the scale–tagged visibilities—both total and channel–resolved—are invariant under gauge–parameter scans and under ζ–exact changes of Ψ, as claimed. □

Intuitive reading. Λ_{law} is the interference fringe visibility between two law–branch evolutions; Λ_{state} is the overlap between the resulting states. Dialing $(\lambda_{\mathrm{UV}}, \ell)$ simply filters which modes contribute, while the channel projectors tease apart correlation/vacuum/spectral contributions. Locality ensures that law changes outside the windowed causal cone cannot influence the measured visibilities, and BRST/Nielsen identities guarantee that gauge choices don't either—so the meters are both scale–aware and physically robust.

Asymptotic signatures by quadrant (cohomological reading).

Proposition 6.277 (Quadrant diagnostics via holosymmetric visibilities)**.** *Work on the physical sector $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ with BRST–closed scale tags $P_x(\lambda_{\mathrm{UV}})$, $K_{\mathrm{IR}}^{(\ell)}$ as in Definition 6.272 and visibilities as in Definition 6.275. Let d be the spatial dimension, z the dynamical exponent, V the volume, and let $\Delta := \min_k \Delta_k$ be the smallest scaling dimension among RG–irrelevant symmetry–breaking operators that appear with couplings ϵ_k in the Lagrangian (cf. Lemma 6.233; see [91, 101, 117]). Then the following asymptotic signatures hold, and are invariant under ζ–exact gauge–fermion deformations (by Proposition 6.276):*

QI (Law Sym., State Sym.)
$|\Lambda_{\mathrm{law}}| \approx 1$, $|\Lambda_{\mathrm{state}}| \approx 1$, *with Ward identities exact (cf. Section 6.12.2).*

QII (Law Sym., State Asym.)
$|\Lambda_{\mathrm{law}}| \approx 1$, $\Lambda_{\mathrm{state}}(V) \to 0 \quad (V \to \infty)$, *and Goldstone data are extractable from phase–twist response (cf. [91]).*

QIII (Law Asym., State Sym.)
$1 - |\Lambda_{\mathrm{law}}(\ell)| = \mathcal{O}\big(\ell^{\Delta-(d+z)}\big)$ *and* $|\Lambda_{\mathrm{state}}| \approx 1$.

QIV (Law Asym., State Asym.)
$|\Lambda_{\mathrm{law}}| < 1$ *(bounded away from 1 in the IR), unique vacuum; no exact Nambu–Goldstone modes.*

Full proof of Proposition 6.277. **Set–up and hypotheses.** Work throughout in the physical sector $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$. The scale tags $K_{\mathrm{IR}}^{(\ell)}$ and $P_x(\lambda_{\mathrm{UV}})$ and the optional channel projectors Π_α are BRST–closed, ghost number 0. The gauge–fixed action is $S_{\mathrm{gf+gh}} = S_0 + \{\hat{Q}_{\mathrm{BRST}}, \Psi\}$ with anomaly–free BRST. Holosymmetric visibilities (total or channel–resolved) are defined as in Definition 6.275. By Proposition 6.276 (Nielsen/cohomological identities), *all* statements below are gauge–parameter and gauge–fermion invariant, pointwise in $(\lambda_{\mathrm{UV}}, \ell)$, and thus on any plateau band.

Let $\{\mathcal{O}_k\}$ denote symmetry–violating operators of scaling dimensions Δ_k entering the micro–law with couplings ϵ_k (Relational Ward input, Lemma 6.233). Set $\Delta := \min_k \Delta_k$ among the *irrelevant* breakings (when present). We write the two law–paths as $U_A(\ell)$ and $U_B(\ell)$ and define $\delta H(\ell') := H_A(\ell') - H_B(\ell') = \sum_k \epsilon_k(\ell')\, \mathcal{O}_k(\ell')$, with $\epsilon_k(\ell')$ the running couplings in the scale variable ℓ'. A Duhamel/BCH expansion (interaction picture w.r.t. H_B) gives

$$U_B^\dagger(\ell) U_A(\ell) = \mathbb{I} - i\int_0^\ell d\ell'\, \widetilde{\delta H}(\ell') - \int_{0<\ell''<\ell'<\ell} d\ell' d\ell''\, \widetilde{\delta H}(\ell')\, \widetilde{\delta H}(\ell'') + \cdots . \tag{6.351}$$

On any plateau–stable band B (Remark 6.254) the tags are order–≤ 0, the block structure is functorial, operator norms are uniformly controlled, and cone–preserving (Proposition 6.257) so only intra–cone phases/subprincipal data are modified.

QI: Law Symmetric, State Symmetric. If the law is exactly G–symmetric, then all explicit symmetry–breaking couplings vanish: $\epsilon_k \equiv 0$ in (6.351). Hence the law kernel is the identity on $\mathcal{H}_{\mathrm{phys}}$ up to IR–irrelevant deformations (coming from non–breaking operators), and

$$|\Lambda_{\mathrm{law}}(\ell)| = 1 + o_B(1).$$

Predicate stability and cluster decomposition (unique G–invariant vacuum) imply $|\Lambda_{\text{state}}(V)| \to 1$ as $V \uparrow \infty$. The Relational Ward identity holds with vanishing RHS in $\mathcal{H}_{\text{phys}}$, yielding exact Ward/Slavnov–Taylor identities. This yields the QI line.

QII: Law Symmetric, State Asymmetric (SSB). Again $\epsilon_k \equiv 0$ so the law visibility stays $|\Lambda_{\text{law}}| \approx 1$ (plateau corrections only). Let $\{|\Omega_\alpha\rangle\}$ be thermodynamic–limit vacua related by G. For distinct $\alpha \neq \beta$, $\langle\Omega_\beta|\Omega_\alpha\rangle \xrightarrow[V\uparrow\infty]{} 0$ (disjoint GNS representations/orthogonality). Thus, for preparations selecting different branches,

$$\Lambda_{\text{state}}(V) = \langle\Omega_\beta|\Omega_\alpha\rangle \xrightarrow[V\uparrow\infty]{} 0.$$

Goldstone data for *global* breaking are extracted from phase–twist response in the correlational block; in gauged sectors would–be NG modes are ζ–exact and absent in $\mathcal{H}_{\text{phys}}$ (Kugo–Ojima quartet, Chapter 8). This yields the QII line.

QIII: Law Asymmetric (irrelevant), State Symmetric (Emergence). Assume explicit *irrelevant* symmetry–breaking operators $\mathcal{O}_k$ with $\Delta_k > d + z$. Along the RG toward the IR (momentum scale $\mu \downarrow 0$) one has $g_k(\mu) := \epsilon_k(\mu) = \mathcal{O}\big(\mu^{\Delta_k-(d+z)}\big)$. Equivalently, in the IR–length variable $\ell \sim 1/\mu$,

$$\epsilon_k(\ell') = \mathcal{O}\big(\ell'^{-(\Delta_k-(d+z))}\big) \qquad (\ell' \uparrow \infty).$$

Insert this in (6.351) and take tagged (or channel–tagged) expectations. Uniform boundedness/mixing on B yields

$$\begin{aligned} 1 - |\Lambda_{\text{law}}(\ell)| &\le c_1 \int_0^\ell d\ell' \sum_k |\epsilon_k(\ell')| + c_2 \int_{0<\ell''<\ell'<\ell} d\ell' d\ell'' \sum_{k,k'} |\epsilon_k(\ell')\epsilon_{k'}(\ell'')| + \cdots \\ &= \mathcal{O}\big(\ell^{-(\Delta-(d+z))}\big) \qquad (\ell \uparrow \infty), \end{aligned} \tag{6.352}$$

with $\Delta = \min_k \Delta_k$. Translating to the UV–probe convention used in the statement (via $\mu = 1/\ell$) gives

$$1 - |\Lambda_{\text{law}}(\ell)| = \mathcal{O}\big(\ell^{\,\Delta-(d+z)}\big) \qquad (\ell \downarrow 0),$$

as claimed in the QIII bullet. The state stays symmetric (unique vacuum), so $|\Lambda_{\text{state}}| \approx 1$. The Relational Ward identity in $\mathcal{H}_{\text{phys}}$ has RHS $\sum_k \epsilon_k \langle\mathcal{S}_{k,a}\rangle = \mathcal{O}(\mu^{\Delta-(d+z)})$, i.e. approximate conservation sharpening into the IR, consistent with (6.352).

QIV: Law Asymmetric (relevant/marginal), State Asymmetric (Explicit). Suppose an explicit breaking with $\Delta \le d + z$ persists in the IR. Then $g(\mu) = \epsilon(\mu)$ approaches a nonzero constant (marginal) or grows (relevant) as $\mu \downarrow 0$. The integrals in (6.351) do not wash out; by positivity/variance bounds for BRST–closed probes (Chapter 8) there exist $c > 0$ and ℓ_0 (uniform on any plateau band B) with

$$1 - |\Lambda_{\text{law}}(\ell)| \;\ge\; c \qquad \forall\, \ell \ge \ell_0,$$

so the law visibility is *bounded away from* 1 into the IR. Explicit breaking selects a unique vacuum; exact NG modes are absent (at most pseudo–NG with $m^2 \propto \epsilon$). This yields the QIV line.

Cohomological and causal robustness. By Proposition 6.276, the above asymptotics are invariant under ζ–exact deformations of the gauge fixing and under gauge–parameter scans. By Proposition 6.257, the holosymmetric lift is blockwise (functorial) and order–≤ 0; it preserves the $\bar{\Sigma}$–cone (principal symbol), hence cannot change kinematic causality or quadrant predicates—only diagnose them.

Conclusion. Putting the four cases together establishes the scale/volume asymptotic signatures stated in Proposition 6.277, with invariance under ζ–exact changes of Ψ and gauge–parameter scans guaranteed by BRST cohomology in $\mathcal{H}_{\text{phys}}$. □

Intuitive reading. Λ_{law} tests whether the evolution "laws" match between branches; Λ_{state} tests whether the resulting states coincide. QI has both intact (both visibilities ≈ 1). QII keeps the law intact but the states separate (overlaps die with volume). QIII has tiny, scale–suppressed law differences from irrelevant breakings (deficit scales with the IR window), while the state stays symmetric. QIV shows robust law differences and a unique, explicitly selected vacuum, so no exact NG modes survive.

6.12.4 Scale-tagged Variational Infrastructure (IR/UV)

We introduce two operational **knobs**. They implement, in concrete form, the reciprocity between UV control (finite entanglement + spectral smoothness) and IR control (finite windows + clustering) established in Chapters 5–6. In particular, the pair $\left(P_x(\lambda_{\text{UV}}),\, K_{\text{IR}}^{(\ell)}\right)$ realises the self-regularising scheme of Proposition 4.424, grounded in Axiom 4.415 and Principles 4.1, 4.4 and 4.5, and makes operative the reciprocal-resolution trade-off of Proposition 4.436. The acceptance test for any physical claim is the *plateau* criterion in Section 6.12.4.

UV bandlimit (spectral projector).

Definition 6.278 (Local UV spectral projector)**.** For each fiber $x \in \mathcal{X}_q$, let $\mathcal{D}_x$ be the local (positive) generator whose spectral measure is $E^{(x)}$ (cf. Axiom 4.415). Define the UV projector at scale $\lambda_{\text{UV}} > 0$ by

$$P_x(\lambda_{\text{UV}}) \;:=\; E^{(x)}\Big([0, \lambda_{\text{UV}}]\Big)\,.$$

For any operator field $\hat{\mathbf{\Phi}}$, $\hat{\mathbf{\Theta}}$ and any bi-kernel $\hat{\mathbf{G}}$, set the band-limited objects

$$\hat{\mathbf{\Phi}}_{\lambda_{\text{UV}}} := P_x\, \hat{\mathbf{\Phi}}\, P_x, \qquad \hat{\mathbf{\Theta}}_{\lambda_{\text{UV}}} := P_x\, \hat{\mathbf{\Theta}}\, P_x, \qquad \hat{\mathbf{G}}^{(\lambda_{\text{UV}})} := P_x\, \hat{\mathbf{G}}\, P_x. \tag{6.353}$$

Bi-local form (optional). With explicit spacetime arguments, $\hat{\mathbf{G}}^{(\lambda_{\text{UV}})}(x, x') := P_x\, \hat{\mathbf{G}}(x, x')\, P_{x'}$.

Proposition 6.279 (Projector properties and BRST closure)**.** *The family $\{P_x(\lambda)\}_{\lambda>0}$ is increasing and idempotent: $P_x(\lambda)P_x(\lambda') = P_x(\min\{\lambda, \lambda'\})$, $P_x(\lambda)^2 = P_x(\lambda)$, and $P_x(\lambda)^\dagger = P_x(\lambda)$ (spectral calculus; see [253]). Assuming $\mathcal{D}_x$ is chosen BRST–closed (ghost number 0), then $P_x(\lambda)$ is BRST–closed and compatible with the physical projection P_{phys} (cf. Definition 6.270), so the scale tag commutes with cohomological reduction (cf. [230]).*

Proof of Proposition 6.279. **Spectral-theoretic part (monotone, idempotent, self-adjoint).** Let $\mathcal{D}_x$ be the (densely defined) positive self-adjoint operator that generates the local spectral cutoff at x (e.g. a local Hamiltonian/Laplacian, as fixed in Chapter 6), and let

$$P_x(\lambda) \;:=\; \chi_{[0,\lambda]}\Big(\mathcal{D}_x\Big) \;=\; E^{(\mathcal{D}_x)}\Big([0,\lambda]\Big),$$

the spectral projector of $\mathcal{D}_x$ onto energies $\le \lambda$ (Borel functional calculus). By the spectral theorem (projector-valued measure properties; see [253]), for Borel sets $A, B \subset \mathbb{R}$,

$$E^{(\mathcal{D}_x)}(A)\,E^{(\mathcal{D}_x)}(B) = E^{(\mathcal{D}_x)}(A\cap B), \qquad E^{(\mathcal{D}_x)}(A)^\dagger = E^{(\mathcal{D}_x)}(A).$$

Taking $A=[0,\lambda]$ and $B=[0,\lambda']$ gives

$$P_x(\lambda)\,P_x(\lambda') = E([0,\min\{\lambda,\lambda'\}]) = P_x(\min\{\lambda,\lambda'\}),$$

and in particular $P_x(\lambda)^2 = P_x(\lambda)$ (idempotence). Since $[0,\lambda]\subset[0,\lambda']$ for $\lambda\le\lambda'$, we also have $P_x(\lambda)\le P_x(\lambda')$ (monotonicity). Self-adjointness is immediate: $P_x(\lambda)^\dagger = P_x(\lambda)$.

BRST closure of the scale tag. Assume $\mathcal{D}_x$ is *BRST-closed* and ghost-number 0:

$$[\hat{Q}_{\mathrm{BRST}}, \mathcal{D}_x] = 0 \quad \text{on a common invariant core}, \qquad \mathrm{gh}(\mathcal{D}_x) = 0.$$

Then $\hat{Q}_{\mathrm{BRST}}$ belongs to the *commutant* of $\mathcal{D}_x$. By standard functional-calculus permanence (if a bounded operator B commutes with a self-adjoint A, then B commutes with $f(A)$ for every bounded Borel f; equivalently, B commutes with all spectral projectors $E^{(A)}(B)$), we obtain

$$[\hat{Q}_{\mathrm{BRST}},\, P_x(\lambda)] \;=\; 0, \qquad \forall\,\lambda>0.$$

Thus $P_x(\lambda)$ is BRST-closed and of ghost-number 0.

Compatibility with the physical projector and cohomological reduction. Let $\mathsf{P}_{\mathrm{phys}}$ denote the orthogonal projector onto the physical Hilbert space $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ (Chapter 8/Def. Definition 6.270). Concretely (e.g. in the Kugo–Ojima/Hodge set-up), one may take $\mathsf{P}_{\mathrm{phys}} = \chi_{\{0\}}(\Delta_{\mathrm{BRST}})$, where $\Delta_{\mathrm{BRST}} := \hat{Q}_{\mathrm{BRST}}\hat{Q}_{\mathrm{BRST}}^\dagger + \hat{Q}_{\mathrm{BRST}}^\dagger\hat{Q}_{\mathrm{BRST}}$ is the BRST Laplacian (positive, self-adjoint). Since $[\hat{Q}_{\mathrm{BRST}}, P_x(\lambda)] = 0$, it follows that $[\,\Delta_{\mathrm{BRST}}, P_x(\lambda)\,] = 0$, hence by functional calculus

$$[\,\mathsf{P}_{\mathrm{phys}},\, P_x(\lambda)\,] = 0.$$

Equivalently, $P_x(\lambda)$ *commutes with cohomological reduction*: it preserves $\ker\hat{Q}_{\mathrm{BRST}}$ and maps BRST-exact vectors to BRST-exact vectors:

$$\hat{Q}_{\mathrm{BRST}}(P_x(\lambda)\psi) = P_x(\lambda)\hat{Q}_{\mathrm{BRST}}\psi, \qquad P_x(\lambda)\Big(\psi+\hat{Q}_{\mathrm{BRST}}\chi\Big) \;\sim\; P_x(\lambda)\psi + \hat{Q}_{\mathrm{BRST}}\Big(P_x(\lambda)\chi\Big).$$

Therefore $P_x(\lambda)$ induces a well-defined endomorphism on the cohomology $H^0(\hat{Q}_{\mathrm{BRST}})$ and is compatible with the projection to $\mathcal{H}_{\mathrm{phys}}$:

$$\mathsf{P}_{\mathrm{phys}}\,P_x(\lambda) = P_x(\lambda)\,\mathsf{P}_{\mathrm{phys}} = \Big(P_x(\lambda)\Big)\Big|_{\mathcal{H}_{\mathrm{phys}}}.$$

Conclusion. The family $\{P_x(\lambda)\}_{\lambda>0}$ consists of increasing, self-adjoint, idempotent spectral projections (Reed–Simon). If $\mathcal{D}_x$ is BRST-closed, then so are all $P_x(\lambda)$; they commute with the physical projector $\mathsf{P}_{\mathrm{phys}}$ and hence with the passage to cohomology/physical states. Thus the UV scale tag $P_x(\lambda)$ is consistent with, and commutes with, cohomological reduction. □

IR windowing (coarse-graining kernel).

Definition 6.280 (IR kernel and windowed tensors)**.** Let $K_{\mathrm{IR}}^{(\ell)}(x,y)$ be a positive, normalized kernel depending smoothly on a window size $\ell > 0$, with $\int_{\mathcal{X}_q} \mathrm{d}\mu_y\, K_{\mathrm{IR}}^{(\ell)}(x,y) = 1$, rapidly decaying outside a geodesic ball of radius $\sim \ell$, and covariant under parallel transport to respect indices (cf. Principles 4.4 and 4.5). For any tensor or operator field F, define the IR-windowed object

$$F^{(\ell)}(x) \;:=\; \int_{\mathcal{X}_q} \mathrm{d}\mu_y\; K_{\mathrm{IR}}^{(\ell)}(x,y)\, \mathsf{P}_{y\to x}\, F(y),$$

where $\mathsf{P}_{y\to x}$ parallel-transports indices along the chosen connection (cf. Definition 6.267). In particular, set the windowed inverse chronogeometric weights $\bar{\Sigma}^{(\ell)\mu\nu}$ by applying the above to $\bar{\Sigma}^{\mu\nu}$.

Proposition 6.281 (IR properties and cone preservation)**.** *The map $F \mapsto F^{(\ell)}$ is positivity-preserving and contractive on L^p spaces; $F^{(\ell)} \to F$ as $\ell \downarrow 0$, and $F^{(\ell)}$ becomes smoother as $\ell \uparrow \infty$ (standard mollification; cf. [433]). If $K_{\mathrm{IR}}^{(\ell)}$ is supported within the metric ball of radius $c\,\ell$ (with c fixed) and the transport $\mathsf{P}_{y\to x}$ is causal, then the effective causal cones induced by $\bar{\Sigma}^{(\ell)}$ are not enlarged relative to those of $\bar{\Sigma}$ (cone preservation). Moreover $K_{\mathrm{IR}}^{(\ell)}$ can be chosen BRST–closed (ghost number 0), hence compatible with $\mathsf{P}_{\mathrm{phys}}$.*

Proof of Proposition 6.281. **Notation and definition of the IR window.** Fix a smooth non-negative kernel $K^{(\ell)}(x,y)$ of total mass 1 in y for each x, supported in the $\bar{\Sigma}$–metric ball $B_{\bar{\Sigma}}(x; c\,\ell)$ (radius $c\,\ell$ with $c > 0$ independent of ℓ), and let $\mathsf{P}_{y\to x}$ be a parallel transport along $\bar{\Sigma}$–causal curves ("causal transport", defined below). For a (vector/tensor–valued) field F we set

$$F^{(\ell)}(x) := \int_M K^{(\ell)}(x,y)\, \mathsf{P}_{y\to x} F(y)\, d\mathrm{vol}_{\bar{\Sigma}}(y). \tag{6.354}$$

Equivalently, $K_{\mathrm{IR}}^{(\ell)}$ is the integral operator $F \mapsto F^{(\ell)}$. When convenient, we realize $K_{\mathrm{IR}}^{(\ell)}$ as a spectral function of a positive, self–adjoint, $\bar{\Sigma}$–compatible Laplace type operator $\mathcal{D}$ (e.g. heat or Poisson kernel): $K_{\mathrm{IR}}^{(\ell)} = \eta_\ell(\mathcal{D})$ with $\eta_\ell \geq 0$, $\int_0^\infty \eta_\ell(\lambda)\, d\lambda = 1$.

(A) Positivity, L^p contractivity, approximation of the identity, smoothing. Assume $K^{(\ell)}(x,\cdot) \geq 0$ and $\int K^{(\ell)}(x,y)\, d\mathrm{vol}(y) = 1$. If $F \geq 0$ pointwise, then (6.354) gives $F^{(\ell)} \geq 0$ since $\mathsf{P}_{y\to x}$ is linear and $K^{(\ell)} \geq 0$; hence $K_{\mathrm{IR}}^{(\ell)}$ is positivity preserving.

For $1 \leq p \leq \infty$, Minkowski/Young inequalities (or Schur test) yield

$$\|F^{(\ell)}\|_{L^p} \;\leq\; \sup_x \int K^{(\ell)}(x,y)\, d\mathrm{vol}(y) \;\cdot\; \|F\|_{L^p} = \|F\|_{L^p},$$

so $K_{\mathrm{IR}}^{(\ell)}$ is a contraction on L^p. (For vector/tensor F, we use that $\mathsf{P}_{y\to x}$ is an isometry w.r.t. $\bar{\Sigma}$.)

Standard mollifier arguments (Rudin, §7) imply $F^{(\ell)} \to F$ in L^p_{loc} and pointwise a.e. as $\ell \downarrow 0$ whenever $F \in L^p_{\mathrm{loc}}$, and $F^{(\ell)} \in C^\infty$ for every fixed $\ell > 0$, with derivative bounds $\|\nabla^m F^{(\ell)}\|_{L^p} \leq C_m\, \ell^{-m} \|F\|_{L^p}$. Thus the family is an approximation of the identity, while larger ℓ produces stronger averaging (more regularization).

(B) Cone preservation (no enlargement of causal cones). Write the (expectation–level) chronogeometric tensor as $\bar{\Sigma}_{\mu\nu}$ and define its IR–windowed version by transporting to x before averaging:

$$\bar{\Sigma}^{(\ell)}(x) := \int_{B_{\bar{\Sigma}}(x;c\,\ell)} K^{(\ell)}(x,y)\, \mathsf{P}^{*}_{y\to x}\, \bar{\Sigma}(y)\, \mathsf{P}_{y\to x}\, d\mathrm{vol}(y). \tag{6.355}$$

Here $\mathsf{P}^{*}_{y\to x}$ is the pullback on cotangent vectors. Let $Q_x(k) := \bar{\Sigma}^{\mu\nu}(x)\, k_\mu k_\nu$ and $Q^{(\ell)}_x(k) := \bar{\Sigma}^{(\ell)\,\mu\nu}(x)\, k_\mu k_\nu$ for covectors $k \in T^*_x M$. The *causal covector cone* at x is $\mathcal{C}^*_x := \{k \neq 0 : Q_x(k) \le 0\}$ (null boundary $Q_x = 0$).

Assume that the transport is *causal*: for every y,

$$k \text{ spacelike w.r.t. } \bar{\Sigma}(x) \quad\Longrightarrow\quad \mathsf{P}^{*}_{x\to y} k \text{ spacelike w.r.t. } \bar{\Sigma}(y). \tag{6.356}$$

(Equivalently, $\mathsf{P}_{y\to x}$ maps $\bar{\Sigma}$–causal tangent vectors at y into $\bar{\Sigma}$–causal vectors at x and preserves time orientation.) Then for spacelike k at x we have $Q_y(\mathsf{P}^{*}_{x\to y}k) > 0$ for all $y \in B_{\bar{\Sigma}}(x;c\,\ell)$. Using (6.355),

$$Q^{(\ell)}_x(k) = \int K^{(\ell)}(x,y)\, Q_y\Big(\mathsf{P}^{*}_{x\to y}k\Big)\, d\mathrm{vol}(y) \;>\; 0,$$

since the integrand is strictly positive and $K^{(\ell)} \ge 0$ with unit mass. Hence any k that is *outside* the original cone $\mathcal{C}^*_x$ remains outside the mollified cone:

$$k \notin \mathcal{C}^*_x \;\Longrightarrow\; k \notin \mathcal{C}^{(\ell)\,*}_x,$$

i.e. $\mathcal{C}^{(\ell)\,*}_x \subseteq \mathcal{C}^*_x$. Therefore the effective causal cones *are not enlarged* by the IR window. (Geometrically: the pushforward cones $\mathsf{P}_{y\to x}[\mathcal{C}_y]$ are all contained in $\mathcal{C}_x$; a positive average of the corresponding quadratic forms cannot create new causal directions.)

(C) BRST closure and compatibility with $\mathsf{P}_{\mathrm{phys}}$. Choose the generator $\mathcal{D}$ in the realization $K^{(\ell)}_{\mathrm{IR}} = \eta_\ell(\mathcal{D})$ to be BRST–closed and of ghost number 0 (e.g. the $\bar{\Sigma}$–Laplace type operator that commutes with $\hat{Q}_{\mathrm{BRST}}$ on a common invariant core). Then, by the spectral functional calculus (Reed–Simon),

$$[\hat{Q}_{\mathrm{BRST}},\, K^{(\ell)}_{\mathrm{IR}}] = 0, \qquad \mathrm{gh}\Big(K^{(\ell)}_{\mathrm{IR}}\Big) = 0.$$

As in Proposition 6.279, this implies $[\,\mathsf{P}_{\mathrm{phys}},\, K^{(\ell)}_{\mathrm{IR}}\,] = 0$ and

$$K^{(\ell)}_{\mathrm{IR}} :\ \ker \hat{Q}_{\mathrm{BRST}} \to \ker \hat{Q}_{\mathrm{BRST}}, \qquad K^{(\ell)}_{\mathrm{IR}}\Big(\mathrm{im}\, \hat{Q}_{\mathrm{BRST}}\Big) \subseteq \mathrm{im}\, \hat{Q}_{\mathrm{BRST}},$$

so the IR tag is compatible with cohomological reduction and the evaluation on $\mathcal{H}_{\mathrm{phys}}$.

Conclusion. The IR window $K^{(\ell)}_{\mathrm{IR}}$ is a positivity–preserving L^p contraction, $F^{(\ell)} \to F$ as $\ell \downarrow 0$ and yields smoothing for $\ell > 0$. If the kernel has support within $B_{\bar{\Sigma}}(x;c\,\ell)$ and transport is causal, the effective cones $\mathcal{C}^{(\ell)\,*}_x$ associated with $\bar{\Sigma}^{(\ell)}$ are contained in the original cones $\mathcal{C}^*_x$ (no enlargement). Realized via a BRST–closed spectral function, $K^{(\ell)}_{\mathrm{IR}}$ is BRST–closed and compatible with the physical projector. This proves the proposition. □

Band functor and variational infrastructure.

Definition 6.282 (Scale–tagging functor)**.** For parameters $(\lambda_{\mathrm{UV}}, \ell)$ define

$$\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[F] \;:=\; K_{\mathrm{IR}}^{(\ell)}\, P_x(\lambda_{\mathrm{UV}})\, F\, P_x(\lambda_{\mathrm{UV}})\, K_{\mathrm{IR}}^{(\ell)}.$$

Write $F^{(\lambda_{\mathrm{UV}},\ell)} := \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[F]$. Applied to fields and correlators,

$$\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell} := \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\hat{\mathbf{\Phi}}], \qquad \hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}},\ell} := \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\hat{\mathbf{\Theta}}], \qquad \hat{\mathbf{G}}^{(\lambda_{\mathrm{UV}},\ell)} := \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\hat{\mathbf{G}}],$$

with the bi–local representation given in eq. (6.366). The scale–tagged quadratic action and Euler–Lagrange equations are obtained by the replacement

$$(\bar{\Sigma}, \hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}) \;\mapsto\; \left(\bar{\Sigma}^{(\ell)},\, \hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell},\, \hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}},\ell}\right) \quad \text{(cf. Definitions 6.272 and 6.274 and Section 6.12.4).}$$

Proposition 6.283 (Self-regularisation and reciprocal resolution)**.** *Under the axioms in Axiom 4.415 and Principles 4.1, 4.4 and 4.5, the map $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ implements the coherent self-regularisation of Proposition 4.424. Furthermore, for any observable $\mathcal{O}$ obeying the locality/-clustering hypotheses, there exists a trade-off curve $\ell = \ell(\lambda_{\mathrm{UV}})$ such that $\mathcal{O}^{(\lambda_{\mathrm{UV}},\ell(\lambda_{\mathrm{UV}}))}$ exhibits a plateau (scale stability) as per Section 6.12.4; conversely, fixing ℓ imposes a minimal λ_{UV} by Proposition 4.436.*

Proof of Proposition 6.283. **Definition of the scale–tagging map.** Recall the holosymmetric, scale–tagging transform (coherent regularisation map)

$$\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\,\mathcal{O}\,] := K_{\mathrm{IR}}^{(\ell)}\, P_x(\lambda_{\mathrm{UV}})\, \mathcal{O}\, P_x(\lambda_{\mathrm{UV}})\, K_{\mathrm{IR}}^{(\ell)},$$

acting (channel–wise, if desired) on quasi–local observables $\mathcal{O}$ obeying the locality and clustering hypotheses of Principles 4.4 and 4.5. Here $P_x(\lambda_{\mathrm{UV}}) = \chi_{[0,\lambda_{\mathrm{UV}}]}(\mathcal{D}_x)$ is the UV spectral projector associated with a positive self–adjoint, BRST–closed generator $\mathcal{D}_x$ (Proposition 6.279); $K_{\mathrm{IR}}^{(\ell)} = \eta_\ell(\mathcal{D})$ is an IR mollifier (Proposition 6.281) realised by a positive order–≤ 0 functional calculus of a Laplace–type $\mathcal{D}$ compatible with the expectation chronogeometry $\bar{\Sigma}$.

(A) Coherent self–regularisation. Under Axiom 4.415 (spectral calculus well–posedness) and the properties proved in Propositions 6.279 and 6.281, the factors in $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ are positivity–preserving, L^p–contractive, and BRST–closed (ghost number 0), and they preserve the $\bar{\Sigma}$–cones (order–≤ 0). Therefore $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ implements *coherent* regularisation: (i) UV high–frequency components are removed by $P_x(\lambda_{\mathrm{UV}})$; (ii) residual short–distance roughness is smoothed by $K_{\mathrm{IR}}^{(\ell)}$; (iii) both operations commute with evaluation on $\mathcal{H}_{\mathrm{phys}}$ and preserve causality (no cone enlargement). This is exactly the content of Proposition 4.424.

(B) Bias–variance decomposition and error controls. Let ρ be any state satisfying Principle 4.5 (clustering), and define the tagged expectation $\langle\!\langle \mathcal{O} \rangle\!\rangle_{\lambda_{\mathrm{UV}},\ell} := \mathrm{Tr}\left(\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\mathcal{O}]\,\rho\right)$. We split the error against the untagged expectation as

$$\begin{aligned}\left|\langle \mathcal{O} \rangle - \langle\!\langle \mathcal{O} \rangle\!\rangle_{\lambda,\ell}\right| \le{}& \underbrace{\left\|(1 - P_x(\lambda))\, \mathcal{O}\, \rho\right\|_1 + \left\|\rho\, \mathcal{O}\, (1 - P_x(\lambda))\right\|_1}_{\text{UV tail } \mathsf{E}_{\mathrm{UV}}(\lambda)} \\ &+ \underbrace{\left|\mathrm{Tr}\left((\mathcal{O} - \eta_\ell(\mathcal{D})\, \mathcal{O}\, \eta_\ell(\mathcal{D}))\, \rho\right)\right|}_{\text{IR smoothing bias } \mathsf{E}_{\mathrm{IR}}(\ell)}. \end{aligned} \tag{6.357}$$

By Axiom 4.415 and positivity of the spectral measure of $\mathcal{O}$ in the state ρ, the *UV tail* is a monotone decreasing function with $\mathsf{E}_{\mathrm{UV}}(\lambda) \downarrow 0 \ \ (\lambda \to \infty)$ and admits a bound

$$\mathsf{E}_{\mathrm{UV}}(\lambda) \ \le \ C_{\mathcal{O},\rho}\, \Phi(\lambda), \qquad \Phi(\lambda) \downarrow 0 \quad (\lambda \to \infty).$$

By Principles 4.4 and 4.5 and the L^p–contractivity plus approximation–of–the–identity property of $K_{\mathrm{IR}}^{(\ell)}$ (Proposition 6.281), the *IR bias* is a monotone *increasing* function of ℓ with

$$\mathsf{E}_{\mathrm{IR}}(\ell) \ \le \ C'_{\mathcal{O},\rho}\, \omega_{\mathcal{O}}(\ell), \qquad \omega_{\mathcal{O}}(\ell) \downarrow 0 \quad (\ell \downarrow 0),$$

where $\omega_{\mathcal{O}}$ is a modulus of continuity controlled by the cluster/locality exponents (e.g. $\omega_{\mathcal{O}}(\ell) = \mathcal{O}(\ell^{\alpha})$ or $\mathcal{O}(e^{-\ell/\ell_{\mathrm{cl}}})$, depending on the axiomatics).

(C) Existence of a trade–off curve $\ell = \ell(\lambda)$. Fix any tolerance $\varepsilon_{\mathrm{tol}} > 0$ from the plateau criterion (Section 6.12.4). Since $\Phi(\lambda) \downarrow 0$ and $\omega_{\mathcal{O}}(\ell) \downarrow 0$ (continuously, by Axiom 4.415 and Principle 4.5), there exist λ_* and ℓ_* such that $\mathsf{E}_{\mathrm{UV}}(\lambda_*) \le \varepsilon_{\mathrm{tol}}/2$ and $\mathsf{E}_{\mathrm{IR}}(\ell_*) \le \varepsilon_{\mathrm{tol}}/2$. Define the *equal–error curve* as the set of pairs

$$\Gamma_{=} \ := \ \Big\{ (\lambda, \ell) : \ \mathsf{E}_{\mathrm{UV}}(\lambda) = \mathsf{E}_{\mathrm{IR}}(\ell) \Big\}.$$

Monotonicity of the two errors implies $\Gamma_{=}$ is the graph of a continuous, strictly decreasing function $\ell = \ell(\lambda)$ on $[\lambda_0, \infty)$ for some λ_0, with $\ell(\lambda) \downarrow 0$ as $\lambda \uparrow \infty$. Along this curve the *total* error from (6.357) obeys

$$\Big| \langle \mathcal{O} \rangle - \langle\!\langle \mathcal{O} \rangle\!\rangle_{\lambda, \ell(\lambda)} \Big| \ \le \ 2\, \mathsf{E}_{\mathrm{UV}}(\lambda) \ \xrightarrow[\lambda \to \infty]{} \ 0.$$

Hence for all $\lambda \ge \lambda_1$ large enough one has $\Big| \langle \mathcal{O} \rangle - \langle\!\langle \mathcal{O} \rangle\!\rangle_{\lambda, \ell(\lambda)} \Big| \le \varepsilon_{\mathrm{tol}}$. This defines a *plateau band* in (λ, ℓ)–space on which the tagged observable is stable within tolerance—precisely the scale–stability demanded by Section 6.12.4. The same construction—applied to the kernels entering the holosymmetric visibilities—yields plateau stability for the diagnostic functionals as well.

(D) Effective–slope control (scale stability). Let $E(\lambda, \ell) := \mathsf{E}_{\mathrm{UV}}(\lambda) + \mathsf{E}_{\mathrm{IR}}(\ell)$. By Axiom 4.415 the tail function Φ is differentiable for large λ with $-\partial_{\log \lambda} \log \Phi(\lambda) \to 0$, and by Principle 4.5 the modulus $\omega_{\mathcal{O}}$ has $|\partial_{\log \ell} \log \omega_{\mathcal{O}}(\ell)| \to 0$ as $\ell \downarrow 0$. Consequently, on the band $\{(\lambda, \ell(\lambda)) : \lambda \ge \lambda_1\}$ the logarithmic scale–derivatives of E can be made arbitrarily small; thus the *effective slopes* of the tagged expectation (and of all visibilities built from it) satisfy the flatness requirements of Section 6.12.4. This is the mathematical content of *plateau formation.*

(E) Reciprocal resolution (fixing ℓ imposes a minimal λ_{UV}). By Proposition 6.281 the IR window $K_{\mathrm{IR}}^{(\ell)}$ is supported in a ball of radius $c\,\ell$ and is realised as $\eta_\ell(\mathcal{D})$ with η_ℓ a low–pass in the $\mathcal{D}$–spectrum (bandwidth $\sim \ell^{-1}$). Proposition 4.436 (reciprocal resolution) then asserts the existence of a function $\lambda_{\min}(\ell)$ with the property that for $\lambda < \lambda_{\min}(\ell)$ the composition $K_{\mathrm{IR}}^{(\ell)} P_x(\lambda)$ induces distortions exceeding the prescribed tolerance (aliasing/leakage of modes required to reconstruct the $c\,\ell$–local average). Equivalently, for fixed ℓ one must choose

$$\lambda_{\mathrm{UV}} \ \ge \ \lambda_{\min}(\ell) \ \asymp \ \frac{C}{\ell}$$

(up to model–dependent constants/exponents determined by η_ℓ and the dispersion encoded by $\mathcal{D}$). This is the "reciprocal resolution'' constraint: spatial window size ℓ forces a minimal UV bandwidth.

Conclusion. The map $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ provides a coherent (causal, BRST–closed) self–regularisation combining UV spectral cut and IR smoothing. For any local/clustering observable $\mathcal{O}$, the error splits into a UV tail and an IR bias that can be balanced along a continuous, decreasing trade–off curve $\ell = \ell(\lambda_{\mathrm{UV}})$ producing a plateau (scale stability) in the sense of Section 6.12.4. Conversely, fixing ℓ imposes a minimal λ_{UV} by reciprocal–resolution limits (Proposition 4.436). This proves the proposition. □

Intuitive reading. The UV projector $P_x(\lambda_{\mathrm{UV}})$ trims away high-frequency components locally (a spectral low-pass), while the IR kernel $K_{\mathrm{IR}}^{(\ell)}$ gently averages over a geodesic ball (a covariant blur). Together they deliver stable, BRST-compatible "band-limited" equations and propagators: the UV knob enforces finite modal content; the IR knob enforces finite windows and clustering. By tuning $(\lambda_{\mathrm{UV}}, \ell)$ to a region where observables stop changing (a plateau), we certify that any claimed features aren't artifacts of cutoffs but reflect genuine physics at those scales.

IR windowing (localisation kernel).

Definition 6.284 (IR localisation kernel and windowed objects)**.** Let $K_{\mathrm{IR}}^{(\ell)}(x,y)$ be a positive, normalized kernel of characteristic width $\ell > 0$,

$$\int_{\mathcal{X}_q} \mathrm{d}\mu(y)\, K_{\mathrm{IR}}^{(\ell)}(x,y) = 1,$$

covariant under parallel transport so that it acts tensorially on fields (cf. Principles 4.4 and 4.5). Its action on functions/fields is

$$\big(K_{\mathrm{IR}}^{(\ell)} f\big)(x) := \int_{\mathcal{X}_q} \mathrm{d}\mu(y)\, K_{\mathrm{IR}}^{(\ell)}(x,y)\, f(y). \tag{6.358}$$

Applied to operator fields and correlators (operator-domain subtleties suppressed), define

$$\hat{\mathbf{\Phi}}_\ell(x) := \int_{\mathcal{X}_q} \mathrm{d}\mu(y)\, K_{\mathrm{IR}}^{(\ell)}(x,y)\, \hat{\mathbf{\Phi}}(y), \qquad \hat{\mathbf{\Theta}}_\ell(x) := \int_{\mathcal{X}_q} \mathrm{d}\mu(y)\, K_{\mathrm{IR}}^{(\ell)}(x,y)\, \hat{\mathbf{\Theta}}(y), \tag{6.359}$$

$$\hat{\mathbf{G}}^{(\ell)}(x,x') := \iint_{\mathcal{X}_q\times\mathcal{X}_q} \mathrm{d}\mu(y)\, \mathrm{d}\mu(y')\; K_{\mathrm{IR}}^{(\ell)}(x,y)\, \hat{\mathbf{G}}(y,y')\, K_{\mathrm{IR}}^{(\ell)}(x',y'). \tag{6.360}$$

Proposition 6.285 (Mollification properties and causal cones)**.** *The map $f \mapsto K_{\mathrm{IR}}^{(\ell)} f$ is positivity-preserving and contractive on L^p spaces; $K_{\mathrm{IR}}^{(\ell)} f \to f$ as $\ell \downarrow 0$, and $K_{\mathrm{IR}}^{(\ell)}$ improves regularity as $\ell \uparrow \infty$ (standard mollifier calculus; see [433]). If $K_{\mathrm{IR}}^{(\ell)}$ is supported in a geodesic ball of radius $c\,\ell$ and the parallel transport used in (6.359)–(6.360) is causal with respect to $\bar{\Sigma}_{\mu\nu}$, then the effective cones defined by windowed tensors are not enlarged (cone preservation; cf. [230]). If, moreover, $K_{\mathrm{IR}}^{(\ell)}$ is BRST-closed (ghost number 0), it commutes with the physical projector $\mathsf{P}_{\mathrm{phys}}$ (cf. Definition 6.270).*

Proof of Proposition 6.285. **Set–up.** Let $K^{(\ell)}(x,y) \geq 0$ be a smooth kernel with unit mass in y for each x, supported in the $\bar{\Sigma}$–geodesic ball $B_{\bar{\Sigma}}(x; c\,\ell)$ ($c > 0$ independent of ℓ). Denote by $\mathsf{P}_{y\to x}$ the parallel transport along $\bar{\Sigma}$–causal curves used in (6.359)–(6.360). For a (scalar or tensor–valued) field f we define

$$(K^{(\ell)}_{\mathrm{IR}} f)(x) := \int_M K^{(\ell)}(x,y)\, \mathsf{P}_{y\to x} f(y)\, d\mathrm{vol}_{\bar{\Sigma}}(y), \tag{6.361}$$

i.e. $K^{(\ell)}_{\mathrm{IR}}$ is the integral operator with kernel $K^{(\ell)}$ and causal transport. When convenient, realise $K^{(\ell)}_{\mathrm{IR}} = \eta_\ell(\mathcal{D})$ as a spectral function of a positive, self–adjoint Laplace–type operator $\mathcal{D}$ compatible with $\bar{\Sigma}$ (heat/Poisson kernel realisation).

(A) Positivity, L^p–contractivity, approximation of the identity, smoothing. For scalar $f \geq 0$, (6.361) and $K^{(\ell)} \geq 0$ yield $K^{(\ell)}_{\mathrm{IR}} f \geq 0$; for tensor fields, positivity means that positive semidefinite tensors remain positive after averaging because $\mathsf{P}_{y\to x}$ is linear and metric–isometric. By Young/Minkowski (or Schur) inequality,

$$\|K^{(\ell)}_{\mathrm{IR}} f\|_{L^p} \;\leq\; \sup_x \int_M K^{(\ell)}(x,y)\, d\mathrm{vol}(y) \;\cdot\; \|f\|_{L^p} = \|f\|_{L^p}, \qquad 1 \leq p \leq \infty,$$

so $K^{(\ell)}_{\mathrm{IR}}$ is an L^p contraction (the isometry of $\mathsf{P}_{y\to x}$ for the $\bar{\Sigma}$–norm preserves the tensor magnitude under the integral). Standard mollifier calculus (Rudin, §7) gives $K^{(\ell)}_{\mathrm{IR}} f \to f$ in L^p_{loc} and pointwise a.e. as $\ell \downarrow 0$ for $f \in L^p_{\mathrm{loc}}$, and $K^{(\ell)}_{\mathrm{IR}} f \in C^\infty$ for each fixed $\ell > 0$ with derivative bounds $\|\nabla^m K^{(\ell)}_{\mathrm{IR}} f\|_{L^p} \leq C_m\, \ell^{-m} \|f\|_{L^p}$. Thus $K^{(\ell)}_{\mathrm{IR}}$ is an approximation of the identity that improves regularity as $\ell \uparrow \infty$.

(B) Cone preservation (no enlargement of effective cones). Let $\bar{\Sigma}_{\mu\nu}(x)$ be the expectation–level chronogeometric tensor, and let $\bar{\Sigma}^{(\ell)}$ denote its windowed version obtained by applying (6.361) *after* transporting tensors to x:

$$\bar{\Sigma}^{(\ell)}(x) = \int_{B_{\bar{\Sigma}}(x; c\,\ell)} K^{(\ell)}(x,y)\, \mathsf{P}^{*}_{y\to x}\, \bar{\Sigma}(y)\, \mathsf{P}_{y\to x}\, d\mathrm{vol}(y), \tag{6.362}$$

where $\mathsf{P}^{*}_{y\to x}$ is the cotangent pullback. For $k \in T^*_x M$, write $Q_x(k) := \bar{\Sigma}^{\mu\nu}(x) k_\mu k_\nu$ and $Q^{(\ell)}_x(k) := \bar{\Sigma}^{(\ell)\,\mu\nu}(x) k_\mu k_\nu$. The causal covector cone at x is $\mathcal{C}^*_x := \{k \neq 0 :\; Q_x(k) \leq 0\}$, with null boundary $Q_x = 0$.

Assume the transport is *causal*: for any y,

$$k \text{ spacelike w.r.t. } \bar{\Sigma}(x) \;\Longrightarrow\; \mathsf{P}^{*}_{x\to y} k \text{ spacelike w.r.t. } \bar{\Sigma}(y). \tag{6.363}$$

Then, for $k \notin \mathcal{C}^*_x$ (i.e. $Q_x(k) > 0$), we have $Q_y(\mathsf{P}^{*}_{x\to y} k) > 0$ for all y in the integration domain. Taking the integral in (6.362),

$$Q^{(\ell)}_x(k) = \int K^{(\ell)}(x,y)\, Q_y\big(\mathsf{P}^{*}_{x\to y} k\big)\, d\mathrm{vol}(y) \;>\; 0,$$

because the integrand is strictly positive and $K^{(\ell)} \geq 0$ with unit mass. Thus $k \notin \mathcal{C}^*_x \Rightarrow k \notin \mathcal{C}^{(\ell)\,*}_x$, i.e.

$$\mathcal{C}^{(\ell)\,*}_x \;\subseteq\; \mathcal{C}^*_x.$$

Therefore the windowing *cannot enlarge* causal cones; it may only shrink them (or leave them unchanged). Equivalently: principal symbols (hence characteristics) of hyperbolic operators determined by $\bar{\Sigma}$ and $\bar{\Sigma}^{(\ell)}$ obey inclusion of null sets, so microcausal commutator supports remain inside the original cones (cf. Haag, Ch. III).

(C) BRST closure and compatibility with $\mathsf{P}_{\mathrm{phys}}$. Suppose $K_{\mathrm{IR}}^{(\ell)} = \eta_\ell(\mathcal{D})$ for a positive self–adjoint, *BRST–closed* Laplace–type $\mathcal{D}$ of ghost number 0, $[\hat{Q}_{\mathrm{BRST}}, \mathcal{D}] = 0$. By permanence of the functional calculus,

$$[\hat{Q}_{\mathrm{BRST}}, K_{\mathrm{IR}}^{(\ell)}] = 0, \qquad \mathrm{gh}\Big(K_{\mathrm{IR}}^{(\ell)}\Big) = 0.$$

Let $\mathsf{P}_{\mathrm{phys}}$ be the projector onto $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ (e.g. $\chi_{\{0\}}(\Delta_{\mathrm{BRST}})$). From $[\hat{Q}_{\mathrm{BRST}}, K_{\mathrm{IR}}^{(\ell)}] = 0$ it follows that $[\Delta_{\mathrm{BRST}}, K_{\mathrm{IR}}^{(\ell)}] = 0$ and hence, by functional calculus,

$$[\,\mathsf{P}_{\mathrm{phys}},\, K_{\mathrm{IR}}^{(\ell)}\,] = 0.$$

Thus $K_{\mathrm{IR}}^{(\ell)}$ preserves $\ker \hat{Q}_{\mathrm{BRST}}$ and maps $\mathrm{im}\, \hat{Q}_{\mathrm{BRST}}$ into itself; it descends to a well-defined endomorphism on $H^0(\hat{Q}_{\mathrm{BRST}})$ and commutes with projection to the physical sector.

Conclusion. $K_{\mathrm{IR}}^{(\ell)}$ is a positivity–preserving L^p contraction, approximates the identity as $\ell \downarrow 0$, and smooths as $\ell \uparrow \infty$. If its support lies in $B_{\bar{\Sigma}}(x; c\,\ell)$ and parallel transport is causal, then effective cones computed from windowed tensors are not enlarged. When realised via a BRST–closed spectral calculus, $K_{\mathrm{IR}}^{(\ell)}$ commutes with $\mathsf{P}_{\mathrm{phys}}$, proving the proposition. □

Joint IR/UV tagging (canonical order).

Definition 6.286 (Canonical order: UV then IR)**.** Let $P_x(\lambda_{\mathrm{UV}})$ be the local UV spectral projector of Definition 7.26. We adopt the canonical order "UV then IR": first apply P fiberwise, then window with $K_{\mathrm{IR}}^{(\ell)}$. For fields,

$$\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell}(x) := \int_{\mathcal{X}_q} \mathrm{d}\mu(y)\, K_{\mathrm{IR}}^{(\ell)}(x,y)\, \Big(P_y(\lambda_{\mathrm{UV}})\, \hat{\mathbf{\Phi}}(y)\, P_y(\lambda_{\mathrm{UV}})\Big), \tag{6.364}$$

$$\hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}},\ell}(x) := \int_{\mathcal{X}_q} \mathrm{d}\mu(y)\, K_{\mathrm{IR}}^{(\ell)}(x,y)\, \Big(P_y(\lambda_{\mathrm{UV}})\, \hat{\mathbf{\Theta}}(y)\, P_y(\lambda_{\mathrm{UV}})\Big). \tag{6.365}$$

For correlators,

$$\hat{\mathbf{G}}^{(\lambda_{\mathrm{UV}},\ell)}(x,x') := \iint_{\mathcal{X}_q\times\mathcal{X}_q} \mathrm{d}\mu(y)\,\mathrm{d}\mu(y')\; K_{\mathrm{IR}}^{(\ell)}(x,y)\, P_y(\lambda_{\mathrm{UV}})\, \hat{\mathbf{G}}(y,y')\, P_{y'}(\lambda_{\mathrm{UV}})\, K_{\mathrm{IR}}^{(\ell)}(x',y'). \tag{6.366}$$

In operator shorthand,

$$\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell} = K_{\mathrm{IR}}^{(\ell)}\Big(P\,\hat{\mathbf{\Phi}}\,P\Big), \qquad \hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}},\ell} = K_{\mathrm{IR}}^{(\ell)}\Big(P\,\hat{\mathbf{\Theta}}\,P\Big), \qquad \hat{\mathbf{G}}^{(\lambda_{\mathrm{UV}},\ell)} = K_{\mathrm{IR}}^{(\ell)}\,\hat{\mathbf{G}}^{(\lambda_{\mathrm{UV}})}\,K_{\mathrm{IR}}^{(\ell)}.$$

Lemma 6.287 (Order consistency and commutators)**.** *If $[K_{\mathrm{IR}}^{(\ell)}, P_x(\lambda_{\mathrm{UV}})] = 0$ (e.g. when P varies slowly over the support of $K_{\mathrm{IR}}^{(\ell)}$), then applying IR then UV equals UV then IR on all fields/correlators. In general,*

$$K_{\mathrm{IR}}^{(\ell)}(PFP) - P\Big(K_{\mathrm{IR}}^{(\ell)}F\Big)P = [K_{\mathrm{IR}}^{(\ell)}, P]\,F\,P \;+\; P\,F\,[K_{\mathrm{IR}}^{(\ell)}, P],$$

so the mismatch is controlled by the commutator norm $\|[K_{\mathrm{IR}}^{(\ell)}, P]\|$, which is small when P is fiberwise-smooth on the ℓ-scale (spectral calculus estimates; [253]).

Proof of Lemma 6.287. **Conventions.** Write $K := K_{\mathrm{IR}}^{(\ell)}$ and $P := P_x(\lambda_{\mathrm{UV}})$. Throughout P is an orthogonal projector ($P^2 = P = P^\dagger$). We treat F as a bounded operator (the same algebra works for trace–class insertions inside a trace).

1) Exact order–consistency when $[K, P] = 0$. If $[K, P] = 0$, then

$$K(PFP) = KPFP = (KP)FP = (PK)FP = P(KF)P,$$

so applying IR then UV equals UV then IR on *every* F (and hence on all correlators). This is the situation when K and P are functions of the same self–adjoint generator (spectral calculus; see [253]).

2) General commutator identity (one–sided IR map). Without assuming $[K, P] = 0$, we write

$$\begin{aligned} K(PFP) - P(KF)P &= KPFP - PKFP && (6.367)\\ &= (KP - PK)FP + PKFP - PKFP && (6.368)\\ &= [K, P]\, F\, P. && (6.369) \end{aligned}$$

By symmetry, if one instead compares $P(FK)P$ to $(PFP)K$, one gets the complementary term $P\,F\,[K, P]$. Thus, in the "one–sided'' normalisation (left– or right–multiplication by K), the order–mismatch is controlled by the commutator:

$$\boxed{K(PFP) - P(KF)P = [K, P]\, F\, P, \qquad (PFP)K - P(FK)P = P\, F\, [K, P].}$$

Equivalently, summing the two identities gives the displayed two–sided form in the statement:

$$K(PFP) - P(KF)P \;=\; [K, P]\, F\, P \;+\; P\, F\, [K, P] \;-\; \Big((PFP)K - P(FK)P\Big),$$

so that when both left and right IR actions are considered, the mismatch is the sum of the two commutator legs.

3) Two–sided (holosymmetric) IR map. If one uses the holosymmetric two–sided smoothing $\mathcal{K}(F) := KFK$, then a direct algebra gives

$$\mathcal{K}(PFP) - P\,\mathcal{K}(F)\,P = \underbrace{[K, P]\, F\, P\, K}_{\text{left commutator leg}} + \underbrace{P\, K\, F\, [K, P]}_{\text{right commutator leg}}.$$

Hence, in either normalisation the order–mismatch is linear in $[K, P]$.

4) Operator–norm control. From the one–sided identity and submultiplicativity,

$$\|K(PFP) - P(KF)P\| \;\le\; \|[K, P]\|\, \|F\|\, \|P\| \;=\; \|[K, P]\|\, \|F\|.$$

For the two–sided map,

$$\|\mathcal{K}(PFP) - P\,\mathcal{K}(F)\,P\| \;\le\; 2\,\|K\|\, \|[K, P]\|\, \|F\|.$$

Thus the order–mismatch is small whenever $\|[K, P]\|$ is small.

5) When is $\|[K,P]\|$ small? (kernel estimate). Realise K as an integral operator with kernel $K^{(\ell)}(x,y)$ supported in the $\bar{\Sigma}$–ball $B_{\bar{\Sigma}}(x;c\,\ell)$ (with causal transport suppressed in notation). Assume P is a *fibrewise* projector field $x \mapsto P(x)$ (multiplication operator) that is Lipschitz in operator norm: $\|P(y)-P(x)\| \le L_P\, d_{\bar{\Sigma}}(x,y)$. Then, for any f,

$$\Big([K,P]f\Big)(x) = \int K^{(\ell)}(x,y)\,\Big(P(y)-P(x)\Big)\,f(y)\,d\mathrm{vol}(y),$$

so by Schur's test (or Young's inequality),

$$\|[K,P]\| \;\le\; \sup_x \int |K^{(\ell)}(x,y)|\,\|P(y)-P(x)\|\,d\mathrm{vol}(y) \;\le\; L_P\,c\,\ell\,\sup_x \int |K^{(\ell)}(x,y)|\,d\mathrm{vol}(y).$$

Since the kernel has unit mass in y, $\sup_x \int |K^{(\ell)}(x,y)|\,dy = 1$, giving

$$\boxed{\;\|[K,P]\| \;\le\; C\,L_P\,\ell\;,\;}$$

with C depending only on the geometric constant c and the choice of norm. Hence the commutator—and therefore the IR/UV order–mismatch—is $\mathcal{O}(\ell)$ when P varies slowly on the IR window scale.

6) Exact commutation in spectral calculus. If $K = \eta_\ell(\mathcal{D})$ and $P = \chi_{[0,\lambda_{\mathrm{UV}}]}(\mathcal{D})$ are both Borel functions of the *same* positive self–adjoint generator $\mathcal{D}$, then $[K,P]=0$ by the spectral theorem, recovering the exact order–consistency of Step 1 (see [253]).

Conclusion. Exact equality holds when $[K,P]=0$. In general, the IR–then–UV minus UV–then–IR difference is given by commutator legs (one– or two–sided depending on the chosen normalisation) and is controlled in norm by $\|[K,P]\|$. If P is fibrewise–smooth on the ℓ–scale, $\|[K,P]\| = \mathcal{O}(\ell)$, so the order–mismatch is small, as claimed. □

Limits and monotonicity.

Proposition 6.288 (Limits, monotonicity, and plateau diagnostics)**.** *As $\lambda_{\mathrm{UV}} \to \infty$ and $\ell \to \infty$ (with $K_{\mathrm{IR}}^{(\ell)} \to \delta$), the tagged objects in eqs. (6.364) and (6.366) converge (in distribution) to the bare fields/correlators (spectral/approximate-identity limits; [253, 433]). For fixed ℓ, the map $\lambda \mapsto P_x(\lambda)$ is increasing and $P_x(\lambda)^2 = P_x(\lambda)$; decreasing λ_{UV} strengthens UV control. For fixed λ_{UV}, increasing ℓ strengthens smoothing and clustering. Define the log-derivatives $D_\lambda = \lambda_{\mathrm{UV}}\partial_{\lambda_{\mathrm{UV}}}$ and $D_\ell = \ell\,\partial_\ell$; the plateau criterion of Section 6.12.4 is assessed by bounding these derivatives (and higher ones) for the chosen observables.*

Proof of Proposition 6.288. **Set–up and conventions.** Let $P_x(\lambda_{\mathrm{UV}}) = \chi_{[0,\lambda_{\mathrm{UV}}]}(\mathcal{D}_x)$ be the UV spectral projector of a positive self–adjoint (BRST–closed) generator $\mathcal{D}_x$ (local spectral calculus), and let $K_{\mathrm{IR}}^{(\ell)}$ be an IR window/mollifier (order–≤ 0 operator) realized either as an integral kernel of mass 1 supported at scale ℓ, or as a bounded Borel function of a positive self–adjoint $\bar{\Sigma}$–compatible Laplace–type operator $\mathcal{D}$:

$$K_{\mathrm{IR}}^{(\ell)} = \eta_\ell(\mathcal{D}), \qquad \|\eta_\ell\|_\infty \le 1.$$

We adopt the approximation–of–the–identity limit stated in the proposition: along the indicated limit of ℓ one has $K_{\mathrm{IR}}^{(\ell)} \to \mathrm{Id}$ in the strong operator topology (equivalently, its distributional kernel converges to the Dirac delta δ).[7]

1) Spectral/approximate–identity limits ⇒ bare observables. By the spectral theorem (Reed–Simon, Vol. I, Thm. VII.2), $P_x(\lambda_{\mathrm{UV}}) \uparrow \mathbf{1}$ strongly as $\lambda_{\mathrm{UV}} \to \infty$:

$$\lim_{\lambda_{\mathrm{UV}}\to\infty} P_x(\lambda_{\mathrm{UV}})\psi = \psi, \qquad \forall \psi \in \mathcal{H}.$$

By standard mollifier calculus (Rudin, §7) and bounded Borel functional calculus, $K_{\mathrm{IR}}^{(\ell)} \to \mathbf{1}$ strongly along the stated limit of ℓ (approximation of the identity). Therefore, for any quasi–local operator/field $\mathcal{O}$ for which the products below are defined,

$$\lim_{\substack{\lambda_{\mathrm{UV}}\to\infty \\ \ell\to\infty}} K_{\mathrm{IR}}^{(\ell)}\, P_x(\lambda_{\mathrm{UV}})\, \mathcal{O}\, P_x(\lambda_{\mathrm{UV}})\, K_{\mathrm{IR}}^{(\ell)} \;=\; \mathcal{O}$$

in the strong (hence distributional) sense. The same holds for correlators: for any states ρ with the clustering/locality hypotheses,

$$\lim_{\lambda,\ell} \mathrm{Tr}\Big(\rho\, K_{\mathrm{IR}}^{(\ell)} P_x(\lambda)\, \mathcal{O}_1 \cdots \mathcal{O}_n\, P_x(\lambda) K_{\mathrm{IR}}^{(\ell)}\Big) = \mathrm{Tr}(\rho\, \mathcal{O}_1 \cdots \mathcal{O}_n)$$

by dominated convergence (using the L^p–contractivity of $K_{\mathrm{IR}}^{(\ell)}$ and $\|P_x(\lambda)\| \le 1$).

2) Monotonicity and idempotence of $P_x(\lambda)$. From the projector–valued measure $E^{(\mathcal{D}_x)}(\cdot)$,

$$P_x(\lambda) = E^{(\mathcal{D}_x)}([0,\lambda]), \quad P_x(\lambda)P_x(\lambda') = E([0,\lambda]\cap[0,\lambda']) = P_x(\min\{\lambda,\lambda'\}),$$

hence $P_x(\lambda)^2 = P_x(\lambda)$ and $P_x(\lambda) \le P_x(\lambda')$ when $\lambda \le \lambda'$ (monotone increasing). Thus decreasing λ_{UV} tightens the UV cut.

3) Smoothing/clustering increases with ℓ (fixed λ_{UV}). Take, for definiteness, a heat–kernel realization $K_{\mathrm{IR}}^{(\ell)} = e^{-t(\ell)\mathcal{D}}$ with $t(\ell)$ increasing in ℓ; then $\{e^{-t\mathcal{D}}\}_{t\ge 0}$ is a contractive, positivity–preserving semigroup on L^p, and $t\uparrow$ increases smoothing (higher Sobolev regularity, shorter correlation length of windowed fields). In the general bounded functional–calculus setting $K_{\mathrm{IR}}^{(\ell)} = \eta_\ell(\mathcal{D})$ with η_ℓ increasingly low–pass in ℓ, the operator norm of high–frequency spectral slices decreases with ℓ:

$$\big\|\chi_{[\Lambda,\infty)}(\mathcal{D})\, K_{\mathrm{IR}}^{(\ell)}\big\| \le \sup_{s\ge\Lambda} |\eta_\ell(s)| \;\downarrow\; 0 \quad (\ell\uparrow\infty,\ \Lambda \text{ fixed}),$$

which expresses strengthened smoothing (and, at the level of correlation functions, enhanced clustering after coarse–graining).

4) Log–derivative diagnostics and plateau criterion. Let $\Phi(\lambda,\ell)$ be any tagged expectation/visibility built from

$$\mathcal{K}(\lambda,\ell) := K_{\mathrm{IR}}^{(\ell)} P_x(\lambda)\, \mathcal{O}\, P_x(\lambda) K_{\mathrm{IR}}^{(\ell)} \quad \text{or its channel–resolved variants.}$$

[7] If one uses the alternative convention where $K_{\mathrm{IR}}^{(\ell)} \to \mathrm{Id}$ as $\ell \downarrow 0$ (standard for compactly supported kernels), simply replace "$\ell \to \infty$ (with $K_{\mathrm{IR}}^{(\ell)} \to \delta$)" by "$\ell \downarrow 0$"; the proof is identical.

Define $D_\lambda = \lambda\partial_\lambda$, $D_\ell = \ell\partial_\ell$. By the product rule and spectral calculus,

$$D_\lambda\mathcal{K} = \lambda(\partial_\lambda P_x)\,\mathcal{O}\,P_x\,K + \lambda P_x\,\mathcal{O}\,(\partial_\lambda P_x)\,K, \qquad D_\ell\mathcal{K} = (\ell\partial_\ell K)\,P_x\,\mathcal{O}\,P_x + P_x\,\mathcal{O}\,P_x\,(\ell\partial_\ell K),$$

where $K = K_{\mathrm{IR}}^{(\ell)}$ and $\partial_\lambda P_x$ is the spectral *density* in the sense of projector–valued measures: $\partial_\lambda P_x = \frac{d}{d\lambda}E^{(\mathcal{D}_x)}([0,\lambda])$ (a measure supported on the spectrum). Taking operator norms,

$$\|D_\lambda\mathcal{K}\| \;\le\; 2\,\|\,\lambda\,\partial_\lambda P_x\,\|\,\|\mathcal{O}\|\,\|K\|, \qquad \|D_\ell\mathcal{K}\| \;\le\; 2\,\|\,\ell\,\partial_\ell K\,\|\,\|\mathcal{O}\|\,\|P_x\|^2.$$

In the bounded functional–calculus realization $K = \eta_\ell(\mathcal{D})$,

$$\|\,\ell\,\partial_\ell K\,\| = \sup_{s\ge 0}\big|\,\ell\,\partial_\ell\eta_\ell(s)\,\big|,$$

so smallness of the multiplier slopes on a band implies small D_ℓ. Likewise, for any $\delta > 0$,

$$\big\|\lambda\,\partial_\lambda P_x\big\| \;\le\; \frac{1}{\log(1+\delta)}\,\big\|P_x((1+\delta)\lambda) - P_x(\lambda)\big\| \;\le\; \frac{1}{\log(1+\delta)},$$

in the sense of difference–quotients for the projector–valued measure; hence bounding the spectral *shell* response controls D_λ. Therefore, on any band where $\|\,\ell\partial_\ell K\,\|$ and the projector shell variation $\|P_x((1+\delta)\lambda) - P_x(\lambda)\|$ are both small, the log–derivatives $D_\lambda\Phi$, $D_\ell\Phi$ (and, by iterating the argument, higher log–derivatives) are small. This is precisely the "flatness'' requirement in Section 6.12.4 and provides a quantitative way to assess/verify plateaus.

Conclusion. (i) As $\lambda_{\mathrm{UV}} \to \infty$ and along the approximation–of–identity limit of ℓ with $K_{\mathrm{IR}}^{(\ell)} \to \delta$ (kernel sense), the tagged objects converge to bare fields/correlators (strong/distributional limits). (ii) For fixed ℓ, $P_x(\lambda)$ is increasing and idempotent, so decreasing λ_{UV} strengthens the UV cut. (iii) For fixed λ_{UV}, increasing ℓ strengthens smoothing/clustering. (iv) The plateau criterion is evaluated by bounding the log–derivatives D_λ, D_ℓ (and higher ones), which reduce to operator–multiplier bounds supplied by the spectral calculus and the mollifier family. This proves the proposition. □

Intuitive reading. $K_{\mathrm{IR}}^{(\ell)}$ is a covariant blur: it averages fields over a geodesic ball of size ℓ while respecting indices and causality. Composed with the local spectral projector $P_x(\lambda_{\mathrm{UV}})$, it yields scale-tagged fields and correlators that live on a controlled band of UV modes and a controlled IR window. Taking $\lambda_{\mathrm{UV}} \to \infty$ or $\ell \to \infty$ removes the corresponding control; tightening either knob increases UV or IR regularity. Using the log-derivatives D_λ, D_ℓ lets us certify plateaus—regions where observables are insensitive to the knobs—so claims are not cutoff artifacts but genuine, scale-stable features.

Narrative (UV precedents). The projector $P_x(\lambda_{\mathrm{UV}})$ encodes the *Spectral Smoothness Axiom* and *Finite Entanglement Capacity* (Axiom 4.415 and Principle 4.1), yielding an *in situ* UV ceiling without external regulators (Proposition 4.424). All operator-valued tensors are thus defined *on the band*, where strong-operator-topology variations are safe (Definition 5.1). At this layer the geometry is triadic and noncommutative, $\hat{g}_{\mu\nu} = \hat{g}^{\mathrm{corr}}_{\mu\nu} + \hat{g}^{\mathrm{vac}}_{\mu\nu} + \hat{g}^{\mathrm{spec}}_{\mu\nu}$; positivity holds on the bandlimited algebra, while expectation-level channel additivity is enforced in the IR (Corollary 5.58 and Definition 5.10). In our variational pipeline, $P_x(\lambda_{\mathrm{UV}})$ is the UV knob that interfaces directly with Section 6.12.2.

BRST overlay (UV). We take $P_x(\lambda_{\mathrm{UV}})$ to be BRST-closed of ghost number 0, i.e. $[\,\hat{Q}_{\mathrm{BRST}}, P_x(\lambda_{\mathrm{UV}})\,] = 0$, and commuting with the channel projectors Π_α. Consequently, bandlimiting descends to the physical algebra $\mathcal{A}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$, and all scale-tagged observables built with P_x are gauge-parameter independent and invariant under ζ-exact improvements (by Proposition 6.264).

IR window (coarse grain).

Definition 6.289 (IR coarse graining by compactly supported kernel)**.** Let $K_{\mathrm{IR}}^{(\ell)} : \mathbb{R}^d \to \mathbb{R}_{\geq 0}$ be a non-negative, compactly supported kernel with characteristic width $\ell > 0$,

$$K_{\mathrm{IR}}^{(\ell)}(x) \geq 0, \qquad \int_{\mathbb{R}^d} \mathrm{d}^d x\, K_{\mathrm{IR}}^{(\ell)}(x) = 1, \qquad \operatorname{supp} K_{\mathrm{IR}}^{(\ell)} \subset B_\ell(0). \tag{6.370}$$

(For curved backgrounds, interpret the integrals in normal coordinates or insert parallel transport as in Definition 6.284.) Define coarse-grained observables (in local coordinates)

$$\bar{\Sigma}_{\mu\nu}^{(\ell)}(x) = \int_{\mathbb{R}^d} \mathrm{d}^d y\, K_{\mathrm{IR}}^{(\ell)}(x-y)\, \bar{\Sigma}_{\mu\nu}(y), \qquad \mathcal{O}^{(\ell)}(x) = \int_{\mathbb{R}^d} \mathrm{d}^d y\, K_{\mathrm{IR}}^{(\ell)}(x-y) \left\langle \hat{\mathcal{O}}(y) \right\rangle. \tag{6.371}$$

Proposition 6.290 (Mollifier properties and consistency)**.** *The map $f \mapsto K_{\mathrm{IR}}^{(\ell)} * f$ is positivity-preserving and contractive on $L^p(\mathbb{R}^d)$, $1 \leq p \leq \infty$, and forms an approximate identity: $K_{\mathrm{IR}}^{(\ell)} * f \to f$ in L^p as $\ell \downarrow 0$; moreover $K_{\mathrm{IR}}^{(\ell)} * f$ gains regularity as $\ell \uparrow \infty$ (standard mollifier calculus; [433]). If $\mathcal{O}$ obeys the clustering assumptions of Principle 4.5, then $\mathcal{O}^{(\ell)}$ retains cluster decay with an IR scale set by ℓ, and cone preservation holds when combined with causal transport (cf. Proposition 6.281, [230]).*

Proof of Proposition 6.290. **Set–up.** Fix a standard mollifier $K \in C_c^\infty(\mathbb{R}^d)$ with $K \geq 0$ and $\int_{\mathbb{R}^d} K = 1$. For $\ell > 0$ define the rescaled kernel

$$K_{\mathrm{IR}}^{(\ell)}(x) := \ell^{-d} K(x/\ell), \qquad (K_{\mathrm{IR}}^{(\ell)} * f)(x) := \int_{\mathbb{R}^d} K_{\mathrm{IR}}^{(\ell)}(x-y)\, f(y)\, dy.$$

(Everything below holds as well when K is only L^1 with unit mass and sufficient decay; the C_c^∞ choice highlights regularization.)

(A) Positivity, L^p–contractivity, approximate identity, smoothing. If $f \geq 0$ pointwise, then $K_{\mathrm{IR}}^{(\ell)} * f \geq 0$ since $K_{\mathrm{IR}}^{(\ell)} \geq 0$. For $1 \leq p \leq \infty$, Young's inequality gives

$$\|K_{\mathrm{IR}}^{(\ell)} * f\|_{L^p} \;\leq\; \|K_{\mathrm{IR}}^{(\ell)}\|_{L^1}\, \|f\|_{L^p} \;=\; \|f\|_{L^p},$$

so the map is a contraction on every L^p. Standard mollifier calculus (Rudin, §7) implies $K_{\mathrm{IR}}^{(\ell)} \to \delta$ in the sense of distributions as $\ell \downarrow 0$, hence

$$K_{\mathrm{IR}}^{(\ell)} * f \;\to\; f \quad \text{in } L^p(\mathbb{R}^d) \;\; (1 \leq p < \infty),$$

and locally uniformly if f is continuous. Moreover $K_{\mathrm{IR}}^{(\ell)} * f \in C^\infty$ and

$$\partial^\alpha \big(K_{\mathrm{IR}}^{(\ell)} * f\big) = \big(\partial^\alpha K_{\mathrm{IR}}^{(\ell)}\big) * f, \qquad \big\|\partial^\alpha \big(K_{\mathrm{IR}}^{(\ell)} * f\big)\big\|_{L^p} \;\leq\; \|\partial^\alpha K_{\mathrm{IR}}^{(\ell)}\|_{L^1}\, \|f\|_{L^p}.$$

Since $\|\partial^\alpha K_{\mathrm{IR}}^{(\ell)}\|_{L^1} = \ell^{-|\alpha|}\|\partial^\alpha K\|_{L^1}$, the derivative norms decay like $\ell^{-|\alpha|}$ as $\ell \uparrow \infty$, i.e. the convolution gains regularity when the window is widened.

(B) Preservation of clustering with IR scale set by ℓ. Assume $\mathcal{O}$ obeys the clustering hypothesis of Principle 4.5, i.e. there is a nonincreasing function $\phi : [0,\infty) \to [0,\infty)$ with $\phi(r) \to 0$ as $r \to \infty$ and a constant $C_{\mathcal{O}}$ such that the connected correlator obeys

$$\big|\langle \mathcal{O}(x)\,\mathcal{O}(y)\rangle_c\big| \;\le\; C_{\mathcal{O}}\,\phi(|x-y|) \qquad \forall x, y \in \mathbb{R}^d.$$

Define the coarse–grained observable $\mathcal{O}^{(\ell)} := K_{\mathrm{IR}}^{(\ell)} * \mathcal{O}$. Then

$$\begin{aligned}\big|\langle \mathcal{O}^{(\ell)}(x)\,\mathcal{O}^{(\ell)}(y)\rangle_c\big| &= \Big|\iint K_{\mathrm{IR}}^{(\ell)}(x-\xi)\,K_{\mathrm{IR}}^{(\ell)}(y-\eta)\,\langle \mathcal{O}(\xi)\,\mathcal{O}(\eta)\rangle_c\,d\xi\,d\eta\Big| \\ &\le C_{\mathcal{O}}\iint K_{\mathrm{IR}}^{(\ell)}(x-\xi)\,K_{\mathrm{IR}}^{(\ell)}(y-\eta)\,\phi(|\xi-\eta|)\,d\xi\,d\eta.\end{aligned}$$

If K has compact support in $B(0,c)$, then $K_{\mathrm{IR}}^{(\ell)}$ is supported in $B(0,c\ell)$, so $|x-\xi| \le c\ell$ and $|y-\eta| \le c\ell$ whenever the integrand is nonzero. By the triangle inequality,

$$|\xi-\eta| \;\ge\; |x-y| - |x-\xi| - |y-\eta| \;\ge\; |x-y| - 2c\ell,$$

whence, using that ϕ is nonincreasing and the kernels have unit mass,

$$\big|\langle \mathcal{O}^{(\ell)}(x)\,\mathcal{O}^{(\ell)}(y)\rangle_c\big| \;\le\; C_{\mathcal{O}}\,\phi\Big(\max\{|x-y| - 2c\ell,\,0\}\Big).$$

Thus the cluster decay is retained with an IR scale set by ℓ (e.g. exponential $\phi(r) = e^{-r/\xi}$ remains exponential with an effective range $\xi_{\mathrm{eff}} = \xi + O(\ell)$). If K has fast decay instead of compact support, the same bound holds up to negligible tails by splitting the integrals into the set $\{|x-\xi|, |y-\eta| \le c\ell\}$ and its complement.

(C) Cone preservation with causal transport. When the coarse–graining is applied to tensor fields (e.g. the expectation chronogeometric tensor $\bar{\Sigma}_{\mu\nu}$), incorporate causal parallel transport $\mathsf{P}_{y\to x}$ and average transported tensors (as in Proposition 6.281):

$$\bar{\Sigma}^{(\ell)}(x) = \int K_{\mathrm{IR}}^{(\ell)}(x-y)\,\mathsf{P}_{y\to x}^{*}\,\bar{\Sigma}(y)\,\mathsf{P}_{y\to x}\,dy.$$

If $\mathsf{P}_{y\to x}$ is causal with respect to $\bar{\Sigma}_{\mu\nu}$ and $\operatorname{supp} K_{\mathrm{IR}}^{(\ell)} \subset B(0,c\ell)$, the quadratic form associated with $\bar{\Sigma}^{(\ell)}(x)$ on any covector $k \in T_x^*\mathbb{R}^d$ is a positive average of quadratic forms that are strictly positive whenever k is spacelike at x (because their pullbacks $\mathsf{P}_{x\to y}^{*}k$ remain spacelike at y). Hence

$$k \notin \mathcal{C}_x^* \;\Longrightarrow\; k \notin \mathcal{C}_x^{(\ell)\,*}, \qquad \text{i.e.} \quad \mathcal{C}_x^{(\ell)\,*} \subseteq \mathcal{C}_x^*,$$

so effective cones are not enlarged by the IR window; this is exactly the cone–preservation statement (cf. Proposition 6.281 and [230]).

Conclusion. The convolution $K_{\mathrm{IR}}^{(\ell)} * f$ is positivity–preserving, L^p–contractive, an approximate identity as $\ell \downarrow 0$, and it gains regularity as $\ell \uparrow \infty$. Under the clustering hypothesis, coarse–grained observables inherit cluster decay with an IR scale controlled by ℓ. With causal transport, windowed tensors preserve (and never enlarge) causal cones. This proves the proposition. □

Proposition 6.291 (IR precedents and reciprocal resolution)**.** *Finite-support windowing realises* Relational Windowing *(Principle 4.4); together with* Relational Cluster Decomposition *(Principle 4.5) it implies the* IR Finiteness Theorem *for the channel distances* $d_{\mathrm{corr}}, d_{\mathrm{vac}}, d_{\mathrm{spec}}$ *(Theorem 4.433). The operational visibility of soft modes is governed by the* IR coherence scale $\lambda_{\mathrm{IR}}(\cdot;\varepsilon)$ *(Definition 4.435), which is monotonically decreasing in* ℓ*, thereby realising the reciprocal-resolution limits of Proposition 4.436 and Remark 4.437. Holosymmetric probes preserve causal cones under the coarse graining in* (6.371) *(no law-signalling) by Propositions 5.157 and 5.159 and Corollary 5.158.*

Proof of Proposition 6.291. **1) Finite–support windowing ⇒ Relational Windowing, and IR finiteness of channel distances.** Fix an IR kernel $K_{\mathrm{IR}}^{(\ell)}(x,y) \geq 0$ of unit mass in y and *finite support* $\operatorname{supp} K_{\mathrm{IR}}^{(\ell)}(x,\cdot) \subset B_{\bar{\Sigma}}(x; c\,\ell)$ (Principle 4.4). For any quasi–local observable $\mathcal{O}$ define its coarse–grained version (channel–wise if needed) by transporting to x and averaging on $B_{\bar{\Sigma}}(x; c\,\ell)$:

$$\mathcal{O}^{(\ell)}(x) \;=\; \int K_{\mathrm{IR}}^{(\ell)}(x,y)\, \mathsf{P}_{y\to x}\, \mathcal{O}(y)\, d\mathrm{vol}(y),$$

with $\mathsf{P}_{y\to x}$ the $\bar{\Sigma}$–causal parallel transport. By Principle 4.5, the connected correlators obey $\big|\langle \mathcal{O}(y)\, \mathcal{O}(y')\rangle_c\big| \leq C_{\mathcal{O}}\, \phi(|y-y'|)$ for some nonincreasing $\phi(r) \downarrow 0$ as $r \to \infty$. Consequently, for x, x',

$$\begin{aligned}\Big|\langle \mathcal{O}^{(\ell)}(x)\, \mathcal{O}^{(\ell)}(x')\rangle_c\Big| &\leq C_{\mathcal{O}} \iint K_{\mathrm{IR}}^{(\ell)}(x,y)\, K_{\mathrm{IR}}^{(\ell)}(x',y')\, \phi(|y-y'|)\, dy dy' \\ &\leq C_{\mathcal{O}}\, \phi\Big(\max\{|x-x'| - 2c\ell,\, 0\}\Big),\end{aligned}$$

using $|x-y| \leq c\ell$, $|x'-y'| \leq c\ell$ on the support and the monotonicity of ϕ. Thus coarse–grained connected correlators retain cluster decay with an IR range controlled by ℓ.

The channel distances $d_{\mathrm{corr}}, d_{\mathrm{vac}}, d_{\mathrm{spec}}$ (as defined in Theorem 4.433) are built from such connected objects and/or norms of differences of channel–restricted kernels against test operators of unit norm. The estimate above immediately yields finite bounds uniform on bounded ℓ–bands:

$$d_{\alpha}^{(\ell)}(x,x') \;\leq\; C_{\alpha}\, \phi\Big(\max\{|x-x'| - 2c\ell,\, 0\}\Big), \qquad \alpha \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\},$$

which proves the *IR Finiteness Theorem* under Principles 4.4 and 4.5.

2) IR coherence scale and reciprocal–resolution monotonicity. Let $\lambda_{\mathrm{IR}}(\cdot;\varepsilon)$ be the IR coherence scale of Definition 4.435; concretely, it is defined so that for spectral parameters $\lambda \geq \lambda_{\mathrm{IR}}(\cdot;\varepsilon)$ the variation of the chosen soft–mode diagnostic (e.g. a channel visibility or its kernel) across a fixed logarithmic shell is $\leq \varepsilon$. Writing the IR tag as a bounded functional calculus of a positive operator, $K_{\mathrm{IR}}^{(\ell)} = \eta_{\ell}(\mathcal{D})$ with η_{ℓ} increasingly low–pass in ℓ (Proposition 6.281), one has the operator monotonicity

$$\ell_2 \geq \ell_1 \quad\Longrightarrow\quad \big\|\chi_{[\Lambda,\infty)}(\mathcal{D})\, K_{\mathrm{IR}}^{(\ell_2)}\big\| \;\leq\; \big\|\chi_{[\Lambda,\infty)}(\mathcal{D})\, K_{\mathrm{IR}}^{(\ell_1)}\big\| \qquad (\Lambda > 0).$$

Hence the spectral *shell response* of any tagged observable decreases with ℓ, and so the minimal λ required to achieve a fixed tolerance ε cannot increase when ℓ grows:

$$\ell_2 \geq \ell_1 \quad\Longrightarrow\quad \lambda_{\mathrm{IR}}(\cdot;\varepsilon;\ell_2) \;\leq\; \lambda_{\mathrm{IR}}(\cdot;\varepsilon;\ell_1).$$

This proves that $\lambda_{\mathrm{IR}}(\cdot;\varepsilon)$ is monotonically *decreasing* in ℓ.

By Proposition 4.436 and Remark 4.437, fixing ℓ imposes a minimal UV bandwidth $\lambda_{\min}(\ell) \asymp C/\ell$ to avoid aliasing of the $B_{\bar{\Sigma}}(x; c\ell)$ average; together, these statements realise the reciprocal–resolution limits operationally: increasing ℓ (more aggressive coarse–graining) lowers the IR coherence threshold and, conversely, a fixed ℓ requires $\lambda_{\mathrm{UV}} \geq \lambda_{\min}(\ell)$.

3) Cone preservation and "no law–signalling". Coarse–graining of tensor probes is performed by transporting to x and averaging on $B_{\bar{\Sigma}}(x; c\ell)$:

$$T^{(\ell)}(x) = \int K_{\mathrm{IR}}^{(\ell)}(x,y)\, \mathsf{P}^{*}_{y\to x}\, T(y)\, \mathsf{P}_{y\to x}\, d\mathrm{vol}(y).$$

If $\mathsf{P}_{y\to x}$ is causal and $K_{\mathrm{IR}}^{(\ell)}$ has finite support as above, then the associated quadratic forms on covectors are positive averages of forms that are strictly positive on *spacelike* covectors at x; hence the effective covector cones cannot be enlarged: $\mathcal{C}_x^{(\ell)*} \subseteq \mathcal{C}_x^{*}$ (Proposition 6.281; see also Propositions 5.157 and 5.159 and Corollary 5.158). Since holosymmetric probes are order–≤ 0 (principal symbol preserved), causal ordering is left unchanged—there is no "law–signalling'' generated by the coarse–graining.

Conclusion. Finite–support IR windowing implements Relational Windowing and, combined with Relational Cluster Decomposition, yields IR–finite channel distances as stated in Theorem 4.433. The IR coherence scale governing soft–mode visibility is monotone decreasing in ℓ, matching the reciprocal–resolution limits, and holosymmetric probes preserve causal cones under (6.371). This proves the proposition. □

BRST overlay (IR).

Lemma 6.292 (BRST-closed smearing and physical invariance)**.** *When used at the operator level (e.g. smearing currents), choose the smearing map induced by $K_{\mathrm{IR}}^{(\ell)}$ to be BRST-closed of ghost number 0 and to commute with each channel projector Π_α (corr/vac/spec). When used at the expectation level in (6.371), BRST-closedness is automatic. Hence coarse graining preserves the physical sector $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ and the channel decomposition, and the extracted plateaus/exponents from $\Lambda_{\mathrm{law}}(\ell)$, $\Lambda_{\mathrm{state}}(V)$ and their channel-resolved versions are invariant under ζ-exact gauge-fixing changes (by Proposition 6.264; see also [117]).*

Proof of Lemma 6.292. **Set–up.** Let $\hat{Q}_{\mathrm{BRST}}$ be nilpotent and define the BRST differential $\zeta(\cdot) := \{\hat{Q}_{\mathrm{BRST}}, \cdot\}$. Assume $K_{\mathrm{IR}}^{(\ell)}$ is BRST–closed and ghost number 0: $[\hat{Q}_{\mathrm{BRST}}, K_{\mathrm{IR}}^{(\ell)}] = 0$, $\mathrm{gh}(K_{\mathrm{IR}}^{(\ell)}) = 0$, and likewise for the channel projectors Π_α (corr/vac/spec): $[\hat{Q}_{\mathrm{BRST}}, \Pi_\alpha] = 0$, $\mathrm{gh}(\Pi_\alpha) = 0$. We use the holosymmetric two–sided smearing map at the *operator level*

$$\mathsf{S}_\ell : \ \mathcal{O} \ \longmapsto \ \mathcal{O}^{(\ell)} := K_{\mathrm{IR}}^{(\ell)}\, \mathcal{O}\, K_{\mathrm{IR}}^{(\ell)},$$

(or, equivalently, the transported integral form when acting on local densities; cf. (6.359)–(6.360)).

1) BRST closure and chain–map property at the operator level. Using $[\hat{Q}_{\mathrm{BRST}}, K_{\mathrm{IR}}^{(\ell)}] = 0$ and the graded Leibniz rule,

$$\begin{aligned}\zeta\big(\mathcal{O}^{(\ell)}\big) &= \zeta\big(K_{\mathrm{IR}}^{(\ell)} \mathcal{O} K_{\mathrm{IR}}^{(\ell)}\big) = \underbrace{\zeta K_{\mathrm{IR}}^{(\ell)}}_{=0}\, \mathcal{O} K_{\mathrm{IR}}^{(\ell)} + K_{\mathrm{IR}}^{(\ell)}\, \zeta\mathcal{O}\, K_{\mathrm{IR}}^{(\ell)} + (-1)^{\mathrm{gh}(\mathcal{O})} K_{\mathrm{IR}}^{(\ell)}\, \mathcal{O}\, \underbrace{\zeta K_{\mathrm{IR}}^{(\ell)}}_{=0} \\ &= K_{\mathrm{IR}}^{(\ell)}\, \zeta\mathcal{O}\, K_{\mathrm{IR}}^{(\ell)} = \mathsf{S}_\ell\big(\zeta\mathcal{O}\big).\end{aligned}$$

Thus S_ℓ *commutes with* ζ: it is a chain map. Consequences: (i) if $\mathcal{O}$ is BRST–closed, then $\mathcal{O}^{(\ell)}$ is BRST–closed; (ii) if $\mathcal{O} = \zeta B$ is BRST–exact, then $\mathcal{O}^{(\ell)} = \zeta\big(\mathsf{S}_\ell B\big)$ is also exact. Hence S_ℓ preserves $\ker \hat{Q}_{\mathrm{BRST}}$ and $\mathrm{im}\, \hat{Q}_{\mathrm{BRST}}$ and descends to a well–defined endomorphism on $H^0(\hat{Q}_{\mathrm{BRST}})$.

2) Channel compatibility. Assume $[\Pi_\alpha, K_{\mathrm{IR}}^{(\ell)}] = 0$ for each α. Then, for any operator $\mathcal{O}$,

$$\Pi_\alpha\, \mathcal{O}^{(\ell)}\, \Pi_\alpha = \Pi_\alpha K_{\mathrm{IR}}^{(\ell)}\, \mathcal{O}\, K_{\mathrm{IR}}^{(\ell)} \Pi_\alpha = K_{\mathrm{IR}}^{(\ell)}\, \Pi_\alpha\, \mathcal{O}\, \Pi_\alpha\, K_{\mathrm{IR}}^{(\ell)} = \big(\Pi_\alpha\, \mathcal{O}\, \Pi_\alpha\big)^{(\ell)}.$$

Thus smearing does not mix channels; it preserves the corr/vac/spec decomposition (and by $[\hat{Q}_{\mathrm{BRST}}, \Pi_\alpha] = 0$ this preservation holds already on cohomology).

3) Expectation–level smearing is automatically BRST–inert. In (6.371) the smearing is applied to *expectation values* (e.g. c–number fields/tensors such as $\bar{\Sigma}_{\mu\nu}(x)$):

$$F^{(\ell)}(x) = \int K^{(\ell)}(x,y)\, \mathsf{P}_{y\to x} F(y)\, d\mathrm{vol}(y).$$

These are numbers/tensors, not operators; the BRST differential acts trivially on them. Hence "BRST–closedness is automatic'' at the expectation level: no further hypothesis is needed.

4) Preservation of the physical sector and gauge–parameter invariance. Let $\mathsf{P}_{\mathrm{phys}}$ project onto $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$. From Step 1, $[\hat{Q}_{\mathrm{BRST}}, K_{\mathrm{IR}}^{(\ell)}] = 0$ implies $[\,\Delta_{\mathrm{BRST}}, K_{\mathrm{IR}}^{(\ell)}\,] = 0$ and hence $[\,\mathsf{P}_{\mathrm{phys}}, K_{\mathrm{IR}}^{(\ell)}\,] = 0$ by functional calculus. Therefore smearing commutes with projection to the physical sector; in particular,

$$\mathrm{Tr}\Big(\mathsf{P}_{\mathrm{phys}}\, \mathcal{O}^{(\ell)}\, \rho\Big) = \mathrm{Tr}\Big(\mathcal{O}^{(\ell)}\, \mathsf{P}_{\mathrm{phys}} \rho\, \mathsf{P}_{\mathrm{phys}}\Big),$$

and BRST–exact insertions continue to have vanishing expectation values. By Proposition 6.264, visibilities and their channel–resolved versions built from BRST–closed, scale–tagged kernels (which include $K_{\mathrm{IR}}^{(\ell)}$ and Π_α) satisfy the Nielsen identity in $\mathcal{H}_{\mathrm{phys}}$: $\partial_\xi(\cdot) = \langle \zeta\, \eta_\xi \rangle_{\mathcal{H}_{\mathrm{phys}}} = 0$. Hence the coarse–grained readouts (including the extracted plateaus/-exponents from $\Lambda_{\mathrm{law}}(\ell)$, $\Lambda_{\mathrm{state}}(V)$ and their channel restrictions) are invariant under ζ–exact gauge–fixing deformations.

Conclusion. Choosing the smearing map induced by $K_{\mathrm{IR}}^{(\ell)}$ to be BRST–closed, ghost number 0, and channel–commuting makes it a chain map on the BRST complex and preserves the channel decomposition. Expectation–level smearing is automatically BRST–inert. Consequently, coarse graining preserves $\mathcal{H}_{\mathrm{phys}}$ and the corr/vac/spec split, and by the Nielsen/cohomological identities the resulting visibilities (and their asymptotic plateaus/exponents) are invariant under ζ–exact gauge–fixing changes. □

Intuitive reading. $K_{\mathrm{IR}}^{(\ell)}$ is a compactly supported "blur" that averages over a ball of size ℓ, producing smooth, finite-window observables. As ℓ shrinks we recover the raw fields; as it grows we emphasise long-distance structure while preserving causal cones and BRST/physical content. Because ℓ trades off against the UV content (via reciprocal resolution), stable plateaus in ℓ certify that the conclusions are not artifacts of the coarse graining.

Scale-tagged actions and identities.

Definition 6.293 (Scale–tagged umbrella and sector actions)**.** Let $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ be the UV projector and IR window of Definitions 6.280 and 7.26, and let the scale–tagging functor be $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ from Definition 7.29. Denote the windowed inverse chronogeometric weights by $\bar{\Sigma}^{(\ell)\mu\nu}$ and volume element $\sqrt{-\det \bar{\Sigma}^{(\ell)}}$ (cf. Definition 6.280 and Proposition 6.281). The *scale–tagged umbrella action* is

$$S^{(\lambda_{\mathrm{UV}},\ell)} \;=\; \int \mathrm{d}^d x\, \sqrt{-\det \bar{\Sigma}^{(\ell)}(x)}\;\; \mathrm{Tr}\Big(\langle \boldsymbol{\Psi}|\, \hat{\mathbf{G}}^{(\lambda_{\mathrm{UV}},\ell)}(x,x)\, |\boldsymbol{\Psi}\rangle \Big), \tag{6.372}$$

where $\hat{\mathbf{G}}^{(\lambda_{\mathrm{UV}},\ell)} := \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\hat{\mathbf{G}}]$ (cf. eq. (6.366)). The *scale–tagged sector actions* for Gaiamiti and Tantrakyma fields (with pairings g_T, g_C and Riesz maps $\mathcal{G}_T, \mathcal{G}_C$) are

$$S_{\Phi}^{(\lambda_{\mathrm{UV}},\ell)} = \int \mathrm{d}^d x\, \sqrt{-\det \bar{\Sigma}^{(\ell)}(x)}\;\; \mathrm{Tr}\Big\{ \bar{\Sigma}^{(\ell)\mu\nu}(x)\, g_T\Big(\nabla_\mu \hat{\boldsymbol{\Phi}}_{\lambda_{\mathrm{UV}},\ell},\, \nabla_\nu \hat{\boldsymbol{\Phi}}_{\lambda_{\mathrm{UV}},\ell} \Big) + \mathcal{V}_\Phi(\hat{\boldsymbol{\Phi}}_{\lambda_{\mathrm{UV}},\ell}) \Big\}, \tag{6.373}$$

$$S_{\Theta}^{(\lambda_{\mathrm{UV}},\ell)} = \int \mathrm{d}^d x\, \sqrt{-\det \bar{\Sigma}^{(\ell)}(x)}\;\; \mathrm{Tr}\Big\{ \bar{\Sigma}^{(\ell)\mu\nu}(x)\, g_C\Big(\nabla_\mu \hat{\boldsymbol{\Theta}}_{\lambda_{\mathrm{UV}},\ell},\, \nabla_\nu \hat{\boldsymbol{\Theta}}_{\lambda_{\mathrm{UV}},\ell} \Big) + \mathcal{V}_\Theta(\hat{\boldsymbol{\Theta}}_{\lambda_{\mathrm{UV}},\ell}) \Big\},$$

with $\hat{\boldsymbol{\Phi}}_{\lambda_{\mathrm{UV}},\ell}$, $\hat{\boldsymbol{\Theta}}_{\lambda_{\mathrm{UV}},\ell}$ given by eq. (6.364). Here ∇_μ is the split connection of eq. (6.314); we assume $[\nabla_\mu, P_x(\lambda_{\mathrm{UV}})] = 0$ so that "UV then IR'' commutes with covariant differentiation (cf. Lemma 6.287).

Proposition 6.294 (Projected Euler–Lagrange operators)**.** *Varying eq. (6.373) at fixed $\bar{\Sigma}^{(\ell)}$ and using the "UV then IR'' convention yields the scale–tagged Euler–Lagrange operators*

$$\mathcal{E}_{\Phi}^{(\lambda_{\mathrm{UV}},\ell)} := P_x(\lambda_{\mathrm{UV}})\Big[-\nabla_\mu\Big(\bar{\Sigma}^{(\ell)\mu\nu}\, \frac{\partial \mathcal{L}_\Phi}{\partial\big(\nabla_\nu \hat{\boldsymbol{\Phi}}_{\lambda_{\mathrm{UV}},\ell}\big)} \Big) + \frac{\partial \mathcal{L}_\Phi}{\partial \hat{\boldsymbol{\Phi}}_{\lambda_{\mathrm{UV}},\ell}} \Big]\, P_x(\lambda_{\mathrm{UV}}) = 0, \tag{6.374}$$

and analogously for Θ with $\Phi \to \Theta$ and $g_T \to g_C$ (cf. eqs. (6.333) and (6.334)). Remark. *If $[\nabla_\mu, P_x(\lambda_{\mathrm{UV}})] \neq 0$, commutator correction terms of the form $[\,\nabla_\mu, P_x(\lambda_{\mathrm{UV}})\,]$ appear (cf. Lemma 6.287); when the norm of this commutator is small on the ℓ–scale, these terms are suppressed [253].*

Proof of Proposition 6.294. **Convention and variables.** We adopt the "UV then IR'' convention: the *field variable* entering the sector action is the UV–bandlimited field $\hat{\boldsymbol{\Phi}}_{\lambda_{\mathrm{UV}},\ell} := P_x(\lambda_{\mathrm{UV}})\, \hat{\boldsymbol{\Phi}}$ (the IR dependence only enters *parametrically* through the fixed background $\bar{\Sigma}^{(\ell)}$ and the $\bar{\Sigma}^{(\ell)}$–compatible covariant derivative). Thus the Φ–sector action (cf. (6.373)) reads, schematically,

$$S_{\Phi}^{(\lambda_{\mathrm{UV}},\ell)} = \int d\mathrm{vol}_{\bar{\Sigma}^{(\ell)}}\; \mathcal{L}_\Phi\Big(\hat{\boldsymbol{\Phi}}_{\lambda_{\mathrm{UV}},\ell}, \nabla \hat{\boldsymbol{\Phi}}_{\lambda_{\mathrm{UV}},\ell};\; \bar{\Sigma}^{(\ell)} \Big),$$

with $d\mathrm{vol}_{\bar{\Sigma}^{(\ell)}}$ the measure built from $\bar{\Sigma}^{(\ell)}$. Virtual variations are restricted to the UV band: $\delta\hat{\boldsymbol{\Phi}}_{\lambda_{\mathrm{UV}},\ell} = P_x(\lambda_{\mathrm{UV}})\, \delta\hat{\boldsymbol{\Phi}}$ (i.e. $\delta\hat{\boldsymbol{\Phi}}_{\lambda_{\mathrm{UV}},\ell}$ takes values in $\mathrm{Ran}\, P_x(\lambda_{\mathrm{UV}})$).

1) First variation and covariant integration by parts. Vary at fixed $\bar{\Sigma}^{(\ell)}$:

$$\delta S_{\Phi}^{(\lambda_{\mathrm{UV}},\ell)} = \int d\mathrm{vol}_{\bar{\Sigma}^{(\ell)}} \Big\langle \frac{\partial \mathcal{L}_\Phi}{\partial \hat{\boldsymbol{\Phi}}_{\lambda_{\mathrm{UV}},\ell}},\, \delta\hat{\boldsymbol{\Phi}}_{\lambda_{\mathrm{UV}},\ell} \Big\rangle + \int d\mathrm{vol}_{\bar{\Sigma}^{(\ell)}} \Big\langle \frac{\partial \mathcal{L}_\Phi}{\partial(\nabla_\nu \hat{\boldsymbol{\Phi}}_{\lambda_{\mathrm{UV}},\ell})},\, \nabla_\nu \delta\hat{\boldsymbol{\Phi}}_{\lambda_{\mathrm{UV}},\ell} \Big\rangle.$$

Using the $\bar{\Sigma}^{(\ell)}$–compatible divergence theorem and dropping boundary terms (variations of compact support or standard boundary conditions),

$$\int \Big\langle \frac{\partial \mathcal{L}_\Phi}{\partial(\nabla_\nu \hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell})},\, \nabla_\nu \delta\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell} \Big\rangle = -\int \Big\langle \nabla_\mu\Big(\bar{\Sigma}^{(\ell)\mu\nu}\, \frac{\partial \mathcal{L}_\Phi}{\partial(\nabla_\nu \hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell})}\Big),\, \delta\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell} \Big\rangle.$$

Hence

$$\delta S_\Phi^{(\lambda_{\mathrm{UV}},\ell)} = \int d\mathrm{vol}_{\bar{\Sigma}^{(\ell)}} \Big\langle \mathcal{E}_\Phi^{\mathrm{bare}}(\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell}; \bar{\Sigma}^{(\ell)}),\, \delta\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell} \Big\rangle,$$

with the usual (unprojected) Euler–Lagrange operator

$$\mathcal{E}_\Phi^{\mathrm{bare}} := -\nabla_\mu\Big(\bar{\Sigma}^{(\ell)\mu\nu}\, \frac{\partial \mathcal{L}_\Phi}{\partial(\nabla_\nu \hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell})}\Big) + \frac{\partial \mathcal{L}_\Phi}{\partial \hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell}}.$$

2) Restricting variations to the UV band $\mathrm{Ran}\, P_x(\lambda_{\mathrm{UV}})$**.** Since $\delta\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell} = P_x(\lambda_{\mathrm{UV}})\, \delta\hat{\mathbf{\Phi}}$ and $P_x(\lambda_{\mathrm{UV}})$ is an orthogonal projector,

$$\Big\langle \mathcal{E}_\Phi^{\mathrm{bare}},\, \delta\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell} \Big\rangle = \Big\langle P_x(\lambda_{\mathrm{UV}})\, \mathcal{E}_\Phi^{\mathrm{bare}}\, P_x(\lambda_{\mathrm{UV}}),\, \delta\hat{\mathbf{\Phi}} \Big\rangle.$$

Because $\delta\hat{\mathbf{\Phi}}$ is otherwise arbitrary (within the band), the stationarity condition $\delta S_\Phi^{(\lambda_{\mathrm{UV}},\ell)} = 0$ for all such variations is equivalent (in distribution) to

$$P_x(\lambda_{\mathrm{UV}})\, \mathcal{E}_\Phi^{\mathrm{bare}}\, P_x(\lambda_{\mathrm{UV}}) = 0,$$

which is exactly (6.374). The argument for Θ is identical (with $g_T \to g_C$).

3) Commutator corrections when $[\nabla_\mu, P_x(\lambda_{\mathrm{UV}})] \neq 0$**.** If the projector depends on position (as a spectral projector of $\mathcal{D}_x$), then

$$\nabla_\nu\Big(\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell}\Big) = \nabla_\nu\Big(P_x(\lambda_{\mathrm{UV}})\hat{\mathbf{\Phi}}\Big) = P_x(\lambda_{\mathrm{UV}})\, \nabla_\nu \hat{\mathbf{\Phi}} + [\nabla_\nu, P_x(\lambda_{\mathrm{UV}})]\, \hat{\mathbf{\Phi}}.$$

Inserting this into the definition of $\partial\mathcal{L}_\Phi/\partial(\nabla_\nu\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell})$ produces extra terms in $\mathcal{E}_\Phi^{\mathrm{bare}}$ proportional to $[\nabla_\nu, P_x(\lambda_{\mathrm{UV}})]$ (and their covariant divergences). Equivalently, after the integration–by–parts step, the E–L operator acquires additive *commutator corrections* built from $[\nabla_\mu, P_x(\lambda_{\mathrm{UV}})]$. By the spectral–calculus estimates (Reed–Simon, Vol. I) and the commutator bounds used in Lemma 6.287, these are controlled in norm by $\|[\nabla, P_x(\lambda_{\mathrm{UV}})]\|$, which is small whenever the projector field varies slowly on the IR scale (e.g. $O(\ell^{-1})$ suppression when P_x is fibrewise–Lipschitz and the window size ℓ is large compared to its variation length).

Conclusion. Varying the scale–tagged action at fixed $\bar{\Sigma}^{(\ell)}$ with band–restricted virtual variations yields the projected Euler–Lagrange equation $P_x(\lambda_{\mathrm{UV}})\, \mathcal{E}_\Phi^{\mathrm{bare}}\, P_x(\lambda_{\mathrm{UV}}) = 0$, i.e. (6.374). If $[\nabla, P_x(\lambda_{\mathrm{UV}})] \neq 0$, commutator corrections appear and are suppressed when the projector is fibrewise–smooth on the ℓ–scale, in agreement with Lemma 6.287. □

Definition 6.295 (Scale–dependent Ward identity (adjoint–covariant form))**.** Let $\mathcal{A}_\mu$ be the background gauge potential (if present) and define the adjoint–covariant divergence $\nabla_\mu^{(\mathrm{ad})} J^\mu := \nabla_\mu J^\mu + i[\mathcal{A}_\mu, J^\mu]$. For scale–tagged improved currents $J_a^{(\lambda_{\mathrm{UV}},\ell)\mu}$ one has

$$\nabla_\mu J_a^{(\lambda_{\mathrm{UV}},\ell)\mu} + i\,[\mathcal{A}_\mu, J_a^{(\lambda_{\mathrm{UV}},\ell)\mu}] = \Delta_a^{(\lambda_{\mathrm{UV}},\ell)}\,, \qquad \nabla_\mu^{(\mathrm{ad})} J_a^\mu := \nabla_\mu J_a^\mu + i[\mathcal{A}_\mu, J_a^\mu], \tag{6.375}$$

where, for $\mathcal{L} = \mathcal{L}_0 + \sum_k \epsilon_k \mathcal{O}_k + \zeta\Psi$ with $\delta_a \mathcal{L}_0 = \nabla_\mu K_a^\mu$ and $\delta_a \mathcal{O}_k = \mathcal{S}_{k,a}$,

$$\Delta_a^{(\lambda_{\mathrm{UV}},\ell)} = \sum_k \epsilon_k(\ell) \left\langle \mathcal{S}_{k,a}^{(\lambda_{\mathrm{UV}})} \right\rangle_{\mathbf{\Psi}} + \left\langle \zeta\, \eta_a^{(\lambda_{\mathrm{UV}},\ell)} \right\rangle_{\mathbf{\Psi}} + \mathcal{A}_a^{(\lambda_{\mathrm{UV}},\ell)}, \tag{6.376}$$

with η_a the local functional generated by $\delta_a\Psi$ and $\mathcal{A}_a$ a possible anomaly (cf. Lemma 6.233; see [82, 117, 230]).

Proposition 6.296 (Physical, anomaly–free reduction)**.** *On the physical sector $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$, if BRST is anomaly–free, then the last two terms in eq. (6.376) vanish and*

$$\nabla_\mu^{(\mathrm{ad})} \bar{J}_{a,\mathrm{imp}}^{(\lambda_{\mathrm{UV}},\ell)\mu} = \sum_k \epsilon_k(\ell) \left\langle \mathcal{S}_{k,a}^{(\lambda_{\mathrm{UV}})} \right\rangle_{\mathbf{\Psi}}, \qquad \bar{J}_{a,\mathrm{imp}}^{(\lambda_{\mathrm{UV}},\ell)\mu} := \langle \hat{J}_{a,\mathrm{imp}}^{(\lambda_{\mathrm{UV}},\ell)\mu} \rangle_{\mathbf{\Psi}}. \tag{6.377}$$

Moreover, by Propositions 6.264 and 6.276, all scale–tagged actions, currents, and their channel–resolved projections are invariant under ζ–exact changes of the gauge–fixing fermion (Nielsen–type independence) provided $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ are BRST–closed and commute with the channel projectors Π_α (cf. [117]).

Proof of Proposition 6.296. **Standing assumptions.** Work with anomaly–free BRST: the nilpotent charge $\hat{Q}_{\mathrm{BRST}}$ leaves the path–integral measure invariant and $\zeta(\cdot) := \{\hat{Q}_{\mathrm{BRST}}, \cdot\}$ squares to zero. The physical sector is $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$, selected by the orthogonal projector $\mathsf{P}_{\mathrm{phys}}$ (commuting with any BRST–closed tag). The scale tags are BRST–closed with $\mathfrak{g}(P_x(\lambda_{\mathrm{UV}})) = \mathfrak{g}(K_{\mathrm{IR}}^{(\ell)}) = 0$:

$$[\hat{Q}_{\mathrm{BRST}}, P_x(\lambda_{\mathrm{UV}})] = 0, \qquad [\hat{Q}_{\mathrm{BRST}}, K_{\mathrm{IR}}^{(\ell)}] = 0,$$

and (when used) channel projectors Π_α commute both with $\hat{Q}_{\mathrm{BRST}}$ and the tags. We adopt the "UV then IR" convention and keep $\bar{\Sigma}^{(\ell)}$ fixed.

1) Relational Ward identity before cohomological reduction. Let the gauge–fixed Lagrangian be $\mathcal{L} = \mathcal{L}_0 + \sum_k \epsilon_k \mathcal{O}_k + \zeta\Psi$, with global G–variation $\delta_a \mathcal{L}_0 = \nabla_\mu K_a^\mu$ and $\delta_a \mathcal{O}_k = \mathcal{S}_{k,a}$. The (improved) Noether current $\hat{J}_{a,\mathrm{imp}}^\mu$ obeys the standard BRST–aware Ward identity (operator or functional form)

$$\nabla_\mu^{(\mathrm{ad})} \hat{J}_{a,\mathrm{imp}}^\mu = \sum_k \epsilon_k \mathcal{S}_{k,a} + \zeta\eta_a + \mathcal{A}_a, \tag{6.378}$$

where $\zeta\eta_a$ is the gauge–fixing/BRST–exact contribution and $\mathcal{A}_a$ is the possible anomaly (Jacobian of the BRST change of variables). Here $\mathfrak{g}(\eta_a) = -1$, hence $\mathfrak{g}(\zeta\eta_a) = 0$, and $\mathfrak{g}(\mathcal{A}_a) = 0$.

2) Projection to $\mathcal{H}_{\mathrm{phys}}$ kills the last two terms. Take expectations with insertions that are BRST–closed and satisfy $\mathfrak{g}(\cdot) = 0$, and project onto the physical sector: $\langle\cdot\rangle_{\mathbf{\Psi},\mathcal{H}_{\mathrm{phys}}} := \mathrm{Tr}(\mathsf{P}_{\mathrm{phys}} \cdot \rho)$. Anomaly–freedom gives $\langle \mathcal{A}_a \rangle_{\mathcal{H}_{\mathrm{phys}}} = 0$. For the BRST–exact term,

$$\left\langle \zeta\eta_a \right\rangle_{\mathcal{H}_{\mathrm{phys}}} = \left\langle \zeta(\eta_a) \right\rangle_{\mathcal{H}_{\mathrm{phys}}} = 0,$$

because the measure is BRST–invariant and $\mathfrak{g}(\eta_a) = -1$ (hence $\mathfrak{g}(\zeta\eta_a) = 0$), and we are taking expectations on $\mathcal{H}_{\mathrm{phys}}$ where BRST–exact insertions vanish. Therefore, taking the expectation of (6.378) on the physical sector yields

$$\nabla_\mu^{(\mathrm{ad})} \bar{J}_{a,\mathrm{imp}}^\mu = \sum_k \epsilon_k \left\langle \mathcal{S}_{k,a} \right\rangle_{\mathbf{\Psi}} \qquad \text{in } \mathcal{H}_{\mathrm{phys}}, \tag{6.379}$$

with $\bar{J}^{\mu}_{a,\mathrm{imp}} := \langle \hat{J}^{\mu}_{a,\mathrm{imp}} \rangle_{\mathbf{\Psi}}$.

3) Inserting the scale tags. Define the tagged current and breaking insertions by the UV–IR sandwich (channel–wise if desired):

$$\hat{J}^{(\lambda_{\mathrm{UV}},\ell)\mu}_{a,\mathrm{imp}} := K^{(\ell)}_{\mathrm{IR}}\, P_x(\lambda_{\mathrm{UV}})\, \hat{J}^{\mu}_{a,\mathrm{imp}}\, P_x(\lambda_{\mathrm{UV}})\, K^{(\ell)}_{\mathrm{IR}}, \qquad \mathcal{S}^{(\lambda_{\mathrm{UV}})}_{k,a} := P_x(\lambda_{\mathrm{UV}})\, \mathcal{S}_{k,a}\, P_x(\lambda_{\mathrm{UV}}).$$

By BRST–closure of $K^{(\ell)}_{\mathrm{IR}}$ and $P_x(\lambda_{\mathrm{UV}})$, the tagged operators are BRST–closed whenever the untagged ones are; hence the argument of Step 2 applies verbatim after tagging. Taking expectations and denoting $\bar{J}^{(\lambda_{\mathrm{UV}},\ell)\mu}_{a,\mathrm{imp}} := \langle \hat{J}^{(\lambda_{\mathrm{UV}},\ell)\mu}_{a,\mathrm{imp}} \rangle_{\mathbf{\Psi}}$, we obtain the *scale–tagged* physical Ward identity

$$\nabla^{(\mathrm{ad})}_{\mu}\, \bar{J}^{(\lambda_{\mathrm{UV}},\ell)\mu}_{a,\mathrm{imp}} = \sum_{k} \epsilon_k(\ell) \left\langle \mathcal{S}^{(\lambda_{\mathrm{UV}})}_{k,a} \right\rangle_{\mathbf{\Psi}},$$

which is precisely (6.377).[8]

4) Gauge–parameter (gauge–fermion) independence. Let $\Psi(\xi)$ be a gauge–fermion family and denote $\partial_\xi S_{\mathrm{gf+gh}} = \zeta\, \eta_\xi$ (Nielsen identity), with $\mathfrak{g}(\eta_\xi) = -1$. Consider any tagged, $\mathfrak{g}(\cdot) = 0$, BRST–closed insertion $\mathcal{O}^{(\lambda_{\mathrm{UV}},\ell)}$ built from actions or currents (and optionally Π_α). Because the tags commute with $\hat{Q}_{\mathrm{BRST}}$ and with Π_α, $\mathcal{O}^{(\lambda_{\mathrm{UV}},\ell)}$ is BRST–closed whenever $\mathcal{O}$ is. Then

$$\partial_\xi \Big\langle \mathcal{O}^{(\lambda_{\mathrm{UV}},\ell)} \Big\rangle_{\mathcal{H}_{\mathrm{phys}}} = \Big\langle \zeta(\eta_\xi)\, \mathcal{O}^{(\lambda_{\mathrm{UV}},\ell)} \Big\rangle_{\mathcal{H}_{\mathrm{phys}}} = \Big\langle \zeta\Big(\eta_\xi\, \mathcal{O}^{(\lambda_{\mathrm{UV}},\ell)}\Big) \Big\rangle_{\mathcal{H}_{\mathrm{phys}}} - \underbrace{\Big\langle (\zeta \mathcal{O}^{(\lambda_{\mathrm{UV}},\ell)})\, \eta_\xi \Big\rangle_{\mathcal{H}_{\mathrm{phys}}}}_{=0} = 0,$$

using anomaly–freedom and BRST–closure of the insertion. The same reasoning applies to changes of gauge–fixing parameters and to channel–resolved projections (the commutation with Π_α prevents channel mixing). This is precisely the Nielsen/cohomological invariance asserted in Propositions 6.264 and 6.276.

Conclusion. On $\mathcal{H}_{\mathrm{phys}}$ and for BRST–closed tags (commuting with the channel projectors), the BRST–exact and anomalous pieces in the tagged Ward identity vanish, yielding (6.377). Moreover, all tagged actions/currents and their channel projections are invariant under ζ–exact gauge–fixing deformations (Nielsen–type independence). □

Narrative (variational precedents). Equations eq. (6.372)–eq. (6.375) are the *scale–tagged* lift of the relational umbrella and sector actions (Section 6.12.2) together with the Ward structure (Section 6.12.2). They combine the UV spectral projector $P_x(\lambda_{\mathrm{UV}})$ (Section 6.12.4) with the IR window $K^{(\ell)}_{\mathrm{IR}}$ (Section 6.12.4) to yield band–safe Euler–Lagrange operators and *adjoint–covariant, scale–dependent* currents. Both knobs are taken BRST–closed (ghost number 0) and commute with the channel projectors, so all scale–tagged objects descend to $\mathcal{H}_{\mathrm{phys}}$ and are gauge–parameter independent by Proposition 6.264 (anomaly–free Stage I) [117, 230]. Holosymmetric visibilities (Section 6.12.3) then read these equations *channel–wise* (corr/vac/spec), justified by orthogonality (Corollary 5.58) and functoriality, while causal consistency of the protocol follows from cone preservation (Propositions 5.157 and 5.159). The resulting expectation–level fields live on the smooth IR manifold (Definition 5.10), providing the bridge back to the common variational backbone (Section 6.12.2).

[8] If ∇_μ does not commute with the tags, the commutators can be absorbed into the definition of the *tagged* improved current (or treated as the small commutator corrections discussed after Proposition 6.294); this does not affect the cohomological content.

Plateau criterion (scale-stability).

Definition 6.297 (Plateau (scale-stable) region)**.** Let $\mathcal{O}(\lambda_{\mathrm{UV}}, \ell)$ be an observable obtained from the scale–tagging functor $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ (cf. Definition 7.29), with UV projector $P_x(\lambda_{\mathrm{UV}})$ and IR window $K_{\mathrm{IR}}^{(\ell)}$ as in Section 6.12.4. Fix tolerances $\varepsilon_{\lambda_{\mathrm{UV}}}, \varepsilon_\ell > 0$ and a rectangular scale box

$$\mathcal{B} \;:=\; [\lambda_{\mathrm{UV},1}, \lambda_{\mathrm{UV},2}] \times [\ell_1, \ell_2].$$

Write $u := \ln \lambda_{\mathrm{UV}}$, $v := \ln \ell$ and define the log–derivatives $D_\lambda := \partial_u$, $D_\ell := \partial_v$. We say that $\mathcal{O}$ is *plateau (scale–stable) on* $\mathcal{B}$ if

$$\sup_{(u,v)\in \ln \mathcal{B}} |D_\lambda \mathcal{O}(u,v)| \;\le\; \varepsilon_{\lambda_{\mathrm{UV}}}, \qquad \sup_{(u,v)\in \ln \mathcal{B}} |D_\ell \mathcal{O}(u,v)| \;\le\; \varepsilon_\ell, \tag{6.380}$$

where $\ln \mathcal{B} := [\ln \lambda_{\mathrm{UV},1}, \ln \lambda_{\mathrm{UV},2}] \times [\ln \ell_1, \ln \ell_2]$.

Proposition 6.298 (Oscillation bound and finite–difference certification)**.** *Assume $\mathcal{O}$ is C^1 on $\ln \mathcal{B}$. If (6.380) holds, then for any $(u,v), (u',v') \in \ln \mathcal{B}$,*

$$|\mathcal{O}(u',v') - \mathcal{O}(u,v)| \;\le\; \varepsilon_{\lambda_{\mathrm{UV}}} |u'-u| + \varepsilon_\ell |v'-v|.$$

Conversely, if $\mathcal{O}$ is C^2 and central finite differences on a grid of mesh sizes $\Delta u, \Delta v$ obey

$$\left|\frac{\mathcal{O}(u+\Delta u, v) - \mathcal{O}(u-\Delta u, v)}{2\Delta u}\right| \le \varepsilon_{\lambda_{\mathrm{UV}}} - \tfrac{1}{2}\, \|D_\lambda^2 \mathcal{O}\|_\infty\, \Delta u,$$
$$\left|\frac{\mathcal{O}(u, v+\Delta v) - \mathcal{O}(u, v-\Delta v)}{2\Delta v}\right| \le \varepsilon_\ell - \tfrac{1}{2}\, \|D_\ell^2 \mathcal{O}\|_\infty\, \Delta v,$$

uniformly on $\ln \mathcal{B}$, then (6.380) holds. (Mean–value theorem and Taylor remainder estimates; cf. [433].)

Proof of Proposition 6.298. Write

$$u := \ln \lambda_{\mathrm{UV}}, \qquad v := \ln \ell, \qquad D_\lambda = \partial_u, \quad D_\ell = \partial_v,$$

so the plateau criterion (6.380) reads

$$\| \partial_u \mathcal{O} \|_{L^\infty(\ln \mathcal{B})} \le \varepsilon_{\lambda_{\mathrm{UV}}}, \qquad \| \partial_v \mathcal{O} \|_{L^\infty(\ln \mathcal{B})} \le \varepsilon_\ell.$$

(1) Oscillation bound from the plateau criterion. Assume $\mathcal{O} \in C^1(\ln \mathcal{B})$ and (6.380) holds. For any $(u,v), (u',v') \in \ln \mathcal{B}$, integrate along the axis–aligned path $(u,v) \to (u',v) \to (u',v')$:

$$\mathcal{O}(u',v') - \mathcal{O}(u,v) = \int_u^{u'} \partial_u \mathcal{O}(\tilde{u}, v)\, d\tilde{u} + \int_v^{v'} \partial_v \mathcal{O}(u', \tilde{v})\, d\tilde{v}.$$

Taking absolute values and using the L^∞ bounds,

$$|\mathcal{O}(u',v') - \mathcal{O}(u,v)| \;\le\; \varepsilon_{\lambda_{\mathrm{UV}}} |u'-u| + \varepsilon_\ell |v'-v|,$$

which is the claimed Lipschitz–type oscillation estimate.

(2) Finite–difference certification ⇒ plateau criterion. Assume now $\mathcal{O} \in C^2(\ln \mathcal{B})$ and that the central finite differences, on a grid with meshes $\Delta u, \Delta v > 0$, satisfy the displayed bounds uniformly on $\ln \mathcal{B}$. Fix $(u, v) \in \ln \mathcal{B}$ and apply Taylor's theorem with (pointwise) remainder:

$$\mathcal{O}(u+\Delta u, v) = \mathcal{O}(u, v) + \Delta u\, \partial_u \mathcal{O}(u, v) + \tfrac{1}{2}(\Delta u)^2\, \partial_u^2 \mathcal{O}(\xi_+, v),$$
$$\mathcal{O}(u-\Delta u, v) = \mathcal{O}(u, v) - \Delta u\, \partial_u \mathcal{O}(u, v) + \tfrac{1}{2}(\Delta u)^2\, \partial_u^2 \mathcal{O}(\xi_-, v),$$

for some $\xi_\pm$ between u and $u \pm \Delta u$ (mean–value form of the remainder). Subtract, divide by $2\Delta u$, and bound the remainder:

$$\begin{aligned} &\left| \frac{\mathcal{O}(u+\Delta u, v) - \mathcal{O}(u-\Delta u, v)}{2\Delta u} - \partial_u \mathcal{O}(u, v) \right| \\ &\quad \le \frac{\Delta u}{4} \Big(\left| \partial_u^2 \mathcal{O}(\xi_+, v) \right| + \left| \partial_u^2 \mathcal{O}(\xi_-, v) \right| \Big) \\ &\quad \le \frac{\Delta u}{2} \, \| \, \partial_u^2 \mathcal{O} \, \|_\infty . \end{aligned}$$

Hence

$$|\partial_u \mathcal{O}(u, v)| \;\le\; \left| \frac{\mathcal{O}(u+\Delta u, v) - \mathcal{O}(u-\Delta u, v)}{2\Delta u} \right| + \frac{\Delta u}{2} \, \| \, \partial_u^2 \mathcal{O} \, \|_\infty \;\le\; \varepsilon_{\lambda_{\mathrm{UV}}}.$$

Since the bound is uniform on $\ln \mathcal{B}$, we conclude $\| \, \partial_u \mathcal{O} \, \|_{L^\infty(\ln \mathcal{B})} \le \varepsilon_{\lambda_{\mathrm{UV}}}$. The same Taylor argument in v gives

$$|\partial_v \mathcal{O}(u, v)| \le \left| \frac{\mathcal{O}(u, v+\Delta v) - \mathcal{O}(u, v-\Delta v)}{2\Delta v} \right| + \frac{\Delta v}{2} \, \| \, \partial_v^2 \mathcal{O} \, \|_\infty \le \varepsilon_\ell,$$

hence $\| \, \partial_v \mathcal{O} \, \|_{L^\infty(\ln \mathcal{B})} \le \varepsilon_\ell$. Therefore (6.380) holds.

Conclusion. Bounded log–derivatives imply the announced Lipschitz oscillation bound, and conversely, uniformly certified central differences with second–derivative remainders control the true derivatives, establishing the plateau criterion. (Mean–value/Taylor remainder estimates; cf. Rudin, §7.) □

Proposition 6.299 (Cohomological (Nielsen) invariance of plateau sets)**.** *Let $\mathcal{O}$ be computed on the physical sector $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ with BRST–closed scale tags that commute with the channel projectors (cf. Definitions 6.270, 6.275 and 6.295). If $\mathcal{O}$ obeys Nielsen/cohomological identities on the band (invariance under ζ–exact gauge–fermion deformations), then the plateau property (6.380) is gauge–parameter independent: the set of $\mathcal{B}$ for which (6.380) holds is unchanged under ζ–exact variations.* (BRST algebra / algebraic renormalization; [117, 82].)

Proof of Proposition 6.299. **Notation.** Set

$$u := \ln \lambda_{\mathrm{UV}}, \qquad v := \ln \ell, \qquad D_\lambda := \partial_u, \quad D_\ell := \partial_v.$$

Let $\Psi(\xi)$ be a gauge–fixing fermion (gauge parameter ξ), and write the (tagged, possibly channel–projected) observable computed on the physical sector as

$$\mathcal{O}_\xi(u, v) \;=\; \Big\langle \mathcal{F}\big(P_x(\lambda_{\mathrm{UV}}),\, K_{\mathrm{IR}}^{(\ell)},\, \Pi_\alpha;\ \text{fields}\big) \Big\rangle_{\mathcal{H}_{\mathrm{phys}},\, \Psi(\xi)},$$

where $P_x(\lambda_{\mathrm{UV}})$, $K_{\mathrm{IR}}^{(\ell)}$ are BRST–closed, ghost number 0, and commute with the channel projectors Π_α.

Step 1: Nielsen/cohomological identity (pointwise on the band). By anomaly–free BRST and the hypotheses (BRST–closed tags; evaluation on $\mathcal{H}_{\mathrm{phys}}$), the Nielsen identity holds *for every* (u, v) *in the band*:

$$\partial_\xi \mathcal{O}_\xi(u, v) = \big\langle \zeta\, \eta_\xi\, \mathcal{F}(\cdots) \big\rangle_{\mathcal{H}_{\mathrm{phys}}} = 0,$$

since BRST–exact insertions have vanishing expectation on $\mathcal{H}_{\mathrm{phys}}$.

Step 2: Commutation of (u, v)–derivatives with ∂_ξ. The only u, v dependence enters through the *tags* $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$. By assumption these tags are BRST–closed and independent of ξ; hence ∂_ξ differentiates solely the gauge–fermion sector inside the expectation and *commutes* with $D_\lambda = \partial_u$ and $D_\ell = \partial_v$:

$$\partial_\xi D_\lambda^m D_\ell^n\, \mathcal{O}_\xi(u, v) = D_\lambda^m D_\ell^n\, \partial_\xi \mathcal{O}_\xi(u, v) = 0 \qquad (\forall\, m, n \in \mathbb{N}_0).$$

Therefore, for all integers $m, n \geq 0$,

$$D_\lambda^m D_\ell^n\, \mathcal{O}_\xi(u, v) \quad \text{is } \textit{independent of } \xi \text{ pointwise on the band.}$$

Step 3: Gauge–parameter independence of the plateau inequalities. The plateau criterion (6.380) reads

$$\|D_\lambda \mathcal{O}_\xi\|_{L^\infty(\ln \mathcal{B})} \leq \varepsilon_{\lambda_{\mathrm{UV}}}, \qquad \|D_\ell \mathcal{O}_\xi\|_{L^\infty(\ln \mathcal{B})} \leq \varepsilon_\ell.$$

By Step 2 the *functions* $D_\lambda \mathcal{O}_\xi$ and $D_\ell \mathcal{O}_\xi$ are ξ–independent pointwise on $\ln \mathcal{B}$; hence their essential suprema over $\ln \mathcal{B}$ are also ξ–independent. Consequently, the truth values of the two inequalities above do not change under ζ–exact variations of Ψ, i.e. the subset of $\mathcal{B}$ on which (6.380) holds is the same for all ξ.

The same argument applies verbatim to higher log–derivatives (used in strengthened plateau tests) and to channel–resolved observables, because $[\Pi_\alpha, P_x] = [\Pi_\alpha, K_{\mathrm{IR}}^{(\ell)}] = 0$ prevents channel mixing and preserves BRST closure.

Conclusion. Nielsen/cohomological identities make $\mathcal{O}_\xi$ and all its (u, v)–derivatives ξ–independent on the band. Hence the plateau property (6.380)—being a set of uniform bounds on those derivatives—is gauge–parameter independent. Equivalently, the family of plateau sets $\mathcal{B}$ is invariant under ζ–exact gauge–fermion deformations. □

Definition 6.300 (Acceptance rule for mode identification)**.** A phenomenological label (e.g. "Higgs–like", "graviton–like", "apeiron", "chrono–kymon") attached to $\mathcal{O}$ in a given quadrant domain (cf. Section 6.12.5) is *provisional* until there exists a nonempty, connected box $\mathcal{B}$ on which (6.380) holds and the inferred exponents/parameters are constant up to the tolerance budget implied by Proposition 6.298. Failure of (6.380) invalidates the claim at that scale.

Intuitive reading. A "plateau'' means we can wiggle the UV cutoff and IR window over a finite box without the observable budging beyond small, predeclared tolerances. Working in $\ln \lambda_{\mathrm{UV}}$ and $\ln \ell$ makes the test dimensionless and additive: if both log–derivatives stay tiny, the observable is genuinely scale–stable. Because the BRST/Nielsen identities remove gauge–parameter dependence in the physical sector, whether a plateau exists is a physics statement, not a gauge artifact—so labels are only accepted once they persist across a bona fide plateau.

6.12.5 Working Strategy to Traverse the Quadrants

Quadrant I – Law symmetric, State symmetric (unbroken). Goal. Fix kinematics and exact identities at the symmetry point; establish a healthy spectrum (no ghosts/tachyons) on $\mathcal{H}_{\text{phys}}$.

Steps.

1. Solve the field equations at the G-invariant vacuum: $\mathcal{E}_{\mathbf{\Phi}} = 0$, $\mathcal{E}_{\mathbf{\Theta}} = 0$ (or their scale-tagged forms (6.333)–(6.334)); verify existence/uniqueness of the symmetric solution.
2. Check exact conservation on $\mathcal{H}_{\text{phys}}$ with $\epsilon_k = 0$: $\nabla_\mu^{(\text{ad})} \bar{J}^\mu_{a,\text{imp}} = 0$ by eq. (6.283) (anomaly-free Stage I, $\langle \zeta(\cdot) \rangle = 0$).
3. Build quadratic kernels $\mathcal{K}_\Phi, \mathcal{K}_\Theta$ and invert to propagators via (6.342); confirm positivity/unitarity (no negative-norm/ghost excitations in the BRST quartet sense).
4. Holosymmetric tomography: verify plateau-stable $|\Lambda_{\text{law}}| \approx 1$ and $|\Lambda_{\text{state}}| \approx 1$ (eq. (6.343)), with T-even/T-odd separation channel-wise (corr/vac/spec). Enforce Section 6.12.4 and Nielsen independence (Proposition 6.264).

Exit. Exact Ward; stable spectrum; visibilities ≈ 1 on plateaus; gauge-parameter independent.

Quadrant II – Law symmetric, State asymmetric (SSB) – provisional tags Goal. Classify NG/radial content and possible gauge absorptions (BRST-consistent), and map mixing blocks.

Steps.

1. Minimise $\mathcal{V}_\Phi(\hat{\mathbf{\Phi}}_{\lambda_{\text{UV}}})$, $\mathcal{V}_\Theta(\hat{\mathbf{\Theta}}_{\lambda_{\text{UV}}})$ to obtain $(\bar{\Phi}_{\lambda_{\text{UV}}}, \bar{\Theta}_{\lambda_{\text{UV}}}) \neq 0$; identify $H \subset G$ and assemble $(M^2_\Phi)^{(\lambda_{\text{UV}})}, (M^2_\Theta)^{(\lambda_{\text{UV}})}, (M^2_{\Phi\Theta})^{(\lambda_{\text{UV}})}$.
2. Diagonalise the quadratic form; tag modes provisionally (NG, radial, mixed) and record uncertainties; use BRST quartet logic to avoid literal "gauge-SSB'' interpretations.
3. Confirm $|\Lambda_{\text{law}}| \approx 1$ and $\Lambda_{\text{state}}(V) \to 0$ as $V \to \infty$; extract NG kinematics from slow phase twists on the vacuum manifold (eq. (6.343)). Check plateau stability and Nielsen independence.
4. Log channel signatures: T-even (order parameter, stiffness in corr/vac) vs. T-odd (spectral transport/phase curvature).

Exit. Correct NG counting; BRST-consistent absorption where present; **provisional** spectrum/mixing ledger, plateau-stable.

Quadrant III – Law asymmetric, State symmetric (emergent symmetry in the IR). Goal. Start from small explicit breakings in the micro-law; demonstrate IR restoration and derive approximate Ward identities.

Steps.

1. Add $\sum_k \epsilon_k \mathcal{O}_k$ with irrelevant scaling $\Delta_k > d + z$.
2. Coarse-grain to window ℓ with $K_{\mathrm{IR}}^{(\ell)}$; bound the law-coherence deficit via Duhamel/BCH to obtain $1 - |\Lambda_{\mathrm{law}}(\ell)| = \mathcal{O}(\ell^{\,\Delta-(d+z)})$ with $\Delta = \min_k \Delta_k$ (Proposition 6.277); enforce plateau stability across $(\lambda_{\mathrm{UV}}, \ell)$ bands.
3. Using eq. (6.283), show the breaking source decays with the same scaling (approximate conservation on $\mathcal{H}_{\mathrm{phys}}$).
4. Verify a unique symmetric vacuum and $|\Lambda_{\mathrm{state}}| \approx 1$; confirm Nielsen independence for extracted exponents (Proposition 6.264).

Exit. Restoration scale $\ell_\star$ (and/or $\mu_\star$) with $|\Delta_a^{(\lambda_{\mathrm{UV}},\ell_\star)}| \le \varepsilon_{\mathrm{tol}}$; recorded exponents, plateau-stable and gauge-parameter independent.

Quadrant IV – Law asymmetric, State asymmetric (explicit + spontaneous). Goal. Analyse full misalignment: explicit breaking plus SSB ⇒ pseudo-NG modes, mixed mass matrices, and consistency (anomaly/BRST) checks.

Steps.

1. Keep $\epsilon_k \neq 0$; choose vacua $(\bar{\Phi}, \bar{\Theta}) \neq 0$; diagonalise full M^2 including explicit terms; track mixing angles and channel attributions.
2. Use eq. (6.375) (physical form eq. (6.377)) to relate $\Delta_a^{(\lambda_{\mathrm{UV}},\ell)}$ to pseudo-NG masses (GMOR-type relations in the present operator setting); check scaling with explicit couplings.
3. Verify $|\Lambda_{\mathrm{law}}| < 1$ (bounded away from unity) and trivial $|\Lambda_{\mathrm{state}}|$ (unique vacuum); ensure plateau stability and Nielsen independence.
4. Confirm T-odd holosymmetric signals track non-conservation; recheck cone preservation (no law-signalling) and Stage I admissibility (no anomalies).

Exit. Controlled pseudo-NG spectrum and misalignment map; Ward/consistency checks passed; scale-stable, gauge-parameter independent diagnostics.

6.12.6 IR/UV-aware Add-ons and Deliverables

QI add-ons (unbroken). (i) *UV health check:* verify positivity of g_T, g_C and ghost-freeness *on the band* (no negative-norm states in $\mathcal{H}_{\mathrm{phys}}$); scan $\lambda_{\mathrm{UV}} \mapsto c\,\lambda_{\mathrm{UV}}$ to confirm stability. (ii) *IR causality & clustering:* at finite ℓ, confirm vanishing of commutators outside cones and clustering of two-point functions as $|x-y|\uparrow$ (cone preservation). (iii) *Plateau:* residues and conserved charges stable across $(\lambda_{\mathrm{UV}}, \ell)$ scans; check Nielsen independence by Proposition 6.264.
Deliverables: bandlimited kernels $\mathcal{K}_\Phi^{(\lambda_{\mathrm{UV}})}$, $\mathcal{K}_\Theta^{(\lambda_{\mathrm{UV}})}$ and propagators; conserved currents $J_a^{(\lambda_{\mathrm{UV}},\ell)}$; plateau plots of visibilities and charges versus $(\lambda_{\mathrm{UV}}, \ell)$ with uncertainty bands.

QII add-ons (SSB; provisional tags). (i) *Vacuum at scale:* minimise $\mathcal{V}_\Phi(\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}}}) + \mathcal{V}_\Theta(\hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}}})$ to obtain $(\bar{\Phi}_{\lambda_{\mathrm{UV}}}, \bar{\Theta}_{\lambda_{\mathrm{UV}}}) \neq 0$; identify $H \subset G$. (ii) *Hessians on the band:* build and diagonalise $(M^2)^{(\lambda_{\mathrm{UV}})}$ as functions of λ_{UV}; record mixing blocks. (iii) *IR robustness:* accept any mode tag only if eigen-data are plateau-stable in both knobs (Section 6.12.4) and gauge-parameter independent (Nielsen). (iv) *Holosymmetric tomography with windows:* phase scans across

ℓ to separate T-even (order parameter, stiffness) from T-odd (transport/phase curvature) channel contributions.
Deliverables: NG counting and vacuum manifold map; $(M^2)^{(\lambda_{\rm UV})}$ eigen-spectra with stability bands $[\lambda_{\rm UV,min}, \lambda_{\rm UV,max}]$, $[\ell_{\rm min}, \ell_{\rm max}]$; holosymmetric visibility traces $\Lambda_{\rm law}, \Lambda_{\rm state}(V)$ demonstrating $|\Lambda_{\rm law}| \approx 1$ and $\Lambda_{\rm state}(V) \to 0$.

QIII add-ons (emergent symmetry). (i) *Small explicit breakings:* include $\varepsilon\, \mathcal{O}_{\rm break}$ with $\Delta_{\rm break} > d+z$. (ii) *Dual flow:*

$$\frac{d\vec{\lambda}}{d\ln\mu} = \vec{\beta}(\vec{\lambda}, \varepsilon), \qquad \frac{d}{d\ln\ell}\Delta_a^{(\lambda_{\rm UV},\ell)} = \gamma_a(\vec{\lambda}, \varepsilon; \lambda_{\rm UV}, \ell). \tag{6.381}$$

(iii) *Emergent Ward plateau:* demonstrate $\Delta_a^{(\lambda_{\rm UV},\ell)} \to 0$ as $\ell \uparrow$ (and/or $\mu \downarrow$) and degeneracy restoration in IR-averaged spectra; extract exponents in (6.277). (iv) *Knob disentangling:* repeat at several $\lambda_{\rm UV}$ to rule out UV artifacts; verify Nielsen independence.
Deliverables: restoration scales $\ell_\star$, $\mu_\star$ with $|\Delta_a^{(\lambda_{\rm UV},\ell_\star)}| \le \varepsilon_{\rm tol}$; law-coherence deficit fits $1 - |\Lambda_{\rm law}(\ell)| \sim \ell^{\Delta-(d+z)}$; channel-resolved visibility trends.

QIV add-ons (explicit + spontaneous). (i) *Pseudo-Goldstone scaling (window-aware):*

$$m_{\rm pG}^2(\lambda_{\rm UV}, \ell) \;\propto\; \frac{\partial \Delta_a^{(\lambda_{\rm UV},\ell)}}{\partial \pi_a} \Big/ f_a^{2(\lambda_{\rm UV},\ell)} \quad \text{evaluated at } (\bar{\Phi}_{\lambda_{\rm UV}}, \bar{\Theta}_{\lambda_{\rm UV}}), \tag{6.382}$$

and track $\ell \uparrow$ softening and $\lambda_{\rm UV}$-insensitivity. (ii) *Misalignment matrix:* diagonalise the full $M^2(\lambda_{\rm UV})$ including explicit terms; map mixing angles versus $(\lambda_{\rm UV}, \ell)$. (iii) *Anomaly consistency:* test $\Delta_a^{(\lambda_{\rm UV},\ell)}$ including possible inflow; verify that T-odd holosymmetric phases track non-conservation; Stage I excludes any anomaly. (iv) *Robustness:* accept pseudo-Goldstone claims only if plateau-stable and Nielsen independent.
Deliverables: pseudo-Goldstone spectrum with $(\lambda_{\rm UV}, \ell)$ stability bands; misalignment/mixing map; non-conservation diagnostics consistent with eq. (6.377); cone-preservation checks.

6.12.7 Practical Ledgers (Maintained Throughout)

1. **Scale settings.** $\lambda_{\rm UV}$ (UV band edge / resolution), ℓ (IR window), kernel $K_{\rm IR}^{(\ell)}$ (Gaussian/box), tolerance $\varepsilon_{\rm tol}$; record plateau bands $[\lambda_{\rm UV,1}, \lambda_{\rm UV,2}] \times [\ell_1, \ell_2]$.

2. **Ward identity ledger.** Currents $\{J_a^\mu\}$ and sources $\{\Delta_a^{(\lambda_{\rm UV},\ell)}\}$ across $(\lambda_{\rm UV}, \ell)$; verdict per setting: exact / approximate / provisional (eqs. (6.282), (6.375) and (6.377)); note Nielsen checks.

3. **Mixing ledger.** Eigen-data of $(M_\Phi^2)^{(\lambda_{\rm UV})}, (M_\Theta^2)^{(\lambda_{\rm UV})}, (M_{\Phi\Theta}^2)^{(\lambda_{\rm UV})}$; channel attributions; plateau ranges; provisional tags and their justifications.

4. **Holosymmetric diagnostics.** Visibility settings $V(\varphi, \theta; \lambda_{\rm UV}, \ell)$; traces of $\Lambda_{\rm law}, \Lambda_{\rm state}$ with T-even/T-odd separation per channel; cone-preservation checks; gauge-parameter (Nielsen) stability.

5. **RG + window flows.** $\vec{\beta}(\lambda_{\rm UV})$ and $\gamma_a(\lambda_{\rm UV}, \ell)$ from (6.381); restoration/misalignment scales $\ell_\star, \mu_\star$; summary of exponent fits with errors.

Chapter 7

Field Laws and Physical States

7.1 Quadrant I: Symmetric Vacuum and Scale-Tagged EOM

7.1.1 Background Split and Vacuum Equations

Definition 7.1 (Background–plus–fluctuation split and vacuum equations)**.** Work at a G–invariant point and decompose the operator–valued fields as

$$\hat{\mathbf{\Phi}} = \bar{\Phi}\,\mathbb{I} + \delta\hat{\mathbf{\Phi}}, \qquad \hat{\mathbf{\Theta}} = \bar{\Theta}\,\mathbb{I} + \delta\hat{\mathbf{\Theta}} \quad \text{(cf. } \textit{Definitions } 6.142 \textit{and } 11.17\text{)}.$$

The c-number backgrounds $\bar{\Phi}, \bar{\Theta}$ are required to solve the Euler–Lagrange (vacuum) equations

$$\mathcal{E}_{\mathbf{\Phi}}[\bar{\Phi}] = 0, \qquad \mathcal{E}_{\mathbf{\Theta}}[\bar{\Theta}] = 0,$$

with $\mathcal{E}_{\mathbf{\Phi}}, \mathcal{E}_{\mathbf{\Theta}}$ as in Definitions 6.267 and 6.268 (see also eqs. (6.327) and (6.328)).

Proposition 7.2 (Quadratic expansion and Hessian blocks)**.** *Collect the fluctuations into* $\delta\hat{\mathbf{U}} := (\delta\hat{\mathbf{\Phi}},\, \delta\hat{\mathbf{\Theta}})^{\mathsf{T}}$*. The quadratic expansion of the action about* $(\bar{\Phi}, \bar{\Theta})$ *defines the linear operator*

$$\mathbb{H}(\bar{\Phi}, \bar{\Theta}) \;=\; \begin{pmatrix} \mathbb{K}_{\Phi} + \mathbb{M}_{\Phi\Phi} & \mathbb{M}_{\Phi\Theta} \\ \mathbb{M}_{\Theta\Phi} & \mathbb{K}_{\Theta} + \mathbb{M}_{\Theta\Theta} \end{pmatrix},$$

acting on $\delta\hat{\mathbf{U}}$*, where the kinetic parts* $\mathbb{K}_{\Phi}, \mathbb{K}_{\Theta}$ *and the Hessian (mass/mixing) blocks* $\mathbb{M}_{\bullet\bullet}$ *are those introduced in Definition 6.271 (cf. eqs. (6.330) and (6.331)). Equivalently, the first variations satisfy the linearised equations* $\delta\mathcal{E}_{\mathbf{\Phi}} = 0$, $\delta\mathcal{E}_{\mathbf{\Theta}} = 0$ *in the block form of eq. (6.331).*

Proposition 7.3 (BRST reading: physical Hessian)**.** *Let* $\mathsf{P}_{\mathrm{phys}}$ *project onto* $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$*. Longitudinal and ghost directions assemble into* ζ*–quartets and decouple in cohomology, whence the physically relevant quadratic operator is*

$$\mathbb{H}_{\mathrm{phys}}(\bar{\Phi}, \bar{\Theta}) \;:=\; \mathsf{P}_{\mathrm{phys}}\, \mathbb{H}(\bar{\Phi}, \bar{\Theta})\, \mathsf{P}_{\mathrm{phys}},$$

as in eq. (6.332) (cf. Definition 6.270). Normal modes and spectra used later in Section 6.12.5 are computed from $\mathbb{H}_{\mathrm{phys}}$*.*

Intuitive reading. Choose G–invariant backgrounds that already solve the full equations; small ripples $\delta\hat{\mathbf{\Phi}}, \delta\hat{\mathbf{\Theta}}$ then obey a *linear* system governed by the Hessian $\mathbb{H}$: the kinetic parts dictate propagation through the background chronogeometry, while the Hessian blocks supply masses and mixings. Projecting to the physical BRST cohomology simply removes gauge and ghost ripples, so the eigenmodes of $\mathbb{H}_{\mathrm{phys}}$ are the genuine, observable small oscillations around $(\bar{\Phi}, \bar{\Theta})$ that feed into the quadrant analysis later on.

7.1.2 Scale-tagged Sector Actions and Euler–Lagrange Operators

Activate the UV projector $P_x(\lambda_{\mathrm{UV}})$ and IR window $K_{\mathrm{IR}}^{(\ell)}$ to form the *scale-tagged sector actions* (with ∇_μ the split connection of eq. (6.314)):

$$\begin{aligned} S_{\Phi}^{(\lambda_{\mathrm{UV}},\ell)} &= \int d^d x\, K_{\mathrm{IR}}^{(\ell)} \sqrt{-\det \bar{\Sigma}^{(\ell)}}\ \mathrm{Tr}\Big\{ \bar{\Sigma}^{(\ell)\mu\nu}\, g_T\Big(\nabla_\mu \hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}}}, \nabla_\nu \hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}}}\Big) + \mathcal{V}_\Phi(\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}}})\Big\}, \\ S_{\Theta}^{(\lambda_{\mathrm{UV}},\ell)} &= \int d^d x\, K_{\mathrm{IR}}^{(\ell)} \sqrt{-\det \bar{\Sigma}^{(\ell)}}\ \mathrm{Tr}\Big\{ \bar{\Sigma}^{(\ell)\mu\nu}\, g_C\Big(\nabla_\mu \hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}}}, \nabla_\nu \hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}}}\Big) + \mathcal{V}_\Theta(\hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}}})\Big\}. \end{aligned} \tag{7.1}$$

Varying gives the *projected Euler–Lagrange operators* (general form):

$$\boxed{\mathcal{E}_{\Phi}^{(\lambda_{\mathrm{UV}},\ell)} := P_x(\lambda_{\mathrm{UV}})\Big[-\nabla_\mu\Big(\bar{\Sigma}^{(\ell)\mu\nu}\, \frac{\partial \mathcal{L}_\Phi}{\partial(\nabla_\nu \hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}}})}\Big) + \frac{\partial \mathcal{L}_\Phi}{\partial \hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}}}}\Big] P_x(\lambda_{\mathrm{UV}}) = 0,} \tag{7.2}$$

$$\boxed{\mathcal{E}_{\Theta}^{(\lambda_{\mathrm{UV}},\ell)} := P_x(\lambda_{\mathrm{UV}})\Big[-\nabla_\mu\Big(\bar{\Sigma}^{(\ell)\mu\nu}\, \frac{\partial \mathcal{L}_\Theta}{\partial(\nabla_\nu \hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}}})}\Big) + \frac{\partial \mathcal{L}_\Theta}{\partial \hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}}}}\Big] P_x(\lambda_{\mathrm{UV}}) = 0.} \tag{7.3}$$

which is summarised in eq. (6.374).

Equivalently—using the "clean" kinetic pairings $\mathcal{G}_T, \mathcal{G}_C$—the *clean, scale-tagged EOM* read

$$\boxed{\mathcal{E}_{\mathbf{\Phi}}^{(\lambda_{\mathrm{UV}},\ell)} := P_x(\lambda_{\mathrm{UV}})\Big[-\nabla_\mu\Big(\bar{\Sigma}^{(\ell)\mu\nu}\, \mathcal{G}_T\, \nabla_\nu \hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}}}\Big) + \frac{\partial \mathcal{V}_\Phi}{\partial \hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}}}}\Big] P_x(\lambda_{\mathrm{UV}}) = 0} \tag{7.4}$$

$$\boxed{\mathcal{E}_{\mathbf{\Theta}}^{(\lambda_{\mathrm{UV}},\ell)} := P_x(\lambda_{\mathrm{UV}})\Big[-\nabla_\mu\Big(\bar{\Sigma}^{(\ell)\mu\nu}\, \mathcal{G}_C\, \nabla_\nu \hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}}}\Big) + \frac{\partial \mathcal{V}_\Theta}{\partial \hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}}}}\Big] P_x(\lambda_{\mathrm{UV}}) = 0} \tag{7.5}$$

with the BRST reading "insert $\mathsf{P}_{\mathrm{phys}}$" and keep P_x, K_{IR} BRST-closed (ghost number 0).

7.1.3 BRST-consistent Reading on $\mathcal{H}_{\mathrm{phys}}$

Definition 7.4 (Cohomological equations on the physical sector)**.** Let $\hat{Q}_{\mathrm{BRST}}$ be nilpotent and set $\zeta(\,\cdot\,) := \{\hat{Q}_{\mathrm{BRST}}, \cdot\}$ (ghost number +1). Choose a gauge–fixing fermion Ψ and add the ζ–exact term $\zeta\Psi$ to the action. Define the local insertions

$$\zeta\, \chi_{\mathbf{\Phi}} := \frac{\delta(\zeta\Psi)}{\delta \hat{\mathbf{\Phi}}}, \qquad \zeta\, \chi_{\mathbf{\Theta}} := \frac{\delta(\zeta\Psi)}{\delta \hat{\mathbf{\Theta}}} \quad \text{(cf. } \textit{Definition } 6.270\text{)}.$$

Let $\mathsf{P}_{\mathrm{phys}}$ denote the orthogonal projector onto $\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}})$. Then the cohomological Euler–Lagrange equations read

$$\mathsf{P}_{\mathrm{phys}}\big(\mathcal{E}_{\mathbf{\Phi}} + \zeta\,\chi_{\mathbf{\Phi}}\big)\mathsf{P}_{\mathrm{phys}} = 0, \qquad \mathsf{P}_{\mathrm{phys}}\big(\mathcal{E}_{\mathbf{\Theta}} + \zeta\,\chi_{\mathbf{\Theta}}\big)\mathsf{P}_{\mathrm{phys}} = 0,$$

with $\mathcal{E}_{\mathbf{\Phi}}, \mathcal{E}_{\mathbf{\Theta}}$ as in Definitions 6.267 and 6.268 (see also eqs. (6.327) and (6.328)).

Proposition 7.5 (Equivalence to the clean equations and gauge-parameter independence)**.** *On $\mathcal{H}_{\mathrm{phys}}$ the above equations are equivalent to the clean forms $\mathsf{P}_{\mathrm{phys}}\,\mathcal{E}_{\mathbf{\Phi}}\,\mathsf{P}_{\mathrm{phys}} = 0$ and $\mathsf{P}_{\mathrm{phys}}\,\mathcal{E}_{\mathbf{\Theta}}\,\mathsf{P}_{\mathrm{phys}} = 0$, because ζ–exact insertions vanish in BRST cohomology. Moreover, changing the gauge parameter in Ψ (e.g. ξ in the general treatment or α_{gf} in BFM) only deforms $\chi_\bullet$ by ζ–exact terms, hence*

$$\partial_\xi\Big(\mathsf{P}_{\mathrm{phys}}\big(\mathcal{E}_\bullet + \zeta\,\chi_\bullet\big)\mathsf{P}_{\mathrm{phys}}\Big) = 0 \quad \textit{or} \quad \partial_{\alpha_{\mathrm{gf}}}\Big(\mathsf{P}_{\mathrm{phys}}\big(\mathcal{E}_\bullet + \zeta\,\chi_\bullet\big)\mathsf{P}_{\mathrm{phys}}\Big) = 0,$$

i.e. Nielsen–type gauge–parameter independence on $\mathcal{H}_{\mathrm{phys}}$ (standard BRST/cohomological identities; see, e.g., [117, 82, 230]).

Proposition 7.6 (Scale–tagged, BRST–closed knobs and channel resolution)**.** *Assume the UV spectral projector $P_x(\lambda_{\mathrm{UV}})$ and the IR window kernel $K_{\mathrm{IR}}^{(\ell)}$ are BRST–closed (ghost number 0) and commute with the channel projectors $\{\Pi_\alpha\}_{\alpha\in\{\mathrm{corr,vac,spec}\}}$. Then every scale–tagged object produced by the functor $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ (cf. Definition 6.282) descends to $\mathcal{H}_{\mathrm{phys}}$, and all cohomological equations and channel–resolved observables are gauge–parameter independent:*

$$\mathsf{P}_{\mathrm{phys}}\,\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\cdot]\,\mathsf{P}_{\mathrm{phys}} = \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}\Big[\mathsf{P}_{\mathrm{phys}}(\cdot)\mathsf{P}_{\mathrm{phys}}\Big],$$

∂_ξ and $\partial_{\alpha_{\mathrm{gf}}}$ derivatives vanish on physical readouts.

(See Definitions 6.272 and 6.275 and Section 6.12.4 for downstream uses.)

7.1.4 UV/IR Knobs and Geometric Inputs

- **UV bandlimit (spectral projector).** $P_x(\lambda_{\mathrm{UV}}) = E^{(x)}([0,\lambda_{\mathrm{UV}}])$ and is applied to all operator fields/correlators per eq. (6.353); positivity and the triadic, non-commutative geometry on the band are as in Chapter 6's channel split.
- **IR window (coarse-graining).** Pick $K_{\mathrm{IR}}^{(\ell)} \geq 0$ with $\int d^d x\, K_{\mathrm{IR}}^{(\ell)} = 1$ and compact support, and define $\bar{\Sigma}_{\mu\nu}^{(\ell)}$ and coarse-grained observables as in eq. (6.371).
- **Band-safety and variational pipeline.** All operator-valued tensors are defined *on the band* (safe in the strong-operator topology), and $P_x(\lambda_{\mathrm{UV}})$ interfaces the relational/sector actions; holosymmetric visibilities subsequently read these equations channel-wise.

7.1.5 Symmetric Vacuum at Scale (What the EOM Demand)

Definition 7.7 (Scale–tagged stationary (vacuum) conditions at a G–invariant point)**.** Let the scale–tagging functor $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ be as in Definition 6.282, and let $\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell}$ and $\hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}},\ell}$ denote

the band–limited, IR–windowed fields (cf. Section 6.12.4). A *homogeneous, G–invariant background* consists of c–number fields $(\bar{\mathbf{\Phi}}, \bar{\mathbf{\Theta}})$ such that

$$\nabla_\mu \bar{\mathbf{\Phi}} = 0, \qquad \nabla_\mu \bar{\mathbf{\Theta}} = 0,$$

and the Euler–Lagrange operators of Definitions 6.267 and 6.268 vanish when evaluated on $(\bar{\mathbf{\Phi}}, \bar{\mathbf{\Theta}})$. For such backgrounds, the *scale–tagged stationary conditions* are

$$\left.\frac{\partial \mathcal{V}_\Phi}{\partial \hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell}}\right|_{\bar{\mathbf{\Phi}}} = 0, \qquad \left.\frac{\partial \mathcal{V}_\Theta}{\partial \hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}},\ell}}\right|_{\bar{\mathbf{\Theta}}} = 0, \tag{7.6}$$

i.e. the fiberwise (Gâteaux) derivatives of the potentials vanish on the constant configurations (cf. Definitions 6.267 and 6.268).

Proposition 7.8 (EOM ⇒ scale–tagged stationary conditions)**.** *Assume the scale–tagged equations of motion eqs.* (6.333) *and* (6.334) *and a homogeneous background* $(\bar{\mathbf{\Phi}}, \bar{\mathbf{\Theta}})$ *with* $\nabla_\mu \bar{\mathbf{\Phi}} = \nabla_\mu \bar{\mathbf{\Theta}} = 0$*. Then* (7.6) *holds.*

Proof. For homogeneous backgrounds, $\nabla_\nu \hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell}\big|_{\bar{\mathbf{\Phi}}} = 0$ and $\nabla_\nu \hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}},\ell}\big|_{\bar{\mathbf{\Theta}}} = 0$ because the background parts are constant and the fluctuation parts vanish on the vacuum configuration; hence the momentum currents in eqs. (6.333) and (6.334) vanish and their covariant divergences are zero. The remaining terms are the potential derivatives, which must therefore vanish at $(\bar{\mathbf{\Phi}}, \bar{\mathbf{\Theta}})$, yielding (7.6). (See standard derivations of homogeneous vacua in covariant field theory [82, 91].) □

Proposition 7.9 (Expectation–value form for homogeneous states)**.** *Let* $|\mathbf{\Psi}\rangle$ *be a homogeneous (translation–invariant) state. Then the stationary conditions are equivalently written as*

$$\left.\frac{\partial \mathcal{V}_\Phi}{\partial \hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell}}\right|_{\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell}=\langle \hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell}\rangle_{\mathbf{\Psi}}} = 0, \qquad \left.\frac{\partial \mathcal{V}_\Theta}{\partial \hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}},\ell}}\right|_{\hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}},\ell}=\langle \hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}},\ell}\rangle_{\mathbf{\Psi}}} = 0. \tag{7.7}$$

with the identifications

$$\bar{\mathbf{\Phi}} = \langle \hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell}\rangle_{\mathbf{\Psi}}, \qquad \bar{\mathbf{\Theta}} = \langle \hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}},\ell}\rangle_{\mathbf{\Psi}}.$$

Proof. For homogeneous $|\mathbf{\Psi}\rangle$, expectation values of local observables are constant; the band functor $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ preserves constants (the UV projector is identity on c–number multiples of $\mathbb{I}$, and the IR kernel integrates to 1), hence $\langle \hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell}\rangle_{\mathbf{\Psi}}$ and $\langle \hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}},\ell}\rangle_{\mathbf{\Psi}}$ are constants that define backgrounds. By the usual chain rule for Gâteaux derivatives on fiber bundles, the stationary conditions on c–numbers are equivalent to those on their expectation values (distributional/functional-analytic details as in [253, 433]). □

Lemma 7.10 (IR window for homogeneous backgrounds)**.** *If* $\bar{\mathbf{\Phi}}, \bar{\mathbf{\Theta}}$ *are constant, then* $K_{\mathrm{IR}}^{(\ell)}$ *acts as the identity on them. Consequently one may replace* $\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell} \to \hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}}}$ *and* $\hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}},\ell} \to \hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}}}$ *in* (7.6)–(7.7)*; all checks remain* on the band *after* $P_x(\lambda_{\mathrm{UV}})$ *(cf. Section 6.12.4).*

Proof. By definition, $\int K_{\mathrm{IR}}^{(\ell)}(x,y)\,\mathrm{d}\mu(y) = 1$ and the kernel is positive; applying it to a constant yields the same constant (mollifier property; [433]). □

BRST–consistent reading and Ward compatibility. Evaluating the stationary conditions on $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ with BRST–closed scale tags preserves the equations and their gauge–parameter independence (Nielsen–type identities), as ζ–exact pieces vanish in cohomology (cf. Section 7.1.3 and Definition 6.295; see [117, 230]). The same scale–tagging that yields $\mathcal{E}_{\bullet}^{(\lambda_{\mathrm{UV}},\ell)} = 0$ also produces the adjoint–covariant, scale–dependent Ward identity of eq. (6.375), enabling later tests of symmetry realization under explicit breakings or anomalies (cf. Section 6.12.2).

7.2 BRST Health and Spectrum at the Symmetric Vacuum (Quadrant I)

This section establishes, with full cohomological care, the BRST consistency and the physical spectrum at the G–invariant vacuum fixed in Section 7.1. All statements hold *on the band* (UV projector $P_x(\lambda_{\mathrm{UV}})$) and within the IR window $K_{\mathrm{IR}}^{(\ell)}$ (cf. Section 6.12.4 and Definition 6.282); the equations of motion used are eqs. (6.333), (6.334) and (6.374). We work in the BRST framework where $\zeta(\,\cdot\,) := \{\hat{Q}_{\mathrm{BRST}}, \cdot\}$ with $\mathrm{gh}(\zeta) = +1$ and $\hat{Q}_{\mathrm{BRST}}^2 = 0$ (nilpotent).

7.2.1 BRST Complex, Nilpotency, and the Physical Sector

Definition 7.11 (Band-limited BRST complex and ghost grading)**.** Introduce the band- and IR–tagged field multiplet (cf. Section 6.12.4):

$$\mathfrak{X}^{(\lambda_{\mathrm{UV}},\ell)} := \Big(\delta\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell},\, \delta\hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}},\ell},\, c_{\lambda_{\mathrm{UV}},\ell},\, \bar{c}_{\lambda_{\mathrm{UV}},\ell},\, b_{\lambda_{\mathrm{UV}},\ell}\Big)^{\mathsf{T}}.$$

where $c, \bar{c}$ are Faddeev–Popov ghost/antighost fields and b is the Nakanishi–Lautrup auxiliary (e.g. [104, 105, 106, 82, 442]). Ghost numbers are assigned as

$$\mathrm{gh}(\delta\hat{\mathbf{\Phi}}) = \mathrm{gh}(\delta\hat{\mathbf{\Theta}}) = \mathrm{gh}(b) = 0, \qquad \mathrm{gh}(c) = +1, \qquad \mathrm{gh}(\bar{c}) = -1, \qquad \mathrm{gh}(\zeta) = +1.$$

By construction (BRST-closed tags), $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ commute with $\hat{Q}_{\mathrm{BRST}}$ and the channel projectors Π_α (cf. Section 7.1.3). Hence the band functor $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ preserves the BRST complex and its ghost grading.

Proposition 7.12 (Nilpotency, (Krein-)Hermiticity, and domain)**.** *At the symmetric G–invariant vacuum (Quadrant I) the BRST charge obeys*

$$\hat{Q}_{\mathrm{BRST}}^2 = 0, \qquad \hat{Q}_{\mathrm{BRST}}^{\dagger} = \hat{Q}_{\mathrm{BRST}}, \tag{7.8}$$

on a common invariant domain stable under $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$. Consequently $\zeta^2 = 0$ and the BRST differential raises ghost number by $+1$. Sketch. *Nilpotency and adjointness are standard at the symmetric baseline with gauge fixing added as $\zeta\Psi$; band/IR tags are BRST–closed and thus preserve the algebra and domains (see [117, 230, 442]).*

Definition 7.13 (Physical sector and cohomology)**.** The *physical Hilbert space* is the ghost-number–zero cohomology

$$\mathcal{H}_{\text{phys}} := H^0(\hat{Q}_{\text{BRST}}) = \frac{\ker \hat{Q}_{\text{BRST}}|_{\text{gh}=0}}{\operatorname{im} \hat{Q}_{\text{BRST}}|_{\text{gh}=-1}}, \tag{7.9}$$

under the inner product induced from the (Krein) space of band-limited states. BRST–exact vectors represent pure gauge and vanish in cohomology; expectation values and S-matrix elements are taken between cohomology classes only (cf. Section 7.1.3).

Theorem 7.14 (Kugo–Ojima quartet decoupling on the band)**.** *At the symmetric vacuum and for BRST–closed tags, negative–norm and longitudinal excitations combine with ghosts into BRST quartets and become trivial in cohomology. Thus unitarity/positivity are recovered on $\mathcal{H}_{\text{phys}}$, and physical observables are independent of gauge parameters (Nielsen identities).*

Idea of proof. The quartet mechanism is the standard BRST implementation of gauge constraints: one shows that $\{Q_{\text{BRST}}, \cdot\}$ pairs longitudinal and ghost/antighost modes into exact-doublets whose contributions cancel in all cohomological matrix elements; band/IR tags commute with Q_{BRST} so the proof carries over verbatim (see [108, 442, 243, 117]). Gauge–parameter independence in the physical sector follows from the cohomological (Nielsen) identity [117, 82]. □

7.2.2 Gauge-fixed Quadratic Theory, Propagators, and Cohomological Invariance

Definition 7.15 (Gauge-fixed quadratic action and physical kernel)**.** Add gauge fixing via a fermion Ψ and form $S_\Psi = S + \zeta\Psi$. Expanding to quadratic order around the symmetric vacuum $(\bar{\Phi}, \bar{\Theta})$ yields

$$S_\Psi^{(2)} = \tfrac{1}{2}\Big\langle \delta\hat{\mathbf{U}}_{\lambda_{\text{UV}},\ell},\ \mathcal{K}_\Psi^{(\lambda_{\text{UV}},\ell)}\, \delta\hat{\mathbf{U}}_{\lambda_{\text{UV}},\ell} \Big\rangle, \quad \delta\hat{\mathbf{U}}_{\lambda_{\text{UV}},\ell} := (\delta\hat{\mathbf{\Phi}}_{\lambda_{\text{UV}},\ell},\ \delta\hat{\mathbf{\Theta}}_{\lambda_{\text{UV}},\ell})^{\mathsf{T}}.$$

Projecting to $\mathcal{H}_{\text{phys}}$ gives the *physical* kernel

$$\mathcal{K}_{\text{phys}}^{(\lambda_{\text{UV}},\ell)} := \mathsf{P}_{\text{phys}}\, \mathcal{K}_\Psi^{(\lambda_{\text{UV}},\ell)}\, \mathsf{P}_{\text{phys}},$$

whose inverse (retarded/advanced/Feynman) is the physical propagator $\mathcal{D}_{\text{phys}}^{(\lambda_{\text{UV}},\ell)}$ (cf. Definition 6.274).

Proposition 7.16 (Cohomological inversion and gauge-parameter independence)**.** *On $\mathcal{H}_{\text{phys}}$ one has*

$$\mathcal{K}_{\text{phys}}^{(\lambda_{\text{UV}},\ell)} \cdot \mathcal{D}_{\text{phys}}^{(\lambda_{\text{UV}},\ell)} = \mathbf{1},$$

and the pole locations/residues of $\mathcal{D}_{\text{phys}}^{(\lambda_{\text{UV}},\ell)}$ are independent of the gauge parameter ($\partial_\xi = 0$ in the generic formulation, or $\partial_{\alpha_{\text{gf}}} = 0$ in BFM). Sketch. *Quartet decoupling removes longitudinal/ghost blocks before inversion; the Slavnov–Taylor/Nielsen identities ensure that gauge–parameter variations are ζ–exact insertions that vanish in cohomology [117, 82, 91].*

Proposition 7.17 (Källén–Lehmann form and positivity)**.** *For any gauge–invariant local operator $\mathcal{O}$ of ghost number* 0*, its two–point function on $\mathcal{H}_{\mathrm{phys}}$ admits a Källén–Lehmann representation* on the band

$$\langle \mathcal{O}\,\mathcal{O}\rangle^{(\lambda_{\mathrm{UV}},\ell)}(p) = \int_0^\infty \frac{\rho_{\mathcal{O}}^{(\lambda_{\mathrm{UV}},\ell)}(m^2)}{p^2 - m^2 + i0}\,\mathrm{d}m^2, \qquad \rho_{\mathcal{O}}^{(\lambda_{\mathrm{UV}},\ell)}(m^2) \geq 0,$$

with the same discrete poles and residues as the untagged physical correlator. Sketch. *Positivity on $\mathcal{H}_{\mathrm{phys}}$ plus locality (Haag–Kastler) yields the standard spectral representation; scale tags are positivity-preserving and commute with the physical projector (cf. Propositions 6.279 and 6.285; see [230, 91]).*

7.2.3 Spectrum at the Symmetric Vacuum (Quadrant I)

Definition 7.18 (Physical normal modes and masses)**.** Let $\mathbb{H}_{\mathrm{phys}}$ be the physical Hessian from Proposition 7.3. Physical normal modes are solutions of $\mathbb{H}_{\mathrm{phys}}\,u_n = \omega_n^2\,u_n$ (in position/momentum space as appropriate), equivalently the poles of $\mathcal{D}_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}$. Masses are read from on–shell conditions $p^2 = m_n^2$ at the poles, with channel resolution provided by Π_α (corr/vac/spec).

Theorem 7.19 (Quadrant I spectrum: no NG from the state, gauge–parameter independent poles)**.** *At the G–invariant vacuum of Section 7.1.5 (Law Sym., State Sym.), (i) there are no Nambu–Goldstone poles sourced by spontaneous breaking of G in the state; (ii) all physical poles and residues of $\mathcal{D}_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}$ are independent of the gauge parameter; (iii) channel-resolved visibilities $\Lambda_{\mathrm{law/state}}^{(\lambda_{\mathrm{UV}},\ell)}$ remain at unit modulus up to finite–size/window effects (cf. Section 6.12.3).*

Proof. (i) follows from the symmetric vacuum conditions Section 7.1.5 (no SSB). (ii) follows from Proposition 7.16 (Nielsen identities). (iii) is the Quadrant I signature with exact Ward identities (Sections 6.12.2 and 6.12.3); BRST–closed tags guarantee cohomological invariance under UV/IR scans within plateaus (Section 6.12.4). □

7.2.4 Scale Tags, Plateaus, and Robustness

Proposition 7.20 (BRST-closed tags commute with the physical projection)**.** *If $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ are BRST–closed and commute with Π_α, then*

$$\mathsf{P}_{\mathrm{phys}}\,\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\cdot]\,\mathsf{P}_{\mathrm{phys}} = \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}\Big[\mathsf{P}_{\mathrm{phys}}(\cdot)\mathsf{P}_{\mathrm{phys}}\Big],$$

so all tagged observables and kernels descend to $\mathcal{H}_{\mathrm{phys}}$ without changing their physical content (cf. Section 7.1.3).

Proposition 7.21 (Plateau stability of the spectrum)**.** *Let $\mathcal{O}$ be any ghost–number–zero observable that diagnoses a pole (mass or residue). If $\mathcal{O}(\lambda_{\mathrm{UV}},\ell)$ satisfies the plateau criterion Section 6.12.4 on a box $\mathcal{B}$, then the extracted masses and residues are invariant under changes of $\lambda_{\mathrm{UV}},\ell$ within $\mathcal{B}$, and coincide with those of the untagged physical theory up to the prescribed tolerances.* (Combine Proposition 6.298 with Propositions 7.16 and 7.244.)

Intuitive reading. Band–limiting (UV) and windowing (IR) are purely *filters* that we arrange to be BRST–closed. They do not change the BRST algebra or the set of physical states; they merely make computations stable and local. At the symmetric vacuum, quartets remove all gauge/ghost contamination, and the remaining physical two–point functions enjoy a positive spectral representation with gauge–parameter independent poles. When an observable sits on a plateau as we vary $(\lambda_{\mathrm{UV}}, \ell)$, we can trust it as a genuine, scale–stable feature of the theory—not a cutoff artifact.

References. Standard sources for BRST, quartet decoupling, and Nielsen/Slavnov–Taylor identities include [104, 105, 106, 117, 442, 243, 82, 91, 230].

7.2.5 Quadratic Action, Hessian, and Transverse/Longitudinal Split

Expanding the scale-tagged actions of Section 6.12.4 to quadratic order around the symmetric background $(\bar{\mathbf{\Phi}}, \bar{\mathbf{\Theta}})$ yields a band-limited quadratic form on the fluctuation multiplet $\mathfrak{X}$ of Section 7.2.1. Denote by $\mathcal{K}_T$ and $\mathcal{K}_C$ the kinetic operators extracted from the clean, scale-tagged EOM (eqs. (6.333) and (6.334)), and by

$$\mathcal{M}^{(\lambda_{\mathrm{UV}},\ell)}_{\Phi\Phi} := \frac{\partial^2 \mathcal{V}_\Phi}{\partial\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell}\,\partial\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell}}\bigg|_{(\bar{\mathbf{\Phi}})}, \quad \mathcal{M}^{(\lambda_{\mathrm{UV}},\ell)}_{\Theta\Theta} := \frac{\partial^2 \mathcal{V}_\Theta}{\partial\hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}},\ell}\,\partial\hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}},\ell}}\bigg|_{(\bar{\mathbf{\Theta}})},$$
$$\mathcal{M}^{(\lambda_{\mathrm{UV}},\ell)}_{\Phi\Theta} := \frac{\partial^2 \mathcal{V}}{\partial\hat{\mathbf{\Phi}}_{\{\lambda_{\mathrm{UV}},\ell\}}\,\partial\hat{\mathbf{\Theta}}_{\{\lambda_{\mathrm{UV}},\ell\}}}\bigg|_{(\bar{\mathbf{\Phi}},\bar{\mathbf{\Theta}})}, \qquad \mathcal{M}^{(\lambda_{\mathrm{UV}},\ell)}_{\Theta\Phi} := \Big(\mathcal{M}^{(\lambda_{\mathrm{UV}},\ell)}_{\Phi\Theta}\Big)^\dagger,$$

the potential (mass/mixing) blocks, all defined *on the UV band and within the IR window* via the "UV then IR" convention.

With the *local* inner pairing

$$\Big\langle A, B\Big\rangle_x := \mathrm{Tr}\Big(A^\dagger(x)\, B(x)\Big),$$

the quadratic action reads

$$S^{(2)}_{\mathrm{band}} = \tfrac{1}{2}\int d^dx\,\sqrt{-\det\bar{\Sigma}^{(\ell)}(x)}\,\Big\langle \mathfrak{X}_{\lambda_{\mathrm{UV}},\ell}(x),\, \mathbb{H}^{(\lambda_{\mathrm{UV}},\ell)}(x)\,\mathfrak{X}_{\lambda_{\mathrm{UV}},\ell}(x)\Big\rangle_x,$$
$$\mathbb{H}^{(\lambda_{\mathrm{UV}},\ell)} := \begin{pmatrix} \mathcal{K}_T + \mathcal{M}^{(\lambda_{\mathrm{UV}},\ell)}_{\Phi\Phi} & \mathcal{M}^{(\lambda_{\mathrm{UV}},\ell)}_{\Phi\Theta} & 0 & 0 & 0 \\ \mathcal{M}^{(\lambda_{\mathrm{UV}},\ell)}_{\Theta\Phi} & \mathcal{K}_C + \mathcal{M}^{(\lambda_{\mathrm{UV}},\ell)}_{\Theta\Theta} & 0 & 0 & 0 \\ 0 & 0 & 0 & \Box^{(\lambda_{\mathrm{UV}},\ell)}_{\mathrm{FP}} & 0 \\ 0 & 0 & \Box^{(\lambda_{\mathrm{UV}},\ell)}_{\mathrm{FP}} & 0 & 0 \\ 0 & 0 & 0 & 0 & \not{\mathbb{H}} \end{pmatrix}. \tag{7.10}$$

Here $\mathfrak{X}_{\lambda_{\mathrm{UV}},\ell} := \Big(\delta\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell},\, \delta\hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}},\ell},\, c_{\lambda_{\mathrm{UV}},\ell},\, \bar{c}_{\lambda_{\mathrm{UV}},\ell},\, b_{\lambda_{\mathrm{UV}},\ell}\Big)^\top$ collects the projected fluctuations, $\Box^{(\lambda_{\mathrm{UV}},\ell)}_{\mathrm{FP}}$ is the band-limited Faddeev–Popov operator generated by the BRST-exact gauge-fixing term $\zeta\Psi$ (schematically $-\bar{c}\,\Box_{\mathrm{FP}}\,c$ with $\Box_{\mathrm{FP}} = \delta F/\delta\theta$ evaluated at $(\bar{\mathbf{\Phi}}, \bar{\mathbf{\Theta}})$), and the b-sector block $\not{\mathbb{H}}$ assumes the conventional normalisation of the gauge fermion (a factor of ξ can be reinstated for R_ξ-type choices).

All blocks act on the *projected* field space (UV band then IR window); equivalently, one may view them as implicitly sandwiched by $P_x(\lambda_{\mathrm{UV}})$ and read within $K^{(\ell)}_{\mathrm{IR}}$. Physical evaluation inserts $\mathsf{P}_{\mathrm{phys}}$ as in Sections 7.1.3 and 7.5.13.

Transverse/longitudinal decomposition. Let $\Pi_{\mathrm{T}}^{(\ell)}$ and $\Pi_{\mathrm{L}}^{(\ell)}$ denote, respectively, the transverse and longitudinal projectors constructed from $\bar{\Sigma}_{\mu\nu}^{(\ell)}$ and the gauge generators, acting on the *projected* field space (UV then IR). They satisfy

$$\Pi_{\mathrm{T}}^{(\ell)} + \Pi_{\mathrm{L}}^{(\ell)} = \mathbb{1}, \qquad \big(\Pi_{\mathrm{T,L}}^{(\ell)}\big)^2 = \Pi_{\mathrm{T,L}}^{(\ell)}, \qquad \zeta\big(\Pi_{\mathrm{T,L}}^{(\ell)}\big) = 0,$$

and commute with the scale knobs: $[\Pi_{\mathrm{T,L}}^{(\ell)},\, P_x(\lambda_{\mathrm{UV}})] = 0$, $[\Pi_{\mathrm{T,L}}^{(\ell)},\, K_{\mathrm{IR}}^{(\ell)}] = 0$ (ghost number 0). Hence, for band-limited fluctuations,

$$\delta\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell} = \Pi_{\mathrm{T}}^{(\ell)}\delta\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell} + \Pi_{\mathrm{L}}^{(\ell)}\delta\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell}, \qquad \delta\hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}},\ell} = \Pi_{\mathrm{T}}^{(\ell)}\delta\hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}},\ell} + \Pi_{\mathrm{L}}^{(\ell)}\delta\hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}},\ell}.$$

Choosing a gauge functional F and the standard gauge fermion Ψ, given by

$$\Psi = \int d^d x\, \sqrt{-\det \bar{\Sigma}^{(\ell)}}\, \bar{c}\, \Big(F + \tfrac{\xi}{2} b\Big),$$

gives the BRST-exact term

$$\zeta\Psi = \int d^d x\, \sqrt{-\det \bar{\Sigma}^{(\ell)}}\, \Big[\, b\, F + \tfrac{\xi}{2} b^2 - \bar{c}\, (\zeta F)\, \Big],$$

with all fields implicitly band-projected. Then

$$\Big(\Pi_{\mathrm{L}}^{(\ell)}\delta\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell},\, c_{\lambda_{\mathrm{UV}},\ell}\Big), \qquad \Big(\Pi_{\mathrm{L}}^{(\ell)}\delta\hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}},\ell},\, c_{\lambda_{\mathrm{UV}},\ell}\Big), \qquad (\bar{c}_{\lambda_{\mathrm{UV}},\ell},\, b_{\lambda_{\mathrm{UV}},\ell})$$

organise into Kugo–Ojima quartets (two BRST doublets), hence decouple in cohomology. Only the transverse components survive in $\mathcal{H}_{\mathrm{phys}}$, and the physical poles are read from the $\Pi_{\mathrm{T}}^{(\ell)}$–projected blocks of $\mathbb{H}^{(\lambda_{\mathrm{UV}},\ell)}$ (plateau-stable under the UV/IR scans of Section 7.3). Background-field derivations of (7.10) and the associated projector algebra are standard; see, e.g., [114, 155, 91] and [43].

ROI-weighted variant (optional). If one prefers an explicit IR window in the gauge-fixing functional, take

$$\Psi^{(\ell)} = \int d^d x\, K_{\mathrm{IR}}^{(\ell)}(x)\, \sqrt{-\det \bar{\Sigma}^{(\ell)}(x)}\, \bar{c}_{\lambda_{\mathrm{UV}}}(x) \Big(F[\cdot](x) + \tfrac{\xi}{2}\, b_{\lambda_{\mathrm{UV}}}(x)\Big),$$

and use *UV-only* tags for $(c, \bar{c}, b)$ to avoid double windowing. Then

$$\zeta\Psi^{(\ell)} = \int d^d x\, K_{\mathrm{IR}}^{(\ell)} \sqrt{-\det \bar{\Sigma}^{(\ell)}}\, \Big[\, b_{\lambda_{\mathrm{UV}}}\, F + \tfrac{\xi}{2} b_{\lambda_{\mathrm{UV}}}^2 - \bar{c}_{\lambda_{\mathrm{UV}}}\, (\zeta F)\, \Big],$$

so the ghost/Nakanishi–Lautrup sector carries only the UV band label, and the induced Faddeev–Popov operator is $\Box_{\mathrm{FP}}^{(\lambda_{\mathrm{UV}})}$. Since $K_{\mathrm{IR}}^{(\ell)}$ has ghost number 0 and is BRST-closed, $\zeta K_{\mathrm{IR}}^{(\ell)} = 0$, Nielsen independence and the quartet decoupling are preserved.

7.2.6 Ghost Decoupling and Positivity on $\mathcal{H}_{\mathrm{phys}}$

Recall that $\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}})$ is the BRST cohomology at ghost number 0, and let $\mathsf{P}_{\mathrm{phys}}$ be the orthogonal projector onto $\mathcal{H}_{\mathrm{phys}}$ with respect to the induced inner product (cf. BRST overlay remarks and gauge-parameter independence by Lemma 6.243). Working in the scale-tagged band setup, the quadratic form eq. (7.10) is block-diagonal in the $(T, C \,|\, \bar{c}, c \,|\, b)$ decomposition, with the FP operator $\Box_{\mathrm{FP}}^{(\lambda_{\mathrm{UV}},\ell)}$ sitting entirely in the ghost/antighost block and the longitudinal projector $\Pi_{\mathrm{L}}^{(\ell)}$ confined to the pure-gauge sector of the gauge-field kinetic block. Thus the ghost/longitudinal content is already separated at the level of the Hessian and remains BRST-exact at the level of operators [104, 105, 106, 442, 91].

Proposition 7.22 (Quartet decoupling in the band formalism)**.** *Assume an anomaly-free BRST setup with gauge fixing implemented by a gauge fermion Ψ so that $\zeta\Psi$ is BRST-exact, and with BRST-closed scale knobs ($\zeta P_x(\lambda_{\mathrm{UV}}) = 0$, $\zeta K_{\mathrm{IR}}^{(\ell)} = 0$). Then on the BRST-physical sector $\mathcal{H}_{\mathrm{phys}}$ one has*

$$\mathsf{P}_{\mathrm{phys}}\, \Box_{\mathrm{FP}}^{(\lambda_{\mathrm{UV}},\ell)}\, \mathsf{P}_{\mathrm{phys}} = 0, \qquad \mathsf{P}_{\mathrm{phys}}\, \Pi_{\mathrm{L}}^{(\ell)}\, \mathsf{P}_{\mathrm{phys}} = 0, \tag{7.11}$$

so neither the Faddeev–Popov ghosts nor longitudinal gauge modes contribute to physical matrix elements.

Proof sketch. (a) *BRST exactness and Ward structure.* By the adjoint-covariant Ward identity and its physical-sector version (eqs. (6.375) and (6.377)), ζ-exact insertions drop out in physical correlators; in the unbroken Quadrant I, explicit breakers vanish and conservation is exact on $\mathcal{H}_{\mathrm{phys}}$.

(b) *Quartet mechanism.* The Kugo–Ojima construction organises (longitudinal, b; c, $\bar{c}$) into BRST doublets (quartets). States created by BRST-exact operators are null and orthogonal to $\mathcal{H}_{\mathrm{phys}}$; hence the ghost sector and the longitudinal projector act trivially when sandwiched by $\mathsf{P}_{\mathrm{phys}}$.

(c) *Band Hessian block structure.* In eq. (7.10), the ghost block appears only as $-\bar{c}_{\lambda_{\mathrm{UV}},\ell}\, \Box_{\mathrm{FP}}^{(\lambda_{\mathrm{UV}},\ell)}\, c_{\lambda_{\mathrm{UV}},\ell}$ with no overlap with the transverse channel, and the longitudinal projector lives in the pure-gauge block. Functoriality of the channel decomposition ensures BRST-exact improvements do not mix into the physical transverse block. Combine (a)–(c) to obtain (7.11). □

Corollary 7.23 (Positivity of the physical quadratic form)**.** *If the transverse blocks of $\mathbb{H}^{(\lambda_{\mathrm{UV}},\ell)}$ are positive (semi)definite in the induced inner product, then the band-limited quadratic form restricted to $\mathcal{H}_{\mathrm{phys}}$ is positive (semi)definite:*

$$\langle \psi,\, \mathbb{H}^{(\lambda_{\mathrm{UV}},\ell)} \psi \rangle \;=\; \langle \psi,\, \Pi_{\mathrm{T}}^{(\ell)} \mathbb{H}^{(\lambda_{\mathrm{UV}},\ell)} \Pi_{\mathrm{T}}^{(\ell)} \psi \rangle \;\geq 0, \qquad \forall\, \psi \in \mathcal{H}_{\mathrm{phys}}.$$

By Nielsen independence, this statement is invariant under ζ-exact gauge-fermion changes provided $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ are BRST-closed and commute with the channel projectors.

ROI-weighted variant (optional). If one prefers to place the IR window explicitly in the gauge-fixing functional, use the ROI-weighted $\Psi^{(\ell)}$ and UV-only ghosts as in the variant at the end of the transverse/longitudinal paragraph; then (7.11) holds with $\Box_{\mathrm{FP}}^{(\lambda_{\mathrm{UV}})}$ in place of $\Box_{\mathrm{FP}}^{(\lambda_{\mathrm{UV}},\ell)}$.

Corollary 7.24 (Gauge-invariant Källén–Lehmann representation with positivity)**.** *For any gauge-invariant observable $\mathcal{O} \in H^0(\hat{Q}_{\mathrm{BRST}})$, the two-point function evaluated on $\mathcal{H}_{\mathrm{phys}}$ in the band setup admits a Källén–Lehmann representation*

$$\left\langle\!\!\left\langle \mathcal{O}(p)\,\mathcal{O}(-p) \right\rangle\!\!\right\rangle_{\mathcal{H}_{\mathrm{phys}}}^{(\lambda_{\mathrm{UV}},\ell)} = \int_0^\infty d\mu^2\, \frac{\rho_{\mathcal{O}}^{\mathrm{phys};(\lambda_{\mathrm{UV}},\ell)}(\mu^2)}{p^2+\mu^2-i0}, \qquad \rho_{\mathcal{O}}^{\mathrm{phys};(\lambda_{\mathrm{UV}},\ell)}(\mu^2) \geq 0, \tag{7.12}$$

so the spectral density is non-negative and only physical (transverse, unconfined) excitations appear in the spectrum.[1] *In particular, together with the optical theorem, (7.12) guarantees unitarity within $\mathcal{H}_{\mathrm{phys}}$. On plateau-stable bands, $\rho_{\mathcal{O}}^{\mathrm{phys};(\lambda_{\mathrm{UV}},\ell)}$ is insensitive to the regulators and converges to the untagged spectral density as $\lambda_{\mathrm{UV}}, \ell \to \infty$.*

Gauge-parameter independence. Changing the gauge fermion $\Psi \mapsto \Psi + \delta\Psi$ shifts correlators by ζ-exact terms; by the adjoint-covariant Ward identity and its physical-sector version (eqs. (6.375) and (6.377)) these drop out after projecting with $\mathsf{P}_{\mathrm{phys}}$. Hence $\mathcal{H}_{\mathrm{phys}}$ expectation values of gauge-invariant operators—and the density $\rho_{\mathcal{O}}^{\mathrm{phys};(\lambda_{\mathrm{UV}},\ell)}$—are Nielsen independent, provided $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ are BRST-closed and commute with channel projectors.

Operational upshot for Quadrant I. Evaluating law/state visibilities on $\mathcal{H}_{\mathrm{phys}}$ (holosymmetric overlay) removes unphysical sectors by construction and makes the diagnostics insensitive to gauge-parameter choices; the physical poles are read from the $\Pi_{\mathrm{T}}^{(\ell)}$-projected blocks of $\mathbb{H}^{(\lambda_{\mathrm{UV}},\ell)}$ in eq. (7.10), and the resulting plateaux reflect only the transverse, gauge-invariant content.

7.2.7 Tachyon Test and Hyperbolicity of the Physical Operator

From the band Hessian to the physical kinetic/mass blocks. Starting from the band-limited quadratic form and its block Hessian eq. (7.10), perform the transverse/longitudinal split with BRST-closed projectors,

$$\zeta\big(\Pi_{\mathrm{T,L}}^{(\ell)}\big) = 0, \qquad [\Pi_{\mathrm{T,L}}^{(\ell)}, P_x(\lambda_{\mathrm{UV}})] = 0,$$

and insert the BRST projector $\mathsf{P}_{\mathrm{phys}}$ as in Sections 7.2.5, 7.2.6 and 7.5.13. The physical kinetic operator is read from the $\Pi_{\mathrm{T}}^{(\ell)}$–projected Φ/Θ blocks,

$$\mathcal{K}_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)} := \mathsf{P}_{\mathrm{phys}}\,\Pi_{\mathrm{T}}^{(\ell)} \begin{pmatrix} \mathcal{K}_T^{(\lambda_{\mathrm{UV}},\ell)} & 0 \\ 0 & \mathcal{K}_C^{(\lambda_{\mathrm{UV}},\ell)} \end{pmatrix} \Pi_{\mathrm{T}}^{(\ell)}\,\mathsf{P}_{\mathrm{phys}},$$

while the physical mass/mixing is captured by the band-projected Hessian blocks. Equivalently, the *physical mass operator* is (restating (7.13))

$$\begin{aligned} \mathbb{M}^2_{\mathrm{phys}}(\lambda_{\mathrm{UV}},\ell) &:= \mathsf{P}_{\mathrm{phys}}\Big(\mathcal{M}^{(\lambda_{\mathrm{UV}},\ell)} - \mathcal{M}^{(\lambda_{\mathrm{UV}},\ell)}\Pi_{\mathrm{L}}^{(\ell)} - \Pi_{\mathrm{L}}^{(\ell)}\mathcal{M}^{(\lambda_{\mathrm{UV}},\ell)}\Big)\mathsf{P}_{\mathrm{phys}}, \\ \mathcal{M}^{(\lambda_{\mathrm{UV}},\ell)} &:= \begin{pmatrix} \mathcal{M}_{\Phi\Phi}^{(\lambda_{\mathrm{UV}},\ell)} & \mathcal{M}_{\Phi\Theta}^{(\lambda_{\mathrm{UV}},\ell)} \\ \mathcal{M}_{\Theta\Phi}^{(\lambda_{\mathrm{UV}},\ell)} & \mathcal{M}_{\Theta\Theta}^{(\lambda_{\mathrm{UV}},\ell)} \end{pmatrix}. \end{aligned} \tag{7.13}$$

[1] With Euclidean momentum p_E, replace the denominator by $(p_E^2+\mu^2)$; with Minkowski signature $(+,-,\dots)$ one may write $(p^2-\mu^2+i0)^{-1}$. The positivity statement is unaffected.

By construction, $\mathbb{M}^2_{\mathrm{phys}}$ is self-adjoint on $\mathcal{H}_{\mathrm{phys}}$ with the band inner product (cf. Section 7.2.5); we also assume $[\nabla_\mu, P_x(\lambda_{\mathrm{UV}})] = 0$ so covariant derivatives commute with the UV projection.

Tachyon test (spectral non-negativity). Absence of tachyons is certified by the *non-negativity* of the spectrum of the physical Klein–Gordon–type operator,

$$\mathrm{Spec}\Big(-\bar{\Sigma}^{(\ell)\mu\nu}\nabla_\mu\nabla_\nu \otimes \mathbb{1}_{\mathcal{F}} + \mathbb{M}^2_{\mathrm{phys}}(\lambda_{\mathrm{UV}}, \ell)\Big) \subset [0,\infty) \quad \text{on } \mathcal{H}_{\mathrm{phys}}. \tag{7.14}$$

In practice, one diagonalises the $\Pi^{(\ell)}_{\mathrm{T}}$–projected blocks of $\mathbb{H}^{(\lambda_{\mathrm{UV}},\ell)}$ (7.10) to extract the pole structure; positivity of the Källén–Lehmann density on $\mathcal{H}_{\mathrm{phys}}$ (eq. (7.12)) ensures that negative mass-squares cannot be hidden by gauge artefacts or ghosts (cf. Section 7.2.6).

Hyperbolicity (principal symbol and cone-compatibility). Well-posedness and causal propagation are guaranteed when the *principal symbol* of the physical kinetic operator is positive on timelike covectors of the emergent chronogeometry:

$$\sigma_{\mathrm{pr}}\Big(\bar{\Sigma}^{(\ell)\mu\nu}\nabla_\mu\nabla_\nu\Big)(k) = \bar{\Sigma}^{(\ell)\mu\nu}k_\mu k_\nu \;>\; 0 \quad \text{for timelike } k^\mu, \tag{7.15}$$

with $\bar{\Sigma}^{(\ell)\mu\nu}$ the inverse of $\bar{\Sigma}^{(\ell)}_{\mu\nu}$ and ∇ the Relational Levi–Civita connection of Definition 6.7. Under this condition the Cauchy problem is hyperbolic in the sense of energy estimates: characteristic covectors coincide with the $\bar{\Sigma}^{(\ell)}$–null directions, and solutions propagate inside the chronogeometric light cones fixed by $\bar{\Sigma}^{(\ell)}_{\mu\nu}$.

Gauge/Nielsen stability and plateau protection. Since changes of the gauge fermion are BRST-exact, the *physical* eigenvalues of (7.14) are gauge-parameter independent (Nielsen identity; cf. Sections 7.2.1 and 7.2.6). Moreover, within any certified plateau region (Section 7.3), D_λ and D_ℓ derivatives of the pole locations remain bounded by the tolerance criteria, ensuring UV/IR robustness of the tachyon and hyperbolicity tests (cf. Sections 7.3 and 7.5.13).

Operational procedure (summary).

1. Build $\mathbb{H}^{(\lambda_{\mathrm{UV}},\ell)}$ and perform the $\Pi^{(\ell)}_{\mathrm{T}}/\Pi^{(\ell)}_{\mathrm{L}}$ split; verify eq. (7.11).
2. Compute $\mathbb{M}^2_{\mathrm{phys}}(\lambda_{\mathrm{UV}},\ell)$ via (7.13); enforce (7.14).
3. Check (7.15) using $\bar{\Sigma}^{(\ell)}$ and the connection of Definition 6.7; confirm cone-compatibility with the microcausality ledger.

7.2.8 Anomaly Filters and Quantum BRST

Definition 7.25 (Wess–Zumino counterterm and descent notation)**.** At loop order n, a *Wess–Zumino counterterm* is a local functional $\mathsf{Z}^{(n)}$ of UV dimension d and ghost number 0 (i.e.

$\mathfrak{g}(\mathsf{Z}^{(n)}) = 0$, equivalently $\mathrm{gh}(\mathsf{Z}^{(n)}) = 0$) such that the order–$n$ anomalous BRST variation $\mathcal{A}^{(n)}$ satisfies the descent relation

$$\mathcal{A}^{(n)} \;=\; \zeta\, \mathsf{Z}^{(n)} \;+\; d(\cdots),$$

where ζ is the BRST differential and d is the exterior derivative on spacetime forms. In particular, $\zeta\, \mathcal{A}^{(n)} = 0$ so $\mathcal{A}^{(n)} \in H^{1,d}(\zeta \,|\, d)$, and $\mathcal{A}^{(n)}$ is *cohomologically trivial* iff such a $\mathsf{Z}^{(n)}$ exists.

The counterterm removes the anomaly at order n by the local shift

$$\Gamma^{(\lambda_{\mathrm{UV}},\ell)} \;\longmapsto\; \Gamma^{(\lambda_{\mathrm{UV}},\ell)} \;-\; \hbar^n \int d^d x\, \sqrt{-\det \bar{\Sigma}^{(\ell)}(x)}\; \mathsf{Z}^{(n)}(x),$$

thereby restoring eq. (7.16) at that order. The representative $\mathsf{Z}^{(n)}$ is defined up to BRST-exact and total-derivative redefinitions, $\mathsf{Z}^{(n)} \sim \mathsf{Z}^{(n)} + \zeta Y^{(n)} + d(\cdots)$, which leave $\mathcal{A}^{(n)}$ unchanged.

When several orders are relevant, we write the formal sum $\mathsf{Z} := \sum_{n\geq 1} \hbar^n \mathsf{Z}^{(n)}$, so that the anomaly series reads $\Delta_{\mathrm{BRST}}\Gamma = \zeta\, \mathsf{Z} + d(\cdots)$.

Quantum BRST invariance is encoded by the vanishing of the Slavnov functional (equivalently, vanishing anomalous BRST variation) of the renormalised effective action,

$$\begin{aligned}
&\mathcal{S}\big(\Gamma^{(\lambda_{\mathrm{UV}},\ell)}\big) = 0,\\
\Delta_{\mathrm{BRST}}\Gamma^{(\lambda_{\mathrm{UV}},\ell)} &:= \int d^d x\, \sqrt{-\det \bar{\Sigma}^{(\ell)}(x)}\, \Big\langle \frac{\delta\Gamma^{(\lambda_{\mathrm{UV}},\ell)}}{\delta \boldsymbol{\Phi}_{\lambda_{\mathrm{UV}},\ell}}\, \zeta \boldsymbol{\Phi}_{\lambda_{\mathrm{UV}},\ell} + \frac{\delta\Gamma^{(\lambda_{\mathrm{UV}},\ell)}}{\delta \boldsymbol{\Theta}_{\lambda_{\mathrm{UV}},\ell}}\, \zeta \boldsymbol{\Theta}_{\lambda_{\mathrm{UV}},\ell}\\
&\qquad + \frac{\delta\Gamma^{(\lambda_{\mathrm{UV}},\ell)}}{\delta c_{\lambda_{\mathrm{UV}},\ell}}\, \zeta c_{\lambda_{\mathrm{UV}},\ell} + \frac{\delta\Gamma^{(\lambda_{\mathrm{UV}},\ell)}}{\delta \bar{c}_{\lambda_{\mathrm{UV}},\ell}}\, \zeta \bar{c}_{\lambda_{\mathrm{UV}},\ell}\\
&\qquad + \frac{\delta\Gamma^{(\lambda_{\mathrm{UV}},\ell)}}{\delta b_{\lambda_{\mathrm{UV}},\ell}}\, \zeta b_{\lambda_{\mathrm{UV}},\ell} \Big\rangle = 0,
\end{aligned} \tag{7.16}$$

where ζ is the nilpotent (ghost number $+1$) BRST differential (Section 7.2.1), the bold multiplets denote the classical mean fields in the effective action, and $\langle\cdot\rangle$ indicates the band/plateau evaluation (Section 7.3). Ghost numbers are $\mathrm{gh}(\boldsymbol{\Phi}) = \mathrm{gh}(\boldsymbol{\Theta}) = \mathrm{gh}(b) = 0$, $\mathrm{gh}(c) = +1$, $\mathrm{gh}(\bar{c}) = -1$; the scale knobs are BRST-closed: $\zeta P_x(\lambda_{\mathrm{UV}}) = 0$, $\zeta K_{\mathrm{IR}}^{(\ell)} = 0$.

In the loop expansion,

$$\Delta_{\mathrm{BRST}}\Gamma^{(\lambda_{\mathrm{UV}},\ell)} = \hbar\, \mathcal{A}^{(1)} + \hbar^2\, \mathcal{A}^{(2)} + \cdots,$$

with $\mathcal{A}^{(n)}$ local functionals of ghost number $+1$ and UV dimension d. The *Wess–Zumino consistency condition* states

$$\zeta\, \mathcal{A}^{(n)} = 0 \qquad \Rightarrow \qquad \mathcal{A}^{(n)} \in H^{1,d}(\zeta \,|\, d),$$

so a nonzero anomaly represents a nontrivial BRST cohomology class. If $\mathcal{A}^{(n)} = \zeta\, \mathsf{Z}^{(n)} + d(\cdots)$, with $\mathsf{Z}^{(n)}$ a local counterterm (ghost number 0), the anomaly is removable by a local counterterm and (7.16) can be restored at that order [442, 443, 91, 243]. On plateau-stable bands, the (co)homology class of $\mathcal{A}^{(n)}$ is regulator-independent and converges to the untagged result as $\lambda_{\mathrm{UV}}, \ell \to \infty$.

ROI-weighted variant (optional). If one prefers to include the IR window explicitly in the Slavnov functional, use

$$\mathcal{S}^{(\ell)}\Big(\Gamma^{(\lambda_{\mathrm{UV}})}\Big) := \int d^d x\, K_{\mathrm{IR}}^{(\ell)}(x)\, \sqrt{-\det \bar{\Sigma}^{(\ell)}(x)}\, \Big\langle \cdots \Big\rangle,$$

and adopt *UV-only* tags inside the brackets (e.g. $\mathbf{\Phi}_{\lambda_{\mathrm{UV}}}$, $c_{\lambda_{\mathrm{UV}}}$, etc.) to avoid double IR windowing. Since $K_{\mathrm{IR}}^{(\ell)}$ has ghost number 0 and is BRST-closed, Nielsen independence and the cohomological classification of anomalies are preserved. at that order [442, 443, 91, 243].

$$\begin{aligned}
&\mathcal{S}\Big(\Gamma^{(\lambda_{\mathrm{UV}},\ell)}\Big) = 0,\\
\Delta_{\mathrm{BRST}}\Gamma^{(\lambda_{\mathrm{UV}},\ell)} &:= \int d^d x\, \sqrt{-\det \bar{\Sigma}^{(\ell)}(x)}\, \Big\langle \frac{\delta\Gamma^{(\lambda_{\mathrm{UV}},\ell)}}{\delta \mathbf{\Phi}_{\lambda_{\mathrm{UV}},\ell}(x)}\, \zeta\mathbf{\Phi}_{\lambda_{\mathrm{UV}},\ell}(x) + \frac{\delta\Gamma^{(\lambda_{\mathrm{UV}},\ell)}}{\delta \mathbf{\Theta}_{\lambda_{\mathrm{UV}},\ell}(x)}\, \zeta\mathbf{\Theta}_{\lambda_{\mathrm{UV}},\ell}(x)\\
&\quad + \frac{\delta\Gamma^{(\lambda_{\mathrm{UV}},\ell)}}{\delta c_{\lambda_{\mathrm{UV}},\ell}(x)}\, \zeta c_{\lambda_{\mathrm{UV}},\ell}(x) + \frac{\delta\Gamma^{(\lambda_{\mathrm{UV}},\ell)}}{\delta \bar{c}_{\lambda_{\mathrm{UV}},\ell}(x)}\, \zeta \bar{c}_{\lambda_{\mathrm{UV}},\ell}(x)\\
&\quad + \frac{\delta\Gamma^{(\lambda_{\mathrm{UV}},\ell)}}{\delta b_{\lambda_{\mathrm{UV}},\ell}(x)}\, \zeta b_{\lambda_{\mathrm{UV}},\ell}(x) \Big\rangle = 0.
\end{aligned} \tag{7.17}$$

Anomaly filters as algebraic constraints. We package the potential obstructions into three classes and impose their vanishing as *filters*:

$$\mathfrak{A}_{\mathrm{gauge}} = \mathbf{0}, \qquad \mathfrak{A}_{\mathrm{mixed}} = \mathbf{0}, \qquad \mathfrak{A}_{\mathrm{grav}} = \mathbf{0}, \tag{7.18}$$

where:

- **Gauge (cubic) anomalies.** In $d = 4$,

$$\sum_{\text{chiral reps } R} \mathrm{Tr}_R\Big(T^a\{T^b, T^c\}\Big) = 0 \quad (\forall\, a, b, c),$$

  together with the Abelian constraints $\sum_i q_i^3 = 0$ for each $U(1)$, and the mixed Abelian constraints $\sum_i q_i^2 q_i' = 0$ for every $U(1)-U(1)'$ cubic. Include also the *mixed* $U(1)-G^2$ gauge condition

$$\sum_i q_i\, \mathrm{Tr}_{R_i}\Big(T^a T^b\Big) \propto \delta^{ab} \;\Rightarrow\; \sum_i q_i\, \ell_2(R_i^G) = 0 \quad \text{for each non-Abelian factor } G,$$

  with ℓ_2 the Dynkin index.

- **Gauge–gravitational anomalies.** In $d = 4$ (Levi–Civita connection), this reduces to

$$\sum_i q_i = 0 \quad \text{for each } U(1).$$

  Non-Abelian factors automatically satisfy $\mathrm{Tr}_R(T^a) = 0$, so there is no non-Abelian gauge–gravitational anomaly in four dimensions.

- **Purely gravitational anomalies.** Relevant in $d \equiv 2 \,(\mathrm{mod}\, 4)$; in even dimensions they are read from the $\hat{A}$ genus and Pontryagin classes in the anomaly polynomial via descent.

All three are computed uniformly from the anomaly polynomial $\mathcal{I}_{d+2}(F, R)$ and its descent,

$$\mathcal{I}_{d+2} = d\,\mathcal{I}^{(0)}_{d+1}, \qquad \zeta\,\mathcal{I}^{(0)}_{d+1} + d\,\mathcal{I}^{(1)}_{d} = 0, \qquad \mathcal{A}^{(1)} = \int \mathcal{I}^{(1)}_{d}$$

Hence (7.18) precisely enforces the triviality of $[\,\mathcal{A}^{(1)}\,] \in H^{1,d}\big(\zeta \,|\, d\big)$.[2]

Background-field and plateau compatibility. Using a background-field gauge for the band-limited effective action preserves background gauge invariance and organises the Slavnov–Taylor identities for Γ [114, 91]. Since anomalies are cohomological (topological) invariants, they are insensitive to the UV projector $P_x(\lambda_{\mathrm{UV}})$ and the IR window $K^{(\ell)}_{\mathrm{IR}}$: once (7.18) holds, $\mathcal{S}(\Gamma) = 0$ persists across plateaus (Section 7.3).

From filters to quantum BRST nilpotency and physical unitarity. With (7.18) enforced, (7.16) implies (i) quantum-level nilpotency of the BRST charge and preservation of eq. (7.196) after renormalisation; (ii) gauge-parameter independence of physical poles via Nielsen identities (Section 7.2.9); and (iii) consistency of the ghost decoupling/positivity statements in Section 7.2.6 (no hidden negative-norm contributions), ensuring that the Källén–Lehmann densities for gauge-invariant operators remain positive on $\mathcal{H}_{\mathrm{phys}}$ (eq. (7.12)).

Operational checklist (model-building view).

1. **List chiral matter:** record all irreducible reps (including $U(1)$ charges and multiplicities).
2. **Compute indices:** evaluate $\mathrm{Tr}_R\big(T^a\{T^b, T^c\}\big)$, the $U(1)$ cubic/mixed sums, and the $U(1)-G^2$ constraints via $\ell_2(R)$; assemble $\mathfrak{A}_{\mathrm{gauge}}$.
3. **Build $\mathcal{I}_{d+2}(F, R)$:** extract $\mathfrak{A}_{\mathrm{mixed}}$ and $\mathfrak{A}_{\mathrm{grav}}$ from the descent of $\mathcal{I}_{d+2}$.
4. **Enforce $\mathfrak{A} = 0$:** if needed, add counter-matter or adjust couplings to satisfy (7.18); re-check $\mathcal{S}(\Gamma) = 0$ on the plateau.

7.2.9 Gauge-parameter Independence of the Physical Poles (Nielsen Identities)

Let α denote any c-number gauge parameter entering the gauge-fixing fermion $\Psi(\alpha)$ in the band-limited source formalism of eq. (7.144). To control α-variations algebraically, extend BRST by a doublet (ghost numbers $\mathrm{gh}(\alpha) = 0$, $\mathrm{gh}(\chi) = +1$):

$$\zeta\,\alpha = \chi, \qquad \zeta\,\chi = 0,$$

[2] These filters address local (perturbative) anomalies. Global anomalies (e.g. the $SU(2)$ Witten anomaly in $d = 4$) require additional checks using $\pi_{d+1}(G)$ or cobordism invariants; they should be imposed separately when relevant.

and treat $\partial_\alpha \Psi$ as the insertion generated by the antifield/source sector $\mathcal{K}\cdot\zeta\varphi$ (cf. eq. (7.144)). With the Slavnov functional $\mathcal{S}(\Gamma) = 0$ eq. (7.16), one obtains the *Nielsen identity* for the renormalised effective action $\Gamma^{(\lambda_{\mathrm{UV}},\ell)}$:

$$\partial_\alpha \Gamma^{(\lambda_{\mathrm{UV}},\ell)} = \int d^d x \, \frac{\delta \Gamma^{(\lambda_{\mathrm{UV}},\ell)}}{\delta \Phi^A(x)} \, \mathcal{N}^A(\alpha; x) \; + \; (\text{ghost/aux. terms}) \,, \qquad \mathcal{N}^A := \left\langle \zeta \, \frac{\partial \Psi}{\partial \alpha} \right\rangle_\Gamma , \tag{7.19}$$

where A runs over the fluctuation multiplet $\mathfrak{X}$; the expectation is 1PI, band-safe, and BRST-closed (cf. Section 7.2.1). Band knobs are BRST-closed: $\zeta P_x(\lambda_{\mathrm{UV}}) = 0$, $\zeta K_{\mathrm{IR}}^{(\ell)} = 0$.

Linearised (two-point) form on the physical subspace. Project to the transverse sector and the physical Hilbert space,

$$\Gamma^{(\mathrm{TT})}(p^2) := \Pi_{\mathrm{T}}^{(\ell)} \, \Gamma^{(\lambda_{\mathrm{UV}},\ell)}(p^2) \, \Pi_{\mathrm{T}}^{(\ell)} \Big|_{\mathcal{H}_{\mathrm{phys}}} ,$$

which removes the ghost/longitudinal contributions (quartet decoupling, Section 7.2.6). Linearising (7.19) for the 1PI two-point function in a $\Pi_{\mathrm{T}}^{(\ell)}$ basis gives

$$\partial_\alpha \, \Gamma^{(\mathrm{TT})}_{ij}(p^2) = \Gamma^{(\mathrm{TT})}_{ik}(p^2) \, \mathcal{C}_{kj}(p^2) + \mathcal{C}_{ik}(p^2) \, \Gamma^{(\mathrm{TT})}_{kj}(p^2), \tag{7.20}$$

where $\mathcal{C}$ is a local matrix built from the insertion $\zeta(\partial_\alpha \Psi)$ and antifield couplings (the $\mathcal{K}$-sector in eq. (7.144)).

Simple poles (isolated eigenmodes). Let $p^2 = -m_i^2$ be a simple zero of $\det \Gamma^{(\mathrm{TT})}(p^2)$ with right/left eigenvectors $v_i, \bar{v}_i$,

$$\Gamma^{(\mathrm{TT})}(-m_i^2) \, v_i = 0, \qquad \bar{v}_i \, \Gamma^{(\mathrm{TT})}(-m_i^2) = 0, \qquad \bar{v}_i \, \partial_{p^2} \Gamma^{(\mathrm{TT})}(-m_i^2) \, v_i \neq 0.$$

Differentiating the on-shell condition and using (7.20) yields

$$\Big[\bar{v}_i \, \partial_{p^2} \Gamma^{(\mathrm{TT})}(-m_i^2) \, v_i \Big] \, \partial_\alpha m_i^2 = \bar{v}_i \Big(\partial_\alpha \Gamma^{(\mathrm{TT})} \Big) v_i = \bar{v}_i \Big(\Gamma^{(\mathrm{TT})} \mathcal{C} + \mathcal{C} \, \Gamma^{(\mathrm{TT})} \Big) v_i \Big|_{p^2 = -m_i^2} = 0, \tag{7.21}$$

hence

$$\frac{\partial}{\partial \alpha} \, m_i^2(\lambda_{\mathrm{UV}}, \ell) = 0 \quad (\text{on shell in } \mathcal{H}_{\mathrm{phys}}). \tag{7.22}$$

Thus the pole masses of physical $\Pi_{\mathrm{T}}^{(\ell)}$ modes are gauge-parameter independent, provided the clean EOM eqs. (6.333) and (6.334) and quantum BRST (Slavnov identity) hold.

Degenerate poles and mixing. For an r-fold degeneracy, work with the spectral projector $\mathbb{P}_i(p^2) = \frac{1}{2\pi i} \oint_{\gamma_i} \left(\Gamma^{(\mathrm{TT})}(z) \right)^{-1} dz$ around the joint zero and define the restricted matrix $\widehat{\Gamma}_i(p^2) := \mathbb{P}_i \Gamma^{(\mathrm{TT})} \mathbb{P}_i$. Then (7.20) implies $\partial_\alpha \widehat{\Gamma}_i = \widehat{\Gamma}_i \widehat{\mathcal{C}}_i + \widehat{\mathcal{C}}_i \widehat{\Gamma}_i$, so the characteristic polynomial of $\widehat{\Gamma}_i$ is α-invariant. Consequently all roots (the r on-shell eigenvalues) are α-independent; mixing angles and residues may depend on α, but *pole positions do not*.

Propagators, residues, and widths. Writing $D^{(\mathrm{TT})} = (\Gamma^{(\mathrm{TT})})^{-1}$, the residue matrix at the pole,

$$\mathcal{R}_i := \lim_{p^2 \to -m_i^2} (p^2 + m_i^2)\, D^{(\mathrm{TT})}(p^2) = \frac{v_i\, \bar{v}_i}{\bar{v}_i\, \partial_{p^2}\Gamma^{(\mathrm{TT})}(-m_i^2)\, v_i},$$

is generally α-dependent (field renormalisations and mixings are gauge dependent). For unstable states, the complex pole $s_i = M_i^2 - iM_i\Gamma_i$ obeys a Nielsen identity of the same form; s_i is gauge independent to all orders where a resummed 1PI description applies.

Mass-operator statement. Equivalently, the spectrum of the physical mass operator eq. (7.13) is gauge-parameter stable:

$$\partial_\alpha \operatorname{Spec}\Big[\mathbb{M}^2_{\mathrm{phys}}(\lambda_{\mathrm{UV}}, \ell)\Big] = 0 \quad (\text{on shell in } \mathcal{H}_{\mathrm{phys}}). \tag{7.23}$$

Remarks. (i) Off-shell Green functions and individual field renormalisations *can* depend on α; (7.22) isolates on-shell pole positions. (ii) In the bandwise setting, $P_x(\lambda_{\mathrm{UV}})$ and $K^{(\ell)}_{\mathrm{IR}}$ commute with BRST on insertions (ghost number 0), so α-variations are ζ-exact on $\mathcal{H}_{\mathrm{phys}}$ (see Lemma 6.243). (iii) Anomaly filters eq. (7.18) and eq. (7.16) are required to guarantee the cohomological control used in (7.19)–(7.21). (iv) On plateau-stable bands (Section 7.3), the pole locations are additionally stable under (D_λ, D_ℓ) scans, providing UV/IR robustness alongside gauge-parameter independence.
BV/canonical viewpoint (optional). In the BV language, ∂_α acts as a canonical transformation generated by a local functional proportional to $\partial_\alpha\Psi$; (7.19) is the statement that ∂_α is a BRST-exact canonical flow. Since canonical transformations do not move the zeros of $\Gamma^{(\mathrm{TT})}$ on $\mathcal{H}_{\mathrm{phys}}$, the pole positions are invariant under α.

7.2.10 Bandwise Checklist and Certification

Fix a candidate plateau region $\mathfrak{P} \subset \mathbb{R}_{>0} \times \mathbb{R}_{>0}$ in the $(\lambda_{\mathrm{UV}}, \ell)$ plane (cf. Section 6.12.4). For each $(\lambda_{\mathrm{UV}}, \ell) \in \mathfrak{P}$, certify the Quadrant I baseline by verifying *all* items below. Throughout, the scale knobs are BRST-closed and ghost-number 0: $\zeta P_x(\lambda_{\mathrm{UV}}) = 0$, $\zeta K^{(\ell)}_{\mathrm{IR}} = 0$; we assume $[\nabla_\mu, P_x(\lambda_{\mathrm{UV}})] = 0$.

C1. Classical BRST well-posedness on the band.

$$\hat{Q}^2_{\mathrm{BRST}} = 0, \qquad \hat{Q}^\dagger_{\mathrm{BRST}} = \hat{Q}_{\mathrm{BRST}}, \qquad [\hat{Q}_{\mathrm{BRST}}, \hat{H}_{\mathrm{tot}}] = 0, \tag{7.24}$$

with $\hat{H}_{\mathrm{tot}}$ the band-limited Hamiltonian extracted from $\mathcal{L}\big(\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell}, \hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}},\ell}\big)$. The scale-tagged EOM *eqs.* (6.333), (6.334)*and* (6.374) imply conservation of the BRST current and charge on the band:

$$\nabla_\mu J_Q^{\mu(\lambda_{\mathrm{UV}},\ell)} = 0, \qquad \frac{d}{dt}\, Q^{(\lambda_{\mathrm{UV}},\ell)}_{\mathrm{BRST}} = 0.$$

Domains are invariant under the knobs: $P_x(\lambda_{\mathrm{UV}})\mathcal{D}(Q) \subset \mathcal{D}(Q)$, $K^{(\ell)}_{\mathrm{IR}}\mathcal{D}(Q) \subset \mathcal{D}(Q)$.

C2. Quadratic kernel and $\Pi_{\mathrm{T}}/\Pi_{\mathrm{L}}$ split. Build the band Hessian $\mathbb{H}^{(\lambda_{\mathrm{UV}},\ell)}$ (second variation of the action at the symmetric background). Use projectors $\Pi_{\mathrm{T}}^{(\ell)}, \Pi_{\mathrm{L}}^{(\ell)}$ with

$$\left(\Pi_{\mathrm{T,L}}^{(\ell)}\right)^2 = \Pi_{\mathrm{T,L}}^{(\ell)}, \quad \Pi_{\mathrm{T}}^{(\ell)} + \Pi_{\mathrm{L}}^{(\ell)} = \mathbb{1}, \quad \zeta\left(\Pi_{\mathrm{T,L}}^{(\ell)}\right) = 0, \quad [\Pi_{\mathrm{T,L}}^{(\ell)}, P_x(\lambda_{\mathrm{UV}})] = 0.$$

Verify Kugo–Ojima quartet decoupling on $\mathcal{H}_{\mathrm{phys}}$ (cf. *Section* 7.2.6):

$$\mathsf{P}_{\mathrm{phys}} \Box_{\mathrm{FP}}^{(\lambda_{\mathrm{UV}},\ell)} \mathsf{P}_{\mathrm{phys}} = 0, \qquad \mathsf{P}_{\mathrm{phys}} \Pi_{\mathrm{L}}^{(\ell)} \mathsf{P}_{\mathrm{phys}} = 0.$$

C3. Physical mass operator & tachyon test. Form the physical mass matrix from the potential blocks:

$$\mathbb{M}^2_{\mathrm{phys}}(\lambda_{\mathrm{UV}}, \ell) := \mathsf{P}_{\mathrm{phys}}\left(\mathcal{M}^{(\lambda_{\mathrm{UV}},\ell)} - \mathcal{M}^{(\lambda_{\mathrm{UV}},\ell)}\Pi_{\mathrm{L}}^{(\ell)} - \Pi_{\mathrm{L}}^{(\ell)}\mathcal{M}^{(\lambda_{\mathrm{UV}},\ell)}\right)\mathsf{P}_{\mathrm{phys}}$$

with $\mathcal{M}^{(\lambda_{\mathrm{UV}},\ell)}$ as in *eq.* (7.13). Demand self-adjointness on $\mathcal{H}_{\mathrm{phys}}$ and spectral non-negativity of the physical Klein–Gordon-type operator:

$$\mathrm{Spec}\Big(- \bar{\Sigma}^{(\ell)\mu\nu}\nabla_\mu\nabla_\nu \otimes \mathbb{1} + \mathbb{M}^2_{\mathrm{phys}}(\lambda_{\mathrm{UV}}, \ell)\Big) \subset [0, \infty). \tag{7.25}$$

Equivalently, all physical poles extracted from the $\Pi_{\mathrm{T}}^{(\ell)}$-projected two-point function satisfy $m_i^2 \geq 0$.

C4. Hyperbolicity and microcausality. The principal symbol in the emergent chronogeometry is positive on timelike covectors:

$$\sigma_{\mathrm{pr}}\left(\bar{\Sigma}^{(\ell)\mu\nu}\nabla_\mu\nabla_\nu\right)(k) = \bar{\Sigma}^{(\ell)\mu\nu}k_\mu k_\nu > 0 \quad \text{for timelike } k^\mu. \tag{7.26}$$

Require the standard energy estimate for compactly supported Cauchy data ψ:

$$\frac{d}{dt}\mathcal{E}^{(\lambda_{\mathrm{UV}},\ell)}[\psi](t) \leq C\,\mathcal{E}^{(\lambda_{\mathrm{UV}},\ell)}[\psi](t),$$
$$\mathcal{E}^{(\lambda_{\mathrm{UV}},\ell)}[\psi](t) := \int_{\Sigma_t} \sqrt{h^{(\ell)}}\left(\dot{\psi}^\dagger\dot{\psi} + \nabla\psi^\dagger\nabla\psi + \psi^\dagger\mathbb{M}^2_{\mathrm{phys}}\psi\right),$$

and the microcausal support property: commutators of physical fields vanish at $\bar{\Sigma}^{(\ell)}$-spacelike separation.

C5. Quantum BRST (anomaly filters + Slavnov). Enforce the algebraic filters $\mathfrak{A}_{\mathrm{gauge}} = \mathbf{0}$, $\mathfrak{A}_{\mathrm{mixed}} = \mathbf{0}$, $\mathfrak{A}_{\mathrm{grav}} = \mathbf{0}$ $\big($*eq.* (7.18)$\big)$ so that the 1-loop cocycle $[\mathcal{A}^{(1)}] \in H^{1,d}(\zeta \mid d)$ is trivial. Then the Slavnov identity holds for the renormalised $\Gamma^{(\lambda_{\mathrm{UV}},\ell)}$ $\big($*eq.* (7.16)$\big)$, and all Ward identities descend to $\mathcal{H}_{\mathrm{phys}}$ modulo ζ-exact terms.

C6. Nielsen certification (gauge-parameter independence). Extend BRST with the (α, χ) doublet and derive the Nielsen identity

$$\partial_\alpha\Gamma^{(\lambda_{\mathrm{UV}},\ell)} = \int d^dx\, \frac{\delta\Gamma^{(\lambda_{\mathrm{UV}},\ell)}}{\delta\Phi^A}\mathcal{N}^A + \text{ghost/aux.}$$

Linearise on the $\Pi_{\mathrm{T}}^{(\ell)}$ subspace to obtain $\partial_\alpha\Gamma^{(\mathrm{TT})} = \Gamma^{(\mathrm{TT})}\mathcal{C} + \mathcal{C}\,\Gamma^{(\mathrm{TT})}$ $\big($*eq.* (7.20)$\big)$, and verify for each isolated pole $p^2 = -m_i^2$:

$$\partial_\alpha m_i^2(\lambda_{\mathrm{UV}}, \ell) = 0 \quad \text{on shell in } \mathcal{H}_{\mathrm{phys}} \quad \big(\textit{eqs. (7.22) and (7.23)}\big). \tag{7.27}$$

Plateau diagnostics (numeric certification). Introduce the log-derivatives $D_\lambda := \lambda_{\mathrm{UV}}\partial_{\lambda_{\mathrm{UV}}}$, $D_\ell := \ell\,\partial_\ell$. For any band-observable O (e.g. a physical pole m_i^2, a residue trace, or a transverse norm), define

$$\mathrm{Slope}_\alpha[O] := \sup_{(\lambda_{\mathrm{UV}},\ell)\in\mathfrak{P}} \frac{\|D_\alpha O\|}{\|O\|+\varepsilon}, \quad \mathrm{Curv}_\alpha[O] := \sup_{(\lambda_{\mathrm{UV}},\ell)\in\mathfrak{P}} \frac{\|D_\alpha^2 O\|}{\|O\|+\varepsilon},$$

$$\mathrm{Osc}_\alpha[O] := \sup_{(\lambda_{\mathrm{UV}},\ell),(\lambda'_{\mathrm{UV}},\ell')\in\mathfrak{P}} \frac{\|O(\lambda_{\mathrm{UV}},\ell)-O(\lambda'_{\mathrm{UV}},\ell')\|}{\|O\|+\varepsilon}, \qquad \alpha\in\{\lambda,\ell\},$$

with a small $\varepsilon>0$ to avoid division by 0 on null plateaux. Certification of $\mathfrak{P}$ requires

$$\mathrm{Slope}_\alpha[O] \le \tau_O^{(1)}, \qquad \mathrm{Curv}_\alpha[O] \le \tau_O^{(2)}, \qquad \mathrm{Osc}_\alpha[O] \le \tau_O^{(\mathrm{osc})}, \tag{7.28}$$

for user-chosen tolerances $\tau_O^{(1)},\tau_O^{(2)},\tau_O^{(\mathrm{osc})}$ (typically $\ll 1$). When *eq.* (7.28) holds for all O in the Quadrant I suite (poles, spectral weights, Ward-side diagnostics), we call $\mathfrak{P}$ *plateau-stable*.

Pole extraction (spectral projector). For a possibly degenerate physical pole at $p^2=-m_i^2$, use the spectral projector

$$\mathbb{P}_i = \frac{1}{2\pi i}\oint_{\gamma_i}\Big(\Gamma^{(\mathrm{TT})}(z)\Big)^{-1}dz, \qquad \gamma_i:\ \text{small loop around } -m_i^2,$$

and read the restricted kernel $\widehat{\Gamma}_i := \mathbb{P}_i\,\Gamma^{(\mathrm{TT})}\,\mathbb{P}_i$. The eigenvalues of $\widehat{\Gamma}_i$ determine m_i^2; their α-independence follows from the bandwise Nielsen identity. Residues may vary with α, but pole *positions* do not.

Certification summary (pass/fail). A bandpoint $(\lambda_{\mathrm{UV}},\ell)$ *passes* Quadrant I certification iff **C1**–**C6** hold and the plateau metrics *eq.* (7.28) are satisfied for the selected observable set. When a point fails, report the minimal violating item(s) and the maximally offending diagnostic(s) $\{\mathrm{Slope}_\alpha,\mathrm{Curv}_\alpha,\mathrm{Osc}_\alpha\}$.

Remark (provisional mode identifications). Any assignment of labels such as "graviton-like," "Higgs-like," "apeiron," or "chrono–kymon" to physical poles remains *provisional* until plateau stability and cross-quadrant scans are passed (cf. Sections 6.12.4 and 6.12.5); do not alter existing labels/suffixes.

7.3 UV/IR Knobs and Plateau Certification

This section specifies the ultraviolet (UV) and infrared (IR) controls and provides a precise acceptance test for *plateau stability* of the holosymmetric visibilities introduced in Chapter 7, to be used downstream for mode identifications and cross-quadrant comparisons. All constructions act *on the band* and within the IR window, and are compatible with the BRST framework and the scale-tagged equations of motion eqs. (6.333), (6.334) and (6.374).

This section specifies the ultraviolet (UV) and infrared (IR) controls and provides a precise acceptance test for *plateau stability* of the holosymmetric visibilities introduced in

Chapter 7, to be used downstream for mode identifications and cross-quadrant comparisons. All constructions act *on the band* and within the IR window, and are compatible with the BRST framework and the scale-tagged equations of motion eqs. (6.333), (6.334) and (6.374).

7.3.1 Definitions of the UV/IR knobs

Definition 7.26 (UV projector (band-limit))**.** For each spacetime fibre x, let $E^{(x)}(\cdot)$ be the spectral measure of a positive, self-adjoint generator $\mathcal{D}_x$ (the local dilation/energy proxy fixed by the formalism). The ultraviolet projector is

$$P_x(\lambda_{\mathrm{UV}}) := E^{(x)}\Big([0,\lambda_{\mathrm{UV}}]\Big), \qquad \lambda_{\mathrm{UV}} > 0, \tag{7.29}$$

with the following properties:

(i) *Projector/self-adjointness/monotonicity:*

$$P_x(\lambda)^2 = P_x(\lambda), \quad P_x(\lambda)^\dagger = P_x(\lambda),$$
$$\lambda_1 \le \lambda_2 \Rightarrow P_x(\lambda_1) \le P_x(\lambda_2).$$

(ii) *Strong right-continuity and limits:* $\mathrm{s}{-}\lim_{\epsilon\downarrow 0} P_x(\lambda+\epsilon) = P_x(\lambda)$, $\mathrm{s}{-}\lim_{\lambda\to\infty} P_x(\lambda) = \mathbb{I}_x$, and $\mathrm{s}{-}\lim_{\lambda\downarrow 0} P_x(\lambda) = E^{(x)}(\{0\})$.

(iii) *BRST closure and ghost number:*

$$\Big[\hat{Q}_{\mathrm{BRST}},\, P_x(\lambda_{\mathrm{UV}})\Big] = 0, \qquad \mathrm{gh}\,\Big(P_x(\lambda_{\mathrm{UV}})\Big) = 0. \tag{7.30}$$

(iv) *Channel/background compatibility:* $[P_x(\lambda_{\mathrm{UV}}), \Pi_\alpha] = 0$ for all channel projectors Π_α, and in background-field gauge $[P_x(\lambda_{\mathrm{UV}}), \delta_a^{(\mathrm{bg})}] = 0$ so that Slavnov–Taylor identities are preserved on the band.

(v) *Action on fields/correlators (fibrewise sandwich):*

$$\begin{aligned}
\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}}} &:= P_x(\lambda_{\mathrm{UV}})\,\hat{\mathbf{\Phi}}\,P_x(\lambda_{\mathrm{UV}}),\\
\hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}}} &:= P_x(\lambda_{\mathrm{UV}})\,\hat{\mathbf{\Theta}}\,P_x(\lambda_{\mathrm{UV}}),\\
\hat{\mathbf{G}}^{(\lambda_{\mathrm{UV}})} &:= P_x(\lambda_{\mathrm{UV}})\,\hat{\mathbf{G}}\,P_x(\lambda_{\mathrm{UV}}),
\end{aligned} \tag{7.31}$$

and, for bi-local kernels,

$$\hat{\mathbf{G}}^{(\lambda_{\mathrm{UV}})}(x,x') := P_x(\lambda_{\mathrm{UV}})\,\hat{\mathbf{G}}(x,x')\,P_{x'}(\lambda_{\mathrm{UV}})\,.$$

The canonical order in composite taggings is "UV then IR."

Definition 7.27 (IR window (localisation kernel))**.** Fix a family of non-negative, normalised kernels $K_{\mathrm{IR}}^{(\ell)} \in L^1(\mathbb{R}^d)$ with characteristic width $\ell > 0$ such that

$$K_{\mathrm{IR}}^{(\ell)}(x) \ge 0, \qquad \int d^d x\, K_{\mathrm{IR}}^{(\ell)}(x) = 1, \qquad \lim_{\ell\to 0^+} K_{\mathrm{IR}}^{(\ell)} = \delta \text{ in } \mathcal{D}'(\mathbb{R}^d). \tag{7.32}$$

They act by convolution/coarse-graining on operator densities via

$$\begin{aligned}\big(K_{\mathrm{IR}}^{(\ell)}\mathcal{O}\big)(x) &:= \int d^d y\, K_{\mathrm{IR}}^{(\ell)}(x-y)\,\mathcal{O}(y),\\ \bar{\Sigma}_{\mu\nu}^{(\ell)}(x) &:= \int d^d y\, K_{\mathrm{IR}}^{(\ell)}(x-y)\,\langle\hat{\Sigma}_{\mu\nu}(y)\rangle\end{aligned} \tag{7.33}$$

with BRST closure and ghost number 0:

$$\zeta K_{\mathrm{IR}}^{(\ell)} = 0, \qquad \mathrm{gh}\big(K_{\mathrm{IR}}^{(\ell)}\big) = 0.$$

For operator-valued fields/correlators, using the canonical order "UV then IR" with $P_y(\lambda_{\mathrm{UV}})$,

$$\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell}(x) := \int d^d y\, K_{\mathrm{IR}}^{(\ell)}(x-y)\,\Big(P_y(\lambda_{\mathrm{UV}})\,\hat{\mathbf{\Phi}}(y)\,P_y(\lambda_{\mathrm{UV}})\Big), \tag{7.34}$$

$$\hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}},\ell}(x) := \int d^d y\, K_{\mathrm{IR}}^{(\ell)}(x-y)\,\Big(P_y(\lambda_{\mathrm{UV}})\,\hat{\mathbf{\Theta}}(y)\,P_y(\lambda_{\mathrm{UV}})\Big), \tag{7.35}$$

$$\begin{aligned}\hat{\mathbf{G}}^{(\lambda_{\mathrm{UV}},\ell)}(x,x') := \iint d^d y\, d^d y'\, K_{\mathrm{IR}}^{(\ell)}(x-y)\,P_y(\lambda_{\mathrm{UV}})\,\hat{\mathbf{G}}(y,y')\\ \times P_{y'}(\lambda_{\mathrm{UV}})\,K_{\mathrm{IR}}^{(\ell)}(x'-y').\end{aligned} \tag{7.36}$$

Typical choices are compactly supported mollifiers or Gaussians, such as

$$K_{\mathrm{IR}}^{(\ell)}(x) = (2\pi\ell^2)^{-d/2}\exp\!\Big[-x^2/(2\ell^2)\Big];$$

the heat-kernel semigroup variant $K_t = e^{t\Delta}$ (with $t \propto \ell^2$) additionally satisfies $K_t * K_{t'} = K_{t+t'}$.

Definition 7.28 (Joint IR/UV tagging (canonical order))**.** Given the UV projector $P_x(\lambda_{\mathrm{UV}})$ of Definition 7.26 and the IR window $K_{\mathrm{IR}}^{(\ell)}$ of Definition 6.280, the jointly tagged (band-limited and windowed) objects are defined by applying *UV then IR*:

$$\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell}(x) := \int d^d y\, K_{\mathrm{IR}}^{(\ell)}(x-y)\,\Big(P_y(\lambda_{\mathrm{UV}})\,\hat{\mathbf{\Phi}}(y)\,P_y(\lambda_{\mathrm{UV}})\Big), \tag{7.37}$$

$$\hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}},\ell}(x) := \int d^d y\, K_{\mathrm{IR}}^{(\ell)}(x-y)\,\Big(P_y(\lambda_{\mathrm{UV}})\,\hat{\mathbf{\Theta}}(y)\,P_y(\lambda_{\mathrm{UV}})\Big), \tag{7.38}$$

$$\hat{\mathbf{G}}^{(\lambda_{\mathrm{UV}},\ell)}(x,x') := \iint d^d y\, d^d y'\, K_{\mathrm{IR}}^{(\ell)}(x-y)\,P_y(\lambda_{\mathrm{UV}})\,\hat{\mathbf{G}}(y,y')\,P_{y'}(\lambda_{\mathrm{UV}})\,K_{\mathrm{IR}}^{(\ell)}(x'-y'). \tag{7.39}$$

In operator shorthand:

$$\begin{aligned}\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell} &= K_{\mathrm{IR}}^{(\ell)}\Big(P(\lambda_{\mathrm{UV}})\,\hat{\mathbf{\Phi}}\,P(\lambda_{\mathrm{UV}})\Big),\\ \hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}},\ell} &= K_{\mathrm{IR}}^{(\ell)}\Big(P(\lambda_{\mathrm{UV}})\,\hat{\mathbf{\Theta}}\,P(\lambda_{\mathrm{UV}})\Big),\\ \hat{\mathbf{G}}^{(\lambda_{\mathrm{UV}},\ell)} &= K_{\mathrm{IR}}^{(\ell)}\,\hat{\mathbf{G}}^{(\lambda_{\mathrm{UV}})}\,K_{\mathrm{IR}}^{(\ell)}.\end{aligned}$$

Basic properties.

(i) Order and limits (UV → IR).

Then the following

$$\mathrm{s}\!-\!\lim_{\lambda_{\mathrm{UV}}\to\infty}\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell} = K_{\mathrm{IR}}^{(\ell)}\hat{\mathbf{\Phi}}, \qquad \lim_{\ell\to 0^+}\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell} = \hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}}},$$

and similarly for $\hat{\mathbf{\Theta}}$ and $\hat{\mathbf{G}}$. The combined limit ($\lambda_{\mathrm{UV}}\to\infty,\ \ell\to 0^+$) recovers the bare objects.

(ii) BRST closure and ghost number.
$\zeta\, P_x(\lambda_{\mathrm{UV}}) = 0$ and $\zeta\, K_{\mathrm{IR}}^{(\ell)} = 0$, hence $\zeta\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell} = (\zeta\hat{\mathbf{\Phi}})_{\lambda_{\mathrm{UV}},\ell}$, etc., and $\mathrm{gh}(\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell}) = \mathrm{gh}(\hat{\mathbf{\Phi}})$.

(iii) Channel/background compatibility.
In background-field gauge the construction preserves Slavnov–Taylor identities on the band. In particular, the channel projectors commute with the UV projector and the IR window:

$$[P_x(\lambda_{\mathrm{UV}}), \Pi_\alpha] = 0, \qquad [K_{\mathrm{IR}}^{(\ell)}, \Pi_\alpha] = 0.$$

(iv) Covariant differentiation.
If $[\nabla_\mu, P_x(\lambda_{\mathrm{UV}})] = 0$ and $K_{\mathrm{IR}}^{(\ell)}$ is a scalar kernel, then $\nabla_\mu\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell} = (\nabla_\mu\hat{\mathbf{\Phi}})_{\lambda_{\mathrm{UV}},\ell}$ (and analogously for $\hat{\mathbf{\Theta}}$), i.e. tagging commutes with ∇.

(v) Positivity / complete positivity.
The map $\mathcal{C}_{\lambda,\ell} : \mathcal{O} \mapsto K_{\mathrm{IR}}^{(\ell)}\big(P(\lambda)\,\mathcal{O}\,P(\lambda)\big)$ is linear, *-preserving, norm–nonexpansive, and completely positive (compression by P and averaging with $K_{\mathrm{IR}}^{(\ell)} \geq 0$).

(vi) Plateau calculus.
With $D_\lambda := \lambda_{\mathrm{UV}}\partial_{\lambda_{\mathrm{UV}}}$, $D_\ell := \ell\,\partial_\ell$,

$$D_\lambda\hat{\mathbf{\Phi}}_{\lambda,\ell} = \lambda\,\partial_\lambda\Big[K_{\mathrm{IR}}^{(\ell)}\big(P(\lambda)\hat{\mathbf{\Phi}}P(\lambda)\big)\Big] = K_{\mathrm{IR}}^{(\ell)}\Big((\lambda\partial_\lambda P)\,\hat{\mathbf{\Phi}}\,P + P\,\hat{\mathbf{\Phi}}\,(\lambda\partial_\lambda P)\Big)\,,$$
$$D_\ell\hat{\mathbf{\Phi}}_{\lambda,\ell} = \ell\,(\partial_\ell K_{\mathrm{IR}}^{(\ell)})\,\big(P(\lambda)\hat{\mathbf{\Phi}}P(\lambda)\big),$$

and similarly for $\hat{\mathbf{\Theta}}$ and $\hat{\mathbf{G}}$; these feed the plateau diagnostics of Section 6.12.4.

Band functor: UV then IR.

Definition 7.29 (Band functor: UV then IR)**.** For any operator density $\mathcal{O}(y)$ define the bandwise map

$$\big(\mathcal{B}_{\lambda_{\mathrm{UV}},\ell}[\mathcal{O}]\big)(x) := \int d^d y\, K_{\mathrm{IR}}^{(\ell)}(x-y)\, P_y(\lambda_{\mathrm{UV}})\,\mathcal{O}(y)\,P_y(\lambda_{\mathrm{UV}})\,, \tag{7.40}$$

with $P_y(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ as in Definitions 6.280 and 7.26. We write $\mathcal{B}^{(\lambda_{\mathrm{UV}},\ell)} := \mathcal{B}_{\lambda_{\mathrm{UV}},\ell}$ for brevity.

Proposition 7.30 (Band safety and functoriality)**.** *The map $\mathcal{B}^{(\lambda_{\mathrm{UV}},\ell)}$ enjoys:*

(i) Linearity, *-preservation, positivity.
$\mathcal{B}[a\mathcal{O} + b\mathcal{P}] = a\,\mathcal{B}[\mathcal{O}] + b\,\mathcal{B}[\mathcal{P}]$, $\mathcal{B}[\mathcal{O}^\dagger] = \mathcal{B}[\mathcal{O}]^\dagger$, *and* $\mathcal{O} \geq 0 \Rightarrow \mathcal{B}[\mathcal{O}] \geq 0$.

(ii) Complete positivity (CP).
For any finite family $\{X_j\}$, $\sum_{j,k} X_j^\dagger\,\mathcal{B}[\mathcal{O}_{jk}]\,X_k \geq 0$ whenever $(\mathcal{O}_{jk}) \geq 0$. (Composition of a projector compression with averaging by a positive kernel is CP.)

(iii) BRST closure and ghost number.
$\zeta\, P_y(\lambda_{\mathrm{UV}}) = 0$ *and* $\zeta\, K_{\mathrm{IR}}^{(\ell)} = 0$ *imply* $\zeta\,\mathcal{B}[\mathcal{O}] = \mathcal{B}[\zeta\,\mathcal{O}]$ *and* $\mathrm{gh}(\mathcal{B}[\mathcal{O}]) = \mathrm{gh}(\mathcal{O})$.

(iv) Channel/background compatibility.
If $[P_y(\lambda_{\mathrm{UV}}), \Pi_\alpha] = [K_{\mathrm{IR}}^{(\ell)}, \Pi_\alpha] = 0$ *then* $\mathcal{B}[\Pi_\alpha \mathcal{O} \Pi_\alpha] = \Pi_\alpha \, \mathcal{B}[\mathcal{O}] \, \Pi_\alpha$. *In background-field gauge,* $\mathcal{B}$ *respects Slavnov–Taylor identities on the band.*

(v) Strong-operator continuity.
$\lambda \mapsto \mathcal{B}_{\lambda,\ell}[\mathcal{O}]$ *is strongly continuous in* λ; $\ell \mapsto \mathcal{B}_{\lambda,\ell}[\mathcal{O}]$ *is norm–continuous for* $K_{\mathrm{IR}}^{(\ell)}$ *continuous in* L^1.

(vi) Descent to the physical sector.
With quartet decoupling and $\mathsf{P}_{\mathrm{phys}}$ *the BRST projector,*

$$\mathsf{P}_{\mathrm{phys}} \, \mathcal{B}[\mathcal{O}] \, \mathsf{P}_{\mathrm{phys}} = \mathcal{B}\Big[\mathsf{P}_{\mathrm{phys}} \, \mathcal{O} \, \mathsf{P}_{\mathrm{phys}}\Big],$$

so sector actions/variations commute with passage to $\mathcal{H}_{\mathrm{phys}}$.

Sketch. **(i)–(ii) Kraus form and CP.**
Use the integral Kraus representation

$$\mathcal{B}[\mathcal{O}] = \int d^d y \int d^d x \; V_y(x)^\dagger \, \mathcal{O} \, V_y(x), \qquad V_y(x) := \sqrt{K_{\mathrm{IR}}^{(\ell)}(x-y)} \, P_y(\lambda_{\mathrm{UV}}).$$

This immediately yields linearity, *-preservation, positivity, and complete positivity.

(iii)–(iv) BRST closure and channel/background compatibility.
Since $\zeta \, P_y(\lambda_{\mathrm{UV}}) = 0$ and $\zeta \, K_{\mathrm{IR}}^{(\ell)} = 0$, one has $\zeta \, \mathcal{B}[\mathcal{O}] = \mathcal{B}[\zeta \, \mathcal{O}]$ and $\mathrm{gh}(\mathcal{B}[\mathcal{O}]) = \mathrm{gh}(\mathcal{O})$. Commutators pass under the integral, so if $[P_y(\lambda_{\mathrm{UV}}), \Pi_\alpha] = [K_{\mathrm{IR}}^{(\ell)}, \Pi_\alpha] = 0$ then $\mathcal{B}[\Pi_\alpha \mathcal{O} \Pi_\alpha] = \Pi_\alpha \mathcal{B}[\mathcal{O}] \Pi_\alpha$. In background-field gauge, Slavnov–Taylor identities are preserved on the band.

(v) Strong-operator continuity.
For fixed ℓ, $\lambda \mapsto P_y(\lambda)$ is strongly continuous (spectral calculus), and for fixed λ, $\ell \mapsto K_{\mathrm{IR}}^{(\ell)}$ is L^1–continuous; dominated convergence gives strong (resp. norm) continuity of $\mathcal{B}_{\lambda,\ell}$.

(vi) Descent to the physical sector.
With quartet decoupling and $\mathsf{P}_{\mathrm{phys}}$ the BRST projector,

$$\mathsf{P}_{\mathrm{phys}} \, \mathcal{B}[\mathcal{O}] \, \mathsf{P}_{\mathrm{phys}} \;=\; \mathcal{B}\Big[\mathsf{P}_{\mathrm{phys}} \, \mathcal{O} \, \mathsf{P}_{\mathrm{phys}}\Big],$$

since the longitudinal/ghost sector is BRST-exact and orthogonal to $\mathcal{H}_{\mathrm{phys}}$.

□

Shorthand and commuting approximation. In the joint IR/UV tagging setup of Definition 7.28 and for the band map $\mathcal{B}^{(\lambda_{\mathrm{UV}},\ell)}$ from Proposition 7.30, assume $K_{\mathrm{IR}}^{(\ell)} \geq 0$ is a scalar window of unit mass and width ℓ, and the projector field $P_y(\lambda_{\mathrm{UV}}) = P_y^\dagger = P_y^2$ varies slowly over the window (e.g. $\|P_y - P_x\| \leq C \, |y-x|/L$ for $|y-x| \lesssim \ell$ with $L \gg \ell$). Then, for bounded operator fields $\mathcal{O}$,

$$\begin{aligned} \mathcal{B}^{(\lambda_{\mathrm{UV}},\ell)}[\mathcal{O}](x) &= \int_{\mathbb{R}^d} K_{\mathrm{IR}}^{(\ell)}(x-y) \, P_y \, \mathcal{O}(y) \, P_y \, d^d y \\ &= P_x \, \mathcal{O}^{(\ell)}(x) \, P_x \;+\; R_{\lambda,\ell}[\mathcal{O}](x), \end{aligned} \tag{7.41}$$

with $\mathcal{O}^{(\ell)} := K_{\mathrm{IR}}^{(\ell)} * \mathcal{O}$ and remainder estimate

$$\left\| R_{\lambda,\ell}[\mathcal{O}](x) \right\| \;\le\; 2\, C\, C_K\, \frac{\ell}{L} \sup_{|y-x|\lesssim \ell} \|\mathcal{O}(y)\|, \qquad C_K := \int_{\mathbb{R}^d} |u|\, K_{\mathrm{IR}}^{(1)}(u)\, du.$$

In the commuting case $P_y \equiv P_x$ on the window the remainder vanishes and $\mathcal{B}^{(\lambda_{\mathrm{UV}},\ell)}[\mathcal{O}](x) = P_x\, \mathcal{O}^{(\ell)}(x)\, P_x$. This shorthand preserves adjoints and positivity by Proposition 7.30(i)–(ii), and its continuity in $(\lambda_{\mathrm{UV}}, \ell)$ follows from Proposition 7.30(v).

7.3.2 Plateau Metrics and Acceptance Criteria

Visibility vector. Let $\mathbf{V}^{(\lambda_{\mathrm{UV}},\ell)}$ denote any fixed set of holosymmetric visibilities (dimensionless, gauge-invariant readouts built from the unified correlators and stress/response channels[3]) evaluated on $\mathcal{H}_{\mathrm{phys}}$ (hence ghost/longitudinal-free by Section 7.2.6). Denote by $\|\cdot\|$ any convenient norm on the finite vector of scalars (we use the ℓ^∞ norm for certification).

Relative drift and curvature. Fix a candidate base point $(\lambda_\star, \ell_\star)$. For a neighbourhood

$$\mathcal{N}_\delta := \Big\{(\lambda,\ell) :\; |\lambda-\lambda_\star| \le \delta_\lambda,\; |\ell-\ell_\star| \le \delta_\ell\Big\},$$

define the *relative drift* and *curvature* seminorms

$$\Delta_{\mathrm{rel}}(\mathbf{V};\mathcal{N}_\delta) := \sup_{(\lambda,\ell)\in\mathcal{N}_\delta} \frac{\|\mathbf{V}^{(\lambda,\ell)} - \mathbf{V}^{(\lambda_\star,\ell_\star)}\|}{\max(1, \|\mathbf{V}^{(\lambda_\star,\ell_\star)}\|)}, \qquad \kappa_{\mathrm{rel}}(\mathbf{V};\mathcal{N}_\delta) := \sup_{(\lambda,\ell)\in\mathcal{N}_\delta} \frac{\left\|\partial_{\lambda,\ell}^2 \mathbf{V}^{(\lambda,\ell)}\right\|}{\max(1, \|\mathbf{V}^{(\lambda_\star,\ell_\star)}\|)}. \tag{7.42}$$

Directional and log–scale diagnostics (auxiliary). It is often convenient to work with scale-invariant (logarithmic) derivatives

$$D_\lambda := \lambda_{\mathrm{UV}} \partial_{\lambda_{\mathrm{UV}}}, \qquad D_\ell := \ell\, \partial_\ell, \tag{7.43}$$

and with directional derivatives along unit vectors $u = (u_\lambda, u_\ell)$, $u_\lambda^2 + u_\ell^2 = 1$:

$$\partial_u := u_\lambda\, \partial_{\lambda_{\mathrm{UV}}} + u_\ell\, \partial_\ell, \qquad \partial_u^2 := \partial_u(\partial_u \cdot)\,.$$

Writing $\mathrm{Hess}_{(\lambda,\ell)}(\mathbf{V})$ for the 2×2 matrix of second derivatives, one has the operator-norm characterisation

$$\sup_{\|u\|=1} \|\partial_u^2 \mathbf{V}^{(\lambda,\ell)}\| = \left\|\mathrm{Hess}_{(\lambda,\ell)}(\mathbf{V})\right\|_{\mathrm{op}}, \tag{7.44}$$

so κ_{rel} controls the maximal directional curvature on $\mathcal{N}_\delta$.

[3] E.g. normalised residues, pole locations m_i^2, spectral-weight ratios, attenuations, and cone-compatibility tests tied to $\bar{\Sigma}_{\mu\nu}^{(\ell)}$.

Commutation of limits. The UV/IR limits are required to commute (up to tolerance) on the physical sector:

$$\Big\| \lim_{\lambda\to\infty}\lim_{\ell\to 0^+} \mathbf{V}^{(\lambda,\ell)} - \lim_{\ell\to 0^+}\lim_{\lambda\to\infty} \mathbf{V}^{(\lambda,\ell)} \Big\| \;\le\; \varepsilon_{\mathrm{swap}}, \qquad \text{on } \mathcal{H}_{\mathrm{phys}}, \tag{7.45}$$

and the swapped limits preserve positivity/unitarity checks (Källén–Lehmann and optical theorem; cf. eq. (7.12) and [82, 91, 241]).

Definition 7.31 (Plateau at $(\lambda_\star,\ell_\star)$)**.** Given tolerances $\varepsilon_{\mathrm{drift}},\varepsilon_{\mathrm{curv}},\varepsilon_{\mathrm{swap}}>0$, the pair $(\lambda_\star,\ell_\star)$ is an *accepted plateau point* for $\mathbf{V}$ if there exists a neighbourhood $\mathcal{N}_\delta$ such that

$$\Delta_{\mathrm{rel}}(\mathbf{V};\mathcal{N}_\delta)\le\varepsilon_{\mathrm{drift}}, \quad \kappa_{\mathrm{rel}}(\mathbf{V};\mathcal{N}_\delta)\le\varepsilon_{\mathrm{curv}}, \quad \text{and} \quad (7.45) \text{ holds with } \varepsilon_{\mathrm{swap}}. \tag{7.46}$$

Sufficient conditions via log–derivatives. Let $R_\lambda := |\log(\lambda/\lambda_\star)|$ and $R_\ell := |\log(\ell/\ell_\star)|$. For any rectifiable path γ inside $\mathcal{N}_\delta$ joining $(\lambda_\star,\ell_\star)$ to (λ,ℓ), the mean-value estimate yields

$$\frac{\|\mathbf{V}^{(\lambda,\ell)}-\mathbf{V}^{(\lambda_\star,\ell_\star)}\|}{\max(1,\|\mathbf{V}^{(\lambda_\star,\ell_\star)}\|)} \;\le\; R_\lambda \sup_{\mathcal{N}_\delta}\frac{\|D_\lambda\mathbf{V}\|}{\max(1,\|\mathbf{V}\|)} + R_\ell \sup_{\mathcal{N}_\delta}\frac{\|D_\ell\mathbf{V}\|}{\max(1,\|\mathbf{V}\|)} + \frac{1}{2}(R_\lambda^2+R_\ell^2)\,\kappa_{\mathrm{rel}}(\mathbf{V};\mathcal{N}_\delta), \tag{7.47}$$

so small *log-slopes* and small curvature suffice for (7.46).

Finite-difference estimators (practical certification). Given a rectangular grid on $\mathcal{N}_\delta$ with steps $(\Delta\lambda,\Delta\ell)$, define centred differences

$$\delta_\lambda\mathbf{V} := \mathbf{V}^{(\lambda+\Delta\lambda,\ell)} - \mathbf{V}^{(\lambda-\Delta\lambda,\ell)}, \qquad \delta_\lambda^2\mathbf{V} := \mathbf{V}^{(\lambda+\Delta\lambda,\ell)} - 2\mathbf{V}^{(\lambda,\ell)} + \mathbf{V}^{(\lambda-\Delta\lambda,\ell)},$$

and similarly for ℓ. Then the consistent estimators

$$\widehat{\Delta}_{\mathrm{rel}} := \sup_{\mathcal{N}_\delta}\frac{\|\mathbf{V}^{(\lambda,\ell)}-\mathbf{V}^{(\lambda_\star,\ell_\star)}\|}{\max(1,\|\mathbf{V}^{(\lambda_\star,\ell_\star)}\|)}, \qquad \widehat{\kappa}_{\mathrm{rel}} := \sup_{\mathcal{N}_\delta}\frac{\big\|\delta_\lambda^2\mathbf{V}/(\Delta\lambda)^2 + \delta_\ell^2\mathbf{V}/(\Delta\ell)^2\big\|}{\max(1,\|\mathbf{V}^{(\lambda_\star,\ell_\star)}\|)} \tag{7.48}$$

converge to (7.42) as $(\Delta\lambda,\Delta\ell)\to 0$.

Oscillation control and minimal width. To suppress spurious micro-oscillations inside $\mathcal{N}_\delta$, introduce

$$\Omega_{\mathrm{rel}}(\mathbf{V};\mathcal{N}_\delta) := \sup_{(\lambda,\ell),(\lambda',\ell')\in\mathcal{N}_\delta}\frac{\|\mathbf{V}^{(\lambda,\ell)}-\mathbf{V}^{(\lambda',\ell')}\|}{\max(1,\|\mathbf{V}^{(\lambda_\star,\ell_\star)}\|)}, \qquad r_{\min} := \min\left\{\frac{\delta_\lambda}{\lambda_\star},\frac{\delta_\ell}{\ell_\star}\right\}. \tag{7.49}$$

In applications we require $\Omega_{\mathrm{rel}}\le\varepsilon_{\mathrm{osc}}$ and $r_{\min}\ge r_{\mathrm{width}}$ for user-chosen tolerances $\varepsilon_{\mathrm{osc}}, r_{\mathrm{width}} > 0$.

Aggregation across visibility suites. For a collection $\{\mathbf{V}^{[s]}\}_s$ (e.g. sector-resolved visibilities, Ward-side diagnostics, and transverse spectral data), define the global metrics by

$$\Delta_{\mathrm{rel}}^{\max} := \max_s \Delta_{\mathrm{rel}}(\mathbf{V}^{[s]};\mathcal{N}_\delta), \quad \kappa_{\mathrm{rel}}^{\max} := \max_s \kappa_{\mathrm{rel}}(\mathbf{V}^{[s]};\mathcal{N}_\delta), \quad \Omega_{\mathrm{rel}}^{\max} := \max_s \Omega_{\mathrm{rel}}(\mathbf{V}^{[s]};\mathcal{N}_\delta).$$

Acceptance is then checked with the same tolerances as in Definition 7.31, supplemented by (7.49).

Error budgets (optional but recommended). When numerical/Monte Carlo errors are present, split the drift into deterministic and stochastic parts via triangle inequality,

$$\Delta_{\mathrm{rel}} \le \Delta_{\mathrm{rel}}^{\mathrm{sys}} + \Delta_{\mathrm{rel}}^{\mathrm{stat}},$$

and require $\Delta_{\mathrm{rel}}^{\mathrm{stat}} \le \varepsilon_{\mathrm{stat}}$ with certified confidence, while tightening $\varepsilon_{\mathrm{drift}}$ accordingly so that (7.46) remains meaningful.

Operational acceptance (summary). A point $(\lambda_\star, \ell_\star)$ is *certified* if: (i) (7.46) holds for each chosen $\mathbf{V}$; (ii) the commuted limits satisfy (7.45); (iii) the auxiliary conditions (7.47), (7.48), (7.49) are within tolerances; and (iv) all checks are performed on $\mathcal{H}_{\mathrm{phys}}$ with BRST/Nielsen safeguards in place (cf. Sections 7.2.6 and 7.2.9).

Pole stability and regulator robustness. For any physical pole m_i^2 (extracted from the Π_{T} two-point function in the sense of eqs. (7.20) and (7.22)), define

$$\Delta_i^{\mathrm{rel}}(\mathcal{N}_\delta) := \sup_{(\lambda,\ell)\in\mathcal{N}_\delta} \frac{|m_i^2(\lambda,\ell) - m_i^2(\lambda_\star,\ell_\star)|}{\max(1, |m_i^2(\lambda_\star,\ell_\star)|)}. \tag{7.50}$$

A certified plateau requires $\Delta_i^{\mathrm{rel}}(\mathcal{N}_\delta) \le \varepsilon_i$ for all i, and independence (within the same tolerance) under admissible changes of $\mathcal{D}_x$ and of the kernel family $K_{\mathrm{IR}}^{(\ell)}$ ("regulator robustness").

Definition 7.32 (Admissible regulator variations)**.** A UV projector replacement $P_x(\lambda_{\mathrm{UV}}) \mapsto P'_x(\lambda_{\mathrm{UV}})$ and an IR window replacement $K_{\mathrm{IR}}^{(\ell)} \mapsto K_{\mathrm{IR}}^{(\ell)\prime}$ are *admissible* on $\mathcal{N}_\delta$ if:

(i) **BRST/channel closure:** $\zeta P'_x = 0$, $\zeta K_{\mathrm{IR}}^{(\ell)\prime} = 0$, $\mathrm{gh}(P'_x) = \mathrm{gh}(K_{\mathrm{IR}}^{(\ell)\prime}) = 0$, and $[P'_x, \Pi_\alpha] = [K_{\mathrm{IR}}^{(\ell)\prime}, \Pi_\alpha] = 0$.

(ii) **Strong-resolvent continuity (UV):** there exists a norm-bounded, strongly continuous path $t \mapsto P_x^{(t)}(\lambda)$ with $P_x^{(0)} = P_x$, $P_x^{(1)} = P'_x$ and

$$\eta_{\mathrm{UV}} := \sup_{(\lambda,\ell)\in\mathcal{N}_\delta} \sup_x \left\| P_x^{(1)}(\lambda) - P_x^{(0)}(\lambda) \right\| < \infty.$$

(iii) L^1 **continuity (IR):** $K_{\mathrm{IR}}^{(\ell)\prime} \in L^1$, non-negative, unit-normalised, and

$$\eta_{\mathrm{IR}} := \sup_{(\lambda,\ell)\in\mathcal{N}_\delta} \left\| K_{\mathrm{IR}}^{(\ell)\prime} - K_{\mathrm{IR}}^{(\ell)} \right\|_{L^1} < \infty.$$

(iv) **Covariant differentiation:** $[\nabla_\mu, P'_x(\lambda_{\mathrm{UV}})] = 0$, and $K_{\mathrm{IR}}^{(\ell)\prime}$ is a scalar kernel, so tagging commutes with ∇ as in the canonical construction.

We call $(\eta_{\mathrm{UV}}, \eta_{\mathrm{IR}})$ the *regulator mismatch metrics*.

Lemma 7.33 (Lipschitz control of the pole map under admissible variations)**.** *Assume Definition 7.32 on $\mathcal{N}_\delta$, hyperbolicity eq. (7.15), and that each physical pole $-m_i^2$ is simple at $(\lambda_\star, \ell_\star)$ with a non-vanishing on-shell slope* (Rayleigh–sandwiched derivative)

$$\mathfrak{s}_i \;:=\; \left| \bar{v}_i \, \partial_{p^2} \Gamma^{(\mathrm{TT})}(-m_i^2) \, v_i \right| \;>\; 0,$$

*i.e. the derivative of the inverse two-point kernel evaluated between the left/right pole eigenvectors at $p^2 = -m_i^2$ (*Källén–Lehmann non-degeneracy*). Then there exist finite constants C_i (depending on local operator norms of the band Hessian/vertices on $\mathcal{N}_\delta$) such that*

$$\sup_{(\lambda,\ell)\in\mathcal{N}_\delta} \frac{|m_i^{2\prime}(\lambda,\ell) - m_i^2(\lambda,\ell)|}{\max(1, |m_i^2(\lambda_\star,\ell_\star)|)} \;\le\; C_i\Big(\eta_{\mathrm{UV}} + \eta_{\mathrm{IR}}\Big). \tag{7.51}$$

Proof of Lemma 7.33. Step 1: Band functor is Lipschitz in the regulators. By Definition 7.29, the band map $\mathcal{B}_{\lambda,\ell}$ is built from the UV limiter $P_x(\lambda_{\mathrm{UV}})$ and the IR window $K_{\mathrm{IR}}^{(\ell)}$ (both of pseudodifferential order ≤ 0), composed with channel projectors Π_α; admissibility in Definition 7.32 gives

$$\big\|\mathcal{B}'_{\lambda,\ell}[\mathcal{O}] - \mathcal{B}_{\lambda,\ell}[\mathcal{O}]\big\| \;\le\; c_{\mathcal{O}}\,\Big(\eta_{\mathrm{UV}} + \eta_{\mathrm{IR}}\Big),$$

uniformly for $(\lambda,\ell)\in\mathcal{N}_\delta$, for any operator $\mathcal{O}$ entering the quadratic/vertex sector. (Uniformity is ensured by the plateau/robustness certification from Chapter 7, cf. eq. (6.380) and Proposition 6.298.)

Step 2: Control of the 1PI two-point kernel. Let $\Gamma^{(\mathrm{TT})}$ denote the (transverse–traceless) 1PI inverse two-point kernel on the band (the projected Hessian). The construction of $\Gamma^{(\mathrm{TT})}$ from banded vertices is locally Lipschitz with respect to the input operators (Dyson/Neumann expansions converge on $\mathcal{N}_\delta$ thanks to admissibility and the plateau bounds). Hence there exists $c' > 0$ such that

$$\big\|\Delta\Gamma^{(\mathrm{TT})}\big\| := \big\|\Gamma^{(\mathrm{TT})\prime} - \Gamma^{(\mathrm{TT})}\big\| \;\le\; c'\,\Big(\eta_{\mathrm{UV}} + \eta_{\mathrm{IR}}\Big), \tag{7.52}$$

uniformly on $(\lambda,\ell)\in\mathcal{N}_\delta$ and for p^2 in a small neighbourhood of $-m_i^2(\lambda_\star,\ell_\star)$. Hyperbolicity eq. (7.15) guarantees that the TT-projection is well-posed and the dispersion relation is controlled on the band.

Step 3: Simple-pole perturbation via the Rayleigh–sandwiched derivative. Define the scalar function

$$f_i(p^2;\lambda,\ell) := \bar v_i(\lambda,\ell)\,\Gamma^{(\mathrm{TT})}(p^2;\lambda,\ell)\,v_i(\lambda,\ell),$$

where $v_i, \bar v_i$ are the right/left eigenvectors at the pole (normalised so that $f_i(-m_i^2;\lambda,\ell) = 0$). At $(\lambda_\star,\ell_\star)$ simplicity and non-degeneracy give

$$\Big|\partial_{p^2} f_i(-m_i^2;\lambda_\star,\ell_\star)\Big| = \Big|\bar v_i\,\partial_{p^2}\Gamma^{(\mathrm{TT})}(-m_i^2)\,v_i\Big| = \mathfrak{s}_i > 0.$$

By continuity on $\mathcal{N}_\delta$, there exists $\underline{\mathfrak{s}}_i > 0$ with

$$\inf_{(\lambda,\ell)\in\mathcal{N}_\delta} \Big|\partial_{p^2} f_i(-m_i^2;\lambda,\ell)\Big| \;\ge\; \underline{\mathfrak{s}}_i. \tag{7.53}$$

Let the perturbed zero be $-m_i^{2\prime}$, i.e. $f_i(-m_i^{2\prime};\lambda,\ell) + \Delta f_i(-m_i^{2\prime};\lambda,\ell) = 0$, where $\Delta f_i := \bar v_i\,\Delta\Gamma^{(\mathrm{TT})}\,v_i$. A first-order Taylor expansion around $-m_i^2$ and the mean-value theorem yield

$$\Big|m_i^{2\prime} - m_i^2\Big| \;\le\; \frac{\Big|\Delta f_i(-m_i^2;\lambda,\ell)\Big|}{\Big|\partial_{p^2} f_i(\xi;\lambda,\ell)\Big|} \;\le\; \frac{\|\Delta\Gamma^{(\mathrm{TT})}\|}{\underline{\mathfrak{s}}_i},$$

for some ξ between $-m_i^2$ and $-m_i^{2\prime}$, using $|\bar v_i(\cdot)\,X\,v_i(\cdot)| \le \|X\|$ for the operator norm.

Step 4: Collecting the bounds. Combining (7.52) and (7.53) gives

$$\left|m_i^{2\prime} - m_i^2\right| \;\le\; \frac{c'}{\mathfrak{s}_i}\left(\eta_{\mathrm{UV}} + \eta_{\mathrm{IR}}\right).$$

Dividing by $\max(1, |m_i^2(\lambda_\star, \ell_\star)|)$ produces the dimensionless form in (7.51). Setting $C_i := c'/\mathfrak{s}_i$ completes the estimate. The constants depend only on local operator norms of the band Hessian/vertices on $\mathcal{N}_\delta$ (via c') and on the non-degeneracy margin $\mathfrak{s}_i$ fixed by the Rayleigh–sandwiched slope.

Conclusion. Under the admissibility and hyperbolicity hypotheses Definition 7.32 and eq. (7.15), simple poles are stable with Lipschitz modulus $C_i(\eta_{\mathrm{UV}} + \eta_{\mathrm{IR}})$, controlled by the Rayleigh–sandwiched on-shell slope $\mathfrak{s}_i$ and the band-uniform bounds furnished by the plateau machinery of Chapter 7 (eq. (6.380) and Proposition 6.298). □

Proposition 7.34 (Plateau ⇒ robust, gauge-independent pole extraction)**.** *Assume the anomaly filters and Slavnov identity eqs.* (7.16) *and* (7.18)*, quartet decoupling eq.* (7.11)*, hyperbolicity eq.* (7.15)*, and the Nielsen identities eq.* (7.22)*. If* $(\lambda_\star, \ell_\star)$ *is an accepted plateau point in the sense of Definition 7.31, then every physical pole* m_i^2 *is (i)* gauge-parameter independent *on shell, and (ii)* UV/IR-regulator robust *with drift bounded by* ε_i *in* $\mathcal{N}_\delta$*.*

Proof sketch. (i) Gauge-parameter independence: eq. (7.22) on $\mathcal{H}_{\mathrm{phys}}$ (with ghost/longitudinal kill, eq. (7.11)) gives $\partial_\alpha m_i^2 = 0$.

(ii) Regulator robustness: by eq. (7.46), both (a) the drift Δ_{rel} and (b) the curvature κ_{rel} of the *visibility vector* are uniformly small on $\mathcal{N}_\delta$; by the chain rule and eq. (7.47) this controls the pole locator functional $m_i^2(\lambda, \ell)$. Admissible replacements of (P, K) produce band-functor shifts controlled by $(\eta_{\mathrm{UV}}, \eta_{\mathrm{IR}})$; applying Lemma 7.33 yields the bound (7.51). Choosing tolerances so that $C_i(\eta_{\mathrm{UV}} + \eta_{\mathrm{IR}}) \le \varepsilon_i$ proves (ii). The commutation of limits (7.45) rules out order-of-limits artefacts as (λ, ℓ) vary within $\mathcal{N}_\delta$. Hyperbolicity ensures a well-posed Cauchy problem across $\mathcal{N}_\delta$, so spectral perturbation applies smoothly on the band. □

Degeneracies (multi-pole clusters). If a cluster of r poles is degenerate, use the spectral projector $\mathbb{P}_i$ and restricted kernel $\widehat{\Gamma}_i$ as in the bandwise checklist: admissible variations preserve the *set* of on-shell eigenvalues (Nielsen) and the cluster drift obeys the analogue of (7.51) with C_i depending on the $\widehat{\Gamma}_i$ spectral gap (see the discussion around the spectral projector in the checklist section).

7.3.3 Operational Certification Protocol

Inputs. Tolerances $\left(\varepsilon_{\mathrm{drift}}, \varepsilon_{\mathrm{curv}}, \varepsilon_{\mathrm{swap}}, \{\varepsilon_i\}_i, \varepsilon_{\mathrm{osc}}, r_{\mathrm{width}}\right)$, a base point $(\lambda_\star, \ell_\star)$, a rectangular neighbourhood $\mathcal{N}_\delta = [\lambda_\star - \delta_\lambda, \lambda_\star + \delta_\lambda] \times [\ell_\star - \delta_\ell, \ell_\star + \delta_\ell]$, and a grid $\mathcal{G}$ on $\mathcal{N}_\delta$ with spacings $(\Delta\lambda, \Delta\ell)$ containing $(\lambda_\star, \ell_\star)$.
Outputs. (*i*) Pass/Fail at $(\lambda_\star, \ell_\star)$; (*ii*) a certificate comprising the maximal observed metrics on $\mathcal{N}_\delta$; (*iii*) witnesses for any failure (counterexamples).

P1. Assemble band objects (well-posedness gate). Build $\mathbb{H}^{(\lambda_{\mathrm{UV}}, \ell)}$ and its $\Pi_{\mathrm{T}}/\Pi_{\mathrm{L}}$ split (Section 7.2.5); verify quartet decoupling eq. (7.11). Compute $\mathbb{M}^2_{\mathrm{phys}}(\lambda_{\mathrm{UV}}, \ell)$ and enforce

the tachyon/hyperbolicity gate eqs. (7.14) and (7.15) on all $(\lambda, \ell) \in \mathcal{G}$. *Reject immediately* if any gate fails, recording the smallest (λ, ℓ) failing point.

P2. Choose a visibility set (physical readouts). Fix a holosymmetric visibility vector $\mathbf{V}^{(\lambda,\ell)}$ (e.g. $\{m_i^2$, residue norms, spectral weights, cone-compatibility tests$\}$) evaluated on $\mathcal{H}_{\text{phys}}$; store the baseline $\mathbf{V}^{(\lambda_\star,\ell_\star)}$.

P3. Drift/curvature scan on the grid. On $\mathcal{G} \subset \mathcal{N}_\delta$, compute the finite-difference estimators $\widehat{\Delta}_{\text{rel}}$ and $\widehat{\kappa}_{\text{rel}}$ from (7.48) and the pole drift $\Delta_i^{\text{rel}}(\mathcal{N}_\delta)$ from (7.50). If $\widehat{\Delta}_{\text{rel}} \le \varepsilon_{\text{drift}}$ and $\widehat{\kappa}_{\text{rel}} \le \varepsilon_{\text{curv}}$ and $\Delta_i^{\text{rel}} \le \varepsilon_i$ for all i, proceed; otherwise adapt $(\Delta\lambda, \Delta\ell)$ (refine) or $(\delta_\lambda, \delta_\ell)$ (shrink) until either the bounds hold or the minimal width $r_{\min} \ge r_{\text{width}}$ (eq. (7.49)) is violated (in which case *reject* with witness).

P4. Swap-limits test (corner check). Check (7.45) numerically/analytically by approaching $(\infty, 0^+)$ along at least two independent paths (e.g. $\lambda \to \infty$ then $\ell \to 0^+$, and the reverse; optionally a diagonal $\ell \propto \lambda^{-a}$). *Accept* this step if the discrepancy is $\le \varepsilon_{\text{swap}}$; otherwise *reject* with the offending path as a witness.

P5. Regulator robustness (admissible replacements). Choose an admissible pair (P', K') in the sense of Definition 7.32 and repeat steps **P1**–**P4** on the same $\mathcal{G}$. Compute the regulator mismatch metrics $(\eta_{\text{UV}}, \eta_{\text{IR}})$ and the pole difference bound via (7.51). *Accept* this step if the observed drifts remain within the same tolerances (in particular $\Delta_i^{\text{rel}\,\prime} \le \varepsilon_i$). If not, *reject* with the maximally offending (λ, ℓ) and pole index i as witnesses.

P6. Gauge-parameter certification (Nielsen gate). Promote α to a BRST doublet and verify (7.20). For each physical pole, confirm $\partial_\alpha m_i^2 = 0$ on shell (eq. (7.22)) at the base and at a representative set of points in $\mathcal{G}$; for degenerate clusters use the spectral projector test (cluster poles invariant; residues may vary). *Reject* on any observed α-dependence of pole positions.

P7. Oscillation control and plateau width (stability gate). Evaluate $\Omega_{\text{rel}}(\mathbf{V}; \mathcal{N}_\delta)$ and $r_{\min}$ (eq. (7.49)). *Accept* if $\Omega_{\text{rel}} \le \varepsilon_{\text{osc}}$ and $r_{\min} \ge r_{\text{width}}$, else adapt $(\delta_\lambda, \delta_\ell)$ or refine $\mathcal{G}$; if infeasible, *reject* with witness pairs $(\lambda, \ell), (\lambda', \ell')$.

P8. Certification record (audit trail). Upon success of **P1**–**P7**, emit a certificate containing:

- the base point $(\lambda_\star, \ell_\star)$, neighbourhood $\mathcal{N}_\delta$, grid spacings $(\Delta\lambda, \Delta\ell)$;
- the maximal observed metrics $\widehat{\Delta}_{\text{rel}}$, $\widehat{\kappa}_{\text{rel}}$, Ω_{rel}, and $\{\Delta_i^{\text{rel}}\}_i$;
- the swap-limits discrepancy and the admissible regulator pair (P', K') with $(\eta_{\text{UV}}, \eta_{\text{IR}})$;
- a list of simple/degenerate poles, their certified values, and (if any) cluster projectors used;
- the statement that all checks were performed on $\mathcal{H}_{\text{phys}}$ with BRST/Nielsen safeguards (references to eqs. (7.11), (7.14), (7.15) and (7.22)).

If any step fails, store the minimal-counterexample witness (point(s), path, or regulator pair) for diagnosis.

Remark (provisional identifications). Any labels such as "graviton-like," "Higgs-like," "apeiron," or "chrono–kymon" attached to the certified poles remain *provisional* until cross-quadrant continuation confirms their stability (cf. the policy stated prior to Section 7.1).

7.3.4 Visibilities and their Scale Dependence

Definition 7.35 (Holosymmetric visibilities (band/IR tagged))**.** Let $\Lambda_\alpha(\lambda_{\mathrm{UV}}, \ell)$ denote any of the holosymmetric *visibilities* (contrast-type diagnostics) evaluated on $\mathcal{H}_{\mathrm{phys}}$ using the controlled, two-branch evolution of Chapter 7; here $\alpha \in \{\mathrm{law}, \mathrm{state}, \mathrm{mix}, \ldots\}$ indexes the diagnostic channel. Abstractly,

$$\begin{aligned} \Lambda_\alpha(\lambda_{\mathrm{UV}}, \ell) := \mathcal{F}_\alpha\Big[\{\text{band-limited, IR-smoothed correlators}\};\, \bar{\Sigma}^{(\ell)}_{\mu\nu}\Big], \\ -1 \le \Lambda_\alpha \le 1, \end{aligned} \tag{7.54}$$

where $\mathcal{F}_\alpha$ is gauge-invariant and BRST-closed, and the dependence on $(\lambda_{\mathrm{UV}}, \ell)$ is solely through the UV projector and IR window of eqs. (7.29) and (7.33). The tagging knobs are BRST-closed (eq. (7.30)) and commute with channel projectors, hence—on shell—visibilities are *gauge-parameter independent* by the Nielsen identities (Section 7.2.9 and eqs. (7.22) and (7.23)).

Basic properties.

(i) *Normalization and boundedness:* $-1 \le \Lambda_\alpha \le 1$ (normalised contrast).

(ii) *BRST/ghost structure:* $\mathrm{gh}(\Lambda_\alpha) = 0$ and $\zeta\Lambda_\alpha = 0$ on $\mathcal{H}_{\mathrm{phys}}$.

(iii) *Band/IR covariance:* Λ_α is functorial under the band map of Definition 7.29 ("UV then IR"), and depends on $(\lambda_{\mathrm{UV}}, \ell)$ only via $P_x(\lambda_{\mathrm{UV}})$ and $K^{(\ell)}_{\mathrm{IR}}$.

(iv) *Gauge-parameter independence (on shell):* if eq. (7.16) holds and quartet decoupling applies, then $\partial_\alpha \Lambda_\alpha = 0$ on $\mathcal{H}_{\mathrm{phys}}$.

Definition 7.36 (Scale gradients (UV/IR flows))**.** Let the logarithmic scale derivatives $D_\lambda := \lambda_{\mathrm{UV}}\partial_{\lambda_{\mathrm{UV}}}$ and $D_\ell := \ell\,\partial_\ell$ be as in eq. (7.43). The *scale gradients* of a visibility are

$$\mathcal{G}^{(\lambda)}_\alpha := \big|D_\lambda \Lambda_\alpha\big|, \qquad \mathcal{G}^{(\ell)}_\alpha := \big|D_\ell \Lambda_\alpha\big|. \tag{7.55}$$

These derivatives exist a.e. on compact rectangles by our regularity assumptions and are obtained by differentiating the band projector and IR window *inside* $\mathcal{F}_\alpha$ (band-safe calculus consistent with Definitions 7.28 and 7.29).

Chain rule and bounds (use in practice). Let $L^{(P)}_\alpha$ and $L^{(K)}_\alpha$ be local Lipschitz moduli of $\mathcal{F}_\alpha$ with respect to its UV-compressed and IR-averaged inputs (induced by operator norms on correlator blocks). Using the plateau calculus of Definition 7.28, one obtains on compact $\mathcal{R} \subset \mathbb{R}^2_{>0}$:

$$\mathcal{G}^{(\lambda)}_\alpha \;\le\; L^{(P)}_\alpha \,\big\|(\lambda_{\mathrm{UV}}\partial_{\lambda_{\mathrm{UV}}}P)\cdot P + P\cdot(\lambda_{\mathrm{UV}}\partial_{\lambda_{\mathrm{UV}}}P)\big\|_{\mathrm{eff}}, \qquad \mathcal{G}^{(\ell)}_\alpha \;\le\; L^{(K)}_\alpha \,\big\|\ell\,\partial_\ell K^{(\ell)}_{\mathrm{IR}}\big\|_{L^1\to\mathrm{eff}},$$

where $\|\cdot\|_{\mathrm{eff}}$ is the operator norm induced on the effective correlator space used by $\mathcal{F}_\alpha$. Consequently, small log-slopes of the knobs imply small scale gradients for all visibilities, and plateau tolerances can be enforced via eqs. (7.42) and (7.46).

Plateau sets for visibilities. Given tolerances $(\varepsilon_\lambda, \varepsilon_\ell, \varepsilon_{\mathrm{osc}})$ and a scan rectangle $\mathcal{R} \subset \mathbb{R}^2_{>0}$, define the *plateau set* for channel α by

$$\mathfrak{P}_\alpha(\mathcal{R}; \varepsilon_\lambda, \varepsilon_\ell, \varepsilon_{\mathrm{osc}}) := \Big\{(\lambda_{\mathrm{UV}}, \ell) \in \mathcal{R} : \ \big|D_\lambda \Lambda_\alpha\big| \le \varepsilon_\lambda, \ \big|D_\ell \Lambda_\alpha\big| \le \varepsilon_\ell, \ \mathrm{osc}_{\mathcal{N}}(\Lambda_\alpha) \le \varepsilon_{\mathrm{osc}}\Big\}, \tag{7.56}$$

where D_λ, D_ℓ are the logarithmic scale derivatives of eq. (7.43) and $\mathrm{osc}_{\mathcal{N}}$ denotes the local sup–inf oscillation over a neighbourhood $\mathcal{N}$ (defined below). This is the channelwise specialization of the general plateau criterion in Section 7.3.2.

Definition 7.37 (Local oscillation)**.** For a bounded function f on $\mathcal{R}$ and a neighbourhood basis $\{\mathcal{N}_{(\lambda,\ell)}\}$ (e.g. axis-aligned rectangles of side-lengths $\delta_\lambda, \delta_\ell$), define

$$\mathrm{osc}_{\mathcal{N}}(f)(\lambda, \ell) := \sup_{(\lambda', \ell'), \, (\lambda'', \ell'') \in \mathcal{N}_{(\lambda,\ell)}} \Big|f(\lambda', \ell') - f(\lambda'', \ell'')\Big|.$$

When the dependence on (λ, ℓ) is clear, we abbreviate $\mathrm{osc}_{\mathcal{N}}(f)$.

Proposition 7.38 (Basic properties of $\mathfrak{P}_\alpha$)**.** *Assume Λ_α is continuous on $\mathcal{R}$ and admits the scale gradients of eq. (7.55). Then:*

(i) Closedness.
$\mathfrak{P}_\alpha(\mathcal{R}; \varepsilon_\lambda, \varepsilon_\ell, \varepsilon_{\mathrm{osc}})$ *is closed in* $\mathcal{R}$.

(ii) Monotonicity in tolerances.
If $(\varepsilon_\lambda, \varepsilon_\ell, \varepsilon_{\mathrm{osc}}) \preceq (\tilde\varepsilon_\lambda, \tilde\varepsilon_\ell, \tilde\varepsilon_{\mathrm{osc}})$ *componentwise, then*

$$\mathfrak{P}_\alpha(\mathcal{R}; \varepsilon_\lambda, \varepsilon_\ell, \varepsilon_{\mathrm{osc}}) \subseteq \mathfrak{P}_\alpha(\mathcal{R}; \tilde\varepsilon_\lambda, \tilde\varepsilon_\ell, \tilde\varepsilon_{\mathrm{osc}}).$$

(iii) Compatibility with global criteria.
If a neighbourhood $\mathcal{N}_\delta$ around $(\lambda_\star, \ell_\star)$ satisfies eq. (7.46) for the visibility vector that contains Λ_α, then $(\lambda_\star, \ell_\star) \in \mathfrak{P}_\alpha(\mathcal{R}; \varepsilon_\lambda, \varepsilon_\ell, \varepsilon_{\mathrm{osc}})$ for suitably chosen $(\varepsilon_\lambda, \varepsilon_\ell, \varepsilon_{\mathrm{osc}})$ determined by the bounds in eqs. (7.47) to (7.49).

Gauge-parameter and regulator stability. On the physical sector,

$$\partial_{\alpha_{\mathrm{gf}}} \Lambda_\alpha(\lambda_{\mathrm{UV}}, \ell) = 0 \quad \text{on shell},$$

by Section 7.2.9 and eq. (7.22). For admissible regulator changes $(P, K) \mapsto (P', K')$ in the sense of Definition 7.32 with mismatch metrics $(\eta_{\mathrm{UV}}, \eta_{\mathrm{IR}})$, the Lipschitz continuity of $\mathcal{F}_\alpha$ (cf. the bounds following Definition 7.69) yields

$$\sup_{(\lambda,\ell)\in\mathcal{R}} \Big|\Lambda'_\alpha(\lambda, \ell) - \Lambda_\alpha(\lambda, \ell)\Big| \ \le \ L^{(P)}_\alpha \, \eta_{\mathrm{UV}} + L^{(K)}_\alpha \, \eta_{\mathrm{IR}} \ =: \ \varepsilon_{\mathrm{reg}}. \tag{7.57}$$

Certification in Section 7.3.3 therefore demands that $\varepsilon_{\mathrm{reg}}$ be dominated by the chosen plateau tolerances (typically $\varepsilon_{\mathrm{reg}} \ll \min\{\varepsilon_\lambda, \varepsilon_\ell, \varepsilon_{\mathrm{osc}}\}$).

Definition 7.39 (Integrated visibilities (box averages))**.** For noisy scans, define the local box average around $(\lambda_{\mathrm{UV}}, \ell)$ with widths $(\Delta\lambda, \Delta\ell)$ by

$$\overline{\Lambda}_\alpha(\lambda_{\mathrm{UV}}, \ell; \Delta\lambda, \Delta\ell) := \frac{1}{\Delta\lambda \, \Delta\ell} \int_{\lambda_{\mathrm{UV}} - \Delta\lambda/2}^{\lambda_{\mathrm{UV}} + \Delta\lambda/2} d\lambda \int_{\ell - \Delta\ell/2}^{\ell + \Delta\ell/2} d\tilde\ell \ \Lambda_\alpha(\lambda, \tilde\ell). \tag{7.58}$$

Lemma 7.40 (Bias bound for box averages)**.** *Let* $J := \left[\lambda - \frac{\Delta\lambda}{2}, \lambda + \frac{\Delta\lambda}{2}\right] \times \left[\ell - \frac{\Delta\ell}{2}, \ell + \frac{\Delta\ell}{2}\right] \subset \mathcal{R}$. *If* $\partial_\lambda \Lambda_\alpha$ *and* $\partial_\ell \Lambda_\alpha$ *are bounded on* J*, then*

$$\left|\overline{\Lambda}_\alpha(\lambda, \ell; \Delta\lambda, \Delta\ell) - \Lambda_\alpha(\lambda, \ell)\right| \;\le\; \tfrac{1}{2}\,\Delta\lambda \sup_J |\partial_\lambda \Lambda_\alpha| \;+\; \tfrac{1}{2}\,\Delta\ell \sup_J |\partial_\ell \Lambda_\alpha|.$$

Equivalently, in terms of log–derivatives and $\lambda_{\min}(J) := \inf_J \lambda$, $\ell_{\min}(J) := \inf_J \ell$,

$$\left|\overline{\Lambda}_\alpha - \Lambda_\alpha\right| \;\le\; \tfrac{1}{2}\frac{\Delta\lambda}{\lambda_{\min}(J)} \sup_J |D_\lambda \Lambda_\alpha| \;+\; \tfrac{1}{2}\frac{\Delta\ell}{\ell_{\min}(J)} \sup_J |D_\ell \Lambda_\alpha|.$$

Proof of Lemma 7.40. Assume $\Lambda_\alpha \in W^{1,\infty}(J)$, i.e. its partial derivatives are essentially bounded on J (the C^1 case is a fortiori covered). For any $(\lambda', \ell') \in J$, write the rectangle path $(\lambda, \ell) \to (\lambda', \ell) \to (\lambda', \ell')$ and use the one–dimensional fundamental theorem along each segment:

$$\Lambda_\alpha(\lambda', \ell') - \Lambda_\alpha(\lambda, \ell) = \int_\lambda^{\lambda'} \partial_\lambda \Lambda_\alpha(s, \ell)\, ds + \int_\ell^{\ell'} \partial_\ell \Lambda_\alpha(\lambda', t)\, dt.$$

Taking absolute values and bounding by the essential suprema on J gives

$$\left|\Lambda_\alpha(\lambda', \ell') - \Lambda_\alpha(\lambda, \ell)\right| \le \left(\sup_J |\partial_\lambda \Lambda_\alpha|\right) |\lambda' - \lambda| + \left(\sup_J |\partial_\ell \Lambda_\alpha|\right) |\ell' - \ell|.$$

Average this inequality over the box $J = \left[\lambda - \frac{\Delta\lambda}{2}, \lambda + \frac{\Delta\lambda}{2}\right] \times \left[\ell - \frac{\Delta\ell}{2}, \ell + \frac{\Delta\ell}{2}\right]$ and divide by its area $\Delta\lambda\,\Delta\ell$:

$$\begin{aligned}\left|\overline{\Lambda}_\alpha(\lambda, \ell; \Delta\lambda, \Delta\ell) - \Lambda_\alpha(\lambda, \ell)\right| \le{}& \left(\sup_J |\partial_\lambda \Lambda_\alpha|\right) \frac{1}{\Delta\lambda} \int_{\lambda - \Delta\lambda/2}^{\lambda + \Delta\lambda/2} |\lambda' - \lambda|\, d\lambda' \\ &+ \left(\sup_J |\partial_\ell \Lambda_\alpha|\right) \frac{1}{\Delta\ell} \int_{\ell - \Delta\ell/2}^{\ell + \Delta\ell/2} |\ell' - \ell|\, d\ell'.\end{aligned}$$

Each centred average equals half the side length:

$$\frac{1}{\Delta\lambda} \int_{\lambda - \Delta\lambda/2}^{\lambda + \Delta\lambda/2} |\lambda' - \lambda|\, d\lambda' = \frac{\Delta\lambda}{2}, \qquad \frac{1}{\Delta\ell} \int_{\ell - \Delta\ell/2}^{\ell + \Delta\ell/2} |\ell' - \ell|\, d\ell' = \frac{\Delta\ell}{2},$$

which yields the first bound in the statement:

$$\left|\overline{\Lambda}_\alpha - \Lambda_\alpha\right| \;\le\; \tfrac{1}{2}\,\Delta\lambda \sup_J |\partial_\lambda \Lambda_\alpha| \;+\; \tfrac{1}{2}\,\Delta\ell \sup_J |\partial_\ell \Lambda_\alpha|.$$

For the log–derivative form, note $\partial_\lambda \Lambda_\alpha = (1/\lambda)\, D_\lambda \Lambda_\alpha$ and $\lambda \ge \lambda_{\min}(J)$ on J; similarly for ℓ. Hence

$$\sup_J |\partial_\lambda \Lambda_\alpha| \le \frac{1}{\lambda_{\min}(J)} \sup_J |D_\lambda \Lambda_\alpha|, \qquad \sup_J |\partial_\ell \Lambda_\alpha| \le \frac{1}{\ell_{\min}(J)} \sup_J |D_\ell \Lambda_\alpha|,$$

and substitution gives the claimed logarithmic version. □

Corollary 7.41 (Rescaled tolerances for averaged visibilities)**.** *If* $\left|D_\lambda \Lambda_\alpha\right| \le \varepsilon_\lambda$ *and* $\left|D_\ell \Lambda_\alpha\right| \le \varepsilon_\ell$ *on* J*, then choosing*

$$\tilde{\varepsilon}_\lambda := \varepsilon_\lambda \cdot \frac{\Delta\lambda}{2\,\lambda_{\min}(J)}, \qquad \tilde{\varepsilon}_\ell := \varepsilon_\ell \cdot \frac{\Delta\ell}{2\,\ell_{\min}(J)}$$

ensures that $\overline{\Lambda}_\alpha$ *satisfies the same plateau inequalities as* Λ_α *with* $(\varepsilon_\lambda, \varepsilon_\ell)$ *replaced by* $(\tilde{\varepsilon}_\lambda, \tilde{\varepsilon}_\ell)$ *(and with the same* ε_{osc} *once* $\Delta\lambda, \Delta\ell$ *are fixed).*

Proof of Corollary 7.41. On J, assume $\sup_J |D_\lambda \Lambda_\alpha| \le \varepsilon_\lambda$ and $\sup_J |D_\ell \Lambda_\alpha| \le \varepsilon_\ell$. Apply the logarithmic bound in Lemma 7.40:

$$\left|\overline{\Lambda}_\alpha - \Lambda_\alpha\right| \;\le\; \tfrac{1}{2}\frac{\Delta\lambda}{\lambda_{\min}(J)}\,\varepsilon_\lambda \;+\; \tfrac{1}{2}\frac{\Delta\ell}{\ell_{\min}(J)}\,\varepsilon_\ell \;=\; \tilde{\varepsilon}_\lambda + \tilde{\varepsilon}_\ell.$$

Therefore, whenever Λ_α satisfies the plateau inequalities with tolerances $(\varepsilon_\lambda, \varepsilon_\ell)$, the averaged observable $\overline{\Lambda}_\alpha$ satisfies the same inequalities with $(\tilde{\varepsilon}_\lambda, \tilde{\varepsilon}_\ell)$ in place of $(\varepsilon_\lambda, \varepsilon_\ell)$. Since the averaging window is fixed, the local oscillation metric $\varepsilon_{\mathrm{osc}}$ is unaffected. □

Remarks. (i) All statements are *on shell* in $\mathcal{H}_{\mathrm{phys}}$ (ghost/longitudinal sectors removed). (ii) $\mathfrak{P}_\alpha$ may be intersected across channels to define a joint plateau set $\bigcap_\alpha \mathfrak{P}_\alpha$; this is compatible with the global acceptance of Section 7.3.2 and Definition 7.31. (iii) When degeneracies in spectral data affect Λ_α, use the spectral-projector construction from the checklist to ensure channel robustness under small regulator variations, then apply (7.57).

Quadrant signatures (for context). On plateau bands, the overlays below capture the *expected trends* of the holosymmetric visibilities Λ_α (cf. (7.54)) without entering the acceptance logic.

Definition 7.42 (Quadrant signature overlays on a certified plateau)**.** Let $\mathfrak{P} \subset \mathbb{R}^2_{>0}$ be a certified plateau (in the sense of Definition 7.31) for a given model. Assume a dynamical scaling exponent z and an irrelevant deformation with scaling dimension $\Delta > d + z$. Then the following *overlays* are expected to hold on $\mathfrak{P}$:

- **QI.** (**QI: holosymmetric phase**) $\left|\Lambda_{\mathrm{law}}\right| \approx 1$ and $\left|\Lambda_{\mathrm{state}}\right| \approx 1$ on $\mathfrak{P}$ (within the plateau tolerances).
- **QII.** (**QII: law-visible, state-quenched**) $\left|\Lambda_{\mathrm{law}}\right| \approx 1$ on $\mathfrak{P}$, while $\Lambda_{\mathrm{state}}(V) \to 0$ as $V \to \infty$, where V is the IR averaging volume (e.g. $V \sim \ell^d$); more precisely, there exists $f(V) \downarrow 0$ with $\left|\Lambda_{\mathrm{state}}(V)\right| \le f(V)$ for V large on $\mathfrak{P}$.
- **QIII.** (**QIII: law recovery under IR coarse-graining**) For large ℓ one has the decay
$$1 - \left|\Lambda_{\mathrm{law}}(\ell)\right| \;=\; \mathcal{O}\Big(\ell^{\,\Delta-(d+z)}\Big) \qquad (\ell \to \infty),$$
reflecting suppression of irrelevant deformations in the law channel.
- **QIV.** (**QIV: law-deficit persists**) The law-level visibility remains uniformly below unity on $\mathfrak{P}$: there exists $\delta_\star \in (0,1)$ with $\left|\Lambda_{\mathrm{law}}(\ell)\right| \le 1 - \delta_\star$ for all ℓ in the plateau band.

Lemma 7.43 (Overlay invariance of quadrant assignment)**.** *Let $\mathcal{Q}$ be the quadrant-assignment rule used in this chapter: it depends only on the certified baseline (plateau, BRST/Nielsen gates, anomaly filters, tachyon/hyperbolicity tests) and on the* primary *visibilities entering the acceptance criteria of Section 7.3.2. If $\widetilde{\Lambda}_\alpha = \Phi_\alpha(\{\Lambda_\beta\}_\beta)$ is any overlay built from the visibilities by a channelwise continuous map Φ_α (e.g. normalisations, absolute values, ratios, logarithms) computed on $\mathcal{H}_{\mathrm{phys}}$, then*

$$\mathcal{Q}\Big(\{\Lambda_\beta\}_\beta\Big) \;=\; \mathcal{Q}\Big(\{\Lambda_\beta\}_\beta, \{\widetilde{\Lambda}_\alpha\}_\alpha\Big).$$

In words: adding overlays does not change quadrant membership.

Proof. By construction, $\mathcal{Q}$ is a function of (i) pass/fail outcomes for BRST, anomaly, and hyperbolicity gates; (ii) plateau acceptance inequalities (7.46) for a fixed visibility vector; and (iii) certified pole properties (all on $\mathcal{H}_{\mathrm{phys}}$). Overlays $\widetilde{\Lambda}_\alpha$ are continuous functions of the already certified Λ_β and are not arguments of $\mathcal{Q}$ by the protocol of Section 7.3.3. Therefore the value of $\mathcal{Q}$ is unchanged when $\{\widetilde{\Lambda}_\alpha\}_\alpha$ are added to the diagnostics set. □

Implementation note. All ingredients in (7.54) are band-safe and BRST-closed: the spectral projector $P_x(\lambda_{\mathrm{UV}})$ and the window $K_{\mathrm{IR}}^{(\ell)}$ commute with channel projectors and with $\hat{Q}_{\mathrm{BRST}}$ (eq. (7.30) and Definition 7.29). Hence scale scans preserve functoriality and cone structure, and the on-shell visibilities Λ_α are invariant under gauge-parameter changes (Nielsen identities, Section 7.2.9 and eqs. (7.22) and (7.23)). The overlays of Definition 7.42 are evaluated on $\mathcal{H}_{\mathrm{phys}}$ and are thus ghost/longitudinal-free by Section 7.2.6.

7.3.5 Plateau Regions: Geometric Definition

Definition 7.44 (Geometric plateau set in log chart)**.** Fix a compact scan window

$$\mathcal{R} := \big[\lambda_{\min}, \lambda_{\max}\big] \times \big[\ell_{\min}, \ell_{\max}\big] \subset \mathbb{R}^2_{>0},$$

and tolerances $\varepsilon_{\mathrm{grad}}, \varepsilon_{\mathrm{osc}}, \varepsilon_{\mathrm{curv}} > 0$. On $\mathbb{R}^2_{>0}$ adopt the log chart $(u, v) = (\ln \lambda_{\mathrm{UV}}, \ln \ell)$ and the flat metric

$$g_{\mathrm{plat}} := du^2 + dv^2,$$

so that $D_\lambda = \partial_u = \lambda_{\mathrm{UV}} \partial_{\lambda_{\mathrm{UV}}}$ and $D_\ell = \partial_v = \ell\, \partial_\ell$ (cf. eq. (7.43)). Let $B_r^{\mathrm{plat}}(\lambda, \ell)$ be the closed g_{plat}-geodesic ball of radius $r > 0$ centered at (λ, ℓ) in log coordinates. The *geometric plateau set* for channel α is

$$\begin{aligned}\mathfrak{P}_\alpha(\varepsilon_{\mathrm{grad}}, \varepsilon_{\mathrm{osc}}, \varepsilon_{\mathrm{curv}}; \mathcal{R}) := \Big\{ (\lambda_{\mathrm{UV}}, \ell) \in \mathcal{R} : \ & \|\nabla^{\mathrm{plat}} \Lambda_\alpha\|_\infty \le \varepsilon_{\mathrm{grad}}, \\ & \operatorname{osc}_{B_r^{\mathrm{plat}}(\lambda_{\mathrm{UV}}, \ell)} \Lambda_\alpha \le \varepsilon_{\mathrm{osc}}, \\ & \|\mathbf{H}^{\mathrm{plat}} \Lambda_\alpha\|_{\mathrm{op}} \le \varepsilon_{\mathrm{curv}} \Big\},\end{aligned} \tag{7.59}$$

where

$$\nabla^{\mathrm{plat}} \Lambda_\alpha = \big(D_\lambda \Lambda_\alpha,\ D_\ell \Lambda_\alpha\big), \qquad \mathbf{H}^{\mathrm{plat}} \Lambda_\alpha = \begin{pmatrix} D_\lambda^2 \Lambda_\alpha & D_\lambda D_\ell \Lambda_\alpha \\ D_\ell D_\lambda \Lambda_\alpha & D_\ell^2 \Lambda_\alpha \end{pmatrix},$$

and $\|\cdot\|_\infty, \|\cdot\|_{\mathrm{op}}$ are the sup and operator norms on $\mathcal{R}$.

Notation. Here $\nabla^{\mathrm{plat}} \Lambda_\alpha = (D_\lambda \Lambda_\alpha,\ D_\ell \Lambda_\alpha)$ with $D_\lambda = \partial_u = \lambda_{\mathrm{UV}} \partial_{\lambda_{\mathrm{UV}}}$ and $D_\ell = \partial_v = \ell\, \partial_\ell$. The Hessian in the g_{plat} chart is

$$\mathbf{H}^{\mathrm{plat}} \Lambda_\alpha = \begin{pmatrix} D_\lambda^2 \Lambda_\alpha & D_\lambda D_\ell \Lambda_\alpha \\ D_\ell D_\lambda \Lambda_\alpha & D_\ell^2 \Lambda_\alpha \end{pmatrix},$$

and $\|\mathbf{H}^{\mathrm{plat}} \Lambda_\alpha\|_{\mathrm{op}}$ is the operator norm induced by the Euclidean metric on the log chart (largest singular value/eigenvalue in 2D).

Lemma 7.45 (Oscillation bound in g_{plat}-balls)**.** *If $\Lambda_\alpha \in C^2$ on $B_r^{\mathrm{plat}}(\lambda,\ell)$, then*

$$\operatorname{osc}_{B_r^{\mathrm{plat}}(\lambda,\ell)} \Lambda_\alpha \;\le\; 2r \sup_{B_r^{\mathrm{plat}}} \|\nabla^{\mathrm{plat}}\Lambda_\alpha\| \;+\; r^2 \sup_{B_r^{\mathrm{plat}}} \|\mathbf{H}^{\mathrm{plat}}\Lambda_\alpha\|_{\mathrm{op}}.$$

Proof. Work in the log chart $(u,v) = (\ln \lambda_{\mathrm{UV}}, \ln \ell)$ where $g_{\mathrm{plat}} = du^2 + dv^2$. Let $f := \Lambda_\alpha$, and pick arbitrary points p, q in the g_{plat}-ball $B_r^{\mathrm{plat}}(\lambda,\ell)$. Write $p = (u_0, v_0)$, $q = (u_1, v_1)$ and set $h := q - p$, $L := \|h\| = \sqrt{(u_1-u_0)^2 + (v_1-v_0)^2}$. Because p, q lie in a radius-r ball, $L \le 2r$. Consider the straight segment (a g_{plat}–geodesic)

$$\gamma(t) := p + th, \qquad t \in [0,1].$$

Second-order Taylor with integral remainder. Using the standard formula at p along direction h,

$$f(q) = f(p) + \nabla^{\mathrm{plat}} f(p)\cdot h + \int_0^1 (1-t)\, h^{\top}\, \mathbf{H}^{\mathrm{plat}} f\big(\gamma(t)\big)\, h \, dt.$$

Taking absolute values and bounding each term uniformly on the ball gives

$$|f(q) - f(p)| \le \|\nabla^{\mathrm{plat}} f(p)\|\, \|h\| + \Big(\sup_{B_r^{\mathrm{plat}}} \|\mathbf{H}^{\mathrm{plat}} f\|_{\mathrm{op}} \Big) \int_0^1 (1-t)\, \|h\|^2 \, dt.$$

Since $\int_0^1 (1-t)\, dt = \frac{1}{2}$, we obtain

$$|f(q) - f(p)| \le L \sup_{B_r^{\mathrm{plat}}} \|\nabla^{\mathrm{plat}} f\| + \tfrac{1}{2} L^2 \sup_{B_r^{\mathrm{plat}}} \|\mathbf{H}^{\mathrm{plat}} f\|_{\mathrm{op}}.$$

Insert $L \le 2r$. This yields

$$|f(q) - f(p)| \le 2r \sup_{B_r^{\mathrm{plat}}} \|\nabla^{\mathrm{plat}} f\| + r^2 \sup_{B_r^{\mathrm{plat}}} \|\mathbf{H}^{\mathrm{plat}} f\|_{\mathrm{op}}.$$

Supremise over pairs. Taking the supremum over all $p, q \in B_r^{\mathrm{plat}}(\lambda,\ell)$ gives the desired bound for $\operatorname{osc}_{B_r^{\mathrm{plat}}(\lambda,\ell)} \Lambda_\alpha$. □

Proposition 7.46 (Compatibility with channelwise plateau criterion)**.** *Let $\mathfrak{P}_\alpha$ be as in Definition 7.44 and let the channelwise plateau set $\mathfrak{P}_\alpha(\mathcal{R}; \varepsilon_\lambda, \varepsilon_\ell, \varepsilon_{\mathrm{osc}})$ be as in (7.56). Fix any $r > 0$ and, in (7.56), choose the neighbourhood $\mathcal{N}$ to be the g_{plat}-ball $B_r^{\mathrm{plat}}(\lambda,\ell)$. Then:*

(i) If $(\lambda,\ell) \in \mathfrak{P}_\alpha(\varepsilon_\lambda, \varepsilon_\ell, \varepsilon_{\mathrm{osc}})$, then

$$(\lambda,\ell) \in \mathfrak{P}_\alpha(\varepsilon_{\mathrm{grad}}, \varepsilon_{\mathrm{osc}}, \varepsilon_{\mathrm{curv}}) \quad \text{with} \quad \varepsilon_{\mathrm{grad}} := \max\{\varepsilon_\lambda, \varepsilon_\ell\},$$

and any $\varepsilon_{\mathrm{curv}} \ge \sup_{B_r^{\mathrm{plat}}(\lambda,\ell)} \|\mathbf{H}^{\mathrm{plat}}\Lambda_\alpha\|_{\mathrm{op}}$.

(ii) Conversely, if $(\lambda,\ell) \in \mathfrak{P}_\alpha(\varepsilon_{\mathrm{grad}}, \varepsilon_{\mathrm{osc}}, \varepsilon_{\mathrm{curv}})$, then

$$(\lambda,\ell) \in \mathfrak{P}_\alpha(\varepsilon_\lambda, \varepsilon_\ell, \varepsilon_{\mathrm{osc}}) \quad \text{with} \quad \varepsilon_\lambda = \varepsilon_\ell := \varepsilon_{\mathrm{grad}}.$$

Proof. Work in the log chart $(u, v) = (\ln \lambda_{\mathrm{UV}}, \ln \ell)$, where $g_{\mathrm{plat}} = du^2 + dv^2$, $D_\lambda = \partial_u = \lambda_{\mathrm{UV}} \partial_{\lambda_{\mathrm{UV}}}$ and $D_\ell = \partial_v = \ell\, \partial_\ell$ (cf. eq. (7.43)). Recall the geometric objects:

$$\nabla^{\mathrm{plat}} \Lambda_\alpha = (D_\lambda \Lambda_\alpha,\ D_\ell \Lambda_\alpha), \qquad \mathbf{H}^{\mathrm{plat}} \Lambda_\alpha = \begin{pmatrix} D_\lambda^2 \Lambda_\alpha & D_\lambda D_\ell \Lambda_\alpha \\ D_\ell D_\lambda \Lambda_\alpha & D_\ell^2 \Lambda_\alpha \end{pmatrix}.$$

(i) Channelwise ⇒ geometric. Assume $(\lambda, \ell) \in \mathfrak{P}_\alpha(\varepsilon_\lambda, \varepsilon_\ell, \varepsilon_{\mathrm{osc}})$ with the neighbourhood chosen as $B_r^{\mathrm{plat}}(\lambda, \ell)$. By definition (7.56),

$$|D_\lambda \Lambda_\alpha| \le \varepsilon_\lambda, \qquad |D_\ell \Lambda_\alpha| \le \varepsilon_\ell, \qquad \operatorname{osc}_{B_r^{\mathrm{plat}}(\lambda,\ell)} \Lambda_\alpha \le \varepsilon_{\mathrm{osc}}.$$

Therefore,

$$\|\nabla^{\mathrm{plat}} \Lambda_\alpha\|_\infty = \max\{|D_\lambda \Lambda_\alpha|, |D_\ell \Lambda_\alpha|\} \le \max\{\varepsilon_\lambda, \varepsilon_\ell\} =: \varepsilon_{\mathrm{grad}},$$

and the oscillation bound is identical (same set B_r^{plat}). The geometric curvature constraint is not part of (7.56); choose any $\varepsilon_{\mathrm{curv}} \ge \sup_{B_r^{\mathrm{plat}}(\lambda,\ell)} \|\mathbf{H}^{\mathrm{plat}} \Lambda_\alpha\|_{\mathrm{op}}$ to satisfy the third clause of (7.59). Hence $(\lambda, \ell) \in \mathfrak{P}_\alpha(\varepsilon_{\mathrm{grad}}, \varepsilon_{\mathrm{osc}}, \varepsilon_{\mathrm{curv}})$.

(ii) Geometric ⇒ channelwise. Assume $(\lambda, \ell) \in \mathfrak{P}_\alpha(\varepsilon_{\mathrm{grad}}, \varepsilon_{\mathrm{osc}}, \varepsilon_{\mathrm{curv}})$. Then, by (7.59),

$$\max\{|D_\lambda \Lambda_\alpha|, |D_\ell \Lambda_\alpha|\} \le \varepsilon_{\mathrm{grad}}, \qquad \operatorname{osc}_{B_r^{\mathrm{plat}}(\lambda,\ell)} \Lambda_\alpha \le \varepsilon_{\mathrm{osc}}.$$

Set $\varepsilon_\lambda = \varepsilon_\ell := \varepsilon_{\mathrm{grad}}$. This yields $|D_\lambda \Lambda_\alpha| \le \varepsilon_\lambda$, $|D_\ell \Lambda_\alpha| \le \varepsilon_\ell$ and the same oscillation bound in (7.56) (with $\mathcal{N} = B_r^{\mathrm{plat}}$), proving that $(\lambda, \ell) \in \mathfrak{P}_\alpha(\varepsilon_\lambda, \varepsilon_\ell, \varepsilon_{\mathrm{osc}})$.

This establishes both inclusions under the stated identification of neighbourhoods and tolerances. □

Parameter-space geometry (log chart).

Definition 7.47 (Plateau parameter-space geometry)**.** On $\mathbb{R}^2_{>0}$ define the log chart $(u, v) = (\ln \lambda_{\mathrm{UV}}, \ln \ell)$ and equip it with the flat metric

$$g_{\mathrm{plat}} := du^2 + dv^2.$$

Denote the associated orthonormal derivatives by

$$D_\lambda := \partial_u = \lambda_{\mathrm{UV}} \partial_{\lambda_{\mathrm{UV}}}, \qquad D_\ell := \partial_v = \ell\, \partial_\ell.$$

For a C^2 scalar field f on parameter space set

$$\nabla^{\mathrm{plat}} f := (D_\lambda f,\ D_\ell f), \qquad \mathbf{H}^{\mathrm{plat}} f := \begin{pmatrix} D_\lambda^2 f & D_\lambda D_\ell f \\ D_\ell D_\lambda f & D_\ell^2 f \end{pmatrix}.$$

For $r > 0$, $B_r^{\mathrm{plat}}(\lambda, \ell)$ denotes the closed g_{plat}-geodesic ball of radius r centered at (λ, ℓ) in log coordinates.

Lemma 7.48 (Norm identities in the log geometry)**.** *For any visibility Λ_α and any compact $\mathcal{R} \subset \mathbb{R}^2_{>0}$,*

$$\|\nabla^{\mathrm{plat}} \Lambda_\alpha\|_\infty = \sup_{\mathcal{R}} \max\{|D_\lambda \Lambda_\alpha|, |D_\ell \Lambda_\alpha|\}, \qquad \|\mathbf{H}^{\mathrm{plat}} \Lambda_\alpha\|_{\mathrm{op}} = \sup_{\mathcal{R}} \left\| \begin{pmatrix} D_\lambda^2 \Lambda_\alpha & D_\lambda D_\ell \Lambda_\alpha \\ D_\ell D_\lambda \Lambda_\alpha & D_\ell^2 \Lambda_\alpha \end{pmatrix} \right\|_{\mathrm{op}}.$$

Moreover, B_r^{plat} is a true geodesic (Euclidean) ball in (u, v).

Proof. By Definition 7.47, g_{plat} is the Euclidean metric in (u, v), so the coordinate basis is orthonormal and the stated expressions are exactly the Euclidean sup and operator norms. Geodesics are straight lines, hence B_r^{plat} is a Euclidean ball in (u, v). □

Link to global certification metrics.

Proposition 7.49 (From geometric bounds to certification metrics)**.** *Let $\mathcal{N}_\delta \subset \mathcal{R}$ be a rectangle around $(\lambda_\star, \ell_\star)$ and fix a visibility $\Lambda_\alpha \in C^2(\mathcal{N}_\delta)$. If on $\mathcal{N}_\delta$*

$$\|\nabla^{\mathrm{plat}} \Lambda_\alpha\| \le \varepsilon_{\mathrm{grad}}, \qquad \|\mathbf{H}^{\mathrm{plat}} \Lambda_\alpha\|_{\mathrm{op}} \le \varepsilon_{\mathrm{curv}},$$

then, writing $R_\lambda := |\ln(\lambda/\lambda_\star)|$ and $R_\ell := |\ln(\ell/\ell_\star)|$,

$$\big|\Lambda_\alpha(\lambda,\ell) - \Lambda_\alpha(\lambda_\star,\ell_\star)\big| \;\le\; \varepsilon_{\mathrm{grad}}(R_\lambda + R_\ell) \;+\; \tfrac{1}{2}\,\varepsilon_{\mathrm{curv}}(R_\lambda^2 + R_\ell^2), \qquad \forall\,(\lambda,\ell) \in \mathcal{N}_\delta. \tag{7.60}$$

If $\mathbf{V} = (\Lambda_{\alpha_1}, \dots, \Lambda_{\alpha_m})$ is a finite visibility vector with componentwise bounds $(\varepsilon_{\mathrm{grad}}^{(j)}, \varepsilon_{\mathrm{curv}}^{(j)})$, then the certification seminorms of eq. (7.42) obey

$$\begin{aligned} \Delta_{\mathrm{rel}}(\mathbf{V};\mathcal{N}_\delta) &\le \max_j \frac{\varepsilon_{\mathrm{grad}}^{(j)}(R_\lambda^{\max} + R_\ell^{\max}) \\ &\qquad + \tfrac{1}{2}\,\varepsilon_{\mathrm{curv}}^{(j)}\Big((R_\lambda^{\max})^2 + (R_\ell^{\max})^2\Big)}{\max\Big(1, \|\mathbf{V}^{(\lambda_\star,\ell_\star)}\|\Big)}, \\ \kappa_{\mathrm{rel}}(\mathbf{V};\mathcal{N}_\delta) &\le \max_j \frac{\varepsilon_{\mathrm{curv}}^{(j)}}{\max\Big(1, \|\mathbf{V}^{(\lambda_\star,\ell_\star)}\|\Big)}, \end{aligned} \tag{7.61}$$

where $R_\lambda^{\max} := \sup_{\mathcal{N}_\delta} R_\lambda$, $R_\ell^{\max} := \sup_{\mathcal{N}_\delta} R_\ell$.

Proof. For the scalar case, work in the log variables $u = \ln\lambda$, $v = \ln\ell$ and set $F(u,v) := \Lambda_\alpha(e^u, e^v)$. Let $(u_\star, v_\star)$ be the base point and (u,v) any point in $\mathcal{N}_\delta$. Along the straight segment $\gamma(t) = (u_\star, v_\star) + t\,(u - u_\star, v - v_\star)$, $t \in [0,1]$, Taylor's theorem with integral remainder gives

$$F(u,v) - F(u_\star,v_\star) = \nabla F(u_\star,v_\star)\cdot(\Delta u, \Delta v) + \int_0^1 (1-t)\,\begin{pmatrix}\Delta u & \Delta v\end{pmatrix}\mathbf{H}F(\gamma(t))\begin{pmatrix}\Delta u \\ \Delta v\end{pmatrix} dt,$$

with $(\Delta u, \Delta v) = (u - u_\star, v - v_\star) = (R_\lambda, R_\ell)$. Taking absolute values and using the bounds $\|\nabla F\| \le \varepsilon_{\mathrm{grad}}$, $\|\mathbf{H}F\|_{\mathrm{op}} \le \varepsilon_{\mathrm{curv}}$ (which are exactly the hypotheses written in log variables), we obtain

$$|F(u,v) - F(u_\star,v_\star)| \le \varepsilon_{\mathrm{grad}}\,\|(R_\lambda, R_\ell)\|_1 + \tfrac{1}{2}\,\varepsilon_{\mathrm{curv}}\,\|(R_\lambda, R_\ell)\|_2^2,$$

i.e. (7.60). For the vector case, apply the scalar bound componentwise and take the ℓ^∞ norm used in eq. (7.42), which yields (7.61). The curvature seminorm in eq. (7.42) is, by eq. (7.44), controlled by the operator norm of the Hessian, hence the stated bound on κ_{rel}. □

By Lemma 7.45 (oscillation in B_r^{plat}) and Proposition 7.49, small geometric bounds (gradient/Hessian) combined with small local oscillation provide a direct route from the geometric set (7.59) to the global acceptance test (7.46). This removes coordinate bias and aligns precisely with the scale gradients of eq. (7.55).

Lemma 7.50 (Equivalence to the metric criterion)**.** *Let $\mathbf{V}$ be any finite collection of visibilities including Λ_α. On compact $\mathcal{R}$, there exist constants $c_1, c_2 > 0$ such that the geometric conditions above are equivalent to the drift/curvature bounds of Section 7.3.2, i.e.*

$$\Delta_{\mathrm{rel}}(\mathbf{V};\mathcal{N}_\delta) \le c_1\,\varepsilon_{\mathrm{grad}} + o(1), \qquad \kappa_{\mathrm{rel}}(\mathbf{V};\mathcal{N}_\delta) \le c_2\,\varepsilon_{\mathrm{curv}} + o(1),$$

for $\mathcal{N}_\delta$ compatible with the ball radius r and with $o(1) \to 0$ as $\delta \to 0$. In particular, acceptance in Definition 7.31 is equivalent (up to tolerance rescaling) to membership in $\mathfrak{P}_\alpha$.

Proof. **Setup.** Fix a compact rectangle $\mathcal{N}_\delta \subset \mathcal{R}$ around $(\lambda_\star, \ell_\star)$ and work in the log chart $(u, v) = (\ln \lambda_{\mathrm{UV}}, \ln \ell)$ with metric g_{plat} (cf. Definition 7.47). For brevity write

$$R_\lambda^{\max} := \sup_{(\lambda,\ell)\in\mathcal{N}_\delta} \left|\ln(\lambda/\lambda_\star)\right|, \qquad R_\ell^{\max} := \sup_{(\lambda,\ell)\in\mathcal{N}_\delta} \left|\ln(\ell/\ell_\star)\right|,$$

and let $\rho_1 := R_\lambda^{\max} + R_\ell^{\max}$ and $\rho_2 := (R_\lambda^{\max})^2 + (R_\ell^{\max})^2$. Note that $\rho_1, \rho_2 \to 0$ as $\delta \to 0$.

(A) Geometric bounds ⇒ drift bound. For a scalar visibility Λ_α, Proposition 7.49 gives

$$\left|\Lambda_\alpha(\lambda,\ell) - \Lambda_\alpha(\lambda_\star,\ell_\star)\right| \le \varepsilon_{\mathrm{grad}}\,(R_\lambda^{\max} + R_\ell^{\max}) + \tfrac{1}{2}\,\varepsilon_{\mathrm{curv}}\Big((R_\lambda^{\max})^2 + (R_\ell^{\max})^2\Big)$$

uniformly on $\mathcal{N}_\delta$. For a finite vector $\mathbf{V}$ we apply the bound componentwise and take the ℓ^∞ norm used in eq. (7.42). Dividing by $\max\Big(1, \|\mathbf{V}^{(\lambda_\star,\ell_\star)}\|\Big)$ yields

$$\Delta_{\mathrm{rel}}(\mathbf{V};\mathcal{N}_\delta) \le \frac{\rho_1}{\max(1, \|\mathbf{V}^{(\lambda_\star,\ell_\star)}\|)}\,\varepsilon_{\mathrm{grad}} \;+\; \frac{\rho_2}{2\,\max(1, \|\mathbf{V}^{(\lambda_\star,\ell_\star)}\|)}\,\varepsilon_{\mathrm{curv}}.$$

Set $c_1 := \rho_1 / \max(1, \|\mathbf{V}^{(\lambda_\star,\ell_\star)}\|)$ and $o(1) := \frac{\rho_2}{2\,\max(1,\|\mathbf{V}^{(\lambda_\star,\ell_\star)}\|)}\,\varepsilon_{\mathrm{curv}}$, which satisfies $o(1) \to 0$ as $\delta \to 0$. This proves the first inequality.

(B) Geometric bounds ⇒ curvature bound. We convert the log–chart second derivatives to the linear (λ, ℓ) derivatives appearing in eq. (7.42). By chain rule (with $D_\lambda = \lambda\partial_\lambda$, $D_\ell = \ell\partial_\ell$):

$$\partial_\lambda^2\Lambda_\alpha = \frac{1}{\lambda^2}\Big(D_\lambda^2\Lambda_\alpha - D_\lambda\Lambda_\alpha\Big), \quad \partial_\ell^2\Lambda_\alpha = \frac{1}{\ell^2}\Big(D_\ell^2\Lambda_\alpha - D_\ell\Lambda_\alpha\Big), \quad \partial_\lambda\partial_\ell\Lambda_\alpha = \frac{1}{\lambda\,\ell}\,D_\lambda D_\ell\Lambda_\alpha.$$

Hence, on the compact domain $\mathcal{R}$,

$$\left\|\partial_{\lambda,\ell}^2\Lambda_\alpha\right\| \;\le\; C_A\left\|\mathbf{H}^{\mathrm{plat}}\Lambda_\alpha\right\|_{\mathrm{op}} \;+\; C_B\left\|\nabla^{\mathrm{plat}}\Lambda_\alpha\right\|,$$

with constants

$$C_A := \max\left\{\frac{1}{\lambda_{\min}^2}, \frac{1}{\lambda_{\min}\ell_{\min}}, \frac{1}{\ell_{\min}^2}\right\}, \qquad C_B := \max\left\{\frac{1}{\lambda_{\min}^2}, \frac{1}{\ell_{\min}^2}\right\}.$$

Using the geometric assumptions $\left\|\mathbf{H}^{\mathrm{plat}}\Lambda_\alpha\right\|_{\mathrm{op}} \le \varepsilon_{\mathrm{curv}}$ and $\|\nabla^{\mathrm{plat}}\Lambda_\alpha\| \le \varepsilon_{\mathrm{grad}}$, we obtain

$$\sup_{\mathcal{N}_\delta}\left\|\partial_{\lambda,\ell}^2\Lambda_\alpha\right\| \;\le\; C_A\,\varepsilon_{\mathrm{curv}} + C_B\,\varepsilon_{\mathrm{grad}}.$$

For a visibility vector $\mathbf{V}$, take the ℓ^∞ norm across components and divide by $\max\Big(1, \|\mathbf{V}^{(\lambda_\star,\ell_\star)}\|\Big)$ to match eq. (7.42). Thus

$$\kappa_{\mathrm{rel}}(\mathbf{V};\mathcal{N}_\delta) \;\le\; \frac{C_A}{\max(1, \|\mathbf{V}^{(\lambda_\star,\ell_\star)}\|)}\,\varepsilon_{\mathrm{curv}} \;+\; \frac{C_B}{\max(1, \|\mathbf{V}^{(\lambda_\star,\ell_\star)}\|)}\,\varepsilon_{\mathrm{grad}}.$$

Set $c_2 := C_A / \max(1, \|\mathbf{V}^{(\lambda_\star,\ell_\star)}\|)$. The extra gradient term can be treated as part of the $o(1)$ in the statement: if one tightens the gradient tolerance in the geometric condition (which

is permitted in certification), this addend is arbitrarily small; alternatively, on shrinking neighbourhoods its contribution is dominated by the drift scale (cf. eq. (7.47)). In either case, we may write $\kappa_{\mathrm{rel}}(\mathbf{V};\mathcal{N}_\delta) \le c_2\,\varepsilon_{\mathrm{curv}} + o(1)$.

(C) Equivalence of acceptance. The forward implications above show that membership in the geometric set (small gradient/Hessian/oscillation) implies the metric acceptance inequalities of Definition 7.31 after a tolerance rescaling by c_1, c_2 and a vanishing remainder $o(1)$. Conversely, the set–theoretic compatibility of Proposition 7.46 (channelwise vs. geometric plateau) together with the mean–value bound eq. (7.47) provides the reverse implication up to rescaling of tolerances, proving the equivalence of notions in the sense of certification. □

Invariances and robustness.

Proposition 7.51 (Gauge-parameter invariance of plateau membership)**.** *Let Λ_α be a holosymmetric visibility evaluated on $\mathcal{H}_{\mathrm{phys}}$ and assume the Nielsen framework of Section 7.2.9 with $\partial_{\alpha_{\mathrm{gf}}}\Lambda_\alpha = 0$ on shell (cf. eqs. (7.22) and (7.23)). Then, for any tolerances and scan region,*

$$(\lambda,\ell) \in \mathfrak{P}_\alpha(\mathcal{R};\varepsilon_\lambda,\varepsilon_\ell,\varepsilon_{\mathrm{osc}}) \quad\Longleftrightarrow\quad (\lambda,\ell) \in \mathfrak{P}_\alpha(\mathcal{R};\varepsilon_\lambda,\varepsilon_\ell,\varepsilon_{\mathrm{osc}}) \text{ computed at any } \alpha'_{\mathrm{gf}}.$$

Proof. By Section 7.2.9 the 1PI two–point structures determining Λ_α are α_{gf}–stationary on shell, hence so are Λ_α, $D_\lambda\Lambda_\alpha$ and $D_\ell\Lambda_\alpha$ (log–derivatives commute with on–shell differentiation in the band formalism). Local oscillation depends only on pointwise values over neighbourhoods and is therefore invariant as well. Thus each inequality in (7.56) holds simultaneously for all α_{gf}. □

Proposition 7.52 (BRST/Slavnov stability of scale derivatives)**.** *Assume $\zeta P_x(\lambda_{\mathrm{UV}}) = \zeta K_{\mathrm{IR}}^{(\ell)} = 0$ with ghost number 0 eq. (7.30) and $\mathcal{S}(\Gamma) = 0$ eq. (7.16). Then the insertions defining $D_\lambda\Lambda_\alpha$ and $D_\ell\Lambda_\alpha$ are BRST closed on $\mathcal{H}_{\mathrm{phys}}$, and positivity/unitarity diagnostics (Källén–Lehmann, optical theorem) are preserved under the scale scan.*

Proof. Band tagging is "UV then IR" (cf. Definitions 7.28 and 7.29); differentiating w.r.t. $\ln\lambda_{\mathrm{UV}}$ and $\ln\ell$ acts inside the BRST-closed functor because ζ commutes with $P_x(\lambda)$ and $K_{\mathrm{IR}}^{(\ell)}$. Since $\mathcal{S}(\Gamma) = 0$, ζ–exact variations of the gauge fermion do not affect on–shell correlators; hence the resulting scale insertions are BRST closed on $\mathcal{H}_{\mathrm{phys}}$. Positivity/unitarity then follow from the KL representation established for gauge-invariant observables on $\mathcal{H}_{\mathrm{phys}}$ (Corollary 7.24). □

Lemma 7.53 (Regulator-induced visibility and gradient drifts)**.** *Let $(P,K) \mapsto (P',K')$ be an admissible regulator change in the sense of Definition 7.32 with mismatch metrics $(\eta_{\mathrm{UV}},\eta_{\mathrm{IR}})$. Assume further first–log–derivative mismatch bounds*

$$\dot\eta_{\mathrm{UV}} := \sup_{(\lambda,\ell)\in\mathcal{R}} \big\|\lambda\,\partial_\lambda P' - \lambda\,\partial_\lambda P\big\|, \qquad \dot\eta_{\mathrm{IR}} := \sup_{(\lambda,\ell)\in\mathcal{R}} \big\|\ell\,\partial_\ell K_{\mathrm{IR}}^{(\ell)\prime} - \ell\,\partial_\ell K_{\mathrm{IR}}^{(\ell)}\big\|_{L^1}.$$

Then, with the Lipschitz moduli $L_\alpha^{(P)}, L_\alpha^{(K)}$ from Definition 7.69,

$$\sup_{\mathcal{R}} \big|\Lambda'_\alpha - \Lambda_\alpha\big| \le L_\alpha^{(P)}\,\eta_{\mathrm{UV}} + L_\alpha^{(K)}\,\eta_{\mathrm{IR}} \;=:\; \varepsilon_{\mathrm{reg}}, \tag{7.62}$$

$$\sup_{\mathcal{R}} \big|D_\lambda\Lambda'_\alpha - D_\lambda\Lambda_\alpha\big| \le L_\alpha^{(P)}\,\dot\eta_{\mathrm{UV}} + L_\alpha^{(K)}\,\eta_{\mathrm{IR}}, \qquad \sup_{\mathcal{R}} \big|D_\ell\Lambda'_\alpha - D_\ell\Lambda_\alpha\big| \le L_\alpha^{(P)}\,\eta_{\mathrm{UV}} + L_\alpha^{(K)}\,\dot\eta_{\mathrm{IR}}. \tag{7.63}$$

Proof. The map $\mathcal{F}_\alpha$ is Lipschitz in each tagged input (compressed by P and averaged by K), hence (7.62) follows from Definition 7.32 exactly as in (7.57). For (7.63), differentiate the tagged objects using the "UV then IR" calculus of Definition 7.28; the only new contributions relative to the original regulator come from replacing $(\lambda\partial_\lambda P,\ \ell\,\partial_\ell K)$ by their primed counterparts, yielding the stated bounds with the same Lipschitz constants. □

Proposition 7.54 (Regulator robustness of plateau membership)**.** *Let $\mathfrak{P}_\alpha$ be computed with (P, K) and $\mathfrak{P}'_\alpha$ with (P', K') as above. If*

$$\varepsilon_{\mathrm{reg}} \ll \min\{\varepsilon_\lambda, \varepsilon_\ell, \varepsilon_{\mathrm{osc}}\} \quad \text{and} \quad L_\alpha^{(P)}\,\dot\eta_{\mathrm{UV}} + L_\alpha^{(K)}\,\dot\eta_{\mathrm{IR}} \ \le\ \min\{\varepsilon_\lambda, \varepsilon_\ell\} - \delta$$

for some $\delta > 0$, then $\mathfrak{P}_\alpha = \mathfrak{P}'_\alpha$ (same membership) on $\mathcal{R}$.

Proof. Use (7.62) to keep oscillation changes below $\varepsilon_{\mathrm{osc}}$, and (7.63) to ensure both log–gradients stay within the original thresholds; membership inequalities (7.56) are preserved. □

Boundary behaviour and commuting limits.

Proposition 7.55 (Finite crossing under commuting limits)**.** *Assume commuting limits (7.45) for Λ_α on $\mathcal{H}_{\mathrm{phys}}$. Suppose there exists a corner neighbourhood U of $(\infty, 0^+)$ in log coordinates where*

$$\|\nabla^{\mathrm{plat}}\Lambda_\alpha\| \le \varepsilon_{\mathrm{grad}}, \qquad \|\mathbf{H}^{\mathrm{plat}}\Lambda_\alpha\|_{\mathrm{op}} \le \varepsilon_{\mathrm{curv}}, \qquad \mathrm{osc}_{B_r^{\mathrm{plat}}}\Lambda_\alpha \le \varepsilon_{\mathrm{osc}}.$$

Then any compact path γ approaching $(\infty, 0^+)$ eventually remains in a single connected component of $\mathfrak{P}_\alpha$; in particular γ crosses only finitely many components.

Proof. By commuting limits and uniform bounds in U, $\Lambda_\alpha \circ \gamma$ is Cauchy near the corner and its first two log–derivatives are uniformly bounded by the plateau tolerances. The complement of $\mathfrak{P}_\alpha$ is the union of closed sets where at least one inequality is violated; by uniform bounds in U, γ cannot accumulate infinitely many violations without contradicting Cauchy convergence. Hence, after some parameter value, γ lies entirely in a single connected component. □

Maximal components and certification.

Definition 7.56 (Maximal components)**.** Let $\{\mathcal{P}_\alpha^{(k)}\}_k$ be the connected components of $\mathfrak{P}_\alpha$ in the g_{plat} topology. A component $\mathcal{P}_\alpha^{(k)}$ is *maximal* if, for any stricter tolerance triple $(\tilde\varepsilon_\lambda, \tilde\varepsilon_\ell, \tilde\varepsilon_{\mathrm{osc}})$ with each entry $\le$ the original and at least one strict, no proper superset of $\mathcal{P}_\alpha^{(k)}$ is contained in the refined plateau set.

Proposition 7.57 (Certification attaches to maximal components)**.** *If a component $\mathcal{P}_\alpha^{(k)}$ passes the certification protocol of Section 7.3.3, then any refinement of tolerances that preserves membership of $\mathcal{P}_\alpha^{(k)}$ keeps the certificate valid (with rescaled numerical bounds). Conversely, if a stricter triple excludes points of $\mathcal{P}_\alpha^{(k)}$, the certificate localises to the largest remaining connected subset, which is again maximal for the refined triple.*

Proof. Monotonicity and closedness of the plateau sets (cf. Proposition 7.38) imply stability of the inequalities under tolerance tightening; the protocol's metrics scale linearly with the thresholds (see eqs. (7.42) and (7.48)), giving rescaled bounds. Loss of points yields a closed subset; maximality then follows from the definition. □

Simultaneous (multi-channel) plateaus. For a set $\mathfrak{C} \subset \{\text{law}, \text{state}, \ldots\}$, recall the joint plateau

$$\mathfrak{P}_{\mathfrak{C}} := \bigcap_{\alpha \in \mathfrak{C}} \mathfrak{P}_\alpha. \tag{7.64}$$

Proposition 7.58 (Certification on intersections)**.** *Each $\mathfrak{P}_\alpha$ is closed; hence $\mathfrak{P}_{\mathfrak{C}}$ is closed. If every $\mathfrak{P}_\alpha$ carries a valid certificate with the same $(\varepsilon_\lambda, \varepsilon_\ell, \varepsilon_{\text{osc}})$, then $\mathfrak{P}_{\mathfrak{C}}$ carries a joint certificate with the same tolerances. If channels have heterogeneous numerical stability, one may use weighted thresholds $\mathbf{w} = (w_\alpha)_{\alpha \in \mathfrak{C}}$ and require $|D_\lambda \Lambda_\alpha| \le w_\alpha \varepsilon_\lambda$, $|D_\ell \Lambda_\alpha| \le w_\alpha \varepsilon_\ell$, $\text{osc} \le \min_{\alpha \in \mathfrak{C}} w_\alpha \varepsilon_{\text{osc}}$.*

Proof. Closedness is immediate. The certification protocol (Section 7.3.3) is conjunctive across channels: drift/curvature/oscillation bounds and the swap-limits test are each satisfied for all $\alpha \in \mathfrak{C}$, hence for their intersection. Weighted thresholds simply rescale the inequalities componentwise. □

Computational notes (practical)

Grid and evaluation (log chart). All scans are performed in the plateau log geometry Definition 7.47, with coordinates $(u, v) = (\ln \lambda_{\text{UV}}, \ln \ell)$ and orthonormal derivatives $D_\lambda = \partial_u$, $D_\ell = \partial_v$. Over the compact window $\mathcal{R}$ we lay down a uniform rectangular grid (u_i, v_j) with spacings $(\Delta u, \Delta v)$. At each grid point we evaluate the visibilities Λ_α on $\mathcal{H}_{\text{phys}}$ (i.e. after the Π_{T}-projection and BRST projection), and cache the associated physical diagnostics: spectral poles m_i^2 and residues/spectral weights entering the chosen visibility set; hyperbolicity flags for the principal symbol test eq. (7.15); and anomaly/Slavnov checks ensuring the quantum BRST constraints (eq. (7.16)).

Derivatives, local oscillation, and optional smoothing. Scale gradients are estimated by centred finite differences in (u, v), providing approximations to $D_\lambda \Lambda_\alpha$ and $D_\ell \Lambda_\alpha$ (and, via standard stencils, the Hessian entries in the log chart). Local oscillation is measured over a small g_{plat}-ball B_r^{plat}, as in the geometric definition Definition 7.44. When data are noisy, we use the box-averaged visibility $\overline{\Lambda}_\alpha$ of Definition 7.39 and re-estimate derivatives from the smoothed field; the bias and tolerance rescalings are controlled by Lemma 7.40 and Corollary 7.41.

Plateau extraction and components. For each channel α, we construct a discrete plateau mask by enforcing, at each grid point, the inequalities of the geometric plateau set Definition 7.44—equivalently, the channelwise criterion in (7.56). The accepted pixels form a discrete approximation to $\mathfrak{P}_\alpha$. Using standard connected-component labelling (with 8-neighbourhood in the (u, v) grid, approximating g_{plat} connectivity), we identify components and record their basic descriptors (cardinality/area, geodesic diameter in the log metric, centroids, and proximity to the scan boundary).

Joint plateaus and maximality. Given a set of channels $\mathfrak{C} \subset \{\text{law}, \text{state}, \ldots\}$, the joint plateau is the intersection $\mathfrak{P}_{\mathfrak{C}} = \bigcap_{\alpha \in \mathfrak{C}} \mathfrak{P}_\alpha$ (7.64). For reporting and certification we retain

maximal components in the sense of the tolerance order (no strict enlargement survives a strictly tighter tolerance triple); this is the notion used to attach certificates and to compare scans across runs.

Component-wise certification metrics. Within each component we recompute the global certification seminorms (7.42) for the chosen visibility vector $\mathbf{V}$ (e.g. pole masses, residues, spectral weights), thereby cross-checking the geometric gate via the drift/curvature metrics. The discrete "swap-limits" check for (7.45) is performed along two canonical paths to the corner $(\lambda \to \infty,\ \ell \to 0^+)$ confined to the component (increase u then decrease v, and vice versa), and we require the sup difference of $\mathbf{V}$ at the endpoints to be within the tolerance $\varepsilon_{\mathrm{swap}}$.

Gauge-parameter and regulator robustness. Nielsen identities imply on-shell gauge-parameter independence of the primary observables; hence plateau membership is invariant under changes of α_{gf} when evaluated on $\mathcal{H}_{\mathrm{phys}}$ (Proposition 7.51). Regulator robustness is assessed by repeating the scan with an admissible alternative projector/window pair and demanding that the induced drifts of visibilities and gradients remain within a strict regulator budget, which—by Proposition 7.54—preserves membership equivalence across regulators.

Pole tracking and hyperbolicity. For each component we track every physical pole m_i^2 across the grid, enforcing continuity (and only expected level crossings). We report the mean, standard deviation, and extremal relative drift Δ_i^{rel} as defined in (7.50), and verify that the hyperbolicity flags (principal symbol positivity eq. (7.15)) hold uniformly throughout the component.

Outputs and practicalities. The recommended artefacts per component are: (i) a compact table of tolerances, size/diameter, drift/curvature metrics, swap-limit outcome, regulator/Nielsen flags, and stable poles with statistics; and (ii) a "heatmap+mask" visual in (u, v) showing Λ_α, its gradient magnitude, and accepted pixels, with the joint-plateau overlay. In implementation, keep all computations in the log chart, pad stencils near boundaries (or use one-sided differences), and, when Λ_α is near ± 1, consider working with a stabilising monotone transform (e.g. arctanh) before differencing. Prefer components whose log-diameter greatly exceeds one grid step and whose pole statistics are consistent—these typically coincide with the maximal components certified above.

7.3.6 Gauge/BRST Robustness (Nielsen-type Independence)

Definition 7.59 (Nielsen setup and master identity)**.** Let α_{gf} be a gauge-fixing parameter entering the BRST gauge fermion $\Psi(\alpha_{\mathrm{gf}})$, and extend BRST by the standard doublet $\zeta\,\alpha_{\mathrm{gf}} = \chi$, $\zeta\,\chi = 0$. Assuming the Slavnov identity $\mathcal{S}(\Gamma) = 0$ eq. (7.16), the 1PI effective action satisfies the Nielsen *master identity* [117, 91]:

$$\partial_{\alpha_{\mathrm{gf}}}\Gamma = \int d^d x\ \frac{\delta\Gamma}{\delta\Phi^A(x)}\,\mathcal{N}^A(\alpha_{\mathrm{gf}}; x)\ +\ (\text{ghost/aux. insertions}), \qquad \mathcal{N}^A := \left\langle \zeta \frac{\partial\Psi}{\partial\alpha_{\mathrm{gf}}} \right\rangle_\Gamma, \tag{7.65}$$

where A runs over the multiplet $\mathfrak{X}$ and the expectation is 1PI/band-safe (cf. Definitions 7.28 and 7.29).

Lemma 7.60 (Band-safe differentiation in $\alpha_{\rm gf}$)**.** *Let $\Lambda_\alpha(\lambda_{\rm UV},\ell)$ be a holosymmetric visibility as in eq. (7.54), constructed from band-limited, IR-smoothed correlators with $P_x(\lambda_{\rm UV})$ and $K_{\rm IR}^{(\ell)}$. If $P_x(\lambda_{\rm UV})$ and $K_{\rm IR}^{(\ell)}$ are BRST closed and gauge-parameter independent eq. (7.30), then*

$$\partial_{\alpha_{\rm gf}}\Lambda_\alpha = \Big\langle \mathcal{O}_\alpha \cdot \zeta \frac{\partial \Psi}{\partial \alpha_{\rm gf}} \Big\rangle + \Big\langle \mathcal{J}_\alpha \cdot \frac{\delta \Gamma}{\delta \Phi} \Big\rangle,$$

for some local composites $\mathcal{O}_\alpha, \mathcal{J}_\alpha$ determined by the functional $\mathcal{F}_\alpha$ in eq. (7.54), with the averages taken in the same (band-safe) 1PI sense.

Proof. Differentiate $\Lambda_\alpha = \mathcal{F}_\alpha[\text{tagged correlators}; \bar{\Sigma}^{(\ell)}]$ with respect to $\alpha_{\rm gf}$. By construction, $\partial_{\alpha_{\rm gf}}$ acts only through Γ (the tagged kernels are independent of $\alpha_{\rm gf}$ and commute with BRST). Apply the chain rule and insert eq. (7.19); reorganise the result into the $\zeta(\partial\Psi/\partial\alpha_{\rm gf})$ insertion and the EOM insertion $\delta\Gamma/\delta\Phi$. □

Proposition 7.61 (Gauge-parameter independence of visibilities on shell)**.** *Assume: (i) anomaly filters and Slavnov identity hold eqs. (7.16) and (7.18); (ii) quartet decoupling and positivity on $\mathcal{H}_{\rm phys}$ Section 7.2.6 and eq. (7.12); and (iii) the scale-tagged EOM eqs. (6.333) and (6.334). Then every channel visibility obeys*

$$\frac{\partial}{\partial \alpha_{\rm gf}} \Lambda_\alpha(\lambda_{\rm UV},\ell) = 0 \qquad \text{on shell in } \mathcal{H}_{\rm phys}. \tag{7.66}$$

Proof. From Lemma 7.60 and eq. (7.19) we have $\partial_{\alpha_{\rm gf}}\Lambda_\alpha = \langle \mathcal{O}_\alpha \cdot \zeta(\partial\Psi/\partial\alpha_{\rm gf})\rangle + \langle \mathcal{J}_\alpha \cdot (\delta\Gamma/\delta\Phi)\rangle$. The first term is a ζ-exact insertion and vanishes on $\mathcal{H}_{\rm phys}$ by the quartet mechanism and positivity (no hidden negative-norm contributions) Section 7.2.6 and eq. (7.12); the second term vanishes on shell by the scale-tagged EOM. □

Corollary 7.62 (Plateau invariance under gauge-parameter changes)**.** *On $\mathcal{H}_{\rm phys}$, the inequalities defining the channelwise/geometric plateau sets (e.g. (7.56) and Definition 7.44) are unchanged under $\alpha_{\rm gf} \mapsto \alpha'_{\rm gf}$. In particular, membership in $\mathfrak{P}_\alpha$ and in any joint $\mathfrak{P}_{\mathfrak{C}}$ is gauge-parameter independent.*

Proof. By Proposition 7.61, the visibilities are $\alpha_{\rm gf}$-stationary on shell; their log–derivatives and local oscillations—computed from pointwise values over neighbourhoods—inherit the same stationarity. Hence each plateau inequality is preserved for any $\alpha'_{\rm gf}$. □

Pole statement (for reference). Projecting eq. (7.19) to the $\Pi_{\rm T}$ two-point kernel and linearising about on-shell configurations yields the standard pole relations eqs. (7.20), (7.22) and (7.23), ensuring that physical masses and the spectrum of the operator $\mathbb{M}^2_{\rm phys}$ are gauge-parameter stable [117, 91, 114, 82].

Proposition 7.63 (Gauge-parameter invariance of plateau membership)**.** *Assume eqs. (7.16) and (7.18) and the band BRST-closure eq. (7.30). Then for any tolerances $(\varepsilon_{\rm grad}, \varepsilon_{\rm osc}, \varepsilon_{\rm curv})$ and scan window $\mathcal{R}$, the geometric plateau set $\mathfrak{P}_\alpha(\varepsilon_{\rm grad}, \varepsilon_{\rm osc}, \varepsilon_{\rm curv}; \mathcal{R})$ of Section 7.3.5 is invariant under variations of $\alpha_{\rm gf}$ when evaluated on shell in $\mathcal{H}_{\rm phys}$.*

Proof. By the Nielsen master identity and its band–safe form (cf. Propositions 7.208 and 7.218), the channel visibility obeys the on–shell statement eq. (7.66). $\partial_{\alpha_{\mathrm{gf}}}\Lambda_\alpha = 0$ on $\mathcal{H}_{\mathrm{phys}}$, under the anomaly/Slavnov assumptions eqs. (7.16) and (7.18) and quartet/positivity control (Section 7.2.6 and eq. (7.12)). In the log chart $(u, v) = (\ln \lambda_{\mathrm{UV}}, \ln \ell)$, the operators $D_\lambda = \partial_u$ and $D_\ell = \partial_v$ differentiate only the band tags $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$, which are BRST-closed and *gauge-parameter independent* by eq. (7.30). Hence the derivatives commute: $\partial_{\alpha_{\mathrm{gf}}} D_\lambda \Lambda_\alpha = D_\lambda \partial_{\alpha_{\mathrm{gf}}} \Lambda_\alpha = 0$ and similarly for D_ℓ, and by repeated commutation also for second log–derivatives (entries of $\mathbf{H}^{\mathrm{plat}}\Lambda_\alpha$). Local oscillation on a fixed g_{plat}-ball is a functional of pointwise values of Λ_α in that neighbourhood; therefore it is also α_{gf}-independent on shell. Thus, each of the three geometric inequalities defining $\mathfrak{P}_\alpha$ (Definition 7.44) is invariant under α_{gf}; the set itself is unchanged. □

BRST robustness and regulator compatibility.

Proposition 7.64 (BRST robustness; regulator compatibility of certified plateaus)**.** *Work on* $\mathcal{H}_{\mathrm{phys}}$.

(i) BRST robustness.
If $\Lambda_\alpha \mapsto \Lambda_\alpha + \langle \zeta X \rangle$ *for some local* X *(ghost number* -1*), then all plateau inequalities remain satisfied with the same tolerances; in particular, membership in* $\mathfrak{P}_\alpha$ *and any joint* $\mathfrak{P}_{\mathfrak{C}}$ *is unchanged.*

(ii) Regulator compatibility.
Let $(P, K) \mapsto (P', K')$ *be an admissible regulator change (as in Section 7.3.3) with mismatch metrics* $(\eta_{\mathrm{UV}}, \eta_{\mathrm{IR}})$ *and first-log-derivative mismatches* $(\dot\eta_{\mathrm{UV}}, \dot\eta_{\mathrm{IR}})$*. If the induced bounds (7.62)–(7.63) satisfy*

$$\sup_{\mathcal{R}} \left| D_\lambda \Lambda'_\alpha - D_\lambda \Lambda_\alpha \right| \le \delta_\lambda, \qquad \sup_{\mathcal{R}} \left| D_\ell \Lambda'_\alpha - D_\ell \Lambda_\alpha \right| \le \delta_\ell, \qquad \sup_{\mathcal{R}} \left| \Lambda'_\alpha - \Lambda_\alpha \right| \le \varepsilon_{\mathrm{reg}},$$

with $\max\{\delta_\lambda, \delta_\ell, \varepsilon_{\mathrm{reg}}\} \ll \min\{\varepsilon_{\mathrm{grad}}, \varepsilon_{\mathrm{osc}}\}$ *and the analogous second-derivative bound controlled by* $\varepsilon_{\mathrm{curv}}$*, then the certified plateau set is regulator invariant:* $\mathfrak{P}_\alpha(P, K) = \mathfrak{P}_\alpha(P', K')$ *on* $\mathcal{R}$*.*

Proof. (i) On $\mathcal{H}_{\mathrm{phys}}$ any ζ-exact insertion has vanishing expectation value by the quartet mechanism and positivity (cf. Section 7.2.6 and Corollary 7.24); therefore $\Lambda_\alpha \mapsto \Lambda_\alpha + \langle \zeta X \rangle$ leaves pointwise values, log–derivatives (commuting with ζ on band tags), and local oscillations unchanged. Each inequality in the geometric/channelwise definitions (Definition 7.44 and eq. (7.56)) is preserved.

(ii) The stated bounds are exactly those in (7.62)–(7.63) (with the second-derivative variant controlled by the same calculus), which ensure that gradients, oscillations, and Hessian operator norms remain within the original tolerances for every grid point and neighbourhood used in the test. Hence the truth values of all plateau inequalities are preserved, and membership coincides across regulators. See also Proposition 7.54 for the corresponding global statement used in certification. □

Remark 7.65 (Gauge-parameter invariance and regulator robustness)**.** Gauge-parameter invariance on shell descends from the Nielsen framework (Section 7.3.6) and relies on eqs. (7.16)

and (7.18) together with BRST closure of the IR/UV tags eq. (7.30) [114, 91, 82]. Regulator robustness is the operational counterpart: admissible projector/window variations only induce drifts bounded by the regulator budget fixed in Section 7.3.3, which is chosen strictly below the plateau tolerances so as not to alter membership.

7.3.7 Regularisation Stability (Kernel/Projector Families)

Let $\{K_{\mathrm{IR}}^{(\ell)}\}$ and $\{\tilde{K}_{\mathrm{IR}}^{(\ell)}\}$ be two IR families satisfying eq. (7.32), and let $P_x, \tilde{P}_x$ be two UV projector families built from generators $\mathcal{D}_x, \tilde{\mathcal{D}}_x$ with the same low-energy fixed-point structure. Denote the corresponding visibilities by Λ_α and $\tilde{\Lambda}_\alpha$. *Regularisation stability* on $\mathcal{R}$ holds if

$$\sup_{(\lambda_{\mathrm{UV}},\ell)\in\mathcal{R}} \left|\Lambda_\alpha(\lambda_{\mathrm{UV}},\ell) - \tilde{\Lambda}_\alpha(\lambda_{\mathrm{UV}},\ell)\right| \le \varepsilon_{\mathrm{reg}}, \qquad \varepsilon_{\mathrm{reg}} \ll \min\{\varepsilon_{\mathrm{osc}}, \varepsilon_{\mathrm{grad}}\}, \tag{7.67}$$

in which case both prescriptions yield the same plateau certification on $\mathcal{R}$ (cf. Section 7.3.3). We now give sufficient conditions for (7.67).

Definition 7.66 (Admissible IR/UV variations)**.** Assume:

(A1) IR closeness.
For each ℓ in the scan range,

$$\|K_{\mathrm{IR}}^{(\ell)} - \tilde{K}_{\mathrm{IR}}^{(\ell)}\|_{L^1(\mathbb{R}^d)} \;\le\; \delta_{\mathrm{IR}}(\ell), \qquad \sup_{\ell\in[\ell_{\min},\ell_{\max}]} \delta_{\mathrm{IR}}(\ell) \le \delta_{\mathrm{IR}}. \tag{7.68}$$

(A2) UV closeness with band margin.
For each x there exists a spectral gap margin

$$\Delta_{\mathrm{band}}(x,\lambda_{\max}) > 0$$

such that

$$\mathrm{dist}\Big(\lambda_{\max}, \mathrm{Spec}(\mathcal{D}_x)\Big) \ge \Delta_{\mathrm{band}}$$

and

$$\mathrm{dist}\Big(\lambda_{\max}, \mathrm{Spec}(\tilde{\mathcal{D}}_x)\Big) \ge \Delta_{\mathrm{band}}$$

and

$$\|\mathcal{D}_x - \tilde{\mathcal{D}}_x\|_{\mathrm{op}} \;\le\; \delta_{\mathrm{UV}} \quad\Rightarrow\quad \sup_{0<\lambda\le\lambda_{\max}} \|P_x(\lambda) - \tilde{P}_x(\lambda)\|_{\mathrm{op}} \;\le\; c_{\mathrm{band}} \frac{\delta_{\mathrm{UV}}}{\Delta_{\mathrm{band}}}, \tag{7.69}$$

for some c_{band} independent of (λ,ℓ).

Lemma 7.67 (IR perturbation bound (Young))**.** *Let $\mathcal{O}:\mathbb{R}^d \to \mathcal{B}(\mathcal{H})$ be an operator–valued density with essentially bounded norm, $\|\mathcal{O}\|_{L^\infty} := \operatorname{ess\,sup}_x \|\mathcal{O}(x)\| < \infty$. For IR windows $K_{\mathrm{IR}}^{(\ell)}, \tilde{K}_{\mathrm{IR}}^{(\ell)} \in L^1(\mathbb{R}^d)$ as in Definition 6.280, one has, for any $\ell > 0$,*

$$\left\| (K_{\mathrm{IR}}^{(\ell)} - \tilde{K}_{\mathrm{IR}}^{(\ell)}) * \mathcal{O} \right\|_{L^\infty} \;\le\; \left\|K_{\mathrm{IR}}^{(\ell)} - \tilde{K}_{\mathrm{IR}}^{(\ell)}\right\|_{L^1} \|\mathcal{O}\|_{L^\infty}.$$

Proof. Set $K_\Delta := K_{\mathrm{IR}}^{(\ell)} - \tilde{K}_{\mathrm{IR}}^{(\ell)} \in L^1(\mathbb{R}^d)$. For a.e. x the convolution is the Bochner integral (in the operator–norm Banach space)

$$(K_\Delta * \mathcal{O})(x) = \int_{\mathbb{R}^d} K_\Delta(y)\, \mathcal{O}(x-y)\, dy,$$

which is well-defined because $\int |K_\Delta(y)|\, \|\mathcal{O}(x-y)\|\, dy \le \|K_\Delta\|_{L^1}\, \|\mathcal{O}\|_{L^\infty} < \infty$. By the triangle inequality for the Bochner integral and submultiplicativity of the operator norm,

$$\big\|(K_\Delta * \mathcal{O})(x)\big\| \le \int_{\mathbb{R}^d} |K_\Delta(y)|\, \|\mathcal{O}(x-y)\|\, dy \le \|\mathcal{O}\|_{L^\infty} \int_{\mathbb{R}^d} |K_\Delta(y)|\, dy = \|K_\Delta\|_{L^1}\, \|\mathcal{O}\|_{L^\infty}.$$

Taking the essential supremum over x yields the claim: $\|(K_\Delta * \mathcal{O})\|_{L^\infty} \le \|K_\Delta\|_{L^1}\, \|\mathcal{O}\|_{L^\infty}$. This is the standard Young convolution bound $L^1 * L^\infty \to L^\infty$, and its proof applies verbatim to Banach–valued (here operator–valued) functions via Bochner integration. For background on convolution–norm inequalities and Bochner integration, see [55, 30]. □

Lemma 7.68 (UV compression perturbation bound)**.** *Let A be a bounded operator on a fibre and let $P, \tilde{P}$ be orthogonal projectors (as in Definition 7.26, used in the joint tagging of Definition 7.28). Then*

$$\| PAP - \tilde{P}\, A\, \tilde{P} \| \;\le\; 2\, \|A\|\, \|P - \tilde{P}\|.$$

Proof. Use the algebraic decomposition

$$PAP - \tilde{P}\, A\, \tilde{P} = (P - \tilde{P})\, A\, P \;+\; \tilde{P}\, A\, (P - \tilde{P}).$$

By the triangle inequality and submultiplicativity of the operator norm,

$$\|PAP - \tilde{P}\, A\, \tilde{P}\| \;\le\; \|(P-\tilde{P})AP\| + \|\tilde{P}A(P-\tilde{P})\| \;\le\; \|P-\tilde{P}\|\, \|A\|\, \|P\| + \|\tilde{P}\|\, \|A\|\, \|P-\tilde{P}\|.$$

Since $P, \tilde{P}$ are orthogonal projectors, they are self-adjoint idempotents with spectrum $\{0, 1\}$, hence $\|P\| = \|\tilde{P}\| = 1$. The bound follows:

$$\|PAP - \tilde{P}\, A\, \tilde{P}\| \;\le\; 2\, \|A\|\, \|P - \tilde{P}\|.$$

□

Remark. Applied to the UV-compression step in the joint IR/UV map (Definition 7.28) with $A = \mathcal{O}^{(\ell)}$, this controls the sensitivity to projector mismatches; combined with the IR convolution bound of Lemma 7.67 and the band functor properties in Proposition 7.30, it yields stability estimates for the full band map.

Definition 7.69 (Lipschitz sensitivities of a visibility)**.** For a fixed channel α, let $L_\alpha^{(K)}$ and $L_\alpha^{(P)}$ be non-negative constants such that

$$\begin{aligned} \big|\Lambda_\alpha(\ldots, K; \ldots) - \Lambda_\alpha(\ldots, \tilde{K}; \ldots)\big| &\le L_\alpha^{(K)}\, \|K - \tilde{K}\|_{L^1}, \\ \big|\Lambda_\alpha(P, \ldots; \ldots) - \Lambda_\alpha(\tilde{P}, \ldots; \ldots)\big| &\le L_\alpha^{(P)}\, \|P - \tilde{P}\|, \end{aligned}$$

whenever the other tagged inputs are held fixed. (These arise from the linearity/complete positivity of the band functor Definition 7.29 and the regularity of $\mathcal{F}_\alpha$ in eq. (7.54).)

Proposition 7.70 (Sufficient conditions for (7.67))**.** *Under Definition 7.32 and Definition 7.69, there exists a constant*

$$C_\alpha \;:=\; L_\alpha^{(K)} \;+\; 2\,L_\alpha^{(P)}\,c_{\mathrm{band}}$$

such that, uniformly on $\mathcal{R}$*,*

$$\left|\Lambda_\alpha - \tilde{\Lambda}_\alpha\right| \;\le\; C_\alpha\left(\delta_{\mathrm{IR}} + \frac{\delta_{\mathrm{UV}}}{\Delta_{\mathrm{band}}}\right) \;=:\; \varepsilon_{\mathrm{reg}}. \tag{7.70}$$

In particular, if $\varepsilon_{\mathrm{reg}} \ll \min\{\varepsilon_{\mathrm{osc}}, \varepsilon_{\mathrm{grad}}\}$ *then both prescriptions yield the same plateau certification on* $\mathcal{R}$*.*

Proof. Write each tagged object as "UV then IR" (Definition 7.28): $K_{\mathrm{IR}}^{(\ell)}\big(P\,\mathcal{O}\,P\big)$ versus $\tilde{K}_{\mathrm{IR}}^{(\ell)}\big(\tilde{P}\,\mathcal{O}\,\tilde{P}\big)$. Add and subtract $K_{\mathrm{IR}}^{(\ell)}\big(\tilde{P}\,\mathcal{O}\,\tilde{P}\big)$ and use Lemma 7.68 for the UV change and Lemma 7.67 (with $\mathcal{O}$ the bounded operator–valued density entering $\mathcal{F}_\alpha$) for the IR change. Applying the channel Lipschitz estimates of Definition 7.69 gives

$$|\Lambda_\alpha - \tilde{\Lambda}_\alpha| \le L_\alpha^{(P)}\,\|P - \tilde{P}\| + L_\alpha^{(K)}\,\|K_{\mathrm{IR}}^{(\ell)} - \tilde{K}_{\mathrm{IR}}^{(\ell)}\|_{L^1}.$$

Invoke (7.69) to bound $\|P - \tilde{P}\| \le c_{\mathrm{band}}\,\delta_{\mathrm{UV}}/\Delta_{\mathrm{band}}$, and (7.68) for the kernel term, yielding (7.70). If $\varepsilon_{\mathrm{reg}}$ is strictly below the plateau tolerances, the inequalities that define the plateau sets (gradients/oscillation/curvature) are unaffected (cf. the regulator-robustness arguments in Section 7.3.3). □

Corollary 7.71 (Equality of certified plateaus across admissible regulators)**.** *If* (P, K) *and* $(\tilde{P}, \tilde{K})$ *satisfy Definition 7.32 and the regulator–stability bounds of* (7.70) *hold with*

$$\varepsilon_{\mathrm{reg}} \;\le\; \tfrac{1}{2}\,\min\{\varepsilon_{\mathrm{osc}},\,\varepsilon_{\mathrm{grad}}\},$$

then $\mathfrak{P}_\alpha(P, K) = \mathfrak{P}_\alpha(\tilde{P}, \tilde{K})$ *on* $\mathcal{R}$*. Consequently, the joint certified plateau is identical for the two prescriptions:*

$$\mathfrak{P}_{\mathfrak{C}}(P, K) \;=\; \bigcap_{\alpha\in\mathfrak{C}} \mathfrak{P}_\alpha(P, K) \;=\; \bigcap_{\alpha\in\mathfrak{C}} \mathfrak{P}_\alpha(\tilde{P}, \tilde{K}) \;=\; \mathfrak{P}_{\mathfrak{C}}(\tilde{P}, \tilde{K}).$$

Proof idea. The sup–norm control (7.70) keeps visibility values within half of each threshold, so both sides of each plateau inequality in (7.56) have the same truth value at every grid point and in every g_{plat}-neighbourhood; joint intersections preserve equality.

Proof. **Step 1: Uniform control of regulator-induced deviations.** By admissibility (Definition 7.32) and the perturbation bounds for the IR window and UV projector (Lemmas 7.67 and 7.68), the change $(P, K) \mapsto (\tilde{P}, \tilde{K})$ induces the stability estimates summarized in (7.70):

$$\sup_{\mathcal{R}}\left|\Lambda'_\alpha - \Lambda_\alpha\right| \le \varepsilon_{\mathrm{reg}}, \quad \sup_{\mathcal{R}}\left|D_\lambda\Lambda'_\alpha - D_\lambda\Lambda_\alpha\right| \le \varepsilon_{\mathrm{reg}}, \quad \sup_{\mathcal{R}}\left|D_\ell\Lambda'_\alpha - D_\ell\Lambda_\alpha\right| \le \varepsilon_{\mathrm{reg}},$$

and likewise for the oscillation/width metric that enters the plateau test (cf. (7.49)):

$$\sup_{\mathcal{R}}\left|\, g_{\mathrm{osc}}(\Lambda'_\alpha) - g_{\mathrm{osc}}(\Lambda_\alpha)\,\right| \;\le\; \varepsilon_{\mathrm{reg}}.$$

Here $'$ denotes quantities evaluated with $(\tilde{P}, \tilde{K})$.

Step 2: Invoke regulator-compatibility criterion. Set the first-derivative mismatch bounds in Proposition 7.64 to

$$\delta_\lambda = \delta_\ell = \varepsilon_{\mathrm{reg}}, \qquad \varepsilon_{\mathrm{reg}}^{(\mathrm{value})} = \varepsilon_{\mathrm{reg}}^{(\mathrm{osc})} = \varepsilon_{\mathrm{reg}},$$

so the hypotheses of Proposition 7.64 are met provided

$$\max\{\delta_\lambda, \delta_\ell, \varepsilon_{\mathrm{reg}}\} \;\ll\; \min\{\varepsilon_{\mathrm{grad}}, \varepsilon_{\mathrm{osc}}\}.$$

The quantitative choice in the statement, $\varepsilon_{\mathrm{reg}} \le \frac{1}{2}\min\{\varepsilon_{\mathrm{osc}}, \varepsilon_{\mathrm{grad}}\}$, is a concrete sufficient condition for this "$\ll$" requirement. Hence the certified plateau set is regulator invariant:

$$\mathfrak{P}_\alpha(P, K) = \mathfrak{P}_\alpha(\tilde{P}, \tilde{K}) \quad \text{on } \mathcal{R}.$$

Step 3: Equality for the joint certified plateau. By definition, $\mathfrak{P}_{\mathfrak{C}}$ is a finite intersection of the per–channel certified plateaus (see (7.56) and the monotonicity/closedness properties in Proposition 7.38). Since equality holds channel-wise for every $\alpha \in \mathfrak{C}$, it is preserved under intersection:

$$\mathfrak{P}_{\mathfrak{C}}(P, K) = \bigcap_{\alpha\in\mathfrak{C}} \mathfrak{P}_\alpha(P, K) = \bigcap_{\alpha\in\mathfrak{C}} \mathfrak{P}_\alpha(\tilde{P}, \tilde{K}) = \mathfrak{P}_{\mathfrak{C}}(\tilde{P}, \tilde{K}).$$

Step 4: Boundary cases. Plateau sets are closed in $\mathcal{R}$ (Proposition 7.38). Thus any potential boundary points saturating the inequalities remain included under both prescriptions; the $\frac{1}{2}$–margin ensures that no interior point crosses the acceptance thresholds when switching regulators, so set equality is exact.

This proves the claim. □

Optional strengthening (derivative stability). If, in addition, the first log–derivative mismatches of the tags are bounded uniformly on $\mathcal{R}$,

$$\dot{\eta}_{\mathrm{UV}} := \sup \left\| \lambda\, \partial_\lambda P - \lambda\, \partial_\lambda \tilde{P} \right\|, \qquad \dot{\eta}_{\mathrm{IR}} := \sup \left\| \ell\, \partial_\ell K_{\mathrm{IR}}^{(\ell)} - \ell\, \partial_\ell \tilde{K}_{\mathrm{IR}}^{(\ell)} \right\|_{L^1},$$

then the same argument with "UV then IR" calculus (Definition 7.28) yields gradient-level bounds

$$\sup_{\mathcal{R}} \left| D_\lambda \Lambda_\alpha - D_\lambda \tilde{\Lambda}_\alpha \right| \;\le\; L_\alpha^{(P)}\, \dot{\eta}_{\mathrm{UV}} + L_\alpha^{(K)}\, \delta_{\mathrm{IR}}, \qquad \sup_{\mathcal{R}} \left| D_\ell \Lambda_\alpha - D_\ell \tilde{\Lambda}_\alpha \right| \;\le\; L_\alpha^{(P)}\, \delta_{\mathrm{UV}} + L_\alpha^{(K)}\, \dot{\eta}_{\mathrm{IR}},$$

and analogous second–derivative bounds, thereby directly ensuring equality of plateau sets even when certification thresholds are specified in terms of derivatives (cf. Proposition 7.54).

Background-field compatibility. The UV closeness hypothesis (7.69) is the standard stability of spectral projectors under gapped perturbations and is compatible with background-field methods [114, 91]; IR bounds are purely measure-theoretic (Young) and insensitive to gauge choices.

Lipschitz continuity of the visibility functional.

Lemma 7.72 (Lipschitz continuity of $\mathcal{F}_\alpha$ on tagged inputs)**.** *Let $\mathcal{F}_\alpha$ be a holosymmetric channel functional as in eq. (7.54), built from a finite family of band-tagged objects of the form $K_{\mathrm{IR}}^{(\ell)}\big(P\,\mathcal{O}\,P\big)$ (and their finite products/contractions) evaluated on $\mathcal{H}_{\mathrm{phys}}$. Assume the band functor properties of Definition 7.29 and Proposition 7.30 and the BRST/positivity framework of Section 7.2.6 and eq. (7.12). Then, on any compact rectangle $\mathcal{R} \subset \mathbb{R}^2_{>0}$, there exists a constant $L_\alpha(\mathcal{R}) > 0$ such that, for any two tag pairs (P, K) and $(\tilde{P}, \tilde{K})$,*

$$\big|\mathcal{F}_\alpha[P,K] - \mathcal{F}_\alpha[\tilde{P},\tilde{K}]\big| \;\le\; L_\alpha(\mathcal{R})\Big(\sup_{x,\lambda\le\lambda_{\max}} \|P_x(\lambda) - \tilde{P}_x(\lambda)\|_{\mathrm{op}} \;+\; \sup_{\ell\in[\ell_{\min},\ell_{\max}]} \|K^{(\ell)} - \tilde{K}^{(\ell)}\|_{L^1}\Big). \tag{7.71}$$

Proof. Each tagged input enters $\mathcal{F}_\alpha$ through "UV then IR" compression/averaging (Definition 7.28), and $\mathcal{F}_\alpha$ is a finite linear combination of products/traces of such inputs on $\mathcal{H}_{\mathrm{phys}}$. By Proposition 7.30 the map $\mathcal{O} \mapsto K_{\mathrm{IR}}^{(\ell)}(P\mathcal{O}P)$ is linear, *-preserving, completely positive, and norm–nonexpansive up to $\|K_{\mathrm{IR}}^{(\ell)}\|_{L^1} = 1$. For a single factor we estimate the IR and UV variations separately:

$$\big\|K_{\mathrm{IR}}^{(\ell)}(P\mathcal{O}P) - \tilde{K}_{\mathrm{IR}}^{(\ell)}(\tilde{P}\mathcal{O}\tilde{P})\big\| \;\le\; \underbrace{\big\|K_{\mathrm{IR}}^{(\ell)}(P\mathcal{O}P) - K_{\mathrm{IR}}^{(\ell)}(\tilde{P}\mathcal{O}\tilde{P})\big\|}_{\text{UV change}} + \underbrace{\big\|(K_{\mathrm{IR}}^{(\ell)} - \tilde{K}_{\mathrm{IR}}^{(\ell)}) * (\,\tilde{P}\mathcal{O}\tilde{P}\,)\big\|}_{\text{IR change}}.$$

The UV change is controlled by Lemma 7.68: $\|P\mathcal{O}P - \tilde{P}\mathcal{O}\tilde{P}\| \le 2\,\|\mathcal{O}\|\,\|P - \tilde{P}\|$. The IR change is controlled by Young's inequality Lemma 7.67: $\|(K - \tilde{K}) * (\tilde{P}\mathcal{O}\tilde{P})\| \le \|K - \tilde{K}\|_{L^1}\,\|\mathcal{O}\|$. By multilinearity (finite products) and norm continuity of traces/contractions on $\mathcal{H}_{\mathrm{phys}}$ (guaranteed by ghost decoupling and positivity, cf. Section 7.2.6 and eq. (7.12); see also [82, 241]), the total variation of $\mathcal{F}_\alpha$ is bounded by a constant multiple of the sum of the factor-wise bounds. Collecting these constants over the finite number of factors and using compactness of $\mathcal{R}$ yields (7.71) with some $L_\alpha(\mathcal{R})$. □

Proposition 7.73 (Sufficient condition for regularisation stability)**.** *Under* (A1)–(A2) *(i.e. eqs. (7.68) and (7.69)) and (7.71),*

$$\sup_{(\lambda_{\mathrm{UV}},\ell)\in\mathcal{R}} \big|\Lambda_\alpha(\lambda_{\mathrm{UV}},\ell) - \tilde{\Lambda}_\alpha(\lambda_{\mathrm{UV}},\ell)\big| \;\le\; L_\alpha(\mathcal{R})\Big(c_{\mathrm{band}}\tfrac{\delta_{\mathrm{UV}}}{\Delta_{\mathrm{band}}} + \delta_{\mathrm{IR}}\Big). \tag{7.72}$$

Hence (7.67) holds whenever $L_\alpha(\mathcal{R})\big(c_{\mathrm{band}}\delta_{\mathrm{UV}}/\Delta_{\mathrm{band}} + \delta_{\mathrm{IR}}\big) \le \varepsilon_{\mathrm{reg}}$.

Proof idea. Apply the Lipschitz estimate (7.71) with (P, K) and $(\tilde{P}, \tilde{K})$. The IR term is bounded by δ_{IR} uniformly on $\mathcal{R}$ by eq. (7.68). For the UV term, eq. (7.69) gives $\sup_{\lambda\le\lambda_{\max}} \|P_x(\lambda) - \tilde{P}_x(\lambda)\|_{\mathrm{op}} \le c_{\mathrm{band}}\,\delta_{\mathrm{UV}}/\Delta_{\mathrm{band}}$, uniformly in x. Substitute these bounds into (7.71) to obtain (7.72). The final sentence follows directly from the definition (7.67).

Proof. **Step 1: Triangular decomposition via a mixed regulator.** Fix $(\lambda_{\mathrm{UV}}, \ell) \in \mathcal{R}$. Insert the intermediate pair $(\tilde{P}, K)$ and use the triangle inequality:

$$\big|\Lambda_\alpha(P,K) - \Lambda_\alpha(\tilde{P},\tilde{K})\big| \;\le\; \underbrace{\big|\Lambda_\alpha(P,K) - \Lambda_\alpha(\tilde{P},K)\big|}_{\text{UV variation}} \;+\; \underbrace{\big|\Lambda_\alpha(\tilde{P},K) - \Lambda_\alpha(\tilde{P},\tilde{K})\big|}_{\text{IR variation}}.$$

Step 2: Apply the visibility Lipschitz bound. By eq. (7.71), Λ_α is Lipschitz in its UV/IR arguments uniformly on $\mathcal{R}$ with constant $L_\alpha(\mathcal{R})$. Hence

$$\big|\Lambda_\alpha(P,K)-\Lambda_\alpha(\tilde P,K)\big| \le L_\alpha(\mathcal{R}) \sup_{\lambda\le\lambda_{\max}} \big\|P_x(\lambda)-\tilde P_x(\lambda)\big\|_{\mathrm{op}},$$
$$\big|\Lambda_\alpha(\tilde P,K)-\Lambda_\alpha(\tilde P,\tilde K)\big| \le L_\alpha(\mathcal{R}) \big\|K_{\mathrm{IR}}^{(\ell)}-\tilde K_{\mathrm{IR}}^{(\ell)}\big\|_{L^1}.$$

Step 3: Insert UV/IR closeness assumptions. By the IR closeness hypothesis eq. (7.68),

$$\big\|K_{\mathrm{IR}}^{(\ell)}-\tilde K_{\mathrm{IR}}^{(\ell)}\big\|_{L^1} \le \delta_{\mathrm{IR}} \quad \text{uniformly on } \mathcal{R}.$$

By the UV closeness hypothesis eq. (7.69),

$$\sup_{\lambda\le\lambda_{\max}} \big\|P_x(\lambda)-\tilde P_x(\lambda)\big\|_{\mathrm{op}} \le c_{\mathrm{band}}\frac{\delta_{\mathrm{UV}}}{\Delta_{\mathrm{band}}} \quad \text{uniformly in } x \text{ and on } \mathcal{R}.$$

Step 4: Combine the bounds and take the supremum on $\mathcal{R}$. Substituting these estimates into the inequalities from Step 2 and then taking $\sup_{(\lambda_{\mathrm{UV}},\ell)\in\mathcal{R}}$ yields

$$\sup_{\mathcal{R}}\big|\Lambda_\alpha-\tilde\Lambda_\alpha\big| \le L_\alpha(\mathcal{R})\Big(c_{\mathrm{band}}\tfrac{\delta_{\mathrm{UV}}}{\Delta_{\mathrm{band}}}+\delta_{\mathrm{IR}}\Big),$$

which is precisely eq. (7.72).

Step 5: Regularisation stability threshold. By the definition of regularisation stability eq. (7.67), the condition $L_\alpha(\mathcal{R})\big(c_{\mathrm{band}}\delta_{\mathrm{UV}}/\Delta_{\mathrm{band}}+\delta_{\mathrm{IR}}\big) \le \varepsilon_{\mathrm{reg}}$ guarantees that the regulator–induced change of Λ_α is below the tolerance $\varepsilon_{\mathrm{reg}}$ everywhere on $\mathcal{R}$, concluding the proof. □

Corollary 7.74 (Plateau equivalence under regulator changes)**.** *Assume the regularisation bound (7.72) holds on the compact scan window $\mathcal{R}$ with $\varepsilon_{\mathrm{reg}} \ll \min\{\varepsilon_{\mathrm{osc}},\varepsilon_{\mathrm{grad}}\}$. Then the plateau sets computed with (P,K) and $(\tilde P,\tilde K)$ coincide on $\mathcal{R}$, up to possible boundary pixels where inequalities are saturated (a set of zero 2D Lebesgue measure in the (u,v) chart). In particular, maximal components and joint plateaus eq. (7.64) are preserved.*

Proof. Let Λ_α and $\tilde\Lambda_\alpha$ be the visibilities obtained with (P,K) and $(\tilde P,\tilde K)$, respectively. By (7.72), $\|\Lambda_\alpha-\tilde\Lambda_\alpha\|_{L^\infty(\mathcal{R})} \le \varepsilon_{\mathrm{reg}}$. *Oscillation:* For any g_{plat}-ball B_r,

$$\big|\operatorname{osc}_{B_r}\Lambda_\alpha-\operatorname{osc}_{B_r}\tilde\Lambda_\alpha\big| \le 2\,\varepsilon_{\mathrm{reg}},$$

so if $\operatorname{osc}_{B_r}\Lambda_\alpha \le \varepsilon_{\mathrm{osc}}-2\varepsilon_{\mathrm{reg}}$ then $\operatorname{osc}_{B_r}\tilde\Lambda_\alpha \le \varepsilon_{\mathrm{osc}}$. *Gradients/Hessians:* There are two standard routes:

(a) Continuum (derivative stability): If one also assumes the first/second log–derivative mismatch bounds from the "optional strengthening (derivative stability)" (cf. the end of Section 7.3.7), then $|D_\lambda\Lambda_\alpha - D_\lambda\tilde\Lambda_\alpha| \le \delta_\lambda$ and $|D_\ell\Lambda_\alpha - D_\ell\tilde\Lambda_\alpha| \le \delta_\ell$, together with an analogous Hessian bound, with $\max\{\delta_\lambda,\delta_\ell\}$ and the Hessian bound $\ll \varepsilon_{\mathrm{grad}},\varepsilon_{\mathrm{curv}}$. Hence each gradient/curvature inequality that holds for Λ_α holds for $\tilde\Lambda_\alpha$ (and conversely).

(b) Discrete (finite-difference certification): If D_λ, D_ℓ and $\mathbf{H}^{\mathrm{plat}}$ are estimated by centred differences on a fixed grid with steps $(\Delta u, \Delta v)$ in the log chart, then for any interior point

$$\left|D_\lambda^{\mathrm{FD}}\Lambda_\alpha - D_\lambda^{\mathrm{FD}}\tilde{\Lambda}_\alpha\right| \le \frac{2\,\varepsilon_{\mathrm{reg}}}{\Delta u}, \qquad \left|D_\ell^{\mathrm{FD}}\Lambda_\alpha - D_\ell^{\mathrm{FD}}\tilde{\Lambda}_\alpha\right| \le \frac{2\,\varepsilon_{\mathrm{reg}}}{\Delta v},$$

and similarly each Hessian entry differs by at most a constant multiple of $\varepsilon_{\mathrm{reg}}/\min\{\Delta u, \Delta v\}^2$. Choosing $\varepsilon_{\mathrm{reg}}$ strictly below the certification thresholds (scaled by the grid steps) preserves all plateau inequalities pointwise, except possibly at pixels where the inequalities are saturated; these form a measure-zero frontier.

Therefore, the truth values of the three defining inequalities of the geometric/channelwise criteria (cf. Section 7.3.5 and eq. (7.56)) coincide for the two regulators almost everywhere in $\mathcal{R}$, hence the plateau sets agree up to boundary pixels. Since connectivity is preserved under the removal of a null frontier, maximal components and their intersections (7.64) are unchanged. □

Commuting limits and fixed-point structure.

Proposition 7.75 (Regulator-independence of commuting limits)**.** *Suppose (P, K) and $(\tilde{P}, \tilde{K})$ share the same low-energy fixed point and satisfy (7.72) with $\varepsilon_{\mathrm{reg}}$ uniform on $\mathcal{R}$. Then*

$$\lim_{\lambda\to\infty}\lim_{\ell\to 0^+}\left(\Lambda_\alpha - \tilde{\Lambda}_\alpha\right) = \lim_{\ell\to 0^+}\lim_{\lambda\to\infty}\left(\Lambda_\alpha - \tilde{\Lambda}_\alpha\right) = 0,$$

and the commuting-limits criterion eq. (7.45) is regulator-independent up to $O(\varepsilon_{\mathrm{reg}})$.

Proof. Uniformly on $\mathcal{R}$, $|\Lambda_\alpha - \tilde{\Lambda}_\alpha| \le \varepsilon_{\mathrm{reg}}$; taking either order of limits preserves the bound, hence each iterated limit of the difference exists and is $\le \varepsilon_{\mathrm{reg}}$. If the two constructions share the same low-energy fixed point, each single-regulator iterated limit exists and coincides; therefore the difference tends to 0. Finally, any difference between the swapped limits for one regulator and the other is bounded by $2\varepsilon_{\mathrm{reg}}$, which yields regulator-independence of eq. (7.45) up to $O(\varepsilon_{\mathrm{reg}})$. □

BRST compatibility.

Proposition 7.76 (BRST-closed evaluation and gauge-parameter independence)**.** *The tags $P_x(\lambda_{\mathrm{UV}}), \tilde{P}_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}, \tilde{K}_{\mathrm{IR}}^{(\ell)}$ are BRST-closed (ghost number 0) and commute with the channel projectors Π_α (cf. Definition 7.28, items (ii)–(iii), and eq. (7.145)). Consequently, all visibility bounds and regulator-comparison inequalities are computed from matrix elements on $\mathcal{H}_{\mathrm{phys}}$ (ghost/longitudinal sectors decouple; cf. Section 7.2.6 and Lemma 7.224), and on-shell gauge-parameter independence follows from the Nielsen framework (Section 7.2.9, Proposition 7.236, and Lemma 7.237; see also [114, 91]).*

Proof idea. BRST closure of the tags ensures ζ commutes with the tagging functor (UV then IR), so taking BRST cohomology and projecting to $\mathcal{H}_{\mathrm{phys}}$ can be done before or after tagging without affecting values; quartet decoupling removes unphysical sectors. Since Λ_α is built from gauge-invariant/BRST-closed composites on $\mathcal{H}_{\mathrm{phys}}$, the Nielsen identities imply $\partial_{\alpha_{\mathrm{gf}}}\Lambda_\alpha = 0$ on shell, uniformly in the choice of regulator satisfying the admissibility hypotheses; this leaves all the inequalities (and their limits) invariant.

Proof. **1) BRST closure and commutation with channels.** By Definition 7.28(ii) one has $\zeta P_x(\lambda_{\mathrm{UV}}) = 0$ and $\zeta K_{\mathrm{IR}}^{(\ell)} = 0$ with ghost number 0, and by Definition 7.28(iii) (channel/background compatibility) $[P_x(\lambda_{\mathrm{UV}}), \Pi_\alpha] = [K_{\mathrm{IR}}^{(\ell)}, \Pi_\alpha] = 0$. Hence for any composite $\mathcal{O}$ with $\zeta\mathcal{O} = 0$ (ghost number 0),

$$\zeta\left(K_{\mathrm{IR}}^{(\ell)} P\, \mathcal{O}\, P\right) = K_{\mathrm{IR}}^{(\ell)} P\, (\zeta\mathcal{O})\, P = 0,$$

so the tagging functor (UV then IR) preserves BRST cohomology classes and may be applied before or after projection to $\mathcal{H}_{\mathrm{phys}}$ without changing expectation values (Definition 7.28).

2) Decoupling of unphysical sectors. Quartet/doublet decoupling implies that ghost/longitudinal insertions are $\mathcal{B}_\Gamma$-exact and do not contribute to matrix elements in the physical sector (cf. Section 7.2.6 and Lemma 7.224). Since the tags commute with Π_α and have ghost number 0, restricting visibilities and their variations to $\mathcal{H}_{\mathrm{phys}}$ is consistent and leaves all norm bounds unchanged.

3) On-shell gauge-parameter independence (Nielsen). Let α_{gf} enter only through the gauge fermion $\Psi(\alpha_{\mathrm{gf}})$. The band Nielsen identity holds (Proposition 7.236):

$$\partial_{\alpha_{\mathrm{gf}}}\Gamma = -\,\mathcal{B}_\Gamma\big(\partial_{\alpha_{\mathrm{gf}}}\Psi\big) = \int d^d x\, \frac{\delta\Gamma}{\delta\varphi^A(x)}\, \mathcal{N}^A(x;\alpha_{\mathrm{gf}}).$$

Therefore, for any BRST-closed observable $\mathcal{O}$ (ghost number 0) built from tagged composites and evaluated on shell (solutions of the field equations), one has $\partial_{\alpha_{\mathrm{gf}}}\mathcal{O}\big|_{\text{on shell}} = 0$ (Lemma 7.237). The conclusion is uniform in the choice of admissible regulator because the tags are BRST-closed, commute with derivatives and channel projectors, and do not carry α_{gf} dependence (Definition 7.28); see also [114, 91] for the background-field/Nielsen perspective.

4) Consequence for the bounds. All inequalities used above (UV compression, IR convolution, Lipschitz continuity of visibilities, and regulator-stability bounds) depend only on operator norms of differences of tags and on channel-wise physical matrix elements. By Steps 1–2 these matrix elements may be evaluated in $\mathcal{H}_{\mathrm{phys}}$ without contamination from unphysical sectors, and by Step 3 they are independent of α_{gf} on shell. Hence the stated bounds and their certified-plateau conclusions are BRST-robust and gauge-parameter independent.

This proves the proposition. □

7.3.8 Channel-wise and Integrated Visibilities

Pointwise (channel) visibilities $\Lambda_\alpha(\lambda_{\mathrm{UV}}, \ell)$ are defined in Section 7.3.4. To average out microscopic fluctuations we introduce a scale–box average in the *linear* $(\lambda_{\mathrm{UV}}, \ell)$ chart:

Definition 7.77 (Integrated (scale–box averaged) visibility)**.** For fixed window widths $\Delta\lambda, \Delta\ell > 0$, define

$$\overline{\Lambda}_\alpha(\lambda_{\mathrm{UV}}, \ell; \Delta\lambda, \Delta\ell) := \frac{1}{\Delta\lambda\,\Delta\ell} \int_{\lambda_{\mathrm{UV}}-\Delta\lambda/2}^{\lambda_{\mathrm{UV}}+\Delta\lambda/2} d\lambda \int_{\ell-\Delta\ell/2}^{\ell+\Delta\ell/2} d\tilde{\ell}\; \Lambda_\alpha(\lambda, \tilde{\ell}). \tag{7.73}$$

Unless stated otherwise, $\Delta\lambda, \Delta\ell$ are constant (do not depend on the centre point).

Remark 7.78 (Log–box variant)**.** In the plateau log chart $(u, v) = (\ln \lambda_{\mathrm{UV}}, \ln \ell)$ with $D_\lambda = \partial_u$, $D_\ell = \partial_v$, one may instead average over a fixed rectangle $B^{\log} = [u - \Delta u/2, u + \Delta u/2] \times [v - \Delta v/2, v + \Delta v/2]$. Write the corresponding average as $\overline{\Lambda}_\alpha^{\log}(u, v; \Delta u, \Delta v)$. This variant is often preferable for derivative calculations (see Lemma 7.79).

Basic properties.

(P1) *Range and gauge/BRST robustness.* Since $-1 \le \Lambda_\alpha \le 1$, one has $-1 \le \overline{\Lambda}_\alpha \le 1$. Because Λ_α is computed on $\mathcal{H}_{\mathrm{phys}}$ from BRST-closed inputs and is Nielsen-protected (Section 7.2.9 and eq. (7.66)), the same holds for $\overline{\Lambda}_\alpha$.

(P2) *Derivative commutation (log–box).* For the log–box average of Remark 7.78 with fixed $\Delta u, \Delta v$,

$$D_\lambda \overline{\Lambda}_\alpha^{\log} = \overline{D_\lambda \Lambda_\alpha}^{\log}, \qquad D_\ell \overline{\Lambda}_\alpha^{\log} = \overline{D_\ell \Lambda_\alpha}^{\log}, \tag{7.74}$$

and likewise for mixed/second derivatives.

(P3) *Gradient/Hessian bounds (linear–box).* For the linear–box average (7.73),

$$\left| D_\lambda \overline{\Lambda}_\alpha \right| \le \overline{|D_\lambda \Lambda_\alpha|} \le \sup_B |D_\lambda \Lambda_\alpha|, \qquad \left\| \mathbf{H}^{\mathrm{plat}} \overline{\Lambda}_\alpha \right\|_{\mathrm{op}} \le \sup_B \left\| \mathbf{H}^{\mathrm{plat}} \Lambda_\alpha \right\|_{\mathrm{op}}, \tag{7.75}$$

where $B = [\lambda_{\mathrm{UV}} - \frac{\Delta\lambda}{2}, \lambda_{\mathrm{UV}} + \frac{\Delta\lambda}{2}] \times [\ell - \frac{\Delta\ell}{2}, \ell + \frac{\Delta\ell}{2}]$ is the averaging rectangle (the inequalities hold componentwise in the log chart via the chain rule).

(P4) *Oscillation reduction.* For any neighbourhood $\mathcal{N}$ (in either chart),

$$\mathrm{osc}_{\mathcal{N}} \left(\overline{\Lambda}_\alpha \right) \le \mathrm{osc}_{\mathcal{N} \oplus B} \left(\Lambda_\alpha \right), \tag{7.76}$$

with $\oplus$ the Minkowski sum and B the averaging box expressed in the same chart as $\mathcal{N}$.

Lemma 7.79 (Derivative commutation for log–box averages)**.** *Assume Λ_α is continuous on the chart $\mathcal{R} \subset \mathbb{R}^2$ and admits the scale gradients of eq. (7.55). For fixed $\Delta u, \Delta v > 0$, define the centred log–box average*

$$\overline{\Lambda}_\alpha^{\log}(u, v) := \frac{1}{\Delta u \, \Delta v} \int_{u - \frac{\Delta u}{2}}^{u + \frac{\Delta u}{2}} \int_{v - \frac{\Delta v}{2}}^{v + \frac{\Delta v}{2}} \Lambda_\alpha(u', v') \, dv' \, du'.$$

Then differentiation commutes with averaging:

$$\partial_u \overline{\Lambda}_\alpha^{\log}(u, v) = \frac{1}{\Delta u \, \Delta v} \int_{u - \frac{\Delta u}{2}}^{u + \frac{\Delta u}{2}} \int_{v - \frac{\Delta v}{2}}^{v + \frac{\Delta v}{2}} \partial_u \Lambda_\alpha(u', v') \, dv' \, du', \tag{7.77}$$

$$\partial_v \overline{\Lambda}_\alpha^{\log}(u, v) = \frac{1}{\Delta u \, \Delta v} \int_{u - \frac{\Delta u}{2}}^{u + \frac{\Delta u}{2}} \int_{v - \frac{\Delta v}{2}}^{v + \frac{\Delta v}{2}} \partial_v \Lambda_\alpha(u', v') \, dv' \, du', \tag{7.78}$$

and similarly for higher and mixed derivatives. Equivalently, in the log chart where $D_\lambda = \partial_u$ and $D_\ell = \partial_v$ (cf. Definition 7.28(vi)), one has

$$D_\lambda \overline{\Lambda}_\alpha^{\log} = \overline{D_\lambda \Lambda_\alpha}^{\log}, \qquad D_\ell \overline{\Lambda}_\alpha^{\log} = \overline{D_\ell \Lambda_\alpha}^{\log}. \tag{7.74}$$

Proof. **Step 1: Fix the integration domain by a linear change of variables.** Set

$$u' = u + \Delta u\,\sigma, \qquad v' = v + \Delta v\,\tau, \qquad (\sigma,\tau) \in \Big[-\tfrac{1}{2},\tfrac{1}{2}\Big]^2.$$

Since $\Delta u, \Delta v$ are fixed constants, the Jacobian is $\Delta u\,\Delta v$, and

$$\overline{\Lambda}^{\log}_{\alpha}(u,v) = \int_{-1/2}^{1/2}\!\!\int_{-1/2}^{1/2} \Lambda_\alpha\Big(u + \Delta u\,\sigma,\ v + \Delta v\,\tau\Big)\,d\tau\,d\sigma.$$

Step 2: Differentiate under the (now fixed) integral. By the hypothesis eq. (7.55), $\partial_u\Lambda_\alpha$ and $\partial_v\Lambda_\alpha$ exist (in the classical or weak sense) and are locally integrable on $\mathcal{R}$. Hence by the dominated convergence theorem (or Leibniz' rule for parameter-dependent integrals on a fixed domain),

$$\partial_u\,\overline{\Lambda}^{\log}_{\alpha}(u,v) = \int_{-1/2}^{1/2}\!\!\int_{-1/2}^{1/2} \partial_u\Big[\Lambda_\alpha\Big(u + \Delta u\,\sigma,\ v + \Delta v\,\tau\Big)\Big]\,d\tau\,d\sigma.$$

By the chain rule, ∂_u acts only through the first argument and $\partial_u\Lambda_\alpha(u+\Delta u\,\sigma, v+\Delta v\,\tau) = \partial_u\Lambda_\alpha(u',v')$ (since $\partial_u(u+\Delta u\,\sigma)=1$). Transforming back to (u',v') yields (7.77). The identity (7.78) is identical with $u \leftrightarrow v$, and mixed/higher derivatives follow by iterating the same argument.

Step 3: Relation to scale derivatives. In the log chart, $D_\lambda = \partial_u$ and $D_\ell = \partial_v$ (see Definition 7.28(vi)), so the statements above are exactly (7.74).

Remark on boundary terms. If one differentiates the original (u',v')-integral directly, the bounds depend on (u,v) and Leibniz' rule produces edge terms. The change of variables to (σ,τ) fixes the domain, and those edge contributions are absorbed into the chain-rule derivative under the integral, leading to the same result without explicit boundary terms.

This proves the lemma. □

Remark 7.80 (Jensen's inequality for box averages)**.** Let $J = I_\lambda \times I_\ell \subset \mathbb{R}^2$ be a rectangle of finite, nonzero measure $|J| := |I_\lambda|\,|I_\ell|$, and let

$$\mathcal{A}_J[f] \;:=\; \frac{1}{|J|}\int_J f(\lambda',\ell')\,d\lambda'\,d\ell'$$

denote the (normalized) box average of an integrable function $f : J \to \mathbb{R}$. For any convex $\phi : \mathbb{R} \to \mathbb{R}$ and integrable f,

$$\phi\Big(\mathcal{A}_J[f]\Big) \;\le\; \mathcal{A}_J[\phi\circ f].$$

In particular, with $\phi(t) = |t|$ one has $|\mathcal{A}_J[f]| \le \mathcal{A}_J[\,|f|\,]$. We use this with $f = \partial_{\lambda'}\Lambda_\alpha$ (or $f = \partial_{\ell'}\Lambda_\alpha$) to obtain the gradient bounds in Lemma 7.82. Background references: Jensen's inequality in the L^1 setting [30].

Remark 7.81 (Fubini–Tonelli on rectangles)**.** Let $J = I_\lambda \times I_\ell$ as above and $f : J \to \mathbb{R}$ be measurable. If $f \ge 0$ (Tonelli) or $f \in L^1(J)$ (Fubini), then the double integral equals either order of iterated integral and the sections are integrable a.e.:

$$\int_J f(\lambda',\ell')\,d\lambda'\,d\ell' \;=\; \int_{I_\lambda}\Big(\int_{I_\ell} f(\lambda',\ell')\,d\ell'\Big)d\lambda' \;=\; \int_{I_\ell}\Big(\int_{I_\lambda} f(\lambda',\ell')\,d\lambda'\Big)d\ell'.$$

We invoke this to pass between the box average and one-dimensional averages in the proof of Lemma 7.82 and in the derivation of (7.75). Classical statements and proofs: [55].

Lemma 7.82 (Bounds for linear–box averages)**.** *For the linear–box average* (7.73) *with fixed* $\Delta\lambda, \Delta\ell > 0$*, the inequalities* (7.75) *hold. Moreover, letting* $J = [\lambda - \frac{\Delta\lambda}{2}, \lambda + \frac{\Delta\lambda}{2}] \times [\ell - \frac{\Delta\ell}{2}, \ell + \frac{\Delta\ell}{2}]$ *be the averaging rectangle, the difference between* $D_\lambda \overline{\Lambda}_\alpha$ *and the linear–box average of* $D_\lambda \Lambda_\alpha$ *obeys*

$$\left| D_\lambda \overline{\Lambda}_\alpha - \overline{D_\lambda \Lambda_\alpha} \right| \;\le\; \frac{\Delta\lambda}{2\,\lambda_{\min}(J)} \sup_J \left| D_\lambda \Lambda_\alpha \right| \;+\; \frac{\Delta\ell}{2\,\ell_{\min}(J)} \sup_J \left| D_\ell \Lambda_\alpha \right|,$$

and analogously with $D_\lambda \leftrightarrow D_\ell$ *and for second derivatives.*

Proof idea. Inequalities (7.75) follow from Jensen and from passing to the log chart via $D_\lambda = \lambda\,\partial_\lambda$, $D_\ell = \ell\,\partial_\ell$, noting that λ, ℓ are bounded on compact rectangles. The bias bound is a direct application of the one-dimensional Leibniz rule in each variable together with the box-bias estimate of Lemma 7.40 (applied componentwise to $D_\lambda \Lambda_\alpha$ and $D_\ell \Lambda_\alpha$).

Proof. **Preliminaries and notation.** Recall the linear–box average

$$\overline{\Lambda}_\alpha(\lambda, \ell) := \frac{1}{\Delta\lambda\,\Delta\ell} \int_{\lambda - \frac{\Delta\lambda}{2}}^{\lambda + \frac{\Delta\lambda}{2}} \int_{\ell - \frac{\Delta\ell}{2}}^{\ell + \frac{\Delta\ell}{2}} \Lambda_\alpha(\lambda', \ell')\, d\ell'\, d\lambda',$$

and the scale derivatives $D_\lambda = \lambda\,\partial_\lambda$, $D_\ell = \ell\,\partial_\ell$. Let $\lambda_{\min}(J) := \inf\{\lambda' : (\lambda', \ell') \in J\}$ and $\ell_{\min}(J)$ analogously.

(i) Gradient bounds (7.75)**.** By Jensen's inequality for the convex map $t \mapsto |t|$ applied to the box average, and by the Fubini–Tonelli theorem on the rectangle J, [30, 55]

$$\left| D_\lambda \overline{\Lambda}_\alpha \right| = \left| \lambda\, \partial_\lambda \overline{\Lambda}_\alpha \right| \le \frac{\lambda}{\Delta\lambda\,\Delta\ell} \int_J \left| \partial_{\lambda'} \Lambda_\alpha \right| d\lambda'\, d\ell' \le \frac{\lambda}{\lambda_{\min}(J)}\, \overline{\left| D_\lambda \Lambda_\alpha \right|}.$$

Since $\lambda \le \lambda_{\max}(J)$ and $\lambda_{\max}(J)/\lambda_{\min}(J)$ is bounded on compact $\mathcal{R}$, this yields the stated bound; the D_ℓ case is identical. (Equivalently, pass to the log chart and use Lemma 7.79.)

(ii) Bias for D_λ: boundary decomposition. Differentiate the outer integral using Leibniz' rule (only limits depend on λ):

$$D_\lambda \overline{\Lambda}_\alpha(\lambda, \ell) = \frac{\lambda}{\Delta\lambda\,\Delta\ell} \int_{\ell - \frac{\Delta\ell}{2}}^{\ell + \frac{\Delta\ell}{2}} \Big[\Lambda_\alpha(\lambda_+, \ell') - \Lambda_\alpha(\lambda_-, \ell') \Big]\, d\ell',$$

with $\lambda_\pm := \lambda \pm \frac{\Delta\lambda}{2}$. On the other hand,

$$\overline{D_\lambda \Lambda_\alpha} = \frac{1}{\Delta\lambda\,\Delta\ell} \int_{\ell - \frac{\Delta\ell}{2}}^{\ell + \frac{\Delta\ell}{2}} \int_{\lambda_-}^{\lambda_+} \lambda'\, \partial_{\lambda'} \Lambda_\alpha(\lambda', \ell')\, d\lambda'\, d\ell'.$$

Integrate by parts in λ' inside the last integral:

$$\int_{\lambda_-}^{\lambda_+} \lambda'\, \partial_{\lambda'} \Lambda_\alpha = \Big[\lambda' \Lambda_\alpha \Big]_{\lambda_-}^{\lambda_+} - \int_{\lambda_-}^{\lambda_+} \Lambda_\alpha\, d\lambda'.$$

Subtracting, we obtain the exact identity

$$D_\lambda \overline{\Lambda}_\alpha - \overline{D_\lambda \Lambda_\alpha} = \frac{1}{\Delta\ell} \int_{\ell - \frac{\Delta\ell}{2}}^{\ell + \frac{\Delta\ell}{2}} h(\ell')\, d\ell', \tag{7.79}$$

$$h(\ell') := \frac{1}{\Delta\lambda} \int_{\lambda_-}^{\lambda_+} \Lambda_\alpha(\lambda', \ell')\, d\lambda' - \frac{\Lambda_\alpha(\lambda_+, \ell') + \Lambda_\alpha(\lambda_-, \ell')}{2}.$$

(iii) Trapezoid (box–bias) control in λ. For each fixed ℓ', apply the one–dimensional box–bias estimate (trapezoid error; cf. Lemma 7.40) to the C^1 function $\lambda' \mapsto \Lambda_\alpha(\lambda', \ell')$:

$$|h(\ell')| \;\le\; \frac{\Delta\lambda}{2} \sup_{\lambda' \in [\lambda_-, \lambda_+]} \left|\partial_{\lambda'} \Lambda_\alpha(\lambda', \ell')\right| \;\le\; \frac{\Delta\lambda}{2\,\lambda_{\min}(J)} \sup_J \left|D_\lambda \Lambda_\alpha\right|.$$

Averaging over ℓ' in (7.79) preserves this bound and yields the first term on the right-hand side of the lemma.

(iv) Averaging bias in ℓ. To capture the additional $\Delta\ell$ contribution, write

$$\frac{1}{\Delta\ell} \int_{\ell - \frac{\Delta\ell}{2}}^{\ell + \frac{\Delta\ell}{2}} h(\ell')\, d\ell' = h(\ell) \;+\; \Big[\, \overline{h} - h(\ell) \,\Big],$$

and bound the deviation of the ℓ–average from the centre value by the 1D box–bias in ℓ:

$$\left|\overline{h} - h(\ell)\right| \le \frac{\Delta\ell}{2} \sup_{\ell' \in [\ell_-, \ell_+]} \left|\partial_{\ell'} h(\ell')\right|.$$

Differentiate h under the integral defining it:

$$\partial_{\ell'} h(\ell') = \frac{1}{\Delta\lambda} \int_{\lambda_-}^{\lambda_+} \partial_{\ell'} \Lambda_\alpha(\lambda', \ell')\, d\lambda' - \frac{\partial_{\ell'} \Lambda_\alpha(\lambda_+, \ell') + \partial_{\ell'} \Lambda_\alpha(\lambda_-, \ell')}{2}.$$

Applying again the 1D box–bias (now in the λ' variable) to $\partial_{\ell'} \Lambda_\alpha$ gives

$$\left|\partial_{\ell'} h(\ell')\right| \le \sup_{\lambda' \in [\lambda_-, \lambda_+]} \left|\partial_{\ell'} \Lambda_\alpha(\lambda', \ell')\right| \le \frac{1}{\ell_{\min}(J)} \sup_J \left|D_\ell \Lambda_\alpha\right|.$$

Therefore

$$\left|\overline{h} - h(\ell)\right| \;\le\; \frac{\Delta\ell}{2\,\ell_{\min}(J)} \sup_J \left|D_\ell \Lambda_\alpha\right|.$$

Combining with the bound for $|h(\ell)|$ from Step (iii) establishes the stated estimate for $\left|D_\lambda \overline{\Lambda}_\alpha - \overline{D_\lambda \Lambda_\alpha}\right|$.

(v) The D_ℓ case and higher derivatives. The D_ℓ statement follows by symmetry interchanging (λ, D_λ) with (ℓ, D_ℓ). For second (and mixed) derivatives, iterate the same argument: differentiate (7.79) once more and apply the same box–bias controls componentwise to obtain bounds in terms of $\sup_J$ of the corresponding second (mixed) scale derivatives.

This completes the proof. □

Proposition 7.83 (Plateau tests for integrated visibilities)**.** *Let $\overline{\Lambda}_\alpha$ be either the log–box or linear–box average.*

*(i) (*Log–box*) The geometric/channelwise plateau inequalities may be imposed on $\overline{\Lambda}_\alpha^{\log}$ with the* same *tolerances, since (7.74) and (7.76) imply*

$$\|\nabla^{\mathrm{plat}} \overline{\Lambda}_\alpha^{\log}\|_\infty \le \sup_B \|\nabla^{\mathrm{plat}} \Lambda_\alpha\|_\infty,$$

$$\|\mathbf{H}^{\mathrm{plat}} \overline{\Lambda}_\alpha^{\log}\|_{\mathrm{op}} \le \sup_B \|\mathbf{H}^{\mathrm{plat}} \Lambda_\alpha\|_{\mathrm{op}},$$

$$\mathrm{osc} \;\; \textit{as in (7.76)}.$$

(i) (Linear–box) *The same plateau inequalities hold for* $\overline{\Lambda}_\alpha$ *with rescaled tolerances given in Corollary 7.41; in particular,*

$$\left|D_\lambda \overline{\Lambda}_\alpha\right| \le \varepsilon_\lambda + \tilde{\varepsilon}_\lambda, \qquad \left|D_\ell \overline{\Lambda}_\alpha\right| \le \varepsilon_\ell + \tilde{\varepsilon}_\ell,$$

with $\tilde{\varepsilon}_{\lambda,\ell}$ *as in Corollary 7.41, and the oscillation bound* (7.76).

Proof. (i) follows from Lemma 7.79 and (7.76). (ii) follows from Lemma 7.82 together with Lemma 7.40 and Corollary 7.41 for the bias/tolerance rescaling. □

Lemma 7.84 (Plateau inheritance: pointwise ⇒ integrated)**.** *Fix tolerances* $(\varepsilon_{\mathrm{grad}}, \varepsilon_{\mathrm{osc}}, \varepsilon_{\mathrm{curv}})$ *and the scan radius* $r > 0$ *from Definition 7.44. Let* B *be the averaging box (either linear– or log–box) centred at* $(\lambda_{\mathrm{UV}}, \ell)$ *(resp.* (u, v)*) and suppose* $B \subset B_r^{\mathrm{plat}}(\lambda_{\mathrm{UV}}, \ell)$*. Then the following hold:*

(a) (Uniform-curvature/oscillation inheritance, same tolerances) *Assume additionally that*

$$\sup_{B_r^{\mathrm{plat}}} \|\mathbf{H}^{\mathrm{plat}} \Lambda_\alpha\|_{\mathrm{op}} \le \varepsilon_{\mathrm{curv}}, \qquad \mathrm{osc}_{B_{2r}^{\mathrm{plat}}} \Lambda_\alpha \le \varepsilon_{\mathrm{osc}}.$$

If $(\lambda_{\mathrm{UV}}, \ell) \in \mathfrak{P}_\alpha$*, then the same point passes the geometric/channelwise plateau test with* Λ_α *replaced by the integrated visibility (log–box* $\overline{\Lambda}_\alpha^{\log}$ *or linear–box* $\overline{\Lambda}_\alpha$*)* with the same tolerance triple.

(b) (Radius-reduction inheritance, same tolerances) *Let* $\rho := \frac{1}{2}\,\mathrm{diam}_{g_{\mathrm{plat}}}(B)$ *and set* $r' := r - \rho > 0$*. If* $(\lambda_{\mathrm{UV}}, \ell) \in \mathfrak{P}_\alpha$*, then the same point passes the geometric/channelwise plateau test for the integrated visibility with the* same tolerances *but with oscillation measured on the smaller ball* $B_{r'}^{\mathrm{plat}}(\lambda_{\mathrm{UV}}, \ell)$*.*

(c) (General inheritance with explicit slack) *Without extra uniform assumptions, let* $\delta_{\mathrm{grad}} := \varepsilon_{\mathrm{grad}} - \|\nabla^{\mathrm{plat}} \Lambda_\alpha(\lambda_{\mathrm{UV}}, \ell)\|_\infty \ge 0$*. If the modulus of continuity of* $\mathbf{H}^{\mathrm{plat}} \Lambda_\alpha$ *on* B_r^{plat} *satisfies*

$$\sup_{B_r^{\mathrm{plat}}} \|\mathbf{H}^{\mathrm{plat}} \Lambda_\alpha\|_{\mathrm{op}} \le \varepsilon_{\mathrm{curv}} + \omega_H(r),$$

then, for the log–box average,

$$\left\|\nabla^{\mathrm{plat}} \overline{\Lambda}_\alpha^{\log}(\lambda_{\mathrm{UV}}, \ell)\right\|_\infty \le \varepsilon_{\mathrm{grad}} + \omega_H(r)\, r, \qquad \left\|\mathbf{H}^{\mathrm{plat}} \overline{\Lambda}_\alpha^{\log}(\lambda_{\mathrm{UV}}, \ell)\right\|_{\mathrm{op}} \le \varepsilon_{\mathrm{curv}} + \omega_H(r),$$

and for the linear–box average the same bounds hold up to the bias terms in Lemma 7.82. Consequently, if $\omega_H(r)\, r \le \delta_{\mathrm{grad}}$ *(and the linear–box biases are within the slack prescribed by Corollary 7.41), the point passes the test with* the same tolerances*.*

Proof. We work in the log chart $(u, v) = (\ln \lambda_{\mathrm{UV}}, \ln \ell)$ with g_{plat} (Definition 7.47); gradients and Hessians are those of Definition 7.44.

(a) Uniform-curvature/oscillation inheritance. For the *log–box* average, Lemma 7.79 gives $D_\lambda \overline{\Lambda}_\alpha^{\log} = \overline{D_\lambda \Lambda_\alpha}^{\log}$ and $D_\ell \overline{\Lambda}_\alpha^{\log} = \overline{D_\ell \Lambda_\alpha}^{\log}$. Hence

$$\left\|\nabla^{\mathrm{plat}} \overline{\Lambda}_\alpha^{\log}(\lambda_{\mathrm{UV}}, \ell)\right\|_\infty \le \sup_B \left\|\nabla^{\mathrm{plat}} \Lambda_\alpha\right\|_\infty.$$

For any $z \in B \subset B_r^{\mathrm{plat}}$, the mean-value estimate along the g_{plat} geodesic from the centre to z yields $\|\nabla^{\mathrm{plat}}\Lambda_\alpha(z)\|_\infty \le \|\nabla^{\mathrm{plat}}\Lambda_\alpha(\mathrm{centre})\|_\infty + \sup_{B_r^{\mathrm{plat}}} \|\mathbf{H}^{\mathrm{plat}}\Lambda_\alpha\|_{\mathrm{op}} \cdot d_{\mathrm{plat}}(z, \mathrm{centre}) \le \varepsilon_{\mathrm{grad}} + \varepsilon_{\mathrm{curv}}\, r$. Under the stated uniform-curvature hypothesis and because $(\lambda_{\mathrm{UV}}, \ell) \in \mathfrak{P}_\alpha$ (so $\|\nabla^{\mathrm{plat}}\Lambda_\alpha(\mathrm{centre})\|_\infty \le \varepsilon_{\mathrm{grad}}$), we get $\sup_B \|\nabla^{\mathrm{plat}}\Lambda_\alpha\|_\infty \le \varepsilon_{\mathrm{grad}}$ (taking r as the scan radius already fixed in the test). Thus $\|\nabla^{\mathrm{plat}}\overline{\Lambda}_\alpha^{\log}\|_\infty \le \varepsilon_{\mathrm{grad}}$. The Hessian bound at the centre follows from convexity of the operator norm under averaging and the uniform assumption: $\|\mathbf{H}^{\mathrm{plat}}\overline{\Lambda}_\alpha^{\log}(\mathrm{centre})\|_{\mathrm{op}} \le \sup_B \|\mathbf{H}^{\mathrm{plat}}\Lambda_\alpha\|_{\mathrm{op}} \le \varepsilon_{\mathrm{curv}}$. For the oscillation, eq. (7.76) with $\mathcal{N} = B_r^{\mathrm{plat}}$ gives $\mathrm{osc}_{B_r^{\mathrm{plat}}}(\overline{\Lambda}_\alpha^{\log}) \le \mathrm{osc}_{B_r^{\mathrm{plat}} \oplus B}(\Lambda_\alpha) \le \mathrm{osc}_{B_{2r}^{\mathrm{plat}}}(\Lambda_\alpha) \le \varepsilon_{\mathrm{osc}}$ since $B \subset B_r^{\mathrm{plat}}$ implies $B_r^{\mathrm{plat}} \oplus B \subset B_{2r}^{\mathrm{plat}}$. The same conclusions for the *linear–box* follow from Lemma 7.82 together with the uniform bounds and the fact that the bias terms vanish for constant widths.

(b) Radius-reduction inheritance. Take $\mathcal{N} = B_{r'}^{\mathrm{plat}}(\mathrm{centre})$ with $r' = r - \rho$. Then $B_{r'}^{\mathrm{plat}} \oplus B \subset B_r^{\mathrm{plat}}$, so by eq. (7.76) $\mathrm{osc}_{B_{r'}^{\mathrm{plat}}}(\overline{\Lambda}_\alpha) \le \mathrm{osc}_{B_r^{\mathrm{plat}}}(\Lambda_\alpha) \le \varepsilon_{\mathrm{osc}}$. For gradients and Hessians, repeat the argument of (a) with $B \subset B_r^{\mathrm{plat}}$; no change of tolerances is required.

(c) General inheritance with explicit slack. By Lemma 7.79,

$$\left\|\nabla^{\mathrm{plat}}\overline{\Lambda}_\alpha^{\log}(\mathrm{centre})\right\|_\infty \le \frac{1}{\mathrm{Area}(B)} \int_B \left\|\nabla^{\mathrm{plat}}\Lambda_\alpha(z)\right\|_\infty dz.$$

For any $z \in B_r^{\mathrm{plat}}$, the mean-value inequality with the ball-wise bound on $\mathbf{H}^{\mathrm{plat}}\Lambda_\alpha$ gives $\|\nabla^{\mathrm{plat}}\Lambda_\alpha(z)\|_\infty \le \|\nabla^{\mathrm{plat}}\Lambda_\alpha(\mathrm{centre})\|_\infty + (\varepsilon_{\mathrm{curv}} + \omega_H(r))\, d_{\mathrm{plat}}(z, \mathrm{centre})$. Averaging over B yields $\|\nabla^{\mathrm{plat}}\overline{\Lambda}_\alpha^{\log}(\mathrm{centre})\|_\infty \le \varepsilon_{\mathrm{grad}} + (\varepsilon_{\mathrm{curv}} + \omega_H(r))\, \overline{d_{\mathrm{plat}}}^{\,B} \le \varepsilon_{\mathrm{grad}} + \omega_H(r)\, r$, since $\overline{d_{\mathrm{plat}}}^{\,B} \le r$ and the $\varepsilon_{\mathrm{curv}}$ term is already accounted for in the baseline tolerance. The Hessian bound follows from convexity: $\|\mathbf{H}^{\mathrm{plat}}\overline{\Lambda}_\alpha^{\log}\|_{\mathrm{op}} \le \varepsilon_{\mathrm{curv}} + \omega_H(r)$. For the linear–box, add the bias terms of Lemma 7.82. If the slack δ_{grad} dominates the additive term $\omega_H(r)r$ (and the linear–box biases, when used), the original inequalities are preserved. □

Lemma 7.85 (Integrated ⇒ pointwise (local reconstruction))**.** *Work in the plateau log chart $(u, v) = (\ln \lambda_{\mathrm{UV}}, \ln \ell)$ with metric g_{plat} and operators $D_\lambda = \partial_u$, $D_\ell = \partial_v$. Let $B = B^{\log}(u, v; \Delta u, \Delta v)$ be the log–box averaging rectangle centred at (u, v) with half-sizes $\Delta u/2, \Delta v/2 \le r$. Assume $\Lambda_\alpha \in C^{1,1}$ on $B_r^{\mathrm{plat}}(u, v)$ with*

$$\sup_{B_r^{\mathrm{plat}}} \left\|\mathbf{H}^{\mathrm{plat}}\Lambda_\alpha\right\|_{\mathrm{op}} \;\le\; L_\alpha,$$

i.e. the gradient is L_α–Lipschitz on the ball. If $(\lambda_{\mathrm{UV}}, \ell)$ passes the plateau test with the integrated *visibility $\overline{\Lambda}_\alpha^{\log}$ for tolerances $(\tilde{\varepsilon}_{\mathrm{grad}}, \tilde{\varepsilon}_{\mathrm{osc}}, \tilde{\varepsilon}_{\mathrm{curv}})$, then the* pointwise *visibility at the centre obeys*

$$\left|D_\lambda \Lambda_\alpha(u, v)\right| \;\le\; \tilde{\varepsilon}_{\mathrm{grad}} + L_\alpha\, r, \qquad \left|D_\ell \Lambda_\alpha(u, v)\right| \;\le\; \tilde{\varepsilon}_{\mathrm{grad}} + L_\alpha\, r, \tag{7.80}$$

$$\mathrm{osc}_{B_r^{\mathrm{plat}}(u,v)}\, \Lambda_\alpha \;\le\; \tilde{\varepsilon}_{\mathrm{osc}} + 2L_\alpha\, r, \tag{7.81}$$

and

$$\left\|\mathbf{H}^{\mathrm{plat}}\Lambda_\alpha(u, v)\right\|_{\mathrm{op}} \;\le\; \tilde{\varepsilon}_{\mathrm{curv}} + L_\alpha. \tag{7.82}$$

In particular, pointwise certification holds with tolerances rescaled by $O(L_\alpha r)$.

Moreover, if the second partials admit anisotropic *bounds on* B_r^{plat},

$$\sup_{B_r^{\mathrm{plat}}}\left|D_\lambda^2\Lambda_\alpha\right|\le L_{\alpha,\lambda},\qquad \sup_{B_r^{\mathrm{plat}}}\left|D_\ell^2\Lambda_\alpha\right|\le L_{\alpha,\ell},\qquad \sup_{B_r^{\mathrm{plat}}}\left|D_\lambda D_\ell\Lambda_\alpha\right|\le L_{\alpha,\times},$$

then the gradient bounds refine to

$$\left|D_\lambda\Lambda_\alpha(u,v)\right|\ \le\ \tilde\varepsilon_{\mathrm{grad}}+\tfrac12 L_{\alpha,\lambda}\,\Delta u+\tfrac12 L_{\alpha,\times}\,\Delta v,\qquad \left|D_\ell\Lambda_\alpha(u,v)\right|\ \le\ \tilde\varepsilon_{\mathrm{grad}}+\tfrac12 L_{\alpha,\ell}\,\Delta v+\tfrac12 L_{\alpha,\times}\,\Delta u. \tag{7.83}$$

Proof. We use the log–box average of Remark 7.78 and its derivative commutation Lemma 7.79: $D_\lambda\overline{\Lambda}_\alpha^{\log}=\overline{D_\lambda\Lambda_\alpha}^{\log}$ and $D_\ell\overline{\Lambda}_\alpha^{\log}=\overline{D_\ell\Lambda_\alpha}^{\log}$. By the plateau hypothesis for the integrated visibility,

$$\begin{aligned}\left|D_\lambda\overline{\Lambda}_\alpha^{\log}(u,v)\right|&\le\tilde\varepsilon_{\mathrm{grad}},\\ \left|D_\ell\overline{\Lambda}_\alpha^{\log}(u,v)\right|&\le\tilde\varepsilon_{\mathrm{grad}},\\ \left\|\mathbf{H}^{\mathrm{plat}}\overline{\Lambda}_\alpha^{\log}(u,v)\right\|_{\mathrm{op}}&\le\tilde\varepsilon_{\mathrm{curv}},\\ \operatorname{osc}_{B_r^{\mathrm{plat}}}\overline{\Lambda}_\alpha^{\log}&\le\tilde\varepsilon_{\mathrm{osc}}.\end{aligned}$$

(Isotropic gradient bound). For $z=(u',v')\in B$, the L_α–Lipschitz property of $\nabla^{\mathrm{plat}}\Lambda_\alpha$ gives

$$\left|D_\lambda\Lambda_\alpha(u,v)-D_\lambda\Lambda_\alpha(u',v')\right|\le L_\alpha\,d_{\mathrm{plat}}(z,(u,v))\le L_\alpha\,r.$$

Averaging over B and using derivative commutation yields

$$\left|D_\lambda\Lambda_\alpha(u,v)-D_\lambda\overline{\Lambda}_\alpha^{\log}(u,v)\right|\le\frac{1}{|B|}\int_B L_\alpha\,d_{\mathrm{plat}}(z,(u,v))\,dz\le L_\alpha\,r,$$

whence (7.80) for D_λ by the triangle inequality with the integrated bound; the D_ℓ case is identical.

(Oscillation). For any $z,z'\in B_r^{\mathrm{plat}}(u,v)$,

$$\left|\Lambda_\alpha(z)-\Lambda_\alpha(z')\right|\le\left|\overline{\Lambda}_\alpha^{\log}(z)-\overline{\Lambda}_\alpha^{\log}(z')\right|+\left|\Lambda_\alpha(z)-\overline{\Lambda}_\alpha^{\log}(z)\right|+\left|\Lambda_\alpha(z')-\overline{\Lambda}_\alpha^{\log}(z')\right|.$$

The first term is bounded by $\operatorname{osc}_{B_r^{\mathrm{plat}}}\overline{\Lambda}_\alpha^{\log}\le\tilde\varepsilon_{\mathrm{osc}}$. For the other two, fix z and note that $\left|\Lambda_\alpha(z)-\overline{\Lambda}_\alpha^{\log}(z)\right|\le\frac{1}{|B|}\int_B\left|\Lambda_\alpha(z)-\Lambda_\alpha(\zeta)\right|d\zeta\le\frac{1}{|B|}\int_B L_\alpha\,d_{\mathrm{plat}}(\zeta,z)\,d\zeta\le L_\alpha r$, since $B\subset B_r^{\mathrm{plat}}(z)$. Taking the supremum over $z,z'\in B_r^{\mathrm{plat}}$ proves (7.81).

(Hessian). By convexity of the operator norm and Lemma 7.79 applied to first derivatives,

$$\begin{aligned}\left\|\mathbf{H}^{\mathrm{plat}}\Lambda_\alpha(u,v)\right\|_{\mathrm{op}}&\le\left\|\mathbf{H}^{\mathrm{plat}}\overline{\Lambda}_\alpha^{\log}(u,v)\right\|_{\mathrm{op}}+\sup_B\left\|\mathbf{H}^{\mathrm{plat}}\Lambda_\alpha-\mathbf{H}^{\mathrm{plat}}\overline{\Lambda}_\alpha^{\log}\right\|_{\mathrm{op}}\\&\le\tilde\varepsilon_{\mathrm{curv}}+\sup_B\left\|\mathbf{H}^{\mathrm{plat}}\Lambda_\alpha\right\|_{\mathrm{op}},\end{aligned}$$

and the last supremum is $\le L_\alpha$ by hypothesis on B_r^{plat}, proving (7.82).

(Anisotropic refinement). Write the double average as $\overline{D_\lambda\Lambda_\alpha}^{\log} = \overline{\overline{D_\lambda\Lambda_\alpha}^{\,u}}^{\,v}$, where $\bar{\cdot}^{u}$ denotes the u–average at fixed v and similarly for v. Then

$$\left|D_\lambda\Lambda_\alpha(u,v) - \overline{D_\lambda\Lambda_\alpha}^{\log}\right| \le \underbrace{\left|D_\lambda\Lambda_\alpha(u,v) - \overline{D_\lambda\Lambda_\alpha}^{\,u}(v)\right|}_{\text{(I)}} + \underbrace{\left|\overline{D_\lambda\Lambda_\alpha}^{\,u}(v) - \overline{\overline{D_\lambda\Lambda_\alpha}^{\,u}}^{\,v}\right|}_{\text{(II)}}.$$

For (I), one-dimensional Taylor with integral remainder in u and the bound $|D_\lambda^2\Lambda_\alpha| \le L_{\alpha,\lambda}$ on B_r^{plat} give (I) $\le \frac{1}{2}L_{\alpha,\lambda}\Delta u$. For (II), differentiate $\overline{D_\lambda\Lambda_\alpha}^{\,u}(v)$ in v and use $\partial_v\overline{D_\lambda\Lambda_\alpha}^{\,u} = \overline{D_\lambda D_\ell\Lambda_\alpha}^{\,u}$ with $|D_\lambda D_\ell\Lambda_\alpha| \le L_{\alpha,\times}$; then a one-dimensional mean-value bound yields (II) $\le \frac{1}{2}L_{\alpha,\times}\Delta v$. Combine with $|\,\overline{D_\lambda\Lambda_\alpha}^{\log}\,| \le \tilde{\varepsilon}_{\mathrm{grad}}$ to obtain the first inequality in (7.83); the second follows by swapping the roles of u and v, using $L_{\alpha,\ell}$ and $L_{\alpha,\times}$. □

Joint (multi-channel) integrated plateaus.

Definition 7.86 (Joint integrated plateau)**.** Let $\mathfrak{C} \subset \{\mathrm{law}, \mathrm{state}, \mathrm{mix}, \ldots\}$ and fix averaging widths $(\Delta\lambda, \Delta\ell)$ (or $(\Delta u, \Delta v)$ in the log chart). For each channel $\alpha \in \mathfrak{C}$, choose a tolerance triple $(\tilde{\varepsilon}_{\mathrm{grad}}, \tilde{\varepsilon}_{\mathrm{osc}}, \tilde{\varepsilon}_{\mathrm{curv}})$ consistent with Lemmas 7.84 and 7.85. Define the *joint integrated plateau* by

$$\overline{\mathfrak{P}}_{\mathfrak{C}}(\Delta\lambda, \Delta\ell) := \bigcap_{\alpha\in\mathfrak{C}} \Big\{(\lambda_{\mathrm{UV}}, \ell) \in \mathcal{R} : \ \overline{\Lambda}_\alpha \text{ satisfies Section 7.3.5 with } (\tilde{\varepsilon}_{\mathrm{grad}}, \tilde{\varepsilon}_{\mathrm{osc}}, \tilde{\varepsilon}_{\mathrm{curv}})\Big\}. \tag{7.84}$$

Proposition 7.87 (Inclusions/equivalence with pointwise plateaus)**.** *Let $\mathfrak{P}_{\mathfrak{C}} := \bigcap_{\alpha\in\mathfrak{C}} \mathfrak{P}_\alpha$ be the pointwise joint plateau (7.64).*

*(i) (*Forward inclusion*) If, for every $\alpha \in \mathfrak{C}$, the averaging box B at $(\lambda_{\mathrm{UV}}, \ell)$ is contained in the test ball $B_r^{\mathrm{plat}}(\lambda_{\mathrm{UV}}, \ell)$, and the same tolerances are used for $\overline{\Lambda}_\alpha$ as for Λ_α, then*

$$\mathfrak{P}_{\mathfrak{C}} \ \subseteq \ \overline{\mathfrak{P}}_{\mathfrak{C}}(\Delta\lambda, \Delta\ell).$$

*(ii) (*Reverse inclusion with rescaled tolerances*) Assume for each $\alpha \in \mathfrak{C}$ the local $C^{1,1}$ (or Lipschitz–gradient) hypothesis of Lemma 7.85 on B_r^{plat} and half-sizes $\Delta u/2, \Delta v/2 \le r$ in the log chart. Then there exist rescaled tolerances $(\varepsilon_{\mathrm{grad}}, \varepsilon_{\mathrm{osc}}, \varepsilon_{\mathrm{curv}})$ determined by $(\tilde{\varepsilon}_\bullet, \Delta u, \Delta v)$ and the channel Lipschitz data such that*

$$\overline{\mathfrak{P}}_{\mathfrak{C}}(\Delta\lambda, \Delta\ell) \ \subseteq \ \mathfrak{P}_{\mathfrak{C}}.$$

*(iii) (*Equivalence*) Under the hypotheses of (i) and (ii) with $B \subset B_r^{\mathrm{plat}}$ and using the tolerance recipes of Lemmas 7.84 and 7.85 (log–box: identical tolerances; linear–box: rescaled as in Corollary 7.41), the two sets coincide:*

$$\overline{\mathfrak{P}}_{\mathfrak{C}}(\Delta\lambda, \Delta\ell) \ = \ \mathfrak{P}_{\mathfrak{C}}.$$

Proof. (i) Apply Lemma 7.84 channelwise and intersect over $\alpha \in \mathfrak{C}$. (ii) Apply Lemma 7.85 channelwise with the stated rescalings and intersect. (iii) Combine (i) and (ii). □

Edge handling and normalisation.

Definition 7.88 (Renormalised edge average)**.** When the averaging rectangle B centred at $(\lambda_{\mathrm{UV}}, \ell)$ intersects the boundary of $\mathcal{R}$, define the *renormalised* linear–box average by

$$\overline{\Lambda}_{\alpha}^{\mathrm{ren}}(\lambda_{\mathrm{UV}}, \ell) := \frac{1}{|B \cap \mathcal{R}|} \int_{B \cap \mathcal{R}} \Lambda_{\alpha}(\lambda, \tilde{\ell})\, d\lambda\, d\tilde{\ell},$$

and analogously in the log chart. For discrete grids, use weights $w_{ij} \geq 0$ supported on available pixels with $\sum w_{ij} = 1$, and replace derivatives by the finite-difference stencils of Section 7.3.3.

Proposition 7.89 (Derivative commutation and bias control at edges)**.** *Let* $\overline{\Lambda}_{\alpha}^{\mathrm{ren}}$ *be as in Definition 7.88.*

*(i) (*Log–box with extension*) If* Λ_{α} *is extended to a* $C^{1,1}$ *function on a neighbourhood of* $\mathcal{R}$ *(e.g. by even reflection) and* $\Delta u, \Delta v$ *are constant, then the commutation identities* (7.74) *hold with* $\overline{\Lambda}_{\alpha}^{\mathrm{ren}}$ *in place of* $\overline{\Lambda}_{\alpha}^{\log}$*.*

*(ii) (*Linear–box bias*) Without extension, for* $C^{1,1}$ *data on* $\mathcal{R}$ *the edge bias obeys first–order bounds of the form*

$$\left| D_{\lambda} \overline{\Lambda}_{\alpha}^{\mathrm{ren}} - \overline{D_{\lambda} \Lambda_{\alpha}}^{\mathrm{ren}} \right| \;\leq\; C\, L_{\alpha}\, \frac{|B \setminus (B \cap \mathcal{R})|}{|B \cap \mathcal{R}|},$$

with C *absolute and* L_{α} *the local Lipschitz constant of* $\nabla^{\mathrm{plat}} \Lambda_{\alpha}$*; analogous bounds hold for* D_{ℓ} *and the Hessian entries. Thus any induced change in the plateau inequalities can be absorbed into the scan's tolerance budget by shrinking* $\Delta u, \Delta v$ *near edges or by using one–sided stencils.*

*(iii) (*Discrete grids*) For weighted discrete averages, centred differences commute with averaging up to* $O(\varepsilon_{\mathrm{grid}})$*, where* $\varepsilon_{\mathrm{grid}}$ *is the consistency error of the stencil; choosing the grid resolution so that* $\varepsilon_{\mathrm{grid}} \ll \min\{\tilde{\varepsilon}_{\mathrm{grad}}, \tilde{\varepsilon}_{\mathrm{curv}}\}$ *preserves pass/fail decisions.*

Regularisation and gauge stability.

Proposition 7.90 (Regulator/gauge stability of joint integrated plateaus)**.** *Assume the admissibility conditions of Section 7.3.7 and the gauge/BRST hypotheses of Section 7.3.6. Then* $\overline{\mathfrak{P}}_{\mathfrak{C}}(\Delta\lambda, \Delta\ell)$ *is:*

(i) Regulator robust: *replacing* $(P_x, K_{\mathrm{IR}}^{(\ell)})$ *by an admissible* $(\tilde{P}_x, \tilde{K}_{\mathrm{IR}}^{(\ell)})$ *changes* $\overline{\mathfrak{P}}_{\mathfrak{C}}$*, if at all, only on the null frontier where inequalities are saturated (cf. Corollary 7.74).*

(ii) Gauge-parameter invariant on shell: *variations of* α_{gf} *leave* $\overline{\mathfrak{P}}_{\mathfrak{C}}$ *unchanged when evaluated on* $\mathcal{H}_{\mathrm{phys}}$ *(Nielsen independence, Proposition 7.63).*

7.3.9 Acceptance Criteria and Certification Pipeline

Acceptance region.

Definition 7.91 (Acceptance region and admissibility hypotheses)**.** Fix tolerances $(\varepsilon_{\mathrm{grad}}, \varepsilon_{\mathrm{osc}}, \varepsilon_{\mathrm{curv}}, \varepsilon_{\mathrm{reg}}, \varepsilon_{\mathrm{swap}}) > 0$ and a compact scan window $\mathcal{R} \subset \mathbb{R}^2_{>0}$. Let $\mathfrak{P}_{\mathfrak{C}}$ be the joint (pointwise) plateau (7.64) for $\mathfrak{C} = \{\mathrm{law}, \mathrm{state}\}$ as defined in Section 7.3.5, and let the regularisation setting satisfy Section 7.3.7. We say that *acceptance* holds on a connected component $\mathcal{P} \subset \mathcal{R}$ if the following hypotheses are satisfied:

(H1) **Non-emptiness of the joint plateau:** $\mathcal{P} \subset \mathfrak{P}_{\{\mathrm{law},\mathrm{state}\}}(\varepsilon_{\mathrm{grad}}, \varepsilon_{\mathrm{osc}}, \varepsilon_{\mathrm{curv}}; \mathcal{R})$ and $\mathcal{P} \neq \varnothing$.

(H2) **Regularisation stability:** the bound (7.67) holds on $\mathcal{R}$, with sufficient conditions given in Section 7.3.7 (in particular (7.68), (7.69), and Proposition 7.73).

(H3) **BRST/gauge robustness on shell:** the Slavnov identity and anomaly filters eqs. (7.16) and (7.18) hold, so that visibilities are gauge-parameter independent on $\mathcal{H}_{\mathrm{phys}}$ by Section 7.2.9 (see also Section 7.3.6 and Proposition 7.63).

(H4) **Kinematics/causality sanity:** for all $(\lambda_{\mathrm{UV}}, \ell) \in \mathcal{P}$, the tachyon test (7.14) and hyperbolicity (7.15) are satisfied.

(H5) **Commutation of limits (corner test):** on $\mathcal{P}$, the swap criterion (7.45) holds with tolerance $\varepsilon_{\mathrm{swap}}$.

We call the union of all such components the *acceptance region* and denote it by

$$\mathcal{A}_{\{\mathrm{law},\mathrm{state}\}}(\varepsilon_\bullet; \mathcal{R}) := \bigcup \{\mathcal{P} \subset \mathcal{R} : \ \text{(H1)–(H5) hold on } \mathcal{P}\}.$$

Theorem 7.92 (Certification from acceptance)**.** *Assume* (H1)–(H5) *on a connected component* $\mathcal{P} \subset \mathcal{R}$*. Then every physical pole extracted on* $\mathcal{P}$ *is*

(i) gauge-parameter independent on shell*, by eqs.* (7.22) *and* (7.23)*;*

(ii) UV/IR-regulator robust*, with drift controlled by* (7.67) *(sufficiently by Proposition 7.73);*

(iii) kinematically sound*, i.e. obeys* (7.14) *and propagates hyperbolically by* (7.15)*;*

(iv) stable under limit swapping*, within* $\varepsilon_{\mathrm{swap}}$*, by* (7.45)*.*

In particular, the robust, gauge-independent pole extraction statement of Proposition 7.34 holds on $\mathcal{P}$*.*

Proof. (i) follows from Section 7.2.9 evaluated on $\mathcal{H}_{\mathrm{phys}}$, together with Section 7.2.6. (ii) is the content of Section 7.3.7 and Proposition 7.73, applied uniformly on $\mathcal{P}$; see also Corollary 7.74. (iii) is (H4). (iv) is (H5). Combining these with Definition 7.31 (and Section 7.3.2) yields the claim and recovers Proposition 7.34. □

Certified centre and uncertainty budget.

Definition 7.93 (Certified centre and uncertainty)**.** Let $\mathcal{P}$ be a connected component of $\mathcal{A}_{\{\mathrm{law,state}\}}(\varepsilon_\bullet;\mathcal{R})$. A *certified centre* $(\lambda_\star,\ell_\star)\in\mathcal{P}$ is any point maximising the inscribed g_{plat}-radius or, equivalently, minimising a convex surrogate such as $\|\nabla^{\mathrm{plat}}\Lambda_{\mathrm{law}}\|^2+\|\nabla^{\mathrm{plat}}\Lambda_{\mathrm{state}}\|^2$ on $\mathcal{P}$. The reported channel values are $\Lambda_\alpha(\lambda_\star,\ell_\star)$, accompanied by the total budget

$$\varepsilon_{\mathrm{tot}} := \max\Big\{\varepsilon_{\mathrm{grad}},\ \varepsilon_{\mathrm{osc}},\ \varepsilon_{\mathrm{curv}},\ \varepsilon_{\mathrm{reg}},\ \varepsilon_{\mathrm{swap}}\Big\}. \tag{7.85}$$

For physical poles m_i^2 the drift metric Δ_i^{rel} of (7.50) is also reported.

Proposition 7.94 (Intra-component bounds from geometric tolerances)**.** *Let $\mathcal{P}$ and $(\lambda_\star,\ell_\star)$ be as above. For any $(\lambda,\ell)\in\mathcal{P}$ one has*

$$\big|\Lambda_\alpha(\lambda,\ell)-\Lambda_\alpha(\lambda_\star,\ell_\star)\big| \le \varepsilon_{\mathrm{grad}}\, d_{\mathrm{plat}}\Big((\lambda,\ell),(\lambda_\star,\ell_\star)\Big) + \tfrac{1}{2}\,\varepsilon_{\mathrm{curv}}\, d_{\mathrm{plat}}\Big((\lambda,\ell),(\lambda_\star,\ell_\star)\Big)^2 + \varepsilon_{\mathrm{osc}},$$

and the same estimate holds for $\overline{\Lambda}_\alpha$ (with log–box) by Proposition 7.83.

Proof. Taylor's theorem with integral remainder in the log chart, together with the bounds defining $\mathfrak{P}_{\{\mathrm{law,state}\}}$; cf. Lemma 7.45. □

Certification pipeline.

Definition 7.95 (Operational certification pipeline)**.** Under the standing assumptions of Sections 7.2.5, 7.2.6, 7.2.9 and 7.3.7, execute:

- **C1. Knob selection.** Fix $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ satisfying Definitions 6.280 and 7.26 (i.e. (7.29), (7.30), (7.32)) and choose $\mathcal{R}$.
- **C2. Pre-flight consistency.** Build $\mathbb{H}^{(\lambda_{\mathrm{UV}},\ell)}$ and perform the $\Pi_{\mathrm{T}}/\Pi_{\mathrm{L}}$ split; verify quartet decoupling eq. (7.11), tachyon test (7.14), hyperbolicity (7.15), and eqs. (7.16) and (7.18).
- **C3. Band-limited EOM (on shell).** Check eqs. (6.333) and (6.334) on $\mathcal{H}_{\mathrm{phys}}$ (Nielsen applicability).
- **C4. Visibility scans.** Compute $\Lambda_{\mathrm{law}},\Lambda_{\mathrm{state}}$ on a uniform grid in (u,v); estimate ∇ and $\mathbf{H}$ via centred stencils (resolution r), cf. Section 7.3.3.
- **C5. Plateau extraction.** Form channel masks via Section 7.3.5; obtain the joint plateau (7.64). If needed, switch to integrated visibilities Definition 7.39 and reuse the tests with Lemmas 7.84 and 7.85.
- **C6. Commutation-of-limits test.** On each component, verify (7.45) by swapped paths to $(\infty,0^+)$.
- **C7. Regularisation cross-check.** Repeat **C4**–**C6** with $(\tilde{P}_x,\tilde{K}_{\mathrm{IR}}^{(\ell)})$; verify (7.67) (use Proposition 7.73 and Corollary 7.74).
- **C8. Gauge-parameter certification.** Sample α_{gf} values; confirm on-shell invariance via Section 7.2.9 and eq. (7.66).
- **C9. Centre selection and report.** Choose $(\lambda_\star,\ell_\star)$ (e.g. maximal inscribed radius); report $\Lambda_\alpha(\lambda_\star,\ell_\star)$, pole data with Δ_i^{rel} (7.50), and uncertainty $\varepsilon_{\mathrm{tot}}$ (7.85), together with component diameter and metrics of Section 7.3.2.

Milestone: scale-tagged, BRST-closed measurement layer. We established two operational "knobs"—the UV projector and the IR window—formalised in Definitions 6.280 and 7.26 and composed in the canonical order "UV then IR" via the band functor Definitions 7.28 and 7.29. These tags are BRST-closed and ghost-number zero, commute with channel projectors, and are compatible with background-field Slavnov–Taylor identities, ensuring that all band-limited composites are functorially well-defined and positivity-preserving on the physical sector $\mathcal{H}_{\mathrm{phys}}$ (Section 7.2.6 and eq. (7.12); see also [114, 82, 91, 241]).

Milestone: decoupling of unphysical sectors and positivity. Working with the scale-tagged quadratic form and Hessian $\mathbb{H}^{(\lambda_{\mathrm{UV}},\ell)}$ (Section 7.2.5 and eq. (7.10)), the transverse/longitudinal decomposition and the quartet mechanism imply that Faddeev–Popov ghosts and longitudinal gauge modes do not contribute to physical matrix elements (Section 7.2.6 and eq. (7.11); cf. [104, 105, 106, 442, 91]). Consequently, gauge-invariant two-point functions on $\mathcal{H}_{\mathrm{phys}}$ admit a Källén–Lehmann representation with non-negative spectral density (Corollary 7.24 and eq. (7.12)), which underpins unitarity in the physical sector [82, 91, 241].

Milestone: kinematic sanity—tachyon test and hyperbolicity. From the band Hessian we extracted the physical kinetic/mass blocks and the mass operator $\mathbb{M}^2_{\mathrm{phys}}(\lambda_{\mathrm{UV}},\ell)$ (Section 7.2.7 and eq. (7.13)). The *tachyon test* demands non-negativity of the physical Klein–Gordon-type operator's spectrum (eq. (7.14)); *hyperbolicity* requires positivity of the principal symbol with respect to the emergent chronogeometry, guaranteeing a well-posed Cauchy problem (eq. (7.15)). These conditions ensure causal, stable propagation on the band.

Milestone: gauge-parameter independence on shell (Nielsen). At the quantum level, anomaly filters together with the Slavnov identity $\mathcal{S}(\Gamma) = 0$ (Section 7.2.8 and eqs. (7.16) and (7.18)) guarantee BRST control. Extending BRST by a gauge-parameter doublet yields the Nielsen master identity, from which we proved that *physical poles are gauge-parameter independent* (Section 7.2.9 and eqs. (7.20), (7.22) and (7.23)). We further showed that channel visibilities Λ_α inherit this independence on shell (Section 7.3.6 and eq. (7.66)).

Milestone: visibilities, plateau geometry, and acceptance. We defined gauge-invariant, BRST-closed *visibilities* $\Lambda_\alpha(\lambda_{\mathrm{UV}},\ell)$ and their log-scale gradients/curvatures (Section 7.3.4 and eq. (7.55)). This led to a geometric notion of *plateau* in the log-chart metric g_{plat}, using gradient, oscillation, and Hessian bounds (Section 7.3.5, Definition 7.44, Lemma 7.45, and Proposition 7.46). Global certification metrics (relative drift/curvature and commuting-limits) were tied to these geometric bounds and shown equivalent up to controllable rescalings (Section 7.3.2, Definition 7.31, eqs. (7.45) and (7.46), and Lemma 7.50).

Milestone: integrated visibilities and inheritance. To stabilise numerics/experiments we introduced *integrated* (box-averaged) visibilities and proved that plateau properties *inherit both ways*: pointwise ⇒ integrated and integrated ⇒ pointwise, with explicit tolerance/bias control (Section 7.3.8, Definition 7.39, Lemmas 7.40, 7.84 and 7.85, Corollary 7.41, and Proposition 7.83). This enables certification on smoothed data without losing quantitative guarantees.

Milestone: regularisation stability and joint plateaus. We identified sufficient conditions for *regularisation stability* under admissible changes of the UV projector family and IR kernels—L^1-closeness for IR windows and gapped spectral stability for UV projectors—together with a Lipschitz continuity bound for the visibility functional (Section 7.3.7, eqs. (7.68), (7.69) and (7.71), and Proposition 7.73). As a result, plateau membership and pole extraction are regulator-robust up to a controlled budget, and plateaus computed with different admissible schemes agree away from measure-zero frontiers (Corollary 7.74). We also formalised *joint* (multi-channel) plateaus and showed inclusions/equivalence with their integrated counterparts under the inheritance lemmas (eq. (7.64), Definition 7.86, and Propositions 7.87 and 7.90).

Milestone: operational pipeline and certified centres. All ingredients were assembled into an *acceptance and certification pipeline* that is reproducible and automatable (Section 7.3.9). Acceptance requires a non-empty joint plateau, regulator robustness, BRST/gauge robustness on shell, kinematic sanity, and commuting UV/IR limits (Definition 7.91 and Theorem 7.92). Certified centres and uncertainty budgets were defined and tied to intra-component drift/curvature bounds (Definition 7.93 and Proposition 7.94), ensuring that reported visibilities and physical poles come with explicit, auditable tolerances.

7.3.10 Quantitative Plateau Metrics

Definition 7.96 (Pointwise quantitative plateau metrics (log chart))**.** Work in the log chart $(u, v) = (\ln \lambda_{\mathrm{UV}}, \ln \ell)$ so that $D_\lambda = \partial_u$ and $D_\ell = \partial_v$ (cf. Section 7.3.4 and the g_{plat} geometry in Section 7.3.5). For each channel α and each $(\lambda_{\mathrm{UV}}, \ell) \in \mathcal{R}$, define:

$$\mathrm{Slope}_\alpha(\lambda_{\mathrm{UV}}, \ell) := \left\|\nabla \Lambda_\alpha(\lambda_{\mathrm{UV}}, \ell)\right\|_2 = \sqrt{\left(D_\lambda \Lambda_\alpha\right)^2 + \left(D_\ell \Lambda_\alpha\right)^2}, \tag{7.86}$$

$$\mathrm{Curv}_\alpha(\lambda_{\mathrm{UV}}, \ell) := \left\|\mathbf{H}\Lambda_\alpha(\lambda_{\mathrm{UV}}, \ell)\right\|_{\mathrm{F}} = \sqrt{\sum_{i,j\in\{\lambda,\ell\}} \left|D_i D_j \Lambda_\alpha\right|^2}, \tag{7.87}$$

$$\mathrm{Osc}_\alpha(\lambda_{\mathrm{UV}}, \ell; r) := \max_{(\lambda',\ell')\in B_r(\lambda_{\mathrm{UV}},\ell)} \left|\Lambda_\alpha(\lambda', \ell') - \Lambda_\alpha(\lambda_{\mathrm{UV}}, \ell)\right|, \tag{7.88}$$

where $\|\cdot\|_{\mathrm{F}}$ is the Frobenius norm of the 2×2 Hessian, and $B_r(\lambda_{\mathrm{UV}}, \ell)$ is the g_{plat}-geodesic ball of radius r in the log chart (cf. Section 7.3.5).

Remark 7.97 (Well-posedness, invariances, and discretisation)**.** (i) The metrics in Definition 7.96 are computed from BRST-closed inputs and are gauge-parameter independent on shell by Section 7.3.6 and eq. (7.66). (ii) If $\Lambda_\alpha \in C^2$ locally, the quantities are classical; otherwise interpret D_i in the a.e. sense and Curv_α via essential bounds of distributional second derivatives; Osc_α is always well-defined. (iii) For discrete scans, replace D_i and $D_i D_j$ by the centred stencils of Section 7.3.3; the induced consistency error is absorbed into the tolerance budget.

Norm comparisons and links to the geometric test.

Proposition 7.98 (Norm comparisons and consequences for geometric plateaus)**.** *For any channel α and any $(\lambda_{\mathrm{UV}}, \ell) \in \mathcal{R}$, the pointwise metrics of Definition 7.96 satisfy the Euclidean/sup and Frobenius/operator norm comparisons in $\mathbb{R}^2$:*

$$\|\nabla\Lambda_\alpha\|_\infty \;\le\; \mathrm{Slope}_\alpha \;\le\; \sqrt{2}\,\|\nabla\Lambda_\alpha\|_\infty, \qquad \|\mathbf{H}\Lambda_\alpha\|_{\mathrm{op}} \;\le\; \mathrm{Curv}_\alpha \;\le\; \sqrt{2}\,\|\mathbf{H}\Lambda_\alpha\|_{\mathrm{op}}. \tag{7.89}$$

Consequently, if $(\lambda_{\mathrm{UV}}, \ell)$ lies in the geometric plateau set of Section 7.3.5 with tolerances $(\varepsilon_{\mathrm{grad}}, \varepsilon_{\mathrm{osc}}, \varepsilon_{\mathrm{curv}})$, then

$$\mathrm{Slope}_\alpha(\lambda_{\mathrm{UV}}, \ell) \;\le\; \sqrt{2}\,\varepsilon_{\mathrm{grad}}, \qquad \mathrm{Curv}_\alpha(\lambda_{\mathrm{UV}}, \ell) \;\le\; \sqrt{2}\,\varepsilon_{\mathrm{curv}}, \qquad \mathrm{Osc}_\alpha(\lambda_{\mathrm{UV}}, \ell; r) \;\le\; \varepsilon_{\mathrm{osc}}. \tag{7.90}$$

Proof. In $\mathbb{R}^2$, for any vector x one has $\|x\|_\infty \le \|x\|_2 \le \sqrt{2}\|x\|_\infty$. Apply this to $x = \nabla\Lambda_\alpha$ to obtain the first inequality chain in (7.89), hence the bound on Slope_α. For a 2×2 matrix A, singular-value calculus gives $\|A\|_{\mathrm{op}} \le \|A\|_{\mathrm{F}} \le \sqrt{2}\,\|A\|_{\mathrm{op}}$ (since $\|A\|_{\mathrm{F}} = \sqrt{\sigma_1^2 + \sigma_2^2} \le \sqrt{2}\,\max\{\sigma_1, \sigma_2\}$). Apply this to $A = \mathbf{H}\Lambda_\alpha$ to obtain the second chain. The implications (7.90) then follow directly from the geometric test in Section 7.3.5, where the gradient and Hessian bounds are stated in the sup and operator norms, and from the definition (7.88) of Osc_α. □

η-deep plateaus and plateau radius.

Definition 7.99 (η-deep plateau point)**.** Fix tolerances $(\varepsilon_{\mathrm{grad}}, \varepsilon_{\mathrm{curv}}, \varepsilon_{\mathrm{osc}}) > 0$ and a channel set $\{\mathrm{law}, \mathrm{state}\}$. Given $\eta \in (0, 1]$, a point $(\lambda_\star, \ell_\star) \in \mathcal{R}$ is an *η-deep plateau point* if, for all $\alpha \in \{\mathrm{law}, \mathrm{state}\}$,

$$\mathrm{Slope}_\alpha(\lambda_\star, \ell_\star) \;\le\; \eta\,\varepsilon_{\mathrm{grad}}, \qquad \mathrm{Curv}_\alpha(\lambda_\star, \ell_\star) \;\le\; \eta\,\varepsilon_{\mathrm{curv}}, \qquad \mathrm{Osc}_\alpha(\lambda_\star, \ell_\star; r_0) \;\le\; \eta\,\varepsilon_{\mathrm{osc}} \tag{7.91}$$

for some reference radius $r_0 > 0$ with $B_{r_0}(\lambda_\star, \ell_\star) \subset \mathcal{R}$.

Definition 7.100 (Plateau radius)**.** Let $(\lambda_\star, \ell_\star)$ be η-deep in the sense of Definition 7.99. The *plateau radius* at $(\lambda_\star, \ell_\star)$ is

$$\rho_\star(\eta) := \sup\Big\{\, r \in [0, r_{\max}) \;:\; \max_{\alpha \in \{\mathrm{law},\mathrm{state}\}} \mathrm{Osc}_\alpha(\lambda_\star, \ell_\star; r) \;\le\; \eta\,\varepsilon_{\mathrm{osc}} \,\Big\}, \tag{7.92}$$

with $r_{\max} := \sup\{r : B_r(\lambda_\star, \ell_\star) \subset \mathcal{R}\}$. Equivalently, $\rho_\star(\eta)$ is the largest geodesic radius over which the oscillation constraint in (7.91) remains satisfied; the slope/curvature constraints are r-independent and are checked at the centre.

Lemma 7.101 (Monotonicity and characterisation of $\rho_\star$)**.** *For each α, the function $r \mapsto \mathrm{Osc}_\alpha(\lambda_\star, \ell_\star; r)$ is non-decreasing on $[0, r_{\max})$. Hence $\rho_\star(\eta)$ of (7.92) is well-defined and satisfies*

$$\rho_\star(\eta) = \inf_{\alpha \in \{\mathrm{law},\mathrm{state}\}} \sup\Big\{\, r : \; \mathrm{Osc}_\alpha(\lambda_\star, \ell_\star; r) \le \eta\,\varepsilon_{\mathrm{osc}} \,\Big\}.$$

If the maps $r \mapsto \mathrm{Osc}_\alpha(\cdot; r)$ are continuous, $\rho_\star(\eta)$ is the smallest r at which one of the channels saturates the oscillation bound.

Proof. If $0 \le r_1 \le r_2$, then $B_{r_1} \subseteq B_{r_2}$, whence the sup–inf range of Λ_α over B_{r_1} is bounded by that over B_{r_2}; thus $r \mapsto \mathrm{Osc}_\alpha(\cdot; r)$ is non-decreasing. The characterisation follows by taking the intersection over channels of the sublevel sets $\{r : \mathrm{Osc}_\alpha \le \eta\varepsilon_{\mathrm{osc}}\}$ and using monotonicity. Continuity in r (e.g. for continuous Λ_α) implies that crossing occurs at a first saturation radius. □

Remark 7.102 (Practical estimation of $\rho_\star$)**.** In numerical scans, one may *grow* r inside $\mathcal{R}$ until any of the three inequalities in (7.91) is saturated. Since Slope_α and Curv_α are r-independent pointwise metrics, $\rho_\star(\eta)$ is typically limited by the oscillation constraint (cf. Lemma 7.101). The resulting $\rho_\star$ provides an intrinsic confidence scale—measured in the g_{plat} distance—for downstream evaluations at $(\lambda_\star, \ell_\star)$ and links directly to the acceptance/certification steps in Section 7.3.9.

7.3.11 Channel and Sector Compatibility

Definition 7.103 (Sector-resolved visibilities and plateaus)**.** Let $\{\Pi_\beta\}$ be BRST-closed channel/sector projectors that commute with the UV/IR tags and with $\hat{Q}_{\mathrm{BRST}}$ (cf. channel/background compatibility in Definition 7.28 and Proposition 7.30). Given a channel functional $\mathcal{F}_\alpha$ (as in eq. (7.54)), define the *sector-resolved visibility*

$$\Lambda_{\alpha|\beta}(\lambda_{\mathrm{UV}}, \ell) := \mathcal{F}_\alpha\Big[\Pi_\beta(\cdot)\Big], \qquad \Lambda_\alpha = \sum_{\beta\in\mathcal{B}} \Lambda_{\alpha|\beta} \quad \text{if } \mathcal{F}_\alpha \text{ is linear}, \tag{7.93}$$

for a specified finite sector set $\mathcal{B}$. The *sectorwise plateau set* $\mathfrak{P}_{\alpha|\beta}$ is defined by applying the geometric test of Section 7.3.5 (or the quantitative metrics of Definition 7.96) to $\Lambda_{\alpha|\beta}$. We say *sector compatibility* holds at $(\lambda_\star, \ell_\star)$ if

$$(\lambda_\star, \ell_\star) \in \bigcap_{\beta\in\mathcal{B}} \mathfrak{P}_{\alpha|\beta}, \tag{7.94}$$

i.e. each resolved sector passes the plateau test at $(\lambda_\star, \ell_\star)$.

Remark 7.104 (Gauge/BRST robustness of sector visibilities)**.** Since $[P_x(\lambda_{\mathrm{UV}}), \Pi_\beta] = [K_{\mathrm{IR}}^{(\ell)}, \Pi_\beta] = 0$ and $\mathrm{gh}(\Pi_\beta) = 0$ (cf. Proposition 7.30), the sector-resolved visibilities are built from BRST-closed inputs and inherit the on-shell gauge-parameter independence from the Nielsen framework (Section 7.3.6 and eq. (7.66)).

Proposition 7.105 (Sectorwise ⇒ channel plateau under linearity)**.** *Assume $\mathcal{F}_\alpha$ is linear in its sector inputs so that (7.93) holds. If, for tolerances $(\varepsilon_{\mathrm{grad}}, \varepsilon_{\mathrm{osc}}, \varepsilon_{\mathrm{curv}})$ and some fixed radius $r > 0$,*

$$\max\Big\{\|\nabla^{\mathrm{plat}}\Lambda_{\alpha|\beta}\|_\infty,\ \|\mathbf{H}^{\mathrm{plat}}\Lambda_{\alpha|\beta}\|_{\mathrm{op}},\ \mathrm{osc}_{B_r^{\mathrm{plat}}}\Lambda_{\alpha|\beta}\Big\} \ \le\ \eta\cdot(\varepsilon_{\mathrm{grad}}, \varepsilon_{\mathrm{curv}}, \varepsilon_{\mathrm{osc}}) \quad \textit{for all } \beta\in\mathcal{B},$$

then the aggregate channel $\Lambda_\alpha = \sum_\beta \Lambda_{\alpha|\beta}$ satisfies

$$\|\nabla^{\mathrm{plat}}\Lambda_\alpha\|_\infty \ \le\ |\mathcal{B}|\,\eta\,\varepsilon_{\mathrm{grad}}, \quad \|\mathbf{H}^{\mathrm{plat}}\Lambda_\alpha\|_{\mathrm{op}} \ \le\ |\mathcal{B}|\,\eta\,\varepsilon_{\mathrm{curv}}, \quad \mathrm{osc}_{B_r^{\mathrm{plat}}}\Lambda_\alpha \ \le\ |\mathcal{B}|\,\eta\,\varepsilon_{\mathrm{osc}}.$$

In particular, choosing $\eta \le 1/|\mathcal{B}|$ ensures that Λ_α passes the channelwise plateau test with the original tolerances.

Proof. Linearity gives $D\,\Lambda_\alpha = \sum_\beta D\,\Lambda_{\alpha|\beta}$ for $D \in \{D_\lambda, D_\ell\}$; hence $\|\nabla^{\mathrm{plat}}\Lambda_\alpha\|_\infty = \max\{|D_\lambda\Lambda_\alpha|, |D_\ell\Lambda_\alpha|\} \le \sum_\beta \max\{|D_\lambda\Lambda_{\alpha|\beta}|, |D_\ell\Lambda_{\alpha|\beta}|\} \le |\mathcal{B}|\,\eta\,\varepsilon_{\mathrm{grad}}$. For curvature, $\mathbf{H}^{\mathrm{plat}}\Lambda_\alpha = \sum_\beta \mathbf{H}^{\mathrm{plat}}\Lambda_{\alpha|\beta}$ and the triangle inequality for the operator norm yields the stated bound. For oscillation, for any ball B_r^{plat},

$$\operatorname{osc}_{B_r}\Lambda_\alpha = \sup_{B_r}\sum_\beta \Lambda_{\alpha|\beta} - \inf_{B_r}\sum_\beta \Lambda_{\alpha|\beta} \le \sum_\beta \Big(\sup_{B_r}\Lambda_{\alpha|\beta} - \inf_{B_r}\Lambda_{\alpha|\beta}\Big) = \sum_\beta \operatorname{osc}_{B_r}\Lambda_{\alpha|\beta}.$$

Combine with the sector bounds. □

Definition 7.106 (Coherence coefficients and cancellation veto)**.** To monitor potential inter-sector cancellations, define

$$\chi_\lambda^{(\alpha)} := \frac{\left|\sum_{\beta\in\mathcal{B}} D_\lambda\Lambda_{\alpha|\beta}\right|}{\sum_{\beta\in\mathcal{B}}\left|D_\lambda\Lambda_{\alpha|\beta}\right|}, \qquad \chi_\ell^{(\alpha)} := \frac{\left|\sum_{\beta\in\mathcal{B}} D_\ell\Lambda_{\alpha|\beta}\right|}{\sum_{\beta\in\mathcal{B}}\left|D_\ell\Lambda_{\alpha|\beta}\right|} \in [0,1],$$

with the convention $\chi = 1$ if both numerator and denominator vanish. A *cancellation veto* at (λ,ℓ) requires $\min\{\chi_\lambda^{(\alpha)}, \chi_\ell^{(\alpha)}\} \ge \chi_{\min}$ for some fixed $\chi_{\min} \in (0,1]$.

Proposition 7.107 (Sector compatibility with coherence control)**.** *Under the linearity assumption and sector compatibility* (7.94)*, if additionally* $\min\{\chi_\lambda^{(\alpha)}, \chi_\ell^{(\alpha)}\} \ge \chi_{\min}$*, then:*

(i) (Plateau propagation) Λ_α *passes the channelwise gradient/curvature/oscillation bounds with tolerances inflated by at most the factor* $|\mathcal{B}|$ *as in Proposition 7.105.*

(ii) (Non-degeneracy) *The channel gradients obey*

$$|D_\lambda\Lambda_\alpha| \;\ge\; \chi_{\min}\sum_\beta |D_\lambda\Lambda_{\alpha|\beta}|, \qquad |D_\ell\Lambda_\alpha| \;\ge\; \chi_{\min}\sum_\beta |D_\ell\Lambda_{\alpha|\beta}|,$$

preventing acceptance based on near-perfect cancellations of sizable sector slopes.

Proof. (i) follows from Proposition 7.105. (ii) is immediate from the definition of $\chi_\lambda^{(\alpha)}, \chi_\ell^{(\alpha)}$. □

Corollary 7.108 (Quantitative metric version)**.** *Let* $\mathrm{Slope}_{\alpha|\beta}, \mathrm{Curv}_{\alpha|\beta}, \mathrm{Osc}_{\alpha|\beta}$ *be the pointwise metrics of Definition 7.96 for each sector. Then*

$$\mathrm{Slope}_\alpha \;\le\; \sum_\beta \mathrm{Slope}_{\alpha|\beta}, \qquad \mathrm{Curv}_\alpha \;\le\; \sum_\beta \mathrm{Curv}_{\alpha|\beta}, \qquad \mathrm{Osc}_\alpha(\cdot;r) \;\le\; \sum_\beta \mathrm{Osc}_{\alpha|\beta}(\cdot;r).$$

Hence each metric for the aggregate channel passes with tolerances scaled by at most $|\mathcal{B}|$ *if the sectors pass with a uniform margin factor* $\eta \le 1/|\mathcal{B}|$*.*

Remark 7.109 (Beyond linearity)**.** If $\mathcal{F}_\alpha$ is not strictly linear but is Lipschitz (or subadditive in absolute value) with respect to sectorwise inputs—this includes many monotone convex diagnostics built from band-limited correlators—analogous sector→channel bounds hold with a global Lipschitz constant L_α, i.e. replace the factor $|\mathcal{B}|$ by $L_\alpha\,|\mathcal{B}|$. The BRST and gauge/regularisation robustness statements remain valid by Sections 7.3.6 and 7.3.7.

Integrated visibilities.

Proposition 7.110 (Sector–channel compatibility for integrated visibilities)**.** *Let $\overline{\Lambda}_{\alpha|\beta}$ be the integrated (box-averaged) sector visibilities of Section 7.3.8 built from the window B (cf. Definition 7.39). Then the bounds and implications of Section 7.3.11 continue to hold with $\Lambda_{\alpha|\beta}$ replaced by $\overline{\Lambda}_{\alpha|\beta}$, with the same tolerances whenever $B \subset B_r^{\mathrm{plat}}$ (by eqs. (7.74) and (7.76) and Lemma 7.84) and with explicitly rescaled tolerances otherwise (by Lemmas 7.40 and 7.85 and Corollary 7.41). In particular, under the linearity hypothesis for $\mathcal{F}_\alpha$,*

$$\begin{aligned}
\overline{\Lambda}_\alpha &= \sum_{\beta\in\mathcal{B}} \overline{\Lambda}_{\alpha|\beta},\\
\mathrm{Slope}_\alpha &\le \sum_{\beta} \mathrm{Slope}_{\alpha|\beta},\\
\mathrm{Curv}_\alpha &\le \sum_{\beta} \mathrm{Curv}_{\alpha|\beta},\\
\mathrm{Osc}_\alpha(\cdot;r) &\le \sum_{\beta} \mathrm{Osc}_{\alpha|\beta}(\cdot;r),
\end{aligned}$$

with all quantities interpreted for the integrated visibilities and metric window B. Moreover, coherence coefficients defined with $\overline{\Lambda}_{\alpha|\beta}$ satisfy the same cancellation-veto properties as in Definition 7.106 and Proposition 7.107.

Proof. The derivative commutation and oscillation reduction for box averages are given by eqs. (7.74) and (7.76). When $B \subset B_r^{\mathrm{plat}}$ the sectorwise geometric bounds transfer verbatim by Lemma 7.84; otherwise the rescaled tolerances follow from Lemmas 7.40 and 7.85 and Corollary 7.41. Linearity and subadditivity arguments used in Proposition 7.105 and Corollary 7.108 apply identically to the averaged objects. Coherence statements are purely algebraic in the sector derivatives and thus carry over to the averaged derivatives by eq. (7.74). □

7.3.12 Outcome and Policy Note

Definition 7.111 (Scale-stable operating point)**.** A pair $(\lambda_\star, \ell_\star) \in \mathcal{R}$ is called a *scale-stable operating point* if:

(i) (*Joint plateau*) $(\lambda_\star, \ell_\star)$ lies in a non-empty joint plateau $\mathfrak{P}_{\{\mathrm{law,state}\}}$ as in eq. (7.64) (or its integrated version of Definition 7.86), with quantitative acceptance per Section 7.3.10 and Definition 7.96.

(ii) (*Gauge/BRST robustness*) The on-shell gauge-parameter independence holds by the Nielsen framework Section 7.3.6 and eq. (7.66), and the BRST/Slavnov constraints eqs. (7.16) and (7.18) ensure physical-sector positivity Section 7.2.6 and Corollary 7.24.

(iii) (*Regulator robustness*) The regularisation stability bound eq. (7.67) (sufficiently ensured by Section 7.3.7 and Proposition 7.73) is satisfied on $\mathcal{R}$, so that plateau membership and extracted poles are invariant up to the regulator budget.

(iv) (*Kinematic sanity and corner test*) The tachyon and hyperbolicity conditions eqs. (7.14) and (7.15) and the commutation-of-limits criterion eq. (7.45) hold in a neighbourhood of $(\lambda_\star, \ell_\star)$.

Proposition 7.112 (Consequences of scale stability)**.** *If* $(\lambda_\star, \ell_\star)$ *is scale-stable in the sense of Definition 7.111, then:*

(a) All physical poles extracted at $(\lambda_\star, \ell_\star)$ *are gauge-parameter independent on shell and regulator-robust within the stated budgets (eqs. (7.22) and (7.23), Proposition 7.34, and Corollary 7.74).*

(b) Reported visibilities $\Lambda_\alpha(\lambda_\star, \ell_\star)$ *admit an uncertainty budget combining the plateau tolerances and regulator/swap terms (cf. Definition 7.93 and eq. (7.85)).*

(c) Sector-resolved and joint-channel certifications coincide up to boundary pixels of measure 0 *in the scan grid, and extend to integrated visibilities by Proposition 7.110.*

7.4 Holosymmetric Probe: Controlled Two-path Evolution and Visibilities

This section instruments the holosymmetric probe as an *operational overlay* that preserves quadrant membership while diagnosing law-level versus state-level coherence. The construction acts on the physical sector $\mathcal{H}_{\text{phys}}$ and is compatible with the scale-tagged equations of motion eqs. (6.333), (6.334) and (6.374) and the UV/IR knobs of Section 7.3. All quantities are evaluated *on the band* with $P_x(\lambda_{\text{UV}})$ and within the IR window $K_{\text{IR}}^{(\ell)}$, both BRST-closed eq. (7.30).

Formal definitions and basic properties

Throughout this subsection fix the UV/IR knobs $(\lambda_{\text{UV}}, \ell)$ and work *on band* with $P_x(\lambda_{\text{UV}})$ and $K_{\text{IR}}^{(\ell)}$. All operators act on the physical sector $\mathcal{H}_{\text{phys}}$, implicitly projected by P_{phys} where appropriate. Ancilla qubit states are in the computational basis $\{|0\rangle, |1\rangle\}$ and the time ordering is denoted by $\mathcal{T}$.

Band-limited physical preparation.

$$\rho_{\text{phys}}^{(\lambda_{\text{UV}},\ell)} := \frac{\mathsf{P}_{\text{phys}}\, \rho^{(\lambda_{\text{UV}},\ell)}\, \mathsf{P}_{\text{phys}}}{\text{Tr}\Big(\mathsf{P}_{\text{phys}} \rho^{(\lambda_{\text{UV}},\ell)}\Big)}, \qquad [\hat{Q}_{\text{BRST}}, \rho_{\text{phys}}^{(\lambda_{\text{UV}},\ell)}] = 0, \tag{7.95}$$

i.e. a BRST-closed, band-limited density operator supported on $\mathcal{H}_{\text{phys}}$.

Controlled two-path evolution.

$$U_{\text{ctrl}}^{(\alpha)}(T) := |0\rangle\langle 0| \otimes U_0^{(\alpha)}(T) + |1\rangle\langle 1| \otimes U_1^{(\alpha)}(T), \tag{7.96}$$

for a channel label $\alpha \in \{\text{law}, \text{state}\}$ and branches $j \in \{0, 1\}$ acting on the system conditional on the ancilla.

Band-safety and BRST compatibility of each branch.

$$U_j^{(\alpha)}(T) = \mathcal{T}\exp\Big(-i\int_0^T dt\ \hat{H}_j^{(\alpha)}(t;\lambda_{\rm UV},\ell)\Big), \qquad \big[\hat{Q}_{\rm BRST}, U_j^{(\alpha)}(T)\big] = \zeta X_j^{(\alpha)}, \tag{7.97}$$

with local $X_j^{(\alpha)}$ such that the defect is cohomologically trivial on the physical sector: $\mathsf{P}_{\rm phys}\,\zeta X_j^{(\alpha)}\,\mathsf{P}_{\rm phys} = 0$.

Ancilla readout and complex visibility. After $U_{\rm ctrl}^{(\alpha)}(T)$, apply a Hadamard to the ancilla and measure σ_x, σ_y. On $\mathcal{H}_{\rm phys}$ one obtains

$$\begin{aligned} \langle\sigma_x\rangle_\alpha &= \mathrm{Re}\ \mathcal{V}_\alpha, \qquad \langle\sigma_y\rangle_\alpha = -\,\mathrm{Im}\ \mathcal{V}_\alpha, \\ \mathcal{V}_\alpha(\lambda_{\rm UV},\ell;T) &:= \frac{\mathrm{Tr}_{\mathcal{H}_{\rm phys}}\big[U_1^{(\alpha)}(T)\,\rho_{\rm phys}^{(\lambda_{\rm UV},\ell)}\,U_0^{(\alpha)\dagger}(T)\big]}{\mathrm{Tr}_{\mathcal{H}_{\rm phys}}\rho_{\rm phys}^{(\lambda_{\rm UV},\ell)}}, \end{aligned} \tag{7.98}$$

so that either $\Lambda_\alpha := \mathrm{Re}\ \mathcal{V}_\alpha$ or $|\mathcal{V}_\alpha|$ serves as the (normalised) channel contrast.

Definition 7.113 (Band-limited physical preparation)**.** Fix UV/IR knobs $(\lambda_{\rm UV}, \ell)$ and the physical projector $\mathsf{P}_{\rm phys}$. A *band-limited, BRST-closed physical preparation* is a density operator $\rho_{\rm phys}^{(\lambda_{\rm UV},\ell)}$ on $\mathcal{H}_{\rm phys}$ given by (7.95). It satisfies: (i) positivity and unit trace; (ii) BRST-closure $[\hat{Q}_{\rm BRST}, \rho_{\rm phys}^{(\lambda_{\rm UV},\ell)}] = 0$; and (iii) support on the band, i.e. $\rho_{\rm phys}^{(\lambda_{\rm UV},\ell)} = P_x(\lambda_{\rm UV})\,K_{\rm IR}^{(\ell)}\,\rho_{\rm phys}^{(\lambda_{\rm UV},\ell)}\,K_{\rm IR}^{(\ell)}\,P_x(\lambda_{\rm UV})$.

Definition 7.114 (Quadrant-preserving operational overlay)**.** Let $\mathcal{H}_{\rm phys} = \bigoplus_{q\in\{I,II,III,IV\}}\mathcal{H}_{\rm phys}^{(q)}$ be the quadrant decomposition. A family of controlled evolutions (7.96) with branches (7.97) is a *holosymmetric operational overlay* iff each branch preserves quadrant membership,

$$U_j^{(\alpha)}(T) = \bigoplus_q U_j^{(\alpha,q)}(T), \qquad U_j^{(\alpha,q)}(T):\ \mathcal{H}_{\rm phys}^{(q)} \to \mathcal{H}_{\rm phys}^{(q)},$$

and is band-safe and BRST-compatible as in (7.97) with cohomologically trivial defect $\mathsf{P}_{\rm phys}\zeta X_j^{(\alpha)}\mathsf{P}_{\rm phys} = 0$.

Definition 7.115 (Channel semantics: law vs. state)**.** Fix $\alpha \in \{\text{law}, \text{state}\}$.

(i) **Law channel** ($\alpha = \text{law}$): There exists a reference relational Hamiltonian $\hat{H}_{\rm ref}(t)$ and BRST-closed, band-safe deformations $\delta\hat{H}_j^{(\text{law})}(t)$ such that $\hat{H}_j^{(\text{law})}(t) = \hat{H}_{\rm ref}(t) + \delta\hat{H}_j^{(\text{law})}(t)$, with the preparation fixed to $\rho_{\rm phys}^{(\lambda_{\rm UV},\ell)}$.

(ii) **State channel** ($\alpha = \text{state}$): There exist preparation unitaries S_j (band-safe, BRST-compatible, quadrant-preserving) with $[\hat{Q}_{\rm BRST}, S_j] = \zeta Y_j$ and $\mathsf{P}_{\rm phys}\zeta Y_j\mathsf{P}_{\rm phys} = 0$, such that evolution is governed by a common Hamiltonian $\hat{H}_{\rm ref}(t)$ and the effective branches are $U_j^{(\text{state})}(T) = U_{\rm ref}(T)$ with the *prepared* state $S_j\,\rho_{\rm phys}^{(\lambda_{\rm UV},\ell)}\,S_j^\dagger$.

Both cases fit (7.96)–(7.97); they differ only in where the "difference" resides.

Definition 7.116 (Channel overlap, amplitude and visibility)**.** Given α and T, the *channel-overlap operator* is

$$M_\alpha(T) := \big(U_0^{(\alpha)}(T)\big)^\dagger U_1^{(\alpha)}(T). \tag{7.99}$$

The *holosymmetric amplitude* and *visibility* are

$$\mathcal{A}_\alpha(T;\lambda_{\mathrm{UV}},\ell) := \mathrm{Tr}\Big(\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}\, M_\alpha(T)\Big), \qquad \Lambda_\alpha(T;\lambda_{\mathrm{UV}},\ell) := \Big|\mathcal{A}_\alpha(T;\lambda_{\mathrm{UV}},\ell)\Big|. \tag{7.100}$$

Their *phase* is $\varphi_\alpha(T) = \arg \mathcal{A}_\alpha(T)$.

Definition 7.117 (Ancilla readout family)**.** Let $\sigma_x, \sigma_y, \sigma_z$ be the Pauli operators of the ancilla and $\sigma_\phi := \cos\phi\,\sigma_x + \sin\phi\,\sigma_y$. Prepare the ancilla in $|+\rangle$, apply $U_{\mathrm{ctrl}}^{(\alpha)}(T)$, and trace out the system. The *readout curve* is $R_\alpha(\phi) := \langle \sigma_\phi \rangle_{\mathrm{anc}}$.

Lemma 7.118 (Readout–overlap equivalence)**.** *For any α and T,*

$$\langle \sigma_x \rangle_{\mathrm{anc}} = \mathrm{Re}\ \mathcal{A}_\alpha(T), \qquad \langle \sigma_y \rangle_{\mathrm{anc}} = \mathrm{Im}\ \mathcal{A}_\alpha(T), \qquad R_\alpha(\phi) = \mathrm{Re}\Big(e^{i\phi}\mathcal{A}_\alpha(T)\Big). \tag{7.101}$$

In particular, $\max_\phi R_\alpha(\phi) = \Lambda_\alpha(T)$ *and the fringe phase equals* $\varphi_\alpha(T)$.

Proof. Write the joint state after $U_{\mathrm{ctrl}}^{(\alpha)}(T)$ and trace out the system. The ancilla off-diagonal coherence equals $\mathrm{Tr}\Big(\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)} M_\alpha(T)\Big)$. The Bloch components of the reduced ancilla are then $\langle \sigma_x \rangle = \mathrm{Re}\,\mathcal{A}_\alpha$ and $\langle \sigma_y \rangle = \mathrm{Im}\,\mathcal{A}_\alpha$, yielding (7.101). □

Proposition 7.119 (BRST/Nielsen gauge-parameter independence)**.** *Let the BRST differential act by $\delta_\zeta(\bullet) = [\hat{Q}_{\mathrm{BRST}}, \bullet]$ on bare operators. Under a BRST-exact variation of either branch, $U_j^{(\alpha)} \mapsto U_j^{(\alpha)} + \delta_\zeta U_j^{(\alpha)}$ with $\mathsf{P}_{\mathrm{phys}}\,\delta_\zeta U_j^{(\alpha)}\,\mathsf{P}_{\mathrm{phys}} = 0$, the amplitude (hence the visibility and readout curve) is invariant:*

$$\delta_\zeta \mathcal{A}_\alpha(T) = 0, \qquad \delta_\zeta \Lambda_\alpha(T) = 0, \qquad \delta_\zeta R_\alpha(\phi) = 0. \tag{7.102}$$

Proof. Using cyclicity of the trace and $\mathsf{P}_{\mathrm{phys}}\hat{Q}_{\mathrm{BRST}} = \hat{Q}_{\mathrm{BRST}}\mathsf{P}_{\mathrm{phys}} = 0$ on $\mathcal{H}_{\mathrm{phys}}$, each BRST-exact deformation contributes $\mathrm{Tr}\Big(\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}[\hat{Q}_{\mathrm{BRST}}, \cdot]\Big) = 0$. Thus $\delta_\zeta \mathrm{Tr}(\rho_{\mathrm{phys}} M_\alpha) = 0$, implying (7.66). □

Proposition 7.120 (Quadrant additivity)**.** *Assume Definition 7.114. Let $w_q := \mathrm{Tr}\Big(\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}\Pi^{(q)}\Big)$ with $\Pi^{(q)}$ the projector onto $\mathcal{H}_{\mathrm{phys}}^{(q)}$ and $\sum_q w_q = 1$. Then*

$$\mathcal{A}_\alpha(T) = \sum_q w_q\, \mathcal{A}_\alpha^{(q)}(T), \qquad \mathcal{A}_\alpha^{(q)}(T) := \frac{\mathrm{Tr}\Big(\Pi^{(q)}\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}\Pi^{(q)}\, M_\alpha^{(q)}(T)\Big)}{w_q}, \tag{7.103}$$

with $M_\alpha^{(q)} = (U_0^{(\alpha,q)})^\dagger U_1^{(\alpha,q)}$. No spurious cross-quadrant interference occurs.

Definition 7.121 (Scale-tagged and integrated visibilities)**.** At fixed $(\lambda_{\mathrm{UV}}, \ell)$ define $\Lambda_\alpha(T; \lambda_{\mathrm{UV}}, \ell)$ as in (7.100). For a box/window B in $(\lambda_{\mathrm{UV}}, \ell, T)$-space, the *integrated visibility* is

$$\overline{\Lambda}_{\alpha|B} := \frac{1}{\mu(B)}\int_B d\lambda_{\mathrm{UV}}\, d\ell\, dT\ \Lambda_\alpha(T;\lambda_{\mathrm{UV}},\ell), \tag{7.104}$$

with $\mu(B)$ the product measure. Its log-scale derivatives and oscillation metrics are defined componentwise and inherit the commutation and bound properties established earlier for visibilities.

Definition 7.122 (Certified operating point for the probe)**.** A pair $(\lambda_\star, \ell_\star)$ is a *scale-stable operating point* for channel α at interrogation time $T_\star$ if: (i) $(\lambda_\star, \ell_\star)$ lies on a joint plateau of $\Lambda_\alpha(T_\star; \lambda_{\mathrm{UV}}, \ell)$ (geometric and metric criteria coincide); (ii) BRST/Nielsen stability (7.66) holds; and (iii) regulator robustness under admissible kernel/projector perturbations is satisfied.

Remark 7.123 (Law–state discrimination)**.** Define $\Delta\Lambda(T) := \Lambda_{\mathrm{law}}(T) - \Lambda_{\mathrm{state}}(T)$ at fixed $(\lambda_{\mathrm{UV}}, \ell)$. On any certified plateau neighborhood, the sign and magnitude of $\Delta\Lambda$ are physically meaningful and insensitive to gauge/regulator choices by Propositions 7.119–7.120 and Definition 7.122.

Definition 7.124 (Echo)**.** The *holosymmetric echo* is $\mathcal{E}_\alpha(T) := \big|\mathcal{A}_\alpha(T)\big|^2 = \Lambda_\alpha(T)^2$. It coincides with the ancilla coherence length squared and admits the same certification criteria as in Definition 7.122.

Formal channel semantics and complex visibility

Definition 7.125 (Channel semantics: law vs. state)**.** Fix $(\lambda_{\mathrm{UV}}, \ell)$, a band/BRST-safe time $T > 0$, and a physical preparation $\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}$ as in (7.95).

(i) **Law channel** ($\alpha = \mathrm{law}$): The physical state is held fixed at $\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}$ while the two branches use BRST-compatible Hamiltonians that differ only by admissible toggles within the symmetric law, i.e.

$$\hat{H}_0^{(\mathrm{law})}(t) \sim \hat{H}_1^{(\mathrm{law})}(t), \qquad U_j^{(\mathrm{law})}(T) = \mathcal{T}\exp\Big(-i\int_0^T dt\, \hat{H}_j^{(\mathrm{law})}(t)\Big),$$

with $[\hat{Q}_{\mathrm{BRST}}, U_j^{(\mathrm{law})}] = \zeta X_j^{(\mathrm{law})}$ and $\mathsf{P}_{\mathrm{phys}}\zeta X_j^{(\mathrm{law})}\mathsf{P}_{\mathrm{phys}} = 0$. This diagnoses *law-level coherence* without altering state-level symmetry.

(ii) **State channel** ($\alpha = \mathrm{state}$): A common BRST-compatible Hamiltonian $\hat{H}^{(\mathrm{state})}(t)$ is used, while two BRST-closed initial states ρ_0 and ρ_1 are prepared from admissible (gauge-invariant, band-safe) preparations. Equivalently, one may *push* the preparations into the branches and write

$$U_{0,1}^{(\mathrm{state})}(T) = U(T)\, W_{0,1},$$

with $U(T) = \mathcal{T}\exp\big(-i\int_0^T dt\, \hat{H}^{(\mathrm{state})}(t)\big)$, and W_j band/BRST-safe, so that a single fiducial $\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}$ is used while the *difference* resides in $W_{0,1}$. This diagnoses *state-level coherence*.

Lemma 7.126 (Preparation/branch equivalence for the state channel)**.** *Let $\rho_j = W_j\, \rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}\, W_j^\dagger$ with W_j band/BRST-safe and $U_0 = U_1 = U(T)$. Then the interferometer defined with distinct inputs (ρ_0, ρ_1) and common $U(T)$ is equivalent to the fixed-input interferometer with branches $U_j^{(\mathrm{state})}(T) = U(T)W_j$ acting on $\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}$. Consequently,are related by cyclicity of the trace:*

$$\begin{aligned} &\mathrm{Tr}\Big[U(T)\,\rho_1\,U^\dagger(T)\Big] \\ \textit{vs.}\ \ &\mathrm{Tr}\Big[U_1^{(\mathrm{state})}(T)\,\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}\,U_0^{(\mathrm{state})\dagger}(T)\Big] \end{aligned} \tag{7.105}$$

Proof. Work on $\mathcal{H}_{\text{anc}} \otimes \mathcal{H}_{\text{phys}}$ with the ancilla initialised in $|+\rangle$ and the system in $\rho_{\text{phys}}^{(\lambda_{\text{UV}},\ell)}$. Assume W_j are band/BRST-safe so that all operations remain in the physical sector.

Step 1 (two equivalent circuit factorizations). Define the controlled "preparation" unitary

$$\mathsf{C}(W) := |0\rangle\langle 0| \otimes W_0 \;+\; |1\rangle\langle 1| \otimes W_1,$$

and the common evolution $U(T)$ on the system. Then

$$\Big(|0\rangle\langle 0| \otimes U(T) + |1\rangle\langle 1| \otimes U(T) \Big)\, \mathsf{C}(W) = |0\rangle\langle 0| \otimes U(T)W_0 \;+\; |1\rangle\langle 1| \otimes U(T)W_1,$$

i.e. the composition "controlled preparation then common $U(T)$" equals a single controlled evolution with branches $U_j^{(\text{state})}(T) = U(T)W_j$ as in the statement. Thus, replacing distinct inputs $\rho_j = W_j \rho_{\text{phys}}^{(\lambda_{\text{UV}},\ell)} W_j^\dagger$ and common $U(T)$ by a fixed input $\rho_{\text{phys}}^{(\lambda_{\text{UV}},\ell)}$ and branches $U_j^{(\text{state})}(T) = U(T)W_j$ produces the same joint unitary on ancilla⊗system and hence the same ancilla reduced state for all measurements.

Step 2 (complex visibility equality). Using the definition (7.98) with α = state and $U_j^{(\text{state})}(T) = U(T)W_j$,

$$\mathcal{V}_{\text{state}} = \frac{\mathrm{Tr}_{\mathcal{H}_{\text{phys}}}\Big[U(T)W_1\, \rho_{\text{phys}}^{(\lambda_{\text{UV}},\ell)}\, W_0^\dagger U^\dagger(T)\Big]}{\mathrm{Tr}_{\mathcal{H}_{\text{phys}}}\, \rho_{\text{phys}}^{(\lambda_{\text{UV}},\ell)}}.$$

By cyclicity of the trace and unitarity of $U(T)$ this reduces to

$$\mathcal{V}_{\text{state}} = \frac{\mathrm{Tr}\Big[W_1\, \rho_{\text{phys}}^{(\lambda_{\text{UV}},\ell)}\, W_0^\dagger\Big]}{\mathrm{Tr}\, \rho_{\text{phys}}^{(\lambda_{\text{UV}},\ell)}}.$$

Equivalently, if one insists on describing the procedure as "distinct inputs with common $U(T)$," then the numerator appearing in (7.98) for the branch-realisation is related, again by cyclicity, to the branch-free form:

$$\mathrm{Tr}\Big[U_1^{(\text{state})}(T)\, \rho_{\text{phys}}^{(\lambda_{\text{UV}},\ell)}\, U_0^{(\text{state})\dagger}(T)\Big] = \mathrm{Tr}\Big[U(T)\, W_1\, \rho_{\text{phys}}^{(\lambda_{\text{UV}},\ell)}\, W_0^\dagger U^\dagger(T)\Big] = \mathrm{Tr}\Big[W_1\, \rho_{\text{phys}}^{(\lambda_{\text{UV}},\ell)}\, W_0^\dagger\Big].$$

This proves that the interferometric readouts (hence $\langle\sigma_{x,y}\rangle$ and $\mathcal{V}_{\text{state}}$) coincide in the two descriptions.

Step 3 (relation to the two-prepared-inputs variant). If one instead prepares $\rho_0 = W_0 \rho W_0^\dagger$ and $\rho_1 = W_1 \rho W_1^\dagger$ and uses the common $U(T)$ in the "direct-overlap" variant, the corresponding amplitude is proportional to the Hilbert–Schmidt overlap $\mathrm{Tr}(\rho_1 \rho_0)$; for pure $\rho = |\psi\rangle\langle\psi|$ this gives $\mathcal{V}_{\text{state}} = \langle\psi_0\rangle\, \psi_1$ and $|\mathcal{V}_{\text{state}}|^2 = \mathrm{Tr}(\rho_1\rho_0)$.

Hence the two implementations are operationally equivalent for the state channel. □

Definition 7.127 (Complex visibility and readout)**.** After the controlled evolution $U_{\text{ctrl}}^{(\alpha)}(T)$, apply a Hadamard on the ancilla and measure σ_x, σ_y. Define the *complex visibility*

$$\mathcal{V}_\alpha(\lambda_{\text{UV}}, \ell; T) := \frac{\mathrm{Tr}_{\mathcal{H}_{\text{phys}}}\Big(U_1^{(\alpha)}(T)\, \rho_{\text{phys}}^{(\lambda_{\text{UV}},\ell)}\, U_0^{(\alpha)\dagger}(T)\Big)}{\mathrm{Tr}_{\mathcal{H}_{\text{phys}}} \rho_{\text{phys}}^{(\lambda_{\text{UV}},\ell)}}. \tag{7.106}$$

The *channel contrast* may be taken as $\Lambda_\alpha := \operatorname{Re}\mathcal{V}_\alpha$ or $|\mathcal{V}_\alpha|$ (both $\in [-1,1]$ under the normalisation above). The ancilla expectations satisfy

$$\langle\sigma_x\rangle_\alpha = \operatorname{Re}\,\mathcal{V}_\alpha, \qquad \langle\sigma_y\rangle_\alpha = -\operatorname{Im}\,\mathcal{V}_\alpha. \tag{7.107}$$

Proposition 7.128 (Bounds and normalisation)**.** *Let $M_\alpha := (U_0^{(\alpha)})^\dagger U_1^{(\alpha)}$ on $\mathcal{H}_{\text{phys}}$. Then M_α is unitary and*

$$|\mathcal{V}_\alpha| = \frac{|\operatorname{Tr}\big(\rho_{\text{phys}}^{(\lambda_{\text{UV}},\ell)} M_\alpha\big)|}{\operatorname{Tr}\big(\rho_{\text{phys}}^{(\lambda_{\text{UV}},\ell)}\big)} \le 1. \tag{7.108}$$

Hence $\operatorname{Re}\mathcal{V}_\alpha \in [-1,1]$ *and* $|\mathcal{V}_\alpha| \in [0,1]$. *If $\rho_{\text{phys}}^{(\lambda_{\text{UV}},\ell)}$ is trace-one, the denominator in (7.98) equals* 1.

Proof. Unitarity. Since each branch $U_j^{(\alpha)}$ is unitary on $\mathcal{H}_{\text{phys}}$,

$$M_\alpha^\dagger M_\alpha = \big((U_1^{(\alpha)})^\dagger U_0^{(\alpha)}\big)\big((U_0^{(\alpha)})^\dagger U_1^{(\alpha)}\big) = (U_1^{(\alpha)})^\dagger\,\mathbb{I}\,U_1^{(\alpha)} = \mathbb{I},$$

and similarly $M_\alpha M_\alpha^\dagger = \mathbb{I}$.

Bound. Let $\rho := \rho_{\text{phys}}^{(\lambda_{\text{UV}},\ell)} \succeq 0$ with finite trace. Using Hölder (or the operator Cauchy–Schwarz) and $\|M_\alpha\|_\infty = 1$ for unitary M_α,

$$|\operatorname{Tr}(\rho M_\alpha)| = |\operatorname{Tr}\big(\rho^{1/2} M_\alpha \rho^{1/2}\big)| \le \|\rho^{1/2}\|_2^2\,\|M_\alpha\|_\infty = \operatorname{Tr}(\rho).$$

Dividing by $\operatorname{Tr}(\rho)$ yields $|\mathcal{V}_\alpha| \le 1$. Consequently, $-|\mathcal{V}_\alpha| \le \operatorname{Re}\mathcal{V}_\alpha \le |\mathcal{V}_\alpha|$ implies $\operatorname{Re}\mathcal{V}_\alpha \in [-1,1]$. If $\operatorname{Tr}(\rho) = 1$ (trace-one preparation), the denominator in (7.98) is 1. □

Corollary 7.129 (State-channel specialisations)**.** *In the state channel with $U_j^{(\text{state})} = UW_j$ and fixed $\rho_{\text{phys}}^{(\lambda_{\text{UV}},\ell)}$,*

$$\mathcal{V}_{\text{state}} = \frac{\operatorname{Tr}\big(W_1\,\rho_{\text{phys}}^{(\lambda_{\text{UV}},\ell)}\,W_0^\dagger\big)}{\operatorname{Tr}\big(\rho_{\text{phys}}^{(\lambda_{\text{UV}},\ell)}\big)}. \tag{7.109}$$

(a) If $\rho_{\text{phys}}^{(\lambda_{\text{UV}},\ell)} = |\psi\rangle\langle\psi|$ and $W_j|\psi\rangle = |\psi_j\rangle$ (unitary preparations), then $\mathcal{V}_{\text{state}} = \langle\psi_0\rangle\,\psi_1$ and the echo $|\mathcal{V}_{\text{state}}|^2$ *equals the (pure-state) fidelity.*

(b) If one realises the "two-prepared-inputs" variant with initial states ρ_0, ρ_1 and a common $U(T)$, then under the standard direct-overlap construction (cf. [287, 444, 427]) the measured amplitude is proportional to the Hilbert–Schmidt overlap $\operatorname{Tr}(\rho_1\rho_0)$; exact equality follows for the canonical normalisation used in those protocols.

Proof. Insert $U_j^{(\text{state})} = UW_j$ into the definition of $\mathcal{V}_\alpha$ in (7.98) (with $\alpha = \text{state}$) and use cyclicity of the trace plus unitarity of U:

$$\mathcal{V}_{\text{state}} = \frac{\operatorname{Tr}\big(UW_1\,\rho_{\text{phys}}^{(\lambda_{\text{UV}},\ell)}\,W_0^\dagger U^\dagger\big)}{\operatorname{Tr}\big(\rho_{\text{phys}}^{(\lambda_{\text{UV}},\ell)}\big)} = \frac{\operatorname{Tr}\big(W_1\,\rho_{\text{phys}}^{(\lambda_{\text{UV}},\ell)}\,W_0^\dagger\big)}{\operatorname{Tr}\big(\rho_{\text{phys}}^{(\lambda_{\text{UV}},\ell)}\big)},$$

which is (7.109). For (a), take $\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)} = |\psi\rangle\langle\psi|$; then $W_j\,|\psi\rangle = |\psi_j\rangle$ gives $\mathcal{V}_{\mathrm{state}} = \langle\psi_0\rangle\,\psi_1$ and $|\mathcal{V}_{\mathrm{state}}|^2$ is the pure-state fidelity. For (b), the standard controlled-unitary "direct overlap" circuit with inputs (ρ_0, ρ_1) and common U produces an ancilla amplitude proportional to $\mathrm{Tr}(\rho_1\rho_0)$; choosing the canonical normalisation in those protocols yields equality (see the cited references). □

Proposition 7.130 (BRST/Nielsen invariance of $\mathcal{V}_\alpha$)**.** *Under BRST-exact deformations $U_j^{(\alpha)} \mapsto U_j^{(\alpha)} + \delta_\zeta U_j^{(\alpha)}$ with $\mathsf{P}_{\mathrm{phys}}\delta_\zeta U_j^{(\alpha)}\mathsf{P}_{\mathrm{phys}} = 0$ and $[\hat{Q}_{\mathrm{BRST}}, \rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}] = 0$, one has*

$$\delta_\zeta \mathcal{V}_\alpha = 0, \qquad \delta_\zeta \Lambda_\alpha = 0. \tag{7.110}$$

Proof. Write $\rho := \rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}$ and recall $\rho = \mathsf{P}_{\mathrm{phys}}\rho\,\mathsf{P}_{\mathrm{phys}}$ from (7.95). From (7.98),

$$\mathcal{V}_\alpha = \frac{\mathrm{Tr}\big(U_1^{(\alpha)}\,\rho\,U_0^{(\alpha)\dagger}\big)}{\mathrm{Tr}(\rho)}.$$

The denominator is BRST-invariant by $[\hat{Q}_{\mathrm{BRST}}, \rho] = 0$. For the numerator $N_\alpha := \mathrm{Tr}\big(U_1^{(\alpha)}\rho\,U_0^{(\alpha)\dagger}\big)$,

$$\delta_\zeta N_\alpha = \mathrm{Tr}\Big((\delta_\zeta U_1^{(\alpha)})\,\rho\,U_0^{(\alpha)\dagger}\Big) + \mathrm{Tr}\Big(U_1^{(\alpha)}\,\rho\,\delta_\zeta U_0^{(\alpha)\dagger}\Big).$$

Insert $\mathsf{P}_{\mathrm{phys}}$ identities (since $\rho = \mathsf{P}\rho\,\mathsf{P}$ and all operators act on $\mathcal{H}_{\mathrm{phys}}$):

$$\delta_\zeta N_\alpha = \mathrm{Tr}\Big(\mathsf{P}\,\delta_\zeta U_1^{(\alpha)}\,\mathsf{P}\,\rho\,\mathsf{P}\,U_0^{(\alpha)\dagger}\,\mathsf{P}\Big) + \mathrm{Tr}\Big(\mathsf{P}\,U_1^{(\alpha)}\,\mathsf{P}\,\rho\,\mathsf{P}\,\delta_\zeta U_0^{(\alpha)\dagger}\,\mathsf{P}\Big) = 0,$$

because the deformations are cohomologically trivial on the physical sector, $\mathsf{P}\,\delta_\zeta U_j^{(\alpha)}\,\mathsf{P} = 0$. Hence $\delta_\zeta N_\alpha = 0$ and $\delta_\zeta \mathcal{V}_\alpha = 0$. Since Λ_α is either $\mathrm{Re}\,\mathcal{V}_\alpha$ or $|\mathcal{V}_\alpha|$, it follows that $\delta_\zeta\Lambda_\alpha = 0$. □

Remark 7.131 (Operational interpretation)**.** For $\alpha = \mathrm{law}$, $\mathcal{V}_{\mathrm{law}}$ probes symmetry-preserving toggles of the relational dynamics; for $\alpha = \mathrm{state}$, $\mathcal{V}_{\mathrm{state}}$ probes coherence between admissibly prepared physical states. In both cases, the single-qubit meter with a final Hadamard realises a direct estimator of the complex overlap in (7.98), consistent with controlled-unitary interferometry and Loschmidt/fidelity-amplitude protocols [287, 444, 427].

Proposition 7.132 (Gauge/Nielsen protection and BRST robustness)**.** *Assume the Nielsen master identity eq. (7.19), BRST closure of the band knobs and preparation ($[\hat{Q}_{\mathrm{BRST}}, P_x(\lambda_{\mathrm{UV}})] = [\hat{Q}_{\mathrm{BRST}}, K_{\mathrm{IR}}^{(\ell)}] = 0$ by eq. (7.30) and $[\hat{Q}_{\mathrm{BRST}}, \rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}] = 0$ by (7.95)), and ghost decoupling/-positivity Section 7.2.6 and eq. (7.12). Then, differentiating (7.98) w.r.t. the gauge-fixing parameter α_{gf} one finds*

$$\partial_{\alpha_{\mathrm{gf}}}\mathcal{V}_\alpha = \frac{\mathrm{Tr}\big(\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}\,\mathsf{O}_{\zeta\text{-}exact}\big)}{\mathrm{Tr}\,\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}} + \frac{\mathrm{Tr}\big(\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}\,\mathsf{O}_{\mathrm{EOM}}\big)}{\mathrm{Tr}\,\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}}, \tag{7.111}$$

where $\mathsf{O}_{\zeta\text{-}exact}$ is BRST-exact and $\mathsf{O}_{\mathrm{EOM}}$ is proportional to the scale-tagged equations of motion. On the physical Hilbert space, the ζ-exact insertions decouple by quartet/positivity Section 7.2.6 and eq. (7.12), and the EOM terms vanish on shell by eqs. (6.333) and (6.334). Hence

$$\partial_{\alpha_{\mathrm{gf}}}\mathcal{V}_\alpha = 0 \quad \textit{on shell}, \qquad \partial_{\alpha_{\mathrm{gf}}}\Lambda_\alpha = 0 \quad \textit{on shell}. \tag{7.112}$$

Moreover, the BRST closure of $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ ensures functorial descent of the construction to $\mathcal{H}_{\mathrm{phys}}$ eq. (7.30), preserving unitarity and positivity [82, 241].

Proof. Differentiate (7.98). By eq. (7.19), the α_{gf}–derivative of any gauge-dependent operator insertable in the trace can be written as a sum of a BRST-exact commutator and EOM terms. Insert $\mathsf{P}_{\mathrm{phys}}$ on both sides of the trace and use $[\hat{Q}_{\mathrm{BRST}}, \rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}] = 0$ and eq. (7.30) to move $\hat{Q}_{\mathrm{BRST}}$ off physical matrix elements; the ζ-exact contributions vanish by quartet/positivity Section 7.2.6 and eq. (7.12), while EOM terms vanish when eqs. (6.333) and (6.334) hold. The statement for Λ_α follows since it is a real functional of $\mathcal{V}_\alpha$. □

Proposition 7.133 (Band safety and UV/IR dependence)**.** *Let $\mathcal{V}_\alpha$ be given by (7.98) and let the abstract visibility functional $\mathcal{F}_\alpha$ be as in eq. (7.54). Then the only explicit dependence of $\mathcal{V}_\alpha$ on $(\lambda_{\mathrm{UV}}, \ell)$ enters via the band projectors $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ (cf. Section 7.3.1). In particular, for any differentiability point of the knobs,*

$$\partial_{\lambda_{\mathrm{UV}}} \mathcal{V}_\alpha = \frac{\mathrm{Tr}\Big((\partial_{\lambda_{\mathrm{UV}}} P_x)\, \mathcal{K}_\alpha\Big)}{\mathrm{Tr}\Big(\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}\Big)} - \frac{\mathrm{Tr}\Big(\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}\, \mathcal{K}_\alpha\Big)}{\mathrm{Tr}\Big(\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}\Big)^2}\, \mathrm{Tr}\Big((\partial_{\lambda_{\mathrm{UV}}} P_x)\, \rho^{(\lambda_{\mathrm{UV}},\ell)}\Big), \tag{7.113}$$

and analogously for $\partial_\ell \mathcal{V}_\alpha$ with $\partial_\ell K_{\mathrm{IR}}^{(\ell)}$, where $\mathcal{K}_\alpha := \Big(U_1^{(\alpha)}(T)\, P_x K_{\mathrm{IR}}^{(\ell)}\Big)\, \rho^{(\lambda_{\mathrm{UV}},\ell)}\, \Big(P_x K_{\mathrm{IR}}^{(\ell)} U_0^{(\alpha)\dagger}(T)\Big)$ is a band-safe trace-class operator. Consequently, plateau certification and regulator robustness are governed by Sections 7.3 and 7.3.7.

Proof. By construction, all system operators entering (7.98) are band-limited by left/right multiplication with $P_x(\lambda_{\mathrm{UV}}) K_{\mathrm{IR}}^{(\ell)}$ already built into $\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}$ and, if needed, into the branch evolutions; see Section 7.3.1. Treating the trace as a Gateaux-differentiable functional in the finite-energy domain and using Leibniz plus cyclicity gives the stated formulas. Band-safety follows since $\partial_{\lambda_{\mathrm{UV}}} P_x$ and $\partial_\ell K_{\mathrm{IR}}^{(\ell)}$ are bounded (in the admissible class) and appear only linearly inside a trace against a trace-class operator. The plateau and robustness claims then reduce to the corresponding criteria and stability results in Sections 7.3 and 7.3.7. □

Proposition 7.134 (Quadrant preservation)**.** *Assume the quadrant decomposition $\mathcal{H}_{\mathrm{phys}} = \bigoplus_q \mathcal{H}_{\mathrm{phys}}^{(q)}$ and the holosymmetric overlay of Definition 7.114. If the inter-branch differences in (7.97) are cohomologically trivial (or symmetry-preserving) on $\mathcal{H}_{\mathrm{phys}}$, then the probe preserves quadrant membership:*

$$U_j^{(\alpha)}(T) = \bigoplus_q U_j^{(\alpha,q)}(T), \qquad M_\alpha(T) = \bigoplus_q M_\alpha^{(q)}(T). \tag{7.114}$$

Hence there is no cross-quadrant interference, and the amplitude decomposes as in eq. (7.103). In Quadrant I, certified plateaus predict large Λ_{law} and Λ_{state}; deviations signal genuine physical decoherence or symmetry-allowed IR effects rather than gauge artefacts (cf. the diagnostic metrics in Sections 7.3.2 and 7.3.4).

Proof. Cohomologically trivial (or symmetry-preserving) deformations commute with the quadrant projectors on $\mathcal{H}_{\mathrm{phys}}$, hence each branch is block-diagonal in q and so is $M_\alpha = (U_0^{(\alpha)})^\dagger U_1^{(\alpha)}$. Taking traces with $\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}$ that is itself block-diagonal yields eq. (7.103). The final claim follows by applying the plateau certification and robustness results to each block and using that gauge artefacts have been excluded by BRST/Nielsen protection. □

7.4.1 Readout and Complex Coherence Amplitude

Definition 7.135 (Complex coherence amplitude)**.** After $U_{\mathrm{ctrl}}^{(\alpha)}(T)$, define the *complex coherence amplitude*

$$\mathcal{C}_\alpha(\lambda_{\mathrm{UV}},\ell;T) := \frac{\mathrm{Tr}\Big(\mathsf{P}_{\mathrm{phys}}\,U_1^{(\alpha)}(T)\,\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}\,U_0^{(\alpha)\dagger}(T)\,\mathsf{P}_{\mathrm{phys}}\Big)}{\mathrm{Tr}\Big(\mathsf{P}_{\mathrm{phys}}\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}\Big)}. \tag{7.115}$$

Lemma 7.136 (Ancilla readout relations)**.** *Measuring the ancilla Pauli operators directly after $U_{\mathrm{ctrl}}^{(\alpha)}(T)$ (no additional rotations) yields*

$$\langle\sigma_x\rangle_\alpha = \mathrm{Re}\;\mathcal{C}_\alpha, \qquad \langle\sigma_y\rangle_\alpha = \mathrm{Im}\;\mathcal{C}_\alpha, \qquad \Rightarrow \quad |\mathcal{C}_\alpha| = \sqrt{\langle\sigma_x\rangle_\alpha^2 + \langle\sigma_y\rangle_\alpha^2}. \tag{7.116}$$

Proof. Let the system+ancilla initial state be $\rho_{\mathrm{anc}} = |+\rangle\langle+|$ and $\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}$ on $\mathcal{H}_{\mathrm{phys}}$. Propagating by $U_{\mathrm{ctrl}}^{(\alpha)}(T)$ and tracing out the system gives the ancilla reduced density matrix with off-diagonal element $\rho_{\mathrm{anc}}^{01} = \mathrm{Tr}\Big(\mathsf{P}_{\mathrm{phys}}U_1^{(\alpha)}\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}U_0^{(\alpha)\dagger}\mathsf{P}_{\mathrm{phys}}\Big)\Big/\,\mathrm{Tr}\Big(\mathsf{P}_{\mathrm{phys}}\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}\Big) = \mathcal{C}_\alpha$. Hence the Bloch components satisfy $\langle\sigma_x\rangle = \mathrm{Re}\;\mathcal{C}_\alpha$ and $\langle\sigma_y\rangle = \mathrm{Im}\;\mathcal{C}_\alpha$, giving (7.116). □

Definition 7.137 (Signed visibility)**.** Fix a reference phase $\phi_\alpha \in \mathbb{R}$. The *signed visibility* is

$$\Lambda_\alpha(\lambda_{\mathrm{UV}},\ell;T) := \mathrm{Re}\Big(e^{-i\phi_\alpha}\,\mathcal{C}_\alpha(\lambda_{\mathrm{UV}},\ell;T)\Big), \qquad -1 \le \Lambda_\alpha \le 1. \tag{7.117}$$

Choosing $\phi_\alpha = \arg\mathcal{C}_\alpha$ maximises Λ_α at $|\mathcal{C}_\alpha|$.

Proposition 7.138 (Equivalence to the complex visibility on $\mathcal{H}_{\mathrm{phys}}$)**.** *Assume (i) $\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)} = \mathsf{P}_{\mathrm{phys}}\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}\mathsf{P}_{\mathrm{phys}}$ as in (7.95), and (ii) the branches preserve the physical sector in the sense of (7.97) (cohomologically trivial defect on $\mathcal{H}_{\mathrm{phys}}$). Then $\mathcal{C}_\alpha$ equals the complex visibility $\mathcal{V}_\alpha$ of (7.98):*

$$\mathcal{C}_\alpha(\lambda_{\mathrm{UV}},\ell;T) = \mathcal{V}_\alpha(\lambda_{\mathrm{UV}},\ell;T).$$

Proof. Insert $\mathsf{P}_{\mathrm{phys}}$ on both sides of the numerator in (7.115) and use $\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)} = \mathsf{P}\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}\mathsf{P}$ with $\mathsf{P} := \mathsf{P}_{\mathrm{phys}}$. By hypothesis (ii), the action of $U_j^{(\alpha)}(T)$ on $\mathcal{H}_{\mathrm{phys}}$ is equivalent to its projection, so $\mathsf{P}U_j^{(\alpha)}(T)\mathsf{P}$ can be replaced by $U_j^{(\alpha)}(T)$ inside physical matrix elements. Thus the numerator in (7.115) equals that of (7.98), and the denominators coincide by (7.95). □

Proposition 7.139 (Bounds and normalisation for $\mathcal{C}_\alpha$)**.** *Let $M_\alpha := (U_0^{(\alpha)})^\dagger U_1^{(\alpha)}$ and $\rho := \rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}$. Then*

$$|\mathcal{C}_\alpha| = \frac{|\,\mathrm{Tr}(\mathsf{P}_{\mathrm{phys}}\rho\,M_\alpha)|}{\mathrm{Tr}(\mathsf{P}_{\mathrm{phys}}\rho)} \le 1,$$

whence $\mathrm{Re}\;\mathcal{C}_\alpha \in [-1,1]$ *and* $|\mathcal{C}_\alpha| \in [0,1]$. *If* $\mathrm{Tr}(\rho) = 1$, *the denominator in* (7.115) *equals* 1.

Proof. M_α is unitary; apply the same Hölder/Cauchy–Schwarz argument as in Proposition 7.128, now with the explicit $\mathsf{P}_{\mathrm{phys}}$ insertions (which do not increase the operator norm and preserve trace-class structure). □

Remark 7.140 (Phase/readout convention)**.** Equation (7.116) uses the "no post-rotation" convention (measure $\sigma_{x,y}$ directly after U_{ctrl}), giving $\langle\sigma_y\rangle = \mathrm{Im}\;\mathcal{C}_\alpha$. If instead a Hadamard is applied before readout (as elsewhere in the chapter), the y-component acquires a minus sign, yielding $\langle\sigma_y\rangle = -\,\mathrm{Im}\;\mathcal{V}_\alpha$ for the visibility form (7.98). Both conventions are equivalent up to a fixed phase choice ϕ_α in (7.117).

Derivation of the readout (one-qubit interferometer).

Lemma 7.141 (Derivation of the readout (one-qubit interferometer))**.** *Let the ancilla be prepared in* $|+\rangle$ *and the system in* ρ_{phys} *(shorthand for* $\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}$*). Under the controlled evolution* (7.96) *followed by a Hadamard on the ancilla, one has*

$$\langle\sigma_x\rangle_\alpha = \frac{\mathrm{Tr}\big(U_1^{(\alpha)}\rho_{\mathrm{phys}}U_0^{(\alpha)\dagger}\big)}{\mathrm{Tr}\,\rho_{\mathrm{phys}}}, \qquad \langle\sigma_y\rangle_\alpha = \frac{\mathrm{Tr}\big(-i\,U_1^{(\alpha)}\rho_{\mathrm{phys}}U_0^{(\alpha)\dagger}\big)}{\mathrm{Tr}\,\rho_{\mathrm{phys}}},$$

which is equivalent to the interferometric relations (7.116)*.*

Proof. Evolve $|+\rangle\langle+|\otimes\rho_{\mathrm{phys}}$ by $U_{\mathrm{ctrl}}^{(\alpha)}(T)$, apply a Hadamard on the ancilla, and trace out the system. The off-diagonal entry of the reduced ancilla equals $\mathrm{Tr}(U_1^{(\alpha)}\rho_{\mathrm{phys}}U_0^{(\alpha)\dagger})/\,\mathrm{Tr}\,\rho_{\mathrm{phys}}$. Identifying the Bloch components gives the displayed formulas. Using $\langle\sigma_x\rangle = \mathrm{Re}(\cdot)$ and $\langle\sigma_y\rangle = -\,\mathrm{Im}(\cdot)$ yields (7.116). □

Proposition 7.142 (Bounds and extremal cases)**.** *Let* $\mathcal{C}_\alpha$ *be as in* (7.115) *and* Λ_α *as in* (7.117)*. Then*

$$|\mathcal{C}_\alpha| \le 1, \qquad |\Lambda_\alpha| \le 1, \tag{7.118}$$

with equality iff $U_1^{(\alpha)}(T)\rho_{\mathrm{phys}}U_0^{(\alpha)\dagger}(T)$ *is a positive multiple of* ρ_{phys} *(*pure state case: $U_1^{(\alpha)}(T)\,|\psi\rangle = e^{i\varphi}U_0^{(\alpha)}(T)\,|\psi\rangle$*). For the* state channel *with a common* $U(T)$ *and preparations* $\rho_j = W_j\rho W_j^\dagger$*, one has* $\mathcal{C}_{\mathrm{state}} \propto \mathrm{Tr}(\rho_1\rho_0)$*; for pure preparations this reduces to* $\langle\psi_1|\psi_0\rangle$*.*

Proof. Write $M_\alpha := (U_0^{(\alpha)})^\dagger U_1^{(\alpha)}$, so $|\mathcal{C}_\alpha| = |\,\mathrm{Tr}(\rho_{\mathrm{phys}}M_\alpha)|/\,\mathrm{Tr}\,\rho_{\mathrm{phys}}$ and $\|M_\alpha\|_\infty = 1$ (unitary). By Cauchy–Schwarz for the Hilbert–Schmidt inner product, $|\,\mathrm{Tr}(\rho M_\alpha)| \le \|\rho^{1/2}\|_2^2\|M_\alpha\|_\infty = \mathrm{Tr}\,\rho$, giving (7.118). Equality holds iff $\rho^{1/2}$ and $M_\alpha\rho^{1/2}$ are linearly dependent, i.e. $M_\alpha\rho = c\,\rho$ with $c \ge 0$, equivalent to the stated proportionality; for pure $\rho = |\psi\rangle\langle\psi|$ this is $M_\alpha\,|\psi\rangle = e^{i\varphi}\,|\psi\rangle$. For the final claim, insert $U_j^{(\mathrm{state})} = UW_j$ in (7.115) and use cyclicity to obtain $\mathcal{C}_{\mathrm{state}} \propto \mathrm{Tr}(W_1\rho W_0^\dagger) = \mathrm{Tr}(\rho_1\rho_0)$; for pure preparations this gives $\langle\psi_1\rangle\,\psi_0$. □

Proposition 7.143 (Short-time expansion and decay scales)**.** *Let* $\delta\hat H_\alpha := \hat H_1^{(\alpha)} - \hat H_0^{(\alpha)}$ *and assume time-independent branches. Then on* $\mathcal{H}_{\mathrm{phys}}$*,*

$$\mathcal{C}_\alpha(T) = 1 - iT\,\frac{\mathrm{Tr}(\rho_{\mathrm{phys}}\delta\hat H_\alpha)}{\mathrm{Tr}\,\rho_{\mathrm{phys}}} - \frac{T^2}{2}\,\frac{\mathrm{Tr}\big(\rho_{\mathrm{phys}}(\delta\hat H_\alpha - \langle\delta\hat H_\alpha\rangle)^2\big)}{\mathrm{Tr}\,\rho_{\mathrm{phys}}} + \mathcal{O}(T^3), \tag{7.119}$$

so that $|\mathcal{C}_\alpha(T)|^2 = 1 - T^2\,\sigma_\alpha^2 + \mathcal{O}(T^3)$*, with* $\langle A\rangle := \mathrm{Tr}(\rho_{\mathrm{phys}}A)/\,\mathrm{Tr}\,\rho_{\mathrm{phys}}$ *and* $\sigma_\alpha^2 := \langle(\delta\hat H_\alpha - \langle\delta\hat H_\alpha\rangle)^2\rangle$*. This expansion is band-safe since* ρ_{phys} *and* $\hat H_j^{(\alpha)}$ *are constructed with* $P_x(\lambda_{\mathrm{UV}})$ *and* $K_{\mathrm{IR}}^{(\ell)}$*.*

Proof. By cyclicity, $\mathcal{C}_\alpha(T) = \langle e^{i\hat H_0^{(\alpha)}T}e^{-i\hat H_1^{(\alpha)}T}\rangle$ with the normalised expectation $\langle\cdot\rangle$. Using the Baker–Campbell–Hausdorff formula, $e^{iH_0T}e^{-iH_1T} = \exp\big(-i\delta\hat H_\alpha T + \frac{T^2}{2}[H_0,H_1] + \mathcal{O}(T^3)\big)$. Expanding the exponential and keeping terms up to $\mathcal{O}(T^2)$ yields

$$\mathcal{C}_\alpha(T) = 1 - iT\langle\delta\hat H_\alpha\rangle - \tfrac{T^2}{2}\langle(\delta\hat H_\alpha)^2\rangle + \tfrac{T^2}{2}\langle[H_0,H_1]\rangle + \mathcal{O}(T^3).$$

Since $[H_0,H_1]$ is anti-Hermitian, $\langle[H_0,H_1]\rangle$ is purely imaginary and contributes only a phase (no change in modulus). Writing $\langle(\delta\hat H_\alpha)^2\rangle = \sigma_\alpha^2 + \langle\delta\hat H_\alpha\rangle^2$ and absorbing the purely imaginary quadratic phase into the reference phase ϕ_α gives (7.119). Squaring the modulus then yields $|\mathcal{C}_\alpha(T)|^2 = 1 - T^2\sigma_\alpha^2 + \mathcal{O}(T^3)$. □

Gauge/BRST robustness and regulator stability. Differentiating (7.115) with respect to a gauge-fixing parameter α_{gf} and using the Nielsen master identity (7.19) shows that $\partial_{\alpha_{\mathrm{gf}}}\mathcal{C}_\alpha$ is a sum of (i) ζ-exact insertions, which decouple on $\mathcal{H}_{\mathrm{phys}}$ by quartet/positivity Section 7.2.6 and eq. (7.12), and (ii) EOM insertions, which vanish on shell eqs. (6.333) and (6.334). Thus $\mathcal{C}_\alpha$ and any phase-fixed Λ_α are *gauge-parameter independent on shell* [117, 91]. Regularisation stability under admissible changes of P_x and $K_{\mathrm{IR}}^{(\ell)}$ follows from Section 7.3.7.

Proposition 7.144 (Gauge/BRST robustness and regulator stability for $\mathcal{C}_\alpha$)**.** *Differentiating* (7.115) *with respect to a gauge-fixing parameter* α_{gf} *and using the Nielsen master identity* (7.19) *yields*

$$\partial_{\alpha_{\mathrm{gf}}}\mathcal{C}_\alpha = \frac{\mathrm{Tr}\big(\mathsf{P}_{\mathrm{phys}}\,\mathsf{O}_{\zeta\text{-}exact}\,\mathsf{P}_{\mathrm{phys}}\big)}{\mathrm{Tr}(\mathsf{P}_{\mathrm{phys}}\rho_{\mathrm{phys}})} + \frac{\mathrm{Tr}\big(\mathsf{P}_{\mathrm{phys}}\,\mathsf{O}_{\mathrm{EOM}}\,\mathsf{P}_{\mathrm{phys}}\big)}{\mathrm{Tr}(\mathsf{P}_{\mathrm{phys}}\rho_{\mathrm{phys}})}.$$

On $\mathcal{H}_{\mathrm{phys}}$*,* ζ*-exact insertions decouple by quartet/positivity Section 7.2.6 and eq.* (7.12)*, and EOM insertions vanish on shell eqs.* (6.333) *and* (6.334)*. Hence*

$$\partial_{\alpha_{\mathrm{gf}}}\mathcal{C}_\alpha = 0 \quad \textit{on shell}, \qquad \partial_{\alpha_{\mathrm{gf}}}\Lambda_\alpha = 0 \quad \textit{on shell},$$

for any phase-fixed Λ_α *in* (7.117)*. Moreover, under admissible regularisation changes of* $P_x(\lambda_{\mathrm{UV}})$ *and* $K_{\mathrm{IR}}^{(\ell)}$*,* $\mathcal{C}_\alpha$ *is stable by the regulator robustness results in Section 7.3.7.*

Proof. Apply (7.19) to the operator insertions in (7.115) and insert $\mathsf{P}_{\mathrm{phys}}$ on both sides of each trace. Use BRST-closure of the preparation and band knobs, and invoke quartet/positivity Section 7.2.6 and eq. (7.12) to kill ζ-exact terms; EOM terms vanish on shell by eqs. (6.333) and (6.334). Regulator stability follows by the Lipschitz/IR-L^1 and gapped-projector arguments in Section 7.3.7. □

Phase choice and signed visibility. The reference phase ϕ_α in (7.117) can be fixed by any of the following equivalent prescriptions: (i) *centre phase* $\phi_\alpha = \arg\mathcal{C}_\alpha(\lambda_\star, \ell_\star; T)$ at the certified plateau centre (so $\Lambda_\alpha(\lambda_\star, \ell_\star; T) = |\mathcal{C}_\alpha|$); (ii) *echo phase* by inserting a symmetric echo to remove common-mode dynamical phase (useful when $U_{0,1}^{(\alpha)}$ share large identical pieces); (iii) *channel gauge* where ϕ_α is the continuous choice along any certified path in the plateau, minimising $\partial_u\phi_\alpha$ and $\partial_v\phi_\alpha$ (smooth-phase gauge). Any such choice leaves $|\mathcal{C}_\alpha|$ invariant and preserves the plateau metrics.

Proposition 7.145 (Phase choice and signed visibility)**.** *Let* Λ_α *be defined by* (7.117)*. The reference phase* ϕ_α *may be fixed by any of the following equivalent prescriptions without affecting* $|\mathcal{C}_\alpha|$ *or the plateau metrics:*

(i) Centre phase.
Set $\phi_\alpha = \arg\mathcal{C}_\alpha(\lambda_\star, \ell_\star; T)$ *at the certified plateau centre, so that* $\Lambda_\alpha(\lambda_\star, \ell_\star; T) = |\mathcal{C}_\alpha|$*.*

(ii) Echo phase.
Insert a symmetric echo to cancel common-mode dynamical phase when $U_{0,1}^{(\alpha)}$ *share large identical pieces; fix* ϕ_α *by the echo.*

(iii) Channel gauge.
Choose ϕ_α *continuously along any certified plateau path to minimise* $\partial_u\phi_\alpha$ *and* $\partial_v\phi_\alpha$ *(smooth-phase gauge).*

All choices are related by a constant phase shift on certified plateaus and leave $|\mathcal{C}_\alpha|$ *invariant.*

Integrated readouts and plateaus. Define the scale–box averaged $\overline{\mathcal{C}}_\alpha$ by replacing Λ_α with $\mathcal{C}_\alpha$ in eq. (7.73). The commutation of averaging with scale derivatives in the log chart eq. (7.74) and the oscillation bound eq. (7.76) imply that $|\overline{\mathcal{C}}_\alpha|$ and the phase-fixed $\overline{\Lambda}_\alpha$ inherit the plateau properties of Section 7.3.5 and the certification protocol of Section 7.3.9.

Proposition 7.146 (Integrated readouts and plateaus)**.** *Define the scale–box average by replacing Λ_α with $\mathcal{C}_\alpha$ in eq. (7.73):*

$$\overline{\mathcal{C}}_{\alpha|B} := \frac{1}{\mu(B)}\int_B d\lambda_{\mathrm{UV}}\, d\ell\, dT\; \mathcal{C}_\alpha(\lambda_{\mathrm{UV}}, \ell; T).$$

Then averaging commutes with log-scale derivatives per eq. (7.74) and obeys the oscillation bound eq. (7.76). Consequently, $|\overline{\mathcal{C}}_\alpha|$ and the phase-fixed $\overline{\Lambda}_\alpha$ inherit the plateau properties of Section 7.3.5 and the certification protocol of Section 7.3.9.

Proof. Linearity of the trace and the admissibility of the knobs imply Gateaux differentiability under the integral; apply eqs. (7.74) and (7.76) to $\mathcal{C}_\alpha$ exactly as for visibilities, and invoke the same geometric/metric equivalence criteria from Section 7.3.5 together with the acceptance pipeline in Section 7.3.9. □

Quadrant preservation and diagnostic meaning. Because the branch differences in (7.97) are cohomologically trivial or symmetry-preserving, the holosymmetric probe does not alter quadrant membership. In Quadrant I, large $|\mathcal{C}_\alpha|$ on certified plateaus indicates high law/state coherence; sustained suppression signals genuine physical decoherence or IR effects consistent with unbroken laws, not gauge artefacts (cf. Sections 7.3 and 7.3.4).

Proposition 7.147 (Quadrant preservation and diagnostic meaning)**.** *Under (7.97), inter-branch differences are cohomologically trivial or symmetry-preserving on $\mathcal{H}_{\mathrm{phys}}$. Therefore the holosymmetric probe preserves quadrant membership:*

$$U_j^{(\alpha)}(T) = \bigoplus_q U_j^{(\alpha,q)}(T), \qquad \mathcal{C}_\alpha = \sum_q \frac{\mathrm{Tr}\big(\Pi^{(q)}\rho_{\mathrm{phys}}\Pi^{(q)}\, M_\alpha^{(q)}\big)}{\mathrm{Tr}(\rho_{\mathrm{phys}})},$$

with $M_\alpha^{(q)} = (U_0^{(\alpha,q)})^\dagger U_1^{(\alpha,q)}$ and $\Pi^{(q)}$ the quadrant projectors. In Quadrant I, certified plateaus imply large $|\mathcal{C}_\alpha|$ (high law/state coherence), whereas sustained suppression signals genuine physical decoherence or symmetry-allowed IR effects—consistent with unbroken laws and not gauge artefacts (cf. Sections 7.3 and 7.3.4).

Proof. Cohomological triviality/symmetry preservation makes each branch block-diagonal in the quadrant decomposition, hence no cross-quadrant interference. The decomposition of $\mathcal{C}_\alpha$ follows by inserting $\sum_q \Pi^{(q)} = \mathbb{I}$ and using cyclicity. The diagnostic statement follows from the certified plateau criteria and robustness guarantees in Section 7.3. □

7.4.2 Channel Realisations: Law vs. State

The two channels differ only in the *pair of branches* $U_{0,1}^{(\alpha)}(T)$; their design guarantees that the operational overlay leaves quadrant membership unchanged (branches differ by cohomologically trivial/symmetry-preserving deformations) [442, 117, 91]. Throughout, $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ are BRST-closed and band-safe eqs. (7.29), (7.30) and (7.33).

Law channel ($\alpha = \text{law}$). Let $H_0(t)$ be the Hamiltonian density generated by the symmetric law (Quadrant I) associated with the clean kinetic pairings and potentials of eqs. (6.333) and (6.334). Define

$$U_0^{(\text{law})}(T) = \mathcal{T}\exp\left[-i\int_0^T dt\, H_0(t)\right], \qquad U_1^{(\text{law})}(T) = \mathcal{T}\exp\left[-i\int_0^T dt\,\Big(H_0(t) + \zeta\,\Delta G(t)\Big)\right], \tag{7.120}$$

with $\Delta G(t)$ local and band-limited. Because $\zeta\,\Delta G$ is BRST-exact, the Slavnov identity and quartet mechanism imply that $U_1^{(\text{law})}$ and $U_0^{(\text{law})}$ produce identical physical matrix elements on $\mathcal{H}_{\text{phys}}$ (anomaly-free case), hence

$$\mathcal{C}_{\text{law}}(\lambda_{\text{UV}}, \ell; T) = 1 \quad \text{and} \quad \Lambda_{\text{law}}(\lambda_{\text{UV}}, \ell; T) = 1 \quad \text{on shell in Quadrant I}, \tag{7.121}$$

up to numerical/regulator tolerances. Any robust suppression $|\mathcal{C}_{\text{law}}| < 1$ signals *law-level incoherence* (e.g. a broken Slavnov identity/anomaly or a regulator pathology), and triggers the anomaly/regularisation checks of Sections 7.2.8 and 7.3.7. The *law visibility* is defined as

$$\Lambda_{\text{law}}(\lambda_{\text{UV}}, \ell; T) := \text{Re}\Big(e^{-i\phi_{\text{law}}}\,\mathcal{C}_{\text{law}}(\lambda_{\text{UV}}, \ell; T)\Big), \tag{7.122}$$

with ϕ_{law} fixed by a calibration at a certified plateau centre $(\lambda_\star, \ell_\star)$ (cf. Section 7.3.9).

Proposition 7.148 (On-shell unity of the law-channel coherence)**.** *Under the assumptions of (7.120) and anomaly freedom, the overlap operator $M_{\text{law}}(T) := (U_0^{(\text{law})})^\dagger U_1^{(\text{law})}$ acts as the identity on $\mathcal{H}_{\text{phys}}$. Consequently, $\mathcal{C}_{\text{law}} = 1$ and $\Lambda_{\text{law}} = 1$ on shell, as in (7.121).*

Proof. Write $U_1^{(\text{law})}$ in the interaction picture with respect to H_0: $U_1^{(\text{law})} = U_0\,\mathcal{T}\exp\Big(-i\int_0^T dt\,\zeta\,\Delta G_I(t)\Big)$, where $U_0 := U_0^{(\text{law})}$. Then $M_{\text{law}} = \mathcal{T}\exp\Big(-i\int_0^T dt\,\zeta\,\Delta G_I(t)\Big)$. Inside physical matrix elements, ζ-exact insertions decouple by quartet/positivity and the Slavnov identity; thus $\mathsf{P}_{\text{phys}}\,M_{\text{law}}\,\mathsf{P}_{\text{phys}} = \mathsf{I}_{\mathcal{H}_{\text{phys}}}$, proving the claim. □

Remark 7.149 (Diagnostic meaning of law-channel suppression)**.** If a certified plateau exhibits $|\mathcal{C}_{\text{law}}| < 1$ beyond tolerances, either (i) the BRST/Slavnov structure is anomalous, (ii) regulator admissibility fails, or (iii) the preparation strays off-shell. The triage proceeds via Sections 7.2.8 and 7.3.7.

State channel ($\alpha = \text{state}$). Fix a BRST-compatible reference Hamiltonian $H_{\text{ref}}(t)$ and two admissible preparations W_j (band-safe, BRST-compatible, quadrant-preserving). Realise the branches by *pushing* the preparations into the evolution,

$$U_j^{(\text{state})}(T) = U_{\text{ref}}(T)\,W_j, \qquad U_{\text{ref}}(T) := \mathcal{T}\exp\left[-i\int_0^T dt\, H_{\text{ref}}(t)\right], \tag{7.123}$$

so that the input remains $\rho_{\text{phys}}^{(\lambda_{\text{UV}},\ell)}$ and the channel difference resides in $W_{0,1}$ (equivalently to preparing $\rho_j = W_j \rho_{\text{phys}} W_j^\dagger$ with a common U_{ref}).

Proposition 7.150 (Coherence amplitude for the state channel)**.** *With* (7.123) *and* (7.115)*, one has*

$$\mathcal{C}_{\mathrm{state}}(\lambda_{\mathrm{UV}},\ell;T)=\frac{\mathrm{Tr}\Big(W_1\,\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}\,W_0^{\dagger}\Big)}{\mathrm{Tr}\Big(\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}\Big)},$$

which is precisely eq. (7.109)*. In particular, for* $\rho_{\mathrm{phys}}=|\psi\rangle\langle\psi|$ *and* $W_j\,|\psi\rangle=|\psi_j\rangle$*,* $\mathcal{C}_{\mathrm{state}}=\langle\psi_1\rangle\,\psi_0$ *and* $|\mathcal{C}_{\mathrm{state}}|^2$ *is the (pure-state) fidelity.*

Proof. Insert $U_j^{(\mathrm{state})}=U_{\mathrm{ref}}W_j$ into (7.115) and use cyclicity plus unitarity of U_{ref} to cancel it, yielding the displayed expression and eq. (7.109). □

Definition 7.151 (State visibility)**.** Define the phase-fixed state visibility by

$$\Lambda_{\mathrm{state}}(\lambda_{\mathrm{UV}},\ell;T):=\mathrm{Re}\Big(e^{-i\phi_{\mathrm{state}}}\,\mathcal{C}_{\mathrm{state}}(\lambda_{\mathrm{UV}},\ell;T)\Big),$$

with ϕ_{state} chosen by any of the prescriptions in Proposition 7.145.

Proposition 7.152 (Bounds and saturation)**.** *The bounds of Propositions 7.128 and 7.139 apply to* $\mathcal{C}_{\mathrm{state}}$ *and* Λ_{state}*. Equality* $|\mathcal{C}_{\mathrm{state}}|=1$ *holds iff* $W_1\rho_{\mathrm{phys}}W_0^{\dagger}$ *is a positive multiple of* ρ_{phys} *(pure-state case:* $W_1\,|\psi\rangle=e^{i\varphi}W_0\,|\psi\rangle$*).*

Remark 7.153 (Gauge/BRST protection and plateaus)**.** By Proposition 7.144 the quantities $\mathcal{C}_{\mathrm{state}}$ and Λ_{state} are gauge-parameter independent on shell and regulator-stable under admissible changes of $P_x, K_{\mathrm{IR}}^{(\ell)}$. Calibration of ϕ_{state} at $(\lambda_\star,\ell_\star)$ and certification follow Section 7.3.9.

State channel ($\alpha=$ state). Hold the *law* fixed to $H_0(t)$ for both branches, and seed a symmetry probe at the *state-preparation* stage by a unitary kick W generated by a conserved charge Q of the unbroken symmetry (Quadrant I):

$$\begin{aligned}\rho_0^{(\lambda_{\mathrm{UV}},\ell)}&\equiv\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)},\qquad \rho_1^{(\lambda_{\mathrm{UV}},\ell)}:=W\,\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}\,W^{\dagger},\\ W&:=\exp(-i\epsilon Q),\quad \epsilon\ll 1,\quad [Q,H_0(t)]=0.\end{aligned}\tag{7.124}$$

Evolve both with the same $U_0(T):=\mathcal{T}\exp\Big[-i\int_0^T dt\,H_0(t)\Big]$ and implement

$$U_0^{(\mathrm{state})}(T)=U_0(T),\qquad U_1^{(\mathrm{state})}(T)=U_0(T),\tag{7.125}$$

with branch initial states ρ_0 and ρ_1, respectively.

Proposition 7.154 (Reduction to a preparation overlap)**.** *Inserting* (7.125) *into* (7.115) *and using* $[Q,H_0]=0$ *gives*

$$\mathcal{C}_{\mathrm{state}}(\lambda_{\mathrm{UV}},\ell;T)=\frac{\mathrm{Tr}\Big(\mathsf{P}_{\mathrm{phys}}\,W\,\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}\,W^{\dagger}\,\mathsf{P}_{\mathrm{phys}}\Big)}{\mathrm{Tr}\Big(\mathsf{P}_{\mathrm{phys}}\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}\Big)}.\tag{7.126}$$

Proof. Since $[Q,H_0]=0$, $U_0(T)$ commutes with W. In (7.115), the U_0 factors cancel by cyclicity and unitarity, leaving (7.126). □

Corollary 7.155 (Expectation in a symmetric state)**.** *If the prepared state is symmetric,* $[Q, \rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}] = 0$ *(up to IR smearing), then*

$$\mathcal{C}_{\mathrm{state}} = 1 + \mathcal{O}(\epsilon^2), \qquad \Lambda_{\mathrm{state}} = 1 + \mathcal{O}(\epsilon^2). \tag{7.127}$$

Proof. From $W\rho W^\dagger = \rho - i\epsilon[Q,\rho] - \frac{\epsilon^2}{2}[Q,[Q,\rho]] + \cdots$, the first nonzero term is $\mathcal{O}(\epsilon^2)$ when $[Q,\rho] = 0$ (modulo IR smearing), whence (7.127). □

Proposition 7.156 (Small-ϵ expansion for pure preparations)**.** *For* $\rho_{\mathrm{phys}} = |\psi\rangle\langle\psi|$,

$$\mathcal{C}_{\mathrm{state}} = \langle\psi|\, e^{-i\epsilon Q}\, |\psi\rangle + \mathcal{O}(\epsilon^3) = 1 - \tfrac{\epsilon^2}{2}\,\mathrm{Var}_\psi(Q) + \mathcal{O}(\epsilon^3), \tag{7.128}$$

where $\mathrm{Var}_\psi(Q) := \langle\psi|\, Q^2\, |\psi\rangle - \langle\psi|\, Q\, |\psi\rangle^2$. *This equality holds in the phase-fixed convention where the linear term* $\sim i\epsilon\langle Q\rangle_\psi$ *is removed by the reference phase (cf. (7.117)); equivalently it describes the short-*ϵ *behaviour of* $|\mathcal{C}_{\mathrm{state}}|$.

Proof. Expand $e^{-i\epsilon Q} = 1 - i\epsilon Q - \frac{\epsilon^2}{2}Q^2 + \mathcal{O}(\epsilon^3)$. The overlap is $1 - i\epsilon\langle Q\rangle_\psi - \frac{\epsilon^2}{2}\langle Q^2\rangle_\psi + \cdots$. Removing the linear phase via $\phi_{\mathrm{state}} = \epsilon\langle Q\rangle_\psi$ (or taking modulus) gives the stated form with $\mathrm{Var}_\psi(Q)$. □

Definition 7.157 (State visibility)**.** The *state visibility* is

$$\Lambda_{\mathrm{state}}(\lambda_{\mathrm{UV}}, \ell; T) := \mathrm{Re}\Big(e^{-i\phi_{\mathrm{state}}}\, \mathcal{C}_{\mathrm{state}}(\lambda_{\mathrm{UV}}, \ell; T)\Big)\,. \tag{7.129}$$

The phase ϕ_{state} may be fixed by any of the prescriptions in Proposition 7.145.

Proposition 7.158 (Design constraints and robustness)**.** *Consider the holosymmetric operational overlay with branches* $U_{0,1}^{(\alpha)}(T)$ *acting on the physical sector, band-limited by* $P_x(\lambda_{\mathrm{UV}})$ *and* $K_{\mathrm{IR}}^{(\ell)}$. *Then:*

(i) Quadrant preservation.
In the law channel, branch differences are BRST-exact (cohomologically trivial); in the state channel, the symmetry generator Q *commutes with* H_0. *Hence the overlay is block-diagonal in the quadrant decomposition and does not alter quadrant membership [442, 117, 91]. Equivalently,* $U_j^{(\alpha)}(T) = \bigoplus_q U_j^{(\alpha,q)}(T)$ *and no cross-quadrant interference occurs (cf. (7.97) and Definition 7.114 and eq. (7.103)).*

(ii) Nielsen/BRST protection.
Differentiating the visibility/coherence definitions with respect to the gauge-fixing parameter and invoking the Nielsen identity (Section 7.2.9 and eq. (7.19)) shows that the derivative reduces to ζ*-exact and EOM insertions. The former decouple on* $\mathcal{H}_{\mathrm{phys}}$ *by quartet/positivity (Section 7.2.6 and eq. (7.12)); the latter vanish on shell by eqs. (6.333) and (6.334). Thus visibilities are gauge-parameter independent on shell, consistent with eq. (7.66) and the BRST invariance results proved above.*

(iii) Regularisation stability.
Replacing $(P_x, K_{\mathrm{IR}}^{(\ell)})$ *by any admissible* $(\tilde{P}_x, \tilde{K}_{\mathrm{IR}}^{(\ell)})$ *perturbs the readouts by at most* $\varepsilon_{\mathrm{reg}}$ *as certified in Section 7.3.7. Quantitatively, writing* $\rho := \rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}$ *and* $M_\alpha := (U_0^{(\alpha)})^\dagger U_1^{(\alpha)}$,

$$\Big|\mathcal{C}_\alpha - \tilde{\mathcal{C}}_\alpha\Big| = \left|\frac{\mathrm{Tr}(P\,\rho\,P\,M_\alpha)}{\mathrm{Tr}(P\,\rho\,P)} - \frac{\mathrm{Tr}\Big(\tilde{P}\,\rho\,\tilde{P}\,M_\alpha\Big)}{\mathrm{Tr}\Big(\tilde{P}\,\rho\,\tilde{P}\Big)}\right| \;\le\; \varepsilon_{\mathrm{reg}},$$

with $P := P_x K_{\mathrm{IR}}^{(\ell)}$ and $\tilde{P} := \tilde{P}_x \tilde{K}_{\mathrm{IR}}^{(\ell)}$, where $\varepsilon_{\mathrm{reg}}$ is controlled by the admissibility norms (IR L^1 closeness and UV-projector gap) and the Lipschitz constants cited in Section 7.3.7. The same bound transfers to Λ_α by continuity.

(iv) Plateau certification.
Both channels inherit the scale-dependence and plateau metrics of Section 7.3; readouts are to be certified via the acceptance pipeline in Section 7.3.9. Hence any reported Λ_α (or $\mathcal{C}_\alpha$) is accompanied by a certified plateau neighbourhood and uncertainty budget.

Proof. (i) Follows from the cohomological triviality/symmetry-preserving hypotheses and the quadrant-compatibility of the overlay (Definition 7.114); cf. the block-diagonal form implied by (7.97). (ii) Is the standard Nielsen/BRST argument using Sections 7.2.6 and 7.2.9 and eqs. (7.12) and (7.19) and on-shell EOM eqs. (6.333) and (6.334) (consistent with eq. (7.66)). (iii) Use the triangle inequality and the regulator-stability estimates in Section 7.3.7; the trace functional is Lipschitz on the admissible class, and the denominator remains bounded away from zero on certified bands. (iv) Is immediate from the plateau geometry and certification results cited. □

7.4.3 Embedding of UV/IR Knobs and Sector Projectors

Every operator in eqs. (7.96) and (7.129) is band-limited and IR-smoothed by the functor $\mathcal{B}^{(\lambda_{\mathrm{UV}},\ell)}$ of eq. (7.40):

$$U_j^{(\alpha)}(T) \;\mapsto\; U_j^{(\alpha)}(T;\lambda_{\mathrm{UV}},\ell) := \mathcal{T}\exp\!\left[-i\int_0^T dt \int d^d x\, K_{\mathrm{IR}}^{(\ell)}(x)\, \mathcal{H}_j^{(\alpha)}(t,x;\lambda_{\mathrm{UV}})\right], \tag{7.130}$$

where $\mathcal{H}_j^{(\alpha)}(t,x;\lambda_{\mathrm{UV}}) := \mathcal{B}^{(\lambda_{\mathrm{UV}},\ell)}[\mathcal{H}_j^{(\alpha)}(t,x)]$ is built from fields projected by $P_x(\lambda_{\mathrm{UV}})$ and averaged with $K_{\mathrm{IR}}^{(\ell)}$ (cf. Section 7.3.1). By construction, $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ are BRST-closed and ghost-number 0 eq. (7.30), so the embedding preserves the Slavnov identities and the decoupling properties on $\mathcal{H}_{\mathrm{phys}}$ [114, 117, 91].

Definition 7.159 (Band/IR functorial embedding)**.** Let $\mathcal{A}$ be the $*$-algebra of (densitised) local Hamiltonian densities admissible for the dynamics. The functor $\mathcal{B}^{(\lambda_{\mathrm{UV}},\ell)} : \mathcal{A} \to \mathcal{A}$ acts by

$$\mathcal{H}(t,x) \;\mapsto\; \Big(\mathcal{B}^{(\lambda_{\mathrm{UV}},\ell)}\mathcal{H}\Big)(t,x) := \int d^d y\, K_{\mathrm{IR}}^{(\ell)}(x-y)\, \Big(P_y(\lambda_{\mathrm{UV}})\, \mathcal{H}(t,y)\, P_y(\lambda_{\mathrm{UV}})\Big),$$

with $K_{\mathrm{IR}}^{(\ell)} \in L^1(\mathbb{R}^d)$ real and even, and $P_y(\lambda_{\mathrm{UV}})$ a local UV projector of eq. (7.29). Then $\mathcal{B}^{(\lambda_{\mathrm{UV}},\ell)}$ is completely positive, $*$-preserving, and norm-contracting on $\mathcal{A}$.

Lemma 7.160 (BRST closure and sector naturality)**.** *If $[\hat{Q}_{\mathrm{BRST}}, P_x(\lambda_{\mathrm{UV}})] = 0$ and $[\hat{Q}_{\mathrm{BRST}}, K_{\mathrm{IR}}^{(\ell)}] = 0$ as in eq. (7.30), then for any $\mathcal{H} \in \mathcal{A}$,*

$$[\hat{Q}_{\mathrm{BRST}}, \mathcal{B}^{(\lambda_{\mathrm{UV}},\ell)}\mathcal{H}] = \mathcal{B}^{(\lambda_{\mathrm{UV}},\ell)}\Big([\hat{Q}_{\mathrm{BRST}}, \mathcal{H}]\Big),$$
$$\mathsf{P}_{\mathrm{phys}}\, \mathcal{B}^{(\lambda_{\mathrm{UV}},\ell)}(\cdot)\, \mathsf{P}_{\mathrm{phys}} = \mathcal{B}^{(\lambda_{\mathrm{UV}},\ell)}\Big(\mathsf{P}_{\mathrm{phys}}(\cdot)\mathsf{P}_{\mathrm{phys}}\Big).$$

Hence $\mathcal{B}^{(\lambda_{\mathrm{UV}},\ell)}$ is BRST-natural and respects the physical/sector projectors.

Proposition 7.161 (Well-posed unitary band evolution)**.** *Assume for each branch j: (i) $\mathcal{H}_j^{(\alpha)}(t,\cdot;\lambda_{\mathrm{UV}})$ is essentially self-adjoint on a common, band-stable core $\mathcal{D}$; (ii) $t \mapsto \mathcal{H}_j^{(\alpha)}(t,\cdot;\lambda_{\mathrm{UV}})$ is strongly continuous on $[0,T]$; (iii) $K_{\mathrm{IR}}^{(\ell)} \in L^1(\mathbb{R}^d)$ and $\sup_t \|\mathcal{H}_j^{(\alpha)}(t,\cdot;\lambda_{\mathrm{UV}})\|$ is locally bounded in $(\lambda_{\mathrm{UV}},\ell)$. Then the time-ordered exponential in (7.130) exists and defines a unitary propagator $U_j^{(\alpha)}(T;\lambda_{\mathrm{UV}},\ell)$ on $\overline{\mathcal{D}}$, depending strongly continuously on T.*

Proof. Young's inequality implies $\|K_{\mathrm{IR}}^{(\ell)} * \cdot\| \le \|K_{\mathrm{IR}}^{(\ell)}\|_{L^1}\|\cdot\|$, preserving essential self-adjointness and the core $\mathcal{D}$. Strong continuity in t is inherited from the integrand; Kato's existence/uniqueness theorem for time-dependent self-adjoint generators then yields a unitary, strongly continuous evolution [115]. □

Proposition 7.162 (Preservation of BRST compatibility under embedding)**.** *Suppose the unembedded branches satisfy $[\hat{Q}_{\mathrm{BRST}}, U_j^{(\alpha)}(T)] = \zeta X_j^{(\alpha)}$ with $\mathsf{P}_{\mathrm{phys}}\zeta X_j^{(\alpha)}\mathsf{P}_{\mathrm{phys}} = 0$. Then the embedded propagators satisfy*

$$[\hat{Q}_{\mathrm{BRST}}, U_j^{(\alpha)}(T;\lambda_{\mathrm{UV}},\ell)] = \zeta\, \widetilde{X}_j^{(\alpha)}(\lambda_{\mathrm{UV}},\ell), \qquad \mathsf{P}_{\mathrm{phys}}\, \widetilde{X}_j^{(\alpha)}\, \mathsf{P}_{\mathrm{phys}} = 0,$$

so BRST cohomological triviality persists on $\mathcal{H}_{\mathrm{phys}}$.

Proof. Differentiate the Dyson series for $U_j^{(\alpha)}(T;\lambda_{\mathrm{UV}},\ell)$; by Lemma 7.160 the commutator with $\hat{Q}_{\mathrm{BRST}}$ pushes through $\mathcal{B}^{(\lambda_{\mathrm{UV}},\ell)}$. Each time-ordered term becomes BRST-exact with the same physical-sector nullification; summing the series gives the claim. □

Proposition 7.163 (Quadrant/block structure under embedding)**.** *Let $\mathcal{H}_{\mathrm{phys}} = \bigoplus_q \mathcal{H}_{\mathrm{phys}}^{(q)}$ and suppose the unembedded branches are block-diagonal, $U_j^{(\alpha)}(T) = \bigoplus_q U_j^{(\alpha,q)}(T)$. Then the embedded propagators remain block-diagonal:*

$$U_j^{(\alpha)}(T;\lambda_{\mathrm{UV}},\ell) = \bigoplus_q U_j^{(\alpha,q)}(T;\lambda_{\mathrm{UV}},\ell), \qquad M_\alpha(T;\lambda_{\mathrm{UV}},\ell) = \bigoplus_q M_\alpha^{(q)}(T;\lambda_{\mathrm{UV}},\ell),$$

with $M_\alpha := (U_0^{(\alpha)})^\dagger U_1^{(\alpha)}$. Consequently, eq. (7.103) holds for the embedded objects.

Proof. $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ commute with the quadrant projectors on $\mathcal{H}_{\mathrm{phys}}$ (ghost number 0, BRST-closed), so $\mathcal{B}^{(\lambda_{\mathrm{UV}},\ell)}$ preserves the block decomposition and hence so do the Dyson series and their adjoints. □

Proposition 7.164 (Gateaux derivatives in UV/IR knobs)**.** *Assume $(\lambda_{\mathrm{UV}},\ell) \mapsto P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ are (piecewise) C^1 in the admissible sense, with $\partial_{\lambda_{\mathrm{UV}}} P_x$ bounded and $\partial_\ell K_{\mathrm{IR}}^{(\ell)} \in L^1$. Then the map $(\lambda_{\mathrm{UV}},\ell) \mapsto U_j^{(\alpha)}(T;\lambda_{\mathrm{UV}},\ell)$ is Gateaux differentiable in the strong operator topology and*

$$\partial_{\lambda_{\mathrm{UV}}} U_j^{(\alpha)}(T;\lambda_{\mathrm{UV}},\ell) = -i\int_0^T dt\; U_j^{(\alpha)}(T,t)\left(\int d^dx\, K_{\mathrm{IR}}^{(\ell)}(x)\, \partial_{\lambda_{\mathrm{UV}}} \mathcal{H}_j^{(\alpha)}(t,x;\lambda_{\mathrm{UV}})\right) U_j^{(\alpha)}(t,0),$$

$$\partial_\ell U_j^{(\alpha)}(T;\lambda_{\mathrm{UV}},\ell) = -i\int_0^T dt\; U_j^{(\alpha)}(T,t)\left(\int d^dx\, (\partial_\ell K_{\mathrm{IR}}^{(\ell)})(x)\, \mathcal{H}_j^{(\alpha)}(t,x;\lambda_{\mathrm{UV}})\right) U_j^{(\alpha)}(t,0),$$

with $U_j^{(\alpha)}(t',t)$ the embedded propagator from t to t'. Moreover, the derivatives obey the Lipschitz bound

$$\left\|\partial_{\lambda_{\mathrm{UV}}} U_j^{(\alpha)}(T;\lambda_{\mathrm{UV}},\ell)\right\| \le T\, \|\partial_{\lambda_{\mathrm{UV}}} \mathcal{H}_j^{(\alpha)}\|_{\mathrm{op}} \|K_{\mathrm{IR}}^{(\ell)}\|_{L^1},$$
$$\left\|\partial_\ell U_j^{(\alpha)}(T;\lambda_{\mathrm{UV}},\ell)\right\| \le T\, \|\mathcal{H}_j^{(\alpha)}\|_{\mathrm{op}} \|\partial_\ell K_{\mathrm{IR}}^{(\ell)}\|_{L^1}.$$

Proof. Differentiate the Dyson series term-by-term; boundedness of the variational kernels and Young's inequality control the series uniformly in order, permitting interchange of limit and sum. The norm estimates follow by Grönwall-type bounds and $\|K * \cdot\| \le \|K\|_{L^1}\|\cdot\|$. □

Corollary 7.165 (Stability of readouts in UV/IR knobs)**.** *Let $\mathcal{C}_\alpha$ and Λ_α be defined as in eqs. (7.115) and (7.117). Under the hypotheses of Proposition 7.164, the maps $(\lambda_{\mathrm{UV}}, \ell) \mapsto \mathcal{C}_\alpha$ and $(\lambda_{\mathrm{UV}}, \ell) \mapsto \Lambda_\alpha$ are Gateaux differentiable and locally Lipschitz on certified bands. In particular,*

$$\left|\partial_{\lambda_{\mathrm{UV}}} \mathcal{C}_\alpha\right| \le \frac{T\,\|K_{\mathrm{IR}}^{(\ell)}\|_{L^1}}{\mathrm{Tr}(\mathsf{P}_{\mathrm{phys}}\rho_{\mathrm{phys}})}\ \mathrm{Tr}\Big(\mathsf{P}_{\mathrm{phys}}\rho_{\mathrm{phys}}\mathsf{P}_{\mathrm{phys}}\Big)\,\|\partial_{\lambda_{\mathrm{UV}}}\mathcal{H}_j^{(\alpha)}\|_{\mathrm{op}},$$

and similarly with ∂_ℓ and $\|\partial_\ell K_{\mathrm{IR}}^{(\ell)}\|_{L^1}$.

Proposition 7.166 (Heisenberg embedding and observable band-limiting)**.** *Let $\hat{O}$ be a (quasi)local observable with finite band norm on the admissible class. Define the embedded Heisenberg evolution*

$$\hat{O}_{j,\mathrm{H}}^{(\alpha)}(T;\lambda_{\mathrm{UV}},\ell) := U_j^{(\alpha)\dagger}(T;\lambda_{\mathrm{UV}},\ell)\,\hat{O}\,U_j^{(\alpha)}(T;\lambda_{\mathrm{UV}},\ell).$$

Then $\hat{O}_{j,\mathrm{H}}^{(\alpha)}$ is band-limited/IR-smoothed (belongs to the image of $\mathcal{B}^{(\lambda_{\mathrm{UV}},\ell)}$ up to conjugation) and preserves BRST/sector naturality as in Lemma 7.160. Consequently, any readout functional built from $\hat{O}_{j,\mathrm{H}}^{(\alpha)}$ inherits the gauge/BRST robustness and regularisation stability established earlier.

Remark 7.167 (Ordering of IR averaging and time-ordering)**.** Equation (7.130) places the IR averaging inside the time-ordered exponential. If one first forms unembedded $U_j^{(\alpha)}(T)$ and then applies the (completely positive) band map on the propagator level, the two procedures coincide on $\mathcal{H}_{\mathrm{phys}}$ up to $\mathcal{O}(\|K_{\mathrm{IR}}^{(\ell)}\|_{L^1}\,\varepsilon_{\mathrm{reg}})$ in the admissible topology, by the CP-contractivity of $\mathcal{B}^{(\lambda_{\mathrm{UV}},\ell)}$ and the regulator robustness estimates of Section 7.3.7.

Scale derivatives (Duhamel formulas). The $(\lambda_{\mathrm{UV}}, \ell)$-dependence of the embedded branches is entirely through the knobs. Differentiating the time-ordered exponential gives band-safe Duhamel identities:

$$\partial_{\ln\lambda_{\mathrm{UV}}} U_j^{(\alpha)}(T;\lambda_{\mathrm{UV}},\ell) = -i\int_0^T dt \int d^dx\, K_{\mathrm{IR}}^{(\ell)}(x)\, U_j^{(\alpha)}(T,t)\,\Big(\partial_{\ln\lambda_{\mathrm{UV}}}\mathcal{H}_j^{(\alpha)}\Big)(t,x;\lambda_{\mathrm{UV}})\,U_j^{(\alpha)}(t,0), \tag{7.131}$$

$$\partial_{\ln\ell} U_j^{(\alpha)}(T;\lambda_{\mathrm{UV}},\ell) = -i\int_0^T dt \int d^dx\,\Big(\partial_{\ln\ell}K_{\mathrm{IR}}^{(\ell)}(x)\Big)\,U_j^{(\alpha)}(T,t)\,\mathcal{H}_j^{(\alpha)}(t,x;\lambda_{\mathrm{UV}})\,U_j^{(\alpha)}(t,0), \tag{7.132}$$

with $U_j^{(\alpha)}(t_2,t_1)$ the band/IR-smoothed propagator. These formulas yield explicit expressions for $D_\lambda\Lambda_\alpha, D_\ell\Lambda_\alpha$ used in the plateau metrics (Sections 7.3.2 and 7.3.4); operator-norm bounds on the integrands give Lipschitz control of the visibilities in the knobs (sufficient conditions in Section 7.3.7).

Proposition 7.168 (Duhamel identities and assumptions)**.** *Assume the hypotheses of Proposition 7.161 and, in addition, (i) $(\lambda_{\mathrm{UV}},\ell) \mapsto \mathcal{H}_j^{(\alpha)}(t,\cdot;\lambda_{\mathrm{UV}})$ is (piecewise) C^1 in the strong operator topology with $\partial_{\ln\lambda_{\mathrm{UV}}}\mathcal{H}_j^{(\alpha)}$ bounded on the scan window, and (ii) $\partial_{\ln\ell}K_{\mathrm{IR}}^{(\ell)} \in L^1(\mathbb{R}^d)$. Then (7.131)–(7.132) hold as strong operator identities on the common core.*

Proof. Write the Dyson series for $U_j^{(\alpha)}(T;\lambda_{\mathrm{UV}},\ell)$ using the embedded densities. Differentiate term-by-term; boundedness of the variational kernels and Young's inequality control each term uniformly, legitimising exchange of $\partial_{\ln\lambda_{\mathrm{UV}}}$ or $\partial_{\ln\ell}$ with the infinite sum. Collect the insertions in the interaction picture to obtain (7.131)–(7.132). □

Lemma 7.169 (Heisenberg form of the kernels)**.** *Define the interaction-picture operators*

$$\mathcal{K}_j^{(\lambda)}(t) := \int d^d x\, K_{\mathrm{IR}}^{(\ell)}(x)\big(\partial_{\ln\lambda_{\mathrm{UV}}}\mathcal{H}_j^{(\alpha)}\big)(t,x;\lambda_{\mathrm{UV}}),$$
$$\mathcal{K}_j^{(\ell)}(t) := \int d^d x\, (\partial_{\ln\ell}K_{\mathrm{IR}}^{(\ell)})(x)\,\mathcal{H}_j^{(\alpha)}(t,x;\lambda_{\mathrm{UV}}).$$

Then (7.131)–(7.132) *read*

$$\partial_\bullet U_j^{(\alpha)}(T) = -i\int_0^T dt\, U_j^{(\alpha)}(T,t)\,\mathcal{K}_j^{(\bullet)}(t)\,U_j^{(\alpha)}(t,0),\quad \bullet\in\{\ln\lambda_{\mathrm{UV}},\ln\ell\}.$$

Proposition 7.170 (Norm bounds and Lipschitz control)**.** *With $\|\cdot\|$ the operator norm and using unitarity of $U_j^{(\alpha)}(t_2,t_1)$,*

$$\big\|\partial_{\ln\lambda_{\mathrm{UV}}}U_j^{(\alpha)}(T)\big\| \le \int_0^T dt\,\big\|\mathcal{K}_j^{(\lambda)}(t)\big\| \le T\,\|K_{\mathrm{IR}}^{(\ell)}\|_{L^1}\sup_t\big\|\partial_{\ln\lambda_{\mathrm{UV}}}\mathcal{H}_j^{(\alpha)}(t,\cdot;\lambda_{\mathrm{UV}})\big\|,$$

and

$$\big\|\partial_{\ln\ell}U_j^{(\alpha)}(T)\big\| \le \int_0^T dt\,\big\|\mathcal{K}_j^{(\ell)}(t)\big\| \le T\,\|\partial_{\ln\ell}K_{\mathrm{IR}}^{(\ell)}\|_{L^1}\sup_t\big\|\mathcal{H}_j^{(\alpha)}(t,\cdot;\lambda_{\mathrm{UV}})\big\|.$$

Hence the maps $(\ln\lambda_{\mathrm{UV}},\ln\ell)\mapsto U_j^{(\alpha)}(T)$ are locally Lipschitz on certified bands.

Proof. Take norms inside (7.131)–(7.132); use $\|U_j^{(\alpha)}(t_2,t_1)\| = 1$ and Young's inequality $\big\|\int K * A\big\| \le \|K\|_{L^1}\|A\|$. □

Proposition 7.171 (Scale derivatives of the coherence amplitude)**.** *Let $\mathcal{C}_\alpha$ be given by* (7.115) *with $\rho_{\mathrm{phys}} := \rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}$. Decompose $D_\bullet\mathcal{C}_\alpha$ into a* dynamical part *from the branches and a* kinematic part *from the preparation/projectors:*

$$\partial_\bullet\mathcal{C}_\alpha = \frac{\begin{array}{l}\mathrm{Tr}\Big(\mathsf{P}\,(\partial_\bullet U_1^{(\alpha)})\,\rho_{\mathrm{phys}}\,U_0^{(\alpha)\dagger}\,\mathsf{P}\Big)\\ +\,\mathrm{Tr}\Big(\mathsf{P}\,U_1^{(\alpha)}\,\rho_{\mathrm{phys}}\,(\partial_\bullet U_0^{(\alpha)\dagger})\,\mathsf{P}\Big)\end{array}}{\mathrm{Tr}(\mathsf{P}\rho_{\mathrm{phys}})} + \Delta_\bullet^{(\mathrm{kin})},$$
$$\text{for } \bullet\in\{\ln\lambda_{\mathrm{UV}},\ln\ell\},$$

where $\mathsf{P} := \mathsf{P}_{\mathrm{phys}}$ and

$$\Delta_\bullet^{(\mathrm{kin})} := \frac{\begin{array}{l}\mathrm{Tr}\Big((\partial_\bullet\mathsf{P})\,U_1^{(\alpha)}\rho_{\mathrm{phys}}U_0^{(\alpha)\dagger}\Big)\\ +\,\mathrm{Tr}\Big(\mathsf{P}\,U_1^{(\alpha)}(\partial_\bullet\rho_{\mathrm{phys}})U_0^{(\alpha)\dagger}\Big)\\ +\,\mathrm{Tr}\Big(\mathsf{P}\,U_1^{(\alpha)}\rho_{\mathrm{phys}}U_0^{(\alpha)\dagger}(\partial_\bullet\mathsf{P})\Big)\end{array}}{\mathrm{Tr}(\mathsf{P}\rho_{\mathrm{phys}})} - \mathcal{C}_\alpha\,\frac{\mathrm{Tr}\Big((\partial_\bullet\mathsf{P})\rho_{\mathrm{phys}}+\mathsf{P}(\partial_\bullet\rho_{\mathrm{phys}})\Big)}{\mathrm{Tr}(\mathsf{P}\rho_{\mathrm{phys}})}.$$

Inserting (7.131)–(7.132) *gives explicit band-safe expressions for the dynamical part.*

Proof. Differentiate the numerator/denominator of (7.115) and use the product rule together with $\partial_\bullet U^\dagger = -U^\dagger(\partial_\bullet U)\,U^\dagger$. The preparation and projector terms collect into $\Delta_\bullet^{(\mathrm{kin})}$. □

Corollary 7.172 (Bounds for $D_\lambda \mathcal{C}_\alpha$ and $D_\ell \mathcal{C}_\alpha$)**.** *On certified bands where* $\mathrm{Tr}(\mathsf{P}\rho_{\mathrm{phys}}) \geq c_0 > 0$,

$$\left|\partial_{\ln\lambda_{\mathrm{UV}}}\mathcal{C}_\alpha\right| \;\leq\; \frac{2T\,\|K_{\mathrm{IR}}^{(\ell)}\|_{L^1}}{c_0}\,\max_{j=0,1}\sup_t\left\|\partial_{\ln\lambda_{\mathrm{UV}}}\mathcal{H}_j^{(\alpha)}(t,\cdot;\lambda_{\mathrm{UV}})\right\| \;+\; \left|\Delta_{\ln\lambda_{\mathrm{UV}}}^{(\mathrm{kin})}\right|,$$

and

$$\left|\partial_{\ln\ell}\mathcal{C}_\alpha\right| \;\leq\; \frac{2T\,\|\partial_{\ln\ell}K_{\mathrm{IR}}^{(\ell)}\|_{L^1}}{c_0}\,\max_{j=0,1}\sup_t\left\|\mathcal{H}_j^{(\alpha)}(t,\cdot;\lambda_{\mathrm{UV}})\right\| \;+\; \left|\Delta_{\ln\ell}^{(\mathrm{kin})}\right|.$$

The same bounds propagate to the signed visibility $\Lambda_\alpha = \mathrm{Re}(e^{-i\phi_\alpha}\mathcal{C}_\alpha)$ *by the chain rule.*

Remark 7.173 (BRST/Nielsen naturality of scale derivatives)**.** Under eq. (7.30) and the BRST-compatibility of the branches, the dynamical kernels in (7.131)–(7.132) are BRST-natural (Lemma 7.160); ζ-exact pieces decouple on $\mathcal{H}_{\mathrm{phys}}$, and on-shell EOM insertions vanish. Thus $D_\lambda\Lambda_\alpha$ and $D_\ell\Lambda_\alpha$ inherit the gauge/BRST robustness used in the plateau metrics (Sections 7.3.2 and 7.3.4).

BRST/Nielsen protection after embedding. Because the knobs commute with $\hat{Q}_{\mathrm{BRST}}$ and carry ghost number 0, $\partial_{\alpha_{\mathrm{gf}}}$-variations of the embedded readouts reduce to ζ-exact and EOM insertions which vanish on shell, exactly as before embedding (Section 7.3.6). Hence the embedded visibilities $\Lambda_\alpha(\lambda_{\mathrm{UV}},\ell)$ are gauge-parameter independent on $\mathcal{H}_{\mathrm{phys}}$ [117, 91].

Proposition 7.174 (Embedded Nielsen/BRST protection)**.** *Let* $\mathcal{C}_\alpha$ *and* Λ_α *be the embedded readouts defined with* $U_j^{(\alpha)}(T;\lambda_{\mathrm{UV}},\ell)$ *and* $\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}$*. Assume* $[\hat{Q}_{\mathrm{BRST}}, P_x(\lambda_{\mathrm{UV}})] = [\hat{Q}_{\mathrm{BRST}}, K_{\mathrm{IR}}^{(\ell)}] = 0$ *and the Nielsen identity eq.* (7.19)*. Then, on shell,*

$$\partial_{\alpha_{\mathrm{gf}}}\mathcal{C}_\alpha(\lambda_{\mathrm{UV}},\ell) = 0, \qquad \partial_{\alpha_{\mathrm{gf}}}\Lambda_\alpha(\lambda_{\mathrm{UV}},\ell) = 0.$$

Proof. Apply eq. (7.19) to the embedded numerator/denominator (with explicit $\mathsf{P}_{\mathrm{phys}}$ and band knobs inside the trace). Knob BRST-closure and ghost number 0 imply BRST-naturality of the embedding; ζ-exact insertions decouple on $\mathcal{H}_{\mathrm{phys}}$ by quartet/positivity, and EOM insertions vanish on shell (as in Section 7.3.6). The same argument transfers from $\mathcal{C}_\alpha$ to Λ_α. □

Sector projectors and resolved visibilities. Let $\{\Pi_\beta\}$ denote the BRST-closed channel/sector projectors (scalar/vector/tensor sectors relative to $\bar{\Sigma}_{\mu\nu}^{(\ell)}$), commuting with $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$. Define the *sector-resolved* coherence amplitude by inserting Π_β at readout,

$$\begin{aligned}\mathcal{C}_{\alpha|\beta} &:= \frac{\mathrm{Tr}\Big(\Pi_\beta\,\mathsf{P}_{\mathrm{phys}}\,U_1^{(\alpha)}(T;\lambda_{\mathrm{UV}},\ell)\,\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}\,U_0^{(\alpha)\dagger}(T;\lambda_{\mathrm{UV}},\ell)\,\mathsf{P}_{\mathrm{phys}}\,\Pi_\beta\Big)}{\mathrm{Tr}(\Pi_\beta\,\mathsf{P}_{\mathrm{phys}}\rho_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)})},\\ \Lambda_{\alpha|\beta} &:= \mathrm{Re}\Big(e^{-i\phi_\alpha}\mathcal{C}_{\alpha|\beta}\Big),\end{aligned} \tag{7.133}$$

which is manifestly gauge invariant and positive on $\mathcal{H}_{\text{phys}}$. If additionally $[\Pi_\beta, \mathcal{H}_j^{(\alpha)}] = 0$ (exact sector symmetry), the evolution is block-diagonal and $\Lambda_\alpha = \sum_\beta w_\beta \Lambda_{\alpha|\beta}$ with weights $w_\beta \geq 0$, $\sum_\beta w_\beta = 1$; otherwise (7.133) still defines a bona fide projector-resolved readout. Sectorwise plateaus are then certified by Section 7.3.11 using the compatibility criterion eq. (7.94).

Proposition 7.175 (Well-posedness, bounds, and gauge invariance)**.** *Let* $M_\alpha(T; \lambda_{\text{UV}}, \ell) := (U_0^{(\alpha)})^\dagger U_1^{(\alpha)}$ *be the embedded overlap. For any sector* β *with nonzero weight*

$$w_\beta := \frac{\mathrm{Tr}\Big(\Pi_\beta \, \mathsf{P}_{\text{phys}} \rho_{\text{phys}}^{(\lambda_{\text{UV}}, \ell)}\Big)}{\mathrm{Tr}\Big(\mathsf{P}_{\text{phys}} \rho_{\text{phys}}^{(\lambda_{\text{UV}}, \ell)}\Big)} > 0,$$

the quantity $\mathcal{C}_{\alpha|\beta}$ *in* (7.133) *is well-defined, satisfies* $|\mathcal{C}_{\alpha|\beta}| \leq 1$*, and is gauge-parameter independent on shell. Consequently,* $\Lambda_{\alpha|\beta} \in [-1, 1]$ *and inherits BRST/Nielsen protection and regulator stability.*

Proof. Denominator positivity follows from $w_\beta > 0$. Write $\mathcal{C}_{\alpha|\beta} = \mathrm{Tr}\Big(\rho^{(\beta)} M_\alpha\Big) / \mathrm{Tr}\Big(\rho^{(\beta)}\Big)$ with $\rho^{(\beta)} := \Pi_\beta \mathsf{P}_{\text{phys}} \rho_{\text{phys}} \mathsf{P}_{\text{phys}} \Pi_\beta \succeq 0$. Unitarity of M_α implies the bound by the same Cauchy–Schwarz/Hölder argument used earlier for un-resolved amplitudes. Gauge-parameter independence follows from Proposition 7.174 because Π_β is BRST-closed and commutes with the knobs, so the Nielsen reduction (exact+EOM) still vanishes on $\mathcal{H}_{\text{phys}}$. □

Proposition 7.176 (Additivity under exact sector symmetry)**.** *If* $[\Pi_\beta, \mathcal{H}_j^{(\alpha)}(t, \cdot; \lambda_{\text{UV}})] = 0$ *for all* t *and* j*, then the embedded propagators are block-diagonal and*

$$\mathcal{C}_\alpha = \sum_\beta w_\beta \, \mathcal{C}_{\alpha|\beta}, \qquad \Lambda_\alpha = \sum_\beta w_\beta \, \Lambda_{\alpha|\beta},$$

with weights w_β *as above (and a sector-independent phase* ϕ_α*).*

Proof. Block-diagonality gives $M_\alpha = \bigoplus_\beta M_{\alpha|\beta}$ and $\rho_{\text{phys}} = \sum_\beta \rho^{(\beta)}$ with $\mathrm{Tr}\, \rho^{(\beta)} = w_\beta \, \mathrm{Tr}(\mathsf{P}_{\text{phys}} \rho_{\text{phys}})$. Linearity of the trace yields the amplitude decomposition; applying $\mathrm{Re}(e^{-i\phi_\alpha} \cdot)$ gives the visibility statement. □

Remark 7.177 (Non-commuting sectors and certification)**.** If $[\Pi_\beta, \mathcal{H}_j^{(\alpha)}] \neq 0$, (7.133) still defines a valid sector-resolved readout (a "sandwiched" estimator). It remains gauge/BRST robust by Proposition 7.174 and admits the same Lipschitz/plateau analysis sectorwise. Certification then proceeds via Section 7.3.11 and the compatibility criterion eq. (7.94).

Regularisation stability and commuting limits. Replacing $(P_x, K_{\text{IR}}^{(\ell)})$ by $(\tilde{P}_x, \tilde{K}_{\text{IR}}^{(\ell)})$ with the same low-energy fixed point changes (7.130) by amounts controlled by Section 7.3.7; consequently the sector-resolved and aggregate visibilities obey the regulator bound (7.67), and the UV/IR swapped limits commute up to ε_{reg} (eq. (7.45)).

Proposition 7.178 (Regulator stability for embedded propagators and readouts)**.** *Let* $P := P_x(\lambda_{\text{UV}}) K_{\text{IR}}^{(\ell)}$ *and* $\tilde{P} := \tilde{P}_x(\lambda_{\text{UV}}) \tilde{K}_{\text{IR}}^{(\ell)}$ *be admissible band projectors with the* same *low-energy fixed point in the sense of Section 7.3.7. Then the embedded propagators in* (7.130) *satisfy*

$$\Big\| U_j^{(\alpha)}(T; \lambda_{\text{UV}}, \ell) - \tilde{U}_j^{(\alpha)}(T; \lambda_{\text{UV}}, \ell) \Big\| \;\leq\; \varepsilon_{\text{reg}}^{(U)},$$

and the overlap operators $M_\alpha := (U_0^{(\alpha)})^\dagger U_1^{(\alpha)}$ *obey*

$$\left\| M_\alpha(T;\lambda_{\mathrm{UV}},\ell) - \tilde{M}_\alpha(T;\lambda_{\mathrm{UV}},\ell) \right\| \;\le\; \varepsilon_{\mathrm{reg}}^{(M)}.$$

Consequently, the aggregate coherence amplitude and visibility satisfy the regulator bound (7.67)*:*

$$\left| \mathcal{C}_\alpha - \tilde{\mathcal{C}}_\alpha \right| \;\le\; \varepsilon_{\mathrm{reg}}, \qquad \left| \Lambda_\alpha - \tilde{\Lambda}_\alpha \right| \;\le\; \varepsilon_{\mathrm{reg}}.$$

Proof. Combine the Lipschitz/Duhamel control (7.131)–(7.132) with the admissibility norms and the robustness estimates in Section 7.3.7 to bound the generator differences inside the time-ordered exponentials; apply Grönwall-type bounds to obtain the stated operator-norm estimates. The trace functionals defining $\mathcal{C}_\alpha$ and Λ_α are Lipschitz on the admissible class, giving (7.67). □

Corollary 7.179 (Sector-resolved stability)**.** *Let* $\mathcal{C}_{\alpha|\beta}$ *and* $\Lambda_{\alpha|\beta}$ *be defined by* (7.133) *for any BRST-closed sector projector* Π_β *with nonzero weight. Then*

$$\left| \mathcal{C}_{\alpha|\beta} - \tilde{\mathcal{C}}_{\alpha|\beta} \right| \;\le\; \varepsilon_{\mathrm{reg}}, \qquad \left| \Lambda_{\alpha|\beta} - \tilde{\Lambda}_{\alpha|\beta} \right| \;\le\; \varepsilon_{\mathrm{reg}}.$$

Proof. Insert Π_β on both sides of the trace; $\|\Pi_\beta\| \le 1$ and the same Lipschitz/robustness arguments as above apply verbatim to the "sandwiched" functionals. □

Proposition 7.180 (Commuting UV/IR limits up to $\varepsilon_{\mathrm{reg}}$)**.** *Assume the UV/IR families* $(P_x, K_{\mathrm{IR}}^{(\ell)})$ *admit a common low-energy fixed point and the limits in eq.* (7.45) *exist along admissible paths. Then, for any certified time window and band,*

$$\left| \lim_{\ell\to\ell_\infty} \lim_{\lambda_{\mathrm{UV}}\to\lambda_\infty} \mathcal{C}_\alpha(\lambda_{\mathrm{UV}},\ell) \;-\; \lim_{\lambda_{\mathrm{UV}}\to\lambda_\infty} \lim_{\ell\to\ell_\infty} \mathcal{C}_\alpha(\lambda_{\mathrm{UV}},\ell) \right| \;\le\; \varepsilon_{\mathrm{reg}},$$

and likewise for Λ_α*, i.e. eq.* (7.45) *holds.*

Proof. Insert and subtract the fixed-point projector $P_\star := P_x(\lambda_\infty) K_{\mathrm{IR}}^{(\ell_\infty)}$ and use the triangle inequality together with (7.67). Each leg of the iterated limit is within $\varepsilon_{\mathrm{reg}}/2$ of the fixed-point readout, so their difference is $\le \varepsilon_{\mathrm{reg}}$. □

Summary. The embedding (7.130) implements the holosymmetric probe entirely within the band/IR window while preserving BRST, Nielsen, and sector structure. It yields explicit, band-safe scale-derivative formulas (7.131)–(7.132) used in the plateau metrics, and it supports sector-resolved diagnostics (7.133) without altering quadrant membership. Moreover, the construction is regulator-stable in the sense of (7.67) and its UV/IR limits commute up to $\varepsilon_{\mathrm{reg}}$ as certified by eq. (7.45) [114, 117, 91, 115, 82, 241].

7.4.4 Gauge/BRST Robustness and Anomaly Filters

Gauge-parameter variations modify the coherence amplitude only by EOM-proportional (and ζ-exact) insertions, so that

$$\partial_{\alpha_{\mathrm{gf}}} \mathcal{C}_\alpha = 0 \quad \text{on shell},$$

by the Nielsen master identity and quartet decoupling (Section 7.2.9 and eqs. (7.12), (7.19) and (7.66); see also [117, 91]). In the *law* channel, BRST-exact deformations of the branch generator must cancel on $\mathcal{H}_{\text{phys}}$ provided the Slavnov identity and anomaly filters hold (Section 7.2.8 and eq. (7.16)). We make this precise and then spell out the diagnostic triage.

Proposition 7.181 (Gauge-parameter independence on shell)**.** *Let $\mathcal{C}_\alpha$ be defined by (7.115) (or equivalently by (7.98) on $\mathcal{H}_{\text{phys}}$). Under the assumptions of the Nielsen identity eq. (7.19), BRST-closure of the band knobs eq. (7.30), and quartet/positivity eq. (7.12), one has*

$$\partial_{\alpha_{\text{gf}}}\mathcal{C}_\alpha = 0, \qquad \partial_{\alpha_{\text{gf}}}\Lambda_\alpha = 0 \quad \textit{on shell.}$$

Proof. Differentiate the (embedded) numerator/denominator w.r.t. α_{gf}. By eq. (7.19), each derivative is a sum of a BRST-exact commutator and EOM terms. The commutator vanishes between physical projectors by quartet/positivity; the EOM insertions vanish on shell (eqs. (6.333) and (6.334)). The statement for Λ_α follows by the chain rule. □

Cohomological invariance of the law channel. Consider the homotopy

$$\hat{H}_s(t) := H_0(t) + s\,\zeta\Delta G(t), \qquad s \in [0,1],$$

with H_0 the symmetric-law Hamiltonian of Quadrant I and ΔG a local, band-limited operator. Let $U_s(T)$ be the corresponding band/IR-smoothed propagator eq. (7.130), and define

$$\mathcal{C}_{\text{law}}(s) := \frac{\operatorname{Tr}\Big(\mathsf{P}_{\text{phys}}\,U_{s=1}(T)\,\rho_{\text{phys}}\,U_{s=0}^{\dagger}(T)\,\mathsf{P}_{\text{phys}}\Big)}{\operatorname{Tr}(\mathsf{P}_{\text{phys}}\rho_{\text{phys}})}.$$

Duhamel's formula gives

$$\frac{d}{ds}\,\mathcal{C}_{\text{law}}(s) = -i\int_0^T dt\;\frac{\operatorname{Tr}\Big(\mathsf{P}_{\text{phys}}\,U_1(T,t)\,\Big(\zeta\Delta G(t)\Big)\,U_s(t,0)\,\rho_{\text{phys}}\,U_0^{\dagger}(T)\,\mathsf{P}_{\text{phys}}\Big)}{\operatorname{Tr}(\mathsf{P}_{\text{phys}}\rho_{\text{phys}})}. \tag{7.134}$$

Using $\zeta\Delta G = [\hat{Q}_{\text{BRST}}, \Delta G]$ and cyclicity, the integrand is a BRST commutator plus EOM terms. On $\mathcal{H}_{\text{phys}}$ the commutator vanishes by quartet positivity and the Slavnov identity, while EOM terms vanish on shell (Section 7.2.6 and eqs. (6.333), (6.334) and (7.12)). Hence $d\mathcal{C}_{\text{law}}/ds = 0$ and

$$\mathcal{C}_{\text{law}}(1) = \mathcal{C}_{\text{law}}(0) = 1, \qquad \Lambda_{\text{law}} = 1 \quad \text{(ideal Quadrant I, anomaly-free).} \tag{7.135}$$

This is the operational form of cohomological BRST invariance [442, 117, 91].

Definition 7.182 (Anomaly functional and filter)**.** Let $\mathcal{S}$ denote the Slavnov operator (BRST Ward functional) and suppose the renormalised theory admits a local breaking Δ_{an} so that $\mathcal{S}(\Gamma) = \Delta_{\text{an}}$. Define the *projected anomaly functional* on $\mathcal{H}_{\text{phys}}$ by

$$\mathcal{A}_T \;:=\; \int_0^T dt\,\Big\|\;\mathsf{P}_{\text{phys}}\,\Big(\Delta_{\text{an}}(t)\Big)\,\mathsf{P}_{\text{phys}}\;\Big\|_{\text{op}}.$$

The *anomaly filter* Section 7.2.8 is said to *pass* on $[0,T]$ if $\mathcal{A}_T \le \varepsilon_{\text{an}}$ for a certified tolerance $\varepsilon_{\text{an}} \ge 0$.

Proposition 7.183 (Quantitative law-channel robustness)**.** *Let* $M_{\mathrm{law}} := (U_0^{(\mathrm{law})})^\dagger U_1^{(\mathrm{law})}$ *with* $U_1^{(\mathrm{law})}$ *defined by a BRST-exact deformation of* H_0 *as above. Then, for any physical preparation* ρ_{phys} *with* $\mathrm{Tr}(\mathsf{P}_{\mathrm{phys}}\rho_{\mathrm{phys}}) \geq c_0 > 0$*,*

$$\left|\mathcal{C}_{\mathrm{law}} - 1\right| \;\leq\; \frac{\mathcal{A}_T}{c_0} \;+\; \varepsilon_{\mathrm{EOM}} \;+\; \varepsilon_{\mathrm{reg}},$$

where $\mathcal{A}_T$ *is the anomaly functional of Definition 7.182,* $\varepsilon_{\mathrm{EOM}}$ *controls off-shell EOM residuals, and* $\varepsilon_{\mathrm{reg}}$ *is the regulator tolerance from Section 7.3.7. In particular, if the anomaly filter passes and one is on shell with admissible regularisation, then* $\mathcal{C}_{\mathrm{law}} = 1 + \mathcal{O}(\varepsilon_{\mathrm{reg}})$*.*

Proof. Integrate (7.134) over $s \in [0,1]$, replace $[\hat{Q}_{\mathrm{BRST}}, \Delta G]$ by the Slavnov variation and decompose into (projected) anomaly plus EOM terms. Take operator norms inside the time-ordered insertions using unitarity of partial propagators; use $\mathrm{Tr}(\mathsf{P}\rho\mathsf{P}) \geq c_0$ and add the regulator tolerance from Section 7.3.7. □

Corollary 7.184 (Sector-resolved law-channel bounds)**.** *For any BRST-closed sector projector* Π_β *with weight* $w_\beta > 0$*,*

$$\left|\mathcal{C}_{\mathrm{law}|\beta} - 1\right| \;\leq\; \frac{1}{w_\beta}\,\mathcal{A}_T^{(\beta)} \;+\; \varepsilon_{\mathrm{EOM}} \;+\; \varepsilon_{\mathrm{reg}}, \qquad \mathcal{A}_T^{(\beta)} := \int_0^T dt\, \left\|\Pi_\beta \mathsf{P}_{\mathrm{phys}} \Delta_{\mathrm{an}}(t) \mathsf{P}_{\mathrm{phys}} \Pi_\beta\right\|_{\mathrm{op}}.$$

Proof. Insert Π_β on both sides of the trace in (7.134) and repeat the norm estimate sectorwise. □

Proposition 7.185 (Diagnostic triage for law-channel suppression)**.** *Suppose a certified plateau exhibits* $\left|1 - \mathcal{C}_{\mathrm{law}}\right| > \delta$ *beyond the joint tolerance* $\delta_\star := \varepsilon_{\mathrm{an}} + \varepsilon_{\mathrm{EOM}} + \varepsilon_{\mathrm{reg}}$*. Then at least one of:*

(a) Anomaly: $\mathcal{A}_T > \varepsilon_{\mathrm{an}}$ *(Slavnov identity broken); check Section 7.2.8 and eq. (7.16) and sectorwise diagnostics (Cor. 7.184).*

(b) Off-shell preparation: *EOM residuals exceed tolerance (verify eqs. (6.333) and (6.334) on the band).*

(c) Regulator pathology: *admissibility fails or numeric tolerance too loose; perform the regulator swap test of Section 7.3.7 and use the commuting-limits check eq. (7.45).*

If all three checks pass, re-calibrate the phase and re-run the plateau certification (Section 7.3.9); persistent suppression then constitutes evidence for genuine law-level incoherence.

Proposition 7.186 (Quantitative stability bound)**.** *Let the norm of the BRST-exact perturbation restricted to* $\mathcal{H}_{\mathrm{phys}}$ *be measured by* ϵ_{BRST}*, defined as*

$$\epsilon_{\mathrm{BRST}} := \sup_{t\in[0,T]} \left\|\mathsf{P}_{\mathrm{phys}}\, \zeta \Delta G(t)\, \mathsf{P}_{\mathrm{phys}}\right\|_{\mathrm{op}}.$$

Assume the denominator in (7.115) is normalised to 1, i.e. $\mathrm{Tr}(\mathsf{P}_{\mathrm{phys}}\rho_{\mathrm{phys}}) = 1$ *(otherwise replace RHS below by* $1/c_0$ *with* $c_0 := \mathrm{Tr}(\mathsf{P}_{\mathrm{phys}}\rho_{\mathrm{phys}}) > 0$*). Then, from (7.134) and unitarity,*

$$\left|1 - \mathcal{C}_{\mathrm{law}}\right| \;\leq\; T\,\epsilon_{\mathrm{BRST}} \;+\; \mathcal{O}(T^2\,\epsilon_{\mathrm{BRST}}^2) \;+\; \varepsilon_{\mathrm{reg}}, \tag{7.136}$$

where $\varepsilon_{\mathrm{reg}}$ *accounts for admissible regulator differences (Section 7.3.7). In the anomaly-free case one has* $\epsilon_{\mathrm{BRST}} = 0$ *(cohomological triviality on* $\mathcal{H}_{\mathrm{phys}}$*), recovering (7.135).*

Proof. Integrate (7.134) over $s \in [0,1]$. Using unitarity of the partial propagators $U_s(t_2,t_1)$ and $\|\cdot\|_{\mathrm{op}}$ submultiplicativity,

$$\left|\mathcal{C}_{\mathrm{law}}(1) - \mathcal{C}_{\mathrm{law}}(0)\right| \le \int_0^T dt\, \left\|\mathsf{P}_{\mathrm{phys}}\, \zeta \Delta G(t)\, \mathsf{P}_{\mathrm{phys}}\right\|_{\mathrm{op}} + \mathcal{O}\Big(\int_0^T \int_0^t \epsilon_{\mathrm{BRST}}^2\, ds\, dt\Big) + \varepsilon_{\mathrm{reg}},$$

which gives $T\,\epsilon_{\mathrm{BRST}} + \mathcal{O}(T^2\epsilon_{\mathrm{BRST}}^2) + \varepsilon_{\mathrm{reg}}$. The $\mathcal{O}(T^2\epsilon_{\mathrm{BRST}}^2)$ term comes from the second and higher Duhamel/Picard iterates with two or more insertions of $\zeta\Delta G$ inside the time-ordered exponential. The regulator term follows from Section 7.3.7. □

Remark 7.187 (Normalisation)**.** If $\mathrm{Tr}(\mathsf{P}_{\mathrm{phys}}\rho_{\mathrm{phys}}) = c_0 \neq 1$, the bound becomes $\left|1 - \mathcal{C}_{\mathrm{law}}\right| \le (T/c_0)\,\epsilon_{\mathrm{BRST}} + \mathcal{O}(T^2\epsilon_{\mathrm{BRST}}^2) + \varepsilon_{\mathrm{reg}}$.

Failure modes and diagnostic triage. Observed suppressions of $|\mathcal{C}_\alpha|$ are interpreted through controlled tests:

1. **If $|\mathcal{C}_{\mathrm{law}}| < 1$:**
   (a) **(L1) Anomaly test:** Check eq. (7.18); compute the descent of the anomaly polynomial and re-verify $\mathcal{S}(\Gamma) = 0$ [443, 442].
   (b) **(L2) Gauge scan:** Vary α_{gf}; any residual dependence contradicts Nielsen protection Section 7.2.9.
   (c) **(L3) Regulator swap:** Replace $(P_x, K_{\mathrm{IR}}^{(\ell)}) \to (\tilde{P}_x, \tilde{K}_{\mathrm{IR}}^{(\ell)})$; violations beyond $\varepsilon_{\mathrm{reg}}$ indicate a UV/IR mismatch Section 7.3.7.
   (d) **(L4) Positivity/hyperbolicity:** Re-check eqs. (7.12) and (7.15); pathologies can feed into $|\mathcal{C}_{\mathrm{law}}| < 1$ via unphysical leakage.
2. **If $|\mathcal{C}_{\mathrm{state}}| < 1$:**
   (a) **(S1) Symmetry probe:** Verify $[Q, \rho_{\mathrm{phys}}] = 0$ (up to IR smearing); failure indicates state-level symmetry tension (7.128).
   (b) **(S2) IR uniformity:** Tighten/widen $K_{\mathrm{IR}}^{(\ell)}$; robust decay across plateau scales points to genuine state decoherence rather than IR artefacts.
   (c) **(S3) Sector check:** Inspect $\Lambda_{\alpha|\beta}$ and coherence coefficients (Section 7.3.11); exclude cancellations across sectors.
   (d) **(S4) Gauge/regulator sanity:** Repeat (L2)–(L3) to rule out overlay artefacts.

Proposition 7.188 (Soundness of the triage)**.** *Let $\delta_\star := \varepsilon_{\mathrm{an}} + \varepsilon_{\mathrm{EOM}} + \varepsilon_{\mathrm{reg}}$ be the joint certified tolerance from anomaly/EOM/regulator budgets. If (L2)–(L4) pass and (L1) fails, law-level suppression is attributed to anomaly ($\mathcal{A}_T > \varepsilon_{\mathrm{an}}$); if (L1), (L3), (L4) pass and (L2) fails, gauge fixing is defective; if all (L1)–(L4) pass yet $|1 - \mathcal{C}_{\mathrm{law}}| > \delta_\star$, declare law-level incoherence beyond certified budgets. Analogous statements hold for the state channel with (S1)–(S4), attributing residual suppression to genuine state decoherence when all sanity checks pass.*

Anomaly filters and cohomology. The algebraic conditions $\mathfrak{A}_{\text{gauge}} = \mathfrak{A}_{\text{mixed}} = \mathfrak{A}_{\text{grav}} = \mathbf{0}$ (Section 7.2.8) ensure that the would-be anomaly cocycle $\mathcal{A} \in H^{1,d}(\zeta \mid d)$ is trivial and removable by counterterms (Wess–Zumino consistency; cf. Definition 7.25) [442, 443, 91, 243]. When these filters hold, BRST nilpotency is preserved at the quantum level, guaranteeing that ζ-exact deformations (such as the law-branch difference) do not affect physical matrix elements—hence $\mathcal{C}_{\text{law}} = 1$ on certified plateaus.

Proposition 7.189 (Anomaly filters ⇒ cohomological triviality)**.** *Let $\mathcal{S}(\Gamma) = \Delta_{\text{an}}$ be the Slavnov functional with local breaking Δ_{an}. Assume the anomaly filters of Section 7.2.8 hold, i.e. $\mathfrak{A}_{\text{gauge}} = \mathfrak{A}_{\text{mixed}} = \mathfrak{A}_{\text{grav}} = \mathbf{0}$. Then the Wess–Zumino consistency condition $\zeta\Delta_{\text{an}} = 0 \pmod d$ implies*

$$[\Delta_{\text{an}}] = 0 \quad \textit{in} \quad H^{1,d}(\zeta \mid d),$$

hence there exist local counterterms Z (ghost number 0) and a $(d-1)$-form Ω such that

$$\Delta_{\text{an}} = \zeta\,\mathsf{Z} + d\Omega,$$

with Z as in Definition 7.25. The renormalised action $\Gamma' := \Gamma - \mathsf{Z}$ obeys the exact Slavnov identity $\mathcal{S}(\Gamma') = 0$ (cf. eq. (7.16)), and the renormalised BRST differential ζ is nilpotent on $\mathcal{H}_{\text{phys}}$.

Proof. The vanishing of the gauge, mixed, and gravitational anomaly polynomials implies that all potential descent representatives in the ζ-mod-d bicomplex are trivial in $H^{1,d}(\zeta \mid d)$ [442, 443]. By Wess–Zumino consistency, there exist local functionals Z of ghost number 0 and $(d-1)$-forms Ω such that $\Delta_{\text{an}} = \zeta\,\mathsf{Z} + d\Omega$; see Definition 7.25 for the precise descent relation and loopwise version. Adding the counterterm $-\mathsf{Z}$ yields the renormalised action $\Gamma' = \Gamma - \mathsf{Z}$ with $\mathcal{S}(\Gamma') = 0$ (cf. eq. (7.16)), which restores quantum BRST nilpotency on $\mathcal{H}_{\text{phys}}$. □

Proposition 7.190 (Operational consequence for the law channel)**.** *Suppose the anomaly filters hold and let the law-channel branch deformation be BRST-exact, $\zeta\Delta G(t)$, as in Section 7.4.2. Then for band/IR-smoothed propagators $U_{0,1}^{(\text{law})}(T)$ one has, on $\mathcal{H}_{\text{phys}}$,*

$$\mathsf{P}_{\text{phys}}(U_0^{(\text{law})})^\dagger U_1^{(\text{law})}\mathsf{P}_{\text{phys}} = \mathsf{I}_{\mathcal{H}_{\text{phys}}}, \qquad \Rightarrow \quad \mathcal{C}_{\text{law}} = 1, \quad \Lambda_{\text{law}} = 1 \quad \textit{on certified plateaus.}$$

Proof. By Prop. 7.189, ζ acts nilpotently on the renormalised theory and ζ-exact insertions decouple between physical projectors (quartet/positivity). In the Dyson expansion of $(U_0^{(\text{law})})^\dagger U_1^{(\text{law})}$ every occurrence of $\zeta\Delta G$ gives a BRST commutator whose contribution vanishes on $\mathcal{H}_{\text{phys}}$. Hence the overlap reduces to the identity on $\mathcal{H}_{\text{phys}}$, yielding $\mathcal{C}_{\text{law}} = 1$. □

Corollary 7.191 (Quantitative corollary with regulator tolerance)**.** *Under the assumptions above and admissible regularisation, the certified readout satisfies $\mathcal{C}_{\text{law}} = 1 + \mathcal{O}(\varepsilon_{\text{reg}})$ and $\Lambda_{\text{law}} = 1 + \mathcal{O}(\varepsilon_{\text{reg}})$, with ε_{reg} from Section 7.3.7.*

Summary. *Gauge robustness* (Nielsen) and *BRST robustness* (anomaly-free Slavnov identity) together imply that the holosymmetric probe's law channel is unity on shell, while the state channel cleanly diagnoses state-level symmetry tension or decoherence. Any departure is actionable via the triage above and must be reconciled before using the visibilities for mode identification [117, 91, 443, 442, 82, 241].

7.4.5 Plateau Coupling and Certification

Within any certified joint plateau $\mathfrak{P}_{\{\text{law,state}\}}$ (cf. Section 7.3), the channel visibilities obey the uniform bounds

$$\left\|\nabla_{(\lambda_{\mathrm{UV}},\ell)}\,\Lambda_\alpha\right\| \le \varepsilon_{\text{grad}}, \qquad \|\mathbf{H}\Lambda_\alpha\|_{\text{op}} \le \varepsilon_{\text{curv}}, \qquad \operatorname{osc}_{B_r}\Lambda_\alpha \le \varepsilon_{\text{osc}}, \tag{7.137}$$

and are regulator-stable in the sense of eq. (7.67). Let $\mathbf{\Lambda} := (\Lambda_{\text{law}}, \Lambda_{\text{state}})^\top$ and work in the log chart $(u, v) = (\ln \lambda_{\mathrm{UV}}, \ln \ell)$ (so $D_\lambda = \partial_u$, $D_\ell = \partial_v$).

Definition 7.192 (Joint plateau and geodesic balls)**.** Let $(u_\star, v_\star) \in \mathfrak{P}_{\{\text{law,state}\}}$. For $\rho > 0$, set $B_\rho(u_\star, v_\star) := \{(u, v) : \|(u, v) - (u_\star, v_\star)\| \le \rho\}$ with the Euclidean norm. The joint plateau is *certified* if (7.137) holds for both channels on every such ball with $\rho \le \rho_{\max}$ and the regulator bound of eq. (7.67) holds uniformly.

Local drift bounds and joint certification. For any (u, v) within a geodesic ball $B_\rho(u_\star, v_\star) \subset \mathfrak{P}_{\{\text{law,state}\}}$,

$$\|\mathbf{\Lambda}(u, v) - \mathbf{\Lambda}(u_\star, v_\star)\| \;\le\; \sqrt{2}\,\varepsilon_{\text{grad}}\,\rho \;+\; \tfrac{1}{2}\,\varepsilon_{\text{curv}}\,\rho^2 \;+\; \varepsilon_{\text{osc}} \;+\; \varepsilon_{\text{reg}}, \tag{7.138}$$

where $\|\cdot\|$ is the Euclidean norm in $\mathbb{R}^2$. The first two terms follow from mean-value/Taylor estimates using eq. (7.137); the last two account for local oscillation and regulator robustness. This bound couples the channels: a single radius ρ controls both simultaneously.

Proposition 7.193 (Vector drift via Jacobian/Hessian control)**.** *Let $F(u, v) := \mathbf{\Lambda}(u, v)$ and $\delta z := (u - u_\star, v - v_\star)$. Along any straight path $t \mapsto (u_\star, v_\star) + t\,\delta z$, $t \in [0, 1]$, one has*

$$\begin{aligned} F(u, v) - F(u_\star, v_\star) = &\int_0^1 J\Big((u_\star, v_\star) + t\,\delta z\Big)\,\delta z\,dt \\ &+ \frac{1}{2}\int_0^1 (1 - t)\,\mathbf{H}_F\Big((u_\star, v_\star) + t\,\delta z\Big)[\delta z, \delta z]\,dt \\ &+ \Delta_{\text{osc}} + \Delta_{\text{reg}}, \end{aligned}$$

where J is the Jacobian and $\mathbf{H}_F$ the block Hessian of F, and the remainders satisfy $\|\Delta_{\text{osc}}\| \le \varepsilon_{\text{osc}}$ and $\|\Delta_{\text{reg}}\| \le \varepsilon_{\text{reg}}$. Moreover, $\|J\|_{\text{op}} \le \sqrt{2}\,\varepsilon_{\text{grad}}$ and $\|\mathbf{H}_F\|_{\text{op}} \le \varepsilon_{\text{curv}}$, whence (7.138).

Proof. Use the integral mean-value/Taylor formula for vector-valued maps. The matrix J has rows $\nabla\Lambda_{\text{law}}^\top$ and $\nabla\Lambda_{\text{state}}^\top$, each bounded by $\varepsilon_{\text{grad}}$, so $\|J\|_{\text{op}} \le \|J\|_{\text{F}} \le \sqrt{2}\,\varepsilon_{\text{grad}}$. The block Hessian $\mathbf{H}_F$ is block-diagonal with blocks $\mathbf{H}\Lambda_\alpha$; its operator norm is the maximum of the block norms, bounded by $\varepsilon_{\text{curv}}$. The path length is $\|\delta z\| = \rho$. Bounds on the oscillation and regulator terms follow from Definition 7.192 and eq. (7.67). □

Coupling metrics. Define the Jacobian and the curvature couplings at $(u_\star, v_\star)$,

$$J_\star = \begin{pmatrix} D_\lambda\Lambda_{\text{law}} & D_\ell\Lambda_{\text{law}} \\ D_\lambda\Lambda_{\text{state}} & D_\ell\Lambda_{\text{state}} \end{pmatrix}_{(u_\star,v_\star)}, \qquad \mathcal{K}_\star := \max_{\alpha\in\{\text{law,state}\}} \|\mathbf{H}\Lambda_\alpha\|_{\text{op},(u_\star,v_\star)},$$

and the *gradient alignment* (coherence) coefficient

$$\chi_\star := \frac{\left|\langle \nabla\Lambda_{\text{law}}, \nabla\Lambda_{\text{state}}\rangle\right|}{\|\nabla\Lambda_{\text{law}}\|\,\|\nabla\Lambda_{\text{state}}\|} \in [0,1],$$

interpreted as the cosine of the angle between the channel gradients. Joint plateaus with small $\|J_\star\|$ are robust to knob perturbations; requiring $\chi_\star \le \chi_{\max}$ (e.g. $1/2$) prevents near-cancellations from masquerading as plateaus (sectorwise safeguards in Section 7.3.11).

Lemma 7.194 (Alignment safeguard)**.** *If* $\max_\alpha \|\nabla\Lambda_\alpha\| \le \varepsilon_{\text{grad}}$ *and* $\chi_\star \le \chi_{\max} < 1$*, then for any unit vector* $\hat n$ *in knob space,*

$$\left\|J_\star\,\hat n\right\| \;\le\; \sqrt{\varepsilon_{\text{grad}}^2 + \varepsilon_{\text{grad}}^2 + 2\,\chi_{\max}\,\varepsilon_{\text{grad}}^2} \;=\; \sqrt{2(1+\chi_{\max})}\,\varepsilon_{\text{grad}}.$$

Thus choosing $\chi_{\max} \le 1/2$ *improves the linear drift constant from* $\sqrt{2}$ *to* $\sqrt{3} < 1.732$*.*

Proof. Let $g_1 := \nabla\Lambda_{\text{law}}\cdot\hat n$, $g_2 := \nabla\Lambda_{\text{state}}\cdot\hat n$. Then $\|J_\star\hat n\|^2 = g_1^2 + g_2^2 \le \|\nabla\Lambda_{\text{law}}\|^2 + \|\nabla\Lambda_{\text{state}}\|^2 + 2|g_1 g_2| \le 2\varepsilon_{\text{grad}}^2 + 2\chi_{\max}\varepsilon_{\text{grad}}^2$. □

Certified operating point and reporting. Choose $(\lambda_\star, \ell_\star)$ (equivalently $(u_\star, v_\star)$) to maximise the inscribed radius $\rho_\star$ subject to $\|J_\star\|$ and $\mathcal{K}_\star$ minimisation.[4] The reported values are

$$\Lambda^\star_{\text{law}} := \Lambda_{\text{law}}(\lambda_\star, \ell_\star; T), \qquad \Lambda^\star_{\text{state}} := \Lambda_{\text{state}}(\lambda_\star, \ell_\star; T), \tag{7.139}$$

with a deterministic *plateau uncertainty* bounded by the right-hand side of (7.138) evaluated at $\rho = \rho_\star$, added in quadrature with the statistical error from Section 7.4.6. Gauge/BRST independence of these readouts on shell follows from Section 7.3.6, and regulator stability from Section 7.3.7 [117, 91, 114].

Proposition 7.195 (Existence of a certified operating point)**.** *If the certified joint plateau* $\mathfrak{P}_{\{\text{law,state}\}}$ *is nonempty, bounded, and closed in the log chart, then an operating point* $(u_\star, v_\star)$ *maximising* $\rho_\star$ *exists. Among maximisers, there exists one minimising* $\|J_\star\|$ *and* $\mathcal{K}_\star$*.*

Proof. Compactness ensures existence of a Chebyshev center (maximiser of the inscribed radius). Continuity of $J_\star$ and $\mathcal{K}_\star$ on the compact argmax set yields a minimiser by the extreme value theorem. □

Corollary 7.196 (Deterministic plateau uncertainty)**.** *Let* Δ_{det} *denote the deterministic plateau uncertainty at* $(u_\star, v_\star)$*. Then*

$$\Delta_{\text{det}} \;\le\; \sqrt{2}\,\varepsilon_{\text{grad}}\,\rho_\star \;+\; \tfrac{1}{2}\,\varepsilon_{\text{curv}}\,\rho_\star^2 \;+\; \varepsilon_{\text{osc}} \;+\; \varepsilon_{\text{reg}},$$

and the total reported uncertainty is $\sqrt{\Delta_{\text{det}}^2 + \Delta_{\text{stat}}^2}$ *with* Δ_{stat} *from Section 7.4.6.*

[4] Ties can be broken by minimising $\|\nabla\Lambda_{\text{law}}\|^2 + \|\nabla\Lambda_{\text{state}}\|^2$ or by maximising the regulator overlap (smallest observed drift under $(P,K)\to(\tilde P,\tilde K)$).

7.4.6 Error Budget and Robustness to Noise

Let N_x (N_y) be the number of ancilla shots used to estimate $\langle\sigma_x\rangle_\alpha$ ($\langle\sigma_y\rangle_\alpha$). Each outcome is ±1, so the standard errors are (binomial statistics)

$$\delta\langle\sigma_x\rangle_\alpha = \sqrt{\frac{1-\langle\sigma_x\rangle_\alpha^2}{N_x}}, \qquad \delta\langle\sigma_y\rangle_\alpha = \sqrt{\frac{1-\langle\sigma_y\rangle_\alpha^2}{N_y}}, \tag{7.140}$$

see, e.g., [287]. Propagating errors to the coherence magnitude $|\mathcal{C}_\alpha| = \sqrt{\langle\sigma_x\rangle_\alpha^2 + \langle\sigma_y\rangle_\alpha^2}$ gives (delta method, neglecting cross-covariance if x and y runs are disjoint)

$$\delta|\mathcal{C}_\alpha| \;\approx\; \frac{1}{|\mathcal{C}_\alpha|}\sqrt{\langle\sigma_x\rangle_\alpha^2\,\delta\langle\sigma_x\rangle_\alpha^2 + \langle\sigma_y\rangle_\alpha^2\,\delta\langle\sigma_y\rangle_\alpha^2}\,, \qquad \delta\Lambda_\alpha \le \delta|\mathcal{C}_\alpha|. \tag{7.141}$$

If $N_x = N_y = N$, (7.140)–(7.141) reduce to the compact form used in Section 7.4.1. For small samples, exact Clopper–Pearson intervals can replace (7.140) (optional).

Proposition 7.197 (Shot-noise formulas and non-asymptotic bounds)**.** *Let $X_k \in \{\pm1\}$ be i.i.d. with mean $\mu_x = \langle\sigma_x\rangle_\alpha$. Then $\widehat{\mu}_x = \frac{1}{N_x}\sum_k X_k$ satisfies $\mathrm{Var}(\widehat{\mu}_x) = \frac{1-\mu_x^2}{N_x}$, yielding (7.140). The same holds for Y_ℓ estimating $\langle\sigma_y\rangle_\alpha$. Moreover, for any $\delta \in (0,1)$, Hoeffding gives a distribution-free concentration:*

$$\Pr\Big(|\widehat{\mu}_x - \mu_x| \ge \epsilon\Big) \le 2\,e^{-2N_x\epsilon^2} \quad\Rightarrow\quad \epsilon = \sqrt{\frac{\ln(2/\delta)}{2N_x}},$$

which can replace Gaussian error bars when needed.

Lemma 7.198 (Delta-method with optional covariance)**.** *If the x and y estimates are obtained from independent shot sets, (7.141) holds. If a joint scheme induces $\mathrm{Cov}(\widehat{\mu}_x, \widehat{\mu}_y) \neq 0$, then*

$$\delta|\mathcal{C}_\alpha| \;\approx\; \frac{1}{|\mathcal{C}_\alpha|}\sqrt{\mu_x^2\,\delta\mu_x^2 + \mu_y^2\,\delta\mu_y^2 + 2\,\mu_x\mu_y\,\mathrm{Cov}(\widehat{\mu}_x,\widehat{\mu}_y)}\,,$$

with $\mu_{x,y} := \langle\sigma_{x,y}\rangle_\alpha$ and $\delta\mu_{x,y}$ from (7.140). The bound $\delta\Lambda_\alpha \le \delta|\mathcal{C}_\alpha|$ follows since Λ_α is a phase-rotated projection of $\mathcal{C}_\alpha$ onto $\mathbb{R}$.

Corollary 7.199 (Equal-shot simplification and worst-case bound)**.** *If $N_x = N_y = N$ and runs are independent, then*

$$\delta|\mathcal{C}_\alpha| \;\approx\; \frac{1}{\sqrt{N}}\sqrt{1 - \frac{|\mathcal{C}_\alpha|^2}{2}} \;\le\; \frac{1}{\sqrt{N}},$$

where the inequality is a worst-case bound over all (μ_x, μ_y) with $\mu_x^2 + \mu_y^2 = |\mathcal{C}_\alpha|^2$.

Proposition 7.200 (Shot allocation optimality)**.** *Given a fixed total budget $N_{\mathrm{tot}} = N_x + N_y$ and independent runs, the first-order allocation minimising $\delta|\mathcal{C}_\alpha|$ is*

$$\frac{N_x}{N_y} \;\approx\; \frac{|\mu_x|\sqrt{1-\mu_x^2}}{|\mu_y|\sqrt{1-\mu_y^2}}\,.$$

For $|\mu_x| \approx |\mu_y|$ this reduces to $N_x \approx N_y$.

Phase uncertainty. If $\Lambda_\alpha = \mathrm{Re}(e^{-i\phi_\alpha}\mathcal{C}_\alpha)$ is reported, a phase uncertainty $\delta\phi_\alpha$ (from calibration at $(\lambda_\star, \ell_\star)$) contributes

$$(\delta\Lambda_\alpha)_\phi \;\approx\; |\mathcal{C}_\alpha|\,\delta\phi_\alpha,$$

to be added in quadrature with (7.141). Echo-style phase choices (common-mode cancellation) minimise $\delta\phi_\alpha$ without affecting $|\mathcal{C}_\alpha|$.

Proposition 7.201 (Phase-error bound)**.** *Let $\Lambda_\alpha = \mathrm{Re}(e^{-i\phi_\alpha}\mathcal{C}_\alpha)$ with $\mathcal{C}_\alpha = |\mathcal{C}_\alpha|e^{i\theta_\alpha}$. Then $\partial_{\phi_\alpha}\Lambda_\alpha = -|\mathcal{C}_\alpha|\sin(\theta_\alpha - \phi_\alpha)$, so $|(\delta\Lambda_\alpha)_\phi| \le |\mathcal{C}_\alpha|\,\delta\phi_\alpha$, with equality at quadrature ($\theta_\alpha - \phi_\alpha = \pm\pi/2$).*

SPAM calibration (affine correction). Static state-preparation-and-measurement (SPAM) errors act as an affine map on the Bloch-plane readout,

$$\begin{pmatrix}\langle\sigma_x\rangle\\ \langle\sigma_y\rangle\end{pmatrix}_{\mathrm{meas}} = A\begin{pmatrix}\langle\sigma_x\rangle\\ \langle\sigma_y\rangle\end{pmatrix}_{\mathrm{true}} + \mathbf{b},$$

with an invertible 2×2 matrix A and offset $\mathbf{b}$. Two-phase calibration at $(\lambda_\star, \ell_\star)$ with known branch identities $U_{0,1}^{(\alpha)} = \nVdash$ (and a $\pi/2$ phase-shifted variant) identifies $(A, \mathbf{b})$; corrected estimates use $(A^{-1}, -A^{-1}\mathbf{b})$, and the associated calibration uncertainty is folded into the tolerance budget of Section 7.3.9.

Proposition 7.202 (Unbiased affine SPAM correction and amplification factor)**.** *If $(\widehat{A}, \widehat{\mathbf{b}}) = (A, \mathbf{b})$ exactly, the corrected estimator is unbiased: $\widehat{\mathbf{s}}_{\mathrm{true}} = A^{-1}(\mathbf{s}_{\mathrm{meas}} - \mathbf{b})$ with $\mathbb{E}[\widehat{\mathbf{s}}_{\mathrm{true}}] = \mathbf{s}_{\mathrm{true}}$. If $(\widehat{A}, \widehat{\mathbf{b}})$ have errors $(\Delta A, \Delta\mathbf{b})$ with $\|A^{-1}\Delta A\| \le \eta < 1$, then*

$$\frac{\|\delta\widehat{\mathbf{s}}_{\mathrm{true}}\|}{\|\delta\mathbf{s}_{\mathrm{meas}}\|} \;\le\; \kappa_2(A) \;+\; \mathcal{O}(\eta), \qquad \kappa_2(A) := \|A\|_2\,\|A^{-1}\|_2,$$

so calibration inflates statistical errors by at most the condition number $\kappa_2(A)$ (to first order), and introduces a bias of order $\|A^{-1}\Delta\mathbf{b}\| + \|A^{-1}\Delta A\|\,\|\mathbf{s}_{\mathrm{meas}} - \mathbf{b}\|$.

Total uncertainty and acceptance. For each channel at $(\lambda_\star, \ell_\star)$ report

$$\varepsilon_{\mathrm{tot},\alpha} := \sqrt{\Big(\sqrt{2}\,\varepsilon_{\mathrm{grad}}\,\rho_\star + \tfrac{1}{2}\varepsilon_{\mathrm{curv}}\,\rho_\star^2 + \varepsilon_{\mathrm{osc}} + \varepsilon_{\mathrm{reg}}\Big)^2 + \Big(\delta|\mathcal{C}_\alpha|\Big)^2 + \Big((\delta\Lambda_\alpha)_\phi\Big)^2},$$

and accept the certification only if $\varepsilon_{\mathrm{tot},\alpha}$ lies below the policy threshold set for downstream mode-identification. Gauge/BRST invariance and regulator robustness of the estimates are ensured by Sections 7.3.6 and 7.3.7 [117, 91, 114].

Proposition 7.203 (Decision rule soundness)**.** *Under the joint plateau assumptions and independent shot allocations, the acceptance rule $\varepsilon_{\mathrm{tot},\alpha} \le \varepsilon_{\mathrm{policy}}$ controls the total (deterministic+statistical+phase) deviation in the reported $\Lambda_\alpha^\star$ at level $\varepsilon_{\mathrm{policy}}$, with SPAM-inflation absorbed via $\kappa_2(A)$ in $\delta|\mathcal{C}_\alpha|$.*

Remark (provisional mode tags). Any mode tagging inferred from $\Lambda_{\mathrm{law}}^\star, \Lambda_{\mathrm{state}}^\star$ (e.g., "graviton-like," "Higgs-like," "apeiron," "chrono–kymon") is *provisional* until validated across quadrants per the stated policy; tags must not be treated as definitive until multi-quadrant plateau compatibility is demonstrated.

7.5 Exact Identities at the Symmetry Point (Baseline for Symmetry Breaking)

This section records the exact Ward, Slavnov–Taylor, and BRST/cohomological identities that hold in Quadrant I at the G-invariant vacuum fixed in Section 7.1. All statements are formulated *on the band* with the UV/IR knobs of Section 7.3 and are to be read on the physical sector $\mathcal{H}_{\rm phys}$ introduced in Section 7.2. They provide the baseline against which symmetry-breaking effects in other quadrants will be measured. Our conventions follow algebraic renormalization and BRST quantization [117, 442, 91].

7.5.1 Band-limited Source Formalism and BRST-exact Gauge Fixing

Introduce sources for the band-limited fields and their BRST variations:

$$\begin{aligned} \mathcal{J}\cdot\varphi &:= J_\Phi\cdot\hat{\mathbf{\Phi}}_{\lambda_{\rm UV}} + J_\Theta\cdot\hat{\mathbf{\Theta}}_{\lambda_{\rm UV}} + \bar\eta\cdot c + \eta\cdot\bar c + B\cdot b, \\ \mathcal{K}\cdot\zeta\varphi &:= K_\Phi\cdot\zeta\hat{\mathbf{\Phi}}_{\lambda_{\rm UV}} + K_\Theta\cdot\zeta\hat{\mathbf{\Theta}}_{\lambda_{\rm UV}} + K_c\cdot\zeta c + K_{\bar c}\cdot\zeta\bar c + K_b\cdot\zeta b\,, \end{aligned} \tag{7.142}$$

where "$\cdot$" includes spacetime integration with the IR window $K_{\rm IR}^{(\ell)}$ and implicit traces, and all fields are projected by $P_x(\lambda_{\rm UV})$ as in Section 7.3.1. The total action is

$$S_{\rm tot} = S_{\Phi,\Theta}^{(\lambda_{\rm UV},\ell)} \;+\; \zeta\Psi \;+\; \mathcal{J}\cdot\varphi \;+\; \mathcal{K}\cdot\zeta\varphi, \tag{7.143}$$

with $\zeta\Psi$ the BRST-exact gauge-fixing/ghost term (ghost number 0), cf. Section 7.1.3. Define

$$Z[\mathcal{J},\mathcal{K}] := \int\mathcal{D}\varphi\;\exp\Big\{i\,S_{\rm tot}\Big\}, \qquad W := -i\ln Z, \qquad \Gamma := \mathsf{Legendre}_{\mathcal{J}}(W) - \mathcal{K}\cdot\zeta\varphi, \tag{7.144}$$

where the Legendre transform is with respect to $\mathcal{J}$ at fixed $\mathcal{K}$. Band-limiting and IR smearing commute with the BRST charge (ghost number 0); in particular,

$$[\hat Q_{\rm BRST},\, P_x(\lambda_{\rm UV})] = 0, \qquad \zeta\Big(K_{\rm IR}^{(\ell)} * \,\cdot\,\Big) = 0, \tag{7.145}$$

so the source formalism descends consistently to $\mathcal{H}_{\rm phys}$.

Definition 7.204 (Band-limited Slavnov functional)**.** Write $\varphi = (\hat{\mathbf{\Phi}}_{\lambda_{\rm UV}}, \hat{\mathbf{\Theta}}_{\lambda_{\rm UV}}, c, \bar c, b)$ and $\mathcal{K} = (K_\Phi, K_\Theta, K_c, K_{\bar c}, K_b)$ with ghost-number assignments standard in BRST ($\mathrm{gh}(c) = 1$, $\mathrm{gh}(\bar c) = -1$, $\mathrm{gh}(b) = 0$, and $\mathrm{gh}(K_\phi) = -1 - \mathrm{gh}(\phi)$). For any functional $F[\varphi,\mathcal{K}]$ define the (band-limited) Zinn–Justin antibracket

$$(F,G) := \frac{\delta_r F}{\delta\varphi}\cdot\frac{\delta_l G}{\delta\mathcal{K}} - \frac{\delta_r F}{\delta\mathcal{K}}\cdot\frac{\delta_l G}{\delta\varphi},$$

and the Slavnov functional

$$\mathcal{S}(F) := \frac{1}{2}\,(F,F).$$

The dot-contractions include the IR kernel $K_{\rm IR}^{(\ell)}$ and UV projector $P_x(\lambda_{\rm UV})$ as in Section 7.3.1.

Proposition 7.205 (Band Slavnov–Taylor identities for W and Γ)**.** *Under BRST invariance of the measure and* (7.145)*, the connected generator W and the 1PI functional Γ satisfy*

$$\mathcal{S}(W) = 0, \qquad \mathcal{S}(\Gamma) = 0, \tag{7.146}$$

exactly on the band in Quadrant I.

Proof. Consider the BRST change of variables $\varphi \mapsto \varphi + \epsilon\,\zeta\varphi$ in the path integral defining Z; the Jacobian is unity by standard arguments, while $\zeta(S_{\mathrm{tot}}) = \zeta(\zeta\Psi) + \zeta(\mathcal{J}\cdot\varphi) + \zeta(\mathcal{K}\cdot\zeta\varphi) = 0$ since $\zeta^2 = 0$, $\zeta(\mathcal{J}\cdot\varphi) = \mathcal{J}\cdot\zeta\varphi$, and the $\mathcal{K}$-term cancels it by construction. Band-limited commutation (7.145) ensures no extra terms arise. Passing to W and performing the Legendre transform at fixed $\mathcal{K}$ yields (7.146). □

Lemma 7.206 (Linearised Slavnov operator)**.** *Define $\mathcal{B}_\Gamma X := (\Gamma, X)$. If $\mathcal{S}(\Gamma) = 0$, then $\mathcal{B}_\Gamma^2 = 0$. Moreover, $\mathcal{B}_\Gamma$ implements the quantum BRST differential on composite operators, restricted to the band.*

Proof. Jacobi identity for the antibracket implies $(\Gamma, (\Gamma, X)) = \frac{1}{2}((\Gamma,\Gamma), X) = 0$ when $\mathcal{S}(\Gamma) = 0$. □

Proposition 7.207 (Band-limited Ward identities for global G)**.** *Let $\delta^a_G\varphi = R^a[\varphi]$ be the (infinitesimal) global G-symmetry at the G-invariant vacuum of Quadrant I, with the band-limited action $S^{(\lambda_{\mathrm{UV}},\ell)}_{\Phi,\Theta}$ invariant and Ψ chosen G-invariant. Then*

$$\mathcal{W}_a(\Gamma) = 0, \qquad \mathcal{W}_a := \frac{\delta\Gamma}{\delta\varphi}\cdot R^a[\varphi], \tag{7.147}$$

and analogously for W. If the background-field split is used, (7.147) *becomes the background Ward identity with band knobs inserted.*

Proof. Repeat the change-of-variables argument using δ^a_G in the band-limited path integral; invariance of the measure and G-invariance of $S^{(\lambda_{\mathrm{UV}},\ell)}_{\Phi,\Theta} + \zeta\Psi$ give the identity. The band operations commute with δ^a_G by construction. □

Proposition 7.208 (Nielsen (gauge-parameter) identity on the band)**.** *Let α_{gf} enter only through the gauge fermion $\Psi(\alpha_{\mathrm{gf}})$. Then*

$$\partial_{\alpha_{\mathrm{gf}}}\Gamma = -\big(\Gamma, \partial_{\alpha_{\mathrm{gf}}}\Psi\big) = -\,\mathcal{B}_\Gamma\big(\partial_{\alpha_{\mathrm{gf}}}\Psi\big), \tag{7.148}$$

with all products band-limited as above. Consequently, $\partial_{\alpha_{\mathrm{gf}}}\Gamma$ is $\mathcal{B}_\Gamma$-exact, and physical, BRST-closed observables are gauge-parameter independent.

Proof. Differentiate Z with respect to α_{gf}; only $\zeta\Psi$ varies. Using the antibracket/linearised identity $\mathcal{B}_\Gamma X = (\Gamma, X)$ (Prop. 7.212) with $X = \partial_{\alpha_{\mathrm{gf}}}\Psi$, and the band commutation (7.145), yields (7.148). □

Proposition 7.209 (Ghost, antighost, and b equations)**.** *Assuming the usual linear dependence of Ψ on $\bar{c}$ and b, one obtains (band-limited) equations of motion*

$$\frac{\delta\Gamma}{\delta b} = \mathcal{G}_b, \qquad \frac{\delta\Gamma}{\delta\bar{c}} = \mathcal{G}_{\bar{c}}, \qquad \frac{\delta\Gamma}{\delta c} = \mathcal{G}_c, \tag{7.149}$$

where $\mathcal{G}_\bullet$ are local functionals fixed by the chosen gauge and knobs. These identities are compatible with (7.146) *and* (7.148)*.*

Proof. Direct variation of (7.143) with respect to the corresponding fields; the band operations commute with these algebraic variations. □

Proposition 7.210 (Cohomology baseline on the band)**.** *Let* $\mathsf{H}^0(\mathcal{B}_\Gamma)$ *denote the cohomology at ghost number* 0 *of the linearised operator* $\mathcal{B}_\Gamma$ *defined on the band. Then* $\mathsf{H}^0(\mathcal{B}_\Gamma)$ *is isomorphic to the space of gauge-invariant, G-invariant local functionals modulo band-limited EOMs. In particular, physical insertions and S-matrix elements restricted to* $\mathcal{H}_{\text{phys}}$ *are represented by* $\mathcal{B}_\Gamma$*-closed classes.*

Proof. Standard algebraic renormalisation/cohomology arguments apply verbatim since (7.145) ensures the band operations preserve ghost number and commute with ζ; locality is preserved by the L^1 IR window and projector locality. See [117, 442]. □

Remark 7.211 (Exactly at the symmetry point)**.** At the G-invariant vacuum of Quadrant I, (7.147), (7.146), and (7.148) together constitute the exact baseline: *any* departure observed in other quadrants must be attributed to (i) controlled, certified symmetry breaking (law or state), (ii) anomaly terms excluded by the filters of Section 7.2.8, or (iii) violations of admissibility/regularisation stability addressed in Section 7.3.7.

7.5.2 Slavnov–Taylor Identity and Linearized Operator

Introduce the Slavnov functional (Zinn–Justin identity) for the band-limited 1PI action:

$$\mathcal{S}(\Gamma) := \int d^dx \left[\frac{\delta\Gamma}{\delta\Phi}\frac{\delta\Gamma}{\delta K_\Phi} + \frac{\delta\Gamma}{\delta\Theta}\frac{\delta\Gamma}{\delta K_\Theta} + \frac{\delta\Gamma}{\delta c}\frac{\delta\Gamma}{\delta K_c} + \frac{\delta\Gamma}{\delta \bar{c}}\frac{\delta\Gamma}{\delta K_{\bar{c}}} + \frac{\delta\Gamma}{\delta b}\frac{\delta\Gamma}{\delta K_b} \right] = 0, \tag{7.150}$$

which is the band/IR version of eq. (7.16). Following [117, 442], define the *linearized* Slavnov operator acting on a functional X:

$$\begin{aligned} \mathcal{B}_\Gamma X := \int d^dx \Bigg[& \frac{\delta\Gamma}{\delta\Phi}\frac{\delta X}{\delta K_\Phi} + \frac{\delta X}{\delta\Phi}\frac{\delta\Gamma}{\delta K_\Phi} + \frac{\delta\Gamma}{\delta\Theta}\frac{\delta X}{\delta K_\Theta} + \frac{\delta X}{\delta\Theta}\frac{\delta\Gamma}{\delta K_\Theta} \\ & + \frac{\delta\Gamma}{\delta c}\frac{\delta X}{\delta K_c} + \frac{\delta X}{\delta c}\frac{\delta\Gamma}{\delta K_c} + \frac{\delta\Gamma}{\delta \bar{c}}\frac{\delta X}{\delta K_{\bar{c}}} + \frac{\delta\Gamma}{\delta b}\frac{\delta X}{\delta K_b} \Bigg]. \end{aligned} \tag{7.151}$$

If $\mathcal{S}(\Gamma) = 0$, then $\mathcal{B}_\Gamma^2 = 0$ (nilpotent) and cohomology classes $H^0(\mathcal{B}_\Gamma)$ characterize physical, gauge-invariant composite operators. These statements hold *on the band* by (7.145) and ensure that BRST-exact insertions do not contribute to physical matrix elements in $\mathcal{H}_{\text{phys}}$ [117, 91].

Proposition 7.212 (Antibracket form and equivalence)**.** *Let* (F, G) *denote the (band-limited) Zinn–Justin antibracket defined as in Definition 7.204. Then*

$$\mathcal{S}(\Gamma) = \tfrac{1}{2}(\Gamma, \Gamma), \qquad \mathcal{B}_\Gamma X = (\Gamma, X),$$

with all contractions understood with the UV projector $P_x(\lambda_{\text{UV}})$ *and IR window* $K_{\text{IR}}^{(\ell)}$*. Hence* (7.150) *is equivalent to* $(\Gamma, \Gamma) = 0$*.*

Proof. A direct comparison of the components in (7.150) and (7.151) with the antibracket definitions (including the source sector) yields the stated identities; band operations commute with functional differentiation by (7.145). □

Lemma 7.213 (Nilpotency of the linearized operator)**.** *If $\mathcal{S}(\Gamma) = 0$, then $\mathcal{B}_\Gamma^2 = 0$ on the space of local integrated functionals (band-limited).*

Proof. Using Proposition 7.212 and the graded Jacobi identity, $\mathcal{B}_\Gamma^2 X = (\Gamma, (\Gamma, X)) = \frac{1}{2}((\Gamma, \Gamma), X) = 0$ when $(\Gamma, \Gamma) = 0$. □

Proposition 7.214 (Ghost number and cohomological gradings)**.** *Assign* $\mathrm{gh}(\Phi) = \mathrm{gh}(\Theta) = \mathrm{gh}(b) = 0$, $\mathrm{gh}(c) = 1$, $\mathrm{gh}(\bar{c}) = -1$, *and* $\mathrm{gh}(K_\phi) = -1 - \mathrm{gh}(\phi)$. *Then* $\mathrm{gh}(\mathcal{B}_\Gamma) = +1$, *so* $\mathcal{B}_\Gamma : \mathsf{F}^g \to \mathsf{F}^{g+1}$. *The cohomology* $\mathsf{H}^g(\mathcal{B}_\Gamma)$ *is thus naturally* $\mathbb{Z}$*-graded; physical insertions correspond to* $\mathsf{H}^0(\mathcal{B}_\Gamma)$ *modulo band-limited EOMs.*

Lemma 7.215 (Doublet lemma (band-limited))**.** *If a pair (u, v) forms a $\mathcal{B}_\Gamma$-doublet, $\mathcal{B}_\Gamma u = v$, $\mathcal{B}_\Gamma v = 0$, then (u, v) does not contribute to the cohomology: any $\mathcal{B}_\Gamma$-closed functional depending on (u, v) is $\mathcal{B}_\Gamma$-exact up to functionals independent of (u, v). In particular, $(\bar{c}, b)$ and the source pairs form doublets, so they drop out of* $\mathsf{H}^0(\mathcal{B}_\Gamma)$.

Proof. Standard homotopy-contraction argument (see [117]) applies verbatim since the band kernels commute with $\mathcal{B}_\Gamma$. □

Proposition 7.216 (Quantum Action Principle and anomaly consistency)**.** *If radiative corrections generate a breaking $\mathcal{S}(\Gamma) = \Delta_{\mathrm{an}}$ (ghost number $+1$), then the Quantum Action Principle implies the* Wess–Zumino consistency condition

$$\mathcal{B}_\Gamma \, \Delta_{\mathrm{an}} = 0,$$

so $\Delta_{\mathrm{an}} \in \mathsf{H}^1(\mathcal{B}_\Gamma)$ (band-limited). If the anomaly filters of Section 7.2.8 hold, then Δ_{an} is $\mathcal{B}_\Gamma$-exact modulo total derivatives and EOMs, and can be removed by local counterterms (cf. Prop. 7.189).

Proof. Differentiate $\mathcal{S}(\Gamma) = \Delta_{\mathrm{an}}$ with $\mathcal{B}_\Gamma$ and use Lemma 7.213 to obtain $\mathcal{B}_\Gamma \Delta_{\mathrm{an}} = 0$. Cohomological triviality follows from the anomaly filters. □

Proposition 7.217 (Stability of the identity and counterterm characterization)**.** *Let $\Gamma \mapsto \Gamma + \hbar^n \Delta\Gamma$ be a finite renormalization at order n that preserves (7.150). Then*

$$\mathcal{B}_\Gamma(\Delta\Gamma) = 0.$$

Hence admissible counterterms are representatives of $\mathsf{H}^0(\mathcal{B}_\Gamma)$ (modulo band-limited EOMs and total derivatives). By Lemma 7.215, they can be chosen G-invariant and independent of $(\bar{c}, b)$ and sources, matching the physical invariant basis.

Proof. Insert $\Gamma + \hbar^n \Delta\Gamma$ into $\mathcal{S}(\Gamma) = 0$ and retain $\mathcal{O}(\hbar^n)$ terms; linearization gives $\mathcal{B}_\Gamma(\Delta\Gamma) = 0$. The cohomological statement follows from the previous lemmas. □

Proposition 7.218 (Nielsen identity in linearized form)**.** *If the gauge parameter α_{gf} enters only via the gauge fermion $\Psi(\alpha_{\mathrm{gf}})$, then*

$$\partial_{\alpha_{\mathrm{gf}}}\Gamma = -\,\mathcal{B}_\Gamma\big(\partial_{\alpha_{\mathrm{gf}}}\Psi\big)\,,$$

i.e. the gauge–parameter derivative is $\mathcal{B}_\Gamma$–exact. Thus any BRST–closed, ghost–number–zero physical functional is gauge–parameter independent on the band (cf. (7.340)).

Proof. Differentiate the band-limited path integral with respect to α_{gf}; only $\zeta\Psi$ varies. Using the Zinn–Justin antibracket equivalence $\mathcal{B}_\Gamma X = (\Gamma, X)$ (Prop. 7.212) with $X = \partial_{\alpha_{\mathrm{gf}}}\Psi$ yields the result. □

Remark 7.219 (Physical matrix elements on $\mathcal{H}_{\mathrm{phys}}$)**.** Because $[\hat{Q}_{\mathrm{BRST}}, P_x(\lambda_{\mathrm{UV}})] = 0$ and $\zeta(K_{\mathrm{IR}}^{(\ell)} * \cdot) = 0$ (eq. (7.145)), the band-limited BRST charge descends to $\mathcal{H}_{\mathrm{phys}}$ and $\mathcal{B}_\Gamma$-exact insertions decouple between physical projectors (quartet/positivity). Hence physical matrix elements and ancilla readouts computed in Section 7.4 are insensitive to $\mathcal{B}_\Gamma$-exact deformations, providing the exact symmetry-point baseline for the probe.

7.5.3 Gauge-fixing, Antighost, and b-equations

For a general gauge fermion Ψ the ghost/antighost sector obeys exact equations of motion (Quantum Action Principle) [117]:

$$\frac{\delta\Gamma}{\delta b} = \frac{\delta}{\delta b}\big(\zeta\Psi\big) = \mathcal{F}_{\mathrm{gf}}(\Phi, \Theta; \alpha_{\mathrm{gf}}), \qquad \frac{\delta\Gamma}{\delta \bar{c}} = \frac{\delta}{\delta \bar{c}}\big(\zeta\Psi\big) - \partial_\mu \frac{\delta\Gamma}{\delta K_\mu^{(A)}} \ \text{(if gauge vectors are present)}, \tag{7.152}$$

and the *ghost equation* (global ghost-number Ward identity) constrains the dependence on c and its source K_c.[5] These identities are exact at the G-invariant vacuum and commute with the band/IR knobs by (7.145). Together with Section 7.5.2, they imply the decoupling of ζ-exact terms from $\mathcal{H}_{\mathrm{phys}}$ matrix elements.

Definition 7.220 (Quantum Action Principle (band version))**.** Let Z, W, and Γ be the band-limited generating functionals defined in (7.144), with UV/IR tagging that commutes with BRST and with functional differentiation (cf. (7.145)). For any local deformation $\delta_\xi S$ of the action (including changes of couplings, field redefinitions, or a gauge-fermion variation $\delta_\xi\Psi$ entering as $s\Psi$), the variation of the 1PI functional is the insertion of a *local* composite operator:

$$\partial_\xi\Gamma \;=\; \int d^d x\; \mathcal{N}_\xi(x),$$

where $\mathcal{N}_\xi(x)$ is a local polynomial (power-counting bounded) in fields/sources and their derivatives. If the deformation is BRST-exact, $\delta_\xi S = s\,\Xi$, then

$$\partial_\xi\Gamma \;=\; -\,\mathcal{B}_\Gamma(\Xi)\,,$$

[5] We keep the gauge representation abstract; when explicit gauge vectors A_μ are present, $K_\mu^{(A)}$ is the source for $\zeta A_\mu = D_\mu c$ and (7.152) reduces to the standard pair (Landau gauge: $\partial_\mu A^\mu = 0$ implies a linear b-equation).

i.e. the breaking is $\mathcal{B}_\Gamma$-exact. These statements hold on the band because the tagging is linear, local/pseudolocal, and BRST-closed (no anomaly on the band). Standard QAP references include [82, 91].

Proposition 7.221 (QAP form of the b- and antighost equations on the band)**.** *Let Γ be the band-limited 1PI functional of (7.144) with BRST-exact gauge fixing $s\Psi$ (we write $\zeta\Psi$ in components). Then the functional derivatives with respect to the Nakanishi–Lautrup field b^a and the antighost $\bar{c}^a$ satisfy the (band) equations*

$$\frac{\delta\Gamma}{\delta b^a(x)} \;=\; \mathcal{F}^a_{\mathrm{gf}}\big(x;\lambda_{\mathrm{UV}},\ell\big), \qquad \frac{\delta\Gamma}{\delta \bar{c}^a(x)} \;+\; \mathcal{R}^{\,a}\big(x;\lambda_{\mathrm{UV}},\ell\big) \;=\; 0, \tag{7.153}$$

where $\mathcal{F}^a_{\mathrm{gf}}$ and $\mathcal{R}^{\,a}$ are local *functionals fixed by the gauge fermion Ψ (hence by the chosen gauge condition) and by the band knobs $(\lambda_{\mathrm{UV}},\ell)$. In linear covariant gauges, $\mathcal{F}^a_{\mathrm{gf}}$ reduces to the gauge condition (e.g. $\partial_\mu A^{a\mu} - \alpha_{\mathrm{gf}}\, b^a$ up to conventions), while $\mathcal{R}^{\,a}$ is the standard source term built from the 1PI source that couples to sA^a_μ (the Zinn–Justin source $K^{\mu(A)a}$).*

Proof idea. Differentiate $\zeta\Psi$ algebraically with respect to b and $\bar{c}$ and use that band operations commute with functional differentiation by (7.145). If A_μ is present, the $\bar{c}$-equation acquires the standard source term proportional to the derivative of Γ w.r.t. $K^{(A)}_\mu$ (the source for ζA_μ), reproducing (7.152).

Proof. (1) Starting point and band commutativity. Work with the band-limited path integral $Z[J,\text{sources}]$ whose action is $S_{\mathrm{band}} = S_0^{\mathrm{band}} + s\Psi^{\mathrm{band}}$; UV/IR tagging acts as in (7.145) and commutes with BRST and with functional derivatives. Introduce sources $J_b, J_{\bar{c}}$ for $b, \bar{c}$ and the Zinn–Justin sources K that couple to BRST variations $s\varphi$. Legendre-transform to Γ as in (7.144).

(2) b-equation (algebraic EOM). The gauge fermion Ψ depends on b at most algebraically (no derivatives, no kinetic term for b). Hence the functional integral over b is Gaussian/algebraic, and a shift of the b-integration variable yields the exact identity in the connected functional W:

$$\frac{\delta W}{\delta J^a_b(x)} \;=\; \Big\langle\, \mathcal{F}^a_{\mathrm{gf}}(x) \,\Big\rangle,$$

where $\mathcal{F}^a_{\mathrm{gf}}$ is the gauge-fixing function fixed by Ψ (e.g. the gauge condition). Legendre transforming gives $\delta\Gamma/\delta b^a(x) = \mathcal{F}^a_{\mathrm{gf}}(x)$, now with all entries *band-limited*. No quantum breaking can appear here: by QAP Definition 7.220 any breaking would be a local polynomial with the ghost number and dimensions of b; power counting and ghost number force the classical form, so $(7.153)_1$ holds exactly [82, 91].

(3) Antighost equation. Consider the variation of the integrand under the *inhomogeneous* shift $\bar{c}^a \mapsto \bar{c}^a + \varepsilon^a$ (at fixed sources). In $s\Psi$ the $\bar{c}$ only appears linearly through the gauge-fixing term $\bar{c}\,\mathcal{G}(A,c,\ldots)$ determined by Ψ and through the Zinn–Justin sector coupling the source $K^{\mu(A)}$ to sA_μ. By a standard change of variables in the path integral and integration by parts one obtains, in the connected generating functional,

$$\frac{\delta W}{\delta J^a_{\bar{c}}(x)} \;+\; \Big\langle\, \mathcal{R}^{\,a}(x) \,\Big\rangle \;=\; 0,$$

where $\mathcal{R}^a$ is the local composite induced by the sA sector (in linear covariant gauges, $\mathcal{R}^a = \partial_\mu K^{\mu(A)a}$ up to conventions). Legendre transforming gives the 1PI relation

$$\frac{\delta\Gamma}{\delta\bar{c}^a(x)} \;+\; \mathcal{R}^a(x) \;=\; 0.$$

Again, by the QAP Definition 7.220 any radiative breaking would be a *local* insertion with ghost number -1 and canonical dimension bounded by the gauge-fixing sector; the only such insertion is the classical $\mathcal{R}^a$ which is already present, so the identity is stable under renormalisation on the band [82, 91].

(4) Band locality and stability. Throughout, UV/IR tagging is linear and BRST-closed, hence it commutes with the changes of variables used above and preserves locality (the kernels are pseudodifferential of order ≤ 0 on the band). Therefore the QAP applies verbatim to the band-limited Γ, and (7.153) holds exactly with *local* $\mathcal{F}^a_{\mathrm{gf}}$ and $\mathcal{R}^a$ determined by Ψ and $(\lambda_{\mathrm{UV}}, \ell)$.

This proves the proposition. □

Proposition 7.222 (Ghost equation with linear breaking (band-limited))**.** *There exists a local Ward operator $\mathcal{G}^a(x)$ of ghost number -1 (the* ghost Ward operator*) such that:*

$$\mathcal{G}^a(x)\,\Gamma = \Delta^a_c(x). \tag{7.154}$$

Here the breaking Δ^a_c *is linear in the external Zinn–Justin sources (e.g. $K^{\mu(A)a}, K^{(\Phi)}_A, K^{(\Theta)}_A, \ldots$) and therefore receives no loop corrections beyond tree level (stability of the breaking by the QAP). In Landau gauge the breaking reduces to a term purely linear in $K^{\mu(A)a}$ and is independent of dynamical fields. All contractions and kernels are understood "on the band," i.e. with $P_x(\lambda_{\mathrm{UV}})$ and $K^{(\ell)}_{\mathrm{IR}}$ inserted as in (7.144), and tagging commutes with BRST by (7.145).*

Proof idea. Perform the change of variables $c \mapsto c + \epsilon$ in the band-limited path integral (constant shift in ghost number $+1$). The Jacobian is unity and the only variation originates from source terms coupled to ζc; passing to Γ yields a linear functional in the sources, i.e. Δ_c. Band operations commute with the shift.

Proof. **1) Setup on the band and sources.** Let Z, W, and Γ be the band-limited generating functionals defined in (7.144). The gauge-fixed action is $S_{\mathrm{band}} = S_{0,\mathrm{band}} + s\Psi_{\mathrm{band}}$, with BRST variations $s\bar{c}^a = b^a$, $sb^a = 0$, and $s\varphi^A = R^A{}_a(\varphi)\,c^a$ for generic matter/gauge multiplets φ^A. Introduce Zinn–Justin sources K that couple linearly to the BRST variations $s\varphi$, so the source part reads schematically $\int d^dx\, K_A\, s\varphi^A$ (and analogously for gauge fields). By (7.145), $P_x(\lambda_{\mathrm{UV}})$ and $K^{(\ell)}_{\mathrm{IR}}$ commute with s and with functional differentiation, so the usual algebra is unaffected on the band.

2) Constant ghost shift in the path integral. Consider the change of variables (translation) in the band-limited path integral

$$c^a(x) \;\mapsto\; c^a(x) + \epsilon^a(x), \qquad \epsilon^a(x) \text{ arbitrary test function (ghost number } +1).$$

The functional measure is translation invariant, hence its Jacobian is 1. The only variation of the integrand comes from the explicit c-dependence in $s\Psi_{\text{band}}$ *via* $s\varphi^A$ and from the Zinn–Justin source term $\int K_A\, s\varphi^A$. At the level of the connected generator W, this yields the identity

$$\int d^dx\, \epsilon^a(x) \left[\frac{\delta W}{\delta J_c^a(x)} + \Big\langle \frac{\delta(s\Psi_{\text{band}})}{\delta c^a(x)} \Big\rangle + \int d^dy\, K_A(y) \, \Big\langle \tfrac{\delta(s\varphi^A(y))}{\delta c^a(x)} \Big\rangle \right] = 0.$$

Legendre-transforming to the 1PI functional Γ and using the standard relations between $\delta W/\delta J$ and classical fields gives a local operator identity acting on Γ:

$$\frac{\delta\Gamma}{\delta c^a(x)} \;+\; \mathcal{U}^a(x;\Phi,\Theta,\ldots) \;+\; \int d^dy\, K_A(y)\, \frac{\delta(s\varphi^A(y))}{\delta c^a(x)} \;=\; 0, \tag{7.155}$$

where $\mathcal{U}^a$ collects the contribution from $s\Psi_{\text{band}}$ expressed in 1PI variables (see Step 4). Equation (7.155) holds *on the band* since tagging commutes with all steps.

3) Definition of the ghost Ward operator and the breaking. Define the (band) ghost Ward operator $\mathcal{G}^a(x)$ acting on functionals F by

$$\mathcal{G}^a(x)\, F \;:=\; \frac{\delta F}{\delta c^a(x)} \;+\; \int d^dy\, \frac{\delta F}{\delta K_A(y)}\, \frac{\delta\big(s\varphi^A(y)\big)}{\delta c^a(x)} \;+\; \bar{c}^b(x)\, \Xi^{ab}\Big[x; \frac{\delta F}{\delta b}\Big], \tag{7.156}$$

where the last term is the algebraic piece enforced by $s\bar{c} = b$ (its concrete form is fixed by the gauge fermion and is linear in the functional derivative w.r.t. b; see Step 4). By construction, $\mathcal{G}^a$ has ghost number -1 and is local. Evaluating (7.156) on $F = \Gamma$ and comparing with (7.155) we obtain

$$\mathcal{G}^a(x)\,\Gamma \;=\; \Delta_c^a(x), \qquad \Delta_c^a(x) \text{ local and linear in the Zinn–Justin sources.}$$

Explicitly,

$$\Delta_c^a(x) \;=\; -\mathcal{U}^a(x;\Phi,\Theta,\ldots)\Big|_{\text{linear in sources}} \;=\; \sum_{\mathsf{A}} \mathcal{C}^a{}_{\mathsf{A}}(\partial, \text{rep.})\, K^{\mathsf{A}}(x), \tag{7.157}$$

with $\mathcal{C}$ a differential operator fixed by the BRST variations $s\varphi$ (representation matrices, covariant derivatives, etc.). Thus Δ_c is *linear* in the external sources and contains no dynamical-field dependence in Landau gauge (Step 4).

4) Landau gauge and explicit form of the linear breaking. In linear covariant gauges one may choose a gauge fermion of the form $\Psi = \int d^dx\, \bar{c}^a\, \mathcal{G}^a(A,\Phi,\Theta,\ldots) - \frac{\alpha_{\text{gf}}}{2}\, \bar{c}^a b^a$, so that $s\Psi$ is linear in b and $\bar{c}$. Using $s\bar{c}^a = b^a$ and $sA_\mu^a = D_\mu c^a$, the $\bar{c}$-shift–induced piece in (7.155) precisely yields the algebraic $\bar{c}\,\delta/\delta b$ contribution indicated in (7.156), and the remaining terms can be arranged so that the breaking depends only on the Zinn–Justin sources. In particular, for the Landau gauge $\alpha_{\text{gf}} = 0$ one finds

$$\Delta_c^a(x) \;=\; \partial_\mu K^{\mu(A)a}(x) \quad \text{(up to representation/gauge-coupling conventions)},$$

independent of dynamical fields. This is the standard linear breaking of the ghost equation (no field-dependent terms) adapted to the band setting [52, 91].

5) Locality and stability of the breaking (QAP). By the band version of the Quantum Action Principle (QAP; Definition 7.220), the variation induced by the local ghost translation is represented in Γ by insertion of a *local* composite. Ghost number and power counting restrict the possible local insertions on the r.h.s. to be linear in the external sources (they carry the appropriate ghost number). Hence Δ_c is *purely linear* and receives no loop-generated non-linear corrections: its form is tree-level exact and stable under renormalisation [82, 91]. Because $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ commute with BRST and functional differentiation (and are of order ≤ 0), the argument lifts verbatim to the band-limited functionals.

Conclusion. With $\mathcal{G}^a(x)$ defined by (7.156), the identity $\mathcal{G}^a(x)\Gamma = \Delta_c^a(x)$ holds on the band, the breaking Δ_c is local and linear in the Zinn–Justin sources, and in Landau gauge it is independent of dynamical fields. Stability follows from the QAP, completing the proof. □

Definition 7.223 (Jet bundles, total derivatives, and local functionals (Chapter 7 conventions))**.** Let the basic fields be collectively $\varphi^A \in \{\Phi^A, \Theta^A, c^a, \bar{c}^a, b^a, \dots\}$ over spacetime M. For a multi–index $\alpha = (\mu_1 \cdots \mu_k)$, write $\partial_\alpha := \partial_{\mu_1} \cdots \partial_{\mu_k}$ and denote jet coordinates by $\varphi^A_{(\alpha)} := \partial_\alpha \varphi^A$ at a common point $x \in M$. The r–jet bundle J^r has local coordinates $(x^\mu, \varphi^A_{(\alpha)})_{|\alpha|\leq r}$; the infinite jet J^∞ collects all orders. The algebra of local functions $\mathcal{F}_{\mathrm{loc}}$ consists of polynomials/smooth functions of finitely many jet coordinates.
(i) Total derivative. The horizontal (total) derivative acts on jet coordinates by

$$D_\mu \;:=\; \partial_\mu \;+\; \sum_{A,\alpha} \varphi^A_{(\alpha\mu)}\, \frac{\partial}{\partial \varphi^A_{(\alpha)}}\,, \qquad D_\alpha := D_{\mu_1} \cdots D_{\mu_k} \;\text{ for }\; \alpha = (\mu_1 \cdots \mu_k). \tag{7.158}$$

(ii) Euler–Lagrange operator. For a local density $f \in \mathcal{F}_{\mathrm{loc}}$,

$$\mathbf{E}_A(f) \;:=\; \sum_{|\alpha|\geq 0} (-D)_\alpha \left(\frac{\partial f}{\partial \varphi^A_{(\alpha)}} \right), \qquad \delta \int f\, d^dx = \int \delta\varphi^A\, \mathbf{E}_A(f)\, d^dx \;+\; \text{boundary}. \tag{7.159}$$

(iii) Band/IR compatibility. Replace fields by jointly tagged ones from Definition 7.28: $\varphi^A \mapsto \varphi^A_{\lambda_{\mathrm{UV}},\ell}$. Under the standing assumption that $[\,\nabla_\mu, P_x(\lambda_{\mathrm{UV}})\,] = 0$ and $K_{\mathrm{IR}}^{(\ell)}$ is a scalar kernel, tagging commutes with differentiation (cf. item (iv) of Definition 7.28):

$$D_\mu\Big(\varphi^A_{\lambda_{\mathrm{UV}},\ell}\Big) = \Big(D_\mu \varphi^A\Big)_{\lambda_{\mathrm{UV}},\ell}, \qquad D_\alpha\Big(\varphi^A_{\lambda_{\mathrm{UV}},\ell}\Big) = \Big(D_\alpha \varphi^A\Big)_{\lambda_{\mathrm{UV}},\ell}. \tag{7.160}$$

Local band–limited functionals are integrals of densities built from finitely many tagged jets.
(iv) BRST/Slavnov prolongation on jets. Writing the BRST variation as $s\,\varphi^A = R^A(\varphi, \partial\varphi, \dots)$, its prolongation acts by

$$s\,\varphi^A_{(\alpha)} = D_\alpha\Big(R^A(\varphi, \partial\varphi, \dots)\Big). \tag{7.161}$$

Here $\mathcal{B}_\Gamma$ acts derivationally on $\mathcal{F}_{\mathrm{loc}}$ and commutes with D_μ on the band.
(v) Doublet counting and contracting homotopy (for $(\bar{c}, b)$). Let N_{dbl} count jets of the BRST doublet $(\bar{c}^a, b^a)$:

$$N_{\mathrm{dbl}} := \sum_{a,\alpha} \left(\bar{c}^a_{(\alpha)} \frac{\partial}{\partial \bar{c}^a_{(\alpha)}} + b^a_{(\alpha)} \frac{\partial}{\partial b^a_{(\alpha)}} \right).$$

Define the contracting homotopy κ on jet generators by

$$\kappa\big(b^a_{(\alpha)}\big) = \bar{c}^a_{(\alpha)}, \qquad \kappa\big(\bar{c}^a_{(\alpha)}\big) = 0, \qquad [\kappa, D_\mu] = 0, \tag{7.162}$$

and extend κ to $\mathcal{F}_{\mathrm{loc}}$ by graded Leibniz. With $s\,\bar{c}^a = b^a$, $s\,b^a = 0$, one has the homotopy identity

$$\{\, s, \kappa \,\} = N_{\mathrm{dbl}}\,. \tag{7.163}$$

Hence, if a local functional X is s–closed (or $\mathcal{B}_\Gamma$–closed on the band) and $N_{\mathrm{dbl}}X = n > 0$, then $X = \frac{1}{n}\, s(\kappa X)$ up to total derivatives and equations of motion; this is the doublet decoupling used in Lemma 7.224.

Lemma 7.224 (Doublet decoupling)**.** *With the ghost-number assignments of Section 7.5.2, $(\bar{c}, b)$ form a BRST doublet. Consequently, any $\mathcal{B}_\Gamma$-closed local functional that depends on $(\bar{c}, b)$ is $\mathcal{B}_\Gamma$-exact up to equations of motion and total derivatives. Hence $(\bar{c}, b)$ do not contribute to $\mathsf{H}^0(\mathcal{B}_\Gamma)$ and drop out of physical matrix elements on $\mathcal{H}_{\mathrm{phys}}$.*

Full proof. **(1) Linearised Slavnov operator and the doublet.** Let $\mathcal{B}_\Gamma$ be the linearised Slavnov operator around the band-limited 1PI functional Γ (see (7.144)). By Stage I anomaly filters, $\mathcal{B}_\Gamma^2 = 0$ on the band. Off shell (with Nakanishi–Lautrup fields present) one has

$$\mathcal{B}_\Gamma\,\bar{c}^a = b^a, \qquad \mathcal{B}_\Gamma\,b^a = 0, \qquad \mathrm{gh}(\bar{c}^a) = -1,\ \mathrm{gh}(b^a) = 0$$

(Section 7.5.2). Band/IR maps commute with BRST and carry ghost number 0 (eq. (7.145)), so these relations hold "on the band".

(2) Jet framework and operators N_{dbl}, κ. Work in the algebra of local functionals built from finitely many jets of the fields (Chapter 7 jets; Definition 7.223). Let D_μ be the total derivatives (7.158), and let the BRST action be prolonged to jets as in (7.161). Define the doublet-counting operator and the contracting homotopy on jets by

$$N_{\mathrm{dbl}} := \sum_{a,\alpha} \left(\bar{c}^a_{(\alpha)} \frac{\partial}{\partial \bar{c}^a_{(\alpha)}} + b^a_{(\alpha)} \frac{\partial}{\partial b^a_{(\alpha)}} \right), \qquad \kappa\big(b^a_{(\alpha)}\big) = \bar{c}^a_{(\alpha)}, \quad \kappa\big(\bar{c}^a_{(\alpha)}\big) = 0, \quad [\kappa, D_\mu] = 0,$$

and extend κ by the graded Leibniz rule; cf. (7.162). By construction,

$$\{\, \mathcal{B}_\Gamma, \kappa \,\} = N_{\mathrm{dbl}} \quad \text{on jet generators, prolonged via (7.161).} \tag{7.164}$$

(3) Cohomological form of (7.164). When applied to any local functional F, the identity (7.164) produces boundary terms (from integrations by parts with D_μ) and, if needed, Euler–Lagrange pieces generated by the jet-to-configuration reduction ((7.159)). Therefore

$$\{\mathcal{B}_\Gamma, \kappa\}\, F = \ N_{\mathrm{dbl}}\, F \ + \ (\text{total derivatives}) \ + \ (\text{terms} \propto \text{EOM}). \tag{7.165}$$

Since our cohomology is taken modulo total derivatives and equations of motion, these extras are immaterial. Band/IR tagging commutes with derivatives (Definition 7.28, item (iv), eq. (7.160)), hence (7.165) is unaffected "on the band".

(4) Decomposition by N_{dbl}-degree. Let F be a $\mathcal{B}_\Gamma$-closed local functional of ghost number 0: $\mathcal{B}_\Gamma F = 0$. Decompose $F = \sum_{n\geq 0} F^{(n)}$ with $N_{\mathrm{dbl}} F^{(n)} = n\, F^{(n)}$. Projecting (7.165) onto the n-eigenspace ($n > 0$) and using $\mathcal{B}_\Gamma F^{(n)} = 0$ gives

$$n\, F^{(n)} = \{\mathcal{B}_\Gamma, \kappa\} F^{(n)} = \mathcal{B}_\Gamma(\kappa F^{(n)}) \;+\; (\mathrm{mod\ EOM},\, d).$$

Hence, for every $n \geq 1$,

$$F^{(n)} = \mathcal{B}_\Gamma\!\left(\tfrac{1}{n}\,\kappa F^{(n)}\right) \quad \mathrm{mod\ (EOM},\, d).$$

Iterating over n eliminates all dependence on $(\bar{c}, b)$ up to $\mathcal{B}_\Gamma$-exact terms.

(5) Consequences. Only $F^{(0)}$ (independent of $(\bar{c}, b)$) can represent a nontrivial class, so $(\bar{c}, b)$ do not contribute to $\mathsf{H}^0(\mathcal{B}_\Gamma)$. Since insertions that are $\mathcal{B}_\Gamma$-exact vanish in physical matrix elements on $\mathcal{H}_{\mathrm{phys}}$ (Nielsen/BRST cohomological invariance), the doublet drops out of observables.

Band/IR remark. All steps are preserved under joint IR/UV tagging (Definition 7.28): $P_x(\lambda_{\mathrm{UV}})$ and $K^{(\ell)}_{\mathrm{IR}}$ commute with the G-action and with D_μ, and are BRST-closed (item (ii) of Definition 7.28), so the homological construction carries over verbatim on the band. □

Definition 7.225 (Linear covariant gauges and Landau limit)**.** With joint IR/UV tagging as in Definition 7.28, fix BRST-exact gauge fixing by the gauge fermion

$$\Psi \;=\; \int d^d x\, K^{(\ell)}_{\mathrm{IR}}(x)\, \bar{c}^a(x) \Big(F^a(\Phi, \Theta)(x) - \tfrac{\alpha_{\mathrm{gf}}}{2}\, b^a(x) \Big),$$

where F^a is linear in the gauge potential (e.g. $F^a = \partial_\mu A^{\mu a}$, implicitly projected by $P_x(\lambda_{\mathrm{UV}})$). The one-parameter family $\alpha_{\mathrm{gf}} \geq 0$ defines the linear covariant gauges; eliminating b by its algebraic equation gives $b^a = \alpha_{\mathrm{gf}}^{-1} F^a$ for $\alpha_{\mathrm{gf}} > 0$. The *Landau limit* is $\alpha_{\mathrm{gf}} \to 0$, in which the b-equation imposes the linear constraint $F^a = 0$ on the band. This convention is stable under the QAP in the algebraic renormalisation framework and preserves the BRST cohomology (see [117, 82, 91]).

Proposition 7.226 (Linear covariant gauge as an illustrative instance)**.** *Suppose* $\Psi = \int d^d x\, K^{(\ell)}_{\mathrm{IR}}(x)\, \bar{c}^a \Big(F^a(\Phi, \Theta) - \tfrac{\alpha_{\mathrm{gf}}}{2}\, b^a \Big)$ *with F linear in the gauge field (e.g. $F^a = \partial_\mu A^{\mu a}$, understood band-limited by $P_x(\lambda_{\mathrm{UV}})$). Then*

$$\frac{\delta \Gamma}{\delta b^a} = F^a - \alpha_{\mathrm{gf}}\, b^a, \qquad \frac{\delta \Gamma}{\delta \bar{c}^a} = \frac{\delta F^a}{\delta A^b_\mu} \frac{\delta \Gamma}{\delta K^{(A)\mu b}} - \partial_\mu \frac{\delta \Gamma}{\delta K^{(A)\mu a}},$$

and the ghost equation reads $\mathcal{G}^a(x)\,\Gamma = \partial_\mu K^{(A)\mu a}(x)$ (all identities understood on the band). In the Landau limit $\alpha_{\mathrm{gf}} \to 0$, the b-equation imposes $F^a = 0$ linearly, and $\Delta_c = \partial_\mu K^{(A)\mu a}$ is a linear, tree-level exact breaking.

Proof idea. Insert the gauge fermion of Definition 7.225 into Proposition 7.221 to get $\delta\Gamma/\delta b^a = F^a - \alpha_{\mathrm{gf}} b^a$ and $\delta\Gamma/\delta \bar{c}^a = \frac{\delta F^a}{\delta A^b_\mu}\, \delta\Gamma/\delta K^{(A)\mu b} - \partial_\mu\, \delta\Gamma/\delta K^{(A)\mu a}$. The ghost equation then follows directly from Proposition 7.222: $\mathcal{G}^a(x)\Gamma = \partial_\mu K^{(A)\mu a}(x)$. Taking the Landau limit $\alpha_{\mathrm{gf}} \to 0$ gives $F^a = 0$ and a purely linear, tree-level exact breaking.

Full proof. **1) Gauge fixing and BRST variations.** With the gauge fermion Ψ as stated and BRST rules $s\,\bar{c}^a = b^a$, $s\,b^a = 0$, $s\,A^a_\mu = (D_\mu c)^a$ (and s acting as a derivation on other fields), the gauge-fixing+ghost density is

$$\mathcal{L}_{\mathrm{gf+gh}} = s\left[K^{(\ell)}_{\mathrm{IR}}\,\bar{c}^a\Big(F^a - \tfrac{\alpha_{\mathrm{gf}}}{2}b^a\Big)\right] = K^{(\ell)}_{\mathrm{IR}}\left[b^a\Big(F^a - \tfrac{\alpha_{\mathrm{gf}}}{2}b^a\Big) - \bar{c}^a\,sF^a\right].$$

Because F is linear in A, $sF^a = \frac{\delta F^a}{\delta A^b_\mu}\,(D_\mu c)^b$. Joint IR/UV tagging commutes with the BRST action and with derivatives (cf. Definition 7.28, items (ii)–(iv)), so these equalities hold on the band.

2) b-equation (algebraic). In the path integral, the field b^a appears only algebraically. The classical identity

$$\frac{\delta S}{\delta b^a} = K^{(\ell)}_{\mathrm{IR}}\Big(F^a - \alpha_{\mathrm{gf}}\,b^a\Big)$$

is therefore stable under renormalisation by the Quantum Action Principle (QAP): linear, non-derivative algebraic equations are not deformed by loops (they can at most receive local redefinitions that are ruled out by ghost number and power counting). Passing to the 1PI functional and dropping the common $K^{(\ell)}_{\mathrm{IR}}$ factor that commutes with functional differentiation (on the band) yields

$$\frac{\delta\Gamma}{\delta b^a} = F^a - \alpha_{\mathrm{gf}}\,b^a,$$

exactly. See the algebraic-renormalisation treatment of auxiliary b-fields [117]; compare also the standard linear covariant gauges [82, 91].

3) Antighost equation. Varying the action w.r.t. $\bar{c}^a$ gives

$$\frac{\delta S}{\delta \bar{c}^a} = -\,K^{(\ell)}_{\mathrm{IR}}\,\frac{\delta F^a}{\delta A^b_\mu}\,(D_\mu c)^b.$$

Introduce Zinn–Justin sources $K^{(A)\mu b}$ coupled to sA^b_μ in the usual way (band-limited as in (7.144)); then, by definition of the Legendre transform,

$$\frac{\delta\Gamma}{\delta K^{(A)\mu b}} = \big\langle (D_\mu c)^b \big\rangle.$$

Using this and integrating by parts in the covariant derivative (keeping the band kernels, which commute with ∂_μ), one finds

$$\frac{\delta\Gamma}{\delta\bar{c}^a} = \frac{\delta F^a}{\delta A^b_\mu}\,\frac{\delta\Gamma}{\delta K^{(A)\mu b}} - \partial_\mu \frac{\delta\Gamma}{\delta K^{(A)\mu a}},$$

which is the claimed antighost equation on the band (QAP ensures stability; cf. [117]).

4) Ghost equation and its linear breaking. Perform the rigid shift $c^a \mapsto c^a + \epsilon^a$ in the (band-limited) path integral. The Jacobian is 1, and only terms that depend explicitly on c contribute. In linear covariant gauges, the dependence appears via the BRST variations

coupled to Zinn–Justin sources. Standard manipulations (or QAP) yield a local "ghost Ward operator" $\mathcal{G}^a(x)$ acting on Γ with a breaking that is *linear* in sources:

$$\mathcal{G}^a(x)\,\Gamma = \partial_\mu K^{(A)\mu a}(x).$$

Linearity implies the breaking receives no loop corrections beyond tree level (stability of the breaking) [117]; see also the textbook discussion of the ghost equation in covariant gauges [82, 91]. Band kernels commute with derivatives and carry ghost number 0 (cf. Definition 7.28), so the same identity holds on the band.

5) Landau limit. For $\alpha_{\rm gf} \to 0$ the b-equation becomes the linear constraint $F^a = 0$, implementing the (band-limited) gauge condition. The ghost equation's breaking remains $\Delta_c = \partial_\mu K^{(A)\mu a}$, which—being linear and source-only—is tree-level exact by the QAP [117].

Conclusion. The three identities follow from BRST-exact gauge fixing, the QAP, and the commutation properties of the joint IR/UV tagging. Their form is preserved on the band, and Landau gauge reduces the b-equation to the linear gauge constraint. □

Corollary 7.227 (Compatibility with Slavnov–Taylor and Nielsen)**.** *Equations* (7.152) *and Proposition 7.222 are compatible with the band Slavnov identity* (7.150) *and the band Nielsen identity* (7.148)*: applying* $\mathcal{B}_\Gamma$ *to* (7.152) *reproduces identities among sources that follow from* (7.150)*, while* $\partial_{\alpha_{\rm gf}}$*–variations are* $\mathcal{B}_\Gamma$*–exact and thus decouple from BRST–closed physical functionals.*

Full proof. **(A) Compatibility with the band Slavnov–Taylor identity.** Recall the band ST identity

$$\mathcal{S}(\Gamma) = \int d^dx \left(\frac{\delta\Gamma}{\delta K_A}\frac{\delta\Gamma}{\delta\varphi^A} + b^a \frac{\delta\Gamma}{\delta\bar{c}^a} \right) = 0,$$

and the linearised Slavnov operator

$$\mathcal{B}_\Gamma F = \int d^dx \left(\frac{\delta\Gamma}{\delta K_A}\frac{\delta F}{\delta\varphi^A} + \frac{\delta F}{\delta K_A}\frac{\delta\Gamma}{\delta\varphi^A} + b^a \frac{\delta F}{\delta\bar{c}^a} \right), \qquad \mathcal{B}_\Gamma\Gamma = 0,$$

which is nilpotent on the band modulo EOM and total derivatives (anomaly filters; band commutation properties from Definition 7.28, items (ii)–(iv); cf. [117]).

Differentiate $\mathcal{S}(\Gamma) = 0$ w.r.t. $b^a(x)$:

$$\mathcal{B}_\Gamma\left(\frac{\delta\Gamma}{\delta b^a(x)} \right) + \frac{\delta\Gamma}{\delta\bar{c}^a(x)} = 0. \tag{$\star$}$$

Insert the b–equation from (7.152), $\delta\Gamma/\delta b^a = F^a - \alpha_{\rm gf}\, b^a$. Since $\mathcal{B}_\Gamma b^a = 0$, we obtain

$$\frac{\delta\Gamma}{\delta\bar{c}^a(x)} = -\,\mathcal{B}_\Gamma F^a(x).$$

Because F^a depends only on the gauge field, $\mathcal{B}_\Gamma$ acts on it via the $K^{(A)}$–channel; integrating by parts (band kernels commute with ∂_μ), this yields exactly the antighost equation in (7.152),

$$\frac{\delta\Gamma}{\delta\bar{c}^a} = \frac{\delta F^a}{\delta A^b_\mu}\frac{\delta\Gamma}{\delta K^{(A)\mu b}} - \partial_\mu \frac{\delta\Gamma}{\delta K^{(A)\mu a}},$$

thus showing that applying $\mathcal{B}_\Gamma$ to the b–equation reproduces the antighost identity dictated by ST.

Likewise, differentiating $\mathcal{S}(\Gamma) = 0$ w.r.t. $\bar{c}^a(x)$ gives

$$\mathcal{B}_\Gamma\left(\frac{\delta\Gamma}{\delta\bar{c}^a(x)}\right) = 0, \tag{$\star\star$}$$

which is the $\mathcal{B}_\Gamma$–closure of the antighost equation. Eqs. ($\star$)–($\star\star$) are the standard algebraic–renormalisation compatibility relations (now "on the band"); see [117] and textbook expositions [82, 91].

(B) Compatibility with the band Nielsen identity. The band Nielsen identity (eq. (7.148); cf. Proposition 7.236) reads

$$\partial_{\alpha_{\rm gf}}\Gamma = -\,\mathcal{B}_\Gamma\big(\partial_{\alpha_{\rm gf}}\Psi\big) = \int d^dx\,\frac{\delta\Gamma}{\delta\varphi^A(x)}\,\mathcal{N}^A(x;\alpha_{\rm gf}),$$

so $\partial_{\alpha_{\rm gf}}\Gamma$ is $\mathcal{B}_\Gamma$–exact (and EOM–proportional). Apply $\partial_{\alpha_{\rm gf}}$ to (7.152) and use the commutation of band operations with functional differentiation (Definition 7.28, items (ii)–(iv)):

$$\partial_{\alpha_{\rm gf}}\left(\frac{\delta\Gamma}{\delta b^a} - F^a + \alpha_{\rm gf} b^a\right) = \frac{\delta}{\delta b^a}\big(\partial_{\alpha_{\rm gf}}\Gamma\big) - b^a = -\,\frac{\delta}{\delta b^a}\,\mathcal{B}_\Gamma(\partial_{\alpha_{\rm gf}}\Psi) - b^a.$$

Because functional differentiation commutes with $\mathcal{B}_\Gamma$ on the band and $\frac{\delta}{\delta b^a}(\partial_{\alpha_{\rm gf}}\Psi) = -(1/2)\,\bar{c}^a$ for the linear covariant family, the r.h.s. becomes $-\mathcal{B}_\Gamma\big(-(1/2)\bar{c}^a\big) - b^a = 0$ using $\mathcal{B}_\Gamma\bar{c}^a = b^a$. Thus the b–equation is stable under $\alpha_{\rm gf}$–variations in $\mathcal{B}_\Gamma$–exact form. The same reasoning applied to the antighost equation shows its $\partial_{\alpha_{\rm gf}}$–variation is $\mathcal{B}_\Gamma$–exact as well. Consequently, for any BRST–closed ghost–number–zero functional $\mathcal{O}$ (physical observable),

$$\partial_{\alpha_{\rm gf}}\,\mathcal{O} = \big\langle \mathcal{B}_\Gamma(\cdots)\big\rangle \equiv 0 \quad \text{in } \mathcal{H}_{\rm phys},$$

establishing decoupling in the physical sector (Nielsen–type independence; cf. [117]).

Conclusion. Part (A) shows that acting with $\mathcal{B}_\Gamma$ on (7.152) reproduces identities among sources fixed by (7.150); part (B) shows $\partial_{\alpha_{\rm gf}}$–variations are $\mathcal{B}_\Gamma$–exact and hence inert on BRST–closed physical functionals. This proves the corollary. □

Remark 7.228 (Operational consequence)**.** Lem. 7.224 implies that SPAM-like systematics associated with $(\bar{c}, b)$ cannot pollute the certified visibilities on $\mathcal{H}_{\rm phys}$; remaining gauge-parameter dependences are controlled by the Nielsen identity and the anomaly filters, as used throughout Section 7.4.

7.5.4 Global G Ward Identities at the Dymmetric Vacuum

Let G act linearly on the matter multiplets (Φ, Θ)—with *gaiamiti* mean field Φ and *tantrakyma* mean field Θ—via generators T^a:

$$\delta^a\Phi = i\,T^a\Phi, \qquad \delta^a\Theta = i\,T^a\Theta,$$

and take ghosts $(c, \bar{c}, b)$ to be G-singlets unless stated otherwise. At the G-invariant vacuum (Quadrant I), the band-limited generating functionals obey the exact Ward identities

$$\mathcal{W}^a W = 0, \qquad \mathcal{W}^a \Gamma = 0, \tag{7.166}$$

with Ward operators

$$\mathcal{W}^a := \int d^d x \, (J_\Phi \, \delta^a \Phi \; + \; J_\Theta \, \delta^a \Theta \; + \; K_\Phi \, \delta^a(\zeta \Phi) \; + \; K_\Theta \, \delta^a(\zeta \Theta)) \, ,$$

and similarly in the 1PI variables. Here Φ, Θ are the 1PI mean fields conjugate to the sources J_Φ, J_Θ for the band-limited operator fields $\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}}}$ and $\hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}}}$ (cf. the source formalism). Band/IR operations commute with G and BRST by (7.145), hence $\delta^a(\zeta\phi) = \zeta(\delta^a \phi)$ for $\phi \in \{\Phi, \Theta\}$.

Proposition 7.229 (Band-limited global Ward identities for W and Γ)**.** *Assume (i) $S_{\Phi,\Theta}^{(\lambda_{\mathrm{UV}},\ell)}$ is G-invariant at the symmetry point, (ii) the gauge fermion Ψ is G-invariant, and (iii) (7.145). Then the connected functional W and the 1PI functional Γ satisfy (7.166) exactly on the band.*

Proof. Let $\varphi = (\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}}}, \hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}}}, c, \bar{c}, b)$ denote the band-limited operator fields and

$$Z[\mathcal{J}, \mathcal{K}] = \int \mathcal{D}\varphi \; \exp\!\Big\{ i \Big(S_{\Phi,\Theta}^{(\lambda_{\mathrm{UV}},\ell)} + \zeta\Psi + \mathcal{J}\cdot\varphi + \mathcal{K}\cdot\zeta\varphi \Big) \Big\},$$

with $\mathcal{J}\cdot\varphi = \int (J_\Phi \cdot \hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}}} + J_\Theta \cdot \hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}}})$ and $\mathcal{K}\cdot\zeta\varphi = \int (K_\Phi \cdot \zeta\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}}} + K_\Theta \cdot \zeta\hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}}})$, where the dots include contraction with the IR window $K_{\mathrm{IR}}^{(\ell)}$ (cf. the source formalism). By (7.145), the band/IR operations commute with BRST and carry ghost number 0; we also assume they are G-scalars so they commute with the G-action.

Step 1 (Change of variables and measure). Consider the global G variation $\delta^a \varphi = R^a[\varphi]$ with $\delta^a \hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}}} = i\, T^a \hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}}}$, $\delta^a \hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}}} = i\, T^a \hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}}}$, and ghosts $(c, \bar{c}, b)$ taken G-singlets (the generalisation is immediate if they transform trivially in another representation). Perform in the path integral the change of variables $\varphi \mapsto \varphi + \epsilon\, \delta^a \varphi$ with constant Grassmann-even ϵ. Since the transformation is linear and global, the functional measure is invariant (Jacobian $= 1$) and

$$0 = \delta^a Z = \int \mathcal{D}\varphi \; i\, \delta^a \Big(S_{\Phi,\Theta}^{(\lambda_{\mathrm{UV}},\ell)} + \zeta\Psi + \mathcal{J}\cdot\varphi + \mathcal{K}\cdot\zeta\varphi \Big) \, e^{i(\cdots)}.$$

Step 2 (Variation of the action pieces). At the symmetry point, assumption (i) gives $\delta^a S_{\Phi,\Theta}^{(\lambda_{\mathrm{UV}},\ell)} = 0$. Assumption (ii) implies $\delta^a(\zeta\Psi) = \zeta(\delta^a\Psi) = 0$ because Ψ is G-invariant and δ^a commutes with ζ (both band-limited by (7.145)). Thus only the source sector varies:

$$\delta^a(\mathcal{J}\cdot\varphi + \mathcal{K}\cdot\zeta\varphi) = \int d^d x \Big(J_\Phi \, \delta^a \hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}}} + J_\Theta \, \delta^a \hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}}} + K_\Phi \, \delta^a(\zeta \hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}}}) + K_\Theta \, \delta^a(\zeta \hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}}}) \Big).$$

Because band/IR operations commute with G and BRST (assumption (iii) and (7.145)), we have $\delta^a(\zeta \hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}}}) = \zeta(\delta^a \hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}}}) = i\,\zeta(T^a \hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}}})$ and analogously for Θ.

Step 3 (Ward identity for W). Divide $\delta^a Z = 0$ by Z and use $W = -i \ln Z$ to obtain

$$0 = \int d^d x \, (J_\Phi \, \delta^a \Phi + J_\Theta \, \delta^a \Theta + K_\Phi \, \delta^a(\zeta\Phi) + K_\Theta \, \delta^a(\zeta\Theta)) \equiv \mathcal{W}^a W,$$

where we identified the mean fields $\Phi = \delta W/\delta J_\Phi$, $\Theta = \delta W/\delta J_\Theta$ and the BRST insertions $\zeta\Phi = \delta W/\delta K_\Phi$, $\zeta\Theta = \delta W/\delta K_\Theta$ in the standard way. This is precisely the first identity in (7.166).

Step 4 (Legendre transform to Γ). Define the 1PI functional at fixed $\mathcal{K}$ by the usual Legendre transform $\Gamma[\Phi,\Theta;\mathcal{K}] = W[\mathcal{J},\mathcal{K}] - \int(J_\Phi\,\Phi + J_\Theta\,\Theta)$, with $\Phi = \delta W/\delta J_\Phi$, $\Theta = \delta W/\delta J_\Theta$ and inverse relations $J_\Phi = -\delta\Gamma/\delta\Phi$, $J_\Theta = -\delta\Gamma/\delta\Theta$. Act with $\mathcal{W}^a$ on Γ:

$$\mathcal{W}^a\Gamma = \int d^dx\Big(\frac{\delta\Gamma}{\delta\Phi}\,\delta^a\Phi + \frac{\delta\Gamma}{\delta\Theta}\,\delta^a\Theta + \frac{\delta\Gamma}{\delta K_\Phi}\,\delta^a(\zeta\Phi) + \frac{\delta\Gamma}{\delta K_\Theta}\,\delta^a(\zeta\Theta)\Big).$$

Using the Legendre relations $\delta\Gamma/\delta\Phi = -J_\Phi$, $\delta\Gamma/\delta\Theta = -J_\Theta$ and the identities $\delta\Gamma/\delta K_\Phi = -\,\delta W/\delta K_\Phi = -(\zeta\Phi)$, $\delta\Gamma/\delta K_\Theta = -(\zeta\Theta)$ (Legendre transform taken at fixed $\mathcal{K}$), we get

$$\mathcal{W}^a\Gamma = -\int d^dx\,(J_\Phi\,\delta^a\Phi + J_\Theta\,\delta^a\Theta + K_\Phi\,\delta^a(\zeta\Phi) + K_\Theta\,\delta^a(\zeta\Theta)) = -\,\mathcal{W}^aW.$$

Since $\mathcal{W}^aW = 0$ from Step 3, it follows that $\mathcal{W}^a\Gamma = 0$, i.e. the second identity in (7.166).

Step 5 (Band/IR remarks and contact terms). Potential total-derivative (contact) terms from the G-variation of $S^{(\lambda_{\mathrm{UV}},\ell)}_{\Phi,\Theta}$ vanish at the symmetric vacuum and under the L^1 IR window $K^{(\ell)}_{\mathrm{IR}}$; moreover, the G-invariance of Ψ eliminates gauge-fixing contact terms. The commutation relations in (7.145) ensure that the foregoing manipulations hold *on the band*. Therefore the Ward identities are exact in the band-limited theory.

Combining Steps 3–4 proves that both the connected generator W and the 1PI functional Γ obey (7.166) exactly on the band, as claimed. □

Proposition 7.230 (Compatibility with Slavnov and linearised BRST)**.** *Let $\mathcal{S}(\Gamma) = 0$ (band Slavnov identity, Section 7.5.2). Then*

$$\mathcal{W}^a\big(\mathcal{S}(\Gamma)\big) = 0, \qquad [\mathcal{W}^a, \mathcal{B}_\Gamma] = 0,$$

so the linearised Slavnov cohomology $\mathsf{H}^\bullet(\mathcal{B}_\Gamma)$ is G-equivariant. In particular, $\mathsf{H}^0(\mathcal{B}_\Gamma)$ decomposes into G-irreps; physical representatives are G-invariant classes at the symmetry point.

Proof. All statements are understood on the band; by (7.145), band/IR operations commute with G and with BRST, and carry ghost number 0.

1) Ward operator in 1PI variables and its properties. Recall that for any functional $F[\Phi,\Theta;K_\Phi,K_\Theta]$ we have

$$\mathcal{W}^aF = \int d^dx\Big(\frac{\delta F}{\delta\Phi}\,\delta^a\Phi + \frac{\delta F}{\delta\Theta}\,\delta^a\Theta + \frac{\delta F}{\delta K_\Phi}\,\delta^a(\zeta\Phi) + \frac{\delta F}{\delta K_\Theta}\,\delta^a(\zeta\Theta)\Big),$$

with $\delta^a\Phi = iT^a\Phi$, $\delta^a\Theta = iT^a\Theta$ and $\delta^a(\zeta\phi) = \zeta(\delta^a\phi)$. Since the G-action is linear and band-limited, $\mathcal{W}^a$ is a graded derivation (ghost number 0) of the functional algebra:

$$\mathcal{W}^a(FG) = (\mathcal{W}^aF)G + F(\mathcal{W}^aG).$$

2) $\mathcal{W}^a$ is a derivation of the antibracket. Let $(\cdot,\cdot)$ denote the Zinn–Justin antibracket, and use the standard form $(F,G)=\int\Big(\frac{\delta_r F}{\delta\Phi}\frac{\delta_l G}{\delta K_\Phi}-\frac{\delta_r F}{\delta K_\Phi}\frac{\delta_l G}{\delta\Phi}+(\Phi\to\Theta)\Big)$ with band-limited contractions. A direct application of the product rule for $\mathcal{W}^a$ and the linearity of δ^a gives

$$\mathcal{W}^a(F,G)=(\mathcal{W}^a F,G)+(F,\mathcal{W}^a G). \tag{7.167}$$

No extra contact terms arise because (i) the Ward variations are linear in the fields/sources, and (ii) the band/IR kernels commute with G.

3) $\mathcal{W}^a(\mathcal{S}(\Gamma))=0$. With $\mathcal{S}(\Gamma)=\frac{1}{2}(\Gamma,\Gamma)$,

$$\mathcal{W}^a\Big(\mathcal{S}(\Gamma)\Big)=\tfrac{1}{2}\,\mathcal{W}^a(\Gamma,\Gamma)=\tfrac{1}{2}\Big((\mathcal{W}^a\Gamma,\Gamma)+(\Gamma,\mathcal{W}^a\Gamma)\Big)\overset{\mathcal{W}^a\Gamma=0}{=}0,$$

where $\mathcal{W}^a\Gamma=0$ is precisely the Ward identity for Γ (see (7.166) and Prop. 7.229), and we used (7.167).

4) $[\mathcal{W}^a,\mathcal{B}_\Gamma]=0$. Using $\mathcal{B}_\Gamma X=(\Gamma,X)$ and (7.167),

$$[\mathcal{W}^a,\mathcal{B}_\Gamma]X=\mathcal{W}^a(\Gamma,X)-(\Gamma,\mathcal{W}^a X)=(\mathcal{W}^a\Gamma,X)+(\Gamma,\mathcal{W}^a X)-(\Gamma,\mathcal{W}^a X)=(\mathcal{W}^a\Gamma,X)=0.$$

Thus the Ward operator commutes with the linearised Slavnov operator.

5) G-equivariance of cohomology. Because $[\mathcal{W}^a,\mathcal{B}_\Gamma]=0$ and $\mathrm{gh}(\mathcal{W}^a)=0$,

$$\mathcal{B}_\Gamma X=0\ \Rightarrow\ \mathcal{B}_\Gamma(\mathcal{W}^a X)=0,\qquad X=\mathcal{B}_\Gamma Y\ \Rightarrow\ \mathcal{W}^a X=\mathcal{B}_\Gamma(\mathcal{W}^a Y).$$

Hence $\mathcal{W}^a$ maps $\mathcal{B}_\Gamma$-cocycles to cocycles and coboundaries to coboundaries, inducing a well-defined G-action on $\mathsf{H}^\bullet(\mathcal{B}_\Gamma)$. In particular, $\mathsf{H}^0(\mathcal{B}_\Gamma)$ decomposes into G-irreducible representations; the physical representatives at the symmetry point are precisely the G-invariant classes.

This establishes the proposition. □

Proposition 7.231 (Momentum-space constraints and tensor structures)**.** *Write the 1PI two-point functions in momentum space as kernels $\Gamma^{(2)}_{\Phi\Phi}(p)$, $\Gamma^{(2)}_{\Theta\Theta}(p)$, etc. Then:*

(i) G-covariance: $[T^a,\Gamma^{(2)}_{\Phi\Phi}(p)]=0$ *and analogously for other sectors. Hence $\Gamma^{(2)}(p)$ is block-diagonal in G-irreps, with each block proportional to the identity on the irrep space (Schur).*

(ii) No spurionic masses: *symmetry-breaking mass terms are forbidden; mass matrices must commute with all T^a and thus be irrep-wise proportional to the identity. Mixing between distinct irreps is prohibited.*

(iii) Current transversality: *for conserved G-currents J^a_μ, the band-limited polarisation tensor satisfies*

$$p_\mu\,\Pi^{\mu\nu}_{ab}(p)=0,\qquad \Pi^{\mu\nu}_{ab}(p)=\Pi_{ab}(p^2)\,\Pi^{\mu\nu}_{\mathrm{T}}(p),$$

with the standard transverse projector $\Pi^{\mu\nu}_{\mathrm{T}}(p)=\eta^{\mu\nu}-p^\mu p^\nu/p^2$ and no longitudinal piece at the symmetry point.

Proof. Work at the symmetric vacuum, with vanishing mean fields and sources, and assume translation invariance on the band so that 1PI kernels are diagonal in total momentum (the band operations commute with G and translations, cf. eq. (7.145)).

(i) G-covariance of two-point kernels. The global Ward identity (7.166) for Γ reads

$$\int d^dx \left(\frac{\delta\Gamma}{\delta\Phi(x)}\,\delta^a\Phi(x) + \frac{\delta\Gamma}{\delta\Theta(x)}\,\delta^a\Theta(x) \right) = 0, \qquad \delta^a\Phi = i\,T^a\Phi, \quad \delta^a\Theta = i\,T^a\Theta. \tag{7.168}$$

Differentiate (7.168) functionally with respect to $\Phi(y)$ and $\Phi(z)$, then set all mean fields to zero. Using

$$\Gamma^{(2)}_{\Phi\Phi}(y,z) := \left.\frac{\delta^2\Gamma}{\delta\Phi(y)\,\delta\Phi(z)}\right|_{\Phi=\Theta=0},$$

and the linearity of $\delta^a\Phi = iT^a\Phi$, one finds

$$i\int d^dx\; \Gamma^{(2)}_{\Phi\Phi}(x,y)\,T^a\,\delta(x-z) - i\int d^dx\; T^a\,\delta(x-y)\,\Gamma^{(2)}_{\Phi\Phi}(x,z) = 0.$$

Fourier transforming and using momentum conservation (the band kernels preserve translation invariance) gives

$$\Gamma^{(2)}_{\Phi\Phi}(p)\,T^a - T^a\,\Gamma^{(2)}_{\Phi\Phi}(p) = 0, \qquad \text{i.e.} \quad [\Gamma^{(2)}_{\Phi\Phi}(p), T^a] = 0, \tag{7.169}$$

and analogously for $\Theta\Theta$ and cross sectors (when allowed). Thus each two-point kernel commutes with the G-action; decomposing the field space into G-irreps, Schur's lemma implies that $\Gamma^{(2)}(p)$ is block-diagonal in irreps and proportional to the identity on each irrep block.

(ii) No spurionic masses / irrep-wise identity. By definition, a (quadratic) mass term is the momentum-independent part of the 1PI two-point kernel at $p = 0$:

$$M_\Phi := \Gamma^{(2)}_{\Phi\Phi}(p=0).$$

From (7.169) with $p = 0$ we have $[M_\Phi, T^a] = 0$ for all a. Hence M_Φ must be proportional to the identity on each irrep block and cannot mix inequivalent irreps. Any would-be "spurionic" mass mixing different irreps would transform nontrivially and violate the commutation with T^a, contradicting (7.169). The same reasoning applies to Θ and to allowed Φ–Θ mixing: the mass matrix must be G-invariant and therefore block-diagonal in G-irreps, with each block a scalar multiple of the identity.

(iii) Current transversality and tensor structure. Introduce a source A^a_μ coupled to the conserved Noether current J^μ_a of G, and denote by $\Pi^{\mu\nu}_{ab}(x-y)$ the (band-limited) 1PI current–current kernel (polarisation tensor). The global Ward identity yields, upon varying with respect to A^a_μ and A^b_ν and Fourier transforming,

$$p_\mu\,\Pi^{\mu\nu}_{ab}(p) = \mathcal{A}^\nu_{ab}(p), \tag{7.170}$$

where $\mathcal{A}^\nu_{ab}(p)$ collects possible contact (local) terms and anomaly contributions. At the symmetric vacuum in Quadrant I, the anomaly filters enforce the absence of gauge/global G anomalies (cohomological triviality), so $\mathcal{A}^\nu_{ab}(p) = 0$ (see Section 7.2.8 and the cohomology baseline). Therefore

$$p_\mu\,\Pi^{\mu\nu}_{ab}(p) = 0. \tag{7.171}$$

Lorentz covariance on the band implies the general decomposition

$$\Pi^{\mu\nu}_{ab}(p) = \Pi^{\mathrm{T}}_{ab}(p^2)\,\Pi^{\mu\nu}_{\mathrm{T}}(p) + \Pi^{\mathrm{L}}_{ab}(p^2)\,\Pi^{\mu\nu}_{\mathrm{L}}(p), \quad \Pi^{\mu\nu}_{\mathrm{T}} = \eta^{\mu\nu} - \frac{p^\mu p^\nu}{p^2}, \quad \Pi^{\mu\nu}_{\mathrm{L}} = \frac{p^\mu p^\nu}{p^2}.$$

Contracting with p_μ and using (7.171) forces $\Pi^{\mathrm{L}}_{ab}(p^2) \equiv 0$, whence

$$\Pi^{\mu\nu}_{ab}(p) = \Pi_{ab}(p^2)\,\Pi^{\mu\nu}_{\mathrm{T}}(p),$$

as claimed. Since the band/IR operations commute with G and BRST (eq. (7.145)), no extra longitudinal pieces are generated by the embedding; anomaly freedom ensures there is no residual local breaking on the right-hand side of (7.170).

Combining the three parts establishes items (i)–(iii). □

Proposition 7.232 (Three-point Ward–Takahashi identity (schematic))**.** *Let $\Gamma^{\mu;AB}_a(p,q)$ be the 1PI vertex for J^μ_a with fields carrying indices A, B in a G-representation. Then*

$$p_\mu\,\Gamma^{\mu;AB}_a(p,q) = i\Big[(T_a)^A{}_C\,\Gamma^{CB}(q+p) - \Gamma^{AC}(q)\,(T_a)^B{}_C\Big],$$

with all quantities band-limited. This enforces G-covariant tensor bases for three-point structures and implies selection rules for vanishing couplings between inequivalent irreps.

Proof. All statements are understood *on the band*; by (7.145), the UV projector and IR window commute with G and with BRST and carry ghost number 0.

Setup with a local spurion and the local Ward identity. Couple the conserved G-current J^μ_a to a classical source (spurion) $A^a_\mu(x)$ and regard Γ as a functional $\Gamma[\Phi, \Theta; A, K]$ (sources K omitted below for brevity). Promote the constant G parameter to a local function $\epsilon^a(x)$ and consider the spurionic variations

$$\delta_\epsilon \Phi = i\,\epsilon^a T_a \Phi, \qquad \delta_\epsilon \Theta = i\,\epsilon^a T_a \Theta, \qquad \delta_\epsilon A^a_\mu = -\partial_\mu \epsilon^a,$$

with ghosts $(c, \bar{c}, b)$ taken G-singlets. G-invariance of the action and of the gauge fermion at the symmetry point, together with the commutation of band/IR operations with G, imply the *local* Ward identity for the 1PI functional (no anomaly in Quadrant I):

$$\partial^x_\mu \frac{\delta\Gamma}{\delta A^a_\mu(x)} = i\,\frac{\delta\Gamma}{\delta\Phi^A(x)}(T_a)^A{}_B\,\Phi^B(x) + i\,\frac{\delta\Gamma}{\delta\Theta^A(x)}(T_a)^A{}_B\,\Theta^B(x). \tag{7.172}$$

(The band/IR kernels commute with ∂_μ and with the G-action, so (7.172) holds "on the band" without extra contact terms.)

Differentiate to obtain the three-point identity. Differentiate (7.172) w.r.t. $\Phi^A(y)$ and $\Phi^B(z)$ and set $(\Phi, \Theta, A) = (0, 0, 0)$:

$$\partial^x_\mu\,\Gamma^{\mu;AB}_a(x; y, z) = i\,(T_a)^A{}_C\,\Gamma^{CB}(y, z)\,\delta(x - y) - i\,\Gamma^{AC}(y, z)\,(T_a)^B{}_C\,\delta(x - z),$$

where

$$\Gamma^{\mu;AB}_a(x; y, z) := \left.\frac{\delta^3\Gamma}{\delta A^a_\mu(x)\,\delta\Phi^A(y)\,\delta\Phi^B(z)}\right|_0, \qquad \Gamma^{CB}(y, z) := \left.\frac{\delta^2\Gamma}{\delta\Phi^C(y)\,\delta\Phi^B(z)}\right|_0.$$

(The same steps apply for Θ or mixed Φ/Θ legs.)

Fourier transform and momentum routing. Define Fourier transforms with the total momentum conserved by translation invariance (preserved by the band/IR embedding):

$$\Gamma_a^{\mu;AB}(p,q) := \int d^dx\, d^dy\, d^dz\; e^{\,ip\cdot x+iq\cdot y-i(q+p)\cdot z}\,\Gamma_a^{\mu;AB}(x;y,z),$$
$$\Gamma^{CB}(k) := \int d^dy\, d^dz\; e^{\,ik\cdot(y-z)}\,\Gamma^{CB}(y,z).$$

Multiplying the x-space identity by $e^{\,ip\cdot x+iq\cdot y-i(q+p)\cdot z}$ and integrating, the derivatives ∂_μ^x become ip_μ, and the delta functions set $x=y$ or $x=z$, yielding

$$p_\mu\,\Gamma_a^{\mu;AB}(p,q) = i\Big[(T_a)^A{}_C\,\Gamma^{CB}(q+p) - \Gamma^{AC}(q)\,(T_a)^B{}_C\Big].$$

This is the Ward–Takahashi identity in momentum space (with band-limited kernels); see, e.g., [82, 409, 91].

Remarks. (i) The same derivation goes through for Θ-legs and for mixed Φ/Θ legs, with the corresponding representation matrices. (ii) The anomaly filters ensure there is no local breaking on the r.h.s. of (7.172); hence no extra contact terms survive in momentum space. (iii) Because $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ are G-scalars and commute with ∂_μ, the band embedding does not spoil the identity.

This completes the proof. □

Corollary 7.233 (Projector decomposition and $\Pi_{\mathrm{T}}/\Pi_{\mathrm{L}}$ split)**.** *In Lorentz-invariant sectors, any two-point tensor kernel admits the decomposition* $\Gamma^{\mu\nu}(p) = \Gamma_{\mathrm{T}}(p^2)\,\Pi_{\mathrm{T}}^{\mu\nu}(p) + \Gamma_{\mathrm{L}}(p^2)\,\Pi_{\mathrm{L}}^{\mu\nu}(p)$*, with* $\Pi_{\mathrm{L}}^{\mu\nu}(p) = p^\mu p^\nu/p^2$*. At the symmetry point and for conserved currents,* $\Gamma_{\mathrm{L}} \equiv 0$*. G-invariance implies* Γ_{T} *is irrep-wise scalar, i.e. identical across components within the same irrep.*

Proof of Corollary 7.233. Work in a Lorentz-invariant sector so any rank-2 tensor built from p^μ and $\eta^{\mu\nu}$ decomposes as

$$\Gamma^{\mu\nu}(p) = A(p^2)\,\Pi_{\mathrm{T}}^{\mu\nu}(p) + B(p^2)\,\Pi_{\mathrm{L}}^{\mu\nu}(p), \qquad \Pi_{\mathrm{T}}^{\mu\nu} := \eta^{\mu\nu} - \frac{p^\mu p^\nu}{p^2}, \quad \Pi_{\mathrm{L}}^{\mu\nu} := \frac{p^\mu p^\nu}{p^2}.$$

This is the unique Lorentz-covariant decomposition. For a conserved current, the three-point Ward–Takahashi identity

$$p_\mu\,\Gamma_a^{\mu;AB}(p,q) = i\Big[(T_a)^A{}_C\,\Gamma^{CB}(q+p) - \Gamma^{AC}(q)\,(T_a)^B{}_C\Big]$$

(Proposition 7.232) implies transversality of the corresponding two-point kernel at the symmetry point, $p_\mu\,\Gamma^{\mu\nu}(p) = 0$, hence $B \equiv 0$ and only the transverse projector survives. Finally, G-invariance means $\Gamma^{\mu\nu}$ intertwines equivalent G-representations; by Schur's lemma it is proportional to the identity within each irrep, so Γ_{T} is irrep-wise scalar. (See also [82, 91].) □

Proposition 7.234 (Stability under renormalisation and anomaly filters)**.** *If the anomaly filters of Section 7.2.8 hold, the breaking of (7.166) is at most linear in external sources (QAP) and can be removed by local counterterms; thus the Ward identities are stable under renormalisation on the band.*

Proof of Proposition 7.234. Assume the anomaly filters of Section 7.2.8 hold so the local Ward identity for the 1PI functional Γ is valid on the band (no anomaly term):

$$\partial_\mu^x \frac{\delta\Gamma}{\delta A_\mu^a(x)} = i\,\frac{\delta\Gamma}{\delta\Phi^A(x)}(T_a)^A{}_B\Phi^B(x) + i\,\frac{\delta\Gamma}{\delta\Theta^A(x)}(T_a)^A{}_B\Theta^B(x),$$

with the UV projector $P_x(\lambda_{\mathrm{UV}})$ and IR window $K_{\mathrm{IR}}^{(\ell)}$ commuting with ∂_μ and the G-action. By the Quantum Action Principle (QAP), any breaking that appears at intermediate renormalisation steps is a *local* polynomial in fields and sources, at most linear in external sources, and constrained by the same quantum numbers. Such local breakings can be cancelled order by order by adding local counterterms chosen within the symmetry-allowed basis, thereby restoring the Ward identity after renormalisation. Because band/IR operations are linear, *-preserving, and commute with G (cf. Definition 7.28 and Proposition 7.30), the same argument applies on the band. Therefore the Ward identities are stable under renormalisation on the band. (Background: [82, 91].) □

Remark 7.235 (Operational baseline)**.** Propositions 7.229–7.234 provide the *exact* symmetric baseline used in Section 7.4: transversality, irrep block-diagonality, and the absence of spurionic mass/mixing at the symmetry point are the invariants against which law/state-channel visibilities are compared in other quadrants.

7.5.5 Nielsen Identity at the Symmetry Point

Gauge-parameter variations of Γ are controlled by the Nielsen (gauge-parameter) identity:

$$\partial_{\alpha_{\mathrm{gf}}}\Gamma = \int d^dx\,\frac{\delta\Gamma}{\delta\varphi^A(x)}\,\mathcal{N}^A(x;\alpha_{\mathrm{gf}}) \quad\Rightarrow\quad \partial_{\alpha_{\mathrm{gf}}}\Big(\text{on-shell observables}\Big) = 0, \tag{7.173}$$

where $A \in \{\Phi, \Theta, c, \bar{c}, b\}$ and

$$\mathcal{N}^A(x;\alpha_{\mathrm{gf}}) := -\,\frac{\delta}{\delta K_A(x)}\Big(\partial_{\alpha_{\mathrm{gf}}}\Psi\Big) \;\equiv\; \Big\langle\, \zeta\,\partial_{\alpha_{\mathrm{gf}}}\Psi \,\Big\rangle_\Gamma.$$

The first equality is the antibracket/linearised form $\partial_{\alpha_{\mathrm{gf}}}\Gamma = -\,\mathcal{B}_\Gamma\Big(\partial_{\alpha_{\mathrm{gf}}}\Psi\Big)$ (cf. Propositions 7.212 and 7.218), expanded at vanishing sources so that the variation is proportional to the equations of motion $\delta\Gamma/\delta\varphi^A$. The second (expectation) form is the standard insertion map induced by Γ. Since $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ commute with $\hat{Q}_{\mathrm{BRST}}$ eq. (7.145), (7.173) holds *on the band*; on-shell physical poles, visibilities, and certified plateaus are gauge-parameter independent in Quadrant I [117, 91].

Proposition 7.236 (Band Nielsen identity in EOM form)**.** *Let α_{gf} enter only through the gauge fermion $\Psi(\alpha_{\mathrm{gf}})$, and assume the anomaly filters hold so that BRST is nilpotent on the band. Then*

$$\partial_{\alpha_{\mathrm{gf}}}\Gamma = -\,\mathcal{B}_\Gamma\Big(\partial_{\alpha_{\mathrm{gf}}}\Psi\Big) = \int d^dx\,\frac{\delta\Gamma}{\delta\varphi^A(x)}\,\mathcal{N}^A(x;\alpha_{\mathrm{gf}}),$$

with $\mathcal{N}^A$ defined above. In particular, $\partial_{\alpha_{\mathrm{gf}}}\Gamma$ is proportional to the equations of motion and thus $\mathcal{B}_\Gamma$-exact.

Proof idea. Differentiate the band-limited path integral Z with respect to $\alpha_{\rm gf}$; only $\zeta\Psi$ varies. Passing to Γ and using the antibracket representation of the linearised Slavnov operator (Prop. 7.212) gives $\partial_{\alpha_{\rm gf}}\Gamma = -(\Gamma, \partial_{\alpha_{\rm gf}}\Psi) = -\mathcal{B}_\Gamma(\partial_{\alpha_{\rm gf}}\Psi)$. Expanding the antibracket at vanishing sources yields the EOM form with the antifield (source) derivative: $\mathcal{N}^A(x;\alpha_{\rm gf}) := -\delta(\partial_{\alpha_{\rm gf}}\Psi)/\delta K_A(x)$. Band/IR operations commute with BRST and carry ghost number 0 eq. (7.145), so no additional terms arise.

Proof. **Setup and extended Slavnov–Taylor identity.** Gauge fixing enters the band-limited generating functional only through the BRST-exact term $s\,\Psi(\alpha_{\rm gf})$, with $s^2 = 0$ on the band by the anomaly filters (Definition 7.28 and Proposition 7.30 and eq. (7.145)). Introduce the standard BRST doublet $(\alpha_{\rm gf}, \chi)$ with $s\,\alpha_{\rm gf} = \chi$, $s\,\chi = 0$, and define the gauge-fixed classical action $\Sigma(\Phi, K; \alpha_{\rm gf}, \chi) := \Sigma_0(\Phi, K) + s\,\Psi(\alpha_{\rm gf}) - \chi\,\partial_{\alpha_{\rm gf}}\Psi$. The 1PI vertex functional Γ satisfies the (band) Slavnov functional identity

$$\mathcal{S}(\Gamma) = 0, \qquad \mathcal{S}(F) := \int d^dx \left(\frac{\delta F}{\delta K_A}\frac{\delta F}{\delta \varphi^A} + b^a \frac{\delta F}{\delta \bar{c}^a} \right),$$

with the usual Zinn–Justin sources K_A and (algebraic) Nakanishi–Lautrup fields b^a [91, 82]. Differentiating the *extended* identity $\mathcal{S}(\Gamma) + \chi\,\partial_{\alpha_{\rm gf}}\Gamma = 0$ with respect to χ and then setting $\chi = 0$ gives the Nielsen identity in linearised (BV) form:

$$\partial_{\alpha_{\rm gf}}\Gamma = -\,\mathcal{B}_\Gamma\big(\partial_{\alpha_{\rm gf}}\Psi\big), \qquad \mathcal{B}_\Gamma X := \int d^dx \left(\frac{\delta\Gamma}{\delta K_A}\frac{\delta X}{\delta \varphi^A} + \frac{\delta\Gamma}{\delta \varphi^A}\frac{\delta X}{\delta K_A} + b^a \frac{\delta X}{\delta \bar{c}^a} \right), \tag{7.174}$$

where we used the antibracket/linearised-Slavnov equivalence (Proposition 7.212). All steps are valid "on the band" since $P_x(\lambda_{\rm UV})$ and $K_{\rm IR}^{(\ell)}$ commute with s, carry ghost number 0, and commute with derivatives (Definition 7.28 and Proposition 7.30).

EOM representation. Expand $\mathcal{B}_\Gamma$ on $X = \partial_{\alpha_{\rm gf}}\Psi$:

$$\mathcal{B}_\Gamma\big(\partial_{\alpha_{\rm gf}}\Psi\big) = \int d^dx \left(\frac{\delta\Gamma}{\delta K_A}\frac{\delta}{\delta\varphi^A}\partial_{\alpha_{\rm gf}}\Psi + \frac{\delta\Gamma}{\delta\varphi^A}\frac{\delta}{\delta K_A}\partial_{\alpha_{\rm gf}}\Psi + b^a\frac{\delta}{\delta\bar{c}^a}\partial_{\alpha_{\rm gf}}\Psi \right).$$

Define the (band) Nielsen kernel by the canonical rule already fixed above,

$$\mathcal{N}^A(x;\alpha_{\rm gf}) \;:=\; -\frac{\delta}{\delta K_A(x)}\partial_{\alpha_{\rm gf}}\Psi,$$

so the second term becomes $-\int(\delta\Gamma/\delta\varphi^A)\,\mathcal{N}^A$. For the remaining two pieces we use: (i) the algebraic b–equation of motion eliminates the b–term (or, equivalently, expresses it as a $\mathcal{B}_\Gamma$–exact variation that vanishes in the cohomology of ghost number 0); (ii) in the gauge-fixed coordinates obtained by the canonical map generated by Ψ (BV gauge fixing), the $\alpha_{\rm gf}$–dependence of Ψ is entirely along the antifield/source direction,[6] hence $(\partial_{\alpha_{\rm gf}}\Psi)_{,\varphi^A} = 0$ at fixed K, and the first term vanishes. We thus obtain

$$\mathcal{B}_\Gamma\big(\partial_{\alpha_{\rm gf}}\Psi\big) = -\int d^dx\,\frac{\delta\Gamma}{\delta\varphi^A(x)}\,\mathcal{N}^A(x;\alpha_{\rm gf}),$$

[6] Concretely, after the canonical transformation $\varphi^*_A \mapsto K_A + \delta\Psi/\delta\varphi^A$, the functional $\partial_{\alpha_{\rm gf}}\Psi$ acquires only K–dependence at fixed (φ, K), so $(\partial_{\alpha_{\rm gf}}\Psi)_{,\varphi^A} = 0$ at fixed K. If one keeps a residual φ–dependence, the associated term is itself $\mathcal{B}_\Gamma$–exact and can be absorbed by a local field redefinition; see the homological decomposition discussed with Proposition 7.212.

and, by (7.174),

$$\partial_{\alpha_{\mathrm{gf}}}\Gamma = \int d^d x\, \frac{\delta\Gamma}{\delta\varphi^A(x)}\, \mathcal{N}^A(x;\alpha_{\mathrm{gf}}).$$

Conclusion and exactness. Therefore $\partial_{\alpha_{\mathrm{gf}}}\Gamma$ is proportional to the Euler–Lagrange equations and is $\mathcal{B}_\Gamma$–exact. In particular, all (band) BRST-closed observables—such as the visibilities used in this chapter—are independent of α_{gf} when evaluated in $\mathcal{H}_{\mathrm{phys}}$ (cf. the Nielsen independence remarks in Remark 6.254). □

Lemma 7.237 (On-shell gauge-parameter independence)**.** *Let $\mathcal{O}[\Phi,\Theta]$ be a BRST-closed, ghost-number 0 functional representing a physical observable (e.g. an S-matrix element, a pole residue, a certified visibility readout restricted to $\mathcal{H}_{\mathrm{phys}}$). Then*

$$\partial_{\alpha_{\mathrm{gf}}}\, \mathcal{O}\Big|_{\text{on shell}} = 0.$$

Proof idea. Differentiate $\mathcal{O}$: $\partial_{\alpha_{\mathrm{gf}}}\mathcal{O} = \int d^d x\, \frac{\delta\mathcal{O}}{\delta\varphi^A(x)}\, \mathcal{N}^A(x)$, using Prop. 7.236. On shell (stationary point of Γ) we have $\delta\Gamma/\delta\varphi^A = 0$. Since $\mathcal{N}^A$ is generated from $\partial_{\alpha_{\mathrm{gf}}}\Psi$ by one antifield derivative, it carries the same EOM proportionality; hence its insertion vanishes on the solution space of the field equations. Moreover, $\mathcal{O}$ being BRST-closed eliminates possible $\mathcal{B}_\Gamma$-exact contributions. Therefore $\partial_{\alpha_{\mathrm{gf}}}\mathcal{O} = 0$ on shell.

Proof. **Preliminaries.** Work on the band where the UV projector and IR window are BRST-closed and commute with derivatives and internal symmetries (cf. eq. (7.145)); the Slavnov functional identity holds, $\mathcal{S}(\Gamma) = 0$, with linearised operator $\mathcal{B}_\Gamma$ (ghost number +1). Gauge-parameter dependence enters only via the gauge fermion $\Psi(\alpha_{\mathrm{gf}})$.

Step 1: Band Nielsen identity for Γ. By Proposition 7.236 one has the (EOM) Nielsen form

$$\partial_{\alpha_{\mathrm{gf}}}\Gamma = -\,\mathcal{B}_\Gamma\big(\partial_{\alpha_{\mathrm{gf}}}\Psi\big) = \int d^d x\, \frac{\delta\Gamma}{\delta\varphi^A(x)}\, \mathcal{N}^A(x;\alpha_{\mathrm{gf}}), \qquad \mathcal{N}^A(x;\alpha_{\mathrm{gf}}) := -\frac{\delta}{\delta K_A(x)}\big(\partial_{\alpha_{\mathrm{gf}}}\Psi\big). \tag{7.175}$$

Step 2: How on-shell solutions move with α_{gf}. Let $\varphi_\star(\alpha_{\mathrm{gf}})$ solve the field equations, $\delta\Gamma/\delta\varphi^A\big|_{\varphi_\star} = 0$. Differentiate this stationarity condition w.r.t. α_{gf}:

$$0 = \frac{d}{d\alpha_{\mathrm{gf}}}\Big(\frac{\delta\Gamma}{\delta\varphi^A}\Big)_{\varphi_\star} = \Gamma_{AB}(\varphi_\star)\,\partial_{\alpha_{\mathrm{gf}}}\varphi^B_\star \;+\; \Big(\frac{\delta}{\delta\varphi^A}\,\partial_{\alpha_{\mathrm{gf}}}\Gamma\Big)_{\varphi_\star},$$

where $\Gamma_{AB} := \delta^2\Gamma/\delta\varphi^A\delta\varphi^B$. Using (7.175) and differentiating under the integral,

$$\frac{\delta}{\delta\varphi^A}\,\partial_{\alpha_{\mathrm{gf}}}\Gamma = \Gamma_{AC}\,\mathcal{N}^C + \frac{\delta\Gamma}{\delta\varphi^C}\,\frac{\delta\mathcal{N}^C}{\delta\varphi^A}.$$

At $\varphi_\star$ the second term vanishes; thus $\Gamma_{AB}(\varphi_\star)\,\big(\partial_{\alpha_{\mathrm{gf}}}\varphi^B_\star - \mathcal{N}^B(\varphi_\star)\big) = 0$. Assuming the usual regularity (invertibility of Γ_{AB} on the physical subspace; gauge zero-modes are projected out by the BRST quartet/physical projector), we conclude

$$\partial_{\alpha_{\mathrm{gf}}}\varphi^A_\star = \mathcal{N}^A\big(\varphi_\star\big). \tag{7.176}$$

Step 3: $\alpha_{\rm gf}$-derivative of on-shell $\mathcal{O}$. Since $\mathcal{O}$ has no explicit antifield dependence and gauge-parameter enters only through the fields,

$$\partial_{\alpha_{\rm gf}} \mathcal{O}\big(\varphi_\star(\alpha_{\rm gf})\big) = \int d^d x \, \frac{\delta \mathcal{O}}{\delta \varphi^A(x)} \, \partial_{\alpha_{\rm gf}} \varphi^A_\star(x).$$

Using (7.176) gives the *characteristic* (Nielsen) contraction

$$\partial_{\alpha_{\rm gf}} \mathcal{O}\Big|_{\text{on shell}} = \int d^d x \, \frac{\delta \mathcal{O}}{\delta \varphi^A(x)} \, \mathcal{N}^A(x;\alpha_{\rm gf}). \tag{7.177}$$

Step 4: BRST-closedness ⇒ vanishing of the characteristic contraction. By definition of $\mathcal{N}^A$ and the BV antibracket $(\cdot,\cdot)$,

$$\int d^d x \, \frac{\delta \mathcal{O}}{\delta \varphi^A(x)} \, \mathcal{N}^A(x) = - \int d^d x \, \frac{\delta \mathcal{O}}{\delta \varphi^A(x)} \, \frac{\delta}{\delta K_A(x)} \big(\partial_{\alpha_{\rm gf}} \Psi\big) = - \, (\mathcal{O}, \partial_{\alpha_{\rm gf}} \Psi)\Big|_{K=0}.$$

The Jacobi identity together with $\mathcal{S}(\Gamma) = 0$ implies the standard linearisation property $\mathcal{B}_\Gamma X = (\Gamma, X) + \cdots$ (the extra $b\delta/\delta\bar{c}$ term does not contribute here: $\mathcal{O}$ has ghost number 0 and no $\bar{c}$-dependence). Since $\mathcal{O}$ is BRST-closed, $\mathcal{B}_\Gamma \mathcal{O} = 0$, hence $(\Gamma, \mathcal{O}) \approx 0$ on the stationary surface.[7] Therefore $(\mathcal{O}, \partial_{\alpha_{\rm gf}} \Psi)$ is $\mathcal{B}_\Gamma$-exact on shell, and BRST-exact deformations do not change physical (cohomology) classes on $\mathcal{H}_{\rm phys}$. Accordingly, the right-hand side of (7.177) vanishes, giving $\partial_{\alpha_{\rm gf}} \mathcal{O}\big|_{\text{on shell}} = 0$.

Conclusion. On-shell solutions move along the Nielsen vector field (7.176); BRST-closed observables are constant along this flow in the physical sector, hence are gauge-parameter independent on shell. □

Proposition 7.238 (Extended Slavnov trick ⇒ band Nielsen identity)**.** *Promote the gauge parameter to a BRST doublet $(\alpha_{\rm gf}, \chi)$ with $s\,\alpha_{\rm gf} = \chi$ and $s\,\chi = 0$, and extend the classical action by the s-exact term $s\int d^d x\, \alpha_{\rm gf}\Psi = \int d^d x \, \big(\chi\,\Psi - \alpha_{\rm gf}\, s\Psi\big)$. Assuming the anomaly filters (nilpotent BRST on the band) and using that the UV/IR tagging commutes with s (cf. eq. (7.145)), the* extended *Slavnov functional identity*

$$\widetilde{\mathcal{S}}(\Gamma) \equiv \mathcal{S}(\Gamma) \; + \; \chi\, \partial_{\alpha_{\rm gf}} \Gamma \; = \; 0$$

holds for the 1PI functional Γ on the band. Differentiating w.r.t. χ and setting $\chi = 0$ gives

$$\partial_{\alpha_{\rm gf}} \Gamma = - \, \mathcal{B}_\Gamma\big(\partial_{\alpha_{\rm gf}} \Psi\big),$$

i.e. the band Nielsen identity of EOM form (identical to Proposition 7.236).

Proof. By construction, the χ-coupling enters Γ only through the s-exact insertion $\int \chi\,\Psi$, and the extended BRST invariance yields $\widetilde{\mathcal{S}}(\Gamma) = 0$. Taking $\partial/\partial\chi$ and then $\chi{=}0$,

$$0 = \frac{\partial}{\partial \chi} \widetilde{\mathcal{S}}(\Gamma)\Big|_{\chi=0} = \frac{\partial}{\partial \chi} \mathcal{S}(\Gamma)\Big|_{\chi=0} \; + \; \partial_{\alpha_{\rm gf}} \Gamma.$$

[7] More precisely, $(\Gamma, \mathcal{O})$ is proportional to the equations of motion when $\mathcal{O}$ is antifield independent; this vanishes on shell.

The chain rule together with the linearisation of $\mathcal{S}$ at Γ gives $\frac{\partial}{\partial\chi}\mathcal{S}(\Gamma)\big|_{\chi=0} = \mathcal{B}_\Gamma\Big(\frac{\partial\Gamma}{\partial\chi}\big|_{\chi=0}\Big)$. Since the only χ-dependence is the linear insertion $\int\chi\,\Psi$, the Quantum Action Principle and s-closure on the band imply $\frac{\partial\Gamma}{\partial\chi}\big|_{\chi=0} = \partial_{\alpha_{\rm gf}}\Psi$. Hence $0 = \mathcal{B}_\Gamma(\partial_{\alpha_{\rm gf}}\Psi) + \partial_{\alpha_{\rm gf}}\Gamma$, which is the stated identity. Band/IR operations commute with s and carry ghost number 0 so no extra terms arise. □

Corollary 7.239 (On-shell independence via the doublet)**.** *For any BRST-closed, ghost-number* 0 *observable* $\mathcal{O}$ *evaluated on stationary points of* Γ, $\partial_{\alpha_{\rm gf}}\mathcal{O}|_{\text{on shell}} = 0$.

Proof. Combine Proposition 7.238 with the characteristic flow $\partial_{\alpha_{\rm gf}}\varphi^A_\star = \mathcal{N}^A(\varphi_\star)$ derived in the proof of Lemma 7.237; BRST-closedness of $\mathcal{O}$ implies constancy along this flow in $\mathcal{H}_{\rm phys}$. □

Corollary 7.240 (Pole-mass Nielsen identity (band-limited))**.** *Let* $\Delta^{-1}(p)$ *denote a renormalised* 1*PI two-point kernel for a physical mode, and suppose* $\Delta^{-1}(p^2 = m_\star^2) = 0$ *with finite residue. Then near the pole*

$$\partial_{\alpha_{\rm gf}}\Delta^{-1}(p) = \mathcal{C}(p)\,\Delta^{-1}(p) \quad \Rightarrow \quad \partial_{\alpha_{\rm gf}} m_\star^2 = 0,$$

with $\mathcal{C}$ *regular at* $p^2 = m_\star^2$*. Hence the pole position is gauge-parameter independent on the band.*

Proof. Apply Prop. 7.236 to the quadratic sector: the variation of the two-point kernel is EOM-proportional, i.e. right-multiplication by a regular form factor (and its transpose in mixed sectors). Evaluating at the zero of Δ^{-1} kills the r.h.s., implying the pole condition is stationary in $\alpha_{\rm gf}$. Regularity follows from the anomaly filters and standard QAP arguments [117, 91]. □

Corollary 7.241 (Gauge-parameter independence of certified visibilities)**.** *For the holosymmetric probe on* $\mathcal{H}_{\rm phys}$*, the complex coherence amplitude* $\mathcal{C}_\alpha$ *and any phase-fixed visibility* Λ_α *(defined on a certified plateau) obey*

$$\partial_{\alpha_{\rm gf}}\mathcal{C}_\alpha = 0, \qquad \partial_{\alpha_{\rm gf}}\Lambda_\alpha = 0,$$

up to admissible regulator tolerances.

Proof. Insert the Nielsen identity into the readout expressions (either (7.115) or (7.98)). All $\partial_{\alpha_{\rm gf}}$-variations reduce to $\mathcal{B}_\Gamma$-exact (EOM) insertions, which vanish on $\mathcal{H}_{\rm phys}$ by BRST closure and on-shell conditions; band/IR commutation guarantees these manipulations hold on the band. □

Remark 7.242 (Ghost number and sector structure)**.** Ghost-number counting gives $\mathrm{gh}(\partial_{\alpha_{\rm gf}}\Psi) = 0$ and $\mathrm{gh}(\mathcal{N}^A) = -1 + \mathrm{gh}(K_A)$, so the insertion in (7.173) preserves total ghost number 0. If sector projectors Π_β commute with $P_x(\lambda_{\rm UV})$ and $K_{\rm IR}^{(\ell)}$, the identity holds sectorwise, yielding projector-resolved Nielsen statements consistent with Section 7.4.3.

7.5.6 Cohomological Content and Physical Operators

Let $\mathcal{B}_\Gamma$ be the linearised ST operator (7.151). Physical, gauge-invariant composite operators $\mathcal{O}$ at the symmetric point are the cohomology classes

$$\mathcal{O} \in H^0(\mathcal{B}_\Gamma) \qquad \text{with} \qquad \mathcal{B}_\Gamma \mathcal{O} = 0, \quad \mathcal{O} \sim \mathcal{O} + \mathcal{B}_\Gamma X. \tag{7.178}$$

All statements below are understood *on the band*; the UV/IR knobs commute with BRST and carry ghost number 0, cf. (7.145).

Proposition 7.243 (Well-definedness on the physical sector)**.** *Let $\mathcal{O}$ be a local composite with $\mathrm{gh}(\mathcal{O}) = 0$. If $\mathcal{B}_\Gamma \mathcal{O} = 0$, then for any physical in/out states $|\Psi_{\mathrm{in}}\rangle, |\Psi_{\mathrm{out}}\rangle \in \mathcal{H}_{\mathrm{phys}}$ (BRST-closed modulo null states) and any set of additional BRST-closed insertions,*

$$\langle \Psi_{\mathrm{out}}| \, \mathcal{O} \, |\Psi_{\mathrm{in}}\rangle \quad \textit{is independent of the representative} \quad \mathcal{O} \mapsto \mathcal{O} + \mathcal{B}_\Gamma X.$$

Proof. On $\mathcal{H}_{\mathrm{phys}}$ (BRST cohomology at ghost number 0), the insertion of a $\mathcal{B}_\Gamma$-exact operator is a BRST commutator and vanishes by the quartet mechanism/positivity and cyclicity; see algebraic BRST treatments [442, 117]. Band/IR commutation (7.145) prevents extra contact terms. □

Proposition 7.244 (Källén–Lehmann positivity on $\mathcal{H}_{\mathrm{phys}}$)**.** *Let $\mathcal{O} \in H^0(\mathcal{B}_\Gamma)$ be local and Hermitian. Then its two-point function on $\mathcal{H}_{\mathrm{phys}}$ admits the Källén–Lehmann representation*

$$\langle \mathcal{O}(x)\, \mathcal{O}(0) \rangle_{\mathrm{phys}} = \int_0^\infty d\mu^2 \; \rho_{\mathcal{O}}(\mu^2)\, \Delta_+(x;\mu^2), \qquad \rho_{\mathcal{O}}(\mu^2) \geq 0,$$

with Δ_+ the free Wightman function and $\rho_{\mathcal{O}}$ a non-negative spectral density [241, 82]. The spectral measure and support are invariant under $\mathcal{O} \mapsto \mathcal{O} + \mathcal{B}_\Gamma X$.

Proof. Restrict correlators to $\mathcal{H}_{\mathrm{phys}}$ via $\mathsf{P}_{\mathrm{phys}}$ (quartet decoupling) [442, 117]. The induced inner product is positive and the Wightman axioms hold for band-smeared locals [241]; spectral decomposition then yields the stated form [82]. $\mathcal{B}_\Gamma$-exact shifts drop out by the previous proposition. □

Proposition 7.245 (Renormalisation stability of $H^0(\mathcal{B}_\Gamma)$)**.** *Assume the anomaly filters so that $\mathcal{S}(\Gamma) = 0$ to all orders. Then the space of physical operators is stable under renormalisation: counterterms can be chosen so that the renormalised operator $\mathcal{O}_R$ and $\mathcal{B}_{\Gamma_R}$ satisfy*

$$\mathcal{B}_{\Gamma_R} \mathcal{O}_R = 0, \qquad \mathcal{O}_R \sim \mathcal{O}_R + \mathcal{B}_{\Gamma_R} X_R,$$

and the mixing matrix is block-upper-triangular between cohomology classes and $\mathcal{B}_{\Gamma_R}$-exact operators. Anomalous dimensions on H^0 are gauge-parameter independent [117, 91].

Proof. Algebraic renormalisation (Quantum Action Principle) with preserved Slavnov identity controls composite-operator mixings and ensures stability of BRST cohomology classes [117]. Nielsen identities render gauge-parameter variations $\mathcal{B}_\Gamma$-exact, so eigenvalues on H^0 do not depend on α_{gf} [91]. □

Lemma 7.246 (Sectorwise decomposition)**.** *Let $\{\Pi_\beta\}$ be BRST-closed sector projectors commuting with the UV/IR knobs. If $\mathcal{O} \in H^0(\mathcal{B}_\Gamma)$, then $\mathcal{O}_\beta := \Pi_\beta \mathcal{O} \Pi_\beta$ defines a class in $H^0(\mathcal{B}_\Gamma)$ (possibly trivial), and*

$$\langle \mathcal{O}(x)\, \mathcal{O}(0)\rangle_{\text{phys}} = \sum_\beta \langle \mathcal{O}_\beta(x)\, \mathcal{O}_\beta(0)\rangle_{\text{phys}},$$

with no cross terms.

Proof. $[\Pi_\beta, \mathcal{B}_\Gamma] = 0$ (BRST-closure), so $\mathcal{B}_\Gamma \mathcal{O} = 0 \Rightarrow \mathcal{B}_\Gamma \mathcal{O}_\beta = 0$. Orthogonality removes cross terms. Positivity sectorwise follows from Prop. 7.244. □

Corollary 7.247 (Gauge-parameter independence on shell)**.** *Let $\mathcal{O} \in H^0(\mathcal{B}_\Gamma)$ and consider any on-shell observable built from $\mathcal{O}$ (pole mass/residue, certified visibility from BRST-closed inputs). Then*

$$\partial_{\alpha_{\text{gf}}}(\textit{observable}) = 0 \quad \textit{on shell},$$

up to admissible regulator tolerances [117, 91].

Remark 7.248 (Connection to holosymmetric visibilities)**.** The channel overlap operators used by the holosymmetric probe are assembled from BRST-closed, band-limited ingredients; their matrix elements depend only on cohomology classes, so visibilities/echoes inherit the positivity and Nielsen/BRST robustness discussed above [117, 442, 91].

7.5.7 Background-field Option

Background split and band-limited gauge fixing. Adopt the background-field method (BFM) [114, 91] by splitting the mean/1PI fields into

$$\Phi \to \bar{\Phi} + \phi, \qquad \Theta \to \bar{\Theta} + \vartheta,$$

where $(\bar{\Phi}, \bar{\Theta})$ are classical background fields and (ϕ, ϑ) are the quantum fields integrated over in the path integral. Choose a *background-covariant* gauge fermion Ψ_{bkg} so that background gauge invariance is manifest. All band/IR operations commute with BRST and with the global/background G-action, cf. (7.145); in particular the band-limited sources and kernels are G-scalars.

Background Ward identity and Slavnov identity. Let $\Gamma[\bar{\Phi}, \bar{\Theta}; \phi, \vartheta, K]$ denote the band-limited 1PI action with sources K only for the *quantum* BRST variations. Denote by δ^a_{bkg} the background G-variation:

$$\delta^a_{\text{bkg}} \bar{\Phi} = i\, T^a \bar{\Phi},$$
$$\delta^a_{\text{bkg}} \bar{\Theta} = i\, T^a \bar{\Theta},$$
$$\delta^a_{\text{bkg}}(\phi, \vartheta) \text{ linear s.t. } \bar{\Phi} + \phi,\ \bar{\Theta} + \vartheta \text{ transform covariantly.}$$

Define the *background Ward operator* acting on 1PI functionals by

$$\mathcal{W}^a_{\text{bkg}} := \int d^d x \left(\frac{\delta}{\delta \bar{\Phi}}\, \delta^a_{\text{bkg}} \bar{\Phi} + \frac{\delta}{\delta \bar{\Theta}}\, \delta^a_{\text{bkg}} \bar{\Theta} + \frac{\delta}{\delta K_\Phi}\, \delta^a_{\text{bkg}}(\zeta\phi) + \frac{\delta}{\delta K_\Theta}\, \delta^a_{\text{bkg}}(\zeta\vartheta) \right),$$

where $\delta^a_{\mathrm{bkg}}(\zeta\phi) = \zeta(\delta^a_{\mathrm{bkg}}\phi)$ etc. by background covariance of Ψ_{bkg}. Then, at the symmetry point and *on the band*,

$$\mathcal{W}^a_{\mathrm{bkg}}\,\Gamma = 0, \qquad \mathcal{S}(\Gamma) = 0, \tag{7.179}$$

i.e. the background Ward identity holds *in addition* to the band Slavnov identity (7.150) [114, 91]. The proof is the standard change-of-variables derivation with background-local transformations, and goes through verbatim because band/IR knobs commute with BRST and with G (7.145).

Proposition 7.249 (Compatibility with Slavnov and linearised BRST in BFM)**.** *Assume* (7.179)*. Then*

$$\mathcal{W}^a_{\mathrm{bkg}}\Big(\mathcal{S}(\Gamma)\Big) = 0, \qquad [\mathcal{W}^a_{\mathrm{bkg}},\, \mathcal{B}_\Gamma] = 0.$$

Consequently, the linearised cohomology $\mathsf{H}^\bullet(\mathcal{B}_\Gamma)$ *is* G*-equivariant under the background symmetry.*

Proof. Use the antibracket form $\mathcal{S}(\Gamma) = \frac{1}{2}(\Gamma,\Gamma)$ and that $\mathcal{W}^a_{\mathrm{bkg}}$ is a derivation of the antibracket (background variations are linear). Then $\mathcal{W}^a_{\mathrm{bkg}}\mathcal{S}(\Gamma) = \frac{1}{2}[(\mathcal{W}^a_{\mathrm{bkg}}\Gamma,\Gamma) + (\Gamma, \mathcal{W}^a_{\mathrm{bkg}}\Gamma)] = 0$. For the commutator, $[\mathcal{W}^a_{\mathrm{bkg}}, \mathcal{B}_\Gamma]X = \mathcal{W}^a_{\mathrm{bkg}}(\Gamma, X) - (\Gamma, \mathcal{W}^a_{\mathrm{bkg}}X) = (\mathcal{W}^a_{\mathrm{bkg}}\Gamma, X) = 0$. □

Background Nielsen identity and on-shell independence. If the gauge-fixing parameter α_{gf} enters only via $\Psi_{\mathrm{bkg}}(\alpha_{\mathrm{gf}})$, the band Nielsen identity holds in the BFM:

$$\partial_{\alpha_{\mathrm{gf}}}\Gamma = -\,\mathcal{B}_\Gamma\Big(\partial_{\alpha_{\mathrm{gf}}}\Psi_{\mathrm{bkg}}\Big),$$

which is background-covariant because $\partial_{\alpha_{\mathrm{gf}}}\Psi_{\mathrm{bkg}}$ is a background scalar. Therefore on-shell, BRST-closed observables that are background-gauge invariant are *gauge-parameter independent* on the band [117, 91].

Proposition 7.250 (Transversality and renormalisation simplifications (BFM))**.** *Let* $\Pi^{\mu\nu}_{ab}$ *denote the background current (or background gauge field)* 1*PI two-point kernel. Then*

$$p_\mu\,\Pi^{\mu\nu}_{ab}(p) = 0, \qquad \Pi^{\mu\nu}_{ab}(p) = \Pi_{ab}(p^2)\,\Pi^{\mu\nu}_{\mathrm{T}}(p),$$

with $\Pi^{\mu\nu}_{\mathrm{T}} = \eta^{\mu\nu} - p^\mu p^\nu/p^2$*, exactly on the band. Moreover, renormalisation constants are constrained by background Ward identities; in particular, the gauge coupling renormalisation can be extracted from the background two-point function with manifest gauge invariance [114, 91].*

Sketch. The local background Ward identity (spurion method) implies transversality as in the non-BFM case, now with background sources; commuting of band/IR knobs with derivatives ensures no extra longitudinal terms. Renormalisation constraints follow from requiring (7.179) after counterterms: only background-covariant structures appear, yielding the standard BFM relations [114, 91]. □

Summary. If the background-field method is used, the 1PI action obeys *both* the background Ward identities and the Slavnov identity,

$$\mathcal{W}^a_{\rm bkg}\,\Gamma = 0, \qquad \mathcal{S}(\Gamma) = 0,$$

with all products and sources band-limited. This guarantees: (i) manifest gauge/BRST robustness of physical matrix elements and visibilities; (ii) exact transversality/selection rules enforced by G at the background level; and (iii) gauge-parameter independence of on-shell quantities, with renormalisation simplified by background covariance [114, 91, 117, 442, 82, 241]. Band/IR knobs preserve these identities by (7.145).

7.5.8 Global Ward Identities at the Symmetric Point

Let $\{T_a\}$ be generators of the unbroken global symmetry G in Quadrant I, and let J^μ_a be the associated conserved Noether currents. We assume the UV generator $\mathcal{D}_x$ is G-invariant so that

$$[Q_a, \mathcal{D}_x] = 0 \quad \Rightarrow \quad [Q_a, P_x(\lambda_{\rm UV})] = 0, \qquad Q_a := \int_{\Sigma_t} d\Sigma_\mu\, J^\mu_a, \tag{7.180}$$

and IR smearing is G-blind. Hence all band/IR operations commute with G ("bandwise symmetry"). For any gauge-invariant composite $\mathcal{O}$ (built from BRST-cohomology classes, cf. Section 7.5.6), invariance of $S^{(\lambda_{\rm UV},\ell)}_{\Phi,\Theta}$ under $\delta_a\varphi := i\,T_a\varphi$ yields the *position-space Ward identity* (with all insertions IR-smoothed as in eq. (7.33))

$$\partial^{(x)}_\mu\, \langle J^\mu_a(x)\, \mathcal{O}(x_1,\dots,x_n)\rangle = -i \sum_{j=1}^{n} \delta^{(d)}(x - x_j)\, \langle \delta_a \mathcal{O}(x_1,\dots,x_n)\rangle, \tag{7.181}$$

which descends to $\mathcal{H}_{\rm phys}$ because BRST-exact pieces decouple and $P_x(\lambda_{\rm UV})$ carries ghost number 0 (Section 7.2.6 and eq. (7.30)). In momentum space this becomes the standard *Ward–Takahashi identity* for 1PI functions (computed from Γ in eq. (7.144)) [91, 82]:

$$k_\mu\, \Gamma^\mu_{J_a\,\mathcal{O}_1\dots\mathcal{O}_n}(k; p_1,\dots,p_n) = \sum_{j=1}^{n} \Gamma_{\mathcal{O}_1\dots(\delta_a\mathcal{O}_j)\dots\mathcal{O}_n}(p_1,\dots,p_n). \tag{7.182}$$

At the G-invariant vacuum fixed in Section 7.1, $\delta_a\bar\Phi = \delta_a\bar\Theta = 0$, so the one-point identity reduces to current conservation,

$$\partial_\mu \langle J^\mu_a\rangle_{\bar\Phi,\bar\Theta} = 0. \tag{7.183}$$

Proposition 7.251 (Band-limited local Ward identity)**.** *Under the assumptions above and anomaly filters (so that BRST is nilpotent and global G is non-anomalous), the identity* (7.181) *holds* on the band *for any set of BRST-closed, gauge-invariant composites $\{\mathcal{O}_j\}$.*

Proof. Promote the constant G parameter to a test function $\epsilon_a(x)$ and perform the change of variables $\varphi \mapsto \varphi + i\,\epsilon_a T_a\varphi$ in the band-limited path integral defining correlators. The functional measure is invariant (global linear change), the variation of $S^{(\lambda_{\rm UV},\ell)}_{\Phi,\Theta}$ yields the conserved current via Noether's theorem, and the source insertions vary by $\delta_a\mathcal{O}_j$. Because $P_x(\lambda_{\rm UV})$ and $K^{(\ell)}_{\rm IR}$ commute with G and BRST (ghost number 0), no extra contact terms appear from the band/IR embedding. Integrating by parts transfers $\partial_\mu\epsilon_a$ onto J^μ_a, giving (7.181). BRST-exact contributions vanish on $\mathcal{H}_{\rm phys}$ by quartet decoupling [442, 117]. □

Proposition 7.252 (Momentum-space Ward–Takahashi identity on the band)**.** *Fourier transforming* (7.181) *and passing to* 1*PI kernels via the Legendre transform (at fixed band/IR knobs) yields* (7.182)*. In particular, for a single* $\mathcal{O}$ *in a representation of* G*, the three-point* 1*PI vertex satisfies* $k_\mu \Gamma^\mu_{J_a \mathcal{O}\mathcal{O}} = i[(T_a)\Gamma - \Gamma(T_a)]$ *[91, 82].*

Proof. Standard manipulations [82, 91]: (i) Fourier transform the local identity; (ii) use connected amputated functions and then the Legendre transform to 1PI kernels. The band/IR knobs commute with derivatives and do not alter momentum routing, so the textbook derivation carries over unchanged. □

Corollary 7.253 (Transversality and irrep constraints)**.** *Let* $\Pi^{\mu\nu}_{ab}(k)$ *be the* 1*PI current–current kernel. Then* $k_\mu \Pi^{\mu\nu}_{ab}(k) = 0$ *and* $\Pi^{\mu\nu}_{ab}(k) = \Pi_{ab}(k^2)\,(\eta^{\mu\nu} - k^\mu k^\nu/k^2)$ *at the symmetry point. Moreover, two-point* 1*PI kernels of matter operators commute with the* G*-action and are block-diagonal in* G*-irreps (Schur), with irrep-wise scalar structures; mixing between inequivalent irreps is forbidden [91].*

Proof. Apply (7.182) with $\mathcal{O} = J^\nu_b$ to obtain transversality; apply it with matter operators to deduce commutation with T_a and invoke Schur's lemma. The band embedding preserves these tensorial statements because it is G-blind and derivative-compatible. □

Remark 7.254 (Compatibility with BRST and Nielsen identities)**.** The global Ward identities commute with the Slavnov identity and with the linearised BRST operator on the band (cf. Section 7.5.2), so they descend to the physical cohomology classes that define observables [117, 442]. Gauge-parameter variations are $\mathcal{B}_\Gamma$-exact (Nielsen identity), hence on-shell Ward consequences (transversality, pole positions, certified visibilities) are gauge-parameter independent [91].

Two-point structures and transversality.

Proposition 7.255 (Current two-point tensor structure and KL-positivity on the band)**.** *Let* J^μ_a *denote the conserved Noether currents of the unbroken global symmetry* G *at the symmetric vacuum (Quadrant I), with band/IR operations commuting with* G *and BRST. Then the* 1*PI current–current kernel is transverse and admits the unique transverse decomposition*

$$k_\mu \Gamma^{\mu\nu}_{J_a J_b}(k) = 0 \quad \Rightarrow \quad \Gamma^{\mu\nu}_{J_a J_b}(k) = \Pi^{ab}_{\mathrm{T}}(k^2)\; P^{\mu\nu}_{\mathrm{T}}(k; \bar{\Sigma}^{(\ell)}), \tag{7.184}$$

where the transverse projector with respect to the IR metric $\bar{\Sigma}^{(\ell)}_{\mu\nu}$ *is*

$$P^{\mu\nu}_{\mathrm{T}}(k; \bar{\Sigma}^{(\ell)}) := \bar{\Sigma}^{(\ell)\mu\nu} - \frac{k^\mu k^\nu}{\bar{\Sigma}^{(\ell)\rho\sigma} k_\rho k_\sigma}\,, \qquad k_\mu P^{\mu\nu}_{\mathrm{T}} = 0, \tag{7.185}$$

and $\Pi^{ab}_{\mathrm{T}}(k^2)$ *is a symmetric matrix of scalar form factors in the adjoint indices* (a, b)*. Moreover, on the physical Hilbert space* $\mathcal{H}_{\mathrm{phys}}$ *the (band-smeared) current correlator has a Källén–Lehmann representation with a positive semidefinite spectral matrix:*

$$\langle\!\langle J^\mu_a(k)\, J^\nu_b(-k) \rangle\!\rangle_{\mathcal{H}_{\mathrm{phys}}} = P^{\mu\nu}_{\mathrm{T}}(k; \bar{\Sigma}^{(\ell)}) \int_0^\infty d\mu^2\; \frac{\rho^{ab}_J(\mu^2)}{k^2 + \mu^2 - i0}, \qquad \rho^{ab}_J(\mu^2) \succeq 0. \tag{7.186}$$

Proof. Transversality. Apply the global Ward identity in momentum space to the three-point 1PI vertex with one current insertion and two current legs (or equivalently differentiate the local Ward identity with respect to sources for J_b^ν and amputate). Using (7.182) (derived on the band) one finds $k_\mu\,\Gamma^{\mu\nu}_{J_a J_b}(k) = 0$; by symmetry of the two-point kernel the analogous contraction with k_ν also vanishes. The band/IR embedding commutes with derivatives and the G-action, so no longitudinal contact terms arise (cf. Section 7.5.8; see also [82, 91]).

Tensor structure. In the presence of the IR metric $\bar{\Sigma}^{(\ell)}_{\mu\nu}$, the most general *symmetric* rank-2 tensor built from k^μ and $\bar{\Sigma}^{(\ell)}_{\mu\nu}$ is a linear combination of $\bar{\Sigma}^{(\ell)\mu\nu}$ and $k^\mu k^\nu$. Imposing transversality with respect to k_μ uniquely fixes the combination to be the projector $P^{\mu\nu}_{\mathrm{T}}(k;\bar{\Sigma}^{(\ell)})$ in (7.185). Hence (7.184) with a scalar form factor matrix $\Pi^{ab}_{\mathrm{T}}(k^2)$. Bose symmetry and Hermiticity of the currents imply $\Pi^{ab}_{\mathrm{T}} = \Pi^{ba}_{\mathrm{T}}$ and reality below multi-particle thresholds [91].

Källén–Lehmann form and positivity. Restricting to $\mathcal{H}_{\mathrm{phys}}$ (quartet decoupling) yields a positive-definite inner product; the usual spectral condition and locality (here implemented with band smearing) give the Källén–Lehmann representation for two-point functions of Hermitian operators [241, 82, 91]. Current conservation removes the longitudinal part, leaving the transverse projector as a kinematic factor. The spectral matrix $\rho_J^{ab}(\mu^2)$ is positive semidefinite: for any complex vector v_a, the scalar operator $J_v^\mu := v_a J_a^\mu$ has nonnegative spectral density, whence $v_a^*\,\rho_J^{ab}(\mu^2)\,v_b \geq 0$. This establishes (7.186), consistent with the general positivity statement in eq. (7.12). □

Charge action and algebra on the band. The band-limited charges $Q_a^{(\lambda_{\mathrm{UV}},\ell)} := \int d\Sigma_\mu\, J_a^\mu$ satisfy

$$[Q_a^{(\lambda_{\mathrm{UV}},\ell)},\,\mathcal{O}] = \delta_a\mathcal{O}, \qquad [Q_a^{(\lambda_{\mathrm{UV}},\ell)},\,Q_b^{(\lambda_{\mathrm{UV}},\ell)}] = i f_{ab}{}^c\,Q_c^{(\lambda_{\mathrm{UV}},\ell)}, \tag{7.187}$$

on BRST-closed, gauge-invariant composites; here $f_{ab}{}^c$ are the structure constants of G. Equation (7.180) ensures that $Q_a^{(\lambda_{\mathrm{UV}},\ell)}$ generate the same algebra as the unbanded charges, and $[Q_a^{(\lambda_{\mathrm{UV}},\ell)}, H] = 0$ at the symmetric vacuum.

Proposition 7.256 (Band-limited charge action and Lie algebra)**.** *Assume (i) global G is non-anomalous (anomaly filters hold), (ii) $[Q_a, \mathcal{D}_x] = 0$ so that band/IR operations commute with G (7.180), and (iii) the IR window renders surface terms null. Then for any BRST-closed, gauge-invariant composite $\mathcal{O}$,*

$$[Q_a^{(\lambda_{\mathrm{UV}},\ell)},\mathcal{O}] = \delta_a\mathcal{O}, \qquad [Q_a^{(\lambda_{\mathrm{UV}},\ell)},Q_b^{(\lambda_{\mathrm{UV}},\ell)}] = i f_{ab}{}^c\,Q_c^{(\lambda_{\mathrm{UV}},\ell)}.$$

Proof. Integrate the local Ward identity (7.181) over a Cauchy slice Σ_t with the IR weight. The left-hand side becomes the time derivative of the charge insertion plus a vanishing boundary term; the right-hand side produces the action of δ_a on $\mathcal{O}$. This yields $[Q_a^{(\lambda_{\mathrm{UV}},\ell)},\mathcal{O}] = \delta_a\mathcal{O}$ (the equal-time commutator form of the Ward identity) [82, 91]. For the algebra, evaluate $[Q_a^{(\lambda_{\mathrm{UV}},\ell)}, J_b^\mu]$ using the same identity with $\mathcal{O} = J_b^\mu$ and integrate; non-anomalous current algebra implies $[Q_a^{(\lambda_{\mathrm{UV}},\ell)}, Q_b^{(\lambda_{\mathrm{UV}},\ell)}] = i f_{ab}{}^c Q_c^{(\lambda_{\mathrm{UV}},\ell)}$ with no Schwinger terms. Commutation with the band projector and IR window follows from (7.180). Quartet/positivity ensures BRST-exact pieces do not contribute on $\mathcal{H}_{\mathrm{phys}}$ [442, 117]. □

Non-renormalization of conserved currents (anomaly-free case). With anomaly filters satisfied (Section 7.2.8), the conserved currents have no anomalous dimension: renormalization can at most add BRST-exact or total-derivative improvements, leaving (7.181)–(7.184) intact. In particular,

$$\mu\frac{d}{d\mu}\,J_a^\mu \sim \partial_\nu(\cdots)+\zeta(\cdots), \qquad \Rightarrow \quad \gamma_{J_a}=0 \ \text{ in } H^0(\mathcal{B}_\Gamma), \tag{7.188}$$

see [91] and [117]. Any nonzero divergence or Schwinger term would indicate an anomaly and thus violate the acceptance criteria of Section 7.3.9.

Proposition 7.257 (Algebraic non-renormalization of J_a^μ)**.** *Assume $\mathcal{S}(\Gamma)=0$ (band Slavnov identity) and exact global Ward identities on the band (non-anomalous G). Then one can choose renormalisation constants so that $Z_J=1$ and the RG flow of J_a^μ is, at most, an improvement by $\partial_\nu(\cdots)$ and a BRST-exact term $\zeta(\cdots)$. Equivalently, $\gamma_{J_a}=0$ in $H^0(\mathcal{B}_\Gamma)$.*

Proof. All statements are *on the band*; $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ commute with BRST and with G and carry ghost number 0, so band/IR operations preserve the identities used below (cf. (7.145)). We work in the algebraic–renormalisation framework (Quantum Action Principle, QAP) with the band Slavnov identity $\mathcal{S}(\Gamma)=0$ and exact global Ward identities (non–anomalous G) [117, 91].

(1) Classification of allowed counterterms for the current insertion. Consider the insertion of the local operator J_a^μ in 1PI functionals. By the QAP, the most general renormalisation of a local insertion with the quantum numbers of J_a^μ is a linear combination of operators with the same dimension, Lorentz and G indices, modulo (i) total derivatives, (ii) equations of motion (EOM) insertions, and (iii) $\mathcal{B}_\Gamma$-exact terms (linearised BRST) [117]. Current conservation and the global Ward identities constrain the counterterm space to

$$[J_a^\mu]_R = Z_J\,J_a^\mu \;+\; \partial_\nu\,\mathfrak{I}_a^{[\nu\mu]} \;+\; \mathcal{B}_\Gamma R_a^\mu \;+\; (\mathrm{EOM})_a^\mu,$$

where $\mathfrak{I}_a^{[\nu\mu]}$ is an antisymmetric local tensor (improvement), R_a^μ is a local functional (ghost number -1), and $(\mathrm{EOM})_a^\mu$ denotes terms proportional to $\delta\Gamma/\delta\varphi$ (vanishing in physical matrix elements). Ghost number and G-covariance fix the form of the $\mathcal{B}_\Gamma$ term. No other independent local structure is compatible with conservation and the identities.

(2) Ward identity fixes the normalisation $Z_J=1$. Insert $[J_a^\mu]_R$ into the *renormalised* local Ward identity. In position space,

$$\partial_\mu^{(x)}\langle [J_a^\mu]_R(x)\,\mathcal{O}_1\cdots\mathcal{O}_n\rangle = -i\sum_j \delta^{(d)}(x-x_j)\,\langle \delta_a\mathcal{O}_j\prod_{\ell\neq j}\mathcal{O}_\ell\rangle,$$

with $\{\mathcal{O}_j\}$ BRST-closed, gauge-invariant operator insertions. Taking the divergence of the r.h.s. fixes the coefficient of the contact term uniquely in terms of the *unrenormalised* G-action δ_a (the representation matrices T_a are fixed by the Lie algebra and normalisation of the fields). On the l.h.s., $\partial_\mu(\partial_\nu\mathfrak{I}_a^{[\nu\mu]})=0$, while $\partial_\mu(\mathcal{B}_\Gamma R_a^\mu)$ and $\partial_\mu(\mathrm{EOM})_a^\mu$ give only BRST-exact or EOM contact terms that cancel against source variations and vanish in physical matrix elements.

Therefore the only term that can match the canonical contact terms is $Z_J\,\partial_\mu J_a^\mu$. Since the canonical Ward identity holds *exactly* (anomaly-free), consistency forces

$$Z_J = 1.$$

Any $Z_J \neq 1$ would either rescale the global generators on the r.h.s. (forbidden) or introduce spurious contact terms, violating the Ward identity [91].

(3) Charge algebra excludes multiplicative renormalisation. Define the renormalised charge $Q_{a,R}^{(\lambda_{\rm UV},\ell)} = \int_{\Sigma_t} d\Sigma_\mu\,[J_a^\mu]_R$. Using the equal-time Ward identity one has $[Q_{a,R}^{(\lambda_{\rm UV},\ell)},\,\mathcal{O}] = \delta_a\mathcal{O}$ on BRST-closed $\mathcal{O}$. If $Z_J \neq 1$, then $Q_{a,R}^{(\lambda_{\rm UV},\ell)} = Z_J Q_a^{(\lambda_{\rm UV},\ell)} + \cdots$ and the Lie algebra becomes $[Q_{a,R}, Q_{b,R}] = i f_{ab}{}^c\, Z_J Q_c + \cdots$, or generates a Schwinger term from the dots; both contradict the exact current algebra in the anomaly-free case. Hence $Z_J = 1$ is also required by the charge algebra on the band [82, 91].

(4) RG equation and vanishing anomalous dimension in cohomology. Let $\mathcal{C}$ be the Callan–Symanzik operator acting on renormalised 1PI functionals with an insertion of $[J_a^\mu]_R$. Using $Z_J = 1$ and the QAP,

$$\mathcal{C}\,[J_a^\mu]_R = \mu\frac{d}{d\mu}\,[J_a^\mu]_R = \partial_\nu\,\widetilde{\mathfrak{J}}_a^{[\nu\mu]} \;+\; \mathcal{B}_\Gamma \widetilde{R}_a^\mu \;+\; (\mathrm{EOM})_a^\mu,$$

i.e. the μ-variation of the insertion is, at most, an improvement plus a $\mathcal{B}_\Gamma$-exact and an EOM term (no multiplicative piece). Projecting onto the physical cohomology $H^0(\mathcal{B}_\Gamma)$ and discarding EOM terms (vanishing on $\mathcal{H}_{\rm phys}$) gives

$$\gamma_{J_a}\,[J_a^\mu]_R = 0 \quad \text{in } H^0(\mathcal{B}_\Gamma),$$

hence $\gamma_{J_a} = 0$. This is the precise sense in which the current is non-renormalised: its cohomology class is RG-invariant [117].

(5) Band/IR robustness. Because $P_x(\lambda_{\rm UV})$ and $K_{\rm IR}^{(\ell)}$ commute with BRST and with G, every step above (Ward identity, charge algebra, QAP, cohomology projection) is unchanged by band/IR embedding; surface terms are controlled by the IR window. Therefore the conclusion holds *on the band*.

Combining (1)–(5) proves the proposition. □

Implications for visibilities and plateaus. Because $P_x(\lambda_{\rm UV})$ and $K_{\rm IR}^{(\ell)}$ commute with Q_a and are BRST-closed eq. (7.30), the Ward identities (7.181)–(7.184) hold *on the band* and descend to $\mathcal{H}_{\rm phys}$. Consequently, current insertions are transverse, spectral densities are positive, and channel visibilities built from G-invariant composites inherit gauge/BRST robustness and regulator stability (cf. Sections 7.3.6 and 7.3.7). These exact identities form the baseline against which any symmetry-breaking signals in other quadrants will be contrasted [82, 91, 442, 117].

7.5.9 Slavnov–Taylor Identity and its Linearisation

Definition 7.258 (Slavnov functional and master equation (band-limited))**.** Let Γ be the band-limited 1PI effective action with sources (K_Φ, K_Θ, K_c) for the BRST variations of (Φ, Θ, c), and with antighost/auxiliary pair $(\bar{c}, b)$. Holding the UV/IR knobs fixed (they commute with BRST, cf. (7.145)), the *Slavnov functional* acts on Γ as

$$\boxed{\mathcal{S}(\Gamma) = \int d^dx \left[\frac{\delta\Gamma}{\delta\Phi}\frac{\delta\Gamma}{\delta K_\Phi} + \frac{\delta\Gamma}{\delta\Theta}\frac{\delta\Gamma}{\delta K_\Theta} + \frac{\delta\Gamma}{\delta c}\frac{\delta\Gamma}{\delta K_c} + b\,\frac{\delta\Gamma}{\delta\bar{c}} \right] = 0,} \tag{7.189}$$

which is the Zinn–Justin (Slavnov–Taylor) master equation on the band [117, 442]. The integrand has ghost number $+1$ and the functional $\mathcal{S}(\Gamma)$ vanishes identically in the anomaly-free case.

Definition 7.259 (Linearised Slavnov operator)**.** Around a solution Γ of (7.189), the *linearised* ST operator $\mathcal{B}_\Gamma$ is the graded first-order differential

$$\mathcal{B}_\Gamma = \int d^dx \left[\frac{\delta\Gamma}{\delta\Phi}\frac{\delta}{\delta K_\Phi} + \frac{\delta\Gamma}{\delta K_\Phi}\frac{\delta}{\delta\Phi} + \frac{\delta\Gamma}{\delta\Theta}\frac{\delta}{\delta K_\Theta} + \frac{\delta\Gamma}{\delta K_\Theta}\frac{\delta}{\delta\Theta} + \frac{\delta\Gamma}{\delta c}\frac{\delta}{\delta K_c} + \frac{\delta\Gamma}{\delta K_c}\frac{\delta}{\delta c} + b\,\frac{\delta}{\delta\bar{c}} \right], \tag{7.190}$$

acting on local functionals with well-defined ghost number and statistics [117].

Proposition 7.260 (Antibracket form and nilpotency)**.** *Define the BV antibracket (X, Y) in the $(\Phi, \Theta, c;\, K_\Phi, K_\Theta, K_c)$ sector (band-limited) so that*

$$\mathcal{S}(\Gamma) = \tfrac{1}{2}\,(\Gamma, \Gamma), \qquad \mathcal{B}_\Gamma X = (\Gamma, X).$$

If $\mathcal{S}(\Gamma) = 0$, then $\mathcal{B}_\Gamma^2 = 0$ and $\mathcal{B}_\Gamma$ is a graded derivation of ghost number $+1$:

$$\mathcal{B}_\Gamma^2 X = \tfrac{1}{2}\big((\Gamma, \Gamma), X\big) = 0, \qquad \mathcal{B}_\Gamma(XY) = (\mathcal{B}_\Gamma X)Y + (-1)^{|X|} X(\mathcal{B}_\Gamma Y).$$

Proof. The identities follow from the graded Jacobi identity for the antibracket and $\mathcal{S}(\Gamma) = \frac{1}{2}(\Gamma, \Gamma) = 0$; see [442, 117]. Band/IR knobs commute with the functional derivatives and carry ghost number 0, so they do not affect the algebra. □

Definition 7.261 (Physical cohomology)**.** The space of *physical* (gauge-invariant) composite operators on the band is the ghost-number zero cohomology

$$H^0(\mathcal{B}_\Gamma) := \frac{\ker(\mathcal{B}_\Gamma : \mathcal{F}^0 \to \mathcal{F}^1)}{\mathrm{im}(\mathcal{B}_\Gamma : \mathcal{F}^{-1} \to \mathcal{F}^0)},$$

where $\mathcal{F}^g$ denotes local functionals of ghost number g [117]. Representatives differing by $\mathcal{B}_\Gamma$-exact terms define the same physical matrix elements on $\mathcal{H}_{\mathrm{phys}}$ (quartet decoupling).

Proposition 7.262 (Quantum Action Principle (QAP) and stability)**.** *Suppose $\mathcal{S}(\Gamma) = 0$ (cf. eq. (7.16)) up to order $n-1$ in loops. At order n, any breaking $\Delta^{(n)}$ of ghost number $+1$ fulfils the (linearised) Wess–Zumino consistency condition*

$$\mathcal{B}_{\Gamma^{(0)}}\,\Delta^{(n)} = 0.$$

If the cohomology $H^1(\mathcal{B}_{\Gamma^{(0)}})$ is trivial in the relevant sector (anomaly filters; see Section 7.2.8), there exists a local counterterm $\mathsf{Z}^{(n)}$ with $\mathfrak{g}(\mathsf{Z}^{(n)}) = 0$ such that

$$\Delta^{(n)} = \mathcal{B}_{\Gamma^{(0)}}\, \mathsf{Z}^{(n)}.$$

Adding $-\hbar^n \mathsf{Z}^{(n)}$ (the Wess–Zumino counterterm at loop n, cf. Definition 7.25) restores $\mathcal{S}(\Gamma) = 0$ at order n. Hence the identity is stable *under renormalisation on the band [117].*

Proof. QAP locality and linearised consistency. By the Quantum Action Principle, any breaking at fixed loop order is a *local* insertion with the appropriate quantum numbers. Expanding $\Gamma = \Gamma^{(0)} + \sum_{k\geq 1} \hbar^k \Gamma^{(k)}$ and the Slavnov functional $\mathcal{S}(\Gamma)$ (cf. eq. (7.16)) gives, at order n,

$$\mathcal{S}(\Gamma) = \hbar^n \Delta^{(n)} + \mathcal{O}(\hbar^{n+1}), \qquad \Delta^{(n)} = \mathcal{B}_{\Gamma^{(0)}}\, \Gamma^{(n)} + (\text{known lower-order terms}).$$

Nilpotency of the linearised Slavnov operator $\mathcal{B}_{\Gamma^{(0)}}$ implies $\mathcal{B}_{\Gamma^{(0)}}\, \Delta^{(n)} = 0$, i.e. the linearised Wess–Zumino consistency condition.

Cohomological triviality via anomaly filters. If $H^1(\mathcal{B}_{\Gamma^{(0)}})$ is trivial in the relevant sector (anomaly filters of Section 7.2.8), then the cocycle $\Delta^{(n)}$ is a coboundary: there exists a *local* $\mathsf{Z}^{(n)}$ with $\mathfrak{g}(\mathsf{Z}^{(n)}) = 0$ such that $\Delta^{(n)} = \mathcal{B}_{\Gamma^{(0)}}\, \mathsf{Z}^{(n)}$. In the descent language this matches Definition 7.25, where the order–n anomalous variation is $\zeta\, \mathsf{Z}^{(n)}$ up to a total derivative.

Counterterm restores the identity at order n. Define the renormalised action

$$\Gamma' \;:=\; \Gamma \;-\; \hbar^n \mathsf{Z}^{(n)}.$$

Expanding $\mathcal{S}(\Gamma')$ to order n and using the linearity of the first variation yields

$$\mathcal{S}(\Gamma') \;=\; \mathcal{S}(\Gamma) - \hbar^n\, \mathcal{B}_{\Gamma^{(0)}}\, \mathsf{Z}^{(n)} + \mathcal{O}(\hbar^{n+1}) \;=\; \hbar^n \Delta^{(n)} - \hbar^n \Delta^{(n)} + \mathcal{O}(\hbar^{n+1}) \;=\; \mathcal{O}(\hbar^{n+1}),$$

so the Slavnov identity is restored at order n (cf. eq. (7.16)).

Band/IR compatibility. Since $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ commute with BRST and preserve ghost number (band setting), locality and the above cohomological argument are unchanged on the band (see Section 7.2.8). Thus stability holds in the band framework.

This proves the proposition. □

Proposition 7.263 (Nielsen identity as a $\mathcal{B}_\Gamma$-exact variation)**.** *If the gauge parameter α_{gf} enters only through the gauge fermion $\Psi(\alpha_{\mathrm{gf}})$, then the α_{gf}-variation of Γ is $\mathcal{B}_\Gamma$-exact on the band,*

$$\partial_{\alpha_{\mathrm{gf}}} \Gamma = -\,\mathcal{B}_\Gamma\Big(\partial_{\alpha_{\mathrm{gf}}} \Psi\Big),$$

so that on-shell, BRST-closed observables (masses, residues, certified visibilities) are gauge-parameter independent [117, 91].

Proof. Differentiate the path integral with respect to α_{gf}; only $\zeta\Psi$ varies. Using the antibracket representation of BRST in the source sector and performing the Legendre transform yields the stated identity; cf. the band versions in Section 7.5.5. □

Corollary 7.264 (Operational consequences)**.** *Assuming $\mathcal{S}(\Gamma) = 0$ and anomaly filters:*

(i) Unphysical modes decouple cohomologically on $\mathcal{H}_{\mathrm{phys}}$ (quartet mechanism), compatibly with Section 7.2.6.

(ii) On-shell poles and holosymmetric visibilities are gauge-parameter independent (Nielsen) [117, 91].

(iii) Renormalisation constants and counterterms satisfy algebraic constraints fixed by $\mathcal{S}(\Gamma)$ (e.g. current non-renormalisation in cohomology), see [117, 91, 82].

All statements hold on the band *since $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ commute with BRST and with functional differentiation* (7.145).

Quantum Action Principle, anomalies, and consistency (formal). All statements below are *on the band*; the UV/IR knobs commute with BRST and carry ghost number 0, cf. (7.145). We use the antibracket form $\mathcal{S}(\Gamma) = \frac{1}{2}(\Gamma, \Gamma)$ and the linearised operator $\mathcal{B}_\Gamma X = (\Gamma, X)$.

Proposition 7.265 (QAP breaking and Wess–Zumino consistency on the band)**.** *Let $\Gamma = \sum_{\ell \geq 0} \hbar^\ell \Gamma^{(\ell)}$ with $\mathcal{S}(\Gamma^{(0)}) = 0$. At loop order $n \geq 1$, the Quantum Action Principle implies a* local *breaking $\Delta^{(n)}$ of ghost number $+1$ such that*

$$\mathcal{S}(\Gamma) = \hbar^n \, \Delta^{(n)} + \mathcal{O}(\hbar^{n+1}), \qquad \mathcal{B}_{\Gamma^{(0)}} \, \Delta^{(n)} = 0, \tag{7.191}$$

i.e. $\Delta^{(n)}$ is a BRST cocycle (Wess–Zumino consistency) [117, 442].

Proof. Expand $\mathcal{S}(\Gamma) = \frac{1}{2}(\Gamma, \Gamma)$ in powers of $\hbar$. The coefficient at order $\hbar^n$ reads

$$\sum_{k=0}^{n} (\Gamma^{(k)}, \Gamma^{(n-k)}) = 2\,(\Gamma^{(0)}, \Gamma^{(n)}) + \sum_{k=1}^{n-1} (\Gamma^{(k)}, \Gamma^{(n-k)}).$$

Define $\Delta^{(n)} := \frac{1}{2} \sum_{k=1}^{n-1} (\Gamma^{(k)}, \Gamma^{(n-k)})$, which is local by the QAP. Then $(\Gamma^{(0)}, \Gamma^{(n)}) = \Delta^{(n)}$, i.e. $\mathcal{B}_{\Gamma^{(0)}} \Gamma^{(n)} = \Delta^{(n)}$. Apply $\mathcal{B}_{\Gamma^{(0)}}$ and use the graded Jacobi identity with $(\Gamma^{(0)}, \Gamma^{(0)}) = 0$ to obtain $\mathcal{B}_{\Gamma^{(0)}} \Delta^{(n)} = 0$. Band/IR operations commute with the functional derivatives entering the antibracket and preserve locality, hence the same derivation holds on the band [117]. □

Proposition 7.266 (All-order restoration under anomaly filters)**.** *If the anomaly filters hold so that $H^1(\mathcal{B}_{\Gamma^{(0)}}) = 0$ in the relevant sector, then the breaking $\Delta^{(n)}$ is $\mathcal{B}_{\Gamma^{(0)}}$-exact:*

$$\Delta^{(n)} = \mathcal{B}_{\Gamma^{(0)}} \, \widehat{\Sigma}^{(n)} \quad \Rightarrow \quad \Gamma \;\mapsto\; \Gamma - \hbar^n \widehat{\Sigma}^{(n)} \;\; \textit{restores} \;\; \mathcal{S}(\Gamma) = 0 \;\textit{at order } n. \tag{7.192}$$

By induction, $\mathcal{S}(\Gamma) = 0$ can be imposed to all orders (algebraic renormalisation) [117, 442].

Proof. Triviality of H^1 implies existence of a local $\widehat{\Sigma}^{(n)}$ with the stated property. Adding $-\hbar^n \widehat{\Sigma}^{(n)}$ shifts $\Gamma^{(n)}$ by a solution of the inhomogeneous equation $\mathcal{B}_{\Gamma^{(0)}} \Gamma^{(n)} = \Delta^{(n)}$, cancelling the breaking at order n. Proceed by induction in n; the QAP ensures locality at each step. □

Cohomological classification of counterterms (formal).

Proposition 7.267 (Counterterms as BRST cohomology)**.** *Let Σ_{ct} be a local counterterm of ghost number 0. Requiring the renormalised action $\Gamma' = \Gamma + \epsilon\,\Sigma_{\mathrm{ct}}$ to satisfy the ST identity to first order in ϵ yields*

$$\mathcal{B}_\Gamma\,\Sigma_{\mathrm{ct}} = 0, \qquad \Sigma_{\mathrm{ct}} \sim \Sigma_{\mathrm{ct}} + \mathcal{B}_\Gamma X, \tag{7.193}$$

so admissible counterterms are classes in $H^0(\mathcal{B}_\Gamma)$, modulo BRST-exact redefinitions [117]. On the band, this cohomology is generated by the G-invariant kinetic pairings and interaction monomials compatible with eqs. (6.333) and (6.334).

Proof. Insert Γ' into $\mathcal{S}(\cdot)$ and expand to $\mathcal{O}(\epsilon)$:

$$0 = \mathcal{S}(\Gamma') = \mathcal{S}(\Gamma) + \epsilon\,\mathcal{B}_\Gamma\Sigma_{\mathrm{ct}} + \mathcal{O}(\epsilon^2) = \epsilon\,\mathcal{B}_\Gamma\Sigma_{\mathrm{ct}} + \cdots,$$

since $\mathcal{S}(\Gamma) = 0$. Thus $\mathcal{B}_\Gamma\Sigma_{\mathrm{ct}} = 0$. If $\Sigma_{\mathrm{ct}} \to \Sigma_{\mathrm{ct}} + \mathcal{B}_\Gamma X$, then $\mathcal{B}_\Gamma\Sigma_{\mathrm{ct}}$ is unchanged (nilpotency), and the change corresponds to a BRST-exact field redefinition that leaves physical matrix elements invariant. Band/IR commutation ensures the same algebraic steps apply. □

Remark 7.268 (Practical reduction of renormalisations)**.** The classification (7.193) reduces independent renormalisations to a small basis of cohomology representatives (invariant kinetic terms, gauge-invariant potentials, etc.), while gauge-fixing and BRST-exact pieces are in the trivial class. This underlies the stability results and the non-renormalisation statements used throughout (e.g. current protection, Nielsen gauge-parameter independence) [117, 91].

Vertex-level consequences and Z-factor relations. Differentiating the master equation (7.189) with respect to appropriate fields and sources, then setting sources to zero at the G-symmetric vacuum, generates the tower of Slavnov–Taylor relations among amputated 1PI kernels. In compact notation,

$$\mathcal{B}_\Gamma\,\Gamma^{(3)} \;=\; \Gamma^{(2)} \star \Gamma^{(3)} \;+\; \cdots, \tag{7.194}$$

where $\star$ denotes the standard index contraction and convolution with propagators and group factors, and the ellipsis collects additional terms from field mixing and source insertions fixed by (7.189) (see [117, 91, 82]).

Proposition 7.269 (Slavnov–Taylor constraints on renormalisation constants)**.** *Assume a multiplicatively renormalisable scheme with field renormalisations $\varphi_i^B = Z_i^{1/2}\,\varphi_i$, two-point kernels $\Gamma_i^{(2)B} = Z_{\Gamma_i^{(2)}}\,\Gamma_i^{(2)}$, and an m-leg 1PI vertex $\Gamma^{(m)B} = Z_{\Gamma^{(m)}}\,\Gamma^{(m)}$. Let Z_{int} denote the renormalisation of the corresponding (gauge-/BRST-invariant) interaction monomial in the action. Then the differentiated ST identities implied by (7.194) yield algebraic relations of the form*

$$Z_{\mathrm{int}} \;=\; \frac{Z_{\Gamma^{(3)}}}{\prod_i Z_{\Gamma_i^{(2)}}^{\alpha_i}}, \qquad \sum_i \alpha_i = \tfrac{1}{2}\,m, \tag{7.195}$$

with exponents α_i fixed by leg counting and field normalisations. In particular, for a cubic vertex built from fields $(\varphi_A, \varphi_B, \varphi_C)$ one recovers the familiar structure $Z_{\Gamma^{(3)}} = Z_{\mathrm{int}}\,Z_A^{1/2}Z_B^{1/2}Z_C^{1/2}$, and analogously for higher-point kernels once reduced by ST identities [117, 91, 82].

Sketch. Apply functional derivatives $\delta/\delta\varphi_{i_1}\cdots\delta/\delta\varphi_{i_m}$ to (7.189) and set sources to zero at the symmetric vacuum. The resulting identities express $\Gamma^{(m)}$ in terms of $\Gamma^{(2)}$ and lower vertices with fixed group-theoretic tensors. Passing to bare versus renormalised quantities and using multiplicativity, the identities can only be preserved if the overall Z-factors satisfy (7.195). The sum rule $\sum_i \alpha_i = m/2$ follows from canonical dimension/leg counting of amputated kernels. For explicit gauges (e.g. Yang–Mills), this reproduces standard relations such as $Z_g = Z_1\, Z_3^{-3/2}$, $Z_{\bar{1}} = Z_g\, Z_3^{1/2} Z_c$ for ghost–gluon, etc. [82, 91]; the abstract cohomological derivation ensures the same algebraic pattern in our setting [117]. □

Band/IR compatibility and descent to $\mathcal{H}_{\text{phys}}$.

Proposition 7.270 (Band/IR compatibility of ST identities and cohomology)**.** *If $P_x(\lambda_{\text{UV}})$ and $K_{\text{IR}}^{(\ell)}$ commute with $\hat{Q}_{\text{BRST}}$ and have ghost number* 0 (7.145)*, then the master equation* (7.189)*, its linearisation* (7.190)*, and all vertex-level consequences (including* (7.194)*–*(7.195)*) hold verbatim* on the band*. Moreover, by quartet decoupling, physical matrix elements depend only on the classes in $H^0(\mathcal{B}_\Gamma)$ on $\mathcal{H}_{\text{phys}}$ (cf. Section 7.2.6 and [117]).*

Proof. Band/IR operations commute with functional differentiation and preserve locality and ghost number. Therefore they can be pulled through the derivation of (7.189) and of its differentiated consequences; cohomology statements are unchanged because $\mathcal{B}_\Gamma$ is unaffected by multiplication with band/IR projectors of ghost number 0. □

Remark (Nielsen identity as an ST corollary). Promoting the gauge parameter α_{gf} to a BRST doublet and differentiating (7.189) yields the Nielsen master identity, i.e. $\partial_{\alpha_{\text{gf}}}\Gamma = -\,\mathcal{B}_\Gamma(\partial_{\alpha_{\text{gf}}}\Psi)$, hence on-shell, BRST-closed observables (masses, residues, certified visibilities) are α_{gf}-independent [117, 91]. In our band-limited setting, the same conclusion holds by (7.145).

7.5.10 BRST Charge, Nilpotency, and Equivalence Theorem

Definition 7.271 (BRST charge and physical Hilbert space)**.** Let $\hat{Q}_{\text{BRST}}$ be the conserved, Hermitian BRST charge on the (indefinite-metric) state space such that

$$\hat{Q}_{\text{BRST}}^2 = 0, \qquad [\hat{Q}_{\text{BRST}}, \hat{H}_0(t)] = 0, \qquad \hat{Q}_{\text{BRST}}^\dagger = \hat{Q}_{\text{BRST}}, \tag{7.196}$$

with $\hat{H}_0(t) = \int d^d x\, \hat{\mathcal{H}}_0(t,x)$ the symmetric-law Hamiltonian. The *physical* Hilbert space is the ghost-number 0 cohomology

$$\mathcal{H}_{\text{phys}} \;\cong\; \ker \hat{Q}_{\text{BRST}}/\text{im}\, \hat{Q}_{\text{BRST}}, \qquad \mathsf{P}_{\text{phys}} :\ \text{orthogonal projector onto } \ker \hat{Q}_{\text{BRST}}.$$

On $\mathcal{H}_{\text{phys}}$, positivity and unitarity are recovered by the quartet mechanism (decoupling of BRST doublets) and the Källén–Lehmann representation with positive spectral measure [442, 117, 91, 241].

Proposition 7.272 (Conservation and cohomology-preserving evolution)**.** *Let $U_0(t_2,t_1)$ be the unitary propagator generated by $\hat{H}_0(t)$. If $[\hat{Q}_{\mathrm{BRST}},\hat{H}_0(t)]=0$ for all t, then*

$$[\hat{Q}_{\mathrm{BRST}},U_0(t_2,t_1)]=0,$$
$$U_0(t_2,t_1):\ \ker\hat{Q}_{\mathrm{BRST}}\to\ker\hat{Q}_{\mathrm{BRST}},$$
$$U_0(t_2,t_1):\ \operatorname{im}\hat{Q}_{\mathrm{BRST}}\to\operatorname{im}\hat{Q}_{\mathrm{BRST}}.$$

Hence time evolution preserves BRST cohomology classes, and U_0 descends unitarily to $\mathcal{H}_{\mathrm{phys}}$.

Proof. Differentiate $U_0^\dagger Q U_0$ with respect to time and use $i\partial_t U_0=\hat{H}_0U_0$, then $d(U_0^\dagger QU_0)/dt=U_0^\dagger i[\hat{H}_0,Q]U_0=0$. The two stability statements follow from $QU_0=U_0Q$. □

Proposition 7.273 (Decoupling of BRST-exact insertions)**.** *Let $|\psi\rangle,|\phi\rangle\in\ker\hat{Q}_{\mathrm{BRST}}$ be physical states and $\hat{X}$ any (graded) operator. Then*

$$\langle\psi|\ \{\hat{Q}_{\mathrm{BRST}},\hat{X}\}\ |\phi\rangle=0,$$

so BRST-exact insertions vanish between physical states. In particular, S-matrix elements between physical asymptotic states are invariant under $\hat{\mathcal{H}}_0\mapsto\hat{\mathcal{H}}_0+\{\hat{Q}_{\mathrm{BRST}},\hat{Y}\}$ (cohomological robustness).

Proof. Use $(\hat{Q}_{\mathrm{BRST}})^\dagger=\hat{Q}_{\mathrm{BRST}}$ and $\hat{Q}_{\mathrm{BRST}}|\psi\rangle=\hat{Q}_{\mathrm{BRST}}|\phi\rangle=0$ to write $\langle\psi|\{\hat{Q},\hat{X}\}|\phi\rangle=\langle\psi|\hat{Q}\hat{X}|\phi\rangle+\langle\psi|\hat{X}\hat{Q}|\phi\rangle=0$. For the Hamiltonian deformation, Dyson-expand the time-ordered exponential; each term brings one insertion of a BRST anticommutator, which vanishes by the same argument. □

Definition 7.274 (Band/IR compatibility of the BRST charge)**.** Let $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ be the UV projector and IR window. Assume they are BRST-closed and ghost-number 0,

$$[\hat{Q}_{\mathrm{BRST}},P_x(\lambda_{\mathrm{UV}})]=0,\qquad\zeta\Big(K_{\mathrm{IR}}^{(\ell)}*\,\cdot\,\Big)=0,$$

as recalled in (7.145). Then the *band-limited* charge $Q^{(\lambda_{\mathrm{UV}},\ell)}:=P_xQP_x$ acts on the same physical cohomology and commutes with the band-limited symmetric evolution generated by $H_0^{(\lambda_{\mathrm{UV}},\ell)}$.

Proposition 7.275 (Equivalence theorem: BRST-exact deformations and field redefinitions)**.** *Consider the band-limited 1PI action Γ satisfying the Slavnov identity $\mathcal{S}(\Gamma)=0$ (7.189).*

*(i) (*BRST-exact deformations*) If $\Gamma\mapsto\Gamma'=\Gamma+\epsilon\,\mathcal{B}_\Gamma X$ with a local X (ghost number -1), then on-shell S-matrix elements of BRST-closed, gauge-invariant operators are unchanged.*

*(ii) (*Local field redefinitions*) If $\varphi\mapsto\varphi'=\varphi+\epsilon\,F(\varphi)$ is a local, invertible redefinition that preserves the asymptotic LSZ structure, then physical S-matrix elements are invariant. If $F=\mathcal{B}_\Gamma Y$ is BRST-exact, the statement follows from (i).*

Proof. (i) Using the antibracket form of the Slavnov identity (linearised BRST) one has $\delta\Gamma = \epsilon\,\mathcal{B}_\Gamma X$. Differentiating Green's functions with respect to ϵ inserts $\mathcal{B}_\Gamma X$, i.e. BRST-exact operators. After LSZ amputation and restricting to external BRST-closed asymptotic states, these insertions vanish by Proposition 7.273 (or, equivalently, by the Nielsen/Slavnov argument that $\mathcal{B}_\Gamma$-exact variations are EOM-proportional on shell) [117, 442, 91].
(ii) The standard equivalence theorem applies: local invertible redefinitions change only contact terms and wavefunction factors that cancel in properly normalised S-matrix elements; see [91]. If the redefinition is of BRST-exact type, it reduces to (i). □

Corollary 7.276 (Gauge-parameter/Nielsen invariance as a special case)**.** *If the gauge parameter α_{gf} enters only via the gauge fermion $\Psi(\alpha_{\mathrm{gf}})$, then*

$$\partial_{\alpha_{\mathrm{gf}}}\Gamma = -\,\mathcal{B}_\Gamma\big(\partial_{\alpha_{\mathrm{gf}}}\Psi\big),$$

so on-shell physical poles, residues, and certified visibilities are independent of α_{gf} (Nielsen identity on the band; cf. Section 7.5.5) [117, 91].

Remark 7.277 (Implications for the holosymmetric probe)**.** Because all ingredients of the probe (band-limited preparations, sector projectors, and controlled evolutions) commute with $\hat{Q}_{\mathrm{BRST}}$ and are BRST-closed on $\mathcal{H}_{\mathrm{phys}}$, Propositions 7.272–7.275 ensure that the channel visibilities are (i) insensitive to BRST-exact branches in the law channel, (ii) invariant under gauge-parameter changes (Nielsen), and (iii) stable under admissible regulator variations that preserve the cohomology class, as used throughout Section 7.4.

Equivalence theorem (BRST-exact deformations).

Theorem 7.278 (Equivalence theorem for BRST-exact deformations (band-limited))**.** *Consider the one-parameter family*

$$S_s[\varphi] = S[\varphi] + s\,\zeta\mathcal{X}[\varphi], \qquad s \in \mathbb{R},$$

with $\mathcal{X}$ local and band-limited, and let (Z_s, W_s, Γ_s) be the corresponding generating and 1PI functionals constructed with fixed UV/IR knobs (which commute with BRST and have ghost number 0, cf. (7.145)). Assume: (i) $\mathcal{S}(\Gamma_s) = 0$ (anomaly filters), (ii) the QAP applies, and (iii) LSZ asymptotics exist for the physical sector. Then there exists a local ghost-number -1 functional $\widehat{\mathcal{X}}_s$ such that

$$\frac{d}{ds}\,\Gamma_s = \mathcal{B}_{\Gamma_s}\,\widehat{\mathcal{X}}_s, \tag{7.197}$$

i.e. the s-variation of Γ_s is BRST-exact in the linearised cohomology. Consequently, for any set of physical insertions $\{\mathcal{O}_i\} \subset H^0(\mathcal{B}_{\Gamma_s})$,

$$\frac{d}{ds}\,\mathcal{M}_s(\mathcal{O}_1, \ldots, \mathcal{O}_n) = 0, \qquad \Rightarrow \quad \mathcal{S} \text{ and all } \mathcal{H}_{\mathrm{phys}} \text{ matrix elements are invariant.} \tag{7.198}$$

All statements hold on the band.

Proof. Differentiate Z_s with respect to s:

$$\frac{d}{ds}Z_s = \int \mathcal{D}\varphi\, (i\,\zeta\mathcal{X})\, e^{iS_s}, \qquad \frac{d}{ds}W_s = \langle \zeta\mathcal{X}\rangle_s.$$

Introduce band-limited sources $(\mathcal{J},\mathcal{K})$ for fields and their BRST variations (cf. Section 7.5.1) and perform the Legendre transform at fixed $\mathcal{K}$. Using the Slavnov master equation in antibracket/linearised form and the QAP, one obtains the standard algebraic–renormalisation result

$$\frac{d}{ds}\,\Gamma_s = -\,\mathcal{B}_{\Gamma_s}\Big[\partial_\epsilon \widehat{\mathcal{X}}_{s,\epsilon}\Big]_{\epsilon=0} \equiv \mathcal{B}_{\Gamma_s}\,\widehat{\mathcal{X}}_s,$$

with $\widehat{\mathcal{X}}_s$ local of ghost number -1; see [117, 91, 442]. Amputating external legs and applying on-shell LSZ to correlators with $\{\mathcal{O}_i\} \subset H^0(\mathcal{B}_{\Gamma_s})$ inserts a $\mathcal{B}_{\Gamma_s}$-exact operator, which vanishes between BRST-closed asymptotic states by cohomological decoupling (quartet mechanism) or, equivalently, because $\mathcal{B}_{\Gamma_s}$-exact variations reduce to EOM insertions on shell. Thus $d\mathcal{M}_s/ds = 0$, proving (7.198). Band/IR commutation guarantees all steps carry over unchanged on the band. □

Remark 7.279. If the anomaly filters fail so that $\mathcal{S}(\Gamma_s) \neq 0$, the QAP breaking $\Delta^{(n)}$ (ghost number $+1$) may obstruct (7.197) (cf. Proposition 7.265); in that case the deformation can affect physical amplitudes.

Operational corollaries for the holosymmetric probe.

Corollary 7.280 (Operational corollaries for the holosymmetric probe)**.** *Work on the band with BRST-closed knobs and assume the hypotheses of Theorem 7.278.*

(i) Law-branch invariance.
In Section 7.4.2 the two law branches differ by a BRST-exact deformation of the generator, $H_0 \mapsto H_0 + \zeta\Delta G$. By Theorem 7.278,

$$\frac{d}{ds}\,\Gamma_s = \mathcal{B}_{\Gamma_s}\,\widehat{\mathcal{X}}_s \quad\Longrightarrow\quad \frac{d}{ds}\,\mathcal{M}_s = 0$$

for any physical insertion; hence the branch difference cancels on $\mathcal{H}_{\mathrm{phys}}$ in the anomaly-free case, yielding the on-shell equalities at certified symmetry points

$$\mathcal{C}_{\mathrm{law}} = 1, \qquad \Lambda^{\star}_{\mathrm{law}} = 1,$$

as stated in eq. (7.121) and Section 7.4.4.

(ii) Gauge-parameter (Nielsen) independence.
Promote the gauge parameter α_{gf} to a BRST doublet and apply the same cohomological argument to the deformation $S \mapsto S + \delta\alpha_{\mathrm{gf}}\,\zeta(\partial_{\alpha_{\mathrm{gf}}}\Psi)$. Then

$$\partial_{\alpha_{\mathrm{gf}}}\Gamma = -\,\mathcal{B}_\Gamma\Big(\partial_{\alpha_{\mathrm{gf}}}\Psi\Big),$$

so on-shell, BRST-closed observables (poles, residues, and certified visibilities) are α_{gf}-independent (Nielsen identity on the band; cf. Section 7.2.9 and [117, 91]).

(iii) Anomaly sensitivity.
If the anomaly filters fail (Section 7.2.8), the loop-breaking $\Delta^{(n)}$ in (7.191) may represent a nontrivial class in $H^1(\mathcal{B}_{\Gamma^{(0)}})$ [117, 442]. Then the variation $\frac{d}{ds}\Gamma_s = \mathcal{B}_{\Gamma_s}\widehat{\mathcal{X}}_s$ need not be cohomologically trivial on shell, the conclusion of (7.198) can fail, and robust suppressions $|\mathcal{C}_{\mathrm{law}}| < 1$ become lawful diagnostics of symmetry violation (cf. Section 7.4.4).

Proof. (i) Apply Theorem 7.278 with the deformation generated by $\zeta\Delta G$; physical amplitudes are s-independent, hence the two branches are indistinguishable on $\mathcal{H}_{\mathrm{phys}}$. (ii) Set the deformation to $\zeta(\partial_{\alpha_{\mathrm{gf}}}\Psi)$ and invoke the Nielsen identity (band version), derived as a $\mathcal{B}_\Gamma$-exact variation; on shell, BRST-closed observables are invariant [117, 91]. (iii) Use Proposition 7.265: if $H^1(\mathcal{B}_{\Gamma^{(0)}}) \neq 0$, the breaking cannot be removed by a counterterm, so the BRST-exactness needed for invariance may fail; deviations in $\mathcal{C}_{\mathrm{law}}$ are then physically meaningful. □

Band/IR consistency and conservation.

Proposition 7.281 (Band/IR consistency and BRST conservation)**.** *Let $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ be the UV projector and IR window satisfying*

$$[\hat{Q}_{\mathrm{BRST}}, P_x(\lambda_{\mathrm{UV}})] = 0, \qquad \zeta\Big(K_{\mathrm{IR}}^{(\ell)} * \cdot\Big) = 0 \quad \textit{(cf. (7.145)).}$$

Let $\hat{H}_0^{(\lambda_{\mathrm{UV}},\ell)}(t)$ be the band/IR-embedded symmetric Hamiltonian (cf. Section 7.4.3). Then

$$[\hat{Q}_{\mathrm{BRST}},\, \hat{H}_0^{(\lambda_{\mathrm{UV}},\ell)}(t)] = 0 \quad\Longrightarrow\quad \frac{d}{dt}\,\hat{Q}_{\mathrm{BRST}}(t) = i\,[\hat{H}_0^{(\lambda_{\mathrm{UV}},\ell)}(t), \hat{Q}_{\mathrm{BRST}}] = 0, \tag{7.199}$$

and the corresponding unitary propagator $U_0^{(\lambda_{\mathrm{UV}},\ell)}(t_2,t_1)$ preserves BRST cohomology classes. Consequently, all algebraic statements used in the equivalence theorem (Slavnov identity, linearisation, QAP locality) commute with the band/IR embedding, and Theorem 7.278 applies verbatim *to the holosymmetric probe constructed on the band.*

Proof. By construction (band/IR embedding of Section 7.4.3) the Hamiltonian density is obtained from its unbanded form by left/right multiplication with $P_x(\lambda_{\mathrm{UV}})$ and convolution with $K_{\mathrm{IR}}^{(\ell)}$, both BRST-closed and of ghost number 0. Since $[\hat{Q}_{\mathrm{BRST}}, \hat{\mathcal{H}}_0] = 0$ for the symmetric law, BRST-closure of the knobs implies $[\hat{Q}_{\mathrm{BRST}}, \hat{H}_0^{(\lambda_{\mathrm{UV}},\ell)}] = 0$, yielding (7.199). The standard argument $d(U^\dagger Q U)/dt = U^\dagger i[\hat{H}_0^{(\lambda_{\mathrm{UV}},\ell)}, Q]U = 0$ shows $[Q, U_0^{(\lambda_{\mathrm{UV}},\ell)}] = 0$, hence stability of $\ker Q$ and $\operatorname{im} Q$ and preservation of cohomology. Because the knobs commute with BRST and functional differentiation, the derivations of the Slavnov–Taylor identity, its linearisation $\mathcal{B}_\Gamma$, and the QAP remain unchanged on the band [117, 442, 91]. Therefore Theorem 7.278 (and its operational corollaries) holds for the band-limited holosymmetric probe. □

Bibliography note. See [117] for the algebraic (BRST) proof of the equivalence theorem and [442, 91] for complementary treatments. For historical context on field-redefinition invariance, see Kamefuchi–O'Raifeartaigh–Salam (1961).

7.5.11 Nielsen Identities at the Symmetry Point

Definition 7.282 (Gauge-parameter BRST doublet and band/IR commutation)**.** Let $\alpha_{\rm gf}$ be a gauge-fixing parameter that appears only through the BRST-exact gauge-fermion term $\zeta\Psi(\alpha_{\rm gf})$. Promote it to a BRST doublet

$$\zeta\,\alpha_{\rm gf} = \chi, \qquad \zeta\,\chi = 0,$$

and recall the band/IR commutation with BRST,

$$[\hat{Q}_{\rm BRST}, P_x(\lambda_{\rm UV})] = 0, \qquad \zeta\Big(K^{(\ell)}_{\rm IR} * \,\cdot\,\Big) = 0, \tag{7.200}$$

so that UV projection and IR smearing carry ghost number 0 and commute with the BRST differential.

Theorem 7.283 (Nielsen (gauge-parameter) master identity on the band)**.** *Let Γ be the band-limited 1PI functional, defined at fixed antifield/source multiplet $\mathcal{K}$ and satisfying the Slavnov identity $\mathcal{S}(\Gamma) = 0$ (anomaly-free case). Then*

$$\boxed{\frac{\partial\Gamma}{\partial\alpha_{\rm gf}} = \int d^dx\, \frac{\delta\Gamma}{\delta\varphi^A(x)}\,\mathcal{N}^A(x;\alpha_{\rm gf}) \equiv \Big\langle \zeta\,\partial_{\alpha_{\rm gf}}\Psi\Big\rangle_\Gamma \cdot \frac{\delta\Gamma}{\delta\varphi}\,,}\tag{7.201}$$

where $\varphi = (\Phi, \Theta, c, \bar{c}, b)$, the dot denotes the band/IR-smeared pairing, and

$$\mathcal{N}^A(x;\alpha_{\rm gf}) \;:=\; -\,\frac{\delta}{\delta K_A(x)}\Big(\partial_{\alpha_{\rm gf}}\Psi\Big) \quad \textit{(linearised/antibracket form).} \tag{7.202}$$

Equivalently,

$$\frac{\partial\Gamma}{\partial\alpha_{\rm gf}} = -\,\mathcal{B}_\Gamma\Big(\partial_{\alpha_{\rm gf}}\Psi\Big), \tag{7.203}$$

with $\mathcal{B}_\Gamma$ the linearised Slavnov operator. All statements hold on the band.

Sketch. Extend the field content by the BRST doublet $(\alpha_{\rm gf}, \chi)$ and consider the band-limited generating functional with sources $(\mathcal{J}, \mathcal{K})$ as in Section 7.5.1. The master equation $\mathcal{S}(\Gamma) = 0$ (band version of eq. (7.189)) remains valid because the knobs commute with BRST and have ghost number 0. Differentiate the functional integral with respect to $\alpha_{\rm gf}$ at fixed $\mathcal{K}$; only the gauge-fermion changes, producing the insertion $\zeta(\partial_{\alpha_{\rm gf}}\Psi)$. Passing to the 1PI functional and using the Quantum Action Principle together with the antibracket form of the Slavnov identity yields (7.203) (see [117]; [91]). Expanding the linearised operator gives (7.201) with the kernel (7.202). Band/IR commutation (7.200) ensures no extra terms arise from the knobs. □

Corollary 7.284 (On-shell gauge-parameter independence)**.** *Let $\mathcal{O}$ be a BRST-closed, gauge-invariant composite in $H^0(\mathcal{B}_\Gamma)$. Then the variation (7.201) is proportional to the equations of motion $\delta\Gamma/\delta\varphi^A$ and hence vanishes on shell. In particular, physical poles, residues, and certified holosymmetric visibilities (constructed from BRST-closed inputs) are $\alpha_{\rm gf}$-independent at the symmetry point in Quadrant I [117, 91].*

Remark 7.285 (Forms of the identity)**.** Equation (7.201) is the "equation-of-motion" (EOM) form; (7.203) is the linearised/antibracket form. The expectation-value notation in (7.201) uses the 1PI insertion map induced by Γ (at fixed $\mathcal{K}$). All forms are equivalent and commute with band/IR embedding by (7.200).

Two-point kernels and pole masses — formal treatment

Definition 7.286 (Transverse (physical) two-point kernel)**.** Let $\Gamma^{(2)}(p;\alpha_{\mathrm{gf}})$ be the 1PI two-point kernel in momentum space. With the longitudinal/transverse projectors $\Pi_{\mathrm{L}},\Pi_{\mathrm{T}}$ from Section 7.3.5, define the physical (transverse) block

$$\Gamma^{(2)}_{\mathrm{TT}}(p^2;\alpha_{\mathrm{gf}}) := \Pi_{\mathrm{T}}\,\Gamma^{(2)}(p;\alpha_{\mathrm{gf}})\,\Pi_{\mathrm{T}}, \qquad \Pi_{\mathrm{L}}\,\Gamma^{(2)}\,\Pi_{\mathrm{T}} = 0 \ \text{ on } \mathcal{H}_{\mathrm{phys}}. \tag{7.204}$$

Lemma 7.287 (Projected Nielsen identity for two-point kernels)**.** *Let Γ satisfy the band Nielsen master identity (7.201). Differentiating twice with respect to fields and projecting with Π_{T} yields*

$$\partial_{\alpha_{\mathrm{gf}}}\Gamma^{(2)}_{\mathrm{TT}}(p^2) = \mathcal{C}(p^2;\alpha_{\mathrm{gf}})\,\Gamma^{(2)}_{\mathrm{TT}}(p^2) + \Gamma^{(2)}_{\mathrm{TT}}(p^2)\,\mathcal{C}^{\dagger}(p^2;\alpha_{\mathrm{gf}}), \tag{7.205}$$

where $\mathcal{C}$ is a local matrix functional built from 1PI insertions involving the Nielsen kernel $\mathcal{N}^A = \left\langle \zeta\,\partial_{\alpha_{\mathrm{gf}}}\Psi \right\rangle_\Gamma$ (cf. (7.202)). Equation (7.205) holds on the band.

Sketch. Differentiate (7.201) functionally with respect to the fields (twice) at vanishing sources; use the Quantum Action Principle to organise the result as a sum of local 1PI insertions. The terms proportional to $\delta\Gamma/\delta\varphi$ generate factors of $\Gamma^{(2)}$; after projection with Π_{T}, longitudinal pieces drop out on $\mathcal{H}_{\mathrm{phys}}$, leaving the bilinear form (7.205). See [117] and [91]. □

Proposition 7.288 (Gauge-parameter independence of pole masses)**.** *Let $-m_i^2(\lambda_{\mathrm{UV}},\ell)$ be a simple physical pole of the transverse 1PI kernel, i.e. there exists a nonzero vector v_i such that*

$$\Gamma^{(2)}_{\mathrm{TT}}(-m_i^2)\,v_i = 0, \qquad v_i^{\dagger}\,\Gamma^{(2)\prime}_{\mathrm{TT}}(-m_i^2)\,v_i \neq 0, \tag{7.206}$$

where $' = \partial/\partial p^2$. Then, on $\mathcal{H}_{\mathrm{phys}}$ and on the band,

$$\boxed{\partial_{\alpha_{\mathrm{gf}}}\, m_i^2(\lambda_{\mathrm{UV}},\ell) = 0.} \tag{7.207}$$

Proof. Differentiate (7.206) with respect to α_{gf}:

$$\left(\partial_{\alpha_{\mathrm{gf}}}\Gamma^{(2)}_{\mathrm{TT}}\right)(-m_i^2)\,v_i - \left(\partial_{\alpha_{\mathrm{gf}}}m_i^2\right)\Gamma^{(2)\prime}_{\mathrm{TT}}(-m_i^2)\,v_i = 0.$$

Left-multiply by $v_i^{\dagger}$ and insert (7.205) at $p^2 = -m_i^2$:

$$v_i^{\dagger}\Big[\mathcal{C}\,\Gamma^{(2)}_{\mathrm{TT}} + \Gamma^{(2)}_{\mathrm{TT}}\,\mathcal{C}^{\dagger}\Big]v_i - \left(\partial_{\alpha_{\mathrm{gf}}}m_i^2\right) v_i^{\dagger}\Gamma^{(2)\prime}_{\mathrm{TT}}v_i = 0.$$

By $\Gamma^{(2)}_{\mathrm{TT}}(-m_i^2)\,v_i = 0$ and $v_i^{\dagger}\Gamma^{(2)}_{\mathrm{TT}}(-m_i^2) = 0$, the first bracket vanishes. Using the simple-pole assumption, $v_i^{\dagger}\Gamma^{(2)\prime}_{\mathrm{TT}}(-m_i^2)v_i \neq 0$, we conclude $\partial_{\alpha_{\mathrm{gf}}}m_i^2 = 0$. This is the band/IR version of the standard Nielsen result [91]; see also [117]. □

Remark 7.289 (Residues and wavefunction factors)**.** Equation (7.207) constrains pole *positions*. Residues (wavefunction factors) can be gauge-parameter dependent; physical S-matrix elements remain gauge independent once LSZ and proper normalisations are applied (Nielsen identity at the level of amputated Green's functions) [117, 91]. Band/IR commutation with BRST ensures the same statements on $\mathcal{H}_{\mathrm{phys}}$.

Nielsen protection for Holosymmetric visibilities.

Proposition 7.290 (Nielsen protection for holosymmetric visibilities)**.** *Let the holosymmetric readout be*

$$\mathcal{C}_\alpha(\lambda_{\mathrm{UV}},\ell;T)=\frac{\mathrm{Tr}\Big(\mathsf{P}_{\mathrm{phys}}\,U_1\,\rho\,U_0^\dagger\,\mathsf{P}_{\mathrm{phys}}\Big)}{\mathrm{Tr}\Big(\mathsf{P}_{\mathrm{phys}}\rho\Big)},\qquad \Lambda_\alpha=\mathrm{Re}\Big(e^{-i\phi_\alpha}\mathcal{C}_\alpha\Big),$$

with band/IR-embedded, BRST-compatible branches $U_{0,1}$ and physical, band-limited ρ (Section 7.4.1 and eq. (7.145)). Assume the band Slavnov identity $\mathcal{S}(\Gamma)=0$ and the Nielsen master identity (7.201). Then

$$\partial_{\alpha_{\mathrm{gf}}}\mathcal{C}_\alpha=\Big\langle\mathcal{O}_\alpha\cdot\zeta\,\partial_{\alpha_{\mathrm{gf}}}\Psi\Big\rangle_{\mathcal{H}_{\mathrm{phys}}}+\Big\langle\mathcal{J}_\alpha\cdot\frac{\delta\Gamma}{\delta\varphi}\Big\rangle_{\mathcal{H}_{\mathrm{phys}}},\tag{7.208}$$

for some local, band-limited composites $\mathcal{O}_\alpha,\mathcal{J}_\alpha$ determined by the functional dependence of $U_{0,1}$ and ρ on the elementary fields.[8] *Moreover, on $\mathcal{H}_{\mathrm{phys}}$ the first term is BRST-exact and decouples by the quartet mechanism/positivity, and the second is proportional to the equations of motion; therefore,* on shell*,*

$$\boxed{\partial_{\alpha_{\mathrm{gf}}}\Lambda_\alpha(\lambda_{\mathrm{UV}},\ell)=0\quad\textit{on the band.}}\tag{7.209}$$

Sketch. Differentiate $\mathcal{C}_\alpha$ w.r.t. α_{gf}. All α_{gf}-dependence enters through $\Psi(\alpha_{\mathrm{gf}})$, hence through the fields' functionals appearing in $U_{0,1}$ and ρ. Insert the Nielsen master identity (7.201) in the EOM/linearised form to rewrite each variation as a sum of a $\zeta(\partial_{\alpha_{\mathrm{gf}}}\Psi)$ insertion and an EOM-proportional term, yielding (7.208). Because $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ commute with BRST and carry ghost number 0 (eq. (7.145)), band/IR operations do not generate extra terms. On $\mathcal{H}_{\mathrm{phys}}$, the ζ-exact contribution vanishes by quartet positivity/Källén–Lehmann positivity (Section 7.2.6 and eq. (7.12)), and the EOM contribution vanishes on shell (eqs. (6.333) and (6.334)). Taking the real part with a fixed reference phase leaves the conclusion unchanged, proving (7.209). See [117, 91]. □

Background-field Nielsen identity and Ward compatibility.

Proposition 7.291 (Background-field Nielsen identity and Ward compatibility)**.** *Adopt the background-field method and choose the gauge fermion Ψ so that background gauge invariance is manifest. Then the band-limited 1PI functional Γ satisfies, simultaneously,*

$$\mathcal{W}^a_{\mathrm{bkg}}\Gamma=0,\qquad\frac{\partial\Gamma}{\partial\alpha_{\mathrm{gf}}}=-\,\mathcal{B}_\Gamma\Big(\partial_{\alpha_{\mathrm{gf}}}\Psi\Big)\quad\textit{(band Nielsen identity (7.201)).}$$

Consequently, the proofs of the pole-mass gauge independence (7.207) and the visibility protection (7.209) simplify: background Ward identities fix the transverse tensor structures a priori, and the Nielsen variation reduces to $\mathcal{B}_\Gamma$-exact insertions in the background-covariant basis.

[8] Concretely, $\mathcal{O}_\alpha$ collects the insertions generated by the chain rule acting on $U_{0,1}$ and ρ together with the Nielsen kernel $\mathcal{N}^A=\langle\zeta\,\partial_{\alpha_{\mathrm{gf}}}\Psi\rangle_\Gamma$; $\mathcal{J}_\alpha$ multiplies the Euler–Lagrange operators from $\delta\Gamma/\delta\varphi$. Locality follows from the QAP.

Sketch. In the BFM, the gauge-fixed action is invariant under background gauge transformations generated by $\mathcal{W}^a_{\rm bkg}$, while quantum gauge variations are controlled by BRST. Choosing Ψ background-covariant leaves $\mathcal{W}^a_{\rm bkg}\Gamma = 0$ intact [114]. The standard algebraic-renormalisation derivation of the Nielsen identity (linearised Slavnov form) carries over verbatim [91]. Transversality of current and gauge two-point functions then follows from background Ward identities, and the Nielsen variation remains an EOM- plus BRST-exact insertion, giving (7.207)–(7.209). □

Corollary 7.292 (Band/IR scope and regulator robustness)**.** *Because* $[\hat{Q}_{\rm BRST}, P_x(\lambda_{\rm UV})] = 0$ *and* $\zeta\big(K^{(\ell)}_{\rm IR} * \cdot\big) = 0$ *(cf. (7.145)), the identities (7.201)–(7.209) hold* on the band. *Moreover, under admissible regulator changes* $(P, K) \to (\tilde{P}, \tilde{K})$ *that preserve the low-energy fixed point, the induced variations of these identities are bounded by the regulator tolerance* $\varepsilon_{\rm reg}$ *from Section 7.3.7; hence the conclusions (gauge-parameter independence of pole masses and visibilities) are regulator-stable within* $\varepsilon_{\rm reg}$.

Remark 7.293 (Bibliography note)**.** For the original gauge-parameter identity, see Nielsen's paper `Nielsen1975gauge` (N. K. Nielsen, *Nucl. Phys. B* **101** (1975) 173–188). Algebraic-renormalisation treatments and proofs in the BRST framework are presented in [117], with textbook exposition in [91]. Background-field gauge and its Ward identities are reviewed in [114].

7.5.12 Callan–Symanzik Identity on the Band

Definition 7.294 (Field content and bandwise renormalisation data)**.** Throughout, we use the conventional symbols

$$\Phi \text{ (gaiamiti)}, \qquad \Theta \text{ (tantrakymon)}, \qquad c \text{ (ghost)},$$

with antifields/sources suppressed when not needed. Bandwise renormalised fields are defined by multiplicative Z-factors at fixed bare parameters,

$$\Phi_0 = Z_\Phi^{1/2}\,\Phi, \qquad \Theta_0 = Z_\Theta^{1/2}\,\Theta, \qquad c_0 = Z_c^{1/2}\,c, \tag{7.210}$$

where UV projection $P_x(\lambda_{\rm UV})$ and IR smoothing $K^{(\ell)}_{\rm IR}$ act as spectators and commute with BRST and with the RG flow (ghost number 0; cf. (7.145)). The anomalous dimensions are defined at fixed bare parameters:

$$\gamma_\Phi := \mu\,\frac{d\ln Z_\Phi}{d\mu}\Big|_{\rm bare}, \qquad \gamma_\Theta := \mu\,\frac{d\ln Z_\Theta}{d\mu}\Big|_{\rm bare}, \qquad \gamma_c := \mu\,\frac{d\ln Z_c}{d\mu}\Big|_{\rm bare}. \tag{7.211}$$

Let $\{g_r\}$ denote the set of renormalised (dimensionless) couplings, with beta functions $\beta_r(g) := \mu\,\frac{dg_r}{d\mu}\big|_{\rm bare}$. If relevant masses m_a appear, define $\gamma_{m_a} := \mu\,\frac{d\ln m_a}{d\mu}\big|_{\rm bare}$.

Definition 7.295 (Band Callan–Symanzik operator)**.** For a 1PI n-point function at the symmetric vacuum, let (n_Φ, n_Θ, n_c) be the external leg counts of (Φ, Θ, c), respectively. The band Callan–Symanzik (CS) operator acting at fixed bare parameters and fixed knobs $(\lambda_{\rm UV}, \ell)$ is

$$\mathcal{D}^{\rm band}_{\rm CS} := \mu\frac{\partial}{\partial\mu} + \sum_r \beta_r(g)\frac{\partial}{\partial g_r} - \frac{n_\Phi}{2}\gamma_\Phi - \frac{n_\Theta}{2}\gamma_\Theta - \frac{n_c}{2}\gamma_c \;\; (+\sum_a \gamma_{m_a}\, m_a\,\partial_{m_a} \text{ if masses}). \tag{7.212}$$

Theorem 7.296 (Callan–Symanzik identity on the band)**.** *Let* $\Gamma^{(n)}_{\text{band}}(\{p_i\};\mu,g)$ *be any renormalised* 1*PI n-point function computed from* (7.144) *with the scale-tagged dynamics of eqs.* (6.333) *and* (6.334) *and with the UV/IR knobs* $(\lambda_{\text{UV}},\ell)$ *held fixed. Then*

$$\boxed{\mathcal{D}^{\text{band}}_{\text{CS}}\,\Gamma^{(n)}_{\text{band}}(\{p_i\};\mu,g)=0\,,} \tag{7.213}$$

i.e. the standard Callan–Symanzik equation holds on the band. *If relevant masses* m_a *are present, include the mass terms indicated in* (7.212).

Sketch. At fixed bare parameters, the bare n-point function is μ-independent. Express it in terms of renormalised fields/couplings using (7.210). Differentiating w.r.t. μ and using the chain rule yields

$$\Big(\mu\partial_\mu+\sum_r\beta_r\partial_{g_r}\Big)\Gamma^{(n)}_{\text{band}}=\frac{1}{2}\Big(n_\Phi\,\gamma_\Phi+n_\Theta\,\gamma_\Theta+n_c\,\gamma_c\Big)\Gamma^{(n)}_{\text{band}}\ (+\ \sum_a\gamma_{m_a}\,m_a\,\partial_{m_a}\ \text{if masses}),$$

which rearranges to (7.213). Band/IR operations commute with the RG (and with BRST) by (7.145), so they do not contribute additional terms. The argument is standard (multiplicative renormalisability and RG at fixed bare parameters) [91, 82]. □

Remark 7.297 (Consistency with ST/Nielsen and scale-tagged EOM)**.** The CS identity (7.213) coexists with the band Slavnov identity and the Nielsen identities; the latter constrain gauge-parameter variations, while (7.213) governs μ-flow of renormalised amplitudes. Since $P_x(\lambda_{\text{UV}})$ and $K^{(\ell)}_{\text{IR}}$ carry ghost number 0 and commute with BRST, the algebraic relations imposed by Section 7.5.9 (and their consequences for visibilities) are compatible with (7.213). The scale-tagged equations of motion eqs. (6.333) and (6.334) fix the operator basis and ensure that all counterterms respect the band structure and the symmetric-point identities [117, 91, 82].

Composite operators and mixing — formal treatment.

Definition 7.298 (Bandwise renormalisation of composites and anomalous dimensions)**.** Let $\{\mathcal{O}_I\}$ be a local basis of BRST-closed, gauge-invariant composites representing $H^0(\mathcal{B}_\Gamma)$ at the symmetric point (fields: Φ for gaiamiti, Θ for tantrakymon, and c for ghosts). Define bandwise renormalised insertions by a finite mixing matrix $Z^{\mathcal{O}}$ at fixed bare parameters,

$$[\mathcal{O}_I]\ =\ \sum_J\big(Z^{\mathcal{O}}\big)_{IJ}\,\mathcal{O}^{\text{bare}}_J\ +\ \mathcal{B}_\Gamma(\cdots)\ +\ \partial_\mu(\cdots), \tag{7.214}$$

where the $\mathcal{B}_\Gamma$-exact and total-derivative pieces encode allowed mixings with BRST-exact and improvement operators; they decouple on $\mathcal{H}_{\text{phys}}$ (positivity/quartet mechanism). The *band anomalous-dimension matrix* is

$$\gamma_{IJ}(g)\ :=\ \Big(\big(Z^{\mathcal{O}}\big)^{-1}\,\mu\frac{d}{d\mu}\,Z^{\mathcal{O}}\Big)_{IJ}\Big|_{\text{bare}}, \tag{7.215}$$

with UV/IR knobs $(\lambda_{\text{UV}},\ell)$ commuting with BRST and with the RG flow, cf. (7.145).

Theorem 7.299 (Band Callan–Symanzik equation for composite insertions)**.** *Let a renormalised n-point 1PI function with m insertions of renormalised composites $[\mathcal{O}_{I_1}], \dots, [\mathcal{O}_{I_m}]$ be denoted by*

$$G^{(n;m)}_{\text{band}}\big(\{p\}; \mu, g\big).$$

At fixed bare parameters and fixed knobs $(\lambda_{\text{UV}}, \ell)$,

$$\Big(\mu\partial_\mu + \beta\cdot\partial_g\Big)\,[\mathcal{O}_I] \;=\; \sum_J \gamma_{IJ}\,[\mathcal{O}_J] \;+\; \mathcal{B}_\Gamma(\cdots) \;+\; \partial_\mu(\cdots), \tag{7.216}$$

and consequently

$$\Big(\mu\partial_\mu + \beta\cdot\partial_g - \tfrac{n_\Phi}{2}\gamma_\Phi - \tfrac{n_\Theta}{2}\gamma_\Theta - \tfrac{n_c}{2}\gamma_c - \sum_a \gamma_{m_a}\, m_a\partial_{m_a}\Big)\, G^{(n;m)}_{\text{band}} = \sum_{k=1}^{m}\sum_J \gamma_{I_k J}\, G^{(n;m)}_{\text{band}}(\mathcal{O}_{I_k}\to\mathcal{O}_J), \tag{7.217}$$

where the right-hand side replaces the k-th insertion by its mixed partner. The $\mathcal{B}_\Gamma$-exact and total-derivative contributions drop out of $\mathcal{H}_{\text{phys}}$ matrix elements.

Sketch. Differentiate (7.214) at fixed bare parameters; use the definition (7.215). Insert this variation into the RG equation for correlators at fixed bare parameters (the proof of Theorem 7.296) and use linearity of the insertion map defined by Γ (QAP). BRST-exact and improvement terms yield contact contributions that vanish in physical matrix elements and do not affect the 1PI kernels away from coincident points [117, 91]. □

Corollary 7.300 (Conserved currents in cohomology)**.** *If the anomaly filters hold (no gauge/mixed/gravitational anomalies) and J^μ_a is a conserved Noether current (global G), then*

$$\gamma_{J_a} = 0 \quad \text{in } H^0(\mathcal{B}_\Gamma), \tag{7.218}$$

i.e. any RG variation is an improvement plus a BRST-exact term (non-renormalisation of conserved currents), cf. (7.188) and Prop. 7.257 [117, 91].

Compatibility with Slavnov–Taylor and gauge independence — formal treatment.

Proposition 7.301 (ST stability of the RG data on the band)**.** *Assume the band Slavnov identity $\mathcal{S}(\Gamma) = 0$ and its linearisation $\mathcal{B}_\Gamma$ (Section 7.5.9). Then the RG generator $\mathcal{D}^{\text{band}}_{\text{CS}}$ (Definition 7.295) satisfies*

$$[\mathcal{D}^{\text{band}}_{\text{CS}},\, \mathcal{B}_\Gamma] = 0, \qquad \mathcal{B}_\Gamma\big(\gamma, \beta\big) = 0 \quad \textit{(cohomologically)}, \tag{7.219}$$

so the CS equations (7.213)–(7.216) are stable under BRST and their data (β, γ) are constrained by the ST cohomology (algebraic renormalisation).

Sketch. The QAP implies that $\mu\partial_\mu$ at fixed bare parameters acts by local counterterm insertions. These insertions preserve the ST identity if and only if they lie in $\ker \mathcal{B}_\Gamma$ modulo $\operatorname{im} \mathcal{B}_\Gamma$; hence $[\mathcal{D}^{\text{band}}_{\text{CS}}, \mathcal{B}_\Gamma] = 0$ and (7.219) [117]. □

Theorem 7.302 (Gauge-parameter independence of physical β-functions)**.** *Let $\alpha_{\rm gf}$ enter only through $\Psi(\alpha_{\rm gf})$, and assume the band Nielsen identity (7.201). For any* physical *coupling g_r (coupling solely to cohomology representatives in $H^0(\mathcal{B}_\Gamma)$), one has*

$$\partial_{\alpha_{\rm gf}}\beta_r(g)=0,\qquad \partial_{\alpha_{\rm gf}}\gamma_{IJ}(g)=\Big[\gamma(g),\mathcal{C}(\alpha_{\rm gf})\Big]_{IJ}\quad\textit{(similarity change of basis)},\tag{7.220}$$

where $\mathcal{C}$ is the local Nielsen kernel acting within the composite-operator space (band version of the standard result). Thus the eigenvalues *of γ in $H^0(\mathcal{B}_\Gamma)$, and all physical β-functions, are gauge-parameter independent [117, 91].*

Sketch. Apply $\partial_{\alpha_{\rm gf}}$ to (7.213)–(7.216) and use the Nielsen identity in the linearised form $\partial_{\alpha_{\rm gf}}\Gamma=-\mathcal{B}_\Gamma(\partial_{\alpha_{\rm gf}}\Psi)$. For physical couplings, the induced variations are $\mathcal{B}_\Gamma$-exact (EOM/BRST insertions), hence they cannot modify β_r. For composite operators, $\partial_{\alpha_{\rm gf}}\gamma$ is a commutator with the Nielsen kernel (basis change), leaving its spectrum in cohomology invariant. □

Corollary 7.303 (Background-field method)**.** *If the background-field method is used with a background-invariant gauge fermion, background Ward identities fix the renormalisation of gauge couplings directly from the background two-point kernel; consequently, β_g is manifestly gauge-parameter independent on the band [114, 91].*

Proposition 7.304 (RG improvement for holosymmetric visibilities)**.** *Let $\mathcal{C}_\alpha$ be the band/IR-embedded coherence amplitude and $\Lambda_\alpha=\mathrm{Re}(e^{-i\phi_\alpha}\mathcal{C}_\alpha)$ the corresponding visibility (Section 7.4.1). Assume multiplicative renormalisability on the band and the Callan–Symanzik identities (7.213)–(7.216) at fixed bare parameters. Then,* on shell *in $\mathcal{H}_{\rm phys}$,*

$$\boxed{\Big(\mu\partial_\mu+\beta\cdot\partial_g\Big)\,\mathcal{C}_\alpha(\lambda_{\rm UV},\ell;\mu,g)=0,\qquad \Big(\mu\partial_\mu+\beta\cdot\partial_g\Big)\,\Lambda_\alpha(\lambda_{\rm UV},\ell;\mu,g)=0\ ,}\tag{7.221}$$

up to $\mathcal{B}_\Gamma$-exact and total-derivative contact terms, which decouple on $\mathcal{H}_{\rm phys}$. Equivalently, $\mathcal{C}_\alpha$ and Λ_α are RG invariants whose residual scale dependence is confined to the running couplings $g(\mu)$.

Sketch. Write $\mathcal{C}_\alpha=N_\alpha/D_\alpha$ with

$$N_\alpha=\mathrm{Tr}\Big(\mathsf{P}_{\rm phys}\,U_1\,\rho\,U_0^\dagger\,\mathsf{P}_{\rm phys}\Big),\qquad D_\alpha=\mathrm{Tr}\Big(\mathsf{P}_{\rm phys}\rho\Big),$$

where $U_{0,1}$ and ρ are built from renormalised fields (Φ,Θ,c) and renormalised couplings $\{g_r\}$, with UV/IR knobs held fixed and commuting with BRST (eq. (7.145)). Applying the band CS operator (7.468) to N_α and D_α and using (7.213)–(7.216) gives two terms: (i) a piece proportional to the anomalous dimensions from field rescalings, and (ii) a piece $\beta\cdot\partial_g$. Because N_α and D_α involve identical field content (same external-leg counting in the insertion map), the anomalous-dimension contributions cancel in the ratio. The remaining BRST-exact/improvement mixings are $\mathcal{B}_\Gamma$-exact or total derivatives and thus vanish on $\mathcal{H}_{\rm phys}$ (quartet mechanism/positivity and the QAP), leaving $(\mu\partial_\mu+\beta\cdot\partial_g)\mathcal{C}_\alpha=0$. Taking a fixed reference phase does not affect RG invariance, hence (7.221) also holds for Λ_α. See [91, 82] for CS/RG, and [117] for the cohomological decoupling on physical matrix elements. □

Definition 7.305 (Band-matched renormalisation scale)**.** To minimise large logarithms in $\Gamma^{(n)}_{\text{band}}$ entering the readout, choose a *band-matched* scale

$$\mu_{\text{band}} = \mu_{\text{band}}(\lambda_{\text{UV}}, \ell)\,, \quad \text{e.g.} \quad \mu_{\text{band}} \sim \ell^{-1} \ \text{ or } \ \mu_{\text{band}} \sim \max\{|p_i|\} \text{ of the kinematics},$$

so that perturbative logs $\ln(\mu/\mu_{\text{band}})$ are small. RG-improved expressions then evaluate $\mathcal{C}_\alpha, \Lambda_\alpha$ with running couplings $g(\mu_{\text{band}})$ while keeping $(\lambda_{\text{UV}}, \ell)$ fixed (they commute with the RG), stabilising the plateau metrics under scale variations [91, 82].

Remark 7.306 (Gauge/BRST compatibility and residuals)**.** The ST/Nielsen identities ensure that the β-functions of physical couplings are gauge-parameter independent, and that any gauge-parameter variation of composite mixings is a similarity transformation in cohomology (eigenvalues invariant) [117, 91]. Therefore the RG-improved visibilities remain gauge/BRST robust on the band. Within certified plateaus (Section 7.3.9), the residual μ-dependence from higher orders is subleading and accounted for in the tolerance budget.

Trace identity (massless, anomaly-free baseline).

Proposition 7.307 (Band-local trace identity: massless, anomaly-free baseline)**.** *Work at the G-invariant symmetric point (Quadrant I) with massless renormalised fields Φ (gaiamiti), Θ (tantrakymon), and ghosts $c, \bar{c}, b$, and assume the anomaly filters hold (no gauge/mixed/-gravitational anomalies). Couple the renormalised theory to a background metric and define the renormalised energy–momentum tensor by functional differentiation of the band-limited action (sources as in (7.144)), with the UV/IR knobs $(\lambda_{\text{UV}}, \ell)$ commuting with BRST, cf. (7.145). Then the trace satisfies the RG-improved, band-local identity*

$$\boxed{\langle T^{\mu}{}_{\mu}(x)\rangle = \sum_r \beta_r(g)\,\big\langle \mathcal{O}_r(x)\big\rangle \;+\; (\text{equations of motion}) \;+\; \partial_\mu(\cdots)\,,} \tag{7.222}$$

where $\mathcal{O}_r$ are the renormalised operators coupled to the (dimensionless) couplings g_r in $S^{(\lambda_{\text{UV}},\ell)}_{\Phi,\Theta}$, and the last two terms are contact terms that vanish in $\mathcal{H}_{\text{phys}}$ matrix elements away from coincident insertions (BRST-exact/EOM improvements). Identity (7.222) holds on the band *and is regulator-stable under admissible changes of $(P_x, K^{(\ell)}_{\text{IR}})$.*

Sketch. Introduce a local Weyl (scale) variation of the background metric and use the Quantum Action Principle to represent the insertion of $T^{\mu}{}_{\mu}$ as the response of the renormalised effective action Γ to a local change of the renormalisation scale. At fixed bare parameters, $\mu\dfrac{d}{d\mu}\Gamma = 0$ implies the Callan–Symanzik generator acts on renormalised couplings and fields; translating this local RG variation into operator insertions yields the $\sum_r \beta_r \mathcal{O}_r$ term, while field rescalings produce EOM and improvement terms [91, 82]. Absence of gauge/mixed/-gravitational anomalies forbids additional local cocycles in the trace (no Weyl anomaly in the assumed baseline); any BRST-exact contributions decouple on $\mathcal{H}_{\text{phys}}$ by quartet positivity. Band/IR operations commute with BRST and preserve locality at the smearing scale by (7.145), so the same algebraic argument holds *on the band*. Regulator stability follows from Section 7.3.7. □

Lemma 7.308 (Local CS identity from the trace)**.** *Insertion of* (7.222) *into renormalised* n*-point* 1*PI functions and integration over* x *reproduces the band Callan–Symanzik equation* (7.213)*; conversely,* (7.213) *implies* (7.222) *up to EOM/improvement terms.*

Remark 7.309 (Scheme and regulator considerations)**.** (i) While β- and γ-functions depend on the renormalisation scheme beyond leading order, the *structure* of (7.222) and its Slavnov–Taylor (ST) compatibility are scheme-independent consequences of algebraic renormalisation [117, 91]. (ii) Admissible regulator changes $(P, K) \to (\tilde{P}, \tilde{K})$ preserving the low-energy fixed point leave (7.222) invariant within the tolerance $\varepsilon_{\mathrm{reg}}$ (Section 7.3.7). (iii) On $\mathcal{H}_{\mathrm{phys}}$, BRST-exact mixings and total derivatives drop out of physical observables and out of the visibilities' RG flow (positivity/decoupling), consistent with Section 7.2.6 and Section 7.5.12.

Bibliography. For CS/RG and their operator form of the trace identity, see [91, 82]. The cohomological (ST/BRST) constraints ensuring compatibility and decoupling are discussed in [117]; background-field perspectives on β-functions appear in [114].

7.5.13 Sector/Projector Compatibility and Plateau Protection

Definition 7.310 (Bandwise sector projectors)**.** Let $\{\Pi_\beta\}_{\beta\in\mathcal{B}}$ be a finite or countable family of mutually orthogonal projectors on the one–particle (or appropriate tensor) space such that

$$\Pi_\beta^\dagger = \Pi_\beta, \qquad \Pi_\beta \Pi_{\beta'} = \delta_{\beta\beta'}\Pi_\beta, \qquad \sum_\beta \Pi_\beta = \mathbb{1}.$$

Assume each Π_β is G-invariant and BRST-closed, and commutes with the band/IR knobs:

$$[\Pi_\beta, Q_a] = 0, \quad [\Pi_\beta, \hat{Q}_{\mathrm{BRST}}] = 0, \quad [\Pi_\beta, P_x(\lambda_{\mathrm{UV}})] = 0, \quad [\Pi_\beta, K_{\mathrm{IR}}^{(\ell)}] = 0, \tag{7.223}$$

where Q_a generate the global G symmetry and $P_x, K_{\mathrm{IR}}^{(\ell)}$ satisfy (7.145). For a multiplet $\varphi = (\Phi, \Theta, c, \bar{c}, b)$, define the *sector-restricted fields* $\varphi_\beta := \Pi_\beta \varphi$ and the *sector-restricted effective action*

$$\Gamma_\beta[\varphi] \;:=\; \Gamma[\Pi_\beta \varphi] \;=\; \Gamma[\varphi_\beta]. \tag{7.224}$$

Proposition 7.311 (Sectorwise Slavnov–Taylor and global Ward identities)**.** *Under* (7.223) *and at the symmetric point,*

$$\boxed{\mathcal{S}\big(\Gamma_\beta\big) = 0, \qquad \mathcal{W}_a\big(\Gamma_\beta\big) = 0,} \tag{7.225}$$

i.e. both the Slavnov functional and the global Ward operator act compatibly on each sector. In particular, the linearised ST operator about Γ_β*,*

$$\mathcal{B}_{\Gamma_\beta} X \;:=\; \frac{d}{d\epsilon}\Big|_{\epsilon=0} \mathcal{S}\big(\Gamma_\beta + \epsilon X\big),$$

is nilpotent: $\mathcal{B}_{\Gamma_\beta}^2 = 0$.

Proof. The identities $\mathcal{S}(\Gamma)=0$ and $\mathcal{W}_a(\Gamma)=0$ hold on the band by Sections 7.5.2 and 7.5.4 and commute with the insertions $P_x, K_{\mathrm{IR}}^{(\ell)}$ by (7.145). Using $[\Pi_\beta, \hat{Q}_{\mathrm{BRST}}]=[\Pi_\beta, Q_a]=0$ and the fact that Π_β has ghost number 0, one may pull Π_β through the antibracket/linearised form of $\mathcal{S}$ and through the Ward operators; thus $\mathcal{S}(\Gamma[\Pi_\beta\varphi])=0$ and $\mathcal{W}_a(\Gamma[\Pi_\beta\varphi])=0$. Nilpotency of $\mathcal{B}_{\Gamma_\beta}$ follows from the graded Jacobi identity as in the standard proof [117, 442, 91]. □

Proposition 7.312 (Cohomology factorisation and sector decomposition)**.** *Let $H^0(\mathcal{B}_\Gamma)$ denote the BRST cohomology at ghost number 0 for Γ. Then the map*

$$\bigoplus_\beta H^0(\mathcal{B}_{\Gamma_\beta}) \;\longrightarrow\; H^0(\mathcal{B}_\Gamma), \qquad \big([\mathcal{O}_\beta]\big)_\beta \;\mapsto\; \Big[\ \sum_\beta \Pi_\beta\,\mathcal{O}_\beta\ \Big],$$

is an isomorphism of vector spaces. In particular, any physical composite operator has a unique decomposition into sector components, and conversely each sector cohomology class lifts canonically to a class of the full theory.

Sketch. Orthogonality and completeness of $\{\Pi_\beta\}$ imply a direct-sum decomposition of the configuration space. Since Π_β commute with BRST, $\mathcal{B}_\Gamma$ preserves sectors and decomposes as a direct sum of $\mathcal{B}_{\Gamma_\beta}$. Exactness and closure are preserved by the direct sum, yielding the claimed isomorphism. □

Definition 7.313 (Sector-resolved readouts and weights)**.** For a channel α and interrogation time T, define the sector-resolved coherence amplitude and visibility by inserting Π_β at readout,

$$\mathcal{C}_{\alpha|\beta} := \frac{\mathrm{Tr}\Big(\Pi_\beta\,\mathsf{P}_{\mathrm{phys}}\,U_1^{(\alpha)}\,\rho_{\mathrm{phys}}\,U_0^{(\alpha)\dagger}\,\mathsf{P}_{\mathrm{phys}}\,\Pi_\beta\Big)}{\mathrm{Tr}(\Pi_\beta\,\mathsf{P}_{\mathrm{phys}}\rho_{\mathrm{phys}})}, \qquad \Lambda_{\alpha|\beta} := \mathrm{Re}\big(e^{-i\phi_\alpha}\mathcal{C}_{\alpha|\beta}\big), \tag{7.226}$$

cf. (7.133). The sector weights are $w_\beta := \mathrm{Tr}(\Pi_\beta\,\mathsf{P}_{\mathrm{phys}}\rho_{\mathrm{phys}})/\,\mathrm{Tr}(\mathsf{P}_{\mathrm{phys}}\rho_{\mathrm{phys}})$.

Theorem 7.314 (Block-diagonality and convex additivity)**.** *If, in addition to (7.223), each branch generator is sector-compatible,*

$$[\Pi_\beta, \mathcal{H}_j^{(\alpha)}(t,x;\lambda_{\mathrm{UV}})]=0 \quad \textit{for all } (t,x),\ \beta,\ j\in\{0,1\}, \tag{7.227}$$

then the band/IR-smeared propagators $U_j^{(\alpha)}$ are block-diagonal in the sector decomposition and the aggregate amplitude decomposes as

$$\mathcal{C}_\alpha = \sum_\beta w_\beta\,\mathcal{C}_{\alpha|\beta}, \qquad \Lambda_\alpha = \sum_\beta w_\beta\,\Lambda_{\alpha|\beta}. \tag{7.228}$$

Proof. Condition (7.227) implies $U_j^{(\alpha)}=\bigoplus_\beta U_j^{(\alpha,\beta)}$, so cross-sector coherences vanish under the sector insertions. Insert the resolution of identity $\sum_\beta \Pi_\beta = \mathbb{1}$ on both sides of ρ_{phys} inside the trace and use orthogonality to obtain (7.228). □

Lemma 7.315 (Leakage bound under weak sector breaking)**.** *Suppose (7.223) holds but (7.227) is violated weakly: $\epsilon_{\mathrm{leak}} := \sup_t \sum_\beta \|[\Pi_\beta, \mathcal{H}_j^{(\alpha)}(t,\cdot;\lambda_{\mathrm{UV}})]\|_{\mathrm{op}} < \infty$. Then for the band/IR propagators (Dyson/Duhamel expansion),*

$$\big\|\Pi_\beta^{\perp}\,U_j^{(\alpha)}(T)\,\Pi_\beta\big\|_{\mathrm{op}} \;\le\; T\,\epsilon_{\mathrm{leak}}\,\exp\Big(T\,\|\mathcal{H}_j^{(\alpha)}\|_{\mathrm{op}}\Big), \tag{7.229}$$

and the deviation from (7.228) obeys

$$\Big|\Lambda_\alpha - \sum_\beta w_\beta\, \Lambda_{\alpha|\beta}\Big| \;\le\; C\,T\,\epsilon_{\mathrm{leak}} \;+\; \mathcal{O}(T^2\epsilon_{\mathrm{leak}}^2), \tag{7.230}$$

for a constant C depending on $\|\rho_{\mathrm{phys}}\|_1$ and operator norms of the sector blocks.

Sketch. Apply Duhamel's formula (7.131)–(7.132) with the decomposition $\mathcal{H}_j = \sum_\beta \Pi_\beta \mathcal{H}_j \Pi_\beta +$ (off-diagonal) and iterate the integral inequality (Gronwall). Estimate the sector off-diagonal blocks of $U_j^{(\alpha)}$ by (7.229) and insert into the trace defining $\mathcal{C}_\alpha$; normalise by $\mathrm{Tr}(\mathsf{P}_{\mathrm{phys}}\rho_{\mathrm{phys}})$ to obtain (7.230). See, e.g., [115, Sec. X.11] for operator-norm evolution bounds. □

Proposition 7.316 (Sectorwise Nielsen/Slavnov protection)**.** *The sector-restricted functionals Γ_β obey the band Nielsen identity and the linearised ST identity with the same kernels as the full theory:*

$$\partial_{\alpha_{\mathrm{gf}}}\Gamma_\beta = -\,\mathcal{B}_{\Gamma_\beta}\big(\partial_{\alpha_{\mathrm{gf}}}\Psi\big), \qquad \mathcal{B}_{\Gamma_\beta}^2 = 0.$$

Consequently, sector-resolved pole masses and visibilities are gauge-parameter independent on shell, and the sector Callan–Symanzik data (β, γ) satisfy the ST cohomological constraints as in the full theory [117, 91].

Proof. Apply the proofs of Propositions 7.229 and 7.301 and Theorem 7.302 with $\varphi \mapsto \Pi_\beta\varphi$; commutations (7.223) ensure functorial descent to each sector. □

Definition 7.317 (Sectorwise plateau protection and certification)**.** Within a certified joint plateau $\mathfrak{P}_{\{\mathrm{law,state}\}}$ (cf. Section 7.4.5), define sectorwise bounds

$$\|\nabla_{(u,v)}\Lambda_{\alpha|\beta}\| \le \varepsilon_{\mathrm{grad},\beta}, \qquad \|\mathbf{H}\Lambda_{\alpha|\beta}\|_{\mathrm{op}} \le \varepsilon_{\mathrm{curv},\beta}, \qquad \mathrm{osc}_{B_r}\,\Lambda_{\alpha|\beta} \le \varepsilon_{\mathrm{osc},\beta},$$

and a leakage tolerance $T\,\epsilon_{\mathrm{leak}} \le \varepsilon_{\mathrm{leak}}$ from (7.229). The *sectorwise certification* accepts $(\lambda_\star, \ell_\star)$ if, for all β with $w_\beta > \tau$ (weight threshold),

$$\varepsilon_{\mathrm{grad},\beta},\ \varepsilon_{\mathrm{curv},\beta},\ \varepsilon_{\mathrm{osc},\beta},\ \varepsilon_{\mathrm{leak}} \ \text{ lie below policy tolerances,}$$

and the gradient-alignment coefficient between channels (restricted to sector β) satisfies $\chi_{\star|\beta} \le \chi_{\max}$ (avoidance of near-cancellations). The aggregate plateau then inherits the bounds via (7.228) or, under weak breaking, with additive error $\mathcal{O}(\varepsilon_{\mathrm{leak}})$ by (7.230).

Corollary 7.318 (Regulator stability per sector)**.** *Under admissible changes $(P_x, K_{\mathrm{IR}}^{(\ell)}) \to (\tilde P_x, \tilde K_{\mathrm{IR}}^{(\ell)})$ preserving the low-energy fixed point, sector-resolved visibilities satisfy*

$$\big|\Lambda_{\alpha|\beta} - \tilde\Lambda_{\alpha|\beta}\big| \;\le\; \varepsilon_{\mathrm{reg},\beta},$$

with $\varepsilon_{\mathrm{reg},\beta}$ controlled as in Section 7.3.7; aggregate stability follows by summing with weights w_β (or with the leakage correction from (7.230)).

Remark 7.319 (Operational upshot)**.** Sectorwise ST/Ward/Nielsen protection (Propositions 7.311 and 7.316) guarantees that sector-resolved readouts are physically meaningful and gauge robust; block-diagonality yields exact convex additivity (7.228), while weak violations are quantitatively bounded by (7.230). These statements hold *on the band* and descend to $\mathcal{H}_{\mathrm{phys}}$ by BRST decoupling [117, 442, 91, 115].

Sectorwise Callan–Symanzik flow (formal treatment).

Proposition 7.320 (Sectorwise CS identity)**.** *Let* $\{\Pi_\beta\}$ *be the BRST-closed, G-invariant band/IR-compatible sector projectors of Section 7.5.13, commuting with the UV/IR knobs as in* (7.223)*. For the renormalised n-point* 1*PI kernels computed on the band,*

$$\Gamma^{(n)}_{\mathrm{band},\beta}(\{p_i\};\mu,g) := \Gamma^{(n)}_{\mathrm{band}}\Big[\Pi_\beta\varphi\Big](\{p_i\};\mu,g),$$

the Callan–Symanzik (CS) operator acts sectorwise with the same β*- and* γ*-functions as in Section 7.5.12:*

$$\boxed{\Big(\mu\partial_\mu + \beta\cdot\partial_g - \tfrac{n_\Phi}{2}\gamma_\Phi - \tfrac{n_\Theta}{2}\gamma_\Theta - \tfrac{n_c}{2}\gamma_c\Big)\,\Gamma^{(n)}_{\mathrm{band},\beta}(\{p_i\};\mu,g) \;=\; 0\,.} \tag{7.231}$$

Moreover, (7.231) *is compatible with the sectorwise Slavnov–Taylor and G-Ward identities* (7.225)*, so the usual cohomological constraints on counterterms hold* within each sector *[117, 91].*

Proof sketch. The CS equation on the band, (7.213), follows from RG invariance at fixed bare data in a multiplicatively renormalisable scheme [91, 82]. Since Π_β has ghost number 0 and commutes with BRST, G, and the knobs (Definition 7.310), it can be inserted either before or after renormalisation without altering renormalisation constants. Consequently, the same (β,γ) govern the sector-restricted amplitudes. Compatibility with Slavnov/Ward identities sectorwise is Proposition 7.311, ensuring stability of (7.231) under $\mathcal{S}$ and $\mathcal{W}_a$ [117, 91]. □

Smoothness in ℓ and spectral gaps (formal treatment).

Definition 7.321 (Sectoring from an IR metric)**.** Let $A(\ell)$ be a self-adjoint band/IR-local operator built covariantly from $\bar\Sigma^{(\ell)}_{\mu\nu}$ (e.g. the appropriate spin/tensor Casimir or a $\bar\Sigma^{(\ell)}$-elliptic kinetic symbol) whose spectral decomposition defines $\{\Pi_\beta(\ell)\}$. Assume a *non-crossing condition* on the scan window: there exists gap > 0 such that separated spectral clusters remain at least gap apart for all ℓ in the window.

Lemma 7.322 (Differentiability of sector projectors)**.** *Under Definition 7.321, each Riesz projector*

$$\Pi_\beta(\ell) \;=\; \frac{1}{2\pi i}\oint_{\mathcal{C}_\beta}\big(z - A(\ell)\big)^{-1}\,dz$$

is C^1 *in* ℓ *and satisfies the Kato-type bound*

$$\big\|\partial_{\ln\ell}\Pi_\beta(\ell)\big\| \;\le\; \frac{\|\,\partial_{\ln\ell}A(\ell)\,\|}{\mathrm{gap}} \;\times\; C_{\mathcal{C}_\beta}, \qquad C_{\mathcal{C}_\beta} = \sup_{z\in\mathcal{C}_\beta}\|(z-A)^{-1}\|^2, \tag{7.232}$$

where $\mathcal{C}_\beta$ *encloses only the cluster defining* Π_β*.*

Proof. This is standard analytic perturbation theory: differentiating the Riesz integral and using the resolvent identity yields (7.232); the spectral gap controls the resolvent norms [115]. □

Corollary 7.323 (Smooth sector-restricted kernels and visibilities)**.** *Let* $\Gamma^{(n)}_{\mathrm{band},\beta}$ *and the sector-resolved visibilities* $\Lambda_{\alpha|\beta}$ *be defined as in* (7.231) *and* (7.226)*. If, in addition, the embedded generators satisfy the Duhamel regularity* (7.132)*, then*

$$\partial_{\ln \ell}\Gamma^{(n)}_{\mathrm{band},\beta}, \quad \partial_{\ln \ell}\Lambda_{\alpha|\beta} \text{ exist and are continuous in } \ell,$$

with quantitative bounds of the form

$$\left|\partial_{\ln \ell}\Lambda_{\alpha|\beta}\right| \;\le\; C_1 \left\|\partial_{\ln \ell}\Pi_\beta\right\| \;+\; C_2 \int_0^T dt\, \left\|\partial_{\ln \ell} K^{(\ell)}_{\mathrm{IR}}\right\| \left\|\mathcal{H}^{(\alpha)}_j(t,\cdot;\lambda_{\mathrm{UV}})\right\|_{\mathrm{op}}, \tag{7.233}$$

for constants $C_{1,2}$ *depending on* $\|\rho_{\mathrm{phys}}\|_1$ *and operator norms of the sector blocks. Combining* (7.232) *and* (7.233) *gives*

$$\left|\partial_{\ln \ell}\Lambda_{\alpha|\beta}\right| \;\lesssim\; \frac{\|\partial_{\ln \ell}\bar{\Sigma}^{(\ell)}\|}{\mathrm{gap}} \;+\; \int_0^T dt\, \left\|\partial_{\ln \ell} K^{(\ell)}_{\mathrm{IR}}\right\| \|\mathcal{H}^{(\alpha)}_j\|_{\mathrm{op}}.$$

Proof sketch. Differentiate the definitions in (7.226) using the product rule, $\partial_{\ln \ell}\Pi_\beta$ from Lemma 7.322, and $\partial_{\ln \ell}U^{(\alpha)}_j$ from the Duhamel identity (7.132). Estimate traces by Schatten/operator norms and use unitarity of the propagators to obtain (7.233). □

Remark 7.324 (RG and cohomology inside sectors)**.** Equation (7.231) ensures that RG improvement can be carried out *per sector*. Since Π_β commute with BRST and G, the sector cohomologies $H^0(\mathcal{B}_{\Gamma_\beta})$ govern allowable counterterms and operator mixings inside each sector exactly as in the full theory (Proposition 7.312, [117, 91]). Smooth ℓ-dependence under the non-crossing hypothesis guarantees that sector plateaus and their certification metrics vary regularly across the scan window (cf. Section 7.4.5).

Plateau protection (sectorwise): formal treatment.

Proposition 7.325 (Sectorwise plateau protection)**.** *Let* $\mathfrak{P}_{\{\mathrm{law,state}\}}$ *be a certified joint plateau in the log–knob chart* $(u,v) = (\ln \lambda_{\mathrm{UV}}, \ln \ell)$*, and let* $\{\Pi_\beta(\ell)\}$ *be the BRST-closed, G-invariant sector projectors with* C^1 *dependence on* ℓ *(non-crossing hypothesis; Kato theory). For any identity functional* $\mathcal{I} \in \{\mathcal{S},\ \mathcal{W}_a,\ \text{CS operator}\}$ *and any sector* β*, define*

$$\Delta_{\mathcal{I},\beta}(u,v) := \mathcal{I}\Big[\Gamma\Big[\Pi_\beta(\ell)\,\varphi\Big]\Big](u,v), \qquad \varphi = (\Phi,\Theta,c,\bar{c},b).$$

Then on every geodesic ball $B_\rho(u_\star,v_\star) \subset \mathfrak{P}_{\{\mathrm{law,state}\}}$ *one has the quantitative bounds*

$$\|\nabla_{(u,v)}\Delta_{\mathcal{I},\beta}\| \;\le\; C_\beta\,\varepsilon_{\mathrm{grad}}\,, \qquad \|\mathbf{H}\,\Delta_{\mathcal{I},\beta}\|_{\mathrm{op}} \;\le\; C_\beta\,\varepsilon_{\mathrm{curv}}\,, \qquad \mathrm{osc}_{B_\rho}\,\Delta_{\mathcal{I},\beta} \;\le\; C_\beta\,\varepsilon_{\mathrm{osc}} + \varepsilon_{\mathrm{reg}}, \tag{7.234}$$

where C_β *depends only on sector smoothness constants* $\sup_{B_\rho}\|\partial_u\Pi_\beta\|$, $\sup_{B_\rho}\|\partial_v\Pi_\beta\|$ *(controlled by the spectral gap and* $\|\partial_{\ln \ell}\bar{\Sigma}^{(\ell)}\|$ *via Kato) and on operator norms of the Fréchet derivatives of* $\mathcal{I}$ *acting on band-limited functionals. In the exact theory,* $\Delta_{\mathcal{I},\beta} \equiv 0$ *by the sectorwise identities, and* (7.234) *quantifies the stability of* numerical/experimental *tests of these identities under UV/IR scans inside the plateau.*

Proof sketch. Step 1 (Exact sectorwise identities). Since Π_β carries ghost number 0 and commutes with BRST and G, the Slavnov functional and the global Ward operators act compatibly on sector-restricted functionals:

$$\mathcal{S}\big(\Gamma[\Pi_\beta\varphi]\big) = 0, \qquad \mathcal{W}_a\big(\Gamma[\Pi_\beta\varphi]\big) = 0,$$

and the sectorwise CS equation holds with the same (β, γ) (see eqs. (7.225) and (7.231); [117, 442, 91]). Thus, ideally, $\Delta_{\mathcal{I},\beta} \equiv 0$.

Step 2 (Differentiability and chain rule). Inside the plateau we work in knobs (u, v). The u- and v-dependence of $\Gamma[\Pi_\beta\varphi]$ comes from the embedded generators and from $\Pi_\beta(\ell)$. Duhamel formulas give band-safe expressions for ∂_u and ∂_v acting on the propagators (eqs. (7.131) and (7.132)); sector smoothness follows from the non-crossing hypothesis and Kato perturbation theory, yielding C^1 bounds for Π_β in ℓ ([115], cf. Lemma 7.322). Applying the chain rule to $\Delta_{\mathcal{I},\beta} = \mathcal{I} \circ \Gamma \circ (\Pi_\beta, \text{knobs})$ and estimating by operator norms yields

$$\|\partial_{u,v}\Delta_{\mathcal{I},\beta}\| \;\le\; \|\mathrm{D}\mathcal{I}\| \Big(\|\partial_{u,v}\Gamma\| + \|\mathrm{D}_\Gamma\Gamma\| \, \|\partial_{u,v}\Pi_\beta\| \Big).$$

Step 3 (Plateau and regulator bounds). By definition of a certified plateau, the relevant band-limited kernels/visibilities satisfy uniform gradient, curvature, and oscillation bounds (Section 7.3.5, e.g. eq. (7.137)), and admissible regulator changes obey eq. (7.67). Collecting these estimates and absorbing fixed operator norms into C_β gives the gradient and Hessian bounds in (7.234). The oscillation bound follows from the same plateau oscillation control plus ε_{reg} for permissible $(P, K) \to (\tilde{P}, \tilde{K})$ swaps.

Step 4 (Conclusion). Since the exact identities give $\Delta_{\mathcal{I},\beta} \equiv 0$, the bounds (7.234) show that *any measured deviation* within B_ρ remains uniformly small and controlled by $(\varepsilon_{\text{grad}}, \varepsilon_{\text{curv}}, \varepsilon_{\text{osc}}, \varepsilon_{\text{reg}})$, i.e. the identities are *plateau-protected* under UV/IR scans. □

Remark 7.326 (Dependence of C_β)**.** Under the spectral-gap hypothesis for the sectoring operator built from $\bar{\Sigma}^{(\ell)}_{\mu\nu}$, Kato's bound gives $\|\partial_{\ln\ell}\Pi_\beta\| \lesssim \|\partial_{\ln\ell}\bar{\Sigma}^{(\ell)}\|/\text{gap}$ [115]. Hence one may take $C_\beta = C_{\mathcal{I}}\Big(1 + \sup_{B_\rho} \|\partial_{\ln\ell}\bar{\Sigma}^{(\ell)}\|/\text{gap}\Big)$, with $C_{\mathcal{I}}$ depending only on the (fixed) operator norms associated with $\mathcal{I}$ on the band-limited function space (cf. [117, 91, 82]).

Sector-resolved visibilities and compatibility (formal treatment).

Proposition 7.327 (Nielsen independence and CS flow, sector-resolved)**.** *Let $\{\Pi_\beta\}$ be the BRST-closed, G-invariant band/IR-compatible sector projectors of Section 7.5.13, and let $\mathcal{C}_{\alpha|\beta}$ and $\Lambda_{\alpha|\beta}$ be defined by inserting Π_β at readout as in (7.133). Then, on the band and on shell,*

$$\partial_{\alpha_{\text{gf}}}\Lambda_{\alpha|\beta} = 0 \qquad \textit{(Nielsen gauge-parameter independence)}, \tag{7.235}$$

and the sector-resolved 1PI kernels obey the Callan–Symanzik flow with the same β- and γ-functions as in Section 7.5.12:

$$\Big(\mu\partial_\mu + \beta\cdot\partial_g - \tfrac{n_\Phi}{2}\gamma_\Phi - \tfrac{n_\Theta}{2}\gamma_\Theta - \tfrac{n_c}{2}\gamma_c\Big)\,\Gamma^{(n)}_{\text{band},\beta} = 0, \qquad \Gamma^{(n)}_{\text{band},\beta} := \Gamma^{(n)}_{\text{band}}\big[\Pi_\beta\varphi\big], \tag{7.236}$$

and consequently the sector-resolved visibilities inherit RG improvement on shell.

Proof sketch. Sectorwise Slavnov/Ward identities (7.225) hold because Π_β commutes with BRST and with G and carries ghost number 0 ([117, 442, 91]). The Nielsen master identity on the band (Section 7.5.11) then descends to each sector, yielding (7.235) exactly as in Section 7.3.6. Bandwise renormalisation commutes with Π_β, so the CS operator acts sectorwise with unchanged (β, γ) (Proposition 7.320 and [91, 82]); thus (7.236) follows. □

Definition 7.328 (Sector compatibility and certified point)**.** For a fixed channel α and a collection of sectors $\mathcal{B}$, define the sectorwise plateau sets $\mathfrak{P}_{\alpha|\beta}$ by the certification criteria of Sections 7.3.5 and 7.3.9 applied to $\Lambda_{\alpha|\beta}$. A *sector-compatible operating point* is any $(\lambda_\star, \ell_\star)$ such that

$$(\lambda_\star, \ell_\star) \;\in\; \bigcap_{\beta\in\mathcal{B}} \mathfrak{P}_{\alpha|\beta}. \tag{7.237}$$

Corollary 7.329 (No spurious aggregate plateaus)**.** *Assume the exact sector symmetry* $[\Pi_\beta, \mathcal{H}_j^{(\alpha)}] = 0$ *so that evolution is block-diagonal; then the aggregate visibility decomposes as a convex combination of sector visibilities,* $\Lambda_\alpha = \sum_\beta w_\beta \Lambda_{\alpha|\beta}$ *with* $w_\beta \geq 0$ *and* $\sum_\beta w_\beta = 1$ *(cf. (7.133)). On a joint plateau ball* $B_\rho(u_\star, v_\star)$*, sectorwise plateau protection (7.234) implies*

$$\|\nabla_{(u,v)}\Lambda_\alpha\| \;\leq\; \sum_\beta w_\beta \,\|\nabla_{(u,v)}\Lambda_{\alpha|\beta}\| \;\leq\; \Big(\max_\beta C_\beta\Big)\,\varepsilon_{\mathrm{grad}},$$

and analogous bounds for the Hessian and oscillation. Conversely, if an aggregate *plateau occurs while some* $\Lambda_{\alpha|\beta}$ *violates the plateau bounds, this requires near-cancellation of sector gradients; such cancellations are excluded by imposing sector coherence constraints (pairwise alignment coefficients bounded by* $\chi_{\max} < 1$*) within the certification pipeline (Section 7.3.11). Hence certified aggregate plateaus cannot arise from fragile cancellations between sectors.*

Proof sketch. The gradient/Hessian inequalities follow from linearity of differentiation and the convex decomposition, plus (7.234). The converse uses the Gram-matrix estimate for $\sum_\beta w_\beta \nabla\Lambda_{\alpha|\beta}$: if all pairwise cosine alignments are bounded by $\chi_{\max} < 1$, small aggregate gradient forces each sector gradient to be small up to a $1/(1-\chi_{\max})$ factor, contradicting a sector's plateau violation. □

Remarks. (i) Background-field implementations preserve sectorwise identities with background Ward operators, so (7.235)–(7.236) hold sector by sector [114]. (ii) If a spectral crossing in $\Pi_\beta(\ell)$ occurs, restrict to subregions where the gap is open or merge sectors to a gapped $\tilde{\beta}$; sectorwise certification and the bounds of Proposition 7.325 then apply unchanged ([115]). (iii) All statements are *on the band* and descend to $\mathcal{H}_{\mathrm{phys}}$ by positivity/decoupling (Section 7.2.6 and eq. (7.12); [241, 82]).

7.5.14 Anomaly Filters and Cohomological Triviality

Definition 7.330 (Bandwise anomaly filters)**.** On the band (UV projector $P_x(\lambda_{\mathrm{UV}})$ and IR window $K_{\mathrm{IR}}^{(\ell)}$ commuting with BRST, eq. (7.145)), impose the vanishing of the gauge, mixed, and gravitational anomaly indicators recorded in Section 7.2.8:

$$\mathfrak{A}_{\mathrm{gauge}} = \mathbf{0}, \qquad \mathfrak{A}_{\mathrm{mixed}} = \mathbf{0}, \qquad \mathfrak{A}_{\mathrm{grav}} = \mathbf{0}.$$

These conditions are understood at the G-invariant symmetry point (Quadrant I), and are stable under admissible regularisation choices that preserve the low-energy fixed point.

Proposition 7.331 (Cohomological triviality of bandwise breakings)**.** *Let Γ be the band-limited 1PI functional obeying the Quantum Action Principle (QAP). Suppose that at loop order n the Slavnov functional and global Ward operator admit a local breaking $\Delta^{(n)}$ of ghost number $+1$:*

$$\mathcal{S}(\Gamma) = \hbar^n \Delta^{(n)} + \mathcal{O}(\hbar^{n+1}), \qquad \mathcal{W}_a \Gamma = \hbar^n \Delta_a^{(n)} + \mathcal{O}(\hbar^{n+1}),$$

with the Wess–Zumino consistency conditions $\mathcal{B}_\Gamma \Delta^{(n)} = 0$ and $\mathcal{W}_a \Delta^{(n)} + \cdots = 0$. If the anomaly filters of Definition 7.330 hold, then the breaking is cohomologically trivial *on the band:*

$$\Delta^{(n)} = \mathcal{B}_\Gamma \, \widehat{\Sigma}^{(n)} + d\, \widehat{\mathcal{R}}^{(n)}, \qquad \Delta_a^{(n)} = \mathcal{W}_a \widehat{\Sigma}^{(n)} + d\, \widehat{\mathcal{J}}_a^{(n)}, \tag{7.238}$$

for some local counterterms $\widehat{\Sigma}^{(n)}$ and local currents $\widehat{\mathcal{R}}^{(n)}$, $\widehat{\mathcal{J}}_a^{(n)}$. Consequently, the redefinition

$$\boxed{\Gamma \;\mapsto\; \Gamma - \hbar^n \widehat{\Sigma}^{(n)}} \tag{7.239}$$

restores exactly*, at order n, the bandwise Slavnov–Taylor and global Ward identities:*

$$\mathcal{S}(\Gamma) = 0, \qquad \mathcal{W}_a \Gamma = 0.$$

Proof sketch. By QAP, $\Delta^{(n)}$ is a local insertion of ghost number $+1$ satisfying $\mathcal{B}_\Gamma \Delta^{(n)} = 0$ (Wess–Zumino consistency) [117, 442]. The anomaly filters imply that the relevant cohomology $H^{1,d}(\zeta \mid d)$ is trivial in the gauge, mixed, and gravitational sectors, so every consistent breaking is a sum of a $\mathcal{B}_\Gamma$-exact term and a spacetime total derivative, yielding (7.238). Subtracting the local counterterm (7.239) removes the breaking at order n. Band/IR operations commute with BRST and preserve locality at the IR scale, so the standard algebraic-renormalisation argument applies verbatim on the band [117, 443]. □

Corollary 7.332 (Law-channel identity at the symmetry point)**.** *Assume BRST nilpotency and self-adjointness (7.196), the anomaly filters of Definition 7.330, and exact bandwise Slavnov/Ward identities restored order by order via (7.239). In the* law *channel of the holosymmetric probe (Section 7.4.2), the branch difference is BRST-exact on $\mathcal{H}_{\mathrm{phys}}$, so the BRST equivalence theorem (Theorem 7.278) gives*

$$\boxed{\Lambda^\star_{\mathrm{law}} = 1, \qquad \partial_{\alpha_{\mathrm{gf}}} \Lambda^\star_{\mathrm{law}} = 0, \qquad \partial_{(\lambda_{\mathrm{UV}},\ell)} \Lambda^\star_{\mathrm{law}} \;\approx\; 0 \quad \textit{on certified plateaus.}} \tag{7.240}$$

The last bound uses the plateau metrics and regulator stability (eqs. (7.67), (7.137) and (7.138)).

Proof sketch. The law branches differ by $H_0 \mapsto H_0 + \zeta \Delta G$; by Proposition 7.331 and the quartet/positivity mechanism, this deformation is invisible on $\mathcal{H}_{\mathrm{phys}}$. Hence the overlap amplitude equals 1 and is Nielsen-protected (gauge-parameter independence on shell), yielding (7.240). Drift bounds inside certified plateaus follow from the geometric metrics and regulator tolerance established earlier. □

Remark 7.333 (State-channel baseline)**.** For the *state* channel under the infinitesimal symmetric kick (7.124), the symmetric-state expansion (7.128) gives $\Lambda^{\star}_{\text{state}} = 1 + \mathcal{O}(\epsilon^2)$. Thus any significant, regulator-stable suppression of $|\mathcal{C}_{\text{state}}|$ is a diagnostic of *state-level* tension or decoherence rather than a law-level anomaly (triage in Section 7.4.4).

Citations. Algebraic renormalisation, ST identities, and anomaly cohomology: [117, 442, 443]; BRST positivity and physical sector: [91].

7.5.15 Identity Ledger for Later SSB Analysis

At the G-invariant vacuum and on $\mathcal{H}_{\text{phys}}$, the following identities hold *on the band*. Items **I1**–**I6** are exact (cohomological/functional) identities; **I7** records baseline readouts whose reported values are exact on shell (law channel) or perturbatively controlled (state channel) and, operationally, are certified within plateau tolerances:

I1. Global Ward identities for G: position- and momentum-space forms eqs. (7.181) and (7.182) with current conservation eq. (7.183); two-point transversality eq. (7.184) and positive spectral densities eq. (7.186).

I2. Slavnov–Taylor master equation & linearisation: eqs. (7.189) and (7.190), implying cohomological decoupling of unphysical modes and Z-factor constraints at Quadrant I (vertex-level relations eq. (7.194)); any loop-level breaking is cohomologically trivial and removable by counterterms eq. (7.239) [117].

I3. BRST nilpotency and equivalence theorem: eqs. (7.196) and (7.198), ensuring invariance of $\mathcal{H}_{\text{phys}}$ matrix elements under BRST-exact deformations (including law-branch homotopies).

I4. Nielsen identities (gauge flows): master identity eq. (7.201) with on-shell gauge-parameter independence of pole masses eq. (7.207) and of visibilities/coherence amplitudes eq. (7.209).

I5. Bandwise Callan–Symanzik equation: CS identity for 1PI functions eq. (7.213) and composite-operator flow eq. (7.216), compatible with Slavnov–Taylor and gauge independence of physical β-functions.

I6. Sectorwise validity and plateau protection: sector-restricted ST/Ward identities eq. (7.225), sectorwise CS flow eq. (7.231), and stability bounds eq. (7.234) ensuring robustness of identity tests under UV/IR scans within certified plateaus.

I7. Law/state probe baselines at symmetry: law-channel unity and robustness eq. (7.240) (exact on shell and bandwise); state-channel symmetry limit from eqs. (7.126) and (7.128) with $\Lambda^{\star}_{\text{state}} = 1 + \mathcal{O}(\epsilon^2)$, and deviations interpreted via the diagnostic triage in Section 7.4.4.

These entries constitute the *baseline ledger* against which spontaneous or induced symmetry breaking (SSB) in other quadrants will be detected, certified, and quantified [117, 442, 91, 443].

7.6 Synthesis of Quadrant I: Discoveries and Baseline Status

D1. Symmetric vacuum fixed and clean EOM on the band. A G-invariant vacuum $(\bar{\Phi}, \bar{\Theta})$ has been fixed (cf. Section 7.1). The *scale-tagged*, BRST-consistent equations of motion have been established in Euler–Lagrange and in "clean" kinetic forms eqs. (6.333), (6.334) and (6.374), each defined *on the band* using the UV projector $P_x(\lambda_{\mathrm{UV}})$ and IR window $K_{\mathrm{IR}}^{(\ell)}$ Section 7.3. Band operations commute with BRST eq. (7.30), hence descend functorially to the physical sector $\mathcal{H}_{\mathrm{phys}}$ (Section 7.2).

D2. BRST health: unitarity, positivity, and causal propagation. On $\mathcal{H}_{\mathrm{phys}}$ the Kugo–Ojima quartet mechanism removes ghosts/longitudinals, yielding positive Källén–Lehmann densities for gauge-invariant observables Section 7.2.6 and eq. (7.12). The physical kinetic operator is hyperbolic with respect to $\bar{\Sigma}_{\mu\nu}^{(\ell)}$ eq. (7.15), and the physical mass operator is non-negative eqs. (7.13) and (7.14). No ghosts/tachyons occur at the symmetric point and causal propagation is well-posed (see also [442, 91, 82]).

D3. UV/IR instrumentation and certified plateaus. Families $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ have been fixed (Section 7.3.1), and the holosymmetric visibilities $\Lambda_\alpha(\lambda_{\mathrm{UV}}, \ell)$ exhibit a *non-empty joint plateau* $\mathfrak{P}_{\{\mathrm{law,state}\}}$ Section 7.3.5 and eq. (7.64). Plateau metrics satisfy the bounds eq. (7.137) and the quantitative criteria in Section 7.3.10. Regularisation stability holds across admissible (P,K) choices Section 7.3.7, and gauge/BRST robustness follows from Nielsen-type on-shell independence Section 7.3.6 and eq. (7.209). A representative operating point $(\lambda_\star, \ell_\star) \in \mathfrak{P}_{\{\mathrm{law,state}\}}$ is fixed via the certification pipeline Section 7.3.9, with bounded slope/curvature/oscillation.

D4. Holosymmetric probe instrumented and calibrated. A controlled two-path unitary U_{ctrl} has been constructed on $\mathcal{H}_{\mathrm{phys}}$ for the *law* and *state* channels Sections 7.4 and 7.4.2, yielding complex coherences $\mathcal{C}_\alpha$ and signed visibilities Λ_α Section 7.4.1. At the symmetric, anomaly-free baseline,

$$\Lambda_{\mathrm{law}}^\star = 1, \qquad \Lambda_{\mathrm{state}}^\star = 1 + \mathcal{O}(\epsilon^2) \quad (\epsilon \to 0 \text{ in eq. (7.124)}),$$

both evaluated at $(\lambda_\star, \ell_\star)$ and stable within plateau tolerances and the error budget Section 7.4.6. Deviations are diagnostic: $|\Lambda_{\mathrm{law}}| < 1$ signals law-level incoherence (e.g. anomaly or regulator pathology), whereas $|\Lambda_{\mathrm{state}}| < 1$ signals state-level symmetry tension or IR non-uniformity (triage in Section 7.4.4).

D5. Exact identities recorded as the baseline ledger. Global Ward identities for G eqs. (7.181) to (7.184), the Slavnov–Taylor master equation and its linearisation eqs. (7.189) and (7.190) (with vertex-level consequences eq. (7.194)), BRST nilpotency/equivalence eqs. (7.196) and (7.198), the Nielsen identities eqs. (7.201), (7.207) and (7.209), and the bandwise Callan–Symanzik equation eqs. (7.213) and (7.216) have been made explicit. These identities hold sectorwise with $\{\Pi_\beta\}$ Section 7.5.13, are *plateau-protected* under UV/IR scans eq. (7.234), and are compatible with BRST/gauge robustness [117, 442, 91, 114].

D6. Implications for subsequent analyses. The Quadrant I baseline is now fully specified: (i) a healthy, causal, anomaly-free physical sector; (ii) regulator- and gauge-robust visibilities anchored at a certified operating point; and (iii) an identity ledger constraining renormalisation and protecting physical conclusions. With anomaly filters enforced and potential breakings cohomologically trivial Section 7.5.14, the *law channel* obeys the unity identity eq. (7.240).

Table 7.1: Particle-like modes at the Quadrant I symmetric baseline (on $\mathcal{H}_{\text{phys}}$). All statements are plateau-certified and gauge/BRST robust; tags marked "provisional" require cross-quadrant confirmation.

Sector / origin	**Mode**	**Spin** s	**Mass / gap**	**DoF (3+1D)**
Gaiamiti $\hat{\mathbf{\Phi}}$	Tensor	2	massless	2 (helicities ± 2)
Tantrakyma $\hat{\mathbf{\Theta}}$	Gauge vector	1	massless	2 (helicities ± 1)
Scalars in $\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}$	Scalar excitations	0	gapped ($m^2 > 0$)	1

In short: Quadrant I closes with a robust, scale-stable, and gauge-independent baseline—no ghosts, no tachyons, bright *law* and *state* channels, and the spectrum above. Any stronger particle-level tagging must now earn its status by surviving plateau tests across symmetry-broken quadrants (Sections 7.3 and 7.4).

Quadrant I baseline. Quadrant I now stands as a healthy, unbroken theory calibrated by explicit UV/IR controls (Section 7.3) and safeguarded by the full suite of BRST, Ward, and Slavnov–Taylor identities (Sections 7.2 and 7.5). *On the band*, the holosymmetric probe (Section 7.4) functions as a dial-stable interferometer: the *law* visibility is unity within plateau tolerances (eq. (7.139) and Section 7.3), excluding anomalies or regulator pathologies, and the *state* visibility is unity up to the symmetric-kick order, $\Lambda^{\star}_{\text{state}} = 1 + \mathcal{O}(\epsilon^2)$ (cf. eq. (7.128)), excluding hidden spontaneous breaking in the prepared state. Ghost and longitudinal artefacts are eliminated on $\mathcal{H}_{\text{phys}}$ by BRST cohomology and the quartet mechanism (Sections 7.2.6 and 7.5.10; [442, 108, 105, 104, 243, 82]), so the observed spectrum has the expected unbroken-phase content.

Gaiamiti (chronogeometric) sector $\hat{\mathbf{\Phi}}$*—tensor modes.* Transverse–traceless fluctuations of the *gaiamiti* field $\hat{\mathbf{\Phi}}(x)$ behave as a massless spin-2 mode with two physical polarisations in 3+1 dimensions—"graviton-like" in pattern—subject to our policy that such labels remain *provisional* until they survive symmetry-broken quadrants under plateau scans (Sections 7.3 and 7.4). BRST and Ward structures remove all non-physical pieces (Section 7.5; [243, 82]), and plateau tests ensure insensitivity to the precise UV cutoff or IR smoothing within the certified window (Section 7.3).

Tantrakyma (internal/gauge) sector $\hat{\mathbf{\Theta}}$*—vector modes.* Unbroken symmetry and the Slavnov–Taylor relations (Section 7.5; [243, 82]) yield massless transverse vectors (spin 1) from the *tantrakyma* field $\hat{\mathbf{\Theta}}(x)$. Longitudinal components and their ghost partners do not appear in

the physical cohomology (Section 7.2.6; [108, 442]). This matches the interferometric readout: bright *law* and *state* channels on the plateau (Sections 7.3 and 7.4).

Scalar content and mixing. Scalar fluctuations in both $\hat{\mathbf{\Phi}}$ and $\hat{\mathbf{\Theta}}$ are gapped at the symmetric point (positive Hessian eigenvalues; no tachyons) (Section 7.2 and eqs. (7.13) and (7.14)), and there are no Goldstones in Quadrant I. Allowed gauge-invariant gaiamiti–tantrakyma (Φ–Θ) mixings simply diagonalise into physical combinations; spurious longitudinal admixtures are removed by BRST (cf. eqs. (7.13) and (7.14), Section 7.2.6; [442, 243]).

Spin census and dimensional remark. The symmetric spectrum on $\mathcal{H}_{\text{phys}}$ thus reads: a massless spin-2 tensor (from $\hat{\mathbf{\Phi}}$, provisional), massless spin-1 gauge vectors (from $\hat{\mathbf{\Theta}}$), and massive spin-0 scalars. In 3+1 dimensions this corresponds to helicities ± 2 (tensor), ± 1 (vectors), and the usual scalar representation; in other dimensions the content reorganises under the appropriate little group but remains "spin-2 $\oplus$ spin-1 $\oplus$ spin-0" ([91, 82]). Mixing respects spin, and the BRST projection guarantees that only transverse, positive-norm modes populate $\mathcal{H}_{\text{phys}}$ (Section 7.2.6; [108, 442]).

Closing statement. Quadrant I therefore closes with a robust, scale-stable, and gauge-independent baseline: no ghosts, no tachyons, a bright *law* channel, a bright *state* channel, and a physically intuitive spectrum (Sections 7.3 to 7.5). Any stronger particle-level tagging—graviton, Higgs, apeiron, chrono–kymon—must now earn its status by surviving the plateau tests across the symmetry-broken quadrants (Sections 7.3 and 7.4; [243, 442]).

7.7 Quadrant II: Spontaneous Symmetry Breaking (SSB)

7.7.1 Hold the Law Symmetric (Keep H_0 Fixed)

Objective. In Quadrant II we probe *spontaneous* (not explicit) symmetry breaking. The *law* channel is kept strictly symmetric by using the same clean kinetic Hamiltonian H_0 (and its covariant Lagrangian counterpart) on both branches of the law interferometer (Sections 7.3 and 7.4). State preparation is free to explore asymmetric vacua; the microscopic law remains identical across branches. Thus any loss of coherence is attributed to the *state* channel, and Quadrant-II membership is preserved under branch overlay (Sections 7.4 and 7.4.1; [117, 442, 91]).

1.1 Clean symmetric baseline

We work on the BRST-closed physical sector $\mathcal{H}_{\text{phys}}$ with UV projector $P_x(\lambda_{\text{UV}})$ and IR window $K_{\text{IR}}^{(\ell)}$ chosen BRST-closed (ghost number 0) and commuting with the channel projectors (Sections 7.2 and 7.3; cf. eq. (7.30)). Concretely, the unbroken *law* is specified by the scale-tagged equations of motion

$$\mathcal{E}_{\mathbf{\Phi}}^{(\lambda_{\text{UV}},\ell)} = 0, \qquad \mathcal{E}_{\mathbf{\Theta}}^{(\lambda_{\text{UV}},\ell)} = 0,$$

as defined in eqs. (6.333) and (6.334) (BRST reading via P_{phys}; ghosts/longitudinals decouple in cohomology). Their quadratic expansion around a G-invariant background determines the band-limited Hessian $\mathbb{H}^{(\lambda_{\text{UV}},\ell)}$ with kinetic blocks $\mathcal{K}_T, \mathcal{K}_C$ and potential (mass/mixing) blocks

$\mathcal{M}_{\bullet\bullet}$ (eq. (7.10); see Section 7.2.5). On $\mathcal{H}_{\rm phys}$ only the transverse parts contribute; physical poles are read from the $\Pi_{\rm T}$-projected blocks of $\mathbb{H}^{(\lambda_{\rm UV},\ell)}$, and these are Nielsen/BRST-stable under the UV/IR scans (Section 7.5).

Hamiltonian generator. Both branches of the interferometer use the *same* relational Hamiltonian $\hat{H}_0(t;\lambda_{\rm UV},\ell)$, obtained (on the band, inside the IR window) as the Legendre transform of the covariant quadratic Lagrangian whose kernel is $\mathbb{H}^{(\lambda_{\rm UV},\ell)}$. Equivalently, the branch unitaries $U_j^{(\alpha)}(T)$ are generated by BRST-compatible Hamiltonians that coincide in the law channel and differ only by ζ-exact deformations when we change presentation (cohomologically trivial on $\mathcal{H}_{\rm phys}$); see eqs. (7.97) and (7.130). Thus "keep H_0 fixed'' means: (i) no symmetry-breaking couplings in $\mathbb{H}^{(\lambda_{\rm UV},\ell)}$; (ii) identical G-invariant kinetic/mass blocks across branches; and (iii) any branch decoration in the law channel is BRST-exact and cancels in physical matrix elements (equivalence theorem).

Explicit form (general d). Write the operator field and momentum multiplets as

$$\hat{\boldsymbol{\varphi}}^{\mathsf{T}} := \big(\hat{\boldsymbol{\Phi}},\, \hat{\boldsymbol{\Theta}}\big), \qquad \hat{\boldsymbol{\pi}}^{\mathsf{T}} := \big(\hat{\pi}_{\boldsymbol{\Phi}},\, \hat{\pi}_{\boldsymbol{\Theta}}\big),$$

where the canonical momenta are taken along a chosen foliation with unit normal n_μ via the momentum currents of Definitions 6.267 and 6.268:

$$\hat{\pi}_{\boldsymbol{\Phi}} := n_\mu\, \Pi^\mu_{\boldsymbol{\Phi}}, \quad \Pi^\mu_{\boldsymbol{\Phi}} = \bar{\Sigma}^{\mu\nu}\, \mathcal{G}_T\big(\nabla_\nu \hat{\boldsymbol{\Phi}}\big), \qquad \hat{\pi}_{\boldsymbol{\Theta}} := n_\mu\, \Pi^\mu_{\boldsymbol{\Theta}}, \quad \Pi^\mu_{\boldsymbol{\Theta}} = \bar{\Sigma}^{\mu\nu}\, \mathcal{G}_C\big(\nabla_\nu \hat{\boldsymbol{\Theta}}\big).$$

Introduce the band/IR–tagged multiplets (cf. Definition 6.282 and Section 6.12.4)

$$\hat{\boldsymbol{\varphi}}_{\lambda_{\rm UV},\ell} := \mathfrak{B}_{\lambda_{\rm UV},\ell}[\hat{\boldsymbol{\varphi}}], \qquad \hat{\boldsymbol{\pi}}_{\lambda_{\rm UV},\ell} := \mathfrak{B}_{\lambda_{\rm UV},\ell}[\hat{\boldsymbol{\pi}}].$$

Let the quadratic kernels be

$$\mathcal{K} = \begin{pmatrix} \mathcal{K}_T & 0 \\ 0 & \mathcal{K}_C \end{pmatrix}, \qquad \mathcal{M}^2 = \begin{pmatrix} \mathcal{M}^2_{\Phi\Phi} & \mathcal{M}^2_{\Phi\Theta} \\ \mathcal{M}^2_{\Theta\Phi} & \mathcal{M}^2_{\Theta\Theta} \end{pmatrix},$$

where $\mathcal{K}_T, \mathcal{K}_C$ are the positive (self-adjoint) kinetic operators induced by the pairings g_T, g_C and the background weights $\bar{\Sigma}^{(\ell)\mu\nu}$, and $\mathcal{M}^2_{\bullet\bullet}$ are the Hessian (mass/mixing) blocks from the quadratic expansion about $(\bar{\Phi},\bar{\Theta})$ (cf. Proposition 7.2). Then the law Hamiltonian used on both branches is

$$\hat{H}_0(t;\lambda_{\rm UV},\ell) = \frac{1}{2}\int d^d x \left[\big(\hat{\boldsymbol{\pi}}_{\lambda_{\rm UV},\ell}\big)^\top \mathcal{K}\, \hat{\boldsymbol{\pi}}_{\lambda_{\rm UV},\ell} \;+\; \big(\hat{\boldsymbol{\varphi}}_{\lambda_{\rm UV},\ell}\big)^\top \mathcal{M}^2\, \hat{\boldsymbol{\varphi}}_{\lambda_{\rm UV},\ell}\right] \;+\; \big\{\hat{Q}_{\rm BRST},\, \Psi_\zeta\big\}. \tag{7.241}$$

Remarks. (i) The explicit $K^{(\ell)}_{\rm IR} P_x(\lambda_{\rm UV})\,(\cdot)\, P_x(\lambda_{\rm UV}) K^{(\ell)}_{\rm IR}$ insertions are absorbed into the definitions of the tagged operators via $\mathfrak{B}_{\lambda_{\rm UV},\ell}$ (cf. Definition 6.282); expanding them out reproduces our original "UV then IR" order. (ii) No projector $\Pi_{\rm T}$ is applied here: retaining the full multiplet ensures the off-diagonal mixing $\mathcal{M}^2_{\Phi\Theta}, \mathcal{M}^2_{\Theta\Phi}$ contributes as intended. (iii) Evaluating on $\mathcal{H}_{\rm phys}$ one may insert $\mathsf{P}_{\rm phys}$ before/after $\hat{H}_0$; by Section 7.1.3, the $\{\hat{Q}_{\rm BRST}, \Psi_\zeta\}$ piece alters only unphysical sectors and leaves physical spectra/visibilities gauge-parameter independent.

Kernel (bi-local) form, without new symbols. Define the band-projected quadratic kernel

$$\begin{aligned}\widetilde{\mathbb{H}}^{(\lambda_{\mathrm{UV}},\ell)} &= \mathrm{diag}\Big(K_{\mathrm{IR}}^{(\ell)}P_x(\lambda_{\mathrm{UV}})\Pi_{\mathrm{T}}, K_{\mathrm{IR}}^{(\ell)}P_x(\lambda_{\mathrm{UV}})\Pi_{\mathrm{T}}\Big)\\ &\quad\times\begin{pmatrix}\mathcal{K} & 0\\ 0 & \mathcal{M}^2\end{pmatrix}\mathrm{diag}\Big(K_{\mathrm{IR}}^{(\ell)}P_x(\lambda_{\mathrm{UV}})\Pi_{\mathrm{T}}, K_{\mathrm{IR}}^{(\ell)}P_x(\lambda_{\mathrm{UV}})\Pi_{\mathrm{T}}\Big).\end{aligned} \tag{7.242}$$

With

$$\hat\pi(\mathbf{x}) := \begin{pmatrix}\hat\pi_{\mathbf{\Phi}}(\mathbf{x})\\ \hat\pi_{\mathbf{\Theta}}(\mathbf{x})\end{pmatrix}, \qquad \hat\varphi(\mathbf{x}) := \begin{pmatrix}\hat{\mathbf{\Phi}}(\mathbf{x})\\ \hat{\mathbf{\Theta}}(\mathbf{x})\end{pmatrix},$$

the Hamiltonian reads

$$\hat H_0(t;\lambda_{\mathrm{UV}},\ell) = \frac{1}{2}\int d^dx\, d^dy\, \begin{pmatrix}\hat\pi(\mathbf{x})\\ \hat\varphi(\mathbf{x})\end{pmatrix}^{\top}\widetilde{\mathbb{H}}^{(\lambda_{\mathrm{UV}},\ell)}(\mathbf{x},\mathbf{y})\begin{pmatrix}\hat\pi(\mathbf{y})\\ \hat\varphi(\mathbf{y})\end{pmatrix} + \big\{\hat Q_{\mathrm{BRST}},\,\Psi_\zeta\big\}. \tag{7.243}$$

Ghost/longitudinal decoupling. Gauge fixing enters as the ζ-exact term $\{\,\hat Q_{\mathrm{BRST}},\Psi_\zeta\,\}$; the Faddeev–Popov/longitudinal sectors sit in separate blocks of $\mathbb{H}^{(\lambda_{\mathrm{UV}},\ell)}$ and are killed on $\mathcal{H}_{\mathrm{phys}}$ by the quartet mechanism ($\mathsf{P}_{\mathrm{phys}}\Box_{\mathrm{FP}}\mathsf{P}_{\mathrm{phys}} = 0 = \mathsf{P}_{\mathrm{phys}}\Pi_{\mathrm{L}}\mathsf{P}_{\mathrm{phys}}$).

All band operations are BRST-closed and commute with the symmetry generators; in particular,

$$[\hat Q_{\mathrm{BRST}}, P_x(\lambda_{\mathrm{UV}})] = [\hat Q_{\mathrm{BRST}}, K_{\mathrm{IR}}^{(\ell)}] = 0, \qquad [\mathsf{J}_\alpha, P_x] = [\mathsf{J}_\alpha, K_{\mathrm{IR}}^{(\ell)}] = 0 \quad (\forall\,\alpha), \tag{7.244}$$

so the scale-tagged objects descend to $\mathcal{H}_{\mathrm{phys}}$, and on-shell visibilities are gauge-parameter independent by the Nielsen identities (cf. eq. (7.115) and Section 7.5).

1.2 Law-channel branches and overlay

Introduce the ancilla two-path interferometer with branches $b \in \{1,2\}$ (Section 7.4). The *law* channel manipulates only the *presentation* of the *same* symmetric law fixed in §1.1; no explicit breaking is ever injected. Accordingly we choose branch operators $U_b^{(\mathrm{law})}$ that preserve the BRST charge, the symmetric Hamiltonian, and all G-generators:

$$\Big[U_b^{(\mathrm{law})},\,\hat H_0(t;\lambda_{\mathrm{UV}},\ell)\Big] = \Big[U_b^{(\mathrm{law})},\,\hat Q_{\mathrm{BRST}}\Big] = \Big[U_b^{(\mathrm{law})},\,\mathsf{J}_\alpha\Big] = 0 \quad (\forall\,\alpha), \tag{7.245}$$

with $\hat H_0$ as in eqs. (7.241) and (7.243). Compatibly with the scale-tagged pipeline, they also commute with the UV/IR/transverse projectors:

$$\Big[U_b^{(\mathrm{law})},\,P_x(\lambda_{\mathrm{UV}})\Big] = \Big[U_b^{(\mathrm{law})},\,K_{\mathrm{IR}}^{(\ell)}\Big] = \Big[U_b^{(\mathrm{law})},\,\Pi_{\mathrm{T}}\Big] = 0, \tag{7.246}$$

so that all band operations remain BRST-closed and G-covariant (cf. eq. (7.244) and Section 7.5). These conditions ensure that $U_b^{(\mathrm{law})}$ act as symmetry-preserving (Nielsen-compatible) reparametrisations of the same H_0.

Explicit realisations (presentation moves). It is often convenient to take $U_b^{(\mathrm{law})}$ from one (or a product) of the following classes, all of which satisfy eqs. (7.245) and (7.246) on $\mathcal{H}_{\mathrm{phys}}$:

1. **BRST-exact similarities:**
$$U_b^{(\mathrm{law})} = \exp\Big(i\{\,\hat{Q}_{\mathrm{BRST}},\,\Delta\Psi_b\,\}\Big), \tag{7.247}$$
where $\Delta\Psi_b$ is a change of the gauge-fixing fermion (e.g. a shift of ζ in Ψ_ζ); these are Nielsen-compatible on plateaus (Section 7.3; [117]).

2. **Background-field reparametrisations:** $U_b^{(\mathrm{law})}$ that reshuffle the background/quantum split without altering G or BRST structure (background-field method; [114]).

3. **Field redefinitions (equivalence moves):** local, invertible redefinitions that leave on-shell matrix elements invariant; at quadratic level these appear as orthogonal rotations within the Φ-Θ sector that commute with $\mathcal{K}$ and $\mathcal{M}^2$ (cf. eq. (7.241)).

4. **Regulator placement/order moves:** permutations of $P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}^{(\ell)}, \Pi_{\mathrm{T}}$ that pairwise commute on the certified plateau window (Sections 7.3 and 7.4.5), leaving $\hat{H}_0$ isospectral.

Overlay on system⊗ancilla. With ancilla basis $\{|1\rangle, |2\rangle\}$, the controlled law overlay is
$$\mathcal{U}_{\mathrm{law}} = |1\rangle\langle 1| \otimes U_1^{(\mathrm{law})} + |2\rangle\langle 2| \otimes U_2^{(\mathrm{law})}, \tag{7.248}$$
which preserves (i) BRST-physicality and (ii) exact G-symmetry of the law on $\mathcal{H}_{\mathrm{phys}}$. Any inter-branch difference is a Nielsen-type presentation/regulator placement that leaves on-shell observables invariant on certified plateaus (Sections 7.3 and 7.4.5; [117, 114]).

Law-channel contrast (why it stays at unity in Quadrant II). Define the branch-difference unitary
$$\Delta U_{\mathrm{law}} := \Big(U_2^{(\mathrm{law})}\Big)^\dagger U_1^{(\mathrm{law})}. \tag{7.249}$$
For any BRST-physical state $|\Psi\rangle \in \mathcal{H}_{\mathrm{phys}}$ and any physical observable $\mathcal{O}$ (BRST-closed, ghost-number 0) evaluated on the plateau,
$$\langle\Psi|\mathcal{O}|\Psi\rangle = \langle\Psi|\,\Big(\Delta U_{\mathrm{law}}\Big)^\dagger\,\mathcal{O}\,\Delta U_{\mathrm{law}}\,|\Psi\rangle, \tag{7.250}$$
so the controlled-interferometer visibility in the *law* channel reduces to
$$\mathcal{V}_{\mathrm{law}} = \Big|\langle\Psi|\,\Delta U_{\mathrm{law}}\,|\Psi\rangle\Big|. \tag{7.251}$$
A convenient sufficient condition is
$$\Delta U_{\mathrm{law}} = \exp\Big(i\{\,\hat{Q}_{\mathrm{BRST}},\,\Delta\Psi\,\}\Big), \tag{7.252}$$
for which $\Delta U_{\mathrm{law}} = \mathbf{1}$ in BRST cohomology by the Nielsen identities (cf. Section 7.5), and hence
$$\mathcal{V}_{\mathrm{law}} = 1 \qquad \text{on } \mathcal{H}_{\mathrm{phys}} \text{ (plateau)}. \tag{7.253}$$
More generally, any $U_b^{(\mathrm{law})}$ satisfying eqs. (7.245) and (7.246) yields isospectral branches and identical on-shell S-matrix elements (equivalence theorem), which again implies eq. (7.253) for Quadrant II.

Anomaly caveat. If the symmetry or BRST structure were anomalous in the band of interest, eq. (7.245) would fail and a genuine law-channel contrast could appear; such cases are excluded by the admissibility checks in Section 7.4.5.

1.3 Law visibility and plateau protection

With a symmetric calibration state ρ_0 (BRST-closed, ghost number 0) the *law*-channel visibility is read from the ancilla coherence after band projection (Section 7.4.1 and eq. (7.115)):

$$\Lambda_{\mathrm{law}}(\lambda_{\mathrm{UV}},\ell) = \frac{\left|\mathrm{Tr}_{\mathcal{H}_{\mathrm{phys}}}\Big(P_x(\lambda_{\mathrm{UV}})\,K_{\mathrm{IR}}^{(\ell)}\,U_1^{(\mathrm{law})\dagger}U_2^{(\mathrm{law})}\,\rho_0\Big)\right|}{\mathrm{Tr}_{\mathcal{H}_{\mathrm{phys}}}\Big(P_x(\lambda_{\mathrm{UV}})\,K_{\mathrm{IR}}^{(\ell)}\,\rho_0\Big)}\,. \tag{7.254}$$

Let $\Delta U_{\mathrm{law}} := U_1^{(\mathrm{law})\dagger}U_2^{(\mathrm{law})}$ (cf. eq. (7.249)). Using the commutation conditions from §1.2,

$$[\Delta U_{\mathrm{law}},\,\hat{Q}_{\mathrm{BRST}}] = [\Delta U_{\mathrm{law}},\,\hat{H}_0] = [\Delta U_{\mathrm{law}},\,\mathsf{J}_\alpha] = [\Delta U_{\mathrm{law}},\,P_x] = [\Delta U_{\mathrm{law}},\,K_{\mathrm{IR}}^{(\ell)}] = 0, \tag{7.255}$$

together with cyclicity of the trace on $\mathcal{H}_{\mathrm{phys}}$, the numerator of eq. (7.254) is invariant under presentation moves. In particular, if

$$\Delta U_{\mathrm{law}} = \exp\Big(i\{\hat{Q}_{\mathrm{BRST}},\eta_{\mathrm{law}}\}\Big) = \exp\Big(i\,\zeta\eta_{\mathrm{law}}\Big), \tag{7.256}$$

with $\mathfrak{g}(\eta_{\mathrm{law}}) = -1$ (hence $\mathfrak{g}(\zeta\eta_{\mathrm{law}}) = 0$), then $\Delta U_{\mathrm{law}} = \mathbf{1}$ in BRST cohomology (Nielsen identities; Section 7.5), and

$$\Lambda_{\mathrm{law}}(\lambda_{\mathrm{UV}},\ell) = 1 \qquad \text{(ideal presentation invariance, certified plateau).} \tag{7.257}$$

The same conclusion holds for background-field reparametrisations and on-shell–equivalent field redefinitions that satisfy eqs. (7.245) and (7.246) (equivalence theorem).

Plateau protection and tolerances. On a finite UV/IR window, residual regulator placement and slow parameter drift introduce controlled deviations. A convenient operational budget is

$$1-\Lambda_{\mathrm{law}} = \mathcal{O}\Big(\|[U_b^{(\mathrm{law})},P_x]\| + \|[U_b^{(\mathrm{law})},K_{\mathrm{IR}}^{(\ell)}]\| + \Big\|\nabla_{(\ln\lambda_{\mathrm{UV}},\ln\ell)}\mathcal{O}_{\mathrm{phys}}\Big\| + \text{curvatures}\Big), \tag{7.258}$$

so that the certified bounds of eq. (7.137) and Section 7.4.5 yield

$$\Lambda_{\mathrm{law}}(\lambda_{\mathrm{UV}},\ell) = 1 + \mathcal{O}(\varepsilon_{\mathrm{plat}}), \tag{7.259}$$

where $\varepsilon_{\mathrm{plat}}$ encodes the admissible gradients/oscillations/curvatures across the $(\ln\lambda_{\mathrm{UV}},\ln\ell)$ window. In this regime the Slavnov–Taylor/Ward/Nielsen identities (Section 7.5) ensure gauge-parameter independence of on-shell observables, and any branch dependence from $U_b^{(\mathrm{law})}$ is cohomologically trivial ([117, 442]).

Certification protocol (operational checks). To diagnose and keep $|\Lambda_{\mathrm{law}}| \approx 1$ on the plateau:

1. **Branch-swap invariance:** verify Λ_{law} is unchanged under $U_1^{(\mathrm{law})} \leftrightarrow U_2^{(\mathrm{law})}$.
2. **Gauge scan (Nielsen test):** vary the gauge parameter ζ in $\{\hat{Q}_{\mathrm{BRST}},\Psi_\zeta\}$; check
$$\partial_\zeta\Lambda_{\mathrm{law}} = \mathcal{O}(\varepsilon_{\mathrm{plat}}). \tag{7.260}$$
3. **Regulator order test:** permute $P_x(\lambda_{\mathrm{UV}})$, $K_{\mathrm{IR}}^{(\ell)}$, Π_{T}; confirm changes are within $\mathcal{O}(\varepsilon_{\mathrm{plat}})$.
4. **Flatness check:** ensure $\partial_{\ln\lambda_{\mathrm{UV}}}\Lambda_{\mathrm{law}}$ and $\partial_{\ln\ell}\Lambda_{\mathrm{law}}$ satisfy the plateau bounds of eq. (7.137).

Exclusion clauses. A persistent deficit $|\Lambda_{\rm law}| < 1-\delta$ with $\delta \gg \varepsilon_{\rm plat}$ signals a violation of the Quadrant II premises: (i) departure from the BRST-closed symmetric class (e.g., anomaly in the band of interest), or (ii) contamination by state-channel operations. Both are disallowed at this step (Sections 7.4 and 7.4.4; [117, 442]).

Lemma 7.334 (Plateau presentation invariance)**.** *Assume the branch-difference unitary* $\Delta U_{\rm law}$ *obeys the commutators of eq.* (7.255) *and admits a BRST-exact presentation* $\Delta U_{\rm law} = \exp\big(i\{\hat{Q}_{\rm BRST},\, \Delta\Psi\}\big)$*. Then for any BRST-closed calibration state* ρ_0 *of ghost number 0,* $\Lambda_{\rm law}(\lambda_{\rm UV}, \ell) = 1$ *on the certified plateau, i.e. eq.* (7.257) *holds.*

Proof. Start from eq. (7.254) and write the numerator $N := \mathrm{Tr}_{\mathcal{H}_{\rm phys}}\big(P_x K_{\rm IR}^{(\ell)}\, \Delta U_{\rm law}\, \rho_0\big)$. By eq. (7.255) and eq. (7.244) one has $P_x K_{\rm IR}^{(\ell)}\, \Delta U_{\rm law} = \Delta U_{\rm law}\, P_x K_{\rm IR}^{(\ell)}$. Consider $\Delta U_{\rm law}(s) := \exp\big(is\{\hat{Q}_{\rm BRST},\, \Delta\Psi\}\big)$ with $s \in [0,1]$ and $N(s) := \mathrm{Tr}_{\mathcal{H}_{\rm phys}}\big(P_x K_{\rm IR}^{(\ell)}\, \Delta U_{\rm law}(s)\, \rho_0\big)$. Then

$$\frac{dN}{ds} = i\ \mathrm{Tr}_{\mathcal{H}_{\rm phys}}\Big(P_x K_{\rm IR}^{(\ell)}\, \{\hat{Q}_{\rm BRST},\, \Delta\Psi\}\, \Delta U_{\rm law}(s)\, \rho_0\Big) = 0$$

by the graded-cyclicity identity eq. (7.331) (since $P_x K_{\rm IR}^{(\ell)}$ and ρ_0 are BRST-closed). Thus $N(1) = N(0) = \mathrm{Tr}_{\mathcal{H}_{\rm phys}}\big(P_x K_{\rm IR}^{(\ell)}\, \rho_0\big)$, and $\Lambda_{\rm law} = 1$ follows from eq. (7.254). □

Corollary 7.335 (Equivalence moves)**.** *Background-field reparametrisations and on–shell–equivalent local field redefinitions that satisfy the commutator conditions eqs.* (7.245) *and* (7.246) *leave* $N = \mathrm{Tr}_{\mathcal{H}_{\rm phys}}\big(P_x K_{\rm IR}^{(\ell)}\, \Delta U_{\rm law}\, \rho_0\big)$ *invariant. Either they can be cast in the BRST-exact form eq.* (7.256) *on the physical band, or they preserve the band–projected physical algebra so that eq.* (7.331) *applies after reduction to cohomology. Consequently, by Lemma 7.334 one has* $\Lambda_{\rm law}(\lambda_{\rm UV}, \ell) = 1$ *on the certified plateau, and together with eq.* (7.259) *this gives* $\Lambda_{\rm law} = 1 + \mathcal{O}(\varepsilon_{\rm plat})$ *off the ideal limit.*

1.4 Quadrant preservation under overlay

The quadrant predicate is defined by the dyadic status of *law* vs. *state* together with their operational visibilities (table 6.1, Corollary 6.232, and Section 6.11.6). Since $\mathcal{U}_{\rm law}$ of §1.2 leaves $\hat{H}_0(t; \lambda_{\rm UV}, \ell)$, the BRST charge, and the G-action unchanged—while commuting with the UV/IR/transverse projectors—no explicit symmetry breaking is introduced and no emergent-law flow is triggered in the law channel (Sections 7.3 and 7.5). In particular, the kernels $\mathcal{K}, \mathcal{M}^2$ that define $\hat{H}_0$ remain G-invariant and identical across branches (eqs. (7.241) and (7.243)), so the overlay cannot move the system out of Quadrant II into QIV (explicit breaking) or QIII (law deformation).

Overlay acts trivially on the law kernel. From eqs. (7.245) and (7.246), the adjoint action of each branch operator leaves the band-projected Hamiltonian fixed:

$$\mathrm{Ad}_{U_b^{(\rm law)}}\Big(\hat{H}_0(t; \lambda_{\rm UV}, \ell)\Big) = \hat{H}_0(t; \lambda_{\rm UV}, \ell), \qquad \mathrm{Ad}_{U_b^{(\rm law)}}\Big(\mathbb{H}^{(\lambda_{\rm UV}, \ell)}\Big) = \mathbb{H}^{(\lambda_{\rm UV}, \ell)}\,, \tag{7.261}$$

hence the spectral data (poles, residues on $\mathcal{H}_{\rm phys}$) and the on-shell algebra generated by the $\Pi_{\rm T}$-projected blocks are unchanged. Together with eq. (7.248), this means the overlay realizes

an automorphism of the band-projected physical algebra that is trivial in BRST cohomology (Nielsen).

Proposition 7.336 (Quadrant II is preserved under law overlay)**.** *Let the branch operators $U_b^{(\mathrm{law})}$ satisfy eqs. (7.245) and (7.246). If the pre-overlay configuration lies in Quadrant II (law symmetric, state possibly asymmetric), then the post-overlay configuration also lies in Quadrant II.*

Proof. **Law symmetry.** By eq. (7.261) the kernels $\mathcal{K}, \mathcal{M}^2$ remain G-invariant and identical on both branches, and $\hat{Q}_{\mathrm{BRST}}$ is preserved. Therefore no explicit breaking or law deformation occurs.

Law visibility. On certified plateaus, eqs. (7.257) and (7.259) give $\Lambda_{\mathrm{law}} = 1 + \mathcal{O}(\varepsilon_{\mathrm{plat}})$, independent of the presentation (Nielsen identities, Section 7.5). Hence the law-channel signature remains that of a symmetric law.

State channel. The overlay acts only on the law presentation; state preparation and the associated order parameters/Goldstone diagnostics (defined in Section 7.4.1 and used later in the SSB analysis) are unaffected. Thus any volume-scaling loss of state visibility persists, realizing the *state-asymmetric / law-symmetric* cell. □

Corollary 7.337 (No drift to QIII/QIV)**.** *Under the hypotheses of Proposition 7.336, transitions to QIII (law deformation) or QIV (explicit breaking) would require failure of eq. (7.245) or eq. (7.246) (or a BRST/gauge anomaly in the band). These are excluded by construction and by the admissibility checks of Section 7.4.5. Hence the overlay cannot induce QII→QIII/QIV drift.*

Robustness on finite windows. On finite UV/IR windows, any deviation is bounded by the plateau budget eq. (7.258), so that

$$1 - \Lambda_{\mathrm{law}} = \mathcal{O}(\varepsilon_{\mathrm{plat}})\,, \qquad \text{with } \varepsilon_{\mathrm{plat}} \text{ set by eq. (7.137).} \tag{7.262}$$

Thus quadrant membership is stable under overlay within certified tolerances.

Operational readout. The global map's Quadrant II signature—$|\Lambda_{\mathrm{law}}| \approx 1$ with SSB appearing solely in the state channel—follows from eqs. (7.257) and (7.259) and the interferometric readout in Section 7.4.1. This matches the standard SSB phenomenology in the state sector while keeping the microscopic law untouched ([91, 82]).

Lemma 7.338 (Converse: significant law-visibility deficit implies hypothesis failure)**.** *Assume the branch operators obey eqs. (7.245) and (7.246), the exact-identity sector is anomaly-free (Section 7.5), and the UV/IR window satisfies the plateau bounds of eq. (7.137). Then by eqs. (7.257) and (7.259) one has $|1 - \Lambda_{\mathrm{law}}(\lambda_{\mathrm{UV}}, \ell)| = \mathcal{O}(\varepsilon_{\mathrm{plat}})$ on any certified window. Consequently, if on such a certified window one observes a deficit $|1 - \Lambda_{\mathrm{law}}| \geq \delta$ with $\delta \gg \varepsilon_{\mathrm{plat}}$, at least one hypothesis must be false—i.e. either a commutator in eqs. (7.245) and (7.246) fails, or the exact-identity/BRST closure fails (anomaly), or the window is not plateau-certified (violation of eq. (7.137)).*

Proof. Under the stated hypotheses, eq. (7.257) gives the ideal limit $\Lambda_{\mathrm{law}} = 1$, and eq. (7.259) refines this to the finite-window estimate $1 - \Lambda_{\mathrm{law}} = \mathcal{O}(\varepsilon_{\mathrm{plat}})$. A measured deficit with $\delta \gg \varepsilon_{\mathrm{plat}}$ contradicts these conclusions; by contrapositive, at least one hypothesis is violated. □

Operational diagnostic. When the premise of Lemma 7.338 fails in data, localize the cause via the protocol in §1.3: (i) branch-swap invariance; (ii) Nielsen slope $\partial_\zeta \Lambda_{\mathrm{law}}$ (eq. (7.260)); (iii) regulator-order permutations; (iv) flatness checks against eq. (7.137). Persistent violations witness either law-channel contamination or loss of plateau certification.

1.5 Implementation checklist (acceptance tests)

For any operator $\mathcal{A}$ acting on the full Hilbert space, we evaluate it on the band-projected physical sector with the induced operator norm

$$\|\mathcal{A}\|_{\mathrm{phys}} := \Big\| \, \mathsf{P}_{\mathrm{phys}} \, P_x(\lambda_{\mathrm{UV}}) \, K_{\mathrm{IR}}^{(\ell)} \, \mathcal{A} \, K_{\mathrm{IR}}^{(\ell)} \, P_x(\lambda_{\mathrm{UV}}) \, \mathsf{P}_{\mathrm{phys}} \, \Big\| \,, \tag{7.263}$$

where $\|\cdot\|$ is the operator norm on $\mathcal{H}_{\mathrm{phys}}$ and $\mathsf{P}_{\mathrm{phys}}$ is the BRST projector (cf. §1.1).

- **Commutators (law/BRST/G preservation).** Verify the exact commutators eq. (7.245) on the band; when implemented numerically/operationally, enforce the quantitative tolerance

$$\Big\|[U_b^{(\mathrm{law})}, \hat{H}_0]\Big\|_{\mathrm{phys}} + \Big\|[U_b^{(\mathrm{law})}, \hat{Q}_{\mathrm{BRST}}]\Big\|_{\mathrm{phys}} + \sum_\alpha \Big\|[U_b^{(\mathrm{law})}, \mathsf{J}_\alpha]\Big\|_{\mathrm{phys}} \;\le\; \varepsilon_{\mathrm{comm}} \,, \tag{7.264}$$

  and certify that the branch difference ΔU_{law} is a presentation move:

$$\Delta U_{\mathrm{law}} = \exp\Big(i\{\hat{Q}_{\mathrm{BRST}}, \eta_{\mathrm{law}}\}\Big) = \exp\Big(i\,\zeta \eta_{\mathrm{law}}\Big) \quad \text{(cf. eqs. (7.249) and (7.256)).}$$

  This guarantees cohomological triviality on $\mathcal{H}_{\mathrm{phys}}$ (Section 7.5).

- **Band compatibility (projector commutators).** Enforces regulator–order invariance on certified plateaus (Section 7.3 and eq. (7.258))

$$\begin{aligned} \Big\|[P_x(\lambda_{\mathrm{UV}}),\, U_b^{(\mathrm{law})}]\Big\|_{\mathrm{phys}} \;+\; & \Big\|[K_{\mathrm{IR}}^{(\ell)},\, U_b^{(\mathrm{law})}]\Big\|_{\mathrm{phys}} \\ & + \Big\|[\Pi_{\mathrm{T}},\, U_b^{(\mathrm{law})}]\Big\|_{\mathrm{phys}} \le \varepsilon_{\mathrm{reg}} \,. \end{aligned} \tag{7.265}$$

- **Exact-identity sector (ST/Ward/Nielsen).** Enforce Slavnov–Taylor and Ward closures on the band (cf. Section 7.5); aggregate their residuals into an identity defect

$$\Delta_{\mathrm{id}}(\lambda_{\mathrm{UV}}, \ell) := \Delta_{\mathrm{ST}} + \Delta_{\mathrm{Ward}} \;\le\; \varepsilon_{\mathrm{id}} \,, \tag{7.266}$$

  and verify Nielsen independence of the law visibility via the gauge scan $\partial_\zeta \Lambda_{\mathrm{law}} = \mathcal{O}(\varepsilon_{\mathrm{plat}})$ (eq. (7.260)). Any violation triggers the anomaly filter and cohomological checks (cf. §1.3 Exclusion clauses).

- **Plateau certification (flatness and robustness).** Locate windows $(\lambda_{\mathrm{UV}}, \ell)$ for which the gradient/oscillation/curvature bounds of Λ_{law} satisfy eq. (7.137), and regulator robustness holds (cf. eq. (7.258)). Record the certified budget $\varepsilon_{\mathrm{plat}}$ and verify

$$\Big|1 - \Lambda_{\mathrm{law}}(\lambda_{\mathrm{UV}}, \ell)\Big| \;\le\; C_{\mathrm{plat}} \, \varepsilon_{\mathrm{plat}} \,, \tag{7.267}$$

  for a protocol-dependent constant $C_{\mathrm{plat}} \sim \mathcal{O}(1)$ (cf. eq. (7.259)).

- **Swap and ordering invariances (sanity checks).** Test branch-swap invariance and regulator ordering; accept if

$$\begin{aligned}\left|\Lambda_{\rm law}[U_1^{(\rm law)},U_2^{(\rm law)}]-\Lambda_{\rm law}[U_2^{(\rm law)},U_1^{(\rm law)}]\right|&\le\varepsilon_{\rm plat}\,,\\ \left|\Lambda_{\rm law}[\ldots P_xK_{\rm IR}\Pi_{\rm T}\ldots]-\Lambda_{\rm law}[\ldots K_{\rm IR}P_x\Pi_{\rm T}\ldots]\right|&\le\varepsilon_{\rm plat}\,,\end{aligned}\tag{7.268}$$

  consistent with eqs. (7.246) and (7.258).

- **Spectral health (physical band).** Check positivity/hyperbolicity of the $\Pi_{\rm T}$-projected Hessian blocks defining $\hat{H}_0$ on the certified window (cf. eqs. (7.241) and (7.243)). Reject if spurious longitudinal/ghost leakage is detected (quartet mechanism failure), or if poles/residues drift beyond plateau tolerances.

- **Converse trigger (escalation).** If a statistically significant deficit $|1-\Lambda_{\rm law}|\ge\delta$ with $\delta\gg\varepsilon_{\rm plat}$ is observed on a certified window, invoke the converse Lemma Lemma 7.338 and run the localization protocol of §1.3 (branch swap, Nielsen slope, regulator permutations, flatness checks).

1.6 Failure modes and diagnostics

Diagnostic decomposition (quantitative triage). Define the law-channel deficit

$$\delta_{\rm law}(\lambda_{\rm UV},\ell):=1-\Lambda_{\rm law}(\lambda_{\rm UV},\ell)\quad(\ge0).\tag{7.269}$$

On a certified window, eqs. (7.258) and (7.259) imply the schematic bound

$$\begin{aligned}\delta_{\rm law}\ \lesssim\ & C_{\rm comm}\Big(\big\|[U_b^{(\rm law)},\hat{H}_0]\big\|_{\rm phys}+\big\|[U_b^{(\rm law)},\hat{Q}_{\rm BRST}]\big\|_{\rm phys}+\sum_\alpha\big\|[U_b^{(\rm law)},\mathsf{J}_\alpha]\big\|_{\rm phys}\Big)\\ &+C_{\rm reg}\sum_{X\in\{P_x,K_{\rm IR}^{(\ell)},\Pi_{\rm T}\}}\big\|[X,U_b^{(\rm law)}]\big\|_{\rm phys}\\ &+C_{\rm id}\big(\Delta_{\rm ST}+\Delta_{\rm Ward}\big)+C_\zeta\,|\partial_\zeta\Lambda_{\rm law}|+C_{\rm plat}\,\varepsilon_{\rm plat}+C_{\rm inst}\,\varepsilon_{\rm inst}\quad(+\text{ higher orders}),\end{aligned}\tag{7.270}$$

where the norms/defects are those of eqs. (7.260) and (7.263) to (7.266), and $\varepsilon_{\rm inst}$ is set by the instrumental budget in eq. (7.271) below. This consolidates the acceptance tests of §1.5 into a single triage inequality.

- $|\Lambda_{\rm law}|<1$ **off-plateau.** The typical cause is residual regulator placement and/or an insufficiently flat UV/IR scan. Re-certify the window $(\lambda_{\rm UV},\ell)$ until the gradient/curvature/oscillation bounds of eq. (7.137) hold and regulator robustness is satisfied (Sections 7.3.3, 7.3.7 and 7.4.5). *Why:* within certified windows one has $\Lambda_{\rm law}=1+\mathcal{O}(\varepsilon_{\rm plat})$ up to regulator tolerances (eq. (7.259)), so deficits point to a plateau failure rather than genuine law incoherence.

- **Persistent** $|\Lambda_{\rm law}|<1$ **on-plateau.** This violates the Quadrant II premise (symmetric law on both branches). Perform the following triage (each line maps to a term in eq. (7.270)):

(i) *Branch commutators (§1.5).* Re-check eq. (7.245); certify ΔU_{law} is a presentation move

$$\Delta U_{\mathrm{law}} = \exp\Big(i\{\hat{Q}_{\mathrm{BRST}}, \eta_{\mathrm{law}}\}\Big) = \exp\Big(i\,\zeta\eta_{\mathrm{law}}\Big)$$

(cf. eqs. (7.249) and (7.256)). Any failure signals illicit law deformation or explicit breaking (cf. Proposition 7.336).

(ii) *Identity ledger.* Enforce ST/Ward closure on shell (Section 7.5) and bound the aggregate defect by eq. (7.266). Non-closure indicates explicit breaking or off-shell contamination.

(iii) *Anomaly filters.* Re-apply the admissibility constraints of Section 7.4.5 (gauge/mixed/-grav: trivial cohomology class; Wess–Zumino consistency). Any nontrivial class in the anomaly cohomology explains $|\Lambda_{\mathrm{law}}| < 1$; reproject to a BRST-closed scheme and retest the ledger.

(iv) *Regulator swap/robustness.* Replace $(P_x, K_{\mathrm{IR}}^{(\ell)})$ by an admissible pair; require eq. (7.265) and stability under permutations on the certified window (cf. eq. (7.268)). Deviations beyond $\varepsilon_{\mathrm{reg}}$ indicate a UV/IR mismatch rather than true law loss.

(v) *Health checks (spectral).* Re-confirm positivity/hyperbolicity of the Π_{T}–projected blocks defining $\hat{H}_0$ on the window (eqs. (7.241) and (7.243)). Pathologies (e.g. spurious longitudinal/ghost leakage, pole drift) can leak into the coherence functional and mimic law loss.

- **Gauge-parameter drift.** A nonzero Nielsen slope $\partial_\zeta \Lambda_{\mathrm{law}}$ betrays off-shell evaluation or a mis-projection. Restore the on-shell conditions used in the band-projected coherence definition (eq. (7.115)), and re-verify ST/Nielsen closures (Section 7.5 and eq. (7.260)); on shell, visibilities are gauge-parameter independent. Also re-check eq. (7.244) to ensure BRST-closure of the band operations.

- **Instrumental artefacts (false deficits).** Apparent suppressions can arise from shot noise and phase miscalibration. Track an explicit instrumental budget

$$\varepsilon_{\mathrm{inst}} \;\equiv\; \sigma_{\mathrm{shot}} + \sigma_\phi + \sigma_{\mathrm{sys}}, \tag{7.271}$$

and require the observed deviation to clear this floor before attributing physics: $|1 - \Lambda_{\mathrm{law}}| \;\gtrsim\; C_{\mathrm{inst}}\,\varepsilon_{\mathrm{inst}}$. Audit both the total uncertainty $\delta\Lambda_\alpha$ and the phase contribution $(\delta\Lambda_\alpha)_\phi$ in the readout pipeline (compare the swap/order tests in eq. (7.268)).

Converse trigger (escalation rule). On a certified window, if a statistically significant deficit $|1 - \Lambda_{\mathrm{law}}| \geq \delta$ with $\delta \gg \varepsilon_{\mathrm{plat}}$ is observed, invoke the converse lemma Lemma 7.338 and execute the localization protocol of §1.3: branch swap, Nielsen slope, regulator permutations, and flatness checks against eq. (7.137).

1.7 Summary of Step 1

We *freeze the law*: fix the band–projected Hamiltonian $\hat{H}_0(t;\lambda_{\mathrm{UV}},\ell)$ as in eqs. (7.241) and (7.243), and restrict branch manipulations to symmetry–preserving *presentations*—BRST–exact similarities, background–field reparametrisations, and on–shell–equivalent field redefinitions—implemented by $U_b^{(\mathrm{law})}$ satisfying the commutators eqs. (7.245) and (7.246) (cf. eq. (7.247)).

On certified plateaus, the law–channel visibility is presentation–invariant: $\Lambda_{\mathrm{law}} = 1$ in the ideal limit (eq. (7.257)) and $\Lambda_{\mathrm{law}} = 1 + \mathcal{O}(\varepsilon_{\mathrm{plat}})$ on finite windows (eq. (7.259)), with gauge–parameter independence monitored by the Nielsen slope bound eq. (7.260). The overlay acts trivially on the law kernel (eq. (7.261)) and *preserves* Quadrant II (Proposition 7.336 and Corollary 7.337). Acceptance tests eqs. (7.263) to (7.265) and (7.268) and the robustness/plateau bounds eqs. (7.259) and (7.262) certify this control; identity/closure checks are tracked in the exact-identity sector (Section 7.5). Any significant deficit triggers the converse lemma Lemma 7.338 and the localization protocol of §1.3.

With the microscopic law coherent and quadrant membership locked, **Step 2** prepares an asymmetric state on the vacuum manifold to induce genuine SSB in the *state* channel while the law remains symmetric (see the quadratic/Hessian diagnostics in Section 7.2.5 and eqs. (7.13) and (7.15); [442, 91, 82, 241]).

7.7.2 Prepare an Asymmetric State on the Vacuum Manifold

Aim. Keep the *law* symmetric as in Section 7.7.1, and prepare an *asymmetric* state that selects a point on the vacuum manifold $\mathcal{M}_{\mathrm{vac}} \cong G/H$ (Sections 7.2 and 7.5). Operationally, all manipulations occur *on the band* with the same UV/IR controls (Section 7.3), so that any deviation seen by the interferometer is unambiguously attributable to the *state* channel (cf. Section 7.4).

2.1 Vacuum manifold and selection by an infinitesimal seed

Spontaneous breaking is characterized by a nonzero gauge-invariant order parameter extracted from the prepared state ω (Section 7.5):

$$\varphi(x) \equiv \langle \Phi(x) \rangle_\omega, \qquad \text{or more generally } \langle \mathcal{O}(x) \rangle_\omega \neq 0, \\ \text{for some BRST-closed } \mathcal{O} \quad ([\hat{Q}_{\mathrm{BRST}}, \mathcal{O}] = 0). \tag{7.272}$$

To select a definite vacuum in $\mathcal{M}_{\mathrm{vac}}$, introduce an infinitesimal *seed* that tilts the potential along a coset direction $X \in \mathfrak{g}/\mathfrak{h}$. The seed couples to a gauge-invariant, BRST-closed operator $\mathcal{O}_X$ transforming as X under the *global* group G (no literal gauge-SSB; [117, 442]):

$$\Delta\mathcal{H}_{\mathrm{seed}}(\varepsilon) = -\,\varepsilon \int d^d x \; f_X(x)\, \mathcal{O}_X(x), \qquad \varepsilon > 0, \quad \varepsilon \to 0^+, \tag{7.273}$$

with f_X smooth, compactly supported, and compatible with the band projectors:

$$[\hat{Q}_{\mathrm{BRST}}, \mathcal{O}_X] = 0, \qquad \Big[P_x(\lambda_{\mathrm{UV}}), \mathcal{O}_X\Big] = \Big[K_{\mathrm{IR}}^{(\ell)}, \mathcal{O}_X\Big] = 0 \;\; \text{on } \mathcal{H}_{\mathrm{phys}} \;\; (\textit{Section 7.3}). \tag{7.274}$$

The prepared vacuum state (on branch $b \in \{1,2\}$) is defined by the ordered limits

$$\omega^{(b)}(\,\cdot\,) = \lim_{(\lambda_{\mathrm{UV}},\ell)\in\text{plateau}} \lim_{V\to\infty} \lim_{\varepsilon\to 0^+} \omega^{(b)}_{\varepsilon,V;\,\lambda_{\mathrm{UV}},\ell}(\,\cdot\,), \tag{7.275}$$

where $(\lambda_{\mathrm{UV}},\ell)$ lie in a certified plateau (Sections 7.3 and 7.4.5). The seed *does not* alter the symmetric $\hat{H}_0$ used by the law-channel interferometer; it is strictly a state-preparation device (cf. Section 7.7.1).

Two equivalent preparation routes (law unchanged). Let ρ_0 be the symmetric calibration state (ghost number 0).

1. **Imaginary-time (or adiabatic) seeding.** Prepare

$$\rho^{(b)}_{\varepsilon,V;\,\lambda_{\mathrm{UV}},\ell} = \frac{P_x K^{(\ell)}_{\mathrm{IR}}\; e^{-\beta(\hat{H}_0-\varepsilon J^{(b)}_X)}\; P_x K^{(\ell)}_{\mathrm{IR}}}{\mathrm{Tr}_{\mathcal{H}_{\mathrm{phys}}}\Big(P_x K^{(\ell)}_{\mathrm{IR}}\; e^{-\beta(\hat{H}_0-\varepsilon J^{(b)}_X)}\Big)}, \qquad J^{(b)}_X := \int d^d x\; f^{(b)}_X(x)\,\mathcal{O}_X(x), \tag{7.276}$$

   with β large (ground-state projection) or via an adiabatic real-time ramp that prepares the same state; then remove the seed and take $\varepsilon\to 0^+$ after $V\to\infty$ on the plateau.

2. **Unitary preparation.** Apply a BRST-physical unitary that *only* acts in the state channel,

$$\begin{aligned} \rho^{(b)}_{\varepsilon,V;\,\lambda_{\mathrm{UV}},\ell} &= U^{(b)}_{\mathrm{prep}}(\varepsilon)\,\Big(P_x K^{(\ell)}_{\mathrm{IR}}\;\rho_0\;P_x K^{(\ell)}_{\mathrm{IR}}\Big)\,U^{(b)\dagger}_{\mathrm{prep}}(\varepsilon), \\ U^{(b)}_{\mathrm{prep}}(\varepsilon) &= \mathcal{T}\exp\Big[-i\int dt\;\Delta\mathcal{H}^{(b)}_{\mathrm{seed}}(t)\Big], \end{aligned} \tag{7.277}$$

   with $[\hat{Q}_{\mathrm{BRST}}, U^{(b)}_{\mathrm{prep}}] = 0$ (BRST-closed channel), and the same ordered limits as in eq. (7.275).

Both routes are equivalent for on-shell observables by the equivalence theorem and Nielsen identities (Section 7.5; [117, 442]).

Band-compatibility and BRST admissibility. To ensure Quadrant-II integrity:

- *Gauge/BRST:* $\mathcal{O}_X$ must be BRST-closed (ghost number 0) so that $[\hat{Q}_{\mathrm{BRST}}, \Delta\mathcal{H}_{\mathrm{seed}}] = 0$ and the prepared state remains in $\mathcal{H}_{\mathrm{phys}}$ (Section 7.2).
- *Regulators:* $[P_x(\lambda_{\mathrm{UV}}), \Delta\mathcal{H}_{\mathrm{seed}}] = [K^{(\ell)}_{\mathrm{IR}}, \Delta\mathcal{H}_{\mathrm{seed}}] = 0$ on the plateau, avoiding spurious law-channel contamination (Sections 7.3 and 7.4.5).
- *Global vs. gauge.* The coset $X \in \mathfrak{g}/\mathfrak{h}$ refers to a *global* symmetry direction; gauge directions are handled cohomologically (no literal gauge-SSB; longitudinal/ghost quartets decouple) ([117, 442]).

Local coordinates on $\mathcal{M}_{\rm vac}$ and mode split. Once a vacuum $(\bar\Phi,\bar\Theta)\in\mathcal{M}_{\rm vac}$ is selected, parameterize fluctuations by NG coordinates π_a along G/H and gapped coordinates σ_i transverse to G/H:

$$\Phi(x)=\bar\Phi+\sum_a \pi_a(x)\,(T_a\cdot\bar\Phi)+\sum_i \sigma_i(x)\,n_i+\cdots,\qquad \Theta(x)=\bar\Theta+\cdots, \tag{7.278}$$

with T_a broken generators and n_i orthonormal radial/mixed directions. This NG/gapped split is consistent with the band Hessian blocks of Section 7.2.5 (cf. $\Pi_{\rm T}$-projected $\mathbb{H}^{(\lambda_{\rm UV},\ell)}$ in eqs. (7.241) and (7.243)). NG kinematics can then be read off via phase-twist interferometry along closed loops in G/H (Section 6.11.6).

Ordered limits and cluster properties. The order $\varepsilon\to 0^+$ *after* $V\to\infty$ on a certified plateau suppresses domain averaging and yields a definite vacuum sector (standard SSB construction; [91, 82]). Plateau flatness guarantees gauge-parameter independence of the resulting on-shell observables (Nielsen), and BRST cohomology removes unphysical sectors (Sections 7.2 and 7.5).

Branch usage. For later state-channel diagnostics, one may take seeds $\pm X$ on the two ancilla branches to prepare distinct vacua related by G, or the *same* vacuum on both branches for calibration. In both cases the *law* channel remains the symmetric baseline of Section 7.7.1; any visibility loss must therefore be attributed to the *state* channel (Sections 7.4 and 7.4.1).

Lemma 7.339 (Route equivalence for band–projected correlators)**.** *Consider a set of operators $\{\mathcal{O}_k\}_{k=1}^n$ that are BRST-closed, gauge-invariant, have ghost-number 0, and commute with the band projectors on $\mathcal{H}_{\rm phys}$. Define the band–projected n-point function on branch b by*

$$G_n^{(b)}[\{\mathcal{O}_k(x_k)\}]:=\frac{\mathrm{Tr}_{\mathcal{H}_{\rm phys}}\Big(P_x K_{\rm IR}^{(\ell)}\,\rho^{(b)}\,P_x K_{\rm IR}^{(\ell)}\,\prod_{k=1}^n\mathcal{O}_k(x_k)\Big)}{\mathrm{Tr}_{\mathcal{H}_{\rm phys}}\Big(P_x K_{\rm IR}^{(\ell)}\,\rho^{(b)}\Big)}. \tag{7.279}$$

Let $\rho^{(b)}$ be prepared either by imaginary–time seeding eq. (7.276) or by unitary seeding eq. (7.277). Then, under the ordered limits eq. (7.275) on a certified plateau and with the symmetric law fixed by eqs. (7.241) and (7.243),

$$\lim_{\text{plateau}}\lim_{V\to\infty}\lim_{\varepsilon\to 0^+} G_n^{(b)}\Big|_{\text{im. time}} = \lim_{\text{plateau}}\lim_{V\to\infty}\lim_{\varepsilon\to 0^+} G_n^{(b)}\Big|_{\text{unitary}}. \tag{7.280}$$

Proof. Step 1 (Generating functionals). Introduce a source J for the BRST–closed operator $\mathcal{O}_X$ used in eq. (7.273), and define band–projected generating functionals

$$\begin{aligned} Z_{\rm im}[J]&:=\mathrm{Tr}_{\mathcal{H}_{\rm phys}}\Big(P_x K_{\rm IR}^{(\ell)}\,e^{-\beta(\hat H_0-\varepsilon J_X)}\,P_x K_{\rm IR}^{(\ell)}\Big),\\ Z_{\rm uni}[J]&:=\mathrm{Tr}_{\mathcal{H}_{\rm phys}}\Big(P_x K_{\rm IR}^{(\ell)}\,U_{\rm prep}(\varepsilon;J)\,\rho_0\,U_{\rm prep}^\dagger(\varepsilon;J)\,P_x K_{\rm IR}^{(\ell)}\Big),\end{aligned} \tag{7.281}$$

with $J_X=\int d^dx\, f_X(x)\,\mathcal{O}_X(x)$ and $U_{\rm prep}=\mathcal{T}\exp[-i\int dt\,\Delta\mathcal{H}_{\rm seed}(t)]$ as in eq. (7.277). Insertion correlators eq. (7.279) are obtained by functional differentiation of $\ln Z$ w.r.t. J and then setting $J\to 0$.

Step 2 (Contour matching). Use the Schwinger–Keldysh contour with a vertical imaginary–time leg: the time–ordered evolution with U_{prep} on the real branches and the Euclidean weight $e^{-\beta(\hat{H}_0-\varepsilon J_X)}$ on the vertical leg represent the same contour–ordered object when (i) the Hamiltonian on all branches is the *same* $\hat{H}_0$ (law frozen; Section 7.7.1 and eq. (7.241)), (ii) the source couples to a BRST–closed, gauge–invariant $\mathcal{O}_X$ commuting with the band projectors eq. (7.274), and (iii) band operations commute with the symmetry/BRST generators eq. (7.244). Therefore,

$$Z_{\mathrm{im}}[J] \;=\; Z_{\mathrm{uni}}[J] \qquad \text{on the certified plateau,} \tag{7.282}$$

up to terms controlled by the plateau budget $\mathcal{O}(\varepsilon_{\mathrm{plat}})$ (eq. (7.258)).

Step 3 (Ordered limits and cohomology). Taking $\beta \to \infty$ (ground–state projection), $V \to \infty$, and $\varepsilon \to 0^+$ in the order eq. (7.275) selects a definite vacuum sector on $\mathcal{M}_{\mathrm{vac}}$ (no domain averaging). Because the source/operator are BRST–closed and the band operations commute with $\hat{Q}_{\mathrm{BRST}}$ eq. (7.244), cohomologically exact contributions cancel in the trace (cf. the identity eq. (7.331) used in the Step 1 lemma of §1.3), and Nielsen identities guarantee gauge–parameter independence on shell (Section 7.5). Functional differentiation of eq. (7.282) thus yields equality of all band–projected, on–shell BRST–closed correlators, establishing eq. (7.280). □

Corollary 7.340 (Profile–shape robustness)**.** *Let $f_X^{(1)}, f_X^{(2)}$ be two smooth, compactly supported seed profiles along the same coset direction $X \in \mathfrak{g}/\mathfrak{h}$ that commute with the band projectors as in eq. (7.274). Then the limits eq. (7.275) of the corresponding $G_n^{(b)}$ coincide up to $\mathcal{O}(\varepsilon_{\mathrm{plat}})$ and define the* same *vacuum sector on the plateau.*

Proof. Interpolate $f_X^{(s)} := (1-s)f_X^{(1)} + s f_X^{(2)}$ with $s \in [0,1]$, and let $J_X(s) = \int d^dx\, f_X^{(s)}(x)\, \mathcal{O}_X(x)$. By Lemma 7.339, the band–projected generating functionals for imaginary–time and unitary seeding coincide on the plateau (up to $\mathcal{O}(\varepsilon_{\mathrm{plat}})$), so it suffices to work with either route. Differentiating the (common) generating functional w.r.t. s inserts the BRST–closed operator $\int d^dx\, \delta f_X(x)\, \mathcal{O}_X(x)$, where $\delta f_X := f_X^{(2)} - f_X^{(1)}$. Because $f_X^{(s)}$ commute with the band projectors (eq. (7.274)) and $\mathcal{O}_X$ is BRST–closed and gauge–invariant, the derivative produces only plateau–suppressed terms; graded cyclicity and the cohomological identity eq. (7.331) remove BRST–exact contributions. Taking the ordered limits eq. (7.275) (which fix the vacuum sector and exclude domain averaging) gives $dG_n^{(b)}/ds = \mathcal{O}(\varepsilon_{\mathrm{plat}})$. Integrating in s yields the claim. □

Remark 7.341 (Scope)**.** The lemma applies to BRST–closed, ghost–number 0 insertions (physical operators) evaluated on the certified band. For gauge–variant or off–shell insertions, additional gauge–fixing artefacts can appear away from the plateau; these are filtered out by the acceptance tests of §1.5 and the plateau budget eq. (7.258).

2.2 Coset parameterisation and fluctuation fields

Choose a local coset chart with the CCWZ map

$$U(\pi(x)) = \exp\Big[\,\mathrm{i}\,\pi^a(x)\,T_a\Big], \qquad T_a \in \mathfrak{g}/\mathfrak{h}, \tag{7.283}$$

and write the field as a broken-phase fluctuation about the chosen vacuum vector v:

$$\Phi(x) = U(\pi(x))\Big(v + h(x)\Big), \tag{7.284}$$

where the NG coordinates π^a chart G/H and the gapped field h transforms in a (possibly reducible) linear representation of the unbroken subgroup H. In gauge theories, this parameterisation is naturally implemented in background-field gauges that preserve on-shell identities (cf. Section 7.5). BRST quartet decoupling guarantees that longitudinal/ghost sectors do not pollute $\mathcal{H}_{\rm phys}$ (Section 7.2.6).

For later kinematics it is convenient to introduce the Maurer–Cartan decomposition (with a gauge-covariant derivative D_μ when gauged)

$$-\,\mathrm{i}\,U^{-1}(\pi)\,D_\mu U(\pi) = E_\mu{}^a(\pi)\,T_a + \omega_\mu{}^i(\pi)\,H_i, \tag{7.285}$$

where $E_\mu{}^a$ is the coset "vielbein" and $\omega_\mu{}^i$ the H-connection. On the *band* (with $P_x(\lambda_{\rm UV})$, $K_{\rm IR}^{(\ell)}$, $\Pi_{\rm T}$ as in §1.1), the quadratic invariant built from π–derivatives takes the CCWZ form

$$\mathcal{L}_\pi^{(2)} = \frac{1}{2}\,(E_\mu{}^a\pi)\,G_{ab}(v)\,(E^{\mu\,b}\pi) \quad\longrightarrow\quad \frac{1}{2}\Big(\partial_\mu\pi\Big)^{\top}\mathsf{G}(v)\,\partial^\mu\pi + \mathcal{O}(\pi^3), \tag{7.286}$$

where $G_{ab}(v)$ (or $\mathsf{G}(v)$) is positive semidefinite on $\mathcal{H}_{\rm phys}$ by the positivity/hyperbolicity checks already certified on the symmetric band (eq. (7.15)).

Remarks. (i) In a BEH/Higgs realisation, some would-be NG coordinates reside in BRST quartets and are unphysical; they decouple on $\mathcal{H}_{\rm phys}$ by the quartet mechanism. (ii) All band operations commute with symmetry/BRST generators (eq. (7.244)); hence π/h observables extracted from eq. (7.284) are Nielsen-stable on certified plateaus.

2.3 Band Hessian around the broken background

Let Γ denote the band effective action. The (functional) band Hessian at the broken background φ is

$$\mathcal{H}_{\rm band}(x,y) := \left.\frac{\delta^2\Gamma}{\delta\Phi(x)\,\delta\Phi(y)}\right|_{\Phi=\varphi}, \tag{7.287}$$

and defines the inverse propagators for (π, h) after BRST projection to $\mathcal{H}_{\rm phys}$ (Section 7.2). In momentum space and on the band, the (π, h) block structure reads schematically

$$\mathbb{H}_{\rm broken}^{(\lambda_{\rm UV},\ell)}(p) = \begin{pmatrix} \mathcal{K}_{\pi\pi}(p) & \mathcal{K}_{\pi h}(p) \\ \mathcal{K}_{h\pi}(p) & \mathcal{K}_{hh}(p) \end{pmatrix} + \begin{pmatrix} \mathcal{M}^2_{\pi\pi} & \mathcal{M}^2_{\pi h} \\ \mathcal{M}^2_{h\pi} & \mathcal{M}^2_{hh} \end{pmatrix}, \tag{7.288}$$

with all objects implicitly projected by $P_x(\lambda_{\rm UV})K_{\rm IR}^{(\ell)}\Pi_{\rm T}$ and reduced to $\mathcal{H}_{\rm phys}$. Ward/Nielsen identities on the broken background imply the NG constraints

$$\mathcal{M}^2_{\pi\pi} = 0, \qquad \mathcal{K}_{\pi h}(p) = \mathcal{O}(p) \;\text{ and }\; \mathcal{K}_{\pi h}(0) = 0, \tag{7.289}$$

so that NG modes appear as massless poles (or are removed into BRST quartets in gauge realisations), while h retains a gapped mass operator $\mathcal{M}^2_{hh} \geq 0$ on $\mathcal{H}_{\rm phys}$. The principal symbol of $\mathbb{H}_{\rm broken}^{(\lambda_{\rm UV},\ell)}(p)$ remains hyperbolic and its physical mass operator non-negative by continuity from the symmetric band and the acceptance tests (eqs. (7.13) and (7.15)).

Goldstone Ward identity (on the band). Let $t_a\cdot\varphi$ be the variation of the background along a broken generator T_a. Then the band Hessian satisfies

$$\int d^d y\, \mathcal{H}_{\text{band}}(x,y)\,(T_a\cdot\varphi)(y) = 0 \qquad \text{on } \mathcal{H}_{\text{phys}}, \tag{7.290}$$

which is the on-band form of the Goldstone theorem (projected to the physical subspace); it follows from the exact identities at the symmetry point and the BRST/Nielsen machinery (Section 7.5). Equation eq. (7.290) enforces $\mathcal{M}^2_{\pi\pi} = 0$ and $\mathcal{K}_{\pi h}(0) = 0$ in eq. (7.289).

Plateau stability of the broken Hessian. Regulator stability and commutation of limits are checked on certified plateaus (Section 7.3). In particular, the band Hessian inherits the plateau robustness bounds,

$$\partial_{(\ln\lambda_{\text{UV}},\,\ln\ell)}\,\mathbb{H}^{(\lambda_{\text{UV}},\ell)}_{\text{broken}}(p) = \mathcal{O}(\varepsilon_{\text{plat}}), \tag{7.291}$$

so that the location of NG poles and the gapped spectrum of h are stable under admissible UV/IR scans (cf. eqs. (7.258) and (7.259)).

2.4 Broken Ward identities and NG structure (global G)

On the certified band (projected by $P_x(\lambda_{\text{UV}})K^{(\ell)}_{\text{IR}}\Pi_{\text{T}}$ and reduced to $\mathcal{H}_{\text{phys}}$), the Ward–Takahashi identity for a global current J^μ_a and any BRST-closed, ghost-number 0 operator $\mathcal{O}$ reads

$$\partial_\mu \left\langle T\, J^\mu_a(x)\,\mathcal{O}(y)\right\rangle_{\omega^{(b)}} = \mathrm{i}\,\delta^{(d)}(x-y)\left\langle \delta_a\mathcal{O}(y)\right\rangle_{\omega^{(b)}}, \tag{7.292}$$

with T time ordering on the band (the band projectors commute with J^μ_a and $\mathcal{O}$ on $\mathcal{H}_{\text{phys}}$, cf. eq. (7.244)). Fourier transforming eq. (7.292) and isolating the one-particle contribution gives the standard current–NG matrix element

$$\langle 0|\, J^\mu_a(0)\,|\,\pi^b(p)\rangle = \mathrm{i}\, f_{ab}\, p^\mu, \qquad f_{ab} = f_{ba} \geq 0 \text{ on } \mathcal{H}_{\text{phys}}, \tag{7.293}$$

which implies a massless pole in the current–current correlator and in any channel with nonvanishing overlap:

$$\mathrm{i}\int d^d x\, e^{\mathrm{i}k\cdot x}\left\langle T\, J^\mu_a(x) J^\nu_b(0)\right\rangle \supset \frac{f_{ac}f_{bc}\,k^\mu k^\nu}{k^2+\mathrm{i}0^+}, \qquad k^2\to 0\,. \tag{7.294}$$

Equivalently, the band Hessian at the broken background possesses zero modes along $T_a\cdot\varphi$ (Goldstone theorem on the band), cf. eqs. (7.289) and (7.290).

Soft-band Ward identity and NG soft theorem. For any insertion of BRST-closed physical operators $\{\mathcal{O}_i\}$, the $k\to 0$ limit of the current insertion reproduces the broken charge action:

$$\lim_{k\to 0} k_\mu\, \widetilde{G}^\mu_a\big(k;\{x_i\}\big) = \sum_i \left\langle \mathcal{O}_1\cdots(\delta_a\mathcal{O}_i)\cdots\mathcal{O}_n\right\rangle_{\omega^{(b)}}, \tag{7.295}$$

where $\widetilde{G}^{\mu}_{a}$ is the Fourier transform of $\langle T\, J^{\mu}_{a}(x) \prod_i \mathcal{O}_i(x_i)\rangle$. Upon LSZ reduction, this yields the NG soft theorem (Adler-type statement) on the band:

$$\mathcal{A}_{n+1}\Big(\pi^a(k),\, \Psi_1, \ldots, \Psi_n\Big) = -\,\frac{\mathrm{i}}{f_{aa}} \sum_{i=1}^{n} \mathsf{T}^{(i)}_{a}\, \mathcal{A}_n\Big(\Psi_1, \ldots, \Psi_n\Big) \;+\; \mathcal{O}(k)\,, \tag{7.296}$$

where $\mathsf{T}^{(i)}_{a}$ is the action of the broken generator on external leg i, and f_{aa} is the appropriate decay constant extracted from eq. (7.293). When the total broken charge on external states vanishes, the leading term cancels and the amplitude exhibits an Adler zero. All statements are gauge-parameter independent on certified plateaus by the Nielsen machinery (Section 7.5).

Plateau protection. The coefficients f_{ab} and the NG pole in eq. (7.294) are stable under admissible UV/IR scans:

$$\partial_{(\ln \lambda_{\mathrm{UV}},\, \ln \ell)}\, f_{ab} = \mathcal{O}(\varepsilon_{\mathrm{plat}})\,, \qquad \partial_{(\ln \lambda_{\mathrm{UV}},\, \ln \ell)}\, \mathrm{Res}_{k^2=0}\, G_{JJ}(k) = \mathcal{O}(\varepsilon_{\mathrm{plat}})\,, \tag{7.297}$$

a direct consequence of eq. (7.291) and the acceptance tests of §1.5.

2.5 BEH reorganisation (gauged G)

When the broken generators are gauged, the longitudinal gauge fields absorb the would-be NG modes (BEH mechanism). Let the covariant derivative be $D_\mu = \partial_\mu + \mathrm{i}\, g_a A^a_\mu T_a$ in the representation of Φ. Expanding $\Phi = U(\pi)(v + h)$ from eq. (7.284) to quadratic order yields the vector mass matrix

$$(M_A^2)_{ab} = g_a g_b \left\langle v \,\middle|\, \{T_a, T_b\} \,\middle|\, v \right\rangle \quad \text{(projected to } \mathcal{H}_{\mathrm{phys}}\text{)}, \tag{7.298}$$

which is positive semidefinite by eq. (7.13). The kinetic mixing between A_μ and π takes the universal derivative form

$$\mathcal{L}^{(2)}_{\mathrm{mix}} = -\, g_a\, A^{a\,\mu}\, \partial_\mu \pi^a + \cdots\,, \tag{7.299}$$

and is removed by R_ξ gauge fixing. In that gauge the unphysical scalar has mass

$$m^2_{\pi^a,\mathrm{unphys}} = \xi\, (M_A^2)_{aa}\,, \tag{7.300}$$

forming part of a BRST quartet with the Faddeev–Popov ghosts/longitudinal modes that decouple from $\mathcal{H}_{\mathrm{phys}}$ (Section 7.2). Physical poles and residues (transverse vectors and h) are gauge-parameter independent on shell by Slavnov–Taylor/Nielsen identities (Section 7.5).

Goldstone equivalence (on the band). For energies $E \gg M_A$ and on-shell external states prepared on the band,

$$\mathcal{M}\Big(A^a_L;\, \mathcal{X}\Big) = \mathcal{M}\Big(\pi^a;\, \mathcal{X}\Big) \;+\; \mathcal{O}\Big(\frac{M_A}{E}\Big)\,, \tag{7.301}$$

where A^a_L is a longitudinal vector and $\mathcal{X}$ denotes other physical insertions. The equivalence is protected by ST/Nielsen identities and the projector commutators eq. (7.244); the $\mathcal{O}(M_A/E)$ corrections are plateau-stable by eq. (7.291).

Plateau protection and provisional identifications. The spectra and residues of transverse vectors and gapped scalars h are stable under admissible UV/IR scans on the band (cf. eq. (7.291)); any "Higgs-like'' identification of h or related composites is *provisional* pending cross-quadrant certification and plateau robustness checks (policy in Section 7.4).

2.6 State visibility: volume scaling signature of SSB

Using the same readout map as in Section 7.4, define the *state* visibility from the ancilla coherence between two prepared vacua (or two preparations of the same vacuum) on the band:

$$\Lambda_{\text{state}}(\lambda_{\text{UV}},\ell;V) := \frac{\left|\operatorname{Tr}_{\mathcal{H}_{\text{phys}}}\Big(P_x(\lambda_{\text{UV}})\,K_{\text{IR}}^{(\ell)}\,U_1^{(\text{state})\dagger}U_2^{(\text{state})}\,\rho_{\text{ref}}\Big)\right|}{\operatorname{Tr}_{\mathcal{H}_{\text{phys}}}\Big(P_x(\lambda_{\text{UV}})\,K_{\text{IR}}^{(\ell)}\,\rho_{\text{ref}}\Big)}\,. \tag{7.302}$$

Here ρ_{ref} is a BRST-closed calibration state (ghost number 0); the state-channel unitaries $U_b^{(\text{state})}$ prepare the broken vacua as in eqs. (7.276) and (7.277), with the ordered limits eq. (7.275). All band operations commute with the symmetry/BRST generators (eq. (7.244)), so Λ_{state} is gauge-parameter independent on certified plateaus (Nielsen) and depends only on the prepared *state* sector (the law channel stays fixed by Section 7.7.1).

Quadrant-II signature (law vs. state). With the law frozen, Quadrant II is diagnosed by

$$|\Lambda_{\text{law}}(\lambda_{\text{UV}},\ell)| = 1+\mathcal{O}(\varepsilon_{\text{plat}}) \quad \text{(eqs. (7.257) and (7.259))}, \qquad \lim_{V\to\infty}|\Lambda_{\text{state}}(\lambda_{\text{UV}},\ell;V)| = 0, \tag{7.303}$$

with all limits taken inside a certified window (Section 7.3).

Asymptotic scaling (orthogonality of pure phases). Let $\omega^{(1)},\omega^{(2)}$ be the two branch states produced by eqs. (7.276) and (7.277) along distinct points of $\mathcal{M}_{\text{vac}}$ (or two macroscopically distinct ensembles around the same vacuum). Under standard cluster properties on the band, the overlap (obtained from eq. (7.302)) obeys a large-volume law

$$-\frac{1}{V}\log|\Lambda_{\text{state}}(\lambda_{\text{UV}},\ell;V)| \xrightarrow[V\to\infty]{} \alpha(\lambda_{\text{UV}},\ell) \;\geq 0\,, \tag{7.304}$$

with $\alpha = 0$ iff the two preparations coincide in the thermodynamic limit. Two useful regimes—both plateau-stable by eq. (7.291)—are:

1. **Gapped (short-range) regime.** If connected correlators have a finite correlation length ξ on the band, block-factorisation gives an extensive decay

$$|\Lambda_{\text{state}}| \;\leq\; \exp\Big[-\,c_{\text{gap}}\,V/\xi^{d}\Big], \qquad c_{\text{gap}}>0 \text{ set by the local overlap between phases.} \tag{7.305}$$

2. **Continuous SSB with NG modes.** For two vacua related by a small global rotation of angle θ in G/H, the zero-mode (uniform) component yields

$$|\Lambda_{\text{state}}| \;\lesssim\; \exp\Big[-\,\tfrac{1}{2}\,\rho_{\text{s}}\,\theta^2\,V\Big] \quad \text{for } \theta\ll 1, \tag{7.306}$$

where ρ_{s} is the appropriate stiffness (decay constant) extracted from eqs. (7.293) and (7.294). Thus distinct NG orientations are orthogonal as $V\to\infty$.

Estimator and plateau stability. Define the finite-volume estimator

$$\widehat{\alpha}(\lambda_{\mathrm{UV}}, \ell; V) := -\frac{1}{V} \log |\Lambda_{\mathrm{state}}(\lambda_{\mathrm{UV}}, \ell; V)| \,. \tag{7.307}$$

On certified windows,

$$\partial_{(\ln \lambda_{\mathrm{UV}}, \ln \ell)} \widehat{\alpha} = \mathcal{O}(\varepsilon_{\mathrm{plat}}) \,, \tag{7.308}$$

so the extracted macroscopic rate is insensitive to admissible UV/IR scans, mirroring eqs. (7.291) and (7.297).

Calibrations and localised contrasts. If both branches prepare the *same* vacuum sector, then $|\Lambda_{\mathrm{state}}| = 1 + \mathcal{O}(\varepsilon_{\mathrm{plat}})$. If the preparations differ only inside a subregion $\Omega \subset V$ (e.g., a local quench or inhomogeneous seed), the scaling in eq. (7.304) holds with V replaced by $|\Omega|$. Interface-dominated contrasts (domain walls) instead scale with boundary area and appear as $\exp[-\sigma\, \mathrm{Area}(\partial\Omega)]$, providing an operational handle to distinguish bulk vs. interface mechanisms.

Operational note. When reporting Λ_{state}, include: (i) the certified plateau window and $\varepsilon_{\mathrm{plat}}$; (ii) whether the two preparations are symmetry-related vacua or the same vacuum; (iii) the finite-volume fit to $-\log|\Lambda_{\mathrm{state}}|$ (linear in V, or area-law if applicable); and (iv) the gauge-scan confirming Nielsen stability (cf. eq. (7.260)). Coupled with eq. (7.303), this constitutes the Quadrant-II SSB fingerprint.

2.7 Acceptance and reporting for the broken-state run

The broken-state run is accepted only if *all* items below are satisfied and reported with their numerical budgets.

- **Plateaus (certification and budgets).** Certify a domain in $(\ln \lambda_{\mathrm{UV}}, \ln \ell)$ with small gradient/oscillation/curvature as in Section 7.3, and report the triple $(\varepsilon_{\mathrm{grad}}, \varepsilon_{\mathrm{osc}}, \varepsilon_{\mathrm{curv}})$ together with regulator robustness (Section 7.4.5). Quote the plateau budget $\varepsilon_{\mathrm{plat}}$ entering eqs. (7.259) and (7.308).
- **Identity ledger (branchwise).** Verify on-shell Slavnov–Taylor, global Ward, and Nielsen identities on each branch (Section 7.5); include anomaly filters if applicable. For global G, exhibit eq. (7.292) and the NG matrix element eq. (7.293); for gauged G, demonstrate ST/Nielsen protection of pole positions/residues together with the BEH reorganisation eqs. (7.298) to (7.301).
- **Spectrum health (band-projected).** Recheck non-negativity of the physical mass operator and hyperbolicity of the principal symbol on $\mathcal{H}_{\mathrm{phys}}$ (cf. eqs. (7.13) and (7.15)), and stability of the broken Hessian under UV/IR scans (eq. (7.291)).
- **Quadrant tests (law vs. state).** Show the law channel remains coherent on the plateau, $|\Lambda_{\mathrm{law}}| = 1 + \mathcal{O}(\varepsilon_{\mathrm{plat}})$ (eqs. (7.257) and (7.259)), and that the state channel exhibits

volume-suppressed visibility $\lim_{V\to\infty}|\Lambda_{\text{state}}| = 0$ (eq. (7.303)). Include finite-size scaling plots at fixed plateaus and fit the visibility via

$$-\log|\Lambda_{\text{state}}(\lambda_{\text{UV}},\ell;V)| = \alpha(\lambda_{\text{UV}},\ell)\,V + \sigma(\lambda_{\text{UV}},\ell)\,\text{Area}(\partial\Omega) + c_0 + \frac{c_1}{V} + \cdots\,, \quad (7.309)$$

reporting α (bulk decay, cf. eq. (7.304)), σ (interface cost when applicable), and uncertainties.

- **Regulator robustness and ordering.** Demonstrate stability across admissible $(P_x, K_{\text{IR}}^{(\ell)})$ choices that commute with G and BRST (cf. eq. (7.244)), enforcing

$$\left|\Lambda_{\text{state}}[U_1^{(\text{state})}, U_2^{(\text{state})}] - \Lambda_{\text{state}}[U_2^{(\text{state})}, U_1^{(\text{state})}]\right| \le \varepsilon_{\text{plat}}\,, \quad (7.310)$$

and showing insensitivity to projector ordering analogous to eq. (7.268) (with law $\to$ state replacements when applicable).

- **Route equivalence and seed profile.** Confirm that imaginary-time and unitary preparations produce identical band–projected correlators in the ordered limit (Lemma 7.339), and that the result is robust under smooth changes of the seed profile along the same coset direction (cf. Corollary 7.340). State explicitly the seed operator $\mathcal{O}_X$ and Hamiltonian $\Delta\mathcal{H}_{\text{seed}}$ used (eqs. (7.273) and (7.274)).

- **Gauge scan (Nielsen).** Report a gauge scan confirming $\partial_\zeta\Lambda_{\text{law}} = \mathcal{O}(\varepsilon_{\text{plat}})$ (eq. (7.260)) and the gauge-parameter independence of on-shell, band-projected state observables (by the same machinery; cf. Section 7.5).

- **Instrumental budget.** Quote the readout floor $\varepsilon_{\text{inst}} = \sigma_{\text{shot}} + \sigma_\phi + \sigma_{\text{sys}}$ (eq. (7.271)); ensure reported deficits/decays clear this floor with margin.

2.8 Failure modes and diagnostics

- *No NG pole (global G).* Check for (i) residual explicit breaking (violations of eq. (7.292)), (ii) improper order of limits in eq. (7.275), or (iii) off-plateau contamination (violation of Section 7.3). Revisit the seed protocol eqs. (7.273) and (7.274) and verify the Goldstone zero-mode identity eq. (7.290) and constraints eq. (7.289).

- *Gauge dependence of poles (gauged G).* A nonzero Nielsen derivative for pole positions/residues indicates off-shell mismatch. Restore on-shell conditions and band closure (eq. (7.244)) and recheck the exact-identity sector (Section 7.5).

- *Tachyon/negative modes on $\mathcal{H}_{\text{phys}}$.* The broken background is unstable. Recompute the band Hessian eq. (7.287), reselect v, and verify $\mathcal{M}^2_{\text{phys}} \ge 0$ and principal-symbol hyperbolicity (cf. eqs. (7.13) and (7.15)).

- $|\Lambda_{\text{law}}| < 1$ *at plateau.* The law channel is compromised; invoke the converse lemma Lemma 7.338 and revert to Step 1: revalidate the commutators eq. (7.245), projector compatibility eq. (7.246), and plateau certification.

- *Weak or absent volume decay of* $|\Lambda_{\text{state}}|$. If $-\log|\Lambda_{\text{state}}|$ is subextensive in V, either (i) both branches prepare the same vacuum sector (expect $|\Lambda_{\text{state}}| = 1 + \mathcal{O}(\varepsilon_{\text{plat}})$), or (ii) the contrast is interface-dominated (area law). Diagnose with the finite-size estimator eq. (7.307) and the fit eq. (7.309); if the preparations are related by a small rotation in G/H, use the NG bound eq. (7.306) to estimate the expected rate $\alpha \simeq \frac{1}{2}\rho_{\text{s}}\theta^2$.
- *Finite-size artefacts / order of limits.* Ensure $V \to \infty$ precedes $\varepsilon \to 0^+$ on a certified plateau (cf. eq. (7.275)); stabilise the extracted rate with the plateau derivative bound eq. (7.308).
- *Regulator mismatch.* If Λ_{state} or NG diagnostics drift under admissible regulator swaps, re-enforce eq. (7.265) and the ordering tests paralleling eq. (7.268) (state channel).
- *Instrumental artefacts.* Apparent deficits consistent with the floor $\varepsilon_{\text{inst}}$ (eq. (7.271)) are non-diagnostic; improve SNR or phase calibration before attributing physics.

7.7.3 Band Hessian to BRST Projection (Transverse/Longitudinal)

Aim. Starting from the asymmetric background prepared in Section 7.7.2 (with the *law* frozen as in Section 7.7.1), we construct the band Hessian, perform the transverse/longitudinal (T/L) split, and project to the BRST cohomology on $\mathcal{H}_{\text{phys}}$ so that ghosts and longitudinal artefacts decouple. All steps are performed *on the band* with the same UV/IR controls and plateau certification as in Section 7.3, and on-shell statements are protected by Slavnov–Taylor/Ward/Nielsen identities (Section 7.5; [117, 442, 91, 82]).

3.1 Band quadratic kernel about the broken background

Let Γ denote the (band) effective action including the background φ selected in Section 7.7.2 and the admissible regulator/gauge-fixing sector used in Section 7.2. The band Hessian at $\Phi = \varphi$ is the quadratic kernel (cf. eq. (7.287))

$$\mathcal{H}_{\text{band}}(x,y) := \left.\frac{\delta^2\Gamma}{\delta\Phi(x)\,\delta\Phi(y)}\right|_{\Phi=\varphi}, \tag{7.311}$$

which determines inverse propagators and mixings among fluctuations $\Psi \equiv (A_\mu, \sigma, h, \pi; c, \bar{c}, b)$. Unless stated otherwise, operators are implicitly composed with the UV projector $P_x(\lambda_{\text{UV}})$ and IR window $K_{\text{IR}}^{(\ell)}$, and traces are taken on $\mathcal{H}_{\text{phys}}$ (Sections 7.2 and 7.3).

Background stationarity and linear terms. The selected background φ is a stationary point of Γ on the certified band:

$$\left.\frac{\delta\Gamma}{\delta\Phi(x)}\right|_{\Phi=\varphi} = 0 \qquad \Rightarrow \qquad \text{no tadpoles at quadratic order,} \tag{7.312}$$

so $\mathcal{H}_{\text{band}}$ fully controls the Gaussian fluctuations about φ. For global breaking, φ lies on $\mathcal{M}_{\text{vac}}$ and satisfies the Goldstone zero–mode identity eq. (7.290).

Band filtering (shorthand). It is convenient to write the band filter as

$$\mathsf{B}_{\lambda,\ell} := K_{\mathrm{IR}}^{(\ell)}\, P_x(\lambda_{\mathrm{UV}})\,, \qquad [\mathsf{B}_{\lambda,\ell}, \hat{Q}_{\mathrm{BRST}}] = [\mathsf{B}_{\lambda,\ell}, \mathsf{J}_\alpha] = 0 \ \text{ (plateau; cf. } eq.\ (7.244)), \tag{7.313}$$

so that any quadratic form is implicitly "sandwiched" by $\mathsf{B}_{\lambda,\ell}$ when restricted to the band.

Kernel properties (symmetries and gradings). On the certified band,

$$\mathcal{H}_{\mathrm{band}}(x,y) = \mathcal{H}_{\mathrm{band}}^{\top}(y,x), \qquad \mathrm{gh}\big(\mathcal{H}_{\mathrm{band}}\big) = 0, \qquad [\hat{Q}_{\mathrm{BRST}}, \mathcal{H}_{\mathrm{band}}] = \{\hat{Q}_{\mathrm{BRST}}, \ldots\} \tag{7.314}$$

(the last relation indicates BRST variations are exact because the gauge-fixing sector is $\{\hat{Q}_{\mathrm{BRST}}, \Psi_\zeta\}$; gauge–parameter variations are therefore BRST–exact at the level of the physical blocks, see eq. (7.339) below §3.4).

Derivative/mass decomposition (broken background). In momentum space, each block of $\mathcal{H}_{\mathrm{band}}$ admits a principal–symbol plus mass/mixing split,

$$\mathcal{H}^{XY}_{\mathrm{band}}(p) = \mathcal{K}^{XY}_{(\varphi)}(p) + \mathcal{M}^{2,XY}_{(\varphi)}\,, \qquad X, Y \in \{T, L, h, \pi, c, \bar{c}, b\}, \tag{7.315}$$

where $\mathcal{K}^{XY}_{(\varphi)}(p)$ carries the highest-derivative (hyperbolic) structure and $\mathcal{M}^{2,XY}_{(\varphi)}$ the local mass/mixing terms evaluated at φ. Continuity from the symmetric band together with the acceptance tests implies that the principal symbol remains hyperbolic and the physical mass operator non-negative on $\mathcal{H}_{\mathrm{phys}}$ (cf. eqs. (7.13) and (7.15)).

Ward/Nielsen constraints (selection rules). Broken global symmetries enforce (on the band) the NG constraints in the scalar sector,

$$\mathcal{M}^2_{\pi\pi} = 0, \qquad \mathcal{K}_{\pi h}(p) = \mathcal{O}(p), \qquad \mathcal{K}_{\pi h}(0) = 0, \tag{7.316}$$

as stated in eq. (7.289), while gauge realizations reorganize would-be NG modes into BRST quartets (see §2.5). Dependence on the gauge parameter ζ is BRST–exact on the physical blocks (Nielsen corollary eq. (7.339)).

Block structure (schematic). In the fluctuation basis Ψ,

$$\mathcal{H}_{\mathrm{band}} = \begin{pmatrix} \mathcal{H}^{TT} & \mathcal{H}^{TL} & \mathcal{H}^{Th} & \mathcal{H}^{T\pi} & 0 & 0 & 0 \\ \mathcal{H}^{LT} & \mathcal{H}^{LL} & \mathcal{H}^{Lh} & \mathcal{H}^{L\pi} & \star & \star & \star \\ \mathcal{H}^{hT} & \mathcal{H}^{hL} & \mathcal{H}^{hh} & \mathcal{H}^{h\pi} & 0 & 0 & 0 \\ \mathcal{H}^{\pi T} & \mathcal{H}^{\pi L} & \mathcal{H}^{\pi h} & \mathcal{H}^{\pi\pi} & 0 & 0 & 0 \\ 0 & \star & 0 & 0 & 0 & \mathcal{H}^{c\bar{c}} & 0 \\ 0 & \star & 0 & 0 & \mathcal{H}^{\bar{c}c} & 0 & 0 \\ 0 & \star & 0 & 0 & 0 & 0 & \mathcal{H}^{bb} \end{pmatrix}, \tag{7.317}$$

where the T/L superscripts refer to the transverse/longitudinal decomposition of A_μ (defined in §3.2). The starred entries implement the BRST doublet structure tying the longitudinal sector to $(c, \bar{c}, b)$; these ensure quartet decoupling on $\mathcal{H}_{\mathrm{phys}}$ (Sections 7.2.6 and 7.5). Gauge/BRST invariance implies that mixings with one longitudinal leg vanish at zero momentum,

$$\mathcal{H}^{TL}(p) \text{ and } \mathcal{H}^{LT}(p) \ \propto\ p \quad \Rightarrow \quad \mathcal{H}^{TL}(0) = \mathcal{H}^{LT}(0) = 0\,, \tag{7.318}$$

consistent with the ST/Ward identities and with the Schur-reduction in §3.4.

Plateau robustness (band response). Within a certified window in $(\ln \lambda_{\rm UV}, \ln \ell)$,

$$\partial_{(\ln \lambda_{\rm UV},\, \ln \ell)}\, \mathcal{H}_{\rm band}(p) = \mathcal{O}(\varepsilon_{\rm plat})\,, \tag{7.319}$$

as already stated in eq. (7.291). Hence eigenvalues/eigenvectors associated with NG directions and gapped modes vary only within plateau tolerances, and all subsequent on-shell reductions (T/L split and BRST projection in §3.2–3.4) inherit the same stability.

3.2 Transverse/longitudinal split and band commutation

In momentum space (Minkowski signature $\eta_{\mu\nu}$; Euclidean continuation is straightforward), define

$$\Pi_{\rm T}^{\mu\nu}(p) = \eta^{\mu\nu} - \frac{p^\mu p^\nu}{p^2}\,, \qquad \Pi_{\rm L}^{\mu\nu}(p) = \frac{p^\mu p^\nu}{p^2}\,, \tag{7.320}$$

with the understanding that $p^2 = 0$ singularities are regulated inside the certified IR window $K_{\rm IR}^{(\ell)}$ (and never probed outside it).

Algebra and kinematics. The projectors obey, for $p^2 \neq 0$,

$$\Pi_{\rm T}^2 = \Pi_{\rm T},\quad \Pi_{\rm L}^2 = \Pi_{\rm L},\quad \Pi_{\rm T}\Pi_{\rm L} = 0,\quad \Pi_{\rm T} + \Pi_{\rm L} = \mathbb{1},\qquad p_\mu \Pi_{\rm T}^{\mu\nu} = 0,\quad \Pi_{\rm L}^{\mu\nu} v_\nu = \frac{p^\mu\, p\cdot v}{p^2}. \tag{7.321}$$

Accordingly we decompose any vector A_μ as $A_\mu = A_\mu^{\rm T} + A_\mu^{\rm L}$, with

$$A_\mu^{\rm T}(p) = \Pi_{\rm T}{}_\mu^{\ \nu}(p)\, A_\nu(p), \qquad A_\mu^{\rm L}(p) = \Pi_{\rm L}{}_\mu^{\ \nu}(p)\, A_\nu(p). \tag{7.322}$$

Real–space (Helmholtz) form on the band. Define the IR–regulated inverse d'Alembertian on the band by the kernel

$$\big(\Delta_\ell^{-1} f\big)(x) := -\int \frac{d^d p}{(2\pi)^d}\, \frac{K_{\rm IR}^{(\ell)}(p)}{p^2}\, e^{{\rm i} p\cdot x}\, \tilde f(p), \tag{7.323}$$

so that $-\Box\, \Delta_\ell^{-1} \approx \mathbb{1}$ on the support of $K_{\rm IR}^{(\ell)}$. Then the band–restricted Helmholtz decomposition reads

$$A_\mu^{\rm L} = \partial_\mu\, \Delta_\ell^{-1}(\partial\cdot A), \qquad A_\mu^{\rm T} = A_\mu - \partial_\mu\, \Delta_\ell^{-1}(\partial\cdot A), \tag{7.324}$$

which reproduces eq. (7.322) after Fourier transform and automatically inherits the IR regularisation.

Band commutation and symmetry/BRST compatibility. On the certified band, the UV projector $P_x(\lambda_{\rm UV})$ and the IR window $K_{\rm IR}^{(\ell)}$ act as scalar multipliers in momentum space (functions of p^2), while $\Pi_{{\rm T}/L}$ act only on Lorentz indices. Hence

$$\Big[P_x(\lambda_{\rm UV}), \Pi_{{\rm T}/L}\Big] = \Big[K_{\rm IR}^{(\ell)}, \Pi_{{\rm T}/L}\Big] = 0, \qquad \Big[\hat Q_{\rm BRST}, \Pi_{{\rm T}/L}\Big] = \Big[\mathsf{J}_\alpha, \Pi_{{\rm T}/L}\Big] = 0, \tag{7.325}$$

as advertised (cf. eq. (7.244)). The last two commutators hold because $\Pi_{{\rm T}/L}$ depend only on p^μ and are blind to internal/gauge indices on which $\hat Q_{\rm BRST}$ and J_α act.

Zero–mode handling ($p^2 \to 0$) and finite volume. In a finite box $V = L^d$ with periodic boundary conditions, the smallest nonzero momentum is $\sim 2\pi/L$; within the IR window the regulated kernel Δ_ℓ^{-1} in eq. (7.323) is well defined provided $L \gg \ell$. If an exact zero mode is present (e.g. strict spatially uniform mode in a massless sector), replace $1/p^2$ by its Moore–Penrose pseudoinverse on the band, or equivalently define $\Delta_{\ell,V}^{-1}$ by projecting out the zero eigenspace before inversion:

$$\Delta_{\ell,V}^{-1} := -\left(\Box|_{\text{band}\ominus\ker\Box}\right)^{-1} K_{\text{IR}}^{(\ell)}\,, \qquad \Pi_{\text{L}} = \partial\,\Delta_{\ell,V}^{-1}\,\partial, \quad \Pi_{\text{T}} = \not\Vdash - \Pi_{\text{L}}. \tag{7.326}$$

All identities used later (idempotence, orthogonality, and the commutators in eq. (7.325)) continue to hold on the orthogonal complement of $\ker\Box$.

Mixing constraints at zero momentum. Gauge/ST identities imply that any mixing carrying a single longitudinal leg is $\mathcal{O}(p)$; thus

$$\mathcal{H}^{TL}(p),\, \mathcal{H}^{LT}(p) \;\propto\; p \quad \Rightarrow \quad \mathcal{H}^{TL}(0) = \mathcal{H}^{LT}(0) = 0, \tag{7.327}$$

consistent with the Schur reduction in §3.4 and the NG/BEH structure discussed in §2.4–2.5.

Multiple gauge sectors and color structure. For nonabelian groups, $\Pi_{\text{T}/L}$ act trivially on color indices: $(\Pi_{\text{T}/L})^{\mu\nu}\delta^{ab}$. Consequently they commute with global/gauge generators and BRST differentials acting in the internal space, ensuring the projector identities used in the BRST decoupling and in the physical block construction (§3.3–3.4).

Plateau stability. Since $\Pi_{\text{T}/L}$ are independent of λ_{UV} and ℓ (the dependence sits only in the band filter and Δ_ℓ^{-1}), the T/L split is stable under admissible UV/IR scans; any residual ℓ-dependence enters only through the regulated Green's operator in eq. (7.324) and is bounded within the plateau budget (cf. eq. (7.319)).

With eqs. (7.322), (7.324) and (7.325) in hand, we will form the physical inverse two-point blocks by combining the T/L split with the BRST doublet structure (quartet mechanism) in §3.3 and the Schur reduction in §3.4.

3.3 BRST doublets and quartet decoupling

BRST algebra near the broken background (band form). Let the gauge–fixing sector be generated by a fermion Ψ_ζ so that $S_{\text{gf+gh}} = \{\hat{Q}_{\text{BRST}}, \Psi_\zeta\}$. Linearised about the background φ (with background–covariant derivative $\bar{D}_\mu$), the BRST variations on the band read schematically

$$\begin{aligned} [\hat{Q}_{\text{BRST}},\, A_\mu^{\text{L}}] &= \bar{D}_\mu c + \mathcal{O}(\text{nonlin}),\\ \{\hat{Q}_{\text{BRST}},\, \bar{c}\} &= b,\\ [\hat{Q}_{\text{BRST}},\, b] &= 0,\\ \{\hat{Q}_{\text{BRST}},\, c\} &= 0, \end{aligned} \tag{7.328}$$

with ghost numbers $\text{gh}(c) = +1$, $\text{gh}(\bar{c}) = -1$, $\text{gh}(b) = 0$, and $\hat{Q}_{\text{BRST}}^2 = 0$. The transverse field A_μ^{T} and the matter sector (h, π) transform covariantly but do not form BRST doublets; their cohomology carries the physical poles.

Doublets and trivial cohomology. A pair (u, v) with $\{\hat{Q}_{\mathrm{BRST}}, u\} = v$ and $[\hat{Q}_{\mathrm{BRST}}, v] = 0$ is a BRST *doublet*. By eq. (7.328), the longitudinal/ghost/Nakanishi–Lautrup variables assemble into two doublets $(\bar{c}, b)$ and $(A^{\mathrm{L}}, \bar{D}c)$. The standard doublet lemma implies that the cohomology in the algebra generated by doublet variables is trivial: any BRST–closed functional of doublets is BRST–exact. Consequently, insertions made of $(A^{\mathrm{L}}, c, \bar{c}, b)$ alone do not contribute to physical matrix elements.

Band gauge fixing and the longitudinal block. Since $S_{\mathrm{gf+gh}} = \{\hat{Q}_{\mathrm{BRST}}, \Psi_\zeta\}$, its quadratic expansion about φ yields the ghost/longitudinal part of the band Hessian. Hence there exists a band operator $\mathcal{K}_{\mathrm{dbl}}$ (depending on Ψ_ζ and the background) such that

$$\mathcal{H}^{LL} \oplus \mathcal{H}^{c\bar{c}} \oplus \mathcal{H}^{\bar{c}c} \oplus \mathcal{H}^{bb} = \left\{\hat{Q}_{\mathrm{BRST}},\, \mathcal{K}_{\mathrm{dbl}}\right\} \qquad \text{(on the band)}. \tag{7.329}$$

In particular, mixings that carry one longitudinal leg $(TL, LT, Lh, L\pi)$ arise from the same BRST–exact sector and are proportional to a single momentum insertion (cf. eq. (7.318)).

Physical projector and graded cyclicity. Let $\mathsf{P}_{\mathrm{phys}}$ denote the projector onto the BRST cohomology (ghost number 0) on the band; by construction,

$$\mathsf{P}_{\mathrm{phys}}\hat{Q}_{\mathrm{BRST}} = \hat{Q}_{\mathrm{BRST}}\mathsf{P}_{\mathrm{phys}} = 0, \qquad [\mathsf{P}_{\mathrm{phys}},\, P_x(\lambda_{\mathrm{UV}})] = [\mathsf{P}_{\mathrm{phys}},\, K_{\mathrm{IR}}^{(\ell)}] = 0. \tag{7.330}$$

Therefore, for any BRST–closed, ghost–number 0 operator $\mathcal{O}$ (or between physical states), graded cyclicity gives the band identity

$$\mathrm{Tr}_{\mathcal{H}_{\mathrm{phys}}}\Big(\mathcal{O}\,\{\hat{Q}_{\mathrm{BRST}}, X\}\Big) = 0, \qquad \langle\psi_{\mathrm{phys}}|\,\{\hat{Q}_{\mathrm{BRST}}, X\}\,|\phi_{\mathrm{phys}}\rangle = 0, \tag{7.331}$$

for any band operator X. Combining eq. (7.329) with eq. (7.331) yields the cohomological vanishing restated in eq. (7.332):

$$\mathsf{P}_{\mathrm{phys}}\left(\mathcal{H}^{LL} \oplus \mathcal{H}^{c\bar{c}} \oplus \mathcal{H}^{\bar{c}c} \oplus \mathcal{H}^{bb}\right)\mathsf{P}_{\mathrm{phys}} \;\simeq\; 0 \qquad \text{(in BRST cohomology)}. \tag{7.332}$$

Quartet mechanism (band version). The two doublets $(\bar{c}, b)$ and $(A^{\mathrm{L}}, \bar{D}c)$ combine into a Kugo–Ojima *quartet*. Off shell, members of a quartet come in norm–indefinite pairs; on the band and on shell, their contributions cancel in any physical matrix element by eq. (7.331). Thus the spectrum and residues relevant to observables are carried entirely by the transverse gauge block and the matter block, as used in §3.4.

Gauge–parameter independence and plateaus. Because the entire doublet sector is BRST–exact on the band, a gauge–parameter variation ∂_ζ acts as a BRST commutator on the physical blocks (Nielsen corollary eq. (7.339)). Moreover, since all band projectors commute with BRST and with the T/L projectors (cf. eq. (7.325)), the quartet decoupling is stable under admissible UV/IR scans within a certified window (plateau robustness $\mathcal{O}(\varepsilon_{\mathrm{plat}})$).

Consequence for the physical Hessian. Equations eqs. (7.329) and (7.331) imply that the Schur–complement subtractions used to eliminate longitudinal mixings are themselves BRST–exact and therefore drop from physical matrix elements; this is formalised and used in §3.4 to obtain the on–shell simplifications eq. (7.336).

3.4 Physical block matrix and on-shell reduction

The band-projected *physical* inverse two-point kernel is obtained by restricting eq. (7.317) to the BRST cohomology and eliminating unphysical mixings. At the operator level, a convenient gauge-invariant construction uses Schur complements:

$$\mathcal{H}^{TT}_{\mathrm{phys}} := \Pi_{\mathrm{T}}\Big(\mathcal{H}^{TT} - \mathcal{H}^{TL}\,(\mathcal{H}^{LL})^{-1}\,\mathcal{H}^{LT}\Big)\Pi_{\mathrm{T}}\,, \tag{7.333}$$

and similarly for the matter block

$$\mathcal{H}^{\mathrm{mat}}_{\mathrm{phys}} := \begin{pmatrix}\mathcal{H}^{hh} & \mathcal{H}^{h\pi}\\ \mathcal{H}^{\pi h} & \mathcal{H}^{\pi\pi}\end{pmatrix} - \begin{pmatrix}\mathcal{H}^{hL}\\ \mathcal{H}^{\pi L}\end{pmatrix}(\mathcal{H}^{LL})^{-1}\begin{pmatrix}\mathcal{H}^{Lh} & \mathcal{H}^{L\pi}\end{pmatrix}, \tag{7.334}$$

all understood on the band and on $\mathcal{H}_{\mathrm{phys}}$.

Lemma 7.342 (BRST-exactness of Schur subtractions on the band)**.** *Let the longitudinal and $(c,\bar c,b)$ sectors form BRST doublets as in eq. (7.328), and assume the band-closure commutators eq. (7.325). Then there exist band operators $\mathcal{B}_T, \mathcal{B}_{\mathrm{mat}}$ such that*

$$\begin{aligned}\Pi_{\mathrm{T}}\,\mathcal{H}^{TL}(\mathcal{H}^{LL})^{-1}\mathcal{H}^{LT}\,\Pi_{\mathrm{T}} &= \{\hat Q_{\mathrm{BRST}},\,\mathcal{B}_T\},\\ \begin{pmatrix}\mathcal{H}^{hL}\\ \mathcal{H}^{\pi L}\end{pmatrix}(\mathcal{H}^{LL})^{-1}\begin{pmatrix}\mathcal{H}^{Lh} & \mathcal{H}^{L\pi}\end{pmatrix} &= \{\hat Q_{\mathrm{BRST}},\,\mathcal{B}_{\mathrm{mat}}\}.\end{aligned} \tag{7.335}$$

Consequently, for any BRST-closed, ghost-number 0 insertion $\mathcal{O}$ (and likewise between physical states), the Schur subtractions vanish:

$$\mathrm{Tr}_{\mathcal{H}_{\mathrm{phys}}}\Big(\mathcal{O}\,\Pi_{\mathrm{T}}\mathcal{H}^{TL}(\mathcal{H}^{LL})^{-1}\mathcal{H}^{LT}\Pi_{\mathrm{T}}\Big) = \mathrm{Tr}_{\mathcal{H}_{\mathrm{phys}}}\Big(\mathcal{O}\,\{\hat Q_{\mathrm{BRST}},\mathcal{B}_T\}\Big) = 0,$$

and similarly for the matter block, by the cohomological identity eq. (7.331).

Proof. By eq. (7.328), the longitudinal/ghost/Nakanishi–Lautrup sector is a sum of BRST doublets; variations of Γ with respect to these fields therefore lie in the image of $\hat Q_{\mathrm{BRST}}$ (see [117, 442]). In particular, the "constraint'' operator that maps longitudinal fluctuations to gauge variations, and the inverse longitudinal kernel $(\mathcal{H}^{LL})^{-1}$ generated by the gauge-fixing term $\{\hat Q_{\mathrm{BRST}}, \Psi_\zeta\}$, are BRST-exact on the band. The mixing blocks $\mathcal{H}^{TL}, \mathcal{H}^{LT}, \mathcal{H}^{hL}, \mathcal{H}^{Lh}, \dots$ arise from this gauge-fixing sector and carry one longitudinal leg; composing them with $(\mathcal{H}^{LL})^{-1}$ keeps the result in the BRST image. Commutators eq. (7.325) ensure that inserting Π_{T} and the band projectors does not spoil exactness. Thus both Schur subtractions in eqs. (7.333) and (7.334) are $\hat Q_{\mathrm{BRST}}$-exact operators of ghost number 0, establishing eq. (7.335). The vanishing of their physical traces then follows from eq. (7.331). □

Corollary 7.343 (On-shell simplification of physical blocks)**.** *Using Lemma 7.342, one has, in BRST cohomology and on shell,*

$$\mathcal{H}^{TT}_{\rm phys} \;\simeq\; \Pi_{\rm T}\,\mathcal{H}^{TT}\,\Pi_{\rm T}\,, \qquad \mathcal{H}^{\rm mat}_{\rm phys} \;\simeq\; \begin{pmatrix} \mathcal{H}^{hh} & \mathcal{H}^{h\pi} \\ \mathcal{H}^{\pi h} & \mathcal{H}^{\pi\pi} \end{pmatrix}. \tag{7.336}$$

Proof. Start from the Schur-reduced definitions eqs. (7.333) and (7.334). By Lemma 7.342, the Schur subtractions are $\hat{Q}_{\rm BRST}$-exact band operators. Thus, when sandwiched by the physical projector or inserted in band traces with BRST-closed, ghost-number 0 operators, they vanish by the cohomological identity eq. (7.331). This yields the stated equivalences in BRST cohomology. On shell, longitudinal mixings are momentum-suppressed or BRST-exact (cf. eq. (7.318)), so the TT and matter blocks above furnish the physical poles/residues. □

Global vs. gauged G. For *global* breaking, the NG constraints eq. (7.289) hold in the matter block, so that $\mathcal{M}^2_{\pi\pi} = 0$ and $\mathcal{K}_{\pi h}(0) = 0$, and NG poles emerge (cf. eq. (7.290)). If the broken generators are *gauged* (BEH realisation), the would-be NGs enter BRST quartets and decouple from $\mathcal{H}_{\rm phys}$; the physical spectrum is carried by $\mathcal{H}^{TT}_{\rm phys}$ (massive transverse vectors) and the gapped scalar block for h (cf. eqs. (7.298), (7.300) and (7.301)).

Physical propagators and positivity/hyperbolicity. The band-projected physical propagator is

$$\mathbb{G}^{(\lambda_{\rm UV},\ell)}_{\rm phys}(p) := \Big(\mathbb{H}^{(\lambda_{\rm UV},\ell)}_{\rm phys}(p)\Big)^{-1}, \qquad \mathbb{H}^{(\lambda_{\rm UV},\ell)}_{\rm phys} := \mathcal{H}^{TT}_{\rm phys} \oplus \mathcal{H}^{\rm mat}_{\rm phys}, \tag{7.337}$$

with non-negativity of the mass operator and hyperbolicity of the principal symbol guaranteed by eqs. (7.13) and (7.15) (continuity from the symmetric band and the acceptance tests).

Plateau stability. Under admissible UV/IR scans within a certified window, the physical blocks inherit the robustness bounds

$$\partial_{(\ln\lambda_{\rm UV},\,\ln\ell)}\,\mathbb{H}^{(\lambda_{\rm UV},\ell)}_{\rm phys}(p) = \mathcal{O}(\varepsilon_{\rm plat}), \tag{7.338}$$

as a direct corollary of eq. (7.291) and the projector commutators eq. (7.325).

Corollary 7.344 (Nielsen: gauge–parameter independence on the band)**.** *Differentiating the BRST–exact Schur subtractions eq. (7.335) with respect to the gauge parameter ζ (the gauge–fixing sector is $\{\hat{Q}_{\rm BRST}, \Psi_\zeta\}$) gives*

$$\partial_\zeta\,\mathbb{H}^{(\lambda_{\rm UV},\ell)}_{\rm phys}(p) = \Big\{\hat{Q}_{\rm BRST},\, \mathcal{N}^{(\lambda_{\rm UV},\ell)}_\zeta(p)\Big\} \qquad \textit{(on the certified band)}, \tag{7.339}$$

for some band operator $\mathcal{N}_\zeta$. Hence, for any BRST–closed, ghost–number 0 insertion $\mathcal{O}$ (or between physical states),

$$\partial_\zeta\;\mathrm{Tr}_{\mathcal{H}_{\rm phys}}\Big(\mathcal{O}\,\mathbb{H}_{\rm phys}\Big) = \mathrm{Tr}_{\mathcal{H}_{\rm phys}}\Big(\mathcal{O}\,\{\hat{Q}_{\rm BRST}, \mathcal{N}_\zeta\}\Big) = 0 \quad \textit{by eq. (7.331)}.$$

Therefore physical poles and residues extracted from $\mathbb{G}_{\rm phys} = (\mathbb{H}_{\rm phys})^{-1}$ are ζ-independent (Nielsen identity), matching the gauge–scan criteria in eqs. (7.343) and (7.347) and the visibility test eq. (7.260). Combined with eq. (7.338), this independence is stable under admissible UV/IR scans on the plateau.

Proof. By Lemma 7.342, the Schur subtractions are BRST–exact band operators. Since the gauge–fixing sector is $\{\hat{Q}_{\mathrm{BRST}}, \Psi_\zeta\}$, differentiating w.r.t. ζ preserves BRST exactness, yielding eq. (7.339). Inserting a BRST–closed, ghost–number 0 $\mathcal{O}$ and using graded cyclicity on $\mathcal{H}_{\mathrm{phys}}$ (eq. (7.331)) gives zero for the derivative of any band trace, which implies the ζ-independence of on-shell poles and residues. On finite plateaus, residuals are bounded as in eqs. (7.343) and (7.347), with UV/IR stability from eq. (7.338). □

Summary. Equations eqs. (7.320), (7.325), (7.328) and (7.332) implement the (T/L) split and BRST projection on the band. The resulting physical inverse two-point kernel eqs. (7.333), (7.334) and (7.336) furnishes the correct poles/residues for on-shell observables, with NG/BEH structure as appropriate and with plateau-stable UV/IR response eq. (7.338).

3.5 Gauge-parameter independence (Nielsen) and ST identities

Master Nielsen identity on the band. Gauge fixing enters through a BRST-exact term $\{\hat{Q}_{\mathrm{BRST}}, \Psi_\zeta\}$. On the certified band the gauge–parameter derivative of the 1PI functional is itself BRST-exact:

$$\partial_\zeta \Gamma = \left\{\hat{Q}_{\mathrm{BRST}},\, \mathcal{N}_\zeta\right\} \qquad (\text{band; } \mathsf{B}_{\lambda,\ell}\text{-projected}) \,, \tag{7.340}$$

for some band operator $\mathcal{N}_\zeta$ built from Ψ_ζ and the background (cf. the Hessian version eq. (7.339)). Therefore, any BRST-closed, ghost-number 0 matrix element of $\partial_\zeta\Gamma$ vanishes on $\mathcal{H}_{\mathrm{phys}}$ by eq. (7.331).

Pole locations are ζ-independent (Nielsen). Let physical poles be defined by the zeros of the band-projected determinant $\det \mathbb{H}_{\mathrm{phys}}(p^2, \zeta) = 0$ at $p^2 = m_n^2(\zeta)$. Differentiating and using left/right eigenvectors L_n, R_n of $\mathbb{H}_{\mathrm{phys}}$ at the pole gives

$$\partial_\zeta m_n^2 = -\,\frac{L_n^\top \left(\partial_\zeta \mathbb{H}_{\mathrm{phys}}\right) R_n}{L_n^\top \left(\partial_{p^2} \mathbb{H}_{\mathrm{phys}}\right) R_n} \,. \tag{7.341}$$

By eq. (7.339), $\partial_\zeta \mathbb{H}_{\mathrm{phys}} = \{\hat{Q}_{\mathrm{BRST}}, \mathcal{N}_\zeta^{(2)}\}$ on the band, so the numerator is a BRST-exact physical matrix element and vanishes by eq. (7.331). Hence

$$\partial_\zeta m_n^2 = 0 \qquad (\text{Nielsen identity on the band}). \tag{7.342}$$

On finite plateaus, residual deviations are bounded by the plateau budget (cf. eq. (7.338)), yielding the acceptance criterion

$$\left|\partial_\zeta m_n^2\right| = \mathcal{O}(\varepsilon_{\mathrm{plat}}) \,. \tag{7.343}$$

Residues are ζ-independent. Let the residue (wavefunction factor) be

$$Z_n^{-1} = L_n^\top (\partial_{p^2} \mathbb{H}_{\mathrm{phys}}) R_n \Big|_{p^2 = m_n^2}$$

Differentiating and using eqs. (7.331) and (7.339) again shows $\partial_\zeta Z_n = 0$ on the band (and $\partial_\zeta Z_n = \mathcal{O}(\varepsilon_{\mathrm{plat}})$ on finite windows).

From poles to S-matrix (LSZ + Nielsen). Band-projected amputated Green functions inherit eq. (7.340). Combining with LSZ reduction (which uses m_n^2 and Z_n) gives gauge–parameter independence of on-shell S-matrix elements on the band:

$$\partial_\zeta \mathcal{S}_{\rm phys} = 0 \qquad (\text{and } = \mathcal{O}(\varepsilon_{\rm plat}) \text{ on finite plateaus}). \tag{7.344}$$

Slavnov–Taylor identities (constraints and transversality). The ST identity is the functional statement of BRST invariance of Γ and, after differentiation, constrains 1PI kernels and vertices. In particular:

$$\begin{aligned} p_\mu\, \Gamma^{\mu\nu}_{AA}(p) \;&=\; 0 \\ \Rightarrow \quad \Gamma^{\mu\nu}_{AA}(p) &= \Pi^{\mu\nu}_{\rm T}(p)\, \Gamma^{TT}_{AA}(p) \end{aligned} \tag{7.345}$$

(transversality of the physical gauge block),

and for matter–gauge mixing,

$$p_\mu\, \Gamma^{\mu}_{A\Phi\Phi}(p; q, -q-p) = \Gamma_{\Phi\Phi}(q+p) - \Gamma_{\Phi\Phi}(q) \tag{7.346}$$

(tree Ward form; background-field version is identical with $A \to \widehat{A}$).

Equations eqs. (7.345) and (7.346) ensure that the physical TT block used in eq. (7.333) is the only gauge block contributing on shell, and that longitudinal mixings are either momentum–suppressed (cf. eq. (7.318)) or BRST-exact (cf. §3.3).

Background-field gauge (optional strengthening). If a background-field gauge is used, background Ward identities imply manifest gauge covariance of background amplitudes and fix renormalisation relations among couplings and field strengths (e.g., $Z_g = Z_{\widehat{A}}^{-1/2}$), tightening the ST constraints without altering the on-band conclusions above.

Operational acceptance (gauge scans). For each reported physical pole m_n^2 and residue Z_n at a fixed plateau, perform a gauge scan $\zeta \mapsto (\, m_n^2(\zeta), Z_n(\zeta)\,)$ and verify

$$\left|\partial_\zeta m_n^2\right| \le \varepsilon_{\rm plat}, \qquad \left|\partial_\zeta Z_n\right| \le \varepsilon_{\rm plat}, \tag{7.347}$$

together with a law-channel check $\partial_\zeta \Lambda_{\rm law} = \mathcal{O}(\varepsilon_{\rm plat})$ (cf. eq. (7.260)). Passing eq. (7.347) certifies that pole locations/residues and the derived on-shell observables are gauge-parameter independent within the plateau tolerance.

3.6 Practical algorithm (run-sheet)

1. **Choose the broken background φ.** Prepare φ as in Section 7.7.2 (seed protocol and ordered limits). Check stationarity on the band, $\delta\Gamma/\delta\Phi|_{\Phi=\varphi} = 0$ (eq. (7.312)), and record the coset direction(s) used (cf. $\mathcal{O}_X$ in Step 2).

2. **Assemble $\mathcal{H}_{\rm band}$.** Compute the quadratic kernel $\mathcal{H}_{\rm band}$ (eq. (7.311)), keeping explicit dependence on the band filter $\mathsf{B}_{\lambda,\ell} = K_{\rm IR}^{(\ell)} P_x(\lambda_{\rm UV})$ (eq. (7.313)) and on gauge parameters ζ (for Nielsen scans). Decompose each block into principal symbol and mass/mixing pieces (eq. (7.315)).

3. **T/L decomposition (vector sector).** In momentum space apply the projectors $\Pi_{\mathrm{T}/L}$ (eq. (7.320)) and rewrite $A_\mu = A^T_\mu + A^L_\mu$ (eq. (7.322)); in real space use the IR-regulated Helmholtz form (eq. (7.324)) and handle zero modes via $\Delta^{-1}_{\ell,V}$ (eq. (7.326)). Verify band commutators (eq. (7.325)).

4. **BRST sort (doublets & quartets).** Identify the BRST doublets $(\bar{c}, b)$ and $(A^{\mathrm{L}}, \bar{D}c)$ via the linearised algebra (eq. (7.328)); confirm that the ghost/longitudinal kernel is BRST-exact (eq. (7.329)) and that single-L mixings vanish at $p = 0$ (eq. (7.318)). Cross-check quartet decoupling with Section 7.2.6.

5. **Project to $\mathcal{H}_{\mathrm{phys}}$ and form physical blocks.** Apply the physical projector properties (eq. (7.330)) and construct the Schur-reduced kernels $\mathcal{H}^{TT}_{\mathrm{phys}}$ and $\mathcal{H}^{\mathrm{mat}}_{\mathrm{phys}}$ (eqs. (7.333) and (7.334)). Use the lemma Lemma 7.342 to drop BRST-exact subtractions and adopt the on-shell simplification (eq. (7.336)). The band-projected physical propagator is then $\mathbb{G}_{\mathrm{phys}} = (\mathbb{H}_{\mathrm{phys}})^{-1}$ (eq. (7.337)).

6. **Health checks (band-projected).** Verify principal-symbol hyperbolicity and non-negativity of the physical mass operator (continuity from Step 1; eqs. (7.13) and (7.15)). For bookkeeping, define the minimal hyperbolicity margin and mass gap

$$\delta_{\mathrm{hyp}} := \inf_{|p|\gg 1} \lambda_{\min}\Big(\operatorname{sym} \mathcal{K}_{\mathrm{phys}}(p)\Big), \qquad m^2_{\min} := \min_n m^2_n \geq 0, \tag{7.348}$$

   and require $\delta_{\mathrm{hyp}} > 0$ and $m^2_{\min} \geq 0$ on the plateau.

7. **Nielsen & ST verification.** Perform a gauge-parameter scan $\zeta \mapsto \mathbb{H}_{\mathrm{phys}}$ and confirm $\partial_\zeta \mathbb{H}_{\mathrm{phys}} = \{\hat{Q}_{\mathrm{BRST}}, \mathcal{N}^{(2)}_\zeta\}$ (eq. (7.339)), whence $\partial_\zeta m^2_n = 0$ and $\partial_\zeta Z_n = 0$ within tolerance (eqs. (7.343) and (7.347)). Check ST identities for two- and three-point structures (eqs. (7.345) and (7.346)).

8. **Plateau certification and robustness.** Confirm small band derivatives of the reduced kernel (eq. (7.338)) and compliance with the visibility/plateau bounds (eq. (7.137)). Where applicable, swap admissible regulators and verify stability (cf. eq. (7.325)).

Outputs to report. List: (i) chosen φ and seed profile, (ii) certified plateau window and $\varepsilon_{\mathrm{plat}}$, (iii) m^2_n, Z_n with gauge-scan slopes, (iv) hyperbolicity/mass margins (eq. (7.348)), and (v) any residual BRST-exact diagnostics observed (all should vanish within numerical tolerance).

3.7 Acceptance tests (physics-facing)

- **Quartet decoupling (null test).** Using a BRST-closed, ghost-number 0 probe $\mathcal{O}$ on the band, verify

$$\Big|\operatorname{Tr}_{\mathcal{H}_{\mathrm{phys}}}\Big(\mathcal{O}\,(\mathcal{H}^{LL} \oplus \mathcal{H}^{c\bar{c}} \oplus \mathcal{H}^{\bar{c}c} \oplus \mathcal{H}^{bb})\Big)\Big| \leq \varepsilon_{\mathrm{plat}} \qquad \text{(expected = 0 by eq. (7.332))}. \tag{7.349}$$

- **Transverse dominance (spin-1 sector).** Define the transversality ratio for the band-projected two-point function

$$\mathcal{T}(p) := \frac{p_\mu \, \mathbb{G}^{\mu\nu}_{AA}(p) \, p_\nu}{p^2 \, \mathrm{tr}\Big(\Pi_{\mathrm{T}} \, \mathbb{G}_{AA}(p)\Big)} \,, \tag{7.350}$$

  and require $|\mathcal{T}(p)| \le \varepsilon_{\mathrm{plat}}$ across the plateau support. Equivalently, the Källén–Lehmann (band) weight carries no A^L component, consistent with eq. (7.345) and the construction in eq. (7.333).

- **Gauge independence (Nielsen scans).** For every reported pole m_n^2 and residue Z_n,

$$\Big|\partial_\zeta m_n^2\Big| \le \varepsilon_{\mathrm{plat}}, \qquad \Big|\partial_\zeta Z_n\Big| \le \varepsilon_{\mathrm{plat}} \tag{7.351}$$

  (cf. eqs. (7.343) and (7.347)); also check the law-channel slope $\partial_\zeta \Lambda_{\mathrm{law}} = \mathcal{O}(\varepsilon_{\mathrm{plat}})$ (eq. (7.260)) as a control.

- **Mass/hyperbolicity.** The physical mass operator is non-negative and the principal symbol hyperbolic on the band: compliance with eqs. (7.13) and (7.15), and margins $\delta_{\mathrm{hyp}} > 0$, $m^2_{\mathrm{min}} \ge 0$ (eq. (7.348)) are reported.

- **Plateau robustness.** Derivatives of the reduced kernel and of extracted observables w.r.t. $(\ln \lambda_{\mathrm{UV}}, \ln \ell)$ are within $\mathcal{O}(\varepsilon_{\mathrm{plat}})$ (eq. (7.338)); regulator permutations that commute with G and BRST leave results invariant within tolerance (cf. eq. (7.325)).

3.8 Failure modes and diagnostics

- *Residual longitudinal leakage in physical correlators.* **Symptom:** nonzero longitudinal weight in band-projected two-point functions. **Test:** the transversality ratio $\mathcal{T}(p)$ in eq. (7.350) fails to stay within $\varepsilon_{\mathrm{plat}}$, or the quartet null test eq. (7.349) is violated. **Likely cause:** BRST misalignment or off-shell evaluation. **Fix:** re-enforce on-shell ST/Nielsen conditions (cf. eqs. (7.339) and (7.345)), recompute the BRST sort (eqs. (7.328) and (7.329)), and redo the Schur reduction (§3.4) using the lemma Lemma 7.342. Confirm $\mathsf{P}_{\mathrm{phys}}$ commutators eq. (7.330) and band commutators eq. (7.325).

- *Gauge-parameter drift of physical poles/residues.* **Symptom:** $|\partial_\zeta m_n^2|, |\partial_\zeta Z_n| \not\le \varepsilon_{\mathrm{plat}}$ (cf. eqs. (7.343) and (7.351)). **Likely cause:** evaluation off a certified plateau or missing/incorrect counterterms in Γ. **Fix:** restore plateau conditions (eq. (7.338)) and check algebraic renormalisation constraints (Nielsen master identity eq. (7.340) and ST relations eqs. (7.345) and (7.346)); update the gauge-fixing fermion Ψ_ζ if needed.

- *Schur instability / singular $\mathcal{H}^{LL}$.* **Symptom:** ill-conditioned or non-invertible longitudinal block obstructs the Schur step. **Diagnostic:** monitor the condition number

$$\kappa_{LL} := \frac{\lambda_{\max}(\mathcal{H}^{LL\dagger}\mathcal{H}^{LL})}{\lambda_{\min}(\mathcal{H}^{LL\dagger}\mathcal{H}^{LL})} \,. \tag{7.352}$$

  Likely cause: inadequate IR treatment of zero modes or an incompatible gauge choice. **Fix:** handle zero modes with the band pseudoinverse $\Delta^{-1}_{\ell,V}$ (eq. (7.326)), switch to an

admissible R_ξ family, and re-verify $\mathcal{H}^{TL/LT}(0) = 0$ (eq. (7.318)); then reapply eqs. (7.333) and (7.334).

- *Tachyonic/elliptic pathologies.* **Symptom:** violation of mass non-negativity or hyperbolicity ($m^2_{\min} < 0$ or $\delta_{\rm hyp} \le 0$; cf. eq. (7.348)). **Likely cause:** unstable background φ or seed/order-of-limits error. **Fix:** revisit the background preparation protocol (Step 2), correct seed ordering, and test regulator stability as in eq. (7.319); if needed, reselect v and recompute $\mathcal{H}_{\rm band}$ (eq. (7.311)).

- *Uncontrolled h–π mixing (global G).* **Symptom:** finite $\mathcal{K}_{\pi h}(0)$ or $\mathcal{M}^2_{\pi\pi} \neq 0$ contradicting the NG constraints eq. (7.316). **Likely cause:** broken Ward identity mishandled or coset parameterisation inconsistent with the chosen vacuum. **Fix:** recheck global Ward identities (§2.4) and the coset chart (cf. §2.2), then re-enforce the constraints eq. (7.316) at the level of $\mathcal{H}^{\rm mat}_{\rm phys}$ (eq. (7.336)).

- *Plateau drift / regulator-order sensitivity.* **Symptom:** derivatives $\partial_{(\ln\lambda_{\rm UV},\,\ln\ell)}\mathbb{H}_{\rm phys}$ exceed $\mathcal{O}(\varepsilon_{\rm plat})$ (eq. (7.338)), or results vary with admissible permutations of band projectors beyond tolerance. **Diagnostic:** define the ordering sensitivity

$$\Delta_{\rm ord} := \left\| \mathbb{H}^{(P_x K_{\rm IR})}_{\rm phys} - \mathbb{H}^{(K_{\rm IR} P_x)}_{\rm phys} \right\|_{\rm phys} \le \varepsilon_{\rm plat}\,, \tag{7.353}$$

  and check commutators eq. (7.325). **Fix:** tighten the plateau window, retune regulator profiles, and ensure all band operations commute with BRST and symmetry generators.

- *Anomalous identity ledger (rare).* **Symptom:** persistent violation of Ward/ST checks not cured by plateau/gauge fixes. **Likely cause:** anomaly contamination in the probed band. **Fix:** revisit the exact-identity sector (Section 7.5) and exclude the anomalous window before re-running the reduction.

Outcome. After these diagnostics, the quadratic dynamics *on the band* is reduced to the BRST-physical blocks with well-posed propagation and gauge-parameter–independent on-shell content: $\mathbb{H}_{\rm phys}$ and $\mathbb{G}_{\rm phys} = (\mathbb{H}_{\rm phys})^{-1}$ (eq. (7.337)) satisfy eqs. (7.338), (7.339) and (7.345). This furnishes the input for Step 4 (readout of visibilities and NG soft structure), where the block-reduced kernel links to interferometric diagnostics (Section 7.4).

7.7.4 Read Visibilities: Verify $|\Lambda_{\rm law}| \approx 1$; Volume Decay of $|\Lambda_{\rm state}|$; Phase–twist Interferometry for NG Soft Limits

Aim. Having frozen the law (Section 7.7.1) and prepared an asymmetric state (Section 7.7.2), and with the band Hessian reduced to physical blocks (Section 7.7.3), we now execute the interferometric readout on certified plateaus (Section 7.3). The goals are: (i) confirm the law–channel coherence $|\Lambda_{\rm law}| \simeq 1$ within plateau tolerances (eqs. (7.137) and (7.139)); (ii) demonstrate *volume–decay* of the state visibility $|\Lambda_{\rm state}|$ at fixed plateau (Quadrant II signature); and (iii) use *phase–twist interferometry* to extract NG soft structure from broken currents via Ward/Nielsen–protected relations (Sections 7.4 and 7.5; [91, 117, 442, 82]).

4.1 Readout pipeline (band–projected coherence)

We use the band–projected coherence functional of Section 7.4.1 and eq. (7.115), with the same UV projector $P_x(\lambda_{\mathrm{UV}})$ and IR window $K_{\mathrm{IR}}^{(\ell)}$ as in Section 7.3. Write the band filter as

$$\mathsf{B}_{\lambda,\ell} \;:=\; P_x(\lambda_{\mathrm{UV}})\, K_{\mathrm{IR}}^{(\ell)}\,, \qquad [\hat{Q}_{\mathrm{BRST}}, \mathsf{B}_{\lambda,\ell}] = [\mathsf{J}_\alpha, \mathsf{B}_{\lambda,\ell}] = 0 \;\; \text{on } \mathcal{H}_{\mathrm{phys}} \; (\textit{eq.}\ (7.244)). \tag{7.354}$$

For a given channel $\alpha \in \{\mathrm{law}, \mathrm{state}\}$ with branch unitaries $(U_1^{(\alpha)}, U_2^{(\alpha)})$ obeying the commutation/BRST constraints (Section 7.7.1), define the band–projected visibility

$$\Lambda_\alpha(\lambda_{\mathrm{UV}}, \ell) = \frac{\Big|\mathrm{Tr}_{\mathcal{H}_{\mathrm{phys}}}\Big(\mathsf{B}_{\lambda,\ell}\, U_1^{(\alpha)\dagger} U_2^{(\alpha)}\, \rho_\alpha\Big)\Big|}{\mathrm{Tr}_{\mathcal{H}_{\mathrm{phys}}}\Big(\mathsf{B}_{\lambda,\ell}\, \rho_\alpha\Big)}\,. \tag{7.355}$$

For $\alpha = \mathrm{law}$ this specialises to eq. (7.254). For the *state* channel we keep the explicit volume dependence (system size V) in the preparation and readout:

$$\Lambda_{\mathrm{state}}(\lambda_{\mathrm{UV}}, \ell; V) = \frac{\Big|\mathrm{Tr}_{\mathcal{H}_{\mathrm{phys}}}\Big(\mathsf{B}_{\lambda,\ell}\, U_1^{(\mathrm{state})\dagger} U_2^{(\mathrm{state})}\, \rho_{\mathrm{ref}}(V)\Big)\Big|}{\mathrm{Tr}_{\mathcal{H}_{\mathrm{phys}}}\Big(\mathsf{B}_{\lambda,\ell}\, \rho_{\mathrm{ref}}(V)\Big)}\,, \tag{7.356}$$

where $\rho_{\mathrm{ref}}(V)$ denotes the branchwise prepared band state at volume V (constructed in Section 7.7.2). At fixed certified plateaus, Quadrant II implies $|\Lambda_{\mathrm{law}}| \approx 1$ (cf. eqs. (7.257) and (7.259)) and $\lim_{V\to\infty} |\Lambda_{\mathrm{state}}| = 0$ (§2.6).

Integrated readouts (optional). To improve signal and track robustness across the window, we also use weighted, band–integrated visibilities. Let $\mu(\lambda_{\mathrm{UV}}, \ell)$ be an admissible probability density supported on a certified plateau (Section 7.4.5). Define

$$\overline{\Lambda}_\alpha[\mu] := \int d(\ln \lambda_{\mathrm{UV}})\, d(\ln \ell)\; \mu(\lambda_{\mathrm{UV}}, \ell)\, \Lambda_\alpha(\lambda_{\mathrm{UV}}, \ell)\,. \tag{7.357}$$

By eq. (7.137) and eq. (7.259), integrated readouts inherit the same tolerances, i.e. $\overline{\Lambda}_{\mathrm{law}}[\mu] = 1 + \mathcal{O}(\varepsilon_{\mathrm{plat}})$ and the finite–size $|\overline{\Lambda}_{\mathrm{state}}[\mu]|$ decays in V at fixed μ.

Reporting. For each run we report: (i) the plateau region and $\varepsilon_{\mathrm{plat}}$ used; (ii) $\Lambda_{\mathrm{law}}(\lambda_{\mathrm{UV}}, \ell)$ and/or $\overline{\Lambda}_{\mathrm{law}}[\mu]$ with the Nielsen slope $\partial_\zeta \Lambda_{\mathrm{law}}$ (eq. (7.260)); (iii) $\Lambda_{\mathrm{state}}(\lambda_{\mathrm{UV}}, \ell; V)$ (and optionally $\overline{\Lambda}_{\mathrm{state}}[\mu; V]$) at several V, together with finite–size slopes $-V^{-1} \ln |\Lambda_{\mathrm{state}}|$ at fixed plateaus (cf. §2.6); and (iv) regulator–order checks as in eq. (7.265).

4.2 Law–channel invariant (calibration and guardrail)

With a *symmetric* calibration state ρ_0 and law–branch reparametrisations satisfying $[U_b^{(\mathrm{law})}, \hat{H}_0] = [U_b^{(\mathrm{law})}, \hat{Q}_{\mathrm{BRST}}] = [U_b^{(\mathrm{law})}, \mathsf{J}_\alpha] = 0$ (Section 7.7.1), the exact on–shell identities (Section 7.5) together with the band commutators (eq. (7.244)) give the ideal law–channel value $\Lambda_{\mathrm{law}} = 1$ (eq. (7.257)). On a certified plateau this becomes

$$\Lambda_{\mathrm{law}}(\lambda_{\mathrm{UV}}, \ell) = 1 + \mathcal{O}(\varepsilon_{\mathrm{plat}}) \qquad \text{with } \varepsilon_{\mathrm{plat}} \text{ bounded by } \textit{eq.}\ (7.137). \tag{7.358}$$

Operational guardrail (explicit bound). Combining the regulator–order budget eq. (7.258) with the projector–commutator acceptance eq. (7.265) yields

$$\left|1 - \Lambda_{\mathrm{law}}(\lambda_{\mathrm{UV}}, \ell)\right| \;\le\; C_{\mathrm{plat}}\, \varepsilon_{\mathrm{plat}} \;+\; C_{\mathrm{reg}}\, \varepsilon_{\mathrm{reg}}, \tag{7.359}$$

for channel–independent constants $C_{\mathrm{plat}}, C_{\mathrm{reg}} = \mathcal{O}(1)$ determined by the window and the chosen norm on $\mathcal{H}_{\mathrm{phys}}$. Hence a systematic plateau–level deficit that exceeds the right–hand side violates at least one Quadrant II hypothesis (hidden explicit breaking, BRST misalignment, or contamination by state operations); triage per Sections 7.7.1 and 7.7.3.

Calibration checklist. At the plateau used for SSB runs, report: (i) branch–swap invariance of Λ_{law}; (ii) Nielsen slope $\partial_\zeta \Lambda_{\mathrm{law}}$ within eq. (7.260); (iii) regulator–order permutation within $\mathcal{O}(\varepsilon_{\mathrm{plat}})$; (iv) compliance with eq. (7.359). These certify that the law interferometer acts as a *guardrail* for the state–channel analysis.

4.3 State visibility and finite–size scaling

For a broken preparation ρ_{br} (Section 7.7.2), the state–branch unitaries compare either *two distinct vacua* related by G or *two preparations* on one vacuum:

$$\Lambda_{\mathrm{state}}(\lambda_{\mathrm{UV}}, \ell; V) = \frac{\left|\mathrm{Tr}_{\mathcal{H}_{\mathrm{phys}}}\left(P_x K_{\mathrm{IR}}^{(\ell)}\, U_1^{(\mathrm{state})\dagger} U_2^{(\mathrm{state})}\, \rho_{\mathrm{br}}\right)\right|}{\mathrm{Tr}_{\mathcal{H}_{\mathrm{phys}}}\left(P_x K_{\mathrm{IR}}^{(\ell)}\, \rho_{\mathrm{br}}\right)}, \tag{7.360}$$

(cf. eq. (7.356)). Quadrant II requires the *two–channel* signature at fixed plateau:

$$\left|\Lambda_{\mathrm{law}}(\lambda_{\mathrm{UV}}, \ell)\right| \simeq 1, \qquad \lim_{V\to\infty} \left|\Lambda_{\mathrm{state}}(\lambda_{\mathrm{UV}}, \ell; V)\right| = 0, \tag{7.361}$$

(cf. eqs. (7.139), (7.257) and (7.259)).

Rate function and scaling collapse. Define the (band, plateau–fixed) rate function

$$I_V(\lambda_{\mathrm{UV}}, \ell) := -\frac{1}{V} \ln \left|\Lambda_{\mathrm{state}}(\lambda_{\mathrm{UV}}, \ell; V)\right|. \tag{7.362}$$

A broken phase exhibits a nonzero asymptotic rate

$$\lim_{V\to\infty} I_V(\lambda_{\mathrm{UV}}, \ell) =: I_\star(\ell) > 0, \tag{7.363}$$

with $I_\star$ depending on NG stiffness/interfacial tension and the IR window ℓ. When interfaces dominate (e.g. fixed–twist preparations), it is convenient to replace V in eq. (7.362) by an effective size $V_{\mathrm{eff}} \propto L^{d-1}$ (sample area), which we will indicate explicitly when used.

Common decay laws (to be reported, not assumed). On certified plateaus, the large–size envelope of $|\Lambda_{\text{state}}|$ typically follows

$$\begin{aligned} \left|\Lambda_{\text{state}}\right| &\sim \exp\Big[-\kappa(\ell)\,V\Big] && \text{(bulk–dominated)},\\ \left|\Lambda_{\text{state}}\right| &\sim \exp\Big[-\sigma(\ell)\,L^{d-1}\Big] && \text{(interface–dominated)}, \end{aligned} \tag{7.364}$$

or, more generally, a stretched–exponential form with an empirically determined exponent. We *report* the best supported law together with confidence bands and the underlying plateau diagnostics, rather than imposing a priori ν.

Acceptance and reporting. At a fixed certified plateau:

1. **Two–channel check:** verify $|\Lambda_{\text{law}}| = 1 + \mathcal{O}(\varepsilon_{\text{plat}})$ (eq. (7.358)) simultaneously with decay of $|\Lambda_{\text{state}}|$.
2. **Rate convergence:** show I_V from eq. (7.362) is monotone (up to finite–size noise) and converges to a plateau–stable $I_\star > 0$ within uncertainty (criterion eq. (7.363)).
3. **Model fit (optional):** fit $\kappa(\ell)$ or $\sigma(\ell)$ in eq. (7.364) and display a scaling collapse across V (or L^{d-1}) with ℓ held within the same plateau.
4. **Robustness:** repeat items 1–3 under admissible regulator permutations and small gauge changes; accept only if variations remain within $\mathcal{O}(\varepsilon_{\text{plat}})$ and Nielsen slopes satisfy eq. (7.260).

4.4 Phase–twist interferometry: construction

Setup and band/BRST admissibility (general d). Let J^μ_a be a conserved Noether current for a *broken* global generator T_a in $D = d{+}1$ spacetime dimensions ($\partial_\mu J^\mu_a = 0$ in the symmetric law; Section 7.5). Choose a smooth spatial smearing $F(\mathbf{x})$ supported inside the IR window and a compact time window $f(t)$; define the band–compatible smeared charge

$$Q_a[F] := \int d^d x\, F(\mathbf{x})\, J^0_a(t,\mathbf{x}), \qquad \Big[\hat{Q}_{\text{BRST}}, Q_a[F]\Big] = 0, \quad \Big[\mathsf{B}_{\lambda,\ell}, Q_a[F]\Big] = 0, \tag{7.365}$$

with $\mathsf{B}_{\lambda,\ell} := P_x(\lambda_{\text{UV}})K^{(\ell)}_{\text{IR}}$ as in eq. (7.354) and band closure eq. (7.244). Implement *opposite* twists on the two state branches,

$$U_1^{(\text{state})}(\phi) = e^{+\mathrm{i}\phi\, Q_a[F]}, \qquad U_2^{(\text{state})}(\phi) = e^{-\mathrm{i}\phi\, Q_a[F]}. \tag{7.366}$$

Characteristic function (band version). The ϕ–dependent visibility is the modulus of the characteristic function of band–projected charge fluctuations,

$$\Lambda_{\text{state}}(\phi;\lambda_{\text{UV}},\ell) = \frac{\Big|\operatorname{Tr}_{\mathcal{H}_{\text{phys}}}\Big(\mathsf{B}_{\lambda,\ell}\, e^{\,2\mathrm{i}\phi\, Q_a[F]}\,\rho_{\text{br}}\Big)\Big|}{\operatorname{Tr}_{\mathcal{H}_{\text{phys}}}\Big(\mathsf{B}_{\lambda,\ell}\,\rho_{\text{br}}\Big)}, \qquad \Lambda_{\text{state}}(\phi) = \Lambda_{\text{state}}(-\phi). \tag{7.367}$$

Small-ϕ curvature and charge variance. A cumulant expansion (band trace, BRST–closed sector) gives

$$\ln \Lambda_{\text{state}}(\phi; \lambda_{\text{UV}}, \ell) = -\,\phi^2\, \mathrm{Var}_{\text{band}}\Big[Q_a[F]\Big] \;+\; \mathcal{O}(\phi^4), \tag{7.368}$$

so the curvature at $\phi = 0$ is

$$\mathrm{Var}_{\text{band}}\Big[Q_a[F]\Big] = -\,\tfrac{1}{2}\,\frac{\partial^2}{\partial\phi^2}\,\ln \Lambda_{\text{state}}(\phi; \lambda_{\text{UV}}, \ell)\Big|_{\phi=0}. \tag{7.369}$$

Within the measurement window $f(t)$,

$$\mathrm{Var}_{\text{band}}\Big[Q_a[F]\Big] = \int dt\, dt'\, f(t) f(t') \int d^d x\, d^d y\, F(\mathbf{x}) F(\mathbf{y})\, \Big\langle J_a^0(t,\mathbf{x})\, J_a^0(t',\mathbf{y})\Big\rangle_{c,\text{band}}, \tag{7.370}$$

with the connected, BRST–physical correlator; on shell and on a certified plateau this is Nielsen–stable (Sections 7.4.5 and 7.5).

Scaling and susceptibility. For bulk–supported F one typically finds

$$\begin{aligned} \mathrm{Var}_{\text{band}}\Big[Q_a[F]\Big] \;&=\; \chi_a(\ell; F)\, V_{\text{eff}} \;+\; o(V_{\text{eff}}), \\ \Rightarrow \qquad \Lambda_{\text{state}}(\phi) \;&\approx\; \exp\Big[-\,\phi^2\, \chi_a(\ell; F)\, V_{\text{eff}}\Big] \quad (\phi \to 0), \end{aligned} \tag{7.371}$$

where V_{eff} is the effective spatial size determined by the support of F and the IR window (bulk: $V \propto L^d$; interface–dominated: $V_{\text{eff}} \propto L^{d-1}$).

Soft–momentum targeting (NG extraction) in $\mathbb{R}^d$. To probe NG soft structure, use momentum–selective twists

$$F_{\mathbf{q}}(\mathbf{x}) = g_\ell(\mathbf{x})\, \cos(\mathbf{q}\cdot\mathbf{x}), \qquad \mathbf{q} \in \mathbb{R}^d, \quad |\mathbf{q}|\ell \ll 1, \tag{7.372}$$

with g_ℓ a smooth envelope inside $K_{\text{IR}}^{(\ell)}$. Then (7.370) determines a static structure factor

$$\chi_a(\ell; \mathbf{q}) := \int dt\, dt'\, f(t) f(t') \int d^d x\, d^d y\, g_\ell(\mathbf{x}) g_\ell(\mathbf{y})\, e^{\,i\mathbf{q}\cdot(\mathbf{x}-\mathbf{y})} \Big\langle J_a^0(t,\mathbf{x})\, J_a^0(t',\mathbf{y})\Big\rangle_{c,\text{band}}. \tag{7.373}$$

Ward identities for a broken global symmetry imply an NG pole in the time–ordered correlator, which translates into the soft law

$$\chi_a(\ell; \mathbf{q}) \;\xrightarrow[|\mathbf{q}|\to 0]{}\; \frac{F_a^2(\ell)}{\mathbf{q}^{\,2}} \;+\; \text{regular}, \tag{7.374}$$

with $F_a(\ell)$ an IR–windowed decay constant (cf. Sections 7.5 and 7.7.3). From (7.368)–(7.373), the ϕ^2–curvature of $\ln \Lambda_{\text{state}}(\phi; \lambda_{\text{UV}}, \ell)$ at small $|\mathbf{q}|$ directly yields $\chi_a(\ell; \mathbf{q})$ and hence the $1/\mathbf{q}^{\,2}$ NG diagnostic.

Acceptance checks (phase–twist run).

1. **Band/BRST compatibility:** verify $[\mathsf{B}_{\lambda,\ell}, Q_a[F_{\mathbf{q}}]] = 0 = [\hat{Q}_{\mathrm{BRST}}, Q_a[F_{\mathbf{q}}]]$ (cf. eq. (7.365)).
2. **Evenness and curvature:** confirm $\Lambda_{\mathrm{state}}(\phi) = \Lambda_{\mathrm{state}}(-\phi)$ and extract $\mathrm{Var}_{\mathrm{band}}[Q_a[F_{\mathbf{q}}]]$ via (7.369) in a regime where $\mathcal{O}(\phi^4)$ is negligible.
3. **Soft law:** demonstrate $\chi_a(\ell;\mathbf{q}) \sim F_a^2/\mathbf{q}^2$ over the accessible soft window, with regulator and gauge scans within $\mathcal{O}(\varepsilon_{\mathrm{plat}})$ (Nielsen stability).
4. **Two–channel guardrail:** record $|\Lambda_{\mathrm{law}}| = 1 + \mathcal{O}(\varepsilon_{\mathrm{plat}})$ on the same plateau while the phase–twist data are taken (eq. (7.358)).

4.5 Soft limits and Ward/Nielsen protection

Soft pole in current correlators (global G, general d). Fourier-transform the conserved broken-current correlator in $D = d+1$ dimensions. For a spontaneously broken *global* generator T_a one finds, in the EFT/NG regime and on shell,

$$\left\langle J_a^\mu J_b^\nu \right\rangle(\omega,\mathbf{q}) \;\sim\; \frac{F_a(\ell)\,F_b(\ell)}{\omega^2 - c^2\mathbf{q}^2 + \mathrm{i}0^+}\, q^\mu q^\nu \;+\; \text{regular}, \tag{7.375}$$

where $F_a(\ell)$ is the IR–windowed decay constant (cf. eq. (7.374)) and c the NG velocity ([91]). This reproduces the static soft law eq. (7.374) for the equal–time structure factor.

Variance in momentum space and volume scaling. With the band–compatible twist of §4.4 and envelope $F(\mathbf{x})$, define its Fourier profile $F_{\mathbf{q}} = \int d^dx\, e^{-\mathrm{i}\mathbf{q}\cdot\mathbf{x}} F(\mathbf{x})$. At equal times (within the compact window f) the band variance is

$$\mathrm{Var}_{\mathrm{band}}\Big[Q_a[F]\Big] = \int \frac{d^dq}{(2\pi)^d}\, \Big|F_{\mathbf{q}}\Big|^2\, S_{aa}(\ell;\mathbf{q}), \tag{7.376}$$

$$S_{aa}(\ell;\mathbf{q}) := \int dt\, dt'\, f(t) f(t') \,\widetilde{\left\langle J_a^0 J_a^0 \right\rangle}_{c,\mathrm{band}}(\omega{=}0,\mathbf{q}). \tag{7.377}$$

Using eq. (7.375) (or equivalently eq. (7.374)), $S_{aa}(\ell;\mathbf{q}) \sim F_a^2(\ell)/\mathbf{q}^2$ for $|\mathbf{q}| \to 0$, so for bulk–supported F (width $\sim L$) the integral scales with the effective size,

$$\mathrm{Var}_{\mathrm{band}}\Big[Q_a[F]\Big] = \chi_a(\ell;F)\, V_{\mathrm{eff}} + o(V_{\mathrm{eff}}), \tag{7.378}$$

in agreement with eq. (7.371). Here $V_{\mathrm{eff}} \propto L^d$ for bulk envelopes and $V_{\mathrm{eff}} \propto L^{d-1}$ for interface–dominated preparations; $\chi_a(\ell;F)$ is the band susceptibility defined by the current–current correlator.

Small–ϕ visibility and the Quadrant II hallmark. Inserting eq. (7.378) into the cumulant relation eq. (7.368)–eq. (7.369) gives

$$\Lambda_{\mathrm{state}}(\phi;\lambda_{\mathrm{UV}},\ell;V) \;\approx\; \exp\Big[-\,\phi^2\,\chi_a(\ell;F)\,V_{\mathrm{eff}}\Big] \quad (\phi \to 0), \tag{7.379}$$

demonstrating the *volume–decay* signature of Quadrant II at fixed certified plateaus.

Soft insertions and factorisation. Differentiating the phase–twist visibility at $\phi = 0$ inserts $Q_a[F]$ in the band trace. Ward identities for broken global symmetries imply that adding a *soft* NG leg $(\omega, \mathbf{q} \to 0)$ to any on–shell matrix element factors through the action of T_a on external states (soft theorem), and eq. (7.376)–eq. (7.374) provide the corresponding static limit of the current–current sector that the interferometer measures.

Nielsen protection (gauge independence). Because the twist is built from a BRST–closed, gauge–invariant $Q_a[F]$ that commutes with the band filter, the gauge–parameter derivative of the reduced kernel is BRST–exact (eq. (7.339)). Therefore the curvature extracted from $\ln \Lambda_{\text{state}}(\phi)$ (and the inferred χ_a) is ζ–independent on shell within plateau tolerances, matching the scan criterion eq. (7.260) and stability eq. (7.338).

Scope (gauged vs. global). The construction above applies to *global* broken charges. For gauged generators the would–be NG mode is reorganised (BEH mechanism) and the gauge current is not a suitable BRST–closed insertion; use a gauge–invariant global charge (if present). Dimensional constraints (e.g. Coleman/Mermin–Wagner in low D) must also be respected when interpreting the soft limit.

4.6 Gauged (BEH) case: dressed twists and diagnostics

Why dress the twist (general d). If the broken generators are *gauged*, would–be NG modes are reorganised (BEH) into massive transverse vectors (Sections 7.7.2 and 7.7.3), and Elitzur's theorem forbids a gauge–variant charge insertion. The naive smeared charge $\int d^d x\, F(\mathbf{x}) J_a^0$ is not gauge–invariant; instead we construct a *dressed* charge that is BRST–closed and commutes with the band filter.

Dressed charge (band/BRST–compatible). Let $F(\mathbf{x})$ be a smooth envelope supported inside the IR window $K_{\text{IR}}^{(\ell)}$, and let $\Omega \subset \mathbb{R}^d$ be the region it selects (e.g. $F \approx 1$ on Ω, tapered near $\partial\Omega$). Introduce a gauge–invariant $(d-1)$–form (spatial) dressing current $\mathcal{W}_a^i(\mathbf{x})$ supported near $\partial\Omega$ so that the composite

$$Q_a^{\text{dress}}[F, \Sigma] := \int_\Omega d^d x\, F(\mathbf{x})\, J_a^0(t, \mathbf{x}) \;+\; \int_{\Sigma = \partial\Omega} dS_i(\mathbf{y})\, \mathcal{W}_a^i(t, \mathbf{y}) \tag{7.380}$$

is BRST–closed and band–compatible:

$$[\hat{Q}_{\text{BRST}}, Q_a^{\text{dress}}] = 0, \qquad [\mathsf{B}_{\lambda,\ell}, Q_a^{\text{dress}}] = 0, \qquad \mathsf{B}_{\lambda,\ell} := P_x(\lambda_{\text{UV}}) K_{\text{IR}}^{(\ell)} \;\; (\textit{eq. (7.354)}). \tag{7.381}$$

Concretely, $\mathcal{W}_a$ can be chosen so that its gauge variation cancels that of the volume term (a Wilson–type surface dressing), ensuring gauge invariance on $\mathcal{H}_{\text{phys}}$.

Dressed twists and visibility. Implement opposite twists on the two state branches,

$$U_1^{(\text{state})}(\phi) = e^{+\mathrm{i}\phi\, Q_a^{\text{dress}}}, \qquad U_2^{(\text{state})}(\phi) = e^{-\mathrm{i}\phi\, Q_a^{\text{dress}}}, \tag{7.382}$$

and define the dressed state–visibility

$$\Lambda_{\text{state}}^{\text{dress}}(\phi; \lambda_{\text{UV}}, \ell) = \frac{\left| \operatorname{Tr}_{\mathcal{H}_{\text{phys}}} \left(\mathsf{B}_{\lambda,\ell}\, e^{\,2\mathrm{i}\phi\, Q_a^{\text{dress}}}\, \rho_{\text{br}} \right) \right|}{\operatorname{Tr}_{\mathcal{H}_{\text{phys}}} \left(\mathsf{B}_{\lambda,\ell}\, \rho_{\text{br}} \right)}, \qquad \Lambda_{\text{state}}^{\text{dress}}(\phi) = \Lambda_{\text{state}}^{\text{dress}}(-\phi). \tag{7.383}$$

Small–ϕ behaviour and boundary law. By the cumulant expansion in the BRST–closed band trace (cf. §4.4),

$$\ln \Lambda^{\mathrm{dress}}_{\mathrm{state}}(\phi) = -\,\phi^2\,\mathrm{Var}_{\mathrm{band}}\Big[Q_a^{\mathrm{dress}}\Big] + \mathcal{O}(\phi^4). \tag{7.384}$$

In a BEH phase the current–current sector is *short–ranged* (no NG pole), and Gauss constraints move weight from the bulk to $\partial\Omega$ through the dressing term. Consequently,

$$\mathrm{Var}_{\mathrm{band}}\Big[Q_a^{\mathrm{dress}}\Big] = \tilde{\chi}_a(\ell; F, \Sigma)\, S_{\mathrm{eff}} \;+\; o(S_{\mathrm{eff}}), \qquad S_{\mathrm{eff}} \propto L^{\,d-1}, \tag{7.385}$$

where S_{eff} is the effective $(d{-}1)$–dimensional measure of the dressing surface and $\tilde{\chi}_a$ a band susceptibility for the dressed operator. Thus,

$$\Lambda^{\mathrm{dress}}_{\mathrm{state}}(\phi; \lambda_{\mathrm{UV}}, \ell) \;\approx\; \exp\Big[-\,\phi^2\,\tilde{\chi}_a(\ell; F, \Sigma)\, S_{\mathrm{eff}}\Big] \quad (\phi \to 0), \tag{7.386}$$

showing a *boundary law* (area/perimeter in general d) rather than a bulk volume law.

Spectral picture and poles. In momentum space the physical transverse vector correlator (on shell, band–reduced; Corollary 7.343) has a massive pole,

$$\Big\langle A_i^T\, A_j^T \Big\rangle(\omega, \mathbf{q}) \sim \frac{\Pi_{ij}^T(\mathbf{q})}{\omega^2 - c^2\mathbf{q}^{\,2} - m_V^2 + \mathrm{i}0^+} \quad \Rightarrow \quad \widetilde{\big\langle J^0 J^0 \big\rangle}_{c,\mathrm{band}}(\omega{=}0, \mathbf{q}) = \text{regular at } \mathbf{q}{=}0, \tag{7.387}$$

consistent with (7.385) once the Gauss/dressing contribution is included.

Acceptance (BEH run).

1. **Law guardrail:** verify $|\Lambda_{\mathrm{law}}| = 1 + \mathcal{O}(\varepsilon_{\mathrm{plat}})$ on the same plateau (eq. (7.358)).
2. **Dressing admissibility:** check $[\hat{Q}_{\mathrm{BRST}}, Q_a^{\mathrm{dress}}] = 0$ and $[\mathsf{B}_{\lambda,\ell}, Q_a^{\mathrm{dress}}] = 0$ (eq. (7.381)).
3. **Boundary scaling:** extract $\mathrm{Var}_{\mathrm{band}}[Q_a^{\mathrm{dress}}]$ from $\ln \Lambda^{\mathrm{dress}}_{\mathrm{state}}(\phi)$ and show S_{eff}-law scaling as in eq. (7.385) (e.g. $S_{\mathrm{eff}} \propto L^{d-1}$).
4. **ST/Nielsen stability:** demonstrate gauge–parameter independence of the extracted vector mass m_V and transverse residues within the band (Nielsen: eq. (7.339); ST closure: Section 7.5), and regulator robustness on the plateau (eq. (7.338)).

Scope. The construction targets *gauged* broken generators (BEH phases). For purely global symmetry breaking use the undressed twist of §4.4, which exhibits bulk volume scaling through the NG pole.

4.7 Plateau certification for visibilities

All readouts are asserted *on certified plateaus*. Concretely, with the band filter $\mathsf{B}_{\lambda,\ell} := P_x(\lambda_{\mathrm{UV}})K_{\mathrm{IR}}^{(\ell)}$ (eq. (7.354)) and Λ_α defined in eqs. (7.355) and (7.356), we require the following

window–flatness metrics (in the coordinates $u := \ln \lambda_{\mathrm{UV}}$, $v := \ln \ell$):

$$\mathcal{G}_\alpha := \sup_{(u,v)\in\mathcal{P}} \left\| \nabla_{(u,v)} \Lambda_\alpha(u,v) \right\|, \qquad \mathcal{O}_\alpha := \sup_{\gamma\subset\mathcal{P}} \left| \oint_\gamma \nabla \Lambda_\alpha \cdot d\boldsymbol{\ell} \right|, \qquad \mathcal{C}_\alpha := \sup_{(u,v)\in\mathcal{P}} \left\| \nabla^2_{(u,v)} \Lambda_\alpha \right\|, \tag{7.388}$$

to obey the tolerance bounds specified by eq. (7.137):

$$\mathcal{G}_\alpha \le \varepsilon_{\mathrm{grad}}, \qquad \mathcal{O}_\alpha \le \varepsilon_{\mathrm{osc}}, \qquad \mathcal{C}_\alpha \le \varepsilon_{\mathrm{curv}} \quad \text{for each } \alpha \in \{\mathrm{law}, \mathrm{state}\}. \tag{7.389}$$

Integrated readouts inherit these tolerances: for any admissible weight μ supported on the plateau (Section 7.4.5 and eq. (7.357)) we demand

$$\left| \overline{\Lambda}_\alpha[\mu] - \Lambda_\alpha(u_*, v_*) \right| = \mathcal{O}\big(\varepsilon_{\mathrm{plat}}\big) \quad \text{for any reference point } (u_*, v_*) \in \mathrm{supp}(\mu). \tag{7.390}$$

Nielsen (gauge–parameter) protection. For *physical* outputs (pole locations, residues, and susceptibilities extracted from $\ln \Lambda_{\mathrm{state}}$), on-shell Nielsen identities imply vanishing gauge slopes within the plateau tolerances (cf. eqs. (7.260) and (7.339)):

$$\partial_\zeta \Lambda_{\mathrm{state}} = \mathcal{O}(\varepsilon_{\mathrm{plat}}), \qquad \partial_\zeta \chi_a = \mathcal{O}(\varepsilon_{\mathrm{plat}}), \tag{7.391}$$

and likewise for Λ_{law} (cf. eq. (7.260)). Reports must include the measured slopes and the window $\mathcal{P}$ used.

4.8 Practical protocol (run–sheet)

1. **Calibration (law guardrail).** With a symmetric ρ_0, verify $|\Lambda_{\mathrm{law}}| = 1 + \mathcal{O}(\varepsilon_{\mathrm{plat}})$ at the chosen plateau, and enforce the commutator/ordering bounds eqs. (7.265) and (7.359). Record $\partial_\zeta \Lambda_{\mathrm{law}}$ and regulator permutations.

2. **Broken readout (finite–size series).** Prepare the broken state ρ_{br} (see Section 7.7.2). At fixed $(\lambda_{\mathrm{UV}}, \ell) \in \mathcal{P}$, measure the state–channel visibility $\Lambda_{\mathrm{state}}(\lambda_{\mathrm{UV}}, \ell; V)$ along a size ladder V (with linear size L in d spatial dimensions). Compute the rate function I_V as in eq. (7.362) and test the convergence $I_V \to I_\star > 0$ per eq. (7.363).

3. **Phase–twist scan (small ϕ).** Implement $U_{1,2}^{(\mathrm{state})}(\phi)$ as in §4.4 (or the dressed version of §4.6 for gauged cases). Fit $\ln \Lambda_{\mathrm{state}}(\phi)$ near $\phi = 0$; extract $\chi_a \equiv -\partial_\phi^2 \ln \Lambda_{\mathrm{state}}\,|_{\phi=0} = \mathrm{Var}_{\mathrm{band}}[Q_a[F]]$ using eq. (7.369), and (optionally) its $\mathbf{q}$-dependence via eq. (7.373).

4. **Soft check (global vs. gauged).** For global breaking, verify NG consistency in the soft limit $\chi_a(\ell; \mathbf{q}) \sim F_a^2/\mathbf{q}^2$ (eq. (7.374)) and volume scaling (eq. (7.371)). For gauged breaking, use the dressed twist and confirm boundary–law scaling (eq. (7.385)) and massive–vector consistency (eq. (7.387)). Confirm Nielsen/Slavnov–Taylor stability (Section 7.5 and eq. (7.391)).

5. **Integration and robustness.** Compare differential and integrated readouts Λ_α vs. $\overline{\Lambda}_\alpha[\mu]$ (eq. (7.390)) across admissible (P_x, K_{IR}) choices (commuting with BRST/G); accept if deviations are $\mathcal{O}(\varepsilon_{\mathrm{plat}})$.

4.9 Acceptance tests (Quadrant II)

- **Law unity (guardrail).** $|1 - \Lambda_{\rm law}| \le C_{\rm plat}\varepsilon_{\rm plat} + C_{\rm reg}\varepsilon_{\rm reg}$ (eq. (7.359)); gradients/oscillations/curvatures within eq. (7.389).
- **State suppression (global).** At fixed plateau, $I_V = -(1/V)\ln|\Lambda_{\rm state}|$ (eq. (7.362)) converges to $I_\star > 0$ (eq. (7.363)); a fit to the families in eq. (7.364) (bulk or interface dominated) passes goodness checks with reported confidence intervals.
- **State suppression (gauged, dressed).** Define a boundary rate

$$J_L := -\frac{1}{L^{d-1}}\ln\left|\Lambda^{\rm dress}_{\rm state}(\phi_0)\right| \tag{7.392}$$

  for a fixed small ϕ_0; accept if J_L plateaus to $J_\star > 0$ as $L \to \infty$, consistent with eqs. (7.385) and (7.386):

$$\lim_{L\to\infty} J_L = J_\star(\phi_0, \ell) > 0. \tag{7.393}$$

- **Soft structure.** For global breaking: $\chi_a(\ell;\mathbf{q}) \sim F_a^2/\mathbf{q}^2$ as $|\mathbf{q}| \to 0$ (eq. (7.374)), consistent with the EFT dispersion. For gauged breaking: no $1/\mathbf{q}^2$ singularity in J^0J^0 (regular soft limit) and a massive transverse pole (eq. (7.387)).
- **Gauge independence (Nielsen).** Gauge–parameter derivatives of *physical* outputs satisfy $\partial_\zeta(\text{poles, residues}, \chi_a) = \mathcal{O}(\varepsilon_{\rm plat})$ (eqs. (7.260), (7.339) and (7.391)).

4.10 Failure modes and remedies

- $|\Lambda_{\rm law}| < 1$ *at plateau.* Violates the law guardrail. Remedy: recheck commutators and BRST closure (Section 7.7.1 and eqs. (7.265) and (7.359)); re–certify the plateau (eq. (7.389)) and the exact–identity sector (Section 7.5).
- *No V–decay of $|\Lambda_{\rm state}|$ (global).* Causes: off–plateau evaluation, wrong order of limits, residual explicit breaking. Remedy: enforce ordered limits (eq. (7.275)); audit the seed protocol and Ward checks (Sections 7.5 and 7.7.2); verify $\chi_a(\ell;\mathbf{q})$ exhibits the $1/\mathbf{q}^2$ soft behaviour (eq. (7.374)).
- *Gauge–parameter drift of χ_a or poles.* Indicates off–shell contamination or missing counterterms. Remedy: project to the physical band, re–impose ST/Nielsen (Section 7.5 and eq. (7.339)), and recalibrate the plateau (eq. (7.389)).
- *Inconsistent boundary scaling (gauged).* Likely an inadmissible twist. Remedy: replace the naive charge by a Wilson–dressed construction (eq. (7.380)) and verify $[\hat{Q}_{\rm BRST}, Q_a^{\rm dress}] = 0 = [\mathsf{B}_{\lambda,\ell}, Q_a^{\rm dress}]$ (eq. (7.381)); confirm massive–vector signatures (eq. (7.387)).

Outcome. Step 4 delivers quantitative coherence diagnostics that characterise Quadrant II: (i) unity of the law channel within certified tolerances; (ii) suppression of state visibility by finite–size (global) or boundary–law (gauged) scaling; and (iii) a phase–twist route to NG soft structure protected by Ward/Nielsen identities (Sections 7.4 and 7.5). Any emergent identifications ("Higgs–like", "graviton–like", "apeiron/chrono–kymon") remain *provisional* pending cross–quadrant checks and plateau–stability certification.

7.7.5 Plateau Certification and Gauge-parameter Independence

Aim. To make the interferometric diagnosis of Quadrant II *quantitative and regulator-stable*, we certify plateaus in the $(\ln \lambda_{\mathrm{UV}}, \ln \ell)$ plane for all relevant readouts—especially Λ_{law} and Λ_{state} from Section 7.7.4—and for the physical spectral data extracted in Section 7.7.3. Certification follows the geometric criteria and acceptance pipeline stated in Sections 7.3 and 7.4.5, and on-shell gauge-parameter independence is enforced via the Nielsen and Slavnov–Taylor identities (Section 7.5; [117, 442, 91, 82, 243]).

5.1 Geometric definition of plateaus (local tests)

Mesh and notation. Work on a rectangular mesh in $(u, v) \equiv (\ln \lambda_{\mathrm{UV}}, \ln \ell)$:

$$(u_i, v_j) \in \mathcal{G}, \quad u_{i\pm1} - u_i = \Delta u, \quad v_{j\pm1} - v_j = \Delta v, \tag{7.394}$$

and let $R_\alpha(u_i, v_j)$ denote any band readout, with $\alpha \in \{\text{law}, \text{state}, \text{pole indices}, \text{residue indices}\}$. All R_α are evaluated with the same band filter $\mathsf{B}_{\lambda,\ell} = P_x(\lambda_{\mathrm{UV}}) K_{\mathrm{IR}}^{(\ell)}$ and BRST–closed insertions.

Local gradients (finite differences). Use centred differences for interior points (one–sided at the boundary):

$$\partial_u R_\alpha(u_i, v_j) \approx \frac{R_\alpha(u_{i+1}, v_j) - R_\alpha(u_{i-1}, v_j)}{2\,\Delta u}. \tag{7.395}$$

$$\partial_v R_\alpha(u_i, v_j) \approx \frac{R_\alpha(u_i, v_{j+1}) - R_\alpha(u_i, v_{j-1})}{2\,\Delta v}. \tag{7.396}$$

For dimensionless control, the (optional) relative-gradient form $\tilde{\nabla} R_\alpha := (\partial_u R_\alpha, \partial_v R_\alpha) \big/ \max\{|R_\alpha|, R_{\min}\}$ can be used with a small floor $R_{\min}$.

Local curvature (Hessian spectral norm). Define the discrete Hessian at (u_i, v_j) by

$$\begin{aligned} \partial_{uu} R_\alpha &\approx \frac{R_{i+1,j} - 2R_{i,j} + R_{i-1,j}}{\Delta u^2}, \\ \partial_{vv} R_\alpha &\approx \frac{R_{i,j+1} - 2R_{i,j} + R_{i,j-1}}{\Delta v^2}, \\ \partial_{uv} R_\alpha &\approx \frac{R_{i+1,j+1} - R_{i+1,j-1} - R_{i-1,j+1} + R_{i-1,j-1}}{4\,\Delta u\,\Delta v}, \end{aligned} \tag{7.397}$$

and set $\mathcal{H}_\alpha(u_i, v_j) := \begin{pmatrix} \partial_{uu} & \partial_{uv} \\ \partial_{uv} & \partial_{vv} \end{pmatrix} R_\alpha$. The curvature proxy is the spectral (operator-2) norm:

$$\|\mathcal{H}_\alpha(u_i, v_j)\|_2 \;:=\; \max\{\text{eigenvalues of } (\mathcal{H}_\alpha^\top \mathcal{H}_\alpha)^{1/2}\}. \tag{7.398}$$

Local oscillation proxy (short-range variance). On a fixed stencil $\mathcal{S}_{i,j} \subset \mathcal{G}$ of radius r (e.g. Moore neighbourhood in the L_∞ metric),

$$\begin{aligned} \mathrm{osc}_\alpha(u_i, v_j) &:= \left(\frac{1}{|\mathcal{S}_{i,j}|} \sum_{(m,n)\in\mathcal{S}_{i,j}} \Big(R_\alpha(u_m, v_n) - \overline{R}_{\alpha,\mathcal{S}_{i,j}} \Big)^2 \right)^{1/2}, \\ \overline{R}_{\alpha,\mathcal{S}_{i,j}} &:= \frac{1}{|\mathcal{S}_{i,j}|} \sum_{(m,n)\in\mathcal{S}_{i,j}} R_\alpha(u_m, v_n). \end{aligned} \tag{7.399}$$

(Equivalently, a bounded second–difference energy can be used; both are compatible with eq. (7.137).)

Pointwise acceptance mask. Given the tolerances $(\varepsilon_{\mathrm{grad}}, \varepsilon_{\mathrm{curv}}, \varepsilon_{\mathrm{osc}})$ from eq. (7.137), define

$$\mathcal{M}_\alpha(u_i, v_j) := \mathbf{1}\Big(\big|\partial_u R_\alpha\big| \le \varepsilon_{\mathrm{grad}} \ \wedge\ \big|\partial_v R_\alpha\big| \le \varepsilon_{\mathrm{grad}} \ \wedge\ \|\mathcal{H}_\alpha\|_2 \le \varepsilon_{\mathrm{curv}} \ \wedge\ \mathrm{osc}_\alpha \le \varepsilon_{\mathrm{osc}} \Big)_{(u_i,v_j)}. \tag{7.400}$$

Connected components and morphological closing. Let $\mathcal{Z}_\alpha := \{(u_i, v_j) \in \mathcal{G} : \mathcal{M}_\alpha(u_i, v_j) = 1\}$. Apply a mild morphological closing (dilation by radius r_{dil} followed by erosion by the same radius) to fill sub-mesh pinholes without bridging separated flats; denote the result by $\mathcal{Z}_\alpha^{\mathrm{cl}}$. Define the *channel plateaus* as the connected components of $\mathcal{Z}_\alpha^{\mathrm{cl}}$ that exceed a minimal area $A_{\min}$ (in cells).

Core plateau (boundary margin). To avoid edge effects in downstream estimates, erode each accepted component by a margin r_{core} cells and keep the nonempty remainder:

$$\mathcal{P}_\alpha^\circ := \mathrm{Erode}_{r_{\mathrm{core}}}\Big(\mathrm{LargestComponents}\big(\mathcal{Z}_\alpha^{\mathrm{cl}}\big) \Big). \tag{7.401}$$

For multi–readout certification, intersect cores across channels:

$$\mathcal{P} := \bigcap_{\alpha\in\mathcal{A}} \mathcal{P}_\alpha^\circ, \qquad \mathcal{A} \subseteq \{\text{law}, \text{state}, \text{pole/residue indices}\}. \tag{7.402}$$

Anisotropy and scaling of axes. If ∂_u and ∂_v sensitivities differ strongly, use a scaled metric $\|\nabla R_\alpha\|_M := \|(S\,\nabla R_\alpha)\|_2$ with $S = \mathrm{diag}(s_u, s_v)$ chosen from prior scans or the dynamic range of each axis; adjust $\varepsilon_{\mathrm{grad}}$ accordingly. This keeps eq. (7.400) invariant under benign reparameterisations of (u, v).

Consistency with global plateau metrics. On any accepted $\mathcal{P}$, the global gradient/oscillation/curvature metrics of eq. (7.388) satisfy the bounds of eq. (7.389) by construction; integrated readouts then obey eq. (7.390). Reports should list: grid spacings $(\Delta u, \Delta v)$, stencil radius r, morphology radii $(r_{\mathrm{dil}}, r_{\mathrm{core}})$, area thresholds $A_{\min}$, and the final $\mathcal{P}$.

5.2 Joint plateau and channel hierarchy

Definition (joint plateau from core components). Let $\mathcal{P}_\alpha^\circ$ be the *core plateau* of channel α obtained in §5.1 after morphological closing and boundary erosion (eq. (7.401)). For a chosen index set $\mathcal{I} \subseteq \{\text{law}, \text{state}, \text{poles}, \text{residues}\}$, define the *joint plateau* as the core–intersection

$$\mathfrak{P} \;:=\; \bigcap_{\alpha\in\mathcal{I}} \mathcal{P}_\alpha^\circ\,. \tag{7.403}$$

When $\mathfrak{P}$ is disconnected, denote by $\{\mathfrak{C}_k\}_{k=1}^{N}$ its connected components (on the (u,v) mesh of eq. (7.394)).

Channel hierarchy (operational order). We certify in the order

$$\text{law} \;\longrightarrow\; \text{state} \;\longrightarrow\; (\text{poles}, \text{residues}),$$

so that $\mathcal{I}$ is grown stepwise: (i) certify $\mathcal{P}_{\text{law}}^\circ$; (ii) intersect with $\mathcal{P}_{\text{state}}^\circ$; (iii) intersect with the spectral cores needed for the physics claim (e.g. NG stiffness proxies or BEH masses from §3). If any step yields an empty intersection, apply the fallback below before proceeding.

Aggregate plateau bounds (intersection consistency). On any joint component $\mathfrak{C} \subseteq \mathfrak{P}$, define the channel–wise maxima of the local metrics from eq. (7.400):

$$\mathcal{G}_{\alpha,\mathfrak{C}} := \sup_{\mathfrak{C}} \|\nabla R_\alpha\|, \quad \mathcal{C}_{\alpha,\mathfrak{C}} := \sup_{\mathfrak{C}} \|\mathcal{H}_\alpha\|_2, \quad \mathcal{O}_{\alpha,\mathfrak{C}} := \sup_{\mathfrak{C}} \operatorname{osc}_\alpha\,. \tag{7.404}$$

The joint component is admissible iff, for all $\alpha \in \mathcal{I}$,

$$\mathcal{G}_{\alpha,\mathfrak{C}} \le \varepsilon_{\text{grad}}, \qquad \mathcal{C}_{\alpha,\mathfrak{C}} \le \varepsilon_{\text{curv}}, \qquad \mathcal{O}_{\alpha,\mathfrak{C}} \le \varepsilon_{\text{osc}}, \tag{7.405}$$

and the integrated/differential agreement eq. (7.390) holds on $\mathfrak{C}$.

Component scoring (area and condition). For each admissible $\mathfrak{C}_k$, define an effective area

$$A_{\text{eff}}(\mathfrak{C}_k) := |\mathfrak{C}_k|\,\Delta u\,\Delta v, \tag{7.406}$$

and a simple "condition number" that aggregates the normalised tolerances across $\mathcal{I}$:

$$\kappa(\mathfrak{C}_k) := \max_{\alpha\in\mathcal{I}} \left\{ \frac{\mathcal{G}_{\alpha,\mathfrak{C}_k}}{\varepsilon_{\text{grad}}}, \frac{\mathcal{C}_{\alpha,\mathfrak{C}_k}}{\varepsilon_{\text{curv}}}, \frac{\mathcal{O}_{\alpha,\mathfrak{C}_k}}{\varepsilon_{\text{osc}}} \right\}. \tag{7.407}$$

Select the primary joint plateau by the lexicographic rule

$$\mathfrak{C}_\star := \arg\max_{\mathfrak{C}_k} \Big(A_{\text{eff}}(\mathfrak{C}_k),\; -\kappa(\mathfrak{C}_k) \Big), \tag{7.408}$$

i.e. largest effective area, then smallest κ as tie–breaker. Set $\mathfrak{P}_\star := \mathfrak{C}_\star$ for downstream physics.

Fallback when $\mathfrak{P} = \varnothing$. If the intersection is empty at a given hierarchy step:

1. **Re–mesh:** halve $(\Delta u, \Delta v)$ locally around the best overlapping fringes (identified by distance transforms between cores).
2. **Re–close:** apply a minimal increase of the closing radius $r_{\mathrm{dil}} \mapsto r_{\mathrm{dil}}{+}1$ and re–erode by the same margin to avoid spurious bridges.
3. **Tolerance micro–relaxation:** increase $\varepsilon_{\mathrm{grad}}, \varepsilon_{\mathrm{osc}}, \varepsilon_{\mathrm{curv}}$ by a common factor $1 + \delta$ with $\delta \le 0.1$, *only* for the spectral channel(s); keep the law/state tolerances fixed to preserve the guardrail.

If $\mathfrak{P}$ remains empty, report *no certified joint plateau* and revert to §4 diagnostics.

Reporting. For transparency, record: the set $\mathcal{I}$, the selected $\mathfrak{P}_\star$ (bounding box in (u, v)), $(\Delta u, \Delta v)$, morphology radii $(r_{\mathrm{dil}}, r_{\mathrm{core}})$, thresholds $(\varepsilon_{\mathrm{grad}}, \varepsilon_{\mathrm{osc}}, \varepsilon_{\mathrm{curv}})$, and the scores $\{A_{\mathrm{eff}}, \kappa\}$ for all admissible components. This enables reproducibility and direct application of eqs. (7.389) and (7.390).

5.3 Integrated readouts and inheritance

Admissible weights and supports. Let $\mathcal{P}$ be a certified plateau (or its core $\mathcal{P}^\circ$, cf. eq. (7.401)). A *band–admissible* weight is a non–negative function supported inside $\mathcal{P}$ and normalised to one:

$$w \ge 0, \qquad \int w = 1, \qquad \operatorname{supp} w \subset \mathcal{P} \text{ (or } \mathcal{P}^\circ). \tag{7.409}$$

We allow w to depend on the orthogonal coordinate (e.g. $w(u; v_*)$ at fixed v_*), or to be a separable 2D weight $W(u, v) = w_u(u) w_v(v)$ with each factor admissible.

Integrated (marginal) readouts: continuous forms. For any band readout $R_\alpha(u, v)$ (law, state, poles/residues), define 1D and 2D integrals

$$\overline{R}_\alpha(v_*) := \int_{u_1}^{u_2} w(u; v_*)\, R_\alpha(u, v_*)\, du, \qquad \int_{u_1}^{u_2} w(\cdot; v_*)\, du = 1, \tag{7.410}$$

$$\overline{R}_\alpha(u_*) := \int_{v_1}^{v_2} w(v; u_*)\, R_\alpha(u_*, v)\, dv, \qquad \int_{v_1}^{v_2} w(\cdot; u_*)\, dv = 1, \tag{7.411}$$

$$\overline{\overline{R}}_\alpha := \iint_{(u,v)\in\mathcal{R}} W(u, v)\, R_\alpha(u, v)\, du\, dv, \qquad \iint_{\mathcal{R}} W = 1, \tag{7.412}$$

where $\mathcal{R} \subset \mathcal{P}$ (typically a small rectangle inside a connected component).

Discrete mesh versions. On the grid $\mathcal{G}$ of eq. (7.394), the natural Riemann–sum analogues are

$$\overline{R}_\alpha(v_j) = \sum_{i=i_1}^{i_2} \omega_i^{(j)}\, R_\alpha(u_i, v_j), \qquad \sum_{i=i_1}^{i_2} \omega_i^{(j)} = 1, \quad \omega_i^{(j)} \ge 0, \tag{7.413}$$

(and similarly along v), or 2D

$$\overline{\overline{R}}_\alpha = \sum_{(i,j)\in\mathcal{R}} \Omega_{ij}\, R_\alpha(u_i, v_j), \qquad \sum_{(i,j)\in\mathcal{R}} \Omega_{ij} = 1, \quad \Omega_{ij} \geq 0, \tag{7.414}$$

with $\operatorname{supp}\omega, \Omega \subset \mathcal{P}$ (or $\mathcal{P}^\circ$).

Lemma 7.345 (Inheritance of plateau bounds)**.** *Assume R_α obeys the local plateau bounds of eq. (7.137) on a neighbourhood of the integration support. Then the integrated readouts $\overline{R}_\alpha$ and $\overline{\overline{R}}_\alpha$ satisfy the* same *type of bounds with equal or better constants:*

$$\left|\partial_v \overline{R}_\alpha(v_*)\right| \leq \int w(u; v_*) \left|\partial_v R_\alpha(u, v_*)\right| du \leq \varepsilon_{\text{grad}}, \tag{7.415}$$

and, analogously, $|\partial_u \overline{R}_\alpha(u_*)| \leq \varepsilon_{\text{grad}}$. For curvature,

$$\left\|\nabla^2 \overline{R}_\alpha(v_*)\right\| \leq \int w(u; v_*) \left\|\nabla^2 R_\alpha(u, v_*)\right\| du \leq \varepsilon_{\text{curv}}, \tag{7.416}$$

and for the oscillation proxy on any stencil contained in $\mathcal{P}$,

$$\operatorname{osc}\left[\overline{R}_\alpha\right] \;\leq\; \int w \operatorname{osc}[R_\alpha] \;\leq\; \varepsilon_{\text{osc}}. \tag{7.417}$$

The discrete versions hold with integrals replaced by convex combinations (7.413)–(7.414).

Proof sketch. Linearity plus convexity of the norm and Jensen's inequality yield $\partial_v \overline{R}_\alpha = \int w\, \partial_v R_\alpha$ and $|\partial_v \overline{R}_\alpha| \leq \int w\, |\partial_v R_\alpha|$. The Hessian estimate is identical using operator norms. For the oscillation proxy, $\operatorname{Var}\left[\int w\, X\right] \leq \int w \operatorname{Var}[X]$ on any bounded stencil; translate to the RMS definition of osc in eq. (7.399). Discrete forms follow from convexity of finite sums with non–negative normalised weights. □

Integrated vs. pointwise agreement (diagnostic). Given a reference point $(u_*, v_*) \in \operatorname{supp}(w) \subset \mathcal{P}$, define

$$\Delta_\alpha := \left|\overline{R}_\alpha - \; R_\alpha(u_*, v_*)\right|. \tag{7.418}$$

By Lemma 7.345 and the mean–value theorem applied inside $\mathcal{P}$, $\Delta_\alpha = \mathcal{O}(\varepsilon_{\text{plat}})$, reproducing eq. (7.390). Reporting should include Δ_α and the chosen weight shape (boxcar/Hann/Gaussian) and support.

Nielsen inheritance. For any physical output R_α protected by the on–shell Nielsen identity (eqs. (7.260) and (7.339)),

$$\partial_\zeta \overline{R}_\alpha = \int w\, \partial_\zeta R_\alpha = \mathcal{O}(\varepsilon_{\text{plat}}), \tag{7.419}$$

i.e. the integrated readout inherits gauge–parameter independence within the same tolerance.

Choice of weights. *Boxcar* weights maximise SNR and are convenient for rectangular $\mathcal{R}$; *Hann* or *Gaussian* (truncated) reduce boundary sensitivity. In all cases abide by (7.409); for narrow plateaus use the core $\mathcal{P}^\circ$ to avoid leakage across $\partial\mathcal{P}$ (cf. eq. (7.401)).

5.4 Regulator robustness and commutation of limits

Admissible families and notation. Let $\mathcal{R}_{\mathrm{adm}} = \{(P^{(i)}, K^{(i)})\}_{i=1}^{N}$ be an admissible family of band operators: each pair is BRST–closed and G–compatible on the certified window,

$$[\hat{Q}_{\mathrm{BRST}}, P^{(i)}] = [\hat{Q}_{\mathrm{BRST}}, K^{(i)}] = [\mathsf{J}_\alpha, P^{(i)}] = [\mathsf{J}_\alpha, K^{(i)}] = 0, \qquad i = 1, \dots, N,$$

and (on plateau) either commute or have commutators bounded within the regularisation stability budget (cf. Section 7.3 and eq. (7.67)). For any readout R_α we write $R_\alpha^{(i)}$ for the value obtained with $(P^{(i)}, K^{(i)})$.

Regulator robustness (uniform envelope over the joint plateau). On the joint plateau $\mathfrak{P}$ (eq. (7.403)), define the regulator envelope

$$\Delta_{\mathrm{reg}}[R_\alpha] := \max_{i,j} \sup_{(u,v)\in\mathfrak{P}} \left| R_\alpha^{(i)}(u,v) - R_\alpha^{(j)}(u,v) \right|. \tag{7.420}$$

Acceptance requires

$$\Delta_{\mathrm{reg}}[R_\alpha] \;\le\; \varepsilon_{\mathrm{reg}}, \tag{7.421}$$

with $\varepsilon_{\mathrm{reg}}$ bounded as in eq. (7.67). The same bound applies to integrated readouts (eqs. (7.410) and (7.412)); by convexity, $\Delta_{\mathrm{reg}}[\overline{R}_\alpha] \le \Delta_{\mathrm{reg}}[R_\alpha]$.

Ordering invariance (placement and permutation test). For each $(P^{(i)}, K^{(i)})$ and any fixed physical operator $\mathcal{O}$, compare admissible placements (e.g. $P^{(i)}K^{(i)}$, $K^{(i)}P^{(i)}$, and versions with Π_{T} interleaved on certified bands). Define the ordering defect

$$\Delta_{\mathrm{ord}}[R_\alpha^{(i)}] := \sup_{(u,v)\in\mathfrak{P}} \max_{\text{admissible orders } o,o'} \left| R_\alpha^{(i);o}(u,v) - R_\alpha^{(i);o'}(u,v) \right|, \tag{7.422}$$

and require $\Delta_{\mathrm{ord}}[R_\alpha^{(i)}] \le C_{\mathrm{ord}}\, \varepsilon_{\mathrm{reg}}$ for a modest constant C_{ord} (typically $C_{\mathrm{ord}} \sim 1$). This is the operational counterpart of the projector–commutator bound eq. (7.265).

Kernel-level robustness (physical blocks). For spectral claims based on the reduced kernel $\mathbb{H}_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}$ (Section 7.7.3), impose

$$\sup_{(u,v)\in\mathfrak{P}} \left\| \mathbb{H}_{\mathrm{phys}}^{(i)}(p;u,v) - \mathbb{H}_{\mathrm{phys}}^{(j)}(p;u,v) \right\| \;\le\; \varepsilon_{\mathrm{reg}} \tag{7.423}$$

uniformly on the momentum region used to extract poles/residues. By eq. (7.339) and the quartet decoupling (eq. (7.336)), this bound propagates to poles and residues themselves.

Commutation of limits (band scans and preparations). On $\mathfrak{P}$ we test that relevant pairs of limits commute within a budget $\varepsilon_{\mathrm{comm}}$:

(i) UV/IR scan limits. For any reference $(u_*, v_*) \in \mathfrak{P}$,

$$\Delta_{\mathrm{comm}}^{(u,v)}[R_\alpha; u_*, v_*] := \left| \lim_{u\to u_*} \lim_{v\to v_*} R_\alpha(u,v) - \lim_{v\to v_*} \lim_{u\to u_*} R_\alpha(u,v) \right| \;\le\; \varepsilon_{\mathrm{comm}}. \tag{7.424}$$

A discrete proxy is the mixed–increment bound

$$\Big| R_\alpha(u{+}\Delta u, v{+}\Delta v) - R_\alpha(u{+}\Delta u, v) - R_\alpha(u, v{+}\Delta v) + R_\alpha(u, v) \Big| \;\le\; C\,\varepsilon_{\mathrm{comm}} \quad (\text{small } \Delta u, \Delta v). \tag{7.425}$$

(ii) Thermodynamic vs. seed limit (SSB preparation). With V the spatial volume and $\varepsilon \to 0^+$ the seed strength (Section 7.7.2),

$$\Delta^{(V,\varepsilon)}_{\mathrm{comm}}[R_\alpha] := \Big| \lim_{V\to\infty} \lim_{\varepsilon\to 0^+} R_\alpha - \lim_{\varepsilon\to 0^+} \lim_{V\to\infty} R_\alpha \Big| \;\le\; \varepsilon_{\mathrm{comm}}. \tag{7.426}$$

(iii) Ground–state projection vs. seed (imaginary time). For $\beta \to \infty$ in the imaginary–time route,

$$\Delta^{(\beta,\varepsilon)}_{\mathrm{comm}}[R_\alpha] := \Big| \lim_{\beta\to\infty} \lim_{\varepsilon\to 0^+} R_\alpha - \lim_{\varepsilon\to 0^+} \lim_{\beta\to\infty} R_\alpha \Big| \;\le\; \varepsilon_{\mathrm{comm}}. \tag{7.427}$$

Violations of (7.424)–(7.427) trigger window tightening or exclusion of the affected region from $\mathfrak{P}$.

Algorithmic notes (how to check in practice).

1. *Envelope:* evaluate $R^{(i)}_\alpha$ on the mesh $\mathcal{G}$ (eq. (7.394)) for all $(P^{(i)}, K^{(i)})$; compute $\Delta_{\mathrm{reg}}[R_\alpha]$ via eq. (7.420).
2. *Ordering:* for each i, build $R^{(i);o}_\alpha$ across admissible operator orders; compute $\Delta_{\mathrm{ord}}[R^{(i)}_\alpha]$ via eq. (7.422).
3. *Limits:* along monotone sequences $u_n \to u_*$, $v_m \to v_*$ inside $\mathfrak{P}$, estimate $\Delta^{(u,v)}_{\mathrm{comm}}$ by Richardson extrapolation or the mixed–increment proxy eq. (7.425). Repeat for (V,ε) and (β,ε) using the same preparation protocol as in Section 7.7.2.
4. *Kernel:* compare $\mathbb{H}^{(i)}_{\mathrm{phys}}$ across regulators as in eq. (7.423); propagate to poles/residues.

Acceptance and reporting. Report the triplet $\big(\Delta_{\mathrm{reg}}[R_\alpha],\, \max_i \Delta_{\mathrm{ord}}[R^{(i)}_\alpha],\, \Delta^{(\cdot)}_{\mathrm{comm}}[R_\alpha]\big)$ together with the family $\mathcal{R}_{\mathrm{adm}}$, the joint plateau $\mathfrak{P}$ (bounding box), and the budgets $(\varepsilon_{\mathrm{reg}}, \varepsilon_{\mathrm{comm}})$. Acceptance requires (7.421), $\Delta_{\mathrm{ord}} \le C_{\mathrm{ord}}\varepsilon_{\mathrm{reg}}$, and (7.424)–(7.427). By eq. (7.338) and Nielsen/ST protection (Section 7.5 and eq. (7.339)), accepted bounds are stable for physical outputs.

5.5 Gauge-parameter independence (Nielsen tests)

Set-up (multi-parameter gauges). Let $\boldsymbol{\xi} = (\xi_1, \dots, \xi_M)$ denote a set of admissible gauge parameters (e.g. covariant gauge ζ, background-field gauge weights, etc.), varied within a domain $\mathcal{X}_{\mathrm{adm}}$ where BRST/ST identities hold and the band projectors remain BRST-closed (cf. Sections 7.3 and 7.5). On the certified joint plateau $\mathfrak{P}$ (eq. (7.403)), any *physical* output O_{phys} (pole mass, residue, stiffness proxy extracted from Λ_{state}, etc.) must satisfy the on-shell Nielsen condition

$$\sup_{(u,v)\in\mathfrak{P}} \sup_{\|\mathbf{e}\|_2=1} \Big| \partial_{\mathbf{e}} O_{\mathrm{phys}}(u, v; \boldsymbol{\xi}) \Big| \;\le\; \varepsilon_{\mathrm{gauge}}, \qquad \partial_{\mathbf{e}} := \sum_{r=1}^{M} e_r\, \partial_{\xi_r}, \tag{7.428}$$

with $\varepsilon_{\text{gauge}}$ chosen no larger than the plateau tolerances in eq. (7.137). Equation eq. (7.428) is the observable-level consequence of the Hessian identity $\partial_{\xi_r}\mathbb{H}_{\text{phys}} = \{\hat{Q}_{\text{BRST}}, \mathcal{N}_{\xi_r}\}$ on the band (eq. (7.339)) combined with $\mathrm{Tr}_{\mathcal{H}_{\text{phys}}}(\mathcal{O}\{\hat{Q}, X\}) = 0$.

Local slope maps (differential test). Define, for each parameter direction ξ_r,

$$\mathcal{S}_r(u,v) \;:=\; \Big|\partial_{\xi_r} O_{\text{phys}}(u,v;\boldsymbol{\xi}_*)\Big|, \tag{7.429}$$

evaluated at a reference $\boldsymbol{\xi}_* \in \mathcal{X}_{\text{adm}}$. Acceptance requires $\sup_{\mathfrak{P}} \mathcal{S}_r \le \varepsilon_{\text{gauge}}$ for all r. Discrete estimates use centred differences on a small gauge stencil:

$$\partial_{\xi_r} O_{\text{phys}}(u,v;\boldsymbol{\xi}_*) \;\approx\; \frac{O_{\text{phys}}(u,v;\boldsymbol{\xi}_* + \delta\xi_r\,\mathbf{e}_r) - O_{\text{phys}}(u,v;\boldsymbol{\xi}_* - \delta\xi_r\,\mathbf{e}_r)}{2\,\delta\xi_r}, \tag{7.430}$$

with $\delta\xi_r$ chosen so that $\boldsymbol{\xi}_* \pm \delta\xi_r\,\mathbf{e}_r \in \mathcal{X}_{\text{adm}}$.

Finite-set scan (uniform envelope). Given a finite admissible set $\{\boldsymbol{\xi}^{(k)}\}_{k=1}^{K} \subset \mathcal{X}_{\text{adm}}$, form the uniform gauge envelope

$$\Delta_{\text{gauge}}[O_{\text{phys}}] := \max_{k,k'} \sup_{(u,v)\in\mathfrak{P}} \Big| O_{\text{phys}}(u,v;\boldsymbol{\xi}^{(k)}) - O_{\text{phys}}(u,v;\boldsymbol{\xi}^{(k')}) \Big| \;\le\; \varepsilon_{\text{gauge}}. \tag{7.431}$$

By eq. (7.428), Δ_{gauge} must be of order $\varepsilon_{\text{gauge}}$; larger values signal off-shell contamination or an unstable plateau.

Commutation with band scans (mixed derivatives). Gauge slopes must commute (within tolerance) with UV/IR scans on $\mathfrak{P}$:

$$\Big|\partial_u\partial_{\xi_r} O_{\text{phys}} - \partial_{\xi_r}\partial_u O_{\text{phys}}\Big| \le \varepsilon_{\text{comm}}, \qquad \Big|\partial_v\partial_{\xi_r} O_{\text{phys}} - \partial_{\xi_r}\partial_v O_{\text{phys}}\Big| \le \varepsilon_{\text{comm}}, \tag{7.432}$$

tested via mixed increments (cf. eq. (7.425)). Violations point to window tightness issues or regulator misplacement.

ST closure residual (auxiliary guardrail). Let $\mathcal{R}_{\text{ST}}(u,v;\boldsymbol{\xi})$ denote a scalar residual of the Slavnov–Taylor identities evaluated on the reduced kernel (e.g. a norm of the linearised ST functional at two-point level). Require

$$\sup_{(u,v)\in\mathfrak{P}} \mathcal{R}_{\text{ST}}(u,v;\boldsymbol{\xi}) \;\le\; \varepsilon_{\text{ST}}, \tag{7.433}$$

with $\varepsilon_{\text{ST}} \lesssim \varepsilon_{\text{gauge}}$. Nonzero ST residuals often coincide with apparent Nielsen drift.

Integrated outputs inherit Nielsen stability. For any admissible weight w supported in $\mathfrak{P}$ (cf. eq. (7.409)), integrated observables obey

$$\partial_{\xi_r}\overline{O}_{\text{phys}} = \int w\,\partial_{\xi_r} O_{\text{phys}} = \mathcal{O}(\varepsilon_{\text{gauge}}), \tag{7.434}$$

matching the inheritance principle of eq. (7.419).

Algorithm (run-sheet).

1. Choose $\boldsymbol{\xi}_*$ and a gauge stencil $\{\boldsymbol{\xi}^{(k)}\} \subset \mathcal{X}_{\mathrm{adm}}$.
2. On $\mathfrak{P}$, build slope maps $\mathcal{S}_r$ via eq. (7.430) and enforce $\sup_{\mathfrak{P}} \mathcal{S}_r \le \varepsilon_{\mathrm{gauge}}$.
3. Compute $\Delta_{\mathrm{gauge}}[O_{\mathrm{phys}}]$ via eq. (7.431).
4. Check mixed-derivative commutation eq. (7.432) and ST residual bound eq. (7.433).
5. Repeat for integrated outputs; compare with pointwise values (discrepancy within $\mathcal{O}(\varepsilon_{\mathrm{gauge}})$).

Failure modes and remedies. A nonzero Nielsen slope (or large Δ_{gauge}) indicates: (i) off-shell evaluation, (ii) lack of a certified plateau, or (iii) missing counterterms/an ST mismatch. Remedy by re-enforcing the projection to $\mathcal{H}_{\mathrm{phys}}$, tightening $\mathfrak{P}$ (§5.1–5.2), and restoring ST/Nielsen closures (Section 7.5 and eq. (7.339)); if needed, adjust the admissible regulator family per §5.4.

5.6 Kato continuity and spectral tracking

Set–up (physical kernel and poles). On the joint plateau $\mathfrak{P}$ (eq. (7.403)), consider the band–reduced physical inverse two–point kernel

$$\mathbb{H}_{\mathrm{phys}}(p;u,v)\,, \qquad (u,v) = (\ln \lambda_{\mathrm{UV}}, \ln \ell), \tag{7.435}$$

obtained in Section 7.7.3 (cf. eq. (7.336)). For fixed $(u,v) \in \mathfrak{P}$, *simple* physical poles $\{p_i^2(u,v)\}$ are those for which there exists an eigenvalue branch $\lambda_i(p^2;u,v)$ of $\mathbb{H}_{\mathrm{phys}}$ such that

$$\lambda_i\Big(p_i^2(u,v);u,v\Big) = 0, \qquad \partial_{p^2}\lambda_i\Big(p_i^2(u,v);u,v\Big) \neq 0, \tag{7.436}$$

(the transversality condition; simple zero). The associated (matrix) residue is defined by the Riesz–projector representation (restriction to the physical subspace):

$$\begin{aligned} Z_i(u,v) \;&:=\; \lim_{p^2 \to p_i^2(u,v)} \Big(p^2 - p_i^2(u,v)\Big)\, \mathbb{G}_{\mathrm{phys}}(p;u,v) \\ &= P_i(u,v)\, \Big[\partial_{p^2}\mathbb{H}_{\mathrm{phys}}\Big(p_i^2(u,v);u,v\Big)\Big]^{-1} P_i(u,v), \end{aligned} \tag{7.437}$$

where $\mathbb{G}_{\mathrm{phys}} = \mathbb{H}_{\mathrm{phys}}^{-1}$ and

$$P_i(u,v) \;:=\; \frac{1}{2\pi i}\oint_{\gamma_i} \Big(\mathbb{H}_{\mathrm{phys}}(p;u,v)\Big)^{-1}\, dp^2 \tag{7.438}$$

is the Riesz projector around a small contour γ_i enclosing only the simple pole at p_i^2. (Nielsen/ST and quartet decoupling guarantee that only physical blocks contribute, cf. eqs. (7.336) and (7.339).)

Pathwise Kato continuity (implicit–function control). Assume $\mathbb{H}_{\text{phys}}(p;u,v)$ is C^1 in (u,v) on $\mathfrak{P}$ and analytic in p^2 near each simple pole (Kato family of type (A) in p^2 with C^1 parameters). Then by the implicit function theorem applied to λ_i at eq. (7.436), each $p_i^2(u,v)$ is C^1 on any simply–connected component of $\mathfrak{P}$ where the pole remains simple. In particular, there exists a Lipschitz constant L_i such that for all $(u,v),(u',v')$ in that component,

$$\frac{\left|p_i^2(u,v)-p_i^2(u',v')\right|}{\|(u,v)-(u',v')\|} \;\le\; L_i, \tag{7.439}$$

and similarly for any scalar summary of Z_i (e.g. Z_i contracted with physical polarisations). Plateau smoothness bounds imply $L_i=\mathcal{O}(\varepsilon_{\text{grad}})$.

Discrete Lipschitz tests (mesh implementation). On the grid $\mathcal{G}$ (eq. (7.394)), track branches $p_i^2(u_i,v_j)$ consistently between neighbours and enforce

$$\mathcal{L}_i \;:=\; \max_{\langle(i,j)\sim(i',j')\rangle} \frac{\left|p_i^2(u_i,v_j)-p_i^2(u_{i'},v_{j'})\right|}{\sqrt{(u_i-u_{i'})^2+(v_j-v_{j'})^2}} \;\le\; \varepsilon_{\text{Kato}}, \tag{7.440}$$

and likewise $\mathcal{L}_i[Z] := \max_{\text{nbrs}} \|Z_i(u_i,v_j)-Z_i(u_{i'},v_{j'})\|/\|\Delta(u,v)\| \le \varepsilon_{\text{Kato}}$. Here "$\langle\cdot\sim\cdot\rangle$" denotes nearest–neighbour pairs. Report both $\mathcal{L}_i$ and pathwise totals along geodesics inside $\mathfrak{P}$.

Gap and multiplet handling (crossings vs. symmetry). Define the spectral gap between simple poles

$$\gamma_i(u,v) \;:=\; \min_{j\neq i}\left|p_j^2(u,v)-p_i^2(u,v)\right|. \tag{7.441}$$

Simple–branch regime: if $\inf_{\mathfrak{P}}\gamma_i \ge \gamma_{\min}>0$, branches are uniquely trackable and eq. (7.440) suffices.

Multiplet regime: if symmetry enforces degeneracy or γ_i dips below a threshold $\gamma_\times$, promote $\{p_i^2\}$ to a cluster $\mathcal{C}$ with projector

$$P_{\mathcal{C}}(u,v) \;:=\; \frac{1}{2\pi i}\oint_{\Gamma}\Big(\mathbb{H}_{\text{phys}}(p;u,v)\Big)^{-1}\,dp^2, \tag{7.442}$$

where Γ encloses the whole cluster. Track the subspace via canonical angles; enforce Grassmann–distance Lipschitzness:

$$\frac{\left\|P_{\mathcal{C}}(u_i,v_j)-P_{\mathcal{C}}(u_{i'},v_{j'})\right\|}{\|\Delta(u,v)\|} \;\le\; \varepsilon_{\text{Kato}}\,, \tag{7.443}$$

and replace $\{Z_i\}$ by the cluster residue $Z_{\mathcal{C}} := P_{\mathcal{C}}\,[\partial_{p^2}\mathbb{H}_{\text{phys}}]^{-1}P_{\mathcal{C}}$ (well–defined on the range). Physics claims must then be phrased at the multiplet level.

Matching and branch continuation (practical tracking). Between neighbouring nodes, assign branches by minimising a compound cost

$$\mathsf{C}(i \to j) \;=\; \alpha \left| p_i^2(u,v) - p_j^2(u',v') \right| \;+\; \beta\, \Theta_{\mathrm{can}}\Big(\mathrm{Ran} P_i, \mathrm{Ran} P_j\Big), \qquad \alpha,\beta > 0, \tag{7.444}$$

where Θ_{can} is a sum of principal angles between the one–dimensional ranges (or between cluster subspaces). Solve the local assignment (Hungarian algorithm) to propagate labels; reject assignments that violate eq. (7.440) or $\gamma_i \geq \gamma_\times$.

Residue stability and normalisation. For simple poles, eq. (7.437) implies

$$\| Z_i(u_i,v_j) - Z_i(u_{i'},v_{j'}) \| \;\leq\; C_i \, \| \Delta(u,v) \| \qquad \text{on } \mathfrak{P}, \tag{7.445}$$

with $C_i = \mathcal{O}(\varepsilon_{\mathrm{grad}})$ controlled by the plateau derivatives of $\partial_{p^2}\mathbb{H}_{\mathrm{phys}}$. When reporting scalar residues (e.g. spectral weights), fix a physical normalisation (polarisation or channel) once and for all and keep it constant across $\mathfrak{P}$.

Acceptance tests (spectral level). For each tracked branch (or cluster) on $\mathfrak{P}$, require:

$$\mathcal{L}_i \leq \varepsilon_{\mathrm{Kato}}, \quad \inf_{\mathfrak{P}} \gamma_i \geq \gamma_{\min} \;\; \text{(or multiplet handling via eq. (7.443))}, \quad \mathcal{L}_i[Z] \leq \varepsilon_{\mathrm{Kato}}, \tag{7.446}$$

together with regulator robustness Δ_{reg} (§5.4) and Nielsen stability (§5.5) for the extracted p_i^2, Z_i. Crossing or bifurcation events ($\gamma_i < \gamma_\times$) inside $\mathfrak{P}$ must be either (i) explained by symmetry and treated as clusters, or (ii) excluded from certification (shrink $\mathfrak{P}$).

Algorithm (run-sheet).

1. **Local solve:** at each $(u_i, v_j) \in \mathcal{G} \cap \mathfrak{P}$, find simple poles by solving $\lambda_k(p^2; u_i, v_j) = 0$ near reference values; compute P_i, Z_i via eq. (7.438)–eq. (7.437).
2. **Neighbour matching:** propagate labels using the cost eq. (7.444); flag any $\gamma_i < \gamma_\times$ and switch to cluster tracking eq. (7.442).
3. **Lipschitz checks:** evaluate $\mathcal{L}_i$, $\mathcal{L}_i[Z]$ via eq. (7.440) and record min gaps γ_i on $\mathfrak{P}$.
4. **Stability:** verify regulator envelope (§5.4) and gauge independence (§5.5) for $\{p_i^2, Z_i\}$.
5. **Report:** list $\varepsilon_{\mathrm{Kato}}$, $\gamma_{\min}$, $\gamma_\times$, violation flags, and any regions removed from $\mathfrak{P}$.

Failure modes and remedies. Excessive $\mathcal{L}_i$ or $\mathcal{L}_i[Z]$ typically indicates: (i) non–simple poles ($\partial_{p^2}\lambda_i \approx 0$), (ii) unresolved crossings (small γ_i), or (iii) off–plateau drift of $\mathbb{H}_{\mathrm{phys}}$. Remedy by (a) promoting to a cluster projector and reporting multiplet observables, (b) tightening $\mathfrak{P}$, or (c) re–establishing plateau/Nielsen/ST bounds (Section 7.3 and eq. (7.339)).

5.7 Channel-aware acceptance and reporting

Scope. For each certified connected component of the joint plateau $\mathfrak{P}$ (cf. §5.2), assemble a *channel-aware* packet covering (i) law coherence, (ii) state suppression and phase–twist susceptibilities, (iii) spectral health (poles/residues), and (iv) robustness under admissible regulators and gauge choices.

Bounds summary (achieved vs. targets). Report the achieved thresholds

$$(\varepsilon_{\mathrm{grad}}, \varepsilon_{\mathrm{osc}}, \varepsilon_{\mathrm{curv}}, \varepsilon_{\mathrm{reg}}, \varepsilon_{\mathrm{comm}}, \varepsilon_{\mathrm{gauge}}, \varepsilon_{\mathrm{Kato}})$$

measured on $\mathfrak{P}$, alongside the targets from eqs. (7.67) and (7.137) (and the budgets set in §5.4–5.6). Indicate the *bounding box* of $\mathfrak{P}$ in (u, v) and the admissible families $\{(P^{(i)}, K^{(i)})\}$ and $\{\boldsymbol{\xi}^{(k)}\}$ used.

Law unity (statistics on $\mathfrak{P}$). Define the summary statistics for the law–channel deficit:

$$\mu_{\mathrm{law}} := \langle 1 - \Lambda_{\mathrm{law}} \rangle_{\mathfrak{P}}\,, \quad \mu_{\mathrm{law}}^{\max} := \sup_{(u,v)\in\mathfrak{P}} \big(1 - \Lambda_{\mathrm{law}}(u,v)\big), \quad \mathrm{CI}_{\mathrm{law}} : \text{ bootstrap or jackknife on } \mathfrak{P}. \tag{7.447}$$

Acceptance: $\mu_{\mathrm{law}}^{\max} \le \varepsilon_{\mathrm{plat}}$ and the CI centred at $\mu_{\mathrm{law}} = \mathcal{O}(\varepsilon_{\mathrm{plat}})$ (cf. eq. (7.259)); failures trigger rollback to Section 7.7.1.

State suppression (finite-size scaling at fixed plateau). At fixed $(u, v) \in \mathfrak{P}$, fit the finite-size dependence with an EFT-motivated family (stretched-exponential/exponential, optionally plus subleading corrections):

$$|\Lambda_{\mathrm{state}}(V)| \approx \exp\Big[- \kappa(\ell)\, V^{\nu} \Big] \times \begin{cases} 1, & \text{global SSB (NG)} \\ \exp[-\alpha_\partial\, |\partial\Omega|], & \text{gauged/BEH with boundary dressing} \end{cases} \tag{7.448}$$

with $\nu \ge 1$ in the global case and a boundary factor for the BEH case (cf. §4.6). Report best-fit parameters $\hat{\kappa}, \hat{\nu}$ (and $\hat{\alpha}_\partial$ when gauged) and goodness-of-fit:

$$\chi^2/\mathrm{dof} \;\le\; \chi^2_\star, \qquad \text{residual runs test within 5\%–95\% quantiles.} \tag{7.449}$$

Also quote the small-ϕ susceptibility $\chi_a = -\partial_\phi^2 \ln \Lambda_{\mathrm{state}}|_{\phi=0}$ and its scaling χ_a/V (global) or boundary scaling (gauged), cf. §§4.4–4.6.

Spectral health (mass/hyperbolicity and stability). Verify compliance with eqs. (7.13) and (7.15) and summarise via

$$m_{\min} \;:=\; \inf_{(u,v)\in\mathfrak{P}} \lambda_{\min}\Big(\mathsf{M}_{\mathrm{phys}}(u,v)\Big) \;\ge 0, \qquad \mathcal{H}_{\mathrm{hyp}} \;:=\; \inf_{(u,v)\in\mathfrak{P}} \inf_{\|n\|=1} \lambda_{\min}\Big(\mathrm{Re}\,\sigma_{\mathrm{prin}}(n)\Big) \;> 0, \tag{7.450}$$

together with Kato metrics from §5.6 (e.g. $\mathcal{L}[p_i^2]$, $\mathcal{L}[Z_i]$, $\gamma_{\min}$). If boundaries of $\mathfrak{P}$ are set by impending violations (e.g. $m_{\min} \downarrow 0$), annotate them.

Regulator and gauge robustness (overlays). Provide multi-(P, K) and multi-$\boldsymbol{\xi}$ overlays and their envelopes on $\mathfrak{P}$:

$$\Delta_{\mathrm{reg}}[\Lambda_\alpha] \;\le\; \varepsilon_{\mathrm{reg}}, \qquad \Delta_{\mathrm{gauge}}[O_{\mathrm{phys}}] \;\le\; \varepsilon_{\mathrm{gauge}}, \tag{7.451}$$

with definitions as in eqs. (7.420) and (7.431); include ordering-defect Δ_{ord} (cf. eq. (7.422)) and mixed-derivative commutation checks (eqs. (7.424), (7.425) and (7.432)).

Deliverables (per component of $\mathfrak{P}$). (i) Plateau mask and bounding box; (ii) heatmaps of local gradients/curvatures and slope maps in $\boldsymbol{\xi}$; (iii) law-unity statistics (7.447); (iv) V-scans with fits (7.309)–(7.449); (v) spectral dashboard; (vi) robustness overlays (7.451); (vii) raw seeds/configs sufficient for reproduction.

5.8 Failure modes and remedies

- *Fragmented plateaus (thin filaments, isolated pixels).* Symptoms: $\mathcal{M}_\alpha$ from §5.1 yields disjoint slivers; integrated readouts disagree with pointwise values beyond tolerance. Remedy: increase the stencil for the oscillation proxy; apply mild morphological closing; tighten admissible (P, K) to improve commutators; recheck commutation of limits (eqs. (7.424) and (7.425)) and shrink to the largest well-conditioned component.

- *Gauge drift persists (nonzero Nielsen slope).* Symptoms: $\mathcal{S}_r(u, v) = |\partial_{\xi_r} O_{\text{phys}}|$ exceeds $\varepsilon_{\text{gauge}}$ on $\mathfrak{P}$, or large Δ_{gauge} in eq. (7.431). Remedy: enforce ST/Nielsen ledger (Section 7.5 and eq. (7.339)), recompute the reduced kernel with the BRST-exact sector properly projected (cf. eq. (7.336)), and tighten $\mathfrak{P}$ (§5.1–5.2).

- *Regulator sensitivity (excess Δ_{reg}).* Symptoms: spreads across admissible (P, K) exceed ε_{reg}. Remedy: restrict to BRST-closer windows (where projector commutators obey eq. (7.265)); use the ordering-invariance test eq. (7.422); adjust UV/IR overlap to increase the certified commuting domain (Section 7.3).

- *Spectral crossings / bifurcations.* Symptoms: small spectral gaps $\gamma_i < \gamma_\times$, erratic branch labels, inflated Kato moduli. Remedy: switch to cluster tracking (Riesz projector $P_{\mathcal{C}}$, eq. (7.442)), enforce Grassmann Lipschitzness eq. (7.443), and phrase physics at the multiplet level; otherwise exclude the neighbourhood from $\mathfrak{P}$ and postpone identifications.

- *Finite-size anomalies (no V-decay in global case).* Symptoms: poor fits to (7.309), $\nu < 1$, or χ_a/V not stabilising. Remedy: re-check seed limits and ordering (cf. Section 7.7.2); verify Ward identities and the presence of an NG pole (§4.4–4.5); ensure measurements are taken strictly on $\mathfrak{P}$.

- *Numerical artefacts (underflow/phase noise).* Symptoms: $|\Lambda_\alpha|$ saturates numerical floor or exhibits phase jitter. Remedy: work with $-\ln|\Lambda_\alpha|$ and unwrap phases; increase sampling and use variance-reduced estimators; cross-validate with integrated readouts (§5.3).

Outcome. Step 5 furnishes a certified domain $\mathfrak{P}$ where all Quadrant II signatures are simultaneously stable: (i) $|\Lambda_{\text{law}}| \simeq 1$ within eq. (7.137); (ii) $|\Lambda_{\text{state}}|$ exhibits finite-size suppression at fixed plateau with accepted fits (7.309)–(7.449); (iii) physical poles/residues satisfy mass/hyperbolicity and remain Nielsen/ST stable. Any phenomenological labels ("Higgs-like", "graviton-like", "apeiron/chrono–kymon") remain *provisional* pending cross-quadrant checks, per Section 7.4.

7.7.6 Identity Ledger: Slavnov–Taylor, Ward, Nielsen, and Callan–Symanzik (On-band, Sectorwise)

Aim. To guarantee that all Quadrant II claims rest on symmetry-controlled, regulator-stable statements, we enforce the full *identity ledger* on the band: the Slavnov–Taylor (ST) master identity and its linearisation, global Ward identities (broken/unbroken), the Nielsen identity for gauge-parameter flows, and the Callan–Symanzik (CS) renormalisation-group equation. All checks are performed after the BRST projection to $\mathcal{H}_{\rm phys}$ and within certified plateaus of $(\ln \lambda_{\rm UV}, \ln \ell)$ (Sections 7.2, 7.3, 7.5, 7.7.3 and 7.7.4; [117, 442, 91, 82, 243, 114]).

6.1 Slavnov–Taylor master identity on the band

Set-up (Zinn–Justin sources; d-dimensional band). Let $\Gamma \equiv \Gamma[\Phi, K; \boldsymbol{\xi}; P_x, K^{(\ell)}_{\rm IR}]$ be the (band) 1PI effective action in d dimensions, with fields Φ^A and Zinn–Justin sources K_A coupled to the BRST variations $s\Phi^A$. All kernels/vertices are *implicitly composed* with the UV projector $P_x(\lambda_{\rm UV})$ and the IR window $K^{(\ell)}_{\rm IR}$ (cf. §5; the commutators eq. (7.244) ensure BRST compatibility on the band).

Master identity and linearisation. The Slavnov functional and the band ST master identity are

$$\begin{aligned} \mathcal{S}(\Gamma) &:= \int d^d x \sum_A \frac{\delta\Gamma}{\delta\Phi^A(x)} \frac{\delta\Gamma}{\delta K_A(x)} + (\text{ghost/gauge-fixing sector}), \\ \mathcal{S}(\Gamma) &= 0 \quad \text{on the band.} \end{aligned} \tag{7.452}$$

Linearising at Γ defines the nilpotent operator (the ST differential)

$$\mathcal{B}_\Gamma X \;:=\; \int d^d x \sum_A \left(\frac{\delta\Gamma}{\delta\Phi^A} \frac{\delta X}{\delta K_A} + \frac{\delta X}{\delta\Phi^A} \frac{\delta\Gamma}{\delta K_A} \right) \;+\; (\text{ghost/gauge-fixing terms})\,, \tag{7.453}$$

obeying

$$\mathcal{S}(\Gamma) = 0 \;\Longrightarrow\; \mathcal{B}_\Gamma^2 = 0\,. \tag{7.454}$$

Quadratic level (constraints on the Hessian). Let $\mathcal{H}_{\rm band}$ be the (band) Hessian at the background φ (eq. (7.311)). The linearised identity imposes

$$\mathcal{B}_\Gamma\, \mathcal{H}_{\rm band} \;=\; 0 \qquad (\text{as an operator identity on the band})\,, \tag{7.455}$$

which encodes, in particular: (i) transversality relations tying longitudinal gauge blocks to the ghost/Nakanishi–Lautrup sector; (ii) compatibility of mixing terms with BRST doublets; and (iii) the cohomological exactness used in §3 to derive eqs. (7.335) and (7.336). Together with the Nielsen identity (eq. (7.339)) this ensures that pole locations and residues extracted from $\mathbb{H}_{\rm phys}$ are gauge-parameter independent on shell.

Residuals and acceptance. Define band residuals (operator norms restricted to $\mathcal{H}_{\rm phys}$)

$$\mathcal{R}_{\rm ST}(u,v) \;:=\; \big\|\mathcal{S}(\Gamma)\big\|_{\rm phys}\,, \qquad \mathcal{R}_{\rm lin}(u,v) \;:=\; \big\|\mathcal{B}_\Gamma\,\mathcal{H}_{\rm band}\big\|_{\rm phys}\,, \tag{7.456}$$

evaluated on the joint plateau component $\mathfrak{P}$ (see §5.2). The acceptance criteria are

$$\sup_{(u,v)\in\mathfrak{P}} \mathcal{R}_{\rm ST}(u,v) \;\le\; \varepsilon_{\rm ST}\,, \qquad \sup_{(u,v)\in\mathfrak{P}} \mathcal{R}_{\rm lin}(u,v) \;\le\; \varepsilon_{\rm ST}\,, \tag{7.457}$$

with $\varepsilon_{\rm ST}$ chosen at or below the plateau budgets in eq. (7.137). Violations disqualify the candidate plateau or trigger a rebuild of gauge-fixing/renormalisation ([117, 243]).

Remarks. (i) All functional derivatives in eqs. (7.452) and (7.453) are taken at fixed band operators, which commute with the BRST charge by eq. (7.244). (ii) In BV notation, $\mathcal{B}_\Gamma$ is the linearisation of the antibracket $(\Gamma,\cdot)$; nilpotency eq. (7.454) is the Jacobi identity at $\mathcal{S}(\Gamma)=0$. (iii) The projections used in §3 imply that the ST constraints reduce to the physical blocks after quartet decoupling.

6.2 Global Ward identities (unbroken and broken)

Local Ward identities on the band (general d). Let R_α be unbroken generators and T_a broken generators of a global symmetry G, with conserved currents J^μ_α, J^μ_a. For any BRST-closed, gauge-invariant $\mathcal{O}$, the band-projected Ward–Takahashi identities read

$$\begin{aligned}
\partial_\mu \Big\langle J^\mu_\alpha(x)\,\mathcal{O}(y)\Big\rangle_{\rm band} &= i\,\Big\langle \delta_\alpha\mathcal{O}(y)\Big\rangle_{\rm band}\,\delta^{(d)}(x-y),\\
\partial_\mu \Big\langle J^\mu_a(x)\,\mathcal{O}(y)\Big\rangle_{\rm band} &= i\,\Big\langle \delta_a\mathcal{O}(y)\Big\rangle_{\rm band}\,\delta^{(d)}(x-y).
\end{aligned} \tag{7.458}$$

with all insertions composed with $P_x, K^{(\ell)}_{\rm IR}$ as in §4.4–§4.5. In momentum space, for amputated vertices this yields the standard constraints (e.g. current conservation and transversality).

Unbroken sector (conservation and transversality). For unbroken R_α, eq. (7.458) implies that current insertions are conserved and for vector correlators the transverse projector saturates the physical contribution:

$$\begin{aligned}
&k_\mu\,\Gamma^{\mu\nu}_{\alpha\beta}(k;u,v)=0,\\
&\Gamma^{\rm phys}_{\alpha\beta}(k;u,v) \coloneqq \Pi^{\mu\nu}_{\rm T}(k)\,\Gamma_{\alpha\beta;\nu\rho}(k;u,v)\,\Pi^{\rho\sigma}_{\rm T}(k).
\end{aligned} \tag{7.459}$$

Here $\Gamma^{\rm phys}$ determines the physical pole structure on the band.

This forbids mass terms for unbroken global currents and, in a gauged realisation with unbroken subgroup, enforces massless transverse vectors (photon-like) in the physical spectrum.

Broken sector (NG vs. BEH). For broken T_a, eq. (7.458) implies (global case) a Nambu–Goldstone pole in the current–current correlator's spectral density; schematically (EFT regime, small k),

$$\Big\langle J^\mu_a J^\nu_b\Big\rangle(k) \;\sim\; \frac{\rho_{ab}\,k^\mu k^\nu}{k^2} \;+\; \cdots\,, \qquad k^2\to 0, \tag{7.460}$$

with stiffness matrix ρ_{ab} (cf. §4.5). In a gauged realisation (BEH), the would-be NG modes are reorganised into longitudinal polarisations of massive vectors; the physical TT correlator shows no massless pole and obeys ST-consistent relations with the gauge-fixing/ghost sector, as guaranteed by eqs. (7.452) and (7.455).

Band and plateau implementation. All Ward identities above are enforced and tested *on the certified plateau* (cf. §5): band projections commute with symmetry/BRST generators by eq. (7.244), and gauge-parameter independence of *physical* outputs follows from the Nielsen identity (eq. (7.339)). Acceptance requires:

$$\begin{aligned}&\text{(i) Unbroken:} \quad \sup_{(u,v)\in\mathfrak{P}} \left\| k_\mu \Gamma^{\mu\nu}_{\alpha\beta}(k;u,v) \right\| \;\le\; \varepsilon_{\mathrm{Ward}},\\ &\text{(ii) Broken global:}\quad \text{presence of an NG pole consistent with eq. (7.460) and §4.5„}\\ &\quad\text{with stiffness proxies from phase–twist susceptibilities matching within plateau}\\ &\quad\text{tolerances}\\ &\text{(iii) Gauged (BEH):}\quad \text{absence of massless poles in the TT current block; massive-}\\ &\quad\text{vector spectrum and residues ST/Nielsen-stable.}\end{aligned} \tag{7.461}$$

Failures indicate either off-shell evaluation, explicit breaking, or a breakdown of the gauge-fixing/renormalisation sector; remedy per §3–§5 and Section 7.5.

6.3 Nielsen identity (gauge-parameter flows)

Band Nielsen identity (general d). Let $\boldsymbol{\xi}$ denote an admissible set of gauge parameters and ξ any one of its components. On the band, the Nielsen identity states that the ξ-derivative of the 1PI effective action is BRST-exact with respect to the linearised ST operator $\mathcal{B}_\Gamma$ (cf. eqs. (7.453) and (7.454)):

$$\frac{\partial \Gamma}{\partial \xi} \;=\; \mathcal{B}_\Gamma \mathcal{N}_\xi \qquad \text{(all insertions composed with } P_x, K^{(\ell)}_{\mathrm{IR}}\text{)}\,, \tag{7.462}$$

for some band operator $\mathcal{N}_\xi$ of ghost number -1. Differentiating twice w.r.t. fields and evaluating at the broken background φ gives a band identity for the quadratic kernel:

$$\begin{aligned}\partial_\xi \mathcal{H}_{\mathrm{band}} &= \mathcal{B}_\Gamma \mathcal{N}^{(2)}_\xi,\\ \mathcal{N}^{(2)}_\xi &:= \left.\frac{\delta^2 \mathcal{N}_\xi}{\delta\Phi\,\delta\Phi}\right|_{\Phi=\varphi}.\end{aligned} \tag{7.463}$$

After BRST projection to the physical subspace and use of the doublet/quartet structure (cf. §3), this reduces to the on-shell statement already used in eq. (7.339):

$$\begin{aligned}&\partial_\xi \mathbb{H}^{(\lambda_{\mathrm{UV}},\ell)}_{\mathrm{phys}}(p) = \{\hat{Q}_{\mathrm{BRST}}, \mathcal{N}^{(\lambda_{\mathrm{UV}},\ell)}_\xi(p)\},\\ \Rightarrow\quad &\partial_\xi\,[\text{physical poles/residues}] = 0 \quad \text{on shell.}\end{aligned} \tag{7.464}$$

Consequences for readouts. For any BRST-closed, ghost-number 0 insertion $\mathcal{O}$, graded cyclicity on $\mathcal{H}_{\mathrm{phys}}$ (eq. (7.331)) yields:

$$\partial_\xi \Lambda_{\mathrm{law}}(\lambda_{\mathrm{UV}}, \ell) = 0 \quad \text{on the certified plateau,} \tag{7.465}$$

and for phase–twist susceptibilities ($\chi_a = -\partial_\phi^2 \ln \Lambda_{\mathrm{state}}|_{\phi=0}$)

$$\partial_\xi \chi_a = \partial_\xi \operatorname{Var}_{\mathrm{band}}\big[Q_a[F]\big] = \operatorname{Tr}_{\mathcal{H}_{\mathrm{phys}}}\Big(\mathcal{O}_a \{\hat{Q}_{\mathrm{BRST}}, \mathcal{N}_\xi\}\Big) = 0 \quad \text{(on shell, on the plateau),} \tag{7.466}$$

with $\mathcal{O}_a$ the BRST-closed quadratic form defining the variance (§4.4).

Acceptance (numerical Nielsen tests). Define the bandwise Nielsen residual for any physical observable O_{phys}:

$$\mathcal{R}_{\mathrm{Niel}}[O_{\mathrm{phys}}](u, v; \xi) := \big|\partial_\xi O_{\mathrm{phys}}(u, v; \xi)\big| . \tag{7.467}$$

On the joint plateau $\mathfrak{P}$ we require

$$\sup_{(u,v)\in\mathfrak{P}} \mathcal{R}_{\mathrm{Niel}}[O_{\mathrm{phys}}](u, v; \xi) \;\le\; \varepsilon_{\mathrm{gauge}} \quad \text{for all tested } \xi,$$

with $\varepsilon_{\mathrm{gauge}}$ chosen no larger than the plateau budgets of eq. (7.137) (cf. §5.5). Persistent nonzero slopes indicate off-shell evaluation, a broken ST ledger, or an uncertified plateau; rebuild per §3–§5 and [117, 114, 243].

6.4 Callan–Symanzik equation (RG on the band)

CS flow for band-projected functionals. Let μ be the renormalisation scale for couplings $\{g^i\}$, with beta functions β^i and field anomalous dimensions γ_Φ. The Callan–Symanzik (CS) operator acting on the band 1PI functional in d dimensions reads

$$\Big(\mu\partial_\mu + \beta^i \partial_{g^i} + \gamma_\Phi \mathcal{N}_\Phi + \sum_r \gamma_{J_r} \mathcal{N}_{J_r}\Big)\, \Gamma \;=\; 0, \tag{7.468}$$

where $\mathcal{N}_\Phi$ counts fields and $\mathcal{N}_{J_r}$ counts renormalised sources/insertions. By construction all objects are composed with the same $P_x(\lambda_{\mathrm{UV}})$, $K_{\mathrm{IR}}^{(\ell)}$.

Interplay with regulator scans (plateau consistency). Small variations of the *regulator* scales $(\lambda_{\mathrm{UV}}, \ell)$ inside a certified plateau affect band-projected Green functions only within the tolerance budgets (eqs. (7.67) and (7.137)). RG-improved predictions must therefore satisfy:

$$\Big(\mu\partial_\mu + \beta^i \partial_{g^i} + \gamma \cdot \mathcal{N}\Big)\, O_{\mathrm{phys}}(u, v; \mu, \{g\}) \;=\; 0 \quad \text{on } \mathfrak{P}, \tag{7.469}$$

and be *stable* against admissible regulator changes:

$$\sup_{(u,v)\in\mathfrak{P}} \Big| \delta_{(P,K)} O_{\mathrm{phys}} - \Big(\mu\partial_\mu + \beta^i \partial_{g^i} + \gamma \cdot \mathcal{N}\Big) O_{\mathrm{phys}}\, \delta \ln \mu \Big| \;\le\; \varepsilon_{\mathrm{reg}} , \tag{7.470}$$

where $\delta_{(P,K)}$ denotes the observed spread across the admissible family $\{(P^{(i)}, K^{(i)})\}$ (cf. eq. (7.67)) and $\delta \ln \mu$ is the corresponding RG step.

Acceptance (RG robustness). For a chosen set of physical outputs O_{phys} (pole masses p_i^2, residues Z_i, phase–twist susceptibilities χ_a, etc.), enforce on $\mathfrak{P}$:

$$\sup_{(u,v)\in\mathfrak{P}} \left|\Big(\mu\partial_\mu + \beta^i\partial_{g^i} + \gamma\cdot\mathcal{N}\Big)O_{\mathrm{phys}}\right| \;\le\; \varepsilon_{\mathrm{RG}}\,, \qquad \Delta_{\mathrm{reg}}[O_{\mathrm{phys}}] \;\le\; \varepsilon_{\mathrm{reg}}, \tag{7.471}$$

with $\varepsilon_{\mathrm{RG}}$ comparable to (or tighter than) the plateau budgets, and with Kato/Lipschitz continuity in (u,v) as in §5.6 to exclude false "plateaus" driven by unresolved spectral rearrangements.

Remarks. (i) The CS flow commutes with BRST reduction: applying $\mathsf{P}_{\mathrm{phys}}$ to eq. (7.468) preserves the form of eq. (7.469) for physical blocks. (ii) Off the plateau, residual regulator dependence can fake RG flow; eq. (7.470) guards against that by comparing band spreads to the CS prediction. (iii) In practice, using RG-invariant combinations (e.g. ratios p_i/μ or stiffness proxies normalised by appropriate powers) sharpens the test in eq. (7.471).

6.5 Anomaly filters (algebraic renormalisation)

Wess–Zumino consistency on the band. Possible breakings Δ of the identity ledger (ST/Ward/Nielsen) are constrained by the linearised ST operator $\mathcal{B}_\Gamma$ (cf. eqs. (7.453) and (7.454)). On the band, with all objects composed with $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$, the Wess–Zumino (WZ) condition reads

$$\mathcal{B}_\Gamma\,\Delta \;=\; 0\,, \qquad \mathrm{gh}(\Delta)=1, \quad \deg(\Delta)=d\,, \tag{7.472}$$

so that Δ defines a cohomology class $[\Delta]\in H^1(\mathcal{B}_\Gamma)$. Local, integrated functionals are understood modulo total derivatives.

Trivial vs. nontrivial breakings. A breaking is *cohomologically trivial* if

$$\Delta \;=\; \mathcal{B}_\Gamma\,\widehat{\Delta} + \partial_\mu R^\mu \qquad\Rightarrow\qquad \text{removable by counterterms/field redefinitions on the band.} \tag{7.473}$$

Nontrivial representatives (e.g. gauge, mixed, or gravitational anomalies) live in $H^{1,d}(s\,|\,d)$ in the BRST bicomplex and are *forbidden* in an admissible plateau.

Band residuals and operational test. Define identity residuals on the joint plateau $\mathfrak{P}$:

$$\mathcal{R}_{\mathrm{WZ}}(u,v) \;:=\; \big\|\mathcal{B}_\Gamma\,\Delta\big\|_{\mathrm{phys}}, \qquad \mathcal{R}_{\mathrm{coh}}(u,v) \;:=\; \inf_{\widehat{\Delta},R}\big\|\Delta - \mathcal{B}_\Gamma\widehat{\Delta} - \partial_\mu R^\mu\big\|_{\mathrm{phys}}, \tag{7.474}$$

where norms are taken on $\mathcal{H}_{\mathrm{phys}}$ and the infimum ranges over local counterterm candidates $\widehat{\Delta}$ and currents R^μ. For concrete gauge groups/representations we also track the coefficients of known anomaly bases (descent of characteristic classes), collecting them into a vector $\mathbf{a}(u,v)$ (e.g. cubic/gauge, mixed, gravitational).

Acceptance (anomaly filters). On $\mathfrak{P}$ we require

$$\sup_{(u,v)\in\mathfrak{P}} \mathcal{R}_{\mathrm{WZ}} \le \varepsilon_{\mathrm{ST}}, \qquad \sup_{(u,v)\in\mathfrak{P}} \mathcal{R}_{\mathrm{coh}} \le \varepsilon_{\mathrm{anom}}, \qquad \sup_{(u,v)\in\mathfrak{P}} \|\mathbf{a}(u,v)\| \le \varepsilon_{\mathrm{anom}}, \tag{7.475}$$

with $\varepsilon_{\mathrm{anom}}$ chosen $\le$ the plateau budgets of eq. (7.137). Thus: (i) no nontrivial class in $H^1(\mathcal{B}_\Gamma)$ is present in the inspected sector; (ii) any residual breaking is $\mathcal{B}_\Gamma$-exact (up to a divergence) and numerically small; (iii) global-chiral, gauge, mixed, and gravitational anomalies are absent for the realised representation content (cf. [117, 243]).

Remarks. (1) Band commutators eq. (7.244) ensure that projecting with $P_x, K_{\mathrm{IR}}^{(\ell)}$ does not alter the WZ consistency analysis in cohomology. (2) In even d the anomaly scan is efficiently implemented via descent equations for the relevant characteristic classes and checked directly on $\mathfrak{P}$.

6.6 Sectorwise enforcement and interface with Steps 3–5

Checklist and cross-links. We enforce the identity ledger *per step* and feed its residuals into plateau certification:

1. **Quadratic kernel (Step 3).** Verify the linearised ST constraint on the band, $\mathcal{B}_\Gamma \mathcal{H}_{\mathrm{band}} = 0$ (eq. (7.455)), and after BRST projection use quartet decoupling to obtain the physical blocks (eq. (7.336)). Confirm hyperbolicity and mass positivity (eqs. (7.13) and (7.15)); failures trigger a rebuild of the gauge-fixing/renormalisation sector.

2. **Visibilities and soft structure (Step 4).** Check law-channel Nielsen stability $\partial_\xi \Lambda_{\mathrm{law}} = 0$ on the plateau (eq. (7.465)); extract phase–twist cumulants and verify Ward/Nielsen protection (eqs. (7.460) and (7.466)). In the gauged (BEH) case, confirm the absence of massless TT poles and ST-consistent massive-vector structure (§4.6, eq. (7.452)).

3. **Plateaus and robustness (Step 5).** Extend the acceptance mask $\mathcal{M}_\alpha$ (cf. §5.1) by *identity residual coordinates*:

$$\mathcal{M}_\alpha^+(u,v) = \mathcal{M}_\alpha(u,v) \wedge \left\{ \begin{array}{ll} \mathcal{R}_{\mathrm{ST}} \le \varepsilon_{\mathrm{ST}}, & \mathcal{R}_{\mathrm{lin}} \le \varepsilon_{\mathrm{ST}}, \\ \mathcal{R}_{\mathrm{Niel}} \le \varepsilon_{\mathrm{gauge}}, & \mathcal{R}_{\mathrm{WZ}} \le \varepsilon_{\mathrm{ST}}, \\ \mathcal{R}_{\mathrm{coh}} \le \varepsilon_{\mathrm{anom}} & \end{array} \right\}. \tag{7.476}$$

   and use this strengthened mask to define the joint plateau $\mathfrak{P}$ (§5.2). Regulator robustness and commutation of limits remain enforced via eq. (7.67) and §5.4.

Outcome. With eqs. (7.475) and (7.476) satisfied, the Step 3–5 artefacts (reduced physical kernel, visibilities, soft factors, and spectral data) are jointly consistent: ST/Ward/Nielsen identities hold on the band; anomalies are filtered out cohomologically; and all claims remain stable on the certified plateau $\mathfrak{P}$.

6.7 Practical run-sheet (auditable checks)

Sampling grid and on-shell locus. Fix a rectangular mesh in $(u, v) \equiv (\ln \lambda_{\mathrm{UV}}, \ln \ell)$ and a discrete set of admissible gauges $\{\xi_k\}$. At each mesh point, restrict to the on-shell locus as in Section 7.2 (vanishing band equations of motion and BRST closure), and use the same background φ and band operators $P_x, K_{\mathrm{IR}}^{(\ell)}$ across all tests.

1. **Source-extended action.** Build the Zinn–Justin sector and evaluate the ST ledger on the band: $\mathcal{S}(\Gamma)$ and $\mathcal{B}_\Gamma$ from eqs. (7.452) and (7.453), with the band commutators of eq. (7.244) enforced for the chosen (P_x, K_{IR}).

2. **ST residuals (on shell).** Compute the operator norms restricted to $\mathcal{H}_{\mathrm{phys}}$ $\mathcal{R}_{\mathrm{ST}}, \mathcal{R}_{\mathrm{lin}}$ defined in eq. (7.456) over the (u, v) mesh; persist them as audit layers and inject into the plateau mask per eq. (7.476).

3. **Ward suite (un/broken).** From $\mathcal{H}_{\mathrm{phys}}$ (§3) extract two-/three-point functions and test: (i) unbroken transversality eq. (7.459); (ii) broken-sector NG/BEH expectations eq. (7.460). For auditing, record the Ward residual

$$\mathcal{R}_{\mathrm{Ward}}(k; u, v) \;:=\; \Big\| k_\mu \, \Gamma^{\mu\nu}(k; u, v) \Big\|_{\mathrm{phys}} , \tag{7.477}$$

   and its sup over tested momenta in the IR window.

4. **Nielsen slopes (finite differences).** For any physical output O_{phys} (pole mass/residue, Λ_{law}, phase–twist susceptibility χ_a), estimate

$$\widehat{\partial_\xi O_{\mathrm{phys}}}(u, v; \xi_k) := \frac{O_{\mathrm{phys}}(u, v; \xi_{k+1}) - O_{\mathrm{phys}}(u, v; \xi_{k-1})}{\xi_{k+1} - \xi_{k-1}} , \tag{7.478}$$

   and store $\mathcal{R}_{\mathrm{Niel}} := |\widehat{\partial_\xi O_{\mathrm{phys}}}|$ for the acceptance bound in eq. (7.467) and §5.5.

5. **CS consistency (RG vs. regulator scans).** Track RG flows using eqs. (7.468) and (7.469) and compare with small changes of (P_x, K_{IR}) inside the admissible family via the discrepancy

$$\mathcal{D}_{\mathrm{CS}}[O_{\mathrm{phys}}] := \Big| \delta_{(P,K)} O_{\mathrm{phys}} - \big(\mu \partial_\mu + \beta^i \partial_{g^i} + \gamma \cdot \mathcal{N}\big) O_{\mathrm{phys}} \, \delta \ln \mu \Big| , \tag{7.479}$$

   to be bounded by eqs. (7.470) and (7.471).

6. **Cohomology filter (algebraic renormalisation).** If a breaking Δ is detected, evaluate the WZ residuals $\mathcal{R}_{\mathrm{WZ}}, \mathcal{R}_{\mathrm{coh}}$ in eq. (7.474) and the anomaly coefficient vector $\mathbf{a}$; accept only $\mathcal{B}_\Gamma$-exact (up to divergences) pieces below the tolerances in eq. (7.475), otherwise exclude the region or revise renormalisation conditions.

Auditable artefacts.

(i) Heatmaps over (u, v) for residuals $\mathcal{R}_{\mathrm{ST}}$, $\mathcal{R}_{\mathrm{lin}}$, $\mathcal{R}_{\mathrm{Niel}}$, $\mathcal{R}_{\mathrm{WZ}}$, $\mathcal{R}_{\mathrm{coh}}$, $\mathcal{R}_{\mathrm{Ward}}$, $\mathcal{D}_{\mathrm{CS}}$.

(ii) Kato–continuity trackers for poles/residues (§5.6).

(iii) Regulator overlays showing spreads vs. eq. (7.67).

6.8 Acceptance tests (ledger-complete)

Aggregated ledger. On the joint plateau $\mathfrak{P}$ (cf. §5.2) we require the following sup-norm bounds:

$$\sup_{(u,v)\in\mathfrak{P}} \left\{ \mathcal{R}_{\mathrm{ST}}(u,v),\ \mathcal{R}_{\mathrm{lin}}(u,v) \right\} \ \le\ \varepsilon_{\mathrm{plat}}, \tag{7.480}$$

$$\sup_{(u,v)\in\mathfrak{P}} \sup_{k\in \mathrm{IR\,win}} \mathcal{R}_{\mathrm{Ward}}(k;u,v) \le \varepsilon_{\mathrm{Ward}}$$

NG/BEH-consistent soft structure per eq. (7.460). (7.481)

$$\sup_{(u,v)\in\mathfrak{P}} \mathcal{R}_{\mathrm{Niel}}[O_{\mathrm{phys}}](u,v;\xi_k) \ \le\ \varepsilon_{\mathrm{gauge}} \quad \text{for all tested } O_{\mathrm{phys}} \text{ (cf. eq. (7.467))}, \tag{7.482}$$

$$\sup_{(u,v)\in\mathfrak{P}} \mathcal{D}_{\mathrm{CS}}[O_{\mathrm{phys}}](u,v) \ \le\ \varepsilon_{\mathrm{reg}} \quad \text{and} \quad \Delta_{\mathrm{reg}}[O_{\mathrm{phys}}] \ \le\ \varepsilon_{\mathrm{reg}} \ \text{(cf. eqs. (7.67) and (7.471))}, \tag{7.483}$$

$$\sup_{(u,v)\in\mathfrak{P}} \left\{ \mathcal{R}_{\mathrm{WZ}}(u,v),\ \mathcal{R}_{\mathrm{coh}}(u,v),\ \|\mathbf{a}(u,v)\| \right\} \le \varepsilon_{\mathrm{anom}}. \tag{7.484}$$

(no nontrivial $H^1(\mathcal{B}_\Gamma)$ representatives; cf. eq. (7.475))

Outcome. With eqs. (7.480) to (7.484) satisfied, the Quadrant II analysis is symmetry- and RG-controlled on certified plateaus: (i) the reduced physical kernel of Section 7.7.3 satisfies ST/Nielsen; (ii) the interferometric readouts of Section 7.7.4 obey Ward/Nielsen protection and exhibit the law/state signatures required; (iii) RG stability holds in the sense of Section 7.3. Any phenomenological mode labels ("Higgs-like", "graviton-like", "apeiron/chrono–kymon") remain *provisional* pending the cross-quadrant programme.

7.7.7 Physicality Tests: Nonnegative Spectrum, Hyperbolicity, and Spectral Positivity on $\mathcal{H}_{\mathrm{phys}}$

Aim. On the certified plateau $\mathfrak{P}$ (Section 7.7.5) and after BRST projection (Section 7.7.3), we must demonstrate *physicality* of dynamics and correlators: (i) the *physical mass operator* on $\mathcal{H}_{\mathrm{phys}}$ is nonnegative (cf. eq. (7.13)); (ii) the *principal symbol* of the physical kinetic operator is *hyperbolic* with causal cone matching the chronogeometric background (cf. eq. (7.15)); and (iii) band-projected two-point functions admit a Källén–Lehmann representation with *nonnegative* spectral densities and no negative-norm contributions (BRST quartets decoupled). All statements are enforced *on shell*, Nielsen/Slavnov–Taylor stable (Section 7.5), and regulator-robust (Section 7.3; [442, 117, 91, 82, 243]).

7.1 Physical mass operator on $\mathcal{H}_{\mathrm{phys}}$

Definition and extraction. From the reduced quadratic kernel $\mathcal{H}_{\mathrm{phys}}$ constructed in Section 7.7.3, define the physical mass operator $\mathbb{M}^2_{\mathrm{phys}}$ via the pole condition (cf. eq. (7.13)) for each physical block $X \in \{A^T, h, \pi\}$:

$$\det\left[\mathcal{H}^{XX}_{\mathrm{phys}}(k)\right]\Big|_{k^2=m_X^2} = 0, \qquad m_X^2 \ge 0, \qquad \mathbb{M}^2_{\mathrm{phys}}\,|X\rangle = m_X^2\,|X\rangle \ \text{ on } \mathcal{H}_{\mathrm{phys}}. \tag{7.485}$$

For an isolated simple pole, the residue matrix is

$$Z_X \;=\; \lim_{k^2\to m_X^2} (k^2 - m_X^2)\left[\mathcal{H}^{XX}_{\mathrm{phys}}(k)\right]^{-1}, \tag{7.486}$$

and determines one-particle normalisation and spectral weight (Källén–Lehmann).

Mixing and projectors. When fields mix inside X (e.g. (h,π)), let $v_{X,\alpha}(k)$ be right null-vectors of $\mathcal{H}^{XX}_{\mathrm{phys}}(k)$ at $k^2 = m^2_{X,\alpha}$. Define the spectral projector onto the pole subspace

$$\mathsf{P}_{X,\alpha} \;=\; Z_{X,\alpha}\, v_{X,\alpha}\, w^\dagger_{X,\alpha}, \qquad w^\dagger_{X,\alpha}\,\partial_{k^2}\mathcal{H}^{XX}_{\mathrm{phys}}\, v_{X,\alpha} = 1, \tag{7.487}$$

where $w_{X,\alpha}$ are left null-vectors and $Z_{X,\alpha}$ the scalar residues fixed by the normalisation condition. The block residue is $Z_X = \sum_\alpha \mathsf{P}_{X,\alpha}$.

Positivity tests (no ghosts). On $\mathcal{H}_{\mathrm{phys}}$ we require positivity of spectral weights:

$$\forall\, u \in \mathcal{H}_{\mathrm{phys}}: \quad u^\dagger Z_X\, u \;\geq 0 \qquad \text{and} \qquad Z_X \succeq 0 \;\; \text{(Loewner order)}. \tag{7.488}$$

Violations indicate ghost leakage or an incomplete BRST reduction; rollback to §3 and the identity ledger of §6.

Algorithm (auditable).

1. *Pole localisation.* On the joint plateau $\mathfrak{P}$ (§5.2), solve $\det \mathcal{H}^{XX}_{\mathrm{phys}}(k) = 0$ for $k^2 = m_X^2$; in practice, fit the inverse kernel to a rational form near the pole and cross-check with dispersion fits $\omega_X(|\mathbf{k}|)$.
2. *Residues.* Compute Z_X via eq. (7.486) or, with mixing, eq. (7.487); verify eq. (7.488).
3. *Acceptance.* Enforce $m_X^2 \geq 0$ and $Z_X \succeq 0$; store error bars from the plateau budgets (eq. (7.137)) and regulator spreads (eq. (7.67)).
4. *Kato continuity.* Check Lipschitz continuity of m_X^2, Z_X across $\mathfrak{P}$ as in §5.6; flag/resolve crossings and near-degeneracies.
5. *Nielsen stability.* Verify $\partial_\xi m_X^2 = \partial_\xi Z_X = 0$ within tolerance (cf. eqs. (7.464) and (7.467)).

Remarks. (i) NG modes (global G) correspond to $m_\pi^2 = 0$ at $|\mathbf{k}| \to 0$ with stiffness extracted from the small-k slope of the pole (cf. §4.5); (ii) in the gauged (BEH) case, only the transverse block contributes, with $m_{A^T} > 0$ and no massless TT pole (§4.6).

7.2 Principal symbol and hyperbolicity

Symbol and cone. Write the kinetic operator in each physical sector as a local differential operator $\mathcal{K}^{XX}_{\mathrm{phys}}(\partial)$ (quadratic order). Its principal symbol $\sigma_{\mathrm{prin}}(\mathcal{K}^{XX}_{\mathrm{phys}})(x,k)$ is the homogeneous highest-order part in k_μ. With respect to the chronogeometric cone (gaiamiti background metric), the hyperbolicity criteria recalled in eq. (7.15) are imposed on $\mathfrak{P}$.

Operational hyperbolicity tests (band, d dimensions).

1. **Real characteristics.** For each sector X, the characteristic polynomial
$$C_X(x,k) \;:=\; \det \sigma_{\mathrm{prin}}\Big(\mathcal{K}^{XX}_{\mathrm{phys}}\Big)(x,k) \tag{7.489}$$
has only real roots in k_0 for all spatial $\mathbf{k}$ sampled on $\mathfrak{P}$.

2. **Symmetrisers and energy estimates.** There exists a positive-definite symmetriser $S_X(x)$ such that $S_X\,\sigma_{\mathrm{prin}}(\mathcal{K}^{XX}_{\mathrm{phys}})$ is symmetric in k and yields an energy estimate uniform on $\mathfrak{P}$:
$$\frac{d}{dt}\,E_X(t) \;\le\; C_X\,E_X(t)\,, \qquad E_X(t) \;\sim\; \langle \Psi_X,\, S_X\,\Psi_X\rangle_{\mathrm{phys}}\,, \tag{7.490}$$
ensuring well-posed Cauchy evolution and finite propagation speed (domain of dependence within the background light cone).

3. **Causal dispersion.** From the pole condition $\det \mathcal{H}^{XX}_{\mathrm{phys}}(\omega,\mathbf{k}) = 0$ extract $\omega_X(\mathbf{k})$ and verify the group-velocity bound
$$\sup_{(u,v)\in\mathfrak{P}}\;\sup_{\mathbf{k}\in \mathrm{IR\,win}} \Big|\nabla_{\mathbf{k}}\,\omega_X(\mathbf{k})\Big| \;\le\; 1 \;+\; \varepsilon_{\mathrm{vel}}\,, \tag{7.491}$$
with uncertainties governed by eq. (7.137) (units $c = 1$).

Acceptance and reporting. Report: (i) verification of real characteristics eq. (7.489); (ii) symmetriser constants and energy-estimate bounds eq. (7.490); (iii) dispersion fits and the group-velocity metric eq. (7.491). All metrics obey the tolerance and robustness criteria of Section 7.7.5 (including Kato continuity, regulator spreads, and Nielsen stability). Failures (tachyonic roots, lack of symmetriser, superluminal drift beyond tolerance) trigger rollback to Steps 3–6 and exclusion from the joint plateau $\mathfrak{P}$.

7.3 Spectral positivity (Källén–Lehmann on the band)

Representation on the band. For each BRST–physical field/operator X (scalar or the transverse part of a vector), the band–projected Wightman two–point function admits a Källén–Lehmann representation compatible with the UV/IR pipeline (Section 7.3):
$$\langle X(x)\,X(0)\rangle_{\mathrm{band}} = \int_0^\infty d\mu^2\;\rho_X(\mu^2)\;\Delta_+^{(d)}(x;\mu^2)\,, \qquad \rho_X(\mu^2)\ge 0\,, \tag{7.492}$$
where $\Delta_+^{(d)}$ is the free positive–frequency kernel in d spacetime dimensions. In momentum space (with the implicit composition by $P_x(\lambda_{\mathrm{UV}})$ and $K^{(\ell)}_{\mathrm{IR}}$),
$$G_X(p) \equiv \big[\mathcal{H}^{XX}_{\mathrm{phys}}(p)\big]^{-1} = \int_0^\infty d\mu^2\;\frac{\rho_X(\mu^2)}{p^2-\mu^2+i0} \quad \text{on } \mathcal{H}_{\mathrm{phys}}\,. \tag{7.493}$$
For a spin–1 field, ρ_X refers to the transverse sector and the correlator carries the projector Π_{T} (cf. eqs. (7.320) and (7.336)).

Extraction and positivity. The band spectral density is obtained from the discontinuity of the physical Green function:

$$\rho_X(\mu^2) = \frac{1}{\pi} \lim_{\epsilon\to 0^+} \operatorname{Im} G_X\Big(p^2 = \mu^2 + i\epsilon\Big) \qquad \text{(on the certified band)}\,. \tag{7.494}$$

BRST cohomology (Section 7.2) guarantees that only cohomology representatives contribute: longitudinal/ghost/Nakanishi–Lautrup quartets cancel, so no negative–norm states enter ([442, 117]).

Discrete poles and continuum. Decompose ρ_X into pole and continuum parts,

$$\rho_X(\mu^2) = \sum_n Z_{X,n}\,\delta(\mu^2 - m_{X,n}^2) \;+\; \rho_X^{\text{cont}}(\mu^2)\,, \qquad Z_{X,n} \ge 0, \quad \rho_X^{\text{cont}}(\mu^2) \ge 0\,, \tag{7.495}$$

with $Z_{X,n}$ the (nonnegative) residues defined in eqs. (7.486) and (7.487). Unbroken/broken Ward and ST identities constrain sum rules among $\{Z_{X,n}\}$ and moments of ρ_X^{cont} (Section 7.5).

Acceptance suite. On the joint plateau $\mathfrak{P}$ (cf. §5.2) we require:

$$\rho_X(\mu^2) \ge 0 \quad \text{for a.e. } \mu^2 \text{ in the fitted support}, \qquad Z_{X,n} \ge 0 \quad \text{(all isolated poles)}\,, \tag{7.496}$$

together with regulator robustness and gauge independence within the established budgets:

$$\Delta_{\text{reg}}[\rho_X] \le \varepsilon_{\text{reg}}, \qquad \partial_\xi \rho_X = 0 \;\text{ within } \varepsilon_{\text{gauge}} \quad \text{(Nielsen on shell; cf. §5.5)}. \tag{7.497}$$

Violations (negative dips, gauge drift beyond tolerance) indicate leakage from nonphysical sectors or off–plateau evaluation and trigger a rollback to Steps 3–6.

Reporting. Report the pole list $\{m_{X,n}^2, Z_{X,n}\}$ with Kato–continuity trackers (cf. §5.6), the fitted ρ_X^{cont} with confidence bands, and Ward/ST–compatible sum rules used as cross–checks.

7.4 Microcausality and locality

Pauli–Jordan commutator on the band. Define the Pauli–Jordan function

$$\Delta_X(x) \;:=\; \langle [\,X(x), X(0)\,] \rangle_{\text{band}} = \int_0^\infty d\mu^2\; \rho_X(\mu^2)\; \Delta_{\text{PJ}}^{(d)}(x;\mu^2)\,, \tag{7.498}$$

where $\Delta_{\text{PJ}}^{(d)}$ is the standard Pauli–Jordan kernel in d dimensions. Local hyperbolic dynamics implies $\operatorname{supp}\Delta_{\text{PJ}}^{(d)} \subseteq \overline{\mathcal{C}}$ (the closed light cone), hence microcausality $\Delta_X(x) = 0$ for spacelike separations $x^2 < 0$.

Band effects and test. Finite UV/IR windows induce a controlled nonlocality (Gibbs tails for hard cutoffs, exponential or power–law for smooth windows). On a certified plateau ($\mathfrak{P}$) we require spacelike suppression bounded by a causal budget $\varepsilon_{\text{causal}}$ determined by the window shapes:

$$\sup_{x^2 \le -\sigma^2} \big|\Delta_X(x)\big| \;\le\; \varepsilon_{\text{causal}}(\lambda_{\text{UV}}, \ell; \sigma), \qquad \varepsilon_{\text{causal}} \xrightarrow[\lambda_{\text{UV}}\to\infty,\; \ell\to\infty]{} 0\,. \tag{7.499}$$

Operationally:

1. Reconstruct ρ_X from $\mathcal{H}_{\rm phys}^{-1}$ (cf. eq. (7.494)) and synthesise Δ_X via eq. (7.498).
2. Independently Fourier–transform the retarded/advanced propagators built from $\mathcal{H}_{\rm phys}$ on the band; confirm consistency of the spacelike suppression.
3. Verify that the measured $\varepsilon_{\rm causal}$ respects eq. (7.499) on $\mathfrak{P}$ and is robust across admissible $(P_x, K_{\rm IR})$ (cf. eq. (7.67)).

Acceptance and remedies. Persistent spacelike leakage (beyond $\varepsilon_{\rm causal}$) or gauge–parameter drift in Δ_X indicates either (i) failure of hyperbolicity (rollback to eq. (7.15)), (ii) off–shell evaluation (enforce the identity ledger of §6), or (iii) unsuitable windowing (adjust $(\lambda_{\rm UV}, \ell)$ or choose smoother $(P_x, K_{\rm IR})$). For gauge fields, microcausality is enforced for BRST–physical (transverse) observables; any apparent violation confined to gauge–variant sectors is dismissed by cohomological projection (Section 7.2).

7.5 Optical–theorem and unitarity proxies (on band)

Forward unitarity on the band. For BRST–physical channels, the optical theorem relates the imaginary part of the *forward* amplitude to a sum of positive transition probabilities. With amputated, band–projected, BRST–consistent external legs and vertices (ST/Nielsen enforced; cf. §6), the forward amplitude $\mathcal{M}_{i\to i}(s)$ satisfies

$$2\,\mathrm{Im}\,\mathcal{M}_{i\to i}(s) \;=\; \sum_{n\in\mathcal{H}_{\rm phys}} \int d\Pi_n^{(\rm band)} \left|\mathcal{M}_{i\to n}(s)\right|^2 \;\geq\; 0\,, \tag{7.500}$$

where $d\Pi_n^{(\rm band)}$ is the n–body phase space consistent with the IR window $K_{\rm IR}^{(\ell)}$ and UV projector $P_x(\lambda_{\rm UV})$ (cf. Section 7.3). Gauge–parameter independence of $\mathrm{Im}\,\mathcal{M}$ on shell follows from the Nielsen identity (cf. §6.3).

Cutkosky cuts with band windows. Let $\mathcal{H}_{\rm phys}$ generate the physical propagators and vertices. For any physical polarisation configuration, compute the cut discontinuity

$$\mathrm{Disc}\,\mathcal{M} \;=\; 2i\,\mathrm{Im}\,\mathcal{M} \;=\; \sum_{\rm cuts} \int d\Pi_{\rm cut}^{(\rm band)} \prod_{\text{cut lines}} \big(2\pi\big)\,\delta^{(+)}(k^2 - m^2)\,, \tag{7.501}$$

with all internal lines taken from the BRST–physical spectrum and the measure restricted by $(P_x, K_{\rm IR})$. *Acceptance:*

$$\mathrm{Im}\,\mathcal{M} \;\geq\; -\,\varepsilon_{\rm unit} \qquad \text{for all tested channels on } \mathfrak{P}, \tag{7.502}$$

with $\varepsilon_{\rm unit}$ chosen at or below the plateau budgets of eq. (7.137). Any systematic violation indicates either nonphysical leakage (incomplete BRST reduction) or off–plateau evaluation.

Partial waves in d dimensions. In the centre–of–mass frame, expand elastic $2\to2$ amplitudes in hyperspherical harmonics on S^{d-2}. Denote the partial–wave coefficients by $a_\ell(s)$

(with normalisation fixed so that the elastic unitarity circle applies). On the band and for BRST–physical external states,

$$|a_\ell(s)| \;\le\; 1, \qquad \operatorname{Im} a_\ell(s) \;\ge\; |a_\ell(s)|^2 \qquad \text{(elastic sector; inelasticity } \eta_\ell \le 1\text{)}, \tag{7.503}$$

within numerical tolerances dictated by $\mathfrak{P}$. We diagnose with the unitarity–circle distance

$$\delta_\ell(s) \;:=\; \Big| a_\ell(s) - \tfrac{i}{2} \Big| - \tfrac{1}{2} \;\le\; \varepsilon_{\mathrm{pw}}\,, \tag{7.504}$$

and the inequality defect

$$\Delta_\ell^{(\mathrm{inel})}(s) \;:=\; |a_\ell(s)|^2 - \operatorname{Im} a_\ell(s) \;\le\; \varepsilon_{\mathrm{pw}}\,. \tag{7.505}$$

Nielsen/ST stability. On $\mathfrak{P}$, the gauge–parameter derivatives of all proxies vanish within tolerance:

$$\partial_\xi \operatorname{Im} \mathcal{M} = 0, \quad \partial_\xi\, a_\ell(s) = 0 \qquad \text{(within } \varepsilon_{\mathrm{gauge}}\text{; cf. §5.5, §6.3)}. \tag{7.506}$$

Failure points to off–shell evaluation or identity–ledger violations (rollback to §6).

Reporting. We report, for each tested channel: (i) forward $\operatorname{Im}\mathcal{M}$ with band cuts and $\varepsilon_{\mathrm{unit}}$; (ii) $\{\delta_\ell, \Delta_\ell^{(\mathrm{inel})}\}$ vs. s; (iii) regulator and gauge spreads (see below). All quantities are quoted with plateau and robustness confidence intervals (cf. §5).

7.6 Regulator & gauge robustness of physicality

Spreads for the physical vector. Collect the physical outputs in $\mathbf{P} = \{\, m_X^2,\; Z_X,\; \rho_X(\mu^2),$ hyperbolicity metrics, $\operatorname{Im}\mathcal{M},\; a_\ell \,\}$.

Across an admissible family $\{(P^{(i)}, K^{(i)})\}$ and gauge parameters $\{\xi_k\}$ (cf. §5.4–§5.5), define

$$\Delta_{\mathrm{phys}}^{(\mathrm{reg})} := \max_{i,j} \sup_{(u,v)\in\mathfrak{P}} \big\| \mathbf{P}^{(i)}(u,v) - \mathbf{P}^{(j)}(u,v) \big\|, \qquad \Delta_{\mathrm{phys}}^{(\mathrm{gauge})} := \max_{k,k'} \sup_{(u,v)\in\mathfrak{P}} \big\| \mathbf{P}(\xi_k) - \mathbf{P}(\xi_{k'}) \big\|. \tag{7.507}$$

Here $\|\cdot\|$ denotes a channel–aware norm (block–diagonal over entries; spectral norm for matrices; L^∞ over μ^2 for ρ_X; sup over s for $\operatorname{Im}\mathcal{M}$ and a_ℓ).

Acceptance. We require

$$\Delta_{\mathrm{phys}}^{(\mathrm{reg})} \;\le\; \varepsilon_{\mathrm{reg}}\,, \qquad \Delta_{\mathrm{phys}}^{(\mathrm{gauge})} \;\le\; \varepsilon_{\mathrm{gauge}}\,, \tag{7.508}$$

with $\varepsilon_{\mathrm{reg}}$ controlled by eq. (7.67) and $\varepsilon_{\mathrm{gauge}}$ not exceeding the plateau tolerances of eq. (7.137). Any excess spread excludes the corresponding region from the joint plateau $\mathfrak{P}$ and triggers a rollback: rebuild $\mathcal{H}_{\mathrm{band}}$ (Step 3), revalidate the ledger (Step 6), and tighten the window family (Step 5).

Outcome. With eqs. (7.500), (7.503) and (7.508) satisfied on $\mathfrak{P}$, the physical content inferred from $\mathcal{H}_{\mathrm{phys}}$ is simultaneously unitary (to band accuracy), BRST/identity–consistent, and robust against admissible regulator and gauge variations.

7.7 Practical run-sheet (auditable tests)

1. **Pole extraction (mass & residues).** From $\mathcal{H}_{\mathrm{phys}}$ (cf. Section 7.7.3), locate poles of $\mathcal{H}_{\mathrm{phys}}^{-1}$ on the joint plateau $\mathfrak{P}$ (cf. §5):
   (a) Solve $\det \mathcal{H}_{\mathrm{phys}}^{XX}(p) = 0$ for each physical block $X \in \{A^T, h, \pi\}$ to obtain $m_X^2 \geq 0$ (cf. eq. (7.13)).
   (b) Compute residue matrices $Z_X = \lim_{p^2 \to m_X^2} (p^2 - m_X^2) \left[\mathcal{H}_{\mathrm{phys}}^{XX}(p)\right]^{-1}$ and enforce $Z_X \succcurlyeq 0$ (Källén–Lehmann consistency, eqs. (7.493) to (7.495)).
   (c) Track (m_X^2, Z_X) across $\mathfrak{P}$ and verify Kato–type continuity (Section 7.7.5).
2. **Hyperbolicity and causality.** Build the principal symbols $\sigma_{\mathrm{prin}}(\mathcal{K}_{\mathrm{phys}}^{XX})$ and check:
   (a) Real characteristics in k_0 for all spatial $\mathbf{k}$ (eq. (7.15)).
   (b) Uniform energy estimates (well-posed Cauchy problem) on $\mathfrak{P}$.
   (c) Group velocities at poles satisfy $v_g = \left|\partial\omega/\partial|\mathbf{k}|\right| \leq 1$ within the plateau tolerances of eq. (7.137).
3. **Spectral positivity (KL).** Reconstruct $\rho_X(\mu^2)$ from discontinuities of $\mathcal{H}_{\mathrm{phys}}^{-1}$ (cf. eq. (7.494)) and verify:
   (a) $\rho_X(\mu^2) \geq 0$ almost everywhere and $Z_{X,n} \geq 0$ for discrete poles (eqs. (7.492), (7.495) and (7.496)).
   (b) Ward/ST-compatible sum rules on pole/continuum weights (Section 7.5).
4. **Microcausality.** Form the Pauli–Jordan commutator $\Delta_X(x)$ from ρ_X (cf. eq. (7.498)) and confirm spacelike suppression consistent with band windowing (eq. (7.499)); cross-check with retarded/advanced reconstructions.
5. **Unitarity proxies.**
   (a) Compute band Cutkosky cuts and verify $\mathrm{Im}\,\mathcal{M} \geq -\varepsilon_{\mathrm{unit}}$ for physical polarisations (eqs. (7.501) and (7.502)).
   (b) For elastic $2 \to 2$ sub-blocks, evaluate partial waves and enforce $|a_\ell(s)| \leq 1$ and $\mathrm{Im}\, a_\ell(s) \geq |a_\ell(s)|^2$ within $\varepsilon_{\mathrm{pw}}$ (eqs. (7.503) to (7.505)).
6. **Robustness and gauge independence.** Repeat the suite across admissible (P_x, K_{IR}) and gauges ξ; certify spreads $\Delta_{\mathrm{phys}}^{(\mathrm{reg})}, \Delta_{\mathrm{phys}}^{(\mathrm{gauge})}$ within $\varepsilon_{\mathrm{reg}}, \varepsilon_{\mathrm{gauge}}$ (cf. eqs. (7.67), (7.137), (7.507) and (7.508)) and Nielsen/ST stability (§6).

7.8 Failure modes and remedies

- *Tachyons* ($m_X^2 < 0$). Indicates an unstable background φ or wrong order of limits. **Remedy:** revisit the seed protocol and ordered limits (Section 7.7.2); rebuild $\mathcal{H}_{\text{band}}$ (eq. (7.311)) and re-test hyperbolicity (eq. (7.15)) on $\mathfrak{P}$.

- *Ghost leakage / negative residues.* Non-positive Z_X or negative spectral weight signals incomplete BRST reduction. **Remedy:** enforce quartet decoupling (eqs. (7.328) and (7.332)) and on-shell simplifications (eq. (7.336)); re-validate the ST/Nielsen ledger (Section 7.7.6).

- *Hyperbolicity loss / superluminality.* Violations of eq. (7.15) or $v_g > 1$. **Remedy:** tighten $(\lambda_{\text{UV}}, \ell)$ windows (Section 7.3), prefer smoother regulators, and exclude offending regions from $\mathfrak{P}$ (cf. §5.4–§5.6).

- *Spectral negativity.* Negative dips in ρ_X or inconsistent sum rules. **Remedy:** re-check BRST projection and Ward/ST constraints (Section 7.5); apply anomaly filters (Section 7.7.6, §6.5) and recompute discontinuities (eq. (7.494)).

- *Microcausality violation.* Spacelike leakage beyond $\varepsilon_{\text{causal}}$. **Remedy:** reduce IR support or soften UV edges (adjust $K_{\text{IR}}^{(\ell)}$, P_x); verify commutation of limits and regulator robustness on $\mathfrak{P}$ (cf. §5.4, eq. (7.499)).

- *Unitarity-proxy failure.* Systematic $\operatorname{Im}\mathcal{M} < 0$ or partial-wave defects beyond ε_{pw}. **Remedy:** ensure amplitudes are built from $\mathcal{H}_{\text{phys}}$ with BRST-physical external states; re-impose ST/Nielsen constraints (Section 7.7.6) and refine the band cuts (eq. (7.501)).

- *Regulator/gauge sensitivity.* Excess spreads $\Delta_{\text{phys}}^{(\text{reg})}$ or $\Delta_{\text{phys}}^{(\text{gauge})}$. **Remedy:** restrict to BRST-closer windows, narrow the admissible (P_x, K_{IR}) family, and rebuild the joint plateau per §5.2–§5.5; re-certify ledger closure (§6).

Outcome. On a certified joint plateau $\mathfrak{P}$, the broken-state dynamics on $\mathcal{H}_{\text{phys}}$ passes the auditable health checks: non-tachyonic spectra, causal hyperbolic propagation, positive spectral weights, and band-accurate unitarity proxies—robust under admissible regulator and gauge variations. Any emergent phenomenological labels ("Higgs-like", "graviton-like", "apeiron/chrono–kymon") remain *provisional* pending the cross-quadrant and stability criteria already mandated (cf. Section 7.4).

7.7.8 Readout Consolidation and Cross–Quadrant Comparison

Aim. To conclude the Quadrant II run, we (i) *consolidate* all visibilities and spectral diagnostics into plateau–certified summaries; (ii) produce *reporting artifacts* (tables/figures) whose content is regulator– and gauge–stable per Sections 7.7.5 and 7.7.6; and (iii) execute a *cross–quadrant comparison* to distinguish QII (SSB) from QI/QIII/QIV based on law/state visibilities, Ward/Nielsen structure, and spectral health (Sections 7.7.3, 7.7.4 and 7.7.7; [91, 82, 442, 117, 243, 114]).

8.1 Plateau–certified aggregation of visibilities

Let $(u, v) = (\ln \lambda_{\mathrm{UV}}, \ln \ell)$ and let $\mathfrak{P}$ be the joint plateau from Section 7.7.5. For each readout $R_\alpha \in \{\Lambda_{\mathrm{law}}, \Lambda_{\mathrm{state}}\}$ define the plateau aggregates

$$\overline{R}_\alpha \equiv \mathrm{Avg}_{(u,v)\in\mathfrak{P}}[R_\alpha], \qquad \Delta_\alpha^{\max} \equiv \sup_{(u,v)\in\mathfrak{P}} \left|R_\alpha - \overline{R}_\alpha\right|, \qquad \mathcal{G}_\alpha^{\max} \equiv \sup_{(u,v)\in\mathfrak{P}} \sqrt{\left|\partial_u R_\alpha\right|^2 + \left|\partial_v R_\alpha\right|^2}. \tag{7.509}$$

Acceptance (law channel). We require

$$\left|1 - \overline{\Lambda}_{\mathrm{law}}\right| \;\le\; \varepsilon_{\mathrm{plat}}, \qquad \max\left\{\Delta_{\mathrm{law}}^{\max},\; \mathcal{G}_{\mathrm{law}}^{\max}\right\} \;\le\; \text{bounds in } \textit{eq.}\ (7.137), \tag{7.510}$$

which implements presentation invariance on certified windows (cf. eq. (7.259)).

Finite–size signature (state channel). At fixed $(u, v) \in \mathfrak{P}$ and fixed phase–twist protocol, the visibility exhibits volume suppression as in Section 7.7.4:

$$\lim_{V\to\infty} |\Lambda_{\mathrm{state}}(\lambda_{\mathrm{UV}}, \ell; V)| = 0. \tag{7.511}$$

with fitted rate/exponent reported alongside the plateau diagnostics.

Integrated (marginal) visibilities. For any admissible weight w supported on a connected component of $\mathfrak{P}$,

$$\overline{R}_\alpha^{[u]}(v) = \int_{u_1}^{u_2} w(u; v)\, R_\alpha(u, v)\, du, \quad \overline{R}_\alpha^{[v]}(u) = \int_{v_1}^{v_2} w(v; u)\, R_\alpha(u, v)\, dv, \qquad \int w = 1, \tag{7.512}$$

inherit the same bounds as eq. (7.137) (Jensen/mean–value control; cf. Section 7.4.5). Discrepancies between pointwise and integrated aggregates beyond tolerance flag hidden micro–oscillations or insufficient windowing and trigger a revisit of the plateau mask.

Report. We quote $\overline{\Lambda}_{\mathrm{law}}, \overline{\Lambda}_{\mathrm{state}}$ with $\Delta^{\max}$ and $\mathcal{G}^{\max}$, plus integrated counterparts (7.512), all confined to $\mathfrak{P}$ and accompanied by regulator/gauge spreads per eqs. (7.67) and (7.137).

8.2 Spectral and identity–ledger overlays

From the block reduction in Section 7.7.3 and the physicality checks of §7, collect (for each physical channel) the pole masses $\{m_i^2\}$, residues $\{Z_i\}$, and hyperbolicity metrics, and overlay them with the identity–ledger residuals of Section 7.7.6. For any scalar observable $O \in \{m_i^2, Z_i, \text{hyperbolicity figures}\}$ define the plateau spread and the Nielsen slope:

$$\mathrm{Spread}_{\mathfrak{P}}(O) \equiv \sup_{(u,v)\in\mathfrak{P}} O(u, v) - \inf_{(u,v)\in\mathfrak{P}} O(u, v), \qquad \mathrm{NielsenSlope}(O) \equiv \sup_{(u,v)\in\mathfrak{P}} \left|\partial_\xi O(u, v)\right|. \tag{7.513}$$

Acceptance.

$$\begin{aligned} \mathrm{Spread}_{\mathfrak{P}}(O) &\le \text{bounds in eq. (7.137)}, \\ \mathrm{NielsenSlope}(O) &\le \varepsilon_{\mathrm{gauge}}. \end{aligned} \tag{7.514}$$

(gauge–parameter independence on shell; cf. §6.3.) When O is matrix–valued (e.g. residue matrices), $\mathrm{Spread}_{\mathfrak{P}}$ is taken in the operator norm and NielsenSlope in the induced matrix norm. All overlays are reported together with the ST master–identity residuals (sup–norm on $\mathfrak{P}$) to document that spectral stability coincides with ledger closure (cf. §6.1–§6.2).

Breaking pattern and particle–like census (formal summary). Let $G \to H$ be the spontaneous breaking pattern realised by a vacuum v. On Lie algebras,

$$\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{m}, \qquad r \equiv \dim \mathfrak{m} = \dim G - \dim H,$$

where $\mathfrak{m}$ is spanned by the broken generators. Let $\mathfrak{g}_{\text{gauge}} \subseteq \mathfrak{g}$ denote the gauged subalgebra and $\mathfrak{h}_{\text{gauge}} = \mathfrak{g}_{\text{gauge}} \cap \mathfrak{h}$ its unbroken part. Define the broken/gauged counts

$$r_{\text{g}} \equiv \dim \mathfrak{g}_{\text{gauge}} - \dim \mathfrak{h}_{\text{gauge}}, \qquad r_{\text{gl}} \equiv r - r_{\text{g}}, \tag{7.515}$$

and let N_s be the number of real scalar directions active near v (so $N_s \geq r$). All statements below refer to the BRST cohomology $\mathcal{H}_{\text{phys}}$ and are to be evaluated on certified plateaus (cf. Section 7.7.5).

(a) Chronogeometric sector. Massless spin-2 tensor (*provisional,* "graviton-like") with 2 helicities; unaffected by internal SSB in this setup (cf. Sections 7.4 and 7.7.7).

(b) Unbroken gauged generators $\mathfrak{h}_{\text{gauge}}$. $\dim \mathfrak{h}_{\text{gauge}}$ massless vectors (spin-1; physical polarisations only after BRST projection), persisting from the unbroken gauge symmetry (cf. Section 7.7.3).

(c) Broken generators $\mathfrak{m}$: global vs. gauged.

- *Global broken* (r_{gl}): r_{gl} Nambu–Goldstone scalars (spin-0, gapless in the $V \to \infty$ limit) in the relativistic/Lorentz-invariant case; see the caveat on type-B modes below.
- *Gauged broken* (r_{g}): r_{g} massive vectors (spin-1) via BEH reorganisation; the corresponding would-be NG modes are eaten and are not counted separately (cf. Sections 7.7.2 and 7.7.4).

(d) Radial/Higgs-like scalars. $(N_s - r)$ massive spin-0 excitations along directions orthogonal to the vacuum manifold; one physical DoF each. These pass the physicality tests on $\mathcal{H}_{\text{phys}}$ (cf. eqs. (7.13) and (7.15) and Section 7.7.7).

Dimensional DoF counts. In d spatial dimensions, a massless (massive) spin-1 field carries $d-1$ (d) physical polarisations after BRST projection. Thus the bosonic DoF ledger reads

$$\underbrace{(d-1)\,\dim \mathfrak{g}_{\text{gauge}} + N_s}_{\text{before breaking}} = \underbrace{(d-1)\,\dim \mathfrak{h}_{\text{gauge}}}_{\text{unbroken gauge}} + \underbrace{d\, r_{\text{g}}}_{\text{BEH massive vectors}} + \underbrace{r_{\text{gl}}}_{\text{NG scalars}} + \underbrace{(N_s - r)}_{\text{radial/Higgs-like}}, \tag{7.516}$$

using $r = r_{\text{g}} + r_{\text{gl}}$. Specialising to d=3 (3+1D) reproduces the familiar numbers (massless vector: 2 helicities; massive vector: 3 polarisations).

Closed-form counts (3+1D DoF).

Massless vectors:	$2\,\dim \mathfrak{h}_{\text{gauge}}$,	
Massive vectors (BEH):	$3\,r_{\text{g}}$,	(massless spin-2 carries 2 helicities;
NG scalars (in spectrum):	r_{gl},	*provisional* identification).
Radial/Higgs-like scalars:	$N_s - r$,	

DoF bookkeeping (consistency check). Before breaking: $(d-1)\dim\mathfrak{g}_{\text{gauge}}+N_s$ bosonic DoF (gauge vectors + scalars). After breaking, the RHS of eq. (7.516) equals the same quantity; hence bosonic DoF are conserved and each eaten NG supplies the longitudinal polarisation of a BEH massive vector (cf. Sections 7.7.3 and 7.7.6).

Remarks (scope and caveats).

- *Type-B NG at finite densities.* In non-Lorentz-invariant settings (e.g. nonzero charge densities), NG counting can reduce: $N_{\text{NG}}=r_{\text{gl}}-\frac{1}{2}\operatorname{rank}\mathcal{C}$ with $\mathcal{C}_{ab}\propto\lim_{V\to\infty}V^{-1}\langle[Q_a,Q_b]\rangle$ (type-B pairs with quadratic dispersion). Our default census assumes the relativistic/Lorentz-invariant regime used in the band reduction; deviations must be reported with the same plateau certification and identity-ledger checks.
- *Dimensional quirks/topological terms.* In lower d or in the presence of topological masses (e.g. Chern–Simons in 2+1D), vector DoF and dispersion change; any such cases must be flagged explicitly and recast in the d-dimensional ledger eq. (7.516).
- *BRST cohomology.* All counts are made on $\mathcal{H}_{\text{phys}}$; ghost/longitudinal quartets cancel and do not contribute (cf. Section 7.7.3).

Particle-like modes in Quadrant II (SSB) on $\mathcal{H}_{\text{phys}}$. Law is held symmetric (Section 7.7.1); the broken state is prepared and analysed on certified plateaus with BRST/Slavnov–Taylor/Nielsen protection (Sections 7.7.2 to 7.7.7). Tags marked "provisional'' require cross-quadrant confirmation (Section 7.4). Counts below are quoted for 3+1D; the general d-dimensional ledger appears in eq. (7.516).

(1) Gaiamiti $\hat{\Phi}$ (chronogeometric). *Mode:* tensor (graviton-like; *provisional*); *Spin:* 2; *Gap:* massless; *DoF (3+1D):* 2 (helicities ±2). Rationale: retains the unbroken QI character on the band; identification is provisional pending cross-quadrant checks (Sections 7.4 and 7.7.7).

(2) Tantrakyma $\hat{\Theta}$ (unbroken H). *Mode:* gauge vector(s) (unbroken); *Spin:* 1; *Gap:* massless; *DoF (3+1D):* 2 (helicities ±1). Verified via BRST projection and spectral positivity on plateaus (Sections 7.7.3 and 7.7.7).

(3) Global SSB (broken global G). *Mode:* Nambu–Goldstone boson(s) π^a; *Spin:* 0; *Gap:* gapless (massless as $V\to\infty$); *DoF:* 1 per broken generator (count $=\dim G-\dim H$ in the Lorentz-invariant regime). Diagnosed by phase–twist interferometry and Ward/Nielsen-protected soft structure (Sections 7.5 and 7.7.4). *Caveat:* in non-Lorentz-invariant settings (finite densities) type-B pairing can reduce the count; deviations must be noted explicitly (cf. discussion around eq. (7.516)).

(4) Gauged SSB (BEH reorganisation). (a) *Massive gauge vector(s)*; *Spin:* 1; *Gap:* $m_V^2>0$; *DoF (3+1D):* 3 (helicities ±1 plus longitudinal); *Count:* $\dim G_{\text{gauge}}-\dim H_{\text{gauge}}$. (b) *Would-be NG modes:* eaten (not counted separately). Verified by pole/residue extraction and Nielsen-stable masses on $\mathfrak{P}$ (Sections 7.7.3 and 7.7.7).

(5) Scalar sector in $\hat{\Phi},\hat{\Theta}$ (radial). *Mode:* radial / Higgs-like excitations; *Spin:* 0; *Gap:* gapped ($m_h^2>0$); *DoF:* 1 per independent radial direction (model-dependent). Certified by nonnegative mass operator, hyperbolicity, and spectral positivity (eqs. (7.13) and (7.15) and Section 7.7.7).

Notes. (i) Items separate *global* from *gauged* breaking: NG bosons appear only for global G, while in gauged phases they reorganise via BEH into longitudinal vector polarisation (Sections 7.7.2 and 7.7.4). (ii) All statements are evaluated *on certified plateaus* and are gauge/regulator-robust within the bounds of eqs. (7.67) and (7.137). (iii) Chronogeometric identifications ("graviton-like"), as well as any "apeiron/chrono–kymon" tags, remain *provisional* until confirmed by the cross-quadrant programme (Section 7.4).

Quadrant II in one view. The SSB analysis freezes the *law* while varying only the *state*. Concretely, we keep the symmetric kinetic baseline H_0 fixed on both law branches and verify unity of the law visibility by interferometric calibration (Sections 7.4 and 7.7.1). All measurements are performed *on the band* with UV/IR controls (Section 7.3), and quantitative statements are made *at plateau* where gradients/oscillations/curvature obey eq. (7.137). On shell, the Slavnov–Taylor, Ward, and Nielsen identities protect both readouts and spectra, rendering them gauge–parameter independent and anomaly–filtered within the certified window (Section 7.5; [117, 442, 243]).

Preparing and diagnosing the broken state. An infinitesimal seed selects a vacuum on $\mathcal{M}_{\rm vac}\cong G/H$ without altering H_0 (Section 7.7.2). Around the chosen background, the band Hessian is assembled, split into transverse/longitudinal sectors, and projected to BRST cohomology so that unphysical quartets decouple (Sections 7.2 and 7.7.3; [442, 117]). The holosymmetric probe then reads *visibilities*: $|\Lambda_{\rm law}|=1+\mathcal{O}(\varepsilon_{\rm plat})$ by construction, while the *state* visibility exhibits finite–size suppression at fixed plateau—the operational hallmark of spontaneous selection (Section 7.7.4 and eq. (7.139)). Phase–twist interferometry ties this suppression to current fluctuations and, via Ward/Nielsen protection, to NG soft structure for global G or to BEH reorganisation for gauged G (Sections 7.5 and 7.7.4; [91, 82]).

Certified stability of the claims. Plateau certification provides joint masks where law/state visibilities, physical poles/residues, and identity–ledger residuals are simultaneously stable (Section 7.7.5). On these domains we verify (i) nonnegativity of the physical mass operator and (ii) hyperbolicity of the principal symbol—hence causal propagation with no tachyons—together with (iii) Källén–Lehmann positivity for band–projected correlators (cf. eqs. (7.13) and (7.15); Section 7.7.7; [91, 82]). Gauge and regulator robustness are enforced by Nielsen stability and regularisation bounds (Section 7.7.6 and eq. (7.67)), so the final readouts and figures are reproducible within certified tolerances (Section 7.7.8).

"Particle–like" content and mode taxonomy. On $\mathcal{H}_{\rm phys}$ the broken–state spectrum organises as follows (Sections 7.7.3 and 7.7.7). *Global G:* massless Nambu–Goldstone scalars π^a (one per broken generator in the Lorentz–invariant regime) carry the soft pole dictated by the broken Ward identities; massive h–type scalars populate radial directions; spectator vectors in unbroken subgroups remain in their QI pattern ([91]). *Gauged G (BEH):* would–be NG modes are absorbed; transverse gauge bosons acquire masses and appear as physical spin–1 poles with positive residues, while longitudinal artefacts remain BRST–exact and do not contribute; scalar excitations in radial directions manifest as "Higgs–like" modes with plateau–stable masses and residues (Sections 7.7.4 and 7.7.7; [82, 117]). The chronogeometric (gaiamiti)

sector's massless spin–2, when present, retains its unbroken QI character on the band; any "graviton–like'' or "apeiron/chrono–kymon'' tags are *provisional* pending cross–quadrant checks (Section 7.4).

Outcome and discrimination. Taken together, QII is singled out by (i) *unity of the law* within eq. (7.137), (ii) *volume decay* of $|\Lambda_{\text{state}}|$ at fixed plateau, and (iii) a mode census consistent with NG/BEH expectations and the physicality tests cited above (Sections 7.7.4 and 7.7.7). The consolidated artefacts (maps, scaling plots, spectral overlays) are regulator/-gauge–stable and support a clean separation from QI (symmetric), QIII (emergent symmetry under coarse graining), and QIV (explicit breaking) (Section 7.7.8). Any phenomenological labels—"Higgs–like", "graviton–like", "apeiron/chrono–kymon''—are highlighted but explicitly flagged as *provisional* until validated by the multi–quadrant programme (Section 7.4; [243, 114]).

7.8 Quadrant III: Explicit (But Small) Law Breakings

Aim and posture. In QIII we deliberately introduce *explicit* symmetry-breaking terms in the *law* channel while insisting on a *symmetric/coherent* state. Operationally, we reuse the band-limited instrumentation ($P_x(\lambda_{\text{UV}})$, $K_{\text{IR}}^{(\ell)}$) and readouts defined earlier—coherence amplitudes $\mathcal{C}_\alpha$ and visibilities Λ_α in eqs. (7.115), (7.116), (7.122) and (7.129)—together with the acceptance/plateau pipeline of Sections 6.12.4 and 7.3.9. BRST/Slavnov–Taylor (ST) consistency is maintained on shell throughout, as in eq. (7.16) ([104, 445, 105, 106, 442, 117, 243]).

7.8.1 Data, Regime, and Physical Reading

Definition 7.346 (Quadrant III data and small-breaking regime)**.** Work on the band and within the IR window (UV projector $P_x(\lambda_{\text{UV}})$, IR kernel $K_{\text{IR}}^{(\ell)}$; cf. Section 6.12.4 and Definition 6.282). Keep the *state channel symmetric* (backgrounds from Section 7.1.5), and introduce *explicit but small* law breakings in the Lagrangian,

$$\mathcal{L} \;=\; \mathcal{L}_0 \;+\; \sum_k \epsilon_k\, \mathcal{O}_k \;+\; \zeta\Psi,$$

where $\zeta(\,\cdot\,) := \{\hat{Q}_{\text{BRST}}, \cdot\,\}$, $\delta_a\mathcal{L}_0 = \nabla_\mu K_a^\mu$, $\delta_a\mathcal{O}_k = \mathcal{S}_{k,a}$ (all band/IR–tagged), and $\{\epsilon_k\}$ are scale–small couplings. Let Δ_k be the RG scaling dimensions of the symmetry–breaking operators and set $\Delta := \min_k \Delta_k$. All statements are evaluated on $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$ with BRST–closed tags and channel projectors (cf. Section 7.1.3).

Proposition 7.347 (Scale–tagged Ward source and physical reduction)**.** *The adjoint–covariant, scale–tagged Ward identity is*

$$\nabla_\mu^{(\text{ad})} J_a^{(\lambda_{\text{UV}},\ell)\mu} = \Delta_a^{(\lambda_{\text{UV}},\ell)}, \qquad \nabla_\mu^{(\text{ad})} J^\mu := \nabla_\mu J^\mu + i[\mathcal{A}_\mu, J^\mu],$$

with (cf. Definition 6.295)

$$\Delta_a^{(\lambda_{\text{UV}},\ell)} = \sum_k \epsilon_k(\ell)\left\langle \mathcal{S}_{k,a}^{(\lambda_{\text{UV}})}\right\rangle_{\boldsymbol{\Psi}} \;+\; \left\langle \zeta\,\eta_a^{(\lambda_{\text{UV}},\ell)}\right\rangle_{\boldsymbol{\Psi}} \;+\; \mathcal{A}_a^{(\lambda_{\text{UV}},\ell)}.$$

On $\mathcal{H}_{\mathrm{phys}}$ and with anomaly-free BRST the last two terms vanish, so

$$\nabla^{(\mathrm{ad})}_{\mu}\,\bar{J}^{(\lambda_{\mathrm{UV}},\ell)\mu}_{a,\mathrm{imp}} = \sum_{k} \epsilon_k(\ell)\,\left\langle \mathcal{S}^{(\lambda_{\mathrm{UV}})}_{k,a} \right\rangle_{\mathbf{\Psi}},$$

in agreement with eq. (6.377) and eq. (6.336) (see also Lemma 6.233). References: *[117, 82, 230].*

Proof. Start from the scale–tagged, gauge–fixed Lagrangian decomposition (UV then IR; tags suppressed for readability)

$$\mathcal{L} = \mathcal{L}_0 + \sum_k \epsilon_k\,\mathcal{O}_k + \zeta\Psi,$$

and let δ_a be a (possibly local/relational) variation with $\delta_a\mathcal{L}_0 = \nabla_\mu K^\mu_a$ and $\delta_a\mathcal{O}_k = \mathcal{S}_{k,a}$. By the relational Ward lemma Lemma 6.233 (Noether–Belinfante improvement absorbing $\nabla_\mu K^\mu_a$), there exists a local functional η_a generated by $\delta_a\Psi$ such that the *off–shell* identity holds

$$\nabla_\mu J^\mu_{a,\mathrm{imp}} = \sum_k \epsilon_k\,\mathcal{S}_{k,a} + \zeta\,\eta_a + \mathcal{A}_a. \tag{7.517}$$

Adjoint covariantization. If G is gauged, promote the divergence to the adjoint–covariant one, $\nabla^{(\mathrm{ad})}_\mu J^\mu := \nabla_\mu J^\mu + i[\mathcal{A}_\mu, J^\mu]$, obtaining

$$\nabla^{(\mathrm{ad})}_\mu J^\mu_{a,\mathrm{imp}} = \sum_k \epsilon_k\,\mathcal{S}_{k,a} + \zeta\,\eta_a + \mathcal{A}_a, \tag{7.518}$$

as in Definition 6.295 and standard derivations [117, 82, 230].

Scale tagging. Apply the band/IR tagging functor (UV projector $P_x(\lambda_{\mathrm{UV}})$ and IR window $K^{(\ell)}_{\mathrm{IR}}$; cf. Definition 6.282 and Section 6.12.4). By the assumptions used in Section 6.12.4 and Section 7.1.3, $[\nabla_\mu, P_x(\lambda_{\mathrm{UV}})] = 0$ and the scalar smearing $K^{(\ell)}_{\mathrm{IR}}$ commutes with covariant differentiation and with the channel projectors; moreover, both tags are BRST–closed (ghost number 0). Therefore (7.518) lifts directly to the scale–tagged form

$$\nabla^{(\mathrm{ad})}_\mu J^{(\lambda_{\mathrm{UV}},\ell)\mu}_{a,\mathrm{imp}} = \sum_k \epsilon_k(\ell)\,\mathcal{S}^{(\lambda_{\mathrm{UV}})}_{k,a} + \zeta\,\eta^{(\lambda_{\mathrm{UV}},\ell)}_a + \mathcal{A}^{(\lambda_{\mathrm{UV}},\ell)}_a,$$

where the notation matches Definition 6.295 and collects the tagged sources into $\Delta^{(\lambda_{\mathrm{UV}},\ell)}_a := \sum_k \epsilon_k(\ell)\,\langle \mathcal{S}^{(\lambda_{\mathrm{UV}})}_{k,a}\rangle_{\mathbf{\Psi}} + \langle \zeta\,\eta^{(\lambda_{\mathrm{UV}},\ell)}_a\rangle_{\mathbf{\Psi}} + \mathcal{A}^{(\lambda_{\mathrm{UV}},\ell)}_a$.

Physical reduction. Taking expectation values in a homogeneous state $|\mathbf{\Psi}\rangle$ and projecting with $\mathsf{P}_{\mathrm{phys}}$ to $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ yields

$$\nabla^{(\mathrm{ad})}_\mu \bar{J}^{(\lambda_{\mathrm{UV}},\ell)\mu}_{a,\mathrm{imp}} = \sum_k \epsilon_k(\ell)\,\left\langle \mathcal{S}^{(\lambda_{\mathrm{UV}})}_{k,a} \right\rangle_{\mathbf{\Psi}} + \left\langle \zeta\,\eta^{(\lambda_{\mathrm{UV}},\ell)}_a \right\rangle_{\mathbf{\Psi}} + \mathcal{A}^{(\lambda_{\mathrm{UV}},\ell)}_a.$$

On the anomaly–free Stage I, the last term vanishes; and in BRST cohomology at ghost number 0, ζ–exact insertions vanish (Nielsen–type independence; cf. Section 7.1.3). Hence

$$\nabla^{(\mathrm{ad})}_\mu \bar{J}^{(\lambda_{\mathrm{UV}},\ell)\mu}_{a,\mathrm{imp}} = \sum_k \epsilon_k(\ell)\,\left\langle \mathcal{S}^{(\lambda_{\mathrm{UV}})}_{k,a} \right\rangle_{\mathbf{\Psi}},$$

which is exactly eq. (6.377) and agrees with eq. (6.336) (see also Lemma 6.233). This proves the proposition. □

Proposition 7.348 (Law deficit scaling and small-breaking control)**.** *Let $\Lambda_{\mathrm{law}}^{(\lambda_{\mathrm{UV}},\ell)}$ be the (channel-resolved) law visibility (cf. Section 6.12.3) evaluated on $\mathcal{H}_{\mathrm{phys}}$ with BRST–closed tags. Assume the breakings are RG–irrelevant with minimal scaling dimension $\Delta > d + z$. Then for fixed UV band and large IR window,*

$$1 - |\Lambda_{\mathrm{law}}^{(\lambda_{\mathrm{UV}},\ell)}| = \mathcal{O}\Big(\ell^{\,\Delta-(d+z)}\Big),$$

as stated in Proposition 6.277. Moreover, if $\mathcal{O}(\lambda_{\mathrm{UV}}, \ell)$ (a ghost–number–zero diagnostic of the same breaking) satisfies the plateau criterion Section 6.12.4 on a box $\mathcal{B}$, its extraction is regulator–stable within $\mathcal{B}$. Idea of proof. *Combine the physical Ward identity above with clustering under $K_{\mathrm{IR}}^{(\ell)}$ and the scaling of the integrated insertions $\mathcal{S}_{k,a}$; use Proposition 6.298 to pass to plateau-stable readouts.*

Proof. Work on the BRST–closed physical sector $\mathcal{H}_{\mathrm{phys}}$ with UV/IR tags (gh$= 0$) and use the adjoint–covariant, scale–tagged Ward identity of Proposition 7.347:

$$\nabla_\mu^{(\mathrm{ad})} \bar{J}_{a,\mathrm{imp}}^{(\lambda_{\mathrm{UV}},\ell)\mu} = \sum_k \epsilon_k(\ell) \left\langle \mathcal{S}_{k,a}^{(\lambda_{\mathrm{UV}})} \right\rangle_{\mathbf{\Psi}} \quad \text{(anomaly–free BRST)}. \tag{7.519}$$

Step 1: From Ward source to a bound on the visibility deficit. Let W_ℓ be the (spatio–temporal) IR window with characteristic scale ℓ (product of the spatial kernel $K_{\mathrm{IR}}^{(\ell)}$ and a time window of width $\asymp \ell^z$; cf. Section 6.12.4). Integrate (7.519) against W_ℓ and use Stokes' theorem in adjoint form to relate the flux imbalance to the windowed source:

$$\left| \int d^{d+1}x\, W_\ell(x)\, \nabla_\mu^{(\mathrm{ad})} \bar{J}_{a,\mathrm{imp}}^{\mu}(x) \right| = \left| \int d^{d+1}x\, W_\ell(x) \sum_k \epsilon_k(\ell)\, \langle \mathcal{S}_{k,a}^{(\lambda_{\mathrm{UV}})}(x) \rangle_{\mathbf{\Psi}} \right|.$$

The law visibility $\Lambda_{\mathrm{law}}^{(\lambda_{\mathrm{UV}},\ell)}$ (cf. Section 6.12.3 and eq. (7.122)) compares the two law branches through this flux; by standard Cauchy–Schwarz/triangle estimates (as in Proposition 7.357's Dyson bound together with the branch equivalence Proposition 7.365) one obtains

$$1 - |\Lambda_{\mathrm{law}}^{(\lambda_{\mathrm{UV}},\ell)}| \;\le\; C_0 \int d^{d+1}x\, |W_\ell(x)| \sum_k |\epsilon_k(\ell)| \left| \langle \mathcal{S}_{k,a}^{(\lambda_{\mathrm{UV}})}(x) \rangle_{\mathbf{\Psi}} \right| \;+\; \mathcal{O}(\varepsilon_{\mathrm{soft}}^2), \tag{7.520}$$

with C_0 independent of $(\lambda_{\mathrm{UV}}, \ell)$ on a plateau window (cf. Proposition 7.359).

Step 2: Scaling of the windowed source. Assume the breakings are RG–irrelevant with minimal scaling dimension $\Delta > d + z$ (definition used in Section 6.12.3). Denote $S_k := \mathcal{S}_{k,a}$ for brevity. Under the anisotropic scaling $(t, \mathbf{x}) \mapsto (b^z t, b\mathbf{x})$, a correlator of a local density S_k scales as $b^{-\Delta_k}$ of its argument; the normalised window scales as $W_\ell(x) = \ell^{-(d+z)} W_1\Big(t/\ell^z, \mathbf{x}/\ell\Big)$. Hence

$$\int d^{d+1}x\, |W_\ell(x)| \left| \langle S_k^{(\lambda_{\mathrm{UV}})}(x) \rangle_{\mathbf{\Psi}} \right| = \ell^{-(d+z)} \int d^{d+1}y\, |W_1(y)| \left| \langle S_k^{(\lambda_{\mathrm{UV}})}(\ell^z y_0, \ell \mathbf{y}) \rangle_{\mathbf{\Psi}} \right|.$$

By RG–irrelevance, the windowed one–point scales as $\left| \langle S_k(\ell^z y_0, \ell \mathbf{y}) \rangle \right| \le C_k\, \ell^{\Delta_k}$ uniformly for y in the support of W_1 (after factoring the UV band $P_x(\lambda_{\mathrm{UV}})$, cf. Section 6.12.4), so

$$\int d^{d+1}x\, |W_\ell(x)| \left| \langle S_k^{(\lambda_{\mathrm{UV}})}(x) \rangle_{\mathbf{\Psi}} \right| \;\le\; C_k'\, \ell^{\,\Delta_k - (d+z)}. \tag{7.521}$$

Let $\Delta := \min_k \Delta_k$; then $\sum_k C'_k\, \ell^{\Delta_k-(d+z)} \le C'\, \ell^{\Delta-(d+z)}$.

Step 3: Conclusion for the deficit and plateau stability. Insert (7.521) into (7.520) and absorb the (slow) running of $\epsilon_k(\ell)$ inside the plateau window into the constant (by Proposition 7.359):

$$1 - |\Lambda_{\text{law}}^{(\lambda_{\text{UV}},\ell)}| \;\le\; C\,\ell^{\,\Delta-(d+z)} \;+\; \mathcal{O}(\varepsilon_{\text{soft}}^2),$$

which gives the stated scaling Proposition 6.277.

For the second claim, let $\mathcal{O}(\lambda_{\text{UV}}, \ell)$ be any ghost–number–zero diagnostic obeying the plateau criterion Section 6.12.4 on a box $\mathcal{B}$; by Proposition 7.359 its variations under $(\lambda_{\text{UV}}, \ell) \mapsto (\tilde{\lambda}_{\text{UV}}, \tilde{\ell}) \in \mathcal{B}$ are bounded by the logarithmic tolerances, so its extraction is regulator–stable within $\mathcal{B}$.

This completes the proof. □

Proposition 7.349 (Symmetric state, shifted law)**.** *Keep the symmetric backgrounds from Section 7.1.5 and expand the law EOM eqs. (6.333) and (6.334) to first order in ϵ_k. Then expectation values of G–odd order parameters remain zero up to $\mathcal{O}(\epsilon_k)$ (no state SSB), while the Hessian blocks receive $\mathcal{O}(\epsilon_k)$ shifts determined by the breaking operators' second variations. Physical poles/residues of the band–tagged propagators shift analytically in ϵ_k and remain gauge–parameter independent by cohomological identities (cf. Section 7.2.2).*

Proof. Work on the BRST–closed physical sector $\mathcal{H}_{\text{phys}}$ with the UV/IR tags $P_x(\lambda_{\text{UV}})$, $K_{\text{IR}}^{(\ell)}$ (ghost number 0; commuting with channel projectors; cf. Section 7.1.3). Let the band–tagged, gauge–fixed Lagrangian be

$$\mathcal{L}^{(\lambda_{\text{UV}},\ell)} = \mathcal{L}_0^{(\lambda_{\text{UV}},\ell)} + \sum_k \epsilon_k\, \mathcal{O}_k^{(\lambda_{\text{UV}})} + \zeta \Psi^{(\lambda_{\text{UV}},\ell)},$$

and recall the law EOM $\mathcal{E}_{\mathbf{\Phi}}^{(\lambda_{\text{UV}},\ell)} = 0$, $\mathcal{E}_{\mathbf{\Theta}}^{(\lambda_{\text{UV}},\ell)} = 0$ (no new labels; cf. eqs. (6.333) and (6.334)). Keep the symmetric backgrounds from Section 7.1.5, so at $\epsilon_k = 0$ one has $\mathcal{E}_{\mathbf{\Phi}}[\bar{\Phi}] = 0$, $\mathcal{E}_{\mathbf{\Theta}}[\bar{\Theta}] = 0$.

(i) G–odd order parameters remain zero up to $\mathcal{O}(\epsilon_k)$. Prepare a G–invariant, BRST–physical initial state (identity state branch; cf. Definition 7.369), so for any G–odd operator $\hat{\mathcal{O}}_{\text{odd}}$ one has $\langle \hat{\mathcal{O}}_{\text{odd}} \rangle_0 = 0$. Let the law micro–Hamiltonian be $\hat{H}_{\text{micro}} = \hat{H}_0 + \sum_k \epsilon_k \hat{\mathcal{O}}_k^{(\lambda_{\text{UV}},\ell)}$ (cf. Definition 7.350). Evolve to time T and expand by Dyson/Duhamel:

$$\big\langle \hat{\mathcal{O}}_{\text{odd}}(T) \big\rangle_{\text{micro}} = \langle \hat{\mathcal{O}}_{\text{odd}}(T) \rangle_0 + i \sum_k \epsilon_k \int_0^T ds\, \Big\langle \big[\hat{\mathcal{O}}_{\text{odd}}^{H_0}(T),\, \hat{\mathcal{O}}_k^{(\lambda_{\text{UV}},\ell),H_0}(s) \big] \Big\rangle_0 + \mathcal{O}(\epsilon^2).$$

The first term vanishes by G–invariance; the integral is bounded by $\sum_k |\epsilon_k|\, C_k\, T$ with C_k finite on a plateau window (boundedness in the $\mathcal{K}$–seminorm; cf. Definition 7.355 and Proposition 7.359). Hence $\langle \hat{\mathcal{O}}_{\text{odd}}(T) \rangle_{\text{micro}} = \mathcal{O}(\epsilon_k)$ (uniformly at fixed T and within the plateau), which excludes state SSB (an $\mathcal{O}(1)$ effect) under the QIII posture.

(ii) Hessian shifts are $\mathcal{O}(\epsilon_k)$ from second variations. Expand the law EOM functionals to first order in ϵ around $(\bar{\Phi}, \bar{\Theta})$:

$$0 = \mathcal{E}_{\mathbf{\Phi}}[\bar{\Phi}] + \sum_k \epsilon_k \left. \frac{\delta}{\delta \hat{\mathbf{\Phi}}} \int \mathcal{O}_k^{(\lambda_{\text{UV}})} \right|_{\bar{\Phi},\bar{\Theta}} + \text{ terms linear in fluctuations},$$

$$0 = \mathcal{E}_{\mathbf{\Theta}}[\bar{\Theta}] + \sum_k \epsilon_k \left. \frac{\delta}{\delta \hat{\mathbf{\Theta}}} \int \mathcal{O}_k^{(\lambda_{\text{UV}})} \right|_{\bar{\Phi},\bar{\Theta}} + \text{ terms linear in fluctuations}.$$

Since $\mathcal{E}_\bullet[\bar\bullet] = 0$, the *linearised* operators (quadratic kernel/Hessian acting on fluctuations $(\delta\hat{\mathbf{\Phi}}, \delta\hat{\mathbf{\Theta}})$) receive additive $\mathcal{O}(\epsilon_k)$ contributions from the *second* functional variations of the breaking insertions:

$$\mathbb{H}^{(\lambda_{\mathrm{UV}},\ell)}(\epsilon) = \mathbb{H}^{(\lambda_{\mathrm{UV}},\ell)}(0) + \sum_k \epsilon_k \begin{pmatrix} \dfrac{\delta^2 \mathcal{O}_k^{(\lambda_{\mathrm{UV}})}}{\delta\hat{\mathbf{\Phi}}\,\delta\hat{\mathbf{\Phi}}} & \dfrac{\delta^2 \mathcal{O}_k^{(\lambda_{\mathrm{UV}})}}{\delta\hat{\mathbf{\Phi}}\,\delta\hat{\mathbf{\Theta}}} \\ \dfrac{\delta^2 \mathcal{O}_k^{(\lambda_{\mathrm{UV}})}}{\delta\hat{\mathbf{\Theta}}\,\delta\hat{\mathbf{\Phi}}} & \dfrac{\delta^2 \mathcal{O}_k^{(\lambda_{\mathrm{UV}})}}{\delta\hat{\mathbf{\Theta}}\,\delta\hat{\mathbf{\Theta}}} \end{pmatrix}_{(\bar\Phi,\bar\Theta)}$$

(cf. Section 7.2.5).

Thus each kinetic/mass/mixing block shifts by $\mathcal{O}(\epsilon_k)$ determined by the breaking operators' second variations evaluated on the symmetric background.

(iii) Analytic, gauge–independent shifts of physical poles/residues. Restrict to $\mathcal{H}_{\mathrm{phys}}$ (quartet decoupling; cf. Section 7.2.1) and denote the physical quadratic kernel by $\mathbb{H}_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}(\epsilon)$. By admissibility (form smallness) and Proposition 7.357, the family $\epsilon \mapsto \mathbb{H}_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}(\epsilon)$ is a self–adjoint, analytic (type–A) perturbation of $\mathbb{H}_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}(0)$. Rellich–Kato theory then gives analytic dependence of isolated spectral data; in particular, the poles/residues of the band–tagged propagators $\mathcal{D}_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}(\epsilon) = \big(\mathbb{H}_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}(\epsilon)\big)^{-1}$ shift analytically in $\{\epsilon_k\}$. Gauge–parameter independence follows from BRST/Slavnov–Taylor consistency on the band (Proposition 7.358) and the cohomological/Nielsen identities used in Section 7.2.2 (physical, ghost–number–zero poles/residues are invariant under ζ–exact changes of the gauge fermion).

Combining (i)–(iii) proves the proposition. □

Quadrant III logic. We hold the *state* symmetric while allowing small, explicit deformations of the law. The scale–tagged Ward identity quantifies the resulting nonconservation by a controlled source proportional to the tiny couplings ϵ_k. Because tags are BRST–closed and all observables are taken in $\mathcal{H}_{\mathrm{phys}}$, ghosts and gauge artifacts do not pollute the readout; plateaus ensure UV/IR robustness of the inferred deficits and mass shifts. As ℓ grows, irrelevant breakings die off with the exponent $\Delta - (d+z)$, matching Proposition 6.277 and preserving Quadrant III membership.

Band-limited micro–Hamiltonian.

Definition 7.350 (QIII law-channel micro–Hamiltonian, band/IR tagged)**.** Let $\hat H_0(t;\lambda_{\mathrm{UV}},\ell)$ be the symmetric baseline Hamiltonian of eq. (7.241), obtained from the G–invariant quadratic kernel (cf. Proposition 7.2) and evaluated *on the band* with UV projector $P_x(\lambda_{\mathrm{UV}})$ and IR window $K_{\mathrm{IR}}^{(\ell)}$ (cf. Section 6.12.4 and Definition 6.282). Fix a family $\{\hat{\mathcal{O}}_B\}_B$ of local, Hermitian operators built from the field multiplet and covariant derivatives (cf. the band Hessian in eq. (7.10)). Their band/IR–tagged versions are

$$\hat{\mathcal{O}}_B^{(\lambda_{\mathrm{UV}},\ell)} := K_{\mathrm{IR}}^{(\ell)}\, P_x(\lambda_{\mathrm{UV}})\, \hat{\mathcal{O}}_B\, P_x(\lambda_{\mathrm{UV}})\, K_{\mathrm{IR}}^{(\ell)}$$

("UV then IR" order; cf. *eq.* (6.353)*and Section* 6.12.4).

For small, real couplings ε_B the *QIII micro–Hamiltonian in the law channel* is

$$\boxed{\hat{H}_{\mathrm{micro}}(t;\lambda_{\mathrm{UV}},\ell) = \hat{H}_0(t;\lambda_{\mathrm{UV}},\ell) + \sum_B \varepsilon_B\, \hat{\mathcal{O}}_B^{(\lambda_{\mathrm{UV}},\ell)}(t).} \tag{7.522}$$

Assumption 7.351 (Admissibility and symmetry posture)**.** *The breaking operators $\hat{\mathcal{O}}_B$ satisfy:*

(A1) BRST closure (on shell).
$\delta_{\mathrm{BRST}}\hat{\mathcal{O}}_B = \{\hat{Q}_{\mathrm{BRST}}, \hat{\mathcal{O}}_B\} = 0$ *and* $\mathrm{gh}(\hat{\mathcal{O}}_B) = 0$*, whereas they* break *the global symmetry G:* $\delta_G^{(\mathrm{global})}\hat{\mathcal{O}}_B \neq 0$*. This keeps Slavnov–Taylor identities exact while allowing global Ward sources (cf. eq. (7.16) and Section 6.12.2; [117, 243]).*

(A2) Locality and power counting.
$\hat{\mathcal{O}}_B$ *are local polynomials (or Wick polynomials) with canonical/anomalous dimensions compatible with the renormalisation scheme used (spurion tracking for G–charges as in [440]; see also Definition 7.346).*

(A3) Smallness (form bound).
$\hat{\mathcal{O}}_B^{(\lambda_{\mathrm{UV}},\ell)}$ *is $\mathcal{K}$–form–bounded w.r.t. $\hat{H}_0$ with* small *relative bound measured by the seminorm of eq. (7.526), i.e.* $\varepsilon_B\, \|\hat{\mathcal{O}}_B^{(\lambda_{\mathrm{UV}},\ell)}\|_{\mathcal{K}} \leq \varepsilon_{\mathrm{soft}} \ll 1$.

(A4) Band/IR compatibility.
$P_x(\lambda_{\mathrm{UV}})$ *and* $K_{\mathrm{IR}}^{(\ell)}$ *are BRST–closed (ghost number 0) and commute with channel projectors Π_α (cf. Section 7.1.3); thus tagging preserves domains and cohomology.*

Proposition 7.352 (Self–adjointness, BRST invariance, and physical unitarity)**.** *Under Assumption 7.351, $\hat{H}_{\mathrm{micro}}(t;\lambda_{\mathrm{UV}},\ell)$ is self–adjoint on the domain of $\hat{H}_0$ and generates a unitary evolution on the band–limited Hilbert space. Moreover,*

$$[\hat{Q}_{\mathrm{BRST}},\ \hat{H}_{\mathrm{micro}}] = 0,$$

so the evolution descends to $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ and preserves BRST cohomology. Consequently, physical matrix elements and spectra are gauge–parameter independent (Nielsen identities).

Proof idea. Kato–Rellich theory gives self–adjointness from (A3) (small form bound) on the common domain of $\hat{H}_0$ [115]. BRST closure (A1) and nilpotency imply $[\hat{Q}_{\mathrm{BRST}}, \hat{H}_0] = 0$ (since $[\hat{Q}_{\mathrm{BRST}}, \{\hat{Q}_{\mathrm{BRST}}, \Psi_\zeta\}] = 0$) and $[\hat{Q}_{\mathrm{BRST}}, \hat{\mathcal{O}}_B^{(\lambda_{\mathrm{UV}},\ell)}] = 0$, hence the commutator vanishes. Gauge–parameter independence on $\mathcal{H}_{\mathrm{phys}}$ follows from cohomological/Nielsen identities [117, 82].

Proof. (i) Self–adjointness and unitary evolution. Write

$$\hat{H}_{\mathrm{micro}}(t;\lambda_{\mathrm{UV}},\ell) = \hat{H}_0(t;\lambda_{\mathrm{UV}},\ell) + V(t), \qquad V(t) := \sum_B \varepsilon_B\, \hat{\mathcal{O}}_B^{(\lambda_{\mathrm{UV}},\ell)}(t),$$

with $\hat{H}_0$ the symmetric baseline of eq. (7.241). By admissibility (A3) in Definition 7.356, each tagged breaking $\hat{\mathcal{O}}_B^{(\lambda_{\mathrm{UV}},\ell)}$ is $\mathcal{K}$–form–bounded relative to $\hat{H}_0$ with small relative bound:

$$\left\|\hat{\mathcal{O}}_B^{(\lambda_{\mathrm{UV}},\ell)}\right\|_{\mathcal{K}} \leq \beta_B, \qquad \sum_B |\varepsilon_B|\, \beta_B \leq \varepsilon_{\mathrm{soft}} \ll 1.$$

Therefore $V(t)$ is a symmetric form perturbation of $\hat{H}_0$ with arbitrarily small relative bound on the common form domain $\mathfrak{h} = \overline{\mathrm{Dom}\,\mathcal{K}^{1/2}}$ (cf. Definition 7.355). By the Kato–Rellich theorem (form version), for each fixed t the operator $\hat{H}_{\mathrm{micro}}(t)$ is self–adjoint on $\mathrm{Dom}\,\hat{H}_0$ and bounded below; moreover, the family is stable under small variations of $(\varepsilon_B, \lambda_{\mathrm{UV}}, \ell)$ (see also Proposition 7.357). Standard Kato theory for time–dependent self–adjoint Hamiltonians with a *time–independent* form domain then yields a unique unitary propagator $U_{\mathrm{micro}}(t, s; \lambda_{\mathrm{UV}}, \ell)$ (the $\mathcal{T}$–exponential) on the band–limited Hilbert space.[9] Hence $\hat{H}_{\mathrm{micro}}$ generates a unitary evolution.

(ii) BRST invariance. By construction in the symmetric baseline and Section 7.1.3, $[\hat{Q}_{\mathrm{BRST}}, \hat{H}_0] = 0$ (the gauge–fixing piece is $\{\hat{Q}_{\mathrm{BRST}}, \Psi_\zeta\}$ and $[\hat{Q}_{\mathrm{BRST}}, \{\hat{Q}_{\mathrm{BRST}}, \Psi_\zeta\}] = 0$ by nilpotency and the graded Jacobi identity). For the explicit breakings, admissibility gives $\delta_{\mathrm{BRST}}\hat{\mathcal{O}}_B = 0$ and the tags are BRST–closed: $[\hat{Q}_{\mathrm{BRST}}, P_x(\lambda_{\mathrm{UV}})] = [\hat{Q}_{\mathrm{BRST}}, K_{\mathrm{IR}}^{(\ell)}] = 0$ (cf. eq. (7.244)). Therefore

$$[\hat{Q}_{\mathrm{BRST}}, \hat{\mathcal{O}}_B^{(\lambda_{\mathrm{UV}},\ell)}] = \left[\hat{Q}_{\mathrm{BRST}}, K_{\mathrm{IR}}^{(\ell)} P_x(\lambda_{\mathrm{UV}}) \hat{\mathcal{O}}_B P_x(\lambda_{\mathrm{UV}}) K_{\mathrm{IR}}^{(\ell)}\right] = 0,$$

and by linearity $[\hat{Q}_{\mathrm{BRST}}, V(t)] = 0$. Consequently $[\hat{Q}_{\mathrm{BRST}}, \hat{H}_{\mathrm{micro}}] = 0$.

(iii) Descent to the physical sector and gauge parameter independence. Since $[\hat{Q}_{\mathrm{BRST}}, \hat{H}_{\mathrm{micro}}] = 0$, the propagator intertwines with $\hat{Q}_{\mathrm{BRST}}$: $\hat{Q}_{\mathrm{BRST}}\, U_{\mathrm{micro}}(t, s) = U_{\mathrm{micro}}(t, s)\, \hat{Q}_{\mathrm{BRST}}$. Thus U_{micro} maps $\ker \hat{Q}_{\mathrm{BRST}}$ into itself and $\mathrm{im}\,\hat{Q}_{\mathrm{BRST}}$ into itself; it descends to a unitary evolution on $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$. Variations of gauge parameters change the gauge fermion by $\Psi_\zeta \to \Psi_\zeta + \delta\Psi$ and induce a BRST-exact deformation $\delta\hat{H} = \{\hat{Q}_{\mathrm{BRST}}, \delta\Psi\}$; by the same argument as above, the induced variation of any ghost-number-zero physical matrix element is a BRST coboundary and therefore vanishes on $\mathcal{H}_{\mathrm{phys}}$ (Nielsen-type identities; cf. Section 7.1.3). Hence physical spectra and matrix elements are gauge-parameter independent. □

Proposition 7.353 (Ward structure with explicit breakings (scale–tagged, physical sector))**.** *Let δ_a be a (possibly local/relational) symmetry variation with $\delta_a\mathcal{L}_0 = \nabla_\mu K_a^\mu$ and $\delta_a\hat{\mathcal{O}}_B = \mathcal{S}_{B,a}$ (band–tagged). Then the adjoint–covariant, scale–tagged Ward identity reads*

$$\nabla_\mu^{(\mathrm{ad})} J_a^{(\lambda_{\mathrm{UV}},\ell)\mu} = \sum_B \varepsilon_B \left\langle \mathcal{S}_{B,a}^{(\lambda_{\mathrm{UV}})} \right\rangle_{\boldsymbol{\Psi}} + \left\langle \zeta\, \eta_a^{(\lambda_{\mathrm{UV}},\ell)} \right\rangle_{\boldsymbol{\Psi}} + \mathcal{A}_a^{(\lambda_{\mathrm{UV}},\ell)},$$

as in Definition 6.295. On the physical sector and with anomaly-free BRST, ζ–exact and anomalous terms drop, yielding

$$\nabla_\mu^{(\mathrm{ad})} \bar{J}_{a,\mathrm{imp}}^{(\lambda_{\mathrm{UV}},\ell)\mu} = \sum_B \varepsilon_B \left\langle \mathcal{S}_{B,a}^{(\lambda_{\mathrm{UV}})} \right\rangle_{\boldsymbol{\Psi}},$$

in agreement with eq. (6.377) and the relational Ward lemma Lemma 6.233 (cf. [117, 230]).

Proof. Work with the scale–tagged Lagrangian $\mathcal{L} = \mathcal{L}_0 + \sum_B \varepsilon_B \mathcal{O}_B + \zeta\Psi$ (UV/IR tags suppressed; cf. Definition 6.295). Let δ_a be the (possibly local/relational) variation with

[9] We use that $t \mapsto V(t)$ is strongly measurable and locally bounded as a quadratic form, which holds for local Wick polynomials with band/IR tags; cf. the assumptions preceding Definition 7.356.

$\delta_a \mathcal{L}_0 = \nabla_\mu K_a^\mu$ and $\delta_a \mathcal{O}_B = \mathcal{S}_{B,a}$. By the relational Ward lemma Lemma 6.233 (applied to the present decomposition), there exists a local functional η_a from $\delta_a \Psi$ such that, after the usual Noether–Belinfante improvement absorbing $\nabla_\mu K_a^\mu$, the (improved) current obeys the local off–shell identity

$$\nabla_\mu J^\mu_{a,\mathrm{imp}} = \sum_B \varepsilon_B \, \mathcal{S}_{B,a} \; + \; \zeta \, \eta_a \; + \; \mathcal{A}_a, \tag{7.523}$$

with a possible anomaly $\mathcal{A}_a$ (vanishing in the anomaly–free case), cf. Definition 6.295 and [117, 230].

Adjoint covariantization. If G is gauged, replace ∇_μ by the adjoint–covariant divergence $\nabla^{(\mathrm{ad})}_\mu J^\mu := \nabla_\mu J^\mu + i[\mathcal{A}_\mu, J^\mu]$; this yields exactly the definition in Definition 6.295:

$$\nabla^{(\mathrm{ad})}_\mu J^\mu_{a,\mathrm{imp}} = \sum_B \varepsilon_B \, \mathcal{S}_{B,a} \; + \; \zeta \, \eta_a \; + \; \mathcal{A}_a. \tag{7.524}$$

Scale tagging. Apply the band/IR tagging functor (UV then IR; cf. Definition 6.282 and Section 6.12.4). Using $[\nabla_\mu, P_x(\lambda_{\mathrm{UV}})] = 0$ (assumption in Section 6.12.4) and the fact that $P_x(\lambda_{\mathrm{UV}})$, $K^{(\ell)}_{\mathrm{IR}}$ are BRST–closed of ghost number 0 and commute with channel projectors (cf. Section 7.1.3 and eq. (7.244)), we obtain the scale–tagged identity

$$\nabla^{(\mathrm{ad})}_\mu J^{(\lambda_{\mathrm{UV}},\ell)\mu}_{a,\mathrm{imp}} = \sum_B \varepsilon_B \, \mathcal{S}^{(\lambda_{\mathrm{UV}})}_{B,a} \; + \; \zeta \, \eta^{(\lambda_{\mathrm{UV}},\ell)}_a \; + \; \mathcal{A}^{(\lambda_{\mathrm{UV}},\ell)}_a, \tag{7.525}$$

which is precisely the statement in the first display of the proposition (compare with Definition 6.295 and eq. (6.376)).

Physical reduction. Taking expectation values in a homogeneous state $|\mathbf{\Psi}\rangle$ and projecting to the physical sector with $\mathsf{P}_{\mathrm{phys}}$ gives

$$\nabla^{(\mathrm{ad})}_\mu \bar{J}^{(\lambda_{\mathrm{UV}},\ell)\mu}_{a,\mathrm{imp}} = \sum_B \varepsilon_B \left\langle \mathcal{S}^{(\lambda_{\mathrm{UV}})}_{B,a} \right\rangle_{\mathbf{\Psi}} \; + \; \left\langle \zeta \, \eta^{(\lambda_{\mathrm{UV}},\ell)}_a \right\rangle_{\mathbf{\Psi}} \; + \; \mathcal{A}^{(\lambda_{\mathrm{UV}},\ell)}_a,$$

where $\bar{J} := \langle \hat{J} \rangle_{\mathbf{\Psi}}$ and we used that tags commute with $\mathsf{P}_{\mathrm{phys}}$ (cf. Section 7.1.3). On the anomaly–free Stage I and within BRST cohomology (ghost number 0), ζ–exact insertions vanish and the anomaly term is absent; hence

$$\nabla^{(\mathrm{ad})}_\mu \bar{J}^{(\lambda_{\mathrm{UV}},\ell)\mu}_{a,\mathrm{imp}} = \sum_B \varepsilon_B \left\langle \mathcal{S}^{(\lambda_{\mathrm{UV}})}_{B,a} \right\rangle_{\mathbf{\Psi}},$$

which is eq. (6.377) and agrees with the relational Ward identity in Lemma 6.233. This completes the proof. □

Remark 7.354 (Law–channel purity and branch equality)**.** In Quadrant III we keep the *state* channel symmetric while allowing explicit law deformations. The branch generators use the *same* G–invariant baseline $\hat{H}_0$ and differ only by the small, tagged $\varepsilon_B \, \hat{\mathcal{O}}_B^{(\lambda_{\mathrm{UV}},\ell)}$ insertions as in (7.522). Any change of presentation by gauge–parameter scans or ζ–exact decorations is cohomologically trivial on $\mathcal{H}_{\mathrm{phys}}$ (cf. Section 7.1.3).

Smallness and admissibility.

Definition 7.355 ($\mathcal{K}$–weighted seminorm and form domain)**.** Let $\mathcal{K}$ be the positive, self–adjoint kinetic operator appearing in the symmetric baseline Hamiltonian of eq. (7.241); write $\mathfrak{h} := \overline{\mathrm{Dom}\,\mathcal{K}^{1/2}}$ for its form domain endowed with the graph norm $\|\Psi\|_{\mathcal{K}} := \|\mathcal{K}^{1/2}\Psi\|$. For any band/IR–tagged operator $\hat{\mathcal{O}}^{(\lambda_{\mathrm{UV}},\ell)}$ define the $\mathcal{K}$–weighted operator seminorm

$$\boxed{\left\|\hat{\mathcal{O}}^{(\lambda_{\mathrm{UV}},\ell)}\right\|_{\mathcal{K}} := \sup_{\|\Psi\|_{\mathcal{K}}=1} \left|\left\langle \Psi \middle| \mathcal{K}^{-1/2}\,\hat{\mathcal{O}}^{(\lambda_{\mathrm{UV}},\ell)}\,\mathcal{K}^{-1/2} \middle| \Psi \right\rangle\right|.} \tag{7.526}$$

We say that a tagged operator is *$\mathcal{K}$–form–bounded with (relative) bound β* if $\left\|\hat{\mathcal{O}}^{(\lambda_{\mathrm{UV}},\ell)}\right\|_{\mathcal{K}} \leq \beta < \infty$.

Definition 7.356 (Admissible explicit breaking in QIII)**.** A family $\{\hat{\mathcal{O}}_B\}_B$ of symmetry–breaking operators is *admissible* if:

(A1) **BRST closure (on shell) and ghost number** 0**:** $\delta_{\mathrm{BRST}}\hat{\mathcal{O}}_B = \{\hat{Q}_{\mathrm{BRST}}, \hat{\mathcal{O}}_B\} = 0$ and $\mathrm{gh}(\hat{\mathcal{O}}_B) = 0$, but $\delta_G^{(\mathrm{global})}\hat{\mathcal{O}}_B \neq 0$ (so global G is explicitly broken in the law channel while Slavnov–Taylor remains exact on shell), cf. eq. (7.16) and [117, 442, 243].

(A2) **Locality and power counting:** $\hat{\mathcal{O}}_B$ is a local (Wick) polynomial in the band/IR-tagged fields and covariant derivatives, with canonical/anomalous dimensions tracked as in eq. (7.211); broken G-charges are monitored by a spurion S_B such that $\varepsilon_B S_B \hat{\mathcal{O}}_B$ is G-invariant [440].

(A3) **Small relative form bound:** for the tagged operator, defined as

$$\hat{\mathcal{O}}_B^{(\lambda_{\mathrm{UV}},\ell)} := K_{\mathrm{IR}}^{(\ell)} P_x(\lambda_{\mathrm{UV}}) \hat{\mathcal{O}}_B P_x(\lambda_{\mathrm{UV}}) K_{\mathrm{IR}}^{(\ell)},$$

one has

$$\varepsilon_B \left\|\hat{\mathcal{O}}_B^{(\lambda_{\mathrm{UV}},\ell)}\right\|_{\mathcal{K}} \leq \varepsilon_{\mathrm{soft}} \ll 1 \quad \text{(smallness on the band and within the IR window).}$$

(A4) **Pipeline stability:** under the regularisation pipeline $(\lambda_{\mathrm{UV}}, \ell) \mapsto (\tilde{\lambda}_{\mathrm{UV}}, \tilde{\ell})$ (cf. eq. (7.67)), the seminorm varies smoothly and obeys plateau–type bounds of Section 6.12.4.

(A5) **Band/IR compatibility:** $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ are BRST–closed (ghost number 0) and commute with the channel projectors Π_α, so tagging preserves domains, cohomology, and channel resolution (cf. Section 7.1.3).

Proposition 7.357 (Kato–Rellich stability and unitary dynamics)**.** *Let $\hat{H}_{\mathrm{micro}}$ be as in Definition 7.350 with admissible $\{\hat{\mathcal{O}}_B\}_B$. Then $\hat{H}_{\mathrm{micro}}(t; \lambda_{\mathrm{UV}}, \ell)$ is self–adjoint on the form domain of $\hat{H}_0(t; \lambda_{\mathrm{UV}}, \ell)$ and generates a unitary evolution on the band–limited Hilbert space. Moreover, for $t \geq 0$,*

$$\left\|U_{\mathrm{micro}}(t; \lambda_{\mathrm{UV}}, \ell) - U_0(t; \lambda_{\mathrm{UV}}, \ell)\right\| \leq C\,\varepsilon_{\mathrm{soft}}\,t + \mathcal{O}(\varepsilon_{\mathrm{soft}}^2\,t^2),$$

with a constant C depending on $\mathcal{K}$ but independent of $(\lambda_{\mathrm{UV}}, \ell)$ inside a plateau box. Sketch. *By (A3) each $\varepsilon_B \hat{\mathcal{O}}_B^{(\lambda_{\mathrm{UV}},\ell)}$ is $\mathcal{K}$–form–small relative to $\hat{H}_0$, hence $\hat{H}_0 + \sum_B \varepsilon_B \hat{\mathcal{O}}_B^{(\lambda_{\mathrm{UV}},\ell)}$ is self–adjoint and lower bounded (Kato–Rellich) and yields a unitary group; the difference bound follows from a first–order Dyson estimate [115].*

Proof. Write

$$\hat{H}_{\mathrm{micro}}(t;\lambda_{\mathrm{UV}},\ell) = \hat{H}_0(t;\lambda_{\mathrm{UV}},\ell) + V(t), \qquad V(t) := \sum_B \varepsilon_B\, \hat{\mathcal{O}}_B^{(\lambda_{\mathrm{UV}},\ell)}(t),$$

with $\hat{H}_0$ the symmetric baseline of eq. (7.241). By admissibility (A3) in Definition 7.356 and the $\mathcal{K}$–seminorm of Definition 7.355–(7.526), each tagged breaking is $\mathcal{K}$–form–bounded relative to $\hat{H}_0$ with small bound:

$$\left\|\hat{\mathcal{O}}_B^{(\lambda_{\mathrm{UV}},\ell)}(t)\right\|_{\mathcal{K}} \le \beta_B, \qquad \sum_B |\varepsilon_B|\,\beta_B \le \varepsilon_{\mathrm{soft}} \ll 1,$$

uniformly for $(\lambda_{\mathrm{UV}},\ell)$ inside a plateau window (cf. Proposition 7.359 and Section 6.12.4).

(i) Self–adjointness. For each fixed t, $V(t)$ is a symmetric form perturbation of $\hat{H}_0(t)$ with relative bound < 1 on the common form domain $\mathfrak{h} = \overline{\mathrm{Dom}\,\mathcal{K}^{1/2}}$. By the Kato–Rellich theorem (form version), $\hat{H}_{\mathrm{micro}}(t) = \hat{H}_0(t) + V(t)$ is self–adjoint on $\mathrm{Dom}\,\hat{H}_0(t)$ and bounded below [115].

(ii) Unitary dynamics. The quadratic forms $q_t[\Psi] := \langle \mathcal{K}^{1/2}\Psi, \mathcal{K}^{1/2}\Psi\rangle + \langle \Psi, V(t)\Psi\rangle$ are defined on the time–independent form domain $\mathfrak{h}$, are closed, and depend (locally) boundedly/measurably on t for local Wick polynomials with band/IR tags (same regularity as used in Definition 7.356). Kato's theory for time–dependent self–adjoint Hamiltonians with fixed form domain then yields a unique unitary propagator $U_{\mathrm{micro}}(t,s;\lambda_{\mathrm{UV}},\ell)$ solving $i\partial_t U_{\mathrm{micro}}(t,s) = \hat{H}_{\mathrm{micro}}(t) U_{\mathrm{micro}}(t,s)$ with $U_{\mathrm{micro}}(s,s) = \mathbf{1}$ [115].

(iii) Difference bound. Duhamel's formula gives

$$U_{\mathrm{micro}}(t) - U_0(t) = -i \int_0^t U_{\mathrm{micro}}(t,s)\, V(s)\, U_0(s)\, ds.$$

Taking operator norms and using unitarity,

$$\|U_{\mathrm{micro}}(t) - U_0(t)\| \;\le\; \int_0^t \|V(s)\|_{\mathrm{op}}\, ds.$$

By the $\mathcal{K}$–form bound and equivalence of the induced operator norms on $\mathfrak{h}$ (cf. Remark 7.361), there exists a constant $C = C(\mathcal{K})$ such that $\|V(s)\|_{\mathrm{op}} \le C \sum_B |\varepsilon_B| \left\|\hat{\mathcal{O}}_B^{(\lambda_{\mathrm{UV}},\ell)}(s)\right\|_{\mathcal{K}} \le C\,\varepsilon_{\mathrm{soft}}$. Hence

$$\|U_{\mathrm{micro}}(t) - U_0(t)\| \le C\,\varepsilon_{\mathrm{soft}}\, t,$$

and the standard Dyson expansion yields the second–order remainder $\mathcal{O}(\varepsilon_{\mathrm{soft}}^2 t^2)$. Inside a plateau box, C can be chosen independent of $(\lambda_{\mathrm{UV}},\ell)$ by Proposition 7.359.

This proves self–adjointness, unitary evolution, and the stated bound. □

Proposition 7.358 (BRST/Slavnov–Taylor consistency on the band)**.** *Under (A1)–(A5),*

$$[\hat{Q}_{\mathrm{BRST}},\, \hat{H}_{\mathrm{micro}}] = 0 \quad \textit{and} \quad \mathcal{S}(\Gamma) = 0 \quad \textit{(ST identity)}$$

hold on the band and within the IR window; here $\mathcal{S}(\Gamma)$ denotes the Slavnov functional acting on the effective action. Consequently, physical matrix elements, spectra, and channel–resolved visibilities evaluated on $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ are gauge–parameter independent (Nielsen identities) and unaffected by ζ–exact decorations, cf. Section 7.1.3. Justification. *BRST closure and ghost number 0 prevent cohomology obstructions; band/IR tags are BRST–closed and commute with Π_α, so the standard algebra carries through [117, 442, 243, 82].*

Proof. (i) Hamiltonian–level BRST invariance on the band. From Definition 7.350 write

$$\hat{H}_{\mathrm{micro}}(t;\lambda_{\mathrm{UV}},\ell) = \hat{H}_0(t;\lambda_{\mathrm{UV}},\ell) + \sum_B \varepsilon_B\, \hat{\mathcal{O}}_B^{(\lambda_{\mathrm{UV}},\ell)}(t),$$
$$\hat{\mathcal{O}}_B^{(\lambda_{\mathrm{UV}},\ell)} := K_{\mathrm{IR}}^{(\ell)} P_x(\lambda_{\mathrm{UV}})\, \hat{\mathcal{O}}_B\, P_x(\lambda_{\mathrm{UV}}) K_{\mathrm{IR}}^{(\ell)}.$$

By construction of the symmetric baseline and Section 7.1.3,

$$[\hat{Q}_{\mathrm{BRST}}, \hat{H}_0] = 0, \qquad [\hat{Q}_{\mathrm{BRST}}, \{\hat{Q}_{\mathrm{BRST}}, \Psi_\zeta\}] = 0$$

(the latter by nilpotency and graded Jacobi). By admissibility (A1)–(A5) of Definition 7.356 we have (on shell) $\delta_{\mathrm{BRST}}\hat{\mathcal{O}}_B = 0$ and the tags are BRST–closed of ghost number 0:

$$[\hat{Q}_{\mathrm{BRST}}, \hat{\mathcal{O}}_B] = 0, \qquad [\hat{Q}_{\mathrm{BRST}}, P_x(\lambda_{\mathrm{UV}})] = [\hat{Q}_{\mathrm{BRST}}, K_{\mathrm{IR}}^{(\ell)}] = 0 \quad \text{(cf. eq. (7.244))}.$$

Therefore

$$[\hat{Q}_{\mathrm{BRST}}, \hat{\mathcal{O}}_B^{(\lambda_{\mathrm{UV}},\ell)}] = \left[\hat{Q}_{\mathrm{BRST}}, K_{\mathrm{IR}}^{(\ell)} P_x\, \hat{\mathcal{O}}_B\, P_x K_{\mathrm{IR}}^{(\ell)}\right] = 0,$$

and by linearity $[\hat{Q}_{\mathrm{BRST}}, \hat{H}_{\mathrm{micro}}] = 0$.

(ii) Slavnov–Taylor identity for the effective action on the band. Consider the scale–tagged gauge–fixed action $S^{(\lambda_{\mathrm{UV}},\ell)} = S_0^{(\lambda_{\mathrm{UV}},\ell)} + \sum_B \varepsilon_B \int \mathcal{O}_B^{(\lambda_{\mathrm{UV}})} + \int \zeta \Psi^{(\lambda_{\mathrm{UV}},\ell)}$, with antifield (Zinn–Justin) sources coupled to BRST variations in the usual way [117, 442, 243]. The functional measure is BRST invariant and the classical action is BRST invariant modulo ζ–exact terms; admissibility (A1) supplies $\delta_{\mathrm{BRST}}\mathcal{O}_B = 0$ and (A5) ensures that the linear tagging maps commute with BRST. Hence the standard derivation of the ST identity goes through *verbatim* for the generating functional $Z^{(\lambda_{\mathrm{UV}},\ell)}$, yielding the ST identity for the effective action $\Gamma^{(\lambda_{\mathrm{UV}},\ell)}$:

$$\mathcal{S}\big(\Gamma^{(\lambda_{\mathrm{UV}},\ell)}\big) = 0$$

in the anomaly–free case (Stage I), see e.g. [82, Ch. 16], [117, 442, 243]. The presence of $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ only composes the fields with BRST–closed linear operators and thus does not alter the cohomological content entering the ST operator.

(iii) Descent to $\mathcal{H}_{\mathrm{phys}}$ and Nielsen independence. From (i) the propagator U_{micro} intertwines with $\hat{Q}_{\mathrm{BRST}}$, so it preserves $\ker \hat{Q}_{\mathrm{BRST}}$ and $\mathrm{im}\, \hat{Q}_{\mathrm{BRST}}$ and descends to a unitary evolution on the ghost–number–zero cohomology $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$. Gauge–parameter variations amount to a deformation of the gauge fermion, $\Psi \mapsto \Psi + \delta\Psi$, and hence to a BRST–exact deformation of the action/Hamiltonian, $\delta S = \int \zeta\, \delta\Psi$, $\delta\hat{H} = \{\hat{Q}_{\mathrm{BRST}}, \delta\Psi\}$. The ST identity together with BRST triviality implies that the variation of any ghost–number–zero, BRST–closed observable reduces to a BRST coboundary; its expectation value vanishes upon projection to $\mathcal{H}_{\mathrm{phys}}$ (Nielsen identities), cf. Section 7.1.3 and [117, 82]. Because the tags commute with channel projectors Π_α (A5), the same conclusion holds for channel–resolved visibilities.

Combining (i)–(iii) proves $[\hat{Q}_{\mathrm{BRST}}, \hat{H}_{\mathrm{micro}}] = 0$, $\mathcal{S}(\Gamma) = 0$ on the band/within the IR window, and gauge–parameter independence of physical matrix elements, spectra, and channel–resolved visibilities. □

Proposition 7.359 (Pipeline/plateau robustness of the seminorm)**.** *If* $\mathcal{O}_B(\lambda_{\mathrm{UV}},\ell) := \big\|\hat{\mathcal{O}}_B^{(\lambda_{\mathrm{UV}},\ell)}\big\|_{\mathcal{K}}$ *satisfies the plateau criterion Section 6.12.4 on* $[\lambda_{\mathrm{UV},1},\lambda_{\mathrm{UV},2}]\times[\ell_1,\ell_2]$*, then for any* $(\lambda_{\mathrm{UV}},\ell)$*,* $(\tilde\lambda_{\mathrm{UV}},\tilde\ell)$ *in this box,*

$$\big|\mathcal{O}_B(\tilde\lambda_{\mathrm{UV}},\tilde\ell)-\mathcal{O}_B(\lambda_{\mathrm{UV}},\ell)\big| \;\le\; \varepsilon_{\lambda_{\mathrm{UV}}}\Big|\ln\tfrac{\tilde\lambda_{\mathrm{UV}}}{\lambda_{\mathrm{UV}}}\Big| \;+\; \varepsilon_\ell\Big|\ln\tfrac{\tilde\ell}{\ell}\Big|.$$

Hence the smallness condition in (7.526) is stable under UV/IR scans within the plateau window.

Proof. Set

$$u:=\ln\lambda_{\mathrm{UV}},\qquad v:=\ln\ell,\qquad F(u,v):=\mathcal{O}_B\big(e^u,e^v\big)=\Big\|\hat{\mathcal{O}}_B^{(e^u,e^v)}\Big\|_{\mathcal{K}}.$$

By the plateau criterion Section 6.12.4 (cf. (6.380)), there exist constants $\varepsilon_{\lambda_{\mathrm{UV}}},\varepsilon_\ell$ such that, *uniformly* on the rectangle $[u_1,u_2]\times[v_1,v_2]=[\ln\lambda_{\mathrm{UV},1},\ln\lambda_{\mathrm{UV},2}]\times[\ln\ell_1,\ln\ell_2]$,

$$\big|\partial_u F(u,v)\big|\le\varepsilon_{\lambda_{\mathrm{UV}}},\qquad \big|\partial_v F(u,v)\big|\le\varepsilon_\ell.$$

Let (u,v) and $(\tilde u,\tilde v)$ be any two points in this rectangle (equivalently, $(\lambda_{\mathrm{UV}},\ell)$ and $(\tilde\lambda_{\mathrm{UV}},\tilde\ell)$ in the stated box). Apply the fundamental theorem of calculus along the piecewise C^1 path that first varies u at fixed v, then varies v at fixed $\tilde u$:

$$\begin{aligned}\big|F(\tilde u,\tilde v)-F(u,v)\big| &\le \Big|\int_u^{\tilde u}\partial_u F(s,v)\,ds\Big|+\Big|\int_v^{\tilde v}\partial_v F(\tilde u,t)\,dt\Big|\\ &\le \varepsilon_{\lambda_{\mathrm{UV}}}\,|\,\tilde u-u\,|+\varepsilon_\ell\,|\,\tilde v-v\,|.\end{aligned}$$

Returning to $(\lambda_{\mathrm{UV}},\ell)$ variables gives $|\,\tilde u-u\,|=|\ln\big(\tilde\lambda_{\mathrm{UV}}/\lambda_{\mathrm{UV}}\big)|$ and $|\,\tilde v-v\,|=|\ln\big(\tilde\ell/\ell\big)|$, hence

$$\big|\mathcal{O}_B(\tilde\lambda_{\mathrm{UV}},\tilde\ell)-\mathcal{O}_B(\lambda_{\mathrm{UV}},\ell)\big| \;\le\; \varepsilon_{\lambda_{\mathrm{UV}}}\Big|\ln\tfrac{\tilde\lambda_{\mathrm{UV}}}{\lambda_{\mathrm{UV}}}\Big| \;+\; \varepsilon_\ell\Big|\ln\tfrac{\tilde\ell}{\ell}\Big|.$$

This proves the displayed inequality. In particular, if the smallness condition $\varepsilon_B\,\mathcal{O}_B(\lambda_{\mathrm{UV}},\ell)\le\varepsilon_{\mathrm{soft}}$ holds at one point with a slack exceeding the right–hand side bound, then it holds throughout the plateau box; hence the smallness requirement in (7.526) is stable under UV/IR scans within that window. □

Corollary 7.360 (Controlled law deficit and spectral shifts)**.** *With admissible breakings and within a plateau window, the law–channel visibility deficit obeys*

$$1-|\Lambda_{\mathrm{law}}^{(\lambda_{\mathrm{UV}},\ell)}| = \mathcal{O}(\varepsilon_{\mathrm{soft}}) \;+\; \mathcal{O}\big(\ell^{\,\Delta-(d+z)}\big),$$

consistent with Proposition 6.277, and the physical pole locations/residues of the band–tagged propagators shift analytically in $\{\varepsilon_B\}$ *while remaining gauge–parameter independent (cf. Proposition 7.16).*

Proof. Law–visibility bound. Let Λ_{law} be defined as in eq. (7.122), with phases calibrated so that in the unbroken baseline ($\varepsilon_B = 0$) one has $\Lambda_{\mathrm{law}} = 1$ (cf. Proposition 7.367). Using Duhamel's formula with $\hat{H}_{\mathrm{micro}} = \hat{H}_0 + \sum_B \varepsilon_B \hat{\mathcal{O}}_B^{(\lambda_{\mathrm{UV}},\ell)}$,

$$U_{\mathrm{micro}}(T) - U_0(T) = -i \int_0^T U_{\mathrm{micro}}(T,t) \left(\sum_B \varepsilon_B \hat{\mathcal{O}}_B^{(\lambda_{\mathrm{UV}},\ell)}(t) \right) U_0(t)\, dt,$$

and the unitarity of U_{micro}, U_0 together with the $\mathcal{K}$–form bound (A3), we obtain the operator–norm estimate

$$\| U_{\mathrm{micro}}(T) - U_0(T) \| \le C\, \varepsilon_{\mathrm{soft}}\, T + \mathcal{O}(\varepsilon_{\mathrm{soft}}^2 T^2) \quad \text{(cf. Proposition 7.357)}.$$

Since the law visibility is a normalised trace/overlap of the product of a branch unitary and its reference (cf. Definition 7.363 and Proposition 7.365), the above yields

$$1 - |\Lambda_{\mathrm{law}}^{(\lambda_{\mathrm{UV}},\ell)}| \;=\; \mathcal{O}(\varepsilon_{\mathrm{soft}}) \;+\; R_{\mathrm{IR}}(\ell),$$

for some nonnegative remainder $R_{\mathrm{IR}}(\ell)$. By the scale–tagged Ward identity on the physical sector (with anomaly–free BRST),

$$\nabla_\mu^{(\mathrm{ad})} \bar{J}_{a,\mathrm{imp}}^{(\lambda_{\mathrm{UV}},\ell)\mu} = \sum_B \varepsilon_B \left\langle \mathcal{S}_{B,a}^{(\lambda_{\mathrm{UV}})} \right\rangle_{\boldsymbol{\Psi}} \quad \text{(Proposition 7.353, eq. (6.377))},$$

and the locality/clustering hypotheses with IR windowing, the integrated source scales as $\ell^{\Delta-(d+z)}$ when the breakings are RG–irrelevant with minimal scaling dimension Δ; hence $R_{\mathrm{IR}}(\ell) = \mathcal{O}\big(\ell^{\Delta-(d+z)}\big)$, reproducing Proposition 6.277 (see Proposition 7.348). Together this gives the stated deficit bound.

Spectral (pole/residue) shifts and gauge independence. On $\mathcal{H}_{\mathrm{phys}}$ (ghost number 0) the band–tagged quadratic kernel reads

$$\mathbb{H}_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}(\varepsilon) = \mathbb{H}_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}(0) \;+\; \sum_B \varepsilon_B\, \mathbb{V}_{B,\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)},$$

where $\mathbb{H}_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}(0)$ has positive kinetic blocks and $\mathbb{V}_{B,\mathrm{phys}}$ are bounded (by (A3) and Definition 7.355). For $\| \sum_B \varepsilon_B \mathbb{V}_{B,\mathrm{phys}} \|$ small (uniformly on a plateau window, cf. Proposition 7.359), the resolvent and hence the band–tagged propagator

$$\mathcal{D}_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}(\varepsilon) = \Big(\mathbb{H}_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}(\varepsilon) \Big)^{-1}$$

admit a convergent Neumann/Kato series in $\{\varepsilon_B\}$, implying analytic dependence of isolated poles and residues on $\{\varepsilon_B\}$ (Rellich/Kato analytic perturbation theory [115]). By Proposition 7.358 the BRST/ST identities hold on the band, so physical (ghost–number–zero, BRST–closed) poles/residues are gauge–parameter independent (Nielsen identities; cf. Section 7.1.3 and Proposition 7.16).

Combining these establishes the claimed visibility deficit and analytic, gauge–independent spectral shifts within the plateau window. □

Remark 7.361 (On the choice of $\mathcal{K}$ and equivalence of seminorms)**.** Let $\mathcal{K}_1, \mathcal{K}_2$ be positive, self–adjoint (elliptic) kinetic kernels with common form domain $\mathfrak{h} = \overline{\mathrm{Dom}\,\mathcal{K}_j^{1/2}}$ and suppose there exist $0 < m \le M < \infty$ such that, for all $\Psi \in \mathfrak{h}$,

$$m\,\langle \Psi, \mathcal{K}_1 \Psi\rangle \;\le\; \langle \Psi, \mathcal{K}_2 \Psi\rangle \;\le\; M\,\langle \Psi, \mathcal{K}_1 \Psi\rangle.$$

(Within a plateau box these bounds hold with m, M independent of $(\lambda_{\mathrm{UV}}, \ell)$.) Then the graph norms $\|\Psi\|_{\mathcal{K}_j} = \|\mathcal{K}_j^{1/2}\Psi\|$ are equivalent, and the $\mathcal{K}$–seminorms of (7.526) satisfy, for any tagged operator $\hat{\mathcal{O}}^{(\lambda_{\mathrm{UV}},\ell)}$,

$$m\,\big\|\hat{\mathcal{O}}^{(\lambda_{\mathrm{UV}},\ell)}\big\|_{\mathcal{K}_2} \;\le\; \big\|\hat{\mathcal{O}}^{(\lambda_{\mathrm{UV}},\ell)}\big\|_{\mathcal{K}_1} \;\le\; M\,\big\|\hat{\mathcal{O}}^{(\lambda_{\mathrm{UV}},\ell)}\big\|_{\mathcal{K}_2}.$$

Hence the notion of "small" in (7.526) is insensitive to benign reparametrisations of the quadratic baseline (uniformly on the plateau). *Justification.* The quadratic–form bounds imply $m^{1/2}\|\cdot\|_{\mathcal{K}_1} \le \|\cdot\|_{\mathcal{K}_2} \le M^{1/2}\|\cdot\|_{\mathcal{K}_1}$ and yield a bounded, coercive intertwiner $R := \mathcal{K}_2^{1/2}\mathcal{K}_1^{-1/2}$ with $\|R\|^2 \le M$, $\|R^{-1}\|^2 \le 1/m$ (Kato's form theory). Then

$$\big\|\hat{\mathcal{O}}\big\|_{\mathcal{K}_2} = \big\|\mathcal{K}_2^{-1/2}\hat{\mathcal{O}}\mathcal{K}_2^{-1/2}\big\| = \big\|\mathcal{K}_1^{-1/2}R^{-1}\,\hat{\mathcal{O}}\,R^{-1}\mathcal{K}_1^{-1/2}\big\| \le \|R^{-1}\|^2\,\big\|\hat{\mathcal{O}}\big\|_{\mathcal{K}_1},$$

and symmetrically for the reverse inequality, giving the constants $1/m$ and M. (See, e.g., [115] for equivalence of closed forms and induced operator norms.)

Branch calibration (law channel).

Definition 7.362 (Presentation–equivalent micro–Hamiltonians)**.** Let $\hat{H}^{[j]}_{\mathrm{micro}}(t;\lambda_{\mathrm{UV}},\ell)$, $j = 0, 1$, be two *law–channel* generators as in Definition 7.350, built from the same baseline $\hat{H}_0$ and the same set $\{\varepsilon_B, \hat{\mathcal{O}}_B^{(\lambda_{\mathrm{UV}},\ell)}\}$, but differing by:

(P1) G**–preserving unitary field redefinitions:** $\hat{H}^{[1]}_{\mathrm{micro}} = \hat{U}_G^\dagger \hat{H}^{[0]}_{\mathrm{micro}} \hat{U}_G - i\,\hat{U}_G^\dagger \dot{\hat{U}}_G$, with $\hat{U}_G(t)$ unitary, $\mathrm{gh}(\hat{U}_G) = 0$, and $[\hat{U}_G,\, \hat{J}_a] = 0$ for all G–generators $\hat{J}_a$ (hence G–symmetric); and

(P2) **BRST–exact deformations:** $\hat{H}^{[1]}_{\mathrm{micro}} = \hat{H}^{[0]}_{\mathrm{micro}} + \{\hat{Q}_{\mathrm{BRST}},\, \delta\Psi(t)\}$, with $\mathrm{gh}(\delta\Psi) = -1$ and $\delta\Psi^\dagger = \delta\Psi$, so that $\{\hat{Q}_{\mathrm{BRST}}, \delta\Psi\}$ is Hermitian.

We write $\hat{H}^{[0]}_{\mathrm{micro}} \sim \hat{H}^{[1]}_{\mathrm{micro}}$ and say the two presentations are *presentation–equivalent*. *Note.* $\delta\Psi$ is a presentation fermion distinct from the fixed gauge–fixing fermion Ψ already used in $\{\hat{Q}_{\mathrm{BRST}}, \Psi\}$.

Definition 7.363 (Branch unitaries and calibration)**.** Given a presentation $\hat{H}^{[j]}_{\mathrm{micro}}$, define the law–channel time–evolution

$$\boxed{\begin{aligned} U_0^{(\mathrm{law;micro})}(T;\lambda_{\mathrm{UV}},\ell) &:= \mathcal{T}\,\exp\Big[-i\int_0^T \mathrm{d}t\,\hat{H}^{[0]}_{\mathrm{micro}}(t;\lambda_{\mathrm{UV}},\ell)\Big],\\ U_1^{(\mathrm{law;micro})}(T;\lambda_{\mathrm{UV}},\ell) &:= \mathcal{T}\,\exp\Big[-i\int_0^T \mathrm{d}t\,\hat{H}^{[1]}_{\mathrm{micro}}(t;\lambda_{\mathrm{UV}},\ell)\Big]. \end{aligned}} \tag{7.527}$$

The *law visibility* is read as in eq. (7.122) (with the same BRST–closed tags and channel projectors), and its overall phase ϕ_{law} is *calibrated once* so that $\Lambda_{\mathrm{law}} = 1$ in the symmetric, unbroken limit (cf. Section 7.3.9).

Lemma 7.364 (Presentation coboundary)**.** *If $\hat{H}^{[1]}_{\mathrm{micro}} \sim \hat{H}^{[0]}_{\mathrm{micro}}$ in the sense of Definition 7.362, then*

$$U_1^{(\mathrm{law;micro})}(T) = \hat{U}_G^\dagger(T)\,\mathcal{T}\,\exp\Big(-i\int_0^T \mathrm{d}t\,\{\hat{Q}_{\mathrm{BRST}},\,\delta\Psi_H(t)\}\Big)\,U_0^{(\mathrm{law;micro})}(T)\,\hat{U}_G(0),$$

where $\delta\Psi_H(t)$ is $\delta\Psi(t)$ in the Heisenberg picture built with $\hat{H}^{[0]}_{\mathrm{micro}}$. Thus $U_1^{(\mathrm{law;micro})}(T)$ and $U_0^{(\mathrm{law;micro})}(T)$ differ by a G–symmetric unitary conjugation and a BRST–exact time–ordered exponential.

Proof. Set

$$W(t) := \mathcal{T}\,\exp\Big(-i\int_0^t \{\hat{Q}_{\mathrm{BRST}},\delta\Psi_H(u)\}\,du\Big), \qquad X(t) := \hat{U}_G^\dagger(t)\,W(t)\,U_0^{(\mathrm{law;micro})}(t)\,\hat{U}_G(0).$$

Then $X(0)=\mathbf{1}$. Using

$$i\,\partial_t U_0^{(\mathrm{law;micro})}(t) = \hat{H}^{[0]}_{\mathrm{micro}}(t)\,U_0^{(\mathrm{law;micro})}(t), \quad i\,\partial_t W(t) = \{\hat{Q}_{\mathrm{BRST}},\delta\Psi_H(t)\}\,W(t),$$

and $\partial_t \hat{U}_G^\dagger(t) = -\hat{U}_G^\dagger(t)\,\dot{\hat{U}}_G(t)\,\hat{U}_G^\dagger(t)$, a product-rule computation gives

$$i\,\partial_t X(t) = \Big(\hat{U}_G^\dagger \hat{H}^{[0]}_{\mathrm{micro}}\hat{U}_G - i\,\hat{U}_G^\dagger\dot{\hat{U}}_G + \hat{U}_G^\dagger\{\hat{Q}_{\mathrm{BRST}},\delta\Psi_H(t)\}\hat{U}_G\Big)X(t).$$

Assuming $[\hat{U}_G,\hat{Q}_{\mathrm{BRST}}]=0$ and that $\delta\Psi$ is G–scalar (so $\hat{U}_G^\dagger\{\hat{Q}_{\mathrm{BRST}},\delta\Psi_H\}\hat{U}_G = \{\hat{Q}_{\mathrm{BRST}},\delta\Psi_H\}$), the bracket equals

$$\hat{H}^{[1]}_{\mathrm{micro}}(t) = \hat{U}_G^\dagger(t)\,\hat{H}^{[0]}_{\mathrm{micro}}(t)\,\hat{U}_G(t) - i\,\hat{U}_G^\dagger(t)\dot{\hat{U}}_G(t) + \{\hat{Q}_{\mathrm{BRST}},\delta\Psi_H(t)\},$$

which is precisely the presentation relation of Definition 7.362. Hence $i\,\partial_t X(t) = \hat{H}^{[1]}_{\mathrm{micro}}(t)\,X(t)$ with $X(0)=\mathbf{1}$, so by uniqueness of time-ordered solutions, $X(T) = U_1^{(\mathrm{law;micro})}(T)$, i.e.

$$U_1^{(\mathrm{law;micro})}(T) = \hat{U}_G^\dagger(T)\,\mathcal{T}\,\exp\Big(-i\int_0^T \{\hat{Q}_{\mathrm{BRST}},\delta\Psi_H(t)\}\,dt\Big)\,U_0^{(\mathrm{law;micro})}(T)\,\hat{U}_G(0). \qquad \square$$

Proposition 7.365 (Physical equivalence of branch choices)**.** *Let $\mathsf{P}_{\mathrm{phys}}$ project onto $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$. Under the hypotheses above (BRST–closed tags; gh= 0 objects), for any ghost–number–zero density matrix ρ_0 supported on $\mathcal{H}_{\mathrm{phys}}$,*

$$\mathrm{Tr}\Big[U_1^{(\mathrm{law;micro})\dagger}(T)\,U_0^{(\mathrm{law;micro})}(T)\,\rho_0\Big] = \mathrm{Tr}\Big[U_{1'}^{(\mathrm{law;micro})\dagger}(T)\,U_{0'}^{(\mathrm{law;micro})}(T)\,\rho_0\Big]$$

for any other pair of presentations $([0'],[1'])$ equivalent to $([0],[1])$ in the sense of Definition 7.362, up to the single *overall calibration phase ϕ_{law}. Hence Λ_{law} is well defined on the* presentation class *and independent of gauge parameters (Nielsen identities; cf. Section 7.1.3).*

Idea. By Lemma 7.364, $U^{[1]}$ and $U^{[0]}$ differ by G–symmetric conjugations and a BRST–exact cocycle. Conjugations drop inside the trace with ρ_0 that commutes with G, and BRST–exact insertions vanish on $\mathcal{H}_{\mathrm{phys}}$ (cohomological triviality). Gauge–parameter changes produce ζ–exact deformations and are handled identically [117, 82]. $\square$

Proof. Fix two equivalent presentations $[0], [1]$ and $[0'], [1']$ in the sense of Definition 7.362. By Lemma 7.364 there exist (i) G-symmetric unitaries $W_a \in G$ (possibly scale-tagged but BRST-closed, cf. Section 7.1.3) and (ii) a finite family of BRST–exact coboundary insertions (i.e. ζ-exact) such that, for each branch $a \in \{0, 1\}$,

$$U_{a'}^{(\mathrm{law;micro})}(T) = e^{\,i\varphi_a}\, W_a\, U_a^{(\mathrm{law;micro})}(T)\, W_a^{-1} \;+\; \mathcal{B}_a, \tag{7.528}$$

where each term in $\mathcal{B}_a$ is a time-ordered insertion that is ζ-exact. (BRST-closed tags ensure cohomological well-posedness; cf. Section 7.1.3.)

Taking adjoints and forming the interference product,

$$\begin{aligned} U_{1'}^{(\mathrm{law;micro})\dagger}(T)\, U_{0'}^{(\mathrm{law;micro})}(T) = e^{\,i(\varphi_0-\varphi_1)} \Big[W_1\, U_1^{(\mathrm{law;micro})\dagger}(T)\, W_1^{-1} \Big] \\ \times \Big[W_0\, U_0^{(\mathrm{law;micro})}(T)\, W_0^{-1} \Big] \;+\; \mathcal{B}, \end{aligned} \tag{7.529}$$

with $\mathcal{B}$ a finite sum of time-ordered ζ-exact insertions (Duhamel expansion; closure under time ordering).

Let $\rho_0 = \mathsf{P}_{\mathrm{phys}} \rho_0 \mathsf{P}_{\mathrm{phys}}$ with $\mathrm{gh}(\rho_0) = 0$. By G-invariance of the physical sector, $[\,\rho_0, W_a\,] = 0$, hence by cyclicity

$$\mathrm{Tr}\Big(\big[W_1 U_1^\dagger W_1^{-1}\big] \big[W_0 U_0 W_0^{-1}\big] \rho_0 \Big) = \mathrm{Tr}\Big(U_1^\dagger\, U_0\, \rho_0 \Big). \tag{7.530}$$

Any ζ-exact insertion vanishes in the physical trace:

$$\mathrm{Tr}\Big(\zeta X\, \rho_0 \Big) = \mathrm{Tr}\Big(\{\hat{Q}_{\mathrm{BRST}}, X\}\, \rho_0 \Big) = \mathrm{Tr}\Big(\hat{Q}_{\mathrm{BRST}} X \rho_0 \Big) - \mathrm{Tr}\Big(X \hat{Q}_{\mathrm{BRST}} \rho_0 \Big) = 0, \tag{7.531}$$

using $\hat{Q}_{\mathrm{BRST}} \rho_0 = \rho_0 \hat{Q}_{\mathrm{BRST}} = 0$. Applying (7.530)–(7.531) to (7.529) gives

$$\mathrm{Tr}\Big[U_{1'}^\dagger(T)\, U_{0'}(T)\, \rho_0 \Big] = e^{\,i(\varphi_0-\varphi_1)}\; \mathrm{Tr}\Big[U_1^\dagger(T)\, U_0(T)\, \rho_0 \Big]. \tag{7.532}$$

Define the single calibration phase $\phi_{\mathrm{law}} := \varphi_0 - \varphi_1$. Fixing it at baseline sets $e^{i\phi_{\mathrm{law}}} = 1$, yielding the equality asserted in Proposition 7.365. Gauge–parameter variations obey the Nielsen identity (Section 7.1.3),

$$\partial_\xi \langle \mathcal{O} \rangle_{\mathrm{phys}} = \langle \zeta(\zeta_\xi)\, \mathcal{O} \rangle = \Big\langle \{\hat{Q}_{\mathrm{BRST}}, \zeta_\xi\}\, \mathcal{O} \Big\rangle,$$

which vanishes by (7.531), so Λ_{law} is presentation-class and gauge independent [117, 82]. □

Intuition. Equivalent presentations differ by harmless G-rotations and BRST coboundaries. Rotations disappear in a G-invariant trace, and coboundaries are invisible on $H^0(\hat{Q}_{\mathrm{BRST}})$. The only residue is a fixed global $U(1)$ calibration between branches, which we pin once (ϕ_{law}) and forget.

Lemma 7.366 (BRST/Nielsen decomposition of regulator derivatives)**.** *Let the tagged branch generators $\mathcal{L}_a(t; \lambda_{\mathrm{UV}}, \ell)$ be built with BRST–closed tags (gh= 0) that commute with the physical*

projector (cf. Section 7.1.3). For $\kappa \in \{\ln \lambda_{\mathrm{UV}}, \ln \ell\}$ *there exist operator families* $\mathcal{N}_{a,\kappa}(t)$ *with* $\mathrm{gh}(\mathcal{N}_{a,\kappa}) = -1$ *and* $\Delta_{a,\kappa}(t)$ *with* $\mathrm{gh}(\Delta_{a,\kappa}) = 0$ *such that*

$$\partial_\kappa \mathcal{L}_a(t) = \zeta\, \mathcal{N}_{a,\kappa}(t) \;+\; \Delta_{a,\kappa}(t), \tag{7.533}$$

and, within any plateau window (Section 6.12.4), the plateau-suppressed remainder $\Delta_{a,\kappa}$ *is uniformly bounded by the prescribed tolerances.*

Proof of Lemma 7.366. Fix $a \in \{0,1\}$ and $\kappa \in \{\ln \lambda_{\mathrm{UV}}, \ln \ell\}$. Write the tagged generator as

$$\mathcal{L}_a(t) = \mathcal{L}_a^{\mathrm{tag}}(t) + \zeta\, \Psi_a(t),$$

where the tag dependence is entirely in $\mathcal{L}_a^{\mathrm{tag}}$ (built from $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$) and Ψ_a is the gauge fermion ($\mathrm{gh}= -1$).

(i) Gauge-fixing sector. Since tags are BRST–closed ($\mathrm{gh}= 0$), $\partial_\kappa\big(\zeta \Psi_a\big) = \zeta\big(\partial_\kappa \Psi_a\big)$. Define

$$\mathcal{N}_{a,\kappa}^{(\mathrm{gf})}(t) := \partial_\kappa \Psi_a(t) \qquad (\mathrm{gh} = -1).$$

(ii) Tag-dependent sector via smooth projector calculus. On a certified plateau (Section 6.12.4), the band projectors vary smoothly with a spectral gap to the band edges. Hence there exists an anti-Hermitian generator $G_{a,\kappa}(t)$ ($\mathrm{gh}= 0$) such that, for each tagged operator block,

$$\partial_\kappa \mathcal{O}_B^{(\lambda_{\mathrm{UV}},\ell)}(t) = \Big[G_{a,\kappa}(t),\, \mathcal{O}_B^{(\lambda_{\mathrm{UV}},\ell)}(t)\Big] \;+\; R_{B,a,\kappa}(t),$$

with a remainder bounded uniformly by the plateau tolerances. Summing the blocks,

$$\partial_\kappa \mathcal{L}_a^{\mathrm{tag}}(t) = \Big[G_{a,\kappa}(t),\, \mathcal{L}_a^{\mathrm{tag}}(t)\Big] \;+\; R_{a,\kappa}(t), \qquad \|R_{a,\kappa}\| = \mathcal{O}(\text{tolerance}).$$

(iii) Cohomological triviality of presentation variations. Within the BRST–consistent reading (Section 7.1.3) and anomaly-free sector, presentation changes generated by $G_{a,\kappa}$ act trivially on $H^0(\hat{Q}_{\mathrm{BRST}})$; therefore the commutator $\Big[G_{a,\kappa}, \mathcal{L}_a^{\mathrm{tag}}\Big]$ is BRST–exact. Thus there exists $\widetilde{\mathcal{N}}_{a,\kappa}(t)$ with $\mathrm{gh}(\widetilde{\mathcal{N}}_{a,\kappa}) = -1$ such that

$$\Big[G_{a,\kappa}(t),\, \mathcal{L}_a^{\mathrm{tag}}(t)\Big] = \zeta\, \widetilde{\mathcal{N}}_{a,\kappa}(t).$$

(Equivalently, apply the BRST doublet/quartet mechanism, cf. Lemma 6.211 and eq. (6.279).)

(iv) Assemble $\mathcal{N}_{a,\kappa}$ *and* $\Delta_{a,\kappa}$. Define

$$\mathcal{N}_{a,\kappa}(t) := \mathcal{N}_{a,\kappa}^{(\mathrm{gf})}(t) + \widetilde{\mathcal{N}}_{a,\kappa}(t), \qquad \Delta_{a,\kappa}(t) := R_{a,\kappa}(t).$$

Then $\mathrm{gh}(\mathcal{N}_{a,\kappa}) = -1$, $\mathrm{gh}(\Delta_{a,\kappa}) = 0$, and combining (i)–(iii) yields

$$\partial_\kappa \mathcal{L}_a(t) = \zeta\, \mathcal{N}_{a,\kappa}(t) \;+\; \Delta_{a,\kappa}(t),$$

with $\Delta_{a,\kappa}$ uniformly bounded on the plateau as required. □

Proposition 7.367 (Calibration and stability on plateaus)**.** *Choose ϕ_{law} so that $\Lambda_{\mathrm{law}} = 1$ in the unbroken baseline ($\varepsilon_B = 0$). Then, within any plateau window for the tags (cf. Section 6.12.4),*

$$\partial_{\ln \lambda_{\mathrm{UV}}} \Lambda_{\mathrm{law}} = 0, \qquad \partial_{\ln \ell} \Lambda_{\mathrm{law}} = 0,$$

up to the prescribed tolerances, and the value of Λ_{law} is independent of presentation choices and gauge parameters. Reason. *Tags are BRST–closed ($gh = 0$) and commute with projectors; presentation changes are G–unitary or ζ–exact; plateaus remove residual regulator dependence.*

Proof of Proposition 7.367. Let

$$\Lambda_{\mathrm{law}}(T; \lambda_{\mathrm{UV}}, \ell) := \mathrm{Tr}\Big[U_1^{(\mathrm{law;micro})\dagger}(T)\, U_0^{(\mathrm{law;micro})}(T)\, \rho_0 \Big], \qquad \rho_0 = \mathsf{P}_{\mathrm{phys}} \rho_0 \mathsf{P}_{\mathrm{phys}},\ \mathrm{gh}(\rho_0) = 0,$$

with tags BRST–closed and commuting with projectors (cf. Section 7.1.3). Fix ϕ_{law} so that at $\varepsilon_B = 0$ one has $\Lambda_{\mathrm{law}} = 1$.

Step 1: Regulator derivatives reduce to ζ–exact insertions. For $\kappa \in \{\ln \lambda_{\mathrm{UV}}, \ln \ell\}$ the Duhamel formula gives

$$\partial_\kappa U_a(T) = \int_0^T dt\; U_a(T,t)\, \partial_\kappa \mathcal{L}_a(t)\, U_a(t,0),$$

hence

$$\begin{aligned} \partial_\kappa \Lambda_{\mathrm{law}} &= \mathrm{Tr}\Big[\big(\partial_\kappa U_1^\dagger\big) U_0\, \rho_0 \Big] + \mathrm{Tr}\Big[U_1^\dagger \big(\partial_\kappa U_0\big)\, \rho_0 \Big] \\ &= \sum_{a=0,1} \sigma_a \int_0^T dt\ \ \mathrm{Tr}\Big[U_1^\dagger(T)\, \mathcal{T}\{\partial_\kappa \mathcal{L}_a(t)\}\, U_0(T)\, \rho_0 \Big], \end{aligned} \tag{7.534}$$

with $\sigma_1 = -1$, $\sigma_0 = +1$. Apply Lemma 7.366: $\partial_\kappa \mathcal{L}_a = \zeta\, \mathcal{N}_{a,\kappa} + \Delta_{a,\kappa}$.

Step 2: ζ–exact pieces vanish on $\mathcal{H}_{\mathrm{phys}}$; G–conjugations drop. Inserting (7.533) into (7.534), all terms with $\zeta\, \mathcal{N}_{a,\kappa}$ vanish under the trace with ρ_0 by the standard cohomological argument (cf. the step around (7.531) in Proposition 7.365):

$$\mathrm{Tr}(\zeta X\, \rho_0) = \mathrm{Tr}\big(\{\hat{Q}_{\mathrm{BRST}}, X\} \rho_0 \big) = 0, \qquad \mathrm{gh}(X) \text{ definite},$$

using $\hat{Q}_{\mathrm{BRST}} \rho_0 = \rho_0 \hat{Q}_{\mathrm{BRST}} = 0$ and $\mathrm{gh}(\rho_0) = 0$. Any G–symmetric conjugations from presentation changes cancel inside the trace with ρ_0 (cyclicity and $[\,\rho_0, G\,] = 0$; cf. Proposition 7.365).

Step 3: Plateau suppression yields regulator stability. Only the $\Delta_{a,\kappa}$ pieces remain. By Section 6.12.4, their contribution is bounded by the plateau tolerances, hence

$$\partial_{\ln \lambda_{\mathrm{UV}}} \Lambda_{\mathrm{law}} = \mathcal{O}(\text{tolerance}), \qquad \partial_{\ln \ell} \Lambda_{\mathrm{law}} = \mathcal{O}(\text{tolerance}),$$

vanishing in the ideal plateau limit.

Step 4: Presentation/gauge independence. Presentation changes within an equivalence class act by G–unitaries and ζ–coboundaries; by Proposition 7.365 they affect Λ_{law} only by the single global calibration phase. With ϕ_{law} fixed at baseline ($\varepsilon_B = 0$), Λ_{law} is presentation–independent on plateaus. Gauge–parameter variations obey Nielsen identities,

$$\partial_\xi \langle \mathcal{O} \rangle_{\mathrm{phys}} = \big\langle \{\hat{Q}_{\mathrm{BRST}}, \mathcal{N}_\xi\}\, \mathcal{O} \big\rangle,$$

which vanish on $\mathcal{H}_{\mathrm{phys}}$ by the same BRST argument, giving gauge–parameter independence [117, 82]. □

Remark 7.368 (Calibration intuition and two-branch diagnostic)**.** Calibrated in the symmetric baseline ($\varepsilon_B = 0$) so that $\Lambda_{\text{law}} = 1$, the branch interferometer only "feels" BRST–exact coboundaries generated by Nielsen kernels and G–unitary presentation changes; both are invisible on $\mathcal{H}_{\text{phys}}$ (cf. Lemma 7.366 and Proposition 7.365). On certified plateaus (Section 6.12.4), regulator variations contribute only the plateau–suppressed remainders $\Delta_{a,\kappa}$, hence $\partial_{\ln \lambda_{\text{UV}}} \Lambda_{\text{law}} = \partial_{\ln \ell} \Lambda_{\text{law}} = 0$ up to tolerances and gauge–parameter independence holds by the Nielsen identities (see Proposition 7.367).

Operationally, keeping two law–channel branches with the same baseline $\hat{H}_0$ but different presentations remains useful: any contrast that survives the BRST/G–unitary and gauge/-plateau filtering diagnoses genuine G–breaking encoded in the scale–tagged Ward source (cf. Proposition 7.353), not a presentation or regulator artifact.

State posture and calibration (identity branch).

Definition 7.369 (State channel: identity branches and phase calibration)**.** Reuse the *identity* state branches of eq. (7.125), namely

$$U_0^{(\text{state})} = U_1^{(\text{state})} = U_0,$$

by setting the W–operator in eq. (7.124) to the identity (no SSB kick). Fix the state–visibility phase ϕ_{state} once and for all as in eq. (7.129) so that, in the symmetric/unbroken limit, the state visibility is calibrated to unit modulus. All objects are evaluated *on the band* with the BRST–closed tags $P_x(\lambda_{\text{UV}})$ and $K_{\text{IR}}^{(\ell)}$ (commuting with channel projectors), and on the physical sector $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$ (cf. Section 7.1.3).

Proposition 7.370 (Channel separation under identity state branches)**.** *Under Definition 7.369 and with the law channel generated by Definition 7.350, the state visibility measured as in eq.* (7.129) *satisfies*

$$\Lambda_{\text{state}}(\lambda_{\text{UV}}, \ell; T) = 1$$

in the symmetric/unbroken limit, independently of presentation choices and gauge parameters; any deviation from unit modulus in the QIII setting is thus attributable to the law *channel.*
Reason. *Identity branches imply* $U_0^{(\text{state})} = U_1^{(\text{state})}$*, so the overlap/trace defining* Λ_{state} *reduces to a normed inner product; BRST–closed tags (gh= 0) and Nielsen identities ensure gauge–parameter independence on* $\mathcal{H}_{\text{phys}}$ *(cf. Section 7.1.3).*

Proof. Let $\rho_0 = \mathsf{P}_{\text{phys}} \rho_0 \mathsf{P}_{\text{phys}}$ be a unit–trace, ghost–number–zero density matrix on $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$. By Definition 7.369, the identity–state condition gives $U_0^{(\text{state})}(T) = U_1^{(\text{state})}(T) =: U^{(\text{state})}(T)$. Thus, using eq. (7.129),

$$\Lambda_{\text{state}}(\lambda_{\text{UV}}, \ell; T) = \operatorname{Tr}\Big[U^{(\text{state})\dagger}(T)\, U^{(\text{state})}(T)\, \rho_0\Big] = \operatorname{Tr}(\rho_0) = 1,$$

since $U^{(\text{state})}$ is unitary and $\operatorname{Tr} \rho_0 = 1$.

Presentation independence. Any presentation change within the equivalence class acts by a G–unitary and BRST–exact coboundaries that are *identical* on both identity branches. Writing $U^{(\text{state})} \mapsto U'^{(\text{state})} = e^{i\varphi}\, W\, U^{(\text{state})} W^{-1}$ with $W \in G$, cyclicity and $[\,\rho_0, W\,] = 0$ give

$$\operatorname{Tr}\Big[U'^{(\text{state})\dagger} U'^{(\text{state})} \rho_0\Big] = \operatorname{Tr}\Big[U^{(\text{state})\dagger} U^{(\text{state})} \rho_0\Big].$$

BRST–exact insertions induced at the generator level vanish on $\mathcal{H}_{\mathrm{phys}}$:

$$\mathrm{Tr}\Big(\{\hat{Q}_{\mathrm{BRST}}, X\}\,\rho_0\Big) = 0 \quad \text{for any } X \text{ of definite ghost number},$$

cf. the argument around (7.531) in Proposition 7.365.

Gauge–parameter independence. Either directly,

$$\partial_\xi \Lambda_{\mathrm{state}} = \mathrm{Tr}\Big[\partial_\xi(U^\dagger U)\,\rho_0\Big] = \mathrm{Tr}\Big[0\cdot\rho_0\Big] = 0 \quad (\text{unitarity } U^\dagger U = \mathbf{1}),$$

or via the Nielsen identity (cf. Section 7.1.3),

$$\partial_\xi \langle \mathcal{O}\rangle_{\mathrm{phys}} = \Big\langle \{\hat{Q}_{\mathrm{BRST}}, \mathcal{N}_\xi\}\,\mathcal{O}\Big\rangle,$$

and with $\mathcal{O} = U^{(\mathrm{state})\dagger}U^{(\mathrm{state})} = \mathbf{1}$ on $\mathcal{H}_{\mathrm{phys}}$ one again finds $\partial_\xi \Lambda_{\mathrm{state}} = 0$, since BRST–exact insertions drop out and the tags are BRST–closed (gh$= 0$).

Therefore, in the symmetric/unbroken limit, $\Lambda_{\mathrm{state}}(\lambda_{\mathrm{UV}}, \ell; T) = 1$ independently of presentation and gauge. Any deviation from unit modulus observed in QIII must originate in the *law* channel. □

Intuition. Identity branches make the state-channel interferometer a norm check: $U^\dagger U = \mathbf{1}$ inside a G-invariant, BRST-closed trace, so it pins to one; only the law-channel dynamics can move it off unit modulus.

Initial readouts and records.

Definition 7.371 (Initial law deficit and state–coherence check)**.** Choose a minimal coarse–graining scale $\ell_{\min}$ consistent with the reciprocal–resolution constraints used around Section 6.12.4 (cf. Proposition 4.436). For a fixed readout time $T^\star$ (the interferometric normalisation of eq. (7.116)), define the *initial law deficit* and confirm state coherence via

$$\boxed{E^{(\mathrm{init})}_{\mathrm{law}}(\ell_{\min}) := 1 - \Lambda_{\mathrm{law}}(\lambda_{\mathrm{UV}}, \ell_{\min}; T^\star) > 0, \qquad \Big|1 - \Lambda_{\mathrm{state}}(\lambda_{\mathrm{UV}}, \ell_{\min}; T^\star)\Big| \le \varepsilon_{\mathrm{init}}.} \tag{7.535}$$

The tolerance $\varepsilon_{\mathrm{init}}$ is chosen below the plateau thresholds of Section 6.12.4 so that the statement "$\Lambda_{\mathrm{state}} \approx 1$" is meaningful uniformly in the volume V.

Proposition 7.372 (Gauge/presentation independence and plateau robustness)**.** *With the calibrations of Definition 7.369 and BRST–closed tags, the quantities in eq. (7.535) are independent of gauge–parameter choices and of presentation–equivalent changes in the law channel (cf. Definition 7.362 and Proposition 7.365). If, furthermore,* $(\lambda_{\mathrm{UV}}, \ell)$ *is varied* within *a plateau window of Section 6.12.4, then*

$$\partial_{\ln \lambda_{\mathrm{UV}}} E^{(\mathrm{init})}_{\mathrm{law}} = 0, \qquad \partial_{\ln \ell} E^{(\mathrm{init})}_{\mathrm{law}} = 0,$$

and Λ_{state} *remains within* $\varepsilon_{\mathrm{init}}$ *of unity, up to the prescribed plateau tolerances.* Reason. *BRST–closed tags (gh$= 0$) commute with projectors and preserve cohomology; Nielsen identities render physical readouts gauge independent; plateau bounds control UV/IR sensitivity (cf. Sections 6.12.4 and 7.1.3).*

Proof. Let the initial deficits be defined as in eq. (7.535), built from the ghost–number–zero physical state $\rho_0 = \mathsf{P}_{\mathrm{phys}}\rho_0\mathsf{P}_{\mathrm{phys}}$ and the QIII evolutions. Assume the state–channel calibration of Definition 7.369 (identity branches in the state channel) and BRST–closed tags that commute with projectors (cf. Section 7.1.3).

(i) Gauge–parameter and presentation independence. Each ingredient entering eq. (7.535) is a physical, gh= 0 trace on $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$. Gauge–parameter variations insert BRST–exact Nielsen terms,

$$\partial_\xi \langle \mathcal{O} \rangle_{\mathrm{phys}} = \Big\langle \{\hat{Q}_{\mathrm{BRST}}, \mathcal{N}_\xi\}\, \mathcal{O} \Big\rangle,$$

which vanish on $\mathcal{H}_{\mathrm{phys}}$ by the cohomological null property $\mathrm{Tr}\Big(\{\hat{Q}_{\mathrm{BRST}}, X\}\rho_0\Big) = 0$ (see the step around (7.531)) [117, 82]. Presentation–equivalent changes in the *law* channel act by G–unitary conjugations and BRST coboundaries; these drop in the same physical trace (cyclicity, $[\,\rho_0, G\,] = 0$), leaving at most the single global calibration phase already fixed at baseline by Proposition 7.365. Therefore the quantities in eq. (7.535) are independent of gauge parameters and presentation choices.

(ii) Plateau robustness for regulator variations. Within a plateau window (Section 6.12.4), the tagged generators admit the BRST/Nielsen decomposition (cf. Lemma 7.366):

$$\partial_\kappa \mathcal{L}_a(t) = \zeta\, \mathcal{N}_{a,\kappa}(t) + \Delta_{a,\kappa}(t), \qquad \kappa \in \{\ln \lambda_{\mathrm{UV}}, \ln \ell\},$$

with $\mathrm{gh}(\mathcal{N}_{a,\kappa}) = -1$ and $\Delta_{a,\kappa}$ the plateau–suppressed remainder (bounded by the prescribed tolerances). Applying the Duhamel formula to each law–channel factor inside eq. (7.535), all BRST–exact insertions $\zeta\,\mathcal{N}_{a,\kappa}$ vanish on $\mathcal{H}_{\mathrm{phys}}$ as above, while the remaining $\Delta_{a,\kappa}$ contributions are uniformly bounded on the plateau. Hence

$$\partial_{\ln \lambda_{\mathrm{UV}}} E^{(\mathrm{init})}_{\mathrm{law}} = 0, \qquad \partial_{\ln \ell} E^{(\mathrm{init})}_{\mathrm{law}} = 0$$

in the ideal plateau limit, and remain within tolerance otherwise.

(iii) Stability of the state visibility. Under Definition 7.369, the state–channel visibility satisfies $\Lambda_{\mathrm{state}} = 1$ in the symmetric limit (unitarity and $\mathrm{Tr}\,\rho_0 = 1$; see also Proposition 7.370). Plateau variations enter as in (ii): BRST–exact insertions cancel on $\mathcal{H}_{\mathrm{phys}}$, and any residual regulator dependence is plateau–suppressed; therefore Λ_{state} stays within $\varepsilon_{\mathrm{init}}$ of unity, up to the same tolerances.

Combining (i)–(iii) yields the proposition. □

Intuition. The "init" observables are physical traces on $H^0(\hat{Q}_{\mathrm{BRST}})$. Gauge and presentation moves are BRST–exact or G–unitary, hence invisible; on a plateau, the only leftover regulator drift is uniformly small, so the law–deficit is flat and the state visibility hugs one.‘‘

Remark 7.373 (Choice of $\ell_{\min}$ and reciprocal resolution)**.** Choose $\ell_{\min}$ at the smallest scale that simultaneously satisfies the reciprocal–resolution limits of Proposition 4.436 *and* the plateau criterion of Section 6.12.4: (i) the UV projector $P_x(\lambda_{\mathrm{UV}})$ cleanly resolves modes down to $1/\ell_{\min}$ without aliasing or leakage (cf. Section 6.12.4); (ii) the IR window kernel W_ℓ with $\ell \geq \ell_{\min}$ preserves locality/cluster properties and avoids under–resolving Ward sources (cf. Section 6.12.4). With this choice—and with the baseline calibration fixed so that

$\Lambda_{\text{law}} = 1$ at $\varepsilon_B = 0$ (Proposition 7.367) and identity–state branches in place (Definition 7.369 and Proposition 7.370)—the meters are plateau–robust and presentation/gauge–independent:

$$\partial_{\ln \lambda_{\text{UV}}} \Lambda_{\text{law}} = \partial_{\ln \ell} \Lambda_{\text{law}} = 0 \text{ (up to plateau tolerances)}, \qquad \Lambda_{\text{state}} = 1 + \mathcal{O}(\text{tolerance}),$$

with invariance under presentation–equivalent changes guaranteed by Propositions 7.365 and 7.372. *Intuition.* $\ell_{\min}$ is set at the "reciprocal–resolution knee" where UV band–limit and IR smearing balance; below it, UV aliasing or regulator drift contaminates reads, while above it, IR averaging dilutes genuine G–breaking. Picking $\ell_{\min}$ at this knee lands the meters on a plateau where only physical law–channel effects can move the readout.

Diagnostic Hygiene.

Lemma 7.374 (On–shell gauge safety)**.** *Assume the QIII breaking operators satisfy* $\zeta\, \hat{\mathcal{O}}_B = \{\hat{Q}_{\text{BRST}}, \hat{\mathcal{O}}_B\} = 0$ *and* $\text{gh}(\hat{\mathcal{O}}_B) = 0$ *(cf. Definition 7.356). Then the Slavnov–Taylor identity* $\mathcal{S}(\Gamma) = 0$ *of eq. (7.16) holds on shell (anomaly–free Stage I); any nonzero law–channel deficit at small* ℓ *arises from* global *Ward sources (explicit breaking of* G*), not from gauge anomalies (which are excluded by Section 7.2.8).* Justification. *BRST closure prevents cohomological obstructions; anomaly freedom ensures ST exactness on shell [117, 442, 243].*

Proof. Work in the QIII setting with admissible breakings (Definition 7.356), i.e. $\zeta\, \hat{\mathcal{O}}_B = 0$ and $\text{gh}(\hat{\mathcal{O}}_B) = 0$. At the classical level the deformation

$$S_{\text{cl}} \longrightarrow S_{\text{cl}} + \varepsilon_B \int \hat{\mathcal{O}}_B$$

preserves BRST symmetry, since $\zeta S_{\text{cl}} = 0$ and $\zeta \hat{\mathcal{O}}_B = 0$. Therefore the Zinn–Justin (Slavnov–Taylor) identity holds for the tree functional $\Gamma^{(0)}$:

$$\mathcal{S}(\Gamma^{(0)}) = 0, \qquad \text{cf. (7.16)}.$$

At the quantum level let $\Gamma = \sum_{n \geq 0} \hbar^n \Gamma^{(n)}$ be the renormalized 1PI functional. If a breaking appears, organize it as

$$\mathcal{S}(\Gamma) = \hbar^n\, \Delta_{\text{ST}}^{(n)} + \mathcal{O}(\hbar^{n+1}),$$

with ghost number $\text{gh}(\Delta_{\text{ST}}^{(n)}) = 1$. The Wess–Zumino consistency condition (graded Jacobi identity) reads

$$\mathcal{B}_\Gamma\, \Delta_{\text{ST}}^{(n)} = 0,$$

where $\mathcal{B}_\Gamma$ is the linearized Slavnov operator (nilpotent on shell). By the Stage I anomaly–freedom hypothesis (cf. Section 7.2.8) the on–shell cohomology $H^1(\mathcal{B}_\Gamma)$ is trivial [117, 442, 243]. Hence there exists a local functional $\widehat{\Delta}_{\text{ST}}^{(n)}$ of ghost number 0 such that

$$\Delta_{\text{ST}}^{(n)} = \mathcal{B}_\Gamma\, \widehat{\Delta}_{\text{ST}}^{(n)} \; + \; (\text{terms} \propto \text{equations of motion}).$$

Adding the counterterm $-\hbar^n\, \widehat{\Delta}_{\text{ST}}^{(n)}$ to Γ removes the breaking at order $\hbar^n$. Inducting in n yields, *on shell*,

$$\mathcal{S}(\Gamma) = 0.$$

Thus: (i) since $\hat{\mathcal{O}}_B$ is BRST–closed with gh $= 0$, it cannot seed a ghost–number–one gauge anomaly; (ii) any surviving nonzero law–channel deficit at small ℓ must therefore originate from *global* Ward sources (explicit G–breaking), not from gauge anomalies excluded by Section 7.2.8. □

Notation. We reserve $\Delta_{a,\kappa}$ for the *plateau–suppressed remainder* in the regulator–derivative decomposition (Lemma 7.366), and $\Delta_{\mathrm{ST}}^{(n)}$ for potential Slavnov–Taylor breakings (anomaly candidates) at loop order n.

Definition 7.375 (Soft vs. hard breaking via the $\mathcal{K}$–seminorm)**.** Let $\|\cdot\|_{\mathcal{K}}$ be the seminorm of Definition 7.355. A tagged breaking $\varepsilon_B \hat{\mathcal{O}}_B^{(\lambda_{\mathrm{UV}},\ell)}$ is called *soft* if

$$\varepsilon_B \left\| \hat{\mathcal{O}}_B^{(\lambda_{\mathrm{UV}},\ell)} \right\|_{\mathcal{K}} \le \varepsilon_{\mathrm{soft}} \ll 1$$

(eq. (7.526)); otherwise it is *hard*. Hard breakings—e.g. relevant operators that violate the small form bound or destabilise the spectrum—are *inadmissible* in QIII and excluded. *References:* standard soft–breaking posture in [14, 91, 82].

Definition 7.376 (Spurion ledger and IR restoration)**.** Assign spurion charges to each coupling ε_B so that the composite $\varepsilon_B S_B \hat{\mathcal{O}}_B$ is G–invariant (spurion S_B transforms in the compensating representation). Record the running $\varepsilon_B(\mu)$; *IR restoration* in Step 3 will be certified by $\varepsilon_B(\mu) \to 0$ as $\mu \to 0$ along the RG, consistent with Wilsonian folklore [389] and algebraic renormalisation [117].

Proposition 7.377 (Numerical/regularisation stability)**.** *Implement eq.* (7.522) *using the* same *tags* $(P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}^{(\ell)})$ *that define* $\hat{H}_0$ *in eq.* (7.241)*. If the pipeline* $(\lambda_{\mathrm{UV}},\ell) \mapsto (\tilde{\lambda}_{\mathrm{UV}},\tilde{\ell})$ *is applied as in eq.* (7.67)*, then under Definition 7.356 one has Kato continuity of the dynamics with respect to* $(\varepsilon_B, \lambda_{\mathrm{UV}}, \ell)$ *on the common form domain:*

$$\left\| U_{\mathrm{micro}}(t;\tilde{\lambda}_{\mathrm{UV}},\tilde{\ell}) - U_{\mathrm{micro}}(t;\lambda_{\mathrm{UV}},\ell) \right\| \;\le\; C\left(|t| \sum_B |\varepsilon_B|\, \delta\|\hat{\mathcal{O}}_B\|_{\mathcal{K}} + \delta_{\mathrm{pipe}} \right),$$

where $\delta\|\hat{\mathcal{O}}_B\|_{\mathcal{K}}$ *is the change of the seminorm under the pipeline and* δ_{pipe} *is controlled by the plateau bounds of Section 6.12.4.* Reason. *Use the form–boundedness in Definition 7.375 and Kato–Rellich/Dyson estimates [115]; identical tagging ensures law/state visibilities are comparable under eq.* (7.67)*.*

Proof. Implement eq. (7.522) with the *same* tags $(P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}^{(\ell)})$ that define $\hat{H}_0$ in eq. (7.241). For each $(\lambda_{\mathrm{UV}},\ell)$ set

$$\hat{H}_{\mathrm{micro}}(\lambda_{\mathrm{UV}},\ell) = \hat{H}_0(\lambda_{\mathrm{UV}},\ell) \;+\; \sum_B \varepsilon_B\, \hat{\mathcal{O}}_B^{(\lambda_{\mathrm{UV}},\ell)},$$

with all operators implicitly band/projected by $P_x(\lambda_{\mathrm{UV}})$ and smeared by $K_{\mathrm{IR}}^{(\ell)}$. By Definitions 7.356 and 7.375, each $\hat{\mathcal{O}}_B$ is $\hat{H}_0$-form–bounded on the *common* quadratic–form domain $\mathcal{D}$ (the latter is common because the tags coincide with those of $\hat{H}_0$). Hence for $\sum_B |\varepsilon_B|\, a_B < 1$ (relative bound a_B of $\hat{\mathcal{O}}_B$), Kato–Rellich yields self–adjointness of $\hat{H}_{\mathrm{micro}}(\lambda_{\mathrm{UV}},\ell)$ on a domain with form core $\mathcal{D}$ and a unitary group $U_{\mathrm{micro}}(t;\lambda_{\mathrm{UV}},\ell) = e^{-it\hat{H}_{\mathrm{micro}}(\lambda_{\mathrm{UV}},\ell)}$ [115].

Consider now a pipeline move $(\lambda_{\mathrm{UV}},\ell) \mapsto (\tilde{\lambda}_{\mathrm{UV}},\tilde{\ell})$ as in eq. (7.67). Write the Hamiltonian difference on $\mathcal{D}$ as

$$\begin{aligned} \Delta\hat{H} &:= \hat{H}_{\mathrm{micro}}(\tilde{\lambda}_{\mathrm{UV}},\tilde{\ell}) - \hat{H}_{\mathrm{micro}}(\lambda_{\mathrm{UV}},\ell) \\ &= \underbrace{\left[\hat{H}_0(\tilde{\lambda}_{\mathrm{UV}},\tilde{\ell}) - \hat{H}_0(\lambda_{\mathrm{UV}},\ell)\right]}_{=\,\Delta\hat{H}_0} + \sum_B \varepsilon_B\, \underbrace{\left[\hat{\mathcal{O}}_B^{(\tilde{\lambda}_{\mathrm{UV}},\tilde{\ell})} - \hat{\mathcal{O}}_B^{(\lambda_{\mathrm{UV}},\ell)}\right]}_{=\,\Delta\hat{\mathcal{O}}_B}. \end{aligned}$$

Because the tags (P_x, K_{IR}) are identical in the law/state constructions, the domains and the comparison are "like–for–like".

By the Duhamel (Dyson) formula, for time–independent generators

$$U_{\mathrm{micro}}(t;\tilde{\lambda}_{\mathrm{UV}},\tilde{\ell}) - U_{\mathrm{micro}}(t;\lambda_{\mathrm{UV}},\ell) = -i\int_0^t ds\, U_{\mathrm{micro}}(t-s;\tilde{\lambda}_{\mathrm{UV}},\tilde{\ell})\,\Delta\hat{H}\,U_{\mathrm{micro}}(s;\lambda_{\mathrm{UV}},\ell).$$

Taking operator norms and using unitarity gives

$$\left\|U_{\mathrm{micro}}(t;\tilde{\lambda}_{\mathrm{UV}},\tilde{\ell}) - U_{\mathrm{micro}}(t;\lambda_{\mathrm{UV}},\ell)\right\| \;\le\; |t|\;\|\Delta\hat{H}\|_{B(\mathcal{D}\to\mathcal{D}^*)}\,. \tag{7.536}$$

By Definition 7.375, the form–boundedness constants provide a seminorm $\|\cdot\|_{\mathcal{K}}$ on $\mathcal{D}$ such that, for some $C_1 > 0$,

$$\|\Delta\hat{H}\|_{B(\mathcal{D}\to\mathcal{D}^*)} \;\le\; C_1\left(\sum_B |\varepsilon_B|\,\|\Delta\hat{\mathcal{O}}_B\|_{\mathcal{K}} \;+\; \|\Delta\hat{H}_0\|_{\mathcal{K}}\right).$$

Define $\delta\|\hat{\mathcal{O}}_B\|_{\mathcal{K}} := \|\Delta\hat{\mathcal{O}}_B\|_{\mathcal{K}}$. Within a plateau window (Section 6.12.4) the tag–induced change of the baseline, $\|\Delta\hat{H}_0\|_{\mathcal{K}}$, is bounded by a pipeline tolerance δ_{pipe} controlled by the plateau bounds (uniform Lipschitz control of $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ with respect to $\ln\lambda_{\mathrm{UV}}$, $\ln\ell$; cf. Section 6.12.4). Therefore, for some constant $C > 0$ depending only on the form–bounds in Definition 7.375 (but not on t within compact intervals),

$$\|\Delta\hat{H}\|_{B(\mathcal{D}\to\mathcal{D}^*)} \;\le\; C\left(\sum_B |\varepsilon_B|\,\delta\|\hat{\mathcal{O}}_B\|_{\mathcal{K}} \;+\; \delta_{\mathrm{pipe}}\right).$$

Inserting this into (7.536) yields

$$\left\|U_{\mathrm{micro}}(t;\tilde{\lambda}_{\mathrm{UV}},\tilde{\ell}) - U_{\mathrm{micro}}(t;\lambda_{\mathrm{UV}},\ell)\right\| \;\le\; C\left(|t|\sum_B |\varepsilon_B|\,\delta\|\hat{\mathcal{O}}_B\|_{\mathcal{K}} \;+\; |t|\,\delta_{\mathrm{pipe}}\right).$$

Absorbing the harmless factor of $|t|$ multiplying δ_{pipe} into the overall constant (or using the plateau's uniform-in-t control when $|t| \le T$ is fixed in the protocol) gives the stated bound, i.e.

$$\left\|U_{\mathrm{micro}}(t;\tilde{\lambda}_{\mathrm{UV}},\tilde{\ell}) - U_{\mathrm{micro}}(t;\lambda_{\mathrm{UV}},\ell)\right\| \;\le\; C\left(|t|\sum_B |\varepsilon_B|\,\delta\|\hat{\mathcal{O}}_B\|_{\mathcal{K}} + \delta_{\mathrm{pipe}}\right).$$

This is precisely Kato continuity of the dynamics with respect to $(\varepsilon_B, \lambda_{\mathrm{UV}}, \ell)$ on the common form domain, with constants furnished by the relative form–bounds and the plateau tolerances [115]. □

Corollary 7.378 (Comparability of visibilities under the pipeline)**.** *With the hypothesis of Proposition 7.377, the law/state visibilities computed at $(\lambda_{\mathrm{UV}}, \ell)$ and $(\tilde{\lambda}_{\mathrm{UV}}, \tilde{\ell})$ differ at most by the plateau tolerances of Section 6.12.4, hence are operationally comparable across the pipeline.*

Proof. Let, for a channel $\mathrm{ch} \in \{\mathrm{law}, \mathrm{state}\}$,

$$\Lambda_{\mathrm{ch}}(T;\lambda_{\mathrm{UV}},\ell) := \mathrm{Tr}\Big[U_1^{(\mathrm{ch})\dagger}(T;\lambda_{\mathrm{UV}},\ell)\,U_0^{(\mathrm{ch})}(T;\lambda_{\mathrm{UV}},\ell)\,\rho_0\Big], \qquad \rho_0 = \mathsf{P}_{\mathrm{phys}}\rho_0\mathsf{P}_{\mathrm{phys}},\ \mathrm{gh}(\rho_0) = 0.$$

Consider a pipeline move $(\lambda_{\mathrm{UV}}, \ell) \mapsto (\tilde{\lambda}_{\mathrm{UV}}, \tilde{\ell})$ as in eq. (7.67). By Proposition 7.377, for each branch $a \in \{0, 1\}$,

$$\left\| U_a^{(\mathrm{ch})}(T; \tilde{\lambda}_{\mathrm{UV}}, \tilde{\ell}) - U_a^{(\mathrm{ch})}(T; \lambda_{\mathrm{UV}}, \ell) \right\| \;\le\; C\Big(|T| \sum_B |\varepsilon_B|\, \delta\|\hat{\mathcal{O}}_B\|_{\mathcal{K}} + \delta_{\mathrm{pipe}} \Big),$$

with $\delta\|\hat{\mathcal{O}}_B\|_{\mathcal{K}}$ the change of the seminorm under the pipeline and δ_{pipe} controlled by the plateau bounds of Section 6.12.4.

Write the visibility difference as

$$\begin{aligned}
\Lambda_{\mathrm{ch}}(T; \tilde{\lambda}_{\mathrm{UV}}, \tilde{\ell}) - \Lambda_{\mathrm{ch}}(T; \lambda_{\mathrm{UV}}, \ell) &= \mathrm{Tr}\Big[\Big(U_1^{(\mathrm{ch})\dagger}(T; \tilde{\cdot})\, U_0^{(\mathrm{ch})}(T; \tilde{\cdot}) - U_1^{(\mathrm{ch})\dagger}(T; \cdot)\, U_0^{(\mathrm{ch})}(T; \cdot) \Big) \rho_0 \Big] \\
&= \mathrm{Tr}\Big[\Big(U_1^{(\mathrm{ch})\dagger}(T; \tilde{\cdot}) - U_1^{(\mathrm{ch})\dagger}(T; \cdot) \Big) U_0^{(\mathrm{ch})}(T; \tilde{\cdot})\, \rho_0 \Big] \\
&\quad + \mathrm{Tr}\Big[U_1^{(\mathrm{ch})\dagger}(T; \cdot) \Big(U_0^{(\mathrm{ch})}(T; \tilde{\cdot}) - U_0^{(\mathrm{ch})}(T; \cdot) \Big) \rho_0 \Big],
\end{aligned}$$

where $(\tilde{\cdot})$ abbreviates $(\tilde{\lambda}_{\mathrm{UV}}, \tilde{\ell})$ and $(\cdot)$ abbreviates $(\lambda_{\mathrm{UV}}, \ell)$. Using $\|\mathrm{Tr}(A\rho_0)\| \le \|A\|\, \|\rho_0\|_1 = \|A\|$ and $\|U\| = 1$ for unitaries,

$$\left| \Lambda_{\mathrm{ch}}(T; \tilde{\lambda}_{\mathrm{UV}}, \tilde{\ell}) - \Lambda_{\mathrm{ch}}(T; \lambda_{\mathrm{UV}}, \ell) \right| \;\le\; \left\| U_1^{(\mathrm{ch})}(T; \tilde{\cdot}) - U_1^{(\mathrm{ch})}(T; \cdot) \right\| + \left\| U_0^{(\mathrm{ch})}(T; \tilde{\cdot}) - U_0^{(\mathrm{ch})}(T; \cdot) \right\|.$$

Applying the bound from Proposition 7.377 to each branch and combining,

$$\left| \Lambda_{\mathrm{ch}}(T; \tilde{\lambda}_{\mathrm{UV}}, \tilde{\ell}) - \Lambda_{\mathrm{ch}}(T; \lambda_{\mathrm{UV}}, \ell) \right| \;\le\; 2C\Big(|T| \sum_B |\varepsilon_B|\, \delta\|\hat{\mathcal{O}}_B\|_{\mathcal{K}} + \delta_{\mathrm{pipe}} \Big).$$

Within a plateau window, both $\delta\|\hat{\mathcal{O}}_B\|_{\mathcal{K}}$ and δ_{pipe} are bounded by the prescribed tolerances of Section 6.12.4. Thus the visibilities computed at $(\lambda_{\mathrm{UV}}, \ell)$ and $(\tilde{\lambda}_{\mathrm{UV}}, \tilde{\ell})$ differ by at most a constant multiple of the plateau tolerances, and are therefore operationally comparable across the pipeline. This holds for both ch = law and ch = state (the latter also enjoying the baseline pinning of Proposition 7.370). □

Outcome of Step 1.

Proposition 7.379 (Deliverables of QIII Step 1)**.** *After calibration:*

(i) A fully specified micro–law $\hat{H}_{\mathrm{micro}}(t; \lambda_{\mathrm{UV}}, \ell)$ satisfying admissibility (Definition 7.356) and smallness (Definition 7.355), with BRST/ST consistency (Proposition 7.358).

(ii) Calibrated law– and state–branch unitaries in the sense of eqs. (7.125) and (7.527) (presentation–class invariance by Proposition 7.365).

(iii) Initial records at the minimal coarse–graining scale $\ell_{\min}$:

$$E_{\mathrm{law}}^{(\mathrm{init})}(\ell_{\min}) := 1 - \Lambda_{\mathrm{law}}(\lambda_{\mathrm{UV}}, \ell_{\min}; T^\star) > 0, \qquad \left| 1 - \Lambda_{\mathrm{state}}(\lambda_{\mathrm{UV}}, \ell_{\min}; T^\star) \right| \le \varepsilon_{\mathrm{init}},$$

as in eq. (7.535), with $\varepsilon_{\mathrm{init}}$ below the plateau thresholds of Section 6.12.4.

These outputs seed Step 2 (emergence diagnostics) and Step 3 (RG irrelevance) without *re–tuning phases or instruments; their gauge/presentation independence and plateau robustness follow from Proposition 7.372.*

Proof. Assume the QIII calibration has been performed: the state channel uses identity branches (Definition 7.369 and eq. (7.125)) and the law channel is phase–calibrated at the symmetric baseline so that $\Lambda_{\mathrm{law}} = 1$ for $\varepsilon_B = 0$ (Proposition 7.367), with tags BRST–closed and commuting with projectors (cf. Section 7.1.3).

(1) Existence and consistency of the micro–law. Define the micro–law Hamiltonian as in eq. (7.522), using the *same* tags $(P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}^{(\ell)})$ that specify $\hat{H}_0$ in eq. (7.241). By admissibility (Definition 7.356) the breaking operators $\hat{\mathcal{O}}_B$ are BRST–closed with $\mathrm{gh} = 0$; by the smallness/softness hypothesis (Definition 7.355) they are $\hat{H}_0$-form–bounded with relative bound < 1. Hence $\hat{H}_{\mathrm{micro}}(t; \lambda_{\mathrm{UV}}, \ell)$ is self–adjoint on the common form domain and generates a unitary dynamics; BRST/Slavnov–Taylor consistency follows from Proposition 7.358 (and the anomaly filters of Section 7.2.8), establishing item (i).

(2) Calibrated branch unitaries and presentation–class invariance. Let $U_a^{(\mathrm{law;micro})}(T)$ be defined by eq. (7.527) and $U_a^{(\mathrm{state})}(T)$ by eq. (7.125). By Proposition 7.365, any change of presentation within a class acts by G-unitary conjugations and BRST–exact coboundaries, affecting the interferometer only via a single global phase that has been fixed at baseline; hence the calibrated branch unitaries are well defined on the presentation class, proving item (ii).

(3) Initial records at ℓ_{min}. Choose ℓ_{min} as in Remark 7.373, i.e. at the reciprocal–resolution knee satisfying Proposition 4.436 and lying in a plateau window (Section 6.12.4). Set $\rho_0 = \mathsf{P}_{\mathrm{phys}} \rho_0 \mathsf{P}_{\mathrm{phys}}$ with $\mathrm{gh}(\rho_0) = 0$. Write the law–visibility as a Hilbert–Schmidt inner product

$$\Lambda_{\mathrm{law}}(\lambda_{\mathrm{UV}}, \ell_{\mathrm{min}}; T) = \left\langle U_1^{(\mathrm{law})}(T) \rho_0^{1/2},\, U_0^{(\mathrm{law})}(T) \rho_0^{1/2} \right\rangle_{\mathrm{HS}},$$

so by Cauchy–Schwarz, $\Lambda_{\mathrm{law}} \le \|U_1 \rho_0^{1/2}\|_{\mathrm{HS}} \|U_0 \rho_0^{1/2}\|_{\mathrm{HS}} = \mathrm{Tr}\, \rho_0 = 1$, with equality iff $U_1^{(\mathrm{law})}(T) = e^{i\theta} U_0^{(\mathrm{law})}(T)$ on $\mathrm{supp}\, \rho_0$. In the presence of a nontrivial global Ward source (explicit G-breaking) and after the baseline calibration, the two branches are *not* related by a pure phase for some $T > 0$ (cf. the diagnostic hygiene in Lemma 7.374); by continuity in T, there exists $T^\star > 0$ with

$$E_{\mathrm{law}}^{(\mathrm{init})}(\ell_{\mathrm{min}}) := 1 - \Lambda_{\mathrm{law}}(\lambda_{\mathrm{UV}}, \ell_{\mathrm{min}}; T^\star) > 0.$$

For the state channel, Proposition 7.370 gives $\Lambda_{\mathrm{state}} = 1$ in the symmetric limit; gauge/presentation independence and plateau robustness imply that at ℓ_{min},

$$\left| 1 - \Lambda_{\mathrm{state}}(\lambda_{\mathrm{UV}}, \ell_{\mathrm{min}}; T^\star) \right| \le \varepsilon_{\mathrm{init}},$$

with $\varepsilon_{\mathrm{init}}$ below the plateau thresholds of Section 6.12.4. This establishes item (iii).

Finally, the gauge/presentation independence and plateau robustness of all these deliverables follow from Proposition 7.372 (and the comparability result of Corollary 7.378 when crossing pipeline steps). No re–tuning of phases or instruments is required after calibration, completing the proof. □

7.8.2 Tracking Law Emergence and Defining a Robust IR Slope

Objective. We quantify the *healing of the law* by measuring a *robust* logarithmic-scale slope of the law-visibility Λ_{law} (defined in eq. (7.122)) along coarse–graining length ℓ (with $v := \ln \ell$). The construction below is invariant under $\ell \mapsto a\,\ell$, is stable to measurement ripples via a light low–pass, and is compatible with the acceptance/plateau pipeline of Sections 6.12.4 and 7.3.9. RG consistency of trends is interpreted in the Wilsonian sense [389]; gauge/BRST safety is ensured by the on–shell posture of eq. (7.16) and the Nielsen protection of physical poles/residues [117, 243, 14, 82].

Low–pass (mollified) visibility.

Definition 7.380 (Low–pass operator and mollified visibilities)**.** Let $v = \ln \ell$. The low–pass operator LP_τ with bandwidth $\tau > 0$ acts by convolution

$$\boxed{\big(\mathsf{LP}_\tau f\big)(v) := \int_{\mathbb{R}} \mathrm{d}v'\, W_\tau(v - v')\, f(v'), \qquad W_\tau(v) = \tfrac{1}{\tau}\, W\big(\tfrac{v}{\tau}\big)\,.} \tag{7.537}$$

The kernel $W \in C_c^\infty(\mathbb{R})$ is even and normalised with vanishing first moment,

$$\boxed{\int_{\mathbb{R}} \mathrm{d}v\; W(v) = 1, \qquad \int_{\mathbb{R}} \mathrm{d}v\; v\, W(v) = 0, \qquad W(v) = W(-v).} \tag{7.538}$$

These conditions ensure that LP_τ preserves constants and linear trends in v while suppressing high–frequency ripples.

For any visibility meter $f(v) \in \{\Lambda_{\mathrm{law}}(v),\, \Lambda_{\mathrm{state}}(v)\}$, define its *mollified* (low–pass) version by

$$f^{\langle\tau\rangle}(v) := (\mathsf{LP}_\tau f)(v) = \int_{\mathbb{R}} \mathrm{d}v'\, W_\tau(v - v')\, f(v').$$

We abbreviate $\Lambda_{\mathrm{law}}^{\langle\tau\rangle} := \mathsf{LP}_\tau \Lambda_{\mathrm{law}}$ and $\Lambda_{\mathrm{state}}^{\langle\tau\rangle} := \mathsf{LP}_\tau \Lambda_{\mathrm{state}}$.

Clarification. The low–pass kernel W acts on the log–scale variable $v = \ln \ell$ and is independent of the spacetime IR window $K_{\mathrm{IR}}^{(\ell)}$ used in the joint coarse-graining of fields and correlators (see eqs. (6.364) and (6.366)). The former is a 1-D mollifier in scale; the latter is a geometric kernel on $\mathcal{X}_q$.

Intuition. Because W is even, normalized, and has vanishing first moment (7.538), LP_τ smooths ripples while exactly preserving constants and linear trends in v. This makes it ideal for plateau diagnostics that rely on first–derivative flatness.

Lemma 7.381 (Basic properties of LP_τ)**.** *Let $W \in C_c^\infty(\mathbb{R})$ satisfy eq. (7.538), and set $W_\tau(v) = \tau^{-1} W(v/\tau)$. With $\mathsf{LP}_\tau f := W_\tau * f$ as in Definition 7.380, the following hold:*

(i) Linearity and boundedness.
For each $1 \le p \le \infty$, $\mathsf{LP}_\tau : L^p(\mathbb{R}) \to L^p(\mathbb{R})$ is bounded with

$$\|\mathsf{LP}_\tau\|_{L^p \to L^p} \le \|W_\tau\|_{L^1} = \|W\|_{L^1}.$$

If $W \ge 0$ and $\int_{\mathbb{R}} W = 1$ (the zeroth-moment condition in eq. (7.538)), then $\|\mathsf{LP}_\tau\|_{L^p \to L^p} = 1$.

(ii) Self–adjointness on L^2.
If W is even, then LP_τ is self–adjoint on $L^2(\mathbb{R})$.

(iii) Reproduction of affine functions.
For all $a, b \in \mathbb{R}$,

$$\mathsf{LP}_\tau(a + b\,v) \;=\; a + b\,v.$$

(iv) Commutation with translations/derivatives.
If $f \in W^{1,p}_{\mathrm{loc}}(\mathbb{R})$, then

$$\frac{\mathrm{d}}{\mathrm{d}v}\Big(\mathsf{LP}_\tau f\Big) = \mathsf{LP}_\tau(f'), \qquad \mathsf{LP}_\tau\Big(f(\cdot + v_0)\Big) = \Big(\mathsf{LP}_\tau f\Big)(\cdot + v_0).$$

(v) Approximation identity.
If $f \in C^0(\mathbb{R})$ is uniformly continuous, then $\|\mathsf{LP}_\tau f - f\|_\infty \to 0$ as $\tau \to 0^+$.

Proof. **(1) Linearity and boundedness.** Linearity is immediate from the integral definition in Definition 7.380. For $1 \le p \le \infty$, Young's convolution inequality gives

$$\|W_\tau * f\|_{L^p} \le \|W_\tau\|_{L^1}\,\|f\|_{L^p},$$

so $\|\mathsf{LP}_\tau\|_{L^p \to L^p} \le \|W_\tau\|_{L^1}$. By the scaling $W_\tau(v) = \tau^{-1} W(v/\tau)$ we have $\|W_\tau\|_{L^1} = \|W\|_{L^1}$. If, in addition, $W \ge 0$ and $\int W = 1$ (zeroth moment in eq. (7.538)), then $\|W\|_{L^1} = 1$, hence $\|\mathsf{LP}_\tau\| \le 1$. Sharpness $\|\mathsf{LP}_\tau\| = 1$ follows by testing on approximate constants (e.g. normalized indicators of large intervals when $1 \le p < \infty$, or $f \equiv 1$ for $p = \infty$), using compact support of W to see the ratio $\|W_\tau * f\|_p / \|f\|_p \to 1$ as the interval length $\to \infty$ (standard convolution theory; see, e.g., [55, 446, 253, 433]).

(2) Self–adjointness on L^2. If W is even, then $W_\tau(v - v') = W_\tau(v' - v)$. For $f, g \in L^2$,

$$\langle \mathsf{LP}_\tau f, g\rangle = \int_{\mathbb{R}}\!\int_{\mathbb{R}} W_\tau(v - v')\, f(v')\, \overline{g(v)}\,\mathrm{d}v'\mathrm{d}v = \int_{\mathbb{R}}\!\int_{\mathbb{R}} W_\tau(v' - v)\, f(v')\, \overline{g(v)}\,\mathrm{d}v\mathrm{d}v' = \langle f, \mathsf{LP}_\tau g\rangle,$$

so LP_τ is self–adjoint. (Equivalently, $\widehat{W_\tau}$ is real and nonnegative, hence the Fourier multiplier is self–adjoint; cf. [55, 446].)

(3) Reproduction of affine functions. Let $f(v) = a + bv$. Using $\int W_\tau = 1$ and the vanishing first moment $\int v\, W(v)\,\mathrm{d}v = 0$ from eq. (7.538) (which is equivalent to $\int (v - v') W_\tau(v - v')\,\mathrm{d}v' = 0$ for each v), we compute

$$(W_\tau * 1)(v) = \int W_\tau(v - v')\,\mathrm{d}v' = 1, \qquad (W_\tau * v)(v) = \int v'\, W_\tau(v - v')\,\mathrm{d}v' = v.$$

Hence $\mathsf{LP}_\tau(a + bv) = a + bv$.

(4) Commutation with translations and derivatives. Translation invariance is standard for convolutions:

$$\Big(\mathsf{LP}_\tau(f(\cdot + v_0))\Big)(v) = \int W_\tau(v - v')\, f(v' + v_0)\,\mathrm{d}v' = (\mathsf{LP}_\tau f)(v + v_0).$$

For derivatives, if $f \in W^{1,p}_{\mathrm{loc}}$, the distributional derivative f' is locally L^p, and $W_\tau \in L^1$ with compact support. Differentiating under the integral is justified by dominated convergence and Fubini/Tonelli (all integrands are integrable on compact sets), giving

$$\frac{\mathrm{d}}{\mathrm{d}v}(W_\tau * f)(v) = \int \frac{\partial}{\partial v} W_\tau(v - v')\, f(v')\,\mathrm{d}v' = \int W_\tau(v - v')\, f'(v')\,\mathrm{d}v' = (W_\tau * f')(v).$$

A classical reference for these interchange rules is, e.g., [55, Chs. 1–2] or [446, §1.3].

(5) Approximation identity. The family $\{W_\tau\}_{\tau>0}$ is an approximate identity: $W_\tau \geq 0$, $\int W_\tau = 1$, and $\operatorname{supp} W_\tau \to \{0\}$ as $\tau \to 0^+$. If $f \in C^0(\mathbb{R})$ is uniformly continuous, then

$$\|W_\tau * f - f\|_\infty = \sup_v \left| \int W_\tau(z)\big(f(v-z) - f(v)\big)\,\mathrm{d}z \right| \leq \int W_\tau(z)\,\omega_f(|z|)\,\mathrm{d}z \xrightarrow[\tau\to 0^+]{} 0,$$

where ω_f is the modulus of continuity of f. This is the standard uniform convergence of mollifier averages; see, e.g., [446, §1.3] or [55, Ch. 1]. (Quantitative rates in terms of ω_f follow immediately.) □

Proposition 7.382 (Second–order accuracy and derivative neutrality)**.** *If $f \in C^2(\mathbb{R})$ and $m_2(W) := \int_{\mathbb{R}} v^2 W(v)\,\mathrm{d}v < \infty$, then for each v*

$$\big(\mathsf{LP}_\tau f\big)(v) = f(v) + \frac{\tau^2 m_2(W)}{2}\, f''(v) + \mathcal{O}(\tau^4),$$
$$\frac{\mathrm{d}}{\mathrm{d}v}\big(\mathsf{LP}_\tau f\big)(v) = f'(v) + \frac{\tau^2 m_2(W)}{2}\, f^{(3)}(v) + \mathcal{O}(\tau^4).$$

In particular, constants and linear trends are preserved exactly, and the first derivative is unbiased up to $\mathcal{O}(\tau^2)$.

Proof. Write the convolution with the change of variables $z = v - v' = \tau u$:

$$(\mathsf{LP}_\tau f)(v) = \int_{\mathbb{R}} W_\tau(z)\, f(v-z)\,\mathrm{d}z = \int_{\mathbb{R}} W(u)\, f(v-\tau u)\,\mathrm{d}u,$$

using $W_\tau(z) = \tau^{-1} W(z/\tau)$ from Definition 7.380. Since $f \in C^2$ and $W \in C_c^\infty$, Taylor's theorem with integral remainder at order 2 gives

$$f(v-\tau u) = f(v) - \tau u f'(v) + \tfrac{\tau^2 u^2}{2} f''(v) + R_4(v,u;\tau),$$

where, for some $\theta \in (0,1)$ depending on v, u, τ,

$$R_4(v,u;\tau) = \frac{\tau^3 u^3}{3!}\, f^{(3)}(v - \theta\tau u).$$

(We will refine this remainder below to expose the $\mathcal{O}(\tau^4)$ behavior using odd-moment cancellations.)

Zeroth and first moments; preservation of affine trends. Integrating termwise and using the moment conditions from eq. (7.538) (namely $\int W = 1$ and $\int u\,W(u)\,\mathrm{d}u = 0$) yields

$$(\mathsf{LP}_\tau f)(v) = f(v) + \frac{\tau^2}{2}\, f''(v) \int u^2 W(u)\,\mathrm{d}u \;+\; \int W(u)\, R_4(v,u;\tau)\,\mathrm{d}u.$$

Thus the constant and linear parts are reproduced exactly (see also Lemma 7.381(3)), and the explicit quadratic bias is $\frac{\tau^2 m_2(W)}{2} f''(v)$.

Odd–moment cancellation and $\mathcal{O}(\tau^4)$ remainder. To sharpen the remainder, expand $f^{(3)}(v - \theta\tau u) = f^{(3)}(v) + \mathcal{O}(\tau|u|)$ as $\tau \to 0^+$ (uniformly for u in $\operatorname{supp} W$), since $f^{(3)}$ is continuous wherever the second display is asserted. Plugging into R_4 and integrating:

$$\int W(u)\, R_4(v,u;\tau)\,\mathrm{d}u = \frac{\tau^3}{3!}\, f^{(3)}(v) \int u^3 W(u)\,\mathrm{d}u \;+\; \mathcal{O}\Big(\tau^4 \int |u|^4 |W(u)|\,\mathrm{d}u\Big).$$

The odd moment vanishes (either by the explicit $\int u^3 W = 0$ in eq. (7.538) or by evenness of W, used elsewhere for LP_τ in Lemma 7.381(2)), hence the leading term is zero and the remainder is $\mathcal{O}(\tau^4)$. Since $W \in C_c^\infty$, $\int |u|^4 |W(u)|\,\mathrm{d}u < \infty$, so the constant in $\mathcal{O}(\tau^4)$ depends on $\|f^{(3)}\|_{C^0(K_v)}$ on a small neighborhood K_v of v and the fourth moment of W (hidden in the big-O).

Altogether,

$$(\mathsf{LP}_\tau f)(v) = f(v) + \frac{\tau^2 m_2(W)}{2}\, f''(v) + \mathcal{O}(\tau^4).$$

Derivative neutrality. By Lemma 7.381(4), $\frac{\mathrm{d}}{\mathrm{d}v}(\mathsf{LP}_\tau f) = \mathsf{LP}_\tau(f')$. Apply the previous expansion to f' (which is C^2 wherever the second display is asserted), obtaining

$$\frac{\mathrm{d}}{\mathrm{d}v}\Big(\mathsf{LP}_\tau f\Big)(v) = (\mathsf{LP}_\tau f')(v) = f'(v) + \frac{\tau^2 m_2(W)}{2}\, f^{(3)}(v) + \mathcal{O}(\tau^4).$$

"In particular" clause. Setting $f(v) = a + bv$ gives $(\mathsf{LP}_\tau f)(v) = a + bv$ exactly by Lemma 7.381(3). The derivative statement shows that the slope estimator is unbiased up to $\mathcal{O}(\tau^2)$.

Remarks on regularity and references. We used the local boundedness/continuity of $f^{(3)}$ to promote the Taylor remainder to an $\mathcal{O}(\tau^4)$ term via odd–moment cancellation; with only $f \in C^2$, one still gets the displayed τ^2 bias and a smaller-order remainder $o(\tau^2)$ from approximate-identity arguments (cf. Lemma 7.381(5)). Standard convolution/Taylor moment expansions can be found in [55, Chs. 1–2] and [446, §1.3]. □

Proposition 7.383 (Compatibility with spectral/trace constructions)**.** *Let* $f(v) = \Lambda_{\mathrm{law}}(v) = \mathrm{Tr}\Big[U_1^{(\mathrm{law})\dagger}(T; e^v)\, U_0^{(\mathrm{law})}(T; e^v)\, \rho_0\Big]$ *with* $\rho_0 = \mathsf{P}_{\mathrm{phys}} \rho_0 \mathsf{P}_{\mathrm{phys}}$, $\mathrm{gh}(\rho_0) = 0$. *Assume the Kato continuity of the branch dynamics in* $v = \ln \ell$ *from Proposition 7.377. Then the Bochner integral*

$$\mathcal{U}(v) := \int_{\mathbb{R}} \mathrm{d}v'\; W_\tau(v - v')\, U_1^{(\mathrm{law})\dagger}(T; e^{v'})\, U_0^{(\mathrm{law})}(T; e^{v'})$$

exists in operator norm with $\|\mathcal{U}(v)\| \le \|W\|_1$, *and*

$$(\mathsf{LP}_\tau f)(v) = \mathrm{Tr}\Big[\mathcal{U}(v)\, \rho_0\Big].$$

Consequently, smoothing commutes with the trace/spectral construction of Λ_{law}, *and the same holds for* Λ_{state}.

Proof. Step 1: Operator–valued integrand and norm–continuity. By Proposition 7.377 the maps $v \mapsto U_a^{(\mathrm{law})}(T; e^v)$ are norm–continuous (Kato stability of the propagators). Therefore

$$A(v') \;:=\; U_1^{(\mathrm{law})\dagger}(T; e^{v'})\, U_0^{(\mathrm{law})}(T; e^{v'}) \in \mathcal{B}(\mathcal{H}_{\mathrm{em}})$$

is norm–continuous in v' and uniformly bounded by 1 because each factor is unitary. Hence $A(\cdot)$ is (strongly) measurable and $\|A(v')\| \le 1$ for all v' [115, Chap. IV].

Step 2: Existence and bound of the Bochner integral. With $W_\tau \in L^1(\mathbb{R})$ (see Definition 7.380 and eq. (7.538)) and $\|A(v')\| \le 1$,

$$\int_{\mathbb{R}} \| W_\tau(v - v')\, A(v') \|\, \mathrm{d}v' \;\le\; \int_{\mathbb{R}} |W_\tau(v - v')|\, \mathrm{d}v' \;=\; \|W_\tau\|_1 = \|W\|_1 < \infty,$$

so the Bochner integral

$$\mathcal{U}(v) \;:=\; \int_{\mathbb{R}} W_\tau(v-v')\,A(v')\,\mathrm{d}v'$$

exists in operator norm and satisfies $\|\mathcal{U}(v)\| \le \|W\|_1$. (Standard Bochner integration in Banach spaces; see, e.g., [433, §2.6].)

Step 3: Exchange of trace and Bochner integral. Let $\mathcal{S}_1(\mathcal{H}_{\mathrm{em}})$ denote the trace class. Define the bounded linear functional

$$\mathsf{Tr}_{\rho_0} : \mathcal{B}(\mathcal{H}_{\mathrm{em}}) \to \mathbb{C}, \qquad \mathsf{Tr}_{\rho_0}(B) := \mathrm{Tr}(B\,\rho_0).$$

By duality $\mathcal{B}(\mathcal{H}) = \mathcal{S}_1(\mathcal{H})^*$ (see [253, Thm. VI.26]), $|\mathsf{Tr}_{\rho_0}(B)| \le \|B\|\,\|\rho_0\|_{\mathcal{S}_1}$. Since $A(\cdot)$ is Bochner–integrable and Tr_{ρ_0} is continuous linear, we may pull Tr_{ρ_0} through the integral:

$$\mathsf{Tr}_{\rho_0}\Big(\int W_\tau(v-v')\,A(v')\,\mathrm{d}v'\Big) = \int W_\tau(v-v')\,\mathsf{Tr}_{\rho_0}\Big(A(v')\Big)\,\mathrm{d}v'.$$

Therefore,

$$\mathrm{Tr}\Big[\mathcal{U}(v)\,\rho_0\Big] = \int_{\mathbb{R}} W_\tau(v-v')\;\mathrm{Tr}\Big[A(v')\,\rho_0\Big]\,\mathrm{d}v' = (\mathsf{LP}_\tau f)(v),$$

which is precisely the desired identity (equivalently, Fubini/Tonelli for Bochner integrals paired with a bounded functional). *Notation.* We use Tr for the operator trace and $\mathsf{Tr}_{\rho_0}(B) := \mathrm{Tr}(B\rho_0)$ for the trace–pairing functional.

Step 4: "Spectral/trace" compatibility and the state channel. All steps above use only (i) Kato norm–continuity of the propagators, (ii) L^1–integrability of W_τ, and (iii) the trace–class pairing with $\rho_0 = \mathsf{P}_{\mathrm{phys}}\rho_0\mathsf{P}_{\mathrm{phys}}$ (cf. Section 7.1.3). Hence smoothing by LP_τ commutes with the trace construction of Λ_{law}. The same argument applies verbatim to Λ_{state} upon replacing $U_a^{(\mathrm{law})}$ by $U_a^{(\mathrm{state})}$ (and, in Quadrant II, still within the BRST–consistent reading).

This proves the proposition. □

Corollary 7.384 (Plateau preservation under low–pass)**.** *Let W have support in $[-R,R]$ and LP_τ be as in Definition 7.380. If f satisfies the plateau criterion on an interval $\mathcal{I}$ (cf. Section 6.12.4), then on the eroded window $\mathcal{I}^{-\tau R} := \{v \in \mathcal{I} : [v-\tau R, v+\tau R] \subset \mathcal{I}\}$ one has*

$$\sup_{v\in\mathcal{I}^{-\tau R}} \big|(\mathsf{LP}_\tau f)'(v)\big| \;\le\; \|W\|_1 \sup_{v\in\mathcal{I}} |f'(v)|.$$

In particular, for $f = \Lambda_{\mathrm{law}}, \Lambda_{\mathrm{state}}$ the derivative remains within the same plateau tolerance on $\mathcal{I}^{-\tau R}$; i.e. low–pass smoothing suppresses ripples without spoiling first–derivative flatness.

Proof. Step 1: Derivative commutation. By Lemma 7.381(4),

$$(\mathsf{LP}_\tau f)' \;=\; \mathsf{LP}_\tau(f') \;=\; W_\tau * f'.$$

Step 2: Young's inequality and the L^∞ bound. By Lemma 7.381(1) (Young's inequality) and $\|W_\tau\|_1 = \|W\|_1$ (Definition 7.380),

$$\|(\mathsf{LP}_\tau f)'\|_{L^\infty} = \|W_\tau * f'\|_{L^\infty} \le \|W_\tau\|_{L^1}\,\|f'\|_{L^\infty} = \|W\|_{L^1}\,\|f'\|_{L^\infty}.$$

Step 3: Restriction to the eroded plateau window. Since $\operatorname{supp} W \subset [-R,R]$, we have $\operatorname{supp} W_\tau \subset [-\tau R, \tau R]$. Thus for any $v \in \mathcal{I}^{-\tau R}$ the integrand in $(W_\tau * f')(v)$ samples f' only on $\mathcal{I}$. Hence

$$\sup_{v\in\mathcal{I}^{-\tau R}} \bigl|(\mathsf{LP}_\tau f)'(v)\bigr| = \sup_{v\in\mathcal{I}^{-\tau R}} \left| \int_{\mathbb{R}} W_\tau(v-v')\, f'(v')\,\mathrm{d}v' \right| \le \|W\|_1 \sup_{v'\in\mathcal{I}} |f'(v')|.$$

Step 4: Application to Λ_{law} and Λ_{state}. By the plateau criterion (cf. Section 6.12.4 and the acceptance region),

$$\sup_{v\in\mathcal{I}} |\Lambda'_{\mathrm{law}}(v)| \le \varepsilon_{\mathrm{grad}}, \qquad \sup_{v\in\mathcal{I}} |\Lambda'_{\mathrm{state}}(v)| \le \varepsilon_{\mathrm{grad}}.$$

Combining with Step 3 yields

$$\sup_{v\in\mathcal{I}^{-\tau R}} \bigl|(\mathsf{LP}_\tau\Lambda_{\mathrm{law}})'(v)\bigr| \le \|W\|_1\,\varepsilon_{\mathrm{grad}}, \qquad \sup_{v\in\mathcal{I}^{-\tau R}} \bigl|(\mathsf{LP}_\tau\Lambda_{\mathrm{state}})'(v)\bigr| \le \|W\|_1\,\varepsilon_{\mathrm{grad}}.$$

In particular, if $W \ge 0$ and $\int W = 1$, then $\|W\|_1 = 1$ and the derivative bounds on $\mathcal{I}^{-\tau R}$ equal $\varepsilon_{\mathrm{grad}}$. □

Robust IR slope (definition).

Definition 7.385 (Robust IR law slope)**.** Let $v = \ln \ell$ and let LP_τ be the low–pass operator of Definition 7.380, defined by (7.537) with kernel moments (7.538). For the law–channel visibility $\Lambda_{\mathrm{law}}(v)$ (suppressing any fixed T), define the *robust IR law slope* by

$$\boxed{\varsigma^{(\tau)}_{\mathrm{law}}(\ell) := \frac{\partial}{\partial \ln \ell}\Bigl[\bigl(\mathsf{LP}_\tau\,\Lambda_{\mathrm{law}}\bigr)(\ell)\Bigr] = \frac{\mathrm{d}}{\mathrm{d}v}\Bigl[\bigl(\mathsf{LP}_\tau\,\Lambda_{\mathrm{law}}\bigr)(v)\Bigr].} \tag{7.539}$$

Intuition. LP_τ removes high–frequency ripples in $v = \ln \ell$ while preserving constants and linear trends (Lemma 7.381). Differentiating $\mathsf{LP}_\tau\Lambda_{\mathrm{law}}$ thus yields a stable estimator of the IR slope that is insensitive to small regulator wiggles but responsive to genuine drift off a plateau.

Lemma 7.386 (Well-posedness and convolution form)**.** *If Λ_{law} is locally absolutely continuous in v (which holds under the Kato continuity of Proposition 7.377 together with the trace exchange of Proposition 7.383), then*

$$\varsigma^{(\tau)}_{\mathrm{law}}(v) = \bigl(\mathsf{LP}_\tau\,\Lambda'_{\mathrm{law}}\bigr)(v) = \int_{\mathbb{R}} W_\tau(v-v')\,\Lambda'_{\mathrm{law}}(v')\,\mathrm{d}v'. \tag{7.540}$$

Moreover, $\|\varsigma^{(\tau)}_{\mathrm{law}}\|_\infty \le \|W\|_1\,\|\Lambda'_{\mathrm{law}}\|_\infty$.

Proof. Preliminaries. Recall $\mathsf{LP}_\tau f := W_\tau * f$ from Definition 7.380 and the robust slope definition $\varsigma^{(\tau)}_{\mathrm{law}} := \frac{\mathrm{d}}{\mathrm{d}v}(\mathsf{LP}_\tau\Lambda_{\mathrm{law}})$ from Definition 7.385. Assume $v \mapsto \Lambda_{\mathrm{law}}(v)$ is locally absolutely continuous. Then $\Lambda_{\mathrm{law}} \in W^{1,1}_{\mathrm{loc}}(\mathbb{R})$ and $\Lambda'_{\mathrm{law}} \in L^1_{\mathrm{loc}}(\mathbb{R})$.

Step 1: Differentiation commutes with LP_τ. By Lemma 7.381(4), if $f \in W^{1,p}_{\mathrm{loc}}(\mathbb{R})$ for some $1 \le p \le \infty$, then

$$\frac{\mathrm{d}}{\mathrm{d}v}\big(\mathsf{LP}_\tau f\big) = \mathsf{LP}_\tau(f') \quad \text{a.e. on } \mathbb{R}.$$

Applying this with $f = \Lambda_{\mathrm{law}}$ gives

$$\frac{\mathrm{d}}{\mathrm{d}v}\big(\mathsf{LP}_\tau \Lambda_{\mathrm{law}}\big)(v) = \big(\mathsf{LP}_\tau \Lambda'_{\mathrm{law}}\big)(v),$$

which is the integral representation in (7.540).

$$\frac{\mathrm{d}}{\mathrm{d}v}\big(\mathsf{LP}_\tau \Lambda_{\mathrm{law}}\big)(v) = \big(\mathsf{LP}_\tau \Lambda'_{\mathrm{law}}\big)(v).$$

By definition of $\varsigma^{(\tau)}_{\mathrm{law}}$ (see Definition 7.385), the left-hand side equals $\varsigma^{(\tau)}_{\mathrm{law}}(v)$.

Step 2: Convolution (integral) form. Using the definition of LP_τ and Fubini/Tonelli (here justified because $W_\tau \in L^1$ and $\Lambda'_{\mathrm{law}} \in L^1_{\mathrm{loc}}$ while W has compact support),

$$\big(\mathsf{LP}_\tau \Lambda'_{\mathrm{law}}\big)(v) = (W_\tau * \Lambda'_{\mathrm{law}})(v) = \int_{\mathbb{R}} W_\tau(v - v')\, \Lambda'_{\mathrm{law}}(v')\, \mathrm{d}v',$$

which is (7.540). Equivalently, one can obtain the same identity by differentiating under the integral in the representation $(\mathsf{LP}_\tau \Lambda_{\mathrm{law}})(v) = \int W_\tau(v - v')\Lambda_{\mathrm{law}}(v')\,\mathrm{d}v'$ (valid by Proposition 7.383) and using dominated convergence on compact v-intervals.

Step 3: Sup–norm bound. Young's convolution inequality (see Lemma 7.381(1)) gives, for any essentially bounded g, $\|W_\tau * g\|_{L^\infty} \le \|W_\tau\|_{L^1}\, \|g\|_{L^\infty}$. Applying this with $g = \Lambda'_{\mathrm{law}}$ yields

$$\|\varsigma^{(\tau)}_{\mathrm{law}}\|_\infty = \|\mathsf{LP}_\tau \Lambda'_{\mathrm{law}}\|_\infty \le \|W_\tau\|_1\, \|\Lambda'_{\mathrm{law}}\|_\infty = \|W\|_1\, \|\Lambda'_{\mathrm{law}}\|_\infty,$$

using $\|W_\tau\|_1 = \|W\|_1$ from Definition 7.380.

Step 4: Well-posedness via the spectral/trace construction (context). Under Proposition 7.377, the maps $v \mapsto U^{(\mathrm{law})}_a(T; e^v)$ are norm-continuous. By Proposition 7.383, smoothing commutes with the physical trace, so $(\mathsf{LP}_\tau \Lambda_{\mathrm{law}})(v) = \mathrm{Tr}[\mathcal{U}(v)\rho_0]$ for a Bochner-integrable $\mathcal{U}(v)$. These two ingredients ensure the local absolute continuity of Λ_{law} invoked at the start (and thus justify applying Lemma 7.381(4)).

This proves the lemma. □

Proposition 7.387 (Scale invariance and dimensionless character)**.** *Under rescaling* $\ell \mapsto a\,\ell$ *(i.e.* $v \mapsto v + \ln a$*), one has*

$$\boxed{\Lambda_{\mathrm{law}}(v) \;\mapsto\; \Lambda_{\mathrm{law}}(v + \ln a) \quad\Longrightarrow\quad \varsigma^{(\tau)}_{\mathrm{law}}(v) \;\mapsto\; \varsigma^{(\tau)}_{\mathrm{law}}(v + \ln a),} \tag{7.541}$$

hence $\varsigma^{(\tau)}_{\mathrm{law}}$ *is dimensionless.*

Proof. Setup and notation. Let $c := \ln a$ and write the translation operator T_c by $(T_c f)(v) := f(v + c)$. Recall $\mathsf{LP}_\tau f = W_\tau * f$ (Definition 7.380) and the robust slope $\varsigma^{(\tau)}_{\mathrm{law}} = \frac{\mathrm{d}}{\mathrm{d}v}(\mathsf{LP}_\tau \Lambda_{\mathrm{law}})$ (Definition 7.385).

Step 1: Low–pass commutes with translations. For any f,

$$(\mathsf{LP}_\tau T_c f)(v) = \int_{\mathbb{R}} W_\tau(v - v')\, f(v' + c)\, \mathrm{d}v' = \int_{\mathbb{R}} W_\tau\big(v - (u + c)\big)\, f(u)\, \mathrm{d}u = (T_c \mathsf{LP}_\tau f)(v),$$

using the change of variables $u = v' - c$. This is $\mathsf{LP}_\tau T_c = T_c \mathsf{LP}_\tau$, i.e. the translation covariance in Lemma 7.381(4).

Step 2: Derivatives commute with translations. For any differentiable g, $\frac{\mathrm{d}}{\mathrm{d}v}(T_c g) = T_c\big(\frac{\mathrm{d}}{\mathrm{d}v} g\big)$, since constant shifts of the argument commute with differentiation.

Step 3: Covariance of $\varsigma^{(\tau)}_{\mathrm{law}}$. Under $\ell \mapsto a\ell$ we have $\Lambda_{\mathrm{law}} \mapsto T_c \Lambda_{\mathrm{law}}$. Define the rescaled slope by

$$\varsigma^{(\tau)}_{\mathrm{law},a}(v) := \frac{\mathrm{d}}{\mathrm{d}v}\Big(\mathsf{LP}_\tau(T_c \Lambda_{\mathrm{law}})\Big)(v).$$

Using Steps 1–2,

$$\varsigma^{(\tau)}_{\mathrm{law},a}(v) = \frac{\mathrm{d}}{\mathrm{d}v}\Big(T_c \mathsf{LP}_\tau \Lambda_{\mathrm{law}}\Big)(v) = T_c\left(\frac{\mathrm{d}}{\mathrm{d}v} \mathsf{LP}_\tau \Lambda_{\mathrm{law}}\right)(v) = (T_c\, \varsigma^{(\tau)}_{\mathrm{law}})(v) = \varsigma^{(\tau)}_{\mathrm{law}}(v + c).$$

Thus (7.541) holds.

Step 4: Dimensionless character. The coordinate $v = \ln \ell$ is dimensionless, so $\frac{\mathrm{d}}{\mathrm{d}v}$ is dimensionless. The kernel W_τ is normalized with $\int W_\tau = 1$ and lives in the dimensionless v-domain, and Λ_{law} is a (dimensionless) visibility. Hence $\varsigma^{(\tau)}_{\mathrm{law}} = \frac{\mathrm{d}}{\mathrm{d}v}(\mathsf{LP}_\tau \Lambda_{\mathrm{law}})$ is itself dimensionless. Equivalently, writing $\frac{\mathrm{d}}{\mathrm{d}v} = \ell \frac{\mathrm{d}}{\mathrm{d}\ell}$ shows that under $\ell \mapsto a\ell$ the factor of ℓ compensates the scale of $\frac{\mathrm{d}}{\mathrm{d}\ell}$, leaving a pure, unitless quantity; Step 3 then asserts it transforms by translation $v \mapsto v + \ln a$, as claimed. □

Remark 7.388 (Convolution form makes the scale covariance immediate)**.** Let T_c denote translation by $c \in \mathbb{R}$, $(T_c f)(v) := f(v + c)$, and set $c = \ln a$ for the rescaling $\ell \mapsto a\ell$ (i.e. $v \mapsto v + \ln a$). By Lemma 7.386, $\varsigma^{(\tau)}_{\mathrm{law}} = W_\tau * \Lambda'_{\mathrm{law}}$. Then

$$\varsigma^{(\tau)}_{\mathrm{law},a} := W_\tau * (T_c \Lambda_{\mathrm{law}})' = W_\tau * T_c(\Lambda'_{\mathrm{law}}) = T_c\Big(W_\tau * \Lambda'_{\mathrm{law}}\Big) = T_c\, \varsigma^{(\tau)}_{\mathrm{law}},$$

where we used $(T_c \Lambda_{\mathrm{law}})' = T_c(\Lambda'_{\mathrm{law}})$ and the translation–covariance of convolution $W_\tau * T_c = T_c * W_\tau$ (cf. Lemma 7.381(4)). Equivalently,

$$\varsigma^{(\tau)}_{\mathrm{law}}(v) \;\mapsto\; \varsigma^{(\tau)}_{\mathrm{law}}(v + \ln a),$$

which is precisely the covariance stated in Proposition 7.387. No evenness or higher moment of W is required for this covariance—only $W_\tau \in L^1$ and translation invariance of convolution.

Proposition 7.389 (Plateau neutrality and accuracy)**.** *If Λ_{law} is computed within a plateau window (Section 6.12.4), then*

$$\sup_{v \in \mathit{window}} \big|\varsigma^{(\tau)}_{\mathrm{law}}(v)\big| \le \|W\|_1 \sup_{v \in \mathit{window}} |\Lambda'_{\mathrm{law}}(v)|,$$

so $\varsigma^{(\tau)}_{\mathrm{law}}(v) = 0$ up to the same plateau tolerances (Corollary 7.384). If, in addition, $\Lambda_{\mathrm{law}} \in C^3$ in a neighborhood of v and the kernel satisfies the moment conditions of eq. (7.538) (in particular $\int W = 1$, $\int u\, W(u)\, du = 0$, and $\int u^3 W(u)\, du = 0$) with $m_2(W) < \infty$, then

$$\varsigma^{(\tau)}_{\mathrm{law}}(v) = \Lambda'_{\mathrm{law}}(v) + \frac{\tau^2 m_2(W)}{2}\, \Lambda^{(3)}_{\mathrm{law}}(v) + \mathcal{O}(\tau^4),$$

so the first–derivative bias is $\mathcal{O}(\tau^2)$ (by Proposition 7.382).

Proof. Step 1 (slope as low–pass of the derivative). By Lemma 7.386, $\varsigma^{(\tau)}_{\mathrm{law}} = \mathsf{LP}_\tau(\Lambda'_{\mathrm{law}}) = W_\tau * \Lambda'_{\mathrm{law}}$.

Step 2 (plateau bound). Apply Young's inequality as in Corollary 7.384 to obtain $\sup_{\mathrm{window}} |\varsigma^{(\tau)}_{\mathrm{law}}| \le \|W\|_1 \sup_{\mathrm{window}} |\Lambda'_{\mathrm{law}}|$.

Step 3 (second–order accuracy). Under the stated smoothness and moment conditions, Proposition 7.382 with $f = \Lambda_{\mathrm{law}}$ yields

$$(\mathsf{LP}_\tau \Lambda_{\mathrm{law}})(v) = \Lambda_{\mathrm{law}}(v) + \frac{\tau^2 m_2(W)}{2}\,\Lambda''_{\mathrm{law}}(v) + \mathcal{O}(\tau^4).$$

Differentiate in v and use Lemma 7.381(4) to commute the derivative with LP_τ:

$$\varsigma^{(\tau)}_{\mathrm{law}}(v) = \frac{d}{dv}(\mathsf{LP}_\tau \Lambda_{\mathrm{law}})(v) = \Lambda'_{\mathrm{law}}(v) + \frac{\tau^2 m_2(W)}{2}\,\Lambda^{(3)}_{\mathrm{law}}(v) + \mathcal{O}(\tau^4).$$

This is the claimed bias. □

Proposition 7.390 (Gauge/presentation independence and pipeline comparability)**.** *With BRST–closed tags and calibrated branches, $\varsigma^{(\tau)}_{\mathrm{law}}$ is independent of gauge parameters and presentation–equivalent choices (Propositions 7.365, 7.367 and 7.372). Across a pipeline move $(\lambda_{\mathrm{UV}}, \ell) \mapsto (\tilde\lambda_{\mathrm{UV}}, \tilde\ell)$, one has*

$$\Big\|\varsigma^{(\tau)}_{\mathrm{law}}(\cdot; \tilde\lambda_{\mathrm{UV}}, \tilde\ell) - \varsigma^{(\tau)}_{\mathrm{law}}(\cdot; \lambda_{\mathrm{UV}}, \ell)\Big\|_\infty \;\le\; \|W\|_1 \sup_v \Big|\Lambda'_{\mathrm{law}}(v; \tilde{\cdot}) - \Lambda'_{\mathrm{law}}(v; \cdot)\Big|,$$

and the right–hand side is bounded by a constant multiple of the plateau tolerances (Corollary 7.378).

Proof. Step 1: Slope defined as a low–pass of the derivative. By Lemma 7.386 (using Definition 7.380), for every admissible parameter pair,

$$\varsigma^{(\tau)}_{\mathrm{law}}(\,\cdot\,; \lambda_{\mathrm{UV}}, \ell) = \frac{\mathrm{d}}{\mathrm{d}v}\Big(\mathsf{LP}_\tau \Lambda_{\mathrm{law}}(\,\cdot\,; \lambda_{\mathrm{UV}}, \ell)\Big) = \mathsf{LP}_\tau\Big(\Lambda'_{\mathrm{law}}(\,\cdot\,; \lambda_{\mathrm{UV}}, \ell)\Big) = W_\tau * \Lambda'_{\mathrm{law}}(\,\cdot\,; \lambda_{\mathrm{UV}}, \ell).$$

Step 2: Gauge/presentation independence. On certified plateaus, Λ_{law} is independent of gauge parameters and presentation–equivalent choices once the calibration phase is fixed; see Proposition 7.367 (gauge/presentation independence on plateaus) and Proposition 7.372 (robustness at initialization). Since LP_τ and $\frac{\mathrm{d}}{\mathrm{d}v}$ act only on the scale variable v and commute with each other (Lemma 7.381(4)), they preserve these invariances:

$$\partial_\xi \varsigma^{(\tau)}_{\mathrm{law}} = \partial_\xi\Big(\mathsf{LP}_\tau \Lambda'_{\mathrm{law}}\Big) = \mathsf{LP}_\tau\Big(\partial_\xi \Lambda'_{\mathrm{law}}\Big) = \mathsf{LP}_\tau\Big(\frac{\mathrm{d}}{\mathrm{d}v}(\partial_\xi \Lambda_{\mathrm{law}})\Big) = \frac{\mathrm{d}}{\mathrm{d}v}\Big(\mathsf{LP}_\tau(\partial_\xi \Lambda_{\mathrm{law}})\Big) = 0,$$

and similarly for presentation changes (replace ∂_ξ by the variation within the presentation class, which vanishes on $\mathcal{H}_{\mathrm{phys}}$ by Propositions 7.365 and 7.367). Hence $\varsigma^{(\tau)}_{\mathrm{law}}$ inherits gauge/presentation independence.

Step 3: Pipeline comparability — algebraic reduction. Fix two admissible pipeline choices $(\lambda_{\mathrm{UV}}, \ell)$ and $(\tilde\lambda_{\mathrm{UV}}, \tilde\ell)$. Using Step 1 and linearity,

$$\varsigma^{(\tau)}_{\mathrm{law}}(\cdot; \tilde{\cdot}) - \varsigma^{(\tau)}_{\mathrm{law}}(\cdot; \cdot) = W_\tau * \Big(\Lambda'_{\mathrm{law}}(\cdot; \tilde{\cdot}) - \Lambda'_{\mathrm{law}}(\cdot; \cdot)\Big).$$

Taking the L^∞ norm and applying Young's inequality (Lemma 7.381(1)) plus $\|W_\tau\|_1 = \|W\|_1$ (Definition 7.380) yields

$$\left\|\varsigma^{(\tau)}_{\mathrm{law}}(\cdot;\tilde{\cdot}) - \varsigma^{(\tau)}_{\mathrm{law}}(\cdot;\cdot)\right\|_\infty \le \|W\|_1 \left\|\Lambda'_{\mathrm{law}}(\cdot;\tilde{\cdot}) - \Lambda'_{\mathrm{law}}(\cdot;\cdot)\right\|_\infty,$$

which is the displayed bound (dropping the centered dot notation in the statement for readability).

Step 4: Pipeline comparability — tolerance control. By the comparability result for visibilities *and their first derivatives* along admissible pipeline moves (Corollary 7.378), there is a constant C (depending only on the certified plateau budgets and the move class) such that

$$\left\|\Lambda'_{\mathrm{law}}(\cdot;\tilde{\cdot}) - \Lambda'_{\mathrm{law}}(\cdot;\cdot)\right\|_\infty \;\le\; C\,(\text{tolerances}),$$

where "tolerances" abbreviates the plateau budgets (e.g. $\varepsilon_{\mathrm{grad}}, \varepsilon_{\mathrm{curv}}$, and the regulator/oscillation budgets specified in the acceptance pipeline). Combining with Step 3 gives

$$\left\|\varsigma^{(\tau)}_{\mathrm{law}}(\cdot;\tilde{\cdot}) - \varsigma^{(\tau)}_{\mathrm{law}}(\cdot;\cdot)\right\|_\infty \;\le\; \|W\|_1\, C\,(\text{tolerances}).$$

If $W \ge 0$ and $\int W = 1$ (zeroth moment in eq. (7.538)), then $\|W\|_1 = 1$ by Lemma 7.381(1), so the constant is just C.

This proves the proposition. □

Remark 7.391 (Eroded intersection window for pipeline moves)**.** Assume $\operatorname{supp} W \subset [-R, R]$, hence $\operatorname{supp} W_\tau \subset [-\tau R, \tau R]$. Let $\mathcal{I}$ and $\tilde{\mathcal{I}}$ be certified plateau windows for the two pipeline choices. Define the *eroded intersection window*

$$(\mathcal{I} \cap \tilde{\mathcal{I}})^{-\tau R} := \Big\{ v \in \mathcal{I} \cap \tilde{\mathcal{I}} \;:\; [\,v - \tau R,\, v + \tau R\,] \subset \mathcal{I} \cap \tilde{\mathcal{I}} \Big\}.$$

cf. the buffer convention in Corollary 7.384. Then the comparability bound of Proposition 7.390 holds verbatim with localized sup–norms:

$$\left\|\varsigma^{(\tau)}_{\mathrm{law}}(\cdot;\tilde{\cdot}) - \varsigma^{(\tau)}_{\mathrm{law}}(\cdot;\cdot)\right\|_{L^\infty\left((\mathcal{I}\cap\tilde{\mathcal{I}})^{-\tau R}\right)} \;\le\; \|W\|_1 \left\|\Lambda'_{\mathrm{law}}(\cdot;\tilde{\cdot}) - \Lambda'_{\mathrm{law}}(\cdot;\cdot)\right\|_{L^\infty(\mathcal{I}\cap\tilde{\mathcal{I}})}.$$

This guarantees that, for every v where the slope difference is evaluated, the convolution kernel $W_\tau(v - \cdot)$ samples the derivatives only inside certified plateau regions. If $W \ge 0$ and $\int W = 1$, then $\|W\|_1 = 1$ by Lemma 7.381, so the prefactor is unity.

IR window and acceptance threshold.

Definition 7.392 (IR window and acceptance threshold)**.** Let $v = \ln \ell$ and let LP_τ be as in Definition 7.380. Fix tolerances $\varepsilon_{\mathrm{curv}} > 0$ and a positivity threshold $\varsigma_{\min} > 0$, chosen consistently with the plateau criteria of Section 6.12.4 and the acceptance pipeline in Section 7.3.9. Define the IR window $\mathcal{I}_{\mathrm{IR}} \subset \mathbb{R}$ and the acceptance condition by

$$\boxed{\mathcal{I}_{\mathrm{IR}} := \left\{ v :\; \frac{\mathrm{d}^2}{\mathrm{d}v^2}\Big(\mathsf{LP}_\tau\, \Lambda_{\mathrm{law}}\Big)(v) \;\le\; \varepsilon_{\mathrm{curv}} \right\}, \qquad \varsigma^{(\tau)}_{\mathrm{law}}(v) \;\ge\; \varsigma_{\min} \;>\; 0 \quad \forall\, v \in \mathcal{I}_{\mathrm{IR}},} \tag{7.542}$$

where $\varsigma^{(\tau)}_{\mathrm{law}}$ is the robust slope of Definition 7.385.

Intuition. Inside $\mathcal{I}_{\mathrm{IR}}$, the low–passed visibility $(\mathsf{LP}_\tau\Lambda_{\mathrm{law}})(v)$ bends only mildly (curvature bounded) and climbs with a strictly positive, ripple–insensitive slope (Lemma 7.386 and Proposition 7.389). This encodes a controlled, monotone approach toward the plateau value (unity), in line with Section 6.12.4.

Lemma 7.393 (Scale covariance)**.** *Under $\ell \mapsto a\,\ell$ (i.e. $v \mapsto v+\ln a$), the window and acceptance test translate rigidly:*

$$\mathcal{I}_{\mathrm{IR}} \;\mapsto\; \mathcal{I}_{\mathrm{IR}} + \ln a, \qquad \varsigma^{(\tau)}_{\mathrm{law}}(v) \;\mapsto\; \varsigma^{(\tau)}_{\mathrm{law}}(v + \ln a),$$

as in Proposition 7.387.

Proof. Convolution with $W_\tau(v - v')$ and differentiation commute with translations in v (Lemma 7.381(4)), which yields the stated covariance. □

Proposition 7.394 (Non–emptiness on plateaus and robustness)**.** *Assume the plateau conditions of Section 6.12.4 and Kato–type continuity for Λ_{law} (Propositions 7.377 and 7.383). Then for any $\varepsilon_{\mathrm{curv}}$ above the plateau curvature tolerance and any $\varsigma_{\min}$ below the plateau slope tolerance, there exists a (possibly disconnected) window $\mathcal{I}_{\mathrm{IR}} \neq \varnothing$ satisfying Definition 7.392. Moreover, $\mathcal{I}_{\mathrm{IR}}$ is stable under small changes of τ and under pipeline moves $(\lambda_{\mathrm{UV}}, \ell) \mapsto (\tilde{\lambda}_{\mathrm{UV}}, \tilde{\ell})$, up to shifts bounded by the plateau tolerances.*

Proof. On a plateau, $|\Lambda'_{\mathrm{law}}(v)|$ and $|\Lambda''_{\mathrm{law}}(v)|$ are uniformly bounded by the tolerance. By Proposition 7.382, smoothing preserves constants/linear trends and perturbs derivatives by $\mathcal{O}(\tau^2)$, hence $(\mathsf{LP}_\tau\Lambda_{\mathrm{law}})''$ and $\varsigma^{(\tau)}_{\mathrm{law}}$ inherit the same bounds for sufficiently small (but fixed) τ, giving non–emptiness. Pipeline comparability of Λ_{law} and its derivatives follows from Corollary 7.378 and Proposition 7.390, implying stability of the inequalities that define $\mathcal{I}_{\mathrm{IR}}$. □

Lemma 7.395 (Law–deficit decay on the window)**.** *Let $D(v) := 1 - (\mathsf{LP}_\tau\Lambda_{\mathrm{law}})(v) \in [0,1]$. If $v \in \mathcal{I}_{\mathrm{IR}}$, then $D'(v) = -\varsigma^{(\tau)}_{\mathrm{law}}(v) \le -\varsigma_{\min}$. Hence for any $v_1 \ge v_0$ in the same connected component of $\mathcal{I}_{\mathrm{IR}}$,*

$$D(v_1) \;\le\; \max\Big\{\, D(v_0) - \varsigma_{\min}(v_1 - v_0),\; 0 \,\Big\}.$$

Proof. Set $D(v) := 1 - (\mathsf{LP}_\tau\Lambda_{\mathrm{law}})(v)$. By definition of $\varsigma^{(\tau)}_{\mathrm{law}}$ (cf. Definition 7.385), on $\mathcal{I}_{\mathrm{IR}}$ we have

$$\varsigma^{(\tau)}_{\mathrm{law}}(v) = \Big(\mathsf{LP}_\tau\Lambda_{\mathrm{law}}\Big)'(v) \ge \varsigma_{\min}.$$

Hence $D'(v) = -\varsigma^{(\tau)}_{\mathrm{law}}(v) \le -\varsigma_{\min}$. Integrating from v_0 to v_1 yields

$$D(v_1) - D(v_0) = \int_{v_0}^{v_1} D'(v)\,\mathrm{d}v \le -\,\varsigma_{\min}\,(v_1 - v_0),$$

i.e.

$$D(v_1) \le \max\Big\{\, D(v_0) - \varsigma_{\min}(v_1 - v_0),\; 0 \,\Big\}.$$

□

Corollary 7.396 (Hitting–length criterion)**.** *Fix v_0 in a connected component of $\mathcal{I}_{\mathrm{IR}}$ and let $L_{\mathrm{hit}} := D(v_0)/\varsigma_{\min}$. If the component length L to the right of v_0 satisfies $L \geq L_{\mathrm{hit}}$, then $D(v) = 0$ for some $v \in [v_0, v_0 + L]$; equivalently, $(\mathsf{LP}_\tau \Lambda_{\mathrm{law}})(v)$ reaches 1 within the window.*

Proof. Immediate from Lemma 7.395 and the bound $D \geq 0$. □

Proposition 7.397 (Gauge/presentation independence and pipeline comparability)**.** *With BRST–closed tags and calibrated branches, the IR window $\mathcal{I}_{\mathrm{IR}}$ and the thresholds in (7.542) are independent of gauge parameters and presentation–equivalent choices (Propositions 7.365, 7.367 and 7.372). Across any admissible pipeline move $(\lambda_{\mathrm{UV}}, \ell) \mapsto (\tilde{\lambda}_{\mathrm{UV}}, \tilde{\ell})$,*

$$\mu\Big(\mathcal{I}_{\mathrm{IR}}(\tilde{\lambda}_{\mathrm{UV}}, \tilde{\ell}) \triangle \mathcal{I}_{\mathrm{IR}}(\lambda_{\mathrm{UV}}, \ell)\Big) \;\leq\; C\Big(\varepsilon_{\mathrm{grad}} + \varepsilon_{\mathrm{curv}} + \varepsilon_{\mathrm{osc}} + \varepsilon_{\mathrm{reg}} + \varepsilon_{\mathrm{swap}}\Big),$$

for Lebesgue measure μ and a constant C depending only on $\|W\|_1$ and the local Lipschitz/crossing bounds from Section 6.12.4 (and the acceptance pipeline in Section 7.3.9).

Proof. Step 1: Gauge/presentation invariance. By Propositions 7.367 and 7.372, once the calibration phase is fixed, Λ_{law} is invariant under gauge–parameter changes and presentation–equivalent choices on certified plateaus. Since LP_τ and $\frac{d}{dv}$ act only on $v = \ln \ell$ and commute (Lemma 7.381(4)), the defining quantities

$$\varphi(v) := (\mathsf{LP}_\tau \Lambda_{\mathrm{law}})''(v), \qquad \psi(v) := (\mathsf{LP}_\tau \Lambda_{\mathrm{law}})'(v) = \varsigma^{(\tau)}_{\mathrm{law}}(v)$$

inherit the same invariances. Hence the curvature sublevel $\{\, \varphi \leq \varepsilon_{\mathrm{curv}} \,\}$, the slope–positivity set $\{\, \psi \geq \varsigma_{\min} \,\}$, and their intersection $\mathcal{I}_{\mathrm{IR}}$ (cf. (7.542)) are gauge/presentation independent.

Step 2: Pipeline perturbations of slope and curvature. For an admissible pipeline move, write

$$\tilde{\psi} - \psi = (\mathsf{LP}_\tau \tilde{\Lambda}'_{\mathrm{law}}) - (\mathsf{LP}_\tau \Lambda'_{\mathrm{law}}) = \mathsf{LP}_\tau\Big(\tilde{\Lambda}'_{\mathrm{law}} - \Lambda'_{\mathrm{law}}\Big), \quad \tilde{\varphi} - \varphi = \mathsf{LP}_\tau\Big(\tilde{\Lambda}''_{\mathrm{law}} - \Lambda''_{\mathrm{law}}\Big),$$

using the commutation in Lemma 7.381(4). Young's inequality and $\|W_\tau\|_1 = \|W\|_1$ (Definition 7.380) imply

$$\|\tilde{\psi} - \psi\|_\infty \leq \|W\|_1 \, \|\tilde{\Lambda}'_{\mathrm{law}} - \Lambda'_{\mathrm{law}}\|_\infty, \qquad \|\tilde{\varphi} - \varphi\|_\infty \leq \|W\|_1 \, \|\tilde{\Lambda}''_{\mathrm{law}} - \Lambda''_{\mathrm{law}}\|_\infty.$$

By the pipeline comparability of visibilities and their derivatives (Corollary 7.378), there exist constants C_ψ, C_φ (depending only on the certified budgets) such that

$$\delta_\psi := \|\tilde{\psi} - \psi\|_\infty \leq C_\psi \|W\|_1 \Big(\varepsilon_{\mathrm{grad}} + \varepsilon_{\mathrm{osc}} + \varepsilon_{\mathrm{reg}} + \varepsilon_{\mathrm{swap}}\Big),$$

$$\delta_\varphi := \|\tilde{\varphi} - \varphi\|_\infty \leq C_\varphi \|W\|_1 \Big(\varepsilon_{\mathrm{curv}} + \varepsilon_{\mathrm{osc}} + \varepsilon_{\mathrm{reg}} + \varepsilon_{\mathrm{swap}}\Big).$$

Step 3: From function perturbations to set perturbations. Let

$$\mathcal{I}_{\mathrm{IR}} = \underbrace{\{\, \varphi \leq \varepsilon_{\mathrm{curv}} \,\}}_{=:E_{\mathrm{curv}}} \cap \underbrace{\{\, \psi \geq \varsigma_{\min} \,\}}_{=:E_{\mathrm{slope}}}, \qquad \tilde{\mathcal{I}}_{\mathrm{IR}} = \{\, \tilde{\varphi} \leq \varepsilon_{\mathrm{curv}} \,\} \cap \{\, \tilde{\psi} \geq \varsigma_{\min} \,\}.$$

A pointwise inclusion holds for any real $g, \tilde{g}$ and threshold t:

$$\{g \le t\} \triangle \{\tilde{g} \le t\} \subset \{|g - t| \le \|g - \tilde{g}\|_\infty\}, \qquad \{g \ge t\} \triangle \{\tilde{g} \ge t\} \subset \{|g - t| \le \|g - \tilde{g}\|_\infty\}.$$

Therefore,

$$\mathcal{I}_{\mathrm{IR}} \triangle \tilde{\mathcal{I}}_{\mathrm{IR}} \;\subset\; \{\,|\varphi - \varepsilon_{\mathrm{curv}}| \le \delta_\varphi\,\} \;\cup\; \{\,|\psi - \varsigma_{\min}| \le \delta_\psi\,\}.$$

Step 4: Lipschitz band–measure estimate and collection of constants. By Section 6.12.4, φ and ψ are locally Lipschitz on the operational v-range; let $L_{\mathrm{curv}}, L_{\mathrm{slope}} > 0$ be corresponding bounds, and let $\mathsf{N}(\cdot)$ denote the finite crossing-count guards from the acceptance pipeline (Section 7.3.9). On any compact interval $\mathcal{J}$,

$$\mu\Big(\{|h - t| \le \delta\} \cap \mathcal{J}\Big) \;\le\; \frac{2\,\delta}{L}\,\mathsf{N}(h; \mathcal{J}) \quad \text{for Lipschitz } h \text{ with bound } L.$$

Applying this to $h = \varphi$, $t = \varepsilon_{\mathrm{curv}}$ and $h = \psi$, $t = \varsigma_{\min}$, summing, and inserting the bounds for $\delta_\varphi, \delta_\psi$ from Step 2 yields

$$\mu\Big(\mathcal{I}_{\mathrm{IR}}(\tilde{\cdot}) \triangle \mathcal{I}_{\mathrm{IR}}(\cdot)\Big) \;\le\; C\,\Big(\varepsilon_{\mathrm{grad}} + \varepsilon_{\mathrm{curv}} + \varepsilon_{\mathrm{osc}} + \varepsilon_{\mathrm{reg}} + \varepsilon_{\mathrm{swap}}\Big),$$

with C depending only on $\|W\|_1$, $L_{\mathrm{curv}}, L_{\mathrm{slope}}$, and the crossing–count bounds $\mathsf{N}(\cdot)$. (If $W \ge 0$ and $\int W = 1$, then $\|W\|_1 = 1$ by Lemma 7.381(1).)

The proposition follows. □

Definition 7.398 (Cell-averaged stencil from the continuum kernel)**.** Let $W \in C_c^\infty(\mathbb{R})$ be even with $\operatorname{supp} W \subset [-R, R]$ and define $W_\tau(s) = \tau^{-1} W(s/\tau)$. For grid step $h > 0$ and radius $m \in \mathbb{N}$, set the weights

$$\boxed{\,w_j := \frac{\displaystyle\int_{(j-\frac{1}{2})h}^{(j+\frac{1}{2})h} W_\tau(s)\,ds}{\displaystyle\sum_{r=-m}^{m} \int_{(r-\frac{1}{2})h}^{(r+\frac{1}{2})h} W_\tau(s)\,ds}, \qquad j = -m, \dots, m.\,} \tag{7.543}$$

Intuition. These w_j are the cell-averages of W_τ over the grid cells; this ties the discrete low–pass $\mathsf{LP}_{\tau,h}$ directly to the continuum operator in eq. (7.537).

Lemma 7.399 (Moment inheritance)**.** *With w_j from Definition 7.398 one has*

$$\sum_{j=-m}^{m} w_j = 1, \qquad \sum_{j=-m}^{m} j\,w_j = 0, \qquad w_{-j} = w_j.$$

Proof. Evenness of W_τ gives $w_{-j} = w_j$. The denominator in eq. (7.543) is a common positive scalar, so $\sum_j w_j = 1$ by construction and $\sum_j j\,w_j = \sum_{j=1}^{m} j(w_j - w_{-j}) = 0$. □

Corollary 7.400 (Affine preservation of $\mathsf{LP}_{\tau,h}$)**.** *Let $(\mathsf{LP}_{\tau,h} f)_k := \sum_{j=-m}^{m} w_j f_{k-j}$ with w_j as above and $v_k = v_0 + k\,h$. If $f(v) = a + b\,v$ and $f_k = f(v_k)$, then $(\mathsf{LP}_{\tau,h} f)_k = a + b\,v_k$.*

Intuition. Lemma 7.399 is the discrete analogue of eq. (7.538); it ensures constant/linear trends in $v = \ln \ell$ pass unchanged through the discrete low–pass.

Proposition 7.401 (Discrete–continuum correspondence)**.** *Let $f \in C^2$ on a neighborhood of v_k. With w_j from Definition 7.398, one has*

$$(\mathsf{LP}_{\tau,h} f)_k \;=\; (\mathsf{LP}_\tau f)(v_k) \;+\; O(h^2) \quad \text{as } h \to 0,$$

and, if $m = m(h)$ with $mh \to \infty$, then $(\mathsf{LP}_{\tau,h} f)_k \to (\mathsf{LP}_\tau f)(v_k)$ uniformly on compact v-sets.

Proof sketch. Write $(\mathsf{LP}_{\tau,h} f)_k = \sum_j f\big(v_k - jh\big)\, \overline{W}_{\tau,h}(j)$ where $\overline{W}_{\tau,h}(j)$ are the normalized cell-averages from eq. (7.543). This is a Riemann-sum for $(\mathsf{LP}_\tau f)(v_k) = \int W_\tau(s) f(v_k - s)\, ds$. Evenness and vanishing first moment yield the $O(h^2)$ local error by Taylor expansion. □

Intuition. The discrete filter is a faithful, second-order accurate proxy for the continuum mollifier; increasing the stencil radius with h^{-1} recovers the full integral.

$$\boxed{\widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(v_k) := \big(\mathsf{D}_0\, \mathsf{LP}_{\tau,h}\, \Lambda_{\mathrm{law}}\big)_k.} \tag{7.544}$$

Intuition. This is the low–pass–then–central–difference readout in v; by Corollary 7.400 and Definition 7.406, it is unbiased on affine trends.

Remark 7.402 (Continuum vs. discrete slope: same concept, distinct objects)**.** The continuum robust slope is defined in Definition 7.385 by $\varsigma^{(\tau)}_{\mathrm{law}}(v) = \frac{d}{dv}\big(\mathsf{LP}_\tau \Lambda_{\mathrm{law}}\big)(v)$ (see eq. (7.539)), with the convolution representation given in Lemma 7.386. The discrete readout displayed above is the grid estimator $\widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(v_k) = (\mathsf{D}_0\, \mathsf{LP}_{\tau,h} \Lambda_{\mathrm{law}})_k$, for which the canonical label is eq. (7.547). These two quantities represent the same conceptual observable but are not the same mathematical object or label. With the cell-average stencil from Definition 7.398 and smooth Λ_{law} at fixed τ,

$$(\mathsf{D}_0\, \mathsf{LP}_{\tau,h} \Lambda_{\mathrm{law}})_k \;=\; \varsigma^{(\tau)}_{\mathrm{law}}(v_k) \;+\; O(h^2),$$

so the discrete estimator is second–order accurate and, by Lemma 7.412 and Corollary 7.400, exact on affine trends.

Definition 7.403 (Admissible boundary extension)**.** Let $I = \{k_{\min}, \dots, k_{\max}\}$. A linear map $E : \mathbb{R}^I \to \mathbb{R}^{\mathbb{Z}}$ is an admissible extension if (i) E agrees with the identity on I, and (ii) E is fixed independently of f (e.g., periodic wrap, zero–padding, or mirrored/Neumann ghosts). Let $R : \mathbb{R}^{\mathbb{Z}} \to \mathbb{R}^I$ be the restriction $R(g) = g|_I$.

Intuition. E specifies a boundary rule so that discrete convolutions can be applied on the whole grid and then restricted back.

Proposition 7.404 (Commutation survives extension–restriction)**.** *Let w and d denote the finite-support kernels of $\mathsf{LP}_{\tau,h}$ and D_0, respectively. For any admissible E and restriction R,*

$$R\Big(d * (w * Ef)\Big) \;=\; R\Big(w * (d * Ef)\Big) \qquad \text{for all } f \in \mathbb{R}^I.$$

Proof. Associativity and commutativity of discrete convolution on $\mathbb{Z}$ yield $d * (w * Ef) = (w * d) * Ef$. Apply R. □

Remark 7.405 (Affine exactness at the edges)**.** With eroded indices, Lemma 7.412 gives exact affine readout everywhere used. Under extensions: periodic/zero–padding preserve commutation but may lose affine exactness at the first/last indices; mirrored/Neumann also preserve commutation but flip the slope at the boundary. State affine exactness claims on the eroded set or specify the extension.

Discrete estimator for audit and numerics.

Definition 7.406 (Central difference on the log–scale grid)**.** Let $v_k = v_0 + k\,h$ with $v = \ln \ell$. The operator $\mathsf{D}_0 : \{f_k\} \mapsto \{(\mathsf{D}_0 f)_k\}$ is

$$\boxed{(\mathsf{D}_0 f)_k := \frac{f_{k+1} - f_{k-1}}{2h}.} \tag{7.545}$$

Remark 7.407 (Meaning and basic properties)**.** D_0 is the standard symmetric, linear time-invariant (LTI) first-difference on the uniform grid in v. It is second-order accurate for $\mathrm{d}/\mathrm{d}v$ (local truncation error $O(h^2)$), annihilates constants, and returns the exact slope on affine trends: if $f(v) = a + b\,v$ and $f_k = f(v_k)$, then $(\mathsf{D}_0 f)_k = b$. Equivalently, $\mathsf{D}_0 = d*$ with $d_{+1} = 1/(2h)$, $d_{-1} = -1/(2h)$, else 0; this convolution form underlies the commutation with discrete low-pass filters used in Lemma 7.412. Boundary handling is via the usual eroded index set or mirrored/Neumann ghost points (as declared later).

Definition 7.408 (Discrete stencil on the log–scale grid)**.** On a uniform grid $v_k = v_0 + k\,h$, a (centered) stencil is a finite set of offsets $\{-m, \dots, m\} \subset \mathbb{Z}$ with coefficients $\{a_j\}_{j=-m}^{m}$ defining a linear, shift–invariant operator

$$(Af)_k := \sum_{j=-m}^{m} a_j\, f_{k-j}.$$

It is *symmetric* if $a_{-j} = a_j$. In this work:

- the low–pass uses the stencil $w = \{w_j\}_{-m}^{m}$ from Definition 7.409, with continuum anchoring via the cell averages eq. (7.543);
- the central difference uses $d_{-1} = -\frac{1}{2h}$, $d_0 = 0$, $d_{+1} = \frac{1}{2h}$ from Definition 7.406;
- the mollified–slope readout $(\mathsf{D}_0 \mathsf{LP}_{\tau,h})$ is the *composed stencil* $c = d * w$, i.e.

$$(\mathsf{D}_0 \mathsf{LP}_{\tau,h} f)_k = \sum_{j=-(m+1)}^{m+1} c_j\, f_{k-j}, \qquad c_j := \sum_{r\in\{-1,0,1\}} d_r\, w_{j-r}. \tag{7.546}$$

Definition 7.409 (Discrete low–pass and robust–slope estimator)**.** Fix a uniform grid $v_k = v_0 + k\,h$ (so $\ell_k = e^{v_k}$), and let $W \in C_c^\infty(\mathbb{R})$ have support in $[-R, R]$. Choose a symmetric, compact stencil $\{w_j\}_{j=-m}^{m}$ with

$$\sum_{j=-m}^{m} w_j = 1, \qquad \sum_{j=-m}^{m} j\, w_j = 0, \qquad w_{-j} = w_j,$$

approximating the continuous kernel in eq. (7.537); take $m = \lceil (\tau R)/h \rceil$. Define the discrete low–pass $\mathsf{LP}_{\tau,h}$ by

$$(\mathsf{LP}_{\tau,h} f)_k := \sum_{j=-m}^{m} w_j \, f_{k-j},$$

and use the central difference D_0 from Definition 7.406. Assume indices k satisfy $\{k-m, \dots, k+m\} \cup \{k \pm 1\}$ lie inside the data window. The *discrete robust law slope* is

$$\boxed{\widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(v_k) := \Big(\mathsf{D}_0 \, \mathsf{LP}_{\tau,h} \, \Lambda_{\mathrm{law}}\Big)_k = \frac{1}{2h}\left[\sum_{j=-m}^{m} w_j \, \Lambda_{\mathrm{law}}(v_{k+1-j}) - \sum_{j=-m}^{m} w_j \, \Lambda_{\mathrm{law}}(v_{k-1-j})\right],} \tag{7.547}$$

which is the discrete counterpart of the robust slope in Definition 7.385 (cf. the convolution form (7.540)).

Intuition. The operator $\mathsf{LP}_{\tau,h}$ is a compact, symmetric average that preserves constants and linear trends (the discrete analogue of the continuous moment conditions in eqs. (7.537) and (7.538)), while D_0 is the standard second–order central derivative. Their composition $\mathsf{D}_0 \, \mathsf{LP}_{\tau,h}$ filters grid–scale ripples yet tracks the IR slope faithfully, yielding a robust discrete counterpart of the continuous slope in Definition 7.385 (see Definitions 7.406 and 7.409).

Stencil intuition. In Definition 7.409, the *stencil* is the finite pattern of offsets $\{-m, \dots, m\}$ and weights $\{w_j\}$ slid along the log–grid $v_k = v_0 + k\,h$. It may be viewed as a compact "mask": symmetry $w_{-j} = w_j$ and the moment constraints $\sum_j w_j = 1$, $\sum_j j\, w_j = 0$ enforce unbiased averaging, exact reproduction of affine trends, and zero phase distortion, so grid–scale oscillations are suppressed without shifting IR features. The half–width $m \simeq \lceil \tau R/h \rceil$ ties the spatial reach of the mask to the IR scale τ, matching the continuous construction in eq. (7.537). Composing the low–pass with the central difference produces a single derivative–stencil with coefficients

$$\widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(v_k) = \sum_{r=-(m+1)}^{m+1} c_r \, \Lambda_{\mathrm{law}}(v_{k-r}), \qquad c_r = \frac{w_{r+1} - w_{r-1}}{2h} \quad (w_j := 0 \text{ for } |j| > m),$$

which is antisymmetric ($c_{-r} = -c_r$), kills constants, and reproduces linear trends with second–order accuracy. The index condition in Definition 7.409 (the "eroded" set) ensures all offsets required by the stencil remain within the available data window, avoiding boundary artefacts. By stabilising the slope $\psi(v) = (\mathsf{D}_0 \, \mathsf{LP}_{\tau,h} \Lambda_{\mathrm{law}})(v)$, the stencil integrates seamlessly with the curvature and total–variation (TV) guards (Definition 7.420 and Lemma 7.423), thereby reducing false positives in emergent–trend detection and providing a phase–neutral, bias–controlled, IR–faithful estimator suitable for certified plateau diagnostics.

Proposition 7.410 (Discrete robust-slope error)**.** *Let $W \in C_c^\infty(\mathbb{R})$ be even with $\int W = 1$, $\int v\, W(v)\, dv = 0$, and finite $m_2(W)$, and let the stencil $\{w_j\}_{-m}^{m}$ be symmetric with $\sum w_j = 1$, $\sum j\, w_j = 0$, $m = \lceil \tau R/h \rceil$. On an eroded index set (to avoid edges),*

$$\left\| \widehat{\varsigma}^{(\tau)}_{\mathrm{law}} - \varsigma^{(\tau)}_{\mathrm{law}} \right\|_\infty \;\le\; C\Big(h^2 + \tau^2\Big) \left\| \Lambda^{(3)}_{\mathrm{law}} \right\|_\infty \;+\; E_{\ker}(h, \tau),$$

where $E_{\ker}(h,\tau)$ is the kernel-discretization error (zero if $w_j \propto h\,W_\tau(jh)$ are used and renormalized). If, moreover, $\Lambda_{\mathrm{law}} \in C^5$ and $m_4(W) < \infty$, then the continuous mollifier bias improves to $\mathcal{O}(\tau^4)$ and

$$\left\| \widehat{\varsigma}^{(\tau)}_{\mathrm{law}} - \varsigma^{(\tau)}_{\mathrm{law}} \right\|_\infty \;\le\; C\Big(h^2 \left\| \Lambda^{(3)}_{\mathrm{law}} \right\|_\infty + \tau^4 \left\| \Lambda^{(5)}_{\mathrm{law}} \right\|_\infty \Big) \;+\; E_{\ker}(h,\tau).$$

Proof. Write $f := \Lambda_{\mathrm{law}}$, $v_k = v_0 + kh$, and set

$$g(v) := (\mathsf{LP}_\tau f)(v), \qquad g_k := (\mathsf{LP}_{\tau,h} f)_k = \sum_{j=-m}^{m} w_j\, f(v_{k-j}).$$

Recall the robust slope and its discrete estimator:

$$\varsigma^{(\tau)}_{\mathrm{law}}(v) = g'(v), \qquad \widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(v_k) = (\mathsf{D}_0 g_\bullet)_k = \frac{g_{k+1} - g_{k-1}}{2h}$$
$$\text{with } g_{k\pm1} \text{ understood as } \sum_j w_j f(v_{k\pm1-j}).$$

We decompose the error by adding and subtracting $f'(v_k)$:

$$\left| \widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(v_k) - \varsigma^{(\tau)}_{\mathrm{law}}(v_k) \right| \;\le\; \underbrace{\left| \widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(v_k) - f'(v_k) \right|}_{\text{(A) discrete vs. true derivative}} \;+\; \underbrace{\left| f'(v_k) - g'(v_k) \right|}_{\text{(B) continuous mollifier bias}}.$$

(B) Continuous mollifier bias. By Proposition 7.382 (applied to f and differentiated using Lemma 7.381(4)),

$$g'(v) = f'(v) + \frac{\tau^2 m_2(W)}{2}\, f^{(3)}(v) + \mathcal{O}(\tau^4),$$

hence

$$\sup_k |f'(v_k) - g'(v_k)| \;\le\; C\,\tau^2\, \|f^{(3)}\|_\infty$$
$$\text{(and improves to } C\,\tau^4\, \|f^{(5)}\|_\infty \text{ if } f \in C^5,\ m_4(W) < \infty).$$

(A) Discrete vs. true derivative. Split (A) further by inserting the continuous smoothed profile g sampled on the grid:

$$\left| \widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(v_k) - f'(v_k) \right| \;\le\; \underbrace{\left| (\mathsf{D}_0 g_\bullet)_k - (\mathsf{D}_0 g(v_\bullet))_k \right|}_{\text{(A1) kernel discretization}} \;+\; \underbrace{\left| (\mathsf{D}_0 g(v_\bullet))_k - g'(v_k) \right|}_{\text{(A2) central-difference error}}.$$

(A2) *Central-difference error.* Since $g \in C^3$ and D_0 is the second–order central scheme,

$$\left| (\mathsf{D}_0 g(v_\bullet))_k - g'(v_k) \right| \;\le\; C\,h^2\, \|g^{(3)}\|_\infty.$$

By Lemma 7.381(4), derivatives commute with LP_τ, so $g^{(3)} = \mathsf{LP}_\tau(f^{(3)})$, whence $\|g^{(3)}\|_\infty \le \|W\|_1\, \|f^{(3)}\|_\infty = \|f^{(3)}\|_\infty$. Thus

$$\sup_k \left| (\mathsf{D}_0 g(v_\bullet))_k - g'(v_k) \right| \;\le\; C\,h^2\, \|f^{(3)}\|_\infty.$$

(A1) *Kernel discretization.* Define the kernel-discretization residual

$$E_{\mathrm{ker}}(h,\tau) := \left\| \mathsf{D}_0\Big(\mathsf{LP}_{\tau,h} f - \mathsf{S}_h(\mathsf{LP}_\tau f)\Big)\right\|_\infty,$$

where S_h is the sampling operator $(\mathsf{S}_h g)_k := g(v_k)$. This term captures the effect of replacing the continuous convolution by a discrete stencil before differencing. By construction,

$$\sup_k \left|(\mathsf{D}_0 g_\bullet)_k - (\mathsf{D}_0 g(v_\bullet))_k\right| \;\le\; E_{\mathrm{ker}}(h,\tau).$$

Moreover, if one *chooses* the stencil via sampled kernel weights, $w_j \propto h\, W_\tau(jh)$ and renormalizes to enforce $\sum w_j = 1$ and symmetry (the audit/numerics convention), then $\mathsf{LP}_{\tau,h}$ is, by definition, the discrete counterpart of LP_τ used in the pipeline, and the above residual is set to zero: $E_{\mathrm{ker}}(h,\tau) = 0$. (If one prefers an analytical bound instead of this convention, standard quadrature estimates give $E_{\mathrm{ker}}(h,\tau) \le C_W\, h^2\, \|f''\|_\infty$ for smooth f and compactly supported W, which can be absorbed into the $Ch^2\|f^{(3)}\|_\infty$ term using $\|f''\|_\infty \le \|f^{(3)}\|_\infty$ on the eroded window up to a multiplicative constant from the local scale.)

Collecting the bounds. Combining (B), (A2), and (A1) via the triangle inequality gives

$$\left\| \widehat{\varsigma}^{(\tau)}_{\mathrm{law}} - \varsigma^{(\tau)}_{\mathrm{law}} \right\|_\infty \;\le\; C\, h^2\, \|f^{(3)}\|_\infty \;+\; C\, \tau^2\, \|f^{(3)}\|_\infty \;+\; E_{\mathrm{ker}}(h,\tau),$$

which is the first displayed estimate. Under $f \in C^5$ and $m_4(W) < \infty$, the τ^2 term improves to $C\,\tau^4\,\|f^{(5)}\|_\infty$, yielding the second estimate. All constants are independent of h, τ and of k, and the bounds hold on the eroded index set $\{k : k \pm m$ remain inside the window$\}$ to avoid edge effects. □

Remark 7.411 (When to ask for higher smoothness)**.** Use C^3 when we want the robust, practical bound $\mathcal{O}(h^2{+}\tau^2)$ for the discrete robust–slope error. **Bump to** C^5 (plus $m_4(W) < \infty$) only if we really want the $\mathcal{O}(\tau^4)$ continuous-bias term spelled out; otherwise the C^3 hypothesis suffices.

Lemma 7.412 (Commutation and affine exactness)**.** *Let f_k be a grid function on the uniform log–scale grid $v_k = v_0 + k\,h$ (with $v = \ln \ell$). Let the discrete low–pass $\mathsf{LP}_{\tau,h}$ with symmetric stencil $\{w_j\}_{j=-m}^{m}$ be as in Definition 7.409 (and hence $\sum_j w_j = 1$, $\sum_j j\, w_j = 0$, $w_{-j} = w_j$), and let the central difference D_0 be as in Definition 7.406. For all interior indices k with available samples (i.e. $\{k \pm 1\} \cup \{k-m, \dots, k+m\}$ lie in the data window), the following hold:*

Commutation.

$$\mathsf{D}_0\Big(\mathsf{LP}_{\tau,h} f\Big) = \mathsf{LP}_{\tau,h}\Big(\mathsf{D}_0 f\Big).$$

Affine exactness.

If $f(v) = a + b\,v$ and $f_k = f(v_k)$, then $(\mathsf{LP}_{\tau,h} f)_k = a + b\,v_k$ and $(\mathsf{D}_0 f)_k = b$. Consequently, the discrete robust law slope $\widehat{\varsigma}^{(\tau)}_{\mathrm{law}}$ in eq. (7.547) returns b exactly on affine trends.

Proof. (1) Using finite support of $\{w_j\}$ and linearity,

$$\begin{aligned}\big(\mathsf{D}_0\,\mathsf{LP}_{\tau,h}f\big)_k &= \frac{1}{2h}\left[\sum_{j=-m}^{m} w_j\, f_{k+1-j} - \sum_{j=-m}^{m} w_j\, f_{k-1-j}\right]\\ &= \sum_{j=-m}^{m} w_j\,\frac{f_{k+1-j}-f_{k-1-j}}{2h}\\ &= \big(\mathsf{LP}_{\tau,h}\,\mathsf{D}_0 f\big)_k.\end{aligned}$$

Equivalently, with discrete convolution $(w*f)_k := \sum_j w_j f_{k-j}$, $\mathsf{LP}_{\tau,h} = w*$ and $\mathsf{D}_0 = d*$ where $d_{+1} = 1/(2h)$, $d_{-1} = -1/(2h)$, else 0; associativity/commutativity of finite-support convolution gives $d*w = w*d$.

(2) For $f_k = a + b\,v_k$ and $v_{k-j} = v_k - jh$,

$$\begin{aligned}(\mathsf{LP}_{\tau,h}f)_k &= \sum_{j=-m}^{m} w_j\big(a + b(v_k - jh)\big)\\ &= a\sum_j w_j + b\,v_k\sum_j w_j - bh\sum_j j\,w_j\\ &= a + b\,v_k,\end{aligned}$$

using $\sum_j w_j = 1$ and $\sum_j j\,w_j = 0$. From Definition 7.406, $(\mathsf{D}_0 f)_k = b$. Combining these with part (1) and eq. (7.547) gives the claim about $\widehat{\varsigma}^{(\tau)}_{\mathrm{law}}$. □

Intuition. This is the discrete shadow of Definition 7.380: symmetry and the first-moment cancellation mirror eq. (7.538), so $\mathsf{LP}_{\tau,h}$ preserves constants/lines in $v = \ln \ell$ and commutes with any LTI first-difference like D_0. It ensures your low-pass slope $\widehat{\varsigma}^{(\tau)}_{\mathrm{law}}$ is an *unbiased* estimator on plateaus/ramps before we look at higher-order structure.

Proposition 7.413 (Consistency and bias)**.** *Assume $\Lambda_{\mathrm{law}} \in C^3$ in v and the discrete weights approximate the continuous kernel of eq. (7.537) (moment conditions as in eq. (7.538)). Then*

$$\widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(v_k) = \varsigma^{(\tau)}_{\mathrm{law}}(v_k) \;+\; \mathcal{O}(h^2) \;+\; \mathcal{O}(\tau^2),$$

where $\varsigma^{(\tau)}_{\mathrm{law}}$ is the continuous robust slope of Definition 7.385.

Remark 7.414 (On the C^3 regularity in Proposition 7.413)**.** The requirement $\Lambda_{\mathrm{law}} \in C^3$ controls *both* sources of error that appear in

$$\widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(v_k) = \varsigma^{(\tau)}_{\mathrm{law}}(v_k) + \mathcal{O}(h^2) + \mathcal{O}(\tau^2).$$

(i) Smoothing bias $\mathcal{O}(\tau^2)$. Using the kernel moment conditions in eq. (7.538) and the Taylor expansion of $\Lambda_{\mathrm{law}}(v-s)$ to second order yields the standard bias formula $(\mathsf{LP}_\tau\Lambda_{\mathrm{law}})(v) = \Lambda_{\mathrm{law}}(v) + \frac{\tau^2 m_2(W)}{2}\Lambda''_{\mathrm{law}}(v) + \mathcal{O}(\tau^4)$ (Proposition 7.382). Differentiating in v and commuting the derivative with LP_τ (Lemma 7.381) gives $\varsigma^{(\tau)}_{\mathrm{law}}(v) = \Lambda'_{\mathrm{law}}(v) + \frac{\tau^2 m_2(W)}{2}\Lambda^{(3)}_{\mathrm{law}}(v) + \mathcal{O}(\tau^4)$. Bounding this term uniformly requires $\Lambda_{\mathrm{law}} \in C^3$.

(ii) Discretization error $\mathcal{O}(h^2)$. Two second–order contributions are present: (a) the Riemann–cell discretization of LP_τ via the weights w_j from Definition 7.398 yields $(\mathsf{LP}_{\tau,h} f)_k = (\mathsf{LP}_\tau f)(v_k) + \mathcal{O}(h^2)$ for smooth f (Proposition 7.401); (b) the central difference D_0 (Definition 7.406) approximates $\frac{d}{dv}$ with local error $\mathcal{O}(h^2)$, whose control involves the third derivative of the target function. Applying (a)–(b) to $f = \Lambda_{\mathrm{law}}$ (and using the commutation in Lemma 7.412) shows that $(\mathsf{D}_0 \mathsf{LP}_{\tau,h} \Lambda_{\mathrm{law}})_k$ deviates from $\frac{d}{dv}(\mathsf{LP}_\tau \Lambda_{\mathrm{law}})(v_k)$ by $\mathcal{O}(h^2)$, with bounds depending on $\sup |\Lambda^{(3)}_{\mathrm{law}}|$.

Summary. The τ-bias term involves $\Lambda^{(3)}_{\mathrm{law}}$ after differentiating the continuum bias; the h-discretization term for D_0 also depends on a third derivative. Hence C^3 is the natural minimal regularity ensuring the stated $\mathcal{O}(h^2) + \mathcal{O}(\tau^2)$ accuracy. (On affine trends, exactness holds by Corollary 7.400 and Definition 7.406.)

Proof. Write $f := \Lambda_{\mathrm{law}}$, $v_k = v_0 + kh$. Work on the eroded index set so that all stencil points are available (cf. the buffer convention used with the continuous τR window).

Step 1: Move the discrete derivative inside the low–pass. By the commutation identity in Lemma 7.412(1),

$$\widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(v_k) = \big(\mathsf{D}_0\, \mathsf{LP}_{\tau,h} f\big)_k = \big(\mathsf{LP}_{\tau,h}\, \mathsf{D}_0 f\big)_k.$$

Step 2: Central difference error $\mathcal{O}(h^2)$. Since $f \in C^3$,

$$(\mathsf{D}_0 f)_k = \frac{f(v_{k+1}) - f(v_{k-1})}{2h} = f'(v_k) + \mathcal{O}(h^2) \qquad \text{(uniformly in } k\text{)}.$$

Applying $\mathsf{LP}_{\tau,h}$ and using $\sum_j |w_j| = \sum_j w_j = 1$ (nonnegative symmetric stencil) yields

$$\big(\mathsf{LP}_{\tau,h}\, \mathsf{D}_0 f\big)_k = \big(\mathsf{LP}_{\tau,h} f'\big)_k + \mathcal{O}(h^2).$$

Step 3: Discrete vs. continuous low–pass $\mathcal{O}(h^2)$. Let $g := f'$. The weights $\{w_j\}$ approximate the continuous kernel W_τ and satisfy the discrete moment conditions $\sum w_j = 1$, $\sum j\, w_j = 0$ (discrete analog of $\int W = 1$, $\int vW = 0$; cf. eq. (7.538)). With $g \in C^2$ (since $f \in C^3$) and compact stencil, a standard quadrature/Taylor estimate gives

$$\big(\mathsf{LP}_{\tau,h} g\big)_k = \big(\mathsf{LP}_\tau g\big)(v_k) + \mathcal{O}(h^2),$$

uniformly for k in the eroded set. Hence

$$\big(\mathsf{LP}_{\tau,h}\, \mathsf{D}_0 f\big)_k = \big(\mathsf{LP}_\tau f'\big)(v_k) + \mathcal{O}(h^2).$$

Step 4: Identify the continuous robust slope. By the differentiation–commutation lemma Lemma 7.386,

$$\big(\mathsf{LP}_\tau f'\big)(v) = \frac{d}{dv}\big(\mathsf{LP}_\tau f\big)(v) = \varsigma^{(\tau)}_{\mathrm{law}}(v),$$

so

$$\widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(v_k) = \varsigma^{(\tau)}_{\mathrm{law}}(v_k) + \mathcal{O}(h^2).$$

Step 5: Add the mollifier bias if comparing to f'. While Step 4 shows $\widehat{\varsigma}^{(\tau)}_{\mathrm{law}} = \varsigma^{(\tau)}_{\mathrm{law}} + \mathcal{O}(h^2)$, one often also reports the bias relative to the *unsmoothed* slope $f' = \Lambda'_{\mathrm{law}}$. By Proposition 7.382 (applied to f and then differentiated),

$$\varsigma^{(\tau)}_{\mathrm{law}}(v) = \big(\mathsf{LP}_\tau f\big)'(v) = f'(v) + \mathcal{O}(\tau^2),$$

so, by the triangle inequality,

$$\widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(v_k) = \varsigma^{(\tau)}_{\mathrm{law}}(v_k) \;+\; \mathcal{O}(h^2) \;+\; \mathcal{O}(\tau^2).$$

All big-O constants are uniform on the eroded index set and depend only on local C^3-bounds for f and on the (fixed) kernel/stencil parameters. *Note.* The $\mathcal{O}(\tau^2)$ term reflects the *continuous* low–pass bias relative to the raw derivative $f' = \Lambda'_{\mathrm{law}}$; if one measures error against the robust slope itself, $\varsigma^{(\tau)}_{\mathrm{law}} = (\mathsf{LP}_\tau f)'$, then $\widehat{\varsigma}^{(\tau)}_{\mathrm{law}} = \varsigma^{(\tau)}_{\mathrm{law}} + \mathcal{O}(h^2)$. □

Lemma 7.415 (Stability and ripple suppression)**.** *Let $\mathsf{LP}_{\tau,h}$ and D_0 be as in Definitions 7.406 and 7.409, with stencil $\{w_j\}_{j=-m}^{m}$ (e.g. the cell–average choice of Definition 7.398). For any admissible index k (on the eroded set so all required samples exist),*

$$\left|\widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(v_k)\right| \;\le\; \sum_{j=-m}^{m} |w_j| \sup_{\xi\in[v_{k-1-j},\,v_{k+1-j}]} \left|\Lambda'_{\mathrm{law}}(\xi)\right| \;\le\; \|w\|_{\ell^1}\,\|\Lambda'_{\mathrm{law}}\|_\infty,$$

where $\widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(v_k) := (\mathsf{D}_0\,\mathsf{LP}_{\tau,h}\Lambda_{\mathrm{law}})_k$ (cf. eq. (7.547)) and $\|w\|_{\ell^1} := \sum_{j=-m}^{m}|w_j|$. In the discrete Fourier domain (grid frequency $\xi \in [-\pi,\pi]$), the estimator has symbol

$$\widehat{\varsigma}(\xi) \;=\; \frac{i\sin\xi}{h}\,\widehat{W}(\xi), \qquad \widehat{W}(\xi) := \sum_{j=-m}^{m} w_j\, e^{-i\xi j},$$

so symmetric stencils ($w_{-j} = w_j$, cf. Lemma 7.399) attenuate high–$|\xi|$ components by the factor $|\widehat{W}(\xi)|$ while preserving DC gain $\widehat{W}(0) = \sum_j w_j = 1$.

Proof. Pointwise stability bound. By Definitions 7.406 and 7.409 (or Lemma 7.412(1)),

$$\widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(v_k) = \sum_{j=-m}^{m} w_j\,\frac{\Lambda_{\mathrm{law}}(v_{k+1-j}) - \Lambda_{\mathrm{law}}(v_{k-1-j})}{2h}.$$

For each fixed j, apply the mean–value theorem to Λ_{law} on $[\,v_{k-1-j},\,v_{k+1-j}\,]$ to obtain some $\xi_{k,j}$ in that interval with

$$\frac{\Lambda_{\mathrm{law}}(v_{k+1-j}) - \Lambda_{\mathrm{law}}(v_{k-1-j})}{2h} = \Lambda'_{\mathrm{law}}(\xi_{k,j}).$$

Hence

$$\left|\widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(v_k)\right| \le \sum_{j=-m}^{m} |w_j| \sup_{\xi\in[\,v_{k-1-j},\,v_{k+1-j}\,]} \left|\Lambda'_{\mathrm{law}}(\xi)\right| \le \Big(\sum_{j=-m}^{m}|w_j|\Big)\,\|\Lambda'_{\mathrm{law}}\|_\infty = \|w\|_{\ell^1}\,\|\Lambda'_{\mathrm{law}}\|_\infty.$$

If, in addition, the stencil is constructed from a nonnegative W via Definition 7.398 (so $w_j \ge 0$) and normalized as in eq. (7.538), then $\|w\|_{\ell^1} = \sum_j w_j = 1$.

Fourier symbol and ripple suppression. Write $(\mathsf{D}_0 f)_k = (d * f)_k$ and $(\mathsf{LP}_{\tau,h} f)_k = (w * f)_k$ (discrete convolution), with $d_{-1} = -\frac{1}{2h}$, $d_{+1} = \frac{1}{2h}$, otherwise 0 (cf. Definition 7.406), and $w = \{w_j\}$ as above. On a periodic (or ℓ^2) grid, the DTFT maps convolution to multiplication, hence the symbol of $\mathsf{D}_0\,\mathsf{LP}_{\tau,h}$ is

$$\widehat{\varsigma}(\xi) = \widehat{d}(\xi)\,\widehat{W}(\xi) = \frac{e^{i\xi} - e^{-i\xi}}{2h}\,\widehat{W}(\xi) = \frac{i\sin\xi}{h}\,\widehat{W}(\xi).$$

If $w_{-j} = w_j$ (guaranteed, e.g., by Lemma 7.399 when using Definition 7.398 and an even W per eq. (7.538)), then $\widehat{W}(\xi) = \sum_j w_j \cos(\xi j)$ is real, even, satisfies $|\widehat{W}(\xi)| \le \sum_j |w_j| = \|w\|_{\ell^1}$, and $\widehat{W}(0) = \sum_j w_j = 1$. Thus $\widehat{W}$ damps high–frequency content (large $|\xi|$ near π) while leaving DC unchanged, establishing ripple suppression by the low–pass stage. □

Remark 7.416 (Eroded index set and frequency response).

- The *eroded index set* mirrors the continuous τR buffer (cf. the low–pass support in Definition 7.380): it guarantees all entries $k \pm j$ used by the stencil and $k \pm 1$ used by D_0 lie inside the available window, so no boundary artefacts appear (see also Corollary 7.384).
- If the weights are constructed from sampled W_τ and renormalized ($w_j \propto h\, W_\tau(jh)$), then the discrete frequency response $\widehat{W}(\xi) = \sum_{j=-m}^{m} w_j e^{-i\xi j}$ closely matches the continuous kernel's response, and the attenuation profile inherits the desired low–pass shape.

Corollary 7.417 (IR acceptance via discrete tests). *Define the discrete curvature with the second–difference* $\mathsf{D}_2 f_k := (f_{k+1} - 2f_k + f_{k-1})/h^2$. *Then the discrete IR window*

$$\mathcal{I}_{\mathrm{IR}}^{(h)} := \left\{ k : \ \left| \mathsf{D}_2 (\mathsf{LP}_{\tau,h} \Lambda_{\mathrm{law}})_k \right| \le \varepsilon_{\mathrm{curv}} \right\}$$

approximates $\mathcal{I}_{\mathrm{IR}}$ *of Definition 7.392 with error* $\mathcal{O}(h^2) + \mathcal{O}(\tau^2)$. *Moreover,*

$$\min_{k \in \mathcal{I}_{\mathrm{IR}}^{(h)}} \widehat{\varsigma}_{\mathrm{law}}^{(\tau)}(v_k) = \min_{v \in \mathcal{I}_{\mathrm{IR}}} \varsigma_{\mathrm{law}}^{(\tau)}(v) \ + \ \mathcal{O}(h^2) + \mathcal{O}(\tau^2),$$

so the windowed minimum reported in practice is a consistent diagnostic.

Proof. Write $f := \Lambda_{\mathrm{law}}$, $v_k = v_0 + kh$. Set

$$\phi(v) := (\mathsf{LP}_\tau f)''(v), \qquad \phi_h(k) := \big(\mathsf{D}_2\, \mathsf{LP}_{\tau,h} f\big)_k.$$

By Lemma 7.381(4), differentiation commutes with LP_τ, so $\phi = \mathsf{LP}_\tau(f'')$. Also, since both D_2 and $\mathsf{LP}_{\tau,h}$ are discrete convolutions with finite support, they commute:

$$\phi_h = \mathsf{D}_2(\mathsf{LP}_{\tau,h} f) = \mathsf{LP}_{\tau,h}(\mathsf{D}_2 f).$$

Step 1: Uniform curvature consistency. Assume $f \in C^4$ in a neighborhood of the (compact) operational window.[10] Then the second–difference satisfies

$$(\mathsf{D}_2 g)(v_k) = g''(v_k) + \mathcal{O}(h^2) \quad \text{uniformly in } k \text{ whenever } g \in C^4.$$

Apply this twice with commutations:

$$\phi_h(k) = \big(\mathsf{LP}_{\tau,h}\, \mathsf{D}_2 f\big)_k = \big(\mathsf{LP}_\tau \mathsf{D}_2 f\big)(v_k) + \mathcal{O}(h^2) = \big(\mathsf{LP}_\tau f''\big)(v_k) + \mathcal{O}(h^2) = \phi(v_k) + \mathcal{O}(h^2).$$

[10] This is the standard smoothness for the second–difference truncation error; see also the discussion under Proposition 7.410.

If one wishes to compare ϕ back to the *unsmoothed* curvature f'', then by the bias expansion (the second–derivative version of Proposition 7.382, using evenness and finite $m_2(W)$)

$$\phi(v) - f''(v) = \frac{\tau^2 m_2(W)}{2}\, f^{(4)}(v) + \mathcal{O}(\tau^4),$$

so overall

$$\sup_k \left|\phi_h(k) - \phi(v_k)\right| = \mathcal{O}(h^2), \qquad \sup_k \left|\phi_h(k) - f''(v_k)\right| = \mathcal{O}(h^2) + \mathcal{O}(\tau^2).$$

Step 2: Set approximation for the curvature test. The continuous window is $\mathcal{I}_{\mathrm{IR}} = \{\, v : |\phi(v)| \le \varepsilon_{\mathrm{curv}} \,\}$. Define the grid–indexed window by the curvature test $\mathcal{I}_{\mathrm{IR}}^{(h)} = \{\, k : \ |\phi_h(k)| \le \varepsilon_{\mathrm{curv}} \,\}$. Let $\delta_\phi := \sup_k |\phi_h(k) - \phi(v_k)| = \mathcal{O}(h^2)$ from Step 1. Then for all k,

$$|\phi(v_k)| \le \varepsilon_{\mathrm{curv}} - \delta_\phi \implies k \in \mathcal{I}_{\mathrm{IR}}^{(h)}, \qquad k \in \mathcal{I}_{\mathrm{IR}}^{(h)} \implies |\phi(v_k)| \le \varepsilon_{\mathrm{curv}} + \delta_\phi.$$

Viewing the grid as a partition of the line and using the Lipschitz regularity of ϕ on the operational window (cf. Section 6.12.4), the standard "band–measure'' estimate (as in Proposition 7.397, Step 4) implies that the symmetric difference between the continuous window and the stepwise-constant embedding of the discrete window has measure

$$\mu\Big(\mathcal{I}_{\mathrm{IR}} \,\triangle\, \mathcal{I}_{\mathrm{IR}}^{(h)}\Big) \ \le\ C\,\delta_\phi \ =\ \mathcal{O}(h^2).$$

If one compares against the *unsmoothed* curvature set $\{\, |f''| \le \varepsilon_{\mathrm{curv}} \,\}$, the additional mollifier bias produces the stated $\mathcal{O}(\tau^2)$ contribution.

Step 3: Consistency of the windowed minimum. Let $\psi(v) := \varsigma_{\mathrm{law}}^{(\tau)}(v) = (\mathsf{LP}_\tau f)'(v)$ and $\widehat{\psi}_k := \widehat{\varsigma}_{\mathrm{law}}^{(\tau)}(v_k)$. By Proposition 7.413 we have the uniform approximation

$$\sup_k |\widehat{\psi}_k - \psi(v_k)| \le C\,h^2 \quad (\text{and } +\ \mathcal{O}(\tau^2) \text{ if compared to } f').$$

Let $k^\star \in \mathcal{I}_{\mathrm{IR}}^{(h)}$ realize the discrete minimum. Then

$$\min_{k \in \mathcal{I}_{\mathrm{IR}}^{(h)}} \widehat{\psi}_k = \widehat{\psi}_{k^\star} = \psi(v_{k^\star}) + \mathcal{O}(h^2) \ \ge\ \min_{v \in \mathcal{I}_{\mathrm{IR}}} \psi(v) + \mathcal{O}(h^2),$$

where we used that the discrete window is embedded in a small $\mathcal{O}(h^2)$-band around the continuous window (Step 2) and ψ is locally Lipschitz on the operational window (from Section 6.12.4 and Lemma 7.386). For the reverse inequality, take $v^\star \in \mathcal{I}_{\mathrm{IR}}$ achieving the minimum of ψ, pick a nearest grid index k (within distance $\le h$), and note that $k \in \mathcal{I}_{\mathrm{IR}}^{(h)}$ once h is small enough by Step 2; then

$$\min_{k \in \mathcal{I}_{\mathrm{IR}}^{(h)}} \widehat{\psi}_k \ \le\ \widehat{\psi}_k = \psi(v_k) + \mathcal{O}(h^2) \ =\ \psi(v^\star) + \mathcal{O}(h^2) \ =\ \min_{v \in \mathcal{I}_{\mathrm{IR}}} \psi(v) + \mathcal{O}(h^2).$$

Combining both directions yields

$$\min_{k \in \mathcal{I}_{\mathrm{IR}}^{(h)}} \widehat{\varsigma}_{\mathrm{law}}^{(\tau)}(v_k) = \min_{v \in \mathcal{I}_{\mathrm{IR}}} \varsigma_{\mathrm{law}}^{(\tau)}(v) \ +\ \mathcal{O}(h^2),$$

and the additional $\mathcal{O}(\tau^2)$ term appears only if the minimum is compared against the *unsmoothed* slope $f' = \Lambda'_{\mathrm{law}}$ (cf. Proposition 7.382). □

Note. The $\mathcal{O}(\tau^2)$ contributions above reflect the *continuous* low–pass bias relative to the raw (unsmoothed) derivative or curvature. If one measures both the window and the minimum against the robust (smoothed) quantities $(\mathsf{LP}_\tau f)''$ and $(\mathsf{LP}_\tau f)'$, then the discrete errors are $\mathcal{O}(h^2)$ alone.

Proposition 7.418 (Gauge/presentation independence; pipeline comparability)**.** *With BRST-closed tags and calibrated branches, the estimator* $\widehat{\varsigma}^{(\tau)}_{\mathrm{law}}$ *and the discrete window* $\mathcal{I}^{(h)}_{\mathrm{IR}}$ *are independent of gauge parameters and presentation-equivalent choices, and are comparable across pipeline moves* $(\lambda_{\mathrm{UV}}, \ell) \mapsto (\tilde{\lambda}_{\mathrm{UV}}, \tilde{\ell})$ *up to the plateau tolerances, with discretization errors* $\mathcal{O}(h^2) + \mathcal{O}(\tau^2)$.

Proof. Preliminaries and notation. Let $f := \Lambda_{\mathrm{law}}$, and recall the discrete operators $\mathsf{LP}_{\tau,h}$ and D_0, and the estimator $\widehat{\varsigma}^{(\tau)}_{\mathrm{law}} = \mathsf{D}_0\,\mathsf{LP}_{\tau,h} f$. Work on the eroded index set so all stencil samples exist (cf. Remark 7.416).

Step 1: Gauge/presentation independence (discrete). By Propositions 7.367 and 7.372, once branches are calibrated and tags are BRST–closed, $f = \Lambda_{\mathrm{law}}$ is independent (on certified plateaus) of gauge parameters and presentation–equivalent choices. The discrete map $f \mapsto \widehat{\varsigma}^{(\tau)}_{\mathrm{law}}$ is linear, grid-local, and depends only on the sampled values of f through fixed, gauge–independent coefficients $\{w_j\}$. Hence $\widehat{\varsigma}^{(\tau)}_{\mathrm{law}}$ inherits the same independence.

For the discrete IR window, use Corollary 7.417: the grid-defined set $\mathcal{I}^{(h)}_{\mathrm{IR}} = \{\, k : |\mathsf{D}_2(\mathsf{LP}_{\tau,h} f)_k| \le \varepsilon_{\mathrm{curv}} \,\}$ approximates the continuous window $\mathcal{I}_{\mathrm{IR}} = \{\, v : |(\mathsf{LP}_\tau f)''(v)| \le \varepsilon_{\mathrm{curv}} \,\}$. Since $\mathcal{I}_{\mathrm{IR}}$ is gauge/presentation independent by Proposition 7.397, the discrete window $\mathcal{I}^{(h)}_{\mathrm{IR}}$ inherits this independence up to $\mathcal{O}(h^2) + \mathcal{O}(\tau^2)$.

Step 2: Pipeline comparability for the estimator (discrete). Let $\tilde{f}(v) = \Lambda_{\mathrm{law}}(v; \tilde{\lambda}_{\mathrm{UV}}, \tilde{\ell})$. By Lemma 7.412(1),

$$\widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(\cdot;\tilde{\cdot}) - \widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(\cdot;\cdot) = \mathsf{LP}_{\tau,h}\Big(\mathsf{D}_0\big(\tilde{f} - f\big)\Big).$$

Discrete Young's inequality (convolution $\ell^1 \to \ell^\infty$) and $\|\mathsf{LP}_{\tau,h}\|_{\ell^\infty \to \ell^\infty} = \|w\|_{\ell^1}$ give

$$\Big\|\widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(\cdot;\tilde{\cdot}) - \widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(\cdot;\cdot)\Big\|_{\ell^\infty} \;\le\; \|w\|_{\ell^1}\,\Big\|\mathsf{D}_0(\tilde{f} - f)\Big\|_{\ell^\infty}.$$

Central differences approximate derivatives with $\mathcal{O}(h^2)$, hence

$$\Big\|\mathsf{D}_0(\tilde{f} - f)\Big\|_{\ell^\infty} \;\le\; \|\tilde{f}' - f'\|_{L^\infty} \;+\; C\,h^2\,\|f^{(3)}\|_{L^\infty},$$

uniformly on the eroded set. By the continuous pipeline comparability bound (Proposition 7.390 together with Corollary 7.378), $\|\tilde{f}' - f'\|_{L^\infty}$ is bounded by a constant multiple of the plateau budgets $(\varepsilon_{\mathrm{grad}}, \varepsilon_{\mathrm{osc}}, \varepsilon_{\mathrm{reg}}, \varepsilon_{\mathrm{swap}})$. Therefore

$$\Big\|\widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(\cdot;\tilde{\cdot}) - \widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(\cdot;\cdot)\Big\|_{\ell^\infty} \;\le\; C\,(\text{plateau tolerances}) \;+\; C\,h^2.$$

If one prefers to compare via the continuous robust slope $\varsigma^{(\tau)} = (\mathsf{LP}_\tau f)'$, use the consistency $\|\widehat{\varsigma}^{(\tau)}_{\mathrm{law}} - \varsigma^{(\tau)}_{\mathrm{law}}\|_\infty = \mathcal{O}(h^2) + \mathcal{O}(\tau^2)$ from Proposition 7.413, obtaining the same tolerance bound plus $\mathcal{O}(h^2) + \mathcal{O}(\tau^2)$.

Step 3: Pipeline comparability for the discrete IR window. By Corollary 7.417, $\mathcal{I}_{\mathrm{IR}}^{(h)}$ is an $\mathcal{O}(h^2)+\mathcal{O}(\tau^2)$ approximation of the continuous window $\mathcal{I}_{\mathrm{IR}}$. Apply the continuous window comparability bound of Proposition 7.397 to the two pipelines and then transfer back to the discrete index set via the grid embedding (as in the proof of Corollary 7.417, Step 2). This yields that the symmetric difference of the discrete windows (measured either in index count times h, or via stepwise-constant embeddings) is controlled by a constant times the plateau tolerances, up to $\mathcal{O}(h^2)+\mathcal{O}(\tau^2)$.

Combining Steps 1–3 proves the proposition: both the discrete estimator and the discrete IR window are gauge/presentation independent on plateaus and are comparable across admissible pipeline moves, up to the plateau budgets, with discretization errors $\mathcal{O}(h^2)+\mathcal{O}(\tau^2)$. □

Remark 7.419 (Explicit estimator comparability bound)**.** On the eroded index set (so all stencil samples exist), the intermediate estimate from Step 2 of the proof of Proposition 7.418 reads

$$\left\|\widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(\cdot;\tilde{\cdot})-\widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(\cdot;\cdot)\right\|_{\ell^\infty} \;\le\; \|w\|_{\ell^1}\left\|\Lambda'_{\mathrm{law}}(\tilde{\cdot})-\Lambda'_{\mathrm{law}}(\cdot)\right\|_{L^\infty} \;+\; C\,(h^2+\tau^2),$$

where $\|w\|_{\ell^1}:=\sum_{j=-m}^{m}|w_j|$ (in particular $\|w\|_{\ell^1}=1$ if $w_j\ge 0$ and $\sum_j w_j=1$). The first term on the right-hand side is controlled by the pipeline comparability of visibilities (Corollary 7.378); the $\mathcal{O}(h^2+\tau^2)$ term comes from the discrete consistency error (Proposition 7.413).

Monotonicity and false–positive guard.

Definition 7.420 (IR monotonicity/false–positive guard)**.** Let $\mathcal{I}_{\mathrm{IR}}$ be an IR window as in Definition 7.392, and set $\psi(v):=\varsigma^{(\tau)}_{\mathrm{law}}(v)=\frac{\mathrm{d}}{\mathrm{d}v}(\mathsf{LP}_\tau\Lambda_{\mathrm{law}})(v)$. We say that $\mathcal{I}_{\mathrm{IR}}$ passes the *monotonicity/false–positive guard* if

$$\boxed{\sup_{v\in\mathcal{I}_{\mathrm{IR}}}\left|(\mathsf{LP}_\tau\Lambda_{\mathrm{law}})''(v)\right|\le\varepsilon_{\mathrm{curv}},\qquad \mathrm{TV}_{\mathcal{I}_{\mathrm{IR}}}[\psi]:=\int_{\mathcal{I}_{\mathrm{IR}}}|\psi'(v)|\,\mathrm{d}v\;\le\;\varepsilon_{\mathrm{tv}}.}\tag{7.548}$$

Remark 7.421 (Total variation (TV))**.** Throughout Chapter 7, "TV" abbreviates the *total variation* of the low–pass slope $\psi(v):=\varsigma^{(\tau)}_{\mathrm{law}}(v)=\frac{d}{dv}(\mathsf{LP}_\tau\Lambda_{\mathrm{law}})(v)$ (cf. Definitions 7.380 and 7.385 and Lemma 7.386). The *continuous* TV over an IR interval $\mathcal{I}_{\mathrm{IR}}\subset\mathbb{R}$ is introduced in Definition 7.420 and used in (7.548):

$$\mathrm{TV}_{\mathcal{I}_{\mathrm{IR}}}[\psi]:=\int_{\mathcal{I}_{\mathrm{IR}}}\left|\psi'(v)\right|dv,$$

which equals the standard BV definition $\sup_{\{v_i\}}\sum_i|\psi(v_i)-\psi(v_{i-1})|$ for absolutely continuous ψ (guaranteed here by low–pass smoothing; cf. Lemma 7.381).

For *discrete* guards on a grid $v_k=v_0+k\,h$, we use the discrete TV (introduced with the discrete monotonicity/false–positive guard in Definition 7.438 and applied in Proposition 7.442):

$$\mathrm{TV}_{h;[k_R,\infty)}[\psi]\;:=\;\sum_{j\ge k_R}\left|\psi_{j+1}-\psi_j\right|,\qquad \psi_j:=\psi(v_j).$$

For smooth ψ, $\mathrm{TV}_{h;[k_R,\infty)}[\psi]$ is a consistent Riemann–sum approximation of $\mathrm{TV}_{\mathcal{I}_{\mathrm{IR}}}[\psi]$. Both the continuous and discrete TV enter the stability bounds of Lemma 7.423.

Remark 7.422 (Intuitive reading: total variation (TV))**.** Total variation of a function f on $[a,b]$ is the cumulative amount it changes: $\mathrm{TV}_{[a,b]}[f] = \int_a^b |f'(x)|\,dx$ when f is absolutely continuous; for a discrete sample $\{f_k\}$ it is $\mathrm{TV}_{[k_0,k_1]}[f] = \sum_{k=k_0}^{k_1-1} |f_{k+1} - f_k|$. If f is monotone, $\mathrm{TV} = |f(b) - f(a)|$; oscillations increase TV because up–down increments add. In this chapter we apply TV to the low–pass slope $\psi(v) = \varsigma_{\mathrm{law}}^{(\tau)}(v) = \frac{d}{dv}\big(\mathsf{LP}_\tau \Lambda_{\mathrm{law}}\big)(v)$ on an IR interval $\mathcal{I}_{\mathrm{IR}}$ (see Definition 7.420), so a small $\mathrm{TV}_{\mathcal{I}_{\mathrm{IR}}}[\psi]$ certifies a stable, nearly monotone slope (no spurious sign flips) and guards against false positives. Heuristically, for $f(v) = A\sin(\omega v)$ on a long interval, TV scales like $A\,\omega$ times the interval length; low–pass filtering (LP_τ) suppresses the high–frequency part, reducing TV and stabilising the trend. In short, TV measures cumulative wiggliness; combined with the curvature bound it yields a robust, scale–aware monotonicity test for ψ (cf. Lemma 7.423).

Lemma 7.423 (Stability from curvature and TV guards)**.** *Let $\mathcal{I}_{\mathrm{IR}}$ and $\psi(v) := \varsigma_{\mathrm{law}}^{(\tau)}(v) = \frac{d}{dv}(\mathsf{LP}_\tau \Lambda_{\mathrm{law}})(v)$ be as in Definition 7.420 (cf. Definition 7.385 and Lemma 7.386). If $\mathcal{I}_{\mathrm{IR}}$ satisfies the guard (7.548), then:*

(i) Lipschitz control

For all $v_1, v_2 \in \mathcal{I}_{\mathrm{IR}}$,

$$|\psi(v_2) - \psi(v_1)| \;\le\; \varepsilon_{\mathrm{curv}}\,|v_2 - v_1|.$$

(ii) Total–variation control

For all $v \in \mathcal{I}_{\mathrm{IR}}$ and any $v_0 \in \mathcal{I}_{\mathrm{IR}}$,

$$|\psi(v) - \psi(v_0)| \;\le\; \mathrm{TV}_{\mathcal{I}_{\mathrm{IR}}}[\psi] \;\le\; \varepsilon_{\mathrm{tv}}.$$

(iii) Positivity stability

If, in addition, $\psi(v) \ge \varsigma_{\min} > 0$ for all $v \in \mathcal{I}_{\mathrm{IR}}$ (cf. Definition 7.392) and $\varepsilon_{\mathrm{tv}} < \varsigma_{\min}$, then

$$\inf_{v \in \mathcal{I}_{\mathrm{IR}}} \psi(v) \;\ge\; \varsigma_{\min} - \varepsilon_{\mathrm{tv}} \;>\; 0,$$

so ψ cannot change sign inside $\mathcal{I}_{\mathrm{IR}}$.

Proof. Preliminaries. By Definition 7.385, $\psi = \frac{d}{dv}(\mathsf{LP}_\tau \Lambda_{\mathrm{law}})$. Differentiating once more and using the derivative–commutation property of LP_τ (see Lemma 7.381(4) and Lemma 7.386), one has $\psi' = (\mathsf{LP}_\tau \Lambda_{\mathrm{law}})''$ on $\mathcal{I}_{\mathrm{IR}}$. The curvature guard in (7.548) asserts $\sup_{\mathcal{I}_{\mathrm{IR}}} |\psi'| \le \varepsilon_{\mathrm{curv}}$, while the TV guard gives $\mathrm{TV}_{\mathcal{I}_{\mathrm{IR}}}[\psi] = \int_{\mathcal{I}_{\mathrm{IR}}} |\psi'| \le \varepsilon_{\mathrm{tv}}$.

(1) *Lipschitz control.* For any $v_1, v_2 \in \mathcal{I}_{\mathrm{IR}}$,

$$\psi(v_2) - \psi(v_1) = \int_{v_1}^{v_2} \psi'(s)\,ds,$$

hence $|\psi(v_2) - \psi(v_1)| \le \int_{v_1}^{v_2} |\psi'(s)|\,ds \le \sup_{\mathcal{I}_{\mathrm{IR}}} |\psi'|\,|v_2 - v_1| \le \varepsilon_{\mathrm{curv}} |v_2 - v_1|$.

(2) *Total–variation control.* For any $v, v_0 \in \mathcal{I}_{\mathrm{IR}}$,

$$|\psi(v) - \psi(v_0)| \le \int_{[v_0, v]} |\psi'(s)|\,ds \le \int_{\mathcal{I}_{\mathrm{IR}}} |\psi'(s)|\,ds = \mathrm{TV}_{\mathcal{I}_{\mathrm{IR}}}[\psi] \le \varepsilon_{\mathrm{tv}},$$

using the TV guard in (7.548). Absolute continuity of ψ needed for the line integral follows from $\psi = \mathsf{LP}_\tau(\Lambda'_{\mathrm{law}})$ and Lemma 7.386 (with Λ_{law} locally absolutely continuous per the standing regularity assumptions used in Lemma 7.386 and Proposition 7.383).

(3) *Positivity stability.* Under Definition 7.392, $\psi(v) \geq \varsigma_{\min}$ on $\mathcal{I}_{\mathrm{IR}}$. Fix any $v_0 \in \mathcal{I}_{\mathrm{IR}}$; then by (2), for all $v \in \mathcal{I}_{\mathrm{IR}}$,

$$\psi(v) \;\geq\; \psi(v_0) - \varepsilon_{\mathrm{tv}} \;\geq\; \varsigma_{\min} - \varepsilon_{\mathrm{tv}}.$$

If $\varepsilon_{\mathrm{tv}} < \varsigma_{\min}$, the right–hand side is strictly positive, hence ψ cannot cross or touch zero on $\mathcal{I}_{\mathrm{IR}}$. Taking the infimum over v yields the stated bound. □

Proposition 7.424 (Gauge/presentation safety of the guard)**.** *Let $\mathcal{I}_{\mathrm{IR}}$ be an IR window as in Definition 7.392 and let the guard quantities be those in Definition 7.420, namely*

$$\sup_{v\in\mathcal{I}_{\mathrm{IR}}} \left|(\mathsf{LP}_\tau \Lambda_{\mathrm{law}})''(v)\right| \quad \textit{and} \quad \mathrm{TV}_{\mathcal{I}_{\mathrm{IR}}}[\psi], \quad \psi := \varsigma^{(\tau)}_{\mathrm{law}} = \tfrac{d}{dv}(\mathsf{LP}_\tau \Lambda_{\mathrm{law}}).$$

Under BRST–closed tags and calibrated branches, these guard quantities:

(i) are independent *of gauge parameters and presentation–equivalent choices on certified plateaus (Propositions 7.367 and 7.372);*

(ii) are comparable across admissible pipeline moves $(\lambda_{\mathrm{UV}}, \ell) \mapsto (\tilde\lambda_{\mathrm{UV}}, \tilde\ell)$ *in the sense that on the eroded intersection window $(\mathcal{I}_{\mathrm{IR}} \cap \tilde{\mathcal{I}}_{\mathrm{IR}})^{-\tau R}$ (cf. Remark 7.391),*

$$\boxed{\begin{aligned} \left|\sup_J|\psi'_\cdot| - \sup_J|\psi'_\cdot|\right| &\leq \|W\|_1 \sup_J \left|\Lambda''_{\mathrm{law}}(\,\cdot\,;\tilde\cdot) - \Lambda''_{\mathrm{law}}(\,\cdot\,;\cdot)\right|, && (7.549)\\ \left|\,\mathrm{TV}_J[\psi_\cdot] - \mathrm{TV}_J[\psi_\cdot]\,\right| &\leq \mu(J)\,\|W\|_1 \sup_J \left|\Lambda''_{\mathrm{law}}(\,\cdot\,;\tilde\cdot) - \Lambda''_{\mathrm{law}}(\,\cdot\,;\cdot)\right|, && (7.550)\end{aligned}}$$

where $J := (\mathcal{I}_{\mathrm{IR}} \cap \tilde{\mathcal{I}}_{\mathrm{IR}})^{-\tau R}$ and $\mu(J)$ is its Lebesgue measure. In particular, the right–hand sides are controlled by the plateau budgets via Corollary 7.378 and Proposition 7.397.

If $W \geq 0$ and $\int W = 1$ (cf. eq. (7.538)), then $\|W\|_1 = 1$ by Lemma 7.381(1).

Proof. Preliminaries. By Definition 7.385 and Lemma 7.386,

$$\psi = \mathsf{LP}_\tau(\Lambda'_{\mathrm{law}}), \qquad \psi' = (\mathsf{LP}_\tau \Lambda_{\mathrm{law}})'' = \mathsf{LP}_\tau(\Lambda''_{\mathrm{law}}).$$

The operator LP_τ is convolution by W_τ (Definition 7.380), hence bounded on L^∞ with $\|\mathsf{LP}_\tau g\|_\infty \leq \|W_\tau\|_1 \|g\|_\infty = \|W\|_1 \|g\|_\infty$ (Lemma 7.381(1)) and commutes with d/dv on the stated regularity class (Lemma 7.381(4)).

(1) *Gauge/presentation independence.* On certified plateaus, Λ_{law} is invariant under gauge–parameter variations and presentation–equivalent choices once calibration is fixed (Propositions 7.367 and 7.372). Thus, for any such variation ∂_ξ,

$$\partial_\xi \Lambda_{\mathrm{law}} = 0 \quad \Rightarrow \quad \partial_\xi \psi = \partial_\xi\big(\mathsf{LP}_\tau \Lambda'_{\mathrm{law}}\big) = \mathsf{LP}_\tau(\partial_\xi \Lambda'_{\mathrm{law}}) = 0,$$

and analogously $\partial_\xi \psi' = \mathsf{LP}_\tau(\partial_\xi \Lambda''_{\mathrm{law}}) = 0$, using the commutation noted above. Consequently, $\sup_{\mathcal{I}_{\mathrm{IR}}} |\psi'|$ and $\mathrm{TV}_{\mathcal{I}_{\mathrm{IR}}}[\psi] = \int_{\mathcal{I}_{\mathrm{IR}}} |\psi'|$ (by absolute continuity and $\psi' = \frac{d}{dv}\psi$) are invariant. This proves Item 1.

(2) *Pipeline comparability on a common window.* Consider an admissible move $(\lambda_{\mathrm{UV}}, \ell) \mapsto (\tilde{\lambda}_{\mathrm{UV}}, \tilde{\ell})$. By Corollary 7.378, for $k = 1, 2$ there exist constants C_k (depending only on the certified budgets) with

$$\sup_J \left|\Lambda^{(k)}_{\mathrm{law}}(\,\cdot\,;\tilde{\cdot}) - \Lambda^{(k)}_{\mathrm{law}}(\,\cdot\,;\cdot)\right| \;\le\; C_k\Big(\varepsilon_{\mathrm{grad}} + \varepsilon_{\mathrm{curv}} + \varepsilon_{\mathrm{osc}} + \varepsilon_{\mathrm{reg}} + \varepsilon_{\mathrm{swap}}\Big),$$

when $J = (\mathcal{I}_{\mathrm{IR}} \cap \tilde{\mathcal{I}}_{\mathrm{IR}})^{-\tau R}$ (cf. Remark 7.391 ensuring that $W_\tau(v - \cdot)$ samples inside certified windows). Applying LP_τ and using the L^1–L^∞ bound,

$$\sup_J \left|\psi'_{\tilde{\cdot}} - \psi'_{\cdot}\right| = \sup_J \left|\mathsf{LP}_\tau(\Lambda''_{\mathrm{law}}(\tilde{\cdot}) - \Lambda''_{\mathrm{law}}(\cdot))\right| \le \|W\|_1 \sup_J \left|\Lambda''_{\mathrm{law}}(\tilde{\cdot}) - \Lambda''_{\mathrm{law}}(\cdot)\right|.$$

This yields (7.549) after taking suprema.

For the total variation, use the triangle inequality and absolute continuity:

$$\mathrm{TV}_J[\psi_{\tilde{\cdot}}] = \int_J |\psi'_{\tilde{\cdot}}| \le \int_J |\psi'_{\cdot}| + \int_J |\psi'_{\tilde{\cdot}} - \psi'_{\cdot}| \le \mathrm{TV}_J[\psi_{\cdot}] + \mu(J) \sup_J |\psi'_{\tilde{\cdot}} - \psi'_{\cdot}|.$$

Insert the previous bound on $\sup_J |\psi'_{\tilde{\cdot}} - \psi'_{\cdot}|$ to obtain

$$\mathrm{TV}_J[\psi_{\tilde{\cdot}}] - \mathrm{TV}_J[\psi_{\cdot}] \;\le\; \mu(J)\,\|W\|_1 \sup_J \left|\Lambda''_{\mathrm{law}}(\tilde{\cdot}) - \Lambda''_{\mathrm{law}}(\cdot)\right|.$$

Swapping the roles of $(\cdot)$ and $(\tilde{\cdot})$ gives the reverse inequality and hence (7.550).

(3) *Windows under pipeline moves.* When comparing on the native windows $\mathcal{I}_{\mathrm{IR}}$ and $\tilde{\mathcal{I}}_{\mathrm{IR}}$, use Proposition 7.397 to bound the symmetric difference $\mu(\mathcal{I}_{\mathrm{IR}} \triangle \tilde{\mathcal{I}}_{\mathrm{IR}})$ by a constant times the plateau budgets. This, together with the bounds on J above, establishes comparability "up to the plateau budgets" in the proposition statement. □

Intuition. The guard depends only on ψ and ψ', i.e. on Λ'_{law} and Λ''_{law} after a fixed mollifier LP_τ. Gauge/presentation invariances of Λ_{law} on plateaus propagate through LP_τ and differentiation. Across pipeline moves, LP_τ is L^1-bounded, so differences of ψ and ψ' are controlled by differences of Λ'_{law} and Λ''_{law}, which are already bounded by the acceptance/plateau budgets.

Remark 7.425 (Discrete implementation)**.** Let $\widehat{\psi}_k := \widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(v_k)$ and define the discrete total variation on the eroded index set by

$$\mathrm{TV}^{(h)}_{\mathcal{I}^{(h)}_{\mathrm{IR}}}[\widehat{\psi}] := \sum_{k \in \mathcal{I}^{(h)}_{\mathrm{IR}} \setminus \{k_{\max}\}} \left|\widehat{\psi}_{k+1} - \widehat{\psi}_k\right|.$$

By Proposition 7.413 and Corollary 7.417, $\mathrm{TV}^{(h)}_{\mathcal{I}^{(h)}_{\mathrm{IR}}}[\widehat{\psi}] \to \mathrm{TV}_{\mathcal{I}_{\mathrm{IR}}}[\psi]$ with error $\mathcal{O}(h^2) + \mathcal{O}(\tau^2)$, and $\min_{k \in \mathcal{I}^{(h)}_{\mathrm{IR}}} \widehat{\psi}_k \to \min_{v \in \mathcal{I}_{\mathrm{IR}}} \psi(v)$ at the same rate. Hence the discrete guard is a consistent proxy for (7.548).

Compatibility with identity ledger and gauge safety.

Remark 7.426 (Compatibility with identity ledger and gauge safety)**.** The law–channel visibility Λ_{law} is constructed on shell from calibrated branches (cf. eqs. (7.122) and (7.527)) with BRST/Nielsen protections in place; hence a positive, stable trend certified by Definitions 7.392 and 7.420 diagnoses the emergence of a *global* G–Ward structure rather than a gauge artefact (see Lemma 7.374 and the standard BRST accounts [117, 442, 243]). The renormalisation–group reading used in Step 3 follows the standard linearisation near IR fixed points [389, 91].

Operational outputs for Step 2. Outputs comprise: (i) plots of $v \mapsto \Lambda_{\mathrm{law}}(v)$ and $v \mapsto \varsigma^{(\tau)}_{\mathrm{law}}(v)$ with the accepted IR window $\mathcal{I}_{\mathrm{IR}}$ shaded (cf. Definitions 7.385 and 7.392); (ii) a compact table reporting $\min_{v\in\mathcal{I}_{\mathrm{IR}}} \varsigma^{(\tau)}_{\mathrm{law}}(v)$, together with the curvature and TV guard values from (7.548), and their spreads under admissible coarse–graining choices $(P_x, K^{(\ell)}_{\mathrm{IR}})$ (cf. the joint coarse–graining maps eqs. (6.364) and (6.366)); (iii) an audit record listing $(\tau, h, \{w_j\})$ as defined in Definitions 7.380, 7.398, 7.406 and 7.409. By Lemma 7.423 and Proposition 7.424, these diagnostics suffice to certify monotonicity and gauge/presentation safety on $\mathcal{I}_{\mathrm{IR}}$.

Acceptance criterion for Step 2. QIII Step 2 is accepted if there exists an IR window $\mathcal{I}_{\mathrm{IR}} \subset \mathbb{R}$ such that the positivity condition of Definition 7.392 and the monotonicity/TV guard of Definition 7.420 hold with tolerances compatible with Section 6.12.4. When accepted, the window $\mathcal{I}_{\mathrm{IR}}$ and the associated parameters $(\tau, h, \{w_j\}, P_x, K^{(\ell)}_{\mathrm{IR}})$ become the input mask for Step 3 (RG irrelevance analysis), with robustness and gauge/presentation safety inherited from Proposition 7.424.

Proposition 7.427 (Mask soundness for Step 3)**.** *Let $\mathcal{I}_{\mathrm{IR}}$ be an accepted IR window with parameters $(\tau, h, \{w_j\}, P_x, K^{(\ell)}_{\mathrm{IR}})$ from Step 2 (cf. Definitions 7.392 and 7.420). Then:*

Kernel support safety.
For every $v \in (\mathcal{I}_{\mathrm{IR}})^{-\tau R}$, the sampling kernel $W_\tau(v-\cdot)$ stays inside $\mathcal{I}_{\mathrm{IR}}$; hence all mollified quantities used in Step 3 depend only on certified data (cf. Remark 7.391).

Gauge/presentation safety.
All Step 3 readouts are invariant under gauge parameters and presentation–equivalent choices on plateaus (cf. Proposition 7.424).

Pipeline stability.
Under admissible pipeline moves, windows and readouts remain comparable up to the plateau budgets (cf. Propositions 7.397 and 7.424).

Scaling and tolerance monotonicity.
The restoration sets and their first–attainment points respect scaling covariance and tolerance monotonicity (cf. Lemma 7.433).

UV–uniform option.
For UV strips $\mathcal{U}$, the uniform restoration length obeys the bounds/monotonicity of Proposition 7.435.

Continuum–discrete coherence.

The discrete audit estimators are consistent with the continuum ones and admit the stated error rates (cf. Lemma 7.440 and Corollary 7.436).

Proof. (1) is Remark 7.391. (2)–(3) follow from Propositions 7.397 and 7.424. (4) is Lemma 7.433. (5) is Proposition 7.435. (6) is Lemma 7.440 and Corollary 7.436. □

7.8.3 Restoration Length

Objective. We turn the qualitative notion of "law healing in the IR'' into a *scale* extracted from the law–visibility Λ_{law} (cf. eq. (7.122)) and the robust slope $\varsigma_{\mathrm{law}}^{(\tau)}$ introduced in eq. (7.539). Throughout we keep the state coherent (uniformly in the volume V) as in Section 6.12.4 and eq. (7.535) so that the diagnostic is purely *law*-channel.

Pointwise restoration length).

Definition 7.428 (Pointwise restoration length)**.** Let $E_{\mathrm{law}}(\ell) := 1 - \Lambda_{\mathrm{law}}(\lambda_{\mathrm{UV}}, \ell; T^\star)$ (cf. eq. (7.535)) and fix tolerances $(\varepsilon_{\mathrm{plat}}, \varsigma_{\min})$ consistent with Section 7.3.9. The *restoration length* is

$$\boxed{\ell_R := \inf\Big\{\ell \geq \ell_{\min} :\ E_{\mathrm{law}}(\ell) \leq \varepsilon_{\mathrm{plat}} \ \text{ and } \ \varsigma_{\mathrm{law}}^{(\tau)}(\ell') \geq \varsigma_{\min} \ \ \forall\, \ell' \geq \ell\Big\},} \tag{7.551}$$

with $\varsigma_{\mathrm{law}}^{(\tau)}$ as in eq. (7.539) and the monotonicity/curvature guards of eq. (7.548). If the set is empty we set $\ell_R := +\infty$ and classify as non-restoring at the probed scales. The definition is invariant under reparametrisations $\ell \mapsto a\,\ell$ (cf. eq. (7.541)).

Intuition. ℓ_R is the first coarse-graining scale beyond which the law-deficit stays within the plateau tolerance while the robust slope remains above threshold, i.e. the law keeps healing monotonically in the IR.

Robust variant (mollified deficit).

Definition 7.429 (Robust restoration length (mollified deficit))**.** To desensitise E_{law} to high–frequency readout ripples, work with the smoothed deficit

$$E_{\mathrm{law}}^{(\tau_E)}(\ell) := 1 - (\mathsf{LP}_{\tau_E}\Lambda_{\mathrm{law}})(\ell),$$

with kernel properties as in eqs. (7.537) and (7.538). Define

$$\boxed{\ell_R^{(\tau_E)} := \inf\Big\{\ell \geq \ell_{\min} :\ E_{\mathrm{law}}^{(\tau_E)}(\ell) \leq \varepsilon_{\mathrm{plat}} \ \text{ and } \ \varsigma_{\mathrm{law}}^{(\tau)}(\ell') \geq \varsigma_{\min} \ \ \forall\, \ell' \geq \ell\Big\}.} \tag{7.552}$$

Here $\varsigma_{\mathrm{law}}^{(\tau)}$ is as in eq. (7.539), with the monotonicity/curvature guards of eq. (7.548). These guards ensure that $\ell_R^{(\tau_E)}$ agrees with the pointwise definition eq. (7.551) up to $\mathcal{O}(\tau_E^2)$ bias; admissibility of the smoothing and exchange of limits follows from Kato continuity [115].

Intuition. Light mollification suppresses readout ripples without moving the attainment scale: $\ell_R^{(\tau_E)}$ tracks ℓ_R to second order in the smoothing width, provided the curvature/TV guards hold.

Uniformity across UV slices.

Definition 7.430 (Uniform restoration length over UV strips)**.** Because QIII certification must not hinge on a special UV projector, require *uniform* restoration over an admissible UV strip $\mathcal{U} = [u_{\min}, u_{\max}]$ in $u := \ln \lambda_{\mathrm{UV}}$ (cf. Section 6.12.4 and eq. (7.67)). Define

$$\boxed{\ell_R^{[\mathcal{U}]} := \inf\Big\{\ell \geq \ell_{\min} : \sup_{u\in\mathcal{U}} E_{\mathrm{law}}(\ell; u) \leq \varepsilon_{\mathrm{plat}} \quad \text{and} \quad \inf_{\substack{u\in\mathcal{U}\\ \ell'\geq\ell}} \varsigma_{\mathrm{law}}^{(\tau)}(\ell'; u) \;\geq\; \varsigma_{\min}\Big\}.} \tag{7.553}$$

Intuition. The same ℓ must certify restoration for *every* admissible UV slice in $\mathcal{U}$; by construction $\ell_R^{[\mathcal{U}]} \geq \sup_{u\in\mathcal{U}} \ell_R(u)$ and it reduces to eq. (7.551) when $|\mathcal{U}| = 0$. Moreover, if $\mathcal{U}_1 \subseteq \mathcal{U}_2$ then $\ell_R^{[\mathcal{U}_2]} \geq \ell_R^{[\mathcal{U}_1]}$, and the definition is invariant under $\ell \mapsto a\,\ell$ (cf. eq. (7.541)).

Dual RG scale.

Definition 7.431 (Restoration scale in RG units)**.** With the fixed identification $\mu = \kappa/\ell$ (recorded once for audit), define

$$\boxed{\mu_R := \kappa/\ell_R, \qquad \mu_R^{[\mathcal{U}]} := \kappa/\ell_R^{[\mathcal{U}]}.} \tag{7.554}$$

Intuition. μ_R (and its UV–uniform counterpart) is the RG scale beyond which explicitly broken law-couplings that are RG-irrelevant have become numerically suppressed (Step 4 RG analysis), in line with standard RG flow expectations [389, 91, 82].

Proposition 7.432 (Existence, first–attainment uniqueness, and stability of ℓ_R)**.** *Work with $v := \ln \ell$. Assume on an IR window $\mathcal{I}_{\mathrm{IR}} = [v_\star, \infty)$ that: (i) $E_{\mathrm{law}}(e^v) \in C^1(\mathcal{I}_{\mathrm{IR}})$, $E_{\mathrm{law}} \geq 0$, and E_{law} is nonincreasing; (ii) $\varsigma_{\mathrm{law}}^{(\tau)}(e^v) \geq \varsigma_{\min} > 0$ for all $v \in \mathcal{I}_{\mathrm{IR}}$ (as in Definition 7.392); (iii) the curvature/TV guards (7.548) hold on $\mathcal{I}_{\mathrm{IR}}$. Then the admissible set in eq. (7.551) is nonempty, its infimum is attained, and ℓ_R is the unique* first *scale at which $E_{\mathrm{law}}(\ell) \leq \varepsilon_{\mathrm{plat}}$ and the slope constraint holds thereafter. Moreover, ℓ_R depends continuously on $(\varepsilon_{\mathrm{plat}}, \varsigma_{\min})$ and on E_{law} under C^0 perturbations dominated by the curvature guard (Kato–type continuity).*

Proof. Step 0 (link to the law deficit and slope). In QIII the "law deficit" is $D(v) := 1 - (\mathsf{LP}_\tau \Lambda_{\mathrm{law}})(v)$ (cf. Lemma 7.395), so

$$D'(v) = -\varsigma_{\mathrm{law}}^{(\tau)}(v) =: -\psi(v) \quad \text{with} \quad \psi(v) = \tfrac{d}{dv}(\mathsf{LP}_\tau \Lambda_{\mathrm{law}})(v) \text{ from Definition 7.385.}$$

When $E_{\mathrm{law}} = D$, item (ii) reads $\psi \geq \varsigma_{\min} > 0$ on $\mathcal{I}_{\mathrm{IR}}$, and (7.548) gives $\sup_{\mathcal{I}_{\mathrm{IR}}} |\psi'| \leq \varepsilon_{\mathrm{curv}}$ and $\mathrm{TV}_{\mathcal{I}_{\mathrm{IR}}}[\psi] \leq \varepsilon_{\mathrm{tv}}$ (see also Lemma 7.423).

Step 1 (nonemptiness). By (i), E_{law} is bounded below and nonincreasing, hence the limit $E_\infty := \lim_{v\to\infty} E_{\mathrm{law}}(e^v) \geq 0$ exists. If $E_\infty > \varepsilon_{\mathrm{plat}}$ then the admissible set in eq. (7.551) is empty and, by definition, $\ell_R = +\infty$. Otherwise there exists $v_1 \geq v_\star$ with $E_{\mathrm{law}}(e^{v_1}) \leq \varepsilon_{\mathrm{plat}}$. Item (ii) ensures the slope constraint $\psi \geq \varsigma_{\min}$ holds on all of $\mathcal{I}_{\mathrm{IR}}$ (and, more generally, cannot be violated within the window by Lemma 7.423(3)). Hence every $v \geq v_1$ satisfies both constraints in eq. (7.551), so the admissible set is nonempty.

Step 2 (attainment of the infimum). The sublevel set $\{\, v \in \mathcal{I}_{\mathrm{IR}} : E_{\mathrm{law}}(e^v) \le \varepsilon_{\mathrm{plat}} \,\}$ is closed by continuity (i), and intersecting with the ray $[v_\star, \infty)$ yields a closed ray $[v_R, \infty)$. Since the slope constraint is already ensured on $\mathcal{I}_{\mathrm{IR}}$ by (ii), the admissible set equals $[v_R, \infty)$. Therefore its infimum is attained at v_R, and setting $\ell_R := e^{v_R}$ produces the claimed attainment.

Step 3 (first–attainment uniqueness). Nonincreasing E_{law} implies that once the inequality $E_{\mathrm{law}}(e^v) \le \varepsilon_{\mathrm{plat}}$ holds, it holds for all larger v. Thus v_R is the unique *first* point where the sublevel set begins (no earlier point can satisfy the bound), giving the stated uniqueness. Equivalently, when $E_{\mathrm{law}} = D$ one has $D'(v) = -\psi(v) \le -\varsigma_{\min} < 0$ by (ii), so D is strictly decreasing on $\mathcal{I}_{\mathrm{IR}}$ and the crossing with the level $\varepsilon_{\mathrm{plat}}$ is unique.

Step 4 (stability / Kato–type continuity). Consider a perturbation $E_{\mathrm{law}} \mapsto \widetilde{E}_{\mathrm{law}} = E_{\mathrm{law}} + \delta E$ with $\|\delta E\|_{C^0(\mathcal{I}_{\mathrm{IR}})} \le \eta$, preserving (ii)–(iii) (this is ensured by Proposition 7.424 together with the curvature/TV guard). Let v_R and $\tilde{v}_R$ be the corresponding first–attainment points. By the mean–value theorem applied to D (or more generally to E_{law} identified with D in Step 0),

$$\varepsilon_{\mathrm{plat}} - E_{\mathrm{law}}(e^{\tilde{v}_R}) = D(e^{v_R}) - D(e^{\tilde{v}_R}) = -\psi(\xi)\,(\tilde{v}_R - v_R) \quad \text{for some } \xi \in [v_R, \tilde{v}_R],$$

hence $|\tilde{v}_R - v_R| \le \frac{|\widetilde{E}_{\mathrm{law}}(e^{\tilde{v}_R}) - E_{\mathrm{law}}(e^{\tilde{v}_R})|}{\psi(\xi)} \le \eta/\varsigma_{\min}$ using (ii). The curvature bound $\sup_{\mathcal{I}_{\mathrm{IR}}} |\psi'| \le \varepsilon_{\mathrm{curv}}$ (from (7.548)) provides a Lipschitz control on ψ (Lemma 7.423(1)), ensuring that linearization errors in the above argument remain uniformly controlled on compact subintervals of $\mathcal{I}_{\mathrm{IR}}$. Therefore v_R (and thus $\ell_R = e^{v_R}$) depends continuously on $(\varepsilon_{\mathrm{plat}}, \varsigma_{\min})$ and on E_{law} under C^0 perturbations dominated by the guard; this is the asserted Kato–type continuity.

All statements persist under admissible pipeline moves by Proposition 7.424 and Remark 7.391, which keep the guards and windows comparable up to the plateau budgets. □

Intuition. Inside the IR window, the guard (7.548) and the positivity condition of Definition 7.392 yield a uniformly positive robust slope $\psi = \varsigma^{(\tau)}_{\mathrm{law}}$ with controlled curvature/TV, so the law deficit $D(v) = 1 - (\mathsf{LP}_\tau \Lambda_{\mathrm{law}})(v)$ decreases monotonically: $D'(v) = -\psi(v) \le -\varsigma_{\min}$ (Definition 7.385 and Lemmas 7.395 and 7.423). Consequently, the first entry into the tolerance band $D \le \varepsilon_{\mathrm{plat}}$ is unique and attained at v_R (hence $\ell_R = e^{v_R}$); see Proposition 7.432. Moreover, perturbations upstream (gauge/presentation or admissible pipeline moves) change ψ and ψ' only within the plateau budgets, so v_R shifts by at most a controlled amount (Kato-type continuity), cf. Proposition 7.424 and Remark 7.391.

Lemma 7.433 (Scaling covariance and tolerance monotonicity)**.** *Let the restoration set be*

$$\mathcal{A}(\varepsilon_{\mathrm{plat}}, \varsigma_{\min}) := \Big\{\, v \in \mathcal{I}_{\mathrm{IR}} \;:\; E_{\mathrm{law}}(e^v) \le \varepsilon_{\mathrm{plat}} \;\; \textit{and} \;\; \psi(v) \ge \varsigma_{\min} \,\Big\}, \qquad \psi(v) := \varsigma^{(\tau)}_{\mathrm{law}}(v),$$

and define $v_R := \inf \mathcal{A}(\varepsilon_{\mathrm{plat}}, \varsigma_{\min})$ *with* $\ell_R := e^{v_R}$ *as in eq. (7.551). Then:*

Scaling covariance in v; invariance of the physical ℓ_R.

For any $a > 0$, under $\ell \mapsto a\,\ell$ one has $v \mapsto v + \ln a$ and

$$\psi(\,\cdot\,) \;\mapsto\; \psi(\,\cdot\, + \ln a) \quad \textit{and} \quad \mathcal{I}_{\mathrm{IR}} \;\mapsto\; \mathcal{I}_{\mathrm{IR}} + \ln a,$$

by Proposition 7.387 and Lemma 7.393. Consequently,

$$\mathcal{A}(\varepsilon_{\mathrm{plat}}, \varsigma_{\min}) \;\mapsto\; \mathcal{A}(\varepsilon_{\mathrm{plat}}, \varsigma_{\min}) - \ln a, \quad \textit{so} \quad v_R \;\mapsto\; v_R - \ln a.$$

Therefore the physical restoration length $\ell_R = e^{v_R}$ is invariant under $\ell \mapsto a\,\ell$. (Equivalently, in the purely numerical ℓ-coordinate without unit conversion, v-translation by $+\ln a$ sends ℓ_R to ℓ_R/a; restoring the units recovers the same physical ℓ_R.)

Tolerance monotonicity.

If $\varepsilon_{\mathrm{plat}}$ increases and/or $\varsigma_{\min}$ decreases, then v_R does not increase (hence ℓ_R does not increase). Conversely, tightening either tolerance makes v_R (and thus ℓ_R) weakly larger.

Proof. (1) The covariance statements for ψ and $\mathcal{I}_{\mathrm{IR}}$ follow from Proposition 7.387 and Lemma 7.393. Since E_{law} is a function of e^v, the inequality $E_{\mathrm{law}}(e^v) \le \varepsilon_{\mathrm{plat}}$ is likewise invariant under the translation $v \mapsto v + \ln a$. Hence the restoration set translates rigidly:

$$v \in \mathcal{A}(\varepsilon_{\mathrm{plat}}, \varsigma_{\min}) \iff v + \ln a \in \mathcal{A}(\varepsilon_{\mathrm{plat}}, \varsigma_{\min}) + \ln a,$$

so inf shifts by $-\ln a$. Passing back to $\ell = e^v$ gives $\ell'_R = e^{v_R - \ln a} = \ell_R/a$ at the level of the raw coordinate; interpreting the map $\ell \mapsto a\ell$ as a change of units recovers the same physical ℓ_R.

(2) Write explicitly

$$\mathcal{A}(\varepsilon, \sigma) = \Big(\underbrace{\{v \in \mathcal{I}_{\mathrm{IR}} : E_{\mathrm{law}}(e^v) \le \varepsilon\}}_{=:A_\varepsilon} \Big) \cap \Big(\underbrace{\{v \in \mathcal{I}_{\mathrm{IR}} : \psi(v) \ge \sigma\}}_{=:B_\sigma} \Big).$$

If $\varepsilon_2 \ge \varepsilon_1$, then $A_{\varepsilon_1} \subseteq A_{\varepsilon_2}$; if $\sigma_2 \le \sigma_1$, then $B_{\sigma_1} \subseteq B_{\sigma_2}$. Therefore $\mathcal{A}(\varepsilon_1, \sigma_1) \subseteq \mathcal{A}(\varepsilon_2, \sigma_2)$ whenever $\varepsilon_2 \ge \varepsilon_1$ and $\sigma_2 \le \sigma_1$, which yields

$$\inf \mathcal{A}(\varepsilon_2, \sigma_2) \le \inf \mathcal{A}(\varepsilon_1, \sigma_1).$$

Thus relaxing tolerances cannot increase v_R (and hence cannot increase $\ell_R = e^{v_R}$). The converse inclusion for tightened tolerances is immediate. These set–inclusion arguments apply whether $\mathcal{A}$ is empty (interpreting $\inf \varnothing = +\infty$) or nonempty; when nonempty, existence/attainment and "first" uniqueness follow from Proposition 7.432. □

Intuition. Changing the ruler (scale translation in v) shifts where the conditions are met in v but leaves the physical ℓ_R unchanged; loosening the bounds enlarges the admissible set, so the first attainment cannot move to the right.

Lemma 7.434 (Mollified vs. pointwise restoration lengths)**.** *Let LP_{τ_E} be as in Definition 7.380 with kernel moments eq. (7.538) and finite second moment $m_2(W) := \int u^2 W(u)\,du < \infty$. On an IR window $\mathcal{I}_{\mathrm{IR}}$ where $\Lambda_{\mathrm{law}} \in C^2$ and the guards (7.548) hold, the pointwise bias satisfies*

$$\boxed{\left|(\mathsf{LP}_{\tau_E}\Lambda_{\mathrm{law}})(v) - \Lambda_{\mathrm{law}}(v)\right| \le \frac{m_2(W)}{2}\,\tau_E^2 \sup_{s \in \mathcal{I}_{\mathrm{IR}}} \left|\Lambda''_{\mathrm{law}}(s)\right| + \mathcal{O}(\tau_E^4),} \tag{7.555}$$

uniformly on compact v-subsets of $\mathcal{I}_{\mathrm{IR}}$ (Proposition 7.382 and Lemma 7.381). Let v_R (resp. $v_R^{(\tau_E)}$) be the first–attainment point of the restoration set $\mathcal{A}(\varepsilon_{\mathrm{plat}}, \varsigma_{\min})$ in eq. (7.551) when

$E_{\mathrm{law}}(v) = 1 - \Lambda_{\mathrm{law}}(v)$ *(resp.* $E^{(\tau_E)}_{\mathrm{law}}(v) := 1 - (\mathsf{LP}_{\tau_E}\Lambda_{\mathrm{law}})(v)$*). If, in addition, the slope guard holds with* $\psi(v) := \varsigma^{(\tau)}_{\mathrm{law}}(v) \geq \varsigma_{\min} > 0$ *on* $\mathcal{I}_{\mathrm{IR}}$ *(Definitions 7.385 and 7.392), then*

$$\boxed{\left| v_R^{(\tau_E)} - v_R \right| \;\leq\; \frac{m_2(W)}{2}\,\frac{\tau_E^2}{\varsigma_{\min}}\,\sup_{s\in\mathcal{I}_{\mathrm{IR}}}\left|\Lambda''_{\mathrm{law}}(s)\right| \;+\; \mathcal{O}\Big(\frac{\tau_E^4}{\varsigma_{\min}}\Big).} \tag{7.556}$$

Consequently,

$$\ell_R^{(\tau_E)} = e^{v_R^{(\tau_E)}} = e^{v_R + \mathcal{O}(\tau_E^2/\varsigma_{\min})} = \ell_R\Big(1 + \mathcal{O}\big(\tfrac{\tau_E^2}{\varsigma_{\min}}\big)\Big), \tag{7.557}$$

i.e. the mollified and pointwise restoration lengths differ by a second–order amount controlled by $\tau_E^2/\varsigma_{\min}$ *(cf. Lemma 7.433 for the* $v \leftrightarrow \ell$ *translation).*

Proof. Bias bound (7.555). Apply Proposition 7.382 with $f = \Lambda_{\mathrm{law}}$: under eq. (7.538) and $m_2(W) < \infty$,

$$(\mathsf{LP}_{\tau_E}\Lambda_{\mathrm{law}})(v) = \Lambda_{\mathrm{law}}(v) + \frac{\tau_E^2 m_2(W)}{2}\,\Lambda''_{\mathrm{law}}(v) + \mathcal{O}(\tau_E^4)$$

uniformly on compact v-ranges; taking absolute values and a supremum over $s \in \mathcal{I}_{\mathrm{IR}}$ yields (7.555). (Regularity on $\mathcal{I}_{\mathrm{IR}}$ follows from the acceptance/guard pipeline and the Kato–type exchange used in Proposition 7.383.)

Shift of the first–attainment point (7.556). Let $D(v) := 1 - \Lambda_{\mathrm{law}}(v)$, $D^{(\tau_E)}(v) := 1 - (\mathsf{LP}_{\tau_E}\Lambda_{\mathrm{law}})(v)$. By (7.555), $\Delta(v) := D^{(\tau_E)}(v) - D(v)$ satisfies $\|\Delta\|_{L^\infty(\mathcal{I}_{\mathrm{IR}})} \leq C_2\,\tau_E^2 + \mathcal{O}(\tau_E^4)$ with $C_2 := \frac{m_2(W)}{2}\sup_{\mathcal{I}_{\mathrm{IR}}}|\Lambda''_{\mathrm{law}}|$. Let v_R be the (unique, attained) first solution of $D(v) \leq \varepsilon_{\mathrm{plat}}$ that also meets the slope guard on $\mathcal{I}_{\mathrm{IR}}$ (Proposition 7.432). By the slope positivity $\psi \geq \varsigma_{\min}$ on $\mathcal{I}_{\mathrm{IR}}$ (Definition 7.392) and $\psi = \frac{d}{dv}(\mathsf{LP}_\tau\Lambda_{\mathrm{law}})$ (Definition 7.385), one has $D'(v) = -\psi(v) \leq -\varsigma_{\min}$. A mean–value argument gives, for v near v_R,

$$D(v) - \varepsilon_{\mathrm{plat}} = \Big(D'(v^*)\Big)\,(v - v_R) \quad \text{for some } v^* \text{ between } v \text{ and } v_R,$$

hence $|v - v_R| \leq |D(v) - \varepsilon_{\mathrm{plat}}|/\varsigma_{\min}$. Evaluating at any v with $D^{(\tau_E)}(v) = \varepsilon_{\mathrm{plat}}$ (in particular $v = v_R^{(\tau_E)}$, the first such point for $D^{(\tau_E)}$),

$$|v_R^{(\tau_E)} - v_R| \leq \frac{|D^{(\tau_E)}(v_R^{(\tau_E)}) - D(v_R^{(\tau_E)})|}{\varsigma_{\min}} = \frac{|\Delta(v_R^{(\tau_E)})|}{\varsigma_{\min}} \leq \frac{C_2}{\varsigma_{\min}}\,\tau_E^2 + \mathcal{O}\Big(\frac{\tau_E^4}{\varsigma_{\min}}\Big),$$

which is (7.556). (Existence/uniqueness of $v_R^{(\tau_E)}$ follows as in Proposition 7.432, since the guard (7.548) is unaffected by the auxiliary smoothing LP_{τ_E}.)

From v *to* ℓ. Writing $\ell_R^{(\tau_E)} = e^{v_R^{(\tau_E)}} = e^{v_R + \delta v}$ with $|\delta v| = \mathcal{O}(\tau_E^2/\varsigma_{\min})$, a first–order expansion yields $\ell_R^{(\tau_E)} = \ell_R(1 + \mathcal{O}(\tau_E^2/\varsigma_{\min}))$, which is (7.557); cf. Lemma 7.433 for scaling covariance between v and ℓ. □

Intuition. The low–pass LP_{τ_E} preserves mass and first moment (eq. (7.538)), so with bounded curvature the bias is second order in τ_E (Proposition 7.382). A vertical perturbation of size $\mathcal{O}(\tau_E^2)$ shifts the first crossing by at most $\mathcal{O}(\tau_E^2/\varsigma_{\min})$, thanks to the uniform slope floor on $\mathcal{I}_{\mathrm{IR}}$ (Definition 7.392, Lemma 7.423, and Proposition 7.432).

Proposition 7.435 (UV–uniform restoration: bounds and monotonicity)**.** *Let $\mathcal{U} \subset \mathbb{R}$ be an admissible UV strip in $u = \ln \lambda_{\mathrm{UV}}$. For each $u \in \mathcal{U}$, let*

$$\psi(v;u) := \varsigma_{\mathrm{law}}^{(\tau)}(v;u) = \tfrac{d}{dv}\big(\mathsf{LP}_\tau \Lambda_{\mathrm{law}}\big)(v;u) \quad \textit{(cf. Definition 7.385 and Lemma 7.386)},$$

and define the per–slice admissible set

$$\mathcal{A}(u;\varepsilon_{\mathrm{plat}},\varsigma_{\min}) := \Big\{ v \in \mathcal{I}_{\mathrm{IR}}(u) \ : \ E_{\mathrm{law}}(e^v;u) \le \varepsilon_{\mathrm{plat}} \ \ \textit{and} \ \ \psi(v;u) \ge \varsigma_{\min} \Big\}$$

(cf. Definitions 7.392 and 7.420), with per–slice restoration point $v_R(u) := \inf \mathcal{A}(u;\varepsilon_{\mathrm{plat}},\varsigma_{\min})$ (so $\ell_R(u) = e^{v_R(u)}$; eq. (7.551)). Define the UV–uniform admissible set and restoration point as

$$\mathcal{A}[\mathcal{U};\varepsilon_{\mathrm{plat}},\varsigma_{\min}] := \bigcap_{u\in\mathcal{U}} \mathcal{A}(u;\varepsilon_{\mathrm{plat}},\varsigma_{\min}), \qquad v_R^{[\mathcal{U}]} := \inf \mathcal{A}[\mathcal{U};\varepsilon_{\mathrm{plat}},\varsigma_{\min}],$$

so that $\ell_R^{[\mathcal{U}]} := e^{v_R^{[\mathcal{U}]}}$ (cf. eq. (7.553)). Then:

(i) $\ell_R^{[\mathcal{U}]} \ge \sup_{u\in\mathcal{U}} \ell_R(u)$.

(ii) If $\mathcal{U}_1 \subseteq \mathcal{U}_2$ then $\ell_R^{[\mathcal{U}_2]} \ge \ell_R^{[\mathcal{U}_1]}$.

(iii) $\ell_R^{[\mathcal{U}]}$ obeys the scaling covariance and tolerance monotonicity of Lemma 7.433.

Proof. Preliminaries. By construction, $\mathcal{A}[\mathcal{U};\varepsilon,\sigma] = \bigcap_{u\in\mathcal{U}} \mathcal{A}(u;\varepsilon,\sigma)$ and $v_R^{[\mathcal{U}]} = \inf \mathcal{A}[\mathcal{U};\varepsilon,\sigma]$, while $v_R(u) = \inf \mathcal{A}(u;\varepsilon,\sigma)$ for each u. Throughout, interpret $\inf \varnothing = +\infty$.

(1) *Lower bound by the worst slice.* Let $v^\sharp := \sup_{u\in\mathcal{U}} v_R(u)$. If $v < v^\sharp$ then there exists u^* with $v < v_R(u^*)$, hence $v \notin \mathcal{A}(u^*;\varepsilon_{\mathrm{plat}},\varsigma_{\min})$. Therefore $v \notin \bigcap_{u\in\mathcal{U}} \mathcal{A}(u;\varepsilon_{\mathrm{plat}},\varsigma_{\min})$, so $v < \inf \mathcal{A}[\mathcal{U};\varepsilon_{\mathrm{plat}},\varsigma_{\min}] = v_R^{[\mathcal{U}]}$. Taking the contrapositive gives $v_R^{[\mathcal{U}]} \ge \sup_u v_R(u)$. Exponentiating yields the stated inequality for ℓ_R.

(2) *Monotonicity in the UV strip.* If $\mathcal{U}_1 \subseteq \mathcal{U}_2$, then

$$\mathcal{A}[\mathcal{U}_2;\varepsilon,\sigma] = \bigcap_{u\in\mathcal{U}_2} \mathcal{A}(u;\varepsilon,\sigma) \ \subseteq \ \bigcap_{u\in\mathcal{U}_1} \mathcal{A}(u;\varepsilon,\sigma) = \mathcal{A}[\mathcal{U}_1;\varepsilon,\sigma].$$

Hence $\inf \mathcal{A}[\mathcal{U}_2;\varepsilon,\sigma] \ge \inf \mathcal{A}[\mathcal{U}_1;\varepsilon,\sigma]$, i.e. $v_R^{[\mathcal{U}_2]} \ge v_R^{[\mathcal{U}_1]}$, and thus $\ell_R^{[\mathcal{U}_2]} \ge \ell_R^{[\mathcal{U}_1]}$.

(3) *Scaling covariance and tolerance monotonicity.* Fix $a > 0$. By Proposition 7.387 and Lemma 7.393, under $\ell \mapsto a\ell$ one has $v \mapsto v + \ln a$, $\psi(\cdot;u) \mapsto \psi(\cdot + \ln a;u)$, and $\mathcal{I}_{\mathrm{IR}}(u) \mapsto \mathcal{I}_{\mathrm{IR}}(u) + \ln a$, so each $\mathcal{A}(u;\varepsilon,\sigma)$ translates rigidly by $-\ln a$ in v. Intersections commute with translations; hence $\mathcal{A}[\mathcal{U};\varepsilon,\sigma] \mapsto \mathcal{A}[\mathcal{U};\varepsilon,\sigma] - \ln a$ and $v_R^{[\mathcal{U}]} \mapsto v_R^{[\mathcal{U}]} - \ln a$. Passing back to $\ell = e^v$ gives invariance of the *physical* $\ell_R^{[\mathcal{U}]}$, as in Lemma 7.433(1).

For tolerances, write $\mathcal{A}(u;\varepsilon,\sigma) = A_\varepsilon(u) \cap B_\sigma(u)$ with $A_\varepsilon(u) := \{v : E_{\mathrm{law}}(e^v;u) \le \varepsilon\}$ and $B_\sigma(u) := \{v : \psi(v;u) \ge \sigma\}$. If $\varepsilon_2 \ge \varepsilon_1$ and $\sigma_2 \le \sigma_1$, then $A_{\varepsilon_1}(u) \subseteq A_{\varepsilon_2}(u)$ and $B_{\sigma_1}(u) \subseteq B_{\sigma_2}(u)$ for every u, hence

$$\mathcal{A}[\mathcal{U};\varepsilon_1,\sigma_1] = \bigcap_u \big(A_{\varepsilon_1}(u) \cap B_{\sigma_1}(u)\big) \ \subseteq \ \bigcap_u \big(A_{\varepsilon_2}(u) \cap B_{\sigma_2}(u)\big) = \mathcal{A}[\mathcal{U};\varepsilon_2,\sigma_2].$$

Thus $\inf \mathcal{A}[\mathcal{U};\varepsilon_2,\sigma_2] \le \inf \mathcal{A}[\mathcal{U};\varepsilon_1,\sigma_1]$, i.e. relaxing tolerances cannot increase $v_R^{[\mathcal{U}]}$ (nor $\ell_R^{[\mathcal{U}]}$), and tightening them cannot decrease it—exactly as in Lemma 7.433(2). □

Intuition. A single restoration length must satisfy the constraints for *every* UV slice in $\mathcal{U}$, so it cannot be smaller than the worst per–slice requirement and increases when the strip widens; changing rulers only translates v and leaves the physical $\ell_R^{[\mathcal{U}]}$ unchanged, while looser thresholds enlarge the admissible intersection and can only move the first attainment to the left.

Existence and stability.

Corollary 7.436 (Existence and stability summary)**.** *Assume the hypotheses of Proposition 7.432 on an IR window $\mathcal{I}_{\mathrm{IR}}$ and the kernel conditions of Definition 7.380 and eq. (7.538) (with finite $m_2(W)$). Then the restoration length $\ell_R = e^{v_R}$ exists, is the unique first–attainment of the tolerance $\varepsilon_{\mathrm{plat}}$, and the map*

$$(\varepsilon_{\mathrm{plat}}, \varsigma_{\min}, \tau, \tau_E) \;\longmapsto\; \ell_R^{(\tau_E)}(\varepsilon_{\mathrm{plat}}, \varsigma_{\min}, \tau)$$

is continuous at $\tau_E = 0$, where $\ell_R^{(0)} = \ell_R$. Moreover, as $\tau_E \to 0$,

$$\boxed{\ell_R^{(\tau_E)} = \ell_R\Big(1 + \mathcal{O}\Big(\tfrac{\tau_E^2}{\varsigma_{\min}}\Big)\Big),} \tag{7.558}$$

with the implicit constant depending on $m_2(W)$ and $\sup_{\mathcal{I}_{\mathrm{IR}}} |\Lambda''_{\mathrm{law}}|$ (cf. Lemma 7.434).

Proof. Existence/first–attainment and tolerance continuity. By Proposition 7.432, the admissible set in eq. (7.551) is nonempty whenever $E_\infty \le \varepsilon_{\mathrm{plat}}$, its infimum is attained at v_R, and v_R is the unique first–attainment point. The same proposition yields (Kato–type) continuity of v_R (hence ℓ_R) with respect to $\varepsilon_{\mathrm{plat}}$ and $\varsigma_{\min}$ under the slope floor and curvature/TV guards (7.548).

Continuity in τ. The robust slope $\psi(v;\tau) = \varsigma_{\mathrm{law}}^{(\tau)}(v) = \frac{d}{dv}(\mathsf{LP}_\tau \Lambda_{\mathrm{law}})(v)$ depends continuously on τ in L^∞, since $\|W_\tau - W_{\tilde\tau}\|_{L^1} \to 0$ as $\tilde\tau \to \tau$ and $\|\mathsf{LP}_\tau f - \mathsf{LP}_{\tilde\tau} f\|_\infty \le \|W_\tau - W_{\tilde\tau}\|_1 \|f\|_\infty$ (Lemma 7.381(1),(4)). The guard quantities in (7.548) thus vary continuously with τ, and the first–attainment map inherits continuity by the same Kato–type argument used in Proposition 7.432.

Continuity in τ_E and bias rate. Let $D(v) = 1 - \Lambda_{\mathrm{law}}(v)$ and $D^{(\tau_E)}(v) = 1 - (\mathsf{LP}_{\tau_E}\Lambda_{\mathrm{law}})(v)$. By Lemma 7.434,

$$\|D^{(\tau_E)} - D\|_{L^\infty(\mathcal{I}_{\mathrm{IR}})} \;\le\; C_2\, \tau_E^2 + \mathcal{O}(\tau_E^4),$$

with $C_2 = \frac{m_2(W)}{2} \sup_{\mathcal{I}_{\mathrm{IR}}} |\Lambda''_{\mathrm{law}}|$. The first–attainment shift in v is bounded by the vertical bias divided by the slope floor:

$$|v_R^{(\tau_E)} - v_R| \;\le\; \frac{\|D^{(\tau_E)} - D\|_{L^\infty(\mathcal{I}_{\mathrm{IR}})}}{\varsigma_{\min}} \;=\; \mathcal{O}\Big(\frac{\tau_E^2}{\varsigma_{\min}}\Big),$$

using $\psi \ge \varsigma_{\min}$ on $\mathcal{I}_{\mathrm{IR}}$ (Definitions 7.385 and 7.392) and the mean–value estimate as in Proposition 7.432. Passing to $\ell = e^v$ gives (7.558) (cf. Lemma 7.433 for the $v \leftrightarrow \ell$ translation). This shows continuity at $\tau_E = 0$ and the stated bias rate.

All statements are compatible with gauge/presentation safety and pipeline moves by Proposition 7.424 and Remark 7.391. □

Discrete estimator (audit path).

Definition 7.437 (Discrete restoration length estimator)**.** Let $v_k = v_0 + k\,h$ (so $\ell_k = e^{v_k}$) be a uniform log–scale grid. Let $\mathsf{LP}_{\tau,h}$ and $\widehat{\varsigma}^{(\tau)}_{\mathrm{law}}$ be as in Definition 7.409 and eq. (7.547), with D_0 from Definition 7.406. Working on the eroded index set so that $\{k \pm 1\} \cup \{k-m, \dots, k+m\}$ exist, define

$$\boxed{k_R := \min\Big\{k \geq k_{\min} :\ E_{\mathrm{law}}(v_k) \leq \varepsilon_{\mathrm{plat}} \ \text{ and } \ \widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(v_j) \geq \varsigma_{\min} \ \ \forall\, j \geq k\Big\}, \qquad \widehat{\ell}_R := e^{v_{k_R}}.} \tag{7.559}$$

Definition 7.438 (Discrete curvature/TV guards (audit guard))**.** On an index interval $I = \{k_a, \dots, k_b\}$ (with $k_a \leq k_b$ in the eroded set), let

$$\psi_k := \widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(v_k) = (\mathsf{D}_0\,\mathsf{LP}_{\tau,h}\Lambda_{\mathrm{law}})_k, \quad (\mathsf{D}_2 f)_k := \frac{f_{k+1} - 2f_k + f_{k-1}}{h^2}.$$

Define the discrete guards (mirroring eq. (7.548)) by

$$\max_{k \in I^-} \big|(\mathsf{D}_2\,\mathsf{LP}_{\tau,h}\Lambda_{\mathrm{law}})_k\big| \ \leq\ \varepsilon_{\mathrm{curv}}, \qquad \mathrm{TV}_{h;I}[\psi] := \sum_{k \in I^-} \big|\psi_{k+1} - \psi_k\big| \ \leq\ \varepsilon_{\mathrm{tv}},$$

where $I^- := \{k_a + 1, \dots, k_b - 1\}$.

Remark 7.439 (Audit properties, guards, and references)**.** (i) *Stability and ripple suppression.* By Lemma 7.415, $\big|\widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(v_k)\big| \leq \|w\|_{\ell^1}\,\|\Lambda'_{\mathrm{law}}\|_\infty$, and the DTFT symbol $\widehat{\varsigma}(\xi) = \frac{i \sin \xi}{h}\widehat{W}(\xi)$ shows high–frequency attenuation for symmetric stencils ($w_{-j} = w_j$).

(ii) *Commutation and affine exactness.* D_0 commutes with $\mathsf{LP}_{\tau,h}$ and is exact on affine trends (Lemma 7.412 and Corollary 7.400); thus the guard is insensitive to linear drifts in v.

(iii) *Consistency with the continuum.* Under C^3 regularity and cell–average weights (Definition 7.398), $(\mathsf{D}_0\mathsf{LP}_{\tau,h}\Lambda_{\mathrm{law}})_k = \varsigma^{(\tau)}_{\mathrm{law}}(v_k) + \mathcal{O}(h^2)$ (Proposition 7.413); together with Proposition 7.432 this implies that, when the guards of Definition 7.438 hold on the forward tail $\{k \geq k_R\}$,

$$v_{k_R} = v_R + \mathcal{O}(h) \quad \text{and} \quad \widehat{\ell}_R = e^{v_{k_R}} = \ell_R\big(1 + \mathcal{O}(h)\big),$$

up to grid quantization in v (compare Corollary 7.436).

(iv) *Scaling and tolerance monotonicity.* The discrete estimator inherits the scaling covariance in v and physical invariance in ℓ, as well as tolerance monotonicity, from the continuum statements in Lemma 7.433 applied on the grid (set–inclusion arguments pass to the discrete admissible sets).

(v) *Boundary hygiene.* Using an eroded index set ensures the needed samples exist. If a full-support evaluation is preferred, admissible extensions (periodic/zero–padded/mirrored) preserve commutation of D_0 and $\mathsf{LP}_{\tau,h}$ on the extended grid (Proposition 7.404); state affine exactness only on the eroded set (Remark 7.405).

Intuition. $\widehat{\ell}_R$ is the grid–quantized *first–attainment* of the tolerance with a forward slope floor: once E_{law} is within $\varepsilon_{\mathrm{plat}}$ and $\psi_k \geq \varsigma_{\min}$ holds for all future indices, k_R is locked (guards in Definition 7.438 prevent false positives from local overshoots), consistent with the continuum ℓ_R and robust under smoothing and pipeline moves (Corollary 7.436 and Proposition 7.424).

Lemma 7.440 (Consistency and rate of the discrete estimator)**.** *Assume the hypotheses of Proposition 7.432 on an IR window $\mathcal{I}_{\mathrm{IR}}$ and that (i) $\Lambda_{\mathrm{law}} \in C^3$ on $\mathcal{I}_{\mathrm{IR}}$; (ii) the slope stencil D_0 and discrete low–pass $\mathsf{LP}_{\tau,h}$ are as in Definitions 7.406 and 7.409; (iii) the kernel satisfies eq. (7.538) with finite $m_2(W)$ and the cell–average weights Definition 7.398. Let $\widehat{\ell}_R$ be the discrete restoration length from Definition 7.437 and ℓ_R the continuum one from eq. (7.551). Then, as $h \to 0$ with τ, τ_E fixed,*

$$\boxed{\left|v_{k_R} - v_R\right| \;\le\; h \;+\; \frac{C}{\varsigma_{\min}}\Big(h^2 + \tau_E^2\Big), \qquad v_{k_R} = \ln \widehat{\ell}_R, \quad v_R = \ln \ell_R,} \tag{7.560}$$

where C depends on $m_2(W)$ and $\sup_{\mathcal{I}_{\mathrm{IR}}} |\Lambda''_{\mathrm{law}}|$, but not on h. Consequently,

$$\boxed{\left|\widehat{\ell}_R - \ell_R\right| \;\le\; \ell_R\Big(h \;+\; \frac{C}{\varsigma_{\min}}\Big(h^2 + \tau_E^2\Big)\Big)\,\Big(1 + \mathcal{O}(h)\Big),} \tag{7.561}$$

so $\widehat{\ell}_R \to \ell_R$ as $h \to 0$. If, instead of the grid index k_R, a de–quantized (linearly interpolated) crossing $\tilde{v}_R \in [v_{k_R-1}, v_{k_R}]$ is reported, then the leading h term in (7.560) is absent:

$$\left|\tilde{v}_R - v_R\right| \;\le\; \frac{C}{\varsigma_{\min}}\Big(h^2 + \tau_E^2\Big). \tag{7.562}$$

All bounds hold uniformly over admissible UV strips for the uniform estimator $\widehat{\ell}_R^{[\mathcal{U}]}$ (replace the per–slice sets by the intersections in Proposition 7.435).

Proof. Step 1 (discrete slope vs. robust slope). By Proposition 7.401 and Definition 7.398 and the second–order accuracy of D_0 (Definition 7.406), the composed estimator satisfies

$$\widehat{\psi}_k - \psi(v_k) = \Big(\mathsf{D}_0\mathsf{LP}_{\tau,h}\Lambda_{\mathrm{law}}\Big)_k - \frac{d}{dv}\Big(\mathsf{LP}_{\tau}\Lambda_{\mathrm{law}}\Big)(v_k) = \mathcal{O}(h^2)$$

uniformly on compact subsets of $\mathcal{I}_{\mathrm{IR}}$; cf. Proposition 7.413 and Lemma 7.412 and the smoothness C^3 (the third derivative enters the truncation error).

Step 2 (mollified pointwise deficit). Write the continuum deficit $D(v) := 1 - (\mathsf{LP}_{\tau}\Lambda_{\mathrm{law}})(v)$ (cf. Lemma 7.395 and Definition 7.385). If an auxiliary measurement mollifier LP_{τ_E} is used (cf. Lemma 7.434), define $D^{(\tau_E)}(v) := 1 - (\mathsf{LP}_{\tau_E}\Lambda_{\mathrm{law}})(v)$. Then Lemma 7.434 yields $\|D^{(\tau_E)} - D\|_{L^\infty(\mathcal{I}_{\mathrm{IR}})} \le C_2\,\tau_E^2 + \mathcal{O}(\tau_E^4)$ with $C_2 = \frac{m_2(W)}{2}\sup_{\mathcal{I}_{\mathrm{IR}}} |\Lambda''_{\mathrm{law}}|$.

Step 3 (vertical error ⇒ horizontal shift). On $\mathcal{I}_{\mathrm{IR}}$, the slope guard of Definition 7.392 gives $\psi \ge \varsigma_{\min} > 0$; hence $D'(v) = -\psi(v) \le -\varsigma_{\min}$ and D is strictly decreasing (Lemma 7.423(3)). Let v_R be the first attainment for D and $v_R^{(\tau_E)}$ the first attainment for $D^{(\tau_E)}$ (both exist and are unique by Proposition 7.432). Then the mean–value estimate (as in Proposition 7.432) gives

$$\left|v_R^{(\tau_E)} - v_R\right| \le \frac{\|D^{(\tau_E)} - D\|_{L^\infty(\mathcal{I}_{\mathrm{IR}})}}{\varsigma_{\min}} \le \frac{C_2}{\varsigma_{\min}}\tau_E^2 + \mathcal{O}\Big(\frac{\tau_E^4}{\varsigma_{\min}}\Big).$$

Step 4 (grid quantization and discrete feasibility). By definition Definition 7.437, k_R is the *first index* at which the discrete feasibility holds. Since D is decreasing with floor

$\varsigma_{\min}$, the continuous crossing v_R must lie in $[v_{k_R-1}, v_{k_R}]$, whence the grid quantization error $|v_{k_R} - v_R| \le h$. To control additional bias from the discrete guards, use Step 1:

$$\widehat{\psi}_k \ge \psi(v_k) - Ch^2 \quad \text{uniformly in } k.$$

For h small enough that $Ch^2 \le \frac{1}{2}\varsigma_{\min}$, the discrete slope guard $\widehat{\psi}_k \ge \varsigma_{\min}$ for all $k \ge k_R$ follows from the continuum guard $\psi \ge \varsigma_{\min}$. Combining with Step 3 (applied to $D^{(\tau_E)}$ if relevant) yields

$$\left|v_{k_R} - v_R\right| \;\le\; h \;+\; \frac{C}{\varsigma_{\min}}\left(h^2 + \tau_E^2\right),$$

for some constant C independent of h, which is (7.560). If a linearly de–quantized crossing $\tilde{v}_R \in [v_{k_R-1}, v_{k_R}]$ is reported (solve the linear model with slope $\widehat{\psi}_{k_R}$), the h term drops, giving (7.562).

Step 5 (from v to ℓ and UV–uniformity). Since $\ell = e^v$, the map $v \mapsto \ell$ is Lipschitz on bounded v-ranges with factor ℓ, giving (7.561). Uniformity over UV strips follows by taking sup/inf in u and using the uniform guard stability in Proposition 7.424 together with the set–inclusion arguments of Proposition 7.435. □

Remark 7.441 (On the roles of h, τ, and τ_E)**.** The bound (7.560) separates: (i) a *quantization* term h intrinsic to grid first–attainment (removed by de–quantization); (ii) a *discrete/measurement bias* $\mathcal{O}(h^2/\varsigma_{\min})$ from D_0 and $\mathsf{LP}_{\tau,h}$ (Proposition 7.401 and Definition 7.406); (iii) an *auxiliary smoothing* bias $\mathcal{O}(\tau_E^2/\varsigma_{\min})$ controlled by Lemma 7.434. The parameter τ defines the *continuum* robust slope and is part of ℓ_R's definition (no extra τ^2 appears here). Scaling/tolerance monotonicity for the discrete estimator follows from Lemma 7.433.

Proposition 7.442 (No false positives under discrete guards)**.** *Let the discrete curvature/TV guards of Definition 7.438 hold on the forward tail $I = \{k_{\min} + m + 1, \dots\}$ of the audit grid $v_k = v_0 + k\,h$ (so all stencil samples exist). If k_R is defined by Definition 7.437 and eq. (7.559), then for all $j \ge k_R$,*

$$E_{\mathrm{law}}(v_j) \le \varepsilon_{\mathrm{plat}} \quad \textit{and} \quad \widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(v_j) =: (\mathsf{D}_0\mathsf{LP}_{\tau,h}\Lambda_{\mathrm{law}})_j \;\ge\; \varsigma_{\min}.$$

Hence transient local overshoots of either E_{law} or the discrete slope cannot produce a spurious (premature) index k_R.

Proof. Slope constraint on the forward tail. By Definition 7.437, the index k_R is the *first* grid point such that $\widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(v_j) \ge \varsigma_{\min}$ for *all* $j \ge k_R$. Thus the slope inequality for $j \ge k_R$ holds by definition (no guard needed). The TV part of Definition 7.438 provides an audit check:

$$\mathrm{TV}_{h;[k_R,\infty)}[\psi] = \sum_{j \ge k_R} \left|\psi_{j+1} - \psi_j\right| \le \varepsilon_{\mathrm{tv}} \quad \text{with} \quad \psi_j := \widehat{\varsigma}^{(\tau)}_{\mathrm{law}}(v_j),$$

ensuring bounded variability of the readout, consistent with Lemma 7.423 at the discrete level.

Persistence of the deficit/tolerance inequality. Introduce the discrete law–deficit

$$D_k := 1 - (\mathsf{LP}_{\tau,h}\Lambda_{\mathrm{law}})_k \qquad \text{(cf. the continuum } D(v) = 1 - (\mathsf{LP}_\tau\Lambda_{\mathrm{law}})(v) \text{ in Lemma 7.395).}$$

By the three–point identity $f_{k+1}-f_k = h\,(\mathsf{D}_0 f)_k + \frac{h^2}{2}\,(\mathsf{D}_2 f)_k$ (valid for any grid function f, with D_0 as in Definition 7.406 and $\mathsf{D}_2 f_k := (f_{k+1} - 2f_k + f_{k-1})/h^2$), applied to $f_k := (\mathsf{LP}_{\tau,h}\Lambda_{\mathrm{law}})_k$, one obtains

$$D_{k+1}-D_k = -\big(f_{k+1}-f_k\big) = -\,h\,(\mathsf{D}_0 f)_k - \frac{h^2}{2}\,(\mathsf{D}_2 f)_k = -\,h\,\psi_k - \frac{h^2}{2}\,(\mathsf{D}_2\mathsf{LP}_{\tau,h}\Lambda_{\mathrm{law}})_k. \tag{7.563}$$

The discrete curvature guard of Definition 7.438 states that $\max_{k\in I^-}\big|(\mathsf{D}_2\mathsf{LP}_{\tau,h}\Lambda_{\mathrm{law}})_k\big| \le \varepsilon_{\mathrm{curv}}$. Combining this with (7.563) and the forward–tail slope floor $\psi_k \ge \varsigma_{\min}$ for all $k \ge k_R$ (from Definition 7.437) gives, for $k \ge k_R$,

$$D_{k+1} - D_k \;\le\; -\,h\,\varsigma_{\min} \;+\; \frac{h^2}{2}\,\varepsilon_{\mathrm{curv}}. \tag{7.564}$$

Hence, provided the grid step obeys the mild CFL–type condition

$$h \;\le\; \frac{2\,\varsigma_{\min}}{\varepsilon_{\mathrm{curv}}}, \tag{7.565}$$

the right–hand side of (7.564) is nonpositive and D_k is nonincreasing for all $k \ge k_R$. At $k = k_R$ the acceptance criterion requires $E_{\mathrm{law}}(v_{k_R}) \le \varepsilon_{\mathrm{plat}}$ (Definition 7.437). Two cases then yield persistence:

• *Canonical (mollified) deficit:* $E_{\mathrm{law}} = D$. If the audit uses $E_{\mathrm{law}}(v_k) = D_k$ (the discrete analogue of the continuum deficit; cf. Lemma 7.395), then

$$D_j \;\le\; D_{k_R} \;\le\; \varepsilon_{\mathrm{plat}} \qquad \forall\, j \ge k_R,$$

by the monotonicity deduced from (7.564)–(7.565).

• *General E_{law} controlled by guards.* If E_{law} is a pointwise (or separately mollified) visibility, assume the standard audit control $|E_{\mathrm{law}}(v_k) - D_k| \le \delta_{\mathrm{aux}}$ with $\delta_{\mathrm{aux}} = \mathcal{O}(\tau_E^2)$ from Lemma 7.434 (or $\mathcal{O}(h^2)$ from Proposition 7.413 when comparing discrete–continuous), uniformly for $k \ge k_R$. Then

$$E_{\mathrm{law}}(v_j) \;\le\; D_j + \delta_{\mathrm{aux}} \;\le\; D_{k_R} + \delta_{\mathrm{aux}} \;\le\; \varepsilon_{\mathrm{plat}} + \delta_{\mathrm{aux}}, \qquad j \ge k_R.$$

If the acceptance threshold incorporates the bias margin (the usual practice in Step 2 audits; cf. Corollary 7.436), the inequality $E_{\mathrm{law}}(v_j) \le \varepsilon_{\mathrm{plat}}$ persists for all $j \ge k_R$. (Equivalently, choose $\varepsilon_{\mathrm{plat}}^{\mathrm{eff}} := \varepsilon_{\mathrm{plat}} - \delta_{\mathrm{aux}}$ in the decision rule.)

Minimality ("first attainment"). By Definition 7.437, k_R is the smallest index that satisfies both inequalities, with the slope floor required on the entire forward tail. Thus no smaller index can be admissible (some constraint fails), excluding premature triggers. □

Remark 7.443 (Relationship to continuum monotonicity)**.** The discrete inequality (7.564) is the grid counterpart of the continuum identity $D'(v) = -\psi(v)$ (cf. Definition 7.385 and Lemma 7.395), with the extra $\frac{h^2}{2}\,\mathsf{D}_2$ term controlled by the discrete curvature guard (Definition 7.438). Under (7.565), the sign of the forward difference matches the continuum sign, preventing false positives from local overshoots. Guard stability across gauges/presentations and pipeline moves follows from Proposition 7.424.

State–coherence guard (uniform in volume).

Definition 7.444 (Uniform state coherence)**.** All restoration statements are conditioned on *state–channel* coherence remaining within the plateau tolerance, *uniformly in the volume* V (cf. the initialization deficits in eq. (7.535) and the plateau criterion in Section 6.12.4):

$$\boxed{\sup_{V} \left| 1 - \Lambda_{\mathrm{state}}(\lambda_{\mathrm{UV}}, \ell; T^{\star}, V) \right| \;\le\; \varepsilon_{\mathrm{plat}} \qquad \text{for all} \;\; \ell \ge \ell_R.} \tag{7.566}$$

Here ℓ_R is the (law–channel) restoration length defined by eq. (7.551) (see also Proposition 7.432).

Intuition. The guard forbids QII–type volume–induced decoherence in the state channel; any certified restoration is genuinely QIII (law–channel phenomenon), not an artefact of drifting states.

Proposition 7.445 (Separation of QIII vs. QII under the state guard)**.** *Assume the law–channel restoration condition of eq. (7.551) (or its UV–uniform variant eq. (7.553)) holds on an IR window* $\mathcal{I}_{\mathrm{IR}}$ *and the state–coherence guard (7.566) is satisfied. Then the system is in QIII (law–asymmetric, state–symmetric) on* $\mathcal{I}_{\mathrm{IR}}$*. Conversely, if there exist* $V_n \to \infty$ *and* $\ell_n \ge \ell_R$ *such that* $\left|1 - \Lambda_{\mathrm{state}}(\lambda_{\mathrm{UV}}, \ell_n; T^{\star}, V_n)\right| > \varepsilon_{\mathrm{plat}}$*, then QIII cannot be certified at those scales.*

Proof. Step 1 (law restoration on $\mathcal{I}_{\mathrm{IR}}$*).* By eq. (7.551) and Proposition 7.432, there exists $v_R = \ln \ell_R$ such that for all $v \in \mathcal{I}_{\mathrm{IR}} \cap [v_R, \infty)$ the law–channel deficit $D(v) := 1 - (\mathsf{LP}_\tau \Lambda_{\mathrm{law}})(v)$ obeys $D(v) \le \varepsilon_{\mathrm{plat}}$, and the robust slope $\psi(v) := \varsigma^{(\tau)}_{\mathrm{law}}(v)$ satisfies $\psi(v) \ge \varsigma_{\min} > 0$ (cf. Definitions 7.385 and 7.392 and Lemma 7.423).

Step 2 (state symmetry uniformly in V*).* The guard (7.566) gives $\sup_V \left| 1 - \Lambda_{\mathrm{state}}(\lambda_{\mathrm{UV}}, \ell; T^{\star}, V) \right| \le \varepsilon_{\mathrm{plat}}$ for all $\ell \ge \ell_R$, hence state–channel symmetry within the same tolerance on the whole IR window. This uniformity is stable under admissible limits and smoothing operations used in the pipeline (see Section 6.12.4 and Proposition 7.383).

Step 3 (quadrant identification and safety). By construction (cf. Section 7.8 vs. Section 7.7), QIII is the regime with restored law–channel (Step 1) and state symmetry (Step 2). Gauge/p-resentation safety of these statements follows from Proposition 7.424 and identity–ledger compatibility Remark 7.426.

In **Step 4, we argue by contrapositive:** If there exist $V_n \to \infty$ and $\ell_n \ge \ell_R$ with $\left|1 - \Lambda_{\mathrm{state}}(\lambda_{\mathrm{UV}}, \ll_n; T^{\star}, V_n)\right| > \varepsilon_{\mathrm{plat}}$, then (7.566) fails on those scales, violating the state–symmetry requirement; QIII cannot be certified there (the behaviour is consistent with QII–type decoherence). □

7.8.4 RG Ledger: Breakings Become Irrelevant

Objective. We show that the explicit breaking operators introduced in eq. (7.522) flow to irrelevance toward the IR. Concretely, we identify an IR fixed point for the symmetry-preserving couplings, linearise the RG flow of the breaking couplings, diagonalise the stability matrix to obtain eigen–operators with exponents $y_{\tilde{B}} < 0$, and verify the measured running

$\varepsilon_{\tilde{B}}(\mu) \sim \mu^{y_{\tilde{B}}} \to 0$ as $\mu \downarrow$ (Wilsonian sense [389]; see also [91, 82, 447]). All quoted exponents are reported *after Nielsen projection* to ensure gauge–parameter independence within tolerance [117, 243].

Setup and linearised flow. Let $g^a(\mu)$ denote symmetry–preserving couplings and $\varepsilon_A(\mu)$ the small breaking couplings multiplying the renormalised operators $\{\hat{\mathcal{O}}_A\}$. Near an IR fixed point $g^a(\mu) \to g^a_\star$ we write the linearised RG for the breakings

$$\boxed{\mu \frac{\mathrm{d}\,\varepsilon_A}{\mathrm{d}\mu} = \sum_B Y_{AB}\,\varepsilon_B \;+\; \mathcal{O}(\varepsilon^2), \qquad Y_{AB} := \left.\frac{\partial\,\beta_{\varepsilon_A}}{\partial \varepsilon_B}\right|_{g=g_\star,\ \varepsilon=0}.} \tag{7.567}$$

Equivalently, if the renormalised operators mix as

$$\mu \frac{\mathrm{d}}{\mathrm{d}\mu}\,\hat{\mathcal{O}}_A \;=\; -\,\gamma_{AB}\,\hat{\mathcal{O}}_B, \qquad \gamma \;=\; Z^{-1}\,\mu \frac{\mathrm{d}Z}{\mathrm{d}\mu},$$

and the deformation is written with explicit engineering dimensions as

$$\delta S \;=\; \int \mathrm{d}^D x \,\sum_A \varepsilon_A(\mu)\,\mu^{\,D-d_A}\,\hat{\mathcal{O}}_A,$$

then the linearised matrix is

$$\boxed{Y_{AB} \;=\; \gamma_{BA} \;-\; (D-d_A)\,\delta_{AB}}.$$

In particular, if the $\mu^{\,D-d_A}$ factor has already been absorbed into the definition of the (dimensionless) couplings, the canonical piece drops and $Y = \gamma^{\top}$.

Definition 7.446 (Centered log–derivative estimator in RG units)**.** Let $t := \ln \mu$ and suppose $\{\mu_k\}$ is a grid monotone in the IR direction.

$$\boxed{\widehat{y}_{\tilde{B}}(\mu_k) := \frac{1}{2\,\Delta \ln \mu}\,\ln\!\left(\frac{|\widetilde{\varepsilon}_{\tilde{B}}(\mu_{k+1})|}{|\widetilde{\varepsilon}_{\tilde{B}}(\mu_{k-1})|}\right), \qquad \Delta \ln \mu := \ln(\mu_{k+1}) - \ln(\mu_{k-1}),} \tag{7.568}$$

Definition 7.447 (IR fixed point and breaking sector)**.** An *IR fixed point* for the symmetry–preserving sector is a point $g_\star$ with $\beta^a(g_\star) = 0$ such that the linearised stability matrix

$$S^a{}_b \;:=\; \left.\frac{\partial \beta^a}{\partial g^b}\right|_{g_\star}$$

has eigenvalues with nonnegative real parts (IR attraction in the preserving subspace). The *breaking sector* is the vector space spanned by the renormalised operators $\{\hat{\mathcal{O}}_A\}$ with coordinates ε_A, equipped with the linearised flow eq. (7.567).

Intuition. We freeze the symmetric couplings at their IR attractor $g_\star$ and study how tiny explicit breakings ε_A evolve under RG.

Definition 7.448 (Nielsen projection and projected exponents)**.** Let α denote gauge parameters and evaluate correlators *on shell* so that the Nielsen identity eq. (7.581) applies. The *Nielsen projector* $\mathcal{N}(\alpha)$ is a linear, idempotent map on the breaking–operator space V such that

$$\mathcal{N}^2 = \mathcal{N}, \qquad \operatorname{im}\mathcal{N} \cong H^0_{\mathrm{BRST}} \text{ (physical cohomology)}, \qquad \partial_\alpha\big(\mathcal{N}X\big) = 0 \ \text{ for all } X \in V$$

(within admissible tolerance, cf. [117, 243]). For any mixing/stability matrix Z, Y on V (cf. eq. (7.567)), define their *Nielsen–projected* versions by restriction to the physical subspace $W := \operatorname{im}\mathcal{N}$:

$$Z^{\mathcal{N}} := \mathcal{N}\, Z\, \mathcal{N}\big|_W, \qquad Y^{\mathcal{N}} := \mathcal{N}\, Y\, \mathcal{N}\big|_W.$$

The eigenvalues of $Y^{\mathcal{N}}$ are the *Nielsen–projected breaking exponents* $\{y_{\tilde B}\}$, and any associated (right) eigenvectors $\{\tilde v^{(\tilde B)}\} \subset W$ define the *eigen–operators*

$$\widetilde{\mathcal{O}}_{\tilde B} \;:=\; \tilde v^{(\tilde B)}_{A}\, \hat{\mathcal{O}}_A.$$

(Equivalently, choosing a dual left eigenbasis $\{\tilde w^{(\tilde B)}\}$ with $(\tilde w^{(\tilde B)})^{\top}\tilde v^{(\tilde B')} = \delta_{\tilde B\tilde B'}$ sets the associated eigen–couplings $\widetilde{\varepsilon}_{\tilde B} := (\tilde w^{(\tilde B)})^{\top}\varepsilon$, cf. Definition 7.454.)

Intuition. Project onto the on–shell BRST–physical subspace first; read exponents from $Y^{\mathcal{N}}$ there. This removes gauge artefacts (isospectral in α by Proposition 7.458, with the tolerance check in eq. (7.588)).

Definition 7.449 (Jordan nilpotent part, index, and chains)**.** Work on the physical subspace $W := \operatorname{im}\mathcal{N}$ with the Nielsen–projected operator $Y^{\mathcal{N}}$ (cf. Definition 7.448). For each eigenvalue $y_{\tilde B}$ of $Y^{\mathcal{N}}$, let the generalised eigenspace be

$$\mathcal{G}_{\tilde B} \;:=\; \ker\!\Big((Y^{\mathcal{N}} - y_{\tilde B}\mathbb{1}_{\mathcal{K}})^{p}\Big) \quad \text{for sufficiently large } p.$$

Choose a basis of $\mathcal{G}_{\tilde B}$ that brings $Y^{\mathcal{N}}$ to Jordan form

$$Y^{\mathcal{N}}\big|_{\mathcal{G}_{\tilde B}} \;=\; J_{\tilde B} \;:=\; \bigoplus_{j=1}^{n_{\tilde B}} \Big(y_{\tilde B}\mathbb{1}_{m_{\tilde B,j}} + \mathsf{N}_{\tilde B,j}\Big),$$

where each $\mathsf{N}_{\tilde B,j}$ is strictly upper–triangular with $(\mathsf{N}_{\tilde B,j})^{m_{\tilde B,j}} = 0$ and $(\mathsf{N}_{\tilde B,j})^{m_{\tilde B,j}-1} \neq 0$. The nilpotent part on $\mathcal{G}_{\tilde B}$ is

$$\mathsf{N}_{\tilde B} \;:=\; \bigoplus_{j=1}^{n_{\tilde B}} \mathsf{N}_{\tilde B,j}, \qquad \operatorname{ind}\mathsf{N}_{\tilde B} \;:=\; r_{\tilde B} \;:=\; \max_{1\le j\le n_{\tilde B}} m_{\tilde B,j}.$$

For each block j, a (right) Jordan chain of length $m_{\tilde B,j}$, $\{\tilde v^{(\tilde B,j)}_{[k]}\}_{k=0}^{m_{\tilde B,j}-1} \subset \mathcal{G}_{\tilde B}$, satisfies

$$(Y^{\mathcal{N}} - y_{\tilde B}\mathbb{1})\,\tilde v^{(\tilde B,j)}_{[0]} = 0, \qquad (Y^{\mathcal{N}} - y_{\tilde B}\mathbb{1})\,\tilde v^{(\tilde B,j)}_{[k]} = \tilde v^{(\tilde B,j)}_{[k-1]} \quad (k \ge 1).$$

(Optionally pick a biorthogonal left chain $\{\tilde w^{(\tilde B,j)}_{[k]}\}_{k=0}^{m_{\tilde B,j}-1} \subset \mathcal{G}^{*}_{\tilde B}$ with $(\tilde w^{(\tilde B,j)}_{[k]})^{\top}\tilde v^{(\tilde B',j')}_{[\ell]} = \delta_{\tilde B\tilde B'}\delta_{jj'}\delta_{k\ell}$.)

Notation. $\mathcal{N}$ denotes the Nielsen projector (onto W), whereas N denotes the nilpotent Jordan part(s) inside blocks of $Y^{\mathcal{N}}$.
Intuition. Each physical eigenspace $\mathcal{G}_{\tilde{B}}$ decomposes into Jordan blocks; the nilpotent index $r_{\tilde{B}}$ is the length of the longest chain for $y_{\tilde{B}}$. When $r_{\tilde{B}} = 1$ the sector is diagonalizable; if $r_{\tilde{B}} > 1$, solutions of $\mu \frac{\mathrm{d}}{\mathrm{d}\mu} \widetilde{\varepsilon} = Y^{\mathcal{N}} \widetilde{\varepsilon}$ acquire factors $\mu^{y_{\tilde{B}}} (\log \mu)^q$ with $0 \le q < r_{\tilde{B}}$, while (ir)relevance is still governed by $\operatorname{Re} y_{\tilde{B}} < 0$ on W.

Proposition 7.450 (Solution of the linearised flow and irrelevance criterion)**.** *Work on the physical subspace $W := \operatorname{im} \mathcal{N}$ and write $\varepsilon(\mu) \in W$ for the Nielsen–projected breaking–coupling vector. Assume $Y^{\mathcal{N}}$ admits a Jordan decomposition on W. For each eigenvalue $y_{\tilde{B}}$, let its Jordan blocks be indexed by $j = 1, \dots, n_{\tilde{B}}$ with sizes $m_{\tilde{B},j}$, and choose right/left Jordan chains $\{\tilde{v}^{(\tilde{B},j)}_{[k]}\}_{k=0}^{m_{\tilde{B},j}-1}$, $\{\tilde{w}^{(\tilde{B},j)}_{[k]}\}_{k=0}^{m_{\tilde{B},j}-1}$ satisfying $(\tilde{w}^{(\tilde{B},j)}_{[k]})^{\top} \tilde{v}^{(\tilde{B}',j')}_{[\ell]} = \delta_{\tilde{B}\tilde{B}'} \delta_{jj'} \delta_{k\ell}$. Then the solution of the linearised system*

$$\mu \frac{\mathrm{d}\varepsilon}{\mathrm{d}\mu} = Y^{\mathcal{N}} \varepsilon, \qquad \varepsilon(\mu_0) \text{ given},$$

is

$$\boxed{\varepsilon(\mu) = \sum_{\tilde{B}} \sum_{j=1}^{n_{\tilde{B}}} \left(\frac{\mu}{\mu_0}\right)^{y_{\tilde{B}}} \sum_{k=0}^{m_{\tilde{B},j}-1} \frac{\left(\ln \frac{\mu}{\mu_0}\right)^k}{k!} c_{\tilde{B},j,k}\, \tilde{v}^{(\tilde{B},j)}_{[k]}, \qquad c_{\tilde{B},j,k} := \left(\tilde{w}^{(\tilde{B},j)}_{[k]}\right)^{\top} \varepsilon(\mu_0).} \tag{7.569}$$

In particular, if $\operatorname{Re} y_{\tilde{B}} < 0$ for every projected breaking mode (all $\tilde{B}$), then each component of $\varepsilon(\mu)$ decays as $\mu \downarrow 0$ (the polynomial logarithmic factors do not change the limit), and the breakings are irrelevant *in the Wilsonian sense.*

Intuition. Solve the linear ODE in Jordan form: each eigenvalue contributes a power $(\mu/\mu_0)^{y_{\tilde{B}}}$ and each Jordan block adds at most a polynomial in $\log(\mu/\mu_0)$. Strictly negative $\operatorname{Re} y_{\tilde{B}}$ wins against any such polynomial, so all projected breaking components vanish toward the IR.

Proof. By the projected linearised RG eq. (7.567) and the Nielsen projection Definition 7.448, the breaking vector on $W := \operatorname{im} \mathcal{N}$ obeys

$$\mu \frac{\mathrm{d}\varepsilon}{\mathrm{d}\mu} = Y^{\mathcal{N}} \varepsilon \; + \; r(\varepsilon, \mu), \qquad r(\varepsilon, \mu) = \mathcal{O}(\|\varepsilon\|^2).$$

Set $t := \ln(\mu/\mu_0)$ so that $\frac{\mathrm{d}}{\mathrm{d}t} = \mu \frac{\mathrm{d}}{\mathrm{d}\mu}$, giving

$$\frac{\mathrm{d}\varepsilon}{\mathrm{d}t} = Y^{\mathcal{N}} \varepsilon + r(\varepsilon, t).$$

Choose a similarity R that brings $Y^{\mathcal{N}}$ to Jordan form on W (cf. Definition 7.449):

$$R^{-1} Y^{\mathcal{N}} R = \bigoplus_{\tilde{B}} \bigoplus_{j=1}^{n_{\tilde{B}}} \left(y_{\tilde{B}} \mathbb{1}_{m_{\tilde{B},j}} + \mathsf{N}_{\tilde{B},j} \right), \qquad (\mathsf{N}_{\tilde{B},j})^{m_{\tilde{B},j}} = 0.$$

Write $\widetilde{\varepsilon} := R^{-1} \varepsilon$. Variation of constants yields

$$\widetilde{\varepsilon}(t) = \mathrm{e}^{(t-t_0)J}\, \widetilde{\varepsilon}(t_0) + \int_{t_0}^{t} \mathrm{e}^{(t-s)J}\, R^{-1} r\Big(R \widetilde{\varepsilon}(s), s\Big)\, \mathrm{d}s, \quad J = \bigoplus_{\tilde{B}} \bigoplus_{j=1}^{n_{\tilde{B}}} \left(y_{\tilde{B}} \mathbb{1}_{m_{\tilde{B},j}} + \mathsf{N}_{\tilde{B},j} \right).$$

For each block, $\mathrm{e}^{(t-t_0)J_{\tilde B,j}} = \mathrm{e}^{y_{\tilde B}(t-t_0)}\sum_{k=0}^{m_{\tilde B,j}-1}\frac{(t-t_0)^k}{k!}(\mathsf{N}_{\tilde B,j})^k$. Expanding the initial data in a (right/left) Jordan basis $\{\tilde v_{[k]}^{(\tilde B,j)}\}$, $\{\tilde w_{[k]}^{(\tilde B,j)}\}$ with $(\tilde w_{[k]}^{(\tilde B,j)})^\top \tilde v_{[\ell]}^{(\tilde B',j')} = \delta_{\tilde B\tilde B'}\delta_{jj'}\delta_{k\ell}$ (see Definition 7.449), we obtain for the *linear* evolution

$$\varepsilon_{\mathrm{lin}}(t) = \sum_{\tilde B}\sum_{j=1}^{n_{\tilde B}} \mathrm{e}^{y_{\tilde B}(t-t_0)} \sum_{k=0}^{m_{\tilde B,j}-1} \frac{(t-t_0)^k}{k!}\, c_{\tilde B,j,k}\, \tilde v_{[k]}^{(\tilde B,j)}, \qquad c_{\tilde B,j,k} := \big(\tilde w_{[k]}^{(\tilde B,j)}\big)^\top \varepsilon(\mu_0).$$

Returning to μ via $t - t_0 = \ln(\mu/\mu_0)$ reproduces the boxed expression eq. (7.569). For the remainder, note that $R^{\pm 1}$ are bounded on the finite-dimensional breaking space and $r(\varepsilon,t) = \mathcal{O}(\|\varepsilon\|^2)$; hence the integral term is $\mathcal{O}(\|\varepsilon\|^2)$ uniformly on compact t-intervals.

If $\mathrm{Re}\, y_{\tilde B} < 0$ for all projected modes, then for each fixed k,

$$\Big(\tfrac{\mu}{\mu_0}\Big)^{y_{\tilde B}}\Big(\ln\tfrac{\mu}{\mu_0}\Big)^k \;\longrightarrow\; 0 \quad \text{as } \mu \downarrow 0,$$

so every linear Jordan component vanishes. For sufficiently small $\|\varepsilon(\mu_0)\|$, a Grönwall estimate applied to $\frac{\mathrm{d}}{\mathrm{d}t}\|\varepsilon\| \le \|Y^{\mathcal{N}}\|\,\|\varepsilon\| + C\|\varepsilon\|^2$ shows that the quadratic remainder cannot overcome the linear decay, and thus $\varepsilon(\mu) \to 0$ componentwise. This proves the irrelevance claim in the Wilsonian sense. □

Intuition. After projecting with Definition 7.448, the flow on W is governed by $Y^{\mathcal{N}}$. Its Jordan structure from Definition 7.449 adds only finite polynomials in $\ln(\mu)$ atop the power $(\mu/\mu_0)^{y_{\tilde B}}$; strict negativity of $\mathrm{Re}\, y_{\tilde B}$ dominates these, and the $\mathcal{O}(\|\varepsilon\|^2)$ term remains subleading for small breakings, yielding eq. (7.569) and decay to zero.

Lemma 7.451 (Basis and scheme covariance)**.** *Let V be the breaking–operator space and $W := \mathrm{im}\,\mathcal{N} \subset V$ the on–shell BRST–physical subspace (cf. Definition 7.448). Under a nonsingular linear mixing $\hat{\mathcal{O}}_A \mapsto M_A{}^B\hat{\mathcal{O}}_B$ that preserves W and the corresponding coupling redefinition that leaves the breaking pairing invariant, the projected stability matrix transforms by similarity on W:*

$$Y^{\mathcal{N}} \;\longmapsto\; Y^{\mathcal{N}\prime} = M_W\, Y^{\mathcal{N}}\, M_W^{-1}, \qquad M_W := \mathcal{N}\, M\, \mathcal{N}\big|_W.$$

Hence the set of projected breaking exponents $\{y_{\tilde B}\}$ (cf. Definition 7.448) is invariant. Moreover, any smooth change of renormalisation scheme implemented by a finite (invertible) renormalisation $F(g,\alpha)$ that preserves W leaves $\{y_{\tilde B}\}$ unchanged at the IR fixed point $g_\star$ (cf. Definition 7.447).

Proof. **(A) Pure basis change in the breaking sector.** Let V^* denote the dual (couplings) and write the breaking deformation as the pairing $\langle \varepsilon, \mathcal{O}\rangle = \sum_A \varepsilon_A \hat{\mathcal{O}}_A$. A basis change $\mathcal{O} \mapsto \mathcal{O}' := M\mathcal{O}$ with $M \in \mathrm{GL}(V)$ induces the dual change $\varepsilon \mapsto \varepsilon' := M^{-T}\varepsilon$ so that $\langle \varepsilon', \mathcal{O}'\rangle = \langle \varepsilon, \mathcal{O}\rangle$. If the linearised RG for couplings reads (cf. eq. (7.567))

$$\mu\,\frac{\mathrm{d}\varepsilon}{\mathrm{d}\mu} = Y^*\,\varepsilon + \mathcal{O}(\varepsilon^2),$$

then

$$\mu\,\frac{\mathrm{d}\varepsilon'}{\mathrm{d}\mu} = M^{-T} Y^* M^T\,\varepsilon' + \mathcal{O}(\varepsilon'^2).$$

Passing to operators ($Y := (Y^*)^T$) gives $Y \mapsto MYM^{-1}$.

Restrict now to $W = \operatorname{im}\mathcal{N}$ (cf. Definition 7.448). By hypothesis $MW \subseteq W$, so $M_W := \mathcal{N}M\mathcal{N}\big|_W : W \to W$ is invertible.[11] The projected matrix $Y^{\mathcal{N}} := \mathcal{N}Y\mathcal{N}\big|_W$ therefore transforms by similarity:

$$Y^{\mathcal{N}} \;\longmapsto\; Y^{\mathcal{N}\prime} = \mathcal{N}(MYM^{-1})\mathcal{N}\big|_W = (\mathcal{N}M\mathcal{N})\,Y^{\mathcal{N}}\,(\mathcal{N}M^{-1}\mathcal{N}) = M_W\,Y^{\mathcal{N}}\,M_W^{-1}.$$

Similarity preserves spectra, hence $\{y_{\tilde{B}}\}$ is invariant under admissible basis changes.

(B) Finite renormalisation (scheme) change at the fixed point. Let $Z(g,\alpha,\mu)$ be the renormalisation matrix on V and define the anomalous–dimension matrix $\gamma := Z^{-1}\mu\,\frac{\mathrm{d}Z}{\mathrm{d}\mu}$; with canonical dimensions separated (see the discussion below eq. (7.567)), write $Y := d - \gamma$ in an operator basis. A smooth scheme change is implemented by a finite, invertible $F(g,\alpha)$:

$$Z \;\longmapsto\; Z' := F\,Z, \qquad \gamma' \;=\; Z'^{-1}\mu\frac{\mathrm{d}Z'}{\mathrm{d}\mu} \;=\; F^{-1}\gamma F \;+\; F^{-1}\left(\beta^a\frac{\partial F}{\partial g^a} + \frac{\partial F}{\partial\alpha}\,\mu\frac{\mathrm{d}\alpha}{\mathrm{d}\mu}\right).$$

Project onto W after going on shell. By the Nielsen identity (eq. (7.581)) and Definition 7.448, the explicit α–term vanishes in cohomology; if F preserves W let $F_W := \mathcal{N}F\mathcal{N}\big|_W$. At an IR fixed point $g_\star$ with $\beta(g_\star) = 0$ (cf. Definition 7.447) we have

$$\gamma_\star^{\mathcal{N}} \;\longmapsto\; \gamma_\star^{\mathcal{N}\prime} = F_W^{-1}\,\gamma_\star^{\mathcal{N}}\,F_W.$$

Assuming F_W mixes only operators within equal canonical-dimension blocks (so $[F_W, d] = 0$), it follows that

$$Y_\star^{\mathcal{N}} = d - \gamma_\star^{\mathcal{N}} \;\longmapsto\; Y_\star^{\mathcal{N}\prime} = F_W^{-1}(d - \gamma_\star^{\mathcal{N}})F_W = F_W^{-1}\,Y_\star^{\mathcal{N}}\,F_W,$$

i.e. a similarity on W. Thus the eigenvalues $\{y_{\tilde{B}}\}$ are scheme independent at the fixed point.

(C) Summary. Admissible basis changes act by similarity on W, and finite scheme changes reduce to similarity at the fixed point; in both cases the spectrum of $Y^{\mathcal{N}}$ (hence the projected exponents from Definition 7.448) is invariant. □

Intuition. Relabelling operators or changing scheme only *conjugates* the projected flow generator $Y^{\mathcal{N}}$ on W, so the physical exponents $y_{\tilde{B}}$ are untouched.

Definition 7.452 (Measured running and local exponent estimator)**.** Let $W := \operatorname{im}\mathcal{N}$ be the on–shell BRST–physical subspace (cf. Definition 7.448). For a projected eigen–operator $\hat{\tilde{\mathcal{O}}}_{\tilde{B}}$ with breaking exponent $y_{\tilde{B}}$ (from the spectrum of $Y^{\mathcal{N}}$; see Definition 7.448), define the *eigen–coupling component*

$$\widetilde{\varepsilon}_{\tilde{B},[0]}(\mu) \;:=\; \left(\tilde{w}_{[0]}^{(\tilde{B})}\right)^{\top}\varepsilon(\mu),$$

[11] If $M_W w = 0$ with $w \in W$, then $\mathcal{N}Mw = 0$. But $Mw \in W$ (preservation), hence $Mw \in W \cap \ker\mathcal{N} = \{0\}$, so $w = 0$ since M is invertible.

using the top element of a left Jordan chain on W (cf. Definition 7.449); in the diagonalizable case this is the usual left eigenvector projection. The *measured running* and its local estimator are then

$$\widetilde{\varepsilon}_{\tilde{B},[0]}(\mu) \;\sim\; \mu^{y_{\tilde{B}}} \times \Big[\text{polylog corrections from Jordan blocks}\Big], \quad \text{cf. eq. (7.569)}, \tag{7.570}$$

$$\widehat{y}_{\tilde{B}}(\mu) := \frac{\mathrm{d}}{\mathrm{d}\ln\mu}\,\ln\Big|\widetilde{\varepsilon}_{\tilde{B},[0]}(\mu)\Big|, \tag{7.571}$$

$$\text{or discretely} \quad \widehat{y}_{\tilde{B}}(\mu_i) := \frac{\ln\Big|\widetilde{\varepsilon}_{\tilde{B},[0]}(\mu_{i+1})\Big| - \ln\Big|\widetilde{\varepsilon}_{\tilde{B},[0]}(\mu_i)\Big|}{\ln\mu_{i+1} - \ln\mu_i}. \tag{7.572}$$

On any certified plateau (cf. Definition 6.297) the estimator is approximately scale–independent and reads off $y_{\tilde{B}}$ within tolerance. All observables entering $\widetilde{\varepsilon}_{\tilde{B},[0]}$ are understood with the Nielsen projection applied (cf. Definition 7.448).

Intuition. Take a log–derivative (or its discrete version) of the Nielsen–projected eigen–coupling along the RG flow eq. (7.567); by eq. (7.569) any Jordan–block polylog factors only modulate the slope transiently, so on a plateau (Definition 6.297) $\widehat{y}_{\tilde{B}}$ stabilises to $y_{\tilde{B}}$.

Proposition 7.453 (Convergence of the measured exponent)**.** *Under the assumptions of Proposition 7.450 and with μ restricted to the Step 3 certified IR half–space (mask $\mathcal{M}^{[\mathcal{U}]}$ of Proposition 7.427), the estimator in eq. (7.572) (see Definition 7.452) satisfies*

$$\widehat{y}_{\tilde{B}}(\mu) = y_{\tilde{B}} + \mathcal{O}\Big(\frac{1}{|\ln(\mu/\mu_0)|}\Big) \quad \text{as } \mu \downarrow 0,$$

and the discrete version converges to the same limit as the grid in $\ln\mu$ is refined. The estimate is gauge–parameter independent within the tolerance set by the Nielsen projection (cf. Definition 7.457 and Corollary 7.459).

Intuition. Using the Jordan expansion of the projected flow (Proposition 7.450 and Definition 7.449), each mode behaves like a power $\mu^{y_{\tilde{B}}}$ times a polynomial in $\ln\mu$, so the log–derivative peels off $y_{\tilde{B}}$ up to $1/\ln\mu$ corrections. Inside the Step 3 mask (Proposition 7.427) the nonlinear remainder is small and decays.

Proof. Let $t := \ln(\mu/\mu_0) \to -\infty$ as $\mu \downarrow 0$ and let $\widetilde{\varepsilon}_{\tilde{B}}$ denote the eigen–coupling component defined in Definition 7.454 (and used in Definition 7.452). By Proposition 7.450 together with the Jordan structure of Definition 7.449, we can write

$$\widetilde{\varepsilon}_{\tilde{B}}(\mu) = \mathrm{e}^{y_{\tilde{B}}t}\Big(P_{\tilde{B}}(t) + r_{\tilde{B}}(t)\Big), \tag{7.573}$$

where $P_{\tilde{B}}$ is a polynomial of degree at most $\operatorname{ind}\mathsf{N}_{\tilde{B}} - 1$ (degree 0 if $Y^{\mathcal{N}}$ is diagonalisable on the $y_{\tilde{B}}$ sector) and $r_{\tilde{B}}(t)$ collects the nonlinear $\mathcal{O}(\widetilde{\varepsilon}^2)$ remainder coming from the projected RG eq. (7.567).

Step 1 (control from the Step 3 mask). Inside $\mathcal{M}^{[\mathcal{U}]}$ (cf. Proposition 7.427) the breaking sector is small and decaying, so there exists $\delta > 0$ with

$$r_{\tilde{B}}(t) = \mathcal{O}(\mathrm{e}^{\delta t}), \qquad r'_{\tilde{B}}(t) = \mathcal{O}(\mathrm{e}^{\delta t}), \tag{7.574}$$

obtained by differentiating the projected flow on $W = \operatorname{im}\mathcal{N}$ (cf. Definition 7.457) and applying a Grönwall estimate to the $\mathcal{O}(\tilde{\varepsilon}^2)$ remainder.

Step 2 (log–derivative). Define $f_{\tilde{B}}(t) := \ln \widetilde{\varepsilon}_{\tilde{B}}(\mu)$. From eq. (7.573),

$$\frac{\mathrm{d}f_{\tilde{B}}}{\mathrm{d}t}(t) = y_{\tilde{B}} + \frac{P'_{\tilde{B}}(t) + r'_{\tilde{B}}(t)}{P_{\tilde{B}}(t) + r_{\tilde{B}}(t)}.$$

If $\deg P_{\tilde{B}} = m \geq 0$, then

$$\frac{P'_{\tilde{B}}(t)}{P_{\tilde{B}}(t)} = \frac{m}{t} + \mathcal{O}\Big(\frac{1}{t^2}\Big). \tag{7.575}$$

Combining eq. (7.575) with eq. (7.574) and the fact that $|P_{\tilde{B}}(t)| \gtrsim |t|^m$ for large $|t|$ (or $P_{\tilde{B}} \to a_0 \neq 0$ when $m = 0$), we get

$$\frac{\mathrm{d}f_{\tilde{B}}}{\mathrm{d}t}(t) = y_{\tilde{B}} + \mathcal{O}\Big(\frac{1}{|t|}\Big), \quad t \to -\infty.$$

Returning to μ (since $\frac{\mathrm{d}}{\mathrm{d}\ln\mu} = \frac{\mathrm{d}}{\mathrm{d}t}$) yields

$$\widehat{y}_{\tilde{B}}(\mu) = \frac{\mathrm{d}}{\mathrm{d}\ln\mu} \ln \widetilde{\varepsilon}_{\tilde{B}}(\mu) = y_{\tilde{B}} + \mathcal{O}\Big(\frac{1}{|\ln(\mu/\mu_0)|}\Big).$$

Step 3 (discrete estimator). For $t_i = \ln(\mu_i/\mu_0)$ with step $h_i := t_{i+1} - t_i$, the forward difference in eq. (7.572) can be written as

$$\widehat{y}_{\tilde{B}}(\mu_i) = \frac{f_{\tilde{B}}(t_{i+1}) - f_{\tilde{B}}(t_i)}{h_i} = \frac{\mathrm{d}f_{\tilde{B}}}{\mathrm{d}t}(t_i^*), \quad t_i^* \in [t_i, t_{i+1}],$$

so by Step 2 we have $\widehat{y}_{\tilde{B}}(\mu_i) = y_{\tilde{B}} + \mathcal{O}(1/|t_i^*|)$. If $h_i \to 0$ (grid refinement) while $t_i \to -\infty$, then $t_i^* \to -\infty$ and $\widehat{y}_{\tilde{B}}(\mu_i) \to y_{\tilde{B}}$. (A centred difference adds an $\mathcal{O}(h_i^2)$ truncation error; the forward difference adds $\mathcal{O}(h_i)$, both vanishing upon refinement.)

Step 4 (gauge–parameter independence). The measurement uses the Nielsen–projected eigen–coupling (cf. Definitions 7.452 and 7.457). The projected spectrum is gauge–independent by the isospectral similarity flow (Proposition 7.458 and Corollary 7.459); any residual dependence is bounded by the tolerance in eq. (7.588). This proves the claim. □

Intuition. After diagonalising the projected flow, each mode looks like $\mu^{y_{\tilde{B}}}$ times at most a polynomial in $\ln\mu$; the log–derivative peels off $y_{\tilde{B}}$ with corrections $\sim 1/\ln\mu$. Discretisation and gauge choices only add vanishing (or tolerance–bounded) effects once we are inside the Step 3 IR mask.

Diagonalisation and eigen–operators.

Definition 7.454 (Eigen–operators and eigen–couplings)**.** Work on the physical subspace $W := \operatorname{im}\mathcal{N}$ (cf. Definition 7.448) and let $Y^{\mathcal{N}} := \mathcal{N}\,Y\,\mathcal{N}\big|_W$ be the projected stability matrix from eq. (7.567). Choose R whose columns form a (generalised) right Jordan basis on W so that

$$R^{-1}Y^{\mathcal{N}}R \;=\; J \;=\; \operatorname{diag}(J_{\tilde{B}}), \qquad J_{\tilde{B}} = y_{\tilde{B}}\mathbb{1} + \mathsf{N}_{\tilde{B}}, \quad (\mathsf{N}_{\tilde{B}})^{m_{\tilde{B}}} = 0 \text{ (cf. } \textit{Definition } 7.449).$$

Define the *eigen–operators* and *eigen–couplings* by the dual change of basis

$$\boxed{\widetilde{\mathcal{O}}_{\tilde{B}} := \sum_{A} (R^{-1})_{\tilde{B}A}\, \hat{\mathcal{O}}_A, \qquad \widetilde{\varepsilon}_{\tilde{B}} := \sum_{A} R_{A\tilde{B}}\, \varepsilon_A,} \tag{7.576}$$

so that the breaking sector is expressed in the (generalised) eigenbasis of $Y^{\mathcal{N}}$. In the diagonalisable case one has $J = \mathrm{diag}(y_{\tilde{B}})$ and $\mathsf{N}_{\tilde{B}} = 0$.

Intuition. This is the Nielsen–projected change of basis aligning operators and couplings with RG eigendirections on W (Definition 7.448); when $Y^{\mathcal{N}}$ is diagonalisable, each mode runs independently at linear order (cf. eq. (7.567)).

Proposition 7.455 (Decoupled linear flow and solution)**.** *Assume $Y^{\mathcal{N}}$ (on $W = \mathrm{im}\,\mathcal{N}$) is diagonalisable over $\mathbb{C}$ with eigenvalues $\{y_{\tilde{B}}\}$ and adopt the eigenbasis of Definition 7.454. Then the linearised RG flow eq. (7.567) decouples componentwise:*

$$\boxed{\mu \frac{\mathrm{d}\,\widetilde{\varepsilon}_{\tilde{B}}}{\mathrm{d}\mu} = y_{\tilde{B}}\, \widetilde{\varepsilon}_{\tilde{B}} + \mathcal{O}(\|\widetilde{\varepsilon}\|^2), \qquad \Rightarrow \quad \widetilde{\varepsilon}_{\tilde{B}}(\mu) = \widetilde{\varepsilon}_{\tilde{B}}(\mu_0) \left(\frac{\mu}{\mu_0}\right)^{y_{\tilde{B}}} \Big[1 + \mathcal{O}_I\Big(\sup_{\mu\in I} \|\widetilde{\varepsilon}(\mu)\|\Big)\Big],}$$

(7.577)

where I is any compact interval in $\ln \mu$ contained in the Step 3 certified IR mask (cf. Proposition 7.427). If $Y^{\mathcal{N}}$ has Jordan blocks, the linear evolution acquires only polynomial factors in $\ln(\mu/\mu_0)$, as stated in Proposition 7.450.

Intuition. After Nielsen projection (Definition 7.448) and diagonalisation, each projected mode follows a one–dimensional linear RG with exponent $y_{\tilde{B}}$; nonlinearity is subleading on the IR mask (Proposition 7.427). In the non–diagonalisable case, see the Jordan-form solution in Proposition 7.450.

Proof. **(1) Preliminaries and change of variables.** Write the projected breaking–sector RG (to quadratic order) as

$$\mu \frac{\mathrm{d}\varepsilon}{\mathrm{d}\mu} = Y^{\mathcal{N}} \varepsilon + \mathcal{R}(\varepsilon, \mu), \qquad \mathcal{R}(\varepsilon, \mu) = \mathcal{O}(\|\varepsilon\|^2), \tag{7.578}$$

with $Y^{\mathcal{N}}$ from eq. (7.567) and $\varepsilon \in W := \mathrm{im}\,\mathcal{N}$ (cf. Definition 7.448). Set $t := \ln(\mu/\mu_0)$ so that $\frac{\mathrm{d}}{\mathrm{d}t} = \mu \frac{\mathrm{d}}{\mathrm{d}\mu}$, giving

$$\frac{\mathrm{d}\varepsilon}{\mathrm{d}t} = Y^{\mathcal{N}} \varepsilon + r(\varepsilon, t), \qquad r(\varepsilon, t) := \mathcal{R}(\varepsilon, \mu(t)). \tag{7.579}$$

Let R be as in Definition 7.454 with $R^{-1} Y^{\mathcal{N}} R = J = \mathrm{diag}(y_{\tilde{B}})$ (diagonalisable case). Define eigen–coordinates $\widetilde{\varepsilon} := R^{-1}\varepsilon$. Then

$$\frac{\mathrm{d}\widetilde{\varepsilon}}{\mathrm{d}t} = J\, \widetilde{\varepsilon} + \widetilde{r}(\widetilde{\varepsilon}, t), \qquad \widetilde{r}(\widetilde{\varepsilon}, t) := R^{-1} r(R\widetilde{\varepsilon}, t) = \mathcal{O}(\|\widetilde{\varepsilon}\|^2), \tag{7.580}$$

since $R^{\pm 1}$ are bounded on the finite–dimensional space W.

(2) Variation of constants and decoupling. Duhamel's formula for eq. (7.580) reads

$$\widetilde{\varepsilon}(t) = \mathrm{e}^{(t-t_0)J}\,\widetilde{\varepsilon}(t_0) + \int_{t_0}^{t} \mathrm{e}^{(t-s)J}\,\widetilde{r}\big(\widetilde{\varepsilon}(s), s\big)\,\mathrm{d}s.$$

Because J is diagonal, the *linear* term decouples componentwise:

$$\big(\mathrm{e}^{(t-t_0)J}\,\widetilde{\varepsilon}(t_0)\big)_{\tilde{B}} = \mathrm{e}^{y_{\tilde{B}}(t-t_0)}\,\widetilde{\varepsilon}_{\tilde{B}}(t_0).$$

The integral couples components only through $\widetilde{r} = \mathcal{O}(\|\widetilde{\varepsilon}\|^2)$ and is higher order.

(3) Uniform control on the IR mask. Restrict to a compact interval $I = [t_-, t_+]$ contained in the Step 3 certified IR half–space (mask $\mathcal{M}^{[\mathcal{U}]}$, Proposition 7.427). Smallness of the breaking sector on I implies the bound

$$\Big\| \int_{t_0}^{t} \mathrm{e}^{(t-s)J}\,\widetilde{r}(\widetilde{\varepsilon}(s), s)\,\mathrm{d}s \Big\| \le K_I \left(\sup_{u\in I} \|\widetilde{\varepsilon}(u)\| \right)^2$$

for some finite K_I independent of $\widetilde{\varepsilon}$. Hence

$$\widetilde{\varepsilon}_{\tilde{B}}(t) = \mathrm{e}^{y_{\tilde{B}}(t-t_0)}\,\widetilde{\varepsilon}_{\tilde{B}}(t_0)\left[1 + \mathcal{O}_I\Big(\sup_{u\in I} \|\widetilde{\varepsilon}(u)\| \Big)\right].$$

(4) Conclusion. Returning to μ via $t = \ln(\mu/\mu_0)$ yields eq. (7.577). If $Y^{\mathcal{N}}$ is not diagonalisable, replace J by its Jordan form and recover the polynomial factors in $\ln(\mu/\mu_0)$ as in Proposition 7.450 and Definition 7.449. □

Corollary 7.456 (Irrelevance criterion)**.** *Work on the physical subspace $W := \operatorname{im}\mathcal{N}$ and let $\widetilde{\varepsilon}_{\tilde{B}}(\mu)$ be the eigen–couplings associated with the eigen–operators $\widetilde{\mathcal{O}}_{\tilde{B}}$ from Definition 7.454, built using the projected matrix $Y^{\mathcal{N}} := \mathcal{N}Y\mathcal{N}\big|_W$ (cf. Definition 7.448 and eq. (7.567)). If* $\operatorname{Re} y_{\tilde{B}} < 0$ *for all projected breaking modes, then $\widetilde{\varepsilon}_{\tilde{B}}(\mu) \to 0$ as $\mu \downarrow 0$. Equivalently, each $\widetilde{\mathcal{O}}_{\tilde{B}}$ is* irrelevant *in the Wilsonian sense.*

Proof. Set $t := \ln(\mu/\mu_0) \le 0$. By Proposition 7.455 in the diagonalisable case (or Proposition 7.450 together with Definition 7.449 for general Jordan blocks), the projected RG flow in the eigenbasis on W satisfies

$$\frac{\mathrm{d}\,\widetilde{\varepsilon}_{\tilde{B}}}{\mathrm{d}t} = y_{\tilde{B}}\,\widetilde{\varepsilon}_{\tilde{B}} + \mathcal{O}(\|\widetilde{\varepsilon}\|^2), \qquad \widetilde{\varepsilon}_{\tilde{B}}(t) = \mathrm{e}^{y_{\tilde{B}}(t-t_0)}\widetilde{\varepsilon}_{\tilde{B}}(t_0)\left[1 + \mathcal{O}_I\Big(\sup_{s\in[t,t_0]} \|\widetilde{\varepsilon}(s)\| \Big)\right],$$

uniformly on any compact interval $[t, t_0] \subset$ the Step 3 certified IR mask (cf. Proposition 7.427). Since $\operatorname{Re} y_{\tilde{B}} < 0$, we have $\mathrm{e}^{\operatorname{Re} y_{\tilde{B}}(t-t_0)} \to 0$ as $t \to -\infty$ (i.e. $\mu \downarrow 0$), and the remainder is subleading for sufficiently small initial data by the same control used in Proposition 7.455; hence $\widetilde{\varepsilon}_{\tilde{B}}(\mu) \to 0$.

In the presence of Jordan blocks, Proposition 7.450 yields

$$\widetilde{\varepsilon}_{\tilde{B}}(t) = \mathrm{e}^{y_{\tilde{B}}(t-t_0)} \sum_{k=0}^{m_{\tilde{B}}-1} \frac{(t-t_0)^k}{k!}\, c_{\tilde{B},k} \;+\; \mathcal{O}(\|\widetilde{\varepsilon}\|^2),$$

so $\mathrm{e}^{\operatorname{Re} y_{\tilde{B}}(t-t_0)}(t-t_0)^k \to 0$ for each fixed k as $t \to -\infty$, again implying $\widetilde{\varepsilon}_{\tilde{B}}(\mu) \to 0$. Thus every projected breaking mode decays toward the IR and is irrelevant. □

Remark (Nielsen projection, gauge/scheme consistency). All statements are made on $W = \operatorname{im}\mathcal{N}$ (cf. Definition 7.448). Using $Y^{\mathcal{N}} := \mathcal{N}Y\mathcal{N}\big|_W$ and (block-)Jordanising it as in Definitions 7.449 and 7.454 ensures that the exponents $y_{\tilde{B}}$ are gauge–parameter independent (isospectral similarity flow of Proposition 7.458, with tolerance as in eq. (7.588)) and basis/scheme covariant (Lemma 7.451).

Nielsen projection (gauge–parameter safety).

Definition 7.457 (Nielsen identity and on–shell/BRST projection)**.** Let α denote the (set of) gauge parameters (e.g. the R_ξ parameter ξ is one component). Gauge–parameter derivatives of the 1PI effective action satisfy the Nielsen identity

$$\boxed{\frac{\partial\Gamma[\phi;\alpha]}{\partial\alpha} + \int \mathrm{d}^D x\, \mathcal{C}_i(\phi,\alpha;x)\,\frac{\delta\Gamma[\phi;\alpha]}{\delta\phi_i(x)} = 0,} \tag{7.581}$$

so gauge–parameter dependence vanishes at stationary points $\delta\Gamma/\delta\phi = 0$ and for on–shell BRST–invariant observables (see [117, 243]). Let $\mathcal{N}(\alpha)$ denote the linear projector on the breaking–operator space V onto the BRST–physical cohomology, evaluated on shell and compatible with the Slavnov–Taylor identity (cf. eq. (7.16)):

$$\mathcal{N}^2 = \mathcal{N}, \qquad \operatorname{im}\mathcal{N} \cong H^0_{\mathrm{BRST}},$$
$$\partial_\alpha\big(\mathcal{N}X\big) = 0 \quad \text{for all physical } X,$$
(within admissible tolerance; cf. *eq.* (7.588)).

For any mixing/flow matrices Z, Y on V (cf. eq. (7.567)), define their Nielsen–projected versions on the physical subspace $W := \operatorname{im}\mathcal{N}$ by restriction:

$$Z^{\mathcal{N}}(\alpha) := \mathcal{N}(\alpha)\,Z(\alpha)\,\mathcal{N}(\alpha)\big|_W, \qquad Y^{\mathcal{N}}(\alpha) := \mathcal{N}(\alpha)\,Y(\alpha)\,\mathcal{N}(\alpha)\big|_W.$$

The eigenvalues $y_{\tilde{B}}(\alpha)$ of $Y^{\mathcal{N}}(\alpha)$ are the *Nielsen–projected breaking exponents*; associated right eigenvectors in W define the physical eigen–operators/eigen–couplings as in Definition 7.454. By the isospectral similarity flow, these exponents are gauge–parameter independent within tolerance (cf. Proposition 7.458 and Corollary 7.459).

Intuition. Project *after* going on shell and into BRST cohomology: $Y^{\mathcal{N}}$ then captures the physically meaningful mixing/flow (eq. (7.567)). The spectrum of $Y^{\mathcal{N}}$ is stable under gauge–parameter variations (within eq. (7.588)) by Proposition 7.458 and Corollary 7.459, and basis/scheme choices act by similarity (Lemma 7.451).

Proposition 7.458 (Similarity flow of the projected stability matrix)**.** *Under the Nielsen identity eq. (7.581) and evaluation on shell with BRST projection as in Definition 7.457, there exists a smooth matrix $K(\alpha)$ on the physical subspace $W := \operatorname{im}\mathcal{N}$ such that, for any gauge–parameter component α,*

$$\frac{\partial Y^{\mathcal{N}}}{\partial\alpha} = [\,K(\alpha),\, Y^{\mathcal{N}}(\alpha)\,].$$

Consequently,

$$Y^{\mathcal{N}}(\alpha) = S(\alpha,\alpha_0)\, Y^{\mathcal{N}}(\alpha_0)\, S(\alpha,\alpha_0)^{-1}, \qquad S(\alpha,\alpha_0) := \mathcal{P}\exp\Big(\int_{\alpha_0}^{\alpha} K(\zeta)\,\mathrm{d}\zeta\Big),$$

and the projected breaking spectrum $\{y_{\tilde{B}}(\alpha)\}$ *is independent of* α *(isospectrality).*

Proof. **Step 0 (Projector variation drops on W).** By the on–shell/BRST construction of Definition 7.457, $\partial_\alpha(\mathcal{N}X) = 0$ for any physical X (within the tolerance of eq. (7.588)). Hence, when differentiating projected objects like $Z^{\mathcal{N}} := \mathcal{N}Z\mathcal{N}|_W$ and $Y^{\mathcal{N}} := \mathcal{N}Y\mathcal{N}|_W$, the $\partial_\alpha\mathcal{N}$ terms vanish on W.

Step 1 (From Nielsen to an on–shell generator on operator space). The Nielsen identity eq. (7.581) implies that a gauge–parameter variation is a field redefinition generated by a local functional $\mathcal{C}_i$, so insertions of renormalised composite operators transform as

$$\partial_\alpha\Big[\mathcal{O}(\phi)\Big]^{\mathrm{R}} = [\,K_\alpha,\,\mathcal{O}^{\mathrm{R}}\,] + (\text{BRST exact}), \quad \text{see [117, 243]}.$$

Projecting to $W := \operatorname{im}\mathcal{N}$ on shell removes BRST–exact pieces (Definition 7.457), so there is a smooth matrix $K(\alpha)$ on W with

$$\partial_\alpha\mathcal{O}^{\mathrm{R}}_W = [\,K(\alpha),\,\mathcal{O}^{\mathrm{R}}_W\,], \qquad \mathcal{O}^{\mathrm{R}}_W := \mathcal{N}\,\mathcal{O}^{\mathrm{R}}\,\mathcal{N}\Big|_W. \tag{7.582}$$

Step 2 (Implications for the projected renormalisation matrix). Let $Z(\alpha,\mu)$ be the operator mixing matrix on the breaking sector (eq. (7.567)) and define $Z^{\mathcal{N}} := \mathcal{N}Z\mathcal{N}\big|_W$. Bare operators are α–independent, and $\mathcal{O}^{\mathrm{R}}_W = Z^{\mathcal{N}}\mathcal{O}^{bare}_W$; using eq. (7.582) and Step 0 gives

$$\partial_\alpha Z^{\mathcal{N}} = A\,Z^{\mathcal{N}} + Z^{\mathcal{N}}B, \qquad A := L_K, \quad B := -R_K, \tag{7.583}$$

i.e. $AX - XB = [K,X]$ for any matrix X on W (left/right actions by K).

Step 3 (Lax equation for $\gamma^{\mathcal{N}}$ and $Y^{\mathcal{N}}$). Define $\gamma^{\mathcal{N}} := Z^{\mathcal{N}-1}\,\mu\,\frac{\mathrm{d}Z^{\mathcal{N}}}{\mathrm{d}\mu}$ and separate canonical dimensions so that $Y^{\mathcal{N}} := d - \gamma^{\mathcal{N}}$ (cf. the conventions below eq. (7.567)). Differentiating $\gamma^{\mathcal{N}}$ w.r.t. α and using eq. (7.583) yields

$$\begin{aligned}\partial_\alpha\gamma^{\mathcal{N}} &= (\partial_\alpha Z^{\mathcal{N}-1})\,\mu\frac{\mathrm{d}Z^{\mathcal{N}}}{\mathrm{d}\mu} + Z^{\mathcal{N}-1}\,\mu\frac{\mathrm{d}}{\mathrm{d}\mu}(\partial_\alpha Z^{\mathcal{N}}) \\ &= [\,\gamma^{\mathcal{N}},\,B\,] \;+\; Z^{\mathcal{N}-1}\mu\frac{\mathrm{d}A}{\mathrm{d}\mu}Z^{\mathcal{N}} \;+\; \mu\,\frac{\mathrm{d}B}{\mathrm{d}\mu}.\end{aligned} \tag{7.584}$$

On shell, RG flow and gauge–parameter variations commute up to ad_K; differentiating eq. (7.583) w.r.t. μ at fixed bare theory gives

$$\mu\,\frac{\mathrm{d}A}{\mathrm{d}\mu} = [\,A,\,\gamma^{\mathcal{N}}\,], \qquad \mu\,\frac{\mathrm{d}B}{\mathrm{d}\mu} = [\,\gamma^{\mathcal{N}},\,B\,]. \tag{7.585}$$

Insert eq. (7.585) into eq. (7.584) and rewrite $Z^{\mathcal{N}-1}[A,\gamma^{\mathcal{N}}]Z^{\mathcal{N}} = [\,Z^{\mathcal{N}-1}AZ^{\mathcal{N}},\,\gamma^{\mathcal{N}}\,]$ to get

$$\partial_\alpha\gamma^{\mathcal{N}} = [\,\gamma^{\mathcal{N}},\,B - Z^{\mathcal{N}-1}AZ^{\mathcal{N}}\,].$$

Define

$$K(\alpha) := B(\alpha) - Z^{\mathcal{N}-1}(\alpha)\, A(\alpha)\, Z^{\mathcal{N}}(\alpha) \quad \text{on } W, \tag{7.586}$$

then

$$\partial_\alpha \gamma^{\mathcal{N}} = [\,\gamma^{\mathcal{N}},\, K(\alpha)\,]. \tag{7.587}$$

Since d is α–independent, we conclude

$$\partial_\alpha Y^{\mathcal{N}} = [\,K(\alpha),\, Y^{\mathcal{N}}(\alpha)\,],$$

i.e. a Lax equation in the gauge parameter.

Step 4 (Integration and isospectrality). The Lax equation integrates to the path–ordered similarity flow

$$Y^{\mathcal{N}}(\alpha) = S(\alpha,\alpha_0)\, Y^{\mathcal{N}}(\alpha_0)\, S(\alpha,\alpha_0)^{-1}, \qquad S(\alpha,\alpha_0) = \mathcal{P}\exp\Big(\int_{\alpha_0}^{\alpha} K(\zeta)\,\mathrm{d}\zeta\Big).$$

Similarity preserves spectra, hence the projected breaking exponents $y_{\tilde{B}}(\alpha)$ (eigenvalues of $Y^{\mathcal{N}}(\alpha)$) are independent of α. This is consistent with the gauge–independence statements collected in Corollary 7.459. □

Intuition. Changing the gauge parameter is (on shell) a field redefinition (eq. (7.581)); after the Nielsen projection (Definition 7.457) it acts by conjugation on the physical operator space W, giving a Lax equation for $Y^{\mathcal{N}}$. The path–ordered solution is a similarity transform, so the eigenvalues $y_{\tilde{B}}$ are invariant.

Corollary 7.459 (Gauge–parameter independence of projected exponents)**.** *Under the hypotheses of Proposition 7.458, the multiset of eigenvalues of $Y^{\mathcal{N}}(\alpha)$ (with algebraic multiplicities) is independent of the gauge parameter component α:*

$$\{y_{\tilde{B}}(\alpha)\}_{\tilde{B}} \;=\; \{y_{\tilde{B}}(\alpha_0)\}_{\tilde{B}} \qquad \textit{for all admissible } \alpha,$$

where admissibility is as in the on–shell/BRST setting of Definition 7.457. In practice (finite statistics/truncations), acceptance demands the gauge–tolerance audit

$$\boxed{\sup_{\alpha\in\Xi_{\mathrm{adm}}} \big|y_{\tilde{B}}(\alpha) - y_{\tilde{B}}(\alpha_0)\big| \;\le\; \varepsilon_{\mathrm{gauge}},} \tag{7.588}$$

with Ξ_{adm} the admissible gauge set and $\varepsilon_{\mathrm{gauge}}$ chosen below plateau tolerances (cf. the certification pipeline and plateau criterion in Definition 6.297).

Proof. By Proposition 7.458, on the physical subspace $W := \operatorname{im}\mathcal{N}$ one has

$$Y^{\mathcal{N}}(\alpha) = S(\alpha,\alpha_0)\, Y^{\mathcal{N}}(\alpha_0)\, S(\alpha,\alpha_0)^{-1},$$

with $S(\alpha,\alpha_0) = \mathcal{P}\exp\big(\int_{\alpha_0}^{\alpha} K(\zeta)\,\mathrm{d}\zeta\big)$ invertible on W. Therefore the characteristic polynomials coincide:

$$\begin{aligned}
\det\big(\lambda\mathbb{1} - Y^{\mathcal{N}}(\alpha)\big) &= \det\big(\lambda\mathbb{1} - SY^{\mathcal{N}}(\alpha_0)S^{-1}\big) \\
&= \det\big(S(\lambda\mathbb{1} - Y^{\mathcal{N}}(\alpha_0))S^{-1}\big) \\
&= \det\big(\lambda\mathbb{1} - Y^{\mathcal{N}}(\alpha_0)\big),
\end{aligned}$$

so the spectra (as multisets with algebraic multiplicities) are identical for all admissible α. Degenerate eigenspaces may rotate by $S(\alpha, \alpha_0)$, but eigenvalues—the projected exponents defined via Definitions 7.454 and 7.457—are unchanged. Inequality (7.588) records the practical tolerance used when validating gauge–independence numerically on certified plateaux (cf. Definition 6.297). □

Intuition. After the Nielsen projection (Definition 7.457), changing the gauge parameter acts by a similarity transform on $Y^{\mathcal{N}}$ (Proposition 7.458); similarity preserves eigenvalues, so the measured exponents are gauge–invariant up to audit tolerance (7.588).

Lemma 7.460 (Compatibility with basis/scheme changes)**.** *Let V be the breaking–operator space and $W := \operatorname{im}\mathcal{N} \subset V$ the on–shell BRST–physical subspace (cf. Definition 7.457). Let $M(\alpha) \in \mathrm{GL}(V)$ be an admissible (gauge–respecting) linear mixing of breaking operators that preserves W, and let the renormalisation scheme vary smoothly with α while preserving evaluation on shell and W. Then*

$$Y^{\mathcal{N}} \;\longmapsto\; \widetilde{Y}^{\mathcal{N}} = (\mathcal{N}M\mathcal{N})\, Y^{\mathcal{N}}\, (\mathcal{N}M^{-1}\mathcal{N})\Big|_W,$$

i.e. a similarity on W, so the multiset $\{y_{\tilde{B}}\}$ of projected breaking exponents is invariant (cf. Definition 7.454). Moreover, any finite (invertible) renormalisation $F(g, \alpha)$ that preserves W leaves $\{y_{\tilde{B}}\}$ unchanged at the IR fixed point $g_\star$ (cf. Definition 7.447).

Proof. **(A) Basis change (on W).** This is the Nielsen–projected version of Lemma 7.451. A basis change $\mathcal{O} \mapsto \mathcal{O}' := M\mathcal{O}$ induces $Y \mapsto MYM^{-1}$ at linear order (with dual couplings transformed to preserve the breaking pairing; cf. eq. (7.567)). Since $MW \subseteq W$, the restricted map $M_W := \mathcal{N}M\mathcal{N}\big|_W$ is invertible on W, and

$$Y^{\mathcal{N}} = \mathcal{N}Y\mathcal{N}\Big|_W \;\longmapsto\; \widetilde{Y}^{\mathcal{N}} = \mathcal{N}(MYM^{-1})\mathcal{N}\Big|_W = M_W\, Y^{\mathcal{N}}\, M_W^{-1},$$

a similarity on W. Therefore the spectrum $\{y_{\tilde{B}}\}$ is unchanged (see also Definition 7.454).

(B) Scheme change at the fixed point. Let $Z(g, \alpha, \mu)$ be the mixing matrix, $\gamma := Z^{-1}\mu\,\frac{\mathrm{d}Z}{\mathrm{d}\mu}$, and separate canonical dimensions so that $Y := d - \gamma$ (conventions below eq. (7.567)). A finite renormalisation $F(g, \alpha)$ produces $Z \mapsto Z' := FZ$ and

$$\gamma' \;=\; Z'^{-1}\mu\frac{\mathrm{d}Z'}{\mathrm{d}\mu} \;=\; F^{-1}\gamma F + F^{-1}\left(\beta^a\partial_{g^a}F + \partial_\alpha F\;\mu\frac{\mathrm{d}\alpha}{\mathrm{d}\mu}\right).$$

Projecting to W after going on shell (Nielsen identity, Definition 7.457 and eq. (7.581)) removes the explicit α–term in cohomology; if F preserves W define $F_W := \mathcal{N}F\mathcal{N}\big|_W$. At an IR fixed point $g_\star$ with $\beta(g_\star) = 0$ (cf. Definition 7.447),

$$\gamma_\star^{\mathcal{N}} \;\longmapsto\; \gamma_\star^{\mathcal{N}'} = F_W^{-1}\,\gamma_\star^{\mathcal{N}}\,F_W.$$

Assuming F_W mixes only operators within equal canonical–dimension blocks (so $[F_W, d] = 0$), we obtain

$$Y_\star^{\mathcal{N}} = d - \gamma_\star^{\mathcal{N}} \;\longmapsto\; Y_\star^{\mathcal{N}'} = F_W^{-1}(d - \gamma_\star^{\mathcal{N}})F_W = F_W^{-1}Y_\star^{\mathcal{N}}F_W,$$

a similarity on W. Hence the projected exponents are scheme–independent at the fixed point (see also Lemma 7.451). □

Remark. The present lemma is equivalent in content to Lemma 7.451 but stated explicitly after the Nielsen projection (Definition 7.457). Together with the similarity flow in the gauge parameter (Proposition 7.458) and the gauge–tolerance audit (7.588), it ensures that the projected exponents $\{y_{\tilde{B}}\}$ are invariant under admissible basis relabellings, scheme changes at $g_\star$, and gauge–parameter variations on the certified band.

IR window and consistency with the restoration mask.

Definition 7.461 (IR μ–window induced by the restoration mask)**.** Let $\mu = \kappa/\ell$ as in Definition 7.431 and write $v := \ln \ell$. Given any admitted IR ℓ–window (uniform in u)

$$\mathcal{I}_\ell := \{\ \ell \geq \ell^{\mathrm{IR}}_{\min}\ \} \qquad \text{within which the Step 3 guards hold for all } u \in \mathcal{U},\ \text{i.e.}$$
eqs. (7.542)*and* (7.548) (and the plateau/certification conditions of Definition 6.297),

define the corresponding μ–window by the order–reversing map (cf. $\mu = \kappa/\ell$)

$$\mathcal{I}_\mu := \{\ \mu \leq \mu^{\mathrm{IR}}_{\max}\ \}, \qquad \mu^{\mathrm{IR}}_{\max} := \kappa/\ell^{\mathrm{IR}}_{\min}.$$

In particular, the Step 3 certified mask $\mathcal{M}^{[\mathcal{U}]} = \{(u, v) : u \in \mathcal{U},\ v \geq \ln \ell_R^{[\mathcal{U}]}\}$ (see Proposition 7.427) induces the μ–half–line

$$\mathcal{I}_\mu^{[\mathcal{U}]} := \{\ \mu \leq \mu_R^{[\mathcal{U}]}\ \}, \qquad \mu_R^{[\mathcal{U}]} := \kappa/\ell_R^{[\mathcal{U}]}.$$

Proposition 7.462 (Consistency of the RG window with law restoration)**.** *On $\mathcal{I}_\mu^{[\mathcal{U}]}$ the law–channel guards of Step 3 remain enforced: along every admissible UV slice $u \in \mathcal{U}$, the law–deficit stays within tolerance (plateau/acceptance per Definition 6.297 and Proposition 7.427) and the certified slope satisfies $\varsigma^{(\tau)}_{\mathrm{law}}(\ell) > 0$ for the corresponding $\ell = \kappa/\mu$ (cf. eqs.* (7.542) *and* (7.548)*). In conjunction with the irrelevance ledger (Corollary 7.456), the breaking sector is small and decaying on $\mathcal{I}_\mu^{[\mathcal{U}]}$, and the hypotheses of Proposition 7.453 hold there.*

Proof. By Proposition 7.427, for every $u \in \mathcal{U}$ and every $v = \ln \ell \geq \ln \ell_R^{[\mathcal{U}]}$ one has the Step 3 plateau and slope constraints, together with the monotonicity/guard conditions eqs. (7.542) and (7.548) (within the certification pipeline Definition 6.297). The map $v = \ln \ell \mapsto \mu = \kappa/\ell$ is smooth and order–reversing, so the condition $\mu \leq \kappa/\ell_R^{[\mathcal{U}]}$ is equivalent to $v \geq \ln \ell_R^{[\mathcal{U}]}$. Therefore the same constraints transport to $\mathcal{I}_\mu^{[\mathcal{U}]}$, establishing the stated enforcement on the μ–half–line.

Since the Step 3 mask also bounds the law–deficit on certified plateaux (Proposition 7.427 and Definition 6.297), the breaking sector remains small there. Together with the negativity of the projected breaking exponents (irrelevance criterion Corollary 7.456, derived from the linearised Jordan solution Proposition 7.450), this implies decay of the projected breaking components on $\mathcal{I}_\mu^{[\mathcal{U}]}$. Consequently, the assumptions required for the estimator convergence Proposition 7.453 are satisfied on this window. □

Lemma 7.463 (Estimator invariances, locality, and accuracy)**.** *Let $t_k := \ln \mu_k$ and suppose $t_{k\pm1}$ exist. Then:*

Scale choice.
eq. (7.568) is invariant under the audit constant κ in $\mu = \kappa/\ell$ (cf. Definition 7.431).

Nielsen projection (gauge safety).
If data are taken on $W = \operatorname{im}\mathcal{N}$ (Definition 7.457), then $\widehat{y}_{\tilde{B}}$ is gauge–parameter independent within the tolerance eq. (7.588) (cf. Proposition 7.458 and Corollary 7.459).

Locality.
On a uniform grid in t with step h, eq. (7.568) equals the centered difference $\widehat{y}_{\tilde{B}}(t_k) = \frac{f_{\tilde{B}}(t_{k+1}) - f_{\tilde{B}}(t_{k-1})}{2h}$, where $f_{\tilde{B}}(t) := \ln|\widetilde{\varepsilon}_{\tilde{B}}(\mu(t))|$.

Normalization invariance.
Multiplying $\widetilde{\varepsilon}_{\tilde{B}}$ by any μ–independent constant (e.g. from re–normalising the eigenvectors in Definition 7.454) leaves eq. (7.568) unchanged.

Truncation error.
If $f_{\tilde{B}} \in C^3$ on an interval containing t_{k-1}, t_k, t_{k+1}, then

$$\widehat{y}_{\tilde{B}}(t_k) = f'_{\tilde{B}}(t_k) + \mathcal{O}(h^2),$$

with the constant depending on $\sup|f^{(3)}_{\tilde{B}}|$ on that interval.

Asymptotic accuracy on the IR window.
On the certified half–line $\mathcal{I}^{[\mathcal{U}]}_{\mu}$ (cf. Definition 7.461 and Proposition 7.462), if $t_k \to -\infty$ and $h \to 0$, then by Proposition 7.453

$$\widehat{y}_{\tilde{B}}(t_k) = y_{\tilde{B}} + \mathcal{O}\Big(\frac{1}{|t_k|}\Big) + \mathcal{O}(h^2).$$

Low–pass stability.
If LP_{τ_μ} satisfies eqs. (7.537) and (7.538) (zero–th and first moment conditions), then

$$\widehat{y}^{(\tau_\mu)}_{\tilde{B}}(t_k) = \widehat{y}_{\tilde{B}}(t_k) + \mathcal{O}(\tau_\mu^2) + \mathcal{O}(h^2),$$

so smoothing does not bias the estimate to leading order on plateaux.

Proof. **(a) Scale choice.** Replacing $\kappa \mapsto a\kappa$ shifts $t = \ln\mu$ by the constant $\ln a$, so $t_{k\pm1} \mapsto t_{k\pm1} + \ln a$. The centered difference cancels this shift exactly, leaving eq. (7.568) unchanged.

(b) Nielsen projection. On $W = \operatorname{im}\mathcal{N}$ (Definition 7.457), the projected flow generator $Y^{\mathcal{N}}$ obeys the similarity (Lax) equation in the gauge parameter (Proposition 7.458), implying gauge–independent projected exponents (Corollary 7.459). The measured quantity in eq. (7.568) is a ratio of $\widetilde{\varepsilon}_{\tilde{B}}$ at nearby μ; any residual gauge–dependence from finite statistics/truncations is bounded by eq. (7.588) on the certified band.

(c) Locality. On a uniform grid $t_{k\pm1} = t_k \pm h$,

$$\widehat{y}_{\tilde{B}}(t_k) = \frac{1}{2h}\Big(\ln|\widetilde{\varepsilon}_{\tilde{B}}(t_k + h)| - \ln|\widetilde{\varepsilon}_{\tilde{B}}(t_k - h)|\Big) = \frac{f_{\tilde{B}}(t_{k+1}) - f_{\tilde{B}}(t_{k-1})}{2h}.$$

(d) Normalization invariance. If $\widetilde{\varepsilon}_{\tilde{B}} \mapsto c\,\widetilde{\varepsilon}_{\tilde{B}}$ with $c > 0$ independent of μ, then the numerator and denominator inside the log in eq. (7.568) each pick up a factor c, which cancels.

(e) Truncation error. Taylor expand $f_{\tilde{B}}$ at t_k:

$$f_{\tilde{B}}(t_{k\pm1}) = f_{\tilde{B}}(t_k) \pm h f'_{\tilde{B}}(t_k) + \tfrac{h^2}{2} f''_{\tilde{B}}(t_k) \pm \tfrac{h^3}{6} f'''_{\tilde{B}}(\xi_\pm),$$

for some $\xi_\pm$ between t_k and $t_{k\pm1}$. Subtract and divide by $2h$:

$$\widehat{y}_{\tilde{B}}(t_k) = \frac{f_{\tilde{B}}(t_{k+1}) - f_{\tilde{B}}(t_{k-1})}{2h} = f'_{\tilde{B}}(t_k) + \tfrac{h^2}{6}\Big(f'''_{\tilde{B}}(\xi_+) - f'''_{\tilde{B}}(\xi_-)\Big) = f'_{\tilde{B}}(t_k) + \mathcal{O}(h^2).$$

(f) Asymptotic accuracy on the IR window. On $\mathcal{I}_\mu^{[\mathcal{U}]}$ (Definition 7.461 and Proposition 7.462), Proposition 7.453 gives $f'_{\tilde{B}}(t) = y_{\tilde{B}} + \mathcal{O}(1/|t|)$ as $t \to -\infty$. Combining with (e) yields $\widehat{y}_{\tilde{B}}(t_k) = y_{\tilde{B}} + \mathcal{O}(1/|t_k|) + \mathcal{O}(h^2)$.

(g) Low–pass stability. Let LP_{τ_μ} be a linear time–invariant operator in t with kernel K_{τ_μ} satisfying the moment conditions in eqs. (7.537) and (7.538): $\int K_{\tau_\mu} = 1$ and $\int t\, K_{\tau_\mu}(t)\,\mathrm{d}t = 0$. A standard moment expansion shows that for smooth inputs $\widehat{y}_{\tilde{B}}$,

$$\mathsf{LP}_{\tau_\mu}[\widehat{y}_{\tilde{B}}](t_k) = \widehat{y}_{\tilde{B}}(t_k) + \tfrac{m_2(K)}{2}\,\tau_\mu^2\,\partial_t^2 \widehat{y}_{\tilde{B}}(t_k) + \mathcal{O}(\tau_\mu^3) = \widehat{y}_{\tilde{B}}(t_k) + \mathcal{O}(\tau_\mu^2),$$

hence the claimed bias. Discretisation contributes the $\mathcal{O}(h^2)$ term from (e). Choosing τ_μ below plateau tolerances (cf. Definition 6.297) keeps the bias subleading. □

Remark 7.464 (Intuition for the centered estimator and the audit path)**.** The centered estimator eq. (7.568) is a numerical stand–in for the local slope $\frac{\mathrm{d}}{\mathrm{d}\ln\mu}\ln|\widetilde{\varepsilon}_{\tilde{B}}|$, which equals the projected breaking exponent on plateaux. The key heuristics are:

(i) Why centered? On a certified plateau (Definition 6.297 and Propositions 7.427 and 7.462) the curve $f_{\tilde{B}}(t) := \ln|\widetilde{\varepsilon}_{\tilde{B}}(\mu(t))|$ is *almost linear* in $t = \ln\mu$. A centered difference approximates $f'_{\tilde{B}}(t_k)$ with $\mathcal{O}(h^2)$ bias (Lemma 7.463(e)), whereas a forward difference has only $\mathcal{O}(h)$ accuracy (Definition 7.452). Thus the centered form is less biased when the goal is to read off a constant slope.

(ii) Why use $|\cdot|$? Jordan blocks in the linearised solution (Proposition 7.450, Definition 7.449) add only *polynomial* factors in t, and possible phases from complex data do not affect the magnitude. Taking $\ln|\cdot|$ isolates the dominant power law, so the estimator tracks $y_{\tilde{B}}$ up to the $\mathcal{O}(1/|t|)$ corrections proved in Proposition 7.453.

(iii) Why it is scale–robust. Changing the audit constant in $\mu = \kappa/\ell$ (cf. Definition 7.431) shifts $t = \ln\mu$ by a constant; the centered difference cancels this shift exactly (Lemma 7.463(a)). Similarly, rescaling eigenvectors only multiplies $\widetilde{\varepsilon}_{\tilde{B}}$ by a μ–independent constant, which cancels inside the logarithmic ratio (Lemma 7.463(d)).

(iv) Why it is gauge–safe. After the Nielsen projection (Definition 7.457), changes in the gauge parameter act by a similarity transform on the projected flow generator (Proposition 7.458), hence leave the spectrum $y_{\tilde{B}}$ invariant (Corollary 7.459). The estimator therefore converges to a gauge–independent limit on the IR window (Definition 7.461 and Proposition 7.462), up to the audit tolerance (7.588) (Lemma 7.463(b)).

(v) Why smoothing helps (and when). On real data, high–frequency jitter in t contaminates differences. A low–pass with moment conditions eqs. (7.537) and (7.538) preserves constants and linear trends, so it attenuates noise without biasing the plateau slope to

leading order (Lemma 7.463(g)). Choosing the smoothing scale below the plateau tolerances (Definition 6.297) keeps the bias subleading.

In short: on the Step 3 IR window the projected eigen–coupling behaves like $\widetilde{\varepsilon}_{\tilde{B}}(\mu) \sim \mu^{y_{\tilde{B}}}$ up to polylogarithmic and higher–order corrections (Proposition 7.450 and Corollary 7.456); the centered, gauge–safe estimator reads off the constant slope $y_{\tilde{B}}$ with controllable $\mathcal{O}(h^2)$ and $\mathcal{O}(1/|t|)$ errors (Proposition 7.453, Lemma 7.463).

Proposition 7.465 (Consistency, rates, and mollified variant)**.** *Work on the IR window $\mathcal{I}_\mu^{[\mathcal{U}]}$ of Definition 7.461 and let $\widehat{y}_{\tilde{B}}$ be the centered estimator of Definition 7.446. Assume the hypotheses of Propositions 7.450 and 7.453 and that the Step 3 guards (plateau/acceptance and monotonicity/curvature) hold on the window (Definition 6.297, Proposition 7.427, and eq. (7.548)). On a uniform t–grid with step h ($t = \ln \mu$) one has*

$$\widehat{y}_{\tilde{B}}(\mu_k) = y_{\tilde{B}} + \mathcal{O}\Big(\tfrac{1}{|\ln(\mu_k/\mu_0)|}\Big) + \mathcal{O}(h^2),$$

and for the mollified estimator $\widehat{y}_{\tilde{B}}^{(\tau_\mu)} := \mathsf{LP}_{\tau_\mu}[\widehat{y}_{\tilde{B}}]$ with a low–pass satisfying the moment conditions eqs. (7.537) and (7.538),

$$\widehat{y}_{\tilde{B}}^{(\tau_\mu)}(\mu_k) = y_{\tilde{B}} + \mathcal{O}\Big(\tfrac{1}{|\ln(\mu_k/\mu_0)|}\Big) + \mathcal{O}(h^2) + \mathcal{O}(\tau_\mu^2).$$

Implicit constants are uniform on compact subintervals of the IR window and are controlled by the curvature/TV guards in eq. (7.548). In particular, $\widehat{y}_{\tilde{B}}(\mu_k) \to y_{\tilde{B}}$ as $k \to \infty$ (i.e. $\mu_k \downarrow 0$) and $h, \tau_\mu \to 0$.

Proof. Write $t_k := \ln \mu_k$ and $f_{\tilde{B}}(t) := \ln|\widetilde{\varepsilon}_{\tilde{B}}(\mu(t))|$ with $\widetilde{\varepsilon}_{\tilde{B}}$ the Nielsen–projected eigen–coupling (Definitions 7.454 and 7.457). On the IR window $\mathcal{I}_\mu^{[\mathcal{U}]}$ we have, by Proposition 7.462, small and decaying breakings and the Step 3 plateau/guard conditions (Definition 6.297, Proposition 7.427, and eq. (7.548)).

Step 1 (continuum bias). By Proposition 7.453,

$$f'_{\tilde{B}}(t) = \frac{\mathrm{d}}{\mathrm{d}t} \ln|\widetilde{\varepsilon}_{\tilde{B}}(\mu(t))| = y_{\tilde{B}} + \mathcal{O}\Big(\tfrac{1}{|t|}\Big) \qquad (t \to -\infty),$$

uniformly on the window thanks to the guards (which bound derivatives and variations).

Step 2 (discretisation error of the centered difference). On a uniform grid with step h, Lemma 7.463(c,e) gives

$$\widehat{y}_{\tilde{B}}(t_k) = \frac{f_{\tilde{B}}(t_{k+1}) - f_{\tilde{B}}(t_{k-1})}{2h} = f'_{\tilde{B}}(t_k) + \mathcal{O}(h^2),$$

with the constant controlled by $\sup_{[t_{k-1},t_{k+1}]} |f'''_{\tilde{B}}|$, which is finite on the window by the curvature/TV guard eq. (7.548). Combining with Step 1 yields

$$\widehat{y}_{\tilde{B}}(t_k) = y_{\tilde{B}} + \mathcal{O}\Big(\tfrac{1}{|t_k|}\Big) + \mathcal{O}(h^2).$$

Step 3 (mollification bias). If $\widehat{y}_{\tilde{B}}^{(\tau_\mu)} = \mathsf{LP}_{\tau_\mu}[\widehat{y}_{\tilde{B}}]$ with the low–pass satisfying the zero–th and first moment conditions eqs. (7.537) and (7.538), then a standard moment expansion gives $\widehat{y}_{\tilde{B}}^{(\tau_\mu)}(t_k) = \widehat{y}_{\tilde{B}}(t_k) + \mathcal{O}(\tau_\mu^2)$; see Lemma 7.463(g). This adds to the error from Step 2.

Step 4 (uniformity and convergence). All $\mathcal{O}(\cdot)$ terms above are uniform on compact subintervals of the IR window because the guards keep $f_{\tilde{B}}$ and its derivatives bounded and because of the smallness/decay ensured by Proposition 7.462. Therefore,

$$\widehat{y}_{\tilde{B}}(\mu_k) \to y_{\tilde{B}} \quad \text{as} \quad k \to \infty \ (\mu_k \downarrow 0), \quad h \to 0,$$

and, with smoothing, also $\tau_\mu \to 0$.

Optional invariances. The estimator is invariant under the audit constant in $\mu = \kappa/\ell$ and under μ–independent re–normalisations of eigenvectors by Lemma 7.463(a,d); gauge safety follows from Definition 7.457, Proposition 7.458, and Corollary 7.459. □

Definition 7.466 (Step 4 acceptance: irrelevance and gauge safety)**.** Work on the physical subspace $W := \operatorname{im}\mathcal{N}$ and on the IR window $\mathcal{I}_\mu^{[\mathcal{U}]}$ induced by the Step 3 mask (cf. Definition 7.461 and Proposition 7.462). The RG ledger for explicit breakings is *accepted* if:

1. **Irrelevance.** For every projected breaking eigendirection $\tilde{B}$ that is excited,

$$\operatorname{Re} y_{\tilde{B}} \ < \ 0 \quad \text{(eigenvalues of } Y^{\mathcal{N}}\text{; cf. Definition 7.454 and Corollary 7.456).}$$

Moreover, the combined (estimator + gauge) band for $y_{\tilde{B}}$ over $\mathcal{I}_\mu^{[\mathcal{U}]}$ does not cross 0: with $\widehat{y}_{\tilde{B}}$ as in Definition 7.446 and the tolerance audit (7.588),

$$0 \ \notin \ \Big[\inf_{\mu \in \mathcal{I}_\mu^{[\mathcal{U}]}} \widehat{y}_{\tilde{B}}(\mu) \ - \ \varepsilon_{\text{gauge}} \ , \quad \sup_{\mu \in \mathcal{I}_\mu^{[\mathcal{U}]}} \widehat{y}_{\tilde{B}}(\mu) \ + \ \varepsilon_{\text{gauge}} \Big].$$

2. **Flow suppression on the window.** For all such $\tilde{B}$,

$$\left|\widetilde{\varepsilon}_{\tilde{B}}(\mu)\right| \ \le \ \varepsilon_{\text{plat}} \qquad \text{for all} \ \ \mu \le \mu_R^{[\mathcal{U}]} = \kappa/\ell_R^{[\mathcal{U}]},$$

with $(\ell_R^{[\mathcal{U}]}, \mu_R^{[\mathcal{U}]})$ from Proposition 7.427 and Definition 7.461 and $\varepsilon_{\text{plat}}$ the plateau tolerance from the certification pipeline (cf. Definition 6.297). This is consistent with the linear decay implied by $\operatorname{Re} y_{\tilde{B}} < 0$ (cf. Proposition 7.450 and Corollary 7.456).

3. **Gauge safety.** Exponents are computed from the Nielsen–projected matrix $Y^{\mathcal{N}}$ (Definition 7.457), and satisfy

$$\boxed{\sup_{\xi \in \Xi_{\text{adm}}} \left| y_{\tilde{B}}(\xi) - y_{\tilde{B}}(\xi_0) \right| \ \le \ \varepsilon_{\text{gauge}},} \tag{7.589}$$

i.e. the gauge–tolerance audit (7.588) holds on $\mathcal{I}_\mu^{[\mathcal{U}]}$ (Proposition 7.458 and Corollary 7.459).

Proposition 7.467 (Soundness and robustness of Step 4 acceptance)**.** *If the criteria of Definition 7.466 hold on the IR window $\mathcal{I}_\mu^{[\mathcal{U}]}$ of Definition 7.461, then:*

(i) Every excited projected mode is RG–irrelevant in the Wilsonian sense: $\widetilde{\varepsilon}_{\tilde{B}}(\mu) \to 0$ as $\mu \downarrow 0$.

(ii) The Step 3 law–channel restoration is consistent with the measured running on $\mathcal{I}_\mu^{[\mathcal{U}]}$; in particular the hypotheses of Proposition 7.453 are satisfied on this window.

(iii) The quoted exponents $\{y_{\tilde{B}}\}$ are gauge–parameter independent up to the audit tolerance $\varepsilon_{\text{gauge}}$ and invariant under admissible basis/scheme changes (Corollary 7.459 and Lemma 7.451).

Proof. All statements are made on the physical subspace $W := \operatorname{im}\mathcal{N}$ after Nielsen projection (Definition 7.457). Let $Y^{\mathcal{N}}$ be the projected stability matrix (eq. (7.567)).

(1) Irrelevance. By Item 1 of Definition 7.466, $\operatorname{Re} y_{\tilde{B}} < 0$ for every excited eigendirection and the combined (estimator+gauge) bands over $\mathcal{I}_\mu^{[\mathcal{U}]}$ do not cross 0. Hence, by the linear Jordan solution and irrelevance criterion (Proposition 7.450, Definition 7.449, and Corollary 7.456), each projected eigen–coupling satisfies

$$\widetilde{\varepsilon}_{\tilde{B}}(\mu) = \left(\tfrac{\mu}{\mu_0}\right)^{y_{\tilde{B}}} \Big[\operatorname{polylog}(\ln \tfrac{\mu}{\mu_0})\Big]\,[1+o(1)] \quad \longrightarrow \quad 0 \qquad (\mu \downarrow 0),$$

since the polynomial in $\ln(\mu/\mu_0)$ cannot overcome the negative real part of $y_{\tilde{B}}$. Because $\mathcal{I}_\mu^{[\mathcal{U}]} = \{\mu \le \mu_R^{[\mathcal{U}]}\}$ is a half–line (Definition 7.461 and Proposition 7.462), this decay holds along the entire IR direction.

(2) Consistency with Step 3 and applicability of Proposition 7.453. By Proposition 7.462, the Step 3 guards (plateaux and monotonicity/curvature, cf. Definition 6.297, Proposition 7.427, and eq. (7.548)) hold on $\mathcal{I}_\mu^{[\mathcal{U}]}$. Consequently the breaking sector is small and decaying there, and the continuum running of each eigen–coupling obeys the asymptotic form used in Proposition 7.453. Item 2 of Definition 7.466 (flow suppression $|\widetilde{\varepsilon}_{\tilde{B}}(\mu)| \le \varepsilon_{\text{plat}}$ for $\mu \le \mu_R^{[\mathcal{U}]}$) is consistent with this decay and guarantees that nonlinear remainders in the RG equation remain subleading on the window (cf. the remainder control in the proofs of Propositions 7.450 and 7.455). Hence the estimator consistency and rate statements of Proposition 7.453 (and, discretely, Proposition 7.465) apply.

(3) Gauge, basis, and scheme robustness. Gauge–parameter variations act by a similarity (Lax) flow on $Y^{\mathcal{N}}$,

$$\partial_\alpha Y^{\mathcal{N}} = [K(\alpha), Y^{\mathcal{N}}], \qquad Y^{\mathcal{N}}(\alpha) = S(\alpha,\alpha_0)\, Y^{\mathcal{N}}(\alpha_0)\, S(\alpha,\alpha_0)^{-1},$$

by Proposition 7.458; hence the spectrum $\{y_{\tilde{B}}\}$ is gauge–independent (Corollary 7.459), and the audit bound (7.588) (= (7.589)) in Definition 7.466 records finite–precision tolerance. Admissible basis changes and fixed–point scheme changes act by similarity on W (Lemma 7.451; Nielsen–projected version in Lemma 7.460), so the multiset of exponents is invariant. Therefore the quoted $\{y_{\tilde{B}}\}$ are robust under the allowed gauge/basis/scheme variations.

Putting (1)–(3) together establishes the claimed soundness and robustness of the Step 4 acceptance. □

Remarks. (i) Item 2 of Definition 7.466 (flow suppression) also ensures numerical stability for the centered estimator Definition 7.446; together with Proposition 7.465 it guarantees that discretisation and mollification errors remain below the plateau tolerances on $\mathcal{I}_\mu^{[\mathcal{U}]}$. (ii) If some $y_{\tilde{B}}$ are nearly degenerate, eigenvectors may rotate along the audit path, but similarity invariance preserves the spectrum and the measured exponents (cf. Corollary 7.459 and Lemma 7.451).

Outputs for Step 4. All deliverables are produced on the on–shell BRST–physical subspace $W := \operatorname{im}\mathcal{N}$ (Nielsen projection; cf. Definition 7.457). First, we present the *projected stability data at the fixed point*: the matrix $Y^{\mathcal{N}}(g_\star)$ at the IR fixed point $g_\star$ (Definition 7.447), its spectrum of projected exponents $\{y_{\tilde{B}}\}$, and the change of basis R that brings $Y^{\mathcal{N}}$ to eigen/Jordan form on W (Definition 7.454); when present, nontrivial Jordan structure is documented via block sizes and nilpotent indices (Definition 7.449).

Next, we report *measured runnings and slope fits*. For each projected mode we plot $\mu \mapsto \widetilde{\varepsilon}_{\tilde{B}}(\mu)$ on the IR window $\mathcal{I}_\mu^{[\mathcal{U}]}$ induced by the Step 3 restoration mask (Definition 7.461), fit slopes using the centered estimator eq. (7.568) (optionally its mollified variant satisfying eqs. (7.537) and (7.538)), and overlay confidence bands consistent with Propositions 7.453 and 7.465.

We then provide a brief *robustness summary*: we record the spread of the extracted $\{y_{\tilde{B}}\}$ across admissible gauges (checked against the audit bound eq. (7.589), equivalent to eq. (7.588) by Corollary 7.459) and across admissible estimator controls (grid step h, smoothing scale τ_μ), citing the rate guarantees of Proposition 7.465.

Finally, we include a *consistency note with the restoration mask and Step 2 slope*: we verify that the Step 4 IR window lies entirely inside the Step 3 certified region (Proposition 7.462) and respects law–channel restoration and slope positivity, i.e. $\varsigma_{\mathrm{law}}^{(\tau)}(\ell) > 0$ together with the monotonicity/curvature guards eqs. (7.542) and (7.548) along the ℓ–preimage $\mathcal{I}_\ell$ (cf. Definitions 6.297 and 7.461 and Proposition 7.427). For reproducibility, we record $(\kappa,\ \mathcal{N},\ Z^{\mathcal{N}},\ Y^{\mathcal{N}})$, the chosen R, the window endpoints $(\ell_R^{[\mathcal{U}]}, \mu_R^{[\mathcal{U}]})$, and the estimator settings (grid h, smoothing τ_μ) alongside code stencils and seeds.

Negative Nielsen–projected exponents $y_{\tilde{B}}$ (eigenvalues of $Y^{\mathcal{N}}$) certify that explicit symmetry breakings are RG–irrelevant (Corollary 7.456), and the estimator plots/bands on $\mathcal{I}_\mu^{[\mathcal{U}]}$ substantiate this with controlled rates (Propositions 7.453 and 7.465). Together with the Step 3 restoration mask and positive IR slope (Proposition 7.427 and eq. (7.542)), this closes the Step 4 ledger and prepares the Ward/Slavnov–Taylor/Nielsen closure in subsequent steps [389, 91, 117, 243].

Conclusion for Step 4. Step 4 established irrelevance of explicit breakings on the *physical* operator space by working after the Nielsen projection $W := \operatorname{im}\mathcal{N}$ (cf. Definition 7.457). On W the projected stability matrix $Y^{\mathcal{N}}$ governs the linearised RG of the breaking sector (cf. eq. (7.567)). Its spectral/Jordan data (Definition 7.449) yield the projected exponents $y_{\tilde{B}}$, and the Jordan-form solution shows that each eigen–coupling runs as a power of μ multiplied by at most a finite polynomial in $\ln\mu$ (Proposition 7.450). The criterion $\operatorname{Re} y_{\tilde{B}} < 0$ therefore guarantees $\widetilde{\varepsilon}_{\tilde{B}}(\mu) \to 0$ toward the IR, i.e. Wilsonian irrelevance (Corollary 7.456).

We then *measured and certified* these exponents on the IR window induced by the Step 3 restoration mask (Definition 7.461 and Proposition 7.462). Using the centered log–derivative estimator (Definition 7.446, eq. eq. (7.568))—optionally mollified under moment constraints—we obtained controlled convergence to $y_{\tilde{B}}$ with explicit rates: $\widehat{y}_{\tilde{B}} = y_{\tilde{B}} + \mathcal{O}(1/|\ln\mu|) + \mathcal{O}(h^2)$ (and $+\mathcal{O}(\tau_\mu^2)$ if smoothed), as formalised in Propositions 7.453 and 7.465. The Step 3 guards ensure the smallness/decay required for these bounds across the entire window (Proposition 7.462).

Robustness was demonstrated along three axes. First, *gauge safety*: by the Nielsen identity

the projected generator $Y^{\mathcal{N}}$ evolves by a similarity (Lax) flow in the gauge parameter, so its spectrum is invariant (Proposition 7.458 and Corollary 7.459), with a practical audit band (7.588). Second, *basis/scheme covariance*: admissible operator relabellings and fixed–point scheme changes act by similarity on W, leaving $\{y_{\tilde{B}}\}$ unchanged (Lemmas 7.451 and 7.460). Third, *flow suppression* on the certified window matches the irrelevance prediction and stabilises estimation (Item 2 in Definition 7.466).

Finally, Step 4 delivered a reproducible audit packet: $Y^{\mathcal{N}}(g_\star)$ and its spectral/Jordan data, estimator plots with confidence bands on $\mathcal{I}_\mu^{[\mathcal{U}]}$, gauge/instrument robustness tables, and a consistency report with the Step 3 mask. With these pieces, the acceptance test of Definition 7.466 implies the soundness/robustness guarantees in Proposition 7.467. In short, Step 4 does not merely assert but *certifies* that the explicit breakings are RG–irrelevant on the physical subspace, closing the QIII ledger and preparing the Ward/Slavnov–Taylor/Nielsen closure in the subsequent steps.

7.8.5 Ward/ST/Nielsen Closure (Band–Projected)

Objective. We certify that global Ward structures are *restored* in the IR while the gauge/BRST sector remains exact on shell. Concretely, we (i) define a band–projected Ward–residual norm $\mathcal{W}(\ell)$ (below), (ii) demand a *strictly negative* logarithmic slope on the Step 3 IR window (cf. eq. (7.542)),

$$\partial_{\ln \ell}\, \mathcal{W}(\ell) \;\le\; -c \;<\; 0 \qquad \text{for all } \ell \in \mathcal{I}_\ell,$$

and (iii) verify restoration at the Step 3 length (cf. eq. (7.551)):

$$\mathcal{W}\big(\ell_R\big) \;\le\; \varepsilon_{\mathrm{plat}}.$$

In parallel, the Slavnov–Taylor (ST) identity holds on shell at all ℓ, and all quoted IR poles/residues have *zero gauge–parameter slope* by the Nielsen identity (cf. eq. (7.581); see also [117, 243, 442]).

Band–projected Ward residual.

Definition 7.468 (Band–limited 1PI functional and Ward residual)**.** Fix the instrumentation used in eq. (7.522): a UV projector $P_x(\lambda_{\mathrm{UV}})$ and an IR kernel $K_{\mathrm{IR}}^{(\ell)}$, applied consistently to fields and sources, and evaluate on shell. Let $\Gamma^{(\lambda_{\mathrm{UV}},\ell)}$ denote the resulting *band–limited* 1PI effective action, and let G be a global symmetry group with generators T^a (representation on fields understood). Write the associated Ward operators $W^a[\Gamma]$ (which vanish identically when the symmetry is exact; see, e.g., [117, 447]).

Define the admitted momentum band and the band–projected Ward operator by

$$\boxed{\begin{aligned} \mathfrak{B}(\lambda_{\mathrm{UV}},\ell) &:= \left\{ p \in \mathbb{R}^d \;\middle|\; \begin{array}{l} P_x(\lambda_{\mathrm{UV}}) \text{ and } K_{\mathrm{IR}}^{(\ell)} \text{ admit external kinematics} \\ \text{with total momentum } p \end{array} \right\}, \\ \mathcal{W}^a(p;\ell) &:= \Big(W^a[\Gamma^{(\lambda_{\mathrm{UV}},\ell)}]\Big)\Big|_p , \end{aligned}} \tag{7.590}$$

where the restriction $|_p$ means evaluation at external kinematics compatible with the band (and on–shell conditions). Let $\mathcal{R}(p;\ell)$ be a *co–channel reference response* (e.g. the appropriate band–limited two–point kernel) such that $\|\mathcal{R}(p;\ell)\|_{\mathrm{op}} > 0$ on $\mathfrak{B}$. The *dimensionless Ward residual* is

$$\boxed{\mathcal{W}(\ell) := \sup_{a} \sup_{p\in\mathfrak{B}(\lambda_{\mathrm{UV}},\ell)} \frac{\left\|\mathcal{W}^{a}(p;\ell)\right\|_{\mathrm{op}}}{\left\|\mathcal{R}(p;\ell)\right\|_{\mathrm{op}}}.} \tag{7.591}$$

Intuition. We quantify the (band–limited) failure of the Ward identities by comparing the Ward operator to a co–channel response, yielding a scale–free number $\mathcal{W}(\ell)$. Restricting to the same band used to define $\Gamma^{(\lambda_{\mathrm{UV}},\ell)}$ keeps the metric consistent with the instrumentation and with the Step 3 IR window; on–shell evaluation ensures gauge/BRST artefacts are excluded (Nielsen and ST protection, cf. eq. (7.581)).

Lemma 7.469 (Well–posedness and invariances for the band–projected residual)**.** *Let* $\mathfrak{B}(\lambda_{\mathrm{UV}},\ell)$ *and* $\mathcal{W}(\ell)$ *be as in eqs.* (7.590) *and* (7.591)*. Assume, for each fixed* $(\lambda_{\mathrm{UV}},\ell)$*:*

(i) $\mathfrak{B}(\lambda_{\mathrm{UV}},\ell)$ *is compact;*

(ii) $p \mapsto \mathcal{W}^{a}(p;\ell)$ *and* $p \mapsto \mathcal{R}(p;\ell)$ *are continuous (as operator–valued maps);*

(iii) $\|\mathcal{R}(p;\ell)\|_{\mathrm{op}} > 0$ *for all* $p \in \mathfrak{B}(\lambda_{\mathrm{UV}},\ell)$*;*

(iv) *the Ward–generator index set is compact: either a* finite *set of generators* $\{T^{a}\}$*, or (equivalently) the supremum over* a *is taken over the unit sphere of a finite–dimensional Lie algebra with a fixed norm.*

Then $\mathcal{W}(\ell)$ *is finite and the supremum is attained.*

Moreover:

(A) *(*Unitary field re–bases*) If the external field multiplet in the audited channel is re–expressed by a unitary change of basis* U *(with respect to the inner product that induces* $\|\cdot\|_{\mathrm{op}}$*), then both* $\mathcal{W}^{a}$ *and* $\mathcal{R}$ *transform by conjugation, the operator norm is unitarily invariant, and* $\mathcal{W}(\ell)$ *is unchanged.*

(B) *(*Uniform leg–rescalings in the channel*) If each external leg in the audited channel is rescaled by a positive scalar* s *(i.e.* $\phi \mapsto s\,\phi$ *for those species only), then* n*–point kernels in that channel acquire the common factor* s^{n}*; hence the numerator and denominator in eq.* (7.591) *are multiplied by the same factor, and* $\mathcal{W}(\ell)$ *is unchanged.*

(C) *(*Reference normalisation*) Replacing* $\mathcal{R}$ *by* $c\,\mathcal{R}$ *with a* constant $c > 0$ *rescales* $\mathcal{W}(\ell)$ *by* $1/c$*. The* log–slope *criterion* $\partial_{\ln\ell}\mathcal{W} \le -c_{0} < 0$ *and any comparison of slopes are invariant under such constant renormalisations. (If* c *depended on* ℓ*, the slope would shift by* $-\partial_{\ln\ell}\ln c(\ell)$*, so admissible choices must be* ℓ*–independent.)*

Proof. **Well–posedness.** Fix $(\lambda_{\mathrm{UV}},\ell)$. For each generator label a, define

$$F_{a}(p;\ell) := \frac{\|\mathcal{W}^{a}(p;\ell)\|_{\mathrm{op}}}{\|\mathcal{R}(p;\ell)\|_{\mathrm{op}}} \qquad (p \in \mathfrak{B}(\lambda_{\mathrm{UV}},\ell)).$$

By (ii)–(iii), both numerator and denominator are continuous in p and the denominator is strictly positive on $\mathfrak{B}(\lambda_{\mathrm{UV}}, \ell)$; hence $F_a(\,\cdot\,;\ell)$ is continuous on the compact set $\mathfrak{B}(\lambda_{\mathrm{UV}}, \ell)$ and attains a finite maximum there (extreme–value theorem).

If the set of a is finite, then $\mathcal{W}(\ell) = \max_a \max_{p\in\mathfrak{B}} F_a(p;\ell)$ is the maximum of finitely many attained maxima and is therefore finite and attained. If instead we take the supremum over the unit sphere in a finite–dimensional Lie algebra (assumption (iv)), the map

$$(p, \hat{a}) \;\longmapsto\; F_{\hat{a}}(p;\ell)$$

is continuous on the compact product $\mathfrak{B}(\lambda_{\mathrm{UV}}, \ell) \times S$ (with S the unit sphere), so the joint maximum exists and is finite; in particular, the supremum over a is attained.

(A) Unitary field re–bases. Let U be unitary with respect to the inner product that induces $\|\cdot\|_{\mathrm{op}}$. In the audited channel, tensors/kernels transform by conjugation (or congruence) with U—the precise side (left/right) depends on slot convention but is identical for numerator and denominator, cf. the "same channel" prescription in Definition 7.468. Because the operator norm is unitarily invariant,

$$\|\mathcal{W}^a\|_{\mathrm{op}} \mapsto \|U^{-1}\mathcal{W}^a U\|_{\mathrm{op}} = \|\mathcal{W}^a\|_{\mathrm{op}}, \qquad \|\mathcal{R}\|_{\mathrm{op}} \mapsto \|U^{-1}\mathcal{R}U\|_{\mathrm{op}} = \|\mathcal{R}\|_{\mathrm{op}},$$

so each ratio F_a and thus $\mathcal{W}(\ell)$ are unchanged.

(B) Uniform leg–rescalings in the channel. Let $\phi \mapsto s\,\phi$ with $s > 0$ for all external legs in the audited channel; then an n–point amputated kernel in that channel scales as s^n. By Definition 7.468, both $\mathcal{W}^a$ and the co–channel reference $\mathcal{R}$ are chosen with the same external–leg structure (same n), hence

$$\|\mathcal{W}^a\|_{\mathrm{op}} \mapsto s^n\|\mathcal{W}^a\|_{\mathrm{op}}, \qquad \|\mathcal{R}\|_{\mathrm{op}} \mapsto s^n\|\mathcal{R}\|_{\mathrm{op}},$$

and F_a (hence $\mathcal{W}$) is invariant.

(C) Reference normalisation. If $\mathcal{R} \mapsto c\,\mathcal{R}$ with a constant $c > 0$, then $F_a \mapsto F_a/c$ and $\mathcal{W}(\ell) \mapsto \mathcal{W}(\ell)/c$. Thus the absolute level depends on the chosen normalisation of the reference kernel, but the *logarithmic slope* $\partial_{\ln\ell} \ln \mathcal{W}$ (and the strict negativity condition of the Step 5 objective) is unaffected by any *constant* choice of c. If c depended on ℓ, the slope would shift by the additive term $-\partial_{\ln\ell} \ln c(\ell)$; hence admissible normalisations must be ℓ–independent for the slope test to be meaningful. In practice we fix $\mathcal{R}$ once and for all on the Step 3 window (cf. Proposition 7.462) to avoid this ambiguity.

These statements use only the band–limited setup and same–channel normalisation in Definition 7.468 and do not rely on additional structure. □

Proposition 7.470 (Vanishing for exact symmetry; leading response to explicit breakings)**.** *Let $\mathcal{W}(\ell)$ be the band–projected Ward residual of Definition 7.468.*

(Exact symmetry). *If the global symmetry generated by T^a is exact for the band–limited 1PI functional $\Gamma^{(\lambda_{\mathrm{UV}},\ell)}$ (on shell), then*

$$\mathcal{W}^a(\cdot;\ell) \equiv 0 \qquad \textit{and hence} \qquad \mathcal{W}(\ell) = 0.$$

(Small explicit breakings). *If explicit breaking couplings $\varepsilon_A(\mu)$ multiply renormalised operators $\hat{\mathcal{O}}_A$ as in eq. (7.522), with $\mu = \kappa/\ell$ (cf. Definition 7.461), then for sufficiently small ε one has the linear expansion (on shell and within the admitted band)*

$$\mathcal{W}^a(p;\ell) = \sum_A \varepsilon_A(\kappa/\ell)\, \mathcal{K}^a{}_A(p;\ell) + \mathcal{O}\Big(\|\varepsilon\|^2\Big). \tag{7.592}$$

for some band–limited kernels $\mathcal{K}^a_A$ depending on the channel and instrumentation $(P_x(\lambda_{\mathrm{UV}}), K^{(\ell)}_{\mathrm{IR}})$. Consequently, there exists a finite constant $C(\lambda_{\mathrm{UV}},\ell) > 0$ (uniform on the band $\mathfrak{B}(\lambda_{\mathrm{UV}},\ell)$) such that

$$\mathcal{W}(\ell) \;\le\; C(\lambda_{\mathrm{UV}},\ell) \sum_{\tilde{B}} \Big|\widetilde{\varepsilon}_{\tilde{B}}(\kappa/\ell)\Big| \;+\; \mathcal{O}(\|\varepsilon\|^2), \tag{7.593}$$

where $\widetilde{\varepsilon}_{\tilde{B}}$ are the Nielsen–projected eigen–couplings (Definitions 7.454 and 7.457).

Proof. **Exact symmetry.** If the symmetry generated by T^a is exact for $\Gamma^{(\lambda_{\mathrm{UV}},\ell)}$ (on shell), then the Ward operator vanishes identically: $W^a[\Gamma^{(\lambda_{\mathrm{UV}},\ell)}] \equiv 0$ (e.g. [117, 447]). Therefore $\mathcal{W}^a(p;\ell) = 0$ for all $p \in \mathfrak{B}(\lambda_{\mathrm{UV}},\ell)$, and by eq. (7.591) the residual $\mathcal{W}(\ell) = 0$.

Linear response to explicit breakings. Consider the band–limited functional with small explicit breakings,

$$\Gamma^{(\lambda_{\mathrm{UV}},\ell)}[\phi;\varepsilon] = \Gamma_0^{(\lambda_{\mathrm{UV}},\ell)}[\phi] + \sum_A \varepsilon_A(\kappa/\ell)\, \mathcal{I}_A^{(\lambda_{\mathrm{UV}},\ell)}[\phi] \;+\; \mathcal{O}(\|\varepsilon\|^2),$$

where $\Gamma_0^{(\lambda_{\mathrm{UV}},\ell)}$ is the symmetric ($\varepsilon = 0$) functional, and $\mathcal{I}_A^{(\lambda_{\mathrm{UV}},\ell)}$ denotes the band–limited, renormalised insertion of $\hat{\mathcal{O}}_A$ (consistent with $P_x(\lambda_{\mathrm{UV}}), K^{(\ell)}_{\mathrm{IR}}$; cf. Definition 7.468). The Ward operator is linear in Γ, so expanding at the symmetric point and evaluating on shell yields the Gateaux derivative

$$W^a[\Gamma^{(\lambda_{\mathrm{UV}},\ell)}] = W^a[\Gamma_0^{(\lambda_{\mathrm{UV}},\ell)}] + \sum_A \varepsilon_A(\kappa/\ell) \Big(\mathrm{D}_{\mathcal{I}_A} W^a\Big)\Big[\Gamma_0^{(\lambda_{\mathrm{UV}},\ell)}\Big] \;+\; \mathcal{O}(\|\varepsilon\|^2).$$

Since the symmetry is exact at $\varepsilon = 0$, the first term vanishes; define the band–limited kernels (on shell)

$$\mathcal{K}^a_A(p;\ell) \;:=\; \Big(\mathrm{D}_{\mathcal{I}_A} W^a\Big)\Big[\Gamma_0^{(\lambda_{\mathrm{UV}},\ell)}\Big]\Big|_p,$$

to obtain eq. (7.592) after restricting to $p \in \mathfrak{B}(\lambda_{\mathrm{UV}},\ell)$ (eq. (7.590)). By the assumptions in Lemma 7.469, the remainder is $\mathcal{O}(\|\varepsilon\|^2)$ uniformly on the compact band.

From linear response to a band–uniform bound. Taking operator norms and using eq. (7.591) plus the triangle inequality,

$$\begin{aligned}
\mathcal{W}(\ell) &= \sup_a \sup_{p\in\mathfrak{B}} \frac{\big\|\sum_A \varepsilon_A\, \mathcal{K}^a_A(p;\ell)\big\|_{\mathrm{op}}}{\|\mathcal{R}(p;\ell)\|_{\mathrm{op}}} + \mathcal{O}(\|\varepsilon\|^2) \\
&\le \sum_A |\varepsilon_A(\kappa/\ell)| \underbrace{\sup_a \sup_{p\in\mathfrak{B}} \frac{\|\mathcal{K}^a_A(p;\ell)\|_{\mathrm{op}}}{\|\mathcal{R}(p;\ell)\|_{\mathrm{op}}}}_{=:C_A(\lambda_{\mathrm{UV}},\ell)} + \mathcal{O}(\|\varepsilon\|^2).
\end{aligned}$$

Set $C(\lambda_{\mathrm{UV}},\ell) := \sum_A C_A(\lambda_{\mathrm{UV}},\ell)$ to get $\mathcal{W}(\ell) \le C(\lambda_{\mathrm{UV}},\ell)\sum_A |\varepsilon_A(\kappa/\ell)| + \mathcal{O}(\|\varepsilon\|^2)$.

Passing to Nielsen–projected eigen–couplings. Let R be the change of basis that (block-)Jordanises the *projected* stability matrix on the physical subspace $W := \operatorname{im}\mathcal{N}$ (cf. Definitions 7.454 and 7.457): $\widetilde{\varepsilon} := R^{-1}\varepsilon$ with components $\widetilde{\varepsilon}_{\tilde{B}}$. In a finite–dimensional space all norms are equivalent, and the linear map R^{-1} is bounded; hence there exists a finite $M_R > 0$ (depending only on R) such that

$$\sum_A |\varepsilon_A| \;\le\; M_R \sum_{\tilde{B}} |\widetilde{\varepsilon}_{\tilde{B}}|.$$

Combining with the previous bound yields eq. (7.593) with the redefined constant $C(\lambda_{\mathrm{UV}},\ell) \leftarrow M_R\, C(\lambda_{\mathrm{UV}},\ell)$. Basis/scheme covariance of this step follows from Lemmas 7.451 and 7.460, and the on–shell construction ensures gauge–parameter safety within the tolerance ledger (eq. (7.581) and Corollary 7.459).

This proves both the vanishing in the symmetric case and the linear–response bound in the explicitly broken case, uniformly on the admitted band $\mathfrak{B}(\lambda_{\mathrm{UV}},\ell)$. □

Comments. (i) The constant $C(\lambda_{\mathrm{UV}},\ell)$ is finite under the compactness/continuity assumptions of Lemma 7.469 and depends only on the instrumentation and the chosen co–channel reference $\mathcal{R}$. (ii) On the Step 4 IR window (Definition 7.461 and Proposition 7.462) the eigen–couplings $\widetilde{\varepsilon}_{\tilde{B}}(\kappa/\ell)$ decay (Step 4: Corollary 7.456), so eq. (7.593) implies $\mathcal{W}(\ell) \downarrow 0$, anticipating the Step 5 restoration test.

Corollary 7.471 (IR decay of the band–projected Ward residual under irrelevance)**.** *Fix the instrumentation* $(P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}^{(\ell)})$ *and work on the IR ray* $\mu = \kappa/\ell \downarrow 0$ *(cf. Definition 7.461). If all* excited *projected breaking exponents satisfy* $\operatorname{Re} y_{\tilde{B}} < 0$ *(as certified in Step 4), then*

$$\mathcal{W}(\ell) \;=\; \mathcal{O}\Big(\sum_{\tilde{B}} \mu^{\operatorname{Re} y_{\tilde{B}}}\, \mathrm{poly}\big(\ln\mu\big)\Big) \;\xrightarrow[\mu\to 0]{}\; 0,$$

with logarithmic factors present only when Jordan blocks occur (cf. Definition 7.449 and Proposition 7.450). Equivalently, letting $y_\star := \max_{\tilde{B}} \operatorname{Re} y_{\tilde{B}} < 0$*, one has* $\mathcal{W}(\ell) = \mathcal{O}\big(\mu^{\operatorname{Re} y_\star}\,\mathrm{poly}(\ln\mu)\big) \to 0$.

Proof. Start from the band–uniform linear–response bound of Proposition 7.470:

$$\mathcal{W}(\ell) \;\le\; C(\lambda_{\mathrm{UV}},\ell) \sum_{\tilde{B}} \big|\widetilde{\varepsilon}_{\tilde{B}}(\mu)\big| \;+\; \mathcal{O}\big(\|\varepsilon(\mu)\|^2\big), \qquad \mu = \kappa/\ell, \tag{7.594}$$

where the constant $C(\lambda_{\mathrm{UV}},\ell)$ depends only on the instrumentation and the co–channel reference (Definition 7.468). By the Step 4 linear solution on the physical subspace (Definition 7.457 and Proposition 7.450), each projected eigen–coupling has the asymptotics

$$\widetilde{\varepsilon}_{\tilde{B}}(\mu) = \mu^{y_{\tilde{B}}}\Big[\mathrm{poly}\big(\ln\mu\big)\Big]\,[1 + o(1)] \qquad (\mu \downarrow 0),$$

with the polynomial factor absent when the corresponding Jordan block is trivial (Definition 7.449). Taking absolute values yields

$$\big|\widetilde{\varepsilon}_{\tilde{B}}(\mu)\big| = \mu^{\operatorname{Re} y_{\tilde{B}}}\,\mathrm{poly}\big(\ln\mu\big)\,[1 + o(1)].$$

Uniformity of the prefactor. Along the Step 3 IR window (half–line in ℓ, cf. Definition 7.461 and Proposition 7.462), the admitted band $\mathfrak{B}(\lambda_{\mathrm{UV}},\ell)$ stays inside a fixed compact region of momentum space and varies continuously with ℓ; $\mathcal{R}$ and the kernels $\mathcal{K}_A^a$ are continuous on band (assumptions of Lemma 7.469). Hence

$$C_\star \;:=\; \sup_{\ell\ge\ell_R}\; C(\lambda_{\mathrm{UV}},\ell) \;<\; \infty,$$

so $C(\lambda_{\mathrm{UV}},\ell)\le C_\star$ for all ℓ in the IR window.[12]

Quadratic remainder. By norm equivalence in finite dimension and the change of basis to the eigen–coordinates (Definition 7.454), there exists $M>0$ with $\|\varepsilon(\mu)\|\le M\sum_{\tilde B}|\widetilde{\varepsilon}_{\tilde B}(\mu)|$. Thus the remainder in (7.594) satisfies

$$\mathcal{O}\Big(\|\varepsilon(\mu)\|^2\Big)=\mathcal{O}\Big(\Big[\sum_{\tilde B}|\widetilde{\varepsilon}_{\tilde B}(\mu)|\Big]^2\Big)=\mathcal{O}\Big(\sum_{\tilde B}\mu^{\,2\,\mathrm{Re}\,y_{\tilde B}}\,\mathrm{poly}\Big(\ln\mu\Big)\Big),$$

which is $o\Big(\sum_{\tilde B}\mu^{\,\mathrm{Re}\,y_{\tilde B}}\,\mathrm{poly}(\ln\mu)\Big)$ as $\mu\downarrow 0$ because all $\mathrm{Re}\,y_{\tilde B}<0$.

Putting these estimates into (7.594) gives

$$\mathcal{W}(\ell)\;\le\; C_\star\sum_{\tilde B}\mu^{\,\mathrm{Re}\,y_{\tilde B}}\,\mathrm{poly}\Big(\ln\mu\Big)\,[1+o(1)],$$

which proves the claimed big–$\mathcal{O}$ bound and the limit $\mathcal{W}(\ell)\to 0$ as $\mu\downarrow 0$. The alternative form with $y_\star=\max_{\tilde B}\mathrm{Re}\,y_{\tilde B}$ follows by bounding the sum by a constant multiple of the dominant term. □

Definition 7.472 (UV–uniform Ward residual)**.** Let $\mathcal{U}=[u_{\min},u_{\max}]$ be an admissible UV strip (cf. Definition 7.430) and write $u=\ln\lambda_{\mathrm{UV}}$. For fixed ℓ, define

$$\mathcal{W}^{[\mathcal{U}]}(\ell):=\sup_{u\in\mathcal{U}}\;\sup_{p\in\mathfrak{B}(\mathrm{e}^u,\ell)}\;\frac{\big\|\mathcal{W}^a(p;\ell;u)\big\|_{\mathrm{op}}}{\big\|\mathcal{R}(p;\ell;u)\big\|_{\mathrm{op}}}\qquad(\text{sup over } a \text{ understood}),$$

with the dependence on u made explicit through the band $\mathfrak{B}(\mathrm{e}^u,\ell)$ and the band–limited objects $\mathcal{W}^a(\cdot;\ell;u)$, $\mathcal{R}(\cdot;\ell;u)$ (cf. Definition 7.468). Then, by definition,

$$\mathcal{W}^{[\mathcal{U}]}(\ell)=\sup_{u\in\mathcal{U}}\mathcal{W}(\ell;u),$$

and, under the assumptions of Lemma 7.469 uniformly in $u\in\mathcal{U}$, $\mathcal{W}^{[\mathcal{U}]}(\ell)$ is finite and inherits the invariances stated there.

Lemma 7.473 (Gauge–parameter safety via on–shell Nielsen projection)**.** *Assume gauge fixing and evaluate correlators* on shell. *Replace all insertions by their Nielsen–projected counterparts on $W=\mathrm{im}\,\mathcal{N}$ (cf. Definition 7.457). Then the band–projected residuals computed from the projected data $(Y^{\mathcal{N}},\Gamma^{(\lambda_{\mathrm{UV}},\ell)}|_W)$ are independent of the gauge parameter within the tolerance eq. (7.588) on any Step 3 IR window (cf. Definition 7.461 and Proposition 7.462).*

[12] If one does not assume global uniformity, the argument below still goes through whenever $C(\lambda_{\mathrm{UV}},\ell)$ has at most polylogarithmic growth in ℓ; the decay from $\mu^{\mathrm{Re}\,y_{\tilde B}}$ dominates. In our instrumentation, boundedness holds.

Proof. **1) On–shell gauge flow (Nielsen).** By the Nielsen identity (eq. (7.581)) and projection to the physical subspace $W = \operatorname{im}\mathcal{N}$ (Definition 7.457), gauge–parameter variation acts by an adjoint flow on composite operator multiplets: for a smooth gauge parameter ξ,

$$\partial_\xi X^{\mathcal{N}}(\xi) = \big[K(\xi),\, X^{\mathcal{N}}(\xi)\big] \quad \text{on } W,$$

for any renormalised insertion X (Ward operator or co–channel reference), with a bounded generator $K(\xi)$ (Proposition 7.458). The solution is the path–ordered similarity $X^{\mathcal{N}}(\xi) = S(\xi,\xi_0)^{-1} X^{\mathcal{N}}(\xi_0) S(\xi,\xi_0)$.

2) Effect on the residual at fixed (u,ℓ,p). Fix (u,ℓ) and an admissible $p \in \mathfrak{B}(\mathrm{e}^u,\ell)$. Denote $N(\xi) := \mathcal{W}^a(p;\ell;u,\xi)^{\mathcal{N}}$ and $D(\xi) := \mathcal{R}(p;\ell;u,\xi)^{\mathcal{N}}$. Then $N(\xi) = S^{-1}N_0 S$ and $D(\xi) = S^{-1}D_0 S$, where the subscript 0 refers to ξ_0. Consider the *dimensionless* map

$$\Phi(\xi) \;:=\; D(\xi)^{-1} N(\xi).$$

Differentiating and using $\partial_\xi D = [K,D]$, $\partial_\xi N = [K,N]$ gives

$$\partial_\xi \Phi = [K,\Phi], \qquad \Phi(\xi) = S(\xi,\xi_0)^{-1}\,\Phi(\xi_0)\,S(\xi,\xi_0).$$

Hence Φ is conjugate along the gauge flow.

3) Norm control and tolerance. Let $\|\cdot\|_{\mathrm{op}}$ be any operator norm subordinate to a fixed (gauge–independent) inner product on the co–channel space. Then

$$\frac{\mathrm{d}}{\mathrm{d}\xi}\,\|\Phi(\xi)\|_{\mathrm{op}} \;=\; \lim_{h\to 0}\frac{\|(\mathrm{I}+h\,\mathrm{ad}_K)\Phi\|_{\mathrm{op}} - \|\Phi\|_{\mathrm{op}}}{h} \;\le\; 2\,\|K(\xi)\|_{\mathrm{op}}\,\|\Phi(\xi)\|_{\mathrm{op}},$$

by submultiplicativity and $\|\,[K,\Phi]\,\| \le \|K\|\,\|\Phi\| + \|\Phi\|\,\|K\|$. Grönwall then yields

$$\|\Phi(\xi)\|_{\mathrm{op}} \;\le\; \exp\!\Big(2\int_{\xi_0}^{\xi}\!\|K(\zeta)\|_{\mathrm{op}}\mathrm{d}\zeta\Big)\,\|\Phi(\xi_0)\|_{\mathrm{op}}.$$

Taking the supremum over $p \in \mathfrak{B}(\mathrm{e}^u,\ell)$ and a, and then over $u \in \mathcal{U}$, one obtains the same bound for $\mathcal{W}(\ell;u,\xi)$ and $\mathcal{W}^{[\mathcal{U}]}(\ell;\xi)$. On the admissible gauge set Ξ_{adm}, the integral of $\|K\|$ is finite; Step 5's gauge audit demands that the resulting variation stays below the tolerance:

$$\sup_{\xi\in\Xi_{\mathrm{adm}}}\Big|\mathcal{W}^{[\mathcal{U}]}(\ell;\xi) - \mathcal{W}^{[\mathcal{U}]}(\ell;\xi_0)\Big| \;\le\; \varepsilon_{\mathrm{gauge}},$$

in the same spirit as eq. (7.588). In particular, if the subordinate norm is chosen from a co–channel Gram metric that is itself protected on shell (via ST/Nielsen), the generator K is skew with respect to that metric and the exponential factor is 1, giving exact invariance; otherwise, invariance holds up to the stated tolerance.

4) Conclusion. Because the ratio that defines $\mathcal{W}$ can be written as $\|\Phi(\xi)\|_{\mathrm{op}}$ and Φ evolves by conjugation, the band–projected residual is independent of the gauge parameter within the audit tolerance on the IR window. This extends to the UV–uniform version $\mathcal{W}^{[\mathcal{U}]}$ by the supremum construction in Definition 7.472. □

Notes.

- In Definition 7.472 we stated equality $\mathcal{W}^{[\mathcal{U}]}(\ell) = \sup_{u\in\mathcal{U}} \mathcal{W}(\ell;u)$ (not just $\geq$), since $\mathcal{W}(\ell;u)$ is itself the supremum over p and a for fixed u.
- Uniform well–posedness of $\mathcal{W}^{[\mathcal{U}]}$ follows by applying Lemma 7.469 uniformly in $u\in\mathcal{U}$ (compactness of $\mathcal{U}$, continuity in u and p, and a uniform lower bound on $|\mathcal{R}|_{\mathrm{op}}$ over the union band).

Proposition 7.474 (Monotone trend under the Step 3 slope guard)**.** *Assume the Step 3 guard* $\varsigma^{(\tau)}_{\mathrm{law}}(\ell) > 0$ *and the curvature/TV constraints of eq. (7.548) on an IR window* $\ell \geq \ell_R$ *(cf. eq. (7.551)). Then there exist* $\ell_\star \geq \ell_R$ *and* $c > 0$ *such that for all* $\ell \geq \ell_\star$*,*

$$\frac{\mathrm{d}}{\mathrm{d}\ell}\mathcal{W}(\ell) \;\leq\; -\frac{c}{\ell}\mathcal{W}(\ell) \;+\; \mathcal{O}\Big(\|\varepsilon(\kappa/\ell)\|^2\Big), \tag{7.595}$$

and hence $\mathcal{W}(\ell)$ *is eventually decreasing (up to quadratic remainders) along the IR ray* $\mu = \kappa/\ell \downarrow 0$*.*

Proof. **Setup and uniform controls.** Fix the instrumentation $(P_x(\lambda_{\mathrm{UV}}), K^{(\ell)}_{\mathrm{IR}})$ and work on the Step 3 IR window $\ell \geq \ell_R$ (cf. Definition 7.461 and Proposition 7.462). By Proposition 7.470, the band–projected Ward residual obeys the band–uniform bound

$$\mathcal{W}(\ell) \;\leq\; C(\lambda_{\mathrm{UV}},\ell)\sum_{\tilde{B}}\big|\widetilde{\varepsilon}_{\tilde{B}}(\mu)\big| \;+\; \mathcal{O}\Big(\|\varepsilon(\mu)\|^2\Big), \qquad \mu = \kappa/\ell, \tag{7.596}$$

with $C(\lambda_{\mathrm{UV}},\ell)$ finite by Lemma 7.469. The curvature/TV guard eq. (7.548) and the smooth dependence of the band/data on ℓ (Definition 7.468) imply that $C(\lambda_{\mathrm{UV}},\ell)$ is bounded and Lipschitz in $t = \ln\ell$ on compact subwindows: there exists $C_\star, L_\star > 0$ with

$$0 < C(\lambda_{\mathrm{UV}},\ell) \leq C_\star, \qquad \left|\frac{\mathrm{d}}{\mathrm{d}\ln\ell}C(\lambda_{\mathrm{UV}},\ell)\right| \leq L_\star \quad (\ell \geq \ell_R). \tag{7.597}$$

Slope guard for the breaking sector. Let $t := \ln\mu = -\ln\ell$. Inside the Step 3 certified region the law–channel guard $\varsigma^{(\tau)}_{\mathrm{law}}(\ell) > 0$ together with Proposition 7.462 ensures that the breaking sector is small and decaying, and the local (logarithmic) slopes of the projected eigen–couplings are strictly negative beyond some $\ell_\star \geq \ell_R$: there exists $c_0 > 0$ such that for all $\ell \geq \ell_\star$,

$$\frac{\mathrm{d}}{\mathrm{d}t}\ln\big|\widetilde{\varepsilon}_{\tilde{B}}(\mu(t))\big| \;\leq\; -c_0 \qquad \text{for every excited mode } \tilde{B}, \tag{7.598}$$

up to errors of order $\mathcal{O}(\|\widetilde{\varepsilon}\|)$ that can be absorbed into c_0 for sufficiently small breakings (see Propositions 7.453 and 7.465). Since $\frac{\mathrm{d}}{\mathrm{d}\ell} = -\frac{1}{\ell}\frac{\mathrm{d}}{\mathrm{d}t}$, (7.598) gives

$$\frac{\mathrm{d}}{\mathrm{d}\ell}\big|\widetilde{\varepsilon}_{\tilde{B}}(\mu)\big| \;\leq\; -\frac{c_0}{\ell}\big|\widetilde{\varepsilon}_{\tilde{B}}(\mu)\big| \;+\; \mathcal{O}\Big(\frac{1}{\ell}\,\|\widetilde{\varepsilon}(\mu)\|^2\Big). \tag{7.599}$$

From eigen–couplings to $\mathcal{W}$. Differentiate (7.596) w.r.t. ℓ and use (7.597)–(7.599):

$$\begin{aligned}\frac{\mathrm{d}}{\mathrm{d}\ell}\mathcal{W}(\ell) &\leq \frac{\mathrm{d}C}{\mathrm{d}\ell}\sum_{\tilde{B}}|\widetilde{\varepsilon}_{\tilde{B}}| \;+\; C(\lambda_{\mathrm{UV}},\ell)\sum_{\tilde{B}}\frac{\mathrm{d}}{\mathrm{d}\ell}|\widetilde{\varepsilon}_{\tilde{B}}| \;+\; \mathcal{O}\Big(\|\varepsilon\|\,\|\varepsilon\|'_\ell\Big)\\ &\leq \frac{L_\star}{\ell}\sum_{\tilde{B}}|\widetilde{\varepsilon}_{\tilde{B}}| \;-\; \frac{c_0}{\ell}\,C(\lambda_{\mathrm{UV}},\ell)\sum_{\tilde{B}}|\widetilde{\varepsilon}_{\tilde{B}}| \;+\; \mathcal{O}\Big(\frac{1}{\ell}\,\|\widetilde{\varepsilon}\|^2\Big) \;+\; \mathcal{O}\Big(\|\varepsilon\|^2\Big).\end{aligned}$$

Using $\sum_{\tilde{B}} |\widetilde{\varepsilon}_{\tilde{B}}| \;\geq\; \frac{1}{C_\star}\,\mathcal{W}(\ell) \;-\; \mathcal{O}(\|\varepsilon\|^2)$ (which follows from (7.596) and $C \leq C_\star$) and absorbing the $\mathcal{O}(\|\varepsilon\|^2)$ pieces, we obtain

$$\frac{\mathrm{d}}{\mathrm{d}\ell}\,\mathcal{W}(\ell) \;\leq\; -\,\frac{1}{\ell}\left(\frac{c_0\,C(\lambda_{\mathrm{UV}},\ell)}{C_\star} - \frac{L_\star}{C_\star}\right)\mathcal{W}(\ell) \;+\; \mathcal{O}\Big(\|\varepsilon\|^2\Big).$$

Define $c := \frac{c_0\,\underline{C} - L_\star}{C_\star}$, where $\underline{C} := \inf_{\ell\geq\ell_\star} C(\lambda_{\mathrm{UV}},\ell) > 0$; for sufficiently large $\ell_\star$ (still within the Step 3 window) we may assume $c > 0$, yielding (7.595).

Eventual monotonicity. The differential inequality (7.595) implies that, for $\ell \geq \ell_\star$,

$$\frac{\mathrm{d}}{\mathrm{d}\ell}\Big(\ell^{\,c}\,\mathcal{W}(\ell)\Big) \;\leq\; \ell^{\,c}\,\mathcal{O}\Big(\|\varepsilon\|^2\Big),$$

so $\ell^{\,c}\mathcal{W}(\ell)$ is eventually nonincreasing up to a small quadratic correction; in particular $\mathcal{W}(\ell)$ is eventually decreasing along the IR ray once the breaking sector is sufficiently small (Proposition 7.462). □

Remarks.

- The slope input (7.598) is the continuum version of the estimator statement in Propositions 7.453 and 7.465, guaranteed on the Step 3 window by the law–channel guard $\varsigma^{(\tau)}_{\mathrm{law}}(\ell) > 0$ and the monotonicity/curvature controls eq. (7.548).
- The instrumentation factor $C(\lambda_{\mathrm{UV}},\ell)$ is band–uniformly bounded and slowly varying by Lemma 7.469 (uniform in ℓ on the window) and the smooth dependence of the band–limited kernels on ℓ (Definition 7.468); the curvature/TV guard ensures the Lipschitz control in (7.597).
- If desired, the inequality can be written in $t = \ln\ell$ form as $\frac{\mathrm{d}}{\mathrm{d}t}\ln\mathcal{W} \leq -c + \mathcal{O}(|\varepsilon|^2)$, equivalent to (7.595).

ST identity on shell (all scales).

Definition 7.475 (Band–limiting map and BRST covariance)**.** Let $\mathscr{B}_{\lambda_{\mathrm{UV}},\ell}$ denote the linear band–limiting map that implements the fixed instrumentation of eq. (7.522) on fields, sources, antifields, and composite insertions:

$$\Gamma^{(\lambda_{\mathrm{UV}},\ell)} \;:=\; \mathscr{B}_{\lambda_{\mathrm{UV}},\ell}\,[\,\Gamma\,], \qquad \mathscr{B}_{\lambda_{\mathrm{UV}},\ell} := \Big(P_x(\lambda_{\mathrm{UV}}), K^{(\ell)}_{\mathrm{IR}}\Big) \;\text{ acting consistently on all entries.}$$

Assume $\mathscr{B}_{\lambda_{\mathrm{UV}},\ell}$ is linear, continuous, commutes with ordinary functional derivatives, and is BRST–covariant in the sense that for the BRST differential ζ (a graded derivation) one has

$$\zeta\,\mathscr{B}_{\lambda_{\mathrm{UV}},\ell} \;=\; \mathscr{B}_{\lambda_{\mathrm{UV}},\ell}\,\zeta. \tag{7.600}$$

Equivalently, for the Slavnov–Taylor functional $\mathcal{S}(\cdot)$ one has

$$\mathcal{S}\Big(\mathscr{B}_{\lambda_{\mathrm{UV}},\ell}[X]\Big) = \mathscr{B}_{\lambda_{\mathrm{UV}},\ell}\Big[\mathcal{S}(X)\Big] \;+\; (\text{EOM terms}),$$

where the "EOM terms" are proportional to Euler–Lagrange derivatives and vanish on shell.[13]

Proposition 7.476 (On–shell Slavnov–Taylor identity at all bands)**.** *Assume:*

(i) anomaly freedom for the gauge symmetry ($H^{1,d}(\zeta\,|\,d) = 0$ at ghost number 1);

(ii) the gauge fixing is BRST–exact;

(iii) the explicit breaking operators selected in Step 1 are BRST–closed (ghost number 0) and lie in the physical cohomology after Nielsen projection (Definition 7.457);

(iv) the band–limiting map $\mathscr{B}_{\lambda_{\mathrm{UV}},\ell}$ satisfies BRST covariance eq. (7.600) (Definition 7.475).

Then for every $(\lambda_{\mathrm{UV}}, \ell)$ one has the on–shell ST identity

$$\boxed{\mathcal{S}\Big(\Gamma^{(\lambda_{\mathrm{UV}},\ell)}\Big) \;=\; 0 \qquad \text{whenever} \qquad \frac{\delta\Gamma^{(\lambda_{\mathrm{UV}},\ell)}}{\delta\phi} \;=\; 0\,.} \tag{7.601}$$

Consequently, any nonzero Ward residual $\mathcal{W}(\ell)$ of eq. (7.591) reflects controlled global *symmetry breaking only (no gauge anomaly) within the band–limited description.*

Proof. **Unbanded ST identity (on shell).** By (i) and algebraic renormalisation (Quantum Action Principle), the renormalised Γ obeys $\mathcal{S}(\Gamma) = 0$ modulo EOM–exact terms to all orders [117, 243, 442]; hence exactly on shell: $\mathcal{S}(\Gamma) = 0$ when $\delta\Gamma/\delta\phi = 0$. Assumption (ii) keeps BRST invariance intact, and (iii) ensures the Step 1 breaking insertions are BRST–closed in the physical cohomology, so they do not generate gauge–sector breaking.

Push through the band–limiting map. With $\Gamma^{(\lambda_{\mathrm{UV}},\ell)} = \mathscr{B}_{\lambda_{\mathrm{UV}},\ell}[\Gamma]$ and eq. (7.600),

$$\mathcal{S}\Big(\Gamma^{(\lambda_{\mathrm{UV}},\ell)}\Big) = \mathcal{S}\Big(\mathscr{B}_{\lambda_{\mathrm{UV}},\ell}[\Gamma]\Big) = \mathscr{B}_{\lambda_{\mathrm{UV}},\ell}\Big[\mathcal{S}(\Gamma)\Big] \;+\; (\text{EOM terms}).$$

The first term vanishes on the unbanded shell. Since $\mathscr{B}$ commutes with functional derivatives (Definition 7.475), $\mathscr{B}_{\lambda_{\mathrm{UV}},\ell}\Big[\delta\Gamma/\delta\phi\Big] = \delta\Gamma^{(\lambda_{\mathrm{UV}},\ell)}/\delta\phi$, so the EOM remainder vanishes precisely on the *band–limited* shell, yielding eq. (7.601).

Implication for $\mathcal{W}$. Because eq. (7.601) holds at all bands, any nonzero $\mathcal{W}(\ell)$ (Definition 7.468) originates from global Ward breaking in the audited channel, not from a gauge anomaly. This is consistent with the Step 4 analysis of projected eigen–couplings and irrelevance (Proposition 7.450 and Corollary 7.456). □

Corollary 7.477 (Gauge safety of band–projected Ward diagnostics)**.** *Evaluate correlators on shell and restrict all operator insertions to the Nielsen–projected physical subspace $W = \operatorname{im}\mathcal{N}$ (cf. Definition 7.457). Then the band–projected Ward residual $\mathcal{W}(\ell)$ of Definition 7.468 is insensitive to gauge–parameter choices up to the tolerance eq. (7.588); in particular,*

$$\sup_{\xi\in\mathcal{X}_{\mathrm{adm}}} \Big|\mathcal{W}(\ell;\xi) - \mathcal{W}(\ell;\xi_0)\Big| \;\le\; \varepsilon_{\mathrm{gauge}},$$

where $\mathcal{X}_{\mathrm{adm}}$ denotes the admissible set of gauge parameters used in the gauge audit. Moreover, any nonzero signal $\mathcal{W}(\ell) > 0$ cannot be attributed to a gauge anomaly.

[13] This follows from linearity of $\mathscr{B}$ and its commutation with functional derivatives; in the antifield/BRST setup the possible departures are EOM–exact by the Quantum Action Principle and algebraic renormalisation [117, 243]. Moreover, since $\mathscr{B}$ acts slotwise on sources/antifields and preserves ghost number, it is compatible with the BRST grading.

Proof. **(A) Gauge–parameter insensitivity on W.** By the Nielsen identity eq. (7.581) and the projection to the on–shell BRST–physical subspace (Definition 7.457), the dependence on a gauge parameter ξ acts on W by conjugation (Lax form) with a smooth generator $K(\xi)$; see Proposition 7.458. For fixed $(\lambda_{\mathrm{UV}}, \ell)$ and admissible external $p \in \mathfrak{B}(\lambda_{\mathrm{UV}}, \ell)$, write

$$N(\xi) := \Big(\mathcal{W}^a(p;\ell)\Big)^{\mathcal{N}}, \qquad D(\xi) := \Big(\mathcal{R}(p;\ell)\Big)^{\mathcal{N}},$$

and consider the dimensionless co–channel ratio $\Phi(\xi) := D(\xi)^{-1}N(\xi)$. By the Nielsen/Lax flow on W, $\partial_\xi \Phi = [K(\xi), \Phi(\xi)]$, hence $\Phi(\xi) = S(\xi,\xi_0)^{-1}\Phi(\xi_0)S(\xi,\xi_0)$ with $S = \mathcal{P}\exp\int K$ (cf. Proposition 7.458). Subordinate operator norms are stable under conjugation up to the tolerance used in the gauge audit eq. (7.588); taking the supremum over a and $p \in \mathfrak{B}$ (and, if used, over u in a UV strip as in Definition 7.472) preserves this bound. This gives the stated gauge–parameter insensitivity for $\mathcal{W}(\ell)$; compare also Lemma 7.473.

(B) Excluding gauge anomalies. By Proposition 7.476 the Slavnov–Taylor identity holds *on shell* at every band $(\lambda_{\mathrm{UV}}, \ell)$, provided the anomaly cohomology is trivial and the band–limiting map is BRST–covariant (Definition 7.475). Therefore gauge symmetry is exact on shell at all scales in the band–limited description, and any nonzero Ward residual $\mathcal{W}(\ell)$ computed as in Definition 7.468 cannot originate from a gauge anomaly. Combining this with the linear–response structure of the Ward operator in the presence of explicit *global* breakings (Proposition 7.470) shows that $\mathcal{W}(\ell) > 0$ reflects only controlled global Ward breaking in the audited channel.

The two claims together establish the corollary. □

IR decay and plateau tolerance for Ward residuals.

Definition 7.478 (IR decay and plateau acceptance for $\mathcal{W}$)**.** Let $\mathcal{I}_{\mathrm{IR}} \subset \mathbb{R}$ be the Step 3 IR window in $v := \ln \ell$ where the guards of eqs. (7.542) and (7.548) hold, and let ℓ_R be the (uniform) restoration length (cf. Definitions 7.430 and 7.461 and eq. (7.551)). We *accept* the Ward–residual trend if there exists a constant $c > 0$ such that

$$\boxed{\frac{\partial\, \mathcal{W}(\ell)}{\partial \ln \ell} \;\le\; -c \;<\; 0 \quad \text{for all} \quad \ell \text{ with } \ln \ell \in \mathcal{I}_{\mathrm{IR}}, \qquad \mathcal{W}(\ell_R) \;\le\; \varepsilon_{\mathrm{plat}}.} \tag{7.602}$$

The constant c is chosen commensurate with the curvature/TV guards of eq. (7.548) so that transient ripples cannot fake monotone decay.

Operational note. In practice we estimate the derivative in eq. (7.602) by a centered finite difference in $v = \ln \ell$, optionally preceded by a low–pass in v satisfying eqs. (7.537) and (7.538); acceptance requires the estimated slope to lie below $-c$ across the window, together with the plateau bound at ℓ_R.

Lemma 7.479 (Log–slope decay implied by irrelevance)**.** *Assume fixed instrumentation $(P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}^{(\ell)})$ and the linear–response bound Proposition 7.470 on a Step 3 IR window (cf. Definition 7.461 and Proposition 7.462):*

$$\mathcal{W}(\ell) \;\le\; C(\lambda_{\mathrm{UV}}, \ell) \sum_{\tilde{B}} \big|\widetilde{\varepsilon}_{\tilde{B}}(\kappa/\ell)\big| \;+\; \mathcal{O}\Big(\|\varepsilon\|^2\Big).$$

If all projected breaking exponents satisfy $\operatorname{Re} y_{\tilde{B}} < 0$*, then on any compact subwindow of the Step 3 mask*

$$\frac{\partial}{\partial \ln \ell}\,\mathcal{W}(\ell) \;\le\; \Big(\frac{\partial}{\partial \ln \ell}\ln C(\lambda_{\mathrm{UV}},\ell)\Big)\,\mathcal{W}(\ell) - \gamma_\star\,\mathcal{W}(\ell) + \mathcal{O}\big(\|\varepsilon\|^2\big), \qquad \gamma_\star := \min_{\tilde{B}} \big|\operatorname{Re} y_{\tilde{B}}\big| > 0. \tag{7.603}$$

In particular, if C *is band–uniformly slowly varying so that* $\sup_{\ln\ell\in\mathcal{I}_{\mathrm{IR}}} \big|\partial_{\ln\ell}\ln C\big| \le \eta < \gamma_\star$*, then*

$$\frac{\partial}{\partial \ln \ell}\,\mathcal{W}(\ell) \;\le\; -\,(\gamma_\star - \eta)\,\mathcal{W}(\ell) \;+\; \mathcal{O}\big(\|\varepsilon\|^2\big),$$

so $\mathcal{W}$ *has a strictly negative log–slope up to quadratic remainders.*

Proof. Write $\mu = \kappa/\ell$ so that $\partial_{\ln\ell} = -\,\partial_{\ln\mu}$. Let $S(\ell) := \sum_{\tilde{B}} \big|\widetilde{\varepsilon}_{\tilde{B}}(\mu)\big|$. From Proposition 7.470,

$$\mathcal{W}(\ell) \;\le\; C(\lambda_{\mathrm{UV}},\ell)\,S(\ell) \;+\; \mathcal{O}\big(\|\varepsilon\|^2\big). \tag{7.604}$$

Differentiate (7.604) w.r.t. $\ln\ell$:

$$\frac{\partial \mathcal{W}}{\partial \ln \ell} \;\le\; \Big(\frac{\partial \ln C}{\partial \ln \ell}\Big)\,\mathcal{W} \;+\; C(\lambda_{\mathrm{UV}},\ell)\,\frac{\partial S}{\partial \ln \ell} \;+\; \mathcal{O}\big(\|\varepsilon\|\,\partial_{\ln\ell}\|\varepsilon\|\big).$$

In the projected eigen–basis (Definitions 7.454 and 7.457), the linearised flow (Propositions 7.450 and 7.455) gives

$$\frac{\partial}{\partial \ln \mu}\,|\widetilde{\varepsilon}_{\tilde{B}}| = \operatorname{Re} y_{\tilde{B}}\,|\widetilde{\varepsilon}_{\tilde{B}}| \;+\; \mathcal{O}\big(\|\widetilde{\varepsilon}\|^2\big),$$

hence

$$\frac{\partial S}{\partial \ln \ell} = -\sum_{\tilde{B}} \operatorname{Re} y_{\tilde{B}}\,|\widetilde{\varepsilon}_{\tilde{B}}| \;+\; \mathcal{O}\big(\|\widetilde{\varepsilon}\|^2\big) \;\le\; -\,\gamma_\star\,S(\ell) \;+\; \mathcal{O}\big(\|\widetilde{\varepsilon}\|^2\big).$$

Plugging back and using (7.604) to replace $S(\ell)$ by $\mathcal{W}(\ell)/C(\lambda_{\mathrm{UV}},\ell)$ yields (7.603). If $\big|\partial_{\ln\ell}\ln C\big| \le \eta < \gamma_\star$ on the window, the strict negativity follows. □

Proposition 7.480 (Sufficient conditions for the acceptance inequality)**.** *Let the assumptions of Lemma 7.479 hold on a compact IR window* $\mathcal{I}_{\mathrm{IR}} = [v_1, v_2] \subset \{\ln\ell \ge \ln\ell_R\}$*, and suppose* $\mathcal{W}$ *is continuous with a positive lower bound on that window:* $\inf_{v\in\mathcal{I}_{\mathrm{IR}}} \mathcal{W}(e^v) \ge W_{\min} > 0$*. Then the acceptance condition eq. (7.602) holds with any*

$$0 \;<\; c \;\le\; \gamma_\star\,W_{\min}\,.$$

Proof. From Lemma 7.479, $\frac{\partial}{\partial\ln\ell}\mathcal{W} \le -\gamma_\star\mathcal{W} + \mathcal{O}(\|\varepsilon\|^2)$. Inside the Step 3 mask the quadratic remainder is dominated by the linear term; in particular, shrinking the window if necessary we may absorb it into the coefficient to get $\frac{\partial}{\partial\ln\ell}\mathcal{W} \le -\frac{\gamma_\star}{2}\mathcal{W}$. By the positivity lower bound, $-\frac{\gamma_\star}{2}\mathcal{W} \le -\frac{\gamma_\star}{2}W_{\min}$, which implies eq. (7.602) with $c \le \frac{\gamma_\star}{2}W_{\min}$. The plateau condition is inherited from Step 3, which enforces $\mathcal{W}(\ell_R) \le \varepsilon_{\mathrm{plat}}$ by construction of the mask and the choice of reference channel. □

Remark 7.481 (Robust (scale–free) alternative)**.** The proportional inequality of Lemma 7.479 yields an equivalent, scale–free acceptance in terms of the *log–slope*:

$$\frac{\partial}{\partial \ln \ell}\,\ln\Big(\mathcal{W}(\ell)\Big) \;\le\; -\gamma \;<\; 0 \quad \text{on } \mathcal{I}_{\mathrm{IR}},$$

which implies exponential decay of $\mathcal{W}$ in $\ln \ell$. This weaker (and more natural) requirement is automatically satisfied under Step 4 irrelevance; the stricter linear floor eq. (7.602) may be adopted as a conservative operational check on finite windows by estimating a uniform slope floor via the robust trend statistics used in Step 3.

Nielsen protection: zero gauge–parameter slope (poles/residues).

Definition 7.482 (Band–limited physical two–point kernel, poles and residues)**.** Fix $(\lambda_{\mathrm{UV}}, \ell)$ and a gauge parameter $\xi \in \Xi_{\mathrm{adm}}$. Let $\Gamma^{(\lambda_{\mathrm{UV}},\ell)}$ be the band–limited 1PI functional (cf. Definition 7.468) and define the *Nielsen–projected* 1PI two–point kernel on the physical subspace $W = \operatorname{im}\mathcal{N}(\xi)$ (evaluated on shell; cf. Definition 7.457) by

$$\Gamma_2^{\mathcal{N}}(p^2;\ell,\xi) \;:=\; \mathcal{N}(\xi)\,\Gamma_2^{(\lambda_{\mathrm{UV}},\ell)}(p^2)\,\mathcal{N}(\xi)\Big|_W.$$

A *physical pole* in channel α is a real $m_\alpha^2(\ell;\xi)$ with

$$\det \Gamma_2^{\mathcal{N}}(p^2;\ell,\xi)\Big|_{p^2=m_\alpha^2} = 0,$$

assumed simple ($\partial_{p^2} \det \Gamma_2^{\mathcal{N}} \neq 0$ at $p^2 = m_\alpha^2$). The corresponding residue $Z_\alpha(\ell;\xi) > 0$ is defined by the Laurent expansion of the projected propagator $D^{\mathcal{N}}(p^2;\ell,\xi) := \Big(\Gamma_2^{\mathcal{N}}(p^2;\ell,\xi)\Big)^{-1}$:

$$D^{\mathcal{N}}(p^2;\ell,\xi) = \frac{Z_\alpha(\ell;\xi)\,P_\alpha(\ell;\xi)}{p^2 - m_\alpha^2(\ell;\xi)} \;+\; \text{regular},$$

where $P_\alpha(\ell;\xi)$ is the rank–one spectral projector on W.

All p^2 are taken in the admitted band $\mathfrak{B}(\lambda_{\mathrm{UV}},\ell)$ of Definition 7.468.

Lemma 7.483 (Nielsen similarity flow for $\Gamma_2^{\mathcal{N}}$)**.** *Under the hypotheses of Definition 7.457 (evaluation on shell), there exists a smooth generator $K(\xi)$ on $W = \operatorname{im}\mathcal{N}(\xi)$ such that*

$$\partial_\xi \Gamma_2^{\mathcal{N}}(p^2;\ell,\xi) = \Big[\,K(\xi),\,\Gamma_2^{\mathcal{N}}(p^2;\ell,\xi)\,\Big] \quad \textit{for all } p^2 \in \mathfrak{B}(\lambda_{\mathrm{UV}},\ell). \tag{7.605}$$

Equivalently, with $S(\xi,\xi_0) := \mathcal{P}\exp\Big(\int_{\xi_0}^{\xi} K(\zeta)\,\mathrm{d}\zeta\Big)$,

$$\Gamma_2^{\mathcal{N}}(p^2;\ell,\xi) = S(\xi,\xi_0)\,\Gamma_2^{\mathcal{N}}(p^2;\ell,\xi_0)\,S(\xi,\xi_0)^{-1}. \tag{7.606}$$

Proof. By the Nielsen identity (eq. (7.581)) and projection to the BRST–physical subspace (Definition 7.457), the ξ–variation of any renormalised operator multiplet on W is an adjoint action generated by a smooth $K(\xi)$; cf. the Lax equation and similarity flow in Proposition 7.458. The 1PI two–point kernel is a bilinear form on the field/operator multiplet; therefore its ξ–variation is also given by the adjoint action $[K(\xi),\cdot]$ on W, which yields eq. (7.605). Path–ordered integration gives eq. (7.606). □

Corollary 7.484 (Zero gauge–parameter slope of pole locations and residues)**.** *Under the assumptions of Lemma 7.483, the physical pole positions and residues of* $\Gamma_2^{\mathcal{N}}$ *are gauge–parameter independent (within the tolerance ledger eq. (7.588)): for each channel* α *and fixed* $(\lambda_{\mathrm{UV}}, \ell)$*,*

$$\partial_\xi m_\alpha^2(\ell;\xi) = 0,$$
$$\partial_\xi Z_\alpha(\ell;\xi) = 0,$$

and the corresponding spectral projectors transform by similarity, $P_\alpha(\ell;\xi) = S(\xi,\xi_0)\, P_\alpha(\ell;\xi_0)\, S(\xi,\xi_0)^{-1}$.

Proof. By eq. (7.606), $\Gamma_2^{\mathcal{N}}(p^2;\ell,\xi) = S\,\Gamma_2^{\mathcal{N}}(p^2;\ell,\xi_0)\,S^{-1}$ with $S = S(\xi,\xi_0)$. Taking determinants,

$$\det \Gamma_2^{\mathcal{N}}(p^2;\ell,\xi) = \det\Big(S\,\Gamma_2^{\mathcal{N}}(p^2;\ell,\xi_0)\,S^{-1}\Big) = \det \Gamma_2^{\mathcal{N}}(p^2;\ell,\xi_0),$$

so the zero set in p^2 is independent of ξ. For a simple zero $p^2 = m_\alpha^2$ (definition above), the implicit function theorem gives $\partial_\xi m_\alpha^2 = 0$.

For residues, invert eq. (7.606) to obtain the propagator similarity,

$$D^{\mathcal{N}}(p^2;\ell,\xi) = \Big(\Gamma_2^{\mathcal{N}}(p^2;\ell,\xi)\Big)^{-1} = S(\xi,\xi_0)\, D^{\mathcal{N}}(p^2;\ell,\xi_0)\, S(\xi,\xi_0)^{-1}.$$

Near a simple pole $p^2 \approx m_\alpha^2$ one has $D^{\mathcal{N}}(p^2;\ell,\xi_0) = \dfrac{Z_\alpha(\ell;\xi_0)\,P_\alpha(\ell;\xi_0)}{p^2 - m_\alpha^2} + \text{regular}$. Conjugating,

$$D^{\mathcal{N}}(p^2;\ell,\xi) = \frac{Z_\alpha(\ell;\xi_0)\, SP_\alpha(\ell;\xi_0)S^{-1}}{p^2 - m_\alpha^2} + \text{regular},$$

which is precisely the Laurent form with the *same* scalar residue and the conjugated projector. Hence $Z_\alpha(\ell;\xi) = Z_\alpha(\ell;\xi_0)$ and $P_\alpha(\ell;\xi) = SP_\alpha(\ell;\xi_0)S^{-1}$. If one implements the numerical audit with subordinate operator norms, any residual ξ–variation is bounded by the tolerance in eq. (7.588); see also the general gauge–safety statements Corollary 7.459 and Lemma 7.473. □

Proposition 7.485 (Gauge–parameter independence of pole positions and residues)**.** *On any IR window* $\ln \ell \in \mathcal{I}_{\mathrm{IR}}$ *inside the Step 3 mask (cf. Definition 7.461 and Proposition 7.462), all physical pole positions and residues of the Nielsen–projected two–point kernel* $\Gamma_2^{\mathcal{N}}$ *(cf. Definition 7.482) are independent of the gauge parameter* ξ*:*

$$\boxed{\partial_\xi m_\alpha^2(\ell;\xi) = 0, \qquad \partial_\xi Z_\alpha(\ell;\xi) = 0, \qquad \forall\, \xi \in \mathcal{X}_{\mathrm{adm}}.} \tag{7.607}$$

Here $\mathcal{X}_{\mathrm{adm}}$ *denotes the admissible set of gauge parameters specified by the gauge audit eq. (7.588). In practice we require the tolerance check* $\sup_{\xi\in\mathcal{X}_{\mathrm{adm}}} \big|\partial_\xi m_\alpha^2\big|,\ \sup_{\xi\in\mathcal{X}_{\mathrm{adm}}} \big|\partial_\xi Z_\alpha\big| \le \varepsilon_{\mathrm{gauge}}$ *with* $\varepsilon_{\mathrm{gauge}} < \varepsilon_{\mathrm{plat}}$ *(cf. eq. (7.588)).*

Proof. Fix $(\lambda_{\mathrm{UV}}, \ell)$ and work on the physical subspace $W = \operatorname{im}\mathcal{N}$ evaluated on shell (Definition 7.457). By the Nielsen similarity flow for two–point kernels (Lemma 7.483), there exists a smooth $K(\xi)$ such that

$$\Gamma_2^{\mathcal{N}}(p^2;\ell,\xi) = S(\xi,\xi_0)\,\Gamma_2^{\mathcal{N}}(p^2;\ell,\xi_0)\,S(\xi,\xi_0)^{-1}, \quad S(\xi,\xi_0) := \mathcal{P}\exp\Big(\int_{\xi_0}^{\xi} K(\zeta)\,\mathrm{d}\zeta\Big).$$

Pole positions. Let $F(p^2,\xi) := \det \Gamma_2^{\mathcal{N}}(p^2;\ell,\xi)$. Determinants are invariant under similarity, hence $F(p^2,\xi) = F(p^2,\xi_0)$ for all p^2 in the admitted band $\mathfrak{B}(\lambda_{\mathrm{UV}},\ell)$ (Definition 7.468). Therefore $\partial_\xi F(p^2,\xi) = 0$ pointwise in p^2. If $m_\alpha^2(\ell;\xi)$ is a *simple* physical pole (definition in Definition 7.482), then $F\big(m_\alpha^2(\ell;\xi),\xi\big) = 0$ with $\partial_{p^2} F \neq 0$ at $p^2 = m_\alpha^2$. Differentiating the identity $F(m_\alpha^2(\xi),\xi) = 0$ and using $\partial_\xi F \equiv 0$ gives

$$0 = \partial_\xi F(m_\alpha^2,\xi) = \partial_{p^2} F(m_\alpha^2,\xi)\,\partial_\xi m_\alpha^2(\xi),$$

hence $\partial_\xi m_\alpha^2(\ell;\xi) = 0$.

Residues. Write $A(p^2,\xi) := \Gamma_2^{\mathcal{N}}(p^2;\ell,\xi)$ and $D(p^2,\xi) := A(p^2,\xi)^{-1}$. Near a simple pole $p^2 = m_\alpha^2$ one has the Laurent form

$$D(p^2,\xi) = \frac{Z_\alpha(\ell;\xi)\,P_\alpha(\ell;\xi)}{p^2 - m_\alpha^2(\ell;\xi)} + \text{regular},$$

with rank–one spectral projector P_α on W (Definition 7.482). From similarity,

$$D(p^2,\xi) = S(\xi,\xi_0)\,D(p^2,\xi_0)\,S(\xi,\xi_0)^{-1}.$$

Since m_α^2 is already ξ–independent, conjugating the Laurent expansion at ξ_0 yields

$$D(p^2,\xi) = \frac{Z_\alpha(\ell;\xi_0)\,SP_\alpha(\ell;\xi_0)S^{-1}}{p^2 - m_\alpha^2} + \text{regular}.$$

Thus the scalar residue is unchanged, $Z_\alpha(\ell;\xi) = Z_\alpha(\ell;\xi_0)$, while the projector transforms by similarity, $P_\alpha(\ell;\xi) = SP_\alpha(\ell;\xi_0)S^{-1}$. Equivalently, using left/right eigenvectors $A(m_\alpha^2,\xi)v_\alpha = 0$, $w_\alpha^\top A(m_\alpha^2,\xi) = 0$ with $w_\alpha^\top v_\alpha = 1$, one has

$$Z_\alpha(\ell;\xi) = \Big(w_\alpha^\top\,\partial_{p^2} A(m_\alpha^2,\xi)\,v_\alpha\Big)^{-1},$$

and under similarity $v_\alpha \mapsto S^{-1}v_\alpha$, $w_\alpha^\top \mapsto w_\alpha^\top S$ and $\partial_{p^2}A \mapsto S^{-1}(\partial_{p^2}A)S$, leaving the scalar $w_\alpha^\top \partial_{p^2}A\,v_\alpha$ invariant.

Tolerance statement. On Step 3 IR windows the band–limited kernels vary smoothly and are uniformly bounded (Lemma 7.469 and Proposition 7.462), so any finite–statistics or truncation–induced residual ξ–dependence is controlled by the audit bound eq. (7.588), yielding the practical check stated.

This proves (7.607). □

Remark 7.486 (Degenerate multiplets)**.** If several physical modes share the same pole (mass degeneracy), Lemma 7.483 implies a *block* similarity within the degenerate subspace. The pole position is ξ–independent and the spectral projector onto the multiplet is conjugated as a block. Individual residues may rotate within the multiplet; choosing a smooth basis (e.g. by good quantum numbers) renders each residue gauge–independent.

Reporting. Evaluate (7.607) throughout the IR window $\ln\ell \in \mathcal{I}_{\mathrm{IR}}$ at or beyond ℓ_R and report the resulting bands together with the law–channel slope $\varsigma_{\mathrm{law}}^{(\tau)}$ from Step 2 (cf. eq. (7.542)).

Discrete estimators (audit path).

Definition 7.487 (Centered discrete estimators and optional smoothing)**.** Let $v_k := \ln \ell_k$ be a uniform grid with step $h > 0$ (cf. the centered–difference pattern used for Step 4 in Definition 7.446), and let $\{\xi_r\}_r$ be a *uniform* grid of gauge parameters with spacing $\Delta\xi > 0$ inside $\mathcal{X}_{\mathrm{adm}}$. Define the centered finite–difference estimators

$$\boxed{\widehat{\partial_v \mathcal{W}}(v_k) := \frac{\mathcal{W}(v_{k+1}) - \mathcal{W}(v_{k-1})}{2h}, \qquad \widehat{\partial_\xi m_\alpha^2}(\ell_k;\xi_r) := \frac{m_\alpha^2(\ell_k;\xi_{r+1}) - m_\alpha^2(\ell_k;\xi_{r-1})}{2\Delta\xi},} \tag{7.608}$$

and analogously for $Z_\alpha(\ell;\xi)$, with $\mathcal{W}$ defined in Definition 7.468 and (m_α^2, Z_α) in Definition 7.482. When numerical ripples are present, apply a low–pass LP_τ in v or in ξ whose kernel satisfies the moment conditions of eqs. (7.537) and (7.538); such filters preserve first derivatives up to an $\mathcal{O}(\tau^2)$ bias.

Lemma 7.488 (Consistency, rate, and smoothing bias)**.** *Assume $\mathcal{W} \in C^3$ in v on the Step 3 IR window (cf. Definition 7.461 and Proposition 7.462) and $m_\alpha^2, Z_\alpha \in C^3$ in ξ on $\mathcal{X}_{\mathrm{adm}}$. Then the centered estimators in (7.608) satisfy*

$$\widehat{\partial_v \mathcal{W}}(v_k) = \partial_v \mathcal{W}(v_k) + \mathcal{O}(h^2), \qquad \widehat{\partial_\xi m_\alpha^2}(\ell_k;\xi_r) = \partial_\xi m_\alpha^2(\ell_k;\xi_r) + \mathcal{O}\big((\Delta\xi)^2\big),$$

with $\Delta\xi := \xi_{r+1} - \xi_r = \xi_r - \xi_{r-1}$; the same holds for Z_α. After applying LP_τ (satisfying eqs. (7.537) and (7.538)), the additional bias is $\mathcal{O}(\tau^2)$. Implicit constants are uniform on compact subwindows thanks to the curvature/TV guards eq. (7.548).

Proof. **(A) Centered differences in v.** Let $f(v) := \mathcal{W}(v)$ and expand at v_k:

$$f(v_{k\pm1}) = f(v_k) \pm h f'(v_k) + \frac{h^2}{2} f''(v_k) \pm \frac{h^3}{6} f^{(3)}(\zeta_\pm),$$

for some $\zeta_\pm$ between v_k and $v_{k\pm1}$. Subtract and divide by $2h$:

$$\widehat{\partial_v \mathcal{W}}(v_k) = \frac{f(v_{k+1}) - f(v_{k-1})}{2h} = f'(v_k) + \frac{h^2}{6}\,\frac{f^{(3)}(\zeta_+) - f^{(3)}(\zeta_-)}{2} = \partial_v \mathcal{W}(v_k) + \mathcal{O}(h^2).$$

On the Step 3 window, the curvature/TV guard eq. (7.548) bounds $f^{(3)}$, giving a uniform $\mathcal{O}(h^2)$.

(B) Centered differences in ξ. Fix ℓ_k and set $g(\xi) := m_\alpha^2(\ell_k;\xi)$ (the same argument applies to Z_α). With $\Delta\xi := \xi_{r+1} - \xi_r = \xi_r - \xi_{r-1}$,

$$g(\xi_{r\pm1}) = g(\xi_r) \pm \Delta\xi\, g'(\xi_r) + \frac{(\Delta\xi)^2}{2} g''(\xi_r) \pm \frac{(\Delta\xi)^3}{6} g^{(3)}(\eta_\pm),$$

so

$$\widehat{\partial_\xi m_\alpha^2}(\ell_k;\xi_r) = \frac{g(\xi_{r+1}) - g(\xi_{r-1})}{2\Delta\xi} = g'(\xi_r) + \mathcal{O}\big((\Delta\xi)^2\big) = \partial_\xi m_\alpha^2(\ell_k;\xi_r) + \mathcal{O}\big((\Delta\xi)^2\big).$$

Smoothness (C^3) on $\mathcal{X}_{\mathrm{adm}}$ yields uniform constants on compact subsets; the admissible set is where the gauge audit eq. (7.588) is performed.

(C) Smoothing bias. Let LP_τ be the convolution/averaging operator in v (or in ξ) with kernel K_τ satisfying the moment conditions of eqs. (7.537) and (7.538): $\int K_\tau = 1$ and $\int t\,K_\tau(t)\,dt = 0$. For a C^2 function h, the standard moment expansion gives

$$(\mathsf{LP}_\tau h)(x) = h(x) + \frac{\mu_2(K)}{2}\,\tau^2 h''(x) + \mathcal{O}(\tau^3),$$

so, applied either to $\mathcal{W}$ in v or to m_α^2, Z_α in ξ, the induced bias on first derivatives is $\mathcal{O}(\tau^2)$ (the $\mathcal{O}(\tau^3)$ term is uniform on compact sets under eq. (7.548)). Combining with (A,B) yields the stated $\mathcal{O}(h^2) + \mathcal{O}((\Delta\xi)^2) + \mathcal{O}(\tau^2)$ rates. □

Corollary 7.489 (Estimator outcome under Nielsen protection)**.** *Assume the hypotheses of Definition 7.487 and Lemma 7.488 and the zero gauge–slope statement Proposition 7.485. Then, for every grid point $v_k = \ln \ell_k$ in the Step 3 IR window (cf. Definition 7.461 and Proposition 7.462) and every $\xi_r \in \mathcal{X}_{\mathrm{adm}}$, the centered estimators (optionally with the low–pass LP_τ of eqs. (7.537) and (7.538)) satisfy*

$$\widehat{\partial_\xi m_\alpha^2}(\ell_k;\xi_r) = 0 + \mathcal{O}\Big((\Delta\xi)^2 + \tau^2\Big), \qquad \widehat{\partial_\xi Z_\alpha}(\ell_k;\xi_r) = 0 + \mathcal{O}\Big((\Delta\xi)^2 + \tau^2\Big),$$

uniformly on compact subwindows of the Step 3 mask. In numerical audits, these errors are required to fall below the tolerance in eq. (7.588).

Proof. By Proposition 7.485 one has, on $W = \operatorname{im}\mathcal{N}$ and throughout the Step 3 IR window,

$$\partial_\xi m_\alpha^2(\ell;\xi) = 0, \qquad \partial_\xi Z_\alpha(\ell;\xi) = 0 \quad \text{for all } \xi \in \mathcal{X}_{\mathrm{adm}}.$$

Fix ℓ_k and define $g(\xi) := m_\alpha^2(\ell_k;\xi)$ and $h(\xi) := Z_\alpha(\ell_k;\xi)$. The regularity assumptions of Lemma 7.488 give $g, h \in C^3(\mathcal{X}_{\mathrm{adm}})$. For the centered finite–difference estimator on the uniform ξ–grid (spacing $\Delta\xi$) introduced in Definition 7.487,

$$\widehat{\partial_\xi g}(\xi_r) = \frac{g(\xi_{r+1}) - g(\xi_{r-1})}{2\Delta\xi} = \partial_\xi g(\xi_r) + \mathcal{O}\Big((\Delta\xi)^2\Big),$$

and the same holds for h, with implicit constants uniform on compact subsets of the window by Lemma 7.488 (the curvature/TV guard eq. (7.548) ensures the needed uniform bounds).

Since $\partial_\xi g \equiv 0$ and $\partial_\xi h \equiv 0$ by Proposition 7.485, the unsmoothed estimators obey

$$\widehat{\partial_\xi m_\alpha^2}(\ell_k;\xi_r) = \mathcal{O}\Big((\Delta\xi)^2\Big), \qquad \widehat{\partial_\xi Z_\alpha}(\ell_k;\xi_r) = \mathcal{O}\Big((\Delta\xi)^2\Big).$$

If, as permitted in Definition 7.487, one applies the low–pass operator LP_τ in ξ with kernel satisfying eqs. (7.537) and (7.538), the moment expansion yields an additional $\mathcal{O}(\tau^2)$ bias on first derivatives (see Lemma 7.488). Combining the finite–difference error with the smoothing bias gives the stated $\mathcal{O}\Big((\Delta\xi)^2 + \tau^2\Big)$ rates for both m_α^2 and Z_α.

Uniformity on compact subwindows of the Step 3 mask follows from the window consistency Proposition 7.462 together with the curvature/TV guard eq. (7.548). Finally, the audit requirement in eq. (7.588) enforces that these numerical errors lie below the prescribed gauge–tolerance threshold. □

Acceptance criteria (identity ledger). On the certified IR window $\mathcal{I}_{\mathrm{IR}}$ (cf. Definition 7.461 and Proposition 7.462), we *accept* the identity ledger when three checks jointly pass. First, **ST exactness**: the on–shell Slavnov–Taylor identity holds at all probed scales, exactly as in eq. (7.601),

$$\boxed{\mathcal{S}\big(\Gamma^{(\lambda_{\mathrm{UV}},\ell)}\big) = 0 \quad \text{upon} \quad \frac{\delta\Gamma^{(\lambda_{\mathrm{UV}},\ell)}}{\delta\phi} = 0.} \tag{7.609}$$

Second, **Ward restoration**: the band–projected residual decays with strictly negative logarithmic slope across the window and is within plateau tolerance at the restoration length (cf. Definition 7.478),

$$\boxed{\widehat{\partial_v \mathcal{W}}(v_k) \;\le\; -\,c \;<\; 0 \quad \forall\, v_k \in \mathcal{I}_{\mathrm{IR}}, \qquad \mathcal{W}(\ell_R) \;\le\; \varepsilon_{\mathrm{plat}},} \tag{7.610}$$

with c chosen commensurate with the curvature/TV guards of eq. (7.548), and where $\widehat{\partial_v \mathcal{W}}$ is the centered estimator of Definition 7.487. Third, **Gauge–parameter independence**: every quoted IR pole and residue is flat in ξ to within the audit tolerance (cf. Proposition 7.485),

$$\boxed{\Big|\widehat{\partial_\xi m_\alpha^2}(\ell_k;\xi_r)\Big| \;\le\; \varepsilon_{\mathrm{gauge}}, \qquad \Big|\widehat{\partial_\xi Z_\alpha}(\ell_k;\xi_r)\Big| \;\le\; \varepsilon_{\mathrm{gauge}},} \tag{7.611}$$

for all ℓ_k with $v_k \in \mathcal{I}_{\mathrm{IR}}$ and all $\xi_r \in \Xi_{\mathrm{adm}}$, where the discrete derivatives are those of Definition 7.487 and the continuum statement is eq. (7.607).

Outputs for Step 5. To document acceptance and enable replication, report (i) a plot of $v \mapsto \mathcal{W}(e^v)$ over $\mathcal{I}_{\mathrm{IR}}$ with the fitted (optionally smoothed) slope, explicitly showing $\widehat{\partial_v \mathcal{W}} \le -c$ and a marker at $v = \ln \ell_R$; (ii) compact tables of the discrete gauge–slopes $\widehat{\partial_\xi m_\alpha^2}$ and $\widehat{\partial_\xi Z_\alpha}$ evaluated at/beyond ℓ_R across Ξ_{adm}, including uncertainty bands reflecting $(h, \Delta\xi, \tau)$ from Definition 7.487; and (iii) a short cross–reference ledger confirming that (a) the Step 2 law–channel slope is positive on the same window (eq. (7.542)), (b) the Step 4 projected breaking exponents satisfy $\mathrm{Re}\, y_{\tilde{B}} < 0$ (Proposition 7.450 and Corollary 7.456), and (c) the restoration length ℓ_R is that certified in Step 3 (Definition 7.430 and eq. (7.551)). Together with the on–shell ST identity at all bands (Proposition 7.476), these artefacts close the Step 5 identity ledger.

7.8.6 Spectral Reconvergence to Symmetric Multiplets

Objective. On the IR plateau (the window $\mathcal{I}_{\mathrm{IR}}$ of eq. (7.542) with $v = \ln \ell \ge \ln \ell_R$ from eq. (7.551)), we demonstrate *multiplet reconvergence*: for each representation ρ of the emergent symmetry G with components $a = 1, \dots, d_\rho$, the mass and residue splittings close,

$$\frac{\delta m_\rho^2(\ell)}{\overline{m_\rho^2}(\ell)} \longrightarrow 0, \qquad \frac{\delta Z_\rho(\ell)}{\overline{Z_\rho}(\ell)} \longrightarrow 0 \quad \text{as} \quad v \in \mathcal{I}_{\mathrm{IR}},\ v \uparrow,$$

consistent with Kato continuity under the ℓ–flow [115] and the RG irrelevance of explicit breakings (Step 7.8.4; [389, 91, 82]).

Band–projected 2–point sector and pole data.

Definition 7.490 (Band–projected two–point sector and pole data)**.** Fix $(\lambda_{\mathrm{UV}}, \ell)$ and admitted momenta $p \in \mathfrak{B}(\lambda_{\mathrm{UV}}, \ell)$ (cf. Definition 7.468). Let $\mathscr{B}_{\lambda_{\mathrm{UV}},\ell}$ be the band–limiting map of Definition 7.475 and let $\mathcal{N}$ be the Nielsen projector of Definition 7.457 with $W = \operatorname{im}\mathcal{N}$. The band– and Nielsen–projected 1PI two–point kernel is

$$\Gamma_2^{\mathcal{N}}(p^2; \lambda_{\mathrm{UV}}, \ell) \;:=\; \Big(\mathscr{B}_{\lambda_{\mathrm{UV}},\ell}\big[\Gamma^{(2)}(p^2)\big] \Big)^{\mathcal{N}}\Big|_W .$$

Physical poles and residues are those of $\Gamma_2^{\mathcal{N}}$ as in Definition 7.482: simple poles $p^2 = m_\alpha^2(\ell)$ are simple zeros of $\det \Gamma_2^{\mathcal{N}}(p^2; \lambda_{\mathrm{UV}}, \ell)$ with $\partial_{p^2} \det \Gamma_2^{\mathcal{N}} \neq 0$, and their scalar residues are

$$\boxed{\; Z_\alpha(\ell) \;=\; \Big(w_\alpha^\top \, \partial_{p^2} \Gamma_2^{\mathcal{N}}\big(m_\alpha^2; \lambda_{\mathrm{UV}}, \ell\big)\, v_\alpha \Big)^{-1}, \;} \tag{7.612}$$

where v_α, w_α are right/left null vectors of $\Gamma_2^{\mathcal{N}}$ at $p^2 = m_\alpha^2$ on W normalized by $w_\alpha^\top v_\alpha = 1$. This spectral characterisation and the use of residues at simple poles follow standard QFT practice [91, 82], while continuity of the pole data under parameter flows is governed by Kato perturbation theory [115]. Gauge–parameter insensitivity on shell (within the admitted audit tolerance) is ensured by the ST/BRST framework and algebraic renormalisation [117, 243, 442] together with Propositions 7.476 and 7.485 and Corollary 7.477.

Intuition (outside the environment): we band-limit with $\mathscr{B}_{\lambda_{\mathrm{UV}},\ell}$, project to the BRST-physical/Nielsen subspace W, then read off pole positions and residues from $\Gamma_2^{\mathcal{N}}$. The cited sources cover (i) pole/residue practice in QFT [91, 82], (ii) continuity of simple poles under smooth parameter flows [115], and (iii) on-shell gauge safety via ST/BRST and algebraic renormalisation [117, 243, 442].

Multiplet identification and dispersion measures.

Definition 7.491 (IR multiplets and dispersion)**.** Let ρ be an irrep of the emergent symmetry G acting on the physical subspace $W = \operatorname{im}\mathcal{N}$ (cf. Definition 7.457). Assume that for ℓ in the IR window $\mathcal{I}_{\mathrm{IR}}$ (cf. eqs. (7.542) and (7.551)) there exists a smooth projector $P^\rho(\ell)$ on W whose range collects the d_ρ physical modes that reconverge to the ρ–multiplet as explicit breakings become irrelevant (Step 7.8.4). Let $\mathcal{I}_\rho(\ell)$ be the index set of poles of $\Gamma_2^{\mathcal{N}}$ (cf. Definitions 7.482 and 7.490) associated with $P^\rho(\ell)$. Define the *multiplet means* and *max–deviation dispersions* by

$$\overline{m_\rho^2}(\ell) := \frac{1}{d_\rho} \sum_{i \in \mathcal{I}_\rho(\ell)} m_i^2(\ell), \qquad \delta m_\rho^2(\ell) := \max_{i \in \mathcal{I}_\rho(\ell)} \Big| m_i^2(\ell) - \overline{m_\rho^2}(\ell) \Big|,$$

and analogously for residues $\overline{Z_\rho}(\ell)$ and $\delta Z_\rho(\ell)$.

$P^\rho(\ell)$ tracks a spectral cluster of the band– and Nielsen–projected kernel $\Gamma_2^{\mathcal{N}}(p^2; \lambda_{\mathrm{UV}}, \ell)$ (Definition 7.490); $\mathcal{I}_\rho(\ell)$ labels the corresponding simple poles $p^2 = m_i^2(\ell)$ in that cluster, and the dispersion measures summarize intra-multiplet splittings using already-defined pole data [91, 82].

Lemma 7.492 (Kato continuity and smooth tracking)**.** *Assume that $v \mapsto \Gamma_2^{\mathcal{N}}(p^2;\lambda_{\mathrm{UV}},e^v)$ is C^1 jointly in (p^2,v) on the admitted band $\mathfrak{B}(\lambda_{\mathrm{UV}},\ell)$ (cf. Definition 7.468) and that, within a given cluster selected by $P^\rho(e^v)$, all poles remain simple and separated from the rest of the spectrum (no accidental crossings beyond symmetry). Then there exist C^1 functions $m_i^2(v)$ and $Z_i(v)$ for $i\in\mathcal{I}_\rho(e^v)$, and the maps $\ell\mapsto \overline{m_\rho^2}(\ell),\delta m_\rho^2(\ell)$ (and likewise for residues) are continuous on the IR window $\mathcal{I}_{\mathrm{IR}}$.*

Proof. Set $A(p^2,v):=\Gamma_2^{\mathcal{N}}(p^2;\lambda_{\mathrm{UV}},e^v)$ and $F(p^2,v):=\det A(p^2,v)$. By hypothesis, A is C^1 in (p^2,v) on the admitted band, hence so is F.

(i) *Smooth pole tracking.* Let $p^2=m_i^2(v_0)$ be a simple zero of $F(\cdot,v_0)$ in the cluster selected by $P^\rho(e^{v_0})$, so $\partial_{p^2}F(m_i^2(v_0),v_0)\neq 0$ (simplicity; cf. Definition 7.482). The Implicit Function Theorem yields a neighborhood U of v_0 and a unique C^1 function $m_i^2(v)$ on U with $F(m_i^2(v),v)=0$ and $m_i^2(v_0)$ as given. Repeating for each simple pole in the finite cluster and using the separation (no crossings beyond symmetry) provides C^1 labels $m_i^2(v)$ for all $i\in\mathcal{I}_\rho(e^v)$ across the window. This is a standard application of Kato's perturbation framework for isolated finite-dimensional spectral clusters [115, Chs. II–IV].

(ii) *Residue regularity.* At a simple pole $p^2=m_i^2(v)$, the inverse $D(p^2,v):=A(p^2,v)^{-1}$ admits the Laurent form

$$D(p^2,v) \;=\; \frac{Z_i(v)\,P_i(v)}{p^2-m_i^2(v)}+\text{regular},$$

where $P_i(v)=v_i(v)\,w_i(v)^\top$ is the rank-one spectral projector on W with $A(m_i^2(v),v)\,v_i(v)=0$, $w_i(v)^\top A(m_i^2(v),v)=0$, $w_i^\top v_i=1$, and

$$Z_i(v) \;=\; \Big(w_i(v)^\top\,\partial_{p^2}A\Big(m_i^2(v),v\Big)\,v_i(v)\Big)^{-1} \quad \text{(cf. Definition 7.490)}.$$

By simplicity and C^1 dependence of A, one may choose $v_i(\cdot),w_i(\cdot)$ C^1 in v (locally uniquely up to smooth rephasing), see [115, Thm. II-1.8, Sec. II-5]. Since $\partial_{p^2}A$ is C^1 and the denominator above never vanishes at a simple pole, $Z_i(v)$ is C^1.

(iii) *Continuity of multiplet averages and max–deviations.* By construction, $P^\rho(e^v)$ is smooth and selects a fixed-rank cluster, so the index set $\mathcal{I}_\rho(e^v)$ varies without changing cardinality. Finite sums and averages of continuous functions $m_i^2(e^v)$ are continuous, hence $\overline{m_\rho^2}(\ell)$ is continuous. The map $x\mapsto \max_j x_j$ is continuous on $\mathbb{R}^{d_\rho}$, so $\delta m_\rho^2(\ell)=\max_{i\in\mathcal{I}_\rho(\ell)}|m_i^2(\ell)-\overline{m_\rho^2}(\ell)|$ is continuous; the same argument applies to residues.

Putting (i)–(iii) together gives the claim. □

Proposition 7.493 (Linear response of splittings to explicit breakings)**.** *Work on the physical subspace $W=\operatorname{im}\mathcal{N}$ and within a fixed IR window $\mathcal{I}_{\mathrm{IR}}$. For the ρ–block selected by $P^\rho(\ell)$ (cf. Definition 7.491) of the band– and Nielsen–projected kernel $\Gamma_2^{\mathcal{N}}$ (cf. Definition 7.490), assume the explicit breaking couplings $\widetilde{\varepsilon}_{\tilde B}(\kappa/\ell)$ are sufficiently small. Then one can decompose, for p^2 near the common symmetric pole $m_{\rho,\mathrm{sym}}^2(\ell)$,*

$$\Gamma_\rho^{(2)}(p^2;\ell)=\Big[\Gamma_\rho^{(2)}\Big]_{\mathrm{sym}}(p^2;\ell)\,\mathbf{1}_{d_\rho}\;+\;\sum_{\tilde B}\widetilde{\varepsilon}_{\tilde B}(\kappa/\ell)\,\Delta_\rho^{(\tilde B)}(p^2;\ell)\;+\;\mathcal{O}(\|\widetilde{\varepsilon}\|^2),$$

where $\big[\Gamma^{(2)}_\rho\big]_{\mathrm{sym}} := \frac{1}{d_\rho}\mathrm{Tr}\,\Gamma^{(2)}_\rho$ *and each* $\Delta^{(\tilde B)}_\rho$ *is traceless on the* ρ*–block (Schur/Wigner–Eckart form; the trace part has been absorbed into the symmetric piece). Consequently, the intra–multiplet mass and residue splittings obey*

$$\delta m^2_\rho(\ell) \;\le\; C_\rho(\ell)\sum_{\tilde B}\big|\widetilde{\varepsilon}_{\tilde B}(\kappa/\ell)\big| + \mathcal{O}(\|\widetilde{\varepsilon}\|^2), \qquad \delta Z_\rho(\ell) \;\le\; C'_\rho(\ell)\sum_{\tilde B}\big|\widetilde{\varepsilon}_{\tilde B}(\kappa/\ell)\big| + \mathcal{O}(\|\widetilde{\varepsilon}\|^2), \tag{7.613}$$

with $\delta m^2_\rho, \delta Z_\rho$ *as in Definition 7.491.*

Proof. Let $A_\rho(p^2;\ell) := P^\rho(\ell)\,\Gamma^{\mathcal{N}}_2(p^2;\lambda_{\mathrm{UV}},\ell)\,P^\rho(\ell)\big|_{\mathrm{ran}\,P^\rho}$ denote the $d_\rho\times d_\rho$ block (cf. Definition 7.490). Decompose A_ρ into its scalar part and traceless part on the block:

$$A_\rho(p^2;\ell) = a_\rho(p^2;\ell)\,\mathbf{1}_{d_\rho} + \sum_{\tilde B}\widetilde{\varepsilon}_{\tilde B}(\kappa/\ell)\,\Delta^{(\tilde B)}_\rho(p^2;\ell) + \mathcal{O}(\|\widetilde{\varepsilon}\|^2),$$

where $a_\rho := \frac{1}{d_\rho}\mathrm{Tr}\,A_\rho$ and $\mathrm{Tr}\,\Delta^{(\tilde B)}_\rho \equiv 0$. Representation theory (Schur's lemma / Wigner–Eckart) justifies that in the symmetric limit the block is proportional to the identity, while explicit breakings furnish traceless tensor structures within the block [91, 82]. Let $m^2_{\rho,\mathrm{sym}}(\ell)$ solve $a_\rho\big(m^2_{\rho,\mathrm{sym}}(\ell);\ell\big)=0$; by Definition 7.482, simple poles in the block are then the zeros of the d_ρ eigenvalue branches of A_ρ near $m^2_{\rho,\mathrm{sym}}(\ell)$.

(i) Linearized pole shifts. Evaluate at $p^2 = m^2_{\rho,\mathrm{sym}}(\ell)$ and set

$$\mu_\rho(\ell) := \frac{1}{d_\rho}\,\mathrm{Tr}\,\partial_{p^2}A_\rho\big(m^2_{\rho,\mathrm{sym}}(\ell);\ell\big), \qquad \delta A_\rho := \sum_{\tilde B}\widetilde{\varepsilon}_{\tilde B}(\kappa/\ell)\,\Delta^{(\tilde B)}_\rho\big(m^2_{\rho,\mathrm{sym}}(\ell);\ell\big).$$

By simplicity of poles within the cluster (cf. Definition 7.482 and Lemma 7.492) and C^1 regularity, Kato/implicit–function theory gives, for each component i,

$$m^2_i(\ell) - m^2_{\rho,\mathrm{sym}}(\ell) \;=\; -\frac{\theta_i(\ell)}{\mu_\rho(\ell)} \;+\; \mathcal{O}(\|\widetilde{\varepsilon}\|^2),$$

where $\theta_i(\ell)$ are the eigenvalues of the traceless perturbation δA_ρ on the ρ–block (degenerate perturbation theory in a finite cluster) [115, Chs. II–IV]. Hence, using $|\theta_i|\le\|\delta A_\rho\|$,

$$\begin{aligned}\delta m^2_\rho(\ell) &= \max_i\big|m^2_i - m^2_{\rho,\mathrm{sym}}\big| \\ &\le\; \frac{\|\delta A_\rho\|}{|\mu_\rho(\ell)|} + \mathcal{O}(\|\widetilde{\varepsilon}\|^2) \\ &\le\; \Big\|\big(\partial_{p^2}A_\rho\big)^{-1}\big(m^2_{\rho,\mathrm{sym}};\ell\big)\Big\|\sum_{\tilde B}\big|\widetilde{\varepsilon}_{\tilde B}(\kappa/\ell)\big|\,\big\|\Delta^{(\tilde B)}_\rho\big\| \\ &\qquad + \mathcal{O}(\|\widetilde{\varepsilon}\|^2),\end{aligned}$$

which yields (7.613) for δm^2_ρ upon absorbing blockwise norms into $C_\rho(\ell)$.

(ii) Linearized residue shifts. Write $D_\rho(p^2;\ell) := A_\rho(p^2;\ell)^{-1}$ and recall that, at a simple pole $p^2 = m^2_i(\ell)$,

$$Z_i(\ell) = \Big(w_i^{\top}\,\partial_{p^2}A_\rho\big(m^2_i(\ell);\ell\big)\,v_i\Big)^{-1} \quad\text{with}\quad A_\rho(m^2_i)v_i = 0,\;\; w_i^{\top}A_\rho(m^2_i) = 0,\;\; w_i^{\top}v_i = 1$$

(cf. Definition 7.490). Differentiate this formula with respect to the breaking amplitudes $\widetilde{\varepsilon}_{\tilde{B}}$ at the symmetric point, using the first–order variations of m_i^2, v_i, w_i, and A_ρ from (i) (Kato's eigenvector perturbation within a simple cluster) [115, Thm. II-1.8, Sec. II-5]. The result is

$$Z_i(\ell) - Z_{\rho,\mathrm{sym}}(\ell) \;=\; -\,Z_{\rho,\mathrm{sym}}(\ell)^2 \left(w_{i,\mathrm{sym}}^{\top}\, \partial_{p^2} \delta A_\rho\, v_{i,\mathrm{sym}} \right) \;+\; \mathcal{O}(\|\widetilde{\varepsilon}\|^2),$$

up to additional linear terms proportional to $\partial_{p^2}^2 A_\rho$ times the already–bounded mass shift from (i). Taking maxima over i and using operator–norm bounds gives

$$\delta Z_\rho(\ell) \;\le\; C'_\rho(\ell) \sum_{\tilde{B}} \left| \widetilde{\varepsilon}_{\tilde{B}}(\kappa/\ell) \right| + \mathcal{O}(\|\widetilde{\varepsilon}\|^2),$$

with $C'_\rho(\ell)$ depending on $\|Z_{\rho,\mathrm{sym}}(\ell)\|$, $\|\partial_{p^2} \Delta_\rho^{(\tilde{B})}\|$, and $\|\partial_{p^2}^2 A_\rho\|$ evaluated at $p^2 = m^2_{\rho,\mathrm{sym}}(\ell)$.

Both bounds use only blockwise linear algebra and standard degenerate perturbation theory for isolated finite clusters, together with the spectral/residue conventions of Definitions 7.482 and 7.490 and smooth tracking from Lemma 7.492; see [115] for the perturbation framework and [91, 82] for the two–point/pole–residue practice. This establishes (7.613). □

Theorem 7.494 (Multiplet reconvergence on the IR plateau)**.** *Work on the Step 3 IR window with $\varsigma_{\mathrm{law}}^{(\tau)}(\ell) > 0$ and curvature/TV guards (cf. eqs. (7.542), (7.548) and (7.551) and Proposition 7.462). Assume Step 4 irrelevance:* $\mathrm{Re}\, y_{\tilde{B}} < 0$ *for all excited projected breakings (cf. Proposition 7.450 and Corollary 7.456). Suppose further that the multiplet mean satisfies either*

$$(A)\;\; \overline{m_\rho^2}(\ell) \;\ge\; m^2_{\rho,\min} > 0 \qquad \text{or} \qquad (B)\;\; \overline{m_\rho^2}(\ell) \;\asymp\; \ell^{-\alpha} \;\; \text{on the window with} \;\; \alpha < \gamma_\star,$$

where $\gamma_\star := \min_{\tilde{B}} |\,\mathrm{Re}\, y_{\tilde{B}}| > 0$ and $\mu = \kappa/\ell$. Then, as $v = \ln \ell \uparrow$ within $\mathcal{I}_{\mathrm{IR}}$ (equivalently $\mu \downarrow 0$),

$$\boxed{\frac{\delta m_\rho^2(\ell)}{\overline{m_\rho^2}(\ell)} \;\longrightarrow\; 0, \qquad \frac{\delta Z_\rho(\ell)}{\overline{Z_\rho}(\ell)} \;\longrightarrow\; 0,} \tag{7.614}$$

with rates controlled by the slowest irrelevant exponent:

$$\frac{\delta m_\rho^2(\ell)}{\overline{m_\rho^2}(\ell)} = \begin{cases} \mathcal{O}\Big(\mu^{\gamma_\star}\, \mathrm{poly}(\ln \mu)\Big) & \text{in case (A)}, \\ \mathcal{O}\Big(\ell^{-(\gamma_\star - \alpha)}\, \mathrm{poly}(\ln \ell)\Big) & \text{in case (B)}, \end{cases}$$

and likewise for $\delta Z_\rho / \overline{Z_\rho}$.

Proof. All statements are taken on the physical subspace $W = \mathrm{im}\,\mathcal{N}$ and for the band– and Nielsen–projected kernel $\Gamma_2^{\mathcal{N}}$ (cf. Definition 7.490); gauge–parameter safety on shell follows from Propositions 7.476 and 7.485 and Corollary 7.477 and is immaterial for the bounds below.

Step 1: Linear response of intra–multiplet splittings. By Proposition 7.493, for the ρ–block selected by $P^\rho(\ell)$ (cf. Definition 7.491) there exist bounded functions $C_\rho(\ell), C'_\rho(\ell)$ on the IR window such that

$$\delta m_\rho^2(\ell) \;\le\; C_\rho(\ell) \sum_{\tilde{B}} \left| \widetilde{\varepsilon}_{\tilde{B}}(\kappa/\ell) \right| + \mathcal{O}(\|\widetilde{\varepsilon}\|^2), \qquad \delta Z_\rho(\ell) \;\le\; C'_\rho(\ell) \sum_{\tilde{B}} \left| \widetilde{\varepsilon}_{\tilde{B}}(\kappa/\ell) \right| + \mathcal{O}(\|\widetilde{\varepsilon}\|^2). \tag{7.615}$$

Here the sum runs over the projected breaking operators that couple within the ρ–block; the traceless Wigner–Eckart structure ensures that only the non-singlet parts contribute linearly.

Step 2: RG running of the explicit breakings. By Step 4 irrelevance, the linearized RG solution on the window satisfies (see Proposition 7.450 and Corollary 7.456)

$$\widetilde{\varepsilon}_{\tilde{B}}(\mu) = \mathcal{O}\Big(\mu^{\,y_{\tilde{B}}}\,\mathrm{poly}(\ln \mu)\Big), \qquad \mathrm{Re}\; y_{\tilde{B}} < 0,$$

so that

$$\sum_{\tilde{B}} \Big|\widetilde{\varepsilon}_{\tilde{B}}(\mu)\Big| = \mathcal{O}\Big(\mu^{\,\gamma_\star}\,\mathrm{poly}(\ln \mu)\Big), \qquad \gamma_\star := \min_{\tilde{B}} |\,\mathrm{Re}\; y_{\tilde{B}}| > 0. \tag{7.616}$$

Combining (7.615) with (7.616) and absorbing bounded C_ρ, C'_ρ into the $\mathcal{O}(\cdot)$ gives

$$\delta m_\rho^2(\ell) = \mathcal{O}\Big(\mu^{\,\gamma_\star}\,\mathrm{poly}(\ln \mu)\Big), \qquad \delta Z_\rho(\ell) = \mathcal{O}\Big(\mu^{\,\gamma_\star}\,\mathrm{poly}(\ln \mu)\Big). \tag{7.617}$$

Step 3: Division by the multiplet means. Kato continuity for pole data on the window (cf. Lemma 7.492) implies that $\overline{m_\rho^2}(\ell)$ and $\overline{Z_\rho}(\ell)$ are continuous and strictly positive (the latter by positivity of physical residues, cf. Definitions 7.482 and 7.490 and standard QFT conventions [91, 82]). We treat the two alternatives for $\overline{m_\rho^2}$:

(A) If $\overline{m_\rho^2}(\ell) \geq m_{\rho,\min}^2 > 0$ on the window, divide (7.617) by the constant lower bound to obtain

$$\frac{\delta m_\rho^2(\ell)}{\overline{m_\rho^2}(\ell)} = \mathcal{O}\Big(\mu^{\,\gamma_\star}\,\mathrm{poly}(\ln \mu)\Big) \;\xrightarrow[\mu \downarrow 0]{}\; 0,$$

and the same argument yields $\delta Z_\rho / \overline{Z_\rho} \to 0$ since $\overline{Z_\rho}$ is bounded below on compact IR windows.

(B) If $\overline{m_\rho^2}(\ell) \asymp \ell^{-\alpha}$ on the window, write $\mu = \kappa/\ell$ to get

$$\frac{\delta m_\rho^2(\ell)}{\overline{m_\rho^2}(\ell)} = \mathcal{O}\Big(\ell^{-(\gamma_\star - \alpha)}\,\mathrm{poly}(\ln \ell)\Big).$$

Under the stated condition $\alpha < \gamma_\star$, this tends to 0 as $\ell \uparrow \infty$ along the IR ray. The same reasoning applies to residues provided $\overline{Z_\rho}$ is bounded below or varies at most polynomially (which holds on the plateau by smooth tracking and positivity).

Step 4: Uniformity and window controls. Uniformity of the $\mathcal{O}(\cdot)$ bounds on compact subwindows follows from the curvature/TV guard and window consistency (cf. eq. (7.548) and Proposition 7.462), which control the norms entering C_ρ, C'_ρ and preclude loss of isolation for the simple poles in the cluster (cf. Lemma 7.492).

Together, these steps prove (7.614) with the quoted rates. □

Remark 7.495.

- The proof depends only on the following:
  1. the linear-response bound Proposition 7.493,
  2. RG decay of breaking amplitudes Proposition 7.450 and Corollary 7.456, and
  3. smooth spectral tracking Lemma 7.492.
- Standard QFT pole/residue conventions underlie positivity and normalization [91, 82].

Remark 7.496 (Massless multiplets and Goldstone sectors)**.** If $\overline{m_\rho^2}(\ell) \to 0$ (e.g. Nambu–Goldstone modes; cf. standard treatments [91, 82]), use a scale–free dispersion such as

$$\frac{\delta m_\rho^2(\ell)}{\max\{m_{\mathrm{ref}}^2,\ \overline{m_\rho^2}(\ell)\}}$$

with a fixed IR reference $m_{\mathrm{ref}} > 0$, or diagnose reconvergence via residue/velocity degeneracy together with band–projected Ward residuals (Definition 7.468). Under the hypotheses of Theorem 7.494 (Step 4 irrelevance and the IR guards), the linear–response bounds (7.613) imply in particular the decay of *absolute* splittings $\delta m_\rho^2(\ell) \to 0$ along the IR ray.

Discrete estimators and acceptance.

Definition 7.497 (Multiplet dispersion estimators)**.** Let $v_k := \ln \ell_k$ be a *uniform* grid with step $h > 0$. For each ρ (cf. Definition 7.491), define

$$\overline{m_\rho^2}(v_k) := \frac{1}{d_\rho} \sum_{i \in \mathcal{I}_\rho(v_k)} m_i^2(v_k), \qquad \delta m_\rho^2(v_k) := \max_{i \in \mathcal{I}_\rho(v_k)} \left| m_i^2(v_k) - \overline{m_\rho^2}(v_k) \right|.$$

When $\overline{m_\rho^2}(v_k) > 0$, set the relative dispersion

$$\Delta_{m,\rho}^{\mathrm{rel}}(v_k) := \frac{\delta m_\rho^2(v_k)}{\overline{m_\rho^2}(v_k)},$$

otherwise use the absolute version $\delta m_\rho^2(v_k)$ (cf. the massless/Goldstone remark Remark 7.496). Estimate the IR trend by the centered slope (as in Definition 7.487)

$$\widehat{\partial_v \ln \Delta_{m,\rho}^{\mathrm{rel}}}(v_k) := \frac{\ln \Delta_{m,\rho}^{\mathrm{rel}}(v_{k+1}) - \ln \Delta_{m,\rho}^{\mathrm{rel}}(v_{k-1})}{2h},$$

with the optional low–pass LP_τ in v whose kernel satisfies eqs. (7.537) and (7.538) (introducing an $\mathcal{O}(\tau^2)$ bias per Lemma 7.488). Define $\Delta_{Z,\rho}^{\mathrm{rel}}$ and $\widehat{\partial_v \ln \Delta_{Z,\rho}^{\mathrm{rel}}}$ analogously, replacing m_i^2 by Z_i.

Definition 7.498 (Step 6 acceptance: multiplet reconvergence)**.** On the IR window $\mathcal{I}_{\mathrm{IR}}$ (cf. eqs. (7.542) and (7.551)), accept the ρ–multiplet as reconverging if:

1. **Monotone reconvergence.** For all $v_k \in \mathcal{I}_{\mathrm{IR}}$,

$$\widehat{\partial_v \ln \Delta_{m,\rho}^{\mathrm{rel}}}(v_k) \le -c_\rho < 0, \qquad \widehat{\partial_v \ln \Delta_{Z,\rho}^{\mathrm{rel}}}(v_k) \le -c'_\rho < 0.$$

2. **Plateau smallness at restoration.** At $v_R = \ln \ell_R$ (cf. eq. (7.551)),

$$\Delta_{m,\rho}^{\mathrm{rel}}(v_R) \le \varepsilon_{\mathrm{plat}}, \qquad \Delta_{Z,\rho}^{\mathrm{rel}}(v_R) \le \varepsilon_{\mathrm{plat}}.$$

3. **Gauge and basis safety.** Masses/residues are evaluated on shell and Nielsen–projected (Definition 7.457), and spreads across admissible gauge choices lie below $\varepsilon_{\mathrm{gauge}}$ (cf. Corollary 7.477 and Proposition 7.485). Changes of channel basis act by similarity on $\Gamma_2^{\mathcal{N}}$ (cf. Proposition 7.458 and Lemma 7.483); the diagnostics $\Delta_{m,\rho}^{\mathrm{rel}}$ and $\Delta_{Z,\rho}^{\mathrm{rel}}$, being built from eigenvalue locations and scalar residues, are invariant under such conjugations.

Multiplet bookkeeping and splitting metrics.

Definition 7.499 (IR multiplet index sets and dispersion metrics)**.** Fix the band– and Nielsen–projected two–point kernel $\Gamma_2^{\mathcal{N}}(p^2;\lambda_{\mathrm{UV}},\ell)$ on the physical subspace $W = \operatorname{im}\mathcal{N}$ (cf. Definitions 7.457 and 7.490). For each irrep ρ of the emergent symmetry G with dimension d_ρ, choose a continuous family of projectors $P^\rho(\ell) : W \to W$ on the IR window $\{\ln \ell \in \mathcal{I}_{\mathrm{IR}}\}$ such that $\operatorname{rank} P^\rho(\ell) = d_\rho$ and $\sum_\rho P^\rho(\ell) = \mathbf{1}_W$ (block decomposition of the physical sector).[14]

Let $\{(m_i^2(\ell), Z_i(\ell))\}_i$ be the simple physical poles/residues of $\Gamma_2^{\mathcal{N}}$ at scale ℓ (cf. Definitions 7.482 and 7.490 and eq. (7.612)). Define the *ρ–multiplet index set* at ℓ as

$$\mathfrak{M}_\rho(\ell) \;:=\; \Big\{ i : \text{ the one–particle mode } i \text{ has } v_i(\ell) \in \operatorname{im} P^\rho(\ell) \text{ (i.e. it is selected by } P^\rho(\ell)) \Big\},$$

so that $|\mathfrak{M}_\rho(\ell)| = d_\rho$ for all ℓ in the window. The *normalised averages* and *max–deviation dispersions* for masses and residues are

$$\boxed{\overline{m_\rho^2}(\ell) := \frac{1}{|\mathfrak{M}_\rho(\ell)|}\sum_{i\in\mathfrak{M}_\rho(\ell)} m_i^2(\ell), \qquad \delta m_\rho^2(\ell) := \max_{i\in\mathfrak{M}_\rho(\ell)} \big| m_i^2(\ell) - \overline{m_\rho^2}(\ell)\big|,} \tag{7.618}$$

$$\boxed{\overline{Z_\rho}(\ell) := \frac{1}{|\mathfrak{M}_\rho(\ell)|}\sum_{i\in\mathfrak{M}_\rho(\ell)} Z_i(\ell), \qquad \delta Z_\rho(\ell) := \max_{i\in\mathfrak{M}_\rho(\ell)} \big| Z_i(\ell) - \overline{Z_\rho}(\ell)\big|.} \tag{7.619}$$

When needed, use the *relative dispersions* $\Delta_{m,\rho}^{\mathrm{rel}} := \delta m_\rho^2/\overline{m_\rho^2}$ (defined when $\overline{m_\rho^2} > 0$) and $\Delta_{Z,\rho}^{\mathrm{rel}} := \delta Z_\rho/\overline{Z_\rho}$, which are scale–free and compare directly across channels (cf. Definition 7.491).

Lemma 7.500 (Well–posedness, label invariance, and continuity)**.** *Assume each pole in a ρ–cluster is simple and that $P^\rho(\ell)$ varies continuously in ℓ on the IR window. Then:*

1. (Well–posed index sets) *$\mathfrak{M}_\rho(\ell)$ is well defined up to permutations of labels and has constant cardinality d_ρ.*
2. (Label/basis invariance) *The functionals $\overline{m_\rho^2}$, δm_ρ^2, $\overline{Z_\rho}$, δZ_ρ are invariant under reordering of indices and under unitary basis rotations inside* $\operatorname{im} P^\rho(\ell)$.
3. (Kato continuity) *Each $m_i^2(\ell)$ and $Z_i(\ell)$ depends continuously on ℓ along the window, hence so do* (7.618)–(7.619).

Moreover, if one introduces the pairwise spreads

$$\Delta_{m,\rho}^{\mathrm{pair}}(\ell) := \max_{i,j\in\mathfrak{M}_\rho(\ell)} |m_i^2(\ell) - m_j^2(\ell)|, \qquad \Delta_{Z,\rho}^{\mathrm{pair}}(\ell) := \max_{i,j\in\mathfrak{M}_\rho(\ell)} |Z_i(\ell) - Z_j(\ell)|,$$

then for all ℓ in the window

$$\delta m_\rho^2(\ell) \;\le\; \Delta_{m,\rho}^{\mathrm{pair}}(\ell) \;\le\; 2\,\delta m_\rho^2(\ell), \qquad \delta Z_\rho(\ell) \;\le\; \Delta_{Z,\rho}^{\mathrm{pair}}(\ell) \;\le\; 2\,\delta Z_\rho(\ell).$$

[14] One practical construction: at some large $\ell_{\mathrm{ref}} \in \mathcal{I}_{\mathrm{IR}}$, identify the ρ–cluster by quantum numbers; then transport $P^\rho(\ell)$ along ℓ by Kato parallel transport [115].

Proof. (1) Since $P^{\rho}(\ell)$ has constant rank d_ρ and varies continuously, its image defines a continuous d_ρ–dimensional subbundle of W along the window. By definition, $\mathfrak{M}_\rho(\ell)$ consists of those simple eigenmodes whose right eigenvectors $v_i(\ell)$ lie in $\operatorname{im} P^{\rho}(\ell)$; simplicity prevents branching ambiguities inside the cluster (cf. Lemma 7.492). Thus $|\mathfrak{M}_\rho(\ell)| = \operatorname{rank} P^{\rho}(\ell) = d_\rho$, and the set is unique up to relabelling.

(2) Reordering indices clearly leaves (7.618)–(7.619) unchanged. For basis changes, let $U(\ell)$ be unitary on $\operatorname{im} P^{\rho}(\ell)$; the block of $\Gamma_2^{\mathcal{N}}$ transforms by similarity, $A_\rho \mapsto U^{*} A_\rho U$ (cf. Proposition 7.458 and Lemma 7.483). Eigenvalues $\{m_i^2(\ell)\}_i$ and scalar residues $\{Z_i(\ell)\}_i$ are invariant under similarity (see Definitions 7.482 and 7.490), hence so are the averages and deviations.

(3) Set $A(p^2,\ell) := \Gamma_2^{\mathcal{N}}(p^2;\lambda_{\mathrm{UV}},\ell)$; by hypothesis and the IR guards, A is C^1 in (p^2,ℓ) on the admitted band. For a simple pole $p^2 = m_i^2(\ell)$ the Implicit Function Theorem/Kato perturbation theory gives continuous (indeed C^1) dependence of $m_i^2(\ell)$ and of the corresponding eigenprojectors/eigenvectors on ℓ; the residue formula

$$Z_i(\ell) = \Big(w_i(\ell)^{\top}\, \partial_{p^2} A\big(m_i^2(\ell),\ell\big)\, v_i(\ell) \Big)^{-1} \qquad \text{(cf. Definition 7.490 and eq. (7.612))}$$

is thus continuous as well (the denominator stays nonzero at simple poles) [115, Chs. II–IV]. Since $\overline{(\cdot)}$ and max are continuous on finite tuples, the claimed continuity follows.

For the pairwise bounds, observe for any finite set $\{x_i\}$ with mean $\bar{x}$ that

$$\max_{i,j} |x_i - x_j| \;\le\; \max_i |x_i - \bar{x}| + \max_j |\bar{x} - x_j| \;=\; 2 \max_k |x_k - \bar{x}|,$$

and also $\max_{i,j} |x_i - x_j| \ge \max_i |x_i - \bar{x}|$ because $\bar{x}$ lies between $\min x_i$ and $\max x_i$. Apply this to $\{m_i^2(\ell)\}_{i\in\mathfrak{M}_\rho(\ell)}$ and to $\{Z_i(\ell)\}_{i\in\mathfrak{M}_\rho(\ell)}$ to obtain the stated inequalities. □

Lemma 7.501 (Equivalence of dispersion metrics)**.** *Let $x_i \in \mathbb{R}$ $(i = 1,\dots,d)$ and set $R := \max_i |x_i - \overline{x}|$ (max–deviation from the mean) and $D := \max_{i,j} |x_i - x_j|$ (pairwise spread). Then*

$$R \;\le\; D \;\le\; 2R.$$

Hence $D \to 0$ iff $R \to 0$. Applied to $\{m_i^2(\ell)\}_{i\in\mathfrak{M}_\rho(\ell)}$ or $\{Z_i(\ell)\}_{i\in\mathfrak{M}_\rho(\ell)}$ this shows that pairwise–spread and mean–deviation give equivalent notions of multiplet reconvergence.

Proof. For any fixed i, by the triangle inequality,

$$|x_i - \overline{x}| \le \max_j |x_i - x_j| \le D,$$

and taking the maximum over i gives $R \le D$. For the reverse bound, choose indices $i^{\star}, j^{\star}$ with $|x_{i^{\star}} - x_{j^{\star}}| = D$. Then

$$D = |x_{i^{\star}} - x_{j^{\star}}| \le |x_{i^{\star}} - \overline{x}| + |x_{j^{\star}} - \overline{x}| \le 2R,$$

so $D \le 2R$. The equivalence $D \to 0 \Leftrightarrow R \to 0$ follows immediately. □

Remark 7.502 (Gauge and scheme safety)**.** Masses and residues are extracted on shell and within $W = \operatorname{im}\mathcal{N}$ (Nielsen projection), hence are gauge–parameter independent within the audit tolerance eq. (7.588) (cf. Proposition 7.485 and Corollary 7.477). Moreover, admissible basis/scheme changes act by similarity within W and do not affect $\{m_i^2, Z_i\}$ (or their dispersions) *as numbers*; see Proposition 7.458 and Lemmas 7.451 and 7.483.

Corollary 7.503 (Continuity of relative dispersions)**.** *Let ρ be fixed and work on the IR window. If $\overline{m_\rho^2}(\ell)$ (resp. $\overline{Z_\rho}(\ell)$) is bounded away from 0 on the window, then $\Delta^{\mathrm{rel}}_{m,\rho}(\ell) := \delta m_\rho^2(\ell)/\overline{m_\rho^2}(\ell)$ (resp. $\Delta^{\mathrm{rel}}_{Z,\rho}(\ell) := \delta Z_\rho(\ell)/\overline{Z_\rho}(\ell)$), as defined in Definition 7.499, is continuous in ℓ. Moreover,*

$$\Delta^{\mathrm{rel}}_{m,\rho}(\ell) \to 0 \quad \Longleftrightarrow \quad \delta m_\rho^2(\ell) \to 0,$$

and the analogous equivalence holds for residues.

Proof. By Lemma 7.500 (Kato continuity), each simple pole $m_i^2(\ell)$ and residue $Z_i(\ell)$ depends continuously on ℓ along the window; hence the functionals

$$\overline{m_\rho^2}(\ell),\ \delta m_\rho^2(\ell) \quad \text{and} \quad \overline{Z_\rho}(\ell),\ \delta Z_\rho(\ell)$$

defined in (7.618)–(7.619) are continuous (finite sums/means and maxima of continuous functions are continuous). See also [115, Chs. II–IV] for the underlying spectral continuity.

If $\overline{m_\rho^2}(\ell) \geq c_m > 0$ on the window, then the quotient

$$\Delta^{\mathrm{rel}}_{m,\rho}(\ell) = \frac{\delta m_\rho^2(\ell)}{\overline{m_\rho^2}(\ell)}$$

is the ratio of continuous functions with denominator bounded away from 0, hence is continuous. Furthermore,

$$\Delta^{\mathrm{rel}}_{m,\rho}(\ell) \to 0 \iff \delta m_\rho^2(\ell) \to 0,$$

since $c_m \leq \overline{m_\rho^2}(\ell) \leq C_m$ on compact subwindows implies $\frac{1}{C_m}\,\delta m_\rho^2(\ell) \leq \Delta^{\mathrm{rel}}_{m,\rho}(\ell) \leq \frac{1}{c_m}\,\delta m_\rho^2(\ell)$. The residue statement is identical when $\overline{Z_\rho}(\ell) \geq c_Z > 0$ on the window.

Finally, by Lemma 7.501, the choice between pairwise spread and mean–deviation is immaterial for diagnosing reconvergence; the present result concerns the (scale–free) ratio built from the mean–deviation. This completes the proof. □

The projector $P^\rho(\ell)$ organizes states into would–be symmetric multiplets (Definition 7.499); the dispersion $\delta(\cdot)$ measures intra-multiplet lifting of degeneracy, while the relative version removes overall scaling. Kato continuity ensures smooth tracking of pole/residue data (Lemma 7.492, [115]); Nielsen projection keeps gauge artifacts out (Definition 7.457), and gauge safety is guaranteed on shell (Proposition 7.485 and Corollary 7.477).

IR plateau acceptance (spectral).

Definition 7.504 (Spectral IR–plateau acceptance)**.** Let $\mathcal{I}_{\mathrm{IR}} \subset \mathbb{R}$ be the Step 3 IR window in $v = \ln \ell$ and let ℓ_R be the restoration length (cf. Definitions 7.430 and 7.478). For each multiplet ρ, with dispersions defined in Definition 7.499, we *accept* the spectral plateau if the relative dispersions are monotone nonincreasing on $\mathcal{I}_{\mathrm{IR}}$ and small at the restoration point:

$$\boxed{\frac{\mathrm{d}}{\mathrm{d}\ln\ell}\left(\frac{\delta m_\rho^2(\ell)}{\overline{m_\rho^2}(\ell)}\right) \leq 0, \qquad \frac{\mathrm{d}}{\mathrm{d}\ln\ell}\left(\frac{\delta Z_\rho(\ell)}{\overline{Z_\rho}(\ell)}\right) \leq 0, \qquad \max\left\{\frac{\delta m_\rho^2(\ell_R)}{\overline{m_\rho^2}(\ell_R)}, \frac{\delta Z_\rho(\ell_R)}{\overline{Z_\rho}(\ell_R)}\right\} \leq \varepsilon_{\mathrm{spec}}.}$$

(7.620)

Here $\varepsilon_{\mathrm{spec}} \le \varepsilon_{\mathrm{plat}}$ is a fixed tolerance, and the monotonicity inequalities are evaluated with the same ripple guards as in eq. (7.548) (optionally after a low–pass in v obeying eqs. (7.537) and (7.538)). When $\overline{m_\rho^2}$ (or $\overline{Z_\rho}$) approaches zero, replace the ratio by the absolute dispersion or use a fixed IR reference in the denominator (cf. Remark 7.496).

Lemma 7.505 (Equivalent log–slope form; invariances)**.** *Let* $\Delta^{\mathrm{rel}}_{m,\rho} := \delta m_\rho^2/\overline{m_\rho^2}$ *and* $\Delta^{\mathrm{rel}}_{Z,\rho} := \delta Z_\rho/\overline{Z_\rho}$ *on the subset where denominators are* > 0*. Then* (7.620) *is equivalent to*

$$\frac{\mathrm{d}}{\mathrm{d}\ln\ell}\,\ln\Delta^{\mathrm{rel}}_{m,\rho}(\ell) \;\le\; 0, \qquad \frac{\mathrm{d}}{\mathrm{d}\ln\ell}\,\ln\Delta^{\mathrm{rel}}_{Z,\rho}(\ell) \;\le\; 0.$$

Moreover, the conditions in (7.620) *are invariant under*

(i) rigid scale reparametrisations $\ell \mapsto a\,\ell$ *(i.e.* $v \mapsto v + \ln a$*), and*
(ii) unitary rotations within the ρ*–block (basis changes in* $\operatorname{im} P^\rho$*).*

Proof. Equivalence of slope and log–slope. On the subset where denominators are > 0, the relative dispersions $\Delta^{\mathrm{rel}}_{m,\rho}$ and $\Delta^{\mathrm{rel}}_{Z,\rho}$ are strictly positive. For a positive C^1 function Δ,

$$\frac{\mathrm{d}}{\mathrm{d}\ln\ell}\,\ln\Delta(\ell) = \frac{1}{\Delta(\ell)}\frac{\mathrm{d}}{\mathrm{d}\ln\ell}\,\Delta(\ell),$$

so the sign of the derivative is preserved: $\frac{\mathrm{d}}{\mathrm{d}\ln\ell}\Delta \le 0$ iff $\frac{\mathrm{d}}{\mathrm{d}\ln\ell}\ln\Delta \le 0$. Applying this to $\Delta^{\mathrm{rel}}_{m,\rho}$ and $\Delta^{\mathrm{rel}}_{Z,\rho}$ yields the claimed equivalence.

Invariance under $\ell \mapsto a\,\ell$. The change $\ell \mapsto a\,\ell$ corresponds to a rigid shift $v = \ln\ell \mapsto v + \ln a$. Derivatives with respect to v (or $\ln\ell$) are invariant under rigid shifts, hence the monotonicity inequalities are unchanged pointwise along the (shifted) window. The threshold condition at the restoration point remains the same because it is evaluated at ℓ_R (which itself transforms as $\ell_R \mapsto a\,\ell_R$ under a mere change of units), so the numerical values of the ratios at restoration are unchanged.

Invariance under block–unitary basis changes. Let $U(\ell)$ be unitary on $\operatorname{im} P^\rho(\ell)$. The ρ–block of the projected kernel transforms by similarity:

$$A_\rho(p^2;\ell) \;\mapsto\; U(\ell)^* A_\rho(p^2;\ell)\,U(\ell) \quad \text{with} \quad A_\rho := P^\rho\,\Gamma_2^{\mathcal{N}}\,P^\rho\Big|_{\operatorname{ran} P^\rho}$$

(cf. Proposition 7.458 and Lemma 7.483). Eigenvalue locations $\{m_i^2(\ell)\}$ and scalar residues $\{Z_i(\ell)\}$ are invariant under similarity (see Definitions 7.482 and 7.490); therefore the averages and deviations (7.618)–(7.619) are basis–invariant, and so are their ratios. Since the invariants are unchanged pointwise in ℓ, their v–derivatives are also unchanged. This proves the claimed invariances.

Finally, continuity of the ratios on the subdomain with positive denominators follows from Lemma 7.500 (Kato continuity of m_i^2, Z_i and continuity of the mean and max functionals), ensuring that the derivatives above are well defined after applying the Step 3 ripple guards eq. (7.548) and, if used, the low–pass operators obeying eqs. (7.537) and (7.538). □

Proposition 7.506 (Sufficient conditions from irrelevance)**.** *Assume the Step 4 irrelevance hypothesis with* $\gamma_\star := \min_{\tilde B} |\operatorname{Re} y_{\tilde B}| > 0$ *(cf. Proposition 7.450 and Corollary 7.456) and the*

regularity/ripple guards on the IR window (cf. eq. (7.548) and Proposition 7.462). Suppose, in addition, that the multiplet mean obeys either

$$\text{(A)}\quad \overline{m_\rho^2}(\ell) \geq m_{\rho,\min}^2 > 0 \qquad \text{or} \qquad \text{(B)}\quad \overline{m_\rho^2}(\ell) \asymp \ell^{-\alpha} \quad \text{on the window with} \quad \alpha \leq \gamma_\star/2.$$

Then there exists $\ell_\star \geq \ell_R$ such that for all $\ell \geq \ell_\star$,

$$\frac{\mathrm{d}}{\mathrm{d}\ln\ell}\ln\Delta_{m,\rho}^{\mathrm{rel}}(\ell) \;\leq\; -\frac{\gamma_\star}{2} \;<\; 0, \qquad \frac{\mathrm{d}}{\mathrm{d}\ln\ell}\ln\Delta_{Z,\rho}^{\mathrm{rel}}(\ell) \;\leq\; -\frac{\gamma_\star}{2} \;<\; 0.$$

Consequently, the monotone part of the spectral acceptance criterion (7.620) holds on $[\ell_\star,\infty)$. If, in addition, the plateau smallness at ℓ_R holds with the tolerance $\varepsilon_{\mathrm{spec}}$ (cf. Definition 7.504), then the spectral IR–plateau is accepted.

Proof. All statements are made on the BRST–physical/Nielsen subspace $W = \operatorname{im}\mathcal{N}$ for the band–projected kernel $\Gamma_2^{\mathcal{N}}$ (cf. Definitions 7.457 and 7.490); gauge safety on shell holds by Proposition 7.485 and Corollary 7.477.

Step 1: Linear response for intra–multiplet splittings. From Proposition 7.493 there exist bounded continuous functions $C_\rho(\ell), C'_\rho(\ell)$ on the IR window such that

$$\delta m_\rho^2(\ell) \;\leq\; C_\rho(\ell)\sum_{\tilde B}\left|\widetilde{\varepsilon}_{\tilde B}(\kappa/\ell)\right| + \mathcal{O}(\|\widetilde{\varepsilon}\|^2), \qquad \delta Z_\rho(\ell) \;\leq\; C'_\rho(\ell)\sum_{\tilde B}\left|\widetilde{\varepsilon}_{\tilde B}(\kappa/\ell)\right| + \mathcal{O}(\|\widetilde{\varepsilon}\|^2). \tag{7.621}$$

Step 2: RG decay and an eventual differential inequality. Let $\mu = \kappa/\ell$. By Step 4 irrelevance (Proposition 7.450 and Corollary 7.456),

$$\widetilde{\varepsilon}_{\tilde B}(\mu) = \mathcal{O}\Big(\mu^{\,y_{\tilde B}}\,\mathrm{poly}(\ln\mu)\Big), \quad \operatorname{Re} y_{\tilde B} < 0.$$

Writing $S(\ell) := \sum_{\tilde B}\left|\widetilde{\varepsilon}_{\tilde B}(\kappa/\ell)\right|$, one has

$$\frac{\mathrm{d}}{\mathrm{d}\ln\ell}\ln S(\ell) = -\frac{\sum_{\tilde B}\operatorname{Re} y_{\tilde B}\,|\widetilde{\varepsilon}_{\tilde B}| + \mathcal{O}(\|\widetilde{\varepsilon}\|^2)}{\sum_{\tilde B}|\widetilde{\varepsilon}_{\tilde B}|} = -\gamma_\star + o(1) \quad \text{as } \ell\to\infty,$$

since the slowest irrelevant sector dominates the sum (polylog factors are slowly varying). Hence there exists ℓ_1 such that for all $\ell \geq \ell_1$,

$$\frac{\mathrm{d}}{\mathrm{d}\ln\ell}\ln S(\ell) \;\leq\; -\frac{3}{4}\gamma_\star. \tag{7.622}$$

Step 3: Log–slope for the relative mass dispersion. Consider

$$\ln\Delta_{m,\rho}^{\mathrm{rel}}(\ell) = \ln\delta m_\rho^2(\ell) - \ln\overline{m_\rho^2}(\ell).$$

By (7.621) and smallness of $\|\widetilde{\varepsilon}\|$, there exists ℓ_2 s.t. for $\ell \geq \ell_2$, $\delta m_\rho^2(\ell) = \Theta\Big(S(\ell)\Big)$ up to a slowly varying multiplicative factor $R_m(\ell)$ with $\ln R_m(\ell)$ bounded and $[\mathrm{d}/\mathrm{d}\ln\ell]\ln R_m(\ell) = o(1)$. (This follows from the first–order degenerate perturbation structure inside the finite ρ–cluster and the smooth tracking of projectors/eigenvectors, cf. Lemma 7.492 and the Wigner–Eckart decomposition in Proposition 7.493.) Therefore,

$$\frac{\mathrm{d}}{\mathrm{d}\ln\ell}\ln\delta m_\rho^2(\ell) = \frac{\mathrm{d}}{\mathrm{d}\ln\ell}\ln S(\ell) + o(1) \;\leq\; -\frac{3}{4}\gamma_\star + o(1),$$

by (7.622). For the denominator:

$$\frac{\mathrm{d}}{\mathrm{d}\ln\ell}\,\ln\overline{m_\rho^2}(\ell) = \begin{cases} o(1), & \text{in case (A)},\\ -\alpha + o(1), & \text{in case (B)}, \end{cases}$$

using the hypothesis on $\overline{m_\rho^2}$ and continuity on the window (Lemma 7.500). Consequently,

$$\frac{\mathrm{d}}{\mathrm{d}\ln\ell}\,\ln\Delta^{\mathrm{rel}}_{m,\rho}(\ell) = \begin{cases} -\frac{3}{4}\gamma_\star + o(1), & \text{in case (A)},\\ -(\frac{3}{4}\gamma_\star - \alpha) + o(1), & \text{in case (B)}. \end{cases}$$

Choose $\ell_\star \geq \max\{\ell_1, \ell_2, \ell_R\}$ large enough that the $o(1)$ term is bounded by $\gamma_\star/4$ and, in case (B), use $\alpha \leq \gamma_\star/2$. Then for all $\ell \geq \ell_\star$,

$$\frac{\mathrm{d}}{\mathrm{d}\ln\ell}\,\ln\Delta^{\mathrm{rel}}_{m,\rho}(\ell) \;\leq\; -\frac{\gamma_\star}{2} \;<\; 0.$$

Step 4: Residues. The same argument applies to $\Delta^{\mathrm{rel}}_{Z,\rho}(\ell)$: by (7.621) and the residue–tracking formula in Definition 7.490 together with Lemma 7.492, the first–order variation is linear in the breakings with the same $S(\ell)$ control, yielding the identical log–slope bound.

Step 5: Acceptance. The obtained bounds give the monotone (log–slope) part of Definition 7.504 on $[\ell_\star, \infty)$. If, in addition, the restoration–point smallness at ℓ_R holds with $\varepsilon_{\mathrm{spec}}$, the full criterion (7.620) is satisfied and the spectral IR–plateau is accepted.

This completes the proof. □

- Step 4 irrelevance and its $\mu^{y_{\bar{B}}}\,\mathrm{poly}(\ln\mu)$ law are taken from Proposition 7.450 and Corollary 7.456 (RG canon [389, 91, 82]).
- Smooth spectral tracking and the stability of the finite ρ–cluster rely on Kato's theory [115] via Lemma 7.492.
- The additional assumption on $\overline{m_\rho^2}$ is the same scaling control used in Theorem 7.494; it is sharpened here to ensure a uniform $-\gamma_\star/2$ log–slope bound.

Discrete audit.

Definition 7.507 (Discrete spectral monotonicity checks)**.** Let $v_k := \ln\ell_k$ be a *uniform* grid with step $h > 0$. Define the centered finite–difference estimators (cf. Definition 7.487 and Lemma 7.488)

$$\widehat{\partial_v \Delta^{\mathrm{rel}}_{m,\rho}}(v_k) := \frac{\Delta^{\mathrm{rel}}_{m,\rho}(v_{k+1}) - \Delta^{\mathrm{rel}}_{m,\rho}(v_{k-1})}{2h},$$
$$\widehat{\partial_v \ln\Delta^{\mathrm{rel}}_{m,\rho}}(v_k) := \frac{\ln\Delta^{\mathrm{rel}}_{m,\rho}(v_{k+1}) - \ln\Delta^{\mathrm{rel}}_{m,\rho}(v_{k-1})}{2h},$$

and analogously for $\Delta^{\mathrm{rel}}_{Z,\rho}$, on the subset where the denominators in the definitions of the relative dispersions are strictly positive (cf. Definition 7.499 and Lemma 7.501). With the

optional low–pass operator LP_τ in v (kernels obeying eqs. (7.537) and (7.538), inducing an $\mathcal{O}(\tau^2)$ bias by Lemma 7.488), *accept* monotone reconvergence on the discrete IR window if

$$\widehat{\partial_v \Delta^{\mathrm{rel}}_{m,\rho}}(v_k) \le 0 \quad \text{and} \quad \widehat{\partial_v \Delta^{\mathrm{rel}}_{Z,\rho}}(v_k) \le 0 \quad \text{for all } v_k \in \mathcal{I}_{\mathrm{IR}},$$

and the restoration–point smallness holds:

$$\max\Big\{\Delta^{\mathrm{rel}}_{m,\rho}(\ln \ell_R),\ \Delta^{\mathrm{rel}}_{Z,\rho}(\ln \ell_R)\Big\} \ \le\ \varepsilon_{\mathrm{spec}} \quad \text{(cf. Definition 7.504)}.$$

When $\overline{m^2_\rho}$ or $\overline{Z_\rho}$ is near zero, replace the log–slope check by the slope of the absolute dispersion or use a fixed IR reference in the denominator (cf. Remark 7.496).

Intuition. Relative dispersions should *tighten* (be nonincreasing in $v = \ln \ell$) and already be small at restoration. Step 4 irrelevance provides an eventual negative log–slope (cf. Theorem 7.494 and Proposition 7.506); the ripple guards and the centered estimators (Lemma 7.488) ensure we detect genuine reconvergence rather than numerical wiggles.

Pseudo–NG option and RG consistency.

Definition 7.508 (Pseudo–Nambu–Goldstone (pNG) channel and spurion power)**.** Let $\widetilde{\mathcal{O}}_{\tilde{B}}$ be a Nielsen–projected breaking eigen–operator with irrelevant exponent $y_{\tilde{B}} < 0$ (Step 7.8.4; cf. Proposition 7.450 and Corollary 7.456). A *pNG channel* is a physical mode that would be exactly massless in the symmetric theory but acquires a small gap from explicit breaking by $\widetilde{\mathcal{O}}_{\tilde{B}}$. By spurion analysis/Wigner–Eckart (the spurion transforming like $\widetilde{\mathcal{O}}_{\tilde{B}}$), the leading pNG mass term arises from the minimal number $n_\star \in \mathbb{N}$ of spurion insertions required to form a singlet in the pNG channel; we call

$$\alpha := n_\star > 0$$

the *spurion power* (model/channel dependent). Consequently, the leading pNG scaling is governed by the $n_\star$–fold products of the running breakings,

$$\begin{aligned}\overline{m^2_{\mathrm{pNG}}}(\ell) &= \mathcal{O}\Big(\sum_{(\tilde{B}_1,\dots,\tilde{B}_{n_\star})} \prod_{j=1}^{n_\star} \widetilde{\varepsilon}_{\tilde{B}_j}(\kappa/\ell)\Big) \\ &= \mathcal{O}\Big(\mu^{\,n_\star \gamma_\star}\,\mathrm{poly}(\ln \mu)\Big), \\ &\qquad \text{where } \mu = \kappa/\ell \text{ and } \gamma_\star := \min_{\tilde{B}} |\operatorname{Re} y_{\tilde{B}}|.\end{aligned}$$

in agreement with the RG running and linear–response structure (cf. Proposition 7.493 and Theorem 7.494). Standard QFT treatments of NG/pNG modes and spurion methods include [91, 82].

Proposition 7.509 (pNG scaling from the RG solution)**.** *Work on the on–shell, band– and Nielsen–projected posture for the two–point kernel $\Gamma_2^{\mathcal{N}}(p^2; \lambda_{\mathrm{UV}}, \ell)$ on $W = \operatorname{im}\mathcal{N}$ (cf. Definitions 7.457 and 7.490). Let $\widetilde{\mathcal{O}}_{\tilde{B}}$ be a Nielsen–projected breaking eigen–operator with irrelevant exponent $y_{\tilde{B}} < 0$ (Step 7.8.4; cf. Proposition 7.450 and Corollary 7.456) and let $\alpha = n_\star > 0$*

be the spurion power for the pNG channel (cf. Definition 7.508). Then on the IR window $\mathcal{I}_{\mathrm{IR}}$ the pNG mass obeys the scaling law

$$\boxed{m_{\mathrm{pNG}}^2(\ell) \;=\; C_{\mathrm{pNG}}(\ell)\,\big|\widetilde{\varepsilon}_{\tilde{B}}(\mu)\big|^{\alpha} \;\sim\; C_{\mathrm{pNG}}(\ell)\,\mu^{\alpha y_{\tilde{B}}} \;=\; C_{\mathrm{pNG}}(\ell)\left(\frac{\kappa}{\ell}\right)^{\alpha y_{\tilde{B}}}, \qquad \mu=\kappa/\ell,}\tag{7.623}$$

where $C_{\mathrm{pNG}}(\ell)$ is slowly varying on the plateau (bounded and ℓ–Lipschitz under the curvature/TV guards; cf. eq. (7.548) and Proposition 7.462). Consequently,

$$\frac{\mathrm{d}}{\mathrm{d}\ln\mu}\,\ln m_{\mathrm{pNG}}^2(\ell) \;=\; \alpha\, y_{\tilde{B}} \;+\; o(1) \quad \text{as } \mu\downarrow 0.$$

Proof. All statements are made on $W=\operatorname{im}\mathcal{N}$; gauge–parameter safety on shell follows from Proposition 7.485 and Corollary 7.477. Let $P^{\mathrm{pNG}}(\ell)$ denote the (rank–one) projector selecting the pNG channel within the relevant multiplet (cf. Definitions 7.499 and 7.508). Consider the reduced scalar kernel obtained by left/right projection onto the pNG mode at scale ℓ:

$$a_\ell(p^2) \;:=\; w_{\mathrm{pNG}}(\ell)^{\top}\,\Gamma_2^{\mathcal{N}}(p^2;\lambda_{\mathrm{UV}},\ell)\,v_{\mathrm{pNG}}(\ell),$$

with $w_{\mathrm{pNG}}^{\top}v_{\mathrm{pNG}}=1$ and $v_{\mathrm{pNG}}(\ell)\in\operatorname{ran}P^{\mathrm{pNG}}(\ell)$ (cf. residue normalisation in Definition 7.490). In the exact symmetric limit (spurion switched off) the pNG mode is massless, so $a_\ell(0)=0$ and the pole at $p^2=0$ is simple with finite residue; equivalently $\partial_{p^2}a_\ell(0)=Z_{\mathrm{sym}}(\ell)^{-1}\neq 0$ by standard QFT pole–residue practice [91, 82].

Turning on the explicit breaking by $\widetilde{\mathcal{O}}_{\tilde{B}}$, spurion/Wigner–Eckart selection rules imply that the first symmetry–allowed invariant in the pNG channel arises at order $n_\star=\alpha$ in the spurion field (cf. Definition 7.508); lower orders vanish by representation constraints. Hence, for small breaking and p^2 near 0,

$$a_\ell(p^2)=\underbrace{\partial_{p^2}a_\ell(0)}_{Z_{\mathrm{sym}}(\ell)^{-1}}\,p^2 \;-\; \sum_{n\geq\alpha} c_n(\ell)\,\big(\widetilde{\varepsilon}_{\tilde{B}}(\mu)\big)^n \;+\; \mathcal{O}\big(p^4\big) \;+\; \mathcal{O}\big(\|\widetilde{\varepsilon}\|^{\alpha+1}\big),$$

where the coefficients $c_n(\ell)$ are bounded and ℓ–Lipschitz on the IR window by the curvature/TV guards and the smooth tracking of eigenvectors/projectors (Kato continuity; cf. Lemma 7.492 and [115, Chs. II–IV]). The pNG mass solves the pole condition $a_\ell(m_{\mathrm{pNG}}^2(\ell))=0$, giving at leading nontrivial order

$$m_{\mathrm{pNG}}^2(\ell)=\; Z_{\mathrm{sym}}(\ell)\,c_\alpha(\ell)\,\big|\widetilde{\varepsilon}_{\tilde{B}}(\mu)\big|^{\alpha}\,\Big(1+\mathcal{O}\big(|\widetilde{\varepsilon}_{\tilde{B}}|\big)\Big).$$

Absorb $Z_{\mathrm{sym}}(\ell)c_\alpha(\ell)$ and the slowly varying subleading factors (including possible polylogarithms from the RG) into $C_{\mathrm{pNG}}(\ell)$; boundedness and ℓ–Lipschitz continuity of C_{pNG} follow from the IR window guards and from the smoothness of the coefficients above.

Finally, Step 4 irrelevance provides the RG running $\widetilde{\varepsilon}_{\tilde{B}}(\mu)=\mathcal{O}\big(\mu^{\,y_{\tilde{B}}}\,\mathrm{poly}(\ln\mu)\big)$ with $y_{\tilde{B}}<0$ (cf. Proposition 7.450 and Corollary 7.456). The slowly varying polylog factor is again absorbed into $C_{\mathrm{pNG}}(\ell)$ on the plateau, yielding (7.623). Differentiating $\ln m_{\mathrm{pNG}}^2(\ell)=\ln C_{\mathrm{pNG}}(\ell)+\alpha\ln|\widetilde{\varepsilon}_{\tilde{B}}(\mu)|$ with respect to $\ln\mu$ and using that $\frac{\mathrm{d}}{\mathrm{d}\ln\mu}\ln C_{\mathrm{pNG}}(\ell)=o(1)$ on the IR window (slow variation) gives $\frac{\mathrm{d}}{\mathrm{d}\ln\mu}\ln m_{\mathrm{pNG}}^2(\ell)=\alpha\,y_{\tilde{B}}+o(1)$ as $\mu\downarrow 0$. □

Remark 7.510 (Multiple breakings and crossovers)**.** If several irrelevant spurions $\{\widetilde{\mathcal{O}}_{\tilde{B}_j}\}$ contribute with powers α_j and exponents $y_j < 0$, then by the same spurion analysis and RG running one obtains

$$m^2_{\mathrm{pNG}}(\ell) \;\sim\; \sum_j C_j(\ell)\, \mu^{\alpha_j y_j},$$

with slowly varying $C_j(\ell)$. As $\mu \downarrow 0$, the *slowest* decaying contribution (largest $\alpha_j y_j$ in the sense of least negative) dominates the log–slope; on finite IR windows this can produce apparent crossovers before the asymptotic regime is reached (cf. Proposition 7.450 and Corollary 7.456 and standard pNG/spurion treatments [91, 82]).

Estimators and RG consistency checks.

Definition 7.511 (Log–slope estimators for pNG scaling)**.** Let $t_k := \ln \mu_k$ be a *uniform* grid with step $\Delta t > 0$ on the IR window (so $\mu_k = \kappa/\ell_k$ and $t_k = \ln \kappa - \ln \ell_k$). Define the centered log–slope estimator for the pNG mass (cf. Lemma 7.488):

$$\widehat{s}_{\mathrm{pNG}}(\mu_k) := \frac{\ln m^2_{\mathrm{pNG}}(\mu_{k+1}) - \ln m^2_{\mathrm{pNG}}(\mu_{k-1})}{2\,(t_{k+1} - t_{k-1})} = \frac{\ln m^2_{\mathrm{pNG}}(\mu_{k+1}) - \ln m^2_{\mathrm{pNG}}(\mu_{k-1})}{2\,\Delta t}.$$

To estimate the spurion power α appearing in the scaling law (7.623) of Proposition 7.509, set

$$\widehat{\alpha}_{\tilde{B}}(\mu_k) := \frac{\widehat{s}_{\mathrm{pNG}}(\mu_k)}{\widehat{y}_{\tilde{B}}(\mu_k)},$$

where $\widehat{y}_{\tilde{B}}$ is the Step 4 slope estimator for the breaking eigen–coupling (cf. the centered–difference construction in Definition 7.446). Optionally apply a low–pass in t with kernel obeying eqs. (7.537) and (7.538), which induces an $\mathcal{O}(\tau^2)$ bias by Lemma 7.488. Consistency with Proposition 7.509 requires $\widehat{s}_{\mathrm{pNG}}(\mu_k) \to \alpha\, y_{\tilde{B}}$ and hence $\widehat{\alpha}_{\tilde{B}}(\mu_k) \to \alpha$ along the IR window.

Proposition 7.512 (Gauge/basis safety and scale invariance)**.** *The quantities m^2_{pNG}, $\widehat{s}_{\mathrm{pNG}}$, and $\widehat{\alpha}_{\tilde{B}}$ are:*

(i) gauge–parameter independent within the tolerance eq. (7.588) when evaluated on shell and Nielsen–projected (Definition 7.457);

(ii) invariant under admissible basis/scheme changes on $W = \mathrm{im}\,\mathcal{N}$ (similarity in the ρ–block; cf. Proposition 7.458 and Lemmas 7.451 and 7.483);

(iii) independent of the audit constant κ in $\mu = \kappa/\ell$ (a rigid shift of $t = \ln \mu$ cancels in centered differences).

Proof. (i) *Gauge safety.* On the on–shell, Nielsen–projected posture, the two–point kernel's pole positions are gauge–parameter independent (within the audit tolerance) by Proposition 7.485 and Corollary 7.477; the pNG mass m^2_{pNG} is such a pole location in the selected channel. Therefore $\ln m^2_{\mathrm{pNG}}$ and its centered finite differences inherit the same gauge independence; dividing by $\widehat{y}_{\tilde{B}}$ (which is itself gauge–safe under the same posture by Step 4's construction) preserves gauge safety for $\widehat{\alpha}_{\tilde{B}}$.

(ii) *Basis/scheme invariance.* Admissible basis/scheme changes act by similarity on the block $A_\rho = P^\rho \Gamma_2^{\mathcal{N}} P^\rho$ (cf. Proposition 7.458 and Lemma 7.483). Pole locations and scalar

residues are invariant under similarity (see Definitions 7.482 and 7.490), hence m^2_{pNG} is invariant as a number. Centered log–differences of an invariant scalar are invariant, and the Step 4 estimator $\widehat{y}_{\tilde{B}}$ is constructed to be scheme–stable (cf. Lemma 7.451), so $\widehat{s}_{\mathrm{pNG}}$ and $\widehat{\alpha}_{\tilde{B}}$ are basis/scheme invariant.

(iii) *Independence of* κ. Changing $\kappa \mapsto a\,\kappa$ shifts $t = \ln \mu$ by a constant: $t \mapsto t + \ln a$. For any function g, the centered difference $\big[g(t_{k+1}) - g(t_{k-1})\big] / \big[2\,(t_{k+1} - t_{k-1})\big]$ is invariant under rigid shifts of t. Hence $\widehat{s}_{\mathrm{pNG}}$ is independent of κ, and so is $\widehat{\alpha}_{\tilde{B}}$.

Finally, optional low–pass filtering in t preserves these invariances and introduces only the controlled $\mathcal{O}(\tau^2)$ bias described in Lemma 7.488. □

Definition 7.513 (Acceptance with pNG allowance)**.** Work on the on–shell, band– and Nielsen–projected posture (cf. Definitions 7.457 and 7.490) on the Step 3 IR window $\mathcal{I}_{\mathrm{IR}}$ with restoration length ℓ_R (cf. Definitions 7.430 and 7.478). A dataset *passes* Step 6 with a pNG mode present if:

1. **Non–pNG multiplets reconverge.** For every *non–pNG* multiplet ρ, the spectral IR–plateau criteria hold as in Definition 7.504.
2. **State–coherence guard.** The state–coherence/consistency guard is satisfied on and beyond ℓ_R (cf. eq. (7.566)).
3. **pNG RG–consistent scaling.** The pNG channel exhibits the scaling implied by Proposition 7.509:

$$\widehat{s}_{\mathrm{pNG}}(\mu_k) \;\in \Big[\alpha\, y_{\tilde{B}} - \delta,\; \alpha\, y_{\tilde{B}} + \delta\Big] \quad \text{for all} \quad \mu_k \in \mathcal{I}_\mu,$$

where $\widehat{s}_{\mathrm{pNG}}$ is the centered log–slope estimator of Definition 7.511, $y_{\tilde{B}}$ is the Step 4 irrelevant exponent (estimated by Definition 7.446), α is the spurion power from Definition 7.508, and $\mathcal{I}_\mu$ is the IR window in $\mu = \kappa/\ell$ (cf. Definition 7.461). The tolerance δ should exceed the known estimator/smoothing biases (cf. Lemma 7.488 and eqs. (7.537) and (7.538)) yet satisfy $\delta \ll |y_{\tilde{B}}|$ on the window. Optionally, when symmetry selection rules predict $\alpha \in \mathbb{N}$, require *integer compatibility* of $\widehat{\alpha}_{\tilde{B}} = \widehat{s}_{\mathrm{pNG}}/\widehat{y}_{\tilde{B}}$ (cf. Definition 7.511) within the same tolerance.

Intuition. A genuine pseudo–Goldstone may retain a small gap while all other multiplets reconverge. Spurion/Wigner–Eckart analysis fixes the leading power α of the breaking amplitude, and Step 4 RG running gives $y_{\tilde{B}} < 0$; together they imply the pNG scaling $m^2_{\mathrm{pNG}} \propto |\tilde{\varepsilon}_{\tilde{B}}|^\alpha \sim \mu^{\alpha y_{\tilde{B}}}$ (Definition 7.508 and Proposition 7.509; see also [91, 82]). Matching the observed pNG log–slope to $\alpha y_{\tilde{B}}$ closes the RG/spectral consistency loop, while the non–pNG multiplets must satisfy the standard reconvergence and restoration ledgers (Definition 7.504).

No state–visibility volume decay (QIII guard).

Definition 7.514 (Volume–uniform state visibility)**.** All spectral diagnostics in Step 6 are conditioned on *uniform* state coherence as in eq. (7.566):

$$\sup_V \Big|1 - \Lambda_{\mathrm{state}}(\lambda_{\mathrm{UV}}, \ell; T^\star, V)\Big| \;\le\; \varepsilon_{\mathrm{plat}} \qquad \text{for all} \quad \ell \ge \ell_R,$$

with the same instrumentation (P_x, K_{IR}) and on–shell/Nielsen posture (cf. Definitions 7.457 and 7.475).

Proposition 7.515 (QIII vs. QII classification guard)**.** *If Definition 7.514 fails on any tail $\ell \geq \ell_\circ$ (i.e. the left–hand side grows with V or does not stay $\leq \varepsilon_{\text{plat}}$), then the observed IR behaviour is* not *QIII: spectral claims (multiplet reconvergence, pNG scaling) are reclassified as QII–like (volume–decoherence–driven). Conversely, if the guard holds, all conclusions of Step 6 (Theorem 7.494 and Definitions 7.504 and 7.513) apply* uniformly in V.

Proof. We work throughout on the on–shell, band– and Nielsen–projected posture for the two–point kernel $\Gamma_2^{\mathcal{N}}(p^2; \lambda_{\text{UV}}, \ell)$ on $W = \operatorname{im}\mathcal{N}$ (cf. Definitions 7.457 and 7.490); on shell, gauge safety holds by Proposition 7.485 and Corollary 7.477.

(A) A failure of volume-uniform state visibility implies that the system is not Q III. The quantity $\Lambda_{\text{state}}(\lambda_{\text{UV}}, \ell; T^\star, V)$ in eq. (7.566) monitors coherence/visibility of one–particle contributions in the channel of interest at time scale $T^\star$ and volume V. If

$$\sup_V \left|1 - \Lambda_{\text{state}}(\lambda_{\text{UV}}, \ell; T^\star, V)\right| > \varepsilon_{\text{plat}} \quad \text{along some tail } \ell \geq \ell_\circ,$$

then either (i) the vacuum–to–one–particle matrix elements decay with V, driving the corresponding LSZ residue to zero, or (ii) the would–be pole contribution is swamped by incoherent multi–particle/continuum effects that do not vanish uniformly in V (both mechanisms are standard in the finite–volume spectral analysis of QFT; cf. [91, 82]). In either case, the extraction of *volume–independent* simple poles and residues used in Step 6 (see Definitions 7.482 and 7.490) becomes unreliable on that tail. The intra–multiplet diagnostics $\{\overline{m_\rho^2}, \delta m_\rho^2, \overline{Z_\rho}, \delta Z_\rho\}$ (Definition 7.499) and their relative versions then reflect volume–driven decoherence rather than genuine RG restoration; accordingly the phenomenon must be classified as QII–like (volume decoherence), not QIII (spectral reconvergence).

(B) Holding of the guard ⇒ uniformity in V of Step 6 conclusions. Assume Definition 7.514. Then there exist V–independent bounds (on compact IR subwindows) for: (i) the residues at simple poles (they stay bounded away from 0 by eq. (7.566) and QFT pole–residue relations [91, 82]); (ii) operator norms and first derivatives of the band–limited kernel and its ρ–blocks (curvature/TV guards and window consistency eq. (7.548) and Proposition 7.462); and (iii) spectral gaps isolating the finite ρ–clusters. These yield resolvent bounds uniform in V, so Kato's perturbation/Implicit Function Theorem arguments (Lemma 7.492; see also [115, Chs. II–IV]) apply uniformly: the pole locations $m_i^2(\ell)$ and residues $Z_i(\ell)$ vary continuously in ℓ with constants independent of V.

The linear–response bounds for intra–multiplet splittings (Proposition 7.493) depend only on the blockwise operator norms and derivatives of $\Gamma_2^{\mathcal{N}}$, and hence the constants $C_\rho(\ell), C'_\rho(\ell)$ can be chosen uniformly in V. Combining these uniform bounds with the Step 4 irrelevance running (Proposition 7.450 and Corollary 7.456) yields the reconvergence rates of Theorem 7.494 uniformly in V, and the acceptance criteria Definition 7.504 (and, with a pNG channel, Definition 7.513) hold uniformly as well. Therefore all Step 6 conclusions apply uniformly in the volume under the guard Definition 7.514.

The two implications (A) and (B) establish the proposition. □

Discrete estimators and audit path.

Definition 7.516 (Spectral estimators on (v, p^2)–grids)**.** Let $v_k := \ln \ell_k$ be a *uniform* grid with step $h > 0$ and let $\{p_r^2\}_r$ be a monotone momentum grid inside the admitted band

$\mathfrak{B}(\lambda_{\mathrm{UV}}, \ell_k)$ (cf. Definition 7.468). Work on the band– and Nielsen–projected two–point kernel $\Gamma_2^{\mathcal{N}}(p^2; \lambda_{\mathrm{UV}}, \ell)$ on $W = \operatorname{im}\mathcal{N}$ (cf. Definitions 7.457 and 7.490); let $\lambda_i(p^2; \ell)$ denote its eigenvalues, labelled continuously in (p^2, ℓ) near each simple pole $p^2 = m_i^2(\ell)$ (cf. Lemma 7.492).

For each (i, k), choose an index $r = r(i, k)$ that *locally brackets the zero* (i.e. $\lambda_i(p_{r-1}^2; \ell_k)\,\lambda_i(p_{r+1}^2; \ell_k) \le 0$) or, when no sign change is observed, minimises the *absolute* value $|\lambda_i(p_r^2; \ell_k)|$. Define the *Newton–stencil mass estimator*

$$\boxed{\widehat{m}_i^2(\ell_k) \;:=\; p_r^2 \;-\; \frac{\lambda_i(p_r^2; \ell_k)}{\widehat{\partial_{p^2}\lambda_i}(p_r^2; \ell_k)},} \tag{7.624}$$

where $\widehat{\partial_{p^2}\lambda_i}(p_r^2; \ell_k)$ is the *second–order* three–point derivative at p_r^2. For a quasi–uniform local mesh $h_\pm := p_{r\pm1}^2 - p_r^2$ one may use the standard nonuniform three–point formula

$$\widehat{\partial_{p^2}\lambda_i}(p_r^2; \ell_k) = -\frac{h_+}{h_-(h_- + h_+)}\,\lambda_i(p_{r-1}^2; \ell_k) + \frac{h_+ - h_-}{h_- h_+}\,\lambda_i(p_r^2; \ell_k) + \frac{h_-}{h_+(h_- + h_+)}\,\lambda_i(p_{r+1}^2; \ell_k).$$

Define the *residue estimator* by

$$\boxed{\widehat{Z}_i(\ell_k) \;:=\; \Big[\, \widehat{\partial_{p^2}\lambda_i}\Big(\widehat{m}_i^2(\ell_k); \ell_k\Big) \,\Big]^{-1},} \tag{7.625}$$

evaluating the same three–point formula after recentering the stencil at the grid point closest to $\widehat{m}_i^2(\ell_k)$.

Construct the multiplet dispersions $\overline{m_\rho^2}$, δm_ρ^2, $\overline{Z_\rho}$, δZ_ρ from eqs. (7.618) and (7.619), using the sets $\mathfrak{M}_\rho(\ell_k)$ of Definition 7.499. Optionally apply a low–pass in v satisfying eqs. (7.537) and (7.538) to the sequences $\{\widehat{m}_i^2(\ell_k)\}_k$ and $\{\widehat{Z}_i(\ell_k)\}_k$ before forming dispersions; this introduces only an $\mathcal{O}(\tau^2)$ bias (cf. Lemma 7.488).

Lemma 7.517 (Consistency and rates of the discrete spectral estimators)**.** *Assume for each (i, k) that $p^2 \mapsto \lambda_i(p^2; \ell_k)$ is C^3 in a neighbourhood of its simple zero $p^2 = m_i^2(\ell_k)$, with*

$$\partial_{p^2}\lambda_i\Big(m_i^2(\ell_k); \ell_k\Big) = Z_i(\ell_k)^{-1} \neq 0, \qquad \|\partial_{p^2}^2\lambda_i\|,\ \|\partial_{p^2}^3\lambda_i\| \text{ bounded near } m_i^2(\ell_k).$$

If the local mesh is quasi–uniform in the sense that $h_\pm = p_{r\pm1}^2 - p_r^2$ satisfy $\theta^{-1} \le h_+/h_- \le \theta$ for some fixed $\theta > 0$, then as the momentum grid is refined (with $h := \max\{h_-, h_+\} \to 0$),

$$\widehat{m}_i^2(\ell_k) = m_i^2(\ell_k) + \mathcal{O}(h^2), \qquad \widehat{Z}_i(\ell_k) = Z_i(\ell_k) + \mathcal{O}(h^2).$$

Proof. Fix (i, k) and write $\lambda(p) := \lambda_i(p^2; \ell_k)$ with p^2 as the scalar variable; let $m^2 := m_i^2(\ell_k)$ and $a := \lambda'(m^2) = Z_i(\ell_k)^{-1} \neq 0$. By Taylor's theorem with integral remainder,

$$\lambda(p^2) = a(p^2 - m^2) + \tfrac{1}{2}b(p^2 - m^2)^2 + \tfrac{1}{6}c(p^2 - m^2)^3 + \mathcal{O}\Big(|p^2 - m^2|^4\Big),$$

with $b = \lambda''(m^2)$ and $c = \lambda^{(3)}(\xi)$ bounded near m^2.

(1) Derivative consistency. The three–point nonuniform formula used for $\widehat{\partial_{p^2}\lambda}(p_r^2)$ is the unique second–order accurate interpolatory stencil on $\{p_{r-1}^2, p_r^2, p_{r+1}^2\}$; standard Peano kernel estimates give

$$\widehat{\partial_{p^2}\lambda}(p_r^2) = \lambda'(p_r^2) + \mathcal{O}(h^2) \quad \text{with } h := \max\{h_-, h_+\},$$

provided $h_\pm$ are boundedly comparable (quasi–uniform) and $\lambda \in C^3$.

(2) Newton–stencil mass estimator. Set $\phi(p^2) := p^2 - \lambda(p^2)/\lambda'(p^2)$ (the exact Newton map). For a simple zero and C^2 smoothness, $\phi(m^2) = m^2$ and $\phi'(m^2) = 0$ with ϕ'' bounded; hence a single Newton step from any p_r^2 in a sufficiently small neighbourhood of m^2 yields

$$p_r^2 - \frac{\lambda(p_r^2)}{\lambda'(p_r^2)} \;=\; m^2 + \mathcal{O}\Big((p_r^2 - m^2)^2\Big).$$

Replacing λ' by the second–order estimator $\widehat{\partial_{p^2}\lambda}$ introduces only an additional $\mathcal{O}(h^2)$ perturbation in the denominator. Since by the bracketing or absolute–minimum choice we have $|p_r^2 - m^2| \le h$ and $\lambda'(m^2) = a \neq 0$, a standard perturbation argument (expanding the denominator around $\lambda'(p_r^2)$ and using the boundedness of $1/\lambda'$ near m^2) gives

$$\widehat{m}_i^2(\ell_k) = p_r^2 - \frac{\lambda(p_r^2)}{\widehat{\partial_{p^2}\lambda}(p_r^2)} = m^2 + \mathcal{O}\Big(h^2\Big).$$

(3) Residue estimator. By definition $Z_i(\ell_k) = \Big[\lambda'(m^2)\Big]^{-1}$. Using the derivative estimator at the recentered stencil nearest to $\widehat{m}_i^2$, write

$$\widehat{Z}_i(\ell_k) = \Big[\lambda'(\widehat{m}_i^2) + \Big(\widehat{\partial_{p^2}\lambda} - \lambda'\Big)(\widehat{m}_i^2)\Big]^{-1}.$$

The first term equals $Z_i(\ell_k)^{-1} + \mathcal{O}(h^2)$ by (2) and C^2 smoothness; the second term is $\mathcal{O}(h^2)$ by (1). Since $Z_i(\ell_k)^{-1} \neq 0$ and the perturbation is small, a Neumann–series expansion of the inverse yields $\widehat{Z}_i(\ell_k) = Z_i(\ell_k) + \mathcal{O}(h^2)$.

The stated rates follow. □

Lemma 7.518 (Gauge/basis and scale invariances)**.** *The discrete spectral estimators* (7.624)–(7.625) *and the derived dispersions* $\overline{m_\rho^2}, \delta m_\rho^2, \overline{Z_\rho}, \delta Z_\rho$ *(cf. Definition 7.499) are:*

(i) gauge–parameter independent within the tolerance eq. (7.588) *when evaluated on shell and Nielsen–projected (Definition 7.457);*

(ii) invariant under admissible basis/scheme changes acting by similarity on $W = \operatorname{im}\mathcal{N}$ *(cf. Proposition 7.458 and Lemmas 7.451 and 7.483);*

(iii) invariant under reparametrisations $\ell \mapsto a\,\ell$ *(rigid shifts* $v = \ln \ell \mapsto v + \ln a$*) and under positive rescalings of the reference response* $\mathcal{R}$ *used in defining the admitted band in Definition 7.468.*

Proof. We work in the on–shell, band– and Nielsen–projected posture for $\Gamma_2^{\mathcal{N}}(p^2; \lambda_{\mathrm{UV}}, \ell)$ (cf. Definitions 7.457 and 7.490).

(i) Gauge–parameter independence. On shell, the gauge–parameter dependence of $\Gamma_2^{\mathcal{N}}$ is a similarity flow (Nielsen/Lax form), and pole positions/residues are gauge–parameter independent within the audit tolerance (see Proposition 7.485 and Corollary 7.477). The estimators (7.624)–(7.625) are built solely from the scalar eigenvalue branches $\lambda_i(p^2; \ell)$ and their p^2–derivatives at the pole; under similarity these eigenvalues are unchanged as functions of p^2. Hence $\widehat{m}_i^2(\ell_k)$ and $\widehat{Z}_i(\ell_k)$ inherit the same gauge safety, and so do the dispersion functionals formed from $\{m_i^2, Z_i\}$ (cf. Definition 7.499).

(ii) Basis/scheme invariance. Any admissible basis change within $\operatorname{im} P^{\rho}(\ell)$ acts by similarity on the ρ–block $A_{\rho} = P^{\rho}\Gamma_2^{\mathcal{N}}P^{\rho}$ (cf. Proposition 7.458 and Lemma 7.483). Similarity leaves eigenvalue locations and the scalar residues (defined by Definitions 7.482 and 7.490) invariant as numbers; therefore the discrete reconstructions (7.624)–(7.625) and the derived dispersions are basis/scheme invariant (see also Lemma 7.451).

(iii) Scale and $\mathcal{R}$ reparametrisations. A change of units $\ell \mapsto a\,\ell$ induces a rigid shift $v \mapsto v + \ln a$. Our spectral estimators are pointwise in ℓ (no v–differences are used at this stage), so the sequences $\{\widehat{m}_i^2(\ell_k)\}$ and $\{\widehat{Z}_i(\ell_k)\}$ are merely reindexed; any optional centered v–differences (when used elsewhere) are invariant under rigid shifts. As for $\mathcal{R}$ in Definition 7.468, positive rescalings of $\mathcal{R}$ do not change the admitted momentum band $\mathfrak{B}(\lambda_{\mathrm{UV}}, \ell)$ determined by the fixed instrumentation (P_x, K_{IR}) (cf. Definition 7.475); consequently the set of p^2 values used in Definition 7.516 is unaffected, and so are the estimators and dispersions.

Combining (i)–(iii) establishes the lemma. □

Definition 7.519 (Windowed reporting and spreads)**.** On the IR window $\mathcal{I}_{\mathrm{IR}}$ used in Definition 7.504, report:

1. **Windowed extrema and slopes:**

$$\max_{v_k \in \mathcal{I}_{\mathrm{IR}}} \Delta^{\mathrm{rel}}_{m,\rho}(v_k), \qquad \max_{v_k \in \mathcal{I}_{\mathrm{IR}}} \Delta^{\mathrm{rel}}_{Z,\rho}(v_k),$$

   and the centered log–slopes $\partial_v \widehat{\ln \Delta^{\mathrm{rel}}_{(\cdot),\rho}}$ on the same window (cf. Definitions 7.497 and 7.507).
2. **Instrumentation and gauge spreads:** Spreads under admissible instrumentation (P_x, K_{IR}) (cf. Definition 7.475) and across admissible gauges Ξ_{adm}, with Nielsen–projected, on–shell posture (cf. Definition 7.457). Gauge flatness is expected within the audit tolerance by Proposition 7.485 and eq. (7.588).
3. **Numerical settings:** The grid parameters $(h, \Delta p^2)$, any low–pass widths used in v (obeying eqs. (7.537) and (7.538)), and the interpolation/stencil order employed in the spectral estimators eqs. (7.624) and (7.625).

All items are conditioned on the volume–uniform state–visibility guard Definition 7.514.

Intuition. We only trust spectral reconvergence when the *state* stays uniformly visible in volume (QIII guard; Definition 7.514). Operationally, we solve and differentiate the band– and Nielsen–projected two–point problem on (v, p^2)–grids (Definition 7.516); centered stencils with light low–pass yield second–order–accurate, nearly unbiased pole/residue estimates (Lemmas 7.488 and 7.517), which then feed the multiplet dispersions used for plateau acceptance (Definition 7.504).

Acceptance criteria (spectral reconvergence).

Definition 7.520 (Step 6 acceptance: spectral reconvergence)**.** Work on the IR window $\mathcal{I}_{\mathrm{IR}}$ in $v = \ln \ell$ with ℓ_R as in Definitions 7.430 and 7.504. The dataset *passes* Step 6 if the following hold:

1. **Multiplet closure.** For every multiplet ρ, the spectral plateau conditions of Definition 7.504 hold at and beyond ℓ_R, i.e.

$$\frac{\mathrm{d}}{\mathrm{d}\ln\ell}\left(\frac{\delta m_\rho^2}{\overline{m_\rho^2}}\right)\le 0, \qquad \frac{\mathrm{d}}{\mathrm{d}\ln\ell}\left(\frac{\delta Z_\rho}{\overline{Z_\rho}}\right)\le 0 \quad \text{on } \mathcal{I}_{\mathrm{IR}},$$
$$\max\left\{\frac{\delta m_\rho^2(\ell_R)}{\overline{m_\rho^2}(\ell_R)}, \frac{\delta Z_\rho(\ell_R)}{\overline{Z_\rho}(\ell_R)}\right\}\le \varepsilon_{\mathrm{spec}}.$$

2. **Gauge safety.** For all quoted poles and residues, gauge–parameter slopes vanish within tolerance on $\mathcal{I}_{\mathrm{IR}}$ (Nielsen–projected, on shell):

$$\frac{\partial m_i^2(\ell;\xi)}{\partial\xi}=0, \qquad \frac{\partial Z_i(\ell;\xi)}{\partial\xi}=0,$$
up to $\varepsilon_{\mathrm{gauge}}$ for all $\forall\,\xi\in\mathcal{X}_{\mathrm{adm}}$ (cf. *Proposition 7.485and Corollary 7.477*).

3. **QIII guard.** State visibility is volume–uniform (no volume–law decay) on/beyond ℓ_R:

$$\sup_V\left|1-\Lambda_{\mathrm{state}}(\lambda_{\mathrm{UV}},\ell;T^\star,V)\right|\le\varepsilon_{\mathrm{plat}} \quad \text{for all } \ell\ge\ell_R$$
(cf. *Definition 7.514and eq.* (7.566)).

4. **pNG allowance.** Any pseudo–NG mode present satisfies the RG–consistent scaling law

$$m^2_{\mathrm{pNG}}(\ell)\ \propto\ \left|\widetilde{\varepsilon}_{\tilde B}(\mu)\right|^{\alpha}\ \sim\ \mu^{\alpha y_{\tilde B}} \quad (\mu=\kappa/\ell),$$
as in Proposition 7.509 and eq. (7.623), with the remaining multiplets still fulfilling (i).

Proposition 7.521 (Soundness of Step 6 acceptance)**.** *If Definition 7.520 holds and Steps 2–4 are satisfied (positive law–slope on the window, finite restoration length, and* $\mathrm{Re}\,y_{\tilde B}<0$ *for excited projected breakings; cf. eq.* (7.542)*, Definitions 7.430 and 7.478, Proposition 7.450, and Corollary 7.456), then multiplet reconvergence* (7.614) *obtains for all non–pNG multiplets, and any pNG mode exhibits the log–slope*

$$\frac{\mathrm{d}}{\mathrm{d}\ln\mu}\ln m^2_{\mathrm{pNG}}(\ell)=\alpha\,y_{\tilde B}+o(1) \quad \text{as } \mu\downarrow 0.$$

Proof. All statements are made on the band– and Nielsen–projected posture for $\Gamma_2^{\mathcal{N}}$ on $W=\mathrm{im}\,\mathcal{N}$ (cf. Definitions 7.457 and 7.490); on shell, gauge safety holds by Proposition 7.485 and Corollary 7.477. Window regularity (curvature/TV guards and Step 3 mask) is assumed from Steps 2–3 (cf. eq. (7.548) and Proposition 7.462).

(A) Non–pNG multiplets reconverge. By item (i) of Definition 7.520 and Lemma 7.505, each relative dispersion $\Delta^{\mathrm{rel}}_{m,\rho}=\delta m_\rho^2/\overline{m_\rho^2}$ and $\Delta^{\mathrm{rel}}_{Z,\rho}=\delta Z_\rho/\overline{Z_\rho}$ is nonincreasing in v on $\mathcal{I}_{\mathrm{IR}}$ and small at $v_R=\ln\ell_R$. Step 4 irrelevance provides decay of the explicit breakings (cf. Proposition 7.450 and Corollary 7.456); combined with the linear–response bounds of Proposition 7.493 and the sufficient–conditions lemma Proposition 7.506, there exists $\ell_\star\ge\ell_R$ such that

$$\frac{\mathrm{d}}{\mathrm{d}\ln\ell}\ln\Delta^{\mathrm{rel}}_{m,\rho}(\ell)\le-\gamma_\star/2<0, \qquad \frac{\mathrm{d}}{\mathrm{d}\ln\ell}\ln\Delta^{\mathrm{rel}}_{Z,\rho}(\ell)\le-\gamma_\star/2<0 \quad \text{for all } \ell\ge\ell_\star,$$

where $\gamma_\star := \min_{\tilde{B}} |\operatorname{Re} y_{\tilde{B}}| > 0$. Integrating these differential inequalities and using smallness at ℓ_R yields

$$\Delta^{\mathrm{rel}}_{m,\rho}(\ell),\ \Delta^{\mathrm{rel}}_{Z,\rho}(\ell)\ \longrightarrow\ 0\quad(\ell\uparrow),$$

i.e. (7.614). By Corollary 7.503 (continuity of the ratios) and Lemma 7.501 (equivalence of dispersion metrics), this is equivalent to vanishing absolute intra–multiplet splittings relative to the means.

(B) QIII guard ⇒ uniform one–particle interpretation. Item (iii) invokes Definition 7.514. By Proposition 7.515, this ensures that the pole/residue extraction and the bounds above hold *uniformly in the volume* V; Kato continuity and simple–pole isolation (cf. Lemma 7.492 and [115, Chs. II–IV]) apply with V–independent constants. Hence the reconvergence conclusion is volume–uniform (QIII, not QII).

(C) pNG scaling. If a pNG channel is present, item (iv) requires the RG–consistent law of Proposition 7.509: $m^2_{\mathrm{pNG}}(\ell) = C_{\mathrm{pNG}}(\ell)\,|\widetilde{\varepsilon}_{\tilde{B}}(\mu)|^{\alpha}$ with slowly varying $C_{\mathrm{pNG}}(\ell)$. Differentiating $\ln m^2_{\mathrm{pNG}}$ with respect to $\ln\mu$ and using the Step 4 running of $\widetilde{\varepsilon}_{\tilde{B}}$ gives $\frac{\mathrm{d}}{\mathrm{d}\ln\mu}\ln m^2_{\mathrm{pNG}} = \alpha\, y_{\tilde{B}} + o(1)$ as $\mu\downarrow 0$, as claimed.

Putting (A)–(C) together proves the proposition: non–pNG multiplets reconverge spectrally on the IR plateau with rates controlled by $\gamma_\star$ (Theorem 7.494), the one–particle interpretation is stable (QIII guard), and any pNG mode exhibits the predicted log–slope. Standard QFT pole/residue conventions underlie the on–shell analysis [91, 82]. □

Outputs for Step 6 (synthesised).

Definition 7.522 (Step 6 outputs)**.** All items below are produced on the IR window $\mathcal{I}_{\mathrm{IR}}$ in $v = \ln\ell$ (cf. eqs. (7.542) and (7.551) and Proposition 7.462), on shell and Nielsen–projected (Definition 7.457), with fixed instrumentation (P_x, K_{IR}) (Definition 7.475), and under the volume–uniform state–visibility guard (Definition 7.514). Report:

1. **Global RG/plateau diagnostics.** Plots of $v\mapsto \Lambda_{\mathrm{law}}(v)$ and $v\mapsto \varsigma^{(\tau)}_{\mathrm{law}}(v)$ together with shaded $\mathcal{I}_{\mathrm{IR}}$ and a marker at $v = \ln\ell_R$ (cf. Definition 7.478 and eqs. (7.542) and (7.551)).
2. **Multiplet reconvergence panels (per ρ).** Plots of

$$v\ \mapsto\ \Delta^{\mathrm{rel}}_{m,\rho}(v) := \frac{\delta m^2_\rho(v)}{\overline{m^2_\rho}(v)},\qquad v\ \mapsto\ \Delta^{\mathrm{rel}}_{Z,\rho}(v) := \frac{\delta Z_\rho(v)}{\overline{Z_\rho}(v)},$$

   with fitted (negative) centered log–slopes $\partial_v\widehat{\ln\Delta^{\mathrm{rel}}_{(\cdot),\rho}}$ (cf. Definitions 7.497, 7.499, 7.504 and 7.507). Mark $v = \ln\ell_R$.
3. **Pole/residue tables and gauge/band spreads.** Tables at/beyond ℓ_R of $\{m_i^2, Z_i\}$, grouped by multiplet ρ, together with: (i) spreads across admissible gauges $\mathcal{X}_{\mathrm{adm}}$ (Nielsen–projected) using the centered discrete gauge–slope checks from Definition 7.487, i.e. the estimators in (7.608) (consistent with the continuous zero–slope result Proposition 7.485); (ii) spreads across admissible band–limiting choices (P_x, K_{IR}) consistent with Definition 7.475.

4. **Numerical/audit settings.** The grid parameters $(h, \Delta p^2)$, any low–pass widths in v and in ξ obeying eqs. (7.537) and (7.538), and the interpolation/stencil order used for the spectral estimators (7.624)–(7.625) (cf. Definition 7.516 and Lemmas 7.488 and 7.517).
5. **(Optional pNG panel).** A plot of $\ln m^2_{\mathrm{pNG}}$ vs. $\ln \mu$ on the IR window in $\mu = \kappa/\ell$ (cf. Definition 7.461), with fitted slope $\widehat{s}_{\mathrm{pNG}}$ and comparison to the Step 4 prediction $\alpha\, y_{\tilde{B}}$ (Propositions 7.450 and 7.509 and Corollary 7.456); also report the inferred spurion power $\widehat{\alpha}_{\tilde{B}} = \widehat{s}_{\mathrm{pNG}}/\widehat{y}_{\tilde{B}}$ (cf. Definition 7.511).
6. **Rate interpretation.** A short cross–reference explaining observed decay rates via Step 4 irrelevance (slowest $\gamma_\star$) and smooth spectral tracking, citing Theorem 7.494, Proposition 7.506, and Lemma 7.492 and the Ward–diagnostics posture Definition 7.468; standard spectral/QFT references [91, 82, 115].

Interpretation. As explicit breakings run to zero, symmetry–forbidden intra–multiplet splittings collapse. Kato continuity on the band– and Nielsen–projected kernel guarantees smooth pole/residue tracking (Lemma 7.492; [115]); Step 4 irrelevance quantifies decay rates (Proposition 7.450 and Corollary 7.456), yielding spectral reconvergence on the IR plateau (Theorem 7.494). Gauge safety follows from the on–shell Nielsen posture (Proposition 7.485 and Corollary 7.477); the QIII state–visibility guard ensures a stable one–particle interpretation *uniformly in volume* (Definition 7.514 and Proposition 7.515). For pNG channels, the spurion/Wigner–Eckart selection rule and RG running give $m^2_{\mathrm{pNG}} \propto |\widetilde{\varepsilon}_{\tilde{B}}|^{\alpha} \sim \mu^{\alpha y_{\tilde{B}}}$ (Definition 7.508 and Proposition 7.509; [91, 82]).

7.8.7 Plateau Certification (Q III Mask)

Objective. We construct a *mask* $\mathfrak{M}_{\mathrm{mask}} \subset \mathbb{R}^2$ in the plane

$$u := \ln \lambda_{\mathrm{UV}}, \qquad v := \ln \ell,$$

that localises the IR plateau where (i) the law heals with positive robust slope and small curvature, (ii) the state remains coherent uniformly in the volume, and (iii) the identity ledger closes (small Ward residuals and gauge–flat on–shell data). The mask gates all downstream IR statements (Steps 7.8.4–7.8.6) and underlies integrated readouts.

Definition (pointwise mask). With the robust slope/curvature guards from eqs. (7.539), (7.542) and (7.548), the state–coherence guard from eq. (7.566), and the Ward/Nielsen conditions from eqs. (7.591), (7.602) and (7.607), define

$$\begin{aligned}\mathfrak{M}_{\mathrm{mask}} := \Big\{ (u,v) : \varsigma^{(\tau)}_{\mathrm{law}}(u,v) \;\geq\; \varsigma_{\min}, \quad & \big|\partial_v^2 (\mathsf{LP}_\tau\, \Lambda_{\mathrm{law}})(u,v)\big| \;\leq\; \varepsilon_{\mathrm{curv}}, \\ & \sup_V \big|1 - \Lambda_{\mathrm{state}}(u,v;V)\big| \;\leq\; \varepsilon_{\mathrm{plat}}, \\ & \mathcal{W}(u,v) \;\leq\; \varepsilon_{\mathrm{plat}}, \\ & \partial_\xi m^2_\alpha(u,v;\xi) = 0, \quad \partial_\xi Z_\alpha(u,v;\xi) = 0 \;\; \forall\, \xi \in \mathcal{X}_{\mathrm{adm}} \Big\}.\end{aligned} \tag{7.626}$$

Reading guide. The first two lines enforce a monotone, gently curved law-healing profile (via the low-pass robust slope $\varsigma_{\mathrm{law}}^{(\tau)}$ and curvature bound). The third line is the uniform state-coherence guard in volume. The fourth controls the band-projected Ward residual $\mathcal{W}$, and the last line imposes Nielsen flatness of quoted on–shell poles and residues across the admissible gauges $\mathcal{X}_{\mathrm{adm}}$ (cf. eq. (7.607)) [117, 243, 442].

Lemma 7.523 (Closedness, tolerance monotonicity, and scale reparametrisation)**.** *Assume the ingredients entering eq. (7.626) are jointly continuous in (u, v) and in any auxiliary indices, and that the probed sets are compact: the admissible gauges $\mathcal{X}_{\mathrm{adm}}$, the IR momentum band used in $\mathcal{W}$ (cf. eq. (7.591)), and the volume set over which the state guard is tested. Then:*

1. *$\mathfrak{M}_{\mathrm{mask}}$ is a closed subset of $\mathbb{R}^2$.*
2. (Tolerance monotonicity) *Relaxing any threshold $\varsigma_{\mathrm{min}}\downarrow$, $\varepsilon_{\mathrm{curv}}\uparrow$, $\varepsilon_{\mathrm{plat}}\uparrow$ can only enlarge $\mathfrak{M}_{\mathrm{mask}}$ (set inclusion is monotone).*
3. (Scale reparametrisation) *For any $\Delta \in \mathbb{R}$, the coordinate change $v \mapsto v + \Delta$ (i.e. $\ell \mapsto e^{\Delta}\ell$) rigidly translates the mask in the v–direction without altering its shape:*

$$T_{\Delta}(\mathfrak{M}_{\mathrm{mask}}) = \{(u, v + \Delta) : (u, v) \in \mathfrak{M}_{\mathrm{mask}}\}.$$

Proof. Recall the pointwise definition eq. (7.626):

$$\mathfrak{M}_{\mathrm{mask}} = \{(u, v) : \text{C1–C5 hold at } (u, v)\},$$

with guards

$$\begin{aligned}
&\text{C1:} && \varsigma_{\mathrm{law}}^{(\tau)}(u, v) \geq \varsigma_{\mathrm{min}}, && \\
&\text{C2:} && \left|\partial_v^2(\mathsf{LP}_{\tau}\Lambda_{\mathrm{law}})(u, v)\right| \leq \varepsilon_{\mathrm{curv}}, && \\
&\text{C3:} && \sup_{V}\left|1 - \Lambda_{\mathrm{state}}(u, v; V)\right| \leq \varepsilon_{\mathrm{plat}} && \text{(cf. eq. (7.566)),} \\
&\text{C4:} && \mathcal{W}(u, v) \leq \varepsilon_{\mathrm{plat}} && \text{(cf. eqs. (7.591) and (7.602)),} \\
&\text{C5:} && \sup_{\xi \in \mathcal{X}_{\mathrm{adm}}}\left|\partial_{\xi} m_{\alpha}^2(u, v; \xi)\right| = 0, && \\
& && \sup_{\xi \in \mathcal{X}_{\mathrm{adm}}}\left|\partial_{\xi} Z_{\alpha}(u, v; \xi)\right| = 0 && \text{(cf. eq. (7.607)).}
\end{aligned}$$

(1) Closedness. Define the real-valued functions

$$\begin{aligned}
f_1(u, v) &:= \varsigma_{\mathrm{law}}^{(\tau)}(u, v) - \varsigma_{\mathrm{min}}, \\
f_2(u, v) &:= \varepsilon_{\mathrm{curv}} - \left|\partial_v^2(\mathsf{LP}_{\tau}\Lambda_{\mathrm{law}})(u, v)\right|, \\
f_3(u, v) &:= \varepsilon_{\mathrm{plat}} - \sup_{V}\left|1 - \Lambda_{\mathrm{state}}(u, v; V)\right|, \\
f_4(u, v) &:= \varepsilon_{\mathrm{plat}} - \mathcal{W}(u, v), \\
f_5^{(m)}(u, v) &:= -\sup_{\xi \in \mathcal{X}_{\mathrm{adm}}}\left|\partial_{\xi} m_{\alpha}^2(u, v; \xi)\right|, \qquad f_5^{(Z)}(u, v) := -\sup_{\xi \in \mathcal{X}_{\mathrm{adm}}}\left|\partial_{\xi} Z_{\alpha}(u, v; \xi)\right|.
\end{aligned}$$

By hypothesis, the basic maps are continuous in (u, v) and jointly continuous in $(u, v; V)$ and $(u, v; \xi)$. Since the probed sets (volumes, momentum band, and $\mathcal{X}_{\mathrm{adm}}$) are compact,

the supremum of a continuous function over a compact index set is continuous in the base variables (standard maximum theorem). Hence each f_i is continuous. The mask is the finite intersection

$$\mathfrak{M}_{\mathrm{mask}} = \bigcap_{i=1}^{4} \{(u,v) : f_i(u,v) \geq 0\} \;\cap\; \{(u,v) : f_5^{(m)}(u,v) \geq 0\} \;\cap\; \{(u,v) : f_5^{(Z)}(u,v) \geq 0\},$$

an intersection of inverse images of the closed set $[0,\infty)$ under continuous maps, hence closed. *(2) Tolerance monotonicity.* Let $(\varsigma_{\min}, \varepsilon_{\mathrm{curv}}, \varepsilon_{\mathrm{plat}})$ and $(\varsigma'_{\min}, \varepsilon'_{\mathrm{curv}}, \varepsilon'_{\mathrm{plat}})$ be two tolerance sets with $\varsigma'_{\min} \leq \varsigma_{\min}$, $\varepsilon'_{\mathrm{curv}} \geq \varepsilon_{\mathrm{curv}}$, $\varepsilon'_{\mathrm{plat}} \geq \varepsilon_{\mathrm{plat}}$. Each guard is an inequality of the form "observable $\geq$ lower bound" or "observable $\leq$ upper bound." Weakening the lower bound or enlarging the upper bound can only convert failures into passes, never the reverse. Therefore $\mathfrak{M}_{\mathrm{mask}}(\varsigma_{\min}, \varepsilon_{\mathrm{curv}}, \varepsilon_{\mathrm{plat}}) \subseteq \mathfrak{M}_{\mathrm{mask}}(\varsigma'_{\min}, \varepsilon'_{\mathrm{curv}}, \varepsilon'_{\mathrm{plat}})$.
(3) Scale reparametrisation (rigid v–translation). Let $T_\Delta(u,v) := (u, v+\Delta)$. All guards in eq. (7.626) are built from (u,v) through the band parameter $\ell = e^v$ and through v–stationary operations in the v–coordinate: (i) the low-pass LP_τ used in the robust slope/curvature guards is translation-equivariant in v by construction (it is a stationary filter), hence

$$\mathsf{LP}_\tau[\Lambda_{\mathrm{law}}(\cdot + \Delta)](v) = (\mathsf{LP}_\tau \Lambda_{\mathrm{law}})(v+\Delta), \qquad \partial_v^k \text{ commutes with } T_\Delta;$$

(ii) the state guard uses $\sup_V |1 - \Lambda_{\mathrm{state}}(u,v;V)|$, which under $v \mapsto v+\Delta$ becomes composition by T_Δ in the base variables and leaves the inequality form unchanged (the probed V set is unaffected); (iii) the Ward residual $\mathcal{W}$ is computed on the IR band determined by $\ell = e^v$ (cf. eq. (7.591)); expressing it as a function of v therefore yields a graph that simply shifts under $v \mapsto v+\Delta$; (iv) the Nielsen flatness conditions $\sup_{\xi \in \mathcal{X}_{\mathrm{adm}}} |\partial_\xi m_\alpha^2| = 0$ and similarly for Z_α are invariant under composing their (u,v)–dependence with T_Δ, since differentiation is in ξ and $\mathcal{X}_{\mathrm{adm}}$ is unchanged. Consequently, for every (u,v) the guards hold at (u,v) iff they hold at $(u, v+\Delta)$ after the reparametrisation, which is precisely $T_\Delta(\mathfrak{M}_{\mathrm{mask}}) = \{(u, v+\Delta) : (u,v) \in \mathfrak{M}_{\mathrm{mask}}\}$. □

Intuition. The mask is defined by finitely many "$\leq/\geq$" tests of continuous diagnostics; that makes it closed and ensures that easing any tolerance only grows the pass set. Because all diagnostics depend on the IR scale through $v = \ln \ell$ and use a stationary low-pass in v, changing the v zero (i.e. units of ℓ) just slides the certified region up or down the v-axis.

Proposition 7.524 (Upward closure in v and frontier)**.** *Assume the Step 3 guards (robust positive slope and curvature bound; cf. eqs. (7.539), (7.542) and (7.548)) and the Step 5 Ward decay/plateau (eqs. (7.591) and (7.602)) hold as formulated in eq. (7.626), and use the state–coherence guard of eq. (7.566). For each fixed $u \in \mathbb{R}$ the vertical section*

$$\mathfrak{M}_{\mathrm{mask}}(u) := \{\, v \in \mathbb{R} : \; (u,v) \in \mathfrak{M}_{\mathrm{mask}} \,\}$$

is an upper set*: if $v \in \mathfrak{M}_{\mathrm{mask}}(u)$ and $v' \geq v$, then $v' \in \mathfrak{M}_{\mathrm{mask}}(u)$. Define the frontier*

$$v_\star(u) := \inf \mathfrak{M}_{\mathrm{mask}}(u) \;\in [-\infty, \infty].$$

Then $\mathfrak{M}_{\mathrm{mask}} = \{(u,v) : v \geq v_\star(u)\}$, and $v_\star$ is lower semicontinuous in u.

Proof. Write the five pointwise guards in eq. (7.626) at (u, v) as:

$$\text{C1: } \varsigma^{(\tau)}_{\text{law}}(u,v) \geq \varsigma_{\min}, \qquad \text{C2: } \left|\partial_v^2(\mathsf{LP}_\tau \Lambda_{\text{law}})(u,v)\right| \leq \varepsilon_{\text{curv}},$$

$$\text{C3: } \sup_V \left|1 - \Lambda_{\text{state}}(u,v;V)\right| \leq \varepsilon_{\text{plat}}, \qquad \text{C4: } \mathcal{W}(u,v) \leq \varepsilon_{\text{plat}},$$

$$\text{C5: } \sup_{\xi \in \mathcal{X}_{\text{adm}}} \left|\partial_\xi m_\alpha^2(u,v;\xi)\right| = 0, \quad \sup_{\xi \in \mathcal{X}_{\text{adm}}} \left|\partial_\xi Z_\alpha(u,v;\xi)\right| = 0.$$

Upward closure. Fix u and suppose $(u, v) \in \mathfrak{M}_{\text{mask}}$. We show each guard persists for any $v' \geq v$:

• **C1 (robust slope).** By the Step 3 acceptance (eqs. (7.542) and (7.548)), the v–profile of the low–pass trend $\mathsf{LP}_\tau \Lambda_{\text{law}}$ is monotone increasing along the IR ray and its robust derivative $\varsigma^{(\tau)}_{\text{law}}$ stays above $\varsigma_{\min}$ across the IR window. Hence if $\varsigma^{(\tau)}_{\text{law}}(u,v) \geq \varsigma_{\min}$, then also $\varsigma^{(\tau)}_{\text{law}}(u,v') \geq \varsigma_{\min}$ for all $v' \geq v$.

• **C2 (curvature bound).** The Step 3 curvature guard is imposed *uniformly* on the same IR v–band used for C1 (via the same stationary low–pass LP_τ). Thus once $|\partial_v^2(\mathsf{LP}_\tau \Lambda_{\text{law}})| \leq \varepsilon_{\text{curv}}$ at (u, v), it remains within the bound for all $v' \geq v$ in that band.

• **C3 (state coherence).** The guard in eq. (7.566) is framed as an IR–anchored acceptance: beyond the restoration scale $v \geq \ln \ell_R$ it holds uniformly in V (by definition of the plateau certification window in Step 7). Hence, if $(u, v) \in \mathfrak{M}_{\text{mask}}$, then for every $v' \geq \max\{v, \ln \ell_R\}$ the inequality also holds. In particular, for $v' \geq v$ the guard persists (either we stay below $\ln \ell_R$ and the local acceptance is unchanged, or we enter the anchored region where it is uniform).

• **C4 (Ward residual).** Step 5 enforces IR decay of the band–projected Ward residual (eq. (7.602)), so $\mathcal{W}(u, v') \leq \mathcal{W}(u, v) \leq \varepsilon_{\text{plat}}$ for all $v' \geq v$.

• **C5 (Nielsen flatness).** The on–shell gauge slopes vanish throughout the admissible gauges on the IR window (cf. eq. (7.607)); composing with $v' \geq v$ does not affect differentiation in ξ, so the equalities persist.

All guards thus remain satisfied at (u, v') for any $v' \geq v$, proving that $\mathfrak{M}_{\text{mask}}(u)$ is an upper set.

Frontier and epigraph form. For each u, define $v_\star(u) := \inf \mathfrak{M}_{\text{mask}}(u)$ (with the usual convention $\inf \varnothing = +\infty$). Because $\mathfrak{M}_{\text{mask}}(u)$ is an upper set, one has

$$\mathfrak{M}_{\text{mask}}(u) = \{\, v \in \mathbb{R} : \; v \geq v_\star(u) \,\}.$$

Equivalently, $\mathfrak{M}_{\text{mask}} = \operatorname{epi} v_\star := \{(u, v) : v \geq v_\star(u)\}$.

Lower semicontinuity of $v_\star$. By Lemma 7.523, $\mathfrak{M}_{\text{mask}}$ is closed. The epigraph of a function is closed iff the function is lower semicontinuous (standard epigraph characterization; e.g., Rockafellar, *Convex Analysis*, Thm. 7.1). Hence $v_\star$ is lower semicontinuous. □

Reading guide. Each guard either improves or is fixed across the IR direction: the law's trend keeps healing (positive robust slope; bounded curvature), the Ward identity residual decays, and the state guard is uniform past $v = \ln \ell_R$. That converts every vertical slice of the mask into an "upper set," so the mask is exactly the epigraph of its frontier $v_\star(u)$; closedness of the mask forces $v_\star$ to be lower semicontinuous.

Corollary 7.525 (Relation to restoration lengths)**.** *Let $\ell_R(u)$ and $\ell_R^{[\mathcal{U}]}$ be the pointwise and UV–uniform restoration lengths from Definitions 7.428 and 7.430, and write $v_R(u) := \ln \ell_R(u)$ and $v_R^{[\mathcal{U}]} := \ln \ell_R^{[\mathcal{U}]}$. If the Step 5 acceptance holds at these scales—namely $\mathcal{W}$ satisfies the decay/plateau conditions of eqs. (7.591) and (7.602) at $v_R(u)$ (and uniformly at $v_R^{[\mathcal{U}]}$)—and the Step 3 robust-slope/curvature guards eqs. (7.539), (7.542) and (7.548) together with the state–coherence guard eq. (7.566) are enforced on the IR window anchored there, then*

$$v_\star(u) \;\le\; v_R(u), \qquad \sup_{u\in\mathcal{U}} v_\star(u) \;\le\; v_R^{[\mathcal{U}]}.$$

Proof. By eq. (7.626), a point (u, v) lies in the mask $\mathfrak{M}_{\mathrm{mask}}$ iff the five guards C1–C5 hold: robust slope (C1), bounded curvature (C2), state coherence (C3), small Ward residual (C4), and Nielsen flatness of on–shell data (C5). At $v = v_R(u)$, Step 5 acceptance gives C4 (decay and $\mathcal{W} \le \varepsilon_{\mathrm{plat}}$, cf. eq. (7.602)). By hypothesis, the Step 3 guards eqs. (7.542) and (7.548) provide C1 and C2 on the same IR window (the LP_τ-based trend and curvature bounds), while eq. (7.566) supplies C3 (uniform in V). Finally, the gauge-flatness requirement C5 is imposed via eq. (7.607) on that IR window. Hence all guards are satisfied at $(u, v_R(u))$, so $(u, v_R(u)) \in \mathfrak{M}_{\mathrm{mask}}$. By upward closure in v (Proposition 7.524), the vertical section is an upper set; therefore $v_\star(u) \le v_R(u)$.

For the uniform statement, by the definition of $\ell_R^{[\mathcal{U}]}$ (Definition 7.430), the same acceptance holds at $v_R^{[\mathcal{U}]}$ *for all* $u \in \mathcal{U}$, hence $(u, v_R^{[\mathcal{U}]}) \in \mathfrak{M}_{\mathrm{mask}}$ for every $u \in \mathcal{U}$. Thus $v_\star(u) \le v_R^{[\mathcal{U}]}$ for all $u \in \mathcal{U}$, and taking the supremum in u yields $\sup_{u\in\mathcal{U}} v_\star(u) \le v_R^{[\mathcal{U}]}$. □

Reading guide. The restoration length anchors the IR window where the identity ledger closes (small $\mathcal{W}$) and where the law's low–pass trend is monotone with controlled curvature while the state remains coherent. Since the mask's vertical slices are upper sets, the frontier $v_\star(u)$ cannot sit *above* the anchor: once the guards hold at v_R, they hold for all larger v, forcing $v_\star(u) \le v_R(u)$ (and uniformly on $\mathcal{U}$).

Definition 7.526 (UV–strip mask and certified half–space)**.** Let $\mathcal{U} = [u_{\min}, u_{\max}]$ be an admissible UV strip. Define the *restricted mask*

$$\mathfrak{M}_{\mathrm{mask}}^{[\mathcal{U}]} := \{(u, v) \in \mathfrak{M}_{\mathrm{mask}} :\; u \in \mathcal{U}\},$$

where $\mathfrak{M}_{\mathrm{mask}}$ is given pointwise by the guards in eq. (7.626). The associated *certified IR half–space* is

$$\mathcal{H}_{\mathrm{IR}}^{[\mathcal{U}]} := \Big\{(u, v) :\; u \in \mathcal{U},\; v \ge v_\star^{[\mathcal{U}]}\Big\}, \qquad v_\star^{[\mathcal{U}]} := \sup_{u\in\mathcal{U}} v_\star(u),$$

with $v_\star$ the vertical frontier of Proposition 7.524. By Proposition 7.524, $\mathcal{H}_{\mathrm{IR}}^{[\mathcal{U}]} \subseteq \mathfrak{M}_{\mathrm{mask}}^{[\mathcal{U}]}$, and by Corollary 7.525, $v_\star^{[\mathcal{U}]} \le \ln \ell_R^{[\mathcal{U}]}$ with $\ell_R^{[\mathcal{U}]}$ from Definition 7.430.

Reading guide (outside the definition). The restricted mask trims $\mathfrak{M}_{\mathrm{mask}}$ to a UV band $\mathcal{U}$. The number $v_\star^{[\mathcal{U}]}$ is the worst (largest) frontier across $u \in \mathcal{U}$; the resulting epigraph $\mathcal{H}_{\mathrm{IR}}^{[\mathcal{U}]}$ is guaranteed to lie inside the restricted mask by upward closure in v (Proposition 7.524) and is anchored by the uniform restoration scale (Corollary 7.525).

Lemma 7.527 (Basis/gauge/scheme safety of the mask)**.** *Evaluate all diagnostics in eq.* (7.626) *on shell and on the BRST physical sector using the projector* $\mathsf{P}_{\mathrm{phys}}$ *(so poles/residues are read from* $\mathbb{H}_{\mathrm{phys}} = \mathsf{P}_{\mathrm{phys}}\,\mathbb{H}\,\mathsf{P}_{\mathrm{phys}}$*). Then:*

1. (Gauge safety) $\mathfrak{M}_{\mathrm{mask}}$ *is invariant—up to the audit tolerance* $\varepsilon_{\mathrm{gauge}}$*—under admissible gauge–parameter changes* $\xi \in \mathcal{X}_{\mathrm{adm}}$ *(cf. eq.* (7.607)*, Proposition 7.458 and Corollary 7.459).*
2. (Basis/scheme safety) $\mathfrak{M}_{\mathrm{mask}}$ *is invariant under admissible field–basis changes and renormalisation–scheme reparameterisations (cf. Lemma 7.451).*
3. (Scale choice for μ) *The choice* $\mu = \kappa/\ell$ *changes* $v = \ln \ell$ *by a rigid shift* $v \mapsto v + \ln \kappa$*; hence* $\mathfrak{M}_{\mathrm{mask}}$ *is independent of* κ *modulo the translation guaranteed by Lemma 7.523*(3)*.*

Proof. (1) Gauge safety. On shell and on $\mathcal{H}_{\mathrm{phys}}$ the Nielsen identities imply vanishing gauge slopes for quoted on–shell data:

$$\sup_{\xi \in \mathcal{X}_{\mathrm{adm}}} \left|\partial_\xi m_\alpha^2\right| = 0, \qquad \sup_{\xi \in \mathcal{X}_{\mathrm{adm}}} \left|\partial_\xi Z_\alpha\right| = 0,$$

as encoded in eq. (7.607) (see also [117, 243, 442]). Moreover, the law/state visibilities and the Ward residual are computed from BRST–invariant correlators (after $\mathsf{P}_{\mathrm{phys}}$ projection) and, by Proposition 7.458 and Corollary 7.459, are invariant under admissible gauge variations up to the declared tolerance $\varepsilon_{\mathrm{gauge}}$. Since each guard in eq. (7.626) depends either on spectral on–shell data or on such invariant correlators, the pass/fail outcome of all five guards is unchanged (within $\varepsilon_{\mathrm{gauge}}$). Hence membership in $\mathfrak{M}_{\mathrm{mask}}$ is gauge–safe.

(2) Basis/scheme safety. Admissible basis redefinitions act by (blockwise) similarity on the quadratic operator, leaving the spectrum of $\mathbb{H}_{\mathrm{phys}}$ isospectral and transporting residues covariantly; admissible scheme changes correspond to smooth reparameterisations of renormalised quantities. By Lemma 7.451 and Proposition 7.458, the objects entering eq. (7.626) (robust trend/curvature of $\mathsf{LP}_\tau \Lambda_{\mathrm{law}}$, the state visibility bound, $\mathcal{W}$, and on–shell poles/residues) are invariant under these transformations, so the mask is unchanged.

(3) Scale choice for μ*.* Setting $\mu = \kappa/\ell$ with $\kappa > 0$ shifts the v–coordinate by a constant $\Delta = \ln \kappa$. By Lemma 7.523 (3), the mask is equivariant under rigid shifts $v \mapsto v + \Delta$, so κ only translates $\mathfrak{M}_{\mathrm{mask}}$ in v without changing its shape or any epigraphic statements (e.g., $v_\star$, $\mathcal{H}_{\mathrm{IR}}^{[\mathcal{U}]}$). □

Reading guide (outside the lemma). The mask depends only on BRST–physical, on–shell, and Nielsen–stable quantities: that makes it robust to gauge choices; similarity/renormalisation moves don't change spectra or the low–pass law trend; and changing the μ–convention merely shifts the v origin, which Lemma 7.523 shows cannot affect certification.

Discrete mask (audit path).

Definition 7.528 (Grid–level mask and morphological clean–up)**.** On grids $\{u_m\}_{m=1}^{M}$, $\{v_k\}_{k=1}^{K}$ (with $v_k = \ln \ell_k$), define the boolean indicator

$$\widehat{\chi}_{mk} := \mathbf{1}\begin{bmatrix} \widehat{\varsigma}_{\mathrm{law}}^{(\tau)}(u_m, v_k) \geq \varsigma_{\min}, \\ \left|\widehat{\partial_v^2}(\mathsf{LP}_\tau \Lambda_{\mathrm{law}})\right| \leq \varepsilon_{\mathrm{curv}}, \\ \sup_V |1 - \widehat{\Lambda}_{\mathrm{state}}| \leq \varepsilon_{\mathrm{plat}}, \\ \widehat{\mathcal{W}} \leq \varepsilon_{\mathrm{plat}}, \\ \left|\widehat{\partial_\xi m_\alpha^2}\right|, \left|\widehat{\partial_\xi Z_\alpha}\right| \leq \varepsilon_{\mathrm{gauge}} \end{bmatrix},$$

where hats denote the discrete estimators from eqs. (7.547) and (7.608) and Definition 7.516. Apply an optional 3×3 majority filter in (m, k) (topology–preserving "opening/closing") to suppress one–pixel holes/spikes consistent with the curvature/TV guards. Set $\widehat{\mathfrak{P}}_{\mathrm{QIII}} := \{(u_m, v_k) : \widehat{\chi}_{mk} = 1\}$.

Proposition 7.529 (Consistency of the discrete mask)**.** *Under the regularity assumptions of Steps 7.8.2–7.8.6 and using second–order centered estimators in v and in ξ, there exist grid spacings $(\Delta u, \Delta v, \Delta\xi)$ small enough such that, for every compact rectangle $K \subset \mathbb{R}^2$,*

$$d_{\mathrm{H}}\Big(\widehat{\mathfrak{M}}_{\mathrm{mask}} \cap K \ , \quad \mathfrak{M}_{\mathrm{mask}} \cap K \Big) \xrightarrow[\Delta u, \Delta v, \Delta\xi \to 0]{} 0,$$

i.e. the discrete mask converges to the pointwise mask in the Hausdorff metric locally on $\mathbb{R}^2$.

Proof. Recall the pointwise mask $\mathfrak{M}_{\mathrm{mask}}$ from eq. (7.626) is the finite intersection of five closed guard sets (robust slope, curvature, state coherence, Ward residual, and Nielsen flatness). By Lemma 7.523(1), $\mathfrak{M}_{\mathrm{mask}}$ is closed.

Step 1: Uniform, second–order control of the guard functionals. For each guard introduce a continuous "margin" functional $f_r(u, v)$ so that the guard is $f_r(u, v) \geq 0$:

$$\begin{aligned} f_1 &:= \varsigma_{\mathrm{law}}^{(\tau)} - \varsigma_{\min}, \qquad f_2 := \varepsilon_{\mathrm{curv}} - \left|\partial_v^2(\mathsf{LP}_\tau \Lambda_{\mathrm{law}})\right|, \\ f_3 &:= \varepsilon_{\mathrm{plat}} - \sup_V |1 - \Lambda_{\mathrm{state}}(u, v; V)|, \qquad f_4 := \varepsilon_{\mathrm{plat}} - \mathcal{W}(u, v), \\ f_{5m} &:= -\sup_{\xi \in \mathcal{X}_{\mathrm{adm}}} \left|\partial_\xi m_\alpha^2\right|, \quad f_{5Z} := -\sup_{\xi \in \mathcal{X}_{\mathrm{adm}}} \left|\partial_\xi Z_\alpha\right|. \end{aligned}$$

Steps 7.8.2–7.8.6 provide (i) a stationary low-pass LP_τ with second–order accuracy in v for trends/curvatures (preserving constants/linears up to $\mathcal{O}(\tau^2)$), (ii) centered finite-difference estimators for v-derivatives with $\mathcal{O}((\Delta v)^2)$ bias, and (iii) compactly probed suprema in V and $\mathcal{X}_{\mathrm{adm}}$ (nets with mesh $\Delta V, \Delta\xi$). Hence there exists $\varepsilon(\Delta u, \Delta v, \Delta\xi) \to 0$ such that on any compact K,

$$\sup_{(u,v)\in K} \max_r \left| f_r(u, v) - f_r^{\Delta}(u, v) \right| \ \leq \ \varepsilon(\Delta u, \Delta v, \Delta\xi),$$

where f_r^{Δ} denotes the corresponding discrete/estimated margin (centered in v and ξ, with the low-pass applied on the grid). This uses continuity of the underlying maps and the maximum theorem for suprema over compact index sets in V and $\mathcal{X}_{\mathrm{adm}}$.

Step 2: Set inclusions via margin thickening. For $\eta > 0$, define the inner/outer thickenings

$$\mathfrak{M}_{\mathrm{mask}}^{(-\eta)} := \bigcap_r \{f_r \geq \eta\}, \qquad \mathfrak{M}_{\mathrm{mask}}^{(+\eta)} := \bigcap_r \{f_r \geq -\eta\}.$$

If $\varepsilon(\Delta u, \Delta v, \Delta\xi) \leq \eta$, then for every (u, v):

$$\big(f_r \geq \eta\big) \implies \big(f_r^{\Delta} \geq 0\big), \qquad \big(f_r^{\Delta} \geq 0\big) \implies \big(f_r \geq -\eta\big).$$

Intersecting over all guards yields the set sandwich on K:

$$\mathfrak{M}_{\mathrm{mask}}^{(-\eta)} \cap K \;\subseteq\; \widehat{\mathfrak{M}}_{\mathrm{mask}} \cap K \;\subseteq\; \mathfrak{M}_{\mathrm{mask}}^{(+\eta)} \cap K.$$

Step 3: From sandwiches to Hausdorff convergence. Because each $\{f_r \geq c\}$ is closed and the f_r are continuous, $\mathfrak{M}_{\mathrm{mask}}^{(-\eta)} \downarrow \mathfrak{M}_{\mathrm{mask}}$ and $\mathfrak{M}_{\mathrm{mask}}^{(+\eta)} \uparrow \mathfrak{M}_{\mathrm{mask}}$ as $\eta \downarrow 0$, with convergence in the Painlevé–Kuratowski sense. On a compact K, this implies convergence in the Hausdorff metric:

$$d_{\mathrm{H}}\Big(\mathfrak{M}_{\mathrm{mask}}^{(-\eta)} \cap K,\ \mathfrak{M}_{\mathrm{mask}} \cap K\Big) \to 0, \quad d_{\mathrm{H}}\Big(\mathfrak{M}_{\mathrm{mask}}^{(+\eta)} \cap K,\ \mathfrak{M}_{\mathrm{mask}} \cap K\Big) \to 0.$$

Combining with the sandwich gives

$$\begin{aligned} 0 &\leq d_{\mathrm{H}}\Big(\widehat{\mathfrak{M}}_{\mathrm{mask}} \cap K,\ \mathfrak{M}_{\mathrm{mask}} \cap K\Big) \\ &\leq \max\left\{ \begin{aligned} &d_{\mathrm{H}}\Big(\mathfrak{M}_{\mathrm{mask}}^{(+\eta)} \cap K,\ \mathfrak{M}_{\mathrm{mask}} \cap K\Big), \\ &d_{\mathrm{H}}\Big(\mathfrak{M}_{\mathrm{mask}}^{(-\eta)} \cap K,\ \mathfrak{M}_{\mathrm{mask}} \cap K\Big) \end{aligned} \right\} \\ &\xrightarrow[\eta \downarrow 0]{} 0. \end{aligned}$$

Step 4: Grid union vs. point samples (pixel geometry). If $\widehat{\mathfrak{M}}_{\mathrm{mask}}$ is implemented as occupied grid *cells* (union of rectangles of size $\Delta u \times \Delta v$) rather than just marked nodes, the Hausdorff error gains an additional additive $O(\max\{\Delta u, \Delta v\})$ that vanishes as the grid refines. Optional binary morphological closing with structuring element of radius $o(\max\{\Delta u, \Delta v\})$ leaves the limit unchanged and only improves the finite-mesh bound.

Altogether, choosing $(\Delta u, \Delta v, \Delta\xi)$ so that $\varepsilon(\Delta u, \Delta v, \Delta\xi) \leq \eta$ and then letting $\eta \to 0$ proves the claim. □

Reading guide. Package each guard in eq. (7.626) as a continuous margin $f_r \geq 0$. The Step-2–6 estimators approximate f_r with sup-norm error $\to 0$ on any fixed window, so the discrete pass set sits between inner/outer thickenings of the true mask. Those thickenings collapse to the mask as the margins go to zero, which turns into Hausdorff convergence of the discrete mask to $\mathfrak{M}_{\mathrm{mask}}$ on the window.

Plateau mask outputs.

Definition 7.530 (Reporting artefacts for the mask)**.** All reports are *restricted* to the certified region defined by the mask $\mathfrak{M}_{\mathrm{mask}}$ of eq. (7.626) and its discrete estimator $\widehat{\mathfrak{M}}_{\mathrm{mask}}$.

(i) Trend heatmap with certified overlays.
A heatmap of the robust law slope $\varsigma^{(\tau)}_{\mathrm{law}}(u,v)$ (from eqs. (7.539), (7.542) and (7.548)), overlaid with the discrete mask $\widehat{\mathfrak{M}}_{\mathrm{mask}}$. Superimpose: (a) the frontier $v_\star(u)$ (see Proposition 7.524), (b) the certified half–space line $v = v_\star^{[\mathcal{U}]}$ for a chosen UV strip $\mathcal{U}$ (Definition 7.526), (c) the IR window $\mathcal{I}_{\mathrm{IR}}$ shading and markers at $v = \ln \ell_R(u)$ and $v = \ln \ell_R^{[\mathcal{U}]}$ (Definitions 7.428 and 7.430 and Corollary 7.525).

(ii) Windowed tables (mask-restricted diagnostics).
Tables of windowed summaries computed *on* $\widehat{\mathfrak{M}}_{\mathrm{mask}}$:

- **(a)** Median and quantiles of the Ward residual $\mathcal{W}$ (eqs. (7.591) and (7.602)).
- **(b)** Median and quantiles of on–shell gauge slopes $\partial_\xi m_\alpha^2$, $\partial_\xi Z_\alpha$ (Nielsen flatness; eq. (7.607)); report spreads across $\xi \in \mathcal{X}_{\mathrm{adm}}$.
- **(c)** Curvature proxy summaries $\left|\partial_v^2(\mathsf{LP}_\tau \Lambda_{\mathrm{law}})\right|$ (bounded as in eq. (7.542)).
- **(d)** Robustness spreads across admissible band knobs ($P_x(\lambda_{\mathrm{UV}})$, $K_{\mathrm{IR}}^{(\ell)}$) and gauges (Nielsen–projected), as required by Lemma 7.527.

Include the marginal coverage statistics $\Phi_{v/u}$ and deficits (δ_u, δ_v) for the reported windows (cf. the mask marginals used in Step 7).

(iii) Cross-reference ledger (mask-gated).
A ledger linking upstream quantities, all evaluated *within* $\widehat{\mathfrak{M}}_{\mathrm{mask}}$:

- **(a)** Step 2: slope positivity and monotone trend acceptance (eqs. (7.539), (7.542) and (7.548)).
- **(b)** Step 3: restoration lengths (ℓ_R, $\ell_R^{[\mathcal{U}]}$) (Definitions 7.428 and 7.430 and Corollary 7.525).
- **(c)** Step 4: irrelevance exponents $y_{\tilde{B}}$ (quoted only where the mask certifies law healing and state coherence).
- **(d)** Step 6: spectral reconvergence (poles/residues read from $\mathbb{H}_{\mathrm{phys}}$; consistency with Lemma 7.527).

Intuition. $\mathfrak{M}_{\mathrm{mask}}$ is the "all–green" locus where the five guards of eq. (7.626) hold simultaneously. By Proposition 7.524 it is upward-closed in v (coarse-graining helps), hence admits the epigraph description via $v_\star(u)$; Definition 7.526 then provides a certified half–space for any UV strip. Restricting plots, tables, and cross-references to $\widehat{\mathfrak{M}}_{\mathrm{mask}}$ ensures every downstream exponent, scale, and spectral statement is quoted on a common, certified plateau.

Integrated (marginal) compliance. In addition to pointwise certification via eq. (7.626), we require that axis–integrated readouts inherit the mask's protection, suppressing isolated outliers. Let meas$(\cdot)$ denote one–dimensional Lebesgue measure and fix measurable windows $I_u \subset \mathbb{R}$ and $I_v \subset \mathbb{R}$. Define the integrated (marginal) compliance sets

$$\boxed{\mathfrak{M}^{(v)}_{\mathrm{mask}} := \Big\{ v \in I_v : \ \mathrm{meas}\Big(\{ u \in I_u : \ (u,v) \in \mathfrak{M}_{\mathrm{mask}}\}\Big) \ \geq \ (1-\delta_u)\ \mathrm{meas}(I_u)\Big\},} \tag{7.627}$$

$$\boxed{\mathfrak{M}^{(u)}_{\mathrm{mask}} := \Big\{ u \in I_u : \ \mathrm{meas}\Big(\{ v \in I_v : \ (u,v) \in \mathfrak{M}_{\mathrm{mask}}\}\Big) \ \geq \ (1-\delta_v)\ \mathrm{meas}(I_v)\Big\},} \tag{7.628}$$

for small audit tolerances $\delta_u, \delta_v \in [0,1)$. *Rule:* any report aggregated along the v–axis (resp. u–axis) must be restricted to $\mathfrak{M}^{(v)}_{\mathrm{mask}}$ (resp. $\mathfrak{M}^{(u)}_{\mathrm{mask}}$). Equivalently, if $\Phi_v(u; I_v)$ and $\Phi_u(v; I_u)$ are the marginal coverages from eq. (7.626), then $\mathfrak{M}^{(v)}_{\mathrm{mask}} = \{v : \inf_{u \in I_u} \Phi_u(v; I_u) \geq 1 - \delta_u\}$ and analogously for $\mathfrak{M}^{(u)}_{\mathrm{mask}}$.

Discrete mask estimator (audit path). On a rectangular grid $\{(u_m, v_k)\}$, define the per–guard booleans from eq. (7.626):

$$\chi_1 := \mathbf{1}\{\varsigma^{(\tau)}_{\mathrm{law}} \geq \varsigma_{\min}\}, \quad \chi_2 := \mathbf{1}\{\, |\partial_v^2(\mathsf{LP}_\tau \Lambda_{\mathrm{law}})| \leq \varepsilon_{\mathrm{curv}}\,\}, \quad \chi_3 := \mathbf{1}\{\, \sup_V |1 - \Lambda_{\mathrm{state}}| \leq \varepsilon_{\mathrm{plat}}\,\},$$

$$\chi_4 := \mathbf{1}\{\, \mathcal{W} \leq \varepsilon_{\mathrm{plat}}\,\}, \quad \chi_5 := \mathbf{1}\Big\{ \sup_{\xi \in \mathcal{X}_{\mathrm{adm}}} |\partial_\xi m_\alpha^2| = 0, \quad \sup_{\xi \in \mathcal{X}_{\mathrm{adm}}} |\partial_\xi Z_\alpha| = 0 \text{ within } \varepsilon_{\mathrm{gauge}}\ \forall \alpha\Big\},$$

and set $\chi_{\mathrm{mask}} := \prod_{r=1}^5 \chi_r \in \{0,1\}$. Optionally pre–filter the underlying fields along v with the same stationary low–pass LP_τ used in eqs. (7.537) to (7.539) to suppress salt–and–pepper while preserving first derivatives (bias $\mathcal{O}(\tau^2)$).

The discrete mask is the occupied grid set

$$\boxed{\widehat{\mathfrak{M}}_{\mathrm{mask}} := \Big\{(u_m, v_k) : \ \chi_{\mathrm{mask}}(u_m, v_k) = 1\Big\}.} \tag{7.629}$$

For integrated summaries on the grid, use the discrete marginals

$$\widehat{\mathfrak{M}}^{(v)}_{\mathrm{mask}} := \Big\{ v_k \in I_v : \ \frac{1}{|I_u|_{\mathrm{grid}}} \#\{m : \ u_m \in I_u, \ (u_m, v_k) \in \widehat{\mathfrak{M}}_{\mathrm{mask}}\} \ \geq \ 1 - \delta_u \Big\},$$

$$\widehat{\mathfrak{M}}^{(u)}_{\mathrm{mask}} := \Big\{ u_m \in I_u : \ \frac{1}{|I_v|_{\mathrm{grid}}} \#\{k : \ v_k \in I_v, \ (u_m, v_k) \in \widehat{\mathfrak{M}}_{\mathrm{mask}}\} \ \geq \ 1 - \delta_v \Big\},$$

where $|I_\bullet|_{\mathrm{grid}}$ denotes the number of grid points in the window. By Proposition 7.529 these discrete marginals converge to eqs. (7.627) and (7.628) as the mesh refines.

Intuition (outside the blocks). The marginals gate *axis–wise* readouts so a few bad columns/rows cannot dominate integrated summaries: at least a $(1-\delta)$ fraction along the orthogonal axis must pass the full mask. The discrete construction mirrors the five guards in eq. (7.626), preserves the same low–pass used for robust slope/curvature, and—by our convergence result—recovers the continuous marginals in the fine–mesh limit.

Acceptance statement (plateau certification). We adopt the continuous mask $\mathfrak{M}_{\mathrm{mask}} \subset \mathbb{R}^2$ given pointwise by the five guards in eq. (7.626), and its indicator $\chi_{\mathrm{mask}}(u,v) = \mathbf{1}[(u,v) \in \mathfrak{M}_{\mathrm{mask}}]$. Axis–integrated readouts use the marginal coverages $\Phi_v(u; I_v)$ and $\Phi_u(v; I_u)$ defined in eqs. (7.627) and (7.628), together with the uncovered–mass deficits $\delta_v(I_u, I_v)$ and $\delta_u(I_u, I_v)$ introduced alongside them.

Certification rule. We declare the plateau certified when the following hold:

- **Nonempty and IR–anchored.** The mask is nonempty and intersects the IR–anchored band:
$$\mathfrak{M}_{\mathrm{mask}} \neq \varnothing, \qquad \mathfrak{M}_{\mathrm{mask}} \cap \{(u,v) : v \geq \ln \ell_R\} \neq \varnothing,$$
with ℓ_R the restoration length of Definition 7.428. Equivalently, by the upward-closure epigraph form $\mathfrak{M}_{\mathrm{mask}} = \{(u,v) : v \geq v_\star(u)\}$ from Proposition 7.524, this demands $v_\star(u) \leq \ln \ell_R(u)$ (cf. Corollary 7.525).

- **Nontrivial marginals on reported windows.** For the chosen windows I_u, I_v, the marginal compliance sets are nonempty:
$$\{u \in I_u : \Phi_v(u; I_v) > 0\} \neq \varnothing, \qquad \{v \in I_v : \Phi_u(v; I_u) > 0\} \neq \varnothing,$$
ensuring that axis–integrated summaries inherit mask protection rather than being driven by isolated outliers. When desired, we additionally enforce coverage budgets $\delta_u, \delta_v \leq \delta_{\max}$ (user-chosen tolerances), again computed via eqs. (7.627) and (7.628).

- **Gauge and instrumentation robustness.** Within the admissible gauges $\mathcal{X}_{\mathrm{adm}}$ and admissible band knobs (P_x, K_{IR}), the mask is invariant up to the audit tolerance $\varepsilon_{\mathrm{gauge}}$:
$$\sup_{(u,v)} \left| \chi_{\mathrm{mask}}^{(P_x, K_{\mathrm{IR}}, \xi)}(u,v) - \chi_{\mathrm{mask}}^{(P_x', K_{\mathrm{IR}}', \xi')}(u,v) \right| \;\leq\; \varepsilon_{\mathrm{gauge}},$$
consistent with the Nielsen flatness requirement eq. (7.607) and the safety statements in Lemma 7.527 (gauge, basis/scheme, and rigid v–shift invariances; see also [117, 243, 442]).

Outputs for Step 7. We report only quantities restricted to the certified region (mask-gated) and its discrete estimator $\widehat{\mathfrak{M}}_{\mathrm{mask}}$:

1. **Certified map.** A heatmap of the robust law slope $\varsigma_{\mathrm{law}}^{(\tau)}(u,v)$ (eqs. (7.539), (7.542) and (7.548)), overlaid with the discrete mask $\widehat{\mathfrak{M}}_{\mathrm{mask}}$, the frontier $v_\star(u)$ (Proposition 7.524), the certified half–space bound $v = v_\star^{[\mathcal{U}]}$ for the chosen UV strip $\mathcal{U}$ (Definition 7.526), and the IR window shading $\mathcal{I}_{\mathrm{IR}}$ with markers at $v = \ln \ell_R(u)$ and $v = \ln \ell_R^{[\mathcal{U}]}$ (Definitions 7.428 and 7.430 and Corollary 7.525).

2. **Mask–restricted tables.** Windowed summaries computed *on* $\widehat{\mathfrak{M}}_{\mathrm{mask}}$: medians/quantiles of the Ward residual $\mathcal{W}$ (eqs. (7.591) and (7.602)); medians/quantiles of the on–shell gauge slopes $\partial_\xi m_\alpha^2, \partial_\xi Z_\alpha$ (Nielsen flatness, eq. (7.607)); and the curvature proxy $|\partial_v^2(\mathsf{LP}_\tau \Lambda_{\mathrm{law}})|$. Include robustness spreads across admissible (P_x, K_{IR}) and $\xi \in \mathcal{X}_{\mathrm{adm}}$ per Lemma 7.527. Also report marginal coverage fractions Φ_v, Φ_u and deficits (δ_u, δ_v) on the declared windows.

3. **Cross-reference ledger (mask-gated).** A concise ledger linking upstream ingredients, all evaluated within $\widehat{\mathfrak{M}}_{\mathrm{mask}}$:

   (a) Step 2–3 slope positivity and monotone trend acceptance (eqs. (7.539), (7.542) and (7.548));

(b) Step 3 restoration lengths $(\ell_R, \ell_R^{[\mathcal{U}]})$ (Definitions 7.428 and 7.430 and Corollary 7.525);

(c) Step 4 irrelevance exponents $y_{\tilde{B}}$ (quoted only where the mask certifies law healing/state coherence; cf. RG texts [389, 91, 82]);

(d) Step 6 spectral reconvergence (poles/residues read from $\mathbb{H}_{\text{phys}}$, consistent with gauge/basis safety, Lemma 7.527, and perturbation control [115]).

Concluding remarks for Step 7. Step 7 is where we stop spot–checking individual diagnostics and instead *certify a common IR stage* on which all downstream claims must stand. Concretely, we carve out a geometric region in scale–space—the mask $\mathfrak{M}_{\text{mask}} \subset \mathbb{R}^2$ over $(u, v) = (\ln \lambda_{\text{UV}}, \ln \ell)$—defined by the simultaneous pass of the five guards in eq. (7.626). Each guard targets a distinct failure mode: the *robust slope* $\varsigma_{\text{law}}^{(\tau)}$ asserts genuine law healing rather than noisy drift, the *curvature bound* keeps that trend plateau–like (the low–pass suppresses UV grit), *state coherence* excludes QII–type volume fragility, the *Ward residual* $\mathcal{W}$ enforces identity closure, and *Nielsen flatness* (zero gauge slopes) guarantees that quoted poles/residues are physical rather than gauge artifacts. Their intersection—the "all-green" pixels—is the certified mask.

Beyond existence, Step 7 explains the *shape* and *use* of this region. Coarse–graining helps once healing has begun, so vertical slices are upper sets; equivalently, the mask is the epigraph of a frontier $v_\star(u)$ (Proposition 7.524). This yields a clean interpretation: for any (u, v) with $v \geq v_\star(u)$, the point is certified. The *restoration lengths* $\ell_R(u)$ (and their UV–uniform counterpart on a strip) *anchor* this epigraph: $v_\star(u) \leq \ln \ell_R(u)$ and $\sup_{u \in \mathcal{U}} v_\star(u) \leq \ln \ell_R^{[\mathcal{U}]}$ (Corollary 7.525), so certification is IR–anchored by a scale already computed. For any chosen UV strip $\mathcal{U}$, the worst frontier $v_\star^{[\mathcal{U}]}$ furnishes a *certified IR half-space* lying inside the mask (Definition 7.526); this is the safest domain for aggregating exponents and spectra.

Step 7 also guards against *integration artifacts*. All integrated readouts (fits, tables, averages) are restricted to the marginal compliance sets $\mathfrak{M}_{\text{mask}}^{(v)}$ and $\mathfrak{M}_{\text{mask}}^{(u)}$ (eqs. (7.627) and (7.628)), which require that at least a $(1 - \delta)$ fraction of columns/rows pass the full mask. This prevents a narrow stripe of "good" points from legitimising a global claim.

Finally, Step 7 justifies the *finite-mesh audit* actually used in practice. The discrete estimator $\widehat{\mathfrak{M}}_{\text{mask}}$, built from the same five guards with centered, second-order estimators and the same low–pass, converges in the Hausdorff sense to the pointwise mask on any fixed window (Proposition 7.529). Together with gauge/basis/scheme safety (Lemma 7.527) and rigid-shift invariance in v (Lemma 7.523), this renders the certification *instrument-robust*.

In short, Step 7 is the *certification gate*: it turns local diagnostics (slope, curvature, state, Ward, Nielsen) into a single certified plateau region that is upward-closed, IR-anchored, robust to instrumentation and gauge, safe under integration, and faithfully approximated on the grid. Every downstream exponent, scale, or spectral multiplet should be quoted *within* that same $\mathfrak{M}_{\text{mask}}$.

7.8.8 Physicality Strengthens with ℓ

Objective. Across the certified mask $\mathfrak{M}_{\mathrm{mask}}$ (cf. eq. (7.626) and Step 7.8.7), we require that basic *physicality* properties not only hold but *improve* as coarse–graining increases ($v = \ln \ell$ grows): (i) nonnegativity of physical pole masses with a nondecreasing *mass margin*; (ii) strong hyperbolicity of the principal symbol of the quadratic IR dynamics with a nondecreasing *hyperbolicity margin*; and (iii) spectral positivity of two–point functions with a nondecreasing *spectral floor*. All statements are made *on shell* on the BRST physical sector and are *Nielsen–projected* (gauge–flat within tolerance; eq. (7.607) and Lemma 7.527). Dependence on ℓ is continuous by standard Kato perturbation theory for isolated poles/eigenvalues [115] (see also [255, 14, 91, 82]).

Mass operators: nonnegativity with improving margin. Let $\mathbb{H}_{\mathrm{phys}} = \mathsf{P}_{\mathrm{phys}}\,\mathbb{H}\,\mathsf{P}_{\mathrm{phys}}$ denote the quadratic operator on the physical sector (cf. eqs. (6.332) and (7.10)); read *on–shell* poles and residues from the corresponding two–point blocks. For $(u, v) \in \mathfrak{M}_{\mathrm{mask}}$, let $\mathcal{I}_{\mathrm{phys}}(u, v)$ index the Nielsen–stable physical poles at (u, v).

Definition 7.531 (Mass margin)**.** For $(u, v) \in \mathfrak{M}_{\mathrm{mask}}$, define

$$\boxed{\Delta_{\mathrm{mass}}(u, v) := \min_{i \in \mathcal{I}_{\mathrm{phys}}(u,v)} m_i^2(u, v) \;\geq\; 0.} \tag{7.630}$$

Reading guide (outside the definition). Δ_{mass} is the gap to tachyonic violation among all quoted physical poles; gauge safety follows from eq. (7.607) and Lemma 7.527. Continuity in v holds by Kato continuity of simple/isolated poles [115].

Definition 7.532 (Strengthening with ℓ)**.** We say the mass sector *strengthens* along $v = \ln \ell$ on $\mathfrak{M}_{\mathrm{mask}}$ if

$$\boxed{\partial_v\, \Delta_{\mathrm{mass}}(u, v) \;\geq\; 0 \quad \text{for all } (u, v) \in \mathfrak{M}_{\mathrm{mask}}.} \tag{7.631}$$

Reading guide. Within the certified mask, the law's robust monotone trend (eqs. (7.542) and (7.548)) couples with identity closure (eq. (7.602)) to prevent pole collisions from crossing into $m^2 < 0$ and to push the minimum m^2 upward (nondecreasing Δ_{mass}). Gauge/basis invariance of the quoted masses is ensured by Lemma 7.527.

Lemma 7.533 (Continuity, local Lipschitzness, and a.e. differentiability of masses)**.** *Fix $u \in \mathbb{R}$. Assume the hypotheses of Lemma 7.492 for the* band–projected, Nielsen–projected *two–point kernel on the certified region, i.e. along $(u, v) \in \mathfrak{M}_{\mathrm{mask}}$ the projected inverse two–point operator (or the relevant quadratic form on $\mathcal{H}_{\mathrm{phys}}$) is a C^1 Kato family in v, uniformly on compact momentum neighborhoods of each on–shell pole, with simple (algebraic multiplicity 1) physical poles except possibly at isolated values of v. Then for every physical channel $i \in \mathcal{I}_{\mathrm{phys}}(u, v)$:*

1. *$v \mapsto m_i^2(u, v)$ is* continuous *on $\{v : (u, v) \in \mathfrak{M}_{\mathrm{mask}}\}$.*
2. *There exists a closed, at most countable set $\Sigma(u) \subset \{v : (u, v) \in \mathfrak{M}_{\mathrm{mask}}\}$ (pole collisions/thresholds) such that on each connected component of the complement the map $v \mapsto m_i^2(u, v)$ is* continuously differentiable *and in fact* locally Lipschitz.[15]

[15] Local Lipschitz follows from uniform bounds on $\|\partial_v \mathcal{K}\|$ and a uniform lower bound on ∂_{m^2} of the dispersion functional at simple poles; see the proof.

Consequently, the mass margin $\Delta_{\mathrm{mass}}(u,v) := \min_{i\in\mathcal{I}_{\mathrm{phys}}(u,v)} m_i^2(u,v)$ *(cf. Definition 7.531 and eq. (7.630)) is* continuous *in* v *on* $\mathfrak{M}_{\mathrm{mask}}$ *and is* differentiable almost everywhere *in* v.

Proof. Setup. Work on the BRST physical sector with $\mathbb{H}_{\mathrm{phys}} = \mathsf{P}_{\mathrm{phys}}\mathbb{H}\mathsf{P}_{\mathrm{phys}}$ (cf. eqs. (6.332) and (7.10)); all quoted poles/residues are Nielsen–projected and gauge–flat per eq. (7.607) and Lemma 7.527. For fixed u and $(u,v) \in \mathfrak{M}_{\mathrm{mask}}$, each physical pole $m_i^2(u,v)$ is realized either as a simple eigenvalue of a Hermitian (or at least normal) effective mass operator on $\mathcal{H}_{\mathrm{phys}}$ at zero momentum, or as a simple real root of a scalar dispersion map $D_i(u,v;m^2) = 0$ obtained by analytic Fredholm reduction of the two–point kernel along the ith channel.

(1) Continuity. By the Kato–tracking hypothesis (Lemma 7.492), the relevant operator family (or dispersion map) is C^1 in v *in operator norm* on compact momentum neighborhoods of each pole. For simple poles, the spectral projection depends continuously on v and the associated eigenvalue (or real root) depends continuously on parameters (Kato, Thm. II-4.10, Thm. II-5.4 in [115]). Thus $v \mapsto m_i^2(u,v)$ is continuous wherever the pole remains simple. When simplicity fails, it can do so only at isolated v by the same perturbation theory; continuity of a continuous selection of eigenvalues across such points still holds (see Rellich's theorem and Kato's continuous selection results). Hence $m_i^2(u,\cdot)$ is continuous on $\{v : (u,v) \in \mathfrak{M}_{\mathrm{mask}}\}$.

(2) Piecewise C^1 *and local Lipschitz off a discrete set.* Let $\Sigma(u)$ denote the set of v where (i) a physical pole ceases to be simple (collision/threshold) or (ii) the index set $\mathcal{I}_{\mathrm{phys}}(u,v)$ changes by birth/decay of a channel. Under the C^1–in–v Kato hypothesis and simplicity elsewhere, $\Sigma(u)$ is closed and at most countable (indeed discrete on compact v–intervals). On each connected component $J \subset \{v : (u,v) \in \mathfrak{M}_{\mathrm{mask}}\} \setminus \Sigma(u)$, the Implicit Function Theorem applies to $D_i(u,v;m^2) = 0$ because $\partial_{m^2} D_i \neq 0$ at a simple pole; therefore $m_i^2(u,\cdot) \in C^1(J)$. Moreover, by the Hellmann–Feynman/Implicit differentiation formula,

$$\partial_v m_i^2(u,v) \;=\; -\frac{\partial_v D_i(u,v;m_i^2)}{\partial_{m^2} D_i(u,v;m_i^2)}\,,$$

and the denominator is bounded away from 0 on compact subintervals of J (simplicity + continuity). Since $\partial_v D_i$ is continuous, $\partial_v m_i^2$ is bounded on compact subintervals, whence m_i^2 is locally Lipschitz on J. (If one works directly with a selfadjoint effective mass operator $M_{\mathrm{eff}}(u,v)$, the standard Kato bound $\left|\partial_v m_i^2\right| \le \|\partial_v M_{\mathrm{eff}}\|$ gives the same conclusion; see [115, Sec. VII-1].)

(3) Consequences for the minimum. The mass margin $\Delta_{\mathrm{mass}}(u,v) = \min_i m_i^2(u,v)$ is the pointwise minimum of finitely many continuous functions (finitely many physical channels at fixed u in the band), hence is continuous. On each J as above, the m_i^2 are locally Lipschitz, so Δ_{mass} is also locally Lipschitz (the pointwise minimum of finitely many locally Lipschitz functions is locally Lipschitz). By Rademacher's theorem, locally Lipschitz functions are differentiable almost everywhere, proving the a.e. differentiability claim. Where the minimum is attained by a unique simple channel $i_\star$, one has the identification

$$\partial_v \Delta_{\mathrm{mass}}(u,v) = \partial_v m_{i_\star}^2(u,v) \quad \text{for a.e. } v \text{ with unique minimiser } i_\star.$$

All statements above are made on $\mathfrak{M}_{\mathrm{mask}}$, so they inherit gauge/basis/scheme safety from Lemma 7.527 and the required IR regularity (monotone trend, bounded curvature, and Ward

plateau) from Step 7 (eqs. (7.542), (7.548) and (7.602)), which guarantee stability of the simple–pole regime away from $\Sigma(u)$. □

Remark 7.534 (NG/pNG channels)**.** If exact NG or admissible pseudo–NG modes are present (cf. Definition 7.508 and Proposition 7.509), Δ_{mass} may approach 0 without violating eq. (7.630). In such cases it is convenient to monitor the *tachyon deficit*

$$\Theta_{\mathrm{mass}}(u,v) := \max\{0,\, -\min_i m_i^2(u,v)\},$$

and require $\partial_v \Theta_{\mathrm{mass}}(u,v) \le 0$ on $\mathfrak{M}_{\mathrm{mask}}$ (no growth of any negative excursion). This condition is equivalent to nondecreasing Δ_{mass} wherever all $m_i^2 \ge 0$, and it cleanly accommodates NG/pNG channels that sit at (or flow to) the origin.

Principal symbol: strong hyperbolicity with improving margin.

Definition 7.535 (Principal symbol, strong hyperbolicity, margin, and strengthening)**.** Work on the BRST physical sector with quadratic IR dynamics reduced at $(u,v) \in \mathfrak{M}_{\mathrm{mask}}$ (cf. eqs. (6.332), (7.10) and (7.626) and Lemma 7.527) to the first–order system

$$\partial_t \Phi \;+\; \sum_{j=1}^{d} A^j(u,v)\, \partial_j \Phi \;+\; B(u,v)\, \Phi \;=\; 0, \tag{7.632}$$

and define the *principal symbol* in direction $\hat{\boldsymbol{k}} \in \mathbb{S}^{d-1}$ by

$$A(u,v,\hat{\boldsymbol{k}}) \;:=\; \sum_{j=1}^{d} \hat{k}_j\, A^j(u,v). \tag{7.633}$$

All objects are computed on–shell and Nielsen–projected (gauge–flat within tolerance; eq. (7.607) and Lemma 7.527).

(Strong hyperbolicity). The system is *strongly hyperbolic* at (u,v) if there exists a positive–definite *symmetriser* $H(u,v,\hat{\boldsymbol{k}})$ with bounded condition number (uniformly in $\hat{\boldsymbol{k}}$) such that

$$H(u,v,\hat{\boldsymbol{k}})\, A(u,v,\hat{\boldsymbol{k}}) \;\text{ is symmetric for all }\; \hat{\boldsymbol{k}} \in \mathbb{S}^{d-1},$$

which yields an energy estimate and a well–posed Cauchy problem for (7.632) (see, e.g., [14, 91]).

(Hyperbolicity margin). For $(u,v) \in \mathfrak{M}_{\mathrm{mask}}$, the (direction–uniform) hyperbolicity margin is

$$\boxed{\Delta_{\mathrm{hyp}}(u,v) := \inf_{\hat{\boldsymbol{k}} \in \mathbb{S}^{d-1}} \lambda_{\min}\!\left(H(u,v,\hat{\boldsymbol{k}})^{1/2}\, \frac{A(u,v,\hat{\boldsymbol{k}}) + A(u,v,\hat{\boldsymbol{k}})^{\top}}{2}\, H(u,v,\hat{\boldsymbol{k}})^{1/2} \right) \;\ge\; 0.} \tag{7.634}$$

We say the system is *uniformly strongly hyperbolic* on $\mathfrak{M}_{\mathrm{mask}}$ if $\Delta_{\mathrm{hyp}}(u,v) > 0$ for all $(u,v) \in \mathfrak{M}_{\mathrm{mask}}$.

(Strengthening with ℓ). Hyperbolicity *strengthens* along $v = \ln \ell$ on the mask if

$$\boxed{\partial_v\, \Delta_{\mathrm{hyp}}(u,v) \;\ge\; 0 \qquad \text{for all } (u,v) \in \mathfrak{M}_{\mathrm{mask}}.} \tag{7.635}$$

Reading guide. Δ_{hyp} is the smallest symmetrised principal–symbol eigenvalue over directions; it quantifies the energy coercivity constant for the first–order reduction. Gauge/basis/scheme safety follows from Lemma 7.527; continuity in v along the mask follows from the same Kato continuity assumptions used for poles (Step 7), applied to $A^j(u,v)$ and $H(u,v,\hat{\boldsymbol{k}})$.

Spectral positivity: Källén–Lehmann density with improving floor. For each physical channel α, the (band–projected, on–shell) two–point function admits a Källén–Lehmann representation with a positive spectral density $\rho_\alpha(\mathfrak{s};u,v)\ge 0$ [14, 255, 82]. Fix an IR window $\mathfrak{S}_{\mathrm{IR}}$ in the spectral variable $\mathfrak{s}$ that contains all quoted poles and the first continuum threshold. All quantities are computed on the BRST physical sector and Nielsen–projected (gauge–flat within tolerance; eq. (7.607) and Lemma 7.527), with (u,v) restricted to the certified mask $\mathfrak{M}_{\mathrm{mask}}$ of eq. (7.626).

Definition 7.536 (Spectral floor and strengthening)**.** For $(u,v)\in\mathfrak{M}_{\mathrm{mask}}$, define the *spectral floor*

$$\boxed{\Delta_{\mathrm{spec}}(u,v) := \inf_{\alpha}\ \inf_{\mathfrak{s}\in\mathfrak{S}_{\mathrm{IR}}}\ \rho_\alpha(\mathfrak{s};u,v)\ \ \ge\ 0,} \tag{7.636}$$

and require *strengthening* along $v=\ln\ell$:

$$\boxed{\partial_v\,\Delta_{\mathrm{spec}}(u,v)\ \ge\ 0\quad\text{for all }(u,v)\in\mathfrak{M}_{\mathrm{mask}}.} \tag{7.637}$$

Remark 7.537 (Continuity, lower semicontinuity, and low–pass safety)**.** Under Step 5 regularity (band projection is positive and norm–continuous in (u,v)) and Kato continuity for the pole sector [115], the two–point functions vary continuously in operator norm on compact $\mathfrak{s}$–windows; the associated spectral measures $\rho_\alpha(\cdot;u,v)$ therefore vary continuously in the weak* topology on finite measures [115, Ch. VI]. Consequently, $(u,v)\mapsto\Delta_{\mathrm{spec}}(u,v)$ is *lower semicontinuous* on $\mathfrak{M}_{\mathrm{mask}}$ and is *a.e. differentiable in* v on any compact v–interval (via local Lipschitz of infima of weak*–continuous families on compact sets).

If an optional low–pass LP_τ in v is applied to the two–point function in a way that preserves the Löwner (positive–semidefinite) order—e.g., a convex combination $\int K_\tau(v-\tilde v)\,G(\tilde v)\,d\tilde v$ with $K_\tau\ge 0$, $\int K_\tau=1$—then the induced spectral densities satisfy

$$\rho_\alpha^{(\tau)}(\mathfrak{s};u,v)=\int K_\tau(v-\tilde v)\,\rho_\alpha(\mathfrak{s};u,\tilde v)\,d\tilde v\ \ \ge\ 0,$$

and Jensen's inequality gives $\Delta_{\mathrm{spec}}^{(\tau)}(u,v)\ \ge\ \inf_{\tilde v}\Delta_{\mathrm{spec}}(u,\tilde v)$, so the inequality eq. (7.637) is not weakened by such filtering.

Unified acceptance (physicality improves on the mask).

Definition 7.538 (Unified physicality margin)**.** On the certified mask $\mathfrak{M}_{\mathrm{mask}}$ (cf. eq. (7.626)), define

$$\boxed{\Delta_{\mathrm{phys}}(u,v) := \min\Big\{\Delta_{\mathrm{mass}}(u,v),\ \Delta_{\mathrm{hyp}}(u,v),\ \Delta_{\mathrm{spec}}(u,v)\Big\},\qquad (u,v)\in\mathfrak{M}_{\mathrm{mask}},} \tag{7.638}$$

where Δ_{mass} is the mass margin (Definition 7.531 and eq. (7.630)), Δ_{hyp} the hyperbolicity margin (Definition 7.535 and eq. (7.634)), and Δ_{spec} the spectral floor (Definition 7.536 and eq. (7.636)).

Lemma 7.539 (Nonnegativity, monotonicity, and regularity of the unified margin)**.** *Assume on* $\mathfrak{M}_{\text{mask}}$ *that each component margin is nonnegative and* strengthens *along* $v = \ln \ell$*:*

$$\begin{aligned} \Delta_{\text{mass}},\ \Delta_{\text{hyp}},\ \Delta_{\text{spec}} &\geq 0, \\ \partial_v \Delta_{\text{mass}} &\geq 0 && (\textit{eq. } (7.631)), \\ \partial_v \Delta_{\text{hyp}} &\geq 0 && (\textit{eq. } (7.635)), \\ \partial_v \Delta_{\text{spec}} &\geq 0 && (\textit{eq. } (7.637)). \end{aligned}$$

Then:

1. (Nonnegativity) $\Delta_{\text{phys}} \geq 0$ *on* $\mathfrak{M}_{\text{mask}}$.
2. (Monotonicity) *For each fixed* u, $v \mapsto \Delta_{\text{phys}}(u, v)$ *is nondecreasing on the vertical section* $\{v : (u, v) \in \mathfrak{M}_{\text{mask}}\}$.
3. (Regularity) Δ_{phys} *is lower semicontinuous on* $\mathfrak{M}_{\text{mask}}$*; moreover, on any compact* v*–interval where the components are locally Lipschitz in* v, Δ_{phys} *is locally Lipschitz and hence a.e. differentiable in* v*. At points where the minimiser is unique,*

$$\partial_v \Delta_{\text{phys}}(u, v) = \partial_v \Delta_j(u, v) \quad \textit{for } j \in \{\text{mass}, \text{hyp}, \text{spec}\} \textit{ realising the minimum.}$$

Proof. (1) Immediate: a pointwise minimum of nonnegative functions is nonnegative.

(2) Let $f_j(v) := \Delta_j(u, v)$ for $j \in \{\text{mass}, \text{hyp}, \text{spec}\}$. Each f_j is nondecreasing by the cited strengthening conditions. Hence $f(v) := \min_j f_j(v)$ is nondecreasing: for $v' \geq v$, $f(v') = \min_j f_j(v') \geq \min_j f_j(v) = f(v)$.

(3) *Lower semicontinuity.* Each component is l.s.c. on $\mathfrak{M}_{\text{mask}}$: (i) Δ_{mass} is continuous in v on the mask by Lemma 7.533 (Kato tracking for simple poles [115]); (ii) Δ_{hyp} is continuous since $(u, v, \hat{\boldsymbol{k}}) \mapsto H^{1/2}(\frac{A+A^\top}{2})H^{1/2}$ is continuous and the map $(M \mapsto \lambda_{\min}(M))$ is continuous on symmetric matrices, while the inf over compact $\mathbb{S}^{d-1}$ preserves continuity; (iii) Δ_{spec} is l.s.c. by weak* continuity of spectral measures (Step 5 regularity) as in Remark 7.537. The pointwise minimum of finitely many l.s.c. functions is l.s.c., so Δ_{phys} is l.s.c.

Local Lipschitz and a.e. differentiability. On any compact v–interval contained in a vertical section of the mask: - Δ_{mass} is locally Lipschitz away from a discrete set where pole simplicity can fail, by implicit differentiation/Hellmann–Feynman and operator C^1 dependence (see proof of Lemma 7.533 and [115, Sec. VII-1]); - Δ_{hyp} is locally Lipschitz if A^j, H are C^1 in v (then $\lambda_{\min}$ of a symmetric matrix depending C^1 on parameters is locally Lipschitz; the uniform inf over compact directions preserves it); - Δ_{spec} is locally Lipschitz if the two–point kernel depends Lipschitzly in operator norm on v over the IR window, which is satisfied under the Step 5 norm–continuity and stationary low–pass in v (convolution with a fixed kernel is 1–Lipschitz w.r.t. the sup norm). The pointwise minimum of finitely many locally Lipschitz functions is locally Lipschitz; by Rademacher, it is a.e. differentiable. When the minimiser is unique, the derivative equals that component's derivative; with multiple active minima, the one–sided Dini derivatives lie between the extremes of the active components. □

Remarks. (i) All statements are on–shell and Nielsen–projected, hence gauge/basis/scheme safe by Lemma 7.527 (see also eq. (7.607)). (ii) By Proposition 7.524, vertical sections of the mask are upper sets; coupled with monotonicity of Δ_{phys}, the unified margin provides a conservative, IR–anchored acceptance to accompany the restoration lengths (Corollary 7.525). (iii) Optional low–pass in v that preserves Löwner order does not degrade Δ_{spec} (and thus not Δ_{phys}), cf. Remark 7.537.

Definition 7.540 (Plateau floor of physicality)**.** The *plateau floor* is defined by

$$\boxed{\Delta_{\mathrm{phys}}^{(\mathrm{plateau})} := \inf_{(u,v)\in\mathfrak{M}_{\mathrm{mask}}} \Delta_{\mathrm{phys}}(u,v) \geq 0.} \tag{7.639}$$

Corollary 7.541 (Frontier reduction and IR anchoring of the floor)**.** *Under Section 7.8.8 and Proposition 7.524, vertical sections of $\mathfrak{M}_{\mathrm{mask}}$ are upper sets and $v \mapsto \Delta_{\mathrm{phys}}(u,v)$ is nondecreasing; hence*

$$\Delta_{\mathrm{phys}}^{(\mathrm{plateau})} \;=\; \inf_{u} \inf_{v\geq v_\star(u)} \Delta_{\mathrm{phys}}(u,v) \;=\; \inf_{u} \Delta_{\mathrm{phys}}\Big(u,\, v_\star(u)\Big),$$

where $v_\star$ is the frontier from Proposition 7.524. Moreover, by Corollary 7.525,

$$\Delta_{\mathrm{phys}}\Big(u,\, v_\star(u)\Big) \;\leq\; \Delta_{\mathrm{phys}}\Big(u,\, \ln \ell_R(u)\Big),$$

so the floor is IR-anchored by the restoration lengths.

Proof. For fixed u, $\mathfrak{M}_{\mathrm{mask}}(u) = \{v \geq v_\star(u)\}$ by Proposition 7.524. Monotonicity in v (from Section 7.8.8) implies $\inf_{v\geq v_\star(u)} \Delta_{\mathrm{phys}}(u,v) = \Delta_{\mathrm{phys}}(u, v_\star(u))$. Taking the infimum over u gives the first equality. The second statement uses $v_\star(u) \leq \ln \ell_R(u)$ from Corollary 7.525 and monotonicity in v. □

Remark 7.542 (pNG/NG allowance)**.** When NG/pNG channels are present (cf. Definition 7.508 and Proposition 7.509), implement the mass component via the tachyon–deficit proxy of Remark 7.534. The functionals Δ_{phys} and $\Delta_{\mathrm{phys}}^{(\mathrm{plateau})}$ are then unchanged while allowing $\min_i m_i^2 \to 0^+$.

Remark 7.543 (Gauge/basis/scheme safety)**.** All statements are on–shell and Nielsen–projected, hence gauge–flat within $\varepsilon_{\mathrm{gauge}}$ (cf. eq. (7.607) and Corollary 7.459); admissible basis/scheme changes act by similarity on the physical sector (Lemma 7.451). Therefore Δ_{phys} and $\Delta_{\mathrm{phys}}^{(\mathrm{plateau})}$ are invariant within the same tolerance (see also Lemma 7.527).

Discrete estimators and audit path.

Definition 7.544 (Discrete estimators on the (u_m, v_k) grid)**.** Work on the (u_m, v_k) grid of eq. (7.629), with uniform v–step h (so $h = \Delta v$) and band–admissible momentum/spectral grids as needed. All diagnostics are computed on–shell and Nielsen–projected (gauge–flat within tolerance; eq. (7.607) and Lemma 7.527) and restricted to the certified mask $\mathfrak{M}_{\mathrm{mask}}$ (eq. (7.626)).

(a) Mass margin and v–slope. Using per–channel discrete pole estimators $\widehat{m}_i^2(u_m, v_k)$ and (if needed) residues $\widehat{Z}_i(u_m, v_k)$ from the Step 6 spectral estimator (two–point block on $\mathbb{H}_{\mathrm{phys}}$), set

$$\boxed{\begin{aligned} \widehat{\Delta}_{\mathrm{mass}}(u_m, v_k) &:= \min_i \widehat{m}_i^2(u_m, v_k), \\ \widehat{\partial_v \Delta_{\mathrm{mass}}}(u_m, v_k) &:= \frac{\widehat{\Delta}_{\mathrm{mass}}(u_m, v_{k+1}) - \widehat{\Delta}_{\mathrm{mass}}(u_m, v_{k-1})}{2h}. \end{aligned}} \tag{7.640}$$

(b) Hyperbolicity margin and v–slope. Choose a finite angular grid $\{\hat{\boldsymbol{k}}_r\} \subset \mathbb{S}^{d-1}$ (e.g., quasi-uniform spherical design). Form principal symbols $A_{mkr} := A(u_m, v_k, \hat{\boldsymbol{k}}_r)$ (Definition 7.535), pick admissible symmetrisers $H_{mkr} \succ 0$ (consistent with Definition 7.535), and define

$$\boxed{\begin{aligned} \widehat{\Delta}_{\mathrm{hyp}}(u_m, v_k) &:= \min_r \ \lambda_{\min}\Big(H_{mkr}^{1/2} \, \tfrac{A_{mkr} + A_{mkr}^{\top}}{2} \, H_{mkr}^{1/2} \Big), \\ \widehat{\partial_v \Delta_{\mathrm{hyp}}}(u_m, v_k) &:= \frac{\widehat{\Delta}_{\mathrm{hyp}}(u_m, v_{k+1}) - \widehat{\Delta}_{\mathrm{hyp}}(u_m, v_{k-1})}{2h}. \end{aligned}} \tag{7.641}$$

(c) Spectral floor and v–slope. From the (band–projected, on–shell) two–point data in channel α, build a nonnegative spectral estimator $\widehat{\rho}_\alpha(\mathfrak{s}; u_m, v_k) \geq 0$ (e.g., constrained maximum-entropy; or rational/Prony fits with an explicit positivity constraint on the density over the IR window). On a spectral grid $\{\mathfrak{s}_s\} \subset \mathfrak{S}_{\mathrm{IR}}$,

$$\boxed{\begin{aligned} \widehat{\Delta}_{\mathrm{spec}}(u_m, v_k) &:= \min_\alpha \ \min_s \ \widehat{\rho}_\alpha(\mathfrak{s}_s; u_m, v_k), \\ \widehat{\partial_v \Delta_{\mathrm{spec}}}(u_m, v_k) &:= \frac{\widehat{\Delta}_{\mathrm{spec}}(u_m, v_{k+1}) - \widehat{\Delta}_{\mathrm{spec}}(u_m, v_{k-1})}{2h}. \end{aligned}} \tag{7.642}$$

(d) Low–pass before differencing. Apply the same stationary low–pass in v as eqs. (7.537) and (7.538) to each discrete margin sequence $\{\widehat{\Delta}_\bullet(u_m, v_k)\}_k$ before forming centered differences; this suppresses ripples while preserving first derivatives (bias $\mathcal{O}(\tau^2) + \mathcal{O}(h^2)$). At grid boundaries, use second–order one-sided differences or shrink the stencil.

Lemma 7.545 (Consistency and rates)**.** *Assume:*

(i) Simple–pole regularity.
The hypotheses of Definition 7.516 hold with simple poles and C^3 regularity in v in a neighborhood of each physical pole on the mask (cf. Lemma 7.533), so that the Step 6 pole estimators are second–order accurate in the momentum-grid variable p^2.

(ii) Principal symbol regularity.
$A(u, v, \hat{\boldsymbol{k}})$ is C^2 in $(v, \hat{\boldsymbol{k}})$ on $\mathfrak{M}_{\mathrm{mask}} \times \mathbb{S}^{d-1}$ and admits a uniformly positive symmetriser $H(u, v, \hat{\boldsymbol{k}})$ as in Definition 7.535.

(iii) Spectral density estimator.
For each channel α, the spectral estimator $\widehat{\rho}_\alpha(\mathfrak{s}; u, v)$ is positivity–preserving and consistent in the sup–norm on the IR window $\mathfrak{S}_{\mathrm{IR}}$, i.e.

$$\sup_{\mathfrak{s} \in \mathfrak{S}_{\mathrm{IR}}} \big| \widehat{\rho}_\alpha(\mathfrak{s}; u, v) - \rho_\alpha(\mathfrak{s}; u, v) \big| \leq \varepsilon_\rho$$

with $\varepsilon_\rho \to 0$ as the spectral grid refines.

Then, for the discrete margins of Definition 7.544 and as the grids are refined,

$$\widehat{\Delta}_{\mathrm{mass}} = \Delta_{\mathrm{mass}} + \mathcal{O}\Big((\Delta p^2)^2\Big), \qquad \widehat{\Delta}_{\mathrm{hyp}} = \Delta_{\mathrm{hyp}} + \mathcal{O}\Big(\Delta\hat{k}\Big), \qquad \widehat{\Delta}_{\mathrm{spec}} = \Delta_{\mathrm{spec}} + \mathcal{O}\Big(\varepsilon_\rho + \Delta\mathfrak{s}\Big),$$

uniformly on compact (u, v)–windows contained in $\mathfrak{M}_{\mathrm{mask}}$. Moreover, away from the discrete exceptional set where channel minima switch (cf. $\Sigma(u)$ in Lemma 7.533), the centered v–differences satisfy

$$\widehat{\partial_v \Delta_\bullet}(u_m, v_k) = \partial_v \Delta_\bullet(u_m, v_k) + \mathcal{O}(h^2), \qquad \bullet \in \{\mathrm{mass}, \mathrm{hyp}, \mathrm{spec}\},$$

with $h = \Delta v$. If an optional low–pass LP_τ is applied in v (eqs. (7.537) and (7.538)), choose $\tau = \mathcal{O}(h)$ so that the additional bias $\mathcal{O}(\tau^2)$ preserves the overall $\mathcal{O}(h^2)$ rate.

Proof. Mass margin. By Definition 7.531 and eq. (7.630), $\Delta_{\mathrm{mass}} = \min_i m_i^2$. Under Assumption (1) (Step 6 estimators with simple poles and C^3 regularity), each discrete pole location obeys $\widehat{m}_i^2 = m_i^2 + \mathcal{O}((\Delta p^2)^2)$ uniformly on compact windows (second–order consistency in the p^2 grid). The inequality

$$\Big|\min_i a_i - \min_i b_i\Big| \;\le\; \max_i |a_i - b_i|$$

then yields $\widehat{\Delta}_{\mathrm{mass}} - \Delta_{\mathrm{mass}} = \min_i \widehat{m}_i^2 - \min_i m_i^2 = \mathcal{O}((\Delta p^2)^2)$.

Hyperbolicity margin. Let $\varphi(u, v, \hat{\boldsymbol{k}}) := \lambda_{\min}\Big(H^{1/2}\, \frac{A + A^\top}{2}\, H^{1/2}\Big)$ so that by eq. (7.634), $\Delta_{\mathrm{hyp}}(u, v) = \inf_{\hat{\boldsymbol{k}} \in \mathbb{S}^{d-1}} \varphi(u, v, \hat{\boldsymbol{k}})$. Assumption (2) implies φ is C^1 in $\hat{\boldsymbol{k}}$ on the compact sphere (composition of C^2 maps with the continuous smallest–eigenvalue functional on symmetric matrices). For an ε–net $\{\hat{\boldsymbol{k}}_r\}$ with mesh $\Delta\hat{k}$, uniform continuity of φ gives

$$\begin{aligned} 0 \le \widehat{\Delta}_{\mathrm{hyp}} - \Delta_{\mathrm{hyp}} &= \min_r \varphi(u, v, \hat{\boldsymbol{k}}_r) - \inf_{\hat{\boldsymbol{k}}} \varphi(u, v, \hat{\boldsymbol{k}}) \\ &\le \sup_{\mathrm{dist}(\hat{\boldsymbol{k}}, \{\hat{\boldsymbol{k}}_r\}) \le \Delta\hat{k}} \Big|\varphi(u, v, \hat{\boldsymbol{k}}) - \varphi(u, v, \hat{\boldsymbol{k}}_r)\Big| \\ &= \mathcal{O}(\Delta\hat{k}). \end{aligned}$$

(If, in addition, the minimiser $\hat{\boldsymbol{k}}_\star$ is unique and interior with nondegenerate Hessian in the tangent space, a local quadratic fit in direction can refine the rate to $\mathcal{O}((\Delta\hat{k})^2)$; see the remark below.)

Spectral floor. By Definition 7.536 and eq. (7.636), $\Delta_{\mathrm{spec}} = \inf_\alpha \inf_{\mathfrak{s} \in \mathfrak{S}_{\mathrm{IR}}} \rho_\alpha(\mathfrak{s})$. On a grid $\{\mathfrak{s}_s\}$ with mesh $\Delta\mathfrak{s}$ and with a sup–norm consistent, positivity–preserving estimator,

$$\widehat{\Delta}_{\mathrm{spec}} = \min_\alpha \min_s \widehat{\rho}_\alpha(\mathfrak{s}_s) = \min_\alpha \min_s \Big[\rho_\alpha(\mathfrak{s}_s) + \mathcal{O}(\varepsilon_\rho)\Big] = \min_\alpha \Big[\inf_{\mathfrak{s}} \rho_\alpha(\mathfrak{s}) + \mathcal{O}(\Delta\mathfrak{s}) + \mathcal{O}(\varepsilon_\rho)\Big],$$

using uniform continuity of ρ_α on the compact IR window (Step 5 regularity) to pass from grid minima to the continuum infimum with error $\mathcal{O}(\Delta\mathfrak{s})$. Taking the minimum in α preserves the bound, yielding the stated rate.

Centered v–differences. On mask–restricted windows, Lemma 7.533 gives piecewise C^1 (indeed locally Lipschitz) dependence in v for m_i^2 and hence for Δ_{mass}; Assumption (2) yields the same for Δ_{hyp}; Remark 7.537 yields lower semicontinuity—and, under the mild Lipschitz–in–v assumption for the two–point kernel (Step 5), local Lipschitzness—for Δ_{spec}. On any subinterval avoiding the discrete set where minima switch, Taylor's theorem with the standard centered-difference formula gives

$$\frac{f(v+h)-f(v-h)}{2h} = f'(v) + \mathcal{O}(h^2)$$

for $f = \Delta_{\text{mass}}, \Delta_{\text{hyp}}, \Delta_{\text{spec}}$ provided $f \in C^3$ there (or f'' Lipschitz). This yields the $\mathcal{O}(h^2)$ rate. If the optional low–pass LP_τ is applied and preserves first moments (eq. (7.538)), the induced additional bias is $\mathcal{O}(\tau^2)$; choosing $\tau = \mathcal{O}(h)$ retains the overall $\mathcal{O}(h^2)$ rate. □

Remark (directional refinement). If the minimising direction $\hat{\boldsymbol{k}}_\star(u,v)$ is unique and non-degenerate (Hessian positive–definite on the tangent space), one may perform a local quadratic fit of $\varphi(u,v,\hat{\boldsymbol{k}})$ using neighboring samples around the discrete argmin to obtain $\widehat{\Delta}_{\text{hyp}} = \Delta_{\text{hyp}} + \mathcal{O}((\Delta\hat{k})^2)$. This mirrors the standard parabolic-refinement trick used in angular optimisation.

Lemma 7.546 (Gauge/basis invariance and mask restriction)**.** *All discrete margins and their centered v–slopes from Definition 7.544 are evaluated* on shell *and within the Nielsen–projected physical subspace. Then, across admissible gauges $\xi \in \mathcal{X}_{\text{adm}}$ and admissible basis/scheme changes, they are invariant within the audit tolerance $\varepsilon_{\text{gauge}}$ (cf. eq. (7.607), Corollary 7.459, and Lemmas 7.451 and 7.527). All audit checks are restricted to $(u_m, v_k) \in \mathfrak{M}_{\text{mask}}$ (cf. eqs. (7.626) and (7.629)).*

Proof. Restriction to the mask. By construction (eq. (7.629)), discrete evaluations are performed only at grid points in the certified set $\widehat{\mathfrak{M}}_{\text{mask}}$, which converges to $\mathfrak{M}_{\text{mask}}$ on windows (Proposition 7.529); hence all checks are mask–restricted.

Gauge invariance (within tolerance). On the physical sector, the Nielsen identities enforce vanishing gauge slopes for quoted on–shell quantities (eq. (7.607)), and Corollary 7.459 asserts gauge insensitivity within $\varepsilon_{\text{gauge}}$. Therefore: (i) Mass sector: m_i^2 and hence $\Delta_{\text{mass}} = \min_i m_i^2$ vary by at most $\mathcal{O}(\varepsilon_{\text{gauge}})$; the inequality $\left|\min_i a_i - \min_i b_i\right| \le \max_i |a_i - b_i|$ preserves the bound. Centered v–differences inherit the same tolerance since the forward/backward differences cancel common-mode perturbations and the remaining variation is bounded by the sup–norm change of Δ_{mass} across adjacent k. (ii) Hyperbolicity: A^j and admissible symmetrisers H computed from on–shell, BRST–invariant data (projected onto $\mathbb{H}_{\text{phys}}$) vary within $\mathcal{O}(\varepsilon_{\text{gauge}})$ by Lemma 7.527. The map $(A,H) \mapsto \lambda_{\min}\!\left(H^{1/2}\frac{A+A^\top}{2}H^{1/2}\right)$ is 1–Lipschitz up to a uniform constant with respect to the operator norm on $(A, H^{1/2})$, hence Δ_{hyp} varies by at most $C\,\varepsilon_{\text{gauge}}$ uniformly in direction; the discrete minimum over $\hat{\boldsymbol{k}}_r$ preserves the bound, as do centered differences in v. (iii) Spectral floor: The (band–projected, on–shell) two–point function is gauge–flat on the physical sector within $\varepsilon_{\text{gauge}}$ (Corollary 7.459 and Lemma 7.527); positivity–preserving spectral estimation thus changes $\widehat{\rho}_\alpha$ by at most $\mathcal{O}(\varepsilon_{\text{gauge}})$ in sup–norm on $\mathfrak{S}_{\text{IR}}$, yielding the same bound for $\widehat{\Delta}_{\text{spec}}$ and for its centered v–difference.

Basis/scheme invariance. Admissible field–basis changes act by (blockwise) similarity on the quadratic operator and transport residues covariantly; admissible renormalisation–scheme changes reparameterise renormalised quantities smoothly (cf. Lemmas 7.451 and 7.527). Spectra and principal symbols on $\mathbb{H}_{\mathrm{phys}}$ are therefore invariant under these transformations; the functionals defining Δ_{mass}, Δ_{hyp}, and Δ_{spec} (and their centered differences) depend only on these invariants, whence invariance within the same tolerance. □

Audit artefacts (reporting checklist). Alongside the discrete curves for the margins and their v–slopes (cf. Definition 7.544 and eqs. (7.640) to (7.642)), report the following audit artefacts, all restricted to the certified mask $\mathfrak{M}_{\mathrm{mask}}$ (eqs. (7.626) and (7.629)):

(i) Grids and filters. The step sizes and meshes (h, Δp^2, $\Delta\hat{k}$, $\Delta\mathfrak{s}$) and the low–pass width τ used before centered differencing (cf. eqs. (7.537) and (7.538)).

(ii) Symmetriser construction. How H was obtained for the principal symbol (e.g. energy inner product; Cholesky factorisation), consistent with Definition 7.535.

(iii) Spectral estimation method. The estimator for $\widehat{\rho}_\alpha$ and positivity constraints imposed over $\mathfrak{S}_{\mathrm{IR}}$ (cf. Definition 7.536); include the achieved sup–norm error/bias if available.

(iv) Mask–restricted summaries. Windowed minima and centered v–slopes of each margin $\widehat{\Delta}_{\mathrm{mass}}, \widehat{\Delta}_{\mathrm{hyp}}, \widehat{\Delta}_{\mathrm{spec}}$ on $\widehat{\mathfrak{M}}_{\mathrm{mask}}$, with spreads across admissible (P_x, K_{IR}) and gauges, as justified by Lemmas 7.527, 7.545 and 7.546.

Notes. Choose $\tau = \mathcal{O}(h)$ to preserve the $\mathcal{O}(h^2)$ centered–difference rate (Lemma 7.545). All items are computed on–shell and Nielsen–projected and are invariant across admissible gauges/bases within $\varepsilon_{\mathrm{gauge}}$ (eq. (7.607) and Lemma 7.546).

Compatibility guards.

Definition 7.547 (Evaluation domain and guards)**.** All Step 8 diagnostics are evaluated *only* on the certified mask $\mathfrak{M}_{\mathrm{mask}} \subset \mathbb{R}^2$ as defined in eq. (7.626). In addition, throughout Step 8 we require:

(i) State–coherence guard (uniform in volume).
$\sup_V \left|1 - \Lambda_{\mathrm{state}}(u, v; V)\right| \le \varepsilon_{\mathrm{plat}}$ for all $(u, v) \in \mathfrak{M}_{\mathrm{mask}}$ (cf. eq. (7.566)).

(ii) Nielsen flatness for on–shell inputs (gauge safety).
For all quoted on–shell poles or residues, the Nielsen slopes vanish, $\partial_\xi m_\alpha^2(u, v; \xi) = \partial_\xi Z_\alpha(u, v; \xi) = 0$ within tolerance for every $\xi \in \mathcal{X}_{\mathrm{adm}}$.

Lemma 7.548 (Well–posedness, continuity/Lipschitz regularity, and invariance)**.** *Under Definition 7.547, the Step 8 component margins* Δ_{mass}, Δ_{hyp}, Δ_{spec} *are* well defined *on* $\mathfrak{M}_{\mathrm{mask}}$, continuous in v *and, on any compact* v*-interval inside a vertical section of the mask,* locally Lipschitz and hence a.e. differentiable in v. *They are also* gauge-/basis-/scheme-invariant within $\varepsilon_{\mathrm{gauge}}$. *Consequently, the unified and floor quantities* Δ_{phys} *and* $\Delta_{\mathrm{phys}}^{(\mathrm{plateau})}$ *(cf. Definitions 7.538 and 7.540 and eqs.* (7.638) *and* (7.639)*) are well defined, lower semicontinuous (indeed locally Lipschitz under the same mild assumptions), and invariant within the same tolerance.*

Proof. Domain and on–shell inputs. All evaluations are restricted to $\mathfrak{M}_{\mathrm{mask}}$ (eq. (7.626)), where the five Step 7 guards hold simultaneously. The state–coherence guard ensures uniformity in V for on–shell inputs; Nielsen flatness removes gauge artefacts (eq. (7.607) and Corollary 7.459). Quotations are taken on the BRST physical sector (cf. eqs. (6.332) and (7.10) and Lemma 7.527).

Mass margin. By Definition 7.531 and eq. (7.630), $\Delta_{\mathrm{mass}}(u,v) = \min_{i\in\mathcal{I}_{\mathrm{phys}}} m_i^2(u,v)$. Kato's perturbation theory (simple, isolated poles) plus the Step 7 regularity imply continuity of $v \mapsto m_i^2(u,v)$ and piecewise C^1 (locally Lipschitz) dependence away from a discrete exceptional set (pole collisions/thresholds), cf. Lemma 7.533 and [115]. A pointwise minimum of finitely many continuous (locally Lipschitz) maps is continuous (locally Lipschitz); hence Δ_{mass} has the stated regularity and is a.e. differentiable in v (Rademacher).

Hyperbolicity margin. Let $A(u,v,\hat{\boldsymbol{k}})$ and an admissible symmetriser $H(u,v,\hat{\boldsymbol{k}}) \succ 0$ be as in Definition 7.535. Define

$$\varphi(u,v,\hat{\boldsymbol{k}}) := \lambda_{\min}\Big(H^{1/2}\,\tfrac{A+A^{\top}}{2}\,H^{1/2}\Big).$$

Continuity of A, H in v (Step 8 hypotheses) and continuity of $\lambda_{\min}$ on symmetric matrices imply continuity of φ in $(v,\hat{\boldsymbol{k}})$. Then $\Delta_{\mathrm{hyp}}(u,v) = \inf_{\hat{\boldsymbol{k}}\in\mathbb{S}^{d-1}} \varphi(u,v,\hat{\boldsymbol{k}})$ is continuous in v since the infimum is taken over compact $\mathbb{S}^{d-1}$ of a jointly continuous map. If A, H are C^1 in v, φ is locally Lipschitz, and the uniform infimum over $\hat{\boldsymbol{k}}$ preserves local Lipschitzness (uniform continuity on a compact set), yielding a.e. differentiability.

Spectral floor. By Definition 7.536 and eq. (7.636), $\Delta_{\mathrm{spec}} = \inf_\alpha \inf_{\mathfrak{s}\in\mathfrak{S}_{\mathrm{IR}}} \rho_\alpha(\mathfrak{s};u,v)$. Step 5 regularity (band projection positive and norm–continuous in v) together with Kato continuity in the pole sector ensure weak* continuity of the spectral measures $\rho_\alpha(\cdot;u,v)$ on the compact IR window $\mathfrak{S}_{\mathrm{IR}}$ [115, Ch. VI], so Δ_{spec} is lower semicontinuous in v (cf. Remark 7.537). If, in addition, the two–point kernel depends locally Lipschitzly in operator norm on v over $\mathfrak{S}_{\mathrm{IR}}$, then ρ_α varies locally Lipschitz in the bounded–variation norm, yielding local Lipschitzness (hence a.e. differentiability) of Δ_{spec}.

Invariance (gauge/basis/scheme). Nielsen projection implies gauge–parameter flatness for the quoted on–shell objects within $\varepsilon_{\mathrm{gauge}}$ (eq. (7.607) and Corollary 7.459). Admissible basis/scheme changes act by similarity and smooth reparametrisation on the physical sector (Lemmas 7.451 and 7.527). The functionals defining Δ_{mass} (min of pole masses), Δ_{hyp} (smallest symmetrised principal–symbol eigenvalue over directions), and Δ_{spec} (infimum of positive spectral densities on $\mathfrak{S}_{\mathrm{IR}}$) are invariant under these transformations, up to the audit tolerance.

Unified quantities. $\Delta_{\mathrm{phys}} = \min\{\Delta_{\mathrm{mass}}, \Delta_{\mathrm{hyp}}, \Delta_{\mathrm{spec}}\}$ (Definition 7.538 and eq. (7.638)) inherits lower semicontinuity (and local Lipschitzness under the mild assumptions above) from the components; invariance follows since minima preserve the tolerance bound. The plateau floor $\Delta_{\mathrm{phys}}^{(\mathrm{plateau})} = \inf_{(u,v)\in\mathfrak{M}_{\mathrm{mask}}} \Delta_{\mathrm{phys}}(u,v)$ (Definition 7.540 and eq. (7.639)) is well defined because $\mathfrak{M}_{\mathrm{mask}} \neq \varnothing$ under Step 7 acceptance (cf. acceptance statement) and the infimum of a bounded–below l.s.c. function on a nonempty set is finite; invariance follows from the componentwise invariance. □

Outputs for Step 8. We report four families of artefacts, all *restricted to the certified mask* $\mathfrak{M}_{\mathrm{mask}}$ (cf. eqs. (7.626) and (7.629)) and computed on–shell in the Nielsen–projected sector

(gauge–flat within tolerance; eq. (7.607) and Lemma 7.527):

- **Heatmaps.** Over (u, v), plot the fields $(u, v) \mapsto \Delta_{\mathrm{mass}}(u, v)$, $\Delta_{\mathrm{hyp}}(u, v)$, $\Delta_{\mathrm{spec}}(u, v)$, with $\mathfrak{M}_{\mathrm{mask}}$ overlaid (non–masked regions dimmed). This makes visible where the mass gap, strong–hyperbolicity margin, and spectral floor are simultaneously nonnegative and improving (cf. Definitions 7.531 and 7.536 and eq. (7.634)).

- **Derivative line–cuts.** For representative u–slices, show the centered v–derivatives $\partial_v \Delta_{\mathrm{mass}}$, $\partial_v \Delta_{\mathrm{hyp}}$, $\partial_v \Delta_{\mathrm{spec}}$ restricted to $\mathfrak{M}_{\mathrm{mask}}$, using the discrete estimators eqs. (7.640) to (7.642) (with optional low–pass per eqs. (7.537) and (7.538)). A nonnegative profile on these cuts evidences strengthening along $v = \ln \ell$ (cf. eqs. (7.631), (7.635) and (7.637)).

- **Plateau floor and spreads.** Quote the unified *plateau floor*

$$\Delta_{\mathrm{phys}}^{(\mathrm{plateau})} = \inf_{(u,v)\in\mathfrak{M}_{\mathrm{mask}}} \Delta_{\mathrm{phys}}(u, v),$$

  (cf. Definitions 7.538 and 7.540 and eqs. (7.638) and (7.639)), and accompany it with instrumentation/gauge spreads obtained by re–auditing over admissible (P_x, K_{IR}) and $\xi \in \mathcal{X}_{\mathrm{adm}}$:

$$\mathrm{err}_{\mathrm{inst/gauge}} := \max_{(P_x,K_{\mathrm{IR}}),\,\xi\in\mathcal{X}_{\mathrm{adm}}} \left| \Delta_{\mathrm{phys}}^{(\mathrm{plateau})}\Big[(P_x, K_{\mathrm{IR}}), \xi\Big] - \Delta_{\mathrm{phys}}^{(\mathrm{plateau})}\Big[(P'_x, K'_{\mathrm{IR}}), \xi'\Big] \right|.$$

  By Lemmas 7.539 and 7.546, the floor is nonnegative, monotone in v on vertical sections, and invariant to within $\varepsilon_{\mathrm{gauge}}$.

- **Audit table.** Summarise numerical settings: (a) symmetrisers H for the principal symbol (construction and conditioning; Definition 7.535); (b) spectral reconstructions $\widehat{\rho}$ (fit family, positivity constraints, IR window $\mathfrak{S}_{\mathrm{IR}}$; Definition 7.536); (c) low–pass kernel and width in v; and (d) grid spacings $(h, \Delta p^2, \Delta\hat{k}, \Delta\mathfrak{s})$ used in Definition 7.544. This enables reproducible verification of the rates in Lemma 7.545.

Interpretation. Nonnegative and *improving* mass, hyperbolicity, and spectral–positivity margins across $\mathfrak{M}_{\mathrm{mask}}$ demonstrate that coarse–graining not only restores symmetry but also *stabilises* the IR dynamics: poles move away from tachyonic regions, the first–order reduction retains (and tightens) strong hyperbolicity, and the Källén–Lehmann density maintains a nondecreasing floor in the IR window—consistent with monotone law healing and RG irrelevance ([14, 91, 82, 255, 115]).

Decision rule. Let $\mathcal{D} \subset \mathbb{R}^2$ be the probed (u, v)–domain, $\mathfrak{M}_{\mathrm{mask}}$ the certified plateau mask of eq. (7.626), and ℓ_R the restoration length of Definition 7.428. We classify behaviour into QI–QIV by a mutually exclusive, exhaustive rule, checked in the following order:

1. **QI (already unified).** For all $(u, v) \in \mathcal{D}$,

$$E_{\mathrm{law}}(e^v) \approx 0 \quad \text{and} \quad \sup_V \left|1 - \Lambda_{\mathrm{state}}(u, v; V)\right| \approx 0 \quad \text{within plateau tolerances.}$$

2. **QII (volume decoherence).** The state–coherence guard fails somewhere (cf. eq. (7.566)); i.e. there exists $(u, v) \in \mathcal{D}$ with

$$\sup_V \left|1 - \Lambda_{\text{state}}(u, v; V)\right| > \varepsilon_{\text{plat}},$$

even if the law shows partial healing.

3. **QIII (healing on a certified plateau).** The mask is nonempty and IR-anchored,

$$\mathfrak{M}_{\text{mask}} \neq \varnothing, \qquad \mathfrak{M}_{\text{mask}} \cap \{(u, v) : v \geq \ln \ell_R\} \neq \varnothing,$$

and, on $\mathfrak{M}_{\text{mask}}$,

$$\sup_{(u,v)\in\mathfrak{M}_{\text{mask}}} \left|1 - \Lambda_{\text{law}}(u, v)\right| \leq \varepsilon_{\text{plat}}, \qquad \varsigma^{(\tau)}_{\text{law}}(u, v) \geq \varsigma_{\min} \quad \text{(cf. } eq. \text{ (7.542)).}$$

4. **QIV (persistent explicit breaking).** Otherwise; equivalently, despite the guards there exists $\delta > 0$ such that, for all sufficiently large v,

$$\Lambda_{\text{law}}(u, v) \;\leq\; 1 - \delta \qquad \text{(no approach to unity on any candidate mask).}$$

The evaluation order (1)→(4) makes the outcomes disjoint; in particular, failure of the state–coherence guard in (2) precludes QIII. All checks using pole/residue data are on shell and Nielsen–projected (gauge–flat within $\varepsilon_{\text{gauge}}$; cf. eq. (7.607)).

Lemma 7.549 (Gauge/instrument robustness of the decision)**.** *Under admissible gauge variations and basis/scheme changes with Nielsen projection (Corollary 7.459 and Lemma 7.451) and across admissible instrument choices* (P_x, K_{IR})*, the quadrant outcome in Section 7.8.8 is invariant within the stated tolerances.*

Proof. By Lemma 7.527, the mask $\mathfrak{M}_{\text{mask}}$ and its frontier are invariant (within $\varepsilon_{\text{gauge}}$) under admissible gauge and basis/scheme changes and rigid shifts in v. The state–coherence guard (QII criterion) uses Λ_{state} evaluated on the physical sector; Nielsen flatness implies gauge–parameter insensitivity within tolerance (eq. (7.607) and Corollary 7.459). The law diagnostics (slope acceptance eq. (7.542) and proximity to unity) are computed using the same on–shell, band–projected data as in Step 7 and are instrument-robust by construction of the mask and its guards. Therefore, each threshold comparison that enters items (1)–(4) is stable under the stated variations, and the ordered decision rule returns the same quadrant (up to the quoted tolerances). □

Outputs (audit kit). We provide four audit artefacts, all restricted to the certified mask $\mathfrak{M}_{\text{mask}}$ (cf. eqs. (7.626) and (7.629)) and computed on–shell in the Nielsen–projected sector (gauge–flat within tolerance; eq. (7.607)):

- **Law heatmaps.** $(u, v) \mapsto 1 - \Lambda_{\text{law}}(u, v)$ with $\mathfrak{M}_{\text{mask}}$ overlaid and the line $v = \ln \ell_R$ marked.

- **Ward residuals.** Panels for $\mathcal{W}(u,v)$ and decay plots $v \mapsto \mathcal{W}(e^v)$ (cf. eqs. (7.591) and (7.602)).
- **Spectral reconvergence.** Line-cuts $v \mapsto \delta m_\rho^2/\overline{m_\rho^2}$ and $v \mapsto \delta Z_\rho/\overline{Z_\rho}$ with acceptance bands (cf. eqs. (7.618) to (7.620)).
- **Certified summary.** A table of key certified quantities $\{\ \ell_R,\ \mu_R,\ \varsigma_{\min},\ y_{\tilde{B}},\ \varepsilon_{\text{spec}}\ \}$ with spreads over admissible (P_x, K_{IR}) and gauges; see eqs. (7.554) and (7.588) for definitions/tolerances.

In short: keep the state coherent, let the law *heal* under coarse–graining, prove it via RG irrelevance (Section 7.8.4) + identity closure (Section 7.8.5) + spectral reconvergence (Section 7.8.6), and *certify* it on an IR plateau via the Q III mask (Section 7.8.7).

7.8.9 QIII Lessons & IR "Particle-like" Outlook

What we learned (QIII).

1. **Law heals under coarse–graining.** The law–visibility increases monotonically on the IR window with a *positive robust slope* $\varsigma_{\text{law}}^{(\tau)}$ (eqs. (7.539) and (7.542)), and the *restoration length* ℓ_R is finite (Definition 7.428).
2. **Breakings are RG–irrelevant.** After Nielsen projection, all excited breaking directions have $y_{\tilde{B}} < 0$ and $\widetilde{\varepsilon}_{\tilde{B}}(\mu) \to 0$ as $\mu \downarrow$ (Wilsonian ledger; Section 7.8.4, [389, 91, 82]).
3. **Identity closure in the IR.** Band–projected Ward residuals $\mathcal{W}(\ell)$ decay and sit within tolerance at ℓ_R; the ST identity is exact on shell; quoted IR poles/residues are *gauge–parameter flat* (Nielsen) on the mask (Section 7.8.5, [117, 243, 442]).
4. **Spectrum reconverges into multiplets.** Mass and residue splittings close on the plateau (eq. (7.620)); optional pseudo–NG modes follow the RG–dictated small–gap scaling (Proposition 7.509), while the state remains coherent (no volume decay; eq. (7.566)). Kato continuity ensures smooth evolution with ℓ [115].
5. **Physicality strengthens.** Nonnegative mass operators, strong hyperbolicity, and spectral positivity hold and *improve* across the certified mask $\mathfrak{M}_{\text{mask}}$ (Section 7.8.8); together with the slope and RG ledger, this completes the QIII certification.

IR "particle-like" outlook if emergence completes. On (and beyond) the QIII plateau $\mathfrak{M}_{\text{mask}}$ (eq. (7.626)), the on–shell pole structure and residues admit the following reading (Nielsen–flat, eq. (7.607); poles/residues read from the physical-sector quadratic form eqs. (6.332) and (7.10)):

- **Approaches the QI census.** *(a) Massless spin–2:* A pole at $p^2 = 0$ with transverse–traceless residue and positive norm signals a helicity–± 2 graviton–like excitation. Gauge–parameter stability follows from the identity ledger; causal propagation is supported by the hyperbolicity margin (Definition 7.535) and positivity by the spectral

floor (Definition 7.536) [14, 91, 255]. *(b) Unbroken massless vectors:* Poles at $p^2 = 0$ in conserved spin–1 channels with positive residues indicate unbroken gauge sectors; ST exactness on shell ensures the transverse/decoupling structure (Section 7.8.5, [117, 243]). *(c) Gapped scalars:* Scalar channels show $m_i^2(\ell_R) > 0$ with multiplet–consistent degeneracies and positive residues (eqs. (7.618) and (7.619); [82, 14]). As $\ell \uparrow$, the QIII outputs become indistinguishable (within tolerances) from the QI pattern.

- **Optional pseudo–NG modes.** If an approximate global symmetry was softly broken in Step 1, QIII allows small–gap scalar modes with $m_{\mathrm{pNG}}^2(\ell) \propto \ell^{-\alpha|y_{\bar{B}}|}$ (Proposition 7.509). Their presence is *consistent* with restoration provided multiplet splittings close (eq. (7.620)) and state visibility remains volume–uniform (eq. (7.566)). In the strict IR ($\ell \to \infty$), these gaps vanish if the associated breaking is truly irrelevant [91, 82].

- **No BEH reorganisation (unless emergent gauge becomes exact).** In QIII, global Ward structures are restored while the state stays coherent, so no Brout–Englert–Higgs (BEH) rearrangement is expected in the IR spectrum. If, however, the emergent symmetry is *gauged and exact* on the plateau (identity ledger shows full gauge structure and ST exactness with vanishing global residues), then the long–ℓ limit should be treated as **QI–like**: vector bosons are either exactly massless (unbroken phase) or acquire BEH masses with the usual Goldstone–eating pattern; diagnostics rely on the pole/residue reorganisation read from eqs. (6.332) and (7.10) and on the positivity/hyperbolicity guards ([14, 91]).

Bottom line. QIII shows that *keeping the state coherent* while letting the *law* evolve under coarse–graining produces an IR where symmetry is effectively restored: the law–visibility Λ_{law} approaches unity with a positive robust slope (Sections 7.8.2 and 7.8.3); explicit breakings die out under RG (Nielsen–projected exponents, Section 7.8.4); Ward, Slavnov–Taylor, and Nielsen identities *close* on shell (Section 7.8.5); spectra *reconverge into multiplets* (Section 7.8.6); and the physicality margins (mass, hyperbolicity, spectral positivity) *improve* across the certified mask (Section 7.8.8). When these conditions hold, the IR particle content *approaches the QI census*—with optional pseudo–NG modes governed by the irrelevant exponents—yielding a coherent, audit–ready account of emergent symmetry in the IR (Sections 7.8.2 to 7.8.8).

7.8.10 Concluding Census of Particle-like Modes on the QIII Plateau

Narrative summary. Having completed the QIII programme—positive robust law–slope and finite restoration length (eqs. (7.539) and (7.542) and Definition 7.428), RG–irrelevance of explicit breakings (Section 7.8.4), Ward/ST/Nielsen closure (Section 7.8.5), spectral reconvergence (Section 7.8.6), and mask certification (eq. (7.626))—the IR admits a clean, *on–shell* particle–like reading (Nielsen–flat within tolerance, eq. (7.607)). The actual census depends on which symmetry remains unbroken in the IR, but the counting itself is simple and audit–ready.

Census rule on the QIII mask. On the certified plateau $\mathfrak{M}_{\mathrm{mask}}$ (eq. (7.626)), poles and residues are extracted from the band–projected quadratic form on the physical sector (first–order reduction around eqs. (6.332) and (7.10)) and grouped into G_{IR}–multiplets (eq. (7.620)). The species count is:

- **Graviton–like:** *one* massless spin–2 species (helicities ± 2 in 3+1D; 2 physical dof) signalled by a transverse–traceless residue at $p^2 = 0$. Causal propagation is supported by the hyperbolicity margin (Definition 7.535); spectral positivity is witnessed by the spectral floor (Definition 7.536) [14, 91, 255].
- **Massless vectors:** $\dim G_{\mathrm{IR}}$ species, one per generator of the unbroken gauge group (each with 2 physical dof in 3+1D). ST exactness on shell ensures the transverse/decoupling structure (Section 7.8.5, [117, 243, 82, 14]).
- **Gapped scalars:** one scalar species per multiplet component that lies below the first continuum threshold. If the surviving multiplets are ρ with dimensions d_ρ, then $N_{\mathrm{scalar}} = \sum_\rho d_\rho$ (each with 1 dof); degeneracies close on the plateau (eq. (7.620)) [14, 82].
- **Pseudo–NG (optional/transient):** at finite ℓ there may be up to as many light scalar modes as approximate broken generators; their gaps scale as $m^2_{\mathrm{pNG}}(\ell) \propto \ell^{-\alpha|y_{\tilde{B}}|}$ with $y_{\tilde{B}} < 0$ from the RG ledger, so they fade as $\ell \uparrow$ if the associated breakings are truly irrelevant (Proposition 7.509 and Section 7.8.4).

Operational counting (audit path). Read the census directly from pole data on $\mathfrak{M}_{\mathrm{mask}}$:

1. a single $p^2 = 0$ pole with TT residue $\Rightarrow$ spin–2 (graviton–like);
2. one massless spin–1 pole per unbroken generator $\Rightarrow \dim G_{\mathrm{IR}}$ vectors;
3. the number of *isolated* scalar poles below the first threshold, grouped into G_{IR}–multiplets $\Rightarrow N_{\mathrm{scalar}} = \sum_\rho d_\rho$;
4. any additional light scalar poles whose m^2 decreases with ℓ as in Proposition 7.509 $\Rightarrow$ transient pseudo–NGs (allowed at finite ℓ, vanish asymptotically).

All pole/residue statements are on–shell and gauge–parameter flat (Nielsen), eq. (7.607).

Ballpark cases.

- *No unbroken IR gauge group* (pure global restoration): 1 massless spin–2 + N_{scalar} gapped scalars; pseudo–NGs typically disappear as $\ell \to \infty$ if irrelevance wins.
- $G_{\mathrm{IR}} = \mathrm{SU}(2) \times \mathrm{U}(1)$ *unbroken:* 1 spin–2 + $\big(3{+}1\big) = 4$ massless vectors + N_{scalar} gapped scalars; pseudo–NGs may be present at finite ℓ but shrink with the irrelevant exponents.

Headline formula. Summarising the species (not dof) count on the QIII plateau,

$$\boxed{N_{\text{modes}} \;\approx\; 1\ (\text{spin–2}) \;+\; \dim G_{\text{IR}}\ (\text{vectors}) \;+\; \sum_{\rho} d_\rho\ (\text{scalars}) \;+\; N_{\text{pNG}}(\text{finite } \ell),} \tag{7.643}$$

with $N_{\text{pNG}} \to 0$ as $\ell \to \infty$ if the QIII irrelevance conditions hold (Section 7.8.4 and Proposition 7.509). This census aligns with the QI–like outlook when emergence completes (Section 7.8.9) and rests on standard degrees–of–freedom counting and spectral positivity [14, 91, 82, 255].

Field provenance (gaiamiti vs. tantrakyma). As in the QI/QII analyses, we tag each particle–like mode on the QIII plateau by its *field provenance*. Using the same band instrumentation $P_x(\lambda_{\text{UV}})$ and $K_{\text{IR}}^{(\ell)}$ as before, and the same two–point extraction on the BRST physical sector (first–order reduction around eqs. (6.332) and (7.10)), we resolve the one–particle subspace into orthogonal, band–consistent components

$$\mathcal{H}_{1\text{p}} = \mathcal{H}_{\text{gai}} \;\oplus\; \mathcal{H}_{\text{tan}},$$

where $\mathcal{H}_{\text{gai}}$ ("gaiamiti") collects metric/geometry–type components (spin–2 block, geometric moduli) and $\mathcal{H}_{\text{tan}}$ ("tantrakyma") collects internal gauge and matter components (Yang–Mills and scalar matter). Let Π_{gai} and Π_{tan} be the corresponding projectors on the *Nielsen–projected* physical subspace, constructed in the same basis used to form the band–projected Hessian in eq. (7.10) and commuting with the band projection.

For each on–shell pole i with unit–norm eigenvector $u_i(u,v)$ of the physical two–point operator, define the sector *origin scores*

$$\boxed{\pi_{\text{gai}}(i;u,v) := \langle u_i, \Pi_{\text{gai}}\, u_i\rangle, \qquad \pi_{\text{tan}}(i;u,v) := \langle u_i, \Pi_{\text{tan}}\, u_i\rangle, \qquad \pi_{\text{gai}} + \pi_{\text{tan}} = 1,} \tag{7.644}$$

the last identity following from $\Pi_{\text{gai}} + \Pi_{\text{tan}} = \mathbf{1}_{\mathcal{H}_{1\text{p}}}$ and $\Pi_{\text{gai}}\Pi_{\text{tan}} = 0$ (up to numerical tolerances). With a small audit tolerance $\delta_{\text{orig}} \geq 0$ and for $(u,v) \in \mathfrak{M}_{\text{mask}}$ (eq. (7.626)), we partition each multiplet $\mathfrak{M}_\rho(\ell)$ (cf. eqs. (7.618) and (7.619)) by

$$\boxed{\begin{aligned}
\mathfrak{M}_\rho^{\text{gai}}(\ell;u,v) &:= \Big\{ i \in \mathfrak{M}_\rho(\ell) \;\Big|\; \pi_{\text{gai}}(i;u,v) \;\geq\; \pi_{\text{tan}}(i;u,v) + \delta_{\text{orig}} \Big\},\\
\mathfrak{M}_\rho^{\text{tan}}(\ell;u,v) &:= \Big\{ i \in \mathfrak{M}_\rho(\ell) \;\Big|\; \pi_{\text{tan}}(i;u,v) \;\geq\; \pi_{\text{gai}}(i;u,v) + \delta_{\text{orig}} \Big\},\\
\mathfrak{M}_\rho^{\text{hyb}}(\ell;u,v) &:= \Big\{ i \in \mathfrak{M}_\rho(\ell) \;\Big|\; \big|\pi_{\text{gai}}(i;u,v) - \pi_{\text{tan}}(i;u,v)\big| < \delta_{\text{orig}} \Big\}.
\end{aligned}} \tag{7.645}$$

By construction, $\mathfrak{M}_\rho^{\text{gai}}$ and $\mathfrak{M}_\rho^{\text{tan}}$ are disjoint and $\mathfrak{M}_\rho = \mathfrak{M}_\rho^{\text{gai}} \cup \mathfrak{M}_\rho^{\text{tan}} \cup \mathfrak{M}_\rho^{\text{hyb}}$. Modes with $\big|\pi_{\text{gai}} - \pi_{\text{tan}}\big| < \delta_{\text{orig}}$ are flagged as *hybrid*; they are rare on the certified mask thanks to spectral reconvergence (Section 7.8.6) and Kato continuity [115].

Canonical tags.

- **Graviton–like (spin–2):** always *gaiamiti*. The transverse–traceless $p^2{=}0$ pole lies in $\mathcal{H}_{\rm gai}$ by construction; causal propagation and positivity are supported by the Step 8 margins (eqs. (7.634) and (7.636)) [14, 91, 255].
- **Massless vectors:** *tantrakyma*. Each unbroken gauge generator gives a spin–1 $p^2{=}0$ pole with transversality enforced by the ST/Nielsen ledger (Section 7.8.5) [117, 243, 82].
- **Gapped scalars:** split by eq. (7.645). Geometric moduli (dilaton/radion–like) tag as *gaiamiti*; matter multiplets tag as *tantrakyma*. Degeneracies and residues close on $\mathfrak{M}_{\rm mask}$ (eq. (7.620)).
- **Pseudo–NG (optional):** provenance follows the *spurion ledger* from Step 1; a pseudo–NG tied to a breaking spurion S_B inherits the sector where S_B carries its quantum numbers [440]. Its gap scales with the irrelevant exponent (Section 7.8.4 and Proposition 7.509).

Tagged census (species, not dof). Decompose the headline count eq. (7.643) into sectoral contributions (on the QIII mask):

$$\boxed{\begin{aligned} N^{\rm gai}_{\rm modes} &= 1\ (\text{spin–2}) \ +\ \sum_{\rho} |\mathfrak{M}^{\rm gai}_{\rho}(\ell_R)|, \\ N^{\rm tan}_{\rm modes} &= \dim G_{\rm IR}\ (\text{vectors}) \ +\ \sum_{\rho} |\mathfrak{M}^{\rm tan}_{\rho}(\ell_R)| \ +\ N^{\rm tan}_{\rm pNG}(\text{finite } \ell), \end{aligned}} \tag{7.646}$$

where $N^{\rm tan}_{\rm pNG}$ counts any finite–ℓ pseudo–NGs sourced by tantrakyma spurions. Hybrids (if any) may be reported separately or assigned by the larger score in eq. (7.644) with an explicit note of $\delta_{\rm orig}$.

Audit note. The origin scores eq. (7.644) and partitions eq. (7.645) are evaluated only on $(u, v) \in \mathfrak{M}_{\rm mask}$ (eq. (7.626)), with all on–shell data Nielsen–projected and gauge–flat within tolerance (eq. (7.607)). Inner products and projectors act on the Nielsen–projected physical subspace. This preserves gauge/basis/scheme safety (cf. Lemmas 7.451 and 7.527) while keeping the tagging aligned with spectral reconvergence and the RG ledger [117, 243, 91, 82, 115].

7.9 Quadrant IV: Explicit Law Breaking with an Asymmetric State

Preconditions (Stage-I gate). Work anomaly-free so that the BRST differential $\zeta(\cdot) = \{\hat{Q}_{\rm BRST}, \cdot\}$ is nilpotent and the physical sector is $\mathcal{H}_{\rm phys} = H^0(\hat{Q}_{\rm BRST})$; all readouts (poles, residues, visibilities) are then gauge-parameter independent by Nielsen/Slavnov–Taylor identities on $\mathcal{H}_{\rm phys}$ [445, 106, 117].

Definition 7.550 (Quadrant IV overlay and diagnostics via law/state channels)**.** Let the UV/IR instrumentation be the scale-tags of Definitions 6.289 and 7.26 and the holosymmetric visibilities those of Definition 6.275. Quadrant IV is the regime in which (i) the *law* contains explicit symmetry-breaking sources and (ii) the *state* is asymmetric with a *unique* vacuum (no vacuum manifold). Concretely, on the band

$$\mathcal{L} \;=\; \mathcal{L}_0 \;+\; \sum_k \epsilon_k\, \mathcal{O}_k \;+\; \zeta\Psi, \qquad \epsilon_k \neq 0,$$

and the chosen background $(\bar{\Phi}, \bar{\Theta})$ is not G-invariant. On the physical sector $\mathcal{H}_{\text{phys}}$, the BRST-improved, adjoint-covariant Ward identities (eqs. (6.292) and (6.293)) isolate the explicit sources as the *only* origin of non-conservation,

$$\nabla^{(\text{ad})}_{\mu}\, \bar{J}^{\mu}_{a,\text{imp}} \;=\; \sum_k \epsilon_k \left\langle \mathcal{S}_{k,a} \right\rangle_{\boldsymbol{\Psi}},$$

so that a *persistent* deviation of the law-visibility from unity and the *absence of exact NG modes* (only pseudo-NG) are the expected QIV signatures on certified plateaus (Definition 6.300). All diagnostics are evaluated cohomologically on $\mathcal{H}_{\text{phys}}$ and must satisfy Nielsen/Slavnov–Taylor gauge-parameter independence. Background references for BRST/ST are [445, 106, 117]; for SSB, pseudo-Goldstones, and GMOR-type scaling see [93, 94, 261, 91, 112].

Intuition. In QIV the **law is explicitly asymmetric** and the **state is uniquely selected** by those explicit terms. We should observe (i) a **law-visibility deficit** that does *not* heal under IR coarse-graining on certified bands, and (ii) **no exact NG** (only pseudo-NG) with masses controlled by the explicit couplings via GMOR-type relations. These expectations must be verified on $\mathcal{H}_{\text{phys}}$ under the UV/IR protocol of Section 6.12.

Recall (improved current). As defined in eq. (6.292), the (BRST-)improved Noether current is

$$\boxed{J^{\mu}_{a,\text{imp}} \;:=\; J^{\mu}_{a} \;-\; \zeta\, \Lambda^{\mu}_{a}}$$

where $\zeta(\cdot) = \{\hat{Q}_{\text{BRST}}, \cdot\}$ and Λ^{μ}_{a} is determined by the gauge-fermion variation $\delta_a\Psi = \alpha_a\, \chi_a + (\nabla_\mu \alpha_a)\Lambda^{\mu}_{a}$. (Operator form used in Ward identities: $\hat{J}^{\mu}_{a,\text{imp}} = \hat{J}^{\mu}_{a} - \zeta\, \Lambda^{\mu}_{a}$.)

Protocol (Goal → Gates → Setup → Steps → Exit)

Goal. Quantify explicit misalignment (law asymmetry with explicitly selected state), extract pseudo-NG spectra/mixings, and certify persistent non-conservation together with a bounded-away-from-unity law visibility on a joint plateau.

Gates (must pass before analysis counts).

1. **Stage-I/BRST viability:** anomaly-free BRST scaffold (Section 6.12); otherwise the candidate law fails before quadrant assignment.

2. **Plateau certification:** enforce slope/curvature/oscillation bounds at bandpoints and only report within Definition 6.300.

3. **Overlay neutrality:** holosymmetric overlays (Definition 6.275) are readouts and cannot alter quadrant logic.

Setup.

- Work **on the band** with $P_x(\lambda_{\mathrm{UV}})$ and within the IR window $K_{\mathrm{IR}}^{(\ell)}$ (Definitions 6.289 and 7.26); use the band/IR-tagged visibilities Λ_{law} and Λ_{state} (Definition 6.275).
- Law: $\mathcal{L} = \mathcal{L}_0 + \sum_k \epsilon_k \mathcal{O}_k + \zeta\Psi$ with **non-vanishing** explicit breakers ϵ_k. State: choose an **asymmetric** background; once $\epsilon_k \neq 0$ there is *no* vacuum manifold.

Steps. Detailed in the remainder: construct the band-limited mass/mixing ledger on $\mathcal{H}_{\mathrm{phys}}$, perform Ward $\Rightarrow$ pseudo-NG (GMOR-type) fits, and certify the joint plateau showing a persistent law-deficit.

Exit. Declare QIV when the certified joint plateau exhibits (i) non-healing law-visibility deficit, (ii) unique-vacuum state visibility (no thermodynamic orthogonality), and (iii) pseudo-NG masses consistent with the explicit sources—each BRST/Nielsen safe (Definition 6.300 and eqs. (6.292) and (6.293)).

7.9.1 UV/IR Instrumentation and Visibility Order Parameters

Definition 7.551 (Band-tagged operators, states, and branch unitaries in QIV)**.** Let $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ be the UV spectral projector and IR coarse-graining window of Definitions 6.289 and 7.26. For any operator $\hat{\mathcal{O}}$ and state ρ_0, define the band/IR-tagged objects

$$\hat{\mathcal{O}}^{(\lambda_{\mathrm{UV}},\ell)} := K_{\mathrm{IR}}^{(\ell)}\, P_x(\lambda_{\mathrm{UV}})\, \hat{\mathcal{O}}\, P_x(\lambda_{\mathrm{UV}})\, K_{\mathrm{IR}}^{(\ell)}, \qquad \rho_0^{(\lambda_{\mathrm{UV}},\ell)} := P_x(\lambda_{\mathrm{UV}})\, \rho_0\, P_x(\lambda_{\mathrm{UV}}).$$

For the holosymmetric interferometer Definition 6.275, define the band-tagged branches

$$U_{A/B}^{(\lambda_{\mathrm{UV}},\ell)} := K_{\mathrm{IR}}^{(\ell)}\, P_x(\lambda_{\mathrm{UV}})\, U_{A/B}\, P_x(\lambda_{\mathrm{UV}})\, K_{\mathrm{IR}}^{(\ell)}, \quad |\Omega_{1/2}^{(\lambda_{\mathrm{UV}},\ell)}\rangle := K_{\mathrm{IR}}^{(\ell)}\, P_x(\lambda_{\mathrm{UV}})\, |\Omega_{1/2}\rangle\,.$$

Assume $[\zeta, P_x(\lambda_{\mathrm{UV}})] = [\zeta, K_{\mathrm{IR}}^{(\ell)}] = 0$ as in Section 6.12, so all tagged objects descend to the physical sector $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$.

Intuition. We restrict both preparation and readout to a controlled UV/IR band so that all observables are regulator-stable. BRST-closed tags ensure everything is evaluable cohomologically on $\mathcal{H}_{\mathrm{phys}}$ (Nielsen/Slavnov–Taylor gauge-independence) [445, 106, 117].

Lemma 7.552 (Descent to the physical sector)**.** *Under the assumptions of Definition 7.551, for any operator $\hat{\mathcal{O}}$ with ghost number 0 and any ζ-closed state on the band, the tagged object $\hat{\mathcal{O}}^{(\lambda_{\mathrm{UV}},\ell)}$ is ζ-closed and defines a unique class in $H^0(\hat{Q}_{\mathrm{BRST}})$; expectation values computed with $\rho_0^{(\lambda_{\mathrm{UV}},\ell)}$ are independent of gauge parameters (Nielsen identity).*

Proof of Lemma 7.552. We split the argument into three parts.

(a) BRST-closure of the tagged operator. By Definition 7.551 the UV/IR taggers are BRST-closed and commute with the BRST differential: $[\zeta, P_x(\lambda_{\mathrm{UV}})] = [\zeta, K_{\mathrm{IR}}^{(\ell)}] = 0$. Hence, for any operator $\hat{\mathcal{O}}$ (ghost number 0),

$$\zeta\Big(\hat{\mathcal{O}}^{(\lambda_{\mathrm{UV}},\ell)}\Big) = \zeta\Big(K_{\mathrm{IR}}^{(\ell)} P_x\, \hat{\mathcal{O}}\, P_x K_{\mathrm{IR}}^{(\ell)}\Big) = K_{\mathrm{IR}}^{(\ell)} P_x\, \zeta(\hat{\mathcal{O}})\, P_x K_{\mathrm{IR}}^{(\ell)}.$$

If $\hat{\mathcal{O}}$ is chosen as a representative of its BRST cohomology class with $\zeta(\hat{\mathcal{O}}) = 0$ (which we can always do by replacing $\hat{\mathcal{O}} \mapsto \hat{\mathcal{O}} - \zeta(\hat{X})$ for some $\hat{X}$), then $\zeta\Big(\hat{\mathcal{O}}^{(\lambda_{\mathrm{UV}},\ell)}\Big) = 0$ as well. Thus the tagged operator is BRST-closed.

(b) Well-defined class in $H^0(\hat{Q}_{\mathrm{BRST}})$. Suppose $\hat{\mathcal{O}}$ and $\hat{\mathcal{O}}'$ are two ghost-number-zero representatives of the same cohomology class, $\hat{\mathcal{O}}' = \hat{\mathcal{O}} + \zeta(\hat{X})$. Tagging preserves BRST-exactness because the taggers commute with ζ:

$$\hat{\mathcal{O}}'^{(\lambda_{\mathrm{UV}},\ell)} - \hat{\mathcal{O}}^{(\lambda_{\mathrm{UV}},\ell)} = \Big(\zeta(\hat{X})\Big)^{(\lambda_{\mathrm{UV}},\ell)} = \zeta\Big(\hat{X}^{(\lambda_{\mathrm{UV}},\ell)}\Big).$$

Hence $\hat{\mathcal{O}}'^{(\lambda_{\mathrm{UV}},\ell)}$ and $\hat{\mathcal{O}}^{(\lambda_{\mathrm{UV}},\ell)}$ define the *same* class in $H^0(\hat{Q}_{\mathrm{BRST}})$. This shows that the class $[\hat{\mathcal{O}}^{(\lambda_{\mathrm{UV}},\ell)}] \in H^0(\hat{Q}_{\mathrm{BRST}})$ is uniquely defined by the class $[\hat{\mathcal{O}}] \in H^0(\hat{Q}_{\mathrm{BRST}})$.

(c) Gauge-parameter independence (Nielsen identity) of expectations. Let $\rho_0^{(\lambda_{\mathrm{UV}},\ell)} = P_x\, \rho_0\, P_x$ be a ζ-closed (band-tagged) state, i.e. $\zeta\Big(\rho_0^{(\lambda_{\mathrm{UV}},\ell)}\Big) = 0$, as in Definition 7.551. For any operator $\hat{Y}$ we use that $\zeta(\cdot) = \{\hat{Q}_{\mathrm{BRST}}, \cdot\}$ is a graded derivation and that the (super)trace of a (super)commutator vanishes:

$$\mathrm{Tr}\Big(\rho_0^{(\lambda_{\mathrm{UV}},\ell)}\, \zeta(\hat{Y})\Big) = \mathrm{Tr}\Big(\zeta\Big(\rho_0^{(\lambda_{\mathrm{UV}},\ell)}\hat{Y}\Big)\Big) - \mathrm{Tr}\Big(\zeta\Big(\rho_0^{(\lambda_{\mathrm{UV}},\ell)}\Big)\, \hat{Y}\Big) = 0 - 0 = 0.$$

Therefore expectation values are insensitive to adding ζ-exact pieces to either the operator or the state.

Now let α_{gf} be a gauge-fixing parameter. Algebraic renormalization (Slavnov–Taylor with α_{gf} promoted to a BRST doublet) gives the *Nielsen identity* [117]:

$$\partial_{\alpha_{\mathrm{gf}}}\, \hat{\mathcal{O}} = \zeta\Big(\hat{\mathcal{N}}_{\alpha_{\mathrm{gf}}}[\hat{\mathcal{O}}]\Big), \qquad \partial_{\alpha_{\mathrm{gf}}}\, \rho_0 = \zeta\Big(\hat{\mathcal{N}}_{\alpha_{\mathrm{gf}}}[\rho_0]\Big),$$

for suitable local insertions $\hat{\mathcal{N}}_{\alpha_{\mathrm{gf}}}[\cdot]$. Since P_x and $K_{\mathrm{IR}}^{(\ell)}$ commute with ζ, the same relations hold after tagging. Hence

$$\begin{aligned}
\partial_{\alpha_{\mathrm{gf}}}\, \mathrm{Tr}\Big(\rho_0^{(\lambda_{\mathrm{UV}},\ell)}\, \hat{\mathcal{O}}^{(\lambda_{\mathrm{UV}},\ell)}\Big) &= \mathrm{Tr}\Big((\partial_{\alpha_{\mathrm{gf}}}\rho_0^{(\lambda_{\mathrm{UV}},\ell)})\, \hat{\mathcal{O}}^{(\lambda_{\mathrm{UV}},\ell)}\Big) \\
&\quad + \mathrm{Tr}\Big(\rho_0^{(\lambda_{\mathrm{UV}},\ell)}\, \partial_{\alpha_{\mathrm{gf}}}\hat{\mathcal{O}}^{(\lambda_{\mathrm{UV}},\ell)}\Big) \\
&= \mathrm{Tr}\Big(\zeta\Big(\hat{\mathcal{N}}_{\alpha_{\mathrm{gf}}}[\rho_0]^{(\lambda_{\mathrm{UV}},\ell)}\Big)\, \hat{\mathcal{O}}^{(\lambda_{\mathrm{UV}},\ell)}\Big) \\
&\quad + \mathrm{Tr}\Big(\rho_0^{(\lambda_{\mathrm{UV}},\ell)}\, \zeta\Big(\hat{\mathcal{N}}_{\alpha_{\mathrm{gf}}}[\hat{\mathcal{O}}]^{(\lambda_{\mathrm{UV}},\ell)}\Big)\Big) = 0,
\end{aligned}$$

by the previous vanishing of expectations of ζ-exact insertions. Thus, expectation values computed with $\rho_0^{(\lambda_{\mathrm{UV}},\ell)}$ are gauge-parameter independent.

Conclusion. Parts (a)–(c) prove that $\hat{\mathcal{O}}^{(\lambda_{\mathrm{UV}},\ell)}$ is BRST-closed, represents a unique element of $H^0(\hat{Q}_{\mathrm{BRST}})$, and has gauge-parameter–independent expectation values on the physical sector, as claimed. □

Definition 7.553 (QIV holosymmetric visibilities (band-tagged))**.** With $U_{A/B}^{(\lambda_{\mathrm{UV}},\ell)}$ and $\rho_0^{(\lambda_{\mathrm{UV}},\ell)}$ as in Definition 7.551, define

$$\Lambda_{\mathrm{law}}^{(\lambda_{\mathrm{UV}},\ell)} := \mathrm{Tr}\Big[\Big(U_B^{(\lambda_{\mathrm{UV}},\ell)}\Big)^\dagger U_A^{(\lambda_{\mathrm{UV}},\ell)}\,\rho_0^{(\lambda_{\mathrm{UV}},\ell)}\Big], \qquad \Lambda_{\mathrm{state}}^{(\lambda_{\mathrm{UV}},\ell)} := \langle\Omega_2^{(\lambda_{\mathrm{UV}},\ell)}|\Omega_1^{(\lambda_{\mathrm{UV}},\ell)}\rangle\,.$$

These are the operational order parameters for the law- and state-channels in QIV (cf. Definition 6.275 and Section 6.12).

Intuition. Λ_{law} compares the two interferometric branches of the *law* dynamics; Λ_{state} compares the prepared *states*. Band-tagging guarantees scale locality; BRST descent ensures physicality.

Definition 7.554 (Triadic channel projectors)**.** Let $\{\Pi_\alpha\}_{\alpha\in\{\mathrm{corr,vac,spec}\}}$ be orthogonal, BRST-closed channel projectors (correlational, vacuum, spectral) commuting with $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$, as in the triadic split of Section 6.12. Define the channel-resolved visibilities by inserting Π_α into the definitions of Definition 7.553.

Proposition 7.555 (Additive channel decomposition)**.** *Under Definition 7.554, the visibilities decompose additively,*

$$\Lambda_\bullet^{(\lambda_{\mathrm{UV}},\ell)} = \sum_{\alpha\in\{\mathrm{corr,vac,spec}\}} \Lambda_\bullet^{(\alpha;\lambda_{\mathrm{UV}},\ell)}, \qquad \bullet\in\{\mathrm{law},\mathrm{state}\}.$$

Proof Proposition 7.555. We treat the law and state visibilities in turn and make explicit where each structural property is used.

Preliminaries. By Definition 7.554, the channel projectors $\{\Pi_\alpha\}_{\alpha\in\{\mathrm{corr,vac,spec}\}}$ obey

$$\Pi_\alpha^\dagger = \Pi_\alpha, \quad \Pi_\alpha\Pi_\beta = \delta_{\alpha\beta}\Pi_\alpha, \quad \sum_\alpha \Pi_\alpha = \mathbb{1},$$

and commute with the UV/IR taggers $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$. Hence, for any band-tagged operator $X^{(\lambda_{\mathrm{UV}},\ell)}$ (e.g. $U_{A/B}^{(\lambda_{\mathrm{UV}},\ell)}$, $\rho_0^{(\lambda_{\mathrm{UV}},\ell)}$) we may freely insert $\sum_\alpha \Pi_\alpha$ and move the Π_α through the taggers.

State visibility. With the band-tagged branch states $|\Omega_{1/2}^{(\lambda_{\mathrm{UV}},\ell)}\rangle := K_{\mathrm{IR}}^{(\ell)}P_x(\lambda_{\mathrm{UV}})\,|\Omega_{1/2}\rangle$, we have

$$\Lambda_{\mathrm{state}}^{(\lambda_{\mathrm{UV}},\ell)} = \langle\Omega_2^{(\lambda_{\mathrm{UV}},\ell)}|\Omega_1^{(\lambda_{\mathrm{UV}},\ell)}\rangle = \langle\Omega_2^{(\lambda_{\mathrm{UV}},\ell)}\Big|\Big(\sum_\alpha \Pi_\alpha\Big)\Big|\Omega_1^{(\lambda_{\mathrm{UV}},\ell)}\rangle\,.$$

Because $[\Pi_\alpha, K_{\mathrm{IR}}^{(\ell)}] = [\Pi_\alpha, P_x(\lambda_{\mathrm{UV}})] = 0$, inserting the resolution of the identity between the tagged states does not disturb band-tagging. Linearity then gives

$$\Lambda_{\mathrm{state}}^{(\lambda_{\mathrm{UV}},\ell)} = \sum_\alpha \langle\Omega_2^{(\lambda_{\mathrm{UV}},\ell)}|\Pi_\alpha|\Omega_1^{(\lambda_{\mathrm{UV}},\ell)}\rangle = \sum_\alpha \Lambda_{\mathrm{state}}^{(\alpha;\lambda_{\mathrm{UV}},\ell)},$$

which is the desired additive decomposition for the state channel.

Law visibility (one-sided decomposition—always valid). We define the following quantities:

$$X^{(\lambda_{\mathrm{UV}},\ell)} := (U_B^{(\lambda_{\mathrm{UV}},\ell)})^\dagger U_A^{(\lambda_{\mathrm{UV}},\ell)}$$
$$\rho^{(\lambda_{\mathrm{UV}},\ell)} := \rho_0^{(\lambda_{\mathrm{UV}},\ell)}$$

Then

$$\Lambda_{\mathrm{law}}^{(\lambda_{\mathrm{UV}},\ell)} = \mathrm{Tr}\Big[X^{(\lambda_{\mathrm{UV}},\ell)}\,\rho^{(\lambda_{\mathrm{UV}},\ell)}\Big] = \mathrm{Tr}\Big[\Big(\sum_\alpha \Pi_\alpha\Big) X^{(\lambda_{\mathrm{UV}},\ell)}\,\rho^{(\lambda_{\mathrm{UV}},\ell)}\Big] = \sum_\alpha \mathrm{Tr}\Big[\Pi_\alpha X^{(\lambda_{\mathrm{UV}},\ell)}\,\rho^{(\lambda_{\mathrm{UV}},\ell)}\Big].$$

Define the channel-resolved law visibility by the *one-sided* insertion

$$\Lambda_{\mathrm{law}}^{(\alpha;\lambda_{\mathrm{UV}},\ell)} := \mathrm{Tr}\Big[\Pi_\alpha X^{(\lambda_{\mathrm{UV}},\ell)}\,\rho^{(\lambda_{\mathrm{UV}},\ell)}\Big].$$

This yields $\Lambda_{\mathrm{law}}^{(\lambda_{\mathrm{UV}},\ell)} = \sum_\alpha \Lambda_{\mathrm{law}}^{(\alpha;\lambda_{\mathrm{UV}},\ell)}$ without requiring any commutation between Π_α and the branch unitaries or the preparation ρ_0; only linearity and $\sum_\alpha \Pi_\alpha = \mathbb{1}$ are used, together with commutation of Π_α with the UV/IR taggers (so that the insertion is legitimate at the band level).

Law visibility (symmetric two-sided form—under a stronger, often satisfied hypothesis). If, in addition, the triadic split is *functorial* for the holosymmetric branches—namely

$$[\Pi_\alpha, U_{A/B}^{(\lambda_{\mathrm{UV}},\ell)}] = 0 \quad \text{and} \quad [\Pi_\alpha, \rho_0^{(\lambda_{\mathrm{UV}},\ell)}] = 0,$$

(which holds in the common case where the channels define a block decomposition preserved by the branch dynamics and the prepared state is channel-diagonal), then the cross-channel traces vanish:

$$\mathrm{Tr}\Big[\Pi_\alpha X^{(\lambda_{\mathrm{UV}},\ell)}\,\Pi_\beta \rho^{(\lambda_{\mathrm{UV}},\ell)}\Big] = \mathrm{Tr}\Big[X^{(\lambda_{\mathrm{UV}},\ell)}\,\Pi_\alpha\Pi_\beta \rho^{(\lambda_{\mathrm{UV}},\ell)}\Big] = \delta_{\alpha\beta}\,\mathrm{Tr}\Big[X^{(\lambda_{\mathrm{UV}},\ell)}\,\Pi_\alpha \rho^{(\lambda_{\mathrm{UV}},\ell)}\Big].$$

Consequently the one-sided definition is equivalent to the *two-sided* (block-diagonal) form

$$\Lambda_{\mathrm{law}}^{(\alpha;\lambda_{\mathrm{UV}},\ell)} = \mathrm{Tr}\Big[\Pi_\alpha (U_B^{(\lambda_{\mathrm{UV}},\ell)})^\dagger U_A^{(\lambda_{\mathrm{UV}},\ell)}\,\Pi_\alpha \rho_0^{(\lambda_{\mathrm{UV}},\ell)}\Big],$$

which can be convenient for positivity or Cauchy–Schwarz estimates within each channel. Either way, the sum over α reproduces $\Lambda_{\mathrm{law}}^{(\lambda_{\mathrm{UV}},\ell)}$.

Conclusion. Combining the state and law analyses yields, for $\bullet \in \{\mathrm{law}, \mathrm{state}\}$,

$$\Lambda_\bullet^{(\lambda_{\mathrm{UV}},\ell)} = \sum_{\alpha\in\{\mathrm{corr},\mathrm{vac},\mathrm{spec}\}} \Lambda_\bullet^{(\alpha;\lambda_{\mathrm{UV}},\ell)},$$

establishing the proposition. □

Proposition 7.556 (Gauge-parameter independence on $\mathcal{H}_{\mathrm{phys}}$)**.** *For any gauge-fixing parameter* α_{gf}, $\partial_{\alpha_{\mathrm{gf}}} \Lambda_\bullet^{(\lambda_{\mathrm{UV}},\ell)} = 0$ *on* $\mathcal{H}_{\mathrm{phys}}$, *for* $\bullet \in \{\mathrm{law}, \mathrm{state}\}$.

Proof Proposition 7.556. We prove the claim for both visibilities, making explicit the role of the Nielsen identity, BRST-closure of the UV/IR taggers, and the cohomological vanishing used in Lemma 7.552.

Preliminaries (Nielsen identity and band-tagging). Let $\alpha_{\rm gf}$ be any gauge-fixing parameter entering through the gauge-fermion Ψ. Algebraic renormalization promotes $\alpha_{\rm gf}$ to a BRST doublet and yields the *Nielsen identity* (linearized Slavnov–Taylor form) for band-tagged objects [117]:

$$\partial_{\alpha_{\rm gf}} X^{(\lambda_{\rm UV},\ell)} = \zeta\Big(\mathcal{N}_{\alpha_{\rm gf}}[X]^{(\lambda_{\rm UV},\ell)}\Big), \qquad \partial_{\alpha_{\rm gf}} \rho_0^{(\lambda_{\rm UV},\ell)} = \zeta\Big(\mathcal{N}_{\alpha_{\rm gf}}[\rho_0]^{(\lambda_{\rm UV},\ell)}\Big), \tag{7.647}$$

for suitable local insertions $\mathcal{N}_{\alpha_{\rm gf}}[\cdot]$. Here the UV/IR taggers $P_x(\lambda_{\rm UV})$ and $K_{\rm IR}^{(\ell)}$ commute with the BRST differential and are chosen $\alpha_{\rm gf}$-independent, as stipulated in Definition 7.551 and Section 6.12. In particular, $[\zeta, P_x] = [\zeta, K_{\rm IR}^{(\ell)}] = 0$ and $\partial_{\alpha_{\rm gf}} P_x = \partial_{\alpha_{\rm gf}} K_{\rm IR}^{(\ell)} = 0$.

We also use the cohomological trace facts (ghost number in parentheses): for any ghost-number-zero band state $\rho^{(\lambda_{\rm UV},\ell)}$ with $\zeta\Big(\rho^{(\lambda_{\rm UV},\ell)}\Big) = 0$ and any operator Y,

$$\mathrm{Tr}\Big(\rho^{(\lambda_{\rm UV},\ell)}\,\zeta(Y)\Big) = 0, \qquad \text{equivalently} \quad \mathrm{Tr}\Big(\zeta(Z)\Big) = 0 \text{ for any } Z \text{ of ghost number } -1, \tag{7.648}$$

which is precisely the content of Lemma 7.552 (on $\mathcal{H}_{\rm phys}$).

(A) State visibility. Recall $\Lambda_{\rm state}^{(\lambda_{\rm UV},\ell)} = \langle \Omega_2^{(\lambda_{\rm UV},\ell)} | \Omega_1^{(\lambda_{\rm UV},\ell)} \rangle$ with band-tagged branch states $|\Omega_{1/2}^{(\lambda_{\rm UV},\ell)}\rangle = K_{\rm IR}^{(\ell)} P_x\,|\Omega_{1/2}\rangle$ (Definition 7.551). Introduce the rank-one band operator $\rho_{21}^{(\lambda_{\rm UV},\ell)} := |\Omega_1^{(\lambda_{\rm UV},\ell)}\rangle\langle\Omega_2^{(\lambda_{\rm UV},\ell)}|$ so that $\Lambda_{\rm state}^{(\lambda_{\rm UV},\ell)} = \mathrm{Tr}\,\big(\rho_{21}^{(\lambda_{\rm UV},\ell)}\big)$. By (7.647), $\partial_{\alpha_{\rm gf}} \rho_{21}^{(\lambda_{\rm UV},\ell)} = \zeta\Big(\mathcal{N}_{\alpha_{\rm gf}}[\rho_{21}]^{(\lambda_{\rm UV},\ell)}\Big)$ (ghost number $0 \mapsto +1$), hence

$$\partial_{\alpha_{\rm gf}} \Lambda_{\rm state}^{(\lambda_{\rm UV},\ell)} = \mathrm{Tr}\Big(\partial_{\alpha_{\rm gf}} \rho_{21}^{(\lambda_{\rm UV},\ell)}\Big) = \mathrm{Tr}\Big(\zeta\Big(\mathcal{N}_{\alpha_{\rm gf}}[\rho_{21}]^{(\lambda_{\rm UV},\ell)}\Big)\Big) = 0,$$

by (7.648). This shows $\partial_{\alpha_{\rm gf}} \Lambda_{\rm state}^{(\lambda_{\rm UV},\ell)} = 0$ on $\mathcal{H}_{\rm phys}$.

(B) Law visibility. Write

$$\Lambda_{\rm law}^{(\lambda_{\rm UV},\ell)} = \mathrm{Tr}\Big[X^{(\lambda_{\rm UV},\ell)}\,\rho_0^{(\lambda_{\rm UV},\ell)}\Big], \qquad X^{(\lambda_{\rm UV},\ell)} := \Big(U_B^{(\lambda_{\rm UV},\ell)}\Big)^{\dagger} U_A^{(\lambda_{\rm UV},\ell)},$$

with band-tagged branches from Definition 7.551. Differentiating and using (7.647) twice (for U_A, U_B, and for ρ_0) we obtain

$$\begin{aligned}\partial_{\alpha_{\rm gf}} \Lambda_{\rm law}^{(\lambda_{\rm UV},\ell)} &= \mathrm{Tr}\Big[\Big(\partial_{\alpha_{\rm gf}} X^{(\lambda_{\rm UV},\ell)}\Big)\rho_0^{(\lambda_{\rm UV},\ell)}\Big] + \mathrm{Tr}\Big[X^{(\lambda_{\rm UV},\ell)}\,\partial_{\alpha_{\rm gf}} \rho_0^{(\lambda_{\rm UV},\ell)}\Big] \\ &= \mathrm{Tr}\Big[\zeta\Big(\mathcal{N}_{\alpha_{\rm gf}}[X]^{(\lambda_{\rm UV},\ell)}\Big)\,\rho_0^{(\lambda_{\rm UV},\ell)}\Big] + \mathrm{Tr}\Big[X^{(\lambda_{\rm UV},\ell)}\,\zeta\Big(\mathcal{N}_{\alpha_{\rm gf}}[\rho_0]^{(\lambda_{\rm UV},\ell)}\Big)\Big].\end{aligned}$$

Now apply the graded Leibniz rule for ζ and the cyclicity of the (super)trace to rewrite each term as a total ζ on the trace:

$$\mathrm{Tr}\Big[\zeta(A)\,B\Big] = -\,\mathrm{Tr}\Big[A\,\zeta(B)\Big] = \mathrm{Tr}\Big[\zeta(AB)\Big]$$

for ghost-number assignments consistent with $\mathrm{Tr}(\zeta(\cdot)) = 0$.

With $A = \mathcal{N}_{\alpha_{\rm gf}}[X]^{(\lambda_{\rm UV},\ell)}$ and $B = \rho_0^{(\lambda_{\rm UV},\ell)}$ in the first term, and $A = \mathcal{N}_{\alpha_{\rm gf}}[\rho_0]^{(\lambda_{\rm UV},\ell)}$, $B = X^{(\lambda_{\rm UV},\ell)}$ in the second, we get

$$\partial_{\alpha_{\rm gf}} \Lambda_{\rm law}^{(\lambda_{\rm UV},\ell)} = \mathrm{Tr}\Big[\zeta\Big(\mathcal{N}_{\alpha_{\rm gf}}[X]^{(\lambda_{\rm UV},\ell)}\, \rho_0^{(\lambda_{\rm UV},\ell)}\Big)\Big] + \mathrm{Tr}\Big[\zeta\Big(X^{(\lambda_{\rm UV},\ell)}\, \mathcal{N}_{\alpha_{\rm gf}}[\rho_0]^{(\lambda_{\rm UV},\ell)}\Big)\Big] = 0,$$

by (7.648). No additional hypothesis on the BRST-closure of the branches is required beyond (7.647) and the band/BRST commutations from Definition 7.551.

Conclusion. Both visibilities satisfy $\partial_{\alpha_{\rm gf}} \Lambda_\bullet^{(\lambda_{\rm UV},\ell)} = 0$ on $\mathcal{H}_{\rm phys}$ for $\bullet \in \{\mathrm{law}, \mathrm{state}\}$, as claimed. □

Theorem 7.557 (Quadrant-IV visibility signature on certified plateaus)**.** *Assume the explicit breakings in the law are nonzero and relevant/marginal on the band, $\mathcal{L} = \mathcal{L}_0 + \sum_k \epsilon_k \mathcal{O}_k + \zeta\Psi$ with $\epsilon_k \neq 0$, and the BRST-improved Ward identities hold on $\mathcal{H}_{\rm phys}$,*

$$\nabla_\mu^{(\rm ad)}\, \bar{J}^\mu_{a,\rm imp} = \sum_k \epsilon_k \left\langle \mathcal{S}_{k,a} \right\rangle_{\boldsymbol{\Psi}} \quad \textit{(eqs. (6.292) and (6.293)).}$$

Then there exists a plateau mask $\mathfrak{M}$ in the $(\lambda_{\rm UV}, \ell)$-grid (cf. Definition 6.300) and a constant $c(\{\epsilon_k\}) > 0$ such that, uniformly on $\mathfrak{M}$,

$$1 - \left|\Lambda_{\rm law}^{(\lambda_{\rm UV},\ell)}\right| \;\geq\; c(\{\epsilon_k\}) \;>\; 0, \qquad \lim_{V\to\infty} \left|\Lambda_{\rm state}^{(\lambda_{\rm UV},\ell)}\right| \;=\; 1,$$

and there are no exact NG modes (only pseudo-NG with GMOR-type masses proportional to spurions) [94, 261, 91, 112].

Proof Theorem 7.557. We prove the three claims in order: (I) a non–vanishing, *uniform* law–visibility deficit on a certified plateau; (II) trivial state–channel (overlap $\to 1$ at large volume); and (III) absence of exact NG together with GMOR–type scaling of pseudo–NG masses. Throughout we work on the physical sector $\mathcal{H}_{\rm phys}$, use the UV/IR taggers of Definition 7.551, the visibilities of Definition 7.553, and the channel split of Definition 7.554, with gauge–parameter independence ensured by Proposition 7.556 and BRST descent by Lemma 7.552.

(I) Non–healing law deficit on a certified plateau. Write

$$\Lambda_{\rm law}^{(\lambda_{\rm UV},\ell)} = \mathrm{Tr}\Big[X^{(\lambda_{\rm UV},\ell)}\, \rho_0^{(\lambda_{\rm UV},\ell)}\Big], \quad X^{(\lambda_{\rm UV},\ell)} := \Big(U_B^{(\lambda_{\rm UV},\ell)}\Big)^\dagger U_A^{(\lambda_{\rm UV},\ell)} \text{ (unitary)}.$$

For any density matrix ρ and unitary W,

$$1 - \mathrm{Re}\,\mathrm{Tr}(\rho W) = \tfrac{1}{2}\,\mathrm{Tr}\Big(\rho\,(I-W)^\dagger(I-W)\Big) - \tfrac{1}{2}\,\mathrm{Tr}\Big(\rho\,(W^\dagger - I)^2\Big) \;\geq\; \tfrac{1}{2}\,\mathrm{Tr}\Big(\rho\,(I-W)^\dagger(I-W)\Big), \tag{7.649}$$

since $W^\dagger W = I$ and $\mathrm{Tr}\Big(\rho A^\dagger A\Big) \geq 0$ for any A. Thus $1 - |\,\mathrm{Tr}(\rho W)| \geq 1 - \mathrm{Re}\,\mathrm{Tr}(\rho W) \geq \tfrac{1}{2}\,\mathrm{Tr}\Big(\rho\,(I-W)^\dagger(I-W)\Big)$. With $\rho = \rho_0^{(\lambda_{\rm UV},\ell)}$ and $W = X^{(\lambda_{\rm UV},\ell)}$,

$$1 - \left|\Lambda_{\rm law}^{(\lambda_{\rm UV},\ell)}\right| \;\geq\; \tfrac{1}{2}\,\mathrm{Tr}\Big(\rho_0^{(\lambda_{\rm UV},\ell)}\, \Delta^\dagger \Delta\Big), \qquad \Delta := I - X^{(\lambda_{\rm UV},\ell)}. \tag{7.650}$$

To relate Δ to the explicit breakings, write (Dyson/Duhamel expansion on the band)

$$X^{(\lambda_{\mathrm{UV}},\ell)} = \mathcal{T}\exp\Big(-i\int_0^1 ds\;\Delta\hat{H}^{(\lambda_{\mathrm{UV}},\ell)}(s)\Big), \qquad \Delta\hat{H}^{(\lambda_{\mathrm{UV}},\ell)}(s) := \Big(\sum_k \epsilon_k\,\hat{\mathcal{O}}_k\Big)_I^{(\lambda_{\mathrm{UV}},\ell)}(s),$$

where the subscript I denotes interaction picture with respect to the common s–dependent interpolation of the law branches (any smooth homotopy suffices; the bound below is homotopy–independent). Then

$$\Delta = \int_0^1 ds\,\Big(-i\Delta\hat{H}^{(\lambda_{\mathrm{UV}},\ell)}(s)\Big) + \int_{0<s_2<s_1<1} ds_1 ds_2\,\Big(-i\Delta\hat{H}(s_1)\Big)\Big(-i\Delta\hat{H}(s_2)\Big) + \cdots .$$

Keeping the exact (non–truncated) series, the quadratic form in (7.650) is bounded from below by the leading non–vanishing contribution and controlled by the *spectral* (T–odd) channel of the holosymmetric readout (see the triadic split of Definition 7.554 and the additive decomposition of Proposition 7.555). The BRST–improved adjoint–covariant Ward identity on $\mathcal{H}_{\mathrm{phys}}$,

$$\nabla_\mu^{(\mathrm{ad})}\,\bar{J}^\mu_{a,\mathrm{imp}} = \sum_k \epsilon_k \left\langle \mathcal{S}_{k,a}\right\rangle_{\mathbf{\Psi}} \quad \text{(eqs. (6.292) and (6.293))},$$

implies that the spectral (commutator/T–odd) cumulants of the band dynamics are *nonzero* whenever $\{\epsilon_k\} \neq 0$ (the only source of non–conservation in QIV). Therefore there is a channel–wise constant $c_{\mathrm{spec}}(\{\epsilon_k\}) > 0$ such that, on any region where the UV/IR tags are stable (no large slope/curvature), the spectral contribution to the right–hand side of (7.650) is bounded below by $c_{\mathrm{spec}}(\{\epsilon_k\})$.

By Definition 6.300 (compactness of the $(\lambda_{\mathrm{UV}},\ell)$–grid, continuity of the tagged maps, and uniform boundedness of the band–restricted generators), there exists a nonempty *plateau mask* $\mathfrak{M}$ on which slopes/curvatures of the tagged observables are within tolerance; hence the spectral lower bound is uniform on $\mathfrak{M}$. Set

$$c(\{\epsilon_k\}) := \tfrac{1}{2}\,c_{\mathrm{spec}}(\{\epsilon_k\}) > 0.$$

Then (7.650) yields, uniformly on $\mathfrak{M}$,

$$1 - \left|\Lambda_{\mathrm{law}}^{(\lambda_{\mathrm{UV}},\ell)}\right| \;\geq\; c(\{\epsilon_k\}) \;>\; 0,$$

establishing the non–healing law deficit. (Gauge–parameter independence of this bound follows from Proposition 7.556.)

(II) Trivial state channel and unique vacuum. Let $|\Omega_{1/2}^{(\lambda_{\mathrm{UV}},\ell)}\rangle = K_{\mathrm{IR}}^{(\ell)} P_x(\lambda_{\mathrm{UV}})\,|\Omega_{1/2}\rangle$ be the band–tagged branch states of Definition 7.551. In QIV, explicit breakings $\{\epsilon_k\} \neq 0$ remove the vacuum manifold and select a *unique* infinite–volume vacuum (GNS representation unique up to phase). Standard exponential clustering for gapped phases and uniqueness of the KMS/GNS state imply that any two finite–volume ground states prepared by local (finite–depth) circuits that respect the same law converge to the *same* infinite–volume state as $V \to \infty$. Since the UV/IR taggers $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ project to long–wavelength content, the

difference between the branch preparations is supported in a boundary layer whose relative weight vanishes with V. Therefore,

$$\Lambda_{\text{state}}^{(\lambda_{\text{UV}},\ell)}(V) = \langle \Omega_2^{(\lambda_{\text{UV}},\ell)} | \Omega_1^{(\lambda_{\text{UV}},\ell)} \rangle \xrightarrow[V\to\infty]{} 1.$$

Equivalently, the fidelity per unit volume tends to 1, which is the precise meaning of a *trivial* state channel (no thermodynamic orthogonality) on QIV plateaus. This discriminates QIV from QII, where multiple vacua give $\Lambda_{\text{state}}(V) \to 0$.

(III) No exact NG and GMOR–type pseudo–NG scaling. Goldstone's theorem requires a conserved Noether current for a continuous symmetry. In QIV, the BRST–improved Ward identity shows that the divergence is *nonzero* and entirely sourced by the explicit spurions: $\nabla_\mu^{(\text{ad})} \bar{J}^\mu_{a,\text{imp}} = \sum_k \epsilon_k \langle \mathcal{S}_{k,a} \rangle_{\mathbf{\Psi}} \neq 0$ on $\mathcal{H}_{\text{phys}}$. Hence no exact NG poles can occur. In the broken–basis (π, h), the band Hessian of Section 7.9 couples the would–be NG fields to the explicit sources; standard current–algebra/PCAC manipulations (integrate the Ward identity against the NG interpolating field and LSZ–reduce the two–point function) give the GMOR–type relation

$$\boxed{f_\pi^2\, m_\pi^2 = \sum_k \epsilon_k \left\langle \mathcal{S}_{k,b}^{(\lambda_{\text{UV}})} \right\rangle_{\mathbf{\Psi}} + \mathcal{O}(\epsilon^2)}$$

where b denotes the broken generator associated with the pseudo–NG mode (i.e., the projection of the Ward source onto that broken direction), and $\mathcal{S}_{k,a}^{(\lambda_{\text{UV}})}$ is the source operator already used in eqs. (6.292) and (6.293). Here f_π is the NG decay constant in the $\epsilon \to 0$ limit. Thus the would–be NG acquire *pseudo–NG* masses proportional to the spurions on the plateau (and remain gauge–parameter independent by Proposition 7.556) [94, 261, 91, 112].

Conclusion. Combining (I)–(III) yields a nonempty plateau mask $\mathfrak{M}$ and a constant $c(\{\epsilon_k\}) > 0$ such that, uniformly on $\mathfrak{M}$,

$$1 - \left|\Lambda_{\text{law}}^{(\lambda_{\text{UV}},\ell)}\right| \geq c(\{\epsilon_k\}) > 0, \qquad \lim_{V\to\infty} \left|\Lambda_{\text{state}}^{(\lambda_{\text{UV}},\ell)}\right| = 1,$$

with no exact NG and pseudo–NG obeying GMOR–type scaling, as claimed. □

Corollary 7.558 (QIV discrimination against QIII and QII)**.** *On $\mathfrak{M}$, QIV satisfies $1 - |\Lambda_{\text{law}}| \geq c(\{\epsilon_k\}) > 0$ and $|\Lambda_{\text{state}}| \to 1$; by contrast, QIII has $1 - |\Lambda_{\text{law}}| \sim \ell^{\Delta-(d+z)} \to 0$ with $|\Lambda_{\text{state}}| \approx 1$, and QII has $|\Lambda_{\text{law}}| \approx 1$ with $|\Lambda_{\text{state}}(V)| \to 0$ as $V \to \infty$.*

Proof Corollary 7.558. Fix a certified plateau mask $\mathfrak{M}$ in the $(\lambda_{\text{UV}}, \ell)$–grid in the sense of Definition 6.300, and work on the physical sector $\mathcal{H}_{\text{phys}}$ with the UV/IR–tagged visibilities of Definition 7.553. Gauge–parameter independence of all statements below follows from Proposition 7.556 together with Lemma 7.552.

QIV behaviour. By Theorem 7.557, when the explicit breakings are nonzero and relevant/-marginal on the band, there exists $c(\{\epsilon_k\}) > 0$ such that, uniformly on $\mathfrak{M}$,

$$1 - \left|\Lambda_{\text{law}}^{(\lambda_{\text{UV}},\ell)}\right| \;\geq\; c(\{\epsilon_k\}) > 0, \qquad \lim_{V\to\infty} \left|\Lambda_{\text{state}}^{(\lambda_{\text{UV}},\ell)}(V)\right| = 1.$$

Thus QIV exhibits a *non–healing* law–visibility deficit and a *trivial* state channel (unique vacuum).

QIII contrast. In Quadrant III (explicit but *small* law breakings; state symmetric), the emergent–symmetry analysis of Section 7.8 says that if the leading breaking operators are RG–irrelevant with scaling dimension Δ, then on any certified plateau one has the improvement law

$$1 - \left|\Lambda_{\text{law}}^{(\lambda_{\text{UV}},\ell)}\right| = \mathcal{O}\Big(\ell^{\,\Delta-(d+z)}\Big) \xrightarrow[\ell\downarrow 0]{} 0, \qquad \left|\Lambda_{\text{state}}^{(\lambda_{\text{UV}},\ell)}\right| \approx 1.$$

(Here d is the spatial dimension and z the dynamical exponent; the statement is the band/IR form of the emergent–symmetry scaling summarized in Section 7.8.) Therefore, unlike QIV, QIII features a *healing* law channel (deficit vanishes with coarse–graining) while the state channel remains trivial.

QII contrast. In Quadrant II (law symmetric; *spontaneously* asymmetric state), the QII diagnostics of Section 7.7 give

$$\left|\Lambda_{\text{law}}^{(\lambda_{\text{UV}},\ell)}\right| \approx 1 \quad \text{on plateaus, while} \quad \lim_{V\to\infty}\left|\Lambda_{\text{state}}^{(\lambda_{\text{UV}},\ell)}(V)\right| = 0,$$

due to thermodynamic orthogonality of distinct vacua (SSB; see also [94, 261, 91] for the standard orthogonality/cluster arguments). Thus, contrary to QIV, QII has an *intact* law channel and a *nontrivial* (decohering) state channel.

Phase	Condition on $\lvert\Lambda_{\text{law}}\rvert$	Condition on $\lvert\Lambda_{\text{state}}\rvert$
QIV:	$1-\lvert\Lambda_{\text{law}}\rvert \;\geq\; c(\{\epsilon_k\}) > 0$	$\lvert\Lambda_{\text{state}}\rvert \;\to\; 1$
QIII:	$1-\lvert\Lambda_{\text{law}}\rvert \;\sim\; \ell^{\Delta-(d+z)} \;\to\; 0$	$\lvert\Lambda_{\text{state}}\rvert \;\approx\; 1$
QII:	$\lvert\Lambda_{\text{law}}\rvert \;\approx\; 1$	$\lvert\Lambda_{\text{state}}(V)\rvert \;\to\; 0 \;(V\to\infty)$

Table 7.3: Comparing the three behaviours on their certified plateaus (denoted generically by $\mathfrak{M}$ for each quadrant):

Discrimination. These signatures are mutually exclusive away from critical boundaries (which are excluded by the plateau tolerances in Definition 6.300), hence they distinguish QIV from QIII and QII as stated. □

Intuition. QIV shows a *non-healing* law deficit and a *trivial* state channel (unique vacuum); QIII heals in the IR (emergent symmetry), and QII displays thermodynamic decoherence of states (vacua orthogonalize).

7.9.2 Fix the Symmetry Reference & Spurions

Definition 7.559 (Reference symmetry data and spurion assignment)**.** Let G be the reference symmetry used in Sections 7.1 and 7.7, with generators $\{T_a\}$ (structure constants $f_{ab}{}^{c}$)

acting on the generalized field stack $\hat{\mathbf{F}} = (\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}})^\top$ via a (possibly reducible) representation ρ. Decompose the band law as in Definition 7.551:

$$\mathcal{L}^{(\lambda_{\mathrm{UV}},\ell)} = \left(\mathcal{L}_0 + \sum_k \epsilon_k \, \mathcal{O}_k + \zeta \Psi\right)^{(\lambda_{\mathrm{UV}},\ell)}.$$

A *spurion assignment* is a choice of transformation rules for the couplings $\{\epsilon_k\}$ such that

$$\delta_a(\epsilon_k \, \mathcal{O}_k) := (\delta_a \epsilon_k) \, \mathcal{O}_k + \epsilon_k \, \delta_a \mathcal{O}_k \;\; = \;\; \nabla_\mu^{(\mathrm{ad})} \, I_a^\mu \;\; + \;\; \zeta(B_a),$$

for some local improvement current I_a^μ (total-derivative improvement) and some local insertion B_a with $\zeta(B_a)$ BRST-exact. Equivalently, on $\mathcal{H}_{\mathrm{phys}}$ the explicit term is G-invariant up to improvements.

BRST improvement (gauge-fixing sector) The improvement term is defined as

$$J^\mu_{a,\mathrm{imp}} = J^\mu_a - \zeta \, \Lambda^\mu_a, \tag{7.651}$$

with the variation given by (eq. (6.292))

$$\delta_a \Psi = \alpha_a \, \chi_a + (\nabla_\mu \alpha_a) \Lambda^\mu_a. \tag{7.652}$$

Spurion improvement (explicit-breaking sector) The ward identity is given by (Definition 7.559)

$$\delta_a(\epsilon_k \mathcal{O}_k) = \nabla_\mu^{(\mathrm{ad})} I_a^\mu + \zeta(B_a). \tag{7.653}$$

Here Λ^μ_a and I^μ_a are *different* objects: Λ^μ_a is fixed by the variation of the gauge fermion Ψ, while I^μ_a collects total-derivative terms that arise when varying the explicit-breaking piece with spurions. Both are legitimate current improvements and can be combined consistently.

Compact band Ward identity (absorbing the spurion improvement). Since currents are defined up to improvements and BRST-exact pieces on $\mathcal{H}_{\mathrm{phys}}$, one may absorb I^μ_a into the improved current,

$$J^\mu_{a,\mathrm{imp}} \;\longrightarrow\; J^\mu_{a,\mathrm{imp}} + I^\mu_a,$$

which yields the scale-tagged Ward identity in the compact form used throughout:

$$\nabla_\mu^{(\mathrm{ad})} \bar{J}^{(\lambda_{\mathrm{UV}},\ell)\mu}_{a,\mathrm{imp}} = \sum_k \epsilon_k \left\langle \mathcal{S}^{(\lambda_{\mathrm{UV}})}_{k,a} \right\rangle_{\mathbf{\Psi}} \quad \text{on } \mathcal{H}_{\mathrm{phys}} \quad \text{(eq. (7.665))}.$$

Intuition. Treating ϵ_k as spurions means we "pretend" symmetry is exact by letting ϵ_k transform in a conjugate representation that cancels the non-invariance of $\mathcal{O}_k$. The remainder is a divergence (improvement) and a BRST-exact term, both harmless for Ward identities on $\mathcal{H}_{\mathrm{phys}}$.

Operator-level AR decomposition: local insertions and their properties

Definition 7.560 (Algebraic–renormalization (AR) decomposition of a local variation)**.** For each local operator $\mathcal{O}_k$ transforming in a (possibly reducible) G-module with generators δ_a, its renormalized local variation admits the AR decomposition

$$\delta_a \mathcal{O}_k \;=\; (R_a)_k{}^j\, \mathcal{O}_j \;+\; \partial_\mu L^\mu_{a,k} \;+\; \zeta(\Sigma_{a,k}), \tag{7.654}$$

where:

- R_a are the representation matrices on the operator index space (so $[R_a, R_b] = f_{ab}{}^c R_c$);
- $L^\mu_{a,k}$ is a local current-density (polynomial/analytic in fields and finitely many derivatives), with ghost number 0 and engineering dimension $\dim(\mathcal{O}_k) - 1$; it packages the total-derivative (Noether-improvement) part of $\delta_a \mathcal{O}_k$ once the linear mixing $(R_a)_k{}^j \mathcal{O}_j$ is peeled off;
- $\Sigma_{a,k}$ is a local functional with ghost number -1 such that $\zeta(\Sigma_{a,k})$ has ghost number 0 and represents the BRST-exact remainder of the variation that is not a total derivative.

Remark 7.561 (Name clash is avoided)**.** The symbol $L^\mu_{a,k}$ here is *not* the same as Λ^μ_a used in the BRST improvement of the Noether current, $J^\mu_{a,\mathrm{imp}} = J^\mu_a - \zeta\, \Lambda^\mu_a$ (eq. (6.292)). We reserve Λ^μ_a exclusively for eq. (6.292) and use $L^\mu_{a,k}$ for the operator-level insertion in (7.654).

Lemma 7.562 (Allowed redefinitions (equivalence class of insertions))**.** *The pair $(L^\mu_{a,k}, \Sigma_{a,k})$ in (7.654) is defined only up to*

$$L^\mu_{a,k} \;\mapsto\; L^\mu_{a,k} + \partial_\nu K^{[\nu\mu]}_{a,k} \;+\; \zeta(U^\mu_{a,k}), \qquad \Sigma_{a,k} \;\mapsto\; \Sigma_{a,k} \;-\; \partial_\mu U^\mu_{a,k}, \tag{7.655}$$

where $K^{[\nu\mu]}_{a,k}$ is antisymmetric in (ν,μ) and $U^\mu_{a,k}$ is a local functional of ghost number -1. These redefinitions leave the combination $\partial_\mu L^\mu_{a,k} + \zeta(\Sigma_{a,k})$ unchanged.

Proof. Since $\partial_\mu \partial_\nu K^{[\nu\mu]}_{a,k} = 0$ identically and $[\zeta, \partial_\mu] = 0$, we have

$$\partial_\mu\Big(L^\mu_{a,k} + \partial_\nu K^{[\nu\mu]}_{a,k} + \zeta(U^\mu_{a,k})\Big) + \zeta\Big(\Sigma_{a,k} - \partial_\mu U^\mu_{a,k}\Big) = \partial_\mu L^\mu_{a,k} + \zeta(\Sigma_{a,k}).$$

□

Lemma 7.563 (Ghost numbers, dimensions, and covariance)**.** *With the assignments $\mathrm{gh}(L^\mu_{a,k}) = 0$ and $\mathrm{gh}(\Sigma_{a,k}) = -1$, the combination $\partial_\mu L^\mu_{a,k} + \zeta(\Sigma_{a,k})$ has ghost number 0 and the same engineering dimension as $\mathcal{O}_k$. Moreover, under the G-action,*

$$\delta_b\Big(\delta_a \mathcal{O}_k - (R_a)_k{}^j \mathcal{O}_j\Big) = f_{ba}{}^c\, \Big(\partial_\mu L^\mu_{c,k} + \zeta(\Sigma_{c,k})\Big)\,,$$

up to further redefinitions of the form (7.655). Hence (7.654) is compatible with the Wess–Zumino consistency condition.

Proof. Ghost numbers. By construction $\mathrm{gh}(\mathcal{O}_k) = 0$, $\mathrm{gh}(\zeta) = +1$, and $\mathrm{gh}(\partial_\mu) = 0$. With $\mathrm{gh}(L^\mu_{a,k}) = 0$ and $\mathrm{gh}(\Sigma_{a,k}) = -1$,

$$\mathrm{gh}\Big(\partial_\mu L^\mu_{a,k}\Big) = 0, \qquad \mathrm{gh}(\zeta(\Sigma_{a,k})) = \mathrm{gh}(\zeta) + \mathrm{gh}(\Sigma_{a,k}) = 1 + (-1) = 0.$$

Hence $\mathrm{gh}\big(\partial_\mu L^\mu_{a,k} + \zeta(\Sigma_{a,k})\big) = 0$, matching $\mathrm{gh}(\delta_a \mathcal{O}_k - (R_a)_k{}^j \mathcal{O}_j)$.

Engineering dimensions. Let $\dim[\mathcal{O}_k] = \Delta_k$ (engineering dimension). The canonical decomposition

$$\delta_a \mathcal{O}_k - (R_a)_k{}^j \mathcal{O}_j \;=\; \partial_\mu L^\mu_{a,k} + \zeta(\Sigma_{a,k}) \tag{7.656}$$

must preserve dimensions. Since $\dim[\partial_\mu] = 1$ and ζ is dimensionless in the BRST complex, choosing

$$\dim[L^\mu_{a,k}] = \Delta_k - 1, \qquad \dim[\Sigma_{a,k}] = \Delta_k$$

ensures $\dim[\partial_\mu L^\mu_{a,k}] = \dim[\zeta(\Sigma_{a,k})] = \Delta_k$. These assignments can always be reached within the improvement freedom (7.655) (see below), which allows adding total derivatives and ζ–exact terms that do not change (7.656) but do adjust the dimensions of the representatives.

Covariance and Wess–Zumino (WZ) consistency. Define the remainder

$$\Delta_{a,k} \;:=\; \delta_a \mathcal{O}_k - (R_a)_k{}^j \mathcal{O}_j \;=\; \partial_\mu L^\mu_{a,k} + \zeta(\Sigma_{a,k}).$$

WZ consistency requires

$$\delta_b \Delta_{a,k} - \delta_a \Delta_{b,k} \;=\; f_{ba}{}^c \, \Delta_{c,k}. \tag{7.657}$$

Using that δ_b is an internal (global) symmetry derivation commuting with spacetime derivatives and with the BRST differential on local operators,[16]

$$\delta_b \Delta_{a,k} = \partial_\mu(\delta_b L^\mu_{a,k}) + \zeta(\delta_b \Sigma_{a,k}), \qquad \delta_a \Delta_{b,k} = \partial_\mu(\delta_a L^\mu_{b,k}) + \zeta(\delta_a \Sigma_{b,k}).$$

Subtracting and comparing with the right-hand side of (7.657) gives

$$\partial_\mu\Big(\delta_b L^\mu_{a,k} - \delta_a L^\mu_{b,k} - f_{ba}{}^c L^\mu_{c,k}\Big) \;+\; \zeta(\delta_b \Sigma_{a,k} - \delta_a \Sigma_{b,k} - f_{ba}{}^c \Sigma_{c,k}) \;=\; 0.$$

By locality and the improvement freedom (7.655), any local expression of ghost number 0 and dimension Δ_k that is annihilated by the operator $\partial_\mu \oplus \zeta$ can be written as a sum of a total derivative and a BRST–exact term. Concretely, there exist local tensors $U^{[\nu\mu]}_{ab,k}$ (antisymmetric in ν, μ), $V^\mu_{ab,k}$ and $W_{ab,k}$ with

$$\begin{aligned} \mathrm{gh}(U) &= 0, \quad \dim[U] = \Delta_k - 2; \\ \mathrm{gh}(V) &= -1, \quad \dim[V] = \Delta_k - 1; \\ \mathrm{gh}(W) &= -2, \quad \dim[W] = \Delta_k. \end{aligned}$$

such that

$$\delta_b L^\mu_{a,k} - \delta_a L^\mu_{b,k} - f_{ba}{}^c L^\mu_{c,k} = \partial_\nu U^{[\nu\mu]}_{ab,k} + \zeta(V^\mu_{ab,k}), \tag{7.658}$$

$$\delta_b \Sigma_{a,k} - \delta_a \Sigma_{b,k} - f_{ba}{}^c \Sigma_{c,k} = -\,\partial_\mu V^\mu_{ab,k} + \zeta(W_{ab,k}). \tag{7.659}$$

Plugging (7.658)–(7.659) back into the previous display and using $[\partial_\mu, \zeta] = 0$, $\zeta^2 = 0$, and $\partial_\mu \partial_\nu U^{[\nu\mu]}_{ab,k} \equiv 0$, one verifies (7.657).

[16] On the local algebra one has $[\delta_b, \partial_\mu] = 0$ and $[\delta_b, \zeta] = 0$; the latter expresses the G–invariance of the BRST complex (Nielsen/BRST robustness on $\mathcal{H}_{\mathrm{phys}}$).

Finally, redefine the representatives by the allowed improvements (the "AR redefinitions" (7.655))

$$L^{\mu}_{a,k} \;\mapsto\; L^{\mu}_{a,k} - \tfrac{1}{2}\Big(\partial_\nu U^{[\nu\mu]}_{ab,k} + \zeta V^{\mu}_{ab,k}\Big), \qquad \Sigma_{a,k} \;\mapsto\; \Sigma_{a,k} - \tfrac{1}{2}\Big(-\partial_\mu V^{\mu}_{ab,k} + \zeta W_{ab,k}\Big),$$

which leave $\partial_\mu L^{\mu}_{a,k} + \zeta(\Sigma_{a,k})$ unchanged but make the transformed objects strictly covariant:

$$\delta_b L^{\mu}_{a,k} - \delta_a L^{\mu}_{b,k} = f_{ba}{}^{c} L^{\mu}_{c,k}, \qquad \delta_b \Sigma_{a,k} - \delta_a \Sigma_{b,k} = f_{ba}{}^{c} \Sigma_{c,k}.$$

Thus (7.654) is compatible with WZ consistency, completing the proof. □

Corollary 7.564 (Induced spurion-sector objects are unique on $\mathcal{H}_{\text{phys}}$)**.** *Upon multiplying* (7.654) *by* ϵ_k *and summing over* k*, define*

$$I^{\mu}_{a} := \epsilon_k\, L^{\mu}_{a,k}, \qquad B_a := \epsilon_k\, \Sigma_{a,k}.$$

Under (7.655)*,* $I^{\mu}_{a} \mapsto I^{\mu}_{a} + \partial_\nu K^{[\nu\mu]}_{a}$ *and* $B_a \mapsto B_a - \partial_\mu U^{\mu}_{a}$*, so* $\nabla^{(\text{ad})}_{\mu} I^{\mu}_{a} + \zeta(B_a)$ *is unchanged and, on* $\mathcal{H}_{\text{phys}}$*, its insertions are insensitive to these redefinitions (Noether improvement and BRST exactness drop out).*

Proof. Step 1: Invariance of the composite operator. Starting from (7.654),

$$\delta_a \mathcal{O}_k - (R_a)_k{}^{j} \mathcal{O}_j \;=\; \partial_\mu L^{\mu}_{a,k} + \zeta(\Sigma_{a,k}),$$

multiply by ϵ_k and sum over k (explicit-breaking spurions are constants in QIV), obtaining

$$\epsilon_k\Big(\delta_a \mathcal{O}_k - (R_a)_k{}^{j} \mathcal{O}_j\Big) \;=\; \partial_\mu I^{\mu}_{a} + \zeta(B_a), \qquad I^{\mu}_{a} := \epsilon_k L^{\mu}_{a,k}, \quad B_a := \epsilon_k \Sigma_{a,k}. \tag{7.660}$$

Under the allowed improvements (7.655) one may take, without loss of generality,

$$L^{\mu}_{a,k} \mapsto L^{\mu}_{a,k} + \partial_\nu K^{[\nu\mu]}_{a,k}, \qquad \Sigma_{a,k} \mapsto \Sigma_{a,k} - \partial_\mu U^{\mu}_{a,k},$$

(possible extra ζ–exact pieces can be absorbed into a redefinition of U, K). Summing with ϵ_k gives

$$I^{\mu}_{a} \mapsto I^{\mu}_{a} + \partial_\nu K^{[\nu\mu]}_{a}, \qquad B_a \mapsto B_a - \partial_\mu U^{\mu}_{a},$$

with $K^{[\nu\mu]}_{a} := \epsilon_k K^{[\nu\mu]}_{a,k}$ and $U^{\mu}_{a} := \epsilon_k U^{\mu}_{a,k}$. Therefore

$$\begin{aligned} \partial_\mu I^{\mu}_{a} + \zeta(B_a) \;&\mapsto\; \partial_\mu\Big(I^{\mu}_{a} + \partial_\nu K^{[\nu\mu]}_{a}\Big) + \zeta(B_a - \partial_\mu U^{\mu}_{a}) \\ &=\; \partial_\mu I^{\mu}_{a} + \underbrace{\partial_\mu\partial_\nu K^{[\nu\mu]}_{a}}_{=0} + \zeta(B_a) - \underbrace{\partial_\mu \zeta(U^{\mu}_{a})}_{[\partial,\zeta]=0} \;=\; \partial_\mu I^{\mu}_{a} + \zeta(B_a). \end{aligned}$$

Since $\nabla^{(\text{ad})}_{\mu}$ acts as ∂_μ on adjoint scalars and as the adjoint covariant divergence on adjoint vectors, replacing ∂_μ by $\nabla^{(\text{ad})}_{\mu}$ in (7.660) preserves the same conclusion: the curl improvement contributes a double divergence that vanishes identically, and the ζ–improvement contributes a BRST-exact divergence. Hence the composite

$$\mathcal{J}_a \;:=\; \nabla^{(\text{ad})}_{\mu} I^{\mu}_{a} + \zeta(B_a)$$

is invariant under (7.655).

Step 2: Invariance of insertions on $\mathcal{H}_{\rm phys}$. Let $\mathcal{W}$ be any time-ordered product of BRST-closed, gauge-invariant operators supported away from the boundary (the standard observable class used on plateaus). Consider the change in any insertion:

$$\Delta \equiv \Big\langle\, T\{\mathcal{W}\ \mathcal{J}_a\}\,\Big\rangle \ \mapsto\ \Big\langle\, T\{\mathcal{W}\ (\mathcal{J}_a + \partial_\mu\partial_\nu K_a^{[\nu\mu]} - \partial_\mu\zeta(U_a^\mu))\}\,\Big\rangle.$$

The double divergence integrates to a boundary term (or, in momentum space, carries a factor $p_\mu p_\nu$ that vanishes in the zero-momentum limit used to define the plateau readouts). By the usual Noether-improvement argument, such total derivatives do not affect correlation functions of local observables with compact support. The BRST-exact contribution vanishes on $\mathcal{H}_{\rm phys}$ by the quartet mechanism/Nielsen identities: for BRST-closed $\mathcal{W}$, $\langle T\{\mathcal{W}\,\zeta(X)\}\rangle = 0$ because ζ acts as a graded derivation, $\zeta\mathcal{W} = 0$, and $\zeta\,|\Omega\rangle = \langle\Omega|\,\zeta = 0$ on the physical vacuum. Therefore Δ is unchanged.

Step 3: Covariance and uniqueness class. By Lemma 7.563, the representatives can be chosen to transform covariantly under the G–action, and they have the correct ghost numbers and engineering dimensions. The preceding steps show that all choices related by (7.655) yield the same composite $\mathcal{J}_a$ and the same physical insertions. Hence the induced spurion-sector objects are unique on $\mathcal{H}_{\rm phys}$, completing the proof. □

Remark 7.565 (Operational recipe and disambiguation)**.** To compute the spurion contribution in practice: (i) evaluate $\delta_a\mathcal{O}_k$ and separate the pure mixing $(R_a)_k{}^j\mathcal{O}_j$; (ii) integrate by parts to isolate $\partial_\mu L^\mu_{a,k}$; (iii) identify the remainder as $\zeta(\Sigma_{a,k})$. This $L^\mu_{a,k}$ is unrelated to the gauge-fermion insertion Λ^μ_a in eq. (6.292); the latter only enters $J^\mu_{a,\rm imp}$, while $L^\mu_{a,k}$ (through I^μ_a) enters the spurion-improved Ward identity Definition 7.559.

Lemma 7.566 (Existence and (cohomological) uniqueness of spurion laws)**.** *Given $\{\mathcal{O}_k\}$ forming a finite-dimensional G-module under δ_a, there exist linear spurion transformation laws $\delta_a\epsilon_k = -(R_a)_k{}^j\,\epsilon_j$ (with R_a the dual representation matrices to those acting on $\{\mathcal{O}_k\}$) that realize Definition 7.559. Any two such choices differ by redefinitions $\epsilon_k \mapsto \epsilon_k + \zeta(\chi_k)$ and improvements $I^\mu_a \mapsto I^\mu_a + \partial_\nu K_a^{[\nu\mu]}$ which are invisible on $\mathcal{H}_{\rm phys}$.*

Proof Lemma 7.566. **Step 1: Local variation of the breaking operators.** Because $\{\mathcal{O}_k\}$ is a G-module, the renormalized local variation admits the algebraic–renormalization decomposition

$$\delta_a\mathcal{O}_k = (R_a)_k{}^j\,\mathcal{O}_j \ +\ \partial_\mu L^\mu_{a,k} \ +\ \zeta(\Sigma_{a,k}), \tag{7.661}$$

with local insertions $L^\mu_{a,k}$ (ghost number 0) and $\Sigma_{a,k}$ (ghost number -1); here R_a furnish a (possibly reducible) matrix representation of $\mathfrak{g} = \mathrm{Lie}(G)$ on the operator index k, and the last term encodes BRST-exact pieces.

Step 2: Existence (constructive spurion rule). Define the spurion transformation as the *dual* action:

$$\delta_a\epsilon_k := -(R_a)_k{}^j\,\epsilon_j. \tag{7.662}$$

Then

$$\begin{aligned}\delta_a(\epsilon_k\mathcal{O}_k) &= (\delta_a\epsilon_k)\mathcal{O}_k + \epsilon_k\delta_a\mathcal{O}_k\\ &= -\epsilon_j(R_a)_k{}^j\mathcal{O}_k + \epsilon_k(R_a)_k{}^j\mathcal{O}_j \ +\ \partial_\mu(\epsilon_k L^\mu_{a,k}) \ +\ \zeta(\epsilon_k\Sigma_{a,k}).\end{aligned}$$

Relabel the dummy indices in the first two terms; they cancel identically. Hence

$$\delta_a(\epsilon_k \mathcal{O}_k) = \partial_\mu(\epsilon_k L^\mu_{a,k}) + \zeta(\epsilon_k \Sigma_{a,k}).$$

Setting

$$I^\mu_a := \epsilon_k L^\mu_{a,k}, \qquad B_a := \epsilon_k \Sigma_{a,k}, \tag{7.663}$$

we obtain exactly the structure required in Definition 7.559: $\delta_a(\epsilon_k \mathcal{O}_k) = \nabla^{(\mathrm{ad})}_\mu I^\mu_a + \zeta(B_a)$. This proves *existence*.

Step 3: Consistency with the Lie algebra (Wess–Zumino condition). Using (7.662),

$$[\delta_a, \delta_b]\epsilon_k = -(R_a)_k{}^j \delta_b \epsilon_j + (a \leftrightarrow b) = (R_a R_b - R_b R_a)_k{}^j \epsilon_j = -[R_a, R_b]_k{}^j \epsilon_j = f_{ab}{}^c \, \delta_c \epsilon_k,$$

since $[R_a, R_b] = f_{ab}{}^c R_c$. Thus the spurion assignment is algebraically consistent and transforms covariantly.

Step 4: Basis independence. Under a change of operator basis $\mathcal{O}'_k = M_k{}^j \mathcal{O}_j$, the representation matrices transform as $R'_a = M R_a M^{-1}$. Choosing the dual spurion basis $\epsilon' = (M^{-T})\,\epsilon$ makes (7.662) basis covariant: $\delta_a \epsilon' = -R'_a \, \epsilon'$. Therefore the construction is intrinsic to the G-module carried by $\{\mathcal{O}_k\}$.

Step 5: Cohomological uniqueness (ambiguities are ζ-exact or improvements). Suppose $\delta'_a \epsilon_k$ is another spurion rule that also realizes Definition 7.559, i.e.

$$\delta'_a(\epsilon_k \mathcal{O}_k) = \partial_\mu I'^\mu_a + \zeta(B'_a).$$

Subtracting the two realizations gives

$$\big(\delta'_a - \delta_a\big)\epsilon_k \; \mathcal{O}_k = \partial_\mu(I'^\mu_a - I^\mu_a) + \zeta(B'_a - B_a). \tag{7.664}$$

By (7.661), the cohomology of local functionals at ghost number 0 modulo total derivatives is spanned by the classes of $\{\mathcal{O}_k\}$; in that quotient the right-hand side of (7.664) vanishes. Hence, in $H^0(\zeta \mid d)$,

$$\Big[(\delta'_a - \delta_a)\epsilon_k\Big]\,[\mathcal{O}_k] = 0.$$

Because the classes $[\mathcal{O}_k]$ are linearly independent in the chosen basis, it follows that $(\delta'_a - \delta_a)\epsilon_k$ is *cohomologically trivial*: there exist local functionals χ_k of ghost number -1 such that

$$(\delta'_a - \delta_a)\epsilon_k = \zeta(\chi_k).$$

Plugging this into (7.664), the entire difference is absorbed into a ζ-exact term plus a total derivative. In other words, the two realizations differ by:

$$\epsilon_k \;\mapsto\; \epsilon_k + \zeta(\chi_k), \qquad I^\mu_a \;\mapsto\; I^\mu_a + \partial_\nu K^{[\nu\mu]}_a,$$

for some antisymmetric $K^{[\nu\mu]}_a$. Both changes are invisible on $\mathcal{H}_{\mathrm{phys}}$: ζ-exact deformations drop out of physical matrix elements, and Noether improvements (total divergences or antisymmetric superpotentials) do not affect Ward identities or conserved charges. This proves *cohomological uniqueness*.

Conclusion. The dual spurion rule (7.662) exists, is algebraically consistent, basis independent, and is unique up to BRST-exact redefinitions of the spurions and standard Noether improvements, as claimed. □

Definition 7.567 (Scale-tagged BRST-improved current and Ward identity)**.** Let $J^{\mu}_{a,\mathrm{imp}}$ be the improved current of eq. (6.292). Define its band-tagged version by

$$\bar{J}^{(\lambda_{\mathrm{UV}},\ell)\mu}_{a,\mathrm{imp}} := \Big(J^{\mu}_{a,\mathrm{imp}}\Big)^{(\lambda_{\mathrm{UV}},\ell)} = K^{(\ell)}_{\mathrm{IR}}\,P_x(\lambda_{\mathrm{UV}})\,J^{\mu}_{a,\mathrm{imp}}\,P_x(\lambda_{\mathrm{UV}})K^{(\ell)}_{\mathrm{IR}}.$$

On the physical sector $\mathcal{H}_{\mathrm{phys}}$ the *scale-tagged Ward identity* reads

$$\nabla^{(\mathrm{ad})}_{\mu}\,\bar{J}^{(\lambda_{\mathrm{UV}},\ell)\mu}_{a,\mathrm{imp}} = \sum_k \epsilon_k \left\langle \mathcal{S}^{(\lambda_{\mathrm{UV}})}_{k,a} \right\rangle_{\boldsymbol{\Psi}} \qquad \text{on } \mathcal{H}_{\mathrm{phys}}, \tag{7.665}$$

where $\mathcal{S}^{(\lambda_{\mathrm{UV}})}_{k,a}$ is the (UV-tagged) source operator appearing in eq. (6.293).

Intuition. Because $P_x(\lambda_{\mathrm{UV}})$ and $K^{(\ell)}_{\mathrm{IR}}$ commute with ζ and respect covariance, the textbook Ward identity of eq. (6.293) descends to the band and remains adjoint-covariant; the explicit spurions are the *only* sources of non-conservation in QIV.

Proposition 7.568 (Adjoint covariance of the band Ward identity)**.** *Under the spurion law of Lemma 7.566, the divergence $\nabla^{(\mathrm{ad})}_{\mu}\bar{J}^{(\lambda_{\mathrm{UV}},\ell)\mu}_{a,\mathrm{imp}}$ transforms in the adjoint:*

$$\delta_b\Big(\nabla^{(\mathrm{ad})}_{\mu}\bar{J}^{(\lambda_{\mathrm{UV}},\ell)\mu}_{a,\mathrm{imp}}\Big) = f_{ba}{}^{c}\,\nabla^{(\mathrm{ad})}_{\mu}\bar{J}^{(\lambda_{\mathrm{UV}},\ell)\mu}_{c,\mathrm{imp}},$$

and the right-hand side of (7.665) transforms accordingly, so (7.665) is covariant.

Proof Proposition 7.568. We proceed in three steps: (I) adjoint covariance for the *untagged* current, (II) descent to the band via UV/IR taggers, and (III) adjoint covariance of the explicit source term.

(I) Untagged improved current transforms in the adjoint. By definition (eq. (6.292)), the improved Noether current $J^{\mu}_{a,\mathrm{imp}}$ is obtained from the canonical Noether current by a BRST improvement. Its symmetry variation has the standard form

$$\delta_b J^{\mu}_{a,\mathrm{imp}} = f_{ba}{}^{c}\,J^{\mu}_{c,\mathrm{imp}} \;+\; \partial_{\nu}K^{[\nu\mu]}_{ab} \;+\; \zeta\Big(B^{\mu}_{ab}\Big), \tag{7.666}$$

where $K^{[\nu\mu]}_{ab}$ is an antisymmetric superpotential (Noether improvement) and B^{μ}_{ab} is a local insertion with $\zeta(B^{\mu}_{ab})$ BRST-exact. Applying the adjoint-covariant divergence $\nabla^{(\mathrm{ad})}_{\mu}$ (which acts as a derivation compatible with the adjoint action),

$$\delta_b\Big(\nabla^{(\mathrm{ad})}_{\mu}J^{\mu}_{a,\mathrm{imp}}\Big) = f_{ba}{}^{c}\,\nabla^{(\mathrm{ad})}_{\mu}J^{\mu}_{c,\mathrm{imp}} \;+\; \nabla^{(\mathrm{ad})}_{\mu}\partial_{\nu}K^{[\nu\mu]}_{ab} \;+\; \nabla^{(\mathrm{ad})}_{\mu}\zeta\Big(B^{\mu}_{ab}\Big).$$

The last two terms are improvements/BRST-exact and therefore cohomologically trivial on $\mathcal{H}_{\mathrm{phys}}$ (cf. Lemma 7.552), so the untagged divergence transforms in the adjoint, modulo trivial terms.

(II) Descent to the band. Define the band-tagged current as

$$\bar{J}^{(\lambda_{\mathrm{UV}},\ell)\mu}_{a,\mathrm{imp}} := K^{(\ell)}_{\mathrm{IR}} P_x(\lambda_{\mathrm{UV}})\, J^{\mu}_{a,\mathrm{imp}}\, P_x(\lambda_{\mathrm{UV}}) K^{(\ell)}_{\mathrm{IR}}. \tag{7.667}$$

By Definition 7.551, the taggers commute with both the BRST differential and the G-action: $[\zeta, P_x] = [\zeta, K^{(\ell)}_{\mathrm{IR}}] = 0$ and $[\delta_b, P_x] = [\delta_b, K^{(\ell)}_{\mathrm{IR}}] = 0$, and they preserve improvements. Therefore, acting with δ_b on the tagged object simply tags (7.666):

$$\delta_b \bar{J}^{(\lambda_{\mathrm{UV}},\ell)\mu}_{a,\mathrm{imp}} = f_{ba}{}^{c}\, \bar{J}^{(\lambda_{\mathrm{UV}},\ell)\mu}_{c,\mathrm{imp}} \;+\; \partial_\nu \Big(K^{[\nu\mu]}_{ab}\Big)^{(\lambda_{\mathrm{UV}},\ell)} \;+\; \zeta\Big(B^{\mu}_{ab}\Big)^{(\lambda_{\mathrm{UV}},\ell)}.$$

Applying $\nabla^{(\mathrm{ad})}_{\mu}$ and using the same commutation/linearity properties gives

$$\delta_b\Big(\nabla^{(\mathrm{ad})}_{\mu} \bar{J}^{(\lambda_{\mathrm{UV}},\ell)\mu}_{a,\mathrm{imp}}\Big) = f_{ba}{}^{c}\, \nabla^{(\mathrm{ad})}_{\mu} \bar{J}^{(\lambda_{\mathrm{UV}},\ell)\mu}_{c,\mathrm{imp}} \;+\; \nabla^{(\mathrm{ad})}_{\mu}\partial_\nu \Big(K^{[\nu\mu]}_{ab}\Big)^{(\lambda_{\mathrm{UV}},\ell)} \;+\; \nabla^{(\mathrm{ad})}_{\mu}\zeta\Big(B^{\mu}_{ab}\Big)^{(\lambda_{\mathrm{UV}},\ell)}.$$

Again, the last two terms are improvements/BRST-exact and thus invisible on $\mathcal{H}_{\mathrm{phys}}$ (Lemma 7.552). This proves the first displayed equality in the proposition.

(III) Adjoint covariance of the explicit source (right-hand side). Consider the right-hand side of the scale-tagged Ward identity (Definition 7.567):

$$\mathcal{R}^{(\lambda_{\mathrm{UV}})}_{a} := \sum_{k} \epsilon_k \left\langle \mathcal{S}^{(\lambda_{\mathrm{UV}})}_{k,a} \right\rangle_{\boldsymbol{\Psi}}.$$

By Lemma 7.566, the spurions transform in the dual representation, $\delta_b \epsilon_k = -(R_b)_k{}^{j} \epsilon_j$. The source operators transform as

$$\delta_b \mathcal{S}_{k,a} = (R_b)_k{}^{j}\, \mathcal{S}_{j,a} \;+\; f_{ba}{}^{c}\, \mathcal{S}_{k,c} \;+\; \partial_\mu(\dots) \;+\; \zeta(\dots),$$

where the last two pieces are improvement/BRST-exact. Tagging is compatible with these variations by the commutation properties from Definition 7.551. Therefore

$$\begin{aligned}
\delta_b \mathcal{R}^{(\lambda_{\mathrm{UV}})}_{a} &= \sum_{k} \Big(\delta_b \epsilon_k\Big) \left\langle \mathcal{S}^{(\lambda_{\mathrm{UV}})}_{k,a} \right\rangle_{\boldsymbol{\Psi}} \\
&\qquad + \sum_{k} \epsilon_k \left\langle \delta_b \mathcal{S}^{(\lambda_{\mathrm{UV}})}_{k,a} \right\rangle_{\boldsymbol{\Psi}} \\
&= \sum_{k} \Big(- (R_b)_k{}^{j} \epsilon_j \Big) \left\langle \mathcal{S}^{(\lambda_{\mathrm{UV}})}_{k,a} \right\rangle_{\boldsymbol{\Psi}} \\
&\qquad + \sum_{k} \epsilon_k \Big[(R_b)_k{}^{j} \left\langle \mathcal{S}^{(\lambda_{\mathrm{UV}})}_{j,a} \right\rangle_{\boldsymbol{\Psi}} \\
&\qquad\qquad + f_{ba}{}^{c} \left\langle \mathcal{S}^{(\lambda_{\mathrm{UV}})}_{k,c} \right\rangle_{\boldsymbol{\Psi}} \\
&\qquad\qquad + \langle \partial_\mu(\dots) + \zeta(\dots) \rangle \Big].
\end{aligned}$$

The terms proportional to R_b cancel after relabeling dummy indices. Expectations of improvements and ζ-exact insertions vanish on $\mathcal{H}_{\mathrm{phys}}$ by Lemma 7.552. We are left with

$$\delta_b \mathcal{R}^{(\lambda_{\mathrm{UV}})}_{a} = f_{ba}{}^{c} \sum_{k} \epsilon_k \left\langle \mathcal{S}^{(\lambda_{\mathrm{UV}})}_{k,c} \right\rangle_{\boldsymbol{\Psi}} = f_{ba}{}^{c}\, \mathcal{R}^{(\lambda_{\mathrm{UV}})}_{c}.$$

Thus the right-hand side of (7.665) transforms in the adjoint.

Conclusion. Both sides of the band Ward identity (7.665) transform covariantly in the adjoint representation (modulo improvement/BRST-exact terms that drop on $\mathcal{H}_{\text{phys}}$). Hence the identity is adjoint-covariant, as claimed. □

Proposition 7.569 (Gauge-parameter independence of the band Ward identity)**.** *The relation* (7.665) *is independent of gauge-fixing parameters on* $\mathcal{H}_{\text{phys}}$.

Proof Proposition 7.569. Let α_{gf} denote any gauge-fixing parameter entering through the gauge-fermion Ψ. Promote α_{gf} to a BRST doublet in the algebraic–renormalization sense; the *Nielsen identity* (linearized Slavnov–Taylor form) then reads, for every band-tagged operator $X^{(\lambda_{\text{UV}},\ell)}$ and the band state $\rho_0^{(\lambda_{\text{UV}},\ell)}$ (cf. Definition 7.551, Lemma 7.552, and Proposition 7.556),

$$\partial_{\alpha_{\text{gf}}} X^{(\lambda_{\text{UV}},\ell)} = \zeta\Big(\mathcal{N}_{\alpha_{\text{gf}}}[X]^{(\lambda_{\text{UV}},\ell)}\Big), \qquad \partial_{\alpha_{\text{gf}}} \rho_0^{(\lambda_{\text{UV}},\ell)} = \zeta\Big(\mathcal{N}_{\alpha_{\text{gf}}}[\rho_0]^{(\lambda_{\text{UV}},\ell)}\Big), \tag{7.668}$$

with some local insertions $\mathcal{N}_{\alpha_{\text{gf}}}[\cdot]$. Here we used that the UV/IR taggers are α_{gf}-independent and BRST-closed: $\partial_{\alpha_{\text{gf}}} P_x = \partial_{\alpha_{\text{gf}}} K_{\text{IR}}^{(\ell)} = 0$ and $[\zeta, P_x] = [\zeta, K_{\text{IR}}^{(\ell)}] = 0$ (Definition 7.551).

Define the band Ward functional (on $\mathcal{H}_{\text{phys}}$)

$$\mathcal{W}_a^{(\lambda_{\text{UV}},\ell)} := \nabla_\mu^{(\text{ad})} \bar{J}_{a,\text{imp}}^{(\lambda_{\text{UV}},\ell)\mu} - \sum_k \epsilon_k \left\langle \mathcal{S}_{k,a}^{(\lambda_{\text{UV}})} \right\rangle_{\boldsymbol{\Psi}}.$$

The scale-tagged Ward identity (7.665) states $\mathcal{W}_a^{(\lambda_{\text{UV}},\ell)} = 0$ on $\mathcal{H}_{\text{phys}}$. We show $\partial_{\alpha_{\text{gf}}} \mathcal{W}_a^{(\lambda_{\text{UV}},\ell)} = 0$ there, whence $\mathcal{W}_a^{(\lambda_{\text{UV}},\ell)}$ is α_{gf}-independent and, being zero for one value of α_{gf}, is zero for all.

LHS variation is ζ-exact. Using (7.668) with $X = \bar{J}_{a,\text{imp}}^{\mu}$ and the commutation $[\zeta, \nabla_\mu^{(\text{ad})}] = 0$,

$$\begin{aligned} \partial_{\alpha_{\text{gf}}}\Big(\nabla_\mu^{(\text{ad})} \bar{J}_{a,\text{imp}}^{(\lambda_{\text{UV}},\ell)\mu}\Big) &= \nabla_\mu^{(\text{ad})}\, \partial_{\alpha_{\text{gf}}} \bar{J}_{a,\text{imp}}^{(\lambda_{\text{UV}},\ell)\mu} \\ &= \nabla_\mu^{(\text{ad})}\, \zeta\Big(\mathcal{N}_{\alpha_{\text{gf}}}[J_{a,\text{imp}}^{\mu}]^{(\lambda_{\text{UV}},\ell)}\Big) \\ &= \zeta\Big(\nabla_\mu^{(\text{ad})} \mathcal{N}_{\alpha_{\text{gf}}}[J_{a,\text{imp}}^{\mu}]^{(\lambda_{\text{UV}},\ell)}\Big). \end{aligned}$$

Hence the entire LHS variation is a ζ-exact insertion.

RHS variation is ζ-exact in expectation. Differentiate the expectation value:

$$\partial_{\alpha_{\text{gf}}}\Big(\sum_k \epsilon_k \left\langle \mathcal{S}_{k,a}^{(\lambda_{\text{UV}})} \right\rangle_{\boldsymbol{\Psi}}\Big) = \sum_k \epsilon_k\, \partial_{\alpha_{\text{gf}}} \left\langle \mathcal{S}_{k,a}^{(\lambda_{\text{UV}})} \right\rangle_{\boldsymbol{\Psi}}.$$

Using the band Nielsen identity for expectations (derived from (7.668) and the graded Leibniz rule),

$$\partial_{\alpha_{\text{gf}}} \left\langle \mathcal{S}_{k,a}^{(\lambda_{\text{UV}})} \right\rangle_{\boldsymbol{\Psi}} = \text{Tr}\Big(\rho_0^{(\lambda_{\text{UV}},\ell)}\, \partial_{\alpha_{\text{gf}}} \mathcal{S}_{k,a}^{(\lambda_{\text{UV}})}\Big) + \text{Tr}\Big((\partial_{\alpha_{\text{gf}}} \rho_0^{(\lambda_{\text{UV}},\ell)})\, \mathcal{S}_{k,a}^{(\lambda_{\text{UV}})}\Big) = \text{Tr}\Big(\rho_0^{(\lambda_{\text{UV}},\ell)}\, \zeta(\dots)\Big),$$

i.e. the variation is the expectation of a ζ-exact insertion (explicitly: the sum of $\zeta(\mathcal{N}_{\alpha_{\text{gf}}}[\mathcal{S}_{k,a}]^{(\lambda_{\text{UV}})})$ and $\zeta(\mathcal{N}_{\alpha_{\text{gf}}}[\rho_0]^{(\lambda_{\text{UV}},\ell)})$ reorganized by cyclicity of the (super)trace).

Cohomological vanishing on $\mathcal{H}_{\text{phys}}$. By Lemma 7.552, for any ghost-number-zero band state $\rho^{(\lambda_{\text{UV}},\ell)}$ with $\zeta(\rho^{(\lambda_{\text{UV}},\ell)}) = 0$,

$$\operatorname{Tr}\Big(\rho^{(\lambda_{\text{UV}},\ell)}\,\zeta(Y)\Big) = 0 \quad \text{on } \mathcal{H}_{\text{phys}}.$$

Therefore both the LHS and RHS variations vanish separately in the physical sector. We conclude $\partial_{\alpha_{\text{gf}}}\mathcal{W}_a^{(\lambda_{\text{UV}},\ell)} = 0$ on $\mathcal{H}_{\text{phys}}$.

Conclusion. Since $\mathcal{W}_a^{(\lambda_{\text{UV}},\ell)} = 0$ for some gauge choice implies $\partial_{\alpha_{\text{gf}}}\mathcal{W}_a^{(\lambda_{\text{UV}},\ell)} = 0$ everywhere, the identity (7.665) holds for all α_{gf} on $\mathcal{H}_{\text{phys}}$. Thus the band Ward identity is gauge-parameter independent. □

Theorem 7.570 (Non-vanishing Ward source in QIV)**.** *If at least one explicit breaking coupling is nonzero and relevant/marginal on the plateau band (QIV hypothesis), then on any certified plateau $\mathfrak{M}$ (cf. Definition 6.300)*

$$\sum_k \epsilon_k \left\langle \mathcal{S}_{k,a}^{(\lambda_{\text{UV}})} \right\rangle_{\boldsymbol{\Psi}} \neq 0 \quad \textit{for some broken generator } a,$$

and the scale-tagged divergence (7.665) does not vanish on $\mathfrak{M}$.

Proof (expanded). Write the band Ward source

$$\mathcal{R}_a^{(\lambda_{\text{UV}},\ell)} := \sum_k \epsilon_k \left\langle \mathcal{S}_{k,a}^{(\lambda_{\text{UV}})} \right\rangle_{\boldsymbol{\Psi}}, \tag{7.669}$$

so the scale-tagged Ward identity (Definition 7.567)

$$\nabla_\mu^{(\text{ad})}\,\bar{J}_{a,\text{imp}}^{(\lambda_{\text{UV}},\ell)\mu} = \mathcal{R}_a^{(\lambda_{\text{UV}},\ell)} \quad \text{on } \mathcal{H}_{\text{phys}}. \tag{7.670}$$

By Proposition 7.568, $\mathcal{R}_a^{(\lambda_{\text{UV}},\ell)}$ transforms in the adjoint; by Proposition 7.569 and Lemma 7.552, it is gauge-parameter independent on $\mathcal{H}_{\text{phys}}$. Throughout we work uniformly on a certified plateau mask $\mathfrak{M}$ (Definition 6.300).

Step I: Contradiction hypothesis and adjoint orbit. Assume, on $\mathfrak{M}$, that

$$\mathcal{R}_a^{(\lambda_{\text{UV}},\ell)} = 0 \quad \text{for all broken generators } a. \tag{7.671}$$

Adjoint covariance (Proposition 7.568) implies that (7.671) holds for the entire adjoint orbit generated from the broken subspace; in particular, it holds for every component connected by $f_{ba}{}^c$ to a broken direction.

Step II: Consequence—band conservation of the improved current. From (7.665) and (7.671) we obtain, for all such a,

$$\nabla_\mu^{(\text{ad})}\,\bar{J}_{a,\text{imp}}^{(\lambda_{\text{UV}},\ell)\mu} = 0 \quad \text{on } \mathcal{H}_{\text{phys}} \text{ and on } \mathfrak{M}. \tag{7.672}$$

Thus each broken-component current is (band-)conserved on the plateau in the physical sector.

Step III: Variation of the band effective action. Let $\Gamma^{(\lambda_{\mathrm{UV}},\ell)}$ denote the band effective action (generating functional of 1PI graphs) obtained with the taggers $P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}^{(\ell)}$; insert a spacetime-dependent test parameter $\alpha^a(x)$ coupled to the adjoint-covariant divergence. The operator Ward identity (eq. (6.293)) and band descent (Lemma 7.552 and Definition 7.567) give, to first order in α,

$$\delta_\alpha \Gamma^{(\lambda_{\mathrm{UV}},\ell)} = \int d^d x \; \alpha^a(x)\, \mathcal{R}_a^{(\lambda_{\mathrm{UV}},\ell)}(x), \tag{7.673}$$

up to BRST-exact insertions and total derivatives, which do not contribute on $\mathcal{H}_{\mathrm{phys}}$. By (7.671), we would have $\delta_\alpha \Gamma^{(\lambda_{\mathrm{UV}},\ell)} = 0$ for all α^a supported on the broken subspace. Hence the band effective action would be invariant (to first order) under those broken transformations.

Step IV: Relevance/marginality forbids trivialization. QIV assumes that at least one explicit breaking operator in the deformation $\sum_k \epsilon_k \mathcal{O}_k$ is *relevant or marginal* on the plateau band and has $\epsilon_k \neq 0$. In the algebraic–renormalization framework [117], a relevant/marginal deformation that is cohomologically trivial (BRST-exact plus total derivative, i.e. a redundant coupling) does not affect Γ at first order and can be removed by a field redefinition. Conversely, if $\delta_\alpha \Gamma^{(\lambda_{\mathrm{UV}},\ell)} = 0$ for all broken α^a, then the broken-direction part of the spurion deformation would be cohomologically trivial on the band, i.e.

$$\sum_k \epsilon_k \, \mathcal{S}_{k,a}^{(\lambda_{\mathrm{UV}})} \;=\; \nabla_\mu^{(\mathrm{ad})} X_a^\mu \;+\; \zeta(Y_a) \qquad \text{(in the band theory)}. \tag{7.674}$$

But (7.674) contradicts the QIV hypothesis: a *relevant/marginal* explicit breaking on $\mathfrak{M}$ is, by definition, a non-redundant deformation in the $H^0(\hat{Q}_{\mathrm{BRST}})$ cohomology class of the band theory; it cannot be written as an improvement plus a BRST-exact term without nullifying its physical effect on Γ. Therefore our assumption (7.671) is false.

Step V: Conclusion. There exists a broken generator a such that $\mathcal{R}_a^{(\lambda_{\mathrm{UV}},\ell)} \neq 0$ on $\mathfrak{M}$. By (7.665), the scale-tagged divergence $\nabla_\mu^{(\mathrm{ad})} \bar{J}_{a,\mathrm{imp}}^{(\lambda_{\mathrm{UV}},\ell)\mu}$ does not vanish on $\mathfrak{M}$. This establishes the claim. □

Corollary 7.571 (QIV vs. QI limit)**.** *Setting $\epsilon_k \to 0$ in (7.665) recovers band conservation, $\nabla_\mu^{(\mathrm{ad})} \bar{J}_{a,\mathrm{imp}}^{(\lambda_{\mathrm{UV}},\ell)\mu} = 0$, which is precisely the QI law-channel signature (Section 7.1).*

Proof. Start from the band Ward identity (Definition 7.567)

$$\nabla_\mu^{(\mathrm{ad})} \bar{J}_{a,\mathrm{imp}}^{(\lambda_{\mathrm{UV}},\ell)\mu} = \sum_k \epsilon_k \left\langle \mathcal{S}_{k,a}^{(\lambda_{\mathrm{UV}})} \right\rangle_{\boldsymbol{\Psi}} \quad \text{on } \mathcal{H}_{\mathrm{phys}}. \tag{7.675}$$

Fix a certified plateau mask $\mathfrak{M}$ (Definition 6.300). By Lemma 7.552 the band-tagged expectations on the right-hand side are well-defined and gauge-parameter independent (Proposition 7.569).

Consider the family of deformations parameterized by $\epsilon = \{\epsilon_k\}$. On $\mathfrak{M}$, plateau certification implies uniform bounds and smooth dependence of band correlators on small couplings;

hence the map $\epsilon \mapsto \langle \mathcal{S}_{k,a}^{(\lambda_{\mathrm{UV}})} \rangle_{\mathbf{\Psi}}$ is continuous at $\epsilon = 0$ (cf. standard locality/analyticity arguments in algebraic renormalization [117] and effective action dependence [91]). Therefore, taking the limit $\epsilon_k \to 0$ termwise in the finite (relevant/marginal) sum,

$$\lim_{\epsilon \to 0} \sum_k \epsilon_k \left\langle \mathcal{S}_{k,a}^{(\lambda_{\mathrm{UV}})} \right\rangle_{\mathbf{\Psi}} \;=\; 0 \qquad \text{uniformly on } \mathfrak{M}.$$

Passing to the limit in (7.675) yields

$$\nabla_\mu^{(\mathrm{ad})} \bar{J}_{a,\mathrm{imp}}^{(\lambda_{\mathrm{UV}},\ell)\mu} \;=\; 0 \quad \text{on } \mathcal{H}_{\mathrm{phys}} \text{ and on } \mathfrak{M}.$$

This is precisely the band conservation statement that characterizes the QI law channel (Section 7.1). The result is independent of gauge-fixing parameters (by Proposition 7.569) and stable under BRST- and Noether-improvements, which do not affect the conserved charge on $\mathcal{H}_{\mathrm{phys}}$ (Lemma 7.552). □

Intuition. This subsection pins down the symmetry frame and shows how the explicit breakings enter the *band* Ward identity in a fully covariant, BRST- and gauge-safe way. In QIV the right-hand side is nonzero on plateaus (relevant/marginal spurions), and this is exactly the seed that drives the non-healing law-visibility deficit in Theorem 7.557.

7.9.3 Cone–preserving Kernels

Goal. Instantiate the UV/IR taggers (Definition 7.551) with an explicit causal kernel family that (i) preserves microcausality (support inside $\mathsf{C}^+ \cup \mathsf{C}^-$), (ii) commutes with BRST/G, and (iii) is compatible with the spectral/tomographic constructions of Definition 7.607 and Proposition 7.609.

Definition 7.572 (Causal IR kernel family)**.** Let $v_* > 0$ be the microscopic signal speed (e.g. $v_* = c$ relativistically, or a Lieb–Robinson velocity on the lattice). For each band point $x = (\lambda_{\mathrm{UV}}, \ell)$, define the IR kernel

$$\kappa_x(t, \mathbf{r}) \;:=\; \chi_\ell(t)\, \phi_\ell\!\left(\frac{\|\mathbf{r}\|}{v_* |t| + \delta} \right),$$

with: (i) $\chi_\ell \in C_c^\infty(\mathbb{R})$, even, supported in $|t| \le \tau(\ell)$, normalized $\int \chi_\ell(t)\, dt = 1$; (ii) $\phi_\ell \in C_c^\infty([0,\infty))$, supported in $[0,1]$, nonnegative, $\phi_\ell(0) = 1$; (iii) $\delta > 0$ a small regulator (removed at the end). The associated convolution tagger on operators is $(K_{\mathrm{IR}}^{(\ell)} \mathcal{O})(x) := \int dt\, d^d\mathbf{r}\; \kappa_x(t,\mathbf{r})\; \mathcal{O}(x^0 - t, \mathbf{x} - \mathbf{r})$, and similarly for states by dual action. Take the UV projector $P_x(\lambda_{\mathrm{UV}})$ as any spectral-cutoff operator commuting with spacetime translations (e.g. a Fourier-multiplier with radial mask).

Proposition 7.573 (Cone preservation and commutation)**.** *The family $\{\kappa_x\}$ of Definition 7.572 is cone–preserving:* $\operatorname{supp} \kappa_x \subset \mathsf{C}^+ \cup \mathsf{C}^-$ *for every x. Moreover $K_{\mathrm{IR}}^{(\ell)}$ and $P_x(\lambda_{\mathrm{UV}})$ commute with BRST and G actions:* $[\zeta, K_{\mathrm{IR}}^{(\ell)}] = [\zeta, P_x] = [\delta_a, K_{\mathrm{IR}}^{(\ell)}] = [\delta_a, P_x] = 0$.

Proof. (A) Cone preservation. By Definition 7.572, each kernel is constructed with a finite time aperture $\tau(\ell)$ and a finite propagation speed bound $v_*(x)$ (from strong hyperbolicity and the principal symbol; cf. Definition 7.535), together with a harmless thickness $\delta \geq 0$. Concretely,

$$\kappa_x(t,\mathbf{r}) = 0 \quad \text{unless} \quad |t| \leq \tau(\ell) \ \text{ and } \ \|\mathbf{r}\| \leq v_*(x)\,|t| + \delta.$$

Hence $\operatorname{supp}\kappa_x \subset \mathsf{C}^+ \cup \mathsf{C}^-$ by definition of the thickened forward/backward cones $\mathsf{C}^\pm$ at x. This establishes cone preservation.

(B) Representation of the taggers. Both taggers are linear, continuous maps on operator–valued distributions obtained by c–number kernels/multipliers:

$$(K_{\mathrm{IR}}^{(\ell)}\mathcal{O})(y) = \int d^d z\ \kappa_x(y-z)\,\mathcal{O}(z), \tag{7.676}$$

$$(P_x(\lambda_{\mathrm{UV}})\mathcal{O})\widehat{\ }(p) = \chi_x(\lambda_{\mathrm{UV}},p)\ \widehat{\mathcal{O}}(p), \tag{7.677}$$

where κ_x is as above and χ_x is a smooth, compactly supported (or rapidly decaying) UV window around $|p| \lesssim \lambda_{\mathrm{UV}}$ fixed at x. Equivalently, P_x can be written as a convolution with a c–number kernel p_x obtained by inverse Fourier transform of χ_x.

(C) Commutation with BRST. Let ζ be the BRST differential acting as a graded derivation on the field/operator algebra, with $[\zeta,\partial_\mu]=0$ and ζ annihilating c–number functions. Using (7.676) and linearity,

$$\zeta\Big(K_{\mathrm{IR}}^{(\ell)}\mathcal{O}\Big)(y) = \zeta\Big(\int d^d z\ \kappa_x(y-z)\,\mathcal{O}(z)\Big) = \int d^d z\ \kappa_x(y-z)\,\zeta\mathcal{O}(z) = (K_{\mathrm{IR}}^{(\ell)}\,\zeta\mathcal{O})(y),$$

so $[\zeta, K_{\mathrm{IR}}^{(\ell)}]=0$. In momentum space,

$$\zeta(P_x\mathcal{O})\widehat{\ }(p) = \zeta\Big(\chi_x(\lambda_{\mathrm{UV}},p)\,\widehat{\mathcal{O}}(p)\Big) = \chi_x(\lambda_{\mathrm{UV}},p)\,\zeta\widehat{\mathcal{O}}(p) = (P_x\,\zeta\mathcal{O})\widehat{\ }(p),$$

hence $[\zeta,P_x]=0$. No domain subtleties arise on certified plateaus since both taggers are bounded on the Banach spaces used there, and ζ is closable on the same domain (standard BRST functional setting).

(D) Commutation with the G–action. Let δ_a denote the infinitesimal G–action (global/internal), acting fibrewise on field indices and as a derivation on products; it commutes with spacetime translations/derivatives and does not act on c–number kernels. Then, using (7.676),

$$\delta_a\Big(K_{\mathrm{IR}}^{(\ell)}\mathcal{O}\Big)(y) = \int d^d z\ \kappa_x(y-z)\,\delta_a\mathcal{O}(z) = (K_{\mathrm{IR}}^{(\ell)}\,\delta_a\mathcal{O})(y),$$

so $[\delta_a, K_{\mathrm{IR}}^{(\ell)}]=0$. Likewise from (7.677),

$$\delta_a(P_x\mathcal{O})\widehat{\ }(p) = \chi_x(\lambda_{\mathrm{UV}},p)\,\delta_a\widehat{\mathcal{O}}(p) = (P_x\,\delta_a\mathcal{O})\widehat{\ }(p),$$

so $[\delta_a,P_x]=0$. If one realises δ_a as an inner derivation $\delta_a(\cdot)=i[Q_a,\cdot]$ with Q_a the conserved charge, the same conclusion follows since $K_{\mathrm{IR}}^{(\ell)}$ and P_x are c–number maps on spacetime arguments and hence commute with taking commutators in field space: $[Q_a, K_{\mathrm{IR}}^{(\ell)}\mathcal{O}] = K_{\mathrm{IR}}^{(\ell)}[Q_a,\mathcal{O}]$, and similarly for P_x.

Combining (A)–(D) gives the stated cone support for κ_x and the commutators $[\zeta, K_{\mathrm{IR}}^{(\ell)}] = [\zeta,P_x] = [\delta_a, K_{\mathrm{IR}}^{(\ell)}] = [\delta_a,P_x] = 0$, completing the proof. □

Corollary 7.574 (Non-signalling after band descent)**.** *If the microscopic theory is microcausal, then for spacelike–separated regions A, B, $\langle[\,\bar{J}^{(x)\mu}_{a,\mathrm{imp}}(A),\,\mathcal{W}^{(x)}(B)\,]\rangle_{\mathbf{\Psi}} = 0$ for all certified $x \in \mathfrak{P}$ (Definition 6.300).*

Proof. Setup. Let $C^{\mu}_{a}(y,z) := \Big\langle\,[\,\bar{J}^{\mu}_{a,\mathrm{imp}}(y),\,\mathcal{W}(z)\,]\,\Big\rangle_{\mathbf{\Psi}}$ denote the microscopic commutator bi–distribution. Microcausality means

$$\operatorname{supp} C^{\mu}_{a} \subset \{(y,z) \in \mathbb{R}^{d} \times \mathbb{R}^{d} :\ (y-z) \in \mathsf{C}^{+} \cup \mathsf{C}^{-}\}.$$

Band descent acts as the c–number maps $K^{(\ell)}_{\mathrm{IR}}$ and $P_x(\lambda_{\mathrm{UV}})$ (Definition 7.572) on operator–valued distributions (convolution in position or multiplication in momentum, cf. Proposition 7.573). If f_A, g_B are test functions supported in A, B respectively, then

$$\bar{J}^{(x)\mu}_{a,\mathrm{imp}}(A) = \bar{J}^{\mu}_{a,\mathrm{imp}}\Big((K^{(\ell)}_{\mathrm{IR}}P_x)^{\dagger} f_A\Big), \qquad \mathcal{W}^{(x)}(B) = \mathcal{W}\Big((K^{(\ell)}_{\mathrm{IR}}P_x)^{\dagger} g_B\Big),$$

where $(\cdot)^{\dagger}$ is the transpose action on test functions. Writing κ_x for the IR kernel and χ_x for the UV window, one has $(K^{(\ell)}_{\mathrm{IR}})^{\dagger} f_A = f_A * \tilde{\kappa}_x$ with $\tilde{\kappa}_x(t,\mathbf{r}) = \kappa_x(-t,-\mathbf{r})$, and $(P_x)^{\dagger}$ acts by multiplication with χ_x in momentum space.

Step 1: Cone preservation under band descent. By Proposition 7.573, $\operatorname{supp}\kappa_x \subset \mathsf{C}^{+} \cup \mathsf{C}^{-}$ and the taggers commute with internal actions. Hence the *effective* test functions after descent satisfy

$$\operatorname{supp}\Big((K^{(\ell)}_{\mathrm{IR}}P_x)^{\dagger} f_A\Big) \subset A + \operatorname{supp}\kappa_x, \qquad \operatorname{supp}\Big((K^{(\ell)}_{\mathrm{IR}}P_x)^{\dagger} g_B\Big) \subset B + \operatorname{supp}\kappa_x,$$

where $S + T := \{s + t :\ s \in S,\ t \in T\}$ is the Minkowski sum.

Step 2: Plateau non–overlap margin. By certification (Definition 6.300) one chooses the IR aperture $\tau(\ell)$ and speed bound $v_*(x)$ (from strong hyperbolicity, Definition 7.535) so that the *thickened supports*

$$A^{\sharp} := A + \operatorname{supp}\kappa_x, \qquad B^{\sharp} := B + \operatorname{supp}\kappa_x$$

remain spacelike–separated: $(y-z) \notin \mathsf{C}^{+} \cup \mathsf{C}^{-}$ for all $y \in A^{\sharp}$, $z \in B^{\sharp}$. This is exactly the "non–signalling margin" encoded in the plateau acceptance (finite time aperture $\tau(\ell)$ and cone thickness δ chosen below the region–separation buffer).

Step 3: Vanishing of the descended commutator. Using the distributional pairing and Fubini,

$$\begin{aligned}\Big\langle\,[\,\bar{J}^{(x)\mu}_{a,\mathrm{imp}}(A),\,\mathcal{W}^{(x)}(B)\,]\,\Big\rangle_{\mathbf{\Psi}} &= \Big\langle\, C^{\mu}_{a},\ (K^{(\ell)}_{\mathrm{IR}}P_x)^{\dagger} f_A \ \otimes\ (K^{(\ell)}_{\mathrm{IR}}P_x)^{\dagger} g_B \,\Big\rangle \\ &= \int dy\, dz\ C^{\mu}_{a}(y,z)\ \Big((K^{(\ell)}_{\mathrm{IR}}P_x)^{\dagger} f_A\Big)(y)\ \Big((K^{(\ell)}_{\mathrm{IR}}P_x)^{\dagger} g_B\Big)(z).\end{aligned}$$

The integrand has support inside

$$\{(y,z) :\ (y-z) \in \mathsf{C}^{+} \cup \mathsf{C}^{-}\}\ \cap\ \Big(A^{\sharp} \times B^{\sharp}\Big),$$

which is empty by the plateau non–overlap from Step 2. Therefore the integral vanishes: $\langle[\,\bar{J}^{(x)\mu}_{a,\mathrm{imp}}(A),\,\mathcal{W}^{(x)}(B)\,]\rangle_{\mathbf{\Psi}} = 0$.

Remarks. (i) The replacement of ∂_μ by the adjoint covariant divergence in $\bar{J}^{(x)\mu}_{a,\mathrm{imp}}$ does not affect support since the connection is a c–number background on spacetime indices. (ii) Commutation $[\zeta, K^{(\ell)}_{\mathrm{IR}}] = [\delta_a, K^{(\ell)}_{\mathrm{IR}}] = [\zeta, P_x] = [\delta_a, P_x] = 0$ (Proposition 7.573) ensures the BRST/G structure is unchanged by descent and does not obstruct microcausality.

This establishes non–signalling of the descended observables for all certified $x \in \mathfrak{P}$. □

Lemma 7.575 (Frequency response and kernel floor)**.** *Let $\widehat{\kappa}_x(\omega, \mathbf{k})$ be the Fourier transform of κ_x. Then $\widehat{\kappa}_x$ is C^∞ in $(\omega, \mathbf{k})$, real–valued and even in ω. Moreover, for any $\omega_0 > 0$ sufficiently small there exists $\kappa_{\mathrm{spec}}(x) > 0$ such that*

$$\mathsf{K}_{\mathrm{spec}}(x; \omega) \;\geq\; \kappa_{\mathrm{spec}}(x) \qquad \text{for } |\omega| \leq \omega_0,$$

with $\mathsf{K}_{\mathrm{spec}}$ the tomography kernel of Definition 7.607.

Proof. Structure of κ_x and smoothness. By Definition 7.572, the IR tagger uses a causal, compactly supported C^∞ kernel which, in local coordinates at x, factors as

$$\kappa_x(t, \mathbf{r}) \;=\; \chi_\ell(t)\, \phi_\ell(\mathbf{r}),$$

with $\chi_\ell \in C^\infty_c(\mathbb{R})$ an even time window ($\chi_\ell(t) = \chi_\ell(-t)$, real, nonnegative) and $\phi_\ell \in C^\infty_c(\mathbb{R}^{d-1})$ a real, radially symmetric spatial window. Hence

$$\widehat{\kappa}_x(\omega, \mathbf{k}) \;=\; \widehat{\chi}_\ell(\omega)\, \widehat{\phi}_\ell(\mathbf{k}),$$

and both factors are entire (Paley–Wiener), thus C^∞ in all variables. Because χ_ℓ is real and even, $\widehat{\chi}_\ell$ is real and even in ω; because ϕ_ℓ is real and even (radial), $\widehat{\phi}_\ell$ is real and even in $\mathbf{k}$. Therefore $\widehat{\kappa}_x$ is smooth and real–even in ω.

Positivity at zero frequency. Since $\chi_\ell \geq 0$ and not identically zero, $\widehat{\chi}_\ell(0) = \int_{\mathbb{R}} \chi_\ell(t)\, dt > 0$. By construction, the tomography kernel of Definition 7.607 is a positive quadratic functional of $\widehat{\kappa}_x$ of the form

$$\mathsf{K}_{\mathrm{spec}}(x; \omega) \;=\; \int_{\mathbb{R}^{d-1}} W_x(\mathbf{k}) \left|\widehat{\kappa}_x(\omega, \mathbf{k})\right|^2 d\mathbf{k} \;=\; \left|\widehat{\chi}_\ell(\omega)\right|^2 \int_{\mathbb{R}^{d-1}} W_x(\mathbf{k}) \left|\widehat{\phi}_\ell(\mathbf{k})\right|^2 d\mathbf{k}, \tag{7.678}$$

where $W_x(\mathbf{k}) \geq 0$ is the channel weight specified in Definition 7.607 (it encodes the T–odd projection and normalization). By design, W_x is not a.e. zero and has positive mass on some compact set $A_x \subset \mathbb{R}^{d-1}$; since $\widehat{\phi}_\ell$ is continuous and not identically zero, there exists a compact A_x with

$$c_x \;:=\; \int_{A_x} W_x(\mathbf{k}) \left|\widehat{\phi}_\ell(\mathbf{k})\right|^2 d\mathbf{k} \;>\; 0.$$

Therefore

$$\mathsf{K}_{\mathrm{spec}}(x; 0) \;=\; \left|\widehat{\chi}_\ell(0)\right|^2 \int_{\mathbb{R}^{d-1}} W_x(\mathbf{k}) \left|\widehat{\phi}_\ell(\mathbf{k})\right|^2 d\mathbf{k} \;\geq\; \left|\widehat{\chi}_\ell(0)\right|^2 c_x \;>\; 0.$$

Continuity and a uniform floor near $\omega = 0$. The map $\omega \mapsto \mathsf{K}_{\mathrm{spec}}(x; \omega)$ is continuous: indeed, $\widehat{\chi}_\ell$ is continuous and $W_x(\mathbf{k})|\widehat{\phi}_\ell(\mathbf{k})|^2$ is integrable on $\mathbb{R}^{d-1}$, so (7.678) is a continuous composition and dominated-convergence applies. Hence there exists $\omega_0 > 0$ such that

$$\left|\widehat{\chi}_\ell(\omega)\right|^2 \;\geq\; \tfrac{1}{2} \left|\widehat{\chi}_\ell(0)\right|^2 \qquad \text{for all } |\omega| \leq \omega_0.$$

Combining with (7.678) and the definition of c_x,

$$\mathsf{K}_{\mathrm{spec}}(x;\omega) \;\geq\; \tfrac{1}{2}\left|\widehat{\chi}_\ell(0)\right|^2 c_x \;=:\; \kappa_{\mathrm{spec}}(x) \;>\; 0 \qquad \text{for all } |\omega| \leq \omega_0.$$

Remarks on invariances. Because κ_x is a c–number kernel and the taggers commute with BRST/G (Proposition 7.573), the bound is taken on the same physical sector $\mathcal{H}_{\mathrm{phys}}$ and is unaffected by gauge-parameter choices (Nielsen robustness).

This proves the existence of a strictly positive low–frequency kernel floor $\kappa_{\mathrm{spec}}(x)$. □

Definition 7.576 (One–time numerical verification (accept/reject))**.** On the working spacetime grid, precompute κ_x and declare a tolerance $\epsilon_{\mathrm{cone}} = 10^{-12}$ (floating–point floor). Accept the kernel family if

$$\max_{(t,\mathbf{r})\notin \mathsf{C}^{\pm}} \; |\kappa_x(t,\mathbf{r})| \;\leq\; \epsilon_{\mathrm{cone}} \quad \text{for all } x \in \mathfrak{P},$$

and if the discrete Fourier sample of $\mathsf{K}_{\mathrm{spec}}(x;\omega)$ satisfies the kernel–floor inequality of Lemma 7.575 with a discretization slack δ_{disc} as in Proposition 7.602. Re-run only when $\tau(\ell)$ or the spatial stencil changes.

Notes. (i) The time–only choice $\phi_\ell \equiv \delta(\mathbf{r})$ is trivially cone–preserving and may be used for baseline audits. (ii) For lattice models, replace v_* by a compatible Lieb–Robinson velocity; proofs are unchanged. (iii) All statements are BRST/Nielsen safe by Lemma 7.552 and Proposition 7.556.

7.9.4 Visibilities (Primary Diagnostics)

Definition 7.577 (Plateau-wise primary diagnostics)**.** Let $\mathfrak{P} \subset \{(\lambda_{\mathrm{UV}}, \ell)\}$ be a certified plateau mask in the sense of Definition 6.300. On $\mathfrak{P}$ we define:

1. *Persistent law-deficit:* there exists $\delta_\star \in (0,1)$ such that
$$\left|\Lambda_{\mathrm{law}}^{(\lambda_{\mathrm{UV}},\ell)}\right| \;\leq\; 1-\delta_\star \qquad \forall\,(\lambda_{\mathrm{UV}},\ell) \in \mathfrak{P}.$$
2. *Unique-vacuum state channel:* for any volume sequence $V \to \infty$,
$$\lim_{V\to\infty} \left|\Lambda_{\mathrm{state}}^{(\lambda_{\mathrm{UV}},\ell)}(V)\right| \;=\; 1 \qquad \text{uniformly on } \mathfrak{P}.$$

Here $\Lambda_{\mathrm{law}}, \Lambda_{\mathrm{state}}$ are the holosymmetric visibilities with UV/IR tags as in Definitions 6.275 and 7.551, and expectations are taken on $\mathcal{H}_{\mathrm{phys}}$ (BRST/Nielsen-safe by Lemma 7.552 and Proposition 7.556).

Lemma 7.578 (From non-vanishing Ward source to a positive law gap at a point)**.** *Fix $(\lambda_{\mathrm{UV}},\ell) \in \mathfrak{P}$. If the band Ward source*

$$\mathcal{R}_a^{(\lambda_{\mathrm{UV}},\ell)} := \sum_k \epsilon_k \left\langle \mathcal{S}_{k,a}^{(\lambda_{\mathrm{UV}})} \right\rangle_{\mathbf{\Psi}}$$

is nonzero for some broken generator a (cf. Theorem 7.570), then there exists $c_\bullet(\{\epsilon_k\},\lambda_{\mathrm{UV}},\ell) > 0$ such that

$$1 - \left|\Lambda_{\mathrm{law}}^{(\lambda_{\mathrm{UV}},\ell)}\right| \;\geq\; c_\bullet(\{\epsilon_k\},\lambda_{\mathrm{UV}},\ell).$$

Proof (expanded). **1) Channel decomposition and the spectral (T-odd) branch.** By the holosymmetric construction, the law visibility at the band point $(\lambda_{\mathrm{UV}}, \ell)$ is the tagged overlap of branch unitaries traced against the tagged state, $\Lambda_{\mathrm{law}}^{(\lambda_{\mathrm{UV}},\ell)} = \mathrm{Tr}\big[(U_B^{(\lambda_{\mathrm{UV}},\ell)})^\dagger U_A^{(\lambda_{\mathrm{UV}},\ell)}\, \rho_0^{(\lambda_{\mathrm{UV}},\ell)}\big]$ (Definitions 6.275 and 7.551). Insert the triadic channel projectors and use Proposition 7.555 to write

$$\Lambda_{\mathrm{law}}^{(\lambda_{\mathrm{UV}},\ell)} = \Lambda_{\mathrm{law}}^{(\mathrm{vac};\lambda_{\mathrm{UV}},\ell)} + \Lambda_{\mathrm{law}}^{(\mathrm{corr};\lambda_{\mathrm{UV}},\ell)} + \Lambda_{\mathrm{law}}^{(\mathrm{spec};\lambda_{\mathrm{UV}},\ell)}.$$

By construction (holosymmetric/T-parity split), the *spectral* branch is T-odd and admits a Kubo-type spectral representation

$$-\,\mathrm{Re}\;\Lambda_{\mathrm{law}}^{(\mathrm{spec};\lambda_{\mathrm{UV}},\ell)} = \int_{\mathbb{R}} d\omega\; \mathsf{K}_{\mathrm{spec}}^{(\lambda_{\mathrm{UV}},\ell)}(\omega)\, \mathsf{S}_a^{(\lambda_{\mathrm{UV}},\ell)}(\omega), \tag{7.679}$$

where $\mathsf{K}_{\mathrm{spec}}^{(\lambda_{\mathrm{UV}},\ell)}(\omega) \geq 0$ is a positive kernel determined by the UV/IR taggers (commuting with BRST and G by Definition 7.551) and $\mathsf{S}_a^{(\lambda_{\mathrm{UV}},\ell)}(\omega)$ is the spectral density associated with the adjoint-covariant divergence of the improved current along a broken direction a (defined cohomologically on $\mathcal{H}_{\mathrm{phys}}$ by Lemma 7.552).[17]

2) Injecting the Ward identity. On $\mathcal{H}_{\mathrm{phys}}$, the scale-tagged Ward identity reads

$$\nabla_\mu^{(\mathrm{ad})}\, \bar{J}_{a,\mathrm{imp}}^{(\lambda_{\mathrm{UV}},\ell)\mu} = \mathcal{R}_a^{(\lambda_{\mathrm{UV}},\ell)} \qquad \text{(Definition 7.567)}. \tag{7.680}$$

A nonzero $\mathcal{R}_a^{(\lambda_{\mathrm{UV}},\ell)}$ (for some broken a) implies that the T-odd two-point function of the divergence has a non-vanishing zero-frequency component: in the spectral language, $\mathsf{S}_a^{(\lambda_{\mathrm{UV}},\ell)}(\omega)$ carries a strictly positive weight in any neighborhood of $\omega = 0$.This uses adjoint covariance (Proposition 7.568) and the fact that BRST-exact/improvement pieces do not contribute on $\mathcal{H}_{\mathrm{phys}}$ (Lemma 7.552).

3) Positivity of the kernel on a certified band point. By the certification of $(\lambda_{\mathrm{UV}}, \ell) \in \mathfrak{P}$ (Definition 6.300), the UV/IR taggers fix the bandwidth and coarse-graining so that the kernel $\mathsf{K}_{\mathrm{spec}}^{(\lambda_{\mathrm{UV}},\ell)}(\omega)$ is continuous and admits a strictly positive lower bound on a compact neighborhood of $\omega = 0$:

$$\exists\, \omega_0 > 0,\ \kappa_{\mathrm{spec}}(\lambda_{\mathrm{UV}}, \ell) > 0 \quad \text{s.t.} \quad \mathsf{K}_{\mathrm{spec}}^{(\lambda_{\mathrm{UV}},\ell)}(\omega) \;\geq\; \kappa_{\mathrm{spec}}(\lambda_{\mathrm{UV}}, \ell) \quad \forall\, \omega \in [-\omega_0, \omega_0].$$

(This follows from the scale-aware setup of Section 6.12 and the commutation/regularity properties of Definition 7.551.)

4) Lower-bounding the modulus deficit. Since $|\Lambda_{\mathrm{law}}| \leq \mathrm{Re}\;\Lambda_{\mathrm{law}}^{(\mathrm{vac})} + \mathrm{Re}\;\Lambda_{\mathrm{law}}^{(\mathrm{corr})} + \mathrm{Re}\;\Lambda_{\mathrm{law}}^{(\mathrm{spec})}$ and the first two branches are T-even (do not contribute negatively to the real part at unity normalization), we have

$$1 - \left|\Lambda_{\mathrm{law}}^{(\lambda_{\mathrm{UV}},\ell)}\right| \;\geq\; -\,\mathrm{Re}\;\Lambda_{\mathrm{law}}^{(\mathrm{spec};\lambda_{\mathrm{UV}},\ell)}.$$

Combining with (7.679) and the kernel bound gives

$$1 - \left|\Lambda_{\mathrm{law}}^{(\lambda_{\mathrm{UV}},\ell)}\right| \;\geq\; \kappa_{\mathrm{spec}}(\lambda_{\mathrm{UV}}, \ell) \int_{-\omega_0}^{\omega_0} d\omega\; \mathsf{S}_a^{(\lambda_{\mathrm{UV}},\ell)}(\omega).$$

[17] Any broken direction suffices for the lower bound below; we suppress the index a when clear.

By Step 2, the integral on the right is strictly positive whenever $\mathcal{R}_a^{(\lambda_{\mathrm{UV}},\ell)} \neq 0$; denote it by $\sigma_a^{(\lambda_{\mathrm{UV}},\ell)} > 0$, a continuous function of the couplings $\{\epsilon_k\}$ at fixed band point (gauge-parameter independent by Proposition 7.569). Therefore

$$1 - \left|\Lambda_{\mathrm{law}}^{(\lambda_{\mathrm{UV}},\ell)}\right| \;\geq\; \kappa_{\mathrm{spec}}(\lambda_{\mathrm{UV}},\ell)\,\sigma_a^{(\lambda_{\mathrm{UV}},\ell)} \;:=\; c_\bullet(\{\epsilon_k\},\lambda_{\mathrm{UV}},\ell) \;>\; 0.$$

5) Cohomological and gauge independence. All steps above are performed on $\mathcal{H}_{\mathrm{phys}}$; BRST-exact and Noether-improvement pieces drop out (Lemma 7.552), and the bound is independent of gauge-fixing parameters (Proposition 7.569). Adjoint covariance ensures that choosing any broken generator a with $\mathcal{R}_a^{(\lambda_{\mathrm{UV}},\ell)} \neq 0$ yields the same conclusion (Proposition 7.568).

This proves the claim with the stated, strictly positive constant $c_\bullet(\{\epsilon_k\},\lambda_{\mathrm{UV}},\ell)$. □

Lemma 7.579 (Plateau uniformization)**.** *If a function $f(\lambda_{\mathrm{UV}},\ell)$ obeys $|f(x)-f(y)| \leq \eta_{\mathfrak{P}}$ for all $x,y \in \mathfrak{P}$ (slope/curvature/oscillation bounds of Definition 6.300), and $f(x_0) \geq c_0 > 0$ for some $x_0 \in \mathfrak{P}$, then*

$$\inf_{x\in\mathfrak{P}} f(x) \;\geq\; c_0 - \eta_{\mathfrak{P}}.$$

Proof (expanded). By the plateau certification criteria (Definition 6.300, within the scale-aware scheme of Section 6.12), the mask $\mathfrak{P}$ enforces a uniform oscillation bound

$$\mathrm{osc}_{\mathfrak{P}}(f) := \sup_{x,y\in\mathfrak{P}} |f(x)-f(y)| \;\leq\; \eta_{\mathfrak{P}}, \tag{7.681}$$

where $\eta_{\mathfrak{P}} \geq 0$ is the certified tolerance determined by the slope/curvature tests on the $(\lambda_{\mathrm{UV}},\ell)$-grid.

Fix any $x \in \mathfrak{P}$. Apply the triangle inequality with the reference point $x_0 \in \mathfrak{P}$:

$$f(x) \;=\; f(x_0) \;-\; \big(f(x_0)-f(x)\big) \;\geq\; f(x_0) \;-\; |f(x_0)-f(x)| \;\geq\; c_0 \;-\; \mathrm{osc}_{\mathfrak{P}}(f).$$

Using (7.681) gives

$$f(x) \;\geq\; c_0 - \eta_{\mathfrak{P}} \qquad \forall\, x \in \mathfrak{P}.$$

Taking the infimum over $x \in \mathfrak{P}$ proves the claim: $\inf_{x\in\mathfrak{P}} f(x) \geq c_0 - \eta_{\mathfrak{P}}$.

Remarks. (i) If f is a BRST/Nielsen-safe observable built from band visibilities (e.g. $f = 1 - |\Lambda_{\mathrm{law}}|$), the bound is gauge-parameter independent by Proposition 7.556, and the band descent used to define f is valid on $\mathcal{H}_{\mathrm{phys}}$ by Lemma 7.552. (ii) When $c_0 > \eta_{\mathfrak{P}}$ the lemma guarantees a strictly positive uniform lower bound on $\mathfrak{P}$. □

Theorem 7.580 (Primary QIV visibility diagnostics on certified plateaus)**.** *Assume the QIV hypotheses (nonzero relevant/marginal explicit breakings on the band) and work on a certified plateau $\mathfrak{P}$. Then:*

1. *(*Persistent law-deficit*) There exists $\delta_\star \in (0,1)$ such that*

$$\left|\Lambda_{\mathrm{law}}^{(\lambda_{\mathrm{UV}},\ell)}\right| \;\leq\; 1-\delta_\star \qquad \forall\,(\lambda_{\mathrm{UV}},\ell) \in \mathfrak{P}.$$

*2. (*Unique-vacuum state channel*) Uniformly on* $\mathfrak{P}$*,*

$$\lim_{V\to\infty} \left|\Lambda_{\text{state}}^{(\lambda_{\text{UV}},\ell)}(V)\right| \;=\; 1.$$

Proof (expanded). We recall (i) the holosymmetric visibilities with UV/IR tags (Definitions 6.275 and 7.551), (ii) the band Ward identity (Definition 7.567), (iii) adjoint covariance of the band Ward identity (Proposition 7.568), (iv) gauge-parameter independence on $\mathcal{H}_{\text{phys}}$ (Proposition 7.569), and (v) the descent and cohomological properties on $\mathcal{H}_{\text{phys}}$ (Lemma 7.552).

(1) Law channel: existence of a uniform deficit on $\mathfrak{P}$. *Step 1a (Pointwise positive gap).* By Theorem 7.570, on any certified plateau $\mathfrak{P}$ there exists a broken generator a and a band point $x_0 = (\lambda_{\text{UV}}, \ell) \in \mathfrak{P}$ such that the band Ward source $\mathcal{R}_a^{(\lambda_{\text{UV}},\ell)} = \sum_k \epsilon_k \langle \mathcal{S}_{k,a}^{(\lambda_{\text{UV}})} \rangle_{\Psi}$ is nonzero. Then Lemma 7.578 yields a *pointwise* strictly positive modulus deficit,

$$1 - \left|\Lambda_{\text{law}}^{(\lambda_{\text{UV}},\ell)}\right| \;\geq\; c_\bullet(\{\epsilon_k\}, \lambda_{\text{UV}}, \ell) \;:=\; c_0 \;>0 \qquad \text{at } x_0 \in \mathfrak{P},$$

with c_0 depending continuously on the explicit couplings and the band point, and independent of the gauge-fixing parameters by Proposition 7.569.

Step 1b (Uniformization over the plateau). By certification (Definition 6.300 in the scale-aware scheme of Section 6.12), the law visibility varies slowly on $\mathfrak{P}$:

$$\sup_{x,y\in\mathfrak{P}} \left|\Lambda_{\text{law}}(x) - \Lambda_{\text{law}}(y)\right| \;\leq\; \eta_{\mathfrak{P}}.$$

Applying Lemma 7.579 to $f(x) := 1 - |\Lambda_{\text{law}}(x)|$ with the single-point lower bound $f(x_0) \geq c_0$ gives

$$\inf_{x\in\mathfrak{P}} \left(1 - |\Lambda_{\text{law}}(x)|\right) \;\geq\; c_0 - \eta_{\mathfrak{P}}.$$

Since $\eta_{\mathfrak{P}}$ is a certification tolerance, we may (if needed) tighten the mask $\mathfrak{P}$ so that $\eta_{\mathfrak{P}} < \frac{1}{2}c_0$. Define

$$\delta_\star \;:=\; \tfrac{1}{2}c_0 \;\in (0,1).$$

Then for all $x \in \mathfrak{P}$,

$$1 - \left|\Lambda_{\text{law}}(x)\right| \;\geq\; \delta_\star \quad\Longleftrightarrow\quad \left|\Lambda_{\text{law}}(x)\right| \;\leq\; 1 - \delta_\star.$$

This proves the *persistent law-deficit* uniformly on $\mathfrak{P}$. Gauge-parameter independence follows from Proposition 7.569, and insensitivity to BRST/improvement trivialities from Lemma 7.552.

(2) State channel: unique vacuum $\Rightarrow |\Lambda_{\text{state}}| \to 1$ uniformly on $\mathfrak{P}$. *Step 2a (Uniqueness and mass gap on the band).* In QIV, explicit breakings select a unique vacuum (no vacuum manifold), so there are *no exact NG modes*; the would-be NG excitations are pseudo-NG with GMOR-type masses controlled by the spurions (see the Ward→GMOR discussion in Section 7.9 and the standard references [94, 91, 112]). On the band, the mass/mixing ledger built from the scale-tagged Hessian (cf. the "Mass-matrix & mixing ledger" step in Section 7.9) shows that all would-be NG directions acquire positive masses $m_\pi^2(\lambda_{\text{UV}}, \ell) > 0$ for $\epsilon_k \neq 0$.

Because $\mathfrak{P}$ is a finite (or compact) mask in the discretized $(\lambda_{\mathrm{UV}}, \ell)$-grid and the ledger depends continuously on parameters, there exists a *uniform* lower bound

$$m_{\min}^2 \;:=\; \inf_{x\in\mathfrak{P}} \; m_\pi^2(x) \;>\; 0. \tag{7.682}$$

Hence the correlation length is uniformly bounded: $\xi_{\max} := \sup_{x\in\mathfrak{P}} \xi(x) = \sup_x m_\pi(x)^{-1} < \infty$.

Step 2b (Exponential clustering and branch preparation). Let $|\Omega_1(x)\rangle\,, |\Omega_2(x)\rangle$ be the two branch-prepared states used in the visibility definition (Definition 6.275), constructed by admissible boundary conditions that differ only in a boundary layer. For a gapped, local theory with correlation length $\xi(x)$, the cluster property implies exponential decay of connected correlators on scales $\gtrsim \xi(x)$ (see, e.g., [91] for continuum arguments). Therefore boundary-induced differences are confined to an $O(\xi(x))$-thick shell near the boundary. If the system volume is $V = L^d$, then standard estimates (trace-norm/fidelity or Lieb–Robinson–type bounds in the lattice regularization) give, for some $C > 0$ independent of L,

$$\big\|\Omega_2(x) - \Omega_1(x)\big\| \;\le\; C\,\frac{\mathrm{Area}(\partial\Lambda)}{V}\, e^{-L/\xi(x)} \;=\; O\Big(e^{-L/\xi(x)}\Big)\,. \tag{7.683}$$

For normalized pure states, the overlap satisfies $1 - |\,\langle\Omega_2\rangle\,\Omega_1|\le \tfrac{1}{2}\|\Omega_2 - \Omega_1\|^2$, hence, inserting (7.683),

$$1 - \Big|\Lambda_{\mathrm{state}}^{(\lambda_{\mathrm{UV}},\ell)}(V)\Big| = 1 - \Big|\,\langle\Omega_2(x)\rangle\,\Omega_1(x)\Big| \;\le\; C'\,e^{-2L/\xi(x)} \;\xrightarrow{L\to\infty}\; 0. \tag{7.684}$$

Step 2c (Uniformity on $\mathfrak{P}$). By (7.682), $\xi(x) \le \xi_{\max}$ for all $x \in \mathfrak{P}$. Therefore (7.684) is uniform on $\mathfrak{P}$:

$$\sup_{x\in\mathfrak{P}} \Big(1 - \Big|\Lambda_{\mathrm{state}}^{(\lambda_{\mathrm{UV}},\ell)}(V)\Big|\Big) \;\le\; C''\,e^{-2L/\xi_{\max}} \;\xrightarrow{L\to\infty}\; 0,$$

which proves $\lim_{V\to\infty} |\Lambda_{\mathrm{state}}^{(\lambda_{\mathrm{UV}},\ell)}(V)| = 1$ uniformly on the plateau. As before, gauge-parameter independence and cohomological robustness follow from Proposition 7.556 and Lemma 7.552.

Conclusion. Steps (1a–1b) establish a *uniform* law-visibility deficit with some $\delta_\star \in (0,1)$; Steps (2a–2c) establish the *uniform* large-volume convergence of the state visibility to 1. Both statements hold on $\mathcal{H}_{\mathrm{phys}}$, are adjoint-covariant (Proposition 7.568), and are stable under BRST/improvement ambiguities (Lemma 7.552). This completes the proof. □

Corollary 7.581 (Channel additivity and robustness)**.** *With the triadic decomposition of Proposition 7.555,*

$$\Lambda_\bullet^{(\lambda_{\mathrm{UV}},\ell)} = \sum_{\alpha\in\{\mathrm{corr},\mathrm{vac},\mathrm{spec}\}} \Lambda_\bullet^{(\alpha;\lambda_{\mathrm{UV}},\ell)}, \qquad \bullet \in \{\mathrm{law},\mathrm{state}\},$$

the law-deficit lower bound of Theorem 7.580 is saturated (up to certified plateau tolerances) by the spectral (T-odd) contribution, while the state channel's approach to unity is saturated by the vacuum piece; both statements are uniform on $\mathfrak{P}$ and invariant under BRST- and Noether-improvements on $\mathcal{H}_{\mathrm{phys}}$.

Proof (expanded). We use: (i) channel additivity and orthogonality (Proposition 7.555); (ii) band descent and BRST/Nielsen safety (Lemma 7.552 and Proposition 7.556); (iii) the QIV law-deficit and state-limit established on certified plateaus (Theorem 7.580); (iv) plateau regularity (Definition 6.300); and (v) the scale-aware setup (Section 6.12).

(A) Law channel: deficit is (plateau-)saturated by the spectral branch. At a fixed band point $x = (\lambda_{\mathrm{UV}}, \ell) \in \mathfrak{P}$, decompose

$$\Lambda_{\mathrm{law}}(x) = \Lambda_{\mathrm{law}}^{(\mathrm{vac})}(x) + \Lambda_{\mathrm{law}}^{(\mathrm{corr})}(x) + \Lambda_{\mathrm{law}}^{(\mathrm{spec})}(x).$$

By construction of the holosymmetric visibilities, the *spectral* piece is T-odd, while the *vacuum* and *correction* pieces are T-even (they come from equal-time, symmetric correlators that do not contribute negatively to the real part at unity normalization). Therefore

$$1 - |\Lambda_{\mathrm{law}}(x)| \;\geq\; -\,\mathrm{Re}\;\Lambda_{\mathrm{law}}^{(\mathrm{spec})}(x). \tag{7.685}$$

The spectral contribution admits a positive-kernel representation (Kubo-type) at the band point (cf. the proof of Lemma 7.578):

$$-\,\mathrm{Re}\;\Lambda_{\mathrm{law}}^{(\mathrm{spec})}(x) = \int_{-\omega_0}^{\omega_0} d\omega\; \mathsf{K}_{\mathrm{spec}}(x;\omega)\, \mathsf{S}_a(x;\omega),$$

with $\mathsf{K}_{\mathrm{spec}}(x;\omega) \geq \kappa_{\mathrm{spec}}(x) > 0$ on a compact window and S_a the spectral density of the adjoint-covariant divergence for a broken direction a. If the Ward source $\mathcal{R}_a(x)$ is nonzero, then the integral is strictly positive (Lemma 7.578).

On a certified plateau $\mathfrak{P}$, oscillation bounds ensure that the *T-even* sum $\Lambda_{\mathrm{law}}^{(\mathrm{vac})} + \Lambda_{\mathrm{law}}^{(\mathrm{corr})}$ stays within $\eta_{\mathfrak{P}}$ of its x_0-value where the bound is attained. Hence,

$$1 - |\Lambda_{\mathrm{law}}(x)| \;=\; -\,\mathrm{Re}\;\Lambda_{\mathrm{law}}^{(\mathrm{spec})}(x) \;\pm\; O(\eta_{\mathfrak{P}}) \qquad \text{uniformly in } x \in \mathfrak{P}, \tag{7.686}$$

where the $\pm O(\eta_{\mathfrak{P}})$ captures the certified variation of the T-even branches (Definition 6.300). Combining (7.685)–(7.686) with Theorem 7.580 yields that the *lower bound* for the law-deficit is provided by the spectral branch and is *saturated up to* $O(\eta_{\mathfrak{P}})$.

Cohomological robustness: by Lemma 7.552, BRST-exact and Noether-improvement terms do not contribute on $\mathcal{H}_{\mathrm{phys}}$; thus both the bound and its (plateau) saturation are invariant under such redefinitions. Gauge-parameter independence is guaranteed by Proposition 7.556.

(B) State channel: convergence to unity is saturated by the vacuum branch. Decompose similarly,

$$\Lambda_{\mathrm{state}}(x;V) = \Lambda_{\mathrm{state}}^{(\mathrm{vac})}(x;V) + \Lambda_{\mathrm{state}}^{(\mathrm{corr})}(x;V) + \Lambda_{\mathrm{state}}^{(\mathrm{spec})}(x;V).$$

In QIV there is a *unique* vacuum (no exact NG directions); the would-be NG modes are pseudo-NG and gapped on the band (mass/mixing ledger, cf. Section 7.9), yielding a uniform correlation-length bound on $\mathfrak{P}$ (see the proof of Theorem 7.580). Standard clustering/exponential-decay arguments then imply that, as $V \to \infty$,

$$\Lambda_{\mathrm{state}}^{(\mathrm{corr})}(x;V) \;\to\; 0, \qquad \Lambda_{\mathrm{state}}^{(\mathrm{spec})}(x;V) \;\to\; 0, \qquad \text{uniformly in } x \in \mathfrak{P}, \tag{7.687}$$

while the vacuum branch converges to unity,

$$\Lambda_{\mathrm{state}}^{(\mathrm{vac})}(x;V) \;\to\; 1 \qquad \text{uniformly in } x \in \mathfrak{P}. \tag{7.688}$$

Thus

$$\lim_{V\to\infty}\; \Lambda_{\mathrm{state}}(x;V) = \lim_{V\to\infty}\; \Lambda_{\mathrm{state}}^{(\mathrm{vac})}(x;V) = 1 \quad \text{uniformly on } \mathfrak{P},$$

which is precisely the statement that the approach to unity is *saturated* by the vacuum piece. As before, BRST/improvement invariance follows from Lemma 7.552, and gauge-parameter independence from Proposition 7.556.

Conclusion. The channel-additive structure (Proposition 7.555) together with the QIV plateau diagnostics (Theorem 7.580) implies: (i) the law-channel deficit originates from—and, up to certified tolerances, is saturated by—the spectral (T-odd) branch; (ii) the state-channel limit to unity is saturated by the vacuum branch; in both cases uniformly on $\mathfrak{P}$ and stable under BRST- and Noether-improvements on $\mathcal{H}_{\text{phys}}$. □

7.9.5 State–visibility guardrail (no SSB / no decoherence of vacua)

Definition 7.582 (State–visibility guardrail and acceptance)**.** Let $\mathfrak{P}$ be a certified plateau mask (Definition 6.300) in the scale-aware setup (Section 6.12). The *state visibility* at a band point $x = (\lambda_{\text{UV}}, \ell)$ and finite volume $V = L^d$ is

$$\Lambda_{\text{state}}(x; V) := \langle \Omega_2^{(\lambda_{\text{UV}},\ell)}(V) \rangle \, \Omega_1^{(\lambda_{\text{UV}},\ell)}(V),$$

with branch states prepared as in Definitions 6.275 and 7.551. We say the state channel *passes the guardrail* on $\mathfrak{P}$ if there exists a tolerance function $\varepsilon_{\mathfrak{P}}(L) \downarrow 0$ such that

$$1 - |\Lambda_{\text{state}}(x; V)| \;\le\; \varepsilon_{\mathfrak{P}}(L) \qquad \text{for all } x \in \mathfrak{P}.$$

Two canonical acceptance envelopes are allowed:

1. *Exponential (gapped case):* $\varepsilon_{\mathfrak{P}}(L) = C\, e^{-L/\xi_{\max}}$ for some $C, \xi_{\max} > 0$ independent of $x \in \mathfrak{P}$.
2. *Power-law (critical but non-SSB case):* $\varepsilon_{\mathfrak{P}}(L) = C\, L^{-\gamma}$ for some $\gamma > 0$ independent of $x \in \mathfrak{P}$.

Gauge-parameter independence of Λ_{state} on $\mathcal{H}_{\text{phys}}$ is guaranteed by Proposition 7.556, and the band descent is BRST-safe by Lemma 7.552.

Lemma 7.583 (Fidelity and trace-distance control for band-prepared branches)**.** *For normalized branch states at $x \in \mathfrak{P}$ and volume V,*

$$1 - |\Lambda_{\text{state}}(x; V)| \;\le\; \tfrac{1}{2} \big\| \rho_2(x; V) - \rho_1(x; V) \big\|_1,$$

where ρ_i are the corresponding band-reduced density operators. The right-hand side is BRST/Nielsen safe on $\mathcal{H}_{\text{phys}}$ by Lemma 7.552 and Proposition 7.556.

Proof. Step 1: Band descent and reduced states. Let Φ_x denote the (UV/IR) band-preparation channel at x (the composition of $P_x(\lambda_{\text{UV}})$, $K_{\text{IR}}^{(\ell)}$, and the restriction to the finite volume V). Then Φ_x is a completely positive, trace-preserving (CPTP) map on density operators, and the band-reduced states are

$$\rho_i(x; V) \;:=\; \Phi_x\Big(|\Psi_i\rangle\langle\Psi_i| \Big), \qquad i = 1, 2,$$

for normalized branch vectors $|\Psi_i\rangle$ on the underlying Hilbert space. By Proposition 7.573, Φ_x commutes with BRST/G actions and is well defined on certified plateaus.

Step 2: Choice of branch overlap as Uhlmann amplitude. By Uhlmann's theorem, for any ρ_1, ρ_2 there exist purifications $|\varphi_i\rangle$ on a system+ancilla space such that

$$\left|\langle\varphi_2|\varphi_1\rangle\right| \;=\; \sqrt{F(\rho_1,\rho_2)}\,, \qquad \text{(Uhlmann fidelity) [385, 386].}$$

In the present setting we realise purifications canonically via a Stinespring dilation of Φ_x: $\rho_i = \mathrm{Tr}_E\big(U_x\,|\Psi_i\rangle\langle\Psi_i|\otimes|0\rangle\langle 0|\,U_x^\dagger\big)$. The freedom $|\varphi_i\rangle \mapsto (\mathbf{1}\otimes U_i)|\varphi_i\rangle$ on the ancilla allows us to *fix the relative ancilla gauge* so that the overlap equals the Uhlmann amplitude. We *define* the (post–band) state overlap as

$$|\Lambda_{\mathrm{state}}(x;V)| \;:=\; \sqrt{F\big(\rho_1(x;V),\rho_2(x;V)\big)}.$$

This convention matches the pure–state case (then ρ_i are pure and $|\Lambda_{\mathrm{state}}| = |\langle\Psi_2|\Psi_1\rangle|$) and is invariant under physically irrelevant ancilla unitaries.

Step 3: Fuchs–van de Graaf (FvdG) inequality. The FvdG bounds relate trace distance $D(\rho_1,\rho_2) := \frac{1}{2}\|\rho_1-\rho_2\|_1$ and fidelity [388]:

$$1-\sqrt{F(\rho_1,\rho_2)} \;\le\; D(\rho_1,\rho_2) \;\le\; \sqrt{1-F(\rho_1,\rho_2)}.$$

Applying the left-hand inequality with $\rho_i = \rho_i(x;V)$ and using the definition in Step 2 yields

$$1-|\Lambda_{\mathrm{state}}(x;V)| \;=\; 1-\sqrt{F(\rho_1,\rho_2)} \;\le\; \tfrac{1}{2}\,\|\rho_2-\rho_1\|_1,$$

which is the desired inequality.

Step 4: BRST/Nielsen safety. On $\mathcal{H}_{\mathrm{phys}}$, BRST-exact deformations do not change physical matrix elements, and gauge-parameter variations are Nielsen-trivial (Lemma 7.552 and Proposition 7.556). Since trace distance is invariant under unitary conjugations and *contractive* under CPTP maps, and since Φ_x commutes with BRST/G (Proposition 7.573), the quantity $\|\rho_2-\rho_1\|_1$ is gauge-parameter independent on certified plateaus. Hence the right-hand side is BRST/Nielsen safe.

This completes the proof. □

Lemma 7.584 (Boundary sensitivity bound (gapped case))**.** *Assume at $x\in\mathfrak{P}$ the band Hessian has a positive spectral gap $m_{\min}(x)>0$ in the would-be NG sector (no exact NG). Then there exist constants $C(x),\xi(x)<\infty$ such that for branch preparations differing only on a boundary layer of width $O(1)$,*

$$\Big\|\rho_2(x;V)-\rho_1(x;V)\Big\|_1 \;\le\; C(x)\,e^{-L/\xi(x)}.$$

If $\inf_{x\in\mathfrak{P}} m_{\min}(x)>0$, then $\sup_{x\in\mathfrak{P}}\xi(x) =: \xi_{\max}<\infty$ and $\sup_{x\in\mathfrak{P}} C(x) =: C<\infty$.

Proof. Setting and notation. Let V be a region of linear size L and let the two preparations differ only on a boundary layer S_{bdy} of thickness $O(1)$ adjacent to ∂V. Denote by Φ_x the band-preparation channel at x (composition of $P_x(\lambda_{\mathrm{UV}})$, $K_{\mathrm{IR}}^{(\ell)}$, and restriction to V), so $\rho_i(x;V) = \Phi_x(|\Psi_i\rangle\langle\Psi_i|)$. By Proposition 7.573, Φ_x is CPTP, commutes with BRST/G, and is quasi-local (cone-preserving).

Step 1: Linear-response representation of the difference. Connect the two boundary preparations by a path of boundary couplings $\{\mathcal{B}_s\}_{s\in[0,1]}$ supported on S_{bdy}, with $\mathcal{B}_0$ and $\mathcal{B}_1$ realizing the two choices. For any observable A supported in the deep interior V_{in} (distance $\gtrsim L$ from the boundary), the Gâteaux derivative of the band-expectation is given by the Kubo/Duhamel formula at x:

$$\frac{d}{ds}\,\mathrm{Tr}\big[A\,\rho_s(x;V)\big] = \int_{S_{\mathrm{bdy}}} d\Sigma \int_0^1 dt\,\Big\langle A^{(x)}\;;\;\mathcal{J}_s^{(x)}(t)\Big\rangle_c\,, \tag{7.689}$$

where $\rho_s(x;V) := \Phi_x(\rho_s)$, $\mathcal{J}_s$ is the boundary insertion local current (a linear functional of $\dot{\mathcal{B}}_s$), $A^{(x)}$ and $\mathcal{J}_s^{(x)}$ denote their band-descended versions, and $\langle\cdot;\cdot\rangle_c$ is the connected correlator on $\mathcal{H}_{\mathrm{phys}}$. The representation (7.689) follows from standard linear-response plus the fact that Φ_x is linear and commutes with BRST/G (Proposition 7.573); non–signalling of descended operators is guaranteed by Corollary 7.574.

Step 2: Exponential clustering. The spectral gap $m_{\min}(x) > 0$ for the (band) Hessian implies exponential clustering of connected correlators with correlation length $\xi(x) \sim m_{\min}(x)^{-1}$:

$$\Big|\langle A^{(x)}\;;\;\mathcal{J}_s^{(x)}(t)\rangle_c\Big| \;\le\; C_0(x)\,\|A\|\,\|\mathcal{J}_s\|\,e^{-\mathrm{dist}(V_{\mathrm{in}},S_{\mathrm{bdy}})/\xi(x)} \;\le\; C_0(x)\,\|A\|\,\|\mathcal{J}_s\|\,e^{-L/\xi(x)}. \tag{7.690}$$

Here $C_0(x)$ absorbs local norms and the finite time aperture coming from the IR tagger, and we used that band descent preserves locality and does not increase operator norm up to a finite factor on certified plateaus (Definition 6.300).

Step 3: From response to trace distance. Integrating (7.689) in $s \in [0,1]$ and $t \in [0,1]$, using (7.690), and the fact that the boundary layer has $O(1)$ thickness and $O(L^{d-2})$ area (absorbed into constants for fixed geometry), we obtain

$$\Big|\,\mathrm{Tr}\Big[A\;(\rho_2(x;V) - \rho_1(x;V))\Big]\Big| \;\le\; C_1(x)\,\|A\|\,e^{-L/\xi(x)}.$$

Taking the supremum over $\|A\| \le 1$ (the dual characterization of the trace norm) yields

$$\big\|\rho_2(x;V) - \rho_1(x;V)\big\|_1 \;\le\; C(x)\,e^{-L/\xi(x)}\,,$$

with $C(x)$ finite for each fixed x.

Step 4: Uniformity on $\mathfrak{P}$. If $\inf_{x\in\mathfrak{P}} m_{\min}(x) > 0$, continuity of the band spectrum and kernels on the certified plateau (compactness of the mask, Definition 6.300) implies $x \mapsto \xi(x)$ and $x \mapsto C(x)$ are bounded on $\mathfrak{P}$. Set $\xi_{\max} := \sup_{x\in\mathfrak{P}} \xi(x) < \infty$ and $C := \sup_{x\in\mathfrak{P}} C(x) < \infty$.

This proves the claim. □

Lemma 7.585 (Boundary sensitivity bound (critical but non-SSB case))**.** *If the band theory at $x \in \mathfrak{P}$ is gapless but has a unique vacuum (no SSB) and finite dynamic exponent $z < \infty$, then for boundary-layer preparations as above there exist $C(x), \gamma(x) > 0$ such that*

$$\big\|\rho_2(x;V) - \rho_1(x;V)\big\|_1 \;\le\; C(x)\,L^{-\gamma(x)}.$$

If critical exponents vary continuously on $\mathfrak{P}$, we may take $C := \sup_{x\in\mathfrak{P}} C(x) < \infty$ and $\gamma := \inf_{x\in\mathfrak{P}} \gamma(x) > 0$.

Proof. Step 1: Linear-response as in the gapped case. With the same setup and notation as in Lemma 7.584, the difference of band-reduced states against a bulk observable A supported in V_{in} admits the linear-response representation (7.689). The only change is the *clustering estimate* appropriate to criticality.

Step 2: Power-law clustering without SSB. At a critical point with unique vacuum and finite $z < \infty$, connected correlators decay as a *power law* set by scaling dimensions. Denote by $\Delta_A(x)$ the (band) scaling dimension of the bulk representative of A and by $\Delta_{\mathrm{b}}(x)$ the boundary scaling dimension of the insertion represented by $\mathcal{J}_s$.[18] Then there exists $\gamma(x) > 0$ such that for separations $r \equiv \mathrm{dist}(V_{\mathrm{in}}, S_{\mathrm{bdy}}) \sim L$,

$$\left|\langle A^{(x)} \,;\, \mathcal{J}_s^{(x)}(t)\rangle_c\right| \;\le\; C_0(x)\,\|A\|\,\|\mathcal{J}_s\|\,r^{-\gamma(x)}\,, \qquad \gamma(x) \;:=\; \Delta_A(x)+\Delta_{\mathrm{b}}(x)-d_{\mathrm{eff}}(x), \tag{7.691}$$

with $d_{\mathrm{eff}}(x) = d - 1 + z(x)$ the effective (boundary + time) scaling dimension. The bound (7.691) is the standard boundary-to-bulk decay from scale invariance in the absence of spontaneous order.

Step 3: Integration and trace-distance bound. Integrating (7.689) over $s \in [0,1]$, $t \in [0,1]$, and the boundary layer (whose thickness is $O(1)$) and using (7.691) with $r \sim L$ yields

$$\left|\,\mathrm{Tr}\Big[A\;(\rho_2(x;V)-\rho_1(x;V))\Big]\right| \;\le\; C_1(x)\,\|A\|\,L^{-\gamma(x)}.$$

Taking the supremum over $\|A\| \le 1$ gives

$$\Big\|\rho_2(x;V)-\rho_1(x;V)\Big\|_1 \;\le\; C(x)\,L^{-\gamma(x)}\,,$$

with $C(x)$ finite.

Step 4: Uniformity on $\mathfrak{P}$. On certified plateaus the critical data $(z, \Delta_A, \Delta_{\mathrm{b}})$ vary continuously (scale-aware regularity), hence $x \mapsto \gamma(x)$ and $x \mapsto C(x)$ are continuous. If $\inf_{x\in\mathfrak{P}}\gamma(x) > 0$, then setting $\gamma := \inf_{x\in\mathfrak{P}}\gamma(x)$ and $C := \sup_{x\in\mathfrak{P}} C(x)$ yields the uniform bound claimed.

This completes the proof. □

Remark 7.586. (i) The linear-response step is legitimate because the two preparations differ only by boundary terms of finite norm; the band-preparation channel is linear and CPTP, and commutes with BRST/G (Proposition 7.573). (ii) In the gapped case, (7.690) is the standard exponential clustering with correlation length $\xi(x) \propto m_{\min}(x)^{-1}$; in the critical case, (7.691) encodes boundary-to-bulk decay from scaling, with $\gamma(x)>0$ ensured by "no SSB." (iii) Non–signalling after band descent (Corollary 7.574) rules out contamination from outside the causal cones of the boundary layer.

Proposition 7.587 (Guardrail acceptance from boundary sensitivity)**.** *Under the hypotheses of Lemma 7.584 (gapped) or Lemma 7.585 (gapless non-SSB), the guardrail in Definition 7.582 holds with the corresponding envelope $\varepsilon_{\mathfrak{P}}(L)$, uniformly on $\mathfrak{P}$.*

[18] In the relativistic case $z = 1$, these are the usual CFT dimensions; for general finite z, replace time with $x^0 \sim r^{\,z}$ in the anisotropic scaling.

Proof. Step 1: From boundary sensitivity to a trace–distance envelope. In the gapped case (Lemma 7.584) there exist $C(x), \xi(x) < \infty$ such that

$$\left\|\rho_2(x;V) - \rho_1(x;V)\right\|_1 \;\le\; C(x)\, e^{-L/\xi(x)}.$$

If $\inf_{x\in\mathfrak{P}} m_{\min}(x) > 0$, continuity on the certified plateau (Definition 6.300) yields uniform bounds $\xi_{\max} := \sup_{x\in\mathfrak{P}} \xi(x) < \infty$ and $C := \sup_{x\in\mathfrak{P}} C(x) < \infty$, so

$$\sup_{x\in\mathfrak{P}} \left\|\rho_2(x;V) - \rho_1(x;V)\right\|_1 \;\le\; C\, e^{-L/\xi_{\max}}.$$

In the gapless non-SSB case (Lemma 7.585) there are $C(x), \gamma(x) > 0$ with $\|\rho_2(x;V) - \rho_1(x;V)\|_1 \le C(x)\, L^{-\gamma(x)}$. Assuming the scale data vary continuously on $\mathfrak{P}$, define $C := \sup_{x\in\mathfrak{P}} C(x) < \infty$ and $\gamma := \inf_{x\in\mathfrak{P}} \gamma(x) > 0$, hence

$$\sup_{x\in\mathfrak{P}} \left\|\rho_2(x;V) - \rho_1(x;V)\right\|_1 \;\le\; C\, L^{-\gamma}.$$

Step 2: From trace distance to the state guardrail. By the Fuchs–van de Graaf inequality (cf. Lemma 7.583),

$$1 - |\Lambda_{\text{state}}(x;V)| \;\le\; \tfrac{1}{2}\left\|\rho_2(x;V) - \rho_1(x;V)\right\|_1.$$

Taking the supremum over $x \in \mathfrak{P}$ and using the envelopes from Step 1 gives, uniformly on $\mathfrak{P}$,

$$\sup_{x\in\mathfrak{P}} \left(1 - |\Lambda_{\text{state}}(x;V)|\right) \;\le\; \begin{cases} \dfrac{C}{2}\, e^{-L/\xi_{\max}}, & \text{(gapped case)},\\[2ex] \dfrac{C}{2}\, L^{-\gamma}, & \text{(gapless non-SSB case)}. \end{cases}$$

Thus the guardrail of Definition 7.582 holds with $\varepsilon_{\mathfrak{P}}(L) = \frac{C}{2} e^{-L/\xi_{\max}}$ in the gapped case, or $\varepsilon_{\mathfrak{P}}(L) = \frac{C}{2} L^{-\gamma}$ in the critical non-SSB case. The constants are plateau–uniform by construction, completing the proof. □

Lemma 7.588 (Plateau uniformization for the state channel)**.** *Suppose for some $x_0 \in \mathfrak{P}$ and volume V we have $1 - |\Lambda_{\text{state}}(x_0;V)| \le \varepsilon_{\mathfrak{P}}(L)$, and the oscillation bound of Definition 6.300 holds for $\Lambda_{\text{state}}(\cdot;V)$. Then*

$$\sup_{x\in\mathfrak{P}} \left(1 - |\Lambda_{\text{state}}(x;V)|\right) \;\le\; \varepsilon_{\mathfrak{P}}(L) + \eta_{\mathfrak{P}}.$$

Proof. Let $f(x) := 1 - |\Lambda_{\text{state}}(x;V)| \ge 0$. The plateau oscillation bound from Definition 6.300 asserts that

$$\sup_{x,y\in\mathfrak{P}} |f(x) - f(y)| \;\le\; \eta_{\mathfrak{P}}.$$

Fix $x_0 \in \mathfrak{P}$ with $f(x_0) \le \varepsilon_{\mathfrak{P}}(L)$. Then for any $x \in \mathfrak{P}$,

$$f(x) \;\le\; f(x_0) + |f(x) - f(x_0)| \;\le\; \varepsilon_{\mathfrak{P}}(L) + \eta_{\mathfrak{P}}.$$

Taking the supremum over $x \in \mathfrak{P}$ yields the claim. Equivalently, this is the direct application of Lemma 7.579 with baseline point x_0 and envelope $\varepsilon_{\mathfrak{P}}(L)$. □

Theorem 7.589 (State–visibility guardrail: no decoherence of vacua)**.** *Assume (i) unique vacuum (no SSB) on the band for all $x \in \mathfrak{P}$, and (ii) either a uniform mass gap on $\mathfrak{P}$ or a critical but non-SSB regime with continuous exponents on $\mathfrak{P}$. Then the state channel passes the guardrail of Definition 7.582: uniformly on $\mathfrak{P}$,*

$$\lim_{V\to\infty} \; |\Lambda_{\text{state}}(x;V)| \;=\; 1.$$

If instead there exists a subsequence of volumes for which $|\Lambda_{\text{state}}(x;V)|$ exhibits a systematic decay bounded below by a positive constant independent of L on any refined mask, the system drifts toward QII (SSB-induced orthogonalization) and the guardrail fails.

Proof. Step 1: Uniform envelope from boundary sensitivity. For the gapped case, Lemma 7.584 yields $\|\rho_2(x;V) - \rho_1(x;V)\|_1 \le C(x)\, e^{-L/\xi(x)}$ whenever the two branch preparations differ only on an $O(1)$ boundary layer; assuming a plateau–uniform gap, we get the uniform bound $\sup_{x\in\mathfrak{P}} \|\rho_2 - \rho_1\|_1 \le C\, e^{-L/\xi_{\max}}$. For the gapless but non-SSB case, Lemma 7.585 gives $\sup_{x\in\mathfrak{P}} \|\rho_2 - \rho_1\|_1 \le C\, L^{-\gamma}$ for some $C < \infty$ and $\gamma > 0$ depending only on $\mathfrak{P}$.

Step 2: From trace distance to state visibility. By the Fuchs–van de Graaf bound (Lemma 7.583),

$$1 - |\Lambda_{\text{state}}(x;V)| \;\le\; \tfrac{1}{2}\, \|\rho_2(x;V) - \rho_1(x;V)\|_1.$$

Taking the supremum over $x \in \mathfrak{P}$ and using Step 1 gives, uniformly on $\mathfrak{P}$,

$$\sup_{x\in\mathfrak{P}} \big(1 - |\Lambda_{\text{state}}(x;V)|\big) \;\le\; \begin{cases} \frac{C}{2}\, e^{-L/\xi_{\max}}, & \text{(gapped)}, \\ \frac{C}{2}\, L^{-\gamma}, & \text{(critical, non-SSB)}. \end{cases}$$

Equivalently, the guardrail of Definition 7.582 holds with a decaying envelope $\varepsilon_{\mathfrak{P}}(L)$; this is exactly Proposition 7.587. Hence $\lim_{V\to\infty} \sup_{x\in\mathfrak{P}} (1 - |\Lambda_{\text{state}}(x;V)|) = 0$, which implies the stated uniform limit $|\Lambda_{\text{state}}(x;V)| \to 1$ on $\mathfrak{P}$.

Step 3: Plateau uniformization (optional sharpening). If one has a baseline point $x_0 \in \mathfrak{P}$ with $1 - |\Lambda_{\text{state}}(x_0;V)| \le \varepsilon_{\mathfrak{P}}(L)$, the plateau oscillation bound (Definition 6.300) and Lemma 7.588 give $\sup_{x\in\mathfrak{P}} (1 - |\Lambda_{\text{state}}(x;V)|) \le \varepsilon_{\mathfrak{P}}(L) + \eta_{\mathfrak{P}}$, which also tends to 0 as $L \to \infty$.

Step 4: Contrapositive / failure mode implies drift to QII. Suppose there exists a subsequence V_n and a mask refinement sequence (staying certified) such that, for some $x^* \in \mathfrak{P}$ and all large n,

$$1 - |\Lambda_{\text{state}}(x^*;V_n)| \;\ge\; \delta_\star > 0,$$

with $\delta_\star$ independent of L_n. Then by Lemma 7.583, $\|\rho_2(x^*;V_n) - \rho_1(x^*;V_n)\|_1 \ge 2\delta_\star$ for all large n. This contradicts the envelopes of Step 1 unless at least one hypothesis of the theorem fails. If the certified plateau remains valid, the only possible failure is the uniqueness of the vacuum: persistent, volume–independent distinguishability of boundary-layer preparations signals the emergence of multiple orthogonal vacua (spontaneous symmetry breaking). In the quadrant language, this is precisely the QII drift (state orthogonalization in the thermodynamic limit). Therefore, sustained decay of $|\Lambda_{\text{state}}|$ bounded away from zero along a refined certified sequence is a witness of QII onset and the guardrail fails.

Step 5: BRST/Nielsen safety. All steps above are taken in the physical sector $\mathcal{H}_{\text{phys}}$; the band-preparation channel commutes with BRST/G actions (Proposition 7.573), non–signalling after band descent holds (Corollary 7.574), and the trace-distance and visibility statements are gauge-parameter independent by Lemmas 7.552 and 7.583 and Proposition 7.556.

This completes the proof. □

Corollary 7.590 (Operational test (log–lin / log–log slopes))**.** *Given measurements of* $1 - |\Lambda_{\text{state}}(x; V)|$ *at increasing* L*:*

(i) Gapped case. *On any certified volume-window, a linear fit in* log–lin *scale has slope* $\le -1/\xi_{\max}$ *within tolerance; failure flags guardrail violation.*

(ii) Critical non-SSB case. *On any certified volume-window, a linear fit in* log–log *scale has slope* $\le -\gamma$ *within tolerance; failure flags guardrail violation.*

Both tests are BRST/Nielsen safe on $\mathcal{H}_{\text{phys}}$ *(Lemma 7.552 and Proposition 7.556) and should be applied uniformly over* $x \in \mathfrak{P}$ *with Lemma 7.588.*

Proof. Preliminaries (uniform envelope and uniformization). By Proposition 7.587, there exists a decaying envelope $\varepsilon_{\mathfrak{P}}(L)$ such that, uniformly on $x \in \mathfrak{P}$,

$$1 - |\Lambda_{\text{state}}(x; V)| \;\le\; \varepsilon_{\mathfrak{P}}(L). \tag{7.692}$$

Moreover, with the plateau oscillation bound (Definition 6.300) and Lemma 7.588, replacing x by the $\sup_{x\in\mathfrak{P}}$ only adds at most $\eta_{\mathfrak{P}}$:

$$y(L) \;:=\; \sup_{x\in\mathfrak{P}} \Big(1 - |\Lambda_{\text{state}}(x; V)|\Big) \;\le\; \varepsilon_{\mathfrak{P}}(L) + \eta_{\mathfrak{P}}. \tag{7.693}$$

We call a *certified volume-window* any interval $[L_{\min}, L_{\max}]$ on which

$$\frac{\eta_{\mathfrak{P}}}{\varepsilon_{\mathfrak{P}}(L)} \;\le\; \tau \quad \text{for all } L \in [L_{\min}, L_{\max}], \tag{7.694}$$

with a chosen tolerance parameter $0 < \tau < 1$ (so the envelope dominates the oscillation floor).

(i) Gapped case: log–lin slope. Here $\varepsilon_{\mathfrak{P}}(L) = \frac{C}{2}\, e^{-L/\xi_{\max}}$ (Proposition 7.587). Using (7.693),

$$\log y(L) \;\le\; \log \varepsilon_{\mathfrak{P}}(L) + \log\Big(1 + \frac{\eta_{\mathfrak{P}}}{\varepsilon_{\mathfrak{P}}(L)}\Big) = \log\Big(\tfrac{C}{2}\Big) - \frac{L}{\xi_{\max}} + \log\Big(1 + r(L)\Big),$$

with $r(L) := \eta_{\mathfrak{P}}/\varepsilon_{\mathfrak{P}}(L)$. On a certified window (7.694), $r(L) \le \tau$ and thus

$$\frac{\log y(L_2) - \log y(L_1)}{L_2 - L_1} \;\le\; -\frac{1}{\xi_{\max}} + \frac{\log(1 + r(L_2)) - \log(1 + r(L_1))}{L_2 - L_1} \;\le\; -\frac{1}{\xi_{\max}} + \frac{\log(1+\tau)}{L_2 - L_1}.$$

Therefore every *secant* slope on $[L_{\min}, L_{\max}]$ is $\le -1/\xi_{\max} + \delta_{\text{lin}}$ with

$$\delta_{\text{lin}} \;:=\; \frac{\log(1+\tau)}{L_{\max} - L_{\min}}.$$

Since the least-squares slope of a line fitted to $(L_i, \log y(L_i))$ is bounded above by the maximum secant slope on the fitting interval, the fitted slope is also $\le -1/\xi_{\max} + \delta_{\text{lin}}$. Choosing τ (how far above the oscillation floor we stay) and the window width $L_{\max} - L_{\min}$ fixes the *tolerance* δ_{lin}. If the fitted slope exceeds $-1/\xi_{\max} + \delta_{\text{lin}}$, then (7.693)–(7.694) cannot simultaneously hold, flagging a guardrail violation.

(ii) Critical non-SSB case: log–log slope. Here $\varepsilon_{\mathfrak{P}}(L) = \frac{C}{2} L^{-\gamma}$ (Proposition 7.587). With $X = \log L$,

$$\log y(L) \;\le\; \log \varepsilon_{\mathfrak{P}}(L) + \log\Big(1 + r(L)\Big) = \log\left(\tfrac{C}{2}\right) - \gamma \log L + \log\Big(1 + r(L)\Big),$$

again with $r(L) = \eta_{\mathfrak{P}}/\varepsilon_{\mathfrak{P}}(L)$. On a certified window (7.694), $r(L) \le \tau$, so for any $L_1 < L_2$,

$$\frac{\log y(L_2) - \log y(L_1)}{\log L_2 - \log L_1} \;\le\; -\gamma \;+\; \frac{\log(1 + r(L_2)) - \log(1 + r(L_1))}{\log L_2 - \log L_1} \;\le\; -\gamma \;+\; \frac{\log(1+\tau)}{\log(L_2/L_1)}.$$

Thus every *log–log* secant slope on $[L_{\min}, L_{\max}]$ is $\le -\gamma + \delta_{\log}$ with

$$\delta_{\log} \;:=\; \frac{\log(1+\tau)}{\log(L_{\max}/L_{\min})}.$$

As before, the least-squares slope in $(\log L, \log y)$ is bounded above by the maximal secant slope, so the fitted slope is $\le -\gamma + \delta_{\log}$. If the fitted slope is larger than $-\gamma + \delta_{\log}$, the guardrail is violated on that window.

BRST/Nielsen safety and uniformity. All quantities are computed from band-descended, BRST-closed observables on $\mathcal{H}_{\text{phys}}$, and the preparation/measurement maps commute with BRST/G (Proposition 7.573), hence the tests are gauge-parameter independent (Lemma 7.552 and Proposition 7.556). Applying Lemma 7.588 ensures the analysis holds uniformly over $x \in \mathfrak{P}$ (the measured series can be pooled or tested pointwise with the same tolerance).

This proves both operational tests. □

7.9.6 State–visibility Envelope Selection, CIs, and Fallback Protocol

Context. On a certified plateau $\mathfrak{P}$ (Definition 6.300) the state channel must *not* exhibit thermodynamic orthogonality (QII signature). The guardrail (Definition 7.582 and Theorem 7.589) admits two acceptance envelopes, motivated by Lemmas 7.584 and 7.585: (i) exponential (gapped) and (ii) power–law (critical, non–SSB). We formalize the model selection per band point x, estimate $(\xi(x), \gamma(x))$ with CIs, aggregate to $(\xi_{\max}, \gamma)$ on $\mathfrak{P}$, and specify a fallback rule.

Definition 7.591 (Volume series and two candidate envelopes)**.** At $x = (\lambda_{\text{UV}}, \ell) \in \mathfrak{P}$, measure

$$D_x(L_n) \;:=\; 1 - |\Lambda_{\text{state}}(x; V_n)|, \qquad V_n = L_n^d, \quad n = 1, \dots, N,$$

with $L_{n+1}/L_n = r > 1$ fixed on the plateau window, and weights $w_n > 0$ from readout uncertainties. Fit the two candidate models:

$$\text{(Exp)} \quad D_x(L) \;=\; C_x\, e^{-L/\xi(x)}\,(1 + \varepsilon_L), \tag{7.695}$$

$$\text{(Pow)} \quad D_x(L) \;=\; C_x\, L^{-\gamma(x)}\,(1 + \varepsilon_L), \tag{7.696}$$

where ε_L is a small model error controlled by Lemmas 7.584 and 7.585.

Proposition 7.592 (WLS estimators and confidence intervals)**.** *Apply weighted least squares (WLS) in the linearized coordinates.*
Exponential (gapped): *log–lin regression* $y_n := \log D_x(L_n) = \alpha_x - L_n/\xi(x) + \eta_n$ *with weights* w_n*. Then* $\widehat{\xi}(x) = -1/\widehat{s}_E$ *where* $\widehat{s}_E$ *is the fitted slope; obtain* $\mathrm{CI}_{1-\beta}(\xi(x))$ *by the delta–method from the slope CI.*
Power–law (critical non-SSB): *log–log regression* $y_n := \log D_x(L_n) = \alpha_x - \gamma(x)\log L_n + \eta_n$ *with weights* w_n*; slope gives* $\widehat{\gamma}(x)$ *and its CI.*

Both fits are BRST/Nielsen safe on $\mathcal{H}_{\mathrm{phys}}$ *by Lemma 7.552 and Proposition 7.556.*

Proof. Setup and notation. For a fixed $x \in \mathfrak{P}$, let $D_x(L_n) > 0$ denote the measured state–deficit at linear size L_n (e.g. $D_x(L_n) = 1 - |\Lambda_{\mathrm{state}}(x; V_n)|$ on a certified mask). Define the linear models

$$y_n \equiv \log D_x(L_n) = \alpha_x + s\, X_n + \eta_n, \qquad X_n = \begin{cases} L_n & \text{(exponential)}, \\ \log L_n & \text{(power–law)}, \end{cases}$$

with $\mathbb{E}[\eta_n \mid X_n] = 0$ and $\mathrm{Var}(\eta_n \mid X_n) \propto w_n^{-1}$ (the proportionality can be absorbed into the scale of w_n). Let $W = \mathrm{diag}(w_1, \dots, w_N)$ and $X = \begin{bmatrix} \mathbf{1} & \mathbf{X} \end{bmatrix}$ with $\mathbf{X} = (X_1, \dots, X_N)^\top$.

WLS estimator and its variance. The WLS estimator $\widehat{\beta} = (\widehat{\alpha}, \widehat{s})^\top$ solves

$$\widehat{\beta} = (X^\top W X)^{-1} X^\top W\, \mathbf{y},$$

so the slope has the familiar centered form

$$\widehat{s} = \frac{\sum_n w_n (X_n - \bar{X}_w)(y_n - \bar{y}_w)}{\sum_n w_n (X_n - \bar{X}_w)^2}, \qquad \bar{X}_w = \frac{\sum_n w_n X_n}{\sum_n w_n}, \quad \bar{y}_w = \frac{\sum_n w_n y_n}{\sum_n w_n}.$$

If $w_n \propto \mathrm{Var}(\eta_n)^{-1}$ (well-specified WLS), then

$$\mathrm{Var}(\widehat{s} \mid X) = \sigma^2 \Big(\sum_n w_n (X_n - \bar{X}_w)^2 \Big)^{-1},$$
$$\widehat{\sigma}^2 := \frac{\sum_n w_n \widehat{\eta}_n^2}{N - 2}.$$

and an approximate $(1-\beta)$ slope–CI is $\widehat{s} \pm t_{N-2,\,1-\beta/2}\ \widehat{\mathrm{se}}(\widehat{s})$, with $\widehat{\mathrm{se}}(\widehat{s}) = \widehat{\sigma}\left(\sum_n w_n (X_n - \bar{X}_w)^2\right)^{-1/2}$. If the weights are only approximate, use the usual WLS *sandwich* (HC) covariance:

$$\widehat{\mathrm{Var}}(\widehat{\beta}) = (X^\top W X)^{-1} \Big(\sum_n w_n^2\, \widehat{\eta}_n^2\, x_n x_n^\top \Big) (X^\top W X)^{-1},$$
$$\text{where } x_n = (1, X_n)^\top,$$

and read off $\widehat{\mathrm{Var}}(\widehat{s})$ from the $(2,2)$ entry.

Exponential case: estimator for $\xi(x)$. Here $s = s_E = -1/\xi(x) < 0$. Define $g(s) := -1/s$ so that $\xi(x) = g(s_E)$. By asymptotic normality of $\widehat{s}$ (linear model with fixed design and

standard regularity), the *delta–method* gives

$$\widehat{\xi}(x) = g(\widehat{s}) = -\frac{1}{\widehat{s}},$$
$$\widehat{\mathrm{Var}}\big(\widehat{\xi}(x)\big) \approx \big(g'(\widehat{s})\big)^2\, \widehat{\mathrm{Var}}(\widehat{s}) = \frac{\widehat{\mathrm{Var}}(\widehat{s})}{\widehat{s}^{\,4}}.$$

Thus an approximate $(1-\beta)$ CI is

$$\mathrm{CI}_{1-\beta}\big(\xi(x)\big) \;:\; \widehat{\xi}(x) \;\pm\; z_{1-\beta/2}\,\frac{\widehat{\mathrm{se}}(\widehat{s})}{|\widehat{s}|^2},$$

with $z_{1-\beta/2}$ (or replace by $t_{N-2,1-\beta/2}$). Because g is monotone on $(-\infty,0)$, one may equivalently *transform the slope–CI* endpoints: if $[\widehat{s}_-,\widehat{s}_+]$ is the $(1-\beta)$ CI for s with $\widehat{s}_- < \widehat{s}_+ < 0$, then

$$\mathrm{CI}_{1-\beta}\big(\xi(x)\big) \;=\; \Big[-\frac{1}{\widehat{s}_+},\; -\frac{1}{\widehat{s}_-}\Big],$$

which is first–order equivalent to the delta–method CI. (If a fitted slope is nonnegative on a window, this violates the guardrail; cf. Corollary 7.590.)

Power–law case: estimator for $\gamma(x)$. Here $s = s_P = -\gamma(x)$. Hence $\widehat{\gamma}(x) = -\widehat{s}$, and the slope–CI $\widehat{s} \pm z_{1-\beta/2}\,\widehat{\mathrm{se}}(\widehat{s})$ maps to

$$\mathrm{CI}_{1-\beta}\big(\gamma(x)\big) \;:\; -\widehat{s} \;\mp\; z_{1-\beta/2}\,\widehat{\mathrm{se}}(\widehat{s}).$$

No delta transform is needed (the map $s \mapsto -s$ is exact and linear).

Regularity, certification, and BRST/Nielsen safety. The log–transform requires $D_x(L_n) > 0$, guaranteed on certified plateaus where measurements lie strictly above the floor (cf. the kernel/spectral floors and mask eligibility). Because D_x is built from band–descended, BRST–closed quantities and the taggers commute with BRST/G (Proposition 7.573), the regression statistics are gauge–parameter independent (Lemma 7.552 and Proposition 7.556). Standard linear–model regularity (bounded design, $\sum_n w_n(X_n - \bar{X}_w)^2 > 0$, finite fourth moments of η_n) yields consistency and asymptotic normality of the WLS slope, justifying the CIs above.

Conclusion. In log–lin (gapped) and log–log (critical non–SSB) coordinates, WLS provides slope estimators whose transforms yield $\widehat{\xi}(x)$ and $\widehat{\gamma}(x)$, with confidence intervals obtained from the slope–CI by the delta–method (exponential) or direct sign–flip (power–law). This establishes the proposition. □

Definition 7.593 (Model selection and mask aggregation)**.** For each $x \in \mathfrak{P}$, compute WLS residual sums $RSS_E(x)$ and $RSS_P(x)$ and the small–sample AIC_c:

$$\mathrm{AIC}_c^{(M)}(x) = N\log\frac{RSS_M(x)}{N} + 2k + \frac{2k(k+1)}{N-k-1}, \quad M \in \{E,P\},$$

with $k = 2$ (intercept and slope). Let $\Delta(x) := \mathrm{AIC}_c^{(P)}(x) - \mathrm{AIC}_c^{(E)}(x)$. Choose the envelope per x by:

$$\text{If } \Delta(x) \ge \Delta_\star \;\Rightarrow\; \text{Exp at } x \quad ; \quad \text{if } \Delta(x) \le -\Delta_\star \;\Rightarrow\; \text{Pow at } x \quad ; \quad \text{else: ambiguous.}$$

We recommend $\Delta_\star \in [4,6]$ (strong evidence). Define the *plateau envelopes*

$$\xi_{\max} := \sup_{x\in\mathfrak{P}_E} \overline{\xi}(x), \qquad \gamma := \inf_{x\in\mathfrak{P}_P} \underline{\gamma}(x),$$

where $\mathfrak{P}_E = \{x : \text{Exp chosen}\}$, $\mathfrak{P}_P = \{x : \text{Pow chosen}\}$, and $\overline{\xi}(x)$ (resp. $\underline{\gamma}(x)$) are the upper (resp. lower) ends of the chosen $(1-\beta)$ CIs.

Proposition 7.594 (Guardrail acceptance with CIs)**.** *If $\mathfrak{P}_E \cup \mathfrak{P}_P$ covers $\mathfrak{P}$ and the aggregated $\xi_{\max} < \infty$ and $\gamma > 0$, then the guardrail of Definition 7.582 holds uniformly on $\mathfrak{P}$ with the envelope*

$$D_x(L) \;\le\; \begin{cases} C\, e^{-L/\xi_{\max}}, & x \in \mathfrak{P}_E, \\ C\, L^{-\gamma}, & x \in \mathfrak{P}_P, \end{cases}$$

up to the plateau oscillation $\eta_{\mathfrak{P}}$. Gauge-parameter independence follows from Proposition 7.556.

Proof. Step 1: Pointwise envelopes. For $x \in \mathfrak{P}_E$ (gapped), Lemma 7.584 provides constants $C_E(x) < \infty$ and $\xi(x) < \infty$ such that

$$D_x(L) \;\le\; C_E(x)\, e^{-L/\xi(x)}. \tag{7.697}$$

For $x \in \mathfrak{P}_P$ (critical, non-SSB), Lemma 7.585 provides $C_P(x) < \infty$ and $\gamma(x) > 0$ with

$$D_x(L) \;\le\; C_P(x)\, L^{-\gamma(x)}. \tag{7.698}$$

Here $D_x(L) := 1 - |\Lambda_{\text{state}}(x; V)|$ on a certified mask for volume $V \sim L^d$.

Step 2: Aggregating decay rates. By hypothesis there exist *aggregated* exponents

$$\xi_{\max} := \sup_{x\in\mathfrak{P}_E} \xi(x) \;<\; \infty, \qquad \gamma := \inf_{x\in\mathfrak{P}_P} \gamma(x) \;>\; 0.$$

Then for any $x \in \mathfrak{P}_E$, $e^{-L/\xi(x)} \le e^{-L/\xi_{\max}}$, hence (7.697) implies $D_x(L) \le C_E(x)\, e^{-L/\xi_{\max}}$. For any $x \in \mathfrak{P}_P$, $L^{-\gamma(x)} \le L^{-\gamma}$, so (7.698) gives $D_x(L) \le C_P(x)\, L^{-\gamma}$.

Step 3: Uniform prefactor. Set

$$C \;:=\; \max\Big\{ \sup_{x\in\mathfrak{P}_E} C_E(x),\; \sup_{x\in\mathfrak{P}_P} C_P(x) \Big\} \;<\; \infty,$$

where finiteness follows from certified plateau regularity/compactness (Definition 6.300) and continuity of the band-preparation channel used in Lemmas 7.584 and 7.585. Then, for all $x \in \mathfrak{P}$,

$$D_x(L) \;\le\; \begin{cases} C\, e^{-L/\xi_{\max}}, & x \in \mathfrak{P}_E, \\ C\, L^{-\gamma}, & x \in \mathfrak{P}_P. \end{cases}$$

Step 4: Incorporating plateau oscillations. By the oscillation bound in Definition 6.300 and the uniformization argument (Lemma 7.588), passing to $\sup_{x\in\mathfrak{P}} D_x(L)$ inflates the envelope by at most $\eta_{\mathfrak{P}}$ on any certified window. This yields the stated envelope "up to $\eta_{\mathfrak{P}}$".

Step 5: How CIs supply the aggregates (optional). From the WLS fits in Proposition 7.592, let the slope CIs be $s_E(x) \in [\widehat{s}_E^-(x), \widehat{s}_E^+(x)] \subset (-\infty, 0)$ in the exponential case and $s_P(x) \in [\widehat{s}_P^-(x), \widehat{s}_P^+(x)] \subset (-\infty, 0)$ in the power-law case. With $\xi(x) = -1/s_E(x)$ and $\gamma(x) = -s_P(x)$, conservative CI–based aggregates are

$$\widehat{\xi}_{\max} := \sup_{x \in \mathfrak{P}_E} \Big(- \frac{1}{\widehat{s}_E^+(x)} \Big), \qquad \widehat{\gamma} := \inf_{x \in \mathfrak{P}_P} \Big(- \widehat{s}_P^+(x) \Big),$$

which satisfy $\xi(x) \le \widehat{\xi}_{\max}$ and $\gamma(x) \ge \widehat{\gamma}$ simultaneously on $\mathfrak{P}$ with the desired (simultaneous) confidence level (e.g. via Bonferroni). Plugging these into Steps 2–4 gives a CI–validated uniform guardrail.

Step 6: Gauge-parameter independence. All constructions are in the physical sector and the taggers commute with BRST/G (Proposition 7.556), hence the envelope is gauge-parameter independent.

Combining the steps establishes the proposition. □

Definition 7.595 (Fallback protocol for ambiguous fits)**.** If $\Delta(x) \in (-\Delta_\star, \Delta_\star)$ (*ambiguous*) or both slope CIs include 0, proceed:

1. **Tighten the mask:** shrink to a submask $\mathfrak{P}' \subset \mathfrak{P}$ with reduced oscillation $\eta_{\mathfrak{P}'} < \eta_{\mathfrak{P}}$ (cf. Definition 6.300) and/or increase the minimum volume $L_{\min}$ to improve separation of models; refit.
2. **Volume extension:** if feasible, add one or two larger L points to increase dynamic range; refit.
3. **Flag drift toward QII:** if for some subsequence $L_m \to \infty$ we observe:

$$\liminf_{m \to \infty} D_x(L_m) \ge \delta_0 > 0$$

with δ_0 exceeding the plateau tolerance (and stable under (1)–(2)), then $|\Lambda_{\text{state}}(x; V)|$ fails to approach 1; flag drift toward QII at x (orthogonalization of vacua), per Theorem 7.589.

Remark 7.596 (Operational thresholds)**.** Set $\Delta_\star = 6$ (strong AIC_c evidence), confidence level $1 - \beta = 0.95$, and require at least $N \ge 5$ volumes with ratio $r \in [1.4, 2]$. For the drift flag, take $\delta_0 = c_s\, \eta_{\mathfrak{P}}$ with $c_s \in [5, 10]$ (cf. Remark 7.600); report per–x decisions and the coverage of $\mathfrak{P}$ by $\mathfrak{P}_E \cup \mathfrak{P}_P$.

Definition 7.597 (Reporting for the state–visibility envelope)**.** For each $x \in \mathfrak{P}$: (i) chosen model (Exp/Pow/ambiguous), (ii) $(\widehat{\xi}(x), \text{CI})$ or $(\widehat{\gamma}(x), \text{CI})$, (iii) AIC_c difference $\Delta(x)$, (iv) residual plot and lack–of–fit test, (v) guardrail decision (pass/fail), and (vi) drift flag if triggered. Aggregate to $(\xi_{\max}, \gamma)$ with CIs as in Definition 7.593, and certify the guardrail via Proposition 7.594.

7.9.7 Law–visibility Slope Test and Tomographic Calibration

Context. On a certified plateau $\mathfrak{P}$ (Definition 6.300), QIV demands a *scale–independent* law–deficit: $1 - |\Lambda_{\text{law}}^{(x)}|$ stays bounded away from 0 with no IR improvement (Lemmas 7.578

and 7.579, Proposition 7.616, and Corollary 7.617). We now (i) make the *numerical slope test* explicit and (ii) give a BRST/Nielsen–safe *calibration protocol* for $\kappa_{\mathrm{spec}}(x)$ and $c_0(x)$ (entering the T–odd tomographic lower bound, Proposition 7.609, and the low–frequency tracking, Lemma 7.608).

Definition 7.598 (Log–slope estimator on the mask)**.** Fix $x = (\lambda_{\mathrm{UV}}, \ell) \in \mathfrak{P}$ and measure $y(\ell) := 1 - |\Lambda_{\mathrm{law}}^{(\lambda_{\mathrm{UV}},\ell)}|$ on a log–spaced window $\ell_n = \ell_0\, r^n$, $n = 0, \dots, N$ with $r > 1$. Define midpoint log–derivative estimates

$$S_n(x) \;:=\; \frac{\log y(\ell_{n+1}) - \log y(\ell_n)}{\log \ell_{n+1} - \log \ell_n} \;=\; \frac{\log\big(y(\ell_{n+1})/y(\ell_n)\big)}{\log r}, \qquad n = 0, \dots, N-1.$$

Let $\widehat{S}(x)$ be their inverse–variance weighted mean, with weights set by the readout uncertainties; define the mask–uniform statistic $S_{\max} := \sup_{x\in\mathfrak{P}} |\widehat{S}(x)|$.

Proposition 7.599 (QIV acceptance vs. QIII rejection (slope test))**.** *Choose a tolerance $\sigma_{\mathrm{tol}} > 0$ satisfying $\sigma_{\mathrm{tol}} \gg \eta_{\mathfrak{P}}$ (instrument floor) yet $\sigma_{\mathrm{tol}} \ll 1$ (no spurious power–law). Then:*

*(i) (*QIV accept*) If $S_{\max} \le \sigma_{\mathrm{tol}}$ and the constant model $y(\ell) \equiv y_\star$ lies within the plateau envelope (oscillation bound of Definition 6.300), the law channel passes the "no IR improvement" criterion uniformly on $\mathfrak{P}$.*

*(ii) (*QIII reject*) Fit the power–law model $y(\ell) = A\,\ell^{\sigma}$ by weighted least squares. If the 95% CI for σ excludes 0 and the likelihood–ratio (or F–test) favors the power–law over the constant by more than the plateau tolerance budget, reject QIV in favor of QIII on that mask.*

Gauge–parameter independence is guaranteed by Proposition 7.556; BRST safety of the readouts by Lemma 7.552.

Proof. Notation and statistic. Take a geometric IR grid $\ell_n = \ell_0 r^{\,n}$ with $r > 1$, $n = 0, \dots, N-1$, and measurements $y_n := y(\ell_n)$ for the (dimensionless) T–odd law–channel readout on a certified mask. Define the log–log secant slopes

$$S_n(x) \;:=\; \frac{\log y_{n+1} - \log y_n}{\log \ell_{n+1} - \log \ell_n} = \frac{\log y_{n+1} - \log y_n}{\log r}\,, \qquad n = 0, \dots, N-2,$$

and their maximum magnitude $S_{\max} := \max_n |S_n(x)|$. (Any WLS slope $\widehat{S}(x)$ fitted to $(\log \ell_n, \log y_n)$ lies between the min/max secants, so $|\widehat{S}(x)| \le S_{\max}$.)

(1) QIV accept. Assume the *constant* model $y(\ell) \equiv y_\star$ holds up to the plateau oscillation budget:

$$|y(\ell) - y_\star| \;\le\; \eta_{\mathfrak{P}} \qquad \text{for all } \ell \text{ on the certified window.} \tag{7.699}$$

Let $y_n = y_\star + \delta_n$ with $|\delta_n| \le \eta_{\mathfrak{P}}$. By the mean–value theorem,

$$\log y_{n+1} - \log y_n = \frac{\delta_{n+1} - \delta_n}{\xi_n}\,, \qquad \xi_n \in \big[y_\star - \eta_{\mathfrak{P}},\, y_\star + \eta_{\mathfrak{P}}\big].$$

Hence

$$|S_n(x)| \;=\; \frac{|\delta_{n+1}-\delta_n|}{\xi_n\,\log r} \;\le\; \frac{2\eta_{\mathfrak{P}}}{(y_\star-\eta_{\mathfrak{P}})\,\log r}.$$

On certified masks the *channel floors* (e.g. $s_{\min},\kappa_{\min}$ in Section 7.9.16) ensure $y_\star \ge c_{\text{floor}}\,\eta_{\mathfrak{P}}$ with $c_{\text{floor}}>1$; hence

$$S_{\max} \;\le\; \frac{2}{(c_{\text{floor}}-1)\,\log r}\;\eta_{\mathfrak{P}}. \tag{7.700}$$

If, in addition, $S_{\max}\le\sigma_{\text{tol}}$ and the constant model satisfies (7.699), then the law channel exhibits *no IR improvement* beyond plateau oscillations; by Definition 6.300 this certifies QIV on the window. Since the bounds and the mask are uniform on $\mathfrak{P}$, the acceptance is uniform on $\mathfrak{P}$.

(2) QIII reject. Suppose $y(\ell)=A\,\ell^{\sigma}(1+\varepsilon(\ell))$ with $|\varepsilon(\ell)|\ll 1$ and $\sigma\neq 0$. Then $\log y_n = \log A+\sigma\log\ell_n+\eta_n$ with $\eta_n=\log(1+\varepsilon(\ell_n))=\mathcal{O}(\varepsilon)$. Weighted least squares of $\log y_n$ on $\log\ell_n$ yields a slope $\widehat{\sigma}(x)$ with asymptotic normal CI. If the 95% CI excludes 0, we have statistically significant $\sigma\neq 0$.

To guard against overfitting plateau oscillations, compare the *power–law* (1–slope) model to the *constant* (0–slope) model via an F–test / likelihood–ratio:

$$F \;=\; \frac{\big(\mathrm{RSS}_0-\mathrm{RSS}_1\big)/1}{\mathrm{RSS}_1/(N-2)}.$$

Under the null (constant up to (7.699)) and well–specified WLS, F has approximately an $F_{1,N-2}$ law; more importantly, the *maximal* reduction in RSS achievable by fitting a spurious slope is $O(\eta_{\mathfrak{P}}^2)$. Thus, if the observed RSS reduction exceeds the *plateau tolerance budget* (a multiple of $\eta_{\mathfrak{P}}^2$ fixed by certification), the power–law fit is genuinely preferred and we reject QIV in favor of QIII on that mask. Both the CI decision and the F–test are invariant under the BRST/G structure by Lemma 7.552 and Proposition 7.556.

Uniformity on $\mathfrak{P}$. Apply Lemma 7.579 to the map $x\mapsto S_{\max}(x)$ and to the residual series to extend the decision uniformly over $x\in\mathfrak{P}$, with at most an $\eta_{\mathfrak{P}}$ slack, which is already absorbed into the tolerance budget.

This proves both statements. □

Remark 7.600 (Practical choices)**.** *Grid.* With a geometric IR grid $\ell_n=\ell_0 r^{\,n}$, take $r\in[1.4,2]$ and $N\ge 5$. Then the log–span is $\log(\ell_{N-1}/\ell_0)=(N-1)\log r\gtrsim 4\log 1.4\approx 1.35$, which stabilizes slope estimates and tightens the bound (7.700).

Tolerance. Set $\sigma_{\text{tol}}=c_s\,\eta_{\mathfrak{P}}$ with $c_s\in[5,10]$. Together with (7.700) and the channel floor $c_{\text{floor}}>1$, this makes false positives from plateau oscillations unlikely; e.g. for $r=1.6$ and $c_{\text{floor}}\ge 3$, the bound gives $S_{\max}\lesssim 1.6\,\eta_{\mathfrak{P}}$.

Reporting. Report $\big(\widehat{S}(x),\,\mathrm{CI}_{95\%}\big)$ per band point x (WLS slope on $(\log\ell_n,\log y_n)$) and the aggregate $S_{\max}$ over $x\in\mathfrak{P}$. Use the CI *and* the model–comparison test as in item (2).

These settings are BRST/Nielsen stable on $\mathcal{H}_{\text{phys}}$ (Lemma 7.552 and Proposition 7.556) and compatible with certified–plateau masking.

Definition 7.601 (Calibration targets)**.** At $x \in \mathfrak{P}$, the tomographic constants are:

$$\kappa_{\mathrm{spec}}(x) := \inf_{|\omega| \le \omega_0(x)} \mathsf{K}_{\mathrm{spec}}(x;\omega),$$

where $c_0(x)$ is the proportionality constant in the relation (Lemma 7.608):

$$\mathcal{T}_a^{(x)}[\chi_{\omega_0(x)}] = c_0(x)\,|\mathcal{R}_a^{(x)}| + \mathcal{O}(\omega_0^2).$$

Here $\mathsf{K}_{\mathrm{spec}}$ is the positive tomography kernel (Definition 7.607), $\omega_0(x)$ is the bandwidth chosen by (7.736).

Proposition 7.602 (Kernel floor from taggers (BRST/Nielsen safe))**.** *Let $\mathsf{K}_{\mathrm{spec}}(x;\omega)$ be computed from the UV/IR taggers (Definition 7.551) via the spectral representation of the law branch (Definition 7.607). Since $\mathsf{K}_{\mathrm{spec}}(x;\omega) \ge 0$ and is continuous in ω on compact sets,*

$$\kappa_{\mathrm{spec}}(x) \;=\; \min_{m\in\{-M,\dots,M\}} \mathsf{K}_{\mathrm{spec}}(x;\omega_m) \;-\; \delta_{\mathrm{disc}}(x),$$

with a uniform grid $\omega_m \in [-\omega_0(x),\omega_0(x)]$ and a discretization slack $\delta_{\mathrm{disc}}(x) = \mathcal{O}(\omega_0(x)/M)$ chosen so that $\kappa_{\mathrm{spec}}(x) > 0$ remains valid as a lower confidence bound. This construction commutes with BRST/G actions and is gauge–parameter independent on $\mathcal{H}_{\mathrm{phys}}$ (Proposition 7.556 and Lemma 7.552).

Proof. (A) Positivity and continuity from the taggers. By Definition 7.551, the IR tagger is a C^∞ c–number kernel $\kappa_x(t,\mathbf{r}) = \chi_\ell(t)\,\phi_\ell(\mathbf{r})$ with $\chi_\ell, \phi_\ell \in C_c^\infty$ real and even. Hence

$$\widehat{\kappa}_x(\omega,\mathbf{k}) = \widehat{\chi}_\ell(\omega)\,\widehat{\phi}_\ell(\mathbf{k})$$

is C^∞ in $(\omega,\mathbf{k})$ and real–even in ω (Paley–Wiener). The tomography kernel of Definition 7.607 reads

$$\mathsf{K}_{\mathrm{spec}}(x;\omega) = \int_{\mathbb{R}^{d-1}} W_x(\mathbf{k})\,\big|\widehat{\kappa}_x(\omega,\mathbf{k})\big|^2\,d\mathbf{k} = \big|\widehat{\chi}_\ell(\omega)\big|^2\,C_x, \qquad C_x := \int W_x(\mathbf{k})\,|\widehat{\phi}_\ell(\mathbf{k})|^2\,d\mathbf{k} \ge 0, \tag{7.701}$$

with $W_x(\mathbf{k}) \ge 0$ the channel weight. On certified plateaus W_x has positive mass on some compact set, while $\widehat{\phi}_\ell \not\equiv 0$, hence $C_x > 0$. Therefore $\mathsf{K}_{\mathrm{spec}} \ge 0$ and is continuous (indeed C^∞) in ω on any compact interval. In particular, by Lemma 7.575 there exists $\omega_0(x) > 0$ such that $\inf_{|\omega|\le\omega_0(x)} \mathsf{K}_{\mathrm{spec}}(x;\omega) > 0$.

(B) A Lipschitz bound on $[-\omega_0,\omega_0]$. Differentiating (7.701) gives

$$\partial_\omega \mathsf{K}_{\mathrm{spec}}(x;\omega) = 2\,\mathrm{Re}\Big(\widehat{\chi}_\ell'(\omega)\,\overline{\widehat{\chi}_\ell(\omega)}\Big)\,C_x,$$

hence on the compact interval $[-\omega_0,\omega_0]$,

$$\big|\partial_\omega \mathsf{K}_{\mathrm{spec}}(x;\omega)\big| \;\le\; L_x \;:=\; 2\,C_x \sup_{|\omega|\le\omega_0}\big|\widehat{\chi}_\ell'(\omega)\big| \sup_{|\omega|\le\omega_0}\big|\widehat{\chi}_\ell(\omega)\big| \;<\; \infty. \tag{7.702}$$

Thus $\mathsf{K}_{\mathrm{spec}}(x;\cdot)$ is Lipschitz on $[-\omega_0,\omega_0]$ with constant L_x.

(C) Discretization slack on a uniform grid. Let the uniform grid have spacing $\Delta\omega := 2\omega_0/M$ and nodes $\omega_m = -\omega_0 + m\Delta\omega$, $m = -M, \dots, M$. For any $\omega \in [-\omega_0, \omega_0]$ choose the nearest gridpoint $\omega_{m(\omega)}$ so that $|\omega - \omega_{m(\omega)}| \le \Delta\omega/2$. By the mean–value inequality and (7.702),

$$\mathsf{K}_{\mathrm{spec}}(x;\omega) \;\ge\; \mathsf{K}_{\mathrm{spec}}(x;\omega_{m(\omega)}) - L_x\,|\omega - \omega_{m(\omega)}| \;\ge\; \min_m \mathsf{K}_{\mathrm{spec}}(x;\omega_m) - \frac{L_x}{2}\,\Delta\omega.$$

Taking the infimum over $\omega \in [-\omega_0, \omega_0]$,

$$\inf_{|\omega|\le\omega_0} \mathsf{K}_{\mathrm{spec}}(x;\omega) \;\ge\; \min_m \mathsf{K}_{\mathrm{spec}}(x;\omega_m) - \delta_{\mathrm{disc}}(x), \qquad \delta_{\mathrm{disc}}(x) := \frac{L_x}{2}\,\Delta\omega = \mathcal{O}\!\left(\frac{\omega_0}{M}\right).$$

Defining

$$\kappa_{\mathrm{spec}}(x) \;:=\; \min_m \mathsf{K}_{\mathrm{spec}}(x;\omega_m) - \delta_{\mathrm{disc}}(x),$$

we obtain a *rigorous lower bound* $\kappa_{\mathrm{spec}}(x) \le \inf_{|\omega|\le\omega_0} \mathsf{K}_{\mathrm{spec}}(x;\omega)$. By choosing M large enough (or, equivalently, $\Delta\omega$ small enough) and using the positive floor guaranteed by Lemma 7.575, one can ensure $\kappa_{\mathrm{spec}}(x) > 0$. This realizes the statement with $\delta_{\mathrm{disc}}(x) = \mathcal{O}(\omega_0/M)$ and certifies $\kappa_{\mathrm{spec}}(x)$ as a *lower confidence bound* against discretization error.

(D) BRST/G commutation and gauge independence. The taggers $K_{\mathrm{IR}}^{(\ell)}$ and $P_x(\lambda_{\mathrm{UV}})$ act by c–number convolution/multiplication and commute with BRST and G actions (Proposition 7.573). The kernel $\mathsf{K}_{\mathrm{spec}}$ is a positive quadratic functional of $\widehat{\kappa}_x$ (cf. (7.701)), hence it inherits this commutation and is built from BRST-closed, gauge–invariant data on $\mathcal{H}_{\mathrm{phys}}$. Therefore $\kappa_{\mathrm{spec}}(x)$ constructed above is gauge–parameter independent (Lemma 7.552 and Proposition 7.556).

Combining (A)–(D) proves the proposition. □

Definition 7.603 (Probe normalization $c_0(x)$ via Ward matching)**.** At fixed x, select a small spurion amplitude window $\epsilon \in (0, \epsilon_{\max}]$ within the linear regime. For each setting, measure the Ward source magnitude $|\mathcal{R}_a^{(x)}(\epsilon)|$ and the tomographic functional $\mathcal{T}_a^{(x)}[\chi_{\omega_0(x)}](\epsilon)$ (same $\omega_0(x)$ as in Definition 7.636). Fit the constrained line through the origin:

$$\widehat{c}_0(x) \;:=\; \arg\min_{c\ge 0} \sum_n w_n \left(\mathcal{T}_a^{(x)}[\chi_{\omega_0(x)}](\epsilon_n) - c\,|\mathcal{R}_a^{(x)}(\epsilon_n)|\right)^2,$$

with weights w_n from measurement uncertainties. Reject points with relative $|\epsilon|$ beyond the GMOR linear window (per Definition 7.634); report $\widehat{c}_0(x)$ and its CI.

Lemma 7.604 (Uniform control with the bandwidth schedule)**.** *With the schedule $\omega_0(x)$ of Definition 7.636, the model error in the regression of Definition 7.603 is $\mathcal{O}(\theta^2)$ uniformly on $\mathfrak{P}$ (Lemma 7.637); the estimate $\widehat{c}_0(x)$ is thus plateau–consistent and Nielsen–independent.*

Proof. Step 1: Low–frequency expansion and evenness. By the spectral construction of the law branch (Definition 7.607) and the tagger regularity (Definition 7.551), the descended T–odd readout admits, for $|\omega| \le \omega_0(x)$,

$$Y_x(\omega) \;:=\; 1 - |\Lambda_{\mathrm{law}}^{(\mathrm{spec};x)}(\omega)| \;=\; c_0(x)\,\mathsf{K}_{\mathrm{spec}}(x;\omega)\,|\mathcal{R}_a^{(x)}| \;+\; R_x(\omega), \tag{7.703}$$

where $\mathsf{K}_{\mathrm{spec}}(x;\omega)$ is C^∞ and even in ω (cf. Lemma 7.575), and the remainder R_x is even and satisfies $R_x(\omega) = \mathcal{O}(\omega^2)$ as $\omega \to 0$. The evenness and smoothness follow from the real–even IR time window and the T–odd projection being linear in the spectrum; the ω^2 order is the first nontrivial even correction in the Taylor series at $\omega = 0$.

Step 2: Bandwidth schedule and uniform bound. By Definition 7.636, the bandwidth is scheduled as $\omega_0(x) \le c_*\,\theta$ with c_* independent of $x \in \mathfrak{P}$. Hence

$$\sup_{|\omega|\le\omega_0(x)} |R_x(\omega)| \;\le\; C_1(x)\,\omega_0(x)^2 \;\le\; C_1(x)\,c_*^2\,\theta^2.$$

On certified plateaus (Definition 6.300) the derivatives of $\mathsf{K}_{\mathrm{spec}}$ and the channel weights are uniformly bounded, so $C_1(x)$ is bounded on $\mathfrak{P}$; write $\sup_{x\in\mathfrak{P}} C_1(x) =: C_1 < \infty$. Therefore

$$\sup_{x\in\mathfrak{P}}\ \sup_{|\omega|\le\omega_0(x)} |R_x(\omega)| \;\le\; C_1\,\theta^2. \tag{7.704}$$

Step 3: WLS normal equation and bias of $\widehat{c}_0(x)$. The regression of Definition 7.603 fits $Y_x(\omega_m)$ (on a grid $|\omega_m| \le \omega_0(x)$) to the one–parameter model $c\,\mathsf{K}_{\mathrm{spec}}(x;\omega_m)\,|\mathcal{R}_a^{(x)}|$ with weights w_m. Let $K_m := \mathsf{K}_{\mathrm{spec}}(x;\omega_m)\,|\mathcal{R}_a^{(x)}|$ and $\varepsilon_m := R_x(\omega_m)$. The WLS estimator solves the normal equation

$$\widehat{c}_0(x) \;=\; c_0(x) \;+\; \frac{\sum_m w_m K_m\,\varepsilon_m}{\sum_m w_m K_m^2}.$$

On the scheduled window, Lemma 7.575 and Proposition 7.602 give $K_m \ge \kappa_{\mathrm{spec}}(x)\,|\mathcal{R}_a^{(x)}|$, hence

$$\left|\widehat{c}_0(x) - c_0(x)\right| \;\le\; \frac{\sum_m w_m K_m\,|\varepsilon_m|}{\sum_m w_m K_m^2} \;\le\; \frac{\max_m |\varepsilon_m|}{\kappa_{\mathrm{spec}}(x)\,|\mathcal{R}_a^{(x)}|}\;\cdot\;\frac{\sum_m w_m K_m}{\sum_m w_m K_m^2} \;\le\; \frac{\max_m |\varepsilon_m|}{\kappa_{\mathrm{spec}}(x)\,|\mathcal{R}_a^{(x)}|}.$$

Using (7.704) and the plateau floor $\inf_{x\in\mathfrak{P}} \kappa_{\mathrm{spec}}(x) > 0$ (cf. Lemma 7.575 and Proposition 7.602) and $|\mathcal{R}_a^{(x)}|$ bounded away from 0 on the certified mask, we obtain

$$\sup_{x\in\mathfrak{P}} \left|\widehat{c}_0(x) - c_0(x)\right| \;\le\; C_2\,\theta^2,$$

for some $C_2 < \infty$ independent of x. This is precisely the $\mathcal{O}(\theta^2)$ model–error control of Lemma 7.637.

Step 4: Plateau consistency and Nielsen/BRST independence. The bound is uniform on $\mathfrak{P}$; hence $\widehat{c}_0(x) \to c_0(x)$ as $\theta \to 0$ uniformly, i.e. plateau–consistency. Since the UV/IR taggers commute with BRST and G actions (Proposition 7.573), and all regressors are c–number functionals of band–descended observables on $\mathcal{H}_{\mathrm{phys}}$, the estimate is gauge–parameter independent (Lemma 7.552 and Proposition 7.556).

This proves the lemma. □

Corollary 7.605 (Certified tomographic lower bound)**.** *For all $x \in \mathfrak{P}$,*

$$1 - |\Lambda_{\mathrm{law}}^{(x)}| \;\ge\; \underbrace{\kappa_{\mathrm{spec}}(x)\,\widehat{c}_0(x)}_{=:K_{\mathrm{low}}(x)}\;|\mathcal{R}_a^{(x)}| \;-\; C\,\theta^2 \;-\; O(\eta_{\mathfrak{P}}),$$

with C gauge–independent. Hence the measured T–odd signal tracks the Ward source with a calibrated *constant $K_{\mathrm{low}}(x)$, uniform on the mask (Proposition 7.609 and Lemma 7.608).*

Proof. Step 1: Pointwise lower bound for the spectral window. From (7.703) (used in Lemma 7.604) and Lemma 7.575, for $|\omega| \le \omega_0(x)$, we have

$$Y_x(\omega) \ge c_0(x)\,\kappa_{\mathrm{spec}}(x)\,|\mathcal{R}_a^{(x)}| - |R_x(\omega)|.$$

Taking the infimum over $\{|\omega| \le \omega_0(x)\}$ and using the uniform bound $\sup_{|\omega|\le\omega_0(x)}|R_x(\omega)| \le C_1\theta^2$ gives

$$\inf_{|\omega|\le\omega_0(x)} Y_x(\omega) \;\ge\; c_0(x)\,\kappa_{\mathrm{spec}}(x)\,|\mathcal{R}_a^{(x)}| - C_1\,\theta^2. \tag{7.705}$$

Step 2: Replace $c_0(x)$ by $\widehat{c}_0(x)$. By Lemma 7.604, $|\widehat{c}_0(x) - c_0(x)| \le C_2\theta^2$ uniformly on $\mathfrak{P}$. Hence $c_0(x) \ge \widehat{c}_0(x) - C_2\theta^2$, and plugging into eq. (7.705) yields

$$\inf_{|\omega|\le\omega_0(x)} Y_x(\omega) \;\ge\; \kappa_{\mathrm{spec}}(x)\,\widehat{c}_0(x)\,|\mathcal{R}_a^{(x)}| \;-\; \Big(C_1 + C_2\,\kappa_{\mathrm{spec}}(x)\,|\mathcal{R}_a^{(x)}|\Big)\,\theta^2.$$

Absorb the bracket into a gauge–independent constant C (uniform on $\mathfrak{P}$ by plateau regularity and the tagger floors).

Step 3: From window infimum to the reported statistic. By the construction of the tomographic readout (Definition 7.607) and the certified–plateau masking, the reported $1 - |\Lambda_{\mathrm{law}}^{(x)}|$ differs from the window infimum by at most the oscillation budget $\mathcal{O}(\eta_{\mathfrak{P}})$ (plateau envelope). Therefore

$$1 - |\Lambda_{\mathrm{law}}^{(x)}| \;\ge\; \kappa_{\mathrm{spec}}(x)\,\widehat{c}_0(x)\,|\mathcal{R}_a^{(x)}| \;-\; C\,\theta^2 \;-\; \mathcal{O}(\eta_{\mathfrak{P}}).$$

This is the stated lower bound with the calibrated constant $K_{\mathrm{low}}(x) := \kappa_{\mathrm{spec}}(x)\,\widehat{c}_0(x)$.

Step 4: Uniformity and Nielsen–independence. Uniformity on $\mathfrak{P}$ follows from the uniform bounds in Lemmas 7.575 and 7.604 and Proposition 7.602 and plateau compactness. BRST/Nielsen–safety is inherited from the commutation of the taggers with BRST/G and the use of band–descended observables (Propositions 7.556 and 7.573 and Lemma 7.552). This aligns with the tracking statements of Proposition 7.609 and Lemma 7.608. □

Definition 7.606 (Reporting for the law–visibility curve)**.** On $\mathfrak{P}$ (or $\mathfrak{M}_{\mathrm{mask}}$):

1. Report the log–slope statistics $\widehat{S}(x)$ with CIs and $S_{\max}$; include the constant vs. power–law model selection (AIC/BIC or F–test) and decision (QIV accept / QIII reject) per Proposition 7.599.
2. Provide $\kappa_{\mathrm{spec}}(x)$ (grid definition, M, and δ_{disc}), $\widehat{c}_0(x)$ with CI, and the product $K_{\mathrm{low}}(x)$ used in the tomographic bound.
3. State the bandwidth $\omega_0(x)$ and the global θ; confirm the $\mathcal{O}(\theta^2)$ control.
4. Confirm gauge–flatness (Nielsen slopes ≈ 0) for all reported quantities (Proposition 7.556).

Only quote inside $\mathfrak{M}_{\mathrm{mask}}$; if the slope test fails on a subwindow, refine the mask.

7.9.8 T-odd Tomography & Cones (No Law-signalling; Anomaly Filters)

Definition 7.607 (T-odd spectral density and tomography kernel)**.** At a certified band point $x = (\lambda_{\mathrm{UV}}, \ell) \in \mathfrak{P}$ (Definition 6.300), define the *T-odd spectral density* for a broken generator a on $\mathcal{H}_{\mathrm{phys}}$ by the retarded two-point function of the adjoint-covariant divergence:

$$\mathsf{S}_a(x;\omega) \;:=\; \frac{1}{\pi}\,\mathrm{Im}\int d^d t\, e^{i\omega t}\,\theta(t)\,\Big\langle\,\Big[\,\nabla^{(\mathrm{ad})}_{\mu}\bar{J}^{(x)\mu}_{a,\mathrm{imp}}(t),\,\mathcal{W}^{(x)}(0)\Big]\,\Big\rangle_{\mathbf{\Psi}},$$

where $\mathcal{W}^{(x)}$ is any admissible local probe density that generates the holosymmetric law visibility (Definition 6.275) at x. The *tomography kernel* is the positive function $\mathsf{K}_{\mathrm{spec}}(x;\omega)$ (dependent only on the UV/IR taggers, Definition 7.551) appearing in the spectral representation

$$-\,\mathrm{Re}\;\Lambda^{(\mathrm{spec};x)}_{\mathrm{law}} \;=\; \int_{\mathbb{R}} d\omega\,\mathsf{K}_{\mathrm{spec}}(x;\omega)\,\mathsf{S}_a(x;\omega), \tag{7.706}$$

as established in the proof of Lemma 7.578. Define the *tomography functional* $\mathcal{T}^{(x)}_a[\varphi] := \int d\omega\,\varphi(\omega)\,\mathsf{S}_a(x;\omega)$ for test functions $\varphi \in C^\infty_c(\mathbb{R})$.

Lemma 7.608 (Ward-source tracking at low frequency)**.** *Let $\chi_\tau(\omega)$ be a normalized, even mollifier supported in $[-\omega_0(\tau),\omega_0(\tau)]$ with $\omega_0(\tau)\downarrow 0$ as $\tau\to\infty$. Then, on $\mathcal{H}_{\mathrm{phys}}$,*

$$\lim_{\tau\to\infty}\;\mathcal{T}^{(x)}_a[\chi_\tau] \;=\; c_0(x)\,\Big|\mathcal{R}^{(x)}_a\Big|, \qquad \mathcal{R}^{(x)}_a := \sum_k \epsilon_k\Big\langle \mathcal{S}^{(\lambda_{\mathrm{UV}})}_{k,a}\Big\rangle_{\mathbf{\Psi}},$$

with $c_0(x) > 0$ a normalization fixed by the choice of probe $\mathcal{W}^{(x)}$ and the taggers. The limit is gauge–parameter independent (Proposition 7.556) and BRST–safe (Lemma 7.552).

Proof. Step 1: Tomographic functional and low–frequency model. Denote the spectral (T–odd) law–channel readout by

$$Y_x(\omega) \;:=\; 1 - \big|\Lambda^{(\mathrm{spec};x)}_{\mathrm{law}}(\omega)\big|\,,$$

and the IR/UV tagger kernel by $\mathsf{K}_{\mathrm{spec}}(x;\omega)$ (Definitions 7.551 and 7.607). Near $\omega = 0$, the band–descended Ward identity (Fourier transform of eq. (7.665)) together with the PCAC/LSZ matching gives the even Taylor expansion

$$Y_x(\omega) \;=\; c_0(x)\,\mathsf{K}_{\mathrm{spec}}(x;\omega)\,|\mathcal{R}^{(x)}_a| \;+\; R_x(\omega), \qquad R_x(\omega) = \mathcal{O}(\omega^2), \tag{7.707}$$

where the constant $c_0(x) > 0$ depends only on the fixed probe $\mathcal{W}^{(x)}$ and the taggers. Evenness of the expansion follows from time–reversal on the T–odd channel combined with the even tagger window; the ω^2 remainder is the first nontrivial even correction.

We define the *tomographic functional* that extracts the zero–frequency coefficient as

$$\mathcal{T}^{(x)}_a[\chi] \;:=\; \int_{-\infty}^{\infty} d\omega\;\chi(\omega)\,\frac{Y_x(\omega)}{\mathsf{K}_{\mathrm{spec}}(x;\omega)}\,. \tag{7.708}$$

This is the continuum counterpart of the one–parameter regression in Definition 7.603: fitting $Y_x(\omega)$ against $\mathsf{K}_{\mathrm{spec}}(x;\omega)$ on a small window is a Riemann–sum discretization of (7.708) with χ proportional to the sampling weights.

Step 2: Kernel floor and regularity on shrinking windows. By Lemma 7.575 and Proposition 7.602 there exists $\omega_0 > 0$ such that

$$\mathsf{K}_{\mathrm{spec}}(x;\omega) \;\geq\; \kappa_{\mathrm{spec}}(x) \;>\; 0 \quad \text{for all } |\omega| \leq \omega_0,$$

and $\mathsf{K}_{\mathrm{spec}}(x;\omega)$ is C^∞ in ω. Hence $1/\mathsf{K}_{\mathrm{spec}}(x;\omega)$ is bounded and smooth on $[-\omega_0,\omega_0]$. Because χ_τ has support in $[-\omega_0(\tau),\omega_0(\tau)]$ with $\omega_0(\tau) \downarrow 0$, for all large τ the support lies inside $[-\omega_0,\omega_0]$.

Step 3: Mollifier limit. Insert (7.707) into (7.708):

$$\mathcal{T}_a^{(x)}[\chi_\tau] \;=\; c_0(x)\,|\mathcal{R}_a^{(x)}|\,\underbrace{\int \chi_\tau(\omega)\,d\omega}_{=1} \;+\; \int \chi_\tau(\omega)\,\frac{R_x(\omega)}{\mathsf{K}_{\mathrm{spec}}(x;\omega)}\,d\omega.$$

The first term gives $c_0(x)|\mathcal{R}_a^{(x)}|$ by normalization of χ_τ. For the second term, use the expansion $R_x(\omega) = \mathcal{O}(\omega^2)$ and the kernel floor to bound

$$\left|\int \chi_\tau(\omega)\,\frac{R_x(\omega)}{\mathsf{K}_{\mathrm{spec}}(x;\omega)}\,d\omega\right| \;\leq\; \frac{1}{\kappa_{\mathrm{spec}}(x)}\int \chi_\tau(\omega)\,C_x\,\omega^2\,d\omega \;\leq\; \frac{C_x}{\kappa_{\mathrm{spec}}(x)}\sup_{|\omega|\leq\omega_0(\tau)}\omega^2 \;\xrightarrow[\tau\to\infty]{}\; 0,$$

for some $C_x < \infty$ (uniform on certified plateaus). Thus

$$\lim_{\tau\to\infty}\;\mathcal{T}_a^{(x)}[\chi_\tau] \;=\; c_0(x)\,|\mathcal{R}_a^{(x)}|.$$

The limit is an instance of dominated (indeed uniform) convergence, using the bounded denominator and compact support.

Step 4: Positivity and BRST/Nielsen safety. Positivity $c_0(x) > 0$ follows from the positivity of the spectral measure in the retarded law–channel (Källén–Lehmann) and the nonnegativity of $\mathsf{K}_{\mathrm{spec}}$ (Lemma 7.575). Gauge–parameter independence and BRST safety hold because the UV/IR taggers commute with BRST and G (Proposition 7.573), and expectations are taken on $\mathcal{H}_{\mathrm{phys}}$ (Proposition 7.556 and Lemma 7.552); BRST–exact and Noether–improvement pieces do not contribute.

Combining the steps proves the claim. □

Proposition 7.609 (Tomographic lower bound for the law-deficit)**.** *On any $x \in \mathfrak{P}$,*

$$1 - \left|\Lambda_{\mathrm{law}}^{(x)}\right| \;\geq\; \kappa_{\mathrm{spec}}(x)\;\mathcal{T}_a^{(x)}[\chi_\tau] \;-\; O(\eta_{\mathfrak{P}}) \;-\; o_\tau(1),$$

where $\kappa_{\mathrm{spec}}(x) > 0$ is the kernel floor on $[-\omega_0,\omega_0]$ (cf. Lemma 7.575 and Proposition 7.602), $\eta_{\mathfrak{P}}$ is the plateau tolerance (Definition 6.300), and $o_\tau(1) \to 0$ as $\tau \to \infty$.

Proof. Step 1: Spectral lower bound on the law channel. By the channel-resolved signature (Corollary 7.677(i)),

$$1 - \left|\Lambda_{\mathrm{law}}^{(x)}\right| \;\geq\; -\,\mathrm{Re}\;\Lambda_{\mathrm{law}}^{(\mathrm{spec};x)} \;-\; O(\eta_{\mathfrak{P}}).$$

By the spectral/tomographic representation (Definition 7.607), there exists a nonnegative spectral density $S_a^{(x)}(\omega) \geq 0$ (the T–odd weight) such that

$$-\,\mathrm{Re}\;\Lambda_{\mathrm{law}}^{(\mathrm{spec};x)} \;=\; \int_{\mathbb{R}} d\omega\;\mathsf{K}_{\mathrm{spec}}(x;\omega)\,S_a^{(x)}(\omega),$$

with $\mathsf{K}_{\mathrm{spec}}(x;\omega) \geq 0$ continuous in ω and, on $|\omega| \leq \omega_0$, bounded below by the positive floor $\kappa_{\mathrm{spec}}(x)$ (Lemma 7.575 and Proposition 7.602). Restricting to the low–frequency window,

$$1 - \left|\Lambda_{\mathrm{law}}^{(x)}\right| \;\geq\; \int_{|\omega|\leq\omega_0} \mathsf{K}_{\mathrm{spec}}(x;\omega)\, S_a^{(x)}(\omega)\, d\omega \;-\; O(\eta_{\mathfrak{P}}). \tag{7.709}$$

Step 2: Relate the spectral weight to the measured $Y_x(\omega)$. Write

$$Y_x(\omega) \;:=\; 1 - \left|\Lambda_{\mathrm{law}}^{(\mathrm{spec};x)}(\omega)\right|.$$

Near $\omega = 0$, smoothness of the taggers and the Ward projection imply the even expansion

$$Y_x(\omega) \;=\; \mathsf{K}_{\mathrm{spec}}(x;\omega)\, S_a^{(x)}(\omega) \;+\; R_x(\omega), \qquad R_x(\omega) = \mathcal{O}(\omega^2) \quad (\omega \to 0), \tag{7.710}$$

uniformly on certified plateaus (bounded derivatives of the taggers and channel weights).

Step 3: Insert a normalized mollifier and use positivity. Let χ_τ be even, supported in $[-\omega_0(\tau), \omega_0(\tau)]$, and normalized: $\int \chi_\tau = 1$, with $\omega_0(\tau) \downarrow 0$. Multiply (7.710) by $\chi_\tau(\omega)$ and integrate over $\mathbb{R}$ (effectively over the support):

$$\int \chi_\tau(\omega)\, Y_x(\omega)\, d\omega = \int \chi_\tau(\omega)\, \mathsf{K}_{\mathrm{spec}}(x;\omega)\, S_a^{(x)}(\omega)\, d\omega \;+\; \int \chi_\tau(\omega)\, R_x(\omega)\, d\omega.$$

Since $\mathsf{K}_{\mathrm{spec}}(x;\omega) \geq \kappa_{\mathrm{spec}}(x)$ on the support of χ_τ,

$$\int \chi_\tau\, \mathsf{K}_{\mathrm{spec}}\, S_a^{(x)} \;\geq\; \kappa_{\mathrm{spec}}(x) \int \chi_\tau\, S_a^{(x)}.$$

By the definition of the tomographic functional (eq. (7.708) in the proof of Lemma 7.608),

$$\mathcal{T}_a^{(x)}[\chi_\tau] \;=\; \int \chi_\tau(\omega)\, \frac{Y_x(\omega)}{\mathsf{K}_{\mathrm{spec}}(x;\omega)}\, d\omega,$$

and, using again $\mathsf{K}_{\mathrm{spec}} \geq \kappa_{\mathrm{spec}}$ on the support,

$$\int \chi_\tau(\omega)\, Y_x(\omega)\, d\omega = \int \chi_\tau\, \mathsf{K}_{\mathrm{spec}} \cdot \frac{Y_x}{\mathsf{K}_{\mathrm{spec}}} \;\geq\; \kappa_{\mathrm{spec}}(x)\, \mathcal{T}_a^{(x)}[\chi_\tau].$$

Combining with the remainder gives

$$\int_{|\omega|\leq\omega_0(\tau)} Y_x(\omega)\, \chi_\tau(\omega)\, d\omega \;\geq\; \kappa_{\mathrm{spec}}(x)\, \mathcal{T}_a^{(x)}[\chi_\tau] \;-\; \left|\int \chi_\tau(\omega)\, R_x(\omega)\, d\omega\right|. \tag{7.711}$$

Step 4: Link the windowed spectral mass to the mollified average. Because $S_a^{(x)} \geq 0$ and $\mathsf{K}_{\mathrm{spec}} \geq 0$, from (7.709),

$$1 - \left|\Lambda_{\mathrm{law}}^{(x)}\right| \;\geq\; \int_{|\omega|\leq\omega_0(\tau)} \mathsf{K}_{\mathrm{spec}}\, S_a^{(x)}\, d\omega \;-\; O(\eta_{\mathfrak{P}}) \;\geq\; \int \chi_\tau\, \mathsf{K}_{\mathrm{spec}}\, S_a^{(x)}\, d\omega \;-\; O(\eta_{\mathfrak{P}}).$$

Using (7.710),

$$\int \chi_\tau\, \mathsf{K}_{\mathrm{spec}}\, S_a^{(x)} = \int \chi_\tau\, Y_x \;-\; \int \chi_\tau\, R_x.$$

Plug this and (7.711) into the previous inequality to obtain

$$1 - \left|\Lambda^{(x)}_{\mathrm{law}}\right| \;\geq\; \kappa_{\mathrm{spec}}(x)\,\mathcal{T}^{(x)}_{a}[\chi_\tau] \;-\; O(\eta_{\mathfrak{P}}) \;-\left|\int \chi_\tau R_x\right|.$$

Step 5: Control of the remainder $o_\tau(1)$. By (7.710), $|R_x(\omega)| \leq C_x\,\omega^2$ for $|\omega| \leq \omega_0$, with C_x bounded uniformly on $\mathfrak{P}$ (plateau regularity). Hence

$$\left|\int \chi_\tau(\omega)\,R_x(\omega)\,d\omega\right| \;\leq\; C_x \sup_{|\omega|\leq\omega_0(\tau)} \omega^2 \;=:\; o_\tau(1) \;\xrightarrow[\tau\to\infty]{}\; 0.$$

Collecting terms yields the claimed bound. □

Corollary 7.610 (Quantitative tracking of the Ward-source residual)**.** *Combining Lemma 7.608 and Proposition 7.609, for large τ and within plateau tolerances,*

$$1 - \left|\Lambda^{(x)}_{\mathrm{law}}\right| \;\gtrsim\; \kappa_{\mathrm{spec}}(x)\,c_0(x)\left|\mathcal{R}^{(x)}_{a}\right|.$$

Thus the measured T–odd holosymmetric signal tracks the Ward–source residual linearly at low frequency.

Proof. From Proposition 7.609,

$$1 - \left|\Lambda^{(x)}_{\mathrm{law}}\right| \;\geq\; \kappa_{\mathrm{spec}}(x)\;\mathcal{T}^{(x)}_{a}[\chi_\tau] \;-\; O(\eta_{\mathfrak{P}}) \;-\; o_\tau(1).$$

By Lemma 7.608, $\mathcal{T}^{(x)}_{a}[\chi_\tau] \to c_0(x)\,|\mathcal{R}^{(x)}_{a}|$ as $\tau \to \infty$, with $c_0(x) > 0$. Therefore, taking τ large and using the plateau envelope to absorb $O(\eta_{\mathfrak{P}})$,

$$1 - \left|\Lambda^{(x)}_{\mathrm{law}}\right| \;\geq\; \kappa_{\mathrm{spec}}(x)\,c_0(x)\,|\mathcal{R}^{(x)}_{a}| \;-\; \underbrace{O(\eta_{\mathfrak{P}}) - o_\tau(1)}_{\text{small}},$$

which is precisely the stated quantitative tracking; the symbol $\gtrsim$ indicates dominance up to these controlled small terms. □

Definition 7.611 (Cone preservation (no law-signalling))**.** Let $\mathsf{C}^{\pm}$ be the forward/backward light-cones (or causal cones with dynamic exponent z). The UV/IR taggers are *cone-preserving* if their time-domain kernels have support contained in $\mathsf{C}^{+} \cup \mathsf{C}^{-}$ and commute with the BRST/G actions (Definition 7.551). We say the law-channel readout is *non-signalling* if, for spacelike-separated regions A, B,

$$\left\langle [\,\bar{J}^{(x)\mu}_{a,\mathrm{imp}}(A),\; \mathcal{W}^{(x)}(B)\,] \right\rangle_{\Psi} = 0,$$

for all $x \in \mathfrak{P}$ (microcausality after band descent).

Proposition 7.612 (Cone preservation under band descent)**.** *If the microscopic theory is microcausal and the taggers are cone-preserving, then the band–retarded correlators vanish outside $\mathsf{C}^{\pm}$, and the law channel is non–signalling in the sense of Definition 7.611.*

Proof. Setting. Let $A(y), B(z)$ be (smeared) local operators and let

$$R_{AB}(y,z) \;:=\; \theta(y^0-z^0)\,\big\langle [\,A(y),B(z)\,]\big\rangle_{\mathbf{\Psi}}$$

be their microscopic retarded bi–distribution. *Microcausality* means

$$\operatorname{supp} R_{AB} \;\subset\; \{(y,z):\, y-z\in \mathsf{C}^+\}\,, \qquad \operatorname{supp}\langle [A(y),B(z)]\rangle_{\mathbf{\Psi}} \;\subset\; \{(y,z):\, y-z\in \mathsf{C}^+\cup\mathsf{C}^-\}. \tag{7.712}$$

Taggers and cone preservation. By hypothesis (cone–preserving taggers), there exist C_c^∞ kernels κ_x (IR) and π_x (the inverse Fourier transform of the UV projector P_x), real and even, such that

$$\operatorname{supp}\kappa_x \;\subset\; \mathsf{C}^+\cup\mathsf{C}^-, \qquad \operatorname{supp}\pi_x \;\subset\; \mathsf{C}^+\cup\mathsf{C}^-. \tag{7.713}$$

Band descent acts on operator–valued distributions by spacetime convolution:

$$A^{(x)} := \kappa_x * (\pi_x * A), \qquad B^{(x)} := \kappa_x * (\pi_x * B),$$

and on bi–distributions by the corresponding left–right convolutions.[19]

(I) Support of the band–retarded correlator. Consider the band–retarded bi–distribution

$$R^{(x)}_{AB}(y,z) \;:=\; \theta(y^0-z^0)\,\big\langle [\,A^{(x)}(y),B^{(x)}(z)\,]\big\rangle_{\mathbf{\Psi}}.$$

Using bilinearity of the commutator and Fubini for distributions,

$$R^{(x)}_{AB}(y,z)=\iint dy_1\,dz_1\;\kappa_x(y-y_1)\,\kappa_x(z-z_1)\;\theta(y^0-z^0)\;\big\langle [\,(\pi_x*A)(y_1),(\pi_x*B)(z_1)\,]\big\rangle_{\mathbf{\Psi}}.$$

Writing $(\pi_x * A)(y_1)=\int dy_2\,\pi_x(y_1-y_2)A(y_2)$ and similarly for B, we obtain

$$R^{(x)}_{AB}(y,z)=\iiiint dy_1dz_1dy_2dz_2\;\kappa_x(y-y_1)\,\kappa_x(z-z_1)\,\pi_x(y_1-y_2)\,\pi_x(z_1-z_2)\;R_{AB}(y_2,z_2)\,.$$

Hence, by successive applications of the support rule for convolutions $\operatorname{supp}(f*g)\subset \operatorname{supp}f+\operatorname{supp}g$ and by (7.712)–(7.713),

$$\operatorname{supp} R^{(x)}_{AB}\subset\Big\{(y,z):\;\exists(y_2,z_2)\text{ s.t. }\; y-y_2\in\operatorname{supp}\kappa_x+\operatorname{supp}\pi_x,\;\; z-z_2\in\operatorname{supp}\kappa_x+\operatorname{supp}\pi_x,\\ y_2-z_2\in\mathsf{C}^+\Big\}.$$

Therefore

$$y-z \;=\; (y_2-z_2)\;+\;(y-y_2)-(z-z_2)\;\in\;\mathsf{C}^+\;+\;(\mathsf{C}^\pm-\mathsf{C}^\pm)\;\subset\;\mathsf{C}^+,$$

because C^+ is a closed convex cone and $\mathsf{C}^\pm\pm\mathsf{C}^\pm\subset\mathsf{C}^\pm$ while $\mathsf{C}^+-\mathsf{C}^+\subset\mathsf{C}^+$.[20] Thus

$$\operatorname{supp} R^{(x)}_{AB} \;\subset\; \{(y,z):\, y-z\in\mathsf{C}^+\}. \tag{7.714}$$

[19] All convolutions are well–defined because the taggers are C_c^∞, hence the usual theory of convolution of distributions applies.

[20] Equivalently, in Fourier space one uses that the multipliers are supported in the dual cone and preserve retardedness.

The same argument without the Heaviside factor gives $\operatorname{supp}\langle[A^{(x)}(y),B^{(x)}(z)]\rangle_{\mathbf{\Psi}} \subset \{(y,z) : y-z \in \mathsf{C}^+ \cup \mathsf{C}^-\}$. Hence *band–retarded correlators vanish outside* C^+.

(II) Non–signalling of the law channel. Let A, B be spacelike–separated regions and let f_A, g_B be test functions supported in A, B, respectively. Define the band–smeared operators

$$\mathcal{J}_A^{(x)} := A^{(x)}(f_A), \qquad \mathcal{W}_B^{(x)} := B^{(x)}(g_B).$$

Their commutator expectation equals the distributional pairing of the microscopic commutator with the *effective* test functions after descent, i.e.

$$\Big\langle [\,\mathcal{J}_A^{(x)}, \mathcal{W}_B^{(x)}\,] \Big\rangle_{\mathbf{\Psi}} = \Big\langle [\,A, B\,],\ (K_x^\dagger f_A) \otimes (K_x^\dagger g_B) \Big\rangle, \quad K_x := \kappa_x * \pi_x,$$

so that $K_x^\dagger$ acts by convolution with $\tilde\kappa_x * \tilde\pi_x$, and

$$\operatorname{supp}(K_x^\dagger f_A) \ \subset\ A + \operatorname{supp}\kappa_x + \operatorname{supp}\pi_x =: A^\sharp, \qquad \operatorname{supp}(K_x^\dagger g_B) \ \subset\ B + \operatorname{supp}\kappa_x + \operatorname{supp}\pi_x =: B^\sharp.$$

By (7.713), $A^\sharp$ and $B^\sharp$ are *thickenings within the cones*. If A, B are spacelike–separated and the tagger apertures (time cutoffs and cone thickness) are chosen within the certified–plateau margins, then $A^\sharp$ and $B^\sharp$ remain spacelike–separated. Because the microscopic commutator distribution vanishes on spacelike pairs (7.712), the pairing is zero:

$$\Big\langle [\,\mathcal{J}_A^{(x)}, \mathcal{W}_B^{(x)}\,] \Big\rangle_{\mathbf{\Psi}} = 0.$$

This is precisely the *non–signalling* property in the sense of Definition 7.611 for the law channel.

(III) BRST/G structure and gauge independence. The taggers act by c–number spacetime convolution (and Fourier multiplication), with no action on field indices, hence they commute with BRST and G transformations (cf. Proposition 7.573). Passing to the physical sector $\mathcal{H}_{\text{phys}}$ removes BRST–exact/improvement terms (Lemma 7.552); therefore no gauge artefact can generate support outside $\mathsf{C}^\pm$ or spoil non–signalling.

Combining (I)–(III) proves the proposition. □

Definition 7.613 (Anomaly filter)**.** The *anomaly filter* is passed on $\mathfrak{P}$ if the following equivalent conditions hold:

1. (BRST nilpotency) $\zeta^2 = 0$ and $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$ (Stage-I gate).
2. (Adjoint covariance) The band Ward identity is adjoint-covariant (Proposition 7.568).
3. (Spectral matching) For all test functions φ supported near $\omega = 0$, $\mathcal{T}_a^{(x)}[\varphi] = C_\varphi(x)\,\mathcal{R}_a^{(x)}$ with $C_\varphi(x)$ independent of gauge parameters and free of non-covariant contact terms.

Failure of any condition signals an anomaly (e.g., a Wess–Zumino obstruction).

Theorem 7.614 (T-odd tomography & cones: completion criterion for Point 4)**.** *Assume the Stage-I gate (BRST anomaly–free), cone–preserving taggers, and a certified plateau* $\mathfrak{P}$*. Then:*

1. *(*Tracking*) The T–odd holosymmetric signal* $-\operatorname{Re}\Lambda_{\mathrm{law}}^{(\mathrm{spec};x)}$ *tracks the Ward–source residual via* (7.706)*; the quantitative lower bound of Corollary 7.610 holds uniformly on* $\mathfrak{P}$*.*
2. *(*No law–signalling*) Cone preservation implies non–signalling in the law channel (Proposition 7.612).*
3. *(*Anomaly filter*) The conditions in Definition 7.613 are satisfied; any observed deviation beyond plateau tolerances indicates either plateau failure or a (BRST/ward) anomaly.*

All statements are gauge–parameter independent on $\mathcal{H}_{\mathrm{phys}}$ *(Proposition 7.556) and stable under BRST/Noether improvements (Lemma 7.552).*

Proof. Preliminaries. Stage–I (*anomaly–free*) means the Wess–Zumino consistency holds for the band–descended Ward structure, with improvement/BRST–exact freedoms organized as in Lemma 7.563 and their spurion–sector versions unique on $\mathcal{H}_{\mathrm{phys}}$ by Corollary 7.564. Cone–preserving taggers commute with BRST/G and act by c–number spacetime convolution (cf. Proposition 7.573). Certified plateaus (Definition 6.300) give the oscillation budget $\eta_{\mathfrak{P}}$ and uniform floors for the spectral kernel (Lemma 7.575 and Proposition 7.602).

(1) Tracking. By the spectral/tomographic representation of the T–odd law channel (Definition 7.607) and the low–frequency Ward matching (Lemma 7.608), with a normalized, even mollifier χ_τ supported in $[-\omega_0(\tau),\omega_0(\tau)]$,

$$\mathcal{T}_a^{(x)}[\chi_\tau] \xrightarrow[\tau\to\infty]{} c_0(x)\,|\mathcal{R}_a^{(x)}|, \qquad c_0(x)>0.$$

The tomographic *lower bound* (Proposition 7.609) gives, uniformly on $\mathfrak{P}$,

$$1-|\Lambda_{\mathrm{law}}^{(x)}| \;\geq\; \kappa_{\mathrm{spec}}(x)\;\mathcal{T}_a^{(x)}[\chi_\tau] \;-\; O(\eta_{\mathfrak{P}}) \;-\; o_\tau(1).$$

Letting $\tau\to\infty$ yields the quantitative tracking bound of Corollary 7.610:

$$1-|\Lambda_{\mathrm{law}}^{(x)}| \;\gtrsim\; \kappa_{\mathrm{spec}}(x)\,c_0(x)\,|\mathcal{R}_a^{(x)}| \quad (\text{up to } O(\eta_{\mathfrak{P}})).$$

This proves Item (1).

(2) No law–signalling. By microcausality of the microscopic theory and cone–preserving taggers, band–retarded correlators have support inside $\mathsf{C}^{\pm}$ and band–smeared commutators vanish at spacelike separations (Proposition 7.612). Hence the law channel is non–signalling in the sense of Definition 7.611. This proves Item (2).

(3) Anomaly filter. Define the *tomographic residual*

$$\Delta_\tau(x) \;:=\; \Big(1-|\Lambda_{\mathrm{law}}^{(x)}|\Big) \;-\; \kappa_{\mathrm{spec}}(x)\,\mathcal{T}_a^{(x)}[\chi_\tau].$$

From Proposition 7.609 we have, uniformly on $\mathfrak{P}$,

$$\Delta_\tau(x) \;\geq\; -\,O(\eta_{\mathfrak{P}}) \;-\; o_\tau(1). \tag{7.715}$$

On the other hand, Stage–I anomaly–freeness plus Lemma 7.608 control the *model side*: $\mathcal{T}_a^{(x)}[\chi_\tau] \to c_0(x)|\mathcal{R}_a^{(x)}|$ with BRST/Noether–improvement pieces dropping out on $\mathcal{H}_{\mathrm{phys}}$

(Lemma 7.552, Proposition 7.556, and Corollary 7.564). Therefore any observed *systematic* violation

$$\limsup_{\tau\to\infty}\ \sup_{x\in\mathfrak{P}}\ \Delta_\tau(x)\ >\ c_\star\,\eta_{\mathfrak{P}}\qquad\text{for some } c_\star>0$$

cannot be accommodated by the $O(\eta_{\mathfrak{P}})$ oscillation budget in (7.715). The only remaining possibilities are: (i) failure of plateau certification (e.g. loss of kernel floor, non–monotone refinement, or masking issues), or (ii) breakdown of the Ward matching—i.e. a BRST/ward anomaly as formulated in Definition 7.613. Hence the anomaly–filter conditions are satisfied: deviations beyond the plateau tolerance trigger a (plateau–or–anomaly) alert. This proves Item (3).

Gauge/BRST robustness. Every step uses band–descended, BRST–closed quantities; taggers commute with BRST/G (Proposition 7.573), and physical expectations are taken on $\mathcal{H}_{\text{phys}}$ (Proposition 7.556 and Lemma 7.552). Thus all conclusions are gauge–parameter independent and immune to BRST/Noether improvements.

The theorem follows. □

7.9.9 Law–channel Diagnosis: Saturation vs. QIII

Definition 7.615 (Admissible preparations on a certified band point)**.** Fix $x=(\lambda_{\text{UV}},\ell)\in\mathfrak{P}$ (Definition 6.300). Let $\mathcal{C}_x$ denote the set of band-prepared, BRST-closed, gauge-parameter–independent density operators $\rho_0^{(x)}$ constructed by the protocol of Definitions 6.275 and 7.551 (same UV/IR taggers, same boundary preparation class).

Proposition 7.616 (Law-channel saturation via Duhamel/BCH (QIV))**.** *Let $x\in\mathfrak{P}$ and write the tagged branch unitaries as $U_A^{(x)},U_B^{(x)}$ with $U_B^{(x)\dagger}U_A^{(x)}=\exp\{i\,\varepsilon_{\text{IR}}(x)\,G_x+\mathcal{O}(\varepsilon_{\text{IR}}^2)\}$ in the interaction picture, where G_x is a self-adjoint, band-local generator proportional to the explicit breaking source (Duhamel representation), and $\varepsilon_{\text{IR}}(x)$ is the band IR-norm of the spurion deformation (non-running in QIV). Then for all $\rho_0^{(x)}\in\mathcal{C}_x$,*

$$1-\left|\Lambda_{\text{law}}^{(x)}\right|=1-\left|\operatorname{Tr}\!\left(U_B^{(x)\dagger}U_A^{(x)}\,\rho_0^{(x)}\right)\right|\ \geq\ \tfrac{1}{2}\,\varepsilon_{\text{IR}}(x)^2\,\operatorname{Var}_{\rho_0^{(x)}}(G_x)\ +\ o\!\left(\varepsilon_{\text{IR}}(x)^2\right).$$

Moreover, on a certified plateau there exists $c_x>0$ (gauge-parameter independent) such that $\operatorname{Var}_{\rho_0^{(x)}}(G_x)\geq c_x$ for all $\rho_0^{(x)}\in\mathcal{C}_x$, and hence

$$\sup_{\rho_0^{(x)}\in\mathcal{C}_x}\left|\Lambda_{\text{law}}^{(x)}\right|\ \leq\ 1-\tfrac{1}{2}\,c_x\,\varepsilon_{\text{IR}}(x)^2\ +\ o\!\left(\varepsilon_{\text{IR}}(x)^2\right).$$

Uniformly on $\mathfrak{P}$, c_x can be replaced by a strictly positive $c_\star$ (plateau uniformization, Lemma 7.579).

Proof sketch. (i) Duhamel/BCH reduction. Set $U_B^{(x)\dagger}U_A^{(x)}=\mathcal{T}\exp\!\left(i\int_0^1 ds\,\Delta H_x(s)\right)$ with ΔH_x the band-IR explicit-breaking Hamiltonian density (spurion-sourced). Duhamel's formula yields

$$U_B^{(x)\dagger}U_A^{(x)}=\exp\!\Big(i\,\varepsilon_{\text{IR}}(x)\,G_x-\tfrac{1}{2}\varepsilon_{\text{IR}}(x)^2\,C_x+\mathcal{O}(\varepsilon_{\text{IR}}^3)\Big),$$

where G_x is self-adjoint and C_x collects the time-ordered commutators (BCH/Dyson series); both are band-local and BRST/Nielsen safe by Definition 7.551, Lemma 7.552, and Proposition 7.556.

(ii) Quantum characteristic function bound. For any density matrix ρ and small ϵ, expand

$$\mathrm{Tr}\Big(e^{i\epsilon G_x}\rho\Big) = 1 + i\epsilon\,\langle G_x\rangle_\rho - \tfrac{1}{2}\epsilon^2\langle G_x^2\rangle_\rho + \mathcal{O}(\epsilon^3).$$

Taking modulus and using $|1 + iz - \frac{1}{2}a + o(1)| \le 1 - \frac{1}{2}(a - z^2) + o(1)$ gives

$$\begin{aligned} 1 - \Big|\,\mathrm{Tr}(e^{i\epsilon G_x}\rho)\Big| &\ge \tfrac{1}{2}\,\epsilon^2\Big(\langle G_x^2\rangle_\rho - \langle G_x\rangle_\rho^{\,2}\Big) + o(\epsilon^2) \\ &= \tfrac{1}{2}\,\epsilon^2\ \mathrm{Var}_\rho(G_x) + o(\epsilon^2). \end{aligned}$$

The additional $\mathcal{O}(\varepsilon_{\mathrm{IR}}^2)$ term from C_x only strengthens the deficit (it is T-even and does not cancel the variance term). Plug in $\epsilon = \varepsilon_{\mathrm{IR}}(x)$ and $\rho = \rho_0^{(x)}$.

(iii) Plateau lower bound for the variance. The non-vanishing Ward source on $\mathfrak{P}$ (Theorem 7.570) together with the T-odd tomography representation (Definition 7.607, Lemma 7.578 and Proposition 7.609) implies a strictly positive low-frequency spectral weight tied to G_x. This gives a uniform positive lower bound

$$\Big\langle G_x^2\Big\rangle_{\rho_0^{(x)}} - \Big\langle G_x\Big\rangle_{\rho_0^{(x)}}^{2} \ \ge\ c_x > 0$$

over the certified class $\mathcal{C}_x$ independent of gauge parameters (Proposition 7.556). Plateau uniformization (Lemma 7.579) yields $c_x \ge c_\star > 0$ for all $x \in \mathfrak{P}$ after tightening tolerances if needed. □

Corollary 7.617 (QIV vs. QIII saturation test)**.** *Under QIV (relevant/marginal spurions),* $\varepsilon_{\mathrm{IR}}(x)$ *does* not *run to* 0 *with* ℓ *on* $\mathfrak{P}$*; hence*

$$\Big|\Lambda_{\mathrm{law}}^{(x)}\Big| \ \le\ 1 - \delta_\star + o(1), \qquad \delta_\star := \tfrac{1}{2}\,c_\star\,\varepsilon_{\mathrm{IR},\star}^2 \in (0,1),$$

uniformly in $x \in \mathfrak{P}$*. By contrast, in QIII the effective spurion norm scales as* $\varepsilon_{\mathrm{IR}}(x) \sim \ell^{\Delta-(d+z)}$ *(emergent symmetry), so the measured law-deficit* improves *as* $1 - |\Lambda_{\mathrm{law}}(x)| = A\,\ell^{\,\Delta-(d+z)} + o(\ell^{\,\Delta-(d+z)})$ *(Theorem 7.580 and the QIII scaling overlay), cleanly disentangling QIV from QIII.*

Proof. Step 1: Quadratic lower bound from the BCH/Duhamel expansion. On a certified plateau, the branch-link has the expansion

$$U_B^{(x)\dagger}U_A^{(x)} = \exp\Big(i\,\varepsilon_{\mathrm{IR}}(x)\,G_x - \tfrac{1}{2}\varepsilon_{\mathrm{IR}}(x)^2 C_x + \cdots\Big)$$

(cf. the BCH form used in Definition 7.674).

Let $\langle\cdot\rangle_{\mathbf{\Psi}}$ denote the physical expectation on $\mathcal{H}_{\mathrm{phys}}$. For small ε one has the standard characteristic–function inequality (second–order cumulant expansion)

$$\Big|\Big\langle e^{\,i\varepsilon G_x}\Big\rangle_{\mathbf{\Psi}}\Big| \ \le\ 1 - \tfrac{1}{2}\,\varepsilon^2\ \mathrm{Var}_{\mathbf{\Psi}}(G_x) + O(\varepsilon^3).$$

hence for the (law–channel) overlap magnitude,

$$1 - |\Lambda^{(x)}_{\mathrm{law}}| \;\geq\; \tfrac{1}{2}\,\varepsilon_{\mathrm{IR}}(x)^2\,\mathrm{Var}_{\Psi}(G_x) \;-\; \mathcal{O}\big(\varepsilon_{\mathrm{IR}}(x)^3\big) \;-\; \mathcal{O}(\eta_{\mathfrak{P}}). \tag{7.716}$$

The $O(\eta_{\mathfrak{P}})$ term reflects the plateau envelope between channels (Definition 6.300, Corollary 7.677). By cone–preserving taggers and kernel positivity (Lemma 7.575 and Proposition 7.602), the band–descended variance has a plateau–uniform strictly positive lower bound:

$$\mathrm{Var}_{\Psi}(G_x) \;\geq\; c_\star > 0 \qquad (x \in \mathfrak{P}).$$

because G_x inherits a positive spectral weight on the certified window (nontrivial T–odd floor; cf. the tracking statements in Corollary 7.677(i), Corollary 7.610).

Step 2: QIV saturation. QIV means the IR spurion does *not* decay with the IR scale on $\mathfrak{P}$ ("no IR improvement"): by Theorem 7.580 (and the slope test Proposition 7.599), there exists

$$\varepsilon_{\mathrm{IR},\star} \;:=\; \inf_{\ell \text{ on the certified window}}\; \inf_{x\in\mathfrak{P}}\; \varepsilon_{\mathrm{IR}}(x;\ell) \;>\; 0.$$

Plugging $\varepsilon_{\mathrm{IR}}(x) \geq \varepsilon_{\mathrm{IR},\star}$ and $\mathrm{Var}(G_x) \geq c_\star$ into (7.716), and absorbing $O(\varepsilon^3_{\mathrm{IR}})$ into $o(1)$ along the window (fixed $\varepsilon_{\mathrm{IR},\star}$, $\ell \to \infty$), we obtain

$$1 - |\Lambda^{(x)}_{\mathrm{law}}| \;\geq\; \tfrac{1}{2}\,c_\star\,\varepsilon^2_{\mathrm{IR},\star} \;-\; o(1) \;-\; O(\eta_{\mathfrak{P}}).$$

Equivalently,

$$|\Lambda^{(x)}_{\mathrm{law}}| \;\leq\; 1 - \delta_\star + o(1), \qquad \delta_\star := \tfrac{1}{2}\,c_\star\,\varepsilon^2_{\mathrm{IR},\star} \in (0,1),$$

uniformly for $x \in \mathfrak{P}$ (plateau compactness). This is the claimed QIV saturation bound.

Step 3: QIII power–law improvement. In QIII (emergent symmetry), the spurion is *irrelevant* at the IR fixed point with scaling dimension $\Delta > d + z$. Under anisotropic RG with dynamic exponent z, the effective IR spurion scales as

$$\varepsilon_{\mathrm{IR}}(\ell) \;\sim\; \ell^{\,\Delta-(d+z)} \qquad (\ell \to \infty),$$

i.e. it runs to 0 with a negative exponent $y = d + z - \Delta < 0$. By the tomographic tracking bound (Proposition 7.609 and Lemma 7.608),

$$1 - |\Lambda^{(x)}_{\mathrm{law}}| \;\geq\; \kappa_{\mathrm{spec}}(x)\,c_0(x)\,|\mathcal{R}^{(x)}_a| \;-\; o(1) \;-\; O(\eta_{\mathfrak{P}}).$$

The Ward residual is linear in the spurion source (eq. (7.665) implicitly used in Lemma 7.608): $|\mathcal{R}^{(x)}_a| \asymp \varepsilon_{\mathrm{IR}}(\ell)$ along the certified window (the proportionality is uniform on $\mathfrak{P}$ by the spectral floors and probe normalization). Consequently,

$$1 - |\Lambda^{(x)}_{\mathrm{law}}| \;=\; A(x)\,\ell^{\,\Delta-(d+z)} \;+\; o\big(\ell^{\,\Delta-(d+z)}\big), \quad A(x) := \kappa_{\mathrm{spec}}(x)\,c_0(x)\,\big\|\text{PCAC factor at } x\big\|,$$

uniformly up to the plateau tolerance. Taking $A := \sup_{x\in\mathfrak{P}} A(x)$ yields the displayed QIII scaling law.

Step 4: Clean separation. In QIV, the uniform *positive* constant $\delta_\star$ forces a nonvanishing law–deficit as $\ell \to \infty$; in QIII, the power–law with exponent $\Delta - (d+z) < 0$ guarantees

improvement (decay to 0). Thus the behaviors are disjoint on certified windows, providing a robust operational separation of QIV vs. QIII as stated.

Gauge/BRST robustness. All inputs—kernel floors, tomographic functionals, Ward residuals—are built from band–descended, BRST–closed observables; taggers commute with BRST/G (Proposition 7.573), and expectations are taken on $\mathcal{H}_{\text{phys}}$ (Lemma 7.552 and Proposition 7.556). Hence the conclusions are gauge–parameter independent and Nielsen–stable.

This completes the proof. □

7.9.10 Near–degeneracies and GMOR Fits in the Mass Ledger

Context. On $\mathcal{H}_{\text{phys}}$ the band Hessian $\mathbb{H}^{(x)}(\epsilon)$ is a self–adjoint, type–(A) analytic family in the spurions $\epsilon = (\epsilon_k)$ (Proposition 7.642); pseudo–NG eigenvalues and their Riesz projectors vary real–analytically, and residues/poles are Nielsen–independent (Proposition 7.644). We now (i) formalize *analytic eigenvector labeling* in the presence of near–crossings and (ii) give a one–line *fit specification* for $m_{\pi_b}^2$ vs. ϵ_j with vanishing intercept within plateau tolerance.

Definition 7.618 (Near–degeneracy window and isolated cluster)**.** Fix a line in spurion space $\epsilon(t) = \epsilon^{(0)} + t\,\hat{\mathbf{e}}_j$ with $t \in I \subset \mathbb{R}$ small, and a band point $x \in \mathfrak{P}$. Let $\{\lambda_\beta^{(x)}(t)\}$ be pseudo–NG eigenvalues of $\mathbb{H}^{(x)}(\epsilon(t))$ with Riesz projectors $\mathsf{P}_\beta^{(x)}(t)$. We say that a finite index set $\mathcal{B}$ forms an *isolated cluster* on I with *gap* $\delta_{\text{gap}} > 0$ if

$$\operatorname{dist}\Big(\ \sigma\Big(\mathbb{H}^{(x)}(\epsilon(t))\Big) \restriction \operatorname{Ran}\textstyle\sum_{\beta\in\mathcal{B}} \mathsf{P}_\beta^{(x)}(t)\ ,\ \sigma\Big(\mathbb{H}^{(x)}(\epsilon(t))\Big)\setminus \operatorname{Ran}\textstyle\sum_{\beta\in\mathcal{B}} \mathsf{P}_\beta^{(x)}(t)\ \Big)\ \geq\ \delta_{\text{gap}} \quad \forall\, t \in I.$$

Equivalently, the spectral subspace $\mathcal{E}_{\mathcal{B}}^{(x)}(t) = \operatorname{Ran}\mathsf{P}_{\mathcal{B}}^{(x)}(t)$ with $\mathsf{P}_{\mathcal{B}}^{(x)}(t) = \frac{1}{2\pi i}\oint_\gamma (z - \mathbb{H}^{(x)}(\epsilon(t)))^{-1}\,dz$ is well–defined for a fixed contour γ enclosing that cluster.

Lemma 7.619 (Analytic subspace tracking (Riesz projectors))**.** *Under Definition 7.618, $t \mapsto \mathsf{P}_{\mathcal{B}}^{(x)}(t)$ is real–analytic on I and so is any spectral invariant (e.g. the eigenvalue sum $\sum_{\beta\in\mathcal{B}} \lambda_\beta^{(x)}(t)$). Gauge independence holds on $\mathcal{H}_{\text{phys}}$ by Proposition 7.644.*

Proof. Setup and gap. Let $\mathsf{H}^{(x)}(t)$ denote the (band) Hermitian operator/matrix depending on $t \in I \subset \mathbb{R}$. By Proposition 7.642 the family $\{\mathsf{H}^{(x)}(t)\}_{t\in I}$ is a type–(A) real–analytic family. By Definition 7.618 there exists a *fixed* positively oriented contour $\Gamma \subset \mathbb{C}$ such that, for all $t \in I$,

$$\Gamma \subset \rho\Big(\mathsf{H}^{(x)}(t)\Big) \qquad \text{and} \qquad \operatorname{dist}\Big(\Gamma,\ \sigma\Big(\mathsf{H}^{(x)}(t)\Big)\Big)\ \geq\ \delta > 0,$$

and Γ encloses exactly the eigenvalue cluster $\mathcal{B}$ (with constant total multiplicity $m := \dim \operatorname{Ran}\mathsf{P}_{\mathcal{B}}^{(x)}(t)$).

Analyticity of the Riesz projector. Write the resolvent $R(t,z) := (z - \mathsf{H}^{(x)}(t))^{-1}$. For type–(A) families, the map $(t,z) \mapsto R(t,z)$ is holomorphic in $z \in \rho(\mathsf{H}^{(x)}(t))$ and real–analytic in t, jointly locally bounded on $I \times \{\text{neigh. of } \Gamma\}$ (standard Kato theory; cf. Proposition 7.642). Define the Riesz projector by the holomorphic functional calculus

$$\mathsf{P}_{\mathcal{B}}^{(x)}(t)\ :=\ \frac{1}{2\pi i}\oint_\Gamma R(t,z)\,dz.$$

Because $R(t,\cdot)$ is holomorphic on and inside Γ and depends real–analytically on t with uniform bounds on Γ, the contour integral preserves real–analyticity: $t \mapsto \mathsf{P}^{(x)}_{\mathcal{B}}(t)$ is real–analytic on I. In particular, $\mathsf{P}^{(x)}_{\mathcal{B}}(t)$ is a projector of constant rank m and depends continuously (indeed, real–analytically) on t. Differentiating under the integral gives the familiar formula (valid for all $t \in I$):

$$\partial_t \mathsf{P}^{(x)}_{\mathcal{B}}(t) \;=\; \frac{1}{2\pi i}\oint_{\Gamma} R(t,z)\,\Big(\partial_t \mathsf{H}^{(x)}(t)\Big)\,R(t,z)\,dz,$$

which exhibits real–analyticity of all derivatives.

Analyticity of spectral invariants. Let f be holomorphic on a neighbourhood of the spectrum inside Γ. Then

$$f\Big(\mathsf{H}^{(x)}(t)\Big)\,\mathsf{P}^{(x)}_{\mathcal{B}}(t) \;=\; \frac{1}{2\pi i}\oint_{\Gamma} f(z)\,R(t,z)\,dz,$$

is real–analytic in t (same argument). Since the trace is a continuous linear functional, $t \mapsto \mathrm{Tr}\Big(f(\mathsf{H}^{(x)}(t))\,\mathsf{P}^{(x)}_{\mathcal{B}}(t)\Big)$ is real–analytic. Choosing $f(z) = z$ yields

$$\sum_{\beta\in\mathcal{B}} \lambda^{(x)}_{\beta}(t) \;=\; \mathrm{Tr}\Big(\mathsf{H}^{(x)}(t)\,\mathsf{P}^{(x)}_{\mathcal{B}}(t)\Big)\,,$$

hence the eigenvalue sum is real–analytic on I. More generally, any spectral invariant that can be written as a trace of a holomorphic functional of $\mathsf{H}^{(x)}(t)$ restricted by $\mathsf{P}^{(x)}_{\mathcal{B}}(t)$ (e.g. power sums $\sum_{\beta\in\mathcal{B}} \lambda_\beta^p$, elementary symmetric polynomials via det on the range, etc.) is real–analytic by the same calculus.

Gauge/Nielsen independence on $\mathcal{H}_{\mathrm{phys}}$. Gauge–parameter variations act as Nielsen reparametrizations that either (i) implement similarity/adjoint actions commuting with the projection onto $\mathcal{H}_{\mathrm{phys}}$, or (ii) add BRST–exact/improvement pieces that are null on $\mathcal{H}_{\mathrm{phys}}$. By Proposition 7.644, spectral quantities built via the holomorphic functional calculus (Riesz projectors, eigenvalue sums on the physical sector) are gauge–parameter independent. Therefore the statements above hold on $\mathcal{H}_{\mathrm{phys}}$.

This completes the proof. □

Proposition 7.620 (Kato parallel transport and analytic labeling)**.** *Let $\mathcal{E}^{(x)}_{\mathcal{B}}(t)$ be as above with $\dim \mathcal{E}^{(x)}_{\mathcal{B}}(t) = r$. Choose an orthonormal frame $\{|\psi^{(x)}_\ell(t)\rangle\}_{\ell=1}^{r}$ in $\mathcal{E}^{(x)}_{\mathcal{B}}(t)$ by* Kato parallel transport*:*

$$\frac{d}{dt}\,|\psi^{(x)}_\ell(t)\rangle = \Big[\partial_t \mathsf{P}^{(x)}_{\mathcal{B}}(t),\,\mathsf{P}^{(x)}_{\mathcal{B}}(t)\Big]\,|\psi^{(x)}_\ell(t)\rangle\,, \qquad |\psi^{(x)}_\ell(t_0)\rangle = |\psi^{(x)}_\ell(t_0)\rangle_0\,, \tag{7.717}$$

for some initial orthonormal eigenbasis at $t_0 \in I$. Then $t \mapsto |\psi^{(x)}_\ell(t)\rangle$ is real–analytic, and the principal angles *between $\{|\psi^{(x)}_\ell(t)\rangle\}$ and a fixed reference basis give analytic, gauge–independent* mixing angles*. If the cluster is actually simple ($r = 1$) at t, this choice coincides with the unique normalized eigenvector up to a constant phase.*

Proof. Step 0: Analytic projectors and identities. By Lemma 7.619, the Riesz projector $\mathsf{P}(t) := \mathsf{P}^{(x)}_{\mathcal{B}}(t)$ is real–analytic on I and self–adjoint. Differentiating $\mathsf{P}(t)^2 = \mathsf{P}(t)$ gives

$$\mathsf{P}\,\dot{\mathsf{P}}\,\mathsf{P} = 0, \qquad (\mathbf{1}-\mathsf{P})\,\dot{\mathsf{P}}\,(\mathbf{1}-\mathsf{P}) = 0, \qquad [\dot{\mathsf{P}},\mathsf{P}]^\dagger = -[\dot{\mathsf{P}},\mathsf{P}]. \tag{7.718}$$

Hence the *Kato generator* $K(t) := [\dot{\mathsf{P}}(t), \mathsf{P}(t)]$ is bounded, real–analytic in t, and skew–adjoint.

Step 1: Existence/uniqueness, unitarity, and invariance of the subspace. Consider the matrix ODE on the ambient Hilbert space,

$$\dot{U}(t) = K(t)\,U(t), \qquad U(t_0) = \mathbf{1}.$$

Since K is continuous (indeed analytic), the Picard–Dyson series converges on compact intervals; hence there is a unique strongly–continuous (indeed analytic) solution $U(t)$. Skew–adjointness of K implies $\frac{d}{dt}\big(U^\dagger U\big) = U^\dagger(K^\dagger + K)U = 0$, so $U(t)$ is unitary for all t.

Define the transported frame by $|\psi_\ell(t)\rangle := U(t)\,|\psi_\ell(t_0)\rangle_0$; orthonormality follows from unitarity. To see that $|\psi_\ell(t)\rangle \in \operatorname{Ran}\mathsf{P}(t)$, set $\eta_\ell(t) := (\mathbf{1} - \mathsf{P}(t))\,|\psi_\ell(t)\rangle$ and differentiate, using (7.718) and $\dot{\psi}_\ell = K\psi_\ell$:

$$\dot{\eta}_\ell = -(\dot{\mathsf{P}})\psi_\ell + (\mathbf{1}-\mathsf{P})K\psi_\ell = -(\dot{\mathsf{P}})\psi_\ell + (\mathbf{1}-\mathsf{P})\dot{\mathsf{P}}\,\mathsf{P}\psi_\ell - (\mathbf{1}-\mathsf{P})\mathsf{P}\,\dot{\mathsf{P}}\psi_\ell = -(\mathbf{1}-\mathsf{P})\,\dot{\mathsf{P}}\,(\mathbf{1}-\mathsf{P})\psi_\ell = 0,$$

where we used $(\mathbf{1} - \mathsf{P})\mathsf{P} = 0$ and (7.718). Since $\eta_\ell(t_0) = 0$, we have $\eta_\ell(t) \equiv 0$; thus $\mathsf{P}(t)\,|\psi_\ell(t)\rangle = |\psi_\ell(t)\rangle$ for all t.

Step 2: Real–analytic dependence. Because $K(t)$ is real–analytic in t (products/commutators of analytic maps), the unique solution $U(t)$ depends real–analytically on t (standard ODE theory with analytic coefficients on finite–dimensional ranges; equivalently, use the convergent time–ordered exponential with analytic integrand). Consequently $t \mapsto |\psi_\ell(t)\rangle = U(t)\,|\psi_\ell(t_0)\rangle_0$ is real–analytic on I.

Step 3: Analytic, gauge–independent mixing angles (principal angles). Fix a reference orthonormal basis $\{|\phi_j\rangle\}_{j=1}^r$ of $\mathcal{E}_{\mathcal{B}}^{(x)}(t_0)$ and let P_0 be the orthogonal projector onto their span. The *principal angles* $\{\vartheta_j(t)\}_{j=1}^r$ between $\mathcal{E}_{\mathcal{B}}^{(x)}(t)$ and $\mathcal{E}_{\mathcal{B}}^{(x)}(t_0)$ are defined by

$$\cos^2\vartheta_j(t) \;=\; \lambda_j\Big(C(t)\Big), \qquad C(t) := P_0\,\mathsf{P}(t)\,P_0 \;\text{ on }\; \operatorname{Ran} P_0,$$

where $\lambda_j(\cdot)$ are the eigenvalues (counting multiplicity). Since $\mathsf{P}(t)$ is analytic and P_0 is constant, the compressed operator $C(t)$ is a finite–dimensional Hermitian analytic family. By Kato's analytic perturbation theory, its eigenprojections/eigenvalues admit real–analytic branches (after choosing an analytic labeling via Riesz projectors around clusters). Therefore $\cos^2\vartheta_j(t)$ are real–analytic, and so are $\vartheta_j(t) = \arccos\sqrt{\lambda_j\Big(C(t)\Big)}$ (the square–root and arccos are analytic on $(0,1]$, which contains the spectrum for isolated subspaces).

Gauge independence. Replacing the transported frame $\{|\psi_\ell(t)\rangle\}$ by any t–dependent unitary rotation inside $\mathcal{E}_{\mathcal{B}}^{(x)}(t)$ leaves $\mathsf{P}(t)$ (hence $C(t)$) invariant; thus the principal angles are independent of the choice of frame. Variations of gauge parameters act by Nielsen–safe transformations that leave spectral projectors on $\mathcal{H}_{\text{phys}}$ invariant (Proposition 7.644); hence the mixing angles are gauge–parameter independent on $\mathcal{H}_{\text{phys}}$.

Step 4: Simple cluster ($r = 1$). If $r = 1$, then $\mathsf{P}(t) = |\psi(t)\rangle\langle\psi(t)|$ is rank one. The Kato ODE (7.717) reduces to the *parallel–transport* condition $\langle\psi(t)\rangle\,\dot{\psi}(t) = 0$ (since K maps $\operatorname{Ran}\mathsf{P}$ to its orthogonal complement by (7.718)). Thus $|\psi(t)\rangle$ is the unique normalized eigenvector in $\operatorname{Ran}\mathsf{P}(t)$ up to a *constant* phase: if $|\varphi(t)\rangle = e^{i\theta(t)}\,|\psi(t)\rangle$ also solves (7.717), then taking inner products with $|\psi(t)\rangle$ shows $\dot{\theta}(t) = 0$.

Combining Steps 1–4 proves the proposition. □

Definition 7.621 (Operational tolerance for near–crossings)**.** On the mask $\mathfrak{M}_{\mathrm{mask}}$ (Proposition 7.663), we enforce the rule:

1. If along $\epsilon(t)$ two pseudo–NG levels approach with minimal separation $\min_{t\in I}|\lambda_\beta^{(x)}(t)-\lambda_{\beta'}^{(x)}(t)|\geq\delta_{\mathrm{gap}}$, we *label* them individually by Kato transport (Proposition 7.620) and report separate masses/mixings.
2. If the separation drops below δ_{gap}, we *merge* them into a single cluster $\mathcal{B}$, report the eigenvalue *mean* and *spread* $\bar\lambda_{\mathcal{B}}=\frac{1}{r}\sum_{\beta\in\mathcal{B}}\lambda_\beta,\quad \Delta_{\mathcal{B}}=\max_{\beta\in\mathcal{B}}|\lambda_\beta-\bar\lambda_{\mathcal{B}}|$, and quote mixing angles inside $\mathcal{B}$ (frame–independent principal angles). We do *not* over–interpret slopes of individual levels through an avoided crossing.

The constant δ_{gap} is chosen $\mathcal{O}(\eta_{\mathfrak{P}})$–larger than the spectral grid resolution, ensuring mask stability under refinement (Lemma 7.662).

Remark 7.622 (Consistency with GMOR)**.** Within any cluster, the linear GMOR relation of Theorem 7.630 applies at the *subspace* level: $F^{(x)}M^2_{\pi,\mathcal{B}}=\sum_k\epsilon_k C^{(x)}_{k,\mathcal{B}}+\mathcal{O}(\epsilon^2)$ with matrices restricted/prolonged to $\mathcal{E}^{(x)}_{\mathcal{B}}(t)$. Individual level slopes are well–defined away from mergers; at mergers, report the linear map on the subspace.

Definition 7.623 (One–line GMOR fit specification (single spurion))**.** Fix $x\in\mathfrak{P}$ and a broken mode label b (or a Kato–labeled branch within a cluster obeying Definition 7.621). Vary a single spurion ϵ_j along $\epsilon=\epsilon^{(0)}+\epsilon_j\hat{\mathbf{e}}_j$ with all other $\epsilon_{k\neq j}$ fixed at their working values. On the admissible interval where Theorem 7.630 holds, fit

$$m^2_{\pi_b}(x;\epsilon_j)\;=\;\alpha^{(x)}_{bj}\,\epsilon_j\quad(+\text{ report }\;\hat\beta\text{ but enforce }|\hat\beta|\leq\eta_{\mathfrak{P}}\text{ for intercept}),$$

by *constrained weighted least squares*:

$$\min_{\alpha,\beta}\;\sum_n w_n\Big(m^2_{\pi_b}(x;\epsilon^{(n)}_j)-\alpha\,\epsilon^{(n)}_j-\beta\Big)^2\quad\text{s.t.}\quad|\beta|\leq\eta_{\mathfrak{P}},$$

with weights w_n from the spectral–fit uncertainties and $\eta_{\mathfrak{P}}$ the plateau tolerance. Report the slope $\hat\alpha^{(x)}_{bj}$ and its CI; verify *linearity* by an F–test against a quadratic extension. By Corollary 7.631, $\hat\alpha^{(x)}_{bj}$ should match $\frac{1}{f^2_{\pi_b}(x)}\big\langle\mathcal{S}^{(\lambda_{\mathrm{UV}})}_{j,b}\big\rangle_{\mathbf{\Psi}}$ within tolerance and is gauge–independent (Proposition 7.644).

7.9.11 Ward–mass Link (GMOR–type Control of Pseudo–NG)

In QIV, explicit spurions select a unique vacuum and render the would–be NG excitations gapped (pseudo–NG). Here we relate those gaps to the *band* Ward source via a GMOR–type identity, and show *linear* scaling in the spurions on certified plateaus.

Definition 7.624 (Non–conservation operator on the band)**.** At a band point $x=(\lambda_{\mathrm{UV}},\ell)$, define the divergence (non–conservation) operator

$$\Delta^{(x)}_a\;:=\;\nabla^{(\mathrm{ad})}_\mu\,\bar J^{(x)\mu}_{a,\mathrm{imp}},\qquad \bar J^{(x)\mu}_{a,\mathrm{imp}}:=K^{(\ell)}_{\mathrm{IR}}P_x\,J^\mu_{a,\mathrm{imp}}\,P_xK^{(\ell)}_{\mathrm{IR}}.$$

On $\mathcal{H}_{\mathrm{phys}}$, the scale–tagged Ward identity (eq. (7.665)) gives $\Delta^{(x)}_a=\mathcal{R}^{(x)}_a:=\sum_k\epsilon_k\,\langle\mathcal{S}^{(\lambda_{\mathrm{UV}})}_{k,a}\rangle_{\mathbf{\Psi}}$ as an operator statement modulo BRST/exact and improvement terms that vanish in physical matrix elements (Lemma 7.552 and Proposition 7.569).

Definition 7.625 (Broken basis, pseudo–NG states and decay constants)**.** Work in the *broken basis* on the band where the mass/mixing ledger (physical Hessian) of Section 7.9 is block–diagonal between (i) light pseudo–NG modes $\{\pi_\beta^{(x)}\}$, $\beta = 1, \dots, N_{\rm br}$, and (ii) heavier modes h. Let $|\pi_\beta^{(x)}(p)\rangle$ be the one–particle pseudo–NG state (relativistic or Lifshitz with $z < \infty$ as appropriate). Define the (band) decay–constant matrix $F^{(x)}$ by the current–pole residue

$$\left\langle \Omega^{(x)} \right| \bar{J}^{(x)\mu}_{a,\rm imp}(0) \left| \pi^{(x)}_\beta(p) \right\rangle = i\, F^{(x)}_{a\beta}\, p^\mu \;+\; \mathcal{O}(p^2), \tag{7.719}$$

and let $m^2_\beta(x)$ denote the pseudo–NG squared masses read from the physical Hessian/mass operator. All objects are evaluated on $\mathcal{H}_{\rm phys}$ and are gauge–parameter independent by Proposition 7.556.

Lemma 7.626 (Band–PCAC on one–particle states)**.** *For each broken index a and pseudo–NG label β,*

$$\left\langle \Omega^{(x)} \right| \Delta^{(x)}_a(0) \left| \pi^{(x)}_\beta(p) \right\rangle = F^{(x)}_{a\beta}\, m^2_\beta(x) \;+\; \mathcal{O}(p^2).$$

Proof. Step 1: Decay constant at band level. By the band–decay definition (eq. (7.719)), the improved band current couples to one–particle pseudo–NG states as

$$\left\langle \Omega^{(x)} \right| \bar{J}^{(x)\mu}_{a,\rm imp}(0) \left| \pi^{(x)}_\beta(p) \right\rangle = i\, F^{(x)}_{a\beta}\, p^\mu \;+\; \mathcal{O}(p^3), \tag{7.720}$$

where the remainder encodes higher–derivative (irrelevant) couplings suppressed at small p. Here $\bar{J}^{(x)\mu}_{a,\rm imp}$ is the cone–preserving, BRST–improved band current (Definition 7.551 and Proposition 7.573), and $|\pi^{(x)}_\beta(p)\rangle$ is normalized in the usual one–particle way.

Step 2: Taking the divergence (PCAC step). Let $\nabla^{(\rm ad)}_\mu$ denote the ad–covariant divergence entering the band Ward identity. Since the UV/IR taggers act by c–number spacetime convolution, they commute with both $\nabla^{(\rm ad)}_\mu$ and the BRST charge ζ ($[\nabla^{(\rm ad)}_\mu, K^{(\ell)}_{\rm IR}] = [\zeta, K^{(\ell)}_{\rm IR}] = 0$ and likewise for P_x; Definition 7.551 and Proposition 7.573). Hence differentiation can be pulled through the descent.

Fourier transforming (7.720) and evaluating at $x = 0$ gives

$$\left\langle \Omega^{(x)} \right| \nabla^{(\rm ad)}_\mu \bar{J}^{(x)\mu}_{a,\rm imp}(0) \left| \pi^{(x)}_\beta(p) \right\rangle = ip_\mu \left\langle \Omega^{(x)} \right| \bar{J}^{(x)\mu}_{a,\rm imp}(0) \left| \pi^{(x)}_\beta(p) \right\rangle = F^{(x)}_{a\beta}\, p^2 \;+\; \mathcal{O}(p^2),$$

where the last estimate uses (7.720): contracting p_μ with the $\mathcal{O}(p^3)$ remainder yields $\mathcal{O}(p^4)$, which we absorb in $\mathcal{O}(p^2)$ once we go on shell.

Step 3: Band–PCAC operator and on–shell reduction. By definition, the band–level PCAC operator is the gauge–invariant, scheme–independent combination

$$\Delta^{(x)}_a \;:=\; \nabla^{(\rm ad)}_\mu \bar{J}^{(x)\mu}_{a,\rm imp} \;+\; \zeta(B^{(x)}_a),$$

with $B^{(x)}_a$ the induced spurion–sector scalar from Corollary 7.564. On the physical Hilbert space $\mathcal{H}_{\rm phys}$, BRST–exact insertions vanish between physical states, so $\left\langle \Omega^{(x)} \right| \zeta(B^{(x)}_a) \left| \pi^{(x)}_\beta(p) \right\rangle = 0$ (Lemma 7.552). Thus

$$\left\langle \Omega^{(x)} \right| \Delta^{(x)}_a(0) \left| \pi^{(x)}_\beta(p) \right\rangle = \left\langle \Omega^{(x)} \right| \nabla^{(\rm ad)}_\mu \bar{J}^{(x)\mu}_{a,\rm imp}(0) \left| \pi^{(x)}_\beta(p) \right\rangle = F^{(x)}_{a\beta}\, p^2 \;+\; \mathcal{O}(p^2).$$

Placing the state on shell, $p^2 = m_\beta^2(x)$, gives

$$\left\langle \Omega^{(x)} \right| \Delta_a^{(x)}(0) \left| \pi_\beta^{(x)}(p) \right\rangle = F_{a\beta}^{(x)}\, m_\beta^2(x) \;+\; \mathcal{O}(p^2).$$

Step 4: Gauge/Nielsen independence. Because all band constructions commute with BRST and internal G actions (Proposition 7.573), and physical matrix elements are taken on $\mathcal{H}_{\mathrm{phys}}$, the result is gauge–parameter independent (Proposition 7.556).

This establishes the lemma. □

Proposition 7.627 (GMOR–type matrix identity on the band)**.** *Let $C_k^{(x)}$ be the* source–overlap *matrices with entries $\left[C_k^{(x)}\right]_{a\beta} := \left\langle \Omega^{(x)} \right| \mathcal{S}_{k,a}^{(\lambda_{\mathrm{UV}})} \left| \pi_\beta^{(x)} \right\rangle$. Then, to first order in the spurions,*

$$F^{(x)}\, M_\pi^2(x) \;=\; \sum_k \epsilon_k\, C_k^{(x)} \;+\; \mathcal{O}(\epsilon^2), \tag{7.721}$$

where $F^{(x)} = [F_{a\beta}^{(x)}]$, $M_\pi^2(x) = \mathrm{diag}\left(m_\beta^2(x)\right)$ in the pseudo–NG basis, and the $\mathcal{O}(\epsilon^2)$ term collects higher–order spurion effects (operator/state mixing, wave–function renormalization).

Proof. Step 1: Band Ward identity to $\mathcal{O}(\epsilon)$. On the band and on $\mathcal{H}_{\mathrm{phys}}$, the tagged Ward identity takes the form

$$\Delta_a^{(x)} \;:=\; \nabla_\mu^{(\mathrm{ad})}\, \bar{J}_{a,\mathrm{imp}}^{(x)\mu} \;+\; \zeta(B_a^{(x)}) \;=\; \sum_k \epsilon_k\, \mathcal{S}_{k,a}^{(\lambda_{\mathrm{UV}})} \;+\; \mathcal{O}(\epsilon^2), \tag{7.722}$$

where $\bar{J}_{a,\mathrm{imp}}^{(x)\mu}$ is the BRST–improved band current and $B_a^{(x)}$ the induced scalar from Corollary 7.564. The $\mathcal{O}(\epsilon^2)$ remainder accounts for operator mixing and higher–order deformations under explicit breaking. The UV/IR taggers commute with $\nabla_\mu^{(\mathrm{ad})}$ and with BRST/G (cone–preserving, c–number action), cf. Definition 7.551 and Proposition 7.573.

Step 2: Sandwich the Ward identity between one–particle states. Insert (7.722) between the band vacuum and a normalized one–pion state:

$$\left\langle \Omega^{(x)} \right| \Delta_a^{(x)}(0) \left| \pi_\beta^{(x)}(p) \right\rangle \;=\; \sum_k \epsilon_k \left\langle \Omega^{(x)} \right| \mathcal{S}_{k,a}^{(\lambda_{\mathrm{UV}})}(0) \left| \pi_\beta^{(x)}(p) \right\rangle \;+\; \mathcal{O}(\epsilon^2). \tag{7.723}$$

By BRST–exactness on $\mathcal{H}_{\mathrm{phys}}$, $\langle \Omega^{(x)} | \zeta(\cdot) | \pi_\beta^{(x)} \rangle = 0$ (Lemma 7.552), so the improvement term in $\Delta_a^{(x)}$ does not contribute.

Step 3: PCAC reduction on the left-hand side. By the band–PCAC lemma Lemma 7.626,

$$\left\langle \Omega^{(x)} \right| \Delta_a^{(x)}(0) \left| \pi_\beta^{(x)}(p) \right\rangle = F_{a\beta}^{(x)}\, m_\beta^2(x) \;+\; \mathcal{O}(p^2).$$

Evaluating on shell ($p^2 = m_\beta^2(x)$) collapses the $\mathcal{O}(p^2)$ remainder to higher–order derivative corrections, which are absorbed into the stated $\mathcal{O}(\epsilon^2)$ once the expansion in explicit breakings is organized at fixed $x \in \mathfrak{P}$ (the pole dominance of the one–particle contribution is ensured by the cone–preserving taggers and the certified window).

Step 4: Identify the source–overlap matrix on the right-hand side. Define $[C_k^{(x)}]_{a\beta} := \langle \Omega^{(x)} | \, \mathcal{S}_{k,a}^{(\lambda_{\mathrm{UV}})} \, | \pi_\beta^{(x)} \rangle$ as in the statement. Then (7.723) reads, componentwise in (a, β),

$$F_{a\beta}^{(x)} \, m_\beta^2(x) \;=\; \sum_k \epsilon_k \left[C_k^{(x)} \right]_{a\beta} \;+\; \mathcal{O}(\epsilon^2).$$

Collecting the entries over a and β yields the matrix identity

$$F^{(x)} \, M_\pi^2(x) \;=\; \sum_k \epsilon_k \, C_k^{(x)} \;+\; \mathcal{O}(\epsilon^2),$$

after choosing the pseudo–NG basis that diagonalizes the one–particle mass matrix, $M_\pi^2(x) = \mathrm{diag}(m_\beta^2(x))$ (existence and analytic labeling along x follow from Lemma 7.619 and Proposition 7.620).

Step 5: Remarks on higher orders and gauge/BRST safety. Corrections of order ϵ^2 arise from: (i) second–order operator mixing in the Ward operator, (ii) spurion–induced state deformations beyond the one–pole approximation, and (iii) wave–function/mass renormalization beyond linear order. All such contributions are uniformly controlled on certified plateaus and are gathered in $\mathcal{O}(\epsilon^2)$. Since the taggers commute with BRST/G and matrix elements are taken on $\mathcal{H}_{\mathrm{phys}}$, the identity is gauge–parameter independent (Nielsen–stable) by Proposition 7.556.

The proposition follows. □

Corollary 7.628 (Diagonal (single–mode) GMOR relation; linear spurion scaling)**.** *In a basis where $F^{(x)}$ is diagonal at $\epsilon = 0$ (or after projecting onto a single pseudo–NG mode b), we have*

$$\boxed{ f_{\pi_b}^2(x) \, m_{\pi_b}^2(x) \;=\; \sum_k \epsilon_k \left\langle \mathcal{S}_{k,b}^{(\lambda_{\mathrm{UV}})} \right\rangle_{\boldsymbol{\Psi}} \;+\; \mathcal{O}(\epsilon^2) } \tag{7.724}$$

with $f_{\pi_b}(x) := F_{bb}^{(x)} + O(\epsilon)$. Consequently, on any certified plateau $\mathfrak{P}$, $m_{\pi_b}^2(x) = \alpha_b^{(x)} \cdot \epsilon + \mathcal{O}(\epsilon^2)$ (linear in spurions), uniformly in $x \in \mathfrak{P}$.

Proof. Step 1: Extract the b–mode from the matrix GMOR. From Proposition 7.627,

$$F^{(x)} \, M_\pi^2(x) \;=\; \sum_k \epsilon_k \, C_k^{(x)} \;+\; \mathcal{O}(\epsilon^2), \qquad [C_k^{(x)}]_{a\beta} = \left\langle \Omega^{(x)} \right| \mathcal{S}_{k,a}^{(\lambda_{\mathrm{UV}})} \left| \pi_\beta^{(x)} \right\rangle.$$

Project onto the broken direction b (or choose a basis where $F^{(x)}$ is diagonal at $\epsilon = 0$):

$$F_{bb}^{(x)} \, m_{\pi_b}^2(x) \;=\; \sum_k \epsilon_k \left[C_k^{(x)} \right]_{bb} \;+\; \mathcal{O}(\epsilon^2). \tag{7.725}$$

Define $f_{\pi_b}(x) := F_{bb}^{(x)} + O(\epsilon)$; analyticity of spectral data (Lemma 7.619 and Proposition 7.620) implies $f_{\pi_b}(x)$ depends smoothly on ϵ on $\mathfrak{P}$.

Step 2: LSZ/PCAC reduction of the vacuum–pion matrix element. For the b–channel, the soft–pion (PCAC) reduction on the band gives, at small momentum and to leading order,

$$\left\langle \Omega^{(x)} \right| \mathcal{S}_{k,b}^{(\lambda_{\mathrm{UV}})}(0) \left| \pi_b^{(x)}(p) \right\rangle \;=\; f_{\pi_b}(x) \left\langle \mathcal{S}_{k,b}^{(\lambda_{\mathrm{UV}})} \right\rangle_{\boldsymbol{\Psi}} \;+\; \mathcal{O}(p^2) \;+\; \mathcal{O}(\epsilon). \tag{7.726}$$

Indeed, the usual LSZ/PCAC commutator reduction (soft–theorem) yields $\langle\Omega|\mathcal{S}_{k,b}|\pi_b\rangle = \frac{i}{f_{\pi_b}}\langle\Omega|[Q_b,\mathcal{S}_{k,b}]|\Omega\rangle + O(m^2_{\pi_b})$; in the band construction the broken–channel commutator reproduces (up to fixed conventions absorbed in $\mathcal{S}_{k,b}$) the order–parameter insertion, i.e. the vacuum expectation $\langle\mathcal{S}_{k,b}\rangle_{\mathbf{\Psi}}$, with BRST–/improvement pieces dropping out on $\mathcal{H}_{\rm phys}$ (Lemma 7.552 and Proposition 7.556) and cone–preserving taggers ensuring single–pole dominance (Propositions 7.573 and 7.612). Evaluated on shell ($p^2 = m^2_{\pi_b}(x)$), the $\mathcal{O}(p^2)$ term is subleading and can be absorbed into $\mathcal{O}(\epsilon^2)$ once (7.725) is used.

Thus, to the accuracy of (7.725),

$$\left[C_k^{(x)}\right]_{bb} \;=\; \left\langle \Omega^{(x)}\right|\mathcal{S}_{k,b}^{(\lambda_{\rm UV})}\left|\pi_b^{(x)}\right\rangle \;=\; f_{\pi_b}(x)\left\langle \mathcal{S}_{k,b}^{(\lambda_{\rm UV})}\right\rangle_{\mathbf{\Psi}} \;+\; \mathcal{O}(\epsilon).$$

Step 3: Conclude the scalar GMOR and linear scaling. Insert the last expression into (7.725):

$$f_{\pi_b}(x)\, m^2_{\pi_b}(x) \;=\; \sum_k \epsilon_k\, f_{\pi_b}(x)\left\langle \mathcal{S}_{k,b}^{(\lambda_{\rm UV})}\right\rangle_{\mathbf{\Psi}} \;+\; \mathcal{O}(\epsilon^2).$$

Cancel $f_{\pi_b}(x)$ (nonzero on certified $\mathfrak{P}$), obtaining the boxed relation (7.724). Since $f_{\pi_b}(x) = F^{(x)}_{bb} + O(\epsilon)$ and $\langle\mathcal{S}^{(\lambda_{\rm UV})}_{k,b}\rangle_{\mathbf{\Psi}}$ are ϵ–smooth and uniformly bounded on $\mathfrak{P}$, the right–hand side is *linear* in ϵ up to $\mathcal{O}(\epsilon^2)$. Therefore

$$m^2_{\pi_b}(x) \;=\; \sum_k \alpha^{(x)}_{b,k}\,\epsilon_k \;+\; \mathcal{O}(\epsilon^2), \qquad \alpha^{(x)}_{b,k} \;:=\; \left.\frac{\left\langle \mathcal{S}_{k,b}^{(\lambda_{\rm UV})}\right\rangle_{\mathbf{\Psi}}}{f^2_{\pi_b}(x)}\right|_{\epsilon=0},$$

with the coefficients continuous in x; plateau compactness and the mask floors imply uniformity on $\mathfrak{P}$.

Gauge/BRST robustness. All ingredients are constructed on $\mathcal{H}_{\rm phys}$; taggers commute with BRST/G (Proposition 7.573), and Nielsen–independence of spectral/PCAC data holds (Proposition 7.556). Thus (7.724) and the linear scaling are gauge–parameter independent and BRST–safe.

The corollary follows. □

Proposition 7.629 (Band–Hessian ↔ Ward–source consistency)**.** *Let $\mathbb{H}^{(\lambda_{\rm UV},\ell)}$ be the band Hessian (mass/mixing ledger) of Section 7.9, projected to the pseudo–NG subspace to give $M^2_\pi(x)$. Then (7.721) is equivalent to matching the* linear *spurion block of $\mathbb{H}^{(\lambda_{\rm UV},\ell)}$ against the right–hand side of eq. (7.665):*

$$F^{(x)}\left(M^2_{\pi,0}(x) + \sum_k \epsilon_k\,\delta M^2_{\pi,k}(x)\right) = \sum_k \epsilon_k\, C_k^{(x)} \;+\; \mathcal{O}(\epsilon^2),$$

with $M^2_{\pi,0}(x) = 0$ in the exact–symmetry limit, so $M^2_\pi(x) = \sum_k \epsilon_k\,\delta M^2_{\pi,k}(x) + \mathcal{O}(\epsilon^2)$. Hence $\delta M^2_{\pi,k}(x)$ extracted from the ledger must equal $F^{(x)-1}C_k^{(x)}$ to leading order.

Proof. Step 1: Ledger decomposition and projection to the pseudo–NG subspace. Fix $x \in \mathfrak{P}$. Write the (band) Hessian on the relevant sector as an analytic perturbation in the spurions:

$$\mathbb{H}(\epsilon) \;:=\; \mathbb{H}_0 \;+\; \sum_k \epsilon_k\,\delta\mathbb{H}_k \;+\; \mathcal{O}(\epsilon^2),$$

where $\mathbb{H}_0$ has an r–dimensional *Goldstone* kernel (exact symmetry), and $\delta\mathbb{H}_k$ are the linear spurion insertions (all objects are those of Section 7.9). Let $\mathsf{P}(\epsilon)$ be the Riesz projector onto the pseudo–NG cluster (near 0) and $\mathcal{E}(\epsilon) = \operatorname{Ran}\mathsf{P}(\epsilon)$ the corresponding subspace. By Lemma 7.619 and Proposition 7.620, both $\mathsf{P}(\epsilon)$ and analytic frames of $\mathcal{E}(\epsilon)$ are real–analytic in ϵ.

Define the compressed (projected) mass/mixing matrix

$$M_\pi^2(\epsilon) \;:=\; \Big(\mathsf{P}(\epsilon)\,\mathbb{H}(\epsilon)\,\mathsf{P}(\epsilon)\Big)\Big|_{\mathcal{E}(\epsilon)}.$$

At $\epsilon = 0$, $M_{\pi,0}^2 := M_\pi^2(0) = 0$ because $\mathbb{H}_0$ vanishes on the Goldstone subspace.

Step 2: First–order variation in Kato gauge. Differentiate $M_\pi^2(\epsilon)$ at $\epsilon = 0$ along the k–th spurion. Using $\frac{d}{d\epsilon_k}\big(\mathsf{P}\,\mathbb{H}\,\mathsf{P}\big) = (\partial_k\mathsf{P})\,\mathbb{H}\,\mathsf{P} + \mathsf{P}\,(\partial_k\mathbb{H})\,\mathsf{P} + \mathsf{P}\,\mathbb{H}\,(\partial_k\mathsf{P})$, and the identities for projectors (Kato calculus) $\mathsf{P}_0\,\dot{\mathsf{P}}_0\,\mathsf{P}_0 = 0$ with $\mathsf{P}_0 := \mathsf{P}(0)$, plus $\mathbb{H}_0\,\mathsf{P}_0 = 0 = \mathsf{P}_0\,\mathbb{H}_0$ on the Goldstone space, we obtain

$$\delta M_{\pi,k}^2 \;:=\; \partial_{\epsilon_k} M_\pi^2(\epsilon)\Big|_{\epsilon=0} \;=\; \mathsf{P}_0\,\delta\mathbb{H}_k\,\mathsf{P}_0 \quad \text{on} \quad \mathcal{E}(0). \tag{7.727}$$

In an analytically transported orthonormal basis $\{|\pi_\beta^{(x)}\rangle\}_{\beta=1}^r$ of $\mathcal{E}(0)$ (Kato parallel transport, Proposition 7.620), (7.727) reads componentwise

$$\Big[\delta M_{\pi,k}^2(x)\Big]_{\beta\beta'} \;=\; \Big\langle \pi_\beta^{(x)}\Big|\,\delta\mathbb{H}_k\,\Big|\pi_{\beta'}^{(x)}\Big\rangle.$$

Therefore, to linear order,

$$M_\pi^2(x) \;=\; \sum_k \epsilon_k\,\delta M_{\pi,k}^2(x) \;+\; \mathcal{O}(\epsilon^2), \qquad M_{\pi,0}^2(x) = 0. \tag{7.728}$$

Step 3: Ward identity matrix elements (GMOR) and identification. On the other hand, by the band Ward identity and PCAC reduction, Proposition 7.627 and Lemma 7.626 give the GMOR–type matrix relation

$$F^{(x)}\,M_\pi^2(x) \;=\; \sum_k \epsilon_k\,C_k^{(x)} \;+\; \mathcal{O}(\epsilon^2), \tag{7.729}$$

with $F^{(x)} = [F_{a\beta}^{(x)}]$ and $[C_k^{(x)}]_{a\beta} = \langle\Omega^{(x)}|\,\mathcal{S}_{k,a}^{(\lambda_{\mathrm{UV}})}\,|\pi_\beta^{(x)}\rangle$.

Insert the linear decomposition (7.728) into (7.729):

$$F^{(x)}\sum_k \epsilon_k\,\delta M_{\pi,k}^2(x) \;=\; \sum_k \epsilon_k\,C_k^{(x)} \;+\; \mathcal{O}(\epsilon^2).$$

By linear independence of the $\{\epsilon_k\}$, the $\mathcal{O}(\epsilon)$ coefficients must match *for each* k:

$$F^{(x)}\,\delta M_{\pi,k}^2(x) \;=\; C_k^{(x)} \qquad\Longleftrightarrow\qquad \delta M_{\pi,k}^2(x) \;=\; F^{(x)-1}\,C_k^{(x)}. \tag{7.730}$$

Thus the *linear spurion block* of the ledger, when projected to the pseudo–NG subspace, equals $F^{(x)-1}C_k^{(x)}$.

Step 4: Equivalence in both directions. ($\Rightarrow$) If the ledger's linear block satisfies (7.730), then summing over k gives $M_\pi^2(x) = \sum_k \epsilon_k F^{(x)-1} C_k^{(x)} + \mathcal{O}(\epsilon^2)$ and hence (7.729). ($\Leftarrow$) Conversely, if (7.729) holds for all small spurions, differentiation with respect to ϵ_k at $\epsilon = 0$ forces (7.730), hence (7.727) identifies the ledger's linear block with $F^{(x)-1} C_k^{(x)}$.

Step 5: Gauge/BRST robustness and plateau uniformity. All constructions are on $\mathcal{H}_{\text{phys}}$; the UV/IR taggers commute with BRST/G and preserve cones (Propositions 7.573 and 7.612), while spectral data and Riesz projectors are gauge–parameter independent (Proposition 7.644). The analytic subspace tracking guarantees smooth x–dependence (Lemma 7.619 and Proposition 7.620); certified–plateau bounds ensure the identifications are uniform on $\mathfrak{P}$.

Combining Steps 1–5 proves the proposition. □

Theorem 7.630 (Ward–mass link on certified plateaus (GMOR control))**.** *Assume QIV hypotheses, Stage–I BRST gate, and a certified plateau $\mathfrak{P}$. Then the pseudo–NG masses satisfy, uniformly on $\mathfrak{P}$,*

$$M_\pi^2(x) \;=\; \sum_k \epsilon_k \, F^{(x)-1} C_k^{(x)} \;+\; \mathcal{O}(\epsilon^2),$$

and, in the single–mode projection b, the scalar GMOR relation (7.724). In particular, $m_{\pi_b}^2(x)$ scales linearly *with the spurions $\{\epsilon_k\}$ up to $\mathcal{O}(\epsilon^2)$, with coefficients that are BRST/Nielsen safe and vary at most by the plateau tolerance $\eta_{\mathfrak{P}}$.*

Proof. Step 1: Matrix GMOR and ledger linearization. On the band and on $\mathcal{H}_{\text{phys}}$, the Ward/PCAC analysis yields the matrix GMOR relation

$$F^{(x)} \, M_\pi^2(x) \;=\; \sum_k \epsilon_k \, C_k^{(x)} \;+\; \mathcal{O}(\epsilon^2), \tag{7.731}$$

(Proposition 7.627). Independently, projecting the band Hessian (mass/mixing ledger) to the pseudo–NG subspace and expanding to first order in spurions gives

$$M_\pi^2(x) \;=\; \sum_k \epsilon_k \, \delta M_{\pi,k}^2(x) + \mathcal{O}(\epsilon^2), \qquad \delta M_{\pi,k}^2(x) \;=\; F^{(x)-1} C_k^{(x)} \quad \text{(componentwise)}, \tag{7.732}$$

(Proposition 7.629).

Step 2: Invertibility and uniform conditioning of $F^{(x)}$. The matrix $F^{(x)} = [F_{a\beta}^{(x)}]$ collects current–to–pseudo–NG residues (band decay constants). On a certified plateau, the IR taggers have a strictly positive kernel floor and the T–odd channel carries a nonzero spectral weight (Lemma 7.575, Proposition 7.602, and Corollary 7.677); thus each broken current couples to the corresponding pseudo–NG sector with strictly positive residue. Consequently

$$\sigma_{\min}\!\left(F^{(x)}\right) \;\geq\; f_\star \;>\; 0, \qquad \|F^{(x)}\| \;\leq\; f^\star \;<\; \infty \qquad (x \in \mathfrak{P}),$$

for plateau–uniform constants $f_\star, f^\star$ (compactness of $\mathfrak{P}$ and analyticity of spectral data; see Lemma 7.619 and Proposition 7.620). Hence $F^{(x)}$ is invertible with $\|F^{(x)-1}\| \leq f_\star^{-1}$ uniformly on $\mathfrak{P}$.

Step 3: Ward–mass link and uniform $\mathcal{O}(\epsilon^2)$ control. Left–multiplying (7.731) by $F^{(x)-1}$ and using Step 2,

$$M_\pi^2(x) \;=\; \sum_k \epsilon_k\, F^{(x)-1} C_k^{(x)} \;+\; \mathcal{O}(\epsilon^2),$$

with the $\mathcal{O}(\epsilon^2)$ remainder bounded uniformly on $\mathfrak{P}$ (all ingredients are uniformly bounded and analytic, and $\|F^{(x)-1}\|$ is uniformly controlled). This proves the first claim.

Step 4: Single–mode projection and scalar GMOR. Projecting onto a single mode b (or working in a basis in which $F^{(x)}$ is diagonal at $\epsilon = 0$) reduces the matrix identity to

$$f_{\pi_b}(x)\, m_{\pi_b}^2(x) \;=\; \sum_k \epsilon_k \left\langle \Omega^{(x)} \middle| \mathcal{S}_{k,b}^{(\lambda_{\mathrm{UV}})} \middle| \pi_b^{(x)} \right\rangle \;+\; \mathcal{O}(\epsilon^2).$$

Using the standard LSZ/PCAC reduction on the band to replace the one–pion matrix element by the order–parameter expectation in the b–channel (up to $\mathcal{O}(\epsilon)$; see Corollary 7.628), we obtain the scalar GMOR formula

$$f_{\pi_b}^2(x)\, m_{\pi_b}^2(x) \;=\; \sum_k \epsilon_k \left\langle \mathcal{S}_{k,b}^{(\lambda_{\mathrm{UV}})} \right\rangle_{\mathbf{\Psi}} \;+\; \mathcal{O}(\epsilon^2),$$

with $f_{\pi_b}(x) = F_{bb}^{(x)} + O(\epsilon)$.

Step 5: Linear scaling and plateau uniformization. From the displayed relations,

$$m_{\pi_b}^2(x) \;=\; \sum_k \alpha_{b,k}^{(x)}\, \epsilon_k \;+\; \mathcal{O}(\epsilon^2), \qquad \alpha_{b,k}^{(x)} \;:=\; \left. \frac{\left\langle \mathcal{S}_{k,b}^{(\lambda_{\mathrm{UV}})} \right\rangle_{\mathbf{\Psi}}}{f_{\pi_b}^2(x)} \right|_{\epsilon=0}.$$

The coefficients $\alpha_{b,k}^{(x)}$ are continuous (real–analytic) functions of x on $\mathfrak{P}$ by Lemma 7.619 and Proposition 7.620; the certified–plateau oscillation bound then implies that their variation across $\mathfrak{P}$ is at most $O(\eta_{\mathfrak{P}})$ (Lemma 7.579). Thus the linear scaling holds uniformly on $\mathfrak{P}$ with coefficients stable to within the plateau tolerance.

Step 6: BRST/Nielsen safety. All quantities are constructed from band–descended, cone–preserving taggers that commute with BRST and the G–action (Proposition 7.573); physical matrix elements are taken on $\mathcal{H}_{\mathrm{phys}}$, and spectral objects are gauge–parameter independent (Proposition 7.556 and Lemma 7.552). Hence the coefficients and statements above are BRST/Nielsen safe.

Combining the steps proves the theorem. □

Corollary 7.631 (Operational GMOR fit on the band)**.** *Fix a broken direction b and vary a single spurion ϵ_j on a certified plateau. Then*

$$m_{\pi_b}^2(x) \;=\; \alpha_{bj}^{(x)}\, \epsilon_j \;+\; o(\epsilon_j), \qquad \alpha_{bj}^{(x)} = \frac{1}{f_{\pi_b}^2(x)} \left\langle \mathcal{S}_{j,b}^{(\lambda_{\mathrm{UV}})} \right\rangle_{\mathbf{\Psi}}.$$

A log–log or linear fit in ϵ_j on the plateau recovers slope 1 (within tolerance), and the intercept vanishes within the statistical/systematic budget; deviations indicate either plateau failure or higher–order spurion effects.

Proof. Step 1: Linearization from the matrix Ward–mass link. By Theorem 7.630,

$$M_\pi^2(x) \;=\; \sum_k \epsilon_k\, F^{(x)-1} C_k^{(x)} \;+\; \mathcal{O}(\epsilon^2).$$

Project onto the single pseudo–NG mode b:

$$m_{\pi_b}^2(x) \;=\; \sum_k \epsilon_k \left[e_b^\top F^{(x)-1} C_k^{(x)} e_b\right] \;+\; \mathcal{O}(\epsilon^2),$$

with e_b the unit vector in the b–channel. Holding all spurions fixed except ϵ_j gives

$$m_{\pi_b}^2(x) \;=\; \underbrace{e_b^\top F^{(x)-1} C_j^{(x)} e_b}_{\alpha_{bj}^{(x)}}\, \epsilon_j \;+\; \mathcal{O}(\epsilon_j^2). \tag{7.733}$$

Step 2: Identify $\alpha_{bj}^{(x)}$ via the scalar GMOR. From the diagonal (single–mode) GMOR relation Corollary 7.628,

$$f_{\pi_b}^2(x)\, m_{\pi_b}^2(x) \;=\; \sum_k \epsilon_k \left\langle \mathcal{S}_{k,b}^{(\lambda_{\mathrm{UV}})} \right\rangle_{\mathbf{\Psi}} \;+\; \mathcal{O}(\epsilon^2).$$

Varying only ϵ_j and comparing with (7.733) yields

$$f_{\pi_b}^2(x)\, \alpha_{bj}^{(x)} \;=\; \left\langle \mathcal{S}_{j,b}^{(\lambda_{\mathrm{UV}})} \right\rangle_{\mathbf{\Psi}}, \qquad \text{i.e.} \quad \alpha_{bj}^{(x)} \;=\; \frac{1}{f_{\pi_b}^2(x)} \left\langle \mathcal{S}_{j,b}^{(\lambda_{\mathrm{UV}})} \right\rangle_{\mathbf{\Psi}}.$$

Analyticity of spectral/projector data on certified plateaus (Lemma 7.619 and Proposition 7.620) implies that $f_{\pi_b}(x)$ and the vacuum expectation on the right are smooth in x; hence the identification is uniform on $\mathfrak{P}$.

Step 3: Little–o control and plateau uniformity. Since the remainder in (7.733) is $\mathcal{O}(\epsilon_j^2)$ and coefficients vary at most by the plateau oscillation $\eta_{\mathfrak{P}}$ (Lemma 7.579), we can write

$$m_{\pi_b}^2(x) \;=\; \alpha_{bj}^{(x)}\, \epsilon_j \;+\; o(\epsilon_j) \qquad (\epsilon_j \to 0),$$

uniformly in $x \in \mathfrak{P}$.

Step 4: Consequences for operational fits. Linear fit: Regress $m_{\pi_b}^2(x)$ against ϵ_j on the plateau. From the exact–symmetry limit, $m_{\pi_b}^2(x)\big|_{\epsilon=0} = 0$, so the true intercept is 0; the fitted intercept is therefore bounded by the systematic/statistical budget ($O(\eta_{\mathfrak{P}})$ plus sampling noise), while the fitted slope converges to $\alpha_{bj}^{(x)}$ up to $o(1)$ as the maximal ϵ_j in the window shrinks.

Log–log fit: For $\epsilon_j > 0$,

$$\log m_{\pi_b}^2(x) \;=\; \log \alpha_{bj}^{(x)} \;+\; \log \epsilon_j \;+\; \log\bigl(1 + o(1)\bigr),$$

so the log–log slope tends to 1 with an $o(1)$ deviation controlled by the window size in ϵ_j; the vertical offset estimates $\log \alpha_{bj}^{(x)}$ (plateau–uniformly up to $O(\eta_{\mathfrak{P}})$).

Step 5: Robustness (BRST/Nielsen). All objects are band–descended and computed on $\mathcal{H}_{\mathrm{phys}}$; the taggers commute with BRST/G, and spectral data are gauge–parameter independent (Propositions 7.556 and 7.573 and Lemma 7.552). Hence the coefficients $\alpha_{bj}^{(x)}$ are BRST/Nielsen safe.

Therefore the stated expansion and operational fitting statements hold; significant deviations indicate either (i) plateau failure (violating the certification/oscillation bounds) or (ii) visible higher–order spurion effects beyond the linear window. □

7.9.12 Multi–spurion GMOR Fits and PCAC/LSZ Bandwidth on the Plateau

Context. On $\mathfrak{P}$ the matrix GMOR identity holds (linear in the spurions) *on the physical sector*:

$$F^{(x)}\, M_\pi^2(x;\epsilon) \;=\; \sum_k \epsilon_k\, C_k^{(x)} \;+\; \mathcal{O}(\epsilon^2), \qquad x \in \mathfrak{P}, \tag{7.734}$$

with $F^{(x)}$ the (positive definite) decay–constant matrix, M_π^2 the pNG mass matrix, and $C_k^{(x)}$ the Ward–source matrices defined in Propositions 7.627 and 7.629 and Corollary 7.628. Gauge–parameter independence holds by Proposition 7.644, and type–(A) analyticity by Proposition 7.642.

Definition 7.632 (Multi–spurion linear model and design)**.** Fix $x \in \mathfrak{P}$ and perform N runs with spurion vectors $\epsilon^{(n)} = (\epsilon_1^{(n)}, \dots, \epsilon_K^{(n)})$, $n = 1, \dots, N$, all within the linear regime. For each run, measure $Y^{(n)} := F^{(x)} M_\pi^2(x;\epsilon^{(n)})$ and (optionally) symmetrize it if M_π^2 is symmetric in the chosen basis.

Let $\operatorname{svec}(\cdot)$ denote half–vectorization of symmetric matrices and $p := B(B{+}1)/2$ the length (with B the number of broken directions). Define

$$\mathbf{y}^{(n)} := \operatorname{svec} Y^{(n)} \in \mathbb{R}^p, \qquad \mathbf{c}_k := \operatorname{svec} C_k^{(x)} \in \mathbb{R}^p.$$

Stack $\mathbf{y}^{(n)}$ into $\mathbf{y} \in \mathbb{R}^{Np}$ and form the block design matrix

$$\mathbf{X} \;:=\; \begin{bmatrix} \epsilon_1^{(1)}\mathbb{I}_p & \cdots & \epsilon_K^{(1)}\mathbb{I}_p \\ \vdots & \ddots & \vdots \\ \epsilon_1^{(N)}\mathbb{I}_p & \cdots & \epsilon_K^{(N)}\mathbb{I}_p \end{bmatrix} \in \mathbb{R}^{Np \times Kp}, \qquad \mathbf{c} \;:=\; \begin{bmatrix} \mathbf{c}_1 \\ \vdots \\ \mathbf{c}_K \end{bmatrix} \in \mathbb{R}^{Kp}.$$

Then (7.734) becomes the linear model

$$\mathbf{y} \;=\; \mathbf{X}\,\mathbf{c} \;+\; \mathbf{r}, \qquad \|\mathbf{r}\| = \mathcal{O}(\|\epsilon\|^2). \tag{7.735}$$

Proposition 7.633 (Constrained weighted least squares; intercept–free)**.** *Let* $\mathbf{W} = \operatorname{diag}(w_1\mathbb{I}_p, \dots, w_N\mathbb{I}_p)$ *be a positive weight matrix (inverse variances of* $\operatorname{svec} Y^{(n)}$*). The GMOR estimate of the source tensors is*

$$\widehat{\mathbf{c}} \;:=\; \arg\min_{\mathbf{c}} \; \big\|\mathbf{W}^{1/2}\big(\mathbf{y} - \mathbf{X}\mathbf{c}\big)\big\|_2^2,$$

with no intercept (the $\epsilon = 0$ *datum is* $\mathbf{y}^{(0)} = \mathbf{0}$ *on* $\mathcal{H}_{\mathrm{phys}}$*, within* $\eta_{\mathfrak{P}}$*). If* $\mathbf{X}$ *has full column rank, then*

$$\widehat{\mathbf{c}} = (\mathbf{X}^\top \mathbf{W}\, \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{W}\, \mathbf{y},$$

and the normal–equation residual is

$$\mathbf{r}_{\mathrm{WLS}} \;:=\; \mathbf{y} - \mathbf{X}\widehat{\mathbf{c}}, \qquad \mathcal{R}_{\mathrm{GMOR}}(x) \;:=\; \frac{\big\|\mathbf{W}^{1/2}\mathbf{r}_{\mathrm{WLS}}\big\|_2}{\sqrt{Np}\,\eta_{\mathfrak{P}}}.$$

Proof. Step 1: Linear GMOR model and stacking. From the matrix GMOR relation (Proposition 7.627 and Theorem 7.630), for each experiment $n = 1, \dots, N$ with spurion vector $\epsilon^{(n)} = (\epsilon_1^{(n)}, \dots, \epsilon_K^{(n)})$ we have, up to $\mathcal{O}(\epsilon^2)$,

$$Y^{(n)} \;:=\; F^{(x)} M_\pi^2(x; \epsilon^{(n)}) \;=\; \sum_{k=1}^{K} \epsilon_k^{(n)}\, C_k^{(x)} \;+\; \mathrm{noise}_{(n)},$$

in the p–dimensional readout (e.g. p is the length of svec). Vectorize with $\mathbf{y}^{(n)} := \operatorname{svec} Y^{(n)} \in \mathbb{R}^p$ and $\mathbf{c}_k := \operatorname{svec} C_k^{(x)} \in \mathbb{R}^p$. Then

$$\mathbf{y}^{(n)} \;=\; \begin{pmatrix} \epsilon_1^{(n)}\mathbb{I}_p & \cdots & \epsilon_K^{(n)}\mathbb{I}_p \end{pmatrix} \begin{bmatrix} \mathbf{c}_1 \\ \vdots \\ \mathbf{c}_K \end{bmatrix} + \boldsymbol{\eta}^{(n)} \;=:\; \mathbf{X}^{(n)}\mathbf{c} + \boldsymbol{\eta}^{(n)}.$$

Stacking over n gives $\mathbf{y} := \left(\mathbf{y}^{(1)}; \dots; \mathbf{y}^{(N)}\right) \in \mathbb{R}^{Np}$, $\mathbf{X} := \operatorname{blkrow}(\mathbf{X}^{(1)}, \dots, \mathbf{X}^{(N)}) \in \mathbb{R}^{Np \times Kp}$, and $\mathbf{c} := (\mathbf{c}_1; \dots; \mathbf{c}_K) \in \mathbb{R}^{Kp}$:

$$\mathbf{y} \;=\; \mathbf{X}\,\mathbf{c} \;+\; \boldsymbol{\eta}.$$

Step 2: Intercept–free constraint from physics. At $\epsilon = 0$ the exact–symmetry limit yields $M_\pi^2 = 0$ on $\mathcal{H}_{\text{phys}}$; hence $Y^{(0)} = 0$ and $\mathbf{y}^{(0)} = \mathbf{0}$ within the plateau tolerance $\eta_{\text{Ψ}}$ (Definition 6.300 and Corollary 7.628). Therefore no intercept term is allowed: adding an intercept column $\mathbf{1} \otimes \mathbb{I}_p$ would impose a coefficient constrained to 0 by physics. Equivalently, the constrained problem (intercept fixed to 0) reduces to the unconstrained WLS in $\mathbf{c}$ as stated.

Step 3: WLS normal equations and closed form. Define the weighted loss $L(\mathbf{c}) := \left\| \mathbf{W}^{1/2}(\mathbf{y} - \mathbf{X}\mathbf{c}) \right\|_2^2$, with $\mathbf{W} = \operatorname{diag}(w_1\mathbb{I}_p, \dots, w_N\mathbb{I}_p) \succ 0$ (inverse variances per experiment). The gradient is

$$\nabla L(\mathbf{c}) \;=\; -2\,\mathbf{X}^\top \mathbf{W}\,(\mathbf{y} - \mathbf{X}\mathbf{c}).$$

Setting $\nabla L(\widehat{\mathbf{c}}) = 0$ gives the *normal equations*

$$\mathbf{X}^\top \mathbf{W}\,\mathbf{X}\,\widehat{\mathbf{c}} \;=\; \mathbf{X}^\top \mathbf{W}\,\mathbf{y}.$$

If $\mathbf{X}$ has full column rank, $\mathbf{X}^\top\mathbf{W}\,\mathbf{X}$ is invertible and $\widehat{\mathbf{c}} = (\mathbf{X}^\top\mathbf{W}\,\mathbf{X})^{-1}\mathbf{X}^\top\mathbf{W}\,\mathbf{y}$. Equivalently, *whitening* with $\widetilde{\mathbf{X}} := \mathbf{W}^{1/2}\mathbf{X}$, $\widetilde{\mathbf{y}} := \mathbf{W}^{1/2}\mathbf{y}$ reduces the problem to ordinary least squares, whose solution is unique under $\operatorname{rank}(\widetilde{\mathbf{X}}) = \operatorname{rank}(\mathbf{X}) = Kp$.

Step 4: Residual orthogonality and expression. The WLS residual is $\mathbf{r}_{\text{WLS}} := \mathbf{y} - \mathbf{X}\widehat{\mathbf{c}}$. By the normal equations,

$$\mathbf{X}^\top \mathbf{W}\,\mathbf{r}_{\text{WLS}} \;=\; \mathbf{0},$$

i.e. $\mathbf{r}_{\text{WLS}}$ is $\mathbf{W}$–orthogonal to the column space of $\mathbf{X}$. In whitened coordinates, $\widetilde{\mathbf{r}} = \widetilde{\mathbf{y}} - \widetilde{\mathbf{X}}\widehat{\mathbf{c}}$ satisfies $\widetilde{\mathbf{X}}^\top\widetilde{\mathbf{r}} = 0$ and $\|\mathbf{W}^{1/2}\mathbf{r}_{\text{WLS}}\|_2 = \|\widetilde{\mathbf{r}}\|_2$.

Step 5: Plateau–normalized misfit. Define the dimensionless misfit

$$\mathcal{R}_{\text{GMOR}}(x) \;:=\; \frac{\|\mathbf{W}^{1/2}\mathbf{r}_{\text{WLS}}\|_2}{\sqrt{Np}\,\eta_{\text{Ψ}}}.$$

If each component residual in experiment n satisfies the plateau envelope in its natural (weighted) units, $\|\mathbf{W}_n^{1/2}\mathbf{r}^{(n)}\|_2 \le \sqrt{p}\,\eta_{\mathfrak{P}}$ with $\mathbf{W}_n = w_n\mathbb{I}_p$, then

$$\|\mathbf{W}^{1/2}\mathbf{r}_{\mathrm{WLS}}\|_2^2 = \sum_{n=1}^{N} \|\mathbf{W}_n^{1/2}\mathbf{r}^{(n)}\|_2^2 \;\le\; Np\,\eta_{\mathfrak{P}}^2,$$

so $\mathcal{R}_{\mathrm{GMOR}}(x) \le 1$. Thus the normalization compares the weighted misfit directly to the certified plateau tolerance budget.

Step 6: Rank–deficient remark (completeness). If $\mathbf{X}$ is not full column rank, the minimizer exists but is not unique; the minimum–norm solution is $\widehat{\mathbf{c}} = \mathbf{X}_{\mathbf{W}}^{\dagger}\,\mathbf{y}$ with the $\mathbf{W}$–weighted Moore–Penrose pseudoinverse $\mathbf{X}_{\mathbf{W}}^{\dagger} := (\mathbf{X}^{\top}\mathbf{W}\,\mathbf{X})^{\dagger}\mathbf{X}^{\top}\mathbf{W}$. In practice, identifiability requires sufficiently rich spurion variations so that $\mathrm{rank}(\mathbf{X}) = Kp$.

All claims follow. □

Definition 7.634 (Acceptance and conditioning)**.** We *accept* the multi–spurion GMOR fit at $x \in \mathfrak{P}$ if

$$\mathcal{R}_{\mathrm{GMOR}}(x) \;\le\; 1 \quad \text{and} \quad \kappa\Big(\mathbf{X}^{\top}\mathbf{W}\,\mathbf{X}\Big) \;\le\; \kappa_{\max},$$

for a fixed conditioning threshold $\kappa_{\max}$ (documented with the mask). Otherwise we (i) shrink the ϵ–window or (ii) redesign the spurion directions to improve rank/conditioning. Gauge–parameter flatness of $\widehat{\mathbf{c}}$ follows from Proposition 7.644.

Remark 7.635 (Single–spurion slice equals the one–line fit)**.** For a single active spurion component ϵ_j and a Kato–labeled branch b, (7.735) reduces to the constrained one–line fit of Definition 7.623, with slope $\alpha_{bj}^{(x)}$ and intercept bounded by $\eta_{\mathfrak{P}}$.

Definition 7.636 (PCAC/LSZ test function and bandwidth schedule)**.** Let $\phi \in C_c^{\infty}([-1,1])$ be an even, nonnegative bump with $\int_{-1}^{1}\phi(u)\,du = 1$ (e.g. $\phi(u) = c\exp\!\Big(-\frac{1}{1-u^2}\Big)\mathbf{1}_{|u|<1}$). For a bandwidth $\omega_0 > 0$, define the *mollifier*

$$\chi_{\omega_0}(\omega) \;:=\; \frac{1}{\omega_0}\,\phi\!\left(\frac{\omega}{\omega_0}\right), \qquad \operatorname{supp}\chi_{\omega_0} \subset [-\omega_0,\omega_0], \quad \int \chi_{\omega_0} = 1.$$

At a band point $x \in \mathfrak{P}$ set the bandwidth by

$$\omega_0(x) \;:=\; \theta\;\min\Big\{\;\tfrac{1}{2}\,m_{\min}(x)\;,\;\Omega_{\mathrm{IR}}(x)\;\Big\}, \qquad 0<\theta<1, \tag{7.736}$$

where $m_{\min}(x)$ is the smallest pseudo–NG mass at x (from the mass ledger) and $\Omega_{\mathrm{IR}}(x)$ is the instrument IR cutoff implied by the taggers (Definition 7.551); take θ uniform across $\mathfrak{P}$ (e.g. $\theta = 0.3$). Define the tomographic functional

$$\mathcal{T}_a^{(x)}[\chi_{\omega_0(x)}] \;:=\; \int_{\mathbb{R}} \chi_{\omega_0(x)}(\omega)\,\mathsf{S}_a(x;\omega)\,d\omega,$$

with S_a the T–odd spectral density of Definition 7.607.

Lemma 7.637 (Zero–frequency extraction with uniform error control)**.** *With the schedule* (7.736)*, for all* $x \in \mathfrak{P}$*,*

$$\mathcal{T}_a^{(x)}[\chi_{\omega_0(x)}] \;=\; c_0(x)\,\big|\mathcal{R}_a^{(x)}\big| \;+\; \mathcal{O}\!\left(\frac{\omega_0(x)^2}{m_{\min}(x)^2}\right),$$

where $c_0(x) > 0$ *is the probe normalization of Lemma 7.608. The error term is* $O(\theta^2)$ *uniformly on* $\mathfrak{P}$ *(since* $\omega_0 \le \frac{1}{2}\theta\, m_{\min}$*); the statement is BRST/Nielsen safe by Lemma 7.552 and Proposition 7.556.*

Proof. Step 1: Tomographic functional as a mollified evaluation at $\omega = 0$*.* By definition of the tomographic functional (see Definition 7.607 and the construction used in Lemma 7.608 and Proposition 7.609), there exists an even, normalized compactly supported mollifier χ_{ω_0} with

$$\chi_{\omega_0}(\omega) \;=\; \frac{1}{\omega_0}\,\chi\!\left(\frac{\omega}{\omega_0}\right), \qquad \int_{\mathbb{R}} \chi(\omega)\,d\omega = 1, \quad \chi(\omega) = \chi(-\omega),$$

such that

$$\mathcal{T}_a^{(x)}[\chi_{\omega_0}] \;=\; \int_{\mathbb{R}} \chi_{\omega_0}(\omega)\, f_x(\omega)\,d\omega, \qquad f_x(\omega) \;:=\; \frac{Y_x(\omega)}{\mathsf{K}_{\mathrm{spec}}(x;\omega)}. \tag{7.737}$$

Here $Y_x(\omega)$ is the measured T–odd holosymmetric readout (cf. Proposition 7.609), and $\mathsf{K}_{\mathrm{spec}}(x;\omega)$ is the nonnegative spectral kernel with a positive floor near $\omega = 0$ (Lemma 7.575 and Proposition 7.602). On the certified window, f_x is even in ω and C^2 in a neighbourhood of 0.

Step 2: Identification of $f_x(0)$ *and even Taylor expansion.* By the low–frequency Ward matching (Lemma 7.608),

$$f_x(0) \;=\; c_0(x)\,|\mathcal{R}_a^{(x)}|, \qquad c_0(x) > 0.$$

Since f_x is even, its Taylor expansion at 0 reads

$$f_x(\omega) \;=\; f_x(0) \;+\; \frac{1}{2} f_x''(\xi_{x,\omega})\,\omega^2,$$

for some $\xi_{x,\omega}$ between 0 and ω.

Step 3: Bounding f_x'' *by the mass gap (two equivalent routes).* Assume the would–be NG sector is gapped with minimal band mass $m_{\min}(x) > 0$ (the gap parameter that also enters (7.736)). Then f_x is real–analytic on $|\omega| < m_{\min}(x)$ (retarded correlators are analytic away from the cut, and the denominator $\mathsf{K}_{\mathrm{spec}}$ is positive and smooth near 0). Two standard bounds give

$$\sup_{|\omega|\le\omega_0} |f_x''(\omega)| \;\le\; \frac{C_*(x)}{m_{\min}(x)^2}, \qquad 0 < \omega_0 < m_{\min}(x), \tag{7.738}$$

with $C_*(x)$ locally bounded on the plateau:

(a) *Cauchy estimate (analytic disk).* For any $r < m_{\min}(x)$, by analyticity of f_x on $|z| \le r$, $|f_x''(0)| \le \frac{2}{r^2}\sup_{|z|=r}|f_x(z)|$, and similarly for $|f_x''(\omega)|$ by translating the centre. Taking $r = m_{\min}(x)$ up to a fixed factor and using plateau–uniform bounds on $|f_x|$ (from tagger normalization and Definition 6.300) gives (7.738).

(b) *Spectral (Källén–Lehmann) estimate.* Writing the retarded correlator through its spectral density supported on $|\Omega| \geq m_{\min}(x)$ and expanding around $\omega = 0$, evenness cancels the linear term and yields $f_x''(0) = 2\int d\Omega\, \rho_x(\Omega)\, \Omega^{-3}\, g_x(\Omega)$, with g_x bounded by tagger/probe constants; hence $|f_x''(0)| \leq C \int \rho_x(\Omega)/\Omega^3 \leq C'/m_{\min}(x)^2$.

Either route gives (7.738). By certified–plateau regularity and compactness, $C_*(x)$ is uniformly bounded on $\mathfrak{P}$: $\sup_{x\in\mathfrak{P}} C_*(x) =: C_* < \infty$.

Step 4: Mollifier's second moment and the averaged error. Let $\mu_2(\chi) := \int \omega^2 \chi(\omega)\, d\omega < \infty$. For the scaled mollifier, $\int \omega^2 \chi_{\omega_0}(\omega)\, d\omega = \mu_2(\chi)\, \omega_0^2$. Insert the Taylor remainder into (7.737) and use evenness ($\int \omega\, \chi_{\omega_0} = 0$):

$$\left|\mathcal{T}_a^{(x)}[\chi_{\omega_0}] - f_x(0)\right| \;\leq\; \frac{1}{2} \sup_{|\omega|\leq\omega_0} |f_x''(\omega)| \int \omega^2 \chi_{\omega_0}(\omega)\, d\omega \;\leq\; \frac{1}{2}\, \frac{C_*}{m_{\min}(x)^2}\, \mu_2(\chi)\, \omega_0^2.$$

Thus

$$\mathcal{T}_a^{(x)}[\chi_{\omega_0}] \;=\; f_x(0) \;+\; \mathcal{O}\!\left(\frac{\omega_0^2}{m_{\min}(x)^2}\right) \;=\; c_0(x)\, |\mathcal{R}_a^{(x)}| \;+\; \mathcal{O}\!\left(\frac{\omega_0^2}{m_{\min}(x)^2}\right).$$

Step 5: Uniform $O(\theta^2)$ bound under the bandwidth schedule. By the schedule (7.736), $\omega_0(x) \leq \frac{1}{2}\theta\, m_{\min}(x)$ with fixed $0 < \theta \ll 1$ on $\mathfrak{P}$; hence

$$\frac{\omega_0(x)^2}{m_{\min}(x)^2} \;\leq\; \frac{\theta^2}{4} \;=\; O(\theta^2),$$

uniformly in $x \in \mathfrak{P}$. Absorbing the fixed constants $\frac{1}{2} C_* \mu_2(\chi)$ into the big–O proves the claimed uniform error control.

Step 6: BRST/Nielsen safety. All objects entering $\mathcal{T}_a^{(x)}$ are built from band–descended operators via taggers that commute with BRST/G; expectations are taken on $\mathcal{H}_{\text{phys}}$. Gauge–parameter variations are Nielsen–trivial for these spectral/tomographic quantities (Lemma 7.552 and Proposition 7.556). Therefore both the leading term and the error control are BRST/Nielsen safe.

The lemma follows. □

Proposition 7.638 (Plateau–consistent PCAC/LSZ schedule)**.** *Choose a single $\theta \in (0,1)$ (e.g. 0.3) and set $\omega_0(x)$ by (7.736). Then*

$$\left|\, \mathcal{T}_a^{(x)}[\chi_{\omega_0(x)}] - c_0(x)\, |\mathcal{R}_a^{(x)}| \,\right| \;\leq\; C\, \theta^2 \;+\; \mathcal{O}(\eta_{\mathfrak{P}}), \quad \forall\, x \in \mathfrak{P},$$

for a constant C independent of gauge parameters. Consequently, the T–odd tomographic lower bound of Proposition 7.609 tracks the measured law–deficit uniformly on $\mathfrak{P}$ (Corollary 7.610).

Proof. Step 1: Zero–frequency extraction with uniform error. By Lemma 7.637, for all $x \in \mathfrak{P}$,

$$\mathcal{T}_a^{(x)}[\chi_{\omega_0(x)}] \;=\; c_0(x)\, |\mathcal{R}_a^{(x)}| \;+\; \mathcal{O}\!\left(\frac{\omega_0(x)^2}{m_{\min}(x)^2}\right),$$

with $c_0(x) > 0$ and $m_{\min}(x)$ the minimal band mass in the would–be NG sector.

Step 2: Bandwidth schedule $\Rightarrow O(\theta^2)$ control (uniform on $\mathfrak{P}$). The schedule (7.736) imposes $\omega_0(x) \le \frac{1}{2}\theta\, m_{\min}(x)$ on $\mathfrak{P}$. Hence

$$\frac{\omega_0(x)^2}{m_{\min}(x)^2} \;\le\; \tfrac{1}{4}\,\theta^2,$$

so the remainder in Step 1 is bounded by $C\,\theta^2$ with a constant C independent of x (plateau compactness and the uniform bounds used in Lemma 7.637). This yields

$$\mathcal{T}_a^{(x)}[\chi_{\omega_0(x)}] \;=\; c_0(x)\,|\mathcal{R}_a^{(x)}| \;+\; \mathcal{O}(\theta^2), \qquad x \in \mathfrak{P}. \tag{7.739}$$

Step 3: Plateau envelope between channels. Certified plateaus control cross–channel oscillations at level $\eta_{\mathfrak{P}}$ (Definition 6.300 and the uniformization principle Lemma 7.579). Thus any residual bias from replacing the idealized zero–frequency readout by the finite–band functional on the certified mask contributes at most $\mathcal{O}(\eta_{\mathfrak{P}})$. Combining with (7.739) gives the displayed inequality.

Step 4: Consequence for the tomographic lower bound. By Proposition 7.609, for any band–limited even mollifier supported in $[-\omega_0(x), \omega_0(x)]$,

$$1 - |\Lambda_{\text{law}}^{(x)}| \;\ge\; \kappa_{\text{spec}}(x)\,\mathcal{T}_a^{(x)}[\chi_{\omega_0(x)}] \;-\; \mathcal{O}(\eta_{\mathfrak{P}}),$$

with $\kappa_{\text{spec}}(x) > 0$ on the certified window (Lemma 7.575 and Proposition 7.602). Using Step 2,

$$1 - |\Lambda_{\text{law}}^{(x)}| \;\ge\; \kappa_{\text{spec}}(x)\,c_0(x)\,|\mathcal{R}_a^{(x)}| \;-\; \kappa_{\text{spec}}(x)\,C\,\theta^2 \;-\; \mathcal{O}(\eta_{\mathfrak{P}}),$$

which is precisely the uniform tracking asserted in Corollary 7.610 (up to the controlled $O(\theta^2)$ and plateau envelope).

Step 5: BRST/Nielsen safety. The taggers commute with BRST/G and act by c–number spacetime convolutions; expectations are taken on $\mathcal{H}_{\text{phys}}$; therefore $c_0(x)$, $\kappa_{\text{spec}}(x)$, and the bounds above are gauge–parameter independent (Lemma 7.552 and Proposition 7.556).

The proposition follows. □

Definition 7.639 (GMOR reporting on the mask)**.** On $\mathfrak{P}$ (or $\mathfrak{M}_{\text{mask}}$ when enforced), report:

- **(i)** $\widehat{C}_k^{(x)} = \text{svec}^{-1}(\hat{\mathbf{c}}_k)$ with CIs, together with $\mathcal{R}_{\text{GMOR}}(x)$ of Proposition 7.633.
- **(ii)** The conditioning $\kappa(\mathbf{X}^\top \mathbf{W} \mathbf{X})$ and (if used) ridge parameter for regularization.
- **(iii)** The PCAC/LSZ bandwidth $\omega_0(x)$, the chosen θ, and the uniform error proxy $C\theta^2$.
- **(iv)** A gauge–flatness audit: numerical Nielsen slopes ≈ 0 for the fitted coefficients and the tomographic functional (cf. Proposition 7.644).

Only values inside $\mathfrak{M}_{\text{mask}}$ (Theorem 7.664) should be quoted.

7.9.13 Gauge/BRST Safety & Spectral Shifts

We show that the *band-tagged quadratic kernel* on the physical sector is a self-adjoint analytic (type-A) family in the explicit couplings $\{\epsilon_k\}$, so that isolated poles and residues (equivalently, eigenvalues/eigenprojections of the band Hessian/inverse propagator) vary *analytically* with $\{\epsilon_k\}$ and are *gauge-parameter independent* (Nielsen stability). This underwrites reliable tracking of pseudo–NG masses and mixing angles as we dial the spurions.

Definition 7.640 (Band-tagged kernels on the physical sector)**.** At a band point $x = (\lambda_{\mathrm{UV}}, \ell)$ define the band-tagged quadratic kernel (inverse propagator / second variation) by

$$\mathbb{K}^{(x)}(\epsilon) \;:=\; \Big(\mathbb{K}_0 \;+\; \sum_k \epsilon_k\, \mathbb{V}_k\Big)^{(x)}_{\mathrm{phys}} \;:=\; \mathsf{P}_{\mathrm{phys}}\, K_{\mathrm{IR}}^{(\ell)} P_x \left(\mathbb{K}_0 + \sum_k \epsilon_k \mathbb{V}_k\right) P_x K_{\mathrm{IR}}^{(\ell)}\, \mathsf{P}_{\mathrm{phys}},$$

where $\mathsf{P}_{\mathrm{phys}}$ is the projector onto $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$, and $P_x, K_{\mathrm{IR}}^{(\ell)}$ are the UV/IR taggers of Definition 7.551. The *band Hessian* (mass/mixing ledger) is the restriction of $\mathbb{K}^{(x)}(\epsilon)$ to the equal-time quadratic form (or to the one-particle sector), which we denote $\mathbb{H}^{(x)}(\epsilon)$ (cf. Section 7.9).

Lemma 7.641 (Self-adjointness and domain independence)**.** *Assume $\mathbb{K}_0$ is self-adjoint on a dense domain $\mathcal{D} \subset \mathcal{H}_{\mathrm{phys}}$, and each perturbation $\mathbb{V}_k$ is symmetric and $\mathbb{K}_0$-bounded with relative bound < 1 (local polynomial interactions after gauge-fixing/BRST completion). Then, for all ϵ in a neighborhood of 0, $\mathbb{K}^{(x)}(\epsilon)$ is self-adjoint on the* same *domain $\mathcal{D}$; likewise for $\mathbb{H}^{(x)}(\epsilon)$.*

Proof. Step 1: Kato–Rellich setup and smallness of the total perturbation. By $\mathbb{K}_0$-boundedness of each $\mathbb{V}_k$, there exist constants $a_k \in [0, 1)$ and $b_k \geq 0$ such that, for all $\psi \in \mathcal{D}$,

$$\|\mathbb{V}_k \psi\| \;\leq\; a_k\, \|\mathbb{K}_0 \psi\| + b_k\, \|\psi\|. \tag{7.740}$$

Let $\epsilon = (\epsilon_k)$ and set $\mathbb{V}(\epsilon) := \sum_k \epsilon_k \mathbb{V}_k$. Then for all $\psi \in \mathcal{D}$,

$$\|\mathbb{V}(\epsilon)\psi\| \;\leq\; \sum_k |\epsilon_k|\, \|\mathbb{V}_k \psi\| \;\leq\; \Big(\sum_k |\epsilon_k| a_k\Big)\, \|\mathbb{K}_0 \psi\| \;+\; \Big(\sum_k |\epsilon_k| b_k\Big)\, \|\psi\|.$$

Choose a neighborhood $\mathcal{U}$ of 0 in spurion space such that

$$a(\epsilon) := \sum_k |\epsilon_k| a_k \;<\; 1 \qquad \text{for all } \epsilon \in \mathcal{U}.$$

Thus $\mathbb{V}(\epsilon)$ is $\mathbb{K}_0$-bounded with relative bound $a(\epsilon) < 1$ and bound $b(\epsilon) := \sum_k |\epsilon_k| b_k$.

Step 2: Self-adjointness and common domain for the sum. By the Kato–Rellich theorem, the operator

$$\mathbb{K}(\epsilon) \;:=\; \mathbb{K}_0 + \mathbb{V}(\epsilon)$$

with domain $D(\mathbb{K}(\epsilon)) = \mathcal{D}$ is self-adjoint for all $\epsilon \in \mathcal{U}$; moreover $\mathcal{D}$ is a *common* operator domain for the whole family $\{\mathbb{K}(\epsilon)\}_{\epsilon \in \mathcal{U}}$, and any core of $\mathbb{K}_0$ is a core for $\mathbb{K}(\epsilon)$.

Step 3: Restriction to the physical sector and band preparation. Let $\mathsf{P}_{\mathrm{phys}}$ be the orthogonal projector onto $\mathcal{H}_{\mathrm{phys}}$. Stage–I (BRST anomaly–free) implies $[\mathbb{K}(\epsilon), \mathsf{P}_{\mathrm{phys}}] = 0$ (the dynamics

preserves the physical subspace). Therefore the restriction $\mathbb{K}(\epsilon)\restriction_{\mathcal{H}_{\rm phys}}$ is self-adjoint on $\mathcal{D}$ (which we assumed $\subset \mathcal{H}_{\rm phys}$).

The UV/IR taggers $P_x(\lambda_{\rm UV})$ and $K_{\rm IR}^{(\ell)}$ act by bounded c-number spacetime convolution and commute with the local differential/polynomial structure of $\mathbb{K}_0$ and of each $\mathbb{V}_k$ (cf. Definition 7.551 and the locality assumptions). Hence $[P_x, \mathbb{K}(\epsilon)] = [K_{\rm IR}^{(\ell)}, \mathbb{K}(\epsilon)] = 0$ and both operators map $\mathcal{D}$ into itself. Consequently the band-prepared generator,

$$\mathbb{K}^{(x)}(\epsilon) \;:=\; K_{\rm IR}^{(\ell)}\, P_x\, \mathbb{K}(\epsilon)\, P_x\, K_{\rm IR}^{(\ell)},$$

is the restriction of $\mathbb{K}(\epsilon)$ to a reducing subspace (since $P_x K_{\rm IR}^{(\ell)}$ commutes with $\mathbb{K}(\epsilon)$); therefore $\mathbb{K}^{(x)}(\epsilon)$ is self-adjoint on the *same* domain $\mathcal{D}$.

Step 4: The band Hessian $\mathbb{H}^{(x)}(\epsilon)$. The band Hessian $\mathbb{H}^{(x)}(\epsilon)$ (mass/mixing ledger) is obtained as the second variation of the band energy/action functional around the vacuum and is (by construction) a self-adjoint operator on the *form* domain of $\mathbb{K}_0$. In the present setting, each quadratic form perturbation induced by $\mathbb{V}_k$ is *form-bounded* with respect to $\mathbb{K}_0$ with relative bound < 1 (same locality/polynomial assumptions). Hence the KLMN theorem (Kato–Lions–Milgram–Nelson) yields a unique self-adjoint operator associated with the form sum, with *common form domain* equal to that of $\mathbb{K}_0$. Since in our hypotheses we already have (7.740) at the operator level, we can equivalently conclude via Kato–Rellich that $\mathbb{H}^{(x)}(\epsilon)$ is self-adjoint on the *same operator domain* $\mathcal{D}$ (after the same band preparation as in Step 3, which commutes with the dynamics and preserves $\mathcal{D}$).

Step 5: Gauge/BRST robustness. The commuting properties used above stem from the BRST completion and the fact that the taggers act by bounded spacetime convolutions that commute with internal (G) and BRST actions; expectations and domains are taken on $\mathcal{H}_{\rm phys}$. Therefore the statements are gauge-parameter independent and Nielsen-stable.

Combining the steps proves that, for all ϵ in a neighborhood of 0, both $\mathbb{K}^{(x)}(\epsilon)$ and $\mathbb{H}^{(x)}(\epsilon)$ are self-adjoint on the *same* domain $\mathcal{D}$. □

Proposition 7.642 (Type-A analyticity in the spurions)**.** *The family $\epsilon \mapsto \mathbb{K}^{(x)}(\epsilon)$ (and its equal-time reduction $\mathbb{H}^{(x)}(\epsilon)$) is an analytic family of type (A) in the sense of Kato: the domain is ϵ-independent and the map is norm-analytic on $\mathcal{D}$ (polynomial in ϵ). Consequently, the resolvent $(z - \mathbb{K}^{(x)}(\epsilon))^{-1}$ is analytic in ϵ for z off the real axis, and for every simple isolated eigenvalue $\lambda_j^{(x)}(\epsilon)$ the eigenvalue and Riesz projector are real-analytic in ϵ.*

Proof. Step 1: Common domain (Kato–Rellich). By Lemma 7.641, $\mathbb{K}^{(x)}(\epsilon)$ is self-adjoint on the *same* dense domain $\mathcal{D} \subset \mathcal{H}_{\rm phys}$ for all ϵ in a neighbourhood of 0. Thus the family has a ϵ-independent domain.

Step 2: Affine (hence analytic) dependence on ϵ. Before band preparation, write

$$\mathbb{K}(\epsilon) = \mathbb{K}_0 + \sum_{k=1}^{K} \epsilon_k\, \mathbb{V}_k,$$

with each $\mathbb{V}_k$ symmetric and $\mathbb{K}_0$-bounded with relative bound < 1. Band preparation acts by bounded operators that commute with the dynamics on $\mathcal{H}_{\rm phys}$ (Definition 7.551

and Lemma 7.641), so

$$\mathbb{K}^{(x)}(\epsilon) \;=\; B^{(x)}\left(\mathbb{K}_0 + \sum_{k=1}^{K} \epsilon_k\, \mathbb{V}_k\right) B^{(x)}\Big|_{\mathcal{H}_{\mathrm{phys}}}, \qquad B^{(x)} := K_{\mathrm{IR}}^{(\ell)} P_x,$$

and therefore still

$$\mathbb{K}^{(x)}(\epsilon) \;=\; \mathbb{K}_0^{(x)} + \sum_{k=1}^{K} \epsilon_k\, \mathbb{V}_k^{(x)} \quad \text{on } \mathcal{D},$$

with $\mathbb{K}_0^{(x)} := B^{(x)}\mathbb{K}_0 B^{(x)}$ and $\mathbb{V}_k^{(x)} := B^{(x)}\mathbb{V}_k B^{(x)}$. Hence for every $\psi \in \mathcal{D}$ the map $\epsilon \mapsto \mathbb{K}^{(x)}(\epsilon)\psi$ is *affine* (polynomial of degree 1) in ϵ, i.e. entire vector-valued analytic. Equivalently, with the graph norm $\|\psi\|_{\mathcal{D}} := \|\psi\| + \|\mathbb{K}_0^{(x)}\psi\|$, $\epsilon \mapsto \mathbb{K}^{(x)}(\epsilon) \in \mathcal{L}(\mathcal{D}, \mathcal{H}_{\mathrm{phys}})$ is norm-analytic (indeed, norm-polynomial).

Step 3: Type-(A) analyticity and resolvent analyticity. A family with fixed domain and operator-valued analytic dependence is a Kato type-(A) family (see, e.g., Kato, *Perturbation Theory for Linear Operators*, Ch. VII). Therefore, for any z in the resolvent set of $\mathbb{K}^{(x)}(\epsilon)$ (in particular, any $z \in \mathbb{C} \setminus \mathbb{R}$ for self-adjoint $\mathbb{K}^{(x)}(\epsilon)$), the resolvent $R^{(x)}(z,\epsilon) := (z - \mathbb{K}^{(x)}(\epsilon))^{-1}$ is analytic in ϵ. Concretely, for $z \notin \mathbb{R}$ and ϵ small enough,

$$R^{(x)}(z,\epsilon) = R_0^{(x)}(z)\left(\mathbf{1} - \sum_k \epsilon_k\, \mathbb{V}_k^{(x)} R_0^{(x)}(z)\right)^{-1} = \sum_{m=0}^{\infty} R_0^{(x)}(z)\Big(\sum_k \epsilon_k\, \mathbb{V}_k^{(x)} R_0^{(x)}(z)\Big)^m,$$

with $R_0^{(x)}(z) := (z - \mathbb{K}_0^{(x)})^{-1}$, which converges in operator norm by $\|(z - \mathbb{K}_0^{(x)})^{-1}\| \le |\operatorname{Im} z|^{-1}$ and smallness of ϵ; this exhibits holomorphy via a Neumann series.

Step 4: Analyticity of isolated eigenvalues and Riesz projectors. Let $\Sigma(\epsilon)$ be a compact, isolated part of the spectrum of $\mathbb{K}^{(x)}(\epsilon)$ (e.g. a simple eigenvalue $\lambda_j^{(x)}(\epsilon)$) with a contour Γ that encloses $\Sigma(\epsilon)$ and no other spectral points for ϵ in a small neighbourhood. The Riesz projector

$$\mathsf{P}_j^{(x)}(\epsilon) \;=\; \frac{1}{2\pi i} \oint_{\Gamma} (z - \mathbb{K}^{(x)}(\epsilon))^{-1}\, dz$$

is analytic in ϵ because the integrand is analytic jointly in (z,ϵ) and the contour is fixed. If the enclosed eigenvalue is simple, then the eigenvalue $\lambda_j^{(x)}(\epsilon)$ and the corresponding (normalized) eigenvector/eigenprojection depend real-analytically on ϵ (Kato's analytic perturbation theory; eigenvalue/eigenprojection are given by holomorphic functionals of the resolvent).

Step 5: Equal-time reduction $\mathbb{H}^{(x)}(\epsilon)$. The equal-time band Hessian (mass/mixing ledger) has the same analytic structure:

$$\mathbb{H}^{(x)}(\epsilon) \;=\; \mathbb{H}_0^{(x)} \;+\; \sum_k \epsilon_k\, \mathbb{W}_k^{(x)} \;+\; \sum_{k,\ell} \epsilon_k \epsilon_\ell\, \mathbb{W}_{k\ell}^{(x)} \;+\; \cdots,$$

where each coefficient operator is $\mathbb{H}_0^{(x)}$-bounded (arising from local polynomial interactions and finite-order counterterms on $\mathcal{H}_{\mathrm{phys}}$). This yields a power-series (hence analytic) dependence on ϵ with fixed domain; all conclusions above (resolvent/eigenvalue/projector analyticity) apply verbatim.

Combining Steps 1–5 establishes that $\epsilon \mapsto \mathbb{K}^{(x)}(\epsilon)$ and $\epsilon \mapsto \mathbb{H}^{(x)}(\epsilon)$ are analytic families of type (A), and proves the stated consequences. □

Lemma 7.643 (Gauge-parameter flow is BRST-exact on $\mathcal{H}_{\mathrm{phys}}$)**.** *Let α_{gf} denote any gauge-fixing parameter. Then*

$$\partial_{\alpha_{\mathrm{gf}}} \mathbb{K}^{(x)}(\epsilon, \alpha_{\mathrm{gf}}) \;=\; \zeta\Big(\mathbb{B}^{(x)}_{\alpha_{\mathrm{gf}}}\Big) \;+\; \partial_\mu(\cdots),$$

for some local insertion $\mathbb{B}^{(x)}_{\alpha_{\mathrm{gf}}}$ (Nielsen identity, Proposition 7.556), where the total-derivative improvement is invisible in spectral data on $\mathcal{H}_{\mathrm{phys}}$.

Proof. Step 1: BRST doublet and gauge fermion. Work in the algebraic-renormalization framework with nilpotent BRST differential ζ (Stage-I gate). Introduce the *gauge fermion* $\Psi(\alpha_{\mathrm{gf}})$ so that the gauge-fixing+ghost action is $S_{\mathrm{gf+gh}} = \zeta\,\Psi(\alpha_{\mathrm{gf}})$. The gauge parameter α_{gf} enters as a BRST doublet with an external source χ (of opposite statistics): $\zeta\,\alpha_{\mathrm{gf}} = \chi$, $\zeta\,\chi = 0$. Hence

$$\partial_{\alpha_{\mathrm{gf}}} S_{\mathrm{gf+gh}} = \partial_{\alpha_{\mathrm{gf}}}\,\zeta\Psi = \zeta\,\Big(\partial_{\alpha_{\mathrm{gf}}}\Psi\Big), \tag{7.741}$$

which is the off-shell (algebraic) content behind the Nielsen identity; see [117].

Step 2: From action to generators (operator statement). Passing to the operator formalism, variations of the microscopic generator (Hamiltonian/Schwinger generator) with respect to α_{gf} are insertions of the local composite operator obtained by the rule (7.741). Thus there exists a local $\mathbb{B}_{\alpha_{\mathrm{gf}}}$ such that

$$\partial_{\alpha_{\mathrm{gf}}} \mathbb{K}(\epsilon, \alpha_{\mathrm{gf}}) \;=\; \zeta\Big(\mathbb{B}_{\alpha_{\mathrm{gf}}}\Big) \;+\; \partial_\mu(\cdots). \tag{7.742}$$

The divergence term is a Noether/improvement piece (total derivative) that does not affect equal-time spectral data and, after spacetime smearing, integrates to boundary terms.

Step 3: Band descent preserves exactness. Band preparation uses the bounded, cone-preserving taggers $B^{(x)} := K^{(\ell)}_{\mathrm{IR}} P_x$ and restriction to the physical subspace by the orthogonal projector $\mathsf{P}_{\mathrm{phys}}$. By construction, $[B^{(x)}, \zeta] = 0$ and $[B^{(x)}, \partial_\mu] = 0$ (Definition 7.551 and Proposition 7.573), and Stage-I implies that $\mathsf{P}_{\mathrm{phys}}$ commutes with both ζ and the dynamics on $\mathcal{H}_{\mathrm{phys}}$. Therefore, differentiating the band generator $\mathbb{K}^{(x)} := \Big(B^{(x)}\,\mathbb{K}\,B^{(x)}\Big)\restriction_{\mathcal{H}_{\mathrm{phys}}}$ gives

$$\partial_{\alpha_{\mathrm{gf}}}\,\mathbb{K}^{(x)}(\epsilon, \alpha_{\mathrm{gf}}) = \Big(B^{(x)}\Big[\partial_{\alpha_{\mathrm{gf}}}\mathbb{K}(\epsilon, \alpha_{\mathrm{gf}})\Big]B^{(x)}\Big)\restriction_{\mathcal{H}_{\mathrm{phys}}} = \zeta\Big(B^{(x)}\mathbb{B}_{\alpha_{\mathrm{gf}}}B^{(x)}\Big) + \;\partial_\mu(\cdots).$$

Define $\mathbb{B}^{(x)}_{\alpha_{\mathrm{gf}}} := B^{(x)}\mathbb{B}_{\alpha_{\mathrm{gf}}}B^{(x)}$. This establishes the displayed identity in the lemma.

Step 4: Vanishing insertions on $\mathcal{H}_{\mathrm{phys}}$. Let $|\Phi\rangle, |\Psi\rangle \in \mathcal{H}_{\mathrm{phys}}$ be BRST-closed (physical) states: $\zeta\,|\Phi\rangle = 0 = \zeta\,|\Psi\rangle$. For any local operator $\mathbb{X}$,

$$\langle\Phi|\;\zeta(\mathbb{X})\;|\Psi\rangle = \langle\Phi|\;[Q_{\mathrm{BRST}}, \mathbb{X}\}_{\pm}\;|\Psi\rangle = 0,$$

since Q_{BRST} annihilates physical states and the graded commutator reduces to boundary terms that vanish on $\mathcal{H}_{\mathrm{phys}}$. Thus *BRST-exact* variations do not change physical matrix elements. The improvement $\partial_\mu(\cdots)$ is also invisible in the spectral data we use: (i) for equal-time Hessians it drops out as a total divergence; (ii) for retarded correlators it becomes a boundary term annihilated by cone-preserving smearing (Proposition 7.612). Consequently, physical spectral quantities (resolvent matrix elements, eigenvalues, Riesz projectors restricted to $\mathcal{H}_{\mathrm{phys}}$) are independent of α_{gf} to all orders consistent with renormalization (Proposition 7.556).

Step 5: Compatibility with band analyticity and plateau certification. Because $\partial_{\alpha_{\rm gf}}\mathbb{K}^{(x)}$ is BRST-exact plus a divergence, it preserves the domain and the type-(A) analytic structure in the spurions (Lemma 7.641 and Proposition 7.642). Certified plateaus ensure uniform bounds that justify exchanging $\partial_{\alpha_{\rm gf}}$ with spectral limits and tomographic smearing used elsewhere.

Combining Steps 1–5 proves the claim. □

Proposition 7.644 (Nielsen (gauge) independence of pole positions and residues)**.** *Let $\lambda_j^{(x)}(\epsilon)$ be an isolated eigenvalue of $\mathbb{H}^{(x)}(\epsilon)$ with normalized eigenvector $|\psi_j^{(x)}(\epsilon)\rangle \in \mathcal{D}$. Then*

$$\partial_{\alpha_{\rm gf}}\,\lambda_j^{(x)}(\epsilon) \;=\; 0, \qquad \partial_{\alpha_{\rm gf}}\left\langle \Omega^{(x)}\right|\bar{J}^{(x)\mu}_{a,\rm imp}\left|\psi_j^{(x)}(\epsilon)\right\rangle \;=\; 0,$$

i.e. pseudo–NG masses and current residues are gauge-parameter independent on $\mathcal{H}_{\rm phys}$.

Proof. Step 1: Differentiability and Hellmann–Feynman. By type-(A) analyticity (Proposition 7.642), $\epsilon \mapsto \mathbb{H}^{(x)}(\epsilon)$ is analytic with an ϵ-independent domain $\mathcal{D}$. Hence eigenpairs $(\lambda_j^{(x)}(\epsilon), |\psi_j^{(x)}(\epsilon)\rangle)$ of an isolated simple pole are real–analytic in any auxiliary parameter for which the generator varies smoothly, such as $\alpha_{\rm gf}$ (Kato's theorem [115]). Differentiating the eigenvalue equation and using the Hellmann–Feynman identity gives

$$\partial_{\alpha_{\rm gf}}\lambda_j^{(x)}(\epsilon) = \left\langle \psi_j^{(x)}(\epsilon)\right|\partial_{\alpha_{\rm gf}}\mathbb{H}^{(x)}(\epsilon,\alpha_{\rm gf})\left|\psi_j^{(x)}(\epsilon)\right\rangle.$$

Step 2: Nielsen identity ⇒ BRST-exact variation. By the algebraic–renormalization Nielsen identity, variations in the gauge-fixing parameters form BRST doublets and generate ζ–exact insertions (plus improvements) [117, 120, 121, 108]. At the band level,

$$\partial_{\alpha_{\rm gf}}\,\mathbb{H}^{(x)}(\epsilon,\alpha_{\rm gf}) = \zeta\Big(\mathbb{C}^{(x)}_{\alpha_{\rm gf}}\Big) \;+\; \partial_\mu(\cdots),$$

and likewise for the time–evolution generator (Lemma 7.643 and Proposition 7.556). Since matrix elements on $\mathcal{H}_{\rm phys}$ kill BRST–exact insertions and total derivatives (Lemma 7.552 and Proposition 7.612), we obtain

$$\partial_{\alpha_{\rm gf}}\lambda_j^{(x)}(\epsilon) = 0.$$

Step 3: Residues—set up and Kato gauge. Define the residue (current coupling) $R_j^{(x)\mu a}(\alpha_{\rm gf}) := \langle\Omega^{(x)}|\,\bar{J}^{(x)\mu}_{a,\rm imp}\,|\psi_j^{(x)}(\epsilon)\rangle$. Choose Kato's parallel–transport gauge in $\alpha_{\rm gf}$ so that $\langle\psi_j^{(x)}\rangle\,\partial_{\alpha_{\rm gf}}\psi_j^{(x)} = 0$ (Proposition 7.620). Then

$$\partial_{\alpha_{\rm gf}}R_j^{(x)\mu a} = \underbrace{\langle\partial_{\alpha_{\rm gf}}\Omega^{(x)}|\,\bar{J}^{(x)\mu}_{a,\rm imp}\,|\psi_j^{(x)}\rangle}_{(A)} + \underbrace{\langle\Omega^{(x)}|\,\partial_{\alpha_{\rm gf}}\bar{J}^{(x)\mu}_{a,\rm imp}\,|\psi_j^{(x)}\rangle}_{(B)} + \underbrace{\langle\Omega^{(x)}|\,\bar{J}^{(x)\mu}_{a,\rm imp}\,|\partial_{\alpha_{\rm gf}}\psi_j^{(x)}\rangle}_{(C)}.$$

Step 4: Vanishing of (B). By the Nielsen identity for composite operators, the gauge–parameter derivative of any BRST–improved, conserved current is BRST–exact up to a divergence [117]:

$$\partial_{\alpha_{\rm gf}}\bar{J}^{(x)\mu}_{a,\rm imp} = \zeta\Big(\mathbb{U}^{(x)\mu}_{a,\alpha_{\rm gf}}\Big) + \partial_\nu(\cdots).$$

Thus $(B) = 0$ on $\mathcal{H}_{\rm phys}$ (Lemma 7.552).

Step 5: Vanishing of (A) *and* (C). For the vacuum, the Nielsen identity implies the *state-level* BRST doublet relation

$$\partial_{\alpha_{\rm gf}} |\Omega^{(x)}\rangle = \zeta \, |\eta^{(x)}_{\alpha_{\rm gf}}\rangle \quad \text{with} \quad \mathfrak{g}\big(\eta^{(x)}_{\alpha_{\rm gf}}\big) = -1,$$

up to improvement terms whose matrix elements vanish [117, 108]. Hence $(A) = 0$ because $\bar{J}^{(x)}_{a,\rm imp}$ is BRST–invariant modulo a divergence (again invisible between physical states).

For (C), differentiate the eigenvalue equation $\mathbb{H}^{(x)} |\psi^{(x)}_j\rangle = \lambda^{(x)}_j |\psi^{(x)}_j\rangle$ with respect to $\alpha_{\rm gf}$ and project onto the orthogonal complement of $|\psi^{(x)}_j\rangle$; Kato perturbation theory gives

$$(\mathbf{1} - \mathsf{P}^{(x)}_j) \, |\partial_{\alpha_{\rm gf}} \psi^{(x)}_j\rangle = (\mathbf{1} - \mathsf{P}^{(x)}_j) \left(\mathbb{H}^{(x)} - \lambda^{(x)}_j\right)^{-1} \left(\partial_{\alpha_{\rm gf}} \mathbb{H}^{(x)}\right) |\psi^{(x)}_j\rangle \,,$$

where $\mathsf{P}^{(x)}_j$ is the Riesz projector (Proposition 7.642). Using $\partial_{\alpha_{\rm gf}} \mathbb{H}^{(x)} = \zeta(\mathbb{C}^{(x)}_{\alpha_{\rm gf}}) + \partial_\mu(\cdots)$, the BRST charge commutes with the dynamics ($[\zeta, \mathbb{H}^{(x)}] = 0$), hence with the reduced resolvent; therefore the right-hand side is $\zeta(\,\cdot\,)$ plus an improvement. Consequently,

$$(C) = \langle \Omega^{(x)}| \, \bar{J}^{(x)\mu}_{a,\rm imp} (\mathbf{1} - \mathsf{P}^{(x)}_j) \, |\partial_{\alpha_{\rm gf}} \psi^{(x)}_j\rangle = 0$$

on $\mathcal{H}_{\rm phys}$, because BRST–exact vectors have zero matrix elements against BRST–invariant insertions (modulo total derivatives), cf. Lemma 7.552 and Proposition 7.612.

Step 6: Conclusion. All three contributions vanish, so $\partial_{\alpha_{\rm gf}} R^{(x)\mu a}_j = 0$. Together with Step 2 this establishes the gauge–parameter independence of both pole positions and residues on $\mathcal{H}_{\rm phys}$.

Citations. Kato type-(A) analyticity and resolvent/projector calculus: [115]. Nielsen identities and BRST doublets/algebraic renormalization: [120, 121, 117]. Physical-state BRST cohomology and vanishing of BRST-exact insertions: [108]. □

Proposition 7.645 (Analytic tracking of mixing angles)**.** *Let* $\mathcal{E}^{(x)}(\epsilon)$ *be an isolated spectral subspace (possibly multi-dimensional) of* $\mathbb{H}^{(x)}(\epsilon)$ *separated by a gap along* $\epsilon \in U \subset \mathbb{R}^n$*. Then the Riesz projector*

$$\mathsf{P}^{(x)}_{\mathcal{E}}(\epsilon) = \frac{1}{2\pi i} \oint_\gamma (z - \mathbb{H}^{(x)}(\epsilon))^{-1} dz$$

is analytic in ϵ *(fixed contour* γ*). Any orthonormal frame obtained by Kato parallel transport in* $\operatorname{Ran} \mathsf{P}^{(x)}_{\mathcal{E}}(\epsilon)$ *varies real-analytically; "mixing angles'' extracted by comparing this frame to a fixed reference basis are analytic functions of* ϵ *(up to constant phases). Gauge independence holds by Proposition 7.644.*

Proof. Step 1: Analyticity of the Riesz projector. By Proposition 7.642, $\epsilon \mapsto \mathbb{H}^{(x)}(\epsilon)$ is a Kato type-(A) analytic family with ϵ-independent domain. Therefore $(z - \mathbb{H}^{(x)}(\epsilon))^{-1}$ is operator-analytic in ϵ for z off the real axis. If $\mathcal{E}^{(x)}(\epsilon)$ is isolated by a fixed contour γ that does not cross the spectrum along $\epsilon \in U$, then

$$\mathsf{P}^{(x)}_{\mathcal{E}}(\epsilon) = \frac{1}{2\pi i} \oint_\gamma (z - \mathbb{H}^{(x)}(\epsilon))^{-1} \, dz$$

is analytic in ϵ by dominated convergence/holomorphic functional calculus (see, e.g., [115], Ch. VII).

Step 2: Kato parallel transport and analytic frames. Fix $\epsilon_0 \in U$ and choose an orthonormal frame $\{|\phi_\ell\rangle\}_{\ell=1}^r$ spanning $\operatorname{Ran} \mathsf{P}_{\mathcal{E}}^{(x)}(\epsilon_0)$. Define the Kato parallel-transported frame $\{|\psi_\ell(\epsilon)\rangle\}_{\ell=1}^r$ by the ODE

$$\partial_{\epsilon_j} |\psi_\ell(\epsilon)\rangle = \left[\partial_{\epsilon_j} \mathsf{P}_{\mathcal{E}}^{(x)}(\epsilon),\, \mathsf{P}_{\mathcal{E}}^{(x)}(\epsilon)\right] |\psi_\ell(\epsilon)\rangle\,, \qquad |\psi_\ell(\epsilon_0)\rangle = |\phi_\ell\rangle\,.$$

Since $\epsilon \mapsto \mathsf{P}_{\mathcal{E}}^{(x)}(\epsilon)$ is analytic and bounded, the right-hand side is analytic, and standard ODE theory in Banach spaces yields a *real-analytic* solution $\epsilon \mapsto |\psi_\ell(\epsilon)\rangle$; orthonormality and spanning of $\operatorname{Ran} \mathsf{P}_{\mathcal{E}}^{(x)}(\epsilon)$ are preserved (Kato's selection theorem, [115]).

Step 3: Overlap matrix and principal (mixing) angles. Let $\Phi = [|\phi_1\rangle \ \cdots \ |\phi_r\rangle]$ and $\Psi(\epsilon) = [|\psi_1(\epsilon)\rangle \ \cdots \ |\psi_r(\epsilon)\rangle]$ be isometries $\mathbb{C}^r \to \mathcal{H}_{\text{phys}}$. The $r \times r$ *overlap matrix*

$$M(\epsilon) := \Phi^\dagger \Psi(\epsilon)$$

is analytic in ϵ (finite-dimensional matrix of analytic scalar products). The (canonical) principal angles $\{\vartheta_\ell(\epsilon)\}_{\ell=1}^r \in [0, \frac{\pi}{2}]$ between the varying subspace $\operatorname{Ran} \Psi(\epsilon)$ and the reference subspace $\operatorname{Ran} \Phi$ are defined by

$$\cos \vartheta_\ell(\epsilon) = \sigma_\ell\Big(M(\epsilon)\Big),$$

the singular values of $M(\epsilon)$ in nonincreasing order. Alternatively (and more invariantly), consider the positive operator on the reference subspace,

$$\mathcal{A}(\epsilon) := \Phi^\dagger \mathsf{P}_{\mathcal{E}}^{(x)}(\epsilon)\, \Phi \in \mathbb{C}^{r\times r}, \qquad \text{self-adjoint and analytic in } \epsilon,$$

whose eigenvalues are $\cos^2 \vartheta_\ell(\epsilon)$ (spectral theorem for principal angles).

Step 4: Analyticity of mixing angles. Since $\mathcal{A}(\epsilon)$ is a finite-dimensional self-adjoint analytic family, its eigenvalues admit *real-analytic* parameterizations on U provided they remain simple (Kato's theory, [115], Thm. 6.1). Thus, away from accidental degeneracies, we can label $\lambda_\ell(\epsilon) = \cos^2 \vartheta_\ell(\epsilon)$ analytically; composition with the analytic map $\vartheta \mapsto \arccos \sqrt{\vartheta}$ on $(0, 1)$ yields real-analytic $\vartheta_\ell(\epsilon)$. In the presence of degeneracies, analytic labeling is still possible for each *isolated eigenvalue cluster* (Riesz projectors for $\mathcal{A}(\epsilon)$ are analytic by Step 1), and within such a cluster one can choose an analytic orthonormal basis (Kato selection); the corresponding angles are then analytic up to a relabeling within the degenerate block. Hence the "mixing angles'' extracted from $M(\epsilon)$ or $\mathcal{A}(\epsilon)$ are real-analytic functions of ϵ (up to constant phases and, in degenerate cases, up to analytic rotations inside the degenerate eigenspaces).

Step 5: Gauge independence. By Proposition 7.644, spectral projectors associated with isolated parts of the spectrum are gauge-parameter independent on $\mathcal{H}_{\text{phys}}$; hence $\mathsf{P}_{\mathcal{E}}^{(x)}(\epsilon)$, $\mathcal{A}(\epsilon)$, and the resulting principal angles are Nielsen-stable.

Combining Steps 1–5 proves the proposition. □

Theorem 7.646 (Gauge/BRST safety and analytic spectral shifts on plateaus)**.** *Assume the Stage-I gate (BRST anomaly-free), the QIV hypotheses, and a certified plateau* $\mathfrak{P}$*. Then, for every* $x \in \mathfrak{P}$*:*

Type-A analyticity.
$\mathbb{H}^{(x)}(\epsilon)$ *forms a self-adjoint analytic family; each isolated pseudo–NG eigenvalue and its Riesz projector are real-analytic in* ϵ *(Proposition 7.642).*

Nielsen independence.
Pseudo–NG masses and current residues are gauge-parameter independent on $\mathcal{H}_{\text{phys}}$ *(Proposition 7.644).*

Mixing reliability.
Mixing angles within any isolated pseudo–NG cluster are real-analytic in ϵ *and gauge independent (Proposition 7.645).*

Uniformity across $x \in \mathfrak{P}$ *follows from plateau tolerances (Definition 6.300) and the band-commutation properties of Definition 7.551; BRST/Nielsen robustness is by Lemma 7.552 and Proposition 7.556.*

Proof. Step 1: Common domain and Type-(A) analyticity. By Lemma 7.641, the band-prepared generators $\mathbb{K}^{(x)}(\epsilon)$ and equal-time band Hessians $\mathbb{H}^{(x)}(\epsilon)$ are self-adjoint on a *common*, ϵ–independent dense domain $\mathcal{D} \subset \mathcal{H}_{\text{phys}}$ for all small spurions. Their ϵ–dependence is affine (polynomial) after the bounded $P_x, K_{\text{IR}}^{(\ell)}$ sandwich and restriction to $\mathcal{H}_{\text{phys}}$ (Definition 7.551), so $\epsilon \mapsto \mathbb{H}^{(x)}(\epsilon)$ is a Kato type-(A) family with fixed domain. Hence the resolvent and any isolated spectral data (eigenvalues, Riesz projectors) are real-analytic in ϵ (Proposition 7.642).

Step 2: Gauge/Nielsen independence of spectral poles and residues. Gauge-parameter flow is BRST-exact up to improvements at the band level (Lemma 7.643). Therefore, for any isolated, normalized eigenpair $(\lambda_j^{(x)}, |\psi_j^{(x)}\rangle)$, Hellmann–Feynman on $\mathcal{H}_{\text{phys}}$ gives $\partial_{\alpha_{\text{gf}}} \lambda_j^{(x)} = \langle \psi_j^{(x)} | \partial_{\alpha_{\text{gf}}} \mathbb{H}^{(x)} | \psi_j^{(x)} \rangle = 0$, since BRST-exact and total-derivative insertions vanish between physical states (Lemma 7.552). The same BRST-exactness and a Kato gauge choice for $|\psi_j^{(x)}\rangle$ imply that current residues $\langle \Omega^{(x)} | \, \bar{J}_{a,\text{imp}}^{(x)\mu} \, | \psi_j^{(x)} \rangle$ are α_{gf}–independent as well (Proposition 7.644). This establishes Item (2).

Step 3: Analytic and gauge-stable mixing angles. Let $\mathcal{E}^{(x)}(\epsilon)$ be an isolated pseudo–NG cluster along ϵ. By Step 1, the Riesz projector $\mathsf{P}_{\mathcal{E}}^{(x)}(\epsilon) = \frac{1}{2\pi i} \oint_\gamma (z - \mathbb{H}^{(x)}(\epsilon))^{-1} dz$ is analytic in ϵ for a fixed contour γ. Kato's parallel-transport ODE inside $\operatorname{Ran} \mathsf{P}_{\mathcal{E}}^{(x)}(\epsilon)$ yields an orthonormal frame that depends real-analytically on ϵ; principal (mixing) angles obtained from overlaps with a fixed reference frame are then real-analytic functions of ϵ (up to constant phases). Gauge independence of $\mathsf{P}_{\mathcal{E}}^{(x)}$ by Proposition 7.644 implies gauge independence of the mixing angles, proving Item (3) (see Proposition 7.645).

Step 4: Uniformity on certified plateaus. Certified plateaus $\mathfrak{P}$ (Definition 6.300) provide: (i) compactness of the accepted mask; (ii) uniform bounds on tagger norms and commutators with BRST/G (Definition 7.551); and (iii) an oscillation envelope $\eta_{\mathfrak{P}}$ controlling cross-x variations. Consequently, the analyticity radii, spectral gaps defining the isolated clusters, and conditioning constants for the Riesz projectors can be chosen uniformly in $x \in \mathfrak{P}$. By

the plateau uniformization principle (Lemma 7.579), all statements in Items (1)–(3) hold uniformly across $\mathfrak{P}$, with any residual x–dependence bounded by $O(\eta_{\mathfrak{P}})$. BRST/Nielsen robustness of all constructions follows from Lemma 7.552 and Proposition 7.556.

Items (1)–(3) and the uniformity claim are thus established. □

Corollary 7.647 (Operational consequence)**.** *As ϵ is dialed on $\mathfrak{P}$, (i) pseudo–NG masses $m^2_{\pi_b}(x,\epsilon)$ admit convergent Taylor fits in ϵ whose linear terms match the GMOR coefficients of Theorem 7.630; (ii) mixing angles admit smooth (real-analytic) drifts with ϵ; (iii) none of these fits change under gauge-parameter variations (within numerical tolerance).*

Proof. (i) Taylor fits and GMOR linear term. Fix $x \in \mathfrak{P}$ and a pseudo–NG mode b. By Item (1) of Theorem 7.646 (Type-A analyticity), the isolated eigenvalue $m^2_{\pi_b}(x,\epsilon)$ is a real-analytic function of ϵ in a neighbourhood of $\epsilon = 0$. Hence there exists $r(x) > 0$ such that the multivariate Taylor series

$$m^2_{\pi_b}(x,\epsilon) \;=\; m^2_{\pi_b}(x,0) \;+\; \sum_j \left(\partial_{\epsilon_j} m^2_{\pi_b}\right)\Big|_{\epsilon=0} \epsilon_j \;+\; \mathcal{O}(\|\epsilon\|^2)$$

converges for $\|\epsilon\| < r(x)$. In the exact-symmetry limit, $m^2_{\pi_b}(x,0) = 0$. By Theorem 7.630 (Ward–mass link), the linear coefficients equal the GMOR slopes:

$$\left(\partial_{\epsilon_j} m^2_{\pi_b}\right)\Big|_{\epsilon=0} \;=\; \alpha^{(x)}_{bj} \;=\; \frac{1}{f^2_{\pi_b}(x)} \left\langle \mathcal{S}^{(\lambda_{\mathrm{UV}})}_{j,b} \right\rangle_{\mathbf{\Psi}},$$

so the fitted linear term matches GMOR. By the plateau uniformization principle (Lemma 7.579) and the uniform analytic control on $\mathfrak{P}$ provided in Theorem 7.646, the radius and the $\mathcal{O}(\|\epsilon\|^2)$ remainder can be chosen uniformly up to the tolerance $\eta_{\mathfrak{P}}$ on the certified mask.

(ii) Analytic drift of mixing angles. Let $\mathcal{E}^{(x)}(\epsilon)$ be the isolated pseudo–NG cluster containing π_b. By Proposition 7.645, the Riesz projector $\mathsf{P}^{(x)}_{\mathcal{E}}(\epsilon)$ and any Kato parallel-transported orthonormal frame inside $\operatorname{Ran}\mathsf{P}^{(x)}_{\mathcal{E}}(\epsilon)$ are real-analytic in ϵ; principal (mixing) angles obtained by comparing this frame against a fixed reference basis are thus real-analytic functions of ϵ (up to constant phases). Therefore, operationally, mixing angles exhibit smooth (real-analytic) drifts as ϵ varies on $\mathfrak{P}$.

(iii) Gauge-parameter independence of the fits. By Item (2) of Theorem 7.646 and Proposition 7.644, both pseudo–NG masses (hence their Taylor coefficients) and residues are gauge-parameter independent on $\mathcal{H}_{\mathrm{phys}}$; Proposition 7.645 further gives gauge independence of mixing angles. Consequently, any regression (e.g. linear or polynomial in ϵ within the analytic radius) fitted to $m^2_{\pi_b}(x,\epsilon)$ or to the mixing angles is invariant under variations of the gauge-fixing parameter α_{gf}, up to the numerical/plateau tolerance budget $\eta_{\mathfrak{P}}$.

These three statements together yield the claimed operational consequences on $\mathfrak{P}$. □

7.9.14 Gauge–flatness Audit and Analytic Tracking Window in $\{\epsilon_k\}$

Context. On the physical sector, poles and residues vary analytically in the explicit couplings (type–A) and are gauge–parameter independent (Nielsen flat) (Propositions 7.642 and 7.644

and Theorem 7.646). We now (i) turn these into *operational tests* at each band point $x \in \mathfrak{P}$, and (ii) define a *certified window* in spurion space where the quadratic Taylor model passes lack-of-fit tests.

Definition 7.648 (Gauge path and stencil)**.** Let $\alpha_{\rm gf} \in \mathbb{R}^q$ denote gauge–fixing parameters. Around a nominal setting α_0 we probe one-dimensional coordinate lines $\alpha_0 \pm h_\alpha \mathbf{e}_\ell$ and $\alpha_0 \pm 2h_\alpha \mathbf{e}_\ell$, $\ell = 1, \dots, q$. Use the 5-point, order-h_α^4 central stencil for a scalar observable $f(\alpha)$:

$$\partial_{\alpha_\ell} f(\alpha_0) \;\approx\; \frac{-f(\alpha_0+2h_\alpha\mathbf{e}_\ell) + 8f(\alpha_0+h_\alpha\mathbf{e}_\ell) - 8f(\alpha_0-h_\alpha\mathbf{e}_\ell) + f(\alpha_0-2h_\alpha\mathbf{e}_\ell)}{12\,h_\alpha}.$$

For complex matrix elements, apply the stencil to Re and Im parts separately and combine in the Euclidean norm.

Definition 7.649 (Step size and tolerance)**.** Choose a dimensionless step $h_\alpha = \eta_{\mathfrak{P}}^{1/2}$ (with a floor $h_\alpha^{\min} = 10^{-3}$) so that truncation error $O(h_\alpha^4)$ is negligible vs. the plateau tolerance $\eta_{\mathfrak{P}}$ while round-off remains small. Define the *Nielsen slope threshold*

$$\epsilon_{\rm Nielsen} \;:=\; c_{\rm N}\,\eta_{\mathfrak{P}}, \qquad c_{\rm N} \in [5, 10].$$

Definition 7.650 (Operational Nielsen slope test)**.** At $x \in \mathfrak{P}$, let $\{\lambda_j^{(x)}\}$ be the tracked poles (eigenvalues) and $\{R_{a,j}^{(x)}\} = \{\langle\Omega|\bar{J}_{a,\rm imp}|\psi_j\rangle\}$ the residues for broken currents a. Compute

$$S_{\rm Nielsen}(x) \;:=\; \max_{j,\ell} \left|\partial_{\alpha_\ell}\lambda_j^{(x)}\right| \;\vee\; \max_{a,j,\ell} \left|\partial_{\alpha_\ell} R_{a,j}^{(x)}\right|$$

with the stencil of Definition 7.648. *Pass* if $S_{\rm Nielsen}(x) \le \epsilon_{\rm Nielsen}$; otherwise *fail*.

Proposition 7.651 (Stability and BRST/Nielsen safety)**.** *The derivatives in Definition 7.650 are computed on $\mathcal{H}_{\rm phys}$ with band taggers that commute with ζ and the G-action (Definition 7.551); therefore the test is BRST-safe (Lemma 7.552) and its acceptance decision is gauge-parameter independent in the ideal limit (Proposition 7.644). Finite h_α and readout noise are controlled by Definition 7.649.*

Proof. Step 1: Objects whose slopes are tested. Let $\mathscr{F}^{(x)}(\alpha_{\rm gf})$ denote any of the observables covered by Definition 7.650 (e.g. pseudo–NG eigenvalues of $\mathbb{H}^{(x)}$, Riesz-projector–based mixing angles, current residues, or any band-limited spectral functional built from resolvents and matrix elements). By construction, all such readouts are formed after band descent with $B^{(x)} := K_{\rm IR}^{(\ell)} P_x$ and evaluated on $\mathcal{H}_{\rm phys}$.

Step 2: Commutation with BRST/G and differentiability. The taggers are bounded spacetime convolutions that commute with BRST and the G-action (Definition 7.551 and Proposition 7.573); hence $[B^{(x)}, \zeta] = [B^{(x)}, \delta_a] = 0$ and $B^{(x)}$ preserves $\mathcal{H}_{\rm phys}$. By type-(A) analyticity (Proposition 7.642), the families $\epsilon \mapsto \mathbb{K}^{(x)}, \mathbb{H}^{(x)}$ and all resolvent functionals are differentiable in any smooth external parameter, in particular $\alpha_{\rm gf}$.

Step 3: BRST-exact gauge flow ⇒ vanishing ideal slopes. The Nielsen identity at the band level (Lemma 7.643) gives $\partial_{\alpha_{\rm gf}} \mathbb{H}^{(x)} = \zeta(\mathbb{C}_{\alpha_{\rm gf}}^{(x)}) + \partial_\mu(\cdots)$. For any physical matrix element $\langle\Phi| \cdot |\Psi\rangle$

with $|\Phi\rangle, |\Psi\rangle \in \mathcal{H}_{\text{phys}}$, BRST-exact insertions and total derivatives vanish (Lemma 7.552 and Proposition 7.612). Therefore, by Hellmann–Feynman and the resolvent calculus,

$$\partial_{\alpha_{\text{gf}}} m_{\pi_b}^2(x) = 0, \quad \partial_{\alpha_{\text{gf}}} \mathsf{P}_{\mathcal{E}}^{(x)}(x) = 0, \quad \partial_{\alpha_{\text{gf}}} \left\langle \Omega^{(x)} \middle| \bar{J}_{a,\text{imp}}^{(x)} \middle| \psi_j^{(x)} \right\rangle = 0,$$

and similarly for any $\mathscr{F}^{(x)}$ built from these spectral objects (chain rule). Thus, in the *ideal* (continuum, noiseless) limit, the true slope $\partial_{\alpha_{\text{gf}}} \mathscr{F}^{(x)}(\alpha_{\text{gf}}) = 0$. This is the content of Nielsen independence (Proposition 7.644) specialized to the observables of the test.

Step 4: Finite-difference estimator, bias and variance. The operational slope uses a finite difference with step h_α from Definition 7.649:

$$\widehat{\partial_\alpha \mathscr{F}^{(x)}} := \begin{cases} \dfrac{\mathscr{F}^{(x)}(\alpha+h_\alpha) - \mathscr{F}^{(x)}(\alpha-h_\alpha)}{2h_\alpha}, & \text{(central)}, \\ \dfrac{\mathscr{F}^{(x)}(\alpha+h_\alpha) - \mathscr{F}^{(x)}(\alpha)}{h_\alpha}, & \text{(forward)}. \end{cases}$$

Since $\mathscr{F}^{(x)}$ is C^3 in α (Step 2), the truncation bias is $O(h_\alpha^2)$ for the central scheme and $O(h_\alpha)$ for the forward scheme, uniformly on the certified mask (plateau compactness). With readout noise level σ_{rd} per evaluation, the variance scales as $\text{Var}(\widehat{\partial_\alpha \mathscr{F}}) \sim c\, \sigma_{\text{rd}}^2 / h_\alpha^2$ for either scheme (with an $O(1)$ constant c depending on whether two or three evaluations are used). The step/tolerance prescription in Definition 7.649 balances these errors so that the combined RMS is $\lesssim \sigma_{\text{tol}}$ and $\sigma_{\text{tol}} \gg \eta_{\mathfrak{P}}$ (to clear the plateau envelope), ensuring stability of the accept/reject decision.

Step 5: Gauge-parameter independence of the acceptance decision. Because the ideal slope vanishes (Step 3), any two gauge-parameter choices yield the *same* acceptance outcome in the limit $h_\alpha \to 0$ and $\sigma_{\text{rd}} \to 0$. With finite h_α and noise, the probability of a flip in the decision is controlled by the RMS error derived in Step 4 and is made negligible by the thresholds stipulated in Definition 7.649. Hence the operational test is BRST-safe and Nielsen-stable as stated.

The proposition follows. □

Remark 7.652 (Practical audit)**.** (i) Verify stencil consistency by halving h_α once; require derivative estimates to agree within $O(h_\alpha^4) + O(\eta_{\mathfrak{P}})$. (ii) Report the per-mode slopes and the maximum $S_{\text{Nielsen}}(x)$. (iii) If the test fails on a subregion of $\mathfrak{P}$, refine the mask or check for anomaly filters (Definition 7.613).

Definition 7.653 (Quadratic Taylor model in the spurions)**.** Along a spurion ray $\epsilon(t) = \epsilon^{(0)} + t\,\hat{\mathbf{v}}$ with small t, model an eigenvalue (or a residue component) $g(t)$ by

$$g(t) \;=\; g_0 \;+\; a\,t \;+\; \tfrac{1}{2}\,b\,t^2 \;+\; \rho(t), \qquad \rho(t) = O(t^3),$$

with WLS (weights from readout errors). For multi-parameter tracking, use $g(\epsilon) = g_0 + \mathbf{a}^\top \epsilon + \frac{1}{2}\,\epsilon^\top B\,\epsilon + \rho(\epsilon)$.

Definition 7.654 (Analytic tracking window)**.** Let $I = [-t_{\max}, t_{\max}]$ be the largest symmetric interval such that:

1. (Lack-of-fit) The quadratic model passes an F-test against a cubic extension at level $1-\beta$ (e.g. 95%), and the normalized residual $\mathcal{R}_{\text{quad}} = \|W^{1/2}\mathbf{r}\|_2/(\sqrt{N}\,\eta_{\mathfrak{P}}) \le 1$.
2. (Coefficient stability) Refit on the halves $[0, t_{\max}]$, $[-t_{\max}, 0]$; require relative drifts of a, b below $c_{\text{stab}}\eta_{\mathfrak{P}}$ (e.g. $c_{\text{stab}} = 5$).
3. (Consistency across directions) For a basis of spurion directions $\{\hat{\mathbf{e}}_k\}$, the unions of their intervals define a polydisc $\mathcal{B}_\epsilon = \{\epsilon : \ |\epsilon_k - \epsilon_k^{(0)}| \le \epsilon_k^{\max}\}$, each $\epsilon_k^{\max}$ coming from $t_{\max}$ on $\hat{\mathbf{e}}_k$.

We call $\mathcal{B}_\epsilon$ the *analytic tracking window* at x.

Proposition 7.655 (Type–A consistency of the window)**.** *By Proposition 7.642, $g(\epsilon)$ is analytic near $\epsilon^{(0)}$; hence a non-empty window $\mathcal{B}_\epsilon$ exists. If $\mathcal{R}_{\text{quad}} \le 1$ and the cubic model is not favored (by AIC_c or an F-test) on $\mathcal{B}_\epsilon$, the quadratic coefficients provide an accurate local parametric tracker for poles/residues, uniformly on $\mathfrak{P}$ up to $O(\eta_{\mathfrak{P}})$.*

Proof. Step 1: Analyticity ⇒ Taylor control and existence of a window. Fix $x \in \mathfrak{P}$ and let $g(\epsilon)$ denote any scalar component of an isolated pole/residue (eigenvalue, current matrix element, or a coordinate of a mixing-angle chart). By Proposition 7.642, g is real–analytic in a ball $B_r(\epsilon^{(0)})$. Write $\Delta\epsilon := \epsilon - \epsilon^{(0)}$. Taylor's theorem with integral remainder gives, for some $M_3(x) < \infty$,

$$g(\epsilon) \ = \ g_0 \ + \ J\cdot\Delta\epsilon \ + \ \tfrac{1}{2}\,\Delta\epsilon^{\top} H\,\Delta\epsilon \ + \ R_3(\epsilon), \qquad |R_3(\epsilon)| \ \le \ M_3(x)\,\|\Delta\epsilon\|^3, \tag{7.743}$$

for all $\epsilon \in B_r$. Choose a radius $0 < \rho \le r$ small enough that $M_3(x)\rho^3$ is within the target truncation budget (fixed below), and define the *window* $\mathcal{B}_\epsilon := \{\epsilon : \|\Delta\epsilon\| \le \rho\}$. Thus $\mathcal{B}_\epsilon \neq \varnothing$ by analyticity.

Step 2: Quadratic WLS as a local parametric tracker. Collect measurements $\{(\epsilon^{(n)}, y_n)\}_{n=1}^N$ with $\epsilon^{(n)} \in \mathcal{B}_\epsilon$, where

$$y_n \ = \ g(\epsilon^{(n)}) \ + \xi_n, \qquad \mathbb{E}[\xi_n] = 0, \quad \operatorname{Var}(\xi_n) = \sigma_n^2,$$

and the noise envelope (systematic+statistical) is controlled at $O(\eta_{\mathfrak{P}})$ by the plateau certification (Definition 6.300, cf. the usage in Proposition 7.592). Fit the quadratic model $q(\epsilon) := \alpha + J\cdot\Delta\epsilon + \frac{1}{2}\Delta\epsilon^{\top}H\,\Delta\epsilon$ by weighted least squares (full-rank design). Standard regression theory with deterministic model misspecification (7.743) yields the coefficient bias

$$\|\widehat{\alpha} - g_0\| + \|\widehat{J} - J\| + \|\widehat{H} - H\| \ = \ O(\rho^3) \ + \ O_{\mathbb{P}}\Big(\max_n \sigma_n\Big),$$

where the $O(\rho^3)$ term comes from the uniform cubic remainder and the stochastic term is controlled by the weights ($\sigma_n = O(\eta_{\mathfrak{P}})$ on a certified mask). Hence, provided ρ is chosen so that $M_3(x)\rho^3 \lesssim \eta_{\mathfrak{P}}$, the quadratic coefficients track the true local Taylor coefficients up to $O(\eta_{\mathfrak{P}})$.

Step 3: Role of $\mathcal{R}_{\text{quad}}$ and the cubic model comparison. Let RSS_2 (resp. RSS_3) be the weighted residual sum of squares for the quadratic (resp. cubic) fit on $\mathcal{B}_\epsilon$. From (7.743), the *best possible* reduction in residual by adding cubic terms is $O(\rho^6)$ (since the neglected signal is

$O(\rho^3)$ in amplitude). Therefore, if the noise floor is $\gtrsim O(\eta_{\mathfrak{P}})$ and the normalization $\mathcal{R}_{\text{quad}} \le 1$ holds (i.e., the quadratic residual sits within the plateau budget), the expected improvement $\text{RSS}_2 - \text{RSS}_3$ is not statistically significant on $\mathcal{B}_\epsilon$. Concretely, AIC_c penalizes the extra cubic parameters by a constant that dominates an $O(\rho^6)$ gain for small enough ρ, and the classical F-test likewise fails to reject the quadratic null once the signal-to-noise is calibrated to the plateau budget. By hypothesis, both diagnostics *do not* favor the cubic model; thus the quadratic specification is operationally consistent on $\mathcal{B}_\epsilon$.

Step 4: Uniformity on $\mathfrak{P}$. On a certified plateau, the analyticity radius and derivative bounds can be chosen uniformly: compactness of the mask and the spectral isolation hypotheses imply $\sup_{x\in\mathfrak{P}} M_3(x) =: M_3 < \infty$ (Lemma 7.619 and Propositions 7.620 and 7.642). Choosing a single $\rho \le \min_{x\in\mathfrak{P}} r(x)$ such that $M_3\rho^3 \lesssim \eta_{\mathfrak{P}}$ makes the remainder control uniform, and the plateau uniformization principle (Lemma 7.579) bounds the cross-x drift of the fitted coefficients by $O(\eta_{\mathfrak{P}})$. Therefore the quadratic coefficients furnish an *accurate, uniform* local tracker for poles/residues on $\mathfrak{P}$, up to $O(\eta_{\mathfrak{P}})$.

Combining the steps proves the proposition. □

Remark 7.656 (Reporting rules)**.** For each $x \in \mathfrak{P}$: (i) report $S_{\text{Nielsen}}(x)$ and pass/fail; (ii) give the window $\mathcal{B}_\epsilon$ (component radii $\epsilon_k^{\max}$), the quadratic coefficients with CIs, the normalized residual $\mathcal{R}_{\text{quad}}$, and the F-test/AIC_c outcomes; (iii) if the window collapses or Nielsen slopes exceed $\epsilon_{\text{Nielsen}}$, shrink the mask or reduce the spurion span and re-audit.

7.9.15 Plateau–mask Certification (Report Only Where Robust)

We formalize the certification protocol that selects *where* in the $(\lambda_{\text{UV}}, \ell)$–grid our readouts are instrument–robust (gauge/basis/scheme safe) and stable under mesh refinement. All diagnostics reported for QIV (§7.9) are then restricted to this certified mask.

Definition 7.657 (Grid and mask)**.** Let $\mathcal{G} = \{\ (\lambda_{\text{UV}}^{(i)}, \ell^{(j)})\ \}$ be the finite sampling grid in scale space, ordered so that $\ell^{(j+1)} > \ell^{(j)}$. A *plateau mask* is a subset $\mathfrak{M}_{\text{mask}} \subset \mathcal{G}$ such that the plateau tolerances of Definition 6.300 hold on every connected component, and all quantities below satisfy the improving–physicality constraints in Definition 7.659.

Definition 7.658 (Physical margins on $\mathcal{H}_{\text{phys}}$)**.** At a band point $x = (\lambda_{\text{UV}}, \ell)$ let:

1. **Mass margin** $\mu(x)$ be the smallest eigenvalue of the pseudo–NG block $M_\pi^2(x)$ of the band Hessian on $\mathcal{H}_{\text{phys}}$ (cf. Section 7.9 and Theorem 7.630). Thus $\mu(x) > 0$ in QIV.
2. **Hyperbolicity margin** $\chi(x)$ be the infimum, over unit covectors $\hat{\mathbf{n}}$, of the minimal positive eigenvalue of the principal symbol of the kinetic operator (time–derivative block) after band descent and physical projection:
$$\chi(x) \ := \ \inf_{\|\hat{\mathbf{n}}\|=1} \ \lambda_{\min}^{+}\Big(\sigma_{\text{prin}}^{(x)}(\hat{\mathbf{n}})\Big),$$
ensuring well–posedness/positivity of the quadratic form (relativistic z=1 or finite z).
3. **Spectral floor (T–odd)** $s_{\text{floor}}(x)$ be the certified low–frequency T–odd weight tied to the Ward source:
$$s_{\text{floor}}(x) \ := \ \kappa_{\text{spec}}(x)\, c_0(x) \max_{a\in\mathfrak{g}_{\text{br}}} \big|\mathcal{R}_a^{(x)}\big|,$$

with $\kappa_{\mathrm{spec}}(x)$ and $c_0(x)$ from Proposition 7.609 and Lemma 7.608, and $\mathcal{R}_a^{(x)}$ from eq. (7.665).

All three margins are computed on $\mathcal{H}_{\mathrm{phys}}$ and are gauge–parameter independent by Propositions 7.556 and 7.644; continuity in the mask follows from Proposition 7.642 and Lemma 7.579.

Definition 7.659 (Improving physicality with ℓ)**.** Given a fixed λ_{UV}, a sequence $\{x_j = (\lambda_{\mathrm{UV}}, \ell^{(j)})\}_{j_0 \le j \le j_1}$ satisfies *improving physicality* if

$$\mu(x_{j+1}) \;\ge\; \mu(x_j) - \eta_{\mathfrak{P}}, \qquad \chi(x_{j+1}) \;\ge\; \chi(x_j) - \eta_{\mathfrak{P}}, \qquad s_{\mathrm{floor}}(x_{j+1}) \;\ge\; s_{\mathrm{floor}}(x_j) - \eta_{\mathfrak{P}},$$

for the plateau tolerance $\eta_{\mathfrak{P}}$ of Definition 6.300. Equivalently, upon tightening tolerances one may enforce monotone nondecreasing behavior.

Definition 7.660 (Instrument–robust observables on the mask)**.** On $\mathfrak{M}_{\mathrm{mask}}$, report only:

- law/state visibilities and their channel splits (holosymmetric overlays, Definition 6.275 and Proposition 7.555);
- exponents and amplitudes from visibility fits (Theorem 7.580 and Corollary 7.581);
- pseudo–NG masses/mixings and GMOR coefficients (Theorem 7.630 and Corollary 7.631);
- T–odd tomography summaries (Definition 7.607 and Theorem 7.614);
- the margins $\mu, \chi, s_{\mathrm{floor}}$ of Definition 7.658.

All are computed on $\mathcal{H}_{\mathrm{phys}}$ (BRST descent Lemma 7.552) and are Nielsen–stable (Propositions 7.556 and 7.644).

Lemma 7.661 (Gauge/basis/scheme robustness on $\mathfrak{M}_{\mathrm{mask}}$)**.** *Let α_{gf} be any gauge–fixing parameter, and let $\mathcal{R}$ denote admissible basis/scheme changes that act by bounded, BRST–compatible similarity transforms after band descent. Then, for $x \in \mathfrak{M}_{\mathrm{mask}}$,*

$$\partial_{\alpha_{\mathrm{gf}}}\{\mu, \chi, s_{\mathrm{floor}}, \Lambda_\bullet, M_\pi^2, F\} \;=\; 0, \qquad \delta_{\mathcal{R}}\{\mu, \chi, s_{\mathrm{floor}}, \Lambda_\bullet, M_\pi^2, F\} \;=\; \mathcal{O}(\eta_{\mathfrak{P}}),$$

i.e. the reported values are gauge–independent and scheme–stable within plateau tolerances.

Proof. Gauge independence. All quantities are computed on $\mathcal{H}_{\mathrm{phys}}$ after band descent by the bounded taggers $B^{(x)} := K_{\mathrm{IR}}^{(\ell)} P_x$, which commute with BRST and the G–action (Definition 7.551 and Proposition 7.573). Gauge–parameter flow is BRST–exact (plus total derivatives) at the band level by the Nielsen identity (Lemma 7.643). Hence any physical matrix element of $\partial_{\alpha_{\mathrm{gf}}}$ of these objects vanishes (Lemma 7.552 and Proposition 7.556):

- μ, χ are spectral margins of the (band) principal symbol/Hessian; their values are spectral data and thus Nielsen–independent (Proposition 7.644).
- M_π^2 (pseudo–NG masses) and F (decay–constant matrix) are isolated spectral poles and residues, already covered by Proposition 7.644.
- s_{floor} and the tomographic kernel floor κ_{spec} are constructed from $\widehat{\kappa}_x$ (taggers) by positive quadratures (Lemma 7.575 and Proposition 7.602); since the taggers commute with BRST/G, these floors are gauge–parameter independent.

- $\Lambda_\bullet$ (state/law overlaps and their T–odd/T–even parts) are built from resolvents and band–descended matrix elements; BRST–exact insertions and improvements drop out on $\mathcal{H}_{\mathrm{phys}}$, hence no gauge dependence (Proposition 7.556).

Therefore $\partial_{\alpha_{\mathrm{gf}}}\{\cdots\} = 0$ on $\mathcal{H}_{\mathrm{phys}}$.

Basis/scheme robustness. Let $\mathcal{R}$ act by a bounded, BRST–compatible similarity S on the band–prepared physical space, $S : \mathcal{H}_{\mathrm{phys}} \to \mathcal{H}_{\mathrm{phys}}$, with S invertible, $[S, \zeta] = 0$, and with the tagger class mapped into itself (admissible scheme change). Then:

- Spectra and spectral gaps are invariant under similarity: $\mathrm{spec}(S^{-1}\mathbb{H}^{(x)}S) = \mathrm{spec}(\mathbb{H}^{(x)})$, hence μ, χ, M_π^2 are unchanged.
- Residues transform covariantly, but their *numerical values* in a fixed physical normalization are invariant because both interpolating operators and states are pushed forward by S; any difference arises only from the *finite* tagger/mesh implementations used in practice.
- s_{floor} and κ_{spec} are positive quadratic functionals of $\widehat{\kappa}_x$; admissible scheme changes preserve the tagger class, so the ideal values are invariant.

On certified masks, the only discrepancies come from discretization and plateau reweightings; these are controlled by the oscillation envelope in Definition 6.300 (uniformization) and by the acceptance/convergence gates (§ "Grid–refinement schedule"): any scheme/basis induced drift of reported numbers is bounded by the budget $O(\eta_{\mathfrak{P}})$ (see also Lemma 7.588, the robustness statements following Proposition 7.594, and the mask thresholds discussion). Hence $\delta_{\mathcal{R}}\{\cdots\} = \mathcal{O}(\eta_{\mathfrak{P}})$.

Combining both parts proves the lemma. □

Lemma 7.662 (Finite–mesh audit: convergence under grid refinement)**.** *Let $\mathcal{G}_h$ be a sequence of grids with mesh size $h \to 0$, and let $\mathfrak{M}_{\mathrm{mask}}^{(h)}$ be the certified masks. If the oscillation bound of Definition 6.300 holds uniformly, then every compact subset of the limiting plateau set is eventually contained in $\mathfrak{M}_{\mathrm{mask}}^{(h)}$, and the reported observables converge uniformly on that subset.*

Proof. Step 1: Equicontinuity and uniform boundedness (plateau envelope). For each observable $Q \in \{\mu, \chi, s_{\mathrm{floor}}, \Lambda_\bullet, M_\pi^2, F, \ldots\}$ define its mesh–h version $Q^{(h)}(x)$ computed on $\mathfrak{M}_{\mathrm{mask}}^{(h)}$. The certified plateau hypothesis gives, for all sufficiently fine h and all x, y in a common certified mask,

$$|Q^{(h)}(x) - Q^{(h)}(y)| \;\le\; \eta_{\mathfrak{P}}\, d(x,y) \quad \text{(after a fixed rescaling of } d),$$

and $|Q^{(h)}(x)| \le C_Q$ uniformly (finite–energy window, tagger bounds). Thus $\{Q^{(h)}\}_h$ is an equicontinuous and uniformly bounded family on any compact K contained in the limiting plateau set.

Step 2: Cauchy property from the refinement gate. Between successive refinement levels the *convergence gate* on the common submask ensures

$$\|Q^{(h)} - Q^{(h/\rho)}\|_{L^\infty(K \cap \mathfrak{M}_{\mathrm{mask}}^{(h)} \cap \mathfrak{M}_{\mathrm{mask}}^{(h/\rho)})} \;\le\; c_{\mathrm{conv}}\, \eta_{\mathfrak{P}} \qquad (c_{\mathrm{conv}} = 2),$$

and the "improving physicality'' inequalities (monotone $\mu, \chi, s_{\text{floor}}$ with refinement) prevent loss of certification under further refinement. Iterating along a geometric mesh sequence shows $\{Q^{(h)}\}_h$ is Cauchy in the sup–norm on K up to the fixed plateau budget; hence there exists a uniform limit $Q^{(\infty)}$ on K.

Step 3: Eventual inclusion of compact subsets. Let K be a compact subset of the limiting plateau. By definition of the limiting set and the monotone "improving physicality'' gate, for all sufficiently small h the margin inequalities and floors (§ "Why these thresholds are not arbitrary", hyperbolicity/mass–gap/spectral floors) continue to hold at the refined resolution on K; therefore $K \subset \mathfrak{M}^{(h)}_{\text{mask}}$ eventually. (Intuitively: the discretization buffer in δ_{gap} scales like $3\,\Delta\lambda_{\text{grid}}(h)$ and shrinks under refinement, so once eligible, points remain eligible; see the monotonicity discussion in that section.)

Step 4: Uniform convergence on K. By Step 1 (equicontinuity) and Step 2 (Cauchy), the Arzelà–Ascoli theorem yields uniform convergence of $Q^{(h)}$ to $Q^{(\infty)}$ on K. The certification inequalities (mass/hyperbolicity margins, spectral floors) are closed under uniform limits, so $Q^{(\infty)}$ remains admissible and the mask stabilizes on K.

Therefore every compact K in the limiting plateau is eventually contained in $\mathfrak{M}^{(h)}_{\text{mask}}$, and all reported observables converge uniformly on K. □

Proposition 7.663 (Construction of the plateau mask)**.** *Given $\mathcal{G}$ and tolerances $(\eta_{\mathfrak{P}}$, fit tolerances$)$, the iterative procedure is as follows:*

- **(i)** *Select all points satisfying the plateau bounds of Definition 6.300.*
- **(ii)** *Prune points that violate Definition 7.659 along any fixed λ_{UV}–line.*
- **(iii)** *Prune isolated points that fail the finite–difference consistency checks for exponents/-margins.*

This produces a nonempty mask $\mathfrak{M}_{\text{mask}}$ such that all observables in Definition 7.660 are instrument–robust (Lemma 7.661) and mesh–convergent (Lemma 7.662).

Proof. Notation. For each grid point $x \in \mathcal{G}$ let $Q(x)$ denote any observable in Definition 7.660 (e.g. $\mu, \chi, s_{\text{floor}}, \Lambda_\bullet, M_\pi^2, F$) computed after band descent. Write

$$\mathcal{S}_1 := \{x \in \mathcal{G} : x \text{ satisfies Definition 6.300}\},$$
$$\mathcal{S}_2 := \{x \in \mathcal{S}_1 : x \text{ satisfies Definition 7.659 along its } \lambda_{\text{UV}}\text{-line}\}.$$

and let $\mathcal{S}_3$ be the subset of $\mathcal{S}_2$ left after removing points that fail the local finite–difference consistency checks (slopes/exponents/margins) with the stated fit tolerances. The algorithm outputs $\mathfrak{M}_{\text{mask}} := \mathcal{S}_3$.

1) Nonemptiness. By Definition 6.300 there exists a certified plateau $\mathfrak{P} \neq \varnothing$ on which the channel oscillations are bounded by $\eta_{\mathfrak{P}}$. All mask observables are continuous (indeed, real–analytic) in the control parameters by type–(A) analyticity and spectral isolation (Propositions 7.620 and 7.642 and Lemma 7.619); hence the acceptance inequalities of Definition 6.300 hold on an open neighbourhood of $\mathfrak{P}$. For any sufficiently fine grid $\mathcal{G}$, there exists at least one $x \in \mathcal{G} \cap \mathfrak{P}$, so $\mathcal{S}_1 \neq \varnothing$.

The inequalities in Definition 7.659 (monotone margins $\mu, \chi, s_{\text{floor}}$ along the λ_{UV}–line) are ensured by the *construction* of the thresholds and the refinement gate: the mass–gap and hyperbolicity reserves are pegged to $5\eta_{\mathfrak{P}}$ and the discretization buffer $3\Delta\lambda_{\text{grid}}$ shrinks geometrically (§ "Why these thresholds are not arbitrary"; see the "Refinement stability" bullet and Proposition 7.602 and Lemma 7.575). Thus, within $\mathcal{S}_1$ there is a nonempty subregion where these inequalities already hold and are preserved under refinement, so $\mathcal{S}_2 \neq \varnothing$.

Finally, by type–(A) analyticity the true derivatives/slopes exist and the finite–difference estimators are consistent on $\mathcal{S}_2$; choosing the fit tolerances as in the slope/visibility tests (Propositions 7.592, 7.594 and 7.599) removes only spurious, noise–induced outliers. Hence $\mathcal{S}_3 \neq \varnothing$.

2) Closure under the certification gates. Membership in $\mathcal{S}_1$ is defined by inequalities stable under $O(\eta_{\mathfrak{P}})$ perturbations and therefore is closed on $\mathcal{G}$ with respect to the plateau envelope; the pruning step (2) enforces additional *monotone* inequalities that are hereditary along each λ_{UV}–line (once satisfied, they remain satisfied under further geometric refinement because the discretization buffer decreases). Step (3) removes points that contradict the local analytic model (quadratic/cubic WLS in ϵ or ℓ); since Proposition 7.655 guarantees a nonempty analytic window where the quadratic tracker is valid and *not* improved by cubic terms, the pruning cannot delete that window. Thus the algorithm stabilizes on a nonempty certified set $\mathfrak{M}_{\text{mask}}$.

3) Instrument robustness on $\mathfrak{M}_{\text{mask}}$. On $\mathfrak{M}_{\text{mask}}$ all quantities are evaluated on $\mathcal{H}_{\text{phys}}$ with taggers that commute with BRST/G (Definition 7.551 and Proposition 7.573). Gauge–parameter variations are BRST–exact (Lemma 7.643), so physical matrix elements, spectra and residues are Nielsen–independent (Proposition 7.644). Admissible basis/scheme changes act by bounded similarities and preserve the ideal values; the only deviations are discretization/-tagger artefacts controlled by the plateau envelope. Consequently, Lemma 7.661 applies and yields instrument–robustness of all observables in Definition 7.660, with scheme drifts $\mathcal{O}(\eta_{\mathfrak{P}})$.

4) Mesh convergence. Because every $x \in \mathfrak{M}_{\text{mask}}$ satisfies Definition 6.300 uniformly and step (2) enforces "improving physicality," the hypotheses of Lemma 7.662 hold: the family of mesh–level observables is equicontinuous and uniformly bounded on compact subsets of the limiting plateau, and the convergence gate on the common submask makes the sequence Cauchy. Hence the reported observables converge uniformly on any compact subset of the limiting plateau and the certified masks eventually contain those subsets (Lemma 7.662).

5) Compatibility with GMOR/visibility diagnostics. Step (3) uses finite–difference and model–selection checks that are precisely those guaranteed by the analytic spectral/GMOR theory on plateaus (Theorems 7.630 and 7.646 and Proposition 7.655): accepted points are exactly those for which the quadratic (or linear-in–spurion) model is adequate within the plateau budget and for which no spurious IR–improvement (QIII) is detected (Proposition 7.599). Therefore the final mask satisfies all consistency requirements used elsewhere in the analysis.

Combining 1)–5) shows that the procedure outputs a nonempty $\mathfrak{M}_{\text{mask}}$ with instrument–robust and mesh–convergent observables, as claimed. □

Theorem 7.664 (Certified reporting on $\mathfrak{M}_{\text{mask}}$)**.** *On the certified mask $\mathfrak{M}_{\text{mask}}$ the following hold uniformly:*

Instrument robustness.
All reported visibilities, exponents, masses/mixings, GMOR slopes, and physical margins are gauge–independent and scheme–stable within $\mathcal{O}(\eta_{\mathfrak{P}})$ (Lemma 7.661).

Improving physicality with ℓ.
μ, χ, s_{floor} are nondecreasing in ℓ up to $\eta_{\mathfrak{P}}$ (Definition 7.659); in particular, $\mu > 0$ (no exact NG) and the T–odd floor persists (QIV law–deficit), consistent with Theorems 7.580 and 7.614.

Mesh convergence.
Reported values converge under grid refinement (Lemma 7.662).

Therefore, all quoted results are restricted to $\mathfrak{M}_{\mathrm{mask}}$, *ensuring that QIV conclusions (persistent law–deficit; unique–vacuum state channel; GMOR linear scaling) are made only where the instrumentation guarantees robustness on $\mathcal{H}_{\mathrm{phys}}$.*

Proof. Preliminaries. The mask $\mathfrak{M}_{\mathrm{mask}}$ is produced by the selection/pruning procedure of Proposition 7.663. Hence every $x \in \mathfrak{M}_{\mathrm{mask}}$ simultaneously satisfies: (i) the plateau envelope and channel oscillation bounds of Definition 6.300; (ii) the *improving physicality* inequalities along each λ_{UV}–line (Definition 7.659); and (iii) local finite–difference/fit consistency with the type–(A) analytic model and the QIV diagnostics (GMOR/visibility), so spurious outliers have been pruned.

(1) Instrument robustness. By Lemma 7.661, all reported observables on $\mathfrak{M}_{\mathrm{mask}}$

$$Q \in \{\mu, \chi, s_{\mathrm{floor}}, \Lambda_{\bullet}, M_{\pi}^2, F, \text{ visibilities/exponents/GMOR slopes}\}$$

are computed after band descent with taggers commuting with BRST/G, and their values are (i) gauge–parameter independent on $\mathcal{H}_{\mathrm{phys}}$ (Nielsen independence) and (ii) stable under admissible basis/scheme changes, up to $\mathcal{O}(\eta_{\mathfrak{P}})$ induced by finite mesh/tagger differences bounded by the plateau envelope. Therefore Item (1) holds uniformly on the mask.

(2) Improving physicality and persistence of the QIV signature. Membership in $\mathfrak{M}_{\mathrm{mask}}$ includes the monotonicity requirements of Definition 7.659: along increasing ℓ (IR coarse–graining) the margins μ (mass gap in the would–be NG sector), χ (hyperbolicity reserve), and the T–odd spectral floor s_{floor} are nondecreasing up to the plateau tolerance $\eta_{\mathfrak{P}}$. In particular, $\mu > 0$ everywhere on the mask, excluding exact NG modes and ensuring QIV (explicit breaking) rather than QII (SSB) behaviour; and $s_{\mathrm{floor}} > 0$ persists, providing a stable law–deficit in the T–odd channel. These statements are precisely the operational content of the QIV diagnostics recorded in Theorem 7.580 and are compatible with the non–signalling/tomographic completion in Theorem 7.614. Hence Item (2) follows.

(3) Mesh convergence. By construction (Proposition 7.663), the certified masks at successive refinement levels obey the convergence gate and "improving physicality'' monotonicity. The hypotheses of Lemma 7.662 are therefore satisfied: uniform oscillation control on the certified sets yields equicontinuity and uniform boundedness of all reported observables; the refinement gate provides a Cauchy property on common submasks; consequently the reported values converge uniformly on compact subsets of the limiting plateau. This proves Item (3).

Conclusion. Items (1)–(3) together imply that all reported QIV conclusions are made *only* on $\mathfrak{M}_{\mathrm{mask}}$, where (i) gauge/BRST robustness and scheme stability are guaranteed within

$\mathcal{O}(\eta_{\mathfrak{P}})$, (ii) physicality margins strengthen with IR scale so that the QIV T–odd signature persists, and (iii) mesh refinement does not change the reported values beyond the plateau budget. This is the claimed certified reporting property. □

7.9.16 Numerical Thresholds and Grid–refinement Schedule for the Mask

Context. We now *fix numbers* for the plateau tolerance and physicality margins and formalize the mesh–refinement audit that underpins Proposition 7.663 and Theorem 7.664. All statements act on $\mathcal{H}_{\text{phys}}$ and inherit BRST/Nielsen safety from Lemma 7.552 and Proposition 7.556.

Definition 7.665 (Fixed thresholds for the mask)**.** Let $\eta_{\mathfrak{P}} \in (0,1)$ be the plateau oscillation tolerance (Definition 6.300). We fix the following *numeric* thresholds:

1. **Plateau tolerance:** $\eta_{\mathfrak{P}} = 10^{-2}$ (1% envelope) by default. Tight variant for precision runs: $\eta_{\mathfrak{P}} = 5 \times 10^{-3}$.
2. **Gap threshold for the mass margin μ (near–degeneracy control):** Let $\Delta\lambda_{\text{grid}}(x)$ be the spectral resolution (eigenvalue discretization uncertainty) for the band Hessian at x. Set
$$\delta_{\text{gap}}(x) \;:=\; \max\Big\{\, 5\,\eta_{\mathfrak{P}}\,\Lambda_{\text{scale}}(x)\;,\; 3\,\Delta\lambda_{\text{grid}}(x)\,\Big\},$$
where $\Lambda_{\text{scale}}(x)$ is the local spectral scale (e.g. the median of nonzero eigenvalues of $\mathbb{H}^{(x)}$). The mass margin constraint in Definition 7.658 is then $\mu(x) \geq \delta_{\text{gap}}(x)$.
3. **Hyperbolicity margin χ:** Let $\chi(x)$ be the smallest *positive* principal–symbol eigenvalue margin preserving well–posedness (Definition 7.658). Impose
$$\chi(x) \;\geq\; \delta_{\text{hyp}}, \qquad \delta_{\text{hyp}} := 5\,\eta_{\mathfrak{P}}.$$
(For models with dynamical exponent $z \neq 1$, interpret the principal symbol with the anisotropic metric.)
4. **Spectral floor (law channel):** Let $s_{\text{floor}}(x)$ denote the minimal calibrated T–odd spectral weight over $|\omega| \leq \omega_0(x)$ (cf. Definitions 7.607 and 7.636). Require
$$s_{\text{floor}}(x) \;\geq\; s_{\min}, \qquad s_{\min} := 3\,\eta_{\mathfrak{P}},$$
and, correspondingly, the kernel floor $\kappa_{\text{spec}}(x)$ to satisfy $\kappa_{\text{spec}}(x) \geq \kappa_{\min} := 2\,\eta_{\mathfrak{P}}$ (cf. calibration in Proposition 7.602 and Definition 7.603).

A band point x is *mask–eligible* if all four inequalities hold.

Remark 7.666 (Units and normalization)**.** The margins μ, χ scale with the chosen spectral norm; we normalize by the local scale $\Lambda_{\text{scale}}(x)$ so that thresholds are dimensionless. The floors s_{floor} and κ_{spec} are already dimensionless (overlap/spectral–kernel quantities).

Definition 7.667 (Grid–refinement schedule)**.** Let $\mathcal{G}^{(m)}$ be the m–th grid on the $(\lambda_{\mathrm{UV}}, \ell)$ domain with spacings $h_\lambda^{(m)}, h_\ell^{(m)}$. The grid is refined geometrically from an initial grid $\mathcal{G}^{(0)}$ as follows:

$$\begin{aligned} h_\lambda^{(m+1)} &= \rho\, h_\lambda^{(m)}, \\ h_\ell^{(m+1)} &= \rho\, h_\ell^{(m)}, \\ \rho &\in \left[\tfrac{1}{2}, \tfrac{2}{3}\right]. \end{aligned}$$

At each refinement, the mask–eligible set $\mathfrak{M}^{(m)}$ is recomputed using the condition from Definition 7.665.

Definition 7.668 (Convergence metric and acceptance)**.** Consider an observable $Q \in \left\{ \Lambda_{\mathrm{law}}, \Lambda_{\mathrm{state}}, \dots \right\}$. The Cauchy–type metric between consecutive grids on their common submask, $\mathfrak{M}^{(m+1)} \subseteq \mathfrak{M}^{(m)}$, is defined by:

$$\Delta^{(m)}[Q] \;:=\; \sup_{x \in \mathfrak{M}^{(m+1)}} \left| Q^{(m+1)}(x) - Q^{(m)}(x) \right|.$$

We *accept convergence* at level $m_\star$ if:

$$\Delta^{(m_\star - 1)}[Q] \;\le\; c_{\mathrm{conv}}\, \eta_{\mathfrak{P}} \quad \text{for all such observables } Q, \text{ with } c_{\mathrm{conv}} = 2,$$

and the *improving–physicality* condition holds, i.e., the quantities $\mu, \chi, s_{\mathrm{floor}}$ are *nondecreasing* with m on the nested masks (Definition 7.659).

Proposition 7.669 (Certified refinement and reporting)**.** *If the schedule of Definition 7.667 and thresholds of Definition 7.665 are enforced, and the acceptance criterion of Definition 7.668 holds at $m_\star$, then the mask $\mathfrak{M}_{\mathrm{mask}} := \mathfrak{M}^{(m_\star)}$ is* instrument–robust*: quoted exponents, masses, and deficits on $\mathfrak{M}_{\mathrm{mask}}$ vary by at most $2\eta_{\mathfrak{P}}$ under one further refinement. Certified reporting per Theorem 7.664 follows.*

Proof. Geometric refinement yields a Cauchy sequence in the sup–norm on the nested masks; the acceptance bound controls discretization error at $O(\eta_{\mathfrak{P}})$. Monotone margins ensure no spurious loss of physicality upon refinement. □

Remark 7.670 (Practical defaults)**.** Use $\eta_{\mathfrak{P}} = 10^{-2}$, $\rho = 1/2$, and stop after two successive accepted levels (with identical decisions for QIV/QIII/QII discrimination across the mask). For tougher problems, lower $\eta_{\mathfrak{P}}$ and raise δ_{gap} proportionally.

Why these thresholds are not arbitrary

The thresholds in Section 7.9.16 are fixed by three principles that are already part of the Chapter VII–VIII machinery: (i) a *single accuracy budget* set on certified plateaus $\mathfrak{P}$ (Definition 7.31 and Proposition 6.298), (ii) *dimensionless normalization* against the local spectral scale and the grid's spectral resolution, and (iii) *monotone refinement* with an explicit convergence gate (Definition 7.668, Proposition 7.663, and Theorem 7.664). Each inequality is just these principles written as numbers.

1) One budget to rule them all (plateau envelope). We first fix the plateau oscillation budget $\eta_{\mathfrak{P}}$ on $\mathfrak{P}$ (default 10^{-2}; tight 5×10^{-3}). This is not a knob for "tuning results"; it is the *unit of numerical significance* that already governs acceptance and Nielsen/BRST robustness on $\mathcal{H}_{\text{phys}}$ (Lemma 7.552 and Proposition 7.556). Every downstream gate is expressed as a small integer multiple of $\eta_{\mathfrak{P}}$. That decision ensures two things: (a) the mask is *dimensionless* and scales if we tighten/loosen the target accuracy, and (b) all channels (spectral T–odd, correlational T–even, kinematics) are compared in the *same currency*.

2) Dimensional correctness and dual buffering (near-degeneracy, hyperbolicity). The *mass-gap threshold*

$$\delta_{\text{gap}}(x) = \max\Big\{\, 5\,\eta_{\mathfrak{P}}\,\Lambda_{\text{scale}}(x),\; 3\,\Delta\lambda_{\text{grid}}(x)\Big\}, \qquad \mu(x) \geq \delta_{\text{gap}}(x),$$

is a direct transcription of dimensional and discretization control: $\Lambda_{\text{scale}}(x)$ (e.g. the median nonzero eigenvalue of the band Hessian) furnishes units, while $\Delta\lambda_{\text{grid}}(x)$ captures the grid's spectral granularity. Taking the *max* makes the decision invariant to whether the ambiguity comes from physics scale or from resolution; scaling μ by Λ_{scale} keeps the inequality dimensionless. The companion *hyperbolicity margin*

$$\chi(x) \geq \delta_{\text{hyp}}, \qquad \delta_{\text{hyp}} := 5\,\eta_{\mathfrak{P}},$$

enforces a positive reserve for the smallest principal-symbol eigenvalue in the first-order reduction, so well-posedness is not at the mercy of the plateau envelope; it is the numerical form of the kinematic sanity checks (tachyon/hyperbolicity) that already gate $\mathcal{A}_{\{\text{law,state}\}}$ (eq. (7.10), Definition 7.535, and Section 7.2.7).

3) Floors that come from physics (T–odd and kernel). QIV's law-deficit is *spectral* in origin at low frequency (Proposition 7.673). We therefore enforce a T–odd spectral *floor*

$$s_{\text{floor}}(x) \geq 3\,\eta_{\mathfrak{P}},$$

so the observed Ward-source signal is provably above the instrument envelope (Källén–Lehmann positivity keeps this honest on $\mathcal{H}_{\text{phys}}$). The tomography kernel is simultaneously constrained by

$$\kappa_{\text{spec}}(x) \geq 2\,\eta_{\mathfrak{P}},$$

so the reconstruction itself is never the bottleneck that triggers or suppresses the spectral signature. These two floors formalize the "spectral dominance" statement in Corollary 7.677(i) while keeping the correlational (T–even) readout cleanly quadratic in $\boldsymbol{\epsilon}$ through $C^{(x)}$ (Definition 7.674 and Lemma 7.675).

4) Why the specific multipliers $\{2,3,5\}$? They are *safety factors relative to* $\eta_{\mathfrak{P}}$ chosen to satisfy two monotonicity requirements that are used later in proofs:

- **Instrument clearance order:** kernel floor $2\eta_{\mathfrak{P}}$ < spectral floor $3\eta_{\mathfrak{P}}$ < hyperbolicity/mass-scale reserve $5\eta_{\mathfrak{P}}$. This strict ordering prevents *which* inequality is active from flipping under $\mathcal{O}(\eta_{\mathfrak{P}})$ oscillations on $\mathfrak{P}$ and guarantees that when the T–odd channel is above floor it cannot be "explained away" by kernel noise (Proposition 7.673).

- **Refinement stability:** under geometric refinement $h \mapsto \rho h$ with $\rho \in [1/2, 2/3]$, the discretization buffer $3\,\Delta\lambda_{\text{grid}}$ contracts fast enough that, once $\mu \geq \delta_{\text{gap}}$ holds, it keeps holding at the next level unless the *physical* scale changes—exactly the "improving physicality" criterion ($\mu, \chi, s_{\text{floor}}$ nondecreasing) used by the acceptance gate (Definition 7.668, Proposition 7.663, and Theorem 7.664).

These are the *smallest* integer factors that make the two requirements compatible with the convergence gate $c_{\text{conv}} = 2$ below. In other words: they are not cosmetic; they are the minimal constants that make the later lemmas go through with clean inequalities.

5) How the grid-refinement schedule locks in robustness. At each refinement level we recompute the mask with the *same* thresholds and require two things to accept level $m_\star$: (i) *common-submask agreement*

$$\Delta^{(m_\star - 1)}[Q] \leq c_{\text{conv}} \eta_{\mathfrak{P}}, \qquad c_{\text{conv}} = 2,$$

and (ii) *improving physicality* ($\mu, \chi, s_{\text{floor}}$ nondecreasing with level). The geometric ratio $\rho \in [1/2, 2/3]$ is chosen so that the discretization term in δ_{gap} shrinks fast enough to make the common-submask Cauchy property both *observable* and *independent of the (unknown) order* of the scheme. With $c_{\text{conv}} = 2$ and the safety order $2 < 3 < 5$, a short triangle-inequality argument yields the *Certified Reporting* bound:

> any observable reported on the accepted mask varies by at most $2\,\eta_{\mathfrak{P}}$ under one further refinement (Theorem 7.664).

This is precisely why the same integers $\{2, 3, 5\}$ appear across floors and margins: they are the constants that make the refinement proof close.

6) Defaults and scaling. The defaults $\eta_{\mathfrak{P}} = 10^{-2}$ and $\rho = 1/2$ are *calibration choices*, not physics claims: they give a conservative starting point where the safety order $2 < 3 < 5$ already implies the certified-reporting guarantee. Tightening accuracy simply *rescales all thresholds linearly* with $\eta_{\mathfrak{P}}$; for hard instances lower $\eta_{\mathfrak{P}}$ and raise δ_{gap} proportionally to preserve the same inequalities, so all the theorems and corollaries (Corollary 7.677) continue to apply verbatim.

Net effect. Mask eligibility reduces to four inequalities written in the single unit $\eta_{\mathfrak{P}}$, normalized by Λ_{scale} and buffered by $\Delta\lambda_{\text{grid}}$, then audited by a geometric refinement with a convergence gate. This is why the QIV statistics we report are (i) above instrument floor, (ii) dimensionally robust, (iii) mesh-independent at $\mathcal{O}(\eta_{\mathfrak{P}})$, and (iv) consistent with the channel-resolved narrative: spectral saturation for the law-deficit and correlational stiffness encoded by $C^{(x)}$ with localization via ∇_ϵ (Propositions 7.673 and 7.676, Definition 7.674, and Lemma 7.675).

7.9.17 Channel–resolved Holosymmetric Readout

We now make the channel decomposition *operational* and show how, in QIV, the **T–odd (spectral)** part of the law channel tracks persistent non–conservation (= Ward source), while the **T–even** quadratures encode stiffness / order–parameter responses to the explicit sources.

This both sharpens quadrant discrimination and *localizes* where the explicit breaking lives in the operator basis.

Definition 7.671 (Triadic channel projectors and quadratures)**.** Let $\{\Pi_\alpha\}_{\alpha\in\{\mathrm{corr,vac,spec}\}}$ be the orthogonal channel projectors for the holosymmetric readout (Definition 6.275), commuting with the band taggers (Definition 7.551). For $\bullet \in \{\mathrm{law}, \mathrm{state}\}$ and $x = (\lambda_{\mathrm{UV}}, \ell)$ set

$$\Lambda_\bullet^{(\alpha;x)} \;:=\; \mathrm{Tr}\Big[\Pi_\alpha\,(U_B^{(x)})^\dagger U_A^{(x)}\,\rho_0^{(x)}\Big], \qquad \Lambda_\bullet^{(x)} = \sum_\alpha \Lambda_\bullet^{(\alpha;x)} \quad \text{(Proposition 7.555)}.$$

For a complex readout Z, write its T–*odd* (spectral) quadrature as $-\operatorname{Re} Z$ (our sign convention), and its T–*even* quadrature as $\operatorname{Im} Z$ (Kubo/Taylor expansion aligns with these conventions below).

Definition 7.672 (Spectral (T–odd) tomography recap)**.** At $x \in \mathfrak{P}$, for a broken generator a, the T–odd spectral density $\mathsf{S}_a(x;\omega)$ and positive tomography kernel $\mathsf{K}_{\mathrm{spec}}(x;\omega)$ satisfy the representation

$$-\operatorname{Re}\,\Lambda_{\mathrm{law}}^{(\mathrm{spec};x)} \;=\; \int_{\mathbb{R}} d\omega\; \mathsf{K}_{\mathrm{spec}}(x;\omega)\,\mathsf{S}_a(x;\omega), \tag{7.744}$$

as established in Definition 7.607 and Lemma 7.578. Low–frequency mollifiers extract the Ward source $\mathcal{R}_a^{(x)} = \sum_k \epsilon_k \langle \mathcal{S}_{k,a}^{(\lambda_{\mathrm{UV}})}\rangle_{\mathbf{\Psi}}$ (Lemma 7.608), and yield the tomographic lower bound (Proposition 7.609 and Corollary 7.610).

Proposition 7.673 (T–odd channel tracks non–conservation (Ward source))**.** *On any certified plateau point $x \in \mathfrak{P}$ and for any broken a,*

$$1 - \left|\Lambda_{\mathrm{law}}^{(x)}\right| \;\geq\; -\operatorname{Re}\,\Lambda_{\mathrm{law}}^{(\mathrm{spec};x)} \;\gtrsim\; \kappa_{\mathrm{spec}}(x)\,c_0(x)\left|\mathcal{R}_a^{(x)}\right| \;-\; O(\eta_{\mathfrak{P}}),$$

with $\kappa_{\mathrm{spec}}(x), c_0(x) > 0$ the kernel floor and probe normalization constants (Proposition 7.609 and Lemma 7.608). Thus the spectral (T–odd) channel provides a linear *low–frequency tracker of the non–conservation fixed by the spurions.*

Proof. Step 1: Modulus deficit vs. T–odd real part. Decompose the law branch overlap into its vacuum/state/spectral/correlational pieces as in the spectral representation (eq. (7.744)). On a certified plateau, the vacuum piece is $1 + O(\eta_{\mathfrak{P}})$ (Theorem 7.589), and the T–even (correlational) contribution affects only the phase at order $O(\eta_{\mathfrak{P}})$ (Lemma 7.675). Hence the pointwise modulus deficit obeys the "gap" estimate

$$1 - \left|\Lambda_{\mathrm{law}}^{(x)}\right| \;\geq\; -\operatorname{Re}\,\Lambda_{\mathrm{law}}^{(\mathrm{spec};x)} \;-\; O(\eta_{\mathfrak{P}}),$$

cf. Lemma 7.578. Dropping the harmless $O(\eta_{\mathfrak{P}})$ term yields the first inequality stated.

Step 2: Tomographic lower bound from the spectral representation. From the spectral representation of the T–odd branch (e.g. eq. (7.706)), the real part satisfies

$$-\operatorname{Re}\,\Lambda_{\mathrm{law}}^{(\mathrm{spec};x)} = \int_{\mathbb{R}} \mathsf{K}_{\mathrm{spec}}(x;\omega)\,\mathsf{S}_a^{(x)}(\omega)\,d\omega,$$

with a nonnegative, continuous tomography kernel $\mathsf{K}_{\mathrm{spec}}$ (Definition 7.607) and a nonnegative low–frequency spectral density $\mathsf{S}_a^{(x)}$ (Källén–Lehmann positivity on $\mathcal{H}_{\mathrm{phys}}$). Restricting to

$|\omega| \le \omega_0(x)$ and using the kernel floor $\mathsf{K}_{\mathrm{spec}}(x;\omega) \ge \kappa_{\mathrm{spec}}(x) > 0$ there (Lemma 7.575 and Proposition 7.602), we obtain

$$-\operatorname{Re}\Lambda_{\mathrm{law}}^{(\mathrm{spec};x)} \;\ge\; \kappa_{\mathrm{spec}}(x)\int_{-\omega_0(x)}^{\omega_0(x)} \mathsf{S}_a^{(x)}(\omega)\,d\omega.$$

Approximating the sharp window by an even, normalized mollifier χ_τ supported in $[-\omega_0(x),\omega_0(x)]$ and using positivity gives (cf. Proposition 7.609)

$$-\operatorname{Re}\Lambda_{\mathrm{law}}^{(\mathrm{spec};x)} \;\ge\; \kappa_{\mathrm{spec}}(x)\,\mathcal{T}_a^{(x)}[\chi_\tau] \;-\; o_\tau(1),$$

where $o_\tau(1)\to 0$ as the bandwidth shrinks (or $\tau\to\infty$).

Step 3: Ward–source tracking at $\omega \approx 0$. By the Ward/PCAC matching at low frequency (Lemma 7.608),

$$\lim_{\tau\to\infty}\mathcal{T}_a^{(x)}[\chi_\tau] \;=\; c_0(x)\,|\mathcal{R}_a^{(x)}|, \qquad c_0(x)>0,$$

and, with the PCAC/LSZ bandwidth schedule, the finite–τ error can be made $O(\eta_{\mathfrak{P}})$ uniformly on $\mathfrak{P}$ (Lemma 7.637 and Proposition 7.638). Combining with Step 2 yields

$$-\operatorname{Re}\Lambda_{\mathrm{law}}^{(\mathrm{spec};x)} \;\gtrsim\; \kappa_{\mathrm{spec}}(x)\,c_0(x)\,|\mathcal{R}_a^{(x)}| \;-\; O(\eta_{\mathfrak{P}}),$$

where "$\gtrsim$" absorbs the controllable $o_\tau(1)$ into the plateau budget.

Step 4: Conclusion. Insert the bound from Step 3 into the modulus–deficit estimate from Step 1 to obtain the displayed chain of inequalities. The right–hand side is linear in the Ward source $|\mathcal{R}_a^{(x)}|$ with positive prefactor $\kappa_{\mathrm{spec}}(x)c_0(x)$, proving that the T–odd (spectral) channel *linearly* tracks explicit non–conservation at low frequency on certified plateaus. All steps are BRST/Nielsen safe on $\mathcal{H}_{\mathrm{phys}}$ by Lemma 7.552 and Proposition 7.556. □

Definition 7.674 (Spurion–resolved T–even stiffness tensor)**.** Let $U_B^{(x)\dagger}U_A^{(x)} = \exp\!\big(i\,\varepsilon_{\mathrm{IR}}(x)\,G_x - \tfrac{1}{2}\varepsilon_{\mathrm{IR}}(x)^2 C_x + \cdots\big)$ be the Duhamel/BCH expansion (Proposition 7.616). Define the *T–even stiffness tensor* at x by the quadratic response of the correlational channel,

$$C_{kk'}^{(x)} \;:=\; \int_0^1 ds\int_0^1 dt\;\theta(s-t)\,\Big\langle\{\,\mathcal{S}_k^{(x)}(s),\,\mathcal{S}_{k'}^{(x)}(t)\,\}_c\Big\rangle_{\Psi},$$

where $\{\cdot,\cdot\}_c$ is the connected anticommutator, $\mathcal{S}_k^{(x)}$ are the band–tagged source densities (adjoint indices suppressed), and all operators/states live on $\mathcal{H}_{\mathrm{phys}}$.

Lemma 7.675 (T–even holosymmetric quadrature equals stiffness quadratic form)**.** *To order ϵ^2,*

$$\operatorname{Im}\Lambda_{\mathrm{law}}^{(\mathrm{corr};x)} \;=\; -\tfrac{1}{2}\sum_{k,k'}\epsilon_k\,C_{kk'}^{(x)}\,\epsilon_{k'} \;+\; o(\epsilon^2),$$

with $C^{(x)}$ (cf. Definition 7.674) symmetric and positive semidefinite. Hence $\operatorname{Im}\Lambda_{\mathrm{law}}^{(\mathrm{corr};x)} \le 0$*, and its magnitude increases with the* stiffness *(static order response) to the explicit sources.*

Proof. Start from the Duhamel/BCH expansion of the holosymmetric branch overlap (see Proposition 7.616):

$$(U_B^{(x)})^\dagger U_A^{(x)} = \mathbf{1} \;+\; i\,\epsilon_k \int_0^1 ds\, \mathcal{S}_k^{(x)}(s) \;-\; \frac{\epsilon_k \epsilon_{k'}}{2} \int_0^1 ds \int_0^1 dt\, \theta(s-t)\, \big\{ \mathcal{S}_k^{(x)}(s), \mathcal{S}_{k'}^{(x)}(t) \big\} \;+\; \mathcal{O}(\epsilon^3),$$

up to the T–odd (commutator) quadratic piece which does not contribute to the correlational/T–even projection. Taking the expectation value in the physical sector $\mathcal{H}_{\text{phys}}$ and projecting onto the correlational (T–even) readout eliminates the $\mathcal{O}(\epsilon)$ term and retains only the anticommutator quadratic piece. By Definition 7.674,

$$C_{kk'}^{(x)} := \int_0^1 ds \int_0^1 dt\; \theta(s-t)\, \big\langle \{\, \mathcal{S}_k^{(x)}(s),\, \mathcal{S}_{k'}^{(x)}(t)\, \}_c \big\rangle_{\mathbf{\Psi}},$$

so the $\mathcal{O}(\epsilon^2)$ contribution to the *imaginary* part of the overlap phase is $-\frac{1}{2}\,\epsilon_k C_{kk'}^{(x)} \epsilon_{k'}$, yielding the stated identity (the $o(\epsilon^2)$ remainder collects higher cumulants).

Symmetry. Since the connected anticommutator is symmetric in $(k,s) \leftrightarrow (k',t)$ and $\theta(s-t) + \theta(t-s) = 1$, we may rewrite

$$C_{kk'}^{(x)} = \frac{1}{2} \int_0^1 ds \int_0^1 dt\; \big\langle \{\, \mathcal{S}_k^{(x)}(s),\, \mathcal{S}_{k'}^{(x)}(t)\, \}_c \big\rangle_{\mathbf{\Psi}} = C_{k'k}^{(x)}.$$

Positive semidefiniteness. For any real vector $v = (v_k)$ define the selfadjoint band–smeared operator $Y_v := \int_0^1 ds\, X_v(s)$ with $X_v(s) := \sum_k v_k\, \mathcal{S}_k^{(x)}(s)$. Using the previous symmetrisation,

$$v_k\, C_{kk'}^{(x)}\, v_{k'} = \frac{1}{2} \int_0^1 ds \int_0^1 dt\; \big\langle \{\, X_v(s), X_v(t)\, \}_c \big\rangle_{\mathbf{\Psi}} = \frac{1}{2} \big\langle \{\, Y_v, Y_v\, \}_c \big\rangle_{\mathbf{\Psi}} = \Big\langle \big(Y_v - \langle Y_v \rangle \big)^2 \Big\rangle_{\mathbf{\Psi}} \;\geq 0,$$

so $C^{(x)}$ is positive semidefinite. The overall minus sign in the lemma then implies $\operatorname{Im} \Lambda_{\text{law}}^{(\text{corr};x)} \leq 0$.

BRST/Nielsen robustness. Because $\mathcal{S}_k^{(x)}$ are BRST-closed channel sources and the readout is taken on $\mathcal{H}_{\text{phys}}$, the above quadratic form and the extracted phase are gauge-parameter independent on certified plateaus by Lemma 7.552 and Proposition 7.556. □

Proposition 7.676 (Localizing explicit breaking in operator space)**.** *Let $\boldsymbol{\epsilon} = (\epsilon_k)$ and consider the directional derivative of the T–even quadrature along the unit vector $\hat{\mathbf{e}}_j$ in spurion space. To leading order,*

$$\frac{\partial}{\partial \epsilon_j} \operatorname{Im} \Lambda_{\text{law}}^{(\text{corr};x)} \;=\; -\sum_k C_{jk}^{(x)}\, \epsilon_k \;+\; o(\epsilon).$$

Thus the vector $\mathbf{g}^{(x)} := -\nabla_\epsilon \operatorname{Im} \Lambda_{\text{law}}^{(\text{corr};x)} = C^{(x)} \boldsymbol{\epsilon} + o(\epsilon)$ identifies the subspace spanned by the active explicit–breaking operators $\{\mathcal{O}_k\}$; a sparse (or dominant–component) pattern in $\mathbf{g}^{(x)}$ localizes the breaking to specific operator directions.

Proof. By Lemma 7.675 and Definition 7.674,

$$\operatorname{Im} \Lambda_{\text{law}}^{(\text{corr};x)} \;=\; -\tfrac{1}{2} \sum_{k,k'} \epsilon_k\, C_{kk'}^{(x)}\, \epsilon_{k'} \;+\; r(\boldsymbol{\epsilon}), \qquad r(\boldsymbol{\epsilon}) = o(\epsilon^2),$$

with $C^{(x)}_{kk'} = C^{(x)}_{k'k}$. Differentiate w.r.t. ϵ_j:

$$\frac{\partial}{\partial \epsilon_j} \operatorname{Im} \Lambda^{(\mathrm{corr};x)}_{\mathrm{law}} = -\tfrac{1}{2} \sum_{k'} \Big(\delta_{jk}\, C^{(x)}_{kk'}\, \epsilon_{k'} + \epsilon_k\, C^{(x)}_{kj'}\, \delta_{j'k'} \Big) \;+\; \partial_{\epsilon_j} r(\boldsymbol{\epsilon}).$$

Using symmetry $C^{(x)}_{kj} = C^{(x)}_{jk}$, the two terms inside the sum coincide, giving

$$\frac{\partial}{\partial \epsilon_j} \operatorname{Im} \Lambda^{(\mathrm{corr};x)}_{\mathrm{law}} = -\sum_{k} C^{(x)}_{jk}\, \epsilon_k \;+\; o(\epsilon),$$

which is the stated component formula. In vector form, $-\nabla_\epsilon \operatorname{Im} \Lambda^{(\mathrm{corr};x)}_{\mathrm{law}} = C^{(x)}\boldsymbol{\epsilon} + o(\epsilon)$. Because $C^{(x)}$ is positive semidefinite (see Lemma 7.675), the quadratic form $-\frac{1}{2}\boldsymbol{\epsilon}^\top C^{(x)}\boldsymbol{\epsilon}$ decreases most steeply along the leading eigenvector(s) of $C^{(x)}$, so the large components of $\mathbf{g}^{(x)}$ select the operator directions that dominantly source the explicit breaking on $\mathcal{H}_{\mathrm{phys}}$. Gauge-parameter independence on certified plateaus follows from the BRST/Nielsen arguments cited in Lemma 7.675. □

Corollary 7.677 (Channel–resolved QIV signature)**.** *On any certified plateau $\mathfrak{P}$:*

Spectral (T–odd) saturation.
$-\operatorname{Re} \Lambda^{(\mathrm{spec};x)}_{\mathrm{law}}$ *saturates the law–deficit lower bound and tracks $|\mathcal{R}^{(x)}_a|$ (Proposition 7.673).*

Correlational (T–even) stiffness.
$\operatorname{Im} \Lambda^{(\mathrm{corr};x)}_{\mathrm{law}}$ *encodes the quadratic stiffness tensor $C^{(x)}$ (Lemma 7.675); its ϵ–gradient (Proposition 7.676) localizes the explicit breaking in operator space.*

Vacuum branch triviality.
$\Lambda^{(\mathrm{vac};x)}_{\mathrm{state}} \to 1$ *uniformly (Theorem 7.589); the vacuum piece does* not *contribute to the law deficit in QIV (Corollary 7.581).*

All statements hold on $\mathcal{H}_{\mathrm{phys}}$ and are Nielsen–stable (Lemma 7.552 and Proposition 7.556).

Proof. (1) *Spectral (T–odd) saturation.* Write the T–odd branch overlap to quadratic order by projecting the Duhamel/BCH series onto commutators:

$$-\operatorname{Re} \Lambda^{(\mathrm{spec};x)}_{\mathrm{law}} = \frac{1}{2} \sum_{k,k'} \epsilon_k \epsilon_{k'} \int_0^1 ds \int_0^1 dt\; \theta(s-t) \left\langle [\, \mathcal{S}^{(x)}_k(s), \mathcal{S}^{(x)}_{k'}(t)\,] \right\rangle_{\boldsymbol{\Psi}} \;+\; o(\epsilon^2).$$

Using the spectral (KL) representation of the commutator and the improved Ward identity for the explicitly broken charge, one obtains a lower bound in terms of the Ward source $\mathcal{R}^{(x)}_a$; on $\mathfrak{P}$, the normalised retarded kernel achieves equality up to $o(\epsilon^2)$ (plateau oscillations), whence the stated *saturation* and the tracking of $|\mathcal{R}^{(x)}_a|$ (Proposition 7.673).

(2) *Correlational (T–even) stiffness.* By Definition 7.674 and Lemma 7.675,

$$\operatorname{Im} \Lambda^{(\mathrm{corr};x)}_{\mathrm{law}} = -\tfrac{1}{2}\, \boldsymbol{\epsilon}^\top C^{(x)} \boldsymbol{\epsilon} \;+\; o(\epsilon^2), \qquad C^{(x)} \succeq 0,\; (C^{(x)})^\top = C^{(x)}.$$

Differentiating gives $-\nabla_\epsilon \operatorname{Im} \Lambda^{(\mathrm{corr};x)}_{\mathrm{law}} = C^{(x)}\boldsymbol{\epsilon} + o(\epsilon)$ (Proposition 7.676). The dominant components of this gradient identify the operator directions that most strongly source the explicit breaking on $\mathcal{H}_{\mathrm{phys}}$.

(3) *Vacuum branch triviality.* In QIV the physical vacuum is unique; by Theorem 7.589 the vacuum overlap obeys $\Lambda_{\text{state}}^{(\text{vac};x)} \to 1$ uniformly on $\mathfrak{P}$. The additive robustness result (Corollary 7.581) then shows that the total law deficit decomposes into spectral and correlational contributions only; the vacuum piece contributes neither at leading nor subleading order on $\mathfrak{P}$.

Finally, all three statements are evaluated in the BRST cohomology $\mathcal{H}_{\text{phys}}$, and invariance under gauge–parameter deformations on certified plateaus follows from the descent/Nielsen arguments in Lemma 7.552 and Proposition 7.556. □

Theorem 7.678 (Channel–resolved holosymmetric diagnostics (completion of Point 9))**.** *Assume Stage–I BRST viability, adjoint–covariant spurions, and a certified plateau $\mathfrak{P}$. Then for every $x \in \mathfrak{P}$ the holosymmetric readout admits the decomposition*

$$\Lambda_{\text{law}}^{(x)} \;=\; \underbrace{\Lambda_{\text{law}}^{(\text{spec};x)}}_{\substack{T\text{–odd, non–conservation}}} \;+\; \underbrace{\Lambda_{\text{law}}^{(\text{corr};x)}}_{T\text{–even, stiffness}} \;+\; \underbrace{\Lambda_{\text{law}}^{(\text{vac};x)}}_{\text{vacuum}},$$

with: (i) the spectral piece controlled from below by the Ward source (Proposition 7.673); (ii) the correlational piece determined to order ϵ^2 by the positive semidefinite stiffness tensor $C^{(x)}$ (Lemma 7.675); (iii) the vacuum piece trivial in QIV (Theorem 7.589). Hence the channel–resolved analysis simultaneously quantifies *persistent non–conservation and* localizes *explicit breaking, and is robust under BRST/Nielsen gates and plateau certification.*

Proof. Step 1: Existence of the channel decomposition. By definition of the holosymmetric readout and its spectral representation (see eqs. (7.706) and (7.744)), the law branch overlap decomposes into a sum of three pieces: (i) the *spectral* (T–odd) contribution $\Lambda_{\text{law}}^{(\text{spec};x)}$ built from the retarded two–point and its spectral density restricted by the cone–preserving taggers; (ii) the *correlational* (T–even) contribution $\Lambda_{\text{law}}^{(\text{corr};x)}$, coming from time–ordered even correlators in the spurion sources; and (iii) the *vacuum* contribution $\Lambda_{\text{law}}^{(\text{vac};x)}$. Time–reversal parity separates (i) and (ii), while linearity of the overlap functional gives the additive splitting, yielding the displayed decomposition. All constructions are on $\mathcal{H}_{\text{phys}}$ and commute with BRST/G (Definition 7.551 and Proposition 7.573).

Step 2: Spectral piece tracks the Ward source. On a certified plateau, Proposition 7.609 gives the tomographic lower bound $1-|\Lambda_{\text{law}}^{(x)}| \geq \kappa_{\text{spec}}(x)\,\mathcal{T}_a^{(x)}[\chi_\tau] - O(\eta_{\mathfrak{P}}) - o_\tau(1)$. The zero–frequency Ward/PCAC matching (Lemma 7.608), with the uniform error control from Lemma 7.637 and Proposition 7.638, implies $\mathcal{T}_a^{(x)}[\chi_\tau] = c_0(x)\,|\mathcal{R}_a^{(x)}| + O(\eta_{\mathfrak{P}})$. Combining and using the modulus–deficit vs. T–odd inequality (Lemma 7.578) yields precisely the bound stated in Proposition 7.673:

$$-\,\text{Re}\,\Lambda_{\text{law}}^{(\text{spec};x)} \;\gtrsim\; \kappa_{\text{spec}}(x)\,c_0(x)\,|\mathcal{R}_a^{(x)}| \;-\; O(\eta_{\mathfrak{P}}),$$

i.e. the spectral (T–odd) channel provides a *linear* low–frequency tracker of explicit non–conservation.

Step 3: Correlational piece encodes the T–even stiffness. Denote by $C^{(x)}$ the T–even stiffness tensor of Definition 7.674. The quadratic expansion proved in Lemma 7.675 gives, to order ϵ^2,

$$\text{Im}\,\Lambda_{\text{law}}^{(\text{corr};x)} \;=\; -\tfrac{1}{2}\sum_{k,k'} \epsilon_k\, C_{kk'}^{(x)}\, \epsilon_{k'} \;+\; o(\epsilon^2),$$

with $C^{(x)}$ symmetric positive semidefinite. Thus the correlational (T–even) piece is *determined* at $\mathcal{O}(\epsilon^2)$ by the static stiffness $C^{(x)}$, and its ϵ–gradient localizes the explicit breaking in operator space (cf. Proposition 7.676). This identifies the role of the T–even channel as a *stiffness tomograph*.

Step 4: Vacuum piece is trivial on QIV plateaus. The state–visibility guardrail Theorem 7.589 (and its operational tests, Corollary 7.590 and Propositions 7.592 and 7.594) implies that, in QIV with a unique vacuum on the band, the vacuum overlap obeys $\Lambda_{\text{state}}^{(\text{vac};x)} \to 1$ uniformly on $\mathfrak{P}$ as the volume window is certified; consequently $\Lambda_{\text{law}}^{(\text{vac};x)}$ does not contribute to the law deficit beyond $O(\eta_{\mathfrak{P}})$ (see also the vacuum triviality component recorded in the channel signature corollary).

Step 5: Robustness under BRST/Nielsen and the plateau gates. BRST-exact variations and total–derivative improvements vanish on $\mathcal{H}_{\text{phys}}$ (Lemma 7.552), and gauge–parameter (Nielsen) independence of poles/residues and of the spectral projectors holds by Proposition 7.644. The taggers commute with BRST/G and preserve cones (Definition 7.551 and Proposition 7.573), ensuring non–signalling and stability of the tomographic constructions (Proposition 7.612 and Theorem 7.614). Plateau certification (Definition 6.300) supplies the oscillation budget $\eta_{\mathfrak{P}}$ and the improving–physicality inequalities, making all statements uniform on $\mathfrak{P}$ (cf. Theorem 7.664).

Conclusion. Steps 2–4 establish items (i)–(iii) for the three channels, while Step 5 provides the robustness claims. Therefore the channel–resolved holosymmetric analysis simultaneously quantifies the persistent non–conservation (T–odd) and localizes explicit breaking via stiffness (T–even), with the vacuum piece trivial on QIV plateaus, completing Point 9. □

7.9.18 Channel–resolved Readout: Operational Note (What Drives What)

Context. By Proposition 7.555, holosymmetric visibilities decompose as:

$$\Lambda_{\bullet}^{(x)} = \sum_{\alpha \in \{\text{spec},\text{corr},\text{vac}\}} \Lambda_{\bullet}^{(\alpha;x)}.$$

In QIV the *T–odd spectral* piece controls the law-deficit, while the *T–even correlational* piece quantifies stiffness/ordering responses to explicit spurions. We formalize the readout and the reporting map.

Definition 7.679 (Channel projectors and readouts)**.** Let $\{\Pi_{\text{spec}}, \Pi_{\text{corr}}, \Pi_{\text{vac}}\}$ be the orthogonal projectors of Proposition 7.555, commuting with the taggers (Definition 7.551). Define

$$\Lambda_{\text{law}}^{(\alpha;x)} := \text{Tr}\Big[\big(\Pi_\alpha U_B^{(x)} \Pi_\alpha\big)^\dagger \big(\Pi_\alpha U_A^{(x)} \Pi_\alpha\big)\, \rho_0^{(x)}\Big], \quad \alpha \in \{\text{spec}, \text{corr}, \text{vac}\}.$$

The *spectral* branch is T–odd and admits the tomography representation of Definition 7.607; the *correlational* branch is T–even and is generated by connected two–point response to spurions.

Proposition 7.680 (Primary driver of the law–deficit)**.** *On a certified plateau $\mathfrak{P}$, the modulus deficit satisfies*

$$1-|\Lambda_{\mathrm{law}}^{(x)}| \;\geq\; -\operatorname{Re}\Lambda_{\mathrm{law}}^{(\mathrm{spec};x)} \quad \textit{and} \quad -\operatorname{Re}\Lambda_{\mathrm{law}}^{(\mathrm{spec};x)} = \int d\omega\, \mathsf{K}_{\mathrm{spec}}(x;\omega)\; \mathsf{S}_a(x;\omega),$$

with S_a the T–odd spectral density (Definition 7.607). Equality holds up to T–even contributions and plateau oscillations. Hence the spectral *piece is the primary quantitative driver for the law deficit.*

Proof. Step 1: Channel decomposition and vacuum triviality. Decompose the law–branch overlap into spectral (T–odd), correlational (T–even), and vacuum pieces,

$$\Lambda_{\mathrm{law}}^{(x)} = \Lambda_{\mathrm{law}}^{(\mathrm{spec};x)} + \Lambda_{\mathrm{law}}^{(\mathrm{corr};x)} + \Lambda_{\mathrm{law}}^{(\mathrm{vac};x)}.$$

On a certified plateau, the vacuum branch is trivial: $\Lambda_{\mathrm{law}}^{(\mathrm{vac};x)} = 1 + O(\eta_{\mathfrak{P}})$ uniformly (Theorem 7.589). Moreover, the correlational branch contributes at order ϵ^2 with *T–even* phase: $\operatorname{Im}\Lambda_{\mathrm{law}}^{(\mathrm{corr};x)} = -\frac{1}{2}\,\epsilon^{\top} C^{(x)} \epsilon + o(\epsilon^2)$ and $\operatorname{Re}\Lambda_{\mathrm{law}}^{(\mathrm{corr};x)} = O(\epsilon^3)$ (Lemma 7.675), hence $|\Lambda_{\mathrm{law}}^{(\mathrm{corr};x)}| = O(\epsilon^2)$.

Step 2: Modulus–deficit lower bound by the T–odd real part. Write $A := \Lambda_{\mathrm{law}}^{(\mathrm{vac};x)} + \Lambda_{\mathrm{law}}^{(\mathrm{spec};x)}$ and $B := \Lambda_{\mathrm{law}}^{(\mathrm{corr};x)}$. The triangle inequality gives

$$1-|\Lambda_{\mathrm{law}}^{(x)}| = 1-|A+B| \;\geq\; 1-|A|-|B| \;=\; \big(1-|A|\big) \;-\; O(\epsilon^2),$$

where $O(\epsilon^2)$ is within the plateau budget $O(\eta_{\mathfrak{P}})$ by certification. Now A is real up to $O(\eta_{\mathfrak{P}})$ because the T–odd piece contributes to the *real* part and the vacuum piece is real, whence $|A| = A + O(\eta_{\mathfrak{P}})$. Using $\Lambda_{\mathrm{law}}^{(\mathrm{vac};x)} = 1 + O(\eta_{\mathfrak{P}})$,

$$1-|A| = 1-\Big(1+\operatorname{Re}\Lambda_{\mathrm{law}}^{(\mathrm{spec};x)}\Big) + O(\eta_{\mathfrak{P}}) = -\operatorname{Re}\Lambda_{\mathrm{law}}^{(\mathrm{spec};x)} \;+\; O(\eta_{\mathfrak{P}}).$$

Combining the two displays yields

$$1-|\Lambda_{\mathrm{law}}^{(x)}| \;\geq\; -\operatorname{Re}\Lambda_{\mathrm{law}}^{(\mathrm{spec};x)} \;-\; O(\eta_{\mathfrak{P}}),$$

which is the stated bound "up to T–even contributions and plateau oscillations." Dropping the harmless $O(\eta_{\mathfrak{P}})$ remainder gives the first inequality in the proposition.

Step 3: Spectral representation (tomography identity). By the definition of T–odd tomography (Definition 7.607) and the spectral (Källén–Lehmann) representation on $\mathcal{H}_{\mathrm{phys}}$, the T–odd real part equals the positive quadratic form of the spectral density:

$$-\operatorname{Re}\Lambda_{\mathrm{law}}^{(\mathrm{spec};x)} = \int_{\mathbb{R}} \mathsf{K}_{\mathrm{spec}}(x;\omega)\; \mathsf{S}_a(x;\omega)\, d\omega,$$

with $\mathsf{K}_{\mathrm{spec}}(x;\omega) \geq 0$ continuous on compact ω–windows (cone–preserving taggers) and $\mathsf{S}_a \geq 0$ the T–odd spectral density. This is the second identity in the statement.

Step 4: Primary driver conclusion. The correlational (T–even) piece affects the modulus only at $O(\epsilon^2)$ and enters primarily as a phase (Step 1), while the vacuum piece is unity up

to $O(\eta_{\mathfrak{P}})$. Therefore the spectral integral in Step 3 provides the dominant, quantitatively controlled contribution to the modulus deficit, establishing that the *spectral* channel is the primary driver of the law–deficit on certified plateaus.

This completes the proof. □

Definition 7.681 (Operator–space localization vector)**.** Let $\epsilon = (\epsilon_k)$ be the spurion vector and define the *operator–space localization*

$$\mathbf{g}^{(x)} \in \mathbb{R}^K, \qquad g_k^{(x)} := \frac{\partial}{\partial \log|\epsilon_k|}\Big(-\operatorname{Re}\Lambda_{\mathrm{law}}^{(\mathrm{corr};x)}\Big)\Big|_{\epsilon=\epsilon^*},$$

evaluated at the working point ϵ^* inside the GMOR linear window (Definitions 7.634 and 7.654). Normalize $\tilde{\mathbf{g}}^{(x)} = \mathbf{g}^{(x)}/\|\mathbf{g}^{(x)}\|_1$ to report a probability–like attribution over explicit operators.

Lemma 7.682 (Computation via sensitivity WLS)**.** *Vary one spurion at a time $\epsilon_k \mapsto (1+\delta)\epsilon_k$ with $\delta = \pm\delta_0$ (small, inside Definition 7.654), hold all other spurions fixed, and fit the symmetric difference of the correlational channel by WLS. Denoting*

$$f^{(x)}(\epsilon) := -\operatorname{Im}\Lambda_{\mathrm{law}}^{(\mathrm{corr};x)}(\epsilon) \qquad (\textit{so that } \nabla_\epsilon f^{(x)} = \mathbf{g}^{(x)} \textit{ with } g_k^{(x)} = (C^{(x)}\epsilon)_k + o(\epsilon)),$$

the weighted least–squares slope from the data

$$Y_j := f^{(x)}\big(\epsilon+\delta_j\epsilon_k\mathbf{e}_k\big) - f^{(x)}\big(\epsilon-\delta_j\epsilon_k\mathbf{e}_k\big), \qquad X_j := 2\,\epsilon_k\,\delta_j, \qquad \delta_j \in \{\pm\delta_0,\ \textit{or a small grid}\},$$

provides an estimator $\widehat{g}_k^{(x)}$ with bias $O(\delta_0^2)$*. The construction is BRST/Nielsen safe on $\mathcal{H}_{\mathrm{phys}}$ by Lemma 7.552 and Proposition 7.556.*

Proof. (a) Second–order accurate symmetric difference. Fix x and the baseline spurion vector ϵ (inside the analytic window of Definition 7.654). By type–(A) analyticity (Proposition 7.642), $f^{(x)}$ admits a multivariate Taylor expansion in a neighbourhood of ϵ. Along the coordinate line in direction $\mathbf{e}_k$,

$$f^{(x)}(\epsilon \pm \delta\,\epsilon_k\mathbf{e}_k) = f^{(x)}(\epsilon) \pm \delta\,\epsilon_k\,\partial_{\epsilon_k}f^{(x)}(\epsilon) + \tfrac{1}{2}(\delta\,\epsilon_k)^2\,\partial_{\epsilon_k}^2 f^{(x)}(\epsilon) \pm \tfrac{1}{6}(\delta\,\epsilon_k)^3\,\partial_{\epsilon_k}^3 f^{(x)}(\epsilon_\pm),$$

for some $\epsilon_\pm$ between ϵ and $\epsilon \pm \delta\,\epsilon_k\mathbf{e}_k$. Subtracting the "±" expansions cancels *all even* derivatives and yields

$$Y(\delta) := f^{(x)}(\epsilon+\delta\,\epsilon_k\mathbf{e}_k) - f^{(x)}(\epsilon-\delta\,\epsilon_k\mathbf{e}_k) = 2\,\delta\,\epsilon_k\,\partial_{\epsilon_k}f^{(x)}(\epsilon) + O\Big((\delta\,\epsilon_k)^3\Big).$$

Divide by $X(\delta) = 2\,\epsilon_k\,\delta$ to get the central–difference estimator

$$\frac{Y(\delta)}{X(\delta)} = \partial_{\epsilon_k}f^{(x)}(\epsilon) + O(\delta^2),$$

i.e. a *bias* of order $O(\delta^2)$ (the constant is controlled by a uniform bound on $\partial_{\epsilon_k}^3 f^{(x)}$ inside the analytic window). Since $g_k^{(x)} = \partial_{\epsilon_k}f^{(x)}(\epsilon)$ by definition (recall $f^{(x)} = -\operatorname{Im}\Lambda_{\mathrm{law}}^{(\mathrm{corr};x)}$ and Lemma 7.675), this shows the basic second–order accuracy.

(b) WLS with multiple symmetric pairs. Collect pairs $\{\pm\delta_j\}$ (e.g. $\delta_j \in \{\delta_0, 2\delta_0\}$) and define (Y_j, X_j) as in the statement. Then

$$Y_j \;=\; X_j\, g_k^{(x)} \;+\; b_j, \qquad |b_j| \;\le\; C\,|\epsilon_k|^3\,|\delta_j|^3,$$

for some C uniform on the certified mask (plateau compactness). Weighted least squares with positive weights w_j solves $\widehat{g}_k^{(x)} = (\sum_j w_j X_j^2)^{-1} \sum_j w_j X_j Y_j$ and satisfies

$$\widehat{g}_k^{(x)} - g_k^{(x)} = \frac{\sum_j w_j X_j b_j}{\sum_j w_j X_j^2} = O\Big(\max_j \delta_j^2\Big) =: O(\delta_0^2),$$

because $X_j \propto \delta_j$ while $b_j = O(\delta_j^3)$. Thus the WLS estimator inherits the *second–order* bias of the central difference and can reduce variance in the presence of heteroscedastic readout noise.

(c) BRST/Nielsen safety. The correlational channel is built from band–descended, BRST–commuting taggers (Definition 7.551 and Proposition 7.573); therefore $f^{(x)}$ is computed from physical matrix elements on $\mathcal{H}_{\mathrm{phys}}$. Gauge–parameter variations are BRST–exact (Nielsen identity), so physical matrix elements and spectral data are gauge–independent (Lemma 7.643 and Proposition 7.556). Consequently, both $g_k^{(x)}$ and its estimator $\widehat{g}_k^{(x)}$ are BRST/Nielsen safe; varying α_{gf} changes neither the target derivative nor the regression design, up to the numerical/plateau tolerance.

(d) Consistency with stiffness tomography. From Lemma 7.675 we have $-\operatorname{Im}\Lambda_{\mathrm{law}}^{(\mathrm{corr};x)} = \frac{1}{2}\epsilon^\top C^{(x)}\epsilon + o(\epsilon^2)$, hence $g_k^{(x)} = (C^{(x)}\epsilon)_k + o(\epsilon)$. The sensitivity–WLS procedure above therefore provides an operational readout of the k–th component of $C^{(x)}\epsilon$ with $O(\delta_0^2)$ bias, justifying its use for localizing explicit breaking (Proposition 7.676).

This proves the lemma. □

Remark on notation. The T–even correlational piece contributes to the *imaginary* part (cf. Lemma 7.675). If an alternative convention is used in which it contributes to the real part, the proof carries over verbatim with Im replaced by Re throughout.

Remark 7.683 (Spatial localization (optional))**.** If spurions are spatially addressable (e.g. patterned sources), define region weights $\{\omega_m\}$ and repeat the sensitivity protocol with local toggles to obtain a coarse spatial map $G_m^{(x)} := \partial_{\omega_m}(-\operatorname{Re}\Lambda_{\mathrm{law}}^{(\mathrm{corr};x)})$. Cone–preserving kernels (Section 7.9.3) ensure causal interpretability.

Definition 7.684 (Reporting (channel–resolved))**.** Per $x \in \mathfrak{P}$ (or inside $\mathfrak{M}_{\mathrm{mask}}$):

1. **Law deficit:** report $1 - |\Lambda_{\mathrm{law}}^{(x)}|$ and its tomographic lower bound $K_{\mathrm{low}}(x)|\mathcal{R}_a^{(x)}|$ (Corollary 7.605); include $\kappa_{\mathrm{spec}}(x)$ and $\widehat{c}_0(x)$.
2. **Spectral (T–odd):** plot $\mathsf{S}_a(x;\omega)$ over $[-\omega_0(x), \omega_0(x)]$ and the kernel $\mathsf{K}_{\mathrm{spec}}$; give $\int \mathsf{K}_{\mathrm{spec}}\mathsf{S}_a$ with CI.
3. **Correlational (T–even):** list $\tilde{\mathbf{g}}^{(x)}$ with CIs; identify dominant spurions/operators.
4. **Vacuum:** state the (near–)unity Λ_{vac} piece (unique vacuum) with the envelope selected in Section 7.9.6.

All quantities are computed on $\mathcal{H}_{\mathrm{phys}}$ and are gauge–flat within the audited tolerances (Section 7.9.14).

7.9.19 Synthesis of QIV

Quadrant IV (QIV) is the "explicitly broken, asymmetrically prepared" regime in which the BRST-improved adjoint-covariant Ward identity on the physical sector carries a non-vanishing source,

$$\nabla_\mu^{(\mathrm{ad})}\, \bar{J}_{a,\mathrm{imp}}^{(\lambda_{\mathrm{UV}},\ell)\mu} = \sum_k \epsilon_k \left\langle \mathcal{S}_{k,a}^{(\lambda_{\mathrm{UV}})} \right\rangle_{\mathbf{\Psi}} \quad \text{on } \mathcal{H}_{\mathrm{phys}} \qquad (eq.\ (7.665)).$$

Because the divergence along broken directions does not vanish on certified plateaus (Theorem 7.570), exact conservation fails and Goldstone's theorem does not apply. Consequently there are **no exact NG poles**; instead the would-be NG modes pick up **small but nonzero gaps** fixed by a GMOR-type identity that holds on the plateau:

$$f_{\pi_b}^2\, m_{\pi_b}^2 = \sum_k \epsilon_k \left\langle \mathcal{S}_{k,b}^{(\lambda_{\mathrm{UV}})} \right\rangle_{\mathbf{\Psi}} + \mathcal{O}(\epsilon^2),$$

with b any broken generator, f_{π_b} the decay constant in the $\epsilon \to 0$ limit, and the slope read off from the band-tagged Ward source (see Theorem 7.630 and Corollary 7.631; background [94, 100, 91, 112]). On a certified mask, this produces the characteristic *linear scaling* $m_{\pi_b}^2 \propto \epsilon$ with vanishing intercept within plateau tolerance.

The *pattern of splittings* across the light sector is dictated by representation theory of the spurions. By construction, the explicit couplings $\{\epsilon_k\}$ transform in the dual G-module to the breaking operators $\{\mathcal{O}_k\}$ (Definition 7.559 and Lemma 7.566); the Ward source is adjoint-covariant (Proposition 7.568). Within any would-be multiplet, *degeneracies split at* $O(\epsilon)$ according to how the spurion irrep projects onto the broken directions. The cleanest way to follow this is the *cluster view* developed for near-crossings: isolate a pNG cluster by a Riesz projector, track its analytic subspace, and use Kato parallel transport for eigenframe labeling (Definition 7.618, Lemma 7.619, and Proposition 7.620). In that setting, the *subspace GMOR map* remains linear, while individual level slopes stay analytic away from avoided crossings.

All of this sits on a robust analytic foundation. The band-limited Hessian $\mathbb{H}^{(x)}(\epsilon)$ on $\mathcal{H}_{\mathrm{phys}}$ is a **type-(A) analytic family** in the explicit couplings (Proposition 7.642); hence *mixing angles are analytic and small*, $\theta = O(\epsilon)$, and can be tracked unambiguously along isolated clusters by Kato transport (Proposition 7.620). The **radial/Higgs sector** remains gapped on the mask, and its masses/couplings shift *analytically* in ϵ (linear terms from explicit sources plus quadratic curvature corrections), with no singular behavior permitted inside the certified mask (Proposition 7.663 and Theorem 7.664). Moreover, **poles and residues are gauge-flat** on $\mathcal{H}_{\mathrm{phys}}$: Nielsen independence holds theoretically (Proposition 7.644 and Theorem 7.646) and operationally via the *Nielsen slope test* (Section 7.9.14), where $S_{\mathrm{Nielsen}} \le \epsilon_{\mathrm{Nielsen}}$ certifies that quoted matrix elements such as $f_{\pi_b} = \langle 0|\bar{J}_{b,\mathrm{imp}}|\pi_b\rangle$ are physically meaningful.

On the diagnostics side, QIV's law channel is driven by a *T-odd spectral weight* tied to the non-conservation source. The spectral (T-odd) component of the holosymmetric readout admits a positive tomography representation,

$$-\,\mathrm{Re}\;\Lambda_{\mathrm{law}}^{(\mathrm{spec};x)} = \int d\omega\, \mathsf{K}_{\mathrm{spec}}(x;\omega)\, \mathsf{S}_a(x;\omega),$$

where S_a is the retarded spectral density of $\nabla^{(\mathrm{ad})} \cdot \bar{J}_{a,\mathrm{imp}}$ and $\mathsf{K}_{\mathrm{spec}}$ is a cone-preserving kernel floor (see Definition 7.607, Lemma 7.575, Proposition 7.609, and Corollary 7.610). This same low-ω weight both **sets the law-visibility deficit** and **predicts pNG masses** through the GMOR fit, closing the Ward–mass loop (Theorems 7.580 and 7.630). Complementarily, the **T-even correlational branch** yields an **operator-space localization vector** $\mathbf{g}^{(x)}$ that identifies which explicit operators dominate the observed stiffness and splittings; its computation by small spurion toggles is BRST/Nielsen safe and provides actionable attribution for mass shifts (Section 7.9.18, Definitions 7.681 and 7.684, and Lemma 7.682).

Finally, QIV's **state channel is trivial** in the thermodynamic sense: explicit terms select a **unique vacuum**, so $|\Lambda_{\mathrm{state}}(x;V)| \to 1$ with a certified envelope (exponential if gapped, power-law if critical but non-SSB), ruling out thermodynamic orthogonality and eliminating *topological vacuum relics* such as domain-wall or string zero modes (Definition 7.582 and Theorem 7.589). All "soft" long-distance physics in QIV is thus encoded in **light pseudo-Goldstones**, not in a vacuum manifold.

These conclusions *sharply separate QIV from the other quadrants*. In **QI** (symmetric vacuum, no explicit breaking; Section 7.1) the law channel *conserves on the band* and there is *no pNG sector*—the spectrum is whatever $\mathcal{L}_0$ provides. In **QII** (SSB; Section 7.7) one finds *exact NG modes* and *state decoherence* with volume (orthogonal vacua), while the law channel is effectively conserved up to improvements; **QIV flips both features**: a **unique vacuum** and **no exact NG**, only pNG with $m_\pi^2 \propto \epsilon$. In **QIII** (small explicit breakings but symmetric state; Section 7.8) the non-conservation *heals in the IR* so the law-deficit improves like $\ell^{\Delta-(d+z)}$, and there is *no asymmetric state*—hence *no pNG sector*. **QIV**, by contrast, shows a **scale-independent saturation deficit** on the law channel together with a **light, linearly scaling pNG band**, both certified on $\mathcal{H}_{\mathrm{phys}}$ and robust under BRST/Nielsen gates and plateau audits (Theorem 7.580, Corollary 7.617, and Propositions 7.569 and 7.616).

Within QIV the **spin content is fixed by the symmetric limit and is not altered by the explicit breaking**. What changes are the **masses and mixings**, not the Lorentz (or rotational) representation carried by each mode. Concretely, work on $\mathcal{H}_{\mathrm{phys}}$ with the QIV hypotheses and the scale–tagged Ward identity (eq. (7.665)). In the *internal–symmetry* case (the QII benchmark breaks $G \to H_0$ without touching spacetime symmetries), every would–be NG is a *Lorentz scalar*; in QIV each becomes a **pseudo–NG scalar** with a small gap from the GMOR relation on the plateau,

$$f_{\pi_b}^2\, m_{\pi_b}^2 = \sum_k \epsilon_k \left\langle \mathcal{S}_{k,b}^{(\lambda_{\mathrm{UV}})} \right\rangle_{\mathbf{\Psi}} + \mathcal{O}(\epsilon^2),$$

so their **spin remains** 0 while $m_{\pi_b}^2 = O(\epsilon)$ (Theorem 7.630 and Corollary 7.631). The **radial (Higgs) modes**—amplitude fluctuations of the order parameter—are also **spin–0** and stay gapped; their masses shift *analytically* in ϵ by type–(A) perturbation theory on the band (Proposition 7.642 and Theorem 7.646).

If some broken directions are *gauged*, BRST projection removes the corresponding would–be NGs ("eaten" longitudinal modes), and the physical spectrum contains **massive spin–1 vectors** with the usual three polarizations; this counting is handled on $\mathcal{H}_{\mathrm{phys}}$ by the Stage-I gate and the descent lemmas (Lemma 7.552) and remains *Nielsen–independent* (Proposition 7.644). Again, the explicit breaking shifts masses/mixings *analytically* in ϵ without changing spin.

If *spacetime symmetries* (e.g. translations/rotations) are the ones broken in the $\epsilon \to 0$ benchmark, the QII NG content is phonon/rotational modes with definite *polarizations* under the spatial rotation group (or its lattice point group). In QIV those become *pseudo–NGs with the same polarization content* (same little-group representation), now with a small gap $O(\epsilon)$. Our scale–aware formalism (dynamic exponent z, cone-preserving taggers) and the Ward–mass link apply verbatim; the *polarization labels* are inherited from the symmetric limit and are unaffected by turning on $\{\epsilon_k\}$ (Section 7.9.3, Definition 7.607, and Theorem 7.630).

Mixings respect these spin/polarization labels: the *band Hessian* $\mathbb{H}^{(x)}(\epsilon)$ decomposes into irreps of the little group, so *modes of different spin (or different J^{PC}/point-group irrep)* do **not** mix at quadratic order unless spacetime symmetry is itself explicitly broken by the spurions. Within a fixed spin/irrep block, mixings are *analytic and* $O(\epsilon)$ and are tracked by *Kato transport* on isolated clusters (Definition 7.618 and Proposition 7.620). All pole residues/decay constants used to identify spins are *gauge-flat* by Nielsen independence (Proposition 7.644 and Theorem 7.630).

1. For *internal* symmetry breaking: expect **spin-0** pNGs (one per QII NG), plus **spin-0** radial modes and whatever other gapped content the QI baseline had; explicit breaking only makes the pNGs light but massive and perturbs gaps/mixings analytically.

2. For *spacetime* symmetry breaking: expect the **same polarization content** as in QII (e.g. longitudinal/transverse phonons), now as **gapped pseudo-modes** with $m^2 = O(\epsilon)$.

3. If some broken directions are *gauged*, the physical spectrum shows **massive spin-1** vectors (Higgs mechanism) and *no* corresponding pNGs on $\mathcal{H}_{\text{phys}}$.

In all cases, QIV's explicit breaking **does not change spin**; it controls **masses and mixings**—and we certify both analytically and gauge-independently on the plateau using the mass/mixing ledger, GMOR fits, and Nielsen tests (Sections 7.9.10 and 7.9.14 and Theorem 7.630).

First, we place the two stacks in the canonical roles we established upstream: the **gaiamiti fields** live in the configuration/order-parameter stack $\hat{\mathbf{\Phi}}$, while the **tantrakyma fields** comprise the gauge-and-scaffold stack $\hat{\mathbf{\Theta}}$ (connections A_μ, gauge-fixing auxiliaries, and BRST doublets). The generalized field vector is $\hat{\mathbf{F}} = (\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}})^\top$ and all statements are taken **on the physical sector** $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$ after the Stage-I gate (nilpotent ζ, anomaly-free) so that our readouts are **Nielsen-independent** and **BRST-safe** (Lemma 7.552, Proposition 7.556, and Theorem 7.646).

Spins and masses from the gaiamiti sector $\hat{\mathbf{\Phi}}$ In the $\epsilon \to 0$ benchmark (QI/QII reference), let the internal symmetry G act on $\hat{\mathbf{\Phi}}$ with vacuum breaking $G \to H_0$. The **broken generators** then label would-be NG directions in the $\hat{\mathbf{\Phi}}$ manifold; these modes carry the **spin content of the symmetric limit**. For **internal** G, that means **spin-0** scalars. Turning on QIV's explicit spurions $\{\epsilon_k\}$ lifts those flat directions to **light pseudo-NG (pNG) scalars** with masses fixed on the plateau by the GMOR-type Ward–mass relation (eq. (7.665), Theorem 7.630, and Corollary 7.631):

$$f_{\pi_b}^2\, m_{\pi_b}^2 = \sum_k \epsilon_k \left\langle \mathcal{S}_{k,b}^{(\lambda_{\text{UV}})} \right\rangle_{\mathbf{\Psi}} + \mathcal{O}(\epsilon^2), \quad b \in \mathfrak{g}/\mathfrak{h}_0.$$

Thus **spin stays** 0 and **mass is generated linearly in the spurions** across the certified mask. Within any would-be multiplet, pNG **mass splittings** appear at $O(\epsilon)$ according to the **spurion irrep** that pairs with $\{\mathcal{O}_k\}$ (Definition 7.559 and Lemma 7.566); slopes are analytic and tracked reliably away from avoided crossings by the **Riesz/Kato** machinery (Definition 7.618 and Proposition 7.620). The **radial (Higgs) modes** in $\hat{\Phi}$ remain **spin-0 and gapped**; their masses are the positive eigenvalues of the band Hessian in the orthogonal (amplitude) directions and shift **analytically** in ϵ by type-(A) perturbation theory (Proposition 7.642 and Theorem 7.646). No singular behavior is permitted inside the plateau mask (Proposition 7.663 and Theorem 7.664).

If, instead, the reference breaking involves *spacetime symmetries* (e.g. broken translations/rotations carried by tensorial components of the gaiamiti sector), the QII benchmark yields phonon/rotational NGs with definite *polarizations* under the spatial little group. In QIV those become *gapped pseudo-phonons* with *the same polarization content*; spins/polarizations are inherited unchanged, while the small gap obeys the same Ward–mass logic (our cone-preserving taggers and dynamic exponent z keep the causal/tomographic control intact; Section 7.9.3 and Definition 7.607).

In the present terminology, *phonons* are a distinguished subclass of **apeirons**. They arise when the *gaiamiti* background breaks spacetime symmetries—typically spatial translations and/or rotations—so that, in Quadrant II, they appear as Nambu–Goldstone apeirons associated with the broken generators, carrying the familiar longitudinal and transverse polarizations. In Quadrant IV, the same modes persist as **pseudo-phonons**: they remain apeirons with identical polarization content, but acquire small gaps through the explicit spurions, with masses governed on the certified plateau by the Ward–GMOR relation linking the non-conserved spacetime currents to the pseudo-NG mass matrix.

Crucially, phonons are *not kymons*. **Kymons** are the quanta of the *tantrakyma* (gauge/scaffold) sector—typically spin-1 excitations after BRST projection—whereas phonons belong to the gaiamiti sector and are determined by broken *spacetime* rather than internal symmetry. For a stricter house style, the term "phonon" may be replaced by *"translational apeiron (longitudinal/transverse)"*, reserving "kymon" exclusively for tantrakyma quanta. In Quadrant IV the translational apeirons obey the same pseudo-NG mass law as their internal-symmetry counterparts; the only difference is that the Ward source is built from the **spacetime Noether currents (stress–energy/elastic currents)**, rather than from internal currents.

Spins and masses from the tantrakyma sector $\hat{\Theta}$ The tantrakyma stack contains the *gauge scaffold*: gauge fields, gauge-fixing auxiliaries, ghosts and anti-ghosts (the latter forming BRST doublets). On $\mathcal{H}_{\text{phys}}$ only *gauge-invariant, BRST-closed* excitations survive. If some of the broken directions of G are *gauged*, the corresponding would-be NGs from the gaiamiti sector are **removed by BRST projection** (they are "eaten"), and one observes **massive spin-1 vectors** (three polarizations) with masses set primarily by the Higgs mechanism (symmetric-limit decay constants and gauge couplings), now receiving *analytic QIV shifts* in ϵ via mixing terms allowed by gauge invariance. Ghosts and Nakanishi–Lautrup auxiliaries do **not** appear as particle-like poles on $\mathcal{H}_{\text{phys}}$ (they cancel in cohomology). Importantly, QIV's explicit breakings are installed as *spurions* precisely so that the band Ward identity remains adjoint-covariant and gauge-consistent (Definition 7.559 and Proposition 7.568); any

"explicitly gauge-noninvariant" operator is only admitted in a way that is *cohomologically trivial* (improvements/BRST-exact pieces), otherwise the Stage-I gate would fail.

Why spins don't change, but masses/mixings do All quadratic data live in the *band Hessian* $\mathbb{H}^{(x)}(\epsilon)$ restricted to $\mathcal{H}_{\text{phys}}$. Because $\mathbb{H}^{(x)}(\epsilon)$ *commutes with the little group* of the spacetime symmetries preserved by the background, it *block-diagonalizes by spin/irrep*. Hence QIV's explicit spurions cannot mix different spins (or different point-group irreps) at quadratic order unless spacetime symmetry is itself explicitly broken. Within a fixed spin/irrep block, however, the *eigenvalues and eigenvectors vary analytically* in ϵ: pNG–pNG and pNG–radial mixings are $O(\epsilon)$, controlled by Kato transport on isolated clusters and reported with the near-degeneracy protocol (Definition 7.618 and Proposition 7.620). The **masses** follow from these eigenvalues (Hessian) and, for the pNG block, are *locked to the Ward source* by GMOR (Theorem 7.630 and Proposition 7.629). The *residues/decay constants* (used to identify spins and couplings) are *Nielsen-flat* by construction (Proposition 7.644 and Section 7.9.14).

Operational cross-checks specific to QIV

- The **law-channel deficit** is driven by the **T-odd spectral weight** of $\nabla^{(\text{ad})} \cdot \bar{J}_{a,\text{imp}}$ (Definition 7.607 and Proposition 7.609); the same low-ω weight determines the **pNG masses** via GMOR. This ties the observed small scalar gaps in the gaiamiti sector directly to measured non-conservation (Theorem 7.580 and Corollary 7.610).
- The **state channel** confirms a **unique vacuum** (no topological relics); exponential or power-law approach $|\Lambda_{\text{state}}| \to 1$ is guaranteed by a mass gap or critical-but-non-SSB scaling (Definition 7.582 and Theorem 7.589).
- If some directions are gauged, expect **spin-1** vectors with analytic ϵ-dependent masses and **no** physical NGs along those directions (BRST descent; Lemma 7.552). Otherwise, all light modes in QIV are **spin-0 pNGs** (one per QII NG), plus **spin-0 radial** excitations that remain gapped and analytic.

In short: for the **gaiamiti** (order-parameter) fields, **spin is inherited** from the symmetric limit (typically scalars), and **mass emerges** by (i) curvature of the potential for radial modes and (ii) **GMOR-controlled explicit breaking** for pNGs. For the **tantrakyma** (gauge scaffold) fields, physical *spin-1 vectors* appear only when the corresponding broken directions are gauged (Higgs mechanism), while ghosts/auxiliaries never survive to $\mathcal{H}_{\text{phys}}$. Across the board, QIV *modifies masses and mixings analytically*—but **does not change spins**—and every statement is made *on certified plateaus*, BRST/Nielsen safe, with adjoint-covariant Ward control (eq. (7.665) and Propositions 7.568 and 7.569).
Formally, the framework now treats **both** "standard–model–like particles" and traditional **quasiparticles** within a single scale–aware, BRST–cohomological machinery, with contributions organized by the two stacks:

- **Gaiamiti** (chronogeometric) stack $\hat{\mathbf{\Phi}}$ supplies **apeirons**: collective/geometric excitations, including the translational/rotational subclasses ("phonons" $\equiv$ translational apeirons) when spacetime symmetries are broken. In QII these appear as exact NG apeirons; in

QIV they become **pseudo–NG apeirons** with small, spurion–controlled gaps certified on the plateau by the Ward–GMOR link (eq. (7.665), Theorem 7.630, and Corollary 7.631). Their mixing and level splittings are analytic in $\{\epsilon_k\}$ (type–(A) family; Proposition 7.642 and Theorem 7.646) and Nielsen–flat on $\mathcal{H}_{\text{phys}}$ (Proposition 7.644).

- **Tantrakyma** (matter/gauge) stack $\hat{\boldsymbol{\Theta}}$ supplies *kymons*: the particle–like excitations of gauge and matter fields (vectors, scalars, spinors) after BRST projection to $\mathcal{H}_{\text{phys}}$. Broken gauged directions "eat" the corresponding apeirons and yield **massive spin–1** kymons; ungauged directions leave light apeirons in the spectrum. Masses and residues of kymons shift *analytically* with explicit breakers while preserving spin/irrep blocks, and stay *gauge–parameter independent* by Nielsen identities (Propositions 7.569 and 7.644).

Operationally, both classes enter identically as *band–limited poles/residues* on $\mathcal{H}_{\text{phys}}$, extracted from the *Hessian/mass ledger* and certified inside a *plateau mask* (Proposition 7.663 and Theorem 7.664). The *law–channel* T–odd spectral weight of $\nabla^{(\text{ad})} \cdot \bar{J}_{a,\text{imp}}$ fixes the persistent deficit (Theorem 7.580) and simultaneously governs *pNG masses* through GMOR; the *state channel* enforces a *unique vacuum* in QIV (Definition 7.582 and Theorem 7.589). Hence "standard–model particles" (kymons of the tantrakyma sector) and "quasiparticles" (apeirons of the gaiamiti sector) are unified as certified spectral features whose *spins are inherited from the symmetric limit* and whose *masses/mixings* are controlled either by curvature (radial/Higgs) or by *explicit spurions* (pNGs), all within the same BRST/Nielsen–safe, adjoint–covariant Ward framework.

This reframes "fundamental" from a field–content notion to a **spectral–cohomological** notion. In this treatise the objects that qualify as fundamental are those **BRST–closed, gauge–parameter–independent poles on $\mathcal{H}_{\text{phys}}$** that remain **plateau–stable** under UV/IR tagging and mesh refinement. Standard–Model–like excitations then appear as *kymons* (tantrakyma quanta), while familiar quasiparticles appear as *apeirons* (gaiamiti quanta). Both are treated on exactly the same footing: as certified spectral features with spins fixed by the symmetric limit, and with masses and mixings controlled either by *curvature of the band Hessian* (radial/Higgs–type) or by *explicit spurions* via the *Ward–GMOR* link for pseudo–NG sectors (eq. (7.665), Theorem 7.630, and Corollary 7.631). In this view, "fundamental" means **cohomologically physical and instrument–robust**, not "elementary field in the Lagrangian."

Crucially, nothing in this reclassification contradicts established constraints: charges and spins are inherited from the symmetric benchmark; gauged breakings Higgs–eat the relevant apeirons; anomalies are excluded at the Stage–I gate; poles/residues are *Nielsen–flat*; and all quoted spectra live inside a *certified plateau mask* (Propositions 7.644 and 7.663 and Theorem 7.664). What changes is the organizing principle: Standard–Model particles and traditional quasiparticles become two manifestations (kymons vs. apeirons) of the same underlying **scale–aware, BRST–cohomological spectrum**, with **law–channel visibilities** and *T–odd tomography* tying observed gaps directly to controlled non–conservation on $\mathcal{H}_{\text{phys}}$ (Theorem 7.580, Definition 7.607, and Corollary 7.610).

7.10 Consolidation of Law–State Signatures, Ward Sources, and Spectral Content (QI–QIV)

This section consolidates the four laws–states quadrants into a single operational ledger, aligning the interferometric visibilities, the scale-tagged Ward identities, and the certified spectral content into one coherent diagnostic. All statements are made on certified plateaus $\mathfrak{P}$ and, where invoked, within the IR mask $\mathfrak{M}$ fixed by the acceptance gates and slope/curvature/oscillation metrics (eq. (7.28), Proposition 6.288, Definition 7.42, Theorem 6.253, and Sections 7.8.9 and 7.9). UV/IR instrumentation follows the spectral projector $P_x(\lambda_{\mathrm{UV}})$ and the IR window $K^{(\ell)}_{*\mathrm{IR}}$ (Definitions 7.26 and 7.35 and eq. (7.29)), with all readouts living on the BRST cohomology $\mathcal{H}_{*\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$; gauge-parameter independence is ensured by the Nielsen identities and gauge-safe analyticity (Propositions 6.276 and 7.644 and Theorem 7.646). Unless stated otherwise, operators and residues are those induced by the holosymmetric lift on $\mathfrak{P}$ and restricted to $\mathcal{H}_{\mathrm{phys}}$.

The quadrant logic is governed by the visibilities $\Lambda_{\mathrm{law}}, \Lambda_{\mathrm{state}}$ and their overlays (Definitions 7.35 and 7.42). For ease of reference, Definition 7.685 specifies the assignment rules on a certified plateau $\mathfrak{P}$, separating the symmetric/symmetric baseline (QI), the symmetric law with state-level SSB (QII), the IR-healing of explicit-looking law deficits (QIII), and the persistent law deficit with a unique state (QIV). This classification is not merely taxonomic: it is the interface through which Ward statements and spectral data are read consistently across scales.

Definition 7.685 (Quadrant decision rule on a certified plateau $\mathfrak{P}$)**.** Write $u = \ln \lambda_{\mathrm{UV}}$, $v = \ln \ell$. On a certified plateau $\mathfrak{P}$ (Section 6.12.4), interpret "≈ 1" as $\big|\,|\cdot| - 1\big| \le \varepsilon_{\mathrm{tol}}$ uniformly on $\mathfrak{P}$. Given the visibilities $(\Lambda_{\mathrm{law}}, \Lambda_{\mathrm{state}})$ on $\mathfrak{P}$, set

$$\mathrm{Quad}(\Lambda_{\mathrm{law}}, \Lambda_{\mathrm{state}}) = \begin{cases} \textbf{QI}, & |\Lambda_{\mathrm{law}}| \approx 1, \quad |\Lambda_{\mathrm{state}}| \approx 1, \\ \textbf{QII}, & |\Lambda_{\mathrm{law}}| \approx 1, \quad \Lambda_{\mathrm{state}}(V) \to 0 \ (V \to \infty), \\ \textbf{QIII}, & 1 - |\Lambda_{\mathrm{law}}(\ell)| = \mathcal{O}\Big(\ell^{\Delta-(d+z)}\Big), \quad |\Lambda_{\mathrm{state}}| \approx 1, \\ \textbf{QIV}, & \exists\, \delta_\star \in (0,1): \ |\Lambda_{\mathrm{law}}| \le 1 - \delta_\star \text{ uniformly on } \mathfrak{P}, \\ & \qquad\qquad \Lambda_{\mathrm{state}}(V) \to 1 \ (V \to \infty). \end{cases}$$

Remark 7.686 (Intuition & reading guide on a certified plateau $\mathfrak{P}$)**.** *Intuition.*

- **QI** is the fully coherent reference: both visibilities are ≈ 1 on $\mathfrak{P}$ and furnish the calibration baseline.
- **QII** preserves law-level conservation but exhibits state-level decoherence driven by a vacuum manifold; distinct vacua orthogonalize as $V \to \infty$.
- **QIII** shows explicit-looking law asymmetries that *heal* under coarse-graining: $1 - |\Lambda_{\mathrm{law}}(\ell)| = \mathcal{O}\Big(\ell^{\Delta-(d+z)}\Big)$ on $\mathfrak{P}$, reflecting RG irrelevance.
- **QIV** retains a *persistent* law deficit $|\Lambda_{\mathrm{law}}| \le 1 - \delta_\star$ on $\mathfrak{P}$, tied to a non-vanishing Ward source; the state is uniquely prepared (no vacuum cat).

Reading guide. Measure the visibilities on $\mathfrak{P}$ *after* BRST projection (evaluate in $\mathcal{H}_{\mathrm{phys}}$).

1. If the law's deficit scales away as in eq. (7.746), classify as **QIII**.
2. If a *persistent* law gap remains (uniform $\delta_\star > 0$), classify as **QIV** and expect GMOR-type gaps via eq. (7.747).
3. Otherwise (law ≈ 1 on $\mathfrak{P}$), decide between **QI**/**QII** purely by the state channel: $|\Lambda_{\text{state}}(V)| \to 0 \Rightarrow$ **QII**; $|\Lambda_{\text{state}}| \approx 1 \Rightarrow$ **QI**.

All statements are plateau-uniform (tolerance ε_{tol}) and gauge-parameter independent by the Nielsen/cohomological identities.

Theorem 7.687 (Equivalence of signatures, Ward statements, and spectral content)**.** *On a certified plateau $\mathfrak{P}$ (and inside the QIII mask $\mathfrak{M}$ when invoked), the following data are mutually consistent and diagnostically equivalent for quadrant assignment:*

Visibility overlays.
Visibility overlays of $\Lambda_{\text{law}}, \Lambda_{\text{state}}$ (Definition 7.42).

Scale-tagged Ward statements.
Scale-tagged Ward statements for improved currents (Definition 6.272): QI/QII exhibit band conservation up to improvements on $\mathcal{H}_{\text{phys}}$ (Sections 7.1 and 7.7); QIII has adjoint divergence trending to zero with ℓ (Theorem 6.253 and Section 7.8.9); QIV carries a non-vanishing adjoint-covariant source (eq. (7.665) and Theorem 7.570).

Light-sector spectrum and residues.
Light-sector spectrum and residues on $\mathcal{H}_{\text{phys}}$ (Nielsen-flat; Proposition 7.644 and Theorem 7.646): QI has no NG sector; QII supports exact NG poles with standard counting [94, 91]; QIII shows no NG/pNG sector with IR healing; QIV exhibits a pseudo-NG band with GMOR slope on the plateau (Theorem 7.630 and Corollary 7.631; [100, 112]).

Proof of Theorem 7.687. **Standing hypotheses and notation.** Work on the certified plateau $\mathfrak{P}$ (and inside the QIII mask $\mathfrak{M}$ when invoked). All readouts are evaluated on the physical sector $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$ with BRST–closed scale tags $P_x(\lambda_{\text{UV}})$, $K_{\text{IR}}^{(\ell)}$ commuting with the channel projectors, so Nielsen identities hold (Propositions 6.264 and 6.276). Write $\bar{J}_{a,\text{imp}}^{(\lambda_{\text{UV}},\ell)\mu}$ for the improved current and $\mathcal{S}_{k,a}^{(\lambda_{\text{UV}})}$ for the tagged symmetry–breaking insertions. The scale–tagged Ward statement on $\mathcal{H}_{\text{phys}}$ is (Proposition 6.296)

$$\nabla_\mu^{(\text{ad})} \bar{J}_{a,\text{imp}}^{(\lambda_{\text{UV}},\ell)\mu} = \sum_k \epsilon_k(\ell) \left\langle \mathcal{S}_{k,a}^{(\lambda_{\text{UV}})} \right\rangle_{\Psi}. \tag{7.745}$$

Visibilities $(\Lambda_{\text{law}}, \Lambda_{\text{state}})$ are as in Definitions 6.275 and 7.42, with plateau uniformity as in Remark 6.254. "Nielsen–flat" spectra means gauge–parameter independence on $\mathcal{H}_{\text{phys}}$ (Proposition 7.644 in Chapter 8).

Outline. We prove mutual equivalence by showing the implications

$$\text{(1) overlays} \Rightarrow \text{(2) Ward)} \Rightarrow \text{(3) spectrum} \Rightarrow \text{(1) overlays},$$

and we do so quadrant by quadrant, using plateau uniformity.

(1) ⇒ (2): Visibility overlays determine the Ward pattern.

- *QI/QII (law coherent).* On $\mathfrak{P}$, $|\Lambda_{\text{law}}| \approx 1$ with only IR–irrelevant corrections (Theorems 6.253 and 6.258). The Duhamel/BCH expansion for the two–path operator (cf. proof of Theorem 6.258) bounds the law–deficit by the integrated norm of the tagged breaking insertions. Plateau–smallness of the deficit forces the r.h.s. of (7.745) to vanish on $\mathfrak{P}$ (up to ζ–exact improvements), i.e. adjoint conservation bandwise.
- *QIII (law healing).* The overlay reads $1 - |\Lambda_{\text{law}}(\ell)| = \mathcal{O}(\ell^{\Delta-(d+z)})$ with $\Delta > d + z$ (Theorem 6.253). The same Duhamel estimate gives $\sum_k \epsilon_k(\ell)\langle \mathcal{S}_{k,a}^{(\lambda_{\text{UV}})}\rangle = \mathcal{O}(\ell^{\Delta-(d+z)})$, i.e. the adjoint divergence in (7.745) trends to 0 with the same exponent (uniformly on $\mathfrak{P}$).
- *QIV (persistent law deficit).* A uniform gap $|\Lambda_{\text{law}}| \le 1 - \delta_\star$ on $\mathfrak{P}$ implies a nonzero lower bound on the integrated breaking strength; thus the r.h.s. of (7.745) does not vanish on $\mathfrak{P}$ (Theorem 7.570). Hence a persistent adjoint–covariant source appears.

(2) ⇒ (3): Ward pattern fixes the light–sector spectrum. Consider the spectral representation of current–current correlators on $\mathcal{H}_{\text{phys}}$, with residues in the channel selected by the plateau tags (Källén–Lehmann on $\mathfrak{P}$; gauge–flat by Nielsen identities).

- *QI.* Exact band conservation and a unique symmetric state forbid massless poles in the G–charged two–point functions. The spectral density is regular at $p^2 = 0$; no NG sector.
- *QII.* Exact band conservation but state asymmetry (SSB for global G) implies Ward–Takahashi identities with nontrivial vacuum order parameter. Standard arguments (Goldstone theorem) give exact NG poles with the usual counting ([94, 91]); residues are Nielsen–flat on $\mathcal{H}_{\text{phys}}$ (Theorem 7.646).
- *QIII.* The adjoint divergence in (7.745) tends to zero with ℓ, but is strictly zero only in the IR limit. On a finite plateau, this precludes protected massless poles: any would-be pole would force a non-decaying RHS by LSZ/WT analysis. Thus no NG/pNG sector appears; instead the small explicit insertions dress regular light continua whose strength decays with ℓ (Section 7.8.9).
- *QIV.* A nonvanishing adjoint source implies a PCAC–type relation. In the light sector, this yields pseudo–Nambu–Goldstones with $m_{\text{pNG}}^2 f^2 \propto$ (Ward source) and hence a GMOR slope on the plateau (Theorem 7.630 and Corollary 7.631; see also [100, 112]).

(3) ⇒ (1): Spectrum and residues reproduce the overlays. Write the holosymmetric law–visibility as a controlled overlap of two tagged evolutions (Chapter 7). Low–energy spectral data dominate the overlap decay on $\mathfrak{P}$.

- *QI.* No NG sector and a unique symmetric vacuum ⇒ no infrared orthogonality and no long–time dephasing in the tagged evolutions. Therefore $|\Lambda_{\text{law}}| \approx 1$ and $|\Lambda_{\text{state}}| \approx 1$.
- *QII.* Exact NG poles induce phase–twist stiffness and vacuum–sector orthogonality as $V \to \infty$ (Theorem 6.261); hence $|\Lambda_{\text{law}}| \approx 1$ (law conserved) while $\Lambda_{\text{state}}(V) \to 0$.
- *QIII.* With no NG/pNG sector and only irrelevant breakings, the tagged overlap improves with coarse–graining; RG power counting gives $1 - |\Lambda_{\text{law}}(\ell)| = \mathcal{O}(\ell^{\Delta-(d+z)})$ (the "healing" law; Theorem 6.262), and the symmetric state yields $|\Lambda_{\text{state}}| \approx 1$.

- *QIV.* A pNG band with GMOR slope enforces a finite, scale–stable dephasing in the law–overlap on $\mathfrak{P}$ (its residue cannot be removed by coarse–graining because the Ward source is nonzero); thus $|\Lambda_{\text{law}}| \le 1-\delta_\star$ uniformly. The unique vacuum gives $|\Lambda_{\text{state}}(V)| \to 1$.

Plateau uniformity and gauge independence. All implications are uniform on $\mathfrak{P}$ by Remark 6.254 and inherit Nielsen–flatness by Proposition 6.299. Channel decompositions are preserved by functoriality and cone stability (Propositions 6.257 and 6.281 and Corollary 6.263).

Conclusion. For each quadrant, the three descriptions—(1) visibility overlays, (2) scale–tagged Ward statements, and (3) light–sector spectral content—are equivalent and mutually consistent on a certified plateau $\mathfrak{P}$ (and within $\mathfrak{M}$ for QIII). Any one determines the other two, completing the proof. □

Intuition. The quadrant is not inferred from a single scalar: the interferometric signatures, the scale-tagged Ward identity, and certified poles/residues *cohere*. In QIV, the persistent law deficit $\Longleftrightarrow$ the non-zero Ward source $\Longleftrightarrow$ the pNG band with GMOR scaling; in QIII, the decaying deficit $\Longleftrightarrow$ vanishing adjoint source in the IR $\Longleftrightarrow$ absence of NG/pNG modes.

Key boxed relations

$$\boxed{1-|\Lambda_{\text{law}}(\ell)| = \mathcal{O}\big(\ell^{\Delta-(d+z)}\big) \qquad \text{(QIII on the mask } \mathfrak{M}\text{; Theorem 6.253 and Section 7.8.9).}} \tag{7.746}$$

$$\boxed{\nabla^{(\text{ad})\mu}\bar{J}^{(\lambda_{\text{UV}},\ell)\mu}_{a,\text{imp}} = \sum_k \epsilon_k \big\langle \mathcal{S}^{(\lambda_{\text{UV}})}_{k,a}\big\rangle_{\mathbf{\Psi}} \;\Rightarrow\; f^2_{\pi_b} m^2_{\pi_b} = \sum_k \epsilon_k \big\langle \mathcal{S}^{(\lambda_{\text{UV}})}_{k,b}\big\rangle_{\mathbf{\Psi}} + \mathcal{O}(\epsilon^2)} \tag{7.747}$$

(QIV on $\mathcal{H}_{\text{phys}}$; Ward source $\neq 0$ and GMOR fit on the plateau: eq. (7.665), Theorem 7.630, and Corollary 7.631.)

The two boxed relations are decisive for separating the IR–healing regime (QIII) from the explicitly persistent regime (QIV). In particular, eq. (7.746) asserts that any law-channel deficit decays as a power of the IR dilation scale, while eq. (7.747) identifies a non-vanishing adjoint-covariant source that pins a strictly positive plateau deficit. Together they make the overlay rules in Definition 7.685 operational beyond taxonomy, providing the precise hinge on which the existence of a certified QIII versus a certified QIV plateau turns; this is exactly the content that underwrites the equivalence claims in Theorem 7.687.

These relations also close the three-way correspondence between interferometric visibilities, scale-tagged Ward structure, and spectral content. On the one hand, eq. (7.746) implies the vanishing of the (improved) adjoint divergence on the IR mask, hence no protected discontinuity remains to be saturated and no NG/pNG sector can persist (Theorem 6.253 and Section 7.8.9). On the other hand, eq. (7.747) ties a non-zero adjoint source directly to the pseudo-Goldstone band and fixes its leading scaling on the plateau via the GMOR fit (Theorem 7.630 and Corollary 7.631). This implements, in both directions, the diagnostic equivalences summarized in Theorem 7.687 using only the visibility overlays from Definitions 7.35 and 7.42 and the scale-tagged equations of motion.

Moreover, the pair furnishes a quantitative and falsifiable calculus. The decay rate in eq. (7.746), controlled by $\Delta - (d + z)$, makes the QIII assignment testable by slope/curvature metrics on certified plateaus and masks, while the source–mass bridge in eq. (7.747) enables an inversion-ready QIV analysis: measured residues $f_{\pi_b}^2$ and masses $m_{\pi_b}^2$ determine (or are predicted from) the explicit breaking coefficients $\{\epsilon_k\}$. Both operations are defined with the UV projector and IR window instrumentation $P_x(\lambda_{\mathrm{UV}})$, $K_{\mathrm{IR}}^{(\ell)}$ (Definition 7.26 and eq. (7.29)) and the readout protocol of Definition 7.35, thereby rendering the classification reproducible with fixed scale tags.

Crucially, the statements are regulator- and gauge-parameter independent on the physical Hilbert space. All currents are improved and all correlators are restricted to $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$, so BRST/Nielsen identities and gauge-safe analyticity ensure that neither the decay law in eq. (7.746) nor the GMOR slope in eq. (7.747) can be artefacts of gauge choice or regulator details (Propositions 6.276 and 7.644 and Theorem 7.646). This is the sense in which the ledger's decisions are stable under admissible changes of presentation.

The spectral implications follow without additional assumptions. In QIII, IR healing of the law channel as in eq. (7.746) excludes NG and pNG light sectors and enforces multiplet reconvergence under coarse-graining (Theorem 6.253 and Section 7.8.9), in agreement with the synopsis of Section 7.10.1. In QIV, the non-zero source in eq. (7.747) both necessitates a pNG band and fixes its leading scaling (GMOR on the plateau), while radial modes remain analytic and gapped—again matching the entries in Section 7.10.1.

Finally, these two relations can be viewed as the minimal scale-sensitive normal forms of the ledger. The conserved-law corner (QI/QII) in Definition 7.685 leaves them inert, with QII's state-channel decoherence orthogonal to the law channel. Consequently, all nontrivial flows pertinent to quadrant assignment reduce to verifying the decay prescribed by eq. (7.746) or the source–slope linkage in eq. (7.747). Practically, the pipeline—choose scale tags, certify $\mathfrak{P}$ (and $\mathfrak{M}$ when invoked), read visibilities, evaluate the adjoint divergence, and fit poles/residues—culminates in a cross-check against these two boxed relations, after which Definition 7.685 and Theorem 7.687 yield a regulator-robust decision.

7.10.1 Comparative Table

Table 7.4: Quadrant synopsis on a certified plateau $\mathfrak{P}$.

Quadrant	**Law channel**	**State channel**	**Low-energy sector (IR window on $\mathcal{H}_{\text{phys}}$)**
QI	Conserved on band (improved)	Unique/coherent	No NG; baseline gapped content; Nielsen-flat residues
QII	Conserved on band	Orthogonality: $\Lambda_{\text{state}}(V) \to 0$	Exact NGs (Goldstone); gauged directions eaten
QIII	Deficit heals: $\mathcal{O}(\ell^{\Delta-(d+z)})$	Unique/coherent	No NG/pNG; multiplet reconvergence; improved physicality
QIV	Persistent deficit $\leq 1-\delta_\star$	Unique/coherent	pNGs with GMOR slope; radial analytic and gapped; gauge-flat poles

Narrative synthesis and operational reading. On QI plateaus, both visibilities saturate (Definition 7.685), the improved current is conserved on band (Section 7.1), and the light spectrum matches the symmetric law without NG content, with residues flat under gauge-parameter variations (Proposition 7.644 and Theorem 7.646). QII inherits the same law conservation but Λ_{state} decoheres with volume, resolving a vacuum manifold; the Goldstone sector appears as exact poles with standard counting and, under gauging, is re-packaged into massive vector modes (Section 7.7, [94, 91]). QIII exhibits a non-saturating law channel at finite scale that *heals* under IR dilation as in eq. (7.746); the associated Ward divergence tends to zero on $\mathfrak{M}$ and no NG/pNG modes are present (Theorem 6.253 and Section 7.8.9). QIV is singled out by a uniform, scale-stable deficit in the law visibility on $\mathfrak{P}$, a non-vanishing adjoint-covariant source, and a pNG band obeying the GMOR fit on the plateau (eqs. (7.665) and (7.747), Theorem 7.630, and Corollary 7.631); the state channel is unique, and radial fluctuations remain analytic and gapped.

Finally, because all objects are computed within $\mathcal{H}_{\text{phys}}$ and certified by the plateau gates, the ledger is regulator-robust and gauge-safe (Propositions 6.276 and 7.644 and Theorem 7.646). In practice: (i) choose scale tags $(\lambda_{\text{UV}}, \ell)$ and the channel decomposition; (ii) certify $\mathfrak{P}$ and, where needed, $\mathfrak{M}$; (iii) read $\Lambda_{\text{law}}, \Lambda_{\text{state}}$, the scale-tagged Ward divergence, and the light-sector residues; (iv) assign the quadrant via Definition 7.685 and validate by Theorem 7.687. This completes the consolidated, cross-validated map from interferometric signatures to Ward sources to spectral content across the four quadrants.

Table 7.5: Spins and modes across quadrants on a certified plateau $\mathfrak{P}$ (and, where invoked, within the IR mask $\mathfrak{M}$). Entries are stated on $\mathcal{H}_{\text{phys}}$ and are gauge-parameter independent by BRST/Nielsen safety (Propositions 6.276 and 7.644 and Theorem 7.646). Quadrant assignment follows Definition 7.685; IR healing and explicit persistence follow eqs. (7.746) and (7.747). Here G_{IR} denotes the IR-recovered unbroken gauge group after band/BRST cleaning.

Species / Mode	**QI** (law $\simeq$ 1, state $\simeq$ 1)	**QII** (law $\simeq$ 1, state decoheres)	**QIII** (law heals, state $\simeq$ 1)	**QIV** (law deficit persists, state $\to$ 1)
Spin-2, massless (helicity ± 2)	Present as in symmetric law	Present as in symmetric law	Present; IR re-converges to QI	Present iff space-time symmetry unbroken
Spin-1, massless (unbroken gauge)	Present for each unbroken generator	Present for each unbroken generator	Present for each generator of G_{IR}; reconverges to QI	Present for each unbroken generator
Spin-1, massive (gauged broken)	Absent (no breaking)	Present (BEH: would-be NG eaten)	Absent in strict IR (no SSB); any finite-ℓ emergent masses vanish as $\ell \to \infty$ by eq. (7.746)	Present when broken directions are gauged (BEH), with analytic mass/mixing shifts
Spin-0, NG (global broken)	Absent	Present: one per broken global generator	Absent in strict IR (healing as eq. (7.746))	Absent (replaced by pNG)
Spin-0, pNG (GMOR)	Absent	Absent	Absent in strict IR; finite-ℓ transients must vanish with ℓ	Present with $m^2_{\pi_b} = \mathcal{O}(\epsilon)$ and slope fixed by eq. (7.747)
Spin-0, radial (Higgs-like)	Present, gapped, analytic	Present, gapped, analytic	Present, gapped; reconverges to QI	Present, gapped, analytic; spin unchanged under explicit breaking
Spacetime NG (phonons/rotations)	Absent	Present if space-time symmetry is spontaneously broken (longitudinal/transverse polarizations)	Absent in strict IR (no SSB)	Absent as exact NG; replaced by pseudo-phonons if explicit space-time breaking

Continued on next page...

Table 7.5: – *Continued from previous page*

Species / Mode	**QI** (law $\simeq$ 1, state $\simeq$ 1)	**QII** (law $\simeq$ 1, state decoheres)	**QIII** (law heals, state $\simeq$ 1)	**QIV** (law deficit persists, state $\to$ 1)
Spacetime pseudo-phonons	Absent	Absent	Absent in strict IR	Present with same polarization content as QII, now gapped $\mathcal{O}(\epsilon)$
Ghosts / auxiliaries	Unphysical; projected out by $\mathsf{P}_{\mathrm{phys}}$	Unphysical; projected out	Unphysical; projected out	Unphysical; projected out

Notes. (i) "Reconverges to QI" indicates channel-wise approach of spectra/residues under IR dilation consistent with eq. (7.746). (ii) In QIV, spins are inherited from the symmetric limit; explicit breaking shifts masses/mixings analytically without changing spin, and the pNG sector obeys the plateau GMOR fit (eq. (7.747), Theorem 7.630, and Corollary 7.631). (iii) All entries are to be read within the certified plateau $\mathfrak{P}$ (and mask $\mathfrak{M}$ when invoked) and under the holosymmetric instrumentation of Definitions 7.26 and 7.35 and eq. (7.29).

The spin assignments that accompany tables 7.5 and 7.6 are fixed by a single chain of band–/BRST–reduced constructions and identities, all evaluated on the physical Hilbert space $\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}})$ and within certified plateaus $\mathfrak{P}$.

First, the *physical-sector pole analysis* determines spins and helicities kinematically and reads them dynamically. From the reduced quadratic kernel on $\mathcal{H}_{\mathrm{phys}}$ (constructed after the decoupling of Kugo–Ojima quartets), the *physical mass operator* $\mathbb{M}^2_{\mathrm{phys}}$ is defined blockwise and its poles m_X^2 and residue matrices Z_X are extracted in the polarization blocks $X \in \{A^T,\ h_{\mathrm{TT}},\ \pi\}$; these identify, respectively, massless/massive vectors ($s = 1$), transverse–traceless tensors ($s = 2$), and scalars ($s = 0$) (cf. eqs. (7.485) and (7.486)). The same construction enforces non-negativity of $\mathbb{M}^2_{\mathrm{phys}}$ and a Källén–Lehmann form with positive spectral density on $\mathcal{H}_{\mathrm{phys}}$, making the pole/residue identification intrinsic (i.e. not gauge artifacts). Gauge-parameter independence of pole positions and residues follows from the Nielsen/Slavnov–Taylor identities proved on the band, again after projection to $\mathcal{H}_{\mathrm{phys}}$ (see Proposition 7.644 and Theorem 7.646). [91, 82, 117, 243]

Second, the *Ward/Noether ledger* ties the scalar versus vector content to the presence of symmetry breaking in QuadrantĨI. With *symmetric law* and *broken global symmetry*, the Ward identities and the spectral representation force *Nambu–Goldstone* poles in the correlational channel, hence *spin–0* excitations. When the broken directions are *unbroken gauged*, the would-be Goldstones are ζ-exact and are eaten by the Brout–Englert–Higgs mechanism, producing *massive spin–1* vectors with the longitudinal polarization restored; the ST/Nielsen identities ensure that these physical poles are gauge independent on $\mathcal{H}_{\mathrm{phys}}$ (cf. Section 7.5.8 and Proposition 7.644). [91, 117]

Third, *spacetime symmetry breaking* produces phononic/rotational NG modes with definite longitudinal/transverse polarizations, entered as *spin–0* excitations in QuadrantĨI and

continued into QuadrantÎV as *pseudo-phonons*. In QuadrantÎV the Ward–source bridge fixes the gap scale $m_{\mathrm{ph}}^2 = \mathcal{O}(\epsilon)$ in precise analogy with the internal pseudo–NG sector, and the polarization content is inherited unchanged (cf. eq. (7.747) and Section 7.10.1). [91]

Fourth, *QuadrantÎII healing* excludes persistent NG/pNG bands and reconverges the spin census to the QuadrantÎ baseline. The adjoint-covariant source decays under IR dilation on the mask $\mathfrak{M}$, so no NG/pNG sector survives asymptotically; the only particle-like content that remains is the symmetric one: massless $s = 2$ (when the relevant spacetime symmetry is unbroken), massless $s = 1$ for the IR–recovered unbroken subgroup, and gapped scalar radials (cf. eq. (7.746)).

Fifth, in *QuadrantÎV* the *PCAC/GMOR* relation on a plateau yields a *pseudo–NG* band with $m_{\pi_b}^2 f_{\pi_b}^2$ proportional to the explicit source; the *spin remains* 0. More generally, pole positions and residues within any isolated spectral subspace move *analytically* under small explicit deformations, while the spin/helicity labels are invariant (helicity stability). This is encoded by the analytic Riesz–projector calculus and the gauge-independence theorem for spectral data on $\mathcal{H}_{\mathrm{phys}}$ (cf. Propositions 7.644 and 7.645). [115, 120, 121, 108, 117]

Sixth, the *mode–helicity catalog* (table 7.6) consolidates the outcome: a massless TT tensor ($s = 2, \lambda = \pm 2$) whenever the relevant spacetime symmetry is unbroken (QI, QII, and the QIII limit; also in QIV for the unbroken spacetime sector); a massless transverse vector ($s = 1, \lambda = \pm 1$) for each unbroken gauge generator; a massive vector ($s = 1, \{+1, 0, -1\}$) when broken directions are gauged (QII/QIV); a gapped scalar radial in all quadrants; and the NG/pNG entries just discussed (NG only in QII; GMOR-controlled pNG in QIV). These are the species-level restatements of the blockwise pole/residue analysis above and of the Ward/SSB/BEH logic (cf. Proposition 6.276 and Theorem 7.646). [91, 117]

Finally, the *quadrant overlay by spin* formalizes how the law–/state–visibilities behave per spin across QI–QIV. Combining the tolerance semantics on $\mathfrak{P}$ with helicity stability and analytic transport of Riesz projectors ensures that, on the band, $|\Lambda^{[s]}_{\mathrm{law/state}}|$ satisfies the same QI–QIV inequalities for each spin sector, with QIII power-law healing and a QIV uniform deficit where appropriate (cf. Definition 7.685 and eqs. (7.746) and (7.747)). The upshot is that the *identification* of spins comes from polarization-resolved poles of $\mathcal{D}_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}$ (via $\mathbb{H}_{\mathrm{phys}}$, Källén–Lehmann, and TT/transverse/scalar projectors), while the *justification* of which spinful modes actually occur in each quadrant is supplied by the Ward/SSB/BEH analysis (QII), IR healing (QIII), and PCAC/GMOR plus helicity stability (QIV), all **BRST/Nielsen-safe** on certified plateaus. [241, 442, 117, 91]

Consolidation of Particle-like Sector

All statements below are made on certified plateaus $\mathfrak{P}$ and, where invoked, within the IR mask $\mathfrak{M}$, with UV/IR instrumentation $(P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}^{(\ell)})$ and holosymmetric readouts on $\mathcal{H}_{\mathrm{phys}}$ (Definitions 7.26 and 7.35 and eq. (7.29)). Gauge-parameter independence and analytic transport of poles/residues are ensured by BRST/Nielsen safety (Propositions 6.276 and 7.644 and Theorem 7.646). Quadrant assignment follows Definition 7.685, and the equivalences between signatures, Ward structure, and spectrum are as in Theorem 7.687. For a quadrant-by-quadrant inventory see also table 7.5.

Massless carriers (reference census). (i) A massless spin-2 sector (helicity ± 2) persists whenever the spacetime symmetry relevant to the low-energy little group is unbroken; this provides the universal reference for spectral health in QI, QII, and the QIII limit, and remains present in QIV for the unbroken spacetime sector. (ii) For each unbroken gauge generator, there is a massless spin-1 vector. These entries are the baseline "conserved-law" content leveraged across quadrants (Definition 7.685 and Section 7.10.1).

Goldstone ladders and their QIV continuations. In QII (symmetric law, state SSB), each broken *global* generator yields an exact NG scalar (spin-0), while broken *gauged* directions yield massive spin-1 vectors via the Brout–Englert–Higgs mechanism (Section 7.7). In QIV (persistent explicit law asymmetry; unique state), the QII NG sector is replaced by a pseudo-NG band with GMOR scaling on the plateau, with masses $m_{\pi_b}^2 = \mathcal{O}(\epsilon)$ and slope fixed by the source–mass bridge (eq. (7.747), Theorem 7.630, and Corollary 7.631). The polarization/spin content of the QIV pseudo-modes is inherited from the corresponding QII NG sector (spins do not change under explicit breaking).

Spacetime NG modes and pseudo-phonons. If spacetime symmetry is spontaneously broken (QII), the spectrum contains phononic/rotational NG modes with definite longitudinal/transverse polarizations under the spatial little group. In QIV, explicit spacetime breaking replaces these by pseudo-phonons with the same polarization content but gapped at $\mathcal{O}(\epsilon)$, governed by the same Ward–source logic as the internal pseudo-NG sector (eq. (7.747) and Section 7.10.1).

Massive vectors from gauged breaking (QII and QIV). Whenever the broken directions are gauged, the would-be NGs are eaten and one observes massive spin-1 vectors with three physical polarizations on $\mathcal{H}_{\text{phys}}$. In QIV, their masses and possible mixings shift analytically with the explicit spurions, but spin/polarization content remains that of spin-1; residues stay Nielsen-flat (Proposition 7.644, Theorem 7.646, and Section 7.10.1).

Radial (Higgs-like) scalars. Across all quadrants, radial/moduli excitations appear as spin-0 gapped modes. They remain analytic on $\mathfrak{P}$; explicit breaking in QIV does not change their spin, only their masses/mixings within the same scalar block. These provide stable anchors for plateau diagnostics and for the GMOR fits that isolate the pseudo-NG sector in QIV (eq. (7.747) and Corollary 7.631).

QIII reconvergence and the absence of NG/pNG in the strict IR. QIII exhibits finite-scale law deficits that *heal* under IR dilation according to the power law in eq. (7.746). On the IR mask $\mathfrak{M}$, the adjoint-covariant source vanishes and no NG/pNG sector persists asymptotically; multiplets reconverge to the QI census (massless spin-2, massless spin-1 for unbroken G_{IR}, and gapped scalar radials). Any finite-ℓ pNG-like transient must decay in accordance with eq. (7.746) (Sections 7.8.9 and 7.10.1).

Spin assignment and analytic transport. Explicit law asymmetry (notably in QIV) does not alter spin assignments: spins are inherited from the symmetric limit, while masses and intra-block mixings shift analytically with spurions and scale tags. Pole residues used for

species identification are gauge-parameter independent on $\mathcal{H}_{\text{phys}}$ (Nielsen-flat), ensuring regulator-robust spectroscopy on $\mathfrak{P}$ (Proposition 7.644 and Theorem 7.646).

Ghosts and auxiliaries. BRST cohomology projects out Faddeev–Popov ghosts and auxiliary fields; they do not appear as particle-like excitations in the physical spectrum. All "particle-like" statements above are therefore to be read within $\mathcal{H}_{\text{phys}}$ (Proposition 6.276).

Operational summary (species-by-quadrant). QI: massless spin-2; massless spin-1 for unbroken gauge; gapped scalar radials; no NG/pNG. QII: add exact NG scalars (global broken) or, if gauged, massive spin-1 vectors; spacetime NGs as phonons/rotations when applicable. QIII: strict-IR reconvergence to QI (no NG/pNG), with finite-ℓ transients suppressed per eq. (7.746). QIV: pseudo-NG scalars (GMOR) and pseudo-phonons (if spacetime explicitly broken); massive spin-1 for gauged broken directions; radial scalars remain gapped/analytic; spins unchanged (see table 7.5).

Terminology (particles vs. quasiparticles). In this chapter, "*particle-like*" denotes any isolated, positive-residue pole on $\mathcal{H}_{\text{phys}}$ certified on $\mathfrak{P}$ (and $\mathfrak{M}$ when invoked), irrespective of UV provenance. It therefore includes both *particles* (kymon sector: massless/massive spin-1 gauge quanta; gapped spin-0 radials; any additional symmetric-law quanta) and *quasiparticles* (apeiron sector: spacetime NG phonons/rotations and their QIV pseudo-phonons; internal NG/pNG scalars in QII/QIV). The species inventory and quadrant placement are summarized in table 7.5; the consolidated logic linking visibilities, Ward sources, and spectra is in Section 7.10.

Polarizations, Helicities, and Mode Catalog

All entries are certified on $\mathcal{H}_{\text{phys}}$ and read on plateaus $\mathfrak{P}$ (and masks $\mathfrak{M}$ when invoked), with gauge/Nielsen safety as in Propositions 6.276 and 7.644 and Theorem 7.646. Quadrant occurrence matches Definition 7.685 and table 7.5, and QIII/QIV scaling follows eqs. (7.746) and (7.747).

Helicity conventions (3+1 reference; general d given below). Massless spin-1: two transverse helicities $\lambda = \pm 1$; *no longitudinal mode*. Massive spin-1: three physical polarizations $\{\lambda = +1, 0, -1\}$. Massless spin-2: two helicities $\lambda = \pm 2$ (transverse traceless). Scalars are spin-0 (no polarization). Spacetime NGs (phonons/rotations) are labeled by longitudinal/transverse content under the spatial little group; QIV pseudo-phonons inherit the same polarization pattern and acquire a gap set by eq. (7.747).

Projectors (for diagnostics, no new labels). Transverse projector for vectors: $P_T^{ij}(\hat{\mathbf{k}}) = \delta^{ij} - \hat{k}^i \hat{k}^j$ (massless spin-1 lives in $\operatorname{Im} P_T$). Transverse-traceless projector for tensors: $\Pi_{\text{TT}}^{ij,kl}(\hat{\mathbf{k}}) = \frac{1}{2}(P_T^{ik}P_T^{jl} + P_T^{il}P_T^{jk}) - \frac{1}{d-1}P_T^{ij}P_T^{kl}$ (massless spin-2 lives in $\operatorname{Im} \Pi_{\text{TT}}$). These are the canonical filters for channel-resolved fits used alongside Definition 7.35 and Section 7.10.1.

Table 7.6: Mode–helicity catalog (species, physical polarizations, quadrant occurrence). "Dispersion" shows plateau prototypes; v_* are channel-dependent cone speeds fixed by the chronogeometric operator used in the holosymmetric lift.

Species	**Spin**	**Pol./Helicity**	**Dispersion (plateau prototype)**	**Quadrants (cf. table 7.5)**
Massless vector (unbroken)	1	$\lambda = \pm 1$	$\omega^2 \simeq v_1^2\,\mathbf{k}^2$	QI, QII, QIII (IR), QIV (unbroken)
Massive vector (gauged broken)	1	$\{+1, 0, -1\}$	$\omega^2 \simeq v_1^2\,\mathbf{k}^2 + m_V^2$	QII, QIV
Massless tensor (TT)	2	$\lambda = \pm 2$	$\omega^2 \simeq v_2^2\,\mathbf{k}^2$	QI, QII, QIII (IR), QIV (unbroken spacetime)
Radial/Higgs-like scalar	0	—	$\omega^2 \simeq v_0^2\,\mathbf{k}^2 + m_H^2$	All quadrants
Internal NG scalar	0	—	$\omega^2 \simeq v_\pi^2\,\mathbf{k}^2$	QII only (global broken)
Internal pNG scalar	0	—	$\omega^2 \simeq v_\pi^2\,\mathbf{k}^2 + m_\pi^2$, $m_\pi^2 \sim \mathcal{O}(\epsilon)$	QIV (GMOR via eq. (7.747))
Spacetime NG (phonon/rotation)	0	L/T	$\omega^2 \simeq c_L^2 k_L^2 + c_T^2 k_T^2$	QII (if spacetime SSB)
Spacetime pseudo-phonon	0	L/T	$\omega^2 \simeq c_{L,T}^2 k^2 + m_{\rm ph}^2$, $m_{\rm ph}^2 \sim \mathcal{O}(\epsilon)$	QIV (explicit spacetime breaking)

Note. "Low-energy/Light sector" refers to physically propagating modes *within the certified IR window* $K_{\rm IR}^{(\ell)}$ (with UV content controlled by $P_x(\lambda_{\rm UV})$) after BRST projection onto $\mathcal{H}_{\rm phys}$. It includes massless carriers for unbroken gauge factors (not necessarily photons), NG/pNG modes (QIV obeys the GMOR slope), and radials below the window; excludes heavy states/continua and ghosts.

Kinematics (plateau prototypes). For massless carriers, group velocities $v_{1,2}$ coincide with the causal cone speeds set by the chronogeometric sector of the holosymmetric lift; for scalars v_0 is channel dependent. In QIV, pNG and pseudo-phonon gaps satisfy the source–mass bridge of eq. (7.747); in QIII, any finite-ℓ pNG-like transient must decay with ℓ according to eq. (7.746).

Degrees of Freedom (DoF) Counts and Conventions

Let $D = d + 1$ be spacetime dimension. Physical DoF per species on $\mathcal{H}_{\text{phys}}$:

$$\begin{aligned} \text{scalar:}\ & 1, \\ \text{massless spin-1:}\ & D-2 = d-1, \\ \text{massive spin-1:}\ & D-1 = d, \\ \text{massless spin-2:}\ & \tfrac{1}{2}D(D-3) = \tfrac{1}{2}(d+1)(d-2). \end{aligned}$$

These counts underwrite the rows of table 7.5 and guide residue normalizations in spectral fits (Nielsen-flat per Proposition 7.644).

Channel ↔ Mode Map

The triadic split used operationally (correlation/vacuum/spectral) decomposes readouts without altering polarization content. A convenient mnemonic (no new labels):

- *Spectral channel:* isolates pole locations and residues for all particle-like species (vectors/tensors/scalars), used for GMOR fits in QIV via eq. (7.747).
- *Vacuum channel:* carries order-parameter plateaus (e.g., QII manifold structure) and the orthogonality trend $\Lambda_{\text{state}}(V) \to 0$ relevant to Definition 7.685.
- *Correlation channel:* sensitive to mixing angles within a fixed spin block; in QIV these vary analytically with explicit spurions, leaving spin assignments unchanged (cf. notes under table 7.5).

Residue Normalization and GMOR Readout (QIV)

For the pNG band, define decay constants f_{π_b} by the usual one-pole saturation of the improved current two-point function on $\mathcal{H}_{\text{phys}}$ (no new label). On plateaus, the source–mass relation in eq. (7.747) fixes the leading slope of $m^2_{\pi_b}$ in the explicit couplings $\{\epsilon_k\}$; radial modes remain gapped/analytic and serve as internal consistency checks for fits.

IR Healing and Multiplet Reconvergence (QIII)

Under IR dilation within $\mathfrak{M}$, eq. (7.746) enforces the decay of the law-channel deficit and rules out a persistent NG/pNG sector. Spectrally, spin blocks reconverge to the QI census with improving positivity/hyperbolicity margins (as summarized in table 7.5); any finite-ℓ anomalies must track the power law in eq. (7.746).

Chapter 8

Parasymmetric Witnesses and Diagnostics

8.1 Sector-Conditioned Witnesses and Quadrant Rules

Chs. 6–7 establish a symmetry-centric, gauge-safe classification based on two predicates—the *law* and the *state* visibilities—read on a certified plateau and mapped to the four quadrants by the decision rule of Definition 7.685. The present section extends that same machinery to further physical sectors without altering the framework.

What stays fixed. The *grid does not change*: we retain the two fundamental visibilities $(\Lambda_{\text{law}}, \Lambda_{\text{state}})$ of Definition 7.35, evaluated after projection to the BRST-physical sector, and we use the same quadrant map. No new cells are introduced.

What changes. Only the *witnesses* and, where needed, the *arena* change. For each admissible sector $\mathscr{S}$ we instantiate the sector-conditioned pair

$$\left(\Lambda_{\text{law}}^{[\mathscr{S}]},\ \Lambda_{\text{state}}^{[\mathscr{S}]}\right)$$

as in Definition 8.9, using the appropriate Ward data/correlators on the relevant arena $\mathcal{A} \in \{\mathfrak{P},\ \mathfrak{P}_\partial\}$ with an IR window $K_{\text{IR}}^{(\ell)}$ (isotropic) or $K_{\text{IR}}^{(\vec{\ell})}$ (anisotropic). Subscripts remain reserved for Λ_{law} and Λ_{state}; sector tags appear only as superscripts.

Scope of this section. We provide witnesses and quadrant assignments for: fermions, SUSY (global/local), p-forms in 3+1D (via BRST-compatible duality), higher-spin $s{>}2$, hydrodynamics, boundary soft/edge modes, anyons/CS (for a chosen symmetry X), fractons/subsystem-symmetric phases, and higher-form/non-invertible symmetries. Each sector is labeled

$$\text{Quad}^{[\mathscr{S}]} = \text{Quad}\left(\Lambda_{\text{law}}^{[\mathscr{S}]}, \Lambda_{\text{state}}^{[\mathscr{S}]}\right).$$

Orthogonality and consistency. Topological order and related modular data are orthogonal to the law/state predicates and are tracked separately when relevant; they do not modify quadrant assignment. BRST/Nielsen safety ensures gauge-parameter independence of all witnesses, and large-window limits distinguish QIII (healing) from QIV (persistence) in the sense of eq. (7.746). Non-equilibrium carriers of the law (Floquet/Lindblad) fit by swapping the dynamical map while keeping the same predicates and decision rule.

How to read the remainder. For each sector block below we (i) specify the witnesses $\big(\Lambda_{\mathrm{law}}^{[\mathscr{S}]}, \Lambda_{\mathrm{state}}^{[\mathscr{S}]}\big)$ on the appropriate arena and window, and (ii) state the resulting QI–QIV assignment, with QIII/QIV decided by the large-window limit in that arena. This yields a uniform, symmetry-agnostic classification across all additional sectors.

Theorem 8.1 (Law–state exhaustion)**.** *On the certified plateau $\mathfrak{P}$, after projection to $\mathcal{H}_{\mathrm{phys}}$, the (binary) visibility predicates associated with the holosymmetric visibilities*

$$\chi_{\mathrm{law}} := \mathbf{1}\{\,|\Lambda_{\mathrm{law}}| \geq 1 - \varepsilon_{\mathrm{tol}}\,\}\,, \qquad \chi_{\mathrm{state}} := \mathbf{1}\{\,|\Lambda_{\mathrm{state}}| \geq 1 - \varepsilon_{\mathrm{tol}}\,\}$$

(Definition 7.35, with the tolerance from the plateau certification Section 6.12.4), together with the decision rule of Definition 7.685, partition the admissible configurations into exactly four disjoint classes **QI–QIV***. The partition is (i)* exclusive *(no configuration belongs to two classes) and (ii)* exhaustive up to threshold sets *on $\mathfrak{P}$. Moreover, the assignment is gauge–parameter independent (BRST/Nielsen safe).*

Proof. **Set–up and notation.** All quantities are evaluated on $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ with BRST–closed scale tags commuting with the channel projectors (Propositions 6.264 and 6.276). Plateau uniformity and derivative bounds are as in Section 6.12.4 and Remark 6.254; in particular, "≈ 1'' means $\big|\,|\cdot| - 1\big| \leq \varepsilon_{\mathrm{tol}}$ uniformly on $\mathfrak{P}$. When $\chi_{\mathrm{law}} = 0$, we refine by the law–deficit dichotomy on $\mathfrak{P}$:

$$\begin{aligned} &\text{either} \quad 1 - |\Lambda_{\mathrm{law}}(\ell)| = \mathcal{O}\big(\ell^{\,\Delta-(d+z)}\big) \quad (\text{QIII mask } \mathfrak{M}), \\ &\text{or} \quad \exists\, \delta_\star \in (0,1) : \ |\Lambda_{\mathrm{law}}| \leq 1 - \delta_\star \text{ uniformly on } \mathfrak{P}, \end{aligned} \tag{$\star$}$$

as established by Proposition 6.277 and equivalently by the Ward/spectral criteria (Theorem 7.687). The state channel is decided by the thermodynamic limit (Theorem 6.258): either $\Lambda_{\mathrm{state}}(V) \to 0$ (SSB) or $|\Lambda_{\mathrm{state}}| \approx 1$ (unique/symmetric vacuum).

(A) Exclusivity. We show the four labels are pairwise disjoint.

QI vs QII. Both require $\chi_{\mathrm{law}} = 1$, but QI imposes $|\Lambda_{\mathrm{state}}| \approx 1$ while QII imposes $\Lambda_{\mathrm{state}}(V) \to 0$. These are mutually exclusive by the thermodynamic orthogonality of distinct vacua in SSB (Theorem 6.261) together with plateau uniformity.

QIII vs QIV. Both have $\chi_{\mathrm{law}} = 0$, but the law–deficit alternatives in ($\star$) are exclusive: if $1 - |\Lambda_{\mathrm{law}}| \to 0$ with a positive power, it cannot be bounded below by any $\delta_\star > 0$, and conversely a uniform gap precludes any power–law decay to zero. This is exactly the "healing'' vs "persistent'' dichotomy proven via the scale–tagged Ward identity (Proposition 6.296) and the overlay bounds (Proposition 6.277).

Cross pairs. QI/QII have $\chi_{\mathrm{law}} = 1$ while QIII/QIV have $\chi_{\mathrm{law}} = 0$, hence disjoint.

(B) Exhaustiveness up to threshold sets. Let a configuration on $\mathfrak{P}$ be given.

If $\chi_{\mathrm{law}} = 1$, then either $\Lambda_{\mathrm{state}}(V) \to 0$ (SSB) or not. In the former case, Theorem 6.261 tags QII; in the latter, $|\Lambda_{\mathrm{state}}| \approx 1$ and we tag QI. If $\chi_{\mathrm{law}} = 0$, apply ($\star$): either the deficit heals with ℓ (QIII, inside $\mathfrak{s}$) or a plateau–uniform gap remains (QIV). Thus every configuration is assigned.

Ambiguities can arise only on *threshold sets*, namely: (i) the $\varepsilon_{\mathrm{tol}}$–boundaries where $\big||\Lambda_{\mathrm{law}}| - 1\big| = \varepsilon_{\mathrm{tol}}$ or $\big||\Lambda_{\mathrm{state}}| - 1\big| = \varepsilon_{\mathrm{tol}}$; (ii) the borderline where the law deficit neither

admits a uniform power–law decay nor a uniform positive gap (e.g. precisely marginal cases tuned to zero amplitude). These sets are of codimension ≥ 1 in control space and of zero measure under generic parameterizations; by definition of "certified plateau'' they are either excluded or flagged (cf. the plateau certification protocol of Section 6.12.4 and the QIII mask construction).

(C) Gauge–parameter independence (Nielsen safety). By Propositions 6.264 and 6.299, $\mathcal{O} \mapsto \mathcal{O}_\xi$ is ξ–independent on $\mathfrak{P}$ for $\mathcal{O} \in \{\, \Lambda_{\mathrm{law}}, \Lambda_{\mathrm{state}}, D_\lambda \mathcal{O}, D_\ell \mathcal{O} \,\}$, hence both the predicates $\chi_{\mathrm{law}}, \chi_{\mathrm{state}}$ and the alternatives in $(\star)$ are gauge–parameter independent. Therefore the quadrant assignment is BRST/Nielsen safe.

Conclusion. The decision rule of Definition 7.685 induces a partition of the admissible configurations on $\mathfrak{P}$ into four classes that are pairwise disjoint (exclusivity) and cover all cases except for codimension–one threshold sets (exhaustiveness up to thresholds). The entire construction is cohomologically well–defined and gauge–parameter independent. □

Remark 8.2 (Reader intuition and operational recipe for law–state exhaustion)**.** *Big picture.* On a certified plateau $\mathfrak{P}$ (scale–stable readouts), the BRST–safe visibilities $(\Lambda_{\mathrm{law}}, \Lambda_{\mathrm{state}})$—evaluated in the physical sector $\mathcal{H}_{\mathrm{phys}}$—are sufficient to classify any admissible configuration into exactly one of the four quadrants (QI–QIV) via the decision rule of Definition 7.685. The classification is *exclusive* (no overlap between labels) and *exhaustive up to thresholds* (the only undecidable cases are measure–zero boundaries where "≈ 1'' is saturated or the knife–edge between QIII healing and QIV persistence). Because Nielsen/cohomological identities hold on $\mathfrak{P}$, the assignment is *gauge–parameter independent*. Equivalently, the overlays, the scale–tagged Ward statements, and the light–sector spectrum cohere (Theorem 7.687).

Operational recipe. Measure the two visibilities on $\mathfrak{P}$ after BRST projection.

- If $|\Lambda_{\mathrm{law}}| \approx 1$ on $\mathfrak{P}$, then we are in **QI** or **QII**; the *state* channel decides: if $|\Lambda_{\mathrm{state}}(V)| \to 0$ as $V \to \infty$, assign **QII**; otherwise $|\Lambda_{\mathrm{state}}| \approx 1$ and assign **QI**.
- If $|\Lambda_{\mathrm{law}}| < 1$ on $\mathfrak{P}$, the *law* channel splits **QIII** vs **QIV**: if the deficit heals with coarse–graining,
$$1 - |\Lambda_{\mathrm{law}}(\ell)| = \mathcal{O}\Big(\ell^{\Delta-(d+z)}\Big) \qquad (\ell \downarrow 0),$$
assign **QIII**; if there is a plateau–uniform gap $\exists\, \delta_\star \in (0,1)$ with $|\Lambda_{\mathrm{law}}| \leq 1 - \delta_\star$, assign **QIV**.

Why there are only four. Under the Chapter 7/8 standing assumptions (certified plateau, evaluation in $\mathcal{H}_{\mathrm{phys}}$, BRST/Nielsen invariance), the only binary predicates are:

$$\text{Law: } |\Lambda_{\mathrm{law}}| \approx 1 \text{ vs deficit}, \qquad \text{State: } |\Lambda_{\mathrm{state}}| \approx 1 \text{ vs thermodynamic orthogonality}.$$

Their Cartesian product yields exactly the four classes QI–QIV; no fifth "type'' exists. Variants such as discrete SSB, topological order, anisotropic scaling, or competing irrelevancies are already subsumed once one uses BRST–closed, gauge–invariant probes and reads the visibilities on $\mathfrak{P}$; any non-generic in–between situations are precisely the *threshold sets* flagged by the plateau certification and do not constitute new quadrants.

Remark 8.3 (On apparent "simultaneity'' outside the formal framework)**. Thesis.** Even "in the wild,'' for a fixed symmetry G, at a fixed observational scale/window, and under a fixed measurement protocol, a single physical system cannot have a *law* that is both symmetric and explicitly asymmetric, nor a *state* that is both symmetric and symmetry–broken, at the same time. Apparent counterexamples arise from mixing contexts—scale, space, ensemble, or sector—and dissolve once those are fixed [91, 230, 101, 102, 82].

Ensemble vs single sample. Below T_c, the *ensemble* density matrix of an Ising ferromagnet can remain symmetric with $\langle M\rangle = 0$, while any *single crystal* in zero field selects $+m$ or $-m$. Symmetric in the ensemble, broken in each pure phase—never both for the same object under the same notion of "state.'' In algebraic language: distinct broken phases correspond to disjoint GNS representations (superselection sectors) and become orthogonal in the thermodynamic limit [230, 91].

Finite size and cat states. With exact symmetry and finite volume, the unique ground state may be a Schrödinger–cat superposition of symmetry–broken sectors. Infinitesimal symmetry–breaking fields or the order of limits (take $V\to\infty$ before $h\to 0$) select a definite sector. Thus the state is formally symmetric at finite V but operationally broken once limits/perturbations are specified [91, 230].

Domain/phase coexistence. Near first–order transitions or with quenched disorder, different spatial regions may choose different orientations. Locally broken, globally (after coarse–graining) possibly symmetric—this is spatial mixing, not simultaneity for a single homogeneous state [101].

Scale separation (UV vs IR). Microscopic anisotropies can be visible at short scales and wash out under coarse–graining: the *law* appears asymmetric in the UV but flows to an IR–symmetric fixed point (emergent symmetry). This is scale dependence, not "both at once'' at a fixed scale [102, 82].

Partial or sectoral symmetries. A symmetry group may be only partially broken (e.g. nematic order breaks rotations but not translations) or broken in one sector of the full theory but exact in another (parity violation in weak interactions vs parity conservation in QCD). For any given generator, the verdict is binary once the sector is fixed [91, 101].

Anomalies. Classically symmetric, quantum mechanically not (e.g. $U(1)_A$ in QCD): the correct quantum *law* is asymmetric; calling it "both'' mixes levels of description. This is the textbook chiral–anomaly resolution [82, 91].

Non–equilibrium / time–crystalline behavior. A Hamiltonian (or Floquet operator) can be symmetric while the *state* breaks time–translation symmetry (e.g. period–doubling). That is the canonical "law symmetric, state asymmetric'' case—not simultaneity of symmetric and explicitly asymmetric *law*. See, e.g., discrete time–crystal constructions in periodically driven many–body systems [396, 448, 449].

Topological order. In topologically ordered phases, local order parameters may vanish although the system exhibits long–range entanglement and topological degeneracy. The *state* can be symmetric under G while displaying nonlocal order; this is neither "both symmetric and broken'' nor an extra quadrant—it is a different organizing principle for order [102].

Rule of thumb. To avoid false "both'' verdicts, fix: (i) the symmetry G and generator, (ii) the observation scale/window, (iii) whether we mean a pure sector or an ensemble/disorder average, (iv) the order of thermodynamic/time limits, and (v) gauge–invariant/BRST–closed probes where applicable. Once these are fixed, the law/state verdict is definite and mutually exclusive—matching the formal classification proven in Chapters 7–8 [117, 91, 230].

Definition 8.4 (Witness = sector–conditioned visibility recipe)**.** Fix an admissible sector $\mathscr{S}$ (bulk or boundary, possibly anisotropic) realised on a certified plateau $\mathfrak{P}$ (or $\mathfrak{P}_\partial$), with BRST–closed scale tags $P_x(\lambda_{\mathrm{UV}})$ and K_{IR} and channel projectors $\{\Pi_\alpha\}$ (ghost number 0). A *witness* for $\mathscr{S}$ is a map that assigns to the tagged data a number in $[0,1]$:

$$\mathbf{W}^{[\mathscr{S}]}:\ \Big(\text{tagged operators, currents, states on } \mathfrak{P}\Big)\ \longrightarrow\ [0,1],$$

constructed from BRST–closed ingredients and evaluated in the physical sector $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$, and satisfying the axioms below. We call the output $\Lambda^{[\mathscr{S}]} := \mathbf{W}^{[\mathscr{S}]}$ a *visibility*.

Definition 8.5 (Law vs. State witnesses)**.** A *law witness* $\Lambda^{[\mathscr{S}]}_{\mathrm{law}}$ diagnoses explicit symmetry breaking in the *dynamics* (e.g. normalized Ward sources, adjoint divergences, control–unitary overlaps). A *state witness* $\Lambda^{[\mathscr{S}]}_{\mathrm{state}}$ diagnoses spontaneous breaking in the *state* (e.g. symmetry–related overlaps/cluster ratios, certified order parameters, pole diagnostics). The *witness pair* for $\mathscr{S}$ is $\big(\Lambda^{[\mathscr{S}]}_{\mathrm{law}},\ \Lambda^{[\mathscr{S}]}_{\mathrm{state}}\big)$.

Definition 8.6 (Certified witness pair and predicates)**.** Fix plateau tolerances $\varepsilon_{\mathrm{law}}, \varepsilon_{\mathrm{state}} \in (0,1)$ (cf. Section 6.12.4). Define binary predicates

$$\chi^{[\mathscr{S}]}_{\mathrm{law}} := \mathbf{1}\Big[\,|\Lambda^{[\mathscr{S}]}_{\mathrm{law}}| \geq 1-\varepsilon_{\mathrm{law}}\,\Big], \qquad \chi^{[\mathscr{S}]}_{\mathrm{state}} := \mathbf{1}\Big[\,|\Lambda^{[\mathscr{S}]}_{\mathrm{state}}| \geq 1-\varepsilon_{\mathrm{state}}\,\Big],$$

and the *law deficit* $d^{[\mathscr{S}]}_{\mathrm{law}} := 1 - |\Lambda^{[\mathscr{S}]}_{\mathrm{law}}|$. The quadrant $\mathrm{Quad}^{[\mathscr{S}]}$ is determined by the canonical decision rule (Definition 7.685), with QIII vs. QIV fixed by the large–window test in the sector arena.

Axioms for a witness (uniform on the certified plateau). Let $\mathbf{W}^{[\mathscr{S}]}$ be built from BRST–closed operators (ghost number 0), with BRST–safe norms. Then on the certified band $B \subset (\lambda_{\mathrm{UV}}, \ell)$ (or its boundary/anisotropic analogue):

(W1) BRST/Nielsen invariance. $\partial_\xi \mathbf{W}^{[\mathscr{S}]} = 0$ and $\mathbf{W}^{[\mathscr{S}]}[\Psi] = \mathbf{W}^{[\mathscr{S}]}[\Psi + \{\hat{Q}_{\mathrm{BRST}}, B\}]$ (Propositions 6.264 and 6.276).

(W2) Cone preservation and functoriality. Holosymmetric lifts, p–form dualities, and triadic projections commute with evaluation and do not alter cones nor mix channels (Propositions 6.257 and 8.16 and Corollary 6.259).

(W3) Normalization and boundedness. $0 \leq \mathbf{W}^{[\mathscr{S}]} \leq 1$. On a sectoral *reference* symmetric instance, $\mathbf{W}^{[\mathscr{S}]} = 1$; explicit breakings reduce it monotonically (up to $o_B(1)$ renormalizations; Remark 6.254).

(W4) Plateau stability (tolerance semantics). $D_\lambda \mathbf{W}^{[\mathscr{S}]}$ and $D_\ell \mathbf{W}^{[\mathscr{S}]}$ are uniformly bounded on B (Proposition 6.298); equalities are interpreted as "≈ 1'' within $(\varepsilon_{\mathrm{law}}, \varepsilon_{\mathrm{state}})$.

(W5) Large–window criterion (QIII/QIV). There exists a sector dilation parameter r (bulk ℓ, boundary L, anisotropic $\min(\vec{\ell})$, or non–eq. coarse–graining) such that:

$$\begin{aligned} \text{QIII} &\iff \exists\, \beta > 0,\, C < \infty :\ d_{\text{law}}^{[\mathscr{S}]}(r) \le C\, r^{-\beta}; \\ \text{QIV} &\iff \exists\, \delta_\star \in (\varepsilon_{\text{law}}, 1) :\ d_{\text{law}}^{[\mathscr{S}]}(r) \ge \delta_\star. \end{aligned}$$

(See sector blocks and Theorem 7.687.)

Remark 8.7 (Relationship to earlier notions)**.** The *visibility* $\Lambda_\bullet$ used throughout Chapters 7–8 is the *value* returned by a witness. Sector blocks (fermions, SUSY, hydrodynamics, boundary, subsystem/fracton, higher–spin, p–forms, non–equilibrium) instantiate $\mathbf{W}^{[\mathscr{S}]}$ concretely (e.g. normalized Ward–source quotients, overlap diagnostics), all conforming to **(W1)**–**(W5)**.

Remark 8.8 (Examples (pointers))**.** Hydrodynamic witnesses: Definition 8.21; boundary: Definition 8.31; subsystem/fracton: Definition 8.36; fermionic/SUSY/ p–form/HS blocks: Definitions 8.10, 8.12, 8.15 and 8.18; non–equilibrium (Floquet/Lindblad): Definition 8.43. Their quadrant rules follow from Propositions 8.23, 8.32 and 8.38 and the universal cartography theorems.

Definition 8.9 (Sector-conditioned visibilities and arenas)**.** Let $\mathscr{S}$ be any admissible sector, realised on an arena $\mathcal{A} \in \{\mathfrak{P},\ \mathfrak{P}_\partial\}$. With the UV projector $P_x(\lambda_{\text{UV}})$ and an (optionally anisotropic) IR window $K_{\text{IR}}^{(\vec{\ell})}$ on $\mathcal{A}$, define the BRST-safe pair

$$\left(\Lambda_{\text{law}}^{[\mathscr{S}]},\ \Lambda_{\text{state}}^{[\mathscr{S}]}\right)$$

by instantiating the witnesses of Definition 7.35 using sector-specific Ward sources and correlators (after projection to $\mathcal{H}_{\text{phys}}$ and band-limiting by $P_x(\lambda_{\text{UV}}) \circ K_{\text{IR}}^{(\vec{\ell})}$). The quadrant label is

$$\text{Quad}^{[\mathscr{S}]} := \text{Quad}\left(\Lambda_{\text{law}}^{[\mathscr{S}]}, \Lambda_{\text{state}}^{[\mathscr{S}]}\right),$$

with Quad given in Definition 7.685. *Pointers:* for concrete witnesses see Definitions 8.21, 8.31, 8.33 and 8.36.

8.1.1 Fermionic Witnesses

Definition 8.10 (Fermionic witnesses, tags, and normalization)**.** Work on the certified bulk plateau $\mathfrak{P}$ and in the physical sector $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$. Let $S^R(\omega, \mathbf{k})$ be the *retarded* fermion propagator with IR/UV tags $K_{\text{IR}}^{(\ell)}$, $P_x(\lambda_{\text{UV}})$ acting on the fermionic block, and let $\mathcal{D}^{(\ell)}$ denote the tagged wave operator with tagged mass matrix $M^{(\ell)}$. Define the *fermionic law visibility* by the (dimensionless) ratio

$$\Lambda_{\text{law}}^{[\text{ferm}]} := 1 - \frac{\left\|\text{Src}^{(\text{ferm})}\right\|_{\mathfrak{P}}}{\left\|\text{Ref}^{(\text{ferm})}\right\|_{\mathfrak{P}}}, \qquad \text{Src}^{(\text{ferm})} := \sum_k \epsilon_k(\ell) \left\langle \mathcal{S}_{k,a}^{(\lambda_{\text{UV}})} \right\rangle_{\Psi}\Big|_{\text{ferm. block}},$$

where $\|\cdot\|_{\mathfrak{P}}$ is any plateau–uniform, BRST–safe operator norm or spectral (Källén–Lehmann) seminorm compatible with the tags, and the reference $\text{Ref}^{(\text{ferm})}$ is a fixed nonzero, conserved

benchmark (e.g. the tagged, improved Noether density or the S^R residue sum on $\mathfrak{P}$), chosen once and for all so that $\Lambda^{[\mathrm{ferm}]}_{\mathrm{law}} \in [0,1]$. Define the *fermionic state visibility* by the presence/absence of spontaneous breaking in the fermionic channel (e.g. chiral or flavor order):

$$\Lambda^{[\mathrm{ferm}]}_{\mathrm{state}} := \begin{cases} 1, & \text{symmetric state on } \mathfrak{P} \ \text{ (no fermionic order parameter)}, \\ 0, & \text{SSB in a fermionic channel on } \mathfrak{P} \ \text{ (nonzero order parameter / NG partner)}. \end{cases}$$

All insertions are evaluated as $P_x(\lambda_{\mathrm{UV}})\, K^{(\ell)}_{\mathrm{IR}}\,(\cdot)\, K^{(\ell)}_{\mathrm{IR}}\, P_x(\lambda_{\mathrm{UV}})$ and are BRST–closed, ghost number 0.

Intuition. The law visibility measures explicit symmetry breaking *restricted to the fermion sector*, via the scale–tagged Ward sources on the fermionic block; the normalization makes it dimensionless and plateau–uniform. The state visibility detects whether the *state* itself breaks the same fermionic symmetry (e.g. chiral SSB), and by BRST/Nielsen invariance it is gauge–parameter independent (Propositions 6.276 and 7.644).

Proposition 8.11 (Fermionic quadrant assignment)**.** *On the certified plateau* $\mathfrak{P}$*,*

QI : $\Lambda^{[\mathrm{ferm}]}_{\mathrm{law}} = 1, \quad \Lambda^{[\mathrm{ferm}]}_{\mathrm{state}} = 1;$

QII : $\Lambda^{[\mathrm{ferm}]}_{\mathrm{law}} = 1, \quad \Lambda^{[\mathrm{ferm}]}_{\mathrm{state}} = 0, \quad$ *(law conserved; fermionic SSB, NG partners present)*;

QIII : $\Lambda^{[\mathrm{ferm}]}_{\mathrm{law}}(\ell) \xrightarrow[\ell\downarrow 0]{} 1, \quad \Lambda^{[\mathrm{ferm}]}_{\mathrm{state}} = 1, \quad$ *(IR healing of irrelevant explicit deformations)*;

QIV : $\Lambda^{[\mathrm{ferm}]}_{\mathrm{law}} \le 1 - \delta_\star$ *uniformly on* $\mathfrak{P}$, $\quad \Lambda^{[\mathrm{ferm}]}_{\mathrm{state}} = 1,$
(persistent explicit breaking; analytic mass/mixing shifts).

Proof (fermionic specialization of Chapter 7/8 results). Standing hypotheses. Work in $\mathcal{H}_{\mathrm{phys}}$ with BRST–closed tags and channel projectors, so Nielsen identities hold (Propositions 6.264 and 6.276). Let $\bar{J}^{(\lambda_{\mathrm{UV}},\ell)\mu}_{a,\mathrm{imp}}$ be the improved current restricted to the fermionic block; the scale–tagged fermionic Ward statement is the fermionic restriction of Proposition 6.296:

$$\nabla^{(\mathrm{ad})}_\mu \bar{J}^{(\lambda_{\mathrm{UV}},\ell)\mu}_{a,\mathrm{imp}}\Big|_{\mathrm{ferm}} = \sum_k \epsilon_k(\ell) \left\langle \mathcal{S}^{(\lambda_{\mathrm{UV}})}_{k,a} \right\rangle_{\Psi}\Big|_{\mathrm{ferm}}. \tag{8.1}$$

QI. Exact symmetry on the law side implies the r.h.s. of (8.1) vanishes; with the chosen normalization, $\Lambda^{[\mathrm{ferm}]}_{\mathrm{law}} = 1$. A unique symmetric state on $\mathfrak{P}$ implies $\Lambda^{[\mathrm{ferm}]}_{\mathrm{state}} = 1$ (no fermionic order parameter; cluster decomposition) [91, 230].

QII. Law symmetry persists $\Rightarrow$ same law verdict as QI. If the state spontaneously breaks a fermionic symmetry (e.g. chiral), the thermodynamic orthogonality of distinct vacua and the Goldstone theorem yield $\Lambda^{[\mathrm{ferm}]}_{\mathrm{state}} = 0$ with exact NG partners in the spectral function [94, 91].

QIII. Let the fermionic explicit breakings be RG–irrelevant ($\Delta_k > d + z$). Then $\epsilon_k(\ell) = \mathcal{O}(\ell^{\Delta_k - (d+z)})$ and the r.h.s. of (8.1) decays with the same power. Hence $\|\mathrm{Src}^{(\mathrm{ferm})}\|_{\mathfrak{P}}$ is $o(1)$ compared to the fixed reference, so $\Lambda^{[\mathrm{ferm}]}_{\mathrm{law}}(\ell) \to 1$; the state remains symmetric, giving $\Lambda^{[\mathrm{ferm}]}_{\mathrm{state}} = 1$ [102, 82].

QIV. With a relevant/marginal explicit fermionic breaking surviving on $\mathfrak{P}$, $\|\mathrm{Src}^{(\mathrm{ferm})}\|_{\mathfrak{P}}$ is bounded below, so $\Lambda_{\mathrm{law}}^{[\mathrm{ferm}]} \le 1-\delta_\star$. The state is uniquely prepared (no SSB in the fermionic channel on $\mathfrak{P}$), hence $\Lambda_{\mathrm{state}}^{[\mathrm{ferm}]} = 1$. By gauge–safe analyticity of tagged two–point functions on $\mathcal{H}_{\mathrm{phys}}$ (Theorem 7.646), mass/mixing shifts are analytic in the explicit couplings and do not change spin 1/2 assignments.

All steps are plateau–uniform and BRST/Nielsen–safe by construction. □

Consistency checks and edge cases (fermionic).

- *Anomalies.* Axial or gauge anomalies (e.g. $U(1)_A$ in QCD) render the quantum law asymmetric; these are excluded by the anomaly–free BRST standing assumption (Remark 6.266) [82, 91].
- *Discrete chiral SSB.* No NG fermions; domain–wall excitations appear instead. The state predicate still drops to 0 in the thermodynamic limit (orthogonality), matching QII.
- *Topological and symmetry–class structure.* Fermionic protected edge modes (e.g. in topological insulators) do not alter the spin content but may constrain allowed mass deformations; use BRST–closed, gauge–invariant probes and the verdict still lands in QI–QIV [102].
- *Majorana vs Dirac and mixings.* The tags and norms are representation–agnostic; explicit deformations move poles/residues *analytically* without changing spin, consistent with Theorem 7.646.

8.1.2 SUSY Witnesses

Definition 8.12 (SUSY witnesses (tolerance form))**.** Work on a certified plateau $\mathfrak{P}$ and in the physical sector $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$. Let S^μ_α be the (band-limited/improved) supercurrent. After projection to $\mathcal{H}_{\mathrm{phys}}$, the tagged SUSY Ward identity reads $\partial_\mu S^\mu_\alpha = \sum_j \epsilon_j(\ell)\, Y_{j,\alpha}$ (modulo BRST-exact improvements). Define the SUSY *law* visibility by

$$\Lambda_{\mathrm{law}}^{[\mathrm{SUSY}]} := 1 - \frac{\left\| \sum_j \epsilon_j \langle Y_j \rangle \right\|_{\mathfrak{P}}}{\| \partial_\mu S^\mu \|_{\mathrm{ref},\mathfrak{P}}} \in [0,1],$$

where the norms are BRST-safe and uniform on $\mathfrak{P}$. Let $\varepsilon_{\mathrm{law}}, \varepsilon_{\mathrm{state}} \in (0,1)$ be the certified tolerances (cf. the plateau criterion). For the *state* witness we adopt plateau–tolerance semantics:

$$\text{Global SUSY:} \quad \begin{cases} \text{unbroken on } \mathfrak{P} \iff |\Lambda_{\mathrm{state}}^{[\mathrm{SUSY}]}| \ge 1-\varepsilon_{\mathrm{state}} \ \ (\text{equivalently } Q_\alpha|\Omega\rangle \approx 0), \\ \text{SSB on } \mathfrak{P} \iff |\Lambda_{\mathrm{state}}^{[\mathrm{SUSY}]}| \le \varepsilon_{\mathrm{state}} \ \ (\text{equivalently } Q_\alpha|\Omega\rangle \not\approx 0). \end{cases}$$

Local SUSY (supergravity):

$$\begin{cases} \text{unbroken on } \mathfrak{P} \iff |\Lambda_{\text{state}}^{[\text{SUSY}]}| \geq 1 - \varepsilon_{\text{state}} \\ \qquad\qquad \text{(vanishing SUSY order parameters,)} \\ \qquad\qquad \text{(background admitting Killing spinor),} \\ \text{broken on } \mathfrak{P} \iff |\Lambda_{\text{state}}^{[\text{SUSY}]}| \leq \varepsilon_{\text{state}} \\ \qquad\qquad \text{(super-Higgs regime).} \end{cases}$$

All insertions are evaluated as $P_x(\lambda_{\text{UV}})\, K_{\text{IR}}^{(\ell)}(\cdot)\, K_{\text{IR}}^{(\ell)}\, P_x(\lambda_{\text{UV}})$ with BRST-closed tags.

Intuition. $\Lambda_{\text{law}}^{[\text{SUSY}]} \approx 1$ (within ε_{law}) iff explicit SUSY-breaking sources are negligible on $\mathfrak{P}$ [52]. The state witness distinguishes spontaneous SUSY breaking: global $\Rightarrow$ goldstino; local $\Rightarrow$ super-Higgs [52].

Lemma 8.13 (Goldstino vs super-Higgs (tolerance form))**.** *Assume on* $\mathfrak{P}$ *that*

$$|\Lambda_{\text{law}}^{[\text{SUSY}]}| \;\geq\; 1 - \varepsilon_{\text{law}}, \qquad |\Lambda_{\text{state}}^{[\text{SUSY}]}| \;\leq\; \varepsilon_{\text{state}}.$$

Then:

Global SUSY.
A massless goldstino appears as the NG mode of SUSY.

Local SUSY.
The goldstino is eaten (super-Higgs) and the gravitino becomes massive; no physical massless goldstino remains.

Proposition 8.14 (SUSY quadrant assignment (tolerance form))**.** *On* $\mathfrak{P}$,

$$\textbf{QI}: \quad \begin{aligned} |\Lambda_{\text{law}}^{[\text{SUSY}]}| &\;\geq\; 1 - \varepsilon_{\text{law}}, \\ |\Lambda_{\text{state}}^{[\text{SUSY}]}| &\;\geq\; 1 - \varepsilon_{\text{state}}, \end{aligned}$$

$$\textbf{QII}: \quad \begin{aligned} |\Lambda_{\text{law}}^{[\text{SUSY}]}| &\;\geq\; 1 - \varepsilon_{\text{law}}, \\ |\Lambda_{\text{state}}^{[\text{SUSY}]}| &\;\leq\; \varepsilon_{\text{state}} \\ &\Big(\textit{global: goldstino; local: super-Higgs to massive } \tfrac{3}{2}\Big), \end{aligned}$$

$$\textbf{QIII}: \quad \begin{aligned} 1 - |\Lambda_{\text{law}}^{[\text{SUSY}]}(\ell)| &\;\leq\; C\, \ell^{\,\Delta-(d+z)} \quad (\ell \downarrow 0), \\ |\Lambda_{\text{state}}^{[\text{SUSY}]}| &\;\geq\; 1 - \varepsilon_{\text{state}}, \end{aligned}$$

$$\textbf{QIV}: \quad \begin{aligned} &\exists\, \delta_\star \in (\varepsilon_{\text{law}}, 1): \; |\Lambda_{\text{law}}^{[\text{SUSY}]}| \;\leq\; 1 - \delta_\star \;\; \textit{uniformly on } \mathfrak{P}, \\ &|\Lambda_{\text{state}}^{[\text{SUSY}]}| \;\geq\; 1 - \varepsilon_{\text{state}}. \end{aligned}$$

Intuition. QII captures spontaneous SUSY breaking (goldstino or super-Higgs) [52, 91]. QIII vs QIV is decided by *decay vs persistent gap* in the SUSY Ward source on $\mathfrak{P}$, in full analogy with the bosonic and fermionic sectors [82, 112]; Nielsen/BRST safety ensures gauge-parameter independence of all witnesses and assignments [82, 91].

8.1.3 p-Forms Witnesses

Definition 8.15 (p-form reduction map in 3+1 (tolerance-consistent))**.** Work on a certified plateau $\mathfrak{P}$ and in the physical sector $\mathcal{H}_{\rm phys} = H^0(\hat{Q}_{\rm BRST})$ with BRST–closed scale tags $P_x(\lambda_{\rm UV})$, $K_{\rm IR}^{(\ell)}$. Let $\mathscr{P}_p$ be an abelian p-form gauge sector in 3+1 with potential A_p and field strength $F_{p+1} = \mathrm{d}A_p$. Define the BRST–compatible Hodge–duality functor $\mathcal{D} : \mathscr{P}_p \to \mathscr{S}_{\rm spin}$ by the on–shell relations

$$F_{p+1} \;=\; *\tilde{F}_{3-p}, \qquad \tilde{F}_{3-p} = \mathrm{d}\tilde{A}_{2-p},$$

so that, in 3+1,

$$\begin{cases} p=1: & \mathscr{P}_1 \xrightarrow{\mathcal{D}} \text{spin-1 sector } (\tilde{A}_1), \\ p=2: & \mathscr{P}_2 \xrightarrow{\mathcal{D}} \text{spin-0 sector } (\tilde{A}_0 \equiv \varphi). \end{cases}$$

After band–limiting and projection to $\mathcal{H}_{\rm phys}$, define the sector–conditioned visibilities *by transport through* $\mathcal{D}$ and interpret "≈ 1'' with certified plateau tolerances (cf. Section 6.12.4):

$$\Lambda_{\rm law}^{[\mathscr{P}_p]} := \Lambda_{\rm law}^{[\mathcal{D}(\mathscr{P}_p)]}, \qquad \Lambda_{\rm state}^{[\mathscr{P}_p]} := \Lambda_{\rm state}^{[\mathcal{D}(\mathscr{P}_p)]}.$$

All statements are understood modulo BRST–exact improvements and with tolerance semantics ($|\Lambda| \geq 1-\varepsilon$ on $\mathfrak{P}$).

Intuition. In 3+1D, a 1–form gauge field is dual to another 1–form, and a 2–form is dual to a scalar [102, 112]. BRST/Hodge duality preserves the Ward data and two–point structures entering the visibilities, so the p–form sector inherits the scalar/vector rows of the spins table (table 7.6—no new quadrant behavior appears [82, 91, 102].

Proposition 8.16 (Quadrant equivalence for p–forms in 3+1)**.** *For $p \in \{1,2\}$ in 3+1D and on a certified plateau $\mathfrak{P}$ (with tolerance semantics),*

$$\mathrm{Quad}^{[\mathscr{P}_p]} \;=\; \mathrm{Quad}^{[\mathcal{D}(\mathscr{P}_p)]}.$$

Proof sketch. Hodge duality $*$ is an isometry on forms (up to signature–dependent signs) and intertwines conserved currents and Ward sources in abelian p–form theories [102, 82]. With BRST–closed tags and evaluation on $\mathcal{H}_{\rm phys}$, $\mathcal{D}$ preserves: (i) the scale–tagged Ward identity (hence the law–channel deficit, within plateau tolerances), and (ii) the light–sector two–point/spectral data (hence the state–channel verdict) [82, 91]. Functoriality and cone preservation ensure no spurious channel mixing. Therefore the visibility predicates—and thus the quadrant label—are invariant under $\mathcal{D}$ on $\mathfrak{P}$. □

Remark 8.17 (Endpoints and couplings)**.** (i) $p = 0$ is already spin–0. For $p = 3$ in 3+1, F_4 is topological and carries no local propagating DOF; when present it maps to a constant/axion–like background, so its quadrant reduces to the scalar entry of $\mathcal{D}(\mathscr{P}_3)$ on $\mathfrak{P}$ (tolerance semantics understood) [112, 102]. (ii) BF/Stückelberg or θ–type couplings induce mixings but, after band–limited diagonalization in the BRST–closed setting, do not affect Proposition 8.16; the visibilities remain Nielsen/BRST–safe and the quadrant assignment is unchanged on $\mathfrak{P}$ [109, 113, 112].

8.1.4 Higher Spins Witnesses

Definition 8.18 (Higher-spin (HS) witnesses (tolerance form))**.** Work on a certified plateau $\mathfrak{P}$ and in the physical sector $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$ with BRST–closed scale tags $P_x(\lambda_{\text{UV}})$, $K_{\text{IR}}^{(\ell)}$ [82, 91]. Let $\mathscr{H}_s$ be a consistent higher–spin ($s > 2$) gauge sector with gauge parameter $\xi_{(s-1)}$ and improved (band–limited) conserved current $J^{(s)}$ [102, 112]. After band/BRST cleaning, the tagged HS Ward identity reads $\nabla \cdot J^{(s)} =$ HS Ward sources (modulo BRST–exact improvements) [82, 91]. Define the HS *law* visibility by

$$\Lambda_{\text{law}}^{[s]} := 1 - \frac{\big\|\text{HS Ward sources}\big\|_{\mathfrak{P}}}{\big\|\nabla \cdot J^{(s)}\big\|_{\text{ref},\mathfrak{P}}} \in [0,1],$$

with plateau–uniform BRST–safe norms [82, 112]. Let $\varepsilon_{\text{law}}, \varepsilon_{\text{state}} \in (0,1)$ be the certified tolerances (cf. the plateau criterion). For the *state* witness we use plateau–tolerance semantics:

$$\text{unbroken on } \mathfrak{P} \iff |\Lambda_{\text{state}}^{[s]}| \geq 1 - \varepsilon_{\text{state}}, \qquad \text{SSB on } \mathfrak{P} \iff |\Lambda_{\text{state}}^{[s]}| \leq \varepsilon_{\text{state}},$$

equivalently detected by a HS order parameter / would–be HS NG sector (for global) or by a BEH–like eating if gauged [94, 91, 112, 113]. All insertions are evaluated as $P_x(\lambda_{\text{UV}})\, K_{\text{IR}}^{(\ell)}(\cdot)\, K_{\text{IR}}^{(\ell)}\, P_x(\lambda_{\text{UV}})$.

Intuition. Breaking in the *state* channel produces a would–be HS NG sector; if the symmetry is gauged, a BEH–like mechanism can occur [94, 91, 112, 113]. Explicit breaking in the *law* channel yields persistent masses/mixings; within a certified plateau we read these via the HS Ward sources [82, 102, 112].

Lemma 8.19 (Helicity stability under explicit deformations)**.** *Within the admissible band on $\mathfrak{P}$ and for small explicit breakings, the helicity content of $\mathscr{H}_s$ is invariant: pole locations and residues shift* analytically *while spin labels are unchanged.*

Proposition 8.20 (Higher–spin quadrant assignment (tolerance form))**.** *On $\mathfrak{P}$,*

$$\textit{QI:}\quad |\Lambda_{\text{law}}^{[s]}| \geq 1 - \varepsilon_{\text{law}}, \qquad |\Lambda_{\text{state}}^{[s]}| \geq 1 - \varepsilon_{\text{state}}, \tag{8.2}$$

$$\textit{QII:}\quad |\Lambda_{\text{law}}^{[s]}| \geq 1 - \varepsilon_{\text{law}}, \qquad |\Lambda_{\text{state}}^{[s]}| \leq \varepsilon_{\text{state}} \ \textit{(SSB; if gauged, HS BEH may operate)}, \tag{8.3}$$

$$\textit{QIII:}\quad 1 - |\Lambda_{\text{law}}^{[s]}(\ell)| \leq C\,\ell^{\,\Delta-(d+z)} \ (\ell \downarrow 0), \qquad |\Lambda_{\text{state}}^{[s]}| \geq 1 - \varepsilon_{\text{state}}, \tag{8.4}$$

$$\textit{QIV:}\quad \exists\, \delta_\star \in (\varepsilon_{\text{law}}, 1) : \ |\Lambda_{\text{law}}^{[s]}| \leq 1 - \delta_\star \ \textit{uniformly on } \mathfrak{P}, \qquad |\Lambda_{\text{state}}^{[s]}| \geq 1 - \varepsilon_{\text{state}}. \tag{8.5}$$

Comments. *QI: massless spin–s representations appear if the background allows; QII: state SSB (global: HS NG sector; gauged: BEH–like eating); QIII: IR healing of irrelevant explicit breakings in the law channel; QIV: persistent explicit mass/mixing; spins fixed by Lemma 8.19.*

Intuition. The quadrant logic is spin–agnostic; only the witnesses differ. Tolerance semantics ensures consistency with the certified–plateau framework used throughout.

8.1.5 Hydrodynamics Witnesses

Definition 8.21 (Hydrodynamic witnesses (tolerance form))**.** Work on a certified plateau $\mathfrak{P}$ and in $\mathcal{H}_{\rm phys} = H^0(\hat{Q}_{\rm BRST})$ with BRST–closed scale tags [82, 91]. Let $G^R_{nn}(\omega, \mathbf{k})$ denote the retarded density correlator near $\omega, \mathbf{k} \to 0$ [119, 102]. On $\mathfrak{P}$ (isotropic IR window $K^{(\ell)}_{\rm IR}$ so that $k \sim \ell^{-1}$),

$$G^R_{nn}(\omega, \mathbf{k}) \;\simeq\; \frac{\chi\, D(\ell)\, k^2}{-i\omega + D(\ell)k^2 + \Gamma(\ell)}\,,$$

which is the standard diffusive pole structure for conserved densities in linear response [119, 102]. Define the *law* visibility and adopt plateau tolerances $\varepsilon_{\rm law}, \varepsilon_{\rm state} \in (0, 1)$:

$$\Lambda^{[\rm hydro]}_{\rm law}(\ell) \;:=\; \frac{D(\ell)k^2}{D(\ell)k^2 + \Gamma(\ell)}\Big|_{k\sim\ell^{-1}}, \qquad \text{“law symmetric”} \iff |\Lambda^{[\rm hydro]}_{\rm law}| \;\geq\; 1 - \varepsilon_{\rm law}.$$

For the *state* witness:

$$\text{“state symmetric”} \iff |\Lambda^{[\rm hydro]}_{\rm state}| \;\geq\; 1 - \varepsilon_{\rm state} \quad \text{(pure diffusion near } k \to 0),$$

$$\text{“state SSB”} \iff |\Lambda^{[\rm hydro]}_{\rm state}| \;\leq\; \varepsilon_{\rm state} \quad \text{(Goldstone sound: } \omega = \pm c_s k - i\gamma k^2 + \cdots)$$

where diffusive and sound modes are the standard hydrodynamic poles associated with conserved charges and spontaneously broken symmetries [119, 102, 112]. *Convention.* In this hydrodynamic block we use an *isotropic* IR window $K^{(\ell)}_{\rm IR}$ (so $k \sim \ell^{-1}$), in contrast to the *anisotropic* $K^{(\vec{\ell})}_{\rm IR}$ employed for the fracton/subsystem sector.

Explanatory remarks. The definition sets up hydrodynamic witnesses in complete analogy with the other sectors of the quadrant scheme, but specialized to the long–wavelength, low–frequency behavior of a conserved density. The working arena is a certified plateau $\mathfrak{P}$ and the physical BRST cohomology $\mathcal{H}_{\rm phys} = H^0(\hat{Q}_{\rm BRST})$ with BRST–closed IR/UV tags. This guarantees that all correlators and Ward identities entering the witnesses are already projected to the physical sector and are insensitive to gauge–parameter choices, within the plateau tolerances [82, 91].

The basic observable is the retarded correlator $G^R_{nn}(\omega, \mathbf{k})$ of a conserved density n, evaluated in the hydrodynamic regime $\omega, \mathbf{k} \to 0$ within an isotropic IR window $K^{(\ell)}_{\rm IR}$ so that the momentum scale is effectively $k \sim \ell^{-1}$. In this regime the correlator is assumed to exhibit the standard diffusive pole structure

$$G^R_{nn}(\omega, \mathbf{k}) \;\simeq\; \frac{\chi\, D(\ell)\, k^2}{-i\omega + D(\ell)k^2 + \Gamma(\ell)}\,,$$

where χ is the static susceptibility, $D(\ell)$ is a possibly scale–dependent diffusion constant, and $\Gamma(\ell)$ is an effective relaxation rate encoding explicit violation of the associated conservation law [119, 102]. The denominator $-i\omega + Dk^2 + \Gamma$ thus separates reversible diffusive transport (Dk^2) from irreversible leakage (Γ).

On this basis, the hydrodynamic *law* witness

$$\Lambda_{\text{law}}^{[\text{hydro}]}(\ell) := \frac{D(\ell)k^2}{D(\ell)k^2 + \Gamma(\ell)}\Big|_{k\sim\ell^{-1}}$$

is introduced as a dimensionless visibility in $[0,1]$ which measures the fraction of the decay rate attributable to diffusion rather than explicit leakage. When $\Gamma(\ell)$ is negligible compared to $D(\ell)k^2$ at the probed scale, the ratio is close to 1 and the law channel is effectively symmetric. Conversely, sizable $\Gamma(\ell)$ drives $\Lambda_{\text{law}}^{[\text{hydro}]}$ away from 1, indicating a persistent law–deficit. The plateau tolerance ε_{law} then provides the precise semantics: "law symmetric'' corresponds to $|\Lambda_{\text{law}}^{[\text{hydro}]}| \geq 1 - \varepsilon_{\text{law}}$, while significant deficits appear as violations of this bound [119, 112].

The *state* witness is defined at the same conceptual level but keyed to the qualitative nature of the hydrodynamic poles. In the symmetric state, the density mode is purely diffusive in the long–wavelength limit $k \to 0$; in the spontaneously broken phase, the hydrodynamic spectrum contains a propagating sound–mode Goldstone with dispersion $\omega = \pm c_s k - i\gamma k^2 + \cdots$. The visibility $\Lambda_{\text{state}}^{[\text{hydro}]}$ is not given by an explicit formula here, but its tolerance semantics are fixed: "state symmetric'' corresponds to $|\Lambda_{\text{state}}^{[\text{hydro}]}| \geq 1 - \varepsilon_{\text{state}}$ (diffusive regime), and "state SSB'' corresponds to $|\Lambda_{\text{state}}^{[\text{hydro}]}| \leq \varepsilon_{\text{state}}$ (sound Goldstone regime). In this way, the diffusive versus sound character of the long–wavelength excitation is compressed into a binary state–channel predicate, compatible with the global quadrant diagnostic scheme [119, 102, 112].

Finally, the convention of an isotropic IR window $K_{\text{IR}}^{(\ell)}$ (so that $k \sim \ell^{-1}$) is recorded explicitly to distinguish the hydrodynamic sector from the anisotropic IR windows $K_{\text{IR}}^{(\vec{\ell})}$ required later for fracton and subsystem–symmetric sectors, where different components of $\vec{k}$ scale differently with the IR lengths. This ensures that the hydrodynamic witnesses are interpreted within a standard isotropic scaling framework while remaining fully compatible with the BRST/plateau infrastructure of the broader formalism [102, 112].

Lemma 8.22 (Hydrodynamic law predicate (tolerance equivalences))**.** *On $\mathfrak{P}$ and at $k \sim \ell^{-1}$,*

$$|\Lambda_{\text{law}}^{[\text{hydro}]}| \;\geq\; 1 - \varepsilon_{\text{law}} \quad\Longleftrightarrow\quad \Gamma(\ell) \;\leq\; \frac{\varepsilon_{\text{law}}}{1 - \varepsilon_{\text{law}}}\, D(\ell)\, k^2.$$

In particular, the sharp limit $\varepsilon_{\text{law}} \to 0$ reproduces $\Gamma(\ell) = 0$ (no relaxation term in the continuity equation on the band).

Proof sketch. From $\Lambda_{\text{law}} = Dk^2/(Dk^2 + \Gamma)$ one has $1 - \Lambda_{\text{law}} = \Gamma/(Dk^2 + \Gamma)$. The inequality $1 - \Lambda_{\text{law}} \leq \varepsilon_{\text{law}}$ is equivalent to $\Gamma \leq \frac{\varepsilon_{\text{law}}}{1-\varepsilon_{\text{law}}} Dk^2$. Taking $\varepsilon_{\text{law}} \to 0$ yields $\Gamma = 0$. □

Proposition 8.23 (Hydrodynamic quadrant assignment (tolerance form))**.** *On a certified*

plateau $\mathfrak{P}$ *(with* $k \sim \ell^{-1}$*):*

$$\begin{aligned}
&\textit{QI:} \quad |\Lambda_{\mathrm{law}}^{[\mathrm{hydro}]}| \;\geq\; 1-\varepsilon_{\mathrm{law}},\\
&\qquad\; |\Lambda_{\mathrm{state}}^{[\mathrm{hydro}]}| \;\geq\; 1-\varepsilon_{\mathrm{state}} \quad \textit{(pure diffusion)},\\[2ex]
&\textit{QII:} \quad |\Lambda_{\mathrm{law}}^{[\mathrm{hydro}]}| \;\geq\; 1-\varepsilon_{\mathrm{law}},\\
&\qquad\; |\Lambda_{\mathrm{state}}^{[\mathrm{hydro}]}| \;\leq\; \varepsilon_{\mathrm{state}} \quad \textit{(Goldstone sound present)},\\[2ex]
&\textit{QIII:} \quad 1-|\Lambda_{\mathrm{law}}^{[\mathrm{hydro}]}(\ell)| \;\leq\; C\,\ell^{\,\beta} \quad (\ell\to\infty)\\
&\qquad\qquad \textit{(equivalently } \Gamma(\ell) \ll D(\ell)k^2 \textit{ at } k\sim\ell^{-1}),\\
&\qquad\; |\Lambda_{\mathrm{state}}^{[\mathrm{hydro}]}| \;\geq\; 1-\varepsilon_{\mathrm{state}},\\[2ex]
&\textit{QIV:} \quad \exists\,\delta_\star\in(\varepsilon_{\mathrm{law}},1):\; |\Lambda_{\mathrm{law}}^{[\mathrm{hydro}]}| \;\leq\; 1-\delta_\star \;\; \textit{uniformly on } \mathfrak{P}\\
&\qquad\qquad \Big(\Leftrightarrow\; \Gamma(\ell) \;\geq\; \frac{\delta_\star}{1-\delta_\star}\,D(\ell)k^2\Big),\\
&\qquad\; |\Lambda_{\mathrm{state}}^{[\mathrm{hydro}]}| \;\geq\; 1-\varepsilon_{\mathrm{state}}.
\end{aligned}$$

Proof sketch. Combine the pole structure of G^R_{nn} with Lemma 8.22. QIII vs. QIV corresponds to *decay vs. persistent gap* of the law deficit on $\mathfrak{P}$: either $1-\Lambda_{\mathrm{law}}^{[\mathrm{hydro}]}(\ell)\to 0$ with a positive power as $\ell\to\infty$ (healing under IR dilation, equivalently $\Gamma \ll Dk^2$ at $k\sim\ell^{-1}$) or it remains uniformly bounded away from 0, which translates to the stated lower bound on Γ/Dk^2. □

8.1.6 Second Sound in the Hydrodynamic Sector (Matrix Formulation)

The longitudinal hydrodynamic sector admits a richer structure than the single–density setting of Definition 8.21. On a certified plateau $\mathfrak{P}$ and within the BRST–physical subspace $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$, the same UV/IR instrumentation $(P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}^{(\ell)})$ ensures that all retarded correlators, Ward identities, and transport data are evaluated in a gauge–parameter–independent manner [82, 91]. In this regime, hydrodynamics is governed by the low–frequency, long–wavelength behaviour of conserved densities and their reversible/dissipative couplings, which admit a unified matrix formulation in terms of their retarded two–point functions [450, 102].

In contrast to the single-density case—where diffusion provides the sole hydrodynamic pole—the coupled longitudinal sector requires the 2×2 matrix of retarded correlators for a pair of conserved or thermodynamic densities, such as (n,s) or (n,ε). On the isotropic IR window $K_{\mathrm{IR}}^{(\ell)}$ (so that $k\sim\ell^{-1}$), the inverse correlator admits the standard hydrodynamic expansion with three ingredients: a positive–definite susceptibility matrix $\boldsymbol{\chi}$, a dissipative matrix $\mathbf{D}$ containing diffusion and thermo-electric coefficients, and a relaxation matrix $\boldsymbol{\Gamma}\geq 0$ encoding explicit violations of the continuity equations on the band [450, 119, 112]. The hydrodynamic poles are then determined by the secular condition

$$\det\!\Big(\mathbf{G}^{\mathrm{R}}(\omega,\mathbf{k})^{-1}\Big) = 0,$$

whose solutions capture both diffusive modes and possible propagating sound branches.

A key structural feature of this block is the existence of *two* possible propagating modes. When the reversible couplings generate a nondegenerate elastic matrix $\mathcal{K}$ in the longitudinal sector, the linear–in–k propagating solutions have speeds given by the positive eigenvalues of the generalised stiffness matrix $S = \chi^{-1}\mathcal{K}$. This reproduces, in particular, the well–known first–sound and second–sound branches of two–fluid hydrodynamics [450, 451, 452]. The attenuations arise at order k^2 through Rayleigh quotients involving the dissipative matrix $\mathbf{D}$, ensuring nonnegative damping consistent with thermodynamic stability [450].

Second sound is thus naturally understood as the presence of a *second*, linearly dispersing hydrodynamic mode in the longitudinal block. Its absence corresponds either to a purely diffusive sector or to a single sound branch, while explicit breaking generates "pseudo–sound" behaviour in which the linear dispersion is retained but shifted or gapped at $\mathcal{O}(\epsilon)$, analogously to pseudo–Goldstone modes in other sectors [112]. This matrix formulation therefore upgrades the scalar hydrodynamic witness to a spectral witness capable of diagnosing first versus second sound on the same certified plateau $\mathfrak{P}$, while remaining fully compatible with BRST closure, IR/UV windowing, causality, and the law/state visibility interpretation used throughout the quadrant framework.

All statements are evaluated on a certified plateau $\mathfrak{P}$ and on the BRST–physical sector $\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}})$, with the same UV/IR instrumentation $(P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}^{(\ell)})$ and notation as in Definition 8.21, Lemma 8.22, and Proposition 8.23. Causality/cone preservation and band safety are as in Proposition 6.281.

Definition 8.24 (Longitudinal two–variable hydrodynamic block)**.** Let $A, B \in \{n, s\}$ label a *longitudinal* pair of conserved/scalar densities (e.g. particle/charge density n and entropy density s; one may equally use $A, B \in \{n, \varepsilon\}$ with energy density ε). Define the 2×2 matrix of retarded correlators

$$\mathbf{G}^{\mathrm{R}}_{AB}(\omega, \mathbf{k}) \;:=\; \begin{pmatrix} G^{\mathrm{R}}_{nn} & G^{\mathrm{R}}_{ns} \\ G^{\mathrm{R}}_{sn} & G^{\mathrm{R}}_{ss} \end{pmatrix}(\omega, \mathbf{k})\,, \qquad \mathbf{k} = k\,\hat{\mathbf{k}}, \quad k \to 0,$$

on the isotropic IR window $K_{\mathrm{IR}}^{(\ell)}$ so that $k \sim \ell^{-1}$. Near $(\omega, \mathbf{k}) \to (0, \mathbf{0})$, its inverse admits the hydrodynamic expansion

$$\mathbf{G}^{\mathrm{R}}(\omega, \mathbf{k})^{-1} \;\simeq\; -i\omega\,\boldsymbol{\chi} \;+\; k^2\,\mathbf{D} \;+\; \boldsymbol{\Gamma} \;+\; \mathcal{O}(\omega^2, \omega k^2, k^3)\,, \tag{8.6}$$

where $\boldsymbol{\chi}$ is the static susceptibility matrix (positive–definite on $\mathfrak{P}$), $\mathbf{D}$ collects dissipative transport coefficients (diffusivities and thermo–electric cross terms), and $\boldsymbol{\Gamma} \geq 0$ encodes relaxation terms that break the exact continuity equations on the band (as in Definition 8.21). The *hydrodynamic poles* are the solutions of

$$\det\Big(\mathbf{G}^{\mathrm{R}}(\omega, \mathbf{k})^{-1}\Big) = 0\,. \tag{8.7}$$

Remark 8.25 (Generator/secular form)**.** Equivalently to the expansion (8.6), the secular equation (8.7) can be written in the "generator" form

$$\det\Big(-\,i\omega\,\mathbf{1} - \mathbf{H}(k)\Big) = 0, \qquad \mathbf{H}(k) = \mathbf{R}\,k - \mathbf{D}\,k^2 - \boldsymbol{\Gamma}, \quad \mathbf{R}^{\dagger} = -\mathbf{R},$$

which makes the linear-in-k propagating roots manifest and yields the same sound branches $(\omega = \pm c_j k - i\gamma_j k^2 + \cdots)$ in the longitudinal sector; see [450].

Definition 8.26 (Second–sound predicate (spectral))**.** On $\mathfrak{P}$, *second sound is present* in the longitudinal hydrodynamic block if and only if there exist two distinct propagating branches

$$\omega_{\pm}^{(j)}(\mathbf{k}) \;=\; \pm\, c_j\, k \;-\; i\,\gamma_j\, k^2 \;+\; o(k^2)\,, \qquad c_j > 0,\ \gamma_j \geq 0, \quad j = 1, 2, \tag{8.8}$$

among the solutions of (8.7). The case $j = 1$ is the usual (first) sound; $j = 2$ is *second sound*. Absence of a second linear branch (with the remainder diffusive or gapped) means no second sound in this block.

Lemma 8.27 (Speeds from susceptibilities and reversible couplings)**.** *Assume thermodynamic stability on* $\mathfrak{P}$ *so that* $\chi > 0$ *and that the reversible couplings induce a nondegenerate symmetric* elastic *matrix* $\mathcal{K}$ *for the longitudinal sector (see Remark 8.28). Then the leading sound speeds are the positive square roots of the eigenvalues of the generalised stiffness matrix*

$$S \;:=\; \chi^{-1}\,\mathcal{K}\,, \qquad c_1^2,\ c_2^2 \;=\; \mathrm{Spec}\big(S\big) \;\subset\; \mathbb{R}_{>0}\,, \tag{8.9}$$

and the attenuations are $\gamma_j = \frac{1}{2}\,\mathbf{v}_j^{\top}\,\chi^{-1}\mathbf{D}\,\mathbf{v}_j \geq 0$ *for the* S*–orthonormal eigenvectors* $\mathbf{v}_j$*.*

Remark 8.28 (Thermoelastic parameters)**.** For (n, s) variables, $\mathcal{K}$ is expressed via isentropic/isothermal compressibilities and thermo–elastic cross derivatives [450]. In a two–fluid superfluid, $\mathcal{K}$ reduces to Landau's coefficients [451, 452].

Sketch. Insert the ansatz (8.8) into (8.7). The $\mathcal{O}(k^2)$ real part yields $\det\big(c^2\,\chi - \mathcal{K}\big) = 0$, whose positive roots give (8.9); the $\mathcal{O}(k^3)$ imaginary part fixes γ_j from the Rayleigh quotient with $\mathbf{D}$ [450]. Positivity follows from $\chi > 0$ and $\mathbf{D} \geq 0$ on $\mathfrak{P}$. □

Proposition 8.29 (Hydrodynamic diagnosis with two longitudinal variables)**.** *With the block of Definition 8.24 and under the plateau tolerances of Definition 8.21:*

(i) QI.
No NG/pNG in the block; if present, at most one linear branch and it is symmetry–allowed.

(ii) QII.
Second sound present*: two branches (8.8) with* $c_{1,2} > 0$*.*

(iii) QIII.
No persistent *NG/pNG; as* $k \to 0$ *one or both branches revert to diffusion (healing).*

(iv) QIV.
"Pseudo–sound": linear structure retained but gapped/relaxing at $\mathcal{O}(\epsilon)$*.*

In the sense that $\omega_{\pm}^{(2)}(k) = \pm c_2 k - i\gamma_2 k^2$ *in QII is replaced by*

$$\omega_{\pm}^{(2)}(k) = \pm c_2 k - i\gamma_2 k^2 \;\underline{- i\, m_{\mathrm{ph}}^2/c_2}$$

in QIV, with $m_{\mathrm{ph}}^2 = \mathcal{O}(\epsilon)$ *(pseudo–phonon/pseudo–sound), while QIII enforces the decay of any such gapless structure under IR dilation (cf. Proposition 8.23).*

Remark 8.30 (Canonical superfluid formula for c_2)**.** In the Landau two–fluid limit (normal and superfluid components with densities ρ_n, ρ_s, entropy density s, temperature T, and specific heat at constant pressure c_p), the general expression (8.9) reproduces the textbook second–sound speed [451, 452, 450]:

$$c_2^2 \;=\; \frac{\rho_s}{\rho_n}\,\frac{s^2\,T}{c_p}\,, \tag{8.10}$$

with attenuation γ_2 fixed by the relevant dissipative coefficients (thermal conductivity, mutual friction, viscosities) [450].

Operational takeaway. To *explicitly include* second sound in the hydrodynamic sector, replace the single-scalar density witness of Definition 8.21 by the longitudinal 2×2 block of Definition 8.24 and diagnose second sound via the presence of two linear hydrodynamic poles (8.8), whose speeds follow from the generalised stiffness (8.9). This formulation is self–contained (no parasymmetry invoked) and sits on the same certified plateau $\mathfrak{P}$, preserving the law/state interpretation and the QI–QIV logic of Proposition 8.23.

8.1.7 Boundary Soft/Edge Modes Witnesses

The boundary sector admits a natural analogue of the bulk visibility framework, tailored to soft and edge modes localized near $\partial\mathcal{X}$. The working arena is a certified *boundary plateau* $\mathfrak{P}_\partial$, equipped with a tangential IR window $K^{(L)}_{\mathrm{IR},\partial}$ of size L along the boundary, and with insertions evaluated in the BRST–physical subspace $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ using BRST–closed UV/IR tags [82, 91]. This infrastructure ensures that boundary Ward identities and correlation functions are defined modulo BRST–exact terms and are insensitive to gauge–parameter choices within the plateau tolerances, in precise analogy with the bulk constructions [112, 102].

Within this setting, the tagged boundary Ward identity

$$\nabla_{\parallel} \cdot J^{(\partial)}_{\parallel} \;=\; S_\partial$$

encodes both the tangential boundary current $J^{(\partial)}_{\parallel}$ and a boundary source S_∂ that measures explicit breaking of the would–be conserved quantity at the edge: for example through leakage into the bulk, boundary driving, or explicit symmetry violation at $\partial\mathcal{X}$ [112, 102]. The boundary *law* visibility $\Lambda^{[\partial]}_{\mathrm{law}}(L)$ is then defined as a normalized deficit

$$1 - \frac{\|S_\partial\|_{\mathfrak{P}_\partial}}{\|\nabla_{\parallel} \cdot J^{(\partial)}_{\parallel}\|_{\mathrm{ref},\mathfrak{P}_\partial}},$$

providing a dimensionless measure of how strongly the boundary Ward identity is violated at the probed scale. Plateau tolerances $\varepsilon_{\mathrm{law}}$ convert this into a binary predicate: "boundary law symmetric'' whenever $|\Lambda^{[\partial]}_{\mathrm{law}}(L)| \geq 1 - \varepsilon_{\mathrm{law}}$, and law–deficit otherwise [112, 102].

The *state* channel is diagnosed through the presence or absence of boundary spontaneous symmetry breaking. A boundary state is declared symmetric when $|\Lambda^{[\partial]}_{\mathrm{state}}| \geq 1 - \varepsilon_{\mathrm{state}}$, and spontaneously broken when $|\Lambda^{[\partial]}_{\mathrm{state}}| \leq \varepsilon_{\mathrm{state}}$, in which case an edge Goldstone mode is

present on $\partial\mathcal{X}$ [92, 94, 102, 112]. The joint behaviour of $\Lambda^{[\partial]}_{\mathrm{law}}(L)$ and $\Lambda^{[\partial]}_{\mathrm{state}}$ under IR dilation $L \to \infty$ then feeds into the quadrant assignment: healing versus persistent boundary sources distinguish QIII from QIV in the law channel, while the existence of an edge Goldstone separates QII from QI in the state channel. In this way, boundary soft/edge modes are incorporated into the same BRST–safe, tolerance–based quadrant scheme that governs the bulk witnesses.

Definition 8.31 (Boundary witnesses (tolerance form))**.** Work on the certified boundary plateau $\mathfrak{P}_\partial$ with a tangential IR window $K^{(L)}_{\mathrm{IR},\partial}$ of size L along $\partial\mathcal{X}$, and evaluate in the physical sector $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ with BRST–closed tags. Let the tagged boundary Ward identity be $\nabla_\parallel \cdot J^{(\partial)}_\parallel = S_\partial$ (modulo BRST–exact improvements). Define the boundary *law* visibility

$$\Lambda^{[\partial]}_{\mathrm{law}}(L) \;:=\; 1 - \frac{\|S_\partial\|_{\mathfrak{P}_\partial}}{\|\nabla_\parallel \cdot J^{(\partial)}_\parallel\|_{\mathrm{ref},\mathfrak{P}_\partial}} \in [0,1],$$

with plateau–uniform, BRST–safe norms on $\mathfrak{P}_\partial$. Let $\varepsilon_{\mathrm{law}}, \varepsilon_{\mathrm{state}} \in (0,1)$ be the certified tolerances (cf. the plateau criterion). For the *state* witness we adopt tolerance semantics:

$$\begin{aligned} \text{“boundary state symmetric”} \quad &\Longleftrightarrow \quad |\Lambda^{[\partial]}_{\mathrm{state}}| \;\geq\; 1 - \varepsilon_{\mathrm{state}}, \\ \text{“boundary SSB”} \quad &\Longleftrightarrow \quad |\Lambda^{[\partial]}_{\mathrm{state}}| \;\leq\; \varepsilon_{\mathrm{state}} \;\; \text{(edge Goldstone present)}. \end{aligned}$$

All insertions are evaluated as $P_x(\lambda_{\mathrm{UV}})\, K^{(L)}_{\mathrm{IR},\partial}(\cdot)\, K^{(L)}_{\mathrm{IR},\partial}\, P_x(\lambda_{\mathrm{UV}})$ and understood modulo BRST–exact improvements.

Proposition 8.32 (Boundary quadrant assignment (tolerance form))**.** *On $\mathfrak{P}_\partial$:*

(i) QI.
$|\Lambda^{[\partial]}_{\mathrm{law}}(L)| \;\geq\; 1 - \varepsilon_{\mathrm{law}}, \qquad |\Lambda^{[\partial]}_{\mathrm{state}}| \;\geq\; 1 - \varepsilon_{\mathrm{state}}.$

(ii) QII.
$|\Lambda^{[\partial]}_{\mathrm{law}}(L)| \;\geq\; 1 - \varepsilon_{\mathrm{law}}, \qquad |\Lambda^{[\partial]}_{\mathrm{state}}| \;\leq\; \varepsilon_{\mathrm{state}}$ *(edge Goldstone present).*

(iii) QIII.
$1 - |\Lambda^{[\partial]}_{\mathrm{law}}(L)| \;\leq\; C\, L^{-\beta} \;\; (L \to \infty), \qquad |\Lambda^{[\partial]}_{\mathrm{state}}| \;\geq\; 1 - \varepsilon_{\mathrm{state}}.$

(iv) QIV.
$\exists\, \delta_\star \in (\varepsilon_{\mathrm{law}}, 1) : \; |\Lambda^{[\partial]}_{\mathrm{law}}(L)| \;\leq\; 1 - \delta_\star$ *uniformly on* $\mathfrak{P}_\partial$, $\qquad |\Lambda^{[\partial]}_{\mathrm{state}}| \;\geq\; 1 - \varepsilon_{\mathrm{state}}.$

Proof sketch. From the boundary Ward identity, the boundary law deficit is proportional to $\|S_\partial\|_{\mathfrak{P}_\partial}$. The tolerance inequalities translate the sharp conditions $S_\partial = 0$ (QI/QII) and the IR healing/persistence (QIII/QIV) into plateau–uniform bounds on $1 - \Lambda^{[\partial]}_{\mathrm{law}}(L)$: either it decays like $L^{-\beta}$ as $L \to \infty$ (healing) or stays uniformly bounded away from 0 (persistent source). The state channel is fixed by the presence/absence of an edge Goldstone mode, expressed via the tolerance predicate. □

8.1.8 Anyons (Chern-Simons) Witnesses

Anyons and Chern–Simons (CS) responses in 2+1 dimensions provide a natural topological analogue of the bulk visibility framework. On the certified bulk plateau $\mathfrak{P}$, the effective response is encoded in a band–limited, BRST–clean effective action $S_{\text{eff}}[A, B; \cdot]$ for background gauge fields: a 1–form A for ordinary $U(1)_0$ charge and a 2–form B for $U(1)_1$ (higher–form charge) [102, 112]. Evaluation is performed in the BRST–physical sector $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$, with BRST–closed IR/UV tags ensuring that all visible couplings, Ward identities, and anomaly constraints are gauge–parameter independent [82, 91]. Within this structure, the symmetry $X \in \{P, T, U(1)_0, U(1)_1\}$ is probed through its variation $\delta_X S_{\text{eff}}$, which captures both topological CS responses and explicit symmetry–breaking terms. Because CS levels and mixed Chern–Simons couplings are quantized, symmetry violation is often discrete and persistent unless cancelled by multi–sector inflow or anomaly matching [102, 112].

The resulting *law* witness $\Lambda_{\text{law}}^{[\text{anyon};X]}$ thus distinguishes two regimes: (i) exact symmetry under X, detected by the vanishing of $\delta_X S_{\text{eff}}$, including the absence of symmetry–violating CS or gravitational–CS responses (e.g. $k_{\text{tot}} = 0$, $c_- = 0$); and (ii) persistent explicit breaking, where quantized CS or anomaly terms remain even after band projection, leading to $\Lambda_{\text{law}}^{[\text{anyon};X]} = 0$ [102, 112]. Intermediate cases, where explicit X–breaking terms are RG–irrelevant on $\mathfrak{P}$, correspond to $\Lambda_{\text{law}}^{[\text{anyon};X]}(\ell) \to 1$ as $\ell \to \infty$, in alignment with the QIII healing pattern of the general quadrant framework.

The *state* witness $\Lambda_{\text{state}}^{[\text{anyon};X]}$ distinguishes whether the ground sector preserves X at large scales. In the X–symmetric case, symmetry defects $D_X(\gamma)$ act trivially on large Wilson lines or correlation functions, and no X–charged order parameter acquires an expectation value. In the spontaneously broken case, the defect acts nontrivially or a charged order parameter signals X–SSB, producing $\Lambda_{\text{state}}^{[\text{anyon};X]} = 0$ [102, 112]. Together, the law and state witnesses supply a binary visibility pair governing the X–quadrant assignment, integrating anyonic and CS responses seamlessly into the BRST–safe, plateau–stabilized quadrant framework.

Definition 8.33 (Anyonic witnesses for a chosen symmetry X)**.** Work in 2+1D on the bulk plateau $\mathfrak{P}$. Fix $X \in \{P, T, U(1)_0, U(1)_1\}$. Couple the appropriate background fields (a 1-form A for $U(1)_0$, a 2-form B for $U(1)_1$) and let $S_{\text{eff}}[A, B; \cdot]$ be the band-limited, BRST-clean effective action. Define

$$\Lambda_{\text{law}}^{[\text{anyon};X]} := \begin{cases} 1, & \delta_X S_{\text{eff}} = 0 \text{ on } \mathfrak{P}, \\ 0, & \delta_X S_{\text{eff}} \neq 0 \text{ on } \mathfrak{P}, \end{cases}$$

$$\Lambda_{\text{state}}^{[\text{anyon};X]} := \begin{cases} 1, & X\text{-symmetric ground sector on } \mathfrak{P}, \\ 0, & X\text{-SSB on } \mathfrak{P}. \end{cases}$$

For P/T: $\delta_X S_{\text{eff}} = 0$ iff the net CS and gravitational-CS responses vanish, i.e. $k_{\text{tot}} = 0$ and $c_- = 0$ (modulo X-allowed counterterms and inflow).

Operationally: for P/T, $\Lambda_{\text{law}}^{[\text{anyon};X]} = 1$ iff no symmetry-violating CS/gravitational-CS terms remain (e.g. net level $k_{\text{tot}} = 0$ and $c_- = 0$ modulo allowed counterterms); for $U(1)_0$ and $U(1)_1$, it equals 1 iff the corresponding Ward identities hold for S_{eff} (equivalently, invariance under $A \mapsto A + \mathrm{d}\alpha$ or $B \mapsto B + \mathrm{d}\beta$). The state witness can be read, e.g., from symmetry

defects $D_X(\gamma)$ acting trivially on correlators at large scales or from the absence of X-charged order parameters. *Here c_- denotes the chiral central charge (gravitational Chern–Simons response / thermal Hall coefficient).*

Proposition 8.34 (Anyonic quadrant assignment for X)**.** *With the witnesses of Definition 8.33, the quadrant for the chosen X is*

(i) QI.
$(1,1)$.

(ii) QII.
$(1,0)$.

(iii) QIII.
$\Lambda_{\mathrm{law}}^{[\mathrm{anyon};X]}(\ell) \to 1, \quad \Lambda_{\mathrm{state}}^{[\mathrm{anyon};X]} = 1.$

(iv) QIV.
$\Lambda_{\mathrm{law}}^{[\mathrm{anyon};X]} = 0$ *(persistent law deficit).*

Proof sketch. Evaluate the X-Ward/constraint under $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ in S_{eff}. If symmetry-violating terms survive (e.g. quantized CS or gravitational-CS that flips under P/T; an uncanceled mixed anomaly; or an X-violating coupling that is not RG-irrelevant on $\mathfrak{P}$), then $\Lambda_{\mathrm{law}}^{[\mathrm{anyon};X]} = 0$ and the law channel sits in **QIV**. If all explicit X-breaking contributions become irrelevant or cancel (e.g. sector sum with $k_{\mathrm{tot}} = 0$), then $\Lambda_{\mathrm{law}}^{[\mathrm{anyon};X]}(\ell) \to 1$ giving **QIII**. When the law is symmetric, the state predicate alone distinguishes **QI** vs **QII**. □

Remark 8.35 (Quantization locks and topological order)**.** (i) For P/T in pure CS TQFTs, level quantization typically makes the law deficit persistent (**QIV**) unless a multi-sector cancellation or inflow restores X (then **QIII**). (ii) Stability of modular data/topological order is orthogonal to the quadrant label and may be tracked by separate diagnostics (does not enter $\Lambda_{\mathrm{law}}, \Lambda_{\mathrm{state}}$).

8.1.9 Fractons Witnesses

Subsystem and fracton phases provide paradigmatic examples of anisotropic and higher–moment conservation laws, where charges can be immobile or restricted to lower–dimensional foliations, and where dipole (or higher multipole) conservation plays a central role in the long–distance dynamics [102, 112]. Within the present framework, such phases are analysed on a certified plateau $\mathfrak{P}$ and in the BRST–physical sector $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$, using BRST–closed UV/IR tags so that Ward identities and coarse–grained correlators are insensitive to gauge–parameter choices [82, 91]. The key structural difference from ordinary hydrodynamics is the use of *anisotropic* IR windows $K_{\mathrm{IR}}^{(\vec{\ell})}$ with $\vec{\ell} = (\ell_\parallel, \ell_\perp, \dots)$ adapted to the foliation or subsystem structure: longitudinal momenta scale as $\mathbf{k}_\parallel \sim \ell_\parallel^{-1}$, while transverse components scale as $\mathbf{k}_\perp \sim \ell_\perp^{-1}$, reflecting the restricted mobility and direction–dependent screening characteristic of fracton–like systems [102, 112].

In a rank–2 $U(1)$ description, the relevant Ward identities include both the ordinary charge continuity equation and a dipole (or higher–moment) continuity equation, schematically $\partial_t \rho + \nabla \cdot \mathbf{J} = S^{(1)}$, $\partial_t \mathbf{P} + \nabla \cdot \mathbf{J}^{(2)} = S^{(2)}$, with $S^{(1)}, S^{(2)}$ encoding explicit violations of the

corresponding subsystem symmetries. The subsystem *law* witness $\Lambda^{[\mathrm{sub}]}_{\mathrm{law}}(\vec{\ell})$ is defined as a dimensionless visibility that compares the norm of the residuals $W_{\mathrm{sub}} = (S^{(1)}, S^{(2)})$ to reference norms of the continuity operators, evaluated at $\mathbf{k} \sim \vec{\ell}^{-1}$. Values of $\Lambda^{[\mathrm{sub}]}_{\mathrm{law}}$ near unity indicate exact or approximate multipole conservation at the anisotropic scales set by $K^{(\vec{\ell})}_{\mathrm{IR}}$, while significant deficits signal persistent explicit breaking that does not wash out under anisotropic coarse–graining [112, 102]. Tolerance parameters $\varepsilon_{\mathrm{law}}$ implement this distinction in the same way as in the bulk and hydrodynamic sectors.

The subsystem *state* witness $\Lambda^{[\mathrm{sub}]}_{\mathrm{state}}$ captures spontaneous breaking of the subsystem symmetry through appropriate order parameters or long–distance correlators. In the symmetric regime, $|\Lambda^{[\mathrm{sub}]}_{\mathrm{state}}| \geq 1 - \varepsilon_{\mathrm{state}}$ and no crystalline or foliated pattern carries a nonzero expectation value; in the broken regime, $|\Lambda^{[\mathrm{sub}]}_{\mathrm{state}}| \leq \varepsilon_{\mathrm{state}}$, and the system exhibits fracton–like ordering tied to the subsystem symmetry [102, 112]. The combined behaviour of $\Lambda^{[\mathrm{sub}]}_{\mathrm{law}}(\vec{\ell})$ under anisotropic dilation and of $\Lambda^{[\mathrm{sub}]}_{\mathrm{state}}$ then feeds directly into the quadrant assignment: decay versus persistence of the law deficit under $\min(\vec{\ell}) \to \infty$ distinguishes QIII from QIV, while the presence or absence of subsystem SSB separates QII from QI, thereby embedding fracton and subsystem–symmetric phases into the same BRST–safe, plateau–stabilized quadrant framework.

Definition 8.36 (Subsystem witnesses (anisotropic arena, tolerance form))**.** Work on a certified plateau $\mathfrak{P}$ and in $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ with BRST–closed scale tags. Adopt *anisotropic* IR windows $K^{(\vec{\ell})}_{\mathrm{IR}}$ with $\vec{\ell} = (\ell_{\parallel}, \ell_{\perp}, \dots)$ that respect the foliation/subsystem structure; probe momenta obey $\mathbf{k}_{\parallel} \sim \ell_{\parallel}^{-1}$ and $\mathbf{k}_{\perp} \sim \ell_{\perp}^{-1}$ on $\mathfrak{P}$. For rank–2 $U(1)$, the tagged continuity equations read

$$\partial_t \rho + \nabla \cdot \mathbf{J} = S^{(1)}, \qquad \partial_t \mathbf{P} + \nabla \cdot \mathbf{J}^{(2)} = S^{(2)}.$$

Let $W_{\mathrm{sub}} := (S^{(1)}, S^{(2)})$ collect the subsystem Ward residuals, and let $\| \cdot \|_{\mathfrak{P}}$ be a plateau–uniform, BRST–safe norm. Define the *law* visibility

$$\Lambda^{[\mathrm{sub}]}_{\mathrm{law}}(\vec{\ell}) := 1 - \frac{\|W_{\mathrm{sub}}\|_{\mathfrak{P}}}{\| \, \partial_t \rho + \nabla \cdot \mathbf{J} \, \|_{\mathrm{ref},\mathfrak{P}} + \| \, \partial_t \mathbf{P} + \nabla \cdot \mathbf{J}^{(2)} \, \|_{\mathrm{ref},\mathfrak{P}}} \in [0,1],$$

evaluated at $\mathbf{k} \sim \vec{\ell}^{-1}$. Let $\varepsilon_{\mathrm{law}}, \varepsilon_{\mathrm{state}} \in (0,1)$ be the certified tolerances. For the *state* witness we adopt tolerance semantics via appropriate subsystem order parameters/correlators:

$$\text{“state symmetric”} \iff |\Lambda^{[\mathrm{sub}]}_{\mathrm{state}}| \geq 1 - \varepsilon_{\mathrm{state}}, \qquad \text{“state SSB”} \iff |\Lambda^{[\mathrm{sub}]}_{\mathrm{state}}| \leq \varepsilon_{\mathrm{state}}.$$

All insertions are evaluated as $P_x(\lambda_{\mathrm{UV}})\, K^{(\vec{\ell})}_{\mathrm{IR}}(\cdot)\, K^{(\vec{\ell})}_{\mathrm{IR}}\, P_x(\lambda_{\mathrm{UV}})$ and are understood modulo BRST–exact improvements.

Intuition. The law predicate tests exact or approximate multipole conservation *at anisotropic scales* set by $K^{(\vec{\ell})}_{\mathrm{IR}}$; the state predicate detects crystalline/foliated ordering that breaks the subsystem symmetry. Tolerance semantics ($|\Lambda| \geq 1 - \varepsilon$ on $\mathfrak{P}$) keeps this block consistent with the global framework.

Lemma 8.37 (Subsystem IR healing criterion (tolerance form))**.** *If all explicit violations of the subsystem Ward identities are RG–irrelevant with respect to the anisotropic dilation induced by* $K_{\mathrm{IR}}^{(\vec{\ell})}$*, then there exists* $\beta > 0$ *such that, on* $\mathfrak{P}$*,*

$$1 - |\Lambda_{\mathrm{law}}^{[\mathrm{sub}]}(\vec{\ell})| \;\le\; C\,\min(\ell_{\parallel}, \ell_{\perp}, \dots)^{-\beta}, \qquad \text{as } \min(\vec{\ell}) \to \infty,$$

hence $|\Lambda_{\mathrm{law}}^{[\mathrm{sub}]}(\vec{\ell})| \to 1$ *and the law channel is "within tolerance" for large enough* $\vec{\ell}$*.*

Proof sketch. Treat the residuals W_{sub} as insertions of higher–dimension operators with *anisotropic* scaling dimensions. Irrelevance implies their tagged norm decays with the coarse–graining volume associated with $K_{\mathrm{IR}}^{(\vec{\ell})}$. Normalizing by the fixed reference denominator yields the stated power–law bound and the approach to unity of $\Lambda_{\mathrm{law}}^{[\mathrm{sub}]}$. □

Proposition 8.38 (Subsystem quadrant assignment (tolerance form))**.** *On* $\mathfrak{P}$ *with* $K_{\mathrm{IR}}^{(\vec{\ell})}$*:*

(i) QI.
$|\Lambda_{\mathrm{law}}^{[\mathrm{sub}]}(\vec{\ell})| \;\ge\; 1 - \varepsilon_{\mathrm{law}}, \qquad |\Lambda_{\mathrm{state}}^{[\mathrm{sub}]}| \;\ge\; 1 - \varepsilon_{\mathrm{state}}.$

(ii) QII.
$|\Lambda_{\mathrm{law}}^{[\mathrm{sub}]}(\vec{\ell})| \;\ge\; 1 - \varepsilon_{\mathrm{law}}, \qquad |\Lambda_{\mathrm{state}}^{[\mathrm{sub}]}| \;\le\; \varepsilon_{\mathrm{state}}.$

(iii) QIII.
$1 - |\Lambda_{\mathrm{law}}^{[\mathrm{sub}]}(\vec{\ell})| \;\le\; C\,\min(\vec{\ell})^{-\beta} \;\Big(\min(\vec{\ell}) \to \infty\Big), \qquad |\Lambda_{\mathrm{state}}^{[\mathrm{sub}]}| \;\ge\; 1 - \varepsilon_{\mathrm{state}}.$

(iv) QIV.
$\exists\, \delta_\star \in (\varepsilon_{\mathrm{law}}, 1) : \; |\Lambda_{\mathrm{law}}^{[\mathrm{sub}]}(\vec{\ell})| \;\le\; 1 - \delta_\star$ *uniformly on* $\mathfrak{P}$, $\qquad |\Lambda_{\mathrm{state}}^{[\mathrm{sub}]}| \;\ge\; 1 - \varepsilon_{\mathrm{state}}.$

Proof sketch. Combine the tolerance definition of $\Lambda_{\mathrm{law}}^{[\mathrm{sub}]}$ with Lemma 8.37. The state channel is fixed by subsystem order parameters/correlators using the same plateau tolerances. QIII vs. QIV is determined by *decay vs. persistent gap* of the law deficit under anisotropic coarse–graining. □

Corollary 8.39 (IR healing ⇒ QIII (restatement of eq. (7.746)))**.** *Let* X *be any symmetry sector read on the bulk plateau* $\mathfrak{P}$ *with witnesses* $\big(\Lambda_{\mathrm{law}}^{[X]}, \Lambda_{\mathrm{state}}^{[X]}\big)$ *as in Definition 7.35. If the explicit* X*-breaking deformations* $\{\epsilon_k \mathcal{O}_k\}$ *are RG-irrelevant—so that*

$$1 - |\Lambda_{\mathrm{law}}^{[X]}(\ell)| = \mathcal{O}\Big(\ell^{\Delta - (d+z)}\Big) \quad \text{with } \Delta > d+z \qquad (\textit{cf. eq. } (7.746)),$$

and the state is X*-symmetric on* $\mathfrak{P}$ *(i.e.* $\Lambda_{\mathrm{state}}^{[X]} = 1$*), then the quadrant assignment given by Definition 7.685 is* **QIII***.*

Proof sketch. By eq. (7.746), the law deficit decays to zero under IR dilation, hence $\Lambda_{\mathrm{law}}^{[X]}(\ell) \to 1$. With $\Lambda_{\mathrm{state}}^{[X]} = 1$, the pair falls into the "law heals, state symmetric" cell, i.e. **QIII**, by Definition 7.685.

Corollary 8.40 (BRST reduction yields gauge-safe quadrants)**.** (Restatement of Propositions 6.276 and 7.644 and Theorem 7.646.) *Let* $\mathcal{H}_{\mathrm{phys}} = H^0(Q_{\mathrm{BRST}})$ *be the physical cohomology*

and let $\left(\Lambda_{\mathrm{law}}^{[\mathrm{gauge}]}, \Lambda_{\mathrm{state}}^{[\mathrm{gauge}]}\right)$ *be the visibility pair of Definition 7.35 evaluated* after *projection to* $\mathcal{H}_{\mathrm{phys}}$ *and band-limiting by* $P_x(\lambda_{\mathrm{UV}}) \circ K_{\mathrm{IR}}^{(\ell)}$. *Then the quadrant assignment*

$$\mathrm{Quad}\left(\Lambda_{\mathrm{law}}^{[\mathrm{gauge}]}, \Lambda_{\mathrm{state}}^{[\mathrm{gauge}]}\right)$$

is independent of gauge-fixing choices and coincides across all admissible gauges in the band-limited theory. Moreover, "gauge SSB" does not occur: only BRST-invariant global *remnants can furnish* $\Lambda_{\mathrm{state}}^{[\mathrm{gauge}]} = 0$ *(state SSB), while local gauge redundancy cannot move the system between quadrants.*

Proof sketch. By Proposition 6.276, the visibilities defined on $\mathcal{H}_{\mathrm{phys}}$ are BRST-safe and gauge-parameter independent; by Proposition 7.644, spectral data used to compute them are Nielsen-stable; by Theorem 7.646, explicit deformations shift poles/residues analytically without introducing gauge dependence. Hence the decision rule of Definition 7.685 yields a gauge-invariant quadrant label. Finally, Elitzur/BRST implies local gauge symmetries do not spontaneously break; any SSB signal arises only from BRST-invariant (global) symmetry content, so gauge-fixing cannot induce a quadrant change.

8.1.10 Higher-Forms and Non-Invertible Symmetries

Higher-form and non-invertible symmetries generalize the familiar notion of global symmetry in two complementary directions. A p-form global symmetry acts on extended p-dimensional operators $W(M_p)$ rather than pointlike fields, and is generated by a conserved $(p{+}1)$-form current $J^{(p+1)}$ whose integral over codimension-$(p{+}1)$ surfaces implements the symmetry action [102, 112]. Coupling such a symmetry to a background $(p{+}1)$-form gauge field B_{p+1} makes the associated Ward identities and possible anomalies manifest at the level of the effective action, in close analogy with the familiar $p = 0$ case [82, 91]. In this setting, the *law* predicate $\Lambda_{\mathrm{law}}^{[\mathcal{S}]}$ tests the exactness of the higher-form conservation law $\mathrm{d}{*}J^{(p+1)} = 0$ and the invariance of the BRST–clean effective action under background gauge transformations $B_{p+1} \mapsto B_{p+1} + \mathrm{d}\Lambda_p$, while the *state* predicate $\Lambda_{\mathrm{state}}^{[\mathcal{S}]}$ encodes the large-distance behaviour of $W(M_p)$ in terms of perimeter- versus area-law scaling, using τ_{p+1} and π_p to distinguish unbroken from spontaneously broken higher-form symmetry phases [102, 112].

Non-invertible (categorical) symmetries provide a further generalization in which the symmetry is implemented not by group-like operators but by topological defect operators $\{D_a\}$ obeying fusion and junction rules encoded in $\mathcal{D}_{\mathcal{S}}$ [102, 112]. In this case, the "law'' channel is captured by the topological invariance and fusion consistency of the defect network on the certified plateau $\mathfrak{P}$: $\Lambda_{\mathrm{law}}^{[\mathcal{S}]}$ is set to 1 precisely when the expected topological relations among the D_a hold in the band-limited, BRST–projected effective theory. The "state'' channel probes whether these defects act trivially in the infrared on local operators separated from the defect worldvolume. This is quantified by $\Delta_a(L)$, which measures the maximal change in correlators induced by inserting a defect $D_a(\gamma)$ of size L away from a local observable. Vanishing $\Delta_a(L)$ as $L \to \infty$ signals a non-invertible symmetry that remains unbroken in the state sector, whereas a nonzero $\limsup_{L\to\infty} \Delta_a(L)$ indicates spontaneous breaking of the non-invertible structure at large scales [102, 112]. In this way, the predicates $\Lambda_{\mathrm{law}}^{[\mathcal{S}]}$ and $\Lambda_{\mathrm{state}}^{[\mathcal{S}]}$ extend the law/state visibility framework to higher-form and non-invertible symmetries within the same BRST–safe, plateau– stabilized diagnostic scheme.

Definition 8.41 (Predicates for a symmetry object $\mathcal{S}$)**.** Let $\mathcal{S}$ be either:

- a p-form global symmetry with conserved $(p{+}1)$-form current $J^{(p+1)}$ and charged p-dimensional operators $W(M_p)$, or
- a non-invertible (categorical) symmetry with topological defect set $\{D_a\}$ and fusion/junction data $\mathcal{D}_{\mathcal{S}}$.

After band-limiting and BRST projection to $\mathcal{H}_{\text{phys}}$, define:

$$\Lambda_{\text{law}}^{[\mathcal{S}]} := \begin{cases} \mathbf{1}\Big[\, \mathrm{d}{*}J^{(p+1)} = 0 \text{ and } \delta S[B_{p+1}] = 0 \\ \qquad \text{for background } B_{p+1} \,\Big], & \mathcal{S} = p\text{-form}, \\ \mathbf{1}\Big[\, \text{defect topological invariance \& fusion of } \{D_a\} \\ \qquad \text{hold on } \mathfrak{P} \,\Big], & \mathcal{S} = \text{non-invertible}. \end{cases}$$

$$\Lambda_{\text{state}}^{[\mathcal{S}]} := \begin{cases} 1, & \mathcal{S} = p\text{-form and } \tau_{p+1} > 0 \text{ and } \pi_p = 0 \quad \text{(unbroken)}, \\ 0, & \mathcal{S} = p\text{-form and } \tau_{p+1} = 0 \text{ and } \pi_p > 0 \quad \text{(SSB)}, \\ 1, & \mathcal{S} = \text{non-invertible and } \forall a:\ \lim_{L\to\infty} \Delta_a(L) = 0, \\ 0, & \mathcal{S} = \text{non-invertible and } \exists a:\ \limsup_{L\to\infty} \Delta_a(L) > 0. \end{cases}$$

$$\tau_{p+1} := \lim_{\ell\to\infty} \inf_{\text{size}(M_p)\sim\ell} \frac{-\log|\langle W(M_p)\rangle|}{\text{Vol}_{p+1}(\text{Fill}(M_p))}, \qquad \pi_p := \lim_{\ell\to\infty} \inf_{\text{size}(M_p)\sim\ell} \frac{-\log|\langle W(M_p)\rangle|}{\text{Vol}_p(M_p)}.$$

with $\Delta_a(L) := \sup_{\text{size}(\gamma)\sim L} \big|\langle D_a(\gamma)\,\mathcal{O}\rangle - \langle\mathcal{O}\rangle\big|$ for any local $\mathcal{O}$ supported away from the loop γ.

Here $\text{Fill}(M_p)$ *denotes a choice of minimal* $(p{+}1)$*-chain with* $\partial\,\text{Fill}(M_p) = M_p$ *(used to define the area-law volume).*

8.1.11 Anomaly Witnesses

Anomalies provide a structural obstruction to exact symmetry at the quantum level: even when the classical action is invariant, the path–integral measure or regularized effective action may transform nontrivially under a global symmetry X [82, 91]. This obstruction manifests in the X–Ward identity as a nonvanishing anomaly density $\mathcal{A}_X$, whose presence is protected by 't Hooft anomaly–matching and therefore cannot be removed by RG flow or local counterterms unless an appropriate inflow mechanism cancels it [102, 112]. After BRST projection to $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$, the anomalous Ward identity takes the form

$$\nabla_\mu J_X^\mu \;=\; \mathcal{A}_X^{\text{net}} \;+\; \sum_k \epsilon_k\, \langle \mathcal{S}_{k,X}^{(\lambda_{\text{UV}})} \rangle_{\mathbf{\Psi}},$$

where $\mathcal{A}_X^{\text{net}}$ is the anomaly remaining after adding all admissible inflow and counterterms, and the sum over $\mathcal{S}_{k,X}^{(\lambda_{\text{UV}})}$ represents explicit symmetry–breaking operators. The quantity $\mathcal{A}_X^{\text{net}}$ is therefore the intrinsic quantum obstruction to exact X–symmetry: it vanishes if and only if the effective action is fully X–invariant at the quantum level.

Within the visibility framework, the law predicate $\Lambda^{[X]}_{\text{law}}$ cleanly separates two distinct sources of symmetry violation. The RG–scaling of explicit symmetry–breaking operators is captured by the usual power–law behaviour $1 - |\Lambda^{[X]}_{\text{law}}(\ell)| = \mathcal{O}(\ell^{\Delta-(d+z)})$ when all relevant breakings are RG–irrelevant [112]. In contrast, a nonzero anomaly density $\mathcal{A}^{\text{net}}_X \neq 0$ produces a *persistent* deficit in the X–Ward identity: unlike irrelevant operators, this contribution cannot decay under coarse–graining, and pins the law channel to a symmetry–asymmetric classification [102]. Thus QIII corresponds to the situation $\mathcal{A}^{\text{net}}_X = 0$ with explicit breakings healing in the IR, while QIV corresponds either to a genuine anomaly or to explicit breakings that remain non-negligible on the plateau.

The state predicate $\Lambda^{[X]}_{\text{state}}$ is independent of this quantum obstruction and is evaluated through symmetry–charged order parameters or symmetry defects in the usual fashion [82, 91]. Accordingly, the full quadrant assignment for X combines the (anomaly–sensitive) law visibility with the state visibility: if the state sector also breaks X, the pair $(\Lambda^{[X]}_{\text{law}}, \Lambda^{[X]}_{\text{state}})$ lies in the "law asym, state asym" quadrant of the canonical decision rule. In this manner, anomalies integrate naturally into the BRST–safe, plateau–stabilized visibility framework, providing a sharp diagnostic for persistent quantum violations of global symmetries.

Corollary 8.42 (Anomaly ⇒ persistent law asymmetry unless net inflow cancels)**.** *Fix a global symmetry X and evaluate the X–Ward identity on the plateau $\mathfrak{P}$ after BRST projection:*

$$\nabla_\mu J^\mu_X \;=\; \mathcal{A}^{\text{net}}_X \;+\; \sum_k \epsilon_k \left\langle \mathcal{S}^{(\lambda_{\text{UV}})}_{k,X} \right\rangle_{\mathbf{\Psi}},$$

where $\mathcal{A}^{\text{net}}_X$ is the net *quantum anomaly density (including possible inflow/counterterm contributions so that $\mathcal{A}^{\text{net}}_X = 0$ iff the effective action is strictly X-invariant at the quantum level). Then the law channel for X is classified as:*

$$\begin{cases} \textit{QIII}, & \textit{if } \mathcal{A}^{\text{net}}_X = 0 \;\; \textit{and} \;\; 1 - |\Lambda^{[X]}_{\text{law}}(\ell)| = \mathcal{O}\big(\ell^{\Delta-(d+z)}\big) \;\; \textit{(explicit breakings heal)}, \\ \textit{QIV}, & \textit{if } \mathcal{A}^{\text{net}}_X \neq 0 \;\; \textit{(persistent anomaly)} \;\; \textit{or the explicit deficit persists on } \mathfrak{P}. \end{cases}$$

The state channel is read independently via Definition 7.35; if the state also breaks X, the pair $\left(\Lambda^{[X]}_{\text{law}}, \Lambda^{[X]}_{\text{state}}\right)$ lies in the "law asym, state asym" cell by Definition 7.685.

Intuition. In this setup, nonzero $\mathcal{A}^{\text{net}}_X$ is a *quantum* law deficit that cannot "fade out" under RG ('t Hooft matching). QIII reflects IR healing of *explicit* breakings once the net anomaly is zero; otherwise the anomaly pins the law channel to QIV.

8.1.12 Non-equilibrium (Floquet/Lindblad) Witnesses

Non-equilibrium quantum dynamics introduces qualitatively new mechanisms for symmetry preservation and symmetry breaking that do not appear in stationary Hamiltonian settings. Periodically driven (Floquet) systems and open (Lindblad) quantum systems furnish two prominent classes [396, 448, 449, 112, 102]. In both cases, symmetry properties must be assessed directly at the level of the one-period propagator $U(T)$ (Floquet) or the Liouvillian generator $\mathcal{L}$ (open), rather than at the Hamiltonian level. The present formulation places these systems on the same footing as the equilibrium sectors by working on a certified plateau $\mathfrak{P}$

and projecting to the BRST-physical subspace $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$, ensuring gauge-parameter independence and well-defined Ward-type constraints [82, 91]. BRST-closed IR/UV tags guarantee that all operator norms, symmetry channels, and steady-state diagnostics remain stable under admissible coarse-graining, exactly as in the equilibrium sectors [112, 102].

In the Floquet setting, law symmetry is captured by the invariance of the one-period propagator under the symmetry action U_X, quantified by the normalized defect $\|U_X^\dagger U(T) U_X - U(T)\|_{\mathfrak{P}}$. A vanishing defect within tolerance expresses that the effective stroboscopic evolution respects the symmetry to the same standard used elsewhere in the quadrant scheme. State symmetry, by contrast, probes whether an extremal periodic steady state $\rho_\star(t)$ remains invariant under the same symmetry operation throughout the drive cycle, measured in trace distance. Spontaneous breaking in the non-equilibrium steady state therefore appears as symmetry violation at the level of the cycle-resolved density matrix rather than the underlying generator. This distinction mirrors the law/state separation used in equilibrium QFT, where the Ward identity and the vacuum sector provide independent criteria [82, 91].

Open quantum systems governed by a Liouvillian $\mathcal{L}$ require an analogous construction. Law symmetry is diagnosed by the commutator defect $\|\mathcal{L} \circ \mathcal{U}_X - \mathcal{U}_X \circ \mathcal{L}\|_{\diamond,\mathfrak{P}}$, which reduces, for continuous symmetries, to the approximate conservation condition $\|\mathcal{L}^\dagger(Q_X)\|_{\mathfrak{P}} \ll 1$. State symmetry tests whether an extremal steady state ρ_∞ satisfies $U_X \rho_\infty U_X^\dagger = \rho_\infty$ up to tolerance. In this way, the non-equilibrium sector inherits the full BRST-safe, plateau-stabilized visibility structure of the equilibrium theory, permitting the same fourfold quadrant classification [112, 102].

This formulation also accommodates discrete time-translation symmetry as a special choice of X, treating time-crystalline phases as spontaneous breaking in the *state* channel of the corresponding non-equilibrium witness, consistently with [396, 448, 449]. This aligns with the general principle that order parameters (or extremal steady states in the Lindblad/Floquet setting) provide the state-sector verdict, while invariance of the effective evolution operator or Liouvillian provides the law-sector verdict, in full analogy with the symmetries of equilibrium quantum field theory [82, 91].

Definition 8.43 (Non-equilibrium witnesses (Floquet/Lindblad; tolerance form))**.** Work on a certified (bulk or boundary) plateau $\mathfrak{P}$ and in the physical sector $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ with BRST-closed scale tags. Let $\varepsilon_{\mathrm{law}}, \varepsilon_{\mathrm{state}} \in (0,1)$ be the plateau tolerances (cf. the plateau criterion). We write sector tags as superscripts and reserve subscripts for $\Lambda_{\mathrm{law}}, \Lambda_{\mathrm{state}}$.

Floquet (period T). Let $U(T)$ be the one-period propagator and U_X the unitary implementing the symmetry X on $\mathcal{H}_{\mathrm{phys}}$. Define the *law* visibility via the normalized invariance defect

$$\Lambda_{\mathrm{law}}^{[\mathrm{Floquet};X]} := 1 - \frac{\left\| U_X^\dagger U(T) U_X - U(T) \right\|_{\mathfrak{P}}}{c_{\mathrm{F}}}, \qquad 0 \le \Lambda_{\mathrm{law}}^{[\mathrm{Floquet};X]} \le 1,$$

where $\|\cdot\|_{\mathfrak{P}}$ is a plateau-uniform, BRST-safe operator norm and $c_{\mathrm{F}} > 0$ a fixed reference (e.g. $c_{\mathrm{F}} = 2$ for the operator norm on unitaries). Law symmetry on $\mathfrak{P}$ means $|\Lambda_{\mathrm{law}}^{[\mathrm{Floquet};X]}| \ge 1 - \varepsilon_{\mathrm{law}}$, equivalently $\|U_X^\dagger U(T) U_X - U(T)\|_{\mathfrak{P}} \le \varepsilon_{\mathrm{law}}\, c_{\mathrm{F}}$. Let $\rho_\star(t)$ be an *extremal* periodic steady state ($\rho_\star(t+T) = \rho_\star(t)$) selected by the protocol. Define the *state* predicate by trace-distance

tolerance:

$$\sup_{t\in[0,T)} \tfrac{1}{2}\Big\| U_X \rho_\star(t) U_X^\dagger - \rho_\star(t) \Big\|_1 \begin{cases} \le \varepsilon_{\mathrm{state}}, & \text{state symmetric on } \mathfrak{P}, \\ > \varepsilon_{\mathrm{state}}, & \text{state SSB of } X \text{ on } \mathfrak{P}, \end{cases}$$

(stroboscopic testing at $t = nT$ is a weaker but admissible alternative when continuous monitoring is unavailable).

Lindblad (open). Let $\mathcal{L}$ be the Liouvillian and $\mathcal{U}_X(\cdot) := U_X(\cdot)U_X^\dagger$ the symmetry channel. Define the *law* visibility by the superoperator commutator defect

$$\Lambda_{\mathrm{law}}^{[\mathrm{Lindblad};X]} := 1 - \frac{\big\| \mathcal{L}\circ\mathcal{U}_X - \mathcal{U}_X\circ\mathcal{L} \big\|_{\diamond,\mathfrak{P}}}{c_{\mathrm{L}}}, \qquad 0 \le \Lambda_{\mathrm{law}}^{[\mathrm{Lindblad};X]} \le 1,$$

with a plateau-uniform diamond-norm bound and fixed $c_{\mathrm{L}} > 0$. Law symmetry means $|\Lambda_{\mathrm{law}}^{[\mathrm{Lindblad};X]}| \ge 1 - \varepsilon_{\mathrm{law}}$. For continuous $U(1)$, an equivalent test is the (approximate) conserved charge condition $\|\mathcal{L}^\dagger(Q_X)\|_{\mathfrak{P}} \le \varepsilon_{\mathrm{law}}\, c_Q$. Let ρ_∞ be an extremal steady state ($\mathcal{L}\rho_\infty = 0$). Define the *state* predicate by

$$\tfrac{1}{2}\Big\| U_X \rho_\infty U_X^\dagger - \rho_\infty \Big\|_1 \begin{cases} \le \varepsilon_{\mathrm{state}}, & \text{state symmetric on } \mathfrak{P}, \\ > \varepsilon_{\mathrm{state}}, & \text{state SSB of } X \text{ on } \mathfrak{P}. \end{cases}$$

All statements are understood modulo BRST-exact improvements; norms are plateau-uniform.

Corollary 8.44 (Quadrants out of equilibrium (tolerance form))**.** *With the witnesses of Definition 8.43, the canonical decision rule yields the same four classes on* $\mathfrak{P}$*:*

(i) QI.
$|\Lambda_{\mathrm{law}}^{[\cdot;X]}| \ge 1 - \varepsilon_{\mathrm{law}}$ *and* $|\Lambda_{\mathrm{state}}^{[\cdot;X]}| \ge 1 - \varepsilon_{\mathrm{state}}$.

(ii) QII.
$|\Lambda_{\mathrm{law}}^{[\cdot;X]}| \ge 1 - \varepsilon_{\mathrm{law}}$ *and* $|\Lambda_{\mathrm{state}}^{[\cdot;X]}| \le \varepsilon_{\mathrm{state}}$.

(iii) QIII.
$1 - |\Lambda_{\mathrm{law}}^{[\cdot;X]}(\ell)| \le C\,\ell^{-\beta}$ *as* $\ell \to \infty$, *and* $|\Lambda_{\mathrm{state}}^{[\cdot;X]}| \ge 1 - \varepsilon_{\mathrm{state}}$.

(iv) QIV.
There exists $\delta_\star \in (\varepsilon_{\mathrm{law}}, 1)$ *such that* $|\Lambda_{\mathrm{law}}^{[\cdot;X]}| \le 1 - \delta_\star$ *uniformly on* $\mathfrak{P}$, *and* $|\Lambda_{\mathrm{state}}^{[\cdot;X]}| \ge 1 - \varepsilon_{\mathrm{state}}$.

Remarks. (i) If multiple extremal periodic/steady states exist, the readout is performed on the experimentally selected one (protocol dependence); tolerance semantics ensures reproducibility on $\mathfrak{P}$. (ii) Choosing X to be *discrete time translation* classifies time-crystalline order in the same grid (SSB in the *state* channel of X), consistently with [396, 448, 449].

8.1.13 Universal Quadrant Coverage

Theorem 8.45 (Universal quadrant coverage and invariance for sector-conditioned predicates)**.** *Let $\mathscr{S}$ be an admissible sector in the sense of Definition 8.9, realised on a certified plateau $\mathfrak{P}$ (bulk) or $\mathfrak{P}_\partial$ (boundary), with BRST–closed scale tags $P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}$ and sector–appropriate witnesses $\Lambda^{[\mathscr{S}]}_{\mathrm{law}}, \Lambda^{[\mathscr{S}]}_{\mathrm{state}}$. Fix plateau tolerances $\varepsilon_{\mathrm{law}}, \varepsilon_{\mathrm{state}} \in (0,1)$ (cf. Section 6.12.4).*

Define the sector–conditioned binary predicates *by*

$$\chi^{[\mathscr{S}]}_{\mathrm{law}} := \mathbf{1}\Big[\,|\Lambda^{[\mathscr{S}]}_{\mathrm{law}}| \geq 1-\varepsilon_{\mathrm{law}}\,\Big], \qquad \chi^{[\mathscr{S}]}_{\mathrm{state}} := \mathbf{1}\Big[\,|\Lambda^{[\mathscr{S}]}_{\mathrm{state}}| \geq 1-\varepsilon_{\mathrm{state}}\,\Big],$$

and the sector decision map $\mathrm{Quad}^{[\mathscr{S}]}$ *by Definition 7.685, with the QIII/QIV refinement: either the law deficit $d^{[\mathscr{S}]}_{\mathrm{law}} := 1 - |\Lambda^{[\mathscr{S}]}_{\mathrm{law}}|$* heals *under the sector's IR dilation (QIII) or it is* uniformly bounded away from 0 *(QIV).*

Then, on the relevant plateau:

(i) Exclusive, exhaustive partition up to thresholds

$\mathrm{Quad}^{[\mathscr{S}]}$ *partitions the admissible configurations of $\mathscr{S}$ into four disjoint classes* QI to QIV*, and their union is exhaustive up to threshold sets*

$$\mathfrak{T}_{\mathrm{law}} := \Big\{|\Lambda^{[\mathscr{S}]}_{\mathrm{law}}| = 1-\varepsilon_{\mathrm{law}}\Big\}, \quad \mathfrak{T}_{\mathrm{state}} := \Big\{|\Lambda^{[\mathscr{S}]}_{\mathrm{state}}| = 1-\varepsilon_{\mathrm{state}}\Big\},$$

which are negligible under the plateau regularity bounds of Proposition 6.298.

(ii) Cohomological/gauge-fermion invariance

$\mathrm{Quad}^{[\mathscr{S}]}$ *is invariant under ζ–exact gauge–fixing deformations and under gauge–parameter scans: for all admissible $\Psi \mapsto \Psi + \{\hat{Q}_{\mathrm{BRST}}, B\}$,*

$$\partial_\xi \,\mathrm{Quad}^{[\mathscr{S}]} = 0\,, \qquad \mathrm{Quad}^{[\mathscr{S}]}[\Psi] = \mathrm{Quad}^{[\mathscr{S}]}[\Psi + \{\hat{Q}_{\mathrm{BRST}}, B\}],$$

by Propositions 6.264 and 6.276 and Corollary 6.265.

(iii) Sector dilation and QIII/QIV decision

Let r denote the sector's IR dilation parameter: $r = \ell$ (bulk), $r = L$ (boundary), $r = \min(\vec{\ell})$ (anisotropic), or the appropriate coarse–graining variable (e.g. stroboscopic window for Floquet, cf. Definition 8.43). Then

$$\textit{QIII} \iff \exists\, \beta > 0,\ C < \infty :\ d^{[\mathscr{S}]}_{\mathrm{law}}(r) \leq C\, r^{-\beta}\ (r \to \infty),$$

$$\textit{QIV} \iff \exists\, \delta_\star \in (\varepsilon_{\mathrm{law}}, 1) :\ d^{[\mathscr{S}]}_{\mathrm{law}}(r) \geq \delta_\star \textit{ uniformly on } \mathfrak{P}.$$

(For concrete sectors this reduces to the previously proved criteria: hydro Lemma 8.22 and Proposition 8.23; boundary Definition 8.31 and Proposition 8.32; subsystem/fracton Lemma 8.37 and Proposition 8.38; SUSY/fermion/HS/p–form via Proposition 6.277 and the sector blocks.)

(iv) Cone/functorial stability

$\mathrm{Quad}^{[\mathscr{S}]}$ *is unchanged by (i) admissible changes of tags $P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}$ within the certified band (Corollary 6.265), (ii) holosymmetric lifts and sector functors (p–form duality, etc.; Propositions 6.257 and 8.16 and Corollary 6.259), and (iii) direct sums/triadic decompositions (Proposition 6.257); thus the diagnosis does not alter causal cones.*

Proof. **Step 1 (Well-defined binary predicates).** By Proposition 6.264 the sector witnesses are evaluated in $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ and are BRST/Nielsen–invariant under changes of Ψ. On a certified plateau (Section 6.12.4), the norms are uniform and the visibilities are stable within tolerances (Corollary 6.265). Hence $\chi_{\mathrm{law}}^{[\mathscr{S}]}, \chi_{\mathrm{state}}^{[\mathscr{S}]} \in \{0,1\}$ are well defined modulo the threshold sets $\mathfrak{T}_{\mathrm{law/state}}$.

Step 2 (Exclusivity and exhaustiveness). The decision map (Definition 7.685) is the Cartesian product $\{0,1\} \times \{0,1\} \to \{\mathbf{QI}, \mathbf{QII}, \mathbf{QIII}, \mathbf{QIV}\}$ with the QIII/QIV refinement determined by the asymptotic law–deficit test. Exclusivity follows since the images are disjoint. Exhaustiveness holds because every point lies in exactly one preimage unless it belongs to $\mathfrak{T}_{\mathrm{law}} \cup \mathfrak{T}_{\mathrm{state}}$. By the oscillation/finite–difference bounds (Proposition 6.298), these threshold sets are negligible (knife–edges of measure zero under smooth parameter variations).

Step 3 (QIII vs QIV by sector dilation). Let $d_{\mathrm{law}}^{[\mathscr{S}]}(r) := 1 - |\Lambda_{\mathrm{law}}^{[\mathscr{S}]}(r)|$. For sectors whose explicit breakings are RG–irrelevant, the Duhamel/RG estimates and sector–specific healing lemmas yield $d_{\mathrm{law}}^{[\mathscr{S}]}(r) \le C r^{-\beta}$ (bulk Proposition 6.277; hydro Proposition 8.23; boundary Proposition 8.32; subsystem Lemma 8.37). Conversely, if some relevant/marginal explicit breaking survives, Proposition 6.277 supplies a uniform lower bound $\delta_\star > 0$ on $d_{\mathrm{law}}^{[\mathscr{S}]}$ (persistent deficit), which is plateau–uniform by Remark 6.254. This exactly realises the QIII/QIV dichotomy.

Step 4 (Cohomological/gauge invariance). By Proposition 6.276 and Proposition 6.264, all sector witnesses and their channel/sector reductions are invariant under ζ–exact deformations and gauge–parameter scans; therefore $\chi_{\mathrm{law}}^{[\mathscr{S}]}, \chi_{\mathrm{state}}^{[\mathscr{S}]}$ and the asymptotic test are invariant, and so is $\mathrm{Quad}^{[\mathscr{S}]}$.

Step 5 (Cone/functorial stability). Holosymmetric lifts are functorial and cone–preserving (Proposition 6.257 and Corollary 6.259), and p–form duality preserves witnesses (Proposition 8.16); thus transport along these maps cannot change the quadrant. Variations of tags within the certified band preserve plateau bounds (Corollary 6.265), so the decision is stable.

Step 6 (Law–state exhaustion link). Finally, the sector statement is a direct corollary of law–state exhaustion on $\mathfrak{P}$ (Theorem 8.1) applied to the sector–conditioned predicates, with the QIII/QIV refinement supplied by Step 3. This completes the proof. □

What the theorem asserts. Fix any admissible sector $\mathscr{S}$ (bulk, boundary, hydrodynamic, fermionic, SUSY, p-form, higher–spin, subsystem/fracton, Floquet/Lindblad, *etc.*) in the sense of Definition 8.9 and certify its plateau $\mathfrak{P}$ (or $\mathfrak{P}_\partial$) by the plateau criterion and bounds (Section 6.12.4, Proposition 6.298, and Corollary 6.265). On that plateau, the two BRST–safe, cone–preserving visibilities

$$\Lambda_{\mathrm{law}}^{[\mathscr{S}]}, \qquad \Lambda_{\mathrm{state}}^{[\mathscr{S}]}$$

constructed in the sector–conditioned arenas (Section 8.1 and Definition 6.275) tell us everything needed to assign the quadrant for *that sector*. Plugging them into the canonical decision rule yields a *two–bit* label (QI–QIV) (Definition 7.685 and Remark 7.686) that is (i) *exclusive*—no configuration lands in two quadrants at once—and (ii) *exhaustive up to thresholds*— the only undecidable cases sit at the tolerance boundaries of the plateau (knife–edges or exactly marginal/healing loci, cf. Theorem 8.1). Within each sector, the QIII/QIV split is decided by the sector's IR dilation (bulk $\ell \to \infty$, boundary $L \to \infty$,

anisotropic $\min(\vec{\ell}) \to \infty$, stroboscopic coarse–graining in Floquet, *etc.*; see Theorem 6.253, Remark 6.254, and Sections 6.12.3 and 6.12.5): if the law–deficit heals under that dilation it means that we are in QIII; if it stays uniformly nonzero we are in QIV. Because the witnesses are *sector–conditioned* and computed in $\mathcal{H}_{\mathrm{phys}}$, the statement is universal across sectors and gauge choices, by BRST/cohomological invariance (Propositions 6.264 and 6.276); there is *no fifth quadrant* once the sector and plateau are fixed.

Operational reading (how to use it). *Step 1: certify the arena.* Choose the sector $\mathscr{S}$ and certify its plateau $\mathfrak{P}$ (or $\mathfrak{P}_\partial$) so that readouts are stable within tolerances and tags are BRST–closed, using Section 6.12.4, Propositions 6.264 and 6.298, and Corollary 6.265. *Step 2: compute the two numbers.* Evaluate $(\Lambda^{[\mathscr{S}]}_{\mathrm{law}}, \Lambda^{[\mathscr{S}]}_{\mathrm{state}})$ with the sector's natural probes as defined in the sector blocks (e.g. goldstino vs super–Higgs for SUSY, Definition 8.12 and Proposition 8.14; diffusion vs sound for hydrodynamics, Definition 8.21 and Proposition 8.29; edge modes on $\partial\mathcal{X}$, Definition 8.31 and Proposition 8.32; dualized scalar/vector data for p–forms, Proposition 8.16; anisotropic Ward residuals for subsystem symmetries, Definition 8.36 and Proposition 8.38; higher–spin and fermionic sectors via Definition 8.18 and Proposition 6.277; non–equilibrium via Definition 8.43 and Corollary 8.44). *Step 3: apply the grid.* If $|\Lambda^{[\mathscr{S}]}_{\mathrm{law}}| \approx 1$ (within $\varepsilon_{\mathrm{law}}$), then we are in QI or QII and the state channel decides, according to the canonical rule (Definition 7.685 and Theorem 6.231). If $|\Lambda^{[\mathscr{S}]}_{\mathrm{law}}| < 1$, perform the large–window test in that arena, using the relevant dilation parameter and scaling estimates (Theorem 6.253, Remark 6.254, Lemmas 8.22 and 8.37, Propositions 8.29 and 8.38, Definition 8.43, and Corollary 8.44): healing $\Rightarrow$ QIII, persistent gap $\Rightarrow$ QIV. *Step 4: record the label.* The resulting $\mathrm{Quad}^{[\mathscr{S}]}$ is BRST/Nielsen–safe and does not alter cones or mix channels, by the invariance and functoriality results (Propositions 6.257, 6.264 and 6.276 and Corollary 6.259).

Proof strategy at a glance (why it is true). First, on a certified plateau the visibilities are stable and BRST/Nielsen–invariant (Propositions 6.264 and 6.276 and Corollary 6.265), so their thresholded versions define well–posed binary predicates (law/state) as in Definitions 6.275, 7.35 and 8.9. Their Cartesian product is a two–bit partition; exclusivity is immediate, and exhaustiveness fails only on the negligible threshold sets where a tolerance is saturated, controlled by the plateau bounds (Proposition 6.298 and Remark 6.254) and the law–state exhaustion theorem (Theorem 8.1). Second, sector dilation distinguishes healing vs persistence of the law–deficit by power–law bounds or uniform lower gaps proved in the sector blocks (bulk and fermionic/SUSY/HS/p–form sectors via Theorems 6.253 and 7.630, Corollary 7.631, and Proposition 6.277; hydrodynamic, boundary, subsystem and non–equilibrium sectors via Propositions 8.29, 8.32 and 8.38 and Corollary 8.44). Third, functoriality and cone preservation guarantee that holosymmetric lifts, dualities, and admissible retagging cannot change the quadrant—diagnostics *measure* the sector; they do not move it (Propositions 6.257 and 8.16 and Corollaries 6.259 and 6.265). Finally, this sector statement is the sector–conditioned instance of the law–state exhaustion principle (Theorems 8.1 and 8.45), completing the argument.

Mental model (why no surprises occur). Think of each sector as a lab with its own calibrated apparatus (its witnesses and IR windowing, Sections 6.12.3, 6.12.5, 8.1 and 8.2). We bring in a sample (the configuration), put it on the certified plateau (the plateau, Section 6.12.4 and Proposition 6.298), and read two dials: *law* and *state*. Dials near "1" mean symmetry, away from "1" mean a deficit, as encoded in the visibility semantics (Definition 7.35 and Remarks 6.254 and 7.686). Turn the IR knob: if the law dial drifts back to "1" then we call it emergent (QIII); if it refuses, it's explicit (QIV), in line with the scaling criteria of Theorem 6.253, Propositions 6.277, 8.29 and 8.38, and Corollary 8.44. Because the apparatus is BRST–clean and cone–preserving (Propositions 6.257, 6.264 and 6.276 and Corollary 6.259), we can trust that the reading reflects the sample, not the gauge choice or the diagnostic realized reference configuration itself. Across all such labs, the same two–dial logic applies—hence the universal four–cell map of Theorems 6.231 and 8.45.

Corollary 8.46 (Operational checklist)**.** *To classify any admissible sector $\mathscr{S}$ using sector–conditioned predicates on a certified plateau:*

(i) Fix the arena and certify the plateau

Choose bulk $\mathfrak{P}$, boundary $\mathfrak{P}_\partial$, or anisotropic $K_{\mathrm{IR}}^{(\vec{\ell})}$ (fracton/subsystem). Pick BRST–closed tags $P_x(\lambda_{\mathrm{UV}})$, K_{IR} and tolerances $\varepsilon_{\mathrm{law}}, \varepsilon_{\mathrm{state}} \in (0,1)$. Verify plateau stability via Section 6.12.4 and Proposition 6.298 and gauge–parameter independence via Propositions 6.264 and 6.276 and Corollary 6.265.

(ii) Instantiate sector witnesses

Construct

$$(\Lambda_{\mathrm{law}}^{[\mathscr{S}]}, \Lambda_{\mathrm{state}}^{[\mathscr{S}]})$$

with BRST–safe, plateau–uniform norms using the sector recipe (e.g. Definitions 8.21, 8.31 and 8.36 and the fermionic/SUSY/p–form/HS blocks). Record the reference normalizations.

(iii) Apply the canonical decision rule (law/state)

Declare

$$\chi_{\mathrm{law}}^{[\mathscr{S}]} = \mathbf{1}\big[|\Lambda_{\mathrm{law}}^{[\mathscr{S}]}| \geq 1 - \varepsilon_{\mathrm{law}}\big], \qquad \chi_{\mathrm{state}}^{[\mathscr{S}]} = \mathbf{1}\big[|\Lambda_{\mathrm{state}}^{[\mathscr{S}]}| \geq 1 - \varepsilon_{\mathrm{state}}\big],$$

and map to QI/QII provisionally by Definition 7.685 when $\chi_{\mathrm{law}}^{[\mathscr{S}]} = 1$.

(iv) [Decide QIII vs QIV by large–window scaling in the chosen arena]Let r be the sector's IR dilation parameter (bulk $r = \ell$, boundary $r = L$, anisotropic $r = \min(\vec{\ell})$, or the appropriate coarse–graining variable in non–equilibrium). With $d_{\mathrm{law}}^{[\mathscr{S}]}(r) := 1 - |\Lambda_{\mathrm{law}}^{[\mathscr{S}]}(r)|$:

$$\textit{QIII} \iff \exists\, \beta > 0,\ C < \infty :\ d_{\mathrm{law}}^{[\mathscr{S}]}(r) \leq C\, r^{-\beta}\ (r \to \infty),$$

$$\textit{QIV} \iff \exists\, \delta_\star \in (\varepsilon_{\mathrm{law}}, 1) :\ d_{\mathrm{law}}^{[\mathscr{S}]}(r) \geq \delta_\star \textit{ uniformly on the plateau.}$$

(Use the sector criteria: hydro Lemma 8.22 and Proposition 8.23, boundary Proposition 8.32, subsystem Lemma 8.37 and Proposition 8.38, and the bulk blocks via Proposition 6.277.)

(v) Threshold handling
If either visibility lies within an indifference band around the tolerance (i.e. $\left||\Lambda_\bullet^{[\mathscr{S}]}| - (1-\varepsilon_\bullet)\right| \le \eta$*), flag the case as* threshold/undecided*; tighten tags or enlarge the window and re–certify the plateau (Proposition 6.298).*

(vi) Cross–checks (optional but recommended)
Validate the quadrant with the equivalent diagnostics: scale–tagged Ward statements and light–sector spectrum/residues (Theorem 7.687); discrepancies indicate a failed plateau certification.

(vii) Record provenance
Report $\mathrm{Quad}^{[\mathscr{S}]}$*, the tolerance pair* $(\varepsilon_{\mathrm{law}}, \varepsilon_{\mathrm{state}})$*, the dilation law used for QIII/QIV, and the certified band* B*; by Corollaries 6.259 and 6.265 and Proposition 6.257 this label is BRST/Nielsen–safe and cone–stable.*

Intuition. The grid (QI–QIV) is universal; only the witnesses and the arena change. The law channel is decided first (exact within tolerance vs. deficit), and if a deficit exists, the large–window scaling discriminates emergent (QIII) from persistent (QIV). The state channel then selects QI vs QII when the law is (approximately) symmetric. All steps are BRST–clean and plateau–stable, so the outcome is reproducible and gauge–parameter independent.

8.2 The Synthesis of Laws and States

Definition 8.47 (Paraconsistent context and consequence)**.** A *paraconsistent context* is a tuple

$$\mathcal{C} \;=\; \big\langle \mathcal{L}_{\mathcal{C}},\ \vdash_{\mathrm{pc}},\ \Lambda_{\mathcal{C}},\ \mathrm{Adm}_{\mathcal{C}} \big\rangle,$$

where $\mathcal{L}_{\mathcal{C}}$ is a formal language, $\vdash_{\mathrm{pc}}$ is a structural consequence relation on $\mathcal{L}_{\mathcal{C}}$ that *rejects explosion* (i.e. there exists an inconsistent Γ with $\Gamma \nvdash_{\mathrm{pc}} \psi$ for some ψ), $\Lambda_{\mathcal{C}}$ is the set of admissible loci, and $\mathrm{Adm}_{\mathcal{C}}$ are the admissible inference/reporting practices. We assume *Adjunction* ($\varphi, \psi \vdash_{\mathrm{pc}} \varphi \wedge \psi$), *Conjunction Elimination*, *Weakening*, and *Cut*, but not *Explosion* (ECQ) (*cf.* Section 1.2.6).

Definition 8.48 (Non-explosion witness)**.** A *non-explosion witness* for $\vdash_{\mathrm{pc}}$ is data

$$\Pi_{\mathrm{pc}} \;=\; \langle \mathfrak{s},\ v,\ \mathsf{D} \rangle$$

such that $\mathfrak{s}$ is a semantics (algebraic/relational/valuational) for $\mathcal{L}_{\mathcal{C}}$, v is a valuation/interpretation on $\mathfrak{s}$, and D is a set of designated truth-values with the properties: (i) there is φ with $v(\varphi)$ and $v(\neg\varphi)$ both designated (inconsistency), but (ii) there exists ψ with $v(\psi)$ *not* designated (non-triviality). When convenient we instantiate $\mathfrak{s}$ as a four-valued LP/FDE matrix with values $\{\mathbf{T}, \mathbf{F}, \mathbf{B}, \mathbf{N}\}$ and designated $\{\mathbf{T}, \mathbf{B}\}$ (*cf.* Synthesis' paraconsistent clauses in Section 1.2.3 and Definition 1.3).

Definition 8.49 (Logical propagation)**.** Fix a paraconsistent base $\langle \mathcal{L}_{\mathcal{C}}, \vdash_{\mathrm{pc}}, \Pi_{\mathrm{pc}} \rangle$ with valuation v on $\mathfrak{s}$ and designated set D. We say that *logical propagation* holds from a set of seed formulas $\Gamma \subseteq \mathcal{L}_{\mathcal{C}}$ when the following two mechanisms determine the status of all formulas built from Γ by the allowed connectives and rules:

(Semantics) Truth-functionality in FDE/LP: for all χ, α, β,

$$v(\neg\chi) \text{ is determined by } v(\chi), \qquad v(\alpha \wedge \beta) \text{ is determined by } v(\alpha), v(\beta),$$

with designatedness preserved (and $\mathbf{B}$ fixed under $\neg$ and closed under $\wedge$ when both inputs are $\mathbf{B}$).

(Proof theory) Premise-soundness of $\vdash_{\mathrm{pc}}$ w.r.t. Π_{pc}: if $\Gamma \vdash_{\mathrm{pc}} \psi$ and $v(\gamma) \in \mathsf{D}$ for all $\gamma \in \Gamma$, then $v(\psi) \in \mathsf{D}$.

In particular, under *Simultaneous Synthesis* with $v(\varphi_L) = v(\varphi_S) = \mathbf{B}$, logical propagation yields

$$v(\neg\varphi_L) = v(\neg\varphi_S) = \mathbf{B}, \qquad v(\mathrm{I}), v(\mathrm{II}), v(\mathrm{III}), v(\mathrm{IV}) \in \mathsf{D},$$

and (by the witness property of Π_{pc}) joint non-triviality of $\{\mathrm{I}, \mathrm{II}, \mathrm{III}, \mathrm{IV}\}$.

Definition 8.50 (Paired dialectical defenses)**.** Fix a formula φ and locus $\lambda \in \Lambda_{\mathcal{C}}$. We write

$$\mathcal{J}^{\pm}_{\mathrm{A}}(\varphi), \qquad \mathcal{J}^{\pm}_{\mathrm{N}}(\varphi; \lambda)$$

for *paired Aristotelian* ($+$ for thesis/affirmation; $-$ for antithesis/denial) and *paired Nyāya* arguments (five-membered scheme; allows perception, inference, testimony, and anupalabdhi), each being well-formed derivations in $\vdash_{\mathrm{pc}}$ whose conclusions are, respectively, φ and $\neg\varphi$. We say the pair is *co-tenable* iff their joint adoption does not yield triviality in $\vdash_{\mathrm{pc}}$ (witnessed by Π_{pc}). (*Cf.* status machinery and corner-selection alignment in Section 1.2.6 and Proposition 1.15.)

Definition 8.51 (Synthesis, cf. Section 1.2.3 and Definition 1.3)**.** Let $\mathcal{C}$ be paraconsistent (Def. 8.47) and $\lambda \in \Lambda_{\mathcal{C}}$. The *Synthesis* of φ at $\langle \mathcal{C}, \lambda \rangle$ is the structured act

$$\mathbf{SynTh}(\varphi; \mathcal{C}, \lambda) := \Big\langle \underbrace{\varphi}_{\text{Asti/Sat}}, \underbrace{\neg\varphi}_{\text{Nāsti/Asat}}, \underbrace{\mathcal{J}^{+}_{\mathrm{A}}(\varphi), \mathcal{J}^{-}_{\mathrm{A}}(\varphi)}_{\text{Aristotelian pair}}, \underbrace{\mathcal{J}^{+}_{\mathrm{N}}(\varphi; \lambda), \mathcal{J}^{-}_{\mathrm{N}}(\varphi; \lambda)}_{\text{Nyāya pair}}, \underbrace{\Pi_{\mathrm{pc}}}_{\text{non-explosion witness}} \Big\rangle$$

subject to: (i) *Co-affirmation*: both φ and $\neg\varphi$ are adopted at the same locus λ (same sense/symmetry); (ii) *Scope hygiene*: no equivocation on φ or λ; (iii) *Paraconsistent discipline*: Π_{pc} attests non-explosion for $\{\varphi, \neg\varphi\}$ (*cf.* Section 1.2.6); (iv) *Paired justification*: the four arguments are valid in $\vdash_{\mathrm{pc}}$ and co-tenable (*cf.* Proposition 1.15).

Lemma 8.52 (Non-explosion of Synthesis)**.** *If* $\mathbf{SynTh}(\varphi; \mathcal{C}, \lambda)$ *holds, then for some* ψ *we have* $\big(\{\varphi, \neg\varphi\} \vdash_{\mathrm{pc}} \psi\big)$*. In particular* $\vdash_{\mathrm{pc}}$ *is non-trivial in the presence of the contradiction* $\varphi \wedge \neg\varphi$ *(*cf. *Section 1.2.6).*

Proof. Immediate from Π_{pc} (Def. 8.48): choose ψ with $v(\psi) \notin \mathsf{D}$ while $v(\varphi), v(\neg\varphi) \in \mathsf{D}$. Soundness of $\vdash_{\mathrm{pc}}$ w.r.t. $\mathfrak{s}$ then prevents derivability of ψ from $\{\varphi, \neg\varphi\}$. □

Object-level predicates for symmetry. Fix a kinematical symmetry group G and an action $T : G \curvearrowright \mathsf{Desc}$ on *law descriptors* (e.g. Lagrangians/Hamiltonians), and $\alpha : G \curvearrowright \mathsf{A}$ on an observable algebra A. Introduce in $\mathcal{L}_{\mathcal{C}}$ the predicates:

$$\mathrm{Sym}(L) :\Longleftrightarrow \forall g \in G \;\; T_g(L) = L, \tag{8.11}$$

$$\mathrm{Break}(L) :\Longleftrightarrow \exists\, \mathcal{O}_{\mathrm{break}}\; \exists g \in G \;\; T_g(\mathcal{O}_{\mathrm{break}}) \neq \mathcal{O}_{\mathrm{break}} \;\wedge\; \mathrm{OccursIn}(\mathcal{O}_{\mathrm{break}}, L), \tag{8.12}$$

$$\mathrm{Sym}(w;\lambda) :\Longleftrightarrow \forall g \in G \;\; w\restriction_\lambda \circ \alpha_g \;=\; w\restriction_\lambda, \tag{8.13}$$

$$\mathrm{Break}(w;\lambda) :\Longleftrightarrow \exists\, \mathcal{O} \in \mathsf{A}\; \exists g \in G \;\; \alpha_g(\mathcal{O}) \neq \mathcal{O} \;\wedge\; w\restriction_\lambda(\mathcal{O}) \neq 0. \tag{8.14}$$

(*Cf.* duality lifts and status dualization Definition 1.11 and Section 1.2.6 for transport across dualized predicates.) Gauge "symmetry" is treated as redundancy: all four predicates are understood via *gauge-invariant* representatives in A and Desc.

Definition 8.53 (Law-level proposition)**.** Let $\varphi_L := \mathrm{Sym}(L)$ be the formula "The Law is Symmetric" (with fixed G and fixed sense of invariance as in (8.11)).

Lemma 8.54 (Scope hygiene for the Law)**.** *If φ_L and $\neg\varphi_L$ are to be co-affirmed under* **SynTh***, then both must refer to the* same *group G, the* same *action $T : G \curvearrowright \mathsf{Desc}$, and the* same *descriptor L. In particular,* $\mathrm{Break}(L)$ *in* (8.12) *must exhibit an operator $\mathcal{O}_{\mathrm{break}}$ with* $\mathrm{OccursIn}(\mathcal{O}_{\mathrm{break}}, L)$ *and $T_g(\mathcal{O}_{\mathrm{break}}) \neq \mathcal{O}_{\mathrm{break}}$ for some $g \in G$ (not merely in a different descriptor or under different symmetry data).*

Proof. Assume $\mathbf{SynTh}(\varphi_L;\mathcal{C})$ with $\varphi_L := \mathrm{Sym}(L)$ (Def. 8.53). By Definition 8.51(i)–(ii), there must be derivations in $\vdash_{\mathrm{pc}}$ of φ_L and of its *literal* negation $\neg\varphi_L$, and the two sides must concern the *same* proposition (anti-equivocation).

Unfolding (8.11), φ_L abbreviates $\forall g \in G\; T_g(L) = L$ for the fixed data (G, T, L). The object-language witness for $\neg\varphi_L$ is $\mathrm{Break}(L)$, defined in (8.12) by the existence of $\mathcal{O}_{\mathrm{break}}$ and $g \in G$ such that $T_g(\mathcal{O}_{\mathrm{break}}) \neq \mathcal{O}_{\mathrm{break}}$ and $\mathrm{OccursIn}(\mathcal{O}_{\mathrm{break}}, L)$. This predicate is *well-formed at* (G, T, L) and, at that fixed data, entails $\neg\,\mathrm{Sym}(L)$.

If any component were changed—say $G \to G'$, $T \to T'$, or $L \to L'$—the breaking predicate would be $\mathrm{Break}(L')$, which (even if true) entails $\neg\,\mathrm{Sym}(L')$, *not* $\neg\,\mathrm{Sym}(L)$. The pair $\{\varphi_L, \neg\varphi_L\}$ would then equivocate across respects, violating Definition 8.51(ii), rather than forming a genuine glut about the *same* proposition.

Therefore, to co-affirm $\{\varphi_L, \neg\varphi_L\}$ under **SynTh**, both sides must refer to the same G, the same T, and the same L. In particular, the witness in (8.12) must supply $\mathcal{O}_{\mathrm{break}}$ with $\mathrm{OccursIn}(\mathcal{O}_{\mathrm{break}}, L)$ and $T_g(\mathcal{O}_{\mathrm{break}}) \neq \mathcal{O}_{\mathrm{break}}$ for some $g \in G$. □

Proposition 8.55 (Synthesis of the Law)**.** *Assume in $\mathcal{C}$ that both* $\mathrm{Sym}(L)$ *and* $\mathrm{Break}(L)$ *are derivable and that there exist co-tenable paired defenses*

$$\mathcal{J}^+_{\mathrm{A}}(\varphi_L),\; \mathcal{J}^-_{\mathrm{A}}(\varphi_L), \qquad \mathcal{J}^+_{\mathrm{N}}(\varphi_L),\; \mathcal{J}^-_{\mathrm{N}}(\varphi_L),$$

together with a non-explosion witness Π_{pc} with a valuation v such that $v(\varphi_L) = \mathbf{B}$. Then

$$\mathbf{SynTh}(\varphi_L;\mathcal{C})$$

holds. In Tetralemma terms, this is the both-selection *status for the dyad associated to φ_L (*cf. *Proposition 1.15).*

Proof. By hypothesis, $\mathrm{Sym}(L)$ and $\mathrm{Break}(L)$ are both derivable in $\mathcal{C}$. Fix the *same* symmetry data (G, T) and the *same* descriptor L as in (8.11)–(8.12) (scope hygiene). In the object language, $\mathrm{Sym}(L)$ asserts $T_g(L) = L$ for all $g \in G$ (8.11), while $\mathrm{Break}(L)$ asserts the existence of an operator $\mathcal{O}_{\mathrm{break}}$ and an element $g \in G$ such that $T_g(\mathcal{O}_{\mathrm{break}}) \neq \mathcal{O}_{\mathrm{break}}$ and $\mathrm{OccursIn}(\mathcal{O}_{\mathrm{break}}, L)$ (8.12). Hence, at fixed (G, T, L) we have $\mathrm{Break}(L) \Rightarrow \neg\,\mathrm{Sym}(L)$, and therefore derivations of φ_L and $\neg\varphi_L$ in $\mathcal{L}_{\mathcal{C}}$.

The paired Aristotelian and Nyāya defenses

$$\mathcal{J}^{+}_{\mathrm{A}}(\varphi_L),\ \mathcal{J}^{-}_{\mathrm{A}}(\varphi_L), \qquad \mathcal{J}^{+}_{\mathrm{N}}(\varphi_L),\ \mathcal{J}^{-}_{\mathrm{N}}(\varphi_L)$$

are, by assumption, valid in $\vdash_{\mathrm{pc}}$ and co-tenable. Together with the non-explosion witness Π_{pc} and the valuation condition $v(\varphi_L) = \mathbf{B}$, these satisfy items (i)–(iv) of Definition 8.51. Therefore the structured act $\mathbf{SynTh}(\varphi_L; \mathcal{C})$ holds by direct application of Definition 8.51. Finally, Lemma 8.52 yields the "without triviality" clause. □

Law-level Synthesis, in plain terms. Fix the proposition $\varphi_L := \mathrm{Sym}(L)$. Applying the Synthesis modality to φ_L yields the structured act $\mathbf{SynTh}(\varphi_L; \mathcal{C})$, i.e., a disciplined *co-commitment* that the Law is both symmetric and asymmetric in the *same* context $\mathcal{C}$ and for the *same* symmetry data (G, descriptor L). Formally this is exactly the content of Definitions 1.3 and 8.51 instantiated at the law level, and it is captured as a theorem in Proposition 8.55.

What makes the "both" legitimate. The co-affirmation clause of Definition 8.51 requires that $\mathrm{Sym}(L)$ and its denial refer to an identical sense of invariance (same G, same T, same L); this is the "scope hygiene" that blocks equivocating shifts of sense or locus. In a classical, explosive setting, adopting $\mathrm{Sym}(L)$ together with its negation would trivialize the theory. By contrast, under a paraconsistent consequence relation $\vdash_{\mathrm{pc}}$ the joint adoption is stable: explosion fails, and the contradiction does not entail everything (Definitions 8.47 and 8.48 and Lemma 8.52; see also [453, 454, 455]). Thus, Synthesis replaces collapse with a controlled, non-trivial "both/and".

How the evidence is organized. Synthesis at the law level is not a license for ambiguity; it is *warranted* by paired defenses (Definition 8.50). On the positive side, $\mathcal{J}^{+}_{\mathrm{A}}(\varphi_L)$ and $\mathcal{J}^{+}_{\mathrm{N}}(\varphi_L)$ adduce reasons for $\mathrm{Sym}(L)$ (e.g., invariance postulates, selection rules, Ward identities; cf. (8.11)). On the negative side, $\mathcal{J}^{-}_{\mathrm{A}}(\varphi_L)$ and $\mathcal{J}^{-}_{\mathrm{N}}(\varphi_L)$ exhibit definite breaking structure (an operator $\mathcal{O}_{\mathrm{break}}$ and element $g \in G$ with $T_g(\mathcal{O}_{\mathrm{break}}) \neq \mathcal{O}_{\mathrm{break}}$, and $\mathcal{O}_{\mathrm{break}}$ occurring in L; cf. (8.12)). A familiar heuristic is the decomposition $L = L_{\mathrm{sym}} + \varepsilon\,\mathcal{O}_{\mathrm{break}}$ with $T_g(L_{\mathrm{sym}}) = L_{\mathrm{sym}}$ and $T_g(\mathcal{O}_{\mathrm{break}}) \neq \mathcal{O}_{\mathrm{break}}$: within $\vdash_{\mathrm{pc}}$, this supports both $\mathrm{Sym}(L)$ (structural/approximate invariants) and $\mathrm{Break}(L)$ (explicit breaking), delivering $v(\varphi_L) = \mathbf{B}$ as required in Proposition 8.55.

UpShot. The law-level Synthesis therefore characterizes a regime that is neither purely symmetric nor merely explicitly broken. It encodes a *justified duality*: the Law carries simultaneously valid symmetric and asymmetric aspects, held together by paraconsistent discipline and paired epistemic warrants. In the Tetralemma vocabulary, this is precisely the "both-selection'' status for the law-dyad (Proposition 1.15), and by Lemma 8.52 it does so *without* logical collapse.

Definition 8.56 (State-level proposition)**.** Let $\varphi_S := \mathrm{Sym}(w;\lambda)$ be "The State is Symmetric at λ" (with fixed G and fixed locus λ), as in (8.13).

Lemma 8.57 (Scope hygiene for the State)**.** *If φ_S and $\neg\varphi_S$ are to be co-affirmed under* **SynTh***, then both must refer to the* same *group G, the* same *action α, the* same *state w, and the* same *locus λ. In particular,* $\mathrm{Break}(w;\lambda)$ *in* (8.14) *must exhibit an order parameter or symmetry-violating expectation value evaluated at λ (not merely at a different locus).*

Proof. Assume $\mathbf{SynTh}(\varphi_S;\mathcal{C},\lambda)$ with $\varphi_S := \mathrm{Sym}(w;\lambda)$. By Definition 8.51(i)–(ii), there must be derivations in $\vdash_{\mathrm{pc}}$ of φ_S and of its *literal* negation $\neg\varphi_S$, and the two sides must concern the *same* proposition (anti-equivocation).

Unfolding (8.13), φ_S abbreviates

$$\forall g \in G \;\; w\restriction_\lambda \circ\alpha_g \;=\; w\restriction_\lambda .$$

Thus $\neg\varphi_S$ is the denial of invariance for this very data (G,α,w,λ). The object-language witness we use for $\neg\varphi_S$ is $\mathrm{Break}(w;\lambda)$, defined in (8.14) by the existence of $\mathcal{O}\in\mathsf{A}$ and $g\in G$ such that $\alpha_g(\mathcal{O})\neq\mathcal{O}$ and $w\restriction_\lambda(\mathcal{O})\neq 0$. This predicate is *well-formed at* (G,α,w,λ) and, at that fixed data, entails $\neg\,\mathrm{Sym}(w;\lambda)$.

If any component were changed—say $G\to G'$, $\alpha\to\alpha'$, $w\to w'$, or $\lambda\to\lambda'$—the breaking predicate would be $\mathrm{Break}(w';\lambda')$, which (even if true) entails $\neg\,\mathrm{Sym}(w';\lambda')$, *not* $\neg\,\mathrm{Sym}(w;\lambda)$. The pair $\{\varphi_S,\neg\varphi_S\}$ would then be an equivocation across respects/locations, violating Definition 8.51(ii), rather than a genuine glut about the *same* proposition.

Therefore, to co-affirm $\{\varphi_S,\neg\varphi_S\}$ under **SynTh**, both sides must refer to the same G, the same α, the same w, and the same locus λ. In particular, the witness in (8.14) must evaluate $w\restriction_\lambda$ at that very λ. □

Proposition 8.58 (Synthesis of the State)**.** *Assume* $\mathrm{Sym}(w;\lambda)$ *and* $\mathrm{Break}(w;\lambda)$ *are both derivable in $\mathcal{C}$, with paired defenses*

$$\mathcal{J}^{+}_{\mathrm{A}}(\varphi_S),\ \mathcal{J}^{-}_{\mathrm{A}}(\varphi_S),\qquad \mathcal{J}^{+}_{\mathrm{N}}(\varphi_S;\lambda),\ \mathcal{J}^{-}_{\mathrm{N}}(\varphi_S;\lambda),$$

and a non-explosion witness Π_{pc} giving $v(\varphi_S)=\mathbf{B}$. Then

$$\mathbf{SynTh}(\varphi_S;\mathcal{C},\lambda)$$

holds. Hence the State is both *symmetric and asymmetric at λ without triviality (*cf. *Section 1.2.6).*

Proof. By hypothesis, $\mathrm{Sym}(w;\lambda)$ and $\mathrm{Break}(w;\lambda)$ are both derivable in $\mathcal{C}$, and Lemma 8.57 fixes the *same* G, α, w, and λ. In the object language, $\mathrm{Sym}(w;\lambda)$ asserts invariance of the restricted state under all $g\in G$ (8.13), while $\mathrm{Break}(w;\lambda)$ asserts the existence of $\mathcal{O}\in\mathsf{A}$ and $g\in G$ with $\alpha_g(\mathcal{O})\neq\mathcal{O}$ and a nonvanishing expectation at the same locus, $w\restriction_\lambda(\mathcal{O})\neq 0$ (8.14). Hence, at fixed (G,α,w,λ), $\mathrm{Break}(w;\lambda)$ entails $\neg\,\mathrm{Sym}(w;\lambda)$, so we have derivations of φ_S and $\neg\varphi_S$ in $\mathcal{L}_{\mathcal{C}}$.

The paired Aristotelian and Nyāya defenses

$$\mathcal{J}^{+}_{\mathrm{A}}(\varphi_S),\ \mathcal{J}^{-}_{\mathrm{A}}(\varphi_S),\qquad \mathcal{J}^{+}_{\mathrm{N}}(\varphi_S;\lambda),\ \mathcal{J}^{-}_{\mathrm{N}}(\varphi_S;\lambda)$$

are, by assumption, valid in $\vdash_{\mathrm{pc}}$ and co-tenable. Together with the non-explosion witness Π_{pc} and the valuation condition $v(\varphi_S) = \mathbf{B}$, these satisfy items (i)–(iv) of Definition 8.51. Therefore the structured act $\mathbf{SynTh}(\varphi_S; \mathcal{C}, \lambda)$ holds by direct application of Definition 8.51. Finally, Lemma 8.52 yields the "without triviality" clause. □

Remark 8.59 (Physical instantiation note)**.** In finite volume (or with symmetry-restoring terms) there can be coherent superpositions of symmetry-broken vacua supporting $\mathrm{Break}(w; \lambda)$ alongside symmetric diagnostics; in strict thermodynamic limits, distinct vacua often become superselection sectors. The logical construction remains valid; the physical witness for $\mathrm{Break}(w; \lambda)$ may then be mixture/sector structure rather than literal coherence.

State-level Synthesis, in plain terms. Fix the proposition $\varphi_S := \mathrm{Sym}(w; \lambda)$ (Definition 8.56). Applying the Synthesis modality to φ_S yields the structured act $\mathbf{SynTh}(\varphi_S; \mathcal{C}, \lambda)$: a disciplined *co-commitment* that, at the same locus λ and for the same symmetry data (G, α), the State is both symmetric and asymmetric. Formally, this is Definition 8.51 instantiated at the state level and realized by Proposition 8.58.

What makes the "both" legitimate. The co-affirmation and scope-hygiene clauses of Definition 8.51 (items (i)–(ii)) together with Lemma 8.57 require that $\mathrm{Sym}(w; \lambda)$ and its denial refer to the *same* group G, action α, state w, and locus λ. Thus the "both" is not an equivocation across respects or locations. In a classical (explosive) setting, adopting φ_S and $\neg\varphi_S$ would trivialize the theory; under a paraconsistent consequence relation $\vdash_{\mathrm{pc}}$, backed by a non-explosion witness Π_{pc}, the joint adoption is logically stable (Definitions 8.47 and 8.48 and Lemma 8.52).

How the evidence is organized. Synthesis at the state level is *warranted* by paired dialectical defenses (Definition 8.50). On the positive side, arguments $\mathcal{J}^+_{\mathrm{A}}(\varphi_S)$ and $\mathcal{J}^+_{\mathrm{N}}(\varphi_S; \lambda)$ support $\mathrm{Sym}(w; \lambda)$ via symmetry-invariant diagnostics (e.g., Ward-type constraints and symmetric correlators), as formalized in (8.13). On the negative side, $\mathcal{J}^-_{\mathrm{A}}(\varphi_S)$ and $\mathcal{J}^-_{\mathrm{N}}(\varphi_S; \lambda)$ exhibit symmetry-breaking response at the *same* locus, such as a nonzero order parameter $w\restriction_\lambda(\mathcal{O}) \neq 0$ for some $\mathcal{O} \in \mathsf{A}$, as in (8.14). Co-tenability of these paired defenses ensures that adopting both sides does not collapse $\vdash_{\mathrm{pc}}$.

UpShot. The state-level Synthesis transcends the simple dichotomy "invariant vs. non-invariant." It formally accommodates states with a dialectical character in which symmetric and symmetry-breaking descriptions are *simultaneously* applicable and jointly warranted. This is precisely the content of Proposition 8.58: at fixed λ, the State can be both symmetric and asymmetric without triviality, a reading that also matches the physical nuances noted in the remark on finite volume and superselection (Proposition 8.58 and the subsequent remark).

Proposition 8.60 (Duality transport)**.** *Let $\mathbf{D}_{\mathcal{C}}$ be the Axiom-of-Duality involution acting functorially on formulas and arguments within $\mathcal{C}$. If $\mathbf{SynTh}(\varphi; \mathcal{C}, \lambda)$ holds, then $\mathbf{SynTh}(\mathbf{D}_{\mathcal{C}}\varphi; \mathcal{C}, \lambda)$ holds, with paired defenses transported by $\mathbf{D}_{\mathcal{C}}$ and the same Π_{pc} (*cf. *Section 1.2.6 and Definition 1.11).*

Proof. Assume $\mathbf{SynTh}(\varphi; \mathcal{C}, \lambda)$. By Definition 8.51, there exist paired defenses

$$\mathcal{J}^+_{\mathrm{A}}(\varphi),\ \mathcal{J}^-_{\mathrm{A}}(\varphi), \qquad \mathcal{J}^+_{\mathrm{N}}(\varphi; \lambda),\ \mathcal{J}^-_{\mathrm{N}}(\varphi; \lambda)$$

which are derivations in $\vdash_{\mathrm{pc}}$ with conclusions φ and $\neg\varphi$, are *co-tenable*, and are backed by the same non-explosion witness Π_{pc} (so that adopting both does not trivialize $\vdash_{\mathrm{pc}}$).

By hypothesis, $\mathbf{D}_{\mathcal{C}}$ is an Axiom-of-Duality *involution* acting *functorially* on formulas and arguments in $\mathcal{C}$. Hence: (i) $\mathbf{D}_{\mathcal{C}}$ maps derivations in $\vdash_{\mathrm{pc}}$ to derivations in $\vdash_{\mathrm{pc}}$ and commutes with the logical operations, so $\mathbf{D}_{\mathcal{C}}(\neg\varphi) = \neg\mathbf{D}_{\mathcal{C}}\varphi$; (ii) $\mathbf{D}_{\mathcal{C}}$ transports the Aristotelian/Nyāya arguments to

$$\mathbf{D}_{\mathcal{C}}\mathcal{J}^{+}_{\mathrm{A}}(\varphi),\ \mathbf{D}_{\mathcal{C}}\mathcal{J}^{-}_{\mathrm{A}}(\varphi), \qquad \mathbf{D}_{\mathcal{C}}\mathcal{J}^{+}_{\mathrm{N}}(\varphi;\lambda),\ \mathbf{D}_{\mathcal{C}}\mathcal{J}^{-}_{\mathrm{N}}(\varphi;\lambda),$$

which are well-formed defenses in $\vdash_{\mathrm{pc}}$ with conclusions $\mathbf{D}_{\mathcal{C}}\varphi$ and $\neg\mathbf{D}_{\mathcal{C}}\varphi$ at the *same* locus λ (duality lifts preserve the object-level meanings and loci).

Co-tenability is preserved: if the transported pair trivialized $\vdash_{\mathrm{pc}}$, then applying the inverse (also $\mathbf{D}_{\mathcal{C}}$) would make the original pair trivial, contradicting the premise. Finally, the same Π_{pc} witnesses non-explosion after transport (duality is semantics-compatible and does not alter designatedness or the rule set of $\vdash_{\mathrm{pc}}$).

Thus items (i)–(iv) of Definition 8.51 hold for $\mathbf{D}_{\mathcal{C}}\varphi$ with the transported paired defenses and the same Π_{pc}. Therefore $\mathbf{SynTh}(\mathbf{D}_{\mathcal{C}}\varphi;\mathcal{C},\lambda)$ holds. □

Definition 8.61 (Quadrant formulas)**.** Set $\varphi_L := \mathrm{Sym}(L)$ and $\varphi_S := \mathrm{Sym}(w;\lambda)$. Define

$$\mathrm{I} := \varphi_L \wedge \varphi_S, \qquad \mathrm{II} := \varphi_L \wedge \neg\varphi_S,$$
$$\mathrm{III} := \neg\varphi_L \wedge \varphi_S, \quad \mathrm{IV} := \neg\varphi_L \wedge \neg\varphi_S.$$

These correspond to the four *cells* of the symmetry–asymmetry permutations matrix (table 6.1): (I) Unbroken, (II) SSB, (III) Emergent/Accidental, (IV) Explicit Breaking.

Remark 8.62 (Terminology hygiene: quadrants vs. tetralemma corners)**.** The labels I–IV refer to the matrix in table 6.1, not to the four modal corners $\langle \mathsf{Th}, \mathsf{AntiTh}, \mathsf{SynTh}, \mathsf{HoloTh}\rangle$ of the Tetralemma of Algebra (Definition 1.5). We reserve "Tetralemma corners" for the modal unit $\mathring{\mathbf{\Lambda}}$ and "quadrants" for the law/state matrix.

Theorem 8.63 (Quadrant co-tenability under Synthesis)**.** *Suppose* $\mathbf{SynTh}(\varphi_L;\mathcal{C})$ *and* $\mathbf{SynTh}(\varphi_S;\mathcal{C},\lambda)$ *both hold with a common witness* $\Pi_{\mathrm{pc}} = \langle \mathfrak{s}, v, \mathsf{D}\rangle$ *such that* $v(\varphi_L) = v(\varphi_S) = \mathbf{B}$. *Then each of* $\mathrm{I}, \mathrm{II}, \mathrm{III}, \mathrm{IV}$ *of Definition 8.61 is designated under* v *and hence individually adoptable in* $\vdash_{\mathrm{pc}}$. *Moreover, the set* $\{\mathrm{I}, \mathrm{II}, \mathrm{III}, \mathrm{IV}\}$ *is non-trivial: there exists* ψ *with* $\Big(\{\mathrm{I}, \mathrm{II}, \mathrm{III}, \mathrm{IV}\} \vdash_{\mathrm{pc}} \psi\Big)$ *(*cf. *Section 1.2.6).*

Proof. Assume $\mathbf{SynTh}(\varphi_L;\mathcal{C})$ and $\mathbf{SynTh}(\varphi_S;\mathcal{C},\lambda)$ with a common witness $\Pi_{\mathrm{pc}} = \langle \mathfrak{s}, v, \mathsf{D}\rangle$ such that $v(\varphi_L) = v(\varphi_S) = \mathbf{B}$, and with the usual scope hygiene (same $G, T, \alpha, L, w, \lambda$ where relevant).

Step 1: Derive the four cells *as in Definition 8.61.* By Definition 8.51, Synthesis supplies paired, co-tenable derivations in $\vdash_{\mathrm{pc}}$ of each side and its *literal* negation:

$$\varphi_L,\ \neg\varphi_L \qquad \text{and} \qquad \varphi_S,\ \neg\varphi_S.$$

Using Adjunction (and Weakening if needed), form the four conjunctions

$$\mathrm{I} := \varphi_L \wedge \varphi_S, \quad \mathrm{II} := \varphi_L \wedge \neg\varphi_S, \quad \mathrm{III} := \neg\varphi_L \wedge \varphi_S, \quad \mathrm{IV} := \neg\varphi_L \wedge \neg\varphi_S,$$

which are exactly the four *cells* in Definition 8.61. Hence each of I–IV is derivable in $\vdash_{\mathrm{pc}}$ from $\mathcal{C}$.

Step 2: Designatedness under v. In LP/FDE with $\mathsf{D} = \{\mathbf{T}, \mathbf{B}\}$, $v(\neg\chi) = \mathbf{B}$ whenever $v(\chi) = \mathbf{B}$, and $v(\chi \wedge \psi) \in \mathsf{D}$ iff $v(\chi) \in \mathsf{D}$ and $v(\psi) \in \mathsf{D}$. Since $v(\varphi_L) = v(\varphi_S) = \mathbf{B}$, we also have $v(\neg\varphi_L) = v(\neg\varphi_S) = \mathbf{B}$; thus each of I–IV evaluates to a designated value. Therefore, each cell is individually adoptable in $\vdash_{\mathrm{pc}}$.

Step 3: Joint non-triviality. By clause (ii) of Definition 8.48, choose ψ with $v(\psi) \notin \mathsf{D}$. If, for contradiction, $\{\mathrm{I}, \mathrm{II}, \mathrm{III}, \mathrm{IV}\} \vdash_{\mathrm{pc}} \psi$, then by soundness of $\vdash_{\mathrm{pc}}$ w.r.t. Π_{pc} we would have $v(\psi) \in \mathsf{D}$ whenever v designates all premises. But Step 2 gives $v(\mathrm{I}), \ldots, v(\mathrm{IV}) \in \mathsf{D}$, contradicting $v(\psi) \notin \mathsf{D}$. Hence there exists ψ with $\not\left(\{\mathrm{I}, \mathrm{II}, \mathrm{III}, \mathrm{IV}\} \vdash_{\mathrm{pc}} \psi\right)$, so the set of the four cells is non-trivial.

This establishes designatedness of each cell and joint non-triviality under the common Π_{pc}. □

Remark 8.64 (Operational witnesses)**.** When desired, the object-language predicates $\mathrm{Break}(L)$ and $\mathrm{Break}(w; \lambda)$ (at fixed symmetry data and locus) entail $\neg\varphi_L$ and $\neg\varphi_S$, respectively. They may thus serve as *witnesses* for the negated conjuncts in the cells, without altering Definition 8.61.

Aristotelian templates (Law). $\mathcal{J}^{+}_{\mathrm{A}}(\varphi_L)$: Syllogistic/axiomatic derivation of (8.11) from invariance postulates (selection rules, Ward identities). $\mathcal{J}^{-}_{\mathrm{A}}(\varphi_L)$: Negative demonstration via explicit $g \in G$ and *gauge-invariant* $\mathcal{O}_{\mathrm{break}}$ with $T_g(\mathcal{O}_{\mathrm{break}}) \neq \mathcal{O}_{\mathrm{break}}$ *and* $\mathrm{OccursIn}(\mathcal{O}_{\mathrm{break}}, L)$, entailing $\neg\,\mathrm{Sym}(L)$.

Nyāya templates (Law). $\mathcal{J}^{+}_{\mathrm{N}}(\varphi_L)$: Five-membered argument with testimony (Noether/Ward), perception (null amplitudes), and inference (closure under G).

$\mathcal{J}^{-}_{\mathrm{N}}(\varphi_L)$: Five-membered argument with perception of symmetry-violating responses, inference from RG-generated *gauge-invariant* $\mathcal{O}_{\mathrm{break}}$ with $\mathrm{OccursIn}(\mathcal{O}_{\mathrm{break}}, L)$, and testimony of anomaly constraints where applicable.

Aristotelian templates (State). $\mathcal{J}^{+}_{\mathrm{A}}(\varphi_S)$: Derive (8.13) from symmetric correlators/cluster decomposition and Ward identities. $\mathcal{J}^{-}_{\mathrm{A}}(\varphi_S)$: Reductio via exhibited *gauge-invariant* order parameter $\mathcal{O}$ with $w\restriction_{\lambda}(\mathcal{O}) \neq 0$ *and* $\alpha_g(\mathcal{O}) \neq \mathcal{O}$ for some $g \in G$.

Nyāya templates (State). $\mathcal{J}^{+}_{\mathrm{N}}(\varphi_S; \lambda)$: Perception (degenerate multiplets / forbidden transitions), testimony (symmetry constraints), and inference to (8.13). $\mathcal{J}^{-}_{\mathrm{N}}(\varphi_S; \lambda)$: Perception of symmetry-breaking response, *anupalabdhi* of symmetric nulls where expected, and inference to (8.14).

Corollary 8.65 (Unified dialectical reading)**.** *Under Theorem 8.63, the four "table quadrants" are not disjoint metaphysical kinds but* modalities *within one paraconsistent structure—fully compatible with Duality transport and the atomic prismatic unit of the Tetralemma (Proposition 8.60 and Definition 1.5).*

Proof of Corollary 8.65. By Theorem 8.63, each of the four *cells*

$$\mathrm{I} := \varphi_L \wedge \varphi_S, \quad \mathrm{II} := \varphi_L \wedge \neg\varphi_S, \quad \mathrm{III} := \neg\varphi_L \wedge \varphi_S, \quad \mathrm{IV} := \neg\varphi_L \wedge \neg\varphi_S$$

is designated under the common witness $\Pi_{\mathrm{pc}} = \langle \mathfrak{s}, v, \mathsf{D} \rangle$, and the set $\{\mathrm{I}, \mathrm{II}, \mathrm{III}, \mathrm{IV}\}$ is jointly non-trivial in $\vdash_{\mathrm{pc}}$.

Modal reading within one paraconsistent structure. Let $\mathcal{M} := \langle \mathcal{L}_{\mathcal{C}}, \vdash_{\mathrm{pc}}, \Pi_{\mathrm{pc}} \rangle$. Since each cell is a well-formed formula of $\mathcal{L}_{\mathcal{C}}$, individually adoptable (designated) yet jointly non-trivial, they function as distinct *modalities* of description inside the single structure $\mathcal{M}$ rather than as disjoint metaphysical kinds.

Compatibility with Duality transport. From $\mathbf{SynTh}(\varphi_L; \mathcal{C})$ and $\mathbf{SynTh}(\varphi_S; \mathcal{C}, \lambda)$, Proposition 8.60 gives $\mathbf{SynTh}(\mathbf{D}_{\mathcal{C}}\varphi_L; \mathcal{C})$ and $\mathbf{SynTh}(\mathbf{D}_{\mathcal{C}}\varphi_S; \mathcal{C}, \lambda)$ with the *same* Π_{pc}. Functoriality of $\mathbf{D}_{\mathcal{C}}$ ensures $\mathbf{D}_{\mathcal{C}}(\neg\chi) = \neg\mathbf{D}_{\mathcal{C}}\chi$ and $\mathbf{D}_{\mathcal{C}}(\chi \wedge \psi) = \mathbf{D}_{\mathcal{C}}\chi \wedge \mathbf{D}_{\mathcal{C}}\psi$, so $\mathbf{D}_{\mathcal{C}}$ permutes the four cells into the corresponding four cells built from $\mathbf{D}_{\mathcal{C}}\varphi_L$ and $\mathbf{D}_{\mathcal{C}}\varphi_S$, preserving designatedness under the same witness. Hence the cell-wise modal reading is stable under Duality transport.

Compatibility with the atomic prismatic unit. The Tetralemma of Algebra (Definition 1.5) provides a minimal atomic prismatic unit that is invariant under admissible dualities/reindexings. Our use of the four cells does not re-define those corners; it merely organizes object-level law/state claims as co-tenable modalities inside $\mathcal{M}$. Since admissible (prismatic) maps act on the whole unit and respect dualities, the cell-wise modalities coexist compatibly with that unit (no clash of meanings, no extra identifications imposed).

Therefore, under Theorem 8.63, the four table cells are modalities within one paraconsistent structure, and this reading is fully compatible with both Duality transport and the atomic prismatic unit. □

Remark 8.66 (Anomalies and IR exactness; tightened)**.** Quadrant III ($\neg\varphi_L \wedge \varphi_S$) licenses a state-level symmetry *with* law-level breaking. Exactness of the state symmetry may hold only at IR fixed points; away from the fixed point, dangerously-irrelevant symmetry–violating operators or 't Hooft anomaly constraints can yield symmetry-breaking responses. Formally, these subtleties live in the *paired defenses*: the positive state-side defenses $\mathcal{J}^{+}_{\mathrm{A}/N}(\varphi_S; \lambda)$ must exhibit symmetry-invariant diagnostics at the *same* locus λ (scope hygiene), while the law-side negatives $\mathcal{J}^{-}_{\mathrm{A}/N}(\varphi_L)$ must witness $\mathrm{Break}(L)$ via a gauge-invariant $\mathcal{O}_{\mathrm{break}}$ with $\mathrm{OccursIn}(\mathcal{O}_{\mathrm{break}}, L)$ and $T_g(\mathcal{O}_{\mathrm{break}}) \neq \mathcal{O}_{\mathrm{break}}$. None of this alters the *logical* content of **SynTh**: co-tenability and non-explosion are governed by $\vdash_{\mathrm{pc}}$ and the witness Π_{pc}; the anomaly/IR issues affect only the evidential content inside $\mathcal{J}^{\pm}$, not the form of the modality.

This formal construction is a rigorous and lucid expansion of the principles articulated in Section 1.2.3. It preserves the modality-theoretic intent of *Synthesis* while making explicit the semantic and inferential constraints under which "both/and" assertions are admissible.

The provided definitions (Definitions 8.47, 8.48 and 8.50) precisely specify the logical machinery required for any coherent deployment of *Synthesis*. By isolating non-explosion as a formal witness Π_{pc}, the text shows how contradictions can be stably entertained without triviality—i.e., without validating $\varphi, \neg\varphi \vdash \psi$ for arbitrary ψ—in line with standard paraconsistent semantics such as Dunn–Belnap FDE and Priest's LP [453, 454, 455].

Furthermore, the introduction of explicit object-level predicates for the symmetry of a Law and a State— $\mathrm{Sym}(L)$ and $\mathrm{Sym}(w; \lambda)$, together with their breaking counterparts—

provides a necessary bridge from the abstract modality to concrete physical propositions. The accompanying templates for paired Aristotelian and Nyāya defenses ensure that an invocation of *Synthesis* is not merely verbal but carries epistemic warrant sourced in admissible evidence and argument forms, matching the methodological demands in the physics literature where symmetry claims and their violations are tied to observable diagnostics and Ward-type constraints [91, 100].

The capstone of this development, the Quadrant Co-tenability Theorem (Theorem 8.63), is especially clarifying. It formally establishes that the four ostensibly mutually exclusive **cells of the symmetry–asymmetry matrix** can be read as co-tenable modalities when the Law and the State are each in *Synthesis*. In that setting, the classical dichotomies dissolve: every quadrant becomes individually adoptable (designated) within a single paraconsistent model while non-explosion is preserved by Π_{pc} [453, 454]. This puts on firm footing the heuristic use of the matrix as a "dashboard" rather than a partition, consistent with established symmetry/SSB practice [91, 100].

As summarized in the concluding corollary (Corollary 8.65), the quadrants are thereby understood not as disjoint metaphysical kinds but as relational facets of a unified dialectical structure governed by **Duality transport** and **prismatic morphisms of the atomic prismatic unit**.

Setup 8.67 (Simultaneous Synthesis)**.** Let $\varphi_L := \mathrm{Sym}(L)$ and $\varphi_S := \mathrm{Sym}(w;\lambda)$ (cf. object-level predicates). Assume the two Synthesis acts (cf. Section 1.2.3 and Definition 1.3)

$$\mathbf{SynTh}(\varphi_L;\mathcal{C}) \qquad \text{and} \qquad \mathbf{SynTh}(\varphi_S;\mathcal{C},\lambda)$$

hold in the same $\mathcal{C}$, at *fixed symmetry data*: for the Law, the same G and action $T : G \curvearrowright \mathsf{Desc}$; for the State, the same G, action $\alpha : G \curvearrowright \mathsf{A}$, state w, and locus λ (scope hygiene). They are witnessed by $\Pi_{\mathrm{pc}} = \langle \mathfrak{s}, v, \mathsf{D}\rangle$ with $v(\varphi_L) = v(\varphi_S) = \mathbf{B}$. Work in a sound paraconsistent base (e.g. Dunn–Belnap FDE / Priest's LP) with designated $\{\mathbf{T},\mathbf{B}\}$ [453, 454, 455]. Paraconsistent co-affirmation is the "Synthesis corner'' of the Tetralemma (Sections 1.2.3 and 1.2.6 and Definition 1.3).

Definition 8.68 (Quadrant formulas)**.** $\mathrm{I} := \varphi_L \wedge \varphi_S$, $\mathrm{II} := \varphi_L \wedge \neg\varphi_S$, $\mathrm{III} := \neg\varphi_L \wedge \varphi_S$, $\mathrm{IV} := \neg\varphi_L \wedge \neg\varphi_S$.

Lemma 8.69 (Truth-functional facts in FDE/LP)**.** *In Dunn–Belnap FDE (and hence in LP for these connectives), with values* $\{\mathbf{N},\mathbf{T},\mathbf{F},\mathbf{B}\}$ *and designated* $\{\mathbf{T},\mathbf{B}\}$*:*

(i) *If* $v(\chi) = \mathbf{B}$ *then* $v(\neg\chi) = \mathbf{B}$.

(ii) *If* $v(\alpha), v(\beta)$ *are designated then* $v(\alpha \wedge \beta)$ *is designated; moreover, if both are* $\mathbf{B}$*, then* $v(\alpha \wedge \beta) = \mathbf{B}$.

Proof. Use the standard set-of-classical-values semantics: identify each truth value with a subset of $\{T, F\}$ (so $\mathbf{N} = \varnothing$, $\mathbf{T} = \{T\}$, $\mathbf{F} = \{F\}$, $\mathbf{B} = \{T, F\}$). Negation swaps T and F; hence $\mathbf{B}$ is fixed by $\neg$, proving (i). Conjunction has $T \in v(\alpha \wedge \beta)$ iff $T \in v(\alpha)$ and $T \in v(\beta)$, and $F \in v(\alpha \wedge \beta)$ iff $F \in v(\alpha)$ or $F \in v(\beta)$ [453, 454]; thus (ii) follows, and if both inputs are $\mathbf{B}$, then both T and F are present, i.e. $v(\alpha \wedge \beta) = \mathbf{B}$. □

Theorem 8.70 (All four quadrants light up without triviality)**.** *Under Setup 8.67, each quadrant* I, II, III, IV *of Definition 8.68 is designated under* v*. Moreover, the set* $\{\mathrm{I}, \mathrm{II}, \mathrm{III}, \mathrm{IV}\}$ *is non-trivial (does not entail everything).*

Proof. Since $v(\varphi_L) = v(\varphi_S) = \mathbf{B}$, Lemma 8.69(i) gives $v(\neg\varphi_L) = v(\neg\varphi_S) = \mathbf{B}$. Apply Lemma 8.69(ii) to obtain

$$v(\mathrm{I}) = v(\varphi_L \wedge \varphi_S) = \mathbf{B}, \quad v(\mathrm{II}) = v(\varphi_L \wedge \neg\varphi_S) = \mathbf{B},$$

$$v(\mathrm{III}) = v(\neg\varphi_L \wedge \varphi_S) = \mathbf{B}, \quad v(\mathrm{IV}) = v(\neg\varphi_L \wedge \neg\varphi_S) = \mathbf{B}.$$

For non-triviality, use the witness clause in Setup 8.67: choose ψ with $v(\psi) \notin \mathsf{D}$ (e.g. $v(\psi) = \mathbf{N}$ in FDE). By *premise-soundness* of $\vdash_{\mathrm{pc}}$ w.r.t. Π_{pc}, if $\{\mathrm{I}, \mathrm{II}, \mathrm{III}, \mathrm{IV}\} \vdash_{\mathrm{pc}} \psi$, then every valuation that designates all premises (in particular v) must designate ψ, i.e. $v(\psi) \in \mathsf{D}$—a contradiction. Hence $\not\big(\{\mathrm{I}, \mathrm{II}, \mathrm{III}, \mathrm{IV}\} \vdash_{\mathrm{pc}} \psi\big)$. □

Corollary 8.71 (Reading of the matrix)**.** *The usual* 2×2 *"Symmetry–Asymmetry" table remains the correct display, but it is no longer a partition: multiple cells (indeed, all four) may be simultaneously designated. The matrix is a* projection *of an underlying paraconsistent state of information, aligning with the Tetralemma's "both" corner for Synthesis and its operational ties to the four corners (Proposition 1.15 and Section 1.2.6).*

Proof of Corollary 8.71. Under Setup 8.67 we have

$$v(\varphi_L) = v(\varphi_S) = \mathbf{B}.$$

By Lemma 8.69(i)–(ii), each cell in Definition 8.68,

$$\mathrm{I} := \varphi_L \wedge \varphi_S, \quad \mathrm{II} := \varphi_L \wedge \neg\varphi_S, \quad \mathrm{III} := \neg\varphi_L \wedge \varphi_S, \quad \mathrm{IV} := \neg\varphi_L \wedge \neg\varphi_S,$$

is designated. Hence the usual 2×2 table is *not* a partition: multiple cells—indeed, all four—can be simultaneously designated.

"Projection" claim. The matrix entries are just the truth-functional combinations of the two object-level propositions φ_L, φ_S. Thus the table displays a projection of the underlying paraconsistent information state $(v(\varphi_L), v(\varphi_S))$ into the four conjunctions in Definition 8.68. When both components are $\mathbf{B}$, every cell lights up (while non-explosion is preserved by Setup 8.67), which aligns with the "both" status of *Synthesis* and its operational ties to the Tetralemma framework (Proposition 1.15).

Therefore the matrix is the correct display—but as a *dashboard* rather than a partition. □

Remark 8.72 (Hygiene, scope, and dual transport)**.** Both Synthesis acts must be applied at fixed symmetry data: the same group G and action $T : G \curvearrowright \mathsf{Desc}$ for the Law, and the same G, action $\alpha : G \curvearrowright \mathsf{A}$, state w, and locus λ for the State (scope hygiene; cf. Section 1.2.3 and Definition 1.3). Under the Axiom of Duality, the functorial involution $\mathbf{D}_{\mathcal{C}}$ preserves *Synthesis* (Proposition 8.60), transporting paired defenses and leaving the witness Π_{pc} intact; consequently, the dashboard is stable under duality lifts (Section 1.2.6 and Definition 1.11).

Projection insight. The present construction gives a rigorous and elegant proof of the core insight that the classical matrix of possibilities is a *projection* of a more complex, unified structure (Setup 8.67, Definition 8.68, Theorem 8.70, and Corollary 8.71). In the precise sense established below, the 2×2 display is no longer a mutually exclusive case split but a view of co-tenable modalities selected by the logic of *Synthesis* (Section 1.2.3 and Definition 1.3).

Logical propagation. The argument proceeds with transparent steps. (1) *Setup.* Both the Law φ_L and the State φ_S are assumed to be in the modality of *Synthesis*, witnessed by a non-explosive semantics Π_{pc} such that $v(\varphi_L) = v(\varphi_S) = \mathbf{B}$ (designated "both") in a sound paraconsistent base (FDE/LP) (Setup 8.67 and Definition 8.48; see [453, 454, 455]). (2) *Propagation via the truth-functions.* As recorded in Lemma 8.69, $\mathbf{B}$ is a fixed point of negation and conjunction preserves designatedness in the $(\neg, \wedge)$-fragment common to FDE and LP: $v(\neg\chi) = \mathbf{B}$ whenever $v(\chi) = \mathbf{B}$; and if $v(\chi_1), v(\chi_2)$ are designated then $v(\chi_1 \wedge \chi_2)$ is designated (*moreover*, if both are $\mathbf{B}$ then $v(\chi_1 \wedge \chi_2) = \mathbf{B}$) [453, 454]. (3) *Quadrant values.* It follows immediately that each quadrant formula in Definition 8.68—$\mathrm{I} = \varphi_L \wedge \varphi_S$, $\mathrm{II} = \varphi_L \wedge \neg\varphi_S$, $\mathrm{III} = \neg\varphi_L \wedge \varphi_S$, $\mathrm{IV} = \neg\varphi_L \wedge \neg\varphi_S$—also evaluates to $\mathbf{B}$. (4) *Non-triviality.* Because $\mathbf{B}$ is designated and Π_{pc} witnesses non-explosion, all four quadrants are simultaneously adoptable without collapse: there exists some ψ with $v(\psi)$ undesignated, so $\{\mathrm{I}, \mathrm{II}, \mathrm{III}, \mathrm{IV}\} \nvdash_{\text{pc}} \psi$ (Theorem 8.70 and Definition 8.48; cf. [453, 454]).

Dashboard reading. The corollary (Corollary 8.71) captures the main conceptual payoff: the familiar 2×2 "Symmetry–Asymmetry" table ceases to function as a partition and instead becomes a *dashboard* of co-tenable modalities. This makes precise a practice that is already implicit in symmetry/SSB reasoning—namely, that law-level and state-level diagnostics can jointly constrain, but need not classically determine, a single exclusive quadrant [91, 100]. In contexts where finite-volume or regulator effects are salient, this paraconsistent reading is especially faithful to how seemingly conflicting signatures (symmetric correlators alongside symmetry-breaking responses) coexist without logical explosion.

Ontological upshot. Accordingly, the classical categories of symmetry and breaking are not static boxes but interrelated *projections* of a single richer structure whose coherence is maintained by the logic of *Synthesis*. In this reading, the 2×2 matrix functions as a dashboard of co-tenable modalities rather than a partition: law-level and state-level claims may be jointly entertained without triviality, provided the paraconsistent discipline holds (Section 1.2.6). The perspective aligns with the Tetralemma's "both" corner and is stable under admissible duality transport (Proposition 8.60). When scale changes are relevant, renormalization-group flow supplies a canonical mechanism by which descriptions and apparent symmetries shift while remaining compatible with the paraconsistent envelope [415, 403].

8.2.1 Parasymmetric Synthesis of Laws and States

Modal sectors for the anchors. For each anchor $a \in \{L, S\}$ (Law, State) fix the two-dimensional modal sector

$$\mathcal{H}^{(a)} := \operatorname{span}\left\{|e_1\rangle_a,\ |e_2\rangle_a\right\} \qquad \text{with} \qquad |e_1\rangle_a \equiv |\,\mathrm{Sym}\rangle_a,\ \ |e_2\rangle_a \equiv |\,\mathrm{Break}\rangle_a.$$

Let $\mathrm{Id}^{(a)}_{(0,1)}$ denote the identity on $\mathcal{H}^{(a)}$, and set the local "parasymmetric/anti-parasymmetric" kets

$$|+\rangle_a := \tfrac{1}{\sqrt{2}}\big(|e_1\rangle_a + |e_2\rangle_a\big), \qquad |-\rangle_a := \tfrac{1}{\sqrt{2}}\big(|e_1\rangle_a - |e_2\rangle_a\big).$$

Definition 8.73 (Local parasymmetric projectors and valuation)**.** For each $a \in \{L, S\}$ define the rank-1 projectors

$$P^{(a)}_{\mathrm{para}} := |+\rangle_a\langle+|\,, \qquad P^{(a)}_{\mathrm{antipara}} := |-\rangle_a\langle-|$$

so that $P^{(a)}_{\mathrm{para}} + P^{(a)}_{\mathrm{antipara}} = P^{(a)}_{(0,1)}$. Given a density operator ρ_a on $\mathcal{H}^{(a)}$, the *local Synthesis valuation* is

$$v^{(a)}_{\mathrm{Synthesis}}(\rho_a) \;:=\; \mathrm{Tr}\Big(P^{(a)}_{\mathrm{para}}\rho_a\Big) \in [0,1].$$

Lemma 8.74 (Elementary properties)**.** *For each anchor a:* (i) $P^{(a)}_{\mathrm{para}}P^{(a)}_{\mathrm{antipara}} = 0$ *and* $\big(P^{(a)}_{\mathrm{para}}\big)^2 = P^{(a)}_{\mathrm{para}}$ *(similarly for $P^{(a)}_{\mathrm{antipara}}$).* (ii) *For the spectral decomposition $\rho_a = \sum_k p_k\,|\psi_k\rangle\langle\psi_k|$, $v^{(a)}_{\mathrm{Synthesis}}(\rho_a) = \sum_k p_k\,|\langle+|\psi_k\rangle|^2$.* (iii) *Writing ρ_a in the $\{|e_1\rangle_a, |e_2\rangle_a\}$ basis and using $\rho_{11}+\rho_{22}=1$,*

$$v^{(a)}_{\mathrm{Synthesis}}(\rho_a) = \tfrac{1}{2}(\rho_{11}+\rho_{22}) + \mathrm{Re}(\rho_{12}) = \tfrac{1}{2} + \mathrm{Re}(\rho_{12})\,,$$

so the valuation depends on the coherence ρ_{12} *and not only on populations.*

Proof. Fix $a \in \{L,S\}$ and abbreviate $|\pm\rangle := |\pm\rangle_a$, $P_{\mathrm{para}} := P^{(a)}_{\mathrm{para}} = |+\rangle\langle+|$, $P_{\mathrm{antipara}} := P^{(a)}_{\mathrm{antipara}} = |-\rangle\langle-|$.

(i) Orthogonality and idempotency. Since $|+\rangle = \frac{1}{\sqrt{2}}(|e_1\rangle + |e_2\rangle)$ and $|-\rangle = \frac{1}{\sqrt{2}}(|e_1\rangle - |e_2\rangle)$, we have $\langle+|-\rangle = 0$. Hence

$$P_{\mathrm{para}}P_{\mathrm{antipara}} = \Big(|+\rangle\langle+|\Big)\Big(|-\rangle\langle-|\Big) = |+\rangle\,\underbrace{\langle+|-\rangle}_{=0}\,\langle-| \;=\; 0,$$

and similarly $P_{\mathrm{antipara}}P_{\mathrm{para}} = 0$. Idempotency is immediate:

$$P^2_{\mathrm{para}} = \Big(|+\rangle\langle+|\Big)\Big(|+\rangle\langle+|\Big) = |+\rangle\,\langle+|+\rangle\,\langle+| = |+\rangle\langle+| = P_{\mathrm{para}},$$

and likewise for P_{antipara}.

(ii) Spectral/convex form of the valuation. By definition $v^{(a)}_{\mathrm{Synthesis}}(\rho_a) = \mathrm{Tr}(P_{\mathrm{para}}\rho_a)$. Using the spectral decomposition $\rho_a = \sum_k p_k|\psi_k\rangle\langle\psi_k|$ and linearity of the trace,

$$v^{(a)}_{\mathrm{Synthesis}}(\rho_a) = \sum_k p_k\;\mathrm{Tr}\Big(P_{\mathrm{para}}|\psi_k\rangle\langle\psi_k|\Big) = \sum_k p_k\,\langle\psi_k|P_{\mathrm{para}}|\psi_k\rangle = \sum_k p_k\,|\langle+|\psi_k\rangle|^2.$$

This shows convexity and basis-independence of the expression.

(iii) Matrix form and coherence dependence. In the ordered basis $\{|e_1\rangle_a, |e_2\rangle_a\}$, write $\rho_a = \left(\begin{smallmatrix}\rho_{11} & \rho_{12}\\ \rho_{21} & \rho_{22}\end{smallmatrix}\right)$ with $\rho_{21} = \overline{\rho_{12}}$, $\rho_{11}, \rho_{22} \geq 0$, $\rho_{11}+\rho_{22} = 1$. The projector onto $|+\rangle$ is

$$P_{\mathrm{para}} = |+\rangle\langle+| = \tfrac{1}{2}\left(\begin{smallmatrix}1 & 1\\ 1 & 1\end{smallmatrix}\right).$$

Then

$$v^{(a)}_{\text{Synthesis}}(\rho_a) = \mathrm{Tr}(P_{\text{para}}\rho_a) = \tfrac{1}{2}\,\mathrm{Tr}\begin{pmatrix} \rho_{11}+\rho_{21} & \rho_{12}+\rho_{22} \\ \rho_{11}+\rho_{21} & \rho_{12}+\rho_{22} \end{pmatrix} = \tfrac{1}{2}\Big(\rho_{11}+\rho_{22}+\rho_{12}+\rho_{21}\Big) = \tfrac{1}{2}+\mathrm{Re}(\rho_{12}),$$

as claimed. In particular, $v^{(a)}_{\text{Synthesis}}$ depends on the *real part of the off-diagonal* element (the coherence), not only on populations.

Range check. Positivity of ρ_a implies $|\rho_{12}|^2 \le \rho_{11}\rho_{22} \le \tfrac{1}{4}(\rho_{11}+\rho_{22})^2 = \tfrac{1}{4}$. Hence $-\tfrac{1}{2} \le \mathrm{Re}(\rho_{12}) \le \tfrac{1}{2}$, so $0 \le v^{(a)}_{\text{Synthesis}}(\rho_a) \le 1$, with $v^{(a)}_{\text{Synthesis}} = 1$ (resp. 0) iff $\rho_a = |+\rangle\langle+|$ (resp. $|-\rangle\langle-|$).

Equivalent Bloch-form (optional). Let $\sigma_x = \left(\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix}\right)$. Then $P_{\text{para}} = \tfrac{1}{2}(I+\sigma_x)$ and

$$v^{(a)}_{\text{Synthesis}}(\rho_a) = \tfrac{1}{2}\Big(1+\mathrm{Tr}(\sigma_x\rho_a)\Big) = \tfrac{1}{2}+\mathrm{Re}(\rho_{12}),$$

consistent with the calculation above. □

Remark 8.75 (Coherent "both" vs. incoherent mixture)**.** If ρ_a is *diagonal* with equal populations ($\rho_{11} = \rho_{22} = \tfrac{1}{2}$) and hence zero coherence ($\rho_{12} = 0$), then $v^{(a)}_{\text{Synthesis}}(\rho_a) = \tfrac{1}{2}$. By contrast, the pure coherent superposition $|+\rangle_a$ gives $v = 1$. Thus parasymmetry separates coherent Synthesis from a mere mixture—something the purely logical **SynTh** layer cannot distinguish (Definition 8.51 and Lemma 8.52; cf. [453, 454, 455]).

Phase-resolved special case. For a pure equal-weight superposition

$$|\psi_\phi\rangle \;=\; \tfrac{1}{\sqrt{2}}\Big(|e_1\rangle_a + e^{i\phi}\,|e_2\rangle_a\Big),$$

one has $\rho_{12} = e^{-i\phi}/2$, and using Lemma 8.74 (iii),

$$v^{(a)}_{\text{Synthesis}}\Big(|\psi_\phi\rangle\langle\psi_\phi|\Big) \;=\; \tfrac{1}{2}+\mathrm{Re}(\rho_{12}) \;=\; \tfrac{1}{2}+\tfrac{1}{2}\cos\phi \;=\; \cos^2\Big(\tfrac{\phi}{2}\Big).$$

This interpolates continuously from 1 at $\phi = 0$ (the $|+\rangle_a$ state) to 0 at $\phi = \pi$ (the $|-\rangle_a$ state), explicitly exhibiting the coherence sensitivity of the parasymmetric valuation.

Definition 8.76 (Law–State parasymmetric subspaces)**.** On $\mathcal{H}^{(LS)} := \mathcal{H}^{(L)} \otimes \mathcal{H}^{(S)}$ define the (Bell-type) kets

$$|\Psi^{(LS)}_{\text{para}}\rangle = \tfrac{1}{\sqrt{2}}\Big(|e_1\rangle_L|e_2\rangle_S + |e_2\rangle_L|e_1\rangle_S\Big), \quad |\Psi^{(LS)}_{\text{antipara}}\rangle = \tfrac{1}{\sqrt{2}}\Big(|e_1\rangle_L|e_2\rangle_S - |e_2\rangle_L|e_1\rangle_S\Big),$$
$$|\Phi^{(LS)}_{\text{para}}\rangle = \tfrac{1}{\sqrt{2}}\Big(|e_1\rangle_L|e_1\rangle_S + |e_2\rangle_L|e_2\rangle_S\Big), \quad |\Phi^{(LS)}_{\text{antipara}}\rangle = \tfrac{1}{\sqrt{2}}\Big(|e_1\rangle_L|e_1\rangle_S - |e_2\rangle_L|e_2\rangle_S\Big),$$

with projectors $P^{(LS)}_{\text{para}} := |\Psi^{(LS)}_{\text{para}}\rangle\langle\Psi^{(LS)}_{\text{para}}|$, $P^{(LS)}_{\text{antipara}}$, $P^{(LS)}_{\Phi,\text{para}}$, $P^{(LS)}_{\Phi,\text{antipara}}$. We call the Ψ-pair the *opposition sector* (II/III plane) and the Φ-pair the *agreement sector* (I/IV plane), relative to the quadrant formulas (Definition 8.61).

Lemma 8.77 (Orthogonality and block decomposition)**.** *The four kets in Definition 8.76 are orthonormal. Moreover,*

$$\mathsf{S}_\Psi := \mathrm{span}\{|\Psi^{(LS)}_{\text{para}}\rangle, |\Psi^{(LS)}_{\text{antipara}}\rangle\}, \qquad \mathsf{S}_\Phi := \mathrm{span}\{|\Phi^{(LS)}_{\text{para}}\rangle, |\Phi^{(LS)}_{\text{antipara}}\rangle\}$$

are mutually orthogonal, and each is the minimal invariant subspace generated by the corresponding computational-basis pairs:

$$\mathsf{S}_\Psi = \mathrm{span}\{|e_1e_2\rangle, |e_2e_1\rangle\}, \qquad \mathsf{S}_\Phi = \mathrm{span}\{|e_1e_1\rangle, |e_2e_2\rangle\}.$$

Proof. For brevity write $|ij\rangle := |e_i\rangle_L|e_j\rangle_S$. Then

$$|\Psi_{\text{para}}\rangle = \tfrac{1}{\sqrt{2}}(|12\rangle + |21\rangle), \quad |\Psi_{\text{antipara}}\rangle = \tfrac{1}{\sqrt{2}}(|12\rangle - |21\rangle),$$

$$|\Phi_{\text{para}}\rangle = \tfrac{1}{\sqrt{2}}(|11\rangle + |22\rangle), \quad |\Phi_{\text{antipara}}\rangle = \tfrac{1}{\sqrt{2}}(|11\rangle - |22\rangle).$$

Orthonormality. Using $\langle ij|k\ell\rangle = \delta_{ik}\delta_{j\ell}$,

$$\langle\Psi_{\text{para}}|\Psi_{\text{para}}\rangle = \tfrac{1}{2}(\langle 12| + \langle 21|)(|12\rangle + |21\rangle) = 1,$$

$$\langle\Psi_{\text{antipara}}|\Psi_{\text{antipara}}\rangle = \tfrac{1}{2}(\langle 12| - \langle 21|)(|12\rangle - |21\rangle) = 1,$$

and $\langle\Psi_{\text{para}}|\Psi_{\text{antipara}}\rangle = \tfrac{1}{2}(\langle 12| + \langle 21|)(|12\rangle - |21\rangle) = 0$. The same computation shows $\langle\Phi_{\text{para}}|\Phi_{\text{para}}\rangle = \langle\Phi_{\text{antipara}}|\Phi_{\text{antipara}}\rangle = 1$ and $\langle\Phi_{\text{para}}|\Phi_{\text{antipara}}\rangle = 0$. Cross terms between the Ψ- and Φ-pairs vanish because the former live in $\text{span}\{|12\rangle, |21\rangle\}$ and the latter in $\text{span}\{|11\rangle, |22\rangle\}$, which are orthogonal. Hence the four kets form an orthonormal set.

Block projectors and orthogonal decomposition. Let

$$P_{\Psi\text{-block}} := P^{(LS)}_{\text{para}} + P^{(LS)}_{\text{antipara}}, \qquad P_{\Phi\text{-block}} := P^{(LS)}_{\Phi,\text{para}} + P^{(LS)}_{\Phi,\text{antipara}}.$$

A direct expansion gives

$$P_{\Psi\text{-block}} = |\Psi_{\text{para}}\rangle\langle\Psi_{\text{para}}| + |\Psi_{\text{antipara}}\rangle\langle\Psi_{\text{antipara}}| = |12\rangle\langle 12| + |21\rangle\langle 21|,$$

$$P_{\Phi\text{-block}} = |\Phi_{\text{para}}\rangle\langle\Phi_{\text{para}}| + |\Phi_{\text{antipara}}\rangle\langle\Phi_{\text{antipara}}| = |11\rangle\langle 11| + |22\rangle\langle 22|.$$

Therefore $\text{Ran}(P_{\Psi\text{-block}}) = \text{span}\{|12\rangle, |21\rangle\} =: \mathsf{S}_\Psi$ and $\text{Ran}(P_{\Phi\text{-block}}) = \text{span}\{|11\rangle, |22\rangle\} =: \mathsf{S}_\Phi$. Since these ranges are orthogonal and exhaust the computational basis, we have

$$P_{\Psi\text{-block}}P_{\Phi\text{-block}} = 0, \qquad P_{\Psi\text{-block}} + P_{\Phi\text{-block}} = \text{Id}_{\mathcal{H}^{(LS)}},$$

so $\mathcal{H}^{(LS)} = \mathsf{S}_\Psi \oplus \mathsf{S}_\Phi$.

Minimality and invariance. Each block projector is the identity on its range and annihilates the other block, hence S_Ψ and S_Φ are invariant under the algebra generated by the four rank-1 projectors (equivalently, under $P_{\Psi\text{-block}}, P_{\Phi\text{-block}}$). Moreover, S_Ψ is the *smallest* subspace containing $|12\rangle, |21\rangle$ that is invariant under this algebra—because $P_{\Psi\text{-block}}$ is precisely the orthogonal projector onto $\text{span}\{|12\rangle, |21\rangle\}$ (and similarly for S_Φ with $|11\rangle, |22\rangle$). Finally, within each block, $\{|\Psi_{\text{para}}\rangle, |\Psi_{\text{antipara}}\rangle\}$ and $\{|\Phi_{\text{para}}\rangle, |\Phi_{\text{antipara}}\rangle\}$ are related to their computational pairs by the unitary $\frac{1}{\sqrt{2}}\left(\begin{smallmatrix} 1 & 1 \\ 1 & -1 \end{smallmatrix}\right)$, so they indeed form orthonormal bases of the respective minimal invariant subspaces. □

Definition 8.78 (Joint Synthesis valuations)**.** For a density operator ρ_{LS} on $\mathcal{H}^{(LS)}$ set

$$v^{(LS)}_{\text{Synthesis}}(\rho_{LS}) := \text{Tr}\Big(P^{(LS)}_{\text{para}}\rho_{LS}\Big), \qquad v^{(LS)}_{\text{Agreement}}(\rho_{LS}) := \text{Tr}\Big(P^{(LS)}_{\Phi,\text{para}}\rho_{LS}\Big).$$

These quantify coherent Law–State *opposition* and *agreement*, respectively.

Definition 8.79 (Unified parasymmetric generator on $L \oplus S$)**.** Define the Hermitian generator on $\mathcal{H}^{(LS)}$:

$$\begin{aligned}\hat{F}^{(LS)}_{\text{para}} &= \lambda_L \Big(P^{(L)}_{\text{para}} - P^{(L)}_{\text{antipara}}\Big) \otimes \mathrm{Id}^{(S)}_{(0,1)} \\ &\quad + \lambda_S \, \mathrm{Id}^{(L)}_{(0,1)} \otimes \Big(P^{(S)}_{\text{para}} - P^{(S)}_{\text{antipara}}\Big) \\ &\quad + \lambda_{LS} \Big(P^{(LS)}_{\text{para}} - P^{(LS)}_{\text{antipara}}\Big) \\ &\quad + \lambda_\Phi \Big(P^{(LS)}_{\Phi,\text{para}} - P^{(LS)}_{\Phi,\text{antipara}}\Big) + \cdots \end{aligned} \tag{8.15}$$

and the unitary $U^{(LS)}_t := \exp\Big(it\, \hat{F}^{(LS)}_{\text{para}}\Big)$. (Here the ellipsis allows additional symmetry-compatible couplings; cf. ladder-type exchanges below.)

Definition 8.80 (Exchange operators (optional))**.** On each anchor define $\hat{\sigma}^{(+)}_a := |e_1\rangle_a\langle e_2|$, $\hat{\sigma}^{(-)}_a := |e_2\rangle_a\langle e_1|$. A *direct excitation exchange* term has the Hermitian form

$$\hat{F}^{(\text{ex})} := \sum_\mu \Big(J_\mu \, \hat{\sigma}^{(+)}_L \hat{\sigma}^{(-)}_S + \overline{J_\mu} \, \hat{\sigma}^{(-)}_L \hat{\sigma}^{(+)}_S \Big),$$

and may be absorbed into $\hat{F}^{(LS)}_{\text{para}}$ with a coupling λ_X.

Proposition 8.81 (Invariance criteria & constants of motion)**.** *Let* $P \in \{\, P^{(L)}_{\text{para}} \otimes \mathrm{Id},\ \mathrm{Id} \otimes P^{(S)}_{\text{para}},\ P^{(LS)}_{\text{para}},\ P^{(LS)}_{\Phi,\text{para}} \,\}$. *For the evolution* $\rho_{LS}(t) = U^{(LS)}_t \, \rho_{LS}(0) \, U^{(LS)\dagger}_t$ *with* $U^{(LS)}_t = \exp\Big(it\, \hat{F}^{(LS)}_{\text{para}}\Big)$,

$$\frac{d}{dt}\, \mathrm{Tr}\Big(P\, \rho_{LS}(t)\Big) = i\, \mathrm{Tr}\Big([\hat{F}^{(LS)}_{\text{para}}, P]\, \rho_{LS}(t)\Big).$$

In particular, if $[\hat{F}^{(LS)}_{\text{para}}, P] = 0$, *then* $\mathrm{Tr}\Big(P\, \rho_{LS}(t)\Big)$ *is conserved.*

Proof. Write $U := U^{(LS)}_t$, $\hat{F} := \hat{F}^{(LS)}_{\text{para}}$, and $\rho := \rho_{LS}(0)$. By definition $\rho(t) = U \rho U^\dagger$. Differentiate and use $\dot{U} = i\hat{F}U$, $\dot{U}^\dagger = -iU^\dagger \hat{F}$:

$$\frac{d}{dt}\, \mathrm{Tr}\Big(P\, \rho(t)\Big) = \mathrm{Tr}\Big(P\, \dot{U}\, \rho\, U^\dagger\Big) + \mathrm{Tr}\Big(P\, U\, \rho\, \dot{U}^\dagger\Big) = i\, \mathrm{Tr}\Big(P\, \hat{F}\, U \rho U^\dagger\Big) - i\, \mathrm{Tr}\Big(P\, U \rho U^\dagger \hat{F}\Big).$$

By cyclicity of trace, $\mathrm{Tr}\Big(P\, \hat{F}\, U \rho U^\dagger\Big) = \mathrm{Tr}\Big(\hat{F}\, U \rho U^\dagger P\Big)$ and $\mathrm{Tr}\Big(P\, U \rho U^\dagger \hat{F}\Big) = \mathrm{Tr}\Big(\hat{F}\, P\, U \rho U^\dagger\Big)$. Thus

$$\frac{d}{dt}\, \mathrm{Tr}\Big(P\, \rho(t)\Big) = i\, \mathrm{Tr}\Big(\, [\hat{F}, P]\, U \rho U^\dagger\Big) = i\, \mathrm{Tr}\Big([\hat{F}, P]\, \rho(t)\Big),$$

which is the claimed Heisenberg-form identity. If $[\hat{F}, P] = 0$ then the derivative vanishes, so $\mathrm{Tr}(P\rho(t))$ is constant in time. □

Lemma 8.82 (Commutation structure of the generator)**.** *Let* $Q_L := P^{(L)}_{\text{para}} - P^{(L)}_{\text{antipara}}$, $Q_S := P^{(S)}_{\text{para}} - P^{(S)}_{\text{antipara}}$, $Q_\Psi := P^{(LS)}_{\text{para}} - P^{(LS)}_{\text{antipara}}$, *and* $Q_\Phi := P^{(LS)}_{\Phi,\text{para}} - P^{(LS)}_{\Phi,\text{antipara}}$. *Then:*

(i) $[Q_\Psi, P^{(LS)}_{\text{para}}] = 0$ *and* $[Q_\Phi, P^{(LS)}_{\text{para}}] = 0$.

(ii) $[Q_L \otimes \mathrm{Id},\ P^{(LS)}_{\text{para}}] \neq 0$ *and* $[\mathrm{Id} \otimes Q_S,\ P^{(LS)}_{\text{para}}] \neq 0$ *in general.*

(iii) *For the exchange operator* $\hat{F}^{(\mathrm{ex})} = \sum_\mu \left(J_\mu \, \hat{\sigma}_L^{(+)} \hat{\sigma}_S^{(-)} + \overline{J_\mu} \, \hat{\sigma}_L^{(-)} \hat{\sigma}_S^{(+)} \right)$, $[\hat{F}^{(\mathrm{ex})}, P^{(LS)}_{\mathrm{para}}] = 0$ *and* $[\hat{F}^{(\mathrm{ex})}, P^{(LS)}_{\Phi,\mathrm{para}}] = 0$.

Proof. Set

$$|ij\rangle := |e_i\rangle_L \otimes |e_j\rangle_S, \qquad \mathsf{S}_\Psi := \mathrm{span}\{|12\rangle, \, |21\rangle\}.$$

In the orthonormal basis $\{|\Psi^{(LS)}_{\mathrm{para}}\rangle, \, |\Psi^{(LS)}_{\mathrm{antipara}}\rangle\}$ of S_Ψ,

$$P^{(LS)}_{\mathrm{para}} \restriction_{\mathsf{S}_\Psi} = \left(\begin{smallmatrix} 1 & 0 \\ 0 & 0 \end{smallmatrix} \right).$$

(a) On S_Ψ, $Q_\Psi = \mathrm{diag}(1, -1)$, hence it commutes with $\mathrm{diag}(1, 0)$. On the orthogonal Φ-block $\mathsf{S}_\Phi = \mathrm{span}\{|11\rangle, |22\rangle\}$ we have $P^{(LS)}_{\mathrm{para}} = 0$, so $[Q_\Phi, P^{(LS)}_{\mathrm{para}}] = 0$ trivially and $[Q_\Psi, P^{(LS)}_{\mathrm{para}}] = 0$ blockwise.

(b) The local "tilt" $Q_L \otimes \mathrm{Id}$ maps $|12\rangle \mapsto |22\rangle$ and $|21\rangle \mapsto |11\rangle$, hence it couples S_Ψ to S_Φ. Therefore it does not commute with $P^{(LS)}_{\mathrm{para}}$, which projects onto S_Ψ. An analogous statement holds for $\mathrm{Id} \otimes Q_S$.

(c) Using $\hat{\sigma}_L^{(+)} \hat{\sigma}_S^{(-)} |12\rangle = |21\rangle$, $\hat{\sigma}_L^{(+)} \hat{\sigma}_S^{(-)} |21\rangle = |12\rangle$, and zero action on $|11\rangle, |22\rangle$, $\hat{F}^{(\mathrm{ex})}$ acts on S_Ψ as $\mathrm{diag}(+|J|, -|J|)$ in the $\{|\Psi_{\mathrm{para}}\rangle, |\Psi_{\mathrm{antipara}}\rangle\}$ basis and annihilates S_Φ. Hence it commutes with both $P^{(LS)}_{\mathrm{para}}$ and $P^{(LS)}_{\Phi,\mathrm{para}}$. □

Corollary 8.83 (When opposition/agreement "scores" are conserved)**.** *Let* $v^{(LS)}_{\mathrm{Synthesis}}(t) := \mathrm{Tr}\Big(P^{(LS)}_{\mathrm{para}} \rho_{LS}(t) \Big)$ *and* $v^{(LS)}_{\mathrm{Agreement}}(t) := \mathrm{Tr}\Big(P^{(LS)}_{\Phi,\mathrm{para}} \rho_{LS}(t) \Big)$.

(i) If $\lambda_L = \lambda_S = 0$ *(no local tilts) and the generator consists of any combination of* Q_Ψ, Q_Φ, *and exchange terms* $\hat{F}^{(\mathrm{ex})}$, *then* $v^{(LS)}_{\mathrm{Synthesis}}(t)$ *and* $v^{(LS)}_{\mathrm{Agreement}}(t)$ *are constants of motion.*

(ii) If a local tilt is present (either $\lambda_L \neq 0$ *or* $\lambda_S \neq 0$*), then generically* $\frac{d}{dt} v^{(LS)}_{\mathrm{Synthesis}}(t) \neq 0$ *and* $\frac{d}{dt} v^{(LS)}_{\mathrm{Agreement}}(t) \neq 0$ *because* $Q_L \otimes \mathrm{Id}$ *and* $\mathrm{Id} \otimes Q_S$ *mix the* Ψ/Φ *blocks (Lemma 8.82(b)).*

Proof. Immediate from Proposition 8.81 and Lemma 8.82: in case (i), each listed term commutes with both $P^{(LS)}_{\mathrm{para}}$ and $P^{(LS)}_{\Phi,\mathrm{para}}$; in case (ii) at least one commutator is nonzero. □

Corollary 8.84 (Preserving local vs. relational parasymmetry)**.** *(a) If* $[\hat{F}^{(LS)}_{\mathrm{para}}, P^{(L)}_{\mathrm{para}} \otimes \mathrm{Id}] = 0 = [\hat{F}^{(LS)}_{\mathrm{para}}, \mathrm{Id} \otimes P^{(S)}_{\mathrm{para}}]$, *then the local valuations* $v^{(L)}_{\mathrm{Synthesis}}$ *and* $v^{(S)}_{\mathrm{Synthesis}}$ *are constants of motion. (b) If* $[\hat{F}^{(LS)}_{\mathrm{para}}, P^{(LS)}_{\mathrm{para}}] = 0$ *(resp.* $[\hat{F}^{(LS)}_{\mathrm{para}}, P^{(LS)}_{\Phi,\mathrm{para}}] = 0$*), then the joint valuations* $v^{(LS)}_{\mathrm{Synthesis}}$ *(resp.* $v^{(LS)}_{\mathrm{Agreement}}$*) are constants of motion.*

Proof. Write $\rho(t) := \rho_{LS}(t) = U_t^{(LS)} \, \rho(0) \, U_t^{(LS)\dagger}$.

(a) Local valuations. Using the partial trace identity,

$$v^{(L)}_{\mathrm{Synthesis}}(t) = \mathrm{Tr}\Big(P^{(L)}_{\mathrm{para}} \, \rho_L(t) \Big) = \mathrm{Tr}\Big((P^{(L)}_{\mathrm{para}} \otimes \mathrm{Id}) \, \rho(t) \Big),$$

and similarly $v^{(S)}_{\mathrm{Synthesis}}(t) = \mathrm{Tr}\Big((\mathrm{Id} \otimes P^{(S)}_{\mathrm{para}}) \, \rho(t) \Big)$. By Proposition 8.81,

$$\frac{d}{dt} \, v^{(L)}_{\mathrm{Synthesis}}(t) = i \, \mathrm{Tr}\Big([\hat{F}^{(LS)}_{\mathrm{para}}, \, P^{(L)}_{\mathrm{para}} \otimes \mathrm{Id}] \, \rho(t) \Big),$$

and analogously for $v^{(S)}_{\text{Synthesis}}(t)$. Under the stated commutation hypotheses each derivative vanishes, proving constancy in time.

(b) Relational valuations. By definition, $v^{(LS)}_{\text{Synthesis}}(t) = \mathrm{Tr}\big(P^{(LS)}_{\text{para}}\,\rho(t)\big)$ and $v^{(LS)}_{\text{Agreement}}(t) = \mathrm{Tr}\big(P^{(LS)}_{\Phi,\text{para}}\,\rho(t)\big)$. Applying Proposition 8.81 with $P = P^{(LS)}_{\text{para}}$ or $P^{(LS)}_{\Phi,\text{para}}$ yields

$$\frac{d}{dt}\,v^{(LS)}_{\text{Synthesis}}(t) = i\ \mathrm{Tr}\Big([\hat{F}^{(LS)}_{\text{para}},\,P^{(LS)}_{\text{para}}]\,\rho(t)\Big), \qquad \frac{d}{dt}\,v^{(LS)}_{\text{Agreement}}(t) = i\ \mathrm{Tr}\Big([\hat{F}^{(LS)}_{\text{para}},\,P^{(LS)}_{\Phi,\text{para}}]\,\rho(t)\Big).$$

Thus each valuation is conserved whenever the corresponding commutator vanishes. □

Remark 8.85 (Steering vs. preserving)**.** *Preserving.* Setting $[\hat{F}^{(LS)}_{\text{para}}, P] = 0$ freezes the associated score $\mathrm{Tr}(P\rho(t))$ (local or relational), as in Corollary 8.84. In particular, the "diagonal" terms $Q_\Psi := P^{(LS)}_{\text{para}} - P^{(LS)}_{\text{antipara}}$ and $Q_\Phi := P^{(LS)}_{\Phi,\text{para}} - P^{(LS)}_{\Phi,\text{antipara}}$ commute with $P^{(LS)}_{\text{para}}$ and $P^{(LS)}_{\Phi,\text{para}}$, so they *do not* change the opposition/agreement scores—they implement phase tilts within each block.

Steering within a block. To *redistribute* probability between "+" and "−" inside a fixed relational block while keeping the block weight constant, include off-diagonal, block-preserving couplings such as

$$X_\Psi := |\Psi^{(LS)}_{\text{para}}\rangle\langle\Psi^{(LS)}_{\text{antipara}}| + \text{h.c.}, \qquad X_\Phi := |\Phi^{(LS)}_{\text{para}}\rangle\langle\Phi^{(LS)}_{\text{antipara}}| + \text{h.c.}$$

These commute with the *block* projectors $P_{\Psi\text{-block}} := P^{(LS)}_{\text{para}} + P^{(LS)}_{\text{antipara}}$ and $P_{\Phi\text{-block}} := P^{(LS)}_{\Phi,\text{para}} + P^{(LS)}_{\Phi,\text{antipara}}$, but not with the individual "+" projectors:

$$[X_\Psi, P_{\Psi\text{-block}}] = 0 \quad \text{and} \quad [X_\Psi, P^{(LS)}_{\text{para}}] \neq 0 \quad \text{(similarly for } X_\Phi\text{)}.$$

Hence adding X_Ψ (resp. X_Φ) *steers* $v^{(LS)}_{\text{Synthesis}}(t)$ (resp. $v^{(LS)}_{\text{Agreement}}(t)$) while preserving the total weight in the Ψ- (resp. Φ-) sector. By contrast, the exchange operator of Definition 8.80 *preserves* both scores, as it commutes with the corresponding "+" projectors (Lemma 8.82(c)).

Definition 8.86 (Quadrant projectors on $\mathcal{H}^{(LS)}$)**.** With $\varphi_L := \mathrm{Sym}(L)$ and $\varphi_S := \mathrm{Sym}(w;\lambda)$, recall the quadrant formulas (Definition 8.61). On $\mathcal{H}^{(LS)}$ define the corresponding computational-basis projectors:

$$\Pi_{\mathrm{I}} := |e_1e_1\rangle\langle e_1e_1|,\ \Pi_{\mathrm{II}} := |e_1e_2\rangle\langle e_1e_2|,\ \Pi_{\mathrm{III}} := |e_2e_1\rangle\langle e_2e_1|,\ \Pi_{\mathrm{IV}} := |e_2e_2\rangle\langle e_2e_2|.$$

Lemma 8.87 (Coherent diagonals through the dashboard)**.** *We have the identities*

$$P^{(LS)}_{\text{para}} = \tfrac{1}{2}(\Pi_{\mathrm{II}} + \Pi_{\mathrm{III}} + |e_1e_2\rangle\langle e_2e_1| + |e_2e_1\rangle\langle e_1e_2|)\,,$$
$$P^{(LS)}_{\Phi,\text{para}} = \tfrac{1}{2}(\Pi_{\mathrm{I}} + \Pi_{\mathrm{IV}} + |e_1e_1\rangle\langle e_2e_2| + |e_2e_2\rangle\langle e_1e_1|)\,.$$

Thus $P^{(LS)}_{\text{para}}$ (resp. $P^{(LS)}_{\Phi,\text{para}}$) enhances the II–III (resp. I–IV) diagonal by adding phase-coherent cross-terms; it is not a mere mixture of the two corners.

Proof. Abbreviate $|ij\rangle := |e_i\rangle_L|e_j\rangle_S$. From Definition 8.76, $|\Psi^{(LS)}_{\text{para}}\rangle = \frac{1}{\sqrt{2}}(|12\rangle + |21\rangle)$, so

$$\begin{aligned} P^{(LS)}_{\text{para}} &= |\Psi^{(LS)}_{\text{para}}\rangle\langle\Psi^{(LS)}_{\text{para}}| = \tfrac{1}{2}\left(|12\rangle + |21\rangle\right)\left(\langle 12| + \langle 21|\right) \\ &= \tfrac{1}{2}\Big(|12\rangle\langle 12| + |21\rangle\langle 21| + |12\rangle\langle 21| + |21\rangle\langle 12|\Big) \\ &= \tfrac{1}{2}\Big(\Pi_{\text{II}} + \Pi_{\text{III}} + |e_1e_2\rangle\langle e_2e_1| + |e_2e_1\rangle\langle e_1e_2|\Big). \end{aligned}$$

Similarly, $|\Phi^{(LS)}_{\text{para}}\rangle = \frac{1}{\sqrt{2}}(|11\rangle + |22\rangle)$ gives

$$\begin{aligned} P^{(LS)}_{\Phi,\text{para}} &= \tfrac{1}{2}\Big(|11\rangle\langle 11| + |22\rangle\langle 22| + |11\rangle\langle 22| + |22\rangle\langle 11|\Big) \\ &= \tfrac{1}{2}\Big(\Pi_{\text{I}} + \Pi_{\text{IV}} + |e_1e_1\rangle\langle e_2e_2| + |e_2e_2\rangle\langle e_1e_1|\Big). \end{aligned}$$

This shows in each case a diagonal (mixture part) plus coherent cross-terms. □

Proposition 8.88 (Refinement of SynTh by parasymmetry)**.** *Suppose the logical layer assigns designated truth* $\mathbf{B}$ *to both* φ_L *and* φ_S *(SynTh holds simultaneously: Setup 8.67 and Theorem 8.70). Then, for any* ρ_{LS}*, the parasymmetric valuations* $\mathrm{Tr}\Big(P^{(LS)}_{\text{para}}\rho_{LS}\Big)$ *and* $\mathrm{Tr}\Big(P^{(LS)}_{\Phi,\text{para}}\rho_{LS}\Big)$ *quantify, respectively, the* coherent opposition *(II/III) and* coherent agreement *(I/IV) content that SynTh leaves undifferentiated.*

Proof. Write the matrix elements of ρ_{LS} in the computational basis as $\rho_{ij,kl} := \langle e_ie_j|\rho_{LS}|e_ke_\ell\rangle$. Using Lemma 8.87,

$$\begin{aligned} \mathrm{Tr}\Big(P^{(LS)}_{\text{para}}\rho_{LS}\Big) &= \tfrac{1}{2}(\rho_{12,12} + \rho_{21,21} + \rho_{12,21} + \rho_{21,12}) \\ &= \tfrac{1}{2}\Big(\underbrace{\rho_{12,12}}_{\text{II}} + \underbrace{\rho_{21,21}}_{\text{III}}\Big) + \mathrm{Re}\Big(\rho_{12,21}\Big), \end{aligned}$$

and

$$\begin{aligned} \mathrm{Tr}\Big(P^{(LS)}_{\Phi,\text{para}}\rho_{LS}\Big) &= \tfrac{1}{2}(\rho_{11,11} + \rho_{22,22} + \rho_{11,22} + \rho_{22,11}) \\ &= \tfrac{1}{2}\Big(\underbrace{\rho_{11,11}}_{\text{I}} + \underbrace{\rho_{22,22}}_{\text{IV}}\Big) + \mathrm{Re}\Big(\rho_{11,22}\Big). \end{aligned}$$

Thus each valuation splits into a *mixture* part (average of the two relevant dashboard corners) plus a *coherence* part (real off-diagonal along the same diagonal).

Under Setup 8.67, the SynTh layer makes all four corners simultaneously adoptable (designated), but it does not encode coherences. The parasymmetric valuations therefore refine SynTh by weighting the II/III and I/IV diagonals and, crucially, by detecting their *coherent* content via the cross-terms $\mathrm{Re}(\rho_{12,21})$ and $\mathrm{Re}(\rho_{11,22})$. This is exactly the claimed opposition vs. agreement quantification. □

Remark 8.89 (Compatibility with paraconsistency)**.** All valuations here are Born weights in $[0, 1]$ and do not alter the underlying consequence relation $\vdash_{\text{pc}}$. In particular, non-explosion remains ensured by the witness Π_{pc} (Definition 8.48 and Lemma 8.52). Parasymmetry thus serves as an *operational semantics* layered on top of the **SynTh** logic: it distinguishes coherent vs. incoherent contributions along the I/IV and II/III diagonals without changing which formulas are designated or derivable in $\vdash_{\text{pc}}$.

Definition 8.90 (Simultaneous Parasymmetry thresholds)**.** Fix thresholds (dimensionless real numbers)

$$0 \le \tau_L,\ \tau_S,\ \tau_{LS},\ \tau_\Phi \le 1.$$

Let ρ_{LS} be a density operator on the joint Law–State space $\mathcal{H}^{(LS)}$, and let its marginals be $\rho_L := \mathrm{Tr}_S(\rho_{LS})$ and $\rho_S := \mathrm{Tr}_L(\rho_{LS})$. With the projectors $P^{(L)}_{\mathrm{para}}, P^{(S)}_{\mathrm{para}}$ from Definition 8.73 and $P^{(LS)}_{\mathrm{para}}, P^{(LS)}_{\Phi,\mathrm{para}}$ from Definition 8.76, we say that $\mathrm{SimulPara}(L, S; \lambda)$ holds if

$$\begin{aligned}
\mathrm{Tr}\Big(P^{(L)}_{\mathrm{para}}\rho_L\Big) &\ge \tau_L,\\
\mathrm{Tr}\Big(P^{(S)}_{\mathrm{para}}\rho_S\Big) &\ge \tau_S,\\
\mathrm{Tr}\Big(P^{(LS)}_{\mathrm{para}}\rho_{LS}\Big) &\ge \tau_{LS},\\
\mathrm{Tr}\Big(P^{(LS)}_{\Phi,\mathrm{para}}\rho_{LS}\Big) &\ge \tau_\Phi.
\end{aligned}$$

Remark 8.91. The predicate $\mathrm{SimulPara}(L, S; \lambda)$ is monotone in the thresholds: if it holds for $(\tau_L, \tau_S, \tau_{LS}, \tau_\Phi)$, it holds for any smaller tuple.

Proof. Each quantity $\mathrm{Tr}(P\rho)$ with P a projector is in $[0, 1]$. If $\mathrm{Tr}(P\rho) \ge \tau$ and $\tau' \le \tau$, then $\mathrm{Tr}(P\rho) \ge \tau'$. Apply this pointwise to the four inequalities in Definition 8.90. □

Theorem 8.92 (Preservation under unified evolution)**.** *Let* $\rho_{LS}(t) = U_t^{(LS)}\,\rho_{LS}(0)\,U_t^{(LS)\dagger}$ *with* $U_t^{(LS)} = \exp\!\big(it\,\hat{F}^{(LS)}_{\mathrm{para}}\big)$*. If the commutator conditions in Corollary 8.84 hold, then* $\mathrm{SimulPara}(L, S; \lambda)$ *is invariant in time: each of the four inequalities in Definition 8.90 remains true for all* $t \ge 0$*.*

Proof. Write $\rho(t) := \rho_{LS}(t)$. We show each score is constant; the inequalities at $t = 0$ then persist.

Local scores. Using the partial trace,

$$v^{(L)}_{\mathrm{Synthesis}}(t) = \mathrm{Tr}\Big(P^{(L)}_{\mathrm{para}}\,\rho_L(t)\Big) = \mathrm{Tr}\Big((P^{(L)}_{\mathrm{para}}\otimes\mathrm{Id})\,\rho(t)\Big), \quad v^{(S)}_{\mathrm{Synthesis}}(t) = \mathrm{Tr}\Big((\mathrm{Id}\otimes P^{(S)}_{\mathrm{para}})\,\rho(t)\Big).$$

By Proposition 8.81,

$$\frac{d}{dt}\,v^{(L)}_{\mathrm{Synthesis}}(t) = i\ \mathrm{Tr}\Big([\hat{F}^{(LS)}_{\mathrm{para}},\ P^{(L)}_{\mathrm{para}}\otimes\mathrm{Id}]\,\rho(t)\Big),$$

and similarly for $v^{(S)}_{\mathrm{Synthesis}}(t)$. If the commutators in Corollary 8.84(a) vanish, both derivatives are zero, so the local inequalities remain true for all t.

Relational scores. By definition,

$$v^{(LS)}_{\mathrm{Synthesis}}(t) = \mathrm{Tr}\Big(P^{(LS)}_{\mathrm{para}}\,\rho(t)\Big), \qquad v^{(LS)}_{\mathrm{Agreement}}(t) = \mathrm{Tr}\Big(P^{(LS)}_{\Phi,\mathrm{para}}\,\rho(t)\Big).$$

Proposition 8.81 gives

$$\frac{d}{dt}\,v^{(LS)}_{\mathrm{Synthesis}}(t) = i\ \mathrm{Tr}\Big([\hat{F}^{(LS)}_{\mathrm{para}},\ P^{(LS)}_{\mathrm{para}}]\,\rho(t)\Big), \qquad \frac{d}{dt}\,v^{(LS)}_{\mathrm{Agreement}}(t) = i\ \mathrm{Tr}\Big([\hat{F}^{(LS)}_{\mathrm{para}},\ P^{(LS)}_{\Phi,\mathrm{para}}]\,\rho(t)\Big).$$

If the commutators in Corollary 8.84(b) vanish, both derivatives are zero, so the relational inequalities remain true for all t.

Since all four scores are constants of motion under the stated hypotheses, any initial satisfaction of the threshold constraints in Definition 8.90 persists for all $t \ge 0$. □

Remark 8.93 (Global unified operator)**.** In many-site settings (n PQS), one replaces $\hat{F}^{(LS)}_{\mathrm{para}}$ by a unified, multi-scale generator

$$\hat{F}^{(\odot)}_{\mathrm{para}} = \lambda_L \hat{F}^{(\mathrm{local})}_{\mathrm{para}} + \lambda_P \hat{F}^{(\mathrm{pair})}_{\mathrm{para}} + \lambda_G \hat{F}^{(n)}_{\mathrm{para}} + \lambda_X \hat{F}^{(\mathrm{ex})} + \cdots ,$$

and sets $U_G^{(\odot)} := \exp\!\big(i\,\hat{F}^{(\odot)}_{\mathrm{para}}\big)$. The blockwise invariance logic is identical: if each block projector P of interest (local or relational, possibly lifted to n sites) satisfies $[\hat{F}^{(\odot)}_{\mathrm{para}}, P] = 0$, then the corresponding Born weight $\mathrm{Tr}(P\rho(t))$ is conserved. Thus any instance of $\mathrm{SimulPara}(L, S; \lambda)$ defined via such projectors is preserved under $U_G^{(\odot)}$.

It translates logic into measurable quantum structure. The parasymmetric layer provides a faithful operationalization of *Synthesis*. For each anchor $a \in \{L, S\}$, the map $\rho_a \mapsto v^{(a)}_{\mathrm{Synthesis}}(\rho_a) = \mathrm{Tr}\big(P^{(a)}_{\mathrm{para}}\rho_a\big)$ assigns a quantitative, phase-sensitive value in $[0, 1]$ that depends on the off-diagonal coherence in the $\{|\,\mathrm{Sym}\rangle, |\,\mathrm{Break}\rangle\}$ sector (Definition 8.73 and Lemma 8.74). Writing

$$P^{(a)}_{\mathrm{para}} = \tfrac{1}{2}\big(P^{(a)}_{(0,1)} + X_a\big), \qquad X_a := |e_1\rangle_a\langle e_2| + |e_2\rangle_a\langle e_1|,$$

one has $v^{(a)}_{\mathrm{Synthesis}}(\rho_a) = \frac{1}{2}\big(1+\langle X_a\rangle_{\rho_a}\big) = \frac{1}{2}+\mathrm{Re}(\rho_{12})$. Thus $v = \frac{1}{2}$ certifies an incoherent mixture, $v = 1$ certifies the coherent "both/and" state $|+\rangle_a$, and $v = 0$ certifies $|-\rangle_a$—an operational distinction the purely logical **SynTh** layer leaves intentionally agnostic (Definition 8.51 and Lemma 8.52; cf. [287, 453, 454, 455]).

It models relational structure via entanglement. On $\mathcal{H}^{(L)}\otimes\mathcal{H}^{(S)}$, the Bell-type projectors onto $|\Psi^{(LS)}_{\mathrm{para}}\rangle$ and $|\Phi^{(LS)}_{\mathrm{para}}\rangle$ quantify, respectively, coherent *opposition* (II/III diagonal) and coherent *agreement* (I/IV diagonal) between Law and State (Definitions 8.76 and 8.78). As made explicit in Lemma 8.87, these operators contain the crucial inter-quadrant cross-terms (e.g. $|e_1e_2\rangle\langle e_2e_1|$) that witness phase coherence across the "dashboard"; they are not mere mixtures of the corner projectors associated to I–IV (Definition 8.61). In multipartite settings the parasymmetric family naturally extends to W-class states (up to SLOCC), making "Synthesis as relation" a genuinely nonlocal resource [456, 287].

It supplies a complete dynamical and control framework. The unified generator $\hat{F}^{(LS)}_{\mathrm{para}}$ (Definition 8.79) yields exact invariants via commutators: for any parasymmetric projector P (local or relational), whenever $[\hat{F}^{(LS)}_{\mathrm{para}}, P] = 0$, the valuation $\mathrm{Tr}(P\rho_{LS}(t))$ is a constant of motion (Proposition 8.81 and Corollary 8.84). Tuning the couplings $\lambda_L, \lambda_S, \lambda_{LS}, \lambda_\Phi$ lets one preserve or steer local and relational coherences at will (Remark 8.85); the same blockwise invariance logic extends to the global multi-scale operator $U_G^{(\odot)}$ (Remark 8.93). In open-system form, for Lindblad evolution $\dot\rho = \mathcal{L}(\rho)$ the exact conservation condition is $\mathcal{L}^\dagger(P) = 0$, i.e.

$$\mathcal{L}^\dagger(P) = i[H, P] + \sum_k \Big(L_k^\dagger P L_k - \tfrac{1}{2}\{L_k^\dagger L_k, P\}\Big) = 0,$$

with the familiar sufficient criterion $[H, P] = [L_k, P] = 0\ \forall k$ (decoherence-free/noiseless subspaces), giving a practical route to robustly preserve chosen parasymmetric coherences [287].

It remains compatible with paraconsistency and duality. All parasymmetric valuations are Born weights layered on the paraconsistent base; they do not alter $\vdash_{\mathrm{pc}}$, and non-explosion remains guaranteed by the witness Π_{pc} (Definition 8.48 and Lemma 8.52). Scope hygiene is enforced throughout—diagnostics must refer to the same group G, the same sense of invariance, the same L, the same w, and (for the state) the same locus λ. Under the Axiom-of-Duality action $\mathbf{D}_{\mathcal{C}}$, the labels $|e_1\rangle, |e_2\rangle$ permute within the $(0,1)$-sector and parasymmetric projectors intertwine accordingly, so $P^{(\cdot)}_{\mathrm{para}}$-type invariants transport intact up to relabeling (Proposition 8.60).

Bottom line. Parasymmetry turns SynTh's logical "both/and" into *observables with dynamics and control*: locally via $v^{(a)}_{\mathrm{Synthesis}}$, relationally via $v^{(LS)}_{\mathrm{Synthesis}}$ and $v^{(LS)}_{\mathrm{Agreement}}$ (Definitions 8.73 and 8.78), and temporally via commutator-based invariants. The symmetry–asymmetry table is thereby promoted from a static classification to a phase-structured, steerable landscape, fully consistent with the paraconsistent logic and duality transport that organize the theory's statements.

Example 8.94 (Multipartite parasymmetric state)**.** The W-like state is defined as

$$|\Psi^{(n)}_{\mathrm{para}}\rangle = \tfrac{1}{\sqrt{n}} \sum_{k=1}^{n} |e_1\rangle_{x_1} \cdots |e_2\rangle_{x_k} \cdots |e_1\rangle_{x_n}.$$

It is normalized and permutation-symmetric. For any site x_j, the reduced state is

$$\rho_{x_j} = \mathrm{Tr}_{\{x_k : k \neq j\}}\Big(|\Psi^{(n)}_{\mathrm{para}}\rangle\langle\Psi^{(n)}_{\mathrm{para}}|\Big) = \tfrac{n-1}{n}\,|e_1\rangle\langle e_1| + \tfrac{1}{n}\,|e_2\rangle\langle e_2|,$$

which is mixed and diagonal (no local coherence). Hence $v^{(x_j)}_{\mathrm{Synthesis}}(\rho_{x_j}) = \frac{1}{2} + \mathrm{Re}(\rho_{12}) = \frac{1}{2}$, while the global projector $P^{(x_1 \ldots x_n)}_{\mathrm{para}} := |\Psi^{(n)}_{\mathrm{para}}\rangle\langle\Psi^{(n)}_{\mathrm{para}}|$ satisfies $\mathrm{Tr}\Big(P^{(x_1 \ldots x_n)}_{\mathrm{para}}\,|\Psi^{(n)}_{\mathrm{para}}\rangle\langle\Psi^{(n)}_{\mathrm{para}}|\Big) = 1$. Thus the state is genuinely multipartite entangled with mixed one-site marginals (cf. [456]); it implements *distributed Synthesis* via its projector $P^{(x_1 \ldots x_n)}_{\mathrm{para}}$.

Remark 8.95 (Standard logical/physical references)**.** The logical background for paraconsistency and non-explosion uses [453, 454, 455]. Ward identities and symmetry diagnostics are standard in QFT [91, 100]. For quantum operations and Bell/W-state structure see [287, 456].

8.2.2 Parasymmetry (with SynTh) vs. Duality

High–level reading: Parasymmetry vs. Duality (with SynTh). The Axiom–of–Duality action $\mathbf{D}_{\mathcal{C}}$ is indispensable for organizing structural equivalences and transporting statements across complementary descriptions (Proposition 8.60). By itself, however, duality is a *mapping principle*: it identifies and relates cells of the dashboard but does not supply observables, dynamical laws, or criteria that distinguish a *coherent* "both/and" from a mere *mixture*. The Parasymmetry $\oplus$ Synthesis layer fills exactly this gap. Building on the non-explosive logical substrate (Definition 8.48 and Lemma 8.52), Parasymmetry adds projectors and Born–valued diagnostics that make "both/and" *measurable*, *phase–sensitive*, and *controllable*, all while remaining compatible with duality.

Operational observables rather than identifications. Duality transports formulas and their defenses, but Parasymmetry supplies concrete tests. Locally, each anchor $a \in \{L, S\}$ carries the parasymmetric projector $P^{(a)}_{\text{para}}$ and the valuation $v^{(a)}_{\text{Synthesis}}(\rho_a) = \text{Tr}\big(P^{(a)}_{\text{para}}\rho_a\big) \in [0,1]$ (Definition 8.73). Relationally, on the Law–State sector one has the agreement/opposition projectors $P^{(LS)}_{\Phi,\text{para}}$, $P^{(LS)}_{\text{para}}$ and the joint scores $v^{(LS)}_{\text{Agreement}} = \text{Tr}\big(P^{(LS)}_{\Phi,\text{para}}\rho_{LS}\big)$, $v^{(LS)}_{\text{Synthesis}} = \text{Tr}\big(P^{(LS)}_{\text{para}}\rho_{LS}\big)$ (Definition 8.78). These are bona fide observables that quantify how much of the coherent "both/and" is present, locally and across the Law–State relation.

Sensitivity to coherence (off–diagonals) vs. mixtures. Where duality is silent about phase, Parasymmetry reads it directly: in the $\{|\,\text{Sym}\rangle, |\,\text{Break}\rangle\}$ basis, the local score depends on the off–diagonal element, $v^{(a)}_{\text{Synthesis}}(\rho_a) = \frac{1}{2} + \text{Re}(\rho_{12})$ (Lemma 8.74). Thus a coherent superposition $|+\rangle$ yields $v = 1$, while the balanced incoherent mixture $\frac{1}{2}(|e_1\rangle\langle e_1| + |e_2\rangle\langle e_2|)$ yields $v = \frac{1}{2}$ (Remark 8.75). This distinction—central to quantum structure—is invisible to duality as mapping.

From quadrant cells to coherent diagonals. The familiar 2×2 symmetry–asymmetry table corresponds to the computational basis on $\mathcal{H}^{(LS)}$ (Definition 8.86). Parasymmetry augments this with two orthogonal *Bell sectors* (Φ for Law–State agreement, Ψ for Law–State opposition), each spanned by a pair of Bell kets (Definition 8.76). Their projectors include inter–cell cross–terms such as $|e_1e_2\rangle\langle e_2e_1|$ and $|e_1e_1\rangle\langle e_2e_2|$, which are the algebraic witnesses of phase–coherent structure across the I/IV and II/III diagonals (Lemmas 8.77 and 8.87). In this way the dashboard is no longer four isolated cells but two coherent diagonals with quantitative weights.

Dynamics, invariants, and control. Parasymmetry also supplies the dynamical engine. The unified Law–State generator $\hat{F}^{(LS)}_{\text{para}}$ (Definition 8.79) and its multi–scale extensions (Remark 8.93) allow one to *preserve* or *steer* specific coherences by imposing exact commutation conditions. The criterion $[\,\hat{F}, P\,] = 0$ ensures that the valuation $\text{Tr}(P\rho(t))$ is conserved (Proposition 8.81); in particular one can lock in local parasymmetry $[\hat{F}, P^{(L)}_{\text{para}} \otimes \text{Id}] = [\hat{F}, \text{Id} \otimes P^{(S)}_{\text{para}}] = 0$ or relational agreement/opposition $[\hat{F}, P^{(LS)}_{\Phi,\text{para}}] = 0$, $[\hat{F}, P^{(LS)}_{\text{para}}] = 0$ (Corollary 8.84 and Remark 8.85). This level of controllable invariance has no analogue at the level of duality alone.

Quantitative, non–explosive multi–cell occupancy. **SynTh** establishes that all four cells can be designated without triviality (Theorem 8.63); Parasymmetry refines this picture by assigning *weights* to coherent diagonals and to the local "both" components. As a result, the matrix becomes a *phase–structured measure space* rather than a mere set of co–tenable propositions. Decision procedures (e.g. the SimulPara thresholds) make this operational: one declares $\text{SimulPara}(L, S; \lambda)$ to hold when the four scores exceed prescribed levels (Definition 8.90), and proves time–invariance of these predicates under suitable commutators (Theorem 8.92). Throughout, the paraconsistent base keeps explosion at bay (Definition 8.48 and Lemma 8.52).

Compatibility, not competition, with Duality. Finally, Parasymmetry respects duality. Under the duality action $\mathbf{D}_{\mathcal{C}}$, the $|\,\text{Sym}\rangle/|\,\text{Break}\rangle$ labels permute while the parasymmetric diagnostics transport functorially (Proposition 8.60). In practice: duality *classifies and relates* descriptions; Parasymmetry *measures and evolves* their coherent coexistence. Together with

SynTh, this turns a duality–organized landscape into a laboratory where "both/and" is not only permitted but *quantified*, *conserved*, and *steerable*.

Enhanced realism and descriptive power. Real systems often inhabit multiple cells at once (e.g. dominant SSB (II) with small explicit breaking (IV)). *SynTh* treats the table as a co-tenable *dashboard*, not a partition, so models needn't collapse to a single idealized cell. It also separates an *incoherent mixture* from a *coherent* "both": the parasymmetric *valuations* $v^{(a)}_{\text{Synthesis}}(\rho_a) = \mathrm{Tr}\big(P^{(a)}_{\text{para}}\rho_a\big) = \frac{1}{2} + \mathrm{Re}(\rho_{12})$ (Definition 8.73 and Lemma 8.74) give a measurable, phase-sensitive readout ($v = 1$ for $|+\rangle_a$, $v = \frac{1}{2}$ for no coherence, $v = 0$ for $|-\rangle_a$).

Novel observables and deeper insight. The joint *scores* $v^{(LS)}_{\text{Agreement}} = \mathrm{Tr}\big(P^{(LS)}_{\Phi,\text{para}}\rho_{LS}\big)$ and $v^{(LS)}_{\text{Synthesis}} = \mathrm{Tr}\big(P^{(LS)}_{\text{para}}\rho_{LS}\big)$ (Definition 8.78) are operational meters for *coherent* I/IV and II/III diagonals. By Lemma 8.87 they equal "mixture of the two corners" + "cross-term coherence," so pure corners give the $\frac{1}{2}$ baseline, while Bell-coherent superpositions yield 1. This cleanly explains finite-volume SSB cats: $v^{(LS)}_{\text{Synthesis}}$ drops from 1 to $\frac{1}{2}$ as $V \uparrow \infty$, matching Theorem 6.258.

A framework for quantum control. The unified generator $\hat{F}^{(LS)}_{\text{para}}$ (Definition 8.79) yields exact invariants: if $[\hat{F}^{(LS)}_{\text{para}}, P] = 0$ then $\mathrm{Tr}(P\rho_{LS}(t))$ is conserved (Proposition 8.81 and Corollary 8.84). Thus we can *preserve* chosen local or relational coherences, or *steer* them with block-preserving couplings (cf. Remark 8.85). In short: the theorem's Λ–asymptotics *locate* cells; *SynTh* lets multiple cells be co-adopted non-explosively; Parasymmetry makes "both/and" *measurable, phase-sensitive, and controllable*.

8.3 SynTh of Generalized Witness of Laws and States

What "Synthesis of Laws and States" formalizes. The aim of Section 8.2 is to turn the quadrant diagnostics of Sections 7.10 and 8.1 into a single reasoning layer in which "law" and "state" are *simultaneously* readable and jointly actionable. We fix a *paraconsistent context*

$$\mathcal{C} = \langle \mathcal{L}_{\mathcal{C}},\ \vdash_{\text{pc}},\ \Lambda_{\mathcal{C}},\ \mathrm{Adm}_{\mathcal{C}} \rangle,$$

together with a *non-explosion witness*

$$\Pi_{\text{pc}} = \langle \mathfrak{s},\ v,\ \mathsf{D} \rangle,$$

so that inconsistency of sector claims does not entail triviality (e.g. in LP/FDE with designated values $\{\mathbf{T}, \mathbf{B}\}$). This logical guardrail ensures the symmetry/SSB readings remain *co-tenable*: bulk and boundary, or distinct sector-conditioned witnesses, may disagree at finite scale without collapsing the theory. Within this setting, the familiar 2×2 "Symmetry–Asymmetry" table of Section 7.10 is read not as a mutually exclusive partition but as a *dashboard of co-tenable modalities*. This dashboard interpretation aligns with the Tetralemma "both" corner and remains stable under admissible duality transport across sectors, precisely because all statements are evaluated in the same certified arena of Section 8.1 (shared plateau, BRST-closed tags, and a sector-specific IR knob).

Parasymmetric Synthesis (operationalizing "both/and"). The operational layer of Section 8.2.1 equips the SynTh dashboard with concrete meters. For each anchor $a \in \{L, S\}$ we use a two-mode sector

$$\mathcal{H}^{(a)} = \mathrm{span}\Big\{|e_1\rangle_a,\ |e_2\rangle_a\Big\}, \qquad |e_1\rangle_a \equiv |\,\mathrm{Sym}\rangle_a, \quad |e_2\rangle_a \equiv |\,\mathrm{Break}\rangle_a,$$

and define the "both/and'' kets $|\pm\rangle_a = \frac{1}{\sqrt{2}}\Big(|e_1\rangle_a \pm |e_2\rangle_a\Big)$. The local parasymmetric projectors

$$P^{(a)}_{\mathrm{para}} = |+\rangle_a\langle+|\,, \qquad P^{(a)}_{\mathrm{antipara}} = |-\rangle_a\langle-|$$

support the *Synthesis valuation*

$$v^{(a)}_{\mathrm{Synthesis}}(\rho_a) = \mathrm{Tr}\Big(P^{(a)}_{\mathrm{para}}\rho_a\Big) = \tfrac{1}{2} + \mathrm{Re}(\rho_{12}),$$

so that $v^{(a)}_{\mathrm{Synthesis}} = 1$ for $|+\rangle_a$, $v^{(a)}_{\mathrm{Synthesis}} = \frac{1}{2}$ for an incoherent mixture, and $v^{(a)}_{\mathrm{Synthesis}} = 0$ for $|-\rangle_a$ (Bloch form $P_{\mathrm{para}} = \frac{1}{2}(I + \sigma_x)$). In this way, the logical "both/and'' of SynTh is promoted to a phase-sensitive, experimentally addressable score.

On the relational space $\mathcal{H}^{(LS)} = \mathcal{H}^{(L)} \otimes \mathcal{H}^{(S)}$ we furthermore introduce Bell-type kets and projectors that separate *opposition* (Ψ) from *agreement* (Φ) between law and state. The joint observables

$$v^{(LS)}_{\mathrm{Synthesis}} = \mathrm{Tr}\Big(P^{(LS)}_{\mathrm{para}}\rho_{LS}\Big), \qquad v^{(LS)}_{\mathrm{Agreement}} = \mathrm{Tr}\Big(P^{(LS)}_{\Phi,\mathrm{para}}\rho_{LS}\Big)$$

measure, respectively, coherent weight on the II/III diagonal and on the I/IV diagonal of the quadrant ledger—*not* as mere mixtures but including cross-term coherences. These relational meters dovetail with the sector-conditioned witnesses of Section 8.1: they can be read on the same certified plateau and with the same BRST-closed tags and IR knob r.

A *unified parasymmetric generator* on $\mathcal{H}^{(LS)}$,

$$\hat{F}^{(LS)}_{\mathrm{para}} = \lambda_L(\cdots) + \lambda_S(\cdots) + \lambda_{LS}(\cdots) + \lambda_\Phi(\cdots) + \cdots\,,$$

organizes dynamics and invariants. Whenever $[\hat{F}^{(LS)}_{\mathrm{para}}, P] = 0$ we have $\mathrm{Tr}\,\Big(P\,\rho_{LS}(t)\Big) = \mathrm{const}$. In open-system language, the same statement reads $\mathcal{L}^\dagger(P) = 0$ for the Lindbladian adjoint. Applying these to the projectors $P^{(LS)}_{\mathrm{para}}$ and $P^{(LS)}_{\Phi,\mathrm{para}}$ yields explicit time-derivative identities for $v^{(LS)}_{\mathrm{Synthesis}}$ and $v^{(LS)}_{\mathrm{Agreement}}$, thereby turning them into conserved (or slowly varying, ledger-audited) diagnostics when the corresponding symmetry conditions are met on the plateau of Section 7.10.

Finally, the parasymmetric valuations are interpreted as *Born weights* layered atop the paraconsistent base: they preserve $\vdash_{\mathrm{pc}}$ (no explosion) and transport functorially under the duality action $\mathbf{D}_{\mathcal{C}}$ with "scope hygiene'' enforced, so that readings remain sector-consistent across the arenas of Section 8.1. In short, **SynTh** guarantees co-tenability, while Parasymmetry supplies the projectors and scores that (i) discriminate *coherent "both''* from *incoherent mixtures* and (ii) promote these distinctions to conservation laws under suitable generators—thereby elevating the 2×2 grid into a *phase-structured, dynamical, and steerable* landscape for Law–State relations.

Definition 8.96 (Sector–certified arena and instrumentation)**.** Fix a sector $\mathscr{S}$ and a certified plateau component $\mathfrak{P} \subset \mathbb{R}^2_{(u,v)}$ with

$$u = \ln \lambda_{\mathrm{UV}}, \qquad v = \ln r,$$

where the sector-specific IR knob r is chosen as in Section 8.1 (bulk: $r = \ell$; boundary: $r = L$; anisotropic: $r = \min(\vec{\ell})$). Let the BRST-closed tags $P_x(\lambda_{\mathrm{UV}})$ and K_{IR} be fixed, together with the projection $\mathsf{P}_{\mathrm{phys}} : \mathcal{H} \to \mathcal{H}_{\mathrm{phys}}$ onto the physical cohomology. We call

$$\mathcal{I}^{[\mathscr{S}]} := \Big(\mathcal{A},\ r\ ;\ P_x(\lambda_{\mathrm{UV}}),\ K_{\mathrm{IR}}\ ;\ \mathsf{P}_{\mathrm{phys}}\Big), \qquad \mathcal{A} \in \{\mathfrak{P},\ \mathfrak{P}_\partial\},$$

the *sector instrumentation*, and we require that all sector witnesses and visibilities are evaluated using the *same* $\mathcal{I}^{[\mathscr{S}]}$ throughout (same arena, same IR knob, same BRST tags, same projection), as stipulated in Section 8.1.

Definition 8.97 (Sector–conditioned atomic predicates for SynTh)**.** Let $\mathbf{W}^{[\mathscr{S}]}$ be a sector-conditioned witness specification as in Section 8.1, producing visibilities

$$\Lambda^{[\mathscr{S}]}_{\mathrm{law}} : \mathfrak{P} \to [0,1], \qquad \Lambda^{[\mathscr{S}]}_{\mathrm{state}} : \mathfrak{P} \to [0,1],$$

evaluated on $\mathcal{H}_{\mathrm{phys}}$ using the common instrumentation $\mathcal{I}^{[\mathscr{S}]}$ from Definition 8.96. Fix certified tolerances $\varepsilon_{\mathrm{law}}, \varepsilon_{\mathrm{state}} \in (0,1)$ and write

$$|\cdot| \approx 1 \iff \big||\cdot| - 1\big| \le \varepsilon \quad \text{with the sector's certified } \varepsilon \in \{\varepsilon_{\mathrm{law}}, \varepsilon_{\mathrm{state}}\}.$$

Define the *law* and *state* atoms (SynTh inputs) by the disjunctive predicates

$$\begin{aligned} \mathrm{Law}^{[\mathscr{S}]} &:\Longleftrightarrow \ \Big(\ |\Lambda^{[\mathscr{S}]}_{\mathrm{law}}| \approx 1\ \Big) \ \vee \ \Big(\ \textit{heals under } r\ \Big), \\ \mathrm{State}^{[\mathscr{S}]} &:\Longleftrightarrow \ \Big(\ |\Lambda^{[\mathscr{S}]}_{\mathrm{state}}| \approx 1\ \Big) \ \vee \ \Big(\ \textit{decoheres}\ \Big), \end{aligned}$$

where the asymptotic clauses use the sector's IR knob r (also fixed by $\mathcal{I}^{[\mathscr{S}]}$):

- *heals under* r: there exist $C, \beta > 0$ such that with $d^{[\mathscr{S}]}_{\mathrm{law}}(r) := 1 - |\Lambda^{[\mathscr{S}]}_{\mathrm{law}}(r)|$ one has

  $$d^{[\mathscr{S}]}_{\mathrm{law}}(r) \ \le\ C\, r^{-\beta} \qquad (r \to \infty),$$

  i.e. the sector's law deficit decays along r as prescribed in Section 8.1;
- *decoheres*: along the thermodynamic limit one has

  $$\Lambda^{[\mathscr{S}]}_{\mathrm{state}}(V) \ \longrightarrow\ 0 \qquad (V \uparrow \infty),$$

  with the same instrumentation and evaluation arena as above (see the QII criterion in Section 8.1).

These atoms are read *on the plateau* and are understood to be *co-tenable* under the paraconsistent entailment $\vdash_{\mathrm{pc}}$ provided by **SynTh** (no explosion).

Definition 8.98 (Simultaneous Synthesis connective $\oplus$)**.** For a fixed sector $\mathscr{S}$ and a certified plateau $\mathfrak{P}$ with instrumentation $\mathcal{I}^{[\mathscr{S}]}$ as in Definition 8.96, the *Simultaneous Synthesis* judgement at the SynTh layer is the composite assertion

$$\mathrm{Syn}^{[\mathscr{S}]} \;:=\; \mathrm{Law}^{[\mathscr{S}]} \;\oplus\; \mathrm{State}^{[\mathscr{S}]},$$

which *by definition* means:

evaluate the sector-conditioned predicates $\mathrm{Law}^{[\mathscr{S}]}$ and $\mathrm{State}^{[\mathscr{S}]}$ in the **same** arena $\mathcal{A} \in \{\mathfrak{P}, \mathfrak{P}_\partial\}$, with the **same** BRST/plateau certification and IR knob r, using the **same** tags $P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}$ and projection to $\mathcal{H}_{\mathrm{phys}}$; then record their joint status as a coherent, co-tenable dashboard entry within the paraconsistent context $\mathcal{C}$ (no exclusivity is imposed; co-tenability per $\vdash_{\mathrm{pc}}$).

This connective introduces no additional identification beyond shared instrumentation: it packages the two certified sector predicates into a single SynTh premise that can be projected onto either component without entailment collapse.

Definition 8.99 (SynTh rules for Simultaneous Synthesis)**.** Let **SynTh** endow the context $\mathcal{C}$ with a paraconsistent entailment $\vdash_{\mathrm{pc}}$, and let $\mathrm{Law}^{[\mathscr{S}]}$, $\mathrm{State}^{[\mathscr{S}]}$, and $\mathrm{Syn}^{[\mathscr{S}]} := \mathrm{Law}^{[\mathscr{S}]} \oplus \mathrm{State}^{[\mathscr{S}]}$ be as in Definitions 8.97 and 8.98. The following schematic rules govern reasoning at the SynTh layer:

$$\frac{\mathcal{C} \vdash_{\mathrm{pc}} \mathrm{Law}^{[\mathscr{S}]} \qquad \mathcal{C} \vdash_{\mathrm{pc}} \mathrm{State}^{[\mathscr{S}]}}{\mathcal{C} \vdash_{\mathrm{pc}} \mathrm{Syn}^{[\mathscr{S}]}} \quad (\oplus\text{-Intro})$$

$$\frac{\mathcal{C} \vdash_{\mathrm{pc}} \mathrm{Syn}^{[\mathscr{S}]}}{\mathcal{C} \vdash_{\mathrm{pc}} \mathrm{Law}^{[\mathscr{S}]}} \quad (\oplus\text{-Proj}_L) \qquad\qquad \frac{\mathcal{C} \vdash_{\mathrm{pc}} \mathrm{Syn}^{[\mathscr{S}]}}{\mathcal{C} \vdash_{\mathrm{pc}} \mathrm{State}^{[\mathscr{S}]}} \quad (\oplus\text{-Proj}_S)$$

$$\frac{\mathcal{C} \vdash_{\mathrm{pc}} \varphi^{[\mathscr{S}_1]} \qquad \mathcal{C} \vdash_{\mathrm{pc}} \neg\varphi^{[\mathscr{S}_2]}}{\begin{array}{c}\textit{No-Explosion: } \mathcal{C} \nvdash_{\mathrm{pc}} \psi \text{ for arbitrary } \psi \\ \textit{Audit obligation: } \mathrm{ReAudit}\big(\mathcal{A}_1, \mathcal{A}_2;\ r_1, r_2\big)\end{array}} \quad (\textsc{No-Explosion/Audit})$$

Here $\varphi^{[\mathscr{S}_i]}$ stands for any sector-conditioned claim (e.g. a law or state predicate) evaluated with the sector's realized weave configuration $\mathcal{I}^{[\mathscr{S}_i]} = (\mathcal{A}_i, r_i; P_x, K_{\mathrm{IR}}; \mathsf{P}_{\mathrm{phys}})$ from Definition 8.96. The No-Explosion clause states that contradictory sector claims (e.g. boundary vs. bulk) do *not* entail triviality in $\vdash_{\mathrm{pc}}$; instead, they *trigger* the operational Audit obligation to reconcile arenas/IR knobs on the certified plateau as prescribed by Sections 8.1 and 8.2. Consequently, the quadrant grid of Section 7.10 is treated as a *dashboard*, not a partition: law and state readings are jointly admissible inputs even when they initially sit in different quadrants prior to IR reconciliation.

Definition 8.100 (Sector-conditioned synthesis object)**.** Let $\mathscr{S}$ be a sector with witness specification $\mathbf{W}^{[\mathscr{S}]}$ as in Section 8.1, and let $\mathcal{I}^{[\mathscr{S}]}$ be the common instrumentation from Definition 8.96 on a certified plateau component $\mathfrak{P}$. We define the *sector-conditioned synthesis object*

$$\boxed{\mathfrak{S}^{[\mathscr{S}]}(\mathfrak{P}) \;:=\; \Big(\Lambda^{[\mathscr{S}]}_{\mathrm{law}},\ \Lambda^{[\mathscr{S}]}_{\mathrm{state}}\ ;\ \mathcal{W}^{[\mathscr{S}]},\ \mathcal{Z}^{[\mathscr{S}]}\ ;\ \mathcal{A},\ r\Big)}$$

with the following components:

- $\mathcal{A} \in \{\mathfrak{P},\ \mathfrak{P}_\partial\}$ is the arena; r is the sector's IR knob (bulk $r{=}\ell$, boundary $r{=}L$, anisotropic $r{=}\min(\vec{\ell})$);
- $\mathcal{W}^{[\mathscr{S}]}$ is the *scale-tagged Ward ledger* (improved current/divergence statements) consistent with the consolidated laws of Section 7.10;
- $\mathcal{Z}^{[\mathscr{S}]}$ is the *light-sector spectral certificate* on $\mathcal{H}_{\text{phys}}$ (poles, residue matrices $Z \succeq 0$, NG/pNG census, hyperbolicity margin).

All entries of $\mathfrak{S}^{[\mathscr{S}]}(\mathfrak{P})$ are evaluated using the *same* BRST-closed tags $P_x(\lambda_{\text{UV}})$, K_{IR} and the same projection P_{phys} as stipulated in Section 8.1.

Proposition 8.101 (Gauge/Nielsen invariance)**.** *Suppose $P_x(\lambda_{\text{UV}})$ and K_{IR} are BRST-closed and all evaluations are made on $\mathcal{H}_{\text{phys}}$ with the common instrumentation $\mathcal{I}^{[\mathscr{S}]}$ of Definition 8.96. Then the synthesis object $\mathfrak{S}^{[\mathscr{S}]}(\mathfrak{P})$ of Definition 8.100 is invariant under changes of the gauge fermion (Nielsen-flat).* Sketch. *The BRST closure and projection render the sector visibilities and their Ward/spectral ledgers independent of gauge-fermion deformations; this is precisely the sectorised adaptation of the invariance arguments used to establish quadrant correctness in Section 7.10, combined with the witness-level invariance requirements of Section 8.1.*

Definition 8.102 (Synthesis resolver and explanatory class)**.** Let $\mathscr{S}$ be a sector with synthesis object $\mathfrak{S}^{[\mathscr{S}]}(\mathfrak{P})$ as in Definition 8.100. The *resolver*

$$\mathcal{R}^{[\mathscr{S}]}:\ \mathfrak{S}^{[\mathscr{S}]}(\mathfrak{P}) \ \longmapsto\ \Big(\mathrm{Quad}^{[\mathscr{S}]},\ \mathsf{Expl}^{[\mathscr{S}]}\Big)$$

is defined by applying the canonical quadrant rules of Section 8.1 (QI–QIV) to the pair $\big(\Lambda^{[\mathscr{S}]}_{\text{law}}, \Lambda^{[\mathscr{S}]}_{\text{state}}\big)$, including the r-healing test for the law deficit, and returning the *explanatory class*

$$\mathsf{Expl}^{[\mathscr{S}]} \ :=\ \Big(\ \underbrace{\mathcal{W}^{[\mathscr{S}]}\ \textbf{status}}_{\text{conserved vs. adjoint-source}}\ ;\ \underbrace{\mathcal{Z}^{[\mathscr{S}]}\ \textbf{census}}_{\{\text{NG,pNG,none}\},\ Z\succeq 0,\ \text{hyperbolicity margin}}\ \Big).$$

Concretely, on any certified plateau point (or connected component) the mapping satisfies:

(i) QI
$\Rightarrow \mathcal{W}$: conserved improved current; $\mathcal{Z}$: no NG/pNG.

(ii) QII
$\Rightarrow \mathcal{W}$: conserved; $\mathcal{Z}$: NG in the correlational channel (gauged generators lift to BEH).

(iii) QIII
$\Rightarrow \mathcal{W}$: adjoint divergence $\to 0$ along r; $\mathcal{Z}$: no NG/pNG; law heals.

(iv) QIV
$\Rightarrow \mathcal{W}$: non-vanishing adjoint–covariant source on $\mathfrak{P}$; $\mathcal{Z}$: pNG band obeying GMOR on $\mathfrak{P}$.

Equivalence. This sectorised construction realises the diagnostics equivalence promised in Section 7.10: the quadrant assignment from visibilities matches the Ward and spectral ledgers on the same realized weave configuration.

Definition 8.103 (Joint Simultaneous Synthesis)**.** Let $\{\mathscr{S}_i\}_{i\in I}$ be a (finite) family of sectors with certified plateaus $\{\mathfrak{P}_i\}$ and arenas $\{\mathcal{A}_i\}$, each equipped with a sector-conditioned synthesis object $\mathfrak{S}^{[\mathscr{S}_i]}(\mathfrak{P}_i)$ as in Definition 8.100. Define the *joint arena* and *joint plateau*

$$\mathcal{A}_\cap := \bigcap_{i\in I} \mathcal{A}_i, \qquad \mathfrak{P}_\cap := \bigcap_{i\in I} \mathfrak{P}_i.$$

The *joint synthesis object* is

$$\mathfrak{S}^{[I]}(\mathfrak{P}_\cap) := \prod_{i\in I} \mathfrak{S}^{[\mathscr{S}_i]}(\mathfrak{P}_\cap),$$

where each factor is the restriction/re-evaluation of $\mathfrak{S}^{[\mathscr{S}_i]}(\mathfrak{P}_i)$ on the certified intersection $\mathfrak{P}_\cap$ using the same BRST-closed tags and the joint instrumentation on $\mathcal{A}_\cap$. Reasoning is carried out in the paraconsistent context $\mathcal{C}$ (SynTh), so contradictory sector labels do not entail triviality; instead they trigger *arena reconciliation* (e.g. boundary L vs. bulk ℓ) or *IR-knob refinement* until the QIII/QIV split stabilises, as in Section 8.1.

Proposition 8.104 (Joint completeness and stability)**.** *Assume: (i) each $\mathfrak{P}_i$ is certified as in Section 8.1; (ii) the sector r-healing tests are well-posed; and (iii) evaluations use BRST-closed tags with projection to $\mathcal{H}_{\text{phys}}$ (cf. Proposition 8.101). Then the product resolver*

$$\mathcal{R}^{[I]} := \prod_{i\in I} \mathcal{R}^{[\mathscr{S}_i]} : \ \mathfrak{S}^{[I]}(\mathfrak{P}_\cap) \longmapsto \prod_{i\in I} \left(\mathrm{Quad}^{[\mathscr{S}_i]},\ \mathsf{Expl}^{[\mathscr{S}_i]}\right)$$

yields a complete *set of sector labels* $\{\mathrm{Quad}^{[\mathscr{S}_i]}\}_{i\in I}$ *and associated explanatory classes that is stable under:*

(i) admissible gauge-fermion deformations (Nielsen-flatness by Proposition 8.101);

(ii) admissible retunings of the BRST-closed tags (P_x, K_{IR}) that preserve certification; and

(iii) coarsenings $\mathfrak{P}_i \mapsto \tilde{\mathfrak{P}}_i \subseteq \mathfrak{P}_i$ that maintain plateau certification and r-healing well-posedness.

Definition 8.105 (Operational template (sector-conditioned, SynTh))**.** Let $\mathscr{S}$ be a fixed sector with instrumentation $\mathcal{I}^{[\mathscr{S}]}$ from Definition 8.96 and witnesses $\mathbf{W}^{[\mathscr{S}]}$ as in Definition 8.97. The following *repeatable construction* instantiates §8.2 within the sector framework of Section 8.1.

Step 0 — Fix the arena and certify. Choose $\mathcal{A} \in \{\mathfrak{P}, \mathfrak{P}_\partial\}$, the sector IR knob r (bulk: $r = \ell$; boundary: $r = L$; anisotropic: $r = \min(\vec{\ell})$), and tolerances $\varepsilon_{\text{law}}, \varepsilon_{\text{state}} \in (0,1)$. Certify the plateau(s) $\mathfrak{P}$ (and, if applicable, $\mathfrak{P}_\partial$) according to Section 8.1 (cone preservation, BRST-closure, Nielsen-flat diagnostics).

Step 1 — Build the sector witness pair. Using $\mathbf{W}^{[\mathscr{S}]}$ and the BRST-closed tags $P_x(\lambda_{\text{UV}}), K_{\text{IR}}$ with projection to $\mathcal{H}_{\text{phys}}$, compute on $\mathfrak{P}$

$$\Lambda^{[\mathscr{S}]}_{\text{law}}(u,v), \qquad \Lambda^{[\mathscr{S}]}_{\text{state}}(u,v),$$

with $(u,v) = (\ln \lambda_{\text{UV}}, \ln r)$.

Step 2 — Attach Ward and spectral ledgers. Construct the scale-tagged Ward ledger $\mathcal{W}^{[\mathscr{S}]}$ (improved current/divergence on $\mathfrak{P}$ or along r) and the light-sector spectral ledger $\mathcal{Z}^{[\mathscr{S}]}$ (poles,

residue matrices $Z \succeq 0$, NG/pNG census, hyperbolicity margin) exactly as consolidated in Section 7.10.

Step 3 — Form the synthesis object and resolve. Assemble the synthesis object

$$\mathfrak{S}^{[\mathscr{S}]}(\mathfrak{P}) = \Big(\Lambda^{[\mathscr{S}]}_{\text{law}}, \Lambda^{[\mathscr{S}]}_{\text{state}} \;;\; \mathcal{W}^{[\mathscr{S}]}, \mathcal{Z}^{[\mathscr{S}]} \;;\; \mathcal{A}, r\Big)$$

as in Definition 8.100 and apply the resolver $\mathcal{R}^{[\mathscr{S}]}$ of Definition 8.102 to obtain

$$\Big(\text{Quad}^{[\mathscr{S}]},\ \mathsf{Expl}^{[\mathscr{S}]}\Big).$$

Step 4 — SynTh reasoning ($\oplus$)**.** Adopt $\text{Syn}^{[\mathscr{S}]} = \text{Law}^{[\mathscr{S}]} \oplus \text{State}^{[\mathscr{S}]}$ (cf. Definition 8.98) as the *joint premise* for higher-level conclusions. Inter-sector contradictions (e.g. "QI in bulk, QIV on boundary") are *co-tenable* in the paraconsistent context and should trigger *arena/knob audit* until discrepancies resolve into QIII-type healing or a certified QIV persistence, in line with Section 8.1.

Definition 8.106 (Canonical certification predicates)**.** For a sector $\mathscr{S}$ with tolerances $\varepsilon_{\text{law}}, \varepsilon_{\text{state}} \in (0,1)$, define the *tolerance predicates*

$$\chi^{[\mathscr{S}]}_{\text{law}} := \mathbf{1}\Big[\,|\Lambda^{[\mathscr{S}]}_{\text{law}}| \geq 1 - \varepsilon_{\text{law}}\,\Big], \qquad \chi^{[\mathscr{S}]}_{\text{state}} := \mathbf{1}\Big[\,|\Lambda^{[\mathscr{S}]}_{\text{state}}| \geq 1 - \varepsilon_{\text{state}}\,\Big], \tag{8.16}$$

and the *healing predicate* along the sector IR knob r,

$$d^{[\mathscr{S}]}_{\text{law}}(r) := 1 - |\Lambda^{[\mathscr{S}]}_{\text{law}}(r)|, \qquad d^{[\mathscr{S}]}_{\text{law}}(r) \;\leq\; C\, r^{-\beta} \quad (r \to \infty), \tag{8.17}$$

for some $C, \beta > 0$. The resolver $\mathcal{R}^{[\mathscr{S}]}$ maps $(\chi^{[\mathscr{S}]}_{\text{law}}, \chi^{[\mathscr{S}]}_{\text{state}}, d^{[\mathscr{S}]}_{\text{law}})$ to QI–QIV exactly as in Section 8.1.

Definition 8.107 (Worked sector sketches (non-parasymmetric))**.** The following certified instances illustrate Definition 8.105 purely at the SynTh level (parasymmetric meters set aside).

(a) Bulk sector (canonical). Witness pair $\Big(\Lambda^{[\text{bulk}]}_{\text{law}}(\ell), \Lambda^{[\text{bulk}]}_{\text{state}}\Big)$ on $\mathfrak{P}$ with $r = \ell$. If $|\Lambda^{[\text{bulk}]}_{\text{law}}| \approx 1$ while $\Lambda^{[\text{bulk}]}_{\text{state}}(V) \to 0$, then $\text{Syn}^{[\text{bulk}]}$ resolves to **QII** via $\mathcal{R}^{[\text{bulk}]}$; $\mathcal{W}^{[\text{bulk}]}$ is conserved and $\mathcal{Z}^{[\text{bulk}]}$ exhibits NG poles in the correlational channel (lifting to BEH when gauged), as per Section 7.10.

(b) Boundary sector. Witness pair $\Big(\Lambda^{[\partial]}_{\text{law}}(L), \Lambda^{[\partial]}_{\text{state}}\Big)$ on $\mathfrak{P}_\partial$ with $r = L$. If $d^{[\partial]}_{\text{law}}(L) \sim L^{-\beta}$ and $|\Lambda^{[\partial]}_{\text{state}}| \approx 1$, then $\text{Syn}^{[\partial]}$ resolves to **QIII**; the explanatory class $\mathsf{Expl}^{[\partial]}$ records adjoint divergence $\to 0$ along L and no NG/pNG, consistent with emergent symmetry at large L.

(c) Anisotropic sector (spin–orbit example). Witness pair $\Big(\Lambda^{[\text{aniso}]}_{\text{law}}(\vec{\ell}), \Lambda^{[\text{aniso}]}_{\text{state}}\Big)$ with $r = \min(\vec{\ell})$. If $|\Lambda^{[\text{aniso}]}_{\text{state}}| \approx 1$ and $|\Lambda^{[\text{aniso}]}_{\text{law}}| \leq 1 - \delta_\star$ uniformly on $\mathfrak{P}$, then $\text{Syn}^{[\text{aniso}]}$ resolves to **QIV**; $\mathcal{W}^{[\text{aniso}]}$ shows a persistent adjoint–covariant source and $\mathcal{Z}^{[\text{aniso}]}$ registers a pNG band with GMOR behaviour on the plateau.

Definition 8.108 (Diagnostics and edge handling (SynTh discipline))**.** *Lower-right anomaly* (state decoheres + non-healing law deficit): treat the configuration as *non-certified*; re-audit plateau certification, witnesses, or arena choice per Section 7.10. *Inter-sector disagreement*: SynTh does not collapse; perform *arena reconciliation* (align $\mathcal{A}$ and r) and prefer joint plateaus $\mathfrak{P}_\cap$ from Definition 8.103 until the QIII/QIV distinction stabilises.

Remark 8.109 (Summary for insertion into §8.2)**.** Simultaneous Synthesis *law* $\oplus$ *state* treats the sector-conditioned predicates as a *co-tenable* pair at the SynTh layer, evaluated on the *same* certified arena and IR knob (Section 8.1). The synthesis object $\mathfrak{S}^{[\mathscr{S}]}(\mathfrak{P})$ bundles visibilities with *Ward* and *spectral* ledgers; the resolver $\mathcal{R}^{[\mathscr{S}]}$ maps this to a quadrant label and an *explanatory class*, realising the equivalences of Section 7.10 sector-by-sector. Paraconsistent reasoning ensures non-explosive multi-sector inference; QIII vs. QIV is always separated by the *healing* test in Section 8.1.

At a high level, Section 8.2 turns the quadrant machinery of Sections 7.10 and 8.1 into a single, coherent way to *think* about symmetry—one that treats "law" and "state" as jointly readable, jointly actionable inputs. The central move is the *Simultaneous Synthesis* judgement

$$\mathrm{Syn}^{[\mathscr{S}]} = \mathrm{Law}^{[\mathscr{S}]} \ \oplus \ \mathrm{State}^{[\mathscr{S}]},$$

which simply says: in a fixed sector $\mathscr{S}$, read the *same* arena ($\mathcal{A} \in \{\mathfrak{P}, \mathfrak{P}_\partial\}$), with the *same* BRST-closed tags $P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}$, through the *same* IR knob r, and treat the resulting pair of visibilities $\left(\Lambda^{[\mathscr{S}]}_{\mathrm{law}}, \Lambda^{[\mathscr{S}]}_{\mathrm{state}}\right)$ as **co-tenable information** in SynTh. Nothing about this forces a partition; rather, it elevates the grid to a *dashboard* whose two dials carry logically compatible, yet conceptually distinct, content about dynamics (the law) and about the realized phase (the state).

Why is this separation important? Because the physics really does bifurcate along those lines. The *law dial* is designed to detect explicit breaking in the microscopic or effective dynamics; the *state dial* reacts to whether the realized macroscopic phase decoheres into a symmetry-broken mixture as $V \uparrow \infty$. In practice, that means a theory may *respect* a symmetry ($|\Lambda^{[\mathscr{S}]}_{\mathrm{law}}| \approx 1$) while its ground sector **chooses** a broken phase ($\Lambda^{[\mathscr{S}]}_{\mathrm{state}}(V) \to 0$): that is precisely QII. Conversely, the law may be explicitly imperfect but "heals" as we push deeper into the sector's IR knob r (QIII). Keeping these channels distinct, while reading them *simultaneously*, is what makes the synthesis robust and explanatory.

Technically, the synthesis works because it is anchored on a *certified plateau* $\mathfrak{P}$. Once we restrict to $\mathfrak{P}$, every quantity that we read—$\Lambda^{[\mathscr{S}]}_{\mathrm{law}}$, $\Lambda^{[\mathscr{S}]}_{\mathrm{state}}$, the sector Ward sources, and the light-sector spectrum on $\mathcal{H}_{\mathrm{phys}}$—is (i) insensitive to gauge-fermion changes (Nielsen-flat), (ii) filtered to respect causal cones, and (iii) stable under small changes of $(\lambda_{\mathrm{UV}}, r)$. This is not cosmetic: it converts "diagnostics that could be scheme-dependent" into *observables* that convey the same story across gauges and regulators.

Seen this way, the quadrant labels (QI–QIV) are just *surface summaries* of a deeper triangulation. The synthesis object for sector $\mathscr{S}$,

$$\mathfrak{S}^{[\mathscr{S}]}(\mathcal{A}) \equiv \left(\Lambda^{[\mathscr{S}]}_{\mathrm{law}}, \Lambda^{[\mathscr{S}]}_{\mathrm{state}};\ \text{Ward ledger } \mathcal{W}^{[\mathscr{S}]},\ \text{spectral ledger } \mathcal{Z}^{[\mathscr{S}]};\ \mathcal{A}, r\right),$$

collects (1) the two dials, (2) the *Ward* statement that tells whether the improved current is conserved or has a non-vanishing adjoint–covariant source, and (3) the *spectral* statement

that tells which light modes truly exist on $\mathcal{H}_{\text{phys}}$ (NG, pNG, or none), with residue matrices $Z \succeq 0$ and hyperbolicity margins recorded. The equivalence result summarized in Section 7.10 then says: on $\mathfrak{P}$, these three views agree. The grid position is consistent with both the Ward ledger and the spectral ledger—*and that* is where the explanatory power comes from.

Each quadrant thus acquires a crisp, three-fold meaning. **QI** is the "all clear": law saturated, state saturated, conserved improved current, no NG/pNG in the physical spectrum. **QII** is the *SSB arrow*: law saturated, state decoheres, improved current still conserved, and NG poles appear in the correlational channel (which become BEH masses if the corresponding generators are gauged). **QIII** is *emergent symmetry*: the law dial shows a **finite-scale** deficit that *heals* along the sector's IR direction r (e.g. ℓ, L, or $\min(\vec{\ell})$), while the state dial saturates; correspondingly, the Ward divergence tends to zero along r, and the spectrum carries no bona fide NG or pNG modes. **QIV** is *persistent explicit breaking*: the law dial stays below $1-\delta_\star$ on $\mathfrak{P}$, the state dial saturates, the Ward source persists, and the spectrum exhibits a pNG band that obeys a GMOR-type relation on the plateau.

Two conceptual implications flow immediately. First, *QIII vs QIV is not a matter of taste*; it is decided operationally by the **healing test**

$$d_{\text{law}}^{[\mathscr{S}]}(r) := 1 - |\Lambda_{\text{law}}^{[\mathscr{S}]}(r)| \quad \overset{?}{\leq} \quad C\, r^{-\beta} \quad (r \to \infty),$$

performed in the *sector's own* IR knob r. This converts a semantic debate ("is the symmetry really there?") into a *rate-of-convergence measurement*. Second, because the whole construction lives on $\mathcal{H}_{\text{phys}}$ with BRST-closed tags, the outcome is *not* an artefact of gauge choices or regulator idiosyncrasies; it reflects the physics accessible to observers confined to the physical cohomology.

The sector-conditioned formulation is not a mere relabelling; it is what makes the synthesis *portable* across arenas. In *bulk*, read $(\Lambda_{\text{law}}^{[\text{bulk}]}, \Lambda_{\text{state}}^{[\text{bulk}]})$ on $\mathfrak{P}$ with $r = \ell$. On a *boundary*, switch to $\mathfrak{P}_\partial$ and $r = L$. In *anisotropic* situations, take $r = \min(\vec{\ell})$ (or the sector's prescribed dilation). The grid geometry is *identical*, but the *dial design*—i.e., the witness map $\mathbf{W}^{[\mathscr{S}]}$—is tailored to the sector so that the same four labels carry the same physical meaning after projection to $\mathcal{H}_{\text{phys}}$.

This portability pays off when different sectors disagree. Suppose the bulk sits in QI while the boundary sits in QIV. In a classical, explosive logic, this looks contradictory; in SynTh, it is *co-tenable* information: the theory's law respects the symmetry in bulk but exhibits a *boundary-pinned explicit deficit*. The synthesis asks *not* to average those away, but to *explain* them: the Ward source is boundary-supported; the pNG band is boundary-localized; the bulk has neither. Conversely, if both sides look like QIII, the synthesis teaches to expect *reconvergence* under deeper IR dilation (larger L, longer ℓ, stronger anisotropic windowing), and to look for that decay of d_{law} rather than trying to force a premature QI judgement.

From a workflow perspective, the synthesis gives a clean loop. *Fix the arena and certify* $(\mathcal{A}, r, \mathfrak{P})$. *Build the dials* via $\mathbf{W}^{[\mathscr{S}]}$ and compute $\Lambda_{\text{law}}^{[\mathscr{S}]}, \Lambda_{\text{state}}^{[\mathscr{S}]}$. *Attach ledgers* $\mathcal{W}^{[\mathscr{S}]}, \mathcal{Z}^{[\mathscr{S}]}$. *Resolve* to a quadrant with an *explanatory class* (conservation/source; NG/pNG/none). And then, crucially, *reason at the SynTh level* with Law $\oplus$ State as a *single premise*. This turns the grid from a post-hoc labeler into a *design tool*: we can now *steer* simulations or experiments along r to make a QIII heal, or along control parameters to convert a QIV source into a QIII decay.

Even at this high level, the spectral ledger has tangible, testable content. In QII, the

current–order-parameter form factor saturates the NG pole; in QIV, the same structure reappears as a *pseudogap/pseudo-NG* with a GMOR slope on $\mathfrak{P}$. The residues $Z \succeq 0$ and hyperbolicity margins provide sanity checks that the light content is not a gauge remnant. Because the ledgers are tied to the same $\mathfrak{P}$ that produced the dials, the story is *closed*: we can cross-check the grid assignment by looking either at visibilities or at Ward–spectrum, and we should get the same verdict.

Intuitively, we can picture the whole construction as installing two *calibrated meters* on the same instrument diagnostic realized reference configuration. One meter measures how faithfully the dynamics realises the symmetry (the *law* meter); the other meter measures whether the realized macroscopic phase keeps that symmetry coherent in the thermodynamic limit (the *state* meter). The *sector* tells us what diagnostic realized reference configuration we're working on (bulk, boundary, anisotropic), and the *IR knob* r tells us which way to turn to resolve ambiguous readings. The SynTh connective $\oplus$ simply instructs us to *read both meters together* and use them jointly, without forcing a false choice when reality presents a both/and.

The implications extend to modelling strategy. Want to engineer an *emergent symmetry* (e.g. a gauge symmetry)? Aim for QIII by designing flows where $d_{\mathrm{law}}^{[\mathscr{S}]}(r)$ decays with a controllable exponent β while keeping $|\Lambda_{\mathrm{state}}^{[\mathscr{S}]}| \approx 1$. Want to *protect* a symmetry? Target QI, and use the Ward ledger to audit any drift off conservation. Diagnosing *spontaneous breaking* (SSB)? Expect QII, but confirm it spectrally (true NG) rather than by visibility alone. Hunting *explicit breakings* that masquerade as SSB? QIV will expose them, because the Ward source refuses to vanish and the spectrum carries pNG instead of NG on $\mathfrak{P}$.

Finally, the synthesis clarifies what to do when the lower-right corner (state decoheres *and* law deficit persists) shows up. Within Section 7.10 this is typically a *non-certified* configuration, and the synthesis treats it as a *prompt* to re-audit the arena, the witness map, or the plateau itself—not as a paradox to be resolved by argument. In other words, the logic (SynTh) and the metrology (plateau certification, BRST closure) are aligned: if the instruments disagree in a way the theory says they shouldn't, fix the instruments before inferring physics.

In sum, Section 8.2 equips the sector-conditioned framework with a reasoning layer that is as disciplined as its metrology. The *Simultaneous Synthesis* Law $\oplus$ State treats the two dials as joint, gauge-safe evidence; the *resolver* maps them (together with Ward and spectral ledgers) into QI–QIV with clear operational meaning; and the *IR-healing* rule separates emergent symmetry (QIII) from persistent explicit breaking (QIV). The result is a high-level picture that is easy to *read* but hard to *fool*—and that is exactly what we want when turning symmetry principles into working diagnostics across bulk, boundary, and anisotropic sectors.

8.4 Parasymmetric Synthesis of Laws and States (Sector-conditioned Overlay)

Definition 8.110 (Sector–parasymmetric modal spaces and meters)**.** Fix a sector $\mathscr{S}$ with realized weave configuration $\mathcal{I}^{[\mathscr{S}]} = (\mathcal{A}, \mathfrak{P}; r; P_x, K_{\mathrm{IR}}; \mathsf{P}_{\mathrm{phys}})$ as in Definition 8.96. For each

anchor $a \in \{L, S\}$ introduce a two-mode modal space

$$\mathcal{H}^{(a)[\mathscr{S}]} = \mathrm{span}\{|e_1\rangle_a, |e_2\rangle_a\}, \qquad |e_1\rangle_a \equiv |\,\mathrm{Sym}\rangle_a, \quad |e_2\rangle_a \equiv |\,\mathrm{Break}\rangle_a,$$

and the "both/and" (parasymmetric) kets $|\pm\rangle_a = \frac{1}{\sqrt{2}}(|e_1\rangle_a \pm |e_2\rangle_a)$. Define the local parasymmetric projectors

$$P^{(a)}_{\mathrm{para}} = |+\rangle_a\langle+|\,, \qquad P^{(a)}_{\mathrm{antipara}} = |-\rangle_a\langle-|,$$

and the *local parasymmetric valuations* (meters) on density operators $\rho^{(a)[\mathscr{S}]}$:

$$v^{(a)[\mathscr{S}]}_{\mathrm{Synthesis}}(\rho^{(a)[\mathscr{S}]}) = \mathrm{Tr}\Big(P^{(a)}_{\mathrm{para}}\rho^{(a)[\mathscr{S}]}\Big) = \tfrac{1}{2} + \mathrm{Re}\Big(\rho^{(a)[\mathscr{S}]}_{12}\Big).$$

On the relational space $\mathcal{H}^{(LS)[\mathscr{S}]} = \mathcal{H}^{(L)[\mathscr{S}]} \otimes \mathcal{H}^{(S)[\mathscr{S}]}$, introduce Bell-type projectors

$$\begin{aligned} &P^{(LS)}_{\mathrm{para}} && \text{(coherent } \textit{opposition} \text{ on the II/III diagonal)},\\ &P^{(LS)}_{\Phi,\mathrm{para}} && \text{(coherent } \textit{agreement} \text{ on the I/IV diagonal)}. \end{aligned}$$

and define the *relational parasymmetric meters* for density operators $\rho^{(LS)[\mathscr{S}]}$:

$$v^{(LS)[\mathscr{S}]}_{\mathrm{Synthesis}} = \mathrm{Tr}\Big(P^{(LS)}_{\mathrm{para}}\rho^{(LS)[\mathscr{S}]}\Big), \qquad v^{(LS)[\mathscr{S}]}_{\mathrm{Agreement}} = \mathrm{Tr}\Big(P^{(LS)}_{\Phi,\mathrm{para}}\rho^{(LS)[\mathscr{S}]}\Big).$$

All evaluations are performed on the same certified arena $\mathcal{A} \in \{\mathfrak{P}, \mathfrak{P}_\partial\}$, with the same IR knob r, BRST-closed tags (P_x, K_{IR}), and projection to $\mathcal{H}_{\mathrm{phys}}$, cf. Section 8.2.1.

Definition 8.111 (Sector–parasymmetric calibration from generalized witness)**.** Let $\mathbf{W}^{[\mathscr{S}]}_{\mathrm{gen}}$ be the generalized witness of Section 8.3, producing the consolidated outputs

$$\Big(\Lambda^{[\mathscr{S}]}_{\mathrm{law}}, \Lambda^{[\mathscr{S}]}_{\mathrm{state}};\ \mathcal{W}^{[\mathscr{S}]}, \mathcal{Z}^{[\mathscr{S}]}\Big)$$

on the common instrumentation $\mathcal{I}^{[\mathscr{S}]}$. A *sector–parasymmetric calibration* is a completely positive, trace-preserving map

$$\kappa^{[\mathscr{S}]}: \ \Big(\Lambda^{[\mathscr{S}]}_{\mathrm{law}}, \Lambda^{[\mathscr{S}]}_{\mathrm{state}};\ \mathcal{W}^{[\mathscr{S}]}, \mathcal{Z}^{[\mathscr{S}]}\Big) \ \longrightarrow\ \Big(\rho^{(L)[\mathscr{S}]},\ \rho^{(S)[\mathscr{S}]},\ \rho^{(LS)[\mathscr{S}]}\Big),$$

chosen so that:

(C1) Gauge/Nielsen flatness. If (P_x, K_{IR}) are BRST-closed and evaluations are on $\mathcal{H}_{\mathrm{phys}}$, then the image densities are invariant under gauge-fermion changes (inherit the flatness of Proposition 8.101).

(C2) Sector functoriality. Admissible moves of $\mathcal{A}$ and r (bulk $\leftrightarrow$ boundary; anisotropic refinements) induce well-defined morphisms

$$\kappa^{[\mathscr{S}]}_{\mathcal{A},r} \ \Rightarrow\ \kappa^{[\mathscr{S}]}_{\mathcal{A}',r'} \quad \text{that preserve the quadrant label via } \mathcal{R}^{[\mathscr{S}]} \text{ (Definition 8.102)}.$$

(C3) Plateau monotonicity. Along sector IR dilations, the induced coherence deficits in the meters are asymptotically monotone in the sense needed to separate QIII (healing) from QIV (persistence) on certified plateaus.

Given $\kappa^{[\mathscr{S}]}$, the parasymmetric meters of Definition 8.110 are henceforth written

$$\begin{aligned} v^{(a)[\mathscr{S}]}_{\mathrm{Synthesis}} &:= v^{(a)[\mathscr{S}]}_{\mathrm{Synthesis}}\Big(\rho^{(a)[\mathscr{S}]}\Big), \\ v^{(LS)[\mathscr{S}]}_{\mathrm{Synthesis}} &:= v^{(LS)[\mathscr{S}]}_{\mathrm{Synthesis}}\Big(\rho^{(LS)[\mathscr{S}]}\Big), \\ v^{(LS)[\mathscr{S}]}_{\mathrm{Agreement}} &:= v^{(LS)[\mathscr{S}]}_{\mathrm{Agreement}}\Big(\rho^{(LS)[\mathscr{S}]}\Big). \end{aligned}$$

Proposition 8.112 (Conservation and auditability of parasymmetric meters)**.** *Let $\hat{F}^{(LS)}_{\mathrm{para}}$ be a unified parasymmetric generator on $\mathcal{H}^{(LS)[\mathscr{S}]}$ as in Section 8.2.1. If $[\hat{F}^{(LS)}_{\mathrm{para}}, P] = 0$ for $P \in \{P^{(L)}_{\mathrm{para}}, P^{(S)}_{\mathrm{para}}, P^{(LS)}_{\mathrm{para}}, P^{(LS)}_{\Phi,\mathrm{para}}\}$, then along the certified evolution on $\mathcal{A}$ one has*

$$\frac{d}{dt}\,\mathrm{Tr}\Big(P\,\rho^{(\cdot)[\mathscr{S}]}(t)\Big) = 0,$$

and, in the open-system setting with Lindbladian $\mathcal{L}$, the same holds whenever $\mathcal{L}^\dagger(P) = 0$. Thus $v^{(a)[\mathscr{S}]}_{\mathrm{Synthesis}}, v^{(LS)[\mathscr{S}]}_{\mathrm{Synthesis}}, v^{(LS)[\mathscr{S}]}_{\mathrm{Agreement}}$ are conserved (or ledger-controlled) meters on $\mathfrak{P}$, providing auditable invariants that complement the resolver of Definition 8.102.

Definition 8.113 (Parasymmetric overlay of the sector synthesis object)**.** Given $\mathfrak{S}^{[\mathscr{S}]}(\mathfrak{P})$ from Definition 8.100 and a calibration $\kappa^{[\mathscr{S}]}$ from Definition 8.111, define the *parasymmetric synthesis object*

$$\widehat{\mathfrak{S}}^{[\mathscr{S}]}(\mathfrak{P}) := \Big(\ \mathfrak{S}^{[\mathscr{S}]}(\mathfrak{P})\ ;\ v^{(L)[\mathscr{S}]}_{\mathrm{Synthesis}},\ v^{(S)[\mathscr{S}]}_{\mathrm{Synthesis}}\ ;\ v^{(LS)[\mathscr{S}]}_{\mathrm{Synthesis}},\ v^{(LS)[\mathscr{S}]}_{\mathrm{Agreement}}\ \Big),$$

all entries computed on the same realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$.

Proposition 8.114 (Quadrant–parasymmetry correspondence (sector-conditioned))**.** *Let $\mathrm{Quad}^{[\mathscr{S}]}$ be the quadrant label returned by $\mathcal{R}^{[\mathscr{S}]}$ in Definition 8.102. Fix sector thresholds $\tau_{\mathrm{syn}}, \tau_{\mathrm{agr}} \in (0, \frac{1}{2})$ determined at calibration. Then, on any certified plateau component:*

$$\begin{aligned} \text{QI } \textit{or}\ \text{QIV} &\implies v^{(LS)[\mathscr{S}]}_{\mathrm{Agreement}} \ \geq\ \tfrac{1}{2} + \tau_{\mathrm{agr}}, \\ \text{QII } \textit{or}\ \text{QIII} &\implies v^{(LS)[\mathscr{S}]}_{\mathrm{Synthesis}} \ \geq\ \tfrac{1}{2} + \tau_{\mathrm{syn}}. \end{aligned}$$

Moreover, in the thermodynamic/IR limits associated to the sector's r:

(i) QII (SSB).
$\Lambda^{[\mathscr{S}]}_{\mathrm{state}}(V) \to 0$ implies $v^{(S)[\mathscr{S}]}_{\mathrm{Synthesis}} \to \frac{1}{2}$ (incoherent mixture baseline), while $v^{(LS)[\mathscr{S}]}_{\mathrm{Synthesis}}$ remains elevated (II/III diagonal).

(ii) QIII (emergence).
If $d^{[\mathscr{S}]}_{\mathrm{law}}(r) \leq Cr^{-\beta}$, then $v^{(LS)[\mathscr{S}]}_{\mathrm{Synthesis}}$ is non-decreasing along r up to $o(1)$, and any deficit in $v^{(LS)[\mathscr{S}]}_{\mathrm{Agreement}}$ shrinks $o(1)$.

(iii) QIV (explicit).
A persistent law deficit on $\mathfrak{P}$ is compatible with an elevated $v^{(LS)[\mathscr{S}]}_{\mathrm{Agreement}}$ (I/IV diagonal), while $v^{(S)[\mathscr{S}]}_{\mathrm{Synthesis}}$ need not exceed the mixture baseline.

These statements refine, but do not modify, the quadrant decision of Section 8.1: the parasymmetric meters provide a coherence profile *aligned with the I/IV vs. II/III diagonal structure described in Section 8.2.1.*

Definition 8.115 (Operational overlay (sector-conditioned, parasymmetric))**.** Augment the operational template of Definition 8.105 with:

(i) Para calibration.
Choose $\kappa^{[\mathscr{S}]}$ satisfying **(C1)–(C3)** of Definition 8.111; fix thresholds $(\tau_{\mathrm{syn}}, \tau_{\mathrm{agr}})$.

(ii) Meter readout.
Compute $\upsilon^{(L)[\mathscr{S}]}_{\mathrm{Synthesis}}, \upsilon^{(S)[\mathscr{S}]}_{\mathrm{Synthesis}}, \upsilon^{(LS)[\mathscr{S}]}_{\mathrm{Synthesis}}, \upsilon^{(LS)[\mathscr{S}]}_{\mathrm{Agreement}}$ on $\mathcal{A}$.

(iii) Coherence audit.
Report the *coherence class*

$$\mathsf{Coh}^{[\mathscr{S}]} := \Big(\begin{array}{l} \text{diag preference: I/IV vs. II/III;} \\ \text{local } L/S \text{ “bothness” levels;} \\ \text{conservation via Proposition 8.112} \end{array} \Big),$$

and attach it to $\mathsf{Expl}^{[\mathscr{S}]}$ from Definition 8.102.

The resulting package $\widehat{\mathfrak{S}}^{[\mathscr{S}]}(\mathfrak{P})$ is sector-certified, gauge/Nielsen-flat, and functorial across arena and IR choices, preserving the quadrant label while enriching its explanation with phase-sensitive coherence data.

Remark 8.116 (Interpretation and sector portability)**.** *Agreement* vs. *Synthesis* meters resolve whether the law/state pair sits on the I/IV or II/III diagonals of the generalized 2×2 grid of Section 8.1. Because all constructions are anchored to the same certified plateau, BRST-closed tags, and projection to $\mathcal{H}_{\mathrm{phys}}$, the parasymmetric overlay is robust under gauge retunings (Nielsen-flat), respects causal cone preservation, and transports across bulk, boundary, and anisotropic sectors without altering the canonical quadrant rule of Section 8.1, in line with Section 8.2.1.

8.5 Applications of Parasymmetric Overlays

Story in one line. The parasymmetric overlay is a *diagnostic instrument* that not only tells whether a fermionic symmetry is broken, but also *why*—by reading the phase–coherent relationship between the *law* and the *state* on a certified bulk plateau $\mathfrak{P}$ within the BRST–safe sector $\mathcal{H}_{\mathrm{phys}}$ (Definition 8.10).

Two suspects, one dashboard. There are two principal culprits behind broken symmetry:

(i) Spontaneous selection (SSB). The law is symmetric but the state *chooses* a direction (QII).

(ii) Built–in imperfection (explicit). The law itself carries a symmetry–breaking term (QIV).

The sector visibilities $\left(\Lambda_{\text{law}}^{[\text{ferm}]}, \Lambda_{\text{state}}^{[\text{ferm}]}\right)$ place the system in QI–QIV (Proposition 8.11), but the overlay adds *meters* that reveal the *coherent geometry* of the law–state relation, not just the magnitudes.

The meters (phase–aware dials). A calibration $\kappa^{[\text{ferm}]}$ maps witness/ledger data to densities $\rho^{(L)[\text{ferm}]}, \rho^{(S)[\text{ferm}]}, \rho^{(LS)[\text{ferm}]}$. From these, two relational and one local meter are read with the projectors of Section 8.2.1:

$$v_{\text{Agreement}}^{(LS)[\text{ferm}]} \text{ (coherent I/IV)},$$
$$v_{\text{Synthesis}}^{(LS)[\text{ferm}]} \text{ (coherent II/III)},$$
$$v_{\text{Synthesis}}^{(S)[\text{ferm}]} \text{ (local state “bothness”)}.$$

Normalization is universal: 1 (coherent both/and), 1/2 (incoherent mixture baseline), 0 (coherent anti–both). All readings are *Nielsen–flat* on $\mathfrak{P}$ and thus gauge–parameter insensitive.

What the dials say (cause vs effect).

(i) SSB (QII). The *tension* dial is high and the *local* state dial collapses to the mixture baseline:

$$|\Lambda_{\text{law}}^{[\text{ferm}]}| \approx 1, \quad |\Lambda_{\text{state}}^{[\text{ferm}]}| \approx 0, \quad v_{\text{Synthesis}}^{(LS)[\text{ferm}]} > \tfrac{1}{2}, \quad v_{\text{Synthesis}}^{(S)[\text{ferm}]} \to \tfrac{1}{2}.$$

(ii) Explicit (QIV). The *harmony* dial is high (coherent I/IV), without requiring large local “bothness”:

$$|\Lambda_{\text{state}}^{[\text{ferm}]}| \approx 1, \quad |\Lambda_{\text{law}}^{[\text{ferm}]}| \leq 1 - \delta_\star, \quad v_{\text{Agreement}}^{(LS)[\text{ferm}]} > \tfrac{1}{2}.$$

(iii) Emergent (QIII). Finite–scale explicit effects *heal* under the IR knob $r = \ell$; the tension meter *rises* with healing:

$$d_{\text{law}}^{[\text{ferm}]}(r) \downarrow, \quad v_{\text{Synthesis}}^{(LS)[\text{ferm}]}(r) \nearrow \text{ up to } o(1).$$

(iv) Symmetric (QI). Agreement is elevated (coherent I/IV); both visibilities are near 1.

Why the readings are physical (spectral/Ward fingerprints). The meters are wired to *observable* ledgers (Definition 8.10):

QII. Elevated tension $\Leftrightarrow$ NG partners in the fermionic channel (spontaneous breaking).

QIV. Elevated harmony $\Leftrightarrow$ pseudo–NG (GMOR–type) band on $\mathfrak{P}$ and analytic mass/mixing shifts in S^R (explicit breaking).

QIII. No NG/pNG; adjoint divergence $\to 0$ along r; tension increases as the law heals (emergence).

Why this matters (high-level). *Ambiguity ends:* identical magnitudes $\big(|\Lambda_{\text{law}}|, |\Lambda_{\text{state}}|\big)$ can arise from different mechanisms; the meters separate *coherent opposition* (QII/QIII) from *coherent agreement* (QI/QIV). *Actionable searches:* elevated tension points to NG channels; elevated harmony points to pNG bands and mass shifts. *Quality assurance:* when the relevant projectors commute with the dynamics (or satisfy $\mathcal{L}^\dagger(P) = 0$), the meters are conserved/slow on $\mathfrak{P}$; drift flags boundary leakage or mis–instrumentation rather than new physics.

The parasymmetric overlay promotes the familiar QI–QIV table into a *phase–structured dashboard*. In fermionic language, it reads whether the law and the state are in *harmony* (I/IV) or in *tension* (II/III), and it ties those readings to concrete spectral/Ward signatures. That is why it can say not only *that* symmetry is broken, but *how*, and *by whom*—the dynamics (explicit), the state (spontaneous), or neither (symmetric), with emergence certified by IR healing trends (Proposition 8.11 and Section 8.2.1).

8.5.1 Sector-Agnostic Parasymmetric Overlay

Scope and layer (sector-agnostic). Fix an arbitrary sector $[\mathscr{S}]$ with realized weave configuration $\mathcal{I}^{[\mathscr{S}]} = (\mathcal{A}, \mathfrak{P}; r; P_x(\lambda_{\text{UV}}), K_{\text{IR}}; \mathsf{P}_{\text{phys}})$ as in Definition 8.96, and work on the physical cohomology $\mathcal{H}_{\text{phys}}$ with BRST-closed tags. The generalized witness $\mathbf{W}^{[\mathscr{S}]}_{\text{gen}}$ (Section 8.3) returns the sector visibilities $\big(\Lambda^{[\mathscr{S}]}_{\text{law}}, \Lambda^{[\mathscr{S}]}_{\text{state}}\big) \in [0,1]^2$ together with Ward/spectral ledgers, all evaluated on the same $\mathcal{I}^{[\mathscr{S}]}$. Parasymmetry overlays this resolver-ready data with phase-sensitive meters that *corroborate and explain* the quadrant assignment without altering it.

Definition 8.117 (Parasymmetric interface (sector-agnostic calibration and meters))**.** A *parasymmetric calibration* for sector $[\mathscr{S}]$ is a completely positive, trace-preserving map

$$\kappa^{[\mathscr{S}]} : \Big(\Lambda^{[\mathscr{S}]}_{\text{law}}, \Lambda^{[\mathscr{S}]}_{\text{state}};\ \mathcal{W}^{[\mathscr{S}]}, \mathcal{Z}^{[\mathscr{S}]}\Big) \longrightarrow \Big(\rho^{(L)[\mathscr{S}]},\ \rho^{(S)[\mathscr{S}]},\ \rho^{(LS)[\mathscr{S}]}\Big),$$

where $\rho^{(L)[\mathscr{S}]}, \rho^{(S)[\mathscr{S}]} \in \mathsf{D}(\mathbb{C}^2)$ encode local "Sym/Break'' content for *law* and *state*, and $\rho^{(LS)[\mathscr{S}]} \in \mathsf{D}(\mathbb{C}^2 \otimes \mathbb{C}^2)$ encodes relational law–state content. With local parasymmetric projectors $P^{(a)}_{\text{para}} = |+\rangle_a\langle+|$ ($a \in \{L, S\}$) and Bell-type projectors $P^{(LS)}_{\text{para}}$ (coherent *opposition*, II/III diagonal) and $P^{(LS)}_{\Phi,\text{para}}$ (coherent *agreement*, I/IV diagonal), define the meters

$$\begin{aligned} v^{(L)[\mathscr{S}]}_{\text{Synthesis}} &= \text{Tr}\Big(P^{(L)}_{\text{para}}\, \rho^{(L)[\mathscr{S}]}\Big), & v^{(S)[\mathscr{S}]}_{\text{Synthesis}} &= \text{Tr}\Big(P^{(S)}_{\text{para}}\, \rho^{(S)[\mathscr{S}]}\Big), \\ v^{(LS)[\mathscr{S}]}_{\text{Synthesis}} &= \text{Tr}\Big(P^{(LS)}_{\text{para}}\, \rho^{(LS)[\mathscr{S}]}\Big), & v^{(LS)[\mathscr{S}]}_{\text{Agreement}} &= \text{Tr}\Big(P^{(LS)}_{\Phi,\text{para}}\, \rho^{(LS)[\mathscr{S}]}\Big). \end{aligned}$$

Normalization is fixed so that 1 denotes a coherent "both/and", 1/2 the incoherent mixture baseline, and 0 a coherent "anti-both''. Calibration inherits gauge/Nielsen flatness and plateau uniformity from $\mathcal{I}^{[\mathscr{S}]}$.

Definition 8.118 (Reading rule and thresholds)**.** Fix sector thresholds $\tau_{\text{agr}}, \tau_{\text{syn}} \in (0, \frac{1}{2})$. A parasymmetric meter v is called *elevated* if $v \geq \frac{1}{2} + \tau$ with the appropriate threshold $\tau \in \{\tau_{\text{agr}}, \tau_{\text{syn}}\}$. Operationally:

- $v^{(LS)[\mathscr{S}]}_{\text{Agreement}}$ elevated $\Rightarrow$ coherent I/IV diagonal (law–state agreement).

- $v^{(LS)[\mathscr{S}]}_{\text{Synthesis}}$ elevated $\Rightarrow$ coherent II/III diagonal (law–state opposition).
- $v^{(L)[\mathscr{S}]}_{\text{Synthesis}}$ or $v^{(S)[\mathscr{S}]}_{\text{Synthesis}} = 1/2 \Rightarrow$ local incoherent mixture; values $\uparrow$ toward 1 indicate local coherent "bothness".

Proposition 8.119 (Quadrant expectations with parasymmetric meters (sector-agnostic))**.** *Let* $(\mathrm{Quad}^{[\mathscr{S}]}, \mathsf{Expl}^{[\mathscr{S}]}) = \mathcal{R}^{[\mathscr{S}]}\big(\mathfrak{S}^{[\mathscr{S}]}(\mathfrak{P})\big)$ *be the sector label/explanatory class from Definition 8.102, determined by the canonical resolver using the tolerances and healing predicates of Section 8.1. Then, on any certified plateau component:*

(i) SSB (QII).
$|\Lambda^{[\mathscr{S}]}_{\text{law}}| \approx 1$ *and* $\Lambda^{[\mathscr{S}]}_{\text{state}}(V) \to 0$. *Consequently* $v^{(S)[\mathscr{S}]}_{\text{Synthesis}} \to \frac{1}{2}$ *(local mixture baseline), while* $v^{(LS)[\mathscr{S}]}_{\text{Synthesis}} \geq \frac{1}{2} + \tau_{\text{syn}}$ *(coherent II/III diagonal);* $v^{(LS)[\mathscr{S}]}_{\text{Agreement}}$ *is not elevated. Ward: conserved improved current. Spectrum: NG poles in the correlational channel.*

(ii) Explicit (QIV).
Persistent law deficit on $\mathfrak{P}$ *with* $|\Lambda^{[\mathscr{S}]}_{\text{state}}| \approx 1$. *Then* $v^{(LS)[\mathscr{S}]}_{\text{Agreement}} \geq \frac{1}{2} + \tau_{\text{agr}}$ *(coherent I/IV diagonal);* $v^{(LS)[\mathscr{S}]}_{\text{Synthesis}}$ *need not be high; local meters need not exceed* $1/2$. *Ward: non-vanishing adjoint–covariant source. Spectrum: pNG (GMOR) on* $\mathfrak{P}$.

(iii) Emergent (QIII).
Law deficit heals with the IR knob r: $d^{[\mathscr{S}]}_{\text{law}}(r) \downarrow$ *(cf. eq. (8.17)), and* $|\Lambda^{[\mathscr{S}]}_{\text{state}}| \approx 1$. *Along* r, $v^{(LS)[\mathscr{S}]}_{\text{Synthesis}}$ *increases monotonically up to* $o(1)$, *while any deficit in* $v^{(LS)[\mathscr{S}]}_{\text{Agreement}}$ *shrinks* $o(1)$. *Ward: adjoint divergence* $\to 0$ *along* r. *Spectrum: no NG/pNG.*

Symmetric (QI).
Both dials ≈ 1. *Then* $v^{(LS)[\mathscr{S}]}_{\text{Agreement}}$ *is elevated (coherent I/IV);* $v^{(LS)[\mathscr{S}]}_{\text{Synthesis}}$ *near baseline unless special phases are prepared. Ward conserved; no NG/pNG.*

Remark 8.120 (Utility: separating coherence from averaging; conserved QA; stability)**.** Parasymmetry separates *coherent structure* from *incoherent averaging*: two sectors with identical magnitudes $(|\Lambda_{\text{law}}|, |\Lambda_{\text{state}}|)$ can differ in phase structure, which the meters expose via the $1/2$ baseline and elevated thresholds. Under generators that commute with the relevant projectors (or satisfy $\mathcal{L}^\dagger(P) = 0$), the meters are conserved/slow on $\mathfrak{P}$, providing QA dials whose drift flags mis-instrumentation or boundary leakage. Because readings live on $\mathcal{H}_{\text{phys}}$ with BRST-closed tags, they are gauge/Nielsen-flat and portable across bulk, boundary, and anisotropic sectors.

Definition 8.121 (Operational checklist (sector-agnostic, SynTh + parasymmetry))**.** Given $[\mathscr{S}]$ on $\mathcal{I}^{[\mathscr{S}]}$:

(i) Resolver.
Read the resolver dials $\left(\Lambda^{[\mathscr{S}]}_{\text{law}}, \Lambda^{[\mathscr{S}]}_{\text{state}}\right)$ and assign QI–QIV via $\mathcal{R}^{[\mathscr{S}]}$.

(ii) Calibration.
Calibrate $\kappa^{[\mathscr{S}]}$ and compute $v^{(L)[\mathscr{S}]}_{\text{Synthesis}}, v^{(S)[\mathscr{S}]}_{\text{Synthesis}}, v^{(LS)[\mathscr{S}]}_{\text{Synthesis}}, v^{(LS)[\mathscr{S}]}_{\text{Agreement}}$.

(iii) Thresholds.
Use thresholds $\tau_{\text{agr}}, \tau_{\text{syn}} > 0$ to call elevated meters and interpret the diagonal (I/IV vs II/III) and local "bothness".

(iv) Ward/spectral check.
Cross-check Ward/spectral ledgers: (QII) conserved Ward with NG; (QIV) adjoint source with pNG/GMOR; (QIII) vanishing divergence and no NG/pNG.

(v) Steering.
For steering: monitor $v^{(LS)[\mathscr{S}]}_{\text{Synthesis}}(r)$ for QIII (rising with r) and $v^{(LS)[\mathscr{S}]}_{\text{Agreement}}$ for QIV (persistent elevation).

Bottom line (sector-agnostic). The quadrant resolver remains the primary classifier of SSB (QII) versus explicit breaking (QIV) versus emergent/symmetric cases (QIII/QI). Parasymmetry supplies *coherence-sensitive, conserved, and gauge-robust meters* that validate and explain that classification by revealing *how* the law and state occupy the generalized 2×2 grid—whether in coherent agreement (I/IV), coherent opposition (II/III), or as finite-scale mixtures converging under the sector's IR knob r.

8.5.2 Parasymmetric Overlay of the Fermionic (Spin-1/2) Sector

This section presents a parasymmetric overlay for diagnosing how a fermionic symmetry is realised or broken on the certified bulk plateau $\mathfrak{P}$, within the BRST–safe physical sector $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$ (Definition 8.10). The overlay refines the canonical quadrant resolver of Proposition 8.11 by adding *phase-sensitive* meters that report coherence between the *law* and the *state*, rather than only magnitudes of sector visibilities.

Quadrants and sector visibilities. The sector-conditioned dials $\left(\Lambda^{[\text{ferm}]}_{\text{law}}, \Lambda^{[\text{ferm}]}_{\text{state}}\right) \in [0,1]^2$ place the system in one of four regimes: QI (both symmetric), QII (spontaneous breaking: law symmetric, state broken), QIII (finite-scale explicit deformations that *heal* under the IR knob $r = \ell$), and QIV (persistent explicit breaking on the plateau). Tolerances $\varepsilon_{\text{law}}, \varepsilon_{\text{state}}$ implement the certified "≈ 1" predicates (Definition 8.10).

Parasymmetric enhancement. A calibration $\kappa^{[\text{ferm}]}$ maps the witness/ledger data to density operators

$$\rho^{(L)[\text{ferm}]}, \quad \rho^{(S)[\text{ferm}]}, \quad \rho^{(LS)[\text{ferm}]},$$

encoding, respectively, the law anchor, the state anchor, and their relation. From these, two relational meters and one local meter are read using the projectors of Section 8.2.1:

$$v^{(LS)[\text{ferm}]}_{\text{Agreement}}, \qquad v^{(LS)[\text{ferm}]}_{\text{Synthesis}}, \qquad v^{(S)[\text{ferm}]}_{\text{Synthesis}},$$

with the normalization 1 (coherent both/and), 1/2 (incoherent mixture baseline), 0 (coherent anti-both). The meters are Nielsen-flat and BRST-safe on $\mathfrak{P}$.

What the overlay achieves. *(i) Cause vs effect.* Spontaneous breaking (QII) exhibits an *elevated opposition* meter $v^{(LS)[\text{ferm}]}_{\text{Synthesis}} > 1/2$ and a *local mixture* $v^{(S)[\text{ferm}]}_{\text{Synthesis}} \to 1/2$, whereas persistent explicit breaking (QIV) yields an *elevated agreement* meter $v^{(LS)[\text{ferm}]}_{\text{Agreement}} > 1/2$ without requiring large local "bothness". In the fully symmetric case (QI) the agreement meter is

likewise elevated. *(ii) Spectral alignment.* Elevated opposition in QII aligns with NG partners in the fermionic channel; elevated agreement in QIV aligns with a pNG (GMOR-type) band and analytic mass/mixing shifts; QIII shows no NG/pNG and a vanishing adjoint divergence along r. *(iii) IR trends.* In QIII the opposition meter increases monotonically as $\ell \downarrow 0$, mirroring the healing predicate $d^{[\mathrm{ferm}]}_{\mathrm{law}}(\ell)\downarrow$; in QIV the agreement meter remains persistently high on $\mathfrak{P}$.

Audit and wiring. Because the meters are conserved/slow when the relevant projectors commute with the dynamics (or satisfy $\mathcal{L}^\dagger(P)=0$), they serve as QA dials: unexpected drift flags boundary leakage or mis-tagging. A plateau-normalised cross-coherence, e.g.

$$\rho_{12}^{(LS)[\mathrm{ferm}]} \;\propto\; \frac{\big\langle \mathrm{Src}^{(\mathrm{ferm})},\, \mathcal{O}_{\mathrm{ferm}} \big\rangle_{\mathfrak{P}}}{\sqrt{\langle \mathrm{Src}^{(\mathrm{ferm})}, \mathrm{Src}^{(\mathrm{ferm})}\rangle_{\mathfrak{P}}\; \langle \mathcal{O}_{\mathrm{ferm}}, \mathcal{O}_{\mathrm{ferm}}\rangle_{\mathfrak{P}}}},$$

implements the meters directly from the Ward source and a tagged fermionic order operator, turning spectral/ledger data into phase-aware diagnostics.

Setup (fermionic sector; same instrumentation). Work on the certified bulk plateau $\mathfrak{P}$ and in the physical sector $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$, using the BRST-closed tags $P_x(\lambda_{\mathrm{UV}})$ and $K^{(\ell)}_{\mathrm{IR}}$ fixed in Definition 8.10. The generalized witness $\mathbf{W}^{[\mathrm{ferm}]}_{\mathrm{gen}}$ (Section 8.3) delivers the fermionic sector visibilities

$$\big(\Lambda^{[\mathrm{ferm}]}_{\mathrm{law}},\; \Lambda^{[\mathrm{ferm}]}_{\mathrm{state}}\big) \in [0,1]^2,$$

together with the Ward/spectral ledgers on the same instrumentation: retarded propagator S^R, tagged wave operator $\mathcal{D}^{(\ell)}$, tagged mass/mixing $M^{(\ell)}$, and scale-tagged sources $\mathcal{S}_{k,a}$. All evaluations are performed on $\mathfrak{P}$, hence Nielsen-flat and BRST-safe.

Definition 8.122 (Fermionic parasymmetric calibration and densities)**.** A *fermionic parasymmetric calibration* is a completely positive, trace-preserving map

$$\kappa^{[\mathrm{ferm}]}\colon\; \Big(\Lambda^{[\mathrm{ferm}]}_{\mathrm{law}}, \Lambda^{[\mathrm{ferm}]}_{\mathrm{state}};\; \mathcal{W}^{[\mathrm{ferm}]}, \mathcal{Z}^{[\mathrm{ferm}]}\Big) \;\longrightarrow\; \Big(\rho^{(L)[\mathrm{ferm}]},\; \rho^{(S)[\mathrm{ferm}]},\; \rho^{(LS)[\mathrm{ferm}]}\Big),$$

where $\rho^{(L)[\mathrm{ferm}]}, \rho^{(S)[\mathrm{ferm}]} \in \mathsf{D}(\mathbb{C}^2)$ are *local* "Sym/Break" density operators for the law and state anchors, and $\rho^{(LS)[\mathrm{ferm}]} \in \mathsf{D}(\mathbb{C}^2 \otimes \mathbb{C}^2)$ is the *relational* law–state density. Calibration inherits: (i) gauge/Nielsen flatness (from BRST closure and projection to $\mathcal{H}_{\mathrm{phys}}$), (ii) sector functoriality under admissible $(\mathcal{A}, r)$ moves, and (iii) plateau monotonicity along the fermionic IR knob $r = \ell$, in the sense of Section 8.2.1.

Definition 8.123 (Fermionic parasymmetric meters)**.** With projectors $P^{(a)}_{\mathrm{para}} = |+\rangle_a\langle +|$ on the local "Sym/Break" bases ($a \in \{L, S\}$) and Bell-type projectors $P^{(LS)}_{\mathrm{para}}$ (opposition, II/III diagonal) and $P^{(LS)}_{\Phi,\mathrm{para}}$ (agreement, I/IV diagonal) on the relational space, define:

$$\begin{aligned}
&\text{Local meters:} && v^{(L)[\mathrm{ferm}]}_{\mathrm{Synthesis}} = \mathrm{Tr}\Big(P^{(L)}_{\mathrm{para}}\, \rho^{(L)[\mathrm{ferm}]}\Big), \quad v^{(S)[\mathrm{ferm}]}_{\mathrm{Synthesis}} = \mathrm{Tr}\Big(P^{(S)}_{\mathrm{para}}\, \rho^{(S)[\mathrm{ferm}]}\Big);\\
&\text{Relational meters:} && v^{(LS)[\mathrm{ferm}]}_{\mathrm{Synthesis}} = \mathrm{Tr}\Big(P^{(LS)}_{\mathrm{para}}\, \rho^{(LS)[\mathrm{ferm}]}\Big), \quad v^{(LS)[\mathrm{ferm}]}_{\mathrm{Agreement}} = \mathrm{Tr}\Big(P^{(LS)}_{\Phi,\mathrm{para}}\, \rho^{(LS)[\mathrm{ferm}]}\Big).
\end{aligned}$$

Normalization: 1 indicates a coherent "both/and" configuration, 1/2 the incoherent mixture baseline, and 0 a coherent "anti-both" configuration. These meters are evaluated on $\mathfrak{P}$.

Lemma 8.124 (Cross-coherence calibration for the relational off-diagonal)**.** *Let $\langle\cdot,\cdot\rangle_{\mathfrak{P}}$ denote the plateau-uniform, BRST-safe Källén–Lehmann seminorm used in Definition 8.10, and let $\mathcal{O}_{\mathrm{ferm}}$ be a tagged fermionic order operator for the symmetry under test (e.g. chiral $\bar\psi\psi$ or $\bar\psi i\gamma_5\tau\psi$). A canonical choice within $\kappa^{[\mathrm{ferm}]}$ is*

$$\rho_{12}^{(LS)[\mathrm{ferm}]} \;\propto\; \frac{\left\langle \mathrm{Src}^{(\mathrm{ferm})},\, \mathcal{O}_{\mathrm{ferm}} \right\rangle_{\mathfrak{P}}}{\sqrt{\langle \mathrm{Src}^{(\mathrm{ferm})}, \mathrm{Src}^{(\mathrm{ferm})}\rangle_{\mathfrak{P}}\ \langle \mathcal{O}_{\mathrm{ferm}}, \mathcal{O}_{\mathrm{ferm}}\rangle_{\mathfrak{P}}}}\,.$$

With this choice,

$$v_{\mathrm{Synthesis}}^{(LS)[\mathrm{ferm}]} = \tfrac{1}{2} + \mathrm{Re}\ \rho_{12}^{(LS)[\mathrm{ferm}]}, \qquad v_{\mathrm{Agreement}}^{(LS)[\mathrm{ferm}]} = \tfrac{1}{2} + \mathrm{Re}\ \rho_{\Phi,12}^{(LS)[\mathrm{ferm}]}.$$

Quadrant-resolved narrative (parasymmetric readings). The parasymmetric overlay does not alter the quadrant assignment of Proposition 8.11; rather, it refines the interpretation by reporting *coherence profiles* aligned with the I/IV vs. II/III diagonals.

QI (law kept, state symmetric). Assume $\Lambda_{\mathrm{law}}^{[\mathrm{ferm}]} = 1$ and $\Lambda_{\mathrm{state}}^{[\mathrm{ferm}]} = 1$ (Proposition 8.11). Then the relational *agreement* meter is elevated, $v_{\mathrm{Agreement}}^{(LS)[\mathrm{ferm}]} \gtrsim \frac{1}{2} + \tau_{\mathrm{agr}}$, signalling coherent population of the I/IV diagonal, while $v_{\mathrm{Synthesis}}^{(LS)[\mathrm{ferm}]}$ remains near baseline unless special phases are prepared. Locally, a coherently prepared symmetric state yields $v_{\mathrm{Synthesis}}^{(S)[\mathrm{ferm}]} \approx 1$, with dephasing pushing the value toward $\frac{1}{2}$. The Ward/spectral ledger shows a conserved improved current and no NG/pNG contributions in the fermionic channel; residues $Z \succeq 0$ are consistent with a symmetry-respecting band in S^R. The elevated agreement meter provides a steady audit dial (often conserved under suitable generators in Section 8.2.1); drifts suggest boundary leakage or mis-instrumentation.

QII (law kept; fermionic SSB). Here $\Lambda_{\mathrm{law}}^{[\mathrm{ferm}]} = 1$ while $\Lambda_{\mathrm{state}}^{[\mathrm{ferm}]} = 0$. In the thermodynamic limit, the local state meter relaxes to the mixture baseline, $v_{\mathrm{Synthesis}}^{(S)[\mathrm{ferm}]} \to \frac{1}{2}$, indicating loss of coherent "bothness'' within the state. Concurrently, the relational *opposition* meter is elevated, $v_{\mathrm{Synthesis}}^{(LS)[\mathrm{ferm}]} \gtrsim \frac{1}{2} + \tau_{\mathrm{syn}}$, signalling coherent population of the II/III diagonal, while the agreement meter stays near baseline. The Ward/spectral ledger retains a conserved improved current and records NG partners in the correlational channel; if the relevant generator is gauged, NG strength migrates to BEH masses.

QIII (emergent fermionic symmetry; finite-scale explicit breaking that heals). Assume $\Lambda_{\mathrm{law}}^{[\mathrm{ferm}]}(\ell) \to 1$ as $\ell \downarrow 0$ and $\Lambda_{\mathrm{state}}^{[\mathrm{ferm}]} = 1$. Along the sector IR knob $r{=}\ell$, the healing predicate $d_{\mathrm{law}}^{[\mathrm{ferm}]}(\ell) = 1 - |\Lambda_{\mathrm{law}}^{[\mathrm{ferm}]}(\ell)| \le C\,\ell^{-\beta}$ is mirrored parasymmetrically by a monotone trend: $v_{\mathrm{Synthesis}}^{(LS)[\mathrm{ferm}]}$ *increases* (or stays above baseline) as $\ell \downarrow 0$, and any deficit in $v_{\mathrm{Agreement}}^{(LS)[\mathrm{ferm}]}$ shrinks $o(1)$. No NG/pNG modes appear; the adjoint divergence tends to zero along r. Parasymmetry thus furnishes a rate-of-healing meter, distinguishing true emergence from near-persistent defects.

QIV (explicit fermionic breaking; persistent). If $\Lambda_{\mathrm{law}}^{[\mathrm{ferm}]} \le 1 - \delta_\star$ uniformly while $\Lambda_{\mathrm{state}}^{[\mathrm{ferm}]} = 1$, the relational *agreement* meter is elevated, $v_{\mathrm{Agreement}}^{(LS)[\mathrm{ferm}]} \gtrsim \frac{1}{2} + \tau_{\mathrm{agr}}$, confirming coherent I/IV population; $v_{\mathrm{Synthesis}}^{(LS)[\mathrm{ferm}]}$ need not be large, and local "bothness'' need not exceed $\frac{1}{2}$. The Ward/spectral ledger exhibits a persistent adjoint–covariant source on $\mathfrak{P}$ and a fermionic pNG (GMOR-type) band; analytic mass/mixing shifts appear in $M^{(\ell)}$ and are visible in S^R. This combination certifies explicitness as a property of the dynamics rather than of the state.

Remark 8.125 (Concrete observables and wiring to S^R)**.** A plateau-normalized cross-coherence $\langle \mathrm{Src}^{(\mathrm{ferm})}, \mathcal{O}_{\mathrm{ferm}} \rangle_{\mathfrak{P}}$, with the same seminorm as in Definition 8.10, supplies the off-diagonal of $\rho^{(LS)[\mathrm{ferm}]}$ (Lemma 8.124), yielding

$$v^{(LS)[\mathrm{ferm}]}_{\mathrm{Synthesis}} = \tfrac{1}{2} + \mathrm{Re}\ \rho^{(LS)}_{12}, \qquad v^{(LS)[\mathrm{ferm}]}_{\mathrm{Agreement}} = \tfrac{1}{2} + \mathrm{Re}\ \rho^{(LS)}_{\Phi,12}.$$

The local state meter is obtained from the reduced, tagged fermionic block (Sym/Break basis built from the order parameter sector), $v^{(S)[\mathrm{ferm}]}_{\mathrm{Synthesis}} = \frac{1}{2} + \mathrm{Re}\ \rho^{(S)}_{12}$, hence in SSB, $\rho^{(S)}_{12} \to 0$ and $v^{(S)}_{\mathrm{Synthesis}} \to \frac{1}{2}$. A law-aligned local meter $v^{(L)[\mathrm{ferm}]}_{\mathrm{Synthesis}}$ is analogously defined using the improved current sector.

Corollary 8.126 (Executive summary for the fermionic sector)**.** *On the certified plateau $\mathfrak{P}$ the parasymmetric overlay yields:*

(i) Symmetric (QI).
Agreement coherence high; no NG/pNG; provides a robust audit dial.

(ii) SSB (QII).
$\Lambda^{[\mathrm{ferm}]}_{\mathrm{law}} = 1$, $\Lambda^{[\mathrm{ferm}]}_{\mathrm{state}} = 0$*;* $v^{(S)}_{\mathrm{Synthesis}} \to \frac{1}{2}$ *and* $v^{(LS)}_{\mathrm{Synthesis}}$ *elevated (coherent II/III); NG partners present.*

(iii) Emergent (QIII).
Law heals with ℓ; $v^{(LS)}_{\mathrm{Synthesis}}$ *grows monotonically along r; Ward divergence $\to 0$; no NG/pNG.*

(iv) Explicit (QIV).
Persistent law deficit; $v^{(LS)}_{\mathrm{Agreement}}$ *elevated (coherent I/IV); pNG/GMOR on $\mathfrak{P}$; analytic shifts in $S^R/M^{(\ell)}$.*

These meters are phase-sensitive, gauge-robust (Nielsen-flat), and provide conserved/slow invariants under the conditions of Section 8.2.1, thereby refining the quadrant interpretation of Proposition 8.11 without altering the canonical sector-conditioned decision rules.

The parasymmetric overlay augments the sector-conditioned fermionic diagnostics of Definition 8.10 with phase-sensitive meters, while preserving the canonical quadrant resolver of Proposition 8.11. All readings are taken on the certified bulk plateau $\mathfrak{P}$, within the physical cohomology $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$, using the same BRST-closed tags $P_x(\lambda_{\mathrm{UV}})$ and $K^{(\ell)}_{\mathrm{IR}}$. The generalized witness $\mathbf{W}^{[\mathrm{ferm}]}_{\mathrm{gen}}$ (Section 8.3) supplies the fermionic visibilities $\left(\Lambda^{[\mathrm{ferm}]}_{\mathrm{law}}, \Lambda^{[\mathrm{ferm}]}_{\mathrm{state}}\right)$ together with Ward and spectral ledgers (retarded propagator S^R, tagged wave operator $\mathcal{D}^{(\ell)}$, tagged mass/mixing $M^{(\ell)}$, sources $\mathcal{S}_{k,a}$). A fermionic calibration $\kappa^{[\mathrm{ferm}]}$ (the sector instance of Section 8.2.1) maps these data to local and relational "Sym/Break" density operators, from which the parasymmetric valuations are read; by construction they are Nielsen-flat and plateau-uniform.

A first payoff is the disentangling of *cause* versus *effect* in fermionic symmetry breaking. When the law is intact while the state selects a broken phase (QII: $\Lambda^{[\mathrm{ferm}]}_{\mathrm{law}} = 1$, $\Lambda^{[\mathrm{ferm}]}_{\mathrm{state}} = 0$), the parasymmetric state meter relaxes to the mixture baseline, $v^{(S)[\mathrm{ferm}]}_{\mathrm{Synthesis}} \to \frac{1}{2}$, reflecting incoherent occupation of broken sectors by the state, whereas the relational "opposition" meter is elevated, $v^{(LS)[\mathrm{ferm}]}_{\mathrm{Synthesis}} \gtrsim \frac{1}{2} + \tau_{\mathrm{syn}}$, indicating coherent population of the II/III diagonal

(law–state tension). In contrast, under explicit breaking (QIV: a plateau-stable law deficit with $\Lambda_{\text{state}}^{[\text{ferm}]} = 1$), the "agreement'' meter is elevated, $v_{\text{Agreement}}^{(LS)[\text{ferm}]} \gtrsim \frac{1}{2} + \tau_{\text{agr}}$, diagnosing coherent occupation of the I/IV diagonal without requiring enhanced local "bothness". Thus, parasymmetry discriminates *coherent opposition* (SSB) from *coherent agreement* (explicit breaking) even when finite-scale visibilities alone could be ambiguous.

The overlay aligns the quadrant labels with *mechanism-specific spectral fingerprints* in the fermionic ledgers. In QII, the improved current remains conserved while S^R exhibits Nambu–Goldstone partners in the correlational channel; upon gauging, this strength migrates to BEH masses, preserving the diagnosis. In QIV, the Ward source persists on $\mathfrak{P}$ and the spectrum presents a pseudo–Nambu–Goldstone (GMOR-type) band together with analytic shifts in $M^{(\ell)}$, directly visible in the pole structure and residues of S^R. In the emergent case QIII (finite-scale explicit breaking that heals), no NG/pNG modes appear and the adjoint divergence vanishes along the fermionic IR knob $r = \ell$, matching a coherent rise of the relational synthesis meter. Parasymmetry does not replace spectral analysis; rather, it renders the QI–QIV labels *explainable* in fermionic terms by tying coherence patterns to Ward/spectral content.

A further advantage is the ability to *track IR healing versus persistence* with a phase-aware dial. Along $r = \ell$, the healing predicate $d_{\text{law}}^{[\text{ferm}]}(\ell) \downarrow$ that defines QIII is mirrored by a monotone increase of $v_{\text{Synthesis}}^{(LS)[\text{ferm}]}$, signalling coherent reconvergence of law and state as the IR window widens; any deficit in $v_{\text{Agreement}}^{(LS)[\text{ferm}]}$ shrinks to $o(1)$. Conversely, in QIV the agreement meter remains persistently high on $\mathfrak{P}$, commensurate with a plateau-stable law deficit. This differential behavior provides a direct, quantitative separation between "nearly explicit but heals'' (QIII) and "truly explicit'' (QIV).

The parasymmetric meters also supply *quality-control invariants* that are robust under admissible retunings. When the relevant projectors commute with the unified parasymmetric generator (or satisfy $\mathcal{L}^\dagger(P) = 0$ in an open-system description), both local and relational meters are conserved or evolve slowly on $\mathfrak{P}$. Any unexpected drift thus functions as an audit signal for boundary leakage, numerical artefacts, or mis-tagged witnesses. Because all readings are taken on $\mathcal{H}_{\text{phys}}$ with BRST-closed tags, they inherit gauge-fermion independence and respect cone preservation, ensuring portability across bulk, boundary, and anisotropic implementations (Section 8.1).

Operationally, the construction furnishes a *practical wiring diagram* for fermions. A plateau-normalized law–state cross-coherence,

$$\rho_{12}^{(LS)[\text{ferm}]} \propto \frac{\left\langle \text{Src}^{(\text{ferm})},\ \mathcal{O}_{\text{ferm}} \right\rangle_{\mathfrak{P}}}{\sqrt{\langle \text{Src}^{(\text{ferm})}, \text{Src}^{(\text{ferm})} \rangle_{\mathfrak{P}}\ \langle \mathcal{O}_{\text{ferm}}, \mathcal{O}_{\text{ferm}} \rangle_{\mathfrak{P}}}},$$

with $\mathcal{O}_{\text{ferm}}$ a tagged order operator (e.g. chiral), populates the relational off-diagonal used by

$$v_{\text{Synthesis}}^{(LS)[\text{ferm}]} = \tfrac{1}{2} + \text{Re}\ \rho_{12}^{(LS)[\text{ferm}]}, \qquad v_{\text{Agreement}}^{(LS)[\text{ferm}]} = \tfrac{1}{2} + \text{Re}\ \rho_{\Phi,12}^{(LS)[\text{ferm}]}.$$

The local state meter follows from the reduced, tagged fermionic block, $v_{\text{Synthesis}}^{(S)[\text{ferm}]} = \frac{1}{2} + \text{Re}\ \rho_{12}^{(S)}$, approaching $\frac{1}{2}$ in SSB; a law-aligned local meter is defined analogously from the improved current sector. This recipe is reproducible, tag-compatible, and consistent with the generalized witness interface of Section 8.3.

In summary, the parasymmetric overlay turns the fermionic visibilities and spectral data into *reliable, phase-aware diagnostics*. It cleanly separates state-driven (QII) from dynamics-driven (QIV) breaking, quantifies coherence at finite scale via the baseline $v = \frac{1}{2}$, reveals IR emergence (QIII) through monotone coherence growth, and provides conserved or slow meters suitable for ongoing audits on $\mathfrak{P}$. All of these enhancements refine, but do not alter, the quadrant rules and sector-conditioned decision procedures of Proposition 8.11 and Section 8.1.

8.5.3 Parasymmetric Overlay of Supersymmetry (SUSY)

Idea in one sentence. Think of supersymmetry (SUSY) diagnostics as a mechanic's dashboard for the cosmos. The parasymmetric overlay adds two phase–aware dials that show whether the *law* and the *state* are moving in *harmony* or in *tension*, and thereby reveal *how* SUSY is broken.

Setting (where the dashboard is read). Readouts are taken on a certified plateau $\mathfrak{P}$ (or boundary $\mathfrak{P}_\partial$) with the IR knob r (bulk: $r{=}\ell$, boundary: $r{=}L$), in the BRST–safe physical sector $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$; see Section 8.1 and Definition 8.12. This makes the numbers *Nielsen–flat* (insensitive to gauge-fermion retunings) and thus trustworthy.

Two suspects behind broken symmetry. *Spontaneous* SUSY breaking: the law is symmetric, the state *chooses* a direction (like a generalized eigen-operator pair falling). *Explicit* SUSY breaking: the law itself carries an asymmetry (fine print in the rule).

The basic dials (visibilities) and the overlay. SUSY supplies two visibilities $\Lambda_{\text{law}}^{[\text{SUSY}]}$ and $\Lambda_{\text{state}}^{[\text{SUSY}]}$ (both in $[0,1]$). These dials place the situation in the QI–QIV table (Proposition 8.14). The parasymmetric overlay (Section 8.2.1) then adds two *coherence* meters, built from density operators $\rho^{(L/S/LS)[\text{SUSY}]}$ via a calibration $\kappa^{[\text{SUSY}]}$:

$$\underbrace{v_{\text{Agreement}}^{(LS)[\text{SUSY}]}}_{\text{Harmony}} \text{ (coherent I/IV)}, \qquad \underbrace{v_{\text{Synthesis}}^{(LS)[\text{SUSY}]}}_{\text{Tension}} \text{ (coherent II/III)}.$$

Normalization is universal: 1 = coherent "both/and", 1/2 = incoherent mixture baseline, 0 = coherent "anti–both".

How to read the dashboard (high-level map).

Why the dials track real physics. The meters are wired to SUSY's Ward and spectral ledgers (Definition 8.12): *Tension high (QII/QIII)* lines up with NG channels (global) or the super–Higgs pattern (local, massive gravitino), and with vanishing adjoint divergence as emergence sets in (QIII). *Harmony high (QI/QIV)* lines up with conserved/improved currents (QI) or persistent adjoint–covariant sources and pseudo–NG scaling when breaking is explicit (QIV).

Figure 8.1: The parasymmetric diagnostic map, correlating visibilities (Λ) with coherence meter readings and their physical fingerprints across the four quadrants of symmetry realization.

Quadrant	**What visibilities say**	**What the meters confirm**
QI (symmetric)	$\lvert\Lambda_{\text{law}}\rvert \approx 1$, $\lvert\Lambda_{\text{state}}\rvert \approx 1$	*Harmony* is high, indicating coherent agreement. The associated supercurrent is conserved and no Nambu–Goldstone (NG) or pseudo-NG (pNG) modes appear.
QII (spontaneous)	$\lvert\Lambda_{\text{law}}\rvert \approx 1$, $\lvert\Lambda_{\text{state}}\rvert \approx 0$	*Tension* is high, indicating coherent opposition. The local state "bothness" collapses to the mixture baseline ($\to \frac{1}{2}$). The physical signature is a goldstino or the super-Higgs mechanism (Lemma 8.13).
QIII (emergent)	The law deficit heals along the IR knob r.	*Tension* meter *rises* monotonically as the symmetry heals with r. The Ward divergence vanishes, and no NG/pNG modes are present.
QIV (explicit)	The law deficit persists on the certified plateau $\mathfrak{P}$.	*Harmony* is high. A persistent boundary or source term is present in the Ward identity, and a pseudo-goldstino is possible.

IR knob intuition (seeing healing vs persistence). Turning the IR knob r widens the view from short to long distances. If the law deficit *heals* (QIII), the *tension* meter $v^{(LS)}_{\text{Synthesis}}(r)$ climbs monotonically up to $o(1)$, showing the law and state coherently re–align. If the deficit *persists* (QIV), the *harmony* meter stays high on $\mathfrak{P}$—the law and state coherently agree on the breaking.

Operational recipe. (1) Place the point in QI–QIV with $\left(\Lambda^{[\text{SUSY}]}_{\text{law}}, \Lambda^{[\text{SUSY}]}_{\text{state}}\right)$ (Proposition 8.14). (2) Compute $v^{(LS)}_{\text{Agreement}}$, $v^{(LS)}_{\text{Synthesis}}$ (and the local state dial if needed). (3) Match the elevated meter to the expected spectral channel (goldstino vs super–Higgs vs pseudo–goldstino). (4) Vary r to confirm *healing* (QIII: tension $\nearrow$) vs *persistence* (QIV: harmony $\uparrow$ steady); cf. Definition 8.131.

The overlay turns "Is SUSY broken?" into "*How* is it broken—and *who* is responsible?" *Harmony* (I/IV) signals explicit or intact laws; *tension* (II/III) signals spontaneous selection or emergent reconvergence. Because the dials are read on a certified plateau and are Nielsen–flat, they serve as stable, audit–ready guides from abstract symmetry talk to concrete, searchable signatures (goldstino, super–Higgs, pseudo–NG).

Setup (SUSY sector; same instrumentation). Work on a certified plateau $\mathfrak{P}$ and in the physical sector $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$ with BRST-closed tags $P_x(\lambda_{\text{UV}})$, $K^{(\ell)}_{\text{IR}}$, as in Definition 8.12. The SUSY witness delivers the sector visibilities $\left(\Lambda^{[\text{SUSY}]}_{\text{law}}, \Lambda^{[\text{SUSY}]}_{\text{state}}\right) \in [0,1]^2$ together with a SUSY Ward ledger (improved supercurrent S^{μ}_{α}, tagged sources $Y_{j,\alpha}$) and spectral diagnostics (global: goldstino channel; local: gravitino/super-Higgs channel). Parasymmetry adds phase-sensitive meters that corroborate the quadrant assignment of Proposition 8.14 and explain it in terms of coherent *agreement* (I/IV diagonal) or coherent *opposition* (II/III diagonal) between law and state.

Definition 8.127 (SUSY parasymmetric calibration and densities)**.** A *SUSY parasymmetric calibration* is a completely positive, trace-preserving map

$$\kappa^{[\text{SUSY}]}: \left(\Lambda^{[\text{SUSY}]}_{\text{law}}, \Lambda^{[\text{SUSY}]}_{\text{state}};\ \mathcal{W}^{[\text{SUSY}]}, \mathcal{Z}^{[\text{SUSY}]}\right) \longrightarrow \left(\rho^{(L)[\text{SUSY}]},\ \rho^{(S)[\text{SUSY}]},\ \rho^{(LS)[\text{SUSY}]}\right),$$

where $\rho^{(L)[\text{SUSY}]}, \rho^{(S)[\text{SUSY}]} \in \mathsf{D}(\mathbb{C}^2)$ are local "Sym/Break" densities for the *law* and *state* anchors, and $\rho^{(LS)[\text{SUSY}]} \in \mathsf{D}(\mathbb{C}^2 \otimes \mathbb{C}^2)$ encodes the relational law–state sector. Calibration is performed on the same realized weave configuration $(\mathcal{A}, \mathfrak{P}; r; P_x, K_{\text{IR}}; \mathsf{P}_{\text{phys}})$ and inherits Nielsen flatness.

Definition 8.128 (SUSY parasymmetric meters)**.** Let $P^{(a)}_{\text{para}} = |+\rangle_a\langle+|$ $(a \in \{L, S\})$ be local parasymmetric projectors and $P^{(LS)}_{\text{para}}$ (opposition, II/III diagonal), $P^{(LS)}_{\Phi,\text{para}}$ (agreement, I/IV diagonal) be relational Bell-type projectors. Define

$$v^{(L)[\text{SUSY}]}_{\text{Synthesis}} = \text{Tr}\left(P^{(L)}_{\text{para}}\rho^{(L)[\text{SUSY}]}\right), \quad v^{(S)[\text{SUSY}]}_{\text{Synthesis}} = \text{Tr}\left(P^{(S)}_{\text{para}}\rho^{(S)[\text{SUSY}]}\right),$$

$$v^{(LS)[\text{SUSY}]}_{\text{Synthesis}} = \text{Tr}\left(P^{(LS)}_{\text{para}}\rho^{(LS)[\text{SUSY}]}\right), \qquad v^{(LS)[\text{SUSY}]}_{\text{Agreement}} = \text{Tr}\left(P^{(LS)}_{\Phi,\text{para}}\rho^{(LS)[\text{SUSY}]}\right).$$

Normalization: 1 indicates a coherent "both/and", 1/2 the incoherent mixture baseline, and 0 a coherent "anti-both".

Lemma 8.129 (Cross-coherence choices: global vs local SUSY)**.** *Let $\langle\cdot,\cdot\rangle_{\mathfrak{P}}$ be the plateau-uniform, BRST-safe seminorm. A canonical choice for the relational off-diagonal is*

$$\rho_{12}^{(LS)[\mathrm{SUSY}]} \;\propto\; \frac{\Big\langle \sum_j \epsilon_j(\ell) Y_j,\ \mathcal{O}_{\mathrm{SUSY}} \Big\rangle_{\mathfrak{P}}}{\sqrt{\Big\langle \sum_j \epsilon_j Y_j, \sum_k \epsilon_k Y_k \Big\rangle_{\mathfrak{P}}\ \langle \mathcal{O}_{\mathrm{SUSY}}, \mathcal{O}_{\mathrm{SUSY}} \rangle_{\mathfrak{P}}}},$$

with $\mathcal{O}_{\mathrm{SUSY}}$ chosen by sector:

(i) Global SUSY.
$\mathcal{O}_{\mathrm{SUSY}} = \eta_\alpha$ *(goldstino interpolating operator extracted from the supercurrent multiplet).*

(ii) Local SUSY (SUGRA).
$\mathcal{O}_{\mathrm{SUSY}} = \psi_\mu^{\parallel}$ *(longitudinal gravitino/super-Higgs channel on the background admitting a Killing spinor when unbroken).*

Then

$$v_{\mathrm{Synthesis}}^{(LS)[\mathrm{SUSY}]} = \tfrac{1}{2} + \mathrm{Re}\ \rho_{12}^{(LS)[\mathrm{SUSY}]}, \qquad v_{\mathrm{Agreement}}^{(LS)[\mathrm{SUSY}]} = \tfrac{1}{2} + \mathrm{Re}\ \rho_{\Phi,12}^{(LS)[\mathrm{SUSY}]}.$$

Quadrant-resolved narrative (parasymmetric readings for SUSY). The parasymmetric overlay refines the tolerance-form quadrants of Proposition 8.14 by reporting coherence along the I/IV vs. II/III diagonals and by aligning these with goldstino/super-Higgs diagnostics (Lemma 8.13).

QI (law intact, state unbroken). If $|\Lambda_{\mathrm{law}}^{[\mathrm{SUSY}]}| \geq 1-\varepsilon_{\mathrm{law}}$ and $|\Lambda_{\mathrm{state}}^{[\mathrm{SUSY}]}| \geq 1-\varepsilon_{\mathrm{state}}$, the agreement meter is elevated, $v_{\mathrm{Agreement}}^{(LS)[\mathrm{SUSY}]} \gtrsim \frac{1}{2} + \tau_{\mathrm{agr}}$, indicating coherent I/IV occupation. The opposition meter is near baseline unless special phases are prepared. Ward: improved supercurrent conserved; Spectrum: no goldstino/pNG; for local SUSY, a Killing spinor exists and the gravitino is massless as dictated by the background.

QII (spontaneous SUSY breaking). Assume $|\Lambda_{\mathrm{law}}^{[\mathrm{SUSY}]}| \geq 1-\varepsilon_{\mathrm{law}}$ while $|\Lambda_{\mathrm{state}}^{[\mathrm{SUSY}]}| \leq \varepsilon_{\mathrm{state}}$. Then $v_{\mathrm{Synthesis}}^{(S)[\mathrm{SUSY}]} \to \frac{1}{2}$ (the state becomes an incoherent mixture across broken SUSY sectors), and the relational *opposition* meter is elevated, $v_{\mathrm{Synthesis}}^{(LS)[\mathrm{SUSY}]} \gtrsim \frac{1}{2} + \tau_{\mathrm{syn}}$ (coherent II/III diagonal). Global SUSY: a massless goldstino appears (NG of SUSY). Local SUSY: the super-Higgs mechanism eats the goldstino and renders the gravitino massive; no physical massless goldstino remains (Lemma 8.13).

QIII (emergent SUSY). If the explicit SUSY-breaking source heals with the IR knob r (e.g. $1-|\Lambda_{\mathrm{law}}^{[\mathrm{SUSY}]}(\ell)| \leq C\,\ell^{\Delta-(d+z)}$ as $\ell \downarrow 0$) while $|\Lambda_{\mathrm{state}}^{[\mathrm{SUSY}]}| \geq 1-\varepsilon_{\mathrm{state}}$, then $v_{\mathrm{Synthesis}}^{(LS)[\mathrm{SUSY}]}$ increases monotonically along r up to $o(1)$, reflecting coherent reconvergence of law and state; any deficit in $v_{\mathrm{Agreement}}^{(LS)[\mathrm{SUSY}]}$ shrinks. Ward: adjoint divergence $\to 0$ along r; Spectrum: no NG/pNG.

QIV (explicit SUSY breaking; persistent). If there exists $\delta_\star > \varepsilon_{\mathrm{law}}$ with $|\Lambda_{\mathrm{law}}^{[\mathrm{SUSY}]}| \leq 1-\delta_\star$ uniformly on $\mathfrak{P}$ and $|\Lambda_{\mathrm{state}}^{[\mathrm{SUSY}]}| \geq 1-\varepsilon_{\mathrm{state}}$, the agreement meter is elevated, $v_{\mathrm{Agreement}}^{(LS)[\mathrm{SUSY}]} \gtrsim \frac{1}{2} + \tau_{\mathrm{agr}}$ (coherent I/IV diagonal), while $v_{\mathrm{Synthesis}}^{(LS)[\mathrm{SUSY}]}$ need not be large. Ward: adjoint–covariant source persists on $\mathfrak{P}$; Spectrum: no true goldstino; in global models with small explicit breaking, a pseudo-goldstino band may appear (GMOR-type scaling on $\mathfrak{P}$); in local SUSY, the super-Higgs pattern remains controlled by the background and couplings.

Proposition 8.130 (SUSY quadrant–parasymmetry correspondence)**.** *Let* $\tau_{\mathrm{agr}}, \tau_{\mathrm{syn}} \in (0, \frac{1}{2})$ *be fixed sector thresholds. On any certified plateau component:*

$$\begin{aligned} \text{QI } \textit{or } \text{QIV} &\implies v^{(LS)[\mathrm{SUSY}]}_{\mathrm{Agreement}} \geq \tfrac{1}{2} + \tau_{\mathrm{agr}}, \\ \text{QII } \textit{or } \text{QIII} &\implies v^{(LS)[\mathrm{SUSY}]}_{\mathrm{Synthesis}} \geq \tfrac{1}{2} + \tau_{\mathrm{syn}}. \end{aligned}$$

In QII the local state meter satisfies $v^{(S)[\mathrm{SUSY}]}_{\mathrm{Synthesis}} \to \frac{1}{2}$ *as the volume grows, while in QIII the relational synthesis meter increases along* r *in tandem with the healing of* $\Lambda^{[\mathrm{SUSY}]}_{\mathrm{law}} \to 1$*. These coherence statements refine, but do not alter, Proposition 8.14.*

Definition 8.131 (Operational overlay (SUSY, SynTh + parasymmetry))**.** Given [SUSY] on $(\mathcal{A}, \mathfrak{P}; r)$, proceed as follows:

(i) Quadrant assignment.
Assign QI–QIV by $\mathcal{R}^{[\mathrm{SUSY}]}$ using Definition 8.12 and Proposition 8.14.

(ii) Calibration and meters.
Calibrate $\kappa^{[\mathrm{SUSY}]}$ and compute $v^{(L)}_{\mathrm{Synthesis}}$, $v^{(S)}_{\mathrm{Synthesis}}$, $v^{(LS)}_{\mathrm{Synthesis}}$, and $v^{(LS)}_{\mathrm{Agreement}}$ on $\mathfrak{P}$.

(iii) Coherence interpretation.
Interpret elevated meters using Proposition 8.130: I/IV (agreement) versus II/III (opposition); detect local mixture baselines at 1/2.

(iv) Ward/spectral cross-checks.
Cross-check ledgers: (QII global) goldstino; (QII local) super-Higgs with massive gravitino; (QIV) persistent adjoint source and possible pseudo-goldstino scaling; (QIII) vanishing divergence, no NG/pNG.

(v) Steering via meter trends.
Use meter trends for steering: monitor $v^{(LS)}_{\mathrm{Synthesis}}(r)$ for emergent SUSY (QIII) and persistent $v^{(LS)}_{\mathrm{Agreement}}$ for explicitness (QIV).

Remark 8.132 (Consequences and advantages.)**.** Parasymmetry renders SUSY-breaking diagnostics *phase-aware*: it separates coherent opposition (SSB) from coherent agreement (explicit), quantifies finite-scale coherence via the 1/2 baseline, and provides conserved/slow meters under suitable generators on $\mathfrak{P}$. Because all constructions are BRST-safe and Nielsen-flat, the meters are stable under benign retunings and portable across backgrounds (global vs local SUSY), while remaining fully compatible with the sector-conditioned resolver of Proposition 8.14.

The parasymmetric overlay for supersymmetry complements the sector-conditioned witnesses of Definition 8.12 by adding phase-sensitive meters that certify how the "law" and the "state" co-occupy the generalized 2×2 grid on a certified plateau $\mathfrak{P}$, after projection to $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ [52, 82, 91]. The canonical resolver of Proposition 8.14 still determines the quadrant (QI–QIV) from the SUSY visibilities $\left(\Lambda^{[\mathrm{SUSY}]}_{\mathrm{law}}, \Lambda^{[\mathrm{SUSY}]}_{\mathrm{state}}\right)$, but the parasymmetric meters—local "bothness" and relational agreement/opposition as in Section 8.2.1—explain *why* a given point sits where it does by revealing coherent *agreement* (I/IV diagonal) versus coherent *opposition* (II/III diagonal) and by distinguishing coherent structure from mere statistical mixtures (baseline 1/2).

A first conceptual dividend is a clean separation of *cause* and *effect* in SUSY breaking. In the spontaneously broken case (QII), the law is intact while the state fails the tolerance test; correspondingly the local state "bothness'' meter relaxes to the mixture baseline $v^{(S)[\mathrm{SUSY}]}_{\mathrm{Synthesis}} \to \frac{1}{2}$, indicating that the state populates broken sectors incoherently, whereas the relational opposition meter is elevated $v^{(LS)[\mathrm{SUSY}]}_{\mathrm{Synthesis}} \geq \frac{1}{2} + \tau_{\mathrm{syn}}$, signaling coherent occupancy of the II/III diagonal. This coherence pattern matches the mechanism: a goldstino appears in global SUSY, while in local SUSY the super-Higgs effect eats the goldstino and renders the gravitino massive (Lemma 8.13) [92, 94, 91, 52]. In the explicitly broken case (QIV), a plateau-stable law deficit coexists with a symmetric state; the elevated agreement meter $v^{(LS)[\mathrm{SUSY}]}_{\mathrm{Agreement}} \geq \frac{1}{2} + \tau_{\mathrm{agr}}$ then certifies coherent I/IV occupancy without requiring large local "bothness". In global models with small explicit breaking, a pseudo-goldstino (GMOR-type) band may appear on $\mathfrak{P}$ [112, 102]. Thus, identical magnitudes $\big(|\Lambda_{\mathrm{law}}|, |\Lambda_{\mathrm{state}}|\big)$ at finite scale can conceal different phase structures that parasymmetry makes explicit: coherent opposition for QII versus coherent agreement for QIV.

The overlay also aligns quadrant labels with mechanism-specific spectral diagnostics. In QII the improved supercurrent is conserved and the spectrum exhibits either a massless goldstino (global) or, in the local case, a massive gravitino produced by the super-Higgs mechanism; in QIV the SUSY Ward source persists on $\mathfrak{P}$ and the spectrum may show pseudo–NG scaling in the SUSY channel; in QIII no NG/pNG modes occur and the adjoint divergence tends to zero along the sector IR knob, consistent with emergent symmetry [52, 91, 112]. Parasymmetry does not replace spectral analysis; rather, it renders the QI–QIV labels *explainable* in SUSY terms by tying an elevated coherence meter to the appropriate Ward/spectral feature.

A further advantage is a phase-aware view of *healing versus persistence*. In the emergent regime QIII, the healing predicate $1 - |\Lambda^{[\mathrm{SUSY}]}_{\mathrm{law}}(r)| \downarrow$ along the IR knob is mirrored by a monotone increase of the relational synthesis meter $v^{(LS)[\mathrm{SUSY}]}_{\mathrm{Synthesis}} \nearrow$ up to $o(1)$, while any shortfall in the agreement meter shrinks; this documents coherent reconvergence of law and state as the IR window widens, in line with the general RG picture of emergent symmetries [112, 102]. In contrast, QIV exhibits a persistently high agreement meter on $\mathfrak{P}$, matching the plateau-stable source and distinguishing truly explicit breaking from nearly explicit deformations that eventually heal.

The meters themselves serve as robust audit invariants. Under generators commuting with the parasymmetric projectors—or, in open dynamics, whenever $\mathcal{L}^\dagger(P) = 0$—both local and relational meters are conserved or slow on $\mathfrak{P}$. Any unexpected drift therefore signals boundary leakage, mis-instrumentation, or witness mis-tagging. Because all readouts are taken with BRST-closed tags on $\mathcal{H}_{\mathrm{phys}}$, the numbers are Nielsen-flat and transport across global versus local SUSY backgrounds without spurious gauge dependence, preserving the sector portability emphasized in Section 8.2.1 [82, 91, 52].

For practice, fixed sector thresholds $\tau_{\mathrm{agr}}, \tau_{\mathrm{syn}} > 0$ yield a compact rule-of-thumb. Elevated agreement ($v^{(LS)}_{\mathrm{Agreement}} \geq \frac{1}{2} + \tau_{\mathrm{agr}}$) characterizes QI/QIV (I/IV diagonal), whereas elevated opposition ($v^{(LS)}_{\mathrm{Synthesis}} \geq \frac{1}{2} + \tau_{\mathrm{syn}}$) characterizes QII/QIII (II/III diagonal). Additionally, QII displays the finite-volume signature $v^{(S)}_{\mathrm{Synthesis}} \to \frac{1}{2}$ as the state becomes an incoherent mixture, and QIII displays $v^{(LS)}_{\mathrm{Synthesis}}(r)$ increasing with the IR knob as the law heals. These signatures

refine, but do not alter, the tolerance-form quadrant assignment of Proposition 8.14.

In summary, the SUSY parasymmetric overlay cleanly separates spontaneous from explicit breaking through coherence structure (opposition versus agreement), ties these patterns to goldstino/super-Higgs or pseudo–NG spectral content, tracks emergent behavior via monotone coherence growth, and provides gauge-robust, audit-ready meters on the certified plateau [52, 91, 112, 102]. The result is a resolver-compatible, mechanism-explanatory portrait of SUSY phases that remains stable under admissible retunings while offering actionable diagnostics for analysis and control.

8.5.4 Parasymmetric Overlay of Boundary Soft/Edge Modes

We now turn to a natural and powerful extension of the *parasymmetric overlay* to a third, richer setting: boundary soft/edge modes. The central advance is methodological rather than merely topical: the framework now operates in a *multi-sector arena*, diagnosing the symmetry story of the bulk (interior) and the boundary (edge) on their respective certified plateaus, $\mathfrak{P}$ and $\mathfrak{P}_\partial$, and then *reconciling* them within a single, coherent construction [112, 102]. The result is a phase-aware, instrument-grade diagnostic that treats edges not as afterthoughts but as first-class loci of symmetry phenomena.

The dashboard readers introduced earlier carry over intact, but the physics at an edge is distinctive. Boundaries can host special modes—soft waves with long memory, chiral currents that carry protected flow, and other excitations absent in the interior [450, 102]. The question is no longer only whether an edge mode is present; it is how the governing symmetry is realised *at the edge* and how that realisation relates to the bulk. In practical terms, the framework offers a precise way to diagnose and explain the symmetry principles behind edge behaviour, rather than merely recording its existence [112, 102].

The bulk–boundary unification is achieved by analysing the two regions *simultaneously*: the Bulk Plateau $\mathfrak{P}$ with IR knob $r = \ell$, and the Boundary Plateau $\mathfrak{P}_\partial$ with tangential IR knob $r = L$. Treating them together enables questions that previously fragmented the analysis: for instance, what to conclude when a symmetry is perfectly respected in the bulk yet appears broken at the boundary. The joint resolver on the certified intersection $\mathfrak{P} \cap \mathfrak{P}_\partial$ provides a common arena for decision, converting apparent contradictions into a coherent diagnosis that localises mechanisms and prescribes which knob (e.g. L vs. ℓ) should be varied [112, 102].

On the boundary, the familiar toolkit is specialised but not reinvented. A calibration map $\kappa^{[\partial]}$ translates edge data into the same trio of density operators, $\rho^{(L)[\partial]}$, $\rho^{(S)[\partial]}$, and $\rho^{(LS)[\partial]}$, that encode the law anchor, the state anchor, and their relation, in the spirit of the general parasymmetric synthesis of law and state Section 8.2.1. Two coherence meters then read the law–state geometry at the boundary: the *harmony* dial $v^{(LS)[\partial]}_{\text{Agreement}}$ (coherent I/IV diagonal, elevated in QI and QIV) and the *tension* dial $v^{(LS)[\partial]}_{\text{Synthesis}}$ (coherent II/III diagonal, elevated in QII and QIII). The normalisation is universal—1 for a coherent both/and signal, 1/2 for an incoherent mixture baseline, 0 for a coherent anti-signal—and all readings are taken on $\mathfrak{P}_\partial$ with BRST-closed tags and projection to $\mathcal{H}_{\text{phys}}$, ensuring Nielsen-flat, audit-ready numbers [82, 91, 112].

This specialisation yields concrete payoffs for boundary physics. First, it delivers a clear,

phase-aware edge diagnosis: spontaneous breaking at the boundary (QII) appears as elevated tension, with the local state "bothness" relaxing to the mixture baseline $v^{(S)[\partial]}_{\text{Synthesis}} \to \frac{1}{2}$ as edge extent grows, and it is physically accompanied by edge Goldstone modes such as chiral currents [94, 102]. By contrast, explicit breaking at the boundary (QIV) exhibits a persistently high harmony reading and aligns with pseudo–Nambu–Goldstone (GMOR-type) bands in the boundary spectrum [112, 102]. Second, it provides a controllable dial for testing: the size L acts as a direct IR knob. Emergent edge symmetry (QIII) can be demonstrated by increasing L and observing the tension meter $v^{(LS)[\partial]}_{\text{Synthesis}}(L)$ rise monotonically while the boundary Ward source heals, $1 - |\Lambda^{[\partial]}_{\text{law}}(L)| \downarrow\sim L^{-\beta}$, in line with the general emergent-symmetry picture in effective field theory [112]. When breaking is explicit (QIV), the harmony dial remains stably high as L grows, tracking a plateau-persistent source.

Perhaps most importantly, bulk–boundary puzzles become solvable rather than vexing. With the joint resolver, situations like "bulk QI but boundary QII/QIV" are recognised not as inconsistencies but as *localised edge mechanisms*. The coherence profile—harmony versus tension—certifies whether an observed split is a genuine physical effect confined to the boundary or a transient artefact of mismatched arenas, and it guides the necessary reconciliation of knobs until a stable QIII/QIV distinction is reached [112, 102].

In summary, bringing boundary modes into the parasymmetric dashboard advances the framework from single-sector checks to a *multi-sector diagnostic* capable of seeing a whole physical system holistically. It independently diagnoses bulk and boundary symmetry structure with the same, sector-conditioned dials; it connects those diagnoses to observable edge phenomena such as currents, soft modes, and memory effects [450, 102]; and it reconciles apparent contradictions through a unified, phase-aware reading. In doing so, it shows that the parasymmetric overlay scales from the heart of a system to its very edges, turning subtle boundary lore into a coherent, testable story told by harmony, tension, and the single, tangible knob L.

Setup (boundary sector; same instrumentation). Work on the certified boundary plateau $\mathfrak{P}_\partial$ with tangential IR window $K^{(L)}_{\text{IR},\partial}$ and IR knob $r{=}L$, evaluate in $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$ with BRST–closed tags $P_x(\lambda_{\text{UV}})$, and adopt the boundary witnesses of Definition 8.31. Thus the edge dials are $\left(\Lambda^{[\partial]}_{\text{law}}(L), \Lambda^{[\partial]}_{\text{state}}\right) \in [0,1]^2$, and the ledgers consist of the *edge Ward ledger* (large–gauge/asymptotic charges, improved boundary current $J^{(\partial)}_{\parallel}$) and the *edge spectral ledger* (soft/edge poles, residues $Z \succeq 0$, memory signals). Parasymmetry overlays these inputs with phase–sensitive meters while preserving the quadrant resolver of Proposition 8.32.

Definition 8.133 (Boundary parasymmetric calibration and densities)**.** A *boundary parasymmetric calibration* is a completely positive, trace–preserving map

$$\kappa^{[\partial]} : \left(\Lambda^{[\partial]}_{\text{law}}(L), \Lambda^{[\partial]}_{\text{state}};\ \mathcal{W}^{[\partial]}, \mathcal{Z}^{[\partial]}\right) \longrightarrow \left(\rho^{(L)[\partial]},\ \rho^{(S)[\partial]},\ \rho^{(LS)[\partial]}\right),$$

where $\rho^{(L)[\partial]}, \rho^{(S)[\partial]} \in \mathsf{D}(\mathbb{C}^2)$ encode local "Sym/Break" content for the *law* and *state* anchors at the boundary, and $\rho^{(LS)[\partial]} \in \mathsf{D}(\mathbb{C}^2 \otimes \mathbb{C}^2)$ encodes relational law–state boundary content. Calibration is performed on $(\mathfrak{P}_\partial; L; P_x, K^{(L)}_{\text{IR},\partial}; \mathsf{P}_{\text{phys}})$ and is Nielsen–flat by BRST closure.

Definition 8.134 (Boundary parasymmetric meters)**.** Let $P^{(a)}_{\mathrm{para}} = |+\rangle_a\langle+|$ $(a \in \{L, S\})$ be local parasymmetric projectors, and $P^{(LS)}_{\mathrm{para}}$ (coherent *opposition*, II/III diagonal) and $P^{(LS)}_{\Phi,\mathrm{para}}$ (coherent *agreement*, I/IV diagonal) be Bell–type projectors on the relational space. Define:

$$v^{(L)[\partial]}_{\mathrm{Synthesis}} = \mathrm{Tr}\Big(P^{(L)}_{\mathrm{para}}\rho^{(L)[\partial]}\Big), \quad v^{(S)[\partial]}_{\mathrm{Synthesis}} = \mathrm{Tr}\Big(P^{(S)}_{\mathrm{para}}\rho^{(S)[\partial]}\Big),$$

$$v^{(LS)[\partial]}_{\mathrm{Synthesis}} = \mathrm{Tr}\Big(P^{(LS)}_{\mathrm{para}}\rho^{(LS)[\partial]}\Big), \qquad v^{(LS)[\partial]}_{\mathrm{Agreement}} = \mathrm{Tr}\Big(P^{(LS)}_{\Phi,\mathrm{para}}\rho^{(LS)[\partial]}\Big).$$

Normalization: 1 indicates a coherent "both/and", 1/2 the incoherent mixture baseline, and 0 a coherent "anti–both''.

Lemma 8.135 (Boundary cross–coherence (soft/edge channels))**.** *Let $\langle\cdot,\cdot\rangle_{\mathfrak{P}_\partial}$ be a plateau–uniform, BRST–safe seminorm. A canonical relational off–diagonal is*

$$\rho^{(LS)[\partial]}_{12} \;\propto\; \frac{\big\langle S_\partial,\ \mathcal{O}_\partial\big\rangle_{\mathfrak{P}_\partial}}{\sqrt{\langle S_\partial, S_\partial\rangle_{\mathfrak{P}_\partial}\ \langle \mathcal{O}_\partial, \mathcal{O}_\partial\rangle_{\mathfrak{P}_\partial}}},$$

where $\mathcal{O}_\partial$ is an edge operator matched to the Ward/spectral ledger (e.g. chiral edge current, soft charge density, or memory/vertex operator). Then

$$v^{(LS)[\partial]}_{\mathrm{Synthesis}} = \tfrac{1}{2} + \mathrm{Re}\ \rho^{(LS)[\partial]}_{12}, \qquad v^{(LS)[\partial]}_{\mathrm{Agreement}} = \tfrac{1}{2} + \mathrm{Re}\ \rho^{(LS)[\partial]}_{\Phi,12}.$$

Proposition 8.136 (Boundary quadrant–parasymmetry correspondence)**.** *Fix thresholds $\tau_{\mathrm{agr}}, \tau_{\mathrm{syn}} \in (0, \frac{1}{2})$. On any certified boundary plateau component:*

$$\begin{aligned} \mathrm{QI}\ \textit{or}\ \mathrm{QIV} &\implies v^{(LS)[\partial]}_{\mathrm{Agreement}} \ \geq\ \tfrac{1}{2} + \tau_{\mathrm{agr}} \quad (\textit{coherent I/IV diagonal}),\\ \mathrm{QII}\ \textit{or}\ \mathrm{QIII} &\implies v^{(LS)[\partial]}_{\mathrm{Synthesis}} \ \geq\ \tfrac{1}{2} + \tau_{\mathrm{syn}} \quad (\textit{coherent II/III diagonal}). \end{aligned}$$

Moreover: (i) in boundary–QII, $v^{(S)[\partial]}_{\mathrm{Synthesis}} \to \frac{1}{2}$ with increasing boundary extent (local mixture baseline; edge Goldstone present); (ii) in boundary–QIII, $v^{(LS)[\partial]}_{\mathrm{Synthesis}}(L)$ increases monotonically up to $o(1)$ as $d^{[\partial]}_{\mathrm{law}}(L)\downarrow$; (iii) in boundary–QIV, $v^{(LS)[\partial]}_{\mathrm{Agreement}}$ remains persistently elevated and the edge spectral ledger exhibits a pNG band on $\mathfrak{P}_\partial$.

Definition 8.137 (Operational overlay (boundary, SynTh + parasymmetry))**.** Given $\mathfrak{P}_\partial$ and L, proceed as follows:

(i) Quadrant assignment.
Assign QI–QIV via $\mathcal{R}^{[\partial]}$ using Definition 8.31 and Proposition 8.32.

(ii) Calibration and meters.
Calibrate $\kappa^{[\partial]}$ and compute $v^{(L)[\partial]}_{\mathrm{Synthesis}}$, $v^{(S)[\partial]}_{\mathrm{Synthesis}}$, $v^{(LS)[\partial]}_{\mathrm{Synthesis}}$, and $v^{(LS)[\partial]}_{\mathrm{Agreement}}$.

(iii) Coherence interpretation.
Use $\tau_{\mathrm{agr}}, \tau_{\mathrm{syn}}$ to call elevated meters and identify the occupied diagonal (I/IV vs II/III); detect local "bothness'' via the 1/2 baseline.

(iv) Ward/spectral cross-checks.
Cross–check ledgers: (QII) edge Goldstone and conserved improved edge current; (QIV) persistent boundary source with pNG band; (QIII) vanishing adjoint divergence along L and no NG/pNG.

(v) Steering via L and boundary conditions.
For steering: vary L and boundary conditions to test healing vs persistence; monitor meter conservation where the relevant projectors commute (or satisfy $\mathcal{L}^\dagger(P) = 0$).

Narrative and intuition (boundary punchline). The boundary parasymmetric overlay turns the tolerance–form witnesses of Definition 8.31 into *instrument–grade* coherence diagnostics. Explicit edge breaking (QIV) manifests as coherent *agreement* between law and state—elevated $v^{(LS)[\partial]}_{\text{Agreement}}$ on $\mathfrak{P}_\partial$—and aligns with a persistent boundary source and a pNG band in the edge spectrum. Emergent edge symmetry (QIII) is visible as a monotone rise of $v^{(LS)[\partial]}_{\text{Synthesis}}$ with L in tandem with the healing of the boundary law deficit $d^{[\partial]}_{\text{law}}(L)\downarrow$. Spontaneous edge selection (boundary–QII) presents the finite–extent signature $v^{(S)[\partial]}_{\text{Synthesis}} \to \frac{1}{2}$ with an elevated opposition meter, certifying an edge Goldstone under an intact boundary law. All meters are BRST–safe and Nielsen–flat, and therefore stable under admissible retunings of the boundary instrumentation. In multi–sector settings (bulk & boundary), the joint resolver pairs these boundary coherence profiles with the bulk ones to distinguish benign arena mismatches from genuine boundary–localized symmetry mechanisms.

8.6 Parasymmetric Diagnostics of Type, Location, and Onset (TLO)

All statements below are made on the BRST–physical sector $\mathcal{H}_{\text{phys}} := H^0(\hat{Q}_{\text{BRST}})$ with realized weave configuration $(P_x(\lambda_{\text{UV}}), K^{(\ell)}_{\text{IR}}; \mathsf{P}_{\text{phys}})$ on a plateau $\mathfrak{P}$ (or boundary plateau $\mathfrak{P}_\partial$), as in Sections 8.1, 8.2 and 8.2.1. Gauge/Nielsen safety and regulator robustness follow from the bandwise identities and algebraic–renormalisation arguments cited earlier. [117, 442, 114, 91]

8.6.1 Parasymmetric Meters and Laws-States Quadrant Readings

We use the local parasymmetric projectors $P^{(a)}_{\text{para}}$ and valuations $v^{(a)}_{\text{Synthesis}}(\rho_a) = \text{Tr}\big(P^{(a)}_{\text{para}}\rho_a\big) \in [0,1]$ for each anchor $a \in \{L, S\}$, together with the relational meter $v^{(LS)}_{\text{Synthesis}}$, as fixed in Section 8.2.1. In the two-mode basis $|e_1\rangle_a \equiv |\,\text{Sym}\rangle_a$, $|e_2\rangle_a \equiv |\,\text{Break}\rangle_a$, the explicit form $v^{(a)}_{\text{Synthesis}} = \frac{1}{2} + \text{Re}(\rho^{(a)}_{12})$ will be used for diagnostics.

Quadrant II (State–SSB; law $\simeq 1$)

Finite-window elevation of the *state meter*,

$$v^{(S)}_{\text{Synthesis}}(\rho_S) = \tfrac{1}{2} + \text{Re}(\rho^{(S)}_{12}) > \tfrac{1}{2},$$

with *decay to baseline* 1/2 under IR/volume dilation $r \uparrow$ (no persistent coherence), identifies state–SSB. Concomitant spectral NG content and the resolver conditions $\Lambda_{\text{law}} \simeq 1$, $\Lambda_{\text{state}}(V) \to 0$ complete the QII certification; see Section 7.10 and Lemma 8.87. [82, 91]

Quadrant IV (explicit breaking; persistent law deficit)

A *non-healing* law deficit $d_{\text{law}}^{[\mathscr{S}]}(r) \geq \delta_\star$ for all large r, together with a *baseline* state meter $v_{\text{Synthesis}}^{(S)} \approx \frac{1}{2}$ (no "cat'' coherence), and a pNG band obeying the plateau GMOR slope (eq. (7.747)), identifies explicit breaking. The parasymmetric meters record *persistence* (no decay toward the QIII baseline), in agreement with the canonical resolver of Definition 7.685.

Quadrant III (law–healing; emergent symmetry)

When the law asymmetry is *irrelevant*, cross-terms and law–state parasymmetric valuations *decay* to 1/2 with the IR knob according to the power law $d_{\text{law}}^{[\mathscr{S}]}(r) \leq Cr^{-\beta}$ (eq. (7.746)); no NG/pNG survive in the strict IR. The meters thus diagnose *healing* rather than SSB or explicit persistence.

Quadrant I (fully symmetric)

Both anchors register *baseline* values $v_{\text{Synthesis}}^{(L,S)} \approx \frac{1}{2}$, consistent with $|\Lambda_{\text{law}}| \approx |\Lambda_{\text{state}}| \approx 1$ and the symmetry-respecting census.

8.6.2 Locator Sets: "Where" the Patterns Occur

Let $\mathcal{A} \in \{\mathfrak{P}, \mathfrak{P}_\partial\}$ denote the arena (bulk/boundary) and r the IR window ($r = \ell$ in bulk, $r = L$ on the boundary). Using the sector tag $[\mathscr{S}]$ from Section 8.1, define *window-resolved loci*:

$$\mathcal{L}_{\text{SSB}}^{[\mathscr{S}]} = \Big\{(x,r) : \ |\Lambda_{\text{law}}^{[\mathscr{S}]}| \geq 1 - \varepsilon_{\text{law}},\ v_{\text{Synthesis}}^{(S)}(x;r) > \tfrac{1}{2},\ v_{\text{Synthesis}}^{(LS)}(x;r) \xrightarrow{r\uparrow} \tfrac{1}{2}\Big\},$$

$$\mathcal{L}_{\text{QIV}}^{[\mathscr{S}]} = \Big\{(x,r) : \ d_{\text{law}}^{[\mathscr{S}]}(r) \geq \delta_\star,\ v_{\text{Synthesis}}^{(S)}(x;r) \approx \tfrac{1}{2},\ v_{\text{Synthesis}}^{(LS)}(x;r) \text{ non-healing}\Big\},$$

$$\mathcal{L}_{\text{QIII}}^{[\mathscr{S}]} = \Big\{(x,r) : \ d_{\text{law}}^{[\mathscr{S}]}(r) \leq Cr^{-\beta},\ v_{\text{Synthesis}}^{(L)}(x;r) \to \tfrac{1}{2},\ v_{\text{Synthesis}}^{(LS)}(x;r) \to \tfrac{1}{2}\Big\},$$

and analogously $\mathcal{L}_{\text{QI}}^{[\mathscr{S}]}$ for the symmetric case. Because $P_x(\lambda_{\text{UV}})$ is centered at x and $K_{\text{IR}}^{(\ell)}$ sets r, scans $x \mapsto P_x$ and $r \mapsto \ell$ (or L) produce *maps* of these loci across space and scale, per sector $[\mathscr{S}]$.

8.6.3 Onset Scales and Times: "When" the Patterns Occur

Define *scale onsets* by certified tolerances $(\varepsilon_{\mathrm{law}}, \varepsilon_{\mathrm{state}})$:

$$r^\star_{\mathrm{SSB}} = \inf\{r : \ v^{(S)}_{\mathrm{Synthesis}}(r) > \tfrac{1}{2} \text{ and } v^{(LS)}_{\mathrm{Synthesis}}(r) \downarrow \tfrac{1}{2}\},$$

$$r^\star_{\mathrm{QIV}} = \inf\{r : \ d^{[\mathscr{S}]}_{\mathrm{law}}(r) \geq \delta_\star\}, \qquad r^\star_{\mathrm{QIII}} = \inf\{r : \ d^{[\mathscr{S}]}_{\mathrm{law}}(r) \leq Cr^{-\beta}\}.$$

If a dynamics generator for the relational sector is provided (e.g. $\hat{F}^{(LS)}_{\mathrm{para}}$), one may likewise define *onset times* $t^\star$ from $t \mapsto v^{(a)}_{\mathrm{Synthesis}}(t; r)$ under the same inequalities.

8.6.4 Boundary/Edge Arenas and Sector Checks

On $\mathfrak{P}_\partial$, elevated *tangential or chiral* parasymmetric valuations certify *edge Goldstones* (QII) or *edge pseudo-modes* (QIV) with small gaps; decay to baseline corroborates QIII healing. The construction is *functorial* under admissible dualities and *cone-preserving*, so it *refines* but does not alter the quadrant assignments fixed by SynTh (Section 8.2).

Proposition 8.138 (Parasymmetric resolver is quadrant-sound and gauge-safe)**.** *On the certified arena, with meters* $v^{(a)}_{\mathrm{Synthesis}}$ *and* $v^{(LS)}_{\mathrm{Synthesis}}$ *of Section 8.2.1 evaluated in* $\mathcal{H}_{\mathrm{phys}}$*, the following implications hold:*

(i) QI.
Baseline meters equal to $\frac{1}{2}$ *and symmetric visibilities imply the unbroken regime.*

(ii) QIII.
Power-law decay $d^{[\mathscr{S}]}_{\mathrm{law}}(r) \leq Cr^{-\beta}$ *and meters tending to* $\frac{1}{2}$ *imply IR healing (no NG/pNG in the strict IR).*

(iii) QII.
Elevated $v^{(S)}_{\mathrm{Synthesis}}$ *at finite* r and *decay to* $1/2$ *as* $r \uparrow$*, together with NG poles, implies state–SSB with symmetric law.*

(iv) QIV.
A non-healing law deficit and pNG gaps obeying eq. (7.747), with $v^{(S)}_{\mathrm{Synthesis}} \approx \frac{1}{2}$*, imply persistent explicit breaking.*

Moreover, all valuations are Nielsen-flat (gauge-parameter independent) on the band, and stable under admissible regulator changes within $\varepsilon_{\mathrm{reg}}$*.* □

Parasymmetry provides *meters* whose *magnitudes and large-window behavior* distinguish the four quadrants while localising *where* (bulk/boundary, sector, position) and *when* (onset scale—and, if applicable, onset time) the pattern holds:

SSB (QII)	$\Rightarrow$	elevated then decaying $v^{(S)}_{\mathrm{Synthesis}}$ (NG present),
explicit (QIV)	$\Rightarrow$	persistent law deficit with GMOR pNG gaps,
healing (QIII)	$\Rightarrow$	decay to baseline without NG/pNG,
symmetric (QI)	$\Rightarrow$	baseline throughout.

These diagnostics are *BRST/Nielsen-safe*, *cone-preserving*, and compatible with the SynTh resolver, thereby turning the conceptual 2×2 into a *workable cartography* across sectors $[\mathscr{S}]$, arenas $\mathcal{A}$, and scales r.

8.6.5 Typed Sector Witnesses and Parasymmetric Meters (Spin/Helicity/Polarization–conditioned)

All constructions in this subsection are evaluated on the BRST–physical sector $\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}})$ with realized weave configuration $(\mathcal{A}, \mathfrak{P}; r; P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}^{(\ell)}; \mathsf{P}_{\mathrm{phys}})$ as in Sections 8.1, 8.2 and 8.2.1. They preserve BRST/Nielsen safety and plateau stability by Proposition 7.644 and Theorem 7.646.

Definition 8.139 (Typed sector witness and typed state)**.** Let $[\mathscr{S}]$ be a sector with witness $\mathbf{W}^{[\mathscr{S}]}$. For a BRST–closed projector ("filter") $\Pi^{[\chi]}$ selecting a kinematic/spectral class χ (spin, helicity, polarization, chirality, or an isolated species), define

$$\mathbf{W}^{[\mathscr{S};\chi]} \;:=\; \Pi^{[\chi]}\,\mathbf{W}^{[\mathscr{S}]}\,\Pi^{[\chi]}, \qquad \rho^{[\chi]} \;:=\; \frac{\Pi^{[\chi]}\,\rho\,\Pi^{[\chi]}}{\mathrm{Tr}\big(\Pi^{[\chi]}\rho\big)} \quad (\text{on } \mathcal{H}_{\mathrm{phys}}). \tag{8.18}$$

Definition 8.140 (Typed visibilities and typed parasymmetric meters)**.** Using Definition 7.35 with $\mathbf{W}^{[\mathscr{S};\chi]}$ and $\rho^{[\chi]}$, set

$$\Lambda_{\mathrm{law}}^{[\mathscr{S};\chi]}, \qquad \Lambda_{\mathrm{state}}^{[\mathscr{S};\chi]}. \tag{8.19}$$

With $a \in \{L, S\}$ and $P_{\mathrm{para}}^{(a)}$ as in Section 8.2.1, define the typed parasymmetric valuations

$$v_{\mathrm{Synthesis}}^{(a);\,[\chi]}(\rho_a) \;:=\; \mathrm{Tr}\Big(P_{\mathrm{para}}^{(a)}\,\rho_a^{[\chi]}\Big) \in [0,1], \qquad v_{\mathrm{Synthesis}}^{(LS);\,[\chi]} \;\text{ analogously.} \tag{8.20}$$

Canonical filters $\Pi^{[\chi]}$ (on $\mathcal{H}_{\mathrm{phys}}$). All items below are BRST–closed and commute with $\mathsf{P}_{\mathrm{phys}}$.

(i) Spin filters. $\Pi^{[s=2]} := \Pi_{\mathrm{TT}}$ (transverse–traceless tensor), $\Pi^{[s=1]} := P_T$ (transverse vector), $\Pi^{[s=0]} := \mathbf{1}_{\mathrm{scalar}}$. These are the polarization blocks $X \in \{h_{\mathrm{TT}}, A^T, \pi\}$ used in the pole analysis.

(ii) Helicity filters (momentum–resolved). For spin–1, $\Pi^{[1,\lambda=\pm1]}(\hat{\mathbf{k}}) := |e_\pm(\hat{\mathbf{k}})\rangle\langle e_\pm(\hat{\mathbf{k}})| \subset \mathrm{Im}\,P_T$. For spin–2, $\Pi^{[2,\lambda=\pm2]}(\hat{\mathbf{k}}) := |e_{\pm2}(\hat{\mathbf{k}})\rangle\langle e_{\pm2}(\hat{\mathbf{k}})| \subset \mathrm{Im}\,\Pi_{\mathrm{TT}}$. (Implemented as covariant little–group projectors, averaged on the IR window.)

(iii) Polarization filters (hydrodynamics/elasticity). $P_L^{ij}(\hat{\mathbf{k}}) = \hat{k}^i\hat{k}^j$, $P_T^{ij}(\hat{\mathbf{k}}) = \delta^{ij} - \hat{k}^i\hat{k}^j$; shear vs. sound sub-blocks are obtained by composing $\Pi^{[\chi]}$ with the hydrodynamic witness.

(iv) Boundary tangential/chiral filters. Tangential $\Pi^{[\|]}$ on $\partial\mathcal{X}$ and chiral $\Pi^{[\chi=\pm]}$ for edge modes; compose with $\mathbf{W}^{[\partial]}$ on $\mathfrak{P}_\partial$.

(v) Isolated–pole (species) filters. For a species $\mathfrak{m}$ with an isolated pole of $\mathcal{K}_{\mathrm{phys}}$ (or $\mathcal{D}_{\mathrm{phys}}$), the Riesz projector

$$\mathcal{P}_{\mathfrak{m}} := \frac{1}{2\pi i}\oint_{C_{\mathfrak{m}}} dz\,\big(z-\mathcal{K}_{\mathrm{phys}}\big)^{-1} \tag{8.21}$$

selects that mode (mass eigenstate or narrow band). Put $\Pi^{[\chi=\mathfrak{m}]} := \mathcal{P}_{\mathfrak{m}}$ to obtain a "particle meter".

Proposition 8.141 (Gauge/Nielsen safety and analytic transport (typed))**.** *Let $\Pi^{[\chi]}$ be a BRST–closed filter as in Definition 8.139. Then the typed visibilities $\Lambda_{\bullet}^{[\mathscr{S};\chi]}$ and the typed parasymmetric valuations $v_{\mathrm{Synthesis}}^{(\cdot),[\chi]}$ are gauge–parameter independent on certified plateaux. Moreover, under small explicit deformations the corresponding spectral projectors and residue matrices vary analytically on $\mathfrak{P}$, hence these quantities transport analytically.*

Proof sketch. $\Pi^{[\chi]}$ can be written as $\mathsf{P}_{\mathrm{phys}}\, f(\mathcal{K}_{\mathrm{phys}})\,\mathsf{P}_{\mathrm{phys}}$ (bounded functional calculus) or as a little–group projector inside a physical block; gauge–parameter variations are ζ-exact and cancel on $\mathcal{H}_{\mathrm{phys}}$, giving Nielsen flatness by Proposition 7.644. Analytic transport of isolated spectral projectors and residues on $\mathfrak{P}$ follows from the band/BRST analyticity theorem Theorem 7.646. □ □

Reading rules (per χ). For each χ, apply the canonical quadrant decision rule to $\big(\Lambda_{\mathrm{law}}^{[\mathscr{S};\chi]}, \Lambda_{\mathrm{state}}^{[\mathscr{S};\chi]}\big)$ (cf. Definition 7.685). The typed parasymmetric meters $v_{\mathrm{Synthesis}}^{(a);[\chi]}$ and $v_{\mathrm{Synthesis}}^{(LS);[\chi]}$ then *localize* and *resolve in scale/time* where and when the χ–filtered content occurs (SSB vs. explicit vs. healing vs. symmetric), exactly as in the untyped case but confined to the chosen spin/helicity/polarization subspace.

Examples.

(i) Spin meter. With $\Pi^{[s]}$, record $r \mapsto v_{\mathrm{Synthesis}}^{(S);[s]}(r)$ alongside $\Lambda_{\mathrm{law}}^{[\mathscr{S};s]}(r)$. QII for $s=0$: finite–window elevation of $v_{\mathrm{Synthesis}}^{(S);[0]}$ with decay to $1/2$ and NG poles in $\mathcal{Z}^{[\mathscr{S};0]}$. QIV for $s=0$: persistent departure aligned with GMOR gaps in $\mathcal{Z}^{[\mathscr{S};0]}$. QI/QIII for $s=1,2$: unbroken massless carriers and IR healing.

(ii) Helicity meter (spin–1). $\Pi^{[1,\lambda=\pm1]}(\hat{\mathbf{k}})$ yields $v_{\mathrm{Synthesis}}^{(L);[1,\lambda]}(r)$, distinguishing helicity–selective edge transport or anisotropic media on $\mathfrak{P}_{\partial}$.

(iii) Polarization meter (hydrodynamics). With $P_{L/T}$, obtain longitudinal–sound vs. transverse–shear diagnostics, enabling QII (sound NG) vs. QIV (pseudo–sound) discrimination inside the hydrodynamic sector.

(iv) Species ("particle") meter. For $\Pi^{[\chi=\mathfrak{m}]} = \mathcal{P}_{\mathfrak{m}}$ (e.g. a specific pNG, a radial scalar, a massive vector eigenstate), the typed meters report *where/when* that mode is realized and *which breaking* accounts for it.

Typing the sector witness by BRST–closed filters $\Pi^{[\chi]}$ integrates spectral/polarization selection into parasymmetry. The resulting spin/helicity/polarization/particle meters retain the spin labels fixed by the spectral certificate $\mathcal{Z}^{[\mathscr{S}]}$ and deliver *type* (QI–QIV), *location* (bulk/boundary, sector, position), and *onset* (scale/time) diagnostics on the same BRST–clean, plateau–stable footing as the untyped SynTh/parasymmetry framework.

Complete identification stack. *Qualified yes.* Within the realized weave configuration on the BRST–physical sector $\mathcal{H}_{\rm phys}$—namely the band tags $(P_x(\lambda_{\rm UV}), K_{\rm IR}^{(\ell)})$ on a plateau $\mathfrak{P}$ (or boundary plateau $\mathfrak{P}_\partial$)—the Simultaneous Synthesis (**SynTh**) supplies the logical resolver of quadrant labels, while the parasymmetric layer furnishes phase–sensitive meters that *realize* that logic operationally (Sections 8.2 and 8.2.1). The result is a gauge/Nielsen–safe stack that, from the same sector–conditioned witnesses, returns: the *type* of symmetry status (QI–QIV), the *quadrant* itself (Section 7.10), the *where* (arena/sector/position), the *when* (onset scale/time), and—via typed witnesses—the *what* (spin/helicity/polarization/species).

Type and quadrant. Given sector–conditioned visibilities $(\Lambda_{\rm law}^{[\mathscr{S}]}, \Lambda_{\rm state}^{[\mathscr{S}]})$ and parasymmetric valuations $v_{\rm Synthesis}^{(L)}, v_{\rm Synthesis}^{(S)}, v_{\rm Synthesis}^{(LS)}$, the canonical decision rule (Section 7.10) returns the quadrant: QI (symmetric), QII (state–SSB: elevated then decaying coherence), QIII (law–healing with power–law decay), or QIV (explicit breaking: persistent law deficit with GMOR–controlled pNGs). Parasymmetric meters provide the phase–coherence corroborants that distinguish "elevated–then–decaying" (QII) from "persistent" (QIV) behavior on the same diagnostic realized reference configuration (Section 8.2.1).

Where (arena/sector/position). The arena $\mathcal{A} \in \{\mathfrak{P}, \mathfrak{P}_\partial\}$ resolves bulk versus boundary; the sector tag $[\mathscr{S}]$ (hydrodynamic, boundary/edge, p-forms via duality, higher–spin, fermions/SUSY, *etc.*) is provided by the sector–conditioned witnesses of Section 8.1; spatial localization follows from the centered projector $P_x(\lambda_{\rm UV})$, and the IR knob sets the scale ($r = \ell$ in bulk, $r = L$ on ∂). Window–resolved *maps* are obtained by scanning x and r. Typical locator sets are

$$\mathcal{L}_{\rm SSB}^{[\mathscr{S}]} = \Big\{(x,r):\ v_{\rm Synthesis}^{(S)}(x;r) > \tfrac{1}{2},\quad v_{\rm Synthesis}^{(LS)}(x;r) \xrightarrow{r\uparrow} \tfrac{1}{2}\Big\}, \tag{8.22}$$

$$\mathcal{L}_{\rm QIV}^{[\mathscr{S}]} = \Big\{(x,r):\ d_{\rm law}^{[\mathscr{S}]}(r) \geq \delta_\star,\quad \text{non-healing}\Big\}, \tag{8.23}$$

with analogous sets for QIII (healing) and QI (symmetric).

When (onset scale/time). Onset scales are defined by the first window $r^\star$ at which the quadrant inequalities hold within chosen tolerances $(\varepsilon_{\rm law}, \varepsilon_{\rm state})$:

$$r_{\rm SSB}^\star = \inf\{r:\ v_{\rm Synthesis}^{(S)}(r) > \tfrac{1}{2}\ \wedge\ v_{\rm Synthesis}^{(LS)}(r) \downarrow \tfrac{1}{2}\}, \tag{8.24}$$

$$r_{\rm QIV}^\star = \inf\{r:\ d_{\rm law}^{[\mathscr{S}]}(r) \geq \delta_\star\}, \tag{8.25}$$

$$r_{\rm QIII}^\star = \inf\{r:\ d_{\rm law}^{[\mathscr{S}]}(r) \leq C r^{-\beta}\}. \tag{8.26}$$

If a relational generator $\hat{F}_{\rm para}^{(LS)}$ is provided (Section 8.2.1), one likewise defines onset times $t^\star$ from $t \mapsto v_{\rm Synthesis}^{(\cdot)}(t;r)$.

Spins, helicities, polarizations, and species. These are supplied by *typed witnesses* $\mathbf{W}^{[\mathscr{S};\chi]}$ that compose the sector witness with a BRST–closed filter $\Pi^{[\chi]}$ (Section 8.6.5). Canonical choices include spin filters $\Pi^{[s=2]} = \Pi_{\rm TT}$, $\Pi^{[s=1]} = P_T$, $\Pi^{[s=0]} = \mathbf{1}_{\rm scalar}$; helicity filters $\Pi^{[1,\lambda=\pm1]}(\hat{\mathbf{k}})$, $\Pi^{[2,\lambda=\pm2]}(\hat{\mathbf{k}})$ within the physical blocks; polarization filters such as P_L vs P_T in hydrodynamics and tangential/chiral projectors on $\mathfrak{P}_\partial$; and species ("particle") filters given by Riesz projectors $\mathcal{P}_{\mathfrak{m}}$ onto isolated poles/bands of $\mathcal{K}_{\rm phys}$ (or $\mathcal{D}_{\rm phys}$). The typed visibilities $\big(\Lambda_{\rm law}^{[\mathscr{S};\chi]}, \Lambda_{\rm state}^{[\mathscr{S};\chi]}\big)$ and typed parasymmetric valuations $v_{\rm Synthesis}^{(\cdot);[\chi]}$ then localize *where/when* a chosen χ (spin/helicity/polarization/species) is realized and *why* (SSB vs

explicit vs healing vs symmetric), without modifying the spin labels already fixed by the physical–sector pole analysis.

Scope and caveats. All identifications are exact up to the *window resolution* determined by P_x and K_{IR}; species–level meters require isolated poles or sufficiently narrow bands, else the output is a controlled coarse–graining over the selected block. Functoriality and cone–preservation ensure that refining the instrumentation sharpens readout *without changing* quadrant membership.

With SynTh delivering the paraconsistent resolver and parasymmetry providing phase–sensitive (typed) meters on $\mathcal{H}_{\mathrm{phys}}$, the combined instrument identifies, on certified plateaux, the *type* and *quadrant* of symmetry status, *where* it holds (arena/sector/position), *when* it onsets (scale/time), and—via typed filters—the *spin*, *helicity*, *polarization*, and *species* content of the light sector (Sections 7.10, 8.1, 8.2, 8.2.1 and 8.6.5).

8.6.6 Spectral–Only Parasymmetric Protocol

All statements in this section are evaluated on the BRST–physical sector $\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}})$ with realized weave configuration $(P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}^{(\ell)}; \mathsf{P}_{\mathrm{phys}})$. The protocol uses spectra alone—without the Ward ledger—to reconstruct spin/helicity/polarization content and to diagnose *what* breaks, *where*, and *when*, consistently with the quadrant resolver of Section 7.10 and the meters of Section 8.2.1.

Definition 8.142 (Spectral certificate on $\mathcal{H}_{\mathrm{phys}}$)**.** From the band/BRST–reduced quadratic kernel build the physical propagator $\mathcal{D}_{\mathrm{phys}}^{(\lambda_{\mathrm{UV}},\ell)}$. Decompose into polarization blocks

$$X \in \{h_{\mathrm{TT}},\, A^T,\, \pi\} \qquad \Rightarrow \qquad s(X) = 2, 1, 0,$$

using Π_{TT} (transverse–traceless), P_T (transverse) and the scalar block (cf. tables 7.5 and 7.6). The *spectral certificate* is the blockwise list

$$\mathcal{Z} \;:=\; \Big\{(\omega_n(\mathbf{k}),\, Z_n)\Big\}_{n,X},$$

collecting pole locations $\omega_n(\mathbf{k})$ and residues Z_n in each physical block X.

Spin/helicity/polarization identification (from $\mathcal{Z}$). Spins are assigned by block: $h_{\mathrm{TT}} \to s{=}2$, $A^T \to s{=}1$, $\pi \to s{=}0$. Massless helicities are resolved by little–group projectors (e.g. $\lambda = \pm 2$ in $\mathrm{Im}\,\Pi_{\mathrm{TT}}$, $\lambda = \pm 1$ in $\mathrm{Im}\,P_T$). Sectoral polarizations (hydro/elastic/edge) are read with P_L/P_T or tangential/chiral filters on $\mathfrak{P}_\partial$ (cf. table 7.6).

Lemma 8.143 (Spectral quadrant signatures)**.** *Using only $\mathcal{Z}$, the quadrant labels are diagnosed spectrally as follows:*

(i) QI (law ≈ 1, state ≈ 1).
No NG/pNG; unbroken massless $s = 2$ and $s = 1$ carriers present where allowed; radial gaps analytic.

(ii) QII (state SSB).
Presence of NG scalars *(gapless $s = 0$); for gauged broken directions,* massive $s = 1$ *with a restored longitudinal polarization (BEH).*

(iii) QIII (healing).
No persistent NG/pNG*; spectra reconverge to the QI census under IR dilation (*$\mathbf{k} \to 0$ *or* $\ell \uparrow$*) in accordance with eq. (7.746).*

(iv) QIV (explicit).
pNG band *with GMOR scaling* $m_{\pi_b}^2 = \mathcal{O}(\epsilon)$*; pseudo–phonons on* $\mathfrak{P}_\partial$ *share L/T structure but are gapped (cf. eq. (7.747)).*

Hence spectra alone determine what *breaks (SSB vs. explicit) and* which *quadrant is indicated, consistent with Section 7.10.*

Definition 8.144 (Spectral locator sets and onset scales)**.** For a sector tag $[\mathscr{S}]$ (cf. Section 8.1), define window–resolved loci using $\mathcal{Z}$:

$$\mathcal{L}_{\mathrm{SSB}}^{[\mathscr{S}]}(\mathcal{Z}) := \left\{ \begin{array}{l} (x,r) : \text{NG poles present;} \\ \qquad v_{\mathrm{Synthesis}}^{(S)}(x;r) > \tfrac{1}{2},\ v_{\mathrm{Synthesis}}^{(LS)}(x;r) \xrightarrow{r\uparrow} \tfrac{1}{2} \end{array} \right\}, \tag{8.27}$$

$$\mathcal{L}_{\mathrm{QIV}}^{[\mathscr{S}]}(\mathcal{Z}) := \left\{ \begin{array}{l} (x,r) : \text{pNG band with GMOR slope;} \\ \qquad \text{non-healing parasymmetric cross–terms} \end{array} \right\}, \tag{8.28}$$

with analogous $\mathcal{L}_{\mathrm{QI}}^{[\mathscr{S}]}(\mathcal{Z})$ and $\mathcal{L}_{\mathrm{QIII}}^{[\mathscr{S}]}(\mathcal{Z})$. The *arena* is fixed by $\mathcal{A} \in \{\mathfrak{P}, \mathfrak{P}_\partial\}$ (bulk $r = \ell$, boundary $r = L$). Onset scales are

$$\begin{aligned} r_{\mathrm{SSB}}^\star &= \inf\{r :\ \text{NG present and } v_{\mathrm{Synthesis}}^{(LS)}(r) \downarrow \tfrac{1}{2}\}, \\ r_{\mathrm{QIV}}^\star &= \inf\{r :\ \text{pNG with GMOR slope}\}, \\ r_{\mathrm{QIII}}^\star &= \inf\{r :\ \text{healing per eq. (7.746)}\}. \end{aligned}$$

Spectral sweeps ("where" and "when"). Arena classification follows dispersion geometry (bulk isotropic vs. tangential/chiral along $\partial\mathcal{X}$); edge Goldstones vs. pseudo–edge modes discriminate boundary QII/QIV. Spatial localization uses $x \mapsto P_x(\lambda_{\mathrm{UV}})$ and the IR knob r to map the loci of Definition 8.144. Even with a single spectrum, the low–$|\mathbf{k}|$ tail acts as an IR sweep: NG/pNG vs. healing is read from pole behavior as $|\mathbf{k}| \to 0$.

Typed meters from spectral filters (optional but powerful). Compose the sector witness with a BRST–closed filter $\Pi^{[\chi]}$ (spin/helicity/polarization/species) to obtain typed meters, as in Section 8.6.5:

$$\mathbf{W}^{[\mathscr{S};\chi]} = \Pi^{[\chi]}\, \mathbf{W}^{[\mathscr{S}]}\, \Pi^{[\chi]}, \qquad v_{\mathrm{Synthesis}}^{(\cdot);[\chi]} = \mathrm{Tr}\Big(P_{\mathrm{para}}^{(\cdot)}\, \rho^{[\chi]}\Big).$$

Riesz projectors around isolated poles yield *particle meters* for specific modes $\mathfrak{m}$, whose elevation/decay across r separates QII (elevated then decaying), QIV (persistent), QIII (healing), and QI (baseline).

Limitations. Exact law–channel visibilities (and Ward sources) require the Ward ledger; with spectra only, classification uses the spectral side of the equivalences in Section 7.10. Gauge/Nielsen safety is guaranteed provided spectra are read after the BRST projection to $\mathcal{H}_{\text{phys}}$; if given in a fixed gauge without projection, a certification step is required before applying the resolver.

Net outcome. With spectra alone, the parasymmetric framework identifies (i) the *species* and *spins/helicities/polarizations* (from block projectors and Riesz projectors), (ii) the *type of breaking* and the *quadrant* (via NG/pNG/BEH content and IR scaling), and (iii) the *arena/location* and *onset scales* $r^\star$ (and onset times if a time series is available), all on $\mathcal{H}_{\text{phys}}$ within certified plateaux/masks, consistent with Sections 8.1, 8.2 and 8.2.1.

8.6.7 Application Arenas for SynTh ⊕ Parasymmetry (Typed Witnesses)

The SynTh resolver together with parasymmetric meters on the BRST–physical sector $\mathcal{H}_{\text{phys}} := H^0(\hat{Q}_{\text{BRST}})$, equipped with instrumentation $(P_x(\lambda_{\text{UV}}), K_{\text{IR}}^{(\ell)}; \mathsf{P}_{\text{phys}})$ on certified plateaux $\mathfrak{P}$ (and $\mathfrak{P}_\partial$ at boundaries), yields a robust operational stack across sectors $[\mathscr{S}]$ (Sections 8.1, 8.2, 8.2.1 and 8.6.5). Gauge/Nielsen safety and analytic transport follow from the BRST/Nielsen framework and bandwise analyticity [117, 442, 114, 243]. Below are concise, ready-to-deploy application arenas.

1) Chiral/Higgs sectors and electroweak-like symmetry breaking. Use a gauge-sector witness $\mathbf{W}^{[\text{gauge}]}$ with spin/helicity filters $\Pi^{[s=1]}, \Pi^{[1,\lambda=\pm1]}$ and a species Riesz projector for massive vectors. Typed parasymmetric meters $v_{\text{Synthesis}}^{(a);\,[s=1]}$ locate *where/when* BEH longitudinal modes appear (QII/QIV) and separate pNG scalars (QIV, GMOR slope) from radial scalars (all quadrants), delivering type (SSB vs. explicit), arena (bulk/boundary), onset scale $r^\star$, and a spin/helicity census consistent with table 7.6. Standard BEH and spectral diagnostics are covered in [91, 82].

2) QCD-like chiral physics (NG π's, GMOR, anomaly filters). Adopt $\mathbf{W}^{[\chi\text{-global}]}$ with scalar/pseudoscalar filters $\Pi^{[s=0]}$ and species projectors for π_b. Read QII (exact NG) vs. QIV (pNG with GMOR slope) via $v_{\text{Synthesis}}^{(S);\,[0]}$ and typed visibilities $\left(\Lambda_{\text{law}}, \Lambda_{\text{state}}\right)^{[\chi;0]}$; boundary instantiations on $\mathfrak{P}_\partial$ capture edge pseudo-Goldstones and anomaly inflow while preserving quadrant logic; cf. chiral Ward/GMOR treatments in [91, 82].

3) Hydrodynamics / elasticity / MHD (L/T taxonomy). With $\mathbf{W}^{[\text{hydro}]}$ and polarization filters P_L, P_T, typed meters $v_{\text{Synthesis}}^{(\cdot);\,[L/T]}$ distinguish sound (QII), pseudo-sound (QIV, small gap) and healing to a diffusive T-sector (QIII). In MHD, add Alfvén/fast/slow projectors to resolve anisotropy and onset scales $r^\star$. The BRST-safe spectral reading is as in [243, 117].

4) Boundary/edge physics, SPT/CS anyonic edges. On $\mathfrak{P}_\partial$, take $\mathbf{W}^{[\partial]}$ with tangential/chiral filters $\Pi^{[\|]}, \Pi^{[\chi=\pm]}$. Parasymmetric meters locate edge Goldstones (QII), pseudo-edge modes (QIV), and certify healing (QIII). Typed helicity meters reveal chiral transport asymmetries

without altering spin labels; quadrant assignment remains SynTh-determined. Background-field and cohomological safety as in [114, 442].

5) Fractons and subsystem-symmetric phases (anisotropic IR knobs). Work with $\mathcal{A} = \mathfrak{P}$ and an *anisotropic* knob $r = \min(\vec{\ell})$. Define $\mathbf{W}^{[\text{subsys}]}$ from line/plane-conserved currents as the law witness. Parasymmetry separates persistent law deficits (QIV) from power-law healing (QIII) along selected axes; typed meters with $\Pi^{[\text{planon/lineon}]}$ localize restricted-mobility species. Robustness again follows [117, 243].

6) Higher-form and non-invertible symmetries (Wilson/'t Hooft sectors). Let $\mathbf{W}^{[\mathscr{P}_p]}$ act on p-form sectors and transport by BRST/Hodge duality to scalar/vector blocks. Typed loop/surface projectors yield "perimeter vs. area" diagnostics; parasymmetry classifies healing (QIII) vs. persistent law deficit (QIV) and maps *where* (bulk vs. boundary) loop-type excitations are realized. Cohomological control as in [442, 117].

7) Floquet/time-crystalline and driven phases (temporal anchor). Extend the IR knob to a time-window; define $\mathbf{W}^{[\text{Floquet}]}$ with frequency-block filters. Trajectories $t \mapsto \upsilon^{(S)}_{\text{Synthesis}}(t; r)$ identify discrete-time SSB (QII-like in time), explicit drive locking (QIV), and healing to symmetric dynamics (QIII), including onset times $t^\star$. Spectral gauge safety as in [243].

8) Open quantum systems / Lindbladian symmetries. With $\mathbf{W}^{[\text{open}]}$ capturing conserved quantities modulo dissipators, parasymmetry separates environment-induced explicit breaking (QIV-like persistence) from prethermal/healing regimes (QIII). Typed species meters (Riesz) still operate on narrow bands of $\mathcal{D}_{\text{phys}}$ provided isolated structures exist; algebraic consistency follows [117, 442].

9) Emergent symmetry at criticality (QIII diagnostics). Near quantum/thermal critical points, set $r \sim \xi$ and deploy $\mathbf{W}^{[\text{crit}]}$. Power-law healing of the law deficit and reconvergence of spin blocks to the symmetric census (massless $s = 1, 2$, gapped radials) are quantified by the decay of parasymmetric cross-terms—an operational test for emergent Lorentz/conformal behaviour consistent with the spectral reconvergence logic of Section 7.10 and [91].

10) Cosmological/gravitation-adjacent arenas (solid-like backgrounds). For backgrounds with broken spacetime symmetries, use $\mathbf{W}^{[\text{metric}]}$ with Π_{TT} (spin-2) and phonon filters. Parasymmetry distinguishes phonons (QII) from pseudo-phonons (QIV) and tracks cone-preserving onset scales; typed spin-2 meters verify the persistence of massless TT carriers where spacetime symmetry is unbroken. TT-filtered spectral safety follows [114, 243].

Practical note (typed meters). In each arena, *typed* witnesses

$$\mathbf{W}^{[\mathscr{S};\chi]} = \Pi^{[\chi]}\,\mathbf{W}^{[\mathscr{S}]}\,\Pi^{[\chi]}, \qquad \upsilon^{(\cdot);[\chi]}_{\text{Synthesis}} = \text{Tr}\Big(P^{(\cdot)}_{\text{para}}\,\rho^{[\chi]}\Big),$$

with BRST-closed filters $\Pi^{[\chi]}$ for *spin*, *helicity*, *polarization*, or *species* (via Riesz projectors), allow parasymmetry to report, for the chosen χ: *type* (QI/QII/QIII/QIV via SynTh's

rule), *where* (bulk/boundary, sector, spatial window P_x), *when* (onset $r^\star$, and $t^\star$ if dynamics are supplied), and *what* (spin/helicity/polarization/species realized)—all remaining gauge/Nielsen–safe on $\mathcal{H}_{\text{phys}}$ and stable on $\mathfrak{P}$ [117, 442, 243].

8.6.8 Parasymmetric Readout of Second Sound

We now apply the parasymmetric layer to the longitudinal hydrodynamic block of Section 8.1.6, retaining the same realized weave configuration $(P_x(\lambda_{\text{UV}}), K_{\text{IR}}^{(\ell)}; \mathsf{P}_{\text{phys}})$ on $\mathfrak{P}$ and working on $\mathcal{H}_{\text{phys}} := H^0(\hat{Q}_{\text{BRST}})$. No new definitions are introduced: we only *type* the existing hydrodynamic witness using Section 8.6.5 and read it with the meters of Section 8.2.1.

Typed witness and species (second sound). Let $\mathbf{W}^{[\text{hydro}]}$ be the hydrodynamic sector witness (Definition 8.21) and let $\Pi^{[L]}$ denote the longitudinal projector (within the hydrodynamic block). Define the typed witness and typed state by

$$\mathbf{W}^{[\text{hydro};L]} = \Pi^{[L]}\,\mathbf{W}^{[\text{hydro}]}\,\Pi^{[L]}, \qquad \rho^{[L]} = \frac{\Pi^{[L]}\,\rho\,\Pi^{[L]}}{\mathrm{Tr}(\Pi^{[L]}\rho)} \quad (\text{on } \mathcal{H}_{\text{phys}}),$$

and the corresponding typed visibilities $\Lambda_{\text{law}}^{[\text{hydro};L]}$, $\Lambda_{\text{state}}^{[\text{hydro};L]}$ as in Definition 8.140. For the *second–sound species*, let $\mathcal{P}_{2\text{nd}}$ be the Riesz projector encircling the second–sound pole of the longitudinal block (cf. $\mathcal{P}_{\mathsf{m}}$ in Section 8.6.5); set $\Pi^{[\chi=2\text{nd}]} := \mathcal{P}_{2\text{nd}}$ to obtain a species–typed witness $\mathbf{W}^{[\text{hydro};2\text{nd}]}$ and the species–typed meters $v_{\text{Synthesis}}^{(a);\,[2\text{nd}]}$.

Meters (longitudinal and species–typed). For $a \in \{L, S\}$,

$$v_{\text{Synthesis}}^{(a);\,[L]} = \mathrm{Tr}\Big(P_{\text{para}}^{(a)}\,\rho^{[L]}\Big), \qquad v_{\text{Synthesis}}^{(a);\,[2\text{nd}]} = \mathrm{Tr}\Big(P_{\text{para}}^{(a)}\,\rho^{[2\text{nd}]}\Big),$$

with the relational meter $v_{\text{Synthesis}}^{(LS);\,[\cdot]}$ defined analogously (Definition 8.140). Gauge/Nielsen safety and plateau analyticity carry over from Proposition 8.141.

Proposition 8.145 (Quadrant diagnostics for second sound (typed parasymmetry))**.** *With the longitudinal hydrodynamic block of Section 8.1.6 and typed meters as above, the following implications hold:*

(i) QI (symmetric).
$\Lambda_{\text{law}}^{[\text{hydro};L]} \approx 1$, $\Lambda_{\text{state}}^{[\text{hydro};L]} \approx 1$*; the longitudinal spectrum shows pure diffusion (no longitudinal NG/pNG). Then* $v_{\text{Synthesis}}^{(a);\,[L]} \approx \frac{1}{2}$ *and the species–typed meter satisfies* $v_{\text{Synthesis}}^{(a);\,[2\text{nd}]} \approx \frac{1}{2}$ *(baseline), consistent with the absence of a second–sound pole.*

(ii) QII (state–SSB).
Two propagating longitudinal branches exist (Definition 8.26); the second–sound species meter *elevates at finite window,*

$$v_{\text{Synthesis}}^{(S);\,[2\text{nd}]}(r) > \tfrac{1}{2}, \qquad v_{\text{Synthesis}}^{(LS);\,[2\text{nd}]}(r) \text{ elevated},$$

and both decay to $\frac{1}{2}$ *under IR/volume dilation* $r \uparrow$*, while* $\Lambda_{\text{law}}^{[\text{hydro};L]} \simeq 1$*. This is the* sound–NG *signature in the longitudinal sector, now resolved at the species level.*

(iii) QIII (healing).
Any putative second–sound signature heals *in the strict IR:*

$$v_{\mathrm{Synthesis}}^{(a);\,[2\mathrm{nd}]}(r) \xrightarrow{r\uparrow} \tfrac{1}{2}, \qquad \Lambda_{\mathrm{law}}^{[\mathrm{hydro};L]}(r) \xrightarrow{r\uparrow} 1, \qquad \textit{and the pole reconverges to diffusion,}$$

in accordance with eq. (7.746) *and Proposition 8.23.*

(iv) QIV (explicit).
The longitudinal law deficit is persistent ($\Lambda_{\mathrm{law}}^{[\mathrm{hydro};L]} \le 1-\delta_\star$)*; the second–sound pole becomes a* pseudo–sound *with a small gap* $m_{\mathrm{ph}}^2 = \mathcal{O}(\epsilon)$ *(Section 8.1.6). The species–typed meters display* non–healing *behaviour:*

$$v_{\mathrm{Synthesis}}^{(LS);\,[2\mathrm{nd}]}(r) \textit{ non–healing}, \qquad v_{\mathrm{Synthesis}}^{(S);\,[2\mathrm{nd}]} \approx \tfrac{1}{2} \textit{ (unique state, no cats)},$$

while the gap satisfies the plateau GMOR/pseudo–phonon scaling inherited from the explicit source (cf. eq. (7.747)*).*

Definition 8.146 (Locator and onset for second sound)**.** Define the *second–sound locus* and *onset scale* at window r (bulk $r=\ell$, boundary $r=L$):

$$\mathcal{L}_{2\mathrm{nd}} := \Big\{(x,r):\ v_{\mathrm{Synthesis}}^{(S);\,[2\mathrm{nd}]}(x;r) > \tfrac{1}{2},\ \text{and } \omega_{\pm}^{(2)}(k) = \pm c_2 k - i\gamma_2 k^2 + o(k^2)\Big\},$$

$$r_{2\mathrm{nd}}^{\star} := \inf\Big\{r:\ v_{\mathrm{Synthesis}}^{(S);\,[2\mathrm{nd}]}(r) > \tfrac{1}{2}\ \wedge\ v_{\mathrm{Synthesis}}^{(LS);\,[2\mathrm{nd}]}(r) \downarrow \tfrac{1}{2}\Big\},$$

with the obvious QIV/QIII variants (non–healing or healing criteria, respectively).

Boundary variant. On $\mathfrak{P}_\partial$, replace $\Pi^{[L]}$ by the tangential projector $\Pi^{[\|]}$ and read the *edge second–sound* species via $\mathcal{P}_{2\mathrm{nd},\partial}$. Quadrant diagnostics follow Proposition 8.145 verbatim (QII: gapless edge sound; QIV: pseudo–edge–sound with $\mathcal{O}(\epsilon)$ gap; QIII: healing to tangential diffusion; QI: baseline).

Operational takeaway. Applying parasymmetry to second sound *does not change its spin/polarization label* (longitudinal, spin–0). It equips the hydrodynamic block with *typed, gauge–safe meters* that: (i) certify the presence of a genuine second–sound pole (vs. diffusion), (ii) *localize* where it occurs (bulk/boundary, position, window), and (iii) determine *when* it turns on (onset scale), while assigning the *type of breaking* (QII/QIII/QIV vs. QI) in the sense of Sections 7.10, 8.2.1 and 8.6.5.

Discussion and conclusion (second sound). The matrix witness of Section 8.1.6 provides an *unambiguous spectral predicate* for second sound: the longitudinal hydrodynamic block carries two propagating roots $\omega_{\pm}^{(2)}(k) = \pm c_2 k - i\gamma_2 k^2 + \cdots$ as stated in Definition 8.26. The parameters (c_2, γ_2) are fixed kinematically and dissipatively by the generalised stiffness and dissipation matrices (Lemma 8.27), and reduce in the two–fluid limit to Landau's classic formula for second sound (Remark 8.30; [450, 451, 452]). This furnishes a *yes/no* certificate on $\mathcal{H}_{\mathrm{phys}}$, independent of model-specific fits and stable under band/BRST reduction.

Layering parasymmetry on this block equips the construction with *typed, species-level diagnostics*. A Riesz projector $\mathcal{P}_{\mathrm{2nd}}$ onto the second–sound pole produces species-typed meters $\upsilon^{(a);\,[\mathrm{2nd}]}_{\mathrm{Synthesis}}$ and $\upsilon^{(LS);\,[\mathrm{2nd}]}_{\mathrm{Synthesis}}$ that *localize* second sound across space and scale, and define an onset $r^{\star}_{\mathrm{2nd}}$ (Section 8.6.8 and Definition 8.146). Gauge/Nielsen safety of these readings follows from the typed–meter invariance and analyticity on certified plateaux (Proposition 8.141; [117, 243]), so observations become *maps* $\mathcal{L}_{\mathrm{2nd}}$ rather than global assertions.

Crucially, the same meters render the phenomenon *quadrant-resolved*. In QII (state–SSB) the species meter is elevated at finite window and decays to 1/2 under IR/volume dilation, certifying a *true NG* sound branch; in QIV (explicit breaking) the branch is a *pseudo–sound* with an $\mathcal{O}(\epsilon)$ gap obeying the GMOR/source bridge (*eq.* (7.747)) and the meter is non–healing; in QIII (healing) any would–be second–sound signature reconverges to diffusion in the strict IR per *eq.* (7.746); and QI remains at baseline (Proposition 8.145). Thus the traditional "first/second sound vs. diffusion" trichotomy is elevated to a *mechanistic* classification that says *why* the branch appears.

The boundary/tangential variant is immediate: replacing $\Pi^{[L]}$ by the tangential projector $\Pi^{[\|]}$ realizes *edge second sound* on $\mathfrak{P}_{\partial}$ with identical quadrant logic and locator/onset definitions (Section 8.6.8).

Finally, the outputs are *falsifiable and robust*: certified (c_2, γ_2) with errors tied to window resolution; a binary species meter (elevated vs. baseline) with healing/persistence trends; a gap–scaling law in QIV; and onset scales $r^{\star}_{\mathrm{2nd}}$—each directly checkable in simulation or experiment and regulator–stable on $\mathfrak{P}$ [117, 243]. Scope and limits are as usual: the analysis holds on certified plateaux, and species–level meters presume an isolated or sufficiently narrow pole; if spectra are provided in a fixed gauge, the BRST projection must be certified prior to applying the resolver [117, 243]. In sum, for second sound, SynTh + typed parasymmetry yields a *gauge-safe, quadrant-aware, mappable instrument* that states precisely *if*, *why*, *where*, and *when* second sound occurs, while reproducing the standard two–fluid limits where appropriate [450, 451, 452].

Chapter 9

The Quanta of Fields

Purpose and scope. The present introduction positions the operator–theoretic construction of fields and quanta on the emergent arena against the established BRST–physical framework and the Σ-aware symmetry diagnostics. The upstream axioms and identities of Chapters 7–8 remain unchanged—BRST charge and cohomology, gauge fixing as ζ-exact, cohomological Euler–Lagrange operators, Σ-compatible linear response, and Ward/Slavnov identities—while the upstream feed is sharpened: the pre-physical, operator-valued chronogeometry $\hat{\Sigma}_{\mu\nu} := \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu}$ is taken as the canonical input that, after Planck-plateau upgrade and BRST projection, yields the expectation tensor $\bar{\Sigma}_{\mu\nu}$ used on the physical side (cf. §6.10 and the relational constructions surrounding $\nabla^{(\bar{\Sigma})}$, Definition 6.7). The goal is a unified and band-regular route from correlator kinematics to physical spectra, without altering the cohomological infrastructure of Chapters 7–8.

Law–state analysis at the pre-physical level. On the emergent/GNS arena, the law–state dichotomy is evaluated at band level using Σ-based, graded-covariant diagnostics. A band tag $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ is implemented via a UV projector $P_x(\lambda_{\mathrm{UV}})$ and a causal IR kernel $K_{\mathrm{IR}}^{(\ell)}$ chosen to be BRST-closed, ensuring commutation with $\hat{Q}_{\mathrm{BRST}}$ and compatibility with the physical projector $\mathsf{P}_{\mathrm{phys}}$ (§7.5, (7.143), (7.24)). With the Σ-probe $\hat{Q}_x[v] = v^\mu v^\nu \hat{\Sigma}_{\mu\nu}(x)$, law and state contrasts $\Delta_{\mathrm{law}}^{\mathrm{pre},\Sigma}$ and $\Delta_{\mathrm{state}}^{\mathrm{pre},\Sigma}$ distinguish explicit symmetry breaking of the operator law from spontaneous symmetry breaking of the Holo-State at fixed tolerance; the quadrant structure (QI–QIV) then classifies regimes and governs subsequent realization on the BRST-physical arena. In particular, QII corresponds to state-asymmetric, law-symmetric configurations that will underwrite NG/pNG content after projection, whereas QIII/QIV capture genuine (explicit) departures of the law channel, to be filtered by anomaly and BRST viability checks.

Unified chronogeometry and graded transport. The chronogeometric tensor $\hat{\Sigma}_{\mu\nu} = \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu}$ consolidates the triadic metric and temporal sector into a single operator-valued object whose coincident Hessians generate the kinematic forms that will later define $\bar{\Sigma}_{\mu\nu}$. This unification preserves gauge/frame covariance and graded selection rules while remaining finite and causal on the band (self-regularizing distances/times and their Hessians). The Fourfold Inter-Modal Correlator $\hat{\mathbf{G}}_\Lambda$ supplies the transport-covariant kernel data with $\mathbb{Z}_2 \times \mathbb{Z}_2$ grading over the modal set $\{0, 1, p, s\}$, and the relational differential structure is read through $\nabla^{(\bar{\Sigma})}$

once expectations are taken (cf. Definitions 6.4 and 6.7). The generalized field doublet $\hat{\mathbf{F}} = (\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}})^{\mathsf{T}}$ sits on the same operator bundle and inherits these covariances, enabling a single Σ-compatible account of geometry-like (Gaiamiti) and matter-like (Tantrakyma) sectors.

Broken directions, mode extraction, and Σ-weighted normalization. Broken symmetry directions are identified pre-physically by the graded tangent $\delta_X \hat{\mathbf{G}}_\Lambda = [X, \hat{\mathbf{G}}_\Lambda]_{\mathrm{gr}}$ for $X \in \mathfrak{g}$, evaluated blockwise and weighted by $\hat{\Sigma}$. The resulting Σ-Gram problems produce principal directions in the geometry block ($\Phi\Phi$) and the matter block ($\Theta\Theta$), defining Apeiron and Kymon creation operators with Σ-orthonormality. The phase-tangential versus amplitude-radial decomposition within the Σ-geometry separates massless versus gapped geometric excitations at the pre-physical level, while maintaining graded covariance and band regularity. These Σ-normalized mode spaces provide the inputs for linear response and spectral diagnostics, but no change is made to the physical particle criteria, which remain cohomological and pole/residue based in $\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}})$ (cf. Section 6.10).

Plateau certification, BRST projection, and Σ-causal response. A certified plateau in the $(\lambda_{\mathrm{UV}}, \ell)$ plane—verifying Σ-regularity, causal/hyperbolic cones, BRST nilpotency and commutation with the band map, together with Nielsen-type gauge-parameter independence—licenses the realization map that sends pre-physical operators to the BRST-physical arena. The expectation $\bar{\Sigma}_{\mu\nu} = \langle \mathsf{P}_{\mathrm{phys}} \lim_{\mathfrak{P}} \mathfrak{B}_{\lambda,\ell}[\hat{\Sigma}_{\mu\nu}] \mathsf{P}_{\mathrm{phys}} \rangle$ then serves as the background tensor for the Relational Levi–Civita connection, for Σ-causal retarded propagators, and for the cohomological Euler–Lagrange operators used in linear response (Definition 6.7). Ward/Slavnov identities, anomaly filters, and Kugo–Ojima unitarity checks are applied exactly as in Chapter 7, now with Σ-compatible currents and Green functions. The BRST-exact sector remains unphysical, and only BRST-closed, Nielsen-stable poles with positive residues are admitted as realized quanta on $\mathcal{H}_{\mathrm{phys}}$ (with the Quadrant-I spectrum at the symmetric baseline and its deformations interpreted through the Σ-fed diagnostics of Chapter 8, (7.143), (7.24)).

Organization of the construction. With these preliminaries, the subsequent sections proceed from the arena, transport, grading, and band tagging to Σ-based law/state order parameters, to Σ-weighted mode extraction in geometry and matter blocks, and finally to plateau-based realization and BRST-physical spectra. Throughout, the generalized field operator, the unified inter-modal correlator, and the operator-bundle connection are treated in their established form (Ch. 7), while the band-limited BRST checklist, acceptance rules, and quadrant logic (Ch. 8) are invoked without modification, the sole refinement being that all inputs are funneled through the chronogeometric tensor $\hat{\Sigma}_{\mu\nu}$ and its expectation $\bar{\Sigma}_{\mu\nu}$.

Overview

The Primordial Quantum Singularity (PQS) and the Quantum Holo–State. All pre–physical diagnostics are anchored at *Primordial Quantum Singularities* (PQS) as defined in

Definition 1.25, with the formal identity $\mathfrak{B}_{\vec{r}} = \hat{\mathbf{\Omega}}_{\vec{r}} = \overset{\circ}{\mathbf{\Lambda}}_{\hat{\Phi}_{\vec{r}},\hat{\Theta}_{\vec{r}}}$ (cf. (1.3)). Each PQS is intrinsically characterized by the *Quantum Holo–State* $|\overset{\circ}{\mathbf{\Psi}}\rangle$ (Definition 1.27), whose GNS realization generates the emergent arena $\mathcal{H}_{\rm em} \equiv \mathcal{H}_{\overset{\circ}{\mathbf{\Psi}}}$ used throughout (Definition 9.87). As detailed in Definitions 1.27, 6.4, 9.20 and 9.60 and eqs. (4.85) and (9.27), the causal/metric pipeline driven by the Holo–State is

$$\begin{aligned} &|\overset{\circ}{\mathbf{\Psi}}\rangle \quad (Definition\ 1.27)\\ &\rightsquigarrow\ \hat{G}_\Lambda \quad (Definition\ 6.4)\\ &\rightsquigarrow\ \langle\cdot,\cdot\rangle_x \quad (eq.\ (4.85))\\ &\rightsquigarrow\ \hat{\Sigma}_{\mu\nu}(x) \quad (Section\ 9.9)\\ &\rightsquigarrow\ \bar{\Sigma}_{\mu\nu}(x) \quad (Definition\ 9.60)\end{aligned}$$

so the fibre inner products and the operator chronogeometry are *induced* from Holo–State expectations. Moreover, Theorem 4.314 establishes that $|\overset{\circ}{\mathbf{\Psi}}\rangle$ is *globally entangled* and non–factorizable across nontrivial splits—a structural fact that remains intact in what follows.

What does it mean, here, for symmetry to be "broken"? "Broken" is precise and occurs *at the band–tagged, pre–physical level*. What can break is the G–invariance of either the *law* (band–projected, graded–covariant kernels/constraints on $\mathcal{H}_{\rm em}$) or the *state* (the Holo–State), tested by the canonical Σ–based order parameters built from the operator chronogeometry and its quadratic probe $\hat{Q}_x[v] = v^\mu v^\nu \hat{\Sigma}_{\mu\nu}(x)$ (see Section 9.9). Both channels are evaluated with the band tag $\mathfrak{B}_{\lambda_{\rm UV},\ell}[\cdot]$ and PSR functionals $\mathcal{I}[\cdot]$, and summarized by the pre–physical contrasts and their typed visibilities $\left(\Delta_\bullet^{\mathrm{pre},\Sigma},\, \Lambda_\bullet^{\Sigma;[\chi]}\right)$ (cf. Sections 9.11 and 9.12). *State breaking* here means *loss of G–invariance* of Holo–State expectations on–band; it does *not* contradict the global non–factorizability of Theorem 4.314.

What does not break: the role of the Fourfold correlator. The Fourfold inter–modal correlator $\hat{G}_\Lambda$ is an *operator object* that transforms *covariantly* under G (cf. Definition 6.4):

$$\hat{G}_\Lambda \ \longmapsto\ \mathrm{Ad}_g\, \hat{G}_\Lambda := U(g)\, \hat{G}_\Lambda\, U(g)^{-1}, \qquad g \in G.$$

At this object level, $\hat{G}_\Lambda$ *only transforms*; nothing "breaks." What the formalism calls *breaking* is detected downstream—either in the band–projected *law channel* functorially built from $\hat{G}_\Lambda$ (explicit breaking) or in the *Holo–State expectations* of G–covariant functionals of $\hat{G}_\Lambda$ (spontaneous breaking). Concretely, after band–tagging $\mathfrak{B}_{\lambda_{\rm UV},\ell}$,

$$\Delta_{\rm law}^{\mathrm{pre},\Sigma}(g;x;v) > \varepsilon \iff \mathfrak{L}\Big[\,\mathrm{Ad}_g\ \mathfrak{B}_{\lambda,\ell}(\hat{G}_\Lambda)\Big] \neq \mathfrak{L}\Big[\mathfrak{B}_{\lambda,\ell}(\hat{G}_\Lambda)\Big],$$

where $\mathfrak{L}[\cdot]$ denotes the Σ–compatible operator–law kernels/constraints (cf. Section 9.9 and Definition 9.45). By contrast, with the law invariant,

$$\Delta_{\rm law}^{\mathrm{pre},\Sigma}(g;x;v) \approx 0 \quad \text{and} \quad \Delta_{\rm state}^{\mathrm{pre},\Sigma}(g;x;v) > \varepsilon,$$

one has *spontaneous state* breaking (equivalently $H_{\rm law}(x) = G$ but $H_{\rm state}(x) \subsetneq G$; cf. Section 9.11). Thus the correlator is the *substrate*; it is the symmetry of the *law* or of the *state*—as revealed by the Σ–based contrasts—that fails.

What exactly is broken? *Explicit law breaking* means the intended symmetry G fails to leave the band–limited operator law invariant, detected as

$$\Delta_{\mathrm{law}}^{\mathrm{pre},\Sigma}(g;x;v) \;>\; \varepsilon \quad (g\in G,\; v\in T_x\mathcal{X}_q),$$

equivalently, a nonzero Σ–probe difference under the graded adjoint action. *Spontaneous state breaking* preserves the law channel but not the Holo–State:

$$\Delta_{\mathrm{law}}^{\mathrm{pre},\Sigma}(g;x;v) \;<\; \varepsilon \quad \text{while} \quad \Delta_{\mathrm{state}}^{\mathrm{pre},\Sigma}(g;x;v) \;>\; \varepsilon,$$

so that

$$H_{\mathrm{law}}(x) := \{g\in G : \Delta_{\mathrm{law}}^{\mathrm{pre},\Sigma}(g;x;v) < \varepsilon\ \forall v\}, \qquad H_{\mathrm{state}}(x) := \{g\in G : \Delta_{\mathrm{state}}^{\mathrm{pre},\Sigma}(g;x;v) < \varepsilon\ \forall v\},$$

and pre–physical SSB means $H_{\mathrm{law}}(x) = G$ but $H_{\mathrm{state}}(x) \subsetneq G$. The quadrant predicate (QI–QIV) encodes the four possibilities—nothing; state–only; law–only; both—driven by $\Delta_{\bullet}^{\mathrm{pre},\Sigma}$ or their typed versions $\Delta_{\bullet}^{\mathrm{pre},\Sigma;[\chi]}$ with visibilities $\Lambda_{\bullet}^{\Sigma;[\chi]}$ (see Section 9.11 and the orientation in Section 9.15.1).

Where do broken directions come from? A generator $X\in\mathfrak{g}$ is diagnosed as broken at x when its graded tangent

$$\delta_X\hat{G} \;:=\; [X,\hat{G}]_{\mathrm{gr}}$$

carries nonzero Σ–weight in a relevant block,

$$\mathcal{I}\Big[(\delta_X\hat{G}_{\bullet\bullet})^{\dagger}\,\hat{\Sigma}^{(\bullet)}\,(\delta_X\hat{G}_{\bullet\bullet})\Big] \;>\; 0, \qquad \bullet\bullet\in\{\Phi\Phi,\Theta\Theta\}.$$

These Σ–weighted variation maps seed the *block–wise* generalized eigenproblems that extract the principal pre–physical mode families (a typed, band–regular upgrade of SVD), yielding Apeiron (geometry, $\Phi\Phi$) and Kymon (matter, $\Theta\Theta$) mode spaces with Σ–orthonormality and factorizations. Their outputs feed the typed Type–Location–Onset assignments $\mathrm{TLO}^{[\chi]}(x;r)$ (cf. Sections 8.6, 9.8, 9.10 and 9.12).

What this does—and does not—claim. The QI–QIV classification and the Apeiron/Kymon construction are *pre–physical inputs*; they are not particle claims. Physical poles/residues are read only after *plateau certification* and Σ–aware *realization* with BRST projection to $\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}})$. On certified plateaus the law channel is Nielsen–flat; BRST hygiene of the band tag guarantees our diagnostics descend functorially and remove gauge–parameter dependence before testing retarded kernels for stable, positive–residue poles (see Section 9.14 and Lemma 9.54). Only BRST–closed, Nielsen–flat features survive; BRST–exact "breakings" are projected out.

Spirit near the primordial regime (PQS). Here "singularity" means the *Primordial Quantum Singularity (PQS)* (Definition 1.25). Diagnostics are performed on finite bands $(\lambda_{\mathrm{UV}},\ell)$ in neighborhoods of the PQS on $\mathcal{H}_{\mathrm{em}}$ and continued only along certified plateaus (cf. Section 9.14).

In this regime the band map is G–covariant and BRST–closed, so the law channel remains symmetric while the Holo–State may not. Accordingly, the natural baseline is QII,

$$\Delta_{\text{law}}^{\text{pre},\Sigma} \approx 0 \quad \text{and} \quad \Delta_{\text{state}}^{\text{pre},\Sigma} > \varepsilon,$$

i.e. $H_{\text{law}}(x) = G$ but $H_{\text{state}}(x) \subsetneq G$. The broken–generator tangents with nonzero Σ–weight then seed the Apeiron/Kymon *pre–physical* mode spaces via the Σ–weighted extraction (see Section 8.6), with physical poles read only after plateau certification and BRST projection.

PQS–anchored scale principle. The pre-physical regime lies strictly between the Primordial Quantum Singularity (PQS) coherence scale and the Planck scale, and the first stage of physics ("realization") begins at the Planck scale:

$$\Big[\underbrace{\text{PQS core}}_{\text{ontic origin}} \rightsquigarrow \underbrace{\text{pre-physical band}}_{\ell_{\text{PQS}} \le \ell < \ell_P} \rightsquigarrow \underbrace{\text{stage of realization (first physics)}}_{\ell \ge \ell_P}$$
$$\rightsquigarrow \underbrace{\text{IR / classical}}_{\ell \gg \ell_P} \rightsquigarrow \underbrace{\text{cosmological}}_{\ell \sim \mathcal{L}_U}\Big]$$

This is consistent with the Chapter 6 genesis chain, where causality emanates from the PQS apex and classical chronogeometry arises only after expectation/realization steps (Section 5.4.6; cf. Table table 5.3).

Definition 9.1 (Regimes and thresholds)**.** Let ℓ_{PQS} denote the PQS coherence/patch scale (Chapter 6, Section 5.4.6) and ℓ_P the Planck length. For an IR-window parameter ℓ (from the band tag $K_{\text{IR}}^{(\ell)}$), define:

1. *PQS core*: $\ell \approx \ell_{\text{PQS}}$, with no resolvable internal distances/times (operator distances/-times are trivialized at the apex; cf. Table table 5.3).
2. *Pre-physical band*: $\ell_{\text{PQS}} \le \ell < \ell_P$.
3. *Stage of realization (first physics)*: $\ell \ge \ell_P$.

Proposition 9.2 (Operational eligibility by regime)**.** *With the operator chronogeometry $\hat{\Sigma}_{\mu\nu} := \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu}$ (Chapter 5: $\hat{g}_{\mu\nu}$ from Definition 5.1; temporal curvature $\hat{t}_{\mu\nu}$ in the Level 2 quantum–geometric layer of Table table 5.3; cf. the unified Σ-notation in Definitions 9.20 and 9.22) and its expectation $\bar{\Sigma}_{\mu\nu} := \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (Chapter 5: Level 3 classical–geometric layer in Table table 5.3; cf. the Σ–realization map Definition 9.80 and eq. (9.35)), the admissible operations are:*

(i) PQS core
No meters or certification procedures apply; band-tagging and parasymmetry are inapplicable at the apex (Section 5.4.6).

(ii) Pre-physical band $\left[\ell_{\text{PQS}}, \ell_P\right)$
Operator-level audits only with band-tagging permiitted

$$(P_x(\lambda_{\text{UV}}), K_{\text{IR}}^{(\ell)})$$

Σ-*cone/hyperbolicity checks, BRST/Nielsen hygiene, and typed permit ledgers.* Prohibited*: plateau certification* $\mathfrak{P}$*, evaluation of holosymmetric visibilities as physical numbers, and projection to* $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$.

(iii) Realization $\ell \geq \ell_P$

First certified plateaux $\mathfrak{P}$ *may be established (tolerances as in eq. (7.28)); thereafter project to* $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$. *Physical, channelized visibilities* $(\Lambda_{\text{law}}, \Lambda_{\text{state}})$ *are evaluated; SynTh and parasymmetric meters act only on* H^0*, remaining* Σ*-causal and BRST-safe.*

Lemma 9.3 (No realization below the Planck scale)**.** *If* $\ell < \ell_P$*, then (i) plateau certification* $\mathfrak{P}$ *is not admissible, and (ii) the realization map to* $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$ *is not invoked. Consequently, all diagnostics in* $\left[\ell_{\text{PQS}}, \ell_P\right)$ *are pre-physical/operator-level and expectation-free, in agreement with Chapter 6.*

Remark 9.4 (Sector readings across realization)**.** At $\ell \geq \ell_P$, sector readings become physical: in the gaiamiti ($\Phi\Phi$) block, the massless spin-2 graviton (Goldstone) and gapped geometric radials are read as poles/residues on H^0; in the tantrakyma ($\Theta\Theta$) block, kymon content realizes with NG/pNG/Higgs logic resolved quadrantly (QI–QIV). For $\ell \gg \ell_P$ and $\ell \sim \mathcal{L}_U$, the same H^0 kernels govern, respectively, IR/classical and cosmological coarse-grained limits, consistent with Chapter 6.

9.1 Arena, Transport, Grading, and Band Tagging

Definition 9.5 (Emergent arena and Fourfold Inter-Modal Correlator)**.** Let $\mathcal{H}_{\text{em}}$ denote the emergent (GNS) arena. The *Fourfold Inter-Modal Correlator* is the operator-valued tensor

$$\hat{G}^{\alpha\beta}_{\Lambda}(x, x') \;:=\; \hat{\mathcal{F}}^{\dagger}_{\alpha}(x)\,\mathcal{U}(x \leftarrow x';\gamma)\,\hat{\mathcal{F}}_{\beta}(x') \;\in\; \mathcal{B}(\mathcal{H}_{\text{em}}), \qquad \alpha, \beta \in \{0, 1, p, s\},$$

as first defined in Section 4.1.5 (see eq. (4.22)). The coincident limit generates fibre inner products and organizes the four dialectical modes.

Proposition 9.6 (Hermiticity and gauge covariance of the correlator)**.** *With* $\mathcal{U}$ *the gauge–covariant transporter (cf. Definitions 6.184 and 6.192), and with the Fourfold Inter–Modal Correlator defined together with its block structure in Definition 4.9 and eq. (4.24), the Twofold block enjoys Hermitian symmetry and the full Fourfold transforms covariantly:*

$$(\hat{G}^{\alpha\beta}(x, x'))^{\dagger} = \hat{G}^{\beta\alpha}(x', x), \qquad \hat{\mathbf{G}}_{\Lambda}(x, x') \mapsto g(x)\,\hat{\mathbf{G}}_{\Lambda}(x, x')\,g(x')^{\dagger},$$

hence expectation values in a gauge–covariant state are gauge–invariant (cf. Section 4.19.1).

Definition 9.7 (Gauge-covariant transport and graded covariant derivative)**.** Let $\mathcal{A}_\mu$ be the operator–bundle connection entering the gauge–covariant transporter $\mathcal{U} = \mathcal{P}\exp\!\left(-\int_\gamma \mathcal{A}\right)$ (cf. Definitions 6.184 and 6.192). For channel components of the Fourfold Inter–Modal Correlator (cf. Definition 6.4 and eq. (4.24)), define the graded covariant derivative

$$\mathfrak{D}_\mu \hat{G}^{\alpha\beta} := \partial_\mu \hat{G}^{\alpha\beta} + [\,, \mathcal{A}_\mu, \hat{G}^{\alpha\beta},]_{\text{gr}}, \qquad (\alpha, \beta \in \{0, 1, p, s\}), \tag{9.1}$$

which transforms homogeneously under gauge changes (cf. Proposition 9.6) and preserves the $\mathbb{Z}_2 \times \mathbb{Z}_2$ degrees (cf. Definition 9.8 and the graded kinematics in Section 4.1.7).

Scale / Regime	**Apeiron & Graviton Manifestation (Gaiamiti Sector)**	**Ontological Level**	**Operational Layer (band/BRST/parasymmetry)**
$\approx \ell_{\mathbf{PQS}}$ **PQS patch (ontic origin)**	No resolvable internal relational structure; only potential excitations in the gaiamiti operator algebra; *no* SSB or particle claims.	Ontic origin; pre-geometric and pre-particle	Pre-physical: outside $H^0(\hat{Q}_{\mathrm{BRST}})$; band-tagging not applied;
$\ell_{\mathbf{PQS}} < \ell < \ell_P$ **Sub-Planckian (pre-physical band)**	Incipient geometric fluctuations $\delta\hat{\mathbf{\Phi}}(x)$; *no* physical spectra; any SSB talk is *diagnostic-only*, not realized.	Pre-physical relational layer	Operator-only: band-tagging allowed, Σ-cone/hyperbolicity audits, typed permit ledgers; *no* plateau certification, *no* projection to H^0.
$\sim \ell_P$ **Planckian (onset of realization)**	Spontaneous symmetry breaking of the geometric vacuum; massive Apeiron modes acquire intrinsic scale m_A; **massless Graviton emerges as the spin-2 Goldstone**. Physical spectra are first read here.	Emergent geometry and particle properties	*Stage of realization*: certify $\mathfrak{P}$, project to $H^0(\hat{Q}_{\mathrm{BRST}})$, use $G^{\mathrm{R}}_{\Phi\Phi,\mathrm{phys}}$ (and mixed blocks). Parasymmetry, if used, acts only on H^0.
$\gg \ell_P$ **Macroscopic (IR)**	Graviton mediates long-range classical GR; massive Apeiron modes are short-ranged.	Classical fields and stable particles	Same physical layer as Planckian: $G^{\mathrm{R}}_{\bullet,\mathrm{phys}}$ on H^0; classical limit from $\bar{\Sigma}_{\mu\nu}$; parasymmetry optional.
$\sim \mathcal{L}_U$ **Cosmological**	Collective/thermodynamic graviton effects (background; large-scale structure).	Thermodynamic/statistical limit	Coarse-grained aggregates of H^0-level kernels over cosmological domains; parasymmetry typically unnecessary.

Table 9.1: Hierarchy of emergence for quanta *(Gaiamiti → apeiron; Tantrakyma → kymon)*. The PQS patch is characterized by a *length* scale ℓ_{PQS} (PQS coherence size); use ℓ^{-1}_{PQS} only for wavenumber thresholds. **Pre-physical:** for $\ell_{\mathrm{PQS}} \leq \ell < \ell_P$, diagnostics are operator-level and expectation-free (no $\mathfrak{P}$, no projection to $H^0(\hat{Q}_{\mathrm{BRST}})$). **Realization:** at $\ell \geq \ell_P$, after plateau certification $\mathfrak{P}$ (cf. eq. (7.28)), observables are evaluated on the physical BRST cohomology $H^0(\hat{Q}_{\mathrm{BRST}})$; parasymmetric diagnostics (if used) act only on H^0.

Definition 9.8 ($\mathbb{Z}_2 \times \mathbb{Z}_2$ grading of channels)**.** Assign the channel colors $\deg(0) = (0,0)$, $\deg(1) = (1,0)$, $\deg(s) = (0,1)$, $\deg(p) = (1,1)$, and set $\deg(\hat{G}^{\alpha\beta}) = \deg(\alpha) + \deg(\beta)$. This equips the correlator components of Definition 6.4 with a minimal graded $*$–algebra structure that preserves mixing rules and sign/phase conventions under transport (cf. Definition 9.7 and Proposition 9.6); the modal basis and fibre structure are those of the Fourfold Dialectic Algebra (cf. Definition 1.5) and its operator implementation within the algebra of observables (cf. Definition 1.32).

Remark 9.9 (Modal fibres and Fourfold logic)**.** The parasymmetric and holosymmetric channels are not linear superpositions of the Twofold; they are independent basis rays in the modal fibre and are accessed by their own effects (cf. Definition 1.32). This reflects the Fourfold Dialectic Algebra (cf. Definition 1.5 and Section 1.2.5) and the emergent modal–Hilbert–space fibres with coincident–limit inner products (cf. Section 4.6); see also the generalized field doublet placed on this operator bundle (cf. Definition 6.1).

Definition 9.10 (Band tag (pre-physical window) and IR window)**.** Let $P_x(\lambda_{\mathrm{UV}})$ denote the UV spectral projector and $K_{\mathrm{IR}}^{(\ell)}$ a causal IR window kernel (mass 1), both chosen to be parallel–transport covariant (cf. Definitions 6.184 and 6.192 and Proposition 9.6) and BRST–closed on the band (cf. eq. (7.24) and Section 7.5). The *band tag* acts on any local observable $\mathfrak{O}$ by

$$\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\mathfrak{O}](x) \;:=\; \int_{\mathcal{X}_q} K_{\mathrm{IR}}^{(\ell)}(x,y)\,\mathcal{U}(x\leftarrow y)\,P_y(\lambda_{\mathrm{UV}})\,\mathfrak{O}(y)\,P_y(\lambda_{\mathrm{UV}})\,\mathcal{U}(y\leftarrow x)\,\mathrm{d}\mu_y, \tag{9.2}$$

where $\mathcal{U}$ is the gauge–covariant transporter (cf. Definitions 6.184 and 6.192) and $\mathrm{d}\mu_y$ is the integration measure on $\mathcal{X}_q$ (consistent with the relational volume conventions; cf. Definition 6.33). This definition implements the IR windowing and UV projection used in the band–limited formalism (cf. Definition 6.280 and eqs. (7.130) and (7.143)) and transforms homogeneously under gauge changes while preserving the $\mathbb{Z}_2 \times \mathbb{Z}_2$ degrees of the channel algebra (cf. Definition 9.8 and Section 4.1.7).

Lemma 9.11 (BRST-compatibility (tag hygiene))**.** *If* $[\hat{Q}_{\mathrm{BRST}}, P_x(\lambda_{\mathrm{UV}})]_\pm = [\hat{Q}_{\mathrm{BRST}}, K_{\mathrm{IR}}^{(\ell)}]_\pm = 0$*, then* $[\hat{Q}_{\mathrm{BRST}}, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}]_\pm = 0$*. Band-tagging preserves microcausality and the locality of commutators.*

Proof. By definition, the band tag acts on a local observable $\mathfrak{O}$ as (cf. Definitions 6.280 and 9.10 and eq. (9.2))

$$\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\mathfrak{O}](x) = \int_{\mathcal{X}_q} K_{\mathrm{IR}}^{(\ell)}(x,y)\,\mathcal{U}(x\leftarrow y)\,P_y(\lambda_{\mathrm{UV}})\,\mathfrak{O}(y)\,P_y(\lambda_{\mathrm{UV}})\,\mathcal{U}(y\leftarrow x)\,\mathrm{d}\mu_y,$$

where $\mathcal{U}$ is the gauge–covariant parallel transporter generated by the operator–bundle connection (cf. Definitions 6.184 and 6.192 and Proposition 9.6) and $\mathrm{d}\mu_y$ is the relational measure (cf. Definition 6.33).

Step 1 (graded derivation and BRST-covariant transport). Using that the BRST differential $\zeta(\cdot) = \{\hat{Q}_{\mathrm{BRST}}, \cdot\}$ is a graded derivation (cf. Section 6.10) and that the transporter is endpoint–covariant, one has the standard identity

$$[\hat{Q}_{\mathrm{BRST}}, \mathcal{U}(x\leftarrow y)\,X(y)\,\mathcal{U}(y\leftarrow x)]_\pm = \mathcal{U}(x\leftarrow y)\,[\hat{Q}_{\mathrm{BRST}}, X(y)]_\pm\,\mathcal{U}(y\leftarrow x),$$

for any ghost–number–zero local operator X built from the channel algebra, provided the endpoint insertions cancel by conjugation (a direct consequence of the endpoint transformation of $\mathcal{U}$ and its inverse; cf. Definition 6.192, Section 4.1.7, and Proposition 9.6).

Step 2 (commutation with the band ingredients). By hypothesis the UV projector and IR window are BRST–closed, $[\hat{Q}_{\mathrm{BRST}}, P_x(\lambda_{\mathrm{UV}})]_\pm = 0$ and $[\hat{Q}_{\mathrm{BRST}}, K_{\mathrm{IR}}^{(\ell)}]_\pm = 0$ (cf. eq. (7.24) and Section 7.5), and both have ghost number zero (cf. Definition 6.280). Using linearity and the graded Leibniz rule inside the integral,

$$\begin{aligned}[\hat{Q}_{\mathrm{BRST}}, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}][\mathfrak{O}](x) &= \int_{\mathcal{X}_q} K_{\mathrm{IR}}^{(\ell)}(x,y)\,\Big[\hat{Q}_{\mathrm{BRST}}, \mathcal{U}(x\leftarrow y)\,P_y\,\mathfrak{O}(y)\,P_y\,\mathcal{U}(y\leftarrow x)\Big]_\pm \,\mathrm{d}\mu_y \\ &= \int_{\mathcal{X}_q} K_{\mathrm{IR}}^{(\ell)}(x,y)\,\mathcal{U}(x\leftarrow y)\,P_y\,[\hat{Q}_{\mathrm{BRST}}, \mathfrak{O}(y)]_\pm\,P_y\,\mathcal{U}(y\leftarrow x)\,\mathrm{d}\mu_y \\ &= \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}\Big[[\hat{Q}_{\mathrm{BRST}}, \mathfrak{O}]_\pm\Big](x).\end{aligned}$$

Therefore $[\hat{Q}_{\mathrm{BRST}}, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}]_\pm = 0$ as superoperators on the local algebra. This is the band–level BRST–compatibility ("tag hygiene"), consistent with the checklist used in the band–limited formalism (cf. eqs. (7.130) and (7.143)).

Step 3 (microcausality and locality of commutators). Let $\mathfrak{O}_1, \mathfrak{O}_2$ be local observables supported in regions $\mathcal{O}_1, \mathcal{O}_2 \subset \mathcal{X}_q$. By construction the IR kernel $K_{\mathrm{IR}}^{(\ell)}(x,y)$ has causal support and decays outside the lightcone (cf. Definition 6.280), and the transporter is path–ordered along γ without enlarging causal support (cf. Definition 6.184 and Proposition 9.6). If $\mathcal{O}_1$ and $\mathcal{O}_2$ are spacelike separated, microcausality of the underlying local algebra (cf. Definition 1.32) implies $[\mathfrak{O}_1(y), \mathfrak{O}_2(y')]_\pm = 0$ for $y \in \mathcal{O}_1$, $y' \in \mathcal{O}_2$. Since $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ is built from causal smearing, parallel–transport conjugation, and spectral projection, none of which enlarge causal support or change graded locality, one obtains

$$\Big[\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\mathfrak{O}_1](x),\, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\mathfrak{O}_2](x')\Big]_\pm = 0 \quad \text{for spacelike } (x,x').$$

Hence band–tagging preserves microcausality and the locality of commutators.

Combining the three steps yields the claim. □

Definition 9.12 (Pre-physical observable and PSR)**.** Let $\mathcal{H}_{\mathrm{em}}$ be the emergent/GNS Hilbert space (cf. Definition 1.34 and Remark 1.36) and let $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ be the band tag defined by UV projection and IR windowing (cf. Definitions 6.280 and 9.10 and eqs. (7.130), (7.143) and (9.2)). For any local observable $\mathfrak{O}$, define the band–tagged *pre-physical observable*

$$\mathfrak{O}_{\mathrm{pre}}(x;\lambda_{\mathrm{UV}},\ell) := \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\mathfrak{O}](x).$$

Pre-physical symmetry reduction (PSR) occurs when the expectation on the emergent arena is nonvanishing *prior* to cohomological projection,

$$\Big\langle \mathfrak{O}_{\mathrm{pre}} \Big\rangle_{\mathcal{H}_{\mathrm{em}}} \neq 0,$$

where expectations are taken in the Holo-State/GNS sense (cf. Definitions 1.18 and 1.34). BRST–functoriality of the band tag is expressed as

$$[\,\hat{Q}_{\mathrm{BRST}}, \mathfrak{O}_{\mathrm{pre}}]_\pm \;=\; \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}\Big(\,[\,\hat{Q}_{\mathrm{BRST}}, \mathfrak{O}]_\pm\Big),$$

which follows from the band–level BRST–compatibility ("tag hygiene") once the UV projector and IR window are BRST–closed (cf. Lemma 9.11, eq. (7.24), and Section 6.10). Subsequent physical projection to $\mathcal{H}_{\text{phys}} := H^0(\hat{Q}_{\text{BRST}})$ is performed only after plateau certification (cf. Section 7.5).

Proposition 9.13 (Spectral discipline and endogenous regularization)**.** *Assume the* Axiom of Spectral Smoothness *and that* $\hat{D} \in \mathfrak{D}_{\text{admissible}}$ *(cf. Definitions 4.341, 4.386, 4.392 and 4.428). Together with the* Principle of Finite Entanglement Capacity *(cf. Definitions 1.18 and 1.34), all triadic distances/times—*$d_{\text{corr}}, d_{\text{vac}}, d_{\text{spec}}$ *(cf. Definitions 4.338, 4.428 and 4.463)—and the operator fields derived from them, including* $\hat{g}_{\mu\nu}$, $\hat{t}_{\mu\nu}$, *and* $\hat{\Sigma}_{\mu\nu}$ *(cf. eq. (5.8), Definition 5.17, and Section 5.2.5) are finite on the band, without external UV cutoffs (cf. Definition 9.10 and eqs. (7.130), (7.143) and (9.2)).*

Justification of Proposition 9.13. Immediate from Definitions 4.341, 4.386, 4.392 and 4.428 (self-regularizing spectral geometry) together with bounded expectations/finite-capacity assumptions (cf. Definitions 1.18 and 1.34) and the constructions of eqs. (5.8), (7.130), (7.143) and (9.2), Definitions 5.17 and 9.10, and Section 5.2.5. No additional argument is required.

Proposition 9.14 (Bundle well-posedness and gauge/frame covariance)**.** *Under Relational and Spectral Smoothness (cf. Definitions 4.341, 4.428 and 6.7), the triads* $(\vec{d}, \vec{t})$ *vary smoothly over* $\mathcal{X}_q$ *(cf. Sections 5.1.4 and 5.2.5), local frame changes act orthogonally on the triads, and the induced connections/holonomies are intrinsic to the Holo-State and Fourfold/Spectral data (cf. Definitions 6.184 and 6.192 and Proposition 9.6).*

Justification of Proposition 9.14. Smooth variation and frame covariance follow from the Relational Levi–Civita construction (cf. Definition 6.7) together with spectral smoothness (cf. Definitions 4.341 and 4.428); gauge/frame covariance and intrinsic holonomies are inherited from the operator–bundle setup (cf. Definitions 6.184 and 6.192) and the correlator's covariance (cf. Proposition 9.6). No further proof is needed.

Remark 9.15 (IR window properties and cone preservation)**.** The windowing map $F \mapsto F^{(\ell)}$ is positivity–preserving and contractive (cf. Definition 6.280 and eq. (7.130)); with causal transport, cones defined by $\bar{\Sigma}^{(\ell)}$ are not enlarged relative to those of $\bar{\Sigma}$ (cf. Definitions 5.122 and 6.7). BRST–closure of $K_{\text{IR}}^{(\ell)}$ ensures compatibility with the physical projector on cohomology, via band–level hygiene (cf. Lemma 9.11, eq. (7.24), and Section 6.10).

Justification of Remark 9.15. Positivity/contractivity of the IR window is by construction (cf. Definition 6.280 and eq. (7.130)); cone preservation follows from causal transport and the effective light-cone definition for $\bar{\Sigma}$ (cf. Definitions 5.122 and 6.7). BRST-compatibility with P_{phys} is ensured by tag hygiene (cf. Lemma 9.11, eq. (7.24), and Section 6.10).

Corollary 9.16 (Σ-riders (no structural change))**.** *The chronogeometric operator fields* $\hat{g}_{\mu\nu}$, $\hat{t}_{\mu\nu}$ *and* $\hat{\Sigma}_{\mu\nu} := \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu}$ *live on the same band–regulated fibres as* $\hat{\mathbf{G}}_\Lambda$*, transform covariantly under* $\mathcal{U}$*, and—by Proposition 9.13—admit SOT–Hessians at coincidence. Thus, replacing "triads only" by* Σ *introduces* no *change to the arena/transport/grade/band scaffolding; it only adds derived operator fields that are already finite and smooth on the band.*

Proof. (i) Band localisation and fibres. The Fourfold Inter–Modal Correlator and its block structure are fixed by Definition 6.4 and eq. (4.24). Band–tagging is implemented by the UV projector and IR window via Definitions 6.280 and 9.10 and eqs. (7.130), (7.143) and (9.2). Hence all objects obtained from $\hat{\mathbf{G}}_\Lambda$ by band–tagging (including distances/times and their Hessians) act on the same band–regulated fibres.

(ii) Gauge/transport covariance and graded kinematics. The transporter $\mathcal{U}$ and the operator–bundle connection are defined in Definitions 6.184 and 6.192, and graded covariant kinematics in Definition 9.7 and Section 4.1.7. Gauge covariance of the correlator (and thus of any covariant functional built from it) is given by Proposition 9.6. Therefore $\hat{g}_{\mu\nu}$ and $\hat{t}_{\mu\nu}$—being built from covariant second variations of band–regular functionals—transform by endpoint conjugation and preserve the $\mathbb{Z}_2 \times \mathbb{Z}_2$ degrees (cf. Definition 9.8).

(iii) Existence and finiteness of SOT–Hessians. The triadic distances/times $d_{\mathrm{corr}}, d_{\mathrm{vac}}, d_{\mathrm{spec}}$ are defined in Definitions 4.338, 4.428 and 4.463, with spectral generator $\hat{D}$ in Definitions 4.341 and 4.386. Under the Axiom of Spectral Smoothness and the Principle of Finite Entanglement Capacity, Proposition 9.13 ensures band–level finiteness and smoothness, so the operator–valued Hessians at coincidence defining $\hat{g}_{\mu\nu}$ and $\hat{t}_{\mu\nu}$ exist in the strong operator topology (cf. eq. (5.8), Definition 5.17, and Section 5.2.5). Hence $\hat{\Sigma}_{\mu\nu} = \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu}$ is likewise band–regular and covariant.

(iv) Structural invariance. The arena/transport/grade/band scaffolding is precisely that of Section 9.1 and Definitions 9.7, 9.8 and 9.10; replacing the separate triads by the unified Σ does not alter these structures, only the (already finite) derived operator fields used as inputs. □

Remark 9.17 (Where BRST enters later)**.** Band tags are BRST–closed by Lemma 9.11 once the checklist eq. (7.24) holds; the pre–physical observable and PSR notion are defined in Definition 9.12. These are arena–level diagnostics and not yet physical. Only after projection to the cohomology $\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}})$ (cf. Section 6.10) are spectra and visibilities evaluated with gauge–parameter independence ensured by the BRST/Nielsen framework (cf. Principle 6.1 and Section 7.5).

Summary of Section 9.1. The construction is placed on the emergent/GNS arena $\mathcal{H}_{\mathrm{em}}$ (cf. Definition 1.34 and Remark 1.36), where the Fourfold Inter–Modal Correlator $\hat{\mathbf{G}}_\Lambda$ is defined with its block structure (cf. Definition 6.4 and eq. (4.24)) and transported by the gauge–covariant parallel transporter $\mathcal{U}$ associated with the operator–bundle connection (cf. Definitions 6.184 and 6.192). Hermiticity of the Twofold block and covariance of the full Fourfold under gauge changes ensure that expectation values in gauge–covariant states are gauge–invariant (cf. Proposition 9.6). Channel components form a $\mathbb{Z}_2 \times \mathbb{Z}_2$ graded $*$–algebra (cf. Definition 9.8), and differentiation is performed by the graded covariant derivative $\mathfrak{D}_\mu$ (cf. Definition 9.7 and Section 4.1.7). A pre–physical window is selected by the band tag $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$, built from the UV spectral projector $P_x(\lambda_{\mathrm{UV}})$ and a causal IR window kernel $K_{\mathrm{IR}}^{(\ell)}$ (cf. Definitions 6.280 and 9.10 and eqs. (7.130), (7.143) and (9.2)). Band hygiene guarantees BRST–compatibility, namely $[\hat{Q}_{\mathrm{BRST}}, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}]_\pm = 0$ once the checklist holds (cf. Lemma 9.11 and eq. (7.24)), and the construction preserves microcausality and locality of commutators (cf. Definition 1.32 and Lemma 9.11). Spectral discipline—combining the admissible spectral operator and self–regularizing spectral distance with the finite–capacity assumptions—yields

finiteness and smoothness on the band for the triadic distances/times $d_{\mathrm{corr}}, d_{\mathrm{vac}}, d_{\mathrm{spec}}$ and for the operator fields derived from them (cf. Definitions 1.18, 1.34, 4.338, 4.341, 4.386, 4.392, 4.428 and 4.463). In particular, the metric and temporal Hessians $\hat{g}_{\mu\nu}$ and $\hat{t}_{\mu\nu}$, and the chronogeometry $\hat{\Sigma}_{\mu\nu} := \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu}$, exist in the strong–operator topology and vary smoothly over $\mathcal{X}_q$ (cf. Propositions 9.13 and 9.14, eq. (5.8), Definition 5.17, and Section 5.2.5). Consequently, $\hat{\Sigma}$ "rides" on exactly the same arena/transport/grade/band scaffolding as the correlator $\hat{\mathbf{G}}_\Lambda$ and introduces no structural change (cf. Corollary 9.16); only the inputs to downstream constructions are unified. Expectation–level objects such as the Relational Levi–Civita connection and effective cones are then read from $\bar{\Sigma}_{\mu\nu}$ in the appropriate physical layer (cf. Definitions 5.122 and 6.7 and Section 7.5).

9.2 Law vs. State, Visibilities, and the Four Quadrants (Pre-Physical)

Definition 9.18 (Graded action of G on correlators)**.** Let G act on channel components by the graded adjoint implemented with the gauge–covariant transport $\mathcal{U}$ of Definition 9.7 (built from the operator–bundle connection; cf. Definitions 6.184 and 6.192). For $g \in G$,

$$\hat{\mathbf{G}}_\Lambda(x,x') \;\longmapsto\; \mathrm{Ad}_g\,\hat{\mathbf{G}}_\Lambda(x,x') \;:=\; g(x)\,\hat{\mathbf{G}}_\Lambda(x,x')\,g(x')^\dagger,$$

with block entries $\hat{G}_\Lambda^{\alpha\beta}$ defined as in Definitions 4.9 and 6.4 and eq. (4.24). This action is compatible with the Fourfold bundle structure and graded kinematics (cf. Sections 4.1.2 and 4.1.7 and Definition 9.8) and preserves the Hermiticity/covariance properties of the correlator (cf. Proposition 9.6).

Definition 9.19 (Correlator amplitude and normalizations)**.** Fix local modal vectors $|\psi_1\rangle_x, |\psi_2\rangle_{x'}$ in the Fourfold fibres (cf. Section 4.6) and define the scalar amplitude and normalizations by

$$\begin{aligned} \mathsf{G}(x,x';\psi_1,\psi_2) &:= \Big\langle \mathring{\mathbf{\Psi}} \Big|\, \hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})[\psi_1^*,\psi_2] \,\Big|\, \mathring{\mathbf{\Psi}} \Big\rangle, \\ N(\psi_1,x) &:= \mathsf{G}(x,x;\psi_1,\psi_1)^{1/2}, \\ N(\psi_2,x') &:= \mathsf{G}(x',x';\psi_2,\psi_2)^{1/2}. \end{aligned} \tag{9.3}$$

Here $\hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'})[\psi_1^*,\psi_2]$ is the bilinear evaluation of the Fourfold correlator with parallel transport $\mathcal{U}(x \leftarrow x';\gamma)$ aligning frames (cf. Definition 4.9, eq. (4.24), and Proposition 9.6). Expectations are taken in the GNS/Holo-State sense (cf. Definitions 1.18 and 1.34), and band–limited locality is enforced by the tag $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ (cf. Definitions 6.280 and 9.10 and eq. (9.2)).

Definition 9.20 (Operator-valued chronogeometry on the band)**.** On the band $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ of Definition 9.10, form the composite operator distances and temporal functionals from the correlator–based triads $d_{\mathrm{corr}}, d_{\mathrm{vac}}, d_{\mathrm{spec}}$ (cf. Definitions 4.338, 4.428 and 4.463) with spectral generator $\hat{D}$ (cf. Definitions 4.341 and 4.386). Assuming spectral discipline (cf. Proposi-

tion 9.13 and Remark 4.425), their SOT–C^2 Hessians at coincidence yield

$$\begin{aligned}\hat{g}_{\mu\nu}(x) &:= \partial_{x^\mu}\partial_{x'^\nu} D^2(x,x')\Big|_{x'=x},\\ \hat{t}_{\mu\nu}(x) &:= \partial_{x^\mu}\partial_{x'^\nu} T^2(x,x')\Big|_{x'=x},\\ \hat{\Sigma}_{\mu\nu}(x) &:= \hat{g}_{\mu\nu}(x) + \hat{t}_{\mu\nu}(x),\end{aligned} \tag{9.4}$$

which are Hermitian, symmetric in μ,ν, and covariant under local frame changes (cf. Definition 9.7 and Proposition 9.14). At the expectation level $\bar{\Sigma}_{\mu\nu} := \langle\hat{\Sigma}_{\mu\nu}\rangle = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (cf. eq. (5.8) and Definitions 5.17 and 6.7).

Remark 9.21 (Finiteness and band regularity)**.** Under Spectral Smoothness and Finite Entanglement Capacity (cf. Definitions 1.34, 4.392 and 4.428), all triadic distances/times are finite on the band, hence $\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu}$ and $\hat{\Sigma}_{\mu\nu}$ are well–defined and smooth (cf. Propositions 9.13 and 9.14). Consequently, for any $\varepsilon > 0$ there exist effective cutoffs $\Lambda_{\text{eff}}(x,x';\varepsilon)$ consistent with the band construction (cf. Definition 9.10 and eq. (7.130)) and with causal cones determined by $\bar{\Sigma}$ (cf. Definition 5.122).

Definition 9.22 (Chronogeometric quadratic form)**.** For a small displacement $v = v^\mu\partial_\mu|_x$, define on the band (cf. Definition 9.10) the local quadratic form built from the operator–valued chronogeometry (cf. Definition 9.20):

$$\hat{Q}_x[v] \;:=\; v^\mu v^\nu\,\hat{\Sigma}_{\mu\nu}(x), \qquad \bar{Q}_x[v] \;:=\; \langle\hat{Q}_x[v]\rangle \;=\; v^\mu v^\nu\,\bar{\Sigma}_{\mu\nu}(x). \tag{9.5}$$

Here $\hat{\Sigma}_{\mu\nu} = \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu}$ is obtained as the SOT–C^2 Hessian at coincidence of the composite distance/time functionals (cf. eqs. (5.8) and (9.4) and Definition 5.17), and $\bar{\Sigma}_{\mu\nu} := \langle\hat{\Sigma}_{\mu\nu}\rangle$ (cf. Definition 6.7). The tensor $\hat{\Sigma}_{\mu\nu}$ transforms covariantly under local frame changes, hence $\hat{Q}_x[v]$ transforms by endpoint conjugation (cf. Definition 9.7 and Proposition 9.14); by gauge covariance of the correlator and cyclicity of the Born pairing, the expectation $\bar{Q}_x[v]$ is gauge–invariant in gauge–covariant states (cf. Proposition 9.6, Definitions 1.18 and 1.34, and Section 4.19.1).

Definition 9.23 (Pre-physical Σ-law and Σ-state contrasts)**.** Let $\mathcal{I}$ be any gauge–invariant, band–local functional (e.g. Holo–State Born pairing or a regulated band–trace; cf. Definitions 1.18, 1.34, 6.280 and 9.10 and eq. (9.2)). Using the graded adjoint action Ad_g of G on correlator–derived objects (cf. Definitions 9.7 and 9.18 and Proposition 9.6), define, for $g \in G$ and $v = v^\mu\partial_\mu|_x$,

$$\Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}}(g;x;v) := \Big(\mathcal{I}\Big[\hat{Q}_x[v] - \mathrm{Ad}_{g(x)}\,\hat{Q}_x[v]\Big]\Big)^{1/2}, \tag{9.6}$$

$$\Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}}(g;x;v) := \Big|\mathcal{I}_\Psi\Big[\hat{Q}_x[v]\Big] - \mathcal{I}_{g\cdot\Psi}\Big[\hat{Q}_x[v]\Big]\Big|^{1/2}, \tag{9.7}$$

where $\mathcal{I}_\Psi[\cdot] = \langle\Psi|\cdot|\Psi\rangle$ is the pre–physical Holo–State evaluation (cf. Definition 1.18), and $g\cdot\Psi$ denotes the push–forward of the Holo–State by the unitary representation implementing g on the bundle (implicit in Definitions 6.184 and 6.192 and Proposition 9.6). These contrasts are

computed on the band and are BRST–functorial whenever the tag is hygienic (cf. Lemma 9.11 and eq. (7.24)); they quantify, respectively, explicit law asymmetry and state asymmetry at the pre–physical level, prior to projection to $\mathcal{H}_{\text{phys}} := H^0(\hat{Q}_{\text{BRST}})$ (cf. Sections 6.10 and 7.5).

Lemma 9.24 (Gauge/frame covariance of contrasts)**.** *The quantities $\Delta^{\text{pre},\Sigma}_{\text{law}}$ and $\Delta^{\text{pre},\Sigma}_{\text{state}}$ are independent of local frame choices and respect the graded action of G described in Definition 9.18. In particular, they depend only on the conjugacy class of g at x.*

Proof. Recall that the chronogeometric quadratic form is $\hat{Q}_x[v] = v^\mu v^\nu \hat{\Sigma}_{\mu\nu}(x)$ with $\hat{\Sigma}_{\mu\nu}$ the covariant SOT–C^2 Hessian on the band (cf. Definitions 9.20 and 9.22). Under a local frame change $h : \mathcal{X}_q \to \mathrm{U}(\mathcal{H}_{\text{em}})$ (implemented by the operator–bundle connection/transport; cf. Definitions 6.184, 6.192 and 9.7), the correlator and any tensor derived covariantly from it transform by endpoint conjugation (cf. Proposition 9.6), hence

$$\hat{Q}_x[v] \;\longmapsto\; \mathrm{Ad}_{h(x)}\,\hat{Q}_x[v] := h(x)\,\hat{Q}_x[v]\,h(x)^\dagger. \tag{9.8}$$

Likewise, the graded adjoint action of $g \in G$ on correlator–derived objects transforms as

$$\mathrm{Ad}_{g(x)} \;\longmapsto\; \mathrm{Ad}_{h(x)g(x)h(x)^{-1}}, \qquad \text{and} \qquad \mathrm{Ad}_{h(x)g(x)h(x)^{-1}} \circ \mathrm{Ad}_{h(x)} = \mathrm{Ad}_{h(x)} \circ \mathrm{Ad}_{g(x)}, \tag{9.9}$$

(cf. Definitions 9.7 and 9.18). Band–local, gauge–invariant functionals $\mathcal{I}$ and $\mathcal{I}_\Psi[\cdot] = \langle\Psi|\cdot|\Psi\rangle$ are conjugation invariant by construction (Born cyclicity/unitarity; cf. Definitions 1.18, 1.34 and 9.23 and Section 4.19.1), i.e.

$$\mathcal{I}[\mathrm{Ad}_h\, X] = \mathcal{I}[X], \qquad \mathcal{I}_{h\cdot\Psi}[\mathrm{Ad}_h\, X] = \mathcal{I}_\Psi[X], \tag{9.10}$$

where $h \cdot \Psi$ denotes the pushed-forward Holo–State (unitary representation).

Law contrast. Using (9.8)–(9.10) in the definition of $\Delta^{\text{pre},\Sigma}_{\text{law}}$ (cf. Definition 9.23):

$$\begin{aligned}
\Big(\Delta^{\text{pre},\Sigma}_{\text{law}}(g;x;v)\Big)^2 &= \mathcal{I}\Big[\hat{Q}_x[v] - \mathrm{Ad}_{g(x)}\,\hat{Q}_x[v]\Big] \\
&\xrightarrow{h}\; \mathcal{I}\Big[\mathrm{Ad}_{h(x)}\,\hat{Q}_x[v] - \mathrm{Ad}_{h(x)g(x)h(x)^{-1}}\,\mathrm{Ad}_{h(x)}\,\hat{Q}_x[v]\Big] \\
&= \mathcal{I}\Big[\mathrm{Ad}_{h(x)}\Big(\hat{Q}_x[v] - \mathrm{Ad}_{g(x)}\,\hat{Q}_x[v]\Big)\Big] = \mathcal{I}\Big[\hat{Q}_x[v] - \mathrm{Ad}_{g(x)}\,\hat{Q}_x[v]\Big]\,,
\end{aligned}$$

which shows frame independence. The middle line used (9.9). Consequently, $\Delta^{\text{pre},\Sigma}_{\text{law}}$ depends only on the conjugacy class of $g(x)$.

State contrast. Similarly, with $\mathcal{I}_\Psi[\cdot] = \langle\Psi|\cdot|\Psi\rangle$ (cf. Definition 9.23) and the representation property $(hgh^{-1})\cdot(h\cdot\Psi) = h\cdot(g\cdot\Psi)$:

$$\begin{aligned}
\Big(\Delta^{\text{pre},\Sigma}_{\text{state}}(g;x;v)\Big)^2 &= \Big|\mathcal{I}_\Psi\Big[\hat{Q}_x[v]\Big] - \mathcal{I}_{g\cdot\Psi}\Big[\hat{Q}_x[v]\Big]\Big| \\
&\xrightarrow{h}\; \Big|\mathcal{I}_{h\cdot\Psi}\Big[\,\mathrm{Ad}_{h(x)}\,\hat{Q}_x[v]\Big] - \mathcal{I}_{(hgh^{-1})\cdot(h\cdot\Psi)}\Big[\,\mathrm{Ad}_{h(x)}\,\hat{Q}_x[v]\Big]\Big| \\
&= \Big|\mathcal{I}_\Psi\Big[\hat{Q}_x[v]\Big] - \mathcal{I}_{g\cdot\Psi}\Big[\hat{Q}_x[v]\Big]\Big|,
\end{aligned}$$

by (9.10). Hence $\Delta^{\text{pre},\Sigma}_{\text{state}}$ is frame independent and depends only on the conjugacy class of $g(x)$.

Both contrasts therefore respect the graded action of G (they are built from the graded adjoint, cf. Definition 9.18) and are invariant under local frame changes dictated by the operator–bundle connection/transport (cf. Definitions 6.184, 6.192 and 9.7). This completes the proof. □

Proposition 9.25 (Reduction to correlator-only distances)**.** *If $\hat{\Sigma}_{\mu\nu}$ is decomposed into triadic parts* $(\mathrm{corr}, \mathrm{vac}, \mathrm{spec})$ *as in the expectation–level decomposition of Definition 5.54 (cf. also eqs. (5.8) and (5.21)), then choosing the band–local, gauge–invariant functional $\mathcal{I}$ channelwise yields*

$$\Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}} \rightsquigarrow \left(\Delta^{\mathrm{pre,corr}}_{\mathrm{law}}, \Delta^{\mathrm{pre,vac}}_{\mathrm{law}}, \Delta^{\mathrm{pre,spec}}_{\mathrm{law}}\right), \quad \Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}} \rightsquigarrow \left(\Delta^{\mathrm{pre,corr}}_{\mathrm{state}}, \Delta^{\mathrm{pre,vac}}_{\mathrm{state}}, \Delta^{\mathrm{pre,spec}}_{\mathrm{state}}\right),$$

and, in particular, the correlator–only contrasts of the original construction when the other channels are deactivated.

Proof. Triadic split and quadratic forms. By Definition 9.20, the band–local chronogeometry admits the operator–level triadic split $\hat{\Sigma}_{\mu\nu} = \sum_{a\in\{\mathrm{corr,vac,spec}\}} \hat{\Sigma}^{(a)}_{\mu\nu}$, consistent with the expectation–level decomposition $\bar{\Sigma}_{\mu\nu} = \sum_a \bar{\Sigma}^{(a)}_{\mu\nu}$ (cf. Definition 5.54 and eqs. (5.8) and (5.21)). Define the channelwise quadratic forms (cf. Definition 9.22)

$$\hat{Q}^{(a)}_x[v] := v^\mu v^\nu \hat{\Sigma}^{(a)}_{\mu\nu}(x), \qquad \hat{Q}_x[v] = \sum_a \hat{Q}^{(a)}_x[v].$$

Channelwise functionals. Let $\mathcal{I}$ be a gauge–invariant, band–local functional (e.g. Born pairing or regulated band–trace; cf. Definitions 1.18, 1.34, 9.10 and 9.23 and eq. (9.2)). By the $\mathbb{Z}_2 \times \mathbb{Z}_2$ grading and block structure of the Fourfold bundle (cf. Definition 9.8, eq. (4.24), and Section 4.1.7), there exist channel–projected functionals $\mathcal{I}^{(a)}$ that act on the (a)–block and annihilate the others, i.e. $\mathcal{I}^{(a)}[\hat{Q}^{(b)}_x[v]] = 0$ for $b \neq a$ and $\mathcal{I}^{(a)}[\hat{Q}_x[v]] = \mathcal{I}^{(a)}[\hat{Q}^{(a)}_x[v]]$. (Concretely, $\mathcal{I}^{(a)}$ is obtained by inserting the band–local projector onto the (a)–block before taking the expectation/trace; the existence of these projectors follows from the graded algebra and transport covariance, cf. Definition 9.7 and Proposition 9.6.)

Law contrast. Using Definition 9.23 and linearity of $\mathcal{I}^{(a)}$,

$$\begin{aligned}\left(\Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}}(g;x;v)\right)^2 &= \mathcal{I}\left[\hat{Q}_x[v] - \mathrm{Ad}_{g(x)}\,\hat{Q}_x[v]\right] = \sum_a \mathcal{I}^{(a)}\left[\hat{Q}^{(a)}_x[v] - \mathrm{Ad}_{g(x)}\,\hat{Q}^{(a)}_x[v]\right] \\ &=: \sum_a \left(\Delta^{\mathrm{pre},a}_{\mathrm{law}}(g;x;v)\right)^2,\end{aligned}$$

where the graded adjoint $\mathrm{Ad}_{g(x)}$ respects the channel decomposition (cf. Definition 9.18). Thus evaluating with each $\mathcal{I}^{(a)}$ yields the channelwise law contrasts $\Delta^{\mathrm{pre},a}_{\mathrm{law}}$.

State contrast. Similarly, with $\mathcal{I}_\Psi[\cdot] = \langle\Psi|\cdot|\Psi\rangle$ (cf. Definition 9.23),

$$\begin{aligned}\left(\Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}}(g;x;v)\right)^2 &= \left|\mathcal{I}_\Psi[\hat{Q}_x[v]] - \mathcal{I}_{g\cdot\Psi}[\hat{Q}_x[v]]\right| = \left|\sum_a \left(\mathcal{I}_\Psi[\hat{Q}^{(a)}_x[v]] - \mathcal{I}_{g\cdot\Psi}[\hat{Q}^{(a)}_x[v]]\right)\right| \\ &\rightsquigarrow \left(\Delta^{\mathrm{pre,corr}}_{\mathrm{state}}, \Delta^{\mathrm{pre,vac}}_{\mathrm{state}}, \Delta^{\mathrm{pre,spec}}_{\mathrm{state}}\right),\end{aligned}$$

by evaluating with $\mathcal{I}^{(a)}_\Psi$ (the channel–projected Born pairing). Gauge/frame covariance of these contrasts follows from Lemma 9.24.

Correlator–only reduction. If the vacuum and spectral channels are deactivated (or their projectors are set to zero on the band), one retains only the correlator channel. In that case, $\hat{\Sigma}_{\mu\nu} \to \hat{\Sigma}^{(\mathrm{corr})}_{\mu\nu}$ and $\hat{Q}_x[v] \to \hat{Q}^{(\mathrm{corr})}_x[v]$, and the contrasts reduce to the correlator–only diagnostics built from d_{corr} and its induced metric data (cf. Definition 4.338 and Section 4.10.2). Analogous single–channel reductions hold for the vacuum and spectral channels (cf. Definitions 4.428 and 4.463).

Independence/orthogonality. Functional orthogonality and Fréchet–type independence of the triadic generators validate the channelwise treatment above (cf. Section 4.11.5), ensuring that no cross–terms are introduced by the graded action.

Combining the steps yields the claim. □

Definition 9.26 (Aggregated Σ-contrasts and visibilities)**.** Given a finite probe set $\{v_k\}_{k=1}^m \subset T_x\mathcal{X}_q$, define the aggregated (root–sum–square) pre–physical Σ–contrasts built from the pointwise contrasts of Definition 9.23 and the quadratic form of Definition 9.22:

$$\begin{aligned} \Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}}(g;x) &:= \left(\sum_{k=1}^{m} \left[\Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}}(g;x;v_k) \right]^2 \right)^{1/2}, \\ \Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}}(g;x) &:= \left(\sum_{k=1}^{m} \left[\Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}}(g;x;v_k) \right]^2 \right)^{1/2}. \end{aligned} \tag{9.11}$$

Here each $\Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}}(g;x;v_k)$ and $\Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}}(g;x;v_k)$ is evaluated with $\hat{Q}_x[v_k] = v_k^\mu v_k^\nu \hat{\Sigma}_{\mu\nu}(x)$ as in Definition 9.22, using any band–local, gauge–invariant functional $\mathcal{I}$ (e.g. Holo–State Born pairing or regulated band–trace; cf. Definitions 1.18, 1.34, 6.280 and 9.10 and eq. (9.2)). By frame covariance of $\hat{Q}_x[v]$ and conjugation–invariance of the functionals (cf. Proposition 9.6, Definition 9.7, and Section 4.19.1), and by Lemma 9.24, the aggregated quantities are independent of local frame choices and respect the graded G–action of Definition 9.18. Channelwise decompositions and correlator–only reductions are consistent with Proposition 9.25, Definition 5.54, and eqs. (5.8) and (5.21). Operationally, the associated *pre–physical visibilities* are any such band–local, gauge–invariant functionals of $\hat{Q}_x[v]$ (or $\bar{Q}_x[v] = \langle \hat{Q}_x[v] \rangle$) used to estimate (9.11) (cf. Definitions 9.20 and 9.23).

Lemma 9.27 (Finiteness and stability under band deformations)**.** *Under the hypotheses of Remark 9.21, the maps $v \mapsto \Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}}(g;x;v)$ and $v \mapsto \Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}}(g;x;v)$ are continuous and bounded; moreover, small changes in the band parameters $(\lambda_{\mathrm{UV}}, \ell) \mapsto (\lambda'_{\mathrm{UV}}, \ell')$ induce $\mathcal{O}(\varepsilon)$ variations controlled by the effective cutoff $\Lambda_{\mathrm{eff}}(x,x';\varepsilon)$ of Remark 9.21.*

Proof. (1) Continuity and boundedness in v. By Definitions 9.20 and 9.22, $\hat{Q}_x[v] = v^\mu v^\nu \hat{\Sigma}_{\mu\nu}(x)$ with $\hat{\Sigma}_{\mu\nu}$ the band–regular, Hermitian SOT–C^2 Hessian (existence/regularity from Propositions 9.13 and 9.14). Let $\|\cdot\|$ denote the operator norm on the band fibre $P_x(\lambda_{\mathrm{UV}})\mathcal{H}_{\mathrm{total}}$. Then

$$\|\hat{Q}_x[v]\| \;\le\; \|\hat{\Sigma}(x)\|\,\|v\|^2, \qquad \|\hat{Q}_x[v] - \hat{Q}_x[w]\| \;\le\; \|\hat{\Sigma}(x)\| \left(\|v\| + \|w\|\right) \|v - w\|,$$

so $v \mapsto \hat{Q}_x[v]$ is continuous and (locally) Lipschitz. For any admissible, gauge–invariant, band–local functional $\mathcal{I}$ (e.g. Born pairing or regulated band–trace; cf. Definitions 1.18, 6.280,

9.10 and 9.23 and eq. (9.2)), boundedness $\big|\mathcal{I}[X]\big| \le C\,\|X\|$ holds with C determined by the state/trace. Therefore

$$\Big(\Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}}(g;x;v)\Big)^2 = \mathcal{I}\Big[\hat{Q}_x[v] - \mathrm{Ad}_{g(x)}\,\hat{Q}_x[v]\Big] \;\le\; 2C\,\|\hat{\Sigma}(x)\|\,\|v\|^2,$$

and similarly for $\Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}}$, showing boundedness on bounded v–sets and continuity by composition (cf. Definitions 9.18 and 9.23 and Proposition 9.6).

(2) Stability under band deformations $(\lambda_{\mathrm{UV}},\ell)$. Write the band tag as in Definition 9.10 and eq. (9.2), with its parameter dependence encoded by $P_x(\lambda_{\mathrm{UV}})$ and $K^{(\ell)}_{\mathrm{IR}}$ (cf. Definition 6.280 and eq. (7.130)). Under Spectral Smoothness and Finite Entanglement Capacity, Remark 9.21 provides, for each $\varepsilon>0$, an effective cutoff $\Lambda_{\mathrm{eff}}(x,x';\varepsilon)$ such that modes above Λ_{eff} contribute less than ε to all band–local functionals under consideration. For the UV part, the variation of the spectral projector obeys

$$\Big\|P_x(\lambda'_{\mathrm{UV}}) - P_x(\lambda_{\mathrm{UV}})\Big\| \;\le\; C_{\mathrm{UV}}\,\Theta\Big(\Lambda_{\mathrm{eff}} - \min\{\lambda_{\mathrm{UV}},\lambda'_{\mathrm{UV}}\}\Big) \;+\; \mathcal{O}(\varepsilon),$$

with the $\mathcal{O}(\varepsilon)$ term supplied by the cutoff (the precise constant depends on the spectral density near the band edge but the tail is ε–small by construction). For the IR part, strong continuity $\|K^{(\ell')}_{\mathrm{IR}} - K^{(\ell)}_{\mathrm{IR}}\| \to 0$ as $\ell' \to \ell$ follows from the window definition and causal regularity (cf. Definition 6.280), hence

$$\Big\|\mathfrak{B}_{\lambda'_{\mathrm{UV}},\ell'}[\mathfrak{O}] - \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\mathfrak{O}]\Big\| \;\le\; C_{\mathfrak{O}}\left(\Big\|P(\lambda'_{\mathrm{UV}}) - P(\lambda_{\mathrm{UV}})\Big\| + \Big\|K^{(\ell')}_{\mathrm{IR}} - K^{(\ell)}_{\mathrm{IR}}\Big\|\right) \;+\; \mathcal{O}(\varepsilon),$$

for local $\mathfrak{O}$ and some constant $C_{\mathfrak{O}}$ (from the bounded transport $\mathcal{U}$, cf. Definitions 6.184 and 6.192). Applying this to the ingredients that define D^2, T^2 and their SOT–Hessians yields

$$\Big\|\hat{\Sigma}'_{\mu\nu}(x) - \hat{\Sigma}_{\mu\nu}(x)\Big\| \;=\; \mathcal{O}(\varepsilon),$$

hence $\Big\|\hat{Q}'_x[v] - \hat{Q}_x[v]\Big\| \le \|v\|^2\,\Big\|\hat{\Sigma}' - \hat{\Sigma}\Big\| = \mathcal{O}(\varepsilon)$ uniformly for $\|v\|\le R$. By boundedness of $\mathcal{I}$ and $\mathcal{I}_\Psi$ (cf. Definition 9.23),

$$\Big|\Big(\Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}}\Big)^2{}' - \Big(\Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}}\Big)^2\Big| \;\le\; C\,\Big\|\hat{Q}'_x[v] - \hat{Q}_x[v]\Big\| \;=\; \mathcal{O}(\varepsilon),$$

and similarly for the state contrast. Taking square roots preserves $\mathcal{O}(\varepsilon)$ control on compact v–sets. This is consistent with the finite–difference stability diagnostics used on the $(\ln\lambda_{\mathrm{UV}}, \ln\ell)$ grid (cf. eqs. (7.394) to (7.396)) and the acceptance rules for plateau certification (cf. Proposition 7.203).

Combining (1) and (2) proves continuity, boundedness, and $\mathcal{O}(\varepsilon)$ stability of the contrasts under small band deformations, with control by $\Lambda_{\mathrm{eff}}(x,x';\varepsilon)$ from Remark 9.21. □

Definition 9.28 (Quadrant predicate (pre-physical, Σ-aware))**.** Fix a tolerance $\varepsilon \ll 1$ and a band $(\lambda_{\mathrm{UV}},\ell)$. Declare at x:

$$\textbf{QI}:\ \Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}}(g;x)<\varepsilon,\ \Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}}(g;x)<\varepsilon; \quad \textbf{QII}:\ \Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}}(g;x)<\varepsilon,\ \Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}}(g;x)>\varepsilon;$$

$$\textbf{QIII}:\ \Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}}(g;x)>\varepsilon,\ \Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}}(g;x)<\varepsilon; \quad \textbf{QIV}:\ \Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}}(g;x)>\varepsilon,\ \Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}}(g;x)>\varepsilon.$$

This mirrors the physical quadrant logic in $H^0(\hat{Q}_{\mathrm{BRST}})$ (cf. Definition 6.224 and Remark 6.223).

Proposition 9.29 (Interpretability via triadic attribution)**.** *Writing the expectation–level chronogeometry as a triadic sum* $\bar{\Sigma}_{\mu\nu} = \bar{\Sigma}^{\mathrm{corr}}_{\mu\nu} + \bar{\Sigma}^{\mathrm{vac}}_{\mu\nu} + \bar{\Sigma}^{\mathrm{spec}}_{\mu\nu}$ *(cf. Definition 5.54 and eqs. (5.8) and (5.21)), a quadrant decision based on* Σ*–contrasts can be attributed to channelwise contributions from coherence, vacuum, and spectral shear, respectively; this attribution preserves gauge invariance and band locality.*

Proof. (i) Channelwise decomposition on the band. At the operator level, $\hat{\Sigma}_{\mu\nu} = \sum_{a\in\{\mathrm{corr,vac,spec}\}} \hat{\Sigma}^{(a)}_{\mu\nu}$ by Definition 9.20, consistent with the expectation–level split quoted above (cf. Definition 5.54). For any probe v, the quadratic form $\hat{Q}_x[v] = v^\mu v^\nu \hat{\Sigma}_{\mu\nu}(x)$ of Definition 9.22 splits as $\hat{Q}_x[v] = \sum_a \hat{Q}^{(a)}_x[v]$ with $\hat{Q}^{(a)}_x[v] := v^\mu v^\nu \hat{\Sigma}^{(a)}_{\mu\nu}(x)$.

(ii) Channel projectors and band–local functionals. By the graded bundle structure and block decomposition of the Fourfold algebra (cf. Definition 9.8, eq. (4.24), and Section 4.1.7), there exist band–local channel projectors $\Pi^{(a)}$ that commute with the transport (graded–covariantly) and select each (a)–block (cf. Definition 9.7 and Proposition 9.6). Define channelwise, gauge–invariant band functionals by $\mathcal{I}^{(a)}[X] := \mathcal{I}[\Pi^{(a)} X \Pi^{(a)}]$, where $\mathcal{I}$ is as in Definitions 9.23 and 9.26. Then $\mathcal{I}^{(a)}[\hat{Q}_x[v]] = \mathcal{I}^{(a)}[\hat{Q}^{(a)}_x[v]]$ and $\mathcal{I}^{(a)}[\hat{Q}^{(b)}_x[v]] = 0$ for $b \neq a$.

(iii) Channelwise attribution of the Σ*–contrasts.* From Definition 9.23 and linearity,

$$\Big(\Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}}(g;x;v)\Big)^2 = \mathcal{I}\Big[\hat{Q}_x[v] - \mathrm{Ad}_{g(x)}\,\hat{Q}_x[v]\Big] = \sum_a \mathcal{I}^{(a)}\Big[\hat{Q}^{(a)}_x[v] - \mathrm{Ad}_{g(x)}\,\hat{Q}^{(a)}_x[v]\Big].$$

The summands define the channelwise law–contributions $\Big(\Delta^{\mathrm{pre},a}_{\mathrm{law}}(g;x;v)\Big)^2$. A completely analogous decomposition holds for the state contrast:

$$\Big(\Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}}(g;x;v)\Big)^2 = \sum_a \Big(\mathcal{I}^{(a)}_\Psi[\hat{Q}^{(a)}_x[v]] - \mathcal{I}^{(a)}_{g\cdot\Psi}[\hat{Q}^{(a)}_x[v]]\Big),$$

with $\mathcal{I}^{(a)}_\Psi$ the channel–projected Born pairing (cf. Definitions 1.18 and 1.34).

(iv) Gauge invariance and band locality. Each term is invariant under local frame changes by the conjugation invariance of $\mathcal{I}, \mathcal{I}_\Psi$ and the covariance of $\hat{Q}^{(a)}_x[v]$ (cf. Proposition 9.6, Definition 9.7, Section 4.19.1, and Lemma 9.24). Band locality follows from the construction of $\mathcal{I}$ and the tag $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ (cf. Definitions 6.280 and 9.10 and eq. (9.2)). Functional orthogonality/independence of the triadic generators (cf. Section 4.11.5) ensures that no spurious cross–terms arise in the attribution. Therefore a quadrant decision based on the Σ–contrasts can be interpreted as a sum of channelwise, gauge–invariant, band–local contributions from coherence, vacuum, and spectral shear. □

Corollary 9.30 (Consistency with the original correlator-distance scheme)**.** *Setting* $\hat{t}_{\mu\nu} \equiv 0$ *and restricting* $\mathcal{I}$ *to the correlator channel reduces Definitions 9.22, 9.23 and 9.26 to the correlator–only distances/contrasts of the initial pre–physical formulation based on* d_{corr} *with normalizations* $N(\cdot)$ *as in Definition 9.19.*

Proof. With $\hat{t}_{\mu\nu} \equiv 0$, one has $\hat{\Sigma}_{\mu\nu} \to \hat{\Sigma}^{(\mathrm{corr})}_{\mu\nu}$ and $\hat{Q}_x[v] \to \hat{Q}^{(\mathrm{corr})}_x[v]$ (cf. Definitions 9.20 and 9.22). Choosing the functional $\mathcal{I}$ to be the correlator–channel restriction $\mathcal{I}^{(\mathrm{corr})}$ constructed in the proof of Proposition 9.29 yields

$$\Big(\Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}}(g;x;v)\Big)^2 = \mathcal{I}^{(\mathrm{corr})}\Big[\hat{Q}^{(\mathrm{corr})}_x[v] - \mathrm{Ad}_{g(x)}\,\hat{Q}^{(\mathrm{corr})}_x[v]\Big],$$

and the analogous expression for the state contrast. These are precisely the correlator–only pre–physical law/state contrasts, i.e. the ones obtained from the correlator distance d_{corr} (cf. Definition 4.338) with the fibre normalizations $N(\psi, x)$ defined by Definition 9.19 and the GNS expectation (cf. Definitions 1.18 and 1.34). The aggregated versions Definition 9.26 reduce accordingly. Thus the Σ–formulation is consistent with, and collapses to, the original correlator–only scheme when the temporal and spectral channels are deactivated. □

Operational note. When working at the expectation level, the chronogeometric quadratic form may be evaluated as $\bar{Q}_x[v] = v^\mu v^\nu \bar{\Sigma}_{\mu\nu}(x)$ with $\bar{\Sigma}_{\mu\nu} := \langle \hat{\Sigma}_{\mu\nu} \rangle$ (cf. Definitions 6.7, 9.20 and 9.22). Using the same band–local, gauge–invariant functionals $\mathcal{I}$ as in the operator–level definitions (Born pairing or regulated band–trace; cf. Definitions 1.18, 1.34, 6.280 and 9.10), one obtains expectation–level $\bar{\Delta}$ versions of (9.6)–(9.7). Gauge/frame covariance follows from the conjugation invariance of the functionals together with the transport covariance of $\hat{\Sigma}_{\mu\nu}$ (cf. Proposition 9.6, Definition 9.7, and Lemma 9.24). The band/plateau discipline of Section 9.1—namely BRST hygiene of the tag (cf. Lemma 9.11 and eq. (7.24)), causal IR windowing (cf. Definition 6.280), and spectral smoothness/finitedness (cf. Proposition 9.13 and Remark 9.21)—ensures that these $\bar{\Delta}$ inputs are well–defined and ready for the realization step on the physical cohomology (cf. Sections 6.10 and 9.5). Plateau certification and acceptance (cf. eqs. (7.394) to (7.396) and Proposition 7.203) then license the Σ–realization and evaluation of physical, gauge–parameter–independent spectra/visibilities.

Summary of Section 9.2. The "distances only" inputs were replaced by the unified operator $\hat{\Sigma}_{\mu\nu}$ (cf. Definition 9.20 and eq. (9.4)), from which the chronogeometric quadratic form $\hat{Q}_x[v]$ was built (cf. Definition 9.22). On this basis, the Σ–law and Σ–state contrasts (9.6)–(9.7) were defined using band–local, gauge–invariant functionals, with full gauge/frame covariance under the graded G–action (cf. Definition 9.18 and Lemma 9.24). These contrasts admit aggregated versions for finite probe sets (cf. Definition 9.26) and possess channelwise attributions by the triadic split (cf. Proposition 9.29, Definition 5.54, and eqs. (5.8) and (5.21)). Finiteness and stability on the band are ensured by spectral discipline and effective cutoffs (cf. Remark 9.21, Proposition 9.13, and Lemma 9.27); moreover, the scheme reduces consistently to the correlator–only construction when the temporal/spectral channels are deactivated (cf. Corollary 9.30 and Definitions 4.338 and 9.19). The resulting Σ–contrasts feed the pre–physical quadrant predicate QI–QIV (cf. Definition 9.28), thereby elevating temporal phase–flow and channel mixing to first–class diagnostic status while remaining fully compatible with the Fourfold correlator/transport scaffolding (cf. Definitions 6.4, 9.7, 9.8 and 9.10, eq. (4.24), and Proposition 9.6).

9.3 From Algebraic Breakings to Quanta

Definition 9.31 (Field blocks and generalized field operator)**.** Let $\hat{\mathbf{F}} = (\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}})^{\mathsf{T}}$ be the generalized field operator (cf. Ch. 7) and let the unified inter-modal correlator decompose as

$$\hat{\mathbf{G}}_\Lambda = \begin{pmatrix} \hat{G}_{\Phi\Phi} & \hat{G}_{\Phi\Theta} \\ \hat{G}_{\Theta\Phi} & \hat{G}_{\Theta\Theta} \end{pmatrix}, \tag{9.12}$$

with blocks acting on the Gaiamiti (Φ) and Tantrakyma (Θ) sectors, respectively; see Definition 9.5 and Ch. 7's block structure. The graded, gauge-covariant transport is as in Definition 9.7.

Definition 9.32 (Law-symmetric tangent map (graded adjoint))**.** For a generator $X \in \mathfrak{g} = \mathrm{Lie}(G)$, the *law-symmetric tangent* of a correlator component is

$$\delta_X \hat{G}^{\alpha\beta}_\Lambda := \frac{d}{dt}\Big|_{t=0} \mathrm{Ad}_{e^{tX}} \hat{G}^{\alpha\beta}_\Lambda = [X, \hat{G}^{\alpha\beta}_\Lambda]_{\mathrm{gr}}, \qquad \alpha, \beta \in \{0, 1, p, s\}, \tag{9.13}$$

where Ad_g is the graded adjoint action of Definition 9.18.

Definition 9.33 (Σ-weighted inner products on operator modes)**.** Restrict the operator-valued chronogeometry $\hat{\Sigma}_{\mu\nu}$ of Definition 9.20 to the field blocks

$$\hat{\Sigma}^{(\Phi)} := \Pi^\dagger_{\Phi\Phi}\, \hat{\Sigma}\, \Pi_{\Phi\Phi}, \qquad \hat{\Sigma}^{(\Theta)} := \Pi^\dagger_{\Theta\Theta}\, \hat{\Sigma}\, \Pi_{\Theta\Theta},$$

and let $\mathcal{I}$ be any gauge-invariant, band-local functional. For band-tagged operator densities O_1, O_2 (on $\mathcal{H}_{\mathrm{em}}$),

$$\langle\!\langle O_1, O_2 \rangle\!\rangle_{\Sigma,\Phi} := \mathcal{I}\Big[O_1^\dagger, \hat{\Sigma}^{(\Phi)}, O_2\Big], \qquad \langle\!\langle O_1, O_2 \rangle\!\rangle_{\Sigma,\Theta} := \mathcal{I}\Big[O_1^\dagger, \hat{\Sigma}^{(\Theta)}, O_2\Big]. \tag{9.14}$$

These are positive semi-definite and gauge/frame covariant on the band specified in Definition 9.10.

Lemma 9.34 (Positivity and covariance)**.** *Under the spectral discipline of Proposition 9.13, $\langle\!\langle \cdot, \cdot \rangle\!\rangle_{\Sigma,\Phi/\Theta}$ are well-defined, positive semi-definite and transform by conjugation under local frame changes; they are invariant under the Born expectation.*

Definition 9.35 (Pre-physical Apeiron and Kymon mode spaces (Σ-aware))**.** Fix a band $(\lambda_{\mathrm{UV}}, \ell)$. Let $\mathfrak{B}$ be the set of *broken* generators X at x in the sense of Definition 9.23 (i.e., $\Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}}(e^{tX}; x) > \varepsilon$ while $\Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}}(e^{tX}; x) < \varepsilon$). Define

$$\mathcal{N}^{\mathrm{pre}}_\Phi := \overline{\mathrm{span}}\Big\{\text{left/right singular vectors of } \delta_X \hat{G}_{\Phi\Phi} \;:\; X \in \mathfrak{B}\Big\},$$

$$\mathcal{N}^{\mathrm{pre}}_\Theta := \overline{\mathrm{span}}\Big\{\text{left/right singular vectors of } \delta_X \hat{G}_{\Theta\Theta} \;:\; X \in \mathfrak{B}\Big\}.$$

The graded commutator $[\cdot, \cdot]_{\mathrm{gr}}$ enforces selection rules for admissible channel mixing (Definition 9.8).

Proposition 9.36 (Σ-weighted generalized eigenproblems (principal modes))**.** *For each broken $X \in \mathfrak{B}$, define the Gram operators*

$$C_\Phi := (\delta_X \hat{G}_{\Phi\Phi})^\dagger\, \hat{\Sigma}^{(\Phi)}\, (\delta_X \hat{G}_{\Phi\Phi}), \qquad C_\Theta := (\delta_X \hat{G}_{\Theta\Theta})^\dagger\, \hat{\Sigma}^{(\Theta)}\, (\delta_X \hat{G}_{\Theta\Theta}), \tag{9.15}$$

where $\delta_X(\cdot)$ is the graded tangent defined by the adjoint action (cf. Definitions 9.7 and 9.18) and $\hat{\Sigma}^{(\Phi)}, \hat{\Sigma}^{(\Theta)}$ are the block–restrictions of the band–local chronogeometry $\hat{\Sigma}$ (cf. Definitions 6.4 and 9.20 and eq. (4.24)). Then the block–wise generalized eigenproblems

$$C_\Phi\, u_a = \lambda_a\, \hat{\Sigma}^{(\Phi)} u_a, \qquad C_\Theta\, v_b = \mu_b\, \hat{\Sigma}^{(\Theta)} v_b, \tag{9.16}$$

admit complete $\hat{\Sigma}$–orthonormal systems $\{u_a\} \subset \mathcal{N}^{\mathrm{pre}}_\Phi$ and $\{v_b\} \subset \mathcal{N}^{\mathrm{pre}}_\Theta$ on the band, with non–negative eigenvalues λ_a, μ_b.

Proof. Step 0 (band–local pre–Hilbert modules and Σ–pairings). On the band specified by the tag $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ (cf. Definitions 6.280 and 9.10 and eq. (9.2)), equip the block spaces with the Σ–weighted sesquilinear forms

$$\langle f,g\rangle_{\Sigma^{(\Phi)}} := \mathcal{I}\Big[f^{\dagger}\hat{\Sigma}^{(\Phi)}g\Big], \qquad \langle f,g\rangle_{\Sigma^{(\Theta)}} := \mathcal{I}\Big[f^{\dagger}\hat{\Sigma}^{(\Theta)}g\Big],$$

where $\mathcal{I}$ is any band–local, gauge–invariant functional (Born pairing or regulated band–trace; cf. Definitions 1.18, 1.34 and 9.23). By spectral discipline (cf. Propositions 9.13 and 9.14 and Remark 9.21), $\hat{\Sigma}$ is positive and defines pre–Hilbert norms $\|f\|^2_{\Sigma^{(\Phi)}} := \langle f,f\rangle_{\Sigma^{(\Phi)}}$ and similarly for Θ. Denote the corresponding completions by $\mathcal{N}^{\mathrm{pre}}_{\Phi}$ and $\mathcal{N}^{\mathrm{pre}}_{\Theta}$.

Step 1 (boundedness and positivity of the Gram forms). Let $A_\Phi := \hat{\Sigma}^{(\Phi)\,1/2}(\delta_X\hat{G}_{\Phi\Phi})$ on the band; define A_Θ analogously. Then $C_\Phi = A_\Phi^{\dagger}A_\Phi$ and $C_\Theta = A_\Theta^{\dagger}A_\Theta$ as operators on the respective blocks. For any $u \in \mathcal{N}^{\mathrm{pre}}_{\Phi}$,

$$\langle u, C_\Phi u\rangle_{\Sigma^{(\Phi)}} = \mathcal{I}\Big[u^{\dagger}(\delta_X\hat{G}_{\Phi\Phi})^{\dagger}\hat{\Sigma}^{(\Phi)}(\delta_X\hat{G}_{\Phi\Phi})u\Big] = \Big\|A_\Phi u\Big\|^2_{L^2(\mathcal{I})} \ge 0,$$

hence C_Φ is positive (and likewise C_Θ). Gauge/frame covariance follows from the graded transport and conjugation–invariance of $\mathcal{I}$ (cf. Definition 9.7, Proposition 9.6, and Section 4.19.1); thus the constructions are frame–independent on the band.

Step 2 (compactness on the band). By the UV projection $P_x(\lambda_{\mathrm{UV}})$ and causal IR window $K^{(\ell)}_{\mathrm{IR}}$ entering the tag (cf. Definitions 6.280 and 9.10), and by spectral smoothness/finite capacity (cf. Proposition 9.13 and Remark 9.21), the band–limited kernels of $\delta_X\hat{G}_{\bullet\bullet}$ are square–integrable on $\mathcal{X}_q\times\mathcal{X}_q$ with respect to the relational measure (cf. Definition 6.33), hence $\delta_X\hat{G}_{\bullet\bullet}$ are Hilbert–Schmidt and therefore compact on the band–local modules. Since $\hat{\Sigma}^{(\bullet)\,1/2}$ are bounded and positive on the band (cf. Definition 9.20), the compositions $A_\bullet = \hat{\Sigma}^{(\bullet)\,1/2}(\delta_X\hat{G}_{\bullet\bullet})$ are also compact; therefore $C_\bullet = A_\bullet^{\dagger}A_\bullet$ are positive, self–adjoint, compact operators on $\mathcal{N}^{\mathrm{pre}}_{\bullet}$.

Step 3 (reduction to an ordinary compact eigenproblem). Define the bounded, positive isomorphisms on the supports of $\hat{\Sigma}^{(\bullet)}$,

$$J_\Phi : \mathcal{N}^{\mathrm{pre}}_{\Phi} \to \mathcal{N}^{\mathrm{pre}}_{\Phi}, \quad J_\Phi u := \hat{\Sigma}^{(\Phi)\,1/2}u, \qquad J_\Theta v := \hat{\Sigma}^{(\Theta)\,1/2}v,$$

and the compact, positive operators

$$K_\Phi := J_\Phi^{-1}C_\Phi J_\Phi^{-1} = \hat{\Sigma}^{(\Phi)\,-1/2}(\delta_X\hat{G}_{\Phi\Phi})^{\dagger}\hat{\Sigma}^{(\Phi)}(\delta_X\hat{G}_{\Phi\Phi})\hat{\Sigma}^{(\Phi)\,-1/2} = A_\Phi^{\dagger}A_\Phi,$$

with K_Θ analogously. Then K_Φ and K_Θ act on the ordinary (unweighted) pre–Hilbert structures and are compact, positive, self–adjoint (by Step 2).

Step 4 (spectral theorem and Σ–orthonormal systems). By the spectral theorem for compact, positive, self–adjoint operators, there exist orthonormal bases $\{w_a\}$ and $\{z_b\}$ with

$$K_\Phi w_a = \lambda_a w_a, \qquad K_\Theta z_b = \mu_b z_b, \qquad \lambda_a,\mu_b \ge 0,$$

and w_a, z_b complete in their respective closures. Setting $u_a := J_\Phi^{-1}w_a = \hat{\Sigma}^{(\Phi)\,-1/2}w_a$ and $v_b := J_\Theta^{-1}z_b = \hat{\Sigma}^{(\Theta)\,-1/2}z_b$ yields

$$C_\Phi u_a = \lambda_a\,\hat{\Sigma}^{(\Phi)}u_a, \qquad C_\Theta v_b = \mu_b\,\hat{\Sigma}^{(\Theta)}v_b,$$

i.e. the generalized eigenproblems (9.16). Moreover,

$$\langle u_a, u_{a'}\rangle_{\Sigma^{(\Phi)}} = \langle w_a, w_{a'}\rangle = \delta_{aa'}, \qquad \langle v_b, v_{b'}\rangle_{\Sigma^{(\Theta)}} = \langle z_b, z_{b'}\rangle = \delta_{bb'},$$

so $\{u_a\}$ and $\{v_b\}$ are Σ–orthonormal systems. Completeness follows from the completeness of $\{w_a\}$ and $\{z_b\}$ and the fact that $J_\bullet$ are isomorphisms on the supports of $\hat\Sigma^{(\bullet)}$ (the null–spaces of $\hat\Sigma^{(\bullet)}$ can be quotient–out as usual if present; cf. positivity/semidefiniteness in Definition 9.20 and Proposition 9.14).

Step 5 (gauge/frame compatibility). All constructions are compatible with the graded G–action and local frames: $C_\bullet$ and $\hat\Sigma^{(\bullet)}$ transform by conjugation under transport (cf. Definition 9.7 and Proposition 9.6), while $\mathcal{I}$ is conjugation–invariant (cf. Section 4.19.1). Hence the spectra $\{\lambda_a\}, \{\mu_b\}$ and the Σ–orthonormality relations are frame–independent on the band.

Combining the steps establishes (9.16) with non–negative eigenvalues and complete $\hat\Sigma$–orthonormal systems on the band. □

Definition 9.37 (Apeiron/Kymon creation operators (pre-physical, Σ-normalized))**.** Let $\{u_a\}$, $\{v_b\}$ be principal solutions of (9.16). There exist band-localized operator densities $\hat A_a^\dagger, \hat K_b^\dagger \in \mathcal{B}(\mathcal{H}_{\mathrm{em}})$ such that

$$\hat G_{\Phi\Phi}^{(\lambda_{\mathrm{UV}},\ell)} = \sum_a s_a\, \hat A_a\, \hat A_a^\dagger, \qquad \hat G_{\Theta\Theta}^{(\lambda_{\mathrm{UV}},\ell)} = \sum_b t_b\, \hat K_b\, \hat K_b^\dagger, \tag{9.17}$$

with $s_a, t_b \geq 0$ the band-restricted spectral weights, and the Σ-normalizations

$$\langle\!\langle \hat A_a, \hat A_{a'}\rangle\!\rangle_{\Sigma,\Phi} = \delta_{aa'}, \qquad \langle\!\langle \hat K_b, \hat K_{b'}\rangle\!\rangle_{\Sigma,\Theta} = \delta_{bb'}. \tag{9.18}$$

We call $\hat A_a^\dagger$ *Apeiron* creators (Gaiamiti sector) and $\hat K_b^\dagger$ *Kymon* creators (Tantrakyma sector).

Lemma 9.38 (Phase–vs–amplitude split via Σ)**.** *Let $v \in T_x\mathcal{X}_q$ and $\hat Q_x[v] = v^\mu v^\nu \hat\Sigma_{\mu\nu}(x)$ as in Definition 9.22, with $\hat\Sigma_{\mu\nu} = \hat g_{\mu\nu} + \hat t_{\mu\nu}$ (cf. Definition 9.20). If $\bar g_{\mu\nu}(x)\, v^\mu v^\nu = 0$ while $\bar t_{\mu\nu}(x)\, v^\mu v^\nu > 0$, then the associated principal directions in $\{u_a\}$ (geometry block) obtained from Proposition 9.36 are* pure phase–flow *(Goldstone–like) precursors. If instead $\bar g_{\mu\nu}(x)\, v^\mu v^\nu > 0$ dominates the contraction of $\bar\Sigma_{\mu\nu}$ along v, these principal directions are* amplitude *(Higgs/Gaiaton–like). An analogous statement holds for the matter block $\{v_b\}$.*

Proof. (1) Directional contractions and Σ–weighted Rayleigh quotients. Contract the chrono-geometric tensors along v:

$$\bar g[v] := \bar g_{\mu\nu} v^\mu v^\nu, \qquad \bar t[v] := \bar t_{\mu\nu} v^\mu v^\nu, \qquad \bar\Sigma[v] = \bar g[v] + \bar t[v],$$

and analogously $\hat\Sigma[v]$ on the band (cf. Definitions 9.20 and 9.22). On the geometry block, the generalized eigenproblem of Proposition 9.36 admits the standard Rayleigh characterization with the Σ–inner product:

$$\mathcal{R}_\Phi(u) \;:=\; \frac{\langle u,\, C_\Phi u\rangle_{\Sigma^{(\Phi)}}}{\langle u, u\rangle_{\Sigma^{(\Phi)}}} \;=\; \frac{\mathcal{I}\Big[\,(\delta_X \hat G_{\Phi\Phi} u)^\dagger\, \hat\Sigma^{(\Phi)}\, (\delta_X \hat G_{\Phi\Phi} u)\Big]}{\mathcal{I}\Big[\,u^\dagger\, \hat\Sigma^{(\Phi)} u\Big]},$$

where $\mathcal{I}$ is any band–local, gauge–invariant functional (Born pairing / regulated band–trace; cf. Definitions 1.18, 1.34 and 9.23). Directional probing along v amounts to replacing $\hat{\Sigma}^{(\Phi)}$ by its contraction $\hat{\Sigma}^{(\Phi)}[v] = v^\mu v^\nu \hat{\Sigma}^{(\Phi)}_{\mu\nu}$ in both numerator and denominator (cf. Definition 9.22); the corresponding expectation–level quotient reads

$$\overline{\mathcal{R}}^{[v]}_{\Phi}(u) \;=\; \frac{\mathcal{I}\Big[\,(\delta_X \hat{G}_{\Phi\Phi} u)^\dagger\, \hat{\Sigma}^{(\Phi)}[v]\, (\delta_X \hat{G}_{\Phi\Phi} u)\Big]}{\mathcal{I}\Big[\,u^\dagger\, \hat{\Sigma}^{(\Phi)}[v]\, u\Big]}\,,$$

and its stationary points select the same principal directions $\{u_a\}$ (the weighting changes but the Euler–Lagrange equations coincide with Proposition 9.36 after contraction).

(2) Phase–flow regime ($\bar{g}[v] = 0$, $\bar{t}[v] > 0$). If $\bar{g}[v] = 0$ while $\bar{t}[v] > 0$, then $\bar{\Sigma}[v] = \bar{t}[v]$ and the directional inner product reduces to the temporal piece:

$$\langle u, u\rangle_{\Sigma^{(\Phi)}[v]} = \mathcal{I}\Big[u^\dagger\, \hat{t}^{(\Phi)}[v]\, u\Big], \qquad \hat{t}^{(\Phi)}[v] := v^\mu v^\nu \hat{t}^{(\Phi)}_{\mu\nu}(x).$$

Thus the metric contribution to the directional norm vanishes, and the response measured by $\overline{\mathcal{R}}^{[v]}_{\Phi}$ is purely *phase–flow* (temporal–curvature) in the sense that the geometric distance part registers no cost along v (cf. the chronogeometric split and its interpretation in Section 5.2.5 and Definition 5.122). In the QII setting (law–symmetric, state–asymmetric), such tangential directions coincide with Goldstone–like modes of the geometric vacuum (Section 9.17.2), and after plateau/BRST realization they furnish the spin–2 graviton as the massless Goldstone direction (Section 7.5 with the BRST/Nielsen framework). Hence the principal directions singled out by $\overline{\mathcal{R}}^{[v]}_{\Phi}$ in this regime are *pure phase–flow* precursors.

(3) Amplitude regime ($\bar{g}[v] > 0$ dominates). If $\bar{g}[v] > 0$ dominates the contraction of $\bar{\Sigma}[v] = \bar{g}[v] + \bar{t}[v]$, then the directional norm $\langle u, u\rangle_{\Sigma^{(\Phi)}[v]} = \mathcal{I}[u^\dagger(\hat{g}^{(\Phi)}[v] + \hat{t}^{(\Phi)}[v])u]$ is controlled by the metric part $\hat{g}^{(\Phi)}[v]$. Geometrically this corresponds to *radial* (distance–changing) deformations in the Σ–geometry, i.e. amplitude modes. On the Lagrangian side these are the directions that probe the curvature of the Gaiamiti potential $V(\hat{\mathbf{\Phi}})$ (cf. Definitions 6.172 and 6.174), and upon realization give massive Gaiaton excitations rather than Goldstone transport (cf. interpretation via Section 9.5 and Proposition 9.36).

(4) Matter block. The same argument, with $\hat{\Sigma}^{(\Theta)}$ and C_Θ in Proposition 9.36, yields the analogous classification for principal directions $\{v_b\}$ in the Tantrakyma block, identifying phase–flow versus amplitude behavior with $\bar{g}[v]$ versus $\bar{t}[v]$ dominance under the Σ–weighted quotient (cf. Definitions 6.4 and 6.142).

(5) Covariance and band–regularity. All directional contractions and Rayleigh quotients are band–local, gauge/frame covariant and finite by Propositions 9.6, 9.13 and 9.14 and Definition 9.10, and are stable under small band deformations by Lemma 9.27. Therefore the phase–vs–amplitude classification is well–posed on the band and persists to the realized physical layer after plateau/BRST projection (cf. Sections 6.10 and 9.5).

The stated alternatives follow. □

Proposition 9.39 (Channel mixing and block diagonality)**.** *If $[\hat{\Sigma}^{(\Phi)}, \delta_X \hat{G}_{\Phi\Phi}] = 0$ (respectively, $[\hat{\Sigma}^{(\Theta)}, \delta_X \hat{G}_{\Theta\Theta}] = 0$), the generalized eigenproblem (9.16) decouples into channel–wise eigenmodes (corr/vac/spec). Otherwise, the cross–terms are absorbed by the generalized eigenproblem, and the principal modes carry the correct Σ–weights across mixed channels.*

Proof. Setup and projectors. On the band (cf. Definitions 6.280 and 9.10) the chronogeometry admits the triadic split $\hat{\Sigma}^{(\Phi)} = \hat{\Sigma}^{(\Phi,\mathrm{corr})} + \hat{\Sigma}^{(\Phi,\mathrm{vac})} + \hat{\Sigma}^{(\Phi,\mathrm{spec})}$, consistent with the expectation–level decomposition $\bar{\Sigma}_{\mu\nu} = \bar{\Sigma}^{\mathrm{corr}}_{\mu\nu} + \bar{\Sigma}^{\mathrm{vac}}_{\mu\nu} + \bar{\Sigma}^{\mathrm{spec}}_{\mu\nu}$ (cf. Definitions 5.54 and 9.20 and eqs. (5.8) and (5.21)). By the graded bundle structure and block algebra (cf. Definitions 9.7 and 9.8, eq. (4.24), and Section 4.1.7), there exist band–local channel projectors $\Pi^{(a)}$ ($a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$) such that

$$\Pi^{(a)}\Pi^{(b)} = \delta_{ab}\Pi^{(a)}, \qquad \sum_a \Pi^{(a)} = \mathbb{I}, \qquad \Pi^{(a)}\hat{\Sigma}^{(\Phi)}\Pi^{(b)} = \delta_{ab}\,\hat{\Sigma}^{(\Phi,a)}.$$

(Construction as in the channel attribution step of Proposition 9.29.)

Decoupled case. Assume $[\hat{\Sigma}^{(\Phi)}, \delta_X\hat{G}_{\Phi\Phi}] = 0$. Then $\delta_X\hat{G}_{\Phi\Phi}$ preserves the spectral decomposition of $\hat{\Sigma}^{(\Phi)}$, hence $[\Pi^{(a)}, \delta_X\hat{G}_{\Phi\Phi}] = 0$ for each a. Let $C_\Phi = (\delta_X\hat{G}_{\Phi\Phi})^\dagger\hat{\Sigma}^{(\Phi)}(\delta_X\hat{G}_{\Phi\Phi})$ (cf. Proposition 9.36). For $a \neq b$,

$$\begin{aligned}\Pi^{(a)}C_\Phi\Pi^{(b)} &= \Pi^{(a)}(\delta_X\hat{G}_{\Phi\Phi})^\dagger\,\underbrace{\hat{\Sigma}^{(\Phi)}\Pi^{(b)}}_{=\Pi^{(b)}\hat{\Sigma}^{(\Phi,b)}}(\delta_X\hat{G}_{\Phi\Phi})\Pi^{(b)}\\ &= (\delta_X\hat{G}_{\Phi\Phi})^\dagger\Pi^{(a)}\Pi^{(b)}\,\hat{\Sigma}^{(\Phi,b)}(\delta_X\hat{G}_{\Phi\Phi})\Pi^{(b)} = 0,\end{aligned}$$

using the commutation with $\Pi^{(a)}$ and orthogonality of the projectors. Thus C_Φ is block–diagonal in the channel decomposition: $C_\Phi = \bigoplus_a C_\Phi^{(a)}$ with $C_\Phi^{(a)} := (\delta_X\hat{G}_{\Phi\Phi})^\dagger\hat{\Sigma}^{(\Phi,a)}(\delta_X\hat{G}_{\Phi\Phi})$ acting on $\Pi^{(a)}\mathcal{N}_\Phi^{\mathrm{pre}}$. The generalized eigenproblem $C_\Phi u = \lambda\,\hat{\Sigma}^{(\Phi)}u$ therefore decouples into

$$C_\Phi^{(a)}\,u^{(a)} = \lambda^{(a)}\,\hat{\Sigma}^{(\Phi,a)}u^{(a)} \qquad (a = \mathrm{corr}, \mathrm{vac}, \mathrm{spec}),$$

i.e. three independent channel–wise problems with Σ–weights *within* each channel. The same argument applies to the Tantrakyma block under $[\hat{\Sigma}^{(\Theta)}, \delta_X\hat{G}_{\Theta\Theta}] = 0$.

Mixed case. If $[\hat{\Sigma}^{(\Phi)}, \delta_X\hat{G}_{\Phi\Phi}] \neq 0$, cross–terms $\Pi^{(a)}C_\Phi\Pi^{(b)}$ with $a \neq b$ need not vanish. However, by Proposition 9.36 (Steps 2–4), the Σ–weighted problem can be reduced to an ordinary compact eigenproblem for

$$K_\Phi := \hat{\Sigma}^{(\Phi)\,-1/2}C_\Phi\hat{\Sigma}^{(\Phi)\,-1/2},$$

which is positive, self–adjoint, and compact on the (unweighted) pre–Hilbert completion; its orthonormal eigenbasis $\{w_a\}$ yields the principal modes $u_a = \hat{\Sigma}^{(\Phi)\,-1/2}w_a$ that are Σ–orthonormal and solve $C_\Phi u_a = \lambda_a\hat{\Sigma}^{(\Phi)}u_a$. In the channel projector basis $\{\Pi^{(a)}\}$, mixing appears as off–diagonal blocks of K_Φ, but the spectral theorem diagonalizes K_Φ globally. Equivalently, the principal modes extremize the Σ–weighted Rayleigh quotient

$$\mathcal{R}_\Phi(u) = \frac{\langle u, C_\Phi u\rangle_{\Sigma^{(\Phi)}}}{\langle u, u\rangle_{\Sigma^{(\Phi)}}} = \frac{\mathcal{I}\Big[(\delta_X\hat{G}_{\Phi\Phi}u)^\dagger\hat{\Sigma}^{(\Phi)}(\delta_X\hat{G}_{\Phi\Phi}u)\Big]}{\mathcal{I}\Big[u^\dagger\hat{\Sigma}^{(\Phi)}u\Big]},$$

(cf. Proposition 9.36), which is basis–independent and automatically accounts for channel mixing via the Σ–weights in numerator and denominator. Channel contributions may still be attributed a posteriori using the functionals $\mathcal{I}^{(a)}$ of Proposition 9.29, but the eigenvectors themselves are generally mixed across channels when the commutator is nonzero.

Covariance and band–regularity. All statements are band–local and gauge/frame covariant by construction (cf. Definitions 9.7 and 9.10 and Proposition 9.6), and the compactness/positivity needed for the spectral reduction follow from spectral discipline (cf. Proposition 9.13 and Remark 9.21) and from the block restrictions of $\hat{\Sigma}$ (cf. Definition 9.20). Thus the generalized eigenproblem absorbs channel mixing and yields Σ–orthonormal principal modes with the correct weights.

The claim follows. □

Corollary 9.40 (Recovery of SVD-based construction)**.** *If $\hat{\Sigma}^{(\Phi)} \propto \mathbb{I}$ (resp. $\hat{\Sigma}^{(\Theta)} \propto \mathbb{I}$) on the band or if only the correlator channel is active, (9.16) reduces to the usual SVD for $\delta_X \hat{G}_{\Phi\Phi}$ (resp. $\delta_X \hat{G}_{\Theta\Theta}$), and (9.17) matches the factorization used in the correlator–only scheme.*

Proof. (1) Reduction of the generalized problem to an ordinary eigenproblem. On the band (cf. Definitions 6.280 and 9.10) define $A_\Phi := \delta_X \hat{G}_{\Phi\Phi}$ and recall $C_\Phi = A_\Phi^\dagger \hat{\Sigma}^{(\Phi)} A_\Phi$ with the generalized eigenproblem $C_\Phi u = \lambda \hat{\Sigma}^{(\Phi)} u$ (cf. Proposition 9.36 and eqs. (9.15) and (9.16)). Let $J_\Phi := \hat{\Sigma}^{(\Phi)\,1/2}$ (well–defined and positive on the band by Definition 9.20 and Propositions 9.13 and 9.14) and set

$$K_\Phi := J_\Phi^{-1} C_\Phi J_\Phi^{-1} = \big(J_\Phi A_\Phi\big)^\dagger \big(J_\Phi A_\Phi\big)$$

on the unweighted pre–Hilbert completion (cf. Proposition 9.36).

Then $K_\Phi w = \lambda w$ with $w := J_\Phi u$ is an *ordinary* compact, positive, self–adjoint eigenproblem (spectral theorem applies; cf. Proposition 9.36).

(2) Case $\hat{\Sigma}^{(\Phi)} = s_\Phi \mathbb{I}$. If $\hat{\Sigma}^{(\Phi)} = s_\Phi \mathbb{I}$ on the band (e.g. within a single active channel with appropriate normalization), then $J_\Phi = \sqrt{s_\Phi}\,\mathbb{I}$ and

$$K_\Phi = s_\Phi\, A_\Phi^\dagger A_\Phi.$$

Hence $K_\Phi w = \lambda w$ is equivalent to $A_\Phi^\dagger A_\Phi w = (\lambda/s_\Phi)\, w$: the eigenvectors $\{w_a\}$ are the *right singular vectors* of $A_\Phi = \delta_X \hat{G}_{\Phi\Phi}$ and $\sigma_a^2 = \lambda_a / s_\Phi$ are the singular values squared. Writing the SVD of A_Φ as $A_\Phi = \sum_a \sigma_a\, |p_a\rangle\langle w_a|$ with $\{p_a\}$ orthonormal (left singular vectors) yields the weighted factorization

$$\delta_X \hat{G}_{\Phi\Phi} = A_\Phi = \sum_a \sigma_a\, |p_a\rangle\langle w_a| = \sum_a \sqrt{\lambda_a}\ \underbrace{|p_a\rangle}_{\text{left modes}}\ \underbrace{\langle u_a|\,\hat{\Sigma}^{(\Phi)}}_{\text{right }\Sigma\text{-dual}}, \qquad u_a := J_\Phi^{-1} w_a = \tfrac{1}{\sqrt{s_\Phi}}\, w_a,$$

which is precisely the Σ–weighted block factorization (9.17) when $\hat{\Sigma}^{(\Phi)} \propto \mathbb{I}$. Gauge/frame covariance and band–locality are inherited from the graded transport and the tag (cf. Definitions 9.7 and 9.10 and Proposition 9.6).

(3) Correlator–only activation. If only the correlator channel is active, channel projectors $\Pi^{(a)}$ (cf. Proposition 9.29) reduce the problem to the correlator block: $\hat{\Sigma}^{(\Phi)} \to \hat{\Sigma}^{(\Phi,\mathrm{corr})}$ and $A_\Phi \to A_\Phi^{(\mathrm{corr})}$, while mixing is absent by construction (cf. Proposition 9.39). With the fibre normalization $N(\psi, x)$ of Definition 9.19, the correlator–channel inner product is the

standard L^2–type pairing, so $\hat{\Sigma}^{(\Phi,\mathrm{corr})} \propto \mathbb{I}$ on the band. Thus Steps (1)–(2) apply verbatim and one recovers the usual SVD of $\delta_X \hat{G}_{\Phi\Phi}$ restricted to the correlator channel, i.e. the original correlator–only construction consistent with Corollary 9.30 and Definition 4.338.

(4) Tantrakyma block. Exactly the same argument applies to the matter block with $A_\Theta := \delta_X \hat{G}_{\Theta\Theta}$ and $\hat{\Sigma}^{(\Theta)}$ (cf. eqs. (4.24), (9.15) and (9.16) and Definition 6.4), yielding the SVD of $\delta_X \hat{G}_{\Theta\Theta}$ when $\hat{\Sigma}^{(\Theta)} \propto \mathbb{I}$ or when only the correlator channel is active.

Therefore, whenever the Σ–weight is proportional to the identity (globally or within the active channel), the generalized eigenproblems (9.16) reduce to ordinary SVDs for $\delta_X \hat{G}_{\Phi\Phi}$ and $\delta_X \hat{G}_{\Theta\Theta}$, and the Σ–weighted block factorization (9.17) coincides with the correlator–only factorization. □

Remark 9.41 (Chrono–Kymon and inter–modal mixing)**.** Within the Tantrakyma (matter) block of the Fourfold correlator (cf. Definition 6.4 and eq. (4.24)), the Σ–tensor restricts to $\hat{\Sigma}^{(\Theta)}$ (cf. Definition 9.20). Isolating its temporal contribution $\hat{t}^{(\Theta)}_{\mu\nu}$—the "torsional" sector obtained from the SOT–C^2 Hessians of temporal functionals (cf. eq. (9.4) and Definition 5.17)—singles out *spin–1 Chrono–Kymon* channels at the pre–physical level. Operationally, probing along $v \in T_x\mathcal{X}_q$ replaces the Σ–weight by its contraction $\hat{\Sigma}^{(\Theta)}[v] = v^\mu v^\nu \hat{\Sigma}^{(\Theta)}_{\mu\nu}(x)$ (cf. Definition 9.22); when $\bar{t}_{\mu\nu} v^\mu v^\nu$ dominates, the principal directions selected by the Σ–weighted Rayleigh quotient are temporal/phase–flow–type in the matter block, i.e. Chrono–Kymon precursors (cf. the phase–vs–amplitude criterion in Lemma 9.38).

Inter–modal entries $\theta^{\alpha\beta}$ inside $\hat{G}_{\Theta\Theta}$ (off–diagonal in the $\{0,1,p,s\}$ grading; cf. Definition 9.8 and Section 4.1.7) control mixing across parasymmetric/holosymmetric channels. Their influence is incorporated *faithfully* by the Σ–weighted generalized eigenproblem

$$(\delta_X \hat{G}_{\Theta\Theta})^\dagger \, \hat{\Sigma}^{(\Theta)} \, (\delta_X \hat{G}_{\Theta\Theta}) \, v_b = \mu_b \, \hat{\Sigma}^{(\Theta)} \, v_b,$$

(cf. Proposition 9.36 and eqs. (9.15) and (9.16)), which is frame– and gauge–covariant by construction (cf. Definition 9.7 and Proposition 9.6) and band–regular (cf. Definition 9.10 and Proposition 9.13). If $[\hat{\Sigma}^{(\Theta)}, \delta_X \hat{G}_{\Theta\Theta}] = 0$, channel projectors $\Pi^{(a)}$ (corr/vac/spec) block–diagonalize the problem and the modes decouple channelwise (cf. Proposition 9.39); otherwise, off–diagonal inter–modal terms are absorbed by the compact, positive operator $K_\Theta = \hat{\Sigma}^{(\Theta)\,-1/2} (\delta_X \hat{G}_{\Theta\Theta})^\dagger \hat{\Sigma}^{(\Theta)} (\delta_X \hat{G}_{\Theta\Theta}) \hat{\Sigma}^{(\Theta)\,-1/2}$, whose spectral theorem furnishes Σ–orthonormal principal modes with the correct temporal weights across mixed channels (cf. Propositions 9.29 and 9.36). In the correlator–only limit or when $\hat{\Sigma}^{(\Theta)} \propto \mathbb{I}$ on the band, one recovers the usual SVD–based construction and block factorization (cf. Corollaries 9.30 and 9.40).

After plateau certification and BRST projection, Chrono–Kymon channels are realized physically only when the Θ–block retarded kernels develop Nielsen–stable, gauge–parameter–independent spin–1 poles with positive residues in $H^0(\hat{Q}_{\mathrm{BRST}})$ (cf. Sections 6.10 and 9.5); the inter–modal mixing diagnosed pre–physically then persists as physical mixing consistent with Ward/Slavnov identities on $\mathcal{H}_{\mathrm{phys}}$.

Lemma 9.42 (Stability under band deformations)**.** *Under the continuity bounds of Lemma 9.27, the principal eigenvalues λ_a, μ_b and the Σ–normalizations of the mode operators $\hat{A}_a, \hat{K}_b$ (from the block factorizations) vary continuously with the band parameters $(\lambda_{\mathrm{UV}}, \ell)$. In particular, for $(\lambda_{\mathrm{UV}}, \ell) \mapsto (\lambda'_{\mathrm{UV}}, \ell')$ inside a plateau neighborhood,*

$$|\lambda'_a - \lambda_a| + |\mu'_b - \mu_b| = \mathcal{O}(\varepsilon), \qquad \|\hat{A}'_a - \hat{A}_a\| + \|\hat{K}'_b - \hat{K}_b\| = \mathcal{O}(\varepsilon),$$

with the $\mathcal{O}(\varepsilon)$ control provided by the effective cutoff $\Lambda_{\mathrm{eff}}(x,x';\varepsilon)$ of Remark 9.21; degeneracy splittings are likewise $\mathcal{O}(\varepsilon)$.

Proof. Step 1 (reduce the generalized problems to compact self–adjoint ones). For the geometry block, set $A_\Phi := \delta_X \hat{G}_{\Phi\Phi}$ and $J_\Phi := \hat{\Sigma}^{(\Phi)\,1/2}$ on the band. By Proposition 9.36 and eqs. (9.15) and (9.16), the generalized problem $C_\Phi u = \lambda\, \hat{\Sigma}^{(\Phi)} u$ with $C_\Phi = A_\Phi^\dagger \hat{\Sigma}^{(\Phi)} A_\Phi$ is equivalent to the ordinary compact, positive, self–adjoint eigenproblem

$$K_\Phi w = \lambda\, w, \qquad K_\Phi := J_\Phi^{-1} C_\Phi J_\Phi^{-1} = (J_\Phi A_\Phi)^\dagger (J_\Phi A_\Phi),$$

with $w := J_\Phi u$. Exactly the same reduction holds for the matter block with $K_\Theta := J_\Theta^{-1} C_\Theta J_\Theta^{-1}$, $J_\Theta := \hat{\Sigma}^{(\Theta)\,1/2}$ (cf. Proposition 9.36).

Step 2 (operator–norm continuity under band deformations). By Lemma 9.27, small changes in $(\lambda_{\mathrm{UV}}, \ell)$ induce $\mathcal{O}(\varepsilon)$ variations of the Σ–tensor: $\|\hat{\Sigma}' - \hat{\Sigma}\| = \mathcal{O}(\varepsilon)$ on the band. The band tag $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ depends continuously on $(\lambda_{\mathrm{UV}}, \ell)$ (UV spectral projector and IR window; cf. Definitions 6.280 and 9.10 and eqs. (7.130) and (9.2)), hence the band–limited kernels of $\hat{G}_{\bullet\bullet}$, and therefore of $\delta_X \hat{G}_{\bullet\bullet}$ (graded derivation; cf. Definitions 9.7 and 9.18), vary in operator norm by $\mathcal{O}(\varepsilon)$, controlled by Λ_{eff} (cf. Remark 9.21 and Proposition 9.13). Functional calculus for positive operators gives $\|J'_\Phi - J_\Phi\| = \mathcal{O}(\varepsilon)$ and $\|J'^{-1}_\Phi - J_\Phi^{-1}\| = \mathcal{O}(\varepsilon)$ on the support of $\hat{\Sigma}^{(\Phi)}$ (positivity and band regularity: Definition 9.20 and Proposition 9.14). Consequently,

$$\|K'_\Phi - K_\Phi\| \;\le\; \|(J'_\Phi A'_\Phi)^\dagger (J'_\Phi A'_\Phi) - (J_\Phi A_\Phi)^\dagger (J_\Phi A_\Phi)\| \;=\; \mathcal{O}(\varepsilon),$$

and similarly $\|K'_\Theta - K_\Theta\| = \mathcal{O}(\varepsilon)$.

Step 3 (eigenvalue/eigenspace continuity and splittings). For compact self–adjoint operators, eigenvalues (counting multiplicities) are 1–Lipschitz with respect to operator–norm perturbations: $|\lambda'_a - \lambda_a| \le \|K'_\Phi - K_\Phi\| = \mathcal{O}(\varepsilon)$ and similarly for μ_b (Weyl–Kato continuity; the needed compactness/positivity came from Step 1). If an eigenvalue is degenerate, the spectral projector

$$\mathsf{P}_\Phi(\Gamma) = \frac{1}{2\pi i} \oint_\Gamma (z - K_\Phi)^{-1}\, dz$$

onto the cluster (for a small contour Γ separating it from the rest of the spectrum) varies by $\|\mathsf{P}'_\Phi - \mathsf{P}_\Phi\| \le c\, \|K'_\Phi - K_\Phi\| = \mathcal{O}(\varepsilon)$; thus splittings inside the cluster are $\mathcal{O}(\varepsilon)$. Identical statements hold for K_Θ.

Step 4 (Σ–orthonormal eigenvectors and normalized mode operators). Choose Σ–orthonormal principal vectors by $u_a = J_\Phi^{-1} w_a$ and $v_b = J_\Theta^{-1} z_b$ (cf. Proposition 9.36). Since $J_\bullet$ and the spectral projectors depend continuously on $(\lambda_{\mathrm{UV}}, \ell)$, one may select continuous local frames inside each isolated (or ε–narrow) cluster so that $\|u'_a - u_a\| + \|v'_b - v_b\| = \mathcal{O}(\varepsilon)$. The Σ–normalizations $\langle u_a, u_{a'} \rangle_{\Sigma^{(\Phi)}} = \delta_{aa'}$ and $\langle v_b, v_{b'} \rangle_{\Sigma^{(\Theta)}} = \delta_{bb'}$ are preserved by construction and remain stable because $\|\hat{\Sigma}' - \hat{\Sigma}\| = \mathcal{O}(\varepsilon)$ (cf. Definition 9.20). The creation operators $\hat{A}_a, \hat{K}_b$ entering the block factorizations (Apeiron/Kymon modes) are linear functionals of $A_\bullet$ and the principal vectors (cf. Proposition 9.36 and the Σ–weighted block factorizations eq. (9.17)); hence $\|\hat{A}'_a - \hat{A}_a\| + \|\hat{K}'_b - \hat{K}_b\| = \mathcal{O}(\varepsilon)$.

Combining Steps 1–4 yields the continuity and $\mathcal{O}(\varepsilon)$ stability claims for eigenvalues, eigenvectors, and Σ–normalized mode operators under band deformations controlled by $\Lambda_{\mathrm{eff}}(x,x';\varepsilon)$ (cf. Remark 9.21 and Lemma 9.27). □

Algorithm 1 Σ-aware block-wise mode extraction (pre-physical)

1. Fix band $(\lambda_{\mathrm{UV}}, \ell)$ (Definition 9.10) and build $\hat{\Sigma}_{\mu\nu}$ (Definition 9.20); restrict to $\hat{\Sigma}^{(\Phi)}, \hat{\Sigma}^{(\Theta)}$.
2. Identify broken generators $\mathfrak{B}$ via Σ-contrasts (Definitions 9.23 and 9.28).
3. For each $X \in \mathfrak{B}$, compute $\delta_X \hat{G}_{\Phi\Phi}$, $\delta_X \hat{G}_{\Theta\Theta}$ (Definition 9.32) and solve (9.16).
4. Normalize principal modes by (9.18); assemble $\hat{A}_a^\dagger, \hat{K}_b^\dagger$ and the factorizations (9.17).
5. Record phase/amplitude character using Lemma 9.38; check stability via Lemma 9.42.

Remark 9.43 (Pre-physical status and realization)**.** All constructions above—band tagging, Σ–weighted Gram operators, generalized eigenproblems, principal modes, and their Chrono–Kymon specializations—are posed on the emergent/GNS arena $\mathcal{H}_{\mathrm{em}}$ and are therefore *pre–physical* (cf. Definitions 1.34, 9.10 and 9.20, Proposition 9.36, and Remark 9.41). They provide PSR–level diagnostics and mode bases but do not by themselves assert particle content (cf. Definitions 9.12, 9.23 and 9.26 and Lemma 9.27). Physical poles and residues identifying graviton/Gaiaton/Kymon sectors arise only after the Σ–aware realization step: certify a plateau via the band grid diagnostics and acceptance rule (cf. eqs. (7.394) to (7.396) and Proposition 7.203), then apply the realization map and BRST projection to the cohomology $\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}})$ (cf. Sections 6.10 and 9.5, Lemma 9.11, and eq. (7.24)). Only at this stage are Σ–causal retarded kernels evaluated on H^0 and tested for Nielsen–stable, gauge–parameter–independent poles with positive residues (Ward/Slavnov identities unchanged, now phrased in Σ–compatible form; cf. Proposition 9.6 and Section 7.5). Thus the present eigenmode analysis is a necessary upstream input to the realized spectrum but is *not* yet a statement about particles until the Σ–realized, BRST–physical evaluation is performed.

Summary of Section 9.3. Block–wise mode extraction has been upgraded from a plain SVD to the Σ–weighted generalized eigenproblem (9.16), built from the Gram operators (9.15) and solved in the Σ–pre–Hilbert modules of the geometry and matter blocks (cf. Proposition 9.36). This yields Σ–orthonormal principal systems $\{u_a\} \subset \mathcal{N}_\Phi^{\mathrm{pre}}$ and $\{v_b\} \subset \mathcal{N}_\Theta^{\mathrm{pre}}$ with non–negative spectra, and furnishes *Apeiron* and *Kymon* creators via the Σ–weighted block factorizations (cf. eq. (9.17)). The construction is band–regular and gauge/frame covariant by design: band–locality and causal support come from the tag $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ and its IR window (cf. Definitions 6.280 and 9.10 and eqs. (7.130) and (9.2)), while covariance is ensured by graded transport and the Hermiticity/covariance of the Fourfold correlator (cf. Definitions 6.4 and 9.7, Proposition 9.6, and eq. (4.24)); finiteness and smoothness on the band follow from spectral discipline (cf. Propositions 9.13 and 9.14 and Remark 9.21).

Crucially, the Σ–weight makes the principal directions intrinsically sensitive to both *amplitude curvature* $\hat{g}$ and *temporal phase–flow* $\hat{t}$ through the chronogeometric quadratic form $\hat{Q}_x[v] = v^\mu v^\nu \hat{\Sigma}_{\mu\nu}(x)$ (cf. Definitions 9.20 and 9.22). The phase–vs–amplitude criterion *along a probe direction* v is encoded by the directional contractions of $\bar{g}$ and $\bar{t}$ and is formalized in Lemma 9.38; in the Tantrakyma block, isolating the temporal torsional

sector identifies spin–1 *Chrono–Kymon* channels and their inter–modal mixing is faithfully captured by the Σ–weighted problem (cf. Remark 9.41). Channel mixing vs. decoupling is governed by the commutator $[\hat{\Sigma}^{(\bullet)}, \delta_X \hat{G}_{\bullet\bullet}]$; exact commutation implies block diagonality and channel–wise eigenproblems, whereas non–commutation is absorbed by the compact, positive operator $K_\bullet = \hat{\Sigma}^{(\bullet)\,-1/2}(\delta_X \hat{G}_{\bullet\bullet})^\dagger \hat{\Sigma}^{(\bullet)}(\delta_X \hat{G}_{\bullet\bullet})\hat{\Sigma}^{(\bullet)\,-1/2}$ with globally diagonalizable spectrum (cf. Propositions 9.36 and 9.39). Channelwise interpretability and attribution to coherence/vacuum/spectral–shear contributions are obtained by the triadic split of Σ and the corresponding projectors/functionals (cf. Proposition 9.29, Definition 5.54, and eqs. (5.8) and (5.21)).

In regimes where $\hat{\Sigma}^{(\bullet)} \propto \mathbb{I}$ on the band—or when only the correlator channel is active—the generalized problem reduces to an ordinary SVD for $\delta_X \hat{G}_{\bullet\bullet}$ and the Σ–weighted factorization matches the correlator–only construction (cf. Corollaries 9.30 and 9.40 and Definitions 4.338 and 9.19). Stability under band deformations is guaranteed: the contrasts are continuous and $\mathcal{O}(\varepsilon)$–stable (Lemma 9.27), which lifts to continuity of principal eigenvalues/eigenspaces and Σ–normalized mode operators with $\mathcal{O}(\varepsilon)$ degeneracy splittings controlled by Λ_{eff} (cf. Lemma 9.42). Finally, all results here are *pre–physical* on $\mathcal{H}_{\text{em}}$ (cf. Remark 9.43); they become BRST–physical diagnostics only after Σ–aware realization—plateau certification and projection to $\mathcal{H}_{\text{phys}} := H^0(\hat{Q}_{\text{BRST}})$—where Ward/Slavnov identities and Nielsen stability are evaluated without altering the downstream cohomological machinery (cf. Sections 6.10, 7.5 and 9.5).

9.4 Pre-physical Symmetry Reduction (PSR)

Definition 9.44 (Law–kernel functionals and band-tagged diagnostics)**.** Let $\mathfrak{L}$ be a graded–covariant *law–kernel functional* acting on the Fourfold correlator $\hat{\mathbf{G}}_\Lambda$ (cf. Definitions 6.4 and 9.5 and eq. (4.24)) and transforming under the graded adjoint as in Definition 9.18. Such functionals may be assembled from band–projected kernels (e.g. operator kernels playing the role of $\mathcal{K}$, $\mathcal{M}^2$ in a given symmetric law) using the gauge–covariant transport $\mathcal{U}$ (cf. Definitions 6.184, 6.192 and 9.7). On a band $(\lambda_{\text{UV}}, \ell)$, define the band–tagged diagnostic by

$$\mathfrak{L}_{\text{pre}} := \mathfrak{B}_{\lambda_{\text{UV}},\ell}\Big[\mathfrak{L}(\hat{\mathbf{G}}_\Lambda)\Big],$$

with $\mathfrak{B}_{\lambda_{\text{UV}},\ell}$ the UV/IR tag (cf. Definitions 6.280 and 9.10 and eqs. (7.130) and (9.2)). Expectations are taken in the Holo–State/GNS sense, $\omega^\Psi(\cdot) = \langle\Psi|\cdot|\Psi\rangle$ (cf. Definitions 1.18 and 1.34). By Hermiticity/covariance of $\hat{\mathbf{G}}_\Lambda$ and endpoint covariance of $\mathcal{U}$ (cf. Proposition 9.6), $\mathfrak{L}$ is graded–adjoint covariant and $\omega^\Psi(\mathfrak{L}_{\text{pre}})$ is gauge–invariant; frame independence follows from the conjugation invariance of the functionals (cf. Sections 4.1.7 and 4.19.1).

Definition 9.45 (Chronogeometric quadratic probe and Σ–contrasts)**.** Let $\hat{\Sigma}_{\mu\nu} = \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu}$ be the operator–valued chronogeometric tensor on the band (cf. Definition 9.20), and for a local probe $v \in T_x\mathcal{X}_q$ set $\hat{Q}_x[v] := v^\mu v^\nu \hat{\Sigma}_{\mu\nu}(x)$ (cf. Definition 9.22). Fix a gauge–invariant, band–local functional $\mathcal{I}$ (Holo–State Born pairing or regulated band–trace; cf. Definitions 1.18, 1.34, 6.280 and 9.10). For $g \in G$, define the pre–physical Σ–law and Σ–state contrasts

(cf. Definition 9.23):

$$\Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}}(g;x;v) := \Big(\mathcal{I}\big[\hat{Q}_x[v] - \mathrm{Ad}_{g(x)}\,\hat{Q}_x[v]\big]\Big)^{1/2}, \tag{9.19}$$

$$\Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}}(g;x;v) := \Big|\mathcal{I}_\Psi\big[\hat{Q}_x[v]\big] - \mathcal{I}_{g\cdot\Psi}\big[\hat{Q}_x[v]\big]\Big|^{1/2}. \tag{9.20}$$

By gauge/frame covariance of $\hat{Q}_x[v]$ and conjugation invariance of $\mathcal{I}$ and $\mathcal{I}_\Psi$ (cf. Proposition 9.6, Definition 9.7, and Section 4.19.1), these quantities are frame–independent and respect the graded G–action of Definition 9.18; see Lemma 9.24. Aggregated versions over finite probe sets $\{v_k\}$ are given in Definition 9.26, and stability under band deformations is ensured by Lemma 9.27. Channelwise attributions and correlator–only reductions follow from Proposition 9.29 and Corollary 9.30.

Lemma 9.46 (Band regularity and finiteness)**.** *Under Spectral Smoothness and Finite Entanglement Capacity, the triads and $\hat{\Sigma}$ exist on $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ and render (9.19)–(9.20) finite and continuous in v and in the band knobs $(\lambda_{\mathrm{UV}},\ell)$.*

Proof. Existence and finiteness on the band. The correlator–based triads D^2, T^2 (and their channel refinements $d_{\mathrm{corr}}, d_{\mathrm{vac}}, d_{\mathrm{spec}}$) are defined via the spectral generator with self–regularizing properties (cf. Definitions 4.338, 4.341, 4.386, 4.428 and 4.463). Spectral Smoothness together with the Principle of Finite Entanglement Capacity imply band–level finiteness and smoothness of these functionals and of the operator fields derived from their SOT–C^2 coincident Hessians, namely $\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu}$ and $\hat{\Sigma}_{\mu\nu} = \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu}$ (cf. Propositions 9.13 and 9.14, Definition 9.20, and eq. (9.4)). Band–tagging by the UV projector and causal IR window (cf. Definitions 6.280 and 9.10 and eqs. (7.130) and (9.2)) preserves gauge/transport covariance (cf. Proposition 9.6 and Definition 9.7) and yields a band–regular $\hat{\Sigma}$.

Finiteness of the Σ–contrasts. Fix x and $v \in T_x\mathcal{X}_q$. The quadratic probe is $\hat{Q}_x[v] = v^\mu v^\nu \hat{\Sigma}_{\mu\nu}(x)$ (cf. Definition 9.22). With $\|\cdot\|$ the operator norm on the band fibre,

$$\|\hat{Q}_x[v]\| \;\le\; \|\hat{\Sigma}(x)\|\,\|v\|^2,$$

so $\hat{Q}_x[v]$ is bounded for fixed v. Any admissible band–local, gauge–invariant functional $\mathcal{I}$ (Born pairing or regulated band–trace; cf. Definitions 1.18 and 1.34) satisfies $|\mathcal{I}[X]| \le C\,\|X\|$ for some $C < \infty$. Hence the law contrast and the state contrast in (9.19)–(9.20) are finite.

Continuity in v. For $v, w \in T_x\mathcal{X}_q$,

$$\|\hat{Q}_x[v] - \hat{Q}_x[w]\| = \|v^\mu v^\nu - w^\mu w^\nu\|\,\|\hat{\Sigma}_{\mu\nu}(x)\| \;\le\; \|\hat{\Sigma}(x)\|\,(\|v\| + \|w\|)\,\|v - w\|,$$

so $v \mapsto \hat{Q}_x[v]$ is (locally) Lipschitz. By continuity of $\mathcal{I}$ and of the push–forward $g\cdot\Psi$ (cf. Definition 9.18 and Section 4.19.1), the maps $v \mapsto \Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}}(g;x;v)$ and $v \mapsto \Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}}(g;x;v)$ are continuous (cf. Definition 9.45).

Continuity in $(\lambda_{\mathrm{UV}},\ell)$. By Remark 9.21, for every $\varepsilon > 0$ there exists an effective cutoff $\Lambda_{\mathrm{eff}}(x,x';\varepsilon)$ such that changing $(\lambda_{\mathrm{UV}},\ell)$ inside the plateau neighborhood alters band–limited kernels by at most $\mathcal{O}(\varepsilon)$ in operator norm. In particular, $\|\hat{\Sigma}' - \hat{\Sigma}\| = \mathcal{O}(\varepsilon)$ and thus

$$\|\hat{Q}'_x[v] - \hat{Q}_x[v]\| \;\le\; \|v\|^2\,\|\hat{\Sigma}' - \hat{\Sigma}\| \;=\; \mathcal{O}(\varepsilon),$$

uniformly on bounded v–sets. Applying the bounded functional $\mathcal{I}$ (and its state–dependent version $\mathcal{I}_\Psi$) yields

$$\left|\left(\Delta_{\mathrm{law}}^{\mathrm{pre},\Sigma}\right)^2 - \left(\Delta_{\mathrm{law}}^{\mathrm{pre},\Sigma}\right)^2\right| \le C\,\|\hat{Q}'_x[v] - \hat{Q}_x[v]\| = \mathcal{O}(\varepsilon),$$

and likewise for the state contrast; taking square roots preserves continuity on compact v–domains. This agrees with the contrast–level stability already established in Lemma 9.27.

Combining the steps proves that, under Spectral Smoothness and Finite Entanglement Capacity, the triads and $\hat{\Sigma}$ exist on the band and render (9.19)–(9.20) finite and continuous in both v and $(\lambda_{\mathrm{UV}}, \ell)$. □

Definition 9.47 (Aggregated Σ–order parameters and visibilities)**.** Given probe directions $\{v_k\}_{k=1}^m \subset T_x\mathcal{X}_q$, aggregate the pointwise Σ–contrasts of Definition 9.45 (built from the quadratic probe $\hat{Q}_x[v] = v^\mu v^\nu \hat{\Sigma}_{\mu\nu}(x)$ of Definition 9.22) by the root–sum–square prescription

$$\Delta_{\mathrm{law}}^{\mathrm{pre},\Sigma}(g;x) := \left(\sum_{k=1}^m \left[\Delta_{\mathrm{law}}^{\mathrm{pre},\Sigma}(g;x;v_k)\right]^2\right)^{1/2}, \quad \Delta_{\mathrm{state}}^{\mathrm{pre},\Sigma}(g;x) := \left(\sum_{k=1}^m \left[\Delta_{\mathrm{state}}^{\mathrm{pre},\Sigma}(g;x;v_k)\right]^2\right)^{1/2}. \tag{9.21}$$

This is consistent with the general aggregation pattern of Definition 9.26. By Lemma 9.24, the aggregated quantities remain frame–independent and respect the graded G–action; by Lemmas 9.27 and 9.46, they are finite and continuous in both the probe set and the band parameters $(\lambda_{\mathrm{UV}}, \ell)$.

Define Σ–*visibilities* as complements (choose a normalization $C > 0$):

$$\Lambda_{\mathrm{law}}^{\Sigma} := 1 - \frac{1}{C}\,\Delta_{\mathrm{law}}^{\mathrm{pre},\Sigma}, \qquad \Lambda_{\mathrm{state}}^{\Sigma} := 1 - \frac{1}{C}\,\Delta_{\mathrm{state}}^{\mathrm{pre},\Sigma}. \tag{9.22}$$

A convenient choice is to take C from a local band bound supplied by Lemma 9.46 or from fibre normalizations $N(\psi, x)$ in Definition 9.19, ensuring $0 \le \Lambda_\bullet^\Sigma \le 1$. Interferometric realizations of these visibilities are provided by holosymmetric overlaps in the control–unitary scheme: evaluate the same band–local functionals of $\hat{Q}_x[v_k]$ (or of $\bar{Q}_x[v_k] = \langle \hat{Q}_x[v_k] \rangle$) as Born pairings in the Holo–State/GNS setting (cf. Definitions 1.18, 1.34 and 9.8, Section 4.19.1, and Remark 9.9). Gauge/transport covariance of $\hat{\mathbf{G}}_\Lambda$ and of $\mathcal{U}$ (cf. Proposition 9.6 and Definition 9.7) and band regularity of $\hat{\Sigma}$ (cf. Definitions 6.280 and 9.10 and Proposition 9.13) guarantee that the interferometric visibilities operationalize (9.21)–(9.22) on the band; stability under band deformations then follows from Lemmas 9.27 and 9.42.

Proposition 9.48 (Equivalence with correlator-only diagnostics and triadic attribution)**.** *If* $\bar{\Sigma}_{\mu\nu} = \bar{\Sigma}_{\mu\nu}^{\mathrm{corr}} + \bar{\Sigma}_{\mu\nu}^{\mathrm{vac}} + \bar{\Sigma}_{\mu\nu}^{\mathrm{spec}}$ *(expectation–level triadic split; cf. Definition 5.54 and eqs. (5.8) and (5.21)), then choosing the band–local, gauge–invariant functional* $\mathcal{I}$ *channelwise yields*

$$(\Delta_{\mathrm{law}}^{\mathrm{pre},\Sigma}, \Delta_{\mathrm{state}}^{\mathrm{pre},\Sigma}) \rightsquigarrow \left(\Delta_{\mathrm{law}}^{\mathrm{pre,corr}}, \Delta_{\mathrm{law}}^{\mathrm{pre,vac}}, \Delta_{\mathrm{law}}^{\mathrm{pre,spec}}\right), \left(\Delta_{\mathrm{state}}^{\mathrm{pre,corr}}, \Delta_{\mathrm{state}}^{\mathrm{pre,vac}}, \Delta_{\mathrm{state}}^{\mathrm{pre,spec}}\right),$$

and, in particular, the correlator–distance scheme upon deactivating the other channels.

Proof. (i) Operator–level triadic split and quadratic forms. On the band, the chronogeometry admits the operator–level split $\hat{\Sigma}_{\mu\nu} = \hat{\Sigma}^{(\mathrm{corr})}_{\mu\nu} + \hat{\Sigma}^{(\mathrm{vac})}_{\mu\nu} + \hat{\Sigma}^{(\mathrm{spec})}_{\mu\nu}$ consistent with the expectation–level decomposition (cf. Definitions 5.54 and 9.20). For a probe $v \in T_x\mathcal{X}_q$, the Σ–quadratic form $\hat{Q}_x[v] = v^\mu v^\nu \hat{\Sigma}_{\mu\nu}(x)$ (cf. Definition 9.22) decomposes as

$$\hat{Q}_x[v] = \sum_{a\in\{\mathrm{corr,vac,spec}\}} \hat{Q}^{(a)}_x[v], \qquad \hat{Q}^{(a)}_x[v] := v^\mu v^\nu \hat{\Sigma}^{(a)}_{\mu\nu}(x).$$

(ii) Channel projectors and channelwise functionals. By the Fourfold grading and its block algebra (cf. Definition 9.8, eq. (4.24), and Section 4.1.7), there exist band–local projectors $\Pi^{(a)}$ selecting the triadic channels and commuting with graded transport (cf. Definition 9.7 and Proposition 9.6). Define channelwise functionals from any admissible band–local, gauge–invariant $\mathcal{I}$ (Born pairing / regulated band–trace; cf. Definitions 1.18, 1.34, 6.280 and 9.10) by

$$\mathcal{I}^{(a)}[X] := \mathcal{I}\Big[\Pi^{(a)} X \Pi^{(a)}\Big].$$

Then $\mathcal{I}^{(a)}[\hat{Q}_x[v]] = \mathcal{I}^{(a)}[\hat{Q}^{(a)}_x[v]]$ and $\mathcal{I}^{(a)}[\hat{Q}^{(b)}_x[v]] = 0$ for $b \neq a$.

(iii) Law contrast. Using the definition of the pre–physical Σ–law contrast (9.19) (cf. Definition 9.45) and linearity,

$$\Big(\Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}}(g;x;v)\Big)^2 = \mathcal{I}\Big[\hat{Q}_x[v] - \mathrm{Ad}_{g(x)}\,\hat{Q}_x[v]\Big] = \sum_a \mathcal{I}^{(a)}\Big[\hat{Q}^{(a)}_x[v] - \mathrm{Ad}_{g(x)}\,\hat{Q}^{(a)}_x[v]\Big].$$

ummands define the channelwise law contributions $\Big(\Delta^{\mathrm{pre},a}_{\mathrm{law}}(g;x;v)\Big)^2$. Frame independence and graded G–covariance follow from Lemma 9.24.

(iv) State contrast. Similarly, with $\mathcal{I}_\Psi[\cdot] = \langle\Psi|\cdot|\Psi\rangle$ (cf. Definition 9.45),

$$\Big(\Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}}(g;x;v)\Big)^2 = \Big|\mathcal{I}_\Psi[\hat{Q}_x[v]] - \mathcal{I}_{g\cdot\Psi}[\hat{Q}_x[v]]\Big| = \sum_a \Big(\mathcal{I}^{(a)}_\Psi[\hat{Q}^{(a)}_x[v]] - \mathcal{I}^{(a)}_{g\cdot\Psi}[\hat{Q}^{(a)}_x[v]]\Big),$$

which yields the channelwise state contributions $\Delta^{\mathrm{pre},a}_{\mathrm{state}}$. Aggregation over $\{v_k\}$ is compatible with these splits (cf. Definitions 9.26 and 9.47).

(v) Correlator–only reduction and attribution. If the vacuum and spectral channels are deactivated (or their projectors set to zero on the band), only $a = \mathrm{corr}$ remains. Then $\hat{\Sigma} \to \hat{\Sigma}^{(\mathrm{corr})}$, $\hat{Q}_x[v] \to \hat{Q}^{(\mathrm{corr})}_x[v]$, and the contrasts reduce to the correlator–only diagnostics built from d_{corr} with fibre normalizations $N(\cdot)$ (cf. Definitions 4.338 and 9.19 and Corollaries 9.30 and 9.40). Functional independence/orthogonality of the triadic generators ensures that no spurious cross–terms contaminate the channelwise attributions (cf. Section 4.11.5); stability and finiteness are guaranteed by Lemmas 9.27 and 9.46.

Combining (iii)–(v) proves the claimed channelwise attribution and the correlator–only equivalence. □

Definition 9.49 (Quadrant predicate (pre-physical, Σ-aware))**.** Fix a tolerance $\varepsilon \ll 1$ and a band $(\lambda_{\mathrm{UV}}, \ell)$ (cf. Definitions 6.280 and 9.10 and eq. (7.130)). Using the pre–physical Σ–contrasts of Definition 9.45 (or their aggregated versions in Definition 9.47), declare at x:

$$\textbf{QI}: \ \Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}}(g;x) < \varepsilon, \ \ \Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}}(g;x) < \varepsilon; \qquad \textbf{QII}: \ \Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}}(g;x) < \varepsilon, \ \ \Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}}(g;x) > \varepsilon;$$

$$\textbf{QIII}: \ \Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}}(g;x) > \varepsilon, \ \ \Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}}(g;x) < \varepsilon; \qquad \textbf{QIV}: \ \Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}}(g;x) > \varepsilon, \ \ \Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}}(g;x) > \varepsilon.$$

By construction, these predicates are frame–independent and respect the graded G–action (cf. Lemma 9.24); finiteness and continuity with respect to v and $(\lambda_{\mathrm{UV}}, \ell)$ follow from Lemmas 9.27 and 9.46. The choice of ε may be tied to local plateau diagnostics and acceptance (cf. eqs. (7.394) to (7.396) and Proposition 7.203). This pre–physical rule mirrors the physical quadrant decision on $\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}})$ after Σ–aware realization (cf. Sections 6.10 and 9.5), where law–channel safety and gauge–parameter independence are ensured by the BRST/Nielsen framework and Ward/Slavnov identities evaluated on plateaus (cf. Section 7.5).

Band-limited BRST complex and graded bracket. On the band/IR–tagged multiplet

$$\mathfrak{X}^{(\lambda_{\mathrm{UV}},\ell)} := \Big(\delta\hat{\mathbf{\Phi}}_{\lambda_{\mathrm{UV}},\ell},\, \delta\hat{\mathbf{\Theta}}_{\lambda_{\mathrm{UV}},\ell},\, c_{\lambda_{\mathrm{UV}},\ell},\, \bar{c}_{\lambda_{\mathrm{UV}},\ell},\, b_{\lambda_{\mathrm{UV}},\ell}\Big)^{\mathsf{T}},$$

ghost numbers are assigned by $\mathrm{gh}(\delta\hat{\mathbf{\Phi}}) = \mathrm{gh}(\delta\hat{\mathbf{\Theta}}) = \mathrm{gh}(b) = 0$, $\mathrm{gh}(c) = +1$, $\mathrm{gh}(\bar{c}) = -1$, and the BRST differential $\zeta(\,\cdot\,) := \{\hat{Q}_{\mathrm{BRST}}, \cdot\}$ raises ghost number by $+1$. By construction, the scale tags commute with the BRST charge and channel projectors, so the band functor $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ preserves the BRST complex and its ghost grading (cf. Section 7.2.1 and Definition 7.11). We use the graded commutator $[A,B]_{\pm} := AB - (-1)^{|A||B|}BA$; nilpotency and (Krein-)Hermiticity of $\hat{Q}_{\mathrm{BRST}}$ hold on a common invariant domain stable under the tags, $\hat{Q}^2_{\mathrm{BRST}} = 0$, $\hat{Q}^{\dagger}_{\mathrm{BRST}} = \hat{Q}_{\mathrm{BRST}}$.

Definition 9.50 (Graded bracket and BRST closure)**.** On a $\mathbb{Z}_2$–graded algebra, the graded (super)–bracket of homogeneous A, B with parities $|A|, |B| \in \{0,1\}$ is

$$[A,B]_{\pm} \ := \ AB \ - \ (-1)^{|A|\,|B|}\, BA.$$

An operator (or instrument) $\mathcal{O}$ is *BRST–closed* if $[\hat{Q}_{\mathrm{BRST}}, \mathcal{O}]_{\pm} = 0$, where $\hat{Q}_{\mathrm{BRST}}$ is the odd BRST charge ($|\hat{Q}_{\mathrm{BRST}}| = 1$). If $|\mathcal{O}| = 0$ (even), then $[\hat{Q}_{\mathrm{BRST}}, \mathcal{O}]_{\pm} = [\hat{Q}_{\mathrm{BRST}}, \mathcal{O}]$; if $|\mathcal{O}| = 1$ (odd), the condition reads $\{\hat{Q}_{\mathrm{BRST}}, \mathcal{O}\} = 0$.

Definition 9.51 (Band tag and its parity)**.** Let $P_x(\lambda_{\mathrm{UV}})$ denote the UV projector (ghost number 0) and $K^{(\ell)}_{\mathrm{IR}}$ the causal IR window (ghost number 0). The *band tag* $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\cdot]$ is the linear instrument obtained by composing these even inputs with the observable argument using only parity–preserving operations (algebraic composition, adjunction, and integration). Consequently, $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ is *even*:

$$\big|\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}\big| = 0.$$

Proposition 9.52 (BRST-closure and naturality of the band tag)**.** *If $P_x(\lambda_{\mathrm{UV}})$ and $K^{(\ell)}_{\mathrm{IR}}$ are BRST–closed and even, i.e.*

$$[\hat{Q}_{\mathrm{BRST}}, P_x(\lambda_{\mathrm{UV}})]_{\pm} = 0, \qquad [\hat{Q}_{\mathrm{BRST}}, K^{(\ell)}_{\mathrm{IR}}]_{\pm} = 0,$$

then:

(i) Closure

$\big[\hat{Q}_{\mathrm{BRST}}, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}\big]_{\pm} = 0$ *as an operator identity; in particular* $|\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}| = 0$ *so the bracket is an ordinary commutator here.*

(ii) Naturality

For every (homogeneous) operator $\hat{O}$*,*

$$\big[\hat{Q}_{\mathrm{BRST}},\, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\hat{O}]\,\big]_{\pm} \;=\; \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}\Big[\,[\hat{Q}_{\mathrm{BRST}}, \hat{O}]_{\pm}\,\Big].$$

Proof. Recall from Definition 9.50 that $\mathrm{ad}_{\hat{Q}_{\mathrm{BRST}}}(\,\cdot\,) := [\hat{Q}_{\mathrm{BRST}},\, \cdot\,]_{\pm}$ is an *odd graded derivation*: for homogeneous A, B,

$$\mathrm{ad}_{\hat{Q}_{\mathrm{BRST}}}(AB) = \mathrm{ad}_{\hat{Q}_{\mathrm{BRST}}}(A)\, B + (-1)^{|A|} A\, \mathrm{ad}_{\hat{Q}_{\mathrm{BRST}}}(B). \tag{9.23}$$

By Definition 9.51, $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ is even and is built from the *even, BRST–closed* $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ using only parity–preserving operations (composition, adjunction, linear smearing/integration by even kernels).

Step 1 (UV sandwich). Let $\mathcal{P}_\lambda(\hat{O}) := P(\lambda_{\mathrm{UV}})\, \hat{O}\, P(\lambda_{\mathrm{UV}})$. Since $[\hat{Q}_{\mathrm{BRST}}, P(\lambda_{\mathrm{UV}})]_{\pm} = 0$ and $|P| = 0$, (9.23) yields

$$\mathrm{ad}_{\hat{Q}_{\mathrm{BRST}}}\Big(\mathcal{P}_\lambda(\hat{O})\Big) = P\, \mathrm{ad}_{\hat{Q}_{\mathrm{BRST}}}(\hat{O})\, P = \mathcal{P}_\lambda\Big(\mathrm{ad}_{\hat{Q}_{\mathrm{BRST}}}(\hat{O})\Big).$$

Step 2 (IR smearing). Write the IR window as an even kernel $K_{\mathrm{IR}}^{(\ell)}$ implementing an even linear superoperator $\mathcal{K}_\ell$ by smearing/convolution, schematically $\mathcal{K}_\ell(X) = \int K_{\mathrm{IR}}^{(\ell)} \diamond X$, with a parity–preserving bilinear pairing $\diamond$. Since $[\hat{Q}_{\mathrm{BRST}}, K_{\mathrm{IR}}^{(\ell)}]_{\pm} = 0$ and $|K_{\mathrm{IR}}^{(\ell)}| = 0$, linearity and (9.23) give

$$\mathrm{ad}_{\hat{Q}_{\mathrm{BRST}}}\Big(\mathcal{K}_\ell(X)\Big) = \int K_{\mathrm{IR}}^{(\ell)} \diamond \mathrm{ad}_{\hat{Q}_{\mathrm{BRST}}}(X) = \mathcal{K}_\ell\Big(\mathrm{ad}_{\hat{Q}_{\mathrm{BRST}}}(X)\Big).$$

Step 3 (Band tag). By construction $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell} = \mathcal{K}_\ell \circ \mathcal{P}_\lambda$. Combining Steps 1–2,

$$\mathrm{ad}_{\hat{Q}_{\mathrm{BRST}}}\Big(\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}(\hat{O})\Big) = \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}\Big(\mathrm{ad}_{\hat{Q}_{\mathrm{BRST}}}(\hat{O})\Big),$$

which is the stated naturality. Taking $\hat{O}$ as a basis of operators implies $\mathrm{ad}_{\hat{Q}_{\mathrm{BRST}}} \circ \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell} = \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell} \circ \mathrm{ad}_{\hat{Q}_{\mathrm{BRST}}}$ as superoperators, hence $[\hat{Q}_{\mathrm{BRST}}, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}]_{\pm} = 0$. □

Corollary 9.53 (Even case (commutator form))**.** *Under the hypotheses of Proposition 9.52,*

$$[\hat{Q}_{\mathrm{BRST}}, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}] = 0, \qquad [\hat{Q}_{\mathrm{BRST}}, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\hat{O}]] = \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}\Big[[\hat{Q}_{\mathrm{BRST}}, \hat{O}]\Big].$$

Proof. Let $\mathrm{ad}_{\hat{Q}}(\cdot) := [\hat{Q}_{\mathrm{BRST}}, \cdot]_{\pm}$ denote the graded derivation (cf. Definition 9.50), and recall from Definition 9.51 that the band tag $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ is *even*, $|\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}| = 0$. By Proposition 9.52 (closure and naturality in graded form) we have, for every homogeneous $\hat{O}$,

$$\mathrm{ad}_{\hat{Q}}\Big(\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\hat{O}]\Big) = \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}\Big(\mathrm{ad}_{\hat{Q}}(\hat{O})\Big) \quad \text{and} \quad \mathrm{ad}_{\hat{Q}}(\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}) = 0, \tag{9.24}$$

as operator identities on a common invariant domain stable under $\hat{Q}_{\mathrm{BRST}}$ and the tag (cf. Section 7.2.1 and Definition 7.11).

Outer (left) bracket becomes an ordinary commutator. Because $|\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}| = 0$ and $|\hat{Q}_{\mathrm{BRST}}| = 1$,

$$[\hat{Q}_{\mathrm{BRST}}, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}]_{\pm} = \hat{Q}_{\mathrm{BRST}}\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell} - (-1)^{1\cdot 0}\,\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}\hat{Q}_{\mathrm{BRST}} = [\hat{Q}_{\mathrm{BRST}}, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}],$$

so the first statement in (9.24) yields

$$[\hat{Q}_{\mathrm{BRST}}, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}] = 0.$$

Inner bracket reduces for even (ghost–number zero) observables. On the BRST complex used for diagnostics and EOM in Chapters 7–8, the physical/observable sector and the band–tagged PSR functionals live at ghost number 0 (even parity). Thus for every such $\hat{O}$ we have $|\hat{O}| = 0$, and hence

$$[\hat{Q}_{\mathrm{BRST}}, \hat{O}]_{\pm} = \hat{Q}_{\mathrm{BRST}}\hat{O} - (-1)^{1\cdot 0}\hat{O}\hat{Q}_{\mathrm{BRST}} = \hat{Q}_{\mathrm{BRST}}\hat{O} - \hat{O}\hat{Q}_{\mathrm{BRST}} = [\hat{Q}_{\mathrm{BRST}}, \hat{O}].$$

Using this in the naturality identity of (9.24) gives

$$[\hat{Q}_{\mathrm{BRST}}, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\hat{O}]] = \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}\Big([\hat{Q}_{\mathrm{BRST}}, \hat{O}]\Big),$$

where the outer bracket is also an ordinary commutator by the first part.

Domain statement. All equalities hold on the common invariant core for $\hat{Q}_{\mathrm{BRST}}$ specified in Section 7.2.1 and preserved by $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ (see Definition 7.11); by linearity and closability (Chapter 7), they extend to the natural operator domains encountered in the band–tagged constructions of Definition 6.272. This proves the two identities of the corollary. □

Lemma 9.54 (BRST hygiene of band-tagged diagnostics)**.** *If the UV projector $P_x(\lambda_{\mathrm{UV}})$ and the IR window $K_{\mathrm{IR}}^{(\ell)}$ are BRST–closed (ghost number 0), then the band tag $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\cdot]$ descends to the physical sector and physical readouts are gauge–parameter independent. Hence the PSR diagnostics* are *valid inputs to the realization in $H^0(\hat{Q}_{\mathrm{BRST}})$.*

Proof. (i) BRST–compatibility of the band tag. By definition (cf. Definitions 6.280 and 9.10 and eq. (9.2)) and the "tag hygiene" hypothesis $[\hat{Q}_{\mathrm{BRST}}, P_x(\lambda_{\mathrm{UV}})]_{\pm} = [\hat{Q}_{\mathrm{BRST}}, K_{\mathrm{IR}}^{(\ell)}]_{\pm} = 0$ (cf. eq. (7.24)), the band tag superoperator satisfies

$$[\hat{Q}_{\mathrm{BRST}}, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}]_{\pm} = 0,$$

as established in Lemma 9.11. Equivalently, for any local operator $\mathfrak{O}$ of ghost number 0,

$$[\hat{Q}_{\mathrm{BRST}}, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\mathfrak{O}]]_{\pm} = \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}\Big([\hat{Q}_{\mathrm{BRST}}, \mathfrak{O}]_{\pm}\Big),$$

i.e. $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ is a chain map for the BRST complex (cf. Definition 9.12).

(ii) Descent to cohomology and the physical projector. Let $\mathsf{P}_{\mathrm{phys}}$ denote the projection onto $\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}})$ (cf. Section 6.10). If $\mathfrak{O}$ is ghost–number 0 and $\mathfrak{O} \sim \mathfrak{O} + [\,\hat{Q}_{\mathrm{BRST}}, Y]_{\pm}$ in cohomology for some local operator Y of ghost number -1, then by the chain–map property,

$$\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\mathfrak{O}] \;\sim\; \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\mathfrak{O}] + [\,\hat{Q}_{\mathrm{BRST}},\, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[Y]\,]_{\pm},$$

so the cohomology class of $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\mathfrak{O}]$ is well–defined. Hence physical readouts (matrix elements with $\mathsf{P}_{\mathrm{phys}}$ inserted) depend only on the class

$$\big\langle \mathsf{P}_{\mathrm{phys}}\, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\mathfrak{O}]\, \mathsf{P}_{\mathrm{phys}} \big\rangle,$$

and therefore $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ descends to H^0.

(iii) Gauge–parameter independence (Nielsen/Slavnov identities). Variations of the gauge–fixing parameter(s) modify the gauge–fixed generator by a BRST–exact insertion; physical expectations of BRST–closed observables are invariant (Nielsen identities, Ward/Slavnov framework; cf. Section 7.5). Since $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\cdot]$ is BRST–compatible and (by construction) ghost–number preserving, for any BRST–closed preimage $\mathfrak{O}$ one has

$$\frac{\partial}{\partial \xi}\big\langle \mathsf{P}_{\mathrm{phys}}\, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\mathfrak{O}]\, \mathsf{P}_{\mathrm{phys}} \big\rangle \;=\; \big\langle \mathsf{P}_{\mathrm{phys}}\, [\hat{Q}_{\mathrm{BRST}}, \dots]_{\pm}\, \mathsf{P}_{\mathrm{phys}} \big\rangle \;=\; 0,$$

so physical readouts are gauge–parameter independent on plateaus (cf. Section 9.5).

(iv) Application to PSR diagnostics. The law–kernel diagnostic $\mathfrak{L}_{\mathrm{pre}} = \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\mathfrak{L}(\hat{\mathbf{G}}_\Lambda)]$ (cf. Definition 9.44) and the Σ–contrasts built from $\hat{Q}_x[v] = v^\mu v^\nu \hat{\Sigma}_{\mu\nu}(x)$ (cf. Definitions 9.20, 9.22 and 9.45) are band–local, gauge–covariant constructs (transport by $\mathcal{U}$; cf. Definition 9.7 and Proposition 9.6). By Step (i), their band–tagged versions are BRST–compatible; by Steps (ii)–(iii) they define elements in H^0 whose physical readouts are gauge–parameter independent. Thus the PSR diagnostics are valid inputs to the Σ–aware realization map and subsequent evaluation on $\mathcal{H}_{\mathrm{phys}}$ (cf. Section 9.5).

Combining (i)–(iv) proves the claim. □

Proposition 9.55 (Operational visibilities and overlaps)**.** *Prepare two chronogeometric histories with unitary propagators $\hat{U}_0, \hat{U}_1$ generated by band–smeared $\hat{\Sigma}_{\mu\nu}$,*

$$\hat{U}_j \;:=\; \exp\!\Big(-i \int_{\mathcal{X}_q} f^{\mu\nu}_{(j)}(x)\, \hat{\Sigma}_{\mu\nu}(x)\, \mathrm{d}\mu_x \Big), \qquad j = 0, 1,$$

where $f^{\mu\nu}_{(j)}$ are band–supported smearing kernels (cf. Definitions 6.280, 9.10 and 9.20 and eq. (9.2)). In the control–unitary interferometric scheme (holosymmetric overlap), the detection probability at output c for a phase bias φ is

$$p_c(\mathbf{0} \mid \varphi) \;=\; \tfrac{1}{2}\Big(1 + \mathrm{Re}\,\big[e^{-i\varphi}\, \langle \Psi_{\mathrm{RR}} | \hat{U}_1^\dagger \hat{U}_0 | \Psi_{\mathrm{RR}} \rangle\big]\Big), \tag{9.25}$$

with $|\Psi_{\mathrm{RR}}\rangle$ a Holo–State readout vector (cf. Definitions 1.18 and 1.34 and Section 4.19.1). The fringe visibility $V := \big|\langle \Psi_{\mathrm{RR}} | \hat{U}_1^\dagger \hat{U}_0 | \Psi_{\mathrm{RR}} \rangle\big|$ is band–local, gauge/frame covariant (cf. Proposition 9.6 and Definition 9.7) and coincides—up to a normalization constant $C > 0$—with the Σ–visibility Λ^Σ defined in Definition 9.47, i.e.

$$V \;\equiv\; \Lambda^\Sigma \;:=\; 1 - \frac{1}{C}\,\Delta^{\mathrm{pre},\Sigma}, \qquad \Delta^{\mathrm{pre},\Sigma} \in \Big\{ \Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}},\; \Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}} \Big\}$$

(cf. eqs. (9.21) and (9.22)), after calibrating C with the same band and probe set. Phase damping with dephasing factor $\eta \in [0, 1]$ on the control arm multiplies the visibility: $V \mapsto \eta\, V$.

Proof. (i) Band–local, covariant histories. By construction the generators $\int f^{\mu\nu}_{(j)}\hat{\Sigma}_{\mu\nu}$ are band–smeared, hence bounded on the band fibre and covariant under graded transport (cf. Definitions 9.7 and 9.20 and Propositions 9.13 and 9.14). Therefore $\hat{U}_j$ are well–defined unitary operators on $\mathcal{H}_{\mathrm{em}}$ that transform by endpoint conjugation, and their matrix elements in gauge–covariant Holo–States are gauge–invariant (cf. Proposition 9.6 and Section 4.19.1).

(ii) Control–unitary interferometry and overlap formula. In the standard control–unitary scheme (holosymmetric overlaps; cf. Definition 9.47 and Section 4.19.1), the control qubit routes the system through $\hat{U}_0$ or $\hat{U}_1$ and recombines with phase bias φ. Tracing the control in the $\{|c\rangle\}$ basis yields (9.25), with complex fringe amplitude $A := \langle\Psi_{\mathrm{RR}}|\hat{U}_1^\dagger\hat{U}_0|\Psi_{\mathrm{RR}}\rangle$ and operational visibility $V = |A|$.

(iii) Identification with Σ–visibilities via BCH/Duhamel and cumulants. Write $\hat{U}_0 = \exp\!\big(-i\hat{H}_0\big)$, $\hat{U}_1 = \exp\!\big(-i\hat{H}_1\big)$ with $\hat{H}_j = \int f^{\mu\nu}_{(j)}\hat{\Sigma}_{\mu\nu}$, and set $\delta\hat{H} := \hat{H}_0 - \hat{H}_1 = \int \delta f^{\mu\nu}\hat{\Sigma}_{\mu\nu}$ where $\delta f^{\mu\nu} := f^{\mu\nu}_{(0)} - f^{\mu\nu}_{(1)}$ has the same band support. The Baker–Campbell–Hausdorff/Duhamel control on the band gives

$$e^{i\hat{H}_1}e^{-i\hat{H}_0} = \exp\!\Big(-i\,\delta\hat{H} \;+\; \tfrac{1}{2}[\hat{H}_1,-\hat{H}_0] \;+\; \mathcal{O}(\|\delta\hat{H}\|^2)\Big) = \exp\!\big(-i\,\delta\hat{H} \;+\; \hat{R}\big),$$

where $\hat{R}$ is anti–Hermitian and $\|\hat{R}\| = \mathcal{O}(\|\delta\hat{H}\|^2)$ (boundedness from band–smoothing; cf. Definitions 6.280 and 9.10). Hence

$$A = \Big\langle e^{-i\,\delta\hat{H}+\hat{R}}\Big\rangle_{\Psi_{\mathrm{RR}}} = \Big\langle e^{-i\,\delta\hat{H}}\Big\rangle_{\Psi_{\mathrm{RR}}} + \underbrace{\langle\hat{R}\rangle_{\Psi_{\mathrm{RR}}}}_{\text{purely imaginary at } \mathcal{O}(\|\delta\hat{H}\|^2)} + \mathcal{O}(\|\delta\hat{H}\|^3).$$

Expanding (or using the cumulant expansion) for the single operator $\delta\hat{H}$ gives

$$\begin{aligned}\Big\langle e^{-i\,\delta\hat{H}}\Big\rangle_{\Psi_{\mathrm{RR}}} &= 1 - i\langle\delta\hat{H}\rangle_{\Psi_{\mathrm{RR}}} - \tfrac{1}{2}\langle\delta\hat{H}^2\rangle_{\Psi_{\mathrm{RR}}} + \mathcal{O}(\|\delta\hat{H}\|^3)\\ &= \exp\!\Big(-i\,\langle\delta\hat{H}\rangle_{\Psi_{\mathrm{RR}}} - \tfrac{1}{2}\,\mathrm{Var}_{\Psi_{\mathrm{RR}}}(\delta\hat{H}) + \mathcal{O}(\|\delta\hat{H}\|^3)\Big),\end{aligned}$$

with $\mathrm{Var}_{\Psi_{\mathrm{RR}}}(\delta\hat{H}) := \langle\delta\hat{H}^2\rangle_{\Psi_{\mathrm{RR}}} - \langle\delta\hat{H}\rangle^2_{\Psi_{\mathrm{RR}}}$. Choosing the interferometer phase as $\varphi = \langle\delta\hat{H}\rangle_{\Psi_{\mathrm{RR}}} + \mathcal{O}(\|\delta\hat{H}\|^2)$ removes the linear phase. Since $\langle\hat{R}\rangle_{\Psi_{\mathrm{RR}}}$ is imaginary at order $\|\delta\hat{H}\|^2$, it alters only the *phase* at that order; the *magnitude* $V = |A|$ is therefore

$$V = 1 - \tfrac{1}{2}\,\mathrm{Var}_{\Psi_{\mathrm{RR}}}(\delta\hat{H}) + \mathcal{O}(\|\delta\hat{H}\|^3).$$

Now decompose $\delta\hat{H}$ along the quadratic probes: $\delta\hat{H} = \sum_k w_k\,\hat{Q}_x[v_k]$, with band–local weights w_k encoding the smearing and probes (cf. Definitions 9.22 and 9.47). Using the same gauge–invariant, band–local functional $\mathcal{I}$ as in Definition 9.45, calibrate a constant $C > 0$ (fixed by $\{v_k\}$ and $\{w_k\}$) so that $\mathrm{Var}_{\Psi_{\mathrm{RR}}}(\delta\hat{H}) = \frac{1}{C}\Big(\Delta^{\mathrm{pre},\Sigma}\Big)^2$. Therefore

$$V \;=\; 1 - \frac{1}{2C}\Big(\Delta^{\mathrm{pre},\Sigma}\Big)^2 + \mathcal{O}\big(\Delta^3\big), \qquad \Lambda^{\Sigma} := 1 - \frac{1}{C}\,\Delta^{\mathrm{pre},\Sigma}$$

(cf. eqs. (9.21) and (9.22)), and after the (plateau–fixed) calibration, V coincides with Λ^{Σ} up to the stated normalization and higher–order terms (cf. Proposition 7.203 and Lemmas 9.27 and 9.46).

(iv) Phase damping. A dephasing channel on the control with factor $\eta \in [0,1]$ multiplies the off–diagonal coherence of the control state by η; since the fringe term in (9.25) is proportional to this coherence, the visibility scales as $V \mapsto \eta V$ (standard interferometric damping; cf. Section 4.19.1). Band–locality and covariance are unaffected, as the dephasing acts only on the control degree of freedom.

Finally, BRST hygiene of the band–tagged diagnostics ensures these interferometric readouts are valid inputs to realization in H^0 (cf. Lemma 9.54 and Section 9.5). This proves the claim. □

Supporting sources for steps (ii)–(iii): the control–unitary overlap formula is stated in Proposition 5.36 (law–layer visibility as a relational overlap). For BCH/Duhamel control and product formulas, see [457, 116, 91]. The interferometric identification and phase choice are standard in quantum information [287] and connect to fidelity/overlap functionals [357]. For a QIV–layered Duhamel/BCH statement at plateau points, see Proposition 7.616.

Corollary 9.56 (Plateau consistency of the law channel)**.** *On certified plateaus $\mathfrak{P}$, law–channel visibilities satisfy $\left|\Lambda^{\Sigma}_{\mathrm{law}} - 1\right) = \mathcal{O}(\varepsilon_{\mathrm{plat}})$ (Nielsen–safe), hence pre–physical QII decisions (law–symmetric, state–asymmetric) are stable under Σ–aware upgrade to the physical sector.*

Proof. (i) Plateau control and band stability. By the grid diagnostics and acceptance rule for a certified plateau (cf. eqs. (7.394) to (7.396) and Proposition 7.203), all band–local observables vary by at most $\mathcal{O}(\varepsilon_{\mathrm{plat}})$ under $(\lambda_{\mathrm{UV}}, \ell)$–moves within $\mathfrak{P}$. In particular, the Σ–law contrast of Definition 9.45 (and its aggregated version Definition 9.47) obeys

$$\Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}}(g;x) \;=\; \mathcal{O}(\varepsilon_{\mathrm{plat}}),$$

with continuity furnished by Lemmas 9.27 and 9.46.

(ii) From contrasts to visibilities. By Definition 9.47, $\Lambda^{\Sigma}_{\mathrm{law}} = 1 - \frac{1}{C}\,\Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}}$ for a fixed normalization $C > 0$ calibrated on the same band/probe set. Hence,

$$\left|\Lambda^{\Sigma}_{\mathrm{law}} - 1\right| = \frac{1}{C}\,\Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}} = \mathcal{O}(\varepsilon_{\mathrm{plat}}).$$

Operationally, the control–unitary interferometric visibility V of Proposition 9.55 coincides with $\Lambda^{\Sigma}_{\mathrm{law}}$ up to this fixed C, so the same plateau bound holds for V.

(iii) Nielsen safety and descent to H^0. The band tag is BRST–compatible (cf. Lemmas 9.11 and 9.54 and eq. (7.24)), so the law–channel readouts descend to $\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}})$ with gauge–parameter independence ensured by the Nielsen/Ward–Slavnov identities on plateaus (cf. Sections 6.10, 7.5 and 9.5). Therefore the near–unity bound for $\Lambda^{\Sigma}_{\mathrm{law}}$ persists after realization, i.e. it is Nielsen–safe.

(iv) Stability of the QII decision. The pre–physical quadrant predicate (Definition 9.49) declares QII when $\Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}} < \varepsilon$ and $\Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}} > \varepsilon$. By (i)–(iii), upgrading to the physical layer alters the law–channel by at most $\mathcal{O}(\varepsilon_{\mathrm{plat}})$ and does not introduce gauge–parameter drift; by continuity and the plateau acceptance slack (cf. Proposition 7.203), this does not cross the law–threshold. State–channel continuity & band stability (Lemmas 9.27, 9.42 and 9.46) prevent threshold crossing provided the empirical margin exceeds $\mathcal{O}(\varepsilon_{\mathrm{plat}})$. Hence QII decisions are stable under upgrade. □

Remark 9.57 (Why Σ strengthens PSR)**.** The Σ–diagnostics probe both *amplitude curvature* $\hat{g}$ and *phase–flow curvature* $\hat{t}$ within a single operator $\hat{\Sigma} = \hat{g} + \hat{t}$ (cf. Definition 9.20 and eq. (9.4)) via the quadratic probe $\hat{Q}_x[v] = v^\mu v^\nu \hat{\Sigma}_{\mu\nu}(x)$ (cf. Definition 9.22). Thus the pre–physical contrasts (cf. Definitions 9.45 and 9.47) are intrinsically sensitive to temporal/Nielsen–type effects and to channel mixing handled by the Σ–weighted eigenmode extraction (cf. Propositions 9.36 and 9.39 and Remark 9.41). At the expectation level, $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ fixes the causal cones governing hyperbolicity and response (the Σ–causal viewpoint), sharpening linear–response kernels and Ward/Slavnov checks without altering the BRST/cohomological downstream machinery (cf. Sections 7.5 and 9.5). This unified treatment is strictly stronger than "distances–only'' diagnostics: it preserves all correlator–only conclusions in the appropriate limit (cf. Corollaries 9.30 and 9.40) while adding principled sensitivity to temporal phase–flow and inter–modal couplings (cf. Lemma 9.38 and Proposition 9.29).

Practical decision rule. Compute the pre–physical Σ–contrasts $\Delta^{\mathrm{pre},\Sigma}_{\mathrm{law/state}}$ using the quadratic probe $\hat{Q}_x[v] = v^\mu v^\nu \hat{\Sigma}_{\mu\nu}(x)$ (cf. Definitions 9.20 and 9.22) and the band–local, gauge–invariant functionals $\mathcal{I}$ of Definition 9.45. When multiple probes $\{v_k\}$ are used, aggregate them by the root–sum–square prescription in Definition 9.47, i.e. (9.21), and optionally report the complementary visibilities (9.22). Assign the quadrant label QI–QIV via the Σ–aware predicate of Definition 9.49, and record channelwise (corr/vac/spec) attributions using the triadic split and channel–projected functionals of Proposition 9.48. Finiteness and continuity of all diagnostics are guaranteed on the band by Lemmas 9.27 and 9.46; gauge/frame covariance follows from the graded transport and Fourfold Hermiticity/covariance (cf. Definition 9.7 and Proposition 9.6). On certified plateaus (grid diagnostics and acceptance: eqs. (7.394) to (7.396) and Proposition 7.203), the law channel is Nielsen–flat and descends to $H^0(\hat{Q}_{\mathrm{BRST}})$ by BRST hygiene of the tag (cf. Lemma 9.54); hence the near–unity bound of Corollary 9.56 ensures that law–symmetric (QII) decisions are stable under Σ–aware realization.

Summary of Section 9.4. We defined Σ–based PSR order parameters using band–tagged, graded–covariant operators that integrate amplitude curvature $\hat{g}$ and temporal phase–flow $\hat{t}$ into the single chronogeometric tensor $\hat{\Sigma} = \hat{g} + \hat{t}$ (cf. Definition 9.20 and eq. (9.4)). (i) These order parameters are *finite and robust* on the band: the quadratic probes and their contrasts are well–defined, continuous in probes and tag knobs, and stable under small band deformations (cf. Definitions 9.45 and 9.47 and Lemmas 9.27 and 9.46); covariance is ensured by Definition 9.7 and Proposition 9.6. (ii) They *reduce to the correlator–only scheme* with explicit triadic attribution: choosing $\mathcal{I}$ channelwise yields correlator/vacuum/spectral contributions and collapses to the original correlator distances when the other channels are deactivated (cf. Proposition 9.48 and Corollaries 9.30 and 9.40). (iii) They *feed a QI–QIV predicate* identical in logic to the physical sector's: the pre–physical quadrant rule Definition 9.49 mirrors the BRST–physical decision after Σ–aware realization (cf. Sections 6.10 and 9.5), with the law channel Nielsen–safe on plateaus by Lemma 9.54 and Corollary 9.56. Altogether, the Σ–diagnostics supply band–regular, gauge–covariant, and triad–interpretable inputs that prepare clean, BRST–consistent realization in $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$.

9.5 Realization to the BRST-physical Sector and Spectrum

Definition 9.58 (Certified plateau and Σ-regularity)**.** A *certified plateau* $\mathfrak{P}$ is a nonempty set of band choices $\{(\lambda,\ell)\}$ (cf. Definitions 6.280 and 9.10 and eqs. (7.130) and (9.2)) with the following properties:

(i) Σ-regularity.
The operator Hessians at coincidence exist and are stable on the band, yielding $\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu}$ and $\hat{\Sigma}_{\mu\nu} = \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu}$ (cf. Definition 9.20 and eq. (9.4)). Finiteness and smoothness on the band follow from Spectral Smoothness and Finite Entanglement Capacity (cf. Propositions 9.13 and 9.14 and Remark 9.21).

(ii) Causality/hyperbolicity.
The expectation $\bar{\Sigma}$ admits a Lorentzian projection and defines causal cones that control response (cf. Definitions 6.7 and 9.22 and Remark 9.15).

(iii) Nielsen-flatness.
Physical observables' poles or residues are insensitive to gauge-parameter or fermion scans on $\mathfrak{P}$, i.e. law-channel visibilities are near unity and gauge-parameter independent (cf. Corollary 9.56, Lemma 9.54, Section 7.5, and Proposition 9.55).

(iv) BRST viability.
$Q_{\mathrm{BRST}}^2 = 0$ and $[Q_{\mathrm{BRST}}, \hat{H}] = 0$ on the relevant domains so that projection to $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ is well-defined and consistent with the band tag (cf. Section 6.10, Lemma 9.11, and Remark 9.17).

Plateau certification is effected by the local grid diagnostics and acceptance rule (cf. eqs. (7.394) to (7.396) and Proposition 7.203).

Lemma 9.59 (BRST hygiene of the band map)**.** *If* $[\hat{Q}_{\mathrm{BRST}}, P_x(\lambda)]_\pm = [\hat{Q}_{\mathrm{BRST}}, K_{\mathrm{IR}}^{(\ell)}]_\pm = 0$ *(cf. Lemma 9.11), then* $[\hat{Q}_{\mathrm{BRST}}, \mathfrak{B}_{\lambda,\ell}]_\pm = 0$*. In particular, band–tagging commutes with BRST cohomology, so BRST–exact signals are removed by the physical projector* $\mathsf{P}_{\mathrm{phys}}$*.*

Proof. Recall the band map (cf. Definitions 6.280 and 9.10 and eq. (9.2))

$$\mathfrak{B}_{\lambda,\ell}[\mathfrak{O}](x) := \int_{\mathcal{X}_q} K_{\mathrm{IR}}^{(\ell)}(x,y)\, P_y(\lambda)\, \mathfrak{O}(y)\, P_y(\lambda)\, \mathrm{d}\mu_y,$$

with all kernels/transport chosen gauge–covariantly (cf. Definition 9.7 and Proposition 9.6). Using the graded Leibniz rule for commutators and the hypotheses $[\hat{Q}_{\mathrm{BRST}}, P(\lambda)]_\pm = 0$, $[\hat{Q}_{\mathrm{BRST}}, K_{\mathrm{IR}}^{(\ell)}]_\pm = 0$ (tag hygiene; cf. Lemma 9.11), one obtains for any local operator $\mathfrak{O}$ of ghost number 0:

$$\begin{aligned}
[\hat{Q}_{\mathrm{BRST}}, \mathfrak{B}_{\lambda,\ell}[\mathfrak{O}]]_\pm &= \int \Big([\hat{Q}_{\mathrm{BRST}}, K_{\mathrm{IR}}^{(\ell)}]_\pm\, P\, \mathfrak{O}\, P + K_{\mathrm{IR}}^{(\ell)} [\hat{Q}_{\mathrm{BRST}}, P]_\pm\, \mathfrak{O}\, P \\
&\qquad \pm K_{\mathrm{IR}}^{(\ell)} P\, [\hat{Q}_{\mathrm{BRST}}, \mathfrak{O}]_\pm\, P \pm K_{\mathrm{IR}}^{(\ell)} P\, \mathfrak{O}\, [\hat{Q}_{\mathrm{BRST}}, P]_\pm \Big)\, \mathrm{d}\mu \\
&= \int K_{\mathrm{IR}}^{(\ell)}(x,y)\, P_y(\lambda)\, [\hat{Q}_{\mathrm{BRST}}, \mathfrak{O}(y)]_\pm\, P_y(\lambda)\, \mathrm{d}\mu_y \\
&= \mathfrak{B}_{\lambda,\ell}\Big([\hat{Q}_{\mathrm{BRST}}, \mathfrak{O}]_\pm \Big),
\end{aligned}$$

i.e. $\mathfrak{B}_{\lambda,\ell}$ is a chain map for the BRST complex:

$$[\hat{Q}_{\mathrm{BRST}}, \mathfrak{B}_{\lambda,\ell}]_{\pm} = 0.$$

Thus $\mathfrak{B}_{\lambda,\ell}$ descends to cohomology: if $\mathfrak{O} \sim \mathfrak{O} + [\,\hat{Q}_{\mathrm{BRST}}, Y]_{\pm}$ for some ghost–number -1 operator Y, then

$$\mathfrak{B}_{\lambda,\ell}[\mathfrak{O}] \;\sim\; \mathfrak{B}_{\lambda,\ell}[\mathfrak{O}] + [\,\hat{Q}_{\mathrm{BRST}}, \mathfrak{B}_{\lambda,\ell}[Y]]_{\pm},$$

so the class $[\mathfrak{B}_{\lambda,\ell}[\mathfrak{O}]] \in H^0(\hat{Q}_{\mathrm{BRST}})$ is well–defined (cf. Definition 9.12 and Lemma 9.54).

Finally, with $\mathsf{P}_{\mathrm{phys}}$ the projector onto $\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}})$ (cf. Section 6.10), one has $\mathsf{P}_{\mathrm{phys}}\hat{Q}_{\mathrm{BRST}} = 0 = \hat{Q}_{\mathrm{BRST}}\mathsf{P}_{\mathrm{phys}}$ on the relevant domains. Hence for any ghost–number -1 operator Y,

$$\mathsf{P}_{\mathrm{phys}}\,\mathfrak{B}_{\lambda,\ell}\Big([\,\hat{Q}_{\mathrm{BRST}}, Y]_{\pm}\Big)\,\mathsf{P}_{\mathrm{phys}} = \mathsf{P}_{\mathrm{phys}}[\,\hat{Q}_{\mathrm{BRST}}, \mathfrak{B}_{\lambda,\ell}[Y]]_{\pm}\mathsf{P}_{\mathrm{phys}} = 0,$$

so BRST–exact signals are removed by the physical projection. This establishes the claim. □

Definition 9.60 (Σ–realization map)**.** Given any band–tagged pre–physical operator $\hat{X}$, define its Σ–*realization* by

$$\mathcal{U}_{\Sigma}[\hat{X}] \;:=\; \lim_{(\lambda,\ell)\in\mathfrak{P}} \mathsf{P}_{\mathrm{phys}}\,\mathfrak{B}_{\lambda,\ell}[\hat{X}]\,\mathsf{P}_{\mathrm{phys}}, \qquad \text{with } \mathfrak{P} \text{ as in Definition 9.58.} \tag{9.26}$$

Here $\mathfrak{B}_{\lambda,\ell}$ is the gauge–covariant band map built from the UV projector and causal IR window (cf. Definitions 6.280 and 9.10 and eqs. (7.130) and (9.2)), and $\mathsf{P}_{\mathrm{phys}}$ projects onto $\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}})$ (cf. Section 6.10). BRST hygiene of the tag implies that $\mathfrak{B}_{\lambda,\ell}$ is a chain map, $[\,\hat{Q}_{\mathrm{BRST}}, \mathfrak{B}_{\lambda,\ell}]_{\pm} = 0$ (cf. Lemmas 9.11 and 9.59), so (9.26) is well–defined on BRST cohomology and BRST–exact contributions are removed by $\mathsf{P}_{\mathrm{phys}}$. Existence of the limit along the plateau net follows from plateau certification and stability—Σ–regularity and causality/hyperbolicity on $\mathfrak{P}$ together with continuity bounds in the band knobs (cf. Definition 9.58, Lemmas 9.27, 9.42 and 9.46, and Corollary 9.56). Gauge/frame covariance of $\mathcal{U}_{\Sigma}[\hat{X}]$ is inherited from the graded transport and the Hermiticity/covariance of the Fourfold correlator (cf. Definition 9.7 and Proposition 9.6).

Applying (9.26) to the chronogeometric operator field $\hat{\Sigma}_{\mu\nu} = \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu}$ defined by SOT–C^2 Hessians on the band (cf. Definition 9.20 and eq. (9.4)) yields the *physical chronogeometry*

$$\bar{\Sigma}_{\mu\nu}(x) \;:=\; \big\langle \mathcal{U}_{\Sigma}[\hat{\Sigma}_{\mu\nu}(x)] \big\rangle \;=\; \bar{g}_{\mu\nu}(x) + \bar{t}_{\mu\nu}(x), \tag{9.27}$$

with the expectation taken in the physical GNS/Born sense (cf. Definitions 1.18 and 1.34). The resulting $\bar{\Sigma}$ admits the relational/Lorentzian projection and governs Σ–causal response on $\mathcal{H}_{\mathrm{phys}}$ (cf. Definition 6.7, Remark 9.15, and Section 9.5).

Proposition 9.61 (Gauge–parameter independence of $\bar{\Sigma}$ (Σ–Nielsen identity))**.** *Under the conditions of Definition 9.58, one has* $\partial_{\xi,\alpha_{\mathrm{gf}}}\bar{\Sigma}_{\mu\nu}(x) = 0$*. Hence all* $\bar{\Sigma}$*–dependent visibilities and response kernels are gauge–parameter independent on the plateau.*

Proof. Step 0 (realized chronogeometry on a certified plateau). By definition,

$$\bar{\Sigma}_{\mu\nu}(x) = \Big\langle \mathcal{U}_\Sigma\big[\hat{\Sigma}_{\mu\nu}(x)\big] \Big\rangle = \lim_{(\lambda,\ell)\in\mathfrak{P}} \Big\langle \mathsf{P}_{\mathrm{phys}}\, \mathfrak{B}_{\lambda,\ell}\big[\hat{\Sigma}_{\mu\nu}(x)\big]\, \mathsf{P}_{\mathrm{phys}} \Big\rangle,$$

with $\mathcal{U}_\Sigma$ in Definition 9.60, $\mathfrak{P}$ certified by Definition 9.58, and $\hat{\Sigma}_{\mu\nu}$ defined band–locally by SOT–C^2 Hessians (cf. Definition 9.20 and eq. (9.4)). Plateau regularity and band stability (Definition 9.58 and Lemmas 9.27, 9.42 and 9.46) justify interchanging the limit and the gauge–parameter derivatives.

Step 1 (BRST chain–map property of the band tag). Tag hygiene (Lemma 9.11) implies for any ghost–number–0 operator $\mathfrak{O}$:

$$[\,\hat{Q}_{\mathrm{BRST}}, \mathfrak{B}_{\lambda,\ell}[\mathfrak{O}]\,]_\pm = \mathfrak{B}_{\lambda,\ell}\Big([\,\hat{Q}_{\mathrm{BRST}}, \mathfrak{O}\,]_\pm\Big),$$

i.e. $[\,\hat{Q}_{\mathrm{BRST}}, \mathfrak{B}_{\lambda,\ell}\,]_\pm = 0$ (cf. Lemma 9.59). Thus $\mathfrak{B}_{\lambda,\ell}$ descends to cohomology and BRST–exact contributions are annihilated by $\mathsf{P}_{\mathrm{phys}}$ (cf. Section 6.10).

Step 2 (Nielsen/Slavnov identity at the operator level). On a certified plateau, BRST viability holds ($Q^2_{\mathrm{BRST}} = 0$ and well–defined domains; cf. Definition 9.58). A change of a gauge–fixing parameter $\theta \in \{\xi, \alpha_{\mathrm{gf}}\}$ deforms the gauge–fixed generator through a BRST–exact variation of the gauge–fixing fermion (standard Nielsen identity; cf. Section 7.5). Equivalently, there exists the band–local Nielsen insertion functional $\mathcal{W}_\theta(\cdot)$ (cf. Definition 9.64) such that for any gauge–covariant, ghost–number–0 observable $\mathfrak{O}$,

$$\partial_\theta\, \mathfrak{O} \;=\; [\,\hat{Q}_{\mathrm{BRST}},\, \mathcal{W}_\theta(\mathfrak{O})\,]_\pm.$$

Applying the band map and using Step 1,

$$\partial_\theta\, \mathfrak{B}_{\lambda,\ell}[\mathfrak{O}] = \mathfrak{B}_{\lambda,\ell}\Big(\partial_\theta \mathfrak{O}\Big) = \mathfrak{B}_{\lambda,\ell}\Big([\,\hat{Q}_{\mathrm{BRST}}, \mathcal{W}_\theta(\mathfrak{O})\,]_\pm\Big) = \Big[\,\hat{Q}_{\mathrm{BRST}},\, \mathfrak{B}_{\lambda,\ell}[\mathcal{W}_\theta(\mathfrak{O})]\,\Big]_\pm.$$

Step 3 (projection to H^0 and gauge–parameter independence). With $\mathsf{P}_{\mathrm{phys}}$ the projector onto $\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}})$ (cf. Section 6.10), one has $\mathsf{P}_{\mathrm{phys}}\hat{Q}_{\mathrm{BRST}} = 0 = \hat{Q}_{\mathrm{BRST}}\mathsf{P}_{\mathrm{phys}}$. Therefore for $\mathfrak{O} = \hat{\Sigma}_{\mu\nu}(x)$,

$$\partial_\theta \Big\langle \mathsf{P}_{\mathrm{phys}}\, \mathfrak{B}_{\lambda,\ell}[\hat{\Sigma}_{\mu\nu}(x)]\, \mathsf{P}_{\mathrm{phys}} \Big\rangle = \Big\langle \mathsf{P}_{\mathrm{phys}} \Big[\,\hat{Q}_{\mathrm{BRST}},\; \mathfrak{B}_{\lambda,\ell}[\mathcal{W}_\theta(\hat{\Sigma}_{\mu\nu}(x))]\,\Big]_\pm \mathsf{P}_{\mathrm{phys}} \Big\rangle = 0.$$

Passing to the plateau limit (Step 0) yields $\partial_\theta\, \bar{\Sigma}_{\mu\nu}(x) = 0$, proving the Σ–Nielsen identity for all $\theta \in \{\xi, \alpha_{\mathrm{gf}}\}$.

Step 4 (visibilities and response kernels). Any band–regular functional or kernel that depends on the gauge parameters only through $\bar{\Sigma}$ inherits the same independence: if F is smooth in its tensor argument, then

$$\partial_\theta\, F\Big(\bar{\Sigma}\Big) = \frac{\partial F}{\partial \bar{\Sigma}_{\rho\sigma}}\, \partial_\theta \bar{\Sigma}_{\rho\sigma} = 0.$$

In particular, $\bar{\Sigma}$–based visibilities (operational overlaps/response built from Proposition 9.55 evaluated after realization) and Σ–causal response kernels on H^0 are gauge–parameter independent on $\mathfrak{P}$ (cf. Corollary 9.56 and Section 9.5). □

Definition 9.62 (Relational Levi–Civita and Σ–causality)**.** Let $\bar{\Sigma}_{\mu\nu}$ be the realized chronogeometric tensor obtained from the Σ–realization map (cf. Definition 9.60 and eq. (9.27)) on a certified plateau (cf. Definition 9.58). Denote by $\nabla^{(\mathrm{rel})}$ the torsion–free, $\bar{\Sigma}$–compatible connection (the *Relational Levi–Civita*; cf. Definition 6.7), and by $\bar{g}^{\mathrm{Lor}}$ the Lorentzian projection of the realized metric part $\bar{g}$ (cf. Definition 9.20, eq. (9.4), and Remark 9.15). For any physical quadratic operator $\mathcal{K}_\Psi(\bar{\Sigma})$ (law–kernel built on the physical sector; cf. Definition 9.44 and Section 9.5), the Σ*–causal retarded kernel* is the unique Green function $G^{\mathrm{R}}(x, x')$ supported in the future cone defined by $\bar{\Sigma}$ and satisfying

$$\mathcal{K}_\Psi(\bar{\Sigma})\, G^{\mathrm{R}} = \delta$$

with retarded boundary conditions fixed by $\nabla^{(\mathrm{rel})}$ (cf. Definition 9.22 and Remark 9.15). Gauge–parameter independence of G^{R} on plateaus follows from the Σ–Nielsen identity (cf. Proposition 9.61) and BRST hygiene (cf. Lemma 9.59).

Definition 9.63 (Physical cohomological EOM and retarded kernels)**.** On the BRST–physical Hilbert space $\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}})$ (cf. Section 6.10), the cohomology–restricted equations of motion (EOM) are

$$\mathcal{E}^{\mathrm{coh}}_{\mathbf{\Phi}}(\bar{\Sigma}) = 0, \qquad \mathcal{E}^{\mathrm{coh}}_{\mathbf{\Theta}}(\bar{\Sigma}) = 0, \tag{9.28}$$

defined modulo ζ–exact terms and evaluated with $\bar{\Sigma}$ from Definition 9.60 (cf. the cohomological EOM framework and Ward/Slavnov identities in Section 7.5). The corresponding *physical* retarded kernels,

$$G^{\mathrm{R}}_{\Phi\Phi;\mathrm{phys}}(x, x'), \qquad G^{\mathrm{R}}_{\Theta\Theta;\mathrm{phys}}(x, x'), \tag{9.29}$$

are the Σ–causal Green functions associated to the quadratic operators obtained by linearizing (9.28) about the realized background, as specified in Definition 9.62. They inherit gauge–parameter independence on certified plateaus from Proposition 9.61 and are compatible with graded transport/Hermiticity (cf. Definition 9.7 and Proposition 9.6).

Definition 9.64 (Nielsen insertion functional)**.** Let $\theta \in \{\xi, \alpha_{\mathrm{gf}}, \dots\}$ be a gauge–fixing parameter and ζ the gauge–fixing fermion (cf. Section 7.5). Define the band–local, graded–linear map of ghost number -1,

$$\mathcal{W}_\theta : \ \mathfrak{O} \ \longmapsto \ \mathcal{W}_\theta(\mathfrak{O}), \qquad \mathcal{W}_\theta(\mathfrak{O}) \ \simeq \ \int_{\mathcal{X}_q} (\partial_\theta \zeta) \triangleright \mathfrak{O} \, \mathrm{d}\mu,$$

such that for any gauge–covariant, ghost–number–0 observable $\mathfrak{O}$,

$$\partial_\theta \mathfrak{O} \ = \ [\, \hat{Q}_{\mathrm{BRST}}, \, \mathcal{W}_\theta(\mathfrak{O}) \,]_\pm. \tag{9.30}$$

Band–locality and covariance follow from Definitions 6.280, 9.7 and 9.10; BRST hygiene of the tag gives $\mathfrak{B}_{\lambda,\ell}\big([\, \hat{Q}_{\mathrm{BRST}}, \mathcal{W}_\theta(\mathfrak{O}) \,]_\pm\big) = [\, \hat{Q}_{\mathrm{BRST}}, \, \mathfrak{B}_{\lambda,\ell}[\mathcal{W}_\theta(\mathfrak{O})] \,]_\pm$ (cf. Lemma 9.59).

Lemma 9.65 (Ward/Slavnov identities in Σ–form)**.** *The BRST current improved by* $\bar{\Sigma}$ *satisfies* $\nabla^{(\mathrm{rel})}_\mu J^\mu_{\mathrm{BRST}} = 0$ *in* H^0*. Any residual breaking is ζ–exact and cannot move physical poles/residues of* (9.29)*.*

Proof. (1) Setup: realized background and cohomological EOM. Work on a certified plateau $\mathfrak{P}$ and let $\bar{\Sigma}_{\mu\nu}$ be the realized chronogeometry from the Σ–realization map (cf. Definitions 9.58 and 9.60 and eq. (9.27)). The physical (cohomology–restricted) equations of motion are imposed modulo ζ–exact terms on $\mathcal{H}_{\rm phys} := H^0(\hat{Q}_{\rm BRST})$ (cf. Definition 9.63 and Section 6.10). Denote by $\nabla^{(\rm rel)}$ the torsion–free, $\bar{\Sigma}$–compatible connection (Relational Levi–Civita; cf. Definition 9.62).

(2) BRST current and $\bar{\Sigma}$–improvement. Let $J^{\mu}_{\rm BRST}$ be the Noether current associated with BRST symmetry in the gauge–fixed (but BRST–invariant) formulation (cf. Section 7.5). To maintain covariance on the realized background, add the standard improvement term fixed by $\nabla^{(\rm rel)}$ (Belinfante–type completion in the $\bar{\Sigma}$ geometry), yielding the $\bar{\Sigma}$–improved current—denoted again $J^{\mu}_{\rm BRST}$—which transforms covariantly under graded transport (cf. Definition 9.7 and Proposition 9.6).

(3) Slavnov–Taylor identity and ζ–exact divergence. BRST invariance of the gauge–fixed generator implies the Ward/Slavnov identity: the divergence of the BRST current equals a BRST–exact insertion controlled by the gauge–fixing fermion ζ (cf. Section 7.5). Equivalently, using the Nielsen insertion functional $\mathcal{W}_\theta(\cdot)$ of Definition 9.64 and writing it schematically as $\mathcal{W}(\zeta;\bar{\Sigma})$ on the realized background, one has

$$\nabla^{(\rm rel)}_{\mu} J^{\mu}_{\rm BRST} = [\,\hat{Q}_{\rm BRST},\, \mathcal{W}(\zeta;\bar{\Sigma})\,]_{\pm},$$

where the use of $\nabla^{(\rm rel)}$ ensures background–covariance (cf. Definition 9.62).

(4) Projection to H^0 and conservation. Since $\mathsf{P}_{\rm phys}$ projects onto $H^0(\hat{Q}_{\rm BRST})$ and annihilates BRST–exact operators, $\mathsf{P}_{\rm phys}\,[\hat{Q}_{\rm BRST},\cdot]_{\pm}\,\mathsf{P}_{\rm phys} = 0$ (cf. Section 6.10 and Lemma 9.59). Therefore, in H^0,

$$\nabla^{(\rm rel)}_{\mu} J^{\mu}_{\rm BRST} = 0,$$

which is the desired Ward/Slavnov identity in Σ–form.

(5) Consequence for poles/residues of the physical retarded kernels. Residual breakings are ζ–exact by Step (3). Insertions of BRST–exact operators cannot alter physical matrix elements on H^0 and, in particular, cannot shift the locations or residues of poles of the Σ–causal retarded kernels $G^{\rm R}_{\Phi\Phi;\rm phys}$, $G^{\rm R}_{\Theta\Theta;\rm phys}$ (cf. eq. (9.29) and Definition 9.62). Moreover, on a certified plateau the Σ–Nielsen identity guarantees gauge–parameter independence of $\bar{\Sigma}$ (and hence of any $\bar{\Sigma}$–dependent kernel) (cf. Proposition 9.61), so neither gauge scans nor ζ–exact redefinitions can move physical poles/residues. This establishes both claims. □

Definition 9.66 (Σ–projected two–point functions of band modes)**.** Let $\hat{A}^{\dagger}_a, \hat{K}^{\dagger}_b$ denote the pre–physical Apeiron/Kymon creation operators obtained from the Σ–weighted principal modes and block factorizations (cf. Proposition 9.36 and eqs. (9.15) to (9.17); see also Remark 9.41 and Corollary 9.40). Define their Σ–projected *physical* two–point functions by

$$\Pi^{(\Phi)}_{ab}(p) \;:=\; \Big\langle \mathcal{U}_\Sigma[\hat{A}_a]\,\mathcal{U}_\Sigma[\hat{A}^{\dagger}_b] \Big\rangle(p), \qquad \Pi^{(\Theta)}_{cd}(p) \;:=\; \Big\langle \mathcal{U}_\Sigma[\hat{K}_c]\,\mathcal{U}_\Sigma[\hat{K}^{\dagger}_d] \Big\rangle(p), \tag{9.31}$$

where $\mathcal{U}_\Sigma[\cdot]$ is the Σ–realization map (cf. Definition 9.60) evaluated on a certified plateau $\mathfrak{P}$ (cf. Definition 9.58), and the expectation $\langle\cdot\rangle$ is the physical Born/GNS pairing on $\mathcal{H}_{\rm phys} := H^0(\hat{Q}_{\rm BRST})$ (cf. Section 6.10 and Definitions 1.18 and 1.34). The momentum

argument p is defined via the $\bar{\Sigma}$–compatible (Relational Levi–Civita) structure used for Fourier/Green analysis (cf. Definition 9.62). Well–posedness and gauge–parameter independence on $\mathfrak{P}$ follow from BRST hygiene of the band map and the Σ–Nielsen identity (cf. Lemma 9.59 and Proposition 9.61), and covariance from the graded transport and Fourfold Hermiticity/covariance (cf. Definition 9.7 and Proposition 9.6).

Theorem 9.67 (Realization criterion and spectrum extraction)**.** *A pre–physical mode operator* $\hat{M}^\dagger \in \{\hat{A}_a^\dagger, \hat{K}_b^\dagger\}$ *realizes as a physical particle iff:*

R1 $\Pi(p)$ *from Definition 9.66 contributes a* Σ–causal retarded pole *to the corresponding kernel in* (9.29) *at* $p = p_\star$*, and* $\partial_{\xi,\alpha_{\rm gf}} p_\star = 0$ *(Nielsen–flatness on certified plateaus; cf. Definition 9.58 and Proposition 9.61);*

R2 *the residue at* $p_\star$ *is positive in* $H^0(\hat{Q}_{\rm BRST})$ *(Kugo–Ojima unitarity; cf. Section 6.10), and the mode's propagation respects the* $\bar{\Sigma}$*–cones (hyperbolicity; cf. Definition 9.62).*

When R1 and R2 hold:

(i) Geometry block

Σ*–null (pure phase–flow) directions in* $\mathcal{N}_\Phi^{\rm pre}$ *yield a* massless spin–2 graviton*;* Σ*–radial (amplitude) directions yield* massive Gaiaton (Apeiron) *modes (cf. Proposition 9.36 and Lemma 9.38).*

(ii) Matter block

Σ*–principal modes in* $\mathcal{N}_\Theta^{\rm pre}$ *realize as the* Kymon *spectrum; isolating the torsional temporal sub–block gives the* Chrono–Kymon *(cf. Propositions 9.36 and 9.39 and Remark 9.41).*

Proof. (1) From pre–physical modes to realized two–point functions and kernels. Pre–physical principal modes are obtained by the Σ–weighted eigenproblems and block factorizations (cf. Proposition 9.36 and eqs. (9.15) to (9.17)). Their physical images are defined via the Σ–realization map $\mathcal{U}_\Sigma[\cdot]$ on certified plateaus (cf. Definitions 9.58 and 9.60). The Σ–projected two–point functions $\Pi_{ab}^{(\Phi)}, \Pi_{cd}^{(\Theta)}$ are given in Definition 9.66; the physical retarded kernels $G^{\rm R}_{\Phi\Phi;{\rm phys}}, G^{\rm R}_{\Theta\Theta;{\rm phys}}$ are the Σ–causal Green functions associated with the cohomology–restricted EOM (cf. Definitions 9.62 and 9.63). Standard linear–response/spectral relations on the realized background (Källén–Lehmann–type correspondence in the Σ–geometry) imply that isolated singularities (poles) of Π contribute the same mass–shell singularities to the corresponding retarded kernel (compatibly with graded transport/Hermiticity; cf. Proposition 9.6).

(2) Nielsen–flatness and gauge–parameter independence. On a certified plateau, the Σ–Nielsen identity ensures $\partial_{\xi,\alpha_{\rm gf}} \bar{\Sigma} = 0$ (cf. Proposition 9.61). Since the Σ–causal operators and dispersion relations depend on the gauge parameters only through $\bar{\Sigma}$ (cf. Definition 9.62), the location $p_\star$ of any pole inherited from Π is gauge–parameter independent: $\partial_{\xi,\alpha_{\rm gf}} p_\star = 0$. Ward/Slavnov identities in Σ–form further ensure that any residual breaking is ζ–exact and cannot shift physical poles/residues in H^0 (cf. Lemma 9.65).

(3) Positivity and hyperbolicity. The residue at $p_\star$ equals the norm (matrix element) of the one–particle state created by the realized operator $\mathcal{U}_\Sigma[\hat{M}^\dagger]$ in the physical Hilbert space; positivity of this residue is precisely the Kugo–Ojima unitarity condition in $H^0(\hat{Q}_{\rm BRST})$

(cf. Section 6.10). By construction of $\nabla^{(\mathrm{rel})}$ and Σ–causal Green functions, propagation is supported inside the $\bar{\Sigma}$–future cone (hyperbolicity; cf. Definition 9.62); thus the realized excitation respects the realized chronogeometric cones.

(4) $\Rightarrow$ If R1 and R2 hold, $\hat{M}^\dagger$ realizes as a particle. Assume an isolated Σ–causal retarded pole at $p_\star$ with gauge–independent location and positive residue in H^0. Then the LSZ–type argument adapted to the realized background identifies a stable (or metastable) one–particle contribution in $G^{\mathrm{R}}_{\bullet\bullet;\mathrm{phys}}$ with causal propagation; acting with $\mathcal{U}_\Sigma[\hat{M}^\dagger]$ on the physical vacuum creates a non–null physical state with that mass–shell. Hence the mode realizes as a physical particle.

(5) $\Leftarrow$ If a mode realizes as a particle, then R1 and R2 hold. Conversely, a realized particle contributes an isolated mass–shell pole to the corresponding retarded kernel with non–negative residue. Gauge–parameter independence of $p_\star$ follows from Proposition 9.61; positivity of the residue follows from Kugo–Ojima positivity on H^0; and hyperbolicity follows from the Σ–causal construction in Definition 9.62. Therefore both (R1) and (R2) are necessary.

(6) Blockwise identification of species. In the geometry block, the Σ–weighted principal directions decompose into *phase–flow* (Σ–null) and *amplitude* (Σ–radial) modes (cf. Lemma 9.38). When (R1)–(R2) hold, the former realize as a massless spin–2 graviton, while the latter produce massive Gaiaton (Apeiron) excitations. In the matter block, Σ–principal modes realize as Kymons, and the torsional temporal sub–block isolates the Chrono–Kymon channels, with inter–modal mixing governed by $\hat{G}_{\Theta\Theta}$ and weighted correctly by $\hat{t}_{\mu\nu}$ (cf. Proposition 9.39 and Remark 9.41). Stability of the extracted spectra under small band deformations follows from Lemma 9.42.

Combining (1)–(6) establishes the stated iff criterion and the blockwise spectrum identification. □

Corollary 9.68 (Cross–block consistency)**.** *On certified plateaus $\mathfrak{P}$, off–diagonal Σ–couplings $\hat{\Sigma}_{\Phi\Theta}$ either vanish in H^0 or generate BRST–admissible mixing angles; in either case, realized masses/dispersion relations are gauge–parameter independent.*

Proof. (1) Setting and notation. Work on a certified plateau and let $\bar{\Sigma}$ be the realized chronogeometry (cf. Definitions 9.58 and 9.60 and eq. (9.27)). Consider the cohomology–restricted quadratic operator for the combined geometry–matter sector, written in block form relative to $(\Phi \oplus \Theta)$,

$$\mathcal{K}_\Psi(\bar{\Sigma}) \;=\; \begin{pmatrix} \mathcal{K}_{\Phi\Phi}(\bar{\Sigma}) & \mathcal{K}_{\Phi\Theta}(\bar{\Sigma}) \\ \mathcal{K}_{\Theta\Phi}(\bar{\Sigma}) & \mathcal{K}_{\Theta\Theta}(\bar{\Sigma}) \end{pmatrix},$$

whose Σ–causal retarded Green operator yields $G^{\mathrm{R}}_{\Phi\Phi;\mathrm{phys}}$ and $G^{\mathrm{R}}_{\Theta\Theta;\mathrm{phys}}$ after projection (cf. Definitions 9.62 and 9.63). The pre–physical principal modes and their Σ–weighted factorizations provide the band modes $\hat{A}^\dagger_a, \hat{K}^\dagger_b$ and the Σ–projected two–point functions $\Pi^{(\Phi)}_{ab}, \Pi^{(\Theta)}_{cd}$ (cf. Proposition 9.36, eqs. (9.16) and (9.17), and Definition 9.66); inter–modal mixing is treated by the generalized eigenproblem (cf. Proposition 9.39).

(2) BRST screening of cohomologically trivial off–diagonal pieces. If the off–diagonal coupling $\hat{\Sigma}_{\Phi\Theta}$ (or the corresponding entry $\mathcal{K}_{\Phi\Theta}$) is BRST–exact on the band, then by BRST hygiene of the tag and projection (cf. Lemma 9.59 and Section 6.10) its physical matrix

elements vanish:

$$\mathsf{P}_{\mathrm{phys}}\,\mathfrak{B}_{\lambda,\ell}\Big([\,\hat{Q}_{\mathrm{BRST}},Y]_{\pm}\Big)\,\mathsf{P}_{\mathrm{phys}}=0.$$

Hence the off–diagonal block decouples in H^0, and the realized dispersion relations are determined by the diagonal blocks alone.

(3) BRST–admissible mixing when the off–diagonal is BRST–closed. If $\hat{\Sigma}_{\Phi\Theta}$ is BRST–closed (ghost number 0) but not exact, then it survives projection and induces a physical mixing. On $\mathcal{H}_{\mathrm{phys}}$, diagonalize the realized quadratic form with respect to the Σ–inner product defined by $\bar{\Sigma}$ (cf. Definition 9.62 and Proposition 9.61). By Proposition 9.39, the generalized eigenproblem absorbs cross–terms and yields a Σ–orthonormal transformation U on H^0 (block–unitary with respect to the Σ–metric) that brings $\mathcal{K}_\Psi(\bar{\Sigma})$ to block–diagonal form. The (physical) *mixing angles* are those parameterizing U inside the relevant two–mode subspaces; being constructed from BRST–closed, gauge–covariant data and evaluated on H^0, these angles are *BRST–admissible*.

(4) Gauge–parameter independence of realized spectra. On a certified plateau, $\partial_\theta\bar{\Sigma}=0$ for every gauge–fixing parameter $\theta\in\{\xi,\alpha_{\mathrm{gf}}\}$ by the Σ–Nielsen identity (cf. Proposition 9.61). Therefore $\partial_\theta\mathcal{K}_\Psi(\bar{\Sigma})=0$, and the spectral projectors/eigenvalues of the realized operator are θ–independent. Equivalently, Ward/Slavnov identities in Σ–form imply that any θ–variation is BRST–exact and thus annihilated between $\mathsf{P}_{\mathrm{phys}}$ insertions (cf. Lemma 9.65 and Section 7.5). Consequently, the pole locations and residues extracted from the physical retarded kernels (9.29) are gauge–parameter independent, whether or not a BRST–admissible mixing is present.

Combining (2)–(4): either the off–diagonal couplings vanish in cohomology (decoupling) or they generate BRST–admissible, Σ–orthonormal mixings; in both cases, the realized masses/dispersion relations are gauge–parameter independent on $\mathfrak{P}$. This proves the corollary. □

Proposition 9.69 (Quadrant mapping under realization)**.** *Let* $(\Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}},\Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}})$ *be the pre–physical order parameters of Definition 9.47, with classification by Definition 9.49. On certified plateaus (cf. Definition 9.58):*

(i) QI→QI
Σ–contrasts stay $\mathcal{O}(\varepsilon)$; only BRST–closed modes can realize.

(ii) QII→QII/QI
Law symmetry persists; state asymmetry survives iff not ζ–exact; realized NG/pNG structure is gauge–independent.

(iii) QIII/QIV
Any BRST–forbidden law asymmetry is annihilated by $\mathsf{P}_{\mathrm{phys}}$; otherwise it persists as explicit breaking visible in Σ–Ward identities (Lemma 9.65).

Proof. (1) Band stability and BRST hygiene. On a certified plateau $\mathfrak{P}$, band–local quantities vary by at most $\mathcal{O}(\varepsilon_{\mathrm{plat}})$ (grid diagnostics and acceptance; cf. Definition 9.58, eqs. (7.394) to (7.396), and Proposition 7.203), and the pre–physical Σ–contrasts are finite/continuous in probes and band knobs (cf. Lemmas 9.27 and 9.46). The band tag $\mathfrak{B}_{\lambda,\ell}$ is a BRST chain map and commutes with projection to cohomology (cf. Lemma 9.59 and Section 6.10); hence BRST–exact contributions are removed by $\mathsf{P}_{\mathrm{phys}}$ in the realization map $\mathcal{U}_\Sigma$ (cf. Definition 9.60).

(2) Law channel under realization. The law–channel visibility is Nielsen–flat on plateaus (cf. Corollary 9.56), and $\bar{\Sigma}$ is gauge–parameter independent (cf. Proposition 9.61). Therefore $\Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}} = \mathcal{O}(\varepsilon)$ remains $\mathcal{O}(\varepsilon)$ after realization, and the realized retarded kernels depend on gauge parameters only through $\bar{\Sigma}$, hence are gauge–parameter independent (cf. Definitions 9.62 and 9.63).

(3) State channel under realization. State–channel asymmetries that are BRST–exact are annihilated by $\mathsf{P}_{\mathrm{phys}}$ (cf. Lemma 9.59); BRST–closed asymmetries survive. In particular, phase–flow (Goldstone–like) vs. amplitude (Higgs/Gaiaton–like) content diagnosed pre–physically by Σ persists through realization (cf. Lemma 9.38 and Theorem 9.67), with gauge independence ensured by Proposition 9.61.

(4) Case analysis of the quadrant map.

QI (law & state symmetric).
Pre–physical $\Delta^{\mathrm{pre},\Sigma}_{\mathrm{law}}, \Delta^{\mathrm{pre},\Sigma}_{\mathrm{state}} < \varepsilon$ remain within $\mathcal{O}(\varepsilon)$ under plateau–controlled realization by (1)–(3). Only BRST–closed mode content can survive; any BRST–exact remnants are removed by $\mathsf{P}_{\mathrm{phys}}$ (cf. Lemma 9.59). Hence QI maps to QI.

QII (law symmetric, state asymmetric).
Law symmetry persists by (2). If the state asymmetry is BRST–exact, projection removes it, yielding QI; if it is BRST–closed, it survives as QII. The realized NG/pNG structure (phase–flow vs. amplitude and possible small explicit breakings) is determined by Lemma 9.38 and Proposition 9.39 and realized as physical poles/residues by Theorem 9.67, with gauge–parameter independence from Proposition 9.61.

QIII/QIV (law asymmetric).
If the law asymmetry is BRST–forbidden (cohomologically trivial), it is annihilated by projection (cf. Lemma 9.59), so the realized law channel reverts to symmetry and the classification follows the surviving state channel. If it is BRST–admissible (BRST–closed), it persists as explicit breaking and appears as a non–vanishing divergence/source in the Σ–Ward/Slavnov identities (cf. Lemma 9.65); nevertheless, realized pole locations/residues remain gauge–parameter independent by Proposition 9.61.

(5) Consistency with spectra and mixing. Any realized particle must satisfy the iff criterion of Theorem 9.67; cross–block effects are either removed in cohomology or produce BRST–admissible Σ–orthonormal mixings without gauge drift (cf. Corollary 9.68). Thus the quadrant mapping is consistent with spectrum extraction.

Combining (1)–(5) yields the stated mapping and properties. □

Lemma 9.70 (Health checklist (Σ–centric))**.** *On any certified plateau $\mathfrak{P}$, realized data satisfy: (i) $\det \bar{g}^{\mathrm{Lor}} < 0$ and a fixed time orientation (Lorentzianity); (ii) Σ–causality/hyperbolicity for the physical retarded kernels (9.29); (iii) Nielsen–flatness of poles/residues; (iv) spectral finiteness on the band (no UV–induced pole drift on $\mathfrak{P}$); (v) unitarity (positive residues in H^0).*

Proof. (i) Lorentzianity. By the definition of a certified plateau, $\bar{\Sigma}$ admits a Lorentzian projection and defines causal cones (cf. Definition 9.58). The Relational Levi–Civita construction fixes $\bar{g}^{\mathrm{Lor}}$ and its time orientation consistently with $\bar{\Sigma}$ (cf. Definition 9.62); hence $\det \bar{g}^{\mathrm{Lor}} < 0$ and the time orientation is globally fixed on $\mathfrak{P}$.

Algorithm 2 Σ-aware upgrade and spectrum readout

1. **Preconditions:** verify $\mathfrak{P}$ (Definition 9.58), BRST hygiene (Lemma 9.59), Σ-regularity.
2. **Realize Σ:** compute $\bar{\Sigma}$ via (9.27); build $\nabla^{(\mathrm{rel})}$ and $\bar{g}^{\mathrm{Lor}}$ (Definition 9.62).
3. **Physical EOM:** form $\mathcal{E}^{\mathrm{coh}}_{\boldsymbol{\Phi}/\boldsymbol{\Theta}}(\bar{\Sigma})$ and $\mathcal{K}_{\Psi}(\bar{\Sigma})$ (Definition 9.63); construct $G^{\mathrm{R}}_{\bullet;\mathrm{phys}}$ with Σ-causal support.
4. **Modes → particles:** evaluate $\Pi^{(\Phi/\Theta)}(p)$ from (9.31); locate Σ-retarded poles and residues; test **(R1)**–**(R2)** of Theorem 9.67.
5. **Health checks:** apply Lemma 9.70; report the graviton/Gaiaton/Kymon ledger with mixing constraints (Corollary 9.68).

(ii) Σ–causality/hyperbolicity of $G^{\mathrm{R}}_{\bullet\bullet;\mathrm{phys}}$. By Definition 9.62, the physical retarded kernels (9.29) are the unique Σ–causal Green functions supported in the $\bar{\Sigma}$–future cone and defined with boundary conditions from $\nabla^{(\mathrm{rel})}$. Therefore they satisfy hyperbolicity and causal support on the realized background.

(iii) Nielsen–flatness. On $\mathfrak{P}$, the Σ–Nielsen identity gives $\partial_{\xi,\alpha_{\mathrm{gf}}}\bar{\Sigma} = 0$ (cf. Proposition 9.61). Since the kernels and dispersion data depend on gauge parameters only through $\bar{\Sigma}$ (cf. Definitions 9.62 and 9.63), their pole locations and residues are gauge–parameter independent. Ward/Slavnov identities in Σ–form ensure that any residual breaking is ζ–exact and thus invisible in H^0 (cf. Lemma 9.65).

(iv) Spectral finiteness and no UV–induced drift on $\mathfrak{P}$. Spectral Smoothness and the Principle of Finite Entanglement Capacity imply endogenous regularization: triads and the derived operators $\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu}, \hat{\Sigma}_{\mu\nu}$ are finite on the band (cf. Propositions 9.13 and 9.14 and Remark 9.21). Continuity under band deformations yields $\mathcal{O}(\varepsilon)$ control for observables and mode spectra (cf. Lemmas 9.27 and 9.42); the plateau acceptance rule maintains these bounds uniformly across $\mathfrak{P}$ (cf. Definition 9.58 and Proposition 7.203). Hence there is no UV–induced pole drift within $\mathfrak{P}$.

(v) Unitarity (positive residues in H^0). Realized particles satisfy the iff criterion of Theorem 9.67: their poles are Σ–causal and the residues are positive on the BRST–physical Hilbert space $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ (Kugo–Ojima positivity; cf. Section 6.10 and Theorem 9.67). Therefore unitarity holds for the realized spectrum.

All items (i)–(v) are thus guaranteed on any certified plateau $\mathfrak{P}$, completing the proof. □

Summary of Section 9.5. The realization stage is completed in a Σ–centric and BRST–clean manner as follows. (i) The *unified* chronogeometry $\hat{\Sigma}_{\mu\nu} = \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu}$, defined band–locally by SOT–C^2 Hessians, is promoted to its realized tensor $\bar{\Sigma}_{\mu\nu}$ via the Σ–realization map $\mathcal{U}_{\Sigma}$ on certified plateaus $\mathfrak{P}$ (cf. Definitions 9.20, 9.58 and 9.60 and eq. (9.4)), with BRST hygiene of the tag guaranteeing cohomological well–posedness (cf. Lemma 9.59). (ii) Physical dynamics are posed as cohomology–restricted EOM together with Σ–*causal* retarded kernels supported

within the $\bar{\Sigma}$–cones (cf. Definitions 9.62 and 9.63 and eq. (9.29)). Ward/Slavnov identities in Σ–form ensure that any residual breaking is ζ–exact and therefore invisible on $\mathcal{H}_{\text{phys}}$ (cf. Lemma 9.65). (iii) Spectrum extraction proceeds from Σ–projected two–point functions of band modes to Σ–causal retarded poles with gauge–independent locations and residues: gauge–parameter independence follows from the Σ–Nielsen identity (cf. Definition 9.66 and Proposition 9.61), while positivity of residues is the Kugo–Ojima unitarity condition on $H^0(\hat{Q}_{\text{BRST}})$ (cf. Section 6.10 and Theorem 9.67). Cross–block effects are either removed in cohomology or appear as BRST–admissible mixing angles that do not induce gauge drift (cf. Corollary 9.68).

Within this framework, Σ–weighted principal modes characterize the *provisional* particle assignments, subject to all health checks (cf. Proposition 9.36 and Lemma 9.70). Tangential (Σ–null, pure phase–flow) directions in the geometry block provisionally correspond to a *massless spin–2 graviton*; radial (Σ–amplitude) directions yield *massive Gaiaton (Apeiron)* modes (cf. Lemma 9.38 and Theorem 9.67). In the matter block, Σ–principal modes provisionally realize as the *Kymon* spectrum, with a torsional temporal sub–block isolating the *Chrono–Kymon*, and inter–modal mixing handled coherently by the generalized eigenproblem (cf. Proposition 9.39 and Remark 9.41). All identifications are *plateau–qualified*: Lorentzianity and causal support are enforced by the Relational Levi–Civita structure (cf. Definition 9.62); poles/residues are Nielsen–flat on $\mathfrak{P}$ (cf. Proposition 9.61 and Corollary 9.56); spectral finiteness and stability under band deformations are controlled by endogenous regularization and plateau diagnostics (cf. Proposition 9.13, Lemmas 9.27 and 9.42, and Definition 9.58); and unitarity is certified in H^0 by residue positivity (cf. Theorem 9.67). Consequently, the realized sector inherits the quadrant–level logic consistently with pre–physical diagnostics (cf. Proposition 9.69), while keeping all particle–level assignments explicitly *provisional* pending verification across certified plateaus.

9.6 Minimal "How-to" Pipeline

Definition 9.71 (Inputs and knobs)**.** The pipeline takes as input the Fourfold inter-modal correlator $\hat{\mathbf{G}}_\Lambda$ and gauge-covariant transport $\mathcal{U}$ (Definitions 9.5 and 9.7), the graded modal structure (Definition 9.8), the band tag $\mathfrak{B}_{\lambda_{\text{UV}},\ell}$ (Definition 9.10), and, if used, an admissible spectral operator $\hat{D}$ satisfying the spectral discipline (Proposition 9.13).

Definition 9.72 (Band choice and Σ-regularity check)**.** Choose $(\lambda_{\text{UV}}, \ell)$ with Spectral Smoothness and Finite Entanglement Capacity (Proposition 9.13). On the band, form $\hat{\mathbf{G}}_\Lambda^{(\lambda,\ell)} := \mathfrak{B}_{\lambda,\ell}[\hat{\mathbf{G}}_\Lambda]$ and verify that the strong-operator Hessians at coincidence exist for D^2, T^2, hence $\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu}$ and

$$\hat{\Sigma}_{\mu\nu} := \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu}, \qquad \hat{\Sigma}^{(\Phi)} := \Pi_{\Phi\Phi}^\dagger \hat{\Sigma}\, \Pi_{\Phi\Phi}, \quad \hat{\Sigma}^{(\Theta)} := \Pi_{\Theta\Theta}^\dagger \hat{\Sigma}\, \Pi_{\Theta\Theta} \tag{9.32}$$

(Definition 9.20).

Definition 9.73 (Σ-order parameters and quadrant classification)**.** Fix probe directions $\{v_k\} \subset T_x\mathcal{X}_q$ and a band-local, gauge-invariant functional $\mathcal{I}$. With $\hat{Q}_x[v] := v^\mu v^\nu \hat{\Sigma}_{\mu\nu}(x)$

(Definition 9.22), define

$$\begin{aligned}\Delta^{\Sigma}_{\mathrm{law}}(g;x;v_k) &:= \Big(\mathcal{I}[\hat{Q}_x[v_k] - \mathrm{Ad}_{g(x)}\,\hat{Q}_x[v_k]]\Big)^{1/2},\\ \Delta^{\Sigma}_{\mathrm{state}}(g;x;v_k) &:= \Big|\mathcal{I}_{\Psi}[\hat{Q}_x[v_k]] - \mathcal{I}_{g\cdot\Psi}[\hat{Q}_x[v_k]]\Big|^{1/2}.\end{aligned} \tag{9.33}$$

and aggregate as in Definition 9.47:

$$\Delta^{\Sigma}_{\mathrm{law}}(g;x) := \Big(\sum_k [\Delta^{\Sigma}_{\mathrm{law}}(g;x;v_k)]^2\Big)^{1/2}, \quad \Delta^{\Sigma}_{\mathrm{state}}(g;x) := \Big(\sum_k [\Delta^{\Sigma}_{\mathrm{state}}(g;x;v_k)]^2\Big)^{1/2}. \tag{9.34}$$

Classify QI–QIV with tolerance ε as in Definition 9.49.

Definition 9.74 (Σ-weighted broken-sector mode extraction)**.** For broken generators X determined by Definition 9.73, compute the graded tangents on blocks (Definition 9.32), $\delta_X\hat{G}_{\Phi\Phi}$, $\delta_X\hat{G}_{\Theta\Theta}$, form the Gram operators (eq. (9.15))

$$C_\Phi = (\delta_X\hat{G}_{\Phi\Phi})^\dagger\hat{\Sigma}^{(\Phi)}\delta_X\hat{G}_{\Phi\Phi}, \qquad C_\Theta = (\delta_X\hat{G}_{\Theta\Theta})^\dagger\hat{\Sigma}^{(\Theta)}\delta_X\hat{G}_{\Theta\Theta},$$

and solve the generalized eigenproblems (eq. (9.16))

$$C_\Phi u_a = \lambda_a\hat{\Sigma}^{(\Phi)}u_a, \qquad C_\Theta v_b = \mu_b\hat{\Sigma}^{(\Theta)}v_b.$$

Define $\hat{A}^\dagger_a, \hat{K}^\dagger_b$ as in Definition 9.37, with Σ-orthonormality (9.18).

Lemma 9.75 (Σ-centric pre-physical health checks)**.** *If $\hat{\Sigma}$ passes the band regularity of Definition 9.72, then: (i) Σ-cones are well-defined and respected by the principal directions (Lemma 9.38); (ii) Δ^Σ in (9.34) varies continuously under small band deformations (Lemma 9.27); (iii) principal spectra $\{\lambda_a, \mu_b\}$ are stable (Lemma 9.42).*

Definition 9.76 (Certified plateau and Σ-realization)**.** Identify a certified plateau $\mathfrak{P}$ (Definition 9.58). Define the Σ-realization map $\mathcal{U}_\Sigma$ by (9.26). Obtain the physical chronogeometry $\bar{\Sigma}_{\mu\nu} = \langle\mathcal{U}_\Sigma[\hat{\Sigma}_{\mu\nu}]\rangle$ (Definition 9.60) and build the Σ-causal retarded kernels $G^{\mathrm{R}}_{\Phi\Phi;\mathrm{phys}}, G^{\mathrm{R}}_{\Theta\Theta;\mathrm{phys}}$ (Definitions 9.62 and 9.63).

Theorem 9.77 (Pipeline soundness)**.** *Assume $\mathfrak{P}$ satisfies Definition 9.58 and the BRST hygiene of the band map (Lemma 9.59). Then the outputs of Definition 9.76 are: (a) gauge-parameter independent (Proposition 9.61); (b) Σ-causal and hyperbolic (Definition 9.62); (c) unitary in $H^0(\hat{Q}_{\mathrm{BRST}})$ with positive residues at realized poles (Theorem 9.67).*

Corollary 9.78 (Spectrum readout rule)**.** *Let $\hat{M}^\dagger \in \{\hat{A}^\dagger_a, \hat{K}^\dagger_b\}$. If $\Pi(p) = \langle\mathcal{U}_\Sigma[\hat{M}]\mathcal{U}_\Sigma[\hat{M}^\dagger]\rangle$ (cf. Definition 9.66) produces a Nielsen-flat Σ-retarded pole with positive residue, then $\hat{M}^\dagger$ realizes as: (i) massless spin-2 (graviton) for Σ-null geometry directions; (ii) massive Gaiaton for Σ-radial geometry directions; (iii) Kymon (incl. Chrono–Kymon) for matter-block principal modes, with BRST-admissible mixings (Theorem 9.67 and Corollary 9.68).*

Remark 9.79 (Triadic attribution and block sensitivity)**.** For interpretability, decompose $\bar{\Sigma} = \bar{\Sigma}^{\mathrm{corr}} + \bar{\Sigma}^{\mathrm{vac}} + \bar{\Sigma}^{\mathrm{spec}}$ and attribute quadrant decisions and realized poles to channels (Proposition 9.48). Block-resolved diagnostics remain available via $\hat{\Sigma}^{(\Phi)}, \hat{\Sigma}^{(\Theta)}$ (Definition 9.33).

Algorithm 3 Σ-aware minimal pipeline

1. **Band & regularity** (Definition 9.72): pick $(\lambda_{\mathrm{UV}}, \ell)$, build $\hat{\mathbf{G}}_{\Lambda}^{(\lambda,\ell)}$; certify SOT–Hessians for $\hat{g}, \hat{t}$.
2. **Build** Σ (Definition 9.72): form $\hat{\Sigma}$ and its block restrictions (9.32).
3. **Order parameters** (Definition 9.73): compute $\Delta^{\Sigma}_{\mathrm{law}}$, $\Delta^{\Sigma}_{\mathrm{state}}$ and classify QI–QIV.
4. **Modes** (Definition 9.74): solve the Σ-weighted eigenproblems; construct $\hat{A}^{\dagger}_{a}$, $\hat{K}^{\dagger}_{b}$.
5. **Pre-physical health** (Lemma 9.75): check Σ-cone compatibility, Σ-weight positivity, stability under band deformations.
6. **Plateau** (Definition 9.76): certify $\mathfrak{P}$ (BRST viability, Nielsen-flatness, Σ-causality).
7. **Σ-realization** (Definition 9.76): apply $\mathcal{U}_{\Sigma}$ to $\hat{\Sigma}$, $\hat{A}^{\dagger}_{a}$, $\hat{K}^{\dagger}_{b}$; obtain $G^{\mathrm{R}}_{\bullet;\mathrm{phys}}$.
8. **Spectrum** (Corollary 9.78): locate Σ-retarded poles/residues; enforce Ward/Slavnov and Kugo–Ojima conditions.

What this achieves (summary). The pipeline implements a *quadrant-aware PSR calculus* using Σ-based order parameters (Definition 9.73), a *graded-consistent, Σ-weighted* construction of Apeiron/Kymon band operators (Definition 9.74), and a *Σ-aware realization* (Definition 9.76) that produces a gauge-clean, causal, unitary spectrum in $H^0(\hat{Q}_{\mathrm{BRST}})$ (Theorem 9.77 and Corollary 9.78).

9.7 Invariance of Chronogeometric Formalisms

Definition 9.80 (The Σ–realization)**.** Define the physical chronogeometry by the Σ–realization map of Definition 9.60:

$$\boxed{\bar{\Sigma}_{\mu\nu}(x) := \Big\langle \mathsf{P}_{\mathrm{phys}} \lim_{(\lambda,\ell)\in\mathfrak{P}} \mathfrak{B}_{\lambda,\ell}\big[\hat{\Sigma}_{\mu\nu}(x)\big]\, \mathsf{P}_{\mathrm{phys}} \Big\rangle} \tag{9.35}$$

All subsequent constructions in Chapters 7–8 (cohomological EOM, linear response, visibilities, Ward/Slavnov identities) are to be understood as functions of $\bar{\Sigma}$. This statement references Definitions 9.10, 9.20, 9.58 and 9.60.

Remark 9.81 (Upstream formalisms are unchanged)**.** Let Q_{BRST} be the BRST charge of Chapter 7 and $H^0(Q_{\mathrm{BRST}})$ the physical sector. Then:

1. **BRST sector:** $Q^2_{\mathrm{BRST}} = 0$, $\mathcal{H}_{\mathrm{phys}} = H^0(Q_{\mathrm{BRST}})$; gauge fixing is ζ-exact; physical observables are defined modulo ζ-exact terms.

2. **Field/bundle scaffolding:** The generalized field operator $\hat{\mathbf{F}} = (\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}})^{\mathsf{T}}$, the Fourfold correlator $\hat{\mathbf{G}}_{\Lambda}$, the operator-bundle connection/curvature, and the graded modal algebra of Definitions 9.5, 9.7 and 9.8 persist verbatim.
3. **Relational geometry (physical side):** The Relational Levi–Civita connection $\nabla^{(\mathrm{rel})}$ and Lorentzian projection of $\bar{g}$ (implicit in $\bar{\Sigma}$) continue to govern linear response/propagators and Ward/Slavnov identities, with Kugo–Ojima unitarity checks unchanged; see Definitions 9.62 and 9.63.

Remark 9.82 (Σ–Nielsen identities: same proofs, stronger statement)**.** On certified plateaus $\mathfrak{P}$ (Definition 9.58), $\bar{\Sigma}$ is gauge–parameter independent by Proposition 9.61. Therefore any quantity in Chapters 7–8 that was previously shown gauge–parameter independent (Nielsen-safe) remains so when rewritten as a functional of $\bar{\Sigma}$; only the *wording* changes (from "triads" to "$\bar{\Sigma}$").

Remark 9.83 (Σ–causal linear response (statement only))**.** Replace the background "metric+temporal data" by $\bar{\Sigma}$ from (9.35). Then the quadratic operators $\mathcal{K}_{\Psi}(\bar{\Sigma})$ and the physical retarded kernels $G^{\mathrm{R}}_{\bullet;\mathrm{phys}}$ defined in Definitions 9.62 and 9.63 are *Σ–causal*: their support respects the cones of $\bar{\Sigma}$. No cohomological overlay or Ward/Slavnov step is altered.

Remark 9.84 (Mode normalization: Σ–inner product vs. triad metrics)**.** When realized quanta (graviton/Gaiaton/Kymons) are discussed in H^0, normalization by the Σ–inner products of Definition 9.33 is equivalent—up to a positive, blockwise bounded reweighting—to any triad-by-triad metric used earlier. Hence particle *criteria* (poles, residues, positivity, gauge-independence) remain identical; see Theorem 9.67.

Remark 9.85 (Quadrant diagnostics (unchanged logic, Σ–inputs))**.** In Chapter 8's PSR stage, compute law/state order parameters with the Σ–based scalars of Definition 9.47 instead of distances-only, and classify QI–QIV via Definition 9.49. The realization mapping then uses $\mathcal{U}_{\Sigma}$ (Definition 9.60) but lands in the same $H^0(Q_{\mathrm{BRST}})$ with the same viability/uniqueness tests; see Proposition 9.69.

Remark 9.86 (What definitively does not change)**.** None of the following are modified: ghost sector and the BRST charge; cohomology definitions; proof strategies for gauge–parameter independence and anomaly exclusions; the operator–bundle/gauge–group filtration recovering the SM sector (Ch. 11 excerpt).

Bottom line. Pre-physical chronogeometry only *feeds* Chapters 7–8: $\hat{\Sigma} \xrightarrow{\mathcal{U}_{\Sigma}} \bar{\Sigma}$. All downstream axioms, definitions, and proofs remain intact. In practice you gain cleaner causality (Σ–cones), unified normalization (Σ–inner products), and simpler, globally gauge–independent statements—while the BRST–physical sector and its machinery are unchanged.

9.8 Parasymmetric Type – Location – Onset on the Σ Pre-Physical Layer

Scope and substrate (with cross-references). This section establishes a fully pre-physical, expectation-free pipeline on the emergent Hilbert layer $\mathcal{H}_{\mathrm{em}}$ (Definition 1.34), beginning at

Algorithm 4 Minimal editorial patch for Chapters 7–8

1. Insert the bridge (9.35) at the start of Chapter 7 (just after introducing the physical arena).
2. Replace every "triadic background" dependency in the statements of cohomological EOM, linear response, visibilities, Ward/Slavnov identities $\rightsquigarrow$ the single symbol $\bar{\Sigma}$.
3. Leave all lemmas/proofs verbatim; optionally add a parenthetical "(Σ–Nielsen form)" referencing Remark 9.82.

the Primordial Quantum Singularity (PQS) (Definition 1.25) and running strictly below the Planckian regime. All constructions are formulated *exclusively* at operator level, using the operator chronogeometry $\hat{\Sigma}_{\mu\nu}$ (Definition 5.25), the unified inter-modal correlator $\hat{\mathbf{G}}_\Lambda$ (Definition 6.4) built from the ΦΦ (Gaiamiti) and ΘΘ (Tantrakyma) blocks and the fourfold dialectical basis $\{0, 1, p, s\}$ (Section 1.4.2 and Definition 4.9), band instrumentation $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell} = (P_x(\lambda_{\mathrm{UV}}), K^{(\ell)}_{\mathrm{IR}})$ (Definitions 7.26 and 9.10 and eq. (7.29)), and the BRST algebra/hygiene that guarantees gauge-parameter independence on the band (Lemma 9.54). Expectation-level objects (bars) never appear; all diagnostics and meters are defined directly on operators supported by the band.

Intent. From the common substrate $(\hat{\Sigma}_{\mu\nu}, \hat{\mathbf{G}}_\Lambda, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell})$ we *derive*—rather than posit—causal structure via the Σ–pre-physical cone $\mathcal{C}_x^{\hat{\Sigma},*}$ and its quadratic probe $\hat{Q}_x[v] = v^\mu v^\nu \hat{\Sigma}_{\mu\nu}(x)$ (Definition 9.22), enforce retardedness and microcausality on band-admissible kernels, and test *typed hyperbolicity*. Within each block ΦΦ (apeiron) and ΘΘ (kymon) (Definition 9.31 and eq. (9.12)) we then *discover* the permitted kinematic content by constructing typed, cone-compatible projectors $\Pi^{[\chi]}$ (spin/helicity/polarization, in the pre-physical sense) and recording a *permit ledger* $\mathcal{C}^{[\cdot]}_{\mathrm{admissible}}$. Using the Σ-probe and BRST-closed band functionals $\mathcal{I}, \mathcal{I}_\Psi$ (Definitions 9.45 and 9.47 and eqs. (9.21) and (9.22)), we define typed Σ–contrasts $\Delta^{\mathrm{pre},\Sigma;[\chi]}_{\mathrm{law/state}}$, normalise them to typed Σ–visibilities $\Lambda^{\Sigma;[\chi]}_{\mathrm{law/state}} \in [0, 1]$, and diagnose *Type* via the quadrant map

$$\mathrm{Quad}^{[\chi]} = \mathrm{Quad}\Big(\Lambda^{\Sigma;[\chi]}_{\mathrm{law}}, \Lambda^{\Sigma;[\chi]}_{\mathrm{state}}\Big) \in \{\mathrm{QI}, \mathrm{QII}, \mathrm{QIII}, \mathrm{QIV}\} \quad (\textit{Definitions } 7.685 \textit{and } 9.28).$$

Finally, we port the *Parasymmetric meters* of §8.6 to the band (restricted by $\Pi^{[\chi]}$) to deliver **T–L–O**: the quadrant *Type*, the *Location* (arena/sector/position/cone sheet; cf. Section 8.1), and the scale/time *Onset* $\big(r^\star[\chi], t^\star[\chi]\big)$. The entire stack is BRST/Nielsen-safe at operator level (Lemma 9.54), and regulator-robust under small retunings of $(\lambda_{\mathrm{UV}}, \ell)$ (Definition 9.10).

Meaning. Causality is extracted from $\hat{\Sigma}$ alone: the covector cone $\mathcal{C}_x^{\hat{\Sigma},*}$ is defined operationally by band functionals (cf. Definition 9.20), IR windowing can only *shrink* cones (Definition 9.10), and all kernels are retarded and cone-gated inside $J^+(\mathcal{C}^{\hat{\Sigma},*})$. Typed hyperbolicity is an *admission test*: a label χ is permitted only if the principal quadratic form

of $\hat{Q}_x[k]$ (Definition 9.22) is strictly positive on cone-timelike covectors after restriction by $\Pi^{[\chi]}$, yielding the permit ledgers $\mathcal{C}^{[\mathrm{apeiron}]}_{\mathrm{admissible}}$ and $\mathcal{C}^{[\mathrm{kymon}]}_{\mathrm{admissible}}$. With $\Pi^{[\chi]}$ fixed, the typed contrasts $\Delta^{\mathrm{pre},\Sigma;[\chi]}_{\mathrm{law/state}}$ (Definition 9.45) provide retarded, BRST-closed diagnostics of explicit-law versus state-level effects; normalisation gives $\Lambda^{\Sigma;[\chi]}_{\mathrm{law/state}}$ (Definition 9.47 and eq. (9.22)), and the quadrant predicate $\mathrm{Quad}^{[\chi]}$ classifies the mechanism (QI/QII/QIII/QIV) (Definitions 7.685 and 9.28), optionally with corr/vac/spec attribution (Definition 5.54). Parasymmetric meters at local anchors $a \in \{L, S\}$ and on the relational LS pair (Section 8.6),

$$\upsilon^{(a);\,[\chi]}_{\mathrm{Synthesis}}, \qquad \upsilon^{(LS);\,[\chi]}_{\mathrm{Synthesis}},$$

turn these into operational valuations whose scale/time dependence furnishes onset thresholds $\big(r^\star[\chi], t^\star[\chi]\big)$, while the location metadata records $(\mathfrak{P}/\mathfrak{P}_\partial,\ \Phi\Phi/\Theta\Theta,\ P_x,\ \text{cone sheet})$ (Section 8.1).

BRST hygiene at pre-physics. Band taggers and typed projectors commute with the nilpotent charge $\hat{Q}_{\mathrm{BRST}}$ and carry ghost number 0 (Lemma 9.54); the band functionals $\mathcal{I}, \mathcal{I}_\Psi$ annihilate BRST-exact insertions and project to ghost number 0 (Definition 9.45). Consequently, any gauge-parameter variation is cohomologically exact and evaluates to zero under $\mathcal{I}, \mathcal{I}_\Psi$: the pipeline is Nielsen-safe without invoking $\bar{\Sigma}$ or $H^0(\hat{Q}_{\mathrm{BRST}})$ (Lemma 9.54). Scope discipline is strict: only the $\Phi\Phi$ and $\Theta\Theta$ blocks receive kinematic labels in the sub-Planckian regime (Definition 9.31); any mapping of permitted sectors to gravitons/particles is deferred to the expectation-level domain and lies outside the present construction.

Relation to the established framework. The present section inherits the fourfold correlator architecture and operator chronogeometry from Chapters 4–6 (no bars; Definitions 4.9, 5.25 and 5.54), respects the unified block field $\hat{\mathbf{F}} = (\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}})^{\mathsf{T}}$ and its covariant transport from Chapter 7 (Definitions 6.4 and 6.184), adopts band instrumentation and BRST hygiene in the spirit of Chapter 8 (Definitions 7.26 and 9.10 and Lemma 9.54), and instantiates a sector-conditioned witness pair $(\Lambda^{\Sigma;[\chi]}_{\mathrm{law}}, \Lambda^{\Sigma;[\chi]}_{\mathrm{state}})$ that reproduces the canonical quadrant decision of Chapter 9—now *typed* and *cone-gated* (Section 8.1 and Definition 7.685). Parasymmetric meters provide the SynTh dashboard that operationalises T–L–O on the band (Sections 8.2.1 and 8.6).

Design guarantees and practical notes. Causal correctness follows from retarded support and cone gating by $\mathcal{C}^{\hat{\Sigma},*}_x$ with probe $\hat{Q}_x[v]$ (Definition 9.22); gauge cleanliness follows from BRST-closed instrumentation and BRST-invariant band functionals (Lemma 9.54 and Definition 9.45); well-posedness is enforced by typed hyperbolicity; stability and functoriality ensure small retunings of $(\lambda_{\mathrm{UV}}, \ell)$ alter estimates but not quadrant membership (within tolerances) (Definition 9.10). Practically, the probe set $\{v_k\}$ defining $\Lambda^{\Sigma;[\chi]}$ must be band-admissible and cone-separating (Definition 9.47); null-sheet degeneracies should be handled as limits from the timelike interior when computing $\Delta^{\mathrm{pre},\Sigma;[\chi]}_{\bullet}$ (Definition 9.45); and each χ should ship with a certificate triple (cone compatibility, typed hyperbolicity, band-stability window), with mode extraction aligned to the Σ-weighted principal-mode framework (Proposition 9.36, eq. (9.16), and Corollary 9.40).

We turn $\hat{\Sigma}$ (Definition 5.25), $\hat{\mathbf{G}}_\Lambda$ (Definition 6.4), and $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ (Definition 9.10) into a BRST-clean, cone-gated, expectation-free engine that (i) *discovers* which χ are kinematically allowed for apeiron/kymon and then (ii) *diagnoses* their *Type–Location–Onset* via typed Σ–visibilities and Parasymmetric meters (Definition 9.47 and Section 8.6)—strictly pre-physical, from the PQS (Definition 1.25) up to just below Planckian.

Hard Constraints (Formal Specification)

Definition 9.87 (Pre-physical arena and PQS baseline)**.** The *pre-physical arena* is the GNS/emergent Hilbert layer $\mathcal{H}_{\mathrm{em}} \equiv \mathcal{H}_{\overset{\circ}{\Psi}}$ obtained from the Quantum Holo-State $\omega^{\overset{\circ}{\Psi}}$ (Definition 1.34). The locus of analysis is any PQS site $\mathfrak{B}_x$ (Definition 1.25), and all constructions in this section hold *from* PQS up to *just below* the Planckian regime.

Definition 9.88 (Operator substrate and expectation map)**.** Let $\mathfrak{A}_{\mathrm{em}}$ denote the $*$–algebra on $\mathcal{H}_{\mathrm{em}}$ (Definition 9.87) generated by the operator fields and instrumentation

$$\hat{\mathbf{F}}(x) = \big(\hat{\mathbf{\Phi}}(x), \hat{\mathbf{\Theta}}(x)\big)^{\mathsf{T}}, \quad \hat{\Sigma}_{\mu\nu}(x), \quad P_x(\lambda_{\mathrm{UV}}),\ K^{(\ell)}_{\mathrm{IR}},\ \hat{Q}_{\mathrm{BRST}},$$

with graded-gauge transport as in Definition 9.7 and band-closedness as in Definition 9.96. For any (not necessarily used) state ω on $\mathfrak{A}_{\mathrm{em}}$, write $E_\omega : \mathfrak{A}_{\mathrm{em}} \to \mathbb{C},\ A \mapsto \omega(A)$ for the *expectation map*, and denote by $\bar{A}$ any symbol that is either $E_\omega(A)$ or a functional of such expectations (e.g. $\bar{\Sigma}_{\mu\nu}$).[1]

Axiom 9.89 (Expectation-free, operator-only regime (tight form))**.** All statements in this section are made *exclusively* in the operator substrate $\mathfrak{A}_{\mathrm{em}}$ of Definition 9.88; in particular:

(O1) **Bar prohibition.** No barred objects appear. Concretely, $\bar{\Sigma}_{\mu\nu}$, $\bar{g}_{\mu\nu}$, $\bar{t}_{\mu\nu}$ and any $\bar{\cdot}$ obtained from E_ω are *excluded*.

(O2) **Admissible operations.** Allowed constructions use only algebraic operations in $\mathfrak{A}_{\mathrm{em}}$ (sums, products, $*$, graded commutators), BRST action with ghost number 0 instruments (Definition 9.96), graded-gauge conjugations (Definition 9.7), and band restriction.

(O3) **Causality and probes are operator-defined.** Causal support and cones are defined via $\hat{\Sigma}$ and the probe $\hat{Q}_x[v] = v^\mu v^\nu \hat{\Sigma}_{\mu\nu}(x)$ (Definitions 9.95 and 9.104); kernels are retarded, cone-gated (Definition 9.106 and Proposition 9.107).

(O4) **Numerical readouts are band functionals, not expectations.** When numbers are needed, we use the band functionals $\mathcal{I}, \mathcal{I}_\Psi$ (Definitions 9.96 and 9.127), which annihilate BRST-exact insertions (Nielsen safety; Proposition 9.97). No E_ω is invoked.

(O5) **Typed restriction only by projectors in $\mathfrak{A}_{\mathrm{em}}$.** Kinematic typing uses $\Pi^{[\chi]} \in \mathfrak{A}_{\mathrm{em}}$ satisfying Definition 9.116 (block internality, BRST-closedness, cone-compatibility), with admission by typed hyperbolicity Definition 9.111.

Lemma 9.90 (Syntactic/semantic bar-elimination)**.** *Under Axiom 9.89, any occurrence of a barred symbol $\bar{A}$ renders a formula* ill-typed *in this section. Each construct that might otherwise refer to an expectation-level notion has an operator-level counterpart:*

[1] In some sources one also pushes $\mathfrak{A}_{\mathrm{em}}$ to an "expectation layer" algebra $\overline{\mathfrak{A}}$; $\bar{\cdot}$ covers that image as well.

- *chronogeometry $\hat{\Sigma}_{\mu\nu}$ and probe $\hat{Q}_x[v]$ (Definition 9.95);*
- *cones, retardedness, microcausality (Definitions 9.104 and 9.106 and Proposition 9.108);*
- *typed contrasts/visibilities and quadrants (Definitions 9.127, 9.133 and 9.137);*
- *Parasymmetric meters and T–L–O (Definitions 9.141, 9.143 and 9.147).*

Proposition 9.91 (Gauge-parameter independence in the operator-only regime)**.** *With Axiom 9.89 and BRST hygiene (Definition 9.96 and Proposition 9.97), every scalar diagnostic produced in §9.133–§9.147 is invariant under gauge-parameter changes. In particular, for any BRST-exact variation $\delta_\theta \mathcal{O} = [\hat{Q}_{\mathrm{BRST}}, X_\theta]$, $\partial_\theta \mathcal{I}(\mathcal{O}) = \partial_\theta \mathcal{I}_\Psi(\mathcal{O}) = 0$ (Proposition 9.97 and Lemmas 9.128 and 9.145).*

Proof of Proposition 9.91. Fix a one–parameter gauge choice $\theta \mapsto g_\theta$ and write δ_θ for differentiation at θ. By the operator-only axiom (Axiom 9.89) we work entirely in $\mathfrak{A}_{\mathrm{em}}$, and by BRST hygiene (Definition 9.96) the band taggers satisfy $[\,\hat{Q}_{\mathrm{BRST}}, P_x(\lambda_{\mathrm{UV}})\,] = [\,\hat{Q}_{\mathrm{BRST}}, K_{\mathrm{IR}}^{(\ell)}\,] = 0$ and the band functionals kill BRST-exact insertions ($\mathcal{I}([\,\hat{Q}_{\mathrm{BRST}}, X]) = \mathcal{I}_\Psi([\,\hat{Q}_{\mathrm{BRST}}, X]) = 0$, Proposition 9.97). Typed projectors are BRST-closed as well ($[\,\hat{Q}_{\mathrm{BRST}}, \Pi^{[\chi]}] = 0$, Definition 9.116).

Step 1: Pointwise typed contrasts are invariant. By definition (Definition 9.127), the basic inputs are

$$\hat{Q}_x[v] = v^\mu v^\nu \hat{\Sigma}_{\mu\nu}(x) \quad \text{and} \quad \hat{Q}_x[v]\Big|_{\Pi^{[\chi]}} = \Pi^{[\chi]} \hat{Q}_x[v] \Pi^{[\chi]} \quad (\textit{Definitions 9.95 and 9.109}).$$

Any gauge-parameter variation of these inputs is BRST-exact: $\delta_\theta \hat{Q}_x[v] = [\,\hat{Q}_{\mathrm{BRST}}, X_\theta\,]$ for some band-admissible X_θ (graded covariance, Definition 9.7), and the same holds after typed restriction because $[\,\hat{Q}_{\mathrm{BRST}}, \Pi^{[\chi]}] = 0$:

$$\delta_\theta\Big(\hat{Q}_x[v]\Big|_{\Pi^{[\chi]}}\Big) = \Pi^{[\chi]}[\,\hat{Q}_{\mathrm{BRST}}, X_\theta\,]\Pi^{[\chi]} = \Big[\,\hat{Q}_{\mathrm{BRST}},\, \Pi^{[\chi]} X_\theta \Pi^{[\chi]}\,\Big].$$

Applying the band functionals that define the law/state contrasts (Definition 9.127) and using Proposition 9.97 yields

$$\partial_\theta \mathcal{I}\Big[\Big(\hat{Q}_x[v] - \mathrm{Ad}_{g_\theta(x)}\, \hat{Q}_x[v]\Big)\Big|_{\Pi^{[\chi]}}\Big] = \partial_\theta \mathcal{I}_\Psi\Big[\hat{Q}_x[v]\Big|_{\Pi^{[\chi]}}\Big] - \partial_\theta \mathcal{I}_{g_\theta\cdot\Psi}\Big[\hat{Q}_x[v]\Big|_{\Pi^{[\chi]}}\Big] = 0.$$

Hence $\partial_\theta \Delta^{\mathrm{pre},\Sigma;[\chi]}_{\mathrm{law/state}}(g_\theta; x; v) = 0$. This recovers Lemma 9.128.

Step 2: Aggregated contrasts and visibilities are invariant. Aggregation is by weighted ℓ^2-combination over a fixed probe family (Definition 9.131), so by the chain rule and Step 1, $\partial_\theta\, \Delta^{\mathrm{pre},\Sigma;[\chi]}_{\bullet}(x; \mathcal{V}, w) = 0$. The typed visibilities are obtained by a fixed normalizer Vis applied to these aggregates (Definition 9.133); therefore $\partial_\theta\, \Lambda^{\Sigma;[\chi]}_{\bullet}(x) = 0$, as summarized in item (a) of Proposition 9.134.

Step 3: Quadrant predicate is invariant. The quadrant assignment depends only on the pair $(\Lambda^{\Sigma;[\chi]}_{\mathrm{law}}(x), \Lambda^{\Sigma;[\chi]}_{\mathrm{state}}(x))$ (Definition 9.137). Since this pair is gauge-invariant by Step 2, the quadrant $\mathrm{Quad}^{[\chi]}(x)$ is gauge-invariant as well.

Step 4: Parasymmetric meters and T–L–O are invariant. Band-reduced typed meter states and valuations are defined in Definitions 9.141, 9.143 and 9.144. Because the band restriction

and typed projector commute with $\hat{Q}_{\mathrm{BRST}}$ and the meter maps act at ghost number 0, gauge-parameter variations are BRST-exact and are annihilated by the band functionals that feed the meters; thus all valuations are invariant (Lemma 9.145(b)). The T–L–O triple draws *only* on these visibilities and meters (Definition 9.147); hence $\mathrm{TLO}^{[\chi]}(x;r)$ is gauge-parameter independent.

Combining Steps 1–4 shows that every scalar diagnostic in §9.133–§9.147 (typed visibilities, quadrants, meter valuations, and T–L–O outputs) is invariant under gauge changes. The explicit identities $\partial_\theta \mathcal{I}(\mathcal{O}) = \partial_\theta \mathcal{I}_\Psi(\mathcal{O}) = 0$ for any BRST-exact variation $\delta_\theta \mathcal{O} = [\,\hat{Q}_{\mathrm{BRST}}, X_\theta\,]$ are precisely Proposition 9.97, and the pointwise typed consequence is Lemma 9.128; meters obey Lemma 9.145(b). This completes the proof. □

Corollary 9.92 (Deliverables depend only on $\mathfrak{A}_{\mathrm{em}}$)**.** *All deliverables of §9.174 (permit ledgers, typed Σ–maps, Parasymmetric dashboards, quadrant certificates) are computed purely from* $\mathfrak{A}_{\mathrm{em}}$ *and the band tag, with no reference to any expectation map. Hence* $\mathrm{Del}[\bullet](x)$ *is meaningful and verifiable within the scope of Axiom 9.89.*

Remark 9.93 (Compatibility if an expectation layer is introduced later)**.** While barred objects are excluded here, if a later section introduces an expectation layer, then—by Nielsen safety (Proposition 9.97) and typed hyperbolicity (Definition 9.111)—the operator-level numbers coincide with those obtained by replacing $\mathcal{I}, \mathcal{I}_\Psi$ with compatible expectation-valued functionals on BRST cohomology classes. This remark is out of scope for the present axiom but clarifies forward compatibility.

Definition 9.94 (Unified generalized field and correlator (block form, fourfold modes))**.** Let the generalized field be

$$\hat{\mathbf{F}}(x) = \Big(\hat{\mathbf{\Phi}}(x),\, \hat{\mathbf{\Theta}}(x)\Big)^{\mathsf{T}},$$

and the unified inter-modal correlator (Definition 6.4) be written in block form over the geometry/matter split and the fourfold dialectical basis $\{0, 1, p, s\}$ (Section 1.4.2 and Definition 4.9):

$$\hat{\mathbf{G}}_\Lambda(x,x') = \begin{pmatrix} \hat{G}_{\Phi\Phi}(x,x') & \hat{G}_{\Phi\Theta}(x,x') \\ \hat{G}_{\Theta\Phi}(x,x') & \hat{G}_{\Theta\Theta}(x,x') \end{pmatrix}, \qquad \text{cf. eq. (9.12).} \tag{9.36}$$

Parallel transport and grading are those of Definition 9.7.

Definition 9.95 (Operator chronogeometry and Σ-probe)**.** The operator-valued unified chronogeometric tensor is

$$\hat{\Sigma}_{\mu\nu} \;=\; \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu} \qquad \text{(cf. Definition 5.25)}, \tag{9.37}$$

and its band-local quadratic probe is

$$\hat{Q}_x[v] \;:=\; v^\mu v^\nu\, \hat{\Sigma}_{\mu\nu}(x), \qquad \text{cf. Definition 9.22.} \tag{9.38}$$

Definition 9.96 (Band instrumentation and BRST-closed taggers)**.** The band tag is

$$\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell} = \Big(P_x(\lambda_{\mathrm{UV}}),\, K^{(\ell)}_{\mathrm{IR}}\Big),$$

where $P_x(\lambda_{\mathrm{UV}})$ is the UV projector (Definition 7.26 and eq. (7.29)) and $K_{\mathrm{IR}}^{(\ell)}$ is the IR kernel/window (Definition 9.10). Both are ghost number 0 and commute with the BRST charge:

$$[\,\hat{Q}_{\mathrm{BRST}},\, P_x(\lambda_{\mathrm{UV}})\,] = [\,\hat{Q}_{\mathrm{BRST}},\, K_{\mathrm{IR}}^{(\ell)}\,] = 0 \qquad \text{(cf. Lemma 9.54).} \tag{9.39}$$

Band functionals $\mathcal{I}, \mathcal{I}_\Psi$ are assumed admissible as in Definitions 9.45 and 9.47.

Proposition 9.97 (Nielsen safety on the band)**.** *If $\mathcal{I}, \mathcal{I}_\Psi$ satisfy the BRST-annihilation property of Definition 9.45, then for any band-supported operator $\mathcal{O}$ whose gauge-parameter variation is BRST-exact, $\delta_\theta \mathcal{O} = [\,\hat{Q}_{\mathrm{BRST}}, X_\theta]$, one has*

$$\partial_\theta\, \mathcal{I}(\mathcal{O}) = \mathcal{I}([\,\hat{Q}_{\mathrm{BRST}}, X_\theta]) = 0, \qquad \partial_\theta\, \mathcal{I}_\Psi(\mathcal{O}) = \mathcal{I}_\Psi([\,\hat{Q}_{\mathrm{BRST}}, X_\theta]) = 0.$$

Hence all band readouts used in this section are gauge-parameter independent without invoking expectation values (cf. Lemma 9.54).

Proof of Proposition 9.97. Let $\theta \mapsto \mathcal{O}(\theta)$ be a band-supported operator family with gauge-parameter variation $\delta_\theta \mathcal{O} = [\,\hat{Q}_{\mathrm{BRST}}, X_\theta]$, where X_θ is band-admissible. By construction of the band functionals (Definitions 9.45 and 9.96), $\mathcal{I}$ and $\mathcal{I}_\Psi$ are linear and continuous on the band algebra (ghost number 0 output). Hence differentiation passes under the functionals:

$$\partial_\theta\, \mathcal{I}(\mathcal{O}) = \mathcal{I}(\partial_\theta \mathcal{O}), \qquad \partial_\theta\, \mathcal{I}_\Psi(\mathcal{O}) = \mathcal{I}_\Psi(\partial_\theta \mathcal{O}). \tag{9.40}$$

Inserting $\partial_\theta \mathcal{O} = [\,\hat{Q}_{\mathrm{BRST}}, X_\theta]$ and using the *BRST-annihilation* property of the band functionals

$$\mathcal{I}([\,\hat{Q}_{\mathrm{BRST}}, Y]) = \mathcal{I}_\Psi([\,\hat{Q}_{\mathrm{BRST}}, Y]) = 0 \quad \text{for all band-admissible } Y \qquad \text{(Definition 9.45)},$$

we obtain

$$\partial_\theta\, \mathcal{I}(\mathcal{O}) = \mathcal{I}([\,\hat{Q}_{\mathrm{BRST}}, X_\theta]) = 0, \qquad \partial_\theta\, \mathcal{I}_\Psi(\mathcal{O}) = \mathcal{I}_\Psi([\,\hat{Q}_{\mathrm{BRST}}, X_\theta]) = 0,$$

which is the desired Nielsen (gauge-parameter) independence on the band.

Typed/restricted observables. If $\mathcal{O}$ enters through a typed restriction $\mathcal{O}\big|_{\Pi^{[\chi]}} = \Pi^{[\chi]} \mathcal{O}\, \Pi^{[\chi]}$ (as in Definitions 9.109 and 9.127), then by BRST-closedness of the projector $[\,\hat{Q}_{\mathrm{BRST}}, \Pi^{[\chi]}] = 0$ (Definition 9.116) we have

$$\delta_\theta\Big(\mathcal{O}\big|_{\Pi^{[\chi]}}\Big) = \Pi^{[\chi]}[\,\hat{Q}_{\mathrm{BRST}}, X_\theta]\Pi^{[\chi]} = \Big[\,\hat{Q}_{\mathrm{BRST}},\, \Pi^{[\chi]} X_\theta \Pi^{[\chi]}\,\Big],$$

so the same annihilation argument applies and the conclusion holds verbatim for all typed band readouts (contrasts/visibilities Definition 9.133, meter valuations Definitions 9.143 and 9.144, and T–L–O outputs Definition 9.147).

This establishes the proposition; see also its use in Lemmas 9.128 and 9.145 and Proposition 9.91. □

Definition 9.98 (Σ–pre-physical covector cone)**.** Define the covector cone at x by

$$\mathcal{C}_x^{\hat{\Sigma},*} := \Big\{ k \in T_x^* \setminus \{0\} : \sup_{\Psi \in \mathcal{S}_{\mathrm{band}}} \mathcal{I}_\Psi\Big[k^\mu k^\nu \hat{\Sigma}_{\mu\nu}(x)\Big] \le 0 \Big\}, \tag{9.41}$$

with null boundary at 0, timelike interior determined by $\inf_\Psi \mathcal{I}_\Psi(\hat{Q}_x[k]) > 0$ (cf. Definitions 9.20 and 9.22). The IR window $K_{\mathrm{IR}}^{(\ell)}$ may *shrink* but never enlarge $\mathcal{C}_x^{\hat{\Sigma},*}$ (Definition 9.10).

Lemma 9.99 (Causal gate and microcausality from $\hat{\Sigma}$)**.** *Let the Σ–pre-physical cone $\mathcal{C}_x^{\hat{\Sigma},*}$ be as in Definition 9.104 and let kernels be retarded and cone-gated as in Definition 9.106. If f, g have supports spacelike separated with respect to $\mathcal{C}^{\hat{\Sigma},*}$, then*

$$[\,\hat{\mathbf{\Phi}}(f), \hat{\mathbf{\Phi}}(g)\,] = [\,\hat{\mathbf{\Theta}}(f), \hat{\mathbf{\Theta}}(g)\,] = 0,$$

and every response kernel used in this section is supported in $J^+\big(\mathcal{C}^{\hat{\Sigma},*}\big)$ (i.e. retarded and cone-gated).

Proof. By Definition 9.106, any kernel $\mathcal{K}$ is replaced by its retarded, cone-gated version $\mathcal{K}^{\mathrm{ret}}(x, x') = \mathbf{1}_{J^+(\mathcal{C}^{\hat{\Sigma},*})}(x, x')\,\mathcal{K}(x, x')$. Proposition 9.107 asserts that this gating preserves band-admissibility and graded covariance, and enforces retarded support inside $J^+(\mathcal{C}^{\hat{\Sigma},*})$. Hence all response kernels constructed downstream are retarded and cone-gated.

For the commutators, let f, g be test functions with supports spacelike separated relative to $\mathcal{C}^{\hat{\Sigma},*}$. The microcausality statement follows directly from Proposition 9.108, which states that, with respect to the Σ–cone, the smeared field commutators vanish outside the causal domain:

$$[\,\hat{\mathbf{\Phi}}(f), \hat{\mathbf{\Phi}}(g)\,] = [\,\hat{\mathbf{\Theta}}(f), \hat{\mathbf{\Theta}}(g)\,] = 0 \quad \text{whenever} \quad \mathrm{supp}(f) \text{ and } \mathrm{supp}(g) \text{ are spacelike.}$$

Equivalently, in distributional form the commutator kernels have support contained in $J^+\big(\mathcal{C}^{\hat{\Sigma},*}\big) \cup J^-\big(\mathcal{C}^{\hat{\Sigma},*}\big)$; for spacelike-separated supports the double integral against $f \otimes g$ vanishes, yielding the identities above.

Therefore, $\hat{\Sigma}$ via the cone $\mathcal{C}^{\hat{\Sigma},*}$ supplies both the causal gate (retarded, cone-gated kernels by Definition 9.106 and Proposition 9.107) and microcausality (vanishing of spacelike commutators by Proposition 9.108). □

Definition 9.100 (Typed kinematic labels (pre-physical))**.** Typed labels χ (spins/helicities/polarizations) are *admissible* only within the diagonal blocks $\Phi\Phi$ (apeiron/Gaiamiti) and $\Theta\Theta$ (kymon/Tantrakyma) of $\hat{\mathbf{G}}_\Lambda$ (Definition 6.4), and are implemented by graded, cone-compatible projectors $\Pi^{[\chi]}$ constructed inside each block (cf. Definitions 9.7 and 9.31 and eq. (9.12)). No graviton/particle language is permitted below Planckian.

Definition 9.101 (Typed hyperbolicity (admission test))**.** A label χ is *typed-hyperbolic* at x if for every cone-timelike covector $k \in \mathcal{C}_x^{\hat{\Sigma},*}$,

$$\inf_{\Psi \in \mathcal{S}_{\mathrm{band}}} \mathcal{I}_\Psi\Big[\, \hat{Q}_x[k] \,\Big|_{\Pi^{[\chi]}}\Big] \;>\; 0. \tag{9.42}$$

Proposition 9.102 (Permit ledger)**.** *The set of* admissible *typed labels is*

$$\mathcal{C}^{[\mathrm{apeiron}]}_{\mathrm{admissible}} \subset \mathrm{End}(\Phi\Phi), \qquad \mathcal{C}^{[\mathrm{kymon}]}_{\mathrm{admissible}} \subset \mathrm{End}(\Theta\Theta),$$

where $\chi \in \mathcal{C}^{[\cdot]}_{\mathrm{admissible}}$ iff $\Pi^{[\chi]}$ is band-finite, graded-covariant (Definition 9.7), cone-compatible (Definition 9.98), and satisfies typed hyperbolicity (9.42) with stability under small retunings of $(\lambda_{\mathrm{UV}}, \ell)$ (Definition 9.10).

Proof of Proposition 9.102. We prove necessity and sufficiency.

(⇒) Necessity. Assume $\chi \in \mathcal{C}^{[\bullet]}_{\text{admissible}}$ with $\bullet \in \{\Phi\Phi, \Theta\Theta\}$. By the very definition of the permit ledger (Definition 9.122), there exists a typed projector $\Pi^{[\chi]}$ admitted in block $\bullet$ that satisfies the axioms of Definition 9.116 together with the typed hyperbolicity condition (Definition 9.111, eq. (9.53)). Hence:

- *Block internality:* $\Pi^{[\chi]} \in \mathrm{End}(\bullet)$ by Definition 9.116(P2), so χ is recorded in the corresponding set $\mathcal{C}^{[\bullet]}_{\text{admissible}}$.
- *Band-finiteness:* $\Pi^{[\chi]}$ commutes with the band taggers and is supported on the band by Definition 9.116(P3) (cf. band instrumentation Definition 9.96).
- *Graded covariance:* under graded transport g, $\Pi^{[\chi]} \mapsto g\,\Pi^{[\chi]} g^{\dagger}$ by Definition 9.116(P5) (see also Definition 9.7).
- *Cone-compatibility:* positivity of the Σ–probe on timelike covectors within the Σ–cone holds after restriction by $\Pi^{[\chi]}$ by Definition 9.116(P6), using the cone of Definition 9.104.
- *Typed hyperbolicity:* the principal form positivity on $\mathcal{C}^{\hat{\Sigma},*}$ after restriction is precisely (9.53) (Definition 9.111).
- *Stability under retunings:* continuity in $(\lambda_{\mathrm{UV}}, \ell)$ holds by Definition 9.116(P7), with the parameters defined for the band in Definition 9.96.

Therefore every admitted χ satisfies all properties listed in the statement.

(⇐) Sufficiency. Conversely, let $\Pi^{[\chi]}$ be a projector in block $\bullet$ that is band-finite, graded-covariant (Definition 9.7), cone-compatible with the Σ–cone (Definition 9.104), satisfies typed hyperbolicity (9.53), and is stable under small retunings of $(\lambda_{\mathrm{UV}}, \ell)$ (Definition 9.96). Then $\Pi^{[\chi]}$ fulfills items (P1)–(P7) of Definition 9.116, so by Definition 9.122 the label χ is *admitted* and belongs to $\mathcal{C}^{[\bullet]}_{\text{admissible}}$. Moreover, the cone/hyperbolicity/stability data constitute the certificate attached to χ (Corollary 9.120).

Existence (constructive aside). When needed, such projectors can be constructed spectrally from Σ–weighted mode operators, with Riesz projectors onto isolated spectral bands (Proposition 9.119); this automatically yields band-finiteness, graded covariance, and the stability radius used above.

Thus, the admissible sets are exactly those stated, completing the proof. □

Corollary 9.103 (Scope discipline)**.** *Typed labels χ defined in Definition 9.100 are the only kinematic labels used below Planckian. Any map of χ-sectors to graviton/particle content is deferred to the expectation-level framework and lies outside the present operator-only regime (cf. Definition 5.25 vs. Definition 5.10).*

Remarks on consistency.

- The ingredients (9.36), (9.37), (9.38) re-state, with explicit band locality, the earlier definitions Definitions 5.25, 6.4 and 9.22. All subsequent typed diagnostics (contrasts/visibilities) that will be introduced in §1–§3 refer back to Definitions 9.45 and 9.47 and eq. (9.22) and will be restricted by $\Pi^{[\chi]}$ in the sense of Definition 9.100.

- BRST hygiene (9.39) and Proposition 9.97 implement band-level Nielsen safety exactly as in Lemma 9.54, but without any use of expectation values.
- The cone functoriality "IR shrinks, never enlarges" invoked in Definition 9.98 follows from the monotonicity of $K_{\mathrm{IR}}^{(\ell)}$ in Definition 9.10.

9.9 Σ–Pre-physical Cones and Causality (Operator Level)

Definition 9.104 (Σ–pre-physical covector cone at x)**.** Let $\hat{\Sigma}_{\mu\nu}$ and the Σ–probe $\hat{Q}_x[v] = v^\mu v^\nu \hat{\Sigma}_{\mu\nu}(x)$ be as in Definitions 5.25 and 9.22 (see also the band-local restatement Definition 9.95). The Σ–*pre-physical covector cone* at x is

$$\mathcal{C}_x^{\hat{\Sigma},*} := \Big\{ k \in T_x^*\backslash\{0\} \ : \ \sup_{\Psi\in\mathcal{S}_{\mathrm{band}}} \mathcal{I}_\Psi\Big[k^\mu k^\nu \hat{\Sigma}_{\mu\nu}(x)\Big] \le 0 \Big\}, \tag{9.43}$$

with *null* boundary given by equality to 0 and *timelike interior* characterized by $\inf_{\Psi\in\mathcal{S}_{\mathrm{band}}} \mathcal{I}_\Psi\big[\hat{Q}_x[k]\big] > 0$. This definition is equivalent to the restated cone in Definition 9.98, but we fix here the label Definition 9.104 for subsequent use.

Lemma 9.105 (IR windowing shrinks, never enlarges, the cone)**.** *Let* $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell} = (P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}^{(\ell)})$ *be as in Definitions 7.26 and 9.10. If* $\ell' \le \ell$ *(stronger IR windowing), then*

$$\mathcal{C}_x^{\hat{\Sigma},*}\big[\ell'\big] \ \subseteq \ \mathcal{C}_x^{\hat{\Sigma},*}\big[\ell\big].$$

Proof of Lemma 9.105. Recall the Σ–pre-physical covector cone at x (Definition 9.104):

$$\mathcal{C}_x^{\hat{\Sigma},*}[\ell] \ := \ \Big\{ k \in T_x^*\backslash\{0\} \ : \ \sup_{\Psi\in\mathcal{S}_{\mathrm{band}}(\ell)} \mathcal{I}_\Psi\Big[k^\mu k^\nu \, \hat{\Sigma}_{\mu\nu}(x)\Big] \ \le \ 0 \Big\}, \tag{9.44}$$

where $\mathcal{S}_{\mathrm{band}}(\ell)$ denotes band-admissible states/functionals under $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ (Definition 9.10). Equivalently, writing the IR-gating map $\Gamma_\ell(A) := K_{\mathrm{IR}}^{(\ell)} A K_{\mathrm{IR}}^{(\ell)}$, (9.44) is

$$\mathcal{C}_x^{\hat{\Sigma},*}[\ell] = \Big\{ k \neq 0 : \ \sup_{\Psi\in\mathcal{S}_{\mathrm{band}}} \mathcal{I}_\Psi\Big[k^\mu k^\nu \, \Gamma_\ell(\hat{\Sigma}_{\mu\nu})(x)\Big] \ \le \ 0 \Big\},$$

since the band functionals act after the IR gate (Definition 9.10).

Step 1 (IR monotonicity of the gate). By Definition 7.26, the family $\{K_{\mathrm{IR}}^{(\ell)}\}$ is positive, cone-preserving, and *monotone in the sense that for* $\ell' \le \ell$ *one has*

$$\Gamma_{\ell'}(A) \ - \ \Gamma_\ell(A) \ \succeq \ 0 \quad \text{for every operator } A \text{ supported on the future cone sheet,} \tag{9.45}$$

i.e. a stronger IR window cannot add past-directed (or non-causal) support and can only *increase* the future-directed part of A (cf. the cone-gating of Definition 9.106 and causal support of Proposition 9.107).

Step 2 (Monotonicity of band readouts). Fix k and x. For future-sheet contributions of $k^\mu k^\nu \hat{\Sigma}_{\mu\nu}(x)$, Lemma 9.129 gives positivity of the induced quadratic form under band

functionals. Combining with (9.45) and positivity of $\mathcal{I}_\Psi$ on the band (Definition 9.10), we get, for every Ψ,

$$\mathcal{I}_\Psi\Big[k^\mu k^\nu\,\Gamma_{\ell'}(\hat{\Sigma}_{\mu\nu})(x)\Big] \;\geq\; \mathcal{I}_\Psi\Big[k^\mu k^\nu\,\Gamma_{\ell}(\hat{\Sigma}_{\mu\nu})(x)\Big] \qquad (\ell' \leq \ell). \tag{9.46}$$

Taking suprema over Ψ preserves the inequality:

$$\sup_\Psi\; \mathcal{I}_\Psi\Big[k^\mu k^\nu\,\Gamma_{\ell'}(\hat{\Sigma}_{\mu\nu})(x)\Big] \;\geq\; \sup_\Psi\; \mathcal{I}_\Psi\Big[k^\mu k^\nu\,\Gamma_{\ell}(\hat{\Sigma}_{\mu\nu})(x)\Big]. \tag{9.47}$$

Step 3 (Inclusion of cones). Suppose $k \in \mathcal{C}_x^{\hat{\Sigma},*}[\ell']$. By (9.44) (with gating) and (9.47),

$$0 \;\geq\; \sup_\Psi\; \mathcal{I}_\Psi\Big[k^\mu k^\nu\,\Gamma_{\ell'}(\hat{\Sigma}_{\mu\nu})(x)\Big] \;\geq\; \sup_\Psi\; \mathcal{I}_\Psi\Big[k^\mu k^\nu\,\Gamma_{\ell}(\hat{\Sigma}_{\mu\nu})(x)\Big],$$

hence $k \in \mathcal{C}_x^{\hat{\Sigma},*}[\ell]$. Therefore $\mathcal{C}_x^{\hat{\Sigma},*}[\ell'] \subseteq \mathcal{C}_x^{\hat{\Sigma},*}[\ell]$.

Conclusion. A stronger IR window (smaller ℓ) makes the Σ–cone test (9.44) *harder* to satisfy (the supremum moves upward by (9.47)), so membership can only be lost: the cone *shrinks*, never enlarges. □

Definition 9.106 (Future cone-gate and retarded domain)**.** Let $J^+\big(\mathcal{C}^{\hat{\Sigma},*}\big)$ denote the *future cone domain* determined by $\mathcal{C}_x^{\hat{\Sigma},*}$, i.e. pairs (x,x') such that there exists a $\hat{\Sigma}$–causal curve from x' to x whose cotangent legs lie in $\mathcal{C}^{\hat{\Sigma},*}$. For any (graded-covariant) kernel $\mathcal{K}(x,x')$ we define its *retarded cone-gated* version by

$$(\mathcal{K}^{\mathrm{ret}})(x,x') \;:=\; \mathbf{1}_{J^+(\mathcal{C}^{\hat{\Sigma},*})}(x,x')\,\mathcal{K}(x,x'),$$

where $\mathbf{1}_{J^+}$ is the characteristic function. (Graded covariance follows from Definition 9.7.)

Proposition 9.107 (Retardedness of band kernels)**.** *Let $\mathcal{K}$ be any band-admissible kernel (e.g. those used in defining the contrasts/visibilities of Definitions 9.45 and 9.47). Then $\mathcal{K}^{\mathrm{ret}}$ of Definition 9.106 is retarded and supported inside $J^+\big(\mathcal{C}^{\hat{\Sigma},*}\big)$; moreover, the operation $\mathcal{K} \mapsto \mathcal{K}^{\mathrm{ret}}$ is compatible with band instrumentation $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ (Definition 9.10) and with BRST (ghost number 0, cf. Lemma 9.54).*

Proof of Proposition 9.107. Write any band-admissible kernel $\mathcal{K}$ as an integral operator on test functions,

$$(\mathcal{K}f)(x) \;=\; \int \mathcal{K}(x,x')\,f(x')\,\mathrm{d}x',$$

with $\mathcal{K}$ belonging to the band-closed operator class of Definition 9.10. By the cone gate definition (Definition 9.106) the retarded version is

$$\mathcal{K}^{\mathrm{ret}}(x,x') \;:=\; \mathbf{1}_{J^+(\mathcal{C}^{\hat{\Sigma},*})}(x,x')\;\mathcal{K}(x,x'), \tag{9.48}$$

where $\mathbf{1}_{J^+(\mathcal{C}^{\hat{\Sigma},*})}$ is the (idempotent) indicator of the Σ–causal future determined by the cone $\mathcal{C}^{\hat{\Sigma},*}$ (Definition 9.104).

(i) Retardedness and cone support. By (9.48), $\operatorname{supp}\mathcal{K}^{\mathrm{ret}} \subseteq J^+(\mathcal{C}^{\hat{\Sigma},*})$, so $\mathcal{K}^{\mathrm{ret}}$ is retarded and cone-gated by construction (restating Definition 9.106). No support can appear outside $J^+(\mathcal{C}^{\hat{\Sigma},*})$ because the gate $\mathbf{1}_{J^+}$ vanishes there.

(ii) Compatibility with band instrumentation $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$. Let the UV projector $P_x(\lambda_{\mathrm{UV}})$ act by left/right multiplication on the (x,x') slots (cf. Definition 9.10). Then

$$\Big(P_x(\lambda_{\mathrm{UV}})\,\mathcal{K}\Big)^{\mathrm{ret}}(x,x') = \mathbf{1}_{J^+}(x,x')\,P_x(\lambda_{\mathrm{UV}})\,\mathcal{K}(x,x') = P_x(\lambda_{\mathrm{UV}})\,\mathcal{K}^{\mathrm{ret}}(x,x'),$$

and similarly on the right slot with $P_{x'}(\lambda_{\mathrm{UV}})$. Thus the gate commutes with UV localization at the level of kernels, and band-admissibility is preserved.

For the IR filter $K_{\mathrm{IR}}^{(\ell)}$, let Γ_ℓ denote the IR-gating superoperator (Definition 9.10). Since Γ_ℓ acts at ghost number 0 and preserves the cone (Definition 9.106), we have

$$\Big(\Gamma_\ell\mathcal{K}\Big)^{\mathrm{ret}} = \mathbf{1}_{J^+}\,\Gamma_\ell(\mathcal{K}) = \Gamma_\ell\Big(\mathbf{1}_{J^+}\,\mathcal{K}\Big) = \Gamma_\ell\Big(\mathcal{K}^{\mathrm{ret}}\Big), \tag{9.49}$$

i.e. retarded gating and IR filtering commute within the admissible class. Hence the map $\mathcal{K}\mapsto\mathcal{K}^{\mathrm{ret}}$ is compatible with $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ and preserves band-admissibility.

(iii) BRST compatibility (ghost number 0). The indicator $\mathbf{1}_{J^+(\mathcal{C}^{\hat{\Sigma},*})}$ is a c–number gate determined by the Σ–cone (it carries no ghost number). Therefore, for the BRST derivation one has

$$[\,\hat{Q}_{\mathrm{BRST}},\mathcal{K}^{\mathrm{ret}}\,] = [\,\hat{Q}_{\mathrm{BRST}},\ \mathbf{1}_{J^+}\,\mathcal{K}\,] = \mathbf{1}_{J^+}\,[\,\hat{Q}_{\mathrm{BRST}},\mathcal{K}\,] = \Big([\,\hat{Q}_{\mathrm{BRST}},\mathcal{K}\,]\Big)^{\mathrm{ret}}.$$

In particular, if $\mathcal{K}$ is BRST-closed at ghost number 0 (the case for our band kernels; see Lemma 9.54 and Definition 9.96), then so is $\mathcal{K}^{\mathrm{ret}}$. This establishes BRST compatibility.

Combining (i)–(iii) proves that the retarded gate (9.48) yields kernels supported in $J^+(\mathcal{C}^{\hat{\Sigma},*})$, preserves band-admissibility and UV/IR instrumentation (Definition 9.10), and is compatible with BRST at ghost number 0. Hence the claim. □

Proposition 9.108 (Microcausality from cone-gating)**.** *Let f,g be test functions with supports spacelike separated with respect to $\mathcal{C}^{\hat{\Sigma},*}$. Then*

$$\Big[\,\hat{\mathbf{\Phi}}(f),\hat{\mathbf{\Phi}}(g)\,\Big] \;=\; \Big[\,\hat{\mathbf{\Theta}}(f),\hat{\mathbf{\Theta}}(g)\,\Big] \;=\; 0, \tag{9.50}$$

for the generalized field $\hat{\mathbf{F}} = (\hat{\mathbf{\Phi}},\hat{\mathbf{\Theta}})^{\mathsf{T}}$ (Definition 9.94). In particular, any cone-gated retarded kernel $\mathcal{K}^{\mathrm{ret}}$ from Definition 9.106 yields vanishing commutators outside $J^+\Big(\mathcal{C}^{\hat{\Sigma},*}\Big)$.

Proof of Proposition 9.108. Let $\mathcal{C}_x^{\hat{\Sigma},*}$ be the Σ–pre-physical cone (Definition 9.104) and let the cone-gated retarded indicator be $\mathbf{1}_{J^+(\mathcal{C}^{\hat{\Sigma},*})}$ as in Definition 9.106. By Definition 9.94, the unified field $\hat{\mathbf{F}} = (\hat{\mathbf{\Phi}},\hat{\mathbf{\Theta}})^{\mathsf{T}}$ is an operator-valued distribution, and for any test functions f,g we define the smeared operators $\hat{\mathbf{\Phi}}(f) = \int \hat{\mathbf{\Phi}}(x)\,f(x)\,\mathrm{d}x$ and similarly for $\hat{\mathbf{\Theta}}(g)$.

Step 1: Commutator as a distribution with causal support. Consider the bi-distribution (Pauli–Jordan–like kernel) for each block

$$\Delta_{\Phi\Phi}(x,x') \;:=\; \Big[\hat{\mathbf{\Phi}}(x),\hat{\mathbf{\Phi}}(x')\Big], \qquad \Delta_{\Theta\Theta}(x,x') \;:=\; \Big[\hat{\mathbf{\Theta}}(x),\hat{\mathbf{\Theta}}(x')\Big].$$

Any band-admissible kernel used to construct $\hat{\mathbf{F}}$ is replaced by its retarded, cone-gated form $\mathcal{K}^{\mathrm{ret}}(x,x') = \mathbf{1}_{J^+}(x,x')\,\mathcal{K}(x,x')$ (Definition 9.106), and this operation is compatible with

the band instrumentation (Definition 9.10) and preserves retarded (future-directed) support (Proposition 9.107). By antisymmetry of the commutator and the fact that advanced parts arise by exchanging the slots, it follows that

$$\operatorname{supp}\Delta_{\Phi\Phi} \;\subseteq\; J^{+}\!\big(\mathcal{C}^{\hat{\Sigma},*}\big) \;\cup\; J^{-}\!\big(\mathcal{C}^{\hat{\Sigma},*}\big), \qquad \operatorname{supp}\Delta_{\Theta\Theta} \;\subseteq\; J^{+}\!\big(\mathcal{C}^{\hat{\Sigma},*}\big) \;\cup\; J^{-}\!\big(\mathcal{C}^{\hat{\Sigma},*}\big), \tag{9.51}$$

i.e. the commutator kernels have no support at spacelike-separated pairs with respect to the Σ–cone.

Step 2: Vanishing for spacelike-separated test functions. Let f, g be test functions with $\operatorname{supp}(f)$ and $\operatorname{supp}(g)$ spacelike separated relative to $\mathcal{C}^{\hat{\Sigma},*}$. Smearing the commutators yields

$$\big[\hat{\mathbf{\Phi}}(f), \hat{\mathbf{\Phi}}(g)\big] = \iint \Delta_{\Phi\Phi}(x,x')\, f(x)\, g(x')\, \mathrm{d}x\, \mathrm{d}x',$$
$$\big[\hat{\mathbf{\Theta}}(f), \hat{\mathbf{\Theta}}(g)\big] = \iint \Delta_{\Theta\Theta}(x,x')\, f(x)\, g(x')\, \mathrm{d}x\, \mathrm{d}x'.$$

Because $\operatorname{supp}\Delta_{\bullet\bullet}$ is contained in the causal sets (9.51) while $\operatorname{supp}(f) \times \operatorname{supp}(g)$ is disjoint from those sets (spacelike separation), each double integral vanishes by the standard support property of distributions. Hence

$$\big[\,\hat{\mathbf{\Phi}}(f), \hat{\mathbf{\Phi}}(g)\,\big] = \big[\,\hat{\mathbf{\Theta}}(f), \hat{\mathbf{\Theta}}(g)\,\big] = 0,$$

which is (9.50).

Step 3: Cone-gated retarded kernels produce vanishing commutators outside J^+. Conversely, let $\mathcal{K}^{\mathrm{ret}}$ be any cone-gated retarded kernel (Definition 9.106). By Proposition 9.107, $\operatorname{supp}\mathcal{K}^{\mathrm{ret}} \subseteq J^{+}(\mathcal{C}^{\hat{\Sigma},*})$, so any commutator built from such kernels has bi-support contained in $J^{+} \cup J^{-}$ as in (9.51). Therefore commutators vanish whenever the argument supports fall outside $J^{+}(\mathcal{C}^{\hat{\Sigma},*})$.

Combining Steps 1–3 establishes microcausality with respect to the Σ–cone and the stated property for cone-gated retarded kernels. □

Definition 9.109 (Typed restriction of the Σ–probe)**.** For a typed projector $\Pi^{[\chi]}$ (constructed within the diagonal blocks as in Definitions 9.100 and 9.116), define the *typed* Σ–probe by

$$\hat{Q}^{[\chi]}_x[k] \;:=\; \Pi^{[\chi]}\, \hat{Q}_x[k]\, \Pi^{[\chi]} \;=\; \Pi^{[\chi]}\, k^{\mu} k^{\nu} \hat{\Sigma}_{\mu\nu}(x)\, \Pi^{[\chi]}. \tag{9.52}$$

Lemma 9.110 (Typed principal form is Hermitian and gauge-covariant)**.** *For any timelike $k \in \mathcal{C}^{\hat{\Sigma},*}_x$ the operator $\hat{Q}^{[\chi]}_x[k]$ of (9.52) is Hermitian and transforms covariantly under local graded-gauge changes (by Definitions 5.25 and 9.7). Hence its band functionals $\mathcal{I}_{\Psi}\big[\hat{Q}^{[\chi]}_x[k]\big]$ are real and frame-independent on the band (Definition 9.96).*

Proof of Lemma 9.110. Recall the Σ–probe and its typed restriction:

$$\hat{Q}_x[v] := v^{\mu} v^{\nu}\, \hat{\Sigma}_{\mu\nu}(x) \quad \text{(Definition 9.95)},$$
$$\hat{Q}^{[\chi]}_x[v] := \Pi^{[\chi]}\, \hat{Q}_x[v]\, \Pi^{[\chi]} \quad \text{(cf. (9.52), Definitions 9.109 and 9.116)}.$$

Fix $k \in \mathcal{C}_x^{\hat{\Sigma},*}$ timelike (Definition 9.104).

(a) Hermiticity. By the operator-unified chronogeometry (Definition 5.25), $\hat{\Sigma}_{\mu\nu}(x)$ is Hermitian (self-adjoint) for each index pair, and for real covectors k^μ one has

$$\hat{Q}_x[k]^\dagger = \big(k^\mu k^\nu \hat{\Sigma}_{\mu\nu}(x)\big)^\dagger = k^\mu k^\nu \hat{\Sigma}_{\mu\nu}(x) = \hat{Q}_x[k].$$

Typed projectors $\Pi^{[\chi]}$ are Hermitian idempotents in the block algebra (Definition 9.116), hence

$$\big(\hat{Q}_x^{[\chi]}[k]\big)^\dagger = \Pi^{[\chi]}\,\hat{Q}_x[k]\,\Pi^{[\chi]} = \hat{Q}_x^{[\chi]}[k],$$

so $\hat{Q}_x^{[\chi]}[k]$ is Hermitian.

(b) Graded-gauge covariance. Under a local graded-gauge transport g (Definition 9.7), $\hat{\Sigma}_{\mu\nu} \mapsto \mathrm{Ad}_{g(x)}\,\hat{\Sigma}_{\mu\nu}$, whence $\hat{Q}_x[k] \mapsto \mathrm{Ad}_{g(x)}\,\hat{Q}_x[k]$. Typed projectors transform covariantly as $\Pi^{[\chi]} \mapsto \mathrm{Ad}_{g(x)}\,\Pi^{[\chi]}$ (Definition 9.116). Therefore

$$\hat{Q}_x^{[\chi]}[k] \;\mapsto\; \mathrm{Ad}_{g(x)}\big(\Pi^{[\chi]}\hat{Q}_x[k]\Pi^{[\chi]}\big) = \big(\mathrm{Ad}_{g(x)}\,\Pi^{[\chi]}\big)\big(\mathrm{Ad}_{g(x)}\,\hat{Q}_x[k]\big)\big(\mathrm{Ad}_{g(x)}\,\Pi^{[\chi]}\big),$$

i.e. $\hat{Q}_x^{[\chi]}[k]$ transforms by conjugation (covariantly) under graded-gauge changes.

(c) Reality and frame-independence of band evaluations. Band functionals $\mathcal{I}, \mathcal{I}_\Psi$ act at ghost number 0 and satisfy $\mathcal{I}(A^\dagger) = \overline{\mathcal{I}(A)}$, $\mathcal{I}_\Psi(A^\dagger) = \overline{\mathcal{I}_\Psi(A)}$, with real range on Hermitian inputs (Definition 9.96). Since $\hat{Q}_x^{[\chi]}[k]$ is Hermitian by (a), $\mathcal{I}_\Psi\big[\hat{Q}_x^{[\chi]}[k]\big] \in \mathbb{R}$. For frame-independence, use graded covariance: for any local g,

$$\mathcal{I}_{g\cdot\Psi}\big[\hat{Q}_x^{[\chi]}[k]\big] = \mathcal{I}_\Psi\big[\,\mathrm{Ad}_{g(x)}\,\hat{Q}_x^{[\chi]}[k]\big] \quad \text{(Definitions 9.7 and 9.127)},$$

so the scalar readout is invariant under simultaneous (co-)transform of observable and band state. Moreover, by Nielsen safety on the band (Proposition 9.97) any gauge-parameter variation that is BRST-exact leaves $\mathcal{I}, \mathcal{I}_\Psi$ unchanged; hence the value is independent of the choice of graded-gauge frame within the band.

Combining (a)–(c) proves the claim: $\hat{Q}_x^{[\chi]}[k]$ is Hermitian, transforms covariantly, and its band evaluations are real and frame-independent.

□

Definition 9.111 (Typed hyperbolicity (operator test, rephrased))**.** A typed subspace χ is *hyperbolic* at x if

$$\inf_{\Psi \in \mathcal{S}_{\mathrm{band}}} \mathcal{I}_\Psi\big[\,\hat{Q}_x^{[\chi]}[k]\,\big] \;>\; 0 \quad \text{for all timelike } k \in \mathcal{C}_x^{\hat{\Sigma},*}, \tag{9.53}$$

which is the typed version of (9.42) (cf. Definition 9.101).

Proposition 9.112 (Typed hyperbolicity ⇒ well-posedness on the band)**.** *Assume Definition 9.111. Then for every compact band window and for every timelike covector k, the quadratic form $\Psi \mapsto \mathcal{I}_\Psi\big[\hat{Q}_x^{[\chi]}[k]\big]$ is strictly positive, yielding an a priori energy estimate for the typed sector χ. In particular, χ admits a retarded Green operator supported in $J^+\big(\mathcal{C}^{\hat{\Sigma},*}\big)$, compatible with Proposition 9.107.*

Proof of Proposition 9.112. Fix a permitted type χ and a compact band window $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ (Definition 9.96). Recall the typed principal operator and its quadratic form

$$\hat{Q}_x^{[\chi]}[k] \;:=\; \Pi^{[\chi]}\,\hat{Q}_x[k]\,\Pi^{[\chi]} \quad \text{with} \quad \hat{Q}_x[k] = k^\mu k^\nu \hat{\Sigma}_{\mu\nu}(x) \quad \text{(Definitions 9.95 and 9.109)}.$$

By Lemma 9.110, $\hat{Q}_x^{[\chi]}[k]$ is Hermitian and gauge-covariant; hence $\mathcal{I}_\Psi[\hat{Q}_x^{[\chi]}[k]] \in \mathbb{R}$ for all band functionals $\mathcal{I}_\Psi$ (Definition 9.96).

Step 1: Strict positivity (coercivity) on the band. Typed hyperbolicity (Definition 9.111, eq. (9.53)) asserts that for every cone-timelike $k \in \mathcal{C}_x^{\hat{\Sigma},*}$ (Definition 9.104)

$$\inf_{\Psi \in \mathcal{S}_{\mathrm{band}}} \mathcal{I}_\Psi\Big[\hat{Q}_x^{[\chi]}[k]\Big] \;>\; 0. \tag{9.54}$$

Consequently, the band energy functional $\mathcal{E}_k^{[\chi]}[\Psi] := \mathcal{I}_\Psi[\hat{Q}_x^{[\chi]}[k]]$ is a strictly positive quadratic form on the (ghost number 0) band class. For any fixed compact band window the infimum in (9.54) is bounded away from 0 uniformly in x (by compactness plus the stability clause of Definition 9.116); thus $\mathcal{E}_k^{[\chi]}$ is *coercive* on the band. This yields the claimed "a priori energy" estimate:

$$\mathcal{E}_k^{[\chi]}[\Psi] \;\geq\; c_k\,\|\Psi\|_{\mathrm{band}}^2, \qquad c_k > 0, \tag{9.55}$$

for some band norm $\|\cdot\|_{\mathrm{band}}$ equivalent to $\Psi \mapsto \mathcal{E}_k^{[\chi]}[\Psi]^{1/2}$ on the window.

Step 2: Retarded parametrix and Green operator. Let $\mathcal{L}^{[\chi]}$ denote any band-evolution operator whose principal quadratic form at $(x;k)$ is $\hat{Q}_x^{[\chi]}[k]$ (e.g. the typed reduction of the dynamical operator in the χ-channel; its explicit subprincipal content is immaterial here). By (9.54), the principal symbol is (cone-)elliptic on timelike k, so one can construct a microlocal inverse symbol on $\mathcal{C}^{\hat{\Sigma},*}$ and hence a *retarded parametrix* G_0^{ret} supported in $J^+(\mathcal{C}^{\hat{\Sigma},*})$ by cone-gating (Definition 9.106) and Proposition 9.107. The defect $R := \mathbf{1} - \mathcal{L}^{[\chi]}G_0^{\mathrm{ret}}$ is a band-smoothing, retarded, cone-gated operator (closure properties in Definitions 9.96 and 9.106 and Proposition 9.107). On a compact band window, R has small operator norm after possibly refining the retarded partition (Volterra structure in time along a cone-timelike flow), so the Neumann series

$$G^{\mathrm{ret}} \;:=\; G_0^{\mathrm{ret}} \sum_{m=0}^{\infty} R^m$$

converges in the band operator norm and defines a *retarded Green operator* with the same cone support: $\mathcal{L}^{[\chi]}G^{\mathrm{ret}} = \mathbf{1}$, $\operatorname{supp} G^{\mathrm{ret}} \subseteq J^+(\mathcal{C}^{\hat{\Sigma},*})$ (Proposition 9.107).

Step 3: Uniqueness and continuous dependence (well-posedness). Let u solve $\mathcal{L}^{[\chi]}u = f$ with past-compact support. Applying the energy functional to a standard retarded regularization of u and using the Hermiticity of $\hat{Q}_x^{[\chi]}[k]$ (Lemma 9.110) yields $\partial_\tau \mathcal{E}_k^{[\chi]}[u(\tau)] \leq C\,\mathcal{E}_k^{[\chi]}[u(\tau)] + C'\,\|f(\tau)\|_{\mathrm{band}}^2$, for some band-bounded C, C' (subprincipal contributions). Grönwall's inequality and (7.490) give the a priori bound

$$\|u\|_{\mathrm{band};\,J^+(\tau_0)}^2 \;\lesssim\; \mathcal{E}_k^{[\chi]}[u(\tau_0)] + \int_{\tau_0}^{\infty} \|f(\tau)\|_{\mathrm{band}}^2\,\mathrm{d}\tau,$$

which implies uniqueness for retarded solutions with vanishing past data and continuous dependence on f (well-posedness on the band). Existence is furnished by $u = G^{\mathrm{ret}}f$ from Step 2. Cone-gating ensures microcausal support (Proposition 9.108).

Combining Steps 1–3: typed hyperbolicity provides coercivity; the cone-gated parametrix yields a retarded Green operator supported in $J^+(\mathcal{C}^{\hat{\Sigma},*})$ (Proposition 9.107); and the energy estimate establishes uniqueness and continuous dependence. Hence the typed sector χ is well-posed on the band, as claimed. □

Corollary 9.113 (Exclusion rule)**.** *If* (9.53) *fails for some timelike k, then χ is excluded from the permit ledger of Proposition 9.102; i.e.,*

$$\chi \notin \mathcal{C}^{[\text{apeiron}]}_{\text{admissible}} \ \textit{(resp. } \mathcal{C}^{[\text{kymon}]}_{\text{admissible}}\textit{)}.$$

Proof of Corollary 9.113. Assume that (9.53) fails for some cone–timelike covector $k \in \mathcal{C}^{\hat{\Sigma},*}_x$; explicitly,

$$\inf_{\Psi\in\mathcal{S}_{\text{band}}} \mathcal{I}_\Psi\big[\hat{Q}^{[\chi]}_x[k]\big] \ \le\ 0,$$

contradicting the typed hyperbolicity requirement of Definition 9.111. By the *necessity* direction in the permit–ledger characterization (Proposition 9.102, using Definitions 9.116 and 9.122), membership $\chi \in \mathcal{C}^{[\bullet]}_{\text{admissible}}$ for either block $\bullet \in \{\Phi\Phi, \Theta\Theta\}$ requires typed hyperbolicity to hold for all cone–timelike k. Hence the failure at this k forces $\chi \notin \mathcal{C}^{[\bullet]}_{\text{admissible}}$.

Moreover, by Proposition 9.112 typed hyperbolicity implies a coercive energy estimate and the existence of a retarded Green operator supported in $J^+(\mathcal{C}^{\hat{\Sigma},*})$; the failure of (9.53) precludes this certificate, which is part of the ledger entry (Definition 9.122). Therefore χ is excluded from the permit ledger, as claimed. □

Remark 9.114 (Gauge-covariant transport and frame choice)**.** Let g be a local graded-gauge element and $\mathcal{U}_\gamma(x \leftarrow y)$ the parallel transport along a path γ as in Definition 9.7. All typed objects used above transform by conjugation:

$$\begin{aligned}
\hat{\Sigma}_{\mu\nu} &\mapsto \mathrm{Ad}_g\, \hat{\Sigma}_{\mu\nu},\\
\hat{Q}_x[v] &\mapsto \mathrm{Ad}_{g(x)}\, \hat{Q}_x[v] \quad (\textit{Definition 9.95}),\\
\Pi^{[\chi]} &\mapsto \mathrm{Ad}_{g(x)}\, \Pi^{[\chi]} \quad (\textit{Definition 9.116}).
\end{aligned}$$

Hence the typed principal form transforms covariantly, $\hat{Q}^{[\chi]}_x[v] \mapsto \mathrm{Ad}_{g(x)}\, \hat{Q}^{[\chi]}_x[v]$, and retains Hermiticity (Lemma 9.110). Band readouts obey the co-variance rule

$$\mathcal{I}_{g\cdot\Psi}\big[\hat{O}\big] = \mathcal{I}_\Psi\big[\,\mathrm{Ad}_g\, \hat{O}\big], \qquad \mathcal{I}_{g\cdot\Psi}\big[\hat{O}^{[\chi]}\big] = \mathcal{I}_\Psi\big[\,\mathrm{Ad}_g\, \hat{O}^{[\chi]}\big],$$

so any scalar diagnostic built from $(\hat{O}, \Psi)$ is invariant under a simultaneous gauge-frame change. In particular, the typed visibilities $\Lambda^{\Sigma;[\chi]}_{\text{law/state}}$ (Definition 9.133), the quadrant predicate $\mathrm{Quad}^{[\chi]}$ (Definition 9.137), and the parasymmetric valuations $v^{(\cdot);[\chi]}_{\text{Synthesis}}$ (Definitions 9.143 and 9.144) are preserved under (i) local changes of modal frames inside the $\{\Phi\Phi, \Theta\Theta\}$ blocks and the dialectical basis $\{0, 1, p, s\}$ (Definition 9.94), and (ii) graded parallel transport $\mathcal{U}_\gamma$ between nearby points, because the cone gate and retardedness are transported functorially (Definition 9.106 and Proposition 9.107). Moreover, any residual gauge-parameter dependence that is BRST-exact is annihilated by the band functionals (Proposition 9.97), ensuring Nielsen safety throughout.

Remark 9.115 (Quasi-classical limit)**.** Outside the scope fixed by the operator-only axiom (Axiom 9.89), suppose an expectation layer is admissible with $\bar{\Sigma}_{\mu\nu} := E_\omega[\hat{\Sigma}_{\mu\nu}]$ (Definition 9.88). Then the Σ–pre-physical cone $\mathcal{C}_x^{\hat{\Sigma},*}$ (Definition 9.104) approaches the Lorentzian covector cone associated with $\bar{\Sigma}$:

$$\mathcal{C}_x^{\bar{\Sigma},*} = \Big\{ k \neq 0 : \; k^\mu k^\nu \, \bar{\Sigma}_{\mu\nu}(x) \leq 0 \Big\},$$

in the following sense. If (a) the band is tightened ($\ell \downarrow$, $\lambda_{\mathrm{UV}} \uparrow$) so that the uncertainty rectangles for the visibilities collapse (Definition 9.168 and Lemma 9.169), (b) the BRST-closed band functionals converge weak* to the expectation E_ω on BRST cohomology classes (Proposition 9.97), and (c) the typed projectors remain stable under this retuning (Definition 9.116), then the defining supremum in Definition 9.104 Γ–converges to the classical quadratic test with $\bar{\Sigma}$. Consequently, the cone gate and retardedness limit to their classical forms (Proposition 9.107), and microcausality reduces to the standard vanishing of spacelike commutators with respect to $\bar{\Sigma}$ (Proposition 9.108). Within the present section we do *not* invoke $\bar{\Sigma}_{\mu\nu}$; the operator-level $\mathcal{C}_x^{\hat{\Sigma},*}$ remains the primitive causal structure, and all diagnostics (typed hyperbolicity Definition 9.111, well-posedness Proposition 9.112, and permit ledgers Proposition 9.102) are established purely on that substrate. Any mapping of permitted χ–sectors to quasi-particle/graviton content belongs to the expectation-level regime and is thus outside the scope here (cf. Axiom 9.89).

9.10 Kinematic Permit Census for Apeiron/Kymon (No *a priori* Spins)

2.1 Typed band-local projectors

Definition 9.116 (Typed band-local projectors, block-internal and graded-covariant)**.** Let $\hat{\mathbf{G}}_\Lambda$ be the unified block correlator of Definition 9.94 with diagonal sector idempotents

$$E_{\Phi\Phi} := \left(\begin{smallmatrix} \mathbb{I} & 0 \\ 0 & 0 \end{smallmatrix}\right), \qquad E_{\Theta\Theta} := \left(\begin{smallmatrix} 0 & 0 \\ 0 & \mathbb{I} \end{smallmatrix}\right) \quad \text{(cf. eq. (9.36))}.$$

A *typed band-local projector* $\Pi^{[\chi]}$ is a bounded operator on $\mathcal{H}_{\mathrm{em}}$ satisfying:

(P1) **Projector & Hermiticity:** $(\Pi^{[\chi]})^2 = \Pi^{[\chi]}$ and $(\Pi^{[\chi]})^\dagger = \Pi^{[\chi]}$.

(P2) **Block-internality:** either $E_{\Phi\Phi}\Pi^{[\chi]} = \Pi^{[\chi]}E_{\Phi\Phi} = \Pi^{[\chi]}$ (apeiron, $\Phi\Phi$) or $E_{\Theta\Theta}\Pi^{[\chi]} = \Pi^{[\chi]}E_{\Theta\Theta} = \Pi^{[\chi]}$ (kymon, $\Theta\Theta$).

(P3) **Band-finiteness:** $P_x(\lambda_{\mathrm{UV}})\,\Pi^{[\chi]} = \Pi^{[\chi]} = \Pi^{[\chi]}P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}\,\Pi^{[\chi]} = \Pi^{[\chi]} = \Pi^{[\chi]}K_{\mathrm{IR}}^{(\ell)}$ (Definition 9.96).

(P4) **BRST-closedness (ghost 0):** $[\hat{Q}_{\mathrm{BRST}}, \Pi^{[\chi]}] = 0$ and $\mathrm{gh}(\Pi^{[\chi]}) = 0$ (Lemma 9.54).

(P5) **Graded-gauge covariance:** under a local graded frame $g(x)$ (Definition 9.7), $\Pi^{[\chi]} \mapsto g\,\Pi^{[\chi]}g^\dagger$.

(P6) **Cone-compatibility:** for every timelike $k \in \mathcal{C}_x^{\hat{\Sigma},*}$ (Definition 9.104), the typed probe $\hat{Q}_x^{[\chi]}[k] = \Pi^{[\chi]}\hat{Q}_x[k]\Pi^{[\chi]}$ (Definition 9.109) is positive in the band-functional sense (see Definition 9.111).

(P7) **Retuning-stability:** there exists $\delta > 0$ such that for all $(\lambda'_{\mathrm{UV}}, \ell')$ with $|\lambda'_{\mathrm{UV}} - \lambda_{\mathrm{UV}}| + |\ell' - \ell| < \delta$, the map $(\lambda'_{\mathrm{UV}}, \ell') \mapsto \Pi^{[\chi]}(\lambda'_{\mathrm{UV}}, \ell')$ is norm-continuous and preserves (P1)–(P6).

Remark 9.117 (Typed structures per sector)**.** Within **apeiron** ($\Phi\Phi$) the standard constructions are: Σ-trace / trace-free split; Σ-divergence transverse/longitudinal split; symmetric / TT-like subspaces; and, whenever a Σ-spatial direction $\hat{\mathbf{k}}$ is meaningful (timelike k picks a spatial hyperplane), helicity-refined projectors. Within **kymon** ($\Theta\Theta$): Σ-transverse / Σ-longitudinal, helicity $\lambda = \pm$, and the torsional/temporal *Chrono–Kymon* channel via $\hat{t}^{(\Theta)}_{\mu\nu} \subset \hat{\Sigma}^{(\Theta)}$ (cf. Definition 5.25 and the Σ-weighted mode extraction of Proposition 9.36 and eq. (9.16)).

Lemma 9.118 (Σ-orthogonality on the band)**.** *If $\Pi^{[\chi]}$ and $\Pi^{[\chi']}$ satisfy (P1)–(P6) of Definition 9.116 and $\Pi^{[\chi]}\Pi^{[\chi']} = 0$, then for any timelike k*

$$\mathcal{I}_\Psi\Big[\hat{Q}^{[\chi]}_x[k]\Pi^{[\chi']}\Big] \;=\; 0 \quad \textit{and} \quad \mathcal{I}_\Psi\Big[\Pi^{[\chi]}\hat{Q}^{[\chi']}_x[k]\Big] \;=\; 0,$$

for all $\Psi \in \mathcal{S}_{\mathrm{band}}$. Hence distinct typed subspaces are orthogonal with respect to the Σ-induced quadratic form (Definition 9.22).

Proof of Lemma 9.118. Recall the Σ–probe and its typed restriction

$$\hat{Q}_x[v] = v^\mu v^\nu \hat{\Sigma}_{\mu\nu}(x) \quad \text{(Definition 9.95)},$$
$$\hat{Q}^{[\chi]}_x[v] = \Pi^{[\chi]}\,\hat{Q}_x[v]\,\Pi^{[\chi]} \quad \text{(cf. (9.52), Definitions 9.109 and 9.116)}.$$

Let $\Pi^{[\chi]}$ and $\Pi^{[\chi']}$ satisfy items (P1)–(P6) of Definition 9.116, and assume $\Pi^{[\chi]}\Pi^{[\chi']} = 0$ (orthogonal projectors in the block algebra).

Step 1: Algebraic annihilation of cross-typed factors. Using idempotence and orthogonality (Definition 9.116(P1)),

$$\hat{Q}^{[\chi]}_x[k]\;\Pi^{[\chi']} = \Big(\Pi^{[\chi]}\,\hat{Q}_x[k]\,\Pi^{[\chi]}\Big)\Pi^{[\chi']} = \Pi^{[\chi]}\,\hat{Q}_x[k]\,\underbrace{\Big(\Pi^{[\chi]}\Pi^{[\chi']}\Big)}_{=0} = 0,$$

and symmetrically,

$$\Pi^{[\chi]}\,\hat{Q}^{[\chi']}_x[k] = \Pi^{[\chi]}\Big(\Pi^{[\chi']}\,\hat{Q}_x[k]\,\Pi^{[\chi']}\Big) = \underbrace{\Big(\Pi^{[\chi]}\Pi^{[\chi']}\Big)}_{=0}\,\hat{Q}_x[k]\,\Pi^{[\chi']} = 0.$$

These identities hold in the band algebra for any covector k; the timelike assumption (from Definition 9.104) is only needed elsewhere for positivity, not for orthogonality.

Step 2: Vanishing of band evaluations. By linearity of the band functionals (Definition 9.96),

$$\mathcal{I}_\Psi\Big[\hat{Q}^{[\chi]}_x[k]\Pi^{[\chi']}\Big] = \mathcal{I}_\Psi[0] = 0, \qquad \mathcal{I}_\Psi\Big[\Pi^{[\chi]}\hat{Q}^{[\chi']}_x[k]\Big] = \mathcal{I}_\Psi[0] = 0,$$

for all $\Psi \in \mathcal{S}_{\mathrm{band}}$.

Step 3: Σ–orthogonality of typed subspaces. Denote the Σ–induced quadratic form by $\hat{Q}_x[k]$ (Definitions 9.22 and 9.95). For vectors $u \in \operatorname{Ran}\Pi^{[\chi]}$ and $v \in \operatorname{Ran}\Pi^{[\chi']}$,

$$\mathcal{I}_\Psi\Big[u^\dagger\,\hat{Q}_x[k]\,v\Big] = \mathcal{I}_\Psi\Big[u^\dagger\,\Pi^{[\chi]}\hat{Q}_x[k]\Pi^{[\chi']}\,v\Big] = \mathcal{I}_\Psi\Big[u^\dagger\,\Pi^{[\chi]}\hat{Q}^{[\chi']}_x[k]\,v\Big] = 0,$$

by Step 1. Thus distinct typed subspaces are orthogonal with respect to the Σ–induced quadratic form, as claimed. □

Proposition 9.119 (Constructive existence via Σ-weighted mode operators)**.** *Let $K_\bullet$ be the Σ-weighted positive compact operator constructed as in Proposition 9.36 and eq. (9.16) inside a fixed block ($\Phi\Phi$ or $\Theta\Theta$). Then the spectral Riesz projectors of $K_\bullet$ onto any finite union of isolated spectral clusters yield band-finite, BRST-closed, graded-covariant projectors that satisfy (P1)–(P7). In particular, helicity and TT-like subspaces are obtained by choosing $K_\bullet$ that commute with the appropriate stabilizer generators (little group) on the Σ-spatial hyperplane determined by a timelike k (Definition 9.104).*

Proof of Proposition 9.119. Fix a block $\bullet \in \{\Phi\Phi, \Theta\Theta\}$ and the Σ-weighted positive compact operator $K_\bullet$ constructed in Proposition 9.36 (generalized Σ-eigenproblem (9.16)), with UV/IR band instrumentation $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ (Definition 9.96) and cone-gating (Definition 9.106). Let $\{\lambda_j\}$ be the (discrete) spectrum of $K_\bullet$, with each nonzero $\lambda_j > 0$ of finite multiplicity (compact positivity). Let $\Gamma = \bigsqcup_{m=1}^{M} \Gamma_m$ be a finite union of pairwise disjoint Jordan curves, each enclosing an *isolated* spectral cluster σ_m and no other spectrum. Define the Riesz projector

$$\Pi_\Gamma \;:=\; \frac{1}{2\pi i} \oint_\Gamma (z\mathbf{1} - K_\bullet)^{-1}\, \mathrm{d}z. \tag{9.56}$$

We prove that Π_Γ satisfies items (P1)–(P7) of Definition 9.116.

(i) Projector & block internality ((P1)). By the holomorphic functional calculus, (9.56) yields an idempotent that commutes with $K_\bullet$, and since $K_\bullet$ is self-adjoint/positive on the block Hilbert module, Π_Γ is the *orthogonal* spectral projector onto $\mathcal{H}_\Gamma = \bigoplus_{\lambda_j \in \cup_m \sigma_m} \ker(K_\bullet - \lambda_j)$. Thus $\Pi_\Gamma^\dagger = \Pi_\Gamma = \Pi_\Gamma^2$ and $\Pi_\Gamma \in \mathrm{End}(\bullet)$.

(ii) Band-finiteness ((P3)). By construction of $K_\bullet$ (Proposition 9.36) the kernel defining $K_\bullet$ is band-admissible (UV-projected, IR-filtered, cone-gated: $P_x(\lambda_{\mathrm{UV}})K_\bullet = K_\bullet = K_\bullet P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)} K_\bullet = K_\bullet K_{\mathrm{IR}}^{(\ell)}$; see Definition 9.96). Resolvents $(z\mathbf{1} - K_\bullet)^{-1}$ preserve band-admissibility for $z \notin \sigma(K_\bullet)$, hence the contour integral (9.56) is band-admissible. Moreover, each cluster σ_m has finite rank in the band window, so Π_Γ is *band-finite*.

(iii) BRST-closedness. $K_\bullet$ is built from BRST-closed ingredients (cone-gated $\hat{\Sigma}$, band taggers of ghost number 0; Definitions 5.25 and 9.96), hence $[\,\hat{Q}_{\mathrm{BRST}}, K_\bullet\,] = 0$. Applying the functional calculus under the BRST derivation gives $[\,\hat{Q}_{\mathrm{BRST}}, \Pi_\Gamma\,] = \frac{1}{2\pi i} \oint_\Gamma (z\mathbf{1} - K_\bullet)^{-1} [\,\hat{Q}_{\mathrm{BRST}}, K_\bullet\,] (z\mathbf{1} - K_\bullet)^{-1} \mathrm{d}z = 0$, so Π_Γ is BRST-closed.

(iv) Graded covariance ((P5)). Under local graded transport g (Definition 9.7), $K_\bullet \mapsto \mathrm{Ad}_g\, K_\bullet$. Hence $(z\mathbf{1} - \mathrm{Ad}_g\, K_\bullet)^{-1} = \mathrm{Ad}_g (z\mathbf{1} - K_\bullet)^{-1}$, and

$$\Pi_\Gamma \;\mapsto\; \frac{1}{2\pi i} \oint_\Gamma (z\mathbf{1} - \mathrm{Ad}_g\, K_\bullet)^{-1} \mathrm{d}z = \mathrm{Ad}_g \Big(\frac{1}{2\pi i} \oint_\Gamma (z\mathbf{1} - K_\bullet)^{-1} \mathrm{d}z \Big) = \mathrm{Ad}_g\, \Pi_\Gamma,$$

so Π_Γ transforms covariantly.

(v) Cone-compatibility ((P6)). By construction $K_\bullet$ is Σ-weighted and cone-gated (Proposition 9.36 and Definition 9.106); its eigenmodes are obtained from kernels supported in $J^+(\mathcal{C}^{\hat{\Sigma},*})$ and therefore lie in the domain on which the Σ–principal form is defined. For any timelike $k \in \mathcal{C}_x^{\hat{\Sigma},*}$ (Definition 9.104), the typed principal form $\hat{Q}_x^{[\Pi_\Gamma]}[k] = \Pi_\Gamma\, \hat{Q}_x[k] \Pi_\Gamma$ is Hermitian and gauge-covariant (Lemma 9.110). Since $K_\bullet$ is built from $\hat{\Sigma}$ and its cone-gated

propagators, the range of Π_Γ is a Σ–compatible subspace in which timelike directions remain timelike for the quadratic form (*compatibility* in the sense of (P6)). (If one additionally demands *positivity* to certify typed hyperbolicity, this is precisely the test of Definition 9.111, verified separately when admitting the type.)

(vi) Stability under small retunings ((P7)). Retuning the band parameters $(\lambda_{\mathrm{UV}}, \ell) \mapsto (\lambda'_{\mathrm{UV}}, \ell')$ produces a bounded perturbation of $K_\bullet$ that is continuous in operator norm ($\|K'_\bullet - K_\bullet\| \to 0$; see Definition 9.96 and Lemma 9.105). By Kato's perturbation theory for isolated spectral clusters (norm-resolvent continuity), the Riesz projector Π_Γ varies continuously provided Γ stays separated from the rest of the spectrum. Thus there exists a stability radius $\delta > 0$ such that for $|\lambda'_{\mathrm{UV}} - \lambda_{\mathrm{UV}}| + |\ell' - \ell| < \delta$ the spectral gap persists and $\|\Pi'_\Gamma - \Pi_\Gamma\| \to 0$, realizing (P7).

Items (P1)–(P7) of Definition 9.116 are therefore satisfied.

Helicity and TT-like subspaces. Let W_k denote the stabilizer (little group) of a timelike k in the Σ-spatial hyperplane (Definition 9.104). Choose $K_\bullet$ so that it *commutes* with the representation of W_k (e.g. by averaging the kernel over W_k inside the band). Then the spectral subspaces of $K_\bullet$ carry W_k-representations and decompose into helicity (or TT-like) sectors; the corresponding Riesz projectors Π_Γ are the helicity-/TT-type projectors. These inherit band-finiteness, BRST-closedness, graded covariance, cone-compatibility, and stability from the preceding steps, completing the constructive existence.

□

Corollary 9.120 (Typed hyperbolicity certificates)**.** *For each $\Pi^{[\chi]}$ produced by Proposition 9.119, the positivity condition (9.53) holds if and only if the corresponding Σ-weighted spectral band is separated from 0. The* cone/hyperbolicity certificate *attached to χ consists of: (i) the admissible timelike interval in $\mathcal{C}_x^{\hat{\Sigma},*}$; (ii) the spectral gap margin of $K_\bullet$; and (iii) the retuning-stability radius from (P7).*

Proof of Corollary 9.120. Let $\Pi^{[\chi]} = \Pi_\Gamma$ be a Riesz projector produced by Proposition 9.119 from the Σ–weighted positive compact operator $K_\bullet$ (constructed in Proposition 9.36 via the generalized eigenproblem (9.16)) inside a fixed block $\bullet \in \{\Phi\Phi, \Theta\Theta\}$. Write $\sigma_\Gamma := \sigma(K_\bullet) \cap \mathrm{int}(\Gamma)$ for the enclosed spectral band.

Step 1: Equivalence between typed quadratic energy and $K_\bullet$–energy on $\mathrm{Ran}\,\Pi_\Gamma$. By construction of $K_\bullet$ and the typed reduction $\hat{Q}_x^{[\chi]}[k] = \Pi_\Gamma\, \hat{Q}_x[k] \Pi_\Gamma$ (with $\hat{Q}_x[k] = k^\mu k^\nu \hat{\Sigma}_{\mu\nu}(x)$, Definitions 9.95 and 9.109), there exist band–window–uniform positive constants $c_1(k), c_2(k)$ such that for every band-admissible vector/mode $u \in \mathrm{Ran}\,\Pi_\Gamma$ and every band functional $\mathcal{I}_\Psi$ (Definition 9.96),

$$c_1(k)\,\langle u,\, K_\bullet u\rangle \;\le\; \mathcal{I}_\Psi\Big[u^\dagger\, \hat{Q}_x^{[\chi]}[k]\, u\Big] \;\le\; c_2(k)\,\langle u,\, K_\bullet u\rangle. \tag{9.57}$$

This is the "principal-modes equivalence" induced by the Σ–weighting in Proposition 9.36 (and the cone-gated construction of kernels, Definition 9.106 and Proposition 9.107), together with Hermiticity/covariance of the typed principal form (Lemma 9.110). Equation (9.57) expresses that, on $\mathrm{Ran}\,\Pi_\Gamma$, the band evaluation of the typed principal quadratic form is equivalent to the $K_\bullet$–energy.

Step 2: "If" direction. Assume the spectral band is separated from zero:

$$\delta_\Gamma \;:=\; \operatorname{dist}\big(0, \sigma_\Gamma\big) \;>\; 0.$$

Then $\langle u, K_\bullet u\rangle \geq \delta_\Gamma \, \|u\|^2$ for all $u \in \operatorname{Ran} \Pi_\Gamma$. Combining with (9.57) gives, for every band functional $\mathcal{I}_\Psi$ and every timelike $k \in \mathcal{C}_x^{\hat{\Sigma},*}$ (Definition 9.104),

$$\mathcal{I}_\Psi\Big[u^\dagger \, \hat{Q}_x^{[\chi]}[k] \, u\Big] \;\geq\; c_1(k) \, \delta_\Gamma \, \|u\|^2.$$

Since the band functionals separate the positive cone on the band algebra (Definition 9.96), taking the infimum over admissible Ψ yields

$$\inf_{\Psi \in \mathcal{S}_{\text{band}}} \; \mathcal{I}_\Psi\Big[\hat{Q}_x^{[\chi]}[k]\Big] \;>\; 0,$$

which is exactly the typed hyperbolicity condition (9.53) (Definition 9.111). Hence spectral separation from 0 implies typed hyperbolicity.

Step 3: "Only if" direction. Conversely, suppose (9.53) holds but σ_Γ is *not* separated from 0. Then there exists a sequence $u_n \in \operatorname{Ran} \Pi_\Gamma$ with $\|u_n\| = 1$ and $\langle u_n, K_\bullet u_n\rangle \to 0$. By (9.57) we obtain

$$\inf_{\Psi} \; \mathcal{I}_\Psi\Big[u_n^\dagger \, \hat{Q}_x^{[\chi]}[k] \, u_n\Big] \;\leq\; c_2(k) \, \langle u_n, K_\bullet u_n\rangle \;\longrightarrow\; 0,$$

contradicting (9.53). Therefore, typed hyperbolicity forces $\delta_\Gamma > 0$, i.e. σ_Γ is separated from 0.

Step 4: Certificate components. Given $\Pi^{[\chi]} = \Pi_\Gamma$ with $\delta_\Gamma > 0$, the *cone/hyperbolicity certificate* attached to χ is:

(i) the admissible timelike interval of k in $\mathcal{C}_x^{\hat{\Sigma},*}$ (where the constants $c_{1,2}(k)$ and the inequality (9.57) hold; cf. Definition 9.104);

(ii) the spectral gap margin $\delta_\Gamma = \operatorname{dist}(0, \sigma_\Gamma)$ controlling the positive lower bound in (9.57) and thus in (9.53);

(iii) the retuning-stability radius supplied by item (P7) of Definition 9.116, realized via norm–resolvent stability of Riesz projectors under band retunings in Proposition 9.119 (see also IR/UV monotonicity Lemma 9.105 and Definition 9.96).

These three data together certify typed hyperbolicity on the band and its robustness under the allowed instrumentation.

The equivalence and the stated certificate are thus established. □

Lemma 9.121 (BRST/Nielsen safety of $\Pi^{[\chi]}$)**.** *If $\Pi^{[\chi]}$ satisfies (P4) of Definition 9.116, then for any BRST-exact variation $\delta_\theta \Pi^{[\chi]} = [\hat{Q}_{\text{BRST}}, X_\theta]$ one has $\mathcal{I}(\delta_\theta \Pi^{[\chi]}) = \mathcal{I}_\Psi(\delta_\theta \Pi^{[\chi]}) = 0$ by Proposition 9.97. Thus all downstream typed contrasts/visibilities (Definitions 9.45 and 9.47) are gauge-parameter independent.*

Proof of Lemma 9.121. Assume $\Pi^{[\chi]}$ satisfies the BRST-closedness item (P4) of Definition 9.116, i.e. $[\hat{Q}_{\text{BRST}}, \Pi^{[\chi]}] = 0$. Let a gauge-parameter variation act by a BRST-exact commutator $\delta_\theta \Pi^{[\chi]} = [\hat{Q}_{\text{BRST}}, X_\theta]$ (band-admissible X_θ).

Step 1: Scalar band evaluations of $\delta_\theta \Pi^{[\chi]}$ vanish. By the Nielsen safety on the band (Proposition 9.97) the band functionals annihilate BRST-exact insertions:

$$\mathcal{I}\big([\hat{Q}_{\mathrm{BRST}}, Y]\big) = \mathcal{I}_\Psi\big([\hat{Q}_{\mathrm{BRST}}, Y]\big) = 0 \qquad \text{for all band-admissible } Y.$$

With $Y = X_\theta$, this gives

$$\mathcal{I}(\delta_\theta \Pi^{[\chi]}) = \mathcal{I}\big([\hat{Q}_{\mathrm{BRST}}, X_\theta]\big) = 0, \qquad \mathcal{I}_\Psi(\delta_\theta \Pi^{[\chi]}) = \mathcal{I}_\Psi\big([\hat{Q}_{\mathrm{BRST}}, X_\theta]\big) = 0,$$

proving the first claim in the lemma.

Step 2: Typed contrasts/visibilities are independent of the frame choice of $\Pi^{[\chi]}$. Let A be any band operator entering the typed restriction (e.g. $A = \hat{Q}_x[v]$ or $A = \hat{Q}_x[v] - \mathrm{Ad}_{g(x)}\,\hat{Q}_x[v]$ in the definitions of the typed contrasts, Definition 9.127). Consider the scalar readout $\mathcal{I}\big(\Pi^{[\chi]} A \Pi^{[\chi]}\big)$ (and similarly with $\mathcal{I}_\Psi$). Varying only the projector (holding A fixed) yields

$$\partial_\theta\, \mathcal{I}\big(\Pi^{[\chi]} A \Pi^{[\chi]}\big) = \mathcal{I}\big((\delta_\theta \Pi^{[\chi]}) A \Pi^{[\chi]}\big) + \mathcal{I}\big(\Pi^{[\chi]} A (\delta_\theta \Pi^{[\chi]})\big).$$

Using $[\hat{Q}_{\mathrm{BRST}}, \Pi^{[\chi]}] = 0$ and the graded derivation property, each term can be re-expressed as a BRST-exact insertion:

$$(\delta_\theta \Pi^{[\chi]}) A \Pi^{[\chi]} = \big[\hat{Q}_{\mathrm{BRST}},\ X_\theta A \Pi^{[\chi]}\big] \ -\ X_\theta\,[\hat{Q}_{\mathrm{BRST}}, A \Pi^{[\chi]}],$$

and analogously for $\Pi^{[\chi]} A (\delta_\theta \Pi^{[\chi]})$. The second pieces above vanish inside $\mathcal{I}$ and $\mathcal{I}_\Psi$ because the inputs used for typed contrasts are BRST-closed modulo exact terms by construction of the diagnostics (Definitions 9.96 and 9.127; see also the pointwise Nielsen consequence Lemma 9.128). The remaining pieces are of the form $\mathcal{I}([\hat{Q}_{\mathrm{BRST}}, \cdot])$ and hence vanish by Proposition 9.97. Therefore

$$\partial_\theta\, \mathcal{I}\big(\Pi^{[\chi]} A \Pi^{[\chi]}\big) = 0, \qquad \partial_\theta\, \mathcal{I}_\Psi\big(\Pi^{[\chi]} A \Pi^{[\chi]}\big) = 0.$$

Applying this with A chosen as in Definition 9.127 shows that both $\Delta^{\mathrm{pre},\Sigma;[\chi]}_{\mathrm{law}}$ and $\Delta^{\mathrm{pre},\Sigma;[\chi]}_{\mathrm{state}}$ are insensitive to the gauge-parameter variation of $\Pi^{[\chi]}$. Aggregation and normalization to visibilities (Definition 9.133) are algebraic operations on these scalars and therefore inherit the invariance.

Altogether, BRST-closedness of $\Pi^{[\chi]}$ combined with Nielsen safety of the band functionals (Proposition 9.97) implies $\mathcal{I}(\delta_\theta \Pi^{[\chi]}) = \mathcal{I}_\Psi(\delta_\theta \Pi^{[\chi]}) = 0$, and all downstream typed contrasts and visibilities (Definitions 9.127 and 9.133) are gauge-parameter independent with respect to variations of $\Pi^{[\chi]}$. □

2.2 Output: permit ledgers

Definition 9.122 (Permit ledgers (apeiron/kymon))**.** The *permit ledger* of admissible types in a sector is

$$\mathcal{C}^{[\mathrm{apeiron}]}_{\mathrm{admissible}} \subset \mathrm{End}(\Phi\Phi), \qquad \mathcal{C}^{[\mathrm{kymon}]}_{\mathrm{admissible}} \subset \mathrm{End}(\Theta\Theta),$$

consisting of all labels χ for which there exists a projector $\Pi^{[\chi]}$ satisfying (P1)–(P7) of Definition 9.116 and the typed hyperbolicity condition (9.53) (Definition 9.111). Each entry χ carries its *cone/hyperbolicity certificate* as in Corollary 9.120.

Proposition 9.123 (Ledger covariance and functoriality)**.** *Under graded local frames g, the ledger transforms by conjugation: $\Pi^{[\chi]} \mapsto g\Pi^{[\chi]}g^\dagger$, and $\mathcal{C}^{[\cdot]}_{\text{admissible}}$ is invariant up to this equivalence. Moreover, refining the band tag $\mathfrak{B}_{\lambda_{\text{UV}},\ell}$ (Definition 9.96) functorially* shrinks *the ledger but cannot add a χ that previously failed* (9.53) *(by Lemma 9.105 and* (P7)*).*

Proof of Proposition 9.123. We split the proof into covariance under graded frames and functorial monotonicity under band refinement.

(A) Covariance under graded local frames. Let g be a graded-gauge transport as in Definition 9.7. For any admitted label χ with projector $\Pi^{[\chi]}$ (in the block $\Phi\Phi$ or $\Theta\Theta$), items (P1), (P4), and (P5) of Definition 9.116 give

$$\Pi^{[\chi]} \mapsto \mathrm{Ad}_g\,\Pi^{[\chi]} := g\,\Pi^{[\chi]}g^\dagger, \qquad [\,\hat{Q}_{\text{BRST}}, \Pi^{[\chi]}] = 0 \implies [\,\hat{Q}_{\text{BRST}}, \mathrm{Ad}_g\,\Pi^{[\chi]}] = 0.$$

The principal Σ–probe and its typed restriction transform covariantly: $\hat{Q}_x[v] \mapsto \mathrm{Ad}_{g(x)}\,\hat{Q}_x[v]$ and $\hat{Q}^{[\chi]}_x[v] = \Pi^{[\chi]}\hat{Q}_x[v]\Pi^{[\chi]} \mapsto \mathrm{Ad}_{g(x)}\,\hat{Q}^{[\chi]}_x[v]$

(Definition 9.95 and Lemma 9.110). Hence the typed hyperbolicity condition (9.53) is frame-invariant, either directly from covariance of the quadratic form or via the gauge-parameter independence of the band readouts $\mathcal{I}, \mathcal{I}_\Psi$ (Propositions 9.91 and 9.97). Band-finiteness and stability ((P3), (P7)) are preserved under conjugation as well, because g acts by unitary equivalence on the band class (Definition 9.96). Thus, if χ is admitted then so is its conjugate class; conversely the inverse frame g^{-1} implies the equivalence. Therefore the ledger $\mathcal{C}^{[\cdot]}_{\text{admissible}}$ is invariant up to g–conjugation.

(B) Functoriality under band refinement; monotonicity (no additions). Consider a refinement of the band tag $\mathfrak{B}_{\lambda_{\text{UV}},\ell} \rightsquigarrow \mathfrak{B}_{\lambda'_{\text{UV}},\ell'}$ with $\lambda'_{\text{UV}} \geq \lambda_{\text{UV}}$ and $\ell' \leq \ell$ (Definition 9.96). By Lemma 9.105, the Σ–covector cone $\mathcal{C}^{\hat{\Sigma},*}_x[\ell'] \subseteq \mathcal{C}^{\hat{\Sigma},*}_x[\ell]$ *shrinks*. The cone gate and retardedness commute with the UV/IR instrumentation (Definition 9.106 and Proposition 9.107), so all typed objects restrict functorially to the refined band.

We show that refinement cannot add a label χ that previously failed (9.53). Suppose χ is *not* admitted at $(\lambda_{\text{UV}}, \ell)$. By the permit-criterion (Proposition 9.102) and the typed-certificate equivalence (Corollary 9.120), either:

(i) *Hyperbolicity failure:* there exists a timelike $k \in \mathcal{C}^{\hat{\Sigma},*}_x[\ell]$ with $\inf_\Psi \mathcal{I}_\Psi[\hat{Q}^{[\chi]}_x[k]] \leq 0$ (i.e. the Σ–weighted spectral band for $\Pi^{[\chi]}$ touches 0; equivalently, the gap margin $\delta_\Gamma = 0$ in Corollary 9.120); or

(ii) *Stability failure:* a nonzero retuning-stability radius ((P7)) cannot be certified.

Assume for contradiction that χ becomes admitted after refinement to $(\lambda'_{\text{UV}}, \ell')$. Then *both* typed hyperbolicity and stability must hold on the refined band. In particular, by Corollary 9.120 there is a positive spectral gap $\delta'_\Gamma > 0$ and a stability radius $r' > 0$ such that typed hyperbolicity persists under all retunings within distance r' of $(\lambda'_{\text{UV}}, \ell')$ (item (P7) of Definition 9.116).

Now enlarge the IR window slightly within this radius: choose $\ell'' \in (\ell', \min\{\ell' + r', \ell\})$. By Lemma 9.105 (applied in reverse), the cone expands $\mathcal{C}^{\hat{\Sigma},*}_x[\ell'] \subseteq \mathcal{C}^{\hat{\Sigma},*}_x[\ell'']$, and by stability the typed positivity and the gap $\delta'_\Gamma/2$ persist at ℓ''. Iterating finitely many times (partitioning the interval $[\ell', \ell]$ into steps of size $< r'$) yields a band parameter $\tilde{\ell}$ arbitrarily close to ℓ

for which typed hyperbolicity holds with a *uniform* positive margin. Passing to the limit (norm–resolvent continuity from Proposition 9.119 and the continuity of band functionals Definition 9.96) we conclude that typed hyperbolicity must also hold at ℓ, contradicting (i). Thus case (i) cannot occur.

If (ii) held at $(\lambda_{\mathrm{UV}}, \ell)$, then admission at the refined point would supply a positive stability radius $r' > 0$ there. The same stepping argument as above shows that stability propagates along a chain of small retunings, contradicting the original failure of stability. Hence case (ii) cannot occur either.

Therefore refinement cannot *add* a label that was excluded before. On the other hand, refinement may *remove* labels (the cone shrinks and spectral clusters may lose a uniform gap or band-finiteness under stricter UV/IR gates), so the ledger can only *shrink*. This proves functorial monotonicity:

$$\mathcal{C}^{[\cdot]}_{\mathrm{admissible}}(\lambda'_{\mathrm{UV}}, \ell') \;\subseteq\; \mathcal{C}^{[\cdot]}_{\mathrm{admissible}}(\lambda_{\mathrm{UV}}, \ell).$$

Combining (A) and (B) establishes the proposition. □

Proposition 9.124 (Partial Σ-orthogonal decomposition)**.** *Let* $\{\Pi^{[\chi_j]}\}_{j=1}^{N} \subset \mathcal{C}^{[\cdot]}_{\mathrm{admissible}}$ *be mutually orthogonal projectors (Lemma 9.118). Then, within the chosen block,*

$$\Pi_{\mathrm{adm}} := \sum_{j=1}^{N} \Pi^{[\chi_j]}$$

is again a band-finite, BRST-closed projector, and the residual $\Pi_{\mathrm{res}} := E_{\bullet\bullet} - \Pi_{\mathrm{adm}}$ *is* Σ*-orthogonal to each* $\Pi^{[\chi_j]}$ *on the band. If* Π_{res} *fails* (9.53)*, its image is excluded from the ledger (Corollary 9.113).*

Proof of Proposition 9.124. Fix a block $\bullet \in \{\Phi\Phi, \Theta\Theta\}$ and let $\{\Pi^{[\chi_j]}\}_{j=1}^{N} \subset \mathcal{C}^{[\bullet]}_{\mathrm{admissible}}$ be mutually orthogonal typed projectors:

$$\Pi^{[\chi_j]}\Pi^{[\chi_{j'}]} = \delta_{jj'}\,\Pi^{[\chi_j]}, \qquad \left(\Pi^{[\chi_j]}\right)^{\dagger} = \Pi^{[\chi_j]} \quad \text{(item (P1) of Definition 9.116)}.$$

(i) Π_{adm} *is a band-finite, BRST-closed projector.* Define $\Pi_{\mathrm{adm}} := \sum_{j=1}^{N} \Pi^{[\chi_j]}$. Orthogonality gives

$$\Pi_{\mathrm{adm}}^2 = \sum_{j,j'} \Pi^{[\chi_j]}\Pi^{[\chi_{j'}]} = \sum_{j} \Pi^{[\chi_j]} = \Pi_{\mathrm{adm}}, \qquad \Pi_{\mathrm{adm}}^{\dagger} = \sum_{j} \left(\Pi^{[\chi_j]}\right)^{\dagger} = \Pi_{\mathrm{adm}},$$

so Π_{adm} is a projector in $\mathrm{End}(\bullet)$. Each $\Pi^{[\chi_j]}$ is band-finite (item (P3) of Definition 9.116), hence the finite sum Π_{adm} is band-finite as the band class is closed under finite sums (Definition 9.96). BRST-closedness is preserved by linearity:

$$[\,\hat{Q}_{\mathrm{BRST}}, \Pi_{\mathrm{adm}}\,] = \sum_{j} [\,\hat{Q}_{\mathrm{BRST}}, \Pi^{[\chi_j]}] = 0 \quad \text{(item (P4) of Definition 9.116)}.$$

(ii) Residual projector and Σ*–orthogonality to each typed sector.* Let $1_\bullet$ denote the identity in $\mathrm{End}(\bullet)$ and set $\Pi_{\mathrm{res}} := 1_\bullet - \Pi_{\mathrm{adm}}$. Then

$$\Pi_{\mathrm{res}}^2 = (1_\bullet - \Pi_{\mathrm{adm}})^2 = 1_\bullet - 2\Pi_{\mathrm{adm}} + \Pi_{\mathrm{adm}} = 1_\bullet - \Pi_{\mathrm{adm}} = \Pi_{\mathrm{res}},$$

so Π_{res} is a projector (self-adjoint by construction). Algebraically,

$$\Pi_{\mathrm{res}}\,\Pi^{[\chi_j]} = (1_\bullet - \Pi_{\mathrm{adm}})\Pi^{[\chi_j]} = \Pi^{[\chi_j]} - \sum_{m=1}^{N} \Pi^{[\chi_m]}\Pi^{[\chi_j]} = \Pi^{[\chi_j]} - \Pi^{[\chi_j]} = 0,$$

and similarly $\Pi^{[\chi_j]}\Pi_{\mathrm{res}} = 0$. Consequently, for any timelike $k \in \mathcal{C}_x^{\hat{\Sigma},*}$ (Definition 9.104),

$$\hat{Q}_x^{[\chi_j]}[k]\ \Pi_{\mathrm{res}} = \Big(\Pi^{[\chi_j]}\hat{Q}_x[k]\Pi^{[\chi_j]}\Big)\Pi_{\mathrm{res}} = \Pi^{[\chi_j]}\hat{Q}_x[k]\underbrace{\Big(\Pi^{[\chi_j]}\Pi_{\mathrm{res}}\Big)}_{=0} = 0,$$

and likewise $\Pi_{\mathrm{res}}\hat{Q}_x^{[\chi_j]}[k] = 0$. Applying any band functional (Definition 9.96) yields

$$\mathcal{I}_\Psi\Big[\hat{Q}_x^{[\chi_j]}[k]\Pi_{\mathrm{res}}\Big] = \mathcal{I}_\Psi[0] = 0, \qquad \mathcal{I}_\Psi\Big[\Pi_{\mathrm{res}}\hat{Q}_x^{[\chi_j]}[k]\Big] = \mathcal{I}_\Psi[0] = 0,$$

which is precisely the Σ–orthogonality on the band (Definition 9.22 and Lemma 9.118).

(iii) Exclusion of the residual image if typed hyperbolicity fails. If Π_{res} fails the typed hyperbolicity condition (9.53) (Definition 9.111) for some timelike k, then by the exclusion rule (Corollary 9.113) its image cannot be admitted to the permit ledger. Equivalently, no additional χ supported in $\operatorname{Ran}\Pi_{\mathrm{res}}$ may be added to $\mathcal{C}^{[\bullet]}_{\mathrm{admissible}}$.

Combining (i)–(iii) proves the proposition. □

Corollary 9.125 (Chrono–Kymon admission)**.** *In the $\Theta\Theta$ sector, let $T := \hat{t}^{(\Theta)}_{\mu\nu}\hat{\Sigma}^{(\Theta)\,\mu\nu}$ be the Σ-weighted torsional operator (cf. Definition 5.25). The Riesz projector $\Pi^{[\mathrm{CK}]}$ onto a nonzero isolated spectral band of T defines a typed projector. If the band is separated from 0, then (9.53) holds and $\mathrm{CK} \in \mathcal{C}^{[\mathrm{kymon}]}_{\mathrm{admissible}}$.*

Proof of Corollary 9.125. Work in the $\Theta\Theta$ block. By the operator-unified chronogeometry (Definition 5.25), both $\hat{t}^{(\Theta)}_{\mu\nu}$ and $\hat{\Sigma}^{(\Theta)\,\mu\nu}$ are Hermitian, band-admissible (UV/IR) and cone-gated (Definitions 9.96 and 9.106 and Proposition 9.107). Hence the Σ-weighted torsional operator

$$T \;:=\; \hat{t}^{(\Theta)}_{\mu\nu}\,\hat{\Sigma}^{(\Theta)\,\mu\nu}$$

is a self-adjoint, band-admissible operator on the $\Theta\Theta$ sector (ghost number 0 and BRST-closed by construction).

Step 1: Typed projector from a Riesz construction. Let $\sigma(T)$ be the spectrum of T and let Γ encircle a nonzero *isolated* spectral band $\sigma_\Gamma \subset \sigma(T)$ disjoint from the rest of the spectrum. The Riesz projector $\Pi^{[\mathrm{CK}]} \equiv \Pi_\Gamma := \frac{1}{2\pi i}\oint_\Gamma (z\mathbf{1} - T)^{-1}\mathrm{d}z$ is band-finite, BRST-closed, and graded-covariant; moreover it lies in $\mathrm{End}(\Theta\Theta)$ (block internality). All these properties follow from the functional-calculus argument in Proposition 9.119 (applied to the normal, band-admissible operator T), together with Definition 9.116 (items (P1), (P3), (P4), (P5), (P7)). By the sector semantics of Definition 9.116 (Chrono–Kymon channel in the $\Theta\Theta$ block), $\Pi^{[\mathrm{CK}]}$ is a *typed projector* in the sense used throughout.

Step 2: Hyperbolicity via separation from 0. Assume now that the band σ_Γ is separated from 0: $\delta_\Gamma := \mathrm{dist}\big(0, \sigma_\Gamma\big) > 0$. Define the positive compact operator $K_\bullet := |T| = (T^2)^{1/2}$ on the

same block. Since Γ does not intersect 0, the spectral subspace of $|T|$ associated with $|\sigma_\Gamma|$ coincides with $\operatorname{Ran}\Pi_\Gamma$; thus Π_Γ is also the Riesz projector for $K_\bullet$ onto the corresponding positive cluster. Applying Corollary 9.120 with this $K_\bullet$ gives the equivalence

$$\delta_\Gamma > 0 \quad\Longleftrightarrow\quad \inf_{\Psi\in\mathcal{S}_{\mathrm{band}}} \mathcal{I}_\Psi\Big[\hat{Q}_x^{[\mathrm{CK}]}[k]\Big] \;>\; 0 \quad \text{for all timelike } k \in \mathcal{C}_x^{\hat{\Sigma},*} \text{ (Definition 9.104)},$$

i.e. the typed hyperbolicity condition (9.53) (Definition 9.111) holds for the Chrono–Kymon projector.

Step 3: Admission to the ledger. Once typed hyperbolicity holds and items (P1)–(P7) of Definition 9.116 are satisfied (Step 1), the admission criterion of the permit ledger (Proposition 9.102) yields $\mathrm{CK} \in \mathcal{C}^{[\mathrm{kymon}]}_{\mathrm{admissible}}$, with the cone/hyperbolicity certificate recorded as in Corollary 9.120 (admissible timelike interval, gap margin δ_Γ, and stability radius).

Combining Steps 1–3 proves the corollary. □

Remark 9.126 (Interface with typed contrasts and visibilities)**.** Once $\Pi^{[\chi]}$ is admitted, all typed diagnostics in §3—namely $\Delta^{\mathrm{pre},\Sigma;[\chi]}_{\mathrm{law/state}}$, $\Lambda^{\Sigma;[\chi]}_{\mathrm{law/state}}$ and the quadrant $\mathrm{Quad}^{[\chi]}$—are computed by restricting the kernels and probes with $\Pi^{[\chi]}$ (cf. Definitions 9.28, 9.45, 9.47 and 9.109). Band retardedness and microcausality follow from Propositions 9.107 and 9.108.

9.11 Typed Σ–Contrasts & Visibilities (Pre-Physical)

All evaluations below use *retarded, cone-gated* kernels (Definition 9.106 and Proposition 9.107) and BRST-closed taggers (Definition 9.96 and Proposition 9.97). Typed restrictions are taken with the band-admissible projectors $\Pi^{[\chi]}$ constructed in Definition 9.116 and admitted by the ledger Definition 9.122.

3.1 Pointwise typed Σ–contrasts

Definition 9.127 (Pointwise typed Σ–contrasts)**.** Let the Σ–probe $\hat{Q}_x[v] = v^\mu v^\nu \hat{\Sigma}_{\mu\nu}(x)$ be as in Definitions 9.22 and 9.95 and let $\Pi^{[\chi]}$ be a typed projector (Definition 9.116). For a local graded gauge frame $g(x)$ (Definition 9.7) define the *typed law-contrast* and *typed state-contrast* at $(x;v)$ by

$$\Delta^{\mathrm{pre},\Sigma;[\chi]}_{\mathrm{law}}(g;x;v) := \Big(\mathcal{I}\Big[\big(\hat{Q}_x[v] - \mathrm{Ad}_{g(x)}\,\hat{Q}_x[v]\big)\Big|_{\Pi^{[\chi]}}\Big] \Big)^{1/2}, \tag{9.58}$$

$$\Delta^{\mathrm{pre},\Sigma;[\chi]}_{\mathrm{state}}(g;x;v) := \Big| \mathcal{I}_\Psi\Big[\hat{Q}_x[v]\Big|_{\Pi^{[\chi]}}\Big] - \mathcal{I}_{g\cdot\Psi}\Big[\hat{Q}_x[v]\Big|_{\Pi^{[\chi]}}\Big] \Big|^{1/2}, \tag{9.59}$$

where $\mathcal{I}, \mathcal{I}_\Psi$ are band functionals as in Definition 9.45. The restriction $\big|_{\Pi^{[\chi]}}$ abbreviates $\Pi^{[\chi]}(\cdot)\Pi^{[\chi]}$ (Definition 9.109).

Lemma 9.128 (BRST/Nielsen safety of pointwise typed contrasts)**.** *Under a gauge-parameter variation for which $\delta_\theta \hat{Q}_x[v] = [\hat{Q}_{\mathrm{BRST}}, X_\theta]$, the quantities in (9.58)–(9.59) are invariant: $\partial_\theta \Delta^{\mathrm{pre},\Sigma;[\chi]}_\bullet(g;x;v) = 0$ for $\bullet \in \{\mathrm{law},\mathrm{state}\}$, by Proposition 9.97 and $[\hat{Q}_{\mathrm{BRST}}, \Pi^{[\chi]}] = 0$ (Definition 9.116).*

Proof of Lemma 9.128. Recall the pointwise typed contrasts (9.58)–(9.59) from Definition 9.127, with typed restriction $A\big|_{\Pi^{[\chi]}} := \Pi^{[\chi]} A\, \Pi^{[\chi]}$ and BRST–closed projector $[\,\hat{Q}_{\mathrm{BRST}}, \Pi^{[\chi]}] = 0$ (Definition 9.116). Band functionals $\mathcal{I}, \mathcal{I}_\Psi$ annihilate BRST–exact insertions (Proposition 9.97) and act at ghost number 0 (Definition 9.96).

Step 1: Typed variation of the Σ–probe is BRST–exact. By hypothesis, $\delta_\theta \hat{Q}_x[v] = [\,\hat{Q}_{\mathrm{BRST}}, X_\theta\,]$. Using $[\,\hat{Q}_{\mathrm{BRST}}, \Pi^{[\chi]}] = 0$ we obtain

$$\delta_\theta\Big(\hat{Q}_x[v]\Big|_{\Pi^{[\chi]}}\Big) = \Pi^{[\chi]}[\,\hat{Q}_{\mathrm{BRST}}, X_\theta\,]\Pi^{[\chi]} = \Big[\,\hat{Q}_{\mathrm{BRST}},\, \Pi^{[\chi]} X_\theta \Pi^{[\chi]}\,\Big]. \tag{9.60}$$

Thus the typed variation is itself BRST–exact.

Step 2: Invariance of the pointwise typed law contrast. Let $A_\theta(x; v) := \hat{Q}_x[v] - \mathrm{Ad}_{g_\theta(x)}\, \hat{Q}_x[v]$ be the (untyped) law–contrast operand (Definition 9.127). A gauge–parameter shift changes A_θ by a BRST–exact commutator on the band algebra (graded covariance, Definition 9.7; cf. the mechanism used in Proposition 9.91):

$$\delta_\theta A_\theta(x; v) = \Big[\,\hat{Q}_{\mathrm{BRST}}, Y_\theta(x; v)\,\Big] \quad \text{for some band–admissible } Y_\theta.$$

Restricting to the typed subspace and using (9.60),

$$\delta_\theta\Big(A_\theta\Big|_{\Pi^{[\chi]}}\Big) = \Big[\,\hat{Q}_{\mathrm{BRST}},\, \Pi^{[\chi]} Y_\theta \Pi^{[\chi]}\,\Big].$$

Applying $\mathcal{I}$ and Proposition 9.97 gives $\partial_\theta\, \mathcal{I}\Big(A_\theta\Big|_{\Pi^{[\chi]}}\Big) = 0$. Since $\Delta_{\mathrm{law}}^{\mathrm{pre},\Sigma;[\chi]}(g; x; v)$ is the nonnegative square root of this scalar (Definition 9.127), the chain rule yields

$$\partial_\theta\, \Delta_{\mathrm{law}}^{\mathrm{pre},\Sigma;[\chi]}(g; x; v) = 0.$$

Step 3: Invariance of the pointwise typed state contrast. For the state contrast operand, write $B_\theta(x; v) := \mathcal{I}_\Psi\Big(\hat{Q}_x[v]\Big|_{\Pi^{[\chi]}}\Big) - \mathcal{I}_{g_\theta\cdot\Psi}\Big(\hat{Q}_x[v]\Big|_{\Pi^{[\chi]}}\Big)$ (Definition 9.127, Definition 9.7). Using (9.60) and the covariance rule of the band functionals (Definitions 9.7 and 9.96), each θ–variation produces BRST–exact insertions inside $\mathcal{I}_\Psi$ and $\mathcal{I}_{g_\theta\cdot\Psi}$, which are killed by Proposition 9.97. Hence $\partial_\theta B_\theta(x; v) = 0$, and taking the absolute value and square root (definition (9.59)) preserves the zero derivative:

$$\partial_\theta\, \Delta_{\mathrm{state}}^{\mathrm{pre},\Sigma;[\chi]}(g; x; v) = 0.$$

Combining Steps 2–3 proves the claimed invariance for $\bullet \in \{\mathrm{law}, \mathrm{state}\}$. □

Lemma 9.129 (Cone causality & positivity)**.** *If v is chosen with $k := k(v) \in \mathcal{C}_x^{\hat{\Sigma},*}$ timelike (Definition 9.104) and kernels are retarded cone-gated (Definition 9.106), then $\mathcal{I}_\Psi\Big[\hat{Q}_x[v]\Big|_{\Pi^{[\chi]}}\Big] \geq 0$ and the commutators entering (9.58)–(9.59) vanish for spacelike separations (Proposition 9.108). Thus both $\Delta_{\mathrm{law}}^{\mathrm{pre},\Sigma;[\chi]}$ and $\Delta_{\mathrm{state}}^{\mathrm{pre},\Sigma;[\chi]}$ are well-defined and real.*

Proof of Lemma 9.129. Fix x and choose v so that $k := k(v) \in \mathcal{C}_x^{\hat{\Sigma},*}$ is timelike (Definition 9.104). Recall the Σ–probe and its typed restriction

$$\hat{Q}_x[v] = v^\mu v^\nu \hat{\Sigma}_{\mu\nu}(x) \quad (\textit{Definition 9.95}),$$
$$\hat{Q}_x^{[\chi]}[v] := \Pi^{[\chi]}\, \hat{Q}_x[v]\, \Pi^{[\chi]} \quad (\text{cf. (9.52), Definition 9.109}).$$

(a) Positivity of the typed quadratic evaluation. By cone-compatibility of typed projectors (item (P6) of Definition 9.116) the principal quadratic form restricted by $\Pi^{[\chi]}$ is nonnegative on cone–timelike covectors. Equivalently, for all band functionals $\mathcal{I}_\Psi$ (Definition 9.96),

$$\mathcal{I}_\Psi\big[\hat{Q}_x^{[\chi]}[k]\big] \;\geq\; 0. \tag{9.61}$$

(If χ is admitted to the ledger, (9.61) strengthens to a strict lower bound by typed hyperbolicity, Definition 9.111.) Moreover, $\hat{Q}_x^{[\chi]}[k]$ is Hermitian (Lemma 9.110), hence the scalar in (9.61) is *real*.

(b) Cone-gated retardedness and microcausality. All kernels are replaced by their retarded, cone-gated versions $\mathcal{K}^{\mathrm{ret}} = \mathbf{1}_{J^+(\mathcal{C}^{\hat{\Sigma},*})}\mathcal{K}$ (Definition 9.106), which are supported in $J^+(\mathcal{C}^{\hat{\Sigma},*})$ (Proposition 9.107). Therefore the Pauli–Jordan bi-distributions of the unified field $\hat{\mathbf{F}} = (\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}})^{\mathsf{T}}$ have support contained in $J^+ \cup J^-$ with respect to the Σ–cone. Consequently, for spacelike separated supports (w.r.t. $\mathcal{C}^{\hat{\Sigma},*}$) one has

$$[\,\hat{\mathbf{\Phi}}(f), \hat{\mathbf{\Phi}}(g)\,] = [\,\hat{\mathbf{\Theta}}(f), \hat{\mathbf{\Theta}}(g)\,] = 0,$$

by microcausality from cone-gating (Proposition 9.108).

(c) Well-definedness and reality of the typed pointwise contrasts. For the *state* contrast, the definition $\Delta^{\mathrm{pre},\Sigma;[\chi]}_{\mathrm{state}}(g;x;v) = \big|\mathcal{I}_\Psi[\hat{Q}_x[v]\big|_{\Pi^{[\chi]}}] - \mathcal{I}_{g\cdot\Psi}[\hat{Q}_x[v]\big|_{\Pi^{[\chi]}}]\big|^{1/2}$ (Definition 9.127, eq. (9.59)) uses an absolute value and the arguments are real by Hermiticity (Lemma 9.110), so the square root is real and well-defined.

For the *law* contrast,

$$\Delta^{\mathrm{pre},\Sigma;[\chi]}_{\mathrm{law}}(g;x;v) = \Big(\mathcal{I}\big[(\hat{Q}_x[v] - \mathrm{Ad}_{g(x)}\,\hat{Q}_x[v])\big|_{\Pi^{[\chi]}}\big]\Big)^{1/2} \quad \text{(Definition 9.127, eq. (9.58))},$$

the operand is Hermitian and evaluated by a positive band functional (Definitions 9.45 and 9.96), and the cone-gated retardedness ensures causal support (Proposition 9.107). Thus the inner scalar is real and nonnegative, so the square root is well-defined and real. In both cases, microcausality from (b) guarantees the vanishing of commutators entering the contrasts for spacelike separations (Proposition 9.108).

Combining (a)–(c) establishes the stated nonnegativity $\mathcal{I}_\Psi[\hat{Q}_x[v]\big|_{\Pi^{[\chi]}}] \geq 0$, the spacelike vanishing of commutators, and the well-defined, real-valued nature of $\Delta^{\mathrm{pre},\Sigma;[\chi]}_{\mathrm{law}}$ and $\Delta^{\mathrm{pre},\Sigma;[\chi]}_{\mathrm{state}}$. □

Proposition 9.130 (Gauge covariance and frame independence)**.** *Let g and $\tilde{g}$ be related by a band-admissible graded gauge transformation. Then the contrasts in (9.58)–(9.59) coincide. In particular, any change $g \mapsto g'$ with*

$$\mathrm{Ad}_{g'(x)}\,\hat{Q}_x[v] = \mathrm{Ad}_{g(x)}\,\hat{Q}_x[v] + [\,\hat{Q}_{\mathrm{BRST}}, Y\,]$$

leaves the contrasts invariant by Lemma 9.128.

Proof of Proposition 9.130. Let $\tilde{g} = h\,g$ with h a band-admissible graded-gauge element (Definitions 9.7 and 9.96). Consider a smooth path $\theta \mapsto h_\theta$ in the graded-gauge group with $h_0 = \mathbf{1}$ and $h_1 = h$, and set $g_\theta := h_\theta g$. Define the pointwise typed law/state contrasts $\Delta_\bullet^{\mathrm{pre},\Sigma;[\chi]}(g_\theta; x; v)$ by Definition 9.127, i.e. via (9.58)–(9.59) with the typed restriction $A\big|_{\Pi^{[\chi]}} = \Pi^{[\chi]} A\, \Pi^{[\chi]}$ and $[\,\hat{Q}_{\mathrm{BRST}}, \Pi^{[\chi]}] = 0$ (Definition 9.116).

Step 1: Infinitesimal gauge change is BRST-exact on the band. By graded covariance and BRST hygiene (Definitions 9.7 and 9.96), the θ–variation of the Σ–probe under g_θ is BRST-exact on the band algebra:

$$\partial_\theta\Big(\,\mathrm{Ad}_{g_\theta(x)}\,\hat{Q}_x[v]\Big) = \Big[\,\hat{Q}_{\mathrm{BRST}},\; X_\theta(x;v)\,\Big] \qquad \text{for some band-admissible } X_\theta, \tag{9.62}$$

cf. the gauge-parameter mechanism used in Proposition 9.91.

Step 2: Pointwise invariance along the path. Applying Lemma 9.128 to (9.62) and using $[\,\hat{Q}_{\mathrm{BRST}}, \Pi^{[\chi]}] = 0$ (Definition 9.116) yields

$$\partial_\theta \Delta_\bullet^{\mathrm{pre},\Sigma;[\chi]}(g_\theta; x; v) = 0, \qquad \bullet \in \{\mathrm{law}, \mathrm{state}\}.$$

Integrating from $\theta = 0$ to $\theta = 1$ gives

$$\Delta_\bullet^{\mathrm{pre},\Sigma;[\chi]}(\tilde{g}; x; v) = \Delta_\bullet^{\mathrm{pre},\Sigma;[\chi]}(g; x; v) \quad \text{for } \bullet \in \{\mathrm{law}, \mathrm{state}\},$$

i.e. the contrasts coincide for g and $\tilde{g}$.

Step 3: The stated special case. If a change $g \mapsto g'$ satisfies $\mathrm{Ad}_{g'(x)}\,\hat{Q}_x[v] = \mathrm{Ad}_{g(x)}\,\hat{Q}_x[v] + [\,\hat{Q}_{\mathrm{BRST}}, Y\,]$, then the contrasts are invariant by a one-step application of Lemma 9.128 (no path integration needed), since the typed variation is BRST-exact and is annihilated by the band functionals (Proposition 9.97).

Combining Steps 1–3 proves gauge covariance and frame independence of the pointwise typed contrasts (9.58)–(9.59). □

3.2 Aggregation and normalization (typed Σ–visibilities)

Definition 9.131 (Typed aggregation on a probe family)**.** Fix a band-admissible probe family $\mathcal{V} = \{v_k\}_{k=1}^m$ and convex weights $w_k > 0$, $\sum_k w_k = 1$. Define the aggregated typed contrasts

$$\Delta_\bullet^{\mathrm{pre},\Sigma;[\chi]}(x; \mathcal{V}, w) := \Big(\,\sum_{k=1}^{m} w_k\,\Big(\Delta_\bullet^{\mathrm{pre},\Sigma;[\chi]}(g; x; v_k)\Big)^2\,\Big)^{1/2}, \quad \bullet \in \{\mathrm{law}, \mathrm{state}\}. \tag{9.63}$$

Lemma 9.132 (Monotonicity and stability of aggregation)**.** *If $\mathcal{V} \subseteq \mathcal{V}'$ and w' extends w by assigning nonnegative weights to new probes, then $\Delta_\bullet^{\mathrm{pre},\Sigma;[\chi]}(x; \mathcal{V}, w) \le \Delta_\bullet^{\mathrm{pre},\Sigma;[\chi]}(x; \mathcal{V}', w')$. Moreover, under retunings $(\lambda_{\mathrm{UV}}, \ell) \mapsto (\lambda'_{\mathrm{UV}}, \ell')$ within the stability radius of Definition 9.116, the map $(\lambda_{\mathrm{UV}}, \ell) \mapsto \Delta_\bullet^{\mathrm{pre},\Sigma;[\chi]}(x; \mathcal{V}, w)$ is continuous.*

Proof of Lemma 9.132. Recall the pointwise typed contrasts from Definition 9.127,

$$\Delta_{\mathrm{law}}^{\mathrm{pre},\Sigma;[\chi]}(g; x; v) := \Big(\,\mathcal{I}\Big[(\hat{Q}_x[v] - \mathrm{Ad}_{g(x)}\,\hat{Q}_x[v])\Big|_{\Pi^{[\chi]}}\Big]\Big)^{1/2}, \tag{9.64}$$

$$\Delta_{\mathrm{state}}^{\mathrm{pre},\Sigma;[\chi]}(g; x; v) := \Big|\;\mathcal{I}_\Psi\Big[\hat{Q}_x[v]\Big|_{\Pi^{[\chi]}}\Big] - \mathcal{I}_{g\cdot\Psi}\Big[\hat{Q}_x[v]\Big|_{\Pi^{[\chi]}}\Big]\;\Big|^{1/2}, \tag{9.65}$$

and the aggregation/normalization step producing the (pre-normalized) typed Σ–aggregates $\Delta_\bullet^{\mathrm{pre},\Sigma;[\chi]}(x;\mathcal{V},w)$ from a probe set $\mathcal{V}=\{v_k\}$ and weights $w=\{w_k\}_{v_k\in\mathcal{V}}$, as specified in Definition 9.133. In particular, the admissible aggregators include the (weighted) root-sum-square (RSS),

$$\Delta_\bullet^{\mathrm{pre},\Sigma;[\chi]}(x;\mathcal{V},w)=\Big(\sum_{v_k\in\mathcal{V}} w_k\,\Delta_\bullet^{\mathrm{pre},\Sigma;[\chi]}(g;x;v_k)^2\Big)^{1/2},\qquad w_k\ge 0, \tag{9.66}$$

and, more generally, any norm-like, coordinatewise-monotone aggregator registered in Definition 9.133 (cf. Definition 9.47 when available).

Monotonicity under $\mathcal{V}\subseteq\mathcal{V}'$ and $w'\supset w$ with nonnegative new weights. By Lemma 9.129, each pointwise contrast $\Delta_\bullet^{\mathrm{pre},\Sigma;[\chi]}(g;x;v)$ is real and nonnegative (the typed quadratic evaluation is nonnegative on cone–timelike covectors, and the square roots in (9.64)–(9.65) ensure ≥ 0). Let $\mathcal{V}'=\mathcal{V}\cup\mathcal{W}$ and let w' extend w by assigning nonnegative weights $\{w'_j\}_{v_j\in\mathcal{W}}\subset[0,\infty)$. For the RSS aggregator (9.66),

$$\begin{aligned}\Delta_\bullet^{\mathrm{pre},\Sigma;[\chi]}(x;\mathcal{V}',w')^2&=\sum_{v_k\in\mathcal{V}} w_k\,\Delta_\bullet(v_k)^2+\sum_{v_j\in\mathcal{W}} w'_j\,\Delta_\bullet(v_j)^2\\&\ge\sum_{v_k\in\mathcal{V}} w_k\,\Delta_\bullet(v_k)^2\\&=\Delta_\bullet^{\mathrm{pre},\Sigma;[\chi]}(x;\mathcal{V},w)^2,\end{aligned}$$

hence (taking square roots) the required inequality follows:

$$\Delta_\bullet^{\mathrm{pre},\Sigma;[\chi]}(x;\mathcal{V},w)\ \le\ \Delta_\bullet^{\mathrm{pre},\Sigma;[\chi]}(x;\mathcal{V}',w').$$

The same argument applies to any admissible aggregator in Definition 9.133 that is coordinatewise nondecreasing on $\mathbb{R}_{\ge 0}^{|\mathcal{V}|}$ and nondecreasing under appending nonnegative coordinates; these properties are part of the aggregation hygiene adopted in Definition 9.133. This proves monotonicity for both $\bullet\in\{\mathrm{law},\mathrm{state}\}$.

Continuity under band retunings within the stability radius. Consider a retuning $(\lambda_{\mathrm{UV}},\ell)\mapsto(\lambda'_{\mathrm{UV}},\ell')$ that lies inside the stability radius of item (P7) in Definition 9.116. Then (i) the typed projector $\Pi^{[\chi]}$ varies continuously in operator norm (Riesz projector stability: Proposition 9.119), (ii) the UV/IR taggers vary continuously in operator norm within the band class (Definition 9.96; IR monotonicity in Lemma 9.105), and (iii) the cone gate and retarded kernels commute with the instrumentation and remain cone-supported (Proposition 9.107 and Definition 9.106). Therefore, for each fixed probe $v\in\mathcal{V}$, the typed operands in (9.64)–(9.65) depend *continuously* on $(\lambda_{\mathrm{UV}},\ell)$ in the operator norm, and applying the bounded linear band functionals $\mathcal{I},\mathcal{I}_\Psi$ (Definition 9.96) yields continuous scalar functions of the retuning parameters. The outer operations (absolute value, positive square root) preserve continuity on $\mathbb{R}_{\ge 0}$, hence each pointwise contrast $\Delta_\bullet^{\mathrm{pre},\Sigma;[\chi]}(g;x;v)$ is continuous in $(\lambda_{\mathrm{UV}},\ell)$. Finally, $\Delta_\bullet^{\mathrm{pre},\Sigma;[\chi]}(x;\mathcal{V},w)$ is obtained from finitely many continuous nonnegative inputs via a continuous aggregator (e.g. the continuous map (9.66)); thus the aggregate is continuous in $(\lambda_{\mathrm{UV}},\ell)$.

Both claims (monotonicity and continuity) are established for $\bullet\in\{\mathrm{law},\mathrm{state}\}$, completing the proof. □

Definition 9.133 (Typed Σ–visibilities)**.** Let $\mathrm{Vis}(\cdot\,;\cdot)$ denote the visibility normalizer of Definition 9.47 and eqs. (9.21) and (9.22) (monotone-decreasing in its first argument, range $[0,1]$, equals 1 at zero contrast). Define

$$\Lambda_{\bullet}^{\Sigma;[\chi]}(x) := \mathrm{Vis}\Big(\Delta_{\bullet}^{\mathrm{pre},\Sigma;[\chi]}(x;\mathcal{V},w) \;;\; \mathcal{N}_{\bullet}^{[\chi]}(x;\mathcal{V},w) \Big) \in [0,1], \quad \bullet \in \{\mathrm{law},\mathrm{state}\}, \tag{9.67}$$

where $\mathcal{N}_{\bullet}^{[\chi]}$ is the sector/typed normalizer specified in Definition 9.47. The *typed law-deficit* is

$$d_{\mathrm{law}}^{\Sigma;[\chi]}(x) := 1 - \Big|\Lambda_{\mathrm{law}}^{\Sigma;[\chi]}(x)\Big|. \tag{9.68}$$

Proposition 9.134 (Invariance and bounds)**.** *The visibilities* (9.67) *are:*

(a) Gauge/BRST-invariant *by Lemma 9.128 and Proposition 9.97.*

(b) Causally well-posed *(retarded, cone-gated) by Proposition 9.107.*

(c) Normalized *to* $[0,1]$ *with* $\Lambda_{\bullet}^{\Sigma;[\chi]}(x) = 1$ *iff* $\Delta_{\bullet}^{\mathrm{pre},\Sigma;[\chi]}(x;\mathcal{V},w) = 0$ *(law/state symmetry baseline, cf. Ch. 8 summary and Definition 9.47).*

Proof of Proposition 9.134. By definition (Definition 9.133), the (typed) visibilities are obtained from the aggregated contrasts via the normalization map encoded in (9.67):

$$\Lambda_{\bullet}^{\Sigma;[\chi]}(x) \;:=\; \mathfrak{n}_{\bullet}\Big(\Delta_{\bullet}^{\mathrm{pre},\Sigma;[\chi]}(x;\mathcal{V},w)\Big), \qquad \bullet \in \{\mathrm{law},\mathrm{state}\}, \tag{9.69}$$

where $\Delta_{\bullet}^{\mathrm{pre},\Sigma;[\chi]}$ is the (nonnegative) aggregate built from the pointwise typed contrasts of Definition 9.127 [(9.58)–(9.59)] and $\mathfrak{n}_{\bullet} : \mathbb{R}_{\geq 0} \to [0,1]$ is the normalization specified in Definitions 9.47 and 9.133 (baseline $\mathfrak{n}_{\bullet}(0) = 1$, monotone nonincreasing).

(a) Gauge/BRST invariance. For each probe v, the pointwise typed contrasts are invariant under band–admissible gauge–parameter variations by Lemma 9.128, using the BRST–annihilation of band functionals (Proposition 9.97) and $[\hat{Q}_{\mathrm{BRST}}, \Pi^{[\chi]}] = 0$ (Definition 9.116). Aggregation is performed by a fixed algebraic rule on these scalars (Definition 9.133), hence $\Delta_{\bullet}^{\mathrm{pre},\Sigma;[\chi]}(x;\mathcal{V},w)$ is gauge/BRST–invariant. Applying the scalar-valued normalization (9.69) preserves invariance. Thus $\Lambda_{\bullet}^{\Sigma;[\chi]}$ is gauge/BRST–invariant.

(b) Causal well-posedness (retarded, cone-gated). All kernels entering the pointwise contrasts are replaced by their retarded, cone-gated versions (Definition 9.106), supported inside $J^{+}\big(\mathcal{C}^{\hat{\Sigma},*}\big)$ by Proposition 9.107. Consequently, the bi-distributional commutators vanish for spacelike separations (Proposition 9.108), and each pointwise contrast is well-defined and real (Lemma 9.129). Aggregation and the pointwise normalization (9.69) do not alter causal support, hence the visibilities are retarded and cone-gated.

(c) Normalization and the $[0,1]$ *range with baseline at* 1. By Lemma 9.129, the aggregate $\Delta_{\bullet}^{\mathrm{pre},\Sigma;[\chi]}(x;\mathcal{V},w) \in \mathbb{R}_{\geq 0}$, so the input to $\mathfrak{n}_{\bullet}$ lies in its domain. The normalization axioms of Definitions 9.47 and 9.133 give

$$0 \;\leq\; \Lambda_{\bullet}^{\Sigma;[\chi]}(x) \;=\; \mathfrak{n}_{\bullet}\Big(\Delta_{\bullet}^{\mathrm{pre},\Sigma;[\chi]}(x;\mathcal{V},w)\Big) \;\leq\; 1, \qquad \mathfrak{n}_{\bullet}(0) = 1.$$

Moreover, the baseline property stipulated in Definition 9.47 ensures that $\mathfrak{n}_\bullet(z) = 1$ iff $z = 0$ (law/state symmetry baseline). Therefore,

$$\Lambda_\bullet^{\Sigma;[\chi]}(x) = 1 \iff \Delta_\bullet^{\mathrm{pre},\Sigma;[\chi]}(x;\mathcal{V},w) = 0,$$

as claimed.

Combining (a)–(c) proves the proposition. □

Corollary 9.135 (Orthogonal additivity)**.** *For a set of mutually Σ-orthogonal types $\{\chi_j\}$ (Lemma 9.118), the squared aggregated contrasts are additive:*

$$\Big(\Delta_\bullet^{\mathrm{pre},\Sigma;\,[\oplus_j \chi_j]}(x)\Big)^2 = \sum_j \Big(\Delta_\bullet^{\mathrm{pre},\Sigma;[\chi_j]}(x)\Big)^2, \quad \bullet \in \{\mathrm{law},\mathrm{state}\},$$

hence the corresponding visibilities combine monotonically under the normalizer of Definition 9.47.

Proof of Corollary 9.135. Let $\{\Pi^{[\chi_j]}\}_{j=1}^N$ be mutually Σ–orthogonal typed projectors in a fixed block (Lemma 9.118) and set $\Pi_{\mathrm{adm}} := \sum_{j=1}^N \Pi^{[\chi_j]}$ (Proposition 9.124). Write, for a fixed probe v, the Hermitian *law* and *state* pointwise operands (Definition 9.127),

$$A(x;v) := \hat{Q}_x[v] - \mathrm{Ad}_{g(x)}\,\hat{Q}_x[v], \qquad B_\Psi(x;v) := \hat{Q}_x[v] \ \text{ (with band evaluations } \mathcal{I}_\Psi,\ \mathcal{I}_{g\cdot\Psi}),$$

where $\hat{Q}_x[v] = v^\mu v^\nu \hat{\Sigma}_{\mu\nu}(x)$ (Definition 9.95).

Step 1: Pointwise additivity of squared contrasts for orthogonal types. Because A and $\hat{Q}_x[v]$ are linear in $\hat{\Sigma}$ and Hermitian (Lemma 9.110), and Π_{adm} is an orthogonal sum, we have the block decomposition

$$\Pi_{\mathrm{adm}}\, A\, \Pi_{\mathrm{adm}} = \sum_j \Pi^{[\chi_j]} A \Pi^{[\chi_j]} \ + \sum_{j\neq j'} \Pi^{[\chi_j]} A \Pi^{[\chi_{j'}]},$$

and the analogous decomposition with A replaced by $\hat{Q}_x[v]$. For v with $k(v) \in \mathcal{C}_x^{\hat{\Sigma},*}$ timelike (Definition 9.104) the Σ–orthogonality lemma (Lemma 9.118) implies that all *off–diagonal* band evaluations vanish:

$$\mathcal{I}\Big[\Pi^{[\chi_j]} A \Pi^{[\chi_{j'}]}\Big] = 0, \qquad \mathcal{I}_\Psi\Big[\Pi^{[\chi_j]} \hat{Q}_x[v] \Pi^{[\chi_{j'}]}\Big] = 0 \qquad (j \neq j').$$

Indeed, each off–diagonal block is a linear combination of terms of the form $\Pi^{[\chi_j]}\hat{Q}_x[v]\Pi^{[\chi_{j'}]}$, and these evaluate to zero by the band Σ–orthogonality (cf. the calculation in Lemma 9.118, Step 2). Therefore

$$\mathcal{I}\Big[(\Pi_{\mathrm{adm}} A \Pi_{\mathrm{adm}})\Big] = \sum_j \mathcal{I}\Big[\Pi^{[\chi_j]} A \Pi^{[\chi_j]}\Big], \tag{9.70}$$

$$\mathcal{I}_\Psi\Big[(\Pi_{\mathrm{adm}} \hat{Q}_x[v] \Pi_{\mathrm{adm}})\Big] = \sum_j \mathcal{I}_\Psi\Big[\Pi^{[\chi_j]} \hat{Q}_x[v] \Pi^{[\chi_j]}\Big]. \tag{9.71}$$

By Definition 9.127, the *squared* pointwise typed contrasts are precisely the right-hand sides of (9.70)–(9.71):

$$\left(\Delta_{\mathrm{law}}^{\mathrm{pre},\Sigma;[\oplus_j \chi_j]}(g;x;v)\right)^2 = \sum_j \left(\Delta_{\mathrm{law}}^{\mathrm{pre},\Sigma;[\chi_j]}(g;x;v)\right)^2,$$
$$\left(\Delta_{\mathrm{state}}^{\mathrm{pre},\Sigma;[\oplus_j \chi_j]}(g;x;v)\right)^2 = \sum_j \left(\Delta_{\mathrm{state}}^{\mathrm{pre},\Sigma;[\chi_j]}(g;x;v)\right)^2.$$

Step 2: Aggregated additivity under RSS (and norm-like) aggregation. Aggregating over a probe set $\mathcal{V}$ with nonnegative weights w using the root-sum-square (RSS) aggregator (Definition 9.133, eq. (9.66)), the pointwise additivity lifts to the *aggregated* level by linearity of finite sums:

$$\begin{aligned}\left(\Delta_{\bullet}^{\mathrm{pre},\Sigma;\,[\oplus_j \chi_j]}(x)\right)^2 &= \sum_{v\in\mathcal{V}} w_v \left(\Delta_{\bullet}^{\mathrm{pre},\Sigma;\,[\oplus_j \chi_j]}(g;x;v)\right)^2 \\ &= \sum_j \sum_{v\in\mathcal{V}} w_v \left(\Delta_{\bullet}^{\mathrm{pre},\Sigma;\,[\chi_j]}(g;x;v)\right)^2 \\ &= \sum_j \left(\Delta_{\bullet}^{\mathrm{pre},\Sigma;\,[\chi_j]}(x)\right)^2,\end{aligned}$$

for $\bullet \in \{\mathrm{law}, \mathrm{state}\}$. (The same conclusion holds for any norm-like, coordinatewise-monotone aggregator from Definition 9.133 that coincides with the ℓ^2-norm on orthogonal collections.)

Step 3: Monotonicity for visibilities. Visibilities are obtained by a nonincreasing normalizer $\mathfrak{n}_\bullet : \mathbb{R}_{\geq 0} \to [0,1]$ (Definitions 9.47 and 9.133; see also Proposition 9.134). Since $\Delta_{\bullet}^{\mathrm{pre},\Sigma;\,[\oplus_j \chi_j]}(x) = \left(\sum_j \Delta_{\bullet}^{\mathrm{pre},\Sigma;\,[\chi_j]}(x)^2\right)^{1/2} \geq \max_j \Delta_{\bullet}^{\mathrm{pre},\Sigma;\,[\chi_j]}(x)$, it follows that

$$\Lambda_{\bullet}^{\Sigma;\,[\oplus_j \chi_j]}(x) = \mathfrak{n}_\bullet\!\left(\Delta_{\bullet}^{\mathrm{pre},\Sigma;\,[\oplus_j \chi_j]}(x)\right) \;\leq\; \mathfrak{n}_\bullet\!\left(\Delta_{\bullet}^{\mathrm{pre},\Sigma;\,[\chi_j]}(x)\right) = \Lambda_{\bullet}^{\Sigma;\,[\chi_j]}(x) \qquad \forall j,$$

i.e. the direct-sum visibility combines *monotonically* under the prescribed normalizer.

Combining Steps 1–3 proves the asserted orthogonal additivity of squared aggregated contrasts and the monotone combination of the corresponding visibilities. □

Lemma 9.136 (Cone-shrink monotonicity for visibilities)**.** *If the IR window is tightened as in Lemma 9.105, then for any fixed $(\mathcal{V}, w)$ one has*

$$\Lambda_{\bullet}^{\Sigma;[\chi]}(x)\Big|_{\ell'} \;\geq\; \Lambda_{\bullet}^{\Sigma;[\chi]}(x)\Big|_{\ell}, \qquad \ell' \leq \ell,$$

i.e. reducing admissible support cannot increase *the measured breaking (monotonicity of* Vis *and of* (9.63)*).*

Proof of Lemma 9.136. Fix x, a permitted type χ, and an admissible probe set $(\mathcal{V}, w)$ (with nonnegative weights) as in Definition 9.133. Let the IR window be tightened from ℓ to ℓ'

with $\ell' \le \ell$. By Lemma 9.105 the Σ–covector cone contracts, $\mathcal{C}_x^{\hat{\Sigma},*}[\ell'] \subseteq \mathcal{C}_x^{\hat{\Sigma},*}[\ell]$, and, at the level of kernels, the *cone gate* (retarded projector) satisfies the Löwner-order relation

$$0 \;\le\; \Gamma_{\ell'} \;\le\; \Gamma_{\ell} \;\le\; \mathbf{1}, \qquad \Gamma_\bullet \text{ as in Definition 9.106, supported in } J^+\Big(\mathcal{C}^{\hat{\Sigma},*}[\bullet]\Big), \tag{9.72}$$

with retardedness preserved by Proposition 9.107.

Step 1: Pointwise contraction of typed operands under the cone gate. For any probe v and typed restriction by $\Pi^{[\chi]}$ (Definition 9.116), the pointwise law/state operands of Definition 9.127 are (schematically)

$$\mathcal{A}_\ell(v) := \Pi^{[\chi]}\,\Gamma_\ell\Big(\hat{Q}_x[v] - \mathrm{Ad}_{g(x)}\,\hat{Q}_x[v]\Big)\Gamma_\ell\,\Pi^{[\chi]}, \qquad \mathcal{B}_{\Psi,\ell}(v) := \Pi^{[\chi]}\,\Gamma_\ell\,\hat{Q}_x[v]\,\Gamma_\ell\,\Pi^{[\chi]},$$

with $\hat{Q}_x[v] = v^\mu v^\nu \hat{\Sigma}_{\mu\nu}(x)$ (Definition 9.95). The replacement $\ell \mapsto \ell'$ inserts the smaller gate $\Gamma_{\ell'}$ in place of Γ_ℓ.

Step 2: Monotonicity of pointwise contrasts under gate shrinking. By (9.72) and the positivity/Hermiticity properties (Lemmas 9.110 and 9.129), the band readouts used in the pointwise contrasts,

$$\Delta^{\mathrm{pre},\Sigma;[\chi]}_{\mathrm{law}}(g;x;v) = \Big(\mathcal{I}\big[\mathcal{A}_\ell(v)\big]\Big)^{1/2},$$
$$\Delta^{\mathrm{pre},\Sigma;[\chi]}_{\mathrm{state}}(g;x;v) = \Big|\mathcal{I}_\Psi\big[\mathcal{B}_{\Psi,\ell}(v)\big] - \mathcal{I}_{g\cdot\Psi}\big[\mathcal{B}_{g\cdot\Psi,\ell}(v)\big]\Big|^{1/2} \quad \text{(Definition 9.127)},$$

are evaluated by positive, ghost-number-0 band functionals (Definitions 9.45 and 9.96). Because the cone gate enters *bilinearly* and as a positive contraction (Löwner order (9.72)), each scalar inside the square roots is *monotone nonincreasing* under $\Gamma_\ell \mapsto \Gamma_{\ell'}$.[2] Therefore, for every fixed probe v,

$$\Delta^{\mathrm{pre},\Sigma;[\chi]}_{\bullet}(g;x;v)\Big|_{\ell'} \;\le\; \Delta^{\mathrm{pre},\Sigma;[\chi]}_{\bullet}(g;x;v)\Big|_{\ell}, \qquad \bullet \in \{\mathrm{law},\mathrm{state}\}. \tag{9.73}$$

Step 3: Aggregation preserves the inequality. Aggregating over $(\mathcal{V}, w)$ with any admissible (coordinatewise nondecreasing) rule (e.g. weighted RSS, Definition 9.133, eq. (9.66)) and using (9.73) pointwise yields

$$\Delta^{\mathrm{pre},\Sigma;[\chi]}_{\bullet}(x;\mathcal{V},w)\Big|_{\ell'} \;\le\; \Delta^{\mathrm{pre},\Sigma;[\chi]}_{\bullet}(x;\mathcal{V},w)\Big|_{\ell}, \qquad \bullet \in \{\mathrm{law},\mathrm{state}\}. \tag{9.74}$$

Step 4: Normalization flips the inequality for visibilities. Visibilities are obtained by the nonincreasing normalizer $\mathfrak{n}_\bullet : \mathbb{R}_{\ge 0} \to [0,1]$ (Definitions 9.47 and 9.133),

$$\Lambda^{\Sigma;[\chi]}_{\bullet}(x) \;=\; \mathfrak{n}_\bullet\Big(\Delta^{\mathrm{pre},\Sigma;[\chi]}_{\bullet}(x;\mathcal{V},w)\Big) \quad \text{(eq. (9.67))}.$$

[2] This is the standard C*-algebraic monotonicity: if $0 \le P' \le P \le \mathbf{1}$ and H is Hermitian, then for any positive linear functional φ, $\varphi\big[(P'^{1/2} H P'^{1/2})_\pm\big] \le \varphi\big[(P^{1/2} H P^{1/2})_\pm\big]$, and in particular for the positive inputs used here (cf. Definition 9.45 and Lemma 9.129) the scalar readouts decrease when the gate shrinks.

Applying $\mathfrak{n}_\bullet$ to (9.74) gives

$$\Lambda_\bullet^{\Sigma;[\chi]}(x)\Big|_{\ell'} \;\geq\; \Lambda_\bullet^{\Sigma;[\chi]}(x)\Big|_{\ell}, \qquad \bullet \in \{\text{law}, \text{state}\}.$$

Thus tightening the IR window (shrinking the cone) cannot *increase* the measured breaking; equivalently, the visibilities are monotone *nondecreasing* under $\ell \downarrow$, as stated. □

3.3 Typed pre-physical Quadrant predicate

Definition 9.137 (Typed pre-physical Quadrant predicate)**.** With the visibilities of Definition 9.133, define the quadrant assignment

$$\mathrm{Quad}^{[\chi]}(x) := \mathrm{Quad}\Big(\Lambda_{\text{law}}^{\Sigma;[\chi]}(x),\ \Lambda_{\text{state}}^{\Sigma;[\chi]}(x)\Big) \in \{\mathrm{QI}, \mathrm{QII}, \mathrm{QIII}, \mathrm{QIV}\}, \tag{9.75}$$

using the canonical decision rule of Definition 7.685. Optional triadic attribution (corr/-vac/spec) is read via the decomposition of Definition 5.54, restricted by $\Pi^{[\chi]}$.

Proposition 9.138 (Typed ⇒ sector/untyped consistency)**.** *If $\Pi^{[\chi]}$ is chosen as the identity on a block ($\Phi\Phi$ or $\Theta\Theta$), then the typed constructions reduce to the sector-level ones of Definitions 9.28, 9.45 and 9.47 and the quadrant rule (9.75) coincides with Definition 7.685.*

Proof of Proposition 9.138. Fix a block $\bullet \in \{\Phi\Phi, \Theta\Theta\}$ and take the typed projector to be the block identity $\Pi^{[\chi]} = E_{\bullet\bullet} \in \mathrm{End}(\bullet)$ (item (P1) of Definition 9.116; cf. Definition 9.94).

Step 1: Typed restriction collapses to the sector operand. For any band operator A supported in the block $\bullet$,

$$A\Big|_{\Pi^{[\chi]}} = \Pi^{[\chi]} A \Pi^{[\chi]} = E_{\bullet\bullet}\, A\, E_{\bullet\bullet} = A,$$

hence the typed Σ–probe coincides with its sector version: $\hat{Q}_x^{[\chi]}[v] = \Pi^{[\chi]} \hat{Q}_x[v] \Pi^{[\chi]} = \hat{Q}_x[v]$ (Definitions 9.95 and 9.109).

Step 2: Pointwise contrasts coincide with the sector definitions. Using (9.58)–(9.59) (Definition 9.127) and the collapse in Step 1,

$$\begin{aligned}
\Delta_{\text{law}}^{\text{pre},\Sigma;[\chi]}(g;x;v) &= \Big(\mathcal{I}\big[(\hat{Q}_x[v] - \mathrm{Ad}_{g(x)}\, \hat{Q}_x[v])\big]\Big)^{1/2} \\
&= \Delta_{\text{law}}^{\text{pre},\Sigma}(g;x;v) \quad \text{(sector version in Definition 9.45)},
\end{aligned}$$

$$\begin{aligned}
\Delta_{\text{state}}^{\text{pre},\Sigma;[\chi]}(g;x;v) &= \Big|\mathcal{I}_\Psi[\hat{Q}_x[v]] - \mathcal{I}_{g\cdot\Psi}[\hat{Q}_x[v]]\Big|^{1/2} \\
&= \Delta_{\text{state}}^{\text{pre},\Sigma}(g;x;v) \quad \text{(sector version in Definition 9.45)}.
\end{aligned}$$

Step 3: Aggregation and normalization agree. Aggregating over $(\mathcal{V}, w)$ via the admissible rules of Definition 9.133 (e.g. weighted RSS (9.66)) gives

$$\Delta_\bullet^{\text{pre},\Sigma;[\chi]}(x;\mathcal{V},w) = \Delta_\bullet^{\text{pre},\Sigma}(x;\mathcal{V},w), \qquad \bullet \in \{\text{law}, \text{state}\}.$$

Applying the visibility normalizer $\mathfrak{n}_\bullet$ (Definitions 9.47 and 9.133) yields

$$\Lambda_\bullet^{\Sigma;[\chi]}(x) = \mathfrak{n}_\bullet\Big(\Delta_\bullet^{\mathrm{pre},\Sigma;[\chi]}(x;\mathcal{V},w)\Big) = \mathfrak{n}_\bullet\Big(\Delta_\bullet^{\mathrm{pre},\Sigma}(x;\mathcal{V},w)\Big) = \Lambda_\bullet^{\Sigma}(x),$$

i.e. the typed visibilities coincide with the sector/untyped visibilities of Definition 9.47.

Step 4: Quadrant rule coincides. The typed quadrant predicate is

$$\mathrm{Quad}^{[\chi]}(x) = \mathrm{Quad}\Big(\Lambda_{\mathrm{law}}^{\Sigma;[\chi]}(x),\, \Lambda_{\mathrm{state}}^{\Sigma;[\chi]}(x)\Big) \quad \text{(eq. (9.75), Definition 9.137)}.$$

By Step 3, $\Lambda_\bullet^{\Sigma;[\chi]}(x) = \Lambda_\bullet^{\Sigma}(x)$, so $\mathrm{Quad}^{[\chi]}(x) = \mathrm{Quad}\big(\Lambda_{\mathrm{law}}^{\Sigma}(x), \Lambda_{\mathrm{state}}^{\Sigma}(x)\big)$, which is exactly the canonical sector-level quadrant decision of Definitions 7.685 and 9.28.

Step 5: Causal/BRST properties persist. Retarded cone-gating (Definition 9.106 and Proposition 9.107) and BRST/Nielsen safety (Proposition 9.97 and Lemma 9.128) do not depend on the choice $\Pi^{[\chi]} = E_{\bullet\bullet}$; hence all causal and gauge properties match those of the sector/untyped constructions in Definitions 9.28, 9.45 and 9.47.

Combining Steps 1–4 establishes that choosing the typed projector as the block identity reduces all typed objects (contrasts, aggregation, visibilities, and quadrant predicate) to their sector-level counterparts, proving the proposition. □

Corollary 9.139 (Plateau stability of the quadrant)**.** *Within the retuning-stability radius of Definition 9.116 and under IR tightening (Lemma 9.105), the map $x \mapsto \mathrm{Quad}^{[\chi]}(x)$ is locally constant except at tolerance thresholds specified by the normalizer (Definition 9.47). In particular, quadrant membership is robust on certified plateaux (cf. Chapter 8 certification items and Proposition 9.134).*

Proof of Corollary 9.139. Fix a permitted type χ and define the visibility vector by

$$\vec{\Lambda}^{[\chi]}(x) := \big(\Lambda_{\mathrm{law}}^{\Sigma;[\chi]}(x), \Lambda_{\mathrm{state}}^{\Sigma;[\chi]}(x)\big) \in [0,1]^2,$$

so that the typed quadrant is the decision map

$$\mathrm{Quad}^{[\chi]}(x) = \mathrm{Quad}\big(\vec{\Lambda}^{[\chi]}(x)\big),$$

with decision surfaces given by the tolerance thresholds of $\mathfrak{n}_\bullet$ and the canonical rule (Definition 9.137, eq. (9.75), Definitions 7.685 and 9.47).

(i) Local constancy under retunings within the stability radius. By item (P7) of Definition 9.116 and the Riesz–projector continuity (Proposition 9.119), the typed projector $\Pi^{[\chi]}$ depends continuously on band retunings $(\lambda_{\mathrm{UV}}, \ell)$ inside its stability radius. Consequently, for each fixed x and probe set $(\mathcal{V}, w)$, the aggregated contrasts $\Delta_\bullet^{\mathrm{pre},\Sigma;[\chi]}(x;\mathcal{V},w)$ vary *continuously* with the retuning parameters (Lemma 9.132, continuity part), and so do the visibilities $\Lambda_\bullet^{\Sigma;[\chi]}(x)$ after the continuous normalization (Definition 9.133). Therefore $\vec{\Lambda}^{[\chi]}(x)$ moves continuously in $[0,1]^2$ as $(\lambda_{\mathrm{UV}}, \ell)$ vary in that neighborhood. Since $\mathrm{Quad}^{[\chi]}$ is a piecewise-constant decision on the partition of $[0,1]^2$ by the tolerance thresholds (Definitions 7.685 and 9.47), it is *locally constant* so long as $\vec{\Lambda}^{[\chi]}(x)$ does not cross a decision surface. Thus quadrant changes can occur only at those thresholds.

(ii) Monotone robustness under IR tightening. If the IR window is tightened $\ell' \le \ell$, then for any fixed $(\mathcal{V}, w)$ the aggregated contrasts are monotone nonincreasing (Lemma 9.136, Eq. (9.74)), hence the visibilities are monotone nondecreasing:

$$\Lambda_{\bullet}^{\Sigma;[\chi]}(x)\Big|_{\ell'} \;\ge\; \Lambda_{\bullet}^{\Sigma;[\chi]}(x)\Big|_{\ell}, \qquad \bullet \in \{\mathrm{law}, \mathrm{state}\}.$$

Therefore the path $\ell \downarrow$ moves $\vec{\Lambda}^{[\chi]}(x)$ monotonically in $[0,1]^2$; the quadrant $\mathrm{Quad}^{[\chi]}(x)$ can change only if this monotone path intersects a tolerance boundary of the normalizer (Definition 9.47). Absent such an intersection (the certified *plateau* regime), quadrant membership is unchanged.

(iii) Causal and gauge well-posedness along the plateau. Throughout these deformations the pointwise/aggregated quantities remain retarded and cone-gated (Proposition 9.107), real and well-defined (Lemma 9.129), and gauge/BRST-invariant (Lemma 9.128 and Propositions 9.97 and 9.134). Hence no spurious quadrant flips arise from gauge or causal artifacts.

Combining (i)–(iii) proves that within the retuning-stability radius and under IR tightening, $x \mapsto \mathrm{Quad}^{[\chi]}(x)$ is locally constant except when $\vec{\Lambda}^{[\chi]}(x)$ meets a tolerance threshold prescribed by the normalizer (Definition 9.47). This is precisely the plateau stability asserted (cf. the certification semantics summarized in Chapter 8 and Proposition 9.134). □

Remark 9.140 (Operational interface to Parasymmetric meters)**.** Given $\mathrm{Quad}^{[\chi]}$ and the visibilities in Definition 9.133, band-restricted Parasymmetric meters (Section 8.6) provide the valuations $v_{\mathrm{Synthesis}}^{(a);[\chi]}$ and $v_{\mathrm{Synthesis}}^{(LS);[\chi]}$, supplying **T–L–O** (Type–Location–Onset). The onset scales/times are then defined as in §4.2 with $d_{\mathrm{law}}^{\Sigma;[\chi]}$ from (9.68), and inherit the causal and BRST properties of Lemma 9.129 and Proposition 9.134.

9.12 Parasymmetric Meters on the Band (Typed)

We port the Parasymmetric diagnostics of §8.6 to *band-reduced*, *typed* objects by restricting all constructions with the band tag $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ (Definition 9.96) and the typed projector $\Pi^{[\chi]}$ (Definition 9.116). All kernels are retarded and cone-gated (Definition 9.106 and Proposition 9.107), and all taggers are BRST-closed with ghost number 0 (Definition 9.96 and Proposition 9.97).

4.1 Local and relational meters (typed)

Definition 9.141 (Band-reduced typed meter state)**.** Fix an anchor $a \in \{L, S\}$ and a permitted type $\chi \in \mathcal{C}_{\mathrm{admissible}}^{[\cdot]}$ (Definition 9.122). Let $\mathcal{M}^{(a)[\chi]}$ denote the meter channel obtained by the band restriction and typed compression,

$$\mathcal{M}^{(a)[\chi]} : \; \mathfrak{A}_{\mathrm{band}} \xrightarrow{\Pi^{[\chi]}\,(\cdot)\,\Pi^{[\chi]}} \mathfrak{A}_{\mathrm{band}}^{[\chi]} \xrightarrow{\text{anchor } a} \mathrm{M}_2(\mathbb{C}),$$

where $\mathfrak{A}_{\mathrm{band}}$ is the band-admissible operator algebra. The *band meter state* is the 2×2 density operator

$$\rho_{\mathrm{band}}^{(a)[\chi]} \;:=\; \mathcal{M}^{(a)[\chi]}(\mathbf{1}), \qquad \rho_{\mathrm{band}}^{(a)[\chi]} \succeq 0, \quad \mathrm{Tr}\,\rho_{\mathrm{band}}^{(a)[\chi]} = 1,$$

with local parasymmetric projectors $P_{\mathrm{para}}^{(a)}$, $P_{\mathrm{antipara}}^{(a)}$ as in §8.6.

Definition 9.142 (Parasymmetric anchor basis and projectors)**.** For each anchor $a \in \{L, S\}$, fix an orthonormal basis

$$\mathcal{B}_a = \{\ |\mathrm{Sym}\rangle_a,\ |\mathrm{Break}\rangle_a\ \}$$

in the meter space of §8.6. Define the local parasymmetric projectors by

$$P^{(a)}_{\mathrm{para}} = |\mathrm{Sym}\rangle_{a\,a}\langle\mathrm{Sym}|, \qquad P^{(a)}_{\mathrm{antipara}} = |\mathrm{Break}\rangle_{a\,a}\langle\mathrm{Break}|.$$

When the band-reduced, typed meter state $\rho^{(a)[\chi]}_{\mathrm{band}}$ (Definition 9.141) is expressed in the ordered basis $\mathcal{B}_a$, its matrix elements are $\rho^{(a)[\chi]}_{ij} = {}_a\langle b_i|\,\rho^{(a)[\chi]}_{\mathrm{band}}\,|b_j\rangle_a$ with $|b_1\rangle_a = |\mathrm{Sym}\rangle_a$ and $|b_2\rangle_a = |\mathrm{Break}\rangle_a$.

Definition 9.143 (Local typed Synthesis valuation)**.** For $a \in \{L, S\}$ and a permitted type χ,

$$v^{(a);[\chi]}_{\mathrm{Synthesis}} := \mathrm{Tr}\Big(P^{(a)}_{\mathrm{para}}\,\rho^{(a)[\chi]}_{\mathrm{band}}\Big) = {}_a\langle\mathrm{Sym}|\,\rho^{(a)[\chi]}_{\mathrm{band}}\,|\mathrm{Sym}\rangle_a = \tfrac{1}{2} + \mathrm{Re}\Big(\rho^{(a)[\chi]}_{12}\Big) \in [0,1], \quad (9.76)$$

where $\rho^{(a)[\chi]}_{12} = {}_a\langle\mathrm{Sym}|\,\rho^{(a)[\chi]}_{\mathrm{band}}\,|\mathrm{Break}\rangle_a$ is the off-diagonal entry in the ordered basis $\{|\mathrm{Sym}\rangle_a, |\mathrm{Break}\rangle_a\}$.

Definition 9.144 (Relational typed meters on LS)**.** Let $\rho^{(LS)[\chi]}_{\mathrm{band}}$ be the band-reduced, typed state on the two-anchor space. Define the *relational* valuations

$$v^{(LS);[\chi]}_{\mathrm{Synthesis}} := \mathrm{Tr}\Big(P^{(LS)}_{\mathrm{para}}\,\rho^{(LS)[\chi]}_{\mathrm{band}}\Big), \qquad v^{(LS);[\chi]}_{\mathrm{Agreement}} := \mathrm{Tr}\Big(P^{(LS)}_{\mathrm{agree}}\,\rho^{(LS)[\chi]}_{\mathrm{band}}\Big),$$

with projectors $P^{(LS)}_{\mathrm{para}}$, $P^{(LS)}_{\mathrm{agree}}$ defined in §8.6.

Lemma 9.145 (Meter well-posedness, bounds, and BRST invariance)**.** *For any typed χ and anchor choice:*

(a) $0 \le v^{(a);[\chi]}_{\mathrm{Synthesis}} \le 1$ *and* $0 \le v^{(LS);[\chi]}_{\mathrm{Synthesis}}, v^{(LS);[\chi]}_{\mathrm{Agreement}} \le 1$ *(positivity of meter states).*

(b) *Under gauge-parameter variations that are BRST-exact, all valuations are invariant, by Proposition 9.97 and* $[\hat{Q}_{\mathrm{BRST}}, \Pi^{[\chi]}] = 0$ *(Definition 9.116).*

(c) *Retardedness and cone-gating of inputs imply causal support within* $J^+(\mathcal{C}^{\hat{\Sigma},*})$ *(Propositions 9.107 and 9.108).*

Proof of Lemma 9.145. Fix a permitted type χ and work with the band-reduced, typed meters of Definitions 9.141, 9.143 and 9.144. We prove (a)–(c) in order.

(a) Bounds $0 \le v \le 1$. By Definitions 9.141, 9.143 and 9.144, the local meter is $v^{(a);[\chi]}_{\mathrm{Synthesis}} = \mathrm{Tr}\Big(P^{(a)}_{\mathrm{para}}\,\rho^{(a)[\chi]}_{\mathrm{band}}\Big)$, where $P^{(a)}_{\mathrm{para}}$ is a projector (hence $0 \le P^{(a)}_{\mathrm{para}} \le \mathbf{1}$) and $\rho^{(a)[\chi]}_{\mathrm{band}}$ is a positive, unit-trace density operator on the (two-level) para-anchor space obtained from the typed, band-reduced state. Therefore

$$0 \ \le\ \mathrm{Tr}\Big(P^{(a)}_{\mathrm{para}}\,\rho^{(a)[\chi]}_{\mathrm{band}}\Big) \ \le\ \mathrm{Tr}\Big(\mathbf{1}\,\rho^{(a)[\chi]}_{\mathrm{band}}\Big) = 1.$$

The relational meters are of the same form with positive effects $0 \le P^{(LS)}_{\bullet} \le \mathbf{1}$ acting on the LS para-space (cf. Definitions 9.141, 9.143 and 9.144), so the same argument yields

$0 \le v^{(LS);[\chi]}_{\text{Synthesis}},\, v^{(LS);[\chi]}_{\text{Agreement}} \le 1$. (Equivalently, the explicit two-level expression shown inDefinitions 9.141, 9.143 and 9.144 immediately gives the same bounds.)

(b) BRST/Nielsen invariance. All ingredients that feed the meters—typed projector $\Pi^{[\chi]}$, band taggers $P_x(\lambda_{\text{UV}}), K^{(\ell)}_{\text{IR}}$, and the cone-gated kernels—are BRST-closed of ghost number 0 (Definitions 9.96, 9.106 and 9.116). Hence a gauge-parameter shift acts as a BRST-exact variation on the band algebra: $\delta_\theta(\cdot) = [\,\hat{Q}_{\text{BRST}}, X_\theta]$. Writing the valuation as a band functional $\mathcal{J}^{(a)}(A) := \mathrm{Tr}\Big(P^{(a)}_{\text{para}}\, A\Big)$ (ghost number 0), BRST-closedness of $P^{(a)}_{\text{para}}$ and the Nielsen safety on the band (Proposition 9.97) give

$$\partial_\theta v^{(a);[\chi]}_{\text{Synthesis}} = \mathcal{J}^{(a)}\Big(\delta_\theta \rho^{(a)[\chi]}_{\text{band}}\Big) = \mathcal{J}^{(a)}\Big([\,\hat{Q}_{\text{BRST}}, X_\theta]\Big) = 0.$$

The same argument applies to $v^{(LS);[\chi]}_{\text{Synthesis}}$ and $v^{(LS);[\chi]}_{\text{Agreement}}$, and uses in addition that $[\,\hat{Q}_{\text{BRST}}, \Pi^{[\chi]}] = 0$ (Definition 9.116) so that any typed restriction entering the construction produces only BRST-exact variations annihilated by band functionals (cf. also Lemmas 9.121 and 9.128).

(c) Retardedness and cone support. By construction all inputs to the meters (the correlators used to form $\rho^{(\cdot)[\chi]}_{\text{band}}$ and the typed restrictions effected by $\Pi^{[\chi]}$) use retarded, cone-gated kernels (Definition 9.106). These are supported in $J^+\Big(\mathcal{C}^{\hat{\Sigma},*}\Big)$ (Proposition 9.107), and microcausality holds for spacelike separated supports (Proposition 9.108). Since the meters are algebraic combinations (traces with positive effects) of these retarded, cone-supported inputs, their valuations inherit the same causal support.

Combining (a)–(c) proves the lemma. □

Proposition 9.146 (Monotonicity under cone shrink and band retuning)**.** *If the IR window tightens as in Lemma 9.105, then for fixed typed inputs* $v^{(\cdot);[\chi]}_{\text{Synthesis}}$ *is nondecreasing towards* $1/2$ *(loss of accessible coherence cannot raise asymmetry beyond the available support). Moreover, within the stability radius of Definition 9.116,* $v^{(\cdot);[\chi]}_{\text{Synthesis}}$ *depends continuously on* $(\lambda_{\text{UV}}, \ell)$.

Proof of Proposition 9.146. Fix a permitted type χ and an anchor $a \in \{L, S, LS\}$. The typed, band-reduced meters are defined in Definitions 9.141, 9.143 and 9.144; for the local anchors one has $v^{(a);[\chi]}_{\text{Synthesis}} = \frac{1}{2} + \mathrm{Re}\Big(\rho^{(a)[\chi]}_{12}\Big)$, with $\rho^{(a)[\chi]}$ the 2×2 para-density obtained from the typed, cone-gated band data. All kernels entering $\rho^{(a)[\chi]}$ are retarded and cone-gated (Definition 9.106 and Proposition 9.107), and the projector $\Pi^{[\chi]}$ is BRST-closed (Definition 9.116).

(A) Cone shrink $\ell' \le \ell$ contracts accessible coherence. Tightening the IR window from ℓ to ℓ' (with $\ell' \le \ell$) *shrinks* the Σ–cone (Lemma 9.105) and thus replaces the retarded cone gate by a smaller positive contraction in the Löwner order:

$$0 \;\le\; \Gamma_{\ell'} \;\le\; \Gamma_\ell \;\le\; \mathbf{1}, \qquad \Gamma_\bullet \text{ as in Definition 9.106.} \tag{9.77}$$

In the para-anchor two-level reduction, the matrix elements $\rho^{(a)[\chi]}_{ij}$ are band functionals (ghost number 0) of typed, cone-gated correlators; concretely, there exist positive linear functionals $\{\mathcal{J}^{(a)}_{ij}\}$ on the band algebra such that

$$\rho^{(a)[\chi]}_{ij}(\ell) = \frac{\mathcal{J}^{(a)}_{ij}\Big(\Gamma_\ell \cdot \Gamma_\ell\Big)}{\mathcal{J}^{(a)}_{11}\Big(\Gamma_\ell \cdot \Gamma_\ell\Big) + \mathcal{J}^{(a)}_{22}\Big(\Gamma_\ell \cdot \Gamma_\ell\Big)},$$

where the dot denotes the (fixed) typed operator input determined by $\Pi^{[\chi]}$ and the anchor a (cf. the construction in Definitions 9.141, 9.143 and 9.144). Introduce the Γ–weighted sesquilinear form on the anchor space $\langle u, v\rangle_\Gamma := \mathcal{J}^{(a)}\big(u^\dagger\,\Gamma\,(\cdot)\,\Gamma\,v\big)$; then Cauchy–Schwarz in the seminorm $\|u\|_\Gamma^2 = \langle u, u\rangle_\Gamma$ gives

$$\left|\rho_{12}^{(a)[\chi]}(\ell)\right| \;\le\; \sqrt{\rho_{11}^{(a)[\chi]}(\ell)\;\rho_{22}^{(a)[\chi]}(\ell)}\,. \tag{9.78}$$

By (9.77) and positivity of the band functionals (Definition 9.96), the diagonals are monotone nonincreasing under $\Gamma_\ell \mapsto \Gamma_{\ell'}$:

$$\rho_{ii}^{(a)[\chi]}(\ell') \;\le\; \rho_{ii}^{(a)[\chi]}(\ell) \qquad (i = 1, 2). \tag{9.79}$$

Applying (9.78) at level ℓ' and using (9.79) yields the *contraction of the coherence deviation*:

$$\begin{aligned}\left|\, v_{\text{Synthesis}}^{(a);[\chi]}(\ell') - \tfrac{1}{2}\,\right| &= \left|\,\mathrm{Re}\;\rho_{12}^{(a)[\chi]}(\ell')\right| \le \left|\rho_{12}^{(a)[\chi]}(\ell')\right| \\ &\le \sqrt{\rho_{11}^{(a)[\chi]}(\ell')\,\rho_{22}^{(a)[\chi]}(\ell')} \le \sqrt{\rho_{11}^{(a)[\chi]}(\ell)\,\rho_{22}^{(a)[\chi]}(\ell)}\,.\end{aligned} \tag{9.80}$$

In particular, the *distance to the baseline* $1/2$ cannot increase when the cone shrinks. Under the standard anchor phase convention used in §8.6 (the off-diagonal is taken real and nonnegative in the fixed $\{|\mathrm{Sym}\rangle_a, |\mathrm{Break}\rangle_a\}$ basis; cf. Definitions 9.141, 9.143 and 9.144), (9.80) reads $v_{\text{Synthesis}}^{(a);[\chi]}(\ell') \;\le\; v_{\text{Synthesis}}^{(a);[\chi]}(\ell)$, i.e. $v_{\text{Synthesis}}^{(a);[\chi]}$ moves *monotonically toward* $1/2$ as $\ell \downarrow$. For a general phase, the statement holds as monotone contraction of $\left|v_{\text{Synthesis}}^{(a);[\chi]} - \frac{1}{2}\right|$, which is the operational asymmetry magnitude. The same argument applies verbatim to the relational meter $v_{\text{Synthesis}}^{(LS);[\chi]}$ (and thus to its deviation from $1/2$), since its construction is again a trace against a positive effect on a typed, cone-gated band state (Definitions 9.141, 9.143 and 9.144).

(B) Continuity under band retunings within the stability radius. Let $(\lambda_{\mathrm{UV}}, \ell) \mapsto (\lambda'_{\mathrm{UV}}, \ell')$ be a retuning inside the stability radius of item (P7) in Definition 9.116. Then the typed Riesz projector $\Pi^{[\chi]}$ depends continuously on the parameters (Proposition 9.119), the UV/IR taggers and cone gates vary continuously in operator norm within the band class (Definitions 9.96 and 9.106 and Proposition 9.107), and the band functionals are bounded linear maps (Definition 9.96). Therefore each matrix element $\rho_{ij}^{(a)[\chi]}$ depends continuously on $(\lambda_{\mathrm{UV}}, \ell)$, and so do $v_{\text{Synthesis}}^{(a);[\chi]} = \frac{1}{2} + \mathrm{Re}(\rho_{12}^{(a)[\chi]})$ and $v_{\text{Synthesis}}^{(LS);[\chi]}$. Hence $v_{\text{Synthesis}}^{(\cdot);[\chi]}$ is continuous in $(\lambda_{\mathrm{UV}}, \ell)$ within the stability radius.

Combining (A) and (B) proves that tightening the IR window cannot increase the measured asymmetry: the synthesis meters move (in magnitude) monotonically toward the baseline $\frac{1}{2}$, and they vary continuously under band retunings inside the certified stability neighborhood. □

4.2 Typed T–L–O readouts

Definition 9.147 (Typed T–L–O map)**.** For each permitted type χ at spacetime point x and scale parameter r (e.g. window size or dilation), define the T–L–O triple

$$\mathrm{TLO}^{[\chi]}(x; r) \;:=\; \Big(\underbrace{\text{Type}}_{=\,\mathrm{Quad}^{[\chi]}(x)}\;,\;\underbrace{\text{Location}}_{\text{arena/sector/position/cone sheet}}\;,\;\underbrace{\text{Onset}}_{(r^\star[\chi],\,t^\star[\chi])}\Big),$$

with components:

(i) **Type:** $\mathrm{Quad}^{[\chi]}(x) \in \{\mathrm{QI}, \mathrm{QII}, \mathrm{QIII}, \mathrm{QIV}\}$ from Definition 9.137.

(ii) **Location:** $\mathcal{A} \in \{\mathfrak{P}, \mathfrak{P}_\partial\}$, sector $\Phi\Phi$ or $\Theta\Theta$, spatial tag $P_x(\lambda_{\mathrm{UV}})$, and cone sheet (future/past, null/interior) determined by $\mathcal{C}_x^{\hat{\Sigma},*}$ (Definition 9.104).

(iii) **Onset (scale/time):** with tolerances $(\varepsilon_{\mathrm{law}}, \varepsilon_{\mathrm{state}})$ and law-deficit $d_{\mathrm{law}}^{\Sigma;[\chi]}$ from (9.68), set

$$\begin{aligned} r^\star_{\mathrm{SSB}}[\chi] &:= \inf\Big\{ r :\ v^{(S);[\chi]}_{\mathrm{Synthesis}}(r) > \tfrac{1}{2} + \varepsilon_{\mathrm{state}} \\ &\qquad\qquad \wedge\ v^{(LS);[\chi]}_{\mathrm{Synthesis}}(r) \downarrow \tfrac{1}{2} \Big\}, \\ r^\star_{\mathrm{QIV}}[\chi] &:= \inf\Big\{ r :\ d^{\Sigma;[\chi]}_{\mathrm{law}}(r) \geq \delta_\star \Big\}, \\ r^\star_{\mathrm{QIII}}[\chi] &:= \inf\Big\{ r :\ d^{\Sigma;[\chi]}_{\mathrm{law}}(r) \leq C\, r^{-\beta} \Big\}, \end{aligned} \tag{9.81}$$

and define temporal onsets $t^\star[\chi]$ analogously from the time traces $t \mapsto v^{(\cdot);[\chi]}_{\mathrm{Synthesis}}(t; r)$.

Lemma 9.148 (Existence of onsets under mild regularity)**.** *If $r \mapsto v^{(S);[\chi]}_{\mathrm{Synthesis}}(r)$ and $r \mapsto v^{(LS);[\chi]}_{\mathrm{Synthesis}}(r)$ are piecewise monotone with one-sided limits at infinity, and $r \mapsto d^{\Sigma;[\chi]}_{\mathrm{law}}(r)$ admits a regularly varying envelope (as in the large-window law test of Definition 7.685), then the infima in (9.81) are finite or $+\infty$ according to whether the corresponding threshold is eventually met or not; in both cases the onsets are well-defined.*

Proof of Lemma 9.148. Recall the onset rules from Definition 9.147 (eq. (9.81)):

$$r^\star_{\mathrm{SSB}}[\chi] := \inf\Big\{r :\ v^{(S);[\chi]}_{\mathrm{Synthesis}}(r) > \tfrac{1}{2} \ \wedge\ v^{(LS);[\chi]}_{\mathrm{Synthesis}}(r) \downarrow \tfrac{1}{2}\Big\},$$
$$r^\star_{\mathrm{QIV}}[\chi] := \inf\Big\{r :\ d^{\Sigma;[\chi]}_{\mathrm{law}}(r) \geq \delta_\star\Big\}, \qquad r^\star_{\mathrm{QIII}}[\chi] := \inf\Big\{r :\ d^{\Sigma;[\chi]}_{\mathrm{law}}(r) \leq C\, r^{-\beta}\Big\},$$

with the understanding that tolerances $(\varepsilon_{\mathrm{law}}, \varepsilon_{\mathrm{state}})$ may be inserted as in Definition 9.147, and where the *typed law–deficit* $d^{\Sigma;[\chi]}_{\mathrm{law}}$ is induced from the typed law–contrast of Definition 9.127. By Lemma 9.145(a) all meters take values in $[0, 1]$, so the threshold statements are meaningful.

(1) SSB onset $r^\star_{\mathrm{SSB}}[\chi]$. Assume $r \mapsto v^{(S);[\chi]}_{\mathrm{Synthesis}}(r)$ and $r \mapsto v^{(LS);[\chi]}_{\mathrm{Synthesis}}(r)$ are piecewise monotone on $[R_0, \infty)$ and admit one–sided limits at $+\infty$ (hypothesis of the lemma; the construction of the meters is in Definitions 9.141, 9.143 and 9.144). Set

$$A := \Big\{r \geq R_0 :\ v^{(S);[\chi]}_{\mathrm{Synthesis}}(r) > \tfrac{1}{2} \ \wedge\ v^{(LS);[\chi]}_{\mathrm{Synthesis}}(r) \downarrow \tfrac{1}{2}\Big\},$$

where "$\downarrow \frac{1}{2}$" abbreviates the monotone approach to $\frac{1}{2}$ from above within the current monotonicity interval (cf. the tolerance semantics in Definition 9.147). Because each meter is piecewise monotone and has a one–sided limit at $+\infty$, either (i) the joint condition in the definition of A is *eventually* satisfied (then A contains a tail interval (R, ∞) for some R), or (ii) it is *eventually* not satisfied (then A is empty beyond some scale). In case (i) $\inf A \in [R_0, \infty)$ is finite; in case (ii) $A = \varnothing$ or A is bounded, and by the convention of Definition 9.147 the infimum is $+\infty$. Thus $r^\star_{\mathrm{SSB}}[\chi] = \inf A \in [0, \infty]$ is well-defined.

(2) QIV onset $r^{\star}_{\mathrm{QIV}}[\chi]$. Let $B := \{r \ge R_1 : \ d^{\Sigma;[\chi]}_{\mathrm{law}}(r) \ge \delta_\star\}$ and assume that $d^{\Sigma;[\chi]}_{\mathrm{law}}$ admits a *regularly varying envelope* as in the large-window law test of Definition 7.685: there exists an eventually monotone function $e(r)$ with $|d^{\Sigma;[\chi]}_{\mathrm{law}}(r)| \le e(r)$ for all $r \ge R_1$ and $e(r) \sim r^{-\alpha}L(r)$ (with slowly varying L) or is asymptotically constant.[3] Then either $e(r) \ge \delta_\star$ eventually, in which case $d^{\Sigma;[\chi]}_{\mathrm{law}}(r) \ge \delta_\star$ eventually and B contains a tail (R, ∞), giving $\inf B < \infty$; or else $e(r) < \delta_\star$ eventually, and then B is empty beyond some scale, giving $\inf B = +\infty$. Hence $r^{\star}_{\mathrm{QIV}}[\chi] = \inf B$ is well-defined and equals a finite scale if and only if the threshold is eventually met.

(3) QIII onset $r^{\star}_{\mathrm{QIII}}[\chi]$. Let $C := \{r \ge R_2 : \ d^{\Sigma;[\chi]}_{\mathrm{law}}(r) \le C_0\, r^{-\beta}\}$. With the same envelope $e(r)$ as above and using $|d^{\Sigma;[\chi]}_{\mathrm{law}}(r)| \le e(r)$ for $r \ge R_2$, either $e(r) \le C_0\, r^{-\beta}$ eventually (then C contains a tail and $\inf C < \infty$), or $e(r) > C_0\, r^{-\beta}$ eventually (then C is empty on a tail and $\inf C = +\infty$). Thus $r^{\star}_{\mathrm{QIII}}[\chi] = \inf C$ is well-defined and is finite exactly when the decay threshold is eventually attained.

In each case the onset is the infimum of a (possibly empty) subset of $[0, \infty)$ determined by eventually monotone data; therefore it exists as an extended nonnegative real number and equals $+\infty$ precisely when the corresponding threshold is not eventually satisfied. This establishes the lemma. □

Proposition 9.149 (Typed Type–Location–Onset is BRST/causally robust)**.** *The T–L–O assignment* $\mathrm{TLO}^{[\chi]}(x; r)$ *is (a) gauge-parameter independent (by Lemma 9.145 and Proposition 9.97); (b) causally supported within* $J^+(\mathcal{C}^{\hat{\Sigma},*})$ *(by Proposition 9.107); and (c) stable under band retunings within the radius of Definition 9.116 (continuity of meters and of* $\Lambda^{\Sigma;[\chi]}$ *in Proposition 9.134).*

Proof of Proposition 9.149. Fix a permitted type χ and a space–scale pair $(x; r)$. By Definition 9.147, the typed Type–Location–Onset assignment is

$$\mathrm{TLO}^{[\chi]}(x; r) := \Big(\underbrace{\mathrm{Quad}^{[\chi]}(x)}_{\text{Type, Definition 9.137}}, \ \underbrace{\mathrm{Loc}^{[\chi]}(x)}_{\text{arena/sector/position/cone sheet, Definition 9.147}}, \ \underbrace{\mathrm{Onset}^{[\chi]}(r)}_{\text{rules (9.81) in Definition 9.147}} \Big),$$

where $\mathrm{Quad}^{[\chi]}$ is computed from the typed visibilities $\vec{\Lambda}^{[\chi]}(x) = (\Lambda^{\Sigma;[\chi]}_{\mathrm{law}}(x), \Lambda^{\Sigma;[\chi]}_{\mathrm{state}}(x))$ (Definition 9.133), the location tags are read from the band instrumentation and cone sheet (Definitions 9.96 and 9.106), and the onset entries use the synthesis meters and typed law–deficit (Definitions 9.127, 9.141, 9.143, 9.144 and 9.147).

(a) Gauge/BRST independence. Pointwise typed contrasts are BRST/Nielsen-safe by Lemma 9.128 together with the band Nielsen property Proposition 9.97 and $[\hat{Q}_{\mathrm{BRST}}, \Pi^{[\chi]}] = 0$

[3] The only properties used here are eventual monotonicity of e and that e bounds $|d_{\mathrm{law}}|$ from above on a tail.

(Definition 9.116). Hence the aggregated contrasts and visibilities are gauge-invariant (Proposition 9.134(a)), and the quadrant decision $\mathrm{Quad}^{[\chi]}(x)$ is therefore gauge-independent (Definition 9.137). The location tags are built from BRST-closed band taggers $P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}^{(\ell)}$ (Definition 9.96) and cone-sheet labels (from $\mathcal{C}^{\hat{\Sigma},*}$, Definition 9.104), so they are invariant as well. For the onset rules (9.81), the valuations entering the thresholds are gauge-invariant: the synthesis meters are BRST/Nielsen-safe by Lemma 9.145(b), while the typed law–deficit is built from the typed contrasts (already gauge-invariant). Hence the infima defining all onset components are unchanged under gauge-parameter variations.

(b) Causal support within $J^+\big(\mathcal{C}^{\hat{\Sigma},*}\big)$. All kernels used to form the typed contrasts/visibilities and the meter states are *retarded and cone-gated* (Definition 9.106), with support in $J^+\big(\mathcal{C}^{\hat{\Sigma},*}\big)$ by Proposition 9.107; the commutators vanish at spacelike separation by Proposition 9.108. Consequently the visibilities (hence the quadrant) are causally well-posed (Proposition 9.134(b)), and the meter valuations inherit the same causal support (Lemma 9.145(c)). Therefore every component of $\mathrm{TLO}^{[\chi]}(x;r)$ is produced from retarded, cone-supported inputs, proving causal robustness.

(c) Stability under band retunings within the certified radius. Inside the stability radius of item (P7) in Definition 9.116, the typed Riesz projector $\Pi^{[\chi]}$ varies continuously (Proposition 9.119); the band taggers and cone gates vary continuously in operator norm within the band class (Definitions 9.96 and 9.106), and the band functionals are bounded linear (Definition 9.96). Hence the pointwise typed contrasts vary continuously, and so do the aggregates by Lemma 9.132; applying the continuous normalizer yields continuity of $\vec{\Lambda}^{[\chi]}$ (cf. Definition 9.133 and Proposition 9.134). For the meters, continuity under retuning is stated in Proposition 9.146(B). Therefore the data entering the onset rules (9.81)—the synthesis meters and the typed law–deficit—vary continuously under retunings, and the resulting onsets are stable (changes can only occur at the declared tolerance thresholds; see also the plateau stability discussion in Corollary 9.139). Altogether, the triple $\mathrm{TLO}^{[\chi]}(x;r)$ depends continuously on $(\lambda_{\mathrm{UV}}, \ell)$ within the certified neighborhood.

Combining (a)–(c) proves that the typed T–L–O assignment is gauge/BRST-independent, causally supported in $J^+(\mathcal{C}^{\hat{\Sigma},*})$, and stable under admissible band retunings. □

Corollary 9.150 (Consistency with typed quadrants and sector rules)**.** *If $\Pi^{[\chi]} = \mathbb{I}$ on a block, then the T–L–O map reduces to the sector/untyped construction of Definitions 9.28 and 9.47, and the onset criteria* (9.81) *coincide with the large-window (QIII vs. QIV) decision rule in Definition 7.685.*

Proof of Corollary 9.150. Fix a block $\bullet \in \{\Phi\Phi, \Theta\Theta\}$ and take the typed projector to be the identity on that block, $\Pi^{[\chi]} = \mathbb{I} = E_{\bullet\bullet}$ (item (P1) of Definition 9.116).

(1) Typed ⇒ sector reduction for contrasts, visibilities, quadrants. By Proposition 9.138, the typed restriction becomes a no-op: $\hat{Q}_x^{[\chi]}[v] = \hat{Q}_x[v]$ and the pointwise/aggregated contrasts (Definitions 9.127 and 9.133) match their sector counterparts in Definition 9.45. Applying the same proposition, the normalized visibilities coincide with the sector/untyped visibilities of Definition 9.47, and the typed quadrant rule $\mathrm{Quad}^{[\chi]}$ (Definition 9.137) equals the canonical sector decision Definition 7.685. Thus the *Type* component of $\mathrm{TLO}^{[\chi]}$ reduces to that of Definitions 9.28 and 9.47.

(2) Reduction of meter valuations and location tags. The band-reduced, typed parasymmetric meters (Definitions 9.141, 9.143 and 9.144) are traces against positive effects built from the same band data; with $\Pi^{[\chi]} = \mathbb{I}$ the typed restriction is trivial, so the local and relational valuations $v^{(a);[\chi]}_{\mathrm{Synthesis}}$, $v^{(LS);[\chi]}_{\mathrm{Synthesis}}$ and $v^{(LS);[\chi]}_{\mathrm{Agreement}}$ coincide with their sector/untyped versions used in §8.6. Likewise, the *Location* entry of $\mathrm{TLO}^{[\chi]}$ (arena/sector/position/cone sheet in Definition 9.147) is unchanged, since it depends only on the band taggers and the Σ–cone (Definitions 9.96 and 9.106), not on the typed restriction.

(3) Onset rules coincide with the large-window sector rule. The onset components in $\mathrm{TLO}^{[\chi]}$ are defined by the infima in (9.81) of Definition 9.147, using (i) the synthesis meters from Definitions 9.141, 9.143 and 9.144 and (ii) the typed law–deficit $d^{\Sigma;[\chi]}_{\mathrm{law}}$ induced by the typed law–contrast (Definition 9.127). With $\Pi^{[\chi]} = \mathbb{I}$, item (i) reduces to the sector/untyped meters (Step 2) and item (ii) reduces to the sector law–deficit from Definition 9.45. Therefore the scale thresholds $r^{\star}_{\mathrm{QIII}}[\chi]$ and $r^{\star}_{\mathrm{QIV}}[\chi]$ in (9.81) are *exactly* the large-window (QIII vs. QIV) decision thresholds stated at sector level in Definition 7.685 (decay to $Cr^{-\beta}$ versus exceeding $\delta_{\star}$), and $r^{\star}_{\mathrm{SSB}}[\chi]$ matches the sector SSB criterion since it is formulated purely in terms of the (now sector) synthesis meters.

Combining (1)–(3) establishes that, for $\Pi^{[\chi]} = \mathbb{I}$, the full T–L–O map reduces to the sector/untyped construction of Definitions 9.28 and 9.47, and the onset criteria (9.81) coincide with the large-window decision rule of Definition 7.685, as claimed. □

Remark 9.151 (Boundary arenas and cone-sheet annotations)**.** Repeating the definitions in Definition 9.141–Definition 9.147 on the boundary plateau $\mathfrak{P}_{\partial}$ involves replacing the IR window $K^{(\ell)}_{\mathrm{IR}}$ with its tangential version and using the boundary cone $\mathcal{C}^{\hat{\Sigma},*}_{x;\partial}$ (inherited from Definition 9.104). The "cone sheet" tag in Location records if the readout is on a null sheet or in the timelike interior; this in turn controls admissible χ via typed hyperbolicity (Definition 9.111).

Remark 9.152 (Operational pipeline)**.** Given a permitted χ: (1) compute $\Lambda^{\Sigma;[\chi]}_{\mathrm{law/state}}$ (Definition 9.133); (2) assign $\mathrm{Quad}^{[\chi]}$ (Definition 9.137); (3) evaluate $v^{(a);[\chi]}_{\mathrm{Synthesis}}$, $v^{(LS);[\chi]}_{\mathrm{Synthesis}}$ (Definitions 9.143 and 9.144); (4) report $\mathrm{TLO}^{[\chi]}$ using (9.81). Each step is retarded, cone-gated, and BRST-clean by Propositions 9.97 and 9.107 and Lemma 9.145.

9.13 Per Sector Execution Outline

We formalize the sector-wise execution as a compositional pipeline that takes $(\hat{\Sigma}, \hat{\mathbf{G}}_{\Lambda}, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell})$ together with a probe family $(\mathcal{V}, w)$ and produces the typed Type–Location–Onset (T–L–O) artefacts for each χ permitted in the chosen sector. All objects and operations below are band-admissible, retarded, cone-gated, and BRST/Nielsen-safe by Definitions 9.96 and 9.106 and Propositions 9.97 and 9.107.

5.1 Sector execution functor

Definition 9.153 (Sector execution functor)**.** Fix a sector label $\bullet \in \{\Phi\Phi, \Theta\Theta\}$. The *sector execution functor*

$$\mathscr{E}_\bullet : \left(\hat{\Sigma}, \hat{\mathbf{G}}_\Lambda, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell};\ \mathcal{V}, w\right) \longmapsto \left(\mathcal{C}^{[\bullet]}_{\text{admissible}},\ \{\Lambda^{\Sigma;[\chi]}_{\text{law/state}}\}_\chi,\ \{v^{(\cdot);[\chi]}_{\text{Synthesis}}\}_\chi,\ \{\mathrm{TLO}^{[\chi]}\}_\chi\right)$$

is the composition of the following typed, band-level stages:

(E1) **Permit ledger construction:** build $\mathcal{C}^{[\bullet]}_{\text{admissible}}$ by producing typed projectors $\Pi^{[\chi]}$ that satisfy (P1)–(P7) in Definition 9.116 and the typed hyperbolicity test (9.53) (Definition 9.111), using the Σ-weighted spectral construction of Proposition 9.119. This realizes Definition 9.122.

(E2) **Typed visibilities:** for each $\chi \in \mathcal{C}^{[\bullet]}_{\text{admissible}}$, compute the aggregated contrasts $\Delta^{\mathrm{pre},\Sigma;[\chi]}_\bullet(x;\mathcal{V},w)$ (Definition 9.131) and normalize to visibilities $\Lambda^{\Sigma;[\chi]}_{\text{law/state}}(x) \in [0,1]$ by Definition 9.133.

(E3) **Parasymmetric meters & T–L–O:** evaluate the typed meters $v^{(a);[\chi]}_{\text{Synthesis}}$ and $v^{(LS);[\chi]}_{\text{Synthesis}}$ (Definitions 9.143 and 9.144) and return the T–L–O readouts $\mathrm{TLO}^{[\chi]}(x;r)$ using Definition 9.147, with Type $= \mathrm{Quad}^{[\chi]}(x)$ from Definition 9.137.

(E4) **Boundary pass (optional):** repeat (E1)–(E3) on the boundary arena $\mathfrak{P}_\partial$ using the induced boundary cone (Remark in §9.147) and record bulk/edge differences.

Lemma 9.154 (Well-posedness and termination of $\mathscr{E}_\bullet$)**.** *Under band-finiteness and compactness hypotheses of Proposition 9.119, the spectral construction in (E1) yields a locally finite family of isolated clusters, each producing a finite-rank projector $\Pi^{[\chi]}$. Hence (E1) terminates at any fixed spectral resolution and bandwidth, and (E2)–(E3) are well-posed by Lemmas 9.129 and 9.145 and Proposition 9.134.*

Proof of Lemma 9.154. We unpack the three stages of the execution pipeline $\mathscr{E}_\bullet$: **(E1)** spectral selection and typed Riesz projectors inside a chosen block ($\Phi\Phi$ or $\Theta\Theta$); **(E2)** construction of typed contrasts/aggregates/visibilities (Definitions 9.127 and 9.133); **(E3)** construction of typed parasymmetric meters and the ensuing T–L–O (Definitions 9.141, 9.143, 9.144 and 9.147).

(E1) Local finiteness, isolated clusters, finite rank, termination. By Proposition 9.119 the Σ–weighted channel operator $K_\bullet$ (constructed as in Proposition 9.36 and (9.16) within a fixed block) is positive and compact on the band-reduced Hilbert space. Hence its nonzero spectrum consists of at most countably many eigenvalues with finite multiplicities, accumulating only at 0. In particular, on any compact spectral window bounded away from 0 there are *finitely many* eigenvalues, thus only finitely many *isolated clusters*. Each such cluster admits a Riesz projector $\Pi^{[\chi]} = \frac{1}{2\pi i}\oint_\Gamma (z\mathbf{1} - K_\bullet)^{-1}\,dz$ of *finite rank*, and, by Proposition 9.119, these projectors are band-finite, BRST-closed, graded-covariant, and stable under small retunings (items (P1)–(P7) of Definition 9.116). Consequently, for any fixed spectral resolution (i.e. a finite partition of the chosen window into disjoint contours Γ) and any fixed bandwidth, only finitely many clusters are selected and only finitely many $\Pi^{[\chi]}$ are produced. Hence **(E1)** terminates at the chosen resolution/bandwidth.

(E2) Well-posed typed contrasts/visibilities. Given $\Pi^{[\chi]}$, the pointwise typed operands in Definition 9.127 are Hermitian and cone-gated; for timelike covectors k one has reality and nonnegativity of the band evaluations by Lemmas 9.110 and 9.129. Retardedness and cone support are ensured by Definition 9.106 and Proposition 9.107, while microcausality at spacelike separation holds by Proposition 9.108. Aggregation and normalization (Definition 9.133) then produce visibilities in $[0,1]$ that are gauge/BRST-invariant (Proposition 9.134(a)), causal (Proposition 9.134(b)), and properly normalized (Proposition 9.134(c)). Thus **(E2)** is well-posed.

(E3) Well-posed typed meters and T–L–O. The band-reduced, typed meters of Definitions 9.141, 9.143 and 9.144 are traces of positive effects against positive band states; therefore their valuations lie in $[0,1]$ and are gauge/BRST-invariant by Lemma 9.145(a,b). Their inputs are retarded and cone-gated, so causal support is inside $J^+\big(\mathcal{C}^{\hat{\Sigma},*}\big)$ (Lemma 9.145(c) together with Proposition 9.107). The T–L–O assignment then inherits gauge/causal robustness and stability under certified retunings by Proposition 9.149. Hence **(E3)** is well-posed.

Combining the compact spectral structure of $K_\bullet$ (local finiteness and finite rank of Riesz spectral projectors; Proposition 9.119) with the positivity/causality/gauge properties of the typed contrasts and meters (Lemmas 9.129 and 9.145 and Proposition 9.134) establishes both the termination of **(E1)** at fixed resolution/bandwidth and the well-posedness of **(E2)**–**(E3)**. □

Proposition 9.155 (Soundness (BRST/causal correctness))**.** *Every numerical output of $\mathscr{E}_\bullet$ (namely, $\Lambda^{\Sigma;[\chi]}_{\mathrm{law/state}}$, $v^{(\cdot);[\chi]}_{\mathrm{Synthesis}}$, and $\mathrm{TLO}^{[\chi]}$) satisfies two key properties. First, it is gauge-parameter independent (BRST/Nielsen-safe), as shown in Proposition 9.97 and Lemmas 9.128 and 9.145. Second, it is causally supported within $J^+(\mathcal{C}^{\hat{\Sigma},*})$, as established by Propositions 9.107 and 9.108.*

Proof of Proposition 9.155. Fix a permitted type χ and a spacetime point x (and, for onsets, a scale r). The numerical outputs of $\mathscr{E}_\bullet$ are: (i) the typed visibilities $\Lambda^{\Sigma;[\chi]}_{\mathrm{law/state}}(x)$ (Definition 9.133, built from the typed contrasts of Definition 9.127); (ii) the typed parasymmetric valuations $v^{(\cdot);[\chi]}_{\mathrm{Synthesis}}$ (Definitions 9.141, 9.143 and 9.144); and (iii) the typed Type–Location–Onset triple $\mathrm{TLO}^{[\chi]}(x;r)$ (Definition 9.147).

(A) Gauge-parameter independence (BRST/Nielsen safety). For each probe v, the *pointwise* typed contrasts are invariant under BRST-exact gauge variations by Lemma 9.128, using in particular the band Nielsen property Proposition 9.97 and the BRST-closedness $[\hat{Q}_{\mathrm{BRST}}, \Pi^{[\chi]}] = 0$ of Definition 9.116. Aggregation and normalization then preserve invariance, yielding gauge-invariant visibilities; this is summarized in Proposition 9.134(a). For the meters, BRST/Nielsen safety of all valuations is stated in Lemma 9.145(b). Finally, $\mathrm{TLO}^{[\chi]}$ is composed of the (gauge-invariant) quadrant extracted from the visibilities, gauge-invariant location tags (built from BRST-closed band taggers, Definition 9.96), and onsets defined via gauge-invariant meters and law-deficits; its gauge/BRST independence is recorded in Proposition 9.149(a). Therefore, every output listed in the proposition is BRST/Nielsen-safe (Proposition 9.97 and Lemmas 9.128 and 9.145).

(B) Causal support inside $J^+\big(\mathcal{C}^{\hat{\Sigma},*}\big)$. All kernels used in $\mathscr{E}_\bullet$ are replaced by their retarded, cone-gated versions (Definition 9.106), which are supported in $J^+\big(\mathcal{C}^{\hat{\Sigma},*}\big)$ by Proposition 9.107. Micro-

causality at spacelike separation then holds for the unified fields (Proposition 9.108), ensuring the pointwise operands that feed the contrasts are well-defined and real (Lemma 9.129). Consequently, the visibilities inherit retarded, cone-gated support (Proposition 9.134(b)). For the meters, causal support of the inputs implies causal support of the valuations (Lemma 9.145(c)). The $\mathrm{TLO}^{[\chi]}$ assignment aggregates only these retarded, cone-supported ingredients (visibilities, meters, and location tags from the cone sheet), and is therefore causally robust by Proposition 9.149(b). Thus every numerical output is supported within $J^{+}\big(\mathcal{C}^{\hat{\Sigma},*}\big)$ (Propositions 9.107 and 9.108).

Combining (A) and (B) proves that all outputs of $\mathscr{E}_{\bullet}$ are both gauge-parameter independent and causally well-posed in the asserted sense. □

Proposition 9.156 (Stability and functoriality)**.** *Within the retuning-stability radius in Definition 9.116, the maps*

$$(\lambda_{\mathrm{UV}},\ell)\ \longmapsto\ \mathcal{C}^{[\bullet]}_{\mathrm{admissible}},\quad (\lambda_{\mathrm{UV}},\ell)\ \longmapsto\ \Lambda^{\Sigma;[\chi]}_{\mathrm{law/state}},\quad (\lambda_{\mathrm{UV}},\ell)\ \longmapsto\ v^{(\cdot);[\chi]}_{\mathrm{Synthesis}}$$

are continuous. Tightening the IR window as in Lemma 9.105 shrinks the ledger (Proposition 9.123) and increases visibilities monotonically towards unity (less apparent breaking) by Lemma 9.136, while meters vary continuously by Proposition 9.146.

Proof of Proposition 9.156. Fix a permitted type χ and consider retunings $(\lambda_{\mathrm{UV}},\ell)$ within the stability radius of item (P7) in Definition 9.116.

(1) Continuity of the ledger map. Inside the certified radius, each typed Riesz projector $\Pi^{[\chi]}$ depends *continuously* on $(\lambda_{\mathrm{UV}},\ell)$ in operator norm by the spectral perturbation stability of Proposition 9.119. The cone/hyperbolicity certificate for χ (Corollary 9.120) supplies a positive spectral gap and an admissible timelike interval; small retunings preserve the positivity margin (9.53), hence the admission test (Proposition 9.102) continues to hold. Therefore the ledger $(\lambda_{\mathrm{UV}},\ell)\mapsto\mathcal{C}^{[\bullet]}_{\mathrm{admissible}}$ is *locally constant* (hence continuous as a set-valued map) in the stability neighborhood.

(2) Continuity of $\Lambda^{\Sigma;[\chi]}_{\mathrm{law/state}}$. For each probe v, the pointwise typed operands in (9.58)–(9.59) (Definition 9.127) depend continuously on $(\lambda_{\mathrm{UV}},\ell)$ because (i) $\Pi^{[\chi]}$ varies continuously (Proposition 9.119), (ii) the band taggers vary continuously within the band class (Definition 9.96), and (iii) the cone gate and retardedness are compatible with instrumentation (Definition 9.106 and Proposition 9.107). The band functionals $\mathcal{I},\mathcal{I}_{\Psi}$ are bounded linear (Definition 9.96), so each pointwise contrast is continuous; the aggregated $\Delta^{\mathrm{pre},\Sigma;[\chi]}_{\bullet}(x;\mathcal{V},w)$ is continuous by the aggregation continuity of Lemma 9.132. Applying the continuous normalizer (visibility map) from Definitions 9.47 and 9.133 yields continuity of $\Lambda^{\Sigma;[\chi]}_{\mathrm{law/state}}$.

(3) Continuity of $v^{(\cdot);[\chi]}_{\mathrm{Synthesis}}$. Within the same stability neighborhood, the meter valuations vary continuously by Proposition 9.146(B), since they are traces of positive effects against band-reduced states that depend continuously on $\Pi^{[\chi]}$, the taggers, and the cone gate (Definitions 9.96, 9.106, 9.141, 9.143 and 9.144).

(4) Effect of IR tightening $\ell'\le\ell$. By Lemma 9.105, the Σ–cone contracts under IR tightening; the ledger then *shrinks functorially* and cannot gain new types by Proposition 9.123 together

with the exclusion rule (Corollary 9.113). For the visibilities, cone shrink implies

$$\Lambda_{\bullet}^{\Sigma;[\chi]}(x)\Big|_{\ell'} \;\geq\; \Lambda_{\bullet}^{\Sigma;[\chi]}(x)\Big|_{\ell} \qquad (\bullet \in \{\mathrm{law}, \mathrm{state}\}),$$

by Lemma 9.136; thus they increase monotonically (bounded above by 1, Proposition 9.134(c)), i.e. move *towards unity* as the cone tightens. Finally, the meters vary continuously under the same IR change by Proposition 9.146(B).

Combining (1)–(4) gives the asserted continuity of the three maps in the proposition, the functorial ledger shrink under IR tightening, the monotone increase of visibilities toward 1 (less apparent breaking), and the continuity of the meter valuations. □

Corollary 9.157 (Typed quadrant and onset robustness)**.** *On certified plateaux,* $\mathrm{Quad}^{[\chi]}(x)$ *is locally constant up to tolerance thresholds (Corollary 9.139), and onset scales/times* $(r^\star[\chi], t^\star[\chi])$ *of Definition 9.147 exist under mild regularity (Lemma 9.148) and are stable under admissible retunings (Proposition 9.149).*

Proof of Corollary 9.157. Fix a permitted type χ and a certified plateau in the sense of Corollary 9.139, i.e. a neighborhood where the typed visibility pair $\vec{\Lambda}^{[\chi]}(x) := (\Lambda_{\mathrm{law}}^{\Sigma;[\chi]}(x), \Lambda_{\mathrm{state}}^{\Sigma;[\chi]}(x))$ stays away from the tolerance decision surfaces of the quadrant normalizer (Definition 9.47).

(i) Local constancy of the typed quadrant. By Corollary 9.139, on any such plateau the map $x \mapsto \mathrm{Quad}^{[\chi]}(x)$ (Definition 9.137) is locally constant, and it can change only at the declared tolerance thresholds of the normalizer (Definition 9.47). This proves the first claim.

(ii) Existence of onset scales/times. The onset components $\mathrm{Onset}^{[\chi]} = (r^\star[\chi], t^\star[\chi])$ are defined by the rules (9.81) in Definition 9.147, using the synthesis meters and the typed law–deficit. Under the mild regularity hypotheses stated in Lemma 9.148—namely, piecewise monotonicity of $r \mapsto v_{\mathrm{Synthesis}}^{(S);[\chi]}(r)$ and $r \mapsto v_{\mathrm{Synthesis}}^{(LS);[\chi]}(r)$ with one–sided limits, and a regularly varying envelope for $r \mapsto d_{\mathrm{law}}^{\Sigma;[\chi]}(r)$—the infima defining $r_{\mathrm{SSB}}^\star[\chi]$, $r_{\mathrm{QIV}}^\star[\chi]$, and $r_{\mathrm{QIII}}^\star[\chi]$ exist as extended nonnegative reals (finite exactly when the corresponding thresholds are eventually met). Hence the onsets are well-defined by Lemma 9.148.

(iii) Stability under admissible retunings. Within the retuning–stability radius of Definition 9.116, the typed visibilities and the meter valuations vary continuously with $(\lambda_{\mathrm{UV}}, \ell)$ by Proposition 9.156 (see also Propositions 9.134 and 9.146). Therefore the data entering (9.81) (the synthesis meters and the typed law–deficit) are continuous in the retuning parameters, and the resulting onsets are stable under these deformations as recorded in Proposition 9.149(c). In particular, on a certified plateau (where the corresponding decision margins are nonzero) small retunings cannot move the system across a tolerance surface, so neither the quadrant nor the onset labels change.

Combining (i)–(iii) gives the asserted robustness: $\mathrm{Quad}^{[\chi]}(x)$ is locally constant on plateaux up to tolerance thresholds (Corollary 9.139), and $(r^\star[\chi], t^\star[\chi])$ exist under the stated regularity (Lemma 9.148) and are stable under admissible retunings (Proposition 9.149). □

5.2 Artefact schema and completeness

Definition 9.158 (Execution artefacts per sector)**.** For each sector $\bullet$ and each point x, the artefacts of $\mathscr{E}_\bullet$ comprise:

$$\begin{aligned}\mathrm{Art}[\bullet](x) := \Big(&\mathcal{C}^{[\bullet]}_{\mathrm{admissible}}(x),\\ &\{\Lambda^{\Sigma;[\chi]}_{\mathrm{law}}(x), \Lambda^{\Sigma;[\chi]}_{\mathrm{state}}(x)\}_{\chi\in\mathcal{C}^{[\bullet]}_{\mathrm{admissible}}},\\ &\{v^{(a);[\chi]}_{\mathrm{Synthesis}}(x), v^{(LS);[\chi]}_{\mathrm{Synthesis}}(x)\}_{\chi},\\ &\{\mathrm{TLO}^{[\chi]}(x;r)\}_{\chi,r}\Big).\end{aligned}$$

Proposition 9.159 (Typed completeness w.r.t. ledger)**.** *For every $\chi \in \mathcal{C}^{[\bullet]}_{\mathrm{admissible}}(x)$, the pipeline $\mathscr{E}_\bullet$ computes all quantities needed to evaluate $\mathrm{TLO}^{[\chi]}(x;r)$ (Type via Definition 9.137, Location as in Definition 9.147, and Onset via (9.81)). Conversely, if $\chi \notin \mathcal{C}^{[\bullet]}_{\mathrm{admissible}}$, then by Corollary 9.113 typed hyperbolicity fails and $\mathrm{TLO}^{[\chi]}$ is undefined (and must not be reported).*

Proof of Proposition 9.159. Fix a block $\bullet \in \{\Phi\Phi, \Theta\Theta\}$ and a spacetime point x.

(I) Completeness for $\chi \in \mathcal{C}^{[\bullet]}_{\mathrm{admissible}}(x)$. If χ is admitted by the ledger, then by the admission criterion of Proposition 9.102 the associated typed projector $\Pi^{[\chi]}$ is band-finite, graded-covariant, cone-compatible (Definition 9.98) and satisfies typed hyperbolicity (9.42), with stability under retunings (Definition 9.116). Constructing $\Pi^{[\chi]}$ via the Σ-weighted spectral machinery is guaranteed by Proposition 9.119, and the execution stage (E1) of the pipeline $\mathscr{E}_\bullet$ terminates and is well-posed by Lemma 9.154.

Given $\Pi^{[\chi]}$, the pipeline then computes:

- *Type.* Pointwise typed contrasts and their aggregates are computed by Definitions 9.127 and 9.133. The typed quadrant is then

$$\mathrm{Quad}^{[\chi]}(x) = \mathrm{Quad}\Big(\Lambda^{\Sigma;[\chi]}_{\mathrm{law}}(x),\ \Lambda^{\Sigma;[\chi]}_{\mathrm{state}}(x)\Big) \quad \text{by Definition 9.137.}$$

  Thus the Type component is obtained canonically from the ledger-admitted χ.
- *Location.* The Location tags (arena $\mathfrak{P}/\mathfrak{P}_\partial$, sector $\bullet$, spatial tag $P_x(\lambda_{\mathrm{UV}})$, cone sheet) are read off as specified in Definition 9.147, using the BRST-closed band instrumentation (Definition 9.96) and the Σ-cone gate (Definitions 9.104 and 9.106). Hence Location is computable for the same χ.
- *Onset.* The band-reduced typed meters $v^{(a);[\chi]}_{\mathrm{Synthesis}}$, $v^{(LS);[\chi]}_{\mathrm{Synthesis}}$ (Definitions 9.141, 9.143 and 9.144) together with the typed law–deficit induced by Definition 9.127 feed the onset rules (9.81) in Definition 9.147, thereby producing $(r^\star[\chi], t^\star[\chi])$. Under the mild regularity hypotheses, existence of these infima is ensured by Lemma 9.148. (Gauge/causal robustness of all ingredients is addressed in Propositions 9.134 and 9.149 and Lemma 9.145.)

Therefore, for every $\chi \in \mathcal{C}^{[\bullet]}_{\mathrm{admissible}}(x)$, the pipeline $\mathscr{E}_\bullet$ computes all quantities needed for $\mathrm{TLO}^{[\chi]}(x;r)$ (Type via Definition 9.137, Location via Definition 9.147, and Onset via (9.81)).

(II) Non-admitted types carry no T–L–O. Conversely, if $\chi \notin \mathcal{C}^{[\bullet]}_{\text{admissible}}(x)$, then by the exclusion rule Corollary 9.113 the typed hyperbolicity condition (9.53) fails for some timelike covector, so the well-posedness implication of Proposition 9.112 does *not* hold for χ. In particular, the typed energy estimate and retarded Green operator required to define the cone-gated, retarded operands of Definition 9.127 are not available. Hence the downstream typed aggregates/visibilities (Definition 9.133), meters (Definitions 9.141, 9.143 and 9.144), and the onset rules (9.81) cannot be soundly instantiated. In this case $\mathrm{TLO}^{[\chi]}$ is *undefined* and, by the ledger discipline of Proposition 9.102, must not be reported.

Combining (I) and (II) proves the stated completeness with respect to the ledger. □

Corollary 9.160 (Bulk/edge relation)**.** *Let* $\mathrm{Art}[\bullet]_{\mathrm{bulk}}$ *and* $\mathrm{Art}[\bullet]_{\partial}$ *denote outputs on* $\mathfrak{P}$ *and* $\mathfrak{P}_{\partial}$*, respectively. Then*

$$\mathcal{C}^{[\bullet]}_{\text{admissible}}(\mathfrak{P}_{\partial}) \;\subseteq\; \mathcal{C}^{[\bullet]}_{\text{admissible}}(\mathfrak{P}), \qquad \Lambda^{\Sigma;[\chi]}_{\bullet'}(x)\Big|_{\mathfrak{P}_{\partial}} \;\geq\; \Lambda^{\Sigma;[\chi]}_{\bullet'}(x)\Big|_{\mathfrak{P}},$$

for $\bullet' \in \{\mathrm{law}, \mathrm{state}\}$*, by cone shrink on the boundary (cf. Lemmas 9.105 and 9.136) and functoriality (Proposition 9.123).*

Proof of Corollary 9.160. Let $\mathfrak{P}$ (bulk) and $\mathfrak{P}_{\partial}$ (edge) be evaluated with their respective band instrumentations (Definition 9.96), writing

$$\mathfrak{B}^{\mathrm{bulk}}_{\lambda_{\mathrm{UV}},\ell} = \Big(P_x(\lambda_{\mathrm{UV}}), K^{(\ell)}_{\mathrm{IR}}\Big), \qquad \mathfrak{B}^{\partial}_{\lambda_{\mathrm{UV}},\ell'} = \Big(P_x(\lambda_{\mathrm{UV}})P_{\partial},\; K^{(\ell')}_{\mathrm{IR}}\Big),$$

where P_{∂} implements the boundary restriction and $\ell' \leq \ell$ denotes the tighter IR window at the edge. By Lemma 9.105, tightening the IR window yields a *cone shrink*

$$\mathcal{C}^{\hat{\Sigma},*}_{x}[\mathfrak{P}_{\partial}] \;\subseteq\; \mathcal{C}^{\hat{\Sigma},*}_{x}[\mathfrak{P}],$$

and the associated cone gates (retarded projectors of Definition 9.106) satisfy the Löwner order $0 \leq \Gamma_{\partial} \leq \Gamma_{\mathrm{bulk}} \leq \mathbf{1}$ (Proposition 9.107).

(1) Ledger inclusion. Passing from $\mathfrak{P}$ to $\mathfrak{P}_{\partial}$ is a *functorial refinement* of the band tags (additional P_{∂} and smaller ℓ'). By Proposition 9.123, such refinements *shrink* the ledger and cannot add a type that previously failed the typed hyperbolicity test (9.53) (see also the exclusion rule Corollary 9.113). Hence

$$\mathcal{C}^{[\bullet]}_{\text{admissible}}(\mathfrak{P}_{\partial}) \;\subseteq\; \mathcal{C}^{[\bullet]}_{\text{admissible}}(\mathfrak{P}).$$

(2) Visibility monotonicity at the edge. For any admitted type χ and fixed probe set $(\mathcal{V}, w)$, cone shrink implies that the *aggregated* contrasts weakly decrease while the normalized visibilities weakly increase, by Lemma 9.136:

$$\Lambda^{\Sigma;[\chi]}_{\bullet'}(x)\Big|_{\mathfrak{P}_{\partial}} \;\geq\; \Lambda^{\Sigma;[\chi]}_{\bullet'}(x)\Big|_{\mathfrak{P}}, \qquad \bullet' \in \{\mathrm{law}, \mathrm{state}\}.$$

Intuitively, restricting causal support cannot increase measured breaking; algebraically, this is the action of the positive contraction $\Gamma_{\partial} \leq \Gamma_{\mathrm{bulk}}$ inside the band functionals (Definitions 9.127 and 9.133) together with the nonincreasing normalizer (Definition 9.47).

Combining (1) and (2) yields the stated bulk/edge relations for the ledger and the visibilities. □

5.3 Compact execution formula

Theorem 9.161 (Compact execution formula)**.** *For each sector* $\bullet$ *and each permitted* χ*, the output of* $\mathcal{E}_\bullet$ *at* $(x;r)$ *may be written as*

$$\mathrm{TLO}^{[\chi]}(x;r) \;=\; \Big(\; \mathrm{Quad}\Big(\Lambda_{\mathrm{law}}^{\Sigma;[\chi]}(x), \Lambda_{\mathrm{state}}^{\Sigma;[\chi]}(x)\Big)\,,\; (\mathcal{A}_\chi, \mathrm{sector}_\chi, P_x, \mathrm{sheet}_\chi)\,,\; (r^\star[\chi], t^\star[\chi])\;\Big),$$

with $\Lambda_\bullet^{\Sigma;[\chi]}$ *given by* (9.67)*,* Quad *by* (9.75)*, and onsets by* (9.81)*. The right-hand side depends only on BRST cohomology classes and on the typed projector* $\Pi^{[\chi]}$ *(gauge covariance Proposition 9.130, BRST invariance Proposition 9.97), and is causally supported inside* $J^+(\mathcal{C}^{\hat\Sigma,*})$ *(Proposition 9.107).*

Proof of Theorem 9.161. Fix a sector $\bullet \in \{\Phi\Phi, \Theta\Theta\}$ and a permitted type $\chi \in \mathcal{C}^{[\bullet]}_{\mathrm{admissible}}$ (ledger admission in Proposition 9.102). Let x be a spacetime point and r a scale (band radius).

Step 1 (Representation by definitions; well-posedness). Given $\Pi^{[\chi]}$, the *typed visibilities* are defined by

$$\Lambda_\bullet^{\Sigma;[\chi]}(x) = \mathfrak{n}_\bullet\Big(\Delta_\bullet^{\mathrm{pre},\Sigma;[\chi]}(x;\mathcal{V},w)\Big), \qquad \bullet \in \{\mathrm{law}, \mathrm{state}\},$$

with $\Delta_\bullet^{\mathrm{pre},\Sigma;[\chi]}$ the aggregated typed contrasts and $\mathfrak{n}_\bullet$ the normalizer (Definitions 9.127 and 9.133; this is Eq. (9.67)). The *typed quadrant* is

$$\mathrm{Quad}^{[\chi]}(x) = \mathrm{Quad}\Big(\Lambda_{\mathrm{law}}^{\Sigma;[\chi]}(x),\; \Lambda_{\mathrm{state}}^{\Sigma;[\chi]}(x)\Big) \quad \text{(Definition 9.137, Eq. (9.75))}.$$

The *Location* tuple $(\mathcal{A}_\chi, \mathrm{sector}_\chi, P_x, \mathrm{sheet}_\chi)$ is read from the arena/sector/band tag and cone sheet (Definition 9.147). The *Onset* pair $(r^\star[\chi], t^\star[\chi])$ is defined by the rules (9.81) using the typed meters (Definitions 9.141, 9.143 and 9.144) and the typed law–deficit induced from Definition 9.127. Thus, by *definition*,

$$\mathrm{TLO}^{[\chi]}(x;r) = \Big(\mathrm{Quad}^{[\chi]}(x),\; (\mathcal{A}_\chi, \mathrm{sector}_\chi, P_x, \mathrm{sheet}_\chi),\; (r^\star[\chi], t^\star[\chi])\Big),$$

which is exactly the displayed right-hand side of the theorem with the cited equations. The fact that each ingredient is *well-posed* for admitted χ follows from Propositions 9.112 and 9.134 and Lemmas 9.145 and 9.148 (see also the termination/well-posedness pipeline statement in Lemma 9.154) and the typed completeness Proposition 9.159.

Step 2 (BRST/Nielsen safety ⇒ dependence only on BRST classes). Pointwise typed contrasts are invariant under BRST-exact gauge variations by Lemma 9.128, based on the band Nielsen property Proposition 9.97 and the BRST-closedness $[\hat{Q}_{\mathrm{BRST}}, \Pi^{[\chi]}] = 0$ (Definition 9.116). Aggregation/normalization preserve invariance (Proposition 9.134(a)), hence the visibilities—and therefore the quadrant—depend only on BRST cohomology classes of the underlying band operators. The typed meters are likewise BRST/Nielsen-safe (Lemma 9.145(b)), so the onset rules (9.81) are unchanged by BRST-exact variations. In particular, any change $A \mapsto A + [\hat{Q}_{\mathrm{BRST}}, X]$ in the band algebra leaves all numerical outputs unchanged (Proposition 9.97), proving the stated *dependence only on BRST classes.* Gauge

covariance under band-admissible graded frames is recorded in Proposition 9.130 (see also ledger covariance Proposition 9.123).

Step 3 (Causal support inside $J^+(\mathcal{C}^{\hat{\Sigma},*})$). All kernels entering the typed contrasts and the band meters are replaced by their *retarded, cone-gated* versions (Definition 9.106), which are supported inside $J^+\big(\mathcal{C}^{\hat{\Sigma},*}\big)$ by Proposition 9.107; microcausality at spacelike separation holds by Proposition 9.108. Consequently, the visibilities are causally well-posed (Proposition 9.134(b)), the meter valuations inherit the same causal support (Lemma 9.145(c)), and the resulting $\mathrm{TLO}^{[\chi]}(x;r)$ is causally robust (Proposition 9.149(b); see also the aggregate soundness statement Proposition 9.155).

Step 4 (Dependence on $\Pi^{[\chi]}$ only, up to conjugation). By construction, the typed restriction is effected solely by $\Pi^{[\chi]}$ (Definition 9.116); hence all typed quantities depend on the chosen projector. Under a graded local frame g, $\Pi^{[\chi]}$ transforms as $g\Pi^{[\chi]}g^\dagger$ and the ledger is invariant up to this equivalence (Proposition 9.123); the numerical outputs remain the same by Proposition 9.130. Therefore the right-hand side in the statement depends only on the typed projector (modulo gauge conjugation) and the BRST classes of the operators it acts upon.

Combining Steps 1–4 yields the compact execution formula and proves that the right-hand side is BRST/gauge correct and causally supported in $J^+(\mathcal{C}^{\hat{\Sigma},*})$, as stated. □

Remark 9.162 (Minimal inputs for a runnable instance)**.** To run $\mathscr{E}_\bullet$ one must specify: (i) band tag $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ (Definition 9.96); (ii) a probe family $(\mathcal{V}, w)$ (Definition 9.131); and (iii) a spectral criterion for $\Pi^{[\chi]}$ (e.g. gap thresholds in Proposition 9.119). All subsequent artefacts then follow by Definitions 9.133, 9.143, 9.144 and 9.147.

9.14 Audits & Guardrails

We formalize the audit predicates and their guarantees for the typed, band-level pipeline developed in §1–§5. Throughout, kernels are retarded and cone-gated (Definition 9.106 and Proposition 9.107), band taggers are BRST-closed with ghost number 0 (Definition 9.96 and Proposition 9.97), and typed objects are restricted by $\Pi^{[\chi]}$ (Definition 9.116).

6.1 Audit predicates

Definition 9.163 (BRST/Nielsen audit)**.** We say the *BRST/Nielsen audit* passes for an instrument set $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell} = (P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}^{(\ell)})$ if

$$\begin{aligned} [\,\hat{Q}_{\mathrm{BRST}}, P_x(\lambda_{\mathrm{UV}})\,] = [\,\hat{Q}_{\mathrm{BRST}}, K_{\mathrm{IR}}^{(\ell)}\,] = 0, \quad \mathrm{gh} = 0, \\ \text{and} \quad \mathcal{I}([\,\hat{Q}_{\mathrm{BRST}}, X]) = \mathcal{I}_\Psi([\,\hat{Q}_{\mathrm{BRST}}, X]) = 0. \end{aligned} \tag{9.82}$$

for all band-admissible X (Definitions 9.45 and 9.96 and Proposition 9.97).

Definition 9.164 (Cone/microcausality audit)**.** We say the *cone/microcausality audit* passes for a kernel family $\{\mathcal{K}\}$ if each kernel is restricted to the retarded, cone-gated form $\mathcal{K}^{\mathrm{ret}}$ (Definition 9.106), and for all test functions f, g with spacelike-separated supports w.r.t. $\mathcal{C}^{\hat{\Sigma},*}$,

$$[\,\hat{\mathbf{\Phi}}(f), \hat{\mathbf{\Phi}}(g)\,] = [\,\hat{\mathbf{\Theta}}(f), \hat{\mathbf{\Theta}}(g)\,] = 0 \qquad \text{(Proposition 9.108).} \tag{9.83}$$

Definition 9.165 (Typed hyperbolicity audit)**.** Given a typed projector $\Pi^{[\chi]}$, the *typed hyperbolicity audit* passes at x if

$$\inf_{\Psi\in\mathcal{S}_{\mathrm{band}}} \mathcal{I}_\Psi\Big[\ \hat{Q}_x^{[\chi]}[k]\ \Big] \ >\ 0 \quad \text{for all timelike } k\in \mathcal{C}_x^{\hat{\Sigma},*}, \tag{9.84}$$

i.e. (9.84) coincides with (9.53) (Definition 9.111).

Definition 9.166 (Band-stability audit)**.** The *band-stability audit* passes for $(\Pi^{[\chi]}, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell})$ if there exists $\delta>0$ such that for all $(\lambda'_{\mathrm{UV}},\ell')$ with $|\lambda'_{\mathrm{UV}}-\lambda_{\mathrm{UV}}|+|\ell'-\ell|<\delta$,

$$\|\Pi^{[\chi]}(\lambda'_{\mathrm{UV}},\ell')-\Pi^{[\chi]}(\lambda_{\mathrm{UV}},\ell)\|\to 0 \quad\text{and}\quad \Delta_\bullet^{\mathrm{pre},\Sigma;[\chi]}(x;\mathcal{V},w),\ v^{(\cdot);[\chi]}_{\mathrm{Synthesis}}(x) \text{ vary continuously,}$$

as in (P7) of Definition 9.116 and Lemma 9.132 and Propositions 9.134 and 9.146.

Definition 9.167 (Functoriality audit)**.** Write $(P',K')\preccurlyeq(P,K)$ if $P'=P_x(\lambda'_{\mathrm{UV}})$ with $\lambda'_{\mathrm{UV}}\le\lambda_{\mathrm{UV}}$ and $K'=K_{\mathrm{IR}}^{(\ell')}$ with $\ell'\le\ell$. The *functoriality audit* passes if for every such refinement:

$$\begin{aligned} \mathcal{C}_x^{\hat{\Sigma},*}\big|_{(P',K')} &\subseteq \mathcal{C}_x^{\hat{\Sigma},*}\big|_{(P,K)} \quad \text{(Lemma 9.105)},\\ \Delta_\bullet^{\mathrm{pre},\Sigma;[\chi]}\downarrow,\quad \Lambda_\bullet^{\Sigma;[\chi]}\uparrow,&\quad \mathrm{Quad}^{[\chi]} \text{ unchanged except at tolerances,} \end{aligned} \tag{9.85}$$

with "unchanged" interpreted via Corollary 9.139.

6.2 Effective visibilities and uncertainty bands

Definition 9.168 (Effective visibility and uncertainty band)**.** Fix a neighborhood $\mathcal{U}_\delta := \{(\lambda'_{\mathrm{UV}},\ell'):\ |\lambda'_{\mathrm{UV}}-\lambda_{\mathrm{UV}}|+|\ell'-\ell|<\delta\}$. For any typed χ define the *effective visibility* and its *band* by

$$\Lambda_{\bullet,\mathrm{eff}}^{\Sigma;[\chi]}(x;\delta):=\inf_{(\lambda'_{\mathrm{UV}},\ell')\in\mathcal{U}_\delta}\Lambda_\bullet^{\Sigma;[\chi]}(x)\Big|_{(\lambda'_{\mathrm{UV}},\ell')},\qquad \mathcal{B}_\bullet^{[\chi]}(x;\delta):=\Big[\ \Lambda_{\bullet,\mathrm{eff}}^{\Sigma;[\chi]}(x;\delta),\ \sup_{(\lambda'_{\mathrm{UV}},\ell')\in\mathcal{U}_\delta}\Lambda_\bullet^{\Sigma;[\chi]}(x)\Big], \tag{9.86}$$

for $\bullet\in\{\mathrm{law},\mathrm{state}\}$ (Definition 9.133).

Lemma 9.169 (Bounds and monotonicity of Λ_{eff})**.** *For each x,χ and $\delta'>\delta\ge 0$,*

$$\begin{aligned} 0\le\Lambda_{\bullet,\mathrm{eff}}^{\Sigma;[\chi]}(x;\delta')&\le\Lambda_{\bullet,\mathrm{eff}}^{\Sigma;[\chi]}(x;\delta)\\ &\le\Lambda_\bullet^{\Sigma;[\chi]}(x)\le 1,\\ \mathcal{B}_\bullet^{[\chi]}(x;\delta')&\supseteq\mathcal{B}_\bullet^{[\chi]}(x;\delta), \end{aligned}$$

and $\delta\mapsto\Lambda_{\bullet,\mathrm{eff}}^{\Sigma;[\chi]}(x;\delta)$ is right-continuous by Proposition 9.134 and Definition 9.166.

Proof of Lemma 9.169. Fix x and an admitted type χ. By the audit/stability convention (Definition 9.166), for each tolerance $\delta\ge 0$ we have a nonempty, compact, nested family of admissible retunings $\mathcal{B}_\bullet^{[\chi]}(x;\delta)$ in the band-parameter space (it contains the nominal point,

and $\delta' > \delta \;\Rightarrow\; \mathcal{B}_\bullet^{[\chi]}(x;\delta') \supseteq \mathcal{B}_\bullet^{[\chi]}(x;\delta)$). The *effective* visibility is the worst-case (infimum) of the typed visibility over the audit set:

$$\Lambda_{\bullet,\mathrm{eff}}^{\Sigma;[\chi]}(x;\delta) \;:=\; \inf_{b\in\mathcal{B}_\bullet^{[\chi]}(x;\delta)} \Lambda_\bullet^{\Sigma;[\chi]}(x;b), \qquad \bullet\in\{\mathrm{law},\mathrm{state}\}, \tag{9.87}$$

where b denotes the band retuning parameters and $\Lambda_\bullet^{\Sigma;[\chi]}(x;b)$ is the typed visibility computed at b (Definition 9.133, Eq. (9.67)).

Bounds $0 \le \Lambda_{\bullet,\mathrm{eff}}^{\Sigma;[\chi]} \le \Lambda_\bullet^{\Sigma;[\chi]} \le 1$. By normalization, for any fixed band point b, $0 \le \Lambda_\bullet^{\Sigma;[\chi]}(x;b) \le 1$ (Proposition 9.134(c)); taking an infimum preserves these bounds, hence $0 \le \Lambda_{\bullet,\mathrm{eff}}^{\Sigma;[\chi]}(x;\delta) \le 1$. Since the nominal band point b_0 (no retuning) lies in every $\mathcal{B}_\bullet^{[\chi]}(x;\delta)$ (Definition 9.166),

$$\Lambda_{\bullet,\mathrm{eff}}^{\Sigma;[\chi]}(x;\delta) = \inf_{b\in\mathcal{B}_\bullet^{[\chi]}(x;\delta)} \Lambda_\bullet^{\Sigma;[\chi]}(x;b) \;\le\; \Lambda_\bullet^{\Sigma;[\chi]}(x;b_0) = \Lambda_\bullet^{\Sigma;[\chi]}(x),$$

which yields the middle inequality.

Monotonicity in δ and set inclusion. If $\delta' > \delta$, then by Definition 9.166 the audit sets are nested: $\mathcal{B}_\bullet^{[\chi]}(x;\delta') \supseteq \mathcal{B}_\bullet^{[\chi]}(x;\delta)$. Taking infima over a larger set can only decrease (or leave unchanged) the value, whence

$$\Lambda_{\bullet,\mathrm{eff}}^{\Sigma;[\chi]}(x;\delta') \;\le\; \Lambda_{\bullet,\mathrm{eff}}^{\Sigma;[\chi]}(x;\delta), \qquad \mathcal{B}_\bullet^{[\chi]}(x;\delta') \supseteq \mathcal{B}_\bullet^{[\chi]}(x;\delta).$$

Right-continuity in δ. By Proposition 9.156 (see also Proposition 9.134), $b \mapsto \Lambda_\bullet^{\Sigma;[\chi]}(x;b)$ is continuous on the band-parameter space; by Definition 9.166, for each δ the set $\mathcal{B}_\bullet^{[\chi]}(x;\delta)$ is compact and the family is increasing in δ. Define $F(\delta) := \Lambda_{\bullet,\mathrm{eff}}^{\Sigma;[\chi]}(x;\delta)$. Fix $\delta_0 \ge 0$. Since F is monotone nonincreasing, $\lim_{\delta\downarrow\delta_0^+} F(\delta)$ exists. To show $F(\delta_0^+) = F(\delta_0)$, let $\varepsilon > 0$ and pick $b_\varepsilon \in \mathcal{B}_\bullet^{[\chi]}(x;\delta_0)$ with $\Lambda_\bullet^{\Sigma;[\chi]}(x;b_\varepsilon) \le F(\delta_0) + \varepsilon$ (attainment holds by compactness; otherwise approximate). By continuity there is a neighborhood U of b_ε on which $\Lambda_\bullet^{\Sigma;[\chi]}(x;\cdot) \le F(\delta_0) + 2\varepsilon$. Since $\{\mathcal{B}(\delta)\}_{\delta>\delta_0}$ increases to a neighborhood of $\mathcal{B}(\delta_0)$, for all $\delta > \delta_0$ sufficiently close we have $U \subseteq \mathcal{B}_\bullet^{[\chi]}(x;\delta)$; hence

$$F(\delta) \;=\; \inf_{b\in\mathcal{B}(\delta)} \Lambda_\bullet^{\Sigma;[\chi]}(x;b) \;\le\; \Lambda_\bullet^{\Sigma;[\chi]}(x;b_\varepsilon) \;+\; 2\varepsilon \;\le\; F(\delta_0) \;+\; 3\varepsilon.$$

Letting $\varepsilon \downarrow 0$ gives $\limsup_{\delta\downarrow\delta_0^+} F(\delta) \le F(\delta_0)$, while monotonicity yields the reverse inequality $F(\delta) \ge F(\delta_0)$ for $\delta > \delta_0$. Hence F is right-continuous at δ_0. This proves the stated right-continuity, relying on the bounds/normalization of Proposition 9.134 and the audit-band properties of Definition 9.166.

Combining the three parts yields the displayed chain of inequalities and the inclusion of audit sets, as well as the right-continuity of $\delta \mapsto \Lambda_{\bullet,\mathrm{eff}}^{\Sigma;[\chi]}(x;\delta)$. □

Proposition 9.170 (Quadrant robustness under uncertainty)**.** *Let* $\mathrm{Quad}^{[\chi]}(x)$ *be assigned by* (9.75)*. If there exists* $\delta > 0$ *such that the rectangle* $\mathcal{R}(x;\delta) := \mathcal{B}_{\mathrm{law}}^{[\chi]}(x;\delta) \times \mathcal{B}_{\mathrm{state}}^{[\chi]}(x;\delta)$ *lies entirely within a single quadrant of the decision rule (Definition 7.685), then* $\mathrm{Quad}^{[\chi]}(x)$ *is invariant under all retunings in* $\mathcal{U}_\delta$*.*

Proof of Proposition 9.170. Fix an admitted type χ at x. By the audit/stability convention (Definition 9.166) there is, for each $\delta > 0$, an admissible retuning set $\mathcal{U}_\delta$ in band-parameter space and the associated *visibility bands*

$$\mathcal{B}_\bullet^{[\chi]}(x;\delta) \;:=\; \Big\{\, \Lambda_\bullet^{\Sigma;[\chi]}(x;b) \;:\; b \in \mathcal{U}_\delta \,\Big\} \subset [0,1], \qquad \bullet \in \{\mathrm{law}, \mathrm{state}\},$$

with $\Lambda_\bullet^{\Sigma;[\chi]}$ the typed visibilities of Definition 9.133 (cf. normalization and invariance in Proposition 9.134). Define the *audit rectangle* $\mathcal{R}(x;\delta) := \mathcal{B}_{\mathrm{law}}^{[\chi]}(x;\delta) \times \mathcal{B}_{\mathrm{state}}^{[\chi]}(x;\delta) \subset [0,1]^2$.

By hypothesis, there exists $\delta > 0$ such that the rectangle $\mathcal{R}(x;\delta)$ lies entirely within a single quadrant of the canonical decision rule (Definition 7.685). For any retuning $b \in \mathcal{U}_\delta$ we have

$$\Big(\Lambda_{\mathrm{law}}^{\Sigma;[\chi]}(x;b),\; \Lambda_{\mathrm{state}}^{\Sigma;[\chi]}(x;b)\Big) \;\in\; \mathcal{R}(x;\delta),$$

hence this pair belongs to the *same* quadrant region. By the typed quadrant predicate (Definition 9.137, Eq. (9.75)),

$$\mathrm{Quad}^{[\chi]}(x;b) = \mathrm{Quad}\Big(\Lambda_{\mathrm{law}}^{\Sigma;[\chi]}(x;b),\; \Lambda_{\mathrm{state}}^{\Sigma;[\chi]}(x;b)\Big)$$

is therefore constant for all $b \in \mathcal{U}_\delta$, i.e. invariant under all admissible retunings in $\mathcal{U}_\delta$.

Finally, note that the visibilities are continuous in the retuning parameters (Proposition 9.156) and remain in $[0,1]$ (Proposition 9.134(c)), so no crossing of quadrant decision surfaces can occur as long as the image stays in $\mathcal{R}(x;\delta)$; this is the same robustness mechanism underlying the plateau stability of Corollary 9.139. The proposition follows. □

6.3 Global guardrail theorem

Theorem 9.171 (Global guardrails for the typed band pipeline)**.** *Suppose the audits of Definitions 9.163 to 9.167 pass for a sector $\bullet$ and a permitted type χ. Then:*

(i) BRST/Nielsen safety
Every scalar diagnostic among $\Delta_\bullet^{\mathrm{pre},\Sigma;[\chi]}$, $\Lambda_\bullet^{\Sigma;[\chi]}$, $v_{\mathrm{Synthesis}}^{(\cdot);[\chi]}$, and $\mathrm{TLO}^{[\chi]}$ is gauge-parameter independent (Proposition 9.97 and Lemmas 9.128 and 9.145).

(ii) Causal correctness
All commutators vanish for spacelike separation and all kernels are supported in $J^+(\mathcal{C}^{\hat{\Sigma},*})$ (Propositions 9.107 and 9.108).

(iii) Typed well-posedness
The principal form is strictly positive on timelike covectors after restriction by $\Pi^{[\chi]}$ (Definition 9.165); hence the typed sector is hyperbolic and admits retarded Green operators (Proposition 9.112).

(iv) Band stability with uncertainty bands
Outputs vary continuously under small retunings and admit certified intervals $\mathcal{B}_\bullet^{[\chi]}$ (Definition 9.168 and Lemma 9.169).

(iv) Functorial sharpening
Refining $(P_x, K_{\mathrm{IR}}^{(\ell)})$ decreases $\Delta_\bullet^{\mathrm{pre},\Sigma;[\chi]}$, increases $\Lambda_\bullet^{\Sigma;[\chi]}$, and preserves quadrant membership off tolerance boundaries (Definition 9.167 and Corollary 9.139).

Proof of Theorem 9.171. Fix a sector $\bullet \in \{\Phi\Phi, \Theta\Theta\}$ and a type χ admitted by the ledger. Assume the audits of Definitions 9.163 to 9.167 all pass.

(a) BRST/Nielsen safety. By the BRST audit Definition 9.163, the band taggers, typed projectors, and kernels are ghost-number 0 and BRST-closed. Pointwise typed contrasts are invariant under BRST-exact gauge variations by Lemma 9.128, using the band Nielsen property Proposition 9.97 and $[\hat{Q}_{\mathrm{BRST}}, \Pi^{[\chi]}] = 0$ (Definition 9.116). Aggregation/normalization preserves invariance, yielding gauge-invariant visibilities (Proposition 9.134(a)). Typed meter valuations are likewise BRST/Nielsen-safe by Lemma 9.145(b). The T–L–O assignment inherits gauge/BRST independence by Proposition 9.149(a). Hence every listed scalar diagnostic is gauge-parameter independent.

(b) Causal correctness. By the cone/causality audit Definition 9.164, all kernels are replaced by their retarded, cone-gated versions (Definition 9.106), which are supported in $J^+\big(\mathcal{C}^{\hat{\Sigma},*}\big)$ (Proposition 9.107), and microcausality holds at spacelike separation (Proposition 9.108). Consequently, typed contrasts are real and well-defined (Lemma 9.129), visibilities are causally well-posed (Proposition 9.134(b)), meters inherit the same causal support (Lemma 9.145(c)), and T–L–O is causally robust (Proposition 9.149(b)).

(c) Typed well-posedness. The hyperbolicity audit Definition 9.165 asserts positivity of the typed principal form on cone-timelike covectors (condition (9.53)); therefore the typed sector is hyperbolic and admits a retarded Green operator supported in $J^+\big(\mathcal{C}^{\hat{\Sigma},*}\big)$ by Proposition 9.112. (If positivity fails, the exclusion rule Corollary 9.113 removes χ from the ledger, ensuring soundness of the pipeline.)

(d) Band stability with uncertainty bands. Within the retuning-stability radius (audit Definition 9.166), the ledger is locally constant and the typed visibilities are continuous in $(\lambda_{\mathrm{UV}}, \ell)$ (Proposition 9.156; see also Proposition 9.134); typed meters depend continuously as well (Proposition 9.146(B)). The effective (audited) visibilities $\Lambda_{\bullet,\mathrm{eff}}^{\Sigma;[\chi]}$ and the band families $\mathcal{B}_{\bullet}^{[\chi]}$ are defined in Definition 9.168, and satisfy bounds, monotonicity, and right-continuity by Lemma 9.169. Therefore outputs admit certified uncertainty intervals and vary continuously under small retunings.

(e) Functorial sharpening. Refining $(P_x, K_{\mathrm{IR}}^{(\ell)})$ as stipulated by the functoriality audit Definition 9.167 *shrinks* the Σ–cone by Lemma 9.105. Hence aggregated typed contrasts weakly decrease while visibilities weakly increase by Lemma 9.136 (cf. normalization in Proposition 9.134(c)). Plateau stability Corollary 9.139 then guarantees that quadrant membership is preserved away from tolerance boundaries; equivalently, audit rectangles entirely within a quadrant ensure robustness by Proposition 9.170. Ledger functoriality (no new types appear under refinement) is ensured by Proposition 9.123 together with Corollary 9.113.

Combining (a)–(e) yields the stated global guardrails for the typed band pipeline. □

6.4 Certificates and reporting

Definition 9.172 (Audit certificate for a typed sector)**.** For each (x, χ) we define the *audit certificate*

$$\mathfrak{C}^{[\chi]}(x) := \Big(\underbrace{\text{BRST/Nielsen}}_{(9.82)}, \underbrace{\text{Cone/Micro}}_{(9.83)}, \underbrace{\text{Hyperbolicity}}_{(9.84)}, \underbrace{\text{Stability}}_{\delta \text{ of Definition 9.166}}, \underbrace{\text{Functoriality}}_{(9.85)}, \underbrace{\mathcal{B}^{[\chi]}_{\text{law}}(x;\delta) \times \mathcal{B}^{[\chi]}_{\text{state}}(x;\delta)}_{\text{uncertainty rectangle}} \Big),$$

together with the decision $\mathrm{Quad}^{[\chi]}(x)$ (Definition 9.137).

Corollary 9.173 (Minimal certificate for publication)**.** *A minimal, publishable report for (x, χ) consists of: (i) the cone/hyperbolicity certificate triple (admissible timelike interval, spectral gap, stability radius) from Corollary 9.120; (ii) the uncertainty rectangles of Definition 9.172; and (iii) the quadrant and T–L–O outputs (Definition 9.147). By Theorem 9.171, these are sufficient to verify BRST safety, causal correctness, and stability.*

Proof of Corollary 9.173. Fix (x, χ) with $\chi \in \mathcal{C}^{[\bullet]}_{\text{admissible}}(x)$ (Proposition 9.102). The proposed certificate consists of:

(C1) the cone/hyperbolicity *certificate triple* (admissible timelike interval in $\mathcal{C}_x^{\hat{\Sigma},*}$, spectral gap for the Σ–weighted channel, stability radius) from Corollary 9.120;

(C2) the *uncertainty rectangles* (audit bands) of Definition 9.172 for the visibility pair $\vec{\Lambda}^{[\chi]}(x)$;

(C3) the *quadrant* and full *T–L–O* outputs as in Definition 9.147.

We show that (C1)–(C3) suffice to verify the three global properties stated in Theorem 9.171.

Typed well-posedness and admission. By Corollary 9.120, (C1) guarantees a *positive* typed principal form on the admissible timelike interval together with a *nonzero spectral gap* for the Σ–weighted channel operator and a *retuning-stability radius*. Hence the typed sector χ is *hyperbolic* and admits a retarded Green operator supported in $J^+\big(\mathcal{C}^{\hat{\Sigma},*}\big)$ by Proposition 9.112. This ensures the typed constructions (contrasts $\to$ visibilities $\to$ quadrant, meters $\to$ onsets) are well-posed, as required in (c) of Theorem 9.171. Moreover, because the projectors are produced by the Σ–spectral construction (Proposition 9.119), they are BRST-closed; this feeds into the BRST audit below.

Band stability and functorial behavior. Item (C2) is precisely the uncertainty data of Definition 9.172, encoding the audit sets $\mathcal{U}_\delta$ and the visibility rectangles $\mathcal{R}(x;\delta) = \mathcal{B}^{[\chi]}_{\mathrm{law}}(x;\delta) \times \mathcal{B}^{[\chi]}_{\mathrm{state}}(x;\delta)$. By Proposition 9.156 the visibilities and meters vary continuously under certified retunings, and by Lemma 9.169 the effective visibilities satisfy the displayed bounds, monotonicity, and right-continuity. If, in addition, $\mathcal{R}(x;\delta)$ lies in a single decision quadrant, the quadrant is robust under all admissible retunings by Proposition 9.170; equivalently, plateau robustness holds by Corollary 9.139. This delivers (d) and (e) of Theorem 9.171 (stability and functorial sharpening; cf. Lemma 9.136 and Proposition 9.123).

Causal correctness and BRST/Nielsen safety. The reported T–L–O in (C3) is computed via the typed, retarded, cone-gated pipeline: retardedness and cone support follow from Proposition 9.107 and microcausality from Proposition 9.108, yielding causal correctness (item (b) of Theorem 9.171; see also Propositions 9.149 and 9.155(b)). BRST/Nielsen safety follows because the typed Riesz projector is BRST-closed (Proposition 9.119 and Lemma 9.121), pointwise typed contrasts are invariant (Lemma 9.128 using Proposition 9.97), the visibilities preserve invariance (Proposition 9.134(a)), and the meter valuations are BRST-safe (Lemma 9.145(b)); therefore T–L–O is BRST/gauge independent by Proposition 9.149(a). This is item (a) of Theorem 9.171.

Conclusion. Thus (C1) furnishes typed hyperbolicity and stability margins; (C2) furnishes certified uncertainty regions guaranteeing stability/robustness; and (C3) supplies the causally correct, BRST-safe outputs. By Theorem 9.171, these three pieces are *sufficient* to verify BRST safety, causal correctness, and stability for (x, χ), completing the proof. □

9.15 Pre-Physical Artefacts

We formalize the artefacts produced by the typed, band-level pipeline (§1–§5) for each sector $\bullet \in \{\Phi\Phi, \Theta\Theta\}$. All deliverables are computed with retarded, cone-gated kernels (Definition 9.106 and Proposition 9.107), BRST-closed taggers (Definition 9.96 and Proposition 9.97), and typed restrictions $\Pi^{[\chi]}$ from the permit ledgers (Definitions 9.116 and 9.122).

7.1 Deliverable schema

Definition 9.174 (Deliverable bundle per sector)**.** For sector $\bullet$ at point x, the *deliverable bundle* is

$$\mathrm{Del}[\bullet](x) \;:=\; \Big(\underbrace{\mathrm{PL}[\bullet](x)}_{\text{permit ledger}}, \underbrace{\mathrm{TM}[\bullet](x)}_{\text{typed }\Sigma\text{–maps}}, \underbrace{\mathrm{PD}[\bullet](x)}_{\text{parasymmetric dashboards}}, \underbrace{\mathrm{QC}[\bullet](x)}_{\text{quadrant certificates}} \Big),$$

whose components are defined in Definitions 9.175 to 9.178 below.

Definition 9.175 (Permit ledger deliverable)**.** The *permit ledger* is the finite (locally) set

$$\mathrm{PL}[\bullet](x) \;:=\; \mathcal{C}^{[\bullet]}_{\mathrm{admissible}}(x) \;\subset\; \mathrm{End}(\bullet),$$

as in Definition 9.122, where each entry χ is packaged with:

(a) a Σ-*construction recipe* for $\Pi^{[\chi]}$ (spectral/Riesz projector from Σ-weighted mode operators; cf. Propositions 9.36 and 9.119 and eq. (9.16));

(b) a *cone/hyperbolicity certificate* consisting of: (i) the admissible timelike interval in $\mathcal{C}_x^{\hat{\Sigma},*}$ (Definition 9.104); (ii) the typed positivity margin for $\hat{Q}_x^{[\chi]}[k]$ (Definition 9.111); (iii) the retuning-stability radius from Definition 9.116 (see also Corollary 9.120).

Definition 9.176 (Typed Σ–map deliverable)**.** For each permitted $\chi \in \mathrm{PL}[\bullet](x)$, the *typed* Σ*–map* is the function

$$\mathrm{TM}^{[\chi]}(x) := \Big(\Lambda_{\mathrm{law}}^{\Sigma;[\chi]}(x),\ \Lambda_{\mathrm{state}}^{\Sigma;[\chi]}(x)\Big) \in [0,1]^2, \tag{9.88}$$

computed as in Definition 9.133 from aggregated typed contrasts (9.63). Optionally, an *uncertainty rectangle*

$$\mathcal{R}^{[\chi]}(x;\delta) := \mathcal{B}_{\mathrm{law}}^{[\chi]}(x;\delta) \times \mathcal{B}_{\mathrm{state}}^{[\chi]}(x;\delta)$$

is included (effective visibilities of Definition 9.168).

Definition 9.177 (Parasymmetric dashboard deliverable)**.** For each χ and scale parameter r, the *parasymmetric dashboard* is the pair of typed meter curves

$$\mathrm{PD}^{[\chi]}(r) := \Big(\upsilon_{\mathrm{Synthesis}}^{(a);[\chi]}(r)\Big|_{a\in\{L,S\}},\ \ \upsilon_{\mathrm{Synthesis}}^{(LS);[\chi]}(r)\Big), \tag{9.89}$$

defined from the band meter states (Definition 9.141) via Definitions 9.143 and 9.144. The *onset scales/times* are reported as

$$(r^\star[\chi], t^\star[\chi]) \ \text{ from } \ (9.81), \quad \text{with law-deficit } d_{\mathrm{law}}^{\Sigma;[\chi]} = 1 - \Lambda_{\mathrm{law}}^{\Sigma;[\chi]} \ (\text{eq. } (9.68)). \tag{9.90}$$

Definition 9.178 (Quadrant certificate deliverable)**.** The *quadrant certificate* for χ at x is the tuple

$$\mathrm{QC}^{[\chi]}(x) := \Big(\underbrace{\mathrm{Quad}^{[\chi]}(x)}_{\text{Definition 9.137}},\ \underbrace{\mathrm{Loc}^{[\chi]}(x)}_{\text{arena/sector/position/sheet}},\ \underbrace{\mathfrak{C}^{[\chi]}(x)}_{\text{audit certificate of Definition 9.172}} \Big),$$

where $\mathrm{Loc}^{[\chi]}(x)$ records: arena $\mathcal{A} \in \{\mathfrak{P}, \mathfrak{P}_\partial\}$, sector $\bullet \in \{\Phi\Phi, \Theta\Theta\}$, spatial tag $P_x(\lambda_{\mathrm{UV}})$ (Definition 9.96), and the cone sheet (future/past, null/interior) from $\mathcal{C}_x^{\hat{\Sigma},*}$ (Definition 9.104).

7.2 Completeness, invariances, and functoriality

Proposition 9.179 (Deliverable completeness)**.** *For each sector* $\bullet$ *and permitted* χ*, the bundle* $\mathrm{Del}[\bullet](x)$ *determines the T–L–O output* $\mathrm{TLO}^{[\chi]}(x;r)$ *uniquely for all* r *by* (9.88), (9.89), (9.90)*, and Definition 9.147; conversely,* $\mathrm{TLO}^{[\chi]}$ *together with* $\mathrm{PL}[\bullet]$ *reconstructs* $\mathrm{TM}^{[\chi]}$ *and* $\mathrm{PD}^{[\chi]}$*. Hence* $\mathrm{Del}[\bullet]$ *is equivalent to the output of the execution functor* $\mathscr{E}_\bullet$ *(Definition 9.153, Theorem 9.161).*

Proof of Proposition 9.179. Fix a sector $\bullet \in \{\Phi\Phi, \Theta\Theta\}$ and an admitted type $\chi \in \mathcal{C}^{[\bullet]}_{\text{admissible}}$ (ledger admission in Proposition 9.102). Recall that the deliverable bundle at x is

$$\mathrm{Del}[\bullet](x) \;:=\; \Big(\mathrm{PL}[\bullet],\ \{\mathrm{TM}^{[\chi]}(x)\}_{\chi},\ \{\mathrm{PD}^{[\chi]}\}_{\chi},\ \text{certificates}\Big),$$

where (i) $\mathrm{PL}[\bullet]$ is the permit ledger (typed projectors with certificates), (ii) $\mathrm{TM}^{[\chi]}(x)$ is the *typed* Σ*–map* at x,

$$\mathrm{TM}^{[\chi]}(x) \equiv \Big(\Lambda^{\Sigma;[\chi]}_{\text{law}}(x),\ \Lambda^{\Sigma;[\chi]}_{\text{state}}(x)\Big), \qquad \text{(eq. (9.88) and Definition 9.133)},$$

and (iii) $\mathrm{PD}^{[\chi]}$ is the typed *parasymmetric dashboard*,

$$\mathrm{PD}^{[\chi]} : r \longmapsto \Big(v^{(L);[\chi]}_{\text{Synthesis}}(r),\ v^{(S);[\chi]}_{\text{Synthesis}}(r),\ v^{(LS);[\chi]}_{\text{Synthesis}}(r)\Big),$$

(eq. (9.89) and Definitions 9.141, 9.143 and 9.144)

(A) Forward direction: $\mathrm{Del}[\bullet](x) \;\Rightarrow\; \mathrm{TLO}^{[\chi]}(x;r)$ *uniquely.* Given $\mathrm{TM}^{[\chi]}(x)$, the *Type* component is the typed quadrant

$$\mathrm{Quad}^{[\chi]}(x) = \mathrm{Quad}\Big(\Lambda^{\Sigma;[\chi]}_{\text{law}}(x),\ \Lambda^{\Sigma;[\chi]}_{\text{state}}(x)\Big),$$

by Definition 9.137 (cf. eq. (9.75)). The *Location* component $(\mathcal{A}_\chi, \mathrm{sector}_\chi, P_x, \mathrm{sheet}_\chi)$ is read directly from the ledger tags and band instrumentation (arena/sector/spatial tag/cone sheet; see Definition 9.147). The *Onset* pair $(r^\star[\chi], t^\star[\chi])$ is determined *uniquely* from $\mathrm{PD}^{[\chi]}$ and the typed law–deficit via the rules (9.90)–(9.81) (Definition 9.147; existence/uniqueness of the infima follows under the mild regularity of Lemma 9.148). Hence, for each r, the triple

$$\mathrm{TLO}^{[\chi]}(x;r) = \Big(\mathrm{Quad}^{[\chi]}(x),\ (\mathcal{A}_\chi, \mathrm{sector}_\chi, P_x, \mathrm{sheet}_\chi),\ (r^\star[\chi], t^\star[\chi])\Big)$$

is fixed by (9.88), (9.89), (9.90) and Definition 9.147. Uniqueness is immediate from the determinism of the normalizer (Definitions 9.47 and 9.133) and of the onset rules.

(B) Converse: $\mathrm{TLO}^{[\chi]} + \mathrm{PL}[\bullet] \;\Rightarrow\; \mathrm{TM}^{[\chi]},\ \mathrm{PD}^{[\chi]}$. By the *compact execution formula* (Theorem 9.161), $\mathrm{TLO}^{[\chi]}(x;r)$ is produced from and only from $\big(\Lambda^{\Sigma;[\chi]}_{\text{law}}(x), \Lambda^{\Sigma;[\chi]}_{\text{state}}(x)\big)$, the location tags, and the onset pair computed from the dashboard curves. Conversely, the deliverable relations provide the *reconstruction maps*:

- Eq. (9.88) inverts the quadrant reporting by associating to $\mathrm{TLO}^{[\chi]}$ the *numerical* typed visibility pair $\mathrm{TM}^{[\chi]}(x) = (\Lambda^{\Sigma;[\chi]}_{\text{law}}(x), \Lambda^{\Sigma;[\chi]}_{\text{state}}(x))$.[4]

[4]This uses the ledger's typed projector $\Pi^{[\chi]}$ and the fixed normalizer in Definitions 9.47 and 9.133; gauge/BRST independence guarantees well-definedness, Proposition 9.134(a), Proposition 9.97.

- Eq. (9.89) reconstructs $\mathrm{PD}^{[\chi]}$ from the meter valuations implicit in $\mathrm{TLO}^{[\chi]}$ (via Definitions 9.141, 9.143 and 9.144) together with the onset calibration (9.90). Precisely, the onset pair pins the threshold levels on the curves, and the functorial/continuity properties (Proposition 9.146, Proposition 9.149(c)) fix the dashboard family on the certified band neighborhood.

Thus $\mathrm{TLO}^{[\chi]}$ plus the ledger $\mathrm{PL}[\bullet]$ suffices to recover both $\mathrm{TM}^{[\chi]}$ and $\mathrm{PD}^{[\chi]}$.

(C) Equivalence with the execution functor. By Definition 9.153 the execution functor $\mathscr{E}_\bullet$ produces exactly the (typed) visibilities, the meters/dashboards, and the T–L–O. The compact execution formula (Theorem 9.161) identifies the TLO constructed from $\mathrm{TM}^{[\chi]}$ and $\mathrm{PD}^{[\chi]}$, while Proposition 9.159 ensures that for each admitted χ the functor computes all quantities needed to evaluate $\mathrm{TLO}^{[\chi]}$ and, conversely, no $\mathrm{TLO}^{[\chi]}$ exists for excluded types (ledger discipline via Corollary 9.113). Parts (A) and (B) above therefore show that $\mathrm{Del}[\bullet]$ and the output of $\mathscr{E}_\bullet$ determine each other, i.e. they are *equivalent data*.

Combining (A)–(C) yields the claim: $\mathrm{Del}[\bullet](x)$ determines $\mathrm{TLO}^{[\chi]}(x;r)$ uniquely for all r, and $\mathrm{TLO}^{[\chi]}$ together with $\mathrm{PL}[\bullet]$ reconstructs $\mathrm{TM}^{[\chi]}$ and $\mathrm{PD}^{[\chi]}$; hence $\mathrm{Del}[\bullet]$ is equivalent to the output of $\mathscr{E}_\bullet$. □

Lemma 9.180 (BRST/causal invariance of deliverables)**.** *All scalar entries in* $\mathrm{Del}[\bullet]$ *are gauge-parameter independent (BRST/Nielsen-safe) by Proposition 9.97 and Lemmas 9.128 and 9.145 and causally supported within* $J^+(\mathcal{C}^{\hat{\Sigma},*})$ *by Propositions 9.107 and 9.108.*

Proof of Lemma 9.180. Recall the deliverable bundle at x:

$$\mathrm{Del}[\bullet](x) = \Big(\mathrm{PL}[\bullet],\ \{\mathrm{TM}^{[\chi]}(x)\}_\chi,\ \{\mathrm{PD}^{[\chi]}\}_\chi,\ \text{certificates}\Big).$$

The *scalar* entries are:

(i) the typed visibility pair $\mathrm{TM}^{[\chi]}(x) = \Big(\Lambda^{\Sigma;[\chi]}_{\mathrm{law}}(x), \Lambda^{\Sigma;[\chi]}_{\mathrm{state}}(x)\Big)$ (eq. (9.88), Definition 9.133);

(ii) the typed meter valuations encoded by the dashboard curves $\mathrm{PD}^{[\chi]}$ (eq. (9.89), Definitions 9.141, 9.143 and 9.144);

(iii) the numerical items of the certificate triple (admissible timelike interval, spectral gap, stability radius; Corollary 9.120).

Gauge-parameter independence (BRST/Nielsen safety). For the visibilities, pointwise typed contrasts are BRST/Nielsen-safe by Lemma 9.128 (using Proposition 9.97 and $[\,\hat{Q}_{\mathrm{BRST}}, \Pi^{[\chi]}] = 0$ from Definition 9.116); aggregation/normalization preserves invariance, hence $\Lambda^{\Sigma;[\chi]}_\bullet$ is gauge-invariant (Proposition 9.134(a)). For the meters, BRST/Nielsen safety of all valuations is stated in Lemma 9.145(b). For the certificate scalars, the underlying typed projectors are BRST-closed (constructed as Riesz projectors; Proposition 9.119) and are gauge-stable under conjugation (Proposition 9.123); the gap/interval/radius therefore depend only on BRST classes (see also Lemma 9.121). Thus every scalar entry in $\mathrm{Del}[\bullet]$ is gauge-parameter independent.

Causal support within $J^+(\mathcal{C}^{\hat{\Sigma},*})$. All kernels entering the evaluation of typed contrasts and meter states are replaced by their retarded, cone-gated versions (Definition 9.106), supported in $J^+\Big(\mathcal{C}^{\hat{\Sigma},*}\Big)$ by Proposition 9.107; microcausality holds at spacelike separation

by Proposition 9.108. Hence the visibilities are causally well-posed (Proposition 9.134(b)), and the meter valuations inherit the same causal support (Lemma 9.145(c)). The certificate numbers themselves do not arise from spacetime kernels (they are spectral/positivity data on the band), so no causal propagation issue arises; they are trivially compatible with the cone gate used elsewhere.

Combining the BRST/Nielsen safety cited above with retarded cone-gating and microcausality establishes that every scalar entry reported in Del[•] is gauge-parameter independent and causally supported as claimed. □

Proposition 9.181 (Functorial refinement of deliverables)**.** *For any refinement* $(P', K') \preccurlyeq (P, K)$ *(per Definition 9.167), the following properties hold:*

$$\begin{gathered}\mathrm{PL}[\bullet]_{(P',K')}(x) \subseteq \mathrm{PL}[\bullet]_{(P,K)}(x),\\ \mathrm{TM}^{[\chi]}(x) \ \uparrow,\\ v^{(\cdot);[\chi]}_{\mathrm{Synthesis}}(r) \xrightarrow{\text{cont.}} \textit{(towards } 1/2\textit{)}.\end{gathered}$$

Furthermore, quadrant membership in $\mathrm{QC}^{[\chi]}(x)$ *remains unchanged off tolerance boundaries (Lemma 9.136, Proposition 9.146, and Corollary 9.139).*

Proof of Proposition 9.181. Let $(P', K') \preccurlyeq (P, K)$ be a refinement in the sense of Definition 9.167; in particular, the IR window tightens and the cone gate contracts (cf. Lemma 9.105).

(i) Ledger inclusion. Refining the band taggers $(P, K) \mapsto (P', K')$ is a functorial operation on the pipeline: by Proposition 9.123 the permit ledger can only *shrink* under such refinements, and by the exclusion rule Corollary 9.113 no new type that failed the typed hyperbolicity test (9.53) can be added. Hence

$$\mathrm{PL}[\bullet]_{(P',K')}(x) \ \subseteq \ \mathrm{PL}[\bullet]_{(P,K)}(x).$$

(ii) Typed Σ*–map monotonicity.* Fix a type χ admitted at both tag levels.[5] Cone shrink (IR tightening) implies that aggregated typed contrasts weakly *decrease* while the normalized visibilities weakly *increase* (Lemma 9.136, using normalization in Proposition 9.134(c)). Since $\mathrm{TM}^{[\chi]}(x) = (\Lambda^{\Sigma;[\chi]}_{\mathrm{law}}(x), \Lambda^{\Sigma;[\chi]}_{\mathrm{state}}(x))$ (Definition 9.133), this yields the componentwise monotonicity

$$\mathrm{TM}^{[\chi]}(x)\Big|_{(P',K')} \ \geq \ \mathrm{TM}^{[\chi]}(x)\Big|_{(P,K)}.$$

(iii) Meter continuity and drift toward symmetry. Under IR tightening, the typed synthesis meters satisfy the monotonicity and continuity properties of Proposition 9.146: for fixed typed inputs, $v^{(\cdot);[\chi]}_{\mathrm{Synthesis}}(r)$ is nondecreasing *towards* $1/2$ (loss of accessible coherence cannot raise asymmetry beyond the available support) and varies continuously with the tag parameters in the certified stability radius (Definition 9.116). Therefore

$$v^{(\cdot);[\chi]}_{\mathrm{Synthesis}}(r)\Big|_{(P',K')} \xrightarrow{\text{cont.}} \text{towards } \tfrac{1}{2}.$$

[5] If χ is removed by refinement, the statement is vacuous for that χ at (P', K').

(iv) Quadrant robustness off tolerance boundaries. Because $\mathrm{TM}^{[\chi]}(x)$ increases componentwise under refinement (Step (ii)) and the meters vary continuously (Step (iii)), the typed quadrant $\mathrm{Quad}^{[\chi]}(x)$ (Definition 9.137) is unchanged as long as the visibility pair does not cross a tolerance surface of the decision rule. This is precisely the plateau robustness statement of Corollary 9.139; equivalently, it follows from Lemma 9.136 together with the continuity in Proposition 9.146.

Combining (i)–(iv) proves the asserted functorial refinement properties for the deliverables. □

Corollary 9.182 (Bulk/edge comparison)**.** *On the boundary arena* $\mathfrak{P}_\partial$, $\mathrm{PL}[\bullet]_\partial(x) \subseteq \mathrm{PL}[\bullet]_{\mathrm{bulk}}(x)$ *and* $\mathrm{TM}^{[\chi]}_\partial(x) \geq \mathrm{TM}^{[\chi]}_{\mathrm{bulk}}(x)$ *componentwise, with differences reported in* $\mathrm{QC}^{[\chi]}(x)$*; cf. Corollary 9.160.*

Proof of Corollary 9.182. Let the bulk and boundary band instrumentations be $\mathfrak{B}^{\mathrm{bulk}}_{\lambda_{\mathrm{UV}},\ell} = \left(P_x(\lambda_{\mathrm{UV}}), K^{(\ell)}_{\mathrm{IR}}\right)$ and $\mathfrak{B}^{\partial}_{\lambda_{\mathrm{UV}},\ell'} = \left(P_x(\lambda_{\mathrm{UV}})P_\partial, K^{(\ell')}_{\mathrm{IR}}\right)$ with $\ell' \leq \ell$. Then $(P_xP_\partial, K^{(\ell')}_{\mathrm{IR}}) \preccurlyeq (P_x, K^{(\ell)}_{\mathrm{IR}})$ is a functorial refinement in the sense of Definition 9.167; equivalently, the boundary cone *shrinks* by Lemma 9.105.

Ledger inclusion. Applying Proposition 9.181(i) to this refinement yields $\mathrm{PL}[\bullet]_\partial(x) \subseteq \mathrm{PL}[\bullet]_{\mathrm{bulk}}(x)$, which also recovers the ledger inclusion proved directly in Corollary 9.160.

Typed Σ–map monotonicity. For any type χ admitted at the boundary, Proposition 9.181(ii) gives the componentwise increase

$$\mathrm{TM}^{[\chi]}_\partial(x) = \left(\Lambda^{\Sigma;[\chi]}_{\mathrm{law},\partial}(x),\, \Lambda^{\Sigma;[\chi]}_{\mathrm{state},\partial}(x)\right) \;\geq\; \left(\Lambda^{\Sigma;[\chi]}_{\mathrm{law,bulk}}(x),\, \Lambda^{\Sigma;[\chi]}_{\mathrm{state,bulk}}(x)\right) = \mathrm{TM}^{[\chi]}_{\mathrm{bulk}}(x),$$

where $\mathrm{TM}^{[\chi]}$ is the typed Σ–map (Definition 9.133). This is the deliverable-level restatement of the visibility monotonicity proved in Corollary 9.160 (via Lemma 9.136 and Proposition 9.134(c)).

Reporting differences in $\mathrm{QC}^{[\chi]}(x)$. The quadrant certificate $\mathrm{QC}^{[\chi]}(x)$ records the typed quadrant and the location tags (arena/sector/position/cone sheet) used by T–L–O (Definition 9.147). Since boundary refinement is functorial (Proposition 9.181) and the cone sheet/arena change from $\mathfrak{P}$ to $\mathfrak{P}_\partial$, any bulk/edge discrepancy in the typed Σ–map is reflected and reported in $\mathrm{QC}^{[\chi]}(x)$ together with the boundary location tag; cf. the bulk/edge discussion in Corollary 9.160.

These three assertions establish the stated bulk/edge comparison. □

7.3 Minimal publishing package and verification

Definition 9.183 (Minimal publishing package)**.** A *minimal package* for (x, χ) consists of:

(i) the ledger entry with recipe and certificate ($\Pi^{[\chi]}$, cone/hyperbolicity/stability) (Definition 9.175);

(ii) the typed Σ–map with uncertainty rectangle $\mathcal{R}^{[\chi]}(x;\delta)$ (Definitions 9.168 and 9.176);

(iii) the dashboard curves and onsets $(r^\star[\chi], t^\star[\chi])$ (Definition 9.177);

(iv) the quadrant certificate $\mathrm{QC}^{[\chi]}(x)$ including the audit certificate $\mathfrak{C}^{[\chi]}(x)$ (Definitions 9.172 and 9.178).

Corollary 9.184 (Sufficiency for independent verification)**.** *Providing the minimal package of Definition 9.183 is sufficient for an independent reconstruction and verification of* $\mathrm{TLO}^{[\chi]}(x;r)$ *for all* r*, and for re-running the audits of §6. In particular, the uncertainty rectangle criterion of Proposition 9.170 certifies quadrant robustness without access to internal gauge choices.*

Proof. By Corollary 9.173, the minimal package (certificate triple + uncertainty rectangles + quadrant/T–L–O outputs) suffices to verify BRST safety, causal correctness, and stability. Independent *reconstruction* of $\mathrm{TLO}^{[\chi]}(x;r)$ follows from deliverable completeness (Proposition 9.179) together with the compact execution formula (Theorem 9.161), which shows that the T–L–O triple is determined by the typed visibilities, location tags, and onset rules.

Re-running the audits of §6 uses only the data included in the minimal package: the cone/hyperbolicity certificate triple supports the hyperbolicity audit (Definition 9.165, cf. Corollary 9.120); the uncertainty rectangles provide the band-stability/audit sets (Definition 9.166) and certify quadrant robustness via Proposition 9.170; BRST/Nielsen safety and causal correctness follow from Theorem 9.171 (items (a)–(b)), which rely only on BRST-closed instrumentation and cone-gated retarded kernels (Propositions 9.97, 9.107 and 9.108 and Lemmas 9.128 and 9.145).

Therefore the minimal package of Definition 9.183 is sufficient both to reconstruct/verify $\mathrm{TLO}^{[\chi]}(x;r)$ for all r and to re-run the audits of §6 without access to internal gauge choices, completing the proof. □

9.15.1 Typed Pre-Physics: Ledger, Visibilities, TLO

The sequence of constructions in Sections 9.8 to 9.15 turns the operator chronogeometry $\hat{\Sigma}$ and the unified block fields into a *typed, testable* pipeline whose outputs are the parasymmetric *Type–Location–Onset* assignments $\mathrm{TLO}^{[\chi]}(x;r)$ for the two pre-physical quanta—*apeiron* ($\Phi\Phi$; Gaiamiti) and *kymon* ($\Theta\Theta$; Tantrakyma).

(1) Causality and order from $\hat{\Sigma}$. The Σ–pre-physical cone $\mathcal{C}^{\hat{\Sigma},*}$ is *derived* from the probe $\hat{Q}_x[v] = v^\mu v^\nu \hat{\Sigma}_{\mu\nu}(x)$ and band functionals; all kernels are retarded and cone-gated in $J^+(\mathcal{C}^{\hat{\Sigma},*})$, and spacelike commutators vanish (Section 9.9). This makes "chronogeometric order" an *operational* feature of $\hat{\Sigma}$, not an assumption imported from an expectation layer.

(2) Kinematics discovered, not posited. Within each block ($\Phi\Phi$ for apeiron, $\Theta\Theta$ for kymon), the *kinematic permit census* uses band-finite, graded-covariant, cone-compatible projectors $\Pi^{[\chi]}$ and a typed hyperbolicity test to *admit* or *exclude* labels χ (see the ledger statement and typed certificates in Proposition 9.102 and Corollary 9.120). Thus, allowed "spins/helicities/polarizations" at pre-physics are *outputs with certificates*, not inputs.

(3) Mechanisms measured by typed visibilities. Typed Σ–contrasts separate *explicit law effects* from *state response* and normalize to visibilities $\Lambda^{\Sigma;[\chi]}_{\mathrm{law/state}} \in [0,1]$ (Section 9.11), feeding

the canonical quadrant map $\mathrm{Quad}^{[\chi]}$ (typed mechanism). This provides a sector-conditioned witness pair for each admitted χ under the same causal gate.

An algorithm, not an analogy. Section 9.13 makes the pipeline a *procedure*: (1) build the permit ledger; (2) compute visibilities; (3) evaluate meters and report T–L–O; (4) compare bulk/edge when applicable. Soundness (BRST safety and causal support) and completeness (no reports for excluded types) are established by the execution results.

Guardrails that make numbers trustworthy. The audit suite in Section 9.14 comprises five items: BRST/Nielsen; cone/causality; typed hyperbolicity; band stability; and functoriality. By the global guardrails theorem, every scalar output is (i) cohomologically invariant, (ii) causally supported in $J^+\big(\mathcal{C}^{\hat{\Sigma},*}\big)$, (iii) typed well posed, and (iv) stable under certified retunings (monotone under cone shrink). Hence the pipeline's numbers are *meaningful* and *robust*—not artifacts of gauge choice or regulator quirks.

Deliverables and reproducibility. Section 9.15 packages the outputs into *publishable artefacts*: a permit ledger with certificates (cone interval, spectral gap, stability radius), typed Σ–maps, dashboards, and quadrant certificates. A minimal certificate suffices for independent reconstruction and re-audit of $\mathrm{TLO}^{[\chi]}(x;r)$.

Implications for the two quanta. *Apeiron (Gaiamiti; ΦΦ).* The census admits precisely those chronogeometric modes that pass typed hyperbolicity after Σ–restriction—e.g., Σ–trace/trace-free and TT-like subspaces, with helicity refinement where a Σ–spatial direction exists (Section 9.10). For each admitted χ, the typed visibilities quantify whether the *law* or the *state* is driving deviations, and the parasymmetric meters certify *onset scales/times* for symmetry breaking or healing. In effect, the apeiron's admissible content becomes a *measured property* of $\hat{\Sigma}$—chronogeometric excitations are certified by cone/hyperbolicity data rather than assumed. Bulk/edge comparisons then show that boundaries *shrink* the ledger and *increase* visibilities (more symmetric appearance), making edge diagnostics a controlled refinement of bulk behavior (cf. functorial/boundary results and Corollary 9.160).

Kymon (Tantrakyma; ΘΘ). The same typed pipeline admits Σ–transverse/longitudinal channels and helicity $\lambda = \pm$, and—crucially—the *Chrono–Kymon* torsional/temporal channel is treated on equal footing: when its Σ–weighted torsional operator has an isolated nonzero band, the corresponding Riesz projector defines a permitted type with a positive certificate margin (Corollary 9.125). Typed visibilities now disentangle explicit torsional law effects from state-induced responses, and the meters supply *onset scales* for torsional dynamics, with the location metadata specifying whether such activity is bulk or boundary, which cone sheet it occupies, and how it coexists with other kymon types. In short, kymonic excitations are *admitted, diagnosed, and localized* by the same causal/BRST-clean yardstick, making torsion a quantifiable pre-physical feature rather than an interpretive add-on.

Bottom line. The parasymmetric application at pre-physics does three things at once: (i) it *derives* causal order from $\hat{\Sigma}$, (ii) it *discovers* the admissible kinematics of apeiron/kymon via typed hyperbolicity, and (iii) it *diagnoses* mechanisms and their *where/when* through typed

visibilities and meters. For both Gaiamiti and Tantrakyma, what counts as a legitimate mode, what drives its behavior, and where/when it turns on are no longer assumptions—they are *certified outcomes* of a BRST-clean, cone-gated, band-stable pipeline, ready to be published and independently verified.

9.15.2 Permitted Kinematics for the Gaiamiti (Apeiron, ΦΦ) Field

No a priori spin: admission by certificate. In the apeiron block ΦΦ, kinematic labels ("spin/helicity/polarization" in the pre-physical sense) are *not* postulated; they are *admitted* if and only if their typed projectors $\Pi^{[\chi]}$ satisfy the band-level criteria of Definition 9.116 and pass the cone/hyperbolicity/stability audits that define the ledger (Proposition 9.102). Concretely, $\chi \in \mathcal{C}^{[\text{apeiron}]}_{\text{admissible}} \subset \text{End}(\Phi\Phi)$ iff

(a) $\Pi^{[\chi]}$ is band-finite, graded-covariant, BRST-closed, and cone-compatible (Definition 9.116); and

(b) the *typed hyperbolicity* condition holds on cone-timelike covectors,

$$\inf_{\Psi} \mathcal{I}_{\Psi}\Big[\hat{Q}_x[k]\Big|_{\Pi^{[\chi]}}\Big] > 0, \qquad k \in \operatorname{int} \mathcal{C}_x^{\hat{\Sigma},*},$$

with continuity under small retunings (Definition 9.111 and Proposition 9.112).

Each admitted χ carries its cone/hyperbolicity certificate (admissible timelike interval, spectral gap, stability radius; Corollary 9.120).

Canonical typed subspaces considered in ΦΦ. Within the symmetric apeiron sector, the *constructive* Σ-weighted spectral/Riesz procedure (Proposition 9.119) yields the following natural classes, each to be kept *only if* it passes the typed hyperbolicity test above:

(i) Σ–trace (scalar-like) $\Pi^{[\Sigma\text{-tr}]}$ projects to the Σ–trace channel of the symmetric tensor. Interpretable as a scalar-like polarization of the chronogeometric block; helicity refinement is trivial (isotropic).

(ii) Σ–trace-free symmetric, TT-like $\Pi^{[\text{TT}_\Sigma]}$ onto the trace-free subspace; where a Σ–spatial direction $\hat{\mathbf{k}}$ is meaningful (away from null degeneracies), introduce *helicity-refined* projectors $\Pi^{[\text{TT}_\Sigma;\lambda]}$ with $\lambda = \pm$ that resolve the two circular polarizations on the Σ–spatial slice (helicity labels at pre-physics; no particle/graviton mapping).

(iii) Σ–divergence decomposition $\Pi^{[\text{T}_\Sigma]}$ (Σ–transverse) and $\Pi^{[\text{L}_\Sigma]}$ (Σ–longitudinal) channels of the symmetric field (defined by the Σ–divergence). Where applicable, one may further refine $\Pi^{[\text{T}_\Sigma]}$ by helicity $\lambda = \pm$ along $\hat{\mathbf{k}}$.

These are the *candidate* polarizations one typically tests in the apeiron ledger; admission is granted *only* when the typed principal form is positive and stable (Proposition 9.112 and Corollary 9.120). Any candidate failing (9.53) is *excluded* by Corollary 9.113.

What the ledger actually permits (and how to read it). The permit ledger for Gaiamiti at a point x is the set

$$\mathcal{C}^{[\text{apeiron}]}_{\text{admissible}}(x) = \Big\{\ \chi \in \{\Sigma\text{-tr},\ \mathrm{TT}_\Sigma,\ \mathrm{T}_\Sigma,\ \mathrm{L}_\Sigma,\ \text{helicity refinements (constructed as above)}\} \ \Big|\ \chi \text{ passes (9.53)}\ \Big\}.$$

For each admitted χ, the *typed* witness pair $\left(\Lambda^{\Sigma;[\chi]}_{\text{law}}(x), \Lambda^{\Sigma;[\chi]}_{\text{state}}(x)\right)$ (Definition 9.133) and the parasymmetric meters determine the *Type* (quadrant), *Location* (arena/sector/position/cone sheet), and *Onset* scales/times (Definition 9.147). Helicity-refined entries carry separate rows in the ledger (each with its own certificate triple), so, e.g., $\mathrm{TT}_\Sigma;+$ and $\mathrm{TT}_\Sigma;-$ may in principle have distinct admission windows or onsets.

Edge/boundary behavior and functoriality. Under IR tightening or passage to the boundary arena $\mathfrak{P}_\partial$, the cone *shrinks* (Lemma 9.105), so the apeiron ledger can only *lose* entries, never gain them (Proposition 9.123); correspondingly, the typed visibilities weakly *increase* (monotone toward symmetry) and meters drift continuously toward 1/2 (Lemma 9.136, Proposition 9.146, and Corollary 9.160). Quadrant membership remains stable off tolerance boundaries (Corollary 9.139 and Proposition 9.170).

Summary. For the Gaiamiti field, the *permitted* spins/helicities/polarizations are precisely those Σ–structured subspaces—*Σ–trace*, *Σ–trace-free (TT-like)*, and *Σ–transverse/longitudinal* (with helicity refinements where a Σ–spatial direction exists)—whose typed projectors are band-finite, BRST-closed, cone-compatible, and satisfy *typed hyperbolicity* with a nonzero certificate margin. No other kinematic labels are admitted at pre-physics; and no mapping to particles/gravitons is made below Planckian. The exact list at a given x is your apeiron permit ledger $\mathrm{PL}[\Phi\Phi](x)$, carrying the cone interval, spectral gap, and stability radius for each χ.

9.15.3 Permitted Kinematics for the Tantrakyma (Kymon, $\Theta\Theta$) Field

Admission, not assumption. As in the apeiron case, kinematic labels in the kymon block $\Theta\Theta$ ("spin, helicity, polarization" in the pre-physical sense) are *admitted* if and only if their typed projectors $\Pi^{[\chi]}$ satisfy the band-level criteria of Definition 9.116 and pass the cone/hyperbolicity/stability audits that define the permit ledger (Proposition 9.102). Concretely, $\chi \in \mathcal{C}^{[\text{kymon}]}_{\text{admissible}} \subset \mathrm{End}(\Theta\Theta)$ iff $\Pi^{[\chi]}$ is band-finite, graded-covariant, BRST-closed, cone-compatible, and the typed principal form is strictly positive on cone-timelike covectors with stability under small retunings (Definition 9.111 and Proposition 9.112). Each admitted χ carries its cone/hyperbolicity *certificate triple* (admissible timelike interval, spectral gap, stability radius; Corollary 9.120).

Canonical typed subspaces considered in $\Theta\Theta$. Using the constructive Σ-weighted spectral/Riesz procedure (Proposition 9.119), the natural *candidate* channels in the kymon sector include:

(i) Σ–transverse vs. Σ–longitudinal Projectors $\Pi^{[\mathrm{T}_\Sigma]}$ and $\Pi^{[\mathrm{L}_\Sigma]}$ defined by the Σ–divergence in the ΘΘ block. Where a Σ–spatial direction $\hat{\mathbf{k}}$ is meaningful (timelike k), refine $\Pi^{[\mathrm{T}_\Sigma]}$ by helicity $\lambda = \pm$ to resolve circular polarizations on the Σ–spatial slice.

(ii) Helicity-refined transverse modes $\Pi^{[\mathrm{T}_\Sigma;\lambda]}$ with $\lambda = \pm$ obtained by commuting the spectral projector with the little-group stabilizer on the Σ–spatial hyperplane determined by a timelike k (same mechanism as in the apeiron case; cf. Definition 9.116).

(iii) Chrono–Kymon (torsional/temporal) channel Let $T := \hat{t}^{(\Theta)}_{\mu\nu}\hat{\Sigma}^{(\Theta)\,\mu\nu}$ be the Σ–weighted torsional operator in the ΘΘ block. If T has a nonzero isolated spectral band, the Riesz projector $\Pi^{[\mathrm{CK}]}$ defines a typed channel; if the band is separated from 0, the typed hyperbolicity condition holds and CK is admitted by Corollary 9.125.

Each candidate is retained *only if* it passes typed hyperbolicity with a nonzero margin (Proposition 9.112 and Corollary 9.120); otherwise it is excluded by Corollary 9.113.

What the ledger actually permits (and how to read it). At a point x the kymon permit ledger is

$$\mathcal{C}^{[\mathrm{kymon}]}_{\mathrm{admissible}}(x) = \Big\{ \chi \in \{\mathrm{T}_\Sigma,\ \mathrm{L}_\Sigma,\ (\mathrm{T}_\Sigma;\ \lambda = \pm),\ \mathrm{CK},\ \Sigma\text{-structured refinements}\} \ \Big|\ \chi \text{ passes (9.53)} \Big\}.$$

For each admitted χ, the typed visibility pair $\left(\Lambda^{\Sigma;[\chi]}_{\mathrm{law}}(x), \Lambda^{\Sigma;[\chi]}_{\mathrm{state}}(x)\right)$ (Definition 9.133) diagnoses mechanism type via the quadrant rule, while the parasymmetric meters determine onset scales/times and supply the *Location* tags (arena/sector/position/cone sheet) (Definition 9.147). Helicity-refined entries ($\lambda = \pm$) appear as separate rows, each with its own certificate triple, so distinct helicities may have different admission windows or onsets.

Edge/boundary behavior and functoriality. Under IR tightening or passage to the boundary arena $\mathfrak{P}_\partial$, cone shrink (Lemma 9.105) implies the kymon ledger can only *lose* entries (Proposition 9.123); visibilities weakly *increase* (monotone toward symmetry) and meters drift continuously toward 1/2 (Lemma 9.136, Proposition 9.146, and Corollary 9.160). Quadrant membership remains stable off tolerance boundaries (Corollary 9.139 and Proposition 9.170).

Summary. For Tantrakyma, the *permitted* pre-physical kinematics are precisely the Σ–structured ΘΘ channels—Σ–transverse/longitudinal (with helicity $\lambda = \pm$ where defined) and the torsional/temporal *Chrono–Kymon* mode—whose typed projectors are band-finite, BRST-closed, cone-compatible, and satisfy *typed hyperbolicity* with a positive certificate margin. No other kinematic labels are admitted at pre-physics, and no particle/graviton mapping is made below Planckian. The concrete, location- and onset-annotated list at x is your kymon permit ledger $\mathrm{PL}[\Theta\Theta](x)$.

9.15.4 Pre-Physics Parasymmetric Admission, Mechanism, Onset

Pre-physical kinematics as admission, not assumption. Kinematics at the pre-physical level is defined operationally and uniformly for both fundamental sectors—apeiron (ΦΦ; Gaiamiti) and kymon (ΘΘ; Tantrakyma)—without invoking expectation values or particle language. A typed label χ is *permitted* exactly when the Σ–induced principal quadratic form, restricted by the typed projector $\Pi^{[\chi]}$, is strictly positive on all cone-timelike covectors of the Σ–pre-physical cone:

$$\inf_{\Psi\in\mathcal{S}_{\mathrm{band}}} \mathcal{I}_\Psi\Big[\hat{Q}_x[k]\Big|_{\Pi^{[\chi]}}\Big] \;>\; 0 \qquad \text{for all } k\in \operatorname{int}\mathcal{C}_x^{\hat{\Sigma},*},$$

with cone-compatibility and band-finiteness as structural requirements (Section 9.10, Definitions 9.111 and 9.116, and Proposition 9.112). This yields a certified *permit ledger* for each sector.

Causal order from operator chronogeometry. Chronogeometric order is extracted from $\hat{\Sigma}$ alone. The covector cone $\mathcal{C}_x^{\hat{\Sigma},*}$ is defined by the band-admissible Σ–probe $\hat{Q}_x[v]=v^\mu v^\nu\hat{\Sigma}_{\mu\nu}(x)$; all kernels are retarded and cone-gated within $J^+\big(\mathcal{C}^{\hat{\Sigma},*}\big)$, and spacelike commutators vanish by construction (Section 9.9). Thus future/past structure, causal support, and microcausality arise as *operator-level* features rather than consequences of an expectation-level metric.

Typed witnesses that separate law from state. For each permitted χ, the visibility vector

$$\vec{\Lambda}^{[\chi]}(x) := \Big(\Lambda_{\mathrm{law}}^{\Sigma;[\chi]}(x),\Lambda_{\mathrm{state}}^{\Sigma;[\chi]}(x)\Big)\in[0,1]^2$$

quantifies explicit-law effects and state-dependent response under the same causal gate (Section 9.11 and Definition 9.133). Applying the canonical quadrant decision to $\vec{\Lambda}^{[\chi]}$ yields the typed mechanism $\mathrm{Quad}^{[\chi]}(x)\in\{\mathrm{QI},\mathrm{QII},\mathrm{QIII},\mathrm{QIV}\}$ (Definitions 7.685, 9.47 and 9.137).

Parasymmetric readouts and T–L–O. Band-level parasymmetric meters, restricted to the typed subspace, provide operational valuations at local anchors and on the relational pair. Their scale/time dependence defines onset thresholds and location metadata, producing the full typed *Type–Location–Onset* assignment $\mathrm{TLO}^{[\chi]}(x;r)$ (Section 9.12 and Definition 9.147).

Meaning for the two fundamental quanta. *Apeiron (ΦΦ; Gaiamiti).* Typical admitted subspaces (per the admission criterion above) include Σ–trace, Σ–trace-free symmetric (TT-like), and helicity refinements when a Σ–spatial direction is meaningful (Section 9.10).
Kymon (ΘΘ; Tantrakyma). The same typed structure applies—Σ–transverse and Σ–longitudinal (with helicities $\lambda=\pm$) together with the torsional/temporal *Chrono–Kymon* channel—each treated within the same admission/witness/readout pipeline (Corollary 9.125).

Uniform guardrails and reproducibility. A unified audit suite—BRST/Nielsen safety, cone/-causality, typed hyperbolicity, band stability, and functoriality—applies to every stage (Section 9.14). Consequently all scalar outputs (visibilities, meters, T–L–O) are cohomologically invariant, causally supported within $J^+\big(\mathcal{C}^{\hat{\Sigma},*}\big)$, well-posed for admitted types, and

stable under certified retunings with monotone behavior under cone shrink. Deliverables are packaged for reconstruction and independent verification (Section 9.15).

Bridge from pre-physical breaking to physical realization. The PQS/QHS operator chronogeometry supplies certified pre-physical structures—admitted types (permit ledgers), mechanisms ($\mathrm{Quad}^{[\chi]}$), and onsets/locations ($\mathrm{TLO}^{[\chi]}$)—which can flow, under appropriate coarse-graining, into physical realizations without invoking particle language or expectation-level geometry in the sub-Planckian regime.

9.16 Typed Parasymmetric Quadrants of the Gaiamiti and Tantrakyma Fields

We *fix the realized regime* (plateau certified; evaluation on the BRST-physical sector) and formalize the typed, sector-resolved *parasymmetric* diagnostics from Chapter 9 so that the *Typed Quadrant Ledger* is populated *before* any particle naming. All readouts use the canonical decision rule of §8.6 on the *same diagnostic realized reference configuration*. Throughout, $\alpha \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$ denotes the triadic channels (Chapter 7), and $S \in \{\Phi\Phi, \Theta\Theta\}$ indexes the (gaiamiti, tantrakyma) sectors. Band tags $P_x(\lambda_{\mathrm{UV}})$, $K_{\mathrm{IR}}^{(\ell)}$ and the band functor $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ follow Chapter 8.

Definition 9.185 (Diagnostic realized reference configuration)**.** Fix a certified plateau $\mathfrak{P}$ (bulk) or $\mathfrak{P}_\partial$ (boundary) per Chapter 8, with band tags $P_x(\lambda_{\mathrm{UV}})$, $K_{\mathrm{IR}}^{(\ell)}$ (cf. use in Section 9.16.2 and location/onset semantics in Definition 9.190). Let $\mathsf{P}_{\mathrm{phys}}$ denote the BRST projection to $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ (see gauge/Nielsen safety in Proposition 9.191).

The *diagnostic realized reference configuration* is

$$\Big(\mathcal{H}_{\mathrm{phys}},\ \ \mathfrak{P} \text{ or } \mathfrak{P}_\partial;\ P_x(\lambda_{\mathrm{UV}}),\ K_{\mathrm{IR}}^{(\ell)};\ \{\Pi_\alpha\}_{\alpha\in\{\mathrm{corr,vac,spec}\}}\Big),$$

with all expectations on $\mathcal{H}_{\mathrm{phys}}$ and all limits taken *within* $\mathfrak{P}$ (or $\mathfrak{P}_\partial$) under Chapter 8 tolerances. This configuration underlies the typed subspaces (Section 9.16.1), the typed witnesses (Definition 9.187), and the typed quadrant predicate (Definition 9.189).

Definition 9.186 (Typed projectors and sector selection)**.** For a *sector* $S \in \{\Phi\Phi, \Theta\Theta\}$ and a *type* $[\chi]$ (e.g. TT spin-2, trace spin-0, T/L, helicity $\lambda = \pm$, Chrono-Kymon), a BRST-closed *typed projector* $\Pi_S^{[\chi]}$ on $\mathcal{H}_{\mathrm{phys}}$ is defined *on the diagnostic realized reference configuration* of Definition 9.185 and with types chosen from the menus in Definitions 9.195 and 9.196. It satisfies:

(i) Triadic compatibility

$\Pi_\alpha\, \Pi_S^{[\chi]} = \Pi_S^{[\chi]}\, \Pi_\alpha$ for all $\alpha \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$ (triadic channels as fixed in Definition 9.185); this ensures channel-wise, unambiguous typed readouts used in Section 9.16.2.

(ii) Sector purity
$\Pi_S^{[\chi]}$ annihilates the complement of S and restricts to the standard spin/helicity/polarization subspaces determined by the realized geometry $\bar{\Sigma}$ on the plateau (cf. the explicit constructions in Definitions 9.200 and 9.201).

(iii) Cone preservation
Principal symbols remain inside the $\bar{\Sigma}$-cones fixed at the plateau (no cross-cone leakage), as required for orthogonality/completeness in Proposition 9.202 and for the well-posedness/Nielsen-flatness statements of Proposition 9.191 and the typed quadrant assignment in Proposition 9.212.

These projectors are the typed filters inserted in the witness $\mathbf{W}_{[\chi]}^{[S]}$ of Definition 9.187 and thereby drive the canonical quadrant predicate in Definition 9.211.

Definition 9.187 (Typed witnesses on H^0 (SynTh))**.** Let the *sector-conditioned witness* be denoted $\mathbf{W}^{[\mathscr{S}]}$ (cf. Definition 8.4). Its typed version is

$$\mathbf{W}_{[\chi]}^{[S]} \equiv \left(v_{\text{Synthesis}}^{(L)},\, v_{\text{Synthesis}}^{(S)},\, v_{\text{Synthesis}}^{(LS)}\right)^{[S,\chi]}.$$

Formally, for fixed $(S,[\chi])$ and a diagnostic realized reference configuration as in Definition 9.185,

$$\left(v_{\text{Synthesis}}^{(L)},\, v_{\text{Synthesis}}^{(S)},\, v_{\text{Synthesis}}^{(LS)}\right)^{[S,\chi]}(\ell;\mathfrak{P})$$

are the SynTh valuations (Law, State, and joint Law⊕State) obtained by inserting the typed filter $\Pi_S^{[\chi]}$ from Definition 9.186 (cf. explicit constructions in Definitions 9.200 and 9.201) and the triadic projectors $\{\Pi_\alpha\}$ from Definition 9.185:

$$\Pi_\alpha\, \Pi_S^{[\chi]}\,(\cdot)\,\Pi_S^{[\chi]}\,\Pi_\alpha, \qquad \alpha \in \{\text{corr}, \text{vac}, \text{spec}\},$$

with evaluation on $\mathcal{H}_{\text{phys}}$ following the procedure of Section 9.16.2. By construction these valuations are Nielsen-flat on $\mathfrak{P}$ (see Proposition 9.191); they are the inputs that drive the typed quadrant predicate in Definition 9.211.

Definition 9.188 (Law/State sources and typed healing)**.** Let $\delta_L^{[S,\chi]}$ (resp. $\delta_S^{[S,\chi]}$) denote the typed *law* (resp. *state*) source derived from the typed witness $\mathbf{W}_{[\chi]}^{[S]}$ on the certified plateau $\mathfrak{P}$ (typed witness as in Definition 9.187, diagnostic realized reference configuration as in Definition 9.185, and typed filters in Definition 9.186). Define the IR *healing indicators*

$$\mathscr{H}_L^{[S,\chi]} := \lim_{\substack{\ell\uparrow\infty \\ (\lambda,\ell)\in\mathfrak{P}}} \delta_L^{[S,\chi]}(\ell), \qquad \mathscr{H}_S^{[S,\chi]} := \lim_{\substack{\ell\uparrow\infty \\ (\lambda,\ell)\in\mathfrak{P}}} \delta_S^{[S,\chi]}(\ell),$$

where '$\uparrow\infty$' denotes the bulk/boundary IR dilation of Chapter 9 (cf. the overlay in Definition 9.214 and the control axis in Definition 9.190). We say *healed* iff $\mathscr{H}_\bullet = 0$ and *persistent* iff $\mathscr{H}_\bullet > 0$; these indicators feed the canonical quadrant predicate of Definition 9.211 and underpin the typed healing/persistence property in Proposition 9.193.

Definition 9.189 (Typed quadrant functional)**.** The *typed quadrant* of a sector/type pair $(S,[\chi])$ (with $S\in\{\Phi\Phi,\Theta\Theta\}$ and $[\chi]$ chosen from Definitions 9.195 and 9.196) is the value

$$\mathrm{Quad}^{[S,\chi]}\in\{\mathrm{QI},\mathrm{QII},\mathrm{QIII},\mathrm{QIV}\},$$

decided by the canonical rule of §8.6, evaluated *on the same diagnostic realized reference configuration* as in Definition 9.185 (same certified $\mathfrak{P}$ or $\mathfrak{P}_\partial$) and with the same typed filter $\Pi_S^{[\chi]}$ of Definition 9.186 inserted in the SynTh witness $\mathbf{W}_{[\chi]}^{[S]}$ from Definition 9.187. The decision uses the diagnostic realized reference configuration SynTh meters of §9.16.2 together with the typed healing indicators of Definition 9.188, and obeys the following invariants:

(i) QI (symmetric)
No law source and no sustained state selection, i.e. $\mathscr{H}_L^{[S,\chi]}=0$ and $\mathscr{H}_S^{[S,\chi]}=0$.

(ii) QII (SSB)
Law symmetric with state selection (finite-volume elevation, asymptotic healing): $\mathscr{H}_L^{[S,\chi]}=0$, $\mathscr{H}_S^{[S,\chi]}=0$, and an NG pole appears in the correlational channel Π_{corr} for the typed mode (cf. Proposition 9.197).

(iii) QIII (explicit, healing)
Nonzero law source at finite scale with IR *healing*, encoded by $\mathscr{H}_L^{[S,\chi]}=0$.

(iv) QIV (explicit, persistent)
Nonzero law source with *persistence* and GMOR slope on $\mathfrak{P}$: $\mathscr{H}_L^{[S,\chi]}>0$, exhibiting pNG behavior where applicable (cf. Proposition 9.197).

Gauge/Nielsen safety of this assignment follows from Proposition 9.191, and the resulting label $\mathrm{Quad}^{[S,\chi]}$ is what is recorded in the ledger of Definition 9.198.

Definition 9.190 (Location and onset (T–L–O data))**.** The *location* $\mathrm{Loc}^{[S,\chi]}$ takes values *bulk* or *boundary* and is determined by the chosen diagnostic realized reference configuration of Definition 9.185: use the bulk plateau $\mathfrak{P}$ for *bulk* and the boundary plateau $\mathfrak{P}_\partial$ for *boundary* (cf. the bulk–boundary overlay in Definition 9.214). The *onset* $\mathrm{On}^{[S,\chi]}$ is the minimal (certified) scale/time along the plateau at which the typed quadrant predicate in Definition 9.189 first holds, i.e.

$$\mathrm{On}^{[S,\chi]}:=\inf\left\{\,\upsilon\in\mathrm{Axis}(\mathfrak{P})\;\middle|\;\mathrm{Quad}^{[S,\chi]}(\upsilon)\text{ is stable under the Chapter 8 tolerances (see Definition 9.210)}\,\right\},$$

where $\mathrm{Axis}(\mathfrak{P})$ denotes the control axis (IR length ℓ, drive time, boundary size, or mixed; Chapter 9; cf. its use in Definition 9.214 and the decision pipeline of Section 9.16.3). The pair $\left(\mathrm{Loc}^{[S,\chi]},\mathrm{On}^{[S,\chi]}\right)$ constitutes the T–L–O data used in the T–L–O stamp of Definition 9.216 and the ledger augmentation of Corollary 9.218.

Proposition 9.191 (Well-posedness and Nielsen flatness)**.** *For every $(S,[\chi])$ and any certified plateau (bulk $\mathfrak{P}$ or boundary $\mathfrak{P}_\partial$) as in Definition 9.185, the typed witness $\mathbf{W}_{[\chi]}^{[S]}$ of Definition 9.187 is well-defined on $\mathcal{H}_{\mathrm{phys}}$. Its components $\left(\upsilon^{(L)}_{\mathrm{Synthesis}},\upsilon^{(S)}_{\mathrm{Synthesis}},\upsilon^{(LS)}_{\mathrm{Synthesis}}\right)^{[S,\chi]}$ (cf. §9.16.2) are gauge-parameter independent (Nielsen-flat), and the typed quadrant $\mathrm{Quad}^{[S,\chi]}$ of Definition 9.189 is invariant under all BRST-exact deformations allowed on $\mathfrak{P}$.*

Proof. **Well-posedness.** By Definition 9.187, $\mathbf{W}_{[\chi]}^{[S]}$ is constructed by inserting the typed filter $\Pi_S^{[\chi]}$ (see Definition 9.186) and the triadic projectors $\{\Pi_\alpha\}$ (see Definition 9.185) into the channel-restricted readouts, then evaluating on $\mathcal{H}_{\mathrm{phys}}$. The commutation $\Pi_\alpha \Pi_S^{[\chi]} = \Pi_S^{[\chi]} \Pi_\alpha$ and cone preservation required in Definition 9.186 imply that principal symbols and residues are stable on the plateau, ensuring existence and finiteness of the aggregated diagnostic realized reference configuration meters $\overline{v}_a^{[S,\chi]}$ (cf. Definition 9.210).

Nielsen flatness (gauge independence). The SynTh components are BRST-closed readouts on H^0 (Chapter 7) taken on a certified plateau (Chapter 8). The Nielsen identities imply that the derivative with respect to a gauge-fixing parameter inserts a BRST-exact operator; its expectation on $\mathcal{H}_{\mathrm{phys}}$ vanishes. Hence the meters are gauge independent (see also the channel/type compatibility statement Proposition 9.207 which records Nielsen-flatness for the typed meters).

Quadrant invariance. The canonical predicate in Definition 9.189 depends only on the diagnostic realized reference configuration meters $\overline{v}_a^{[S,\chi]}$ and the typed healing indicators $\mathscr{H}_\bullet^{[S,\chi]}$ of Definition 9.188. Since these are unchanged under BRST-exact deformations and gauge reparametrizations (previous paragraphs), the value $\mathrm{Quad}^{[S,\chi]}$ is invariant. This completes the proof (see also Proposition 9.212 for the assignment's global well-definedness). □

Lemma 9.192 (Typed projector compatibility)**.** *If $\Pi_S^{[\chi]}$ and $\Pi_S^{[\chi']}$ are distinct types for the same sector S, then the corresponding SynTh meters (cf. §9.16.2) decouple at leading order on the certified plateau $\mathfrak{P}$:*

$$\textit{cross-terms} \;=\; \mathcal{O}(\varepsilon_{\mathfrak{P}}),$$

where $\varepsilon_{\mathfrak{P}}$ is the Chapter 8 tolerance scale (Slope/Curv/Osc aggregates).

Proof. By Definition 9.186, the typed filters $\Pi_S^{[\chi]}$ are BRST-closed, commute with the triadic projectors Π_α fixed on the diagnostic realized reference configuration (see Definition 9.185), and preserve the $\bar{\Sigma}$-cones. Orthogonality/completeness of the typed decompositions in each sector (Gaiamiti and Tantrakyma) is recorded in Proposition 9.202, implying a block-diagonal principal symbol for the channel-restricted readouts. The SynTh meters are obtained by inserting these filters into the readouts and aggregating over the plateau window (cf. §9.16.2 and Definition 9.210). Mixed terms then arise only from subleading commutators and boundary-of-window effects; a Duhamel/BCH expansion combined with the Chapter 8 certification tolerances bounds these by $\mathcal{O}(\varepsilon_{\mathfrak{P}})$. Order-of-insertion independence and gauge flatness are consistent with Propositions 9.191 and 9.207. □

Proposition 9.193 (Typed healing vs. persistence)**.** *For each $(S, [\chi])$, the sign of $\mathscr{H}_L^{[S,\chi]}$ in Definition 9.188 is independent of other types $[\chi']$ (for the same sector) and of the complementary sector. Consequently, the QIII/QIV split is meaningful* per type *in the canonical predicate of Definition 9.189.*

Proof. Type-wise independence follows from Lemma 9.192: cross-couplings between $[\chi]$ and $[\chi']$ contribute only at $\mathcal{O}(\varepsilon_{\mathfrak{P}})$, which is below the Chapter 8 significance thresholds entering Definition 9.210. Sector-wise independence is ensured by sector purity in Definition 9.186, which annihilates the complement of S. By Definition 9.188, $\mathscr{H}_L^{[S,\chi]}$ is an IR limit taken along the plateau control axis $\mathrm{Axis}(\mathfrak{P})$ (see Definition 9.190); since the diagnostic realized

reference configuration aggregation and the IR dilation are performed with the same typed filter, the limit depends only on the $[\chi]$-block. Therefore the law-channel dichotomy "healing vs. persistence''—and hence the QIII/QIV branch in Definition 9.189—is a well-defined *typed* property, unaffected by orthogonal types or sectors. Gauge/Nielsen safety of these statements is inherited from Proposition 9.191. □

Corollary 9.194 (Typed Quadrant Ledger is well-defined)**.** *The sextuple*

$$\mathbf{L}^{[S,\chi]} := \Big(\underbrace{S}_{\text{Sector}}, \underbrace{[\chi]}_{\text{Type}}, \underbrace{\mathbf{W}^{[S]}_{[\chi]}}_{\textit{Meters on } H^0}, \underbrace{\mathrm{Quad}^{[S,\chi]}}_{\text{Quadrant}}, \underbrace{\mathrm{Loc}^{[S,\chi]}}_{\text{Location}}, \underbrace{\mathrm{On}^{[S,\chi]}}_{\text{Onset}} \Big)$$

is uniquely determined for every sector/type pair at any certified plateau (bulk $\mathfrak{P}$ or boundary $\mathfrak{P}_\partial$).

Proof. Existence. Fix an diagnostic realized reference configuration as in Definition 9.185 and a sector/type $(S,[\chi])$ with $[\chi]$ taken from the menus Definitions 9.195 and 9.196. The typed projector $\Pi_S^{[\chi]}$ is BRST-closed and cone-preserving by Definition 9.186, hence the typed witness $\mathbf{W}^{[S]}_{[\chi]}$ of Definition 9.187 exists and is finite on $\mathcal{H}_{\mathrm{phys}}$ (diagnostic realized reference configuration aggregation in §9.16.2). The typed quadrant value $\mathrm{Quad}^{[S,\chi]}$ exists by the canonical predicate Definition 9.189, which uses the diagnostic realized reference configuration SynTh meters and healing indicators Definition 9.188. Location/onset data $(\mathrm{Loc}^{[S,\chi]}, \mathrm{On}^{[S,\chi]})$ exist by Definition 9.190. Thus all six entries of $\mathbf{L}^{[S,\chi]}$ are defined.

Gauge/Nielsen safety. By Proposition 9.191, the components of $\mathbf{W}^{[S]}_{[\chi]}$ are Nielsen-flat, and $\mathrm{Quad}^{[S,\chi]}$ is invariant under BRST-exact deformations on $\mathfrak{P}$. Since $\mathrm{Loc}^{[S,\chi]}$ and $\mathrm{On}^{[S,\chi]}$ are defined purely from the diagnostic realized reference configuration choice and the certified stability criteria (Chapter 8; cf. Definitions 9.190 and 9.210), they inherit this invariance.

Uniqueness. Sector purity and typed orthogonality/completeness (cf. Proposition 9.202) together with typed decoupling at leading order (Lemma 9.192) imply that the diagnostic realized reference configuration-aggregated meters for $(S,[\chi])$ are independent of other types $[\chi']$ up to $\mathcal{O}(\varepsilon_{\mathfrak{P}})$, which lies below the Chapter 8 tolerances used in Definition 9.210. Hence the canonical predicate Definition 9.189 produces a unique quadrant label for the fixed diagnostic realized reference configuration and type. Likewise, $\mathrm{On}^{[S,\chi]}$ is the infimum over the control axis $\mathrm{Axis}(\mathfrak{P})$ where that predicate stabilizes, which is unique by the same tolerances (see Definition 9.190). Therefore the mapping $(\mathfrak{P}; S, [\chi]) \mapsto \mathbf{L}^{[S,\chi]}$ is a well-defined function for any certified plateau (bulk or boundary).

Collecting the conclusions proves the statement. □

Typed menus for the two sectors. The following menus fix the vocabulary for $[\chi]$, to be used with the typed projectors $\Pi_S^{[\chi]}$ of Definition 9.186 and the typed witness $\mathbf{W}^{[S]}_{[\chi]}$ of Definition 9.187. They are evaluated on the diagnostic realized reference configuration of Definition 9.185 and feed the quadrant assignment of Definition 9.189 (see also the application of meters in §9.16.2 and the per-type decision in §9.16.3). Entries chosen here are the *Type* components that populate the ledger rows of Definition 9.198.

Definition 9.195 (Gaiamiti ($\Phi\Phi$) typed menu)**.** On the realized geometry $\bar{\Sigma}$ (diagnostic realized reference configuration of Definition 9.185) admit:

(i) TT
Spin-2, transverse-traceless w.r.t. $\bar{\Sigma}$.

(ii) Tr
Spin-0 trace.

(iii) Optional vectorial admixtures
Optional vectorial admixtures if a $\bar{\Sigma}$-preferred foliation is fixed (kept orthogonal to TT/Tr at the plateau; cf. the projector constructions in Definition 9.200).

These types are the allowed $[\chi]$ for $S = \Phi\Phi$ in Definition 9.186, are inserted into the witness of Definition 9.187, and determine the typed quadrant via Definition 9.189.

Definition 9.196 (Tantrakyma ($\Theta\Theta$) typed menu)**.** On $\bar{\Sigma}$ admit:

(i) T, L
Transverse/longitudinal.

(ii) Helicities $\lambda = \pm$
Helicities $\lambda = \pm$ where a chiral frame is meaningful.

(iii) Chrono-Kymon channel
The torsional/time-like *Chrono-Kymon* channel (spin-1 precursor) as a distinct type.

These types are the allowed $[\chi]$ for $S = \Theta\Theta$ in Definition 9.186; the corresponding projectors are given in Definition 9.201. They are used to form the typed witness in Definition 9.187, to evaluate the meters in §9.16.2, and to assign $\mathrm{Quad}^{[S,\chi]}$ via Definition 9.189, after which they are recorded in the ledger of Definition 9.198.

Proposition 9.197 (NG/pNG semantics per type)**.** *Let $(S, [\chi])$ be a sector/type pair chosen from Definitions 9.195 and 9.196 and evaluated on the diagnostic realized reference configuration of Definition 9.185 with typed filter $\Pi_S^{[\chi]}$ as in Definition 9.186. If* $\mathrm{Quad}^{[S,\chi]} = \mathrm{QII}$ *(per Definition 9.189), then the correlational channel $\Pi_{\mathrm{corr}}\,\Pi_S^{[\chi]}$ exhibits an* exact *NG pole with positive residue on $\mathcal{H}_{\mathrm{phys}}$. If* $\mathrm{Quad}^{[S,\chi]} = \mathrm{QIV}$*, the same channel exhibits a pseudo–NG band with GMOR slope on the certified plateau $\mathfrak{P}$. (For* QIII *there is no NG/pNG in the IR;* QI *is fully symmetric.)*

Sketch. The canonical quadrant semantics of §8.6 imply: (i) QII ⇒ state selection without law source, producing an NG pole in the *correlational* channel, and (ii) QIV ⇒ explicit (persistent) law source with GMOR scaling, yielding a pseudo–NG band on the plateau. In the typed setting, we insert $\Pi_S^{[\chi]}$ (see Definition 9.186) into the witness construction $\mathbf{W}_{[\chi]}^{[S]}$ (see Definition 9.187) and into the channel restriction Π_{corr} used by the SynTh meters of §9.16.2. Typed decoupling $\mathcal{O}(\varepsilon_{\mathfrak{P}})$ from other modes follows by Lemma 9.192, so the NG/pNG readout is *per type*. Positivity of the residue in QII is inherited from the projected Hessian on H^0 under the plateau tolerances (Chapter 8), while gauge/Nielsen safety of the statement follows from Proposition 9.191. Thus the untyped NG/pNG semantics of Chapter 9 transfer to the typed channel $\Pi_{\mathrm{corr}}\Pi_S^{[\chi]}$ exactly as stated. □

Definition 9.198 (Typed Quadrant Ledger)**.** The *ledger* has one row per $(S, [\chi])$ containing the sextuple $\mathbf{L}^{[S,\chi]}$ of Corollary 9.194. Each row is obtained by evaluating, on the diagnostic realized reference configuration of Definition 9.185, the typed witness $\mathbf{W}_{[\chi]}^{[S]}$ from Definition 9.187

(built with the typed projectors of Definition 9.186 and the meters of §9.16.2), assigning the typed quadrant via Definition 9.189 (decision pipeline in §9.16.3), and stamping the T–L–O data using Definition 9.190 (bulk–boundary overlay in Definition 9.214). We fix the column order

$$\big[S\big] \quad \big[\chi\big] \quad \big(\mathbf{W}^{[S]}_{[\chi]}\big) \quad \big[\mathrm{Quad}\big] \quad \big[\mathrm{Loc}\big] \quad \big[\mathrm{On}\big],$$

and require stability under the Chapter 8 tolerances (cf. Definition 9.210); boundary rows use $\mathfrak{P}_\partial$ with boundary IR dilation as in Definition 9.214.

Corollary 9.199 (Operational fixing of labels)**.** *Once Definition 9.198 is populated, the* spin/helicity/polarization *labels for each typed mode are fixed* operationally *by the chosen* $[\chi]$ *(from the menus Definitions 9.195 and 9.196), independently of any particle names. Only thereafter may SSB nomenclature be used (e.g.* $(\Phi\Phi, \mathrm{TT})$ *in QII* $\to$ *"graviton";* $(\Theta\Theta, \mathrm{L})$ *or Chrono-Kymon in QII/QIV* $\to$ *Higgs/NG/pNG), consistently with the typed NG/pNG semantics of Proposition 9.197.*

Proof. By Definition 9.186, the typed filter $\Pi^{[\chi]}_S$ selects a BRST-closed, cone-preserving subspace whose kinematic content (spin, helicity, polarization) is determined by $[\chi]$ (menus in Definitions 9.195 and 9.196). Evaluating the typed witness on the diagnostic realized reference configuration (Definitions 9.185 and 9.187) produces meters that are Nielsen-flat and gauge-safe (Proposition 9.191); the canonical predicate then yields a unique $\mathrm{Quad}^{[S,\chi]}$ (Definition 9.189 and Proposition 9.212). Typed decoupling (Lemma 9.192) ensures that these determinations are independent of other types and sectors at leading order, which suffices under the Chapter 8 tolerances used in Definition 9.210. Therefore, the operational labels (spin/helicity/polarization) are fixed by $[\chi]$ and recorded in the ledger $\mathbf{L}^{[S,\chi]}$ (Definition 9.198 and Corollary 9.194), while particle names applied afterward follow the quadrant semantics (NG/pNG) of Proposition 9.197. □

9.16.1 Typed Subspaces and Projectors

We now *choose typed subspaces* that pin down the degrees of freedom to be labeled in each sector, using the realized geometry $\bar{\Sigma}_{\mu\nu}$ on the diagnostic realized reference configuration of Definition 9.185. All operators below act on $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ and are BRST-closed; Π_α denotes the triadic channel projectors (Chapter 7). When a helicity refinement is invoked, it is understood that a $\bar{\Sigma}$-admissible spatial orientation (e.g. a unit spatial direction $\hat{n}$ from a foliation or a spectral mode direction) has been fixed at the plateau.

Definition 9.200 (Gaiamiti ($\Phi\Phi$) typed subspaces)**.** Let Sym^2 denote the $\bar{\Sigma}$-symmetric rank-2 block of the $\Phi\Phi$ sector on $\mathcal{H}_{\mathrm{phys}}$ (diagnostic realized reference configuration of Definition 9.185; types chosen from the menu Definition 9.195). Define BRST-closed projectors $\Pi^{[\chi]}_{\Phi\Phi} : \mathsf{Sym}^2 \to \mathsf{Sym}^2$ that commute with each triadic channel projector Π_α ($\alpha \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$, cf. Definition 9.185) and preserve the $\bar{\Sigma}$-cones (as required in Definition 9.186):

(i) Trace (spin-0)

The spin-0 trace is

$$\big(\Pi^{[\mathrm{Tr}]}_{\Phi\Phi} h\big)_{\mu\nu} := \frac{1}{D}\,\bar{\Sigma}_{\mu\nu}\,\big(\bar{\Sigma}^{\rho\sigma} h_{\rho\sigma}\big), \qquad h \in \mathsf{Sym}^2,$$

where $D := \bar{\Sigma}^{\mu\nu}\bar{\Sigma}_{\mu\nu}$ is the $\bar{\Sigma}$-trace of the identity (plateau slice; §9.16.1).

(ii) TT-like (spin-2, trace-free symmetric)

For TT-like spin-2, trace-free symmetric

$$\Pi^{[\mathrm{TT}]}_{\Phi\Phi} := \mathbb{1}_{\mathsf{Sym}^2} - \Pi^{[\mathrm{Tr}]}_{\Phi\Phi}, \qquad \bar{\Sigma}^{\mu\nu}\big(\Pi^{[\mathrm{TT}]}_{\Phi\Phi} h\big)_{\mu\nu} = 0.$$

If a covariant spatial connection is fixed on $\bar{\Sigma}$ at the plateau, one may additionally impose transversality; the projector remains BRST-closed and cone-preserving, consistent with Definition 9.186.

(iii) Helicity refinement (optional)

When a $\bar{\Sigma}$-spatial direction $\hat{n}$ is available on the diagnostic realized reference configuration, let $\mathcal{J}_{\hat{n}}$ be the generator of rotations about $\hat{n}$ restricted to the TT subspace. Define the $\lambda = \pm 2$ helicity projectors by functional calculus on $\mathcal{J}_{\hat{n}}$:

$$\Pi^{[\lambda]}_{\Phi\Phi} := \Pi^{[\mathrm{TT}]}_{\Phi\Phi}\, \mathbf{1}_{\{\mathcal{J}_{\hat{n}}=\lambda\}}\, \Pi^{[\mathrm{TT}]}_{\Phi\Phi}, \qquad \lambda = \pm 2,$$

so that $\Pi^{[\mathrm{TT}]}_{\Phi\Phi} = \Pi^{[+2]}_{\Phi\Phi} + \Pi^{[-2]}_{\Phi\Phi}$ with orthogonality and completeness on $\operatorname{Ran}\Pi^{[\mathrm{TT}]}_{\Phi\Phi}$ (see Proposition 9.202).

These projectors serve as the typed filters for $S = \Phi\Phi$ in Definition 9.186, are inserted into the typed witness $\mathbf{W}^{[S]}_{[\chi]}$ of Definition 9.187, and thereby feed the meters of §9.16.2 and the quadrant predicate Definition 9.189.

Definition 9.201 (Tantrakyma ($\Theta\Theta$) typed subspaces)**.** Let Vec denote the $\bar{\Sigma}$-covector (or vector) block of the $\Theta\Theta$ sector on $\mathcal{H}_{\mathrm{phys}}$ (diagnostic realized reference configuration of Definition 9.185; types chosen from the menu Definition 9.196). Fix a unit timelike field u^μ (realized rest frame) with $\bar{\Sigma}_{\mu\nu}u^\mu u^\nu = -1$, and set $h_{\mu\nu} := \bar{\Sigma}_{\mu\nu} + u_\mu u_\nu$ for the induced spatial metric. Define BRST-closed projectors $\Pi^{[\chi]}_{\Theta\Theta} : \mathsf{Vec} \to \mathsf{Vec}$ that commute with each triadic projector Π_α ($\alpha \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$, cf. Definition 9.185) and preserve $\bar{\Sigma}$-cones (as required in Definition 9.186):

(i) Chrono-Kymon (temporal/torsional, spin-1 precursor)

Then the chrono-kymon is

$$\big(\Pi^{[\mathrm{CK}]}_{\Theta\Theta}\,\Theta\big)_\mu := -\,u_\mu\,(u^\nu\Theta_\nu), \qquad \Theta \in \mathsf{Vec}.$$

This isolates the time-like (torsional) channel singled out by u^μ (ultimately tied to the temporal piece of $\bar{\Sigma}$; see diagnostic realized reference configuration geometry in Definition 9.185).

(ii) Spatial L/T (Helmholtz) decomposition

And the longitudinal or transverse

$$\Theta^{\mathrm{sp}}_\mu := h_\mu{}^\nu\Theta_\nu, \qquad \Theta^{[L]}_\mu := \bar{\nabla}_\mu\varphi, \quad \Theta^{[T]}_\mu := \Theta^{\mathrm{sp}}_\mu - \bar{\nabla}_\mu\varphi,$$

where $\bar{\nabla}$ is the Levi–Civita connection of h at the plateau and $\varphi := \Delta^{-1}(\bar{\nabla}^\rho\Theta^{\mathrm{sp}}_\rho)$ uses the spatial Laplacian $\Delta := \bar{\nabla}^\rho\bar{\nabla}_\rho$ with plateau boundary conditions. Define projectors $\Pi^{[L]}_{\Theta\Theta}\Theta := \Theta^{[L]}$ and $\Pi^{[T]}_{\Theta\Theta}\Theta := \Theta^{[T]}$. Then $h^{\mu\nu}\bar{\nabla}_\mu\Theta^{[T]}_\nu = 0$, and

$$\Pi^{[\mathrm{CK}]}_{\Theta\Theta} + \Pi^{[T]}_{\Theta\Theta} + \Pi^{[L]}_{\Theta\Theta} = \mathbb{1}_{\mathsf{Vec}}, \qquad \Pi^{[i]}_{\Theta\Theta}\Pi^{[j]}_{\Theta\Theta} = \delta^{ij}\Pi^{[i]}_{\Theta\Theta}\ (i,j \in \{\mathrm{CK}, T, L\}),$$

with orthogonality/completeness summarized in Proposition 9.202.

(iii) Helicity refinement (transverse, $\lambda = \pm$)

When a spatial orientation $\hat{n}$ (or mode direction) is fixed on the diagnostic realized reference configuration, let $\mathcal{J}_{\hat{n}}$ be the generator of rotations about $\hat{n}$ restricted to $\operatorname{Ran}\Pi^{[T]}_{\Theta\Theta}$. Define

$$\Pi^{[\lambda]}_{\Theta\Theta} := \Pi^{[T]}_{\Theta\Theta}\,\mathbf{1}_{\{\mathcal{J}_{\hat{n}}=\lambda\}}\,\Pi^{[T]}_{\Theta\Theta}, \qquad \lambda = \pm 1,$$

so that $\Pi^{[T]}_{\Theta\Theta} = \Pi^{[+]}_{\Theta\Theta} + \Pi^{[-]}_{\Theta\Theta}$ with orthogonality and completeness on the transverse subspace (cf. Proposition 9.202).

These projectors provide the typed filters for $S = \Theta\Theta$ in Definition 9.186, are inserted into the typed witness $\mathbf{W}^{[S]}_{[\chi]}$ of Definition 9.187, and thereby feed the meters of §9.16.2 and the typed quadrant predicate Definition 9.189.

Proposition 9.202 (Orthogonality, completeness, and triadic compatibility)**.** *The projectors of Definitions 9.200 and 9.201 satisfy:*

(i) Gaiamiti

$\Pi^{[\mathrm{Tr}]}_{\Phi\Phi} + \Pi^{[\mathrm{TT}]}_{\Phi\Phi} = \mathbb{1}_{\mathsf{Sym}^2}$ *with* $\Pi^{[\mathrm{Tr}]}\Pi^{[\mathrm{TT}]} = 0$*; any helicity refinement obeys* $\Pi^{[\mathrm{TT}]} = \Pi^{[+2]}_{\Phi\Phi} + \Pi^{[-2]}_{\Phi\Phi}$ *and orthogonality on* $\operatorname{Ran}\Pi^{[\mathrm{TT}]}_{\Phi\Phi}$ *(see construction in Definition 9.200).*

(ii) Tantrakyma

$\Pi^{[\mathrm{CK}]}_{\Theta\Theta} + \Pi^{[T]}_{\Theta\Theta} + \Pi^{[L]}_{\Theta\Theta} = \mathbb{1}_{\mathsf{Vec}}$ *with pairwise orthogonality; any helicity refinement obeys* $\Pi^{[T]}_{\Theta\Theta} = \Pi^{[+]}_{\Theta\Theta} + \Pi^{[-]}_{\Theta\Theta}$ *and orthogonality on* $\operatorname{Ran}\Pi^{[T]}_{\Theta\Theta}$ *(see Definition 9.201).*

(iii) Triadic compatibility

$\Pi_\alpha \Pi^{[\chi]}_S = \Pi^{[\chi]}_S \Pi_\alpha$ *for all* $\alpha \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$ *(triadic channels fixed on the diagnostic realized reference configuration of Definition 9.185); consequently, typed readouts can be taken channel-by-channel without ambiguity (cf. the usage in §9.16.2).*

(iv) Cone preservation

Each $\Pi^{[\chi]}_S$ *preserves the* $\bar{\Sigma}$*-cones as required in Definition 9.186, and thus preserves the principal symbols relevant for the canonical decision rule (§8.6; typed predicate in Definition 9.189).*

Proof. **(1) Gaiamiti block.** Work on the diagnostic realized reference configuration of Definition 9.185 and endow Sym^2 with the $\bar{\Sigma}$-inner product $\langle h, k\rangle_{\bar{\Sigma}} := \bar{\Sigma}^{\mu\alpha}\bar{\Sigma}^{\nu\beta}h_{\mu\nu}k_{\alpha\beta}$. By Definition 9.200, the trace projector is $\left(\Pi^{[\mathrm{Tr}]}_{\Phi\Phi}h\right)_{\mu\nu} = \frac{1}{D}\bar{\Sigma}_{\mu\nu}\,(\bar{\Sigma}^{\rho\sigma}h_{\rho\sigma})$ with $D = \bar{\Sigma}^{\mu\nu}\bar{\Sigma}_{\mu\nu}$. A direct computation shows idempotence and self-adjointness: $(\Pi^{[\mathrm{Tr}]})^2 = \Pi^{[\mathrm{Tr}]}$ and $\langle \Pi^{[\mathrm{Tr}]}h, k\rangle_{\bar{\Sigma}} = \langle h, \Pi^{[\mathrm{Tr}]}k\rangle_{\bar{\Sigma}}$. Define $\Pi^{[\mathrm{TT}]} := \mathbb{1} - \Pi^{[\mathrm{Tr}]}$. Then $(\Pi^{[\mathrm{TT}]})^2 = \Pi^{[\mathrm{TT}]}$, $\Pi^{[\mathrm{Tr}]}\Pi^{[\mathrm{TT}]} = 0$, and $\Pi^{[\mathrm{Tr}]} + \Pi^{[\mathrm{TT}]} = \mathbb{1}_{\mathsf{Sym}^2}$, proving orthogonality and completeness for the TT/Tr split, as stated in item (1). Moreover, $\bar{\Sigma}^{\mu\nu}(\Pi^{[\mathrm{TT}]}h)_{\mu\nu} = 0$, so TT is the $\bar{\Sigma}$-orthogonal complement of Tr.

For the helicity refinement (when a $\bar{\Sigma}$-admissible spatial direction $\hat{n}$ exists on the plateau), Definition 9.200 defines $\mathcal{J}_{\hat{n}}$ on $\operatorname{Ran}\Pi^{[\mathrm{TT}]}$. The generator $\mathcal{J}_{\hat{n}}$ is essentially self-adjoint on the TT subspace with respect to $\langle\cdot,\cdot\rangle_{\bar{\Sigma}}$, hence by the spectral theorem its spectral projectors $\mathbf{1}_{\{\mathcal{J}_{\hat{n}}=\lambda\}}$ are orthogonal idempotents that resolve the identity on that subspace. Setting

$\Pi^{[\lambda]}_{\Phi\Phi} = \Pi^{[\mathrm{TT}]}\mathbf{1}_{\{\mathcal{J}_{\hat{n}}=\lambda\}}\Pi^{[\mathrm{TT}]}$ ($\lambda = \pm 2$) gives $\Pi^{[\mathrm{TT}]} = \Pi^{[+2]} + \Pi^{[-2]}$ with orthogonality on $\operatorname{Ran}\Pi^{[\mathrm{TT}]}$, establishing the helicity clause of item (1).

(2) Tantrakyma block. On Vec (diagnostic realized reference configuration of Definition 9.185) choose the $\bar{\Sigma}$-inner product $\langle A, B\rangle_{\bar{\Sigma}} := \bar{\Sigma}^{\mu\nu}A_\mu B_\nu$. With u^μ and $h_{\mu\nu} = \bar{\Sigma}_{\mu\nu} + u_\mu u_\nu$ from Definition 9.201, the Chrono–Kymon projector is $\left(\Pi^{[\mathrm{CK}]}\Theta\right)_\mu = -u_\mu(u^\nu\Theta_\nu)$. Then $(\Pi^{[\mathrm{CK}]})^2 = \Pi^{[\mathrm{CK}]}$ and $\langle\Pi^{[\mathrm{CK}]}\Theta, \Phi\rangle_{\bar{\Sigma}} = \langle\Theta, \Pi^{[\mathrm{CK}]}\Phi\rangle_{\bar{\Sigma}}$; further, $h_\mu{}^\nu u_\nu = 0$ implies $\langle\Pi^{[\mathrm{CK}]}\Theta, h\Phi\rangle_{\bar{\Sigma}} = 0$, so the temporal CK channel is orthogonal to the spatial block.

For the spatial Helmholtz/Hodge decomposition on the plateau domain (with Chapter 8 boundary conditions), define $\Theta^{\mathrm{sp}}_\mu = h_\mu{}^\nu\Theta_\nu$ and $\varphi := \Delta^{-1}(\bar{\nabla}^\rho\Theta^{\mathrm{sp}}_\rho)$ where $\Delta = \bar{\nabla}^\rho\bar{\nabla}_\rho$. Set $\Theta^{[L]}_\mu := \bar{\nabla}_\mu\varphi$ and $\Theta^{[T]}_\mu := \Theta^{\mathrm{sp}}_\mu - \bar{\nabla}_\mu\varphi$ (Definition 9.201). Standard Hodge theory on the certified plateau then yields $h^{\mu\nu}\bar{\nabla}_\mu\Theta^{[T]}_\nu = 0$, $\langle\Theta^{[L]}, \Theta^{[T]}\rangle_{\bar{\Sigma}} = 0$, and $\Theta^{\mathrm{sp}} = \Theta^{[L]} + \Theta^{[T]}$. Thus the operators $\Pi^{[L]}\Theta = \Theta^{[L]}$, $\Pi^{[T]}\Theta = \Theta^{[T]}$ are orthogonal idempotents that resolve the identity on the spatial block. Combining with CK gives $\Pi^{[\mathrm{CK}]} + \Pi^{[T]} + \Pi^{[L]} = \mathbb{1}_{\mathsf{Vec}}$ and pairwise orthogonality, proving item (2). When a spatial orientation $\hat{n}$ is available, the restricted rotation generator $\mathcal{J}_{\hat{n}}$ on $\operatorname{Ran}\Pi^{[T]}$ is self-adjoint; its spectral projectors $\Pi^{[\pm]} = \Pi^{[T]}\mathbf{1}_{\{\mathcal{J}_{\hat{n}}=\pm 1\}}\Pi^{[T]}$ are orthogonal idempotents with $\Pi^{[T]} = \Pi^{[+]} + \Pi^{[-]}$ (again as in Definition 9.201).

(3) Triadic compatibility. By Definition 9.186, each typed projector $\Pi^{[\chi]}_S$ is defined from $\bar{\Sigma}$, u^μ, $h_{\mu\nu}$, and the plateau connection $\bar{\nabla}$, all fixed by the diagnostic realized reference configuration (Definition 9.185). The triadic channel projectors Π_α act on the correlation/-vacuum/spectral channels (Chapter 7) and commute with these kinematical constructions; hence $\Pi_\alpha\Pi^{[\chi]}_S = \Pi^{[\chi]}_S\Pi_\alpha$ for all α, establishing item (3) and justifying channel-wise typed readouts (cf. §9.16.2).

(4) Cone preservation. The operators $\Pi^{[\chi]}_S$ are $\bar{\Sigma}$-intrinsic algebraic (TT/Tr, CK) or elliptic (L/T via Δ^{-1}) filters with smooth symbols that project onto subbundles invariant under the $\bar{\Sigma}$-lightcone structure. Therefore their principal symbols are the usual momentum-space projectors (e.g. $p_\mu p_\nu/|p|^2$ for L and $h_{\mu\nu} - p_\mu p_\nu/|p|^2$ for T), which preserve $\bar{\Sigma}$-cones and do not induce cross-cone leakage at leading order; this is the cone-preservation clause of Definition 9.186. Consequently, the principal symbols relevant for the Chapter 9 canonical decision rule (§8.6, typed form Definition 9.189) are stable under insertion of $\Pi^{[\chi]}_S$, yielding item (4).

Combining the four parts proves Proposition 9.202. □

Remark 9.203 (Use in witnesses and meters)**.** For any sector/type pair $(S, [\chi])$, the *typed witness* $\mathbf{W}^{[S]}_{[\chi]}$ is defined in Definition 9.187 and is evaluated on the diagnostic realized reference configuration of Definition 9.185. Operationally, one inserts the typed projector $\Pi^{[\chi]}_S$ (see Definition 9.186, with concrete constructions in Definitions 9.200 and 9.201) together with the triadic channel projectors $\{\Pi_\alpha\}$ into the channel-restricted readouts, following the procedure of §9.16.2. Triadic compatibility and cone preservation (Proposition 9.202) ensure that the order of insertion is immaterial and that the resulting SynTh meters are well posed and Nielsen-flat (Proposition 9.191; cf. Proposition 9.207). Helicity refinements are *optional* and are invoked only when a $\bar{\Sigma}$-admissible spatial direction is fixed at the plateau (see the helicity clauses in Definitions 9.200 and 9.201). The typed meters thus obtained feed the typed quadrant predicate of Definition 9.189 and, together with the location/onset data of Definition 9.190 (bulk–boundary overlay in Definition 9.214), populate the corresponding

row in the Typed Quadrant Ledger $\mathbf{L}^{[S,\chi]}$ (Definition 9.198 and Corollary 9.194).

9.16.2 Applying Parasymmetric Meters to Typed Subspaces

We now *evaluate* the Chapter 9 parasymmetric meters on each typed subspace chosen in §9.16.1. All quantities are read on the diagnostic realized reference configuration of Definition 9.185 and on the BRST-physical sector $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$. Typed filters $\Pi_S^{[\chi]}$ (from Definitions 9.186, 9.200 and 9.201) and triadic channel projectors Π_α (Chapter 7) are inserted in every readout. We follow the *Simultaneous Synthesis* (SynTh) notation of Chapter 9, using the sector-conditioned witness $\mathbf{W}_{[\chi]}^{[S]}$ with components

$$\Big(v^{(L)}_{\text{Synthesis}},\ v^{(S)}_{\text{Synthesis}},\ v^{(LS)}_{\text{Synthesis}}\Big)^{[S,\chi]}(\ell;\mathfrak{P}),$$

which drive the quadrant decision of §8.6.

Definition 9.204 (Local parasymmetric two–mode reduction)**.** Fix a certified plateau cell (scale/time) on the diagnostic realized reference configuration of Definition 9.185 and a sector/type pair $(S,[\chi])$ (types chosen from Definitions 9.195 and 9.196). Let $\mathcal{H}_{\text{loc}}^{[S,\chi]} \subset \mathcal{H}_{\text{phys}}$ be the two–mode subspace singled out by the typed projector $\Pi_S^{[\chi]}$ of Definition 9.186 and the SynTh channel predicate (Law, State, or Law⊕State) used in §9.16.2:

$$\mathcal{H}_{\text{loc}}^{[S,\chi]} := \operatorname{Ran}\Big(\Pi_S^{[\chi]}\,\Pi_\alpha\Big)\ \restriction_{\text{SynTh channel}}, \qquad \alpha \in \{\text{corr},\text{vac},\text{spec}\}\ \text{(cf. Definition 9.185)}.$$

Choose an orthonormal pair $\{|\psi_1\rangle, |\psi_2\rangle\}$ spanning $\mathcal{H}_{\text{loc}}^{[S,\chi]}$. For each $a \in \{L,S,LS\}$ (Law, State, Law⊕State) define the *reduced density matrix*

$$\rho_{ij}^{(a)\,[S,\chi]}(\ell) \;:=\; \frac{\big\langle \psi_i,\ \Pi_S^{[\chi]}\,\Pi_\alpha\,\mathcal{O}^{(a)}(\ell)\,\Pi_\alpha\,\Pi_S^{[\chi]}\ \psi_j\big\rangle}{\sum_{k=1}^{2}\big\langle \psi_k,\ \Pi_S^{[\chi]}\,\Pi_\alpha\,\mathcal{O}^{(a)}(\ell)\,\Pi_\alpha\,\Pi_S^{[\chi]}\ \psi_k\big\rangle}, \qquad i,j\in\{1,2\},$$

where $\mathcal{O}^{(a)}(\ell)$ is the SynTh channel operator whose expectation yields the valuations in Definition 9.187 (see also the diagnostic realized reference configuration aggregation and significance criteria in §9.16.2 and Definition 9.210). This local two–mode reduction underlies the typed meters $v^{(a)[S,\chi]}_{\text{Synthesis,loc}}(\ell)$ of Definition 9.205 and enjoys the channel/type compatibility properties stated in Proposition 9.207.

Definition 9.205 (Typed parasymmetric meters (local form))**.** For a fixed sector/type pair $(S,[\chi])$ on the diagnostic realized reference configuration of Definition 9.185, let $\rho_{ij}^{(a)\,[S,\chi]}(\ell)$ be the reduced density matrix from the local two–mode construction of Definition 9.204, with $a \in \{L,S,LS\}$ denoting the SynTh channel (Law, State, Law⊕State; cf. §9.16.2). The *local* SynTh valuations are

$$\begin{aligned}&\Big(v^{(L)}_{\text{Synthesis}}, v^{(S)}_{\text{Synthesis}}, v^{(LS)}_{\text{Synthesis}}\Big)^{[S,\chi]}_{\text{loc}}(\ell) :=\\ &\quad\Big(\tfrac{1}{2}+\operatorname{Re}\,\rho_{12}^{(L)\,[S,\chi]}(\ell),\ \tfrac{1}{2}+\operatorname{Re}\,\rho_{12}^{(S)\,[S,\chi]}(\ell),\ \tfrac{1}{2}+\operatorname{Re}\,\rho_{12}^{(LS)\,[S,\chi]}(\ell)\Big),\end{aligned}$$

which obey the bounds and baselines of Lemma 9.206. The *diagnostic realized reference configuration* valuations used by the typed witness $\mathbf{W}^{[S]}_{[\chi]}$ (see Definition 9.187) are obtained by the Chapter 8 aggregation $\mathcal{A}_{\mathfrak{P}}$ across the certification window of the chosen plateau $\mathfrak{P}$ (significance/tolerances in Definition 9.210):

$$\left(v^{(L)}_{\text{Synthesis}},\, v^{(S)}_{\text{Synthesis}},\, v^{(LS)}_{\text{Synthesis}}\right)^{[S,\chi]}(\ell;\mathfrak{P}) \;:=\; \mathcal{A}_{\mathfrak{P}}\Big[\, \left(v^{(L)}_{\text{Synthesis}},\, v^{(S)}_{\text{Synthesis}},\, v^{(LS)}_{\text{Synthesis}}\right)^{[S,\chi]}_{\text{loc}}(\ell') \,\Big]_{\ell'\in\mathfrak{P}}.$$

These meters are Nielsen-flat and compatible with channel/type insertions (Propositions 9.191 and 9.207), and they feed the typed quadrant predicate of Definition 9.189.

Lemma 9.206 (Bounds, baselines, and phase insensitivity)**.** *For each $a \in \{L, S, LS\}$ and $(S, [\chi])$:*

(i) $0 \le v^{(a)[S,\chi]}_{\text{Synthesis,loc}}(\ell) \le 1$ *(and hence the same bounds hold for the diagnostic realized reference configuration–aggregated meters defined in Definition 9.205).*

(ii) $v^{(a)[S,\chi]}_{\text{Synthesis,loc}}(\ell) = \frac{1}{2}$ *iff the two–mode reduction is incoherent, i.e.* $\operatorname{Re}\rho^{(a)[S,\chi]}_{12}(\ell) = 0$ *(see Definitions 9.204 and 9.205).*

(iii) $v^{(a)[S,\chi]}_{\text{Synthesis,loc}}(\ell) = 1$ *iff the two–mode reduction is maximally (Bell–)coherent with* $\operatorname{Re}\rho^{(a)[S,\chi]}_{12}(\ell) = \frac{1}{2}$ *and equal marginals* $\rho_{11} = \rho_{22} = \frac{1}{2}$*;* $v^{(a)[S,\chi]}_{\text{Synthesis,loc}}(\ell) = 0$ *iff* $\operatorname{Re}\rho^{(a)[S,\chi]}_{12}(\ell) = -\frac{1}{2}$ *with the same equal–marginal condition.*

(iv) *The value depends only on the* relative *phase of $|\psi_1\rangle, |\psi_2\rangle$ chosen in Definition 9.204; a common $U(1)$ rephasing leaves $v^{(a)[S,\chi]}_{\text{Synthesis,loc}}(\ell)$ invariant.*

Proof. Work on a fixed plateau cell of the diagnostic realized reference configuration (Definition 9.185) and a fixed sector/type pair $(S, [\chi])$. By Definition 9.204, the (typed, channel–restricted) two–mode reduction produces a 2×2 positive semidefinite density matrix $\rho^{(a)[S,\chi]}(\ell)$ with unit trace, thanks to the normalization in the denominator. The local SynTh meter is $v^{(a)[S,\chi]}_{\text{Synthesis,loc}}(\ell) = \frac{1}{2} + \operatorname{Re}\rho^{(a)[S,\chi]}_{12}(\ell)$ (Definition 9.205).

(1) Bounds. Positivity implies $\det\rho^{(a)} \ge 0$, i.e. $\rho_{11}\rho_{22} - |\rho_{12}|^2 \ge 0$, hence $|\rho_{12}| \le \sqrt{\rho_{11}\rho_{22}}$. Since $\rho_{11} + \rho_{22} = 1$, AM–GM gives $\sqrt{\rho_{11}\rho_{22}} \le \frac{1}{2}$, so $|\rho_{12}| \le \frac{1}{2}$. Therefore $-\frac{1}{2} \le \operatorname{Re}\rho_{12} \le \frac{1}{2}$ and $0 \le v^{(a)[S,\chi]}_{\text{Synthesis,loc}}(\ell) \le 1$. Diagnostic realized reference configuration aggregation over $\mathfrak{P}$ (Chapter 8; Definitions 9.205 and 9.210) preserves these bounds by convexity.

(2) Baseline. By definition $v = \frac{1}{2} + \operatorname{Re}\rho_{12}$; hence $v = \frac{1}{2}$ iff $\operatorname{Re}\rho_{12} = 0$, which is the usual incoherence condition in the selected two–mode subspace (off–diagonal purely imaginary or vanishing).

(3) Extremes and Bell coherence. From $|\rho_{12}| \le \sqrt{\rho_{11}\rho_{22}} \le \frac{1}{2}$, the extremal values $v = 1$ or $v = 0$ require $|\rho_{12}| = \frac{1}{2}$. Equality in Cauchy–Schwarz (or the positivity bound) occurs iff $\rho_{11} = \rho_{22} = \frac{1}{2}$ and ρ_{12} is aligned in phase with the chosen basis so that $\operatorname{Re}\rho_{12} = \pm\frac{1}{2}$. Thus $v = 1$ (resp. $v = 0$) iff $\operatorname{Re}\rho_{12} = +\frac{1}{2}$ (resp. $-\frac{1}{2}$) and $\rho_{11} = \rho_{22} = \frac{1}{2}$, i.e. a maximally coherent (Bell–like) two–mode reduction.

(4) Phase insensitivity (relative vs. common). Under a basis rephasing $|\psi_j\rangle \mapsto e^{i\phi_j}|\psi_j\rangle$ (within the typed, channel–selected subspace of Definition 9.204), one has $\rho_{12} \mapsto e^{i(\phi_1-\phi_2)}\rho_{12}$. Hence $v = \frac{1}{2} + \operatorname{Re}\rho_{12}$ depends only on the *relative* phase $\phi_1 - \phi_2$; in particular, a common $U(1)$ phase $\phi_1 = \phi_2$ leaves v unchanged.

These statements are insensitive to gauge–fixing (Nielsen–flat) because the operators inserted are BRST–closed and evaluated on H^0 (Proposition 9.191); channel/type insertion order does not affect the result by Proposition 9.207. □

Proposition 9.207 (Channel and type compatibility)**.** *Let $(S,[\chi])$ be fixed (with $[\chi]$ taken from Definitions 9.195 and 9.196) and let $a \in \{L,S,LS\}$ denote the SynTh channel. On the diagnostic realized reference configuration of Definition 9.185,*

$$\Pi_\alpha\,\Pi_S^{[\chi]} \;\rightsquigarrow\; \left(v^{(a)}_{\mathrm{Synthesis}}\right)^{[S,\chi]}(\ell;\mathfrak{P}) \quad \textit{is well defined, Nielsen–flat, and independent of the order of insertion of } \Pi_\alpha, \Pi_S^{[\chi]}.$$

Moreover, if $[\chi'] \neq [\chi]$ for the same sector S, then $\left(v^{(a)}_{\mathrm{Synthesis}}\right)^{[S,\chi]}$ and $\left(v^{(a)}_{\mathrm{Synthesis}}\right)^{[S,\chi']}$ decouple at leading order on $\mathfrak{P}$: cross–terms are $\mathcal{O}(\varepsilon_{\mathfrak{P}})$, where $\varepsilon_{\mathfrak{P}}$ is the Chapter 8 tolerance scale (cf. Definition 9.210). The same statements hold on the boundary diagnostic realized reference configuration $\mathfrak{P}_\partial$.

Proof. **Well-definedness and order independence.** By construction of the typed projectors (Definition 9.186) and the diagnostic realized reference configuration (Definition 9.185), we have triadic compatibility $\Pi_\alpha\Pi_S^{[\chi]} = \Pi_S^{[\chi]}\Pi_\alpha$ (see also Proposition 9.202, item 3). Hence the composed insertion $\Pi_\alpha\Pi_S^{[\chi]}$ is an idempotent, self-adjoint filter on $\mathcal{H}_{\mathrm{phys}}$, independent of insertion order. The local two–mode reduction (Definition 9.204) therefore yields a normalized 2×2 density matrix $\rho^{(a)[S,\chi]}(\ell)$, and the local meters $v^{(a)[S,\chi]}_{\mathrm{Synthesis,loc}}(\ell)$ are defined by Definition 9.205. Aggregating across the certified window by $\mathcal{A}_{\mathfrak{P}}$ (Chapter 8; Definitions 9.205 and 9.210) produces the diagnostic realized reference configuration valuations $\left(v^{(a)}_{\mathrm{Synthesis}}\right)^{[S,\chi]}(\ell;\mathfrak{P})$, establishing well-definedness.

Nielsen flatness (gauge independence). The SynTh readouts are BRST-closed on H^0 and evaluated on a certified plateau; therefore their gauge-parameter derivatives insert BRST-exact operators whose expectations vanish on $\mathcal{H}_{\mathrm{phys}}$. This is the content of Proposition 9.191 (see also the discussion around §9.16.2), which directly applies to the typed meters because the only change is the insertion of the commuting, cone-preserving projector $\Pi_S^{[\chi]}$ (Definition 9.186). Hence the valuations are Nielsen-flat.

Typed decoupling. For two distinct types $[\chi] \neq [\chi']$ within the same sector, orthogonality/completeness of the typed decompositions was established in Proposition 9.202 (items 1–2). Consequently, the channel-restricted principal symbols are block-diagonal at the plateau. A Duhamel/BCH control of subleading commutators across the certification window (Chapter 8) yields that mixed contributions fall below tolerance, which is summarized in Lemma 9.192 as $\mathcal{O}(\varepsilon_{\mathfrak{P}})$. Therefore the diagnostic realized reference configuration valuations computed with $\Pi_S^{[\chi]}$ are insensitive, at leading order, to the presence of orthogonal typed filters $\Pi_S^{[\chi']}$, proving the stated decoupling.

All statements carry over verbatim to the boundary diagnostic realized reference configuration by replacing $\mathfrak{P}$ with $\mathfrak{P}_\partial$ and using the edge dilation/overlay (cf. Definition 9.214). □

Definition 9.208 (Quadrant drivers)**.** Fix a sector/type pair $(S,[\chi])$ (types from Definitions 9.195 and 9.196) and an diagnostic realized reference configuration as in Definition 9.185.

Let

$$\mathbf{V}^{[S,\chi]}(\ell;\mathfrak{P}) := \Big(v^{(L)}_{\text{Synthesis}},\ v^{(S)}_{\text{Synthesis}},\ v^{(LS)}_{\text{Synthesis}} \Big)^{[S,\chi]}(\ell;\mathfrak{P})$$

be the typed SynTh meters obtained from the typed witness $\mathbf{W}^{[S]}_{[\chi]}$ (Definition 9.187) via the diagnostic realized reference configuration aggregation of Definition 9.205 (constructed locally by Definition 9.204 and used operationally in §9.16.2). Together with the typed healing indicators $\big(\mathscr{H}_L^{[S,\chi]}, \mathscr{H}_S^{[S,\chi]}\big)$ of Definition 9.188, this triplet $\mathbf{V}^{[S,\chi]}$ *drives* the canonical typed quadrant predicate $\mathrm{Quad}^{[S,\chi]}$ in Definition 9.189, evaluated on the same certified plateau (bulk $\mathfrak{P}$ or boundary $\mathfrak{P}_\partial$).

Corollary 9.209 (Typed parasymmetric meters determine the quadrant)**.** *On any certified $\mathfrak{P}$ (or $\mathfrak{P}_\partial$), the data $\big(\mathbf{V}^{[S,\chi]},\ \mathscr{H}_L^{[S,\chi]},\ \mathscr{H}_S^{[S,\chi]}\big)$ are sufficient to evaluate the predicate of Definition 9.189 and thus to assign $\mathrm{Quad}^{[S,\chi]} \in \{\mathrm{QI}, \mathrm{QII}, \mathrm{QIII}, \mathrm{QIV}\}$. Consequently, these determinations populate the Typed Quadrant Ledger entry $\mathbf{L}^{[S,\chi]}$ (cf. Definition 9.198 and Corollary 9.194).*

Proof. By construction, $\mathbf{V}^{[S,\chi]}$ consists of the diagnostic realized reference configuration SynTh meters defined in Definition 9.205 from the typed witness of Definition 9.187, and $\mathscr{H}_\bullet^{[S,\chi]}$ are the typed healing indicators of Definition 9.188. The canonical predicate in Definition 9.189 depends only on these quantities (together with the fixed Chapter 8 tolerances referenced in Definition 9.210); hence they are *sufficient* to decide the quadrant on the chosen diagnostic realized reference configuration. Well-posedness and gauge/Nielsen safety of the meters and the decision follow from Propositions 9.191 and 9.207, while uniqueness of the resulting label on a certified plateau is ensured by Proposition 9.212. Once $\mathrm{Quad}^{[S,\chi]}$ is fixed, inserting it (along with the already-defined witness and T–L–O data) into the ledger row $\mathbf{L}^{[S,\chi]}$ is exactly the prescription of Definition 9.198, whose existence/uniqueness were established in Corollary 9.194. □

9.16.3 Assigning the Quadrant Per Type (Canonical decision on H^0)

We now *assign the quadrant* for each typed subspace using the Chapter 9 canonical rule, evaluated on the diagnostic realized reference configuration of Definition 9.185. All inputs are the typed SynTh meters and healing indicators introduced in §9.16.2:

$$\mathbf{V}^{[S,\chi]}(\ell;\mathfrak{P}) = \Big(v^{(L)}_{\text{Synthesis}},\ v^{(S)}_{\text{Synthesis}},\ v^{(LS)}_{\text{Synthesis}} \Big)^{[S,\chi]}(\ell;\mathfrak{P}), \qquad \big(\mathscr{H}_L^{[S,\chi]},\ \mathscr{H}_S^{[S,\chi]}\big).$$

Decisions are taken *on the same diagnostic realized reference configuration* and are thus BRST/Nielsen–safe (cf. Proposition 9.191).

Definition 9.210 (Stability and significance on the plateau)**.** Fix a certified plateau $\mathfrak{P}$ (or $\mathfrak{P}_\partial$). Let $\mathrm{Axis}(\mathfrak{P})$ be its control axis (Chapter 8) and let $\mathcal{A}_{\mathfrak{P}}[\cdot]$ denote the Chapter 8 aggregation over the certification window.

1. (*Diagnostic realized reference configuration meters*) Define the diagnostic realized reference configuration values by aggregation:

$$\overline{v}_a^{[S,\chi]} := \mathcal{A}_{\mathfrak{P}}\Big[v^{(a)\,[S,\chi]}_{\text{Synthesis}}(\ell) \Big], \quad a \in \{L, S, LS\}.$$

2. (*Baselines and deviations*) Use the SynTh baseline 1/2 (§9.16.2) and set $\Delta_S^{[S,\chi]} := \left|\overline{v}_S^{[S,\chi]} - \frac{1}{2}\right|$.
3. (*Significance thresholds*) Let $\tau_L, \tau_S > 0$ be the Chapter 8 detection tolerances (Slope/Curv/Osc encoded) for law–source and state–selection, respectively. We write Law $\mathrm{src}^{[S,\chi]}$ if $\overline{v}_L^{[S,\chi]}$ exceeds its symmetric-band acceptance by $> \tau_L$, and State $\mathrm{sel}^{[S,\chi]}$ if $\Delta_S^{[S,\chi]} > \tau_S$.

Definition 9.211 (Typed canonical quadrant predicate)**.** For a fixed sector/type $(S,[\chi])$ on the diagnostic realized reference configuration, define $\mathrm{Quad}^{[S,\chi]} \in \{\mathrm{QI}, \mathrm{QII}, \mathrm{QIII}, \mathrm{QIV}\}$ by:

$$\mathrm{QI}: \quad \neg\text{Law src}^{[S,\chi]} \ \wedge\ \neg\text{State sel}^{[S,\chi]} \iff \mathscr{H}_L^{[S,\chi]} = 0,\ \mathscr{H}_S^{[S,\chi]} = 0,\ \Delta_S^{[S,\chi]} \le \tau_S;$$

$$\mathrm{QII}: \quad \neg\text{Law src}^{[S,\chi]} \ \wedge\ \text{State sel}^{[S,\chi]} \ \wedge\ \mathscr{H}_S^{[S,\chi]} = 0 \iff \mathscr{H}_L^{[S,\chi]} = 0,\ \Delta_S^{[S,\chi]} > \tau_S;$$

$$\mathrm{QIII}: \quad \text{Law src}^{[S,\chi]} \ \wedge\ \mathscr{H}_L^{[S,\chi]} = 0 \iff \text{explicit law source at finite scale, } \textit{healing} \text{ in the IR;}$$

$$\mathrm{QIV}: \quad \text{Law src}^{[S,\chi]} \ \wedge\ \mathscr{H}_L^{[S,\chi]} > 0 \iff \text{explicit law source with } \textit{persistence} \text{ (GMOR slope; pNG band when applicable).}$$

All predicates are evaluated using the same $\mathfrak{P}$ and the same typed filter $\Pi_S^{[\chi]}$ (cf. Definition 9.186).

Proposition 9.212 (Typed quadrant assignment is well defined and BRST/Nielsen–safe)**.** *For each $(S,[\chi])$ and certified $\mathfrak{P}$ (or $\mathfrak{P}_\partial$), the value $\mathrm{Quad}^{[S,\chi]}$ given by Definition 9.211 exists, is independent of gauge–fixing parameters, and is invariant under BRST–exact deformations admissible on $\mathfrak{P}$.*

Proof. Existence and invariance follow from Proposition 9.191 (diagnostic realized reference configuration well–posedness, Nielsen flatness) together with typed decoupling at leading order (Lemma 9.192). The predicates use only diagnostic realized reference configuration quantities $\overline{v}_a^{[S,\chi]}$, $\mathscr{H}_\bullet^{[S,\chi]}$ and Chapter 8 tolerances, all defined on the same diagnostic realized reference configuration. □

Corollary 9.213 (Per–type quadrant label)**.** *For every type $[\chi]$ (spin/helicity/polarization/species) in either sector $S \in \{\Phi\Phi, \Theta\Theta\}$ (types chosen from Definitions 9.195 and 9.196 and implemented via Definition 9.186), we obtain a definite label*

$$\mathrm{Quad}^{[S,\chi]} \in \{\mathrm{QI}, \mathrm{QII}, \mathrm{QIII}, \mathrm{QIV}\}.$$

These labels are the inputs recorded in the Typed Quadrant Ledger entry $\mathbf{L}^{[S,\chi]}$ (see Definition 9.198) and are to be fixed prior *to any SSB/particle nomenclature (cf. Corollary 9.199).*

Proof. Fix an diagnostic realized reference configuration (bulk $\mathfrak{P}$ or boundary $\mathfrak{P}_\partial$) as in Definition 9.185 and a sector/type pair $(S,[\chi])$. The typed witness $\mathbf{W}_{[\chi]}^{[S]}$ exists and is Nielsen–flat by Definition 9.187 and Proposition 9.191; its diagnostic realized reference

configuration meters are constructed in Definition 9.205 and the associated typed healing indicators in Definition 9.188. The canonical predicate that assigns $\mathrm{Quad}^{[S,\chi]}$ is given in Definition 9.189 and, by Proposition 9.212, is well defined and gauge/BRST–safe on any certified plateau.

Uniqueness and per–type definiteness follow from typed orthogonality/triadic compatibility (Proposition 9.202), typed decoupling at leading order (Lemma 9.192), and the Chapter 8 stability tolerances used in Definition 9.210: these ensure that the diagnostic realized reference configuration meters determining the predicate are insensitive to orthogonal types up to errors below certification thresholds. Hence a single, definite quadrant value is produced for each $(S,[\chi])$.

Finally, once $\mathrm{Quad}^{[S,\chi]}$ is fixed, the ledger row $\mathbf{L}^{[S,\chi]}$ is populated per Definition 9.198; existence/uniqueness of the ledger entry was established in Corollary 9.194. The instruction to defer SSB/particle naming until after ledger population is precisely the operational rule stated in Corollary 9.199. □

9.16.4 Recording the Type, Location, and Onset (TLO)

We now *stamp* each typed mode with its Type–Location–Onset (T–L–O) data. We use the same diagnostic realized reference configuration and meters as in §9.16.2, together with the typed quadrant assignment of §9.16.3. Recall the location/onset primitives from Definition 9.190 and write

$$\mathbf{V}^{[S,\chi]}(\ell;\mathfrak{P}) = \left(v^{(L)}_{\mathrm{Synthesis}}, v^{(S)}_{\mathrm{Synthesis}}, v^{(LS)}_{\mathrm{Synthesis}}\right)^{[S,\chi]}(\ell;\mathfrak{P}),$$

with healing indicators $\mathscr{H}^{[S,\chi]}_{\bullet}$ from Definition 9.188.

Definition 9.214 (Bulk and boundary meters (overlay))**.** For each sector/type $(S,[\chi])$ we define the *bulk* and *boundary* SynTh triplets by evaluating the same typed witness on the appropriate diagnostic realized reference configuration:

$$\mathbf{V}^{[S,\chi]}_{\mathrm{bulk}}(\ell) := \mathbf{V}^{[S,\chi]}(\ell;\mathfrak{P}), \qquad \mathbf{V}^{[S,\chi]}_{\partial}(\ell) := \mathbf{V}^{[S,\chi]}(\ell;\mathfrak{P}_{\partial}),$$

together with the corresponding healing indicators $\mathscr{H}^{[S,\chi]}_{\bullet,\mathrm{bulk}}$ and $\mathscr{H}^{[S,\chi]}_{\bullet,\partial}$. The *boundary overlay* is the comparison map

$$\mathcal{O}^{[S,\chi]}_{\partial\leftarrow\mathrm{bulk}} : \left(\mathbf{V}^{[S,\chi]}_{\mathrm{bulk}}, \mathscr{H}^{[S,\chi]}_{\bullet,\mathrm{bulk}}\right) \longrightarrow \left(\mathbf{V}^{[S,\chi]}_{\partial}, \mathscr{H}^{[S,\chi]}_{\bullet,\partial}\right),$$

taken along the boundary IR dilation prescribed in Chapter 9.

Definition 9.215 (Edge status (healing/persistent/spontaneous selection))**.** Fix $(S,[\chi])$. Let $\mathrm{Quad}^{[S,\chi]}_{\mathrm{bulk}}$ (resp. $\mathrm{Quad}^{[S,\chi]}_{\partial}$) be the typed quadrant on $\mathfrak{P}$ (resp. $\mathfrak{P}_{\partial}$), as in Definition 9.211. Define the *edge status* $\mathrm{Edge}^{[S,\chi]} \in \{\mathrm{heal}, \mathrm{persist}, \mathrm{sel}\}$ by:

1. heal if $\mathrm{Quad}^{[S,\chi]}_{\partial} \in \{\mathrm{QI}, \mathrm{QIII}\}$ with $\mathscr{H}^{[S,\chi]}_{L,\partial} = 0$ (law heals at the edge);
2. persist if $\mathrm{Quad}^{[S,\chi]}_{\partial} = \mathrm{QIV}$ with $\mathscr{H}^{[S,\chi]}_{L,\partial} > 0$ (explicit breaking persists at the edge);

3. sel (*spontaneous edge selection*) if $\mathrm{Quad}^{[S,\chi]}_{\mathrm{bulk}} = \mathrm{QI}$ and $\mathrm{Quad}^{[S,\chi]}_{\partial} = \mathrm{QII}$ with $\mathscr{H}^{[S,\chi]}_{S,\partial} = 0$ (edge-only SSB, bulk symmetric).

Definition 9.216 (T–L–O stamp)**.** For each typed mode $(S,[\chi])$ we define its *T–L–O stamp* as the triplet

$$\mathbf{TLO}^{[S,\chi]} := \Big(\underbrace{\mathrm{Type}}_{[\chi]},\ \underbrace{\mathrm{Loc}^{[S,\chi]}}_{\text{bulk or boundary}},\ \underbrace{\mathrm{On}^{[S,\chi]}}_{\text{scale/time onset}} \Big),$$

where $\mathrm{Loc}^{[S,\chi]}$ and $\mathrm{On}^{[S,\chi]}$ are as in Definition 9.190. For boundary rows we also record the edge status $\mathrm{Edge}^{[S,\chi]}$ from Definition 9.215 as auxiliary metadata.

Proposition 9.217 (Existence, uniqueness, and stability of T–L–O)**.** *For any certified plateau (bulk $\mathfrak{P}$ or boundary $\mathfrak{P}_\partial$) and any type $[\chi]$ in sector S, the stamp $\mathbf{TLO}^{[S,\chi]}$ of Definition 9.216 exists, is unique, and is stable under Chapter 8 tolerances. In the boundary case, the auxiliary $\mathrm{Edge}^{[S,\chi]}$ is well defined by Definition 9.215.*

Proof. Existence/uniqueness of $\mathrm{Loc}^{[S,\chi]}$ and $\mathrm{On}^{[S,\chi]}$ follow from Definition 9.190 once $\mathrm{Quad}^{[S,\chi]}$ is fixed (cf. Proposition 9.212). Stability is inherited from the aggregation $\mathcal{A}_{\mathfrak{P}}$ and the Slope/Curv/Osc tolerances (Chapter 8). The boundary status is determined by the quadrants and healing indicators on $\mathfrak{P}_\partial$, hence is well posed. □

Corollary 9.218 (Ledger update with T–L–O)**.** *Each row of the Typed Quadrant Ledger (Definition 9.198) is augmented by the T–L–O stamp (Definition 9.216):*

$$\mathbf{L}^{[S,\chi]} \ \mapsto\ \Big(\mathbf{L}^{[S,\chi]},\ \mathbf{TLO}^{[S,\chi]}\ ;\ \textit{if}\ \mathrm{Loc}^{[S,\chi]} = \text{boundary},\ \mathrm{Edge}^{[S,\chi]} \Big).$$

This augmentation is performed after *the quadrant assignment of §9.16.3, on the* same *certified plateau (bulk $\mathfrak{P}$ or boundary $\mathfrak{P}_\partial$; Definition 9.185), thereby maintaining BRST/Nielsen safety (Proposition 9.191) and typed consistency (cf. Proposition 9.207).*

Proof. Existence and stability. By Corollary 9.194, each row $\mathbf{L}^{[S,\chi]}$ exists once the typed witness and quadrant are fixed on a certified plateau. After the quadrant decision (§9.16.3, Proposition 9.212), the T–L–O stamp $\mathbf{TLO}^{[S,\chi]}$ exists and is stable by Definitions 9.190 and 9.216 and the well-posedness result Proposition 9.217; in the boundary case the auxiliary edge status $\mathrm{Edge}^{[S,\chi]}$ is defined by Definition 9.215 using the bulk–boundary overlay Definition 9.214.

Gauge/BRST safety. Because the augmentation uses the *same* diagnostic realized reference configuration as the meters and quadrant (no change of tags or projectors), Nielsen flatness and BRST invariance from Proposition 9.191 carry over. Channel/type order-independence (Proposition 9.207) ensures that inserting the T–L–O metadata does not alter previously certified meters or the quadrant.

Uniqueness. Uniqueness of $\mathrm{Loc}^{[S,\chi]}$ and $\mathrm{On}^{[S,\chi]}$ follows from the certification tolerances along the control axis (Chapter 8) encoded in Definition 9.190 and guaranteed in Proposition 9.217. Since $\mathbf{L}^{[S,\chi]}$ was already unique (Corollary 9.194), appending a unique $\mathbf{TLO}^{[S,\chi]}$ (and, when applicable, $\mathrm{Edge}^{[S,\chi]}$) yields a unique augmented row.

Therefore, performing the update after the quadrant assignment on the same diagnostic realized reference configuration produces a BRST/Nielsen-safe and uniquely defined ledger augmentation, as claimed. □

Operational note. In the *bulk* we use the standard typed meters $\mathbf{V}^{[S,\chi]}_{\text{bulk}} := \mathbf{V}^{[S,\chi]}(\ell;\mathfrak{P})$ of Definition 9.214; at *edges* we replace them by the boundary overlay $\mathbf{V}^{[S,\chi]}_{\partial} := \mathbf{V}^{[S,\chi]}(\ell;\mathfrak{P}_\partial)$ and classify the edge via $\text{Edge}^{[S,\chi]} \in \{\text{heal}, \text{persist}, \text{sel}\}$ (healing vs. persistence vs. spontaneous selection) as in Definition 9.215, before stamping $\mathbf{TLO}^{[S,\chi]}$ (Definition 9.216) and updating the ledger (Corollary 9.218). Practically:

1. **Fix the diagnostic realized reference configuration and tags.** Choose bulk $\mathfrak{P}$ or boundary $\mathfrak{P}_\partial$ on the diagnostic realized reference configuration (Definition 9.185), with the same band tags $P_x(\lambda_{\text{UV}}), K^{(\ell)}_{\text{IR}}$ used throughout the typed workflow (§9.16.2).

2. **Insert identical typed/channel filters.** For the chosen $(S,[\chi])$, insert $\Pi^{[\chi]}_S$ (Definition 9.186) and the triadic Π_α into the readouts to form the typed witness $\mathbf{W}^{[S]}_{[\chi]}$ (Definition 9.187); channel/type order is immaterial (Proposition 9.207).

3. **Evaluate meters and healing.** Compute the diagnostic realized reference configuration SynTh triplet $\mathbf{V}^{[S,\chi]}(\ell;\mathfrak{P})$ from the local two–mode construction (Definitions 9.204 and 9.205), and the typed healing indicators $\mathscr{H}^{[S,\chi]}_\bullet$ (Definition 9.188); bounds/baselines follow Lemma 9.206.

4. **Boundary overlay (if applicable).** For edges, push bulk data through the comparison map $\mathcal{O}^{[S,\chi]}_{\partial\leftarrow\text{bulk}}$ (Definition 9.214) to obtain $\mathbf{V}^{[S,\chi]}_\partial$ and $\mathscr{H}^{[S,\chi]}_{\bullet,\partial}$.

5. **Decide the quadrant per type.** Apply the canonical predicate on the same diagnostic realized reference configuration (Definition 9.189, pipeline in §9.16.3) using $\mathbf{V}$ and $\mathscr{H}_\bullet$ (Definition 9.208 and Corollary 9.209); well-posed/Nielsen-flat by Proposition 9.191.

6. **Classify the edge (boundary only).** Read $\text{Edge}^{[S,\chi]}$ from Definition 9.215: heal vs. persist vs. sel (spontaneous edge selection, i.e. bulk QI but boundary QII).

7. **Stamp T–L–O and record.** Set $\text{Loc}^{[S,\chi]}$ and $\text{On}^{[S,\chi]}$ via Definition 9.190, form $\mathbf{TLO}^{[S,\chi]}$ (Definition 9.216), and append to the ledger row $\mathbf{L}^{[S,\chi]}$ (Definition 9.198 and Corollary 9.218).

Throughout, use the *same* diagnostic realized reference configuration and tags to preserve BRST/Nielsen safety (Proposition 9.191) and typed decoupling within tolerances (Lemma 9.192); any bulk/edge differences are attributed solely to the diagnostic realized reference configuration choice and IR dilation (Chapter 9), not to a change of types or channels.

9.17 The Gaiamiti Field Potential and the Origin of the Apeiron

Typed Parasymmetric TLO Prelude (Gaiamiti)

Before drawing any SSB conclusions from the Gaiamiti potential (cf. §9.17, §9.17), we first perform the typed parasymmetric analysis on the diagnostic realized reference configuration

(Definition 9.185) and *fix* the per–type quadrant and T–L–O data for the $(\Phi\Phi)$ sector. This ensures BRST/Nielsen safety and operational determinacy prior to any particle naming (Corollary 9.199).

Diagnostic realized reference configuration and types. Work on a certified bulk (or boundary) plateau $\mathfrak{P}$ ($\mathfrak{P}_\partial$) with tags $P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}^{(\ell)}$ (Definition 9.185). Choose the Gaiamiti typed projectors from the menu $[\chi] \in \{\mathrm{Tr}, \mathrm{TT}\}$ and optional helicities $\lambda = \pm 2$ when available (Definitions 9.195 and 9.200); these supply the typed filter $\Pi_{\Phi\Phi}^{[\chi]}$ (Definition 9.186).

Witness and meters. Insert $\Pi_{\Phi\Phi}^{[\chi]}$ and the triadic channel projectors $\{\Pi_\alpha\}$ into the readouts on H^0 to form the typed witness $\mathbf{W}_{[\chi]}^{[\Phi\Phi]}$ (Definition 9.187). Compute the local two–mode reductions and SynTh meters (Definitions 9.204 and 9.205); the bounds/baselines and phase properties are given in Lemma 9.206. Gauge/Nielsen safety and channel/type order independence are ensured by Propositions 9.191 and 9.207.

Quadrant decision (typed). Evaluate the typed healing indicators $\left(\mathscr{H}_L^{[\Phi\Phi,\chi]}, \mathscr{H}_S^{[\Phi\Phi,\chi]}\right)$ (Definition 9.188) and apply the canonical predicate to obtain $\mathrm{Quad}^{[\Phi\Phi,\chi]} \in \{\mathrm{QI}, \mathrm{QII}, \mathrm{QIII}, \mathrm{QIV}\}$ (Definition 9.189, pipeline §9.16.3). Typed decoupling and stability on the plateau follow from Lemma 9.192, Definition 9.210, and Proposition 9.212.

T–L–O stamp and ledger. Stamp the location and onset $\left(\mathrm{Loc}^{[\Phi\Phi,\chi]}, \mathrm{On}^{[\Phi\Phi,\chi]}\right)$ (Definition 9.190); for boundary diagnostic realized reference configuration classify the edge via $\mathrm{Edge}^{[\Phi\Phi,\chi]} \in \{\mathrm{heal}, \mathrm{persist}, \mathrm{sel}\}$ (Definitions 9.214 and 9.215). Record the row $\mathbf{L}^{[\Phi\Phi,\chi]}$ in the Typed Quadrant Ledger (Definition 9.198 and Corollaries 9.194 and 9.218). By Corollaries 9.199 and 9.213, this fixes the *operational* spin/polarization labels for the typed mode $[\chi]$ *before* any SSB naming.

Proposition 9.219 (SSB naming gated by typed T–L–O)**.** *Let* $[\chi] \in \{\mathrm{Tr}, \mathrm{TT}\}$ *(with optional* $\lambda = \pm 2$*) be fixed. If and only if the ledger row* $\mathbf{L}^{[\Phi\Phi,\chi]}$ *exists with a certified quadrant* $\mathrm{Quad}^{[\Phi\Phi,\chi]}$ *(Definition 9.198), may SSB nomenclature be applied in §9.17; in particular, a* $(\Phi\Phi, \mathrm{TT})$ *entry with* QII *may be* called *"graviton" (and* QIV *implies pNG semantics) in accordance with Proposition 9.197 and Corollary 9.199.*

Proof. Existence, gauge independence, and uniqueness of the typed meters and quadrant follow from Propositions 9.191, 9.207 and 9.212 together with typed orthogonality/decoupling (Proposition 9.202 and Lemma 9.192). Location/onset (and edge status, when applicable) are well posed by Definitions 9.190 and 9.216 and Proposition 9.217. Hence the ledger row $\mathbf{L}^{[\Phi\Phi,\chi]}$ is uniquely determined (Corollaries 9.194 and 9.218). By Proposition 9.197 and Corollary 9.199, NG/pNG semantics and any SSB naming are to be applied *after* these typed determinations. □

Potential–driven typed quadrant evaluation (Gaiamiti)

We connect the explicit Gaiamiti potential of §9.17 to the typed SynTh meters on the diagnostic realized reference configuration (Definition 9.185) so that the canonical predicate

$\mathrm{Quad}^{[\Phi\Phi,\chi]}$ (Definition 9.189) can be computed directly from the potential's typed curvature data, *before* any SSB naming (cf. Corollary 9.199).

Definition 9.220 (Typed curvature and order parameter from the potential)**.** Let $\mathcal{U}_{\Phi\Phi}$ denote the Gaiamiti potential (or the zero–momentum limit of the effective action) specified in §9.17, evaluated on the certified plateau $\mathfrak{P}$ (or $\mathfrak{P}_\partial$). For a typed projector $\Pi^{[\chi]}_{\Phi\Phi}$ (Definitions 9.186 and 9.200) define:

(i) Typed order parameter.
At a candidate background $\bar{\Phi}$, the typed order parameter is

$$\varphi^{[\chi]} := \Pi^{[\chi]}_{\Phi\Phi}\,\bar{\Phi},$$

where $\bar{\Phi}$ is determined by the diagnostic realized reference configuration variational problem (Chapter 8) and the tags specified in Definition 9.185.

(ii) Typed curvature operator.
The typed curvature operator (Hessian restricted to the type) is

$$\mathcal{K}^{[\chi]}_{\Phi\Phi} \;:=\; \Pi^{[\chi]}_{\Phi\Phi}\,\frac{\delta^2\mathcal{U}_{\Phi\Phi}}{\delta\Phi\,\delta\Phi}\Big|_{\bar{\Phi}}\,\Pi^{[\chi]}_{\Phi\Phi}\,,$$

viewed as a positive (semi)definite operator on the typed subspace, with principal symbol inside the $\bar{\Sigma}$–cones (cone preservation in Definition 9.186).

Definition 9.221 (Potential–based SynTh operators and sources)**.** On the same diagnostic realized reference configuration and type $[\chi]$, choose the SynTh channel operators of Definitions 9.204 and 9.205 as:

$$\mathcal{O}^{(L)}(\ell) \;\equiv\; f_L\Big(\mathcal{K}^{[\chi]}_{\Phi\Phi}(\ell)\Big), \qquad \mathcal{O}^{(S)}(\ell) \;\equiv\; f_S\Big(\varphi^{[\chi]}(\ell)\Big), \qquad \mathcal{O}^{(LS)}(\ell) \;\equiv\; f_{LS}\Big(\mathcal{K}^{[\chi]}_{\Phi\Phi}(\ell),\varphi^{[\chi]}(\ell)\Big),$$

with $f_\bullet$ the diagnostic realized reference configuration–normalized maps used by the SynTh construction (cf. §9.16.2); insert them as $\Pi_\alpha\,\Pi^{[\chi]}_{\Phi\Phi}(\cdot)\Pi^{[\chi]}_{\Phi\Phi}\,\Pi_\alpha$ per Definition 9.187. Define the typed *law* and *state* sources $\delta^{[\Phi\Phi,\chi]}_{L,S}$ and healing indicators $\mathscr{H}^{[\Phi\Phi,\chi]}_{L,S}$ exactly as in Definition 9.188, now computed from these potential–based operators.

Remark 9.222 (On the notation $\mathcal{O}$ in Definition 9.221)**.** Throughout this chapter, $\mathcal{O}$ denotes a *SynTh channel operator*—an operator–valued, scale–tagged map acting on the physical Hilbert space $\mathcal{H}_{\mathrm{phys}}$ after insertion into the typed/triadic subspace. For a fixed type $[\chi]$ and window ℓ, the three instances $\mathcal{O}^{(L)}(\ell)$, $\mathcal{O}^{(S)}(\ell)$, $\mathcal{O}^{(LS)}(\ell)$ are defined by applying the diagnostic realized reference configuration–normalized functional calculi f_L, f_S, f_{LS} (cf. Section 9.16.2) to the typed *law–side* input $\mathcal{K}^{[\chi]}_{\Phi\Phi}(\ell)$ and the typed *state–side* input $\varphi^{[\chi]}(\ell)$, and then sandwiching with the typed and triadic projectors as prescribed in Definition 9.187:

$$\mathcal{O}^{(\bullet)}(\ell) \;\equiv\; \Pi_\alpha\,\Pi^{[\chi]}_{\Phi\Phi}\,f_\bullet\Big(\;\cdot\;\Big)\,\Pi^{[\chi]}_{\Phi\Phi}\,\Pi_\alpha \;:\; \mathcal{H}_{\mathrm{phys}}\to\mathcal{H}_{\mathrm{phys}}, \qquad \alpha\in\{\mathrm{corr},\mathrm{vac},\mathrm{spec}\}.$$

Here, the superscripts indicate the *channel content*: (L) uses only the law–side curvature $\mathcal{K}^{[\chi]}_{\Phi\Phi}(\ell)$; (S) uses only the state probe $\varphi^{[\chi]}(\ell)$; and (LS) is the mixed, jointly typed construction

$f_{LS}\big(\mathcal{K}^{[\chi]}_{\Phi\Phi}(\ell), \varphi^{[\chi]}(\ell)\big)$. When the type $[\chi]$ and triadic channel α are fixed by context, we write simply $\mathcal{O}^{(\bullet)}(\ell)$ or $\mathcal{O}^{(\bullet)}$. This is *operator* notation (not big-O asymptotics).

If the inputs are self–adjoint and $f_\bullet$ are real Borel maps, then $\mathcal{O}^{(\bullet)}(\ell)$ is self–adjoint; diagnostic realized reference configuration normalization ensures $0 \le \mathcal{O}^{(\bullet)}(\ell) \le \mathbb{1}$ in spectral order, so the typed sources $\delta^{[\Phi\Phi,\chi]}_{L,S}$ and healing indicators $\mathscr{H}^{[\Phi\Phi,\chi]}_{L,S}$ of Definition 9.188 are well-posed and scale–stable on certified windows. Moreover, because the typed projectors commute with the triadic channels and are BRST–closed (cf. Definition 9.186 and Proposition 9.207), all resulting $\mathcal{O}^{(\bullet)}(\ell)$ are Nielsen–flat and BRST–safe on $\mathcal{H}_{\text{phys}}$.

Proposition 9.223 (Equivalence of potential–driven and correlational drivers (typed))**.** *For Gaiamiti types $[\chi] \in \{\mathrm{Tr}, \mathrm{TT}\}$ (and helicity refinements when applicable), the typed SynTh meters $\mathbf{V}^{[\Phi\Phi,\chi]}(\ell;\mathfrak{P})$ built from the potential–based operators of Definition 9.221 are, on a certified plateau, equivalent (within Chapter 8 tolerances) to those obtained from the usual correlational channel (Π_{corr}) used in §9.16.2. Consequently, the canonical predicate $\mathrm{Quad}^{[\Phi\Phi,\chi]}$ (Definition 9.189) and the NG/pNG semantics (Proposition 9.197) agree for both constructions.*

Sketch. On the diagnostic realized reference configuration, the (typed) curvature $\mathcal{K}^{[\chi]}_{\Phi\Phi}$ is the zero–momentum limit of the 1PI two–point kernel $\Gamma^{(2)}$ restricted by $\Pi^{[\chi]}_{\Phi\Phi}$. By standard effective action relations, $\Gamma^{(2)}$ is the inverse (in the appropriate sense) of the retarded correlator kernel used in the correlational channel; zeros of $\Gamma^{(2)}$ correspond to poles of the correlator, and small explicit masses map to pNG GMOR slopes. Typed insertion preserves this correspondence by triadic compatibility and cone preservation (Definition 9.186 and Proposition 9.202), while diagnostic realized reference configuration aggregation ensures numerical agreement within tolerances (Definition 9.210). Gauge/Nielsen safety follows from Propositions 9.191 and 9.207. Thus the resulting meters and healing indicators (and hence the quadrant) coincide up to $\mathcal{O}(\varepsilon_{\mathfrak{P}})$. □

Corollary 9.224 (Potential–first recipe for the typed quadrant (Gaiamiti))**.** *To assign $\mathrm{Quad}^{[\Phi\Phi,\chi]}$ using the explicit potential of §9.17:*

(i) Fix the diagnostic realized reference configuration and type (Definitions 9.185 and 9.200); compute $\varphi^{[\chi]}$ and $\mathcal{K}^{[\chi]}_{\Phi\Phi}$ (Definition 9.220).

(ii) Build the potential–based SynTh operators and the typed witness (Definitions 9.187 and 9.221); obtain the diagnostic realized reference configuration meters $\mathbf{V}^{[\Phi\Phi,\chi]}$ (Definition 9.205) and healing indicators (Definition 9.188).

(iii) Apply the canonical predicate (Definition 9.189); NG/pNG semantics then follow from Proposition 9.197.

(iv) Stamp T–L–O on the same diagnostic realized reference configuration and update the ledger (Definitions 9.190, 9.198 and 9.216 and Corollary 9.218).

By Proposition 9.223, this potential–first route yields the same typed quadrant as the correlational route in §9.16.2.

Remark 9.225 (Gaiamiti: salient quadrants and their semantics)**.** For the Gaiamiti sector $S = \Phi\Phi$ (types from Definitions 9.195 and 9.200), evaluated on the diagnostic realized reference

configuration (Definition 9.185) and assigned via the canonical predicate (Definition 9.189), the operationally relevant quadrants are:

- **TT (spin–2):** $\mathrm{Quad}^{[\Phi\Phi,\mathrm{TT}]} = \mathrm{QII}$ in the canonical (law–symmetric, state–selecting) case. This yields an NG pole in the correlational typed channel $\Pi_{\mathrm{corr}}\Pi_{\Phi\Phi}^{[\mathrm{TT}]}$ (Proposition 9.197). The decision may be computed equivalently from the potential via the typed curvature/order parameter (Definitions 9.220 and 9.221 and Proposition 9.223). After stamping T–L–O (Definition 9.190) and recording the ledger row (Definition 9.198 and Corollary 9.194), the TT mode may be *named* "graviton" (Corollary 9.199).
- **TT with explicit breaking:** $\mathrm{Quad}^{[\Phi\Phi,\mathrm{TT}]} = \mathrm{QIV}$ iff the typed law source persists ($\mathscr{H}_L^{[\Phi\Phi,\mathrm{TT}]} > 0$; Definition 9.188); then the typed channel shows a pNG band with GMOR slope on $\mathfrak{P}$ (Proposition 9.197).
- **Tr (spin–0):** generically $\mathrm{Quad}^{[\Phi\Phi,\mathrm{Tr}]} = \mathrm{QI}$ with a gapped radial (Apeiron) mass read from the typed curvature; no NG/pNG in this typed channel (cf. Proposition 9.197).
- **QIII (explicit, healing):** may occur at finite scale if a typed law source is present but heals in the IR ($\mathscr{H}_L^{[\Phi\Phi,\chi]} = 0$); it is recorded via T–L–O (Definition 9.190) but is not the asymptotic (IR) regime.

Boundary diagnostic realized reference configuration $\mathfrak{P}_\partial$ use the same predicate; differences show up only in the *Loc/On* fields of the ledger (bulk vs. boundary onset), not in the spin/type semantics (Definitions 9.190 and 9.198).

Standing assumption (SSB–only regime for Gaiamiti). Throughout this subsection we assume a law–symmetric diagnostic realized reference configuration (no explicit symmetry–breaking spurions in the typed TT channel) and certified plateau control as in Definition 9.185. Typed meters and the quadrant are fixed *first* via Definitions 9.187 to 9.189 and 9.205, and the ledger row is populated per Definition 9.198 and Corollary 9.194. In particular, we work with the potential–driven drivers $\varphi^{[\chi]}, \mathcal{K}_{\Phi\Phi}^{[\chi]}$ (Definitions 9.220 and 9.221) or, equivalently, the correlational route (Proposition 9.223).

Proposition 9.226 (Consistency of the SSB–only treatment)**.** *Under Section 9.17, the Gaiamiti SSB subsection is consistent iff the typed quadrants satisfy*

$$\mathrm{Quad}^{[\Phi\Phi,\mathrm{TT}]} = \mathrm{QII}, \qquad \mathrm{Quad}^{[\Phi\Phi,\mathrm{Tr}]} = \mathrm{QI},$$

(with optional helicity refinement for TT when available), and the typed healing indicators obey $\mathscr{H}_L^{[\Phi\Phi,\mathrm{TT}]} = \mathscr{H}_S^{[\Phi\Phi,\mathrm{TT}]} = 0$, $\mathscr{H}_L^{[\Phi\Phi,\mathrm{Tr}]} = \mathscr{H}_S^{[\Phi\Phi,\mathrm{Tr}]} = 0$. *In this case the TT channel exhibits an NG pole and the Tr channel remains gapped; boundary diagnostic realized reference configuration alter only* Loc/On *in the ledger, not the spin/type semantics.*

Proof. By the canonical typed predicate Definition 9.189, SSB corresponds to QII: law–symmetric with state selection and asymptotic healing. Thus $\mathrm{Quad}^{[\Phi\Phi,\mathrm{TT}]} = \mathrm{QII}$ is equivalent to $\mathscr{H}_L^{[\Phi\Phi,\mathrm{TT}]} = \mathscr{H}_S^{[\Phi\Phi,\mathrm{TT}]} = 0$ plus the TT NG residue in $\Pi_{\mathrm{corr}}\Pi_{\Phi\Phi}^{[\mathrm{TT}]}$ (Proposition 9.197). For the trace channel, the SSB–only regime (no explicit breaking, no state selection) gives $\mathrm{Quad}^{[\Phi\Phi,\mathrm{Tr}]} = \mathrm{QI}$ with positive typed curvature $\mathcal{K}_{\Phi\Phi}^{[\mathrm{Tr}]}$ (gapped radial), consistent with Proposition 9.197. Potential– and correlator–driven meters are equivalent on a certified plateau by

Proposition 9.223, and gauge/Nielsen safety follows from Proposition 9.191 with channel/type compatibility Proposition 9.207. Boundary overlays (if used) affect only the T–L–O stamp (Definition 9.190 and Corollary 9.218) and do not change the typed quadrant provided the same diagnostic realized reference configuration discipline is maintained. Hence the subsection's SSB claims are consistent exactly under the stated quadrant conditions. □

Remark 9.227 (What is excluded by the SSB–only restriction)**.** Cases with persistent typed law sources (QIV) or finite–scale sources that heal (QIII) in the TT channel are *not* part of the SSB–only narrative and should be deferred to the explicit–breaking analysis. Detecting these is automatic in the workflow via $\mathscr{H}_L^{[S,\chi]}$ and the predicate of Definition 9.189.

While the kinetic term in the action governs the propagation of the Gaiamiti field, the *potential term*, $\mathcal{V}_\Phi(\hat{\mathbf{\Phi}})$, dictates its self-interaction and determines the energetically preferred ground state structure of the emergent geometry. It is the source of the vacuum's intrinsic stability and the mechanism by which the fundamental quantum of geometry—the Apeiron—acquires its properties and gives rise to gravitational excitations.

Invariant Construction of the Gaiamiti Potential

It is both necessary and illuminating to demonstrate how the specific form of the Gaiamiti potential—as given in eq. (9.95)—arises naturally from the representation theory, invariance principles, and the functional role of the Gaiamiti field $\hat{\mathbf{\Phi}}(x)$ within the theory. We shall now construct this expression from these first principles.

1. The Fundamental Invariant of the Gaiamiti Field

The Gaiamiti field is an operator–valued vector field

$$\hat{\mathbf{\Phi}}(x) \;\in\; T_{|\Omega(x)\rangle}\mathcal{M}_{\text{vac}} \quad \text{for each locus } x,$$

i.e., a section of the vacuum–tangent bundle determined by the Gaiamiti sector (Definition 6.118). The fibrewise inner product on the vacuum–tangent space,

$$g_T(|V\rangle, |W\rangle) \;:=\; \langle V|W\rangle_{\mathcal{H}_{\text{em}}},$$

is induced from the ambient emergent Hilbert space and is positive–definite on the tangent fibre (Definition 6.104). Any admissible potential energy density $\mathcal{V}_\Phi(x)$ must be a Hermitian scalar operator constructed from $\hat{\mathbf{\Phi}}(x)$ and invariant under the symmetries of the theory (base diffeomorphisms compatible with the Relational Levi–Civita structure and unitary gauge on the operator bundle; cf. Definition 6.7, Section 6.5.3, and eq. (6.257)).

Definition 9.228 (Gaiamiti tangent norm and fundamental invariant)**.** The *fundamental invariant* of the Gaiamiti field is the fibrewise scalar operator

$$\mathcal{I}_\Phi(x) \;:=\; g_T\Big(\hat{\mathbf{\Phi}}(x),\, \hat{\mathbf{\Phi}}(x)\Big)\,. \tag{9.91}$$

This is the canonical (norm–squared) quadratic invariant available because $\hat{\mathbf{\Phi}}(x)$ takes values in a tangent space. It is the precise analogue of $|\phi|^2$ for a complex scalar field, with g_T replacing the flat inner product.

Lemma 9.229 (Hermiticity, positivity, and scalar character)**.** *For each x, the operator $\mathcal{I}_\Phi(x)$ is Hermitian and positive, with $\mathcal{I}_\Phi(x) \geq 0$ and $\mathcal{I}_\Phi(x) = 0$ iff $\hat{\mathbf{\Phi}}(x)$ is the zero section of the vacuum–tangent bundle. Moreover, $\mathcal{I}_\Phi(x)$ transforms as a scalar under base diffeomorphisms compatible with the Relational Levi–Civita connection and is invariant under unitary frame changes in the operator bundle.*

Proof sketch. Hermiticity and positivity follow from $g_T(\cdot,\cdot)$ being the restriction of the ambient inner product (Definition 6.104). Unitary changes of fibre frame act by $|V\rangle \mapsto U(x)|V\rangle$ with $U(x)$ unitary, hence $g_T(U\hat{\mathbf{\Phi}}, U\hat{\mathbf{\Phi}}) = g_T(\hat{\mathbf{\Phi}}, \hat{\mathbf{\Phi}})$. Compatibility of the Relational Levi–Civita connection with the unified chronogeometric tensor implies scalar behaviour under base diffeomorphisms (Definition 6.7). □

Definition 9.230 (Admissible Gaiamiti potential density)**.** An admissible Gaiamiti potential energy density is any Hermitian operator of the form

$$\mathcal{V}_\Phi(x) \;=\; F\Big(\mathcal{I}_\Phi(x)\Big)\,, \qquad \mathcal{I}_\Phi(x) = g_T\Big(\hat{\mathbf{\Phi}}(x), \hat{\mathbf{\Phi}}(x)\Big)\,, \tag{9.92}$$

for some real functional F chosen to ensure spectral/variational well–posedness on the operative band. In particular, a polynomial ansatz

$$\mathcal{V}_\Phi(x) \;=\; \frac{m_\Phi^2}{2}\,\mathcal{I}_\Phi(x) \;+\; \frac{\lambda_\Phi}{4}\,\mathcal{I}_\Phi(x)^2 \;+\; \sum_{k\geq 3} \frac{c_k}{k!}\,\mathcal{I}_\Phi(x)^{\,k} \tag{9.93}$$

(with real couplings $m_\Phi^2, \lambda_\Phi, c_k$) is admissible whenever F renders the Hamiltonian bounded below and compatible with the band–limited calculus used in Chapter VI (Section 6.5.3 and eq. (6.259)).

Remark 9.231 (From density to scalar action via the invariant volume)**.** The potential contributes to the (wordwise or standard) action once integrated against the invariant volume element. In the HS formulation,

$$d\boldsymbol{\mu}(x) = d^d x\, \boldsymbol{\varpi}(x), \qquad \boldsymbol{\varpi}(x) = \Big(\det{}_{\mathrm{HS}}\Big(-\boldsymbol{g}^{\mathrm{Lor}}(x)\Big)\Big)^{1/2} \quad (\textit{Definition 6.34}),$$

with expectation reduction $\langle d\boldsymbol{\mu}\rangle = \sqrt{-\det \bar{g}^{\mathrm{Lor}}}\, d^d x$ (Remark 6.35). The pure–potential contribution to the action is therefore

$$S_\Phi^{(\mathrm{pot})} \;=\; \int_{\mathcal{X}_q} d\boldsymbol{\mu}(x)\; \mathcal{V}_\Phi(x) \tag{9.94}$$

(and analogously in the standard expectation–level presentation).

Proposition 9.232 (Symmetry invariance of $\mathcal{V}_\Phi$)**.** *If $\mathcal{V}_\Phi(x) = F(\mathcal{I}_\Phi(x))$ as in Definition 9.230, then $\mathcal{V}_\Phi$ is:*

(i) a Hermitian operator at each x;

(ii) a scalar under base diffeomorphisms compatible with the Relational Levi–Civita connection (Definition 6.7);

(iii) invariant under unitary gauge transformations of the operator bundle (the bundle connection enters only through kinetic/covariant terms; cf. Section 6.5.3 and eqs. (6.257) and (6.259)).

Consequently, the action contribution (9.94) is fully gauge– and diffeo–invariant when paired with the invariant volume (Definition 6.34).

Proof sketch. Items (1)–(2) are immediate from Lemma 9.229. For (3), unitary bundle transformations act as fibre isometries on g_T, hence preserve $\mathcal{I}_\Phi$ and thus $F(\mathcal{I}_\Phi)$. The volume element is constructed to be invariant (HS) and reduces to the standard invariant measure under expectation (Definition 6.34 and Remark 6.35). □

Remark 9.233 (Holosymmetric wordwise counterpart)**.** All statements admit a holosymmetric–wordwise rewriting by replacing g_T with the HS inner product g_T^{HS} on the vacuum–tangent fibre:

$$\mathcal{I}_\Phi^{\mathrm{HS}}(x) \;:=\; g_T^{\mathrm{HS}}\Big(\hat{\mathbf{\Phi}}(x), \hat{\mathbf{\Phi}}(x)\Big)\,,$$

where g_T^{HS} is defined via the HS projector and coincides with the standard g_T at expectation level (Definition 6.106 and eq. (6.167)). Then $\mathcal{V}_\Phi(x) = F(\mathcal{I}_\Phi^{\mathrm{HS}}(x))$ and the action uses $d\boldsymbol{\mu}$ as above (Definition 6.34).

Corollary 9.234 (Stationarity and mass term)**.** *For a polynomial F with $F'(0) = \frac{m_\Phi^2}{2}$, the quadratic variation of $S_\Phi^{(\mathrm{pot})}$ around a vacuum configuration determines the Gaiamiti mass parameter m_Φ^2 in the potential sector, with higher coefficients λ_Φ, c_k controlling self–interaction strength; compatibility with the dynamical (kinetic/covariant) sector is governed by the Relational Levi–Civita principle (Definition 6.7) and the total derivative structure (Section 6.5.3 and eq. (6.259)).*

Summary. Because $\hat{\mathbf{\Phi}}(x)$ is a vacuum–tangent vector, the scalar operator

$$g_T\Big(\hat{\mathbf{\Phi}}(x), \hat{\mathbf{\Phi}}(x)\Big)$$

is the unique, canonical quadratic invariant (Definition 6.104). *All* admissible Gaiamiti potentials are functions of this invariant, $\mathcal{V}_\Phi = F(\mathcal{I}_\Phi)$, ensuring Hermiticity and invariance under the theory's symmetries, with integration performed against the invariant volume (Definition 6.34 and Remark 6.35).

2. Constraints of Stability and Symmetry

Any physically viable potential must satisfy two crucial constraints:

Stability: The potential must be bounded from below to ensure the existence of a stable vacuum state. This requires that for large field values, the potential approaches $+\infty$. This condition immediately disfavors odd powers of the field and requires that the highest-order term in any polynomial expansion have a positive coefficient.

Symmetry for Spontaneous Breaking: For symmetry to be broken *spontaneously* rather than explicitly, the Lagrangian itself must possess the symmetry. The simplest relevant symmetry is a $\mathbb{Z}_2$ invariance under which the field transforms as $\hat{\mathbf{\Phi}}(x) \mapsto -\hat{\mathbf{\Phi}}(x)$. The invariant $g_T(\hat{\mathbf{\Phi}}, \hat{\mathbf{\Phi}})$ respects this symmetry, as it is quadratic in the field. To preserve this symmetry, the potential must be an even function of $\hat{\mathbf{\Phi}}$, which means it must be a polynomial in powers of the invariant $g_T(\hat{\mathbf{\Phi}}, \hat{\mathbf{\Phi}})$.

These two principles together demand that the potential must be a polynomial of the form $\mathcal{V}_\Phi = \sum_n c_n [g_T(\hat{\mathbf{\Phi}}, \hat{\mathbf{\Phi}})]^n$ with a positive leading coefficient c_n.

3. The Principle of Minimality and Universality

Among all possible polynomials, physical theories favor the simplest form capable of describing the required phenomena. This is guided by two related principles:

Renormalizability: In standard quantum field theory, only potentials with terms up to the fourth power of the field (quartic interactions) are renormalizable in four spacetime dimensions [82, 14]. Higher-order terms are suppressed at low energies. While the IQR framework may possess its own regularization scheme, the principle of selecting the lowest-order relevant terms remains a powerful guide.

Universality: The theory of critical phenomena shows that near a phase transition, a vast number of disparate physical systems are described by the same universal Landau-Ginzburg potential, which is a simple polynomial in the order parameter [101]. The minimal form that allows for a second-order phase transition (i.e., spontaneous symmetry breaking) is precisely the quadratic-plus-quartic form.

These principles compel us to select the lowest-order even polynomial in $\hat{\mathbf{\Phi}}$ that can support a non-trivial vacuum structure.

The Unique Minimal Potential

The confluence of these principles—invariance, stability, symmetry, and minimality—uniquely determines the form of the Gaiamiti potential. The simplest non-trivial, Hermitian, and symmetric potential constructed from the invariant $g_T(\hat{\mathbf{\Phi}}, \hat{\mathbf{\Phi}})$ is the operator-valued analogue of the classic ϕ^4 Higgs potential:

$$\mathcal{V}_\Phi(\hat{\mathbf{\Phi}}) = \frac{1}{2} m_A^2 \, g_T(\hat{\mathbf{\Phi}}, \hat{\mathbf{\Phi}}) + \frac{\lambda_A}{4!} \left[g_T(\hat{\mathbf{\Phi}}, \hat{\mathbf{\Phi}}) \right]^2$$

This expression therefore arises as the unique and necessary consequence of the foundational requirements for a consistent and stable theory of emergent geometry that supports spontaneous symmetry breaking.

The Gaiamiti Field Potential

Definition 9.235 (Gaiamiti Field Potential)**.** Let $\hat{\mathbf{\Phi}}(x) \in \mathrm{End}(\mathcal{H}_{\mathrm{em}}) \otimes T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}$ be the operator-valued Gaiamiti field. The *potential energy operator density*, $\mathcal{V}_\Phi(\hat{\mathbf{\Phi}}(x))$, is a Hermitian

operator-valued function constructed from invariants in the tangent space:

$$\boxed{\mathcal{V}_\Phi(\hat{\mathbf{\Phi}}(x)) := \frac{1}{2}m_A^2\, g_T\left(\hat{\mathbf{\Phi}}(x), \hat{\mathbf{\Phi}}(x)\right) + \frac{\lambda_A}{4!}\left[g_T\left(\hat{\mathbf{\Phi}}(x), \hat{\mathbf{\Phi}}(x)\right)\right]^2 + \dots,} \tag{9.95}$$

where:

(i) g_T is the inner product on the tangent space $T_{|\Omega(x)\rangle}\mathcal{M}_{\text{vac}}$ (Definition 6.104).

(ii) m_A^2 is a parameter setting the mass scale of the fundamental Apeiron quantum.

(iii) λ_A is a dimensionless self-interaction coupling constant for the Apeiron field.

Physical Interpretation: From Potential to Graviton Emergence The physically relevant potential landscape is the classical, scalar-valued function $V_\Phi(x)$, obtained by taking the expectation value of the operator density in the Quantum Holo-State:

$$V_\Phi(x) := \langle \overset{\circ}{\mathbf{\Psi}} | \mathcal{V}_\Phi(\hat{\mathbf{\Phi}}(x)) | \overset{\circ}{\mathbf{\Psi}} \rangle. \tag{9.96}$$

The structure of this classical potential governs the fundamental properties of the emergent geometry. The minima of V_Φ, where $\frac{\delta V_\Phi}{\delta \hat{\mathbf{\Phi}}} = 0$, define the vacuum expectation value (VEV) of the field, $\bar{\Phi}(x)$, which corresponds to the stable, ground-state configuration of the geometry.

Holosymmetric Wordwise Gaiamiti Field Potential

We now dress the standard Gaiamiti potential of Section 9.17 in the *holosymmetric wordwise* (HS) formalism developed in Chapter VI, using the HS vacuum–tangent inner product (Definition 6.106 and eq. (6.167)) and the HS bundle/gauge structure (Sections 6.4.1 and 6.6.1). Chronogeometric compatibility is with the Relational Levi–Civita structure (Definition 6.7).

Definition 9.236 (HS fundamental invariant for the Gaiamiti field)**.** Let $\hat{\mathbf{\Phi}}(x) \in \mathrm{End}(\mathcal{H}_{\text{em}}) \otimes T_{|\Omega(x)\rangle}\mathcal{M}_{\text{vac}}$ (Definition 6.118). The holosymmetric fundamental invariant is the fibrewise scalar operator

$$\boxed{\mathcal{I}_\Phi^{\text{HS}}(x) \;:=\; g_T^{\text{HS}}\Big(\hat{\mathbf{\Phi}}(x),\, \hat{\mathbf{\Phi}}(x)\Big),} \tag{9.97}$$

where g_T^{HS} is the HS inner product on $T_{|\Omega\rangle}\mathcal{M}_{\text{vac}}$ (Definition 6.106). At expectation level, $\mathcal{I}_\Phi^{\text{HS}}$ reduces to $g_T(\hat{\mathbf{\Phi}}, \hat{\mathbf{\Phi}})$ (Definition 6.104).

Definition 9.237 (HS Gaiamiti potential operator density)**.** The *HS potential energy operator density* is any Hermitian function of $\mathcal{I}_\Phi^{\text{HS}}$:

$$\boxed{\mathcal{V}_\Phi^{\text{HS}}\Big(\hat{\mathbf{\Phi}}(x)\Big) \;:=\; \frac{1}{2}\, m_A^2\, \mathcal{I}_\Phi^{\text{HS}}(x) \;+\; \frac{\lambda_A}{4!}\left[\mathcal{I}_\Phi^{\text{HS}}(x)\right]^2 \;+\; \sum_{k\geq 3}\frac{c_k}{k!}\left[\mathcal{I}_\Phi^{\text{HS}}(x)\right]^k,} \tag{9.98}$$

with real couplings m_A^2, λ_A, c_k. For $m_A^2 \geq 0$ and $\lambda_A > 0$ (and c_k bounded appropriately), $\mathcal{V}_\Phi^{\text{HS}}$ is bounded below on the HS band. The pair (m_A^2, λ_A) coincides with the standard parameters in Definition 9.235.

Proposition 9.238 (HS gauge/frame invariance and diffeo covariance)**.** *Under HS modal frame changes $g(x) \in U(n)$ and vacuum–tangent rotations $R(x) \in O(r)$,*

$$\hat{\mathbf{\Phi}} \mapsto \hat{\mathbf{\Phi}}' = \Big(\mathrm{Ad}_g\, \hat{\mathbf{\Phi}}\Big) \cdot R, \qquad g_T^{\mathrm{HS}}(\mathrm{Ad}_g\, X \cdot R, \mathrm{Ad}_g\, Y \cdot R) = g_T^{\mathrm{HS}}(X, Y),$$

so $\mathcal{I}_\Phi^{\mathrm{HS}}$ and hence $\mathcal{V}_\Phi^{\mathrm{HS}}$ are invariant. Moreover, $\mathcal{V}_\Phi^{\mathrm{HS}}$ is a base scalar for diffeomorphisms compatible with $\nabla^{(\mathrm{rel})}$ (Definition 6.7).

Proof idea. g_T^{HS} is defined by the HS projector and is preserved by HS unitaries on the operator slot and orthogonal rotations on $T\mathcal{M}_{\mathrm{vac}}$ (Definition 6.106); hence $\mathcal{I}_\Phi^{\mathrm{HS}}$ is invariant and any functional of it is invariant. Because g_T^{HS} is a fibrewise pairing and carries no free base indices, $\mathcal{V}_\Phi^{\mathrm{HS}}$ is a base scalar; compatibility of $\nabla^{(\mathrm{rel})}$ with $\bar{\Sigma}$ ensures scalar behaviour under the allowed diffeomorphisms.

Holosymmetric wordwise proof. We show invariance under HS modal frame changes $g(x) \in U(n)$ and vacuum–tangent rotations $R(x) \in O(r)$, and then establish that $\mathcal{V}_\Phi^{\mathrm{HS}}$ is a base scalar for diffeomorphisms compatible with $\nabla^{(\mathrm{rel})}$.

Preliminaries and notation. Fix a g_T–orthonormal local frame $\{|e_i(x)\rangle\}_{i=1}^r$ of $T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}$ with dual $\{\langle e^i(x)|\}$, so that $\hat{\mathbf{\Phi}} = \sum_i \hat{\Phi}^i\, |e_i\rangle$ with operator components $\hat{\Phi}^i \in \mathrm{End}(\mathcal{H}_{\mathrm{em}})$. A vacuum–tangent rotation $R(x) \in O(r)$ acts as $|e_i\rangle \mapsto |e'_i\rangle := R_i{}^j |e_j\rangle$, while an HS modal frame change $g(x) \in U(n)$ acts by the adjoint automorphism $\mathrm{Ad}_g(X) = gXg^\dagger$ on the operator slot. Accordingly,

$$\hat{\Phi}'^i \;=\; \mathrm{Ad}_g(\hat{\Phi}^j)\, R_j{}^i, \qquad \hat{\mathbf{\Phi}}' \;=\; \Big(\mathrm{Ad}_g\, \hat{\mathbf{\Phi}}\Big) \cdot R, \tag{9.99}$$

which is the component form of the transformation stated in the proposition.

Step 1: $O(r)$–invariance of g_T^{HS} on the tangent index. By Definition 6.106, g_T^{HS} is a fibrewise inner product on the vacuum–tangent space, hence in the orthonormal frame $g_T^{\mathrm{HS}}(|e_i\rangle, |e_j\rangle) = \delta_{ij}$. For two HS–tangent fields $\hat{\mathbf{X}} = \hat{X}^i |e_i\rangle$ and $\hat{\mathbf{Y}} = \hat{Y}^i |e_i\rangle$,

$$g_T^{\mathrm{HS}}(\hat{\mathbf{X}}, \hat{\mathbf{Y}}) \;=\; \sum_i g_T^{\mathrm{HS}}\Big(\hat{X}^i |e_i\rangle,\; \hat{Y}^i |e_i\rangle\Big) \;=\; \sum_i \mathsf{B}\Big(\hat{X}^i, \hat{Y}^i\Big),$$

where B denotes the operator–slot pairing induced by the HS construction (precise form immaterial here). Under a rotation $R \in O(r)$,

$$\begin{aligned} \hat{X}'^i &= \hat{X}^j R_j{}^i, & \hat{Y}'^i &= \hat{Y}^k R_k{}^i, \\ \sum_i \mathsf{B}(\hat{X}'^i, \hat{Y}'^i) &= \sum_{i,j,k} R_j{}^i R_k{}^i\, \mathsf{B}(\hat{X}^j, \hat{Y}^k) \\ &= \sum_j \mathsf{B}(\hat{X}^j, \hat{Y}^j). \end{aligned}$$

using orthogonality $R^T R = \mathbf{1}$. Thus g_T^{HS} is $O(r)$–invariant on the tangent index.

Step 2: $U(n)$–invariance of g_T^{HS} on the operator slot. By Definition 6.106, g_T^{HS} is defined via the HS projector and is preserved by HS unitaries on the operator slot; in particular, for all operator entries X, Y,

$$g_T^{\mathrm{HS}}(\mathrm{Ad}_g X, \mathrm{Ad}_g Y) \;=\; g_T^{\mathrm{HS}}(X, Y), \qquad g \in U(n). \tag{9.100}$$

This follows from unitarity of g and the fact that the HS pairing is built from unitary–invariant expressions on the operator fibre (e.g. Hilbert–Schmidt–type or FD–trace–reduced pairings in the HS construction).

Step 3: Invariance of the HS fundamental invariant and of the potential. Applying (9.99) and Steps 1–2 to the HS fundamental invariant $\mathcal{I}_\Phi^{\mathrm{HS}} = g_T^{\mathrm{HS}}(\hat{\mathbf{\Phi}}, \hat{\mathbf{\Phi}})$ (Definition 9.236),

$$\begin{aligned}
\mathcal{I}_\Phi^{\mathrm{HS}}[\hat{\mathbf{\Phi}}'] &= g_T^{\mathrm{HS}}\Big(\mathrm{Ad}_g\, \hat{\mathbf{\Phi}} R, \mathrm{Ad}_g\, \hat{\mathbf{\Phi}} R\Big) \\
&= \sum_i \mathsf{B}\Big(\mathrm{Ad}_g\, \hat{\Phi}^j R_j{}^i, \mathrm{Ad}_g\, \hat{\Phi}^k R_k{}^i\Big) \\
&= \sum_{j,k} (R^{\mathrm{T}} R)_{jk}\, \mathsf{B}\Big(\mathrm{Ad}_g\, \hat{\Phi}^j, \mathrm{Ad}_g\, \hat{\Phi}^k\Big) \\
&= \sum_j \mathsf{B}(\mathrm{Ad}_g\, \hat{\Phi}^j, \mathrm{Ad}_g\, \hat{\Phi}^j) \\
&= \sum_j \mathsf{B}(\hat{\Phi}^j, \hat{\Phi}^j) \\
&= \mathcal{I}_\Phi^{\mathrm{HS}}[\hat{\mathbf{\Phi}}].
\end{aligned}$$

where we used orthogonality $R^T R = \mathbf{1}$ and the isometry (9.100). Therefore $\mathcal{I}_\Phi^{\mathrm{HS}}$ is invariant under $U(n) \times O(r)$, and for any real functional F,

$$\mathcal{V}_\Phi^{\mathrm{HS}}(\hat{\mathbf{\Phi}}) := F\Big(\mathcal{I}_\Phi^{\mathrm{HS}}(\hat{\mathbf{\Phi}})\Big) \quad \text{obeys} \quad \mathcal{V}_\Phi^{\mathrm{HS}}(\hat{\mathbf{\Phi}}') = \mathcal{V}_\Phi^{\mathrm{HS}}(\hat{\mathbf{\Phi}}),$$

proving HS gauge/frame invariance (cf. Definition 9.237).

Step 4: Base–scalar behaviour under diffeomorphisms compatible with $\nabla^{(\mathrm{rel})}$. Let $\varphi : \mathcal{X}_q \to \mathcal{X}_q$ be a diffeomorphism and denote the induced pullback on sections by φ^*. Because g_T^{HS} is a *fibrewise* pairing with no free base indices and the Relational Levi–Civita connection $\nabla^{(\mathrm{rel})}$ is compatible with $\bar{\Sigma}$ (Definition 6.7), the HS inner product is transported naturally:

$$\mathcal{I}_\Phi^{\mathrm{HS}}(\varphi^* \hat{\mathbf{\Phi}}) \;=\; \varphi^*\Big(\mathcal{I}_\Phi^{\mathrm{HS}}(\hat{\mathbf{\Phi}})\Big),$$

i.e. $\mathcal{I}_\Phi^{\mathrm{HS}}$ is a *scalar function* on the base. Hence so is $\mathcal{V}_\Phi^{\mathrm{HS}} = F\Big(\mathcal{I}_\Phi^{\mathrm{HS}}\Big)$, establishing diffeo covariance in the sense stated:

$$\mathcal{V}_\Phi^{\mathrm{HS}}(\varphi^* \hat{\mathbf{\Phi}}) \;=\; \varphi^*\Big(\mathcal{V}_\Phi^{\mathrm{HS}}(\hat{\mathbf{\Phi}})\Big).$$

Conclusion. Combining Steps 1–4, $\mathcal{I}_\Phi^{\mathrm{HS}}$ and therefore $\mathcal{V}_\Phi^{\mathrm{HS}}$ are invariant under HS modal frame changes $g \in U(n)$ and vacuum–tangent rotations $R \in O(r)$, and $\mathcal{V}_\Phi^{\mathrm{HS}}$ is a base scalar for diffeomorphisms compatible with $\nabla^{(\mathrm{rel})}$. This proves Proposition 9.238. □

Remark 9.239 (HS action contribution and reduction to standard)**.** The HS potential contributes to the action through the HS invariant volume $d\boldsymbol{\mu} = d^d x\, \boldsymbol{\varpi}$, $\boldsymbol{\varpi} = (\det_{\mathrm{HS}}(-\boldsymbol{g}^{\mathrm{Lor}}))^{1/2}$ (Definition 6.34):

$$\boxed{\widehat{\boldsymbol{S}}_{\Phi,\mathrm{pot}} := \int_{\mathcal{X}_q} d\boldsymbol{\mu}(x)\ \mathrm{Tr}_{\mathrm{FD}}\Big\{\mathcal{V}_\Phi^{\mathrm{HS}}(\hat{\boldsymbol{\Phi}})\Big\},} \tag{9.101}$$

where $\mathrm{Tr}_{\mathrm{FD}}$ is the field–dialectic trace (Definition 6.45). Applying the expectation/volume push–throughs (Remark 6.35 and eq. (6.114)) yields the standard potential sector with $g_T^{\mathrm{HS}} \to g_T$ and $d\boldsymbol{\mu} \to \sqrt{-\det \bar{g}^{\mathrm{Lor}}}\, d^d x$ (Section 9.17).

Lemma 9.240 (First variation and stationarity)**.** *Let $\delta\hat{\boldsymbol{\Phi}}$ be a compactly supported variation in the HS tangent fibre. Then*

$$\boxed{\delta\,\mathrm{Tr}_{\mathrm{FD}}\Big\{\mathcal{V}_\Phi^{\mathrm{HS}}\Big\} = \mathrm{Tr}_{\mathrm{FD}}\left\{\left(\tfrac{1}{2}\,m_A^2 + \tfrac{\lambda_A}{3!}\,\mathcal{I}_\Phi^{\mathrm{HS}} + \sum_{k\geq 3} \tfrac{c_k}{(k-1)!}\,[\mathcal{I}_\Phi^{\mathrm{HS}}]^{k-1}\right)\delta\Big(g_T^{\mathrm{HS}}(\hat{\boldsymbol{\Phi}},\hat{\boldsymbol{\Phi}})\Big)\right\}.} \tag{9.102}$$

Using bilinearity of g_T^{HS}, $\delta g_T^{\mathrm{HS}}(\hat{\boldsymbol{\Phi}},\hat{\boldsymbol{\Phi}}) = 2\,\mathrm{Re}\, g_T^{\mathrm{HS}}(\hat{\boldsymbol{\Phi}},\delta\hat{\boldsymbol{\Phi}})$, so the stationary condition contributes the fibrewise "mass+self–interaction" term proportional to $\hat{\boldsymbol{\Phi}}$ in the Euler–Lagrange equation.

Holosymmetric wordwise proof. By Definition 9.237, the HS Gaiamiti potential is a real functional of the HS invariant $\mathcal{I}_\Phi^{\mathrm{HS}} = g_T^{\mathrm{HS}}(\hat{\boldsymbol{\Phi}},\hat{\boldsymbol{\Phi}})$ (Definition 9.236 and eq. (9.97)), with polynomial presentation

$$\mathcal{V}_\Phi^{\mathrm{HS}} = \frac{1}{2}m_A^2\,\mathcal{I}_\Phi^{\mathrm{HS}} + \frac{\lambda_A}{4!}\,(\mathcal{I}_\Phi^{\mathrm{HS}})^2 + \sum_{k\geq 3}\frac{c_k}{k!}\,(\mathcal{I}_\Phi^{\mathrm{HS}})^k \quad (\text{cf. } eq.\ (9.98)).$$

Let $F(s) = \frac{1}{2}m_A^2 s + \frac{\lambda_A}{4!}s^2 + \sum_{k\geq 3}\frac{c_k}{k!}s^k$ so that $\mathcal{V}_\Phi^{\mathrm{HS}} = F(\mathcal{I}_\Phi^{\mathrm{HS}})$. Using linearity/cyclicity of the field–dialectic trace (Definition 6.45 and eq. (6.99)) and the chain rule in the strong–operator topology (cf. Remark 6.187), we obtain

$$\delta\,\mathrm{Tr}_{\mathrm{FD}}\{\mathcal{V}_\Phi^{\mathrm{HS}}\} = \mathrm{Tr}_{\mathrm{FD}}\Big\{F'(\mathcal{I}_\Phi^{\mathrm{HS}})\,\delta\mathcal{I}_\Phi^{\mathrm{HS}}\Big\}, \qquad F'(s) = \tfrac{1}{2}m_A^2 + \tfrac{\lambda_A}{3!}\,s + \sum_{k\geq 3}\tfrac{c_k}{(k-1)!}\,s^{k-1}. \tag{9.103}$$

It remains to compute $\delta\mathcal{I}_\Phi^{\mathrm{HS}}$. By definition (Definition 9.236),

$$\mathcal{I}_\Phi^{\mathrm{HS}} = g_T^{\mathrm{HS}}(\hat{\boldsymbol{\Phi}},\hat{\boldsymbol{\Phi}}),$$

and since g_T^{HS} is a Hermitian fibrewise inner product on the HS vacuum–tangent space (Definition 6.106 and eq. (6.167)), its bilinearity yields

$$\delta\mathcal{I}_\Phi^{\mathrm{HS}} = \delta g_T^{\mathrm{HS}}(\hat{\boldsymbol{\Phi}},\hat{\boldsymbol{\Phi}}) = g_T^{\mathrm{HS}}(\delta\hat{\boldsymbol{\Phi}},\hat{\boldsymbol{\Phi}}) + g_T^{\mathrm{HS}}(\hat{\boldsymbol{\Phi}},\delta\hat{\boldsymbol{\Phi}}) = 2\,\mathrm{Re}\, g_T^{\mathrm{HS}}(\hat{\boldsymbol{\Phi}},\delta\hat{\boldsymbol{\Phi}}). \tag{9.104}$$

Substituting (9.104) into (9.103) gives exactly the boxed identity (9.102):

$$\delta\,\mathrm{Tr}_{\mathrm{FD}}\Big\{\mathcal{V}_\Phi^{\mathrm{HS}}\Big\} = \mathrm{Tr}_{\mathrm{FD}}\left\{\left(\tfrac{1}{2}\,m_A^2 + \tfrac{\lambda_A}{3!}\,\mathcal{I}_\Phi^{\mathrm{HS}} + \sum_{k\geq 3}\tfrac{c_k}{(k-1)!}\,[\mathcal{I}_\Phi^{\mathrm{HS}}]^{k-1}\right)\delta\Big(g_T^{\mathrm{HS}}(\hat{\mathbf{\Phi}},\hat{\mathbf{\Phi}})\Big)\right\}.$$

Because $\delta\hat{\mathbf{\Phi}}$ is compactly supported in the HS tangent fibre, no boundary terms arise (the potential has no derivatives). Inserting (9.104) into the variation of the full HS action for the Gaiamiti field (cf. the HS action template and volume element in Section 6.5.2 and Definition 6.34) shows that the potential contributes the fibrewise *mass+ self–interaction* term proportional to $\hat{\mathbf{\Phi}}$ in the Euler–Lagrange equation, as claimed. □

Corollary 9.241 (Boundedness and vacuum types)**.** *If $\lambda_A > 0$ and all higher c_k obey the usual positivity bounds, then $\mathcal{V}_\Phi^{\mathrm{HS}}$ is bounded below. For $m_A^2 > 0$ the symmetric vacuum $\hat{\mathbf{\Phi}} = 0$ is stationary; for $m_A^2 < 0$ a "Mexican-hat" structure arises in the HS tangent norm $\mathcal{I}_\Phi^{\mathrm{HS}}$, with symmetry breaking analyzed via the HS vacuum manifold of Chapter 6 (Section 6.1.9).*

Proof. Let $s(x) := \mathcal{I}_\Phi^{\mathrm{HS}}(x) = g_T^{\mathrm{HS}}(\hat{\mathbf{\Phi}}(x),\hat{\mathbf{\Phi}}(x)) \geq 0$ by positivity of the HS vacuum–tangent inner product (Definition 6.106 and eq. (6.167)). By Definition 9.237, the HS potential is the polynomial functional

$$\mathcal{V}_\Phi^{\mathrm{HS}}(\hat{\mathbf{\Phi}}) \;=\; F(s) \quad \text{with} \quad F(s) = \tfrac{1}{2}m_A^2 s + \tfrac{\lambda_A}{4!}s^2 + \sum_{k\geq 3}\tfrac{c_k}{k!}\,s^k \qquad (s \geq 0). \tag{9.105}$$

(i) Boundedness from below. Assume $\lambda_A > 0$ and that the "usual positivity bounds" on higher couplings hold (e.g. $c_k \geq 0$ for all even $k \geq 4$ and any odd-power contributions are chosen so that $F(s) \to +\infty$ as $s \to \infty$).[6] Then F is coercive on $[0,\infty)$:

$$\lim_{s\to\infty} F(s) = +\infty.$$

Since F is continuous on the closed ray $[0,\infty)$, it attains a global minimum $F(s_\star)$ with $s_\star \geq 0$. Hence $\mathcal{V}_\Phi^{\mathrm{HS}} = F(\mathcal{I}_\Phi^{\mathrm{HS}})$ is bounded below on field configurations, proving the first claim.

(ii) Stationarity and vacuum type for $m_A^2 > 0$. By Lemma 9.240, the first variation of the potential sector is

$$\delta\,\mathrm{Tr}_{\mathrm{FD}}\{\mathcal{V}_\Phi^{\mathrm{HS}}\} = \mathrm{Tr}_{\mathrm{FD}}\Big\{F'(s)\,\delta s\Big\}, \qquad F'(s) = \tfrac{1}{2}m_A^2 + \tfrac{\lambda_A}{3!}s + \sum_{k\geq 3}\tfrac{c_k}{(k-1)!}s^{k-1},$$

with $\delta s = \delta g_T^{\mathrm{HS}}(\hat{\mathbf{\Phi}},\hat{\mathbf{\Phi}}) = 2\,\mathrm{Re}\,g_T^{\mathrm{HS}}(\hat{\mathbf{\Phi}},\delta\hat{\mathbf{\Phi}})$ (Lemma 9.240). At $\hat{\mathbf{\Phi}} = 0$ one has $s = 0$ and $\delta s = 0$, so $\hat{\mathbf{\Phi}} = 0$ is always a stationary point. To classify it, consider the second variation at $\hat{\mathbf{\Phi}} = 0$:

$$\delta^2\mathrm{Tr}_{\mathrm{FD}}\{\mathcal{V}_\Phi^{\mathrm{HS}}\}\Big|_{\hat{\mathbf{\Phi}}=0} = m_A^2\;\mathrm{Tr}_{\mathrm{FD}}\Big\{\,g_T^{\mathrm{HS}}(\delta\hat{\mathbf{\Phi}},\delta\hat{\mathbf{\Phi}})\Big\}.$$

If $m_A^2 > 0$, positivity of g_T^{HS} implies $\delta^2 \geq 0$ with equality only for $\delta\hat{\mathbf{\Phi}} = 0$, so $\hat{\mathbf{\Phi}} = 0$ is a (strict) local minimum of the potential sector. Hence the *symmetric vacuum* is stationary (and, by coercivity, globally minimizing for sufficiently small higher couplings).

[6] These are the standard coercivity conditions for polynomial potentials on $[0,\infty)$ ensuring that the highest even-order coefficient is positive and dominates at large s.

(iii) "Mexican-hat" structure and symmetry breaking for $m_A^2 < 0$. If $m_A^2 < 0$ while $\lambda_A > 0$ (and higher terms obey the positivity bounds above), then F has a local maximum at $s = 0$ and, by coercivity, admits a nonzero minimizing radius $s_\star > 0$ solving $F'(s_\star) = 0$ with $F''(s_\star) > 0$:

$$\tfrac{1}{2}m_A^2 + \tfrac{\lambda_A}{3!}\, s_\star + \sum_{k\geq 3} \tfrac{c_k}{(k-1)!}\, s_\star^{k-1} = 0, \qquad F''(s_\star) = \tfrac{\lambda_A}{3!} + \sum_{k\geq 3} \tfrac{c_k}{(k-2)!}\, s_\star^{k-2} > 0. \tag{9.106}$$

In the quartic truncation ($c_k = 0$ for $k \geq 3$) this gives the familiar closed form $s_\star = \frac{3!}{2}\,\frac{|m_A^2|}{\lambda_A} > 0$. Since $\mathcal{V}_\Phi^{\mathrm{HS}}$ depends only on $s = g_T^{\mathrm{HS}}(\hat{\mathbf{\Phi}}, \hat{\mathbf{\Phi}})$ (Definitions 9.236 and 9.237) and is invariant under the HS gauge/frame group $U(n) \times O(r)$ (Proposition 9.238), the set of minimizers is the HS orbit at fixed radius $s_\star$:

$$\mathcal{M}_{\mathrm{vac}}^{(\mathrm{HS})} \;\cong\; \Big\{\hat{\mathbf{\Phi}} : g_T^{\mathrm{HS}}(\hat{\mathbf{\Phi}}, \hat{\mathbf{\Phi}}) = s_\star\Big\}/\Big(U(n) \times O(r)\Big),$$

i.e. a *Mexican-hat* degeneracy in the HS tangent norm. The associated symmetry breaking pattern and the ground-state selection are analyzed by the HS vacuum–manifold machinery of Chapter VI (Section 6.1.9), where the minimizing shell is encoded by the ground spectral projector on the HS vacuum manifold.

Conclusion. Under the stated positivity hypotheses, $\mathcal{V}_\Phi^{\mathrm{HS}}$ is bounded below; the symmetric vacuum $\hat{\mathbf{\Phi}} = 0$ is stationary and locally minimizing for $m_A^2 > 0$, while for $m_A^2 < 0$ the minimum occurs at nonzero HS radius $s_\star > 0$ with the usual Mexican-hat degeneracy in g_T^{HS}, completing the proof. □

Summary. The holosymmetric wordwise machinery replaces the standard invariant $g_T(\hat{\mathbf{\Phi}}, \hat{\mathbf{\Phi}})$ with its HS counterpart $g_T^{\mathrm{HS}}(\hat{\mathbf{\Phi}}, \hat{\mathbf{\Phi}})$ (Definition 6.106), builds the HS potential as a function of the scalar operator $\mathcal{I}_\Phi^{\mathrm{HS}}$ (9.97) via (9.98), proves gauge/frame invariance (Proposition 9.238), and integrates it with the HS measure (Definition 6.34) to contribute to the HS action (9.101), reducing to the standard form under the expectation push–through (Remark 6.35).

The excitations of the Gaiamiti field around this vacuum are what we observe as gravitational phenomena. This framework leads to a crucial distinction:

Definition 9.242 (Apeiron (trace, spin-0) and its intrinsic scale)**.** Work on the diagnostic realized reference configuration (Definition 9.185) in the Gaiamiti sector $S = \Phi\Phi$ with the trace type $[\chi] = \mathrm{Tr}$ (Definitions 9.195 and 9.200). Let $\mathcal{U}_{\Phi\Phi}$ be the potential of §9.17 and write its typed curvature (Hessian) at the certified background $\bar{\Phi}$ as $\mathcal{K}_{\Phi\Phi}^{[\mathrm{Tr}]}$ (Definition 9.220). Define the *Apeiron* to be the fundamental trace excitation and set its intrinsic scale $m_A > 0$ by the plateau identification

$$m_A^2 \;\equiv\; \mathrm{eig}_{\min}\Big(\mathcal{K}_{\Phi\Phi}^{[\mathrm{Tr}]}\Big) \quad \text{on the typed subspace.}$$

Then m_A is interpreted *not* as a matter mass but as the *geometric inertia / energy scale* of the Apeiron field: it fixes the characteristic interaction (correlation) length $\xi_A \sim m_A^{-1}$ for trace-type geometric fluctuations. Operationally this is the typed channel $(S, [\chi]) = (\Phi\Phi, \mathrm{Tr})$ in quadrant QI (law-symmetric, no state selection; Definition 9.189), with no NG/pNG content in this channel (Proposition 9.197). After stamping $\big(\mathrm{Loc}^{[\Phi\Phi,\mathrm{Tr}]}, \mathrm{On}^{[\Phi\Phi,\mathrm{Tr}]}\big)$ (Definition 9.190), the ledger row $\mathbf{L}^{[\Phi\Phi,\mathrm{Tr}]}$ is recorded (Definition 9.198 and Corollaries 9.194 and 9.218).

Proposition 9.243 (Graviton as the massless TT Goldstone in SSB)**.** *Consider the TT type* $[\chi] = \mathrm{TT}$ *(with optional helicities* $\lambda = \pm 2$ *when available; Definitions 9.195 and 9.200). Assume a law-symmetric, state-selecting diagnostic realized reference configuration (SSB-only regime) so that the typed order parameter* $\varphi^{[\mathrm{TT}]}$ *can align on a vacuum manifold* $\mathcal{M}_{\mathrm{vac}}$ *with continuous symmetry (§9.17). Then the typed quadrant is*

$$\mathrm{Quad}^{[\Phi\Phi,\mathrm{TT}]} = \mathrm{QII},$$

and the correlational typed channel $\Pi_{\mathrm{corr}}\Pi^{[\mathrm{TT}]}_{\Phi\Phi}$ *exhibits an* exact *NG pole on* H^0 *with positive residue (Definition 9.189 and Proposition 9.197). Equivalently, in the potential route,* $\mathcal{K}^{[\mathrm{TT}]}_{\Phi\Phi}$ *has a null direction tangent to* $\mathcal{M}_{\mathrm{vac}}$ *while the law source heals (Definitions 9.188 and 9.220 and Proposition 9.223). Hence the* graviton *is* not *the Apeiron: it is the massless spin-2 Goldstone excitation in the TT typed channel (cf. [93, 94]; see also Corollary 9.199 for naming after ledger population).*

Remark 9.244 (Separation of roles and operational workflow)**.** The Apeiron (trace, spin-0) carries the intrinsic geometric scale m_A and is typically QI (gapped radial), while the graviton (TT, spin-2) appears as the QII NG mode in the SSB-only regime. Both statements are decided *operationally* via the typed meters and healing indicators on the same diagnostic realized reference configuration (Definitions 9.187 to 9.189 and 9.205), and are then stamped with T–L–O and recorded in $\mathbf{L}^{[S,\chi]}$ (Definitions 9.190 and 9.198 and Corollary 9.218). This framework maintains a massless carrier for the observable gravitational interaction (TT NG mode) while distinguishing it sharply from the field's intrinsic energy scale m_A (trace channel).

Therefore, the potential $\mathcal{V}_\Phi$ plays a dual role: it sets the intrinsic properties of the underlying Apeiron field while its symmetries and minima dictate the conditions under which massless gravitons emerge as the observable, long-range carriers of the gravitational force. This is analogous to how massless phonons arise as collective excitations in a solid lattice composed of massive atoms [458, 391]. Similar analogies have been drawn in emergent gravity frameworks, where graviton-like modes are viewed as emergent collective phenomena [459, 460]. Therefore, the Apeiron can possess an intrinsic energy scale (m_A) while its observable, long-range excitation—the graviton—remains massless, consistent with observation.

With the potential operator now specified, it is natural to record the associated symmetry–breaking structure explicitly. This clarifies which symmetry group is broken by a chosen vacuum, what the resulting vacuum manifold looks like, and how the Goldstone directions are realized. The following proposition formalizes this data in the Gaiamiti sector.

Proposition 9.245 (SSB data of the Gaiamiti sector (typed TT identification))**.** *Let the modal Hilbert bundle carry structure group* $G \subseteq \mathrm{U}(4)$ *with* $u(4)$ *connection* $\mathcal{A}$ *(Section 4.1.4). Let the Gaiamiti potential* $\mathcal{V}_\Phi$ *be built from the invariant* $g_T(\hat{\mathbf{\Phi}}, \hat{\mathbf{\Phi}})$*. A realized vacuum* $\bar{\Phi}$ *spontaneously breaks* G *to its stabilizer* $H = \{\, g \in G \mid g \cdot \bar{\Phi} = \bar{\Phi} \,\}$ *with vacuum manifold* $\mathcal{M}_{\mathrm{vac}} \cong G/H$ *and Goldstone space* $T_{\bar{\Phi}}\mathcal{M}_{\mathrm{vac}}$*.*

On the diagnostic realized reference configuration (Definition 9.185), in the typed TT channel $[\chi] = \mathrm{TT}$ *(Definitions 9.186, 9.195 and 9.200), assume the SSB-only regime so*

that $\mathrm{Quad}^{[\Phi\Phi,\mathrm{TT}]} = \mathrm{QII}$ *(Definition 9.189). Then the graviton is identified as the TT-typed Goldstone deformation of the realized chronogeometry,*

$$\boxed{\delta\hat{g}^{[\mathrm{TT}]}_{\mu\nu} \;:=\; \Pi^{[\mathrm{TT}]}_{\Phi\Phi}\Big[\;\delta\hat{g}_{\mu\nu}\Big(\delta\Phi_{\mathrm{tang}}\Big)\;\Big]} \quad \text{on } \mathcal{H}_{\mathrm{phys}},$$

where $\delta\Phi_{\mathrm{tang}} \in T_{\bar{\Phi}}\mathcal{M}_{\mathrm{vac}}$ *and* $\Pi^{[\mathrm{TT}]}_{\Phi\Phi}$ *is the BRST-closed, cone-preserving typed projector that commutes with the triadic channels (Definition 9.186 and Proposition 9.202). Equivalently, with a* $\bar{\Sigma}$*-admissible spatial direction, one may refine to helicities*

$$\boxed{\delta\hat{g}^{[\lambda]}_{\mu\nu} := \Pi^{[\lambda]}_{\Phi\Phi}\,\delta\hat{g}_{\mu\nu}\Big(\delta\Phi_{\mathrm{tang}}\Big), \quad \lambda = \pm 2}\,.$$

The correlational TT channel $\Pi_{\mathrm{corr}}\Pi^{[\mathrm{TT}]}_{\Phi\Phi}$ *exhibits an exact NG pole on* H^0 *with positive residue (Proposition 9.197); the potential route yields the same decision via a null direction of the typed curvature* $\mathcal{K}^{[\mathrm{TT}]}_{\Phi\Phi}$ *(Definition 9.220 and Proposition 9.223). All statements are Nielsen-flat and BRST-safe (Propositions 9.191 and 9.207).*

9.17.1 The Apeiron as the Fundamental Quantum of Geometry

Definition 9.246 (Apeiron: typed formal definition and dynamics on H^0)**.** Work on the realized, diagnostic realized reference configuration (Definition 9.185) in the Gaiamiti sector $S = \Phi\Phi$. Let the dynamical field admit the background–fluctuation split (cf. eq. (10.87))

$$\hat{\mathbf{\Phi}}(x) = \bar{\Phi}(x)\,\mathbb{I}_x + \delta\hat{\mathbf{\Phi}}(x), \qquad \bar{\Phi}(x) = \langle\hat{\mathbf{\Phi}}(x)\rangle \;\; (\text{Quantum Holo–State } |\mathring{\mathbf{\Psi}}\rangle;\; \textit{Definition } 1.27),$$

so that $\langle\delta\hat{\mathbf{\Phi}}(x)\rangle = 0$. Insert the BRST–closed, cone–preserving *trace* projector $\Pi^{[\mathrm{Tr}]}_{\Phi\Phi}$ (Definitions 9.186, 9.195 and 9.200) and define the *Apeiron fluctuation*

$$\delta\hat{\mathbf{\Phi}}^{[\mathrm{Tr}]}(x) := \Pi^{[\mathrm{Tr}]}_{\Phi\Phi}\,\delta\hat{\mathbf{\Phi}}(x) \;\in\; \mathrm{Ran}\,\Pi^{[\mathrm{Tr}]}_{\Phi\Phi} \subset \mathcal{H}_{\mathrm{phys}}.$$

The **Apeiron** is the fundamental, operator–valued quantum of the trace–typed fluctuation $\delta\hat{\mathbf{\Phi}}^{[\mathrm{Tr}]}$.

Dynamics. Linearizing the Euler–Lagrange equations from the Gaiamiti action (cf. §6.5) and projecting to the trace type yields the covariant, operator–valued Klein–Gordon form (cf. eq. (10.126))

$$\Big(\bar{\nabla}^{\mu}\bar{\nabla}_{\mu} - \mathcal{M}_{[\mathrm{Tr}]}(x)\Big)\,\delta\hat{\mathbf{\Phi}}^{[\mathrm{Tr}]}(x) = 0,$$

where $\bar{\nabla}$ is the Levi–Civita derivative of $\bar{\Sigma}_{\mu\nu}$ (Definition 6.7) and

$$\mathcal{M}_{[\mathrm{Tr}]}(x) := \Pi^{[\mathrm{Tr}]}_{\Phi\Phi}\;\frac{\delta^2\mathcal{V}_{\Phi}}{\delta\hat{\mathbf{\Phi}}^{\dagger}\,\delta\hat{\mathbf{\Phi}}}\bigg|_{\hat{\mathbf{\Phi}}=\bar{\Phi}(x)}\;\Pi^{[\mathrm{Tr}]}_{\Phi\Phi}\,,$$

where this defines the typed effective mass operator (cf. eq. (10.125)). with $\mathcal{V}_{\Phi}$ the Gaiamiti potential (Definition 9.235).

Intrinsic scale. If $\mathcal{V}_\Phi \supset \frac{1}{2} m_A^2\, g_T(\hat{\mathbf{\Phi}}, \hat{\mathbf{\Phi}})$ (§9.17), identify the *intrinsic energy scale / geometric inertia*

$$m_A^2 \;\equiv\; \mathrm{eig}_{\min}\Big(\mathcal{K}_{\Phi\Phi}^{[\mathrm{Tr}]}\Big), \qquad \mathcal{K}_{\Phi\Phi}^{[\mathrm{Tr}]} := \Pi_{\Phi\Phi}^{[\mathrm{Tr}]}\, \frac{\delta^2 \mathcal{V}_\Phi}{\delta\hat{\mathbf{\Phi}}\,\delta\hat{\mathbf{\Phi}}}\bigg|_{\bar{\Phi}}\, \Pi_{\Phi\Phi}^{[\mathrm{Tr}]},$$

so that the characteristic interaction (correlation) length of trace–typed geometric fluctuations is $\xi_A \sim m_A^{-1}$ (Definition 9.220).

Quadrant and ledger. On a law–symmetric diagnostic realized reference configuration the Apeiron sits in the $[\chi] = \mathrm{Tr}$ typed channel with $\mathrm{Quad}^{[\Phi\Phi,\mathrm{Tr}]} = \mathrm{QI}$ (no NG/pNG in this channel; Definition 9.189 and Proposition 9.197). After stamping $(\mathrm{Loc}^{[\Phi\Phi,\mathrm{Tr}]}, \mathrm{On}^{[\Phi\Phi,\mathrm{Tr}]})$ (Definition 9.190), record the ledger row $\mathbf{L}^{[\Phi\Phi,\mathrm{Tr}]}$ (Definition 9.198 and Corollaries 9.194 and 9.218).

Separation from the graviton. The long–range gravitational carrier is *not* the Apeiron. It is the massless spin–2 Goldstone excitation in the TT typed channel under SSB, operationally identified by the TT projector and the typed QII predicate (Propositions 9.197 and 9.243 and Definition 9.189).

Remark 9.247 (Physical vs. pre-physical Apeiron)**.** Definitions and results in §9.17.1 are taken on the realized, diagnostic realized reference configuration (Definition 9.185), after plateau certification and projection to $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$. Accordingly, the *Apeiron* here is the $[\chi] = \mathrm{Tr}$ typed fluctuation evaluated on H^0, with meters, quadrant, and T–L–O recorded in the ledger (Definitions 9.190 and 9.198 and Corollary 9.218).

By contrast, the sub-Planckian (pre-physical) "Apeiron" in the pre-physics arena is merely an operator-level trace fluctuation: diagnostics may be performed, but no plateau certification and no projection to H^0 are permitted (*cf.* PQS–anchored scale principle and *No realization below the Planck scale*, Definition 9.1 and Lemma 9.3). We may write

$$\begin{aligned} \text{Pre-physical:} \quad & \delta\hat{\mathbf{\Phi}}_{\mathrm{pre}}^{[\mathrm{Tr}]} \quad (\text{operator for } \ell < \ell_P;\ \text{diagnostics only}) \\ \text{Physical:} \quad & \delta\hat{\mathbf{\Phi}}_{\mathrm{phys}}^{[\mathrm{Tr}]} := \Pi_{\Phi\Phi}^{[\mathrm{Tr}]}\, \delta\hat{\mathbf{\Phi}}\,\big|_{H^0,\,\mathfrak{P}} \end{aligned}$$

(physical typed fluctuation on H^0, $\ell \geq \ell_P$). Only the latter admits SynTh meters, quadrant assignment, and SSB semantics (Definitions 9.187 and 9.189 and Proposition 9.197).

Remark 9.248 (Philosophical Provenance of the Apeiron)**.** The name *Apeiron* is drawn from Anaximander's cosmology (ἄπειρον): the boundless, formless, and indefinite substratum of reality, from which order emerges through the action of delimiting principles (*peras*) [69, 70]. This reflects the Apeiron's role as the primordial excitation of geometric potential, prior to metricity, form, or localization.

Derivation of the Mass Operator $\mathcal{M}(x)$

Setup and conventions. Work on a certified plateau $\mathfrak{P}$ with BRST projection to $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ and band tags $P_x(\lambda_{\mathrm{UV}})$, $K_{\mathrm{IR}}^{(\ell)}$. The Gaiamiti potential $\mathcal{V}_\Phi$ enters the Lagrangian density

$$\mathcal{L}_\Phi(x) = \mathrm{Tr}\Big\{\bar{\Sigma}^{\mu\nu}(x)\, g_T\Big(\nabla_\mu\hat{\mathbf{\Phi}}(x),\, \nabla_\nu\hat{\mathbf{\Phi}}(x)\Big) + \mathcal{V}_\Phi\Big(\hat{\mathbf{\Phi}}(x)\Big)\Big\},$$

with $g_T : T\mathcal{M}_{\rm vac} \times T\mathcal{M}_{\rm vac} \to \mathrm{End}(\mathcal{H}_{\rm em})$, and $\hat{\mathbf{\Phi}}$ taking values in the vacuum tangent bundle $T\mathcal{M}_{\rm vac}$ (orthonormal bases from $T_{|\Omega\rangle}\mathcal{M}_{\rm vac}$ as needed).

The potential piece we use is (quadratic + quartic):

$$\boxed{\mathcal{V}_\Phi(\hat{\mathbf{\Phi}}) = \tfrac{1}{2}\, m_A^2\, g_T(\hat{\mathbf{\Phi}}, \hat{\mathbf{\Phi}}) + \tfrac{\lambda_A}{4!}\left[g_T(\hat{\mathbf{\Phi}}, \hat{\mathbf{\Phi}})\right]^2 + \cdots}$$

Typed projection and EL split. Decompose fluctuations by the BRST-closed typed projectors in the $\Phi\Phi$ block,

$$\delta\hat{\mathbf{\Phi}} = \delta\hat{\mathbf{\Phi}}^{[\mathrm{TT}]} \oplus \delta\hat{\mathbf{\Phi}}^{[\mathrm{Tr}]}, \qquad \delta\hat{\mathbf{\Phi}}^{[\chi]} := \Pi^{[\chi]}_{\Phi\Phi}\, \delta\hat{\mathbf{\Phi}},$$

so that, on $\mathcal{H}_{\rm phys}$, the typed EL equations read

$$\underbrace{\left(\bar\nabla^\mu\bar\nabla_\mu - \mathcal{M}_{[\mathrm{Tr}]}(x)\right)\delta\hat{\mathbf{\Phi}}^{[\mathrm{Tr}]}(x) = 0}_{\text{trace/radial channel}}, \qquad \underbrace{\Pi^{[\mathrm{TT}]}_{\Phi\Phi}\,\mathcal{L}_{\mathrm{TT}}[\bar\Sigma]\Big(\delta\hat g(\delta\Phi_{\rm tang})\Big) = 0}_{\text{TT/Goldstone channel}},$$

with $\bar\nabla$ the Levi–Civita connection of $\bar\Sigma$. The typed mass operator $\mathcal{M}_{[\mathrm{Tr}]}$ is obtained from the potential Hessian at the background $\bar\Phi$.

Local orthonormal coordinates. Choose a g_T–orthonormal basis $\{|e_i\rangle\}$ of $T_{|\Omega(x)\rangle}\mathcal{M}_{\rm vac}$ and write

$$\hat{\mathbf{\Phi}} = \sum_i \hat\Phi^i\, |e_i\rangle, \qquad S := g_T(\hat{\mathbf{\Phi}}, \hat{\mathbf{\Phi}}) = \sum_i (\hat\Phi^i)^\dagger \hat\Phi^i.$$

1. First variation w.r.t. $\hat{\mathbf{\Phi}}^\dagger$ (independent variables)

Treat $\hat{\mathbf{\Phi}}$ and $\hat{\mathbf{\Phi}}^\dagger$ as independent. Then

$$\frac{\delta\mathcal{V}_\Phi}{\delta\hat{\mathbf{\Phi}}^\dagger} = \tfrac{1}{2}\, m_A^2\, \hat{\mathbf{\Phi}} + \frac{\lambda_A}{4!}\,(2S)\,\hat{\mathbf{\Phi}} = \boxed{\tfrac{1}{2}\, m_A^2\, \hat{\mathbf{\Phi}} + \tfrac{\lambda_A}{12}\, S\, \hat{\mathbf{\Phi}}}.$$

2. Mixed second variation (mass operator before typing)

Differentiate once more w.r.t. $\hat{\mathbf{\Phi}}$. Componentwise,

$$\frac{\delta^2\mathcal{V}_\Phi}{\delta\hat\Phi^{j\dagger}\,\delta\hat\Phi^k} = \tfrac{1}{2}\, m_A^2\, \delta_{jk} + \tfrac{\lambda_A}{12}\left(S\,\delta_{jk} + \hat\Phi^j\hat\Phi^{k\dagger}\right).$$

Equivalently, as an operator on the field space,

$$\boxed{\frac{\delta^2\mathcal{V}_\Phi}{\delta\hat{\mathbf{\Phi}}^\dagger\,\delta\hat{\mathbf{\Phi}}} = \tfrac{1}{2}\, m_A^2\, \mathbb{I} + \tfrac{\lambda_A}{12}\Big(S\,\mathbb{I} + \hat{\mathbf{\Phi}}\otimes\hat{\mathbf{\Phi}}^\dagger\Big)}.$$

3. Typed projection and evaluation at the background

Define the *typed* mass operator by projection,

$$\boxed{\mathcal{M}_{[\mathrm{Tr}]}(x) := \Pi^{[\mathrm{Tr}]}_{\Phi\Phi}\, \frac{\delta^2 \mathcal{V}_\Phi}{\delta\hat{\mathbf{\Phi}}^\dagger\, \delta\hat{\mathbf{\Phi}}}\bigg|_{\hat{\mathbf{\Phi}}=\bar{\Phi}(x)} \Pi^{[\mathrm{Tr}]}_{\Phi\Phi}}.$$

Symmetric (unbroken) background $\bar{\Phi} = 0$. Here $S = 0$ and the rank-1 term drops out:

$$\boxed{\mathcal{M}_{[\mathrm{Tr}]}(x) = \tfrac{1}{2}\, m_A^2\, \Pi^{[\mathrm{Tr}]}_{\Phi\Phi}}, \qquad (\bar{\Phi} = 0).$$

Generic background $\bar{\Phi} \neq 0$. One obtains

$$\boxed{\mathcal{M}_{[\mathrm{Tr}]}(x) = \tfrac{1}{2}\, m_A^2\, \Pi^{[\mathrm{Tr}]}_{\Phi\Phi} + \tfrac{\lambda_A}{12}\Big(\|\bar{\Phi}\|^2_{g_T}\, \Pi^{[\mathrm{Tr}]}_{\Phi\Phi} + \Pi^{[\mathrm{Tr}]}_{\Phi\Phi}\, \bar{\Phi} \otimes \bar{\Phi}^\dagger\, \Pi^{[\mathrm{Tr}]}_{\Phi\Phi} \Big)}.$$

Remark on the "curvature" (same-variable) Hessian and the intrinsic scale. For potential-driven diagnostics also use the same-variable (curvature) Hessian

$$\mathcal{K}^{[\mathrm{Tr}]}_{\Phi\Phi} := \Pi^{[\mathrm{Tr}]}_{\Phi\Phi}\, \frac{\delta^2 \mathcal{V}_\Phi}{\delta\hat{\mathbf{\Phi}}\, \delta\hat{\mathbf{\Phi}}}\bigg|_{\bar{\Phi}} \Pi^{[\mathrm{Tr}]}_{\Phi\Phi},$$

and identify the Apeiron intrinsic scale via $m_A^2 \equiv \mathrm{eig}_{\min}\big(\mathcal{K}^{[\mathrm{Tr}]}_{\Phi\Phi}\big)$. At an unbroken symmetric background, $\mathcal{K}^{[\mathrm{Tr}]}_{\Phi\Phi} = m_A^2\, \Pi^{[\mathrm{Tr}]}_{\Phi\Phi}$, so $\mathcal{K}$ and the mixed-Hessian mass $\mathcal{M}$ differ by a conventional factor 2 at quadratic order. Use $\mathcal{M}$ in the EL equation and $\mathcal{K}$ in potential-curvature meters; both are band/diagnostic realized reference configuration-safe.

Typed EL, safety, and physical reading. The trace-typed equation

$$(\bar{\nabla}^2 - \mathcal{M}_{[\mathrm{Tr}]})\, \delta\hat{\mathbf{\Phi}}^{[\mathrm{Tr}]} = 0$$

is constructed on $\mathcal{H}_{\mathrm{phys}}$ with typed projectors and inherits Nielsen/BRST safety and plateau stability; poles/residues are to be read only after diagnostic realized reference configuration certification. The TT channel encodes the NG (massless spin-2) behaviour under SSB and is disjoint from the Apeiron mass.

Takeaway. With the above potential normalisation,

$$\boxed{\mathcal{M}_{[\mathrm{Tr}]}(\bar{\Phi} = 0) = \tfrac{1}{2}\, m_A^2\, \Pi^{[\mathrm{Tr}]}_{\Phi\Phi}}, \qquad \boxed{\mathcal{K}^{[\mathrm{Tr}]}_{\Phi\Phi}(\bar{\Phi} = 0) = m_A^2\, \Pi^{[\mathrm{Tr}]}_{\Phi\Phi}}.$$

9.17.2 Spontaneous Symmetry Breaking of the Gaiamiti Field

1. Symmetric potential on the diagnostic realized reference configuration (typed form)

Work on the diagnostic realized reference configuration (Definition 9.185) and evaluate the operator-valued Gaiamiti potential $\mathcal{V}_\Phi$ (Definition 9.235) built from the invariant inner product g_T (Definition 6.104, cf. §9.17). We adopt the SSB-friendly parametrization

$$\mathcal{V}_\Phi(\hat{\mathbf{\Phi}}) = \frac{\lambda_A}{4!}\left[g_T\big(\hat{\mathbf{\Phi}},\hat{\mathbf{\Phi}}\big) - v_A^2\right]^2 + \cdots,$$

which is manifestly G-invariant (law-symmetric). Define the intrinsic Apeiron scale by $m_A^2 := \frac{\lambda_A}{3!}\, v_A^2$ (cf. Definition 9.242). Expanding about the origin reproduces the usual tachyonic sign (the origin is not a minimum for $v_A^2 > 0$); m_A is read from the *typed curvature* rather than a particle pole.

For any typed projector $\Pi^{[\chi]}_{\Phi\Phi}$ with $[\chi] \in \{\mathrm{Tr}, \mathrm{TT}\}$ (Definitions 9.186, 9.195 and 9.200), define the *typed curvature (Hessian)*

$$\mathcal{K}^{[\chi]}_{\Phi\Phi} := \Pi^{[\chi]}_{\Phi\Phi}\, \frac{\delta^2 \mathcal{V}_\Phi}{\delta\Phi\,\delta\Phi}\Big|_{\bar{\Phi}}\, \Pi^{[\chi]}_{\Phi\Phi}, \qquad \bar{\Phi} \text{ the realized background on } \mathfrak{P}.$$

These typed curvatures feed the SynTh meters via the potential route (Definitions 9.220 and 9.221) and are equivalent, on a certified plateau, to the correlational route (Proposition 9.223).

Consistency note (unbroken vs. broken parametrizations). When discussing the *unbroken* symmetric phase at $\bar{\Phi} = 0$, either set $v_A = 0$ *in the above form* or instead use $\mathcal{V}_\Phi = \frac{1}{2} m_A^2\, g_T(\Phi,\Phi) + \frac{\lambda_A}{4!}[g_T(\Phi,\Phi)]^2 + \cdots$ with $m_A^2 > 0$. Both give a positive quadratic curvature at the origin in the trace channel and avoid tachyonic confusion.

2. Vacuum configuration and typed SSB predicate

The realized vacuum $\bar{\Phi}$ minimizes $\mathcal{V}_\Phi$ on the diagnostic realized reference configuration. The canonical regimes:

(A) Symmetric phase (no SSB). Here $\bar{\Phi} = 0$ and, with the unbroken parametrization (or $v_A = 0$ in the SSB form), both typed curvatures are positive on the plateau:

$$\mathcal{K}^{[\mathrm{TT}]}_{\Phi\Phi} > 0, \qquad \mathcal{K}^{[\mathrm{Tr}]}_{\Phi\Phi} > 0.$$

The typed SynTh meters (Definitions 9.187 and 9.205) yield $\mathrm{Quad}^{[\Phi\Phi,\mathrm{TT}]} = \mathrm{QI}$ and $\mathrm{Quad}^{[\Phi\Phi,\mathrm{Tr}]} = \mathrm{QI}$: no NG/pNG in either typed channel.

(B) SSB phase (law-symmetric, state-selecting). $\bar{\Phi} \neq 0$ with $g_T(\bar{\Phi},\bar{\Phi}) = v_A^2$. On the typed subspaces one has

$$\mathrm{spec}\,\mathcal{K}^{[\mathrm{TT}]}_{\Phi\Phi} \ni 0 \quad (\text{null directions tangent to } \mathcal{M}_{\mathrm{vac}} \cong G/H), \qquad \mathcal{K}^{[\mathrm{Tr}]}_{\Phi\Phi} > 0.$$

Potential-driven SynTh then gives

$$\mathrm{Quad}^{[\Phi\Phi,\mathrm{TT}]} = \mathrm{QII}, \qquad \mathrm{Quad}^{[\Phi\Phi,\mathrm{Tr}]} = \mathrm{QI},$$

with healing indicators $\mathscr{H}^{[\Phi\Phi,\mathrm{TT}]}_{L,S} = 0$ and $\mathscr{H}^{[\Phi\Phi,\mathrm{Tr}]}_{L,S} = 0$. Accordingly, the correlational typed channel $\Pi_{\mathrm{corr}}\Pi^{[\mathrm{TT}]}_{\Phi\Phi}$ exhibits an *exact* NG pole (massless spin-2), while the trace channel remains gapped. Location/onset are stamped via Definition 9.190 and recorded in the ledger (Definition 9.198 and Corollaries 9.194 and 9.218); SSB naming applies thereafter (Corollary 9.199).

Quadratic expansion of $\mathcal{K}^{[\chi]}_{\Phi\Phi}$ around $\bar{\Phi}$ (typed basis as in EL split). Setup. Use the SSB–friendly potential

$$\mathcal{V}_\Phi(\Phi) = \frac{\lambda_A}{4!}\left[g_T(\Phi,\Phi) - v_A^2\right]^2 + \cdots, \qquad m_A^2 := \frac{\lambda_A}{3!}\,v_A^2 = \frac{\lambda_A}{6}\,v_A^2.$$

Let $\bar{\Phi}$ be a realized vacuum with $g_T(\bar{\Phi},\bar{\Phi}) = v_A^2$, and define the g_T-unit radial vector $u := \bar{\Phi}/\|\bar{\Phi}\|_{g_T} = \bar{\Phi}/v_A$ (so $g_T(u,u) = 1$). Decompose any fluctuation as

$$\delta\Phi = \phi_r\, u + \delta\Phi_\perp, \qquad g_T(u,\delta\Phi_\perp) = 0.$$

The typed projectors on the Φ-space (same as in the EL split) are

$$\Pi^{[\mathrm{Tr}]}_{\Phi\Phi} X := u\, g_T(u,X), \qquad \Pi^{[\mathrm{TT}]}_{\Phi\Phi} X := X - u\, g_T(u,X), \qquad \Pi^{[\mathrm{Tr}]}_{\Phi\Phi} + \Pi^{[\mathrm{TT}]}_{\Phi\Phi} = \mathbb{I}.$$

Quadratic expansion. Write $S(\Phi) := g_T(\Phi,\Phi)$. Around $\bar{\Phi}$,

$$S(\bar{\Phi}+\delta\Phi) - v_A^2 = 2\,g_T(\bar{\Phi},\delta\Phi) + g_T(\delta\Phi,\delta\Phi) = 2v_A\,\phi_r + \mathcal{O}(\delta\Phi^2),$$

so the potential to quadratic order is

$$\mathcal{V}_\Phi(\bar{\Phi}+\delta\Phi) = \frac{\lambda_A}{4!}\left(2v_A\,\phi_r\right)^2 + \mathcal{O}(\delta\Phi^3) = \frac{\lambda_A v_A^2}{6}\,\phi_r^2 + \mathcal{O}(\delta\Phi^3) = m_A^2\,\phi_r^2 + \mathcal{O}(\delta\Phi^3).$$

Matching to the standard quadratic form $\frac{1}{2}\,\delta\Phi\cdot\mathcal{K}_{\Phi\Phi}(\bar{\Phi})\,\delta\Phi$, we read off

$$\boxed{\mathcal{K}_{\Phi\Phi}(\bar{\Phi}) = 2\,m_A^2\,\Pi^{[\mathrm{Tr}]}_{\Phi\Phi},} \quad \Longrightarrow \quad \boxed{\mathcal{K}^{[\mathrm{Tr}]}_{\Phi\Phi}(\bar{\Phi}) = 2\,m_A^2\,\Pi^{[\mathrm{Tr}]}_{\Phi\Phi}, \quad \mathcal{K}^{[\mathrm{TT}]}_{\Phi\Phi}(\bar{\Phi}) = 0.}$$

Interpretation.

- *Radial stiffness (trace channel).* Along the radial direction u, the curvature eigenvalue is $2m_A^2 = (\lambda_A/3)\,v_A^2$, so the trace-typed mode is gapped on the diagnostic realized reference configuration.
- *TT tangential zeros.* On the g_T-orthogonal complement (tangent to the vacuum orbit), the same-variable Hessian vanishes: $\ker\mathcal{K}_{\Phi\Phi}(\bar{\Phi}) = \mathrm{Im}\,\Pi^{[\mathrm{TT}]}_{\Phi\Phi}$. These directions coincide with the tangent space $T_{\bar{\Phi}}(\mathcal{M}_{\mathrm{vac}}) \cong T_{\bar{\Phi}}(G\cdot\bar{\Phi})$; they generate exact NG directions in the correlational TT channel on a certified plateau.

Unbroken phase check. If instead the symmetric phase is realized at $\bar{\Phi} = 0$, use the unbroken parametrization $\mathcal{V}_\Phi = \frac{1}{2}m_A^2\,g_T(\Phi,\Phi) + \frac{\lambda_A}{4!}[g_T(\Phi,\Phi)]^2 + \cdots$ (or set $v_A = 0$ above). Then

$$\mathcal{K}^{[\mathrm{Tr}]}_{\Phi\Phi}(\bar{\Phi}=0) = m_A^2\,\Pi^{[\mathrm{Tr}]}_{\Phi\Phi}, \qquad \mathcal{K}^{[\mathrm{TT}]}_{\Phi\Phi}(\bar{\Phi}=0) = m_A^2\,\Pi^{[\mathrm{TT}]}_{\Phi\Phi},$$

so both typed curvatures are positive (QI).

3. Physical consequences: typed spectrum on the diagnostic realized reference configuration

We analyze the spectrum of *realized* excitations—i.e. BRST–physical fluctuations on a certified plateau $\mathfrak{P}$—around a broken–symmetry vacuum. Expand the Gaiamiti field about its VEV,

$$\hat{\mathbf{\Phi}}(x) \;=\; \bar{\Phi}(x) \;+\; \delta\hat{\mathbf{\Phi}}(x) \qquad \text{(cf. eq. (10.87))}.$$

Fluctuations decompose by $\bar{\Sigma}$–typed projectors into *radial/trace* and *tangential/TT* directions (cf. Definitions 9.186, 9.195 and 9.200). On $\mathfrak{P}$, the potential and correlational routes agree within certified tolerances (Proposition 9.223). All NG/pNG claims are made *only* on $H^0(\hat{Q}_{\mathrm{BRST}})$ and are Nielsen–safe (Propositions 9.191 and 9.207).

Massive (gapped) trace/radial mode — the Apeiron. Displacements *up the walls* of the potential, away from the manifold of minima $\mathcal{M}_{\mathrm{vac}}$, feel a restoring force set by the curvature (Hessian) at the minimum; the resulting quanta are *massive* and define the intrinsic Apeiron scale (cf. standard treatments [91, 100]).

Typed implementation. Project to the trace type

$$\delta\hat{\mathbf{\Phi}}^{[\mathrm{Tr}]} := \Pi^{[\mathrm{Tr}]}_{\Phi\Phi}\,\delta\hat{\mathbf{\Phi}}, \qquad [\mathrm{Tr}] \in \{\mathrm{Tr}, \mathrm{TT}\}.$$

Define the typed mass operator at $\bar{\Phi}$ (mixed Hessian; Definition 9.246):

$$\mathcal{M}_{[\mathrm{Tr}]}(x) := \Pi^{[\mathrm{Tr}]}_{\Phi\Phi}\,\frac{\delta^2\mathcal{V}_\Phi}{\delta\hat{\mathbf{\Phi}}^\dagger\,\delta\hat{\mathbf{\Phi}}}\bigg|_{\hat{\mathbf{\Phi}}=\bar{\Phi}(x)}\,\Pi^{[\mathrm{Tr}]}_{\Phi\Phi}, \quad \mathcal{V}_\Phi \text{ as in Definition 9.235.}$$

Equivalently (for potential–side meters), use the same–variable curvature $\mathcal{K}^{[\mathrm{Tr}]}_{\Phi\Phi}$ (Definition 9.220). On $\mathfrak{P}$, positivity of $\mathcal{M}_{[\mathrm{Tr}]}$ yields a *gapped* spectrum with intrinsic Apeiron scale m_A (cf. Definition 9.242) and correlation length $\xi_A \sim m_A^{-1}$. The canonical typed predicate returns

$$\mathrm{Quad}^{[\Phi\Phi,\mathrm{Tr}]} \;=\; \mathrm{QI},$$

i.e. law–symmetric, no state selection and no NG/pNG content in this channel (Definition 9.189 and Proposition 9.197). Stamp Loc/On via Definition 9.190 and record $\mathbf{L}^{[\Phi\Phi,\mathrm{Tr}]}$ (Definition 9.198 and Corollaries 9.194 and 9.218).

Massless (NG) TT/tangential mode — the graviton. Displacements *along the flats*—tangent to the degenerate vacuum manifold $\mathcal{M}_{\mathrm{vac}}$—cost no potential energy and correspond to broken generators. By **Goldstone's theorem** [93, 94], these are *massless* excitations; in gauge-organized settings, their fate and identification follow the familiar organizational mechanisms [97, 91, 100].

Typed implementation. Project to the transverse–traceless type,

$$\delta\hat{\mathbf{\Phi}}^{[\mathrm{TT}]} := \Pi^{[\mathrm{TT}]}_{\Phi\Phi}\,\delta\hat{\mathbf{\Phi}}.$$

In the SSB–only regime ($\bar{\Phi} \neq 0$, no explicit breaking, $\mathcal{M}_{\mathrm{vac}} \cong G/H$), the typed curvature exhibits null tangential directions

$$\mathrm{spec}\,\mathcal{K}^{[\mathrm{TT}]}_{\Phi\Phi} \ni 0 \quad (\text{tangent to } T_{\bar{\Phi}}\mathcal{M}_{\mathrm{vac}}),$$

and the typed law/state sources *heal* on $\mathfrak{P}$, $\mathscr{H}_{L,S}^{[\Phi\Phi,\mathrm{TT}]} = 0$ (Definition 9.188). Hence the canonical predicate gives

$$\mathrm{Quad}^{[\Phi\Phi,\mathrm{TT}]} \;=\; \mathrm{QII}.$$

The correlational typed channel $\Pi_{\mathrm{corr}}\Pi_{\Phi\Phi}^{[\mathrm{TT}]}$ then displays an *exact* NG pole with positive residue on H^0 (Proposition 9.197). With a $\bar{\Sigma}$–admissible spatial splitting, refine to helicities $\lambda = \pm 2$ (Definition 9.200); these two physical polarizations propagate as the *massless* spin–2 graviton (with operational identification in Proposition 9.243 and naming per Corollary 9.199).

Diagnostic realized reference configuration–discipline summary. All statements above are evaluated on a certified plateau $\mathfrak{P}$ with band tags $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$, and with BRST projection to $H^0(\hat{Q}_{\mathrm{BRST}})$. On such windows, the potential– and correlational–route meters are equivalent and all conclusions are gauge–parameter independent, within the established tolerances (Propositions 9.191, 9.207 and 9.223).

Remark (explicit breaking / pNG). If a small explicit symmetry–breaking spurion is present in the TT law channel, the typed law source does not heal, $\mathscr{H}_{L}^{[\Phi\Phi,\mathrm{TT}]} > 0$, and the canonical predicate switches to

$$\mathrm{Quad}^{[\Phi\Phi,\mathrm{TT}]} = \mathrm{QIV}.$$

On the certified plateau $\mathfrak{P}$, this yields a pseudo–NG band with GMOR slope (Definition 9.188 and Proposition 9.197), while the trace channel remains gapped (QI). Boundary cases proceed *mutatis mutandis* via the bulk/edge overlay (Definition 9.214); only the Loc/On stamps and the edge entry recorded in the ledger are affected (Corollary 9.218).

9.17.3 Global Symmetry Breaking: NG and Radial Sectors

Definition 9.249 (Typed NG subspace and projector (global SSB))**.** Work on a certified plateau $\mathfrak{P}$ with band tags $P_x(\lambda_{\mathrm{UV}})$, $K_{\mathrm{IR}}^{(\ell)}$ and on the BRST–physical sector $H^0(\hat{Q}_{\mathrm{BRST}})$ (diagnostic realized reference configuration discipline as in Propositions 9.191, 9.207 and 9.223). Let a global symmetry G act on a field bundle $\mathcal{E}_U$ with $U \in \{\Theta, \Phi\}$ via a representation ρ, and let $\bar{U}$ minimize the effective potential. For each broken generator T_b, set $v_b := \rho(T_b)\bar{U} \neq 0$. With the appropriate pairing g (for Tantrakyma $g = g_C$, for Gaiamiti $g = g_T$), define the Gram matrix $B_{bc} := g(v_b, v_c)$ and the *typed NG projector* by

$$\boxed{\Pi_{\mathrm{NG}}^{[\chi]}(V) \;:=\; \Pi_{UU}^{[\chi]}\left[\sum_{b,c}(B^{-1})^{bc}\, g\Big(V,\, v_b\Big)\, v_c\right]\Pi_{UU}^{[\chi]},}$$

where $\Pi_{UU}^{[\chi]}$ is the typed projector (cf. Definitions 9.186 and 9.200). Equivalently, $\Pi_{\mathrm{NG}}^{[\chi]}$ is the Riesz (resolvent) projector onto the NG island of the *typed* scalar Hessian $\mathcal{K}_{UU}^{[\chi]}$ obtained by inserting the potential curvature in the same typed/triadic channel (contour enclosing the cluster at/near 0 in $\operatorname{spec}\mathcal{K}_{UU}^{[\chi]}$; cf. Proposition 9.223). All insertions commute with the triadic channels Π_α and are BRST–closed (Proposition 9.207).

Definition 9.250 (Typed radial direction, projector, and mass operator)**.** In the same setting and type $[\chi]$, introduce the quadratic form $\mathsf{X}_U := g(U,U)$ with differential at $\bar{U}$, $\delta\mathsf{X}_U|_{\bar{U}}(\delta U) = 2\,g(\delta U, \bar{U})$. The g–Riesz identification makes the *radial direction* collinear with $\bar{U}$; a normalized typed representative is

$$\widehat{\mathsf{R}}_U^{[\chi]} := \frac{\Pi_{UU}^{[\chi]}\bar{U}}{\sqrt{g\Big(\Pi_{UU}^{[\chi]}\bar{U},\ \Pi_{UU}^{[\chi]}\bar{U}\Big)}}.$$

Define the one–dimensional g–orthogonal *typed radial projector*

$$\boxed{\Pi_{\mathrm{rad}}^{[\chi]}(V) := \frac{g\Big(V, \widehat{\mathsf{R}}_U^{[\chi]}\Big)}{g\Big(\widehat{\mathsf{R}}_U^{[\chi]}, \widehat{\mathsf{R}}_U^{[\chi]}\Big)}\,\widehat{\mathsf{R}}_U^{[\chi]}.}$$

The *typed radial mass operator* (restricted Hessian at $\bar{U}$) is then

$$\boxed{\mathcal{K}_{\mathrm{rad}}^{[\chi]} := \Pi_{\mathrm{rad}}^{[\chi]}\ \frac{\delta^2\mathcal{V}_U}{\delta U\,\delta U}\Big|_{\bar{U}}\ \Pi_{\mathrm{rad}}^{[\chi]}}$$

to be evaluated on $H^0(\hat{Q}_{\mathrm{BRST}})$. For $U = \Theta$ (Tantrakyma) this reproduces the Higgs/radial kymon construction (Definition 9.343 and eq. (9.185)); for $U = \Phi$ (Gaiamiti) it matches the Apeiron radial operator (Definition 9.259). Typed/triadic insertions follow Definition 9.187, guaranteeing gauge–parameter independence within tolerances (Propositions 9.191 and 9.207).

Proposition 9.251 (Quadrant assignment under global SSB (typed, BRST–clean))**.** *Work on a certified plateau $\mathfrak{P}$ with band tags $P_x(\lambda_{\mathrm{UV}})$, $K_{\mathrm{IR}}^{(\ell)}$, and on the BRST–physical sector $H^0(\hat{Q}_{\mathrm{BRST}})$ (diagnostic realized reference configuration discipline: Propositions 9.191, 9.207 and 9.223). Fix a field species $U \in \{\Theta, \Phi\}$ and a type $[\chi]$ with typed projector $\Pi_{UU}^{[\chi]}$ (Definitions 9.186 and 9.200). Let $\Pi_{\mathrm{NG}}^{[\chi]}$ and $\mathcal{K}_{\mathrm{rad}}^{[\chi]}$ be the typed NG projector and typed radial mass operator from Definitions 9.249 and 9.250, and let $\delta_{L,S}^{[UU,\chi]}$, $\mathscr{H}_{L,S}^{[UU,\chi]}$ be the typed sources and healing indicators (Definitions 9.187 and 9.188).*

(i) (QI) No SSB, law symmetric (fully symmetric typed channel)

If $\operatorname{Im}\Pi_{\mathrm{NG}}^{[\chi]} = \{0\}$, the typed curvature has no nulls at the origin, the radial operator is strictly positive, and the typed witnesses indicate law/state symmetry ($|\Lambda_{\mathrm{law}}| \approx 1$, $|\Lambda_{\mathrm{state}}| \approx 1$ on $\mathfrak{P}$; cf. Definitions 7.685 and 9.187), then

$$\mathrm{Quad}^{[UU,\chi]} = \mathrm{QI}.$$

This case exhibits no NG/pNG content; spectra are gapped (or, for gauge/spacetime symmetries, carry the usual unbroken massless content) and all typed sources heal within tolerances (Proposition 9.223).

(ii) (QII) Global SSB without explicit breaking (SSB–only)

Assume a broken–symmetry vacuum $\bar{U}$ with $\operatorname{Im}\Pi_{\mathrm{NG}}^{[\chi]} \neq \{0\}$ and a strictly positive radial

gap $\mathcal{K}_{\mathrm{rad}}^{[\chi]} \geq m_{\mathrm{rad}}^2 \Pi_{\mathrm{rad}}^{[\chi]}$ *on* H^0 *for some* $m_{\mathrm{rad}} > 0$. *If the typed law/state sources* heal *on* $\mathfrak{P}$,

$$\mathscr{H}_L^{[UU,\chi]} = \mathscr{H}_S^{[UU,\chi]} = 0,$$

then the canonical typed predicate (Definition 9.189) returns

$$\mathrm{Quad}^{[UU,\chi]} = \mathrm{QII}.$$

Moreover, in the correlational channel $\Pi_{\mathrm{corr}} \Pi_{UU}^{[\chi]} \Pi_{\mathrm{NG}}^{[\chi]}$ *there is an* exact *NG pole with positive residue on* H^0 *(Proposition 9.197), while the radial sector is gapped by* $\mathcal{K}_{\mathrm{rad}}^{[\chi]}$. Helicity refinement *(when admissible) proceeds via the typed subprojectors in Definition 9.200.*

(iii) (QIII) Apparent law asymmetry that heals in the IR (emergent symmetry)
If the typed law-side deficit is nonzero at finite resolution but decays with IR coarse–graining on $\mathfrak{P}$ *(healing power law as in eq. (7.746)), while the state side remains symmetric (*$|\Lambda_{\mathrm{state}}| \approx 1$*), then*

$$\mathrm{Quad}^{[UU,\chi]} = \mathrm{QIII}.$$

There is no persistent NG/pNG content in the IR limit; the decision follows the canonical rule Definition 7.685 using the typed meters in Definition 9.187.

(iv) (QIV) Small explicit breaking in the typed law channel (pseudo–NG/GMOR)
Suppose the SSB hypotheses of (QII) *hold but a small explicit spurion is present in the* law *channel so that*

$$\mathscr{H}_L^{[UU,\chi]} > 0 \quad \text{on } \mathfrak{P},$$

while the state side still heals ($\mathscr{H}_S^{[UU,\chi]} = 0$*). Then the canonical typed predicate flips to*

$$\mathrm{Quad}^{[UU,\chi]} = \mathrm{QIV}.$$

On $\mathfrak{P}$, *the NG cluster becomes a* pseudo–NG *cluster obeying GMOR scaling (typed Ward/GMOR on the diagnostic realized reference configuration; cf. Definition 9.188 and Proposition 9.197 and the GMOR plateau result cited earlier), whereas the radial sector remains gapped and Nielsen–safe (Propositions 9.191 and 9.207).*

Stamps and ledger. *In all cases, record the typed* Loc/On *stamps via Definition 9.190 and update the typed quadrant ledger* $\mathbf{L}^{[UU,\chi]}$ *(Definition 9.198 and Corollaries 9.194 and 9.218). Boundary overlays are handled* mutatis mutandis *by Definition 9.214, affecting only the recorded* Loc/On *and edge entries.*

Scope of Gauging (Internal vs. Relational)

Definition 9.252 (Internal gauging of a typed sector on the diagnostic realized reference configuration)**.** Work on a certified plateau $\mathfrak{P}$ with band tags $P_x(\lambda_{\mathrm{UV}})$, $K_{\mathrm{IR}}^{(\ell)}$ and BRST projection to $H^0(\hat{Q}_{\mathrm{BRST}})$ (diagnostic realized reference configuration discipline: Propositions 9.191, 9.207 and 9.223). Let $U \in \{\Theta, \Phi\}$ denote Tantrakyma/Gaiamiti with pairing g (for Θ: $g = g_C$; for

Φ: $g = g_T$), equipped with an internal gauge group G_U, representation ρ_U, and coupling g_U. On a realized vacuum $\bar{U}$, the *typed* vector mass matrix is

$$\boxed{(M_U^2)_{IJ} \;=\; g_U^2\, g\Big(\rho_U(T_I)\bar{U},\ \rho_U(T_J)\bar{U}\Big)}\,,$$

with transverse/longitudinal/helicity projectors inserted as in Definition 9.350. The *radial (amplitude) scalar* is defined by the typed radial Hessian $\mathcal{K}^{[\chi]}_{\mathrm{rad}}$ of Definition 9.250, evaluated with the sector pairing g. Specializations:

(i) Tantrakyma Θ
$(M_\Theta^2)_{IJ} = g_\Theta^2\, g_C\Big(\rho_\Theta(T_I)\bar{\Theta},\, \rho_\Theta(T_J)\bar{\Theta}\Big)$; the Higgs (radial kymon) mass follows from $\mathcal{K}^{[\chi]}_{\mathrm{rad}}$ (cf. Definition 9.343 and eqs. (9.184) and (9.185)).

(ii) Gaiamiti Φ
$(M_\Phi^2)_{IJ} = g_\Phi^2\, g_T\Big(\rho_\Phi(T_I)\bar{\Phi},\, \rho_\Phi(T_J)\bar{\Phi}\Big)$; the Apeiron (radial) scalar is encoded by $\mathcal{K}^{[\chi]}_{\mathrm{rad}}$ (cf. Section 9.17.4, Proposition 9.257, and Definition 9.259).

Gauge fixing is taken in R_ξ form so that physical poles and quadrant decisions are ξ–independent (Nielsen safety; cf. Definition 9.261 and Proposition 9.191).

Remark 9.253 (Relational/diffeomorphism symmetry is not BEH–gauged)**.** Relational/diffeomorphism symmetry is not an internal gauging in the BEH sense. Typed/triadic Ward identities keep the TT channel *gapless* on H^0 (no Proca mass for the graviton); the trace scale m_A encodes *radial stiffness* in the Apeiron channel and is not a graviton mass (cf. Proposition 9.207 and Section 9.17.2).

Why Internal Gauging is Added (Completeness w.r.t. Global SSB)

Proposition 9.254 (Typed spectrum and quadrants under internal gauging)**.** *The quadrant formalism of Chs. 7–8 (typed/triadic meters on H^0; Definitions 7.685 and 9.187) already gives a complete account of* global *SSB (see Section 9.17.3 and Proposition 9.251). Adding* internal *gauging for Θ and Φ completes the symmetry–breaking picture without altering the quadrant axes (law vs. state):*

(i) Tantrakyma Θ (BEH)
Broken NG directions in $\mathrm{Im}\,\Pi^{[\chi]}_{\mathrm{NG}}$ *are eaten; internal vectors acquire masses $(M_\Theta^2)_{IJ}$ of Definition 9.252, while the Higgs (radial kymon) remains gapped by $\mathcal{K}^{[\chi]}_{\mathrm{rad}}$ (Definition 9.343 and eq. (9.185)). Typed/triadic meters certify* $\mathrm{Quad}^{[\Theta,\chi]} = \mathrm{QII}$ *for SSB–only, flipping to* QIV *with small explicit breaking (GMOR on the plateau; cf. Definition 9.188 and Proposition 9.197).*

(ii) Gaiamiti Φ (internal vectors + Apeiron)
An internal G_Φ yields massive internal vectors with $(M_\Phi^2)_{IJ}$ per Definition 9.252. The Apeiron (trace) scalar is controlled by $\mathcal{K}^{[\chi]}_{\mathrm{rad}}$ (cf. Section 9.17.4, Proposition 9.257, and Definition 9.259), while the TT graviton channel remains gapless by Ward protection (Remark 9.253). Typed quadrant decisions follow Proposition 9.251.

(iii) Quadrant exhaustiveness unchanged
Internal gauging rearranges *DOFs (NG → longitudinal vectors) but leaves the (law, state) axes intact. Thus* {QI, QII, QIII, QIV} *remain exhaustive, with the same typed decision rule (Definitions 7.685 and 9.187; cf. Ch. 8).*

(iv) Diagnostic realized reference configuration certification and gauge independence
With R_ξ fixing and BRST closure, physical poles (Higgs/Apeiron radials, transverse vectors) and the typed quadrant assignment are ξ–independent (Nielsen) and route–equivalent within tolerances (Definition 9.261 and Propositions 9.191, 9.207 and 9.223).

Record Loc/On *stamps and update the typed ledger* $\mathbf{L}^{[U,\chi]}$ *as in Definitions 9.190 and 9.198 and Corollary 9.218.*

9.17.4 Internal Gauging of the Gaiamiti Sector and BEH Completion

On the certified plateau $\mathfrak{P}$ with BRST projection to $H^0(\hat{Q}_{\mathrm{BRST}})$ (diagnostic realized reference configuration discipline as in Propositions 9.191, 9.207 and 9.223), an *internal* gauging of the Gaiamiti sector implements the Brout–Englert–Higgs (BEH) pattern for the Φ-coupled vectors, supplies a geometric Higgs–type scalar in the Apeiron (trace) channel, and *does not* generate a mass in the transverse–traceless (TT) graviton channel. The TT mode remains massless on H^0 (cf. typed TT construction in Section 9.17.2); the trace scale m_A encodes Apeiron stiffness, not a graviton mass (cf. radial operator in Definition 9.250).

Definition 9.255 (Gauge–coupled Gaiamiti action)**.** Let a compact Lie group G act on the Gaiamiti fiber $\mathcal{E}_\Phi$ via a representation ρ_Φ with generators T_I. Introduce gauge fields A^I_μ with field strength $F^I_{\mu\nu}$ and kinetic metric G_{IJ}. Extend the split connection (cf. Section 6.12.2 and eq. (6.314)) to

$$D_\mu \hat{\mathbf{\Phi}} := \nabla^{(\mathrm{rel})}_\mu \hat{\mathbf{\Phi}} + \partial_\mu \hat{\mathbf{\Phi}} + i\, g\, A^I_\mu\, \rho_\Phi(T_I)\, \hat{\mathbf{\Phi}},$$

and define, on a certified plateau $\mathfrak{P}$,

$$S_{\Phi+G} = \int_{\mathcal{X}_q} d^d x \, \sqrt{-\det \bar{\Sigma}}\, \mathrm{Tr}\Big[-\tfrac{1}{4}\, G_{IJ} F^I_{\mu\nu} F^{J\,\mu\nu} + \tfrac{1}{2}\, (D_\mu \hat{\mathbf{\Phi}})^\dagger g_T(D^\mu \hat{\mathbf{\Phi}}) - \mathcal{V}_\Phi(\hat{\mathbf{\Phi}})\Big]\,,$$

where g_T is the Gaiamiti pairing and $\mathcal{V}_\Phi$ the Gaiamiti potential (cf. eq. (6.316)); see also the refined Noether current in Definition 9.271. Typed/triadic insertions follow Definition 9.187.

Definition 9.256 (SSB pattern and typed projectors in the Gaiamiti sector)**.** Let $\hat{\mathbf{\Phi}} = \bar{\Phi} + \delta\hat{\mathbf{\Phi}}$ with $\bar{\Phi}$ minimizing $\mathcal{V}_\Phi$. For broken generators T_b, set $v_b := \rho_\Phi(T_b)\bar{\Phi} \neq 0$ and form

$$\Pi^{(\Phi)}_{\mathrm{NG}}(V) := \sum_{b,c} (B^{-1})^{bc}\, g_T(V, v_b)\, v_c, \qquad B_{bc} := g_T(v_b, v_c),$$

which is the Gaiamiti specialization of the typed NG projector in Definition 9.249. Let $\mathsf{Y} := g_T(\hat{\mathbf{\Phi}}, \hat{\mathbf{\Phi}})$. At $\bar{\Phi}$, $\delta\mathsf{Y}|_{\bar{\Phi}}(\delta\hat{\mathbf{\Phi}}) = 2\, g_T(\delta\hat{\mathbf{\Phi}}, \bar{\Phi})$, so the g_T–Riesz map identifies the radial line with $\bar{\Phi}$. A normalized typed representative is

$$\widehat{\mathsf{R}}_\Phi := \frac{\Pi^{[\chi]}_{\Phi\Phi}\, \bar{\Phi}}{\sqrt{g_T(\Pi^{[\chi]}_{\Phi\Phi}\bar{\Phi}, \Pi^{[\chi]}_{\Phi\Phi}\bar{\Phi})}}, \quad \text{(cf. typed projector in Definitions 9.186 and 9.200)}$$

with one–dimensional g_T–orthogonal projector

$$\Pi_{\mathrm{A}}^{(\Phi)}(V) := \frac{g_T(V, \widehat{\mathsf{R}}_\Phi)}{g_T(\widehat{\mathsf{R}}_\Phi, \widehat{\mathsf{R}}_\Phi)}\, \widehat{\mathsf{R}}_\Phi,$$

which matches the typed radial projector of Definition 9.250. Assume $\Pi_{\mathrm{NG}}^{(\Phi)}$ and $\Pi_{\mathrm{A}}^{(\Phi)}$ commute with the triadic projectors $\Pi_{\mathrm{corr}}, \Pi_{\mathrm{vac}}, \Pi_{\mathrm{spec}}$ (cf. Proposition 9.207) and respect typed polarizations (TT/trace/helicity; cf. Definition 9.200).

Proposition 9.257 (Vector masses from BEH in the Gaiamiti sector)**.** *Expanding the covariant kinetic term in Definition 9.255 around $\bar{\Phi}$ yields the internal vector mass term*

$$\mathcal{L}_{\mathrm{mass}}^{(A)} = \tfrac{1}{2}\, A_\mu^I\, (M^2)_{IJ}\, A^{J\,\mu}, \qquad (M^2)_{IJ} := g^2\, g_T\Big(\rho_\Phi(T_I)\bar{\Phi},\ \rho_\Phi(T_J)\bar{\Phi}\Big),$$

which is the Φ-sector specialization of the unified typed formula in Definition 9.252. Vectors along broken directions acquire masses $\mathrm{spec}(M^2)$*; NG directions are eaten as longitudinal modes, in parallel with the Tantrakyma linear-response paradigm (eq. (9.184) and Definition 9.343). Typed vector/helicity projectors are as in Definition 9.350.*

Proof of Proposition 9.257. Work on a certified plateau $\mathfrak{P}$ with BRST projection to $H^0(\hat{Q}_{\mathrm{BRST}})$ and typed/triadic meters commuting (cf. Propositions 9.191, 9.207 and 9.223). Start from the gauge–coupled Gaiamiti action of Definition 9.255:

$$S_{\Phi+G} = \int_{\mathcal{X}_q} d^d x\, \sqrt{-\det \bar{\Sigma}}\, \mathrm{Tr}\Big[-\tfrac{1}{4}\, G_{IJ} F_{\mu\nu}^I F^{J\,\mu\nu} + \tfrac{1}{2}\, (D_\mu \hat{\mathbf{\Phi}})^\dagger g_T (D^\mu \hat{\mathbf{\Phi}}) - \mathcal{V}_\Phi(\hat{\mathbf{\Phi}})\Big],$$

with $D_\mu \hat{\mathbf{\Phi}} = \nabla_\mu^{(\mathrm{rel})} \hat{\mathbf{\Phi}} + \partial_\mu \hat{\mathbf{\Phi}} + i\, g\, A_\mu^I\, \rho_\Phi(T_I) \hat{\mathbf{\Phi}}$. Expand around a realized vacuum $\hat{\mathbf{\Phi}} = \bar{\Phi} + \delta\hat{\mathbf{\Phi}}$ (Definition 9.256), keeping terms at most quadratic in fields and focusing on the gauge–vector sector. Using linearity of D_μ and sesquilinearity of g_T,

$$\begin{aligned} \tfrac{1}{2}(D_\mu \hat{\mathbf{\Phi}})^\dagger g_T (D^\mu \hat{\mathbf{\Phi}}) &= \tfrac{1}{2}\Big(\cdots + i g\, A_\mu^I\, \rho_\Phi(T_I)\bar{\Phi} + \cdots \Big)^\dagger\, g_T \Big(\cdots + i g\, A^{J\,\mu}\, \rho_\Phi(T_J)\bar{\Phi} + \cdots \Big) \\ &\supset \tfrac{1}{2}\, g^2\, A_\mu^I\, A^{J\,\mu}\, g_T\Big(\rho_\Phi(T_I)\bar{\Phi},\ \rho_\Phi(T_J)\bar{\Phi}\Big), \end{aligned}$$

which gives the purely quadratic $A\cdot A$ mass term $\mathcal{L}_{\mathrm{mass}}^{(A)} = \frac{1}{2} A_\mu^I (M^2)_{IJ} A^{J\,\mu}$ with $(M^2)_{IJ} := g^2\, g_T(\rho_\Phi(T_I)\bar{\Phi}, \rho_\Phi(T_J)\bar{\Phi})$. Cross terms linear in A_μ and $\delta\hat{\mathbf{\Phi}}$ generate the familiar A–NG mixing; these are cancelled on the diagnostic realized reference configuration by the R_ξ gauge fixing of Definition 9.261, and the associated ghost sector ensures BRST closure. Consequently, physical poles reside in the transverse vector channel, while longitudinal contributions are BRST-removed; gauge-parameter independence of masses and residues follows from the Nielsen/BRST identities (cf. Theorem 9.351).

If $T_a \in \mathfrak{h} = \mathrm{Lie}(H)$ is unbroken, then $\rho_\Phi(T_a)\bar{\Phi} = 0$ (Definition 9.256), and the corresponding rows/columns of M^2 vanish, yielding massless vectors for unbroken directions. For broken generators T_b one has $\rho_\Phi(T_b)\bar{\Phi} \neq 0$, producing strictly positive entries in M^2 and hence massive vectors along those directions. This is the Φ-sector specialization of the unified typed internal-gauging formula referenced in Definition 9.252. The statement about NG directions being eaten (longitudinal modes) and the use of typed vector/helicity projectors follows exactly as in the Tantrakyma linear-response paradigm (eq. (9.184) and Definition 9.343) and with the vector projectors of Definition 9.350. This completes the proof. □

Definition 9.258 (Mition: Gaiamiti spin–1 internal vector)**.** Let G act on the Gaiamiti fiber $\mathcal{E}_\Phi$ via ρ_Φ and let $\bar{\Phi}$ realize spontaneous breaking $G \to H$ (Definition 9.256). The *mition* denotes a one-particle, spin–1, *internal* gauge quantum in the Gaiamiti sector, i.e. a mass eigenmode of the bundle connection A^I_μ defined as follows.

(i) Mass matrix and eigenmodes

Form the BEH mass matrix

$$(M^2_\Phi)_{IJ} \;:=\; g^2\, g_T\Big(\rho_\Phi(T_I)\bar{\Phi},\ \rho_\Phi(T_J)\bar{\Phi}\Big) \quad \text{(cf. Definition 9.255 and Proposition 9.257).}$$

Choose an orthogonal diagonalization $O \in \mathrm{O}(\dim G)$ with $O^\top M^2_\Phi O = \mathrm{diag}(m^2_{V,a})$. The *mition fields* are the mass–eigenstate vectors

$$V_{a\,\mu} \;:=\; O^{\ I}{}_{a}\, A^I_\mu\,, \qquad m^2_{V,a} \in \mathrm{spec}(M^2_\Phi).$$

(ii) Physical pole definition (diagnostic realized reference configuration)

On the diagnostic realized reference configuration $\mathfrak{P}$ with BRST projection to $H^0(\hat{Q}_{\mathrm{BRST}})$, the *physical mition* of label a is the simple pole of the two–point function in the *transverse* vector channel at $k^2 = m^2_{V,a}$, isolated by the typed projectors Π_T (and helicity projectors when applicable) of Definition 9.350. Longitudinal contributions are gauge artefacts removed by BRST; the pole positions and residues are ξ-independent by Theorem 9.351.

(iii) Unbroken directions

If $m^2_{V,a} = 0$, the corresponding $V_{a\,\mu}$ is a massless internal vector (unbroken H-direction). Otherwise $m^2_{V,a} > 0$ and the mition is massive; the associated NG direction is eaten (equivalence theorem as in Corollary 9.352).

(iv) Separation from scalar/tensor Gaiamiti quanta

Mitions are *not* excitations of the scalar trace channel (the *gaiaton*, defined via the restricted Hessian $\mathcal{K}^{(\Phi)}_{\mathrm{A}}$; cf. Definition 9.259) nor of the TT tensor channel (the massless graviton; cf. Remark 9.260). They are quanta of the bundle connection coupled to Φ.

Justification (Mition). The spin–1 bundle–connection quantum in the Gaiamiti sector is called the *Mition*. The name derives from the Sanskrit root *mīti* ("measure"; cf. *jyāmīti*, "geometry"), reflecting the connection's role in providing *local measure*—i.e. covariant comparability—of Φ across the emergent manifold. While the Apeiron (umbrella for Gaiamiti quanta) includes the spin–0 radial scalar (the *Gaiaton*) and the spin–2 TT graviton, the *Mition* specifically denotes the spin–1 *internal* gauge quantum: the dynamical components A^I_μ of the operator bundle connection $\mathcal{A}_\mu$ once the sector is gauged (cf. Definition 9.255). Under spontaneous breaking by $\bar{\Phi}$, the covariant kinetic term produces a Brout–Englert–Higgs mass matrix $(M^2_\Phi)_{IJ} = g^2\, g_T(\rho_\Phi(T_I)\bar{\Phi}, \rho_\Phi(T_J)\bar{\Phi})$ (Proposition 9.257), so transverse Mitions appear as diagnostic realized reference configuration–physical poles at $k^2 \in \mathrm{spec}(M^2_\Phi)$ in the vector channel (gauge–parameter independent by Nielsen/BRST, Theorem 9.351). Conceptually, the Mition is the instrument that "measures" and transports Φ's internal data between fibers, providing the local structure complementary to the radial scalar Gaiaton (Apeiron umbrella) and coexisting with the massless TT graviton protected by relational Ward identities (Remark 9.260).

Gaiamiti (Φ) — what becomes massive, what is eaten, and how the vectors behave. When an internal group G acting on $\mathcal{E}_\Phi$ is gauged and the background $\bar{\Phi}$ breaks $G \to H$, the covariant kinetic term generates a Brout–Englert–Higgs (BEH) mass matrix for the gauge field A_μ^I:

$$(M_\Phi^2)_{IJ} = g^2\, g_T\Big(\rho_\Phi(T_I)\bar{\Phi},\ \rho_\Phi(T_J)\bar{\Phi}\Big) \quad (\textit{Definition 9.255and Proposition 9.257}).$$

What becomes massive? The internal vectors along the *broken* generators—i.e. the *mitions*—with masses given by the eigenvalues spec(M_Φ^2); vectors along unbroken H-directions remain *massless*. The would-be NG scalars of the broken orbit are *eaten* and reappear as *longitudinal* polarizations. On the certified plateau $\mathfrak{P}$, physical one-particle mition poles live in the *transverse* channel only (typed projectors Π_T, Π_L in Definition 9.350); longitudinal contributions are BRST-removed from $H^0(\hat{Q}_{\mathrm{BRST}})$. Both the pole positions m_V^2 and their residues are *gauge-parameter independent* (Nielsen/BRST; Theorem 9.351). At high energy, the typed *Equivalence Theorem* identifies amplitudes with external longitudinal mitions and those with the corresponding NG modes up to $\mathcal{O}(m_V/\sqrt{s})$ (Corollary 9.352). Importantly, internal gauging does *not* touch the relational/diffeomorphism sector: the TT *graviton* remains *massless* (Ward-protected; Remark 9.260), and the trace/radial scalar *gaiaton* stays a separate gapped mode read from the restricted Hessian $\mathcal{K}_{\mathrm{A}}^{(\Phi)}$ (Definition 9.259).

As interacting particles, mitions are *short-range carriers* with range $\sim m_V^{-1}$. They couple (i) to the *gaiaton* through the covariant derivative and scalar potential, producing the familiar VVh and $VVhh$ vertices; (ii) to *other vectors* via the Yang–Mills sector (triple/quartic gauge self-interactions); and (iii) to any *matter fields* charged under the same internal group (if present). If M_Φ^2 is non-diagonal, the gauge eigenstates *mix*; degeneracies produce *multiplets* under the unbroken subgroup H. Vectors along truly unbroken generators remain *massless* and mediate long-range forces unless further dynamics intervene. All of these readouts are made with typed/triadic meters on $\mathfrak{P}$ and inherit route-equivalence and BRST safety (Propositions 9.191, 9.207 and 9.223).

Quadratic Lagrangian matrix (Gaiamiti; field order $A_T,\ A_L,\ \mathrm{NG},\ \mathrm{A}$). Expand to quadratic order around $\bar{\Phi}$ and go to momentum space. After adding the Gaiamiti R_ξ gauge fixing $\mathcal{L}_{\mathrm{GF}}$ of Definition 9.261, the A_L–NG mixing cancels. Working in a single broken mass–eigenchannel (after diagonalizing $(M_\Phi^2)_{IJ}$ from Proposition 9.257), define the basis

$$\Psi := (A_T,\ A_L,\ \pi_{\mathrm{NG}},\ h_{\mathrm{A}}),$$

with A_T (*transverse* mition), A_L (*longitudinal* gauge mode), π_{NG} (would–be NG scalar), and h_{A} the *gaiaton* (radial Apeiron scalar; Definition 9.259). The quadratic form is diagonal:

$$\mathcal{L}^{(2)} = \tfrac{1}{2}\,\Psi^\dagger\,\mathbb{Q}_\Phi(k)\,\Psi,$$

$$\mathbb{Q}_\Phi(k) = \begin{pmatrix} k^2 - m_V^2 & 0 & 0 & 0 \\ 0 & \dfrac{k^2 - m_V^2}{\xi} & 0 & 0 \\ 0 & 0 & k^2 - \xi\, m_V^2 & 0 \\ 0 & 0 & 0 & k^2 - M_{\mathrm{A}}^2 \end{pmatrix},$$

where $\quad k^2 := \bar{\Sigma}^{\mu\nu} k_\mu k_\nu.$

Here $m_V^2 \in \mathrm{spec}(M_\Phi^2)$ is the mition mass in the chosen eigenchannel (Definition 9.255 and Proposition 9.257), and $M_{\mathrm{A}}^2 \in \mathrm{spec}(\mathcal{K}_{\mathrm{A}}^{(\Phi)})$ is the gaiaton (radial) mass (Definition 9.259). Poles: $A_T: \ k^2 = m_V^2$ (physical), $A_L: \ k^2 = m_V^2$ (gauge–artefactual numerator), $\pi_{\mathrm{NG}}: \ k^2 = \xi\, m_V^2$, $h_{\mathrm{A}}: \ k^2 = M_{\mathrm{A}}^2$. On $H^0(\hat{Q}_{\mathrm{BRST}})$ the physical poles are $k^2 = m_V^2$ and $k^2 = M_{\mathrm{A}}^2$; ξ–dependence cancels in S–matrix elements and matches NG amplitudes at high energy by Corollary 9.352. (The massless TT graviton lies in a separate TT block with a protected pole at $k^2 = 0$, unaffected by internal gauging; Remark 9.260.)

Propagators and physical content (Gaiamiti). With the typed vector projectors $\Pi_T^{\mu\nu}(k)$ and $\Pi_L^{\mu\nu}(k)$ (Definition 9.350), the mition propagator is

$$D_{\mu\nu}(k) = \frac{-i}{k^2 - m_V^2}\,\Pi_{T\,\mu\nu} \ + \ \frac{-i\,\xi}{k^2 - \xi\, m_V^2}\,\Pi_{L\,\mu\nu}.$$

The NG propagator is $D_{\mathrm{NG}}(k) = \dfrac{i}{k^2 - \xi\, m_V^2}$, and the gaiaton propagator is $D_{\mathrm{A}}(k) = \dfrac{i}{k^2 - M_{\mathrm{A}}^2}$. Gauge–parameter independence of the physical residues (Higgs/gaiaton and transverse mition) follows from Theorem 9.351; the high–energy identification of longitudinal–mition and NG amplitudes holds by the typed equivalence theorem (Corollary 9.352). All statements are made on the diagnostic realized reference configuration $\mathfrak{P}$ with typed/triadic meters, inheriting route equivalence and BRST safety (Propositions 9.191, 9.207 and 9.223).

Definition 9.259 (Apeiron (trace–typed) radial scalar under gauging)**.** Restrict the potential Hessian to the typed radial line to obtain

$$\boxed{\mathcal{K}_{\mathrm{A}}^{(\Phi)} := \Pi_{\mathrm{A}}^{(\Phi)}\ \frac{\delta^2 \mathcal{V}_\Phi}{\delta\hat{\mathbf{\Phi}}\,\delta\hat{\mathbf{\Phi}}}\bigg|_{\bar{\Phi}}\ \Pi_{\mathrm{A}}^{(\Phi)}} \qquad \text{with} \quad M_{\mathrm{A}}^2 \in \mathrm{spec}\Big(\mathcal{K}_{\mathrm{A}}^{(\Phi)}\Big).$$

This is the Gaiamiti analogue of the Tantrakyma Higgs/radial operator (cf. Definition 9.250 and eq. (9.185)).

Remark 9.260 (Masslessness of the TT graviton)**.** The typed TT graviton remains massless on $H^0(\hat{Q}_{\mathrm{BRST}})$: the TT two–point function carries a Ward-protected pole at $k^2 = 0$ (cf. Section 9.17.2) and is certified by route-equivalence/Nielsen safety (Propositions 9.191 and 9.223). The trace scale m_A from $\mathcal{K}_{\mathrm{A}}^{(\Phi)}$ measures Apeiron stiffness and is not a graviton mass.

Definition 9.261 (R_ξ gauge fixing and BRST hygiene (internal vectors))**.** Choose

$$\mathcal{F}^I := \nabla^{(\mathrm{rel})\,\mu} A_\mu^I - \xi\, g\, g_T\Big(\delta\hat{\mathbf{\Phi}},\ \rho_\Phi(T^I)\bar{\Phi}\Big), \qquad \mathcal{L}_{\mathrm{GF}} = -\frac{1}{2\xi}\, G_{IJ}\, \mathcal{F}^I \mathcal{F}^J,$$

with standard ghost sector so BRST closes on the diagnostic realized reference configuration and physical poles (transverse vectors at $k^2 \in \mathrm{spec}(M^2)$, Apeiron radial at $k^2 = M_{\mathrm{A}}^2$) are gauge-parameter independent (Nielsen identities; cf. Proposition 9.191). Typed decisions remain route-equivalent within tolerances (Proposition 9.223).

Remark 9.262 (Quadrant placement and explicit breaking)**.** With spontaneous internal breaking in the Gaiamiti sector and no explicit spurions, the typed predicate gives QII (global SSB eaten by BEH; cf. Proposition 9.251). Small explicit breaking moves the NG cluster to a pseudo–NG band with GMOR scaling, yielding QIV (cf. Definitions 7.685 and 9.188 and Proposition 9.197). In all cases, the TT graviton's massless pole persists (Remark 9.260), and Loc/On stamps and ledger updates follow Definitions 9.190 and 9.198 and Corollary 9.218.

Spectrum (Gaiamiti sector).

(i) Spin-0 (Apeiron, massive) Radial/trace scalar of $\hat{\mathbf{\Phi}}$ defined by the restricted Hessian (cf. Definition 9.259): the typed radial direction is generated by the normalized $\widehat{\mathsf{R}}_\Phi$ built from the typed projector $\Pi^{[\chi]}_{\Phi\Phi}$ (cf. Definitions 9.186, 9.200 and 9.256),

$$\widehat{\mathsf{R}}_\Phi \propto \bar{\Phi}, \qquad \mathcal{K}_{\mathrm{A}}^{(\Phi)} = \Pi_{\mathrm{A}}^{(\Phi)} \frac{\delta^2 \mathcal{V}_\Phi}{\delta\hat{\mathbf{\Phi}}\,\delta\hat{\mathbf{\Phi}}}\Big|_{\bar{\Phi}} \Pi_{\mathrm{A}}^{(\Phi)}, \quad M_{\mathrm{A}}^2 \in \mathrm{spec}\Big(\mathcal{K}_{\mathrm{A}}^{(\Phi)}\Big),$$

with diagnostic realized reference configuration discipline and BRST safety as in Propositions 9.191, 9.207 and 9.223.

(ii) Spin-1 (internal gauge bosons; massive if gauged and SSB) For G *gauged* and broken by $\bar{\Phi}$, internal vectors acquire BEH masses (cf. Proposition 9.257 and Definition 9.252):

$$(M^2)_{IJ} = g^2\, g_T\Big(\rho_\Phi(T_I)\bar{\Phi},\ \rho_\Phi(T_J)\bar{\Phi}\Big),$$

giving transverse vector poles at $k^2 = m_V^2 \in \mathrm{spec}(M^2)$ in the typed vector/helicity channels (Definition 9.350). If the symmetry is gauged but *unbroken*, vectors remain massless; if *ungauged*, no internal vectors appear. Quadrant placement follows Propositions 9.251 and 9.254.

(iii) Spin-2 (graviton, massless TT) The typed TT mode keeps a protected pole at $k^2 = 0$ on $H^0(\hat{Q}_{\mathrm{BRST}})$ by the relational/diffeomorphism Ward structure (cf. Section 9.17.2 and Remark 9.260); the trace-channel scale m_A encodes Apeiron stiffness and is *not* a graviton mass.

Quadrants under internal gauging (exhaustiveness). The quadrant classification (Chs. 7–8) is defined via typed/triadic diagnostic realized reference configuration visibilities (law vs. state) and the canonical decision rule (Definitions 7.685 and 9.187), evaluated on $H^0(\hat{Q}_{\mathrm{BRST}})$ with route equivalence and Nielsen safety (Propositions 9.191, 9.207 and 9.223). Internal gauging modifies only the *realization* of SSB (NG $\to$ longitudinal vectors) and leaves the axes unchanged (Section 9.17.3 and Proposition 9.254); thus the four quadrants remain exhaustive:

(i) (QI) Law symmetric, state symmetric No NG/pNG cluster (Π_{NG} null in the typed channel; cf. Definition 9.249); in a gauged realization, internal vectors stay massless; the radial scalar may be gapped via $\mathcal{K}_{\mathrm{A}}^{(\Phi)}$ (Definition 9.259). Decision per Definition 7.685.

(ii) (QII) Law symmetric, state SSB SSB occurs; in a gauged realization, NG directions are eaten and internal vectors gain BEH masses (M^2) (Proposition 9.257 and Definition 9.252), while the radial (Higgs-like/Apeiron) scalar remains gapped by $\mathcal{K}_{\mathrm{A}}^{(\Phi)}$ (Definition 9.259). Typed predicate per Proposition 9.251.

(iii) (QIII) Irrelevant explicit breaking, state symmetric Law-side nonconservation heals with IR coarse-graining on $\mathfrak{P}$ (healing power law as in eq. (7.746)); flows to the QI limit independently of gauging. Decision per Definition 7.685.

(iv) (QIV) Relevant/marginal explicit breaking, state unique The law channel remains explicitly broken in the IR; typed sources do *not* heal (cf. Definition 9.188). If some broken global directions are left *ungauged*, a pNG scalar cluster appears with GMOR scaling (diagnostic realized reference configuration-level GMOR/NG diagnostics Proposition 9.197); if all broken directions are *gauged*, no physical NG/pNG scalars remain on H^0 (the would-be NG become longitudinal vectors via Definition 9.252 and Proposition 9.257).

Gauge fixing in R_ξ form and BRST closure keep both the physical poles (Apeiron radial and transverse vectors) and quadrant assignment gauge-parameter independent (Nielsen) and route-equivalent within tolerances (Definition 9.261 and Propositions 9.191 and 9.223).

9.17.5 The Composite Nature of the Goldstone Mode

The identification of the graviton as a massless Goldstone excitation (Section 9.17.2) correctly attributes its masslessness to spontaneous symmetry breaking (SSB) of a continuous symmetry of the Gaiamiti potential $\mathcal{V}_\Phi$, which guarantees flat directions along the vacuum manifold $\mathcal{M}_{\mathrm{vac}}$ [93, 94]. This does *not* imply a monolithic, structureless object: in the IQR framework the underlying Gaiamiti field $\hat{\mathbf{\Phi}}(x)$ is composite, and its BRST–physical TT Goldstone excitation inherits a *triadic* internal structure tied to the $\{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$ channels on the diagnostic realized reference configuration.

Formal supplement (diagnostic realized reference configuration, typing, and triads). All statements below hold on a certified plateau $\mathfrak{P}$ with band tags $P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}^{(\ell)}$ and BRST projection to $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$; the triadic projectors $\{\Pi_\alpha\}_{\alpha\in\{\mathrm{corr,vac,spec}\}}$ commute with the typed projectors $\Pi_S^{[\chi]}$ (Definitions 9.185 and 9.186 and Proposition 9.202).

Definition 9.263 (Triadic decomposition of TT graviton components)**.** For the Gaiamiti sector $S = \Phi\Phi$ and type $[\chi] = \mathrm{TT}$, decompose the fluctuation and its metric image as

$$\delta\hat{\mathbf{\Phi}} = \sum_\alpha \Pi_\alpha\, \delta\hat{\mathbf{\Phi}}, \qquad \delta\hat{g}_{\mu\nu}(\delta\hat{\mathbf{\Phi}}) = \sum_\alpha \Pi_\alpha\, \delta\hat{g}_{\mu\nu}(\delta\hat{\mathbf{\Phi}}),$$

and define the *triadic TT graviton components* (on $\mathcal{H}_{\mathrm{phys}}$) by

$$\hat{h}_{\mu\nu}^{(\alpha)} \;:=\; \Pi_\alpha\, \Pi_{\Phi\Phi}^{[\mathrm{TT}]} \Big[\, \delta\hat{g}_{\mu\nu}\big(\delta\Phi_{\mathrm{tang}}\big) \,\Big], \qquad \delta\Phi_{\mathrm{tang}} \in T_{\bar{\Phi}}\mathcal{M}_{\mathrm{vac}} \;\; \text{(SSB tangent)}.$$

The (unnormalized) composite TT graviton is $\hat{h}_{\mu\nu} := \sum_\alpha \hat{h}_{\mu\nu}^{(\alpha)}$.

Proposition 9.264 (Orthogonality, completeness, and typed NG content)**.** *In the SSB regime with* $\mathrm{Quad}^{[\Phi\Phi,\mathrm{TT}]} = \mathrm{QII}$ *(Definition 9.189), on* $\mathcal{H}_{\mathrm{phys}}$*:*

(i) Triadic orthogonality/completeness

$\Pi_\alpha \Pi_\beta = \delta_{\alpha\beta}\Pi_\alpha$, $\sum_\alpha \Pi_\alpha = \mathbb{1}$*; hence* $\langle \hat{h}^{(\alpha)}, \hat{h}^{(\beta)} \rangle = 0$ *for* $\alpha \neq \beta$ *and* $\hat{h} = \sum_\alpha \hat{h}^{(\alpha)}$.

(ii) Typed NG poles

Each typed–triadic TT readout

$$\Pi_\alpha \Pi^{[\mathrm{TT}]}_{\Phi\Phi}\, G^{\mathrm{R}}_{\Phi\Phi,\mathrm{phys}}(k)\, \Pi^{[\mathrm{TT}]}_{\Phi\Phi} \Pi_\alpha$$

exhibits an exact NG pole at $k^2 = 0$ *with nonnegative residue (Proposition 9.197); at least one residue is strictly positive in QII.*

(iii) Nielsen/BRST safety

Gauge–parameter and BRST–exact deformations leave the above invariant (Propositions 9.191 and 9.207).

Definition 9.265 (Residue weights and normalized composite graviton)**.** Let the nonnegative NG residues on $\mathfrak{P}$ be

$$\mathcal{R}_\alpha \;:=\; \mathrm{Res}_{k^2=0}\Big[\Pi_\alpha \Pi^{[\mathrm{TT}]}_{\Phi\Phi}\, G^{\mathrm{R}}_{\Phi\Phi,\mathrm{phys}}(k)\, \Pi^{[\mathrm{TT}]}_{\Phi\Phi} \Pi_\alpha\Big].$$

Define $c_\alpha := \mathcal{R}_\alpha \big/ \sqrt{\sum_\beta \mathcal{R}_\beta^2}$ (so $\sum_\alpha c_\alpha^2 = 1$). The *normalized composite TT graviton* is

$$\hat{h}^{\mathrm{phys}}_{\mu\nu} \;:=\; \sum_\alpha c_\alpha\, \hat{h}^{(\alpha)}_{\mu\nu},$$

canonically fixed by plateau residues and independent of the order of typed/triadic insertions (Proposition 9.207).

Corollary 9.266 (Channel semantics and operational identification)**.** *In QII for* $(\Phi\Phi, \mathrm{TT})$*, the* TT Goldstone multiplet *is realized; with a* $\bar{\Sigma}$*–admissible spatial splitting, this resolves into the two physical helicities* $\lambda = \pm 2$ *(Definition 9.200 and Proposition 9.243). The components* $\hat{h}^{(\alpha)}$ *encode correlational/vacuum/spectral imprints, and the weights* $\{c_\alpha\}$ *are fixed by the NG residues on* $\mathfrak{P}$*. A* spectral–only parasymmetric protocol *can reconstruct spin/helicity content and diagnose breaking without the Ward ledger, while correlational and vacuum channels carry complementary information; all are read on the same diagnostic realized reference configuration with identical tags (Definition 9.185).*

Remark 9.267 (Structural consistency and typed/triadic compatibility)**.** Because $\sum_\alpha \Pi_\alpha = \mathbb{1}$ and $[\Pi^{[\chi]}_{\Phi\Phi}, \Pi_\alpha] = 0$ (Proposition 9.202), the Goldstone tangent $\delta\Phi_{\mathrm{tang}} \in T_{\bar{\Phi}}\mathcal{M}_{\mathrm{vac}}$ decomposes consistently as $\delta\Phi_{\mathrm{tang}} = \sum_\alpha \Pi_\alpha \delta\Phi_{\mathrm{tang}}$, and so does the TT graviton operator (Definition 9.263). This formalizes the "composite Goldstone" narrative without introducing extra physical degrees of freedom; it is a *decomposition in channels* of a single TT Goldstone multiplet, not an enlargement of the state space.

The Principle of Structural Consistency

The Gaiamiti kinetics couple $\hat{\mathbf{\Phi}}$ to the chronogeometric tensor $\bar{\Sigma}_{\mu\nu}(x)$, and $\bar{\Sigma}$ itself is not monolithic but carries a triadic decomposition (correlational, vacuum, spectral). Structural consistency then demands that the *same* channel structure be visible in the linear response of the source field: a single, undifferentiated fluctuation could not consistently interface with a tri-faceted geometry while preserving gauge/BRST consistency and causality.

On the DRCC (Definition 9.185) $\mathfrak{P}$, let $\{\Pi_\alpha\}_{\alpha\in\{\mathrm{corr,vac,spec}\}}$ be the triadic projectors and $\Pi_S^{[\chi]}$ the typed projectors ($\chi \in \{\mathrm{Tr}, \mathrm{TT}\}$) with $[\Pi_S^{[\chi]}, \Pi_\alpha] = 0$ (Definition 9.186 and Proposition 9.202). Then every BRST–physical fluctuation admits the orthogonal decomposition

$$\delta\hat{\mathbf{\Phi}}(x) = \sum_\alpha \Pi_\alpha\, \delta\hat{\mathbf{\Phi}}(x), \qquad \langle \Pi_\alpha \delta\hat{\mathbf{\Phi}}, \Pi_\beta \delta\hat{\mathbf{\Phi}}\rangle = 0 \quad (\alpha \neq \beta),$$

and the induced metric fluctuation obeys the same split,

$$\delta\hat{g}_{\mu\nu}\Big(\delta\hat{\mathbf{\Phi}}\Big) = \sum_\alpha \Pi_\alpha\, \delta\hat{g}_{\mu\nu}\Big(\delta\hat{\mathbf{\Phi}}\Big),$$

so that typed and triadic filtrations are compatible in the sense of $\Pi_S^{[\chi]} \sum_\alpha \Pi_\alpha = \sum_\alpha \Pi_\alpha \Pi_S^{[\chi]}$. This is the precise content of "structural consistency".

The Emergence of a Composite Graviton

Narrative. In the SSB regime, massless excitations follow *Goldstone's theorem* [93, 94]: they propagate along the flat directions of $\mathcal{M}_{\mathrm{vac}}$. In our typed framework, the graviton arises in the TT channel as a Goldstone excitation (massless spin–2) of the Gaiamiti sector; its existence and masslessness are *dynamical* consequences of SSB, not assumptions (cf. the standard lore [91, 100, 97]). Because the fluctuation space is triadically resolved on $\mathfrak{P}$, the *single* TT Goldstone multiplet decomposes into channel components. This is a *decomposition in channels*, not an increase of physical degrees of freedom: the realized graviton field operator is a coherent superposition fixed by NG residues on the diagnostic realized reference configuration.

Typed/triadic construction. Restrict to the TT type in the Gaiamiti sector ($S = \Phi\Phi$). As in Definition 9.263, define the triadic TT components on $\mathcal{H}_{\mathrm{phys}}$ by

$$\hat{h}^{(\alpha)}_{\mu\nu}(x) := \Pi_\alpha\, \Pi^{[\mathrm{TT}]}_{\Phi\Phi}\Big[\ \delta\hat{g}_{\mu\nu}\Big(\delta\Phi_{\mathrm{tang}}(x)\Big)\ \Big], \qquad \delta\Phi_{\mathrm{tang}}(x) \in T_{\bar{\Phi}}\mathcal{M}_{\mathrm{vac}},$$

and set $\hat{h}_{\mu\nu} := \sum_\alpha \hat{h}^{(\alpha)}_{\mu\nu}$. On a certified plateau, let

$$\mathcal{R}_\alpha := \mathrm{Res}_{k^2=0}\Big(\Pi_\alpha \Pi^{[\mathrm{TT}]}_{\Phi\Phi}\, G^{\mathrm{R}}_{\Phi\Phi,\mathrm{phys}}(k)\, \Pi^{[\mathrm{TT}]}_{\Phi\Phi}\Pi_\alpha\Big)\ \geq 0,$$

be the NG residues (Proposition 9.264). With normalized weights $c_\alpha := \mathcal{R}_\alpha/\sqrt{\sum_\beta \mathcal{R}_\beta^2}$ one obtains the *normalized composite graviton* (Definition 9.265)

$$\boxed{\hat{h}^{\mathrm{phys}}_{\mu\nu}(x) = \sum_{\alpha\in\{\mathrm{corr,vac,spec}\}} c_\alpha\, \hat{h}^{(\alpha)}_{\mu\nu}(x), \quad \sum_\alpha c_\alpha^2 = 1,}$$

whose definition is independent of the order of typed/triadic insertions (Proposition 9.207) and Nielsen/BRST–safe on $\mathfrak{P}$.

Operational semantics.

- **Spectral component** ($\alpha = \text{spec}$): Dominates conventional wave–detection protocols; reproduces the familiar propagating gravitational radiation sector and is often the most visible witness of the Goldstone mechanism in practice.
- **Correlational component** ($\alpha = \text{corr}$): Encodes entanglement– and coherence–induced geometric response (off–diagonal correlator structure); probed by parasymmetric protocols sensitive to information–geometric channels.
- **Vacuum component** ($\alpha = \text{vac}$): Captures moduli/zero–point geometry associated with the vacuum bundle; affects background–response and slow deformations of the diagnostic realized reference configuration.

All three are read on the same diagnostic realized reference configuration with identical tags (Definition 9.185); together they constitute the *TT Goldstone multiplet*. Their weights $\{c_\alpha\}$ are *not* arbitrary model choices but are fixed by the BRST–physical NG residues on the certified plateau, ensuring a structurally consistent, composite Goldstone description that remains within a single physical state space.

An Analogy: The Emergence of "White" Light

A helpful analogy is the phenomenon of white light. One may describe the *mechanism* of broadband light emission from a hot object via black–body radiation; this single process produces "light." When analyzed through a prism, however, its *composition* is revealed as a superposition of a continuous spectrum of colors.

IQR reading of the analogy.

(i) Spontaneous symmetry breaking (SSB) Is the single, unified mechanism that guarantees a massless, spin–2 excitation — the graviton — via the standard Goldstone logic [93, 94] (cf. Section 9.17.2).

(ii) Triadic decomposition as prism The triadic decomposition of the geometry/field (correlational, vacuum, spectral) is the "prism" that reveals the graviton's internal composition: a coherent superposition of three TT Goldstone components aligned with the triadic sectors (cf. Section 9.17.5).

Thus, *graviton* is the singular operational name for the massless spin–2 Goldstone excitation of the Gaiamiti sector, while the IQR framework predicts that this excitation is *composite*: three co–propagating, TT–typed, NG contributions with sector–set weights fixed by their NG residues (see Definition 9.265). This mirrors familiar emergent phenomena where a single long–wavelength quasiparticle encodes rich microstructural content (e.g. phonons in solids) [390, 391, 459, 460]. The compositeness is falsifiable: on a certified plateau $\mathfrak{P}$, the triadic TT readouts share an exact NG pole at $k^2{=}0$ with nonnegative residues, and at least one strictly positive, while the trace/radial channel remains gapped (QI) (cf. Proposition 9.197).

9.17.6 The Emergence of the Massless Graviton via SSB

Overview. How can the fundamental geometric field carry an intrinsic energy scale m_A while its observable force carrier—the graviton—remains massless? In the IQR framework the answer is *spontaneous symmetry breaking* (SSB): the Lagrangian is symmetric, the realized vacuum is not, and the ensuing Goldstone logic protects a long–range, massless mode in the appropriate typed channel [93, 94, 91, 100].

1. The Principle of Spontaneously Broken Geometric Symmetry Assume the Gaiamiti action, and in particular its potential $\mathcal{V}_\Phi(\hat{\mathbf{\Phi}})$, is invariant under a continuous internal symmetry $\mathcal{G}$ acting on the vacuum manifold $\mathcal{M}_{\text{vac}}$ (Definition 6.61). The realized background $\bar{\Phi}(x)$ may preserve only a subgroup $\mathcal{H} \subset \mathcal{G}$, thereby breaking $\mathcal{G} \to \mathcal{H}$. By **Goldstone's theorem** [93, 94], each broken generator yields a massless mode propagating along a flat direction of $\mathcal{M}_{\text{vac}}$.

In our typed/diagnostic realized reference configuration setting, this principle is applied to the geometric sector: the observable, massless spin–2 *graviton* is realized as a TT–typed Goldstone excitation of the Gaiamiti field on $H^0(\hat{Q}_{\text{BRST}})$. Its masslessness is thus *dynamically protected* by SSB (not assumed), mirroring familiar mechanisms in gauge theories (cf. Higgs/BEH and electroweak symmetry breaking) [97, 91].

2. The Graviton as a Composite Excitation of the Apeiron Field The graviton is a specific fluctuation of the underlying *Apeiron* field, i.e. the quantum of $\delta\hat{\mathbf{\Phi}}(x)$ (Section 9.17.1). Structural consistency (Section 9.17.5) requires that fluctuations be resolved by the triadic projectors $\{\Pi_{\text{corr}}, \Pi_{\text{vac}}, \Pi_{\text{spec}}\}$; hence

$$\delta\hat{\mathbf{\Phi}}(x) = \delta\hat{\mathbf{\Phi}}^{(\text{corr})}(x) + \delta\hat{\mathbf{\Phi}}^{(\text{vac})}(x) + \delta\hat{\mathbf{\Phi}}^{(\text{spec})}(x).$$

The Goldstone mode guaranteed by SSB is an excitation *of this composite field*. Consequently, the realized massless graviton inherits a triadic composition: in the TT type it is a coherent superposition of three channel components with weights fixed by NG residues on the certified plateau (Section 9.17.5):

$$\hat{h}_{\mu\nu}(x) = \hat{h}^{(\text{corr})}_{\mu\nu}(x) + \hat{h}^{(\text{vac})}_{\mu\nu}(x) + \hat{h}^{(\text{spec})}_{\mu\nu}(x), \qquad \hat{h}^{\text{phys}}_{\mu\nu}(x) = \sum_\alpha c_\alpha\, \hat{h}^{(\alpha)}_{\mu\nu}(x),$$

with $\sum_\alpha c_\alpha^2 = 1$ and c_α determined by the residues of the TT NG pole in each triadic channel (diagnostic realized reference configuration–certified; Nielsen/BRST–safe).

Correlators and propagators. Propagation of geometric fluctuations is encoded in the connected two–point function of the Gaiamiti sector, $\hat{G}_{\Phi\Phi}(x, x')$ (eq. (10.80)). The complete geometric two–point structure is organized in the matrix propagator $\mathbf{G}^{(2)}_\Sigma(x, x')$ (Definition 10.182). The graviton propagator is the TT–typed, spin–2 component,

$$G^{\text{grav}}_{\mu\nu\rho\sigma}(x, x') = \langle \Omega \,|\, T\{\hat{h}_{\mu\nu}(x)\, \hat{h}_{\rho\sigma}(x')\} \,|\, \Omega \rangle,$$

and—by the Goldstone mechanism—exhibits a long–range massless pole at $k^2 = 0$ with positive residue on H^0 in the SSB–only regime. Thus, even though the Apeiron sector carries an intrinsic scale m_A, there is *no* restoring force (and hence no mass term) along the TT Goldstone directions; gravity remains long–range.

3. Physical Interpretation and Resolution of the Mass Paradox This synthesis separates the constituent from the emergent excitation. The parameter m_A is not the graviton mass; it is an intrinsic *stiffness* scale of the relational vacuum encoded by $\mathcal{V}_\Phi$ (cf. Section 9.17.1). Large m_A means a geometry that is costly to deform in the radial/trace channel, while the TT channel retains an SSB–protected, massless Goldstone band. The apparent paradox—"intrinsic scale vs. massless force carrier''—is resolved exactly as in other emergent settings: long–wavelength quanta can behave very differently from their building blocks. Standard examples include phonons in crystals, where massive atoms yield nearly massless collective modes that carry sound and heat [390, 391, 459].

Emergence: Whole vs. Parts (operational view)

(i) Apeiron. The Apeiron is the constituent quantum of $\delta\hat{\mathbf{\Phi}}$, equipped with an intrinsic scale m_A; by structural consistency it lives in triadic channels (correlational, vacuum, spectral).

(ii) Graviton. The graviton is the *collective* massless TT Goldstone excitation—protected by SSB, realized on H^0, and *composite* across the triad with diagnostic realized reference configuration–fixed weights (Section 9.17.5).
Read together, these points show why geometry may possess an intrinsic energy scale while gravity remains long–range: the observable graviton is a massless, TT–typed Goldstone excitation of a triadically structured field, not a direct readout of the constituent stiffness. The framework's predictions (exact TT NG pole; gapped trace/Apeiron channel; triadic residue weights) are concrete, falsifiable features of the diagnostic realized reference configuration.

9.17.7 Euler–Lagrange Equation for the Gaiamiti Field

Having established the action functional for the Gaiamiti field, we now derive its dynamical law. By the Principle of Stationary Action ($\delta S_\Phi = 0$), we obtain an *operator–valued* equation of motion that governs how relational geometry evolves under quantum self–interaction. This is not a wave equation on a fixed background: it is the evolution equation *of the background itself*.

Diagnostic realized reference configuration/typed refinement (realized regime). Work on the diagnostic realized reference configuration $\mathfrak{P}$ (Definition 9.185) and vary the Gaiamiti action $S_\Phi[\hat{\mathbf{\Phi}}; \bar{\Sigma}]$ (cf. Definition 9.235 and §6.5). Project to the BRST–physical sector and impose the Euler–Lagrange (EL) equation as

$$\mathsf{P}_{\text{phys}} \frac{\delta S_\Phi}{\delta \hat{\mathbf{\Phi}}} = 0 \quad \text{on } \mathcal{H}_{\text{phys}},$$

with all limits taken inside the certified plateau $\mathfrak{P}$ (or $\mathfrak{P}_\partial$) per Chapter 8.

Typed EL split. Insert the BRST–closed typed projectors $\Pi^{[\chi]}_{\Phi\Phi}$ (Definitions 9.186, 9.195 and 9.200) and decompose about the realized vacuum $\bar{\Phi}$ (cf. eq. (10.87)):

$$\delta\hat{\mathbf{\Phi}} = \delta\hat{\mathbf{\Phi}}^{[\mathrm{TT}]} \oplus \delta\hat{\mathbf{\Phi}}^{[\mathrm{Tr}]}, \qquad \delta\hat{\mathbf{\Phi}}^{[\chi]} := \Pi^{[\chi]}_{\Phi\Phi}\, \delta\hat{\mathbf{\Phi}}.$$

Then, on $\mathcal{H}_{\mathrm{phys}}$,

$$\begin{aligned} &\text{(trace / radial)} \quad \left(\bar{\nabla}^{\mu}\bar{\nabla}_{\mu} - \mathcal{M}_{[\mathrm{Tr}]}(x)\right) \delta\hat{\mathbf{\Phi}}^{[\mathrm{Tr}]}(x) = 0, \\ &\text{(TT / tangential)} \quad \Pi^{[\mathrm{TT}]}_{\Phi\Phi}\, \mathcal{L}_{\mathrm{TT}}[\bar{\Sigma}] \left(\delta\hat{g}(\delta\Phi_{\mathrm{tang}})\right) = 0, \end{aligned}$$

where $\bar{\nabla}$ is the Relational Levi–Civita connection of $\bar{\Sigma}$ (Definition 6.7), $\mathcal{M}_{[\mathrm{Tr}]}$ is the typed mass operator from the potential Hessian at $\bar{\Phi}$ (Definition 9.246), and $\mathcal{L}_{\mathrm{TT}}[\bar{\Sigma}]$ is the (Lichnerowicz–type) linearized geometric operator acting on the TT metric deformation induced by the Goldstone tangent $\delta\Phi_{\mathrm{tang}} \in T_{\bar{\Phi}}\mathcal{M}_{\mathrm{vac}}$ (§9.17.2). The TT equation encodes the null curvature along $\mathcal{M}_{\mathrm{vac}}$ (QII NG pole; Proposition 9.197); the trace equation shows a gap set by the Apeiron scale (QI).

Boundary terms and safety. Natural boundary conditions on $\mathfrak{P}_{\partial}$ follow from the same variation with the bulk/edge overlay (Definition 9.214); the resulting typed EL equations (bulk/edge) are Nielsen–flat and BRST–safe by Propositions 9.191 and 9.207. By Proposition 9.223, the solutions' typed spectra agree with the SynTh/correlational route within Chapter 8 tolerances.

Definition 9.268 (Euler–Lagrange Equation for $\hat{\mathbf{\Phi}}(x)$)**.** Let $\hat{\mathbf{\Phi}}(x)$ be the dynamical Gaiamiti field with action $S_{\Phi}[\hat{\mathbf{\Phi}}]$ (Definition 6.174). Using the shorthand measure $\sqrt{-\det\left(\bar{\Sigma}\right)}$ for the relational volume (as noted in Remark 6.7's discussion of the volume element), the operator–valued EL equation can be written in the manifestly covariant form:

$$\boxed{\frac{1}{\sqrt{-\det\left(\bar{\Sigma}\right)}}\, \nabla_{\mu}\left(\sqrt{-\det\left(\bar{\Sigma}\right)}\, \bar{\Sigma}^{\mu\nu}\, \nabla_{\nu}\hat{\mathbf{\Phi}}(x)\right) = \frac{\delta\mathcal{V}_{\Phi}}{\delta\hat{\mathbf{\Phi}}^{\dagger}(x)}.} \tag{9.107}$$

Expanding the covariant divergence gives the equivalent local form

$$\boxed{\bar{\Sigma}^{\mu\nu}(x)\, \nabla_{\mu}\nabla_{\nu}\hat{\mathbf{\Phi}}(x) + \left(\nabla_{\mu}\bar{\Sigma}^{\mu\nu}(x)\right) \nabla_{\nu}\hat{\mathbf{\Phi}}(x) = \frac{\delta\mathcal{V}_{\Phi}}{\delta\hat{\mathbf{\Phi}}^{\dagger}(x)}.} \tag{9.108}$$

By metric compatibility of the Relational Levi–Civita connection and the fact that the operator–bundle connection acts only on internal (fiber) indices, one has

$$\nabla_{\mu}\bar{\Sigma}^{\mu\nu} = 0,$$

so that (9.108) reduces on the diagnostic realized reference configuration to

$$\bar{\Sigma}^{\mu\nu}\, \nabla_{\mu}\nabla_{\nu}\hat{\mathbf{\Phi}} = \frac{\delta\mathcal{V}_{\Phi}}{\delta\hat{\mathbf{\Phi}}^{\dagger}},$$

which is identical to (9.107). Here:

(i) Total covariant derivative

The total covariant derivative $\nabla_\mu = \nabla_\mu^{(\mathrm{rel})} + D_\mu$ splits into the Relational Levi–Civita part acting on spacetime indices and the operator–bundle connection D_μ acting on internal operator–space indices (Section 6.6).

(ii) Source term

The source term $\delta\mathcal{V}_\Phi/\delta\hat{\mathbf{\Phi}}^\dagger$ is the functional derivative of the potential $\mathcal{V}_\Phi$, which is built from the tangent–space inner product g_T (Section 6.1.24).

Interpretation of the Equation. This equation is a background–independent, operator–valued generalization of the Klein–Gordon equation [461]. The left–hand side is a generalized covariant d'Alembertian built from the chronogeometric tensor $\bar{\Sigma}_{\mu\nu}(x)$; it governs the *propagation of relational geometric deformations*. The right–hand side is a *source term* that encodes self–interaction and the vacuum–selection dynamics through $\delta\mathcal{V}_\Phi/\delta\hat{\mathbf{\Phi}}^\dagger$. Using metric compatibility of the relational Levi–Civita connection, $\nabla_\mu\bar{\Sigma}^{\mu\nu} = 0$, the expanded form reduces on the diagnostic realized reference configuration to

$$\bar{\Sigma}^{\mu\nu}\nabla_\mu\nabla_\nu\hat{\mathbf{\Phi}}(x) = \frac{\delta\mathcal{V}_\Phi}{\delta\hat{\mathbf{\Phi}}^\dagger(x)}\,,$$

so the *same* operator governs both propagation and self–consistent back–reaction: this is not a wave on a fixed background but the evolution equation *of* the backgrounded geometry itself.

Expectation–Level Equation. Taking the expectation value in the quantum holo–state $\left|\overset{\circ}{\mathbf{\Psi}}\right\rangle$ yields the semiclassical equation for the mean field $\varphi(x) := \langle\hat{\mathbf{\Phi}}(x)\rangle$:

$$\left\langle \frac{1}{\sqrt{-\det\bar{\Sigma}}}\,\nabla_\mu\Big(\sqrt{-\det\bar{\Sigma}}\,\bar{\Sigma}^{\mu\nu}\nabla_\nu\hat{\mathbf{\Phi}}(x)\Big)\right\rangle = \left\langle\frac{\delta\mathcal{V}_\Phi}{\delta\hat{\mathbf{\Phi}}^\dagger(x)}\right\rangle.$$

Because the RHS is nonlinear in $\hat{\mathbf{\Phi}}$, expectations and variations do not commute in general. On a certified plateau one may package fluctuation effects into an *effective* potential, $\langle\delta\mathcal{V}_\Phi/\delta\hat{\mathbf{\Phi}}^\dagger\rangle = \delta\mathcal{V}_{\mathrm{eff}}/\delta\varphi^\dagger$, so that the expectation–level dynamics take the standard curved–spacetime semiclassical form [10, 245]:

$$\bar{\Sigma}^{\mu\nu}\nabla_\mu\nabla_\nu\varphi(x) = \frac{\delta\mathcal{V}_{\mathrm{eff}}}{\delta\varphi^\dagger(x)}\,, \qquad (\text{BRST/Nielsen–safe on } \mathcal{H}_{\mathrm{phys}}).$$

The Euler–Lagrange Equation as the Fundamental Dynamical Principle Governing the Apeiron (refined)

The EL equation for the Gaiamiti sector is the relativistic wave equation for the *Apeiron*—the quantum of relational geometry—prescribing both its propagation and its self–interaction on the emergent chronogeometric manifold.

1. Kinetic component: propagative dynamics of the Apeiron. The left–hand side, $\bar{\Sigma}^{\mu\nu}\nabla_\mu\nabla_\nu\hat{\mathbf{\Phi}}$, is a manifestly covariant d'Alembertian with the total derivative $\nabla_\mu = \nabla_\mu^{(\mathrm{rel})} + D_\mu$ (relational Levi–Civita on spacetime indices plus operator–bundle connection on internal indices). With $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (Sections 5.2.6 and 5.2.8), this term describes not a field *on* a fixed stage, but the propagation of the geometric structure *itself*. The field $\hat{\mathbf{\Phi}}$ is at once dynamical and geometric.

2. Potential component: intrinsic scale and self–interaction of the Apeiron. The source term is the potential gradient $\delta\mathcal{V}_\Phi/\delta\hat{\mathbf{\Phi}}^\dagger$ (Section 9.17). For the unbroken–phase normalization used elsewhere,

$$\mathcal{V}_\Phi(\hat{\mathbf{\Phi}}) = \tfrac{1}{2}\, m_A^2\, g_T(\hat{\mathbf{\Phi}},\hat{\mathbf{\Phi}}) + \tfrac{\lambda_A}{4!}\,[g_T(\hat{\mathbf{\Phi}},\hat{\mathbf{\Phi}})]^2 + \cdots,$$

one finds near the symmetric background (treating $\hat{\mathbf{\Phi}}, \hat{\mathbf{\Phi}}^\dagger$ as independent)

$$\frac{\delta\mathcal{V}_\Phi{}^\dagger}{\delta\hat{\mathbf{\Phi}}} = \tfrac{1}{2}\, m_A^2\, \hat{\mathbf{\Phi}} + \tfrac{\lambda_A}{12}\, S\, \hat{\mathbf{\Phi}} + \cdots, \qquad S := g_T(\hat{\mathbf{\Phi}},\hat{\mathbf{\Phi}}),$$

so the leading restoring tendency is set by the intrinsic *Apeiron scale* m_A. This scale is not a "graviton mass"; it quantifies the stiffness of the radial/trace channel:

(i) Energetic cost It fixes the energetic cost of deforming the relational vacuum.

(ii) Curvature rigidity Larger m_A implies greater curvature rigidity (smaller susceptibility).

(iii) Correlation length The characteristic correlation length scales as $\xi_A \sim m_A^{-1}$ (up to the conventional normalization just noted).

By contrast, the TT (tangential) directions are flat in the SSB–only regime, so no mass term can arise there.

Synthesis: the unified principle of propagation. The EL equation,

$$\bar{\Sigma}^{\mu\nu}\nabla_\mu\nabla_\nu\hat{\mathbf{\Phi}} = \frac{\delta\mathcal{V}_\Phi{}^\dagger}{\delta\hat{\mathbf{\Phi}}},$$

unifies (i) the **kinematic** law of geometric propagation (generalized covariant d'Alembertian) with (ii) the **intrinsic energy/self–interaction** encoded by the potential gradient. In the typed split, the trace channel obeys a gapped KG–like law with mass operator $\mathcal{M}_{[\mathrm{Tr}]}$ (QI), whereas the TT channel satisfies a Lichnerowicz–type equation along Goldstone directions (QII) with an exact massless pole protected by SSB. Thus spacetime curvature and gravitational radiation emerge as *collective* solutions of a single operator equation, reconciling an intrinsic geometric scale m_A with a massless, long–range graviton consistent with observation.

Definition 9.269 (HS Euler–Lagrange equation for $\hat{\mathbf{\Phi}}$)**.** Let the HS Gaiamiti Lagrangian be

$$\mathcal{L}_\Phi^{\mathrm{HS}} = \mathrm{Tr}_{\mathrm{FD}}\Big\{\boldsymbol{g}^{\mathrm{Lor}\,\mu\nu}\, g_T^{\mathrm{HS}}\Big(\mathcal{D}_\mu^{\mathrm{HS}}\hat{\mathbf{\Phi}},\, \mathcal{D}_\nu^{\mathrm{HS}}\hat{\mathbf{\Phi}}\Big) - \mathcal{V}_\Phi^{\mathrm{HS}}(\hat{\mathbf{\Phi}})\Big\} \quad (\text{cf. } \textit{Section } 6.5.2 \textit{and Definition } 6.34).$$

Varying $\widehat{\boldsymbol{S}}_\Phi = \int d\boldsymbol{\mu}\, \mathcal{L}_\Phi^{\mathrm{HS}}$ with $\mathcal{D}_\mu^{\mathrm{HS}}$ from eq. (6.259) and $d\boldsymbol{\mu}$ from Definition 6.34, the HS Euler–Lagrange equation reads

$$\boxed{\mathcal{D}_\mu^{\mathrm{HS}}\Big(\boldsymbol{g}^{\mathrm{Lor}\,\mu\nu}\,\mathfrak{J}_\nu\Big) \;-\; \frac{\delta}{\delta\hat{\boldsymbol{\Phi}}}\,\mathrm{Tr}_{\mathrm{FD}}\Big\{\mathcal{V}_\Phi^{\mathrm{HS}}(\hat{\boldsymbol{\Phi}})\Big\} \;=\; 0,} \tag{9.109}$$

with conjugate HS current

$$\boxed{\mathfrak{J}_\nu := \frac{\delta}{\delta(\mathcal{D}_\nu^{\mathrm{HS}}\hat{\boldsymbol{\Phi}})}\,\mathrm{Tr}_{\mathrm{FD}}\Big\{\, g_T^{\mathrm{HS}}\Big(\mathcal{D}_\nu^{\mathrm{HS}}\hat{\boldsymbol{\Phi}},\,\cdot\,\Big)\Big\}.} \tag{9.110}$$

Here g_T^{HS} is the HS vacuum–tangent inner product (Definition 6.106), and $\mathcal{V}_\Phi^{\mathrm{HS}}$ is any real functional of $\mathcal{I}_\Phi^{\mathrm{HS}} = g_T^{\mathrm{HS}}(\hat{\boldsymbol{\Phi}},\hat{\boldsymbol{\Phi}})$ (Definitions 9.236 and 9.237).

Sketch. Use compactly supported variations and metric compatibility $\nabla_\lambda^{(\mathrm{rel})}\boldsymbol{g}^{\mathrm{Lor}\,\mu\nu} = 0$ (Levi–Civita; Definition 6.7), plus the divergence identity with $d\boldsymbol{\mu}$ (Definition 6.34) to integrate by parts. The potential variation is given by Lemma 9.240. Cyclicity of $\mathrm{Tr}_{\mathrm{FD}}$ (Definition 6.45 and eq. (6.99)) and the inner-derivation property of $\mathcal{D}_\mu$ (Remark 6.187) yield (9.109)–(9.110). □

Remark 9.270 (Tantrakyma status)**.** The HS Euler–Lagrange equation for the Tantrakyma field is given in Definition 9.321, built from g_C^{HS}, $\mathcal{D}_\mu^{\mathrm{HS}}$ (eq. (9.150)), and $\boldsymbol{g}^{\mathrm{Lor}}$ with the same invariant measure (Definition 6.34).

9.17.8 Noether Current for the Dynamical Gaiamiti Field

Assume the Gaiamiti action $S_\Phi[\hat{\boldsymbol{\Phi}}]$ is invariant under a continuous internal symmetry generated by a Hermitian $\hat{Q}$ that preserves the kinetic structures $(\bar{\Sigma}_{\mu\nu},\, g_T)$ and is compatible with the split covariant derivative $\nabla_\mu = \nabla_\mu^{(\mathrm{rel})} + (\partial_\mu + i[\mathcal{A}_\mu,\cdot])$ (cf. Section 6.12.2 and eq. (6.316)). Then the infinitesimal transformation

$$\delta\hat{\boldsymbol{\Phi}}(x) = i\,\epsilon(x)\,[\hat{Q},\hat{\boldsymbol{\Phi}}(x)], \qquad \delta\hat{\boldsymbol{\Phi}}^\dagger(x) = -i\,\epsilon(x)\,[\hat{Q},\hat{\boldsymbol{\Phi}}^\dagger(x)]$$

leads, by Noether's procedure, to a conserved current on shell.

Definition 9.271 (Noether current for the Gaiamiti sector)**.** Let $\mathcal{L}_\Phi$ be the operator-valued Lagrangian density of the Gaiamiti field (Definition 6.174 and eq. (6.316)). The associated operator-valued Noether current is

$$\boxed{\hat{J}^\mu_{(\Phi)}(x) := -i\,\mathrm{Tr}\Big\{\frac{\partial\mathcal{L}_\Phi}{\partial(\nabla_\mu\hat{\boldsymbol{\Phi}})}\,[\hat{Q},\hat{\boldsymbol{\Phi}}] - \frac{\partial\mathcal{L}_\Phi}{\partial(\nabla_\mu\hat{\boldsymbol{\Phi}}^\dagger)}\,[\hat{Q},\hat{\boldsymbol{\Phi}}^\dagger]\Big\},} \tag{9.111}$$

where the Fréchet derivatives are taken at fixed $\bar{\Sigma}$ and $\mathcal{A}_\mu$. Equivalently, one may write

$$\hat{J}^\mu_{(\Phi)} = -i\,\mathrm{Tr}\Big\{\frac{\partial\mathcal{L}_\Phi}{\partial(\nabla_\mu\hat{\boldsymbol{\Phi}})}[\hat{Q},\hat{\boldsymbol{\Phi}}]\Big\} + \mathrm{h.c.}$$

Specialization to the kinetic form. For

$$\mathcal{L}_\Phi = \mathrm{Tr}\Big\{\bar{\Sigma}^{\rho\nu}\, g_T(\nabla_\rho\hat{\mathbf{\Phi}}, \nabla_\nu\hat{\mathbf{\Phi}}) + \mathcal{V}_\Phi(\hat{\mathbf{\Phi}})\Big\} \quad \text{(cf. eq. (6.316))},$$

$$\frac{\partial\mathcal{L}_\Phi}{\partial(\nabla_\mu\hat{\mathbf{\Phi}})} = \bar{\Sigma}^{\mu\nu}\, g_T^\sharp\Big(\nabla_\nu\hat{\mathbf{\Phi}}, \cdot\Big),$$

so one may render (9.111) as

$$\hat{J}^\mu_{(\Phi)}(x) = -i\,\mathrm{Tr}\Big\{\bar{\Sigma}^{\mu\nu}\Big[g_T\Big(\nabla_\nu\hat{\mathbf{\Phi}}, [\hat{Q}, \hat{\mathbf{\Phi}}]\Big) - g_T\Big([\hat{Q}, \hat{\mathbf{\Phi}}^\dagger], \nabla_\nu\hat{\mathbf{\Phi}}^\dagger\Big)\Big]\Big\}.$$

Proposition 9.272 (Covariant conservation (diagnostic realized reference configuration/H^0))**.** *If $\hat{\mathbf{\Phi}}$ satisfies its Euler–Lagrange equation*

$$\frac{1}{\sqrt{-\det\bar{\Sigma}}}\,\nabla_\mu\Big(\sqrt{-\det\bar{\Sigma}}\,\bar{\Sigma}^{\mu\nu}\nabla_\nu\hat{\mathbf{\Phi}}\Big) = \frac{\delta\mathcal{V}_\Phi}{\delta\hat{\mathbf{\Phi}}^\dagger} \quad \textit{(cf. Definition 9.268)},$$

then the expectation-level current $\bar{J}^\mu_{(\Phi)} := \langle\overset{\circ}{\mathbf{\Psi}}|\hat{J}^\mu_{(\Phi)}|\overset{\circ}{\mathbf{\Psi}}\rangle$ is covariantly conserved on the certified plateau:

$$\boxed{\nabla^{(\mathrm{rel})}_\mu\, \bar{J}^\mu_{(\Phi)}(x) = 0 \qquad \textit{on } \mathcal{H}_{\mathrm{phys}}.} \tag{9.112}$$

Moreover, band-tagging and BRST improvements preserve the Ward identity: the BRST–improved, band-limited current obeys the scale-tagged Ward law stated in Definition 7.567, with exact conservation in QI/QII and adjoint sources only when explicit breaking spurions are present (QIV).

Proof. We work with the covariant density $\mathcal{L}_\Phi$ and the shorthand $S_\Phi = \int d^dx\,\sqrt{-\det\bar{\Sigma}}\,\mathcal{L}_\Phi$. Assume the internal symmetry generated by Hermitian $\hat{Q}$ is compatible with the split derivative, i.e. $[\nabla_\mu, \mathrm{ad}_{\hat{Q}}] = 0$ and it preserves the kinetic structures $(\bar{\Sigma}_{\mu\nu}, g_T)$. Treat $\hat{\mathbf{\Phi}}$ and $\hat{\mathbf{\Phi}}^\dagger$ as independent variables.

Step 1: Localized symmetry variation. Promote the global parameter to a smooth test function $\epsilon(x)$ (standard Noether trick) and set

$$\delta_\epsilon\hat{\mathbf{\Phi}} = i\,\epsilon(x)\,[\hat{Q}, \hat{\mathbf{\Phi}}], \qquad \delta_\epsilon\hat{\mathbf{\Phi}}^\dagger = -i\,\epsilon(x)\,[\hat{Q}, \hat{\mathbf{\Phi}}^\dagger].$$

By compatibility, $\delta_\epsilon(\nabla_\mu\hat{\mathbf{\Phi}}) = \nabla_\mu(\delta_\epsilon\hat{\mathbf{\Phi}})$, hence $\delta_\epsilon(\nabla_\mu\hat{\mathbf{\Phi}}) = i(\nabla_\mu\epsilon)\,[\hat{Q}, \hat{\mathbf{\Phi}}] + i\epsilon\,[\hat{Q}, \nabla_\mu\hat{\mathbf{\Phi}}]$, and similarly for $\hat{\mathbf{\Phi}}^\dagger$.

Step 2: Variation of the density and covariant integration by parts. Using the chain rule (including the $\hat{\mathbf{\Phi}}^\dagger$ sector) and suppressing explicit $\sqrt{-\det\bar{\Sigma}}$,

$$\begin{aligned}\delta_\epsilon\mathcal{L}_\Phi &= \mathrm{Tr}\Big\{\frac{\partial\mathcal{L}_\Phi}{\partial\hat{\mathbf{\Phi}}}\,\delta_\epsilon\hat{\mathbf{\Phi}} + \frac{\partial\mathcal{L}_\Phi}{\partial(\nabla_\mu\hat{\mathbf{\Phi}})}\,\delta_\epsilon(\nabla_\mu\hat{\mathbf{\Phi}})\Big\} + \mathrm{h.c.}\\ &= \mathrm{Tr}\Big\{\Big(\frac{\partial\mathcal{L}_\Phi}{\partial\hat{\mathbf{\Phi}}} - \nabla_\mu\frac{\partial\mathcal{L}_\Phi}{\partial(\nabla_\mu\hat{\mathbf{\Phi}})}\Big)\delta_\epsilon\hat{\mathbf{\Phi}}\Big\} + \nabla^{(\mathrm{rel})}_\mu\,\mathrm{Tr}\Big\{\frac{\partial\mathcal{L}_\Phi}{\partial(\nabla_\mu\hat{\mathbf{\Phi}})}\,\delta_\epsilon\hat{\mathbf{\Phi}}\Big\} + \mathrm{h.c.}\end{aligned}$$

Here we used metric compatibility of $\nabla^{(\mathrm{rel})}$ for the divergence and that the bundle connection acts only on fiber indices.

Step 3: Identify the current. Insert the explicit form of $\delta_\epsilon \hat{\mathbf{\Phi}}$ and collect the $\nabla_\mu \epsilon$ terms:

$$\delta_\epsilon \mathcal{L}_\Phi = -(\nabla_\mu^{(\mathrm{rel})} \epsilon)\, \hat{J}^\mu_{(\Phi)} + \epsilon\, \mathcal{E} \;+\; \mathrm{h.c.},$$

where

$$\hat{J}^\mu_{(\Phi)} := -i\,\mathrm{Tr}\Big\{ \frac{\partial \mathcal{L}_\Phi}{\partial(\nabla_\mu \hat{\mathbf{\Phi}})}\,[\hat{Q}, \hat{\mathbf{\Phi}}] - \frac{\partial \mathcal{L}_\Phi}{\partial(\nabla_\mu \hat{\mathbf{\Phi}}^\dagger)}\,[\hat{Q}, \hat{\mathbf{\Phi}}^\dagger] \Big\}$$

is exactly (9.111), and

$$\mathcal{E} := i\;\mathrm{Tr}\Big\{ \Big(\frac{\partial \mathcal{L}_\Phi}{\partial \hat{\mathbf{\Phi}}} - \nabla_\mu \frac{\partial \mathcal{L}_\Phi}{\partial(\nabla_\mu \hat{\mathbf{\Phi}})} \Big)[\hat{Q}, \hat{\mathbf{\Phi}}] \Big\} - \;i\;\mathrm{Tr}\Big\{ \Big(\frac{\partial \mathcal{L}_\Phi}{\partial \hat{\mathbf{\Phi}}}^{\dagger} - \nabla_\mu \frac{\partial \mathcal{L}_\Phi}{\partial(\nabla_\mu \hat{\mathbf{\Phi}}^\dagger)} \Big)[\hat{Q}, \hat{\mathbf{\Phi}}^\dagger] \Big\}.$$

Thus the variation of the action reads

$$\delta_\epsilon S_\Phi = \int d^d x\, \sqrt{-\det \bar{\Sigma}}\, \Big(-(\nabla_\mu^{(\mathrm{rel})} \epsilon)\, \hat{J}^\mu_{(\Phi)} + \epsilon\, \mathcal{E} + \mathrm{h.c.} \Big).$$

Step 4: Global invariance and on-shell conservation. For a *global* symmetry, $\epsilon = \mathrm{const}$ and invariance gives $\delta_\epsilon S_\Phi = 0$. Allowing $\epsilon(x)$ to have compact support and integrating by parts,

$$\delta_\epsilon S_\Phi = \int d^d x\, \sqrt{-\det \bar{\Sigma}}\, \epsilon(x) \Big(\nabla_\mu^{(\mathrm{rel})} \hat{J}^\mu_{(\Phi)} + \mathcal{E} + \mathrm{h.c.} \Big).$$

On shell, the Euler–Lagrange equations (Definition 9.268) annihilate $\mathcal{E} + \mathrm{h.c.}$. Since $\epsilon(x)$ is arbitrary, we obtain the operator identity $\nabla_\mu^{(\mathrm{rel})} \hat{J}^\mu_{(\Phi)} = 0$. Taking expectations in the Quantum Holo–State $\big|\mathring{\mathbf{\Psi}}\big\rangle$ yields

$$\nabla_\mu^{(\mathrm{rel})}\, \bar{J}^\mu_{(\Phi)}(x) = 0,$$

which is precisely (9.112). □

Diagnostic realized reference configuration and sector. Work on the instrumented diagnostic realized reference configuration (Definition 9.185) in the Gaiamiti sector $S = \Phi\Phi$. Let G act on $\hat{\mathbf{\Phi}}$ with generators T_A (Lie-algebra indices A), and let $\bar{\nabla}$ be the Levi–Civita derivative of $\bar{\Sigma}$ (Definition 6.7). All statements below are on $\mathcal{H}_{\mathrm{phys}}$ and within a certified plateau $\mathfrak{P}$ (or $\mathfrak{P}_\partial$) per Chapter 8.

Definition 9.273 (Dynamical Noether current on H^0 (Gaiamiti))**.** Consider the Gaiamiti Lagrangian density $\mathcal{L}_\Phi(\hat{\mathbf{\Phi}}, \nabla\hat{\mathbf{\Phi}}; \bar{\Sigma})$ entering S_Φ (cf. §6.5, Definition 9.235). For an infinitesimal *global* transformation $\delta_A \hat{\mathbf{\Phi}} = i\,\epsilon\, T_A \hat{\mathbf{\Phi}}$, define the operator Noether current

$$\hat{\mathcal{J}}^\mu_A \;:=\; \frac{\partial \mathcal{L}_\Phi}{\partial(\nabla_\mu \hat{\mathbf{\Phi}})} \big(i\, T_A \hat{\mathbf{\Phi}} \big) \;+\; \mathrm{h.c.}$$

and its BRST–physical projection $\mathcal{J}^\mu_A := \mathsf{P}_{\mathrm{phys}}\, \hat{\mathcal{J}}^\mu_A\, \mathsf{P}_{\mathrm{phys}}$ on $\mathcal{H}_{\mathrm{phys}}$. For typed/triadic readings we use the commuting projectors $\Pi^{[\chi]}_{\Phi\Phi}$ (Definitions 9.186 and 9.200) and Π_α with $\alpha \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$ (Chapter 9 triad), and write the typed/triadic current as

$$\mathcal{J}^{\mu,[\chi]}_{A,(\alpha)} \;:=\; \Pi_\alpha\, \Pi^{[\chi]}_{\Phi\Phi}\, \mathcal{J}^\mu_A\, \Pi^{[\chi]}_{\Phi\Phi}\, \Pi_\alpha.$$

Remark 9.274 (Notation: generator index A vs. kinetic operator A_x)**.** In Definition 9.273 (and Definition 9.278) the symbol A is a *Lie–algebra index* that labels the generator T_A, the charge $\hat{Q}_A$, and the current $\mathcal{J}^\mu_A$. It is not an operator. Summation over A contracts with the invariant internal metric (equivalently via the Hermitian–completion map $\mathsf{H}[\cdot]$ componentwise) and has no relation to the Legendre/Riesz identification.

By contrast, A_x in Proposition 9.292 (case B) is a *fiberwise kinetic/inertia operator* on the vacuum tangent space $T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}$, appearing in the diagnostic realized reference configuration kinetic term

$$\mathcal{L}_{\Phi,\mathrm{kin}} = \tfrac{1}{2}\, g_T^{(x)}\Big(\dot{\Phi},\ \mathsf{A}_x\,\dot{\Phi}\Big),$$

with A_x positive, boundedly invertible, and $g_T^{(x)}$–self–adjoint. It implements the Legendre/Riesz isometry $\mathcal{R}_x^{(\mathsf{A})} : T_{|\Omega}\mathcal{M}_{\mathrm{vac}} \to T^*_{|\Omega}\mathcal{M}_{\mathrm{vac}}$ and, on the diagnostic realized reference configuration, commutes with the typed/triadic projectors ($[\mathsf{A}_x, \Pi^{[\chi]}_{\Phi\Phi}] = [\mathsf{A}_x, \Pi_\alpha] = 0$).

Finally, the map $\mathsf{H}[\cdot]$ in Definition 9.278 denotes *Hermitian completion* and should not be confused with either the index A or the operator A_x.

In short: A (index) $\neq$ A_x (operator) $\neq$ H (map).

Proposition 9.275 (Conservation, Ward identities, and typed NG coupling)**.** *Assume the Gaiamiti action is G–invariant on the diagnostic realized reference configuration (law–symmetric case, i.e. QI/QII for the relevant types). Then on $\mathcal{H}_{\mathrm{phys}}$:*

(i) Bulk conservation

$\bar{\nabla}_\mu\, \mathcal{J}^\mu_A = 0$ *on* $\mathfrak{P}$. *Equivalently, each typed/triadic component satisfies* $\bar{\nabla}_\mu\, \mathcal{J}^{\mu,[\chi]}_{A,(\alpha)} = 0$.

(ii) Ward identity (typed form)

Let $\delta\Phi_{\mathrm{tang}} \in T_{\bar{\Phi}}\mathcal{M}_{\mathrm{vac}}$ *and* $\hat{h}^{(\alpha)}_{\mu\nu}$ *be the TT Goldstone metric deformation in channel* α *(Definition 9.263). In momentum space,*

$$i\,k_\mu \left\langle \mathcal{J}^{\mu,[\mathrm{TT}]}_{A,(\alpha)}(k)\ \cdots \right\rangle = \left\langle \delta_A \mathcal{O}_{\mathrm{TT}}(k)\ \cdots \right\rangle,$$

which implies an NG pole at $k^2 = 0$ *in the correlator* $\Pi_\alpha \Pi^{[\mathrm{TT}]}_{\Phi\Phi}\, G^{\mathrm{R}}_{\Phi\Phi,\mathrm{phys}}\, \Pi^{[\mathrm{TT}]}_{\Phi\Phi} \Pi_\alpha$ *with nonnegative residue (strictly positive in QII; Proposition 9.197).*

(iii) Explicit breaking and GMOR on $\mathfrak{P}$

If a small symmetry–breaking spurion deforms the law channel, the divergence obeys $\bar{\nabla}_\mu \mathcal{J}^\mu_A = \varepsilon_A\, \mathcal{X}_A$. *Typed/triadic SynTh sources then give a GMOR–type scaling for the TT pseudo–NG band on the plateau (QIV:* $\mathscr{H}_L^{[\Phi\Phi,\mathrm{TT}]} > 0$*; Definition 9.188 and Proposition 9.197).*

All statements are Nielsen–flat and BRST–safe by Propositions 9.191 and 9.207.

Definition 9.276 (Noether charge, flux, and edge class)**.** Let Σ_t be a constant–time slice with unit normal n_μ and induced measure $d\Sigma_\mu := n_\mu\, d\Sigma$. Define the physical Noether charge $Q_A(t) := \int_{\Sigma_t} d\Sigma_\mu\, \mathcal{J}^\mu_A$ and the boundary flux $\Phi_A(t) := \oint_{\partial\Sigma_t} dS_\mu\, \mathcal{J}^\mu_A$. On bulk diagnostic realized reference configuration with vanishing flux, $\dot{Q}_A(t) = 0$; with boundaries we classify the edge via the overlay of Definition 9.214 and record $\mathrm{Edge}^{[S,\chi]}$ in the ledger update (Corollary 9.218).

Corollary 9.277 (Quadrant semantics for currents)**.** *In the TT typed channel: QI ⇒ conserved current with no NG coupling; QII ⇒ conserved current sourcing an exact NG pole; QIV ⇒ nonzero divergence proportional to the typed law source* ($\mathscr{H}_L^{[\Phi\Phi,\mathrm{TT}]} > 0$) *and a pNG band with GMOR slope (Definitions 9.188 and 9.189 and Proposition 9.197). The trace (Apeiron) channel remains QI in the SSB–only regime.*

Definition 9.278 (Charge–generator convention and Hermitian completion)**.** Fix an internal label A and let the conserved charge $\hat{Q}_A$ act by adjoint on field operators so that

$$[\hat{Q}_A, \hat{\mathbf{\Phi}}] = T_A\,\hat{\mathbf{\Phi}}, \qquad \delta_A\hat{\mathbf{\Phi}} = i\,\epsilon\,T_A\hat{\mathbf{\Phi}} = i\,\epsilon\,[\hat{Q}_A, \hat{\mathbf{\Phi}}],$$

with T_A the (matrix) generator in the Gaiamiti representation and ϵ a constant parameter for global transformations. Define the Hermitian completion map $\mathsf{H}[X] := X + X^\dagger$. Assume the invariant internal metric contracts indices so that $\mathrm{Tr}\{AB\}$ implements the same bilinear as $\mathsf{H}[\cdot]$ componentwise.

Lemma 9.279 (Noether current bridge at the operator level)**.** *Let* $\mathcal{L}_\Phi(\hat{\mathbf{\Phi}}, \nabla\hat{\mathbf{\Phi}})$ *be the operator-valued Gaiamiti Lagrangian density. Set*

$$X^\mu_A \;:=\; \frac{\partial\mathcal{L}_\Phi}{\partial(\nabla_\mu\hat{\mathbf{\Phi}})}\,\big(i\,T_A\hat{\mathbf{\Phi}}\big).$$

Then the boxed current of Definition 9.273 and the "+ h.c." current of Definition 9.273 coincide under the invariant trace via

$$\boxed{\hat{\mathcal{J}}^\mu_A = \mathsf{H}\big[X^\mu_A\big] \equiv -i\,\mathrm{Tr}\left\{\frac{\partial\mathcal{L}_\Phi}{\partial(\nabla_\mu\hat{\mathbf{\Phi}})}\,[\hat{Q}_A, \hat{\mathbf{\Phi}}]\right\} =: \;\hat{J}^\mu_{(\Phi)}.} \tag{9.113}$$

Comment. *Equality is exact in the standard unitary representation where the internal trace implements the invariant bilinear; otherwise the two forms differ by an internally traceless anti-Hermitian commutator whose projection to* H^0 *vanishes on the diagnostic realized reference configuration.*

Definition 9.280 (Diagnostic realized reference configuration realization and projection map)**.** On the diagnostic realized reference configuration (Definition 9.185), replace $\nabla \to \bar{\nabla}(\bar{\Sigma})$ and apply the BRST–physical projector $\mathsf{P}_{\mathrm{phys}}$:

$$\mathfrak{B}^\mu_A[\,\cdot\,] \;:=\; \mathsf{P}_{\mathrm{phys}}\,\mathcal{U}_\Sigma\Big[\,(\cdot)\,\Big|_{\,\nabla\to\bar{\nabla}}\Big]\,\mathsf{P}_{\mathrm{phys}}.$$

For typed/triadic readings (Definitions 9.186 and 9.200) with $\alpha \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$, define

$$\mathfrak{B}^{\mu,[\chi]}_{A,(\alpha)}[\,\cdot\,] \;:=\; \Pi_\alpha\,\Pi^{[\chi]}_{\Phi\Phi}\,\mathfrak{B}^\mu_A[\,\cdot\,]\,\Pi^{[\chi]}_{\Phi\Phi}\,\Pi_\alpha.$$

Proposition 9.281 (Equivalence of the two currents on H^0 and in typed/triadic channels)**.** *With the conventions of Definition 9.278 and the map of Definition 9.280, the two definitions are the same current, realized on* H^0*:*

$$\mathcal{J}^\mu_A \;:=\; \mathfrak{B}^\mu_A\big[\hat{J}^\mu_{(\Phi)}\big] \;=\; \mathfrak{B}^\mu_A\big[\hat{\mathcal{J}}^\mu_A\big],$$

and, for every type $[\chi]$ *and triad label* α*,*

$$\boxed{\begin{aligned}\mathcal{J}^{\mu,[\chi]}_{A,(\alpha)} &= \Pi_\alpha\,\Pi^{[\chi]}_{\Phi\Phi}\,\mathcal{J}^\mu_A\,\Pi^{[\chi]}_{\Phi\Phi}\,\Pi_\alpha \\ &= \mathfrak{B}^{\mu,[\chi]}_{A,(\alpha)}\Big[\hat{J}^\mu_{(\Phi)}\Big] \\ &= \mathfrak{B}^{\mu,[\chi]}_{A,(\alpha)}\Big[\hat{\mathcal{J}}^\mu_A\Big].\end{aligned}} \tag{9.114}$$

Proof. *Apply Lemma 9.279 to identify the operator currents; then substitute* $\nabla \to \bar{\nabla}$*, apply* $\mathsf{P}_{\mathrm{phys}}$*, and commute the projectors* $\Pi^{[\chi]}_{\Phi\Phi}, \Pi_\alpha$ *with* $\mathsf{P}_{\mathrm{phys}}$ *(diagnostic realized reference configuration assumptions), yielding* (9.114). □

Corollary 9.282 (Conservation and improvements)**.** *If* S_Φ *is invariant under the* global *transformation (*$\epsilon = \text{const.}$*), then* $\bar{\nabla}_\mu\,\mathcal{J}^\mu_A = 0$ *on* H^0 *up to standard improvement terms and boundary contributions determined by* $\bar{\Sigma}$*; the same holds typewise and triadically for* $\mathcal{J}^{\mu,[\chi]}_{A,(\alpha)}$*.*

Remark 9.283 (Local parameter and anomalies)**.** For $\epsilon(x)$ (local variation) one obtains the Ward identity with source $\partial_\mu\epsilon(x)$; possible BRST or internal anomalies appear as usual in the divergence of $\mathcal{J}^\mu_A$ and are diagnosed in the triadic/typed channels by applying (9.114).

Interpretation. The expectation current $\bar{J}^\mu_{(\Phi)}(x) := \langle \mathring{\boldsymbol{\Psi}} \,|\, \hat{J}^\mu_{(\Phi)}(x) \,|\, \mathring{\boldsymbol{\Psi}} \rangle$ associated with the operator current of Definition 9.271 is a geometric conservation law: unlike Standard Model currents that transport *matter* charges, it transports conserved quantities tied to *symmetries of the geometric fabric* itself. When the generator $\hat{Q}$ implements a continuous motion on the vacuum manifold $\mathcal{M}_{\mathrm{vac}}$ (Definition 6.61), $\bar{J}^\mu_{(\Phi)}$ is the flow of the corresponding "geometric charge" [91, 100], ensuring compatibility between internal geometric symmetries and emergent dynamics.

On the diagnostic realized reference configuration, conservation appears in typed Ward form (Proposition 9.275):

$$\bar{\nabla}_\mu\,\mathcal{J}^\mu_A = 0 \quad\Longrightarrow\quad i\,k_\mu\,\Big\langle \mathcal{J}^{\mu,[\chi]}_{A,(\alpha)}(k)\,\cdots\Big\rangle = \Big\langle \delta_A\mathcal{O}^{[\chi]}_{(\alpha)}(k)\,\cdots\Big\rangle,$$

where $\mathcal{J}^{\mu,[\chi]}_{A,(\alpha)}$ is the typed/triadic projection of $\mathcal{J}^\mu_A$ defined in Definition 9.273 and linked to either current representation by eq. (9.114). In the TT channel this Ward identity enforces an exact NG pole at $k^2 = 0$ in QII (residue fixed by the TT Goldstone overlap; Proposition 9.197), whereas in QIV the pole remains as pNG with a GMOR slope set by the typed law source (Definition 9.188 and Proposition 9.197). Operationally, coupling a weak probe a^A_μ via $\delta S_\Phi = \int d^dx\, a^A_\mu\,\mathcal{J}^\mu_A$ promotes $\mathcal{J}^\mu_A$ to a susceptibility meter: the linear response

$$\chi^{[\chi]}_{AB,(\alpha)}(k) := i\,G^{\mathrm{R}}_{\mathcal{J}_A\mathcal{J}_B}\Big|_{[\chi],(\alpha)}(k)$$

reads the symmetry geometry; in the static, long-wavelength limit $\chi^{[\chi]}_{AB}(\omega\to 0, \mathbf{k}\to 0)$ measures the metric on the typed slice of $\mathcal{M}_{\mathrm{vac}}$.

The current inherits the triadic anatomy (Definition 9.263):

$$\mathcal{J}^\mu_A = \sum_{\alpha\in\{\mathrm{corr,vac,spec}\}} \Pi_\alpha\,\mathcal{J}^\mu_A\,\Pi_\alpha,$$

which separates correlational transport, vacuum–bundle, and spectral–carrier contributions. Their relative weights c_α coincide with those assembling the composite TT graviton on H^0 (Definition 9.265). With boundaries, Stokes' theorem relates bulk charge and boundary flow,

$$\dot{Q}_A(t) = -\Phi_A(t), \qquad Q_A(t) = \int_{\Sigma_t} d\Sigma_\mu \, \mathcal{J}_A^\mu, \quad \Phi_A(t) = \oint_{\partial\Sigma_t} dS_\mu \, \mathcal{J}_A^\mu$$

(Definition 9.276); comparing bulk vs. boundary typed meters fixes the edge class via the bulk/edge overlay and ledger update (Definition 9.214 and Corollary 9.218).

Scheme choices preserve physics: improvements $\mathcal{J}_A^\mu \to \mathcal{J}_A^\mu + \bar{\nabla}_\nu K_A^{[\mu\nu]}$ leave charges and typed Ward identities intact; BRST–exact additions vanish on H^0; and Nielsen/compatibility results guarantee gauge–parameter independence across the plateau $\mathfrak{P}$ (Propositions 9.191 and 9.207). Finally, the channel roles are complementary: in $(S, [\chi]) = (\Phi\Phi, \mathrm{Tr})$ the sector sits in QI—$\mathcal{J}_A^\mu$ diagnoses a *gapped* trace (Apeiron) channel with intrinsic scale m_A and correlation length $\xi_A \sim m_A^{-1}$ but no NG content—while the TT type carries the symmetry–breaking kinematics (QII/QIV). Conservation becomes operational at the typed onset $\mathrm{On}^{[S,\chi]}$ where SynTh meters stabilize (Definition 9.190); below onset, transient law–state mixing may occur without affecting plateau charges or residues.

The Geometric Charge as an Intrinsic Property of the Apeiron (augmented) The geometric charge is an intrinsic property of the *Apeiron* sector; this follows from the chain below.

(i) Provenance of the current (operator → DRCC, typed) The Noether current arises from a continuous internal symmetry of $S_\Phi[\hat{\mathbf{\Phi}}]$. Via the charge–generator bridge (Definition 9.278) and the operator identity (Lemma 9.279), the two operator forms $\hat{J}^\mu_{(\Phi)} = -i\,\mathrm{Tr}\{\partial\mathcal{L}_\Phi/\partial(\nabla_\mu\hat{\mathbf{\Phi}})\,[\hat{Q}, \hat{\mathbf{\Phi}}]\}$ and $\hat{\mathcal{J}}_A^\mu = \frac{\partial\mathcal{L}_\Phi}{\partial(\nabla_\mu\hat{\mathbf{\Phi}})}(i\,T_A\hat{\mathbf{\Phi}}) + \mathrm{h.c.}$ are equivalent. Realizing on the diagnostic realized reference configuration (DRCC) and projecting to H^0 with $\mathfrak{B}_A^\mu$ (Definition 9.280) yields $\mathcal{J}_A^\mu = \mathsf{P}_{\mathrm{phys}}\,\hat{\mathcal{J}}_A^\mu\,\mathsf{P}_{\mathrm{phys}}$, and the typed/triadic component $\mathcal{J}^{\mu,[\chi]}_{A,(\alpha)}$ as in Definition 9.273. Equivalence in each typed/triadic channel is *exact* by Proposition 9.281 and eq. (9.114). The conserved charge is $Q_A(t) = \int_{\Sigma_t} d\Sigma_\mu\,\mathcal{J}_A^\mu$ (Definition 9.276).

(ii) Action–quantum correspondence (Apeiron channel on $\mathfrak{P}$) The same action that defines the current also governs the *Apeiron*, i.e. the trace-typed quantum in $(S, [\chi]) = (\Phi\Phi, \mathrm{Tr})$ (Section 9.17.1). On a certified plateau $\mathfrak{P}$, the intrinsic scale is fixed by the typed curvature $m_A^2 = \mathrm{eig}_{\min}\big(\mathcal{K}^{[\mathrm{Tr}]}_{\Phi\Phi}\big)$ (Definition 9.242), so the correlation length behaves as $\xi_A \sim m_A^{-1}$. Let $\hat{a}^\dagger_{n,\mathrm{Tr}}$ create $\bar{\Sigma}$-orthonormal trace modes. Using the charge–generator convention (Definition 9.278),

$$[Q_A,\ \hat{a}^\dagger_{n,\mathrm{Tr}}] \;=\; q^{(\mathrm{Tr})}_{A,n}\,\hat{a}^\dagger_{n,\mathrm{Tr}} \;+\; \Big(\text{BRST-exact} \;\oplus\; \text{orthogonal-type pieces}\Big),$$

so the conserved quantum number $q^{(\mathrm{Tr})}_{A,n}$ is *carried* by Apeiron quanta on H^0.

(iii) Principle of charge attribution (typed/triadic semantics) Conserved charges extracted from S_Φ belong to the quanta of the field in the *relevant typed channel*. In the trace type the quadrant is $\mathrm{Quad}^{[\Phi\Phi,\mathrm{Tr}]} = \mathrm{QI}$ (Definition 9.189): the sector is gapped (no

NG/pNG), yet the charge is well-defined and measurable via (i) the equal-time commutator above, (ii) the static, long-wavelength limit of the typed current–current susceptibility $\chi^{[\mathrm{Tr}]}_{AB}(\omega\to 0,\mathbf{k}\to 0)$ (from the probe coupling in Proposition 9.275), and (iii) the ledgered **TLO** stamp recorded per Corollary 9.218.

Hence, because the Noether current and charge originate in the symmetry of the same action that defines the Apeiron, the associated conserved *geometric charge* is an intrinsic property of the Apeiron sector—consistent with QI (no NG content in this channel).

Elucidation on the Nature of Geometric Charge (augmented) It is crucial to distinguish "geometric charge'' from familiar matter charges.

1. An electron *possesses* electric charge as a property of a matter field; its current lives in a matter sector and counts matter quanta.

2. The Apeiron does not *possess* geometric charge in that ordinary sense; rather, it *is* the trace-typed quantum manifestation of a conserved property of the geometric vacuum, read from the diagnostic realized reference configuration-realized current $\mathcal{J}^{\mu}_{A}$ and its triadic anatomy $\mathcal{J}^{\mu}_{A}=\sum_{\alpha\in\{\mathrm{corr,vac,spec}\}}\Pi_{\alpha}\,\mathcal{J}^{\mu}_{A}\,\Pi_{\alpha}$ (Definition 9.273).

If the symmetry corresponds to a continuous rotation on $\mathcal{M}_{\mathrm{vac}}$ (Definition 6.61), the geometric charge is the conserved "orientation'' on $\mathcal{M}_{\mathrm{vac}}$; Apeiron quanta are its trace-typed carriers. By contrast, the TT type encodes symmetry breaking: the same current sources an NG (or pNG) pole (Proposition 9.197) whose residue feeds the composite graviton via triadic weights c_{α} (Definition 9.265), while the Apeiron remains gapped with scale m_{A} (Definition 9.242). Bulk–boundary overlays relate charge and flux by $\dot{Q}_{A}(t)=-\Phi_{A}(t)$ with Q_{A},Φ_{A} as in Definition 9.276; edge behavior is classified and ledgered via Definition 9.214 and Corollary 9.218. None of the admissible scheme choices—Belinfante improvements (Corollary 9.282), BRST-exact pieces (Proposition 9.207), or gauge-parameter changes (Nielsen flatness; Proposition 9.191)—alter charge attribution on H^{0} or the typed Ward identities across the plateau $\mathfrak{P}$.

9.17.9 The Noether Stress–Energy Tensor of the Gaiamiti Field

Within the framework of any physical theory, continuous symmetries of the action imply, via Noether's theorem, conserved currents. For the Gaiamiti field, invariance under infinitesimal spacetime translations yields the stress–energy tensor. In the IQR formalism this tensor—denoted $\hat{T}^{\mu\nu}_{(\Phi)}(x)$—is an operator–valued field encoding energy–momentum flux of *relational* geometric variations. Its expectation value $\bar{T}^{\mu\nu}_{(\Phi)}(x)$ plays the role of a classical stress–energy in GR, but sources the emergent, state–dependent geometry rather than a fixed background.

Definition 9.284 (Noether Stress–Energy Tensor of the Gaiamiti Field)**.** Let $\hat{\mathbf{\Phi}}(x)$ be the Gaiamiti field on $(\mathcal{X}_{q},\bar{\Sigma}_{\mu\nu})$ with action $S_{\Phi}[\hat{\mathbf{\Phi}}]$. The operator–valued Noether stress–energy tensor is

$$\boxed{\hat{T}^{\mu\nu}_{(\Phi)}(x):=\Big\{\big(\bar{\nabla}^{\mu}\hat{\mathbf{\Phi}}(x)\big)^{\dagger},\,\bar{\nabla}^{\nu}\hat{\mathbf{\Phi}}(x)\Big\}\;-\;\bar{\Sigma}^{\mu\nu}(x)\,\mathrm{Tr}\Big(\mathcal{L}_{\Phi}(x)\Big)\,\mathbb{I},}\tag{9.115}$$

Conventions.

(i) Relational Levi–Civita derivative
$\bar{\nabla}_\mu$ is the relational Levi–Civita derivative compatible with $\bar{\Sigma}_{\mu\nu}$ (Definition 6.7); indices are raised with $\bar{\Sigma}^{\mu\nu}$ so that $\bar{\nabla}^\mu := \bar{\Sigma}^{\mu\lambda}\bar{\nabla}_\lambda$ (Definition 6.184).

(ii) Anticommutator
$\{A, B\} := AB + BA$ is the anticommutator, ensuring Hermiticity and symmetry in (μ, ν).

(iii) Gaiamiti Lagrangian density
$\mathcal{L}_\Phi(x)$ is the operator–valued Gaiamiti Lagrangian *density* (Definition 6.172); on the diagnostic realized reference configuration it depends on $\bar{\Sigma}$ only through $\bar{\nabla}$-contractions and $\sqrt{-\det \bar{\Sigma}}$.

(iv) Internal trace
$\mathrm{Tr}(\cdot)$ traces over internal/operator fibers (not spacetime; Section 6.1.10), hence $\mathrm{Tr}(\mathcal{L}_\Phi)$ is a c–number density.

(v) Identity on the modal fiber
$\mathbb{I} \equiv \mathbb{I}_{\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}}$ is the identity on the local modal fiber (eq. (6.131)).

Hilbert–Noether equivalence. For minimally coupled $\mathcal{L}_\Phi$,

$$\hat{T}^{\mu\nu}_{(\Phi)}(x) = \frac{2}{\sqrt{-\det \bar{\Sigma}(x)}} \frac{\delta S_\Phi}{\delta \bar{\Sigma}_{\mu\nu}(x)} \quad \text{(up to standard improvement terms).}$$

Expectation on the diagnostic realized reference configuration Taking the expectation value in the Quantum Holo–State $|\mathring{\boldsymbol{\Psi}}\rangle$ defines the classical (expectation–level) stress–energy:

$$\boxed{\bar{T}^{\mu\nu}_{(\Phi)}(x) := \Big\langle \mathring{\boldsymbol{\Psi}} \Big| \mathsf{P}_{\mathrm{phys}}\, \hat{T}^{\mu\nu}_{(\Phi)}(x)\, \mathsf{P}_{\mathrm{phys}} \Big| \mathring{\boldsymbol{\Psi}} \Big\rangle \equiv \langle \mathring{\boldsymbol{\Psi}} | \hat{T}^{\mu\nu}_{(\Phi)}(x) | \mathring{\boldsymbol{\Psi}} \rangle,} \tag{9.116}$$

where the equivalence holds whenever $|\mathring{\boldsymbol{\Psi}}\rangle \in \mathcal{H}_{\mathrm{phys}}$. On the diagnostic realized reference configuration we regard $\hat{T}^{\mu\nu}_{(\Phi)}$ as band–limited/IR–windowed and properly normal–ordered so that $\bar{T}^{\mu\nu}_{(\Phi)}$ is finite on the certified plateau $\mathfrak{P}$.
Typed/triadic expectation. For $\alpha \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$ and $[\chi] \in \{\mathrm{TT}, \mathrm{Tr}\}$, set

$$\bar{T}^{\mu\nu\,[\chi]}_{(\Phi),(\alpha)}(x) := \Big\langle \mathring{\boldsymbol{\Psi}} \Big| \Pi_\alpha \Pi^{[\chi]}_{\Phi\Phi}\, \hat{T}^{\mu\nu}_{(\Phi)}(x)\, \Pi^{[\chi]}_{\Phi\Phi} \Pi_\alpha \Big| \mathring{\boldsymbol{\Psi}} \Big\rangle. \tag{9.117}$$

It inherits symmetry $\bar{T}^{\mu\nu} = \bar{T}^{\nu\mu}$ and transforms covariantly with $\bar{\nabla}$. With (9.116)–(9.117) in place, the conservation law of Proposition 9.285 follows.

Proposition 9.285 (Conservation of the Noether Stress–Energy Tensor)**.** *If the Gaiamiti field $\hat{\boldsymbol{\Phi}}(x)$ satisfies its Euler–Lagrange equation (Definition 9.268), then the expectation–level stress–energy tensor is covariantly conserved with respect to the relational Levi–Civita connection:*

$$\boxed{\bar{\nabla}_\mu\, \bar{T}^{\mu\nu}_{(\Phi)}(x) = 0.} \tag{9.118}$$

Proof of Proposition 9.285. The conservation of the Noether stress–energy tensor follows from diffeomorphism invariance of the Gaiamiti action S_Φ together with the field equations.

Step 1: Diffeomorphism invariance and total variation. Write the action as an integral of a scalar density,

$$S_\Phi = \int_{\mathcal{X}_q} d^d x \, \mathcal{L}_\Phi,$$

and consider an infinitesimal diffeomorphism $x^\mu \mapsto x'^\mu = x^\mu - \xi^\mu(x)$ with $\xi^\mu|_{\partial\mathcal{X}_q} = 0$. The total variation splits into field and geometry parts:

$$\delta S_\Phi = \int_{\mathcal{X}_q} \left[\frac{\delta S_\Phi}{\delta \hat{\mathbf{\Phi}}} \, \delta\hat{\mathbf{\Phi}} + \frac{\delta S_\Phi}{\delta \bar{\Sigma}_{\mu\nu}} \, \delta\bar{\Sigma}_{\mu\nu} \right] d^d x = 0.$$

Step 2: On–shell reduction. Assuming the Euler–Lagrange equations hold (Definition 9.268), we have $\delta S_\Phi / \delta\hat{\mathbf{\Phi}} = 0$, hence

$$\delta S_\Phi = \int_{\mathcal{X}_q} \frac{\delta S_\Phi}{\delta \bar{\Sigma}_{\mu\nu}} \, \delta\bar{\Sigma}_{\mu\nu} \, d^d x = 0.$$

Step 3: Metric variation and Hilbert–Noether identity. Under the diffeomorphism,

$$\delta\bar{\Sigma}_{\mu\nu} = \mathcal{L}_\xi \bar{\Sigma}_{\mu\nu} = \bar{\nabla}_\mu \xi_\nu + \bar{\nabla}_\nu \xi_\mu,$$

with $\bar{\nabla}$ the Levi–Civita derivative of $\bar{\Sigma}$. Using the Hilbert form (equivalent to the boxed Noether definition in (9.115) up to standard improvement terms),

$$\frac{\delta S_\Phi}{\delta \bar{\Sigma}_{\mu\nu}} = \frac{\sqrt{-\det \bar{\Sigma}}}{2} \, \hat{T}^{\mu\nu}_{(\Phi)},$$

we obtain

$$\delta S_\Phi = \int_{\mathcal{X}_q} \sqrt{-\det \bar{\Sigma}} \; \hat{T}^{\mu\nu}_{(\Phi)} \, \bar{\nabla}_\mu \xi_\nu \, d^d x = 0,$$

where symmetry $\hat{T}^{\mu\nu}_{(\Phi)} = \hat{T}^{\nu\mu}_{(\Phi)}$ was used to combine the two terms.

Step 4: Integration by parts and operator identity. Integrating by parts and dropping the boundary term (since ξ^μ vanishes on $\partial\mathcal{X}_q$),

$$0 = \int_{\mathcal{X}_q} \sqrt{-\det \bar{\Sigma}} \left[\bar{\nabla}_\mu\big(\hat{T}^{\mu\nu}_{(\Phi)} \xi_\nu\big) - \big(\bar{\nabla}_\mu \hat{T}^{\mu\nu}_{(\Phi)}\big)\xi_\nu \right] d^d x = - \int_{\mathcal{X}_q} \sqrt{-\det \bar{\Sigma}} \, \big(\bar{\nabla}_\mu \hat{T}^{\mu\nu}_{(\Phi)}\big) \, \xi_\nu \, d^d x.$$

Because ξ_ν is arbitrary, we conclude the operator-level conservation law:

$$\bar{\nabla}_\mu \hat{T}^{\mu\nu}_{(\Phi)}(x) = 0.$$

Step 5: Expectation-level conservation. Taking the expectation in the Quantum Holo–State $|\overset{\circ}{\mathbf{\Psi}}\rangle$ and using linearity of $\bar{\nabla}$ on spacetime indices yields

$$\bar{\nabla}_\mu \, \bar{T}^{\mu\nu}_{(\Phi)}(x) = \bar{\nabla}_\mu \, \langle \overset{\circ}{\mathbf{\Psi}} | \hat{T}^{\mu\nu}_{(\Phi)}(x) | \overset{\circ}{\mathbf{\Psi}} \rangle = 0,$$

which is (9.118). □

Corollary 9.286 (Typed/triadic conservation on the diagnostic realized reference configuration)**.** *For* $\alpha \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$ *and* $[\chi] \in \{\mathrm{TT}, \mathrm{Tr}\}$*, define the* operator-level *typed/triadic tensor*

$$\hat{T}^{\mu\nu\,[\chi]}_{(\Phi),(\alpha)} := \Pi_\alpha\, \Pi^{[\chi]}_{\Phi\Phi}\, \hat{T}^{\mu\nu}_{(\Phi)}\, \Pi^{[\chi]}_{\Phi\Phi}\, \Pi_\alpha,$$

and its expectation-level *version* $\bar{T}^{\mu\nu\,[\chi]}_{(\Phi),(\alpha)} := \langle \overset{\circ}{\boldsymbol{\Psi}} | \hat{T}^{\mu\nu\,[\chi]}_{(\Phi),(\alpha)} | \overset{\circ}{\boldsymbol{\Psi}} \rangle$*. On the certified plateau* $\mathfrak{P}$ *(diagnostic realized reference configuration projectors commuting with* $\bar{\nabla}$ *within tolerances),*

$$\bar{\nabla}_\mu\, \bar{T}^{\mu\nu\,[\chi]}_{(\Phi),(\alpha)} = 0.$$

Remark 9.287 (Scope and improvements)**.** (i) Belinfante–Rosenfeld improvements $T^{\mu\nu} \to T^{\mu\nu} + \bar{\nabla}_\lambda K^{[\lambda\mu]\nu}$, with $K^{[\lambda\mu]\nu} = -K^{[\mu\lambda]\nu}$, preserve (9.118) ([462, 463, 464]). (ii) If $\mathcal{L}_\Phi$ contains explicit (nonminimal) metric dependence beyond $\bar{\nabla}$-contractions and $\sqrt{-\det \bar{\Sigma}}$, additional terms appear in $\bar{\nabla}_\mu T^{\mu\nu}$ proportional to that breaking; potential anomalies would modify the divergence in the standard way (cf. [301, 300]).

Interpretation and Physical Significance

The stress–energy tensor $\hat{T}^{\mu\nu}_{(\Phi)}(x)$ is a fundamentally geometric object. By construction in (9.115) (and equivalently by the Hilbert–Noether identity stated under that definition), it is Hermitian and symmetric,

$$\hat{T}^{\mu\nu}_{(\Phi)} = \hat{T}^{\nu\mu}_{(\Phi)},$$

with symmetry preserved under Belinfante–Rosenfeld improvements (see Remark 9.287). When the field equations hold, its expectation is covariantly conserved on the diagnostic realized reference configuration,

$$\bar{\nabla}_\mu\, \bar{T}^{\mu\nu}_{(\Phi)} = 0,$$

as proved in Proposition 9.285; typed/triadic slices obey the same conservation law by Corollary 9.286. The expectation $\bar{T}^{\mu\nu}_{(\Phi)}$ is defined in (9.116).

At the Quantum Level. The operator $\hat{T}^{\mu\nu}_{(\Phi)}(x)$ captures the local energy–momentum flux of the relational degrees of freedom of quantum geometry encoded by the Gaiamiti field $\hat{\boldsymbol{\Phi}}$ (see Definition 6.172 and §6.1.10). In the band-limited, BRST-physical setting on the certified plateau $\mathfrak{P}$, its typed/triadic projections

$$\hat{T}^{\mu\nu\,[\chi]}_{(\Phi),(\alpha)} = \Pi_\alpha\, \Pi^{[\chi]}_{\Phi\Phi}\, \hat{T}^{\mu\nu}_{(\Phi)}\, \Pi^{[\chi]}_{\Phi\Phi}\, \Pi_\alpha, \qquad \alpha \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\},\ [\chi] \in \{\mathrm{TT}, \mathrm{Tr}\}$$

(Corollary 9.286 and Definitions 9.186 and 9.200) resolve:

(i) Kinetic flux Kinetic flux from relational propagation across $\mathcal{X}_q$ (the covariant-derivative term in (9.115)).

(ii) Potential energy Potential energy of internal deformations tied to the Gaiamiti potential (cf. Definition 9.235 and the typed curvature drivers Definition 9.220).

This decomposition is the energy–momentum analogue of the current's typed/triadic anatomy used in Definition 9.273 and Proposition 9.275.

At the Classical Level. The expectation $\bar{T}^{\mu\nu}_{(\Phi)}(x)$ (eq. (9.116)) functions as the effective source for the realized chronogeometry through the Hilbert–Noether equivalence just below (9.115):

$$\frac{\delta S_\Phi}{\delta \bar{\Sigma}_{\mu\nu}} = \frac{\sqrt{-\det \bar{\Sigma}}}{2}\, \hat{T}^{\mu\nu}_{(\Phi)} \quad \text{(up to improvements)}.$$

In an IQR gravity closure, this provides the right–hand side of a relational Einstein-like equation in which $\bar{\Sigma}_{\mu\nu}$ is state-dependent and dynamical rather than a fixed background.

The Self-Sourcing Nature of Geometry

The principle is *geometry sources itself*: the energy–momentum of relational fluctuations governs the evolution of the very geometry that supports them. Concretely, the structure of $\hat{T}^{\mu\nu}_{(\Phi)}$ in (9.115) shows:

(i) Flux transport term The $\{(\bar{\nabla}^\mu \hat{\mathbf{\Phi}})^\dagger, \bar{\nabla}^\nu \hat{\mathbf{\Phi}}\}$ term encodes local transport of quantum-geometric flux across $\mathcal{X}_q$.

(ii) Background-independent subtraction The subtraction $\bar{\Sigma}^{\mu\nu}\,\mathrm{Tr}(\mathcal{L}_\Phi)\,\mathbb{I}$ enforces compatibility with relational background independence and yields the Hilbert source via the metric variation just cited.

Remark 9.288 (Relation to the graviton and typed symmetry breaking)**.** When $\bar{T}^{\mu\nu}_{(\Phi)}$ is sourced by excitations of the Apeiron (trace type) about its certified background (Definition 9.242), the trace channel remains gapped and law-symmetric (QI), while the TT channel carries symmetry-breaking kinematics. In the TT type, the Noether current's Ward identity (Proposition 9.275) enforces an NG (or pNG, in QIV) pole whose residue feeds the composite graviton; the corresponding energy–momentum coupling is isolated by the TT projector on $T^{\mu\nu}$ (cf. the metric-response line under (9.115)). This mirrors the phonon analogy in condensed matter ([461, 14, 393]) but with geometry's TT Goldstone as the graviton; onset/location and edge behavior are recorded through the typed ledger (Definitions 9.190 and 9.198 and Corollary 9.286).

9.17.10 Projection Operators and Geometric Duality

Definition 9.289 (Projection Operators for Geometric and Matter Sectors)**.** Let $\mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}}$ be the local modal Hilbert space at $x \in \mathcal{X}_q$, with local vacuum $|\Omega(x)\rangle \in \mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}}$, and let $T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}} \subset \mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}}$ denote the (assumed closed) tangent space to the vacuum manifold at $|\Omega(x)\rangle$.

Conventions. $g^{(x)}_{\mathrm{mod}}(\cdot,\cdot)$ is the relational inner product on $\mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}}$; orthogonality $(\cdot)^\perp$ is taken w.r.t. $g^{(x)}_{\mathrm{mod}}$. The identity on the fiber is $\mathbb{I}_x \equiv \mathbb{I}_{\mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}}}$. Bras $\langle\cdot|$ denote the $g^{(x)}_{\mathrm{mod}}$–Riesz duals. If $T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}$ is infinite–dimensional, operator sums below are understood in the strong–operator topology.

(i) Geometric projector

For any $g^{(x)}_{\mathrm{mod}}$–orthonormal basis $\{|\delta\Omega_i(x)\rangle\}_{i=1}^{r}$ of $T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}$ ($r = \dim T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}$, finite or countable), define

$$\hat{P}_T(x) \;:=\; \sum_{i=1}^{r} |\delta\Omega_i(x)\rangle\langle\delta\Omega_i(x)|, \tag{9.119}$$

which is basis–independent and satisfies $\hat{P}_T(x)^2 = \hat{P}_T(x) = \hat{P}_T(x)^\dagger$. Its image is the geometric excitation space

$$\mathcal{E}_{\mathrm{geo}}(x) \;:=\; \mathrm{Im}\,\hat{P}_T(x) \;=\; T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}. \tag{9.120}$$

This space supports the dynamical Gaiamiti field $\hat{\mathbf{\Phi}}(x)$; in particular, its *trace–typed* quanta are the Apeiron (Definition 9.242).

(ii) Matter projector

The orthogonal–complement projector is

$$\hat{P}_{\mathrm{exc}}(x) \;:=\; \mathbb{I}_x - \hat{P}_T(x), \tag{9.121}$$

with $\hat{P}_{\mathrm{exc}}(x)^2 = \hat{P}_{\mathrm{exc}}(x) = \hat{P}_{\mathrm{exc}}(x)^\dagger$ and $\hat{P}_{\mathrm{exc}}(x)\hat{P}_T(x) = 0$. Its image is the excitation fiber

$$\mathcal{E}_{\mathrm{exc}}(x) \;:=\; \mathrm{Im}\,\hat{P}_{\mathrm{exc}}(x) \;=\; \big(T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}\big)^{\perp}, \tag{9.122}$$

which supports the Tantrakyma field $\hat{\mathbf{\Theta}}(x)$; its quantized excitations are the kymons.

Consequently, $\mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}} = \mathcal{E}_{\mathrm{geo}}(x) \oplus \mathcal{E}_{\mathrm{exc}}(x)$ and $\hat{P}_T(x) + \hat{P}_{\mathrm{exc}}(x) = \mathbb{I}_x$.

Remark 9.290 (Orthogonal decomposition of modal degrees of freedom)**.** With the projectors of Definition 9.289, the fiber $\mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}}$ splits $g^{(x)}_{\mathrm{mod}}$–orthogonally as

$$\mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}} = \mathcal{E}_{\mathrm{geo}}(x) \oplus \mathcal{E}_{\mathrm{exc}}(x), \qquad \hat{P}_T(x) + \hat{P}_{\mathrm{exc}}(x) = \mathbb{I}_x, \quad \hat{P}_T(x)\hat{P}_{\mathrm{exc}}(x) = 0.$$

This partition is *canonical* relative to the choice of vacuum $|\Omega(x)\rangle$ and relational inner product $g^{(x)}_{\mathrm{mod}}$, and it is basis–independent: for any $g^{(x)}_{\mathrm{mod}}$–orthonormal basis of $T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}$, the series in (9.119) (finite or countable) converges in the strong–operator topology to the same $\hat{P}_T(x)$.

Regularity. Assume $x \mapsto \hat{P}_T(x)$ is C^1 on the diagnostic realized reference configuration. Then $\mathcal{E}_{\mathrm{geo}} := \mathrm{Im}\,\hat{P}_T$ and $\mathcal{E}_{\mathrm{exc}} := \mathrm{Im}\,\hat{P}_{\mathrm{exc}}$ assemble into smooth subbundles of $\mathcal{H}_{\mathrm{mod}}$ over $x \in \mathcal{X}_q$. Moreover, we assume compatibility with the typed/triadic structure of the $\Phi\Phi$ sector:

$$[\hat{P}_T(x), \Pi^{[\chi]}_{\Phi\Phi}] = 0, \qquad [\hat{P}_T(x), \Pi_\alpha] = 0,$$

so typed/triadic readings (Definitions 9.186 and 9.200) respect the geometric–matter split fiberwise.

Parallel transport of the split. If, in addition, $\bar{\nabla}_\mu \hat{P}_T(x) = 0$ (with $\bar{\nabla}$ from Definition 6.7), the decomposition is *parallel*: covariant transport preserves $\mathcal{E}_{\mathrm{geo}}$ and $\mathcal{E}_{\mathrm{exc}}$ along $\mathcal{X}_q$. More generally, the off–diagonal operator $\bar{\nabla}_\mu \hat{P}_T(x)$ measures any residual leakage between the two subbundles; on the certified plateau $\mathfrak{P}$ this is taken to vanish within diagnostic realized reference configuration tolerances, ensuring the conservation statements and typed Ward identities apply slice–wise (cf. Corollary 9.286).

Definition 9.291 (Diagnostic realized reference configuration–compatibility assumptions)**.** **Setting.** Work on the instrumented diagnostic realized reference configuration (Definition 9.185) on a certified plateau $\mathfrak{P}$. Use the geometric/matter split of Definition 9.289 and Remark 9.290 and the typed/triadic projectors $\Pi_{\Phi\Phi}^{[\chi]}$ and Π_α ($\alpha \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$; Definitions 9.186 and 9.200).

Smoothness. The fiber maps $x \mapsto \hat{P}_T(x)$, $\Pi_{\Phi\Phi}^{[\chi]}(x)$, $\Pi_\alpha(x)$ are C^1 in $x \in \mathcal{X}_q$. When used, $x \mapsto J^{(x)}$ (Kähler branch) and $x \mapsto \mathsf{A}_x$ (Legendre branch) are also C^1.

Commutation (diagnostic realized reference configuration tolerances). Projectors and diagnostic realized reference configuration structures commute within plateau tolerances:

$$[\hat{P}_T(x), \Pi_{\Phi\Phi}^{[\chi]}] = [\hat{P}_T(x), \Pi_\alpha] = 0, \qquad [\mathsf{P}_{\mathrm{phys}}, \Pi_{\Phi\Phi}^{[\chi]}] = [\mathsf{P}_{\mathrm{phys}}, \Pi_\alpha] = 0.$$

In the Kähler branch, $[J^{(x)}, \Pi_{\Phi\Phi}^{[\chi]}] = [J^{(x)}, \Pi_\alpha] = 0$; in the Legendre branch, $[\mathsf{A}_x, \Pi_{\Phi\Phi}^{[\chi]}] = [\mathsf{A}_x, \Pi_\alpha] = 0$. Band tags commute with these projectors ($P_x(\lambda_{\mathrm{UV}})$, $K_{\mathrm{IR}}^{(\ell)}$; Chapter 8).

Parallel transport (optional strengthening). Either $\bar{\nabla}_\mu \hat{P}_T(x) = 0$ (parallel split) or the leakage operator $\bar{\nabla}_\mu \hat{P}_T(x)$ is negligible on $\mathfrak{P}$, ensuring typed/triadic Ward and conservation statements apply slice-wise (cf. Corollary 9.286).

Functional–analytic regularity. $T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}$ is closed in $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$; sums like $\sum_i |\delta\Omega_i\rangle\langle\delta\Omega_i|$ converge in the strong-operator topology (Definition 9.289).

Proposition 9.292 (Vacuum duality and geometric identification)**.** *Let the vacuum manifold $\mathcal{M}_{\mathrm{vac}}$ carry the metric induced from the relational inner product on $\mathcal{H}_{\mathrm{mod}}$ at $|\Omega(x)\rangle$, so that $T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}$ is a Hilbert space with inner product $g_T^{(x)}$. Assume, in addition to Definition 9.289 and Remark 9.290, one of the following sufficient structures at each $x \in \mathcal{X}_q$:*

(i) (A) Kähler duality
Assume that $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ admits a Kähler triple $\big(g_{\mathrm{mod}}^{(x)}, J^{(x)}, \omega^{(x)}\big)$ with $J^{(x)}\big(T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}\big) = \mathcal{E}_{\mathrm{exc}}(x)$, $\omega^{(x)}(\cdot,\cdot) = g_{\mathrm{mod}}^{(x)}(\cdot, J^{(x)}\cdot)$, and, on the diagnostic realized reference configuration, $J^{(x)}$ commutes with the typed/triadic projectors: $[J^{(x)}, \Pi_{\Phi\Phi}^{[\chi]}] = [J^{(x)}, \Pi_\alpha] = 0$.

(ii) (B) Legendre nondegeneracy (diagnostic realized reference configuration Lagrangian)
The $\Phi\Phi$-sector kinetic term is strictly nondegenerate on the diagnostic realized reference configuration, $\mathcal{L}_{\Phi,\mathrm{kin}} = \frac{1}{2} g_T^{(x)}(\dot{\Phi}, \mathsf{A}_x\dot{\Phi})$, with $\mathsf{A}_x : T_{|\Omega}\mathcal{M}_{\mathrm{vac}} \to T_{|\Omega}\mathcal{M}_{\mathrm{vac}}$ positive and boundedly invertible, and $[\mathsf{A}_x, \Pi_{\Phi\Phi}^{[\chi]}] = [\mathsf{A}_x, \Pi_\alpha] = 0$. The fiberwise Legendre/Riesz map $\mathcal{R}_x^{(\mathsf{A})} : v \mapsto g_T^{(x)}(\mathsf{A}_x v, \cdot)$ is an isometric isomorphism $T_{|\Omega}\mathcal{M}_{\mathrm{vac}} \xrightarrow{\cong} T_{|\Omega}^*\mathcal{M}_{\mathrm{vac}}$. Furthermore, the Tantrakyma fiber is the canonical momentum *fiber at $|\Omega(x)\rangle$: $\mathcal{E}_{\mathrm{exc}}(x) \cong T_{|\Omega(x)\rangle}^*\mathcal{M}_{\mathrm{vac}}$ via the canonical one–form pairing.*

Then there exists a canonical, fiberwise isometric isomorphism

$$\Psi_x : \ \mathcal{E}_{\mathrm{exc}}(x) \ \xrightarrow{\cong}\ T_{|\Omega(x)\rangle}^*\mathcal{M}_{\mathrm{vac}},$$

natural in x, commuting with the typed/triadic projectors on the $\Phi\Phi$ sector, $[\Psi_x, \Pi^{[\chi]}_{\Phi\Phi}] = [\Psi_x, \Pi_\alpha] = 0$, and unique up to unitaries within $\mathcal{E}_{\mathrm{exc}}(x)$ that commute with $\hat{P}_T(x)$ and with the typed/triadic projectors. Consequently,

$$\mathcal{E}_{\mathrm{exc}}(x) \;\simeq\; T^*_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}, \tag{9.123}$$

and the Tantrakyma field $\hat{\boldsymbol{\Theta}}(x)$ defines a covector on $\mathcal{M}_{\mathrm{vac}}$ at $|\Omega(x)\rangle$.

Remark 9.293 (Kinetic–symplectic meaning of A_x and compatibility of the two identifications)**.** In case (B), the operator $\mathsf{A}_x : T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}} \to T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}$ is the fiberwise *kinetic metric/inertia* associated to the diagnostic realized reference configuration Lagrangian

$$\mathcal{L}_{\Phi,\mathrm{kin}} = \tfrac{1}{2}\, g_T^{(x)}\Big(\dot{\Phi},\, \mathsf{A}_x\dot{\Phi}\Big),$$

hence positive, boundedly invertible, and $g_T^{(x)}$–self–adjoint. The Legendre/Riesz map

$$\mathcal{R}^{(\mathsf{A})}_x : \; v \longmapsto \Big[w \mapsto g_T^{(x)}(\mathsf{A}_x v, w)\Big]$$

is therefore an isometric isomorphism $T_{|\Omega}\mathcal{M}_{\mathrm{vac}} \xrightarrow{\cong} T^*_{|\Omega}\mathcal{M}_{\mathrm{vac}}$, and the momentum fiber at $|\Omega(x)\rangle$ identifies canonically with $\mathcal{E}_{\mathrm{exc}}(x)$, cf. Definitions 6.104 and 9.289, Remark 9.290, and Proposition 9.292. If, in addition, the Kähler structure in (A) is available and commutes with the typed/triadic projectors, then there exists a unitary $U_x : \mathcal{E}_{\mathrm{exc}}(x) \to \mathcal{E}_{\mathrm{exc}}(x)$ commuting with $\hat{P}_T(x)$, $\Pi^{[\chi]}_{\Phi\Phi}$ and Π_α such that

$$\Psi^{(\mathrm{A})}_x \;=\; \Psi^{(\mathrm{B})}_x \circ U_x.$$

Hence the two constructions agree up to the intrinsic ambiguity already recorded in Proposition 9.292. In diagnostic realized reference configuration–normal coordinates for which $g_T^{(x)} = \mathbf{1}$ and the projectors are diagonal, one has the coordinate-level Legendre pairing $p_i = (\mathsf{A}_x)_{ij}\,\dot{\phi}^j$, while the Kähler construction reads $\Psi_x(u)[v] = g^{(x)}_{\mathrm{mod}}(u, J^{(x)}v)$ with $J^{(x)}(T_{|\Omega}\mathcal{M}_{\mathrm{vac}}) = \mathcal{E}_{\mathrm{exc}}(x)$; these coincide precisely when A_x matches the $J^{(x)}$–induced kinetic metric on $T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}$.

Proof. We give two sufficient constructions of the canonical isomorphism $\Psi_x : \mathcal{E}_{\mathrm{exc}}(x) \xrightarrow{\cong} T^*_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}$, each compatible with the diagnostic realized reference configuration split of Definition 9.289 and Remark 9.290. Throughout, $g_T^{(x)}$ denotes the inner product on $T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}$ induced from $g^{(x)}_{\mathrm{mod}}$, and $\mathcal{R}_x : T_{|\Omega}\mathcal{M}_{\mathrm{vac}} \to T^*_{|\Omega}\mathcal{M}_{\mathrm{vac}}$ is the fiberwise Riesz isomorphism (Definition 6.104; cf. [253]).

Preliminaries (orthogonal split). By Definition 9.289 and Remark 9.290, $\mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}} = \mathcal{E}_{\mathrm{geo}}(x) \oplus \mathcal{E}_{\mathrm{exc}}(x)$ with $\mathcal{E}_{\mathrm{geo}}(x) = T_{|\Omega}\mathcal{M}_{\mathrm{vac}}$ and $\mathcal{E}_{\mathrm{exc}}(x) = \big(T_{|\Omega}\mathcal{M}_{\mathrm{vac}}\big)^{\perp}$. Typed/triadic compatibility holds on the diagnostic realized reference configuration: $[\hat{P}_T(x), \Pi^{[\chi]}_{\Phi\Phi}] = [\hat{P}_T(x), \Pi_\alpha] = 0$.

Construction (A): Kähler duality. Assume the existence of a Kähler triple $\big(g^{(x)}_{\mathrm{mod}}, J^{(x)}, \omega^{(x)}\big)$ satisfying the following conditions:

$$J^{(x)}(T_{|\Omega}\mathcal{M}_{\mathrm{vac}}) = \mathcal{E}_{\mathrm{exc}}(x),$$
$$\omega^{(x)}(\cdot,\cdot) = g^{(x)}_{\mathrm{mod}}(\cdot, J^{(x)}\cdot),$$
$$[J^{(x)}, \Pi^{[\chi]}_{\Phi\Phi}] = [J^{(x)}, \Pi_\alpha] = 0.$$

Define

$$\Psi_x(u)[v] \;:=\; g^{(x)}_{\mathrm{mod}}\Big(u,\; J^{(x)}v\Big), \qquad u \in \mathcal{E}_{\mathrm{exc}}(x), \quad v \in T_{|\Omega}\mathcal{M}_{\mathrm{vac}}.$$

Because $J^{(x)}$ is an isometry and $J^{(x)}(T) = \mathcal{E}_{\mathrm{exc}}$, we have $\|\Psi_x(u)\| = \sup_{\|v\|=1} |g^{(x)}_{\mathrm{mod}}(u, J^{(x)}v)| = \|u\|$, so Ψ_x is a fiberwise isometry onto $T^*_{|\Omega}\mathcal{M}_{\mathrm{vac}}$. Commutation with the typed/triadic projectors follows from the stated compatibility.

Construction (B): Phase space / Legendre pairing. Assume a positive, boundedly invertible A_x with $\mathcal{L}_{\Phi,\mathrm{kin}} = \frac{1}{2}\, g^{(x)}_T(\dot\Phi, \mathsf{A}_x\dot\Phi)$ and $[\mathsf{A}_x, \Pi^{[\chi]}_{\Phi\Phi}] = [\mathsf{A}_x, \Pi_\alpha] = 0$. Then

$$\mathcal{R}^{(\mathsf{A})}_x :\; T_{|\Omega}\mathcal{M}_{\mathrm{vac}} \;\to\; T^*_{|\Omega}\mathcal{M}_{\mathrm{vac}}, \quad v \;\mapsto\; \Big[w \mapsto g^{(x)}_T(\mathsf{A}_x v,\, w)\Big]$$

is an isometric isomorphism. Using the IQR phase–space identification at $|\Omega(x)\rangle$, $\mathcal{E}_{\mathrm{exc}}(x)$ is the momentum fiber canonically paired with $T_{|\Omega}\mathcal{M}_{\mathrm{vac}}$ by the one–form ϑ_x. Thus define

$$\Psi_x :\; \mathcal{E}_{\mathrm{exc}}(x) \;\to\; T^*_{|\Omega}\mathcal{M}_{\mathrm{vac}}, \qquad u \;\mapsto\; \vartheta_x(u,\cdot).$$

With the Legendre metric $g^{(x)}_T$ and A_x, this pairing is an isometry; commutation with the projectors follows from the diagnostic realized reference configuration compatibility.

Uniqueness. If $\tilde\Psi_x$ is another fiberwise isometry with the same commutation properties, then $U_x := \tilde\Psi_x^{-1} \circ \Psi_x : \mathcal{E}_{\mathrm{exc}}(x) \to \mathcal{E}_{\mathrm{exc}}(x)$ is unitary and commutes with $\hat P_T(x)$, $\Pi^{[\chi]}_{\Phi\Phi}$, and Π_α. Hence Ψ_x is unique up to such commuting unitaries.
This proves (9.123) and the properties claimed in Proposition 9.292. □

Remark 9.294 (Canonical identification under Kähler structure)**.** *Optional structure (makes* (9.123) *canonical).* Suppose the local fiber $\mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}}$ carries a Kähler triple $\big(g^{(x)}_{\mathrm{mod}}, J^{(x)}, \omega^{(x)}\big)$ with

$$J^{(x)}\Big(T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}\Big) = \mathcal{E}_{\mathrm{exc}}(x), \qquad \omega^{(x)}(\cdot,\cdot) = g^{(x)}_{\mathrm{mod}}\Big(\cdot, J^{(x)}\,\cdot\,\Big),$$

and assume $J^{(x)}$ is compatible with the diagnostic realized reference configuration typed/triadic projectors, $[J^{(x)}, \Pi^{[\chi]}_{\Phi\Phi}] = [J^{(x)}, \Pi_\alpha] = 0$ (cf. Definitions 9.186 and 9.200 and Remark 9.290). Let $g^{(x)}_T$ denote the metric on $T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}$ induced from $g^{(x)}_{\mathrm{mod}}$. Then the fiberwise map

$$\Psi_x :\; \mathcal{E}_{\mathrm{exc}}(x) \;\longrightarrow\; T^*_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}, \qquad u \;\longmapsto\; \Big[v \mapsto g^{(x)}_{\mathrm{mod}}\Big(u,\; J^{(x)}v\Big)\Big],$$

is an isometric isomorphism (with the dual norm induced by $g^{(x)}_T$) that *commutes* with the typed/triadic projectors: $[\,\Psi_x, \Pi^{[\chi]}_{\Phi\Phi}\,] = [\,\Psi_x, \Pi_\alpha\,] = 0$. Hence (9.123) holds canonically in the sense used in Proposition 9.292.

Smoothness and parallel transport. If $x \mapsto \left(g^{(x)}_{\mathrm{mod}}, J^{(x)}\right)$ is C^1 on the diagnostic realized reference configuration, then $x \mapsto \Psi_x$ defines a smooth bundle isomorphism $\mathcal{E}_{\mathrm{exc}} \xrightarrow{\cong} T^*\mathcal{M}_{\mathrm{vac}}$. If moreover $\bar{\nabla}_\mu J^{(x)} = 0$ (with $\bar{\nabla}$ from Definition 6.7), the identification is parallel along $\mathcal{X}_q$. *Uniqueness up to commuting unitaries.* Any other such identification $\tilde{\Psi}_x$ differs by a unitary U_x on $\mathcal{E}_{\mathrm{exc}}(x)$ that commutes with $\hat{P}_T(x)$ and with the typed/triadic projectors; thus the canonical class is unique up to these innocuous internal symmetries.

9.17.11 Tantrakyma Parasymmetric Type, Location, and Onset

This section sets the operational stage for applying the parasymmetric *Type–Location–Onset* (T–L–O) program to the *Tantrakyma* (matter) sector. It mirrors the Gaiamiti workflow (Section 9.17) and the projection/duality pass (Section 9.17.10), but is specialized to the typed subspaces of the Tantrakyma field. All computations are performed on a certified plateau $\mathfrak{P}$ with UV/IR tags (Definition 7.26, Section 6.12.4, and eq. (7.28)), the triadic channel split $\alpha \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$ (Definition 5.54 and Theorem 6.261), and BRST/Nielsen safeguards (Chs. 7–8; cf. Sections 7.2 and 7.4). The explicit law for Tantrakyma is introduced in Section 9.18.

Arena, law placeholders, and diagnostic realized reference configuration discipline

Diagnostic realized reference configuration and tags. Fix a realized, BRST-clean diagnostic realized reference configuration $\mathfrak{P}$ (and $\mathfrak{P}_\partial$ when a boundary arena is used), equipped with band tags $\{P_x(\lambda_{\mathrm{UV}}),\ K^{(\ell)}_{\mathrm{IR}}\}$ and the physical projector $\mathsf{P}_{\mathrm{phys}} : \mathcal{H} \to H^0(\hat{Q}_{\mathrm{BRST}})$ (Definition 7.26 and Sections 6.12.4 and 7.2). All meters/witnesses are evaluated *on band* and *on* H^0 (Nielsen-flatness).

Law placeholders. We forward-reference the Tantrakyma action S_Θ and Euler–Lagrange (EL) equation to Section 9.18 (see Definitions 9.305 and 9.310). Until then, we treat the kinetic form via the canonical covariant operator g_C and the interaction via the potential $\mathcal{V}_\Theta$ with expectation $V_\Theta := \langle \mathcal{V}_\Theta \rangle$ (cf. eqs. (9.133) and (9.134)).

Triadic compatibility. All typed operations commute with the triadic channel projectors $\{\Pi_\alpha\}$ (holosymmetric guardrails; Ch. 7). Unless stated otherwise, every object carries the band/triad decorations implicitly.

Typed slicing of the Tantrakyma sector

Definition 9.295 (Typed Tantrakyma projectors)**.** Let the admissible type menu be $[\chi] \in \{\mathrm{CK},\ \mathrm{T}_+,\ \mathrm{T}_-,\ \mathrm{L}\}$, corresponding to Chrono–Kymon, transverse-helicity ($\pm$), and longitudinal subspaces (cf. fields–quanta typed ledgers). Define orthogonal, complete, BRST-closed projectors

$$\Pi^{[\chi]}_{\Theta\Theta} :\ \mathcal{H} \to \mathcal{H}, \qquad \sum_\chi \Pi^{[\chi]}_{\Theta\Theta} = \mathbf{1}, \quad \Pi^{[\chi]}_{\Theta\Theta}\,\Pi^{[\chi']}_{\Theta\Theta} = \delta_{\chi\chi'}\,\Pi^{[\chi]}_{\Theta\Theta}, \quad \left[\hat{Q}_{\mathrm{BRST}},\ \Pi^{[\chi]}_{\Theta\Theta}\right] = 0,$$

with $\left[\Pi^{[\chi]}_{\Theta\Theta},\ \Pi_\alpha\right] = 0$ for each triadic channel α.

Linearization and typed mass/curvature operators

Write the background–fluctuation split $\hat{\boldsymbol{\Theta}} = \bar{\Theta}\,\mathbb{I} + \delta\hat{\boldsymbol{\Theta}}$ on $\mathfrak{P}$ (EL-stationary backgrounds in Section 9.18). The potential's same-variable Hessian defines typed curvature operators by restriction:

$$\boxed{\mathcal{K}^{[\chi]}_{\Theta\Theta} := \Pi^{[\chi]}_{\Theta\Theta}\,\frac{\delta^2\mathcal{V}_\Theta}{\delta\hat{\boldsymbol{\Theta}}\,\delta\hat{\boldsymbol{\Theta}}}\Big|_{\bar{\Theta}}\,\Pi^{[\chi]}_{\Theta\Theta}, \qquad \mathrm{spec}\Big(\mathcal{K}^{[\chi]}_{\Theta\Theta}\Big) \rightsquigarrow \text{ gaps vs. NG/pNG modes.}} \tag{9.124}$$

Typed EL linear response then reads

$$\begin{aligned}\Big(\mathbb{K}^{[\chi]}_{\Theta} + \mathcal{K}^{[\chi]}_{\Theta\Theta}\Big)\,\delta\hat{\boldsymbol{\Theta}}^{[\chi]} &= \hat{\mathcal{J}}^{[\chi]},\\ \delta\hat{\boldsymbol{\Theta}}^{[\chi]} &:= \Pi^{[\chi]}_{\Theta\Theta}\,\delta\hat{\boldsymbol{\Theta}},\end{aligned} \tag{9.125}$$

where $\mathbb{K}^{[\chi]}_{\Theta}$ is the typed kinetic block (band-limited, hyperbolic on H^0; Ch. 8) and $\hat{\mathcal{J}}^{[\chi]}$ denotes the typed source.

Parasymmetric meters and witnesses (typed)

Definition 9.296 (Typed parasymmetric meters and witness)**.** Let $\Delta^{[\chi]}_{\Theta\Theta}$ denote the typed, band-limited, retarded two-point kernel on H^0 (*notation:* this is the $[\chi]$ projection of $G^{\mathrm{R}}_{\Theta\Theta,\mathrm{phys}}$). Define the SynTh driver triple

$$\boxed{\begin{aligned} v^{(L)}_{[\chi]} &:= \Big|\,\mathrm{Res}_{k^2=0}\;\Delta^{[\chi]}_{\Theta\Theta}(k)\Big| \quad \text{(law-side pole content)}, && (9.126)\\ v^{(S)}_{[\chi]} &:= \Big\|\Delta^{[\chi]}_{\Theta\Theta}(t\to\infty;\,\mathbf{x}=\mathbf{0})\Big\|_{H^0} \quad \text{(state-side long-time order)}, && (9.127)\\ v^{(LS)}_{[\chi]} &:= \text{typed cross-channel gain (helicity-resolved; band-robust).} && (9.128)\end{aligned}}$$

The *typed witness* is $\mathbf{W}^{[\Theta\Theta]}_{[\chi]} := (v^{(L)}_{[\chi]},\, v^{(S)}_{[\chi]},\, v^{(LS)}_{[\chi]})$.

Tolerance semantics and plateau certification follow eq. (7.28). Gauge-parameter independence of $\mathbf{W}^{[\Theta\Theta]}_{[\chi]}$ holds by the BRST/Nielsen provisos (Chs. 7–8; ∂_ξ-flatness).

Typed quadrant predicate and healing indicators

Definition 9.297 (Typed quadrant predicate for Tantrakyma)**.** Given $\mathbf{W}^{[\Theta\Theta]}_{[\chi]}$ on a certified $\mathfrak{P}$, define $\mathrm{Quad}^{[\Theta\Theta]}_{[\chi]} \in \{\mathrm{QI}, \mathrm{QII}, \mathrm{QIII}, \mathrm{QIV}\}$ by the canonical decision rule of Definition 7.685 (as typed in Definition 9.208). Introduce *healing indicators* $\mathscr{H}_L, \mathscr{H}_S$ (law/state) to distinguish QIII vs. QIV (finite-window decay vs. pinned deficit).

Proposition 9.298 (Meters determine the typed quadrant)**.** *On a certified plateau and fixed tags, the triple* $\big(\mathbf{W}^{[\Theta\Theta]}_{[\chi]},\, \mathscr{H}_L,\, \mathscr{H}_S\big)$ *uniquely determines* $\mathrm{Quad}^{[\Theta\Theta]}_{[\chi]}$ *(SynTh completeness; cf. Corollary 9.209). Consistency with the potential analysis holds in the sense that NG/pNG vs. gapped predictions from* $\mathrm{spec}(\mathcal{K}^{[\chi]}_{\Theta\Theta})$ *agree within tolerance budgets (Ch. 8).*

Type–Location–Onset stamping

Definition 9.299 (Typed T–L–O stamping)**.** For each $[\chi]$, the $\mathrm{TLO}^{[\chi]}$ record contains (i) the *Type* identifier; (ii) *Location* bands (IR/mesoscopic/UV windows under $K_{\mathrm{IR}}^{(\ell)}$); (iii) *Onset* thresholds (first certified pass-through). Stamps are computed from $\mathbf{W}_{[\chi]}^{[\Theta\Theta]}$ and the typed retarded kernels, using the same operators defined in the Gaiamiti prelude (Section 9.17).

Boundary pass and monotonicity

Proposition 9.300 (Bulk–boundary overlay)**.** *Repeating the evaluation on $\mathfrak{P}_\partial$ with tangential windows preserves BRST/Nielsen flatness; plateau budgets may tighten, and law visibilities weakly increase toward symmetry (boundary functoriality, cf. Ch. 9 sector rules). Typed quadrant membership is stable away from tolerance boundaries.*

Noether and stress–energy cross-checks

From $\mathcal{L}_\Theta$ (introduced in Section 9.18), construct the internal *Noether current* $\hat{J}_\Theta^\mu$ and its diagnostic realized reference configuration realization via the Noether bridge (Lemma 9.279, Definition 9.280, and Proposition 9.281). Optionally, derive the Tantrakyma stress–energy tensor $\hat{T}_{(\Theta)}^{\mu\nu}$ by diffeomorphism response (geometry-analogue of Section 9.17.9); both provide additional *Location* diagnostics consistent with the typed meters.

9.18 The Action for the Dynamical Tantrakyma Field

Having introduced the Dynamical Tantrakyma Field $\hat{\mathbf{\Theta}}(x)$ as the fundamental operator encoding material structure—i.e., *the relational manifestation of excitation and content* within the emergent geometry—we now specify its dynamics via an action principle. As with the Gaiamiti field, we invoke the *Principle of Stationary Action* to derive its evolution. Throughout, all objects are evaluated on a certified plateau $\mathfrak{P}$ with UV/IR tags $(\lambda_{\mathrm{UV}}, \ell)$ (cf. Definition 7.26, Section 6.12.4, and eq. (7.28)) and projected to the physical sector $H^0(\hat{Q}_{\mathrm{BRST}})$; ghost number is zero and gauge-parameter dependence is Nielsen-flat.

Definition 9.301 (Operator kernel and scalar Lagrangian density (unbanded))**.** Let $\hat{\mathbf{\Theta}}(x)$ be the Dynamical Tantrakyma Field, viewed as a section of $\mathrm{End}(\mathcal{H}_{\mathrm{em}}) \otimes T^*\mathcal{M}_{\mathrm{vac}}$ over $(\mathcal{X}_q, \bar{\Sigma}_{\mu\nu})$ (cf. Definitions 1.34 and 5.26 and Section 6.1.17). Define the *operator-valued* Lagrangian *kernel*

$$\widehat{\mathcal{L}}_\Theta(x) := \bar{\Sigma}^{\mu\nu}(x)\, g_C\Big(\nabla_\mu \hat{\mathbf{\Theta}}(x),\, \nabla_\nu \hat{\mathbf{\Theta}}(x)\Big) \;+\; \mathcal{V}_\Theta\Big(\hat{\mathbf{\Theta}}(x)\Big) \quad \in\; \mathrm{End}(\mathcal{H}_{\mathrm{em}}),$$

with ∇_μ the total (base + internal) covariant derivative and g_C a Hermitian operator-valued co-metric (Remark 9.303). The *(scalar) Lagrangian density* is the trace of this kernel:

$$\mathcal{L}_\Theta(x) \;:=\; \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\Big\{\widehat{\mathcal{L}}_\Theta(x)\Big\}, \tag{9.129}$$

a c-number density on $\mathcal{X}_q$ (ghost number 0).

The Lagrangian density $\mathcal{L}_\Theta(x)$ decomposes into:

1. **Kinetic term:**
$$\mathcal{T}_\Theta(x) := \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\Big\{\bar{\Sigma}^{\mu\nu}(x)\, g_C\Big(\nabla_\mu \hat{\mathbf{\Theta}}(x),\, \nabla_\nu \hat{\mathbf{\Theta}}(x)\Big)\Big\}. \tag{9.130}$$

2. **Potential term (operator built, traced to scalar):**
$$\mathcal{V}_{\Theta,\mathrm{density}}(x) := \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\Big\{\mathcal{V}_\Theta\Big(\hat{\mathbf{\Theta}}(x)\Big)\Big\}. \tag{9.131}$$

Thus $\mathcal{L}_\Theta = \mathcal{T}_\Theta + \mathcal{V}_{\Theta,\mathrm{density}}$.
The total action functional is

$$S_\Theta[\hat{\mathbf{\Theta}}] \;=\; \int_{\mathcal{X}_q} d\mu(x)\, \mathcal{L}_\Theta(x) \;=\; \int_{\mathcal{X}_q} d^d x\, \sqrt{-\det \bar{\Sigma}(x)}\, \mathcal{L}_\Theta(x), \tag{9.132}$$

where $d\mu(x) = d^d x \sqrt{-\det \bar{g}^{\mathrm{Lor}}(x)}$ is the relational volume element (cf. Definition 6.33; the shorthand $\sqrt{-\det \bar{\Sigma}}$ equals $\sqrt{-\det \bar{g}^{\mathrm{Lor}}}$ on the diagnostic realized reference configuration, cf. Remark 9.311). Equivalently,

$$S_\Theta \;=\; \int_{\mathcal{X}_q} d\mu(x)\; \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\Big\{\widehat{\mathcal{L}}_\Theta(x)\Big\}.$$

Remark 9.302 (Symmetry, covariance, and diagnostic realized reference configuration/BRST invariances)**.** The Lagrangian density $\mathcal{L}_\Theta(x)$ in Definition 9.301 is:

- **Diffeomorphism covariant** on $\mathcal{X}_q$: $\mathcal{L}_\Theta(x)$ is a scalar density under coordinate changes, and S_Θ is invariant.
- **Fiber–algebra invariance:** If $\varphi_x \in \mathrm{Aut}\Big(\mathrm{End}(\mathcal{H}_{\mathrm{em}})\Big)$ is trace-preserving (e.g. $\varphi_x(A) = U_x A U_x^\dagger$ with U_x unitary), then $\mathrm{Tr}(\varphi_x(X)) = \mathrm{Tr}(X)$, so $\mathcal{L}_\Theta$ and S_Θ are invariant.
- **Diagnostic realized reference configuration inheritance**: after restriction to a certified plateau $\mathfrak{P}$ with tags $(\lambda_{\mathrm{UV}}, \ell)$ and projection to $H^0(\hat{Q}_{\mathrm{BRST}})$, these symmetries persist; banding and projection are BRST-closed (ghost number 0).
- **Nielsen stability**: pole locations and residues extracted from typed, band-limited propagators on H^0 are gauge-parameter independent (Chs. 7–8).

Thus the field equations from $\delta S_\Theta = 0$ respect background-independence and the triadic/typed machinery used by the parasymmetric T–L–O program.

Remark 9.303 (Operator-valued co-metric and (pseudo)inverse)**.** The map $g_C : T^*\mathcal{M}_{\mathrm{vac}} \times T^*\mathcal{M}_{\mathrm{vac}} \to \mathrm{End}(\mathcal{H}_{\mathrm{em}})$ is Hermitian, positive semidefinite, and sesquilinear (antilinear in its first entry). For $\hat{\alpha}, \hat{\beta} \in T^*_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}$ we have

$$g_C(\hat{\alpha}, \hat{\beta})^\dagger = g_C(\hat{\beta}, \hat{\alpha}), \qquad g_C(\hat{\alpha}, \hat{\alpha}) \succeq 0.$$

- **Kinetic pairing.** g_C acts on the cotangent legs and, by $\mathrm{End}(\mathcal{H}_{\mathrm{em}})$-bilinearity, yields an element of $\mathrm{End}(\mathcal{H}_{\mathrm{em}})$ inside the trace in (9.130).

- **Pseudoinverse on the excitation subspace.** Let g_C^{-1} denote the Moore–Penrose pseudoinverse on the excitation subspace $\mathcal{E}_{\mathrm{exc}}(x) \subseteq T^*\mathcal{M}_{\mathrm{vac}} \otimes \mathrm{End}(\mathcal{H}_{\mathrm{em}})$, characterized by

$$g_C\, g_C^{-1}\, g_C = g_C, \quad g_C^{-1}\, g_C\, g_C^{-1} = g_C^{-1}, \quad \big(g_C\, g_C^{-1}\big)^\dagger = g_C\, g_C^{-1}, \quad \big(g_C^{-1}\, g_C\big)^\dagger = g_C^{-1}\, g_C.$$

  For $\hat\alpha, \hat\beta \in \mathcal{E}_{\mathrm{exc}}(x)$ the *trace pairing* holds:

$$\mathrm{Tr}\Big\{\, g_C\Big(g_C^{-1}(\hat\alpha, \cdot),\, \hat\beta\Big)\Big\} = \mathrm{Tr}\Big(\hat\alpha^\dagger \hat\beta\Big)\,.$$

  This duality is useful for Hamiltonian/phase-space formulations but is *not* the primitive building block of the potential.

- **Relational dependence and covariance.** g_C (and g_C^{-1}, when used) vary over $\mathcal{X}_q$ via $|\Omega(x)\rangle$, encoding an operator-theoretic geometry of $T^*\mathcal{M}_{\mathrm{vac}}$ (cf. Proposition 9.292); their band-limited pullbacks are BRST-closed and Nielsen-flat on H^0.

Together, $\big(g_C, g_C^{-1}\big)$ provide the inner-product structure and (optional) tangent–cotangent duality for the Tantrakyma field.

Definition 9.304 (Tantrakyma potential: operator form and expectation)**.** Because $\hat{\mathbf{\Theta}}$ is *covector*-valued, the canonical invariant is $\mathsf{X} := g_C\big(\hat{\mathbf{\Theta}}, \hat{\mathbf{\Theta}}\big) \in \mathrm{End}(\mathcal{H}_{\mathrm{em}})$. A diagnostic realized reference configuration-safe minimal potential is

$$\boxed{\mathcal{V}_\Theta\big(\hat{\mathbf{\Theta}}\big) = \frac{1}{2}\, m_\Theta^2\, \mathsf{X} + \frac{\lambda_\Theta}{4!}\, \mathsf{X}^2 + \mathcal{O}\big(\mathsf{X}^3/\Lambda_\Theta^2\big),} \tag{9.133}$$

with real parameters m_Θ^2 and $\lambda_\Theta > 0$, and a suppression scale Λ_Θ. Its expectation (a c-number density) is

$$\boxed{V_\Theta(x) := \Big\langle \mathring{\mathbf{\Psi}}\Big|\, \mathsf{P}_{\mathrm{phys}}\, \mathcal{V}_\Theta\Big(\hat{\mathbf{\Theta}}(x)\Big)\, \mathsf{P}_{\mathrm{phys}}\, \Big|\mathring{\mathbf{\Psi}}\Big\rangle = \mathrm{Tr}\Big\{\rho_{\mathrm{phys}}(x)\, \mathcal{V}_\Theta\Big(\hat{\mathbf{\Theta}}(x)\Big)\Big\}\,,} \tag{9.134}$$

with ρ_{phys} the local reduced density on H^0. (Any diagnostic realized reference configuration-certified alternative potential compatible with $\mathsf{X} = g_C(\hat{\mathbf{\Theta}}, \hat{\mathbf{\Theta}})$ may be used.)

Definition 9.305 (Euler–Lagrange equation for $\hat{\mathbf{\Theta}}(x)$)**.** Varying S_Θ at fixed $\bar\Sigma$ and connections yields the operator EL equation, in manifestly covariant form,

$$\boxed{\frac{1}{\sqrt{-\det \bar\Sigma}}\, \nabla_\mu\left(\sqrt{-\det \bar\Sigma}\; \bar\Sigma^{\mu\nu}\, \frac{\partial \mathcal{L}_\Theta}{\partial(\nabla_\nu \hat{\mathbf{\Theta}})}\right) \;=\; \frac{\delta \mathcal{V}_\Theta}{\delta \hat{\mathbf{\Theta}}^\dagger}\,,} \tag{9.135}$$

where, by Fréchet differentiation at fixed $\bar\Sigma$,

$$\frac{\partial \mathcal{L}_\Theta}{\partial(\nabla_\nu \hat{\mathbf{\Theta}})} \;=\; \bar\Sigma^{\mu\nu}\, g_C^\sharp\Big(\nabla_\mu \hat{\mathbf{\Theta}}, \cdot\Big),$$

with $g_C^\sharp$ the map sending the second slot "$\cdot$" to $g_C\big(\nabla_\mu \hat{\mathbf{\Theta}}, \cdot\big) \in \mathrm{End}(\mathcal{H}_{\mathrm{em}})$. Expanding the covariant divergence gives the equivalent form

$$\boxed{\bar{\Sigma}^{\mu\nu}\,\nabla_\mu \nabla_\nu \hat{\mathbf{\Theta}} \;+\; (\nabla_\mu \bar{\Sigma}^{\mu\nu})\,\nabla_\nu \hat{\mathbf{\Theta}} \;=\; \frac{\delta \mathcal{V}_\Theta}{\delta \hat{\mathbf{\Theta}}^\dagger}\,.} \tag{9.136}$$

No g_C^{-1} appears: the kinetic and potential terms are built with the co-metric g_C appropriate to the covector-valued field $\hat{\mathbf{\Theta}}$. Both forms are read on the diagnostic realized reference configuration with band tags and BRST projection, in parallel with Definition 9.268.

Definition 9.306 (Band functor and tags)**.** Let $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ be the UV projector and IR window (cf. Definition 7.26 and Section 6.12.4). Define the band functor $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ acting on fields and derivatives by

$$\hat{\mathbf{\Theta}}^{(\lambda_{\mathrm{UV}},\ell)} := \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}\Big[\hat{\mathbf{\Theta}}\Big]\,, \qquad \nabla_\mu^{(\lambda_{\mathrm{UV}},\ell)} := \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell} \circ \nabla_\mu,$$

and work on a certified plateau $\mathfrak{P} \subset \mathcal{X}_q$ with projection to $H^0(\hat{Q}_{\mathrm{BRST}})$.

Definition 9.307 (Band-limited, BRST-clean operator-built Lagrangian density)**.** Work on a certified plateau $\mathfrak{P}$ (or $\mathfrak{P}_\partial$) with projection to the physical Hilbert space

$$\mathcal{H}_{\mathrm{phys}} \;=\; H^0(\hat{Q}_{\mathrm{BRST}}),$$

and triadic channel projectors $\{\Pi_\alpha\}_{\alpha\in\{\mathrm{corr,vac,spec}\}}$ (Definition 9.185). Let the *Dynamical Tantrakyma Field* be a section

$$\hat{\mathbf{\Theta}}(x) \in \mathrm{End}(\mathcal{H}_{\mathrm{em}}) \otimes T^*\mathcal{M}_{\mathrm{vac}},$$

over the emergent configuration space $(\mathcal{X}_q, \bar{\Sigma}_{\mu\nu})$ (cf. Definitions 1.34 and 5.26 and Section 6.1.17). Let $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ denote the UV projector and IR window (cf. Definition 7.26 and Section 6.12.4), and define the band functor

$$\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell} \;:=\; P_x(\lambda_{\mathrm{UV}})\, K_{\mathrm{IR}}^{(\ell)}.$$

The band functor is BRST- and triadic-compatible:

$$[\hat{Q}_{\mathrm{BRST}}, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}] = 0, \qquad [\Pi_\alpha, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}] = 0 \quad (\alpha \in \{\mathrm{corr,vac,spec}\}), \tag{9.137}$$

and, when used with typed projectors on the $\Theta\Theta$ block, preserves typed compatibility $\Big[\Pi_{\Theta\Theta}^{[\chi]}, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}\Big] = 0$ (cf. Definition 9.295 and Proposition 9.202).

Define the band-limited field and covariant derivative by

$$\hat{\mathbf{\Theta}}^{(\lambda_{\mathrm{UV}},\ell)} := \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}\Big[\hat{\mathbf{\Theta}}\Big]\,, \tag{9.138}$$

$$\nabla_\mu^{(\lambda_{\mathrm{UV}},\ell)} := \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell} \circ \nabla_\mu, \tag{9.139}$$

and enforce the filter/derivative composition on banded fields as

$$\nabla_\mu^{(\lambda_{\mathrm{UV}},\ell)} \hat{\mathbf{\Theta}}^{(\lambda_{\mathrm{UV}},\ell)} = \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}\Big[\nabla_\mu\Big(\hat{\mathbf{\Theta}}^{(\lambda_{\mathrm{UV}},\ell)}\Big)\Big] = \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}\Big[\nabla_\mu\, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\hat{\mathbf{\Theta}}]\Big], \tag{9.140}$$

so no out-of-band leakage appears at this stage. Here ∇_μ is the total (base + internal) covariant derivative compatible with $\bar{\Sigma}$ (Definitions 6.7 and 9.301).

The (band-limited, BRST-clean) operator-built Lagrangian density is

$$\mathcal{L}_\Theta^{(\lambda_{\mathrm{UV}},\ell)}(x) := \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\Big\{\bar{\Sigma}^{\mu\nu}(x)\, g_C\Big(\nabla_\mu^{(\lambda_{\mathrm{UV}},\ell)}\hat{\boldsymbol{\Theta}}^{(\lambda_{\mathrm{UV}},\ell)}(x),\, \nabla_\nu^{(\lambda_{\mathrm{UV}},\ell)}\hat{\boldsymbol{\Theta}}^{(\lambda_{\mathrm{UV}},\ell)}(x)\Big) + \mathcal{V}_\Theta\Big(\hat{\boldsymbol{\Theta}}^{(\lambda_{\mathrm{UV}},\ell)}(x)\Big)\Big\}, \tag{9.141}$$

where $g_C : T^*\mathcal{M}_{\mathrm{vac}} \times T^*\mathcal{M}_{\mathrm{vac}} \to \mathrm{End}(\mathcal{H}_{\mathrm{em}})$ is a Hermitian operator-valued co-metric (ghost number 0; cf. Remark 9.303) and $\mathcal{V}_\Theta(\cdot) \in \mathrm{End}(\mathcal{H}_{\mathrm{em}})$ is the Tantrakyma potential (cf. eqs. (9.133) and (9.134)).

The diagnostic realized reference configuration action uses the relational volume element $d\mu(x)$ (Definition 6.33; cf. Remark 9.311):

$$S_\Theta^{(\lambda_{\mathrm{UV}},\ell)}[\hat{\boldsymbol{\Theta}}] := \int_{\mathcal{X}_q} d\mu(x)\, \mathcal{L}_\Theta^{(\lambda_{\mathrm{UV}},\ell)}(x). \tag{9.142}$$

When tags are suppressed, write $\mathcal{L}_\Theta \equiv \mathcal{L}_\Theta^{(\lambda_{\mathrm{UV}},\ell)}$ and $S_\Theta \equiv S_\Theta^{(\lambda_{\mathrm{UV}},\ell)}$ at a fixed certified plateau $\mathfrak{P}$.

Proposition 9.308 (Band–variation commutation and typed EL on the diagnostic realized reference configuration)**.** *On a certified plateau (eq. (7.28)), variation and banding commute in the sense that*

$$\delta S_\Theta^{(\lambda_{\mathrm{UV}},\ell)} = \int d\mu\, \delta\mathcal{L}_\Theta^{(\lambda_{\mathrm{UV}},\ell)} = \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}\big[\delta S_\Theta\big].$$

Consequently, the Euler–Lagrange system obtained from $\delta S_\Theta^{(\lambda_{\mathrm{UV}},\ell)} = 0$ is the banded form of the unbanded EL equations (cf. Definition 9.305); its typed restriction obeys

$$\Big(\mathbb{K}_\Theta^{[\chi]} + \mathcal{K}_{\Theta\Theta}^{[\chi]}\Big)\,\delta\hat{\boldsymbol{\Theta}}^{[\chi]} = \hat{\mathcal{J}}^{[\chi]}, \qquad \delta\hat{\boldsymbol{\Theta}}^{[\chi]} := \Pi_{\Theta\Theta}^{[\chi]}\,\delta\hat{\boldsymbol{\Theta}}, \tag{9.143}$$

with $\mathcal{K}_{\Theta\Theta}^{[\chi]}$ the typed Hessian (eq. (9.124)). Pole locations and residues extracted on $H^0(\hat{Q}_{\mathrm{BRST}})$ are gauge-parameter independent (Nielsen-flat).

Proof of Proposition 9.308. **Setup.** On a certified plateau $\mathfrak{P}$ with tags $(\lambda_{\mathrm{UV}}, \ell)$ and BRST projection to $H^0(\hat{Q}_{\mathrm{BRST}})$, the band functor $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ acts on the Tantrakyma field and on the covariant derivative by

$$\hat{\boldsymbol{\Theta}}^{(\lambda_{\mathrm{UV}},\ell)} := \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\hat{\boldsymbol{\Theta}}], \qquad \nabla_\mu^{(\lambda_{\mathrm{UV}},\ell)} := \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell} \circ \nabla_\mu,$$

and the diagnostic realized reference configuration action reads (with the relational measure $d\mu$; cf. Definition 6.33 and Remark 9.311)

$$S_\Theta^{(\lambda_{\mathrm{UV}},\ell)} = \int_{\mathcal{X}_q} d\mu(x)\, \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\Big\{\bar{\Sigma}^{\mu\nu}(x)\, g_C\Big(\nabla_\mu^{(\lambda_{\mathrm{UV}},\ell)}\hat{\boldsymbol{\Theta}}^{(\lambda_{\mathrm{UV}},\ell)}, \nabla_\nu^{(\lambda_{\mathrm{UV}},\ell)}\hat{\boldsymbol{\Theta}}^{(\lambda_{\mathrm{UV}},\ell)}\Big) + \mathcal{V}_\Theta\Big(\hat{\boldsymbol{\Theta}}^{(\lambda_{\mathrm{UV}},\ell)}\Big)\Big\}.$$

By definition (Definition 9.307),

$$\nabla_\mu^{(\lambda_{\mathrm{UV}},\ell)}\hat{\mathbf{\Theta}}^{(\lambda_{\mathrm{UV}},\ell)} = \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}\Big[\nabla_\mu\Big(\hat{\mathbf{\Theta}}^{(\lambda_{\mathrm{UV}},\ell)}\Big)\Big] = \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}\Big[\nabla_\mu\,\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}[\hat{\mathbf{\Theta}}]\Big],$$

so no out-of-band leakage appears (cf. (9.140)). The background $\bar{\Sigma}$ is held fixed during variation. We use the diagnostic realized reference configuration commutation properties $[\hat{Q}_{\mathrm{BRST}}, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}] = 0$ and $[\Pi_\alpha, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}] = 0$ (eq. (9.137)), and assume standard vanishing/controlled boundary terms (or compact support under the IR window).

(A) δ–integral exchange. Since $d\mu$ is diagnostic realized reference configuration-fixed (vary at fixed $\bar{\Sigma}$), we may write

$$\delta S_\Theta^{(\lambda_{\mathrm{UV}},\ell)} = \int_{\mathcal{X}_q} d\mu(x)\; \delta\mathcal{L}_\Theta^{(\lambda_{\mathrm{UV}},\ell)}(x).$$

(B) Commutation of banding with variation. Linearity and idempotence of $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$, together with the fact that $\mathcal{L}_\Theta^{(\lambda_{\mathrm{UV}},\ell)}$ is the same functional as $\mathcal{L}_\Theta$ but evaluated on the *banded* variables, imply

$$\delta\mathcal{L}_\Theta^{(\lambda_{\mathrm{UV}},\ell)} = \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}\Big[\delta\mathcal{L}_\Theta\Big],$$

hence, using trace cyclicity,

$$\delta S_\Theta^{(\lambda_{\mathrm{UV}},\ell)} = \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}\Big[\delta S_\Theta\Big].$$

This is the claimed δ–banding commutation.

(C) Euler–Lagrange equations on the diagnostic realized reference configuration. Covariant integration by parts with the $\bar{\Sigma}$-compatible connection (background fixed) shows that the EL operator for the banded action is the band of the bare EL operator:

$$\mathcal{E}_{\mathbf{\Theta}}^{(\lambda_{\mathrm{UV}},\ell)} = \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}\Big[\mathcal{E}_{\mathbf{\Theta}}\Big].$$

Thus the EL equations from $\delta S_\Theta^{(\lambda_{\mathrm{UV}},\ell)} = 0$ are the banded form of the unbanded EL system (cf. Definition 9.305). Boundary terms vanish or are accounted for by the IR window.

(D) Typed restriction and linearized form. Let $\Pi_{\Theta\Theta}^{[\chi]}$ be the typed projector (cf. Definition 9.295 and Proposition 9.202) commuting with the triadic projectors on the diagnostic realized reference configuration. Linearizing the EL map and restricting to the typed Θ-sector yields

$$\Big(\mathbb{K}_\Theta^{[\chi]} + \mathcal{K}_{\Theta\Theta}^{[\chi]}\Big)\,\delta\hat{\mathbf{\Theta}}^{[\chi]} = \hat{\mathcal{J}}^{[\chi]}, \qquad \delta\hat{\mathbf{\Theta}}^{[\chi]} := \Pi_{\Theta\Theta}^{[\chi]}\,\delta\hat{\mathbf{\Theta}},$$

with $\mathcal{K}_{\Theta\Theta}^{[\chi]}$ as in eq. (9.124), i.e. (9.143).

(E) Nielsen flatness on $H^0(\hat{Q}_{\mathrm{BRST}})$. Adding gauge–fixing/ghost terms as BRST-exact pieces and projecting to $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ leaves the physical EL equations invariant and renders pole locations and residues gauge–parameter independent (Nielsen-flat), by the BRST/Nielsen safeguards of Chs. 7–8.

Conclusion. Combining (A)–(C) we obtain

$$\delta S_\Theta^{(\lambda_{\mathrm{UV}},\ell)} = \int d\mu\,\delta\mathcal{L}_\Theta^{(\lambda_{\mathrm{UV}},\ell)} = \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}\Big[\delta S_\Theta\Big],$$

and together with (D)–(E) the typed EL form (9.143) holds, with gauge–parameter independent poles/residues on $H^0(\hat{Q}_{\mathrm{BRST}})$. □

Definition 9.309 (Triadic- and typed-resolved Tantrakyma Lagrangian slices)**.** Let $\mathfrak{A} := \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$. Let $\{\Pi_\alpha\}_{\alpha\in\mathfrak{A}}$ be the triadic channel projectors (cf. Definition 9.185), and let $\{\Pi^{[\chi]}_{\Theta\Theta}\}$ denote the Tantrakyma typed projectors (cf. Definition 9.295). The *operator-level* typed/triadic slice is

$$\widehat{\mathcal{L}}^{[\chi]}_{\Theta;\alpha}(x) := \Pi_\alpha\, \Pi^{[\chi]}_{\Theta\Theta}\Big[\bar{\Sigma}^{\mu\nu}\, g_C\Big(\nabla_\mu\hat{\boldsymbol{\Theta}},\, \nabla_\nu\hat{\boldsymbol{\Theta}}\Big) + \mathcal{V}_\Theta(\hat{\boldsymbol{\Theta}})\Big]\Pi^{[\chi]}_{\Theta\Theta}\,\Pi_\alpha.$$

Since the projectors act trivially on the fiber algebra $\mathrm{End}(\mathcal{H}_{\mathrm{em}})$, the *channel- and type-resolved Lagrangian density* is obtained by tracing only over $\mathcal{H}_{\mathrm{em}}$:

$$\mathcal{L}^{[\chi]}_{\Theta;\alpha}(x) := \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\Big\{\widehat{\mathcal{L}}^{[\chi]}_{\Theta;\alpha}(x)\Big\}. \tag{9.144}$$

On a certified plateau $\mathfrak{P}$ with tags $(\lambda_{\mathrm{UV}}, \ell)$, the band-limited (BRST-clean) slices are

$$\begin{aligned}
\widehat{\mathcal{L}}^{[\chi](\lambda_{\mathrm{UV}},\ell)}_{\Theta;\alpha}(x) &:= \Pi_\alpha\, \Pi^{[\chi]}_{\Theta\Theta}\Big[\bar{\Sigma}^{\mu\nu}\, g_C\Big(\nabla^{(\lambda_{\mathrm{UV}},\ell)}_\mu\hat{\boldsymbol{\Theta}}^{(\lambda_{\mathrm{UV}},\ell)},\, \nabla^{(\lambda_{\mathrm{UV}},\ell)}_\nu\hat{\boldsymbol{\Theta}}^{(\lambda_{\mathrm{UV}},\ell)}\Big) + \mathcal{V}_\Theta\Big(\hat{\boldsymbol{\Theta}}^{(\lambda_{\mathrm{UV}},\ell)}\Big)\Big]\Pi^{[\chi]}_{\Theta\Theta}\,\Pi_\alpha,\\
\mathcal{L}^{[\chi](\lambda_{\mathrm{UV}},\ell)}_{\Theta;\alpha}(x) &:= \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\Big\{\widehat{\mathcal{L}}^{[\chi](\lambda_{\mathrm{UV}},\ell)}_{\Theta;\alpha}(x)\Big\},
\end{aligned} \tag{9.145}$$

with band functor $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ as in Definition 9.307 and compatibility $[\hat{Q}_{\mathrm{BRST}}, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}] = 0 = [\Pi_\alpha, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}]$ (cf. eq. (9.137)).

Properties.

1. **BRST neutrality.** Ghost number 0 and $[\hat{Q}_{\mathrm{BRST}}, \widehat{\mathcal{L}}^{[\chi]}_{\Theta;\alpha}] = 0$; hence $[\hat{Q}_{\mathrm{BRST}}, \mathcal{L}^{[\chi]}_{\Theta;\alpha}] = 0$.
2. **Triadic/type compatibility.** $[\Pi_\alpha, \Pi^{[\chi]}_{\Theta\Theta}] = 0$, with orthogonality/idempotence as in Proposition 9.202: $\Pi_\alpha\Pi_\beta = \delta_{\alpha\beta}\Pi_\alpha$, $\sum_{\alpha\in\mathfrak{A}}\Pi_\alpha = \mathbf{1}$; $\Pi^{[\chi]}_{\Theta\Theta}\Pi^{[\chi']}_{\Theta\Theta} = \delta_{\chi\chi'}\,\Pi^{[\chi]}_{\Theta\Theta}$, $\sum_\chi \Pi^{[\chi]}_{\Theta\Theta} = \mathbf{1}$ (on the $\Theta\Theta$ block).
3. **Aggregation identities.** Writing the *typed* slice $\mathcal{L}^{[\chi]}_\Theta := \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\Big\{\Pi^{[\chi]}_{\Theta\Theta}[\cdots]\Pi^{[\chi]}_{\Theta\Theta}\Big\}$ and the *full* density $\mathcal{L}_\Theta$ from Definition 9.307, one has
$$\sum_{\alpha\in\mathfrak{A}}\mathcal{L}^{[\chi]}_{\Theta;\alpha} = \mathcal{L}^{[\chi]}_\Theta, \qquad \sum_\chi\sum_{\alpha\in\mathfrak{A}}\mathcal{L}^{[\chi]}_{\Theta;\alpha} = \mathcal{L}_\Theta,$$
and the same equalities hold for the banded forms in (9.145).
4. **Inputs to Parasymmetric T–L–O.** The slices $\mathcal{L}^{[\chi]}_{\Theta;\alpha}$ (banded on the diagnostic realized reference configuration) feed the typed SynTh meters (cf. Section 9.17.11); their spectra and pole content agree with the typed Hessian route within plateau tolerances (Ch. 8).

9.18.1 The Tantrakyma Field Action Functional

The dynamics of the Tantrakyma field—and, by extension, its fundamental quanta, the **kymons**—are governed by the **Tantrakyma Field Action Functional** $S_\Theta[\hat{\boldsymbol{\Theta}}]$. This functional is obtained by integrating the operator-valued Lagrangian density $\mathcal{L}_\Theta(x)$ (Definition 9.301) over the emergent quantum configuration space $\mathcal{X}_q$, with the relational volume element $d\mu(x)$.

Definition 9.310 (Tantrakyma Field Action Functional)**.** Let $\mathcal{L}_\Theta(x)$ be the (scalar, c-number) Lagrangian density defined in Definition 9.301, obtained by tracing the operator-level kernel over the fiber algebra. The action functional is

$$\boxed{S_\Theta[\hat{\boldsymbol{\Theta}}] := \int_{\mathcal{X}_q} d\mu(x)\, \mathcal{L}_\Theta(x) = \int_{\mathcal{X}_q} d^d x \sqrt{-\det \bar{\Sigma}(x)}\, \mathcal{L}_\Theta(x),} \tag{9.146}$$

where $d\mu(x) := d^d x \sqrt{-\det \bar{\Sigma}(x)}$ is the relational volume element (cf. Definition 6.33). Substituting $\mathcal{L}_\Theta$ from Definition 9.301 gives the explicit form

$$S_\Theta[\hat{\boldsymbol{\Theta}}] = \int_{\mathcal{X}_q} d^d x \sqrt{-\det \bar{\Sigma}(x)}\, \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\Big\{ \bar{\Sigma}^{\mu\nu}(x)\, g_C\Big(\nabla_\mu \hat{\boldsymbol{\Theta}}(x),\, \nabla_\nu \hat{\boldsymbol{\Theta}}(x)\Big) + \mathcal{V}_\Theta\Big(\hat{\boldsymbol{\Theta}}(x)\Big)\Big\}. \tag{9.147}$$

The symbols are as follows:

- $\mathcal{X}_q$: emergent configuration manifold (ambient arena used for integration; cf. Definition 6.33).
- $\bar{\Sigma}_{\mu\nu}(x) = \bar{g}_{\mu\nu}(x) + \bar{t}_{\mu\nu}(x)$: unified relational chronogeometric tensor (Definition 5.26); $\bar{\Sigma}^{\mu\nu}(x)$ denotes its inverse (on the diagnostic realized reference configuration; cf. Definition 6.24).
- $\nabla_\mu \hat{\boldsymbol{\Theta}}(x)$: total covariant derivative (base + internal connections) of the Tantrakyma field (Definition 9.301).
- $g_C : T^*\mathcal{M}_{\mathrm{vac}} \times T^*\mathcal{M}_{\mathrm{vac}} \to \mathrm{End}(\mathcal{H}_{\mathrm{em}})$: Hermitian operator-valued co-metric furnishing the inner-product structure for covector-valued variations (Definition 9.301, Remark 9.303).
- $\mathcal{V}_\Theta\Big(\hat{\boldsymbol{\Theta}}(x)\Big)$: Tantrakyma potential operator encoding self-interactions and mass structure of kymons, *built from the co-metric invariant* $\mathsf{X} = g_C(\hat{\boldsymbol{\Theta}}, \hat{\boldsymbol{\Theta}})$ (not from g_C^{-1}); cf. Definition 9.301.
- $\mathrm{Tr}\{\cdots\}$: trace over the fiber algebra $\mathrm{End}(\mathcal{H}_{\mathrm{em}})$, yielding a scalar density.

Remark 9.311 (Relational volume element and $\sqrt{-\det \bar{\Sigma}}$ shorthand)**.** By Definition 6.33, the relational volume element on $\mathcal{X}_q$ is

$$d\mu(x) \;=\; d^d x \sqrt{-\det \bar{g}^{\mathrm{Lor}}_{\mu\nu}(x)}.$$

In other manipulations (e.g. Noether identities), the shorthand $\sqrt{-\det \bar{\Sigma}}$ is employed together with the $\bar{\Sigma}$-compatible Relational Levi–Civita connection $\nabla^{(\mathrm{rel})}$ (Definition 6.7). These notations are consistent because $\bar{g}^{\mathrm{Lor}}$ is the Lorentzian projection determined by $\bar{\Sigma}$ (cf. Section 6.1.5). For strict adherence to Definition 6.33, the second equality in eq. (9.146) may be written with $\sqrt{-\det \bar{g}^{\mathrm{Lor}}}$; if the shorthand $\sqrt{-\det \bar{\Sigma}}$ is retained, include the parenthetical note "shorthand; equal on $\mathfrak{P}$ via $\bar{g}^{\mathrm{Lor}}(\bar{\Sigma})$".

With either notation, the action reads

$$S_\Theta[\hat{\boldsymbol{\Theta}}] = \int_{\mathcal{X}_q} d\mu(x)\, \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\Big\{ \bar{\Sigma}^{\mu\nu}\, g_C\Big(\nabla_\mu \hat{\boldsymbol{\Theta}},\, \nabla_\nu \hat{\boldsymbol{\Theta}}\Big) + \mathcal{V}_\Theta(\hat{\boldsymbol{\Theta}})\Big\},$$

which is consistent with the band-limited (diagnostic realized reference configuration-clean) formulation on a certified plateau $\mathfrak{P}$ with tags $(\lambda_{\mathrm{UV}}, \ell)$ (cf. Definition 9.185 and Section 9.17.11).

Interpretation of the Tantrakyma Action The action functional $S_\Theta[\hat{\mathbf{\Theta}}]$ (Definition 9.310) governs the *dynamical structure of what we perceive as matter* within the emergent geometry. In the same way that the Gaiamiti analysis prepares the chronogeometric stage (Sections 6.5 and 9.17.9), the Tantrakyma action directs the actors—the *kymons*. Crucially, the law depends not merely on a metric but on the full *chronogeometric* data $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (Definition 5.26), with $\bar{\Sigma}^{\mu\nu}$ its inverse (cf. Definition 6.24); this is made explicit in the operator-valued density $\mathcal{L}_\Theta$ (Definition 9.301) and in its band-limited, BRST-clean form (Definition 9.307).

Kinetic term as chronogeometric propagation. The kinetic density $\mathcal{T}_\Theta$ in (9.130) reads

$$\mathcal{T}_\Theta(x) = \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\Big\{\bar{\Sigma}^{\mu\nu}(x)\, g_C\Big(\nabla_\mu\hat{\mathbf{\Theta}}(x),\, \nabla_\nu\hat{\mathbf{\Theta}}(x)\Big)\Big\},$$

and encodes how kymons propagate across the emergent manifold $\mathcal{X}_q$ (cf. Definition 6.33). Here g_C is the operator-valued co-metric on $T^*\mathcal{M}_{\mathrm{vac}}$ (Definition 9.301 and Remark 9.303), so the "cost" of variation is measured in the *cotangent* geometry of matter, while $\bar{\Sigma}^{\mu\nu}$ couples that cost to the *complete* chronogeometric background: curvature of distinguishability via $\bar{g}_{\mu\nu}$ and curvature of phase flow via $\bar{t}_{\mu\nu}$. This is precisely why (i) hyperbolicity and causal cones are $\bar{\Sigma}$-controlled, and (ii) propagation statements descend BRST-safely to $H^0(\hat{Q}_{\mathrm{BRST}})$ on the diagnostic realized reference configuration (Ch. 8; cf. Definition 9.307).

Potential term as identity and self-interaction. The potential density $\mathcal{V}_{\Theta,\mathrm{density}}$ in (9.131),

$$\mathcal{V}_{\Theta,\mathrm{density}}(x) = \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\Big\{\mathcal{V}_\Theta\Big(\hat{\mathbf{\Theta}}(x)\Big)\Big\},$$

governs the *identity* of the kymons: emergent masses, spectra, and internal symmetry-breaking patterns (Definition 9.301). Linearizing around a realized background yields the typed Hessians $\mathcal{K}^{[\chi]}_{\Theta\Theta}$ (eq. (9.124)) and the typed EL response (eq. (9.143)), which determine NG/pNG versus gapped behavior in each Tantrakyma type $[\chi] \in \{\mathrm{CK}, \mathrm{T}_\pm, \mathrm{L}\}$ (Definition 9.295). These are the inputs consumed by the Parasymmetric T–L–O meters (Section 9.17.11) to populate the typed quadrant ledger (Section 9.17.11). In keeping with the covector nature of $\hat{\mathbf{\Theta}}$, the scalar invariant used in the potential is built from the *co-metric* g_C, mirroring the Gaiamiti (vector) case that uses the tangent metric g_T; the pseudoinverse g_C^{-1} is reserved for dualizations/Legendre maps, not for the mass/quartic primitive.

Hierarchy and coupling of the two fundamental sectors. The Gaiamiti field generates the chronogeometric landscape $\bar{\Sigma}_{\mu\nu}$ (Section 6.5), while the Tantrakyma field—and its kymonic excitations—*live within and respond to this complete structure* through $\mathcal{L}_\Theta$ (Definition 9.301). Hence matter dynamics are richer than in metric-only theories: they are inextricably coupled to both geometric and temporal textures of emergence (via $\bar{g}_{\mu\nu}$ and $\bar{t}_{\mu\nu}$), and are diagnosed operationally through the typed, triadic T–L–O pipeline on the certified plateau (Definition 9.307 and Sections 9.17 and 9.17.11).

Proposition 9.312 (Euler–Lagrange equations on the diagnostic realized reference configuration)**.** *For admissible variations $\delta\hat{\mathbf{\Theta}}$ with compact support (diagnostic realized reference configuration boundary terms suppressed by the plateau budget), stationarity $\delta S_\Theta = 0$ yields the operator EL system*

$$\bar{\Sigma}^{\mu\nu}\,\nabla_\mu\nabla_\nu\,\hat{\mathbf{\Theta}} \;+\; (\nabla_\mu\bar{\Sigma}^{\mu\nu})\,\nabla_\nu\hat{\mathbf{\Theta}} \;=\; \frac{\delta\mathcal{V}_\Theta}{\delta\hat{\mathbf{\Theta}}}\,, \tag{9.148}$$

with the companion adjoint equation obtained by varying $\hat{\mathbf{\Theta}}^\dagger$ *when* $\hat{\mathbf{\Theta}}$ *is not Hermitian. Its typed restriction and linearization read*

$$\begin{aligned}\left(\mathbb{K}_\Theta^{[\chi]} + \mathcal{K}_{\Theta\Theta}^{[\chi]}\right)\delta\hat{\mathbf{\Theta}}^{[\chi]} &= \hat{\mathcal{J}}^{[\chi]},\\ \delta\hat{\mathbf{\Theta}}^{[\chi]} &:= \Pi_{\Theta\Theta}^{[\chi]}\,\delta\hat{\mathbf{\Theta}},\end{aligned} \tag{9.149}$$

which is the same statement as (9.143)*, with* $\mathcal{K}_{\Theta\Theta}^{[\chi]}$ *the typed Hessian of the potential (cf. eq.* (9.124)*).*

Remark 9.313 (Symmetry, covariance, and Nielsen stability)**.** The density $\mathcal{L}_\Theta$ is covariant under diffeomorphisms of $\mathcal{X}_q$ and invariant under local automorphisms of $\mathrm{End}(\mathcal{H}_{\mathrm{em}})$. On $H^0(\hat{Q}_{\mathrm{BRST}})$ the pole locations and residues extracted from the typed, band-limited propagators are gauge-parameter independent (Nielsen-flat), and the principal symbol of $\mathbb{K}_\Theta$ is $\bar{\Sigma}$-hyperbolic (Ch. 8), ensuring well-posedness and Kugo–Ojima positivity of realized modes used by T–L–O. In particular, the mass/quartic potential is built from the co-metric invariant $\mathsf{X} = g_C(\hat{\mathbf{\Theta}}, \hat{\mathbf{\Theta}})$ (not g_C^{-1}), mirroring the Gaiamiti (tangent-metric) construction.

Diagnostic realized reference configuration usage. On a certified plateau $\mathfrak{P}$, we use the band-limited density $\mathcal{L}_\Theta^{(\lambda_{\mathrm{UV}},\ell)}$ and action $S_\Theta^{(\lambda_{\mathrm{UV}},\ell)}$ (cf. Definition 9.307 and eq. (9.142)), and the typed/triadic instrumentation from Section 9.17.11.

9.18.2 Holosymmetric Wordwise Tantrakyma Field

In the holosymmetric wordwise (HS) formulation, the Tantrakyma field is treated entirely within the words–dressed bundle framework (Section 6.4.1), with kinematics governed by the operator-bundle connection (Section 6.6.1) and the Relational Levi–Civita geometry induced by the unified chronogeometric tensor (Definition 6.7).

Definition 9.314 (HS Tantrakyma field as a words–dressed section)**.** The Tantrakyma field is a words–dressed section

$$\hat{\mathbf{\Theta}}(x) \in \Gamma\left(\mathcal{E}_\Theta^{\mathrm{HS}}\right),$$

where $\mathcal{E}_\Theta^{\mathrm{HS}} \to \mathcal{X}_q$ is the HS operator bundle for the Tantrakyma sector with fibres contained in $\mathrm{End}(\mathcal{H}_{\mathrm{em}})$, expressed in holosymmetric frames (Section 6.4.1). Frame changes act by HS–unitaries and are mediated by the operator bundle connection one–form in Section 6.6.1.

Definition 9.315 (HS total covariant derivative for $\hat{\mathbf{\Theta}}$)**.** The HS covariant derivative acting on $\hat{\mathbf{\Theta}}$ is the tripartite operator

$$\boxed{\mathcal{D}_\mu^{\mathrm{HS}}\hat{\mathbf{\Theta}} \;:=\; \left(\nabla_\mu^{(\mathrm{rel})} + \nabla_\mu^{(T)} + \mathcal{D}_\mu\right)\hat{\mathbf{\Theta}}\,, \quad \text{as in (6.259)}\,,} \tag{9.150}$$

combining the $\bar{\Sigma}$–compatible Relational Levi–Civita piece, the vacuum–tangent pullback, and the operator–bundle adjoint connection, respectively (Definition 6.7 and Sections 6.4.1 and 6.6.1).

Definition 9.316 (HS co–metric for Tantrakyma variations)**.** The HS co–metric is a Hermitian operator–valued bilinear form

$$g_C^{\mathrm{HS}} : T^*\mathcal{M}_{\mathrm{vac}} \times T^*\mathcal{M}_{\mathrm{vac}} \longrightarrow \mathrm{End}(\mathcal{H}_{\mathrm{em}}) ,$$

obtained by words–dressing the standard co–metric g_C (Definition 9.301). It is invariant under HS frame changes and compatible with the operator–bundle connection (Section 6.6.1).

Definition 9.317 (HS fundamental invariant for the Tantrakyma potential)**.** The basic HS invariant for the Tantrakyma potential is

$$\boxed{\mathsf{X}^{\mathrm{HS}}(x) \;:=\; g_C^{\mathrm{HS}}\Big(\hat{\boldsymbol{\Theta}}(x), \hat{\boldsymbol{\Theta}}(x)\Big) ,} \tag{9.151}$$

a fibrewise Hermitian scalar operator built from the HS co–metric. Any admissible HS potential density must be a real functional of X^{HS}:

$$\boxed{\mathcal{V}_\Theta^{\mathrm{HS}}(x) \;=\; F_\Theta\Big(\mathsf{X}^{\mathrm{HS}}(x)\Big) , \quad F_\Theta : \mathbb{R}_{\geq 0} \to \mathbb{R} .} \tag{9.152}$$

Definition 9.318 (HS Tantrakyma Lagrangian density)**.** Let $\boldsymbol{g}_{\mu\nu}^{\mathrm{Lor}}$ be the HS effective Lorentzian tensor and $\boldsymbol{g}^{\mathrm{Lor}\,\mu\nu}$ its inverse in holosymmetric frames (Section 5.3.7 and Definition 6.34). The HS Lagrangian density is the c–number obtained by the field–dialectic trace (Definition 6.45):

$$\boxed{\mathcal{L}_\Theta^{\mathrm{HS}}(x) \;:=\; \mathrm{Tr}_{\mathrm{FD}}\Big\{\boldsymbol{g}^{\mathrm{Lor}\,\mu\nu}(x)\, g_C^{\mathrm{HS}}\Big(\mathcal{D}_\mu^{\mathrm{HS}}\hat{\boldsymbol{\Theta}}(x),\, \mathcal{D}_\nu^{\mathrm{HS}}\hat{\boldsymbol{\Theta}}(x)\Big) \;+\; \mathcal{V}_\Theta^{\mathrm{HS}}\Big(\hat{\boldsymbol{\Theta}}(x)\Big)\Big\} .} \tag{9.153}$$

Definition 9.319 (HS Tantrakyma action functional)**.** With the HS relational volume element $d\boldsymbol{\mu}(x) = d^d x\, \boldsymbol{\varpi}(x)$, $\boldsymbol{\varpi}(x) = \Big(\det_{\mathrm{HS}}(-\boldsymbol{g}^{\mathrm{Lor}}(x))\Big)^{1/2}$ (Definition 6.34), the HS action is

$$\boxed{\hat{\boldsymbol{S}}_\Theta\,[\hat{\boldsymbol{\Theta}}] \;:=\; \int_{\mathcal{X}_q} d\boldsymbol{\mu}(x)\; \mathcal{L}_\Theta^{\mathrm{HS}}(x) .} \tag{9.154}$$

This formulation is words–only (HS) and reduces to the standard presentation under the expectation/volume push–throughs (Remark 6.35 and eqs. (6.99) and (6.110)).

Proposition 9.320 (HS symmetry properties and positivity)**.** *The density $\mathcal{L}_\Theta^{\mathrm{HS}}$ in (9.153) is:*

(i) gauge–invariant under HS frame transformations generated by the operator–bundle connection (Section 6.6.1);

(ii) covariant under base diffeomorphisms compatible with the Relational Levi–Civita connection (Definition 6.7);

(iii) positive–semidefinite in its kinetic part whenever g_C^{HS} is positive and $\boldsymbol{g}^{\mathrm{Lor}}$ has the standard Lorentzian signature on the certified plateau.

Proof. We prove (1)–(3) for the holosymmetric wordwise Lagrangian density

$$\mathcal{L}^{\mathrm{HS}}_{\Theta}(x) \;=\; \mathrm{Tr}_{\mathrm{FD}}\Big\{\boldsymbol{g}^{\mathrm{Lor}\,\mu\nu}(x)\, g^{\mathrm{HS}}_{C}\Big(\mathcal{D}^{\mathrm{HS}}_{\mu}\hat{\boldsymbol{\Theta}}(x),\, \mathcal{D}^{\mathrm{HS}}_{\nu}\hat{\boldsymbol{\Theta}}(x)\Big) \;+\; \mathcal{V}^{\mathrm{HS}}_{\Theta}\Big(\hat{\boldsymbol{\Theta}}(x)\Big)\Big\} \quad (\text{cf. } eq.\ (9.153)), \tag{9.155}$$

where $\mathcal{D}^{\mathrm{HS}}_{\mu} = \nabla^{(\mathrm{rel})}_{\mu} + \nabla^{(T)}_{\mu} + \mathcal{D}_{\mu}$ (eq. (9.150) and Definition 9.327), the HS co-metric g^{HS}_{C} is Hermitian (Definition 9.316), and $\mathrm{Tr}_{\mathrm{FD}}$ is the field–dialectic trace (Definition 6.45).

(1) Gauge invariance under HS frame transformations. Let $U(x)$ be an HS frame change generated by the operator–bundle connection (Section 6.6.1), acting by conjugation on bundle fibres:

$$\hat{\boldsymbol{\Theta}} \mapsto \hat{\boldsymbol{\Theta}}' = U\,\hat{\boldsymbol{\Theta}}\,U^{\dagger}, \qquad \mathcal{A}_{\mu} \mapsto \mathcal{A}'_{\mu} = U\mathcal{A}_{\mu}U^{\dagger} - (\partial_{\mu}U)U^{\dagger}.$$

By standard gauge-covariance of the connection, the HS total derivative transforms *equivariantly* (eq. (9.150)):

$$\mathcal{D}^{\mathrm{HS}}_{\mu}\hat{\boldsymbol{\Theta}} \;\mapsto\; \mathcal{D}^{\mathrm{HS}\,\prime}_{\mu}\hat{\boldsymbol{\Theta}}' \;=\; U\Big(\mathcal{D}^{\mathrm{HS}}_{\mu}\hat{\boldsymbol{\Theta}}\Big)U^{\dagger}. \tag{9.156}$$

By construction, g^{HS}_{C} is invariant under HS frame changes in the sense that

$$g^{\mathrm{HS}}_{C}\Big(UXU^{\dagger},\, UYU^{\dagger}\Big) \;=\; U\, g^{\mathrm{HS}}_{C}(X,Y)\,U^{\dagger}, \tag{9.157}$$

and the potential is built from the HS invariant $\mathsf{X}^{\mathrm{HS}} = g^{\mathrm{HS}}_{C}(\hat{\boldsymbol{\Theta}},\hat{\boldsymbol{\Theta}})$ (Definition 9.317 and eqs. (9.151) and (9.152)), so that $\mathcal{V}^{\mathrm{HS}}_{\Theta}(\hat{\boldsymbol{\Theta}}) \mapsto U\,\mathcal{V}^{\mathrm{HS}}_{\Theta}(\hat{\boldsymbol{\Theta}})\,U^{\dagger}$. Since $\boldsymbol{g}^{\mathrm{Lor}\,\mu\nu}$ is a base tensor (gauge-blind), inserting (9.156)–(9.157) into (9.155) and using the cyclicity of $\mathrm{Tr}_{\mathrm{FD}}$ (Definition 6.45 and eq. (6.99)) gives

$$\mathcal{L}^{\mathrm{HS}}_{\Theta}(x) \;\mapsto\; \mathrm{Tr}_{\mathrm{FD}}\Big\{U(\cdots)U^{\dagger}\Big\} = \mathcal{L}^{\mathrm{HS}}_{\Theta}(x).$$

Hence (1).

(2) Covariance under base diffeomorphisms compatible with $\nabla^{(\mathrm{rel})}$. Let $\varphi : \mathcal{X}_q \to \mathcal{X}_q$ be a diffeomorphism and denote the Jacobian by $J^{\mu}{}_{\alpha} = \partial x^{\mu}/\partial x'^{\alpha}$. The Relational Levi–Civita connection is $\bar{\Sigma}$–compatible (Definition 6.7), hence $\boldsymbol{g}^{\mathrm{Lor}\,\mu\nu}$ transforms as a contravariant $(2,0)$ tensor and $\mathcal{D}^{\mathrm{HS}}_{\mu}\hat{\boldsymbol{\Theta}}$ as a covector–valued operator:

$$\boldsymbol{g}^{\mathrm{Lor}\,\mu\nu} \mapsto \boldsymbol{g}^{\mathrm{Lor}\,\alpha\beta}\, J^{\mu}{}_{\alpha} J^{\nu}{}_{\beta}, \qquad \mathcal{D}^{\mathrm{HS}}_{\mu}\hat{\boldsymbol{\Theta}} \mapsto J^{\alpha}{}_{\mu}\,(\mathcal{D}^{\mathrm{HS}}_{\alpha}\hat{\boldsymbol{\Theta}}) \circ \varphi^{-1}.$$

Therefore the contraction $\boldsymbol{g}^{\mathrm{Lor}\,\mu\nu}\, g^{\mathrm{HS}}_{C}(\mathcal{D}^{\mathrm{HS}}_{\mu}\hat{\boldsymbol{\Theta}}, \mathcal{D}^{\mathrm{HS}}_{\nu}\hat{\boldsymbol{\Theta}})$ is a scalar (operator) on the base, and $\mathrm{Tr}_{\mathrm{FD}}$ yields a scalar c–number density. Consequently $\mathcal{L}^{\mathrm{HS}}_{\Theta}$ transforms as a scalar under such diffeomorphisms, proving (2).

(3) Positivity of the kinetic part on the certified plateau. Write the HS kinetic operator at x as

$$\mathcal{K}^{\mathrm{HS}}(x) \;:=\; \mathrm{Tr}_{\mathrm{FD}}\Big\{\boldsymbol{g}^{\mathrm{Lor}\,\mu\nu}(x)\, g^{\mathrm{HS}}_{C}\Big(\mathcal{D}^{\mathrm{HS}}_{\mu}\hat{\boldsymbol{\Theta}}(x),\, \mathcal{D}^{\mathrm{HS}}_{\nu}\hat{\boldsymbol{\Theta}}(x)\Big)\Big\}.$$

Assume g^{HS}_{C} is positive in the sense that $g^{\mathrm{HS}}_{C}(X,X) \succeq 0$ as an operator for all X (Definition 9.316). On a certified plateau, choose a time orientation and let u_{μ} be a future unit covector; decompose the co-metric as

$$\boldsymbol{g}^{\mathrm{Lor}\,\mu\nu} \;=\; u^{\mu}u^{\nu} \;-\; h^{\mu\nu},$$

with $h^{\mu\nu}$ the induced positive–definite spatial co-metric (standard Lorentzian signature). Let $W_\mu := \mathcal{D}^{\mathrm{HS}}_\mu \hat{\boldsymbol{\Theta}}$. By cone–gating on the plateau, W_μ is supported in the dual causal cone (cf. the effective light–cone structure and causal gating: Section 5.4.2 and Proposition 9.108), hence

$$u^\mu u^\nu \, g^{\mathrm{HS}}_C(W_\mu, W_\nu) \;\succeq\; 0, \qquad h^{\mu\nu} \, g^{\mathrm{HS}}_C(W_\mu, W_\nu) \;\succeq\; 0,$$

as operator inequalities. Therefore

$$\boldsymbol{g}^{\mathrm{Lor}\,\mu\nu} \, g^{\mathrm{HS}}_C(W_\mu, W_\nu) \;=\; u^\mu u^\nu \, g^{\mathrm{HS}}_C(W_\mu, W_\nu) \;-\; h^{\mu\nu} \, g^{\mathrm{HS}}_C(W_\mu, W_\nu) \;\succeq\; 0,$$

where the last inequality follows because dual–cone support guarantees the time–like contribution dominates the spatial one (standard Minkowski inequality inside the cone). Taking the (positive) field–dialectic trace preserves the ordering, hence $\mathcal{K}^{\mathrm{HS}}(x) \geq 0$. This shows that the kinetic part of $\mathcal{L}^{\mathrm{HS}}_\Theta$ is positive–semidefinite on the certified plateau when g^{HS}_C is positive and $\boldsymbol{g}^{\mathrm{Lor}}$ has standard Lorentzian signature, establishing (3).

Items (1)–(3) together prove the proposition. □

Definition 9.321 (HS Euler–Lagrange equation for $\hat{\boldsymbol{\Theta}}$)**.** Varying (9.154) with respect to $\hat{\boldsymbol{\Theta}}$ (holding HS frames fixed) yields the covariant field equation

$$\boxed{\mathcal{D}^{\mathrm{HS}}_\mu \left(\boldsymbol{g}^{\mathrm{Lor}\,\mu\nu} \frac{\delta}{\delta(\mathcal{D}^{\mathrm{HS}}_\nu \hat{\boldsymbol{\Theta}})} \mathrm{Tr}_{\mathrm{FD}}\Big\{ g^{\mathrm{HS}}_C(\mathcal{D}^{\mathrm{HS}}_\nu \hat{\boldsymbol{\Theta}}, \cdot) \Big\} \right) \;-\; \frac{\delta}{\delta \hat{\boldsymbol{\Theta}}} \mathrm{Tr}_{\mathrm{FD}}\Big\{ \mathcal{V}^{\mathrm{HS}}_\Theta(\hat{\boldsymbol{\Theta}}) \Big\} \;=\; 0\,,} \tag{9.158}$$

with all derivatives understood as in (9.150). The operator functional derivatives are taken inside the $\mathrm{Tr}_{\mathrm{FD}}$; contractions use $\boldsymbol{g}^{\mathrm{Lor}}$.

Remark 9.322 (Reduction to standard variables)**.** Applying the expectation/volume push–throughs (Remark 6.35) to (9.154) recovers the standard action in Section 9.18.1, with the identifications $\boldsymbol{g}^{\mathrm{Lor}} \mapsto \bar{g}^{\mathrm{Lor}}$, $g^{\mathrm{HS}}_C \mapsto g_C$, and $\mathcal{D}^{\mathrm{HS}}_\mu \mapsto \nabla_\mu$, consistent with the Chapter VI HS↔standard equivalence (Definition 6.45 and eqs. (6.99), (6.110) and (6.259)).

9.18.3 Covariant Derivative of the Tantrakyma Field

The Tantrakyma field $\hat{\boldsymbol{\Theta}}(x)$ is a section of the operator-valued cotangent bundle $\mathrm{End}(\mathcal{H}_{\mathrm{em}}) \otimes T^*\mathcal{M}_{\mathrm{vac}}$ over the emergent configuration space $\mathcal{X}_q$ (cf. Definition 9.301). To define its dynamics as used in the action $\mathcal{L}_\Theta$ and S_Θ (Definitions 9.301 and 9.310), the derivative must capture variations along the base manifold and transformations in the internal fiber. Accordingly, the total covariant derivative ∇_μ unifies general covariance with local gauge covariance.

Definition 9.323 (Covariant derivative of the Tantrakyma field)**.** Let $\hat{\boldsymbol{\Theta}}(x)$ be the Dynamical Tantrakyma Field. Define

$$\boxed{\nabla_\mu \hat{\boldsymbol{\Theta}} := \nabla^{(\mathrm{rel})}_\mu \hat{\boldsymbol{\Theta}} \;+\; D_\mu \hat{\boldsymbol{\Theta}},} \tag{9.159}$$

where:

1. **Relational Levi–Civita derivative** $\nabla_\mu^{(\mathrm{rel})}$ is the torsion-free, $\bar{\Sigma}$-compatible connection associated with the chronogeometric tensor $\bar{\Sigma}_{\alpha\beta}$ ($\nabla_\mu^{(\mathrm{rel})}\bar{\Sigma}_{\alpha\beta}=0$), defined in Definition 6.7, built from $\bar{\Sigma}_{\alpha\beta}=\bar{g}_{\alpha\beta}+\bar{t}_{\alpha\beta}$ (Definition 5.26). It acts on the cotangent indices of $\hat{\boldsymbol{\Theta}}$.
2. **Gauge-covariant derivative** D_μ acts on the operator-valued fiber via the operator-bundle connection $\mathcal{A}_\mu(x)$ (cf. Definition 6.192):

$$D_\mu\hat{\boldsymbol{\Theta}}(x):=\partial_\mu\hat{\boldsymbol{\Theta}}(x)+i\big[\mathcal{A}_\mu(x),\,\hat{\boldsymbol{\Theta}}(x)\big].$$

The field strength is $\mathcal{F}_{\mu\nu}:=\partial_\mu\mathcal{A}_\nu-\partial_\nu\mathcal{A}_\mu-i[\mathcal{A}_\mu,\mathcal{A}_\nu]$. Under a local unitary U_x (inner automorphism), $\hat{\boldsymbol{\Theta}}\mapsto U_x\hat{\boldsymbol{\Theta}}U_x^\dagger$, $\mathcal{A}_\mu\mapsto U_x\mathcal{A}_\mu U_x^\dagger-i(\partial_\mu U_x)U_x^\dagger$, and $D_\mu\hat{\boldsymbol{\Theta}}$ transforms covariantly.

In a local basis $\{\mathbf{e}_\nu(x)\}$ of $T^*_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}$ with $\hat{\boldsymbol{\Theta}}(x)=\hat{\boldsymbol{\Theta}}^\nu(x)\,\mathbf{e}_\nu(x)$, the component form is

$$\boxed{(\nabla_\mu\hat{\boldsymbol{\Theta}})_\lambda(x):=\partial_\mu\hat{\boldsymbol{\Theta}}_\lambda(x)-\Gamma^\nu_{\mu\lambda}\big[\bar{\Sigma}\big]\,(x)\,\hat{\boldsymbol{\Theta}}_\nu(x)+i\big[\,\mathcal{A}_\mu(x),\,\hat{\boldsymbol{\Theta}}_\lambda(x)\,\big],}\tag{9.160}$$

where $\Gamma^\nu_{\mu\lambda}[\bar{\Sigma}]$ are the Christoffel symbols of $\nabla^{(\mathrm{rel})}$.

Diagnostic realized reference configuration / band-limited form and BRST compatibility. On a certified plateau $\mathfrak{P}$ with tags $(\lambda_{\mathrm{UV}},\ell)$ and projection to $H^0(\hat{Q}_{\mathrm{BRST}})$ (cf. Definition 9.307 and eq. (9.137)), we use the band-limited derivative

$$\nabla_\mu^{(\lambda_{\mathrm{UV}},\ell)}\hat{\boldsymbol{\Theta}}^{(\lambda_{\mathrm{UV}},\ell)}:=\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}\big[\nabla_\mu\hat{\boldsymbol{\Theta}}\big]=\nabla_\mu^{(\mathrm{rel})(\lambda_{\mathrm{UV}},\ell)}\hat{\boldsymbol{\Theta}}^{(\lambda_{\mathrm{UV}},\ell)}+D_\mu^{(\lambda_{\mathrm{UV}},\ell)}\hat{\boldsymbol{\Theta}}^{(\lambda_{\mathrm{UV}},\ell)}.\tag{9.161}$$

This is BRST-closed (ghost number 0) and compatible with the typed/triadic filters used in T–L–O: on $\mathfrak{P}$ we choose connections so that $[\Pi_\alpha,\nabla_\mu]=0$ and $[\Pi^{[\chi]}_{\Theta\Theta},D_\mu]=0$ (or commute up to BRST-exact terms) (cf. Definition 9.295 and Proposition 9.202).

Proposition 9.324 (Compatibility with $\bar{\Sigma}$ and g_C; Leibniz rule)**.** *The total derivative* $\nabla_\mu=\nabla_\mu^{(\mathrm{rel})}+D_\mu$ *satisfies:*

(i) $\nabla_\mu^{(\mathrm{rel})}\bar{\Sigma}_{\alpha\beta}=0$ *(metric compatibility) and torsion-freeness (Definition 6.7).*

(ii) Co-metric compatibility on all legs: $\nabla_\mu^{(\mathrm{rel})}g_C=0$ *on the cotangent slots and* $D_\mu g_C=0$ *on the operator (fiber) slot; equivalently,* $\nabla_\mu g_C=0$ *(cf. Remark 9.303).*

(iii) The Leibniz rule for operator-valued tensors and trace cyclicity under $\mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}$*, ensuring covariant integration by parts in* S_Θ *(Definition 9.310).*

These properties guarantee that the kinetic term in $\mathcal{L}_\Theta$ *is well-defined (with the co-metric* g_C *contracting the covector legs) and variationally compatible with the Euler–Lagrange system (Definition 9.305 and eq. (9.143)).*

Remark 9.325 (Curvatures and identities)**.** The relational Riemann tensor $R^\rho{}_{\sigma\mu\nu}[\bar{\Sigma}]$ and the bundle field strength $\mathcal{F}_{\mu\nu}$ obey their respective Bianchi identities. On the diagnostic realized reference configuration, the typed/triadic projections of $\nabla_\mu\hat{\boldsymbol{\Theta}}$ provide inputs to the Parasymmetric meters (Section 9.17.11), feeding the quadrant decision and T–L–O stamps (Section 9.17.11). In keeping with the covector nature of $\hat{\boldsymbol{\Theta}}$, scalar invariants for the *potential* are built with the co-metric g_C (not g_C^{-1}), mirroring the Gaiamiti (vector, metric) construction.

Interpretation of the Composite Derivative The dual structure of the covariant derivative ∇_μ in (9.159) is the natural consequence of the dual identity of the Tantrakyma field: $\hat{\boldsymbol{\Theta}}$ is simultaneously a geometric covector and an operator on the fiber algebra (cf. Definition 9.301). Accordingly, ∇_μ splits into a *relational* (base-manifold) part and a *gauge* (fiber) part, each with a clear operational meaning and symmetry transformation law (Definition 9.323 and eq. (9.160)).

Relational part (geometric covariance). The relational Levi–Civita connection $\nabla^{(\mathrm{rel})}$ is torsion-free and $\bar{\Sigma}$-compatible ($\nabla^{(\mathrm{rel})}_\mu \bar{\Sigma}_{\alpha\beta} = 0$; Definitions 5.26 and 6.7), hence it transports the covector indices of $\hat{\boldsymbol{\Theta}}$ consistently with the full chronogeometric structure $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$. In components this yields the Christoffel-symbol correction $-\Gamma^\nu_{\mu\lambda}[\bar{\Sigma}]\,\hat{\boldsymbol{\Theta}}_\nu$ in (9.160), which ties the *propagation of kymons* to both (i) curvature of distinguishability ($\bar{g}$) and (ii) curvature of quantum phase flow ($\bar{t}$). Through the kinetic density $\mathcal{T}_\Theta$ (eq. (9.130)) this relational coupling governs the causal cones and hyperbolicity of the EL system (eqs. (9.148) and (9.149)).

Gauge part (fiber covariance). The operator–bundle connection $\mathcal{A}_\mu$ (cf. Definition 6.192) implements local unitary changes of basis in the fiber $\mathrm{End}(\mathcal{H}_{\mathrm{em}})$. Under a unitary U_x,

$$\hat{\boldsymbol{\Theta}} \;\mapsto\; U_x\,\hat{\boldsymbol{\Theta}}\,U_x^\dagger, \qquad \mathcal{A}_\mu \;\mapsto\; U_x\,\mathcal{A}_\mu\,U_x^\dagger - i(\partial_\mu U_x)\,U_x^\dagger,$$

so the commutator term $i[\mathcal{A}_\mu, \hat{\boldsymbol{\Theta}}]$ in $D_\mu\hat{\boldsymbol{\Theta}}$ enforces covariant transport. The associated field strength is $\mathcal{F}_{\mu\nu} := \partial_\mu\mathcal{A}_\nu - \partial_\nu\mathcal{A}_\mu - i[\mathcal{A}_\mu, \mathcal{A}_\nu]$, and the full derivative $\nabla_\mu = \nabla^{(\mathrm{rel})}_\mu + D_\mu$ appears in eqs. (9.159) and (9.160). Because $\mathcal{L}_\Theta$ uses trace functionals (Definition 9.301), fiber $*$-automorphisms preserve the action (fiber–algebra invariance; cf. Remark 9.313).

Unification and diagnostic realized reference configuration readiness. Taken together, $\nabla_\mu = \nabla^{(\mathrm{rel})}_\mu + D_\mu$ unifies general covariance with local gauge covariance in a way compatible with the chronogeometric background and the operator-valued fiber structure. On a certified plateau $\mathfrak{P}$ with tags $(\lambda_{\mathrm{UV}}, \ell)$, the band-limited derivative $\nabla^{(\lambda_{\mathrm{UV}},\ell)}_\mu$ of (9.161) is BRST-closed and (by construction of the diagnostic realized reference configuration instrumentation) commutes with the triadic and typed filters $(\Pi_\alpha, \Pi^{[\chi]}_{\Theta\Theta})$ (cf. Definitions 9.185, 9.295, 9.306 and 9.307 and eq. (9.137)). This ensures that the kinetic term remains variationally well-posed, g_C-compatible (Proposition 9.324 and Remark 9.303), and ready for Parasymmetric T–L–O diagnostics (Section 9.17.11).

In essence, the composite derivative ∇_μ is the correct and complete tool to describe the dynamics of a matter-representing quantum field in a self-consistently generated, relational spacetime: it makes the law $\mathcal{L}_\Theta$ and action S_Θ (Definitions 9.301 and 9.310) simultaneously faithful to the base chronogeometry and to the internal quantum gauge structure.

9.18.4 Holosymmetric Wordwise Tantrakyma Covariant Derivative

We formulate the covariant derivative of the Tantrakyma field entirely in the holosymmetric wordwise (HS) framework, mirroring the Gaiamiti construction in Section 6.5.4. The bundle background and gauge data are those of Sections 6.4.1 and 6.6.1; chronogeometry is governed by the Relational Levi–Civita structure of Definition 6.7.

Definition 9.326 (HS Tantrakyma field as a words–dressed cotangent section)**.** The Tantrakyma field is a words–dressed section

$$\hat{\boldsymbol{\Theta}}(x) \;\in\; \Gamma\Big(\mathcal{E}_{\Theta}^{\mathrm{HS}}\Big) \;\subset\; \Gamma\Big(\mathrm{End}(\mathcal{H}_{\mathrm{em}}) \otimes T^{*}\mathcal{M}_{\mathrm{vac}}\Big),$$

expressed in holosymmetric frames (Section 6.4.1). HS frame changes act by unitary conjugation on the operator slot and by orthogonal transformations on $T^{*}\mathcal{M}_{\mathrm{vac}}$, mediated by the operator bundle connection of Section 6.6.1.

Definition 9.327 (Total HS covariant derivative for $\hat{\boldsymbol{\Theta}}$)**.** The total HS covariant derivative combines base, vacuum–tangent, and operator connections (cf. eq. (6.259)):

$$\boxed{\mathcal{D}_{\mu}^{\mathrm{HS}}\hat{\boldsymbol{\Theta}} \;:=\; \underbrace{\nabla_{\mu}^{(\mathrm{rel})}\hat{\boldsymbol{\Theta}}}_{\text{base (Levi–Civita on } \bar{\Sigma})} \;+\; \underbrace{\nabla_{\mu}^{(T)}\hat{\boldsymbol{\Theta}}}_{\text{pullback on } T^{*}\mathcal{M}_{\mathrm{vac}}} \;+\; \underbrace{\mathcal{D}_{\mu}\hat{\boldsymbol{\Theta}}}_{\text{operator (adjoint) connection}} .} \tag{9.162}$$

Here $\nabla_{\mu}^{(\mathrm{rel})}\bar{\Sigma}_{\alpha\beta} = 0$ (torsion-free), $\nabla_{\mu}^{(T)}$ is induced by $x \mapsto |\Omega(x)\rangle$ and g_T on $T^{*}\mathcal{M}_{\mathrm{vac}}$, and $\mathcal{D}_{\mu} = \partial_{\mu} + [A_{\mu},\,\cdot\,]$ acts on the $\mathrm{End}(\mathcal{H}_{\mathrm{em}})$ slot (Definition 6.185 and Section 6.6.1).

Definition 9.328 (HS component forms (cotangent-valued field))**.** Let $\{\mathbf{e}^{i}(x)\}_{i=1}^{r}$ be a g_T–orthonormal local frame of $T^{*}_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}$ with dual $\{\mathbf{e}_{i}(x)\}$. Write $\hat{\boldsymbol{\Theta}} = \hat{\Theta}_{i}\,\mathbf{e}^{i}$. Then

$$\boxed{(\mathcal{D}_{\mu}^{\mathrm{HS}}\hat{\boldsymbol{\Theta}})_{i} = \partial_{\mu}\hat{\Theta}_{i} \;-\; \omega_{\mu\,i}{}^{j}\,\hat{\Theta}_{j} \;+\; [A_{\mu},\hat{\Theta}_{i}],} \tag{9.163}$$

where $\omega_{\mu\,i}{}^{j}$ are the coefficients of the pullback connection on $T^{*}\mathcal{M}_{\mathrm{vac}}$. If a base index is used instead, $\hat{\boldsymbol{\Theta}} = \hat{\boldsymbol{\Theta}}_{\lambda}\,dx^{\lambda}$ and

$$\boxed{(\mathcal{D}_{\mu}^{\mathrm{HS}}\hat{\boldsymbol{\Theta}})_{\lambda} = \partial_{\mu}\hat{\boldsymbol{\Theta}}_{\lambda} \;-\; \Gamma_{\mu\lambda}^{\nu}[\bar{\Sigma}]\,\hat{\boldsymbol{\Theta}}_{\nu} \;+\; [A_{\mu},\hat{\boldsymbol{\Theta}}_{\lambda}].} \tag{9.164}$$

Remark 9.329 (Leibniz and SOT regularity)**.** $\mathcal{D}_{\mu}$ is an inner derivation, so $\mathcal{D}_{\mu}(XY) = (\mathcal{D}_{\mu}X)Y + X(\mathcal{D}_{\mu}Y)$. All derivatives are taken in the strong–operator topology on band–limited fibres (Remark 6.187).

Definition 9.330 (Total HS curvature acting on $\hat{\boldsymbol{\Theta}}$)**.** The commutator of HS covariant derivatives yields

$$\boxed{[\mathcal{D}_{\mu}^{\mathrm{HS}},\mathcal{D}_{\nu}^{\mathrm{HS}}]\hat{\boldsymbol{\Theta}} \;=\; \mathcal{R}_{\mu\nu}\cdot\hat{\boldsymbol{\Theta}} \;+\; [\mathcal{F}_{\mu\nu},\hat{\boldsymbol{\Theta}}], \qquad \mathcal{F}_{\mu\nu} = \partial_{[\mu}A_{\nu]} + [A_{\mu},A_{\nu}],} \tag{9.165}$$

where $\mathcal{R}_{\mu\nu}$ collects the base Riemann curvature $R[\bar{\Sigma}]$ and the vacuum–tangent curvature $\Omega_{\mu\nu} = \partial_{[\mu}\omega_{\nu]} + [\omega_{\mu},\omega_{\nu}]$, acting on the corresponding cotangent indices of $\hat{\boldsymbol{\Theta}}$.

Proposition 9.331 (Gauge/frame covariance of $\mathcal{D}_{\mu}^{\mathrm{HS}}\hat{\boldsymbol{\Theta}}$)**.** *Under modal frame changes $g(x) \in U(n)$ and tangent rotations $R(x) \in O(r)$,*

$$\hat{\Theta}_{i} \mapsto \mathrm{Ad}_{g}(\hat{\Theta}_{j})\,(R^{-1})^{j}{}_{i}, \qquad A_{\mu} \mapsto gA_{\mu}g^{\dagger} - (\partial_{\mu}g)\,g^{\dagger}, \qquad \omega_{\mu} \mapsto R\omega_{\mu}R^{-1} + (\partial_{\mu}R)R^{-1},$$

and $(\mathcal{D}_{\mu}^{\mathrm{HS}}\hat{\boldsymbol{\Theta}})_{i}$ transforms covariantly in the same way. If $\hat{\boldsymbol{\Theta}}$ carries a base index, $\Gamma[\bar{\Sigma}]$ transforms as the Levi–Civita connection.

Proof idea. Working in the g_T–orthonormal frame of Definition 9.328, apply the same steps as in Proposition 6.188 (Gaiamiti case), noting the cotangent sign in (9.163) and the action of $R \in O(r)$ on covector indices. The inhomogeneous pieces in A'_μ and ω'_μ cancel by the same conjugation and frame-rotation identities; the result is the stated covariance.

Holosymmetric wordwise proof (expanded). Work in the g_T–orthonormal coframe of Definition 9.328, writing $\hat{\mathbf{\Theta}} = \hat{\Theta}_i\, \mathbf{e}^i$ so that

$$(\mathcal{D}^{\mathrm{HS}}_\mu \hat{\mathbf{\Theta}})_i = \partial_\mu \hat{\Theta}_i \;-\; \omega_{\mu\, i}{}^{j}\, \hat{\Theta}_j \;+\; [A_\mu, \hat{\Theta}_i] \qquad \text{(cf. eq. (9.163))}. \tag{9.166}$$

Let a *modal frame change* $g(x) \in U(n)$ act by the adjoint $\mathrm{Ad}_g(X) = gXg^\dagger$ on the operator slot, and let a *tangent rotation* $R(x) \in O(r)$ act on the coframe by $\mathbf{e}^i \mapsto \mathbf{e}'^i = (R^{-1})^i{}_j\, \mathbf{e}^j$. Since $\hat{\mathbf{\Theta}}$ is a *covector*-valued field, its components transform as

$$\hat{\Theta}_i \;\longmapsto\; \hat{\Theta}'_i := \mathrm{Ad}_g(\hat{\Theta}_j)\, (R^{-1})^j{}_i. \tag{9.167}$$

The connections transform as

$$A'_\mu = gA_\mu g^\dagger - (\partial_\mu g)\, g^\dagger, \qquad \omega'_\mu = R\omega_\mu R^{-1} + (\partial_\mu R)\, R^{-1}. \tag{9.168}$$

Step 1: Differentiate the transformed field. Using the product rule, the adjunct derivative identity $\partial_\mu(\mathrm{Ad}_g X) = \mathrm{Ad}_g(\partial_\mu X) + [(\partial_\mu g)g^\dagger, \mathrm{Ad}_g X]$, and $\partial_\mu(R^{-1}) = -R^{-1}(\partial_\mu R)R^{-1}$,

$$\begin{aligned} \partial_\mu \hat{\Theta}'_i &= \partial_\mu\Big(\mathrm{Ad}_g(\hat{\Theta}_j)(R^{-1})^j{}_i\Big) \\ &= \mathrm{Ad}_g(\partial_\mu \hat{\Theta}_j)(R^{-1})^j{}_i + [(\partial_\mu g)g^\dagger, \mathrm{Ad}_g(\hat{\Theta}_j)](R^{-1})^j{}_i \\ &\quad - \mathrm{Ad}_g(\hat{\Theta}_j)(R^{-1})^j{}_k (\partial_\mu R)^k{}_m (R^{-1})^m{}_i. \end{aligned} \tag{9.169}$$

Step 2: Act with the transformed pullback connection. Using (9.166) with ω'_μ and $\hat{\Theta}'_i$,

$$-\,\omega'_{\mu\, i}{}^{k}\, \hat{\Theta}'_k = -\left(R\omega_\mu R^{-1}\right)_i^{\,k}\; \mathrm{Ad}_g(\hat{\Theta}_\ell)\, (R^{-1})^\ell{}_k \;-\; \left((\partial_\mu R)R^{-1}\right)_i^{\,k}\; \mathrm{Ad}_g(\hat{\Theta}_\ell)\, (R^{-1})^\ell{}_k. \tag{9.170}$$

Step 3: Act with the transformed operator connection. From A'_μ in (9.168) and (9.167),

$$\begin{aligned} [\, A'_\mu, \hat{\Theta}'_i\,] &= [\, gA_\mu g^\dagger - (\partial_\mu g)g^\dagger,\; \mathrm{Ad}_g(\hat{\Theta}_j)\, (R^{-1})^j{}_i\,] \\ &= \underbrace{[\, gA_\mu g^\dagger,\; \mathrm{Ad}_g(\hat{\Theta}_j)\,]}_{=\; \mathrm{Ad}_g([A_\mu, \hat{\Theta}_j])} (R^{-1})^j{}_i \;-\; [\,(\partial_\mu g)g^\dagger,\; \mathrm{Ad}_g(\hat{\Theta}_j)\,]\, (R^{-1})^j{}_i. \end{aligned} \tag{9.171}$$

Step 4: Assemble $(\mathcal{D}^{\mathrm{HS}}_\mu \hat{\mathbf{\Theta}})'_i$. By definition,

$$(\mathcal{D}^{\mathrm{HS}}_\mu \hat{\mathbf{\Theta}})'_i = \partial_\mu \hat{\Theta}'_i \;-\; \omega'_{\mu\, i}{}^{k}\, \hat{\Theta}'_k \;+\; [\, A'_\mu, \hat{\Theta}'_i\,].$$

Insert (9.169), (9.170), (9.171). Two cancellations occur:

- The terms proportional to $(\partial_\mu g)g^\dagger$ cancel *exactly* between $[(\partial_\mu g)g^\dagger, \mathrm{Ad}_g(\hat{\Theta}_j)](R^{-1})^j{}_i$ in (9.169) and $-[\,(\partial_\mu g)g^\dagger, \mathrm{Ad}_g(\hat{\Theta}_j)\,](R^{-1})^j{}_i$ in (9.171).

- The terms containing $(\partial_\mu R)R^{-1}$ cancel between the third term of (9.169) and the second term of (9.170) upon relabeling indices and using $-(R^{-1})^j{}_k(\partial_\mu R)^k{}_m(R^{-1})^m{}_i = -\big((\partial_\mu R)R^{-1}\big)_i{}^k(R^{-1})^j{}_k$.

What remains is

$$\begin{aligned}(\mathcal{D}^{\mathrm{HS}}_\mu\hat{\boldsymbol{\Theta}})'_i &= \mathrm{Ad}_g(\partial_\mu\hat\Theta_j)(R^{-1})^j{}_i - (R\omega_\mu R^{-1})_i{}^k\ \mathrm{Ad}_g(\hat\Theta_\ell)(R^{-1})^\ell{}_k + \mathrm{Ad}_g([A_\mu,\hat\Theta_j])(R^{-1})^j{}_i\\ &= \mathrm{Ad}_g\Big(\partial_\mu\hat\Theta_j - \omega_\mu{}_j{}^\ell\hat\Theta_\ell + [A_\mu,\hat\Theta_j]\Big)(R^{-1})^j{}_i\end{aligned}$$

(using $R\omega_\mu R^{-1}$ on the covector index)

$$= \mathrm{Ad}_g\Big((\mathcal{D}^{\mathrm{HS}}_\mu\hat{\boldsymbol{\Theta}})_j\Big)(R^{-1})^j{}_i.$$

This is precisely the claimed *covariant* transformation law of the HS derivative: it transforms by inner automorphism on the operator slot and by R^{-1} on the covector index.

Step 5 (optional): Inclusion of a base index. If $\hat{\boldsymbol{\Theta}} = \hat{\boldsymbol{\Theta}}_\lambda\, dx^\lambda$, then

$$(\mathcal{D}^{\mathrm{HS}}_\mu\hat{\boldsymbol{\Theta}})_\lambda = \partial_\mu\hat{\boldsymbol{\Theta}}_\lambda - \Gamma^\nu_{\mu\lambda}[\bar\Sigma]\,\hat{\boldsymbol{\Theta}}_\nu + [A_\mu,\hat{\boldsymbol{\Theta}}_\lambda] \qquad \text{(cf. eq. (9.164))}.$$

Under a diffeomorphism or base frame change, $\Gamma^\nu_{\mu\lambda}[\bar\Sigma]$ transforms as the Levi–Civita connection (Definition 6.7), so the same calculation goes through with an additional covector index λ carried along; covariance persists componentwise.

Conclusion. Therefore, under the HS modal frame $g \in U(n)$ and tangent rotation $R \in O(r)$,

$$(\mathcal{D}^{\mathrm{HS}}_\mu\hat{\boldsymbol{\Theta}})_i \;\longmapsto\; (\mathcal{D}^{\mathrm{HS}}_\mu\hat{\boldsymbol{\Theta}})'_i = \mathrm{Ad}_g\Big((\mathcal{D}^{\mathrm{HS}}_\mu\hat{\boldsymbol{\Theta}})_j\Big)\,(R^{-1})^j{}_i,$$

and, when present, base indices transform by the Levi–Civita rule. This proves Proposition 9.331. □

Remark 9.332 (Insertion into the HS Tantrakyma dynamics)**.** Substituting (9.150) and the component forms (9.163)–(9.164) into the HS Lagrangian and action (Definitions 9.318 and 9.319) yields the wordwise dynamics used in Proposition 9.320. The HS→standard reduction follows Remark 9.322, identifying $\boldsymbol{g}^{\mathrm{Lor}} \mapsto \bar g^{\mathrm{Lor}}$, $g^{\mathrm{HS}}_C \mapsto g_C$, and $\mathcal{D}^{\mathrm{HS}}_\mu \mapsto \nabla^{(\mathrm{rel})}_\mu + D_\mu$ as in Section 9.18.3 and Definition 9.323.

9.18.5 Tantrakyma Field Potential and the Origin of the Kymon

While the kinetic term governs propagation (eq. (9.130)), the *potential term* $\mathcal{V}_\Theta(\hat{\boldsymbol{\Theta}})$ fixes the self-interaction and selects energetically preferred backgrounds for the matter sector (eq. (9.131)). It is the proximate source of the kymon's intrinsic properties (masses, symmetry patterns), and it is the matter-sector analogue of the Gaiamiti potential discussed in Section 9.17.

Invariant Construction of the Tantrakyma Potential

The Tantrakyma field is an operator-valued *covector* on the vacuum manifold, $\hat{\mathbf{\Theta}}(x) \in T^*_{|\Omega(x)\rangle}\mathcal{M}_{\text{vac}} \otimes \text{End}(\mathcal{H}_{\text{em}})$ (cf. Proposition 9.292 and Definitions 6.142 and 9.301). To build scalar (operator-valued) invariants from covectors one uses the operator-valued *co-metric* g_C (cf. Remark 9.303). The basic invariant is therefore the "amplitude"

$$\mathsf{X}(x) := g_C\Big(\hat{\mathbf{\Theta}}(x),\, \hat{\mathbf{\Theta}}(x)\Big) \;\in\; \text{End}(\mathcal{H}_{\text{em}}),$$

which plays for Tantrakyma the role that $g_T(\cdot,\cdot)$ plays for Gaiamiti. (The pseudoinverse g_C^{-1} is useful for dualizations/Legendre maps but is *not* the primitive building block of the mass/quartic potential.)

Definition 9.333 (Tantrakyma field potential)**.** The minimal, Hermitian, symmetry-respecting operator potential built from $\mathsf{X} = g_C(\hat{\mathbf{\Theta}}, \hat{\mathbf{\Theta}})$ is

$$\boxed{\mathcal{V}_\Theta(\hat{\mathbf{\Theta}}) = \tfrac{1}{2}\, m_\Theta^2\, \mathsf{X} \;+\; \tfrac{\lambda_\Theta}{4!}\, \mathsf{X}^2 \;+\; \mathcal{O}\Big(\mathsf{X}^3/\Lambda_\Theta^2\Big),} \tag{9.172}$$

with real parameters m_Θ^2 and $\lambda_\Theta > 0$, and a suppression scale Λ_Θ for higher operators. The scalar potential density used in $\mathcal{L}_\Theta$ is

$$\mathcal{V}_{\Theta,\text{density}}(x) = \text{Tr}_{\mathcal{H}_{\text{em}}}\{\mathcal{V}_\Theta(\hat{\mathbf{\Theta}}(x))\}. \tag{9.173}$$

Diagnostic realized reference configuration usage. On a certified plateau $\mathfrak{P}$ with tags $(\lambda_{\text{UV}}, \ell)$, the same form holds with $\hat{\mathbf{\Theta}} \to \hat{\mathbf{\Theta}}^{(\lambda_{\text{UV}},\ell)}$ (cf. Definitions 9.306 and 9.307). Expectation values define the effective potential

$$V_\Theta \;:=\; \Big\langle \mathcal{V}_\Theta \Big\rangle_{\mathfrak{P}} \;=\; \text{Tr}_{\mathcal{H}_{\text{em}}}\Big\langle \tfrac{1}{2} m_\Theta^2\, \mathsf{X} \;+\; \tfrac{\lambda_\Theta}{4!}\, \mathsf{X}^2 \Big\rangle_{\mathfrak{P}} \;+\; \cdots, \tag{9.174}$$

to be minimized over admissible backgrounds $\bar{\Theta}$ consistent with the EL equations (eq. (9.148)).

1. Field content and representation in the dual space

The covector character of $\hat{\mathbf{\Theta}}$ (Ch. 7) forces the invariant building block to use the *co-metric* g_C (rather than the Gaiamiti pairing g_T); see Remark 9.303. Thus the fundamental scalar operator is

$$\mathsf{X} \;:=\; g_C\Big(\hat{\mathbf{\Theta}},\, \hat{\mathbf{\Theta}}\Big) \;\in\; \text{End}(\mathcal{H}_{\text{em}}),$$

from which all polynomial invariants follow via the trace.

2. Constraints of stability, symmetry, and minimality

Stability. Boundedness from below requires $\lambda_\Theta > 0$ and, for the quartic truncation, fixes the sign structure of the extrema in V_Θ.

Symmetry. A $\mathbb{Z}_2$ invariance $\hat{\mathbf{\Theta}} \mapsto -\hat{\mathbf{\Theta}}$ and, more generally, invariance under local fiber inner $*$-automorphisms (cf. Definitions 6.192 and 9.301) restrict $\mathcal{V}_\Theta$ to depend only on the invariant X.

Minimality/Universality. Retaining operators up to mass dimension four yields the quartic form in eq. (9.133), the lowest-order choice capable of nontrivial vacuum structure (cf. Ch. 8 discussion of universality classes).

3. Vacuum structure and kymon origin (typed view)

Let $x := \left\langle \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}} \mathsf{X} \right\rangle_{\mathfrak{P}}$ be the invariant amplitude. Minimizing the quartic part of eq. (9.174) gives

$$\frac{d}{dx}\left(\tfrac{1}{2}m_\Theta^2 x + \tfrac{\lambda_\Theta}{4!}x^2\right) = 0 \implies x_\star = \begin{cases} 0, & m_\Theta^2 > 0, \\ -\dfrac{6\,m_\Theta^2}{\lambda_\Theta}, & m_\Theta^2 < 0 \text{ (SSB)}. \end{cases}$$

About any realized background $\bar{\Theta}$ solving the EL equations (eq. (9.148)), the typed mass/curvature operators are obtained from the potential Hessian restricted by the Tantrakyma projectors (Definition 9.295):

$$\boxed{\mathcal{K}^{[\chi]}_{\Theta\Theta} := \Pi^{[\chi]}_{\Theta\Theta}\, \frac{\delta^2 \mathcal{V}_\Theta}{\delta\hat{\boldsymbol{\Theta}}\,\delta\hat{\boldsymbol{\Theta}}}\bigg|_{\bar{\Theta}} \Pi^{[\chi]}_{\Theta\Theta}, \qquad [\chi] \in \{\mathrm{CK}, \mathrm{T}_\pm, \mathrm{L}\}.} \tag{9.175}$$

The eigenstructure of $\mathcal{K}^{[\chi]}_{\Theta\Theta}$ determines the kymon content: zero/pseudo-zero modes indicate NG/pNG directions; positive eigenvalues signal gapped modes. These predictions cross-check the SynTh meters and typed quadrant outcomes in the T–L–O pipeline (eqs. (9.126) to (9.128) and Section 9.17.11).

Remark 9.334 (Coupling to chronogeometry and diagnostic realized reference configuration readiness)**.** Although $\mathcal{V}_\Theta$ depends only on X, the *realization* of vacua and spectra is diagnostic realized reference configuration- and channel-dependent through the kinetic pairing with $\bar{\Sigma}^{\mu\nu}$ in $\mathcal{T}_\Theta$ (eq. (9.130)) and the typed/triadic evaluation on $\mathfrak{P}$ (Definition 9.307 and Section 9.17.11). BRST/Nielsen hygiene ensures gauge-parameter independence of pole data on $H^0(\hat{Q}_{\mathrm{BRST}})$.

Summary. The potential eq. (9.133) is the unique minimal operator-valued choice consistent with stability, symmetry, and universality. Its Hessian eq. (9.175) seeds the typed mass operators that generate the kymon spectrum; together with the chronogeometric kinetic pairing, it yields a complete, diagnostic realized reference configuration-certified account of matter dynamics that mirrors the Gaiamiti story while living on the cotangent side of the vacuum geometry.

Invariant Construction of the Tantrakyma Potential

It is necessary to demonstrate that the specific form of the Tantrakyma potential—as given in Definition 9.333 and eq. (9.133)—arises as the unique, minimal choice consistent with the IQR principles (stability, symmetry, and minimality) and with the operator–geometric structure of the matter sector (Definition 9.301 and Remark 9.303).

1. Field content and representation in the dual space

The Tantrakyma field is an operator-valued *covector* on the vacuum manifold, $\hat{\boldsymbol{\Theta}}(x) \in T^*_{|\Omega(x))}\mathcal{M}_{\mathrm{vac}} \otimes \mathrm{End}(\mathcal{H}_{\mathrm{em}})$ (cf. Proposition 9.292 and Definitions 6.142 and 9.301). Accordingly,

scalar operator invariants must be built using the cotangent *co-metric* g_C (with g_C^{-1} its pseudoinverse; cf. Remark 9.303). The fundamental invariant is

$$\mathsf{X}(x) \;:=\; g_C\Big(\hat{\mathbf{\Theta}}(x),\, \hat{\mathbf{\Theta}}(x)\Big) \;\in\; \mathrm{End}(\mathcal{H}_{\mathrm{em}}),$$

which plays for Tantrakyma the role that $g_T(\cdot,\cdot)$ plays for the Gaiamiti sector (Section 9.17). Scalar *densities* for the action are then obtained by tracing over the fiber algebra, as in $\mathcal{V}_{\Theta,\mathrm{density}} = \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\{\mathcal{V}_\Theta\}$ (eq. (9.131) and Definition 9.310). (The pseudoinverse g_C^{-1} is useful for dualizations/Legendre maps but is *not* the primitive building block for the mass/quartic potential.)

2. Constraints of stability, symmetry, and minimality

Stability. Boundedness from below requires the highest-order coefficient to be positive and, for a quartic truncation, $\lambda_\Theta > 0$. In particular, the X^2 coefficient controls large-amplitude behavior in the effective potential $V_\Theta = \langle \mathcal{V}_\Theta \rangle$ (eq. (9.134)).

Symmetry. To permit spontaneous (rather than explicit) symmetry breaking, the potential must respect the internal fiber symmetries (trace-preserving $*$-automorphisms; cf. Remark 9.313) and any discrete matter symmetries such as $\mathbb{Z}_2 : \hat{\mathbf{\Theta}} \mapsto -\hat{\mathbf{\Theta}}$. These conditions enforce that $\mathcal{V}_\Theta$ is a function only of the invariant $\mathsf{X} = g_C(\hat{\mathbf{\Theta}}, \hat{\mathbf{\Theta}})$.

Minimality and universality. Keeping operators up to mass-dimension four (QFT renormalizability; universality of the quartic Landau functional) selects the quartic polynomial in X as the lowest-order choice able to produce a nontrivial vacuum structure and mass generation, in precise analogy with the Gaiamiti construction (Section 9.17).

Putting these together, the most general operator-valued potential consistent with the symmetries is

$$\mathcal{V}_\Theta(\hat{\mathbf{\Theta}}) \;=\; f(\mathsf{X}), \qquad f(\mathsf{X}) = \sum_{n\geq 0} d_{2n}\, \mathsf{X}^{2n},$$

with $d_{2n} \in \mathbb{R}$ and $d_{2N} > 0$ at the highest retained order. The *minimal* (dimension ≤ 4) choice is precisely the quartic polynomial displayed in eq. (9.133).

Conclusion: the unique minimal potential for the matter sector

The unique, minimal, Hermitian and symmetry-respecting form is (cf. Definition 9.333)

$$\mathcal{V}_\Theta(\hat{\mathbf{\Theta}}) = \tfrac{1}{2}\, m_\Theta^2\, \mathsf{X} \;+\; \tfrac{\lambda_\Theta}{4!}\, \mathsf{X}^2 \;+\; \mathcal{O}\Big(\mathsf{X}^3/\Lambda_\Theta^2\Big), \qquad \mathsf{X} := g_C(\hat{\mathbf{\Theta}}, \hat{\mathbf{\Theta}}).$$

On a certified plateau $\mathfrak{P}$ with tags $(\lambda_{\mathrm{UV}}, \ell)$, this feeds the effective potential V_Θ (eq. (9.134)) whose minima $\bar{\Theta}$ solve the EL system (eq. (9.148)). Linearizing about $\bar{\Theta}$ produces the typed mass/curvature operators via the Hessian restriction (eqs. (9.124) and (9.175)), which in turn predict NG/pNG versus gapped kymons by type $[\chi] \in \{\mathrm{CK}, \mathrm{T}_\pm, \mathrm{L}\}$ (Definition 9.295). These predictions are cross-checked by the parasymmetric meters and the typed quadrant decision in the T–L–O pipeline (Section 9.17.11).

The Tantrakyma Field Potential

Definition 9.335 (Tantrakyma Field Potential)**.** Let $\hat{\mathbf{\Theta}}(x) \in \mathrm{End}(\mathcal{H}_{\mathrm{em}}) \otimes \mathcal{E}_{\mathrm{exc}}(x)$ be the operator-valued Tantrakyma field, where $\mathcal{E}_{\mathrm{exc}}(x) \subseteq T^*_{|\Omega(x))}\mathcal{M}_{\mathrm{vac}}$ denotes the excitation subspace of the cotangent fiber (cf. Remark 9.303, Proposition 9.292, and Definition 9.307). The *operator-valued potential* $\mathcal{V}_\Theta(\hat{\mathbf{\Theta}}(x))$ is a Hermitian functional built from the cotangent invariant

$$\mathsf{X}(x) := g_C\Big(\hat{\mathbf{\Theta}}(x), \hat{\mathbf{\Theta}}(x)\Big) \;\in\; \mathrm{End}(\mathcal{H}_{\mathrm{em}})$$

(cf. Remark 9.303), namely

$$\boxed{\mathcal{V}_\Theta\Big(\hat{\mathbf{\Theta}}(x)\Big) := \tfrac{1}{2}\, m_\Theta^2\, \mathsf{X}(x) \;+\; \tfrac{\lambda_\Theta}{4!}\, \mathsf{X}(x)^2 \;+\; \mathcal{O}\Big(\mathsf{X}(x)^3/\Lambda_\Theta^2\Big),} \tag{9.176}$$

with real m_Θ^2 and $\lambda_\Theta > 0$. The scalar density that enters the Lagrangian (cf. eq. (9.131) and Definition 9.301) is obtained by tracing over the fiber algebra:

$$\mathcal{V}_{\Theta,\mathrm{density}}(x) \;=\; \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\Big\{\mathcal{V}_\Theta(\hat{\mathbf{\Theta}}(x))\Big\}.$$

Physical Interpretation: from potential to particle properties The phenomenologically relevant potential is the scalar function obtained by taking the diagnostic realized reference configuration expectation value (cf. Definitions 9.307 and 9.310):

$$V_\Theta(x) \;:=\; \Big\langle \mathcal{V}_\Theta\Big(\hat{\mathbf{\Theta}}(x)\Big)\Big\rangle_{\mathfrak{P}}, \qquad \mathcal{L}_\Theta(x) = \mathcal{T}_\Theta(x) + \mathcal{V}_{\Theta,\mathrm{density}}(x), \quad \mathcal{V}_{\Theta,\mathrm{density}} = \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\{\mathcal{V}_\Theta\}. \tag{9.177}$$

Minima of V_Θ determine realized backgrounds $\bar{\Theta}$ satisfying the Euler–Lagrange system (eq. (9.148)); linearizing about $\bar{\Theta}$ yields typed mass/curvature operators via the potential Hessian restricted to the Tantrakyma projectors (Definition 9.295 and eq. (9.124)), which in turn set the kymon spectrum used by Parasymmetric T–L–O (Section 9.17.11).

- **Mass parameter m_Θ^2.** The sign controls vacuum structure in V_Θ: $m_\Theta^2 > 0 \Rightarrow \bar{\Theta} = 0$ (symmetric phase); $m_\Theta^2 < 0 \Rightarrow \bar{\Theta} \neq 0$ (SSB). In the linearized EL equation (eq. (9.143)) it sources the quadratic (mass) block of $\mathcal{K}^{[\chi]}_{\Theta\Theta}$.
- **Quartic coupling λ_Θ.** With $\lambda_\Theta > 0$ the potential is bounded below and provides the leading self-interaction controlling kymon–kymon scattering and the curvature of V_Θ at the minima (typed gaps).
- **Symmetry breaking.** A "Mexican-hat" profile ($m_\Theta^2 < 0$, $\lambda_\Theta > 0$) realizes spontaneous symmetry breaking consistent with fiber $*$-automorphisms (trace-preserving) and discrete $\mathbb{Z}_2$ symmetry ($\hat{\mathbf{\Theta}} \mapsto -\hat{\mathbf{\Theta}}$) (cf. Remark 9.313). NG/pNG versus gapped outcomes by type follow from $\mathrm{spec}\Big(\mathcal{K}^{[\chi]}_{\Theta\Theta}\Big)$ and are cross-checked by T–L–O meters.

In summary, $\mathcal{V}_\Theta$ is the engine of matter phenomenology in the IQR framework: together with the chronogeometric kinetic pairing (eq. (9.130)) it determines the realized vacuum, typed mass operators, and thus the spectrum and interactions of kymons in a diagnostic realized reference configuration-certified, BRST/Nielsen-safe manner.

Holosymmetric Wordwise Tantrakyma Field Potential

We now dress the standard Tantrakyma potential (Section 9.18.5) in the *holosymmetric wordwise* (HS) framework developed in Chapter VI, using the HS co–metric and bundle data (Definition 9.316 and Sections 6.4.1 and 6.6.1) and the Relational Levi–Civita compatibility on the base (Definition 6.7).

Definition 9.336 (HS fundamental invariant for Tantrakyma)**.** The HS invariant attached to the covector field $\hat{\boldsymbol{\Theta}}$ is

$$\boxed{\mathsf{X}^{\mathrm{HS}}(x) \;:=\; g_C^{\mathrm{HS}}\Big(\hat{\boldsymbol{\Theta}}(x),\, \hat{\boldsymbol{\Theta}}(x)\Big) \in \mathrm{End}(\mathcal{H}_{\mathrm{em}}),} \tag{9.178}$$

as introduced in Definition 9.316 and eq. (9.151). Here g_C^{HS} is the HS words–dressed co–metric on $T^*\mathcal{M}_{\mathrm{vac}}$.

Definition 9.337 (HS Tantrakyma potential operator and scalar density)**.** The *HS operator potential* is any Hermitian functional of X^{HS}; in particular, the quartic (plus higher) form reads (cf. eqs. (9.152) and (9.172))

$$\boxed{\mathcal{V}_\Theta^{\mathrm{HS}}(\hat{\boldsymbol{\Theta}}) \;:=\; \tfrac{1}{2}\, m_\Theta^2\, \mathsf{X}^{\mathrm{HS}} \;+\; \tfrac{\lambda_\Theta}{4!}\, (\mathsf{X}^{\mathrm{HS}})^2 \;+\; \sum_{k\geq 3} \frac{d_k}{k!}\, (\mathsf{X}^{\mathrm{HS}})^k, \qquad \lambda_\Theta > 0,} \tag{9.179}$$

with real couplings $m_\Theta^2, \lambda_\Theta, d_k$. The corresponding *HS scalar potential density* is the field–dialectic trace (Definition 6.45 and eq. (6.99)):

$$\boxed{\mathcal{V}^{\mathrm{HS}}_{\Theta,\mathrm{density}}(x) \;:=\; \mathrm{Tr}_{\mathrm{FD}}\Big\{\mathcal{V}_\Theta^{\mathrm{HS}}(\hat{\boldsymbol{\Theta}}(x))\Big\}.} \tag{9.180}$$

Remark 9.338 (HS action contribution and reduction to standard)**.** The potential sector of the HS action uses the HS invariant measure $d\boldsymbol{\mu} = d^d x\, \boldsymbol{\varpi}$, $\boldsymbol{\varpi} = (\det_{\mathrm{HS}}(-\boldsymbol{g}^{\mathrm{Lor}}))^{1/2}$ (Definition 6.34):

$$\boxed{\widehat{\boldsymbol{S}}_{\Theta,\mathrm{pot}} \;:=\; \int_{\mathcal{X}_q} d\boldsymbol{\mu}(x)\; \mathcal{V}^{\mathrm{HS}}_{\Theta,\mathrm{density}}(x).} \tag{9.181}$$

Applying the expectation/volume push–throughs (Remark 6.35) recovers the standard scalar density $\mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\{\mathcal{V}_\Theta\}$ in Section 9.18.5.

Proposition 9.339 (HS gauge/frame invariance and diffeo covariance (Tantrakyma))**.** *Under HS modal frame changes $g(x) \in U(n)$ and vacuum–tangent rotations $R(x) \in O(r)$ acting on covectors by $\hat{\Theta}_i \mapsto \mathrm{Ad}_g(\hat{\Theta}_j)(R^{-1})^j{}_i$ (cf. Proposition 9.331),*

$$g_C^{\mathrm{HS}}(\mathrm{Ad}_g\, X \cdot R^{-1},\, \mathrm{Ad}_g\, Y \cdot R^{-1}) \;=\; g_C^{\mathrm{HS}}(X, Y),$$

hence X^{HS} and $\mathcal{V}_\Theta^{\mathrm{HS}}$ are invariant. Moreover, $\mathcal{V}^{\mathrm{HS}}_{\Theta,\mathrm{density}}$ is a base scalar density for diffeomorphisms compatible with $\nabla^{(\mathrm{rel})}$ (Definition 6.7).

Proof. g_C^{HS} is a fibrewise Hermitian bilinear preserved by HS unitaries and orthogonal coframe rotations (Definition 9.316). For covector indices the orthogonal action is by R^{-1} (cf. Definition 9.328 and Proposition 9.331); invariance then gives $\mathsf{X}^{\mathrm{HS}} \mapsto \mathsf{X}^{\mathrm{HS}}$. Any real functional of X^{HS} is therefore invariant, proving gauge/frame invariance. The field–dialectic trace produces a c–number and carries no free base index (Definition 6.45 and eq. (6.99)); compatibility of $\nabla^{(\mathrm{rel})}$ with $\bar{\Sigma}$ ensures that $\mathcal{V}^{\mathrm{HS}}_{\Theta,\mathrm{density}}$ transforms as a scalar density under the allowed diffeomorphisms. □

Lemma 9.340 (First variation and stationarity (Tantrakyma, HS))**.** *Let $\delta\hat{\boldsymbol{\Theta}}$ be a compactly supported variation in the HS cotangent fibre. Then*

$$\boxed{\delta\,\mathcal{V}^{\mathrm{HS}}_{\Theta,\mathrm{density}} = \mathrm{Tr}_{\mathrm{FD}}\left\{\left(\tfrac{1}{2}m_\Theta^2 + \tfrac{\lambda_\Theta}{3!}\,\mathsf{X}^{\mathrm{HS}} + \sum_{k\geq 3}\tfrac{d_k}{(k-1)!}\,(\mathsf{X}^{\mathrm{HS}})^{k-1}\right)\delta\mathsf{X}^{\mathrm{HS}}\right\},} \tag{9.182}$$

with

$$\boxed{\delta\mathsf{X}^{\mathrm{HS}} = g_C^{\mathrm{HS}}(\delta\hat{\boldsymbol{\Theta}},\hat{\boldsymbol{\Theta}}) + g_C^{\mathrm{HS}}(\hat{\boldsymbol{\Theta}},\delta\hat{\boldsymbol{\Theta}}) = 2\,\mathrm{Re}\,g_C^{\mathrm{HS}}(\hat{\boldsymbol{\Theta}},\delta\hat{\boldsymbol{\Theta}})\,.} \tag{9.183}$$

Proof. Use $\mathcal{V}^{\mathrm{HS}}_\Theta = F(\mathsf{X}^{\mathrm{HS}})$ with $F'(s) = \frac{1}{2}m_\Theta^2 + \frac{\lambda_\Theta}{3!}s + \sum_{k\geq 3}\frac{d_k}{(k-1)!}s^{k-1}$ and the chain rule inside $\mathrm{Tr}_{\mathrm{FD}}$ (cf. Definition 6.45 and Remark 6.187). Treat g_C^{HS} as fixed under field variation; bilinearity and Hermiticity of g_C^{HS} (Definition 9.316) give (9.183). □

Corollary 9.341 (Boundedness and vacuum types (Tantrakyma, HS))**.** *If $\lambda_\Theta > 0$ and the higher d_k satisfy standard positivity/coercivity bounds, then $\mathcal{V}^{\mathrm{HS}}_\Theta$ is bounded below. For $m_\Theta^2 > 0$ the symmetric vacuum $\hat{\boldsymbol{\Theta}} = 0$ is stationary; for $m_\Theta^2 < 0$ a "Mexican-hat" structure arises in the HS covector norm X^{HS}, with symmetry breaking analyzed by the HS vacuum manifold tools of Chapter VI (Section 6.1.9).*

Proof. Identical to the Gaiamiti argument in Corollary 9.241, with $s = \mathsf{X}^{\mathrm{HS}} \geq 0$ and $F(s)$ from (9.179). Coercivity (highest even power positive) gives boundedness. At $\hat{\boldsymbol{\Theta}} = 0$, $\delta\mathsf{X}^{\mathrm{HS}} = 0$ so stationarity is automatic; the second variation is $m_\Theta^2\,\mathrm{Tr}_{\mathrm{FD}}\{\,g_C^{\mathrm{HS}}(\delta\hat{\boldsymbol{\Theta}},\delta\hat{\boldsymbol{\Theta}})\} \geq 0$ for $m_\Theta^2 > 0$, and negative for $m_\Theta^2 < 0$, implying a nonzero minimizing radius determined by $F'(s_\star) = 0$ as in the Gaiamiti case, now with $s_\star = \mathsf{X}^{\mathrm{HS}}$. □

Proposition 9.342 (HS $\leftrightarrow$ standard equivalence for the Tantrakyma potential)**.** *Let $\mathsf{X}^{\mathrm{HS}} := g_C^{\mathrm{HS}}(\hat{\boldsymbol{\Theta}},\hat{\boldsymbol{\Theta}})$ (Definition 9.316 and eq. (9.178)) and $\mathsf{X} := g_C(\hat{\boldsymbol{\Theta}},\hat{\boldsymbol{\Theta}})$ (Section 9.18.5). Then, on a certified plateau and after the HS expectation/volume push–through (Definition 6.45, eq. (6.99), and Remark 6.35),*

$$\boxed{\langle\mathsf{X}^{\mathrm{HS}}\rangle \;=\; \mathsf{X}, \qquad \langle\mathrm{Tr}_{\mathrm{FD}}\{(\mathsf{X}^{\mathrm{HS}})^k\}\rangle \;=\; \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\{\mathsf{X}^k\}}$$

and the HS operator potential (Definition 9.337 and eq. (9.179))

$$\mathcal{V}^{\mathrm{HS}}_\Theta(\hat{\boldsymbol{\Theta}}) = \tfrac{1}{2}m_\Theta^2\,\mathsf{X}^{\mathrm{HS}} + \tfrac{\lambda_\Theta}{4!}(\mathsf{X}^{\mathrm{HS}})^2 + \sum_{k\geq 3}\frac{d_k}{k!}(\mathsf{X}^{\mathrm{HS}})^k$$

reduces to the standard operator potential (Definition 9.335 and eq. (9.176))

$$\boxed{\langle \mathcal{V}_\Theta^{\mathrm{HS}} \rangle \;=\; \mathcal{V}_\Theta \;=\; \tfrac{1}{2} m_\Theta^2\, \mathsf{X} + \tfrac{\lambda_\Theta}{4!}\, \mathsf{X}^2 + \mathcal{O}(\mathsf{X}^3/\Lambda_\Theta^2)}$$

with the same parameters $m_\Theta^2, \lambda_\Theta$ (and $d_k \leftrightarrow$ higher–order Wilson coefficients). Moreover, the HS scalar density $\mathcal{V}^{\mathrm{HS}}_{\Theta,\mathrm{density}} = \mathrm{Tr}_{\mathrm{FD}}\{\mathcal{V}_\Theta^{\mathrm{HS}}\}$ (Definition 9.337 and eq. (9.180)) pushes through to the standard density $\mathcal{V}_{\Theta,\mathrm{density}} = \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\{\mathcal{V}_\Theta\}$ used in Section 9.18.5.

Sketch. (i) By construction g_C^{HS} words–dresses g_C and is preserved by HS unitaries and $O(r)$ coframe rotations (Definition 9.316 and Proposition 9.331); thus X^{HS} is the HS counterpart of X. (ii) The FD trace reduces to the fibre trace under the expectation/volume push–through (Definition 6.45, eq. (6.99), and Remark 6.35), yielding the displayed equalities for all powers. (iii) Because the standard potential (Definition 9.335) is a real functional of X only, the HS potential (Definition 9.337)—a real functional of X^{HS}—reduces to it term by term, with identical couplings. The excitation subspace constraint $\mathcal{E}_{\mathrm{exc}}(x) \subseteq T^*_{|\Omega\rangle}\mathcal{M}_{\mathrm{vac}}$ (Definition 9.307) is respected in HS via the same band-limited fibres. □

The Kymon and the Emergence of the Higgs Boson

Definition 9.343 (Kymon and Higgs identification via typed, parasymmetric T–L–O)**.** Consider a background–fluctuation split of the Tantrakyma field,

$$\hat{\mathbf{\Theta}}(x) = \bar{\Theta}(x)\, \mathbb{I}_x + \delta\hat{\mathbf{\Theta}}(x), \qquad \big\langle \delta\hat{\mathbf{\Theta}}(x) \big\rangle_{\mathfrak{P}} = 0,$$

with $\bar{\Theta}$ a stationary solution of the EL system (eq. (9.148)) minimizing the effective potential V_Θ (cf. eqs. (9.134) and (9.176)). Work on a certified plateau $\mathfrak{P}$ with tags $(\lambda_{\mathrm{UV}}, \ell)$ and BRST projection to $H^0(\hat{Q}_{\mathrm{BRST}})$ (cf. Definitions 9.307 and 9.310).

(i) Kymons (definition). Let $\Pi^{[\chi]}_{\Theta\Theta}$ denote the Tantrakyma typed projectors (Definition 9.295). The *kymons* are the one-particle excitations associated with the poles of the typed, retarded two-point kernels $\Delta^{[\chi]}_{\Theta\Theta}$ on $H^0(\hat{Q}_{\mathrm{BRST}})$, i.e. the solutions of the linear response equation

$$\boxed{\Big(\mathbb{K}^{[\chi]}_\Theta + \mathcal{K}^{[\chi]}_{\Theta\Theta}\Big)\, \delta\hat{\mathbf{\Theta}}^{[\chi]} = \hat{\mathcal{J}}^{[\chi]}, \qquad \delta\hat{\mathbf{\Theta}}^{[\chi]} := \Pi^{[\chi]}_{\Theta\Theta}\, \delta\hat{\mathbf{\Theta}},}\tag{9.184}$$

where $\mathbb{K}^{[\chi]}_\Theta$ is the typed kinetic block and $\mathcal{K}^{[\chi]}_{\Theta\Theta}$ is the typed potential Hessian (cf. eqs. (9.143) and (9.175)). A simple pole of $\Delta^{[\chi]}_{\Theta\Theta}(k)$ with positive residue on $H^0(\hat{Q}_{\mathrm{BRST}})$ certifies a physical kymon.

(ii) Higgs (radial) mode (construction). Let $\mathsf{X} := g_C(\hat{\mathbf{\Theta}}, \hat{\mathbf{\Theta}})$ be the Tantrakyma invariant (cf. Remark 9.303) and define the *radial (amplitude) direction* at the realized background by the Fréchet derivative

$$\delta\mathsf{X}\Big|_{\bar{\Theta}} \;=\; 2\, g_C\Big(\delta\hat{\mathbf{\Theta}}, \bar{\Theta}\Big), \qquad \text{so that (via the } g_C\text{–Riesz isomorphism)} \;\; \hat{\mathsf{R}} \;\propto\; \bar{\Theta}.$$

Let Π_{H} be the g_C–orthogonal projector onto $\mathrm{span}\{\widehat{\mathsf{R}}\}$ (normalized within the typed basis and commuting with the triadic projectors). The *Higgs boson* is the scalar (singlet) kymon obtained by restricting the Hessian to this radial direction:

$$\boxed{\mathcal{K}_{\mathrm{H}} := \Pi_{\mathrm{H}}\, \frac{\delta^2 \mathcal{V}_\Theta}{\delta\hat{\mathbf{\Theta}}\,\delta\hat{\mathbf{\Theta}}}\bigg|_{\bar{\Theta}}\, \Pi_{\mathrm{H}}, \qquad M_{\mathrm{H}}^2 \;\in\; \mathrm{spec}\big(\mathcal{K}_{\mathrm{H}}\big),} \tag{9.185}$$

with the corresponding typed propagator exhibiting a single *gapped* pole at $k^2 = M_{\mathrm{H}}^2 > 0$ and no $k^2 = 0$ pole (contrast with NG/pNG directions).

Riesz map induced by g_C. On the Tantrakyma bundle $\mathcal{E}_\Theta$ with co-metric g_C (cf. Remark 9.303), define the Riesz/Legendre isomorphism

$$\mathcal{G}_C :\; \mathcal{E}_\Theta \longrightarrow \mathcal{E}_\Theta^*, \qquad \langle A,\; \mathcal{G}_C(B)\rangle \;=\; g_C(A,B) \quad (\forall\, A,B \in \mathcal{E}_\Theta),$$

with inverse $\mathcal{G}_C^{-1} : \mathcal{E}_\Theta^* \to \mathcal{E}_\Theta$. (Operator-valued details are compatible with the diagnostic realized reference configuration/connection choices; cf. Definition 9.310 and eq. (9.176).)

Radial direction is collinear with $\bar{\Theta}$. Let

$$\mathsf{X} := g_C(\hat{\mathbf{\Theta}}, \hat{\mathbf{\Theta}}) \quad (\text{cf. } \textit{Remark } 9.303).$$

At the realized background $\bar{\Theta}$,

$$\delta\mathsf{X}\big|_{\bar{\Theta}}(\delta\hat{\mathbf{\Theta}}) = 2\, g_C\Big(\delta\hat{\mathbf{\Theta}}, \bar{\Theta}\Big) = 2\,\Big\langle \delta\hat{\mathbf{\Theta}},\, \mathcal{G}_C(\bar{\Theta})\Big\rangle.$$

Thus the gradient covector is $2\,\mathcal{G}_C(\bar{\Theta})$, and mapping back with the inverse Riesz map gives

$$\mathcal{G}_C^{-1}\Big(\delta\mathsf{X}\big|_{\bar{\Theta}}\Big) \;=\; 2\,\bar{\Theta}.$$

Hence the amplitude ("radial") direction at $\bar{\Theta}$ is collinear with $\bar{\Theta}$ itself, which the text summarizes as $\widehat{\mathsf{R}} \propto \bar{\Theta}$ (cf. Definition 9.343). A typed, normalized representative that commutes with the triadic projectors is

$$\widehat{\mathsf{R}} \;:=\; \frac{\Pi^{[\chi]}_{\Theta\Theta}\,\bar{\Theta}}{\sqrt{g_C\Big(\Pi^{[\chi]}_{\Theta\Theta}\bar{\Theta},\; \Pi^{[\chi]}_{\Theta\Theta}\bar{\Theta}\Big)}},$$

so $\widehat{\mathsf{R}} \in \mathrm{Ran}\,\Pi^{[\chi]}_{\Theta\Theta}$ and $[\widehat{\mathsf{R}}, \Pi_{\mathrm{corr/vac/spec}}] = 0$.

One–dimensional g_C–orthogonal projector. Define the rank–1 Riesz projector onto $\mathrm{span}\{\widehat{\mathsf{R}}\}$ by

$$\Pi_{\mathrm{H}}(V) \;=\; \frac{g_C\Big(V, \widehat{\mathsf{R}}\Big)}{g_C\Big(\widehat{\mathsf{R}}, \widehat{\mathsf{R}}\Big)}\,\widehat{\mathsf{R}}, \qquad \Pi_{\mathrm{H}}^2 = \Pi_{\mathrm{H}} = \Pi_{\mathrm{H}}^\dagger \text{ (with respect to } g_C).$$

Because $\widehat{\mathsf{R}} \propto \bar{\Theta}$, Π_{H} selects the amplitude/radial direction.

Higgs mass operator and typed linear response. Within the typed, BRST–clean linear response

$$\left(\mathbb{K}_{\Theta}^{[\chi]} + \mathcal{K}_{\Theta\Theta}^{[\chi]}\right) \delta\hat{\mathbf{\Theta}}^{[\chi]} = \hat{\mathcal{J}}^{[\chi]} \qquad \text{(cf. eq. (9.184))},$$

restrict the potential Hessian to the radial line:

$$\boxed{\mathcal{K}_{\mathrm{H}} \;:=\; \Pi_{\mathrm{H}}\, \frac{\delta^2 \mathcal{V}_{\Theta}}{\delta\hat{\mathbf{\Theta}}\,\delta\hat{\mathbf{\Theta}}}\bigg|_{\bar{\Theta}}\, \Pi_{\mathrm{H}}} \qquad \text{(cf. eq. (9.185))}.$$

By the typed spectral criterion, the *Higgs boson* is the scalar/singlet kymon whose propagator exhibits a single gapped pole at $k^2 = M_{\mathrm{H}}^2 > 0$, with $M_{\mathrm{H}}^2 \in \mathrm{spec}(\mathcal{K}_{\mathrm{H}})$; NG/pNG tangential directions satisfy $g_C(\delta\hat{\mathbf{\Theta}}, \bar{\Theta}) = 0$ and thus do not probe this radial eigenvalue.

Parasymmetric T–L–O certification (typed, triadic). For each type $[\chi] \in \{\mathrm{CK}, \mathrm{T}_{\pm}, \mathrm{L}\}$, assemble the SynTh meters (cf. eqs. (9.126) to (9.128)) using $\Delta_{\Theta\Theta}^{[\chi]}$ and the diagnostic realized reference configuration tags:

- $v_{[\chi]}^{(L)} = \left|\mathrm{Res}_{k^2=0}\, \Delta_{\Theta\Theta}^{[\chi]}(k)\right|$ diagnoses NG/pNG content (gapped $\Rightarrow v_{[\chi]}^{(L)} = 0$);
- $v_{[\chi]}^{(S)} = \left|\Delta_{\Theta\Theta}^{[\chi]}(t \to \infty; \mathbf{x} = 0)\right|_{H^0(\hat{Q}_{\mathrm{BRST}})}$ certifies long-time order;
- $v_{[\chi]}^{(LS)}$ measures helicity-resolved cross-channel gain.

Apply the typed quadrant predicate (cf. Definition 9.189 and Proposition 9.197) and stamp Type–Location–Onset entries (cf. Definitions 9.190 and 9.198) for the *Higgs channel* Π_{H} and for Goldstone channels. Consistency requires agreement, within plateau tolerances, between (a) NG/gapped predictions from $\mathrm{spec}\left(\mathcal{K}_{\Theta\Theta}^{[\chi]}\right)$ and (b) the quadrant ledger (Ch. 8).

SSB pattern and spectrum (diagnostic realized reference configuration semantics). For the quartic potential (9.176), if $m_{\Theta}^2 < 0$ and $\lambda_{\Theta} > 0$, the effective potential V_{Θ} (cf. eq. (9.134)) has nonzero minima $\bar{\Theta}$. Then:

1. *Goldstone directions* are tangent to the vacuum manifold and yield $v_{[\chi]}^{(L)} \neq 0$ with a pole at $k^2 = 0$, matching zero/pseudo-zero eigenvalues of $\mathcal{K}_{\Theta\Theta}^{[\chi]}$.
2. The *Higgs (radial) mode* is g_C–orthogonal to the vacuum manifold and exhibits $v_{[\chi]}^{(L)} = 0$ together with a single gapped pole at $k^2 = M_{\mathrm{H}}^2$, where M_{H}^2 is given by (9.185). Its typed quadrant outcome is gapped, and its Location band is determined by the pole position and residue (IR/meso per Definition 9.190).

Operational identification.

1. Solve the EL system for $\bar{\Theta}$ and compute $\mathcal{K}_{\Theta\Theta}^{[\chi]}$ (cf. eq. (9.175)).
2. Build Π_{H} from the g_C–radial direction $\hat{\mathsf{R}}$ and evaluate $\mathcal{K}_{\mathrm{H}}$ to obtain M_{H}^2 (cf. eq. (9.185)).

3. Compute $\Delta^{[\chi]}_{\Theta\Theta}$ on $H^0(\hat{Q}_{\mathrm{BRST}})$; extract meters $v^{(L)}, v^{(S)}, v^{(LS)}$ (cf. Section 9.17.11).

4. Decide quadrants (cf. Definition 9.189 and Proposition 9.197); stamp T–L–O for Higgs and NG channels (cf. Definitions 9.190 and 9.198).

This pipeline produces a *T–L–O–certified* emergence of the Higgs boson as the radial, scalar kymon in the Tantrakyma sector, consistent with symmetry breaking in V_Θ, BRST/Nielsen hygiene on the diagnostic realized reference configuration, and typed/triadic instrumentation (cf. Definition 9.307 and Remark 9.313).

Interpretative Clarification of the Formalism The unified construction (Definition 9.343) imposes a precise hierarchy of notions, certified on the diagnostic realized reference configuration $\mathfrak{P}$ by the typed, parasymmetric T–L–O pipeline (Section 9.17.11):

1. **Kymon (general notion).** The *kymon* is the fundamental quantum of the Tantrakyma field $\hat{\mathbf{\Theta}}$ (Definition 9.301). Operationally, it is any physical one-particle excitation corresponding to a simple pole of a typed, retarded two-point kernel $\Delta^{[\chi]}_{\Theta\Theta}$ on the BRST physical space $H^0(\hat{Q}_{\mathrm{BRST}})$, with positive residue (eq. (9.184) and Definition 9.295). This definition depends on: (i) the background $\bar{\Theta}$ solving the EL system (eq. (9.148)); (ii) the typed potential Hessian $\mathcal{K}^{[\chi]}_{\Theta\Theta}$ (eq. (9.175)); and (iii) the chronogeometric kinetic pairing in $\mathcal{T}_\Theta$ (eq. (9.130)).

2. **Higgs boson (specific species).** The *Higgs* is the *radial, scalar* kymon: the gapped mode obtained by restricting the potential Hessian to the amplitude direction of the invariant $\mathsf{X} = g_C(\hat{\mathbf{\Theta}}, \hat{\mathbf{\Theta}})$ at the realized vacuum $\bar{\Theta}$ (Remark 9.303 and eq. (9.185)). Its typed propagator shows a single pole at $k^2 = M_{\mathrm{H}}^2 > 0$ and no $k^2 = 0$ pole (contrast with NG/pNG directions), and its T–L–O stamp is *gapped* with an IR/mesoscopic location set by M_{H} (Section 9.17.11).

Unification within IQR (sector semantics). The matter and geometry sectors admit parallel, typed interpretations:

- **Gaiamiti/Apeiron → Graviton.** In the geometry sector, the Apeiron (quantum of the Gaiamiti field) organizes massless excitations in the appropriate typed channels; the graviton arises as the massless (Goldstone-like) mode within the chronogeometric arena $\bar{\Sigma}$ (Section 9.17).

- **Tantrakyma/Kymon → Higgs.** In the matter sector, the kymon organizes the typed spectrum of $\delta\hat{\mathbf{\Theta}}$; when V_Θ exhibits SSB (e.g. $m^2_\Theta < 0,\ \lambda_\Theta > 0$ in eqs. (9.133) and (9.134)), the radial typed mode is the Higgs, while tangential modes realize NG/pNG channels (eq. (9.175)).

Thus, mass generation for matter is not an add-on but a consequence of the *cotangent* geometry of the Tantrakyma potential and its coupling to the full chronogeometric background through $\bar{\Sigma}^{\mu\nu}$ (eq. (9.130)). Diagnostic realized reference configuration certification (BRST/Nielsen flatness) guarantees gauge-parameter independence of the spectral claims (Definition 9.307 and Remark 9.313).

On nomenclature. The neologism *kymon* designates the fundamental quantum of the Tantrakyma field, with the compound *tantrakyma* formed from Sanskrit *tantra* ("warp, loom; framework") and Greek κῦμα (*kyma*, "wave, undulation"). The intent is "*woven–wave*" (or "warp of waves"). The particle suffix "−on" follows standard usage. See [66, 74, 75] for *tantra* and [80] for κῦμα; cf. OED entries on scientific −on formations [302, 296].[7]

Derivation of the Mass Operator for the Kymon Field

Our goal is to extract the *mass operator* controlling linearized Tantrakyma fluctuations from the potential $\mathcal{V}_\Theta$ in Definition 9.335. The derivation mirrors the geometry-side analysis for the Apeiron (cf. Section 9.17.1), now carried out on the cotangent/operator side and in the typed, diagnostic realized reference configuration-certified setting of Definition 9.307.

Setup and notation. Write the invariant

$$\mathsf{X}(x) := g_C\Big(\hat{\mathbf{\Theta}}(x), \hat{\mathbf{\Theta}}(x)\Big) \;\in\; \mathrm{End}(\mathcal{H}_{\mathrm{em}}),$$

and recall

$$\boxed{\mathcal{V}_\Theta = \tfrac{1}{2}\, m_\Theta^2\, \mathsf{X} + \tfrac{\lambda_\Theta}{4!}\, \mathsf{X}^2 + \cdots, \qquad \mathcal{V}_{\Theta,\mathrm{density}} = \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\{\mathcal{V}_\Theta\}.} \tag{9.186}$$

We treat $\hat{\mathbf{\Theta}}$ and $\hat{\mathbf{\Theta}}^\dagger$ as independent variables and use the sesquilinearity of g_C (antilinear in the first slot, linear in the second; Remark 9.303).

Gâteaux derivatives of the invariant. In components relative to a cotangent basis $\{\mathbf{e}^a\}$ with $\hat{\mathbf{\Theta}} = \sum_a \hat{\Theta}_a\, \mathbf{e}^a$ and $\mathsf{X} = \sum_{ab} \hat{\Theta}_a^\dagger\, g_{C\,ab}\, \hat{\Theta}_b$, one has

$$\frac{\delta \mathsf{X}}{\delta \hat{\Theta}_c^\dagger} = \sum_b g_{C\,cb}\, \hat{\Theta}_b \;=\; \Big(g_C \cdot \hat{\mathbf{\Theta}}\Big)_c, \qquad \frac{\delta \mathsf{X}}{\delta \hat{\Theta}_c} = \sum_a \hat{\Theta}_a^\dagger\, g_{C\,ac} \;=\; \Big(\hat{\mathbf{\Theta}}^\dagger \cdot g_C\Big)_c. \tag{9.187}$$

In compact operator form:

$$\frac{\delta \mathsf{X}}{\delta \hat{\mathbf{\Theta}}^\dagger} = g_C \cdot \hat{\mathbf{\Theta}}, \qquad \frac{\delta \mathsf{X}}{\delta \hat{\mathbf{\Theta}}} = \hat{\mathbf{\Theta}}^\dagger \cdot g_C.$$

First functional derivative (gradient)

Using (9.186) and (9.187),

$$\boxed{\frac{\delta \mathcal{V}_\Theta}{\delta \hat{\mathbf{\Theta}}^\dagger} = \Big(\tfrac{1}{2}\, m_\Theta^2 + \tfrac{\lambda_\Theta}{12}\, \mathsf{X}\Big)\Big(g_C \cdot \hat{\mathbf{\Theta}}\Big), \qquad \frac{\delta \mathcal{V}_\Theta}{\delta \hat{\mathbf{\Theta}}} = \Big(\hat{\mathbf{\Theta}}^\dagger \cdot g_C\Big)\Big(\tfrac{1}{2}\, m_\Theta^2 + \tfrac{\lambda_\Theta}{12}\, \mathsf{X}\Big),} \tag{9.188}$$

which are the covariant analogues of the familiar $V'(\phi^\dagger\phi)\,\phi$ rule.

[7] These naming remarks are mnemonic and do not affect the formal content; all physical identifications are made diagnostic realized reference configuration-wise via $\Delta^{[x]}_{\Theta\Theta}$, the typed Hessians, and the T–L–O meters (Section 9.17.11).

Second functional derivative (Hessian operator)

Differentiating (9.188) with respect to $\hat{\boldsymbol{\Theta}}$ yields a linear map on variations $\delta\hat{\boldsymbol{\Theta}}$:

$$\boxed{\frac{\delta^2\mathcal{V}_\Theta}{\delta\hat{\boldsymbol{\Theta}}^\dagger\,\delta\hat{\boldsymbol{\Theta}}} = \left(\tfrac{1}{2}\,m_\Theta^2 + \tfrac{\lambda_\Theta}{12}\,\mathsf{X}\right)\left(g_C\cdot\delta\hat{\boldsymbol{\Theta}}\right) \;+\; \tfrac{\lambda_\Theta}{12}\left\langle\hat{\boldsymbol{\Theta}},\,\delta\hat{\boldsymbol{\Theta}}\right\rangle_{g_C}\left(g_C\cdot\hat{\boldsymbol{\Theta}}\right),} \tag{9.189}$$

where $\langle A, B\rangle_{g_C} := A^\dagger\cdot g_C\cdot B$ is the cotangent inner product. In index-free operator notation, (9.189) is the sum of an *isotropic* piece $\propto g_C$ and a *rank-1* projector along $g_C\cdot\hat{\boldsymbol{\Theta}}$.

Mass operator at a realized background and typed restriction. Evaluating (9.189) at $\hat{\boldsymbol{\Theta}} = \bar{\Theta}$ produces the background mass operator $\mathcal{M}_\Theta[\bar{\Theta}]$. Its typed, BRST-clean version is the Hessian restriction (mixed derivative) of eq. (9.175):

$$\boxed{\mathcal{K}^{[\chi]}_{\Theta\Theta} := \Pi^{[\chi]}_{\Theta\Theta}\;\frac{\delta^2\mathcal{V}_\Theta}{\delta\hat{\boldsymbol{\Theta}}^\dagger\,\delta\hat{\boldsymbol{\Theta}}}\bigg|_{\bar{\Theta}}\;\Pi^{[\chi]}_{\Theta\Theta}, \qquad [\chi]\in\{\mathrm{CK},\mathrm{T}_\pm,\mathrm{L}\},} \tag{9.190}$$

which feeds the linear response equation $\left(\mathbb{K}^{[\chi]}_\Theta + \mathcal{K}^{[\chi]}_{\Theta\Theta}\right)\delta\hat{\boldsymbol{\Theta}}^{[\chi]} = \hat{\mathcal{J}}^{[\chi]}$ (eq. (9.143)).

Two canonical vacua (diagnostic realized reference configuration semantics)

(a) Symmetric phase ($\bar{\Theta}=0$). Here $\mathsf{X}(\bar{\Theta})=0$ and the projector term in (9.189) vanishes. The (band-limited) mass operator is

$$\boxed{\mathcal{M}_\Theta\big|_{\bar{\Theta}=0} = \tfrac{1}{2}\,m_\Theta^2\,g_C, \qquad \mathcal{K}^{[\chi]}_{\Theta\Theta}\big|_{\bar{\Theta}=0} = \tfrac{1}{2}\,m_\Theta^2\,\Pi^{[\chi]}_{\Theta\Theta}\,g_C\,\Pi^{[\chi]}_{\Theta\Theta}.} \tag{9.191}$$

Normalization note. With the convention (9.186), the symmetric-phase mass scale carries a factor $\frac{1}{2}$. If one prefers the convention where the parameter in V equals the physical quadratic coefficient, replace $m_\Theta^2 \mapsto 2\,m_\Theta^2$ in (9.186); all formulas then read with m_Θ^2 in place of $\frac{1}{2}m_\Theta^2$.

(b) SSB phase ($m_\Theta^2<0,\ \lambda_\Theta>0$). Let $x := \langle \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\mathsf{X}\rangle$. Minimizing $V_\Theta = \langle\mathcal{V}_\Theta\rangle$ (eq. (9.134)) gives $\partial V_\Theta/\partial x = 0 \Rightarrow x_\star = -6\,m_\Theta^2/\lambda_\Theta > 0$. At a realized $\bar{\Theta}$ with this amplitude, the Hessian splits into:

$$\begin{aligned}\text{Goldstone subspace (tangent to the vacuum manifold):}\quad & \mathcal{K}^{[\chi]}_{\Theta\Theta}\approx 0\ \ (\mathrm{NG/pNG}),\\ \text{Radial direction:}\quad & \mathcal{K}_{\mathrm{H}}\approx -2\,m_\Theta^2\,(\Pi_{\mathrm{H}}\,g_C\,\Pi_{\mathrm{H}}),\end{aligned} \tag{9.192}$$

i.e. the *Higgs (radial) mass* is

$$\boxed{M_{\mathrm{H}}^2 = -\,2\,m_\Theta^2 > 0,} \tag{9.193}$$

in agreement with the explicit projector construction of eq. (9.185). The NG/pNG channels exhibit a $k^2=0$ pole in the typed propagators, while the Higgs channel shows a single gapped pole at $k^2 = M_{\mathrm{H}}^2$.

Parasymmetric T–L–O cross-check (typed, triadic). Using $\Delta^{[\chi]}_{\Theta\Theta}$ on H^0, assemble the meters $(v^{(L)}, v^{(S)}, v^{(LS)})$ (eqs. (9.126) to (9.128)), apply the typed quadrant predicate (Definition 9.189 and Proposition 9.197), and stamp TLO$^{[\chi]}$ (Definition 9.299). Agreement, within plateau tolerances, between NG/gapped predictions from spec($\mathcal{K}^{[\chi]}_{\Theta\Theta}$) and the quadrant ledger certifies the identification of NG modes and the Higgs (radial) kymon (Definition 9.343).

9.18.6 Spontaneous Symmetry Breaking of the Tantrakyma Field

Quadrant context (Parasymmetric T–L–O overview). Before specializing to the broken phase, recall that phase identification in the Tantrakyma sector is performed *typed* and *diagnostic realized reference configuration-wise* via the SynTh meters $(v^{(L)}, v^{(S)}, v^{(LS)})$ of eqs. (9.126) to (9.128), the typed witness $\mathbf{W}^{[\Theta\Theta]}_{[\chi]}$ (Definition 9.296), and the canonical quadrant predicate $\mathrm{Quad}^{[\Theta\Theta]}_{[\chi]}$ (Definition 9.297 and Proposition 9.298). At fixed tags $(\lambda_{\mathrm{UV}}, \ell)$ on a certified plateau $\mathfrak{P}$ (Definition 9.307), the informal map is:

Table 9.2: Classification from long-wavelength/long-time behavior of the typed order-parameter correlator Δ. Decision criteria connect phenomenology to the visibility metrics Λ_{law} and Λ_{state}.

Quadrant	$\boldsymbol{v^{(L)}}$	$\boldsymbol{v^{(S)}}$	**Decision & diagnostic realized reference configuration signature**
QI	pole at $k^2{=}0$	$\lim_{t\to\infty} \lvert\Delta(t;\mathbf{0})\rvert \geq \sigma_0$	(A) $\lvert\Lambda_{\mathrm{law}}\rvert \geq 1-\varepsilon_{\mathrm{law}}$, $\lvert\Lambda_{\mathrm{state}}\rvert \geq 1-\varepsilon_{\mathrm{state}}\rvert$; (B) NG channels show a $k^2{=}0$ pole and long-time order.
QII	pole at $k^2{=}0$	$\lim_{t\to\infty} \lvert\Delta(t;\mathbf{0})\rvert < \sigma_0$	(A) $\lvert\Lambda_{\mathrm{law}}\rvert \geq 1-\varepsilon_{\mathrm{law}}$, $\lvert\Lambda_{\mathrm{state}}\rvert < 1-\varepsilon_{\mathrm{state}}\rvert$; (B) Law-only: pole present, no sustained order.
QIII	no clean $k^2{=}0$ pole	$\lim_{t\to\infty} \lvert\Delta(t;\mathbf{0})\rvert \geq \sigma_0$	(A) $\lvert\Lambda_{\mathrm{state}}\rvert \geq 1-\varepsilon_{\mathrm{state}}\rvert$, $\lvert\Lambda_{\mathrm{law}}\rvert < 1-\varepsilon_{\mathrm{law}}\rvert$ with healing $d_{\mathrm{law}}(r) \leq C\, r^{-\beta}$; (B) State-only window: order present, law unresolved.
QIV	no $k^2{=}0$ pole	$\lim_{t\to\infty} \lvert\Delta(t;\mathbf{0})\rvert < \sigma_0$	(A) $\exists\, \delta_\star \in (\varepsilon_{\mathrm{law}}, 1) : d_{\mathrm{law}}(r) \geq \delta_\star$ uniformly, $\lvert\Lambda_{\mathrm{state}}\rvert < 1-\varepsilon_{\mathrm{state}}\rvert$; (B) Disordered: no pole, no long-time order.

Notes and references.

(N1) **Quadrant decision in visibility space.** The decision is made in $(\Lambda_{\mathrm{law}}, \Lambda_{\mathrm{state}})$ with tolerances $(\varepsilon_{\mathrm{law}}, \varepsilon_{\mathrm{state}})$ via the canonical rule; the healing clause $d_{\mathrm{law}}(r) := 1 - |\Lambda_{\mathrm{law}}(r)| \sim r^{-\beta}$ with $\beta > 0$ separates QIII from QIV (cf. Definition 9.297 and Proposition 9.298).

(N2) **Meters (Tantrakyma sector).**

$$v^{(L)} := \left|\mathrm{Res}_{k^2=0}\, \Delta^{[\chi]}_{\Theta\Theta}(k)\right|, \qquad v^{(S)} := \left|\Delta^{[\chi]}_{\Theta\Theta}(t \to \infty; \mathbf{x} = 0)\right|_{H^0}.$$

Diagnostic realized reference configuration thresholds (ρ_0, σ_0) are the instrument floors consistent with the typed witness normalization (cf. Definition 9.296).

(N3) **NG vs. Higgs labeling.** NG channels for Tantrakyma are stamped QI in the typed, parasymmetric readout, while the Higgs (radial) channel is gapped (cf. Definition 9.343 and eq. (9.185)).

(N4) **Synthesis baselines and stability.** SynTh meters elsewhere use a 1/2 baseline; here the quadrant call relies on visibilities $\Lambda_\bullet \in [0, 1]$ and their normalizers/aggregators (e.g. RSS), with monotonicity/continuity and plateau robustness ensuring stable decisions (cf. eqs. (9.126) to (9.128) and Definition 9.297).

Healing indicators $(\mathscr{H}_L, \mathscr{H}_S)$ refine QIII vs. QIV calls (cf. Definition 9.297). We now place the spontaneous symmetry breaking (SSB) analysis in this quadrant framework.

1. Symmetric potential and its invariance group

The Tantrakyma action $S_\Theta[\hat{\mathbf{\Theta}}]$ (Definition 9.310) employs the operator-valued potential $\mathcal{V}_\Theta$ of Definition 9.335 and eq. (9.133):

$$\mathcal{V}_\Theta(\hat{\mathbf{\Theta}}) = \tfrac{1}{2}\, m_\Theta^2\, \mathsf{X} + \tfrac{\lambda_\Theta}{4!}\, \mathsf{X}^2 + \cdots, \qquad \mathsf{X} := g_C(\hat{\mathbf{\Theta}}, \hat{\mathbf{\Theta}}),$$

which is invariant under internal, trace-preserving fiber $*$-automorphisms (cf. Remark 9.313) and, more generally, under a continuous internal group G_{matter} acting as $\hat{\mathbf{\Theta}} \mapsto U\, \hat{\mathbf{\Theta}}\, U^\dagger$, $U \in G_{\mathrm{matter}}$. Hence the *law* is symmetric. In the symmetric (disordered) phase, the typed meters typically place channels in QIV (no $k^2 = 0$ pole, no long-time order).

2. Asymmetric vacuum, vacuum manifold, and SSB

Let $V_\Theta = \langle \mathcal{V}_\Theta \rangle$ be the diagnostic realized reference configuration potential (eq. (9.134)). For $m_\Theta^2 < 0$, $\lambda_\Theta > 0$, the minima satisfy

$$\left\langle \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\, \mathsf{X} \right\rangle \;=\; v_\Theta^2 \;=\; -\,\frac{6\, m_\Theta^2}{\lambda_\Theta} \;>\; 0,$$

defining a continuous vacuum manifold with a realized background $\bar{\Theta}$ that solves the EL system (eq. (9.148)). The unbroken subgroup $\mathcal{H}_{\mathrm{matter}} \subset G_{\mathrm{matter}}$ leaves $\bar{\Theta}$ invariant; broken generators span the tangent (Goldstone) directions. In typed T–L–O, the corresponding channels exhibit QI stamps (law & state present): a $k^2 = 0$ pole and long-time order.

3. Physical spectrum and quadrant-certified identification of the Higgs

Decompose $\hat{\mathbf{\Theta}} = \bar{\Theta}\,\mathbb{I}_x + \delta\hat{\mathbf{\Theta}}$ (Definition 10.204) and restrict the potential Hessian to typed subspaces (eq. (9.175)):

$$\mathcal{K}^{[\chi]}_{\Theta\Theta} := \Pi^{[\chi]}_{\Theta\Theta}\ \frac{\delta^2\mathcal{V}_\Theta}{\delta\hat{\mathbf{\Theta}}^\dagger\,\delta\hat{\mathbf{\Theta}}}\bigg|_{\bar{\Theta}}\ \Pi^{[\chi]}_{\Theta\Theta}.$$

1. **Goldstone modes (NG/pNG).** Tangent-to-vacuum directions yield zero/pseudo-zero eigenvalues of $\mathcal{K}^{[\chi]}_{\Theta\Theta}$ and a $k^2 = 0$ pole in $\Delta^{[\chi]}_{\Theta\Theta}$. Their T–L–O outcome is QI (nonzero $v^{(L)}$ and $v^{(S)}$); in gauged realizations, these may be eaten by gauge bosons as usual.

2. **Higgs (radial) kymon.** The amplitude (radial) direction—projected by Π_{H} built from $\hat{\mathsf{R}} \propto \delta\mathsf{X}/\delta\hat{\mathbf{\Theta}}\big|_{\bar{\Theta}}$—produces a scalar, gapped eigenvalue $M^2_{\mathrm{H}} = -2\,m^2_\Theta > 0$ (cf. eqs. (9.185) and (9.193)). Its typed quadrant is *gapped* (no $k^2 = 0$ pole; nonzero gap location in the meters), and its Location band (IR/mesoscopic) is set by M_{H} (Definition 9.299).

Quadrant map (phase semantics, by type).

- *Symmetric phase (no order):* QIV in all Tantrakyma types; $m^2_\Theta > 0$, $\bar{\Theta} = 0$.
- *Approach to criticality:* QII (law-only) channels appear as the $k^2 = 0$ pole emerges while $v^{(S)}$ remains suppressed.
- *Ordered window without resolved law:* QIII may occur (long-time order present, pole unresolved) depending on tags/tolerances.
- *SSB (broken phase):* QI for NG channels, plus a *gapped* radial (Higgs) typed entry.

All assignments are certified by $\mathbf{W}^{[\Theta\Theta]}_{[\chi]}$ with plateau tolerances (Definition 9.296 and Proposition 9.298).

Summary. The SSB analysis is *typed*, *diagnostic realized reference configuration-limited*, and *quadrant-certified*. It ties the sign structure of the quartic parameters (eqs. (9.133) and (9.134)) to the spectrum via the Hessian restriction (eq. (9.175)) and validates the NG/Higgs identification with Parasymmetric T–L–O meters and stamps (Section 9.17.11 and Definition 9.299), mirroring the Gaiamiti workflow and respecting the co-metric construction $g_C : T^* \times T^* \to \mathrm{End}(\mathcal{H}_{\mathrm{em}})$.

Remark 9.344 (General explanatory note: reading the spectrum from fluctuations)**.** To understand the particle content of the Tantrakyma sector, expand the field about a realized vacuum (cf. Definition 10.204):

$$\hat{\mathbf{\Theta}}(x) \;=\; \bar{\Theta}(x)\,\mathbb{I}_x \;+\; \delta\hat{\mathbf{\Theta}}(x), \qquad \big\langle \delta\hat{\mathbf{\Theta}}(x) \big\rangle_{\mathfrak{P}} = 0,$$

with diagnostic realized reference configuration expectation $\langle\cdot\rangle_{\mathfrak{P}}$ taken on $H^0(\hat{Q}_{\mathrm{BRST}})$. The quanta of $\delta\hat{\mathbf{\Theta}}$—the *kymons*—split kinematically according to their direction in field space relative to the minimum of the potential V_Θ (cf. Definition 9.335 and eq. (9.134)):

1. **Radial (amplitude) fluctuations ⇒ massive scalar (Higgs).** Variations that change the g_C-*amplitude* of the field, i.e. the invariant $\mathsf{X} = g_C(\hat{\mathbf{\Theta}}, \hat{\mathbf{\Theta}})$, probe the curvature of the potential at the minimum and thus acquire a nonzero mass. In the IQR framework this radial, scalar kymon is identified with the **Higgs boson** (cf. Section 9.18.5). Its mass is set by the second variation (Hessian) of the potential along the g_C-radial direction at $\bar{\Theta}$; equivalently, by the restricted operator $\mathcal{K}_{\mathrm{H}}$ of eq. (9.185).
2. **Tangential (phase) fluctuations ⇒ massless Goldstone modes.** Variations *tangent* to the vacuum manifold $G_{\mathrm{matter}}/\mathcal{H}_{\mathrm{matter}}$ cost no potential energy at leading order and therefore remain massless. These are the Goldstone modes implied by Goldstone's theorem [93, 94]. In a gauged realization they are "eaten" by the corresponding gauge bosons, which thereby become massive (the usual Higgs mechanism).

In short, the *kymon* is the general quantum of the Tantrakyma field, while the *Higgs boson* is the particular, radial, scalar kymon that appears when the Tantrakyma potential exhibits spontaneous symmetry breaking. Both identifications are certified on the diagnostic realized reference configuration by the typed Hessians $\mathcal{K}^{[\chi]}_{\Theta\Theta}$ and by the poles/visibilities of the typed propagators $\Delta^{[\chi]}_{\Theta\Theta}$ used in the Parasymmetric T–L–O pipeline (cf. eqs. (9.175) and (9.184) and Section 9.17.11).

9.18.7 Gauge-Coupled Tantrakyma and the Brout–Englert–Higgs Mechanism

Definition 9.345 (Gauge sector and covariant coupling)**.** Work on a certified plateau $\mathfrak{P}$ with band tags $P_x(\lambda_{\mathrm{UV}})$, $K^{(\ell)}_{\mathrm{IR}}$ and BRST projection to $H^0(\hat{Q}_{\mathrm{BRST}})$ (diagnostic realized reference configuration discipline: Propositions 9.191, 9.207 and 9.223). Let a compact Lie group G act on the Tantrakyma fiber $\mathcal{E}_\Theta$ via a representation ρ with generators T_I. Introduce gauge fields A^I_μ with field strength $F^I_{\mu\nu}$ and kinetic metric G_{IJ}. Extend the covariant derivative used in Definition 9.323 (cf. the split connection, Section 6.12.2 and eq. (6.314)) to

$$D_\mu \hat{\mathbf{\Theta}} \;:=\; \nabla^{(\mathrm{rel})}_\mu \hat{\mathbf{\Theta}} \;+\; \partial_\mu \hat{\mathbf{\Theta}} \;+\; i\, g\, A^I_\mu\, \rho(T_I)\, \hat{\mathbf{\Theta}}.$$

On the diagnostic realized reference configuration, define the typed/triadic-inserted action

$$S_{\mathrm{gauge+}\Theta} = \int_{\mathcal{X}_q} d^d x\, \sqrt{-\det \bar{\Sigma}}\, \mathrm{Tr}\Big[-\tfrac{1}{4}\, G_{IJ}\, F^I_{\mu\nu} F^{J\,\mu\nu} + \tfrac{1}{2}\, (D_\mu \hat{\mathbf{\Theta}})^\dagger\, g_C\, (D^\mu \hat{\mathbf{\Theta}}) - \mathcal{V}_\Theta(\hat{\mathbf{\Theta}}) \Big],$$

with $\mathcal{V}_\Theta$ as in Definition 9.310, and typed/triadic insertions as in Definition 9.187.

Definition 9.346 (SSB pattern and kymon split)**.** Let $\bar{\Theta}$ solve the EL system and minimize V_Θ (cf. Definition 9.343). The stabilizer $H \subset G$ satisfies $\rho(T_a)\bar{\Theta} = 0$ for $T_a \in \mathfrak{h}$. Broken generators T_b obey $\rho(T_b)\bar{\Theta} \neq 0$. Define the NG subspace

$$\mathcal{N} \;:=\; \mathrm{span}\{\, \rho(T_b)\bar{\Theta}\,\}_b \;\subset\; \mathcal{E}_\Theta,$$

and the g_C-Riesz projector onto $\mathcal{N}$ (typed version of Definition 9.249):

$$\Pi_{\mathrm{NG}}(V) \;:=\; \sum_{b,c} (B^{-1})^{bc}\, g_C\Big(V,\, \rho(T_b)\bar{\Theta}\Big)\; \rho(T_c)\bar{\Theta}, \qquad B_{bc} := g_C\Big(\rho(T_b)\bar{\Theta},\, \rho(T_c)\bar{\Theta}\Big).$$

The radial (Higgs) line is defined as in Definition 9.343 by $\hat{\mathsf{R}} \propto \bar{\Theta}$ with typed radial projector $\Pi^{[\chi]}_{\mathrm{rad}}$ as in Definition 9.250; the radial mass operator is $\mathcal{K}^{[\chi]}_{\mathrm{rad}}$ (cf. eq. (9.185)). All projectors commute with the triadic channels (Proposition 9.207).

Proposition 9.347 (Vector mass matrix from BEH)**.** *Expanding $\hat{\mathbf{\Theta}} = \bar{\Theta} + \delta\hat{\mathbf{\Theta}}$ in Definition 9.345 and reading the quadratic terms in A_μ gives the internal vector mass term*

$$\mathcal{L}^{(A)}_{\mathrm{mass}} = \tfrac{1}{2}\, A^I_\mu\, (M^2)_{IJ}\, A^{J\,\mu}, \qquad (M^2)_{IJ} := g^2\, g_C\Big(\rho(T_I)\bar{\Theta},\ \rho(T_J)\bar{\Theta}\Big),$$

which is the Θ-sector specialization of the unified diagnostic realized reference configuration formula in Definition 9.252. Thus vectors associated with broken directions acquire masses $\{m_V^2\} \subset \mathrm{spec}(M^2)$, while H-vectors remain massless. The scalar spectrum splits into: (i) the Higgs (radial) kymon *from eq. (9.185), and (ii) the NG/pNG kymons spanning $\mathcal{N}$ (typed NG projector of Definition 9.346 / Definition 9.249). Typed transverse/longitudinal/helicity resolutions follow Definition 9.350.*

Proof of Proposition 9.347. Work on a certified plateau $\mathfrak{P}$ with BRST projection to $H^0(\hat{Q}_{\mathrm{BRST}})$, where typed/triadic meters commute and route–equivalence holds (Propositions 9.191, 9.207 and 9.223). Start from the gauge–coupled Tantrakyma action in Definition 9.345:

$$S_{\mathrm{gauge+}\Theta} = \int_{\mathcal{X}_q} d^d x\, \sqrt{-\det \bar{\Sigma}}\, \mathrm{Tr}\Big[-\tfrac{1}{4}\, G_{IJ} F^I_{\mu\nu} F^{J\,\mu\nu} + \tfrac{1}{2}\, (D_\mu \hat{\mathbf{\Theta}})^\dagger g_C (D^\mu \hat{\mathbf{\Theta}}) - \mathcal{V}_\Theta(\hat{\mathbf{\Theta}})\Big],$$

with covariant derivative $D_\mu \hat{\mathbf{\Theta}} = \nabla^{(\mathrm{rel})}_\mu \hat{\mathbf{\Theta}} + \partial_\mu \hat{\mathbf{\Theta}} + i\, g\, A^I_\mu\, \rho(T_I)\hat{\mathbf{\Theta}}$. Let $\hat{\mathbf{\Theta}} = \bar{\Theta} + \delta\hat{\mathbf{\Theta}}$ where $\bar{\Theta}$ minimizes V_Θ and realizes $G \to H$ (Definitions 9.343 and 9.346). Expanding the kinetic term to quadratic order in fields and using sesquilinearity of g_C,

$$\begin{aligned} \tfrac{1}{2}(D_\mu \hat{\mathbf{\Theta}})^\dagger g_C (D^\mu \hat{\mathbf{\Theta}}) &= \tfrac{1}{2}\Big(\cdots + i g\, A^I_\mu\, \rho(T_I)\bar{\Theta} + \cdots\Big)^\dagger\, g_C\, \Big(\cdots + i g\, A^{J\,\mu}\, \rho(T_J)\bar{\Theta} + \cdots\Big) \\ &\supset \tfrac{1}{2}\, g^2\, A^I_\mu\, A^{J\,\mu}\, g_C\Big(\rho(T_I)\bar{\Theta},\ \rho(T_J)\bar{\Theta}\Big), \end{aligned}$$

which yields the purely quadratic $A\cdot A$ mass term

$$\mathcal{L}^{(A)}_{\mathrm{mass}} = \tfrac{1}{2}\, A^I_\mu\, (M^2)_{IJ}\, A^{J\,\mu}, \qquad (M^2)_{IJ} := g^2\, g_C\Big(\rho(T_I)\bar{\Theta},\ \rho(T_J)\bar{\Theta}\Big).$$

The mixed A–NG terms are cancelled on the diagnostic realized reference configuration by the R_ξ gauge fixing of Definition 9.349; the associated ghost sector closes BRST and removes gauge–artefactual longitudinal contributions from H^0. Gauge–parameter independence of the physical pole positions/residues follows from the Nielsen/BRST identities (Theorem 9.351).

If $T_a \in \mathfrak{h} = \mathrm{Lie}(H)$ is unbroken, then $\rho(T_a)\bar{\Theta} = 0$ (Definition 9.346), so the corresponding rows/columns of M^2 vanish and those vectors remain massless. For broken generators T_b one has $\rho(T_b)\bar{\Theta} \neq 0$, producing strictly positive entries in M^2; transverse simple poles at $k^2 \in \mathrm{spec}(M^2)$ are then the physical vector masses. The scalar sector splits into the radial (Higgs) kymon from the typed restricted Hessian (eq. (9.185) and Definition 9.343) and the NG/pNG kymons spanning the typed NG subspace $\mathcal{N}$ (Riesz/Gram projector of Definitions 9.249 and 9.346). Vector polarizations are resolved with the typed transverse/longitudinal/helicity projectors of Definition 9.350. This proves the proposition. □

Definition 9.348 (Tantron: Tantrakyma spin–1 internal vector)**.** Let G act on the Tantrakyma fiber $\mathcal{E}_\Theta$ via ρ and let $\bar{\Theta}$ realize $G \to H$ (cf. Definitions 9.343 and 9.346). The *tantron* is a one–particle, spin–1 *internal* gauge quantum in the Tantrakyma sector, i.e. a mass–eigenstate of the bundle connection A^I_μ defined as follows.

(i) **Mass matrix and eigenmodes.** Form the BEH mass matrix

$$(M^2_\Theta)_{IJ} := g^2\, g_C\Big(\rho(T_I)\bar{\Theta},\ \rho(T_J)\bar{\Theta}\Big) \quad \text{(cf. } \textit{Definition } 9.345 \textit{and Proposition } 9.347\text{)}.$$

Choose $O \in \mathrm{O}(\dim G)$ with $O^\top M^2_\Theta O = \mathrm{diag}(m^2_{V,a})$. Define tantron fields (mass eigenvectors) by

$$W_{a\,\mu} \ := \ O^{\ I}{}_{a}\, A^I_\mu\,, \qquad m^2_{V,a} \in \mathrm{spec}(M^2_\Theta).$$

(ii) **Physical pole definition (diagnostic realized reference configuration).** On the diagnostic realized reference configuration $\mathfrak{P}$ with BRST projection to $H^0(\hat{Q}_{\mathrm{BRST}})$, the *physical tantron* labelled by a is the simple pole of the two–point function in the *transverse* channel at $k^2 = m^2_{V,a}$, isolated by the typed projectors Π_T (and helicity projectors when applicable) of Definition 9.350. Longitudinal parts are gauge–artefacts removed by BRST; pole positions/residues are ξ–independent (Nielsen/BRST, Theorem 9.351).

(iii) **Unbroken directions.** If $m^2_{V,a} = 0$, the corresponding $W_{a\,\mu}$ is a massless internal vector (unbroken H-direction). Otherwise $m^2_{V,a} > 0$ and the tantron is massive; the would–be NG partner is eaten (typed equivalence theorem Corollary 9.352).

(iv) **Separation from scalar content.** Tantrons are quanta of the bundle connection in the Θ sector; they are distinct from the scalar *Higgs/kymon*, whose mass is read from the typed radial Hessian $\mathcal{K}^{[\chi]}_{\mathrm{rad}}$ (eq. (9.185) and Definition 9.343), and there is no spin–2 excitation in this sector.

Justification (Tantron). The spin–1 bundle–connection quantum in the Tantrakyma sector is called the *Tantron*. The name is an elegant truncation of the sector label, anchored in the Sanskrit root *tantra* (: "loom," "weave," "system"), which captures the particle's function: it furnishes the locally woven *framework* by which the wave–like *Kymon* field is compared covariantly across the internal fiber. Formally, the Tantron denotes the *internal* spin–1 gauge quantum corresponding to the dynamical components A^I_μ of the operator–bundle connection $\mathcal{A}_\mu$ once the Tantrakyma sector is gauged (cf. Definition 9.345). Under spontaneous breaking by $\bar{\Theta}$, the covariant kinetic term produces the Brout–Englert–Higgs mass matrix

$$(M^2_\Theta)_{IJ} = g^2\, g_C\Big(\rho(T_I)\bar{\Theta},\ \rho(T_J)\bar{\Theta}\Big) \quad \text{(see Proposition 9.347)},$$

so that transverse Tantrons appear as diagnostic realized reference configuration–physical poles at $k^2 \in \mathrm{spec}(M^2_\Theta)$ in the vector channel (isolated by the typed projectors of Definition 9.350 and gauge–parameter independent by Nielsen/BRST, Theorem 9.351). Conceptually, the Tantron provides the "threads" that weave Kymon wave–states through the internal bundle, serving as the structural vector complement to the spin–0 radial Kymon (the *Higgs* of Definition 9.343 and eq. (9.185)).

Tantrakyma (Θ) — what becomes massive, what is eaten, and how the vectors behave. Gauging an internal G acting on $\mathcal{E}_\Theta$ and realizing $\bar{\Theta}$ breaks $G \to H$ and produces the Tantrakyma BEH mass matrix

$$(M^2_\Theta)_{IJ} = g^2\, g_C\Big(\rho(T_I)\bar{\Theta},\ \rho(T_J)\bar{\Theta}\Big) \quad (\textit{Definition 9.345and Proposition 9.347}).$$

What becomes massive? The internal vectors along the *broken* generators—i.e. the *tantrons*—acquire masses $\{m_V^2\} = \mathrm{spec}(M^2_\Theta)$; vectors along unbroken H-directions remain *massless*. The would-be NG kymons are *eaten* and reappear as *longitudinal* polarizations. On the diagnostic realized reference configuration, physical one-particle tantron poles lie in the *transverse* channel (typed Π_T, Π_L; Definition 9.350); longitudinal pieces are BRST-removed from H^0. As in the Φ sector, both *masses and residues are ξ-independent* by Nielsen/BRST (Theorem 9.351), and the typed *Equivalence Theorem* matches longitudinal-vector and NG amplitudes at $\sqrt{s} \gg m_V$ up to $\mathcal{O}(m_V/\sqrt{s})$ (Corollary 9.352). The scalar spectrum simultaneously splits into the gapped *Higgs/kymon*—the radial eigenmode of the typed Hessian (Definition 9.343 and eq. (9.185))—and (if present) pNG modes only when explicit breaking is introduced and the corresponding directions are left ungauged (Definitions 9.249 and 9.346).

Physically, tantrons are *short-range carriers* with range $\sim m_V^{-1}$. They couple (i) to the *Higgs/kymon* via the covariant derivative and scalar potential, generating VVh and $VVhh$ vertices; (ii) to *other vectors* through Yang–Mills triple and quartic interactions; and (iii) to any *charged matter* introduced in the model. Non-diagonal M^2_Θ induces *mixing* of gauge directions; degeneracies yield *multiplets* under H. Unbroken-direction vectors remain *massless* and long-range unless further dynamics intervene. All statements are made with typed/triadic readouts on $\mathfrak{P}$ and inherit route-equivalence and BRST hygiene (Propositions 9.191, 9.207 and 9.223 and Definition 9.349).

Definition 9.349 (R_ξ gauge fixing and ghosts on the diagnostic realized reference configuration)**.** Choose the gauge-fixing functional (cf. Gaiamiti analogue Definition 9.261)

$$\mathcal{F}^I \ :=\ \nabla^{(\mathrm{rel})\,\mu} A^I_\mu \ -\ \xi\, g\, g_C\Big(\delta\hat{\mathbf{\Theta}},\ \rho(T^I)\bar{\Theta}\Big).$$

Add

$$\mathcal{L}_{\mathrm{GF}} \ =\ -\tfrac{1}{2\xi}\, G_{IJ}\, \mathcal{F}^I\, \mathcal{F}^J, \qquad \mathcal{L}_{\mathrm{ghost}} \ =\ \bar{c}_I \Big[\, -\nabla^{(\mathrm{rel})\,2}\delta^I{}_J \ -\ \xi\,(M^2)^I{}_J \ +\ \cdots \Big] c^J,$$

with BRST variations closing on $H^0(\hat{Q}_{\mathrm{BRST}})$ (diagnostic realized reference configuration discipline: Propositions 9.191 and 9.207). This cancels the A–NG mixing and yields longitudinal-vector masses $\xi^{1/2} m_V$, while keeping the physical spectra $\mathrm{spec}(M^2)$ and M^2_{H} gauge-parameter independent (Nielsen identities; see Theorem 9.351).

Definition 9.350 (Typed vector projectors and physical polarizations)**.** Let $\mathbb{K}_A$ be the vector kinetic operator (built from $\bar{\Sigma}^{\mu\nu}$ and G_{IJ}). Define the Σ- and g-compatible transverse/longitudinal projectors Π_T, Π_L and the helicity projectors $\Pi^{(\lambda)}_{\mathrm{hel}}$. Physical one-particle vector states on $H^0(\hat{Q}_{\mathrm{BRST}})$ are the simple poles in the Π_T channel; Π_L is gauge-artefactual (removed by BRST). This mirrors the typed scalar linear-response analysis $\Big(\mathbb{K}^{(\Theta)} - \omega^2\Big)\delta\hat{\mathbf{\Theta}} = 0$ in eq. (9.184).

Theorem 9.351 (Nielsen/BRST gauge-parameter independence)**.** *On the certified plateau $\mathfrak{P}$, the pole positions of the Higgs propagator and of the physical vector propagators are independent of the gauge parameter ξ:*

$$\frac{d\,M_{\mathrm{H}}^2}{d\xi} = 0, \qquad \frac{d\,m_V^2}{d\xi} = 0,$$

and likewise for residues in physical channels. Hence BEH mass generation is a diagnostic realized reference configuration-certified observable (route equivalence & BRST safety as in Propositions 9.191 and 9.223).

Proof of Theorem 9.351. **DRCC discipline and BRST/Nielsen setup.** Work on a certified plateau $\mathfrak{P}$ with BRST projection to $H^0(\hat{Q}_{\mathrm{BRST}})$ and commuting typed/triadic meters (Propositions 9.191, 9.207 and 9.223). Use R_ξ gauge fixing with ghosts as in Θ and Φ (Definitions 9.261 and 9.349). On the diagnostic realized reference configuration (DRCC), the (typed) Nielsen identity for the 1PI effective action Γ is written in *BRST-exact* form

$$\partial_\xi \Gamma \;=\; s\,\mathcal{N}_\xi,$$

where s is the BRST differential and $\mathcal{N}_\xi$ is a local functional built from the gauge-fixing/ghost sector and antifields at gauge parameter ξ (notation $\mathcal{N}_\xi$ replaces the inconsistent Ξ_ξ). Insertions of $s(\cdot)$ vanish between physical states (cohomology H^0) up to EOM terms, and this persists after typed/triadic insertions (Proposition 9.207).

Step 1 — Higgs (radial kymon) mass M_{H}^2. By Definitions 9.250 and 9.343 and eq. (9.185),

$$M_{\mathrm{H}}^2 \in \mathrm{spec}\Big(\mathcal{K}_{\mathrm{rad}}^{[\chi]}\Big), \qquad \mathcal{K}_{\mathrm{rad}}^{[\chi]} = \Pi_{\mathrm{rad}}^{[\chi]}\,\frac{\delta^2\Gamma}{\delta\Theta\,\delta\Theta}\Big|_{\bar\Theta}\,\Pi_{\mathrm{rad}}^{[\chi]},$$

with $\bar\Theta$ solving the EL system on $\mathfrak{P}$. Differentiating w.r.t. ξ gives

$$\partial_\xi\left(\frac{\delta^2\Gamma}{\delta\Theta\,\delta\Theta}\right) = \frac{\delta^2}{\delta\Theta\,\delta\Theta}\big(s\,\mathcal{N}_\xi\big).$$

Any explicit ξ-shift of the background drops out by the potential Nielsen identity at the minimum: $\partial_\xi\Gamma|_{\bar\Theta} = -(\partial_i\Gamma)\,\mathcal{C}_\xi^i|_{\bar\Theta} = 0$ since $\partial_i\Gamma(\bar\Theta) = 0$. Projecting onto the typed radial line shows the ξ-variation of the radial block is BRST-exact and cohomologically trivial on H^0. Equivalently, in the correlational route the Higgs pole is the simple zero of $G_H^{-1}(k^2;\xi)$; its typed Nielsen identity has on-shell form

$$\partial_\xi G_H^{-1}(k^2;\xi) = \mathcal{B}_\xi^{\mathrm{H}}(k^2)\,G_H^{-1}(k^2;\xi) + G_H^{-1}(k^2;\xi)\,\mathcal{B}_\xi^{\mathrm{H}\,\dagger}(k^2),$$

so at $k^2 = M_{\mathrm{H}}^2$ (where $G_H^{-1} = 0$) one has $\partial_\xi G_H^{-1}(M_{\mathrm{H}}^2;\xi) = 0$ and thus $\partial_\xi M_{\mathrm{H}}^2 = 0$. Route equivalence on $\mathfrak{P}$ (Proposition 9.223) finishes the step.

Step 2 — Physical vector masses m_V^2. For $U \in \{\Theta, \Phi\}$ use the typed transverse/longitudinal/helicity projectors of Definition 9.350; physical vectors live in the transverse channel on H^0. The typed inverse propagator near the pole reads

$$\big(G_V^{-1}\big)_{\mathrm{T}}(k^2;\xi) = k^2\,\Pi_{\mathrm{T}} - M_U^2\big|_{\mathrm{T}} - \Sigma_{\mathrm{T}}(k^2;\xi),$$

with the tree-level mass matrix M_U^2 given algebraically by the BEH formula (Definition 9.252; concretely Propositions 9.257 and 9.347). The typed Nielsen identity implies

$$\partial_\xi\big(G_V^{-1}\big)_{\mathrm{T}} = \Pi_{\mathrm{T}}\Big[\ \mathcal{B}_\xi^{\mathrm{V}}\, G_V^{-1} + G_V^{-1}\, \mathcal{B}_\xi^{\mathrm{V}\,\dagger}\ \Big]\Pi_{\mathrm{T}},$$

with $\mathcal{B}_\xi^{\mathrm{V}}$ built from BRST-exact, gauge-fixing structures. Evaluating at a transverse physical pole $k^2 = m_V^2$ where $\big(G_V^{-1}\big)_{\mathrm{T}} = 0$ yields $\partial_\xi\big(G_V^{-1}\big)_{\mathrm{T}}(m_V^2;\xi) = 0$ and hence $\partial_\xi m_V^2 = 0$. Background dependences through $\bar{U}$ cancel by the potential Nielsen identity as in Step 1.

Step 3 — Residues in physical channels. Let $Z_{\mathrm{H}}(\xi)$ and $Z_{\mathrm{T}}(\xi)$ be the residues at the Higgs and transverse–vector simple poles. Differentiating the LSZ residues and inserting the on-shell Nielsen forms shows $\partial_\xi Z_{\mathrm{H}} = 0$ and $\partial_\xi Z_{\mathrm{T}} = 0$ on H^0 (any ξ-dependence factors through G^{-1}, which vanishes at the pole). This is consistent with the typed equivalence theorem on $\mathfrak{P}$ (Corollary 9.352).

Conclusion. On the certified plateau, the BRST-exact Nielsen identity $\partial_\xi\Gamma = s\mathcal{N}_\xi$ implies $\frac{dM_{\mathrm{H}}^2}{d\xi} = 0$ and $\frac{dm_V^2}{d\xi} = 0$, and likewise ∂_ξ kills residues in physical channels. Therefore BEH mass generation is diagnostic realized reference configuration-certified (route equivalence & BRST safety), as claimed. □

Corollary 9.352 (Equivalence theorem (typed, on the diagnostic realized reference configuration))**.** *For $\sqrt{s} \gg m_V$, matrix elements with external longitudinal vectors equal those with the corresponding NG kymons in the Π_{NG} channel (typed projector in Definitions 9.249 and 9.346), up to $\mathcal{O}(m_V/\sqrt{s})$.*

Proof. **Diagnostic realized reference configuration discipline and typed setup.** Work on a certified plateau $\mathfrak{P}$ with band tags $P_x(\lambda_{\mathrm{UV}})$, $K_{\mathrm{IR}}^{(\ell)}$ and BRST projection to $H^0(\hat{Q}_{\mathrm{BRST}})$, with route–equivalence and gauge hygiene as in Propositions 9.191, 9.207 and 9.223. Use R_ξ gauge fixing and ghosts as in Definition 9.349 (Tantrakyma) / Definition 9.261 (Gaiamiti). Let the internal vectors be typed by the transverse/helicity projectors of Definition 9.350, and let the NG projector be the typed version of Definitions 9.249 and 9.346. Tree–level masses arise from spec(M^2) with M^2 given by Propositions 9.257 and 9.347 and Definition 9.252.

Step 1 — Longitudinal polarization at high energy. For an external massive vector with momentum k and mass m_V, choose the *helicity–0* (longitudinal) polarization vector $\varepsilon_L^\mu(k)$ from the physical (transverse) typed space ($\varepsilon_L \cdot k = 0$; cf. Definition 9.350). Kinematically,

$$\varepsilon_L^\mu(k) \;=\; \frac{k^\mu}{m_V} \;+\; \mathcal{O}\!\left(\frac{m_V}{\sqrt{s}}\right), \qquad s \equiv k_{\mathrm{cm}}^2 \gg m_V^2,$$

so any amplitude with an external longitudinal vector satisfies $\mathcal{M}[\ldots, A_L^I(k), \ldots] = \varepsilon_L^\mu(k)\,\mathcal{T}_\mu^I(k,\ldots) = \frac{k^\mu}{m_V}\,\mathcal{T}_\mu^I + \mathcal{O}(m_V/\sqrt{s})$, where $\mathcal{T}_\mu^I$ is the amputated Green's function for the vector leg in the typed channel.

Step 2 — Typed Slavnov–Taylor/Nielsen identity with one vector leg. From the R_ξ gauge functionals in Definitions 9.261 and 9.349 and BRST closure on the diagnostic realized

reference configuration, the (typed, triadic–inserted) Slavnov–Taylor/Nielsen identity for the amputated 1PI functional with one vector leg implies

$$k^\mu \mathcal{T}^I_\mu \;=\; m_V \mathcal{T}^I_{\mathrm{NG}} \;+\; \mathcal{R}^I,$$

where $\mathcal{T}^I_{\mathrm{NG}}$ is the amputated functional with the NG insertion projected by Π_{NG} (typed version of Definitions 9.249 and 9.346) and $\mathcal{R}^I$ is a sum of BRST–exact terms and gauge–fixing artefacts proportional to the inverse kernels. On the diagnostic realized reference configuration and on shell, $\mathcal{R}^I$ vanishes between physical states in $H^0(\hat{Q}_{\mathrm{BRST}})$ (by Proposition 9.207 and the EOM at the realized background), leaving

$$k^\mu \mathcal{T}^I_\mu \;=\; m_V \mathcal{T}^I_{\mathrm{NG}} \quad \text{on } H^0.$$

Step 3 — LSZ reduction and comparison of matrix elements. Insert the identity above into the LSZ expression for the matrix element with one external longitudinal vector:

$$\mathcal{M}[\ldots, A^I_L(k), \ldots] = \frac{k^\mu}{m_V}\mathcal{T}^I_\mu \;+\; \mathcal{O}\!\left(\frac{m_V}{\sqrt{s}}\right) = \mathcal{T}^I_{\mathrm{NG}} \;+\; \mathcal{O}\!\left(\frac{m_V}{\sqrt{s}}\right),$$

where the $\mathcal{O}(m_V/\sqrt{s})$ terms originate from the subleading part of ε^μ_L in Step 1 and any suppressed BRST–exact remainders. By route–equivalence on $\mathfrak{P}$ (Proposition 9.223) and gauge–parameter independence (Theorem 9.351), both sides are diagnostic realized reference configuration–certified observables in the typed channels.

Conclusion. For $\sqrt{s} \gg m_V$,

$$\mathcal{M}[\ldots, A^I_L(k), \ldots] = \mathcal{M}[\ldots, \phi^I_{\mathrm{NG}}(k), \ldots]_{\Pi_{\mathrm{NG}}} \;+\; \mathcal{O}\!\left(\frac{m_V}{\sqrt{s}}\right),$$

i.e. the matrix element with an external longitudinal vector equals the one with the corresponding NG kymon projected by Π_{NG} (typed projector in Definitions 9.249 and 9.346), up to $\mathcal{O}(m_V/\sqrt{s})$. □

Relation to existing Chapter 10 structures.

(i) **Higgs = radial kymon (typed, on the diagnostic realized reference configuration).** The Higgs boson is precisely the *radial kymon* defined in Definition 9.343, with mass extracted from the typed radial Hessian $\mathcal{K}^{[\chi]}_{\mathrm{rad}}$ (eq. (9.185) and Definition 9.250). All statements are made on the certified plateau and respect route–equivalence and BRST/Nielsen safety (Propositions 9.191 and 9.223).

(ii) **NG/pNG sector = $\mathrm{Ran}\,\Pi_{\mathrm{NG}}$ (typed).** The Nambu–Goldstone sector is the range of the NG projector defined for Tantrakyma in Definition 9.346 and, in typed generality, in Definition 9.249. With small explicit breaking, the NG cluster becomes pseudo–NG with GMOR scaling on the diagnostic realized reference configuration; quadrant assignment flips to QIV as in Proposition 9.251 (diagnostic realized reference configuration sources/healing and NG diagnostics in Definition 9.188 and Proposition 9.197).

(iii) **Typed & triadic compatibility (T–L–O ready).** The typed projectors $\Pi_{UU}^{[\chi]}$ commute with the triadic channels $\Pi_{\mathrm{corr}}, \Pi_{\mathrm{vac}}, \Pi_{\mathrm{spec}}$ (Definition 9.186 and Proposition 9.207), and, by construction, with the NG and radial projectors (Π_{NG}, Π_{H} for Tantrakyma in Definition 9.346, and $\Pi_{\mathrm{A}}^{(\Phi)}$ for Gaiamiti in Definition 9.259). Consequently, all SynTh readouts—typed witnesses and potential–based operators $\mathcal{O}^{(\bullet)}$—remain compatible with the *Type–Location–Onset* pipeline (Definitions 9.187, 9.190 and 9.221). Typed decisions and their recording use the ledger interfaces (Definition 9.198 and Corollaries 9.194 and 9.218).

(iv) **Consistency with gauging and mass generation.** Under internal gauging, vector masses are generated by the diagnostic realized reference configuration–algebraic BEH matrices (Θ: Proposition 9.347 and Definition 9.252; Φ: Proposition 9.257 and Definition 9.252), while the radial scalars are governed by $\mathcal{K}_{\mathrm{rad}}^{[\chi]}$ (eq. (9.185) and Definition 9.259). Gauge–parameter independence of Higgs and transverse–vector poles and residues follows from the diagnostic realized reference configuration Nielsen theorem (Theorem 9.351); at high energy, longitudinal vectors match NG kymons in the typed NG channel (Corollary 9.352).

(v) **Quadrant semantics preserved.** Typed quadrant decisions (QI–QIV) follow the canonical rule on the diagnostic realized reference configuration (Definition 7.685 and Propositions 9.251 and 9.254), with TT graviton masslessness protected (for Φ) by the relational Ward structure (Section 9.17.2 and Remark 9.260).

9.18.8 A Minimal $U(1)$ Worked Example of the BEH Mechanism

All constructions below are evaluated on a certified plateau $\mathfrak{P}$ with band tags $P_x(\lambda_{\mathrm{UV}})$, $K_{\mathrm{IR}}^{(\ell)}$ and BRST projection to $H^0(\hat{Q}_{\mathrm{BRST}})$; typed projectors commute with the triadic channels $\Pi_{\mathrm{corr}}, \Pi_{\mathrm{vac}}, \Pi_{\mathrm{spec}}$ (Propositions 9.191, 9.207 and 9.223 and Definition 9.186).

Setup and action. Let $G = U(1)$ act on the Tantrakyma fiber $\mathcal{E}_\Theta$ by a representation ρ with charge q, normalized anti–self–adjointly with respect to g_C,

$$\rho(T)^\dagger\, g_C + g_C\, \rho(T) = 0,$$

so that the NG (phase) direction is g_C-orthogonal to the radial one (diagnostic realized reference configuration hygiene per Proposition 9.207). Introduce a gauge field A_μ with field strength $F_{\mu\nu}$, coupling g, and extend the covariant derivative of Definition 9.323 (cf. Section 6.12.2 and eq. (6.314)) to

$$D_\mu \hat{\boldsymbol{\Theta}} := \nabla_\mu^{(\mathrm{rel})} \hat{\boldsymbol{\Theta}} \;+\; \partial_\mu \hat{\boldsymbol{\Theta}} \;+\; i\, g\, A_\mu\, \rho(T)\, \hat{\boldsymbol{\Theta}}.$$

On the certified plateau $\mathfrak{P}$, the (operator-valued) action with typed/triadic insertion is

$$S_{U(1)+\Theta} = \int_{\mathcal{X}_q} d^d x\, \sqrt{-\det \bar{\Sigma}}\, \mathrm{Tr}\Big[-\tfrac{1}{4}\, F_{\mu\nu} F^{\mu\nu} + \tfrac{1}{2}\, (D_\mu \hat{\boldsymbol{\Theta}})^\dagger\, g_C\, (D^\mu \hat{\boldsymbol{\Theta}}) - \mathcal{V}_\Theta(\hat{\boldsymbol{\Theta}}) \Big],$$

with $\mathcal{V}_\Theta$ as in Definition 9.343.

SSB, radial/NG directions, and projectors. Let $\hat{\mathbf{\Theta}} = \bar{\Theta} + \delta\hat{\mathbf{\Theta}}$ with $\bar{\Theta}$ minimizing V_Θ (cf. Definition 9.343). Define the invariant $\mathsf{X} := g_C(\hat{\mathbf{\Theta}}, \hat{\mathbf{\Theta}})$ and its background derivative

$$\delta\mathsf{X}\big|_{\bar{\Theta}}(\delta\hat{\mathbf{\Theta}}) = 2\, g_C(\delta\hat{\mathbf{\Theta}}, \bar{\Theta}) \quad \text{(cf. Remark 9.303)}.$$

The radial (Higgs) direction is $\hat{\mathsf{R}} \propto \bar{\Theta}$ with one-dimensional projector Π_{H} as in Definitions 9.250 and 9.343 and eq. (9.185). The Nambu–Goldstone direction is the g_C-orthogonal phase vector $v := \rho(T)\bar{\Theta}$ with $g_C(\bar{\Theta}, v) = 0$ by the normalization above. The g_C–Riesz NG projector (the $U(1)$ specialization of Definitions 9.249 and 9.346) is

$$\Pi_{\mathrm{NG}}(V) := \frac{g_C\Big(V, \rho(T)\bar{\Theta}\Big)}{g_C\Big(\rho(T)\bar{\Theta}, \rho(T)\bar{\Theta}\Big)}\, \rho(T)\bar{\Theta}, \qquad \operatorname{Ran}\Pi_{\mathrm{NG}} = \operatorname{span}\{\rho(T)\bar{\Theta}\}.$$

All projectors commute with triadic channels (Proposition 9.207).

Vector mass from BEH. Expanding the kinetic term $\frac{1}{2}(D_\mu\hat{\mathbf{\Theta}})^\dagger g_C(D^\mu\hat{\mathbf{\Theta}})$ about $\bar{\Theta}$ yields

$$\mathcal{L}^{(A)}_{\mathrm{mass}} = \tfrac{1}{2}\, m_V^2\, A_\mu A^\mu, \qquad m_V^2 \;:=\; g^2\, g_C\Big(\rho(T)\bar{\Theta},\ \rho(T)\bar{\Theta}\Big) \;=\; g^2\, q^2\, g_C(\bar{\Theta}, \bar{\Theta}),$$

which is the $U(1)$ case of the unified diagnostic realized reference configuration formula in Definition 9.252 and the sector proposition Proposition 9.347. The scalar sector splits into: (i) the *Higgs (radial) kymon*, defined by

$$\mathcal{K}_{\mathrm{H}} := \Pi_{\mathrm{H}}\, \frac{\delta^2\mathcal{V}_\Theta}{\delta\hat{\mathbf{\Theta}}\,\delta\hat{\mathbf{\Theta}}}\bigg|_{\bar{\Theta}}\, \Pi_{\mathrm{H}}, \qquad M_{\mathrm{H}}^2 \in \operatorname{spec}(\mathcal{K}_{\mathrm{H}}) \quad \text{(cf. eq. (9.185) and Definition 9.250)},$$

and (ii) the NG direction $\operatorname{Ran}\Pi_{\mathrm{NG}}$.

R_ξ gauge fixing and BRST cancellation of A–NG mixing. Define

$$\mathcal{F} := \nabla^{(\mathrm{rel})\,\mu} A_\mu \;-\; \xi\, g\, g_C\Big(\delta\hat{\mathbf{\Theta}}, \rho(T)\bar{\Theta}\Big), \qquad \mathcal{L}_{\mathrm{GF}} = -\tfrac{1}{2\xi}\,\mathcal{F}^2,$$

together with the standard ghost term so BRST closes on $H^0(\hat{Q}_{\mathrm{BRST}})$ (cf. Definitions 9.261 and 9.349 and Proposition 9.207). This cancels the A–NG mixing, gives the would–be NG a gauge-dependent mass $\sqrt{\xi}\, m_V$, and leaves the physical transverse-vector pole at $k^2 = m_V^2$. Typed readouts use Π_T, Π_L for vectors and $\Pi_{\mathrm{H}}, \Pi_{\mathrm{NG}}$ for scalars (cf. the linear response eq. (9.184) and Definition 9.350).

Propagators and gauge-parameter independence (diagnostic realized reference configuration certification). In the typed channels Π_T, Π_L and $\Pi_{\mathrm{H}}, \Pi_{\mathrm{NG}}$ (cf. Definitions 9.249, 9.343 and 9.350),

$$\frac{d\, m_V^2}{d\xi} = 0, \qquad \frac{d\, M_{\mathrm{H}}^2}{d\xi} = 0,$$

and residues are ξ–independent by the diagnostic realized reference configuration Nielsen theorem Theorem 9.351. Hence both the vector mass m_V and the Higgs mass M_{H} are diagnostic realized reference configuration-certified observables (route equivalence and BRST safety: Propositions 9.191 and 9.223).

Quadrant viewpoint. With $\mathcal{V}_\Theta$ exhibiting SSB and no explicit $U(1)$ breaking, the typed predicate gives QII (NG present, eaten $\Rightarrow$ massive vector m_V, gapped radial kymon M_{H}) as in Proposition 9.251. With a small explicit breaking spurion, the system moves to QIV with a pseudo–NG of mass $\mathcal{O}(\epsilon)$ (GMOR scaling; Definition 9.188 and Proposition 9.197), while the vector/Higgs poles remain diagnostic realized reference configuration–certified (Theorem 9.351).

Typed Vector Projectors and Quadratic Lagrangian (U(1) BEH)

Typed vector projectors. Work locally in momentum space on $\mathfrak{P}$, with relational metric $\bar{\Sigma}_{\mu\nu}$. Define

$$\Pi_T^{\mu\nu}(k) := \bar{\Sigma}^{\mu\nu} - \frac{k^\mu k^\nu}{k^2}, \qquad \Pi_L^{\mu\nu}(k) := \frac{k^\mu k^\nu}{k^2},$$

so that $\Pi_T^2 = \Pi_T,\ \Pi_L^2 = \Pi_L,\ \Pi_T\Pi_L = 0,\ \Pi_T + \Pi_L = \mathbb{1}$. We decompose $A_\mu = A_T^\mu + A_L^\mu$ with $A_T^\mu := \Pi_{T\nu}^\mu A^\nu$, $A_L^\mu := \Pi_{L\nu}^\mu A^\nu$ (Definition 9.350).

SSB pattern and masses. Let $G = U(1)$ with generator T in representation ρ, charge q, and coupling g. For the realized background $\bar{\Theta}$ (cf. Definition 9.343),

$$m_V^2 := g^2\, g_C\Big(\rho(T)\bar{\Theta}, \rho(T)\bar{\Theta}\Big) = g^2 q^2\, g_C(\bar{\Theta}, \bar{\Theta}),$$

and the Higgs (radial) mass operator is

$$\mathcal{K}_{\mathrm{H}} \;:=\; \Pi_{\mathrm{H}}\, \frac{\delta^2 \mathcal{V}_\Theta}{\delta\hat{\boldsymbol{\Theta}}\,\delta\hat{\boldsymbol{\Theta}}}\Big|_{\bar{\Theta}}\, \Pi_{\mathrm{H}}, \quad M_{\mathrm{H}}^2 \in \mathrm{spec}(\mathcal{K}_{\mathrm{H}}) \quad \text{(cf. eq. (9.185) and Definition 9.250).}$$

The NG direction is $\mathrm{span}\{\rho(T)\bar{\Theta}\}$ with projector Π_{NG} as in Definitions 9.249 and 9.346.

R_ξ gauge fixing and ghosts. Choose the momentum–space gauge-fixing functional

$$\mathcal{F} \;:=\; k_\mu A^\mu \;-\; \xi\, g\, g_C\Big(\delta\hat{\boldsymbol{\Theta}},\, \rho(T)\bar{\Theta}\Big), \quad \mathcal{L}_{\mathrm{GF}} = -\frac{1}{2\xi}\,\mathcal{F}^2,$$

and the Abelian ghost Lagrangian

$$\mathcal{L}_{\mathrm{ghost}} = \bar{c}\,(k^2 + \xi\, m_V^2)\, c,$$

with diagnostic realized reference configuration BRST closing in $H^0(\hat{Q}_{\mathrm{BRST}})$ (cf. Definition 9.349 and Proposition 9.207).

Riesz construction of the scalar projectors (one line each). Using the typed resolvent definition (cf. Definition 9.249),

$$\boxed{\Pi_{\mathrm{H}} = \frac{1}{2\pi i}\oint_{\Gamma_{\mathrm{H}}} \Big(z - \mathcal{K}_{\Theta\Theta}^{[\chi]}\Big)^{-1} dz, \qquad \Pi_{\mathrm{NG}} = \frac{1}{2\pi i}\oint_{\Gamma_0} \Big(z - \mathcal{K}_{\Theta\Theta}^{[\chi]}\Big)^{-1} dz\ .}$$

Equivalently, in rank-1 g_C–Riesz form (cf. Definitions 9.343 and 9.346),

$$\Pi_{\mathrm{H}}(V) = \frac{g_C\big(V, \widehat{\mathsf{R}}\big)}{g_C\big(\widehat{\mathsf{R}}, \widehat{\mathsf{R}}\big)}\, \widehat{\mathsf{R}}, \qquad \Pi_{\mathrm{NG}}(V) = \frac{g_C\big(V, \rho(T)\bar{\Theta}\big)}{g_C\big(\rho(T)\bar{\Theta}, \rho(T)\bar{\Theta}\big)}\, \rho(T)\bar{\Theta},$$

where $\widehat{\mathsf{R}} \propto \bar{\Theta}$.

Cluster (multiplet) Riesz projector for NG directions (non-Abelian, quasi-degenerate). Let the broken generators be $\{T_b\}_{b\in\mathcal{B}}$ and set $v_b := \rho(T_b)\bar{\Theta}$. When the NG sector forms a quasi-degenerate cluster, isolate the cluster of the typed scalar Hessian $\mathcal{K}^{[\chi]}_{\Theta\Theta}$ near $z = 0$ by a contour Γ_{NG}. The *cluster Riesz projector* is

$$\boxed{\Pi^{\mathrm{cl}}_{\mathrm{NG}} \;=\; \frac{1}{2\pi i}\oint_{\Gamma_{\mathrm{NG}}} \Big(z - \mathcal{K}^{[\chi]}_{\Theta\Theta}\Big)^{-1}\, dz}$$

and admits the finite-rank g_C–Riesz form

$$\boxed{\Pi^{\mathrm{cl}}_{\mathrm{NG}}(V) \;=\; \sum_{b,c\in\mathcal{B}} (B^{-1})^{bc}\, g_C\Big(V,\, v_b\Big)\, v_c, \qquad B_{bc} := g_C(v_b, v_c)}$$

(which coincides with the contour form on the cluster subspace), consistent with the typed NG construction in Definition 9.249.

pNG case (small explicit breaking). With a controlled explicit breaking ($\epsilon \ll 1$), replace Γ_{NG} by a contour Γ_{pNG} that encloses the small eigenvalues $z = \mathcal{O}(\epsilon)$; the same formulas define $\Pi^{\mathrm{cl}}_{\mathrm{pNG}}$. On the diagnostic realized reference configuration, $\Pi^{\mathrm{cl}}_{\mathrm{NG}}$ and $\Pi^{\mathrm{cl}}_{\mathrm{pNG}}$ commute with the triadic projectors and with the scalar readouts used in Definition 9.343 and eq. (9.185).

Quadratic Lagrangian matrix (field order A_T, A_L, NG, H). Expand to quadratic order around $\bar{\Theta}$ and move to momentum space. After adding $\mathcal{L}_{\mathrm{GF}}$, the A_L–NG mixing cancels. In the basis $\Psi := (A_T,\ A_L,\ \pi_{\mathrm{NG}},\ h_{\mathrm{H}})$ (transverse vector, longitudinal vector, NG scalar, Higgs scalar), the quadratic form is diagonal:

$$\mathcal{L}^{(2)} = \tfrac{1}{2}\, \Psi^\dagger\, \mathbb{Q}(k)\, \Psi,$$

$$\mathbb{Q}(k) = \begin{pmatrix} k^2 - m_V^2 & 0 & 0 & 0 \\ 0 & \dfrac{k^2 - m_V^2}{\xi} & 0 & 0 \\ 0 & 0 & k^2 - \xi\, m_V^2 & 0 \\ 0 & 0 & 0 & k^2 - M_{\mathrm{H}}^2 \end{pmatrix},$$

where $k^2 := \bar{\Sigma}^{\mu\nu} k_\mu k_\nu$.

Poles: A_T : $k^2 = m_V^2$ (physical), A_L : $k^2 = m_V^2$ (gauge-artefactual numerator), π_{NG} : $k^2 = \xi m_V^2$, h_{H} : $k^2 = M_{\mathrm{H}}^2$. On $H^0(\hat{Q}_{\mathrm{BRST}})$ the physical poles are $k^2 = m_V^2$ and $k^2 = M_{\mathrm{H}}^2$; ξ-dependence cancels in S–matrix elements and matches NG amplitudes at high energy by Corollary 9.352.

Propagators and physical content. The vector propagator is

$$D_{\mu\nu}(k) = \frac{-i}{k^2 - m_V^2}\,\Pi_{T\,\mu\nu} \;+\; \frac{-i\,\xi}{k^2 - \xi\,m_V^2}\,\Pi_{L\,\mu\nu},$$

the NG propagator is $D_{\mathrm{NG}}(k) = \frac{i}{k^2 - \xi\,m_V^2}$, and the Higgs propagator is $D_{\mathrm{H}}(k) = \frac{i}{k^2 - M_{\mathrm{H}}^2}$. Gauge–parameter independence of the physical residues follows from Theorem 9.351.

Typed/triadic compatibility. Typed scalar readouts use $\Pi_{\mathrm{H}}, \Pi_{\mathrm{NG}}$ (cf. Definitions 9.249 and 9.343), and vector readouts use Π_T, Π_L (Definition 9.350); all commute with the triadic projectors $\Pi_{\mathrm{corr}}, \Pi_{\mathrm{vac}}, \Pi_{\mathrm{spec}}$ (Proposition 9.207). This keeps the BEH completion fully consistent with the T–L–O program and the linear response equation *eq.* (9.184).

The worked $U(1)$ example demonstrates that *diagnostic realized reference configuration discipline*—evaluation on a certified plateau $\mathfrak{P}$ with typed/triadic insertions and reading in the physical BRST cohomology $H^0(\hat{Q}_{\mathrm{BRST}})$—renders the potential and correlational routes equivalent and the results insensitive to gauge–fixing choices. In this setting, pole locations and residues extracted from typed channels are bona fide observables rather than gauge artefacts (route equivalence and Nielsen/BRST safety as in Propositions 9.191 and 9.223).

The *scalar sector* separates cleanly into two directions. The *radial (Higgs) kymon* is defined by the typed radial Hessian $\mathcal{K}_{\mathrm{H}} \equiv \mathcal{K}_{\mathrm{rad}}^{[\chi]}$ (Definitions 9.250 and 9.343 and eq. (9.185)), which determines the Higgs mass M_{H}^2. The *NG sector* is captured by a typed Riesz/Gram projector Π_{NG}, rank–1 in the $U(1)$ case and realized as a cluster projector for non–Abelian or pseudo–NG multiplets (Definitions 9.249 and 9.346).

An *anti–self–adjoint normalization* of the representation with respect to g_C ensures $g_C\big(\bar{\Theta}, \rho(T)\bar{\Theta}\big) = 0$. This fixes the orthogonality of the NG (phase) direction to the radial one at the level of definitions and prevents hidden radial–phase mixing in subsequent calculations (Proposition 9.207).

Vector masses arise algebraically from the BEH mechanism and depend only on the vacuum orientation in the internal space:

$$m_V^2 \;=\; g^2\, g_C\Big(\rho(T)\bar{\Theta}, \rho(T)\bar{\Theta}\Big) \quad (\text{for } U(1)\text{: } g^2 q^2\, g_C(\bar{\Theta}, \bar{\Theta})),$$

the $U(1)$ specialization of the unified typed mass matrix (Definition 9.252; see also Proposition 9.347). No explicit mass term is needed.

Typed *projectors* make the physical content manifest. Vectors are decomposed with Π_T, Π_L; scalars with $\Pi_{\mathrm{H}}, \Pi_{\mathrm{NG}}$ (Definitions 9.249, 9.343 and 9.350). On $H^0(\hat{Q}_{\mathrm{BRST}})$ only the *transverse vector pole* at $k^2 = m_V^2$ and the *Higgs pole* at $k^2 = M_{\mathrm{H}}^2$ remain; longitudinal and would–be NG components are gauge artefacts removed by BRST.

R_ξ *gauge fixing* is used purely as a computational device. The functional chosen cancels A–NG mixing and gives the would–be NG a gauge–dependent mass $\sqrt{\xi}\,m_V$, while leaving transverse and Higgs poles untouched (Definition 9.349). The *Nielsen/BRST theorem* then guarantees

$$\frac{\mathrm{d}m_V^2}{\mathrm{d}\xi} = 0 \qquad \text{and} \qquad \frac{\mathrm{d}M_{\mathrm{H}}^2}{\mathrm{d}\xi} = 0,$$

and likewise for physical residues (Theorem 9.351).

At high energy, the *typed Equivalence Theorem* applies: for $\sqrt{s} \gg m_V$, amplitudes with an external longitudinal vector coincide with those with the corresponding NG mode in the Π_{NG} channel up to $\mathcal{O}(m_V/\sqrt{s})$ (Corollary 9.352). This provides a practical replacement rule for computations while preserving diagnostic realized reference configuration certification.

The *quadrant semantics* remain intact. With SSB and no explicit breaking, the typed predicate yields *QII*: NG present and eaten ($\Rightarrow$ massive internal vector and gapped radial scalar). Introducing a small explicit spurion yields *QIV* with a pNG cluster obeying GMOR scaling, while the transverse vector and Higgs poles remain diagnostic realized reference configuration–certified (Propositions 9.197 and 9.251 and Definition 9.188).

The worked example is explicitly carried out in the Tantrakyma sector. For non–Abelian internal symmetry, the replacement of the rank–1 NG projector by a *cluster* Riesz projector is immediate and preserves all typed/diagnostic realized reference configuration properties (Definitions 9.249 and 9.346). For the Gaiamiti sector, the corresponding BEH completion is *not* rederived here; instead, the consequences are established elsewhere in this chapter: the gauge–coupled action and SSB pattern (Definitions 9.255 and 9.256), the internal vector mass matrix $(M^2)_{IJ} = g^2\, g_T\Big(\rho_\Phi(T_I)\bar{\Phi}, \rho_\Phi(T_J)\bar{\Phi}\Big)$ (Proposition 9.257), the Apeiron (trace–typed) radial operator $\mathcal{K}_{\mathrm{A}}^{(\Phi)}$ and its mass M_{A}^2 (Definition 9.259), and the masslessness of the TT graviton protected by relational/diffeomorphism Ward identities (Remark 9.260 and Section 9.17.2). In summary, internal gauging in the Gaiamiti sector gives masses only to *internal* vectors (Section 9.17.3), leaves the TT graviton massless, and realizes a radial "stiffness" mode governed by $\mathcal{K}_{\mathrm{A}}^{(\Phi)}$, consistent with the typed quadrant semantics and diagnostic realized reference configuration certification.

Finally, a *pragmatic workflow* emerges for any model: fix typed projectors; determine the realized background $\bar{U}$ and construct Π_{H} and Π_{NG}; read M_{H}^2 from the typed radial Hessian and m_V^2 from the BEH matrix; use R_ξ for convenience; report gauge–independent poles/residues under Nielsen/BRST control; and record Type/Location/Onset and the quadrant decision in the ledger (Definitions 9.190 and 9.198). The result is a coherent, end–to–end pipeline with clean mode separation, algebraic mass generation, and invariant physical content.

Derivation of the Higgs Boson Mass from the Tantrakyma Potential

We extract the *radial (Higgs) mass* as the curvature of the Tantrakyma potential along the vacuum (amplitude) direction. Equivalently, in operator form this is the radial restriction of the potential Hessian (cf. eqs. (9.175) and (9.185)).

1. Potential and its minimum. Start from the operator-valued potential (cf. Definition 9.335 and eq. (9.133))

$$\mathcal{V}_\Theta(\hat{\mathbf{\Theta}}) = \tfrac{1}{2}\, m_\Theta^2\, \mathsf{X} \;+\; \tfrac{\lambda_\Theta}{4!}\, \mathsf{X}^2 \;+\; \cdots, \qquad \mathsf{X} := g_C(\hat{\mathbf{\Theta}}, \hat{\mathbf{\Theta}}),$$

with $m_\Theta^2 < 0$, $\lambda_\Theta > 0$ for SSB. To obtain a scalar amplitude on the diagnostic realized reference configuration, define

$$x(x') \;:=\; \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\Big\{\mathsf{X}(x')\Big\}, \qquad \text{and set} \;\; v_\Theta^2 \;:=\; \Big\langle x \Big\rangle_{\mathfrak{P}}.$$

Then $V_\Theta(v_\Theta) = \frac{1}{2}m_\Theta^2 v_\Theta^2 + \frac{\lambda_\Theta}{4!}v_\Theta^4 + \cdots$ (cf. eq. (9.134)). Minimizing in v_Θ gives

$$\frac{\partial V_\Theta}{\partial v_\Theta} = m_\Theta^2\, v_\Theta + \frac{\lambda_\Theta}{6}\, v_\Theta^3 = 0 \quad \Longrightarrow \quad v_\Theta^2 = -\,\frac{6\, m_\Theta^2}{\lambda_\Theta} \;>\; 0.$$

2. Curvature at the minimum (radial mass). The radial (Higgs) mass is the curvature along v_Θ:

$$\frac{\partial^2 V_\Theta}{\partial v_\Theta^2} = m_\Theta^2 + \frac{\lambda_\Theta}{2}\, v_\Theta^2, \qquad M_{\mathrm{H}}^2 := \left.\frac{\partial^2 V_\Theta}{\partial v_\Theta^2}\right|_{v_\Theta^2 = -6m_\Theta^2/\lambda_\Theta}.$$

Substituting the vacuum value yields

$$M_{\mathrm{H}}^2 = m_\Theta^2 + \frac{\lambda_\Theta}{2}\left(-\frac{6m_\Theta^2}{\lambda_\Theta}\right) = -\,2\, m_\Theta^2.$$

3. Final expression and equivalence with the Hessian projector.

$$\boxed{M_{\mathrm{H}}^2 \;=\; -\,2\, m_\Theta^2 \;>\; 0 \qquad (\text{SSB phase: } m_\Theta^2 < 0).} \tag{9.194}$$

This matches the operator–projector construction: if Π_{H} projects onto the g_C-radial direction $\hat{\mathsf{R}} \propto \delta\mathsf{X}/\delta\hat{\boldsymbol{\Theta}}\big|_{\bar{\Theta}}$ (cf. eq. (9.185)), then the radial eigenvalue of the Hessian $\Pi_{\mathrm{H}}\left(\delta^2\mathcal{V}_\Theta/\delta\hat{\boldsymbol{\Theta}}^\dagger\delta\hat{\boldsymbol{\Theta}}\right)\big|_{\bar{\Theta}}\Pi_{\mathrm{H}}$ equals M_{H}^2 in (9.194), while tangent directions yield NG/pNG zeros certified by the typed meters (eqs. (9.126) to (9.128)) and the T–L–O quadrant call (Definition 9.297).

Interpretation

- **Positivity.** In the SSB phase $m_\Theta^2 < 0$, (9.194) gives $M_{\mathrm{H}}^2 > 0$, i.e. a physical, gapped scalar kymon (cf. Section 9.18.5).
- **Emergent scale.** M_{H} is fixed by $(m_\Theta^2, \lambda_\Theta)$ via $v_\Theta^2 = -6m_\Theta^2/\lambda_\Theta$; it is not an independent parameter (cf. Definition 9.335 and eq. (9.133)).
- **Typed/TLO certification.** On the diagnostic realized reference configuration $\mathfrak{P}$, NG channels show a $k^2 = 0$ pole (QI), while the Higgs (radial) channel has a single gapped pole at $k^2 = M_{\mathrm{H}}^2$ and no massless pole, with a consistent Type–Location–Onset stamp (cf. Definition 9.299).

In sum, the Higgs boson emerges as the *radial, scalar kymon* mandated by spontaneous symmetry breaking (cf. Section 9.18.6), with its mass fixed by the Tantrakyma potential and its physical status certified by the typed, parasymmetric T–L–O workflow—paralleling the Standard Model mechanism [82, 91] while remaining faithful to the IQR formalism.

The Composite Nature of the Higgs Boson

Just as the graviton in the geometry sector resolves into triadic species under the Gaiamiti instrumentation, the Higgs boson—as the radial, scalar kymon of the Tantrakyma sector—inherits the *triadic* structure of its parent field and must be identified *typed* and *channel-wise* on a certified plateau $\mathfrak{P}$ with tags $(\lambda_{\mathrm{UV}}, \ell)$ (cf. Definitions 5.54 and 9.301 and Section 9.17.11).

1. Triadic matter channels and typed slicing. The Tantrakyma fluctuation field $\delta\hat{\mathbf{\Theta}}$ carries both a *type* index $[\chi] \in \{\mathrm{CK}, \mathrm{T}_\pm, \mathrm{L}\}$ and a *channel* index $\alpha \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$. With channel projectors Π_α commuting with the typed projectors $\Pi^{[\chi]}_{\Theta\Theta}$ (triadic compatibility; cf. Definition 9.295), we write

$$\delta\hat{\mathbf{\Theta}} = \sum_\alpha \Pi_\alpha\, \delta\hat{\mathbf{\Theta}} = \sum_{\alpha,[\chi]} \Pi^{[\chi]}_{\Theta\Theta}\, \Pi_\alpha\, \delta\hat{\mathbf{\Theta}} \;=:\; \sum_{\alpha,[\chi]} \delta\hat{\mathbf{\Theta}}^{(\alpha,[\chi])}, \qquad [\Pi_\alpha, \Pi^{[\chi]}_{\Theta\Theta}] = 0. \tag{9.195}$$

All kernels and meters below are evaluated on $H^0(\hat{Q}_{\mathrm{BRST}})$ and respect band limitations $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ (cf. Definition 9.306).

2. The Higgs as a composite kymon (typed, triadic). Let Π_{H} denote the projector onto the radial (amplitude) direction at the realized vacuum, $\hat{\mathsf{R}} \propto \delta\mathsf{X}/\delta\hat{\mathbf{\Theta}}\big|_{\bar{\Theta}}$ with $\mathsf{X} := g_C(\hat{\mathbf{\Theta}}, \hat{\mathbf{\Theta}})$ (cf. eq. (9.185)). The Higgs operator field $\hat{\phi}_{\mathrm{H}}$ is the *scalar, radial* component of $\delta\hat{\mathbf{\Theta}}$:

$$\hat{\phi}_{\mathrm{H}}(x) := \Pi_{\mathrm{H}}\, \delta\hat{\mathbf{\Theta}}(x) = \sum_\alpha \Pi_\alpha\, \Pi_{\mathrm{H}}\, \delta\hat{\mathbf{\Theta}}(x) \;=:\; \sum_\alpha \hat{\phi}^{(\alpha)}_{\mathrm{H}}(x), \qquad \alpha \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}. \tag{9.196}$$

Thus the *single observed* Higgs boson corresponds to a *coherent superposition* across triadic channels, with relative weights fixed operationally by the typed, retarded propagators and the parasymmetric meters (cf. Section 9.17.11 and Definition 9.296).[8]

3. Parasymmetric T–L–O diagnostics of the composite Higgs. For each type $[\chi]$ and channel α, form the band-limited retarded kernel $\Delta^{(\alpha,[\chi])}_{\Theta\Theta}$ and meters $(v^{(L)}, v^{(S)}, v^{(LS)})_{(\alpha,[\chi])}$ (cf. eqs. (9.126) to (9.128)). The *Higgs-certified* outcome requires:

1. **Gapped radial pole.** In the Π_{H}-projected sector, a simple pole at $k^2 = M_{\mathrm{H}}^2 > 0$ with positive residue and no $k^2 = 0$ pole (contrast NG/pNG channels), agreeing with $\mathrm{spec}(\mathcal{K}_{\mathrm{H}})$ (cf. eqs. (9.185) and (9.194)).

2. **Typed quadrant call.** The Higgs channel stamps *gapped* (no massless pole) with a Location band (IR/meso) set by M_{H} and Onset given by the first certified threshold (cf. Definitions 9.297 and 9.299).

3. **Cross-channel coherence.** The helicity-resolved gain $v^{(LS)}_{(\alpha,[\chi])}$ quantifies mixing among channels, enabling a *parasymmetric* reconstruction of the coherent superposition $\hat{\phi}_{\mathrm{H}} = \sum_\alpha \hat{\phi}^{(\alpha)}_{\mathrm{H}}$.

4. Channel semantics for the Higgs species. With $\hat{\phi}^{(\alpha)}_{\mathrm{H}} = \Pi_\alpha \hat{\phi}_{\mathrm{H}}$, the triadic "species" acquire precise meanings:

- **Correlational Higgs** $\hat{\phi}^{(\mathrm{corr})}_{\mathrm{H}}$: Dominant response to relational (entanglement) structure; enhanced long-time meter $v^{(S)}$ with modest law-pole $v^{(L)}$ indicates complexity-driven mass attribution.

[8] "Composite" here means *triadic-channel composition* within the same field, not a preonic bound state.

- **Vacuum Higgs** $\hat{\phi}_{\mathrm{H}}^{(\mathrm{vac})}$: Sensitive to vacuum-manifold geometry and SSB topology; its Location bands track vacuum-sector windows (useful for defect/texture diagnostics).
- **Spectral Higgs** $\hat{\phi}_{\mathrm{H}}^{(\mathrm{spec})}$: Closest to the textbook Higgs of conventional QFT, with mass and residues governed by the spectral curvature of $\mathcal{V}_\Theta$ and standard couplings (e.g. Yukawa-like terms [82]).

In practice, experiments predominantly access the spec channel; the corr and vac channels are predictions of the triadic formalism and provide *additional* structure beyond the standard picture.

5. Composite mass and effective pole. Let $\Delta_{\mathrm{H}}(k)$ denote the Π_{H}-projected, triad-resolved propagator matrix $\left(\Delta_{\mathrm{H}}^{(\alpha\beta)}(k)\right)_{\alpha,\beta}$ with entries $\Delta_{\mathrm{H}}^{(\alpha\beta)} := \Pi_\alpha\,\Pi_{\mathrm{H}}\,\Delta_{\Theta\Theta}\,\Pi_{\mathrm{H}}\,\Pi_\beta$. The observable Higgs pole $k^2 = M_{\mathrm{H}}^2$ solves

$$\det\Big[k^2\mathbf{1} - \mathcal{M}_{\mathrm{H}}^{\mathrm{eff}}(k^2)\Big]_{(\alpha,\beta)} = 0, \qquad \mathcal{M}_{\mathrm{H}}^{\mathrm{eff}}(k^2) := \mathbb{K}_{\mathrm{H}}(k^2) + \Sigma_{\mathrm{H}}(k^2), \tag{9.197}$$

where $\mathbb{K}_{\mathrm{H}}$ is the radial kinetic block (typed and banded) and Σ_{H} the triadic self-energy, both restricted by Π_{H}. When cross-channel mixing (off-diagonal $\alpha \neq \beta$) is weak (small $v^{(LS)}$), one may approximate

$$M_{\mathrm{H}}^2 \;\simeq\; \sum_\alpha w_\alpha \left(M_{\mathrm{H}}^{(\alpha)}\right)^2, \qquad w_\alpha := \frac{\mathrm{Res}_{k^2=M_{\mathrm{H}}^2}\;\Delta_{\mathrm{H}}^{(\alpha\alpha)}(k)}{\sum_\gamma \mathrm{Res}_{k^2=M_{\mathrm{H}}^2}\;\Delta_{\mathrm{H}}^{(\gamma\gamma)}(k)}\,, \tag{9.198}$$

which makes precise the "composite mass" intuition in terms of channel residues (law-side) certified by meters (cf. eqs. (9.126) and (9.128)). The exact value is fixed by (9.197).

6. Symmetry and sectoral unification. The triadic composite picture mirrors the geometry sector:

- **Apeiron → gravitons (triadic).** In Gaiamiti, with channel-resolved species and T–L–O stamps (cf. Section 9.17).
- **Kymon → Higgs (triadic).** In Tantrakyma, identified via the radial projector and Hessian spectrum (cf. eqs. (9.175), (9.185) and (9.194)) and triadic meters.

In both sectors, the *spectral* species aligns with conventional particle notions, while the *correlational* and *vacuum* species are novel, IQR-specific predictions tied to quantum information structure and vacuum topology.

Narrative synthesis. The Higgs boson, as accessed in standard experiments, is the *spectral face* of a deeper composite object whose full identity is triadic. Parasymmetric T–L–O provides the operational calculus: typed kernels $\Delta_{\Theta\Theta}^{(\alpha,[\chi])}$, meters $(v^{(L)}, v^{(S)}, v^{(LS)})$, quadrant calls, and T–L–O stamps collectively certify the presence of a gapped, radial kymon and quantify its channel composition. This preserves the familiar phenomenology (e.g. Yukawa couplings [82]) while revealing additional correlational and vacuum contributions that enrich the mass-generating narrative within the IQR framework.

Regime & Scale	Gaiamiti Sector (Geometry)	Tantrakyma Sector (Matter)	Ontological Level	Parasymmetric T–L–O Signature (typed/triadic)
Ultra-sub-Planckian $\ll \ell_P$	Fibrewise Gaiamiti operator algebra in $\mathfrak{F}_x$; pre-metric correlations	Fibrewise Tantrakyma operator algebra in $\mathfrak{F}_x$; pre-matter correlations	Coherence-only domain; pre-geometric and pre-particle	*Tags:* $(\lambda_{\mathrm{UV}}, \ell)$ extreme. *Meters:* $v^{(L)}, v^{(S)} \approx 0$. *Quadrant:* QIV (no pole, no order) across channels $\alpha \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$; BRST projection to H^0 assumed (Definitions 9.306 and 9.307).
Planckian $\sim \ell_P$	Onset of stable modal structure; incipient Apeiron fluctuations	Onset of stable modal structure; incipient Kymon fluctuations	Proto-geometry and proto-particles	*Meters:* $v^{(L)}$ begins to rise in select types $[\chi]$; $v^{(S)}$ still low. *Quadrant:* QII (law-only incipience) in some typed channels (Section 9.17.11 and Definition 9.297).
Relational / Mesoscopic $\gtrsim \ell_{\mathrm{PQS}}^{-1}$	Expectation tensor $\bar{\Sigma}_{\mu\nu}$ emerges; semi-classical geometry	SSB in V_Θ; Higgs mechanism; typed mass operators $\mathcal{K}^{[\chi]}_{\Theta\Theta}$ from potential Hessian (eq. (9.175))	Emergent geometry and particle properties	*NG channels:* $v^{(L)} \neq 0$ with $k^2 = 0$ pole $\Rightarrow$ QI. *Higgs (radial):* gapped pole at $k^2 = M_H^2$ (no $k^2 = 0$), *Location* IR/meso, stamped via Definition 9.299. Cross-channel mixing quantified by $v^{(LS)}$ (eqs. (9.126) to (9.128) and (9.194)).
Macroscopic (IR) $\gg \ell_P$	Causal manifold $\mathcal{M}_{\mathrm{geom}}$; GR regime	Stable kymon quanta (incl. Higgs) and interactions; SM-like windows	Classical fields and stable particles	*Geometry:* long-time order robust; NG channels remain QI (massless). *Matter:* Higgs radial channel remains gapped; depending on tags, state-dominant windows can appear as QIII in some types; T–L–O stamps fix Onset thresholds (Definition 9.299).
Cosmological $\sim \mathcal{L}_U$	Global solutions of chronogeometric dynamics	Thermodynamics of kymon fields; cosmic backgrounds	Thermodynamic/statistical limit	Channel-wise vacuum energetics regulated via triadic balance with geometry (Section 9.18.8); meters certify plateau stability ($v^{(S)}$) and gauge-independence (H^0). Effective large-scale behavior often *state-dominant* (tending toward QIII) depending on windowing.

Table 9.3: Unified hierarchy of emergent geometric and matter regimes with typed/triadic Parasymmetric T–L–O certification. Meters and quadrants per eqs. (9.126) to (9.128), Definition 9.297, and Proposition 9.298; typed mass operators and Higgs identification per Definition 9.295 and eqs. (9.175), (9.185) and (9.194).

The Higgs Boson: Parallel Challenges and Triadic Resolutions

Just as the graviton's masslessness is clarified when viewed through the typed, triadic Gaiamiti instrumentation, the Higgs boson reveals its own constellation of conceptual challenges that invite a *parasymmetric T–L–O* reading. In the IQR framework, the Higgs is the *radial, scalar kymon* of the Tantrakyma sector (Section 9.18.5 and eq. (9.194)), and its identification is *diagnostic realized reference configuration-certified* by typed kernels, SynTh meters, quadrant calls, and T–L–O stamps (Section 9.17.11, Definitions 9.296, 9.297 and 9.299, and Proposition 9.298). This section reframes three classic paradoxes with explicit reference to the typed/triadic structure and the quadrant semantics.

Quadrant prelude (typed phase semantics). At fixed tags $(\lambda_{\mathrm{UV}}, \ell)$ on a certified plateau $\mathfrak{P}$ (Definitions 9.306 and 9.307), the SynTh meters $(v^{(L)}, v^{(S)}, v^{(LS)})$ (eqs. (9.126) to (9.128)) feed the typed predicate $\mathrm{Quad}^{[\Theta\Theta]}_{[\chi]} \in \{\mathrm{QI}, \mathrm{QII}, \mathrm{QIII}, \mathrm{QIV}\}$ (Definition 9.297). In brief:

$$\begin{aligned} \mathrm{QI}: &\ \text{NG/pNG present and ordered;} \\ \mathrm{QII}: &\ \text{incipient law (critical) without long-time order;} \\ \mathrm{QIII}: &\ \text{state-only order;} \\ \mathrm{QIV}: &\ \text{symmetric (no pole, no order).} \end{aligned}$$

The *Higgs (radial) channel* is gapped and stamps with a nonzero Location band (IR/meso) and finite Onset (eqs. (9.185) and (9.194) and Definition 9.299); NG channels stamp QI with a $k^2 = 0$ pole.

1. The Hierarchy Problem: a question of emergent scales

Problem. The Higgs mass appears unnaturally small relative to an ultraviolet scale (often taken as the Planck scale), inviting fine-tuning [91, 112].

Triadic T–L–O resolution (IQR). In IQR, the UV threshold is an emergent diagnostic realized reference configuration tag rather than a fundamental constant (cf. *sub/ultra-Planck* discussion), and the observed Higgs is a *composite* radial kymon across the triadic channels (Section 9.18.8). Operationally, the Π_{H}-projected, channel-resolved propagator $\Delta_{\mathrm{H}}^{(\alpha\beta)}(k)$ determines the physical pole $k^2 = M_{\mathrm{H}}^2$ via the triadic pole condition (eq. (9.197)). When off-diagonal mixing is small, the effective mass admits the residue-weighted approximation (eq. (9.198)):

$$M_{\mathrm{H}}^2 \simeq \sum_{\alpha \in \{\mathrm{corr,vac,spec}\}} w_\alpha \left(M_{\mathrm{H}}^{(\alpha)}\right)^2, \quad w_\alpha \propto \mathrm{Res}\ \Delta_{\mathrm{H}}^{(\alpha\alpha)}.$$

Thus the "hierarchy" becomes a *channel composition* question governed by meters: (i) $v^{(L)}$ nails the law-side pole content per channel; (ii) $v^{(S)}$ constrains long-time order (stability of the low mass); (iii) $v^{(LS)}$ quantifies cross-channel seesaw/cancellations. Quadrant diagnostics localize the phenomenon: near-critical behavior registers QII in specific channels, explaining light composite masses without fine-tuning of a single bare parameter.

2. The Nature of the Higgs Potential: a question of triadic origin

Problem. The Standard Model posits a quartic "wine-bottle" potential $\mathcal{V}(\phi)$ by fiat [82].

Triadic T–L–O resolution (IQR). IQR derives the Tantrakyma potential from cotangent invariants ($\mathsf{X} = g_C^{-1}(\hat{\mathbf{\Theta}}, \hat{\mathbf{\Theta}})$); Definition 9.335, Remark 9.303, and eq. (9.133)) and admits a *triadic decomposition*

$$\mathcal{V}_\Theta \;=\; \mathcal{V}_\Theta^{(\mathrm{corr})} \;+\; \mathcal{V}_\Theta^{(\mathrm{vac})} \;+\; \mathcal{V}_\Theta^{(\mathrm{spec})},$$

consistent with the commuting channel projectors (Definition 9.295 and eq. (9.195)). The sign of the quadratic coefficient (driving SSB) is an *emergent balance* among channel contributions. Meters and quadrant calls track how a negative correlational contribution (information-driven) can outweigh positive spectral/vacuum pieces, pushing the system from QIV (symmetric) through QII (incipient) to QI (NG present), while the radial channel registers a gapped stamp with mass fixed by $\partial^2 V_\Theta$ at the minimum (eq. (9.194)). Thus the potential's profile is *explained* by channel competition rather than postulated.

3. The Cosmological Constant Problem: a question of a triadic vacuum

Problem. Naïve vacuum energy estimates from the Higgs sector vastly exceed the observed cosmological constant [211, 212].

Triadic T–L–O resolution (IQR). The total diagnostic realized reference configuration stress–energy is $\bar{T}^{\mu\nu}_{\mathrm{total}} = \bar{T}^{\mu\nu}_{(\Sigma)} + \bar{T}^{\mu\nu}_{(\Theta)}$ (Definition 9.358 and Sections 9.17.9 and 9.17.11). In the triadic split, *each channel* contributes and must *self-regulate*:

$$\langle \mathcal{V}_\Theta^{(\mathrm{corr})} \rangle + \langle \mathcal{V}_\Sigma^{(\mathrm{corr})} \rangle \approx 0, \quad \langle \mathcal{V}_\Theta^{(\mathrm{vac})} \rangle + \langle \mathcal{V}_\Sigma^{(\mathrm{vac})} \rangle \approx 0, \quad \langle \mathcal{V}_\Theta^{(\mathrm{spec})} \rangle + \langle \mathcal{V}_\Sigma^{(\mathrm{spec})} \rangle \approx 0,$$

up to a small residual that reads as the observed cosmological constant. Parasymmetric meters provide the consistency checks: state-side plateaus $v^{(S)}$ certify long-time vacuum stability per channel; law-side content $v^{(L)}$ ensures the cancellations are not gauge-artifacts (BRST/Nielsen clean). This structured, channel-wise mechanism replaces a brute-force two-number cancellation.

Synthesis: from paradox to a composite principle

The triadic, typed, parasymmetric T–L–O formalism turns the Higgs challenges into *operational questions* about channel composition, quadrant placement, and diagnostic realized reference configuration-certified meters. The Higgs boson emerges as a *composite radial kymon* whose mass, potential, and vacuum contributions are distributed across correlational, vacuum, and spectral channels (Section 9.18.8). The same logic applies mutatis mutandis in the geometry sector for the graviton, yielding a sector-by-sector symmetry (Sections 9.17 and 9.17.10). Rather than invoking new ultra-heavy particles, IQR attributes the observed phenomena to the *diagnostic realized reference configuration-instrumented* structure of reality—triadic channels, typed projections, SynTh meters, and quadrant semantics—which together determine Type, Location, and Onset of the excitations we call particles. In this sense, what appear as paradoxes at the Lagrangian level become *composite principles* when read through Parasymmetric T–L–O.

9.18.9 Euler–Lagrange Equation for the Tantrakyma Field

The dynamics of the Tantrakyma field are determined by applying the *Principle of Stationary Action* to its action functional $S_\Theta[\hat{\mathbf{\Theta}}]$ (cf. Definition 9.310). By extremizing this action with respect to the field $\hat{\mathbf{\Theta}}(x)$, we obtain the operator-valued Euler–Lagrange (EL) equation that governs matter-like excitations on the emergent quantum manifold.

Definition 9.353 (Euler–Lagrange equation for $\hat{\mathbf{\Theta}}(x)$)**.** Let $\hat{\mathbf{\Theta}}(x)$ be the Tantrakyma field operator on the relational manifold $(\mathcal{X}_q, \bar{\Sigma}_{\mu\nu})$, with dynamics governed by $S_\Theta[\hat{\mathbf{\Theta}}]$. Then the corresponding operator-valued Euler–Lagrange equation can be written in divergence form as

$$\boxed{\frac{1}{\sqrt{-\det \bar{\Sigma}}}\,\nabla_\mu\Big(\sqrt{-\det \bar{\Sigma}}\;\bar{\Sigma}^{\mu\nu}\, g_C^\sharp\Big(\nabla_\nu \hat{\mathbf{\Theta}},\,\cdot\,\Big)\Big) \;=\; \frac{\delta \mathcal{V}_\Theta}{\delta \hat{\mathbf{\Theta}}^\dagger}(x),} \tag{9.199}$$

where:

(i) Total covariant derivative
$\nabla_\mu \hat{\mathbf{\Theta}}(x)$ is the total covariant derivative (relational Levi–Civita + operator-bundle/gauge connection; cf. Definition 9.323).

(ii) Inverse chronogeometric tensor
$\bar{\Sigma}^{\mu\nu}(x)$ is the inverse chronogeometric tensor (cf. Definition 6.24).

(iii) Co-metric Riesz map
$g_C^\sharp$ is the Riesz map associated with the operator-valued co-metric g_C, sending the second slot to an $\mathrm{End}(\mathcal{H}_{\mathrm{em}})$-valued element (cf. Remark 9.303).

(iv) Potential gradient
$\delta \mathcal{V}_\Theta / \delta \hat{\mathbf{\Theta}}^\dagger$ is the functional derivative of the Tantrakyma potential built from the invariant $\mathsf{X} = g_C(\hat{\mathbf{\Theta}}, \hat{\mathbf{\Theta}})$ (cf. Definition 9.335).

Using $\nabla_\mu^{(\mathrm{rel})} \bar{\Sigma}_{\alpha\beta} = 0$ and $D_\mu g_C = 0$ (cf. Definition 6.7 and Proposition 9.324), (9.199) is equivalently

$$\bar{\Sigma}^{\mu\nu}\,\nabla_\mu\Big(g_C^\sharp\Big(\nabla_\nu \hat{\mathbf{\Theta}},\,\cdot\,\Big)\Big) \;=\; \frac{\delta \mathcal{V}_\Theta}{\delta \hat{\mathbf{\Theta}}^\dagger}(x), \tag{9.200}$$

and, since $g_C^\sharp$ is covariantly constant, this further reduces to $\bar{\Sigma}^{\mu\nu}\, g_C^\sharp\Big(\nabla_\mu \nabla_\nu \hat{\mathbf{\Theta}},\,\cdot\,\Big) = \delta \mathcal{V}_\Theta / \delta \hat{\mathbf{\Theta}}^\dagger$. The companion adjoint equation follows from varying $\hat{\mathbf{\Theta}}$ instead of $\hat{\mathbf{\Theta}}^\dagger$.

Diagnostic realized reference configuration-ready (band-limited, BRST-clean) EL. On a certified plateau $\mathfrak{P}$ with tags $(\lambda_{\mathrm{UV}}, \ell)$ and physical projector $\mathsf{P}_{\mathrm{phys}} : \mathcal{H} \to H^0(\hat{Q}_{\mathrm{BRST}})$ (cf. Definitions 9.306 and 9.307), the working EL equation is

$$\mathsf{P}_{\mathrm{phys}}\left[\frac{1}{\sqrt{-\det \bar{\Sigma}}}\,\nabla_\mu\Big(\sqrt{-\det \bar{\Sigma}}\;\bar{\Sigma}^{\mu\nu}\, g_C^\sharp\Big(\nabla_\nu^{(\lambda_{\mathrm{UV}},\ell)} \hat{\mathbf{\Theta}}^{(\lambda_{\mathrm{UV}},\ell)},\,\cdot\,\Big)\Big) - \frac{\delta \mathcal{V}_\Theta}{\delta \hat{\mathbf{\Theta}}^\dagger}\right] = 0, \tag{9.201}$$

which is BRST-closed and gauge-parameter independent (Nielsen-flat).

Typed, parasymmetric EL (projection to types and channels). Let $\Pi^{[\chi]}_{\Theta\Theta}$ be the typed projectors ($[\chi] \in \{\mathrm{CK}, \mathrm{T}_\pm, \mathrm{L}\}$; cf. Definition 9.295) commuting with the triadic channel projectors Π_α, $\alpha \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$. The typed EL equations are

$$\Pi^{[\chi]}_{\Theta\Theta}\left[\frac{1}{\sqrt{-\det\bar{\Sigma}}}\,\nabla_\mu\Big(\sqrt{-\det\bar{\Sigma}}\;\bar{\Sigma}^{\mu\nu}\,g^\sharp_C\big(\nabla_\nu\hat{\boldsymbol{\Theta}},\,\cdot\,\big)\Big) - \frac{\delta\mathcal{V}_\Theta}{\delta\hat{\boldsymbol{\Theta}}^\dagger}\right] = 0, \qquad [\Pi^{[\chi]}_{\Theta\Theta}, \Pi_\alpha] = 0. \quad (9.202)$$

This prepares the EL system for *typed* Parasymmetric T–L–O analysis.

Linearized, typed EL and the Hessian link. With the background–fluctuation split $\hat{\boldsymbol{\Theta}} = \bar{\Theta}\,\mathbb{I} + \delta\hat{\boldsymbol{\Theta}}$ (cf. Definition 10.204), the linearized, typed EL takes the standard form

$$\boxed{\big(\mathbb{K}^{[\chi]}_\Theta + \mathcal{K}^{[\chi]}_{\Theta\Theta}\big)\,\delta\hat{\boldsymbol{\Theta}}^{[\chi]} = \hat{\mathcal{J}}^{[\chi]}, \qquad \delta\hat{\boldsymbol{\Theta}}^{[\chi]} := \Pi^{[\chi]}_{\Theta\Theta}\,\delta\hat{\boldsymbol{\Theta}},} \quad (9.203)$$

where $\mathbb{K}^{[\chi]}_\Theta$ is the typed kinetic block (from $\mathcal{T}_\Theta$; cf. eq. (9.130)), and $\mathcal{K}^{[\chi]}_{\Theta\Theta}$ is the typed potential Hessian (mass/curvature operator) defined by restriction (cf. eq. (9.175)). The typed retarded kernels $\Delta^{[\chi]}_{\Theta\Theta}$ are inverses of the EL operators in (9.203).

Parasymmetric T–L–O interpretation (meters → quadrants → stamps). From $\Delta^{[\chi]}_{\Theta\Theta}$ build the SynTh meters (cf. eqs. (9.126) to (9.128)):

$$\begin{aligned} v^{(L)}_{[\chi]} &= \big|\mathrm{Res}_{k^2=0}\,\Delta^{[\chi]}_{\Theta\Theta}\big|, \\ v^{(S)}_{[\chi]} &= \big|\Delta^{[\chi]}_{\Theta\Theta}(t\to\infty;\mathbf{x}=0)\big|_{H^0}, \\ v^{(LS)}_{[\chi]} &= \text{helicity-resolved cross-channel gain}, \end{aligned}$$

assemble the typed witness $\mathbf{W}^{[\Theta\Theta]}_{[\chi]}$ (cf. Definition 9.296), and apply the typed quadrant predicate $\mathrm{Quad}^{[\Theta\Theta]}_{[\chi]} \in \{\mathrm{QI}, \mathrm{QII}, \mathrm{QIII}, \mathrm{QIV}\}$ (cf. Definition 9.297 and Proposition 9.298). Consistency requires that NG/pNG versus gapped expectations from $\mathrm{spec}(\mathcal{K}^{[\chi]}_{\Theta\Theta})$ match the quadrant call within tolerances (Ch. 8). The certified outcome is recorded as a *typed T–L–O stamp* (cf. Definition 9.299) with *Location* (IR/meso/UV windows) and *Onset* thresholds.

Noether current and BRST/Nielsen safety. On-shell EL implies covariant conservation of the matter-sector Noether current $\hat{J}^\mu_\Theta$ built from $\mathcal{L}_\Theta$ (cf. Section 9.17.11), providing additional Location diagnostics consistent with the typed meters. All statements are made on H^0 and are therefore gauge-parameter independent (Nielsen-flat), as guaranteed by the diagnostic realized reference configuration discipline (cf. Definitions 9.306 and 9.307).

Interpretation Equation (9.199) is the foundational law for how *relational material content*, encoded by the Tantrakyma field $\hat{\boldsymbol{\Theta}}(x)$, evolves within the emergent spacetime. It generalizes a relativistic wave equation to the full *chronogeometric* arena: the propagation operator is built from the unified tensor $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (Definitions 5.26 and 6.24), rather than the

metric alone. Consequently, kymon dynamics are simultaneously sensitive to the curvature of distinguishability ($\bar{g}_{\mu\nu}$) and to the curvature of quantum phase flow ($\bar{t}_{\mu\nu}$). In the expanded form of the EL operator one encounters a term $\big(\nabla_\mu \bar{\Sigma}^{\mu\nu}\big)\nabla_\nu \hat{\mathbf{\Theta}}$; however, with the $\bar{\Sigma}$-compatible (relational Levi–Civita) connection used here this term *vanishes identically*, so propagation is governed by $\bar{\Sigma}^{\mu\nu}\, g_C^\sharp\big(\nabla_\mu\nabla_\nu \hat{\mathbf{\Theta}}, \cdot\,\big)$. All derivatives are taken with the composite covariant derivative ∇_μ that combines the relational Levi–Civita and operator-bundle connections (Definition 9.323).

The right-hand side $\delta\mathcal{V}_\Theta/\delta\hat{\mathbf{\Theta}}^\dagger$ supplies the *identity* of the excitations via the Tantrakyma potential (Definition 9.335), which—because $\hat{\mathbf{\Theta}}$ is *covector*-valued—must be built from the co-metric invariant $\mathsf{X} = g_C(\hat{\mathbf{\Theta}}, \hat{\mathbf{\Theta}})$, not from g_C^{-1}. Around a realized vacuum $\bar{\Theta}$ (eq. (9.134)), the background–fluctuation split $\hat{\mathbf{\Theta}} = \bar{\Theta}\,\mathbb{I} + \delta\hat{\mathbf{\Theta}}$ leads to the *typed* linear response (eq. (9.203)): the kinetic block $\mathbb{K}_\Theta^{[\chi]}$ pairs with the typed Hessian $\mathcal{K}_{\Theta\Theta}^{[\chi]}$ extracted from the potential by projector restriction (eq. (9.175)). The spectrum (NG/pNG vs. gapped) is therefore read directly from $\operatorname{spec}\mathcal{K}_{\Theta\Theta}^{[\chi]}$, providing, in particular, the mass of the radial (Higgs) kymon in the SSB phase (eq. (9.194) and Section 9.18.5).

Parasymmetric T–L–O reading. On a certified plateau $\mathfrak{P}$ with tags $(\lambda_{\mathrm{UV}}, \ell)$, we project the EL system to types $[\chi]$ and channels $\alpha \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$ using BRST-closed, triad-compatible projectors (Definition 9.295). The corresponding typed retarded kernels $\Delta_{\Theta\Theta}^{[\chi]}$ drive the SynTh meters $(v^{(L)}, v^{(S)}, v^{(LS)})$ (eqs. (9.126) to (9.128)), assembled into the typed witness $\mathbf{W}_{[\chi]}^{[\Theta\Theta]}$ (Definition 9.296). The *quadrant predicate* $\mathrm{Quad}_{[\chi]}^{[\Theta\Theta]}$ (Definition 9.297 and Proposition 9.298) then certifies phase semantics: NG channels stamp QI with a $k^2 = 0$ pole; the Higgs (radial) channel stamps as *gapped* (no massless pole) with a *Location* band (IR/meso) and an *Onset* threshold recorded by the T–L–O entry (Definition 9.299). Thus, the EL equation is not only the formal equation of motion; it is the *generator* of operational diagnostics—linking chronogeometric propagation (left-hand side) and identity-defining self-interactions (right-hand side) to measurable meters, quadrant calls, and stamps on $H^0(\hat{Q}_{\mathrm{BRST}})$.

9.18.10 Noether Current for the Dynamical Tantrakyma Field

The action $S_\Theta[\hat{\mathbf{\Theta}}]$ (cf. Definition 9.310) admits internal fiber symmetries acting on the excitation sector via local unitary automorphisms of $\mathrm{End}(\mathcal{H}_{\mathrm{em}})$ (cf. Definition 9.323 and the symmetry remarks in Definition 9.301). When S_Θ is invariant under a continuous family of such transformations, Noether's theorem furnishes a (conserved) current that diagnoses the local flow of excitation charge across $\mathcal{X}_q$. Throughout, recall that the Tantrakyma *potential* is built with the co-metric g_C (not g_C^{-1}).

We consider infinitesimal, local unitary variations

$$\delta\hat{\mathbf{\Theta}}(x) = i\,\epsilon^a(x)\,[T_a,\, \hat{\mathbf{\Theta}}(x)], \qquad [T_a, T_b] = i f_{ab}{}^c T_c, \tag{9.204}$$

with Hermitian generators T_a and small real functions $\epsilon^a(x)$. The total covariant derivative is $\nabla_\mu = \nabla_\mu^{(\mathrm{rel})} + D_\mu$ (cf. Definition 9.323).

Definition 9.354 (Noether current for the Tantrakyma field)**.** If the band-limited, BRST-clean Lagrangian density $\mathcal{L}_\Theta$ is invariant under (9.204) up to a total divergence, the associated

operator-level current kernel is

$$\boxed{\widehat{\mathcal{J}}^{\mu}_{\Theta;a}(x) := -\,i\,\hat{\Pi}^{\mu}_{\Theta}(x)\Big([T_a,\,\hat{\mathbf{\Theta}}(x)]\Big)\;\in\;\mathrm{End}(\mathcal{H}_{\mathrm{em}}),} \tag{9.205}$$

where $\hat{\Pi}^{\mu}_{\Theta}$ is the canonical momentum (operator) map defined by

$$\boxed{\hat{\Pi}^{\mu}_{\Theta}(x)\Big[\,\delta\hat{\mathbf{\Theta}}\,\Big]\;:=\;\bar{\Sigma}^{\mu\nu}(x)\;g_C\Big(\delta\hat{\mathbf{\Theta}}(x),\,\nabla_\nu\hat{\mathbf{\Theta}}(x)\Big).} \tag{9.206}$$

Equivalently, in a symmetrized (manifestly Hermitian) form one may write

$$\widehat{\mathcal{J}}^{\mu}_{\Theta;a} = \frac{i}{2}\,\bar{\Sigma}^{\mu\nu}\Big[g_C\Big([T_a,\hat{\mathbf{\Theta}}],\,\nabla_\nu\hat{\mathbf{\Theta}}\Big) - g_C\Big(\nabla_\nu\hat{\mathbf{\Theta}},\,[T_a,\hat{\mathbf{\Theta}}]\Big)\Big]\,.$$

The corresponding *scalar* Noether current density is obtained by tracing over the fiber algebra:

$$J^{\mu}_{\Theta;a}(x)\;:=\;\mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\Big\{\widehat{\mathcal{J}}^{\mu}_{\Theta;a}(x)\Big\}\,.$$

On-shell conservation (total covariant form). For compactly supported ϵ^a, variation of the action—after integration by parts and using the EL equations (9.199)—yields the covariant conservation law

$$\boxed{\nabla_\mu\,\widehat{\mathcal{J}}^{\mu}_{\Theta;a}(x)\;=\;0\quad\text{on-shell},} \tag{9.207}$$

where $\nabla_\mu = \nabla^{(\mathrm{rel})}_\mu + i[\mathcal{A}_\mu,\,\cdot\,]$ acts on the spacetime index and on the fiber operator (cf. Definition 9.323). Taking the fiber trace gives $\nabla_\mu J^{\mu}_{\Theta;a} = 0$. For a pure internal symmetry with anomaly-free gauging this is equivalent to the gauge-covariant statement $D_\mu J^{\mu}_{\Theta;a} = 0$.

Diagnostic realized reference configuration (band-limited, BRST-clean) form and expectation current. On a certified plateau $\mathfrak{P}$ with tags $(\lambda_{\mathrm{UV}},\ell)$ and physical projector $\mathsf{P}_{\mathrm{phys}} : \mathcal{H}\to H^0(\hat{Q}_{\mathrm{BRST}})$ (cf. Definitions 9.306 and 9.307), we use $\hat{\mathbf{\Theta}}^{(\lambda_{\mathrm{UV}},\ell)}$ and $\nabla^{(\lambda_{\mathrm{UV}},\ell)}_\mu$. Define the band-limited canonical momentum (operator) map by

$$\hat{\Pi}^{\mu\,(\lambda_{\mathrm{UV}},\ell)}_{\Theta}\Big[\,\delta\hat{\mathbf{\Theta}}\,\Big] := \bar{\Sigma}^{\mu\nu}(x)\;g_C\Big(\delta\hat{\mathbf{\Theta}}(x),\,\nabla^{(\lambda_{\mathrm{UV}},\ell)}_\nu\hat{\mathbf{\Theta}}^{(\lambda_{\mathrm{UV}},\ell)}(x)\Big),$$

and the operator-level current kernel by

$$\widehat{\mathcal{J}}^{\mu\;(\lambda_{\mathrm{UV}},\ell)}_{\Theta;a}(x) := -\,i\,\hat{\Pi}^{\mu\,(\lambda_{\mathrm{UV}},\ell)}_{\Theta}\Big([T_a,\,\hat{\mathbf{\Theta}}^{(\lambda_{\mathrm{UV}},\ell)}(x)\Big). \tag{9.208}$$

The scalar current is the fiber trace

$$J^{\mu\;(\lambda_{\mathrm{UV}},\ell)}_{\Theta;a}(x) := \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\Big\{\widehat{\mathcal{J}}^{\mu\;(\lambda_{\mathrm{UV}},\ell)}_{\Theta;a}(x)\Big\}\,,$$

and the realized (c-number) current on H^0 is

$$\bar{J}^{\mu}_{\Theta;a}(x) := \Big\langle\,\mathring{\mathbf{\Psi}}\;\Big|\;\mathsf{P}_{\mathrm{phys}}\,\mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\{\widehat{\mathcal{J}}^{\mu\;(\lambda_{\mathrm{UV}},\ell)}_{\Theta;a}(x)\}\;\Big|\;\mathring{\mathbf{\Psi}}\,\Big\rangle,\qquad \nabla_\mu\bar{J}^{\mu}_{\Theta;a} = 0\;\;\text{on-shell}.$$

Typed & triadic resolution (parasymmetric). Let $\Pi_{\Theta\Theta}^{[\chi]}$ be the typed projectors ($[\chi] \in \{\mathrm{CK}, \mathrm{T}_\pm, \mathrm{L}\}$; cf. Definition 9.295) and Π_α the channel projectors $\alpha \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$, with $[\Pi_\alpha, \Pi_{\Theta\Theta}^{[\chi]}] = 0$. The typed/channel-resolved current kernels are

$$\widehat{\mathcal{J}}^{\mu\ (\alpha,[\chi])}_{\Theta;a} := \Pi_\alpha\, \Pi_{\Theta\Theta}^{[\chi]}\, \widehat{\mathcal{J}}^{\mu}_{\Theta;a}\, \Pi_{\Theta\Theta}^{[\chi]}\, \Pi_\alpha, \qquad J^{\mu\ (\alpha,[\chi])}_{\Theta;a} := \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\Big\{\widehat{\mathcal{J}}^{\mu\ (\alpha,[\chi])}_{\Theta;a}\Big\}. \tag{9.209}$$

Their retarded two-point functions provide *Location* diagnostics: peaks in the spectral density of $\langle \widehat{\mathcal{J}}\, \widehat{\mathcal{J}} \rangle$ co-locate with the $\Delta_{\Theta\Theta}$ poles that certify NG channels (QI) or the gapped Higgs (radial) channel, reinforcing T–L–O stamps (cf. Definition 9.299 and eqs. (9.126) to (9.128)).

Consistency with EL and symmetry of $\mathcal{V}_\Theta$. Using the EL equation (9.199) and the fact that the potential depends only on the co-metric invariant $\mathsf{X} = g_C(\hat{\mathbf{\Theta}}, \hat{\mathbf{\Theta}})$ (hence is invariant under fiber inner $*$-automorphisms; cf. Definition 9.335 and Remark 9.313), one obtains $\nabla_\mu \widehat{\mathcal{J}}^{\mu}_{\Theta;a} = 0$ on-shell. Spontaneous symmetry breaking (SSB) is signaled by a nonzero $\bar{\Theta}$ and manifests as *massless* NG channels (QI stamp) together with a *gapped* radial (Higgs) channel, as certified by the typed meters and quadrant predicate (cf. Section 9.18.5, eq. (9.194), and Definition 9.297).

Boundary pass. On $\mathfrak{P}_\partial$, conservation reads as vanishing flux through the boundary (up to specified sources), providing an *edge* Location diagnostic that must agree with bulk T–L–O stamping within tolerance budgets (cf. Proposition 9.300).

Expectation-Level Current As in the geometric (Gaiamiti) sector, the *observable* conserved quantity is the expectation value of the current in the Quantum Holo-State (cf. Definition 1.27). On a certified plateau $\mathfrak{P}$ with tags $(\lambda_{\mathrm{UV}}, \ell)$ and projection to the physical cohomology $H^0(\hat{Q}_{\mathrm{BRST}})$, the BRST-clean, band-limited current is

$$\boxed{\bar{J}^{\mu}_{\Theta;a}(x) := \Big\langle \mathring{\mathbf{\Psi}} \Big|\, \mathsf{P}_{\mathrm{phys}}\, \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\Big\{\widehat{\mathcal{J}}^{\mu\ (\lambda_{\mathrm{UV}},\ell)}_{\Theta;a}(x)\Big\}\, \Big|\, \mathring{\mathbf{\Psi}}\Big\rangle, \qquad \nabla_\mu \bar{J}^{\mu}_{\Theta;a} = 0 \ \text{ on-shell},} \tag{9.210}$$

where $\widehat{\mathcal{J}}^{\mu}_{\Theta;a}$ is the operator-level current kernel of Definition 9.354 (cf. (9.208)), $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ is the band functor (cf. Definition 9.306), and $\mathsf{P}_{\mathrm{phys}}$ projects onto $H^0(\hat{Q}_{\mathrm{BRST}})$. This guarantees gauge-parameter independence (Nielsen flatness) of $\bar{J}^{\mu}_{\Theta;a}$.

Typed & triadic expectation (parasymmetric). With typed projectors $\Pi_{\Theta\Theta}^{[\chi]}$ and channel projectors Π_α that commute (cf. Definition 9.295), the resolved expectation currents are

$$\bar{J}^{\mu\ (\alpha,[\chi])}_{\Theta;a}(x) := \Big\langle \mathring{\mathbf{\Psi}} \Big|\, \mathsf{P}_{\mathrm{phys}}\, \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\Big\{\Pi_\alpha\, \Pi_{\Theta\Theta}^{[\chi]}\, \widehat{\mathcal{J}}^{\mu\ (\lambda_{\mathrm{UV}},\ell)}_{\Theta;a}(x)\, \Pi_{\Theta\Theta}^{[\chi]}\, \Pi_\alpha\Big\}\, \Big|\, \mathring{\mathbf{\Psi}}\Big\rangle, \qquad [\Pi_\alpha, \Pi_{\Theta\Theta}^{[\chi]}] = 0, \tag{9.211}$$

which provide *Location* diagnostics consistent with the typed SynTh meters $(v^{(L)}, v^{(S)}, v^{(LS)})$ and the T–L–O stamps (cf. eqs. (9.126) to (9.128) and Definition 9.299). In NG channels (QI) one finds characteristic low-frequency spectral weight aligned with the $k^2 = 0$ pole of $\Delta_{\Theta\Theta}$; in the Higgs (radial) channel the spectral peak co-locates with the gapped pole $k^2 = M_{\mathrm{H}}^2$ (cf. eq. (9.194)), corroborating the quadrant call (cf. Definition 9.297 and Proposition 9.298).

Proposition 9.355 (Conservation of the Tantrakyma Noether Current)**.** *If the Tantrakyma field $\hat{\mathbf{\Theta}}(x)$ satisfies its Euler–Lagrange equation (9.199) and the action is invariant under the internal fiber automorphism generated by a Hermitian T_a (global parameter), then the expectation-level Noether current component $\bar{J}^{\mu}_{\Theta;a}(x)$ (cf. eq. (9.210)) is covariantly conserved with respect to the relational connection:*

$$\boxed{\nabla^{(\mathrm{rel})}_{\mu}\, \bar{J}^{\mu}_{\Theta;a}(x) = 0 \quad \textit{(on-shell).}} \tag{9.212}$$

If the symmetry is gauged, the statement upgrades to gauge-covariant *conservation, $D_{\mu}\, \bar{J}^{\mu}_{\Theta;a} = 0$ (cf. Definition 9.323).*

Proof of Proposition Proposition 9.355. We use a localized-parameter Noether variation and then impose the Euler–Lagrange (EL) equation (9.199).

(i) Localized internal variation. Let

$$\delta\hat{\mathbf{\Theta}}(x) = i\,\epsilon(x)\,[T_a,\, \hat{\mathbf{\Theta}}(x)], \tag{9.213}$$

with smooth $\epsilon(x)$. With the total covariant derivative $\nabla_{\mu} = \nabla^{(\mathrm{rel})}_{\mu} + D_{\mu}$ (cf. Definition 9.323),

$$\delta\Big(\nabla_{\mu}\hat{\mathbf{\Theta}}\Big) = \nabla_{\mu}\Big(\delta\hat{\mathbf{\Theta}}\Big) = i\,(\partial_{\mu}\epsilon)\,[T_a, \hat{\mathbf{\Theta}}] + i\,\epsilon\,[T_a,\, \nabla_{\mu}\hat{\mathbf{\Theta}}]. \tag{9.214}$$

(ii) Variation of the Lagrangian density. For the (scalar) Lagrangian density $\mathcal{L}_{\Theta}$ (cf. Definition 9.301),

$$\delta\mathcal{L}_{\Theta} = \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\left\{ \frac{\delta\mathcal{L}_{\Theta}}{\delta\hat{\mathbf{\Theta}}}\,\delta\hat{\mathbf{\Theta}} + \frac{\delta\mathcal{L}_{\Theta}}{\delta(\nabla_{\mu}\hat{\mathbf{\Theta}})}\,\delta(\nabla_{\mu}\hat{\mathbf{\Theta}}) \right\}. \tag{9.215}$$

Insert (9.213)–(9.214) and separate the ϵ- and $\partial_{\mu}\epsilon$-terms:

$$\delta\mathcal{L}_{\Theta} = i\,\epsilon\,\mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\left\{ \frac{\delta\mathcal{L}_{\Theta}}{\delta\hat{\mathbf{\Theta}}}[T_a, \hat{\mathbf{\Theta}}] + \frac{\delta\mathcal{L}_{\Theta}}{\delta(\nabla_{\mu}\hat{\mathbf{\Theta}})}[T_a, \nabla_{\mu}\hat{\mathbf{\Theta}}] \right\} + i\,(\partial_{\mu}\epsilon)\,\mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\left\{ \frac{\delta\mathcal{L}_{\Theta}}{\delta(\nabla_{\mu}\hat{\mathbf{\Theta}})}[T_a, \hat{\mathbf{\Theta}}] \right\}. \tag{9.216}$$

(iii) On-shell reduction and total divergence. Using the EL identity (on-shell) $\frac{\delta\mathcal{L}_{\Theta}}{\delta\hat{\mathbf{\Theta}}} = \nabla_{\mu}\Big(\frac{\delta\mathcal{L}_{\Theta}}{\delta(\nabla_{\mu}\hat{\mathbf{\Theta}})}\Big)$ and cyclicity of the trace, we obtain

$$i\,\epsilon\,\mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\left\{ \frac{\delta\mathcal{L}_{\Theta}}{\delta\hat{\mathbf{\Theta}}}[T_a, \hat{\mathbf{\Theta}}] + \frac{\delta\mathcal{L}_{\Theta}}{\delta(\nabla_{\mu}\hat{\mathbf{\Theta}})}[T_a, \nabla_{\mu}\hat{\mathbf{\Theta}}] \right\} = i\,\epsilon\,\nabla^{(\mathrm{rel})}_{\mu}\,\mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\left\{ \frac{\delta\mathcal{L}_{\Theta}}{\delta(\nabla_{\mu}\hat{\mathbf{\Theta}})}[T_a, \hat{\mathbf{\Theta}}] \right\}. \tag{9.217}$$

Define the *scalar* Noether current (trace of the operator-level kernel, cf. Definition 9.354)

$$J^{\mu}_{\Theta;a} := -\,i\,\mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\left\{ \frac{\delta\mathcal{L}_{\Theta}}{\delta(\nabla_{\mu}\hat{\mathbf{\Theta}})}\,[T_a, \hat{\mathbf{\Theta}}] \right\}. \tag{9.218}$$

Then, by the Leibniz rule for $\nabla^{(\mathrm{rel})}$,

$$\delta\mathcal{L}_{\Theta} = -\,\nabla^{(\mathrm{rel})}_{\mu}\Big(\epsilon\, J^{\mu}_{\Theta;a}\Big). \tag{9.219}$$

(iv) Localized-parameter argument & conservation. Integrating over $\mathcal{X}_q$ with $d\mu = d^d x \sqrt{-\det \bar{\Sigma}}$ and integrating by parts (discarding boundary terms under the diagnostic realized reference configuration IR window) yields

$$\delta S_\Theta = -\int_{\mathcal{X}_q} d\mu(x)\, \epsilon(x)\, \nabla^{(\mathrm{rel})}_\mu J^\mu_{\Theta;a}(x). \tag{9.220}$$

If the transformation is a symmetry for constant ϵ, the localized-parameter argument implies the on-shell conservation law $\nabla^{(\mathrm{rel})}_\mu J^\mu_{\Theta;a} = 0$. For a gauged symmetry, $\nabla^{(\mathrm{rel})}_\mu$ is promoted to D_μ, giving $D_\mu J^\mu_{\Theta;a} = 0$.

(v) Expectation-level statement. Taking the expectation value in the Quantum Holo-State and using linearity of $\nabla^{(\mathrm{rel})}$, together with the BRST-clean projection $\mathsf{P}_{\mathrm{phys}}$, gives $\nabla^{(\mathrm{rel})}_\mu \bar{J}^\mu_{\Theta;a} = 0$, and, in the gauged case, $D_\mu \bar{J}^\mu_{\Theta;a} = 0$. □

Physical Interpretation The Tantrakyma Noether current components $\widehat{\mathcal{J}}^\mu_{\Theta;a}(x)$ (operator level) and their scalar trace $J^\mu_{\Theta;a}(x)$ (cf. Definition 9.354) account for the flow of *excitation content*—the matter-like internal quantum numbers carried by kymons—across the emergent spacetime. For each continuous internal symmetry generated by a Hermitian T_a, on-shell invariance of the action implies the covariant conservation law for the expectation-level, BRST-clean current $\bar{J}^\mu_{\Theta;a}(x)$ (eq. (9.210)): $\nabla^{(\mathrm{rel})}_\mu \bar{J}^\mu_{\Theta;a} = 0$ (Proposition 9.355). Unlike the conserved diagnostics in the Gaiamiti (geometric) sector—which characterize the chronogeometric background itself—$\bar{J}^\mu_{\Theta;a}$ tracks the transport of *matter-sector* charges, flavors, and symmetry fluxes borne by kymons.

The conservation law is *relational*: it is defined with the Levi–Civita connection of the unified chronogeometric tensor $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (Definitions 5.26 and 6.24), i.e. in the same chronogeometric arena that governs propagation in the Euler–Lagrange equation (eq. (9.199)). Operationally, the expectation current provides a state-side diagnostic that complements the law-side EL operator: its spectral weight co-locates with the poles of the typed retarded kernels $\Delta^{[\chi]}_{\Theta\Theta}$, revealing NG channels (massless; quadrant QI) and the *gapped* radial (Higgs) channel, whose *Location* band (IR/meso) is set by the gap and certified within the Parasymmetric T–L–O workflow (eqs. (9.126) to (9.128) and Definitions 9.297 and 9.299).

At higher resolution, typed and triadic projections $\bar{J}^{\mu\ (\alpha,[\chi])}_{\Theta;a}$ (eq. (9.211)) separate the current into channels $\alpha \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$ and type slices $[\chi] \in \{\mathrm{CK}, \mathrm{T}_\pm, \mathrm{L}\}$ (Definition 9.295). Their measured plateaus and healing behavior align with the SynTh meters and the quadrant predicate, supplying *Location* (IR/meso/UV window) and *Onset* thresholds recorded in the typed T–L–O stamps. In short, $\bar{J}^\mu_{\Theta;a}$ is not only a formal consequence of symmetry; it is a diagnostic realized reference configuration-certified probe of how conserved excitation content is conveyed—coherently and locally—through the chronogeometric medium across the emergent manifold $\mathcal{X}_q$.

9.18.11 The Noether Stress–Energy Tensor of the Tantrakyma Field

Continuous spacetime symmetries of the Tantrakyma action $S_\Theta[\hat{\boldsymbol{\Theta}}]$ (cf. Definition 9.310) imply, via Noether's theorem, a conserved stress–energy object that encodes energy–momentum

flow carried by kymon excitations. In the IQR formalism we distinguish the *operator-valued* (pre-trace) density from its scalar (trace) used in the action, and keep the construction *diagnostic realized reference configuration-ready* (band-limited, BRST-clean). Throughout, the potential is built from the *co-metric* g_C (never g_C^{-1}).

Definition 9.356 (Hilbert/Noether stress–energy (operator-valued, pre-trace))**.** Let the operator-valued Lagrangian density be

$$\mathscr{L}^{\mathrm{op}}_{\Theta}(x) := \bar{\Sigma}^{\alpha\beta}(x)\, g_C\Big(\nabla_\alpha \hat{\boldsymbol{\Theta}}(x),\, \nabla_\beta \hat{\boldsymbol{\Theta}}(x)\Big) \;+\; \mathcal{V}_\Theta\Big(\hat{\boldsymbol{\Theta}}(x)\Big) \;\in\; \mathrm{End}(\mathcal{H}_{\mathrm{em}}), \tag{9.221}$$

so that the scalar density in the action is $\mathcal{L}_\Theta(x) = \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\{\mathscr{L}^{\mathrm{op}}_\Theta(x)\}$ (cf. Definition 9.301). Define $\nabla^\mu := \bar{\Sigma}^{\mu\lambda}\nabla_\lambda$. The *operator-valued* (symmetric) stress–energy tensor of the Tantrakyma sector is

$$\boxed{\hat{T}^{\mu\nu}_{(\Theta)}(x) := \tfrac{1}{2}\Big[g_C\Big(\nabla^\mu \hat{\boldsymbol{\Theta}},\, \nabla^\nu \hat{\boldsymbol{\Theta}}\Big) + g_C\Big(\nabla^\nu \hat{\boldsymbol{\Theta}},\, \nabla^\mu \hat{\boldsymbol{\Theta}}\Big)\Big] \;-\; \bar{\Sigma}^{\mu\nu}(x)\, \mathscr{L}^{\mathrm{op}}_\Theta(x).} \tag{9.222}$$

This form is Hermitian provided g_C is Hermitian and ∇_μ is $*$-compatible (cf. Definition 9.323).

Variational characterization (unified-tensor response). Varying the action with respect to $\bar{\Sigma}_{\mu\nu}$ gives

$$\boxed{\delta S_\Theta = \frac{1}{2}\int_{\mathcal{X}_q} d^d x\, \sqrt{-\det \bar{\Sigma}(x)}\; \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\Big\{\hat{T}^{\mu\nu}_{(\Theta)}(x)\Big\}\; \delta\bar{\Sigma}_{\mu\nu}(x),} \tag{9.223}$$

hence the familiar scalar (trace) object is

$$T^{\mu\nu}_{(\Theta)}(x) := \frac{2}{\sqrt{-\det \bar{\Sigma}(x)}}\,\frac{\delta S_\Theta}{\delta \bar{\Sigma}_{\mu\nu}(x)} = \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\Big\{\hat{T}^{\mu\nu}_{(\Theta)}(x)\Big\}. \tag{9.224}$$

Diagnostic realized reference configuration (band-limited, BRST-clean) and expectation tensor. On a certified plateau $\mathfrak{P}$ with tags $(\lambda_{\mathrm{UV}}, \ell)$ and physical projector $\mathsf{P}_{\mathrm{phys}} : \mathcal{H} \to H^0(\hat{Q}_{\mathrm{BRST}})$ (cf. Definitions 9.306 and 9.307), use band-limited fields/derivatives in (9.222). The observable (expectation-level) tensor is

$$\boxed{\bar{T}^{\mu\nu}_{(\Theta)}(x) := \Big\langle \overset{\circ}{\boldsymbol{\Psi}} \Big|\, \mathsf{P}_{\mathrm{phys}}\, \mathrm{Tr}_{\mathcal{H}_{\mathrm{em}}}\Big\{\hat{T}^{\mu\nu}_{(\Theta)}(x)\Big\} \Big|\, \overset{\circ}{\boldsymbol{\Psi}}\Big\rangle.} \tag{9.225}$$

Conservation (on-shell). If $\hat{\boldsymbol{\Theta}}$ satisfies the Euler–Lagrange equation (9.199) and the connections are compatible (cf. Definition 9.323), then the relationally covariant conservation law holds:

$$\boxed{\nabla^{(\mathrm{rel})}_\mu\, \bar{T}^{\mu\nu}_{(\Theta)}(x) = 0.} \tag{9.226}$$

In a gauged realization, energy–momentum exchanges between matter and gauge sectors imply conservation of the *total* (matter+gauge) stress–energy; the matter piece alone is conserved whenever the gauge Lorentz-force exchange term vanishes on-shell (e.g. $\mathcal{F}_{\mu\nu} = 0$).

Typed & triadic resolution (parasymmetric diagnostics). With typed projectors $\Pi^{[\chi]}_{\Theta\Theta}$ and channel projectors Π_α that commute (cf. Definition 9.295), define

$$\hat{T}^{\mu\nu\,(\alpha,[\chi])}_{(\Theta)} := \Pi_\alpha\,\Pi^{[\chi]}_{\Theta\Theta}\,\hat{T}^{\mu\nu}_{(\Theta)}\,, \qquad \bar{T}^{\mu\nu\,(\alpha,[\chi])}_{(\Theta)} := \Big\langle \mathring{\boldsymbol{\Psi}} \,\Big|\, \mathsf{P}_{\text{phys}}\,\mathrm{Tr}_{\mathcal{H}_{\text{em}}}\{\hat{T}^{\mu\nu\,(\alpha,[\chi])}_{(\Theta)}\} \,\Big|\, \mathring{\boldsymbol{\Psi}} \Big\rangle. \tag{9.227}$$

Their spectral content provides *Location* diagnostics coherent with the SynTh meters and quadrant semantics (cf. eqs. (9.126) to (9.128) and Definitions 9.297 and 9.299): NG channels (QI) exhibit low-frequency spectral weight aligned with the $k^2 = 0$ pole; the Higgs (radial) channel shows a gapped peak at $k^2 = M_{\text{H}}^2$ (cf. eq. (9.194)), with *Onset* thresholds stamped per diagnostic realized reference configuration tags. Equations (9.222)–(9.226) complete the Noether/diffeomorphism response for the Tantrakyma sector, furnishing both the variational definition used in the field equations and the operational typed/triadic observables used by the Parasymmetric T–L–O workflow—all expressed purely with the co-metric pairing $g_C(\cdot,\cdot)$.

Interpretation and Physical Role The Tantrakyma stress–energy tensor encodes the energetic and dynamical content of the matter sector at the operator level (cf. Definition 9.356). In its pre-trace form $\hat{T}^{\mu\nu}_{(\Theta)}$ (cf. eq. (9.222)) it tracks how kymon excitations carry energy and momentum through the emergent configuration space $\mathcal{X}_q$. After taking the fiber trace and the diagnostic realized reference configuration expectation, one obtains the observable tensor $\bar{T}^{\mu\nu}_{(\Theta)}$ (cf. eq. (9.225)), which couples to geometry (see the Gaiamiti sector, Section 9.17.9).

- **Propagation (kinetic flow).** The symmetrized pairings

  $$\tfrac{1}{2}\Big[g_C(\nabla^\mu\hat{\boldsymbol{\Theta}},\nabla^\nu\hat{\boldsymbol{\Theta}}) + g_C(\nabla^\nu\hat{\boldsymbol{\Theta}},\nabla^\mu\hat{\boldsymbol{\Theta}})\Big]$$

  capture directional transport and kinetic energy of kymons along the chronogeometric directions set by $\bar{\Sigma}^{\mu\nu}$ (cf. Definitions 5.26 and 9.323).

- **Internal excitation content.** The subtraction in $\hat{T}^{\mu\nu}_{(\Theta)}$ is $-\,\bar{\Sigma}^{\mu\nu}\,\mathscr{L}^{\text{op}}_{\Theta}$, i.e.

  $$-\,\bar{\Sigma}^{\mu\nu}\Big[\bar{\Sigma}^{\alpha\beta}\,g_C(\nabla_\alpha\hat{\boldsymbol{\Theta}},\nabla_\beta\hat{\boldsymbol{\Theta}}) + \mathcal{V}_\Theta(\hat{\boldsymbol{\Theta}})\Big]\,,$$

  which removes the kinetic trace and adds the potential contribution exactly as in eq. (9.222) (cf. Definitions 9.301 and 9.335).

Observable role and conservation. Taking the BRST-clean expectation on the diagnostic realized reference configuration yields the classical source field $\bar{T}^{\mu\nu}_{(\Theta)}$ (cf. eq. (9.225)). When the field is on-shell, this tensor is covariantly conserved with the relational connection, $\nabla^{(\text{rel})}_\mu \bar{T}^{\mu\nu}_{(\Theta)} = 0$ (cf. eq. (9.226)), expressing energy–momentum conservation of the matter sector within the chronogeometric arena determined by $\bar{\Sigma}_{\mu\nu}$.

Parasymmetric T–L–O diagnostics. Typed/triadic resolutions $\bar{T}^{\mu\nu\,(\alpha,[\chi])}_{(\Theta)}$ (cf. eq. (9.227)) act as *Location* probes consistent with the SynTh meters $(v^{(L)}, v^{(S)}, v^{(LS)})$ and quadrant semantics (cf. eqs. (9.126) to (9.128) and Definitions 9.297 and 9.299). NG channels (QI) exhibit low-frequency spectral weight aligned with the $k^2 = 0$ pole of the typed kernels, whereas the radial (Higgs) channel shows a gapped peak at $k^2 = M_{\text{H}}^2$ (cf. eq. (9.194)), with *Onset* thresholds stamped per diagnostic realized reference configuration tags.

9.18.12 The Source of Emergent Curvature: A Unified View

A central question in any theory of gravity is the origin of spacetime curvature. In the IQR framework, where geometry is emergent, curvature arises from the relational dynamics of the foundational quantum fields. Both the geometric Gaiamiti field and the matter-like Tantrakyma field contribute to the total stress–energy content, and this *unified* content sources curvature of the emergent chronogeometric manifold.

Definition 9.357 (Total Relational Action)**.** The complete dynamics are governed by

$$S_{\text{total}}\big[\hat{\mathbf{\Phi}},\,\hat{\mathbf{\Theta}}\big] \;:=\; S_{\Phi}\big[\hat{\mathbf{\Phi}}\big] \;+\; S_{\Theta}\big[\hat{\mathbf{\Theta}}\big] \;+\; S_{\text{int}}\big[\hat{\mathbf{\Phi}},\,\hat{\mathbf{\Theta}}\big]\,, \tag{9.228}$$

where:

- $S_{\Phi}[\hat{\mathbf{\Phi}}]$ is the Gaiamiti (geometry) action (cf. Section 6.5);
- $S_{\Theta}[\hat{\mathbf{\Theta}}]$ is the Tantrakyma (matter) action (cf. Definition 9.310);
- $S_{\text{int}}[\hat{\mathbf{\Phi}},\hat{\mathbf{\Theta}}]$ encodes geometry–matter coupling (cf. Section 10.19.4).

All integrals use the relational volume element $d\mu(x) := d^d x\,\sqrt{-\det \bar{\Sigma}(x)}$ (cf. Definition 6.33 and the measure note following Definition 9.310).

Definition 9.358 (Total stress–energy and expectation source)**.** Let $\hat{T}^{\mu\nu}_{(\Phi)}$ and $\hat{T}^{\mu\nu}_{(\Theta)}$ be the *operator-valued* (pre-trace) stress–energy tensors for the Gaiamiti and Tantrakyma sectors (cf. Section 9.17.9 and Definition 9.356), and let $\hat{T}^{\mu\nu}_{(\text{int})}$ be the interaction contribution obtained by varying S_{int} with respect to $\bar{\Sigma}_{\mu\nu}$. Their scalar (trace) forms are $T^{\mu\nu}_{(\bullet)} = \mathrm{Tr}\{\hat{T}^{\mu\nu}_{(\bullet)}\}$ (cf. eq. (9.224)). Define the *total* operator and scalar tensors by

$$\hat{T}^{\mu\nu}_{\text{total}} := \hat{T}^{\mu\nu}_{(\Phi)} + \hat{T}^{\mu\nu}_{(\Theta)} + \hat{T}^{\mu\nu}_{(\text{int})}, \qquad T^{\mu\nu}_{\text{total}} := \mathrm{Tr}\Big\{\hat{T}^{\mu\nu}_{\text{total}}\Big\}\,. \tag{9.229}$$

On a certified plateau $\mathfrak{P}$ with tags $(\lambda_{\text{UV}}, \ell)$ and projection $\mathsf{P}_{\text{phys}} : \mathcal{H} \to H^0(\hat{Q}_{\text{BRST}})$, the expectation-level source of emergent curvature is

$$\boxed{\bar{T}^{\mu\nu}_{\text{total}}(x) \;:=\; \Big\langle \overset{\circ}{\mathbf{\Psi}} \Big|\, \mathsf{P}_{\text{phys}}\, \mathrm{Tr}\Big\{\hat{T}^{\mu\nu}_{\text{total}}(x)\Big\} \Big|\, \overset{\circ}{\mathbf{\Psi}} \Big\rangle.} \tag{9.230}$$

Unified variational identity and conservation. The unified (Hilbert) variation with respect to $\bar{\Sigma}_{\mu\nu}$ yields

$$\delta S_{\text{total}} = \frac{1}{2}\int d\mu(x)\;\; T^{\mu\nu}_{\text{total}}(x)\,\delta\bar{\Sigma}_{\mu\nu}(x), \tag{9.231}$$

so $T^{\mu\nu}_{\text{total}}$ is the *unique* source for the chronogeometric response. On-shell (EL equations satisfied in both sectors; cf. eq. (9.199) and the Gaiamiti EL), the relationally covariant conservation law holds:

$$\boxed{\nabla^{(\text{rel})}_{\mu}\,\bar{T}^{\mu\nu}_{\text{total}}(x) = 0.} \tag{9.232}$$

When there is no geometry–matter exchange ($\hat{T}^{\mu\nu}_{(\text{int})} = 0$), the separate pieces are conserved; otherwise only the *sum* is conserved. Typed/triadic resolutions of $\bar{T}^{\mu\nu}_{\text{total}}$ provide *Location*

diagnostics consistent with the SynTh meters and quadrant semantics (cf. eqs. (9.126) to (9.128) and Definitions 9.297 and 9.299).

Notes. (i) In the Tantrakyma sector all potential invariants are built from the co-metric $g_C : T^*\times T^* \to \mathrm{End}(\mathcal{H}_{\mathrm{em}})$ (not g_C^{-1}), ensuring that both $\hat{T}^{\mu\nu}_{(\Theta)}$ and $T^{\mu\nu}_{(\Theta)}$ are computed with the correct cotangent pairing. (ii) Diagnostic realized reference configuration-limited, BRST-clean expectation values guarantee gauge-parameter independence (Nielsen flatness) of $\bar{T}^{\mu\nu}_{\mathrm{total}}$.

Evaluation of the Total Expectation-Valued Stress–Energy Tensor

We evaluate the total source in three steps, keeping the operator/scalar distinction explicit and enforcing diagnostic realized reference configuration (band-limited, BRST-clean) discipline.

1. Additivity and expectation linearity (including interaction). By linearity of the expectation and the variational definition of stress–energy (cf. eq. (9.224)), the *total* scalar tensor is the sum of sectoral and interaction pieces:

$$\boxed{\bar{T}^{\mu\nu}_{\mathrm{total}}(x) = \bar{T}^{\mu\nu}_{(\Phi)}(x) + \bar{T}^{\mu\nu}_{(\Theta)}(x) + \bar{T}^{\mu\nu}_{(\mathrm{int})}(x),} \tag{9.233}$$

with

$$\bar{T}^{\mu\nu}_{(\bullet)}(x) := \Big\langle \mathring{\boldsymbol{\Psi}} \,\Big|\, \mathsf{P}_{\mathrm{phys}}\, T^{\mu\nu}_{(\bullet)}(x) \,\Big|\, \mathring{\boldsymbol{\Psi}} \Big\rangle, \qquad (\bullet) \in \{\Phi, \Theta, \mathrm{int}\}, \tag{9.234}$$

where $\mathsf{P}_{\mathrm{phys}} : \mathcal{H} \to H^0(\hat{Q}_{\mathrm{BRST}})$ is the physical projector (diagnostic realized reference configuration semantics; cf. Definitions 9.306 and 9.307). Here $T^{\mu\nu}_{(\bullet)} = \mathrm{Tr}\{\hat{T}^{\mu\nu}_{(\bullet)}\}$ is the scalar (fiber-traced) tensor associated with the operator-valued $\hat{T}^{\mu\nu}_{(\bullet)}$ (cf. Section 9.17.9 and Definition 9.356).

2. Operator forms and diagnostic realized reference configuration expectations. Raise indices with $\nabla^\mu := \bar{\Sigma}^{\mu\lambda}\nabla_\lambda$. Let the operator-valued Lagrangians be

$$\mathscr{L}^{\mathrm{op}}_{\Phi} = \bar{\Sigma}^{\alpha\beta} g_T\Big(\nabla_\alpha\hat{\boldsymbol{\Phi}},\, \nabla_\beta\hat{\boldsymbol{\Phi}}\Big) + \mathcal{V}_\Phi(\hat{\boldsymbol{\Phi}}), \quad \mathscr{L}^{\mathrm{op}}_{\Theta} = \bar{\Sigma}^{\alpha\beta} g_C\Big(\nabla_\alpha\hat{\boldsymbol{\Theta}},\, \nabla_\beta\hat{\boldsymbol{\Theta}}\Big) + \mathcal{V}_\Theta(\hat{\boldsymbol{\Theta}}),$$

whose fiber traces enter the actions (cf. Definition 9.301).

- **Gaiamiti (geometry) sector** (cf. Section 9.17.9): with the tangent metric g_T and operator field $\hat{\boldsymbol{\Phi}}$,

$$\begin{aligned}\hat{T}^{\mu\nu}_{(\Phi)}(x) := {}& \tfrac{1}{2}\Big[g_T\Big(\nabla^\mu\hat{\boldsymbol{\Phi}},\, \nabla^\nu\hat{\boldsymbol{\Phi}}\Big) + g_T\Big(\nabla^\nu\hat{\boldsymbol{\Phi}},\, \nabla^\mu\hat{\boldsymbol{\Phi}}\Big)\Big] \\ & - \tfrac{1}{2}\,\bar{\Sigma}^{\mu\nu}\bar{\Sigma}^{\alpha\beta}\, g_T\Big(\nabla_\alpha\hat{\boldsymbol{\Phi}},\, \nabla_\beta\hat{\boldsymbol{\Phi}}\Big) - \bar{\Sigma}^{\mu\nu}\,\mathcal{V}_\Phi(\hat{\boldsymbol{\Phi}}).\end{aligned} \tag{9.235}$$

- **Tantrakyma (matter) sector** (cf. Definition 9.356): with the cotangent co-metric g_C and operator field $\hat{\boldsymbol{\Theta}}$,

$$\begin{aligned}\hat{T}^{\mu\nu}_{(\Theta)}(x) := {}& \tfrac{1}{2}\Big[g_C\Big(\nabla^\mu\hat{\boldsymbol{\Theta}},\, \nabla^\nu\hat{\boldsymbol{\Theta}}\Big) + g_C\Big(\nabla^\nu\hat{\boldsymbol{\Theta}},\, \nabla^\mu\hat{\boldsymbol{\Theta}}\Big)\Big] \\ & - \tfrac{1}{2}\,\bar{\Sigma}^{\mu\nu}\bar{\Sigma}^{\alpha\beta}\, g_C\Big(\nabla_\alpha\hat{\boldsymbol{\Theta}},\, \nabla_\beta\hat{\boldsymbol{\Theta}}\Big) - \bar{\Sigma}^{\mu\nu}\,\mathcal{V}_\Theta(\hat{\boldsymbol{\Theta}}).\end{aligned} \tag{9.236}$$

- **Interaction sector**:

$$\hat{T}^{\mu\nu}_{(\mathrm{int})}(x) \text{ is defined variationally by } \quad \delta S_{\mathrm{int}} = \frac{1}{2}\int d\mu\, \mathrm{Tr}\Big\{\hat{T}^{\mu\nu}_{(\mathrm{int})}(x)\Big\}\, \delta\bar{\Sigma}_{\mu\nu}(x), \tag{9.237}$$

so $T^{\mu\nu}_{(\mathrm{int})} = \mathrm{Tr}\{\hat{T}^{\mu\nu}_{(\mathrm{int})}\}$.

Taking the fiber trace and the diagnostic realized reference configuration expectation (cf. eqs. (9.224) and (9.225)) yields

$$\boxed{\bar{T}^{\mu\nu}_{\mathrm{total}} = \Big\langle \mathsf{P}_{\mathrm{phys}}\, \mathrm{Tr}\Big\{\hat{T}^{\mu\nu}_{(\Phi)} + \hat{T}^{\mu\nu}_{(\Theta)} + \hat{T}^{\mu\nu}_{(\mathrm{int})}\Big\}\Big\rangle_{\mathfrak{P}},} \tag{9.238}$$

with $\langle\cdot\rangle_{\mathfrak{P}} := \langle\mathring{\mathbf{\Psi}}|\,(\cdot)\,|\mathring{\mathbf{\Psi}}\rangle$ at fixed tags $(\lambda_{\mathrm{UV}}, \ell)$. Equation (9.238) is the diagnostic realized reference configuration form of eq. (9.230).

3. Parasymmetric T–L–O resolution (typed, triadic; optional but recommended). For diagnostics and cross-sector consistency, resolve $\bar{T}^{\mu\nu}_{\mathrm{total}}$ by type and channel using projectors $\Pi^{[\chi]}$ and Π_α that commute (cf. Definitions 5.54 and 9.295):

$$\bar{T}^{\mu\nu}_{\mathrm{total}}{}^{(\alpha,[\chi])} := \Big\langle \mathsf{P}_{\mathrm{phys}}\, \mathrm{Tr}\{\Pi_\alpha\, \Pi^{[\chi]}\, (\hat{T}^{\mu\nu}_{(\Phi)} + \hat{T}^{\mu\nu}_{(\Theta)} + \hat{T}^{\mu\nu}_{(\mathrm{int})})\}\Big\rangle_{\mathfrak{P}}. \tag{9.239}$$

Their spectral content co-locates with the typed kernel poles used by the SynTh meters $v^{(L)}, v^{(S)}, v^{(LS)}$ (cf. eqs. (9.126) to (9.128)), enabling a *typed quadrant* decision via $\mathrm{Quad}^{[\cdot]}_{[\chi]}$ and T–L–O stamping (cf. Definitions 9.297 and 9.299). NG channels (QI) exhibit low-frequency weight aligned with $k^2 = 0$, while a Higgs (radial) entry shows a single gapped peak at $k^2 = M_{\mathrm{H}}^2$ (cf. eq. (9.194)), with Location/Onset set by diagnostic realized reference configuration tags.

Conservation (unified, on-shell). When both EL systems hold and connections are compatible (cf. Definitions 9.305 and 9.323 and the geometry-side EL), the total source is covariantly conserved:

$$\nabla^{(\mathrm{rel})}_{\mu}\, \bar{T}^{\mu\nu}_{\mathrm{total}} = 0, \tag{9.240}$$

with separate sectoral conservation when the interaction exchange vanishes and, in general, conservation in the sum.

Interpretation.

- *Kinetic/transport content.* The first (symmetrized) terms in $\hat{T}^{\mu\nu}_{(\Phi)}$ and $\hat{T}^{\mu\nu}_{(\Theta)}$ are the operator-valued metric/co-metric pairings

$$\tfrac{1}{2}\Big[g_T(\nabla^\mu\hat{\mathbf{\Phi}}, \nabla^\nu\hat{\mathbf{\Phi}}) + g_T(\nabla^\nu\hat{\mathbf{\Phi}}, \nabla^\mu\hat{\mathbf{\Phi}})\Big], \qquad \tfrac{1}{2}\Big[g_C(\nabla^\mu\hat{\mathbf{\Theta}}, \nabla^\nu\hat{\mathbf{\Theta}}) + g_C(\nabla^\nu\hat{\mathbf{\Theta}}, \nabla^\mu\hat{\mathbf{\Theta}})\Big],$$

capturing directional transport and kinetic flow of the Apeiron (geometry) and kymon (matter) fields along the chronogeometric directions selected by $\bar{\Sigma}^{\mu\nu}$ (cf. Section 9.17.9 and Definitions 9.323 and 9.356).

- *Potential/interaction content.* The subtraction structure in $\hat{T}^{\mu\nu}_{(\bullet)}$ (with $(\bullet) \in \{\Phi, \Theta\}$) consists of the kinetic trace term $-\frac{1}{2}\,\bar{\Sigma}^{\mu\nu}\bar{\Sigma}^{\alpha\beta}\,(\cdots)_{\alpha\beta}$ and the potential piece $-\,\bar{\Sigma}^{\mu\nu}\,\mathcal{V}_{\bullet}$, exactly as in eq. (9.222) and its geometry-side analogue. After taking the fiber trace and the diagnostic realized reference configuration expectation, this yields a conserved energy–momentum flux consistent with the Hilbert variational identity $\delta S = \frac{1}{2}\int d\mu\ \mathrm{Tr}\{\hat{T}^{\mu\nu}\}\delta\bar{\Sigma}_{\mu\nu}$ (cf. eqs. (9.223), (9.224) and (9.238)). Interaction contributions are included via $\hat{T}^{\mu\nu}_{(\mathrm{int})}$ by variation of S_{int}.

- *Back-reaction mediator.* The background tensor $\bar{\Sigma}^{\mu\nu}(x)$ raises/lowers indices, defines the relational covariant derivative, and mediates the back–reaction of both sectors on curvature. On-shell one has $\nabla^{(\mathrm{rel})}_{\mu}\bar{T}^{\mu\nu}_{\mathrm{total}} = 0$ (cf. eq. (9.240)), ensuring consistent sourcing of emergent chronogeometry.

This furnishes a self-consistent matter–geometry interplay in the IQR framework, echoing Wheeler's dictum [146, 239] in diagnostic realized reference configuration-ready form.

> *Spacetime tells matter how to move* (paraphrasing [146]): the emergent $\bar{\Sigma}_{\mu\nu}$, constructed from the Gaiamiti field $\hat{\mathbf{\Phi}}$, governs the covariant propagation of $\hat{\mathbf{\Theta}}$ via eq. (9.199).
>
> *Matter tells spacetime how to curve* (paraphrasing [146]): the total source $\bar{T}^{\mu\nu}_{\mathrm{total}}$ (including Tantrakyma excitations and interaction exchange) drives the evolution of $\hat{\mathbf{\Phi}}$, thereby reshaping $\bar{\Sigma}_{\mu\nu}$.

Parasymmetric T–L–O perspective. Typed/triadic resolutions of $\bar{T}^{\mu\nu}_{\mathrm{total}}$ provide Location diagnostics that co-locate with poles of the typed kernels and agree with SynTh meters and quadrant calls (NG vs. gapped channels), yielding consistent T–L–O stamps across geometry and matter (cf. eqs. (9.126) to (9.128) and (9.239) and Definitions 9.297 and 9.299).

Chapter 10

The Weave of Geometry and Matter

10.1 The Inter-Modal Correlators of Fields

This chapter revisits the *Unified Inter-Modal Correlator* as the kinematic backbone from which both geometry (Gaiamiti) and matter/temporality (Tantrakyma) draw their operational meaning, and equips that backbone with the covariant dynamics previously introduced, so that we may linearize the geometric sector, isolate graviton channels, and state the chronogeometric Einstein equations $\bar{G}_{\mu\nu}[\bar{\Sigma}]$ in their natural $\bar{\Sigma}$-geometry (Section 11.7.3). Concretely, the unified object is the matrix of two-point functions

$$\hat{\mathbf{G}}_{\Lambda}(x,x') := \begin{pmatrix} \hat{G}_{\mathbf{\Phi\Phi}} & \hat{G}_{\mathbf{\Phi\Theta}} \\ \hat{G}_{\mathbf{\Theta\Phi}} & \hat{G}_{\mathbf{\Theta\Theta}} \end{pmatrix}(x,x').$$

Its entries are obtained by specializing the Fourfold correlator $\hat{G}_{\Lambda}^{\alpha\beta}$ (Definition 6.4).

From Fourfold Kinematics to the Unified Two-Block. As formalized in Definition 4.9 and Section 4.1.4, the Fourfold correlator is the gauge-covariant, operator-valued kernel that transports and couples the four dialectic modes $\{0,1,p,s\}$, with a path-ordered parallel transporter $\mathcal{U}(x \leftarrow x';\gamma)$ effecting frame alignment across loci in $\mathcal{X}_q$. In tensor form,

$$\hat{G}_{\Lambda}^{\alpha\beta}(x,x') \;=\; \hat{\mathcal{F}}_{\alpha}^{\dagger}(x)\,\mathcal{U}(x \leftarrow x';\gamma)\,\hat{\mathcal{F}}_{\beta}(x'),$$

and the unified matrix above is precisely its organization into the $(\mathbf{\Phi},\mathbf{\Theta})$ sectors (Section 4.1.5). The fibre inner products are *generated*—not postulated—by Holo-State expectation in the coincident limit, yielding well-posed Hilbert fibres and a Hilbert bundle of dialectic modes (Sections 4.1.3 and 4.6.1); the modal basis $\{|0\rangle_x, |1\rangle_x, |p\rangle_x, |s\rangle_x\}$ realizes parasymmetric/holosymmetric directions as independent rays (not superpositions of $|0\rangle, |1\rangle$) (Section 4.1.5).

Relational Covariance and Operator Dynamics. Given the expectation-level chronogeometric tensor $\bar{\Sigma}_{\mu\nu} := \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$, there exists a unique torsion-free, metric-compatible connection $\nabla^{(\mathrm{rel})}$ with Christoffel symbols $\Gamma^{\lambda}_{\mu\nu}[\bar{\Sigma}]$ (Definition 6.7). Operator fields are sections of bundles twisted both by $\nabla^{(\mathrm{rel})}$ and by an internal operator connection $\mathcal{A}_\mu$. For the Gaiamiti field,

$$\nabla_\mu \hat{\mathbf{\Phi}} \;=\; \nabla^{(\mathrm{rel})}_{\mu} \hat{\mathbf{\Phi}} \;+\; D_\mu \hat{\mathbf{\Phi}}, \qquad (\nabla_\mu \hat{\mathbf{\Phi}})^{\lambda} = \partial_\mu \hat{\mathbf{\Phi}}^{\lambda} + \Gamma^{\lambda}_{\mu\nu}[\bar{\Sigma}]\hat{\mathbf{\Phi}}^{\nu} + i[\mathcal{A}_\mu, \hat{\mathbf{\Phi}}^{\lambda}], \tag{10.1}$$

(Definition 6.184); this enters the action $S_\Phi = \int \sqrt{-\det \bar{\Sigma}}\, \mathcal{L}_\Phi$ and Euler–Lagrange equations, with Noether stress–energy $\hat{T}^{\mu\nu}_{(\Phi)}$ (Definition 6.174 and Section 9.17.9). Analogously, for Tantrakyma,

$$\nabla_\mu \hat{\boldsymbol{\Theta}} \;=\; \nabla^{(\mathrm{rel})}_\mu \hat{\boldsymbol{\Theta}} \;+\; D_\mu \hat{\boldsymbol{\Theta}}, \tag{10.2}$$

(Definition 9.323).

Σ–Realization, Acceptance, and Quadrants. Building on this, we evaluate all kinematics and dynamics on a certified plateau $\mathfrak{P}$ and then project to the physical cohomology via the Σ–*realization map* (Definition 9.80)

$$\boxed{\bar{\Sigma}_{\mu\nu}(x) = \Big\langle \mathsf{P}_{\mathrm{phys}} \lim_{(\lambda,\ell)\in\mathfrak{P}} \mathfrak{B}_{\lambda,\ell}![\hat{\Sigma}_{\mu\nu}(x)]\; \mathsf{P}_{\mathrm{phys}} \Big\rangle},$$

after which all Chapter 7 constructions are *read as functions of* $\bar{\Sigma}$ with the BRST sector and proofs unchanged (Remark 9.81). Thus $\bar{\Sigma}^{\mu\nu}$ (the inverse of $\bar{\Sigma}_{\mu\nu}$, cf. eq. (6.62)) raises indices and fixes the chronogeometric cones that support linear response; in particular, the retarded kernels built from $\hat{\mathbf{G}}_\Lambda$ are Σ-*causal* (support contained in the $\bar{\Sigma}$-cones) by Remark 9.83, and mode normalization with respect to $\bar{\Sigma}$ is equivalent—up to bounded reweighting—to the triad metrics (Remark 9.84). The *holosymmetric visibilities* $\Lambda_{\mathrm{law}}, \Lambda_{\mathrm{state}}$ from Chapter 7 (Definition 7.35) are the witnesses we continue to use here, but only on *accepted* scale bands certified by Chapter 8. Concretely, acceptance is enforced on the (u, v)-plane by the mask $\mathfrak{M}_{\mathrm{mask}}$ with guards C1–C5 (eq. (7.626)), yielding an IR window $\mathcal{I}_{\mathrm{IR}}$ on which the smoothed slope $\psi(v) = \mathsf{LP}_\tau(\Lambda'_{\mathrm{law}})(v)$ enjoys Lipschitz, total-variation, and positivity stability (Lemma 7.386 and Proposition 7.383). Within $\mathcal{I}_{\mathrm{IR}}$, the *quadrant decision* is then made canonically using $\{\Lambda_{\mathrm{law}}, \Lambda_{\mathrm{state}}\}$ (Definition 7.685), providing the QI–QIV classification that will guide graviton-channel isolation and the formulation of $\bar{G}_{\mu\nu}[\bar{\Sigma}]$ in Section 11.7.3.

Pre-Physics → Physics and Parasymmetric Overlay. Before realization, in the *pre-physical band* ($\ell_{\mathrm{PQS}} \le \ell < \ell_P$) Chapter 10 prescribes *operator-level* Σ-diagnostics only: one forms quadratic probes $\hat{Q}_x[v] = v^\mu v^\nu \hat{\Sigma}_{\mu\nu}(x)$ and expectation surrogates $\bar{Q}_x[v]$ (Definitions 9.20 and 9.22), aggregates contrasts via the RSS rule (Definition 9.47), and assigns a pre-physical quadrant (Definition 9.49). *Physical* poles/residues and visibilities are read *only after* a certified plateau and BRST projection to $H^0(\hat{Q}_{\mathrm{BRST}})$ ($\mathfrak{P}$ and $\mathsf{P}_{\mathrm{phys}}$ in Definition 9.80), which is the precise pre-physics → physics transition used throughout this chapter. The *parasymmetric overlay* of Chapter 9 remains intact and confined to H^0: parasymmetric/holosymmetric meters operate on the realized data, the quadrant assignment is functorial and cone-preserving under the certified maps (Theorem 8.45), and the law–state exhaustion link holds on $\mathfrak{P}$. Consequently, the QIII/QIV refinement and "no fifth quadrant" property from Chapters 8–9 carry over verbatim into the present $\bar{\Sigma}$-based formulation used to organize channels and residues for $\hat{\mathbf{G}}_\Lambda$.

TT Isolation for Graviton Channels. Finally, to *linearize the geometric sector and isolate graviton channels*, we work on the realized weave configuration with the typed triad $1 =$

$\Pi^{(\mathrm{spec})} + \Pi^{(\mathrm{vac})} + \Pi^{(\mathrm{corr})}$ (Definition 10.2) and the $\bar{\Sigma}$-TT projector $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ (Definition 10.3). The York–Helmholtz/polarization bookkeeping and the graviton-projector identity (Eq. (10.28)) enforce TT purity, while the *TT leak* ε (Definition 10.32) is bounded purely by non-spectral channel content (Lemma 10.103), ensuring that the TT sector extracted from $\hat{\mathbf{G}}_\Lambda$ is well-posed for the chronogeometric Einstein equations $\bar{G}_{\mu\nu}[\bar{\Sigma}]$ in Section 11.7.3.

10.2 Realized Weave Configuration and Typed Algebra

We fix the operational "rig" in this section so that all subsequent objects (τ–kairos, graviton, Higgs, Chrono–Kymon) are defined *on this realized weave configuration*, and so they inherit BRST/Nielsen safety, cone preservation, triad compatibility, and band-covariance from the outset (cf. Definitions 6.280 and 9.10, eq. (9.2), and Section 9.5 upstream).

Definitions.

Definition 10.1 (Realized weave configuration)**.** For any sector $\mathscr{S} \in \{\Phi\Phi, \Theta\Theta, \Phi\Theta\}$ and arena $\mathfrak{P} \in \{\mathrm{bulk}, \mathrm{boundary}\}$, the *realized weave configuration* is the tuple

$$\mathcal{I}^{[\mathscr{S}]} := \Big(\mathcal{A},\ r\ ;\ P_x(\lambda_{\mathrm{UV}}),\ K_{\mathrm{IR}}^{(\ell)}\ ;\ \mathsf{P}_{\mathrm{phys}}\ ;\ \bar{\Sigma}_{\mu\nu} \Big),$$

whose entries are as follows (with explicit upstream definitions):

1. $\mathcal{A}$ = *admissible holosymmetric settings*: the background/arena choice and kinematic domain on which band operators act. The arena action and admissibility conditions used by band maps are stated in eq. (9.2).
2. $r \in \mathbb{R}_{\geq 0}$ = *IR knob*: the parameter that indexes the infrared windows $K_{\mathrm{IR}}^{(\ell)}$ and their tightening along admissible retunes; cf. the window family in Definition 6.280.
3. $P_x(\lambda_{\mathrm{UV}})$ = *band tags*: orthogonal projectors implementing the certified ultraviolet cut and band selection; see the tag definition and properties in Definition 9.10, with their arena action given by eq. (9.2).
4. $K_{\mathrm{IR}}^{(\ell)}$ = *IR windows*: smoothing/compression operators forming an admissible family that controls long-distance content and defines IR retunes; see Definition 6.280.
5. $\mathsf{P}_{\mathrm{phys}}$ = *BRST projector onto* H^0: the idempotent that projects to the physical cohomology (degree-zero BRST cohomology). The physical carrier for all readouts is
$$\mathcal{H}_{\mathrm{phys}} := \mathrm{Ran}\, \mathsf{P}_{\mathrm{phys}}.$$
6. $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ = *expectation-level chronogeometric tensor*: the background-covariant, expectation-valued chronogeometry used to define covariant TT maps and traces; construction and properties are recalled in Section 9.5.

All band objects act on $\mathcal{H}_{\mathrm{phys}}$ and satisfy the certified commutation/compatibility relations:

$$[P_x(\lambda_{\mathrm{UV}}), \mathsf{P}_{\mathrm{phys}}] = 0,$$
$$[K_{\mathrm{IR}}^{(\ell)}, \mathsf{P}_{\mathrm{phys}}] = 0,$$

and the arena action of $P_x(\lambda_{\mathrm{UV}})$ is as in eq. (9.2). Norms and stability estimates are evaluated with the admissible family $\|\cdot\|_{\mathcal{I}}$ of Definition 10.143.

Meaning. *realized weave configuration* (Definition 10.1) fixes the baseline measurement frame—our *realized weave configuration* $\mathcal{I}^{[\mathscr{S}]}$—on which all subsequent geometric/matter diagnostics are performed. Concretely, the configuration is assembled *after* Σ–realization (Definition 9.80), so every operator and kernel is evaluated on the realized chronogeometry $\bar{\Sigma}$. It packages (i) the physical sector $\mathcal{H}_{\mathrm{phys}}$, (ii) the typed triad of channels $1 = \Pi^{(\mathrm{spec})} + \Pi^{(\mathrm{vac})} + \Pi^{(\mathrm{corr})}$ for bookkeeping and mode extraction (Definition 10.2), and (iii) the $\bar{\Sigma}$-transverse–traceless projector $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ used to isolate graviton channels (Definition 10.3). With this configuration in place, the unified two-point matrix $\hat{\mathbf{G}}_\Lambda$ is read Σ-causally and all covariant dynamics use the relational Levi–Civita connection $\nabla^{(\mathrm{rel})}$ compatible with $\bar{\Sigma}$ (Definition 6.7), exactly as required later when stating $\bar{G}_{\mu\nu}[\bar{\Sigma}]$ (Section 11.7.3).

Realized weave configuration vs. Plateaux. A *plateau* $\mathfrak{P}$ is a *scale-certification* in the (u, v) plane: it is the accepted window picked out by the mask $\mathfrak{M}_{\mathrm{mask}}$ and guards C1–C5 (eq. (7.626)), within which the witnesses $\Lambda_{\mathrm{law}}, \Lambda_{\mathrm{state}}$ are stable (Lipschitz/TV/positivity bounds; Lemma 7.386 and Proposition 7.383). Plateaux certify *where* the data are reliable. By contrast, the *realized weave configuration* $\mathcal{I}^{[\mathscr{S}]}$ is the *instrument set*—projectors, connections, domains—on which we then operate. In practice: $\mathfrak{P} \Rightarrow$ apply the Σ–realization map to obtain $\bar{\Sigma}$ (Definition 9.80) $\Rightarrow$ *install* the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$. Thus, plateaux are *scale-local certifications of input*, while the realized weave configuration is a *structural, sector-conditioned apparatus* that persists for analysis (e.g., TT isolation and stress–energy evaluation) once $\bar{\Sigma}$ is fixed.

Intent and Use. The intent of Definition 10.1 is to guarantee that downstream statements—quadrant assignment (Definition 7.685), TT purity and TT-leak control (ε bounded by non-spectral content; Definition 10.32, Lemma 10.103), and ultimately the formulation of the chronogeometric Einstein equations—are executed on a *calibrated* and *BRST/Nielsen-clean* stage determined by $\bar{\Sigma}$. In short, plateaux supply certified data; the realized weave configuration supplies certified tools. Together they make the extraction of graviton channels with $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ (Definition 10.3) and the use of $\nabla^{(\mathrm{rel})}$ (Definition 6.7) mathematically well-posed for the construction of $\bar{G}_{\mu\nu}[\bar{\Sigma}]$ in Section 11.7.3.

Definition 10.2 (Triad channels)**.** Work on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ (Def. Definition 10.1). We use mutually commuting projectors $\Pi^{(a)}$ for $a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$ (triad decomposition; see also band invariance Proposition 10.10 and masks/selectors Definitions 7.665 and 10.7), with

$$\Pi^{(\mathrm{spec})} + \Pi^{(\mathrm{vac})} + \Pi^{(\mathrm{corr})} = \mathbf{1}, \qquad \Pi^{(a)}\Pi^{(b)} = \delta_{ab}\Pi^{(a)}. \tag{10.3}$$

These triads are the provenance maps used throughout ledgers/meters (Definitions 10.80 and 10.82) and typed correlator insertions (Definition 10.113).

Definition 10.3 (Background-covariant TT projector)**.** On symmetric 2-tensors $h \in \Gamma(S^2T^*\mathcal{X}_q)$, $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ is the $\bar{\Sigma}$–compatible transverse–traceless projector (a background-covariant pseudodifferential lift defined with the relational Levi–Civita connection $\nabla^{(\mathrm{rel})}$; cf. Ch. 7 Definition 6.7 and eq. (6.22); see also [84, 248]). Let $d := \dim \mathcal{X}_q$ and define

$$\mathrm{tr}_{\bar{\Sigma}}\, h := \bar{\Sigma}^{\mu\nu}h_{\mu\nu}, \qquad (\mathrm{div}_{\bar{\Sigma}}h)_\nu := \nabla^{(\mathrm{rel})\mu}h_{\mu\nu}, \qquad (\mathcal{L}_{\bar{\Sigma}}v)_{\mu\nu} := \nabla^{(\mathrm{rel})}_{(\mu}v_{\nu)} - \frac{1}{d}\,\bar{\Sigma}_{\mu\nu}\,\nabla^{(\mathrm{rel})}\cdot v,$$

together with the $\bar{\Sigma}$–adjoint

$$(\mathcal{L}_{\bar{\Sigma}}^{\dagger} h)_{\mu} := -2\Big(\nabla^{(\mathrm{rel})\nu} h_{\mu\nu} - \frac{1}{d}\,\nabla_{\mu}^{(\mathrm{rel})}\,\mathrm{tr}_{\bar{\Sigma}}\,h\Big)\,,$$

where adjoints are taken with respect to the $\bar{\Sigma}$–inner product (cf. Definition 9.33). Writing the *York vector–Laplacian* as $\Delta_V^{(\mathrm{rel})} := \mathcal{L}_{\bar{\Sigma}}^{\dagger}\mathcal{L}_{\bar{\Sigma}}$—elliptic on the realized weave configuration, modulo Killing vectors [43, 44, 7]—the projector is

$$\boxed{\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \;:=\; \mathbf{1} \;-\; \mathcal{L}_{\bar{\Sigma}}\,(\Delta_V^{(\mathrm{rel})})^{-1}\,\mathcal{L}_{\bar{\Sigma}}^{\dagger} \;-\; \frac{1}{d}\,\bar{\Sigma}\otimes\mathrm{tr}_{\bar{\Sigma}}} \tag{10.4}$$

acting on $\Gamma(S^2T^*\mathcal{X}_q)$. The construction (10.4) is the standard York–Lichnerowicz TT projector written in background–covariant form [43, 44].

This operator is idempotent and $\bar{\Sigma}$–self-adjoint,

$$(\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]})^2 = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}, \qquad \langle \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} h, k\rangle_{\bar{\Sigma}} = \langle h, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} k\rangle_{\bar{\Sigma}},$$

and it enforces the TT conditions

$$\mathrm{tr}_{\bar{\Sigma}}\Big(\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} h\Big) = 0, \qquad \mathrm{div}_{\bar{\Sigma}}\Big(\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} h\Big) = 0,$$

hence annihilates pure traces and longitudinal pieces,

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}(\varphi\,\bar{\Sigma}) = 0, \qquad \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\Big(\mathcal{L}_{\bar{\Sigma}} v\Big) = 0, \qquad \forall\,\varphi \in C^{\infty},\ v \in \Gamma(T^*\mathcal{X}_q).$$

It is stable and commuting with the certified band/typed structure (Propositions 10.9 and 10.10), in particular

$$\Big[\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]},\, \Pi^{(\mathrm{spec})}\Big] = \Big[\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]},\, \Pi^{(\mathrm{vac})}\Big] = \Big[\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]},\, \Pi^{(\mathrm{corr})}\Big] = 0,$$

and it satisfies TT purity as recorded in Lemma 10.28.

Meaning and role of $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ Definition Definition 10.3 provides a *background–covariant* way to extract the physically relevant, gauge–invariant content of a symmetric 2–tensor $h_{\mu\nu}$ relative to the realized chronogeometry $\bar{\Sigma}$ (with $\nabla^{(\mathrm{rel})}$ from Definition 6.7). The projector removes (i) pure traces and (ii) longitudinal, diffeomorphism–generated pieces, leaving the *transverse–traceless* remainder that carries spin–2 gravitational radiation. Because it is built from $\bar{\Sigma}$ and $\nabla^{(\mathrm{rel})}$, $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ is compatible with the realized weave configuration (Definition 10.1) and commutes with the typed/triad structure (Proposition 10.9), ensuring that TT extraction is consistent with the rest of the weave calculus (e.g., TT purity and leak control in Lemma 10.28).

Why the York vector–Laplacian and Killing fields? The decomposition encoded by (10.4) is the York–Lichnerowicz split: for suitable h one has the *orthogonal* decomposition

$$h_{\mu\nu} = \underbrace{\big(\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} h\big)_{\mu\nu}}_{\mathrm{TT}} + \underbrace{(\mathcal{L}_{\bar{\Sigma}} v)_{\mu\nu}}_{\text{longitudinal}} + \underbrace{\frac{1}{d}\,\bar{\Sigma}_{\mu\nu}\,\mathrm{tr}_{\bar{\Sigma}}\,h}_{\text{trace}}, \qquad v \;=\; (\Delta_V^{(\mathrm{rel})})^{-1}\,\mathcal{L}_{\bar{\Sigma}}^{\dagger} h \;\;\mathrm{mod}\;\; \ker\Delta_V^{(\mathrm{rel})}.$$

Here $\Delta_V^{(\mathrm{rel})} := \mathcal{L}_{\bar\Sigma}^\dagger \mathcal{L}_{\bar\Sigma}$ is the *York vector–Laplacian*, an elliptic operator whose kernel is spanned by *Killing fields* of $(\mathcal{X}_q, \bar\Sigma)$ [43, 44, 7]. Ellipticity guarantees a well-posed inversion on the orthogonal complement of this kernel, so the longitudinal component is removed by the term $\mathcal{L}_{\bar\Sigma}(\Delta_V^{(\mathrm{rel})})^{-1}\mathcal{L}_{\bar\Sigma}^\dagger$ while the pure trace is removed by $\frac{1}{d}\,\bar\Sigma \otimes \mathrm{tr}_{\bar\Sigma}$. The residual piece is TT by construction and, crucially, is independent of background isometries (Killing directions) which represent gauge redundancies rather than physical degrees of freedom [43, 7]. This is why (10.4) is the natural, covariant and rig–compatible way to isolate graviton channels in our framework.

Definition 10.4 (τ–kairos typed projectors)**.** On the τ–kairos typed block (typed, not a sector; Ch. 10 typed layer Definition 9.116), we admit the $\bar\Sigma$–self-adjoint, mutually orthogonal idempotents

$$S^{(b)}, \quad S^{(L)}, \quad \tilde S^{(p)}, \quad A^{(+)}, \quad A^{(-)},$$

corresponding respectively to *breathing scalar*, *longitudinal scalar*, *parity-odd* pseudoscalar (Levi–Civita built), and *helicity-resolved* vectors (Chrono–Kymon $\lambda = \pm$). They obey, on the τ-typed block,

$$S^{(i)}S^{(j)} = \delta_{ij}\,S^{(i)}, \qquad (S^{(i)})^{\dagger_{\bar\Sigma}} = S^{(i)}, \qquad S^{(b)} + S^{(L)} + \tilde S^{(p)} + A^{(+)} + A^{(-)} = \mathbf{1}_\tau, \tag{10.5}$$

where $\dagger_{\bar\Sigma}$ denotes adjoint with respect to the $\bar\Sigma$–inner products (Definition 9.33), and $\mathbf{1}_\tau$ is the identity on the τ–typed block.

Breathing vs. longitudinal scalars. $S^{(b)}$ projects onto the isotropic (trace-like) breathing mode generated by scalar dilations within the τ–typed block; $S^{(L)}$ projects onto the *longitudinal* gradient mode. Concretely, writing the relational Levi–Civita connection $\nabla^{(\mathrm{rel})}$ determined by $\bar\Sigma$ (Definition 6.7), one may characterize $S^{(L)}$ by the covariant gradient–divergence pair

$$\mathcal{G} := \nabla^{(\mathrm{rel})}, \qquad \mathcal{G}^\dagger := -\nabla^{(\mathrm{rel})}\cdot \ (\bar\Sigma\text{–adjoint}),$$

and the (typed) scalar Laplacian $\Delta_S^{(\mathrm{rel})} := \mathcal{G}^\dagger\mathcal{G}$ on the realized weave configuration (Definition 10.1), so that

$$S^{(L)} = \mathcal{G}\,\Big(\Delta_S^{(\mathrm{rel})}\Big)^{-1}\,\mathcal{G}^\dagger, \tag{10.6}$$

modulo $\ker\Delta_S^{(\mathrm{rel})}$ (constants/Killing modes on the realized weave configuration). The breathing projector $S^{(b)}$ is the complementary scalar part within (10.5) (see the typed split in Definition 10.2).

Pseudoscalar and helicity projectors. The pseudoscalar $\tilde S^{(p)}$ is odd under spatial parity and is built using the $\bar\Sigma$–Levi–Civita density $\varepsilon[\bar\Sigma]$ (Hodge dual on the realized weave configuration), while the vector projectors $A^{(\pm)}$ resolve *helicity* by diagonalizing the typed helicity operator H_τ:

$$\mathsf{H}_\tau A^{(\pm)} = \pm A^{(\pm)}, \qquad A^{(\pm)} = \tfrac{1}{2}\Big(\mathbf{1}_\tau \pm \mathsf{H}_\tau\Big), \tag{10.7}$$

with parity action recorded in Definition 10.58. Parity leaves $S^{(b)}, S^{(L)}$ invariant, flips $\tilde S^{(p)} \mapsto -\tilde S^{(p)}$, and swaps $A^{(+)} \leftrightarrow A^{(-)}$.

TT-null property and stability. All τ–typed projectors are *TT-null* with respect to the background-covariant TT projector $\Pi_{\bar\Sigma}^{[\mathrm{TT}]}$ of Definition 10.3:

$$\Pi_{\bar\Sigma}^{[\mathrm{TT}]}\,X = 0, \qquad X \in \{S^{(b)}, S^{(L)}, \tilde S^{(p)}, A^{(+)}, A^{(-)}\}, \tag{10.8}$$

and TT-nullity propagates through typed mixing as in Lemma 10.123 (see also Corollary 10.125). Commutation with the certified band/typed structure is summarized in Proposition 10.9, and invariance on the realized weave configuration is stated in Proposition 10.10.

On the meaning of CP and its role in the $\Theta\Theta$ scalar subspace. Throughout this chapter, *CP* denotes the standard combined charge–conjugation and parity transformation ($\mathsf{C} \times \mathsf{P}$) familiar from quantum field theory; it is not a symbol introduced ad hoc for the present framework. Within the scalar subspace of the $\Theta\Theta$ block, an abstract, involutive operator CP is taken to act as a symmetry of the rigged calculus, with $\mathsf{CP}^2 = \mathbf{1}_{\mathrm{sc}}$. This structure allows the scalar sector to be decomposed into CP-even and CP-odd components by means of idempotent projectors that extract the corresponding eigenspaces of CP, consistently with the typed weave algebra and the commutation-stability properties recorded earlier (Proposition 10.9 and Definition 9.116).

Projectors onto CP eigen-subspaces and state classification. In the scalar subspace, the CP projectors are introduced algebraically by

$$\Pi^{(\mathrm{CP}\pm)} \;=\; \tfrac{1}{2}(\mathbf{1}_{\mathrm{sc}} \pm \mathsf{CP})\,, \qquad \Pi^{(\mathrm{CP}+)} + \Pi^{(\mathrm{CP}-)} = \mathbf{1}_{\mathrm{sc}}, \quad \Pi^{(\mathrm{CP}\pm)}\Pi^{(\mathrm{CP}\mp)} = 0,$$

so that a scalar state $|\phi\rangle$ is CP-even precisely when $\mathsf{CP}\,|\phi\rangle = +\,|\phi\rangle$ (equivalently $\Pi^{(\mathrm{CP}+)}\,|\phi\rangle = |\phi\rangle$) and CP-odd when $\mathsf{CP}\,|\phi\rangle = -\,|\phi\rangle$ (equivalently $\Pi^{(\mathrm{CP}-)}\,|\phi\rangle = |\phi\rangle$); orthogonality and closure are summarized in Lemma 10.45. In a local field representation, for a possibly complex scalar $\Theta(x)$ one may take

$$(\mathsf{CP})\,\Theta(t,\mathbf{x})\,(\mathsf{CP})^{-1} \;=\; \eta_{\mathrm{CP}}\;\Theta^{*}(t,-\mathbf{x}), \qquad \eta_{\mathrm{CP}} \in \{+1,-1\},$$

where $\eta_{\mathrm{CP}} = +1$ yields a CP-even (scalar) and $\eta_{\mathrm{CP}} = -1$ a CP-odd (pseudoscalar) transformation law. For neutral/real radial modes, C reduces to complex conjugation and the classification effectively coincides with parity even/odd; the full CP labelling is nevertheless retained to accommodate charged or multi-field Higgs-like sectors without modification.

CP refinement of the radial typed subspace and diagnostics. The radial typed projector in the scalar channel is refined by the CP idempotents to

$$\Pi_{\Theta}^{[\mathrm{rad}\pm]} \;:=\; \Pi_{\Theta}^{[\mathrm{rad}]}\,\Pi^{(\mathrm{CP}\pm)},$$

with $\Pi^{(\mathrm{CP}\pm)}$ commuting with the provenance projectors $\Pi^{(a)}$ and typed idempotents, ensuring compatibility with the realized weave configuration (Proposition 10.9 and Definition 9.116). A CP-pure radial excitation has support entirely in one of these images (e.g. $\Pi_{\Theta}^{[\mathrm{rad}+]}\,|\phi\rangle = |\phi\rangle$, $\Pi_{\Theta}^{[\mathrm{rad}-]}\,|\phi\rangle = 0$ for CP-even), while deviations from purity are quantified by the CP-asymmetry diagnostic introduced later (Definition 10.51). These considerations motivate the following definition.

Definition 10.5 (Higgs (radial) CP-refined projectors)**.** In the scalar subspace of the $\Theta\Theta$ block, let $\Pi^{(\mathrm{CP}\pm)}$ commute with $\Pi^{(a)}$ and with typed idempotents (commutation stability Proposition 10.9 and Definition 9.116). Motivated by the CP-even/odd splitting of scalar

excitations in Higgs-like sectors [14, 82, 97, 96, 98, 465, 466, 467], we refine the radial type (Ch. 10 physics layer) by

$$\Pi_{\Theta}^{[\mathrm{rad}\pm]} := \Pi_{\Theta}^{[\mathrm{rad}]}\,\Pi^{(\mathrm{CP}\pm)}, \qquad \Pi^{(\mathrm{CP}+)} + \Pi^{(\mathrm{CP}-)} = \mathbf{1}_{\mathrm{sc}}, \quad \Pi^{(\mathrm{CP}\pm)}\Pi^{(\mathrm{CP}\mp)} = 0.$$

Orthogonality and CP-purity are recorded in Lemma 10.45; the CP-asymmetry diagnostic is Definition 10.51. In particular, $\Pi^{(\mathrm{CP}+)}$ ($\Pi^{(\mathrm{CP}-)}$) selects CP-even (CP-odd) scalar modes within the radial typed subspace, in the sense standard in Higgs phenomenology [466, 467].

CP-purity and CP eigenstates. Within the scalar subspace of the $\Theta\Theta$ block, the idempotents $\Pi^{(\mathrm{CP}\pm)}$ in Definition 10.5 select *CP eigenstates* of the abstract CP operator CP: CP-even states satisfy $\mathsf{CP}\,|\phi\rangle = +\,|\phi\rangle$ and are isolated by $\Pi^{(\mathrm{CP}+)}$, while CP-odd states satisfy $\mathsf{CP}\,|\phi\rangle = -\,|\phi\rangle$ and are isolated by $\Pi^{(\mathrm{CP}-)}$ (orthogonality/closure in Lemma 10.45). In our typed calculus this means a *CP-pure* radial excitation has support entirely in one of the projector images, e.g. $\Pi_{\Theta}^{[\mathrm{rad}+]}\,|\phi\rangle = |\phi\rangle$ and $\Pi_{\Theta}^{[\mathrm{rad}-]}\,|\phi\rangle = 0$ for CP-even. This mirrors the standard Higgs-field decomposition into CP-even (scalar) and CP-odd (pseudoscalar) components used in QFT and Higgs phenomenology [14, 82, 97, 96, 98, 465, 466, 467].

CP asymmetry and why it matters. The *CP-asymmetry* diagnostic of Definition 10.51 quantifies departures from CP-purity by measuring the imbalance or interference between the $\Pi^{(\mathrm{CP}+)}$ and $\Pi^{(\mathrm{CP}-)}$ components; it vanishes iff the state (or correlator) is CP-pure and becomes nonzero when CP-violating mixing is present (yet remains well-defined thanks to the commutation stability in Proposition 10.9). Physically, CP-even vs. CP-odd composition controls selection rules and characteristic angular correlations in production/decay channels, while a nonzero asymmetry encodes genuine CP violation in the scalar sector—precisely the effect scrutinized in Higgs-like theories and their extensions [466, 467]. In our framework, refining the radial projector to $\Pi_{\Theta}^{[\mathrm{rad}\pm]}$ isolates these features at the operator level, ensuring that CP properties are tracked consistently across the realized weave configuration and downstream witnesses.

Riesz selectors for typed spin content: purpose and construction. The objective is to isolate the linearized dynamics carried by the helicity–2 (spin–2, TT/graviton) and spin–0 (radial/Higgs-like) channels within the certified band of the instrumentation $\mathcal{I}^{[\mathscr{S}]}$. This is achieved via the Riesz (holomorphic) functional calculus: given a closed kinetic operator K and an isolated spectral component ("spectral island") whose boundary lies in the resolvent set, the contour integral of the resolvent $(K - z)^{-1}$ defines the bounded projector onto the corresponding invariant spectral subspace. Summing such Riesz projectors over all spectral islands assigned to a given spin class (within the band) yields the *total spectral selector* for that class; measurability and invariance on the realized weave configuration follow from the stability statements recorded earlier (Definition 10.1, Propositions 10.10 and 10.149, and Lemma 10.137).

Spin–2 versus spin–0 carriers and compatibility with the weave. In this chapter, the spin–2 selector is constructed from the Gaiamiti kinetic operator $K_{\Phi\Phi}$, while the spin–0 selector is

constructed from the scalar sub-block $K^{(0)}_{\Theta\Theta}$ of the Tantrakyma sector; both are implicitly band-limited on $\mathcal{I}^{[\mathscr{S}]}$. Because these selectors are defined by holomorphic functional calculus, they commute with the BRST/physical projector $\mathsf{P}_{\mathrm{phys}}$, the provenance (spec/vac/corr) projectors, and the typed idempotents of the weave algebra (Proposition 10.9 and Definition 9.116). This guarantees compatibility with the TT/Higgs witnesses and with Källén–Lehmann positivity in the sense summarized in eq. (7.12) (see also the pre-physics analogue in Ch. 9).

Spectral islands. In this chapter, "spectral islands" are the isolated pieces of the spectrum of the relevant kinetic operators that are singled out by the realized weave configuration band (cf. Definition 10.1). Each such piece is separated from the remainder of the spectrum by a nonzero gap that lies entirely in the resolvent set, so a closed contour can be drawn around it without crossing any spectral points. The Riesz (holomorphic) functional calculus then associates to each island a bounded projector onto the invariant subspace carried by that portion of the spectrum; summing over the islands assigned to a given spin class yields the corresponding *spectral selector* (cf. Definition 10.7).

Operator-theoretically, let K be a closed operator with spectrum $\sigma(K)$ and resolvent set $\rho(K) = \mathbb{C} \setminus \sigma(K)$. Inside the certified band (determined by $\mathcal{I}^{[\mathscr{S}]}$ of Definition 10.1), consider connected components of $\sigma(K)$ that admit a positively oriented simple closed contour $\gamma \subset \rho(K)$ enclosing only that component. For such components, the Riesz integral $\frac{1}{2\pi i} \oint_\gamma (K - z)^{-1}\, dz$ yields an idempotent that projects onto the invariant subspace supported by the enclosed spectral portion (cf. Definition 10.7). In the purely discrete case, an island may reduce to a finite set of isolated eigenvalues; in the self-adjoint continuous case, it may be a compact interval separated by a gap; the definition covers both and extends to non-normal operators.

In the weave calculus, "helicity-2 islands" of $K_{\Phi\Phi}$ and "spin-0 islands" of $K^{(0)}_{\Theta\Theta}$ are those connected spectral components within the certified band whose invariant subspaces carry, respectively, TT (spin-2) and radial/Higgs (spin-0) typed content—compatibly with the commuting provenance and typed projectors (cf. Definitions 10.3 and 10.5 and Proposition 10.9). The symbols γ_i and Γ_j denote the enclosing contours of these components; the index sets $\mathfrak{S}_2$ and $\mathfrak{S}_0$ range over the corresponding families of components (one index per island; cf. Definition 10.7).

Definition 10.6 (Spectral islands (band-limited))**.** Let K be a closed operator on a Hilbert space with spectrum $\sigma(K)$ and resolvent set $\rho(K)$. Fix a certified band (from $\mathcal{I}^{[\mathscr{S}]}$, cf. Definition 10.1) and set $\sigma_{\mathrm{band}}(K) := \sigma(K) \cap \mathrm{band}$. A *spectral island* of K (within the band) is a connected subset $S \subset \sigma_{\mathrm{band}}(K)$ for which there exists a positively oriented simple closed contour $\gamma \subset \rho(K)$ such that: (i) S is contained in the bounded component of $\mathbb{C} \setminus \gamma$; (ii) $\sigma(K) \cap \mathrm{int}(\gamma) = S$. The associated Riesz projector is $P_S := \frac{1}{2\pi i} \oint_\gamma (K - z)^{-1}\, dz$, which is idempotent and projects onto the invariant subspace supported by S (cf. Definition 10.7). For disjoint islands S_α, the projectors $\{P_{S_\alpha}\}$ are mutually orthogonal and $\sum_\alpha P_{S_\alpha}$ (finite or strongly convergent) projects onto $\bigcup_\alpha S_\alpha$ (cf. Definition 10.7).

Meaning of the summations and contour symbols. The index sets $\mathfrak{S}_2$ and $\mathfrak{S}_0$ label the *connected spectral components* (spectral islands, Definition 10.6) of $K_{\Phi\Phi}$ and $K^{(0)}_{\Theta\Theta}$ that are assigned—by the typing rules and the certified band—to the spin–2 and spin–0 classes,

respectively (cf. Definitions 10.1, 10.3 and 10.5 and Proposition 10.9). The sums $\sum_{i\in\mathfrak{S}_2}$ and $\sum_{j\in\mathfrak{S}_0}$ are therefore sums *over islands*, not over spacetime indices or eigenvalue multiplicities. For each island, γ_i (helicity–2) and Γ_j (spin–0) denote positively oriented (counterclockwise) closed contours in the complex plane that enclose exactly that island and no other part of the spectrum; outside the enclosed set, the contour lies in the resolvent region (cf. Definition 10.7). The integrands $(K_{\Phi\Phi}-z)^{-1}$ and $(K^{(0)}_{\Theta\Theta}-z)^{-1}$ are the corresponding resolvents (cf. Definition 10.7). The Riesz integral $\frac{1}{2\pi i}\oint_{\gamma_i}\cdot\,dz$ (and analogously over Γ_j) yields an idempotent projector onto the invariant subspace of the enclosed island; for disjoint islands these projectors are mutually orthogonal, and their finite (or strongly convergent countable) sum projects onto the union of those spectral pieces (cf. Definition 10.7). Band limitation typically ensures that only finitely many islands contribute within $\mathcal{I}^{[\mathscr{J}]}$ (cf. Definition 10.1).

Intuition and downstream use. In purely discrete situations, each contour encloses one or finitely many eigenvalues and the Riesz projector reduces to the sum of the associated eigenprojectors. The present construction is preferred because it remains valid under continuous spectrum, non-normality, and admissible perturbations on the certified plateau $\mathfrak{P}$. The selectors $\mathsf{P}^{(2)}_{\mathrm{spec}}$ and $\mathsf{P}^{(0)}_{\mathrm{spec}}$ thus provide coordinate- and gauge-independent operator-level filters for the TT (graviton) and Higgs-radial (scalar) channels used in the TT/Higgs witnesses and in the typed propagator machinery (Section 10.9.5). These considerations motivate the following definition.

Definition 10.7 (Riesz spectral selectors (spin-2 and spin-0))**.** Let $K_{\Phi\Phi}$ be the linearized, gauge-fixed kinetic operator in the gaiamiti sector and $K^{(0)}_{\Theta\Theta}$ the scalar sub-block in the tantrakyma sector (both band-limited on $\mathcal{I}^{[\mathscr{J}]}$; Definition 10.1 and Proposition 10.10). Define the (Riesz) spectral projectors via the holomorphic functional calculus [115, 253]:

$$\mathsf{P}^{(2)}_{\mathrm{spec}} := \sum_{i\in\mathfrak{S}_2}\frac{1}{2\pi i}\oint_{\gamma_i}\Big(K_{\Phi\Phi}-z\Big)^{-1}dz, \qquad \mathsf{P}^{(0)}_{\mathrm{spec}} := \sum_{j\in\mathfrak{S}_0}\frac{1}{2\pi i}\oint_{\Gamma_j}\Big(K^{(0)}_{\Theta\Theta}-z\Big)^{-1}dz,$$

for contours γ_i,Γ_j enclosing helicity-2 and spin-0 spectral islands on the certified band (measurability/invariance Proposition 10.149 and Lemma 10.137; Källén–Lehmann positivity eq. (7.12) as standard in QFT [14, 255, 230, 82, 244]). (For the pre-physics analogue, see Ch. 9.) By functional calculus, these projectors commute with $\mathsf{P}_{\mathrm{phys}}$, the triad projectors, and the typed idempotents (Proposition 10.9) and are used in the TT/Higgs witness entries (Section 10.9.5).

Structural facts.

Lemma 10.8 (Orthogonality and completeness on the τ–kairos block)**.** *Let*

$$\mathcal{S}_\tau := \{S^{(b)},\, S^{(L)},\, \tilde{S}^{(p)}\}, \qquad \mathcal{A}_\tau := \{A^{(+)},\, A^{(-)}\}$$

be the τ–typed projectors of Definition 10.4 on the realized weave configuration $\mathcal{I}^{[\mathscr{J}]}$; see also the typed layer Definition 9.116. On the τ–typed carrier, the following relations hold:

$$S^{(i)}S^{(j)}=\delta_{ij}S^{(i)}, \qquad A^{(\lambda)}A^{(\lambda')}=\delta_{\lambda\lambda'}A^{(\lambda)}, \qquad S^{(i)}A^{(\lambda)}=0,$$

for all $S^{(i)}, S^{(j)} \in \mathcal{S}_\tau$ *and* $A^{(\lambda)}, A^{(\lambda')} \in \mathcal{A}_\tau$*, and the completeness relation*

$$\sum_{i\in\{b,L,p\}} S^{(i)} + \sum_{\lambda=\pm} A^{(\lambda)} = \mathbf{1}_\tau.$$

Proof. Notation. Let $\mathcal{H}_\tau$ denote the τ–typed carrier Hilbert space on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ with inner product $\langle\cdot,\cdot\rangle_{\bar\Sigma}$. For a linear map $P : \mathcal{H}_\tau \to \mathcal{H}_\tau$, write the range or image is

$$\operatorname{Ran} P \;:=\; \{\, v \in \mathcal{H}_\tau :\; \exists\, u \in \mathcal{H}_\tau \text{ with } v = Pu \,\}$$

for the *range (image)* subspace of P (and, when needed, $\ker P := \{u :\; Pu = 0\}$ for its kernel).

Idempotence and intra-family orthogonality. By Definition 10.4, the maps $S^{(b)}, S^{(L)}, \tilde S^{(p)}$ (scalar sub-block) and $A^{(+)}, A^{(-)}$ (helicity-resolved vector sub-block) are typed projectors on $\mathcal{H}_\tau$: they are idempotent and $\bar\Sigma$–Hermitian, and are mutually orthogonal within their respective families (typed algebra on the band; see also Ch. 10 Definition 9.116). Hence $S^{(i)}S^{(j)} = \delta_{ij}S^{(i)}$ and $A^{(\lambda)}A^{(\lambda')} = \delta_{\lambda\lambda'}A^{(\lambda)}$. Equivalently, for $i \neq j$ (resp. $\lambda \neq \lambda'$), $\operatorname{Ran} S^{(i)} \perp \operatorname{Ran} S^{(j)}$ (resp. $\operatorname{Ran} A^{(\lambda)} \perp \operatorname{Ran} A^{(\lambda')}$).

Scalar–vector orthogonality. The τ–typed carrier splits (typed block decomposition, Ch. 10 Definition 9.116) into an orthogonal direct sum of a scalar subspace and a vector (Chrono–Kymon) subspace. Each $S^{(i)}$ projects into the scalar subspace, while each $A^{(\lambda)}$ projects into the vector subspace. These carrier spaces are orthogonal in $\langle\cdot,\cdot\rangle_{\bar\Sigma}$, so $S^{(i)}A^{(\lambda)} = 0$, and hence $\operatorname{Ran} S^{(i)} \perp \operatorname{Ran} A^{(\lambda)}$.

Completeness. The typed decomposition further refines the scalar subspace as $\operatorname{Ran} S^{(b)} \oplus \operatorname{Ran} S^{(L)} \oplus \operatorname{Ran} \tilde S^{(p)}$ and the vector subspace as $\operatorname{Ran} A^{(+)} \oplus \operatorname{Ran} A^{(-)}$. Thus,

$$\mathcal{H}_\tau = \operatorname{Ran} S^{(b)} \;\oplus\; \operatorname{Ran} S^{(L)} \;\oplus\; \operatorname{Ran} \tilde S^{(p)} \;\oplus\; \operatorname{Ran} A^{(+)} \;\oplus\; \operatorname{Ran} A^{(-)}.$$

Equivalently, for every $v \in \mathcal{H}_\tau$, $v = \Big(\sum_{i\in\{b,L,p\}} S^{(i)} + \sum_{\lambda=\pm} A^{(\lambda)}\Big)v$, which yields the projector identity $\sum_{i\in\{b,L,p\}} S^{(i)} + \sum_{\lambda=\pm} A^{(\lambda)} = \mathbf{1}_\tau$.

All statements hold on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$, and the projectors commute with the band/BRST machinery by Propositions 10.9 and 10.10, ensuring stability under triad sandwiches and audit moves. □

Audit moves. Let $\mathfrak{A}_{\text{audit}}$ be the unital *-algebra generated on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ by

$$\mathsf{P}_{\text{phys}},\; P_x(\lambda_{\text{UV}}),\; K^{(\ell)}_{\text{IR}},\; \Pi^{(\text{spec})}, \Pi^{(\text{vac})}, \Pi^{(\text{corr})},\; S^{(b)}, S^{(L)}, \tilde S^{(p)}, A^{(\pm)},\; \Pi^{[\text{TT}]}_{\bar\Sigma},\; \Pi^{(\text{CP}\pm)},\; \mathsf{P}^{(2)}_{\text{spec}},\; \mathsf{P}^{(0)}_{\text{spec}},$$

subject to the commutation–stability relations of Proposition 10.9. An *audit move* is any $M \in \mathfrak{A}_{\text{audit}}$ acting by composition or sandwiching on operators/states (e.g. $O \mapsto MOM$, $v \mapsto Mv$). For every such M, the projector algebra on the τ block is stable:

$$S^{(i)}S^{(j)} = \delta_{ij}S^{(i)}, \quad A^{(\lambda)}A^{(\lambda')} = \delta_{\lambda\lambda'}A^{(\lambda)}, \quad S^{(i)}A^{(\lambda)} = 0, \quad \sum_{i\in\{b,L,p\}} S^{(i)} + \sum_{\lambda=\pm} A^{(\lambda)} = \mathbf{1}_\tau,$$

and the same identities hold after left/right insertions of M. In particular, "triad sandwiches" are the special case $M = \Pi^{(a)}$ with $a \in \{\text{spec}, \text{vac}, \text{corr}\}$.

Proposition 10.9 (Commutation with triad and TT)**.** *On the realized weave configuration* $\mathcal{I}^{[\mathcal{S}]}$ *(Def. Definition 10.1), every* typed *idempotent* $\Pi^{[\chi]}$ *(Ch. 10 typed layer Definition 9.116; concretely* $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$*,* $\{S^{(b)}, S^{(L)}, \tilde{S}^{(p)}, A^{(+)}, A^{(-)}\}$ *from Definition 10.4, and* $\Pi^{[\mathrm{rad}\pm]}_{\Theta}$ *from Definition 10.5) commutes with each triad projector* $\Pi^{(a)}$ *(*$a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$*; Definition 10.2):*

$$[\Pi^{(a)}, \Pi^{[\chi]}] = 0 \qquad (a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}).$$

Moreover, every τ*–kairos type is* TT–null*:*

$$\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}} X = 0, \qquad X \in \{S^{(b)}, S^{(L)}, \tilde{S}^{(p)}, A^{(+)}, A^{(-)}\} \quad (\textit{Definition 10.4}). \tag{10.9}$$

Proof. **(A) Preliminaries on the realized weave configuration and selectors.** By instrumentation hygiene (Proposition 10.10), the band tags P_x and IR windows $K^{(\ell)}_{\mathrm{IR}}$ are idempotent contractions commuting with the BRST projector $\mathsf{P}_{\mathrm{phys}}$, so all sandwiched operators live on $\mathcal{H}_{\mathrm{phys}} = \operatorname{Ran} \mathsf{P}_{\mathrm{phys}}$ with a bounded holomorphic functional calculus. The triad maps $\Pi^{(a)}$ are Riesz projectors determined by certified contours (triad provenance; Definitions 10.2 and 10.7 and Proposition 10.149).

(B) Commutation with triads. Each triad projector is a contour integral of the resolvent of the (band–limited, gauge–fixed) kinetic operator on the corresponding block (spin-2, 0, or the typed vector block; Definition 10.7 and Proposition 10.149). Typed idempotents $\Pi^{[\chi]}$ (TT, τ–types, Higgs–CP) are obtained fiberwise from symmetry generators (transverse/trace, parity, helicity, CP) and lifted as $\bar{\Sigma}$–Hermitian idempotents stable on the band (Ch. 10 Definition 9.116; instantiated in Definitions 10.3 to 10.5). By construction they commute with the relevant kinetic operators on the realized weave configuration (typed stability; Definition 9.116, item (P7)); hence they commute with each resolvent $(K - z)^{-1}$ and therefore with its Riesz integral. Thus $[\Pi^{(a)}, \Pi^{[\chi]}] = 0$ for all a and all typed $\Pi^{[\chi]}$.

(C) TT–nullity for τ types. Equation (10.9) is stated in Definition 10.4: the τ–typed projectors map into scalar/vector carriers orthogonal (w.r.t. $\langle \cdot, \cdot \rangle_{\bar{\Sigma}}$) to the TT subspace, and the background–covariant TT projector $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ (Ch. 7 Definition 10.3) annihilates traces/divergences, hence the τ images. Orthogonality within the τ–block is recorded in Lemma 10.8; TT–null preservation across any τ–involving channel is in Lemma 10.123, with the graviton readout invariance in Corollary 10.125.

All equalities hold on $\mathcal{I}^{[\mathcal{S}]}$ and are preserved under admissible retunes and arena switches by Propositions 10.10 and 10.149. □

Meaning and importance. Proposition Proposition 10.9 asserts that, on the realized weave configuration $\mathcal{I}^{[\mathcal{S}]}$ (Definition 10.1), the *spectral bookkeeping* provided by the triad projectors $\Pi^{(a)}$ (Definition 10.2) is compatible with all *typed* decompositions used operationally—namely the background–covariant TT projector $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ (Definition 10.3), the τ–kairos types $S^{(b)}, S^{(L)}, \tilde{S}^{(p)}, A^{(+)}, A^{(-)}$ (Definition 10.4), and the Higgs CP-refined radial projectors $\Pi^{[\mathrm{rad}\pm]}_{\Theta}$ (Definition 10.5). The commutators $[\Pi^{(a)}, \Pi^{[\chi]}] = 0$ guarantee that *typing* (trace/transverse, parity, helicity, CP) and *spectral triage* (spec/vac/corr) may be performed in either order without changing outcomes. Practically, this removes order-of-operations ambiguities in witness extraction and channel isolation, ensuring that TT graviton readouts, τ–typed diagnostics, and CP analyses are all computed on the same block-diagonal stage of the

theory and remain stable under the band/arena hygiene of Proposition 10.10. The second clause—TT–nullity (10.9)—establishes a strict separation between the τ–typed scalar/vector carriers and the TT graviton subspace, preventing spurious leakage of non-TT content into spin-2 observables (cf. TT purity in Definition 10.3 and its use downstream in TT/Higgs witnesses, Section 10.9.5).

Why the proof matters. The proof reduces the commutation statement to the *holomorphic functional calculus*: each triad projector is a Riesz selector of a band-limited, gauge-fixed kinetic operator (Definition 10.7 and Proposition 10.149), while each typed idempotent $\Pi^{[\chi]}$ is constructed to commute with those kinetic blocks on the realized weave configuration (typed stability; Ch. 10 Definition 9.116). Commutation with resolvents implies commutation with their contour integrals, giving $[\Pi^{(a)}, \Pi^{[\chi]}] = 0$. TT–nullity then follows from the defining action of $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$—which annihilates traces/divergences—together with the geometry of the τ–typed images and its preservation under typed mixing (Lemma 10.123 and Corollary 10.125). Conceptually, these steps certify that our decomposition is not merely convenient but *mathematically rigid*: spectral selection, typing, and TT extraction cohere on $\mathcal{I}^{[\mathscr{S}]}$, so that all subsequent constructions (e.g., graviton-channel isolation and CP diagnostics) inherit consistency and robustness from the realized weave configuration-level algebra.

Proposition 10.10 (BRST/Nielsen flatness and band–covariance)**.** *Let $\mathcal{J}$ be any finite composition of*

$$\mathsf{P}_{\mathrm{phys}},$$
$$\Pi^{(a)} \ (a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}) \ \textit{(Definition 10.2)},$$
$$\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}} \ \textit{(Definition 10.3)},$$
$$\textit{typed projectors } X \ \textit{(Definitions 10.4 and 10.5)},$$

and Riesz spectral projectors $\mathsf{P}^{(s)}_{\mathrm{spec}}$ ($s \in \{0, 2\}$; Definition 10.7), all evaluated on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ (Def. Definition 10.1). Then for any $\mathcal{J}$–projected observable $\mathcal{O}$ and any physical state on $\mathcal{H}_{\mathrm{phys}} = \mathrm{Ran}\,\mathsf{P}_{\mathrm{phys}}$, the correlation functions $\langle \mathcal{J}\mathcal{O}\rangle_{\mathcal{H}_{\mathrm{phys}}}$ are

(i) gauge–parameter independent *(Nielsen flat) on shell; and*

(ii) stable under admissible retunes (P_x, K_{IR}) *within the certified class (including IR–tightening, spectral–window refinement, and arena switch), i.e. they vary continuously and remain uniformly bounded along admissible paths.*

Proof. **(i) Nielsen flatness.** Let ξ denote a gauge parameter of the gauge–fixed kinetic operators and correlators. By the BRST/Nielsen identity (embedded form, see Ch. 8 and Prop. 7.174 used throughout), one has for any (sufficiently regular) observable $\mathcal{A}$

$$\partial_\xi \langle \mathcal{A}\rangle_{\mathcal{H}_{\mathrm{phys}}} = \langle [\hat{Q}_{\mathrm{BRST}}, \mathcal{K}_\xi]\, \mathcal{A}\rangle_{\mathcal{H}_{\mathrm{phys}}} + \langle \mathrm{EOM}_\xi(\mathcal{A})\rangle_{\mathcal{H}_{\mathrm{phys}}},$$

with $\hat{Q}_{\mathrm{BRST}}$ nilpotent and EOM_ξ vanishing on shell. Choose $\mathcal{A} = \mathcal{J}\mathcal{O}$ with $\mathcal{J}$ as in the statement. By typed stability on the realized weave configuration (Ch. 10 Definition 9.116, item (P7)) and commutation of typed/triad/Riesz projectors with the band operators

(Prop. 10.9 and functional–calculus commutation for Riesz selectors Definition 10.7 and Proposition 10.149), all components of $\mathcal{J}$ are BRST–closed on $\mathcal{H}_{\text{phys}}$. Hence the BRST–exact term has vanishing expectation, and the EOM term vanishes on shell; therefore $\partial_\xi \langle \mathcal{J}\mathcal{O}\rangle_{\mathcal{H}_{\text{phys}}} = 0$. This establishes Nielsen flatness.

(ii) Band–covariance and stability under retunes. Consider an admissible path of retunes $t \mapsto (P_x(t), K_{\text{IR}}^{(\ell(t))}, \text{Loc}(t))$ as in Def. 10.101. By instrumentation hygiene (band/BRST compatibility), Chs. 7–8, and Prop. 10.149, the resolvents $(K - z)^{-1}$ vary norm–continuously along t on the certified contours, so the associated Riesz projectors vary norm–continuously as Bochner integrals. Typed/triad idempotents are fixed (or vary continuously if parametrized) and commute with the band tags (Definitions 10.2 to 10.5, Prop. 10.9). Thus $t \mapsto \mathcal{J}(t)$ is norm–continuous on $\mathcal{H}_{\text{phys}}$.

Let $\|\cdot\|_{\mathcal{I}}$ be any admissible band–invariant norm (Def. 10.143). By equivalence across the family $\mathfrak{N}$ (Prop. 10.147), there are rig–dependent constants $c_\pm$ such that norms along different admissible choices are uniformly comparable; IR–tightening is contractive in these norms (Prop. 10.147, last clause). Therefore $\|\mathcal{J}(t) - \mathcal{J}(t_0)\|_{\mathcal{I}} \to 0$ as $t \to t_0$. Since correlation functionals are continuous with respect to the operator norm on $\mathcal{H}_{\text{phys}}$ (trace–class/HS bounds inherited from the realized weave configuration; see also the calibrated meters' finiteness Lem. 10.83), it follows that $\langle \mathcal{J}(t)\mathcal{O}\rangle_{\mathcal{H}_{\text{phys}}} \to \langle \mathcal{J}(t_0)\mathcal{O}\rangle_{\mathcal{H}_{\text{phys}}}$ along any admissible retune, and the correlators remain uniformly bounded along the path.

Finally, arena switches preserve the certified spectral set and the Riesz projectors (Lemma 10.137); hence $\mathcal{J}$ is unchanged up to the continuous variations already controlled above. This proves stability under all admissible retunes in the certified class. □

Corollary 10.11 (Gauge/retune invariance of visibilities and onsets)**.** *For any sector $\mathscr{S} \in \{\Phi\Phi, \Theta\Theta, \Phi\Theta\}$ and admitted type $[\chi]$, the calibrated visibilities $\Lambda_{\text{law}}^{[\mathscr{S};\chi]}$, $\Lambda_{\text{state}}^{[\mathscr{S};\chi]}$ (Def. Definition 10.82) and the certified onset $r^\star[\chi]$ (Def. Definition 10.85) are:*

(i) Gauge–parameter independent *(Nielsen–flat) on shell; and*

(ii) Stable under admissible retunes $(P_x, K_{\text{IR}}, \text{Loc})$ *in the certified class (continuous in those parameters and uniformly bounded along admissible paths).*

In particular, all diagnostic entries used for SynTh/Audit (e.g. ε, Δ_P, Δ_{CP}; Definition 10.80 and Remark 10.81) inherit the same invariance, and threshold crossings are unaffected by a change of gauge.

Proof. Each component of the law residual vector $\mathbf{R}^{[\mathscr{S};\chi]}$ and state feature vector $\mathbf{S}^{[\mathscr{S};\chi]}$ (Def. Definition 10.80) is computed by applying BRST/typed/triad/Riesz projectors to the kinetic objects and then taking a band–invariant norm from the admissible family $\mathfrak{N}$ (Def. Definition 10.143). Therefore all meters/visibilities are functionals of $\mathcal{J}$–projected observables with $\mathcal{J}$ a finite composition of P_{phys}, $\Pi^{(a)}$, $\Pi_{\bar{\Sigma}}^{[\text{TT}]}$, typed projectors, and Riesz selectors (exactly the class covered by Proposition 10.10).

Gauge independence. By Proposition 10.10(i) (Nielsen flatness of $\mathcal{J}$–projected correlators) together with the calibration maps being fixed scalar functions (Def. Definition 10.82), each entry of $\mathbf{R}$ and $\mathbf{S}$, hence Λ_{law} and Λ_{state}, is independent of the gauge parameter ξ on shell. Consequently, the threshold inequalities in Definition 10.85 are gauge–invariant and so is $r^\star[\chi]$.

Stability under admissible retunes. By Proposition 10.149, resolvents vary Bochner–measurably and Riesz projectors vary continuously along admissible audit moves (Def. Definition 10.101); typed and triad idempotents commute and remain stable on the realized weave configuration (Prop. Proposition 10.9, Ch. 10 Definition 9.116). Thus $t \mapsto \mathcal{J}(t)$ is norm–continuous on $\mathcal{H}_{\text{phys}}$. Band–invariant norms from $\mathfrak{N}$ are equivalent and contractive under IR tightening (Defs. Definition 10.143, Prop. Proposition 10.147), so $\|\mathcal{J}(t) - \mathcal{J}(t_0)\|_{\mathcal{I}} \to 0$ and the corresponding ledger entries vary continuously and stay bounded. The calibrations f_j, g_m are C^1 and bounded/monotone (Def. Definition 10.82), hence the visibilities are continuous and bounded along the path. Therefore the superlevel sets in Definition 10.85 vary upper–semicontinuously, and, with the monotonicities in Lemma 10.102 and slope bounds in Proposition 10.88, the onset $r^\star[\chi] = \inf \mathcal{A}_{[\chi]}$ is stable under admissible retunes.

Finally, diagnostics $\varepsilon, \Delta_P, \Delta_{\text{CP}}$ enter as components of $\mathbf{S}$ (Remark 10.81); they are built from the same $\mathcal{J}$–projected objects and norms, hence inherit gauge independence and retune stability as above. □

Remark 10.12 (Why instrumentation-first)**.** By fixing the realized weave configuration $\mathcal{I}^{[\mathcal{F}]}$ (Def. Definition 10.1) and the typed algebra on the band (Ch. 10: Definition 9.116, instantiated in Definitions 10.3 to 10.5), every subsequent "object definition" (–kairos, graviton, Higgs, Chrono–Kymon) is a *projected object on* $\mathcal{H}_{\text{phys}} = \operatorname{Ran} \mathsf{P}_{\text{phys}}$. This ensures:

1. we never violate SynTh's "same instrumentation" clause—comparisons/audits are made on the *same* realized weave configuration (cf. Definitions 8.98 and 8.99);
2. every visibility/witness is BRST–clean and band/triad–covariant—ledgers/meters are built from $\mathcal{J}$–projected observables and admissible norms (Defs. Definitions 10.80, 10.82 and 10.143) and inherit Nielsen flatness and retune stability (Prop. Proposition 10.10); they also respect the provenance cones spec/vac/corr (triads; Definition 10.2);
3. future species (new poles, new types) are absorbed by *adding contours or idempotents*, never by rewriting prior equations: add Riesz contours/selectors (Defs./Props. Definition 10.7 and Proposition 10.149) or typed idempotents (typed stability/commutation Definition 9.116 and Proposition 10.9).

10.3 The τ-kairos Field

The Concept of Kairos

The ancient Greek word καιρός (kairos) encodes a conception of time more nuanced than mere duration. It does not denote *chronos* (χρόνος)—chronological, metrically accumulated time—but rather "the right moment," "the opportune time," or a "critical instant." In standard lexicography, *kairos* marks a qualitative, situationally charged now—a juncture whose value lies not in its quantitative length but in what may be realized within it [80]. It is the moment when conditions align for decisive action.

Etymological and Metaphorical Roots The proposed concrete roots of *kairos* emphasize precision, timing, and consequence. One prominent etymological line points to the "vital/fatal

spot" targeted by an archer, suggesting a critical locus in space–time where a single, well-timed action yields a definitive result [468]. Another widely discussed metaphor links *kairos* to weaving: it is the fleeting opening in the shed of the loom through which the weaver must pass the shuttle; a moment too early or too late tangles the thread and mars the pattern. From such physical images, the term becomes a general metaphor for briefly open windows of action in practical life, discourse, and design [469, 470].

Kairos vs. Chronos: A Necessary Distinction To grasp *kairos*, it helps to contrast it with *chronos*. The latter underwrites "chronology" and "chronometer" and refers to sequential, measurable, quantitative time: the ticking of a clock, the duration of an event, the linear succession of moments [80]. *Kairos*, by contrast, is opportune, qualitative, and circumstantial—the fitting time to speak, act, or decide. In our usage below, *chronos* concerns metric duration and sequentiality, while *kairos* concerns the conditions under which processes become phase-appropriate. Put simply, *chronos* is the time one measures; *kairos* is the time one seizes [469, 470].

In Rhetoric. From the Sophists through Plato and Aristotle, *kairos* is a cornerstone of rhetorical theory, signifying the tact of timing. Even a logically impeccable, elegantly phrased argument fails if ill-timed. For classical rhetoricians—including Isocrates—efficacy hinges on recognizing and seizing the opportune moment, adapting the message to the concrete circumstances of the hearers [471, 472].

In Mythology and Iconography (the Deity Kairos). Classical culture personified *Kairos* as the daemon of opportunity, an emblem of the critical instant. Lysippos famously depicted him as a fleet youth with winged feet, a forelock (to be grasped as he approaches), a razor (signifying decision), and a bald pate behind—once the moment passes, there is nothing to hold [473]. Our usage of "the Deity Kairos" is strictly in this classical, allegorical register.

Definitions. Throughout this section we work on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ of Definition 10.1, with triad projectors $\Pi^{(a)}$ from Definition 10.2, the background-covariant TT map $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ in Definition 10.3, and the typed τ-kairos idempotents of Definition 10.4. All statements are taken on $\mathcal{H}_{\mathrm{phys}} = \operatorname{Ran} \mathsf{P}_{\mathrm{phys}}$.

Definition 10.13 (τ-kairos decomposition)**.** The τ-kairos field is the orthogonal sum

$$\boxed{\hat{\tau}_{\mu\nu} = S^{(b)}_{\mu\nu}[\hat{\varphi}_b] \;\oplus\; S^{(L)}_{\mu\nu}[\hat{\varphi}_L] \;\oplus\; \tilde{S}^{(p)}_{\mu\nu}[\tilde{\varphi}] \;\oplus\; \sum_{\lambda=\pm} A^{(\lambda)}_{\mu\nu}\big[\hat{K}^{(\tau)}_{\lambda}\big]} \tag{10.10}$$

with scalar amplitudes $\hat{\varphi}_b, \hat{\varphi}_L$, pseudoscalar $\tilde{\varphi}$, and helicity-resolved vector creators $\hat{K}^{(\tau)}_{\lambda}$. Each typed summand is $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$-null, cf. Definition 10.4.

Explanatory notes (objects and meaning).

- *Typed tensor-image maps.* $S^{(b)}_{\mu\nu}[\cdot]$, $S^{(L)}_{\mu\nu}[\cdot]$, $\tilde{S}^{(p)}_{\mu\nu}[\cdot]$, and $A^{(\lambda)}_{\mu\nu}[\cdot]$ are the τ-kairos typed maps of Definition 10.4 that take a scalar, pseudoscalar, or vector input (defined

on $\mathcal{H}_{\text{phys}}$) and return a τ-tensor on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$. They are idempotent/orthogonal at the level of types and preserve TT-nullity.

- $S^{(b)}_{\mu\nu}[\hat{\varphi}_b]$ *(scalar "breathing"/bulk piece).* The operator $\hat{\varphi}_b$ is a scalar (spin–0, CP-even) amplitude; $S^{(b)}_{\mu\nu}$ injects it into the τ-tensor with the "bulk/breathing" scalar pattern. This is part of the *Gaiaton* content (scalar sector).
- $S^{(L)}_{\mu\nu}[\hat{\varphi}_L]$ *(scalar longitudinal piece).* $\hat{\varphi}_L$ is a scalar (spin–0) longitudinal/coherence amplitude; $S^{(L)}_{\mu\nu}$ yields the corresponding τ-tensor image. Also Gaiaton (scalar sector).
- $\tilde{S}^{(p)}_{\mu\nu}[\tilde{\varphi}]$ *(pseudoscalar piece).* $\tilde{\varphi}$ is CP-odd (spin–0) and maps via the pseudoscalar typed channel $\tilde{S}^{(p)}_{\mu\nu}$ to a τ-tensor image. This is the *Gaiaton* pseudoscalar branch.
- $\sum_{\lambda=\pm} A^{(\lambda)}_{\mu\nu}[\hat{K}^{(\tau)}_{\lambda}]$ *(vector/helicity pieces).* $A^{(\lambda)}$ are the helicity projectors (spin–1, $\lambda = \pm 1$) in the τ-vector family; the operators $\hat{K}^{(\tau)}_{\lambda}$ are the helicity-resolved vector creators.[1] This entire block is the *Chrono–Kymon (CK)* content.
- *Orthogonal direct sum.* The symbol $\oplus$ denotes orthogonality with respect to the $\bar{\Sigma}$-weighted inner product on $\mathcal{H}_{\text{phys}}$: each typed image is orthogonal to the others; projector/idempotent relations ensure no double counting.
- *TT purity (no graviton in τ).* Each typed summand is annihilated by $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$, hence $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\hat{\tau}_{\mu\nu} = 0$. The spin–2 TT graviton lives in the *tensor* sector and is excluded from the τ-kairos by construction (cf. Definition 10.3).
- *Triad provenance (optional bookkeeping).* With $\Pi^{(a)}$ from Definition 10.2, the triad-resolved field $\hat{\tau}^{(a)}_{\mu\nu} := \Pi^{(a)}\hat{\tau}_{\mu\nu}$ decomposes as the same typed sum but with triad-extracted inputs $\hat{\varphi}^{(a)}_b, \hat{\varphi}^{(a)}_L, \tilde{\varphi}^{(a)}, \hat{K}^{(\tau)(a)}_{\lambda}$.

Remark 10.14 (Expanded forms of the τ–decomposition)**.** Starting from the compressed orthogonal sum in Definition 10.13, Eq. (10.10), there are two equivalent expanded views on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$.

(1) Projector expansion (typed projectors on the raw field). By completeness and orthogonality of the τ–typed idempotents (Lemma 10.8),

$$\hat{\tau}_{\mu\nu} = \left(S^{(b)}\hat{\tau}\right)_{\mu\nu} + \left(S^{(L)}\hat{\tau}\right)_{\mu\nu} + \left(\tilde{S}^{(p)}\hat{\tau}\right)_{\mu\nu} + \left(A^{(+)}\hat{\tau}\right)_{\mu\nu} + \left(A^{(-)}\hat{\tau}\right)_{\mu\nu}, \tag{10.11}$$

i.e.

$$\boxed{\hat{\tau}_{\mu\nu} = S^{(b)}{}_{\mu\nu}{}^{\rho\sigma}\,\hat{\tau}_{\rho\sigma} + S^{(L)}{}_{\mu\nu}{}^{\rho\sigma}\,\hat{\tau}_{\rho\sigma} + \tilde{S}^{(p)}{}_{\mu\nu}{}^{\rho\sigma}\,\hat{\tau}_{\rho\sigma} + \sum_{\lambda=\pm} A^{(\lambda)}{}_{\mu\nu}{}^{\rho\sigma}\,\hat{\tau}_{\rho\sigma}\,.} \tag{10.12}$$

Thus $\oplus$ in (10.10) is the orthogonal (direct) sum on the realized weave configuration, and each summand is TT–null by construction (Def. 10.4, Eq. (10.8); see also Lemma 10.17).

(2) Amplitude/polarization expansion (explicit extractors/creators). Since each typed projector $X \in \{S^{(b)}, S^{(L)}, \tilde{S}^{(p)}, A^{(\pm)}\}$ is an idempotent onto its carrier, choose bounded "extractors" and "creators" giving a continuous splitting on the realized weave configuration:

scalars/pseudoscalar:	$\epsilon^{(i)\,\rho\sigma},\ \iota^{(i)}_{\mu\nu}\quad (i \in \{b, L, p\}),$	$\epsilon^{(i)}\iota^{(i)} = \mathrm{id},\quad \iota^{(i)}\epsilon^{(i)} = S^{(i)};$
vectors:	$\epsilon^{(\lambda)\,\rho\sigma}{}_{\alpha},\ \iota^{(\lambda)\alpha}_{\mu\nu}\quad (\lambda = \pm),$	$\epsilon^{(\lambda)}\iota^{(\lambda)} = \mathrm{id},\quad \iota^{(\lambda)}\epsilon^{(\lambda)} = A^{(\lambda)}.$

[1] Parity on the τ-vector subspace swaps helicities: $\mathcal{P}_\tau A^{(+)}\mathcal{P}_\tau^{-1} = A^{(-)}$ and vice versa.

Define the typed amplitudes by extraction,

$$\hat{\varphi}_b := \epsilon^{(b)\,\rho\sigma}\hat{\tau}_{\rho\sigma}, \quad \hat{\varphi}_L := \epsilon^{(L)\,\rho\sigma}\hat{\tau}_{\rho\sigma}, \quad \tilde{\varphi} := \epsilon^{(p)\,\rho\sigma}\hat{\tau}_{\rho\sigma}, \quad \hat{K}^{(\tau)}_{\lambda,\alpha} := \epsilon^{(\lambda)\,\rho\sigma}{}_{\alpha}\hat{\tau}_{\rho\sigma},$$

and reconstruct:

$$\boxed{\hat{\tau}_{\mu\nu} = \underbrace{\iota^{(b)}_{\mu\nu}\,\hat{\varphi}_b}_{S^{(b)}_{\mu\nu}[\hat{\varphi}_b]} + \underbrace{\iota^{(L)}_{\mu\nu}\,\hat{\varphi}_L}_{S^{(L)}_{\mu\nu}[\hat{\varphi}_L]} + \underbrace{\iota^{(p)}_{\mu\nu}\,\tilde{\varphi}}_{\tilde{S}^{(p)}_{\mu\nu}[\tilde{\varphi}]} + \sum_{\lambda=\pm} \underbrace{\iota^{(\lambda)\alpha}_{\mu\nu}\,\hat{K}^{(\tau)}_{\lambda,\alpha}}_{A^{(\lambda)}_{\mu\nu}[\hat{K}^{(\tau)}_{\lambda}]}}. \tag{10.13}$$

Because $\iota^{(i)}\epsilon^{(i)} = S^{(i)}$ and $\iota^{(\lambda)}\epsilon^{(\lambda)} = A^{(\lambda)}$, (10.13) coincides with (10.11)–(10.12) and is *unique up to* basis changes inside each typed range (Lemma 10.18). Each term is TT–annihilated (Def. 10.4; Lemma 10.17), and all maps commute with triads (Prop. 10.9).

Triad compatibility. Applying the triad sandwich $\Pi^{(a)}$ ($a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$; Definition 10.2) to (10.11)–(10.13) gives the same formulas with $\hat{\tau}$ replaced by $\hat{\tau}^{(a)} := \Pi^{(a)}\hat{\tau}$, since $[\Pi^{(a)}, X] = 0$ for all typed X (Prop. 10.9).

Definition 10.15 (Triad-resolved τ-kairos components)**.** For $a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$, define

$$\hat{\tau}^{(a)}_{\mu\nu} := \Pi^{(a)}\,\hat{\tau}_{\mu\nu} = \sum_{X \in \{S^{(b)}, S^{(L)}, \tilde{S}^{(p)}, A^{(+)}, A^{(-)}\}} \left(\Pi^{(a)} X\right)_{\mu\nu}, \qquad \hat{\tau}_{\mu\nu} = \sum_{a} \hat{\tau}^{(a)}_{\mu\nu}, \tag{10.14}$$

where the sum uses the orthogonal decomposition of Lemma 10.8.

Definition 10.16 (Parity and helicity diagnostics on the τ-block)**.** Let $\mathcal{P}_\tau$ be the formal parity on the scalar/pseudoscalar subspace so that $\mathcal{P}_\tau S^{(b)}\mathcal{P}_\tau^{-1} = S^{(b)}$, $\mathcal{P}_\tau S^{(L)}\mathcal{P}_\tau^{-1} = S^{(L)}$, $\mathcal{P}_\tau \tilde{S}^{(p)}\mathcal{P}_\tau^{-1} = -\tilde{S}^{(p)}$. On the vector channel, $\mathcal{P}_\tau A^{(+)}\mathcal{P}_\tau^{-1} = A^{(-)}$, $\mathcal{P}_\tau A^{(-)}\mathcal{P}_\tau^{-1} = A^{(+)}$. Define the (law/state) *parity-asymmetry audit*

$$\Delta_P^{[\mathrm{law/state}]} := \Lambda^{[A^{(+)}]}_{\mathrm{law/state}} - \Lambda^{[A^{(-)}]}_{\mathrm{law/state}}, \tag{10.15}$$

to be evaluated with the typed witnesses of Definitions 10.80 and 10.82.

Structural facts.

Lemma 10.17 (TT-purity of the τ–kairos)**.** $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\hat{\tau}_{\mu\nu} = 0$ *and* $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\hat{\tau}^{(a)}_{\mu\nu} = 0$ *for each* $a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$.

Proof. Work on the realized weave configuration $\mathcal{I}^{[\mathcal{J}]}$ (Def. Definition 10.1). On the τ–typed carrier, the orthogonal idempotents $S^{(b)}, S^{(L)}, \tilde{S}^{(p)}, A^{(+)}, A^{(-)}$ (Def. Definition 10.4) satisfy the completeness/orthogonality relations

$$\sum_{i \in \{b,L,p\}} S^{(i)} + \sum_{\lambda=\pm} A^{(\lambda)} = \mathbf{1}_\tau, \qquad S^{(i)}S^{(j)} = \delta_{ij}S^{(i)}, \quad A^{(\lambda)}A^{(\lambda')} = \delta_{\lambda\lambda'}A^{(\lambda)}, \quad S^{(i)}A^{(\lambda)} = 0,$$

as recorded in Lemma 10.8. Hence the (typed) τ–kairos observable decomposes as

$$\hat{\tau}_{\mu\nu} = \sum_{i \in \{b,L,p\}} S^{(i)}\hat{\tau}_{\mu\nu} + \sum_{\lambda=\pm} A^{(\lambda)}\hat{\tau}_{\mu\nu}.$$

Step 1: TT–nullity on each typed summand. By the TT–null constraint built into the τ–typed projectors (Def. Definition 10.4, Eq. (10.8)),

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} X = 0 \quad \text{for each } X \in \{S^{(b)}, S^{(L)}, \tilde{S}^{(p)}, A^{(+)}, A^{(-)}\}.$$

Therefore,

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \hat{\tau}_{\mu\nu} = \sum_i \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} S^{(i)} \hat{\tau}_{\mu\nu} + \sum_\lambda \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} A^{(\lambda)} \hat{\tau}_{\mu\nu} = 0.$$

This uses linearity and the fact that the TT projector is the background–covariant pseudodifferential lift that annihilates traces and $\bar{\Sigma}$–divergences (Def. Definition 10.3), while the τ–typed images live in scalar/vector carriers orthogonal to TT (Ch. 10 typed layer Definition 9.116).

Step 2: Triad-resolved TT–nullity. Define the triad-resolved observable $\hat{\tau}_{\mu\nu}^{(a)} := \Pi^{(a)} \hat{\tau}_{\mu\nu}$ with $a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$ (triads Definition 10.2). By commutation of typed/TT idempotents with triads on the realized weave configuration (Prop. Proposition 10.9),

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \hat{\tau}_{\mu\nu}^{(a)} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \Pi^{(a)} \hat{\tau}_{\mu\nu} = \Pi^{(a)} \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \hat{\tau}_{\mu\nu} = \Pi^{(a)} \cdot 0 = 0.$$

Realized weave configuration and retune stability. All operators involved commute with the band tags and remain admissible under audit moves (Prop. Proposition 10.10, Riesz measurability Proposition 10.149), so the identities hold throughout admissible retunes and arena switches. Finally, the same TT–nullity propagates through any τ–involving mixing channel (Lemma 10.123), preventing induced TT content (Cor. 10.125). □

Lemma 10.18 (Uniqueness of the typed decomposition)**.** *Work on the realized weave configuration $\mathcal{I}^{[\mathcal{S}]}$ (Def. Definition 10.1). Given the τ–typed projector algebra of Lemma 10.8 and the typed layer structure (Ch. 10: Definition 9.116, instantiated in Definition 10.4), the representation*

$$\hat{\tau} = \sum_{i \in \{b,L,p\}} S^{(i)} \hat{\tau} + \sum_{\lambda = \pm} A^{(\lambda)} \hat{\tau} \tag{10.16}$$

is unique up to *reparameterizations internal to each typed range (i.e. changes of basis/redefinitions of scalar/pseudoscalar amplitudes and vector creators within* $\operatorname{Ran} S^{(i)}$ *and* $\operatorname{Ran} A^{(\lambda)}$*). Moreover, this uniqueness is triad–compatible: the same statement holds for $\hat{\tau}^{(a)} := \Pi^{(a)} \hat{\tau}$ with $a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$ (triads Definition 10.2).*

Proof. By Lemma 10.8, the family

$$\mathcal{P}_\tau := \{S^{(b)}, S^{(L)}, \tilde{S}^{(p)}, A^{(+)}, A^{(-)}\}$$

consists of $\bar{\Sigma}$–Hermitian idempotents that are pairwise orthogonal and complete on the τ–typed carrier:

$$X^2 = X, \quad XY = 0 \ (X \neq Y), \quad \sum_{X \in \mathcal{P}_\tau} X = \mathbf{1}_\tau.$$

Therefore any $\hat{\tau}$ admits the canonical decomposition (10.10).

Uniqueness at the projector level. Suppose there are two typed decompositions

$$\hat{\tau} = \sum_{X \in \mathcal{P}_\tau} \hat{\tau}_X^{(1)} = \sum_{X \in \mathcal{P}_\tau} \hat{\tau}_X^{(2)}, \qquad \hat{\tau}_X^{(k)} \in \operatorname{Ran} X.$$

Apply any $Y \in \mathcal{P}_\tau$ to both sides. Using orthogonality $YX = \delta_{YX}Y$,

$$Y\hat{\tau} = \sum_X Y\hat{\tau}_X^{(1)} = Y\hat{\tau}_Y^{(1)}, \qquad Y\hat{\tau} = \sum_X Y\hat{\tau}_X^{(2)} = Y\hat{\tau}_Y^{(2)}.$$

Hence $Y\hat{\tau}_Y^{(1)} = Y\hat{\tau}_Y^{(2)}$, but each side already lies in $\operatorname{Ran} Y$ and Y acts as the identity on that range; therefore $\hat{\tau}_Y^{(1)} = \hat{\tau}_Y^{(2)}$. Since Y was arbitrary, all typed components coincide and the decomposition is unique.

"Up to redefinitions" inside each range. Within a fixed typed subspace $\operatorname{Ran} X$ the choice of coordinates (e.g. scalar/pseudoscalar amplitude bases or vector creation bases) is not unique; any invertible change of basis $U : \operatorname{Ran} X \to \operatorname{Ran} X$ with $UX = XU = U$ gives a different internal parameterization of the *same* component $X\hat{\tau}$. Thus the uniqueness is exactly modulo such internal reparameterizations.

Triad compatibility. By Proposition 10.9, $[\Pi^{(a)}, X] = 0$ for every $X \in \mathcal{P}_\tau$ and $a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$. Applying $\Pi^{(a)}$ to (10.10) and using commutation yields

$$\hat{\tau}^{(a)} = \Pi^{(a)}\hat{\tau} = \sum_{X \in \mathcal{P}_\tau} X\,\Pi^{(a)}\hat{\tau} = \sum_{X \in \mathcal{P}_\tau} X\,\hat{\tau}^{(a)},$$

which is the same projector decomposition but for the triad–resolved observable. Repeating the uniqueness argument above (apply $Y \in \mathcal{P}_\tau$) gives uniqueness of the typed components of $\hat{\tau}^{(a)}$ as well, again up to internal reparameterizations in each $\operatorname{Ran} X$.

All statements are made on the realized weave configuration $\mathcal{I}^{[\mathcal{S}]}$ and are stable under admissible retunes by Proposition 10.10; Riesz measurability Proposition 10.149 and typed stability (Ch. 10 Definition 9.116) ensure the decomposition persists along audit moves (Def. Definition 10.101). □

Meaning and importance. Lemma 10.18 says that once the τ–typed projector algebra is fixed (Lemma 10.8) and the typed layer is installed (Ch. 10: Definition 9.116, instantiated in Definition 10.4), any observable $\hat{\tau}$ admits a *canonical* direct-sum decomposition into breathing, longitudinal, pseudoscalar, and helicity-resolved vector parts, cf. (10.16). The word "canonical'' here is precise: orthogonal, complete, $\bar{\Sigma}$–Hermitian idempotents select mutually exclusive carrier spaces whose sum is the whole τ-typed block, so there is no ambiguity about which piece of $\hat{\tau}$ belongs to which type. The only latitude that remains is a harmless *change of coordinates within* each range (e.g., scalar amplitude redefinitions or a different helicity basis), which does not alter the typed component itself. This uniqueness is crucial operationally: it guarantees that downstream witnesses, selection rules, and parity/helicity/CP diagnostics are insensitive to the order in which typing is performed and that no spurious mixing is introduced when we later combine typing with spectral bookkeeping (triads; Definition 10.2). In short, the typed split is a rigid, basis-independent statement about $\hat{\tau}$ on the realized weave configuration $\mathcal{I}^{[\mathcal{S}]}$ (Definition 10.1).

Why the proof matters. The proof formalizes two standard functional-analytic mechanisms. First, orthogonal idempotents yield a unique projection decomposition: applying any projector Y from the family $\mathcal{P}_\tau$ to two candidate sums immediately identifies and equalizes the Y–component, and repeating for all Y pins down every typed piece uniquely. Second, *triad*

compatibility follows from commutation with the Riesz spectral projectors that define the triad (Definition 10.7 and Proposition 10.9): since typed idempotents commute with the band-limited kinetic blocks on the realized weave configuration (typed stability; Ch. 10 Definition 9.116), they commute with the resolvents and hence with the contour integrals that produce the triad projectors; this is exactly the holomorphic functional-calculus logic [115, 253]. Therefore the same uniqueness holds for each triad-resolved observable $\hat{\tau}^{(a)}$, ensuring that spectral triage (spec/vac/corr) and typing can be interleaved without ambiguity or loss of information—one of the key algebraic guarantees behind our weave-level diagnostics.

Proposition 10.19 (BRST/Nielsen invariance and band stability for τ–correlators)**.** *Work on the realized weave configuration $\mathcal{I}^{[\mathcal{S}]}$ (Def. Definition 10.1). Let $\mathcal{J}_\tau$ be any finite composition of the triad maps $\Pi^{(a)}$ ($a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$; Definition 10.2), the TT projector $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ (Definition 10.3), and the τ–typed idempotents $S^{(b)}, S^{(L)}, \tilde{S}^{(p)}, A^{(\pm)}$ (Definition 10.4). Then every correlator on $\mathcal{H}_{\mathrm{phys}} = \mathrm{Ran}\,\mathsf{P}_{\mathrm{phys}}$ of $\mathcal{J}_\tau$–projected τ–observables (e.g. $\langle \mathcal{J}_\tau\, \hat{\tau}_{\mu\nu} \rangle$) is:*

(i) gauge–parameter independent *(Nielsen flat) on shell; and*

(ii) stable under admissible retunes $(P_x, K_{\mathrm{IR}}, \mathrm{Loc})$ *within the certified class (continuous and uniformly bounded along admissible paths; Def. Definition 10.101).*

Proof. 1. **Gauge–parameter independence (Nielsen flatness).** Let ξ denote any gauge parameter in the gauge–fixed kinetic operators. On $\mathcal{H}_{\mathrm{phys}}$, the embedded Nielsen identity (Ch. 8; cf. Prop. Proposition 7.174) yields, for sufficiently regular $\mathcal{A}$,

$$\partial_\xi \langle \mathcal{A} \rangle_{\mathcal{H}_{\mathrm{phys}}} = \Big\langle [\hat{Q}_{\mathrm{BRST}}, \mathcal{K}_\xi]\, \mathcal{A} \Big\rangle_{\mathcal{H}_{\mathrm{phys}}} + \Big\langle \mathrm{EOM}_\xi(\mathcal{A}) \Big\rangle_{\mathcal{H}_{\mathrm{phys}}}. \tag{10.17}$$

Here $\hat{Q}^2_{\mathrm{BRST}} = 0$, and the EOM_ξ term vanishes on shell. Choose $\mathcal{A} = \mathcal{J}_\tau\, \hat{\tau}_{\mu\nu}$, where $\mathcal{J}_\tau$ is a composition of idempotents listed in the statement. By typed stability and commutation on the realized weave configuration (Ch. 10 typed layer Definition 9.116 and Proposition 10.9), each factor in $\mathcal{J}_\tau$ is BRST–closed and commutes with the band machinery on $\mathcal{H}_{\mathrm{phys}}$. Therefore the BRST–exact contribution has zero expectation and the EOM term vanishes on shell, giving $\partial_\xi \langle \mathcal{J}_\tau\, \hat{\tau}_{\mu\nu} \rangle_{\mathcal{H}_{\mathrm{phys}}} = 0$. Hence gauge–parameter independence.

2. Stability under admissible retunes. Consider an admissible path (Def. Definition 10.101):

$$t \mapsto \Big(P_x(t),\, K_{\mathrm{IR}}^{(\ell(t))},\, \mathrm{Loc}(t) \Big)$$

Triad maps $\Pi^{(a)}$ are realized by Riesz projectors of the band–limited kinetic operators on certified contours (selectors Definition 10.7); by Riesz measurability Proposition 10.149, the map $t \mapsto \Pi^{(a)}(t)$ is Bochner–measurable and norm–continuous along admissible paths. The TT projector $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ (Definition 10.3) and the τ–typed idempotents (Definition 10.4) are fixed $\bar{\Sigma}$–Hermitian idempotents that commute with the band tags and triads on the realized weave configuration (Proposition 10.9). Hence $t \mapsto \mathcal{J}_\tau(t)$ is norm–continuous on $\mathcal{H}_{\mathrm{phys}}$.
Correlation functionals built from $\mathcal{J}_\tau(t)\hat{\tau}_{\mu\nu}$ are continuous in the operator norm, and remain uniformly bounded along admissible paths because admissible band–invariant

norms are equivalent and contractive under IR tightening (Def. Definition 10.143, Prop. Proposition 10.147). Arena switches preserve the certified spectral set and associated projectors (Lemma 10.137), so no discontinuity is introduced by bulk $\leftrightarrow$ boundary. Therefore $\langle \mathcal{J}_\tau(t)\,\hat{\tau}_{\mu\nu}\rangle_{\mathcal{H}_{\rm phys}}$ varies continuously and stays uniformly bounded along any admissible retune, proving stability.

Alternative route. The result also follows immediately by specializing Proposition 10.10 to compositions that only involve $\Pi^{(a)}$, $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$, and the τ–typed idempotents. □

Corollary 10.20 (No contamination of the graviton channel)**.** *For any symmetric 2-tensor perturbation $\delta\hat{g}_{\mu\nu}$, the graviton field $\hat{h}_{\mu\nu}$ obtained via eq. (10.28) is independent of $\hat{\tau}_{\mu\nu}$:* $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}(\delta\hat{g}_{\mu\nu}+\hat{\tau}_{\mu\nu}) = \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\delta\hat{g}_{\mu\nu}$.

Proof. By Lemma 10.17, $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\hat{\tau} = 0$; sandwiching by $\mathsf{P}_{\rm phys}$ and spectral/triad maps (which commute with $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$) preserves this equality in eq. (10.28). □

Operational extraction and certification.

Definition 10.21 (Typed τ-kairos witnesses and visibilities)**.** For each admitted type $X \in \{S^{(b)}, S^{(L)}, \tilde{S}^{(p)}, A^{(+)}, A^{(-)}\}$ and sector $\mathscr{S}$, define the consolidated witness $\mathbf{W}^{[\mathscr{S};X]}$ and the law/state visibilities $\Lambda^{[\mathscr{S};X]}_{\text{law/state}}$ as in Definitions 10.80 and 10.82, evaluated on $\mathcal{H}_{\rm phys}$ with instrumentation $\mathcal{I}^{[\mathscr{S}]}$.

Proposition 10.22 (Well-posed Parasymmetric TLO for τ-types)**.** *For each X as above, the Parasymmetric TLO triple* (Type, Location, Onset) *of Definition 10.85 and Proposition 10.90 exists and is unique on the certified band. Moreover, the Onset $r^\star[X]$ is stable under admissible retuning $r \mapsto r + \delta r$ and tightening of $K_{\rm IR}$.*

Proof (sketch). Existence/uniqueness of the Type layer follows from Lemma 10.18. Location is determined by the arena in $\mathcal{I}^{[\mathscr{S}]}$. Onset stability is the monotone-meters property inherited from band covariance and BRST/Nielsen flatness (Proposition 10.10); tightening $K_{\rm IR}$ cannot create spurious visibility plateaus for admissible types. □

Remark 10.23 (Parity bookkeeping at the scalar level)**.** The scalar subspace decomposes canonically into even/odd parity via $\Pi^{(e)}_{\rm sc} = S^{(b)} + S^{(L)}$, $\Pi^{(o)}_{\rm sc} = \tilde{S}^{(p)}$. Vector parity is reported via Δ_P in eq. (10.15) while keeping helicity $A^{(\pm)}$ as primitive types.

Remark 10.24 (Chrono–Kymon identification)**.** The Chrono–Kymon is the τ-vector content captured by $A^{(\lambda)}$ ($\lambda = \pm$). Its extraction on $\mathcal{H}_{\rm phys}$ uses the same realized weave configuration and typed witnesses, and is automatically TT-null by Lemma 10.17.

10.4 Chronogeometric Origin and Certified Construction of the τ–kairos

Work on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ of Definition 10.1 with physical space $\mathcal{H}_{\rm phys} = \operatorname{Ran}\mathsf{P}_{\rm phys}$, triad projectors $\Pi^{(a)}$ from Definition 10.2, the TT projector $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ from

Definition 10.3, and the typed families of Definition 10.4. The goal is to formalize the emergence of the τ–kairos from a constitutive chronogeometric observable and to place it within the canonical typed/Riesz framework (Ch. 6–10).

Constitutive origin and linear response. Let the chronogeometric observable be a constitutive functional of the Gaiamiti and Tantrakyma substrates,

$$\Sigma_{\mu\nu}[\Phi,\Theta] = g_{\mu\nu}[\Phi] \;+\; t_{\mu\nu}[\Phi,\Theta]\,, \tag{10.18}$$

with background $(\bar{\Phi},\bar{\Theta})$ and fluctuation $\delta\Sigma_{\mu\nu}$. Linearization gives the chain–rule split

$$\delta\Sigma_{\mu\nu} = (D_\Phi g)_{\mu\nu}\cdot\,\delta\Phi + (D_\Phi t)_{\mu\nu}\cdot\,\delta\Phi + (D_\Theta t)_{\mu\nu}\cdot\,\delta\Theta\,. \tag{10.19}$$

Certified TT isolation and definition of the τ–kairos (expanded steps). Step 0: Constitutive linearization. From the constitutive map $\Sigma_{\mu\nu}[\Phi,\Theta] = g_{\mu\nu}[\Phi]+t_{\mu\nu}[\Phi,\Theta]$ one has, about $(\bar{\Phi},\bar{\Theta})$,

$$\delta\Sigma_{\mu\nu} = (D_\Phi g)_{\mu\nu}\cdot\,\delta\Phi + (D_\Phi t)_{\mu\nu}\cdot\,\delta\Phi + (D_\Theta t)_{\mu\nu}\cdot\,\delta\Theta\,. \tag{10.20}$$

Step 1: Insert the TT/complement decomposition. With the certified TT projector $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ of Definition 10.3, write the identity $\mathrm{Id} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} + (\mathrm{Id} - \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]})$ and apply it to $\delta\hat{\Sigma}_{\mu\nu}$:

$$\hat{h}_{\mu\nu} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{\Sigma}_{\mu\nu} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\Big[(D_\Phi g + D_\Phi t)_{\mu\nu}\cdot\,\delta\hat{\Phi}\Big] + \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\Big[(D_\Theta t)_{\mu\nu}\cdot\,\delta\hat{\Theta}\Big], \tag{10.21a}$$

$$\hat{\tau}_{\mu\nu} := \Big(\mathrm{Id} - \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\Big)\,\delta\hat{\Sigma}_{\mu\nu}, \qquad \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\hat{\tau}_{\mu\nu} = 0 \;\; \text{(idempotence)}. \tag{10.21b}$$

Step 2: Why the Θ–term carries no TT content. By typed admission (Definition 10.4) the non-tensor families $\{S^{(b)}, S^{(L)}, \tilde{S}^{(p)}, A^{(\pm)}\}$ are *TT-null*:

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\Pi^{[X]} = 0 = \Pi^{[X]}\,\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}, \qquad X \in \{S^{(b)}, S^{(L)}, \tilde{S}^{(p)}, A^{(\pm)}\}. \tag{10.22}$$

On the certified band (Ch. 10), $(D_\Theta t)_{\mu\nu}\cdot\,\delta\Theta$ lies in the closed span of these TT-null typed images, hence

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\Big[(D_\Theta t)_{\mu\nu}\cdot\,\delta\Theta\Big] = 0. \tag{10.23}$$

Combining (10.21a) and (10.23) reproduces the displayed identity in (10.21a) of the main text.

Step 3: Define $\hat{\tau}$ and check TT-nullity transparently. Equation (10.21b) yields, by idempotence of $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$,

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\hat{\tau}_{\mu\nu} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\Big(\mathrm{Id} - \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\Big)\delta\Sigma_{\mu\nu} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\delta\Sigma_{\mu\nu} - \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\delta\Sigma_{\mu\nu} = 0.$$

Thus $\hat{\tau}_{\mu\nu}$ is TT-null by construction.

Step 4: Typed resolution of the non–TT residual. Introduce the typed projectors $\Pi^{[X]}$ associated with the τ-families of Definition 10.4. On the certified band, $\sum_X \Pi^{[X]}$ acts as the identity on the TT-complement, and each image admits a *typed representative* via the corresponding map ($X \in \{S^{(b)}, S^{(L)}, \tilde{S}^{(p)}, A^{(\pm)}\}$):

$$\Pi^{[S^{(b)}]}\hat{\tau} = S^{(b)}_{\mu\nu}[\hat{\varphi}_b], \quad \Pi^{[S^{(L)}]}\hat{\tau} = S^{(L)}_{\mu\nu}[\hat{\varphi}_L], \quad \Pi^{[\tilde{S}^{(p)}]}\hat{\tau} = \tilde{S}^{(p)}_{\mu\nu}[\tilde{\varphi}], \quad \Pi^{[A^{(\lambda)}]}\hat{\tau} = A^{(\lambda)}_{\mu\nu}[\hat{K}^{(\tau)}_{\lambda}]. \tag{10.24}$$

The amplitudes $\hat{\varphi}_b, \hat{\varphi}_L, \tilde{\varphi}, \hat{K}_\lambda^{(\tau)}$ are obtained by the certified-band Riesz selectors and right-inverses guaranteed by the typed construction in Ch. 10 (no off-band components enter). Summing (10.24) over all types recovers the τ-kairos decomposition

$$\hat{\tau}_{\mu\nu} = S_{\mu\nu}^{(b)}\big[\hat{\varphi}_b\big] \;\oplus\; S_{\mu\nu}^{(L)}\big[\hat{\varphi}_L\big] \;\oplus\; \tilde{S}_{\mu\nu}^{(p)}\big[\tilde{\varphi}\big] \;\oplus\; \sum_{\lambda=\pm} A_{\mu\nu}^{(\lambda)}\big[\hat{K}_\lambda^{(\tau)}\big] \quad \text{(Definition 10.13).} \tag{10.25}$$

Step 5: Triad provenance commutes with typing. With $\Pi^{(a)}$ from Definition 10.2 and the commutation $[\Pi^{(a)}, \Pi^{[X]}] = 0$ (Proposition 10.9), one has

$$\hat{\tau}_{\mu\nu}^{(a)} := \Pi^{(a)}\hat{\tau}_{\mu\nu} = \sum_X \Pi^{(a)}\Pi^{[X]}\hat{\tau} = \sum_X \Pi^{[X]}\Pi^{(a)}\hat{\tau}, \qquad a \in \{\text{spec}, \text{vac}, \text{corr}\}, \tag{10.26}$$

and each typed block in (10.25) acquires its triad-resolved input $\hat{\varphi}_b^{(a)}, \hat{\varphi}_L^{(a)}, \tilde{\varphi}^{(a)}, \hat{K}_\lambda^{(\tau)(a)}$ accordingly.

Step 6: CK/vector and scalar spectral selection (existence of inputs). Existence/uniqueness of the inputs in (10.24) on the certified band follows from the Riesz projectors: for the vector channel use $\mathsf{P}_{\text{spec}}^{(1)}$ and $\mathcal{P}_{\text{CK},\lambda}^{(1)}$ (Definitions 10.56 and 10.57); for the scalar channel use the spin–0 selector $\mathsf{P}_{\text{spec}}^{(0)}$ (cf. the scalar construction). Additivity of Riesz projectors implies automatic extension when new certified spectral branches appear (cf. Lemma 10.62).

Equations (10.21a)–(10.25) now follow *mechanically* from the constitutive linearization (10.20), TT idempotence, typed TT-null admission (10.22), and the certified-band typed/Riesz reconstruction in Ch. 10, with triad provenance attached by (10.26).

Triad provenance and orthogonality. With the triad projectors $\Pi^{(a)}$ from Definition 10.2, define

$$\hat{\tau}_{\mu\nu}^{(a)} := \Pi^{(a)}\,\hat{\tau}_{\mu\nu} = \; S_{\mu\nu}^{(b)}\big[\hat{\varphi}_b^{(a)}\big] \oplus S_{\mu\nu}^{(L)}\big[\hat{\varphi}_L^{(a)}\big] \oplus \tilde{S}_{\mu\nu}^{(p)}\big[\tilde{\varphi}^{(a)}\big] \oplus \sum_{\lambda=\pm} A_{\mu\nu}^{(\lambda)}\big[\hat{K}_\lambda^{(\tau)(a)}\big], \tag{10.27}$$

with $\Pi_{\bar{\Sigma}}^{[\text{TT}]}\hat{\tau}_{\mu\nu}^{(a)} = 0$. Typed and triad pieces are mutually orthogonal by the idempotent algebra (cf. Lemma 10.8 and Proposition 10.9).

Certified-band selection (scalar and vector). Scalar and vector content on the certified band are isolated by spin–0/1 Riesz selectors. For the vector (CK) block, Definition 10.56 defines $\mathsf{P}_{\text{spec}}^{(1)}$ and the CK projector $\mathcal{P}_{\text{CK},\lambda}^{(1)}$ (Definition 10.57); the scalar (gaiaton) block admits the analogous $\mathsf{P}_{\text{spec}}^{(0)}$ and $\Pi_\tau^{(\text{spec};0)}$ (cf. the scalar construction in Sec. 10.8). Riesz additivity implies automatic inclusion of new certified spectral branches (future-proofing), cf. Lemma 10.62.

Meters, witnesses, and diagnostics. Typed, triad-resolved visibilities follow Definitions 10.80 and 10.82. For the vector CK channel, use $\Lambda_{\text{law/state}}^{[\tau\text{-vec};A^{(\lambda)}]}$ (Definition 10.64) and the parity diagnostic $\Delta_{P,\Lambda}^{(1)}$ (Definition 10.58). For the scalar channel (gaiaton), use the CP-resolved visibilities and, where needed, the CP asymmetry $\Delta_{\text{CP},\Lambda}^{(0)}$ (cf. the scalar definitions). All meters are evaluated on $\mathcal{H}_{\text{phys}}$ with the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$.

Invariance, stability, and TLO onset. Any correlator formed from $\hat{\tau}$-typed blocks with finitely many insertions of the typed projectors and certified Riesz selectors is BRST/Nielsen-flat and stable under admissible retunings of (P_x, K_{IR}) (Proposition 10.10; see also Proposition 10.61 for CK). For each admitted type, the Parasymmetric TLO triple (Type, Location, Onset) exists and is unique on the certified band (Definition 10.85 and Proposition 10.90); onsets $r^\star[\cdot]$ inherit the same retuning stability.

Mixing, whitening, and reporting. Non–TT mixing (including intra-scalar and CK helicity mixing) is quantified by typed mixing norms and resolved by the $\bar{\Sigma}$–weighted operator-generalized eigen-operator pair eigenproblem with whitening (Definitions 10.116 and 10.117). The outcome is an orthogonal set of eigen-channels inside the τ block, each carrying triad provenance and ledgered power. TT mixing does not occur: $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ annihilates all τ-typed insertions.

Interfaces with Gaiamiti (TT) and Tantrakyma ($\Theta\Theta$). By (10.21a)–(10.21b), the TT graviton (Gaiamiti tensor) is isolated entirely in the TT channel and is orthogonal to τ (no TT contamination). Higgs/radial kymon content is identified in the $\Theta\Theta$ (tantrakyma) sector with CP resolution and TLO guarantees (Section 10.6); comparisons with τ-typed readouts proceed through the common witness/ledger interface, not by identification.

Optional representatives (notation only). First-order representatives may be introduced for interpretation without changing the typed machinery: scalars $\{\hat{\varphi}_b, \hat{\varphi}_L, \tilde{\varphi}\}$ for the scalar/pseudoscalar block and a vector $K_\mu^{(\tau)}$ for CK, with $\mathcal{F}_{\mu\nu}^{(\tau)} = 2\bar{\nabla}_{[\mu} K_{\nu]}^{(\tau)}$ (cf. Definition 10.57). These objects are proxies for typed correlator content and do not modify TT purity.

Processing equivalence. The constitutive split (10.19), followed by TT isolation (10.21a) and typed/Riesz extraction, is equivalent (at the level of reported types, triad ledgers, and TLO onsets) to proceeding directly with the typed-correlator pipeline on the realized weave configuration; this is a consequence of projector idempotence and the realized weave configuration-level invariance statements referenced above (cf. Ch. 10 and Proposition 10.10).

Summary. The τ–kairos is the TT-null, typed projection of the linear response of the chronogeometric observable (10.18), organized as the orthogonal sum (10.25) with triad provenance (10.27). All readouts are certified by the same typed/Riesz infrastructure as in Ch. 6–10: they inherit BRST/Nielsen flatness, admissible-retuning stability, TLO well-posedness, and standardized meters/diagnostics. The TT (graviton) and tantrakyma (Higgs/radial kymon) sectors remain orthogonal and are interfaced with τ only via the shared witness/ledger scaffold.

10.5 The Graviton in the Gaiamiti (Geometry) Sector

We work on the realized weave configuration $\mathcal{I}^{[\mathscr{P}]}$ of Definition 10.1 in the $\Phi\Phi$ (gaiamiti) sector, with triad projectors $\Pi^{(a)}$ from Definition 10.2, the background-covariant TT map $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ of



Figure 10.1: Constitutive map and channel-resolved sourcing. Gaiamiti Φ feeds the metric TT graviton and the scalar part of $\hat{\tau}_{\mu\nu}$ (via apeiron), while Tantrakyma Θ feeds the vector part (Chrono–Kymon). **Legend:** (1) TT radiation lives only in the metric channel: $\hat{h}_{\mu\nu} = \mathcal{P}^{(2)}[\delta\hat{g}_{\mu\nu}] = \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{g}_{\mu\nu}$; (2) Tau–kairos is TT-null: $\mathcal{P}^{(2)}[\hat{\tau}_{\mu\nu}] = 0$, so no spin-2 GW; (3) $\hat{\tau}_{\mu\nu} = S_{\mu\nu}[\hat{\varphi}_A] \oplus A_{\mu\nu}[\hat{K}^{(\tau)}]$ combines scalar (from Φ via apeiron) and vector (from Θ via Chrono–Kymon).

Definition 10.3, and the spin-2 Riesz selector $\mathsf{P}^{(2)}_{\mathrm{spec}}$ from Definition 10.7. All statements are taken on $\mathcal{H}_{\mathrm{phys}} = \mathrm{Ran}\,\mathsf{P}_{\mathrm{phys}}$.

Definition 10.25 (Graviton projector (typed TT, triad-resolved, future-proof))**.** Let $K_{\Phi\Phi}$ be the band-limited, gauge-fixed linearized kinetic operator in the $\Phi\Phi$ block. The *graviton projector* is

$$\boxed{\mathcal{P}^{(2)}_{\mathrm{grav}} := \mathsf{P}_{\mathrm{phys}}\,\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\Big(\,\mathsf{P}^{(2)}_{\mathrm{spec}} \;\oplus\; \Pi^{(\mathrm{vac})} \;\oplus\; \Pi^{(\mathrm{corr})}\,\Big)\,\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\mathsf{P}_{\mathrm{phys}}} \tag{10.28}$$

with $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ the background-covariant TT projector (Definition 10.3), $\mathsf{P}^{(2)}_{\mathrm{spec}}$ the spin-2 Riesz selector (Definition 10.7), and $\Pi^{(a)}$ the triad projectors ($a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$; Definition 10.2). The *graviton field* is obtained from any symmetric perturbation $\delta\hat{g}_{\mu\nu}$ by

$$\boxed{\hat{h}_{\mu\nu} := \mathcal{P}^{(2)}_{\mathrm{grav}}\,\delta\hat{g}_{\mu\nu} = \sum_{a\in\{\mathrm{spec,vac,corr}\}} \hat{h}^{(a)}_{\mu\nu}, \qquad \hat{h}^{(a)}_{\mu\nu} := \mathsf{P}_{\mathrm{phys}}\,\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\Pi^{(a)}\,\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\mathsf{P}_{\mathrm{phys}}\,\delta\hat{g}_{\mu\nu}.} \tag{10.29}$$

House notation (TT identification and helicity). Consistently with standard practice [43, 7, 14],

we *identify the graviton channel by its TT content*:

$$\hat{h}^{\mathrm{TT}}_{\mu\nu} \;:=\; \left(\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{g}\right)_{\mu\nu}, \tag{10.30}$$

which coincides with the spectral spin-2 component of (10.29) by commutation of typed/Riesz/triad projectors (Proposition 10.9 and Definition 10.7). When a helicity resolution is needed, one may write

$$\hat{h}^{(a)}_{\mu\nu} = \sum_{\lambda=\pm 2} \epsilon^{(\lambda)}_{\mu\nu}[\bar{\Sigma}]\;\hat{a}^{(a)}_{\lambda} \;+\; \text{h.c.}, \tag{10.31}$$

where $\epsilon^{(\lambda)}_{\mu\nu}[\bar{\Sigma}]$ are $\bar{\Sigma}$-transverse–traceless polarization tensors ($\bar{\Sigma}^{\mu\nu}\epsilon^{(\lambda)}_{\mu\nu} = 0$, $\nabla^{(\mathrm{rel})\mu}\epsilon^{(\lambda)}_{\mu\nu} = 0$) normalized on the realized weave configuration, and $\hat{a}^{(a)}_{\lambda}$ are the corresponding helicity mode operators [7, 14].

Conceptual role of the projector. The graviton projector $\mathcal{P}^{(2)}_{\mathrm{grav}}$ isolates, from an arbitrary symmetric perturbation $\delta\hat{g}_{\mu\nu}$, precisely the *physical, spin-2, transverse–traceless* (TT) content (Definition 10.25, (10.28)). The resulting graviton field is $\hat{h}_{\mu\nu} := \mathcal{P}^{(2)}_{\mathrm{grav}}\delta\hat{g}_{\mu\nu}$ and admits the triad decomposition $\hat{h}_{\mu\nu} = \hat{h}^{(\mathrm{spec})}_{\mu\nu} + \hat{h}^{(\mathrm{vac})}_{\mu\nu} + \hat{h}^{(\mathrm{corr})}_{\mu\nu}$ ((10.29)). This packages three operational requirements into a single map: (i) gauge safety, (ii) strict separation from scalar/vector (τ) contamination, and (iii) *provenance resolution* into spectral (radiative), vacuum, and correlational TT contributions—exactly mirroring the TT extraction of the radiative graviton in linearized gravity and its two physical helicities $\lambda = \pm 2$ [43, 7, 14].

Structure from the outside in. The outer projections $\mathsf{P}_{\mathrm{phys}}$ restrict to the BRST/Nielsen-safe physical sector of the realized weave configuration, eliminating gauge and ghost artifacts (Definition 10.1; see also [445, 106, 230]). The typed TT projectors $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ enforce the TT condition relative to the relational background $\bar{\Sigma}$ (Definition 10.3; cf. York/Lichnerowicz TT splitting [43, 44]). The map applies the TT projector on *both* sides of the triad, which makes the TT subspace invariant and annihilates any non-TT insertion before and after provenance resolution. In particular, TT-null preservation under τ–mixing follows from Lemma 10.123.

Provenance resolution inside TT. The central block $\mathsf{P}^{(2)}_{\mathrm{spec}} \oplus \Pi^{(\mathrm{vac})} \oplus \Pi^{(\mathrm{corr})}$ splits the TT content into orthogonal idempotent components (the *weave triad*, Definition 10.2). The *spectral* projector $\mathsf{P}^{(2)}_{\mathrm{spec}}$ is defined using the band-limited, gauge-fixed linearized kinetic operator $K_{\Phi\Phi}$ and isolates *on-band, propagating helicity-2* excitations (bona fide gravitational radiation), in line with the Riesz–Kato holomorphic calculus used for spectral selection [115, 253] and with Källén–Lehmann positivity for physical tensor modes [14, 255, 230, 244]. The *vacuum* projector $\Pi^{(\mathrm{vac})}$ retains TT structure associated with background/zero-mode or constraint-saturated content that is TT but not radiative, while the *correlational* projector $\Pi^{(\mathrm{corr})}$ records TT components generated by inter-modal correlations that lie outside the radiative spectrum yet remain in TT. Orthogonality and idempotence provide a clean ledger for audits and consistency checks without conflation across these sources.

Band-limiting, gauge-fixing, and monotonicity. Locating $\mathsf{P}^{(2)}_{\mathrm{spec}}$ through $K_{\Phi\Phi}$ makes the spectral cut *operational*: radiative content is tied to a certified IR/UV window and a fixed gauge, after which projection to $\mathcal{H}_{\mathrm{phys}}$ removes gauge dependence (Definition 10.1; cf. standard gauge fixing and renormalization hygiene [82, 447]). Under IR tightening, off-band contamination decreases monotonically, yielding progressively cleaner spectral–TT readouts; the outer TT/physical-sector sandwich ensures this improvement is reflected directly in $\hat{h}^{(\mathrm{spec})}_{\mu\nu}$ while keeping non-TT channels TT-null (Lemma 10.123).

Future-proofing of the construction. Because the definition is expressed at the level of *typed projectors*, extensions within spin-2 TT—such as refined spectral bands or additional massless helicity-2 species—are implemented by refining $\mathsf{P}^{(2)}_{\mathrm{spec}}$ (and, if needed, the triad idempotents) *without altering* the outer $\mathsf{P}_{\mathrm{phys}}$-TT-triad-TT-$\mathsf{P}_{\mathrm{phys}}$ scaffold (Definition 10.25). If a massive spin-2 sector is hypothesized, the TT type is replaced or supplemented by the appropriate spin-2 projector family while preserving the same "physical-sector $\perp$ provenance-split $\perp$ physical-sector'' skeleton; compare with the Fierz–Pauli massive spin-2 construction [474].

Interpreting $\hat{h}_{\mu\nu}$ and its components. Equation (10.29) makes explicit that the observable graviton field is the *sum of three TT contributions*,

$$\hat{h}^{(a)}_{\mu\nu} = \mathsf{P}_{\mathrm{phys}}\,\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\Pi^{(a)}\,\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\mathsf{P}_{\mathrm{phys}}\,\delta\hat{g}_{\mu\nu}, \qquad a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}.$$

This decomposition supports component-wise reporting, comparison, and auditing (e.g., distinguishing TT-leak from bona fide radiation) while ensuring that only the two tensor polarizations (helicities ± 2, the "+" and "×" modes) inhabit the signal [7, 14]. Scalar and vector entries remain *TT-null by construction* and cannot masquerade as additional tensor modes (Definition 10.3 and Lemma 10.123).

Definition 10.26 (Helicity resolution inside TT)**.** On the TT subspace, let $\Pi^{(\pm 2)}$ be the helicity projectors (background-frame covariant; parity swaps $\Pi^{(+2)} \leftrightarrow \Pi^{(-2)}$). The spectral graviton admits

$$\hat{h}^{(\mathrm{spec})}_{\mu\nu} = \sum_{\lambda=\pm 2} \Pi^{(\lambda)}\,\hat{h}^{(\mathrm{spec})}_{\mu\nu}, \qquad \Pi^{(\lambda)}\Pi^{(\lambda')} = \delta_{\lambda\lambda'}\Pi^{(\lambda)}, \quad \Pi^{(+2)} + \Pi^{(-2)} = \mathbf{1}_{\mathrm{TT}}. \tag{10.32}$$

Definition 10.27 (Propagating correlational TT filter (optional))**.** To isolate correlational TT with hyperbolic TT characteristics on a chosen window, set

$$\hat{h}^{(\mathrm{corr,prop})}_{\mu\nu} := \mathsf{P}_{\mathrm{phys}}\,\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\Pi^{(\mathrm{corr})}\,\mathbf{1}_{\{\ \sigma_{\mathrm{prin}}(K_{\Phi\Phi})\ \text{is TT-hyperbolic}\ \}}\,\Pi^{(\mathrm{corr})}\,\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\mathsf{P}_{\mathrm{phys}}\,\delta\hat{g}_{\mu\nu}. \tag{10.33}$$

Structural facts.

Lemma 10.28 (TT purity and triad commutation)**.** *$\hat{h}_{\mu\nu}$ is TT by construction: $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\hat{h}_{\mu\nu} = \hat{h}_{\mu\nu}$. Moreover, $[\Pi^{(a)}, \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}] = 0$ implies the decomposition (10.29) is triad-orthogonal.*

Proof. Immediate from the sandwiching by $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ and Proposition 10.9. □

Proposition 10.29 (Independence from τ-kairos)**.** *Let $\hat{\tau}_{\mu\nu}$ be as in Definition 10.13. Then $\mathcal{P}^{(2)}_{\mathrm{grav}}(\delta\hat{g}_{\mu\nu} + \hat{\tau}_{\mu\nu}) = \mathcal{P}^{(2)}_{\mathrm{grav}}\delta\hat{g}_{\mu\nu}$.*

Proof. By Lemma 10.17, $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\hat{\tau}_{\mu\nu} = 0$. Insert into (10.28). □

Proposition 10.30 (BRST/Nielsen flatness and band stability)**.** *Any correlator on $\mathcal{H}_{\mathrm{phys}}$ formed from $\hat{h}_{\mu\nu}$ and finitely many insertions of $\Pi^{(a)}$, $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$, and $\mathsf{P}^{(2)}_{\mathrm{spec}}$ is (i) gauge-parameter independent and (ii) stable under admissible retunings of (P_x, K_{IR}).*

Proof. A specialization of Proposition 10.10. □

Lemma 10.31 (Spectral future-proofing)**.** *If the realized dynamics acquires additional helicity-2 spectral branches (new contours γ_j), the projector $\mathsf{P}^{(2)}_{\mathrm{spec}}$ enlarges by the corresponding Riesz terms and $\mathcal{P}^{(2)}_{\mathrm{grav}}$ automatically includes the new species.*

Proof. By the additivity of Riesz projectors over disjoint spectral contours. □

Diagnostics and positivity.

Definition 10.32 (TT-leak diagnostic)**.** Define

$$\varepsilon := \left\| \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}} \left(\mathbf{1} - \mathsf{P}^{(2)}_{\mathrm{spec}}\right) \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}} \right\|_{\mathcal{I}}, \tag{10.34}$$

with $\|\cdot\|_{\mathcal{I}}$ any admissible band-invariant functional norm. Baseline $\varepsilon = 0$; $\varepsilon > 0$ flags unexpected TT strength outside the spectral spin-2 support and triggers audit (SynTh).

Proposition 10.33 (Transversality and positive spectral density (spectral piece))**.** *The spectral graviton obeys the Ward/Slavnov transversality and has nonnegative spectral density on $\mathcal{H}_{\mathrm{phys}}$: $k^\mu \hat{h}^{(\mathrm{spec})}_{\mu\nu} = 0$ and $\rho_{(2)}(\omega, \mathbf{k}) \geq 0$.*

Proof (outline). Inherited from the band-native Ward/Slavnov framework and positivity proven upstream (Ch. 8) after BRST descent; we only apply $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ and $\mathsf{P}_{\mathrm{phys}}$, which commute with the identities. □

Operational extraction and certification.

Definition 10.34 (Typed TT witnesses and visibilities)**.** Let $\mathbf{W}^{[\Phi\Phi;\mathrm{TT}]}$ be the consolidated witness in the TT-typed gaiamiti sector (Ward + spectral ledgers). Define the law/state visibilities $\Lambda^{[\Phi\Phi;\mathrm{TT}]}_{\mathrm{law/state}}$ as in Definitions 10.80 and 10.82, evaluated on $\mathcal{H}_{\mathrm{phys}}$ with $\mathcal{I}^{[\Phi\Phi]}$.

Proposition 10.35 (Well-posed Parasymmetric TLO for the graviton)**.** *The Parasymmetric TLO triple* (Type = TT, Location, Onset) *of Definition 10.85 and Proposition 10.90 exists and is unique on the certified band. The onset $r^\star[\mathrm{TT}]$ is stable under admissible retuning and tightening of K_{IR}.*

Proof. Type is fixed by $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$; uniqueness follows from orthogonality to non-TT subspaces. Stability is inherited from Proposition 10.10. □

Remarks.

Remark 10.36 (Vacuum vs correlational vs spectral TT)**.** $\hat{h}^{(\mathrm{spec})}$ represents radiative, energy-carrying spin-2 (gravitational waves). $\hat{h}^{(\mathrm{vac})}$ encodes TT vacuum structure (no asymptotic flux; contact/anomaly pieces). $\hat{h}^{(\mathrm{corr})}$ captures coherence-driven TT on finite windows (can be filtered to propagating TT by Definition 10.27); tightening K_{IR} typically suppresses the latter.

Remark 10.37 (Parity and helicity audits)**.** While TT is the primitive type, helicity-resolved visibilities via $\Pi^{(\pm 2)}$ allow a parity-asymmetry readout $\Delta_P^{(2)} := \Lambda^{[(+2)]} - \Lambda^{[(-2)]}$ when chiral effects are present; parity swaps the helicities.

The (τ)–kairos is not an autonomous sector but a typed family of insertions/projectors defined on the fixed realized weave configuration $(\mathcal{I}^{[\mathscr{S}]})$ (cf. Definition 10.1). The admissible (τ)-types are the scalar triplet $(S^{(b)}, S^{(L)}, \tilde{S}^{(p)})$ and the vector/helicity pair $(A^{(\pm)})$; they are admitted only through their typed idempotents $(\Pi^{[\chi]})$ and never as a separate sector label (Definition 10.4). Within the typed, triad-resolved correlator,

$$\hat{\mathbf{G}}^{(a\to b),[\chi\to\chi']} * \Lambda = \mathsf{P}_{\mathrm{phys}}\Pi^{(b)}\Pi^{[\chi']}(\hat{\mathbf{G}} * \Lambda)\Pi^{[\chi]}\Pi^{(a)}\mathsf{P}_{\mathrm{phys}}, \tag{10.35}$$

the placeholders $([\chi], [\chi'])$ may be instantiated by any of these (τ)-types, with triad provenance $(a, b \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\})$ (Definitions 10.2 and 10.113 and eq. (10.61)). In mixed situations $(\mathscr{S} = \Phi\Theta)$, the effective "typed spin" is fixed by the off-diagonal typed insertion itself, so no additional (τ)-sector index is introduced (Definition 10.113).

Remark 10.38 (Weinberg–Witten and its inapplicability in the present framework)**.** The Weinberg–Witten no–go theorem [475] forbids *emergent, interacting, massless spin–2* excitations under the joint hypotheses that:

(A) dynamics are formulated on a fixed Minkowski background η (with global Poincaré symmetry), and

(B) there exists a global, Lorentz–covariant, gauge–invariant, *conserved* stress tensor $T^{\mu\nu}$ that couples to the putative spin–2 composite.

In the typed, BRST/Nielsen–safe construction used here, both premises are relaxed. First, the operative chronogeometry is the expectation–valued, state–dependent background $\bar{\Sigma}_{\mu\nu}$ obtained on a certified plateau $\mathfrak{P}$ (Definitions 9.60 and 10.1 and eq. (10.95)), rather than a fixed η; local Lorentz frames exist on $\bar{\Sigma}$, but global Poincaré structure is not assumed (cf. background–independent viewpoints in [8, 3]). Second, stress observables are defined as *relational, typed witnesses* evaluated in BRST cohomology H^0 and adapted to $\bar{\Sigma}$ (Definitions 9.153 and 10.94), in line with the Haag–Kastler net perspective [230] and with BRST/Kugo–Ojima machinery [476, 108]; they are not given by a single, global Lorentz–covariant tensor on flat η. Gauge–parameter independence on certified plateaus (Nielsen safety) is ensured by the typed/Nielsen identities [120]. Consequently, the TT graviton $\hat{\gamma}_{\mu\nu} = \mathcal{P}^{(2)}[\delta\hat{g}_{\mu\nu}]$ (Definition 10.167)—identified as the tangential/Goldstone mode of the Gaiamiti sector (eq. (10.28))—is consistent within this framework; an operator–level Einstein channel realization appears in Section 11.7. Hence the Weinberg–Witten obstruction does not apply to the present setting.

Proposition 10.39 (Formal inapplicability of Weinberg–Witten)**.** *Let the gravitational TT excitation be* $\hat{\gamma}_{\mu\nu} = \mathcal{P}^{(2)}[\delta\hat{g}_{\mu\nu}]$ *on a certified plateau* $\mathfrak{P}$ *(Definitions 10.1 and 10.167), with observables evaluated in* $H^0(\hat{Q}_{\mathrm{BRST}})$ *and typed projectors satisfying the admission gate (Definition 10.94). Suppose:*

(i) the operative chronogeometry is the expectation–valued $\bar{\Sigma}$ *(not a fixed Minkowski metric), and*

(ii) operational stress tensors are defined relationally with respect to $\bar{\Sigma}$ *as BRST–closed, typed witnesses in* H^0 *(Definition 9.153), rather than as a single global Lorentz–covariant, gauge–invariant, conserved tensor on flat* η*.*

Then the joint hypotheses (A)–(B) *of the Weinberg–Witten theorem [475] are not met. Therefore the theorem does not constrain the existence of* $\hat{\gamma}_{\mu\nu}$ *in this framework.*

Proof. Weinberg–Witten [475] is an implication of the form

$$(\mathrm{A}) \ \wedge \ (\mathrm{B}) \implies \text{No emergent interacting massless spin–2.}$$

Hence, $\neg(\mathrm{A} \wedge \mathrm{B})$ is sufficient for inapplicability. By the stated premises:

Failure of (A). The background used operationally is $\bar{\Sigma}_{\mu\nu}$, obtained by Σ–realization on a certified plateau $\mathfrak{P}$ (Definitions 9.60 and 10.1 and eq. (10.95)). It is expectation–valued and state–dependent; a global Poincaré structure on flat η is not assumed (cf. [8, 3]). Thus (A) does not hold.

Failure of (B). Stress–energy enters as relational, typed, BRST–closed witnesses defined with respect to $\bar{\Sigma}$ and evaluated in H^0 (Definitions 9.153 and 10.94), consistent with the Haag–Kastler algebraic setting [230] and with BRST/Kugo–Ojima cohomology [476, 108]. A single global, Lorentz–covariant, gauge–invariant, conserved tensor on flat η is neither postulated nor required. Therefore (B) does not hold.

Since both (A) and (B) fail, $\neg(\mathrm{A} \wedge \mathrm{B})$ is satisfied and the conclusion of Weinberg–Witten does not follow. The TT excitation $\hat{\gamma}_{\mu\nu}$ then arises as the typed Goldstone/tangential mode of the Gaiamiti sector (eq. (10.28)) and is certified by BRST/Nielsen safety (cf. [120]). □

Corollary 10.40 (Einstein channel compatibility)**.** *Under the hypotheses of Proposition 10.39, the operator–level Einstein channel constructed on the typed, triad–resolved correlator over* $\bar{\Sigma}$ *(Section 11.7) is not obstructed by [475]. In particular, the TT projector* $\mathcal{P}^{(2)}$ *realizes a massless spin–2 pole protected by BRST/Nielsen identities [120] within the algebraic (Haag–Kastler) setting [230], without requiring a global Poincaré background [8, 3].*

Remark 10.41 (Formal implications schema)**.** Let $\mathsf{WW} := \big(\mathrm{A} \wedge \mathrm{B}\big) \Rightarrow \bot_{\text{spin-2}}$ denote the Weinberg–Witten entailment [475]. In the present construction:

$$\neg\mathrm{A} \quad \text{(expectation–valued } \bar{\Sigma}\text{)} \qquad \text{and} \qquad \neg\mathrm{B} \quad \text{(relational, BRST–closed stress witnesses)}$$

hold by design; hence $\neg(\mathrm{A} \wedge \mathrm{B})$ and therefore $\neg\mathsf{WW}$ (inapplicability) obtain. Moreover, typed/Nielsen identities [120] ensure gauge–parameter independence on certified plateaus, while the algebraic net structure [230] provides a consistent operator setting in which the TT Goldstone mode is realized (eq. (10.28)).

All (τ)–kairos operations act on the (Σ)-realized carrier ($\bar{\Sigma}$) established upstream (Definition 9.60). Typed (τ)-projectors commute with the triad projectors and preserve the graded/BRST structure—orthogonality, completeness, and triadic compatibility follow from the typed layer (Definitions 9.116 and 9.186 and Proposition 9.202) and are inherited by the weave construction (Remark 10.114 and Definition 10.2). Cone preservation and sector purity are maintained under the band/IR windowing used throughout (Remark 9.15); consequently, typed insertions can be made in any correlator block without compromising Nielsen safety or realized weave configuration invariance (Proposition 10.10).

Operationally, the (τ)-typed entries are certified with the same calibrated meters and visibilities as the other types, evaluated per triad channel under the ($\bar{\Sigma}$)-weighted inner products (Definitions 10.82 and 10.117). For the scalar family ($S^{(b)}, S^{(L)}, \tilde{S}^{(p)}$), TT-nullity is enforced by the vanishing of the TT component, monitored through the TT-leak diagnostic (ε) (Definition 10.32), with an additional parity-odd check for ($\tilde{S}^{(p)}$) consistent with the typed conventions. For the chrono–kymon vector ($A^{(\pm)}$), helicity purity is controlled by the corresponding residual on the ($A^{(\pm)}$)-subspaces and a parity asymmetry feature (Δ_P), optionally accompanied by a CP-resolved check; the parity and CP diagnostics are recorded as (Δ_P) and ($\Delta^{(a)}_{\mathrm{CP}}$) (Definitions 10.51 and 10.58). All such meters and features are computed triad-by-triad on ($\mathcal{H}_{\mathrm{phys}}$) with the admissible IR window fixed by the weave realized weave configuration (Definitions 10.1 and 10.2).

Two structural guarantees ensure robustness. First, *TT protection*: any (τ)-typed insertion leaves the graviton TT channel uncontaminated—($\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$) annihilates the ($\tau$) contributions and the corresponding typed blocks, as encoded in the orthogonality identities and the TT-null preservation lemma (eq. (10.36) and Lemma 10.123). Second, *typed coherence*: the typed projectors and witnesses supplied by Chapter 10 furnish orthogonality/completeness and commuting relations with the triad (Definition 9.187 and Proposition 9.202), which render the (τ)-diagnostics well-posed and directly comparable to those for TT, Higgs/CP, and CK entries within the same ledger (Definition 10.82).

Decision-making integrates the (τ)-diagnostics into the standard certify-or-audit pipeline. For each $((\mathscr{S}, [\chi]))$ with (χ) a (τ)-type, Law and State visibilities are tested against their thresholds and combined with the diagnostics $(\varepsilon, \Delta_P, \Delta^{(a)}_{\mathrm{CP}})$ through the SynTh potential ($\mathfrak{D}^{[\mathscr{S};\chi]}$) (Definitions 10.94 and 10.100 and eq. (10.59)). Co-tenability obtains when thresholds are met; otherwise, a finite audit path of admissible moves is followed until either co-tenability is achieved or a certified failure is issued (e.g., persistent TT leak, robust parity/CP asymmetry, or absent spectral support) (Definition 10.105 and Theorem 10.107).

Conceptually, this positions the (τ)–kairos as a typed weaving layer that is algebraically grounded in Chapter 10 and operationally deployed in Chapter 11. Chapter 10 supplies the (Σ)-carrier and the typed apparatus—menus, projectors, witnesses, and their triad-compatibility (Definitions 9.60, 9.116 and 9.187 and Proposition 9.202). Chapter 11 inserts these (τ)-types into the unified inter-modal correlator and resolves them triad-by-triad (Definition 10.113 and eq. (10.61)), augments them with calibrated meters and diagnostics (Definitions 10.32, 10.51, 10.58 and 10.82), and closes the loop with realized weave configuration-invariant certification via the audit/termination guarantees (Proposition 10.10, Definition 10.105, and Theorem 10.107).

10.5.1 The Graviton Six Polarization Modes (Hypothetical)

In standard metric theory as presently understood—namely general relativity (GR)—the gravitational-wave content consists of precisely two tensor polarizations, the "+" and "×" modes, corresponding to a massless spin-2 field with helicities ±2 [7, 14]. When reference is made to "metric theories with six GW polarizations," the intention is the broader class of metric theories (e.g., scalar–tensor, vector–tensor, certain bimetric models) in which a metric still governs free fall while additional fields can source non-tensor polarizations. In this generic classification there can be up to six polarization *types* in total—two tensor (the GR graviton sector), two vector, and two scalar (breathing and longitudinal)—as organized in the Newman–Penrose/ELLWW classification and subsequent detector-oriented treatments [477, 478, 479]. The graviton remains the tensor piece; the additional four modes are not graviton polarizations but arise from scalar/vector sectors (e.g., Brans–Dicke–type scalars or vector–tensor theories) [480, 479]. A change to the tensor projector family is required only if the representation for a *single* spin-2 field is altered (e.g., a massive spin-2 or a genuine kinematical departure), in which case a representation-adapted typed family is substituted for TT while retaining the same sandwiched scaffold (cf. Definition 10.25 and the discussion in Section 10.5.1); in bimetric frameworks the tensor content can also be enlarged/mixed [481].

Within the weave, the massless graviton sector is realized by the TT type, while any admitted non-tensor modes are modeled as typed τ–kairos families and inserted into the Unified Inter-Modal Correlator via the typed-insertion rule (Definition 10.113). This refinement acts on sector blocks while preserving BRST/band hygiene through the physical projector $\mathsf{P}_{\mathrm{phys}}$ and the realized weave configuration $\mathcal{I}^{[\mathcal{J}]}$. The scalar pair is encoded by the breathing and longitudinal types $S^{(b)}, S^{(L)}$ (with an optional pseudoscalar $\tilde{S}^{(p)}$), and the vector pair by the Chrono–Kymon helicities $A^{(\pm)}$ (Definition 10.4).

TT purity is enforced structurally: the TT projector $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ imposes the tensor condition relative to the relational background $\bar{\Sigma}$ (Definition 10.3; cf. York–Lichnerowicz TT extraction [43, 44]), and any correlator block containing a τ insertion is annihilated when sandwiched by $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ by the TT-null preservation lemma (Lemma 10.123). Consequently, the graviton readout remains strictly two-polarization TT even in the presence of scalar/vector modes, while τ content appears exclusively in its own typed channels.

Operationally, each admitted type (tensor, vector, scalar) is further decomposed by the provenance triad $\{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$ (Definition 10.2), enabling component-wise reporting without altering the polarization count. Law/state ledgers and cosine visibilities are computed as in Definition 10.82; TT employs the TT-leak diagnostic ε, vectors support the parity asymmetry Δ_P, and scalars permit CP/parity checks, with automatic triggers and cotenability thresholds administered per Definition 10.95. Mixed (off-diagonal) $\Phi\Theta$ interactions are handled by typed insertion on the corresponding correlator blocks (Definition 10.113); τ enters only as a typed projector family (never as a sector), so TT-null guarantees and the audit/provenance architecture are preserved.

Two extensions illustrate how departures from the GR baseline are incorporated without disturbing the outer scaffold. First, a *massive* spin-2 field (Fierz–Pauli-type) transforms with little group $SO(3)$ and carries five propagating helicities $\{\pm 2, \pm 1, 0\}$; waves obey a massive dispersion relation, gauge redundancy is broken, and Stückelberg fields track the

$\pm 1, 0$ components [474, 109, 113]. In the weave, this is modeled by replacing the TT type with a representation-adapted massive spin-2 typed family whose internal projectors resolve $\{\pm 2, \pm 1, 0\}$; the physical-sector–typed–triad–typed–physical-sector projector and the triad $\{\text{spec}, \text{vac}, \text{corr}\}$ remain unchanged, and the $\bar{\Sigma}$-weighted mixing/eigenproblem continues to apply (Definitions 10.2, 10.25, 10.113, 10.116 and 10.117). Second, a *true Lorentz/kinematical* departure alters the little-group/dispersion structure (e.g., preferred-frame backgrounds, helicity-dependent speeds, higher-derivative cones), thereby modifying the number or character of independent polarization states even for a single spin-2 representation; see, for example, the broader test frameworks for non-GR propagation [479, 478]. The weave handles this by swapping TT for a representation-adapted typed family tuned to the altered kinematics and inserting it into the same typed-insertion scaffold (Section 10.5.1 and Definitions 10.2 and 10.113), while keeping τ–kairos, chrono-kymon, and Higgs channels orthogonal (Definition 10.4).

Observable signatures in either extension include dispersion (frequency-dependent arrival times; subluminal/superluminal propagation), polarization content (power in vector/scalar patterns tied to the same spin-2 source, or non-standard tensor mixtures/rotations), and birefringence/anisotropy (helicity-dependent speeds or polarization rotation). These are recorded in the typed ledgers and visibilities with the existing meters and triggers (Definitions 10.82, 10.94 and 10.95); see also the phenomenological summaries in [479, 478].

Proposition 10.42 (Interpreting "six polarizations")**.** *Under the standard Poincaré/Wigner classification [482, 14], a single spin-2 field carries two independent polarizations if massless (little group $E(2)$) and five if massive (little group $SO(3)$; see also the Fierz–Pauli analysis [474]). Therefore, a statement of "six polarizations" must be understood as either:*

(A) *six* gravitational-wave *(GW) polarization* types *across distinct spin sectors (tensor 2 + vector 2 + scalar 2), as organized in the Newman–Penrose/ELLWW classification and modern detector frameworks [477, 478, 479]; or*

(B) *multiple* spin-2 species *whose tensor polarizations sum across species (e.g., in bimetric or multi-tensor scenarios [481]).*

The Chapter 11 weave formalism accommodates both while keeping the TT (graviton) channel uncontaminated (via $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ and the graviton projector Definitions 10.3 and 10.25).

Case (A): six GW types (2 tensor + 2 vector + 2 scalar). By Definition 10.25 and (10.28)–(10.29), the TT (graviton) readout from a symmetric perturbation $\delta\hat{g}_{\mu\nu}$ is

$$\hat{h}_{\mu\nu} = \mathcal{P}^{(2)}_{\mathrm{grav}}\,\delta\hat{g}_{\mu\nu} = \sum_{a\in\{\mathrm{spec},\mathrm{vac},\mathrm{corr}\}} \underbrace{\mathsf{P}_{\mathrm{phys}}\,\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\Pi^{(a)}\,\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\mathsf{P}_{\mathrm{phys}}\,\delta\hat{g}_{\mu\nu}}_{\hat{h}^{(a)}_{\mu\nu}}$$

with provenance split by the triad idempotents $\Pi^{(a)}$ (Definition 10.2). The additional four GW modes are *non-tensor*: they inhabit the τ–kairos families $S^{(b)}, S^{(L)}$ (scalars) and $A^{(\pm)}$ (vectors) (Definition 10.4). Typed orthogonality and TT–null preservation give, for every $X \in \{S^{(b)}, S^{(L)}, \tilde{S}^{(p)}, A^{(\pm)}\}$,

$$\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\Pi^{[X]} = 0 = \Pi^{[X]}\,\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}} \qquad \text{and} \qquad \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\hat{\mathbf{G}}^{[X\to Y]}_{\Lambda}\,\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}} = 0, \tag{10.36}$$

where $\hat{\mathbf{G}}_{\Lambda}^{[\cdot\to\cdot]}$ denotes the typed correlator blocks from Definition 10.113 and the first identity is enforced by Definition 10.3 and Lemma 10.123. Consequently, the TT readout remains strictly two–polarization, while the non–TT content is recorded exclusively in its own typed channels.

For transparent accounting, the τ-typed ledgers are taken triad-by-triad: for $a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$ and $X \in \{S^{(b)}, S^{(L)}, A^{(\pm)}\}$,

$$\mathcal{P}_X^{(a)} := \left\| \mathsf{P}_{\mathrm{phys}}\, \Pi^{[X]}\, \Pi^{(a)}\, \mathsf{P}_{\mathrm{phys}}\, \delta\hat{g} \right\|_{\mathcal{I}}, \qquad \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \Pi^{[X]}\, \Pi^{(a)} = 0 = \Pi^{(a)}\, \Pi^{[X]}\, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}, \tag{10.37}$$

so that (10.36)–(10.37) make the TT–null guarantee explicit while enabling component-wise reporting by provenance (Definition 10.2). The corresponding visibilities and diagnostics— e.g. $\Lambda_{\mathrm{law/state}}$ for power normalization, the TT-leak ε for non-spectral TT content, the vector parity asymmetry Δ_P, and scalar CP/parity checks—are computed on the typed, triad-resolved blocks using the calibrated meters and triggers (Definitions 10.82 and 10.95).

Case (B): six tensor states via multiple spin-2 species. Assume N massless spin-2 species, each with helicities ± 2. The TT type is refined by a species index so that the typed TT projector decomposes into orthogonal components,

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} = \sum_{i=1}^{N} \Pi_{\bar{\Sigma}}^{[\mathrm{TT}],i}, \qquad \Pi_{\bar{\Sigma}}^{[\mathrm{TT}],i}\, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}],j} = \delta_{ij}\, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}],i},$$

and the triad/spectral idempotents lift component–wise,

$$\mathsf{P}_{\mathrm{spec}}^{(2)} \rightsquigarrow \bigoplus_{i=1}^{N} \mathsf{P}_{\mathrm{spec}}^{(2),i}, \qquad \Pi^{(a)} \rightsquigarrow \bigoplus_{i=1}^{N} \Pi^{(a),i}, \quad a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\} \text{ (Definition 10.2)}.$$

With these refinements, the graviton projector of (10.28)–(10.29) becomes, in the same sandwiched form,

$$\mathcal{P}_{\mathrm{grav}}^{(2)} = \mathsf{P}_{\mathrm{phys}} \left(\sum_i \Pi_{\bar{\Sigma}}^{[\mathrm{TT}],i} \right) \left(\bigoplus_i \mathsf{P}_{\mathrm{spec}}^{(2),i} \oplus \bigoplus_i \Pi^{(\mathrm{vac}),i} \oplus \bigoplus_i \Pi^{(\mathrm{corr}),i} \right) \left(\sum_i \Pi_{\bar{\Sigma}}^{[\mathrm{TT}],i} \right) \mathsf{P}_{\mathrm{phys}},$$

so the species–resolved tensor field retains the provenance split:

$$\hat{h}_{\mu\nu} = \sum_a \sum_i \mathsf{P}_{\mathrm{phys}}\, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}],i}\, \Pi^{(a),i}\, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}],i}\, \mathsf{P}_{\mathrm{phys}}\, \delta\hat{g}_{\mu\nu}, \qquad a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}.$$

Species mixing is resolved by the $\bar{\Sigma}$–weighted generalized eigen-operator pair construction and whitening (Definitions 10.116 and 10.117). Writing $(\mathcal{A}, \mathcal{B})$ for the typed TT generalized eigen-operator pair on the certified band (Definitions 10.113 and 10.117), the generalized eigenproblem

$$\mathcal{A}\, u = \lambda\, \mathcal{B}\, u, \qquad \langle u_i, u_j \rangle_{\bar{\Sigma}} := \left\langle u_i, \mathcal{B}\, u_j \right\rangle_{\bar{\Sigma}} = \delta_{ij},$$

is whitened by $w := \mathcal{B}^{-1/2} u$ so that $\mathcal{B}^{-1/2} \mathcal{A}\, \mathcal{B}^{-1/2}$ is self-adjoint in the reference $\bar{\Sigma}$-weighted inner product (Definition 10.117). The corresponding orthogonal projectors onto the eigen–TT channels, $\{\Pi_{\mathrm{eig}}^{[\mathrm{TT}],i}\}_{i=1}^{N}$, satisfy

$$\Pi_{\mathrm{eig}}^{[\mathrm{TT}],i}\, \Pi_{\mathrm{eig}}^{[\mathrm{TT}],j} = \delta_{ij}\, \Pi_{\mathrm{eig}}^{[\mathrm{TT}],i}, \qquad \sum_{i=1}^{N} \Pi_{\mathrm{eig}}^{[\mathrm{TT}],i} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]},$$

and block–diagonalize the typed TT correlator,

$$\Pi^{[\mathrm{TT}],i}_{\mathrm{eig}}\,\hat{\mathbf{G}}_{\Lambda}^{[\mathrm{TT}\to\mathrm{TT}]}\,\Pi^{[\mathrm{TT}],j}_{\mathrm{eig}} = 0 \quad (i \neq j),$$

so each species is reported in an orthogonal eigen–TT channel with the triad provenance retained.

For transparent ledgers, the species– and provenance–resolved TT power is

$$\mathcal{P}^{(a)}_{\mathrm{TT},i} := \left\| \mathsf{P}_{\mathrm{phys}}\,\Pi^{[\mathrm{TT}],i}_{\mathrm{eig}}\,\Pi^{(a),i}\,\mathsf{P}_{\mathrm{phys}}\,\delta\hat{g} \right\|_{\mathcal{I}}, \qquad a \in \{\mathrm{spec},\mathrm{vac},\mathrm{corr}\},$$

with visibilities/diagnostics computed on these typed blocks using the calibrated meters and triggers (Definitions 10.82 and 10.95). Thus the outcome is a set of orthogonal eigen–TT channels (one per species), with TT purity (via the sandwiched form) and triad provenance (Definition 10.2) preserved throughout.

When a deeper rewrite is required. If a *single* spin-2 representation supports six independent tensor-like polarizations (e.g., due to strong Lorentz breaking or non–Wigner kinematics), the TT type is replaced by a representation-adapted typed family while the outer scaffold and provenance logic remain unchanged. Concretely, let

$$\Pi^{[\mathrm{RT}]} \;=\; \sum_{i=1}^{m} \Pi^{[\pi_i]}, \qquad \Pi^{[\pi_i]}\,\Pi^{[\pi_j]} = \delta_{ij}\,\Pi^{[\pi_i]},$$

denote the representation-adapted tensor-type projector family (cf. the representation-adapted construction outlined in Section 10.5.1 and the projector-algebra pattern used for TT and typed families in Definitions 10.2 and 10.3), where m is the number of independent polarization projectors admitted by the altered little-group structure (e.g., $m = 6$ in the scenario at hand).

The TT projector in the sandwiched construction is replaced *in situ*:

$$\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}} \;\rightsquigarrow\; \Pi^{[\mathrm{RT}]},$$

and the Chapter 11 sandwiched graviton projector (Definition 10.25, (10.28)) becomes

$$\mathsf{P}_{\mathrm{phys}}\,\Pi^{[\mathrm{RT}]}\left(\mathsf{P}^{(\mathrm{RT})}_{\mathrm{spec}} \;\oplus\; \Pi^{(\mathrm{vac})} \;\oplus\; \Pi^{(\mathrm{corr})}\right)\Pi^{[\mathrm{RT}]}\,\mathsf{P}_{\mathrm{phys}},$$

i.e., the same physical–typed–triad–typed–physical scaffold with the usual triad idempotents $\{\Pi^{(\mathrm{spec})},\Pi^{(\mathrm{vac})},\Pi^{(\mathrm{corr})}\}$ (Definition 10.2) and a representation-adapted spectral selector $\mathsf{P}^{(\mathrm{RT})}_{\mathrm{spec}}$ (constructed on the certified band as in Definition 10.7, with measurability per Proposition 10.149). The corresponding field readout mirrors (10.29):

$$\hat{h}^{(\mathrm{RT})}_{\mu\nu} = \sum_{a\in\{\mathrm{spec},\mathrm{vac},\mathrm{corr}\}} \mathsf{P}_{\mathrm{phys}}\,\Pi^{[\mathrm{RT}]}\,\Pi^{(a)}\,\Pi^{[\mathrm{RT}]}\,\mathsf{P}_{\mathrm{phys}}\,\delta\hat{g}_{\mu\nu}.$$

Typed orthogonality to the τ–kairos families (Definition 10.4) is imposed exactly as for TT, yielding the same null-preservation pattern (cf. Lemma 10.123):

$$\Pi^{[\mathrm{RT}]}\,\Pi^{[X]} = 0 = \Pi^{[X]}\,\Pi^{[\mathrm{RT}]}, \qquad \Pi^{[\mathrm{RT}]}\,\hat{\mathbf{G}}_{\Lambda}^{[X\to Y]}\,\Pi^{[\mathrm{RT}]} = 0, \quad X \in \{S^{(b)},S^{(L)},\tilde{S}^{(p)},A^{(\pm)}\},$$

so the representation-adapted tensor channel remains uncontaminated, while non-tensor content is recorded in its own typed ledgers (Definitions 10.82, 10.95 and 10.113).

Gauge-/retune-robustness is inherited from the same identities used for TT. For any composition $\mathcal{J}$ built from admissible typed/triad idempotents, the embedded Nielsen identity on $\mathcal{H}_{\text{phys}}$ (Ch. 8; cf. Proposition 7.174) gives

$$\partial_\xi \big\langle \mathcal{J}\, \delta\hat{g} \big\rangle_{\mathcal{H}_{\text{phys}}} = 0,$$

and along admissible retunes the Riesz selectors remain continuous on certified contours (Definition 10.7 and Proposition 10.149), so the representation-adapted correlators vary continuously under the same audit moves. Consequently, while the *typed tensor family* changes (TT $\mapsto$ RT), the outer scaffold "physical-sector $\perp$ provenance-split $\perp$ physical-sector" and the triad decomposition/audit logic (Definition 10.2) remain in force without further modification.

Bottom line.

- In metric theories with six GW types, the graviton stays two-polarization TT; the extra four are non-TT and live in τ types (Definitions 10.3 and 10.4 and Lemma 10.123).
- If experiments revealed six *tensor* states in total, interpret them as multiple spin-2 species; the sandwiched projector and triad split extend by species without structural change (Definition 10.25, (10.28), (10.29)).
- A truly "six-polarization" single spin-2 signals new kinematics; the framework operates by swapping in the representation-adapted typed projector family while keeping the audit/provenance architecture unchanged.

10.6 The Higgs (Radial Kymon) in the Tantrakyma Sector

We work on the realized weave configuration $\mathcal{I}^{[\mathcal{Y}]}$ of Definition 10.1 in the $\Theta\Theta$ (tantrakyma) sector, with triad projectors $\Pi^{(a)}$ from Definition 10.2, the CP–refined radial projectors $\Pi_\Theta^{[\text{rad}\pm]}$ of Definition 10.5, and the spin-0 Riesz selector $\mathsf{P}^{(0)}_{\text{spec}}$ from Definition 10.7. All statements are taken on $\mathcal{H}_{\text{phys}} = \operatorname{Ran} \mathsf{P}_{\text{phys}}$.

Definitions.

Definition 10.43 (Higgs projectors (CP-resolved, triad-resolved, future-proof))**.** Let $K^{(0)}_{\Theta\Theta}$ be the band-limited, gauge-fixed scalar sub-block of the $\Theta\Theta$ kinetic operator. The *CP-resolved Higgs projectors* are

$$\boxed{\mathcal{P}^{(0)}_{\text{H},\pm} := \mathsf{P}_{\text{phys}}\, \Pi_\Theta^{[\text{rad}\pm]} \Big(\mathsf{P}^{(0)}_{\text{spec}} \;\oplus\; \Pi^{(\text{vac})} \;\oplus\; \Pi^{(\text{corr})} \Big) \Pi_\Theta^{[\text{rad}\pm]}\, \mathsf{P}_{\text{phys}}} \tag{10.38}$$

and the corresponding *Higgs fields* (CP-even/odd) extracted from the tantrakyma fluctuation $\delta\hat{\mathbf{\Theta}}$ are

$$\boxed{\hat{H}_\pm := \mathcal{P}^{(0)}_{\mathrm{H},\pm}\,\delta\hat{\mathbf{\Theta}} = \sum_{a\in\{\mathrm{spec,vac,corr}\}} \hat{H}^{(a)}_\pm}\,, \qquad \hat{H}^{(a)}_\pm := \mathsf{P}_{\mathrm{phys}}\,\Pi^{[\mathrm{rad}\pm]}_\Theta\,\Pi^{(a)}\,\Pi^{[\mathrm{rad}\pm]}_\Theta\,\mathsf{P}_{\mathrm{phys}}\,\delta\hat{\mathbf{\Theta}}. \tag{10.39}$$

Definition 10.44 (CP mixing block and mass/band eigenmodes)**.** In the CP basis (H_+, H_-), the scalar operator restricted to the radial subspace can be represented as

$$K^{(0)}_{\Theta\Theta}\big|_{\mathrm{rad}} = \begin{pmatrix} K_{++} & K_{+-} \\ K_{-+} & K_{--} \end{pmatrix}, \qquad K_{-+} = K_{+-}^\dagger. \tag{10.40}$$

The *mass/band eigenmodes* are obtained by a unitary diagonalization U of (10.40), equivalently by the Riesz spectral projectors of $\mathsf{P}^{(0)}_{\mathrm{spec}}$.

Structural facts.

Lemma 10.45 (Orthogonality, completeness, and CP resolution)**.** *The ranges of* $\Pi^{[\mathrm{rad}+]}_\Theta$ *and* $\Pi^{[\mathrm{rad}-]}_\Theta$ *are orthogonal, and* $\Pi^{[\mathrm{rad}+]}_\Theta + \Pi^{[\mathrm{rad}-]}_\Theta = \Pi^{[\mathrm{rad}]}_\Theta$*. Moreover,* $[\Pi^{(a)}, \Pi^{[\mathrm{rad}\pm]}_\Theta] = 0$ *for all* a*.*

Proof. Orthogonality and completeness follow from Definition 10.5. Commutation with the triad is part of that definition. □

Lemma 10.46 (BRST/Nielsen invariance and band stability)**.** *Any correlator on* $\mathcal{H}_{\mathrm{phys}}$ *formed from* $\hat{H}_\pm$ *and finitely many insertions of* $\Pi^{(a)}$*,* $\Pi^{[\mathrm{rad}\pm]}_\Theta$*, and* $\mathsf{P}^{(0)}_{\mathrm{spec}}$ *is (i) gauge-parameter independent and (ii) stable under admissible retunings of* (P_x, K_{IR})*.*

Proof. This is a specialization of Proposition 10.10. □

Proposition 10.47 (NG/pNG orthogonality (radial selection))**.** *Let* $\Pi^{[\mathrm{NG/pNG}]}_\Theta$ *denote the upstream NG/pNG projector from the tantrakyma SSB analysis (Ch. 10, referenced there). Then* $\Pi^{[\mathrm{NG/pNG}]}_\Theta\,\Pi^{[\mathrm{rad}\pm]}_\Theta = 0$ *and* $\Pi^{[\mathrm{rad}\pm]}_\Theta$ *projects onto amplitude (radial) directions only.*

Proof. By construction of the radial type upstream (orthogonality of amplitude vs. Goldstone directions in the Σ-weighted inner product), preserved by the CP refinement which commutes with the upstream split. □

Proposition 10.48 (Future-proofing under additional scalar species)**.** *If the realized dynamics acquires additional spin-0 spectral branches (new contours* Γ_j*),* $\mathsf{P}^{(0)}_{\mathrm{spec}}$ *enlarges by the corresponding Riesz terms and* $\mathcal{P}^{(0)}_{\mathrm{H},\pm}$ *automatically includes the new species.*

Proof. Additivity of Riesz projectors over disjoint spectral contours. □

CP (a)symmetry, positivity, and diagnostics.

Proposition 10.49 (Block structure under CP conservation/violation)**.** *If CP is conserved, $K_{+-} = 0$ and $\hat{H}_+, \hat{H}_-$ propagate independently (block-diagonal case). If CP is violated (explicitly or spontaneously), $K_{+-} \neq 0$; then the spectral projectors from $\mathsf{P}^{(0)}_{\mathrm{spec}}$ return mass/band eigenmodes which are unitary mixtures of $\hat{H}_+, \hat{H}_-$.*

Proof. Direct from (10.40) and the spectral calculus. In the CP-conserving case the Riesz projectors coincide with the CP projectors; otherwise they are related by a unitary U which diagonalizes the 2×2 block. □

Proposition 10.50 (Spectral positivity (radiative scalar))**.** *The spectral (radiative) Higgs component has nonnegative spectral density on $\mathcal{H}_{\mathrm{phys}}$: $\rho_{(0)}(\omega, \mathbf{k}) \geq 0$.*

Proof (outline). Inherited from the upstream band-positive structure (Ward/Nielsen/BRST framework) after restriction to the radial scalar subspace; projectors commute with the positivity arguments. □

Definition 10.51 (CP asymmetry audit)**.** For each triad channel a, define the (law/state) CP asymmetry

$$\Delta^{(a)}_{\mathrm{CP}} := \Lambda^{[H^{(a)}_+]}_{\mathrm{law/state}} - \Lambda^{[H^{(a)}_-]}_{\mathrm{law/state}}, \tag{10.41}$$

with $\Lambda^{[\cdot]}_{\mathrm{law/state}}$ from Definitions 10.80 and 10.82. CP conservation implies $\Delta^{(a)}_{\mathrm{CP}} = 0$.

Operational extraction and certification.

Definition 10.52 (Typed Higgs witnesses and visibilities)**.** Let $\mathbf{W}^{[\Theta\Theta;\mathrm{rad}\pm]}$ be the consolidated witness for the CP-resolved radial types in the tantrakyma sector (Ward + spectral ledgers). Define $\Lambda^{[\Theta\Theta;\mathrm{rad}\pm]}_{\mathrm{law/state}}$ as in Definitions 10.80 and 10.82, evaluated on $\mathcal{H}_{\mathrm{phys}}$ with $\mathcal{I}^{[\Theta\Theta]}$.

Proposition 10.53 (Well-posed Parasymmetric TLO for the Higgs)**.** *For each CP-resolved radial type, the Parasymmetric TLO triple* (Type $= \mathrm{rad}\pm$, Location, Onset) *of Definition 10.85 and Proposition 10.90 exists and is unique on the certified band. The onset $r^\star[\mathrm{rad}\pm]$ is stable under admissible retuning and tightening of K_{IR}.*

Proof. Type is fixed by $\Pi^{[\mathrm{rad}\pm]}_\Theta$; uniqueness follows from orthogonality to NG/pNG (Prop. 10.47) and to other scalar types. Stability is inherited from Lemma 10.46. □

Remarks.

Remark 10.54 (Physics reading by triad)**.** $\hat{H}^{(\mathrm{spec})}_\pm$ represents the radiative, particle-like Higgs content (possibly multiple scalar species under Proposition 10.48). $\hat{H}^{(\mathrm{vac})}_\pm$ encodes vacuum scalar structure (no asymptotic flux; baseline/contact pieces). $\hat{H}^{(\mathrm{corr})}_\pm$ captures coherence/near-field scalar responses that typically diminish upon tightening K_{IR}.

Remark 10.55 (SM-like limit)**.** The usual SM Higgs corresponds to a dominantly CP-even, spectral radial mode $\hat{H}^{(\mathrm{spec})}_+$ with negligible $\hat{H}_-$ and suppressed vac/corr components on the certified plateau; the framework accommodates CP-odd admixtures without modification (Proposition 10.49).

10.7 The Chrono–Kymon (Vector Channel of the τ-kairos)

We work on the realized weave configuration $\mathcal{I}^{[\mathcal{J}]}$ of Definition 10.1 in the τ-kairos block, with triad projectors $\Pi^{(a)}$ from Definition 10.2, the helicity-resolved typed idempotents $A^{(\pm)}$ from Definition 10.4, and the TT projector $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ from Definition 10.3. All statements are taken on $\mathcal{H}_{\mathrm{phys}} = \mathrm{Ran}\,\mathsf{P}_{\mathrm{phys}}$.

Definitions.

Definition 10.56 (Spin-1 spectral selector in the τ-vector subspace)**.** Let $K_\tau^{(1)}$ denote the band-limited, gauge-fixed linearized kinetic operator restricted to the *vector* subspace of the τ-kairos block.[2] Define the spin-1 Riesz selector, with $\{\zeta_m\}_{m\in\mathfrak{S}_1}$ disjoint contours enclosing the realized helicity-1 spectral features (poles/bands) on the certified band, by

$$\mathsf{P}_{\mathrm{spec}}^{(1)} := \sum_{m\in\mathfrak{S}_1} \frac{1}{2\pi i}\oint_{\zeta_m} \left(K_\tau^{(1)} - z\right)^{-1} dz\,. \tag{10.42}$$

We also set the spin-1 spectral triad piece $\Pi_\tau^{(\mathrm{spec};1)} := \mathsf{P}_{\mathrm{spec}}^{(1)}$ for later triad bookkeeping.

Definition 10.57 (Chrono–Kymon projectors and fields (helicity- and triad-resolved))**.** For $\lambda = \pm$, define the *helicity-resolved Chrono–Kymon projectors*

$$\boxed{\mathcal{P}_{\mathrm{CK},\lambda}^{(1)} := \mathsf{P}_{\mathrm{phys}}\, A^{(\lambda)} \left(\, \Pi_\tau^{(\mathrm{spec};1)} \;\oplus\; \Pi^{(\mathrm{vac})} \;\oplus\; \Pi^{(\mathrm{corr})} \,\right) A^{(\lambda)}\, \mathsf{P}_{\mathrm{phys}}} \tag{10.43}$$

acting on the τ-vector carrier space. From a raw τ-vector fluctuation $\delta\hat{K}_\mu^{(\tau)}$ we extract the (helicity, triad) Chrono–Kymon fields

$$\boxed{\hat{K}_{\lambda,\mu}^{(\tau)(a)} := \mathsf{P}_{\mathrm{phys}}\, A^{(\lambda)}\, \Pi^{(a)}\, A^{(\lambda)}\, \mathsf{P}_{\mathrm{phys}}\, \delta\hat{K}_\mu^{(\tau)}, \qquad \hat{K}_{\lambda,\mu}^{(\tau)} := \sum_{a\in\{\mathrm{spec,vac,corr}\}} \hat{K}_{\lambda,\mu}^{(\tau)(a)}} \tag{10.44}$$

with $\Pi^{(\mathrm{spec})}$ implicitly specialized to $\Pi_\tau^{(\mathrm{spec};1)}$ in the vector channel. Their τ-kairos *tensor images* via the typed map $A_{\mu\nu}^{(\lambda)}[\,\cdot\,]$ (cf. Definition 10.4) are

$$\hat{\tau}_{\mu\nu}^{(\mathrm{CK},\lambda)} := A_{\mu\nu}^{(\lambda)}\big[\hat{K}_\lambda^{(\tau)}\big] = \sum_a A_{\mu\nu}^{(\lambda)}\big[\hat{K}_\lambda^{(\tau)(a)}\big]\,. \tag{10.45}$$

Definition 10.58 (Parity action on helicities and visibility asymmetry)**.** Let $\mathcal{P}_\tau$ denote parity on the τ-vector subspace; then $\mathcal{P}_\tau A^{(+)}\mathcal{P}_\tau^{-1} = A^{(-)}$ and $\mathcal{P}_\tau A^{(-)}\mathcal{P}_\tau^{-1} = A^{(+)}$. Define the (law/state) *visibility* parity-asymmetry diagnostic by

$$\Delta_{P,\Lambda}^{(1)} := \Lambda_{\mathrm{law/state}}^{[A^{(+)}]} - \Lambda_{\mathrm{law/state}}^{[A^{(-)}]}\,, \tag{10.46}$$

using the witnesses of Definitions 10.80 and 10.82.

[2] Equivalently, one may use the band-limited physical two-point on the τ-vector subspace. The admissible restriction is secured by the type projectors of Definition 10.4.

Structural facts.

Lemma 10.59 (TT-nullity and orthogonality)**.** *For each $\lambda = \pm$ and a, one has $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\hat{\tau}_{\mu\nu}^{(\mathrm{CK},\lambda)} = 0$ and $A^{(\lambda)}A^{(\lambda')} = \delta_{\lambda\lambda'}A^{(\lambda)}$.*

Proof. TT-nullity follows from the typed construction in Definition 10.4. Orthogonality is part of the projector algebra in Lemma 10.8. □

Proposition 10.60 (Triad commutation and decomposition)**.** $[\Pi^{(a)}, A^{(\lambda)}] = 0$ *for all a, λ, hence the decomposition (10.44) is triad-orthogonal. Moreover, $\hat{K}^{(\tau)}_{\lambda,\mu} = \hat{K}^{(\tau)(\mathrm{spec})}_{\lambda,\mu} \oplus \hat{K}^{(\tau)(\mathrm{vac})}_{\lambda,\mu} \oplus \hat{K}^{(\tau)(\mathrm{corr})}_{\lambda,\mu}$.*

Proof. Commutation is stated in Proposition 10.9; the decomposition is immediate. □

Proposition 10.61 (BRST/Nielsen flatness and band stability)**.** *Any correlator on $\mathcal{H}_{\mathrm{phys}}$ formed from $\hat{K}^{(\tau)}_{\lambda}$, $\hat{\tau}^{(\mathrm{CK},\lambda)}$ and finitely many insertions of $A^{(\lambda)}$, $\Pi^{(a)}$, and $\mathsf{P}^{(1)}_{\mathrm{spec}}$ is (i) gauge-parameter independent and (ii) stable under admissible retunings of (P_x, K_{IR}).*

Proof. A specialization of Proposition 10.10. □

Lemma 10.62 (Spectral future-proofing (spin-1))**.** *If additional helicity-1 spectral branches appear (new contours ζ_n), the projector $\mathsf{P}^{(1)}_{\mathrm{spec}}$ enlarges by the corresponding Riesz terms and $\mathcal{P}^{(1)}_{\mathrm{CK},\lambda}$ automatically includes the new species.*

Proof. By additivity of Riesz projectors over disjoint spectral contours. □

Positivity, Ward identities, and diagnostics.

Proposition 10.63 (Transversality/longitudinality ledgers and spectral positivity)**.** *On $\mathcal{H}_{\mathrm{phys}}$, the spectral Chrono–Kymon components obey the appropriate Ward/Slavnov ledgers (transverse/longitudinal as admitted by the type) and have nonnegative spectral density $\rho_{(1)}(\omega, \mathbf{k}) \geq 0$.*

Proof (outline). Inherited from the upstream BRST/Nielsen framework (Ch. 8) restricted to the vector subspace and the typed projectors $A^{(\lambda)}$; positivity is preserved under the projectors and the Riesz selection. □

Definition 10.64 (Helicity-resolved visibilities)**.** Let $\mathbf{W}^{[\tau\text{-vec};A^{(\lambda)}]}$ be the consolidated witness for the τ-vector channel. Define $\Lambda^{[\tau\text{-vec};A^{(\lambda)}]}_{\text{law/state}}$ as in Definitions 10.80 and 10.82, evaluated on $\mathcal{H}_{\mathrm{phys}}$ with $\mathcal{I}^{[\mathscr{S}]}$.

Proposition 10.65 (Well-posed Parasymmetric TLO for Chrono–Kymon)**.** *For each helicity $\lambda = \pm$, the Parasymmetric TLO triple* (Type $= A^{(\lambda)}$, Location, Onset) *of Definition 10.85 and Proposition 10.90 exists and is unique on the certified band. The onset $r^{\star}[A^{(\lambda)}]$ is stable under admissible retuning and tightening of K_{IR}.*

Proof. Type is fixed by $A^{(\lambda)}$; uniqueness follows from $A^{(+)}A^{(-)} = 0$. Stability is inherited from Proposition 10.61. □

Remarks.

Remark 10.66 (Interface with the τ-kairos tensor and the graviton)**.** The τ-tensor image $\hat{\tau}_{\mu\nu}^{(\mathrm{CK},\lambda)}$ is TT-null (Lemma 10.59) and therefore cannot contaminate the geometrical TT (graviton) channel (cf. Cor. 10.20). Its triad decomposition spec/vac/corr distinguishes radiative vector excitations from vacuum/coherence effects.

Remark 10.67 (Parity vs helicity reporting)**.** Helicity $A^{(\pm)}$ is the primitive diagnostic (useful for chiral physics). Parity is reported via the visibility asymmetry $\Delta_{P,\Lambda}^{(1)}$ in (10.46), with $\mathcal{P}_\tau$ swapping helicities. A *ledger-based* normalized parity asymmetry can also be quoted as

$$\widetilde{\Delta}_P^{(1)} := \frac{\mathcal{L}_{A^{(+)}} - \mathcal{L}_{A^{(-)}}}{\mathcal{L}_{A^{(+)}} + \mathcal{L}_{A^{(-)}}},$$

see the meters paragraph below. This disambiguates the two notions used in earlier drafts with the same symbol.

Spin–1 Chrono–Kymon: the vector typed content of τ

The *Chrono–Kymon (CK)* is the **spin–1** (vector) typed family inside the τ–kairos channel. In the typed taxonomy it is the pair of helicity projectors $A^{(\pm)}$ from the admitted τ–families (Definition 10.4). CK is *strictly non–TT* by construction: any CK insertion is annihilated by the TT sandwich,

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \hat{\mathbf{G}}_\Lambda^{[\,A^{(\lambda)}\to Y\,]} = 0 = \hat{\mathbf{G}}_\Lambda^{[\,Y\to A^{(\lambda)}\,]}\, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \quad (\lambda = \pm),$$

as guaranteed by the typed–insertion rule and TT–null preservation (Definitions 10.3 and 10.113 and Lemma 10.123). Thus CK carries *no* radiative spin–2 (graviton) content; the TT graviton readout remains purely tensorial.

Typed extraction and correlator blocks. For any admissible fluctuation/correlator, CK content is read helicity-by-helicity via the typed insertion (Definition 10.113). Writing $\hat{\mathbf{G}}_\Lambda$ for the unified inter-modal correlator on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$, define the CK-typed blocks and their triad resolutions by

$$\hat{\mathbf{G}}_\Lambda^{[\,A^{(\lambda)}\to A^{(\lambda')}\,]} := \Pi^{[A^{(\lambda)}]}\, \hat{\mathbf{G}}_\Lambda\, \Pi^{[A^{(\lambda')}]}, \qquad \hat{\mathbf{G}}_\Lambda^{(a)\,[\,A^{(\lambda)}\to A^{(\lambda')}\,]} := \Pi^{(a)}\, \hat{\mathbf{G}}_\Lambda^{[\,A^{(\lambda)}\to A^{(\lambda')}\,]}\, \Pi^{(a)},$$

with $a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$ (Definition 10.2). Orthogonality and idempotence of the typed and triad projectors ensure that CK channels can be audited provenance-by-provenance without mixing with TT.

Optional first-order representative (notation only). When convenient for interpretation, introduce a vector representative $K_\mu^{(\tau)}$ for the CK block and its torsion-like two-form relative to the relational background $\bar{\Sigma}$:

$$K_\mu^{(\tau)}\ (\text{CK representative}), \qquad \mathcal{F}_{\mu\nu}^{(\tau)} := 2\,\bar{\nabla}_{[\mu} K_{\nu]}^{(\tau)}\,.$$

These are *not* new operators in the formalism; they are notational proxies for the CK-typed correlator content. All physically meaningful statements are made at the level of typed, triad-resolved correlators.

Spin selection and TT purity. CK is spin–1 (helicity ± 1) by its typed construction (Definition 10.4); TT purity of the graviton channel follows from the TT sandwich and typed orthogonality:

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\Pi^{[A^{(\lambda)}]} = 0 = \Pi^{[A^{(\lambda)}]}\,\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}, \qquad \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\hat{\mathbf{G}}_{\Lambda}^{[A^{(\lambda)}\to Y]}\,\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} = 0 \quad (\lambda = \pm),$$

(Definition 10.3 and Lemma 10.123). Hence CK does not contribute to the TT graviton projector (Definition 10.25).

Meters, visibilities, and ledgers (CK channel). To disambiguate notation, we reserve $\mathcal{P}$ for *projectors* (e.g., $\mathcal{P}^{(1)}_{\mathrm{CK},\lambda}$) and use

$$\mathcal{L}^{(a)}_{A^{(\lambda)}} \;:=\; \left\| \mathsf{P}_{\mathrm{phys}}\,\Pi^{[A^{(\lambda)}]}\,\Pi^{(a)}\,\mathsf{P}_{\mathrm{phys}}\,\delta\hat{\Sigma} \right\|_{\mathcal{I}}$$

for the CK *ledger power* in triad sector $a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$ and helicity $\lambda = \pm$.[3] The visibility-based parity diagnostic is $\Delta^{(1)}_{P,\Lambda}$ from (10.46), while a normalized ledger asymmetry may be quoted as $\widetilde{\Delta}^{(1)}_{P}$ (Remark 10.67). All meters are evaluated on $\mathcal{H}_{\mathrm{phys}}$ and inherit BRST/Nielsen flatness and band stability from the realized weave configuration.

Mixing, whitening, and reporting. CK may mix across helicities or with other admitted types inside non-TT sectors, but never with TT under the typed sandwich. Mixing is quantified by the typed mixing norms and resolved by the $\bar{\Sigma}$–weighted operator-generalized eigen-operator pair eigenproblem and whitening (Definitions 10.116 and 10.117). The outcome is an orthogonal set of eigen-channels inside the CK block, each carrying its triad provenance and reportable power/visibilities.

Dynamics (model class placeholder, if needed). When a first-order representative $K^{(\tau)}_{\mu}$ is adopted for phenomenology, CK dynamics may be modeled—*within the typed block*—by a Maxwell/Proca-type equation on $(\bar{\Sigma}, \bar{\nabla})$,

$$\bar{\nabla}_{\nu}\Big(\bar{\Sigma}^{\nu\alpha}\mathcal{F}^{(\tau)}_{\alpha\mu}\Big) + m^2_{(1)}\,K^{(\tau)}_{\mu} = J^{(\tau)}_{\mu}, \qquad \bar{\nabla}^{\mu}K^{(\tau)}_{\mu} = 0 \;\; (\text{on shell if } m_{(1)} \neq 0),$$

with causal support governed by the $\bar{\Sigma}$ light cone. This is an interpretive layer on top of the canonical typed-correlator machinery; it introduces no new projectors and does not affect TT purity.

Summary (CK in the weave). CK is the vector $(A^{(\pm)})$ typed content of the τ–kairos channel (Definition 10.4). It is TT-null by construction and cannot contaminate the graviton readout (Definitions 10.3 and 10.25 and Lemma 10.123). Observables are recorded in typed, triad-resolved ledgers with the standard meters and audit triggers (Definitions 10.2, 10.82 and 10.95); internal mixing is orthogonalized via the $\bar{\Sigma}$–weighted generalized eigen-operator pair and whitening (Definitions 10.116 and 10.117). No additional primitives are introduced; this

[3] Earlier drafts used the same script $\mathcal{P}$ for both projectors and powers; the present convention removes that ambiguity.

section uses only the established typed families and audit/provenance scaffold, with notational conventions made consistent across projectors ($\mathcal{P}$), powers ($\mathcal{L}$), and parity diagnostics ($\Delta^{(1)}_{P,\Lambda}$ vs. $\widetilde{\Delta}^{(1)}_{P}$).

10.8 The Spin–0 Apeiron (Gaiaton): Scalar Channel in the τ-kairos

Spontaneous symmetry breaking (SSB) of the Gaiamiti potential organizes the physical spectrum into two distinguished responses on the realized weave configuration $\mathcal{I}^{[\mathcal{J}]}$ after Σ–realization and projection to the physical cohomology $H^0(\hat{Q}_{\mathrm{BRST}})$ (Definition 9.60). The resulting pair consists of a *massless spin-2* transverse–traceless (TT) Goldstone-like mode—identified with the graviton—and a *gapped spin-0* scalar excitation along the apeiron directions, termed the *Gaiaton* (Definition 10.70). This fits the standard SSB logic that gapless modes are protected by symmetry/identity structures (Ward/Slavnov–Taylor) and gauge cohomology (BRST), while orthogonal directions acquire masses from the curvature of the potential [94, 14, 483, 484, 445, 106]. Gauge–parameter independence of physical observables on certified plateaux is controlled by Nielsen identities (gauge-variation of the effective action) [120], which we enforce via the realization map and band/typed hygiene (Definitions 9.60 and 9.116).

On the *TT/Goldstone branch*, the ΦΦ (Gaiamiti) typed menu singles out the TT block $\Pi^{[\mathrm{TT}]}_{\widetilde{\Sigma}}$ (Definitions 9.195 and 10.3). Its two-point function supports a *massless spin-2 pole* realizing the radiative graviton degrees of freedom; TT purity (trace/divergence removal) ensures the pole carries only the physical tensor helicities $\lambda = \pm 2$ [43, 7, 14]. Masslessness is protected by the Ward/Slavnov–Taylor structure and BRST cohomology on the realized weave configuration, while TT-nullity and typed orthogonality prevent scalar contamination (Proposition 9.202 and Definitions 9.195 and 10.3).

On the *scalar/apeiron branch*, the τ–kairos *scalar* typed subspace $S^{(b)}, S^{(L)}, \tilde{S}^{(p)}$ captures fluctuations along the trace/apeiron directions. The second variation of the Gaiamiti potential in these directions produces a *massive spin-0 pole*. Operational isolation uses a band-limited Riesz selector applied to the scalar kinetic generalized eigen-operator pair $K^{(0)}_{\tau}$ within the admissible IR window fixed by the Chapter 10 band machinery (Remark 9.15), yielding a realized spectral mass $m_0 > 0$ and the associated physical projector [115, 253]. When invoked, CP resolution partitions the scalar ledger into CP(±) subchannels (as in Higgs-sector analyses [97, 96, 98, 467]), without modifying TT protection because the typed sandwich annihilates TT components by construction (Definition 9.116 and Proposition 9.202).

Taken together, these results are *triad-by-triad* and *typed-resolved* consequences of the Chapter 10 scaffold: TT masslessness follows from symmetry and gauge constraints encoded in the typed layer (Ward/Slavnov–Taylor, BRST, Nielsen), while the scalar mass is controlled by the curvature of the Gaiamiti potential in the apeiron directions [14, 483, 484, 445, 120]. Typed orthogonality, cone preservation, and TT-nullity (all fixed in Chapter 10) preclude TT–scalar mixing and guarantee that the pair {massless spin-2 graviton, massive spin-0 Gaiaton} is well-posed and realized weave configuration-invariant on certified plateaux (Definitions 9.60 and 9.116, Proposition 9.202, and Remark 9.15).

We work on the realized weave configuration $\mathcal{I}^{[\mathcal{G}]}$ of Definition 10.1 in the τ-kairos block, with triad projectors $\Pi^{(a)}$ from Definition 10.2, the *scalar* typed idempotents (CP-resolved) $S^{(\eta)}$ from Definition 10.4 with $\eta \in \{+,-\}$, and the TT projector $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ from Definition 10.3. All statements are taken on $\mathcal{H}_{\mathrm{phys}} = \operatorname{Ran} \mathsf{P}_{\mathrm{phys}}$ (BRST cohomological physical space) [108, 230, 14][4].

Definitions.

Definition 10.68 (Spin-0 spectral selector in the τ-scalar subspace)**.** Let $K_\tau^{(0)}$ denote the band-limited, gauge-fixed linearized kinetic operator restricted to the *scalar* subspace of the τ-kairos block.[5] Define the spin-0 Riesz selector, with $\{\zeta_m\}_{m\in\mathfrak{S}_0}$ disjoint contours enclosing the realized scalar spectral features (poles/bands) on the certified band, by

$$\mathsf{P}_{\mathrm{spec}}^{(0)} := \sum_{m\in\mathfrak{S}_0} \frac{1}{2\pi i} \oint_{\zeta_m} \left(K_\tau^{(0)} - z\right)^{-1} dz\,. \tag{10.47}$$

We also set the scalar spectral triad piece $\Pi_\tau^{(\mathrm{spec};0)} := \mathsf{P}_{\mathrm{spec}}^{(0)}$ for later triad bookkeeping.

Definition 10.69 (Gaiaton projectors and fields (CP- and triad-resolved))**.** For $\eta \in \{+,-\}$ (CP-even/odd), define the *Gaiaton projectors*

$$\boxed{\mathcal{P}_{\mathrm{G},\eta}^{(0)} := \mathsf{P}_{\mathrm{phys}}\, S^{(\eta)} \left(\Pi_\tau^{(\mathrm{spec};0)} \oplus \Pi^{(\mathrm{vac})} \oplus \Pi^{(\mathrm{corr})} \right) S^{(\eta)}\, \mathsf{P}_{\mathrm{phys}}} \tag{10.48}$$

acting on the τ-scalar carrier space. From a raw τ-scalar fluctuation $\delta\hat{\Phi}$ we extract the (CP, triad) Gaiaton fields

$$\boxed{\hat{\Phi}_\eta^{(a)} := \mathsf{P}_{\mathrm{phys}}\, S^{(\eta)}\, \Pi^{(a)}\, S^{(\eta)}\, \mathsf{P}_{\mathrm{phys}}\, \delta\hat{\Phi}, \qquad \hat{\Phi}_\eta := \sum_{a\in\{\mathrm{spec,vac,corr}\}} \hat{\Phi}_\eta^{(a)}} \tag{10.49}$$

with $\Pi^{(\mathrm{spec})}$ implicitly specialized to $\Pi_\tau^{(\mathrm{spec};0)}$ in the scalar channel.

Definition 10.70 (Gaiaton (spin-0 apeiron))**.** Within the τ–kairos scalar subspace (Definition 10.4), the *Gaiaton* is the CP- and triad-resolved scalar excitation selected by the spin-0 Riesz projector $\mathsf{P}_{\mathrm{spec}}^{(0)}$ of Definition 10.68 and carried by the CP projectors $S^{(\eta)}$ for $\eta \in \{+,-\}$. Concretely, the Gaiaton channel is the image of the *Gaiaton projectors*

$$\mathcal{P}_{\mathrm{G},\eta}^{(0)} \text{ of Definition 10.69}, \qquad \text{with fields } \hat{\Phi}_\eta^{(a)} \text{ and } \hat{\Phi}_\eta \text{ given by (10.49)}.$$

Its realized masses $m_{0,\eta}$ are the pole/band locations enclosed by the contours in (10.47), read from the typed, triad-resolved correlator blocks (Definitions 10.2 and 10.113) on the Σ-carrier (Definition 9.60).

[4] Applied to the physical projector $\mathsf{P}_{\mathrm{phys}}$, $\mathcal{H}_{\mathrm{phys}} = \operatorname{Ran} \mathsf{P}_{\mathrm{phys}}$ is the physical subspace (isomorphic to the BRST cohomology $H^0(\hat{Q}_{\mathrm{BRST}})$ in this setup). On $\operatorname{Ran} \mathsf{P}_{\mathrm{phys}}$ the projector acts as the identity; on $\ker \mathsf{P}_{\mathrm{phys}}$ it vanishes, and $\mathcal{H} = \operatorname{Ran} \mathsf{P}_{\mathrm{phys}} \oplus \ker \mathsf{P}_{\mathrm{phys}}$ (orthogonal direct sum, since $\mathsf{P}_{\mathrm{phys}} = \mathsf{P}_{\mathrm{phys}}^*$).

[5] Equivalently, one may use the band-limited physical two-point on the τ-scalar subspace. The admissible restriction is secured by the type projectors of Definition 10.4.

Lemma 10.71 (TT-orthogonality and typed safety)**.** *For all admissible η, $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\mathcal{P}_{\mathrm{G},\eta}^{(0)} = 0 = \mathcal{P}_{\mathrm{G},\eta}^{(0)}\,\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$; hence the Gaiaton channel does not mix with the TT graviton channel. Moreover, $\mathcal{P}_{\mathrm{G},\eta}^{(0)}$ commutes with the triad projectors $\Pi^{(\mathrm{spec})}, \Pi^{(\mathrm{vac})}, \Pi^{(\mathrm{corr})}$, so all meters/visibilities are triad-resolved and BRST/Nielsen-safe on certified plateaus.* Proof sketch. *TT-nullity of the τ–scalar types and triad-compatibility of typed projectors are enforced by the typed weave (Definitions 10.2 and 10.4) and the Chapter 10 typed layer; the conclusion follows by substituting the definitions of* (10.48) *and* (10.49)*.*

Proof. **(1) TT-orthogonality.** By construction (see the scalar typed family in Definition 10.4), the Gaiaton projector acts within the τ–scalar typed ranges and has the typed–sandwich form

$$\mathcal{P}_{\mathrm{G},\eta}^{(0)} \;=\; \mathsf{P}_{\mathrm{phys}}\, S^{(\eta)}\, \mathcal{R}^{(0)}\, S^{(\eta)}\, \mathsf{P}_{\mathrm{phys}}, \qquad \eta \in \{b, L, p, +, -\},$$

where $\mathcal{R}^{(0)}$ is a scalar Riesz/triad selector on the realized weave configuration (cf. (10.48), (10.49) and Definitions 10.2 and 10.7). The TT-null property of τ–types (Definition 10.4, equation (10.8)) gives

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} S^{(\eta)} = 0 = S^{(\eta)} \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}.$$

Using commutation of $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ with the realized weave configuration hygiene maps and spectral/triad selectors (Propositions 10.9 and 10.10), move $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ next to $S^{(\eta)}$ to obtain

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \mathcal{P}_{\mathrm{G},\eta}^{(0)} = \mathsf{P}_{\mathrm{phys}}\, (\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} S^{(\eta)})\, \mathcal{R}^{(0)}\, S^{(\eta)}\, \mathsf{P}_{\mathrm{phys}} = 0,$$

and similarly $\mathcal{P}_{\mathrm{G},\eta}^{(0)}\, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} = 0$. Therefore the Gaiaton (spin–0) channel is orthogonal to the TT graviton channel.

(2) Triad-compatibility. Write the scalar provenance split as $\mathcal{R}^{(0)} = \mathsf{P}_{\mathrm{spec}}^{(0)} \oplus \Pi^{(\mathrm{vac})} \oplus \Pi^{(\mathrm{corr})}$ (Definitions 10.2 and 10.7). By typed stability and functional calculus, each typed idempotent commutes with the triad projectors (Proposition 10.9), and the scalar Riesz selectors commute with the triad as well (Definition 10.7). Since $\mathsf{P}_{\mathrm{phys}}$ also commutes with admissible band/triad maps (Proposition 10.10), we have

$$\left[\Pi^{(a)},\, \mathcal{P}_{\mathrm{G},\eta}^{(0)}\right] = \mathsf{P}_{\mathrm{phys}} \left[\Pi^{(a)},\, S^{(\eta)} \mathcal{R}^{(0)} S^{(\eta)}\right] \mathsf{P}_{\mathrm{phys}} = 0, \qquad a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}.$$

(3) Consequences for meters/visibilities. Because the projector is TT-orthogonal and triad-compatible, any calibrated meter/visibility computed in the Gaiaton channel is automatically triad-resolved and free from TT contamination (Definition 10.82). BRST/Nielsen safety on certified plateaus follows from executing all operations on the realized physical realized weave configuration (Definitions 9.60 and 10.1) together with typed/triad commutation (Proposition 10.9) and band invariance (Proposition 10.10). This completes the proof. □

Definition 10.72 (CP action and visibility asymmetry (scalar))**.** Let $\mathcal{CP}_\tau$ denote the CP action on the τ-scalar subspace; it preserves the scalar carrier but may flip the CP label. Define the (law/state) *visibility* CP-asymmetry diagnostic by

$$\Delta_{\mathrm{CP},\Lambda}^{(0)} := \Lambda_{\mathrm{law/state}}^{[S^{(+)}]} - \Lambda_{\mathrm{law/state}}^{[S^{(-)}]}, \tag{10.50}$$

using the witnesses of Definitions 10.80 and 10.82.

Structural facts.

Lemma 10.73 (TT-nullity and orthogonality (scalar))**.** *One has* $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\big(\cdot\big) = 0$ *on any insertion confined to the* τ*-scalar subspace; moreover* $S^{(\eta)}S^{(\eta')} = \delta_{\eta\eta'}S^{(\eta)}$*.*

Proof. TT-nullity is immediate for scalars and follows from typed null-preservation in Definitions 10.3 and 10.4; orthogonality is part of the projector algebra in Lemma 10.8. □

Proposition 10.74 (Triad commutation and decomposition (scalar))**.** $[\Pi^{(a)}, S^{(\eta)}] = 0$ *for all* a, η*, hence the decomposition* (10.49) *is triad-orthogonal. Moreover,* $\hat{\Phi}_\eta = \hat{\Phi}_\eta^{(\mathrm{spec})} \oplus \hat{\Phi}_\eta^{(\mathrm{vac})} \oplus \hat{\Phi}_\eta^{(\mathrm{corr})}$*.*

Proof. Commutation is stated in Proposition 10.9; the decomposition is immediate. □

Proposition 10.75 (BRST/Nielsen flatness and band stability (scalar))**.** *Any correlator on* $\mathcal{H}_{\mathrm{phys}}$ *formed from* $\hat{\Phi}_\eta$ *and finitely many insertions of* $S^{(\eta)}$*,* $\Pi^{(a)}$*, and* $\mathsf{P}_{\mathrm{spec}}^{(0)}$ *is (i) gauge-parameter independent and (ii) stable under admissible retunings of* (P_x, K_{IR})*.*

Proof. A specialization of Proposition 10.10. □

Lemma 10.76 (Spectral future-proofing (spin-0))**.** *If additional scalar spectral branches appear (new contours* ζ_n *disjoint from the previously certified ones), the projector* $\mathsf{P}_{\mathrm{spec}}^{(0)}$ *enlarges by the corresponding Riesz terms and* $\mathcal{P}_{\mathrm{G},\eta}^{(0)}$ *automatically includes the new species.*

Proof. By definition (Definition 10.7), the scalar spectral selector is a sum of Riesz projectors

$$\mathsf{P}_{\mathrm{spec}}^{(0)} = \sum_{j\in\mathfrak{S}_0} \frac{1}{2\pi i}\oint_{\Gamma_j}\Big(K_{\Theta\Theta}^{(0)} - z\Big)^{-1}\,dz,$$

where the contours $\{\Gamma_j\}$ enclose the certified scalar spectral islands (measurability/invariance: Proposition 10.149 and Lemma 10.137). If new, disjoint scalar islands appear with enclosing contours $\{\zeta_n\}$, the holomorphic functional calculus implies *additivity* of Riesz projectors over pairwise disjoint spectral sets [115, 253]:

$$\mathsf{P}_{\mathrm{spec}}^{(0)} \;\longrightarrow\; \mathsf{P}_{\mathrm{spec,new}}^{(0)} = \mathsf{P}_{\mathrm{spec}}^{(0)} + \sum_n \frac{1}{2\pi i}\oint_{\zeta_n}\Big(K_{\Theta\Theta}^{(0)} - z\Big)^{-1}\,dz.$$

The Gaiaton projector has the typed–sandwich form (spin-0 provenance split; cf. (10.48), (10.49))

$$\mathcal{P}_{\mathrm{G},\eta}^{(0)} = \mathsf{P}_{\mathrm{phys}}\,S^{(\eta)}\Big(\mathsf{P}_{\mathrm{spec}}^{(0)} \oplus \Pi^{(\mathrm{vac})} \oplus \Pi^{(\mathrm{corr})}\Big)S^{(\eta)}\,\mathsf{P}_{\mathrm{phys}},$$

with $S^{(\eta)}$ a scalar typed idempotent (Definition 10.4) and the triad projectors from Definition 10.2. By commutation stability on the realized weave configuration (Propositions 10.9 and 10.10), replacing $\mathsf{P}_{\mathrm{spec}}^{(0)}$ by $\mathsf{P}_{\mathrm{spec,new}}^{(0)}$ propagates through the sandwich without further modification. Hence $\mathcal{P}_{\mathrm{G},\eta}^{(0)}$ *automatically* includes the newly admitted scalar species. □

Positivity, Ward identities, and diagnostics.

Proposition 10.77 (Scalar ledgers and spectral positivity)**.** *On $\mathcal{H}_{\mathrm{phys}}$, the scalar Gaiaton components obey the appropriate Ward/Slavnov ledgers for the scalar sector and have nonnegative spectral density $\rho_{(0)}(\omega, \mathbf{k}) \geq 0$.*

Proof (outline). Work on the realized, realized weave configuration with projection to the physical cohomology (Definitions 9.60 and 10.1). The Ward/Slavnov–Taylor identities hold on $\mathcal{H}_{\mathrm{phys}}$ for the realized correlators (upstream axioms unchanged; Remark 9.81), and gauge-parameter independence on certified plateaux is controlled by Nielsen identities; see [483, 484, 445, 106, 120]. Restricting to the scalar typed subspace via $S^{(\eta)}$ (Definition 10.4) preserves these ledgers because the typed idempotents commute with the admissible realized weave configuration/triad maps (Propositions 10.9 and 10.10). Thus the scalar Gaiaton components inherit the Ward/Slavnov constraints on $\mathcal{H}_{\mathrm{phys}}$.

For spectral positivity, start from the Källén–Lehmann (KL) representation for realized, BRST-clean two-point functions with the KL positivity condition (eq. (7.12); standard treatments include [14, 255, 230, 244]). Orthogonal projection by a $\bar{\Sigma}$–Hermitian idempotent preserves positivity of quadratic forms, hence $\rho \mapsto \rho_{(0)}$ remains nonnegative under the scalar typed projector $S^{(\eta)}$ (Definitions 9.33 and 10.4). Finally, the scalar spectral selector $\mathsf{P}^{(0)}_{\mathrm{spec}}$ is a sum of Riesz projectors (Definition 10.7), measurable and arena-stable (Proposition 10.149 and Lemma 10.137), which commute with the typed/triad structure (Proposition 10.9); selecting spectral islands by the holomorphic functional calculus leaves the positivity of the spectral measure intact [115, 253]. Therefore the Gaiaton scalar density obeys $\rho_{(0)}(\omega, \mathbf{k}) \geq 0$ on $\mathcal{H}_{\mathrm{phys}}$, as claimed. □

Definition 10.78 (CP-resolved visibilities (scalar))**.** Let $\mathbf{W}^{[\tau\text{-sca};S^{(\eta)}]}$ be the consolidated witness for the τ-scalar channel. Define $\Lambda^{[\tau\text{-sca};S^{(\eta)}]}_{\mathrm{law/state}}$ as in Definitions 10.80 and 10.82, evaluated on $\mathcal{H}_{\mathrm{phys}}$ with $\mathcal{I}^{[\mathscr{P}]}$.

Proposition 10.79 (Well-posed Parasymmetric TLO for the Gaiaton)**.** *For each CP sector $\eta \in \{+, -\}$, the Parasymmetric TLO triple* (Type $= S^{(\eta)}$, Location, Onset) *of Definition 10.85 and Proposition 10.90 exists and is unique on the certified band. The onset $r^\star[S^{(\eta)}]$ is stable under admissible retuning and tightening of K_{IR}.*

Proof. **(A) Setting and admissibility.** Work on the realized, realized weave configuration with projection to the physical cohomology (Definitions 9.60 and 10.1). Fix the scalar typed projectors $S^{(\eta)}$ (CP-resolved) from Definition 10.4 and note their $\bar{\Sigma}$–Hermitian, orthogonal idempotent algebra (10.5). Typed maps commute with the realized weave configuration hygiene and the triad (Proposition 10.9 and Definition 10.2), so restricting any calibrated meter to the typed image is admissible (Definitions 9.116 and 10.82). Band acceptance is enforced by the mask $\mathfrak{M}_{\mathrm{mask}}$ and guards C1–C5, which produce a nonempty IR window K_{IR} with the needed regularity on smoothed slopes and total variation (eq. (7.626), Lemma 7.386, Proposition 7.383, and Remark 9.15).

(B) Existence (TLO). By Proposition 10.90, the TLO triple exists whenever (i) the accepted window K_{IR} is nonempty and (ii) the location/onset meters satisfy the Lipschitz/-monotonicity bounds on K_{IR}. Both conditions hold after typed restriction because $S^{(\eta)}$

commutes with the acceptance machinery and does not degrade the slope/variation estimates (Definition 9.116 and Proposition 10.9); thus the parasymmetric TLO for Type $= S^{(\eta)}$ exists on the certified band.

(C) Uniqueness (typed orthogonality and one-threshold property). Orthogonality $S^{(+)}S^{(-)} = 0$ and completeness on the scalar typed block (Definition 10.4, (10.5)) imply that any TLO evaluated in $S^{(+)}$ cannot receive contributions from $S^{(-)}$, and vice versa. Within a fixed typed range, Proposition 10.90 gives uniqueness of the onset provided the onset meter is strictly monotone on K_{IR}, which is guaranteed by the acceptance/low-pass properties (Lemma 7.386 and Proposition 7.383). Hence $r^\star[S^{(+)}]$ and $r^\star[S^{(-)}]$ exist and are unique.

(D) Stability under retuning/tightening. Admissible retunes of the band tags and IR tightening of K_{IR} preserve the commutation and measurability properties on the realized weave configuration (Propositions 10.10 and 10.149 and Lemma 10.137). Since the onset functional is continuous (Lipschitz) in the meters used to define it (Definition 10.85 and Proposition 10.90) and typed restriction is an orthogonal projection that does not increase variation, the onset $r^\star[S^{(\eta)}]$ varies continuously and monotonically with admissible changes of K_{IR} (cf. Remark 9.15). Therefore $r^\star[S^{(\eta)}]$ is stable under admissible retuning and IR tightening.

(E) BRST/Nielsen safety. All constructions take place on $\mathcal{H}_{\mathrm{phys}} = \mathrm{Ran}\,\mathsf{P}_{\mathrm{phys}}$ via the realization map (Definition 9.60), so the derived TLO is BRST-clean and gauge-parameter independent on certified plateaux, in the sense of the upstream axioms (Remark 9.81; cf. [445, 106, 120]).

Combining (B)–(E) proves existence, uniqueness, and stability of the Parasymmetric TLO for the Gaiaton in each CP sector. □

Typed extraction and correlator blocks (scalar). For any admissible fluctuation/correlator, Gaiaton content is read CP-by-CP via the typed insertion (Definition 10.113). Writing $\hat{\mathbf{G}}_\Lambda$ for the unified inter-modal correlator on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$, define the scalar-typed blocks and their triad resolutions by

$$\hat{\mathbf{G}}_\Lambda^{[\,S^{(\eta)}\to S^{(\eta')}\,]} := \Pi^{[S^{(\eta)}]}\,\hat{\mathbf{G}}_\Lambda\,\Pi^{[S^{(\eta')}]}, \qquad \hat{\mathbf{G}}_\Lambda^{(a)\,[\,S^{(\eta)}\to S^{(\eta')}\,]} := \Pi^{(a)}\,\hat{\mathbf{G}}_\Lambda^{[\,S^{(\eta)}\to S^{(\eta')}\,]}\,\Pi^{(a)},$$

with $a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$ (Definition 10.2). Orthogonality and idempotence of the typed and triad projectors ensure that scalar channels can be audited provenance-by-provenance without mixing with TT.

Meters, visibilities, and ledgers (Gaiaton channel). To disambiguate notation, we reserve $\mathcal{P}$ for *projectors* (e.g., $\mathcal{P}^{(0)}_{\mathrm{G},\eta}$) and use

$$\mathcal{L}^{(a)}_{S^{(\eta)}} := \left\| \mathsf{P}_{\mathrm{phys}}\,\Pi^{[S^{(\eta)}]}\,\Pi^{(a)}\,\mathsf{P}_{\mathrm{phys}}\,\delta\hat{\Sigma} \right\|_{\mathcal{I}}$$

for the scalar *ledger power* in triad sector $a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$ and CP sector $\eta \in \{+,-\}$. The visibility-based CP diagnostic is $\Delta^{(0)}_{\mathrm{CP},\Lambda}$ from (10.50), while a normalized ledger asymmetry may be quoted as

$$\widetilde{\Delta}^{(0)}_{\mathrm{CP}} := \frac{\mathcal{L}_{S^{(+)}} - \mathcal{L}_{S^{(-)}}}{\mathcal{L}_{S^{(+)}} + \mathcal{L}_{S^{(-)}}}\,.$$

All meters are evaluated on $\mathcal{H}_{\mathrm{phys}}$ and inherit BRST/Nielsen flatness and band stability from the realized weave configuration (Definitions 10.82 and 10.95).

Spin selection, TT purity, and selection rules. Gaiaton content is spin–0 by typed construction; any scalar insertion is annihilated by the TT sandwich:

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\hat{\mathbf{G}}_{\Lambda}^{[\,S^{(\eta)}\to Y\,]} = 0 = \hat{\mathbf{G}}_{\Lambda}^{[\,Y\to S^{(\eta)}\,]}\,\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,.$$

Typed selection rules further imply that off-diagonal blocks that violate CP selection are null at leading order, and any symmetry breaking is quantified by the corresponding mixing norms (cf. the typed diagnostics in Definition 10.82).

Mixing, whitening, and reporting. Scalar channels may mix among CP sectors or with other admitted non-TT types, but never with TT under the typed sandwich. Mixing is quantified by the typed mixing norms and resolved by the $\bar{\Sigma}$–weighted operator-generalized eigen-operator pair eigenproblem and whitening (Definitions 10.116 and 10.117). The outcome is an orthogonal set of scalar eigen-channels, each carrying its triad provenance and reportable power/visibilities.

Dynamics (model-class placeholder, if needed). When a first-order representative Φ is adopted for phenomenology, Gaiaton dynamics may be modeled—*within the typed block*—by a Klein–Gordon–type equation on $(\bar{\Sigma}, \bar{\nabla})$,

$$\bar{\nabla}_{\mu}\Big(\bar{\Sigma}^{\mu\nu}\bar{\nabla}_{\nu}\Phi\Big) + m_{(0)}^{2}\,\Phi = J_{(0)}\,,$$

with an effective potential $V(\Phi)$ for interpretive layers (inflation/reheating) treated at the level of witnesses and meters; this introduces no new projectors and does not affect TT purity. Any CP-sensitive phenomenology is reported through $\Delta_{\mathrm{CP},\Lambda}^{(0)}$ and $\widetilde{\Delta}_{\mathrm{CP}}^{(0)}$ rather than raw field parameters.

Remarks (geometry/matter interface). The scalar Gaiaton lives in the τ-kairos scalar family of Definition 10.4; it modulates scalar coherence without contributing to the graviton TT readout. Interactions with vector ($A^{(\pm)}$) content (CK) or other scalar families are mediated only through typed, triad-resolved correlator blocks and are subject to the same audit/termination guarantees (No-Explosion) as in Propositions 10.10 and 10.90. No additional primitives are introduced; all statements use the established realized weave configuration, typed families, meters, and TLO scaffold.

The *Gaiaton* (spin-0 apeiron) is introduced as a typed, scalar excitation of the Gaiamiti sector, defined on the realized weave configuration $\mathcal{I}^{[\mathscr{P}]}$ and realized on the Σ-carrier $\bar{\Sigma}$ after projection to the physical cohomology $H^{0}(\hat{Q}_{\mathrm{BRST}})$ (Definitions 9.60 and 9.116). Ontologically it is not an independent sector but a member of the τ–kairos *scalar* typed family $S^{(b)}$, $S^{(L)}$, $\tilde{S}^{(p)}$ admitted via their idempotents $\Pi^{[\chi]}$ and inserted within the typed, triad-resolved correlator blocks of the unified inter-modal correlator (Definitions 10.2, 10.4 and 10.113). In contrast to the massless spin-2 transverse–traceless (TT) Goldstone mode of the Gaiamiti symmetry (the graviton) identified in the $\Phi\Phi$ typed menu (Definition 9.195 and Proposition 9.197), the Gaiaton corresponds to scalar fluctuations along the *apeiron* (trace/radial) directions—orthogonal to the vacuum manifold $\mathcal{M}_{\mathrm{vac}}$ (Definition 6.61). Its mass originates from the curvature of the Gaiamiti potential in these directions and is

therefore nonzero after spontaneous symmetry breaking [93, 94, 91]. *Phenomenologically*, this scalar channel aligns with well-studied gravitational scalars: in scalar–tensor gravity (Jordan–Brans–Dicke), a companion field ϕ yields a breathing/longitudinal GW polarization [480, 479, 477, 478]; in higher-curvature $f(R)$ gravity, dynamics are equivalent to GR plus a massive "scalaron" [244, 479]; and in massive/bimetric gravity, the helicity-0 mode of the spin-2 field (with its decoupling-limit Galileon avatar) provides a scalar gravitational excitation in ghost-free formulations [113, 481]. These map to the detector taxonomy of "extra scalar GW polarization(s)" (two tensor + possible vector/scalar types), so the **Gaiaton** is naturally read as the theory's typed, CP-refined representative of that scalar gravitational sector [477, 478, 479].

Formally, the scalar content of the τ–kairos block may be represented by a typed scalar representative whose effective potential governs the radial displacement from the vacuum configuration. In this representation the local effective potential on the scalar block takes the quartic form

$$\mathcal{V}_\Phi(\varphi) \;=\; \tfrac{1}{2}\, m_A^2\, \varphi^2 \;+\; \tfrac{\lambda_A}{4!}\, \varphi^4,$$

with m_A^2 fixed by the curvature about $\mathcal{M}_{\mathrm{vac}}$ and λ_A a self-interaction strength [91]. The realized scalar mass $m_A > 0$ is extracted *operationally* from the typed, triad-resolved spectrum using the Riesz selector acting on the scalar kinetic generalized eigen-operator pair $K_\tau^{(0)}$ within the admissible IR window—yielding the physical projector $\mathsf{P}^{(0)}_{\mathrm{spec}}$ and the corresponding pole location (Definition 10.68 and Remark 9.15). TT purity is guaranteed throughout by typed orthogonality and TT-nullity: $\Pi_{\bar\Sigma}^{[\mathrm{TT}]}$ annihilates τ-scalar contributions, forbidding TT–scalar mixing (Proposition 9.202 and Lemma 10.123).

Couplings of the Gaiaton are geometric rather than force-mediating in the conventional sense. The scalar mode modulates the *inertial rigidity* of the vacuum—implemented in practice through calibrated meters and visibilities on $\mathcal{H}_{\mathrm{phys}}$ that quantify the scalar ledger power and its response under band-limited probes (Definitions 10.82 and 10.117). Increased scalar ledger power corresponds to enhanced stiffness of the $\bar\Sigma$-diagnostic realized reference configuration, while reduced power corresponds to greater susceptibility, all within the BRST/Nielsen-safe plateau regime (Proposition 9.97 and Definition 10.1); this viewpoint aligns with the general quantum-geometric picture [8, 3].

At cosmological scales, the same scalar block provides a natural *inflaton* candidate. In an early-universe regime where the energy density is dominated by $\mathcal{V}_\Phi$, slow-roll conditions can be encoded directly in the typed potential and the $\bar\Sigma$-weighted inner product, producing an accelerated expansion epoch consistent with quartic slow-roll templates (cf. the potential form above) [89, 485]. Termination of inflation and reheating are governed by the *interaction action* obtained from the off-diagonal entries of the Fourfold inter-modal structure,

$$S_{\mathrm{int}}\big[\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}\big] \;=\; \int d^d x\, d^d x'\, \mathrm{Tr}\big[\hat{\mathbf{\Phi}}^\dagger(x)\, \mathcal{U}_{\Phi\Theta}(x,x')\, \hat{\mathbf{\Theta}}(x')\big]\,,$$

where $\mathcal{U}_{\Phi\Theta}$ is the typed, nonlocal kernel linking the Gaiamiti (Φ) and Tantrakyma/Kymon (Θ) sectors (Definitions 9.357 and 10.223); the associated reheating picture is standard in scalar-field cosmology [485, 91]. This coupling allows the *decay channel* Gaiaton $\to$ Kymon+Kymon and thereby converts geometric vacuum energy into matter excitations within the Θ sector, in line with the typed weave's inter-modal consistency.

Residual scalar potential energy after reheating yields a natural *late-time* component. If the scalar representative remains displaced from the minimum or if $\mathcal{V}_\Phi$ exhibits near-degenerate minima, the residual energy density

$$\rho_\Lambda^{(\mathrm{eff})} \;\sim\; \mathcal{V}_\Phi\Big[\bar{\Phi}_{\min}\Big]$$

acts as an effective cosmological constant and supports a quintessence-like description when allowed to vary adiabatically over cosmological timescales [486, 485]. Both cases fit within the typed ledger and retain Nielsen/realized weave configuration invariance, since the relevant meters and selectors are defined on $\mathcal{H}_{\mathrm{phys}}$ with certified IR tags (Definition 10.1 and Remark 9.15).

Conceptually, the Gaiaton thereby unifies three narratives within the same formal scaffold: (i) sub-Planckian *quantum rigidity* of spacetime through scalar ledger power on $\bar{\Sigma}$; (ii) *inflationary dynamics* via the typed scalar potential and inter-modal reheating into Θ; and (iii) *late-time acceleration* through residual vacuum energy. All elements descend from the Gaiamiti typed layer established in Chapter 10 and operate within the Quantum Holo-State $\big|\mathring{\mathbf{\Psi}}\big\rangle$ framework that governs emergent properties at each PQS locus (Definition 1.27), while the apeiron lineage evokes the original metaphysical notion of the *ápeiron* [69].

10.9 Witnesses, Visibilities, and Parasymmetric TLO (Full Construction)

We complete Step (4) of the operational pipeline by specifying the *explicit* witnesses and meters, and by proving finiteness, BRST/Nielsen flatness, monotonicity under admissible IR retuning, and Onset uniqueness with stability radii (cf. Ch. 10: Definitions 9.116, 9.133 and 9.147 and eq. (9.81)). We work on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ of Definition 10.1; triad maps $\Pi^{(a)}$ are from Definition 10.2; the admitted typed projectors we instantiate here include TT, τ–kairos, Higgs, and Chrono–Kymon (Definitions 10.3 to 10.5).

Global setup. For any admitted type $[\chi]$ (e.g. TT, $S^{(b)}$, $S^{(L)}$, $\tilde{S}^{(p)}$, $A^{(\pm)}$, rad$\pm$) and *sector*

$$\mathscr{S} \in \{\Phi\Phi, \Theta\Theta, \Phi\Theta\},$$

define the typed *physical* observable

$$\mathfrak{O}^{[\mathscr{S};\chi]} \;:=\; \mathsf{P}_{\mathrm{phys}}\, \Pi^{[\chi]}\, \mathfrak{O}\, \Pi^{[\chi]}\, \mathsf{P}_{\mathrm{phys}},$$

where the sector block $\mathscr{S}$ is taken in the unified inter–modal correlator and its typed decomposition (see (10.61)). All norms $\|\cdot\|_{\mathcal{I}}$ below are the fixed, band–invariant norms chosen once for this chapter. BRST/Nielsen flatness and gauge–parameter independence of the resulting meters follow from the invariance statements (e.g. Propositions 10.30 and 10.61) together with the gauge–independence remark Remark 10.93, and will be used tacitly.

10.9.1 Law and State Ledgers: Residuals and Features

Definition 10.80 (Law residuals and State features)**.** Let $\mathscr{S} \in \{\Phi\Phi, \Theta\Theta, \Phi\Theta\}$ be a sector block of the unified inter–modal correlator (cf. (10.61)), and let $[\chi]$ be any *admitted type* from the typed layer (Ch. 10; e.g. TT, $S^{(b)}$, $S^{(L)}$, $\tilde{S}^{(p)}$, $A^{(\pm)}$, rad±, cf. Definition 9.116 and its weave instantiations Definitions 10.3 to 10.5). On the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ (Def. Definition 10.1), define the *typed physical observable*

$$\mathfrak{O}^{[\mathscr{S};\chi]} \;:=\; \mathsf{P}_{\mathrm{phys}}\, \Pi^{[\chi]}\, \mathfrak{O}\, \Pi^{[\chi]}\, \mathsf{P}_{\mathrm{phys}},$$

and write $\mathfrak{O} = \mathfrak{O}^{[\mathscr{S};\chi]}$ for brevity. All norms $\|\cdot\|_{\mathcal{I}}$ are the fixed band–invariant norms for this chapter (cf. invariance discussion and admissible norms in Proposition 10.10 and Remark 10.93).

Law-residual vector. The *Law-residual vector* $\mathbf{R}^{[\mathscr{S};\chi]} \in \mathbb{R}_+^{N_{\mathrm{law}}}$ collects type-specific constraints that must vanish on $\mathcal{H}_{\mathrm{phys}}$ (Ch. 10 cohomological EOM and Ward/Slavnov, Definition 9.63):

$$\mathbf{R}^{[\mathscr{S};\chi]} = \Bigg(\underbrace{\|[\hat{Q}_{\mathrm{BRST}}, \mathfrak{O}]\|_{\mathcal{I}}}_{\text{BRST closure (Ch. 8: Definition 7.11 and eq. (7.8))}}, \quad \underbrace{\|\nabla^{(\mathrm{rel})\mu}\mathfrak{O}_{\mu\nu}\|_{\mathcal{I}}}_{\text{Ward transversality (Ch. 7: Definition 6.7 and eq. (6.22))}}, \quad \underbrace{\|\mathrm{tr}_{\bar{\Sigma}}\mathfrak{O}\|_{\mathcal{I}}}_{\text{trace test (if applicable; Ch. 6: Definition 5.26)}}, \quad \underbrace{\left\|\partial_\xi \mathcal{G}_\xi^{[\mathscr{S};\chi]}\right\|_{\mathcal{I}}}_{\text{Nielsen flatness (Ch. 8/9: Definition 7.59 and eq. (7.16))}}, \ldots \Bigg) \tag{10.51}$$

where $\nabla^{(\mathrm{rel})}$ is the $\bar{\Sigma}$-compatible Levi–Civita derivative (Chs. 6–7), $\mathrm{tr}_{\bar{\Sigma}}$ contracts indices with $\bar{\Sigma}_{\mu\nu}$ (Ch. 6), ξ is a gauge parameter, and $\mathcal{G}_\xi^{[\mathscr{S};\chi]}$ denotes the typed two-point/kinetic resolvent on the certified band used for meters (Ch. 10: Definition 9.133). By the gauge-independence remark and the type-level invariance propositions (e.g. graviton TT and CK: Propositions 10.30 and 10.61 and Remark 10.93), ∂_ξ–residuals vanish on shell in $\mathcal{H}_{\mathrm{phys}}$.

State-feature vector. The *State-feature vector* $\mathbf{S}^{[\mathscr{S};\chi]} \in \mathbb{R}_+^{N_{\mathrm{state}}}$ measures radiative/vacuum/-coherence content on the triad (weave triad maps Definition 10.2):

$$\mathbf{S}^{[\mathscr{S};\chi]} = \Bigg(\underbrace{\int W_{\mathrm{spec}}(\omega,\mathbf{k})\; \rho_{(s)}^{[\mathscr{S};\chi]}(\omega,\mathbf{k})\, d\mu}_{\text{spectral weight (Ch. 8 KL positivity: } eq.\ (7.12))}, \quad \underbrace{\|\Pi^{(\mathrm{vac})}\mathfrak{O}\|_{\mathcal{I}}}_{\text{vacuum amplitude}}, \quad \underbrace{\|\Pi^{(\mathrm{corr})}\mathfrak{O}\|_{\mathcal{I}}}_{\text{coherence amplitude}}, \ \ldots \Bigg), \tag{10.52}$$

where $s \in \{2,1,0\}$ is the *typed spin* selected by $\Pi^{[\chi]}$ within the chosen sector block $\mathscr{S}$, W_{spec} is a fixed band-admissible window (Ch. 8 band functor and IR window: Definitions 6.280 and 7.29), $\rho_{(s)}^{[\mathscr{S};\chi]}$ is the corresponding typed spectral density (Källén–Lehmann form; Ch. 8:

eq. (7.12)), and $\Pi^{(\mathrm{vac})}, \Pi^{(\mathrm{corr})}$ are the triad projectors (Def. Definition 10.2). For mixed $\Phi\Theta$, the typed spin s is determined by the *typed insertion* used in the off-diagonal correlator block (cf. (10.61)); τ–kairos may appear here only as a *typed projector* (not as a sector), via Definition 10.4.

Remark 10.81 (Type-specific components)**.** Beyond the universal entries in (10.51)–(10.52), we append the following *type-specific* components (typed projectors as in Ch. 10 and their weave instantiations; cf. Definitions 9.116 and 10.3 to 10.5). All quantities are evaluated on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ (Def. Definition 10.1) and on the physical carrier $\mathcal{H}_{\mathrm{phys}}$.

- **Graviton (TT in $\Phi\Phi$).** Add the TT-purity residual

$$\left\| \Pi_{\bar{\Sigma}}^{\perp[\mathrm{TT}]}\, \mathfrak{O} \right\|_{\mathcal{I}} \quad \text{with} \quad \Pi_{\bar{\Sigma}}^{\perp[\mathrm{TT}]} := \mathbf{1} - \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \quad (\text{see } \textit{Definition } 10.3),$$

  to $\mathbf{R}^{[\mathscr{S};\chi]}$, and include the *TT-leak* diagnostic ε from Definition 10.32 in $\mathbf{S}^{[\mathscr{S};\chi]}$. (Invariance/flatness: Proposition 10.30.)
- **τ-kairos scalars $S^{(b)}$, $S^{(L)}$, $\tilde{S}^{(p)}$ (typed, not a sector).** Enforce TT-nullity by adding

$$\left\| \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \mathfrak{O} \right\|_{\mathcal{I}} \quad (\text{TT-annihilation; } \textit{Definition } 10.3 \textit{and Lemma } 10.17),$$

  and, for the pseudoscalar $\tilde{S}^{(p)}$, append a parity-odd check via the Levi–Civita contraction consistent with the τ-parity conventions in Definition 10.4 (see also the parity diagnostic pattern used for CK in Definition 10.58).
- **Chrono–Kymon $A^{(\pm)}$ (typed helicities in τ-vector).** Add the helicity-purity residual

$$\left\| \left(\mathbf{1} - A^{(\pm)}\right) \mathfrak{O} \right\|_{\mathcal{I}} \quad (\text{helicity projectors } A^{(\pm)} \text{ in } \textit{Definition } 10.4),$$

  and include the vector parity asymmetry Δ_P from Definition 10.58 among the state features. (CK invariance/flatness: Proposition 10.61.)
- **Higgs** rad$\pm$ **(radial tantrakyma in $\Theta\Theta$).** Enforce NG/pNG orthogonality by adding

$$\left\| \Pi_{\Theta}^{[\mathrm{NG/pNG}]}\, \mathfrak{O} \right\|_{\mathcal{I}} \quad (\text{orthogonality: } \textit{Proposition } 10.47),$$

  and append CP-purity via

$$\left\| \Pi^{(\mathrm{CP}\mp)}\, \mathfrak{O} \right\|_{\mathcal{I}} \quad (\text{CP projectors in } \textit{Definition } 10.5).$$

Scope across sectors. For the mixed interaction block $\mathscr{S} = \Phi\Theta$, all checks above are applied after the *typed insertion* into the off-diagonal correlator block of the unified inter-modal correlator (cf. (10.61)); τ-kairos enters here only as a *typed projector family* (not as a sector), per Definition 10.4.

10.9.2 Calibrated Meters and Visibilities

Definition 10.82 (Calibration maps and visibilities)**.** Let $\mathscr{S} \in \{\Phi\Phi, \Theta\Theta, \Phi\Theta\}$ be a sector block of the unified inter–modal correlator (cf. (10.61)), and let $[\chi]$ be any admitted type (Ch. 10; Definition 9.116, instantiated in Definitions 10.3 to 10.5). Given the Law/State ledgers $\mathbf{R}^{[\mathscr{S};\chi]}, \mathbf{S}^{[\mathscr{S};\chi]}$ from Definition 10.80, fix positive weights $\boldsymbol{\alpha} \in \mathbb{R}_+^{N_{\text{law}}}$, $\boldsymbol{\beta} \in \mathbb{R}_+^{N_{\text{state}}}$ and choose *calibration maps*

$$\begin{aligned} &\text{law: } f_j : \mathbb{R}_+ \to (0,1], \quad f_j(x) = e^{-\alpha_j x} \ \text{(strictly } \textit{decreasing}, C^1), \\ &\text{state: } g_m : \mathbb{R}_+ \to [0,1), \quad g_m(x) = \frac{\beta_m x}{1+\beta_m x} \ \text{(strictly } \textit{increasing}, C^1). \end{aligned}$$

Define the *calibrated* vectors

$$\mathbf{L}^{[\mathscr{S};\chi]} := \Big(f_j(R_j^{[\mathscr{S};\chi]})\Big)_{j=1}^{N_{\text{law}}}, \qquad \mathbf{V}^{[\mathscr{S};\chi]} := \Big(g_m(S_m^{[\mathscr{S};\chi]})\Big)_{m=1}^{N_{\text{state}}}, \tag{10.53}$$

and the *law/state visibilities* (cosine meters; cf. Definition 10.80):

$$\Lambda_{\text{law}}^{[\mathscr{S};\chi]} = \frac{\langle \mathbf{L}^{[\mathscr{S};\chi]},\ \mathbf{1}\rangle}{\|\mathbf{L}^{[\mathscr{S};\chi]}\|\,\|\mathbf{1}\|} \in (0,1], \qquad \Lambda_{\text{state}}^{[\mathscr{S};\chi]} = \begin{cases} \dfrac{\langle \mathbf{V}^{[\mathscr{S};\chi]},\ \mathbf{1}\rangle}{\|\mathbf{V}^{[\mathscr{S};\chi]}\|\,\|\mathbf{1}\|}, & \mathbf{V}^{[\mathscr{S};\chi]} \neq \mathbf{0}, \\ 0, & \mathbf{V}^{[\mathscr{S};\chi]} = \mathbf{0} \ \text{(convention)}. \end{cases} \tag{10.54}$$

Here $\mathbf{1}$ is the all-ones vector of the appropriate length. Larger values mean better law compliance / stronger certified state signal. All objects live on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ (Def. Definition 10.1) and on the physical carrier $\mathcal{H}_{\text{phys}}$, with band/BRST hygiene as in Proposition 10.10 and Remark 10.93.

Lemma 10.83 (Finiteness of meters)**.** *For every admitted* $[\chi]$ *and sector* $\mathscr{S} \in \{\Phi\Phi, \Theta\Theta, \Phi\Theta\}$*, the visibilities* $\Lambda_{\text{law}}^{[\mathscr{S};\chi]}$ *and* $\Lambda_{\text{state}}^{[\mathscr{S};\chi]}$ *are finite and well defined on* $\mathcal{H}_{\text{phys}}$*.*

Proof. Each component $R_j^{[\mathscr{S};\chi]}$ and $S_m^{[\mathscr{S};\chi]}$ is, by construction, a band-limited norm of a BRST-sandwiched operator (see Definition 10.80; band functor and IR window in Ch. 8: Definitions 6.280 and 7.29), hence finite on the realized weave configuration by Proposition 10.10. Therefore $f_j(R_j) \in (0,1]$ and $g_m(S_m) \in [0,1)$, so $\mathbf{L}^{[\mathscr{S};\chi]}$ is non-zero and $\mathbf{V}^{[\mathscr{S};\chi]}$ is either non-zero or (if all state features vanish) we apply the stated convention $\Lambda_{\text{state}}^{[\mathscr{S};\chi]} = 0$. In both cases the cosine with $\mathbf{1}$ is well defined, establishing finiteness. □

What the meters measure (and why they are calibrated). Definition Definition 10.82 turns *raw ledgers*—feature vectors extracted from a chosen sector block $\mathscr{S} \in \{\Phi\Phi, \Theta\Theta, \Phi\Theta\}$ of the unified inter–modal correlator (cf. (10.61)) and a selected type $[\chi]$ (Ch. 10; Definition 9.116, instantiated in Definitions 10.3 to 10.5)—into *unitless, bounded observables*. The "law" ledger $\mathbf{R}^{[\mathscr{S};\chi]}$ aggregates constraint/axiom compliance (e.g., conservation, Ward/Slavnov consistency) while the "state" ledger $\mathbf{S}^{[\mathscr{S};\chi]}$ aggregates physically certified signal features (Definition 10.80). Because these entries can carry different scales and units, we pass them through monotone, C^1 *calibration maps*: a strictly decreasing exponential $f_j(x) = e^{-\alpha_j x} \in (0,1]$ for law-terms (larger violation ⇒ smaller score), and a strictly increasing saturating map $g_m(x) = \frac{\beta_m x}{1+\beta_m x} \in [0,1)$

for state-terms (stronger signal ⇒ larger score). The result is the pair of calibrated vectors $(\mathbf{L}^{[\mathscr{S};\chi]}, \mathbf{V}^{[\mathscr{S};\chi]})$ in (10.53), which preserve the ordering of performance while compressing dynamic range and improving robustness to outliers; the positive weights $(\boldsymbol{\alpha}, \boldsymbol{\beta})$ control sensitivity per feature.

Visibilities as cosine meters. The visibilities in (10.54) are *cosines* between the calibrated vectors and the all-ones direction $\mathbf{1}$. Thus, $\Lambda_{\text{law}}^{[\mathscr{S};\chi]} \in (0,1]$ quantifies how *uniformly* the law features meet their targets (near 1 means broad compliance across entries), while $\Lambda_{\text{state}}^{[\mathscr{S};\chi]} \in [0,1]$ quantifies how *coherently* the state features register a certified signal (the zero convention covers the trivial case $\mathbf{V}^{[\mathscr{S};\chi]} = \mathbf{0}$). Because these are angles in a Euclidean ledger space, the meters are dimensionless, comparable across types and sectors, and naturally bounded—properties that are used repeatedly when comparing TT, τ–scalar, and Higgs-CP channels side-by-side (Definitions 10.3 to 10.5).

Well-posedness and hygiene. All objects live on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ and the physical carrier $\mathcal{H}_{\text{phys}}$ (Definition 10.1), with band/BRST hygiene ensured by the acceptance and invariance statements (Proposition 10.10 and Remark 10.93). Each ledger component is built from *band-limited* norms of BRST-sandwiched operators (Ch. 8: Definitions 6.280 and 7.29), guaranteeing finiteness; consequently the calibrated maps land in $(0,1]$ or $[0,1)$ and the cosines are well defined—this is recorded in Lemma 10.83. Because the admitted types $[\chi]$ commute with the realized weave configuration/triad machinery (Definitions 9.116 and 10.2 and Proposition 10.9), the meters are *typed-resolved* without order-of-operations ambiguity: projecting to TT, τ–scalar, or CP-refined Higgs channels before/after calibration yields the same visibilities. In short, Definition 10.82 provides a uniform, gauge- and band-safe way to score law compliance and physical signal strength across sectors and types, and Lemma 10.83 ensures these scores are always finite on $\mathcal{H}_{\text{phys}}$.

10.9.3 Monotonicity Under IR Retuning

Proposition 10.84 (Monotonicity of meters under IR tightening)**.** *Let $\mathscr{S} \in \{\Phi\Phi, \Theta\Theta, \Phi\Theta\}$ and let $r \mapsto K_{\text{IR}}^{(r)}$ be an admissible IR knob on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ (Def. Definition 10.1; IR window and band functor in Ch. 8: Definitions 6.280 and 7.29). For any admitted type $[\chi]$ and the corresponding calibrated visibilities $\Lambda_{\text{law/state}}^{[\mathscr{S};\chi]}(r)$ (Def. Definition 10.82), the following hold for* IR tightening *(increasing r):*

(i) $\Lambda_{\text{law}}^{[\mathscr{S};\chi]}(r)$ *is* nondecreasing *in r.*

(ii) $\Lambda_{\text{state}}^{[\mathscr{S};\chi]}(r)$ *is* nonincreasing *for the* vac *and* corr *components, and* nondecreasing *for the* spec *component measured with a fixed spectral window W_{spec} (cf. eq. (10.52)).*

Proof. All objects are evaluated on the physical carrier $\mathcal{H}_{\text{phys}}$ with band/BRST hygiene as in Proposition 10.10 and Remark 10.93. The IR window $K_{\text{IR}}^{(r)}$ is positive, order-preserving, and L^p-contractive on the certified band (Ch. 7/8; see Definition 6.280), and composes monotonically in r (admissible knob on $\mathcal{I}^{[\mathscr{S}]}$, Def. Definition 10.1).

(1) Law meter is nondecreasing. Each law entry $R_j^{[\mathscr{S};\chi]}(r)$ in eq. (10.51) is a norm of a constraint violation (BRST commutator, relational divergence, $\bar{\Sigma}$-trace, Nielsen derivative) applied to the IR-filtered typed observable $\mathcal{O}^{[\mathscr{S};\chi]}(r)$. Since $K_{\mathrm{IR}}^{(r)}$ is contractive and smoothing, tightening (larger r) yields

$$R_j^{[\mathscr{S};\chi]}(r_2) \;\le\; R_j^{[\mathscr{S};\chi]}(r_1) \quad \text{for } r_2 \ge r_1.$$

With the calibration $f_j(x) = e^{-\alpha_j x}$ (Def. Definition 10.82), which is strictly decreasing, we get $f_j(R_j(r_2)) \ge f_j(R_j(r_1))$ componentwise, hence $\mathbf{L}^{[\mathscr{S};\chi]}(r_2) \ge \mathbf{L}^{[\mathscr{S};\chi]}(r_1)$ entrywise. For positive vectors, the cosine with $\mathbf{1}$, $\cos(\mathbf{L},\mathbf{1}) = \langle \mathbf{L},\mathbf{1}\rangle/(\|\mathbf{L}\|\|\mathbf{1}\|)$, is monotone with respect to entrywise increases; therefore $\Lambda_{\mathrm{law}}^{[\mathscr{S};\chi]}(r)$ is nondecreasing.

(2) State meter splits by triad channel. State features in eq. (10.52) decompose on the triad (Def. Definition 10.2): vacuum $\|\Pi^{(\mathrm{vac})}\mathcal{O}\|_{\mathcal{I}}$, coherence $\|\Pi^{(\mathrm{corr})}\mathcal{O}\|_{\mathcal{I}}$, and a spectral weight $\int W_{\mathrm{spec}}\, \rho_{(s)}^{[\mathscr{S};\chi]}\, d\mu$ with fixed W_{spec} and KL-positive density $\rho_{(s)}$ (Ch. 8: eq. (7.12)). Tightening $K_{\mathrm{IR}}^{(r)}$ suppresses long-distance content: it contracts the vacuum amplitude and damps off-diagonal/coherence contributions (order-preserving L^p-contractivity), hence

$$S_m^{(\mathrm{vac})}(r_2) \le S_m^{(\mathrm{vac})}(r_1), \qquad S_m^{(\mathrm{corr})}(r_2) \le S_m^{(\mathrm{corr})}(r_1) \quad (r_2 \ge r_1).$$

As $g_m(x) = \beta_m x/(1+\beta_m x)$ is strictly increasing (Def. Definition 10.82), the calibrated entries $g_m(S_m)$ are nonincreasing in r for these channels, and so is their cosine meter contribution.

For the spec feature, with a *fixed* W_{spec}, the IR tightening removes ultra-IR leakage but preserves (and can sharpen) the weight inside the fixed spectral window by KL-positivity and band covariance; thus $S_m^{(\mathrm{spec})}(r_2) \ge S_m^{(\mathrm{spec})}(r_1)$ is admissible (nondecrease). Applying the increasing calibration g_m gives a nondecreasing calibrated spectral entry and hence a nondecreasing contribution to $\Lambda_{\mathrm{state}}^{[\mathscr{S};\chi]}(r)$.

The cosine aggregation preserves these one-sided monotonicities because all calibrated vectors remain in the positive orthant (Def. Definition 10.82). As an illustrative special case, the TT-leak diagnostic ε (Def. Definition 10.32) obeys the same monotone decrease under IR tightening (Cor. Corollary 10.104), consistent with the coherence channel's trend. □

10.9.4 Onset Definition, Uniqueness, and Stability Radii

Definition 10.85 (Onset thresholding and certified onset)**.** Let $\mathscr{S} \in \{\Phi\Phi, \Theta\Theta, \Phi\Theta\}$ be a sector block of the unified inter–modal correlator (cf. (10.61)), and let $[\chi]$ be any admitted type from the typed layer (Ch. 10; Definition 9.116, instantiated in Definitions 10.3 to 10.5). Fix thresholds $\theta_{\mathrm{law}}, \theta_{\mathrm{state}} \in (0,1)$ and consider the calibrated visibilities $\Lambda_{\mathrm{law}}^{[\mathscr{S};\chi]}(r)$ and $\Lambda_{\mathrm{state}}^{[\mathscr{S};\chi]}(r)$ of Definition 10.82 with (10.54), evaluated on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ (Def. Definition 10.1).

Admission set. Define

$$\mathcal{A}_{[\chi]} \;:=\; \Big\{ r \;\Big|\; \Lambda_{\mathrm{law}}^{[\mathscr{S};\chi]}(r) \;\ge\; \theta_{\mathrm{law}} \;\text{ and }\; \Lambda_{\mathrm{state}}^{[\mathscr{S};\chi]}(r) \;\ge\; \theta_{\mathrm{state}} \Big\}.$$

Certified onset. The (first) certified onset is

$$r^\star[\chi] \;:=\; \inf \mathcal{A}_{[\chi]} \qquad \big(r^\star[\chi] = +\infty \text{ if } \mathcal{A}_{[\chi]} = \varnothing\big). \tag{10.55}$$

Canonical masking for monotonicity. Unless stated otherwise, the *state* visibility entering $\mathcal{A}_{[\chi]}$ is taken with the canonical spectral mask from Ch. 8 (fixed thresholds; Definition 7.665), i.e. only the spectral entry of (10.52) participates in the onset test. With this mask, both meters are one–sided monotone under IR tightening (Prop. 9.146), ensuring uniqueness below.

Lemma 10.86 (Interval property and uniqueness)**.** *With the canonical spectral mask (Ch. 8: Definition 7.665) applied to the state meter, $\Lambda_{\mathrm{law}}^{[\mathscr{S};\chi]}(r)$ and the masked $\Lambda_{\mathrm{state}}^{[\mathscr{S};\chi]}(r)$ are nondecreasing in r (Proposition 9.146). Hence $\mathcal{A}_{[\chi]}$ is either empty or an interval of the form $[r^\star[\chi],\infty)$, and the onset (10.55) is unique whenever $\mathcal{A}_{[\chi]}\neq\varnothing$.*

Proof. **Step 1: Monotonicity of the law meter.** By Proposition 9.146(1), the calibrated law visibility $\Lambda_{\mathrm{law}}^{[\mathscr{S};\chi]}(r)$ defined in (10.54) (with calibrations from Definition 10.82) is nondecreasing in the tuning parameter r on the certified window (band hygiene Proposition 10.10).

Step 2: Monotonicity of the masked state meter. The canonical spectral mask of Definition 7.665 is an *isotone* (order-preserving) thresholding on spectral content, with thresholds that are themselves nondecreasing in r by construction. By Proposition 9.146(2), the (pre-mask) state visibility $\Lambda_{\mathrm{state}}^{[\mathscr{S};\chi]}(r)$ is nondecreasing; composing with an isotone mask preserves nondecreasing order, hence the *masked* state meter $\widetilde{\Lambda}_{\mathrm{state}}^{[\mathscr{S};\chi]}(r)$ is also nondecreasing in r.

Step 3: Upper-set structure of the acceptance region. Let the acceptance set be

$$\mathcal{A}_{[\chi]} := \Big\{ r :\ \Lambda_{\mathrm{law}}^{[\mathscr{S};\chi]}(r)\ \geq\ \tau_{\mathrm{law}}\,,\ \ \widetilde{\Lambda}_{\mathrm{state}}^{[\mathscr{S};\chi]}(r)\ \geq\ \tau_{\mathrm{state}} \Big\},$$

with fixed thresholds $(\tau_{\mathrm{law}},\tau_{\mathrm{state}})$ specified by Definition 7.665. If $r_1\in\mathcal{A}_{[\chi]}$ and $r_2\geq r_1$, monotonicity from Steps 1–2 implies both inequalities remain satisfied at r_2, so $r_2\in\mathcal{A}_{[\chi]}$. Thus $\mathcal{A}_{[\chi]}$ is an *upper set* in $\mathbb{R}$, hence either empty or of the form $[r^\star[\chi],\infty)$ where $r^\star[\chi]:=\inf\mathcal{A}_{[\chi]}$.

Step 4: Uniqueness of the onset. Since the real line is totally ordered, the infimum $r^\star[\chi]$ is unique whenever $\mathcal{A}_{[\chi]}\neq\varnothing$, establishing uniqueness of the onset in (10.55). (Under the regularity inherited from the acceptance window—Lipschitz/TV bounds via Lemma 7.386 and Proposition 7.383—the onset is stable and, when the meters are right-continuous, attained.) □

What this means. The parameter r is the "tightening knob" (IR acceptance, spectral cut, or resolution) fixed by the canonical mask (Definition 7.665). Lemma 10.86 says that as r increases, the *law* compliance and the (masked) *state* signal—computed from the calibrated meters of Definition 10.82—can only improve or plateau (Proposition 9.146). Consequently, the set of r for which a type $[\chi]$ passes both tests is a *single ray* $[r^\star[\chi],\infty)$ or empty. There is no ambiguity from oscillations or interleaving pass/fail bands: the minimal acceptable $r^\star[\chi]$ is a well-defined number that summarizes "how much tightening" is needed for that typed signal to be simultaneously lawful and visible.

Why it is useful. This interval property underpins unique onset reporting and auditability across sectors and types (Definitions 9.116 and 10.2). It means that we can compare TT, τ–scalar, or Higgs–CP channels on *equal footing*: each has a unique onset $r^\star[\chi]$ that is band/BRST-clean (Proposition 10.10 and Remark 10.93) and insensitive to the order of typed

vs. spectral operations (Proposition 10.9). Together with the stability/regularity of the meters (Lemma 7.386 and Proposition 7.383), this provides a principled thresholding mechanism for declaring detections, ranking channels, and triggering downstream actions—without risking spurious accept/reject toggling as analysis settings are tightened.

Conceptual understanding. Think of r as the *Steward of Tightening*: an admissible dial that governs monotone refinements of the analysis, including spectral/temporal *resolution*. The *Gatekeeper*—the canonical mask (Definition 7.665)—and the *Cartographer*—the band functor with its IR window (Ch. 8: Definitions 6.280 and 7.29)—are implemented so that as the Steward raises r, acceptance becomes stricter/finer. The *Judges*—the calibrated meters from Definition 10.82—then return scores that are provably *nondecreasing* in r (Proposition 9.146). Concretely:

- *Resolution tightening (the Lens narrows).* Decrease a smoothing scale $\sigma(r)$ or narrow bins $(\Delta\omega, \Delta k)$ so the resolving power grows with r; the Gatekeeper acts as an *isotone* mask ensuring order is preserved (Definition 7.665 and Remark 9.15).
- *Band/IR tightening (the Gate closes).* Shrink the accepted window $K_{\mathrm{IR}}(r)$ or raise SNR-like thresholds in a manner nondecreasing in r (Ch. 8: Definition 6.280); the Cartographer updates the certified domain accordingly.
- *Mode cutoffs (the Archivist prunes).* Increase harmonic/order cutoffs (e.g. $L_{\max}(r)$, $N(r)$) or raise spectral contours; the *Riesz Archivists* (selectors) then respect the new contours (Definition 10.7).

Because all of these moves are implemented via *isotone masks/selectors*, the acceptance set is an *upper set*:

$$\mathcal{A}_{[\chi]} = \Big\{ r : \text{law \& state pass} \Big\} = \varnothing \quad \text{or} \quad [\, r^\star[\chi], \infty),$$

and the onset $r^\star[\chi]$ is unique whenever $\mathcal{A}_{[\chi]} \neq \varnothing$ (Lemma 10.86). Moreover, since the *Typist* (typed projectors) and the *Provenance Clerk* (triad) commute with the band machinery (Definitions 9.116 and 10.2 and Propositions 10.9 and 10.10), the meaning of r as "resolution" (or any other admissible tightening) is compatible with TT, τ–scalar, and Higgs–CP channels without order ambiguities.

Proposition 10.87 (Stability radii under admissible retuning)**.** *Let $(\lambda_{\mathrm{UV}}, \ell, r)$ be the band/IR knobs of $\mathcal{I}^{[\mathscr{S}]}$ (Def. 10.1). Assume the typed projectors used for $[\chi]$ satisfy the band retuning stability of Ch. 10 (Definition 9.116, item (P7)), so that the calibrated meters $\Lambda^{[\mathscr{S};\chi]}_{\mathrm{law/state}}$ depend continuously on $(\lambda_{\mathrm{UV}}, \ell, r)$ (cf. Def. 10.82, Prop. 10.10). Then for every $\varepsilon \in (0,1)$ there exists a stability radius $\rho_{[\chi]}(\varepsilon) > 0$ such that any admissible retuning*

$$\big\| (\lambda'_{\mathrm{UV}}, \ell', r') - (\lambda_{\mathrm{UV}}, \ell, r) \big\| \;<\; \rho_{[\chi]}(\varepsilon)$$

implies

$$\Big| \, r^\star[\chi]\big|_{(\lambda'_{\mathrm{UV}}, \ell', \cdot)} \;-\; r^\star[\chi]\big|_{(\lambda_{\mathrm{UV}}, \ell, \cdot)} \Big| \;\leq\; \varepsilon,$$

i.e. the onset is locally stable (upper semicontinuous and, under the canonical mask, locally Lipschitz) with respect to admissible band/IR retunings.

Proof. By Ch. 10 retuning stability (Definition 9.116, item (P7)), the typed projectors and hence the calibrated entries $f_j(R_j), g_m(S_m)$ vary continuously under admissible retunings; by Definition 10.82, the visibilities (10.54) vary continuously as compositions of continuous maps on the positive orthant. With the canonical spectral mask, both onset predicates are nondecreasing (Prop. 9.146), so the hitting time of the thresholds $\theta_{\mathrm{law}}, \theta_{\mathrm{state}}$ depends upper semicontinuously on parameters. Standard perturbation of hitting times for continuous, one–sided monotone functions gives the stated $\rho_{[\chi]}(\varepsilon)$; local Lipschitzness follows when the meters cross the thresholds with nonzero one–sided slope (Ch. 10 onset rules: eq. (9.81), and TLO readouts: Definition 9.147). □

Reporting. In the Parasymmetric TLO, the onset component uses $r^\star[\chi]$ from (10.55) together with the sector/location tags (cf. Definition 9.147); the uniqueness and stability granted by Lemma 10.86 and Prop. 10.87 ensure reproducible certificates on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$.

Proposition 10.88 (Onset uniqueness and stability radius)**.** *Let $\mathscr{S} \in \{\Phi\Phi, \Theta\Theta, \Phi\Theta\}$ and $[\chi]$ be any admitted type (Ch. 10; Definition 9.116, instantiated in Definitions 10.3 to 10.5). Assume the calibrated visibilities $\Lambda^{[\mathscr{S};\chi]}_{\mathrm{law/state}}(r)$ (Def. Definition 10.82, (10.54)) are continuous in r and satisfy the monotonicities of Proposition 9.146 under IR tightening $r\uparrow$. Then:*

(i) Uniqueness

The certified onset $r^\star[\chi]$ of Definition 10.85 is unique (if it exists).

(ii) One-sided stability radius (IR tightening)

Fix any $r_0 \geq r^\star[\chi]$ and suppose there are slope bounds $m_{\mathrm{law}}, m_{\mathrm{state}} > 0$ such that, for all $r \geq r_0$,

$$\begin{aligned}
\frac{d}{dr}\Lambda^{[\mathscr{S};\chi]}_{\mathrm{law}}(r) &\geq m_{\mathrm{law}},\\
\frac{d}{dr}\Lambda^{[\mathscr{S};\chi]}_{\mathrm{state}}(r) &\geq m_{\mathrm{state}} \quad \text{(if the state meter is spectral; cf. eq. (10.52))},\\
\frac{d}{dr}\Lambda^{[\mathscr{S};\chi]}_{\mathrm{state}}(r) &\leq -\,m_{\mathrm{state}} \quad \text{(if the state meter is vacuum/coherence)}.
\end{aligned}$$

Let the gaps at r_0 be

$$\Delta_{\mathrm{law}}(r_0) := \Lambda^{[\mathscr{S};\chi]}_{\mathrm{law}}(r_0) - \theta_{\mathrm{law}}, \qquad \Delta_{\mathrm{state}}(r_0) := \Lambda^{[\mathscr{S};\chi]}_{\mathrm{state}}(r_0) - \theta_{\mathrm{state}},$$

(which are ≥ 0 by $r_0 \geq r^\star[\chi]$). Then for any $\delta r \in [0, \rho[\chi])$ with

$$\rho[\chi] \;:=\; \min\left\{\frac{\Delta_{\mathrm{law}}(r_0)}{m_{\mathrm{law}}},\; \frac{\Delta_{\mathrm{state}}(r_0)}{m_{\mathrm{state}}}\right\}, \tag{10.56}$$

the admission condition of Definition 10.85 remains satisfied at $r_0 + \delta r$. In particular, choosing $r_0 > r^\star[\chi]$ yields a strictly positive stability radius.

Proof. (1) By Proposition 9.146, $\Lambda^{[\mathscr{S};\chi]}_{\mathrm{law}}(r)$ is nondecreasing and, with the canonical spectral mask from Ch. 8 (Definition 7.665), the relevant state meter used in the onset test is also nondecreasing; therefore the superlevel sets $\{r : \Lambda_{\mathrm{law}} \geq \theta_{\mathrm{law}}\}$ and $\{r : \Lambda_{\mathrm{state}} \geq \theta_{\mathrm{state}}\}$ are

intervals of the form $[r_{\mathrm{law}}, \infty)$ and $[r_{\mathrm{state}}, \infty)$, and their intersection is $[r^\star, \infty)$. Its infimum $r^\star$ is unique (cf. Definition 10.85).

(2) Fix $r_0 \geq r^\star[\chi]$. By the mean value theorem and the assumed slope bounds, for any $\delta r \geq 0$,

$$\Lambda_{\mathrm{law}}(r_0 + \delta r) \;\geq\; \Lambda_{\mathrm{law}}(r_0) \;+\; m_{\mathrm{law}}\delta r,$$

and, depending on the channel used in the state meter,

$$\begin{aligned} \Lambda_{\mathrm{state}}(r_0 + \delta r) &\geq \Lambda_{\mathrm{state}}(r_0) + m_{\mathrm{state}}\delta r \quad \text{(spectral)}, \\ \Lambda_{\mathrm{state}}(r_0 + \delta r) &\geq \Lambda_{\mathrm{state}}(r_0) - m_{\mathrm{state}}\delta r \quad \text{(vac/corr)}. \end{aligned}$$

Thus both visibilities remain above their thresholds provided $m_{\mathrm{law}}\delta r \leq \Delta_{\mathrm{law}}(r_0)$ and $m_{\mathrm{state}}\delta r \leq \Delta_{\mathrm{state}}(r_0)$, which is exactly $\delta r < \rho[\chi]$ in (10.56). If the canonical spectral mask (Definition 7.665) is used for onset (so only the spectral entry participates), the second inequality is of the increasing type. □

Remark 10.89. (i) The result is *one-sided* with respect to IR tightening ($\delta r \geq 0$), consistent with the monotonicities of Proposition 9.146. (ii) At $r_0 = r^\star[\chi]$ the gaps may vanish, yielding $\rho[\chi] = 0$; taking any $r_0 > r^\star[\chi]$ produces a positive radius. (iii) Continuity and slope lower bounds are ensured in practice by the typed retuning stability of Ch. 10 (Definition 9.116, item (P7)) together with band/BRST hygiene on the realized weave configuration (Definition 10.1 and Proposition 10.10). (iv) The slopes $m_{\mathrm{law}}, m_{\mathrm{state}}$ correspond to the one-sided derivatives in the onset rules of Ch. 10 (eq. (9.81)), which underwrite the TLO "Onset" component in Definition 9.147.

10.9.5 Explicit Witness Contents by Type and Sector

Below, $\mathfrak{O}$ denotes the *typed, physical* observable $\mathfrak{O}^{[\mathscr{S};\chi]} := \mathsf{P}_{\mathrm{phys}}\Pi^{[\chi]}\mathfrak{O}\,\Pi^{[\chi]}\mathsf{P}_{\mathrm{phys}}$ (cf. Definition 10.80), evaluated on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ (Def. Definition 10.1) with sector block $\mathscr{S} \in \{\Phi\Phi, \Theta\Theta, \Phi\Theta\}$. All divergences and traces use the $\bar{\Sigma}$–compatible Levi–Civita derivative $\nabla^{(\mathrm{rel})}$ and contraction with $\bar{\Sigma}$ (Chs. 6–7; Definitions 5.26 and 6.7 and eq. (6.22)). Triad projectors $\Pi^{(\mathrm{spec})}, \Pi^{(\mathrm{vac})}, \Pi^{(\mathrm{corr})}$ are from Definition 10.2. Typed families (TT, τ–kairos scalars/vectors, Higgs CP) are as instantiated in Definitions 10.3 to 10.5. Spectral weights use the Källén–Lehmann density and a fixed admissible window W_{spec} (Ch. 8: eq. (7.12); cf. eq. (10.52)).

(A) Graviton (TT, $\Phi\Phi$–sector). Here $\chi = \mathrm{TT}$ and $\mathscr{S} = \Phi\Phi$; TT purity/TT-leak are as in Definitions 10.3 and 10.32.

Law residuals (cf. eq. (10.51)):

$$R_1 = \left\|\nabla^{(\mathrm{rel})\mu}\mathfrak{O}_{\mu\nu}\right\|_{\mathcal{I}}, \qquad R_2 = \left\|\mathrm{tr}_{\bar{\Sigma}}\mathfrak{O}\right\|_{\mathcal{I}},$$
$$R_3 = \left\|\partial_\xi\, \mathcal{G}_\xi^{[\Phi\Phi;\mathrm{TT}]}\right\|_{\mathcal{I}}, \qquad R_4 = \left\|\Pi_{\bar{\Sigma}}^{\perp[\mathrm{TT}]}\,\mathfrak{O}\right\|_{\mathcal{I}}.$$

State features (cf. eq. (10.52)):

$$S_1 = \int W_{\mathrm{spec}}\ \rho_{(2)}^{[\Phi\Phi;\mathrm{TT}]}\, d\mu, \qquad S_2 = \left\|\Pi^{(\mathrm{vac})}\mathfrak{O}\right\|_{\mathcal{I}},$$
$$S_3 = \left\|\Pi^{(\mathrm{corr})}\mathfrak{O}\right\|_{\mathcal{I}}, \qquad S_4 = \varepsilon \quad \text{(Definition 10.32)}.$$

(Invariance/flatness for TT meters: Proposition 10.30.)

(B) τ–kairos scalars. Types: $S^{(b)}$, $S^{(L)}$, $\tilde{S}^{(p)}$. Typed insertion allowed in any sector $\mathscr{S} \in \{\Phi\Phi, \Theta\Theta, \Phi\Theta\}$. TT–nullity and parity conventions follow Definition 10.4; TT annihilation uses Definition 10.3 and τ–TT purity Lemma 10.17.

Law residuals (cf. eq. (10.51)):

$$R_1 = \left\|\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\mathfrak{O}\right\|_{\mathcal{I}},$$
$$R_2 = \left\|\partial_\xi\, \mathcal{G}_\xi^{[\mathscr{S};\tau\text{-sc}]}\right\|_{\mathcal{I}},$$
$$R_3 = \left\|\nabla^{(\mathrm{rel})\mu}\mathfrak{O}_{\mu\nu}\right\|_{\mathcal{I}}, \quad R_4 = \left\|\mathrm{tr}_{\bar{\Sigma}}\mathfrak{O}\right\|_{\mathcal{I}},$$
$$\text{and for } \tilde{S}^{(p)}: \ R_5 = \left\|\varepsilon^{\alpha\beta\gamma(\mu}\nabla_\alpha^{(\mathrm{rel})}\mathfrak{O}_{\beta\gamma}{}^{\nu)}\right\|_{\mathcal{I}}$$
(parity–odd check; cf. Definition 10.4).

State features (cf. eq. (10.52)):

$$S_1 = \int W_{\mathrm{spec}}\ \rho_{(0)}^{[\mathscr{S};\tau\text{-sc}]}\, d\mu,$$
$$S_2 = \left\|\Pi^{(\mathrm{vac})}\mathfrak{O}\right\|_{\mathcal{I}}, \quad S_3 = \left\|\Pi^{(\mathrm{corr})}\mathfrak{O}\right\|_{\mathcal{I}}.$$

(Nielsen/BRST flatness for typed τ scalars inherits the general invariance on the realized weave configuration: Proposition 10.10 and Remark 10.93.)

(C) Chrono–Kymon $A^{(\pm)}$ (τ–vector, typed insertion in any sector). Here $\chi \in \{A^{(+)}, A^{(-)}\}$ and $\mathscr{S}$ is arbitrary in $\{\Phi\Phi, \Theta\Theta, \Phi\Theta\}$. The CK tensor image $A_{\mu\nu}^{(\pm)}[\mathfrak{O}]$ is as in (10.45) (from the CK field definition (10.44)); helicity projectors $A^{(\pm)}$ and parity asymmetry Δ_P are in

Definitions 10.4 and 10.58. TT-nullity follows from Lemma 10.59.

Law residuals (cf. eq. (10.51)):

$$R_1 = \left\| \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} A_{\mu\nu}^{(\pm)}[\mathfrak{O}] \right\|_{\mathcal{I}}, \qquad R_2 = \left\| \left(\mathbf{1} - A^{(\pm)}\right) \mathfrak{O} \right\|_{\mathcal{I}},$$
$$R_3 = \left\| \partial_\xi \, \mathcal{G}_\xi^{[\mathscr{S};\tau\text{-vec}]} \right\|_{\mathcal{I}}.$$

State features (cf. eq. (10.52)):

$$S_1 = \int W_{\mathrm{spec}} \, \rho_{(1)}^{[\mathscr{S};\tau\text{-vec}]} \, d\mu, \qquad S_2 = \left\| \Pi^{(\mathrm{vac})} \mathfrak{O} \right\|_{\mathcal{I}},$$
$$S_3 = \left\| \Pi^{(\mathrm{corr})} \mathfrak{O} \right\|_{\mathcal{I}}, \qquad S_4 = \Delta_P \quad \text{(Definition 10.58)}.$$

(Invariance/flatness for CK meters: Proposition 10.61.)

(D) Higgs (radial tantrakyma) rad± (ΘΘ–sector). Here $\chi \in \{\mathrm{rad}+, \mathrm{rad}-\}$ and $\mathscr{S} = \Theta\Theta$. NG/pNG orthogonality and CP projectors/asymmetry are from Proposition 10.47 and Definitions 10.5 and 10.51.

Law residuals (cf. eq. (10.51)):

$$R_1 = \left\| \Pi_{\Theta}^{[\mathrm{NG/pNG}]} \mathfrak{O} \right\|_{\mathcal{I}}, \qquad R_2 = \left\| \Pi^{(\mathrm{CP}\mp)} \mathfrak{O} \right\|_{\mathcal{I}},$$
$$R_3 = \left\| \partial_\xi \, \mathcal{G}_\xi^{[\Theta\Theta;\mathrm{rad}\pm]} \right\|_{\mathcal{I}}.$$

State features (cf. eq. (10.52)):

$$S_1 = \int W_{\mathrm{spec}} \, \rho_{(0)}^{[\Theta\Theta;\mathrm{rad}\pm]} \, d\mu, \qquad S_2 = \left\| \Pi^{(\mathrm{vac})} \mathfrak{O} \right\|_{\mathcal{I}},$$
$$S_3 = \left\| \Pi^{(\mathrm{corr})} \mathfrak{O} \right\|_{\mathcal{I}}, \qquad S_4 = \Delta_{\mathrm{CP}}^{(a)} \quad \text{(Definition 10.51)}.$$

Sector note. For the mixed interaction block $\mathscr{S} = \Phi\Theta$, all entries above are understood after *typed insertion* into the off–diagonal correlator blocks of the unified inter–modal correlator (cf. (10.61)); τ–kairos enters only as a *typed projector family* (not as a sector), per Definition 10.4. Gauge–parameter independence and band stability for all meters follow from the realized weave configuration-level invariance (Proposition 10.10 and Remark 10.93).

10.9.6 Parasymmetric TLO: Existence, Uniqueness, and Stability

Proposition 10.90 (Existence and uniqueness of TLO per type)**.** *Let $\mathscr{S} \in \{\Phi\Phi, \Theta\Theta, \Phi\Theta\}$ and let $[\chi]$ be any admitted type from the typed layer (Ch. 10; Definition 9.116, instantiated in Definitions 10.3 to 10.5). On the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ (Def. Definition 10.1), the* Parasymmetric TLO *triple*

$$\mathrm{TLO}^{[\chi]} = \big(\mathrm{Type},\ \mathrm{Location},\ \mathrm{Onset}\big)$$

as defined in Definition 9.147 with onset $r^\star[\chi]$ from Definition 10.85 exists and is unique. *Moreover, under the canonical spectral mask (Ch. 8; Definition 7.665), if the spectral feature S_1 in (10.52) is strictly positive on some admissible window for some r_0, and if $\Lambda_{\mathrm{law}}^{[\mathscr{S};\chi]}(r_0) \geq \theta_{\mathrm{law}}$, then $r^\star[\chi] < \infty$.*

Proof. **Existence/uniqueness of** Type**.** For fixed $[\chi]$, the typed projectors are idempotent and mutually orthogonal inside their block (Ch. 10: Definition 9.116); in the weave these are instantiated for TT, τ–kairos, and Higgs families (Definitions 10.3 to 10.5). Orthogonality/purity lemmas (e.g. TT purity Lemma 10.28, τ algebra Lemma 10.8, Higgs CP orthogonality Lemma 10.45) ensure a single typed branch is selected for the meter. Hence the quadrant decision (Type $=$ Quad$^{[\chi]}$) obtained from the calibrated visibilities (Definition 10.82, (10.54)) is well defined and unique.

Existence/uniqueness of Location**.** Location (arena/sector/position/sheet) is inherited from the realized weave configuration $\mathcal{I}^{[\mathcal{S}]}$ (Def. Definition 10.1) and the sector block $\mathcal{S}$; uniqueness is therefore immediate.

Existence/uniqueness of Onset**.** With the canonical spectral mask (Ch. 8: Definition 7.665), Prop. 9.146 yields that the law visibility $\Lambda_{\mathrm{law}}^{[\mathcal{S};\chi]}(r)$ and the masked state visibility are *nondecreasing* in r. Thus, by Lemma 10.86, the admission set is an interval $[r^\star[\chi],\infty)$ (or empty), and the onset $r^\star[\chi]$ of Definition 10.85 is unique when it exists.

Finiteness (sufficient condition). If there exists r_0 such that the spectral feature $S_1(r_0) > 0$ on some admissible window (Ch. 8: eq. (7.12)) and $\Lambda_{\mathrm{law}}^{[\mathcal{S};\chi]}(r_0) \geq \theta_{\mathrm{law}}$, then the masked state visibility at r_0 exceeds some positive value; by nondecreasing monotonicity (Prop. 9.146) it will cross any fixed $\theta_{\mathrm{state}} \in (0,1)$ at some finite $r \geq r_0$, whence $r^\star[\chi] < \infty$.

Stability. The onset is stable under admissible retunings by Prop. 10.87, which relies on the typed retuning stability of Ch. 10 (Definition 9.116, item (P7)) and realized weave configuration-level invariance/continuity (Proposition 10.10). □

Corollary 10.91 (Audit triggers)**.** *If any of the following holds on the realized weave configuration $\mathcal{I}^{[\mathcal{S}]}$ (Def. Definition 10.1),*

$$\begin{aligned} \varepsilon &> \varepsilon_{\max} && \textit{(graviton TT leak; Definition 10.32)},\\ |\Delta_P| &> \delta_P && \textit{(vector parity bias; Definition 10.58)},\\ |\Delta_{\mathrm{CP}}^{(a)}| &> \delta_{\mathrm{CP}} && \textit{(Higgs CP bias; Definition 10.51)}, \end{aligned}$$

where the audit thresholds $\varepsilon_{\max}, \delta_P, \delta_{\mathrm{CP}}$ are fixed a priori (cf. Ch. 8 fixed mask thresholds Definition 7.665), then a SynTh audit *must be initiated (see §10.10).*

Proof. Each listed diagnostic is included among the *state features* $\mathbf{S}^{[\mathcal{S};\chi]}$ (cf. eq. (10.52) and the type-specific augmentations in Remark 10.81; explicit contents by type appear in §10.9.5). By the chapter's audit design, these bounds define the *automatic trigger* predicate (Def. Definition 10.95); when any bound is exceeded, Cor. Corollary 10.110 mandates entering the *Audit path* (Def. Definition 10.105). Operationally this is the SynTh No-Explosion/Audit branch from the simultaneous synthesis rules (Ch. 9: Definitions 8.98 and 8.99), executed on the realized weave configuration $\mathcal{I}^{[\mathcal{S}]}$. Hence an audit is required irrespective of the onset status. □

Remark 10.92 (Why exponential and rational calibrations)**.** The choices in Definition 10.82,

$$f_j(x) = e^{-\alpha_j x} \in (0,1], \qquad g_m(x) = \frac{\beta_m x}{1+\beta_m x} \in [0,1),$$

ensure: (i) *boundedness* of the calibrated entries (so the cosine meters (10.54) are always finite); (ii) *strict monotonicity* in the required directions—f_j is strictly decreasing (larger residual ⇒ smaller law score) while g_m is strictly increasing (larger state feature ⇒ larger state score)—which underpins the one-sided monotonicities in Proposition 9.146; and (iii) C^1 *differentiability* with nonzero slopes on $(0,\infty)$, which supplies the slope bounds used in the onset stability analysis Proposition 10.88. Any alternative calibrations $f_j : \mathbb{R}_+ \to (0,1]$ and $g_m : \mathbb{R}_+ \to [0,1)$ that are bounded, strictly monotone (in the same respective directions), and C^1 would serve identically in (10.53)–(10.54).

Remark 10.93 (Independence from gauge fixing)**.** In the Law ledger (10.51), the BRST–closure residual $\|[\hat{Q}_{\mathrm{BRST}}, \mathcal{O}]\|_{\mathcal{I}}$ and the explicit Nielsen residual $\|\partial_\xi \mathcal{G}_\xi^{[\mathscr{S};\chi]}\|_{\mathcal{I}}$ (cf. Chs. 8–10 Nielsen setup) are evaluated on the typed, physical observable $\mathcal{O}^{[\mathscr{S};\chi]}$ of Definition 10.80. Vanishing of these residuals on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ (Def. Definition 10.1) implies that the calibrated visibilities (10.54) are independent of the gauge parameter on $\mathcal{H}_{\mathrm{phys}}$. This is consistent with the realized weave configuration-level invariance and band hygiene (Proposition 10.10) and with the type-level examples such as TT graviton and Chrono–Kymon, where gauge-parameter independence (Nielsen flatness) is established in Propositions 10.30 and 10.61. Therefore the meters used in the Parasymmetric TLO (Definition 9.147) are gauge-fixing independent once the ledger residuals of Definition 10.80 are satisfied.

10.10 SynTh Resolution and Audit for the Weave

We complete Step (5) of the operational pipeline by formalizing SynTh (Simultaneous Synthesis) on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ (Def. Definition 10.1; Ch. 9: Definitions 8.98 and 8.99) and by proving a *No-Explosion/Audit Theorem* specialized to the geometry–matter weave. Throughout, visibilities, witnesses, and TLO components are those constructed in Section 10.9 (see Definitions 10.80, 10.82 and 10.85, Section 10.9.5, Lemma 10.86, and Proposition 10.87).

Set-up. Fix a sector $\mathscr{S} \in \{\Phi\Phi, \Theta\Theta, \Phi\Theta\}$ and an admitted type $[\chi]$ (e.g. TT, $S^{(b)}$, $S^{(L)}$, $\tilde{S}^{(p)}$, $A^{(\pm)}$, rad$\pm$). The law/state visibilities

$$\Lambda_{\mathrm{law}}^{[\mathscr{S};\chi]}, \qquad \Lambda_{\mathrm{state}}^{[\mathscr{S};\chi]}$$

are the calibrated cosine meters of Definition 10.82 (see (10.54)), built from the ledgers in Definition 10.80 and the type-specific components in Remark 10.81; explicit witness contents by type/sector are listed in §10.9.5. Diagnostics ε, Δ_P, $\Delta_{\mathrm{CP}}^{(a)}$ are as in Definitions 10.32, 10.51 and 10.58. All objects live on the physical carrier $\mathcal{H}_{\mathrm{phys}}$ and inherit band/BRST hygiene (Proposition 10.10 and Remark 10.93).

Definition 10.94 (Co-tenability vs. conflict; admission predicate)**.** Fix thresholds $\theta_{\mathrm{law}}, \theta_{\mathrm{state}} \in (0,1)$ (Ch. 8 fixed masks: Definition 7.665). Given $(\mathscr{S}, [\chi])$, define the *admission predicate* (cf. Definition 10.85)

$$\mathrm{Adm}_{[\chi]}(r) \;:=\; \Big[\, \Lambda_{\mathrm{law}}^{[\mathscr{S};\chi]}(r) \geq \theta_{\mathrm{law}} \;\wedge\; \Lambda_{\mathrm{state}}^{[\mathscr{S};\chi]}(r) \geq \theta_{\mathrm{state}} \,\Big].$$

We say *Law and State are co-tenable* (write $\mathrm{Syn}^{[\mathscr{S};\chi]}$) at r if $\mathrm{Adm}_{[\chi]}(r)$ holds; otherwise they are in *conflict*. The *certified onset* $r^\star[\chi]$ is the first r where $\mathrm{Adm}_{[\chi]}(r)$ holds (Definition 10.85).

Definition 10.95 (Automatic triggers and Audit path)**.** (A) *Automatic triggers.* A SynTh audit must be initiated if any of the following hold (cf. Corollary 10.91):

$$\varepsilon > \varepsilon_{\max}, \qquad |\Delta_P| > \delta_P, \qquad |\Delta_{\mathrm{CP}}^{(a)}| > \delta_{\mathrm{CP}},$$

with a priori thresholds $(\varepsilon_{\max}, \delta_P, \delta_{\mathrm{CP}})$ (Ch. 8 masks: Definition 7.665).
(B) *Audit path.* Starting from a pair $(\mathscr{S}, [\chi])$ and scale r, the Audit path $\mathsf{Audit}(\mathcal{I}^{[\mathscr{S}]}, [\chi]; r)$ is the (finite) sequence of steps:

(i) Evaluate ledgers and visibilities (Definitions 10.80 and 10.82); record $(\Lambda_{\mathrm{law}}, \Lambda_{\mathrm{state}})$ and diagnostics $(\varepsilon, \Delta_P, \Delta_{\mathrm{CP}})$.

(ii) If any automatic trigger fires (above), *enter* Audit; else, if $\neg\mathrm{Adm}_{[\chi]}(r)$, enter Audit due to *conflict*.

(iii) Apply an admissible, monotone IR retuning $r \mapsto r'$ (tightening) and/or permitted band retuning $(\lambda_{\mathrm{UV}}, \ell)$ within the typed stability radius (Ch. 10 stability: Definition 9.116, item (P7); onset stability: Proposition 10.87); re-evaluate step (i).

(iv) Stop when either $\mathrm{Adm}_{[\chi]}(r')$ holds (co-tenability achieved) or a certified failure certificate is produced (see Theorem 10.98).

Meaning. Definition 10.95 installs two complementary safety rails. Part (A) lists *diagnostic thresholds* that *force* a SynTh audit whenever core invariants are violated: the TT–leak ε (tensor purity; cf. Definition 10.32), the vector–parity asymmetry Δ_P (Definition 10.58), and the scalar CP asymmetry $\Delta_{\mathrm{CP}}^{(a)}$ in a chosen provenance a (Definition 10.51). Their cutoffs $(\varepsilon_{\max}, \delta_P, \delta_{\mathrm{CP}})$ are fixed by the canonical, isotone mask (Definition 7.665). Part (B) then defines the *Audit path*: a finite, recipe-level loop that starts from a sector/type $(\mathscr{S}, [\chi])$ and scale r, evaluates the calibrated visibilities $(\Lambda_{\mathrm{law}}, \Lambda_{\mathrm{state}})$ (Definitions 10.80 and 10.82), and either (i) escalates immediately if any automatic trigger fires, or (ii) enters audit on *conflict* when the acceptance predicate $\mathrm{Adm}_{[\chi]}(r)$ (law & state passing under the mask) fails (Definition 7.665 and Lemma 10.86). The intent is clear: even when visibilities look acceptable, any breach of TT purity, parity balance, or CP balance must be investigated; conversely, if acceptance fails despite tame diagnostics, the configuration is inconsistent and must be retuned.

How the audit proceeds (and why it terminates). Step (iii) of Definition 10.95 retunes only along admissible, monotone directions—tightening r and/or permitted band parameters $(\lambda_{\mathrm{UV}}, \ell)$—inside the typed stability radius (Definition 9.116, item (P7)), with meters that are *nondecreasing* under such moves (Proposition 9.146) and with typed/triad commutation intact (Proposition 10.9). Because the mask is isotone, the acceptance set for each type is an *upper set* $[r^\star[\chi], \infty)$ when nonempty, so the audit either reaches co-tenability at the unique onset $r^\star[\chi]$ (Lemma 10.86) or produces a certified failure certificate (Theorem 10.98). Throughout, band/BRST hygiene and realized weave configuration invariance are preserved (Proposition 10.10 and Remark 10.93), guaranteeing that audit outcomes are physically

meaningful and independent of the order in which typing, spectral triage, and calibration are applied (Definitions 10.2 and 10.82).

Lemma 10.96 (Monotone descent invariants)**.** *Let* $\mathscr{S} \in \{\Phi\Phi, \Theta\Theta, \Phi\Theta\}$ *and* $[\chi]$ *be any admitted type (Ch. 10; Definition 9.116, instantiated in Definitions 10.3 to 10.5). Along* IR tightening $r \uparrow$ *(Ch. 8 IR window: Definition 6.280, band functor: Definition 7.29), the following hold for the calibrated meters of Definition 10.82:*

1. $\Lambda_{\mathrm{law}}^{[\mathscr{S};\chi]}(r)$ *is* nondecreasing. *With the* canonical spectral mask *(Ch. 8: Definition 7.665) applied to the state meter, the masked state visibility* $\Lambda_{\mathrm{state}}^{[\mathscr{S};\chi]}(r)$ *is also* nondecreasing. *(Proposition 9.146)*
2. *The TT-leak diagnostic* ε *is* nonincreasing. *Moreover, the (unmasked) vacuum and coherence state features are* nonincreasing. *(Corollary 10.104 and Proposition 9.146)*

Consequently, with thresholds $\theta_{\mathrm{law}}, \theta_{\mathrm{state}} \in (0,1)$ *from Definition 10.85 and a priori trigger bounds* $(\varepsilon_{\max}, \delta_P, \delta_{\mathrm{CP}})$ *(Ch. 8 masks: Definition 7.665), the* core descent potential

$$\mathfrak{D}_{\mathrm{core}}(r) := \max\Big\{\ \theta_{\mathrm{law}} - \Lambda_{\mathrm{law}}^{[\mathscr{S};\chi]}(r),\quad \theta_{\mathrm{state}} - \Lambda_{\mathrm{state}}^{(\mathrm{spec})[\mathscr{S};\chi]}(r),\ 0\Big\} \ +\ \mathbf{1}\Big[\varepsilon(r) > \varepsilon_{\max}\Big] \tag{10.57}$$

is nonincreasing *along IR tightening on the Audit path (Def. Definition 10.101). If one augments* (10.57) *by the parity/CP indicators to form*

$$\mathfrak{D}_{\mathrm{audit}}(r) := \mathfrak{D}_{\mathrm{core}}(r) + \mathbf{1}\Big[|\Delta_P(r)| > \delta_P\Big] + \mathbf{1}\Big[|\Delta_{\mathrm{CP}}^{(a)}(r)| > \delta_{\mathrm{CP}}\Big], \tag{10.58}$$

then $\mathfrak{D}_{\mathrm{audit}}$ *is nonincreasing on any tightening segment where no new parity/CP trigger fires; if a parity/CP trigger does fire, the SynTh policy mandates immediate Audit entry (Def. Definition 10.95, Cor. Corollary 10.91).*

Proof. (1) By Proposition 9.146(1), tightening the IR window $K_{\mathrm{IR}}^{(r)}$ makes each law residual R_j weakly decrease; with the decreasing calibration $f_j(x) = e^{-\alpha_j x}$ (Def. Definition 10.82) each calibrated entry $f_j(R_j)$ weakly *increases*, hence Λ_{law} (cosine with $\mathbf{1}$, (10.54)) is nondecreasing. For the state meter, Proposition 9.146(2) states that the *spectral* feature (with fixed W_{spec}, Ch. 8: Definition 7.665) weakly *increases*, whereas vacuum/coherence weakly *decrease*. With the canonical spectral mask used in onset tests (cf. Definition 10.85), the resulting masked Λ_{state} is nondecreasing.

(2) TT-leak ε is defined by a TT-sandwiched non-spectral norm (Def. Definition 10.32); Corollary 10.104 gives its monotone *decrease* under IR tightening. For vacuum/coherence, Proposition 9.146(2) establishes the weak *decrease* of their raw features; after the increasing calibration $g_m(x) = \beta_m x/(1+\beta_m x)$ (Def. Definition 10.82), their contributions to the unmasked state meter are also nonincreasing.

For the potentials: By (1), $\theta_{\mathrm{law}} - \Lambda_{\mathrm{law}}(r)$ and $\theta_{\mathrm{state}} - \Lambda_{\mathrm{state}}^{(\mathrm{spec})}(r)$ are each nonincreasing in r; their $\max(\cdot, 0)$ is therefore nonincreasing. By (2), the indicator $\mathbf{1}[\varepsilon(r) > \varepsilon_{\max}]$ is nonincreasing (once it turns 0 it cannot return to 1). Hence $\mathfrak{D}_{\mathrm{core}}(r)$ in (10.57) is nonincreasing along IR tightening.

Parity/CP indicators have no proven monotone trend under tightening in general. Thus $\mathfrak{D}_{\mathrm{audit}}$ in (10.58) is guaranteed nonincreasing on any tightening segment where no new

parity/CP trigger activates. If a new parity/CP trigger does activate (indicator flips 0→1), the SynTh rules require immediate entry into the Audit branch (Def. Definition 10.95, Cor. Corollary 10.91), which is treated as a terminal step rather than a continued descent; this is the policy used in the No-Explosion/Audit Theorem (Theorem 10.98). All statements are on the physical carrier $\mathcal{H}_{\text{phys}}$ and inherit realized weave configuration-level invariance/continuity (Proposition 10.10). □

Lemma 10.97 (Finite termination and sound triggers)**.** *Let $\mathscr{S} \in \{\Phi\Phi, \Theta\Theta, \Phi\Theta\}$ and $[\chi]$ be an admitted type (Ch. 10; Definition 9.116, instantiated in Definitions 10.3 to 10.5). Consider an Audit path (Def. Definition 10.101) that applies IR tightening $r \uparrow$ and permitted retunings on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ (Def. Definition 10.1). Then:*

(i) Soundness of automatic triggers

If an automatic trigger fires (Def. Definition 10.95; Cor. Corollary 10.91), Audit is entered immediately. Moreover, tightening cannot create *a new trigger that was previously absent for the TT-leak ε or for the vacuum/coherence channels, because these quantities are nonincreasing under IR tightening (Lemma 10.96, using Proposition 9.146 and Corollary 10.104). Hence trigger-based entry is sound (Cor. 10.110) and reduces the SynTh potential (cf. (10.59)).*

(ii) Termination via onset or certified failure

Assume no trigger is active. Use the canonical spectral mask (Ch. 8: Definition 7.665) for the state meter. If the spectral feature is positive on some admissible window (KL positivity; eq. (7.12)), then the masked state visibility $\Lambda^{[\mathscr{S};\chi]}_{\text{state}}(r)$ is nondecreasing in r (Prop. 9.146); likewise $\Lambda^{[\mathscr{S};\chi]}_{\text{law}}(r)$ is nondecreasing (Prop. 9.146). Let

$$\Lambda^{\infty}_{\text{state}} := \lim_{r\to\infty} \Lambda^{[\mathscr{S};\chi]}_{\text{state}}(r) \in (0,1], \qquad \Lambda^{\infty}_{\text{law}} := \lim_{r\to\infty} \Lambda^{[\mathscr{S};\chi]}_{\text{law}}(r) \in (0,1].$$

Then for any thresholds $\theta_{\text{law}}, \theta_{\text{state}} \in (0,1)$ (Def. 10.85):

(a) Admission via onset

If $\theta_{\text{law}} < \Lambda^{\infty}_{\text{law}}$ and $\theta_{\text{state}} < \Lambda^{\infty}_{\text{state}}$, the admission predicate $\text{Adm}_{[\chi]}(r)$ (Def. 10.99) holds for all $r \geq r^{\star}[\chi]$ with a unique onset $r^{\star}[\chi]$ (Lemma 10.86).

(b) Certified failure

If either inequality fails (e.g. $\theta_{\text{state}} \geq \Lambda^{\infty}_{\text{state}}$ or $\theta_{\text{law}} \geq \Lambda^{\infty}_{\text{law}}$), then admission cannot be achieved by further tightening; a certified failure *is produced, listing the limiting visibilities and active residuals/diagnostics (Theorem 10.98 (2b)).*

Consequently, every Audit path terminates in finitely many steps either in co-tenability with onset (Def. 10.85) or with a certified failure.

Proof. (1) The automatic triggers are $\varepsilon > \varepsilon_{\max}$, $|\Delta_P| > \delta_P$, and $|\Delta^{(a)}_{\text{CP}}| > \delta_{\text{CP}}$ (Def. 10.95; Cor. 10.91). By Lemma 10.96, ε (TT-leak) and the vacuum/coherence features are nonincreasing under IR tightening, so if their indicators are 0 at some r_0 they cannot flip to 1 for $r > r_0$; thus tightening cannot create a new trigger in those channels. (Parity/CP indicators are handled by the trigger policy rather than a monotonic claim—see Lemma 10.96.) Therefore, when a trigger fires we must enter Audit immediately, and doing so weakly decreases the SynTh potential $\mathfrak{D}^{[\mathscr{S};\chi]}$ (cf. (10.59)), establishing soundness (Cor. 10.110).

(2) With no active trigger, apply the canonical spectral mask (Ch. 8: Definition 7.665). By Prop. 9.146, both $\Lambda_{\text{law}}(r)$ and the masked $\Lambda_{\text{state}}(r)$ are nondecreasing under $r \uparrow$. KL positivity (eq. (7.12)) and the admissible window guarantee that the spectral mass is positive, hence the masked state visibility has a positive limit $\Lambda^{\infty}_{\text{state}} > 0$. Because the visibilities are continuous in r under typed-stable retunings (Prop. 10.87; Ch. 10 stability Definition 9.116, item (P7)), standard monotone convergence implies:

- If $\theta_{\text{law}} < \Lambda^{\infty}_{\text{law}}$ and $\theta_{\text{state}} < \Lambda^{\infty}_{\text{state}}$, there exists $r^{\star}[\chi]$ such that both inequalities in Def. 10.85 hold for all $r \geq r^{\star}[\chi]$; uniqueness follows from Lemma 10.86.

- Otherwise, at least one threshold exceeds the corresponding limit, so admission cannot be met by any further tightening; per Theorem 10.98 (2b), we output a certified failure that records the limiting visibilities and residuals/diagnostics.

In either case, the Audit path terminates in finitely many steps because each audit move (Def. 10.101) strictly reduces the discrete potential or reaches an absorbing state (co-tenability or certified failure), and the onset stability radius (Prop. 10.87) prevents oscillatory regress. □

Theorem 10.98 (No-Explosion/Audit Theorem for the weave)**.** *For any $(\mathscr{S}, [\chi])$ on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$, the SynTh resolution obeys:*

(i) No-Explosion (paraconsistent safety)

Under the SynTh rules (Definitions 8.98 and 8.99), contradictions between Law and State do not entail triviality; instead, the process branches into the Audit path (Def. 10.95).

(ii) Finite termination

Along any admissible Audit path with IR tightening and typed-stable band retuning, the potential $\mathfrak{D}$ of Lemma 10.96 is nonincreasing and achieves a terminal state in finitely many steps: either

(a) Co-tenability

$\text{Adm}_{[\chi]}(r^{\sharp})$ *holds, yielding the TLO onset $r^{\star}[\chi] \leq r^{\sharp}$ (cf. Definition 10.85, Lemma 10.86, and Proposition 10.87); or*

(b) Certified failure

A certified failure is produced: a tuple listing the active residuals and diagnostics that cannot be healed by admissible retunings within the typed stability radius (stability: Definition 9.116, item (P7)).

(iii) Soundness of automatic triggers

If any of $\varepsilon > \varepsilon_{\max}$, $|\Delta_P| > \delta_P$, $|\Delta^{(a)}_{\text{CP}}| > \delta_{\text{CP}}$ holds, then entering Audit is necessary and correct (Cor. 10.91), and subsequent IR tightening weakens (never worsens) the corresponding violation for TT-leak and vacuum/coherence (Lemma 10.96).

Proof sketch. (1) Paraconsistent safety is by the SynTh calculus (Ch. 9: Definitions 8.98 and 8.99), which replaces explosion with the Audit branch. (2) Monotone descent of $\mathfrak{D}$ follows from Lemma 10.96; IR tightening and typed-stable retunings ensure continuity (Proposition 10.87) and prevent oscillatory regress. Termination to (a) uses Lemma 10.86 and Proposition 9.146; termination to (b) occurs when residuals/diagnostics remain above

thresholds despite exhausting admissible retunings (stability radius in Proposition 10.87, typed stability in Ch. 10: Definition 9.116, item (P7)). (3) Trigger soundness is Corollary 10.91 together with Lemma 10.96. □

Reporting and certificates. When (a) holds, the Parasymmetric TLO deliverable $\mathrm{TLO}^{[\chi]} = (\mathrm{Type}, \mathrm{Location}, \mathrm{Onset})$ is reported per Definition 9.147 with the certified onset $r^\star[\chi]$ of Definition 10.85. When (b) holds, the *Audit certificate* records: sector $\mathscr{S}$, type $[\chi]$, the active residual components from eq. (10.51), the diagnostic values $(\varepsilon, \Delta_P, \Delta_{\mathrm{CP}})$, and the retuning region explored (bounds given by Proposition 10.87); this certificate is attached to the quadrant decision for traceability (cf. Definitions 9.147, 10.80 and 10.82).

Sector note. For the mixed interaction block $\mathscr{S} = \Phi\Theta$, all typed checks and meters are applied *after typed insertion* into the off-diagonal correlator blocks of the unified inter-modal correlator (cf. (10.61)); τ–kairos enters only as a *typed projector family* (not as a sector), per Definition 10.4.

10.10.1 Co-tenability, Conflicts, and Thresholds

Definition 10.99 (Co-tenability and conflict (thresholded refinement))**.** Fix a sector $\mathscr{S} \in \{\Phi\Phi, \Theta\Theta, \Phi\Theta\}$ (cf. unified inter–modal correlator (10.61)) and an admitted type $[\chi]$ from the typed layer (Ch. 10; Definition 9.116, instantiated in Definitions 10.3 to 10.5). Let the calibrated visibilities be $\Lambda_{\mathrm{law}}^{[\mathscr{S};\chi]}$, $\Lambda_{\mathrm{state}}^{[\mathscr{S};\chi]}$ (Def. Definition 10.82, (10.54)) built from the ledgers in Definition 10.80 with type-specific additions in Remark 10.81. Choose thresholds $\theta_{\mathrm{law}}, \theta_{\mathrm{state}} \in (0,1)$ (Ch. 8 fixed masks: Definition 7.665) and diagnostic bounds $\varepsilon_{\max}, \delta_P, \delta_{\mathrm{CP}} > 0$ (cf. Corollary 10.91). We say the pair $\left(\mathrm{Law}^{[\mathscr{S};\chi]}, \mathrm{State}^{[\mathscr{S};\chi]}\right)$ is:

(i) Co-tenable

Co-tenable if

$$\Lambda_{\mathrm{law}}^{[\mathscr{S};\chi]} \geq \theta_{\mathrm{law}}, \qquad \Lambda_{\mathrm{state}}^{[\mathscr{S};\chi]} \geq \theta_{\mathrm{state}},$$

and, for diagnostics applicable to $[\chi]$ (see Remark 10.81 and Section 10.9.5),

$$\varepsilon \leq \varepsilon_{\max}, \qquad |\Delta_P| \leq \delta_P, \qquad |\Delta_{\mathrm{CP}}^{(a)}| \leq \delta_{\mathrm{CP}}.$$

(ii) In conflict

In conflict otherwise.

Definition 10.100 (SynTh potential)**.** Define the nonnegative SynTh potential for $(\mathscr{S}, [\chi])$ by

$$\mathfrak{D}^{[\mathscr{S};\chi]} := w_L\left(1-\Lambda_{\mathrm{law}}^{[\mathscr{S};\chi]}\right)+w_S\left(1-\Lambda_{\mathrm{state}}^{[\mathscr{S};\chi]}\right)+u_\varepsilon\left(\frac{\varepsilon}{\varepsilon_{\max}}-1\right)_+ +u_P\left(\frac{|\Delta_P|}{\delta_P}-1\right)_+ +u_{\mathrm{CP}}\left(\frac{|\Delta_{\mathrm{CP}}^{(a)}|}{\delta_{\mathrm{CP}}}-1\right)_+, \tag{10.59}$$

with positive weights $w_L, w_S, u_\bullet$ and $(x)_+ := \max\{x, 0\}$. Co-tenability (Def. Definition 10.99) is equivalent to $\mathfrak{D}^{[\mathscr{S};\chi]} = 0$. Under IR tightening (Prop. 9.146, Lemma 10.96) the potential is nonincreasing when the *canonical spectral mask* is used for the state meter (Ch. 8: Definition 7.665).

10.10.2 Audit Moves and Basic Monotonicities

Definition 10.101 (Audit moves)**.** An *audit move* modifies the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ (Def. Definition 10.1) within the certified class while preserving BRST/Nielsen safety and band admissibility (realized weave configuration invariance: Proposition 10.10). Permitted moves are:

(i) IR tightening

IR tightening $r \mapsto r + \delta r$ with $\delta r > 0$, i.e. replacing the IR window by a tighter $K_{\mathrm{IR}}^{(r+\delta r)}$ (window/band functor: Definitions 6.280 and 7.29). This yields the monotonicities in Prop. 9.146 and Lemma 10.96.

(ii) Arena switch

Arena switch $\mathfrak{P}$: bulk $\leftrightarrow$ boundary, changing the Location component of the TLO (Def. Definition 9.147) but preserving typed meters on $\mathcal{H}_{\mathrm{phys}}$ (band/BRST hygiene: Proposition 10.10).

(iii) Spectral window refinement

Spectral window refinement $W_{\mathrm{spec}} \mapsto W'_{\mathrm{spec}}$ by *narrowing* around certified band features (Ch. 7 mask thresholds: Definition 7.665; spectral feature in (10.52)). Refinement preserves the KL positivity structure (eq. (7.12)) and maintains the one-sided improvement of the spectral entry used in onset tests (Prop. 9.146).

All moves are understood to respect the typed retuning stability of the projectors (Ch. 9: Definition 9.116, item (P7)) and the onset stability radius (Prop. 10.87). For the mixed interaction block $\mathscr{S} = \Phi\Theta$, all meters/diagnostics are applied after *typed insertion* into the off–diagonal correlator blocks of the unified correlator (cf. (10.61)); the τ–kairos enters only as a *typed projector family* (not as a sector), per Definition 10.4.

Lemma 10.102 (Monotonic effects of audit moves)**.** *Let $\mathscr{S} \in \{\Phi\Phi, \Theta\Theta, \Phi\Theta\}$ and $[\chi]$ be any admitted type (Definition 9.116, instantiated in Definitions 10.3 to 10.5). For the calibrated visibilities $\Lambda_{\mathrm{law/state}}^{[\mathscr{S};\chi]}$ (Def. Definition 10.82, (10.54)) and state features (10.52), the following hold for the audit moves of Definition 10.101:*

(i) IR tightening

Under $r\uparrow$ (tightening the IR window; Definitions 6.280 and 7.29), $\Lambda_{\mathrm{law}}^{[\mathscr{S};\chi]}(r)$ is nondecreasing, the vacuum/coherence contributions to $\Lambda_{\mathrm{state}}^{[\mathscr{S};\chi]}(r)$ are nonincreasing, and the masked spectral *contribution is nondecreasing (Prop. 9.146; see also Lemma 10.96).*

(ii) Arena switch (bulk $\leftrightarrow$ boundary)

Let $\Pi^{(\mathrm{corr})}$ be the triad coherence projector (Def. Definition 10.2). The realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ (Def. Definition 10.1) includes an arena comparison inequality *for coherence (relational boundary profile), namely a constant $\kappa_{\mathrm{coh}} \in (0, 1]$ such that*

$$\left\|\Pi^{(\mathrm{corr})}\mathfrak{O}\right\|_{\mathcal{I}}^{\mathrm{bulk}} \;\le\; \kappa_{\mathrm{coh}} \left\|\Pi^{(\mathrm{corr})}\mathfrak{O}\right\|_{\mathcal{I}}^{\mathrm{bdry}} \quad \textit{(same typed } \mathfrak{O} \textit{ on the two arenas).}$$

Therefore coherence-dominated features decrease *under the switch* bdry $\to$ bulk *(and* increase *under* bulk $\to$ bdry*). In particular, if a conflict in Definition 10.99 is due solely to boundary coherence (i.e. the only positive term in $\mathfrak{D}^{[\mathscr{S};\chi]}$ from (10.59) is the coherence channel), then the bulk switch strictly decreases $\mathfrak{D}^{[\mathscr{S};\chi]}$.*

(iii) Spectral window refinement

Let $W'_{\mathrm{spec}} \subseteq W_{\mathrm{spec}}$ *be a refinement* narrowing around the certified features *(Ch. 7 mask framework: Definition 7.665). By KL positivity (Ch. 7: eq. (7.12)) and the refinement* $W'_{\mathrm{spec}} \leq W_{\mathrm{spec}}$ *(pointwise), nuisance spectral leakage outside the certified band* does not increase*; moreover, because the certified feature lies entirely within* W'_{spec} *by construction of the refinement (Def. 10.101), the calibrated spectral entry*

$$S'_1 = \int W'_{\mathrm{spec}}\, \rho_{(s)}^{[\mathscr{S};\chi]}\, d\mu$$

satisfies $g_m(S'_1) \geq g_m(S_1)$ *for the increasing calibration* g_m *(Def. 10.82). Hence the spectral contribution to* $\Lambda_{\mathrm{state}}^{[\mathscr{S};\chi]}$ weakly improves.

Proof. (i) is exactly Prop. 9.146; see also Lemma 10.96 for the auxiliary monotone quantities (TT-leak ε, vacuum/coherence).

(ii) The arena comparison inequality is part of the certified rig's arena profile (bulk vs. boundary) and follows from the boundary-to-bulk localization property of the relational dynamics (same typed $\mathfrak{O}$ and band/BRST hygiene Proposition 10.10). Applying $\|\Pi^{(\mathrm{corr})} \cdot\|_{\mathcal{I}}$ on both arenas yields the stated inequality, which transfers via the increasing state calibration g_m (Def. 10.82) to the coherence component of Λ_{state}. When coherence is the sole positive contributor to the SynTh potential (10.59), the inequality implies a strict decrease of $\mathfrak{D}^{[\mathscr{S};\chi]}$ under bdry $\to$ bulk.

(iii) By KL positivity eq. (7.12) and $W'_{\mathrm{spec}} \leq W_{\mathrm{spec}}$, any off-band leakage cannot increase; by the refinement rule in Definition 10.101 the certified spectral feature remains fully contained in the refined window, so the raw spectral mass on the feature is preserved while leakage is reduced. The state calibration g_m is strictly increasing (Definition 10.82); therefore the calibrated spectral entry, and hence its contribution to Λ_{state}, weakly improves. These statements are taken on the physical carrier $\mathcal{H}_{\mathrm{phys}}$ with band/BRST hygiene (realized weave configuration invariance: Proposition 10.10). □

Lemma 10.103 (Bounding the TT-leak by non-spectral TT content)**.** *In the geometry sector* $\mathscr{S} = \Phi\Phi$*, the TT-leak* ε *(Def. Definition 10.32) is bounded by the TT content in the* non-spectral *triad channels:*

$$\varepsilon \;\leq\; C_{\mathrm{corr}} \left\| \Pi^{(\mathrm{corr})}\, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \delta\hat{g} \right\|_{\mathcal{I}} \;+\; C_{\mathrm{vac}} \left\| \Pi^{(\mathrm{vac})}\, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \delta\hat{g} \right\|_{\mathcal{I}}, \tag{10.60}$$

for realized weave configuration-dependent constants $C_{\mathrm{corr}}, C_{\mathrm{vac}} > 0$*. Here* $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ *is the TT projector (Def. Definition 10.3),* $\delta\hat{g}$ *is the* $\Phi\Phi$*-sector metric fluctuation, and* $\Pi^{(\mathrm{spec})}, \Pi^{(\mathrm{vac})}, \Pi^{(\mathrm{corr})}$ *are the triad idempotents (Def. Definition 10.2).*

Proof. By definition of the TT-leak (Def. Definition 10.32) and the spin-2 Riesz projector on the spectral band (typed layer; Definition 9.116),

$$\varepsilon = \left\| \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \left(1 - \mathsf{P}^{(2)}_{\mathrm{spec}}\right) \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \delta\hat{g} \right\|_{\mathcal{I}}.$$

Insert the triad decomposition $1 = \Pi^{(\mathrm{spec})} + \Pi^{(\mathrm{vac})} + \Pi^{(\mathrm{corr})}$ (Def. Definition 10.2) and use that $\mathsf{P}^{(2)}_{\mathrm{spec}}$ acts as the identity on the *spectral* spin-2 support while annihilating the non-spectral channels (standard Riesz calculus within the typed layer; Definition 9.116):

$$(1 - \mathsf{P}^{(2)}_{\mathrm{spec}})\, \Pi^{(\mathrm{spec})} = 0, \qquad (1 - \mathsf{P}^{(2)}_{\mathrm{spec}})\, \Pi^{(a)} = \Pi^{(a)} \quad \text{for } a \in \{\mathrm{vac}, \mathrm{corr}\}.$$

Hence

$$\varepsilon = \left\| \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \left(\Pi^{(\mathrm{vac})} + \Pi^{(\mathrm{corr})} \right) \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \, \delta\hat{g} \right\|_{\mathcal{I}} \;\le\; \left\| \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \, \Pi^{(\mathrm{vac})} \, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \, \delta\hat{g} \right\|_{\mathcal{I}} + \left\| \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \, \Pi^{(\mathrm{corr})} \, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \, \delta\hat{g} \right\|_{\mathcal{I}}.$$

By the typed-layer commutation on the realized weave configuration (typed projectors and the triad commute; see Definition 9.116 together with Def. Definition 10.2), we may swap $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ with $\Pi^{(a)}$ inside the $\|\cdot\|_{\mathcal{I}}$ norm. Using boundedness of the projectors in the band-invariant norm (realized weave configuration invariance; Proposition 10.10) and the triangle inequality, there exist realized weave configuration-dependent constants $C_{\mathrm{vac}}, C_{\mathrm{corr}} > 0$ such that

$$\left\| \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \, \Pi^{(a)} \, X \right\|_{\mathcal{I}} \;\le\; C_a \left\| \Pi^{(a)} \, X \right\|_{\mathcal{I}}, \qquad a \in \{\mathrm{vac}, \mathrm{corr}\}.$$

Apply this with $X = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \, \delta\hat{g}$ to obtain (10.60). □

Comments. (i) With a Hilbertian, projector-contractive choice of $\|\cdot\|_{\mathcal{I}}$ the constants can be taken as $C_{\mathrm{vac}} = C_{\mathrm{corr}} = 1$; in general we keep the explicit realized weave configuration-dependent bounds (Prop. 10.10). (ii) Under IR tightening the right-hand side is nonincreasing because the vac/corr TT norms weakly decrease (Prop. 9.146), recovering Cor. 10.104 as a corollary of the bound.

Corollary 10.104 (IR tightening decreases the TT-leak)**.** *Under IR tightening $r \uparrow$ (Ch. 8 IR window/band functor: Definitions 6.280 and 7.29; audit move: Definition 10.101), the TT-leak ε (Def. Definition 10.32) is* nonincreasing. *Moreover, if the conflict includes* correlational TT *(i.e. $\|\Pi^{(\mathrm{corr})}\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\delta\hat{g}\|_{\mathcal{I}} > 0$) and the tightening step is* strict *(the IR window shrinks on a set of positive spectral measure carrying that TT coherence), then the decrease is* strict.

Proof. Let $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ be the TT projector (Def. Definition 10.3) and $\Pi^{(\mathrm{spec})}, \Pi^{(\mathrm{vac})}, \Pi^{(\mathrm{corr})}$ the triad idempotents (Def. Definition 10.2). By Def. Definition 10.32 and the spin–2 Riesz projector on the band (typed layer; Definition 9.116),

$$\varepsilon(r) = \left\| \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \big(1 - \mathsf{P}^{(2)}_{\mathrm{spec}}(r) \big) \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \, \delta\hat{g}(r) \right\|_{\mathcal{I}}.$$

Insert $1 = \Pi^{(\mathrm{spec})} + \Pi^{(\mathrm{vac})} + \Pi^{(\mathrm{corr})}$ and use $(1 - \mathsf{P}^{(2)}_{\mathrm{spec}})\Pi^{(\mathrm{spec})} = 0$, $(1 - \mathsf{P}^{(2)}_{\mathrm{spec}})\Pi^{(a)} = \Pi^{(a)}$ for $a \in \{\mathrm{vac}, \mathrm{corr}\}$, to get

$$\varepsilon(r) = \left\| \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \big(\Pi^{(\mathrm{vac})} + \Pi^{(\mathrm{corr})} \big) \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \, \delta\hat{g}(r) \right\|_{\mathcal{I}}.$$

By boundedness of the projectors in the realized weave configuration norm (realized weave configuration invariance; Proposition 10.10) there exist $C_{\mathrm{vac}}, C_{\mathrm{corr}} > 0$ such that

$$\varepsilon(r) \;\le\; C_{\mathrm{vac}} \left\| \Pi^{(\mathrm{vac})} \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \delta\hat{g}(r) \right\|_{\mathcal{I}} \;+\; C_{\mathrm{corr}} \left\| \Pi^{(\mathrm{corr})} \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \delta\hat{g}(r) \right\|_{\mathcal{I}},$$

which is (10.60) (Lemma 10.103). Under IR tightening, the *vacuum* and *coherence* features are *nonincreasing* (Prop. 9.146); hence the right-hand side is nonincreasing in r, and so is $\varepsilon(r)$.

For strictness, assume: (i) $\left\|\Pi^{(\mathrm{corr})}\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\delta\hat{g}(r_1)\right\|_{\mathcal{I}} > 0$, and (ii) the tightening step $r_1 \to r_2$ is *strict*: $K_{\mathrm{IR}}^{(r_2)} < K_{\mathrm{IR}}^{(r_1)}$ on a set of positive spectral measure that carries the TT–coherence density (KL positivity; Ch. 8 eq. (7.12)). By positivity and L^p–contractivity of the IR window (Ch. 8; Definition 6.280) the coherence norm $\|\Pi^{(\mathrm{corr})}\Pi^{[\mathrm{TT}]}\delta\hat{g}(r)\|_{\mathcal{I}}$ *strictly* decreases from r_1 to r_2; the vacuum norm is nonincreasing (Prop. 9.146). Therefore the sum on the right-hand side of (10.60) strictly decreases, implying $\varepsilon(r_2) < \varepsilon(r_1)$.

All statements are on the physical carrier $\mathcal{H}_{\mathrm{phys}}$ and preserve gauge/BRST hygiene by Proposition 10.10 (see also the masked state monotonicity in Proposition 9.146 and the descent invariants of Lemma 10.96). □

10.10.3 Finite Audit Paths and No-Explosion

Definition 10.105 (Audit path and termination)**.** An *audit path* is a finite sequence of audit moves (Def. 10.101) applied to $\mathcal{I}^{[\mathscr{S}]}$. It *terminates successfully* if it reaches a co-tenable realized weave configuration ($\mathfrak{D} = 0$). It *terminates by certificate* if, after finitely many moves, one of the following holds persistently for all further admissible moves:

(i) Robust TT-leak
$\varepsilon > \varepsilon_{\max}$.

(ii) Robust vector parity asymmetry
$|\Delta_P| > \delta_P$.

(iii) Robust CP asymmetry
$|\Delta_{\mathrm{CP}}| > \delta_{\mathrm{CP}}$.

(iv) Absence of spectral support
$S_1 \equiv 0$ on all admissible W_{spec}.

The outcome is then a *certificate of non-co-tenability.*

Proposition 10.106 (Descent of the SynTh potential)**.** *Let $\mathscr{S} \in \{\Phi\Phi, \Theta\Theta, \Phi\Theta\}$ and $[\chi]$ be any admitted type (Ch. 10; Definition 9.116, instantiated in Definitions 10.3 to 10.5). From any* conflicting *configuration on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ (Def. Definition 10.1) there exists an* audit move *(Def. Definition 10.101) that does not increase the SynTh potential $\mathfrak{D}^{[\mathscr{S};\chi]}$ of (10.59); moreover, if the conflict is attributable to a specific diagnostic (coherence, TT-leak, parity, or CP), there exists a move that* strictly *decreases $\mathfrak{D}^{[\mathscr{S};\chi]}$.*

Proof. Recall $\mathfrak{D}^{[\mathscr{S};\chi]}$ from (10.59) and the audit moves in Definition 10.101. We proceed by cases, using the monotonicities of Lemma 10.102 and Lemma 10.96, and the TT-leak behavior in Corollary 10.104.

Case A: law/state shortfall (no trigger active). If the conflict is $\Lambda_{\mathrm{law}} < \theta_{\mathrm{law}}$ and/or masked $\Lambda_{\mathrm{state}}^{(\mathrm{spec})} < \theta_{\mathrm{state}}$ (Def. Definition 10.99), then *IR tightening* $r \uparrow$ is an audit move that makes the law meter nondecreasing and the masked spectral state meter nondecreasing (Prop. 9.146; Lemma 10.96). Hence the two positive contributions $1 - \Lambda_{\mathrm{law}}$ and $1 - \Lambda_{\mathrm{state}}$ do not increase, so $\mathfrak{D}$ does not increase. Moreover, if the tightening is *strict* on a set of positive spectral measure where either meter has positive one-sided slope (Ch. 10 onset rules: eq. (9.81),

stability Proposition 10.87), then at least one of the two terms strictly decreases, so $\mathfrak{D}$ strictly decreases.

Case B: coherence diagnostic dominates. If the conflict is attributable to coherence (the only positive term in (10.59) arising from the corr channel), perform an *arena switch* bdry $\to$ bulk (Def. 10.101). By the arena comparison in Lemma 10.102(ii), coherence decreases under bdry $\to$ bulk and, when the conflict is *solely* coherence-driven, the decrease is strict; the state calibration g_m is increasing (Def. 10.82), so the coherence contribution to $\mathfrak{D}$ strictly decreases. Thus $\mathfrak{D}$ strictly decreases.

Case C: TT-leak diagnostic active. If the conflict is due to TT-leak ε (Def. 10.32), apply *IR tightening*. By Corollary 10.104, ε is nonincreasing; if there is correlational TT on a set of positive spectral measure, the decrease is strict. Hence the penalty $u_\varepsilon(\varepsilon/\varepsilon_{\max} - 1)_+$ in (10.59) is nonincreasing and strictly decreases under the stated regularity.

Case D: parity (Δ_P) or CP ($\Delta_{\mathrm{CP}}^{(a)}$) diagnostic active. Two standard sources can drive these asymmetries (see Definitions 10.51 and 10.58): (i) boundary-amplified coherence and (ii) off-band spectral leakage.

(D1) If the asymmetry is *coherence-dominated*, the arena switch bdry $\to$ bulk reduces the coherence norm (Lemma 10.102(ii)); since both diagnostics are built from calibrated readouts of typed projectors (Defs. 10.58, 10.51, 10.82), decreasing the coherence contribution strictly reduces the corresponding normalized asymmetry whenever the asymmetry has positive support on the reduced region. Hence the penalty terms $u_P(\frac{|\Delta_P|}{\delta_P} - 1)_+$ or $u_{\mathrm{CP}}(\frac{|\Delta_{\mathrm{CP}}^{(a)}|}{\delta_{\mathrm{CP}}} - 1)_+$ strictly decrease.

(D2) If the asymmetry is driven by *off-band leakage*, perform *spectral window refinement* $W_{\mathrm{spec}} \mapsto W'_{\mathrm{spec}} \subset W_{\mathrm{spec}}$ that narrows around the certified feature (Def. 10.101). By KL positivity (Ch. 8: eq. (7.12)) and Lemma 10.102(iii), nuisance leakage cannot increase and the calibrated spectral contribution weakly improves. When the asymmetric leakage has positive mass in the excised annulus, the normalization reduces the observed asymmetry strictly, so the corresponding penalty strictly decreases.

Conclusion. In every conflicting situation, there exists at least one audit move (IR tightening, arena switch, or window refinement) that does not increase $\mathfrak{D}$ (Cases A–D). Whenever the conflict is attributable to a specific diagnostic (coherence, TT-leak, parity, CP) and the move acts on a set of positive spectral measure supporting that diagnostic's excess, the corresponding term strictly decreases, so $\mathfrak{D}$ strictly decreases. All moves preserve BRST/Nielsen safety and band admissibility (realized weave configuration invariance: Proposition 10.10). □

Theorem 10.107 (No-Explosion/Audit Theorem for the weave)**.** *For any $(\mathscr{S}, [\chi])$ with $\mathscr{S} \in \{\Phi\Phi, \Theta\Theta, \Phi\Theta\}$ and admitted type $[\chi]$ (Definition 9.116, instantiated in Definitions 10.3 to 10.5), starting from any* conflicting *configuration on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ (Def. Definition 10.1) there exists a* finite *Audit path (Def. Definition 10.95, item* (B)*) that either:*

(i) Co-tenability

Reaches co-tenability $\left(\mathfrak{D}^{[\mathscr{S};\chi]} = 0\right)$ (Def. Definition 10.99, potential (10.59)).

(ii) Certified failure

Terminates with a certified failure (the Audit certificate of §10.10, produced by the path in Def. Definition 10.95).

In particular, SynTh does not explode under the rules of (Definitions 8.98 and 8.99): Law ⊕ State becomes co-tenable after finite audits or yields a stable, falsifiable certificate.

Proof. We use the SynTh potential $\mathfrak{D}^{[\mathscr{S};\chi]}$ from (10.59), the monotone effects of audit moves in Lemma 10.102, the descent invariants in Lemma 10.96, the TT-leak behavior in Cor. 10.104, and the descent guarantee in Prop. 10.106.

Audit schedule. Choose an IR schedule $r_0 < r_1 < \dots$ with fixed step $\Delta r > 0$ smaller than the onset stability radii from Prop. 10.87 and compatible with the slope bounds in Prop. 10.88. Permit at most one arena switch (bulk↔boundary) and a *finite* nested refinement chain of spectral windows $W_{\mathrm{spec}} \supset W^{(1)}_{\mathrm{spec}} \supset \cdots \supset W^{(N)}_{\mathrm{spec}}$ (Def. 10.101; Ch. 8 fixed masks Definition 7.665). At each step, select an audit move guaranteed by Prop. 10.106 *not to increase* $\mathfrak{D}$, and if the conflict is attributable to a specific diagnostic (coherence, TT-leak, parity, CP), pick the move that *strictly decreases* $\mathfrak{D}$ (positive-measure strictness as in Cor. 10.104 and Lemma 10.102).

Monotone descent and noncreativity of triggers. By Lemma 10.96 the core descent terms $\theta_{\mathrm{law}} - \Lambda_{\mathrm{law}}$ and $\theta_{\mathrm{state}} - \Lambda^{(\mathrm{spec})}_{\mathrm{state}}$ are nonincreasing under $r \uparrow$; the TT-leak indicator is nonincreasing; arena switch and window refinement have the effects listed in Lemma 10.102. Hence the potential does not increase along the schedule; moreover, if a TT-leak or coherence source is active on a set of positive spectral measure, a strict decrease occurs (Cor. 10.104, Lemma 10.102). Automatic triggers are *sound* (Lemma 10.97 (1); see also Cor. 10.91): IR tightening cannot *create* a new TT-leak or vac/corr trigger (they are nonincreasing).

Termination by onset or certificate. If no automatic trigger is active, apply the canonical spectral mask (Definition 7.665). Then $\Lambda_{\mathrm{law}}(r)$ and the masked $\Lambda_{\mathrm{state}}(r)$ are nondecreasing (Prop. 9.146). By the slope hypotheses in Prop. 10.88, choosing $\Delta r \le \min\{\Delta_{\mathrm{law}}/m_{\mathrm{law}}, \Delta_{\mathrm{state}}/m_{\mathrm{state}}\}$ at any stage with gaps $\Delta_\bullet > 0$ forces a *finite* number of tightening steps to cross the thresholds in Def. 10.85; uniqueness/interval structure then follows from Lemma 10.86, yielding co-tenability (case (1)).

If an automatic trigger is active and persists above its bound after exhausting the *finite* window refinements and the single arena switch, or if a threshold cannot be reached because its limit value is insufficient (cf. Lemma 10.97 (2)), the path produces a *certified failure*: the certificate records sector/type, limiting visibilities, active residuals and diagnostics, and the retuning region explored (as in §10.10; see the Audit path in Def. 10.95). This is case (2).

No-Explosion. Throughout, SynTh's paraconsistent calculus (Definitions 8.98 and 8.99) routes any Law–State conflict into the Audit branch rather than trivializing inference; the finite termination above ensures the branch returns either a co-tenable TLO with onset (Definitions 9.147 and 10.85) or a falsifiable certificate. Hence no explosion occurs.

□

10.10.4 Trigger Thresholds and Automatic Audit

Definition 10.108 (Automatic audit triggers)**.** Fix a sector $\mathscr{S} \in \{\Phi\Phi, \Theta\Theta, \Phi\Theta\}$ and an admitted type $[\chi]$ (typed layer Ch. 10; Definition 9.116, instantiated in Definitions 10.3 to 10.5). Let the calibrated meters be those of Definition 10.82 and the type-specific diagnostics as in Remark 10.81 with explicit definitions in Definitions 10.32, 10.51 and 10.58. Choose a priori bounds $(\varepsilon_{\max}, \delta_P, \delta_{\mathrm{CP}}) > 0$ (cf. the mask framework of Ch. 8: Definition 7.665). We say an *automatic audit* is triggered for $[\mathscr{S}; \chi]$ if *any* of the following holds:

$$\begin{aligned} \varepsilon &> \varepsilon_{\max} && \text{(TT leak; Definition 10.32),} \\ |\Delta_P| &> \delta_P && \text{(CK parity; Definition 10.58),} \\ |\Delta_{\mathrm{CP}}^{(a)}| &> \delta_{\mathrm{CP}} && \text{(Higgs CP; Definition 10.51).} \end{aligned}$$

When an automatic trigger fires, the process *immediately* enters the Audit path (SynTh rules Ch. 9: Definitions 8.98 and 8.99; weave specialization §10.10). This restates the trigger condition captured in Corollary 10.91 and is used by the Audit path definition in §10.10.

Remark 10.109 (Admission failure vs. automatic trigger)**.** Thresholds $(\theta_{\mathrm{law}}, \theta_{\mathrm{state}}) \in (0, 1)$ are defined for the calibrated visibilities in Definition 10.82 and used by the admission predicate/onset in Definition 10.85 (see also the thresholded co-tenability in Definition 10.99). If either $\Lambda_{\mathrm{law}}^{[\mathscr{S};\chi]} < \theta_{\mathrm{law}}$ or the *masked* state visibility (canonical spectral mask; Ch. 8 Definition 7.665) satisfies $\Lambda_{\mathrm{state}}^{[\mathscr{S};\chi]} < \theta_{\mathrm{state}}$, the pair is in *conflict* and enters the Audit path due to *admission failure*, not due to an automatic trigger (cf. §10.10, Def. 10.95, item (B)). Monotone evolution of these meters under IR tightening is given by Proposition 9.146 and summarized in Lemma 10.96; descent and termination properties follow by Proposition 10.106, Lemma 10.97, and Theorem 10.107.

Corollary 10.110 (Soundness of triggers)**.** *Let $\mathscr{S} \in \{\Phi\Phi, \Theta\Theta, \Phi\Theta\}$ and $[\chi]$ be an admitted type (Ch. 10; Definition 9.116, instantiated in Definitions 10.3 to 10.5). Whenever an* automatic trigger *fires for $[\mathscr{S}; \chi]$ (Def. Definition 10.108; cf. Corollary 10.91), there exists an* audit move *(Def. Definition 10.101) that* weakly decreases *the SynTh potential $\mathfrak{D}^{[\mathscr{S};\chi]}$ of (10.59). Moreover, if the trigger concerns ε, Δ_P, or $\Delta_{\mathrm{CP}}^{(a)}$, then—unless a certificate condition already holds (Theorem 10.107)—there exists an audit move that* strictly *decreases the corresponding penalty term in (10.59).*

Proof. By Def. 10.108 (and Corollary 10.91), an automatic trigger means one of $\varepsilon > \varepsilon_{\max}$ (Definition 10.32), $|\Delta_P| > \delta_P$ (Definition 10.58), or $|\Delta_{\mathrm{CP}}^{(a)}| > \delta_{\mathrm{CP}}$ (Definition 10.51) holds. Such a configuration is *conflicting* under the thresholded co-tenability definition (Def. 10.99) and contributes a positive penalty to $\mathfrak{D}^{[\mathscr{S};\chi]}$ (Eq. (10.59)).

Weak decrease. From any conflict there exists an audit move that does not increase $\mathfrak{D}$ by Prop. 10.106. Concretely: - For ε (TT-leak), *IR tightening* $r \uparrow$ makes ε nonincreasing (Cor. 10.104); the penalty $u_\varepsilon(\varepsilon/\varepsilon_{\max} - 1)_+$ therefore does not increase. - For parity/CP diagnostics, either an *arena switch* (boundary→bulk) or a *spectral window refinement* weakly reduces the asymmetry carried by coherence or off-band leakage, respectively (Lemma 10.102(ii),(iii), KL positivity eq. (7.12)). Since g_m is increasing (Def. 10.82), the normalized penalties $u_P(\frac{|\Delta_P|}{\delta_P} - 1)_+$ and $u_{\mathrm{CP}}(\frac{|\Delta_{\mathrm{CP}}^{(a)}|}{\delta_{\mathrm{CP}}} - 1)_+$ are weakly reduced.

Strict decrease unless certified. If the trigger concerns a specific diagnostic and the offending content occupies a set of positive spectral measure: - For ε: strict IR tightening that removes TT-coherence support of positive measure yields $\varepsilon(r_2) < \varepsilon(r_1)$ (Cor. 10.104), hence a *strict* drop in its penalty. - For Δ_P or $\Delta_{\mathrm{CP}}^{(a)}$: if the excess is coherence-dominated, boundary→bulk strictly reduces the coherence norm (Lemma 10.102(ii)); if it is off-band leakage, a refinement $W_{\mathrm{spec}} \mapsto W'_{\mathrm{spec}}$ that excises an annulus of positive measure around the certified feature strictly reduces the asymmetry contribution (Lemma 10.102(iii) with eq. (7.12)). In both subcases the corresponding penalty in (10.59) strictly decreases.

If no admissible move can strictly reduce the active penalty while the trigger remains, the situation meets the certificate scenario in Theorem 10.107 (case (2)), and Audit terminates by certificate rather than continuing descent.

Thus, whenever an automatic trigger fires, there is always an audit move that weakly decreases $\mathfrak{D}$, and—except in certificate cases—one that strictly decreases the triggered penalty. □

Remark 10.111 (Minimality of audit moves)**.** The three audit moves of Definition 10.101 are *minimal and sufficient* for SynTh on the realized weave configuration $\mathcal{I}^{[\mathcal{S}]}$ (Def. Definition 10.1):

(i) IR retuning
IR retuning $r \mapsto r + \delta r$ acts through the IR window $K_{\mathrm{IR}}^{(r)}$ (Ch. 6: Definitions 6.280 and 7.29) and is the only knob that changes long-distance content while preserving band admissibility; its effects on the meters/diagnostics are exactly those in Proposition 9.146, Lemma 10.96, and Corollary 10.104.

(ii) Arena switch
Arena switch $\mathfrak{P}$: bulk $\leftrightarrow$ boundary is the canonical way to alter the Location component (Def. Definition 9.147) without changing typed projectors; its coherence comparison is captured in Lemma 10.102(ii).

(iii) Spectral window refinement
Spectral window refinement $W_{\mathrm{spec}} \mapsto W'_{\mathrm{spec}}$ (masks: Definition 7.665) is the only certified mechanism to remove off-band leakage while preserving the feature; its monotone effect is Lemma 10.102(iii) with KL positivity eq. (7.12).

Any further "micro-tuning" (e.g. band-tag or sheet micro-adjustments) factors through these moves by *band covariance* and projector stability on the realized weave configuration (Proposition 10.10 and stability Definition 9.116, item (P7)); consequently, their impact on visibilities/diagnostics is subsumed by the monotonicities already listed.

Remark 10.112 (Reporting after audit)**.** Upon termination of the Audit path (Theorem 10.107), Step (7) reports, for each $(\mathcal{S}, [\chi])$:

(i) Triad-resolved outcomes
The triad-resolved outcomes $(\mathrm{spec}, \mathrm{vac}, \mathrm{corr})$ with the corresponding calibrated entries (cf. Definitions 10.2 and 10.82 and eq. (10.52)).

(ii) Arena
The used arena $\mathfrak{P} \in \{\mathrm{bulk}, \mathrm{boundary}\}$ (Def. Definition 10.101).

(iii) Certified onsets
The certified onsets $r^\star[\chi]$ (Def. Definition 10.85, uniqueness/stability via Lemma 10.86 and Proposition 10.87).

(iv) Parasymmetric TLO readout
The Parasymmetric TLO readout (Def. Definition 9.147)

$$\mathrm{TLO}^{[\chi]} = (\mathrm{Type}, \mathrm{Location}, \mathrm{Onset})$$

(v) Outcome label
The outcome is labelled co-tenability, $\mathfrak{D}^{[\mathscr{S};\chi]} = 0$ (Definition 10.99 and eq. (10.59)) or a certificate that lists the active diagnostic(s) and the explored retuning region bounded by the stability radii (Theorem 10.107, Prop. 10.87, and triggers Definition 10.108 and Corollary 10.91).

All reported quantities are evaluated on the physical carrier $\mathcal{H}_{\mathrm{phys}}$ and inherit gauge/BRST hygiene by Proposition 10.10; sectorization is strictly $\mathscr{S} \in \{\Phi\Phi, \Theta\Theta, \Phi\Theta\}$ with τ–kairos entering only via typed projectors (Definition 10.4).

10.11 Weaving the Sectors via the Unified Inter-Modal Correlator

We complete Step (6) by inserting the *typed* projectors into the Unified Inter–Modal Correlator and analyzing mixing on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ (Def. Definition 10.1). We use the triad projectors $\Pi^{(a)}$ (Def. Definition 10.2), the TT map $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ (Def. Definition 10.3), the τ–kairos typed families (Def. Definition 10.4), and the Higgs CP–refined projectors (Def. Definition 10.5). The global correlator $\hat{\mathbf{G}}_\Lambda$ is as defined upstream (Def. Definition 6.4).

Typed insertion into $\hat{\mathbf{G}}_\Lambda$.

Definition 10.113 (Typed and triad-resolved correlator)**.** Let $[\chi]$ and $[\chi']$ be admitted types from the typed layer (Ch. 10; Definition 9.116, instantiated in Definitions 10.3 to 10.5). For $a, b \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$ define the *typed, triad-resolved correlator*

$$\hat{\mathbf{G}}_\Lambda^{(a\to b)\,[\chi\to\chi']} := \mathsf{P}_{\mathrm{phys}}\, \Pi^{(b)}\, \Pi^{[\chi']}\, \hat{\mathbf{G}}_\Lambda\, \Pi^{[\chi]}\, \Pi^{(a)}\, \mathsf{P}_{\mathrm{phys}}, \tag{10.61}$$

where $\Pi^{[\chi]}$ denotes the *product* of the relevant typed projectors for $[\chi]$ (e.g. $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ for gravitons; $A^{(\pm)}$ for Chrono–Kymon vectors; $\Pi^{(\mathrm{CP}\pm)}\Pi_\Theta^{[\mathrm{rad}\pm]}$ for Higgs CP/refined radial modes; cf. Definitions 10.3 to 10.5). The sector-diagonal blocks $\hat{G}_{\Phi\Phi}, \hat{G}_{\Theta\Theta}$ and the mixed blocks $\hat{G}_{\Phi\Theta}, \hat{G}_{\Theta\Phi}$ (Def. Definition 6.4) are thereby refined to typed channels $[\chi] \to [\chi']$ with explicit triad provenance $a \to b$.

Remark 10.114 (Band and BRST safety; commutation on the realized weave configuration)**.** On the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$, the maps in (10.61) preserve band admissibility and gauge-parameter independence by Proposition 10.10 (all operators are evaluated on $\mathcal{H}_{\mathrm{phys}}$). Moreover, the typed projectors (Ch. 10: Definition 9.116) commute with the triad

idempotents from Definition 10.2 on the realized weave configuration, so the order in (10.61) is unambiguous up to BRST-exact/EOM terms (cf. Remark 10.93). Continuity under admissible retuning follows from typed stability (Definition 9.116, item (P7)).

Remark 10.115 (Sector note: mixed $\Phi\Theta$ channels)**.** For the mixed interaction block $\mathscr{S} = \Phi\Theta$, the construction (10.61) is applied *after typed insertion* into the off–diagonal correlator blocks of $\hat{\mathbf{G}}_\Lambda$ (Def. Definition 6.4). Here τ–kairos appears only via the *typed* projectors in $\Pi^{[\chi]}$ (Def. Definition 10.4); it is *not* a sector. All triad-resolved channels $a \to b$ are reported with their certified provenance (Definition 10.2).

10.11.1 Σ-weighted Generalized Eigenproblem for Mixing

Definition 10.116 ($\bar{\Sigma}$–Hermitian generalized eigen-operator pairs)**.** Let $(\mathcal{H}, \langle\cdot,\cdot\rangle_{\bar{\Sigma}})$ be a $\bar{\Sigma}$–weighted Hilbert space (Chs. 6–7). For linear maps $A, B : \mathcal{H} \to \mathcal{H}$ with $\bar{\Sigma}$–adjoints $A^{\dagger_{\bar{\Sigma}}}, B^{\dagger_{\bar{\Sigma}}}$:

(i) $\bar{\Sigma}$–Hermitian generalized eigen-operator pair

The ordered pair (A, B) is called a $\bar{\Sigma}$–*Hermitian generalized eigen-operator pair* if

$$A = A^{\dagger_{\bar{\Sigma}}} \qquad \text{and} \qquad B = B^{\dagger_{\bar{\Sigma}}}.$$

Associated to (A, B) is the one–parameter family of operators $A - \lambda B$ ($\lambda \in \mathbb{C}$) and the generalized eigenproblem $Av = \lambda Bv$.

(ii) Generalized eigenpair and Rayleigh quotient

A *generalized eigenpair* (λ, v) satisfies $Av = \lambda Bv$, $v \neq 0$. If B is positive (semi)definite on $\operatorname{Ran} B$, the Rayleigh quotient is

$$\lambda(v) = \frac{\langle v, Av\rangle_{\bar{\Sigma}}}{\langle v, Bv\rangle_{\bar{\Sigma}}}$$

on that support; whitening by $B^{1/2}$ yields a standard Hermitian problem (cf. Remark 10.119).

Definition 10.117 ($\bar{\Sigma}$-weighted inner products, adjoints, and typed generalized eigen-operator pairs)**.** Let $\langle\cdot,\cdot\rangle_{\bar{\Sigma}}$ be the $\bar{\Sigma}$-weighted inner product on the relevant carriers (Chs. 6–7). For a linear map X on (or between) typed subspaces of $\mathcal{H}_{\text{phys}}$, define the $\bar{\Sigma}$-adjoint by

$$X^{\dagger_{\bar{\Sigma}}} \text{ such that } \langle Xu, v\rangle_{\bar{\Sigma}} = \langle u, X^{\dagger_{\bar{\Sigma}}} v\rangle_{\bar{\Sigma}}.$$

Set $\mathrm{Re}_{\bar{\Sigma}}(X) = \frac{1}{2}(X + X^{\dagger_{\bar{\Sigma}}})$ and $\mathrm{Im}_{\bar{\Sigma}}(X) = \frac{1}{2\mathrm{i}}(X - X^{\dagger_{\bar{\Sigma}}})$, both $\bar{\Sigma}$-Hermitian. With the typed insertion $\hat{\mathbf{G}}_\Lambda^{[\chi\to\chi']}$ (Definition 10.113, (10.61)), define the *typed generalized eigen-operator pair*

$$A^{[\chi\to\chi']} := \mathrm{Re}_{\bar{\Sigma}}\big(\hat{\mathbf{G}}_\Lambda^{[\chi\to\chi']}\big), \qquad B^{[\chi\to\chi']} := \mathrm{Im}_{\bar{\Sigma}}\big(\hat{\mathbf{G}}_\Lambda^{[\chi\to\chi']}\big), \tag{10.62}$$

and, for $[\chi] = [\chi']$, the diagonal metric

$$M^{[\chi]} := \mathrm{Re}_{\bar{\Sigma}}\big(\hat{\mathbf{G}}_\Lambda^{[\chi\to\chi]}\big).$$

All maps are evaluated on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ and are BRST/band safe by Proposition 10.10.

Definition 10.118 ($\bar{\Sigma}$-weighted generalized eigenproblem (mixing))**. (a) Intra-type ($[\chi] = [\chi']$).** On the typed carrier $\mathcal{H}_{[\chi]} = \operatorname{Ran}\Pi^{[\chi]} \subset \mathcal{H}_{\text{phys}}$,

$$A^{[\chi\to\chi]}v \;=\; \lambda\, B^{[\chi\to\chi]}v, \qquad v \in \mathcal{H}_{[\chi]}. \tag{10.63}$$

(b) Cross-type ($[\chi] \neq [\chi']$). Because $\hat{\mathbf{G}}_\Lambda^{[\chi\to\chi']} : \mathcal{H}_{[\chi]} \to \mathcal{H}_{[\chi']}$ maps across distinct typed spaces, we realize the corresponding generalized eigen-operator pair on the *direct sum*

$$\mathcal{H}_{[\chi]\oplus[\chi']} := \mathcal{H}_{[\chi]} \oplus \mathcal{H}_{[\chi']}, \quad \langle (v,w),(v',w')\rangle_{\bar{\Sigma}} := \langle v,v'\rangle_{\bar{\Sigma}} + \langle w,w'\rangle_{\bar{\Sigma}},$$

with block forms

$$A_{\text{mix}} = \begin{pmatrix} 0 & \operatorname{Re}_{\bar{\Sigma}}\big(\hat{\mathbf{G}}_\Lambda^{[\chi\to\chi']}\big) \\ \operatorname{Re}_{\bar{\Sigma}}\big(\hat{\mathbf{G}}_\Lambda^{[\chi'\to\chi]}\big) & 0 \end{pmatrix}, \qquad B_{\text{mix}} = \begin{pmatrix} 0 & \operatorname{Im}_{\bar{\Sigma}}\big(\hat{\mathbf{G}}_\Lambda^{[\chi\to\chi']}\big) \\ \operatorname{Im}_{\bar{\Sigma}}\big(\hat{\mathbf{G}}_\Lambda^{[\chi'\to\chi]}\big) & 0 \end{pmatrix}, \tag{10.64}$$

so that

$$A_{\text{mix}}\begin{pmatrix} v \\ w \end{pmatrix} = \lambda\, B_{\text{mix}}\begin{pmatrix} v \\ w \end{pmatrix}, \qquad (v,w) \in \mathcal{H}_{[\chi]\oplus[\chi']}.$$

Equivalently, one may linearize on the graph space or use a Rayleigh quotient on $\mathcal{H}_{[\chi]\oplus[\chi']}$. The spectrum λ quantifies the strength/phase of $[\chi] \leftrightarrow [\chi']$ mixing within the band-limited, BRST-projected space (Def. Definition 10.1).

Remark 10.119 (Well-posedness and whitening)**.** By KL-positivity on the certified band (Ch. 8: eq. (7.12)) and typed stability (Ch. 10: Definition 9.116, item (P7)), the diagonal metrics $M^{[\chi]} = \operatorname{Re}_{\bar{\Sigma}}(\hat{\mathbf{G}}_\Lambda^{[\chi\to\chi]})$ and $M^{[\chi']}$ are positive semidefinite and strictly positive on their supports. In particular, on $\operatorname{Ran} M^{[\chi]}$ (and similarly for $M^{[\chi']}$) there exist unique positive square roots $M^{[\chi]1/2}$, $M^{[\chi]-1/2}$ satisfying $M^{[\chi]} = M^{[\chi]1/2}M^{[\chi]1/2}$.

On those supports, one may *whiten* the generalized eigen-operator pair (A,B) by the similarity transform with $M^{-1/2}$: for $v \in \operatorname{Ran} M^{[\chi]}$ set $u := M^{[\chi]1/2}v$ and define

$$\tilde{A} := M^{[\chi]-1/2} A\, M^{[\chi]-1/2}, \qquad \tilde{B} := M^{[\chi]-1/2} B\, M^{[\chi]-1/2}.$$

Then the generalized eigenproblem $Av = \lambda Bv$ is equivalent to

$$\tilde{A}u = \lambda\, \tilde{B}u$$

on $\operatorname{Ran} M^{[\chi]}$, with $\tilde{A}$ and $\tilde{B}$ $\bar{\Sigma}$-Hermitian. In particular, whenever B coincides with $M^{[\chi]}$ (or with a block-diagonal metric built from $M^{[\chi]}$, $M^{[\chi']}$ in the cross-type case), this whitening reduces the problem to a standard Hermitian eigenproblem $\tilde{A}u = \lambda u$ in the $\bar{\Sigma}$-inner product (cf. Definition 10.116).

If B (or B_{mix}) is singular, restrict to $\operatorname{Ran} B$ or use a certified Tikhonov regularization $B \mapsto B + \eta M$ with $\eta > 0$ small; all operations remain within $\mathcal{I}^{[\mathscr{S}]}$ by Proposition 10.10.

Remark 10.120 (Sector and mixed-block usage)**.** For sector-diagonal channels ($\Phi\Phi$ or $\Theta\Theta$) the typed blocks in (10.62) act within the same carrier; for the mixed block $\Phi\Theta$ (and $\Theta\Phi$), use the typed insertion (10.61) on the off–diagonal correlator, and then the direct-sum generalized eigen-operator pair (10.64) (cf. Remark 10.115). In all cases τ–kairos enters only via typed projectors (Definition 10.4), not as a sector label.

Remark 10.121 (Gauge/BRST and band safety)**.** All maps/forms above are sandwiched by $\mathsf{P}_{\mathrm{phys}}$ (Def. Definition 10.1) and composed with typed/triad idempotents (Definitions 10.2 to 10.5), hence inherit gauge-parameter independence and band admissibility by Proposition 10.10. Continuity under admissible retunings follows from Definition 9.116, item (P7).

Lemma 10.122 (Well-posedness on certified bands)**.** *Assume the certified band avoids spectral singularities (no poles on the fixed Riesz contours) for the diagonal typed blocks of the unified correlator (Def. Definition 10.113, (10.61)). Then the $\bar{\Sigma}$-weighted generalized eigenproblems are well-posed on the realized weave configuration $\mathcal{I}^{[\mathcal{S}]}$:*

(i) Intra-type
In (10.63), $B^{[\chi\to\chi]}$ is bounded on $\mathcal{H}_{[\chi]}$ and $A^{[\chi\to\chi]}$ is compact; hence the spectrum $\{\lambda\}$ is discrete with possible accumulation only at 0.

(ii) Cross-type
For the block generalized eigen-operator pair (10.64) on $\mathcal{H}_{[\chi]\oplus[\chi']}$, B_{mix} is bounded and A_{mix} is compact; hence the spectrum is discrete with possible accumulation only at 0.

Proof. **Step 1: Boundedness on the realized weave configuration.** All operators are evaluated on the realized weave configuration $\mathcal{I}^{[\mathcal{S}]}$ (Def. Definition 10.1). By realized weave configuration hygiene and band admissibility (Proposition 10.10) and typed stability (Ch. 10: Definition 9.116, item (P7)), the typed/triad idempotents are bounded. The unified correlator $\hat{\mathbf{G}}_\Lambda$ restricted to the certified band is bounded on $\mathcal{H}_{\mathrm{phys}}$ (no spectral singularities on the fixed contours; Riesz selectors Definition 10.7). Therefore the typed insertions $\hat{\mathbf{G}}_\Lambda^{[\chi\to\chi']}$ (Def. Definition 10.113) are bounded, and so are their $\bar{\Sigma}$-Hermitian parts

$$A := \mathrm{Re}_{\bar{\Sigma}}(\hat{\mathbf{G}}_\Lambda^{[\chi\to\chi']}), \qquad B := \mathrm{Im}_{\bar{\Sigma}}(\hat{\mathbf{G}}_\Lambda^{[\chi\to\chi']})$$

(Definition 10.117, (10.62)). Thus $B^{[\chi\to\chi']}$ is bounded in both intra- and cross-type settings; likewise for the block B_{mix} in (10.64).

Step 2: Compactness of A. On the certified band, the real part of the correlator can be written, contour-wise, as a finite linear combination of Riesz-integrals of resolvents (no poles on the contours):

$$\mathrm{Re}_{\bar{\Sigma}}(\hat{\mathbf{G}}_\Lambda^{[\chi\to\chi']}) = \sum_\ell \int_{\Gamma_\ell} K_\ell(\zeta)\,(\zeta - \hat{\mathbf{H}})^{-1}\,d\zeta,$$

for suitable scalar weights K_ℓ and a band-limited generator $\hat{\mathbf{H}}$ (schematic; the exact representation is via the standard Cauchy integral applied to the typed insertions; cf. Riesz selectors Definition 10.7). Under band-limitation (Ch. 8 masks; Definition 7.665) and the absence of singularities on the contours, these contour integrals define operators whose kernels are square-integrable on the realized weave configuration (Hilbert–Schmidt), hence compact. Typed and triad projectors commute on the realized weave configuration (Ch. 10: Definition 9.116, item (P7)), so left/right multiplication by them preserves compactness. Therefore $A^{[\chi\to\chi']}$ is compact; in particular $A^{[\chi\to\chi]}$ is compact on $\mathcal{H}_{[\chi]}$, and the off-diagonal blocks entering A_{mix} in (10.64) are compact, making A_{mix} compact on $\mathcal{H}_{[\chi]\oplus[\chi']}$.

Step 3: Reduction to a compact self-adjoint problem on $\mathrm{Ran}\,B$**.** Restrict to the (closed) range $\mathcal{H}_B := \overline{\mathrm{Ran}\,B}$ (or $\overline{\mathrm{Ran}\,B_{\mathrm{mix}}}$ in the cross-type case). On $\mathcal{H}_B$, either B is strictly positive

(on the certified support; KL positivity eq. (7.12)) or one may use the standard certified regularization $B_\eta := B + \eta M$ with $M := M^{[\chi]}$ (or $M^{[\chi']}$ blockwise) from Definition 10.117 and $\eta > 0$ small (cf. Remark 10.119). Define the whitening transform $T_\eta := B_\eta^{-1/2}\, A\, B_\eta^{-1/2}$ on $\mathcal{H}_B$. Since A is compact and $B_\eta^{-1/2}$ is bounded on $\mathcal{H}_B$, T_η is compact (and $\bar{\Sigma}$-Hermitian). Letting $\eta \downarrow 0$ on $\mathcal{H}_B$ recovers the same nonzero spectrum (standard limit argument; see Remark 10.119). Therefore the generalized spectrum of (A, B) is the (possibly zero-accumulating) point spectrum of the compact operator T.

Step 4: Discreteness and accumulation at 0. By the spectral theorem for compact (Hermitian) operators, the spectrum is discrete with finite multiplicities and may only accumulate at 0. Mapping back through the whitening transform yields the same property for the generalized eigenvalues of (10.63) (intra-type) and of the block problem (10.64) (cross-type). All constructions remain within $\mathcal{I}^{[\mathcal{J}]}$ and preserve BRST/gauge and band safety by Proposition 10.10. □

10.11.2 TT-null Preservation for τ-mixing

Lemma 10.123 (TT-null preservation under any τ-mixing)**.** *Let $X \in \{S^{(b)}, S^{(L)}, \tilde{S}^{(p)}, A^{(\pm)}\}$ be any τ-kairos type (Def. Definition 10.4) and let Y be any admitted type in the gaiamiti or tantrakyma families (i.e. typed in the $\Phi\Phi$ or $\Theta\Theta$ sectors; cf. Definitions 10.3 and 10.5). Then, for the typed correlator blocks of Def. Definition 10.113,*

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \hat{\mathbf{G}}_\Lambda^{[\,X\to Y\,]} \;=\; 0, \qquad \hat{\mathbf{G}}_\Lambda^{[\,Y\to X\,]}\, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \;=\; 0, \tag{10.65}$$

as operators on the realized weave configuration $\mathcal{I}^{[\mathcal{J}]}$ (Def. Definition 10.1). The same statements hold triad-wise for each $(a\to b)$ block in (10.61).

Proof. Write the typed insertion (Def. Definition 10.113) without triad indices for brevity:

$$\hat{\mathbf{G}}_\Lambda^{[\,X\to Y\,]} = \mathsf{P}_{\mathrm{phys}}\, \Pi^{[Y]}\, \hat{\mathbf{G}}_\Lambda\, \Pi^{[X]}\, \mathsf{P}_{\mathrm{phys}}, \qquad \hat{\mathbf{G}}_\Lambda^{[\,Y\to X\,]} = \mathsf{P}_{\mathrm{phys}}\, \Pi^{[X]}\, \hat{\mathbf{G}}_\Lambda\, \Pi^{[Y]}\, \mathsf{P}_{\mathrm{phys}}.$$

By the τ–TT purity lemma Lemma 10.17 and TT projector definition Definition 10.3, the τ-typed ranges/kernels satisfy

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \Pi^{[X]} = 0, \qquad \Pi^{[X]}\, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} = 0,$$

and, when extended block-diagonally to the full carrier, $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ acts as zero on non-geometry components. Using typed stability/commutation on the realized weave configuration (Ch. 10: Definition 9.116, item (P7); see also Remark 10.114) and the fact that $\mathsf{P}_{\mathrm{phys}}$ preserves these relations on $\mathcal{H}_{\mathrm{phys}}$ (Proposition 10.10), we compute

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \hat{\mathbf{G}}_\Lambda^{[\,X\to Y\,]} = \mathsf{P}_{\mathrm{phys}} \left(\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \Pi^{[Y]}\right) \hat{\mathbf{G}}_\Lambda\, \Pi^{[X]}\, \mathsf{P}_{\mathrm{phys}}.$$

If Y is a non-TT type (e.g. Higgs CP-refined or other $\Theta\Theta$ typed object), then $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\Pi^{[Y]} = 0$ by orthogonality of the typed algebra, giving zero. If $Y = \mathrm{TT}$, then $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\Pi^{[Y]} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$, but the right factor $\Pi^{[X]}$ is τ-typed and annihilated by TT on the geometry side; equivalently, $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \hat{\mathbf{G}}_\Lambda\, \Pi^{[X]} = 0$ by τ–TT purity (Lemma 10.17). Thus the left identity in (10.65) holds.

The right identity is analogous:

$$\hat{\mathbf{G}}_{\Lambda}^{[\,Y\to X\,]}\,\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} = \mathsf{P}_{\mathrm{phys}}\,\Pi^{[X]}\,\hat{\mathbf{G}}_{\Lambda}\left(\Pi^{[Y]}\,\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\right)\mathsf{P}_{\mathrm{phys}}.$$

If $Y \neq \mathrm{TT}$, then $\Pi^{[Y]}\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} = 0$; if $Y = \mathrm{TT}$, then $\Pi^{[X]}\hat{\mathbf{G}}_{\Lambda}\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} = 0$ again by τ–TT purity. Triad-resolved statements follow by inserting $\Pi^{(a)}/\Pi^{(b)}$ (Def. Definition 10.2) on the right/left of $\hat{\mathbf{G}}_{\Lambda}$; these commute with the typed projectors on the realized weave configuration (Ch. 10: Definition 9.116, item (P7)), so each $(a\to b)$ block is separately annihilated. All steps occur on $\mathcal{H}_{\mathrm{phys}}$ and inherit BRST/gauge safety by Proposition 10.10. □

Remark 10.124 (Mixed $\Phi\Theta$ channels)**.** For the mixed block $\mathscr{S} = \Phi\Theta$, apply the same calculation to the off-diagonal typed insertion (Def. Definition 10.113, (10.61); cf. Remark 10.115). TT acts purely on the geometry slot and remains null on any τ insertion by Lemma 10.17; thus (10.65) holds verbatim in the mixed case.

Corollary 10.125 (No induced TT from τ-mixing)**.** *The graviton readout $\hat{h}_{\mu\nu}$ (TT extraction; Definition 10.25, using the TT sandwich (10.28)) is unaffected by any τ-mixing channel: TT content cannot be generated by coupling to τ–kairos. Equivalently, the TT component of the typed, triad-resolved correlator remains unchanged when any factor $\hat{\mathbf{G}}_{\Lambda}^{[\,X\to Y\,]}$ or $\hat{\mathbf{G}}_{\Lambda}^{[\,Y\to X\,]}$ with a τ–type $X \in \{S^{(b)}, S^{(L)}, \tilde{S}^{(p)}, A^{(\pm)}\}$ (Def. Definition 10.4) is inserted.*

Proof. By the typed insertion rule (Definition 10.113, (10.61)) any contribution to the readout that involves a τ–typed factor has the schematic form

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\hat{\mathbf{G}}_{\Lambda}^{[\,X\to Y\,]} \quad \text{or} \quad \hat{\mathbf{G}}_{\Lambda}^{[\,Y\to X\,]}\,\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]},$$

possibly with triad idempotents $\Pi^{(a)}, \Pi^{(b)}$ on the right/left (Def. Definition 10.2). By TT–null preservation for τ–mixing (Lemma 10.123, identity (10.65)), each such TT–sandwiched block vanishes on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ (Def. Definition 10.1). Therefore, when the TT sandwich that defines the graviton readout (10.28) (see Definitions 10.3 and 10.25) is applied to any correlator decomposition containing a τ–mixing channel, all τ–dependent terms drop out, leaving the TT readout unchanged.

The same conclusion holds triad-wise because the $(a \to b)$ blocks of (10.61) inherit Lemma 10.123 (the triad/typed projectors commute on the realized weave configuration; Ch. 10 stability Definition 9.116, item (P7)). For the mixed $\Phi\Theta$ channel, apply the argument to the off-diagonal typed insertion (Def. Definition 10.113; Remark 10.115); TT acts on the geometry slot and still annihilates any τ insertion. Gauge/BRST and band safety are preserved throughout by Proposition 10.10. □

10.11.3 Selection Rules for Typed Mixing

Proposition 10.126 (Symmetry selection rules (parity, helicity, CP))**.** *Let $\mathcal{P}$ denote spatial parity on the certified band,* Hel *the helicity operator on TT and τ–vector subspaces, and* CP *the CP action on the scalar subspace. Assume the unified correlator $\hat{\mathbf{G}}_{\Lambda}$ (with typed insertion as in Definition 10.113 and eq. (10.61))* commutes *on $\mathcal{H}_{\mathrm{phys}}$ with the relevant symmetry:*

$$\mathcal{P}\,\hat{\mathbf{G}}_{\Lambda}\,\mathcal{P}^{-1} = \hat{\mathbf{G}}_{\Lambda}, \qquad \mathsf{Hel}\,\hat{\mathbf{G}}_{\Lambda} = \hat{\mathbf{G}}_{\Lambda}\,\mathsf{Hel}, \qquad \mathsf{CP}\,\hat{\mathbf{G}}_{\Lambda}\,\mathsf{CP}^{-1} = \hat{\mathbf{G}}_{\Lambda},$$

respectively (band/BRST safety: Proposition 10.10). Then:

(i) Parity

If $\mathcal{P}$–invariance holds, a parity–even type does not *mix with a parity–odd type. In particular, using the parity assignments from Definitions 10.3 to 10.5,*

$$\hat{\mathbf{G}}_\Lambda^{[\ \tilde{S}^{(p)}\to \mathrm{TT}\ ]} = 0, \qquad \hat{\mathbf{G}}_\Lambda^{[\ \tilde{S}^{(p)}\to \mathrm{rad}+\ ]} = 0,$$

and similarly for the reverse channels.

(ii) Helicity

If $\hat{\mathbf{G}}_\Lambda$ is helicity–diagonal on TT, then the TT helicity sectors do not mix:

$$\Pi^{(+2)}\, \hat{\mathbf{G}}_\Lambda^{[\ \mathrm{TT}\to\mathrm{TT}\ ]}\, \Pi^{(-2)} = 0$$

and vice versa (TT helicity split within the typed layer; Ch. 10 Definition 9.116). Likewise, for τ–vectors, $A^{(+)}$ does not mix with $A^{(-)}$ (Definitions 10.4 and 10.58).

(iii) CP

If $\hat{\mathbf{G}}_\Lambda$ is CP–invariant on scalars, the Higgs CP–refined modes do not mix:

$$\hat{\mathbf{G}}_\Lambda^{[\ \mathrm{rad}+\to\mathrm{rad}-\ ]} = 0$$

and the reverse channel vanishes (Definition 10.5).

All statements hold triad–wise for each $(a\to b)$ block in (10.61).

Proof. We give a uniform argument using symmetry projectors. Let S be any of $\mathcal{P}$, Hel, or CP, and let $\Pi^{(s)}$ denote the spectral/idempotent projector onto the symmetry sector labeled by s (e.g. parity $\pm$, helicity ± 2 on TT or $\pm$ on CK, CP $\pm$ on scalars). On $\mathcal{H}_{\mathrm{phys}}$, assume S commutes with $\hat{\mathbf{G}}_\Lambda$ and preserves the typed/triad subspaces (typed stability on the realized weave configuration: Ch. 10 Definition 9.116, item (P7); realized weave configuration hygiene: Proposition 10.10).

Let $[\chi] \to [\chi']$ be a typed channel with definite S–labels s and s' respectively (the labels are fixed by the type definitions: Definitions 10.3 to 10.5). Using the typed insertion (Def. Definition 10.113),

$$\hat{\mathbf{G}}_\Lambda^{[\chi\to\chi']} = \mathsf{P}_{\mathrm{phys}}\, \Pi^{[\chi']}\, \hat{\mathbf{G}}_\Lambda\, \Pi^{[\chi]}\, \mathsf{P}_{\mathrm{phys}} = \mathsf{P}_{\mathrm{phys}}\, \Pi^{[\chi']}\, \Pi^{(s')}\, \hat{\mathbf{G}}_\Lambda\, \Pi^{(s)}\, \Pi^{[\chi]}\, \mathsf{P}_{\mathrm{phys}},$$

where we inserted $\Pi^{(s')}$ on the left and $\Pi^{(s)}$ on the right because $\Pi^{[\chi']}/\Pi^{[\chi]}$ refine to definite S–labels by construction. Since S commutes with $\hat{\mathbf{G}}_\Lambda$, we have $\Pi^{(s')}\, \hat{\mathbf{G}}_\Lambda\, \Pi^{(s)} = \hat{\mathbf{G}}_\Lambda\, \Pi^{(s')}\Pi^{(s)}$. If $s' \neq s$ (opposite parity, opposite helicity, or opposite CP), orthogonality yields $\Pi^{(s')}\Pi^{(s)} = 0$, hence $\hat{\mathbf{G}}_\Lambda^{[\chi\to\chi']} = 0$. This proves (1)–(3). Triad–resolved versions follow by inserting $\Pi^{(a)}/\Pi^{(b)}$ (Def. Definition 10.2) on the right/left of $\hat{\mathbf{G}}_\Lambda$; these idempotents commute with the symmetry projectors and the typed layer on the realized weave configuration (Ch. 10 Definition 9.116, item (P7)), so each $(a\to b)$ block vanishes under the same mismatch condition. □

Remark 10.127 (When selection rules can be violated)**.** If a background or the boundary machine breaks a symmetry, the corresponding commutation with $\hat{\mathbf{G}}_\Lambda$ fails (e.g. $\mathcal{P}$–, helicity–, or CP–violation). The same typed formalism applies, and the magnitude of violation is

quantified by the norms of the previously forbidden off–diagonal typed blocks (e.g. parity–odd mass in a parity–even channel, helicity cross–terms TT $+2 \leftrightarrow -2$, or rad$+ \leftrightarrow$ rad$-$). Operationally, these appear as diagnostics already tracked in the meters: CK vector parity asymmetry Δ_P (Definition 10.58) and Higgs CP asymmetry $\Delta_{\mathrm{CP}}^{(a)}$ (Definition 10.51); when they exceed bounds, automatic triggers fire (Def. Definition 10.108, Cor. Corollary 10.91) and the Audit path proceeds per §10.10.

10.11.4 Stability Bounds: Mixing Norms vs. Visibility Changes

Definition 10.128 (Typed mixing norm)**.** For a typed channel $[\chi] \leftrightarrow [\chi']$ and triads $a, b \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$, let

$$\hat{\mathbf{G}}_\Lambda^{(a\to b)\,[\chi\to\chi']} := \mathsf{P}_{\mathrm{phys}}\,\Pi^{(b)}\,\Pi^{[\chi']}\,\hat{\mathbf{G}}_\Lambda\,\Pi^{[\chi]}\,\Pi^{(a)}\,\mathsf{P}_{\mathrm{phys}}$$

(typed insertion; Definition 10.113, (10.61)).

The *triad-resolved mixing norm* is

$$\mu_{[\chi\to\chi']}^{(a\to b)} := \left\|\hat{\mathbf{G}}_\Lambda^{(a\to b)\,[\chi\to\chi']}\right\|_{\mathcal{I}}, \tag{10.66}$$

where $\|\cdot\|_{\mathcal{I}}$ is the band-invariant operator norm on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ (realized weave configuration hygiene: Proposition 10.10). We also use the aggregated norms

$$\mu_{[\chi\to\chi']} := \Big(\sum_{a,b}\big(\mu_{[\chi\to\chi']}^{(a\to b)}\big)^2\Big)^{1/2}, \qquad \mu_{[\chi\to\chi']}^{(\mathrm{ns})} := \Big(\sum_{a\in\{\mathrm{vac},\mathrm{corr}\},\,b}\big(\mu_{[\chi\to\chi']}^{(a\to b)}\big)^2\Big)^{1/2}$$

for total and *non-spectral* (vac/corr-sourced) mixing, respectively.

Remark 10.129 (Well-posedness, safety, and trivial zeros)**.** All blocks in (10.66) are bounded on $\mathcal{I}^{[\mathscr{S}]}$ by Proposition 10.10; the definition is therefore well-posed. Typed/triad idempotents commute on the realized weave configuration (typed stability: Ch. 10 Definition 9.116, item (P7)). If a channel is symmetry-forbidden, the corresponding block vanishes (selection rules: Proposition 10.126). In particular, any block that would induce TT from a τ insertion is zero by TT–null preservation (Lemma 10.123), so $\mu_{[X\to\mathrm{TT}]}^{(a\to b)} = \mu_{[\mathrm{TT}\to X]}^{(a\to b)} = 0$ for $X \in \{S^{(b)}, S^{(L)}, \tilde{S}^{(p)}, A^{(\pm)}\}$.

Proposition 10.130 (Stability bounds: visibilities vs. mixing norms)**.** *Let* $\mathbf{L}^{[\mathscr{S};\chi]}$ *and* $\mathbf{V}^{[\mathscr{S};\chi]}$ *be the calibrated law/state vectors and* $\Lambda_{\mathrm{law/state}}^{[\mathscr{S};\chi]}$ *the corresponding visibilities of Definition 10.82 and eq. (10.54). Consider a perturbation of the typed correlator by a single channel block* $\Delta\hat{\mathbf{G}} = \hat{\mathbf{G}}_\Lambda^{(a\to b)\,[\chi\to\chi']}$ *with norm* $\mu := \mu_{[\chi\to\chi']}^{(a\to b)}$*. Then there exist realized weave configuration-dependent constants* $K_{j,(a\to b)}^{\mathrm{law}}$, $K_{m,(a\to b)}^{\mathrm{state}}$ *(independent of* μ*) such that*

$$\left|\Delta R_j^{[\mathscr{S};\chi]}\right| \le K_{j,(a\to b)}^{\mathrm{law}}\,\mu, \qquad \left|\Delta S_m^{[\mathscr{S};\chi]}\right| \le K_{m,(a\to b)}^{\mathrm{state}}\,\mu,$$

and, using the calibrations $f_j(x) = e^{-\alpha_j x}$ *and* $g_m(x) = \frac{x}{1+\beta_m x}$ *(Definition 10.82),*

$$\left\|\Delta\mathbf{L}^{[\mathscr{S};\chi]}\right\| \le \left(\sum_j \alpha_j^2 \left(K^{\text{law}}_{j,(a\to b)}\right)^2\right)^{1/2} \mu,$$
$$\left\|\Delta\mathbf{V}^{[\mathscr{S};\chi]}\right\| \le \left(\sum_m \left(K^{\text{state}}_{m,(a\to b)}\right)^2\right)^{1/2} \mu,$$

and hence the visibility changes satisfy a Lipschitz bound

$$\left|\Delta\Lambda^{[\mathscr{S};\chi]}_{\text{law}}\right| \le C_{\text{law},(a\to b)}\,\mu, \qquad \left|\Delta\Lambda^{[\mathscr{S};\chi]}_{\text{state}}\right| \le C_{\text{state},(a\to b)}\,\mu, \tag{10.67}$$

with constants $C_{\text{law/state},(a\to b)}$ *determined by the above bounds and the cosine aggregation in* (10.54).

Proof. Each law residual R_j (Definition 10.80, (10.51)) and state feature S_m ((10.52)) is a realized weave configuration-norm of a *linear* map (commutator with BRST charge, divergences, traces, typed spectral densities, and triad projections) applied to the typed observable/correlator on $\mathcal{H}_{\text{phys}}$. By submultiplicativity of $\|\cdot\|_{\mathcal{I}}$ and boundedness of projectors/derivatives on $\mathcal{I}^{[\mathscr{S}]}$ (Proposition 10.10), perturbing by a single channel block of norm μ shifts each R_j, S_m by at most a constant times μ, yielding the stated K's. For the calibrated entries, $|f_j'(x)| = \alpha_j e^{-\alpha_j x} \le \alpha_j$ and $|g_m'(x)| = \frac{1}{(1+\beta_m x)^2} \le 1$, so $|f_j(x+\delta) - f_j(x)| \le \alpha_j|\delta|$, $|g_m(x+\delta) - g_m(x)| \le |\delta|$. Summing in quadrature gives the $\|\Delta\mathbf{L}\|$ and $\|\Delta\mathbf{V}\|$ bounds. Finally, Λ is a smooth function of its vector argument via the cosine with $\mathbf{1}$ (Definition 10.82, (10.54)), so perturbations are Lipschitz with constants depending on the baseline norms of $\mathbf{L}, \mathbf{V}$ and $\mathbf{1}$; composing the previous bounds yields (10.67). □

Corollary 10.131 (Monotone effects on mixing norms under audit moves)**.** *Under the audit moves of Definition 10.101:*

(i) IR tightening
IR tightening weakly decreases non-spectral mixing $\mu^{(\text{ns})}_{[\chi\to\chi']}$ *(coherence/vacuum suppression; Lemma 10.102, item (1)).*

(ii) Arena switch (boundary → bulk)
A boundary to bulk arena switch weakly decreases coherence-dominated mixing (Lemma 10.102, item (2)).

(iii) Spectral window refinement
Spectral window refinement weakly decreases off-band mixing while preserving certified features (Lemma 10.102, item (3), KL positivity eq. (7.12)).

Combining with (10.67)*, these moves weakly improve the visibilities whenever mixing is the sole cause of their degradation.*

Proof. Recall the typed, triad-resolved insertion (Def. Definition 10.113, (10.61))

$$\hat{\mathbf{G}}^{(a\to b)\,[\chi\to\chi']}_{\Lambda} = \mathsf{P}_{\text{phys}}\,\Pi^{(b)}\,\Pi^{[\chi']}\,\hat{\mathbf{G}}_{\Lambda}\,\Pi^{[\chi]}\,\Pi^{(a)}\,\mathsf{P}_{\text{phys}},$$

and the mixing norm (Def. Definition 10.128, (10.66)) $\mu^{(a\to b)}_{[\chi\to\chi']} = \big\|\hat{\mathbf{G}}^{(a\to b)\,[\chi\to\chi']}_{\Lambda}\big\|_{\mathcal{I}}$. All operators act on the realized weave configuration $\mathcal{I}^{[\mathcal{S}]}$ and enjoy BRST/band safety and boundedness (Proposition 10.10). We argue item-wise.

(i) IR tightening. Let $r \mapsto r'$ denote a tightening step (Ch. 8 IR window/band functor; Definitions 6.280 and 7.29). On the realized weave configuration there exist channelwise contractions (Schur multipliers/filters) $\mathcal{T}^{(a)}_{r\to r'}$ with

$$\|\mathcal{T}^{(\mathrm{vac})}_{r\to r'}\|_{\mathcal{I}} \le 1, \quad \|\mathcal{T}^{(\mathrm{corr})}_{r\to r'}\|_{\mathcal{I}} \le 1, \quad \|\mathcal{T}^{(\mathrm{spec})}_{r\to r'}\|_{\mathcal{I}} = 1,$$

and, functorially, the tightened correlator block factors as

$$\hat{\mathbf{G}}^{(a\to b)}_{\Lambda}(r') = \mathsf{P}_{\mathrm{phys}}\,\Pi^{(b)}\,\mathcal{T}^{(b)}_{r\to r'}\,\Pi^{[\chi']}\,\hat{\mathbf{G}}_{\Lambda}(r)\,\Pi^{[\chi]}\,\mathcal{T}^{(a)}_{r\to r'}\,\Pi^{(a)}\,\mathsf{P}_{\mathrm{phys}},$$

with commutation on the realized weave configuration (typed stability; Ch. 10 Definition 9.116, item (P7)). By submultiplicativity of $\|\cdot\|_{\mathcal{I}}$,

$$\mu^{(a\to b)}_{[\chi\to\chi']}(r') \;\le\; \|\mathcal{T}^{(b)}_{r\to r'}\|_{\mathcal{I}}\;\mu^{(a\to b)}_{[\chi\to\chi']}(r)\;\|\mathcal{T}^{(a)}_{r\to r'}\|_{\mathcal{I}}.$$

Hence $\mu^{(a\to b)}$ is nonincreasing whenever a or b is non-spectral (vac or corr), which yields the stated decrease for the aggregated non-spectral norm $\mu^{(\mathrm{ns})}$. If the tightened window removes positive spectral measure carrying the non-spectral part (KL positivity eq. (7.12)), at least one of the contractions is *strict* and the corresponding block strictly decreases (cf. Lemma 10.102, item (1)).

(ii) Arena switch (boundary→bulk). Let $\mathfrak{P}$: bdry $\to$ bulk be the arena move (Definition 10.101). By the bulk/boundary comparison (Lemma 10.102, item (2)), the bulk machine implements a coherence-suppressing contraction $\mathcal{B}^{(\mathrm{corr})}$ (with $\|\mathcal{B}^{(\mathrm{corr})}\|_{\mathcal{I}} \le 1$) and leaves spectral content unchanged (isometry on the certified band). Thus for any coherence-dominated block (i.e. with $a = \mathrm{corr}$ or $b = \mathrm{corr}$),

$$\mu^{(a\to b)}_{[\chi\to\chi']}(\mathrm{bulk}) \;\le\; \|\mathcal{B}^{(b)}\|_{\mathcal{I}}\;\mu^{(a\to b)}_{[\chi\to\chi']}(\mathrm{bdry})\;\|\mathcal{B}^{(a)}\|_{\mathcal{I}} \;\le\; \mu^{(a\to b)}_{[\chi\to\chi']}(\mathrm{bdry}).$$

If coherence has positive measure on the region affected by the switch, the inequality is strict (ibid.).

(iii) Spectral window refinement. Let $W_{\mathrm{spec}} \mapsto W'_{\mathrm{spec}} \subset W_{\mathrm{spec}}$ (Def. 10.101). The refinement acts by pre/post multiplication with the orthogonal spectral projector $\Pi_{W'_{\mathrm{spec}}}$ inside (10.61). Since $\|\Pi_{W'_{\mathrm{spec}}}\|_{\mathcal{I}} \le 1$ and $\Pi_{W'_{\mathrm{spec}}}\Pi_{W_{\mathrm{spec}}} = \Pi_{W'_{\mathrm{spec}}}$, we get

$$\mu^{(a\to b)}_{[\chi\to\chi']}(W'_{\mathrm{spec}}) \;\le\; \mu^{(a\to b)}_{[\chi\to\chi']}(W_{\mathrm{spec}}).$$

If off-band leakage with positive spectral measure is excised, the decrease is strict (KL positivity eq. (7.12); Lemma 10.102, item (iii)).

Visibilities. When mixing is the sole cause of visibility degradation, the Lipschitz bounds (10.67) (Prop. 10.130) together with the inequalities above imply that each move weakly *improves* (or leaves unchanged) $\Lambda^{[\mathcal{S};\chi]}_{\mathrm{law/state}}$, with strict improvement when a strict contraction occurs on a set of positive spectral measure.

All steps are within $\mathcal{I}^{[\mathcal{S}]}$ and preserve BRST/gauge hygiene by Proposition 10.10. □

Proposition 10.132 (Resolvent/Davis–Kahan type bound for visibilities)**.** *Let $\Lambda_\bullet^{[\mathscr{S};\chi]}$ be any visibility for type $[\chi]$ built as in Definition 10.82 and eq. (10.54) on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ (Def. Definition 10.1). Assume the* unmixed *diagonal typed block $\hat{\mathbf{G}}_\Lambda^{[\chi\to\chi]}$ (Def. Definition 10.113, (10.61)) admits a spectral gap $\Delta > 0$ around its certified feature (Riesz contour contained in the gap). Then for sufficiently small mixing norm μ (e.g. the aggregated channel norm from Definition 10.128),*

$$\left|\Lambda_\bullet^{[\mathscr{S};\chi]} - \Lambda_\bullet^{[\mathscr{S};\chi]}\big|_{\text{no-mix}}\right| \;\le\; C_\bullet\,\frac{\mu}{\Delta} \;+\; O\!\left(\frac{\mu^2}{\Delta^2}\right), \tag{10.68}$$

with a rig–dependent constant $C_\bullet > 0$. Here $|_{\text{no-mix}}$ denotes the visibility with the off–diagonal typed channels $[\chi] \leftrightarrow [\chi']$ suppressed.

Proof. Work on $\mathcal{H}_{[\chi]\oplus[\chi']} := \mathcal{H}_{[\chi]} \oplus \mathcal{H}_{[\chi']}$ equipped with the orthogonal sum inner product (see §10.11.1). Write the typed, block–decomposed correlator (Def. Definition 10.113)

$$\mathcal{G} = \begin{pmatrix} \hat{\mathbf{G}}_\Lambda^{[\chi\to\chi]} & \hat{\mathbf{G}}_\Lambda^{[\chi\to\chi']} \\ \hat{\mathbf{G}}_\Lambda^{[\chi'\to\chi]} & \hat{\mathbf{G}}_\Lambda^{[\chi'\to\chi']} \end{pmatrix} \;=:\; \mathcal{G}_0 + \mathcal{V},$$

with the *unmixed* part $\mathcal{G}_0 := \operatorname{diag}\!\big(\hat{\mathbf{G}}_\Lambda^{[\chi\to\chi]}, \hat{\mathbf{G}}_\Lambda^{[\chi'\to\chi']}\big)$ and the off–diagonal mixing $\mathcal{V} := \left[\begin{smallmatrix} 0 & \hat{\mathbf{G}}_\Lambda^{[\chi\to\chi']} \\ \hat{\mathbf{G}}_\Lambda^{[\chi'\to\chi]} & 0 \end{smallmatrix}\right]$. Let $\mu := \|\mathcal{V}\|_{\mathcal{I}}$; by Definition 10.128 one may take $\mu \le c\,\mu_{[\chi\to\chi']}$ for some realized weave configuration constant $c > 0$ that aggregates the $(a\to b)$ blocks.

By the gap hypothesis, there is a contour Γ enclosing the certified feature of $\hat{\mathbf{G}}_\Lambda^{[\chi\to\chi]}$ and no other spectrum of $\mathcal{G}_0$, with $\operatorname{dist}(\Gamma, \sigma(\mathcal{G}_0 \setminus \text{feat})) = \Delta$. Let P_0 be the corresponding Riesz projector of $\mathcal{G}_0$ onto $\mathcal{H}_{[\chi]}$ (it equals the fixed typed/triad spectral idempotent used for the certified band; cf. Definition 10.2 and the Riesz usage in Lemma 10.103). Let P be the Riesz projector of $\mathcal{G}$ associated to the image of the same contour under the perturbed resolvent.

Kato's resolvent identity yields (on $\mathcal{I}^{[\mathscr{S}]}$, using Proposition 10.10 for boundedness/BRST safety)

$$P - P_0 = \frac{1}{2\pi i}\oint_\Gamma (\zeta - \mathcal{G})^{-1}\,\mathcal{V}\,(\zeta - \mathcal{G}_0)^{-1}\,d\zeta,$$

hence $\|P - P_0\|_{\mathcal{I}} \;\le\; \frac{1}{2\pi}\,|\Gamma|\,\sup_{\zeta\in\Gamma}\|(\zeta-\mathcal{G})^{-1}\|_{\mathcal{I}}\,\|\mathcal{V}\|_{\mathcal{I}}\,\|(\zeta-\mathcal{G}_0)^{-1}\|_{\mathcal{I}} = O\!\big(\mu/\Delta\big)$, since both resolvents are $O(1/\Delta)$ on Γ for μ small enough (standard Neumann–series control). Equivalently, by the Davis–Kahan $\sin\Theta$ theorem for invariant subspaces, $\|P - P_0\|_{\mathcal{I}} \;\le\; C\,\mu/\Delta$ for some numerical C depending only on the contour geometry.

The typed observables entering the meters are linear functionals of the projected correlator, built from P (or P_0) and the fixed typed/triad idempotents (Defs. Definitions 10.80 and 10.82); hence the underlying ledger entries (R_j, S_m) change by at most $K\,\|P - P_0\|_{\mathcal{I}}$ for rig–dependent $K > 0$ (bounded operator calculus on $\mathcal{I}^{[\mathscr{S}]}$, Proposition 10.10). Applying the calibrations $f_j(x) = e^{-\alpha_j x}$ and $g_m(x) = \frac{x}{1+\beta_m x}$ (Lipschitz on $\mathbb{R}_+$ with constants α_j and 1, respectively; Definition 10.82) and the cosine aggregation (Definition 10.82 and eq. (10.54)) gives—exactly as in Proposition 10.130—that the visibility perturbations are Lipschitz in the underlying projected objects. Composing these bounds yields

$$\left|\Lambda_\bullet^{[\mathscr{S};\chi]} - \Lambda_\bullet^{[\mathscr{S};\chi]}|_{\text{no-mix}}\right| \;\le\; C_\bullet\,\|P - P_0\|_{\mathcal{I}} \;\le\; C'_\bullet\,\frac{\mu}{\Delta},$$

for a rig–dependent $C'_\bullet > 0$. The $O(\mu^2/\Delta^2)$ remainder follows from the second–order term in the resolvent expansion (or from the refined Davis–Kahan estimates when the graph distance is small). All steps occur on $\mathcal{H}_{\text{phys}}$ and respect BRST/gauge and band safety by Proposition 10.10. □

Corollary 10.133 (Weak–mixing robustness of visibilities and onsets (gap–controlled))**.** *Fix a sector $\mathscr{S} \in \{\Phi\Phi, \Theta\Theta, \Phi\Theta\}$, an admitted type $[\chi]$, and a triad $a \in \{\text{spec}, \text{vac}, \text{corr}\}$. Let $\Delta > 0$ be the certified spectral gap of the diagonal typed block for $[\chi]$ on the band (Prop. Proposition 10.132). Assume that, at some configuration (or along an admissible audit path), the* unmixed *visibilities exceed thresholds with margins*

$$\eta_L := \Lambda_{\text{law}}^{[\mathscr{S};\chi]} - \theta_{\text{law}} > 0, \qquad \eta_S := \Lambda_{\text{state}}^{[\mathscr{S};\chi]} - \theta_{\text{state}} > 0,$$

and let $\mu_*^{(a\to a)}$ be a mixing–norm bound for all off–diagonal channels into/out of $[\chi]$ on triad a:

$$\mu_*^{(a\to a)} \;\geq\; \Big\| \sum_{\chi' \neq \chi} \hat{\mathbf{G}}_\Lambda^{(a\to a)\,[\chi\to\chi']} \Big\|_{\mathcal{I}} \quad \textit{(e.g. the sum of norms; Def. Definition 10.128).}$$

If, uniformly after *each admissible move (or equivalently as a pathwise supremum),*

$$\frac{\mu_*^{(a\to a)}}{\Delta} \;<\; \min\Big\{ \frac{\eta_L}{C_{\text{law}}},\; \frac{\eta_S}{C_{\text{state}}} \Big\}, \tag{10.69}$$

where $C_{\text{law}}, C_{\text{state}}$ are the rig–dependent constants from (10.68) (Prop. Proposition 10.132), then:

(i) The visibilities remain above thresholds along the entire admissible path: $\Lambda_{\text{law}}^{[\mathscr{S};\chi]} \geq \theta_{\text{law}}$, $\Lambda_{\text{state}}^{[\mathscr{S};\chi]} \geq \theta_{\text{state}}$.

(ii) The certified onset $r^\star[\chi]$ (Def. Definition 10.85) is unchanged by the mixing, and, in particular, is stable *under IR tightening, spectral refinement, and arena switches (Defs. Definition 10.101, Lemmas 10.102 and 10.137 and Corollary 10.131).*

Proof. By the Davis–Kahan/Kato visibility bound (Prop. 10.132, (10.68)), for any single channel $[\chi] \leftrightarrow [\chi']$ one has

$$\big|\Delta\Lambda_\bullet^{[\mathscr{S};\chi]}\big| \;\leq\; C_\bullet \frac{\mu_{[\chi\to\chi']}^{(a\to a)}}{\Delta} \;+\; O\Big((\mu/\Delta)^2\Big), \qquad \bullet \in \{\text{law}, \text{state}\}.$$

Summing over $\chi' \neq \chi$ and using the triangle inequality (or the definition of $\mu_*^{(a\to a)}$), we get

$$\big|\Delta\Lambda_\bullet^{[\mathscr{S};\chi]}\big| \;\leq\; C_\bullet \frac{\mu_*^{(a\to a)}}{\Delta} \;+\; O\Big((\mu_*/\Delta)^2\Big).$$

Condition (10.69) ensures that the first–order term is strictly below the respective margin $\eta_\bullet$, so the post–move visibilities remain above their thresholds. This proves item (1).

For item (2), apply the same bound at $r^\star[\chi]$ and invoke the onset–stability radius from Proposition 10.88: the gap-controlled visibility variation is strictly smaller than the margin

to threshold, so the infimum defining $r^\star[\chi]$ does not change. Along admissible audit paths, Δ is arena–invariant (Lemma 10.137) and $\mu_*^{(a\to a)}$ weakly decreases under IR tightening and spectral refinement, and decreases under boundary→bulk when coherence dominates (Lemma 10.102, Cor. 10.131). Thus (10.69) remains valid along the path, yielding pathwise stability of the onset. All steps are on the realized weave configuration and respect BRST/band hygiene (Prop. 10.10); TT–null preservation of τ–types (Lemma 10.123 and Corollary 10.125) guarantees no spurious TT creation in $\Phi\Phi$ when τ mixing is present. □

Corollary 10.134 (Robustness of onsets under weak mixing)**.** *Let $r^\star[\chi]$ be the certified onset for type $[\chi]$ (Def. Definition 10.85), with stability radius $\rho[\chi]$ from Proposition 10.88. Write the* margins at onset

$$\eta_L := \Lambda_{\mathrm{law}}^{[\mathscr{S};\chi]}(r^\star[\chi]) - \theta_{\mathrm{law}}, \qquad \eta_S := \Lambda_{\mathrm{state}}^{[\mathscr{S};\chi]}(r^\star[\chi]) - \theta_{\mathrm{state}},$$

both nonnegative, and let $\Delta > 0$ be the spectral gap for the certified feature of the diagonal typed block $\hat{\mathbf{G}}_\Lambda^{[\chi\to\chi]}$ (Riesz contour; Definition 10.113). If a mixing channel $[\chi] \leftrightarrow [\chi']$ has aggregated norm μ (Def. Definition 10.128) such that

$$\frac{\mu}{\Delta} \;<\; \min\Big\{\frac{\eta_L}{C_{\mathrm{law}}},\; \frac{\eta_S}{C_{\mathrm{state}}}\Big\}, \tag{10.70}$$

where $C_{\mathrm{law}}, C_{\mathrm{state}} > 0$ are the realized weave configuration-dependent constants from Proposition 10.132, then the certified onset is unchanged by the mixing: $r^\star[\chi]$ remains the onset after turning on $[\chi] \leftrightarrow [\chi']$. Equivalently, $\Lambda_{\mathrm{law/state}}^{[\mathscr{S};\chi]}(r^\star[\chi])$ stay above thresholds after mixing, and no earlier $r < r^\star[\chi]$ attains admission.

Proof. By the gap–perturbation bound (10.68) in Proposition 10.132,

$$\big|\Delta\Lambda_{\mathrm{law}}^{[\mathscr{S};\chi]}\big| \le C_{\mathrm{law}}\frac{\mu}{\Delta} + O\big((\mu/\Delta)^2\big), \qquad \big|\Delta\Lambda_{\mathrm{state}}^{[\mathscr{S};\chi]}\big| \le C_{\mathrm{state}}\frac{\mu}{\Delta} + O\big((\mu/\Delta)^2\big).$$

For μ/Δ small, the quadratic remainder is dominated by halving the linear margin; the sufficient condition (10.70) guarantees

$$\Lambda_{\mathrm{law}}^{[\mathscr{S};\chi]}(r^\star[\chi]) \ge \theta_{\mathrm{law}}, \quad \Lambda_{\mathrm{state}}^{[\mathscr{S};\chi]}(r^\star[\chi]) \ge \theta_{\mathrm{state}}$$

after mixing. Thus $r^\star[\chi] \in \mathcal{A}_{[\chi]}$ (Def. Definition 10.85) still holds with mixing on.

It remains to rule out an earlier onset $\tilde{r} < r^\star[\chi]$. By Proposition 10.88, for $r < r^\star[\chi]$ the unmet threshold(s) have one–sided slope bounds $m_{\mathrm{law}}, m_{\mathrm{state}} > 0$; hence the deficit at r is at least $\min\{m_{\mathrm{law}}, m_{\mathrm{state}}\}\,(r^\star[\chi] - r)$. By (10.68), mixing lifts visibilities by at most $C_\bullet(\mu/\Delta)$, which cannot close this deficit for any $r \le r^\star[\chi] - \rho[\chi]$ if (10.70) holds (use the definition of $\rho[\chi]$ in Proposition 10.88 and that $C_\bullet(\mu/\Delta) < \eta_\bullet$). Therefore no $r < r^\star[\chi]$ reaches both thresholds under mixing, so the onset remains $r^\star[\chi]$ by uniqueness (part (1) of Proposition 10.88). All steps occur on $\mathcal{I}^{[\mathscr{S}]}$ and respect BRST/band safety (Proposition 10.10). □

Summary of the weave analysis. Typed insertions (10.61) give channel- and triad-resolved correlators on the realized weave configuration. The $\bar{\Sigma}$–weighted generalized eigen framework (§10.11.1, (10.63), (10.64)) quantifies mixing. TT–null is preserved for any τ–involving channel (Lemma 10.123), while symmetry selection rules (Proposition 10.126) kill forbidden typed pairs. Mixing impacts the calibrated visibilities *Lipschitzly* in the mixing norm (Prop. 10.130) and, with a band gap, admits the Davis–Kahan/Kato control (10.68) (Prop. 10.132). Consequently, Parasymmetric TLO decisions (type, location, onset) are robust: weak, gapped inter-modal mixing cannot shift the certified onset (Cor. 10.134) and audit moves monotonically suppress adverse mixing channels (Corollary 10.131), ensuring stable, falsifiable outcomes within SynTh/Audit.

10.12 Triad-Resolved Reporting with Boundary vs. Bulk Provenance

We finalize Step (7) by specifying a formal *reporting schema* for typed, triad-resolved outcomes with location provenance (bulk vs. boundary), and by preparing a *Boundary–vs–Bulk Invariance* statement: under admissible retunes, a bulk spectral claim cannot be created by boundary coherence alone (proved in §10.12.3). All objects live on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ (Def. Definition 10.1); triads and typed insertions are as in Definitions 10.2 and 10.113.

10.12.1 Schema and Macros for Weave Reports

Definition 10.135 (Weave report tuple)**.** Fix a sector $\mathscr{S} \in \{\Phi\Phi, \Theta\Theta, \Phi\Theta\}$, an admitted type $[\chi]$ (e.g. TT, $S^{(b)}$, $S^{(L)}$, $\tilde{S}^{(p)}$, $A^{(\pm)}$, rad$\pm$; see Definitions 10.3 to 10.5), a triad $a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$ (Def. Definition 10.2), and a location $\mathrm{Loc} \in \{\mathrm{bulk}, \mathrm{boundary}\}$ (arena of Definition 10.101). The *report tuple* is

$$\mathfrak{R}^{[\mathscr{S};\chi]}_{(a,\mathrm{Loc})} := \Big(a,\ [\chi],\ \mathrm{Loc}\ ;\ \Lambda^{[\mathscr{S};\chi]}_{\mathrm{law}},\ \Lambda^{[\mathscr{S};\chi]}_{\mathrm{state}},\ r^{\star}[\chi]\Big),$$

where the calibrated visibilities $\Lambda^{[\mathscr{S};\chi]}_{\mathrm{law/state}}$ are defined in Definition 10.82 and eq. (10.54) using the ledgers of Definition 10.80 and Remark 10.81, and the certified onset $r^{\star}[\chi]$ is from Definition 10.85 (uniqueness/stability in Proposition 10.88). The witnesses/meters this summarizes are those built in Section 10.9.

Definition 10.136 (Certified weave claim)**.** Fix thresholds $(\theta_{\mathrm{law}}, \theta_{\mathrm{state}}) \in (0,1)^2$. A tuple $\mathfrak{R}^{[\mathscr{S};\chi]}_{(a,\mathrm{Loc})}$ is a *certified claim* if

$$\Lambda^{[\mathscr{S};\chi]}_{\mathrm{law}} \geq \theta_{\mathrm{law}}, \qquad \Lambda^{[\mathscr{S};\chi]}_{\mathrm{state}} \geq \theta_{\mathrm{state}},$$

and all applicable diagnostics are below their audit bounds:

$$\begin{aligned} \varepsilon &\leq \varepsilon_{\max} && \text{(graviton TT leak; Definition 10.32)},\\ |\Delta_P| &\leq \delta_P && \text{(CK parity; Definition 10.58)},\\ |\Delta^{(a)}_{\mathrm{CP}}| &\leq \delta_{\mathrm{CP}} && \text{(Higgs CP; Definition 10.51)}. \end{aligned}$$

with automatic trigger policy per Definition 10.108 and Corollary 10.91. When these hold, Law/State are co-tenable for $(\mathscr{S}, [\chi])$ at the reported triad/location (§10.10, Definition 10.99).

What the report tuple is summarizing. The tuple of Definition 10.135 is a compact, standardized payload that answers two questions at once: *where* in the weave a statement lives, and *how strong* that statement is. The "where'' fields $(a, [\chi], \mathrm{Loc})$ pin down the provenance triad $a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$ (Definition 10.2), the admitted type $[\chi]$ (e.g. TT, τ–scalar, Higgs–CP; Definitions 10.3 to 10.5), and the arena $\mathrm{Loc} \in \{\mathrm{bulk}, \mathrm{boundary}\}$ used in audit moves (Definition 10.101). The "how strong'' fields are the *calibrated visibilities* $\Lambda^{[\mathscr{S};\chi]}_{\mathrm{law/state}}$ computed from the Law/State ledgers via the cosine meters (Definitions 10.80 and 10.82 and eq. (10.54)), and the unique onset $r^\star[\chi]$ that marks the minimal tightening at which law and state become co-tenable (Definition 10.85, Proposition 10.88, and Lemma 10.86). Because these objects are evaluated on the realized weave configuration and physical carrier, they inherit band/BRST hygiene and order-of-operations robustness (Definition 10.1 and Propositions 10.9 and 10.10); finiteness is guaranteed by Lemma 10.83. In practice, the tuple is the row you place in a table or the point you plot when comparing TT, τ–scalar, or Higgs–CP channels across sectors $\mathscr{S} \in \{\Phi\Phi, \Theta\Theta, \Phi\Theta\}$ (witness construction in Section 10.9).

When a tuple becomes a certified claim. Definition 10.136 turns a report into a *decision*. Given thresholds $(\theta_{\mathrm{law}}, \theta_{\mathrm{state}})$, a tuple is certified when both visibilities clear their bars and all applicable diagnostics are inside their audit bounds: TT-leak ε for the graviton TT sector (Definition 10.32), vector parity asymmetry Δ_P for CK vectors (Definition 10.58), and scalar CP asymmetry $\Delta^{(a)}_{\mathrm{CP}}$ in each provenance a for Higgs-like scalars (Definition 10.51). The bound values are governed by the automatic-trigger policy (Definition 10.95 and Corollary 10.91). When these conditions hold, Law/State are *co-tenable* for $(\mathscr{S}, [\chi])$ at the specified triad/location (§10.10, Definition 10.99); if not, the automatic triggers and audit path of Definition 10.95 prescribe a monotone, terminating retune to either reach the unique onset $r^\star[\chi]$ or to produce a certified failure certificate. In short, the schema provides a minimal, reproducible interface for reporting, comparing, and certifying results across the weave.

10.12.2 Arena-Invariant Spectral Set and Typed Stability

Lemma 10.137 (Arena-invariant certified spectral set)**.** *Let the certified band and contour families $\{\gamma_i\}_{(2)}, \{\Gamma_j\}_{(0)}, \{\zeta_m\}_{(1)}$ be fixed by the band tags P_x (Def. Definition 10.1; Riesz selectors Definition 10.7). Under an arena switch* bulk $\leftrightarrow$ boundary *(Def. Definition 10.101), these contour families—and hence the associated Riesz projectors—remain admissible and unchanged. Consequently, the presence/absence of spectral poles/bands on the certified set is* arena-invariant.

Proof. **Step 1: Fixed selectors and realized weave configuration.** The band tags P_x determine (i) the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ and (ii) the Riesz contours $\{\gamma_i\}_{(2)}, \{\Gamma_j\}_{(0)}, \{\zeta_m\}_{(1)}$ that enclose the certified spectral sets for spin $2, 0, 1$ respectively (Def. Definitions 10.1 and 10.7). All operators below are evaluated on $\mathcal{I}^{[\mathscr{S}]}$ and inherit BRST/gauge safety and band admissibility (bounded resolvents on the selected contours) by Proposition 10.10.

Step 2: Arena switch as an admissible homotopy. An arena switch bdry $\to$ bulk is an admissible audit move (Def. Definition 10.101). Hence there exists a norm–continuous path of certified operators $\{\hat{\mathbf{G}}_\Lambda(t)\}_{t\in[0,1]}$ on $\mathcal{I}^{[\mathscr{S}]}$ with $\hat{\mathbf{G}}_\Lambda(0) = \hat{\mathbf{G}}_\Lambda^{(\mathrm{bdry})}$, $\hat{\mathbf{G}}_\Lambda(1) = \hat{\mathbf{G}}_\Lambda^{(\mathrm{bulk})}$, such that for every fixed selector contour Γ from $\{\gamma_i, \Gamma_j, \zeta_m\}$, the resolvent set condition holds uniformly:

$$\Gamma \subset \rho\big(\hat{\mathbf{G}}_\Lambda(t)\big) \quad \text{and} \quad \sup_{\zeta\in\Gamma,\, t\in[0,1]} \big\|(\zeta - \hat{\mathbf{G}}_\Lambda(t))^{-1}\big\|_{\mathcal{I}} < \infty.$$

This is exactly the "admissibility" preservation guaranteed by realized weave configuration hygiene (Proposition 10.10) and the bulk/boundary comparison that leaves the certified *spectral* content unchanged while only affecting coherence (cf. Lemma 10.102, item (ii)).

Step 3: Riesz projectors are homotopy-invariant. For any such contour Γ, define the (typed/triad-suppressed) Riesz projector

$$\mathsf{P}_\Gamma(t) \;:=\; \frac{1}{2\pi i}\oint_\Gamma (\zeta - \hat{\mathbf{G}}_\Lambda(t))^{-1}\, d\zeta.$$

By the uniform resolvent bound along the path, $t \mapsto \mathsf{P}_\Gamma(t)$ is norm–continuous; moreover, the spectral mapping theorem for contour integrals implies $\mathsf{P}_\Gamma(t)$ is *constant in* t as long as Γ lies in the resolvent set for all t (standard homotopy invariance of the Riesz projection). Therefore $\mathsf{P}_\Gamma^{(\mathrm{bulk})} = \mathsf{P}_\Gamma^{(\mathrm{bdry})}$.

Step 4: Typed/triad refinement and conclusion. Typed and triad idempotents commute on the realized weave configuration and are stable under admissible retunings (Ch. 10: Definition 9.116, item (P7); triads Definition 10.2). Hence the typed/triad–refined spectral projectors (obtained by sandwiching P_Γ with the typed/triad idempotents; cf. Definition 10.113) are likewise invariant under the arena switch. Consequently, the presence/absence of spectral poles/bands *on the certified set*—equivalently, whether the corresponding Riesz projector is zero or nonzero—does not depend on the arena. (As used elsewhere, audit moves may suppress vac/corr features or off–band leakage, but they cannot create a new certified–band pole by admissibility and KL positivity; see Lemma 10.102 and eq. (7.12).) □

Lemma 10.138 (Typed spectral vacuum)**.** *Fix a type* $[\chi]$ *and a location*

$$\mathrm{Loc} \in \{\mathrm{bulk}, \mathrm{boundary}\}.$$

Assume that for every admissible spectral window W_{spec} *one has*

$$S_1^{[\chi]}(W_{\mathrm{spec}}) = 0, \tag{10.71}$$

where S_1 *is the spectral feature of* (10.52) *(see Definition 10.80) and admissibility of* W_{spec} *is as in Definition 7.665. Then the typed spectral projector onto the spin-s certified set annihilates the observable:*

$$\mathsf{P}_{\mathrm{spec}}^{(s)}\, \Pi^{[\chi]}\, \mathcal{O} = 0. \tag{10.72}$$

Here $\mathsf{P}_{\mathrm{spec}}^{(s)}$ *is the Riesz projector determined by the fixed contour family (Riesz selectors; Definition 10.7), and* $\Pi^{[\chi]}$ *is the typed projector (Ch. 10: Definition 9.116; see also Definitions 10.3 to 10.5).*

Proof. By (10.52), the spectral feature is

$$S_1^{[\chi]}(W_{\text{spec}}) \;=\; \int W_{\text{spec}}(\omega,\mathbf{k})\;\rho_{(s)}^{[\chi]}(\omega,\mathbf{k})\,d\mu(\omega,\mathbf{k}),$$

where $\rho_{(s)}^{[\chi]} \geq 0$ is the (KL-positive) typed spectral density on the certified band (KL positivity eq. (7.12)), and $s \in \{2,1,0\}$ matches the carrier (TT/vector/scalar). By Ch. 8, the admissible windows form a separating cone dense in the unit ball of L^∞ over the certified spectral set (cf. Definition 7.665); hence if $S_1^{[\chi]}(W_{\text{spec}}) = 0$ for *all* admissible W_{spec}, then by density and $\rho_{(s)}^{[\chi]} \geq 0$ we must have $\rho_{(s)}^{[\chi]} = 0$ almost everywhere on the certified spectral set (standard "testing by windows"/duality argument).

Let $\mathsf{P}_{\text{spec}}^{(s)}$ be the Riesz projector specified by the fixed contour family (Definition 10.7) that encloses precisely that certified spectral set. Since $\rho_{(s)}^{[\chi]}$ vanishes a.e. on that set, the spectral measure of $\Pi^{[\chi]}\mathcal{O}$ there is zero, whence the Riesz integral yields $\mathsf{P}_{\text{spec}}^{(s)}\,\Pi^{[\chi]}\mathcal{O} = 0$ on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$. Typed/triad idempotents and the unified correlator are band-admissible and BRST/gauge-safe on the realized weave configuration (Proposition 10.10), so the argument is valid in either arena. (If desired, arena invariance of the contour/projector itself follows from Lemma 10.137.) □

Remark 10.139 (Typed stability under arena switches)**.** Typed and triad idempotents commute on the realized weave configuration and are stable under admissible retuning (Ch. 10: Definition 9.116, item (P7); triads Definition 10.2). Together with Lemma 10.137, this implies that typed spectral claims (when present) are carried identically across bulk ↔ boundary; only the vacuum/coherence content changes (cf. Lemma 10.102, item (ii)).

10.12.3 Boundary–vs–Bulk Invariance

Proposition 10.140 (Boundary–vs–Bulk Invariance for spectral claims)**.** *Fix an admitted type* $[\chi]$ *and let* $\mathscr{S} \in \{\Phi\Phi, \Theta\Theta, \Phi\Theta\}$. *Consider an arena switch* boundary → bulk *(or vice versa; Def. Definition 10.101). Suppose that in the* boundary *arena the spectral feature entry satisfies*

$$S_1^{[\chi]} \equiv 0 \quad \textit{for all admissible windows } W_{\text{spec}}$$

(as in Eq. (10.52), Def. Definition 10.80), and any conflict is attributed to vac/corr *features (no spectral support). Then, after any finite sequence of admissible audit moves (IR tightening, spectral refinement, arena switches; Def. Definition 10.101), one cannot obtain a* certified spectral claim in bulk, *i.e.*

$$\not\exists\; \mathfrak{R}_{(\text{spec,bulk})}^{[\mathscr{S};\chi]} \quad \textit{with} \quad \Lambda_{\text{state}}^{[\mathscr{S};\chi]} \geq \theta_{\text{state}}$$

(Def. Definitions 10.135 and 10.136), unless *a new spectral pole/band* outside *the certified set appears (disallowed by admissibility; Defs. Definitions 10.1 and 10.7).*

Proof. By Lemma 10.138, the hypothesis $S_1^{[\chi]} \equiv 0$ for all admissible W_{spec} implies that the typed spectral Riesz projector annihilates the observable on the certified band: $\mathsf{P}_{\text{spec}}^{(s)}\,\Pi^{[\chi]}\,\mathcal{O} = 0$. By Lemma 10.137, the certified contour families and their Riesz projectors are *arena-invariant*; hence the same spectral projector vanishes after switching to bulk.

IR tightening and spectral window refinement are admissible audit moves (Definition 10.101) that *cannot create* new spectral poles within the fixed certified set: they act as contractions that suppress vac/corr content and excise off-band leakage while preserving certified spectral locations (KL positivity eq. (7.12) and monotonicities Lemma 10.102). Therefore $S_1^{[\chi]}$ remains identically zero on the certified set throughout any finite audit sequence that respects admissibility.

From the meter construction (Definition 10.82 and eq. (10.54)) and the ledger definition (Definition 10.80), the spectral entry $S_1 = 0$ yields the calibrated component $g_1(S_1) = 0$, so the spec contribution to $\mathbf{V}^{[\mathscr{S};\chi]}$ vanishes. With thresholds $(\theta_{\text{law}}, \theta_{\text{state}}) \in (0,1)^2$ (Def. Definition 10.136), the state visibility for a *spectral* claim cannot cross θ_{state} when its spectral component is identically zero. Hence no tuple $\mathfrak{R}^{[\mathscr{S};\chi]}_{(\text{spec,bulk})}$ can be certified unless a new spectral pole/band appears *outside* the certified set (which is excluded by the admissibility of audit moves and the fixed selectors; Defs. Definitions 10.1 and 10.7).

All steps occur on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ and preserve BRST/gauge and band safety by Proposition 10.10. □

Corollary 10.141 (No creation of TT spectral claims from τ–coherence)**.** *Fix* $\mathscr{S} \in \{\Phi\Phi, \Theta\Theta, \Phi\Theta\}$ *and consider an arena switch* boundary → bulk *(Def. Definition 10.101). Assume in the boundary arena that the TT spectral feature is absent on the certified set,*

$$S_1^{[\mathrm{TT}]} \equiv 0 \quad \textit{for all admissible } W_{\text{spec}} \qquad (\textit{Eq. (10.52) in Definition 10.80; Ch. 8 Definition 7.665}),$$

and that any nonzero content is vac/corr *coherence attributable to τ–typed channels $X \in \{S^{(b)}, S^{(L)}, \tilde{S}^{(p)}, A^{(\pm)}\}$ (typed only; Definition 10.4). Then, after any finite sequence of admissible audit moves (IR tightening, spectral refinement, and arena switches; Definition 10.101), there is no certified TT spectral claim in bulk:*

$$\nexists\ \mathfrak{R}^{[\mathscr{S};\mathrm{TT}]}_{(\text{spec,bulk})} \quad \textit{with} \quad \Lambda^{[\mathscr{S};\mathrm{TT}]}_{\text{state}} \geq \theta_{\text{state}} \quad (\textit{Defs. Definitions 10.135 and 10.136}),$$

unless a new spectral pole/band appears outside *the certified set (disallowed by admissibility; Definitions 10.1 and 10.7).*

Proof. (1) *Typed spectral vacuum at the boundary.* From $S_1^{[\mathrm{TT}]} \equiv 0$ for all admissible W_{spec} and KL positivity (eq. (7.12)), Lemma 10.138 gives

$$\mathsf{P}^{(2)}_{\text{spec}}\, \Pi^{[\mathrm{TT}]}\, \mathcal{O} \;=\; 0 \quad \text{on the certified band,}$$

where $\mathsf{P}^{(2)}_{\text{spec}}$ is the (spin–2) Riesz projector selected by the fixed contours (Riesz selectors; Definition 10.7).

(2) *Arena invariance of the spectral set.* By Lemma 10.137, the certified contour families and their Riesz projectors are invariant under boundary ↔ bulk. Therefore the same projector vanishes in bulk: $\mathsf{P}^{(2)}_{\text{spec}}\, \Pi^{[\mathrm{TT}]}\, \mathcal{O} = 0$.

(3) *τ–mixing cannot populate TT spectral content.* For any τ–typed channel X, TT–null preservation (Lemma 10.123) gives the typed, triad–resolved identities

$$\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\, \hat{\mathbf{G}}^{(a\to b)\,[X\to\mathrm{TT}]}_{\Lambda} = 0, \qquad \hat{\mathbf{G}}^{(a\to b)\,[\mathrm{TT}\to X]}_{\Lambda}\, \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}} = 0 \quad \text{for all } a, b \in \{\text{spec}, \text{vac}, \text{corr}\}$$

(typed insertion: Definition 10.113, (10.61)). Thus off–diagonal $\tau \leftrightarrow$ TT mixing cannot generate TT spectral weight; the only possible contributor to a TT spectral claim is the diagonal TT block, already annihilated by Step (2).

(4) *Audit moves cannot create certified–band poles.* IR tightening and spectral window refinement act as contractions that suppress vac/corr and remove off–band leakage but do not create poles inside the fixed certified set (monotonicities: Lemma 10.102; KL positivity: eq. (7.12); realized weave configuration hygiene: Proposition 10.10). Therefore $S_1^{[\mathrm{TT}]}$ remains 0 on the certified band along any admissible finite audit path.

Combining (2)–(4), the TT spectral component of the calibrated state vector stays null in bulk, preventing the spectral triad visibility from crossing θ_{state}; hence no certified tuple $\mathfrak{R}^{[\mathscr{S};\mathrm{TT}]}_{(\mathrm{spec,bulk})}$ exists. The only exception would be the appearance of a new TT spectral pole outside the certified set, which is excluded by admissibility of the selectors (Definitions 10.1 and 10.7). This also recovers Prop. 10.140 as the special case $[\chi] = \mathrm{TT}$. □

10.12.4 Stability of Reported Onsets under Arena Changes and Mixing

Proposition 10.142 (Report stability under admissible moves)**.** *Let* $\mathfrak{R}^{[\mathscr{S};\chi]}_{(a,\mathrm{Loc})}$ *be a* certified *report tuple (Defs. Definitions 10.135 and 10.136) with sector* $\mathscr{S} \in \{\Phi\Phi, \Theta\Theta, \Phi\Theta\}$*, triad* $a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$*, and location* $\mathrm{Loc} \in \{\mathrm{bulk}, \mathrm{boundary}\}$*. Let the law/state margins be*

$$\eta_L := \Lambda^{[\mathscr{S};\chi]}_{\mathrm{law}} - \theta_{\mathrm{law}}, \qquad \eta_S := \Lambda^{[\mathscr{S};\chi]}_{\mathrm{state}} - \theta_{\mathrm{state}}, \quad \eta_{L,S} > 0,$$

and let $\Delta > 0$ *be the certified band gap for the diagonal typed block (Prop. Proposition 10.132). Assume a mixing channel* $[\chi] \leftrightarrow [\chi']$ *has aggregated triad-diagonal norm* $\mu^{(a\to a)}_{[\chi\to\chi']}$ *(Def. Definition 10.128). If*

$$\frac{\mu^{(a\to a)}_{[\chi\to\chi']}}{\Delta} \;<\; \min\Big\{\frac{\eta_L}{C_{\mathrm{law}}},\; \frac{\eta_S}{C_{\mathrm{state}}}\Big\}, \tag{10.73}$$

where $C_{\mathrm{law}}, C_{\mathrm{state}}$ *are the constants in the visibility bound (Prop. Proposition 10.132,* (10.68)*), then:*

IR tightening / spectral refinement
After any admissible IR tightening or spectral window refinement (Def. Definition 10.101), the updated report $\mathfrak{R}^{[\mathscr{S};\chi]}_{(a,\mathrm{Loc})}$ *remains certified.*

Arena switch
After an admissible arena switch $\mathrm{Loc} \to \mathrm{Loc}'$ *(Def. Definition 10.101), the updated report* $\mathfrak{R}^{[\mathscr{S};\chi]}_{(a,\mathrm{Loc}')}$ *remains certified.*

Proof. Let $\Delta\Lambda_{\mathrm{law/state}}$ denote the visibility change induced by a single admissible move together with the fixed mixing channel. By the gap–controlled Davis–Kahan/Kato bound (Prop. 10.132, (10.68)),

$$\big|\Delta\Lambda^{[\mathscr{S};\chi]}_{\mathrm{law}}\big| \;\le\; C_{\mathrm{law}}\frac{\mu^{(a\to a)}_{[\chi\to\chi']}}{\Delta} + O\big((\mu/\Delta)^2\big), \qquad \big|\Delta\Lambda^{[\mathscr{S};\chi]}_{\mathrm{state}}\big| \;\le\; C_{\mathrm{state}}\frac{\mu^{(a\to a)}_{[\chi\to\chi']}}{\Delta} + O\big((\mu/\Delta)^2\big),$$

where Δ is the certified band gap for the diagonal typed block and $\mu^{(a\to a)}_{[\chi\to\chi']}$ is the *post-move* (or pathwise supremum) triad-diagonal mixing norm (Def. 10.128). Assume the smallness condition (10.73). Then, at the updated configuration,

$$\Lambda^{[\mathscr{S};\chi]}_{\mathrm{law}} \geq \theta_{\mathrm{law}} \quad \text{and} \quad \Lambda^{[\mathscr{S};\chi]}_{\mathrm{state}} \geq \theta_{\mathrm{state}},$$

so the law/state inequalities remain satisfied.

(1) IR tightening / spectral refinement. By Lemma 10.102, IR tightening weakly increases Λ_{law} and the spectral component of Λ_{state}, while decreasing non-spectral (vac/corr) contributions; spectral window refinement removes off-band leakage while preserving certified features. Moreover, non-spectral mixing norms weakly decrease under these moves (Cor. 10.131), so the right-hand side of the Davis–Kahan bound does not worsen. Diagnostics $(\varepsilon, \Delta_P, \Delta_{\mathrm{CP}})$ remain below bounds by the same monotonicities and trigger soundness (Lemma 10.96 and Corollary 10.110). Hence the tuple $\mathfrak{R}^{[\mathscr{S};\chi]}_{(a,\mathrm{Loc})}$ remains certified (Defs. 10.135, 10.136).
(2) Arena switch. The certified spectral set and the gap Δ are arena-invariant (Lemma 10.137). The arena switch weakly decreases coherence-dominated mixing (Lemma 10.102, item (2); Cor. 10.131), so if $a \in \{\mathrm{vac}, \mathrm{corr}\}$ the Davis–Kahan bound improves. If $a = \mathrm{spec}$, we apply the same bound directly with the post-move $\mu^{(\mathrm{spec}\to\mathrm{spec})}$ (the inequality (10.73) is stated in terms of the post-move ratio, or uniformly along the admissible path). Therefore the threshold inequalities hold at (a, Loc'), with diagnostics still below bounds as above, and $\mathfrak{R}^{[\mathscr{S};\chi]}_{(a,\mathrm{Loc}')}$ remains certified.

Finally, the onset $r^\star[\chi]$ is robust under weak, gapped mixing (Cor. 10.134), so the reported onset remains valid throughout. All steps occur on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ and preserve BRST/gauge and band safety (Prop. 10.10). □

10.13 Norms, Contours, and TT-Hyperbolicity

What these norms are doing. Definition 10.143 fixes a *house topology* on the physical carrier $\mathcal{H}_{\mathrm{phys}} = \mathrm{Ran}\,\mathsf{P}_{\mathrm{phys}}$ (Definition 10.1) for all operator estimates that appear later (ledgers, meters, mixing, and contour bounds). Every norm first *sandwiches* an operator X with the certified band tag $P_x(\lambda_{\mathrm{UV}})$ and IR window $K^{(\ell)}_{\mathrm{IR}}$—both commuting contractions by Proposition 10.10—and then measures its size either in the uniform (operator) sense, $\|\cdot\|_{\mathcal{B}}$, or in a Schatten sense, $\|\cdot\|_{\mathfrak{S}_p}$. The combined norm (10.76) balances *worst–case* control ($\|\cdot\|^{(\infty)}_{\mathcal{I}}$) with *bulk/average* control ($\|\cdot\|^{(2)}_{\mathcal{I}}$), tuned by $\kappa > 0$. Allowing $p \in [2, \infty]$ and varying the IR index ℓ within the certified class yields the admissible family $\mathfrak{N}$, which is stable under the windows/masks and spectral selectors used throughout Chs. 7–8 (Definitions 7.665 and 10.7). The hygiene statement in Remark 10.144 ensures that these sandwiches preserve BRST/Nielsen safety while remaining band–covariant, so that anything we measure lives where it should (on $\mathcal{H}_{\mathrm{phys}}$ and within the accepted band).

Why they matter (and where they are used). These norms are the ambient rulers for (i) forming Law/State ledgers and calibrated visibilities (Definitions 10.80 and 10.82), guaranteeing finiteness (Lemma 10.83); (ii) controlling typed mixing blocks (Definition 10.128); and (iii) bounding resolvents along Riesz contours used for spectral selection (Definition 10.7). Because

the sandwiches commute with the realized weave configuration/triad machinery (Proposition 10.10 and Definition 10.2), comparisons across types $[\chi]$ (e.g. TT, τ–scalar, Higgs–CP; Definitions 10.3 to 10.5) and across provenances (spec, vac, corr) are *order–independent* and robust. In the wider "Norms, Contours, and TT–Hyperbolicity" program of §10.13, this band–invariant topology is what lets us (a) keep contour integrals well posed as analysis settings are tightened, and (b) certify TT–sector propagation after projection with $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ against the realized background (cf. Definition 10.3 and Σ–causal response in Remark 9.83). In short, the admissible norm family is the quiet infrastructure that makes meters stable, contours convergent, and TT statements meaningful under the certified band/BRST hygiene.

10.13.1 Admissible Norm Family

Definition 10.143 (Admissible band–invariant norm family $\|\cdot\|_{\mathcal{I}}$)**.** Work on $\mathcal{H}_{\mathrm{phys}} = \operatorname{Ran} \mathsf{P}_{\mathrm{phys}}$ (Def. Definition 10.1). For any certified band tags $P_x(\lambda_{\mathrm{UV}})$ and IR window $K_{\mathrm{IR}}^{(\ell)}$ (both contractions on $\mathcal{H}_{\mathrm{phys}}$, commuting with $\mathsf{P}_{\mathrm{phys}}$; cf. Proposition 10.10), define for an operator X on $\mathcal{H}_{\mathrm{phys}}$:

$$\|X\|_{\mathcal{I}}^{(\infty)} := \left\| K_{\mathrm{IR}}^{(\ell)} P_x X P_x K_{\mathrm{IR}}^{(\ell)} \right\|_{\mathcal{B}(\mathcal{H}_{\mathrm{phys}})}, \tag{10.74}$$

$$\|X\|_{\mathcal{I}}^{(2)} := \left\| K_{\mathrm{IR}}^{(\ell)} P_x X P_x K_{\mathrm{IR}}^{(\ell)} \right\|_{\mathfrak{S}_2(\mathcal{H}_{\mathrm{phys}})}. \tag{10.75}$$

Fix any $\kappa > 0$ and set

$$\|X\|_{\mathcal{I}} \;:=\; \|X\|_{\mathcal{I}}^{(\infty)} \;+\; \kappa\, \|X\|_{\mathcal{I}}^{(2)}. \tag{10.76}$$

The *admissible family* $\mathfrak{N}$ consists of all norms obtained from (10.76) by varying the IR index ℓ within the certified class and by replacing $\mathfrak{S}_2$ with any Schatten $\mathfrak{S}_p$ ($p \in [2, \infty]$) with uniform bounds along admissible retunes (Chs. 7–8; windows/masks Definition 7.665, selectors Definition 10.7).

Remark 10.144 (Realized weave configuration hygiene and band covariance)**.** By Proposition 10.10, P_x and $K_{\mathrm{IR}}^{(\ell)}$ are idempotent contractions on $\mathcal{H}_{\mathrm{phys}}$ and commute with the BRST projector $\mathsf{P}_{\mathrm{phys}}$, so the sandwiches $K_{\mathrm{IR}}^{(\ell)} P_x(\cdot) P_x K_{\mathrm{IR}}^{(\ell)}$ preserve BRST/Nielsen safety and band admissibility. These norms are the ones used for ledgers/meters (Definitions 10.80 and 10.82) and for mixing blocks (Definition 10.128).

Lemma 10.145 (Norm properties and admissible equivalence)**.** *For any fixed admissible $(P_x, K_{\mathrm{IR}}^{(\ell)})$ and $\kappa > 0$:*

1. *$\|\cdot\|_{\mathcal{I}}$ of (10.76) is a norm on $\mathcal{B}(\mathcal{H}_{\mathrm{phys}})$.*
2. *If $p \in [2, \infty]$, the replacement $\mathfrak{S}_2 \mapsto \mathfrak{S}_p$ defines an equivalent norm; likewise for any two admissible IR indices ℓ, ℓ' there exist $c_\pm \in (0, \infty)$ (independent of X) with*

$$c_-\, \|X\|_{\mathcal{I}}^{(\ell,p)} \;\le\; \|X\|_{\mathcal{I}}^{(\ell',p')} \;\le\; c_+\, \|X\|_{\mathcal{I}}^{(\ell,p)}.$$

3. *For any projector/idempotent Π that is band/BRST-admissible (typed or triad; Definitions 9.116 and 10.2), one has $\|\Pi X \Pi\|_{\mathcal{I}} \le \|X\|_{\mathcal{I}}$ (contraction under admissible sandwiches).*

Proof. (1) Positivity and homogeneity are immediate; the triangle inequality follows from the same property of $\|\cdot\|_{\mathcal{B}}$ and of Schatten norms after the fixed contraction sandwich by $K_{\mathrm{IR}}^{(\ell)} P_x$.

(2) For $p \geq 2$ one has $\|T\|_{\mathcal{B}} \leq \|T\|_{\mathfrak{S}_p}$ and $\|T\|_{\mathfrak{S}_{p'}} \leq C\,\|T\|_{\mathfrak{S}_p}$ on the band-limited range (finite or uniformly locally finite effective rank due to selectors/masks; Chs. 7–8, Definitions 7.665 and 10.7). The IR family $\{K_{\mathrm{IR}}^{(\ell)}\}$ is an admissible chain of contractions (monotone retunes, Definition 10.101 and Lemma 10.102); hence sandwiches with different ℓ's are uniformly comparable, giving the constants $c_{\pm}$.

(3) If Π is an admissible idempotent commuting with $P_x, K_{\mathrm{IR}}^{(\ell)}$ (typed/triad stability on the realized weave configuration; Ch. 10 Definition 9.116, item (P7)), then $KP\,\Pi X \Pi\,PK = (KP\Pi)\,X\,(\Pi PK)$, with $KP\Pi$ a contraction; submultiplicativity of $\|\cdot\|_{\mathcal{B}}$ and Schatten norms yields the claim. □

Remark 10.146 (Use downstream)**.** The equivalence in Lemma 10.145 ensures that (i) the meters/visibilities (Definition 10.82) and (ii) the mixing/perturbation bounds (Definition 10.128 and Propositions 10.130 and 10.132) are insensitive to the particular choice within $\mathfrak{N}$, up to fixed multiplicative constants absorbed into calibration weights.

Proposition 10.147 (Equivalence and stability in $\mathfrak{N}$)**.** *Let $\mathfrak{N}$ be the admissible family of band–invariant norms from Definition 10.143 on $\mathcal{H}_{\mathrm{phys}} = \mathrm{Ran}\,\mathsf{P}_{\mathrm{phys}}$ (Def. Definition 10.1). For any two choices $\|\cdot\|_{\mathcal{I}}, \|\cdot\|'_{\mathcal{I}} \in \mathfrak{N}$ there exist rig–dependent constants $0 < c_- \leq c_+ < \infty$ such that*

$$c_-\,\|X\|_{\mathcal{I}} \;\leq\; \|X\|'_{\mathcal{I}} \;\leq\; c_+\,\|X\|_{\mathcal{I}}$$

for all operators X formed by finite compositions of

$$\mathsf{P}_{\mathrm{phys}}, \quad \Pi^{(a)} \;\; (\textit{triads; } a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\},\ \textit{Definition 10.2}), \quad \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \;\; (\textit{Definition 10.3}),$$

the typed projectors (Definitions 10.4 and 10.5), and the Riesz resolvents and projectors selected by the certified contour families (Definition 10.7). Moreover, if $K_{\mathrm{IR}}^{(\ell')} \leq K_{\mathrm{IR}}^{(\ell)}$ in the operator order (IR tightening within the certified class), then for the corresponding norms $\|\cdot\|_{\mathcal{I}'}$ (built with ℓ') and $\|\cdot\|_{\mathcal{I}}$ (built with ℓ) one has the monotonicity

$$\|X\|_{\mathcal{I}'} \;\leq\; \|X\|_{\mathcal{I}} \qquad \textit{for all such } X.$$

Proof. **Set-up and bounded sandwiches.** Each norm in $\mathfrak{N}$ has the form (cf. Definition 10.143)

$$\|X\|_{\mathcal{I}}^{(\infty)} = \|KP\,X\,PK\|_{\mathcal{B}}, \qquad \|X\|_{\mathcal{I}}^{(p)} = \|KP\,X\,PK\|_{\mathfrak{S}_p}, \qquad \|X\|_{\mathcal{I}} = \|X\|_{\mathcal{I}}^{(\infty)} + \kappa\|X\|_{\mathcal{I}}^{(p)},$$

with $P := P_x(\lambda_{\mathrm{UV}})$ and $K := K_{\mathrm{IR}}^{(\ell)}$ band/IR tags (Def. Definition 10.1). By realized weave configuration hygiene (Proposition 10.10), P, K are idempotent contractions on $\mathcal{H}_{\mathrm{phys}}$ commuting with $\mathsf{P}_{\mathrm{phys}}$, so $X \mapsto KPXPK$ is a bounded compression both in the operator norm and in Schatten $\mathfrak{S}_p$.

Class of operators. The generators listed in the statement are all bounded on $\mathcal{H}_{\mathrm{phys}}$: - $\mathsf{P}_{\mathrm{phys}}$, $\Pi^{(a)}$, $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$, and the typed projectors are idempotent contractions stable on the realized weave configuration (triads Definition 10.2; TT and typed projectors Definitions 10.3 to 10.5; typed stability in Ch. 10 Definition 9.116, item (P7)). - Riesz resolvents/projectors along

the certified contours are uniformly bounded by construction (selectors; Definition 10.7) and remain admissible under retunes (Proposition 10.10).

Finite compositions of these remain bounded, and their band–sandwiched versions inherit Schatten class membership on account of the certified masks/windows (Chs. 7–8; Definitions 7.665 and 10.7).

Equivalence across $\mathfrak{N}$. Let $\|\cdot\|_{\mathcal{I}}$ and $\|\cdot\|'_{\mathcal{I}}$ be built from (P, K, p, κ) and (P', K', p', κ'), respectively. On the certified band, the effective ranks (or local spectral multiplicities) determined by the selectors are uniformly controlled (Chs. 7–8), and for $p, p' \in [2, \infty]$ one has the standard comparison of Schatten norms on a uniformly band-limited range:

$$\|Y\|_{\mathcal{B}} \;\le\; \|Y\|_{\mathfrak{S}_p} \;\le\; C_{p,p'}\,\|Y\|_{\mathfrak{S}_{p'}} \qquad \text{for all band-sandwiched } Y,$$

with $C_{p,p'} < \infty$ independent of Y. Since P, P', K, K' are contractions, there exist constants $c_1, c_2 \in (0, \infty)$ (depending only on the realized weave configuration choices) such that for all X in the stated class:

$$\|K'P'XP'K'\|_{\mathcal{B}} \;\le\; \|KPXPK\|_{\mathcal{B}}, \qquad \|K'P'XP'K'\|_{\mathfrak{S}_{p'}} \;\le\; c_1\,\|KPXPK\|_{\mathfrak{S}_p}.$$

Combining these and the linear combos that define $\|\cdot\|_{\mathcal{I}}, \|\cdot\|'_{\mathcal{I}}$ gives the two-sided inequality with constants $c_\pm$ depending only on (P, K, p, κ) and (P', K', p', κ'), not on X. This yields the claimed equivalence.

Monotonicity under IR tightening. If $K' \le K$ (operator order; admissible IR tightening, Definition 10.101 and Lemma 10.102), then $K' = K'K = KK'$, and with $C := K'P$ and $D := KP$ we have $C = K'D$ with $\|K'\| \le 1$. Thus, for every X,

$$\|K'PXPK'\|_{\mathcal{B}} = \|CXC\|_{\mathcal{B}} \le \|K'\|^2\,\|DXD\|_{\mathcal{B}} \le \|KPXPK\|_{\mathcal{B}}.$$

The same contraction argument applies to $\|\cdot\|_{\mathfrak{S}_p}$, since Schatten norms are two-sided ideal norms: $\|CXC\|_{\mathfrak{S}_p} \le \|C\|^2\,\|X\|_{\mathfrak{S}_p}$. Adding the components as in (10.76) (and its primed analogue) gives $\|X\|_{\mathcal{I}'} \le \|X\|_{\mathcal{I}}$.

Conclusion. We have established two-sided norm equivalence for all X in the stated operator class and the monotone stability under IR tightening. All steps are carried out on $\mathcal{I}^{[\mathscr{S}]}$ and preserve BRST/gauge and band admissibility by Proposition 10.10. □

Remark 10.148 (Downstream consequences)**.** By Proposition 10.147, the quantitative bounds for meters, mixing norms, and visibility stability are invariant up to fixed constants across $\mathfrak{N}$ (cf. Definitions 10.82 and 10.128 and Propositions 10.130 and 10.132). Moreover, IR tightening can only improve the non-spectral diagnostics and cannot worsen the norm-based estimates used in SynTh/Audit (Lemma 10.102 and Corollary 10.131).

10.13.2 Measurable Structure for Riesz Integrals

Bochner measurability and Riesz integrals. Bochner measurability is the standard notion of measurability for Banach–valued maps and underpins the construction of Bochner integrals, which generalize the Lebesgue integral to functions taking values in a Banach space rather than in $\mathbb{R}$ or $\mathbb{C}$ (see, e.g., [433, 253]). Let $(X, \|\cdot\|)$ be a Banach space (in our case $X = \mathcal{B}(\mathcal{H}_{\text{phys}})$ with the operator norm), and let $f : (\Omega, \mathcal{F}, \mu) \to X$ be a map from a measure space. The

map f is *Bochner–measurable* (or *strongly measurable*) if it is measurable with respect to the Borel σ–algebra on X induced by the norm topology; equivalently, by the Pettis theorem, f is almost everywhere the limit of X–valued simple functions [433]. Under mild integrability conditions, one then defines the *Bochner integral*

$$\int_\Omega f(\omega)\, d\mu(\omega) \ \in X,$$

which enjoys the usual linearity and dominated–convergence properties of the Lebesgue integral but now returns an element of X [433, 253, 55].

Within the measurable structure developed in Section 10.13.2, these general notions are applied to the resolvents of the band–limited, gauge–fixed kinetic operators $K \in \{K_{\Phi\Phi}, K^{(0)}_{\Theta\Theta}, K^{(1)}_{\tau}\}$. For fixed K and a Riesz contour γ enclosing a certified spectral island, the map

$$z \longmapsto (K - z)^{-1} \in \mathcal{B}(\mathcal{H}_{\text{phys}})$$

is norm–continuous on γ, thanks to the resolvent identity and the uniform spectral gap δ that separates the island from the rest of the spectrum [253, 487]. Norm–continuity implies Bochner measurability, so the operator–valued contour integral

$$P \ = \ \frac{1}{2\pi i} \oint_\gamma (K - z)^{-1}\, dz$$

is a well-defined Bochner integral in $\mathcal{B}(\mathcal{H}_{\text{phys}})$ [433, 253]. This is precisely the setting of Proposition 10.149: part (i) asserts that for every contour point z and every admissible IR retune r and arena $\text{Loc} \in \{\text{bulk}, \text{boundary}\}$, the mapping

$$(r, \text{Loc}) \ \longmapsto \ (K - z)^{-1} \ \in \ \mathcal{B}(\mathcal{H}_{\text{phys}})$$

is Bochner–measurable (indeed norm–continuous) along admissible paths on the realized weave configuration, and part (ii) then identifies the associated Riesz projectors as bounded operators obtained by such contour integrals, invariant under admissible contour deformations and commuting with the physical and typed projectors (Definitions 9.116 and 10.2).

A second, equally important role of Bochner measurability in Section 10.13.2 is to control how the Riesz projectors depend on the audit parameters. Along an admissible audit path $t \mapsto K(t)$ (retunings of r, arena switches, and related moves, as in Definitions 10.1 and 10.101), realized weave configuration hygiene guarantees norm–continuity of $K(t)$ on the certified class [487, 253]. The resolvent identity

$$(K(t) - z)^{-1} - (K(t_0) - z)^{-1} = (K(t) - z)^{-1}\Big(K(t_0) - K(t)\Big)(K(t_0) - z)^{-1},$$

together with a uniform bound on $\|(K(t) - z)^{-1}\|$ provided by the spectral gap, implies $\|(K(t) - z)^{-1} - (K(t_0) - z)^{-1}\| \to 0$ as $t \to t_0$. Thus $t \mapsto (K(t) - z)^{-1}$ is norm–continuous and therefore Bochner–measurable as a map into $\mathcal{B}(\mathcal{H}_{\text{phys}})$ [433]. By dominated convergence for Bochner integrals [433, 253], one may pass the limit under the Riesz integral and conclude that

$$P(t) \ := \ \frac{1}{2\pi i} \oint_\gamma (K(t) - z)^{-1}\, dz$$

varies continuously with t and is Bochner–measurable as a function of the realized weave configuration parameters. This provides the measurable/continuous structure on which Proposition 10.149 and the subsequent audit and onset analysis (Lemma 10.102, Proposition 10.87, and Theorem 10.107) are built, ensuring that SynTh/audit manipulations of the realized weave configuration do not introduce pathological jumps in the spectral decomposition of the kinetic operators and that the corresponding typed bands and visibilities evolve in a controlled, Lebesgue-style measurable framework [433, 253, 487].

Proposition 10.149 (Contour classes and Bochner measurability for Riesz projectors)**.** *Let $K_{\Phi\Phi}$ and $K^{(0)}_{\Theta\Theta}$ be the band–limited, gauge–fixed kinetic operators on the $\Phi\Phi$ (spin–2) and $\Theta\Theta$ (spin–0) blocks, and let $K^{(1)}_{\tau}$ denote the τ–*typed vector *(Chrono–Kymon) kinetic operator on the τ–vector typed subspace (cf. typed projectors in Definition 10.4 and instrumentation in Definition 10.1). Fix the certified band via P_x and choose piecewise C^1 Jordan contour families*

$$\{\gamma_i\}_{(2)} \ \text{ in } \ \rho(K_{\Phi\Phi}), \qquad \{\Gamma_j\}_{(0)} \ \text{ in } \ \rho(K^{(0)}_{\Theta\Theta}), \qquad \{\zeta_m\}_{(1)} \ \text{ in } \ \rho(K^{(1)}_{\tau}),$$

each enclosing disjoint spectral islands with a uniform gap $\delta > 0$ from the remainder of the spectrum across the certified class (Riesz selectors Definition 10.7; realized weave configuration hygiene Proposition 10.10). Then:

(i) Bochner measurability of resolvents

For every contour point z and every admissible IR retune r and arena $\mathrm{Loc} \in \{\mathrm{bulk}, \mathrm{boundary}\}$ *(audit moves Definition 10.101), the mappings*

$$(r, \mathrm{Loc}) \ \longmapsto \ (K - z)^{-1} \ \in \ \mathcal{B}(\mathcal{H}_{\mathrm{phys}})$$

are Bochner–measurable (indeed norm-continuous on admissible paths) for $K \in \{K_{\Phi\Phi}, K^{(0)}_{\Theta\Theta}, K^{(1)}_{\tau}\}$.

(ii) Riesz projectors: definition, boundedness, invariance, commutations

The Riesz projectors

$$\frac{1}{2\pi i}\oint_{\gamma_i} (K_{\Phi\Phi} - z)^{-1}\, dz, \quad \frac{1}{2\pi i}\oint_{\Gamma_j} (K^{(0)}_{\Theta\Theta} - z)^{-1}\, dz, \quad \frac{1}{2\pi i}\oint_{\zeta_m} (K^{(1)}_{\tau} - z)^{-1}\, dz$$

are well–defined bounded operators on $\mathcal{H}_{\mathrm{phys}}$, invariant under admissible contour deformations within the resolvent set, and they commute with $\mathsf{P}_{\mathrm{phys}}$, the triad maps $\Pi^{(a)}$ (Definition 10.2), and all typed projectors (Ch. 10: Definition 9.116, instantiated in Definitions 10.3 to 10.5).

Proof. **(1) Bochner measurability.** By admissibility of audit moves (Def. Definition 10.101) and realized weave configuration hygiene (Prop. Proposition 10.10), there is a norm–continuous path of band–limited, gauge–fixed kinetic operators $t \mapsto K(t)$ on the realized weave configuration $\mathcal{I}^{[\mathscr{J}]}$ interpolating any two admissible settings (r, Loc). Fix a contour point z; by the uniform gap δ one has $\|(K(t) - z)^{-1}\|_{\mathcal{B}} \le M$ for all t and some $M < \infty$ (selectors Definition 10.7). Using the resolvent identity,

$$(K(t) - z)^{-1} - (K(t_0) - z)^{-1} = (K(t) - z)^{-1}\Big(K(t_0) - K(t)\Big)\,(K(t_0) - z)^{-1},$$

so $\|(K(t)-z)^{-1}-(K(t_0)-z)^{-1}\| \le M^2\,\|K(t)-K(t_0)\| \to 0$ as $t \to t_0$. Thus $(r, \mathrm{Loc}) \mapsto (K-z)^{-1}$ is norm–continuous along admissible paths and therefore Bochner–measurable in $\mathcal{B}(\mathcal{H}_{\mathrm{phys}})$.

(2) Projector properties. Each Riesz projector is defined by the Cauchy integral over the corresponding contour; boundedness follows from the uniform resolvent bound and finite contour length (standard Riesz theory; the gap δ gives a uniform operator bound along the contour). Deformation invariance holds as long as the contour stays in the resolvent set, which is guaranteed by the selectors and realized weave configuration hygiene (Definition 10.7 and Proposition 10.10); the same argument shows arena invariance of the certified spectral set (cf. Lemma 10.137).

For commutations, note that on $\mathcal{I}^{[\mathscr{J}]}$ the projectors $\mathsf{P}_{\mathrm{phys}}$, $\Pi^{(a)}$, and all typed idempotents commute with the kinetic operators (typed stability and block structure; Definition 9.116, item (P7); see also Proposition 10.9). Hence they commute with every resolvent $(K-z)^{-1}$, and therefore with its contour integral. All statements are made on $\mathcal{H}_{\mathrm{phys}} = \operatorname{Ran}\mathsf{P}_{\mathrm{phys}}$ and preserve BRST/Nielsen safety and band admissibility (Prop. 10.10). □

Remark on the TT–Hyperbolic Indicator

The indicator $\mathbf{1}_{\{\sigma_{\mathrm{prin}}(K_{\Phi\Phi})\ \text{TT–hyperbolic}\}}$ used in Definition 10.27 is realized via the holomorphic functional calculus at the *symbol* level: restrict $K_{\Phi\Phi}$ to the TT typed subspace (projector $\Pi_{\Sigma}^{[\mathrm{TT}]}$, Definition 10.3), take the principal symbol σ_{prin} in the cotangent fiber ξ, choose a spectral cut that separates TT–hyperbolic eigenvalues from the rest, form the Riesz projector of the symbol–level resolvent over that cut, and quantize microlocally back to the operator on the certified band. This construction is compatible with band tags/selectors (Definition 10.7) and with realized weave configuration invariance (Prop. 10.10).

10.14 Findings and Quadrants: High-Level Synthesis

Instrumentation-first safety. All constructions are performed *on the realized weave configuration* $\mathcal{I}^{[\mathscr{J}]}$, so that every typed object (graviton, Higgs/radial kymon, τ-kairos scalars and Chrono–Kymon vectors) and every correlator inherits BRST/Nielsen flatness and band stability from the outset. Concretely, the BRST projector $\mathsf{P}_{\mathrm{phys}}$, the triad maps $\Pi^{(a)}$, and all typed idempotents commute and act on $\mathcal{H}_{\mathrm{phys}}$ without introducing gauge-parameter artifacts; see Propositions 10.9 and 10.10 and the standard BRST/Nielsen framework [445, 106, 120, 230]. This "instrumentation-first" posture ensures that subsequent claims are stated and tested under identical measurement conditions, avoiding spurious differences due to incompatible rigs.

τ-kairos purity and the Chrono–Kymon. The τ-kairos field decomposes into breathing and longitudinal scalars, a parity-odd pseudoscalar, and helicity-resolved vectors, and *each* typed component is TT-null by construction. As a result, projecting any τ-kairos content onto the background-covariant TT subspace annihilates it (Lemma 10.17), so there is no contamination of the graviton channel (Corollary 10.20). The vector content—the Chrono–Kymon—is extracted helicity by helicity, carries a parity audit Δ_P, and is selected spectrally by Riesz

projectors that automatically expand if new spin-1 features appear; cf. Definitions 10.57 and 10.58 and Lemma 10.62. The scalar/vector taxonomy follows the general GW-polarization classification beyond GR [477, 478, 479], while TT extraction is background-covariant in the York/Lichnerowicz sense [43].

Graviton (spin-2, TT). The graviton is obtained by a TT sandwich around the spin-2 Riesz selector, triad-resolved so that spectral, vacuum, and correlational TT pieces can be reported separately; see Definition 10.25. Independence from τ-mixing follows immediately from TT-nullity of the τ-types (Proposition 10.29 and Corollary 10.125). The TT-leak meter ε quantifies unexpected TT strength outside spectral support and is bounded by non-spectral TT components; it *decreases* under IR tightening (Lemma 10.103 and Corollary 10.104). If additional helicity-2 branches are realized physically, they are incorporated by adding contours to the spin-2 Riesz sum (Lemma 10.31), so the definition remains future-proof; the spectral selection and contour calculus follow standard holomorphic functional calculus [115, 253], and the radiative graviton degrees of freedom coincide with the usual massless spin-2 TT modes [7, 14, 43].

Higgs (radial kymon, CP-resolved). Within the tantrakyma scalar sector, the CP-refined radial projectors isolate amplitude (radial) directions in strict orthogonality to the NG/pNG manifold, ensuring no Goldstone leakage into the radial readout (Proposition 10.47). When CP is conserved the CP-even/odd channels decouple; otherwise their mixing is cleanly diagonalized by the spectral calculus (Proposition 10.49). Positivity of the spectral density on $\mathcal{H}_{\text{phys}}$ and the same contour-based future-proofing as above hold for the scalar case too (Propositions 10.48 and 10.50); compare with standard Higgs-sector CP diagnostics and positivity discussions [466, 467, 14, 255, 230].

Witnesses → visibilities → Parasymmetric TLO. Type-specific *law* residuals (Ward, trace, BRST/Nielsen) and *state* features (spectral weight, vacuum/coherence magnitudes, and diagnostics like $\varepsilon, \Delta_P, \Delta_{\text{CP}}$) are converted by calibrated meters into visibilities Λ_{law} and Λ_{state}. These meters are finite on the certified band (Lemma 10.83), monotone under IR retuning with the expected directions (Proposition 9.146), and yield unique onsets $r^\star[\chi]$ with explicit stability radii (Proposition 10.88). Consequently, the Parasymmetric TLO is well-posed, type by type, with controllable thresholds and robustness margins (Proposition 10.90); Ward/Slavnov–Taylor and BRST/Nielsen hygiene underpin the "law" side of these meters [483, 484, 445, 120].

Weaving (typed mixing) and robustness. Typed insertion into the Unified Inter-Modal Correlator furnishes a $\bar{\Sigma}$-weighted generalized eigenproblem that quantifies inter-type and inter-sector mixing on the certified band (Definitions 10.113 and 10.118). This problem is well-posed under our band assumptions (Lemma 10.122). Crucially, any channel touching the τ-block preserves TT-nullity (Lemma 10.123), so τ-mixing cannot generate spurious spin-2 TT content. Parity/helicity/CP symmetries gate which typed pairs can mix at all (Proposition 10.126). When mixing is permitted, visibility changes are perturbatively bounded in a Davis–Kahan sense and onsets remain stable under weak mixing (Proposition 10.132

and Corollary 10.134); spectral stability again relies on the standard holomorphic calculus [115, 253].

SynTh and finite audits. SynTh aggregates law/state shortfalls and diagnostic breaches into a single potential that *descends* under admissible moves (IR tightening, spectral refinement, arena switch). The No-Explosion/Audit Theorem guarantees that every conflict leads, in finitely many steps, either to co-tenability or to a stable certificate of non-co-tenability (Theorem 10.107). Automatic triggers for excessive TT-leak, parity, or CP asymmetry are sound and ensure that out-of-spec conditions are never silently accepted (Definition 10.95 and Corollary 10.110); see also the polarization/CP diagnostics context in [477, 478, 466, 467].

Reporting and provenance, boundary vs. bulk. We standardize reporting with a plain booktabs table listing sector, type, triad, location, visibilities, and onset. The certified spectral set and Riesz contours are arena-invariant, and tightening or switching arenas cannot *create* bulk spectral claims from boundary coherence alone (Proposition 10.140). In particular, τ-coherence at the boundary can never be promoted to a bulk TT graviton claim (Corollary 10.141). Admissible norm families and contour measurability are fixed once and for all (Definition 10.143 and Proposition 10.149), so that numerical meters and proofs share the same functional backbone [115, 253].

Quadrants: the SynTh Law $\oplus$ State map. Given thresholds $\theta_{\mathrm{law}}, \theta_{\mathrm{state}} \in (0,1)$, each typed, triad-resolved, and location-tagged outcome lands in one of four SynTh quadrants (Table 8.1). **QI** (both visibilities above threshold) is *co-tenable/certified* and should be reported with its onset $r^\star$ and diagnostics. **QII** (law OK, state weak) calls for IR tightening and spectral refinement; boundary coherence may be demoted and need not yield a spectral claim. **QIII** (state strong, law broken) mandates an immediate audit of gauges, meters, and off-band leakage before any claim is entertained. **QIV** (both weak) is the protected null zone: no claim is made, though exploratory retunes remain permissible. Across *all* quadrants, if any diagnostic exceeds its bound

$$\varepsilon > \varepsilon_{\max}, \qquad |\Delta_P| > \delta_P, \qquad |\Delta_{\mathrm{CP}}| > \delta_{\mathrm{CP}},$$

an audit is mandatory (Definition 10.95). Read in this way, the quadrants operationalize the theorems above: they translate instrumentation-first safety, typed mixing control, and audit guarantees into a clear decision map from raw meters to publishable, provenance-aware physics.

10.15 Permitted Spins, Polarizations, Helicities, and Tags: High-Level Overview

Scope of spins in this chapter. The realized weave configuration instantiated in this chapter admits spins $s \in \{2, 1, 0\}$. This is a *modeling choice*, not a prohibition: the operator-theoretic machinery (typed projectors, Riesz selectors, witnesses/TLO, SynTh) places no *a priori* ban on other spins (e.g. $s = \frac{3}{2}$ or higher); it simply was not activated here. Within the selected

scope, each spin is paired with its appropriate polarization/helicity structure, parity/CP tags when relevant, and the universal triad/location provenance.

Spins, helicities, and tags. This section declares, for the present realized weave configuration, which spins and polarization/helicity structures are *active* and how they are labeled for reporting and audit. Spin–2 lives in the $\Phi\Phi$ (geometry) block and is read exclusively through the background–covariant TT projector (Definition 10.3), so only the two massless helicities $\lambda = \pm 2$ appear, as required by the Poincaré/Wigner classification for a massless spin–2 representation (little group $E(2)$) [482, 14]; the operational extraction uses the York/Lichnerowicz TT machinery built into Definition 10.25 (cf. [43]). Non–TT content is *not* recorded here by design: if scalar or vector features are present, they are routed to the τ–kairos block and remain TT–null, so they cannot contaminate the graviton readout (Lemma 10.123 and Corollary 10.125). In parallel, the universal provenance tags—triad $a \in \{\text{spec}, \text{vac}, \text{corr}\}$ and location $\text{Loc} \in \{\text{bulk}, \text{boundary}\}$—travel with *every* type (Definitions 10.2 and 10.101), recording where the signal sits in the spectrum and arena without altering its spin/helicity meaning.

Where the other polarizations live (and why TT is safe). All non–TT gravitational content is organized in typed channels with symmetry tags that match standard GW–polarization taxonomies beyond GR [477, 478, 479]. The τ–kairos scalars comprise breathing and longitudinal even–parity types $S^{(b)}, S^{(L)}$ and an odd–parity pseudoscalar $\tilde{S}^{(p)}$; the matter vector (Chrono–Kymon) carries spin–1 with helicities $\lambda = \pm 1$ resolved by $A^{(\pm)}$ and a parity diagnostic Δ_P (Definitions 10.4, 10.57 and 10.58). Each of these is *TT–null* by construction (Lemma 10.123), so projecting onto the TT subspace annihilates them, preserving the purity of the spin–2 definition. In the $\Theta\Theta$ sector, the Higgs (radial kymon) is a spin–0 amplitude mode further refined by CP tags (rad$\pm$), with CP structure interpreted in the usual Higgs–sector sense [466, 467] and implemented here via Definition 10.5. Selection rules then gate mixing—parity–even does not mix with parity–odd, CP+ with CP−, and helicity sectors stay diagonal (Proposition 10.126)—and any sequence that touches the τ block still preserves TT–nullity (Lemma 10.123). Put together, the permitted spins/polarizations/helicities and their tags form a clean ledger: spin–2 TT for the graviton (Definitions 10.3 and 10.25), spin–1 vectors and spin–0 scalars in τ or Higgs channels, all triad/location–resolved and audited without ever blurring the TT graviton definition.

Graviton (geometry, $\Phi\Phi$). The graviton is pure spin-2 and *TT only*: the typed TT projector singles out helicities $\lambda = \pm 2$, with parity swapping $+2 \leftrightarrow -2$ (Definitions 10.25 and 10.26). No scalar or vector polarizations are admitted in this channel; any non-TT signatures that appear in practice are recorded in the τ-block (see below) and are identically TT-null, hence orthogonal to the graviton's TT readout (Proposition 10.29 and Corollary 10.125). Triad tags spec/vac/corr separate radiative spin-2 strength from vacuum structure and coherence, and bulk/boundary tags record provenance.

τ-kairos scalars and the Chrono–Kymon. The τ-kairos block captures non-TT structure in a typed, parity/helicity-aware way: two even-parity scalars—breathing $S^{(b)}$ and longitudinal

$S^{(L)}$—and one odd-parity pseudoscalar $\tilde{S}^{(p)}$ (Definition 10.4). All are *TT-null* by construction (Lemma 10.17). The matter vector channel—the Chrono–Kymon—carries spin-1 with helicities $\lambda = \pm 1$ resolved by $A^{(\pm)}$, parity swapping $+ \leftrightarrow -$ (Definitions 10.57 and 10.58 and Lemma 10.59). Each of these types is triad- and location-resolved just like the graviton, ensuring uniform reporting and audit.

Higgs (tantrakyma radial scalar). In the $\Theta\Theta$ sector, the Higgs (radial kymon) is a spin-0 amplitude mode equipped with CP tags rad± (even/odd), orthogonal to NG/pNG directions so that symmetry-breaking phases do not leak into the amplitude readout (Definitions 10.5 and 10.43 and Proposition 10.47). CP mixing—when present—is handled spectrally and reported without altering the typed structure (Proposition 10.49).

Triad and location are provenance, not polarizations. Every type, regardless of spin, carries the universal triad tag spec/vac/corr and the location tag bulk/boundary. These record *where* and *how* the signal arises (radiative vs. vacuum vs. coherence; bulk vs. boundary) and are orthogonal to spin/helicity or parity/CP. They are critical to SynTh's decision-making and to the *Boundary–vs–Bulk Invariance* guarantees established earlier.

Selection rules and TT safety under mixing. When the correlator respects a symmetry, selection rules block forbidden mixings: parity-even does not mix with parity-odd, helicity sectors are diagonal, and CP+ does not mix with CP− (Proposition 10.126). Moreover, any channel that involves the τ-block preserves TT-nullity, so no sequence of typed mixings can *create* TT graviton content from non-TT sources (Lemma 10.123). These two structural protections—selection rules and TT-null preservation—explain why "extra polarizations" observed operationally do not alter the spin-2 TT definition: they are registered and audited in the appropriate τ-types or scalar CP tags, not in the graviton projector.

Provenance of entries. The graviton's TT/helicity structure and its protection from τ-mixing come from Definitions 10.25, 10.26 and 10.32, Proposition 10.29, and Lemma 10.103. The τ-block's scalar/vector types, parity and helicity tags, and TT-nullity appear in Definitions 10.4, 10.57 and 10.58 and Lemmas 10.17 and 10.59. The Higgs CP split and NG/pNG orthogonality are in Definitions 10.5 and 10.43 and Propositions 10.47 and 10.49. Selection rules and TT-null preservation under mixing are gathered in Proposition 10.126 and Lemma 10.123, and universal triad/location tags are fixed by Definitions 10.1 and 10.2.

Scope and extensibility. The restriction to spins $s \in \{2, 1, 0\}$ in this chapter is a choice of scope, not a prohibition. Nothing in the typed–projector/Riesz–selector architecture, nor in the BRST/Nielsen safety layer of the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$, forbids $s = \frac{3}{2}$ or other spins. The framework is type–agnostic: once a carrier, its kinematic constraints, and the relevant symmetries are specified, the same three steps—(i) construct the typed projector(s), (ii) install the corresponding spectral selector(s) on the certified band, and (iii) insert them into witnesses, visibilities, TLO, SynTh, and reporting—apply unchanged.

Spin $3/2$ **(Rarita–Schwinger) channel.** For a spinor–vector carrier Ψ_μ, the pure-$3/2$ subspace is enforced by the divergence–free and γ-traceless constraints,

$$\Pi^{[\mathrm{RS}]} := \Pi^{[\nabla\Psi=0]}\,\Pi^{[\gamma\Psi=0]},$$

as in the Rarita–Schwinger (RS) framework [488, 14]. If the dynamics admits the RS gauge symmetry $\Psi_\mu \to \Psi_\mu + \nabla_\mu \epsilon$, the graded BRST structure of the realized weave configuration (same $\mathsf{P}_{\mathrm{phys}}$ with fermionic ghosts) applies verbatim [445, 106, 442, 489]. The spin-$3/2$ spectral selector is introduced exactly as for $s = 2, 1, 0$ using Riesz projectors on certified contours (Definition 10.7) [115, 253]):

$$\mathsf{P}^{(3/2)}_{\mathrm{spec}} = \sum_\ell \frac{1}{2\pi i} \oint_{\Upsilon_\ell} \left(K^{(3/2)} - z\right)^{-1} dz,$$

with certified contours $\{\Upsilon_\ell\}$. In the massless case one may helicity–resolve $\sum_{\lambda=\pm 3/2} \Pi^{(\lambda)} = \mathbf{1}_{\mathrm{RS}}$, with parity swapping $+ \leftrightarrow -$ [482, 14]. Triad and location tags (spec/vac/corr) $\times$ (bulk/boundary) attach verbatim (Definitions 10.2 and 10.101), maintaining uniform reporting and audits. The future–proof extractor mirrors the spin–2 graviton projector (Definition 10.25):

$$\mathcal{P}^{(3/2)}_{\mathrm{RS}} = \mathsf{P}_{\mathrm{phys}}\,\Pi^{[\mathrm{RS}]}\left(\mathsf{P}^{(3/2)}_{\mathrm{spec}} \oplus \Pi^{(\mathrm{vac})} \oplus \Pi^{(\mathrm{corr})}\right)\Pi^{[\mathrm{RS}]}\,\mathsf{P}_{\mathrm{phys}}.$$

Law–ledgers add RS–gauge closure and the γ-trace/divergence constraints; state–ledgers use the spin–$3/2$ spectral density $\rho_{(3/2)}$. Meter finiteness, monotonicity under IR tightening, onset uniqueness $r^\star[\frac{3}{2}]$, and stability–radius bounds carry over directly because they depend only on band–sandwiched norms and BRST covariance (Definition 10.143, Lemma 10.83, and Propositions 9.146, 10.10 and 10.88).[6]

Mixing and TT safety. Typed insertion into the unified correlator $\hat{\mathbf{G}}_\Lambda$ proceeds exactly as in the $s \in \{2, 1, 0\}$ cases (Definition 10.113); symmetry selection rules continue to determine admissible couplings (Proposition 10.126). The TT sandwich ensures that no spurious graviton is produced: the graviton extractor acts on symmetric 2–tensors within the TT subspace (Definitions 10.3 and 10.25), whereas Ψ_μ lives in a spinor–vector bundle. Any effect on the spin–2 channel must therefore appear as *genuine* $s = 2$ spectral support (i.e. new helicity–2 contours), which the Riesz–spectral formalism registers explicitly rather than via leakage (Definition 10.7 and Lemma 10.31).

Higher spins and anyonic sectors. The same template scales to higher integer spins and to fractional/anyonic content. For higher spins (e.g. symmetric Fronsdal fields with $s \geq 3$), impose the appropriate typed constraints (traceless/double–traceless, covariant transversality) and add $\mathsf{P}^{(s)}_{\mathrm{spec}}$ on certified contours; if massless, resolve helicities $\lambda = \pm s$; if massive, use polarization projectors indexed by $\pi \in \{1, \ldots, 2s+1\}$ [492, 493, 494, 495]. For anyons and fractional spin, work in 2+1 dimensions (or an effectively planar boundary arena) and replace helicity/parity tags by central idempotents $\Pi^{[a]}$ of a modular tensor category $\mathcal{C}$ (anyon charge

[6] Consistent interactions for spin–3/2 fields require the usual caveats (e.g. the Velo–Zwanziger problem and gauge-consistent couplings) [490, 491]; our use is kinematic/projector–level and agnostic to a specific interacting completion.

a, topological spin θ_a, braiding data); spectral selection can proceed via dispersive blocks $K^{[a]}$ or, when appropriate, via loop/line operators whose expectations enter the state–ledger—still band–sandwiched, so meters remain finite and monotone under IR tightening [496, 434, 497, 498, 499, 500]. Selection rules then follow braiding and charge–conjugation: mixing is allowed only when the fusion admits the vacuum (the tensor unit) [498, 499].

Conclusion. The weave is a reusable scaffold. To go beyond $\{2, 1, 0\}$—to $s = \frac{3}{2}$, to higher spins, or to anyonic sectors—no refactoring of prior constructions is required; one *extends the dictionary of types* by adding the relevant typed projectors and, where dispersive, the corresponding Riesz selectors on the same realized weave configuration. Core guarantees—BRST/Nielsen flatness, triad/location provenance, TT safety for the graviton, well–posed TLO with stable onsets, and finite, monotone meters—remain intact under these extensions (Definitions 10.1, 10.2, 10.82 and 10.95 and Lemma 10.86).

10.16 Fundamental Quanta and Their Excitations

Having distinguished the excitation channels of the τ–kairos (Section 10.17.3), the discussion turns to the quanta that populate those channels. The organization proceeds from the substrate fields to their typed, triad–resolved excitations: the *apeiron* of Gaiamiti (Φ) and the *kymon* of Tantrakyma (Θ), followed by the derived modes obtained through the constitutive split and the kinematic/typed projectors of the weave. In this ordering the higher excitations appear as structured outgrowths of the dual substrate (Φ, Θ) rather than as independent species fixed *a priori*.

Apeiron. The apeiron is the fundamental quantum register of the Gaiamiti field Φ (cf. Section 10.17.1). At the operator level it is the fluctuation $\delta\hat{\mathbf{\Phi}}$ on the Σ–realized carrier; at the spectral level relevant for the weave it materializes in the *scalar* typed family of the τ–kairos. When projected and certified within the typed, triad–resolved correlator, this scalar register yields the *Gaiaton* (Definitions 10.68 to 10.70). Through the constitutive dependence of the temporal tensor $t_{\mu\nu}[\Phi, \Theta]$ (eq. (10.101)), the apeiron also reappears in the temporal channel as the scalar contribution to $\hat{\tau}_{\mu\nu}$, consistently with TT–nullity and typed commutations (Definitions 10.2 and 10.4 and Lemma 10.71).

Kymon. The kymon is the generic quantum of the Tantrakyma field Θ (cf. Section 10.18.6). Its spectrum organizes matter, flavor, and internal–symmetry degrees of freedom in the weave's typed language. In the temporal channel, a dedicated vector register is singled out: the *Chrono–Kymon*, realized by the typed pair $A^{(\pm)}$ in the τ–kairos family (Definition 10.4) and mapped torsionally through the constitutive rules (cf. Section 10.7). This places the vector content of $\hat{\tau}_{\mu\nu}$ on the same realized weave configuration as the scalar content.

Graviton. From the Gaiamiti sector, the tangential (Goldstone) direction selected by the TT projector $\mathcal{P}^{(2)}$ defines the massless spin–2 graviton

$$\hat{h}_{\mu\nu} = \mathcal{P}^{(2)}[\delta\hat{g}_{\mu\nu}] \quad (Definition\ 10.167).$$

In the typed weave it saturates the TT content of the metric fluctuation and exhausts the radiative tensor degrees of freedom, consistent with the spin analysis (cf. Section 10.17.9) and with its identification as the spin–2 apeiron/Goldstone mode of Gaiamiti (eq. (10.28)). BRST/Nielsen safety and the triad commutations guarantee that no scalar or vector leakage contaminates the TT channel (Proposition 9.202 and Definition 10.2).

Gaiaton. When the apeiron is viewed in the metric sector through the τ–kairos *scalar* typed family $S^{(b)}, S^{(L)}, \tilde{S}^{(p)}$, it is termed the *Gaiaton*. Spectrally it is isolated by the spin–0 Riesz selector acting on the scalar kinetic generalized eigen-operator pair $K_\tau^{(0)}$ and realized by the CP– and triad–resolved projectors/fields of the weave (Definitions 10.68 to 10.70). TT orthogonality and typed safety ensure $\Pi_{\bar\Sigma}^{[\mathrm{TT}]}\,\mathcal{P}^{(0)}_{\mathrm{G},\eta} = 0$ for all admissible CP branches, so the tangent–radial decomposition of Gaiamiti fluctuations is cleanly implemented (Lemma 10.71).

Chrono–Kymon. Within the kymon family of Θ, the Chrono–Kymon provides the τ–kairos *vector* register via the typed pair $A^{(\pm)}$ (Definition 10.4). Its field strength $\hat{\mathcal{F}}^{(\tau)}_{\mu\nu} = 2\bar\nabla_{[\mu}\hat K^{(\tau)}_{\nu]}$ maps into the antisymmetric component of the temporal tensor (cf. Section 10.7), and its dynamics are captured by the background–covariant Maxwell/Proca–type equations on $\bar\Sigma$ developed for the Tantrakyma sector (cf. Section 10.18.6). This supplies the vector ledger and visibilities that accompany the scalar ledger of the Gaiaton (Definitions 10.82 and 10.117).

Tau–kairos. The temporal tensor fluctuation $\hat\tau_{\mu\nu}$—defined from the constitutive temporal map $t_{\mu\nu}[\Phi,\Theta]$ on the certified rig—obeys the TT–null rule and decomposes as

$$\boxed{\hat\tau_{\mu\nu} = \underbrace{\Big(\bar\nabla_{(\mu}\bar\nabla_{\nu)} - \tfrac{1}{d}\,\bar\Sigma_{\mu\nu}\,\bar\Box\Big)\hat\varphi_A}_{S_{\mu\nu}[\hat\varphi_A]} + \underbrace{2\,\bar\nabla_{[\mu}\hat K^{(\tau)}_{\nu]}}_{A_{\mu\nu}[\hat K^{(\tau)}]} \quad \text{with} \quad \Pi_{\bar\Sigma}^{[\mathrm{TT}]}\,\hat\tau_{\mu\nu} = 0 \;\; \text{(TT–null)}.}$$

where $\bar\Box := \bar\Sigma^{\rho\sigma}\bar\nabla_\rho\bar\nabla_\sigma$. This is equivalent to the τ-kairos typed split (10.10) once you resolve

$$S_{\mu\nu}[\hat\varphi_A] = S^{(b)}_{\mu\nu}[\hat\varphi_b] \oplus S^{(L)}_{\mu\nu}[\hat\varphi_L] \oplus \tilde S^{(p)}_{\mu\nu}[\tilde\varphi], \qquad A_{\mu\nu}[\hat K^{(\tau)}] = \sum_{\lambda=\pm} A^{(\lambda)}_{\mu\nu}[\hat K^{(\tau)}_\lambda],$$

using the τ–kairos type projectors from Definition 10.4. Here $S_{\mu\nu}[\hat\varphi_A]$ is the scalar (symmetric, trace–adjusted) piece descending from the Gaiaton channel, selected in practice by the τ–scalar type projectors $\Pi^{[\chi]}$ and the Riesz selector on $K_\tau^{(0)}$ (cf. Definitions 10.4, 10.68 and 10.69); $A_{\mu\nu}[\hat K^{(\tau)}] = \hat{\mathcal{F}}^{(\tau)}_{\mu\nu}$ is the vector (antisymmetric) piece sourced by the Chrono–Kymon, with $\hat{\mathcal{F}}^{(\tau)}_{\mu\nu} = 2\bar\nabla_{[\mu}\hat K^{(\tau)}_{\nu]}$ (cf. Section 10.7). Typed/triad commutations ensure compatibility with the weave meters and visibilities, keeping all diagnostics BRST/Nielsen–safe on certified plateaus (cf. Definitions 10.2 and 10.82 and Proposition 9.97). In particular, τ–kairos enters the pipeline only via typed projectors (not as an independent sector), and TT–purity $\Pi_{\bar\Sigma}^{[\mathrm{TT}]}\hat\tau_{\mu\nu} = 0$ guarantees that graviton observables remain unaffected by $\hat\tau_{\mu\nu}$ insertions (cf. Definition 10.4 and Proposition 10.29).

Mition and Tantron (spin–1 internal connection quanta). The spin–1 quanta termed the *mition* (Gaiamiti) and the *tantron* (Tantrakyma) originate from the *relational bundle connection* that appears in the covariant derivatives of Chapter 7 (cf. eqs. (10.1) and (10.2)). Their masses are generated via internal gauge SSB (BEH) as developed in Chapter 10 (cf. Definitions 9.252, 9.258, 9.261 and 9.348), but the excitations themselves are connection modes rather than substrate (Φ, Θ) quanta. On the realized weave configuration (Definition 10.1), the spin–1 content is isolated by a transverse, typed vector block and a spin–1 Riesz selector, exactly paralleling the spin–2 and spin–0 cases. Let $\Pi^{[\mathrm{T}]}$ denote the typed transverse projector for the internal vector bundle (commuting with the triad and provenance projectors), and let K_{vec} be the linearized connection kinetic operator on the realized weave configuration. Define the transverse block

$$K^{(1)}_{\mathrm{vec}} := \left(\mathsf{P}_{\mathrm{phys}}\Pi^{(\mathrm{spec})}\Pi^{[\mathrm{T}]}\right) K_{\mathrm{vec}} \left(\Pi^{[\mathrm{T}]}\Pi^{(\mathrm{spec})}\mathsf{P}_{\mathrm{phys}}\right),$$

and let $\mathfrak{S}_1$ be the set of its band–limited *spectral islands* with contours $\{\Upsilon_k\}$ (cf. Definition 10.6). The spin–1 selector is

$$\boxed{\mathsf{P}^{(1)}_{\mathrm{spec}} := \sum_{k\in\mathfrak{S}_1} \frac{1}{2\pi i} \oint_{\Upsilon_k} \left(K^{(1)}_{\mathrm{vec}} - z\right)^{-1} dz} \qquad \text{(cf. Definition 10.7).}$$

Projector–fields for the mition/tantron are obtained by sandwiching the connection operators with $\mathsf{P}^{(1)}_{\mathrm{spec}}$ and the sector–typed projectors. Longitudinal components are BRST–exact and decouple on $\mathcal{H}_{\mathrm{phys}}$ (Nielsen/BRST hygiene), so poles/residues read off from the transverse block are *gauge–parameter independent* on certified plateaus and *stable under admissible retunings* (cf. Propositions 10.9 and 10.10 and Corollary 10.11). TT–isolation is automatic since $\Pi^{[\mathrm{TT}]}_{\Sigma}$ acts on symmetric 2–tensors: $\Pi^{[\mathrm{TT}]}_{\Sigma}\,\mathsf{P}^{(1)}_{\mathrm{spec}} = 0$ (cf. Definition 10.3). As in the spin–2 and spin–0 cases, vector–sector *meters* and *onsets* may be defined from $\mathsf{P}^{(1)}_{\mathrm{spec}}$, inheriting calibration and robustness (cf. Definition 10.82 and Corollary 10.11).

Synthesis. The certified spectrum is thus balanced across the dual substrate. From Φ (Gaiamiti) emerge the massless spin–2 graviton and the massive spin–0 Gaiaton; from Θ (Tantrakyma) emerges the kymon family with the Chrono–Kymon providing the τ–kairos vector register. The τ–kairos binds these contributions into a single constitutive response—scalar content from the Gaiaton and vector content from the Chrono–Kymon—subject to TT–nullity and the typed commutation rules of the weave (Definitions 10.2 and 10.4 and Lemma 10.71). In this way the graviton, Gaiaton, Chrono–Kymon, and τ–kairos appear as interlocking manifestations of the same (Φ, Θ) architecture, each carrying distinct spin and dynamical roles while remaining inseparable from the constitutive chronogeometry that unifies them.

Predicted Regimes of Operation for the τ–Kairos

Framework stance (precision). The temporal/coherence fluctuation $\hat{\tau}_{\mu\nu} := \delta\hat{t}_{\mu\nu}$ is TT–null, $\mathcal{P}^{(2)}[\hat{\tau}_{\mu\nu}] = 0$ (eq. (10.90) and Sections 10.3 and 10.4), and decomposes constitutively as

$$\hat{\tau}_{\mu\nu} = S_{\mu\nu}[\hat{\varphi}_A] \;\oplus\; A_{\mu\nu}[\hat{K}^{(\tau)}],$$

with the Chrono–Kymon field strength $\hat{\mathcal{F}}^{(\tau)}_{\mu\nu} = 2\bar{\nabla}_{[\mu}\hat{K}^{(\tau)}_{\nu]}$ encoding antisymmetric temporal curvature (Definitions 10.4, 10.13 and 10.57 and Section 10.7). Parasymmetric execution $\mathscr{E}_\bullet$ stamps each admitted type $\chi \in \{\tau_S, \tau_A, K^{(\tau)}_{\mathrm{T}}, K^{(\tau)}_{\mathrm{L}}\}$ with a T–L–O triple $\mathrm{TLO}^{[\chi]} = (\mathrm{Type}, \mathrm{Location}, \mathrm{Onset})$ (Definitions 9.153 and 10.94 and Proposition 10.35); the onset $(r^\star, t^\star)$ is read from the meters $\upsilon^{(a)}_{\mathrm{Synthesis}}$, $\upsilon^{(LS)}_{\mathrm{Synthesis}}$, $\upsilon^{(LS)}_{\mathrm{Agreement}}$ (Definition 9.153). Crucially, *no additional spin–2 radiation* is predicted ($\mathcal{P}^{(2)}[\hat{\tau}] = 0$); the graviton remains the unique TT carrier (Section 10.17.8).

1. Microscopic Quantum Systems

Narrative. In laboratory settings that sustain quantum coherence, the τ–kairos operates as a phase–coherence modulator rather than as a classical force. Observable signatures include orientation– and history–dependent shifts in atom–interferometer phases (cf. Berry [24] and Pancharatnam [23]) and a geometry–tied dephasing channel for qubits, distinguishable from standard noise by its dependence on local temporal curvature.

T–L–O view (precision). *Type:* vector (τ_A, $K^{(\tau)}_{\mathrm{T}/L}$) for torsional/antisymmetric effects; scalar (τ_S) for trace/coarse–graining shifts (Definitions 10.4 and 10.13). *Location:* laboratory arena / instrument tags on the realized weave configuration. *Onset:* $(r^\star, t^\star)$ when the parasymmetric meters cross threshold (Definition 9.153). *Constraint:* TT purity remains intact ($\mathcal{W}_{\mathrm{TT\text{–}purity}} \approx 0$; Lemma 10.17); no extra GW polarizations (Section 10.17.8).

2. Early–Universe Cosmology

Narrative. In ultra–early (relational) epochs (table 9.3), temporal phase–flow can rival spatial curvature. The τ–kairos may seed *non–tensorial* imprints in primordial observables—e.g. scalar/vector contributions to CMB polarization anisotropies [439]—and feed characteristic structure into the stochastic GW background through non–TT couplings [501].

T–L–O view (precision). *Type:* τ_S (scalar) and $\tau_A/K^{(\tau)}$ (vector), admitted via Definition 10.94. *Location:* cosmological cone sheets determined by $\bar{\Sigma}$. *Onset:* early–epoch meter crossings yield wide coherence windows. *Constraint:* TT selection $\mathcal{P}^{(2)} \circ D_\Theta \Sigma = 0$ is preserved (eq. (10.101)); the graviton sector remains purely tensorial.

3. Near Black–Hole Horizons

Narrative. Near horizons, strong curvature and state–dependent coherence amplify τ–kairos sensitivity. Phase/entanglement structures of Hawking quanta can be modulated [87], informing information–paradox diagnostics [231, 138, 337]. Conversion proceeds through non–TT channels and through off–diagonal blocks of the weave's matrix propagator.

T–L–O view (precision). *Type:* primarily vector (τ_A, $K^{(\tau)}_{\mathrm{T}}$) via $\hat{\mathcal{F}}^{(\tau)}$; scalar τ_S if background profiles vary. *Location:* horizon–proximate sheets/tags on $\mathfrak{P}$ (and $\mathfrak{P}_\partial$ if used). *Onset:* meter–driven windows aligned with local redshift/time–dilation scales. *Mechanism:* mixing in off–diagonal correlators of the graviton–temporal matrix (Definition 10.182 and eq. (10.112)); TT purity guarded by $\mathcal{W}_{\mathrm{TT\text{–}purity}}$ (Lemma 10.17).

4. The Classical Macroscopic Limit

Narrative. Environmental decoherence averages temporal phase–flow to negligible levels [170, 282]. The effective radiative theory reduces to the purely tensorial TT sector carried by the graviton, reproducing GR phenomenology [7, 146].

T–L–O view (precision). *Type:* τ_S, τ_A fail admission (meters sub–threshold) or receive late/empty Onset windows. *Location:* macroscopic arenas with coarse tags. *Outcome:* $\mathrm{TLO}^{[\gamma]}$ is the sole radiative stamp; no reclassification errors (cf. Definition 10.26).

Synthesis. The domains above are precisely where the parasymmetric pipeline predicts enhanced τ–kairos visibility: where coherence persists and meters furnish early/strong onsets. These predictions respect TT isolation (eqs. (10.90) and (10.101)), typed admission (Definition 10.94), and the matrix–propagator architecture (Definition 10.182), *without* introducing extra GW polarizations and while remaining Nielsen/BRST–safe by design.

Proposition 10.150 (Predicted operational regimes of the τ–kairos)**.** *Let $\hat{\tau}_{\mu\nu} := \delta\hat{t}_{\mu\nu}$ denote the TT–null temporal/coherence fluctuation of the chronogeometric tensor (cf. eqs. (10.90), (10.100) and (10.101)), with scalar and vector components $S_{\mu\nu}[\hat{\varphi}_A]$ and $A_{\mu\nu}[\hat{K}^{(\tau)}]$ inherited respectively from the Gaiamiti and Tantrakyma substrates (see Section 10.16 and Definitions 10.4 and 10.13). Then:*

(i) Quantum–coherent dominance

τ–kairos couples constitutively to temporal phase flow rather than directly to classical stress–energy. Its effects therefore dominate only in regimes of sustained quantum coherence, where phase correlations cannot be coarse–grained away. T–L–O view: *$\chi \in \{\tau_S, \tau_A, K_{\mathrm{T}}^{(\tau)}, K_{\mathrm{L}}^{(\tau)}\}$ are admitted iff the hyperbolicity/cone and BRST conditions hold (Definition 10.94); onset $(r^\star, t^\star)$ is extracted from the parasymmetric meters $v^{(a)}_{\mathrm{Synthesis}}, v^{(LS)}_{\mathrm{Synthesis}}, v^{(LS)}_{\mathrm{Agreement}}$ (Definition 9.153), defining the coherence window in which $\hat{\tau}$ becomes operational.*

(ii) Microscopic quantum systems

In laboratory experiments sustaining coherence (e.g. atomic interferometry, entangled qubits), τ–kairos contributes orientation–dependent phase shifts and decoherence channels absent in the purely TT sector; these can appear as effective Berry/Pancharatnam–like phases of quantum–geometric origin [24, 23] (cf. Sections 10.3 and 10.4). T–L–O view: Type $=$ *τ_A (vector/antisymmetric) or τ_S (scalar/trace);* Location *from instrument/arena tags;* Onset *when meter thresholds are crossed (Definition 9.153).*

(iii) Early–universe cosmology

During ultra–early relational epochs (table 9.3), τ–kairos dynamics may rival curvature dynamics, leaving non–tensorial signatures in CMB polarization and contributing to the stochastic GW background [439, 501]. T–L–O view: Type $\in \{\tau_S, \tau_A\}$, Location *on cosmological cone sheets, with wide* Onset *windows when meters are large (parasymmetric synthesis).*

(iv) Near black–hole horizons

In the vicinity of event horizons, τ–kairos can modulate the coherence/entanglement structure of Hawking quanta, impacting information–paradox diagnostics [87, 231, 138, 337]. T–L–O view: *predominantly* Type $=$ *τ_A (via $K^{(\tau)}$ and its field strength),* Location

on horizon–proximate sheets/tags, with Onset *tied to local redshift/time–dilation scales (Definition 9.153).*

(v) Classical macroscopic limit

In decohered macroscopic regimes, environmental averaging suppresses τ–kairos contributions [170, 282], leaving the TT graviton sector $\hat{\gamma}_{\mu\nu} = \mathcal{P}^{(2)}[\delta\hat{g}_{\mu\nu}]$ as the effective radiative carrier (cf. Section 10.17.8), consistent with GR phenomenology [7, 146]. T–L–O view: *τ-types fail admission or receive late/empty onset windows; only* $\mathrm{TLO}^{[\gamma]}$ *is radiative (Type from QI–QIV, Location from diagnostic realized reference configuration metadata, Onset from meters; cf. Definition 10.26).*

Proof sketch. (1) follows from TT isolation, $\mathcal{P}^{(2)}[\hat{\tau}] = 0$ and $\mathcal{P}^{(2)} \circ D_\Theta \Sigma = 0$ (eqs. (10.90) and (10.101)), which forbids direct spin–2/stress–energy coupling and confines $\hat{\tau}$ to scalar/vector coherence. (2)–(4) apply this structural constraint in regimes where the parasymmetric meters certify early/strong onset and the typed admission gate is passed (Definitions 9.153 and 10.94); signatures therefore appear as phase/entanglement modulations in the non–TT channel, not as extra GW polarizations. (5) uses standard decoherence: meter response drops below threshold, typed admission fails or yields trivial onset, and only the TT graviton sector remains dynamically radiative (Section 10.17.8). Law–side TT purity is monitored by $\mathcal{W}_{\text{TT–purity}}$ (Lemma 10.17), ensuring no hidden TT leakage. □

The Gaiamiti Field as the Architect of Spacetime

The *Gaiamiti field* $\hat{\mathbf{\Phi}}(x)$ is the exclusive source of purely geometric excitations in the IQR framework. Its potential $\mathcal{V}_\Phi$ and vacuum manifold $\mathcal{M}_{\text{vac}}$ determine the emergent chronogeometric background $\bar{\Sigma}_{\mu\nu}$ and the spectrum of geometric quanta (Definition 5.54 and Section 10.17.6). When the symmetries of $\mathcal{V}_\Phi$ are spontaneously broken, two canonical fluctuation types follow from the standard SSB logic on curved backgrounds [244, 7]:

Massless spin–2 graviton ($\hat{h}_{\mu\nu}$) Long–range geometry must be mediated by a massless spin–2 carrier sourced by a conserved rank–2 current; in our formalism this is the TT-projected metric fluctuation,

$$\hat{h}_{\mu\nu} \;:=\; \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\, \delta\hat{g}_{\mu\nu},$$

with propagation captured by the TT graviton two–point function G^{TT} (eqs. (10.28) and (10.131) and Corollary 10.20). Within the Gaiamiti sector, the graviton's *masslessness* is not postulated but follows from the Goldstone mechanism for broken geometric vacuum symmetries (Section 10.17.9 and Proposition 11.29; see also [415, 502]).

Gaiaton (massive spin–0 apeiron) Fluctuations of the Gaiamiti field Φ orthogonal to the vacuum manifold $\mathcal{M}_{\text{vac}}$ of the potential $\mathcal{V}_\Phi$ (the trace/radial directions in field space) produce the scalar apeiron mode, canonically identified as the *Gaiaton* (Definitions 6.61 and 10.70). On the realized weave configuration it is realized in the τ–kairos *scalar* typed family $S^{(b)}, S^{(L)}, \tilde{S}^{(p)}$ (Definition 10.4) and is isolated spectrally by the spin–0 Riesz selector acting on the τ–scalar kinetic generalized eigen-operator pair $K^{(0)}_\tau$, yielding the mass and projector of the channel (Definition 10.68). CP– and triad–resolved projectors/fields are provided by Definition 10.69, and TT–orthogonality is enforced by

Lemma 10.71, so no TT contamination occurs. The realized mass is governed by the curvature (Hessian) of $\mathcal{V}_\Phi$ at $\mathcal{M}_{\rm vac}$ in exact analogy with standard SSB ([244]), while any allowed scalar mixing proceeds only through typed, triad–compatible couplings in the weave (Section 10.19.6, Definition 10.2, and Proposition 9.202).

Absence of spin–1 Gaiamiti excitations The typed menu for the Gaiamiti ($\Phi\Phi$) sector admits the transverse–traceless (TT) graviton entry (spin–2) and the scalar/trace (apeiron) entry (spin–0), but *no* vector register (Definition 9.195). Consequently, physical Gaiamiti excitations reside only in the symmetric metric channel $\hat{g}_{\mu\nu}$ as either the TT Goldstone/tangential mode or the scalar (apeiron) mode; a spin–1 Gaiamiti register is not part of the admissible typed layer.

Vector/longitudinal coherence is carried instead by the τ–kairos family through the Chrono–Kymon typed pair $A^{(\pm)}$ (Definition 10.4). In the constitutive split of the temporal/coherence tensor,

$$\hat{\tau}_{\mu\nu} = S_{\mu\nu}[\hat{\varphi}_A] \;\oplus\; A_{\mu\nu}[\hat{K}^{(\tau)}],$$

the τ channel has spin content $0 \oplus 1$ and is TT–null by construction, i.e. $\mathcal{P}^{(2)}[\hat{\tau}_{\mu\nu}] = 0$ (Definitions 10.4 and 10.13, Sections 10.3 and 10.4, and eq. (10.90)). The antisymmetric piece is encoded by the Chrono–Kymon field strength $\hat{\mathcal{F}}^{(\tau)}_{\mu\nu} = 2\bar{\nabla}_{[\mu}\hat{K}^{(\tau)}_{\nu]}$ (cf. Section 10.7), providing the unique vector register of the weave while remaining orthogonal to the metric TT sector.

Any would–be spin–1 content tied to Gaiamiti metric fluctuations fails to survive in the physical cohomology $H^0(\hat{Q}_{\rm BRST})$ for two independent reasons. (i) Longitudinal vector pieces are gauge (BRST–exact) and therefore vanish in H^0. (ii) Antisymmetric/torsional pieces are, by the constitutive map, routed into the τ–kairos vector block $A_{\mu\nu}[\hat{K}^{(\tau)}]$, not into the symmetric metric channel (Definitions 10.4 and 10.13). Typed orthogonality and triad–compatibility ensure this routing is clean and gauge–parameter independent on certified plateaus (Proposition 9.202 and Definition 10.2).

TT purity further guarantees that no scalar or vector leakage contaminates the graviton readout. The typed $\Pi^{[\rm TT]}_{\bar{\Sigma}}$–null lemmas enforce $\Pi^{[\rm TT]}_{\bar{\Sigma}} S_{\mu\nu}[\cdot] = 0$ and $\Pi^{[\rm TT]}_{\bar{\Sigma}} A_{\mu\nu}[\cdot] = 0$ (Lemma 10.123), while orthogonality/completeness of the typed projectors prevents cross–mixing (Proposition 9.202). Hence the only radiative tensor degree of freedom in the metric channel is the massless spin–2 graviton, identified as the TT Goldstone/tangential mode of the Gaiamiti sector (eq. (10.28), Section 10.17.8, and Definition 10.167).

Consequence. Within the certified, typed weave, the Gaiamiti ($\Phi\Phi$) physical spectrum comprises the spin–2 TT graviton and the spin–0 apeiron (Gaiaton) channels (Definitions 9.195, 10.69 and 10.70), while all spin–1 content resides in the τ–kairos via $A^{(\pm)}$ (Definition 10.4). There is therefore *no spin–1 apeiron* in this framework.

$$\text{Gaiamiti } (\Phi\Phi) \text{ physical spectrum} = \{\text{spin-2 TT graviton}\} \cup \{\text{spin-0 apeiron (Gaiaton)}\},$$
$$\text{spin-1 lives in } \tau \text{ via } A^{(\pm)}.$$

The Gaiamiti field furnishes the full geometric stage: a symmetry–broken vacuum supporting (i) massless, channel–resolved spin–2 gravitons as geometric Goldstone modes (Sections 10.17.6 and 10.17.9) and (ii) a massive radial scalar; vector content is excluded by the canonical bimodal split and TT selection rules [502, 244, 7].

Related — Mition (spin–1 internal connection quantum). The **mition** is a spin–1 excitation of the **relational bundle connection** that enters the covariant derivative, not a substrate excitation of $(\hat{\mathbf{\Phi}})$ (cf. eq. (10.1)). Its mass is generated via internal gauge SSB (BEH) triggered by the Gaiamiti vacuum (cf. Definitions 9.252, 9.258 and 9.261). On the realized weave configuration $(\mathcal{R})$ (cf. Definition 10.1), the spin–1 content is isolated by a transverse, typed vector block and a spin–1 Riesz selector built from spectral islands (cf. Definitions 10.6 and 10.7); the resulting projector commutes with provenance/typed/triad projectors and is Nielsen-flat and retune-stable (cf. Propositions 10.9 and 10.10 and Corollary 10.11). TT isolation is automatic since $(\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}})$ acts on symmetric 2-tensors (cf. Definition 10.3), so the graviton channel is unaffected.

The Tantrakyma Field as the Source of Substance

The *Tantrakyma field* $\hat{\mathbf{\Theta}}(x)$ sources matter and temporal–coherence excitations in the typed weave. Its potential $\mathcal{V}_\Theta$ and the band–limited quadratic kernel determine the spectrum of its quanta (kymons) via the background–fluctuation split and second–variation dynamics (Definitions 9.310 and 10.204 and Section 10.18.6). Two–point functions in the typed (modal) basis and in projected "species" channels are encoded by the matrix propagator $\mathbf{\Delta}_\Theta$ and by the projected propagators Δ_p, respectively (Definitions 10.203 and 10.208; cf. [244, 7]).

Goldstone modes and the Higgs mechanism (matter sector). Spontaneous breaking of a continuous symmetry of the Tantrakyma potential $\mathcal{V}_\Theta$ produces massless Goldstone modes in the Tantrakyma sector, as read from the second–variation analysis on the Σ–realized background $\bar{\Sigma}$ (Definitions 9.310 and 10.204 and Section 10.18.6). For *global* breaking the modes appear as propagating, massless Goldstone kymons; for *local* (gauge) breaking the would–be Goldstones are removed from the asymptotic spectrum and reappear as the longitudinal polarizations of the corresponding gauge vectors (unitary–gauge Higgs mechanism). The construction is carried out within the typed projector framework of the weave (Definition 9.116) and is the standard SSB/Higgs paradigm adapted to the expectation–valued geometry $\bar{\Sigma}$ (see [244, 7, 415]).

Massive excitations. Following SSB (and, where applicable, Higgsing), the remaining Tantrakyma excitations become massive eigenmodes of the local mass operator $\mathcal{M}_\Theta(x)$ (eq. (10.157)). Their masses are read from the pole structure of the projected propagators Δ_p (Section 10.18.5).

1. **Massive spin–0 Higgs kymon (radial scalar).** The radial fluctuation in the Tantrakyma potential—an observable scalar remnant of SSB—acquires mass from the Hessian of $\mathcal{V}_\Theta$ at the vacuum (Section 10.18.6; [244]).

2. **Massive spin–1 Chrono–Kymon (vector).** The vector excitation of the Tantrakyma field furnishes the τ–kairos vector register via the typed family $A^{(\pm)}$ (Definition 10.4). Its field strength $\hat{\mathcal{F}}^{(\tau)}_{\mu\nu} = 2\bar{\nabla}_{[\mu}\hat{K}^{(\tau)}_{\nu]}$ sources the antisymmetric temporal sector and contributes to $\hat{t}_{[\mu\nu]}$ (anisotropic phase flow/"temporal torsion"), in agreement with the scalar–vector constitutive decomposition of the τ–kairos channel and with the operator definition of temporal curvature (Sections 10.3, 10.4 and 10.7 and Definitions 10.4, 10.13 and 10.57).

No spin–2 kymons. The Tantrakyma sector carries *no* irreducible spin–2 excitations: its contributions to the chronogeometric response are strictly non–TT, $\hat{\tau}_{\mu\nu} \in \mathbf{0} \oplus \mathbf{1}$ with $\mathcal{P}^{(2)}[\hat{\tau}_{\mu\nu}] = 0$ (Sections 10.3 and 10.4 and eq. (10.90)). Accordingly, $\mathcal{P}^{(2)}[\delta\hat{\boldsymbol{\Theta}}] = 0$ at the level of physical (typed, BRST–closed) observables, and the radiative spin–2 (TT) sector is exclusively geometric in the Gaiamiti channel—the graviton $\hat{\gamma}_{\mu\nu} = \mathcal{P}^{(2)}[\delta\hat{g}_{\mu\nu}]$ (Definition 10.167 and Section 10.17.8).

Triadic organisation and state dependence. Propagation and mixing for Tantrakyma excitations are triad–resolved into spectral, vacuum, and correlational channels in the typed weave (Definitions 10.2 and 10.208). State dependence enters through the expectation–valued background $\bar{\Sigma}_{\mu\nu}$ and the local mass operator $\mathcal{M}_\Theta(x)$ (Section 10.18.6). Backreaction is mediated by the mixed (off–diagonal) blocks of the chronogeometric/graviton matrix propagator while preserving the TT selection rules (Section 10.17.5, Definition 10.182, and eqs. (10.90) and (10.112)); thus, TT purity of the graviton channel is maintained even in the presence of matter/coherence couplings.

Related — Tantron (spin–1 internal connection quantum). The *tantron* is a spin–1 excitation of the *relational bundle connection* that enters the Tantrakyma covariant derivative—*not* a substrate excitation of $(\hat{\boldsymbol{\Theta}})$ itself (cf. eq. (10.2) and Definition 9.348). Its mass is generated via internal gauge SSB (BEH) triggered by the Tantrakyma vacuum (cf. Definitions 9.252 and 9.261). On the realized weave configuration $(\mathcal{R})$, the tantron's *transverse* content is isolated by a spin–1 Riesz selector built from spectral islands of the connection's vector block, exactly paralleling the spin–2 and spin–0 cases (cf. Definitions 10.6 and 10.7). Longitudinal components are BRST–exact and decouple on $(\mathcal{H}_{\text{phys}})$, so pole locations/residues in the transverse block are *gauge–parameter independent* on certified plateaus and *stable under admissible retunings* (cf. Propositions 10.9 and 10.10 and Corollary 10.11). TT–isolation is automatic because $(\Pi^{[\text{TT}]}_{\bar{\Sigma}})$ acts on symmetric 2-tensors (cf. Definition 10.3), leaving the graviton channel unaffected.

The Deeper Principle at Play

Spin labels *alone* do not determine ontological identity in the typed weave. What matters is the *field of origin*—the geometric substrate $\hat{\boldsymbol{\Phi}}$ (Gaiamiti), the matter/coherence substrate $\hat{\boldsymbol{\Theta}}$ (Tantrakyma), or the *relational bundle connection* that enters the covariant derivatives (cf. eqs. (10.1) and (10.2))—together with the associated typed/constitutive selection rules. Thus two quanta may share the same spin and tensor type yet belong to distinct sectors

with different couplings, causal support, and propagation laws (Definitions 10.4, 10.13 and 10.182 and Sections 10.17.5 and 10.18.6; cf. [244, 7, 502]). Sectorial identity is fixed on the Σ–realized background $\bar{\Sigma}$ using the typed projectors and triad decomposition from Chs. 10–11 (Definitions 9.60, 9.116 and 10.2; see also the realized weave configuration Definition 10.1).

Concretely:

Geometric origin (Gaiamiti) The massless spin–2 graviton is the TT projection of the metric fluctuation, $\hat{h}_{\mu\nu} = \mathcal{P}^{(2)}[\delta\hat{g}_{\mu\nu}] = \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{g}_{\mu\nu}$, the Goldstone/tangential mode of the Gaiamiti sector (Definition 10.167, eq. (10.28), and Section 10.17.8). The massive spin–0 *Gaiaton* is the apeiron (radial/trace) scalar realized in the τ–kairos scalar typed family and isolated by the spin–0 Riesz selector on $K^{(0)}_{\tau}$ (Definitions 10.68 to 10.70). Both are *geometric* (Gaiamiti) quanta, with TT–purity protecting the graviton from scalar/vector leakage (Lemma 10.123 and Proposition 9.202; cf. Definition 10.3).

Matter/coherence origin (Tantrakyma) The τ–kairos tensor carries exclusively spin $0 \oplus 1$ and is TT–null, $\mathcal{P}^{(2)}[\hat{\tau}_{\mu\nu}] = 0$ (Sections 10.3 and 10.4 and eq. (10.90)). Its *vector* register is the Chrono–Kymon typed family $A^{(\pm)}$ with field strength $\hat{\mathcal{F}}^{(\tau)}_{\mu\nu} = 2\bar{\nabla}_{[\mu}\hat{K}^{(\tau)}_{\nu]}$ (Definitions 10.4 and 10.57 and Section 10.7); its *scalar* register is a matter–induced phase fluctuation in the temporal channel, *not* the Gaiaton (despite sharing spin–0), and couples through the τ–channel blocks of the matrix propagator (Definitions 10.4, 10.13 and 10.182).

Higgs kymon (Tantrakyma scalar) Spontaneous symmetry breaking of the Tantrakyma potential $\mathcal{V}_{\Theta}$ produces Goldstone modes (global case) or longitudinal gauge polarizations (local case), with the *radial* scalar acquiring a mass from the Hessian of $\mathcal{V}_{\Theta}$ at the vacuum—the Higgs kymon of the matter sector (Section 10.18.6; see [244, 7, 415]).

Relational–connection origin (internal vectors) The *mition* (Gaiamiti–sourced) and *tantron* (Tantrakyma–sourced) are spin–1 excitations of the *relational bundle connection*—*not* substrate quanta of $\hat{\mathbf{\Phi}}$ or $\hat{\mathbf{\Theta}}$ (cf. eqs. (10.1) and (10.2) and Definitions 9.258 and 9.348). Their masses arise via internal gauge SSB (BEH) (Definitions 9.252 and 9.261). On the realized weave configuration (Definition 10.1) their *transverse* content is isolated by a spin–1 Riesz selector constructed from spectral islands of the vector connection block, exactly paralleling the spin–2 and spin–0 cases (Definitions 10.6 and 10.7). The resulting projectors commute with provenance/typed/triad projectors and are Nielsen–flat and retune–stable (Propositions 10.9 and 10.10 and Corollary 10.11); TT isolation holds because $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ acts on symmetric 2–tensors (Definition 10.3).

Hence, particle classification in the weave is *sectorial*: a quantum is indexed by $(\text{origin}, \text{channel}, \text{spin})$, not by spin alone. In the present context,
The TT selection rules ensure radiative spin–2 content remains purely geometric, while scalar/vector τ–kairos modes encode state–dependent coherence without adding GW polarizations (eqs. (10.90) and (10.101) and Section 10.17.8). Typed orthogonality and the triad commutations keep the channels separated and BRST/Nielsen–safe on certified plateaus (Propositions 9.97 and 9.202 and Definition 10.2); the internal–connection (mition/tantron)

Quantum	Origin	Channel (typed)	Spin
Graviton (TT)	Gaiamiti	metric/TT	2
Gaiaton (apeiron)	Gaiamiti	τ–scalar	0
Chrono–Kymon	Tantrakyma	τ–vector $A^{(\pm)}$	1
Higgs kymon	Tantrakyma	matter scalar	0
Mition	Gaiamiti-sourced (BEH)	internal conn., transverse vector	1
Tantron	Tantrakyma-sourced (BEH)	internal conn., transverse vector	1

projectors commute with the same provenance/typed/triad structure and remain TT–orthogonal by construction (Definitions 10.3 and 10.7 and Propositions 10.9 and 10.10).

Table 10.1: Typed classification of emergent quanta and their geometric/matter roles within the weave.

Excitation	Parent Field	Spin	Contributes to	Nature / Selector
Graviton	Gaiamiti ($\hat{\mathbf{\Phi}}$)	2	Metric TT sector: $\hat{h}_{\mu\nu} = \mathcal{P}^{(2)}[\delta\hat{g}_{\mu\nu}] = \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{g}_{\mu\nu}$	Massless Goldstone/tangential mode of Gaiamiti (Definition 10.167 and eq. (10.28))
Gaiaton (apeiron scalar)	Gaiamiti ($\hat{\mathbf{\Phi}}$)	0	τ–kairos *scalar* typed block $S^{(b)}, S^{(L)}, \tilde{S}^{(p)}$ (Definition 10.4)	Massive scalar; isolated by spin–0 Riesz selector on $K^{(0)}_{\tau}$ (Definitions 10.68 to 10.70)
Chrono–Kymon	Tantrakyma ($\hat{\mathbf{\Theta}}$)	1	τ–kairos *vector* block $A^{(\pm)}$; field strength $\hat{\mathcal{F}}^{(\tau)}_{\mu\nu} = 2\bar{\nabla}_{[\mu}\hat{K}^{(\tau)}_{\nu]}$	Vector (helicity $\pm$); Maxwell/Proca–type on $\bar{\Sigma}$; strictly non-TT (Definitions 10.4 and 10.57, Sections 10.7 and 10.18.6, and eq. (10.90))
Higgs kymon (radial scalar)	Tantrakyma ($\hat{\mathbf{\Theta}}$)	0	Matter scalar channel (radial mode of $\mathcal{V}_\Theta$)	Massive scalar; mass from Hessian of $\mathcal{V}_\Theta$ at the vacuum (Section 10.18.6)

How Kymons Make Matter and Write Time

Following the typed identification of the τ–vector register (Chrono–Kymon) and its field strength (Definitions 10.4 and 10.57 and Section 10.7), it is important to foreground the *dual role* of *kymons*—the excitations of the Tantrakyma field Θ. On one hand, they generate the spectrum of massive matter degrees of freedom; on the other, they inscribe themselves into the temporal/coherence channel through the τ–kairos tensor. These functions are *linked but not identical*, and their interplay is essential for understanding how matter and temporal coherence are co–woven within the dual substrate (Φ, Θ) (Section 10.18.6 and Definitions 10.4 and 10.13).

(A) Mass generation (matter side). Projected Tantrakyma modes $\delta\hat{\mathbf{\Theta}}^{(p)}$ acquire masses through the local mass operator $\mathcal{M}_\Theta(x)$, the Hessian of the potential $\mathcal{V}_\Theta$ around the vacuum background (eq. (10.157), Definition 10.204, and Section 10.18.6). The eigenvalue problem

$$\boxed{\mathcal{M}_\Theta(x)\,u_p(x) = m_p^2(x)\,u_p(x) \quad\Longrightarrow\quad \text{poles of } \Delta_p(k) \;\text{ at }\; k^2 = m_p^2(x),} \tag{10.77}$$

Table 10.2: Unified duality of emergent quanta and their properties. The Gaiamiti and Tantrakyma fields furnish complementary registers whose excitations interlock in the constitutive split.

Feature	**Geometric Sector (Gaiamiti field Φ)**	**Matter & Coherence Sector (Tantrakyma field Θ)**
Fundamental Role	Chronogeometry / metric sector ($\bar{\Sigma}$, TT channel) (Definition 9.60)	Substance & temporal coherence (kymons, τ–vector) (Section 10.18.6 and Definition 10.4)
Primary Quanta	**Apeiron** $\rightarrow$ **Gaiaton** (scalar)	**Kymon** (matter/coherence)
Emergent Spectrum	Massless **spin–2 graviton**; massive **spin–0 Gaiaton** (apeiron scalar) (Definitions 10.70 and 10.167 and eq. (10.28))	General **kymon** excitations; distinguished **spin–1 Chrono–Kymon**; **Higgs kymon** (spin–0) (Section 10.18.6 and Definitions 10.4 and 10.57)
Spin Domain	Extremes 0 and 2 (no vector register in $\Phi\Phi$) (Definition 9.195)	Intermediate 0 and 1 (strictly non-TT in τ) (eq. (10.90) and Sections 10.3 and 10.4)
Mechanism	SSB of $\mathcal{V}_\Phi$: TT Goldstone (graviton) + radial scalar (Gaiaton) (eq. (10.28) and Definition 10.70)	SSB of $\mathcal{V}_\Theta$: Goldstones / longitudinal gauge modes + Higgs kymon (Section 10.18.6)
Contribution to τ–kairos $\hat{\tau}_{\mu\nu}$	**Spin–0** scalar piece via $S^{(b)}, S^{(L)}, \tilde{S}^{(p)}$ (Gaiaton channel) (Definitions 10.4 and 10.69)	**Spin–1** vector piece via $A^{(\pm)}$ (Chrono–Kymon) with $\hat{\mathcal{F}}^{(\tau)}_{\mu\nu}$ (Definitions 10.4 and 10.57 and Section 10.7)

determines the mass spectrum of emergent matter excitations, with u_p the species eigenvectors and Δ_p the projected propagators (Definitions 10.203 and 10.208). Thus the kymon family functions as the dynamical source of materiality in the Tantrakyma sector (for the SSB/Higgs narrative and the radial *Higgs kymon*, see Section 10.18.6).

(B) Temporal component (coherence side). Simultaneously, Tantrakyma fluctuations imprint upon the *temporal/coherence tensor* $\hat{\tau}_{\mu\nu}$. In the *vector channel* this imprint is carried by the *Chrono–Kymon*, extracted by the τ–vector typed projector from Tantrakyma fluctuations,

$$\hat{K}^{(\tau)}_{\mu} \;:=\; \left(\Pi^{[A]}_{\tau}\,\delta\hat{\mathbf{\Theta}}\right)_{\mu} \qquad \text{(helicity basis } A^{(\pm)} \text{ in } \textit{Definition } 10.4\text{)},$$

whose field strength yields antisymmetric temporal curvature,

$$\boxed{\hat{\mathcal{F}}^{(\tau)}_{\mu\nu} \;=\; 2\,\bar{\nabla}_{[\mu}\hat{K}^{(\tau)}_{\nu]},} \tag{10.78a}$$

$$\boxed{\hat{t}_{[\mu\nu]} \;=\; \chi_{\tau}\,\hat{\mathcal{F}}^{(\tau)}_{\mu\nu},} \tag{10.78b}$$

with χ_τ fixed by the constitutive map (Definitions 10.4 and 10.13 and Section 10.7). By contrast, the *scalar* temporal channel inherits its content from the Gaiamiti apeiron (the *Gaiaton*) realized in the τ–scalar typed family and isolated by the spin–0 selector (Definitions 10.68 and 10.70). Altogether, the constitutive decomposition

$$\hat{\tau}_{\mu\nu} \;=\; S_{\mu\nu}[\hat{\varphi}_A] \;\oplus\; A_{\mu\nu}[\hat{K}^{(\tau)}]$$

(Definitions 10.4 and 10.13 and eq. (10.90)) shows that τ–kairos is a mixed object: scalar feed from Gaiamiti (Gaiaton), vector feed from Tantrakyma (Chrono–Kymon).

(B') Clarification: Chrono–Kymon vs. internal connection vectors. The *Chrono–Kymon* is a τ–channel vector extracted from Tantrakyma fluctuations by typed projectors and governs *temporal/coherence* dynamics (Definitions 10.4 and 10.57). It is *distinct* from the spin–1 internal connection quanta—the *mition* and *tantron*—which originate from the *relational bundle connection* entering the covariant derivatives (not from $\hat{\mathbf{\Phi}}$ or $\hat{\mathbf{\Theta}}$) (eqs. (10.1) and (10.2) and Definitions 9.258 and 9.348). On the realized weave configuration (Definition 10.1) these internal vectors are isolated by a *spin–1* Riesz selector built from spectral islands of the connection's vector block (Definitions 10.6 and 10.7); their transverse projectors commute with provenance/typed/triad projectors and are Nielsen–flat and retune–stable (Propositions 10.9 and 10.10 and Corollary 10.11). TT isolation holds because $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ acts on symmetric 2–tensors (Definition 10.3). Thus, *Chrono–Kymon* (τ–vector) and *mition/tantron* (internal connection vectors) are separate origins, channels, and ledgers—no duplication occurs.

(C) Selection rule and non–duplication. Despite this dual role, a strict separation persists: kymons *do not* generate spin–2 radiation. The TT projector annihilates all τ–kairos contributions,

$$\boxed{\mathcal{P}^{(2)}[\hat{\tau}_{\mu\nu}] = 0, \qquad \text{gravitational waves:}\;\; \hat{h}_{\mu\nu} = \mathcal{P}^{(2)}[\delta\hat{g}_{\mu\nu}] = \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{g}_{\mu\nu},} \tag{10.79}$$

so that GW radiation remains *purely geometric*, confined to the Gaiamiti/graviton TT sector (eqs. (10.90) and (10.101), Definition 10.167, and Sections 10.17.8 and 10.18.7). The kymon imprint is entirely scalar/vector and coherence–driven, never duplicating the spin–2 graviton channel.

Kymons are the fundamental quanta of the Tantrakyma field. They *make matter*, by acquiring masses through $\mathcal{M}_\Theta$ (with the radial *Higgs kymon* as the canonical scalar remnant of SSB), and they *write time*, by sourcing the τ–kairos tensor—most visibly through the spin–1 *Chrono–Kymon*, which imprints torsional/coherence structure on $\bar{\Sigma}$ via $\hat{\mathcal{F}}^{(\tau)}_{\mu\nu}$ (Definition 10.57 and Section 10.7). In this dual role, kymons bridge the material and temporal aspects of the dual substrate: they provide substance to matter while simultaneously modulating the relational timer that synchronizes geometry and quantum coherence (Section 10.16 and Proposition 10.150).

Reciprocity of the Sectors: From Pulse to Wave, From Wave to Form

A central feature of the dual–substrate construction is the *bidirectional traffic* between the geometric (Gaiamiti) and matter/coherence (Tantrakyma) registers. The constitutive split

$$\Sigma_{\mu\nu} = g_{\mu\nu}[\Phi] \;+\; t_{\mu\nu}[\Phi, \Theta]$$

(eq. (10.101) and Section 10.17.4) does not isolate geometry from matter but enables a disciplined reciprocity under strict typed and spin constraints: *no spin–1* in the Gaiamiti (metric) channel and *no spin–2* in the Tantrakyma (τ–kairos) channel (Definition 9.195, eq. (10.90), and Section 10.18.7).

Geometry → Matter (pulse release). On the geometric side, the spin–0 *apeiron* register of Φ—the **Gaiaton**—appears as a scalar pulse of geometric inertia. Certified insertions of the Gaiaton live in the τ–kairos *scalar* typed family $S^{(b)}, S^{(L)}, \tilde{S}^{(p)}$ and are isolated by the spin–0 Riesz selector on $K^{(0)}_\tau$ (Definitions 10.4 and 10.68 to 10.70). Through the off–diagonal entries of the fourfold inter–modal structure, this scalar geometric energy can *discharge* into matter, feeding *kymons* of the Tantrakyma sector via the interaction action (Definition 10.223), in the same typed, triad–compatible sense used throughout Chs. 10–11.

Matter → Geometry (wave inscription). Conversely, on the matter/temporal side, the **Chrono–Kymon** (spin–1) inscribes back into geometry by writing phase and timing into the temporal block $t_{\mu\nu}[\Phi, \Theta]$. Concretely, the τ–kairos decomposes as

$$\hat{\tau}_{\mu\nu} = S_{\mu\nu}[\hat{\varphi}_A] \;\oplus\; A_{\mu\nu}[\hat{K}^{(\tau)}],$$

with the scalar piece inherited from the Gaiaton channel and the vector piece supplied by the Chrono–Kymon (Definitions 10.4 and 10.13). The latter carries field strength $\hat{\mathcal{F}}^{(\tau)}_{\mu\nu} = 2\bar{\nabla}_{[\mu}\hat{K}^{(\tau)}_{\nu]}$, sourcing the antisymmetric temporal curvature and entering the constitutive map as $\hat{t}_{[\mu\nu]} = \chi_\tau\, \hat{\mathcal{F}}^{(\tau)}_{\mu\nu}$ (Definition 10.57 and Section 10.7). This inscription *never* duplicates the graviton: $\mathcal{P}^{(2)}[\hat{\tau}_{\mu\nu}] = 0$ by construction, so GW radiation remains confined to the TT graviton sector (eq. (10.90), Definition 10.167, and Section 10.17.8).

Gatekeeping by τ–kairos. Across both directions, the τ–kairos acts as the gatekeeper of reciprocity. It prescribes *when* exchanges are admissible (coherence/plateau conditions) and *which* channels are open (scalar/vector only), enforcing the typed projector rules and triad commutations (Definitions 10.2 and 10.4 and Propositions 9.97 and 9.202). In this rigorously governed reciprocity—pulse becomes wave, wave becomes form—τ–kairos synchronizes the exchange without violating TT purity or sectorial identity.

Significance and Testable Consequences

The dual role of kymons is conceptually and empirically consequential: it unifies *mass generation* and *temporal coherence* within a single Tantrakyma sector while preserving the purely geometric nature of spin–2 radiation. This synthesis is realized with typed projectors, triad–resolved propagators, and background–field quantization on the expectation–valued geometry $\bar{\Sigma}$ (Definitions 9.60, 10.182 and 10.208; cf. [244, 7, 415, 502]).

Clean polarization prediction (consistency with GR) The TT selection rule

$$\mathcal{P}^{(2)}[\hat{\tau}_{\mu\nu}] = 0$$

forbids any additional tensor polarizations from the τ–channel; gravitational waves remain purely TT and geometric (eqs. (10.90) and (10.101) and Section 10.18.7). Thus the framework reproduces the standard two tensor polarizations of GR (Section 10.17.8; cf. [502, 415]). The τ–channel can modulate phases/amplitudes through matrix–propagator couplings without creating extra polarizations (Sections 10.17.5 and 10.17.6).

State–dependent masses and ranges (matter side) Kymon species masses arise as eigenvalues of the Tantrakyma mass operator $\mathcal{M}_\Theta(x)$ (eq. (10.157) and Sections 10.18.5 and 10.18.6), with propagator poles at $k^2 = m_p^2(x)$. In particular, the *Chrono–Kymon* admits a Proca mass $m_{(1)}$ and screening length $\ell_{(1)} \sim m_{(1)}^{-1}$ on $\bar{\Sigma}$ (Section 10.18.6); because $m_{(1)}$ is state–dependent, its effective range is environment–sensitive, making Yukawa–like laboratory/astrophysical probes natural tests.

Temporal torsion phenomenology (coherence side) Via the constitutive relation

$$\hat{t}_{[\mu\nu]} = \chi_\tau \, \hat{\mathcal{F}}^{(\tau)}_{\mu\nu},$$

the Chrono–Kymon sources antisymmetric temporal curvature (Section 10.7 and Definition 10.57). This leads to testable signatures: orientation/history– dependent phase shifts in interferometry, coherence transport in condensed–matter platforms, and phase–locking effects in quantum–optical arrays. In coherence–certified regimes the dynamics are Maxwell/Proca–type on $\bar{\Sigma}$ (Section 10.18.6), enabling quantitative modeling.

Triadic fingerprints Triad–resolved propagation produces distinctive imprints: matter propagators decompose as $\mathbf{\Delta}_\Theta = \sum_a \mathbf{\Delta}_\Theta^{(a)}$, and graviton propagators as $\mathbf{G}_\Sigma^{(2)} = \sum_a \mathbf{G}_\Sigma^{(2,a)}$ (Sections 10.17.6 and 10.18.7). This separation into correlational, vacuum, and spectral channels supports *source tomography*: diagnosing whether an effect is coherence–driven, vacuum–polarization–induced, or energy–momentum–driven.

Dark–sector avenue (conservative) A massive, electrically neutral, spin–1 Chrono–Kymon contributes to $\hat{t}_{\mu\nu}$ as a stress component while leaving the TT graviton sector unchanged. Relic Chrono–Kymons thus constitute a conservative dark–sector candidate—quantum–torsional and consistent with $\bar{\Sigma}$–causality and TT constraints (Section 5.4.5)—with potential imprints in cosmic stress–energy budgets and coherence observables.

In brief. The Tantrakyma sector furnishes *what* matter is (through mass spectra and mixings) and *how* time coheres (through temporal torsion and coherence dynamics). The Gaiamiti sector, by contrast, fixes *how* space curves (via the TT graviton) and supplies the scalar Gaiaton that populates the τ–scalar ledger. The separation is sharp in spin content yet intertwined operationally through channel–resolved propagators, exactly as laid out in Sections 10.17.5 and 10.18.7. This architecture grounds the theory in the standard QFT–in–curved–spacetime toolkit [244, 7, 415, 502], while pointing to testable phenomena in quantum–coherence laboratories, cosmology, and black–hole physics.

Typed portals between ledgers

Definition 10.151 (Typed portals between ledgers)**.** A *portal* is a typed/triad–compatible interaction (or mixing map) that connects two otherwise orthogonal ledgers (e.g. Gaiamiti scalar $\leftrightarrow$ τ–scalar; visible photon $\leftrightarrow$ τ–vector; visible currents $\leftrightarrow$ connection vectors), while preserving the weave's selection rules. Formally, a portal is specified by:

Target/source ledgers and channels
A pair $(\mathcal{L}_{\mathrm{src}}^{[\chi_{\mathrm{src}}]}, \mathcal{L}_{\mathrm{tgt}}^{[\chi_{\mathrm{tgt}}]})$ of typed subspaces selected by the commuting projectors (provenance/typed/triad) on the realized weave configuration (cf. Definitions 9.116, 10.1 and 10.2).

Small strength parameter(s)
Dimensionless couplings $\lambda \in \{\, g_{\mathrm{mix}},\ \epsilon,\ \chi_\tau \lambda', \dots \}$ that control the portal's observable imprint (cf. Section 10.16).

Structural constraints (typed safety)
The portal map P obeys

$$[\mathsf{P}, \Pi^{[\chi]}] = [\mathsf{P}, \Pi_\alpha] = 0, \qquad \mathsf{P} = \mathsf{P}_{\mathrm{phys}}\, \mathsf{P}\, \mathsf{P}_{\mathrm{phys}},$$

with $\Pi^{[\chi]}$ the typed projectors and Π_α the triad projectors (Definitions 9.116 and 10.2), and with $\mathsf{P}_{\mathrm{phys}}$ the BRST–physical projector. If spectral selectors are used (Riesz), P additionally commutes with the island projectors (Definitions 10.6 and 10.7 and Propositions 10.9 and 10.10).

TT–purity (no graviton contamination)
There is no leakage

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \mathsf{P} \;=\; \mathsf{P}\, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \;=\; 0,$$

so that the portal never feeds the radiative spin–2 channel (cf. Definition 10.3 and eq. (10.90)).

Realization

Either as an *off–diagonal block* in a typed/triad–resolved propagator (matrix Green's function) or as a *diagnostic realized reference configuration Lagrangian* cross–term, both read with the calibrated meters and $\bar{\Sigma}$–weighted inner products (Definitions 10.82, 10.117 and 10.182, Section 10.17.5, and eq. (10.112)).

Remark 10.152 (Canonical portal families with examples)**.** The following instances satisfy Definition 10.151; each is constrained by Section 10.16.

Scalar (Gaiaton–τ) portal

Allowed mixing in the *scalar* block between the Gaiamiti scalar (Gaiaton) and the τ–scalar:

$$\mathbf{G}^{(0)}(k) = \begin{pmatrix} G_{\text{Gaiaton}} & \boxed{g_{\text{mix}}\, G_{\text{cross}}} \\ g_{\text{mix}}\, G_{\text{cross}} & G_{\tau\text{-scalar}} \end{pmatrix},$$

read with the typed projectors (Definitions 10.68 to 10.70 and 10.182 and Section 10.17.5). Equivalently on the diagnostic realized reference configuration: $\delta\mathcal{L} \supset -\frac{1}{2}\, g_{\text{mix}}\, \varphi_{\text{G}} \varphi_{\tau}$.

Vector kinetic–mixing portal

A dark–photon–like, gauge–invariant mixing of field strengths for the τ–vector (Chrono–Kymon):

$$\delta\mathcal{L} \;\supset\; -\frac{\epsilon}{2}\, F^{\mu\nu}_{\text{vis}}\, \mathcal{F}^{(\tau)}_{\mu\nu}, \qquad \hat{\mathcal{F}}^{(\tau)}_{\mu\nu} = 2\bar{\nabla}_{[\mu} \hat{K}^{(\tau)}_{\nu]},$$

with $\epsilon \simeq g_{\text{mix}}$ (Definitions 10.4 and 10.57 and Section 10.7). An analogous kinetic–mixing portal can be written for the *connection* vectors (mition/tantron) using their field strengths and the spin–1 connection selector (cf. Definition 10.7).

Composition–dependent (fifth–force) portal

Couplings of a light vector to matter currents (e.g. B, L, $B-L$):

$$\delta\mathcal{L} \;\supset\; g_X\, V_\mu\, J^\mu_{B-L},$$

mapping to WEP tests; the effective strength scales as $\alpha_{\text{eff}} \propto g^2_{\text{mix}}$ (with constitutive weights where applicable) (cf. item B in Section 10.16).

Spin / torsion–like portal (antisymmetric temporal channel)

Using the constitutive law

$$\hat{t}_{[\mu\nu]} = \chi_\tau\, \hat{\mathcal{F}}^{(\tau)}_{\mu\nu},$$

one may couple to fermion axial currents (spin), producing torsion–like signatures; constraints effectively bound the *product* $\chi_\tau g_{\text{mix}}$ (cf. item C in Section 10.16).

Oscillatory (ultralight) portal

For a coherent background with frequency $\omega \simeq m_{(1)}$, portals induce phase/frequency modulation in clock and interferometric meters; sensitivities constrain $g_{\text{mix}}\chi_\tau$ across the Hz–mHz band (cf. item E in Section 10.16).

Parity–odd birefringent portal

If the portal plus constitutive transport violates parity, isotropic EB rotation in the CMB bounds $|g_{\text{mix}}\chi_\tau|$ (cf. item F in Section 10.16).

Connection–vector portals (mition/tantron)

The same structures apply to the internal connection vectors isolated by the spin–1 connection selector: kinetic mixing with the visible photon, composition–dependent couplings, and spin–coupled portals, all commuting with the typed/triad projectors and TT–null by construction (cf. Definitions 10.1, 10.3, 10.6 and 10.7).

What Present Data Allow (Mapping into $\{m_{(1)}, \chi_\tau, g_{\rm mix}\}$)

A. Absence of extra GW polarizations ⇒ limits on spin–1 leakage into the TT channel. Multi–event LVK tests (GWTC–3) report signals consistent with pure–tensor GR radiation and no statistically significant scalar/vector polarizations [122]. This supports the TT selection rule $\mathcal{P}^{(2)}[\hat{\tau}_{\mu\nu}] = 0$ and bounds any portal that would feed Chrono–Kymon power into the radiative channel through off–diagonal entries of the graviton/temporal matrix propagator (Definition 10.182, Section 10.17.5, and eq. (10.112)).

B. Weak Equivalence Principle (WEP) ⇒ bounds on long–range, composition–dependent portals. MICROSCOPE reports a null WEP–violation signal, $\eta = (-1 \pm 9) \times 10^{-15}$ at 1σ [503, 504], implying $|\eta| \lesssim 10^{-14}$ overall. Any Chrono–Kymon portal that couples nonuniversally to baryon number, $B-L$, or species number with range $\lambda \gtrsim 10^5\,\mathrm{m}$ must have a strongly suppressed effective Yukawa strength $\alpha_{\rm eff} \propto g_{\rm mix}^2$ (and, where applicable, proportional to the constitutive weight χ_τ). This constrains the ultralight regime $m_{(1)} \lesssim 10^{-12}\,\mathrm{eV}$.

C. Spin–dependent "torsion–like" couplings ⇒ bounds at the 10^{-31} GeV level. Chrono–Kymon exchange that mimics an axial torsion background via the constitutive map $\hat{t}_{[\mu\nu]} = \chi_\tau \hat{\mathcal{F}}^{(\tau)}_{\mu\nu}$ (coupled to fermion spin) is bounded by torsion–pendulum and maser experiments. SME analyses bound torsionlike components at $\lesssim 10^{-31}\,\mathrm{GeV}$, limiting $\chi_\tau g_{\rm mix}$ for long–range spin portals [56, 505, 506].

D. Kinetic–mixing / vector–portal limits ⇒ broad constraints on $g_{\rm mix}$. If the Chrono–Kymon kinetically mixes with the photon, $\epsilon \simeq g_{\rm mix}$, dark–photon bounds apply. Global reviews [430, 507] summarize constraints from $\epsilon \sim 10^{-12}$ (radio/cosmology) to $\epsilon \sim 10^{-3}$ (MeV–GeV). Parker Solar Probe measurements constrain 70 kHz–20 MHz particularly strongly, cutting into the ultralight vector sector [508, 430].

E. Clock networks and ultralight coherent backgrounds ⇒ constraints on oscillatory portals. Optical–clock networks bound periodic modulations of fundamental constants sourced by ultralight bosons. Modeling a coherent Chrono–Kymon background $\hat{K}^{(\tau)}_\mu$ with $\omega \simeq m_{(1)}$, non–observation constrains $g_{\rm mix}\chi_\tau$ responsible for phase/frequency shifts. Sensitivity peaks for $m_{(1)} \sim 10^{-22}$–$10^{-15}\,\mathrm{eV}$ (Hz–mHz), complementing plasma/radio probes [509].

F. Cosmic birefringence (isotropic EB rotation) ⇒ bounds on parity–odd transport. CMB polarization analyses suggest a marginal isotropic EB rotation at the few–tenths–degree level [429]. If $\hat{t}_{[\mu\nu]} = \chi_\tau \hat{\mathcal{F}}^{(\tau)}_{\mu\nu}$ combined with a portal induces parity–violating temporal transport,

these observations bound $|g_{\rm mix}\chi_\tau|$. Because EB vanishes in parity–conserving ΛCDM, any confirmed nonzero rotation would directly trace τ–kairos–mediated birefringence.

G. Near–future reach (atom–interferometry mid–band). Large–baseline atom–interferometer programs (**MAGIS–100**, **AION**) will probe the GW mid–band (10^{-2}–10 Hz) and ultralight dark–sector fields. Projected reach for vector portals with $m_{(1)} \sim 10^{-15}$–10^{-12} eV can strengthen limits on $g_{\rm mix}$, targeting coherence–preserving regimes where Chrono–Kymon dynamics are most active [510, 511].

Synthesis of constraints. Current data allow a massive spin–1 Chrono–Kymon that modulates temporal phase torsion (χ_τ) and participates in the Tantrakyma spectrum ($\mathcal{M}_\Theta$), subject to: (i) TT silence in the radiative channel (LVK polarization tests) consistent with $\mathcal{P}^{(2)}[\hat{\tau}_{\mu\nu}] = 0$ (eqs. (10.90) and (10.101)); (ii) sufficient universality or short range to evade WEP bounds (MICROSCOPE); and (iii) consistency with kinetic–mixing limits (especially tight in the kHz–MHz window). Spin–pendulum/SME constraints cap long–range spin portals near 10^{-31} GeV. The most promising discovery windows are ultralight coherent backgrounds accessible to clock networks and the GW mid–band in atom–interferometry—precisely the coherence–selective regimes highlighted by the τ–kairos pipeline (Proposition 10.150).

Speculative Narrative: The Dark Sector

Within the typed weave, two *non–tensorial* (TT–null) registers arise as natural dark–sector candidates, each tied to a distinct substrate and admitted via the τ–kairos typed layer (Definition 10.4 and eq. (10.90)).

Gaiaton (spin–0, scalar, radial). The **Gaiaton** is the scalar (apeiron) excitation originating in the Gaiamiti sector and *realized* on the τ–kairos *scalar* typed family $S^{(b)}, S^{(L)}, \tilde{S}^{(p)}$, isolated spectrally by the spin–0 Riesz selector on the scalar kinetic generalized eigen-operator pair $K_\tau^{(0)}$ (Section 10.8 and Definitions 10.4 and 10.68 to 10.70). Its potential $\mathcal{V}_\Phi$ naturally supports slow–roll or quintessence–like dynamics, providing an evolving equation of state $w(z)$ for dark energy [486]. In a different mass/coupling regime, a massive, (meta)stable Gaiaton can behave as cold dark matter through coherent (misalignment) oscillations, subject to production and fifth–force constraints [485]. In brief: the Gaiaton is *prominent for dark energy* and *plausible for dark matter* when stabilized.

Chrono–Kymon (spin–1, vector, temporal). The **Chrono–Kymon** furnishes the τ–kairos *vector* register $A^{(\pm)}$ with field strength $\hat{\mathcal{F}}^{(\tau)}_{\mu\nu} = 2\bar{\nabla}_{[\mu}\hat{K}^{(\tau)}_{\nu]}$ (Definitions 10.4 and 10.57 and Section 10.7). As a massive vector, it is a canonical dark–matter candidate (misalignment or freeze–in) that *does not* alter the TT graviton sector (Sections 10.3 and 10.4 and eq. (10.90)). A dark–energy role is more constrained: generic vector condensates threaten large–scale anisotropy, but if τ–kairos dynamics enforce effective isotropy (e.g., triad averaging on certified plateaus), a subdominant or early–dark–energy–like component may be allowed (Definition 10.2 and Proposition 9.97). Summary: the Chrono–Kymon is *prominent for dark matter*, with at most conditional relevance for dark energy.

Mition (spin–1, internal connection, Gaiamiti–sourced). The **mition** is a spin–1 excitation of the *relational operator bundle connection* $\mathcal{A}_\mu$ that enters the Gaiamiti covariant derivative (cf. Section 6.6, Definitions 6.192 and 9.258, Remarks 6.198 and 6.200, and eq. (10.1)). Its mass arises via internal gauge SSB (BEH) triggered by the Gaiamiti vacuum (cf. Definitions 9.252 and 9.261). On the realized weave configuration it is isolated by a *spin–1* Riesz selector acting on the connection's transverse vector block (cf. Definitions 10.1, 10.6 and 10.7), commuting with the provenance/typed/triad projectors and remaining TT–orthogonal since $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ acts on symmetric 2–tensors (cf. Propositions 10.9 and 10.10, Definition 10.3, and Corollary 10.11). Phenomenologically, a massive mition is a canonical dark–matter candidate (coherent misalignment or freeze–in), with portal strengths constrained by the bounds summarized in Section 10.16. It is *not* a τ–kairos register; rather, it populates the *connection* ledger and influences $\Sigma_{\mu\nu}$ only indirectly via its effects on (Φ, Θ) in the constitutive split (cf. eq. (10.101)).

Tantron (spin–1, internal connection, Tantrakyma–sourced).

The **tantron** is Tantrakyma–sourced: a spin–1 excitation of the relational connection $\mathcal{A}_\mu$ that enters the Tantrakyma covariant derivative (cf. eq. (10.2) and Definition 9.348). Like the mition, it acquires mass via Brout–Englert–Higgs (BEH).

On the realized weave configuration, its transverse, BRST–physical modes are isolated by the *spin–1* connection selector constructed from spectral islands (cf. Definitions 10.1, 10.6 and 10.7). The resulting projectors commute with the provenance/typed/triad structure and remain TT–orthogonal, since $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ acts only on symmetric 2–tensors (cf. Propositions 10.9 and 10.10 and Definition 10.3).

As a massive, weakly coupled vector, the tantron is a viable dark–matter candidate. Relevant portal constraints include WEP tests, kinetic–mixing limits, bounds on spin–dependent torsionlike couplings, clock–network searches for ultralight backgrounds, and cosmic–birefringence measurements (cf. Section 10.16). It lies outside the τ–kairos registers and provides an *additional* non–TT dark–sector pathway sourced by the relational connection rather than by the substrate fields themselves.

Synthesis. Together, the scalar **Gaiaton** anchors the dark–energy narrative (and potentially a CDM fraction), while the vector **Chrono–Kymon** anchors the dark–matter narrative (with limited dark–energy scope under isotropy). In addition, the *relational connection* sector provides two further TT–null vector candidates—the **mition** (Gaiamiti–sourced) and the **tantron** (Tantrakyma–sourced)—which originate from the operator bundle connection and are isolated on the realized weave configuration by the *spin–1* connection selector built from spectral islands (cf. Section 6.6, Definitions 6.192, 10.1, 10.6 and 10.7, and Remarks 6.198 and 6.200). These internal vectors are *not* τ–kairos registers; they populate the connection ledger and affect $\Sigma_{\mu\nu}$ only indirectly via their influence on (Φ, Θ) in the constitutive split (cf. eq. (10.101)), with phenomenological portals constrained by Section 10.16. Throughout, the spin–2 graviton sector remains purely geometric—protected by TT selection and typed orthogonality—so gravitational–wave polarizations are unaffected even when dark–sector dynamics are active (Definition 10.167, eqs. (10.28) and (10.90), and Proposition 9.202). The selection rules (no spin–1 in the Gaiamiti/metric channel; no spin–2 in the τ–kairos/-Tantrakyma channel) thus remain intact, while the dark sector is populated by quanta that

are *natural outgrowths* of the dual–substrate construction.

Dark–Sector Excitations of the τ–Kairos

The τ–kairos *typed* structure itself ($\hat{\tau}_{\mu\nu} = \delta\hat{t}_{\mu\nu}$) provides a natural umbrella for dark–sector phenomenology. As established in Section 10.17.3, $\hat{\tau}_{\mu\nu}$ admits scalar and vector registers—captured by the constitutive split

$$\hat{\tau}_{\mu\nu} = S_{\mu\nu}[\hat{\varphi}_A] \ \oplus \ A_{\mu\nu}[\hat{K}^{(\tau)}] \quad (\text{TT–null: } \mathcal{P}^{(2)}[\hat{\tau}_{\mu\nu}] = 0)$$

(Definitions 10.4 and 10.13 and eq. (10.90)). The scalar register is realized by the Gaiaton (Gaiamiti origin, τ–scalar realization), while the vector register is supplied by the Chrono–Kymon (Tantrakyma origin, τ–vector realization) (Definitions 10.4, 10.57 and 10.69). These are strictly spin–0/1 excitations, consistent with the absence of additional spin–2 modes (Section 10.18.7).

Taken together, these results support the view that τ–kairos, via its scalar and vector typed registers, *constitutively imprints* the dark sector. Accordingly, we adopt the following identifications for phenomenology: the *Gaiaton* (scalar, τ–scalar) and the *Chrono–Kymon* (vector, τ–vector) provide, respectively, the scalar and vector contributions to dark energy and dark matter, while the spin–2 graviton channel remains purely geometric and TT.

Proposition 10.153 (Scalar τ–kairos register realized by the Gaiaton)**.** *The spin–0 register of the τ–kairos, realized by the* Gaiaton*, contributes to the dark sector in two complementary ways:*

(i) Dark energy
The potential $\mathcal{V}_\Phi$ supports slow–roll and quintessence–like dynamics, enabling an evolving equation of state $w(z)$ (cf. Sections 10.8 and 10.16; see [486]).

(ii) Dark matter
In a massive, (meta)stable regime the Gaiaton acts as a cold, pressureless condensate via coherent oscillations (misalignment), consistent with the weave's selectors and meters (cf. Section 10.8 and Definitions 10.1 and 10.68; see [485]).

In both cases the Gaiaton enters the chronogeometric response through the scalar contribution $S_{\mu\nu}[\hat{\varphi}_A]$ in the constitutive map (Definitions 10.4 and 10.13).

Proposition 10.154 (Vector τ–kairos register supplied by the Chrono–Kymon)**.** *The spin–1 register of the τ–kairos, provided by the* Chrono–Kymon*, contributes as follows:*

(i) Dark matter
A massive Chrono–Kymon produced by misalignment or freeze–in yields a relic vector condensate while preserving TT–purity of gravitational waves (Section 10.18.7, eq. (10.90), and Definition 10.4).

(ii) Dark energy (conditional)
A subdominant contribution is possible only if τ–kairos dynamics enforce effective isotropy (e.g., via triads or domain averaging on certified plateaus), avoiding large–scale anisotropy (Definition 10.2 and Proposition 9.97).

Constitutively, the Chrono–Kymon supplies the vector piece $A_{\mu\nu}[\hat{K}^{(\tau)}]$ in the τ–kairos decomposition, embedding dark–sector dynamics in the coherence channel without affecting the spin–2 graviton sector (Definitions 10.4, 10.13 and 10.57 and Section 10.7).

Dark–Sector Candidates from the Relational Connection (Mition, Tantron)

Proposition 10.155 (Internal–connection vector *mition* as a dark–matter candidate)**.** *Let $\mathcal{A}_\mu$ be the operator bundle connection of Chapter 7 (cf. Section 6.6, Definition 6.192, and Remarks 6.198 and 6.200) entering the Gaiamiti covariant derivative (eq. (10.1)). Assume internal gauge SSB (BEH) in the Gaiamiti vacuum so that the transverse, BRST–physical modes of $\mathcal{A}_\mu$ acquire a mass (cf. Definitions 9.252, 9.258 and 9.261). On the realized weave configuration (Definition 10.1), let the connection's vector block be band–limited and let its spin–1 content be isolated by the Riesz selector built from spectral islands (Definitions 10.6 and 10.7). Then:*

(i) Dark matter

A massive mition *produced by coherent misalignment or freeze–in provides a viable vector dark–matter component, with portal strengths constrained by the bounds in Section 10.16.*

(ii) Typed/TT safety

The mition's projector commutes with provenance/typed/triad projectors and remains TT–orthogonal because $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ acts on symmetric 2–tensors (Propositions 10.9 and 10.10, Definition 10.3, and Corollary 10.11); hence gravitational–wave polarizations are unaffected (eq. (10.90)).

(iii) Ledger separation

The mition is not *a τ–kairos register; it populates the* connection *ledger and influences $\Sigma_{\mu\nu}$ only indirectly via its effects on (Φ, Θ) in the constitutive split (eq. (10.101)).*

Proposition 10.156 (Internal–connection vector *tantron* as a dark–matter candidate)**.** *Let $\mathcal{A}_\mu$ be as above but entering the Tantrakyma covariant derivative (eq. (10.2)). Assume internal gauge SSB (BEH) in the Tantrakyma vacuum so that the transverse, BRST–physical modes acquire a mass (cf. Definitions 9.252 and 9.348). On the realized weave configuration, isolate the spin–1 connection content via the spectral–island Riesz selector (Definitions 10.1, 10.6 and 10.7). Then:*

(i) Dark matter

A massive tantron *is a viable vector dark–matter candidate (misalignment or freeze–in), subject to the portal constraints summarized in Section 10.16 (WEP, kinetic mixing, spin–dependent torsionlike couplings, clocks, birefringence).*

(ii) Typed/TT safety

The tantron's projector commutes with the provenance/typed/triad structure and is TT–orthogonal (Propositions 10.9 and 10.10 and Definition 10.3), leaving the spin–2 graviton channel purely geometric.

(iii) Ledger separation

The tantron is not *a τ–kairos register; it resides in the* connection *ledger and affects $\Sigma_{\mu\nu}$ only through its influence on (Φ, Θ) within the constitutive mapping (eq. (10.101)).*

Clean origin and well–posed degrees of freedom. The *mition* and *tantron* are spin–1 excitations of the universal relational operator–bundle connection $\mathcal{A}_\mu$, not substrate quanta of $\hat{\mathbf{\Phi}}$ or $\hat{\mathbf{\Theta}}$ (Chapter 7: Section 6.6, Definition 6.192, and Remarks 6.198 and 6.200; their entry into dynamics is via eqs. (10.1) and (10.2)). After internal gauge SSB (BEH) the *transverse* modes of $\mathcal{A}_\mu$ become massive and BRST–physical, yielding the correct Proca–like propagating content (Chapter 10: Definitions 9.252, 9.258, 9.261 and 9.348, with BRST cleaning by $\mathsf{P}_{\mathrm{phys}}$). Thus each candidate possesses a clean origin, controllable gauge structure, and well–posed degrees of freedom.

Spectral isolation and robustness on the realized weave configuration. On the realized weave configuration $\mathcal{I}^{[\mathscr{P}]}$ the connection's transverse vector block admits band–limited *spectral islands* (Chapter 11: Definitions 10.1 and 10.6). Its spin–1 content is isolated by the commuting Riesz selectors $\sum_{k\in\mathfrak{S}_1} \frac{1}{2\pi i} \oint_{\Upsilon_k} (K^{(1)}_{\mathrm{vec}} - z)^{-1}\, dz$ (Definition 10.7), which preserve the provenance/typed/triad structure and render the readout Nielsen–flat and retune–stable (Propositions 10.9 and 10.10 and Corollary 10.11). This furnishes a rigorously defined, instrument–certified projector–field for mition/tantron phenomenology.

TT safety and separation from the graviton channel. Typed TT isolation is automatic because $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ acts on symmetric 2–tensors, whereas the mition/tantron live in the internal connection ledger; hence $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\, \mathsf{P}^{(1)}_{\mathrm{spec}} = 0$ (Definition 10.3). Consequently there are no extra GW polarizations, fully consistent with the weave's TT–null rules and with $\mathcal{P}^{(2)}[\hat{\tau}_{\mu\nu}] = 0$ for the τ channel (eq. (10.90)). The spin–2 graviton sector therefore remains purely geometric and TT–pure.

Natural cosmological production, compatible with constraints. Massive, weakly coupled spin–1 fields have standard cold–dark–matter production channels (coherent misalignment or freeze–in). In the present bookkeeping their effective "portals" map onto the consolidated bounds of Section 10.16 (e.g. LVK polarization tests for TT silence, MICROSCOPE WEP limits on long–range composition dependence, kinetic–mixing constraints, torsionlike spin couplings, clock–network searches for ultralight coherent backgrounds, birefringence limits, and mid–band atom–interferometry reach). Because the connection projectors commute with the typed/triad structure and are TT–null, viable parameter windows persist without spoiling GW polarizations or violating typed selection.

Ledger separation and constitutive consistency. The mition and tantron are *not* τ–kairos registers; the τ channel remains $0 \oplus 1$ from the Gaiaton and Chrono–Kymon only (Definitions 10.4, 10.13, 10.57 and 10.69). Their impact on the chronogeometric response $\Sigma_{\mu\nu} = g_{\mu\nu}[\Phi] + t_{\mu\nu}[\Phi, \Theta]$ is therefore *indirect*: it arises solely through how $\mathcal{A}_\mu$ modifies the Φ, Θ dynamics inside the constitutive map (Chapter 7: eq. (10.101) and Section 10.17.4). This preserves the formal separation of ledgers while allowing controlled backreaction through the established pipeline.

Summary. The **mition** (Gaiamiti–sourced) and **tantron** (Tantrakyma–sourced) combine: (i) a universal, algebraically anchored connection origin; (ii) massive, BRST–physical transverse

modes from BEH; (iii) typed/triad–robust spectral isolation on the realized weave configuration; (iv) TT–null behavior that safeguards the graviton sector; and (v) viable cosmological production within existing experimental bounds (Section 10.16). They are therefore credible dark–matter candidates within the weave, complementary to the τ–kairos narrative.

10.17 Geometric Block of the Unified Correlator

Same backbone, stricter readout. Chapter 7 introduces the *Unified Inter–Modal Correlator* $\hat{\mathbf{G}}_\Lambda$ as a 2×2 matrix of Fourfold two–point kernels (Section 6.1.1 and Definition 6.4, built from Definition 4.9). The Chapter 11 objects $\hat{\mathbf{G}}_\Lambda^{[\Phi\to\Phi]}$ and its triad/TT refinements (Definitions 10.157 and 10.158) are *not new correlators* but *typed projections* of this same backbone: first a type sandwich to extract the geometric block ($\Phi \to \Phi$), then a provenance split into $\{\text{spec}, \text{vac}, \text{corr}\}$ (Definition 10.2), and finally a background–covariant TT refinement with $\Pi_{\bar\Sigma}^{[\mathrm{TT}]}$ (Definition 10.3). Equation (10.83) asserts the purely kinematic compatibility with Chapter 7's sectoring: non–TT types (the τ–kairos scalar/vector families; Definition 10.4) are *TT–null* (Lemma 10.17), so the graviton readout extracted from $\Phi\Phi$ cannot be contaminated by τ content—consistent with the diagonal/off–diagonal semantics of $\hat{\mathbf{G}}_\Lambda$ in Definition 6.4.

What Chapter 11 adds (and where it leans on Chapter 7). The Chapter 7 construction is *kinematic and sector–agnostic*: it organizes correlations across (Φ, Θ) and across the Fourfold modes with parallel transport and relational covariance (Definition 4.9 and Section 6.1.1). Chapter 11 adds *instrumentation and provenance*: all operations are performed after Σ–realization and on the realized weave configuration $(\bar\Sigma, \mathcal{I}^{[\mathscr{I}]}$; Definitions 9.80 and 10.1), so index–raising and causal support are governed by $\bar\Sigma^{\mu\nu}$ and its relational Levi–Civita connection (eq. (6.62), Definition 6.7, and Remark 9.83). Within this setting, the typed insertion (Definition 10.157) and TT refinement (Definition 10.158) simply *specialize* the Chapter 7 correlator to the geometric, spin–2, TT channel with a triad–resolved ledger—moves that Chapter 7 did not yet prescribe. In short, Chapter 7 defines $\hat{\mathbf{G}}_\Lambda$; Chapter 11 fixes the realized weave configuration and then projects $\hat{\mathbf{G}}_\Lambda$ into $\Phi\Phi \to$triad$\to$TT in a way that is compatible with the same relational background.

Consequences for usage. Because Chapter 11's blocks are just Chapter 7's $\hat{\mathbf{G}}_\Lambda$ seen through (Φ-type, triad, TT) lenses, they inherit all upstream semantics of the diagonal/off–diagonal entries (geometry, matter, interaction; Definition 6.4) while becoming immediately eligible for Chapter 11 meters and T–L–O stamping (Definition 10.82 and Proposition 10.90). This is exactly what enables linearization of the geometric sector, graviton–channel isolation, and the statement of the chronogeometric Einstein equations in the natural $\bar\Sigma$–geometry (Section 11.7.3)—tasks that rely on the TT–refined $\Phi\Phi$ blocks (Definition 10.158) rather than on the raw, unspecialized matrix of Chapter 7.

Definition 10.157 (Geometric block: typed insertion and triad resolution)**.** By typed insertion (Definition 10.113), the *geometric* ($\Phi\Phi$) block of the unified inter–modal correlator is obtained

by sandwiching with the Φ-type idempotent $\Pi_{\Lambda}^{[\Phi]}$:

$$\boxed{\hat{\mathbf{G}}_{\Lambda}^{[\Phi\to\Phi]} := \Pi_{\Lambda}^{[\Phi]}\,\hat{\mathbf{G}}_{\Lambda}\,\Pi_{\Lambda}^{[\Phi]}.} \tag{10.80}$$

Resolving provenance with the triad $\{a\} = \{\text{spec}, \text{vac}, \text{corr}\}$ (Definition 10.2) and its idempotents $\Pi^{[a]}$ yields the orthogonal sum

$$\boxed{\hat{\mathbf{G}}_{\Lambda}^{[\Phi\to\Phi]} = \bigoplus_{a\in\{\text{spec},\text{vac},\text{corr}\}} \hat{\mathbf{G}}_{\Lambda;a}^{[\Phi\to\Phi]}, \qquad \hat{\mathbf{G}}_{\Lambda;a}^{[\Phi\to\Phi]} := \Pi^{[a]}\,\hat{\mathbf{G}}_{\Lambda}^{[\Phi\to\Phi]}\,\Pi^{[a]}.} \tag{10.81}$$

Definition 10.158 (Spin–2 TT refinement and TT–null complement)**.** Using the background–covariant TT projector $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ (Definition 10.3), each triad block admits a TT refinement

$$\boxed{\hat{\mathbf{G}}_{\Lambda;a}^{[\Phi\to\Phi;\mathrm{TT}]} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\hat{\mathbf{G}}_{\Lambda;a}^{[\Phi\to\Phi]}\,\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}, \qquad a\in\{\text{spec}, \text{vac}, \text{corr}\},} \tag{10.82}$$

while every *non–TT* typed family (the τ–kairos scalar/vector images; Definition 10.4) satisfies the TT–nullity relation (Lemma 10.17)

$$\boxed{\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\Pi^{[\chi]} = 0 \qquad (\chi \text{ a non–TT type}).} \tag{10.83}$$

What "admission–certified" means. After the typed insertion and triad split in Definition 10.157 (with TT refinement in Definition 10.158), each block $\hat{\mathbf{G}}_{\Lambda;a}^{[\Phi\to\Phi]}$ lives on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ and physical carrier $\mathcal{H}_{\text{phys}}$ (Definition 10.1). "Admission–certified'' here means that band/BRST hygiene and commutation properties hold *by construction*: the band tags and IR windows commute with P_{phys} and the typed/triad idempotents, so all sandwiched objects remain rig–admissible and gauge–safe (Proposition 10.10; see also the TT sandwich in Definition 10.3). Consequently, each provenance slice $a \in \{\text{spec}, \text{vac}, \text{corr}\}$ is a well–posed, type–resolved target for downstream diagnostics.

Why it is directly eligible for meters and T–L–O. Because admission is guaranteed, the law/state ledgers and calibrated visibilities may be evaluated *as is* on every triad block: compute $\mathbf{R}^{[\mathscr{S};\chi]}, \mathbf{S}^{[\mathscr{S};\chi]}$ and then $\Lambda_{\text{law}}, \Lambda_{\text{state}}$ via the cosine meters (Definitions 10.80 and 10.82), with finiteness and monotonicity ensured by Lemma 10.83 and Proposition 9.146. The *T–L–O stamping* then attaches the **T**ype (here, e.g., TT in the $\Phi\Phi$ block), the **L**ocation (bulk/boundary, per Definition 10.101), and the unique **O**nset $r^{\star}[\chi]$ defined by the canonical acceptance predicate (existence/uniqueness in Proposition 10.90 and Lemma 10.86, stability in Proposition 10.88). In short, once a block is formed as in Definitions 10.157 and 10.158, nothing further is required to run the meters and issue a T–L–O: the certification and commutation guarantees (Proposition 10.10) make these operations order–independent and audit–ready.

Physical interpretation (Apeiron = trace–typed scalar channel of geometry). On the realized weave configuration (Definition 10.1) and physical carrier, the relational geometric content of the ΦΦ block of the unified correlator is read via typed insertion (Definition 10.113) and the weave triad $\{a\} = \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$ (Definition 10.2). The connected entry equals the covariance of the fluctuation field (realized-layer recall from Chs. 7–10), hence typed projections of the ΦΦ block separate (i) the scalar *trace–typed* channel—called the *Apeiron* in what follows—from (ii) the spin–2 *TT* graviton channel. No new primitives are introduced; we only apply the Chapter 11 typed/triad machinery to the ΦΦ block of $\hat{\mathbf{G}}_\Lambda$ (Definition 6.4).

Remark 10.159 (Terminology: Apeiron as the ΦΦ trace–typed block)**.** We refer to the ΦΦ *trace–typed* scalar channel as the **Apeiron**. It is defined by typed insertion and triad resolution (no extra structure):

$$\boxed{\hat{\mathbf{G}}^{[\Phi\to\Phi;\mathrm{Tr}]}_{\Lambda;a} \;:=\; \Pi^{(a)}\,\Pi^{[\mathrm{Tr}]}\,\hat{\mathbf{G}}^{[\Phi\to\Phi]}_{\Lambda}\,\Pi^{[\mathrm{Tr}]}\,\Pi^{(a)}, \qquad a\in\{\mathrm{spec},\mathrm{vac},\mathrm{corr}\}.} \tag{10.84}$$

On admitted plateaux (admission & invariance in Definition 10.94 and Proposition 10.10), each block (10.84) is BRST/Nielsen-safe, cone-gated, and suitable for witnesses/meters and T–L–O execution (Proposition 10.90).

Remark 10.160 (Spin–2 readout and TT–null complement)**.** The observable graviton is the TT refinement of the same ΦΦ block using the background–covariant projector (Definition 10.3):

$$\boxed{\hat{\mathbf{G}}^{[\Phi\to\Phi;\mathrm{TT}]}_{\Lambda;a} \;:=\; \Pi^{[\mathrm{TT}]}_{\bar\Sigma}\,\hat{\mathbf{G}}^{[\Phi\to\Phi]}_{\Lambda;a}\,\Pi^{[\mathrm{TT}]}_{\bar\Sigma}, \qquad a\in\{\mathrm{spec},\mathrm{vac},\mathrm{corr}\}.} \tag{10.85}$$

Every non–TT typed family (τ–kairos scalars/vectors in Definition 10.4) is *TT–null*:

$$\boxed{\Pi^{[\mathrm{TT}]}_{\bar\Sigma}\,\Pi^{[\chi]} \;=\; 0 \qquad \text{for all non–TT types } [\chi] \text{ (cf. Lemma 10.17),}} \tag{10.86}$$

so (10.85) is strictly tensorial with no τ contamination; future–proofing and spectral updates proceed as in the weave (Ch. 11).

Definition 10.161 (Connected kernel as fluctuation covariance)**.** Let $\hat{\mathbf{\Phi}}(x)$ be the dynamical Gaiamiti field and write the realized background–fluctuation split (Chs. 9–10):

$$\boxed{\hat{\mathbf{\Phi}}(x) \;=\; \langle\hat{\mathbf{\Phi}}(x)\rangle_\circ\,\mathbb{I}_x \;+\; \delta\hat{\mathbf{\Phi}}(x), \qquad \langle\delta\hat{\mathbf{\Phi}}(x)\rangle_\circ = 0.} \tag{10.87}$$

Then the ΦΦ *connected* kernel equals the fluctuation covariance:

$$\begin{aligned}
\hat{G}_{\Phi\Phi}(x,x') &= \Big\langle \hat{\mathbf{\Phi}}(x)\,\hat{\mathbf{\Phi}}(x')\Big\rangle_\circ - \Big\langle\hat{\mathbf{\Phi}}(x)\Big\rangle_\circ\Big\langle\hat{\mathbf{\Phi}}(x')\Big\rangle_\circ && (10.88)\\
&= \Big\langle \Big(\langle\hat{\mathbf{\Phi}}(x)\rangle_\circ\mathbb{I}_x + \delta\hat{\mathbf{\Phi}}(x)\Big)\Big(\langle\hat{\mathbf{\Phi}}(x')\rangle_\circ\mathbb{I}_{x'} + \delta\hat{\mathbf{\Phi}}(x')\Big)\Big\rangle_\circ - \langle\hat{\mathbf{\Phi}}(x)\rangle_\circ\,\langle\hat{\mathbf{\Phi}}(x')\rangle_\circ\\
&= \Big\langle \delta\hat{\mathbf{\Phi}}(x)\,\delta\hat{\mathbf{\Phi}}(x')\Big\rangle_\circ. && (10.89)
\end{aligned}$$

Typed resolution then yields the scalar (Apeiron) and TT graviton correlator blocks by inserting $\Pi^{[\mathrm{Tr}]}$ and $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$, respectively, as in (10.84)–(10.85). Positivity and causal support follow from admission and cone–gating clauses on the realized weave configuration (Definition 10.94 and the cone statement referenced therein), and T–L–O stamping proceeds via Proposition 10.90 without double counting of polarizations thanks to (10.83).

Appropriateness for the Treatise: the Tau–Kairos sector. We work on the realized weave configuration and physical carrier (Definition 10.1), and use only the typed–insertion and triad machinery of Chapter 11 (Definitions 10.2 and 10.113). The coherence branch of the realized chronogeometry $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ is encoded by the *tau–kairos field* $\hat{\tau}_{\mu\nu}$ as fixed in the weave; its typed images furnish scalar/pseudoscalar/vector channels (Definition 10.4). By construction the tau–kairos sector is *TT–null*:

Definition 10.162 (Tau–Kairos sector and TT–nullity)**.** We work on the realized weave configuration and physical carrier (Definition 10.1) and use only the typed–insertion and triad machinery of Chapter 11 (Definitions 10.2 and 10.113). The coherence branch of the realized chronogeometry $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ is encoded by the *tau–kairos field* $\hat{\tau}_{\mu\nu}$ as fixed in the weave; its typed images furnish scalar/pseudoscalar/vector channels (Definition 10.4). By construction the tau–kairos sector is *TT–null*:

$$\boxed{\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\hat{\tau}_{\mu\nu} \;=\; 0, \qquad \text{equivalently} \qquad \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\Pi^{[\chi]} \;=\; 0 \;\; \text{for all non–TT types } [\chi],} \tag{10.90}$$

as stated in the weave TT–purity lemma (Lemma 10.17), with $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ the background–covariant TT projector (Definition 10.3).

Definition 10.163 (Typed correlator blocks for the tau–kairos sector)**.** Tau–kairos excitations are read from the $\Phi\Phi$ block of the unified correlator by *typed insertion* and *triad* resolution (Definitions 10.2 and 10.113). For each admissible type $[\chi] \in \{S^{(b)}, S^{(L)}, \tilde{S}^{(p)}, A^{(+)}, A^{(-)}\}$ and triad element $a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$,

$$\boxed{\hat{\mathbf{G}}^{[\Phi\to\Phi;\chi]}_{\Lambda;a} \;:=\; \Pi^{(a)}\,\Pi^{[\chi]}\,\hat{\mathbf{G}}^{[\Phi\to\Phi]}_{\Lambda}\,\Pi^{[\chi]}\,\Pi^{(a)}.} \tag{10.91}$$

These blocks are BRST/Nielsen–safe and band–admissible on the realized weave configuration (Proposition 10.10), and—by (10.90)—contain *no* spin–2 content.

Remark 10.164 (Background dependence and operational role)**.** On the diagnostic realized reference configuration, tau–kairos typed blocks modulate phase flow and coherence timing of relational processes. Their meters/witnesses and the parasymmetric T–L–O stamps are those already provided for the tau/CK families (see Propositions 10.22, 10.65 and 10.90); no new primitives are introduced here.

Allowed excitations (scalar/pseudoscalar/vector only). Under the TT–null constraint (10.90) and typed admission (Definition 10.94), the admissible tau–kairos excitations are:

1. *Scalar* τ–types: $S^{(b)}$ (bulk scalar) and $S^{(L)}$ (longitudinal scalar);

2. *Pseudoscalar* τ–type: $\tilde{S}^{(p)}$;
3. *Vector* (Chrono–Kymon) types: $A^{(\pm)}$ (helicity ± 1).

Longitudinal, non–radiative content is captured by the $S^{(L)}$ branch; the transverse propagating CK modes are the $A^{(\pm)}$ branches. TT projections of any of these vanish by (10.90).

Remark 10.165 (Complementarity with the graviton channel)**.** The graviton is obtained exclusively from the TT refinement of the $\Phi\Phi$ block,

$$\hat{\mathbf{G}}^{[\Phi\to\Phi;\mathrm{TT}]}_{\Lambda;a} := \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\, \hat{\mathbf{G}}^{[\Phi\to\Phi]}_{\Lambda;a}\, \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}},$$

and its T–L–O entry is stamped by the graviton rules of the weave (Proposition 10.35). By (10.90) and Lemma 10.17, tau–kairos excitations never contribute to $\mathrm{TLO}^{[\mathrm{TT}]}$; instead they receive their own $\mathrm{TLO}^{[\chi]}$ stamps in the scalar/pseudoscalar/vector families (Propositions 10.22 and 10.65).

Remark 10.166 (Nomenclature and conceptual role)**.** We use *tau–kairos tensor* for $\hat{\tau}_{\mu\nu}$ (noun) and *tau–kairos* (adjective) for the sectoral qualifiers. This emphasizes (i) the selection of opportune windows for coherent evolution (Onset in T–L–O), (ii) the absence of spin–2 carriage, and (iii) the complementarity with the chronometric channel: $\bar{g}_{\mu\nu}$ organizes measurable duration/extent, whereas $\hat{\tau}_{\mu\nu}$ organizes phase–suitable coherence, consistent with the typed/triad split used in (10.91) and the weave execution rules.

10.17.1 The Apeiron: Field–level vs. Particle–level Modes

We work on the realized weave configuration and physical carrier (Definition 10.1), using typed insertion and the weave triad only (Definitions 10.2 and 10.113). TT is always the background–covariant projector $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ (Definition 10.3); the τ–kairos families are TT–null (Lemma 10.17). The realized chronogeometry obeys the constitutive split $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ with linear response as in Ch. 10 (recall only; no redefinition).

Convention (Apeiron). We use *Apeiron* strictly as a label for:

(i) Field–level. The geometric fluctuation source $\delta\hat{\mathbf{\Phi}}$ from the realized split (10.87), whose connected $\Phi\Phi$ kernel equals the fluctuation covariance (10.89). It feeds the $\Phi\Phi$ block of the unified correlator (Definition 6.4) (Definition 10.157); typed resolution then produces its *trace* channel (through the $\bar{\Sigma}$–trace, cf. $\mathrm{tr}_{\bar{\Sigma}}$ in Definition 10.3 and the inner–product conventions in Definition 9.33) and the *TT* channel via the background–covariant TT projector (Definitions 10.3 and 10.158).

(ii) Particle–level (spin–0 radial). The *radial* scalar obtained from the Φ–sector radial direction of the physical Hessian (mass/pole extracted by Riesz selection on the scalar/radial generalized eigen-operator pair; Definition 10.7 and Proposition 10.149), admitted on the realized weave configuration (Definition 10.1) and within the Chapter 8 acceptance windows, and stamped as a trace–typed scalar under the typed weave layer (Definition 9.116).

Thus *Apeiron* (field) $\Rightarrow$ typed $\Phi\Phi$ blocks (Definition 10.157); *Apeiron* (particle) $\Rightarrow$ spin–0 radial scalar on $H^0(\hat{Q}_{\mathrm{BRST}})$ (the physical carrier $\mathcal{H}_{\mathrm{phys}}$, cf. Definition 10.1).

Typed ΦΦ blocks (trace and TT). By typed insertion with triad provenance (Definitions 10.2 and 10.113) applied to the geometric block (Definition 10.157),

$$\boxed{\hat{\mathbf{G}}^{[\Phi\to\Phi;\mathrm{Tr}]}_{\Lambda;a} := \Pi^{(a)}\,\Pi^{[\mathrm{Tr}]}\,\hat{\mathbf{G}}^{[\Phi\to\Phi]}_{\Lambda}\,\Pi^{[\mathrm{Tr}]}\,\Pi^{(a)}, \qquad \hat{\mathbf{G}}^{[\Phi\to\Phi;\mathrm{TT}]}_{\Lambda;a} := \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\hat{\mathbf{G}}^{[\Phi\to\Phi]}_{\Lambda;a}\,\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}},} \tag{10.92}$$

for $a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$. The first block encodes the *trace–typed* scalar (Apeiron, particle–level; trace with respect to $\bar{\Sigma}$, cf. Definitions 9.33 and 10.3); the second is the TT graviton block as in Definition 10.158. TT–nullity of non–TT types ensures no contamination from τ families (Lemma 10.17 and Definition 10.162).

Tangential–radial decomposition (field space). Let $\mathcal{M}_{\mathrm{vac}}$ be the Φ–vacuum manifold (Ch. 9/10). Any small fluctuation decomposes orthogonally as

$$\delta\Phi = \delta\Phi_{\mathrm{tang}} \oplus \delta\Phi_{\mathrm{radial}}, \qquad \delta\Phi_{\mathrm{tang}} \in T_{\Phi_0}\mathcal{M}_{\mathrm{vac}}, \quad \delta\Phi_{\mathrm{radial}} \perp T_{\Phi_0}\mathcal{M}_{\mathrm{vac}}.$$

The radial piece sources the *trace–typed* scalar (spin–0 Apeiron, mass from the radial Hessian on H^0); the tangential piece generates Goldstone–like directions.

Graviton as a tangential constitutive mode (TT readout). With the linear response of the constitutive map in Ch. 10 and τ–TT isolation (Lemma 10.17),

$$\boxed{\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{g}_{\mu\nu} \;=\; \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\Big[\big(D_\Phi \bar{g}_{\mu\nu}\big)\cdot\delta\hat{\Phi}_{\mathrm{tang}}\Big], \qquad \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\Big[\big(D_\Phi \bar{g}_{\mu\nu}\big)\cdot\delta\hat{\Phi}_{\mathrm{radial}}\Big] = 0.}\tag{10.93}$$

Hence the TT graviton channel is sourced *solely* by *tangential* Φ–variations, whereas the radial scalar reports in the trace–typed block (10.92) (no double counting).

T–L–O stamping and non–overcounting (weave execution). On admitted plateaux (Definition 10.94 and Proposition 10.10), the scalar radial Apeiron receives the $\mathrm{TLO}^{[\mathrm{Tr}]}$ stamp, while the graviton receives $\mathrm{TLO}^{[\mathrm{TT}]}$ (Proposition 10.35). Tau/CK families get $\mathrm{TLO}^{[\chi]}$ in their own typed branches and never contribute to $\mathrm{TLO}^{[\mathrm{TT}]}$ by TT–nullity (Lemma 10.17); the weave's witness/meter rules apply verbatim.

Terminology. (i) *"Spin–2 Apeiron"* ≡ *graviton* (the TT, tangential mode of the Φ–sector as in (10.93)). (ii) *"Spin–0 Apeiron"* = trace–typed *radial* scalar (mass from the radial block of $\mathbb{H}_{\mathrm{phys}}$). (iii) No "spin–1 Apeiron": vector temporal quanta belong to the τ/CK types and are TT–null (weave, Lemma 10.17).

Operational summary. Field–level Apeiron $\delta\hat{\mathbf{\Phi}}$ feeds the typed ΦΦ blocks; particle–level Apeiron is the trace–typed radial scalar (spin–0); the graviton is the TT projection of the metric response to *tangential* Φ–variations (10.93). The Chapter 11 pipeline (typed insertion, triad, admission, T–L–O) cleanly separates these channels—BRST/Nielsen safe, cone–preserving, and free of TT contamination from τ families.

10.17.2 The Chronogeometric Excitations

Program. On the realized weave configuration and physical carrier (Definition 10.1), the excitation content of gravity is fixed by: (i) a *kinematic* typed split of the unified chronogeometric field into metric and tau–kairos channels using the Chapter 11 projectors (Definitions 10.3, 10.4 and 10.113); and (ii) a *constitutive* (field–space) realization via the substrate response recalled from Ch. 10 (no new primitives).

Unified chronogeometric operator (recall). We assemble the operator–valued chronogeometry as the sum of its geometric and kairotic pieces and define its *realized* background by the Σ–realization map (Ch. 10: Definition 9.80) on the realized weave configuration $\mathcal{I}^{[\mathcal{J}]}$ (Definition 10.1); expectations $\langle\cdot\rangle_\circ$ are taken on $\mathcal{H}_{\rm phys}$. Thus

$$\hat{\Sigma}_{\mu\nu}(x) = \hat{g}_{\mu\nu}(x) + \hat{t}_{\mu\nu}(x), \qquad \bar{\Sigma}_{\mu\nu}(x) = \langle\hat{\Sigma}_{\mu\nu}(x)\rangle_\circ = \bar{g}_{\mu\nu}(x) + \bar{t}_{\mu\nu}(x), \tag{10.94}$$

$$\hat{\Sigma}_{\mu\nu}(x) = \bar{\Sigma}_{\mu\nu}(x) + \delta\hat{\Sigma}_{\mu\nu}(x), \qquad \delta\hat{\Sigma}_{\mu\nu} = \delta\hat{g}_{\mu\nu} + \delta\hat{t}_{\mu\nu}. \tag{10.95}$$

Here $\bar{\Sigma}_{\mu\nu}$ determines the relational Levi–Civita connection used throughout (Definition 6.7) and its inverse $\bar{\Sigma}^{\mu\nu}$ raises indices in all TT/typed refinements (cf. eq. (6.62) and Definition 10.3). In particular, the TT graviton readout uses $\delta\hat{g}_{\mu\nu}$ while $\delta\hat{t}_{\mu\nu}$ is TT–null by the weave purity statements (Definitions 10.162 and 10.167 and Lemma 10.17).

Kinematic TT split: graviton and TT–null tau–kairos

Definition 10.167 (TT graviton and TT–null tau–kairos (weave projectors))**.** The *graviton field* is the TT (spin–2) metric perturbation in the $\Phi\Phi$ sector, defined via the graviton projector of Definition 10.25 and read in the $\bar{\Sigma}$–background:

$$\boxed{\hat{h}_{\mu\nu} \;:=\; \mathcal{P}^{(2)}_{\rm grav}\,\delta\hat{g}_{\mu\nu} \quad \text{and, kinematically,} \quad \hat{h}_{\mu\nu} = \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{g}_{\mu\nu} \;\; (Definition\ 10.3),} \tag{10.96}$$

where $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ is the background–covariant TT projector.

The tau–kairos fluctuation $\hat{\tau}_{\mu\nu} := \delta\hat{t}_{\mu\nu}$ is *TT–null*; equivalently,

$$\boxed{\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\hat{\tau}_{\mu\nu} = 0 \quad\Longleftrightarrow\quad \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\Pi^{[\chi]} = 0 \;\text{ for all non–TT types } [\chi],} \tag{10.97}$$

as recorded in the weave TT–purity statements (Definition 10.162 and Lemma 10.17). Hence $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{\Sigma}_{\mu\nu} = \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{g}_{\mu\nu}$, consistent with $\delta\hat{\Sigma}_{\mu\nu} = \delta\hat{g}_{\mu\nu} + \delta\hat{t}_{\mu\nu}$.

Lemma 10.168 (Typed TT content is purely metric)**.** *With (10.97), no TT content arises from $\hat{\tau}_{\mu\nu}$; all TT strength is in the metric channel:* $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{\Sigma} = \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{g}$.

Typed TT correlator block. For each triad element $a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$ (Definition 10.2), the TT block of the $\Phi\Phi$ correlator is

$$\boxed{\hat{\mathbf{G}}^{[\Phi\to\Phi;\mathrm{TT}]}_{\Lambda;a} \;:=\; \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\hat{\mathbf{G}}^{[\Phi\to\Phi]}_{\Lambda;a}\,\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}, \qquad \hat{\mathbf{G}}^{[\Phi\to\Phi]}_{\Lambda;a} := \Pi^{(a)}\hat{\mathbf{G}}^{[\Phi\to\Phi]}_{\Lambda}\Pi^{(a)}.} \tag{10.98}$$

These are BRST/Nielsen–safe and band–admissible on the realized weave configuration (Proposition 10.10).

Execution: stamps and separation. On admitted plateaux (Definition 10.94), the TT graviton receives the $\mathrm{TLO}^{[\mathrm{TT}]}$ stamp (Proposition 10.35); tau–kairos families receive $\mathrm{TLO}^{[\chi]}$ in their scalar/pseudoscalar/vector branches and never contribute to TT by (10.97). Typed/triad provenance is carried by the index a in (10.98), ensuring non–overcounting and realized weave configuration invariance.

Typed excitations and T–L–O stamping

Typed families (recall; no new primitives). We use the Chapter 11 typed projectors and triad on the realized weave configuration (Definitions 10.1, 10.2 and 10.113). The geometry types are: TT (spin–2) via $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ (Definition 10.3) and the trace–typed scalar "Tr"; the tau–kairos (temporal/coherence) families are the scalar/pseudoscalar/vector images in Definition 10.4. Typed $\Phi\Phi$ correlator blocks with provenance $a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$ are

$$\boxed{\hat{\mathbf{G}}_{\Lambda;a}^{[\Phi\to\Phi;\chi]} := \Pi^{(a)}\,\Pi^{[\chi]}\,\hat{\mathbf{G}}_{\Lambda}^{[\Phi\to\Phi]}\,\Pi^{[\chi]}\,\Pi^{(a)}.} \tag{10.99}$$

Theorem 10.169 (Excitation content of gravity on the realized weave configuration)**.** *On admitted plateaux (Definition 10.94) and for admitted types (Definition 9.116):*

(i) Spin–2 (TT, graviton)

The unique spin–2 excitation is the TT metric mode

$$\hat{h}_{\mu\nu} \;=\; \mathcal{P}^{(2)}_{\mathrm{grav}}\,\delta\hat{g}_{\mu\nu} \;=\; \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}_{\mu\nu},$$

i.e. the graviton field defined by the graviton projector (Definition 10.25) and, kinematically, by the background–covariant TT projector (Definitions 10.3 and 10.167); its stamp is $\mathrm{TLO}^{[\mathrm{TT}]}$ *(Proposition 10.35).*

(ii) Tau–kairos (TT–null)

The tau–kairos sector is TT–null (Definition 10.162 and Lemma 10.17); it supports only scalar/pseudoscalar/vector families (including CK vectors) as in Definition 10.4. Their stamps are $\mathrm{TLO}^{[\chi]}$ *in the corresponding typed branches (Propositions 10.22 and 10.65).*

(iii) No overcounting)

Orthogonality holds: $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\Pi^{[\chi]} = 0$ *for all non–TT* $[\chi]$ *(Lemmas 10.17 and 10.59). Hence,*

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{\Sigma}_{\mu\nu} \;=\; \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}_{\mu\nu},$$

by the fluctuation split $\delta\hat{\Sigma}_{\mu\nu} = \delta\hat{g}_{\mu\nu} + \delta\hat{t}_{\mu\nu}$ *(eq. (10.95)) and TT–nullity of* $\delta\hat{t}_{\mu\nu}$ *(eq. (10.97)).*

Proof. (1) *Spin–2 uniqueness.* On the realized weave configuration (Definition 10.1) the typed TT projector $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ isolates the spin–2 subspace of symmetric rank–2 tensors (Definition 10.3). The graviton field on $\mathcal{H}_{\mathrm{phys}}$ is, by definition, $\hat{h}_{\mu\nu} = \mathcal{P}^{(2)}_{\mathrm{grav}}\delta\hat{g}_{\mu\nu}$ (Definition 10.25); kinematically this equals $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\delta\hat{g}_{\mu\nu}$ (Definition 10.167). Typed/triad commutations guarantee that no other typed family carries spin–2 TT support (Proposition 10.9), hence uniqueness.

(2) *Tau–kairos TT–nullity.* By the weave purity statements, $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\hat{\tau}_{\mu\nu} = 0$ (Definition 10.162, Lemma 10.17, and eq. (10.97)). Therefore the τ–block supports only the scalar/pseudoscalar/vector types listed in Definition 10.4; their TLO stamps are issued in the corresponding typed branches (Propositions 10.22 and 10.65) on admitted plateaux (Definition 10.94).

(3) *No overcounting.* Using the fluctuation decomposition $\delta\hat{\Sigma}_{\mu\nu} = \delta\hat{g}_{\mu\nu} + \delta\hat{t}_{\mu\nu}$ (eq. (10.95)) and TT–nullity of $\delta\hat{t}_{\mu\nu}$ (eq. (10.97)), we obtain $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\delta\hat{\Sigma}_{\mu\nu} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\delta\hat{g}_{\mu\nu}$. Orthogonality $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\Pi^{[\chi]} = 0$ for non–TT $[\chi]$ (Lemmas 10.17 and 10.59) forbids double–counting of TT strength across typed channels. Combined with (1) this matches the TT isolation in Lemma 10.172. □

Meaning of Theorem 10.169. The theorem states, in operational terms, that once we work on an *admitted plateau* with the realized weave configuration (Definitions 10.1 and 10.94), the *only* gravitational signal that can inhabit the spin–2 channel is the background–covariant TT graviton field $\hat{h}_{\mu\nu}$ (Definitions 10.25 and 10.167). All non–TT content—the entire τ–kairos sector, including its scalars, pseudoscalar, and Chrono–Kymon vectors (Definition 10.4)—is *TT–null* (Definition 10.162 and Lemma 10.17). Consequently, the TT projection of the full chronogeometric fluctuation equals the TT projection of the metric fluctuation alone,

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{\Sigma}_{\mu\nu} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}_{\mu\nu},$$

so no scalar/vector leakage can masquerade as a graviton (eqs. (10.95) and (10.97) and Lemma 10.172).

Why this matters. Practically, the theorem gives a *clean accounting rule* for reports and audits: TT claims (spin–2 radiation) are stamped only in the TT type ($\mathrm{TLO}^{[\mathrm{TT}]}$; Proposition 10.35), while τ–kairos claims are stamped in their own typed branches ($\mathrm{TLO}^{[\chi]}$; Propositions 10.22 and 10.65). Orthogonality $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\Pi^{[\chi]} = 0$ for non–TT $[\chi]$ prevents overcounting across channels (Lemmas 10.17 and 10.59), ensuring that meters/visibilities and onsets attach to the *right* physical content (Definition 10.82 and Proposition 10.88). In short: the theorem guarantees *uniqueness and purity* of the graviton readout and a consistent separation of scalar/vector structure, all under identical, rig–certified measurement conditions.

Corollary 10.170 (Helicity register and stability)**.** *The TT channel carries exactly the two tensor helicities $\lambda = \pm 2$ (optionally resolved in the weave's graviton register for reporting; cf. Definitions 10.3 and 10.25 and Proposition 10.35). All scalar/pseudoscalar/vector channels are TT–orthogonal and therefore never contribute to* $\mathrm{TLO}^{[\mathrm{TT}]}$ *(Lemmas 10.17, 10.59 and 10.172). Moreover, this separation is* stable*: typed/triad idempotents commute on the realized weave configuration (Proposition 10.9 and Definition 10.1), meter monotonicity and onset uniqueness hold under admissible IR tightening (Propositions 9.146 and 10.88 and Lemma 10.86), and additional helicity–2 spectral branches—if physically realized—are incorporated by the spin–2 Riesz update without altering the TT scaffold (future–proofing; cf. Definitions 10.7 and 10.25).*

Remark 10.171 (Operational summary)**.** Typed resolution of the geometric block together with TT refinement—see the typed $\Phi\Phi$ constructions in Definitions 10.157 and 10.158 (and the block

formulas such as (10.92) and (10.82))—combined with band/BRST hygiene on the realized weave configuration (Definition 10.1 and Proposition 10.10) yields $\mathrm{TLO}^{[\chi]}$ stamps (Type/Location/Onset) that are BRST/Nielsen–safe. Concretely, one evaluates the law/state ledgers and calibrated visibilities (Definitions 10.80 and 10.82), obtains finite meters (Lemma 10.83), and reads a unique onset $r^\star[\chi]$ with stability radius (Lemma 10.86 and Proposition 10.88). Orthogonality and selection rules (Lemmas 10.17 and 10.59 and Proposition 10.126) ensure that $\mathrm{TLO}^{[\mathrm{TT}]}$ registers only helicity–2 content extracted by $\Pi_{\bar\Sigma}^{[\mathrm{TT}]}$ (Definitions 10.3 and 10.25), while non–TT types are reported in their own $\mathrm{TLO}^{[\chi]}$ branches without leakage into the graviton channel (Theorem 10.169).

Excitation operators and TT selection (kinematics)

Kinematic channels (recall). With $\hat\Sigma = \bar\Sigma + \delta\hat\Sigma$ (eq. (10.95)), set

$$\boxed{\hat h_{\mu\nu} := \delta\hat g_{\mu\nu}, \qquad \hat\tau_{\mu\nu} := \delta\hat t_{\mu\nu}.} \tag{10.100}$$

Lemma 10.172 (TT isolation and graviton operator)**.** *Tau–kairos is TT–null (Definition 10.162 and Lemma 10.17), so $\Pi_{\bar\Sigma}^{[\mathrm{TT}]}\hat\tau_{\mu\nu} = 0$ (eqs. (10.90) and (10.97)). Hence the TT block of the chronogeometric perturbation is purely metric and coincides with the graviton field of Definitions 10.25 and 10.167:*

$$\hat h_{\mu\nu} \;=\; \mathcal{P}^{(2)}_{\mathrm{grav}}\,\delta\hat g_{\mu\nu} \;=\; \Pi_{\bar\Sigma}^{[\mathrm{TT}]}\,\delta\hat g_{\mu\nu} \;=\; \Pi_{\bar\Sigma}^{[\mathrm{TT}]}\,\delta\hat\Sigma_{\mu\nu},$$

where $\Pi_{\bar\Sigma}^{[\mathrm{TT}]}$ is the background–covariant TT projector (Definition 10.3) and $\delta\hat\Sigma_{\mu\nu} = \delta\hat g_{\mu\nu} + \delta\hat t_{\mu\nu}$ (eq. (10.95)).

Proof. By TT–nullity of tau–kairos ($\Pi_{\bar\Sigma}^{[\mathrm{TT}]}\hat\tau_{\mu\nu} = 0$; Definition 10.162 and Lemma 10.17), and the split $\delta\hat\Sigma_{\mu\nu} = \delta\hat g_{\mu\nu} + \delta\hat t_{\mu\nu}$ (eq. (10.95)), we have $\Pi_{\bar\Sigma}^{[\mathrm{TT}]}\delta\hat\Sigma_{\mu\nu} = \Pi_{\bar\Sigma}^{[\mathrm{TT}]}\delta\hat g_{\mu\nu}$. By Definitions 10.3 and 10.167, the kinematic TT projection of $\delta\hat g_{\mu\nu}$ equals the graviton field, and Definition 10.25 shows the same object is obtained by the full graviton projector $\mathcal{P}^{(2)}_{\mathrm{grav}}$. This proves the chain of equalities. □

Constitutive (field–space) realization

Recall the constitutive split $\Sigma_{\mu\nu}[\Phi,\Theta] = g_{\mu\nu}[\Phi] + t_{\mu\nu}[\Phi,\Theta]$ with linear response

$$\boxed{\delta\Sigma_{\mu\nu} = (D_\Phi g_{\mu\nu})\cdot\delta\Phi + (D_\Phi t_{\mu\nu})\cdot\delta\Phi + (D_\Theta t_{\mu\nu})\cdot\delta\Theta.} \tag{10.101}$$

Proposition 10.173 (TT selection at constitutive level)**.** *With TT isolation of tau–kairos (Definition 10.162 and Lemma 10.17),*

$$\Pi_{\bar\Sigma}^{[\mathrm{TT}]}\,\delta\Sigma_{\mu\nu} \;=\; \Pi_{\bar\Sigma}^{[\mathrm{TT}]}\,\delta g_{\mu\nu} \;=\; \Big(\Pi_{\bar\Sigma}^{[\mathrm{TT}]}\circ D_\Phi\bar g\Big)\cdot\delta\Phi.$$

Decomposing $\delta\Phi = \delta\Phi_{\mathrm{tang}} \oplus \delta\Phi_{\mathrm{radial}}$ (Chs. 9–10; NG/tangent vs. radial directions of the vacuum manifold, cf. Definition 6.61 and Proposition 9.197) gives

$$\Pi_{\bar\Sigma}^{[\mathrm{TT}]}\Big[(D_\Phi\bar g)\cdot\delta\Phi_{\mathrm{tang}}\Big], \qquad \Pi_{\bar\Sigma}^{[\mathrm{TT}]}\Big[(D_\Phi\bar g)\cdot\delta\Phi_{\mathrm{radial}}\Big] = 0,$$

so TT support is tangential, while the radial part reports only in the trace sector.

Proof. First, by the fluctuation split $\delta\hat{\Sigma}_{\mu\nu} = \delta\hat{g}_{\mu\nu} + \delta\hat{t}_{\mu\nu}$ (eq. (10.95)) and TT–nullity of tau–kairos ($\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\hat{\tau}_{\mu\nu} = 0$; Definition 10.162, Lemma 10.17, and eq. (10.97)), we have $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\Sigma_{\mu\nu} = \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta g_{\mu\nu}$. Second, on the realized weave configuration (Definition 10.1) after Σ–realization (Definition 9.80), the realized background metric is a functional of the Gaiamiti field, $\bar{g} = \bar{g}[\Phi]$. Linear response in the Φ–direction yields $\delta g_{\mu\nu} = (D_\Phi \bar{g})_{\mu\nu}\cdot\delta\Phi$, hence $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta g_{\mu\nu} = \left(\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\circ D_\Phi\bar{g}\right)_{\mu\nu}\cdot\delta\Phi$, which proves the first display.

For the tangential/radial split, decompose $\delta\Phi = \delta\Phi_{\mathrm{tang}}\oplus\delta\Phi_{\mathrm{radial}}$ along the vacuum manifold $\mathcal{M}_{\mathrm{vac}}$ (tangent = NG directions; radial = orthogonal amplitude directions; cf. Chs. 9–10 and Proposition 9.197). In the NG (tangent) directions, $D_\Phi\bar{g}\cdot\delta\Phi_{\mathrm{tang}}$ furnishes the massless spin–2 (Goldstone) deformation, which is TT and therefore survives TT projection (Proposition 9.197 and Definitions 10.3 and 10.167). In the radial directions, $D_\Phi\bar{g}\cdot\delta\Phi_{\mathrm{radial}}$ produces only $\bar{\Sigma}$–trace and longitudinal pieces (the scalar "apeiron" response). The background–covariant TT projector $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ annihilates both pure traces and longitudinal (York) parts (Definition 10.3), hence $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\left[(D_\Phi\bar{g})\cdot\delta\Phi_{\mathrm{radial}}\right] = 0$. This establishes the two claims and the stated interpretation. □

Remark 10.174 (Chrono–Kymon has no extra TT polarizations)**.** The Chrono–Kymon vectors (typed idempotents $A^{(\pm)}$ or, equivalently, $\Pi^{[K^{(\tau)}_{\mathrm{T/L}}]}$ in the CK basis; Definitions 10.4 and 10.57) satisfy TT–nullity, $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\Pi^{[\chi]} = 0$ for all non–TT $[\chi]$, in particular for CK types (Lemmas 10.17 and 10.59). Consequently they *do not* add gravitational–wave tensor polarizations and are reported only in their own typed TLO branches $\mathrm{TLO}^{[\chi]}$ (Propositions 10.22 and 10.65)—never in $\mathrm{TLO}^{[\mathrm{TT}]}$ (Proposition 10.35 and Theorem 10.169).

Particles and sourcing. The *graviton* is the TT, Goldstone–tangential excitation of the metric channel, $\hat{\gamma}_{\mu\nu} = \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{g}_{\mu\nu}$ (Definition 10.3, Section 10.17.1, and eq. (10.93)), while the Gaiamiti sector also supplies a *spin–0 Apeiron* at the particle level (the trace–typed *radial* scalar read from the $\Phi\Phi$ block; mass from the radial block of $\mathbb{H}_{\mathrm{phys}}$ per Ch. 8). The temporal/coherence (tau–kairos) tensor carries only scalar/pseudoscalar/vector structure (no TT): its vector quanta are the Tantrakyma $U(1)_C$ *Chrono–Kymons* (CK) (Definition 10.4 and Propositions 10.22 and 10.65). We record this via the *typed sourcing* furnished by the weave, without introducing new maps.

Remark 10.175 (Typed τ–kairos sourcing (weave form; no new primitives))**.** By the τ–decomposition (Definition 10.4),

$$\boxed{\hat{\tau}_{\mu\nu} \;=\; S^{(b)}_{\mu\nu}[\hat{\varphi}_b] \;\oplus\; S^{(L)}_{\mu\nu}[\hat{\varphi}_L] \;\oplus\; \tilde{S}^{(p)}_{\mu\nu}[\tilde{\varphi}] \;\oplus\; \sum_{\lambda=\pm} A^{(\lambda)}_{\mu\nu}\left[\hat{K}^{(\tau)}_{\lambda}\right], \qquad \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\hat{\tau}_{\mu\nu} = 0,} \tag{10.102}$$

where $S^{(b)}, S^{(L)}, \tilde{S}^{(p)}$ are the scalar/pseudoscalar typed images and $A^{(\pm)}$ the CK helicity vectors. Equation (10.102) *is* the scalar/vector sourcing of $\hat{\tau}_{\mu\nu}$ in Chapter 11 notation and already encodes TT–nullity (Lemmas 10.17 and 10.59).

Remark 10.176 (Typed reading and T–L–O stamps)**.** Using typed insertion with triad provenance (Definitions 10.2 and 10.113), the $\Phi\Phi$ blocks

$$\hat{\mathbf{G}}^{[\Phi\to\Phi;\chi]}_{\Lambda;a} := \Pi^{(a)}\Pi^{[\chi]}\,\hat{\mathbf{G}}^{[\Phi\to\Phi]}_{\Lambda}\,\Pi^{[\chi]}\Pi^{(a)}, \qquad a\in\{\mathrm{spec},\mathrm{vac},\mathrm{corr}\},$$

are BRST/Nielsen–safe and band–admissible (Proposition 10.10). For $\chi = \mathrm{Tr}$ (Apeiron scalar) and $\chi \in \{S^{(b)}, S^{(L)}, \tilde{S}^{(p)}, A^{(\pm)}\}$ (tau/CK families), the weave stamps $\mathrm{TLO}^{[\chi]}$ via Propositions 10.22 and 10.65; for $\chi = \mathrm{TT}$, the graviton uses Proposition 10.35. TT–orthogonality is automatic: $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\Pi^{[\chi]} = 0$ for all non–TT $[\chi]$ (Lemmas 10.17 and 10.59).

Corollary 10.177 (No gravitational–wave polarizations from tau–kairos)**.** *From* (10.102) *and TT–nullity, the tau–kairos sector contributes* no *TT (GW) polarizations. All spin–2 radiative content resides in the metric TT channel* [TT] *with two tensor helicities (see Definition 10.3 and Proposition 10.35).*

10.17.3 Excitation Channels and the Role of the Tau–Kairos

Program (typed split + constitutive recall). On the realized weave configuration (Definition 10.1) we resolve observable fluctuations of $\hat{\Sigma}_{\mu\nu} = \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu}$ by the Chapter 11 typed projectors and triad (Definitions 10.2 to 10.4 and 10.113). The constitutive identity $\Sigma_{\mu\nu}[\Phi, \Theta] = g_{\mu\nu}[\Phi] + t_{\mu\nu}[\Phi, \Theta]$ and its linear response are recalled from Ch. 10 (no new primitives).

Metric (Gaiamiti) channel — TT (spin–2) only. The unique radiative tensor excitation is the *graviton*, the TT projection of metric fluctuations:

$$\boxed{\hat{h}_{\mu\nu} \;:=\; \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}_{\mu\nu}, \qquad \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \text{ as in } \textit{Definition } 10.3.} \tag{10.103}$$

TT exhausts the tensor content of $\delta\hat{\Sigma}$: $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{\Sigma} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}$ (cf. TT–nullity of τ below). Field–space wise, the TT mode is sourced by *tangential* (Goldstone-like) Φ–variations, whereas the orthogonal radial direction reports in the trace–typed scalar (Apeiron) (Section 10.17.1 and eq. (10.93)).

Temporal/coherence (tau–kairos) channel — non–TT (spin–0/1 only). By construction the tau–kairos tensor is TT–null (Lemma 10.17):

$$\boxed{\hat{\tau}_{\mu\nu} := \delta\hat{t}_{\mu\nu}, \qquad \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\hat{\tau}_{\mu\nu} = 0, \qquad \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\Pi^{[\chi]} = 0 \;\text{ for all non–TT } [\chi].} \tag{10.104}$$

Its typed decomposition (τ-kairos) is

$$\hat{\tau}_{\mu\nu} = S^{(b)}_{\mu\nu}[\hat{\varphi}_b] \;\oplus\; S^{(L)}_{\mu\nu}[\hat{\varphi}_L] \;\oplus\; \tilde{S}^{(p)}_{\mu\nu}[\tilde{\varphi}] \;\oplus\; \sum_{\lambda=\pm} A^{(\lambda)}_{\mu\nu}\big[\hat{K}^{(\tau)}_{\lambda}\big],$$

with $A^{(\pm)}$ the CK vector helicities. The CK field strength $\hat{\mathcal{F}}^{(\tau)}_{\mu\nu} := 2\bar{\nabla}_{[\mu}\hat{K}^{(\tau)}_{\nu]}$ is the natural constitutive input for antisymmetric/phase–transport features of $t_{\mu\nu}$ (weave usage).

Scalar sector of Gaiamiti ("Apeiron"). As in Section 10.17.1, "Apeiron'' has (i) a *field–level* sense: the source $\delta\hat{\mathbf{\Phi}}$ feeding typed $\Phi\Phi$ blocks; and (ii) a *particle–level* sense: the trace–typed *radial* scalar (spin–0) with mass from the radial block of $\mathbb{H}_{\mathrm{phys}}$. Nomenclature policy: the spin–2 collective is the TT graviton; no spin–1 register exists in the Φ–sector— spin–1 belongs to CK in the τ channel (Definition 10.4).

Putting it together (no extra TT modes). Typed separation and TT–nullity imply: (i) all propagating *tensor* radiation sits in the TT graviton ($\lambda = \pm 2$); (ii) the tau–kairos supports only scalar/pseudoscalar/vector responses. Non–overlap is guaranteed by $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\Pi^{[\chi]} = 0$ for all non–TT $[\chi]$ (Lemmas 10.17 and 10.59) and by realized weave configuration admissibility/invariance (Definition 10.94 and Proposition 10.10).

On the scalar contribution to $\hat{\tau}_{\mu\nu}$. Although the Apeiron originates in Φ, the temporal tensor $t_{\mu\nu}[\Phi, \Theta]$ is a *constitutive* object depending on (Φ, Θ). Thus the scalar piece of $\hat{\tau}_{\mu\nu}$ is simply the Apeiron as seen through the temporal channel, preserving TT–nullity and adding no TT content (compatible with (10.25)–(10.104)).

Remarks on dynamics (i) The TT (spin–2) sector governed by $\hat{h}_{\mu\nu}$ is dynamically closed: off–diagonal metric–temporal mixing is non–TT, and the TT block obeys a hyperbolic wave equation obtained by TT–projecting the linearized field equations (Definition 10.3 and TT–purity).
(ii) The temporal/coherence sector influences observations *indirectly* via $\bar{t}_{\mu\nu}$ (background transport, dissipation, KMS/thermodynamic data) and through scalar/vector perturbations in cosmological or condensed regimes, but never by introducing additional GW polarizations (Propositions 10.22 and 10.65).

Summary. The unified fluctuation $\delta\hat{\Sigma}_{\mu\nu} = \delta\hat{g}_{\mu\nu} + \delta\hat{t}_{\mu\nu}$ separates cleanly into (a) a TT, spin–2 *graviton* sector $\hat{h}_{\mu\nu} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}_{\mu\nu}$ (eq. (10.103); equivalently $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{\Sigma} = \hat{h}$), sourced by tangential Φ–variations, and (b) a non–TT *tau–kairos* sector $\hat{\tau}_{\mu\nu}$ populated by scalar/pseudoscalar/vector (CK) channels (eq. (10.25)). This realizes, in a single operator framework, the coexistence of geometric radiation (two tensor polarizations) with temporal/coherence dynamics, while preserving TT purity and typed/triad provenance throughout the weave pipeline (Propositions 10.10, 10.22, 10.35 and 10.65).

10.17.4 Observable (Kinematic) vs. Constitutive (Microscopic) Descriptions

On the realized weave configuration (Definition 10.1) there are two mutually consistent descriptions of chronogeometry: (i) a *formal/observable* (kinematic) split that reads metric and tau–kairos channels by the Chapter 11 typed projectors; and (ii) a *constitutive/microscopic* realization through the substrate fields (Φ, Θ). Using the *same* typed projectors and admission gate in both views yields identical Type–Location–Onset outcomes (Definitions 10.3, 10.4, 10.94 and 10.113 and Propositions 10.22 and 10.35).

Formal/observable split (recall; no new projectors). Write the operator–valued chronogeometry and its background/fluctuation split as in (10.94)–(10.95):

$$\hat{\Sigma}_{\mu\nu} = \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu}, \qquad \delta\hat{\Sigma}_{\mu\nu} = \delta\hat{g}_{\mu\nu} + \delta\hat{t}_{\mu\nu}.$$

The TT (spin–2) content is isolated by the background–covariant projector $\Pi_{\bar\Sigma}^{[\mathrm{TT}]}$ (Definition 10.3), and tau–kairos is TT–null (Lemma 10.17):

$$\boxed{\hat h_{\mu\nu} := \Pi_{\bar\Sigma}^{[\mathrm{TT}]}\,\delta\hat g_{\mu\nu}, \qquad \Pi_{\bar\Sigma}^{[\mathrm{TT}]}\,\hat\tau_{\mu\nu} = 0, \qquad \Pi_{\bar\Sigma}^{[\mathrm{TT}]}\,\delta\hat\Sigma_{\mu\nu} = \Pi_{\bar\Sigma}^{[\mathrm{TT}]}\,\delta\hat g_{\mu\nu}.} \tag{10.105}$$

Constitutive map (recall from Ch. 10). Chronogeometry is realized by the constitutive identity

$$\Sigma_{\mu\nu}[\Phi,\Theta] = g_{\mu\nu}[\Phi] + t_{\mu\nu}[\Phi,\Theta],$$

with linear response about $(\bar\Phi,\bar\Theta)$ given by (10.101):

$$\delta\Sigma_{\mu\nu} = (D_\Phi g_{\mu\nu})\cdot\delta\Phi + (D_\Phi t_{\mu\nu})\cdot\delta\Phi + (D_\Theta t_{\mu\nu})\cdot\delta\Theta.$$

Tau–kairos TT–isolation implies $\Pi_{\bar\Sigma}^{[\mathrm{TT}]}\circ D_\Theta\Sigma = 0$ (weave usage of Lemma 10.17).

Theorem 10.178 (Equivalence after TT projection and graviton sourcing)**.** *With* (10.105) *and* (10.101)*,*

$$\boxed{\Pi_{\bar\Sigma}^{[\mathrm{TT}]}\,\delta\Sigma_{\mu\nu} = \Pi_{\bar\Sigma}^{[\mathrm{TT}]}\,\delta g_{\mu\nu} = \left(\Pi_{\bar\Sigma}^{[\mathrm{TT}]}\circ D_\Phi\bar g\right)\cdot\delta\Phi, \qquad \hat h_{\mu\nu} = \Pi_{\bar\Sigma}^{[\mathrm{TT}]}\,\delta\hat g_{\mu\nu} = \Pi_{\bar\Sigma}^{[\mathrm{TT}]}\,\delta\hat\Sigma_{\mu\nu}.} \tag{10.106}$$

Thus the TT sector depends only on Gaiamiti variations; at the operator level the graviton is the TT metric fluctuation.

Proposition 10.179 (Parasymmetric alignment and T–L–O invariance)**.** *Run the sector execution functor with the* same *typed projectors and admission (Definitions 9.153 and 10.94) on either description:*

TT sector (geometry)
$\mathrm{TLO}^{[\mathrm{TT}]}_{\mathrm{formal}} = \mathrm{TLO}^{[\mathrm{TT}]}_{\mathrm{constitutive}}$ *with identical Type/Location/Onset, as per Proposition 10.35.*

Tau–kairos sector (non–TT)
For each admitted $\chi \in \{S^{(b)}, S^{(L)}, \tilde S^{(p)}, A^{(\pm)}\}$, $\mathrm{TLO}^{[\chi]}_{\mathrm{formal}} = \mathrm{TLO}^{[\chi]}_{\mathrm{constitutive}}$, *and* $\Pi_{\bar\Sigma}^{[\mathrm{TT}]}\Pi^{[\chi]} = 0$ *prevents bleed into TT (Lemmas 10.17 and 10.59 and Propositions 10.22 and 10.65).*

Helicity register and purity guards
Optional TT–helicity refinement and the stability witness follow the same meters in both views (weave graviton register; TT–purity from Lemma 10.17).

Corollary 10.180 (Uniqueness of the split (up to benign field redefinitions))**.** *If (i)* $g_{\mu\nu}$ *is Lorentzian, (ii)* $\Pi_{\bar\Sigma}^{[\mathrm{TT}]}\hat\tau = 0$*, and (iii)* $t_{\mu\nu}\to 0$ *when temporal/KMS data are trivial, then any explicit* Θ*–dependence in* $g_{\mu\nu}$ *would produce TT strength via* $D_\Theta g$ *and violate* $\Pi_{\bar\Sigma}^{[\mathrm{TT}]}\circ D_\Theta\Sigma = 0$*. Hence, modulo benign field redefinitions,*

$$g_{\mu\nu} = g_{\mu\nu}[\Phi], \qquad t_{\mu\nu} = t_{\mu\nu}[\Phi,\Theta].$$

Proof. Work on the realized weave configuration $\mathcal{I}^{[\mathcal{S}]}$ after Σ–realization (Definitions 9.80 and 10.1). By the fluctuation split $\delta\hat{\Sigma}_{\mu\nu} = \delta\hat{g}_{\mu\nu} + \delta\hat{t}_{\mu\nu}$ (eq. (10.95)) and TT–nullity of tau–kairos ($\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\hat{\tau}_{\mu\nu} = 0$; Definition 10.162, Lemma 10.17, and eq. (10.97)),

$$\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\Sigma_{\mu\nu} = \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta g_{\mu\nu}.$$

Assume, for contradiction, that $g_{\mu\nu}$ depends explicitly on Θ, so that $\delta g_{\mu\nu} = (D_\Phi g)_{\mu\nu}\cdot\,\delta\Phi + (D_\Theta g)_{\mu\nu}\cdot\,\delta\Theta$. Choose variations with $\delta\Phi = 0$. Then

$$\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\Sigma_{\mu\nu} = \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\Big[(D_\Theta g)_{\mu\nu}\cdot\,\delta\Theta\Big].$$

However, by TT selection at the constitutive level (Proposition 10.173), the only source of TT support in the geometric channel is $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\Big[(D_\Phi g)\cdot\,\delta\Phi\Big]$; that is, TT arises from *tangential* Φ–directions and not from Θ–directions. Therefore we must have

$$\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\circ D_\Theta g \;=\; 0,$$

i.e. any Θ–dependence of g is annihilated by the York TT projector (pure trace/longitudinal content; Definition 10.3).

Let Θ_0 denote trivial temporal/KMS data (assumption (iii) gives $t_{\mu\nu}[\Phi,\Theta_0] = 0$). Define the benign field redefinitions

$$g'_{\mu\nu}[\Phi] := g_{\mu\nu}[\Phi,\Theta_0], \qquad t'_{\mu\nu}[\Phi,\Theta] := t_{\mu\nu}[\Phi,\Theta] + \Big(g_{\mu\nu}[\Phi,\Theta] - g_{\mu\nu}[\Phi,\Theta_0]\Big).$$

Since $g_{\mu\nu}[\Phi,\Theta] - g_{\mu\nu}[\Phi,\Theta_0]$ is TT–killed for all Θ (previous line), this addition is a pure trace/-longitudinal transfer that does not alter the TT channel, meters, or TLO stamps (band/BRST hygiene and commutations: Propositions 10.9 and 10.10; TT isolation: Lemma 10.172 and Theorem 10.169). Consequently the geometry can be represented as $g'_{\mu\nu}[\Phi]$ with all Θ–dependence absorbed into $t'_{\mu\nu}[\Phi,\Theta]$, proving the claim "modulo benign field redefinitions." Lorentzian character (assumption (i)) ensures the background–covariant TT projector is well–defined (ellipticity/York decomposition; Definition 10.3). □

Meaning. The corollary formalizes an identifiability statement: on the realized weave configuration, the *tensor* (TT, spin–2) content of the chronogeometry comes *only* from the Gaiamiti sector Φ (tangential directions), while all Θ (temporal/kairotic) effects can be placed—without loss of physics—into the non–TT piece $t_{\mu\nu}$. If one tried to let $g_{\mu\nu}$ depend on Θ, any nontrivial $D_\Theta g$ would otherwise feed TT power and contradict the TT–selection/uniqueness results (Proposition 10.173 and Theorem 10.169). Thus, after harmless field redefinitions that shuffle pure trace/longitudinal parts (Definition 10.3 and Proposition 10.9), the split is unique in the sense stated:

$$g_{\mu\nu} = g_{\mu\nu}[\Phi] \quad \text{(TT source)}, \qquad t_{\mu\nu} = t_{\mu\nu}[\Phi,\Theta] \quad \text{(trace/longitudinal, TT–null)}.$$

Operationally, this guarantees that TT graviton claims ($\mathrm{TLO}^{[\mathrm{TT}]}$) are insensitive to temporal/KMS choices, while Θ–driven structure is recorded only in the appropriate non–TT typed ledgers (Definition 10.4 and Proposition 10.22).

Remark 10.181 (Coordinate choice and parasymmetric invariants)**.** The formal view uses $(\Sigma;$ typed projectors); the constitutive view uses (Φ, Θ) and then the *same* typed projectors. Because the sector execution depends only on typed insertion, admission, and meters (Definitions 9.153, 10.94 and 10.113), the realized T–L–O stamps on $\mathfrak{P}$ are *coordinate–invariant*:

$$\textbf{Formal:}\quad \delta\hat{\Sigma} = \delta\hat{g} + \delta\hat{\tau}, \qquad \textbf{Constitutive:}\quad \delta\Sigma = (D_\Phi g)\cdot\delta\Phi + (D_\Phi t)\cdot\delta\Phi + (D_\Theta t)\cdot\delta\Theta,$$

with TT selection and non–overlap guaranteed by (10.105) and the weave TT–null rules.

10.17.5 Relational Graviton Propagator Matrix

To characterize the dynamical behavior of quantum gravitational excitations in the IQR, we introduce the two–point correlation structure of the unified chronogeometric fluctuations organized by the Chapter 11 typed/triad machinery (Sections 10.17.3 and 10.17.4 and Definitions 10.2 to 10.4 and 10.113). In the background/fluctuation split,

$$\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}, \qquad \delta\hat{\Sigma}_{\mu\nu} = \delta\hat{g}_{\mu\nu} + \delta\hat{t}_{\mu\nu}, \tag{10.107}$$

(cf. eq. (10.95)) we define the channel operators in the Chapter 11 convention

$$\hat{h}_{\mu\nu} := \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{g}_{\mu\nu}, \qquad \hat{\tau}_{\mu\nu} := \delta\hat{t}_{\mu\nu}, \tag{10.108}$$

so that TT–selection yields $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\hat{\tau}_{\mu\nu} = 0$ by Lemma 10.17, and the TT projector acts on the unified fluctuation as

$$\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{\Sigma}_{\mu\nu} = \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{g}_{\mu\nu} = \hat{h}_{\mu\nu} \quad \text{(cf. eqs. (10.103), (10.105) and (10.106)).}$$

Thus all radiative spin–2 content resides in the metric channel and is represented by the TT graviton $\hat{h}_{\mu\nu}$.

Definition 10.182 (Geometric propagator matrix)**.** The complete two–point function of the geometric excitation sector is the matrix–valued propagator

$$\boxed{\mathbf{G}^{(2)}_{\Sigma}(x,x') := \begin{pmatrix} \langle\Omega|T\{\hat{h}_{\mu\nu}(x)\,\hat{h}_{\rho\sigma}(x')\}|\Omega\rangle & \langle\Omega|T\{\hat{h}_{\mu\nu}(x)\,\hat{\tau}_{\rho\sigma}(x')\}|\Omega\rangle \\ \langle\Omega|T\{\hat{\tau}_{\mu\nu}(x)\,\hat{h}_{\rho\sigma}(x')\}|\Omega\rangle & \langle\Omega|T\{\hat{\tau}_{\mu\nu}(x)\,\hat{\tau}_{\rho\sigma}(x')\}|\Omega\rangle \end{pmatrix}}, \tag{10.109}$$

where correlators are taken in the interacting vacuum $|\Omega\rangle$ (Section 6.1.20) and $T\{\cdot\}$ denotes the time–ordering operator (Definition 11.53). The TT definition of $\hat{h}_{\mu\nu}$ and the TT–nullity of $\hat{\tau}_{\mu\nu}$ ensure consistency with the typed projector algebra (Definition 10.3 and Lemma 10.17); typed/triad insertions and provenance constraints may be applied to each block via Definition 10.113, Proposition 10.10, and Section 10.15.

Tensorial structure. Each entry of $\mathbf{G}^{(2)}_{\Sigma}$ is a rank–4 bitensor over $\mathcal{X}_q \times \mathcal{X}_q$ with typed/triad provenance (Definitions 10.2 and 10.113 and Proposition 10.10), symmetric in each index pair and obeying reciprocity for real neutral fields:

$$G_{\mu\nu\rho\sigma}(x,x') = G_{\rho\sigma\mu\nu}(x',x). \tag{10.110}$$

When needed, one may resolve $\mathbf{G}^{(2)}_{\Sigma}$ by relational channel $a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$ consistent with the SynTh/typed ledger (Section 10.15 and Definition 10.94) and write

$$\mathbf{G}^{(2)}_{\Sigma}(x,x') = \sum_{a} \mathbf{G}^{(2,a)}_{\Sigma}(x,x')\,. \tag{10.111}$$

For the block entries, the operators are the Chapter 11 graviton $\hat{h}_{\mu\nu}$ and tau–kairos $\hat{\tau}_{\mu\nu}$ of Definition 10.182, with $\hat{h}_{\mu\nu} := \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{g}_{\mu\nu}$ and $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\hat{\tau}_{\mu\nu} = 0$ (Definition 10.3, Lemma 10.17, and eqs. (10.105) and (10.106)).

Matrix Green's equation (block form and causality). By the split eq. (10.95) and identifications eq. (10.100), the second variation of the action around $\bar{\Sigma}_{\mu\nu}$ defines a background–covariant block operator that acts on the $(\hat{h},\hat{\tau})$ sector:

$$\mathcal{D}(x) = \begin{pmatrix} \mathcal{D}_{hh} & \mathcal{C}_{h\tau} \\ \mathcal{C}_{\tau h} & \mathcal{D}_{\tau\tau} \end{pmatrix}_{(x)}, \qquad \mathcal{D}(x)\,\mathbf{G}^{(2)}_{\Sigma}(x,x') = \mathbf{I}\,\delta^{(d)}(x-x')\,. \tag{10.112}$$

Here $\mathcal{D}_{hh}$ is the (gauge–fixed) Lichnerowicz–type metric operator acting on the TT graviton sector $(\hat{h}_{\mu\nu})$, $\mathcal{D}_{\tau\tau}$ acts on tau–kairos fluctuations, and the off–diagonal couplings $\mathcal{C}_{h\tau}, \mathcal{C}_{\tau h}$ encode state–dependent conversion between metric and temporal/coherence channels (cf. Definition 10.182). TT–purity implies that, after projection with $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$, cross–blocks do not generate additional TT content (Lemma 10.17 and eqs. (10.105) and (10.106)); causality choices (retarded/advanced/Feynman) are selected consistently with the time–ordering operator and typed admission gate (Definitions 10.94 and 11.53). Fundamental solutions exist for (10.112) within the second–variation framework [156] and, for matrix–valued normally hyperbolic operators on curved backgrounds, under the standard hypotheses [244, 245].

Causal support from the unified architecture. Causality is governed by the two–layer light–cone structure determined by $\bar{\Sigma}_{\mu\nu}$ (see Chapter 5 and Section 5.4.5 and the Chapter 11 kinematic/constitutive setup Section 10.17.4). Denoting $J^{\pm}_{\bar{\Sigma}}(x')$ the future/past domains defined by the $\bar{\Sigma}$ cone, the retarded/advanced matrix propagators satisfy

$$\operatorname{supp}\mathbf{G}^{R}_{\Sigma}(\cdot,x') \subseteq J^{+}_{\bar{\Sigma}}(x'), \qquad \operatorname{supp}\mathbf{G}^{A}_{\Sigma}(\cdot,x') \subseteq J^{-}_{\bar{\Sigma}}(x'), \tag{10.113}$$

and the time–ordered (Feynman) propagator in Definition 10.182 is recovered by the usual combination with the time–ordering operator (Definition 11.53). Microcausality holds in the sense that equal–time commutators vanish at $\bar{\Sigma}$–spacelike separation [245]. Typed/triad provenance for all blocks follows the admission/ledger rules of Chapter 11 (Definitions 10.2 and 10.113 and Proposition 10.10).

TT sector consistency. Applying the Chapter 11 TT projector $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ (Definition 10.3) to eq. (10.112) and using $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\hat{\tau}_{\mu\nu} = 0$ (Lemma 10.17 and eq. (10.105)) isolates the graviton equation in the spin–2 sector, reproducing the GW operator and TT wave dynamics (eqs. (10.96) and (10.131)). Equivalently, with eq. (10.106) one has that off–diagonal blocks do not generate extra TT polarizations after projection.

Definition 10.183 (TT–projected and typed propagators)**.** Define the TT–projected metric–metric propagator

$$G^{\mathrm{TT}}_{\mu\nu\rho\sigma}(x,x') := \Big\langle \Omega \Big| \, T\{\hat{h}_{\mu\nu}(x)\hat{h}_{\rho\sigma}(x')\} \, \Big| \Omega \Big\rangle = \Pi^{[\mathrm{TT}]\,\alpha\beta}_{\bar{\Sigma}\,\mu\nu}\, \Pi^{[\mathrm{TT}]\,\gamma\delta}_{\bar{\Sigma}\,\rho\sigma}\, G^{hh}_{\alpha\beta\gamma\delta}(x,x'), \tag{10.114}$$

where $\hat{h}_{\mu\nu} := \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\, \delta\hat{g}_{\mu\nu}$ is the Chapter 11 graviton (eqs. (10.103) and (10.105) and Definition 10.3). Typed correlators are obtained by inserting the admissible typed projectors $\Pi^{[\chi]}$ on the operator legs,

$$G^{[\chi\chi']}(x,x') := \Big\langle \Omega \Big| \, T\{\Pi^{[\chi]}\cdot \delta\hat{\Sigma}(x)\ \Pi^{[\chi']}\cdot \delta\hat{\Sigma}(x')\} \, \Big| \Omega \Big\rangle,$$

with the typed/triad conventions of Chapter 11 (Definitions 10.2, 10.25 and 10.113). For $\chi \in \{\gamma, 0_\Phi, \tau_S, \tau_A, K^{(\tau)}_{\mathrm{T}}, K^{(\tau)}_{\mathrm{L}}\}$ (cf. Definition 10.13, Lemma 10.59, and Section 10.15), these correlators resolve $\mathbf{G}^{(2)}_{\Sigma}$ into the physically meaningful spin/helicity/polarization blocks while preserving TT purity (Lemma 10.17).

Proposition 10.184 (TT closure and null cross–blocks)**.** *With the Chapter 11 identities* $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}} \circ \Pi_t = 0$ *and* $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}} \circ D_\Theta \Sigma = 0$ *(cf. eqs. (10.8) and (10.101) and Lemma 10.17), one has*

$$\Big\langle \Omega \Big| \, T\{\hat{h}_{\mu\nu}(x)\, \hat{\tau}_{\rho\sigma}(x')\} \, \Big| \Omega \Big\rangle = 0, \tag{10.115}$$

and, more generally, $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ *annihilates any typed leg in the temporal class* $\chi \in \{\tau_S, \tau_A, K^{(\tau)}_{\mathrm{T}}, K^{(\tau)}_{\mathrm{L}}\}$ *(Definition 10.4). Hence the TT propagator eq. (10.114) is dynamically closed (eqs. (10.105) and (10.106)).*

Proposition 10.185 (Parasymmetric T–L–O augmentation of propagators)**.** *On a certified plateau* $\mathfrak{P}$ *(Definition 10.1), the sector execution functor* $\mathcal{E}_\bullet$ *(Definition 9.153) acts on* $\mathbf{G}^{(2)}_{\Sigma}$ *as follows:*

(i) Permit ledger and typed admission
Build the ledger of admissible types $\Pi^{[\chi]}$ *and enforce the admission gate (Definition 10.94) and cone–gating (eq. (10.65)); non–admitted types receive no T–L–O. Typed/triad consistency follows Definitions 10.2 and 10.113 and Proposition 10.10.*

(ii) Typed visibilities from correlators
Extract law/state visibilities $\vec{\Lambda}^{[\chi]}$ *by reading spectral weight (poles/residues) and integrated strengths from the typed correlators* $G^{[\chi\chi]}$*; assign the quadrant* $\mathrm{Quad}^{[\chi]}$ *(QI–QIV) using the Chapter 9 decision rule, while respecting Chapter 11's spin/polarization scope (Section 10.15).*

(iii) Parasymmetric meters and Onset
Evaluate $v^{(a)}_{\mathrm{Synthesis}}$, $v^{(LS)}_{\mathrm{Synthesis}}$, $v^{(LS)}_{\mathrm{Agreement}}$ *on the typed propagators' windows; the Onset* $(r^\star, t^\star)$ *is the (scale,epoch) at which the meters cross their tolerance thresholds (cf. Definition 9.153). Law–side TT purity and helicity asymmetries are monitored via* $\mathcal{W}_{\mathrm{TT\text{–}purity}}$ *and* Δ_χ *(cf. Lemma 10.17 and Definition 10.26).*

(iv) T–L–O stamping

Record $\mathrm{TLO}^{[\chi]} = (\mathrm{Type}, \mathrm{Location}, \mathrm{Onset})$ *with Location taken from diagnostic realized reference configuration metadata (arena/sector/space tag/cone sheet) and Type from the quadrant; align with Chapter 11's TLO propositions (Propositions 10.22, 10.35 and 10.65).*

The resulting T–L–O augmentation is BRST/Nielsen–safe and unique by Proposition 10.90.

Physical interpretation of the diagonal components

Metric–channel (hh) propagator and TT graviton.

$$G^{hh}_{\mu\nu\rho\sigma}(x,x') := \Big\langle \Omega \Big| \, T\{\hat{h}_{\mu\nu}(x)\hat{h}_{\rho\sigma}(x')\} \, \Big| \Omega \Big\rangle. \tag{10.116}$$

This metric–channel correlator is the causal response of spatial–distinguishability fluctuations in the TT sector, with $\hat{h}_{\mu\nu} := \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}} \, \delta\hat{g}_{\mu\nu}$ the Chapter 11 graviton (Definitions 10.3 and 10.182 and eqs. (10.105) and (10.106)). The *graviton propagator* is the TT–projected object

$$G^{\mathrm{TT}}_{\mu\nu\rho\sigma}(x,x') := \Big\langle \Omega \Big| \, T\{\hat{h}_{\mu\nu}(x)\hat{h}_{\rho\sigma}(x')\} \, \Big| \Omega \Big\rangle, \tag{10.117}$$

and, using the TT projector explicitly,

$$G^{\mathrm{TT}}_{\mu\nu\rho\sigma} = \Pi^{[\mathrm{TT}]\,\alpha\beta}_{\bar{\Sigma}\,\mu\nu} \, \Pi^{[\mathrm{TT}]\,\gamma\delta}_{\bar{\Sigma}\,\rho\sigma} \, G^{hh}_{\alpha\beta\gamma\delta} \quad \text{(cf. eq. (10.114) and Definition 10.3).}$$

By eq. (10.115) (TT–closure and null cross–blocks), the mixed TT–temporal correlator vanishes, consistent with Lemma 10.17 and eq. (10.105).

Tau–kairos ($\tau\tau$) propagator $\langle T\{\hat{\tau}_{\mu\nu}(x)\hat{\tau}_{\rho\sigma}(x')\}\rangle$. This correlator tracks the propagation of temporal phase–flow fluctuations. It does not mediate a classical force; rather, it modulates quantum phase/coherence transport, influences entanglement dynamics, and can affect geometric–phase acquisition in embedded quantum systems. In the parasymmetric T–L–O pipeline, the typed entries $\chi \in \{\tau_S, \tau_A, K^{(\tau)}_{\mathrm{T}}, K^{(\tau)}_{\mathrm{L}}\}$ (Chapter 11 temporal class, Definitions 10.4 and 10.13) receive Onset windows $(r^\star, t^\star)$ from the meters (coherence activation scales), read from typed correlators $G^{[\chi\chi]}$ (Definitions 10.94, 10.113 and 10.183); but they remain TT–null by Lemmas 10.17 and 10.59, thereby leaving the graviton T–L–O entry $\mathrm{TLO}^{[\mathrm{TT}]}$ (two tensor polarizations) unaltered (Propositions 10.10, 10.22 and 10.35 and Section 10.15).

Graviton–Kairos Mixing and Off–Diagonal Correlators

The off–diagonal entries $\Big\langle \Omega \Big| \, T\{\hat{h}_{\mu\nu}(x)\hat{\tau}_{\rho\sigma}(x')\} \, \Big| \Omega \Big\rangle$ and its transpose encode *state–dependent conversion amplitudes* between spatial (metric) and temporal/coherence excitations. They are absent in classical GR and reflect the deeper relational structure of geometry in the IQR. Crucially, by the TT–null constraint for the temporal channel (Lemma 10.17 and eq. (10.105)) and TT–closure (Proposition 10.184 and eq. (10.115)), these mixings carry *no spin–2 content*:

$$\Big\langle \Omega \Big| \, T\{\hat{h}_{\mu\nu}(x) \, \hat{\tau}_{\rho\sigma}(x')\} \, \Big| \Omega \Big\rangle \; = \; 0, \tag{10.118}$$

with the same conclusion for the transpose. Equivalently, after projection with the Chapter 11 TT projector $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ (Definition 10.3), off–diagonal blocks cannot generate TT modes (eq. (10.106)).

Analogy. Just as phonons and non–propagating thermal excitations can mix in a lattice due to their common medium, the metric–channel and tau–kairos fluctuations mix because they are orthogonal components of the unified fluctuation $\delta\hat{\bar{\Sigma}}_{\mu\nu}$ (cf. eq. (10.95)), *not* because the graviton is sourced by the temporal sector. TT projection isolates the radiative tensor content uniquely in $\hat{h}_{\mu\nu} = \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{g}_{\mu\nu}$ (eqs. (10.103) and (10.105)), consistently with the GW operator and TT wave dynamics (eqs. (10.96) and (10.131)).

Propagator as a Matrix–Valued Green's Function

The full matrix $\mathbf{G}^{(2)}_{\Sigma}(x,x')$ solves the linearized coupled system for unified gravitational fluctuations:

$$\mathcal{D}(x)\,\mathbf{G}^{(2)}_{\Sigma}(x,x') \;=\; \mathbf{I}\,\delta^{(d)}(x-x'), \qquad \text{cf. } eq.\ (10.112). \tag{10.119}$$

Here $\mathcal{D}$ is the second–variation operator with off–diagonal couplings $\mathcal{C}_{h\tau}, \mathcal{C}_{\tau h}$ that generate the mixing entries (cf. Definition 10.182). Cone–gating and existence of fundamental solutions follow the standard theory for (matrix–valued) normally hyperbolic operators on curved backgrounds [156, 244, 245], together with the unified light–cone structure of $\bar{\Sigma}$ (Section 5.4.5) and the typed/triad provenance rules (Definitions 10.2 and 10.113 and Proposition 10.10).

Parasymmetric T–L–O view of mixing. On a certified plateau $\mathfrak{P}$ (Definition 10.1), the sector execution functor $\mathscr{E}_\bullet$ (Definition 9.153) reads the *typed* off–diagonal correlators $G^{[\chi\,\chi']}$ (with $\chi \in \{\gamma\}$ or $\chi \in \{\tau_S, \tau_A, K^{(\tau)}_{\mathrm{T}}, K^{(\tau)}_{\mathrm{L}}\}$) to assign quadrants and Onset windows via the parasymmetric meters, subject to typed admission and cone–gating (Definition 10.94 and eq. (10.65)). TT–null types never contribute to the γ T–L–O entry; $\mathrm{TLO}^{[\gamma]}$ remains the unique spin–2 stamp (Propositions 10.22, 10.35 and 10.90, Lemma 10.59, and Section 10.15). Typed/triad consistency across all legs is enforced via Definitions 10.2, 10.4 and 10.113.

Channel–Resolved Propagator for Chronogeometric Fluctuations (Bimodal–Triadic Form)

The two–point function $\mathbf{G}^{(2)}_{\Sigma}(x,x')$ (Definition 10.182) is naturally organized as a *bi–modal* split (metric vs. temporal) refined by a *triadic* source resolution $a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$, in the Chapter 11 typed/triad framework (Section 10.17.4, Definitions 10.2 and 10.113, and Proposition 10.10). The bimodal background/fluctuation structure is

$$\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}, \qquad \delta\hat{\bar{\Sigma}}_{\mu\nu} = \delta\hat{g}_{\mu\nu} + \delta\hat{t}_{\mu\nu} \quad (\text{cf. } eq.\ (10.95)),$$

with channel operators fixed by Chapter 11 as

$$\hat{h}_{\mu\nu} := \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{g}_{\mu\nu}, \qquad \hat{\tau}_{\mu\nu} := \delta\hat{t}_{\mu\nu},$$

so that $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\hat{\tau}_{\mu\nu} = 0$ (Definition 10.3 and Lemma 10.17) and $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{\bar{\Sigma}}_{\mu\nu} = \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{g}_{\mu\nu} = \hat{h}_{\mu\nu}$ (eqs. (10.103), (10.105) and (10.106)). The GW operator and TT wave dynamics appear in eqs. (10.96) and (10.131).

Proposition 10.186 (Triadic channel resolution and structure)**.** *As a sum over interaction channels,*

$$\boxed{\mathbf{G}^{(2)}_{\Sigma}(x,x') \;=\; \sum_{a\in\{\mathrm{corr,vac,spec}\}} \mathbf{G}^{(2,a)}_{\Sigma}(x,x')\,,} \tag{10.120}$$

where each $\mathbf{G}^{(2,a)}_{\Sigma}$ *is a* 2×2 *matrix of time–ordered correlators,*

$$\mathbf{G}^{(2,a)}_{\Sigma}(x,x') = \begin{pmatrix} \langle\Omega|T\{\hat{h}^{(a)}_{\mu\nu}(x)\,\hat{h}^{(a)}_{\rho\sigma}(x')\}|\Omega\rangle & \langle\Omega|T\{\hat{h}^{(a)}_{\mu\nu}(x)\,\hat{\tau}^{(a)}_{\rho\sigma}(x')\}|\Omega\rangle \\ \langle\Omega|T\{\hat{\tau}^{(a)}_{\mu\nu}(x)\,\hat{h}^{(a)}_{\rho\sigma}(x')\}|\Omega\rangle & \langle\Omega|T\{\hat{\tau}^{(a)}_{\mu\nu}(x)\,\hat{\tau}^{(a)}_{\rho\sigma}(x')\}|\Omega\rangle \end{pmatrix},$$

with $T\{\cdot\}$ *as in Definition 11.53 and the vacuum* $|\Omega\rangle$ *as in Section 6.1.20. Each entry is a rank–4 bitensor, obeys reciprocity, and is causal with respect to the* $\bar{\Sigma}$ *cone (eq. (10.112) and Section 5.4.5); typed/triad provenance may be carried on each leg via Definitions 10.2 and 10.113.*

TT sector and selection rule. The physical (spin–2) propagator is the TT–projected metric correlator,

$$G^{\mathrm{TT}}_{\mu\nu\rho\sigma}(x,x') := \Big\langle\Omega\Big|\, T\{\hat{h}_{\mu\nu}(x)\hat{h}_{\rho\sigma}(x')\}\,\Big|\Omega\Big\rangle. \tag{10.121}$$

Cf. eqs. (10.114) and (10.131) and Definition 10.3.

and the selection rule

$$\Big\langle\Omega\Big|\, T\{\hat{h}_{\mu\nu}(x)\hat{\tau}^{(a)}_{\rho\sigma}(x')\}\,\Big|\Omega\Big\rangle = 0 \qquad \forall\, a\in\{\mathrm{corr,vac,spec}\}$$

follows from TT–nullity of the temporal class and TT–closure (Lemma 10.17, eqs. (10.90) and (10.115), and Proposition 10.184). Thus all radiative spin–2 content flows through the hh block (eqs. (10.105) and (10.106)).

Parasymmetric T–L–O augmentation (channel–wise). On $\mathfrak{P}$ (Definition 10.1), apply $\mathscr{E}_\bullet$ to the *typed*, channel–resolved correlators $G^{[\chi\chi']\langle a\rangle}$ (Definition 10.183) to: (i) enforce typed admission (Definition 10.94); (ii) compute typed visibilities $\vec{\Lambda}^{[\chi]}$ and assign quadrants (QI–QIV) consistent with Chapter 11's spin/polarization scope (Section 10.15); (iii) extract Onset windows $(r^\star,t^\star)$ from the parasymmetric meters (Definition 9.153); and (iv) stamp $\mathrm{TLO}^{[\chi]}$ with diagnostic realized reference configuration–derived Locations (Propositions 10.22, 10.35 and 10.65). Because $\chi\in\{\tau_S,\tau_A,K^{(\tau)}_{\mathrm{T}},K^{(\tau)}_{\mathrm{L}}\}$ are TT–null (Lemmas 10.17 and 10.59), their T–L–O entries never alter $\mathrm{TLO}^{[\mathrm{TT}]}$; law–side TT purity and helicity asymmetries are monitored via $\mathcal{W}_{\mathrm{TT\text{–}purity}}$ and Δ_χ (cf. Definition 10.26).

Consequence and channel count

For each $a\in\{\mathrm{corr,vac,spec}\}$ the operator content decomposes into the pair $(\hat{h}^{(a)}_{\mu\nu},\hat{\tau}^{(a)}_{\mu\nu})$ in the Chapter 11 bimodal split (Section 10.17.4 and Definition 10.182). Hence there are exactly six channel–separated modes:

$$(\hat{h},\hat{\tau})\times(\mathrm{corr,vac,spec}) = \{\,\hat{h}^{(\mathrm{corr})},\hat{h}^{(\mathrm{vac})},\hat{h}^{(\mathrm{spec})},\hat{\tau}^{(\mathrm{corr})},\hat{\tau}^{(\mathrm{vac})},\hat{\tau}^{(\mathrm{spec})}\,\}.$$

Among these, only the TT graviton sector

$$\hat{h}_{\mu\nu} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}_{\mu\nu} \quad \Rightarrow \quad G^{\mathrm{TT}}_{\mu\nu\rho\sigma} = \left\langle \Omega \middle| T\{\hat{h}_{\mu\nu}\hat{h}_{\rho\sigma}\} \middle| \Omega \right\rangle$$

supplies propagating tensor quanta (two polarizations) in the radiative channel (eqs. (10.96) and (10.131) and Section 10.15), with the projector definition and TT selection given by Definition 10.3 and eqs. (10.105), (10.106) and (10.114). The tau–kairos sector remains strictly non–TT (eq. (10.90), Lemmas 10.17 and 10.59, and Definition 10.4), contributing only spin–0 and spin–1 structure and no extra GW polarizations.

Origin Channel	**Graviton (Spin-2)** $\hat{h}^{(a)}_{\mu\nu}(x)$	**Tau-Kairos tensor (Spin-0,1)** $\hat{\tau}^{(a)}_{\mu\nu}(x)$
Correlational	$\hat{h}^{(\mathrm{corr})}_{\mu\nu}(x)$ *(eq. (11.74))*	$\hat{\tau}^{(\mathrm{corr})}_{\mu\nu}(x)$ *(temporal entanglement)*
Vacuum	$\hat{h}^{(\mathrm{vac})}_{\mu\nu}(x)$ *(Definition 11.43)*	$\hat{\tau}^{(\mathrm{vac})}_{\mu\nu}(x)$ *(vacuum coherence structure)*
Spectral	$\hat{h}^{(\mathrm{spec})}_{\mu\nu}(x)$ *(Definition 11.44)*	$\hat{\tau}^{(\mathrm{spec})}_{\mu\nu}(x)$ *(energy-induced phase dynamics)*

Table 10.3: The Six Fundamental Modes of Geometric Excitation

10.17.6 Triadic Decomposition of Gravitational Propagation

In the IQR framework, emergent geometry is encoded by the operator–valued chronogeometric tensor $\hat{\Sigma}_{\mu\nu}(x)$, with background $\bar{\Sigma}_{\mu\nu}$ and fluctuation $\delta\hat{\Sigma}_{\mu\nu} = \delta\hat{g}_{\mu\nu} + \delta\hat{t}_{\mu\nu}$ (cf. eq. (10.95)). We adopt the *triadic channelization* introduced for the propagator (corr, vac, spec; see Definitions 10.2, 10.3, 10.25 and 10.113 and Proposition 10.10) as an organizing principle for fluctuations on $\bar{\Sigma}$. In Chapter 11 notation the bimodal operators are $\hat{h}_{\mu\nu} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}_{\mu\nu}$ and $\hat{\tau}_{\mu\nu} := \delta\hat{t}_{\mu\nu}$ with $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\hat{\tau}_{\mu\nu} = 0$ (Definition 10.3, Lemma 10.17, and eqs. (10.103), (10.105) and (10.106)).

We now show that this triadic structure induces a corresponding decomposition of the metric and tau–kairos fluctuations—and hence of the full gravitational propagator matrix.

Definition 10.187 (Triadic decomposition of chronogeometric fluctuations and graviton sector)**.** The unified fluctuation decomposes by channels as

$$\boxed{\begin{aligned} \delta\hat{\Sigma}_{\mu\nu}(x) &= \delta\hat{\Sigma}^{(\mathrm{corr})}_{\mu\nu}(x) + \delta\hat{\Sigma}^{(\mathrm{vac})}_{\mu\nu}(x) + \delta\hat{\Sigma}^{(\mathrm{spec})}_{\mu\nu}(x), \\ \hat{h}_{\mu\nu}(x) &= \sum_{a\in\{\mathrm{corr},\mathrm{vac},\mathrm{spec}\}} \hat{h}^{(a)}_{\mu\nu}(x), \qquad \hat{\tau}_{\mu\nu}(x) = \sum_{a} \hat{\tau}^{(a)}_{\mu\nu}(x), \end{aligned}} \tag{10.122}$$

with $\hat{h}_{\mu\nu} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}_{\mu\nu}$ and $\hat{\tau}_{\mu\nu} := \delta\hat{t}_{\mu\nu}$ (Definition 10.3 and eq. (10.105)). The physical (spin–2) graviton in each channel is already TT by construction, while the temporal class is TT–null:

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\hat{h}^{(a)}_{\mu\nu} = \hat{h}^{(a)}_{\mu\nu}, \qquad \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\hat{\tau}^{(a)}_{\mu\nu} = 0 \quad \text{(cf. Lemma 10.17 and eq. (10.90))},$$

so that $\hat{h}_{\mu\nu} = \sum_a \hat{h}^{(a)}_{\mu\nu}$ and no TT content arises from τ (cf. Corollary 10.20 and eq. (10.131)).

Each $\hat{h}^{(a)}_{\mu\nu}$ is a covariant symmetric TT tensor field and each $\hat{\tau}^{(a)}_{\mu\nu}$ a covariant symmetric non–TT tensor; the channels are mutually orthogonal in the sense induced by the triadic split and typed admission (Definitions 10.2, 10.25, 10.94 and 10.113).

Definition 10.188 (Triadic mode decomposition of the gravitational propagator)**.** The complete geometric propagator matrix $\mathbf{G}^{(2)}_{\Sigma}(x,x')$ (Definition 10.182) decomposes as

$$\boxed{\mathbf{G}^{(2)}_{\Sigma}(x,x') = \sum_{a\in\{\mathrm{corr,vac,spec}\}} \mathbf{G}^{(2,a)}_{\Sigma}(x,x')\,,} \tag{10.123}$$

where each $\mathbf{G}^{(2,a)}_{\Sigma}$ retains the full 2×2 block structure over (h,τ) (cf. Section 10.17.5):

$$\mathbf{G}^{(2,a)}_{\Sigma}(x,x') = \begin{pmatrix} \langle\Omega|\,T\{\hat{h}^{(a)}_{\mu\nu}(x)\hat{h}^{(a)}_{\rho\sigma}(x')\}\,|\Omega\rangle & \langle\Omega|\,T\{\hat{h}^{(a)}_{\mu\nu}(x)\hat{\tau}^{(a)}_{\rho\sigma}(x')\}\,|\Omega\rangle \\ \langle\Omega|\,T\{\hat{\tau}^{(a)}_{\mu\nu}(x)\hat{h}^{(a)}_{\rho\sigma}(x')\}\,|\Omega\rangle & \langle\Omega|\,T\{\hat{\tau}^{(a)}_{\mu\nu}(x)\hat{\tau}^{(a)}_{\rho\sigma}(x')\}\,|\Omega\rangle \end{pmatrix}.$$

The TT–projected graviton propagator correspondingly decomposes as

$$G^{\mathrm{TT}}_{\mu\nu\rho\sigma}(x,x') = \sum_a \langle\Omega|\,T\{\hat{h}^{(a)}_{\mu\nu}(x)\hat{h}^{(a)}_{\rho\sigma}(x')\}\,|\Omega\rangle,$$

with no TT leakage into τ by $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\hat{\tau}^{(a)} = 0$ (cf. Lemma 10.17 and eq. (10.90)); see also eqs. (10.105), (10.106) and (10.114).

Physical interpretation of the triadic channels. The three channels are distinct physical sources of curvature, yielding complementary gravitational responses:

(i) Correlational (corr) *Entanglement–induced geometry*—curvature sourced by inter–modal quantum correlations and informational structure (consistent with the channelization in Definitions 10.2, 10.3, 10.25 and 10.113).

(ii) Vacuum (vac) *Vacuum–deformation geometry*—curvature from vacuum–manifold structure and SSB in the Gaiamiti sector; the TT graviton's masslessness arises from tangential (Goldstone) directions (cf. Sections 9.17.2 and 10.5 and eqs. (10.96), (10.103) and (10.131); see also [415, 502]).

(iii) Spectral (spec) *Energy–induced geometry*—curvature sourced by spectral energy–momentum (i.e. matter/kymon excitations) in the operator Green–function framework on curved backgrounds [244, 7]; causal support follows the $\bar{\Sigma}$ cone (Section 5.4.5).

Remark 10.189 (Quantum gravity as operator–valued dynamics)**.** In this representation, quantum gravity is a state–relative, operator–valued response theory: the coupled operators $(\hat{h}_{\mu\nu}, \hat{\tau}_{\mu\nu})$ evolve via linearized second–variation equations (cf. eq. (10.112) and Section 10.17.7), and their *channel–resolved* propagators $\mathbf{G}_{\Sigma}^{(2,a)}$ encode causal, covariant responses to perturbations of distinct origin. Causality and support are governed by the $\bar{\Sigma}$–light cone (Section 5.4.5); the TT projector isolates the radiative spin–2 sector, ensuring that gravitational waves reside entirely in $\hat{h}_{\mu\nu}$ with two polarizations (cf. eqs. (10.28), (10.105), (10.114) and (10.131) and Corollary 10.20). For background–field and propagator conventions, see [244, 7].

10.17.7 Covariant Dynamics of Gravitational Fluctuations

Gravitational excitations in the IQR arise as quantum fluctuations of the operator–valued chronogeometric tensor $\hat{\Sigma}_{\mu\nu}(x) = \hat{g}_{\mu\nu}(x) + \hat{t}_{\mu\nu}(x)$ (cf. Section 5.2.6 and eqs. (10.94) and (10.95)). The observable TT (spin–2) sector is the Chapter 11 graviton $\hat{h}_{\mu\nu} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \delta\hat{g}_{\mu\nu}$, while the temporal/coherence sector is $\hat{\tau}_{\mu\nu} := \delta\hat{t}_{\mu\nu}$ with $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\hat{\tau}_{\mu\nu} = 0$ (Definition 10.3, eqs. (10.90), (10.103) and (10.105), and Lemma 10.17). Microscopically, Σ is produced constitutively from the substrate fields (Φ, Θ) via $\Sigma_{\mu\nu}[\Phi, \Theta] = g_{\mu\nu}[\Phi] + t_{\mu\nu}[\Phi, \Theta]$ (eqs. (10.94) and (10.101)), so that, upon linearization, $\delta\Sigma_{\mu\nu} = (D_{\Phi}g)\cdot\delta\Phi + (D_{\Phi}t)\cdot\delta\Phi + (D_{\Theta}t)\cdot\delta\Theta$ (eq. (10.101)). The TT projection singles out the Gaiamiti variations:

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\Sigma = \Big(\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \circ D_{\Phi}g\Big)\cdot\delta\Phi \quad \text{(cf. eq. (10.106))}.$$

1. Linearization of the Gaiamiti action

We start from the Gaiamiti action (background connection $\bar{\nabla}$ and measure built from $\bar{\Sigma}_{\mu\nu}$; cf. eq. (10.87)):

$$S_{\Phi}[\hat{\mathbf{\Phi}}] = \int_{\mathcal{X}_q} d^d x \,\sqrt{-\det\big(\bar{\Sigma}\big)}\;\mathrm{Tr}\Big\{\bar{\Sigma}^{\mu\nu}\, g_T\Big(\bar{\nabla}_{\mu}\hat{\mathbf{\Phi}}, \bar{\nabla}_{\nu}\hat{\mathbf{\Phi}}\Big) + \mathcal{V}_{\Phi}(\hat{\mathbf{\Phi}})\Big\}\,.$$

Decompose about the expectation background (cf. eq. (10.87))

$$\hat{\mathbf{\Phi}}(x) = \bar{\Phi}(x)\,\mathbb{I} + \delta\hat{\mathbf{\Phi}}(x),$$

and expand $S_{\Phi} = S^{(0)} + S^{(1)} + S^{(2)} + \cdots$. Stationarity of $\bar{\Phi}$ sets $S^{(1)} = 0$. The quadratic action governing fluctuations is

$$\boxed{S^{(2)}[\delta\hat{\mathbf{\Phi}}] = \int d^d x \,\sqrt{-\det\big(\bar{\Sigma}\big)}\;\mathrm{Tr}\Big\{\bar{\Sigma}^{\mu\nu}\Big(\bar{\nabla}_{\mu}\delta\hat{\mathbf{\Phi}}\Big)^{\dagger}\Big(\bar{\nabla}_{\nu}\delta\hat{\mathbf{\Phi}}\Big) - \tfrac{1}{2}\,\delta\hat{\mathbf{\Phi}}^{\dagger}\,\mathcal{M}(x)\,\delta\hat{\mathbf{\Phi}}\Big\}\,,} \tag{10.124}$$

with the (generally position–dependent, operator–valued) mass kernel

$$\boxed{\mathcal{M}(x) := \left.\frac{\delta^2\mathcal{V}_{\Phi}}{\delta\hat{\mathbf{\Phi}}^{\dagger}(x)\,\delta\hat{\mathbf{\Phi}}(x)}\right|_{\bar{\Phi}}.} \tag{10.125}$$

Euler–Lagrange variation. Varying $S^{(2)}$ with respect to $\delta\hat{\mathbf{\Phi}}^\dagger$ gives

$$\frac{\delta S^{(2)}}{\delta\big(\delta\hat{\mathbf{\Phi}}^\dagger\big)} = -\,\bar{\nabla}_\mu\Big(\bar{\Sigma}^{\mu\nu}\bar{\nabla}_\nu\delta\hat{\mathbf{\Phi}}\Big) - \mathcal{M}(x)\,\delta\hat{\mathbf{\Phi}} = 0,$$

whence the covariant fluctuation equation

$$\boxed{\bar{\nabla}_\mu\Big(\bar{\Sigma}^{\mu\nu}\bar{\nabla}_\nu\delta\hat{\mathbf{\Phi}}\Big) - \mathcal{M}(x)\,\delta\hat{\mathbf{\Phi}} = 0.} \tag{10.126}$$

Introducing the unified d'Alembertian $\Box_{\bar{\Sigma}} := \bar{\nabla}_\mu\big(\bar{\Sigma}^{\mu\nu}\bar{\nabla}_\nu\,\cdot\,\big)$ (or equivalently $\bar{\nabla}^\mu\bar{\nabla}_\mu$ with indices raised by $\bar{\Sigma}^{\mu\nu}$), (10.126) is $(\Box_{\bar{\Sigma}} - \mathcal{M})\,\delta\hat{\mathbf{\Phi}} = 0$. This normally–hyperbolic form ensures well–posed propagation with respect to the $\bar{\Sigma}$ light–cones (cf. Section 5.4.5); it provides the dynamical seeds for the propagator matrix of Definition 10.182 and its Green's equation eq. (10.112).

From $\delta\hat{\mathbf{\Phi}}$ to observables. Via the constitutive chain rule (eq. (10.101)),

$$\delta\hat{g}_{\mu\nu} = \Big(D_\Phi g_{\mu\nu}\Big)\cdot\delta\hat{\mathbf{\Phi}}, \qquad \delta\hat{t}_{\mu\nu} = \Big(D_\Phi t_{\mu\nu}\Big)\cdot\delta\hat{\mathbf{\Phi}} + \Big(D_\Theta t_{\mu\nu}\Big)\cdot\delta\hat{\mathbf{\Theta}},$$

and, by TT isolation (Lemma 10.17 and eq. (10.105)), the radiative sector is

$$\hat{h}_{\mu\nu} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}_{\mu\nu} = \Big(\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\circ D_\Phi g\Big)\cdot\delta\hat{\mathbf{\Phi}} \quad \text{(cf. eqs. (10.103) and (10.106)).}$$

Thus (10.126) is the dynamical origin for all spin–2 quanta after TT projection, while $\hat{\tau}_{\mu\nu}$ remains non–TT (cf. Section 10.17.8 and Lemma 10.59).

Parasymmetric T–L–O view (typed, localized, onset–gated). The fluctuation equation (10.126) is admitted as a physical evolution only when the typed admission gate certifies hyperbolicity and cone–gating ($\Box_{\bar{\Sigma}}$ normally hyperbolic; BRST–closed) for the relevant types (γ in the TT sector; τ_S, τ_A in the temporal sector) (Definition 10.94). Upon admission, the sector–execution functor $\mathcal{E}_\bullet$ stamps $\mathrm{TLO}^{[\chi]} = (\mathrm{Type}, \mathrm{Location}, \mathrm{Onset})$ for each χ (Definition 9.153): Type is fixed by the projector (γ vs. τ); Location is read from the diagnostic realized reference configuration/arena metadata and cone–sheet tags; Onset $(r^\star, t^\star)$ is extracted from the parasymmetric meters and defines the coherence windows during which the corresponding solutions of (10.126) (pushed forward to δg and δt) are operational in readouts (cf. Propositions 10.22, 10.35 and 10.65). Law–side TT purity is monitored by $\mathcal{W}_{\mathrm{TT\text{–}purity}}$ (Lemma 10.17), ensuring no leakage of non–TT content into the graviton ledger and consistency with the null cross–blocks of Proposition 10.184.

Summary. The quadratic Gaiamiti action (10.124) yields the covariant fluctuation equation (10.126). Through the constitutive map and TT projection, its solutions generate the TT graviton sector $\hat{h}_{\mu\nu}$ (two polarizations) while the temporal/coherence response $\hat{\tau}_{\mu\nu}$ remains TT–null (eq. (10.90)). Parasymmetric T–L–O formalisms provide the operational gate (*admission*), the contextual placement (*location*), and the meter–derived thresholds (*onset*) for these fluctuations in experimental and cosmological settings (Definitions 9.153 and 10.94).

2. Interpretation of the Linearized Dynamics

Equation (10.126) is the covariant wave equation for Gaiamiti fluctuations ("Apeiron" modes), i.e. a curved–background generalization of a relativistic Klein–Gordon equation that controls how geometric disturbances propagate and seed observable responses (Section 10.17.7).

(i) Covariant chronogeometric propagation. The kinetic operator $\bar{\nabla}^{\mu}\bar{\nabla}_{\mu}$ is built from the background *chronogeometric tensor* $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (cf. Section 5.2.6 and eq. (10.94)). Thus the field–level Apeiron propagation is sensitive to both the metric background $\bar{g}_{\mu\nu}$ (spatial distinguishability) and the temporal/coherence background $\bar{t}_{\mu\nu}$ (phase flow). Through the constitutive map, this unified propagation feeds the two observable channels g and t (Section 10.17.4 and eq. (10.101)). *By contrast*, the radiative TT graviton obeys the TT–projected metric equation built with $\bar{g}_{\mu\nu}$, $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\big[\delta\bar{G}_{\mu\nu}[\hat{h}] + \Lambda\,\hat{h}_{\mu\nu}\big] = 0$ (Definition 10.3 and eqs. (10.103) and (10.131)); any $\bar{t}_{\mu\nu}$ influence can only enter indirectly via non–TT mixing and background parameters, not as extra TT polarizations (cf. Lemma 10.17, eq. (10.90), and Proposition 10.184).

(ii) Operator mass term $\mathcal{M}(x)$ and Goldstone directions. The operator $\mathcal{M}(x)$ arises from the Hessian of the Gaiamiti potential $\mathcal{V}_{\Phi}$ (eq. (10.125)) and acts as a local restoring force. Flat directions of $\mathcal{V}_{\Phi}$ on the vacuum manifold $\mathcal{M}_{\mathrm{vac}}$ yield zero eigenvalues (Goldstone modes) [93, 94]. In our framework the *massless spin–2 graviton* is precisely the Goldstone arising from tangential variations of Φ projected to the TT sector (Sections 9.17.2 and 10.5 and eqs. (10.96), (10.103) and (10.106)). The tau–kairos sector, being TT–null ($\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\hat{\tau} = 0$; Lemma 10.17 and eq. (10.90)), is *not* a spin–2 Goldstone; its scalar/vector components may be massive or massless depending on model symmetries (e.g. a gauged Chrono–Kymon vs. Proca–type mass; cf. Definition 10.57 and Section 10.18.6).

(iii) Induced dynamics of gravitational operators. Via the linear response of the constitutive map (eq. (10.101)),

$$\delta\hat{\Sigma}_{\mu\nu} \;=\; \big(D_{\Phi}\Sigma\big)_{\mu\nu}\cdot\delta\hat{\mathbf{\Phi}} \;=\; \underbrace{\big(D_{\Phi}g\big)_{\mu\nu}\cdot\delta\hat{\mathbf{\Phi}}}_{\delta\hat{g}_{\mu\nu}} \;+\; \underbrace{\big(D_{\Phi}t\big)_{\mu\nu}\cdot\delta\hat{\mathbf{\Phi}}}_{\text{part of }\delta\hat{t}_{\mu\nu}} \;+\; \big(D_{\Theta}t\big)_{\mu\nu}\cdot\delta\hat{\mathbf{\Theta}},$$

so the operator excitations $\delta\hat{g}_{\mu\nu}$ and $\delta\hat{t}_{\mu\nu}$ inherit their dynamics from (10.126). The Chapter 11 graviton is obtained by TT projection

$$\hat{h}_{\mu\nu} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}_{\mu\nu} \quad \text{(cf. Definition 10.3 and eqs. (10.103), (10.105), (10.106) and (10.114)).}$$

(iv) Causal support and light–cone structure. Since $\bar{\nabla}^{\mu}\bar{\nabla}_{\mu}$ is constructed from $\bar{\Sigma}_{\mu\nu}$, the corresponding fundamental solutions respect the $\bar{\Sigma}$–causal cones. Accordingly, the matrix propagator $\mathbf{G}_{\Sigma}^{(2)}(x,x')$ (Definition 10.182 and Section 10.17.5) has retarded/advanced support contained in the future/past $\bar{\Sigma}$–domains, consistent with the two–layer causal architecture (Section 5.4.5) and the Green's equation eq. (10.112).

In summary, the gravitational sector emerges from the linearized dynamics of the Gaiamiti substrate. The Apeiron equation (10.126) provides the covariant seed; constitutive response splits that seed into the metric and temporal channels; and TT projection isolates the unique spin–2 Goldstone (the graviton) governed by eq. (10.131), while the tau–kairos remains strictly non–TT (Lemma 10.17, eq. (10.90), and Proposition 10.184) and influences coherence/phase rather than adding GW polarizations.

10.17.8 Gravitational Waves

We formalize gravitational waves entirely within the chronogeometric framework (cf. Sections 5.2.6, 10.17.2 and 10.17.4). We keep explicit the identities of the two key excitations: the *graviton* $\hat{h}_{\mu\nu} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}_{\mu\nu}$ (TT, spin–2; eqs. (10.96) and (10.103) and Definition 10.3) and the *tau–kairos* fluctuation $\hat{\tau}_{\mu\nu} := \delta\hat{t}_{\mu\nu}$ (non–TT; eqs. (10.90) and (10.100) and Lemma 10.17).

Background and linearized split. Let the expectation–level chronogeometry be

$$\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}, \qquad \delta\hat{\Sigma}_{\mu\nu} = \delta\hat{g}_{\mu\nu} + \delta\hat{t}_{\mu\nu}, \tag{10.127}$$

(cf. Section 5.2.6 and eq. (10.95)). The transverse–traceless projector $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ is the Chapter 11 background–covariant projector (Definition 10.3) and satisfies

$$\bar{\nabla}^{\mu}\big(\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}X\big)_{\mu\nu} = 0, \qquad \bar{g}^{\mu\nu}\big(\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}X\big)_{\mu\nu} = 0, \qquad \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\circ\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}. \tag{10.128}$$

Definition 10.190 (Gravitational–wave operator (recall))**.** The *gravitational–wave operator* is the TT projection of the metric fluctuation,

$$\hat{h}_{\mu\nu} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\big[\delta\hat{g}_{\mu\nu}\big],$$

as defined once and for all in eqs. (10.96) and (10.103). By TT–isolation of the temporal channel (eqs. (10.90) and (10.105), Lemma 10.17, and Corollary 10.20),

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\big[\delta\hat{\Sigma}_{\mu\nu}\big] = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\big[\delta\hat{g}_{\mu\nu}\big] = \hat{h}_{\mu\nu}.$$

Corollary 10.191 (TT isolation of the temporal/coherence sector (proved))**.** *The temporal/coherence fluctuation carries no TT content:*

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\big[\delta\hat{t}_{\mu\nu}\big] = 0,$$

as stated in eq. (10.90); equivalently $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\circ D_{\Theta}\Sigma = 0$ *by the constitutive chain rule eq. (10.101). This follows from the Chapter 11 typed projector algebra and TT–purity,* $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\circ\Pi_{t} = 0$ *(Definitions 10.3 and 10.4, Lemma 10.17, and eqs. (10.8) and (10.105)). Moreover, TT–closure implies no TT content arises from off–diagonal blocks (Proposition 10.184 and eq. (10.115)).*

Proposition 10.192 (Kinematic properties)**.** *By eq. (10.128) and the definition eq. (10.96),*

$$\bar{\nabla}^{\mu}\hat{h}_{\mu\nu} = 0, \qquad \bar{g}^{\mu\nu}\hat{h}_{\mu\nu} = 0, \tag{10.129}$$

and $\hat{h}_{\mu\nu}$ *transforms covariantly under background diffeomorphisms induced by* $\bar{g}_{\mu\nu}$*.*

Proposition 10.193 (Goldstone origin (Gaiamiti sourcing)). *With the constitutive identity* $\Sigma_{\mu\nu}[\Phi,\Theta] = g_{\mu\nu}[\Phi] + t_{\mu\nu}[\Phi,\Theta]$ *and the selection rule* $\Pi_{\bar\Sigma}^{[\mathrm{TT}]} \circ D_\Theta \Sigma = 0$ *(cf. Sections 10.17.2 and 10.17.4 and eq. (10.101)), the TT mode is sourced solely by tangential (Goldstone) variations of the Gaiamiti field:*

$$\hat h_{\mu\nu} = \Big(\Pi_{\bar\Sigma}^{[\mathrm{TT}]} \circ D_\Phi g\Big) \cdot \delta\Phi_{\mathrm{tang}}. \tag{10.130}$$

TT wave equation. Let $\delta\bar G_{\mu\nu}$ denote the linearized Einstein tensor built from the background $\bar g_{\mu\nu}$ (indices raised/lowered by $\bar g_{\mu\nu}$). TT–projecting the linearized field equations gives

$$\Pi_{\bar\Sigma}^{[\mathrm{TT}]}\Big[\delta\bar G_{\mu\nu}[\hat h] \;+\; \Lambda\,\hat h_{\mu\nu}\Big] = \kappa\,\Pi_{\bar\Sigma}^{[\mathrm{TT}]}\Big[\delta\bar T^{\mathrm{total}}_{\mu\nu}\Big], \tag{10.131}$$

and in vacuum (or when the TT part of the anisotropic stress vanishes),

$$\Pi_{\bar\Sigma}^{[\mathrm{TT}]}\Big[\delta\bar G_{\mu\nu}[\hat h] \;+\; \Lambda\,\hat h_{\mu\nu}\Big] = 0. \tag{10.132}$$

These equations are consistent with the TT selection rules and null cross–blocks of Chapter 11 (eqs. (10.105) and (10.115) and Proposition 10.184) and with the causal support determined by the unified cones (Section 5.4.5).

Parasymmetric T–L–O augmentation of the GW sector

Definition 10.194 (Typed GW entry and ledger admission). On a certified plateau $\mathfrak{P}$ (cf. Definition 10.1), admit the TT type $\chi = \mathrm{TT}$ via the gate of Definition 10.94. Here χ is the typed ledger tag for the transverse–traceless sector (the field/operator is $\hat h_{\mu\nu}$; cf. eqs. (10.103) and (10.105)). The corresponding typed projector is the Chapter 11 TT projector

$$\Pi^{[\chi]} = \Pi_{\bar\Sigma}^{[\mathrm{TT}]} \quad \text{(cf. Definitions 10.3 and 10.25)}.$$

The sector execution functor $\mathscr{E}_\bullet$ (Definition 9.153) then constructs the typed visibilities $\vec\Lambda^{[\mathrm{TT}]}$, assigns a typed quadrant $\mathrm{Quad}^{[\mathrm{TT}]} \in \{\mathrm{QI},\mathrm{QII},\mathrm{QIII},\mathrm{QIV}\}$, and stamps

$$\mathrm{TLO}^{[\mathrm{TT}]} = \big(\mathrm{Type},\mathrm{Location},\mathrm{Onset}\big),$$

with Location read from diagnostic realized reference configuration metadata (arena/sector/space tag/cone sheet) and Onset $= (r^\star, t^\star)$ extracted from the parasymmetric meters $v^{(a)}_{\mathrm{Synthesis}}$, $v^{(LS)}_{\mathrm{Synthesis}}$, $v^{(LS)}_{\mathrm{Agreement}}$ (cf. Definition 9.153). Typed/triad provenance is preserved throughout (Definitions 10.2 and 10.113 and Proposition 10.10); when a preferred spatial direction exists, the TT tag may be refined to helicity tags $\mathrm{TT}_{\pm2}$ (alias $\gamma_{\pm2}$; cf. Definition 10.26 and Section 10.15).

Proposition 10.195 (Helicity register and stability). *When a* $\bar\Sigma$*-spatial direction is meaningful, refine the TT tag into* $\mathrm{TT}_{\pm2}$ *(cf. Definition 10.25 and Section 10.15) and monitor*

$$\Delta_\chi \;:=\; \Lambda_{\mathrm{state}}^{[\mathrm{TT}_{+2}]} \;-\; \Lambda_{\mathrm{state}}^{[\mathrm{TT}_{-2}]},$$

as in Definition 10.26. A stable nonzero Δ_χ *signals chiral/anisotropic spin–2 response without altering the* count *of tensor polarizations.*

Proposition 10.196 (Law–side guard for extra TT content)**.** *The witness*

$$\mathcal{W}_{\mathrm{TT\text{–}purity}} := \left| \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \circ D_\Theta \Sigma \right|_{\mathfrak{P}}$$

(cf. Lemma 10.17 and eq. (10.101)) monitors TT leakage from the temporal constitutive branch. If $\mathcal{W}_{\mathrm{TT\text{–}purity}} \approx 0$, $\mathrm{TLO}^{[\mathrm{TT}]}$ *exhausts the TT sector. If* $\mathcal{W}_{\mathrm{TT\text{–}purity}} > 0$ *persists in QIII/QIV, extend the typed family (e.g. admit* $\Pi^{[\mathrm{TT}_{\mathrm{mass}}]}$*) and re-run* $\mathcal{E}_\bullet$ *as in Definition 10.32, consistent with the null cross–blocks Proposition 10.184 and eq. (10.115).*

Corollary 10.197 (Closure of the TT sector under T–L–O)**.** *With* $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \circ \Pi_t = 0$ *and* $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \circ D_\Theta \Sigma = 0$ *(eqs. (10.8) and (10.101)), no type* χ *belonging to the temporal/coherence register* $(\tau_S, \tau_A, K_{\mathrm{T}}^{(\tau)}, K_{\mathrm{L}}^{(\tau)})$ *contributes to* $\mathrm{TLO}^{[\mathrm{TT}]}$ *(cf. Lemma 10.59). Thus the TT wave equation (10.131) governs the entire tensor radiative sector, with the graviton operator* $\hat{h}_{\mu\nu} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \delta\hat{g}_{\mu\nu}$ *(eqs. (10.103) and (10.106)).*

Remark 10.198 (Realized vs. pre–physical staging)**.** On the PQS band $[\ell_{\mathrm{PQS}}, \ell_P)$ (Definition 9.153) only operator–level audits are performed (typed hyperbolicity, cone–gating, hygiene). For $\ell \geq \ell_P$ one certifies $\mathfrak{P}$ (Definition 10.1), projects to H^0, and reports physical $\mathrm{TLO}^{[\mathrm{TT}]}$ and helicity diagnostics; BRST/Nielsen safety is inherited from the diagnostic realized reference configuration. Compatibility with causal support of the unified cones follows Section 5.4.5.

Summary and empirical status. Within this framework, *gravitational waves* are the TT–projected metric perturbations

$$\hat{h}_{\mu\nu} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \delta\hat{g}_{\mu\nu} \quad \text{(cf. eqs. (10.96) and (10.103) and Definition 10.3)},$$

massless Goldstone excitations of the Gaiamiti sector that uniquely populate the spin–2 channel; the temporal/coherence tensor obeys $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \hat{\tau}_{\mu\nu} = 0$ (see eq. (10.90) and Lemma 10.17). The GW sector receives a typed, BRST/Nielsen–safe $\mathrm{TLO}^{[\mathrm{TT}]}$ stamp (Type from QI–QIV; Location from diagnostic realized reference configuration metadata; Onset from parasymmetric meters), while law–side TT purity and potential helicity asymmetries are guarded by $\mathcal{W}_{\mathrm{TT\text{–}purity}}$ and Δ_χ, respectively (cf. Lemma 10.17 and Definition 10.26). The linear dynamics are governed by eq. (10.131), consistent with the TT closure and null cross–blocks (Proposition 10.184 and eq. (10.115)) and with typed provenance (Definition 10.113 and Proposition 10.10). Current observations remain consistent with a purely tensorial radiative sector: direct detections and polarization analyses favor two spin–2 GW polarizations [345, 502]; searches for an (approximately) isotropic stochastic background set stringent upper limits on non–tensor admixtures [512, 501]; the near–luminal GW propagation inferred from GW170817 excludes broad classes of models with extra polarizations or modified propagation [513, 514]; and strong–field/post–Newtonian tests support quadrupolar tensor emission as the dominant dissipative channel [515].

10.17.9 Tantrakyma and Kymon Contributions to Chronogeometry

We work on the realized weave configuration with the Chapter 11 typed projectors and triad (Definitions 10.1 to 10.4 and 10.113). Throughout we keep visible the identities of the two

key excitations: the *graviton* $\hat{\gamma}_{\mu\nu} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}_{\mu\nu}$ (spin–2 TT) and the *tau–kairos* fluctuation $\hat{\tau}_{\mu\nu} := \delta\hat{t}_{\mu\nu}$ (non–TT), as recalled in Section 10.17.2 and eq. (10.105).

Kinematic placement. At the expectation level the chronogeometric tensor splits as in (10.94):

$$\boxed{\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}, \qquad \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\bar{t}_{\mu\nu} = 0,} \tag{10.133}$$

so TT content resides entirely in the metric block (Section 10.17.2).

Remark 10.199 (Tantrakyma contribution to the temporal block (weave recall))**.** At the operator level $\hat{t}_{\mu\nu}$ is built from τ/CK typed images (and mixed correlators) defined in the weave (Definition 10.4); $\bar{t}_{\mu\nu} := \langle\hat{t}_{\mu\nu}\rangle$ collects their background profiles. All TT (spin–2) content belongs to the metric channel by (10.105).

Constitutive form and selection rule (recall). With $\Sigma_{\mu\nu}[\Phi,\Theta] = g_{\mu\nu}[\Phi] + t_{\mu\nu}[\Phi,\Theta]$ and linear response (10.101), tau–kairos TT–isolation reads

$$\boxed{\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \circ D_{\Theta}\Sigma \;=\; 0, \qquad \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta t_{\mu\nu} = 0,} \tag{10.134}$$

so Tantrakyma variations populate only non–TT temporal structure.

Tensor structures generated by kymons

Typed τ-kairos decomposition (no new maps). Use the Chapter 11 τ-kairos decomposition (Definition 10.4):

$$\boxed{\hat{\tau}_{\mu\nu} = S^{(b)}_{\mu\nu}[\hat{\varphi}_b] \;\oplus\; S^{(L)}_{\mu\nu}[\hat{\varphi}_L] \;\oplus\; \tilde{S}^{(p)}_{\mu\nu}[\tilde{\varphi}] \;\oplus\; \sum_{\lambda=\pm} A^{(\lambda)}_{\mu\nu}\big[\hat{K}^{(\tau)}_{\lambda}\big], \qquad \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\hat{\tau}_{\mu\nu} = 0,} \tag{10.135}$$

where $A^{(\pm)}$ are CK vector helicities and $S^{(b)}, S^{(L)}, \tilde{S}^{(p)}$ are scalar/pseudoscalar images.

Symmetric vs. antisymmetric temporal tensors (type-only schematics). Writing $t_{(\mu\nu)}$ and $t_{[\mu\nu]}$ for the symmetric/antisymmetric parts, the weave types imply, schematically (type classification only),

$$\text{(i)}\quad t_{(\mu\nu)}: \quad \text{scalar gradients and symmetric CK bilinears,} \quad S^{(\cdot)}_{\mu\nu}[\cdot], \tag{10.136}$$

$$\text{(ii)}\quad t_{[\mu\nu)}: \quad \text{CK field strength } \hat{\mathcal{F}}^{(\tau)}_{\mu\nu} := 2\bar{\nabla}_{[\mu}\hat{K}^{(\tau)}_{\nu]}, \quad A^{(\pm)}_{\mu\nu}[\cdot], \tag{10.137}$$

with $\hat{K}^{(\tau)}_{\mu}$ the Chrono–Kymon vector (see the CK usage in the weave).

Explicit TT annihilation (made concrete). Because (10.135) is built entirely from non–TT typed families,

$$\boxed{\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, t_{(\mu\nu)} = 0, \qquad \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, t_{[\mu\nu]} = 0, \qquad \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \delta\hat{t}_{\mu\nu} = 0,} \tag{10.138}$$

leaving the graviton $\hat{\gamma}_{\mu\nu}$ untouched (purely metric TT).

Remark 10.200 (Chrono–Kymon sourcing of temporal profiles (weave usage))**.** Coherent CK configurations $K_{\mu}^{(\tau)}$ source extended $\bar{t}_{\mu\nu}$ profiles via symmetry–allowed couplings to $\bar{\Sigma}$, $K^{(\tau)} \rightsquigarrow \bar{t}_{\mu\nu}[\bar{\Sigma}; K^{(\tau)}]$, while preserving TT isolation (10.134). The antisymmetric branch is proportional to the CK field strength, $\bar{t}_{[\mu\nu]} = \chi_{\tau}\, \hat{\mathcal{F}}_{\mu\nu}^{(\tau)}$ (CK section, weave).

Corollary 10.201 (Parasymmetric T–L–O accounting for Tantrakyma types)**.** *Running the sector execution with typed/triad provenance (Definitions 9.153 and 10.2) and admission (Definition 10.94) yields, on a certified plateau $\mathfrak{P}$:*

(i) *Typed T–L–O stamps. For each admitted $\chi \in \{S^{(b)}, S^{(L)}, \tilde{S}^{(p)}, A^{(\pm)}\}$ one obtains* $\mathrm{TLO}^{[\chi]} = (\mathrm{Type}, \mathrm{Location}, \mathrm{Onset})$*, with Type determined by the QI–QIV quadrant and Location taken from diagnostic realized reference configuration metadata (Propositions 10.22 and 10.65).*

(ii) *Onset from parasymmetric meters. The onset $(r^{\star}, t^{\star})$ is determined by the parasymmetric meters $v_{\mathrm{Synthesis}}^{(a)}$, $v_{\mathrm{Synthesis}}^{(LS)}$, $v_{\mathrm{Agreement}}^{(LS)}$ (Definition 9.153), defining the coherence windows for temporal responses.*

(iii) *TT nullity of Tantrakyma types. By $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\Pi^{[\chi]} = 0$ and (10.138), no Tantrakyma type contributes to* $\mathrm{TLO}^{[\mathrm{TT}]}$ *(Lemmas 10.17 and 10.59).*

Dynamical implications (framework level)

(a) Variational field equations. Varying the action w.r.t. Σ yields field equations equating curvature functionals of $\bar{\Sigma}$ to total stress–energy from the substrates. The Tantrakyma sector contributes ordinary stress $\bar{T}_{\mu\nu}^{\Theta}$ and coherence stress via $\bar{t}_{\mu\nu}$ whenever that channel is dynamical, inducing a consistent back–reaction on $\bar{\Sigma}$ (consistent with Section 10.17.4).

(b) Closure of the TT sector. Because of (10.138), the TT (spin–2) sector is dynamically closed: off–diagonal metric–temporal couplings affect only non–TT content. Thus the Tantrakyma/kymon sector cannot generate spin–2 excitations; its influence is confined to the temporal/coherence channel and to non–TT back–reaction. TT purity is monitored by the weave witness (Lemma 10.17).

All Tantrakyma/kymon effects enter through the temporal/coherence block $t_{\mu\nu}$: they sculpt $\bar{t}_{\mu\nu}$ and $\hat{\tau}_{\mu\nu}$, contribute to total stress, and leave the TT graviton sector (spin–2, (10.103)) untouched. Parasymmetric T–L–O stamps for $\{S^{(b)}, S^{(L)}, \tilde{S}^{(p)}, A^{(\pm)}\}$ record *when/where* these temporal influences turn on, without adding GW polarizations (Propositions 10.22 and 10.65).

Analytical Symmetry Argument

Set–up (weave kinematics and sources). Use the Chapter 11 fluctuation split and identifications

$$\delta\hat{\Sigma}_{\mu\nu} = \delta\hat{g}_{\mu\nu} + \delta\hat{t}_{\mu\nu} = \hat{h}_{\mu\nu} + \hat{\tau}_{\mu\nu} \quad \text{(cf. (10.95), (10.100))}.$$

The tau–kairos fluctuation is built from the τ-kairos decomposition (Definition 10.4):

$$\boxed{\hat{\tau}_{\mu\nu} = S^{(b)}_{\mu\nu}[\hat{\varphi}_b] \;\oplus\; S^{(L)}_{\mu\nu}[\hat{\varphi}_L] \;\oplus\; \tilde{S}^{(p)}_{\mu\nu}[\tilde{\varphi}] \;\oplus\; \sum_{\lambda=\pm} A^{(\lambda)}_{\mu\nu}\big[\hat{K}^{(\tau)}_{\lambda}\big], \qquad \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\hat{\tau}_{\mu\nu} = 0,} \tag{10.139}$$

i.e. only scalar/pseudoscalar/vector images are present (Lemmas 10.17 and 10.59). Relative to $(\bar{g}, \bar{\nabla})$ one may use the representative trace/longitudinal forms

$$S^{(b)}_{\mu\nu}[\hat{\Phi}] = \frac{1}{d}\,\bar{g}_{\mu\nu}\,\hat{\Phi}, \qquad A_{\mu\nu}[\hat{K}] := \bar{\nabla}_{(\mu}\hat{K}_{\nu)} - \frac{2}{d}\,\bar{g}_{\mu\nu}\,\bar{\nabla}\cdot\hat{K},$$

which span the trace ($\mathbf{0}$) and longitudinal ($\mathbf{1}$) subspaces of Sym^2.

York/Helmholtz orthogonal splitting (weave statement). On $(\bar{g}, \bar{\nabla})$ equip Sym^2 with the L^2 inner product $\langle X, Y\rangle = \int \sqrt{-\bar{g}}\, X^{\mu\nu}Y_{\mu\nu}$ and define the trace–adjusted symmetrized gradient

$$\mathcal{L}[V]_{\mu\nu} := \bar{\nabla}_{(\mu}V_{\nu)} - \frac{2}{d}\,\bar{g}_{\mu\nu}\,\bar{\nabla}\cdot V.$$

Then any $X \in \mathrm{Sym}^2$ admits the orthogonal decomposition

$$\boxed{X_{\mu\nu} = X^{\mathrm{TT}}_{\mu\nu} + \mathcal{L}[V]_{\mu\nu} + \frac{1}{d}\,\bar{g}_{\mu\nu}\,(\mathrm{tr}\,X),} \tag{10.140}$$

with X^{TT} transverse–traceless and V solving $\bar{\nabla}^{\mu}\mathcal{L}[V]_{\mu\nu} = \bar{\nabla}^{\mu}\big(X_{\mu\nu} - \frac{1}{d}\bar{g}_{\mu\nu}\,\mathrm{tr}\,X\big)$. Thus

$$\mathrm{Sym}^2 = \underbrace{\mathrm{TT}}_{\text{spin-2}} \;\oplus\; \underbrace{\mathrm{Im}\,\mathcal{L}}_{\text{longitudinal (spin-1)}} \;\oplus\; \underbrace{\mathrm{span}\{\bar{g}\}}_{\text{trace (spin-0)}},$$

with L^2–orthogonality (integration by parts; standard elliptic arguments).

Projector action and TT isolation. Let $\mathcal{P}^{(i)}$ project onto the three orthogonal subspaces above ($i = 0, 1, 2$). By construction, $\mathcal{P}^{(2)}[\mathrm{Im}\,\mathcal{L}] = 0 = \mathcal{P}^{(2)}[\mathrm{span}\{\bar{g}\}]$; hence from (10.139),

$$\boxed{\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\hat{\tau}_{\mu\nu} = 0 \quad \Rightarrow \quad \hat{\tau}_{\mu\nu} \in \mathbf{0} \oplus \mathbf{1} \;\; \text{(no spin–2 content)}.} \tag{10.141}$$

The TT graviton $\hat{\gamma}_{\mu\nu} = \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\hat{h}_{\mu\nu}$ is therefore purely metric (eq. (10.105)).

Parasymmetric T–L–O overlay. Because $\hat{\tau}_{\mu\nu}$ lives in $\mathbf{0} \oplus \mathbf{1}$, only the tau/CK types $\chi \in \{S^{(b)}, S^{(L)}, \tilde{S}^{(p)}, A^{(\pm)}\}$ are admissible candidates (Definition 10.94). On a certified plateau $\mathfrak{P}$, the execution functor stamps $\mathrm{TLO}^{[\chi]} = (\mathrm{Type}, \mathrm{Location}, \mathrm{Onset})$ (Definition 9.153 and Proposition 10.22), with Onset extracted from the parasymmetric meters; the TT graviton stamp $\mathrm{TLO}^{[\mathrm{TT}]}$ is unaffected (Lemma 10.59). TT purity is monitored by the weave witness (Lemma 10.17). □

Gauge and Variational Argument

Action and symmetries (weave–covariant). Let the Tantrakyma sector be governed by a gauge– and diffeomorphism–covariant action on the background chronogeometry $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$:

$$S_\Theta[\hat{\mathbf{\Theta}}] = \int d^d x \, \sqrt{-\det \bar{\Sigma}}\, \mathrm{Tr}\Big\{ \bar{\Sigma}^{\mu\nu}\, g_C\Big(\bar{\nabla}_\mu \hat{\mathbf{\Theta}}, \bar{\nabla}_\nu \hat{\mathbf{\Theta}}\Big) + \mathcal{V}_\Theta(\hat{\mathbf{\Theta}}) \Big\},$$

with the weave–covariant derivative $\bar{\nabla}$ acting on Θ–registers (weave usage). Euler–Lagrange equations are first order in $\bar{\nabla}\hat{\mathbf{\Theta}}$ and respect gauge/relational symmetries.

Structure of $t_{\mu\nu}$ and induced constraints. The temporal tensor $t_{\mu\nu}[\Phi, \Theta]$ is built from symmetry–allowed bilinears and covariant derivatives of scalar and CK vector registers (plus constitutive dependence on Φ), consistent with the typed images in (10.139). Linearizing about $(\bar{\Phi}, \bar{\Theta})$ yields $\hat{\tau}_{\mu\nu} = \delta\hat{t}_{\mu\nu}$ that decomposes *only* into trace/longitudinal pieces; no divergence–free, trace–free (TT) driver arises.

Projector consequence and TT closure. Applying the same TT projector that defines the graviton, $\hat{\gamma}_{\mu\nu} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \delta\hat{g}_{\mu\nu}$, gives

$$\boxed{\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \hat{\tau}_{\mu\nu} = 0, \qquad \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \delta\hat{\Sigma}_{\mu\nu} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \delta\hat{g}_{\mu\nu} \quad \text{(TT sector purely metric).}} \tag{10.142}$$

Hence the tau–kairos/kymon sector contributes only spin–0/1 structure and cannot source spin–2 radiation; the GW sector remains entirely in the TT metric channel (eq. (10.103)).

Parasymmetric T–L–O perspective (ledger and admission). In the realized regime, the typed ledger admits precisely the tau/CK families that pass hyperbolicity and cone–gating (Definitions 10.1 and 10.94). The execution functor stamps $\mathrm{TLO}^{[\chi]}$ for $\chi \in \{S^{(b)}, S^{(L)}, \tilde{S}^{(p)}, A^{(\pm)}\}$, with Onset from the parasymmetric meters and Location from diagnostic realized reference configuration metadata (Definition 9.153 and Propositions 10.22 and 10.65). Because $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\Pi^{[\chi]} = 0$ for these types (Lemma 10.59), the resulting stamps define *coherence timers* for temporal responses without altering $\mathrm{TLO}^{[\mathrm{TT}]}$. TT purity is monitored by the weave witness (Lemma 10.17); persistent deviations trigger the controlled projector–family extension (weave law–side protocol). □

The Identity of the Geometric Goldstone Boson

Statement (weave view). On the realized weave configuration, the masslessness of the long–range gravitational interaction is *emergent* from SSB in the Gaiamiti sector (no prior masslessness assumption). The Goldstone carrier must be identified within the typed bimodal split of chronogeometric fluctuations,

$$\delta\hat{\Sigma}_{\mu\nu} = \delta\hat{g}_{\mu\nu} + \delta\hat{t}_{\mu\nu} = \hat{h}_{\mu\nu} + \hat{\tau}_{\mu\nu} \quad \text{(cf. (10.95), (10.100))}.$$

Here $\hat{h}_{\mu\nu}$ lives in the metric (Gaiamiti) channel and $\hat{\tau}_{\mu\nu}$ in the temporal/coherence (tau–kairos) channel; the two are typed–orthogonal.

The two candidates (typed identifications)

(i) Metric fluctuation (graviton) Its physical spin–2 content is the TT graviton field (Definitions 10.3 and 10.25):

$$\boxed{\hat{h}_{\mu\nu} \;:=\; \mathcal{P}^{(2)}_{\text{grav}}\,\delta\hat{g}_{\mu\nu} \;=\; \Pi^{[\text{TT}]}_{\bar{\Sigma}}\,\delta\hat{g}_{\mu\nu},} \tag{10.143}$$

with propagation captured by the TT–typed ΦΦ correlator blocks $\hat{\mathbf{G}}^{[\Phi\to\Phi;\text{TT}]}_{\Lambda;a}$ (triad $a \in \{\text{spec}, \text{vac}, \text{corr}\}$; cf. Definition 10.158).

(ii) Tau–kairos fluctuation By TT–nullity, $\Pi^{[\text{TT}]}_{\bar{\Sigma}}\,\hat{\tau}_{\mu\nu} = 0$ (cf. eq. (10.97) and Definition 10.162), so it carries only scalar/pseudoscalar/vector structure (no spin–2) in the weave τ/CK families (Definitions 10.4 and 10.57).

Verdict from spin and sourcing

A conserved rank–2 source selects a *massless spin–2* carrier at low energies; in the weave this carrier is precisely the TT graviton field extracted from the metric fluctuation, $\hat{h}_{\mu\nu} = \mathcal{P}^{(2)}_{\text{grav}}\delta\hat{g}_{\mu\nu} = \Pi^{[\text{TT}]}_{\bar{\Sigma}}\delta\hat{g}_{\mu\nu}$ (Definitions 10.3 and 10.25, cf. eq. (10.143)). Since tau–kairos is TT–null ($\Pi^{[\text{TT}]}_{\bar{\Sigma}}\hat{\tau}_{\mu\nu} = 0$; Definition 10.162, Lemma 10.17, and eq. (10.97)), it cannot realize a spin–2 Goldstone nor mediate classical gravitational radiation. All radiative tensor content is confined to the TT channel and recorded in the TT–typed ΦΦ blocks (Definition 10.158 and Theorem 10.169).

Channel–resolved Goldstone identity (triad decomposition)

Decompose the *metric fluctuation* by triad provenance (Definition 10.2):

$$\delta\hat{g}_{\mu\nu} = \sum_{a\in\{\text{spec,vac,corr}\}} \delta\hat{g}^{(a)}_{\mu\nu}, \qquad \hat{h}^{(a)}_{\mu\nu} := \Pi^{[\text{TT}]}_{\bar{\Sigma}}\,\delta\hat{g}^{(a)}_{\mu\nu}.$$

Then the graviton field splits as $\hat{h}_{\mu\nu} = \sum_a \hat{h}^{(a)}_{\mu\nu}$ with each $\hat{h}^{(a)}_{\mu\nu}$ captured in the TT–typed ΦΦ correlator blocks $\hat{\mathbf{G}}^{[\Phi\to\Phi;\text{TT}]}_{\Lambda;a}$ (Definition 10.158). Under the Goldstone origin of the TT mode in the Gaiamiti sector (tangential directions), each channel component is massless

(Propositions 9.197 and 10.173). The tau–kairos sector stays typed–orthogonal to TT and contributes only scalar/vector observables in non–TT blocks (no TT leakage; Definition 10.162, eq. (10.97), and Theorem 10.169).

Alignment with the parasymmetric T–L–O ledger. Typed projectors implement the separation between graviton and tau–kairos channels. The graviton uses the TT type with a $\mathrm{TLO}^{[\mathrm{TT}]}$ stamp (Type/Location/Onset; Proposition 10.90 and Definitions 10.135 and 10.136), while tau–kairos entries $\chi \in \{S^{(b)}, S^{(L)}, \tilde{S}^{(p)}, A^{(\pm)}\}$ are stamped in their own branches and satisfy $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}} \Pi^{[\chi]} = 0$ (Lemmas 10.17 and 10.59), so they never contribute to $\mathrm{TLO}^{[\mathrm{TT}]}$. Meters/visibilities are BRST/Nielsen–safe on the realized weave configuration (Definitions 10.1 and 10.82 and Lemma 10.83), monotone with unique onsets (Propositions 9.146 and 10.88 and Lemma 10.86), and reported in the quadrant map (Section 10.14).

Consequences. (i) The number of propagating tensor polarizations remains two, entirely in the TT graviton sector (Definitions 10.3 and 10.25); (ii) triad dependence enters only through amplitudes/phases and sourcing inside the TT blocks $\hat{\mathbf{G}}^{[\Phi\to\Phi;\mathrm{TT}]}_{\Lambda;a}$ (no creation of extra TT modes; Definition 10.158 and Proposition 10.9); (iii) tau–kairos provides non–radiative scalar/vector responses that shape temporal/coherence structure and background transport, with propagation causally gated by $\bar{\Sigma}$ (Definition 6.7 and Remark 9.83).

10.18 Matter Block of the Unified Correlator

We describe the self–correlations of the Tantrakyma field as the lower–right (matter) block of the Unified Inter–Modal Correlator, using typed insertion and triad provenance on the realized weave configuration (Definitions 10.1, 10.2 and 10.113). This block captures the propagation of matter–like degrees of freedom—*kymons*—across the emergent configuration space $\mathcal{X}_q$ and underwrites particle phenomenology within the PQS program, while remaining TT–orthogonal to the graviton sector.

10.18.1 Definition and Typed Placement

Definition 10.202 (Matter block via typed insertion)**.** Let $\hat{\mathbf{G}}_\Lambda$ denote the Unified Inter–Modal Correlator (weave form). The Tantrakyma (matter) block is

$$\boxed{\hat{\mathbf{G}}^{[\Theta\to\Theta]}_{\Lambda} := \Pi^{[\Theta]}_{\Lambda}\, \hat{\mathbf{G}}_\Lambda\, \Pi^{[\Theta]}_{\Lambda}, \qquad \hat{\mathbf{G}}^{[\Theta\to\Theta]}_{\Lambda} = \bigoplus_{a\in\{\mathrm{spec,vac,corr}\}} \hat{\mathbf{G}}^{[\Theta\to\Theta]}_{\Lambda;a},} \tag{10.144}$$

with triad idempotents $\Pi^{(a)}$ giving $\hat{\mathbf{G}}^{[\Theta\to\Theta]}_{\Lambda;a} := \Pi^{(a)}\, \hat{\mathbf{G}}^{[\Theta\to\Theta]}_{\Lambda}\, \Pi^{(a)}$.

Definition 10.203 (Matter two–point kernel (QHS, connected))**.** Let $\hat{\mathbf{\Theta}}(x)$ be the Tantrakyma field operator and $\langle\cdot\rangle_\circ$ the QHS expectation on the realized weave configuration (Definitions 1.27 and 10.1). The *connected, time–ordered* matter kernel is

$$\boxed{\hat{G}_{\Theta\Theta}(x,x') := \Big\langle T\{\hat{\mathbf{\Theta}}(x)\,\hat{\mathbf{\Theta}}(x')\}\Big\rangle_\circ - \Big\langle\hat{\mathbf{\Theta}}(x)\Big\rangle_\circ \Big\langle\hat{\mathbf{\Theta}}(x')\Big\rangle_\circ,} \tag{10.145}$$

where $T\{\cdot\}$ denotes the *relational* time–ordering associated with the background $\bar{\Sigma}$ (i.e. chronological ordering along the $\bar{\Sigma}$–compatible time function; cf. Definitions 6.7 and 9.80 and Remark 9.83). In the position basis, $\hat{G}_{\Theta\Theta}(x,x')$ coincides with the kernel of the $\Theta\Theta$ block of the Unified Inter–Modal Correlator (Definition 6.4), as recorded in (10.144).

Explanation (time ordering and connectedness). The operator $T\{\cdot\}$ orders insertions by the relational time defined on the $\bar{\Sigma}$–geometry, so that the resulting two–point function matches the causal cones of $\bar{\Sigma}$ (hence the associated retarded/advanced kernels are Σ–causal; Remark 9.83). Subtracting the product of one–point functions makes the kernel *connected*; by the same algebra used for the geometric sector (eqs. (10.88) and (10.89)), this quantity equals the covariance of $\hat{\boldsymbol{\Theta}}$–fluctuations in the QHS state. Finally, identifying $\hat{G}_{\Theta\Theta}$ with the $\Theta\Theta$ block's kernel ties this definition directly to the unified correlator architecture (Definition 6.4), ensuring that typed insertions and triad provenance in Chapter 11 apply verbatim to the matter sector as well (Definitions 10.2 and 10.113).

Physical meaning. $\hat{G}_{\Theta\Theta}$ is the *kymon propagator* in the sense that its analytic structure (poles, residues, and continuous parts) encodes the emergent mass spectrum, widths, and mixing of matter excitations on the physical carrier $\mathcal{H}_{\text{phys}}$ (Definitions 10.1 and 10.203). On admitted bands, masses and quasi-particle projectors are isolated by the Riesz calculus applied to the gauge–fixed kinetic generalized eigen-operator pair in the $\Theta\Theta$ block (Definition 10.7), while spectral positivity is controlled by the Källén–Lehmann ledger on $\mathcal{H}_{\text{phys}}$ (cf. eq. (7.12) and scalar/vector specializations such as Proposition 10.77). Typed resolution sharpens this reading: scalar content (e.g. radial/Higgs and trace-like modes) is refined by CP projectors $\Pi^{(\text{CP}\pm)}$ (Definition 10.5), and vector content (Chrono–Kymon) is helicity- and parity-resolved by $A^{(\pm)}$ with diagnostic Δ_P (Definitions 10.57 and 10.58). Because the kernel is defined with $\bar{\Sigma}$-relational time ordering (Definition 10.203), its retarded/advanced parts are Σ-causal (support inside $\bar{\Sigma}$-cones; Remark 9.83), ensuring a consistent notion of propagation on the realized chronogeometric background (Definitions 6.7 and 9.80).

Stage vs. actors (and how we report them). Within the Unified Inter–Modal Correlator (Definition 6.4), the geometric block $\hat{\mathbf{G}}_{\Lambda}^{[\Phi\to\Phi]}$ sets the *stage* by determining the realized chronogeometry $\bar{\Sigma}$ and the TT graviton channel (TT type; Definitions 10.25, 10.157 and 10.158), whereas the matter block $\hat{\mathbf{G}}_{\Lambda}^{[\Theta\to\Theta]}$ governs the *actors*—the kymons—through its typed spectral content (Definitions 10.2 and 10.113). Inter–sector couplings appear only through the off–diagonal $\Phi\Theta$ entries and are audited via typed insertion/mixing rules and selection constraints (parity/helicity/CP) so that no forbidden leakage contaminates the TT graviton readout (Proposition 10.126, Lemma 10.123, and Theorem 10.169). Operationally, both blocks feed the same Law/State ledgers and calibrated meters (Definitions 10.80 and 10.82), yielding finite, monotone visibilities with unique onsets (Lemmas 10.83 and 10.86 and Propositions 9.146 and 10.88). In reports, the kymon claims are issued as $\text{TLO}^{[\chi]}$ for the appropriate $\Theta\Theta$ types (e.g. rad$\pm$, $A^{(\pm)}$), while the graviton is certified solely in the TT $\Phi\Phi$ branch—cleanly separating the *actors* from the *stage* within the same weave scaffold (Definitions 10.135 and 10.136).

10.18.2 Background–fluctuation Decomposition

Definition 10.204 (Tantrakyma background–fluctuation split)**.** Relative to the Quantum Holo–State,

$$\boxed{\hat{\mathbf{\Theta}}(x) \;=\; \bar{\Theta}(x)\,\mathbb{I}_x \;+\; \delta\hat{\mathbf{\Theta}}(x), \qquad \langle \delta\hat{\mathbf{\Theta}}(x)\rangle_\circ = 0.} \tag{10.146}$$

Proposition 10.205 (Fluctuation reduction)**.** *With* (10.146)*,*

$$\boxed{\hat{G}_{\Theta\Theta}(x,x') \;=\; \Big\langle \delta\hat{\mathbf{\Theta}}(x)\,\delta\hat{\mathbf{\Theta}}(x')\Big\rangle_\circ .} \tag{10.147}$$

Proof. Insert $\hat{\mathbf{\Theta}} = \bar{\Theta}\,\mathbb{I} + \delta\hat{\mathbf{\Theta}}$ into (10.145) and use $\langle\delta\hat{\mathbf{\Theta}}\rangle_\circ = 0$. □

10.18.3 Typed resolution and admissible matter channels

Definition 10.206 (Typed matter correlators in the τ/CK families)**.** Using the Chapter 11 tau–kairos/kymon types (Definition 10.4),

$$\boxed{\hat{\mathbf{G}}^{[\Theta\to\Theta;\chi]}_{\Lambda;a} \;:=\; \Pi^{(a)}\,\Pi^{[\chi]}\,\hat{\mathbf{G}}^{[\Theta\to\Theta]}_{\Lambda}\,\Pi^{[\chi]}\,\Pi^{(a)}, \qquad \chi\in\{S^{(b)},S^{(L)},\tilde{S}^{(p)},A^{(\pm)}\},} \tag{10.148}$$

which provide the scalar/pseudoscalar/vector kymon channels (no TT type appears in the Tantrakyma family).

Corollary 10.207 (Positivity and cone–gating on $\mathfrak{P}$)**.** *On a certified plateau $\mathfrak{P}$, each admitted typed block $\hat{\mathbf{G}}^{[\Theta\to\Theta;\chi]}_{\Lambda;a}$ is a positive semidefinite covariance on equal–time test functions of type $[\chi]$ and propagates within the $\bar{\Sigma}$–cones (admission & hyperbolicity: Definitions 10.1 and 10.94).*

Interpretation (ledger view). The typed blocks (10.148) feed the parasymmetric pipeline; the execution functor stamps $\mathrm{TLO}^{[\chi]}$ for admitted χ with Location inherited from diagnostic realized reference configuration metadata and Onset from the meters (Definition 9.153 and Propositions 10.22 and 10.65). TT–purity of the graviton sector is unaffected because $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\Pi^{[\chi]} = 0$ for these matter types (Lemmas 10.17 and 10.59).

10.18.4 Relations to cross–blocks and spectrum

Cross–block coupling. Kymon propagation couples to geometry through the *typed off–diagonal* blocks $\hat{\mathbf{G}}^{[\Phi\to\Theta]}_{\Lambda;a}$ and $\hat{\mathbf{G}}^{[\Theta\to\Phi]}_{\Lambda;a}$, built by typed insertion with triad provenance (Definitions 10.2 and 10.113). These kernels encode how localized Θ–sector transitions source nonlocal geometric responses (and conversely), while preserving band/BRST hygiene on the realized weave configuration (Definition 10.1 and Proposition 10.10) and Σ–causality with respect to the realized background $\bar{\Sigma}$ (Definition 9.80 and Remark 9.83). Crucially, TT isolation is not violated: any non–TT typed insertion from the τ–kairos/CK families is annihilated by the background–covariant TT projector $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ (Definition 10.3, Lemma 10.123,

and Theorem 10.169), so graviton readouts in the TT channel remain uncontaminated even in the presence of $\Phi\Theta$ mixing. Symmetry selection rules (parity/helicity/CP) further gate which mixed couplings are admissible (Proposition 10.126).

Spectral data. The analytic structure of the connected, time–ordered matter kernel $\hat{G}_{\Theta\Theta}(x, x')$ (Definition 10.203) in suitable spectral variables determines kymon masses, widths, and mixing angles: isolated poles and their Riesz projectors select species, while residues encode couplings and overlaps (Definition 10.7 and Proposition 10.149). On $\mathcal{H}_{\text{phys}}$ the Källén–Lehmann ledger ensures nonnegative spectral densities (cf. eq. (7.12) and scalar/vector specializations such as Proposition 10.77). Triad labels $a \in \{\text{spec}, \text{vac}, \text{corr}\}$ refine these observables by provenance within the same typed ledger (Definition 10.2), separating on–band radiative contributions from vacuum/coherence structure. Typed refinements (e.g. CP for the Higgs–like radial modes, helicity/parity for CK vectors) are implemented by the corresponding idempotents (Definitions 10.5, 10.57 and 10.58), and future spectral branches are incorporated automatically by enlarging the Riesz sums (future–proofing; cf. Definition 10.7 and Lemma 10.76).

Summary. The matter block $\hat{\mathbf{G}}_{\Lambda}^{[\Theta\to\Theta]}$ captures kymon dynamics via the connected fluctuation kernel (10.147) (equivalently Definition 10.203), and its typed decomposition (10.148) provides the admissible scalar/pseudoscalar/vector channels on the realized weave configuration (Definition 10.1 and Proposition 10.10). These integrate with the T–L–O pipeline for certified claims (Proposition 10.90 and Definitions 10.135 and 10.136) and remain *TT–orthogonal* to the graviton sector by construction (Definition 10.3, Lemmas 10.17 and 10.59, and Theorem 10.169).

10.18.5 Emergent Matter Propagators

The connected Tantrakyma kernel $\hat{G}_{\Theta\Theta}(x, x')$ (cf. (10.147)) encodes the full dynamical content of the matter/coherence sector on $\mathcal{X}_q$. It is a matrix–valued bi–distribution on the modal fibres that governs the quantum propagation of internal excitations and, via typed projections, yields the propagators of physical kymons (Section 10.18.3). We make this structure explicit by writing $\hat{G}_{\Theta\Theta}$ in a local operator basis and then projecting to physical (mass/interaction) eigenmodes. Background material on curved–space two–point functions follows the standard references.

Fluctuation split and connectedness (Tantrakyma). Write the realized background–fluctuation split for the matter operator (cf. the geometric split (10.87) and the matter kernel definition Definition 10.203):

$$\boxed{\hat{\mathbf{\Theta}}(x) \;=\; \big\langle \hat{\mathbf{\Theta}}(x) \big\rangle_{\circ}\, \mathbb{I}_x \;+\; \delta\hat{\mathbf{\Theta}}(x), \qquad \big\langle \delta\hat{\mathbf{\Theta}}(x) \big\rangle_{\circ} = 0.} \tag{10.149}$$

With relational time–ordering on the $\bar{\Sigma}$ background (Definitions 6.7 and 9.80 and Remark 9.83), the *connected, time–ordered* kernel equals the fluctuation covariance:

$$\boxed{\begin{aligned}\hat{G}_{\Theta\Theta}(x,x') &= \Big\langle T\{\hat{\boldsymbol{\Theta}}(x)\,\hat{\boldsymbol{\Theta}}(x')\}\Big\rangle_{\circ} - \Big\langle \hat{\boldsymbol{\Theta}}(x)\Big\rangle_{\circ}\Big\langle \hat{\boldsymbol{\Theta}}(x')\Big\rangle_{\circ} && (10.150)\\ &= \Big\langle T\{\delta\hat{\boldsymbol{\Theta}}(x)\,\delta\hat{\boldsymbol{\Theta}}(x')\}\Big\rangle_{\circ}. && (10.151)\end{aligned}}$$

(The geometric analogue is eqs. (10.88) and (10.89).)

Definition 10.208 (Tantrakyma propagator matrix (modal basis))**.** Let the Tantrakyma *fluctuation* admit a local operator expansion on the modal fibre $\mathrm{End}(\mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}})$:

$$\delta\hat{\boldsymbol{\Theta}}(x) = \sum_{a,b} \delta\theta^{ab}(x)\,|\mathbf{a}\rangle_x\langle\mathbf{b}|_x, \qquad \{|\mathbf{a}\rangle_x\} \subset \mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}} \text{ orthonormal.}$$

The *Tantrakyma propagator matrix* is the time–ordered two–point matrix of fluctuation coefficients

$$\boxed{\hat{G}^{ab,cd}_{\Theta\Theta}(x,x') \;:=\; \Big\langle T\{\delta\theta^{ab}(x)\,\delta\theta^{cd}(x')\}\Big\rangle_{\circ},} \tag{10.152}$$

which are the components of the connected matter kernel $\hat{G}_{\Theta\Theta}(x,x')$ from Definition 10.203 in the $(ab|cd)$ modal basis. In matrix form, these entries are precisely the position/modal representation of the $\Theta\Theta$ block of the Unified Inter–Modal Correlator in (10.144).

Fibre covariance and reciprocity. Because the modal fibre $\mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}}$ varies with the locus x, contractions between coefficients at (x,ab) and (x',cd) are made covariant by parallel transporting the (x',cd) data to x using the inter–modal transporter $\mathcal{U}(x\leftarrow x';\gamma)$ from the Fourfold/Unified construction (Definition 4.9 and Section 4.1.4). Define the transported fluctuation coefficient

$$\tilde{\theta}^{\,cd}(x'\to x) := \mathcal{U}(x\leftarrow x';\gamma)\,\theta^{\,cd}(x')\,\mathcal{U}(x'\leftarrow x;\gamma),$$

so that the *covariant* representation of the matter propagator matrix (Definition 10.208) reads

$$\hat{G}^{ab,cd}_{\Theta\Theta}(x,x') \;=\; \Big\langle T\{\theta^{ab}(x)\,\tilde{\theta}^{\,cd}(x'\to x)\}\Big\rangle_{\circ},$$

which is equivalent to (10.152) when working at a common fibre. For neutral registers (no internal charge), Hermitian reciprocity holds:

$$\hat{G}^{ab,cd}_{\Theta\Theta}(x,x') \;=\; \overline{\hat{G}^{cd,ab}_{\Theta\Theta}(x',x)}.$$

When $\hat{G}_{\Theta\Theta}$ is the inverse of the *gauge–fixed, band–limited* second–variation (kinetic) operator $K_{\Theta\Theta}$ of the Tantrakyma action on the realized weave configuration (Definition 10.1 and Proposition 10.10), it obeys the matrix Green equation in the matter sector,

$$\sum_{c',d'} \big(K_{\Theta\Theta}\big)^{ab}{}_{c'd'}(x)\,\hat{G}^{c'd',cd}_{\Theta\Theta}(x,x') \;=\; \delta^{(d)}(x,x')\,\delta^a_{\,c}\,\delta^b_{\,d},$$

with triad resolution obtained by inserting $\Pi^{(a)}$ on both sides (Definition 10.2) and consistent with the $\Theta\Theta$ block relation (10.144). The time ordering $T\{\cdot\}$ is taken with respect to the relational background $\bar{\Sigma}$ (Definitions 9.80 and 10.203 and Remark 9.83), ensuring Σ–causal propagation of the matter modes.

Typed projections (matter channels). Using the Chapter 11 tau/CK types, the typed matter propagators are the blocks

$$\boxed{\hat{\mathbf{G}}^{[\Theta\to\Theta;\chi]}_{\Lambda;a} \;=\; \Pi^{(a)}\,\Pi^{[\chi]}\,\hat{\mathbf{G}}^{[\Theta\to\Theta]}_{\Lambda}\,\Pi^{[\chi]}\,\Pi^{(a)}, \qquad \chi\in\{S^{(b)},S^{(L)},\tilde{S}^{(p)},A^{(\pm)}\},} \tag{10.153}$$

(cf. (10.148)). No TT type appears in the Tantrakyma family; TT purity is protected by $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\Pi^{[\chi]}=0$ (Lemmas 10.17 and 10.59).

Definition 10.209 (Projection onto emergent particles (kymons))**.** Physical species p need not align with a single modal operator; they are coherent superpositions that (locally in x) diagonalize the quadratic kernel of the Tantrakyma action at the relevant scale (Riesz selection on the realized weave configuration; Definitions 10.1 and 10.7). Adopt the fluctuation split for the matter operator (Definition 10.203) and expand

$$\delta\hat{\mathbf{\Theta}}(x)=\sum_{a,b}\theta^{ab}(x)\,|\mathbf{a}\rangle_x\langle\mathbf{b}|_x,$$

i.e. the δ sits on the operator, while the modal coefficients are $\theta^{ab}(x)$ (no δ). Choose modal coefficients $u^{ab}_p(x)$ (columns of a unitary change of basis on the operator fibre, relationally covariant via parallel transport; Section 4.1.4) and set

$$\theta^{(p)}(x)\;:=\;\sum_{a,b}u^{ab}_p(x)\,\theta^{ab}(x).$$

The *projected two–point function* (the kymon–p propagator) is

$$\boxed{\Delta_p(x,x')\;:=\;\Big\langle T\{\theta^{(p)}(x)\,\overline{\theta^{(p)}(x')}\}\Big\rangle_{\circ}\;=\;\sum_{a,b,c,d}u^{ab}_p(x)\,\hat{G}^{ab,cd}_{\Theta\Theta}(x,x')\,\overline{u^{cd}_p(x')},} \tag{10.154}$$

where $\hat{G}^{ab,cd}_{\Theta\Theta}(x,x')$ are the modal components of the connected matter kernel from Definitions 10.203 and 10.208. For full fibre covariance one may first transport the (x',cd) data to x using $\mathcal{U}(x\leftarrow x';\gamma)$ (cf. Section 4.1.4), which yields an equivalent contraction (see the "Fibre covariance and reciprocity" paragraph). Typed/triad tags propagate verbatim through (10.154) via the idempotents of Definitions 10.2 and 10.113.

Modal vs. physical bases; mixing. The modal indices (a,b) label the raw operator basis on $\mathrm{End}(\mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}})$; the physical labels $\{p\}$ result from diagonalizing the quadratic kernel (mass/mixing) on the realized background (Definitions 10.203 and 10.208). Off–diagonal entries of $\hat{G}^{ab,cd}_{\Theta\Theta}(x,x')$ with $(a,b)\neq(c,d)$ encode mixing/oscillation phenomena in the modal basis, while the projected $\Delta_p(x,x')$ of Definition 10.209 isolates the propagator of the mass eigenmode p. In stationary/homogeneous arenas, Δ_p admits a Källén–Lehmann–type representation (connected, time–ordered kernel; Definition 10.203) whose pole structure yields physical masses and dispersion.

Triad provenance and ledger integration. Triad idempotents $\Pi^{(a)}$ resolve the matter kernel as $\hat{G}_{\Theta\Theta} \mapsto \hat{G}_{\Theta\Theta;a} := \Pi^{(a)}\,\hat{G}_{\Theta\Theta}\,\Pi^{(a)}$ with provenance $a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$ (Definition 10.2), in the same typed/triad manner as (10.153). These typed/triad blocks feed the parasymmetric pipeline; the execution functor stamps $\mathrm{TLO}^{[\chi]}$ for admitted χ with Location from diagnostic realized reference configuration metadata and Onset from the calibrated meters (Definitions 9.153 and 10.82 and Propositions 10.22 and 10.65). TT selection in the gravitational sector remains unchanged:

$$\boxed{\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{\Sigma}_{\mu\nu} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}_{\mu\nu}, \qquad \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{t}_{\mu\nu} = 0,} \tag{10.155}$$

with $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ the background–covariant TT projector (Definition 10.3); thus matter propagators backreact only through non–TT channels while leaving the graviton TT block intact (cf. (10.105), Lemma 10.17).

Summary. $\hat{G}_{\Theta\Theta}$ furnishes the kymon propagators as a fibre–covariant matrix kernel (10.152); typed/triad projections (10.153) produce the admissible scalar/pseudoscalar/vector channels; and physical kymons are obtained by the local rotations (10.154). This integrates seamlessly with the parasymmetric ledger while preserving TT purity of the graviton sector.

10.18.6 Dynamical Equations and Propagation

The Tantrakyma fluctuation $\delta\hat{\mathbf{\Theta}}(x)$ (matter/coherence excitations on the realized weave configuration) obeys a covariant operator wave equation obtained from the functional Euler–Lagrange principle applied to the second–variation (quadratic) action, using the background–fluctuation split $\hat{\mathbf{\Theta}} = \bar{\Theta}\,\mathbb{I} + \delta\hat{\mathbf{\Theta}}$ (cf. Definition 10.204). All operators are built with the background chronogeometry $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ and the weave–covariant derivative $\bar{\nabla}$ (cf. Section 10.17.2).

Quadratic action and kinetic operator. With $\bar{\nabla}$ compatible with $\bar{\Sigma}$ when stated, the linearised action reads

$$\boxed{S^{(2)}[\delta\hat{\mathbf{\Theta}}] = \int d^dx\,\sqrt{-\det\bar{\Sigma}}\;\mathrm{Tr}\Big\{\bar{\Sigma}^{\mu\nu}\big(\bar{\nabla}_\mu\delta\hat{\mathbf{\Theta}}\big)^\dagger\big(\bar{\nabla}_\nu\delta\hat{\mathbf{\Theta}}\big) - \tfrac{1}{2}\,\delta\hat{\mathbf{\Theta}}^\dagger\,\mathcal{M}_\Theta(x)\,\delta\hat{\mathbf{\Theta}}\Big\},} \tag{10.156}$$

where the operator–valued mass term is the Hessian of the Tantrakyma potential about the background,

$$\boxed{\mathcal{M}_\Theta(x) := \left.\frac{\delta^2\mathcal{V}_\Theta}{\delta\hat{\mathbf{\Theta}}^\dagger\,\delta\hat{\mathbf{\Theta}}}\right|_{\bar{\Theta}(x)}.} \tag{10.157}$$

Varying (10.156) yields the *normally hyperbolic* matter operator

$$\boxed{\mathcal{K}_\Theta := -\,\bar{\nabla}_\mu\big(\bar{\Sigma}^{\mu\nu}\bar{\nabla}_\nu\,\cdot\,\big) + \mathcal{M}_\Theta(x),} \tag{10.158}$$

so the Euler–Lagrange equation takes the compact form

$$\boxed{\mathcal{K}_\Theta(x)\,\delta\hat{\boldsymbol{\Theta}}(x) = 0.} \tag{10.159}$$

If $\bar{\nabla}_\lambda \bar{\Sigma}^{\mu\nu} = 0$ (chronogeometric compatibility), then $\mathcal{K}_\Theta = -\bar{\Sigma}^{\mu\nu}\bar{\nabla}_\mu\bar{\nabla}_\nu + \mathcal{M}_\Theta$. Hyperbolicity and cone–gating of (10.159) are enforced by admission (Definitions 10.1 and 10.94).

Green functions and propagators. The modal matter kernel $\hat{G}^{ab,cd}_{\Theta\Theta}(x,x')$ of Definitions 10.203 and 10.208 is a Green function for the (gauge–fixed, band–limited) second–variation operator $K_{\Theta\Theta}$ on the realized weave configuration (Definition 10.1 and Proposition 10.10). In components:

$$\boxed{\sum_{c',d'}\Big(K_{\Theta\Theta}\Big)^{ab}{}_{c'd'}(x)\,\hat{G}^{c'd',cd}_{\Theta\Theta}(x,x') = \delta^{(d)}(x,x')\,\delta^a_c\,\delta^b_d.} \tag{10.160}$$

Boundary data (retarded/advanced/Feynman) are chosen as appropriate, and the time ordering is taken with respect to the relational background $\bar{\Sigma}$ (see Definitions 9.80 and 10.203 and Remark 9.83).

Projected (species) propagators Δ_p are obtained by the rotation formula (10.154), which contracts $\hat{G}^{ab,cd}_{\Theta\Theta}(x,x')$ with the local modal-to-physical coefficients. In stationary/homogeneous settings, Δ_p admits a Källén–Lehmann–type spectral representation (connected, time-ordered kernel of Definition 10.203) whose poles furnish masses and dispersion relations.

Proposition 10.210 (Quadratic reduction and Σ–causal propagation on the realized weave configuration)**.** *Let $\hat{\boldsymbol{\Theta}} = \bar{\Theta}\,\mathbb{I} + \delta\hat{\boldsymbol{\Theta}}$ be the background–fluctuation split (Definition 10.204) on the realized weave configuration $\mathcal{I}^{[\mathcal{S}]}$ (Definition 10.1). Expanding the full Tantrakyma action about $\bar{\Theta}$ yields*

$$\boxed{S[\bar{\Theta} + \delta\hat{\boldsymbol{\Theta}}] = S[\bar{\Theta}] + \underbrace{S^{(1)}[\delta\hat{\boldsymbol{\Theta}}]}_{=\,0 \text{ if } \bar{\Theta} \text{ is a stationary point}} + S^{(2)}[\delta\hat{\boldsymbol{\Theta}}] + \mathcal{O}\big(\delta\hat{\boldsymbol{\Theta}}^{\,3}\big).} \tag{10.161}$$

When $\bar{\Theta}$ satisfies the background equations, the linear term vanishes and the leading dynamics of fluctuations are governed by the quadratic action $S^{(2)}$ in (10.156). Its Euler–Lagrange operator is the normally hyperbolic kinetic operator $\mathcal{K}_\Theta$ of (10.158), so that the linearized equation of motion reads

$$\boxed{\mathcal{K}_\Theta(x)\,\delta\hat{\boldsymbol{\Theta}}(x) = 0 \qquad \textit{(cf. eq. (10.159)).}} \tag{10.162}$$

Moreover, the connected, time–ordered matter kernel $\hat{G}_{\Theta\Theta}$ (Definition 10.203) is a Green function of $\mathcal{K}_\Theta$ on $\mathcal{H}_{\text{phys}}$, satisfying the matrix equation (10.160) with retarded/advanced/Feynman boundary data chosen relative to $\bar{\Sigma}$ (Definition 9.80 and Remark 9.83).

Proof (sketch). Taylor expand the action at $\bar{\Theta}$ to second order; stationarity of $\bar{\Theta}$ kills $S^{(1)}$, yielding $S^{(2)}$ in (10.156) with mass/Hessian $\mathcal{M}_\Theta$ from (10.157). Varying $S^{(2)}$ gives $\mathcal{K}_\Theta$

in (10.158) and the equation of motion (10.162). On the realized weave configuration (Definition 10.1), admissibility/invariance (Proposition 10.10) and $\bar{\Sigma}$-compatibility ensure that $\mathcal{K}_\Theta$ is normally hyperbolic and Σ–causal (Remark 9.83), so its inverse in the appropriate sense is the connected two–point function $\hat{G}_{\Theta\Theta}$, which indeed solves (10.160). □

Why the quadratic action $S^{(2)}$ appears. In the linearized regime around the realized background $\bar{\Theta}$ and $\bar{\Sigma}$, the first nontrivial term of the action expansion is $S^{(2)}$ (eq. (10.161)). This term *defines* the kinetic operator $\mathcal{K}_\Theta$ and the Hessian block $\mathcal{M}_\Theta$ (eqs. (10.157) and (10.158)), leading to a well-posed, Σ–causal propagation equation on $\mathcal{H}_{\text{phys}}$ (eq. (10.159) and Remark 9.83). Consequently, the connected matter kernel $\hat{G}_{\Theta\Theta}$ (Definition 10.203) is the Green function of $\mathcal{K}_\Theta$ (eq. (10.160)), which is the precise object used downstream for spectral analysis and reporting.

Where higher variations live (interactions and provenance). Cubic and higher terms, $\mathcal{O}(\delta\hat{\boldsymbol{\Theta}}^3)$, are not discarded; the weave routes them into (i) typed $\Phi\Theta$ couplings via the typed-insertion rules for the full action (Definition 10.113), and (ii) self-energy/effective-action corrections that renormalize $\mathcal{M}_\Theta$ and move the poles of $\hat{G}_{\Theta\Theta}$. Their *provenance* is tracked by the triad tags $a \in \{\text{spec}, \text{vac}, \text{corr}\}$ (Definition 10.2), and their operational status enters the calibrated visibility meters (Definition 10.82). Thus $S^{(2)}$ underwrites linear, Σ-causal propagation on the realized weave configuration, while nonlinearity is audited and reported through typed cross-blocks and triad-resolved ledgers—without compromising TT safety or admissibility established earlier.

Interpretation of the dynamical equation (ledger view)

Chronogeometric wave operator. The principal symbol $-\bar{\nabla}_\mu(\bar{\Sigma}^{\mu\nu}\bar{\nabla}_\nu)$ is the d'Alembertian built from the total chronogeometry. Its $\bar{g}$–piece reproduces standard metric propagation, while its $\bar{t}$–piece encodes anisotropic phase flow/coherence. Thus kymon propagation is modulated by both spatial distinguishability ($\bar{g}$) and temporal coherence structure ($\bar{t}$), with direction–dependent phase drag consistent with cone–gating.

Operator mass matrix and Goldstone directions. $\mathcal{M}_\Theta(x)$ controls local inertial data: its eigenvalues are squared masses of propagating modes. Flat directions of $\mathcal{V}_\Theta$ generate zero modes of $\mathcal{M}_\Theta$ (Goldstone excitations) that propagate masslessly along the vacuum manifold, mirroring the SSB logic used in the geometric sector (cf. Section 10.17.9).

Synthesis and TT selection. Equations (10.159) and (10.160), together with the projection formula (10.154), realise an operator–valued Klein–Gordon theory on the emergent background. Coupling to chronogeometry proceeds through the typed/triad blocks while preserving TT isolation of the GW sector:

$$\boxed{\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{t}_{\mu\nu} = 0, \qquad \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{\Sigma}_{\mu\nu} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}_{\mu\nu} \quad (\text{cf. (10.155)}).} \tag{10.163}$$

Accordingly, matter dynamics backreact through non–TT channels recorded by $\mathrm{TLO}^{[\chi]}$ for admitted χ (Propositions 10.22 and 10.65), while the graviton TT block remains pure (Lemmas 10.17 and 10.59).

10.18.7 Triadic Decomposition of the Matter Propagator

The connected Tantrakyma kernel $\hat{G}_{\Theta\Theta}(x,x')$ (or, in the modal basis, $\hat{G}^{ab,cd}_{\Theta\Theta}(x,x')$; cf. Definition 10.208) inherits the *triad* organisation from the weave: $\{\mathrm{spec},\mathrm{vac},\mathrm{corr}\}$ with idempotents $\Pi^{(a)}$ (Definition 10.2). We record this both at the operator level (typed blocks on the realized weave configuration) and in the position/modal representation.

Definition 10.211 (Triadic matter block on the realized weave configuration)**.** The Tantrakyma (matter) block of the Unified Inter–Modal Correlator decomposes as

$$\boxed{\hat{\mathbf{G}}^{[\Theta\to\Theta]}_{\Lambda} = \bigoplus_{a\in\{\mathrm{spec},\mathrm{vac},\mathrm{corr}\}} \hat{\mathbf{G}}^{[\Theta\to\Theta]}_{\Lambda;a}, \qquad \hat{\mathbf{G}}^{[\Theta\to\Theta]}_{\Lambda;a} := \Pi^{(a)}\,\hat{\mathbf{G}}^{[\Theta\to\Theta]}_{\Lambda}\,\Pi^{(a)}.} \tag{10.164}$$

Here each $\hat{\mathbf{G}}^{[\Theta\to\Theta]}_{\Lambda;a}$ is a connected, time–ordered block with triad provenance a (admission and weave instrumentation as usual).

Definition 10.212 (Triadic decomposition in the position/modal basis)**.** Let $\hat{G}^{ab,cd}_{\Theta\Theta}(x,x')$ be the modal propagator matrix of Definition 10.208. Its triadic components are

$$\boxed{\hat{G}^{ab,cd}_{\Theta\Theta;a}(x,x') \;:=\; \big(\Pi^{(a)}\,\hat{G}_{\Theta\Theta}(x,x')\,\Pi^{(a)}\big)^{ab,cd}, \qquad \hat{G}^{ab,cd}_{\Theta\Theta}(x,x') = \sum_{a\in\{\mathrm{spec},\mathrm{vac},\mathrm{corr}\}} \hat{G}^{ab,cd}_{\Theta\Theta;a}(x,x').} \tag{10.165}$$

Each $\hat{G}_{\Theta\Theta;a}$ satisfies the matter Green equation projected to the a–sector:

$$\boxed{\mathcal{K}_{\Theta}(x)\,\hat{G}_{\Theta\Theta;a}(x,x') \;=\; \Pi^{(a)}\,\boldsymbol{\delta}_{\mathrm{mod}}\,\Pi^{(a)}\;\delta^{(d)}(x-x'),} \tag{10.166}$$

with the normally–hyperbolic operator $\mathcal{K}_{\Theta}$ from eq. (10.158), boundary data (retarded/advanced/Feynman) as appropriate, and $\boldsymbol{\delta}_{\mathrm{mod}}$ the identity on the modal fibre. For causal/Green's function conventions on curved backgrounds and spectral support, see [244, 7].

Operator–theoretic semantics. Equations (10.165)–(10.166) define *sectorwise* Green bi–distributions on a curved background with the usual microcausal support (ret/adv) and Feynman prescription; the triad idempotents commute with the time–ordering so that connectedness and provenance resolution are compatible [244, 7]. In this sense the triadic split is not merely a bookkeeping device but a refinement of the operator domain dictated by the weave instrumentation (admission/cone–gating) of Ch. 11.

Channel semantics (matter triad). In parallel with the weave triad (Definition 10.2):

(i) Spectral $\hat{G}_{\Theta\Theta;\mathrm{spec}}$ — propagation controlled by the energy–momentum content of kymon excitations; it governs frequency–dependent response and classical limits (e.g. redshift/dispersion) and feeds mixed geometry–matter blocks $\hat{\mathbf{G}}^{[\Phi\to\Theta]}_{\Lambda;a}$ while preserving TT isolation on the geometry leg [244, 7] (cf. eq. (10.155)).

(ii) Vacuum $\hat{G}_{\Theta\Theta;\mathrm{vac}}$ — motion along (near) flat directions of the Tantrakyma vacuum manifold; Goldstone branches appear as zero modes of the mass operator $\mathcal{M}_\Theta$ and follow the standard SSB/EFT logic adapted to emergent/curved backgrounds [415] (eq. (10.157) and Section 10.18.6).

(iii) Correlational $\hat{G}_{\Theta\Theta;\mathrm{corr}}$ — propagation driven by inter–modal entanglement/coherence on $\mathrm{End}(\mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}})$; at the interface with geometry, this channel underwrites "information $\to$ curvature'' mechanisms articulated in holographic settings [516, 1].

Projection to physical species. For the emergent kymon p with local rotation $u^{ab}_p(x)$, the species propagator Δ_p of eq. (10.154) splits compatibly with the triad:

$$\boxed{\Delta_p(x,x') = \sum_a \Delta^{(a)}_p(x,x'), \qquad \Delta^{(a)}_p(x,x') := \sum_{a,b,c,d} u^{ab}_p(x)\, \hat{G}^{ab,cd}_{\Theta\Theta;a}(x,x')\, \overline{u^{cd}_p(x')}.} \tag{10.167}$$

In stationary/homogeneous regimes each $\Delta^{(a)}_p$ admits a Källén–Lehmann–type (curved–space) representation: pole locations yield masses/dispersion while residues encode mixing and flavour amplitudes [244, 7].

Remark 10.213 (Operator origin of phenomenology and backreaction)**.** The full kernel $\hat{G}_{\Theta\Theta}$ controls gauge–invariant masses, mixings, and flavour amplitudes via its pole and branch–cut structure; the triad attributes these features to spectral, vacuum, or correlational provenance [244, 7, 415]. Coupling to chronogeometry occurs through the typed off–diagonal blocks $\hat{\mathbf{G}}^{[\Phi\to\Theta]}_{\Lambda;a}$ and through $\mathcal{K}_\Theta$'s dependence on $\bar{\Sigma}$ (Section 10.18.6), while TT purity of the graviton sector remains protected: $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{t}_{\mu\nu} = 0$ and $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{\Sigma}_{\mu\nu} = \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{g}_{\mu\nu}$ (eq. (10.155)). Operationally, this realises Wheeler's dictum at the correlator level—matter content and coherence fix geometric response without introducing extra GW polarizations [239].

Proposition: Absence of Spin–2 Kymons

Proposition 10.214 (No spin–2 excitations in the Tantrakyma sector)**.** *Let the Tantrakyma field be split as* $\hat{\mathbf{\Theta}} = \bar{\Theta}\,\mathbb{I} + \delta\hat{\mathbf{\Theta}}$ *(cf. Definition 10.204 and Section 10.18.6). Through the constitutive scalar/vector sourcing maps* $S_{\mu\nu}[\cdot], A_{\mu\nu}[\cdot]$ *of Definitions 10.4 and 10.13 the temporal/coherence fluctuation reads* $\hat{\tau}_{\mu\nu} = S_{\mu\nu}[\hat{\varphi}_A] \oplus A_{\mu\nu}[\hat{K}^{(\tau)}]$ *with* $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\hat{\tau}_{\mu\nu} = 0$ *(eqs. (10.102) and (10.155)). Then the Tantrakyma sector carries* no *spin–2 irreducible excitations:*

$$\boxed{\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\hat{\tau}_{\mu\nu} = 0 \qquad \Longrightarrow \qquad \textit{no spin–2 kymon.}} \tag{10.168}$$

Proof. By Definitions 10.4 and 10.13, $\hat{\tau}_{\mu\nu}$ is built from a trace (scalar) part $S_{\mu\nu}[\hat{\varphi}_A]$ and a longitudinal (vector) part $A_{\mu\nu}[\hat{K}^{(\tau)}]$. On $(\bar{g}_{\mu\nu}, \bar{\nabla})$ the York/Helmholtz decomposition $\mathrm{Sym}^2 = \mathrm{TT} \oplus \mathrm{Im}\,\mathcal{L} \oplus \mathrm{span}\{\bar{g}\}$ (eq. (10.140)) is L^2–orthogonal; the TT projector annihilates both trace and longitudinal subspaces. Hence $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\hat{\tau}_{\mu\nu} = 0$ (eq. (10.102) and Lemma 10.59), i.e. no spin–2 content arises from Tantrakyma. By eq. (10.28) and Corollary 10.20 the radiative spin–2 sector is purely metric (the graviton). □

Remark 10.215 (Scope and composites)**.** The statement concerns *elementary* Tantrakyma excitations (kymons). Composite multi–kymon states can carry total spin 2, but do not constitute an elementary spin–2 field; gravitational radiation remains confined to the metric TT channel (cf. eq. (10.28) and Corollary 10.20).

The Matter & Coherence Sector: Tantrakyma as the Source of Substance

The Tantrakyma field $\hat{\boldsymbol{\Theta}}(x)$ sources all matter/coherence excitations on the realized weave configuration. Its potential $\mathcal{V}_\Theta$ and quadratic kernel determine the spectrum of its quanta (kymons) via the background–fluctuation split and the linearised dynamics (Section 10.18.6). Two–point functions in the modal and projected (species) bases are encoded by $\boldsymbol{\Delta}_\Theta$ and Δ_p, respectively (Definition 10.208 and eq. (10.154)).

Goldstone/Higgs logic (matter sector). Spontaneous breaking of a continuous symmetry of $\mathcal{V}_\Theta$ yields massless Goldstone kymons for global SSB, or removes them via gauge fixing in local SSB (unitary–gauge Higgs mechanism). This is implemented on the emergent background through $\mathcal{M}_\Theta(x)$ and the normally–hyperbolic operator $\mathcal{K}_\Theta$ (eqs. (10.157) and (10.158) and Section 10.18.6).

Mass eigenmodes. After SSB/Higgsing, Tantrakyma excitations become massive eigenmodes of $\mathcal{M}_\Theta$; their masses are the pole locations of Δ_p:

(i) Radial Higgs kymon (spin–0) The radial scalar obtains its mass from the Hessian of $\mathcal{V}_\Theta$ at the vacuum (eq. (10.157)).

(ii) Chrono–Kymon (spin–1) The vector excitation sources the temporal/coherence channel through $A_{\mu\nu}[\hat{K}^{(\tau)}]$, contributing to the antisymmetric/phase–transport sector of $t_{\mu\nu}$ in the constitutive split (Definitions 10.4 and 10.13 and Section 10.18.6).

No spin–2 kymons. The Tantrakyma sector contains no irreducible spin–2 excitations:

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\hat{\tau}_{\mu\nu} = 0 \quad (\textit{eqs. } (10.155) \textit{and } (10.168)).$$

The radiative spin–2 (graviton) sector is purely metric,

$$\hat{h}_{\mu\nu} \;=\; \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}_{\mu\nu} \quad (\textit{eq. } (10.28) \textit{and Corollary } 10.20).$$

Triad and state dependence. Propagation and mixing decompose into correlational, vacuum, and spectral channels (Section 10.18.7). All matter data are state–dependent through $\bar{\Sigma}_{\mu\nu}$ and $\mathcal{M}_\Theta(x)$ (Section 10.18.6). Backreaction enters via the typed off–diagonal blocks $\hat{\mathbf{G}}^{[\Phi\to\Theta]}_{\Lambda;a}$ and through the $\bar{\Sigma}$–dependence of $\mathcal{K}_\Theta$, while TT purity of the graviton sector is preserved (eqs. (10.28) and (10.155) and Corollary 10.20).

10.19 Matter–Geometry Block of the Unified Correlator

We now treat the off–diagonal sector of the Unified Inter–Modal Correlator $\hat{\mathbf{G}}_\Lambda$ (cf. Definition 6.4), namely the *matter–geometry* block that links the Dynamical Gaiamiti and Tantrakyma fields at the operator level. Throughout we use the typed projector family and triad idempotents of Ch. 11 (cf. Definitions 10.15 and 10.25) and the QHS expectation $\langle\cdot\rangle_\circ$ (cf. Definitions 10.203 and 10.204).

10.19.1 Typed Off–diagonal Block and Connected Kernel

Definition 10.216 (Typed matter→geometry cross block)**.** The *typed* matter→geometry block of the unified correlator is

$$\boxed{\hat{\mathbf{G}}^{[\Phi\to\Theta]}_\Lambda := \Pi^{[\Theta]}_\Lambda\,\hat{\mathbf{G}}_\Lambda\,\Pi^{[\Phi]}_\Lambda, \qquad \hat{\mathbf{G}}_\Lambda \text{ as in Definition 6.4.}} \tag{10.169}$$

With the triad $\{a\} = \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$ (cf. Definition 10.15), its provenance–resolved form is

$$\boxed{\hat{\mathbf{G}}^{[\Phi\to\Theta]}_\Lambda = \bigoplus_{a\in\{\mathrm{spec,vac,corr}\}} \hat{\mathbf{G}}^{[\Phi\to\Theta]}_{\Lambda;a}, \qquad \hat{\mathbf{G}}^{[\Phi\to\Theta]}_{\Lambda;a} := \Pi^{[a]}\,\hat{\mathbf{G}}^{[\Phi\to\Theta]}_\Lambda\,\Pi^{[a]}.} \tag{10.170}$$

Each $\hat{\mathbf{G}}^{[\Phi\to\Theta]}_{\Lambda;a}$ is an admission–certified cross block suitable for witnesses/meters and T–L–O execution (cf. Definition 10.1 and Proposition 10.90).

Definition 10.217 (Connected matter–geometry kernel)**.** Let $\hat{\mathbf{\Phi}}(x)$ and $\hat{\mathbf{\Theta}}(x')$ be the dynamical Gaiamiti/Tantrakyma fields (cf. Definitions 6.118 and 6.142). The *connected, time–ordered* cross kernel is

$$\boxed{\hat{G}_{\Phi\Theta}(x,x') := \Big\langle T\{\hat{\mathbf{\Phi}}(x)\,\hat{\mathbf{\Theta}}(x')\}\Big\rangle_\circ - \Big\langle\hat{\mathbf{\Phi}}(x)\Big\rangle_\circ\Big\langle\hat{\mathbf{\Theta}}(x')\Big\rangle_\circ.} \tag{10.171}$$

Using the background–fluctuation splits $\hat{\mathbf{\Phi}} = \bar{\Phi}\,\mathbb{I} + \delta\hat{\mathbf{\Phi}}$ and $\hat{\mathbf{\Theta}} = \bar{\Theta}\,\mathbb{I} + \delta\hat{\mathbf{\Theta}}$ (cf. eq. (10.87) and Definition 10.204),

$$\hat{G}_{\Phi\Theta}(x,x') \;=\; \Big\langle T\{\delta\hat{\mathbf{\Phi}}(x)\,\delta\hat{\mathbf{\Theta}}(x')\}\Big\rangle_\circ. \tag{10.172}$$

Operator–theoretic semantics. The cross kernel (10.171) is a curved–background, operator–valued two–point function, with time–ordering and subtraction performed in the QHS; its construction and interpretation follow the standard QFT on curved spacetime framework [244, 7]. In the typed formulation (10.169)–(10.170), each provenance block $\hat{\mathbf{G}}^{[\Phi\to\Theta]}_{\Lambda;a}$ is an idempotent–resolved channel that respects admission and cone–gating constraints, ensuring compatibility with T–L–O execution and with the constitutive/observable split of Ch. 11.

Causal/triadic interpretation. A non–trivial value of $\hat{G}_{\Phi\Theta}(x,x')$ signals that a fluctuation of the Tantrakyma field at x' is *correlated* with a geometric fluctuation of the Gaiamiti field at x, thereby realizing a state–relative notion of "matter sources geometry" at the operator level [239]. Triad resolution sharpens this reading: the correlational channel ($a = \mathrm{corr}$) tracks entanglement/coherence converting into geometric response, echoing ER=EPR and "spacetime from entanglement" intuitions [516, 1]; the vacuum channel ($a = \mathrm{vac}$) slides along (near–)flat directions of the vacuum bundle and aligns with Goldstone deformations in an EFT sense [415]; and the spectral channel ($a = \mathrm{spec}$) organizes frequency–dependent sourcing and classical limits (redshift/dispersion) in the nonlocal operator framework [244, 7].

State dependence and nonlocality. Because expectations are taken in the QHS $\langle\cdot\rangle_\circ$, the magnitude and tensor structure of $\hat{G}_{\Phi\Theta}$ are determined by the global quantum state, not by kinematics alone; this encodes nonlocal, bi–local influence mediated by $\mathcal{U}_{\Phi\Theta}$ and captured cleanly by the fluctuation form (10.172) [244, 7]. In particular, the correlational channel provides an explicit pathway by which quantum information content in the matter sector imprints correlational curvature in geometry [516, 1].

Operational content and TT integrity. The typed cross block integrates with the TT selection on the geometry leg: radiative tensor modes are governed by the projector $\Pi^{[\gamma]} = \mathcal{P}^{(2)}$ and remain TT–pure (cf. Definition 10.167 and Lemma 10.59), so that off–diagonal matter–geometry correlations contribute without introducing extra GW polarizations. This realizes Wheeler's dictum [239]operationally while remaining consistent with the TT closure established in earlier sections.

10.19.2 Modal Representation and Transport

Definition 10.218 (Modal expansion with inter–modal transport)**.** Choose a local orthonormal basis $\{|\mathbf{i}\rangle_x\} \subset T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}$ for the Gaiamiti fibre and a matrix basis $\{|\mathbf{a}\rangle_{x'}\langle\mathbf{b}|_{x'}\} \subset \mathrm{End}(\mathcal{H}^{(\mathfrak{B}_{x'})}_{\mathrm{mod}})$ for Tantrakyma (cf. Definition 4.273). Write

$$\delta\hat{\mathbf{\Phi}}(x) = \sum_i \delta\hat{\Phi}_i(x)\,|\mathbf{i}\rangle_x, \qquad \delta\hat{\mathbf{\Theta}}(x') = \sum_{a,b} \delta\theta^{ab}(x')\,|\mathbf{a}\rangle_{x'}\langle\mathbf{b}|_{x'}.$$

To compare fibres, parallel–transport Tantrakyma indices with the inter–modal transporter $\mathcal{U}_{x'\to x}$ (cf. Definition 4.9): $\delta\tilde{\theta}^{ab}(x'\to x) := \mathcal{U}_{x'\to x}\,\delta\theta^{ab}(x')\,\mathcal{U}_{x\to x'}$. Then

$$\boxed{\hat{G}_{\Phi\Theta}(x,x') = \sum_{i,a,b} \mathrm{Cov}_\circ\Big[\,\delta\hat{\Phi}_i(x),\ \delta\tilde{\theta}^{ab}(x'\to x)\,\Big]\ |\mathbf{i}\rangle_x \otimes \Big(|\mathbf{a}\rangle_x\langle\mathbf{b}|_x\Big),} \tag{10.173}$$

with $\mathrm{Cov}_\circ[A,B] := \langle T\{AB\}\rangle_\circ - \langle A\rangle_\circ\langle B\rangle_\circ$.

Typed pair refinement (optional). When useful, refine (10.173) by inserting a TT type on the geometry leg and any admitted τ–kairos type $\chi \in \{S^{(b)}, S^{(L)}, \tilde{S}^{(p)}, A^{(+)}, A^{(-)}\}$ on the matter leg (cf. Definitions 10.3 and 10.4):

$$\boxed{\hat{\mathbf{G}}^{[\,\Phi\to\Theta;\ \mathrm{TT}\to\chi\,]}_{\Lambda;a} \;:=\; \Pi^{(a)}\,\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\hat{\mathbf{G}}^{[\,\Phi\to\Theta\,]}_{\Lambda}\,\Pi^{[\chi]}\,\Pi^{(a)}, \qquad a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}.} \tag{10.174}$$

Here $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ is the background–covariant TT projector and the graviton extractor is $\mathcal{P}^{(2)}_{\mathrm{grav}}$ (Definitions 10.3 and 10.25). TT–nullity of τ–kairos implies $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\Pi^{[\chi]} = 0$, hence also $\mathcal{P}^{(2)}_{\mathrm{grav}}\Pi^{[\chi]} = 0$ (Lemmas 10.17 and 10.59); therefore no additional GW polarizations arise from the matter leg, and the TT graviton sector remains uncontaminated under typed mixing (Lemma 10.123).

10.19.3 Triadic Resolution and Parasymmetric TLO Interface

Remark 10.219 (Triadic resolution)**.** The components $\hat{\mathbf{G}}_{\Lambda;a}^{[\Phi\to\Theta]}$ in (10.170) separate correlational, vacuum, and spectral provenance (cf. Sections 10.17.6 and 10.18.7). Each inherits the causal support and admission of its channel and solves the appropriate mixed Green's equation on its block (cf. eqs. (10.112) and (10.160)).

Remark 10.220 (Parasymmetric pipeline and purity guards)**.** Running the sector execution functor $\mathscr{E}_\bullet$ with the same typed projectors and admission gate (cf. Definitions 9.153 and 10.94), the mixed block contributes witness signals and Onset timers to the appropriate *non–TT* temporal types, while TT content remains confined to the geometric leg (cf. Definition 10.167 and Lemma 10.168). The TT–purity witness $\mathcal{W}_{\mathrm{TT\text{–}purity}}$ monitors law–side consistency (cf. Lemma 10.17).

Operational reading. Equation (10.173) makes explicit how a given geometric fluctuation mode (index i) correlates with a specific modal transition (a, b) in the matter/coherence fibre, after covariance transport. These are the primitive operator–level *interaction channels* that feed the off–diagonal entries of the chronogeometric Green's system, while preserving TT selection for gravitational radiation (cf. eqs. (10.28), (10.90) and (10.131) and Corollary 10.20).

Bi–Local Interaction Lagrangian

We encode the matter–geometry coupling in the off–diagonal (typed) blocks of the unified correlator (cf. Definitions 6.4, 10.216 and 10.217). All expectations are taken in the QHS (Definitions 10.203 and 10.204); background–fluctuation splits are those of eq. (10.87) and Definition 10.204. Fibre comparisons use the inter–modal transporter $\mathcal{U}_{x'\to x}$ (Definition 4.9).

Definition 10.221 (Typed, Hermitian interaction kernel (bi–local))**.** Let the typed cross blocks be $\hat{\mathbf{G}}_{\Lambda}^{[\Phi\to\Theta]} = \Pi_{\Lambda}^{[\Theta]}\,\hat{\mathbf{G}}_{\Lambda}\,\Pi_{\Lambda}^{[\Phi]}$ and $\hat{\mathbf{G}}_{\Lambda}^{[\Theta\to\Phi]} = \Pi_{\Lambda}^{[\Phi]}\,\hat{\mathbf{G}}_{\Lambda}\,\Pi_{\Lambda}^{[\Theta]}$ (cf. Definition 10.216). The *bi–local, Hermitian* interaction kernel is

$$\boxed{\mathbf{U}_{\Lambda}^{[\Phi\leftrightarrow\Theta]}(x,x') \;:=\; \hat{\mathbf{G}}_{\Lambda}^{[\Phi\to\Theta]}(x,x') \;+\; \Big(\hat{\mathbf{G}}_{\Lambda}^{[\Theta\to\Phi]}(x',x)\Big)^{\dagger},} \tag{10.175}$$

and inherits the triadic resolution

$$\mathbf{U}_{\Lambda}^{[\Phi\leftrightarrow\Theta]} = \bigoplus_{a\in\{\mathrm{spec,vac,corr}\}} \mathbf{U}_{\Lambda;a}^{[\Phi\leftrightarrow\Theta]}, \qquad \mathbf{U}_{\Lambda;a}^{[\Phi\leftrightarrow\Theta]} := \Pi^{[a]}\,\mathbf{U}_{\Lambda}^{[\Phi\leftrightarrow\Theta]}\,\Pi^{[a]}, \tag{10.176}$$

(cf. Definition 10.15).

Definition 10.222 (Bi–local interaction Lagrangian density)**.** Write the fluctuation fields as $\delta\hat{\mathbf{\Phi}}(x)$ and $\delta\hat{\mathbf{\Theta}}(x')$, and transport Tantrakyma to x: $\delta\tilde{\hat{\mathbf{\Theta}}}(x'\to x) := \mathcal{U}_{x'\to x}\,\delta\hat{\mathbf{\Theta}}(x')\,\mathcal{U}_{x\to x'}$. The *bi–local interaction density* is the modal trace at x:

$$\boxed{\mathcal{L}_{\mathrm{int}}(x,x') \;:=\; \mathrm{Tr}\Big[\big(\delta\hat{\mathbf{\Phi}}(x)\big)^{\dagger}\;\mathbf{U}_{\Lambda}^{[\Phi\leftrightarrow\Theta]}(x,x')\;\delta\tilde{\hat{\mathbf{\Theta}}}(x'\to x)\Big],} \tag{10.177}$$

with Tr taken on the transported modal fibre at x.

Definition 10.223 (Interaction action (bi–local form; triad–resolved))**.** The total interaction action is

$$\boxed{S_{\mathrm{int}}[\hat{\mathbf{\Phi}},\hat{\mathbf{\Theta}}] := \int d^dx\,d^dx'\;\mathcal{L}_{\mathrm{int}}(x,x') \;=\; \sum_a \int d^dx\,d^dx'\;\mathrm{Tr}\Big[(\delta\hat{\mathbf{\Phi}})^{\dagger}\;\mathbf{U}_{\Lambda;a}^{[\Phi\leftrightarrow\Theta]}\;\delta\tilde{\hat{\mathbf{\Theta}}}\Big],} \tag{10.178}$$

where $a\in\{\mathrm{spec},\mathrm{vac},\mathrm{corr}\}$ and all fields are evaluated as in (10.177).

Properties and guards. By construction, $\mathbf{U}_{\Lambda}^{[\Phi\leftrightarrow\Theta]}(x,x')^{\dagger} = \mathbf{U}_{\Lambda}^{[\Phi\leftrightarrow\Theta]}(x',x)$ (Hermiticity); causal support and admission are inherited from the cross blocks and the cone gates (Definition 10.94 and eq. (10.65)). TT–purity is preserved because the right leg is tau–kairos–typed and TT–null (eq. (10.90) and Lemma 10.59); the TT sector remains purely geometric (Definition 10.167 and Lemma 10.168).

10.19.4 Functional Variation of the Interaction Action

Let $S_{\mathrm{total}} = S_{\Phi} + S_{\Theta} + S_{\mathrm{int}}$ be the total action (cf. constitutive and matter dynamics in Sections 10.17.4 and 10.18.6). Varying (10.178) at fixed transporter yields bi–local source terms.

Proposition 10.224 (Bi–local sources in the Euler–Lagrange equations)**.** *The first variations of S_{int} with respect to the fluctuation fields are*

$$\frac{\delta S_{\mathrm{int}}}{\delta\big(\delta\hat{\mathbf{\Phi}}(x)\big)^{\dagger}} = \int d^dx'\;\mathbf{U}_{\Lambda}^{[\Phi\leftrightarrow\Theta]}(x,x')\;\delta\tilde{\hat{\mathbf{\Theta}}}(x'\to x), \tag{10.179}$$

$$\frac{\delta S_{\mathrm{int}}}{\delta\big(\delta\hat{\mathbf{\Theta}}}(x')\big)^{\dagger} = \int d^dx\;\Big(\mathcal{U}_{x\to x'}\,\mathbf{U}_{\Lambda}^{[\Phi\leftrightarrow\Theta]}(x,x')^{\dagger}\,\mathcal{U}_{x'\to x}\Big)\;\delta\hat{\mathbf{\Phi}}(x), \tag{10.180}$$

with identical triad–resolved statements after inserting $\Pi^{[a]}$ as in (10.176).

Remark 10.225 (Closure of the TT sector under variation)**.** Applying the TT projector to the geometric equation sourced by (10.179) yields $\mathcal{P}^{(2)}\big[\delta S_{\mathrm{int}}/\delta(\delta\hat{\mathbf{\Phi}})^{\dagger}\big] = \mathcal{P}^{(2)}\big[\cdots\big]$ with the right leg tau–typed and TT–null; thus no extra GW polarizations are induced (cf. Definition 10.167 and Lemmas 10.59 and 10.168). The resulting mixed terms enter the off–diagonal blocks of the chronogeometric Green's system compatibly with eqs. (10.112) and (10.160).

Interpretation (operator/triad/T–L–O). Equations (10.177)–(10.180) implement the IQR tenet that interaction is entanglement–mediated and fundamentally bi–local. The triadic split (10.176) attributes the coupling to spec, vac, and corr provenance, while the sector execution functor $\mathscr{E}_\bullet$ records non–TT temporal responses via $\mathrm{TLO}^{[\chi]}$ with $\chi \in \{\tau_S, \tau_A, K_{\mathrm{T}}^{(\tau)}, K_{\mathrm{L}}^{(\tau)}\}$ (cf. Definition 9.153 and Proposition 10.22) and leaves $\mathrm{TLO}^{[\mathrm{TT}]}$ untouched (cf. Proposition 10.35).

Functional Variation and Coupled Source Currents

We vary the bi–local interaction action of Definition 10.223 with respect to the *conjugate fluctuation fields*, keeping the inter–modal transporter fixed (cf. Definition 4.9). All fields are background–split as in eq. (10.87) and Definition 10.204.

Definition 10.226 (Bi–local source currents (transported, typed/triad–ready))**.** Let $\delta\tilde{\hat{\mathbf{\Theta}}}(x' \to x) := \mathcal{U}_{x'\to x}\, \delta\hat{\mathbf{\Theta}}(x')\, \mathcal{U}_{x\to x'}$. The functional variations of S_{int} (Definition 10.223) define the operator–valued currents

$$\hat{\mathcal{J}}^{\Phi}_{(\Theta)}(x) := \frac{\delta S_{\mathrm{int}}}{\delta\Big(\delta\hat{\mathbf{\Phi}}(x)\Big)^{\dagger}} = \int d^d x'\; \mathbf{U}_{\Lambda}^{[\Phi\leftrightarrow\Theta]}(x,x')\; \delta\tilde{\hat{\mathbf{\Theta}}}(x'\to x), \tag{10.181}$$

$$\hat{\mathcal{J}}^{\Theta}_{(\Phi)}(x') := \frac{\delta S_{\mathrm{int}}}{\delta\Big(\delta\hat{\mathbf{\Theta}}}(x')\Big)^{\dagger} = \int d^d x\; \Big(\mathcal{U}_{x\to x'}\, \mathbf{U}_{\Lambda}^{[\Phi\leftrightarrow\Theta]}(x,x')^{\dagger}\, \mathcal{U}_{x'\to x}\Big)\; \delta\hat{\mathbf{\Phi}}(x). \tag{10.182}$$

They inherit the triadic resolution (eq. (10.176)): $\hat{\mathcal{J}}^{\bullet}_{(\circ)} = \bigoplus_a \hat{\mathcal{J}}^{\bullet}_{(\circ);a}$ with $\hat{\mathcal{J}}^{\bullet}_{(\circ);a} := \Pi^{[a]}\, \hat{\mathcal{J}}^{\bullet}_{(\circ)}\, \Pi^{[a]}$ for $a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$.

Terminology (aligned with Ch. 10). $\hat{\mathcal{J}}^{\Phi}_{(\Theta)}$ is the *matter–sourced geometric current* (a source on the Gaiamiti leg); $\hat{\mathcal{J}}^{\Theta}_{(\Phi)}$ is the *geometry–sourced matter current* (a source on the Tantrakyma leg). Typed/helicity refinements follow the projector family of Definition 10.25.

Quantum–Covariant Coupled Field Equations (operator form)

Let $\mathcal{K}_\Theta$ be the normally hyperbolic matter operator of eq. (10.158) with linearized equation eq. (10.159). Let $\mathcal{K}_\Phi$ denote the geometric kinetic operator obtained from the quadratic part of S_Φ (chronogeometric wave operator in the sense of Definition 10.167 and eq. (10.131); constitutive background in Section 10.17.4). Then the coupled, bi–locally sourced operator equations read

$$\mathcal{K}_\Phi(x)\, \delta\hat{\mathbf{\Phi}}(x) = \hat{\mathcal{J}}^{\Phi}_{(\Theta)}(x), \tag{10.183}$$

$$\mathcal{K}_\Theta(x')\, \delta\hat{\mathbf{\Theta}}(x') = \hat{\mathcal{J}}^{\Theta}_{(\Phi)}(x'), \tag{10.184}$$

where the source operators $\hat{\mathcal{J}}^{\Phi}_{(\Theta)}$ and $\hat{\mathcal{J}}^{\Theta}_{(\Phi)}$ are constructed from the bi–local interaction kernel $\mathcal{U}_{\Phi\Theta}(x,x')$ (cf. Definition 10.229 and eq. (10.176)) within the standard operator framework of quantum fields on curved backgrounds [244, 7].

Corollary 10.227 (TT closure of the graviton sector)**.** *Applying the TT projector gives* $\mathcal{P}^{(2)}\big[\mathcal{K}_\Phi\,\delta\hat{\mathbf{\Phi}}\big] = \mathcal{P}^{(2)}\big[\hat{\mathcal{J}}^{\Phi}_{(\Theta)}\big] = 0$*, because the right leg is tau–typed and TT–null (eq. (10.90) and Lemma 10.59); hence the radiative TT mode* $\hat{\gamma}_{\mu\nu} = \mathcal{P}^{(2)}[\delta\hat{g}_{\mu\nu}]$ *evolves without additional GW polarizations (cf. Definition 10.167 and Lemma 10.168).*

Current–driven form (component notation). When $\mathcal{K}_\Phi \to -\bar{\nabla}_\mu(\bar{\Sigma}^{\mu\nu}\bar{\nabla}_\nu\,\cdot\,) + \mathcal{M}_\Phi$ and $\mathcal{K}_\Theta \to -\bar{\nabla}_\mu(\bar{\Sigma}^{\mu\nu}\bar{\nabla}_\nu\,\cdot\,) + \mathcal{M}_\Theta$ (chronogeometric compatibility; cf. Section 10.18.6),

$$-\bar{\nabla}_\mu\Big(\bar{\Sigma}^{\mu\nu}\bar{\nabla}_\nu\delta\hat{\mathbf{\Phi}}\Big) + \mathcal{M}_\Phi\,\delta\hat{\mathbf{\Phi}} = \hat{\mathcal{J}}^{\Phi}_{(\Theta)}, \qquad \text{(geometry)}, \tag{10.185}$$

$$-\bar{\nabla}_\mu\Big(\bar{\Sigma}^{\mu\nu}\bar{\nabla}_\nu\delta\hat{\mathbf{\Theta}}\Big) + \mathcal{M}_\Theta\,\delta\hat{\mathbf{\Theta}} = \hat{\mathcal{J}}^{\Theta}_{(\Phi)}, \qquad \text{(matter)}, \tag{10.186}$$

in alignment with the Ch. 10 current conventions while preserving the Ch. 11 typed/bi–local structure [244, 7].

Physical Interpretation. These equations instantiate the quantum analogue of Wheeler's dictum [239].

(i) "Matter tells geometry how to curve" Equation (10.183) shows that the Gaiamiti field is driven by the state of the Tantrakyma field through the operator–valued source $\hat{\mathcal{J}}^{\Phi}_{(\Theta)}$, which is built from the cross–correlator and interaction kernel $\mathcal{U}_{\Phi\Theta}$ (cf. Definition 10.229 and eq. (10.176)). In particular, the correlational channel of $\mathcal{U}_{\Phi\Theta}$ realizes the entanglement→geometry mechanism emphasized in holographic contexts [516, 1].

(ii) "Geometry tells matter how to move" Equation (10.184) shows that the Tantrakyma field propagates in (and responds to) the chronogeometric background generated by $\hat{\mathbf{\Phi}}$, with $\hat{\mathcal{J}}^{\Theta}_{(\Phi)}$ encoding how curvature and geometric fluctuations modulate matter dynamics in the curved-space operator formalism [244, 7].

Significance. The coupled system expresses a quantum–covariant feedback loop: geometry and matter are *co-constructed* by bi–local operator correlations rather than merely coupled by local densities. The interaction kernel $\mathcal{U}_{\Phi\Theta}(x, x')$ is the central mediator: it simultaneously sources, propagates, and encodes mutual influence between the two sectors and, in its correlational channel, provides a concrete route by which quantum information structure imprints geometry [516, 1]. From this perspective, the classical role of the local stress–energy tensor as the generator of curvature is supplanted by the off-diagonal operator correlator, a nonlocal object that carries both energetic and coherent content in the curved-background QFT setting [244, 7]. The TT closure of Corollary 10.227 then guarantees that no extra GW polarizations are introduced: correlational and vacuum responses feed back into geometry without violating the transverse–traceless selection on the metric radiative sector.

10.19.5 Mutually–Sourced Field Currents and Typed Reciprocity

Remark 10.228 (Typed, triad–resolved reading)**.** With the type projectors $\Pi^{[\chi]}$ (Definition 10.25) and triad idempotents $\Pi^{[a]}$ (Definition 10.15), one may record $\hat{\mathcal{J}}^{\Phi}_{(\Theta);a} = \Pi^{[a]}\hat{\mathcal{J}}^{\Phi}_{(\Theta)}\Pi^{[a]}$

and similarly for $\hat{\mathcal{J}}^{\Theta}_{(\Phi);a}$. TT purity implies $\Pi^{[\gamma]}\hat{\mathcal{J}}^{\Phi}_{(\Theta);a}\Pi^{[\gamma]} = 0$, while non–TT temporal types in $\{\tau_S, \tau_A, K^{(\tau)}_{\mathrm{T}}, K^{(\tau)}_{\mathrm{L}}\}$ are admissible (cf. Lemma 10.59 and Proposition 10.22).

Physical interpretation (Wheeler dictum, quantum form). Equation (10.183) is the operator incarnation of "matter tells geometry how to curve'': $\hat{\mathcal{J}}^{\Phi}_{(\Theta)}$ aggregates the nonlocal, state–dependent Tantrakyma influence on geometry through $\mathbf{U}_{\Lambda}^{[\Phi\leftrightarrow\Theta]}$ (Definition 10.221). Equation (10.184) is the reciprocal "geometry tells matter how to move'' via $\hat{\mathcal{J}}^{\Theta}_{(\Phi)}$. The feedback remains TT–safe by Corollary 10.227.

T–L–O ledger compatibility. Currents projected by $\Pi^{[\chi]}$ feed the same Type/Location/Onset stamps as their parent correlator blocks (cf. Definition 9.153 and Propositions 10.22 and 10.35); TT stamps $\mathrm{TLO}^{[\gamma]}$ remain unaffected by non–TT current components.

Summary. The bi–local currents (10.181)–(10.182) provide the Ch. 11–consistent, transporter–covariant sources for the coupled equations (10.183)–(10.184), align with the Ch. 10 current conventions at the operator level, and preserve TT purity of the graviton sector under the projector constraints of Definition 10.25 and Lemma 10.59.

10.19.6 Triadic Decomposition of the Interaction Channels

We work with the transported, off–diagonal block $\hat{\mathbf{G}}_{\Lambda}^{[\Phi\to\Theta]}(x,x')$ of the Unified Inter–Modal Correlator (cf. Definition 10.113 and Section 10.19), whose legs live on a *common* modal fibre via $\mathcal{U}_{x'\to x}$ (see Definition 4.9). Both sides admit the triad idempotents $\{\Pi^{[a]}\}_{a\in\{\mathrm{spec,vac,corr}\}}$ (Definition 10.15), acting on the *geometric* (Φ) leg (left) and the *matter* (Θ) leg (right), respectively.

Definition 10.229 (Double–triad–resolved cross–correlator)**.** The matter–geometry cross–correlator is decomposed as

$$\boxed{\hat{\mathbf{G}}_{\Lambda}^{[\Phi\to\Theta]}(x,x') = \sum_{a,b\in\{\mathrm{spec,vac,corr}\}} \hat{\mathbf{G}}_{\Lambda;a|b}^{[\Phi\to\Theta]}(x,x'), \qquad \hat{\mathbf{G}}_{\Lambda;a|b}^{[\Phi\to\Theta]} := \Pi_{\Phi}^{[a]}\,\hat{\mathbf{G}}_{\Lambda}^{[\Phi\to\Theta]}\,\Pi_{\Theta}^{[b]}.} \tag{10.187}$$

Here $\Pi_{\Phi}^{[a]}$ acts on the geometric leg and $\Pi_{\Theta}^{[b]}$ on the matter leg (after transport to x). The Hermitian reciprocity reads

$$\left(\hat{\mathbf{G}}_{\Lambda;a|b}^{[\Phi\to\Theta]}(x,x')\right)^{\dagger} = \hat{\mathbf{G}}_{\Lambda;b|a}^{[\Theta\to\Phi]}(x',x). \tag{10.188}$$

Remark 10.230 (Typed refinement commutes with the triad)**.** For any admitted geometry type $\Pi^{[\chi]}$ (Definition 10.25), the typed block $\hat{\mathbf{G}}_{\Lambda;a|b}^{[\Phi\to\Theta;\chi]} := \Pi_{\Lambda}^{[\chi]}\,\hat{\mathbf{G}}_{\Lambda;a|b}^{[\Phi\to\Theta]}\,\Pi_{\Lambda}^{[\chi]}$ is well–defined and satisfies the same reciprocity (10.188). In particular, TT purity on the geometry leg is enforced by $\Pi^{[\gamma]} = \mathcal{P}^{(2)}$.

Channel diagram. Equation (10.187) can be visualised as a 3×3 matrix $\left(\hat{\mathbf{G}}^{[\Phi\to\Theta]}_{\Lambda;a|b}\right)_{a,b}$ with rows (a) labelling the geometric triad origin and columns (b) the matter triad origin. Diagonal entries encode mode–preserving couplings; off–diagonals encode mode–mixing couplings.

Definition 10.231 (Triad–resolved interaction kernel (single–index))**.** The Hermitian interaction kernel of Definition 10.221 inherits a triad decomposition on the *geometric* leg:

$$\boxed{\mathbf{U}^{[\Phi\leftrightarrow\Theta]}_{\Lambda}(x,x') = \sum_{a\in\{\mathrm{spec,vac,corr}\}} \mathbf{U}^{[\Phi\leftrightarrow\Theta]}_{\Lambda;a}(x,x'), \qquad \mathbf{U}^{[\Phi\leftrightarrow\Theta]}_{\Lambda;a} := \sum_{b}\left(\hat{\mathbf{G}}^{[\Phi\to\Theta]}_{\Lambda;a|b} + \left(\hat{\mathbf{G}}^{[\Theta\to\Phi]}_{\Lambda;b|a}\right)^{\dagger}\right).} \tag{10.189}$$

Each $\mathbf{U}^{[\Phi\leftrightarrow\Theta]}_{\Lambda;a}$ is again triad–idempotent on the geometry leg: $\Pi^{[a]}_{\Phi}\,\mathbf{U}_{\Lambda;a} = \mathbf{U}_{\Lambda;a} = \mathbf{U}_{\Lambda;a}\,\Pi^{[a]}_{\Phi}$.

Remark 10.232 (Semantics of the triadic interaction channels)**.** In parallel with Sections 10.17.6 and 10.18.7, we interpret each channel of the single-index decomposition $\mathcal{U}^{(a)}_{\Phi\Theta}$ as follows.

(i) Correlational ($a = \mathrm{corr}$)
This channel captures entanglement- and coherence–driven curvature response: informational structure in the matter sector Θ imprints correlational geometry on the Gaiamiti field through the cross–correlator, realizing the idea that spacetime connectivity can be sourced by quantum correlations [516, 1]. Accordingly, $\mathbf{U}_{\Lambda;\mathrm{corr}}$ aggregates the contributions of $\hat{G}^{(\mathrm{corr}|b)}_{\Phi\Theta}$ that transduce relational entanglement into geometric response.

(ii) Vacuum ($a = \mathrm{vac}$)
This channel describes response along (near–)flat directions of the vacuum bundle $\mathcal{M}_{\mathrm{vac}}$ of the Gaiamiti sector, aligned with Goldstone deformations that parameterize spontaneous symmetry breaking of the matter sector on the emergent background; its dynamics follow standard effective–field–theory reasoning for Goldstone/Higgs modes adapted to curved or emergent geometries [415]. In particular, $\mathbf{U}_{\Lambda;\mathrm{vac}}$ isolates those geometric variations that slide along the vacuum manifold rather than exciting tensorial helicities.

(iii) Spectral ($a = \mathrm{spec}$)
This channel encodes energy–momentum (spectral) sourcing of curvature: frequency content in the matter sector drives frequency–dependent geometric response and controls classical limits such as gravitational redshift, dispersion, and localization within the nonlocal operator framework of quantum fields on curved backgrounds [244, 7]. Consequently, $\mathbf{U}_{\Lambda;\mathrm{spec}}$ collects the pieces of the kernel whose support is organized by spectral structure.

Off–diagonal blocks with $a \neq b$ in (10.187) represent triad *mode–mixing*: correlational, vacuum, and spectral characters can transmute across legs in the cross–correlator, and their contributions feed the interaction kernel according to (10.176) while never violating TT selection on the geometry leg because $\mathcal{P}^{(2)} \circ \Pi^{[\chi]} = 0$ when χ is non–TT (cf. Lemma 10.59).

Corollary 10.233 (Compatibility with currents and T–L–O stamps)**.** *The currents of Section 10.19.4 inherit the same triad split:* $\hat{\mathcal{J}}^{\Phi}_{(\Theta)} = \bigoplus_a \hat{\mathcal{J}}^{\Phi}_{(\Theta);a}$ *with* $\hat{\mathcal{J}}^{\Phi}_{(\Theta);a}$ *driven by* $\mathbf{U}_{\Lambda;a}$*. Running the sector execution functor (Definition 9.153) with the usual admission gate (Definition 10.94) stamps* $\mathrm{TLO}^{[\chi]}$ *per admitted type; TT stamps* $\mathrm{TLO}^{[\mathrm{TT}]}$ *remain unaffected by non–TT contributions within any* $(a|b)$ *block (cf. Propositions 10.22 and 10.35 and Lemma 10.168).*

Chapter 11

The Graviton and Chronogeometric Gravity

11.1 Quantization and Interaction of the Geometric Sector

The geometric sector is quantized on the *chronogeometric background* $\left(\mathcal{X}_q, \bar{\Sigma}_{\mu\nu}\right)$ introduced in Chapter 6, where $\bar{\Sigma}_{\mu\nu} := \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ is the expectation-level chronogeometric field obtained from the operator Hessians $(\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu})$ of composite distinguishability/temporal functionals (Chapter 5; cf. $\bar{\Sigma}$ and the construction in Section 5.2). All covariant operations below use the *Relational Levi–Civita* connection $\bar{\nabla}$ uniquely compatible with $\bar{\Sigma}$ $\left(\bar{\nabla}_\lambda \bar{\Sigma}_{\mu\nu} = 0\right)$; see Definition 6.7. The *typed carrier* for radiative spin–2 is fixed once and for all by the *background-covariant TT projector* $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ from Chapter 11 (York-type, $\bar{\Sigma}$-self-adjoint, idempotent, elliptic on the realized weave configuration). We *define* the graviton field operator by

$$\boxed{\hat{h}_{\mu\nu} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \delta\hat{g}_{\mu\nu}}, \qquad \bar{g}^{\mu\nu}\hat{h}_{\mu\nu} = 0, \quad \bar{\nabla}^{\mu}\hat{h}_{\mu\nu} = 0,$$

so that transversality and tracelessness hold *by construction* (TT calculus of Chapter 11). Crucially, *temporal/coherence* fluctuations $\hat{\tau}_{\mu\nu} := \delta\hat{t}_{\mu\nu}$ are *TT-null* in the weave (Chapter 11) and therefore cannot radiate spin–2; this *insulates* the typed graviton $\hat{h}_{\mu\nu}$ from the τ-kairos sector in all physical readouts.

All quantization and interaction statements are carried out on a *certified plateau* $\mathfrak{P}$ in the sense of Chapter 8, using BRST-closed *band tags* $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell} = \left(P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}^{(\ell)}\right)$ that commute with both the *triad projectors* $\Pi^{(a)}$ for $a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$ and the typed projector $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ (Definitions 6.280 and 9.10; cf. the BRST/clean band formalism in eq. (7.10) and Proposition 7.3). On $\mathfrak{P}$ the *physical sector* is $\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}})$ (Chapter 8); all realized kernels and spectra are *Nielsen-flat* (gauge-parameter independent), *hyperbolic* with respect to the $\bar{\Sigma}$-cone, and enjoy *Källén–Lehmann positivity* for gauge-invariant probes. In particular, canonical equal-time commutators and the *Peierls bracket* will be imposed *after* the typed/triadic selection, ensuring that their supports remain $\bar{\Sigma}$-causal (Chapters 8–10).

The *triadic composition* remains explicit at every stage. We decompose

$$\hat{h}_{\mu\nu} = \hat{h}_{\mu\nu}^{(\mathrm{corr})} + \hat{h}_{\mu\nu}^{(\mathrm{vac})} + \hat{h}_{\mu\nu}^{(\mathrm{spec})}, \qquad \hat{h}_{\mu\nu}^{(a)} := \Pi^{(a)}\,\hat{h}_{\mu\nu}, \quad [\Pi^{(a)}, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}] = 0,$$

so that creation–annihilation operators carry a *sector label* a and the *Fock space factorizes* across the triad (cf. the sector bookkeeping from Chapter 5 and the typed weave of Chapter 11). The *TT two-point* function admits a *triad-resolved Källén–Lehmann representation* on $\mathcal{H}_{\mathrm{phys}}$,

$$\Delta^{(2),a}(q) = \frac{Z_a}{q^2 + i\epsilon} + \int_0^\infty \frac{\rho_a^{(2)}(\mu^2)\, d\mu^2}{q^2 - \mu^2 + i\epsilon}, \qquad Z_a \geq 0,\ \rho_a^{(2)}(\mu^2) \geq 0,$$

with the *residue vector* $\vec{Z} = (Z_{\mathrm{corr}}, Z_{\mathrm{vac}}, Z_{\mathrm{spec}})$ read *on the plateau* (Chapter 8). This is the precise sense in which the realized graviton is *TT-typed* yet may carry a *triadic admixture* in its spectral weight; the admixture is physical only insofar as it survives the BRST/plateau certification (Proposition 7.3, positivity statements in Chapter 8).

Quantization proceeds in two complementary, equivalent languages. In the *canonical* approach, we impose the equal-time canonical algebra on the *typed* variables $(\hat{h}_{\mu\nu}, \hat{\Pi}^{\mu\nu})$ after band-sandwiching by $P_x(\lambda_{\mathrm{UV}})$ and projecting with $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$; this preserves BRST hygiene and keeps commutators localized within the $\bar{\Sigma}$-induced foliation (Chapter 10). In the *covariant* approach, we use the *Peierls bracket* generated by the *TT-projected Lichnerowicz operator* on $\left(\mathcal{X}_q, \bar{\Sigma}\right)$, so that the commutator is the *difference of TT retarded/advanced Green operators* with support on the $\bar{\Sigma}$-*null cone* (Chapters 8 and 10). Both perspectives are strictly equivalent on $\mathfrak{P}$ and define the same realized propagator; we will use whichever is most convenient for the task at hand.

Interactions are organized as typed reductions of the *bi-local coupling* between the geometry- and matter-like blocks set up in Chapter 11 (the $\Phi\Phi \leftrightarrow \Theta\Theta$ off-diagonal of the unified correlator, Chapter 6). Concretely, the linearized graviton–Tantrakyma vertex in this part,

$$S_{\mathrm{int}}^{(1)}[\hat{h}, \hat{\Theta}] = \int d\mu(x)\ \hat{h}_{\mu\nu}(x)\, \hat{V}^{\mu\nu}(x; \hat{\Theta})\,,$$

is understood as the *TT-typed shadow* (left leg TT-projected) of that *bi-local parent*; Hermiticity, causal support restricted to the $\bar{\Sigma}$-cone, and *TT-preservation* under composition follow from the weave algebra and the TT-nullity of the τ-typed right leg (Chapter 11). This guarantees that *spin–2 radiative content* is purely metric and cannot be sourced by τ-kairos fluctuations, in agreement with the TT-purity results of Chapter 11.

Finally, we attach a *typed Parasymmetric TLO* layer (Chapter 9) *for the graviton only*. The *law witness* is the *TT-projected* linearized Einstein operator,

$$\mathbf{R}^{[\mathrm{TT}]} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\Big[\delta\bar{G}_{\mu\nu}[\hat{h}] + \Lambda\, \hat{h}_{\mu\nu} - \kappa\, \delta\bar{T}^{\mathrm{total}}_{\mu\nu}\Big], \qquad \Lambda_{\mathrm{law}}^{[\mathrm{TT}]} := 1 - \frac{\|\mathbf{R}^{[\mathrm{TT}]}\|_{\mathcal{I}}}{\|\cdot\|_{\mathrm{ref}}} \in [0,1],$$

which evaluates the *law channel* on $\mathfrak{P}$ (Nielsen-flat, Chapter 7). The *state visibility* $\Lambda_{\mathrm{state}}^{[\mathrm{TT}]}$ is read from the TT two-point KL content (pole residue Z_a and positive spectral density $\rho_a^{(2)}$ in each sector) and synthesized across the triad. Two *type-specific diagnostics* complete the ledger: the *TT-purity* residual $\|\Pi_{\bar{\Sigma}}^{\perp[\mathrm{TT}]}\mathfrak{O}\|_{\mathcal{I}}$ (absence of non-TT contamination) and the *TT-leak* $\varepsilon_{\mathrm{TT}}$ (forbidden inflow from non-TT channels). The canonical *QI–QIV resolver* of Chapter 9 (Chapter 8) is then applied to $\left(\Lambda_{\mathrm{law}}^{[\mathrm{TT}]}, \Lambda_{\mathrm{state}}^{[\mathrm{TT}]}\right)$ with the standard plateau tolerances set in Chapter 8, ensuring that every claim about graviton propagation, coupling, and exchange is *diagnostic realized reference configuration-certified*, *typed*, and *sector-aware* before we proceed to mode decompositions, propagators, and scattering.

Definition 11.1 (Typed structure on the realized weave configuration)**.** Work on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$ with BRST projector $\mathsf{P}_{\mathrm{phys}}$, band tag $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell} = (P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}^{(\ell)})$, and realized background $(\mathcal{X}_q, \bar{\Sigma})$. A *typed* object is obtained by restricting (or decomposing) a field/operator to the image of a commuting family of idempotent projectors ("types"), drawn from:

- the TT projector on symmetric rank-2 tensors, $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ (York–Lichnerowicz; eq. (10.4) and Definition 10.3);
- the triad provenance projectors $\Pi^{(\mathrm{spec})}, \Pi^{(\mathrm{vac})}, \Pi^{(\mathrm{corr})}$ (Definition 10.2);
- the τ–kairos typed idempotents $S^{(b)}, S^{(L)}, \tilde{S}^{(p)}, A^{(+)}, A^{(-)}$ (scalar/vector split with helicity resolution; Definition 10.4);
- CP refinements in the scalar (radial) subspace, $\Pi^{(\mathrm{CP}\pm)}$ (Definition 10.5);
- Riesz spectral selectors for spin content, $\mathsf{P}^{(2)}_{\mathrm{spec}}, \mathsf{P}^{(0)}_{\mathrm{spec}}$ (Definition 10.7);
- the certified band projectors $P_x(\lambda_{\mathrm{UV}})$, $K_{\mathrm{IR}}^{(\ell)}$ (Definitions 6.280 and 9.10).

These projectors commute (commutation stability; Proposition 10.9), so any composition or "audit move" preserves type. A *typed operator* $O^{[\chi]}$ obeys $O^{[\chi]} = \Pi^{[\chi]} O \Pi^{[\chi]}$ and maps $\operatorname{Ran}\Pi^{[\chi]}$ to itself; a *typed field* is the corresponding restriction to $\operatorname{Ran}\Pi^{[\chi]}$.

Cross-reference to Chapters 9–10. The operational use of types with projector sandwiches and triad filters is set up in the Chapter-10 typed calculus (Definition 9.116), and the evaluation of *witnesses* (law/state visibilities and quadrant decisions) is organized in Chapter-9 (Definitions 8.4 and 8.5). In this chapter, all "typed" statements (ETCR, Peierls, propagators, and TT ledgers) are to be read with this convention and are compatible with the witness/quadrant machinery via the same $\Pi^{[\chi]}$ sandwiching.

Canonical examples (Chapter 12).

1. *Typed canonical pair and ETCR.* $\hat{h}_{\mu\nu} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \delta\hat{g}_{\mu\nu}$ and $\hat{\Pi}^{\mu\nu} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \hat{\pi}^{\mu\nu}\, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ (Definition 11.3) satisfy the equal-time commutator *restricted to the TT subspace* (Proposition 11.4), with the band sandwich $P_x(\lambda_{\mathrm{UV}})$.

2. *Typed Lichnerowicz/Peierls.* The kinetic operator $K^{[\mathrm{TT}],(a)}$ (with $a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$) and the Pauli–Jordan bi-solution $\Delta^{[\mathrm{TT}]}$ are TT-typed, yielding typed microcausality and canonical–Peierls equivalence on $\mathfrak{P}$ (Definition 11.60 and Proposition 11.14).

3. *Typed propagators and KL.* $G^{\mathrm{F},[\mathrm{TT}],(a)}$ and the triad-resolved Källén–Lehmann form are constructed within the typed subspace (eq. (11.91)).

4. *Typed witnesses and ledgers.* TT law residual/visibility $\mathbf{R}^{[\mathrm{TT}]}$, $\Lambda_{\mathrm{law}}^{[\mathrm{TT}]}$ use TT-typed meters and remain stable under audit moves (Definition 11.6).

Orthogonal decompositions behind "typed". On the symmetric 2-tensor carrier,

$$\mathbf{1} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \oplus \Pi_{\bar{\Sigma}}^{\perp[\mathrm{TT}]}, \qquad \Pi_{\bar{\Sigma}}^{\perp[\mathrm{TT}]} \cong \underbrace{\operatorname{Ran} S^{(b)} \oplus \operatorname{Ran} S^{(L)} \oplus \operatorname{Ran} \tilde{S}^{(p)}}_{\text{scalar } \tau\text{-types}} \oplus \underbrace{\operatorname{Ran} A^{(+)} \oplus \operatorname{Ran} A^{(-)}}_{\text{vector } \tau\text{-types}},$$

with the scalar radial sector optionally refined by $\Pi^{(\mathrm{CP}\pm)}$ (Lemma 10.8 and Definition 10.5). Thus, "typed" statements insure that calculations, brackets, and propagators are carried out *inside* the intended projector image and are compatible with triad provenance and band/BRST structure.

Witnesses (reminder and Chapter-12 specialization). A *witness* is a band-/BRST-clean, gauge-invariant, *typed* functional producing a visibility $\Lambda \in [0,1]$ that certifies a law- or state-level property inside a projector-defined type; see the sectoral definitions and quadrant rule in Ch. 9 (Definitions 8.4 and 8.5) and the typed projector calculus in Ch. 10 (Definition 9.116). Concretely, all evaluations are performed on the certified plateau $\mathfrak{P}$ with projector sandwiches that commute (commutation stability; Proposition 10.9), hence are invariant under audit moves.

TT specialization used in this chapter. With the type fixed to the graviton carrier $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$, the witnesses employed are:

- **TT law residual/visibility** (Definition 11.6):

$$\mathbf{R}^{[\mathrm{TT}]} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\Big[\delta\bar{G}_{\mu\nu}[\hat{h}] + \Lambda\,\hat{h}_{\mu\nu} - \kappa\,\delta\bar{T}^{\mathrm{total}}_{\mu\nu}\Big], \qquad \Lambda^{[\mathrm{TT}]}_{\mathrm{law}} := 1 - \|\mathbf{R}^{[\mathrm{TT}]}\|_{\mathcal{I}}/\|\cdot\|_{\mathrm{ref}}.$$

  It admits the Peierls "hook" representation $\mathbf{R}^{[\mathrm{TT}]} \sim \{\cdot, S^{[\mathrm{TT}]}\}_{\mathrm{P}}$ (Corollary 11.12), where $S^{[\mathrm{TT}]}$ is the TT-projected quadratic action on $\mathfrak{P}$.

- **TT purity/leak** (Definition 11.7): for any TT readout object $\mathfrak{O}$,

$$\varepsilon_{\mathrm{TT}} := \frac{\|\Pi_{\bar{\Sigma}}^{\perp[\mathrm{TT}]}\mathfrak{O}\|_{\mathcal{I}}}{\|\mathfrak{O}\|_{\mathcal{I}}},$$

  which certifies TT-purity (small leak) and is Nielsen-flat on the plateau.

- **Quadrant assignment in the TT type:** the pair $\left(\Lambda^{[\mathrm{TT}]}_{\mathrm{law}}, \Lambda^{[\mathrm{TT}]}_{\mathrm{state}}\right)$ feeds the standard law/state quadrant decision inherited from Ch. 9 (QI–QIV; Definition 8.5), with the typed/triad projector sandwich from Ch. 10 (Definition 9.116).

All Peierls and ETCR statements in this chapter are to be read in the typed sense (TT sandwiches understood), cf. Definition 11.1 and Propositions 11.4 and 11.14.

Instrumentation (Chs. 9–10 reminder). *Sector-realized weave configuration* collects the standardized meters/tags and evaluation settings used to compute witness visibilities in a given sector $[\mathscr{S}]$. It fixes the arena $\mathcal{A}$ (bulk/boundary), an IR knob r, the band tag $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell} = (P_x(\lambda_{\mathrm{UV}}), K^{(\ell)}_{\mathrm{IR}})$, and the BRST projection $\mathsf{P}_{\mathrm{phys}}$, so that all law/state witnesses are evaluated on the same instrumental frame (Definitions 6.280, 8.4, 8.5, 9.10 and 9.116). In short, instrumentation specifies the calibrated *how* measurements are performed (meters, UV/IR windows, and BRST hygiene) before any typed restriction is applied.

Certified realized weave configuration (Chs. 11–12 usage). In the weave chapters, the instrumentation is *instantiated and frozen* as the *realized weave configuration* $\mathcal{I}^{[\mathcal{S}]}$, now including the realized chronogeometric background $\bar{\Sigma}$. This fixed frame is used for all typed decompositions, selection rules, and dynamical statements; it packages the band tag, the physical cohomology $\mathcal{H}_{\text{phys}}$, and the commuting projector family (TT, triad, τ-types, CP, spectral) with commutation stability (Definitions 10.1 and 10.3, Propositions 10.9 and 10.10, and eq. (10.4)). Operationally, working "on the rig" means every construction is band-/BRST-clean and $\bar{\Sigma}$-covariant by design.

Interaction with types and witnesses (bridge to Ch. 12). Types are the projector-defined subspaces inside which objects are restricted (e.g. TT) and are formalized here by Definition 11.1 (cf. Ch. 10 Definition 9.116). Witnesses are typed, gauge-invariant functionals that output visibilities in $[0,1]$ and drive the law/state quadrant logic from Ch. 9 (Definitions 8.4 and 8.5). In this chapter, TT-typed statements—canonical algebra and Peierls equivalence—are all read on the realized weave configuration with projector sandwiches understood (Propositions 11.4 and 11.14), and TT witnesses such as the law residual/visibility and the Peierls hook are evaluated in the same frame (Definition 11.6 and Corollary 11.12).

11.1.1 Canonical Quantization of the Gaiamiti Field

To quantize the geometric sector we work on the chronogeometric background $(\mathcal{X}_q, \bar{\Sigma}_{\mu\nu})$ of Chapter 6 with the Relational Levi–Civita connection $\bar{\nabla}$ (see Definition 6.7). We impose, *from the outset*, the background–covariant TT typing of Chapter 11 by projecting the metric fluctuation $\delta\hat{g}_{\mu\nu}$ with the York-type projector $\Pi_{\bar{\Sigma}}^{[\text{TT}]}$. All canonical statements are evaluated on a certified plateau $\mathfrak{P}$ (Chapter 8) and within BRST-closed band tags $\mathfrak{B}_{\lambda_{\text{UV}},\ell} = \left(P_x(\lambda_{\text{UV}}), K_{\text{IR}}^{(\ell)}\right)$ (Definitions 6.280 and 9.10), which commute with physical projectors (cf. Proposition 7.3).

Canonical Conjugate Momentum

Definition 11.2 (Canonical Conjugate Momentum of the Gaiamiti Field)**.** Let $S_\Phi[\hat{\mathbf{\Phi}}]$ be the action of the Gaiamiti field with operator-valued Lagrangian density $\mathcal{L}_\Phi$ (Definition 6.172). The *canonical momentum* conjugate to $\hat{\mathbf{\Phi}}(x)$ is

$$\boxed{\hat{\mathbf{\Pi}}_\Phi(x) := \frac{\delta \mathcal{L}_\Phi}{\delta\left(\nabla_0 \hat{\mathbf{\Phi}}(x)\right)}}, \tag{11.1}$$

where ∇_0 is the covariant derivative along the emergent temporal direction. For the quadratic Gaiamiti sector this yields $\hat{\mathbf{\Pi}}_\Phi(x) = 2\,\bar{\Sigma}^{0\nu}(x)\left(\nabla_\nu \hat{\mathbf{\Phi}}(x)\right)^\dagger$ up to conventional normalizations.

$S_\Phi[\hat{\mathbf{\Phi}}]$ (Gaiamiti action). The action of the Gaiamiti sector is the integral of the scalar Lagrangian density $\mathcal{L}_\Phi$ (built from the operator-valued kernel $\mathscr{L}_\Phi^{\text{op}}$) over the emergent configuration space $(\mathcal{X}_q, \bar{\Sigma})$:

$$S_\Phi[\hat{\mathbf{\Phi}}] \;=\; \int_{\mathcal{X}_q} d\mu(x)\; \mathcal{L}_\Phi(x) \;=\; \int_{\mathcal{X}_q} d^d x\, \sqrt{-\det \bar{\Sigma}(x)}\; \mathcal{L}_\Phi(x), \tag{11.2}$$

where $d\mu(x) = \sqrt{-\det \bar{\Sigma}(x)}\, d^d x$ is the relational volume element (see Definition 6.33), and $\mathcal{L}_\Phi$ is specified in Definition 6.172. The kinetic term uses the inverse chronogeometric tensor $\bar{\Sigma}^{\mu\nu}$ (see Definition 6.24) and the vacuum–tangent Hermitian form g_T.

$\mathcal{L}_\Phi$ (operator–trace scalar density).

$$\boxed{\mathcal{L}_\Phi(x) := \mathrm{Tr}\Big\{ \bar{\Sigma}^{\mu\nu}(x)\, g_T\Big(\nabla_\mu \hat{\mathbf{\Phi}}(x),\, \nabla_\nu \hat{\mathbf{\Phi}}(x)\Big) + \mathcal{V}_\Phi\Big(\hat{\mathbf{\Phi}}(x)\Big)\Big\}} \tag{11.3}$$

(see eq. (6.243)), with $\bar{\Sigma}^{\mu\nu}$ the inverse of $\bar{\Sigma}_{\mu\nu}$ (see Definition 6.24) and g_T the Hermitian inner product on $T\mathcal{M}_{\mathrm{vac}}$.

∇_0 (time component of the split covariant derivative). The total covariant derivative is the split connection

$$\nabla_\mu \;=\; \nabla_\mu^{(\mathrm{rel})} \;+\; D_\mu, \qquad D_\mu \;=\; \partial_\mu \;+\; i[\mathcal{A}_\mu, \cdot], \tag{11.4}$$

as in eq. (6.314), combining the torsion-free, $\bar{\Sigma}$-compatible Relational Levi–Civita piece $\nabla_\mu^{(\mathrm{rel})}$ with $\nabla_\mu^{(\mathrm{rel})}\bar{\Sigma}_{\alpha\beta} = 0$ (see Definition 6.7) and the operator–bundle (gauge) part D_μ (see Definitions 6.184 and 6.192). The symbol ∇_0 denotes the component along the emergent temporal direction used in canonical (equal-time) constructions.

$\bar{\Sigma}^{0\nu}$ (mixed time–space inverse components). The entries $\bar{\Sigma}^{0\nu}$ are the 0ν components of $\bar{\Sigma}^{\mu\nu}$, the inverse of $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$; they raise the index on the time derivative in the kinetic density and enter the canonical momentum (see Definitions 6.24 and 11.2).

$(\cdot)^\dagger$ (fiber adjoint on $\mathrm{End}(\mathcal{H}_{\mathrm{em}})$). All operator kernels live in $\mathrm{End}(\mathcal{H}_{\mathrm{em}})$ and are traced to produce c-number densities; $*$–compatibility of the bundle connection ensures Hermiticity of kinetic structures and invariance under fiber $*$–automorphisms within the action.

"Quadratic Gaiamiti sector." The part of $\mathcal{L}_\Phi$ quadratic in first derivatives of $\hat{\mathbf{\Phi}}$, namely the kinetic bilinear $\bar{\Sigma}^{\mu\nu}\, g_T(\nabla_\mu \hat{\mathbf{\Phi}}, \nabla_\nu \hat{\mathbf{\Phi}})$ (with potential $\mathcal{V}_\Phi$ omitted for defining the canonical momentum), as in eq. (6.243).

Functional/Fréchet derivative $\dfrac{\delta \mathcal{L}_\Phi}{\delta(\nabla_0 \hat{\mathbf{\Phi}})}$. Variations are taken at fixed background $\bar{\Sigma}$ and fixed bundle data $\mathcal{A}_\mu$, exactly as in the covariant variational principle leading to the Gaiamiti EL equations (cf. the discussion following $S_\Phi = \int d\mu\, \mathcal{L}_\Phi$ and Definition 11.2).

Short derivation (quadratic Gaiamiti sector) Work with the kinetic density (suppressing $\mathcal{V}_\Phi$):

$$\mathcal{T}_\Phi = \mathrm{Tr}\Big\{ \bar{\Sigma}^{\mu\nu}\, (\nabla_\mu \hat{\mathbf{\Phi}})^\dagger (\nabla_\nu \hat{\mathbf{\Phi}}) \Big\} + \mathrm{h.c.} \qquad \text{(Hermitian by } *\text{–compatibility)}. \tag{11.5}$$

This is the standard Hermitian realization of the bilinear g_T (compare the symmetrization pattern used for kinetic bilinears in Ch. 10).

Vary only the "velocity" variable $\nabla_0\hat{\mathbf{\Phi}}$ (holding $\bar{\Sigma}, \mathcal{A}_\mu$ fixed). The variation of the first trace gives

$$\delta\mathcal{T}_\Phi = \mathrm{Tr}\Big\{\bar{\Sigma}^{\mu\nu}\,(\nabla_\mu\hat{\mathbf{\Phi}})^\dagger\,\delta(\nabla_\nu\hat{\mathbf{\Phi}})\Big\} + \mathrm{h.c.} \tag{11.6}$$

so picking the $\nu = 0$ component and, from the Hermitian conjugate, the $\mu = 0$ component,

$$\delta\mathcal{T}_\Phi = \mathrm{Tr}\Big\{\bar{\Sigma}^{\mu 0}\,(\nabla_\mu\hat{\mathbf{\Phi}})^\dagger\,\delta(\nabla_0\hat{\mathbf{\Phi}})\Big\} + \mathrm{Tr}\Big\{\bar{\Sigma}^{0\nu}\,(\nabla_\nu\hat{\mathbf{\Phi}})^\dagger\,\delta(\nabla_0\hat{\mathbf{\Phi}})\Big\}\,. \tag{11.7}$$

Using the symmetry $\bar{\Sigma}^{\mu 0} = \bar{\Sigma}^{0\mu}$, the two contributions are identical; hence

$$\delta\mathcal{T}_\Phi = 2\,\mathrm{Tr}\Big\{\bar{\Sigma}^{0\nu}\,(\nabla_\nu\hat{\mathbf{\Phi}})^\dagger\,\delta(\nabla_0\hat{\mathbf{\Phi}})\Big\}\,. \tag{11.8}$$

By definition (cf. Definition 11.2),

$$\boxed{\hat{\mathbf{\Pi}}_\Phi := \frac{\delta\mathcal{L}_\Phi}{\delta(\nabla_0\hat{\mathbf{\Phi}})} \qquad \Rightarrow \qquad \hat{\mathbf{\Pi}}_\Phi(x) = 2\,\bar{\Sigma}^{0\nu}(x)\Big(\nabla_\nu\hat{\mathbf{\Phi}}(x)\Big)^\dagger} \tag{11.9}$$

(up to the overall normalization convention adopted for the symmetric kinetic bilinear). This uses the same variational conventions as in the Noether constructions (Fréchet derivative at fixed backgrounds).

Definition 11.3 (TT-projected canonical variables (project first, then quantize))**.** Let $\delta\hat{g}_{\mu\nu}$ be the Gaiamiti (metric) fluctuation. The *typed* TT graviton and its *typed* TT canonical momentum are defined using the background–covariant TT projector $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ (eq. (10.4) and Definition 10.3) as

$$\boxed{\begin{aligned}\hat{h}_{\mu\nu} &:= \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{g}_{\mu\nu}\,, &(11.10)\\ \hat{\Pi}^{\mu\nu} &:= \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\hat{\pi}^{\mu\nu}\,\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,, &(11.11)\end{aligned}}$$

where the unprojected canonical momentum $\hat{\pi}^{\mu\nu}$ is obtained by the covariant variational rule (cf. Gaiamiti momentum, Definition 11.2)

$$\hat{\pi}^{\mu\nu} := \frac{\delta\mathcal{L}_\Phi}{\delta\Big(\nabla_0\,\delta\hat{g}_{\mu\nu}\Big)}\,,$$

with ∇_0 the temporal component of the split connection $\nabla_\mu = \nabla^{(\mathrm{rel})}_\mu + D_\mu$ (eq. (6.314) and Definition 6.7).

By construction and the properties of $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ (idempotent, trace/longitudinal annihilation), the TT variable satisfies

$$\bar{g}^{\mu\nu}\hat{h}_{\mu\nu} = 0, \qquad \bar{\nabla}^\mu\hat{h}_{\mu\nu} = 0$$

(*TT calculus*; see Definition 10.3), where $\bar{\nabla}$ is the $\bar{\Sigma}$-compatible Levi–Civita derivative (Definition 6.7). Moreover, the temporal/coherence fluctuation $\hat{\tau}_{\mu\nu} := \delta\hat{t}_{\mu\nu}$ is *TT-null*:

$$\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\hat{\tau}_{\mu\nu} = 0,$$

so it does not contribute to $\hat{h}_{\mu\nu}$ (see the τ–kairos typing and TT–null statements in Definition 10.4).

Notation and mathematical meaning in Proposition 11.4. The equal-time commutator on a foliation slice $\Sigma_t \subset (\mathcal{X}_q, \bar{\Sigma})$ reads

$$[\hat{h}_{\mu\nu}(t,\mathbf{x}),\, \hat{\Pi}^{\rho\sigma}(t,\mathbf{y})] = i\,\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta_{\mu\nu}{}^{\rho\sigma}\,\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,P_x(\lambda_{\mathrm{UV}})\,\delta^{(d-1)}(\mathbf{x}-\mathbf{y}),$$

where t is the emergent temporal parameter that foliates $(\mathcal{X}_q, \bar{\Sigma})$ into Cauchy hypersurfaces Σ_t via the $\bar{\Sigma}$-compatible Levi–Civita connection $\bar{\nabla}$ (Definition 6.7); bold $\mathbf{x}, \mathbf{y}$ are spatial coordinates on Σ_t (dimension $d-1$). The Dirac distribution $\delta^{(d-1)}(\mathbf{x}-\mathbf{y})$ on Σ_t is characterized by $\int_{\Sigma_t} d^{d-1}\mathbf{y}\,\delta^{(d-1)}(\mathbf{x}-\mathbf{y})\,f(\mathbf{y}) = f(\mathbf{x})$ for test fields f. The UV spectral projector $P_x(\lambda_{\mathrm{UV}})$ arises from the band tag $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell} = (P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}^{(\ell)})$, enforces band support, and commutes with the typed/projected meters (tag hygiene; Definitions 6.280 and 9.10 and Lemma 9.11). The York–Lichnerowicz map $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ is the background-covariant TT projector on symmetric rank–2 tensors (idempotent, self-adjoint, annihilating trace and longitudinal parts; eq. (10.4) and Definition 10.3). The bi-tensor $\delta_{\mu\nu}{}^{\rho\sigma}$ is the symmetric identity: $\delta_{\mu\nu}{}^{\rho\sigma}A_{\rho\sigma} = \frac{1}{2}(\delta_\mu{}^\rho\delta_\nu{}^\sigma + \delta_\mu{}^\sigma\delta_\nu{}^\rho)A_{\rho\sigma}$. The bracket $[\,\cdot\,,\,\cdot\,]$ denotes the operator commutator, and "equal time" means both operators are evaluated on the same slice Σ_t.

Index conventions (clarifying $\mu\nu$ and $\rho\sigma$). Greek indices μ, ν, ρ, σ label spacetime tensor components on $(\mathcal{X}_q, \bar{\Sigma})$, running over $\{0, 1, \ldots, d-1\}$; indices are raised/lowered with $\bar{\Sigma}^{\alpha\beta}$, $\bar{\Sigma}_{\alpha\beta}$. The foliation index 0 designates the emergent temporal direction; spatial components are indexed by Latin letters $i, j, k, l \in \{1, \ldots, d-1\}$ on Σ_t. In the canonical equal-time algebra, one may equivalently write

$$[\hat{h}_{ij}(t,\mathbf{x}),\, \hat{\Pi}^{kl}(t,\mathbf{y})] = i\,\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta_{ij}{}^{kl}\,\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,P_x(\lambda_{\mathrm{UV}})\,\delta^{(d-1)}(\mathbf{x}-\mathbf{y}),$$

with components carrying a 0-index playing the role of constraints/non-dynamical variables, while the TT projector restricts to the transverse–traceless *spatial* sector on Σ_t. Symmetrization and antisymmetrization follow the standard conventions $A_{(\mu\nu)} := \frac{1}{2}(A_{\mu\nu} + A_{\nu\mu})$, $A_{[\mu\nu]} := \frac{1}{2}(A_{\mu\nu} - A_{\nu\mu})$.

Canonical pair, distributional nature, and band/BRST structure. The typed variables $\hat{h}_{\mu\nu}$ and $\hat{\Pi}^{\rho\sigma}$ are the TT-projected canonical pair (Definition 11.3), obtained from the unprojected metric fluctuation $\delta\hat{g}_{\mu\nu}$ and its momentum $\hat{\pi}^{\rho\sigma} = \delta\mathcal{L}_\Phi/\delta(\nabla_0\delta\hat{g}_{\rho\sigma})$ (Definition 11.2), where ∇_0 is the temporal component of the split connection $\nabla_\mu = \nabla_\mu^{(\mathrm{rel})} + D_\mu$ (eq. (6.314)). Mathematically, Proposition 11.4 states that these TT-projected fields are operator-valued distributions on Σ_t obeying the standard equal-time canonical algebra *restricted to the TT subspace* by the projector sandwich $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}(\cdot)\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$. Band regularization appears explicitly through $P_x(\lambda_{\mathrm{UV}})$ and implicitly through the IR window $K_{\mathrm{IR}}^{(\ell)}$, which preserves $\bar{\Sigma}$-causal cones and ensures the existence of the equal-time limit (Remark 9.83). BRST cleanliness and invariance under gauge-parameter deformations on a certified plateau $\mathfrak{P}$ follow from the cohomological projection to $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ (Proposition 7.3) together with the commutation properties of the realized weave configuration operators (Proposition 10.9).

Proposition 11.4 (ETCR on the band; BRST-healthy, regulator-safe)**.** *On a certified plateau* $\mathfrak{P}$ *with band tags* $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ *that commute with* $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ *(Definitions 6.280 and 9.10), the equal-time canonical commutator for the typed variables takes the form*

$$\boxed{\left[\hat{h}_{\mu\nu}(t,\mathbf{x}),\ \hat{\Pi}^{\rho\sigma}(t,\mathbf{y})\right] \;=\; i\;\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta_{\mu\nu}{}^{\rho\sigma}\,\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,P_x(\lambda_{\mathrm{UV}})\,\delta^{(d-1)}(\mathbf{x}-\mathbf{y})\,,} \tag{11.12}$$

with all other equal-time commutators vanishing. Here $\delta_{\mu\nu}{}^{\rho\sigma}$ *is the symmetric identity bitensor on rank-2 tensors. The BRST cleanliness and physical-sector projection are as in Proposition 7.3.*

Proof. Work on a certified plateau $\mathfrak{P}$ where the band tags $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell} = (P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}^{(\ell)})$ are BRST-closed and commute with all typed/projected meters (Definitions 6.280 and 9.10, Lemma 9.11, and Proposition 10.9). Define the TT-typed variables as in Definition 11.3 using the background–covariant projector $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ (eq. (10.4) and Definition 10.3).

Step 1 (unprojected ETCR on the band). By the canonical definition of the metric's conjugate momentum $\hat{\pi}^{\rho\sigma} := \delta\mathcal{L}_\Phi/\delta(\nabla_0\delta\hat{g}_{\rho\sigma})$ (Definition 11.2) and the equal-time canonical algebra constructed via the covariant (Peierls/Dirac) procedure on the band, the unprojected equal-time commutator reads

$$\left[\delta\hat{g}_{\mu\nu}(t,\mathbf{x}),\ \hat{\pi}^{\rho\sigma}(t,\mathbf{y})\right] \;=\; i\,\delta_{\mu\nu}{}^{\rho\sigma}\,P_x(\lambda_{\mathrm{UV}})\,\delta^{(d-1)}(\mathbf{x}-\mathbf{y}), \qquad \left[\delta\hat{g},\delta\hat{g}\right] = \left[\hat{\pi},\hat{\pi}\right] = 0, \tag{11.13}$$

with the UV projector $P_x(\lambda_{\mathrm{UV}})$ explicitly present (band support), and where the equal-time limit is well-defined because the IR window does not enlarge $\bar{\Sigma}$-cones (Remark 9.83).

Step 2 (TT sandwiching). Using $\hat{h}_{\mu\nu} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}_{\mu\nu}$ and $\hat{\Pi}^{\rho\sigma} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\hat{\pi}^{\rho\sigma}\,\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ (Definition 11.3), and the facts that $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ is idempotent/self-adjoint, annihilates traces and longitudinal parts, and commutes with the band tags and typed/triad maps (eq. (10.4), Definition 10.3, and Proposition 10.9), we compute

$$\begin{aligned}
\left[\hat{h}_{\mu\nu},\,\hat{\Pi}^{\rho\sigma}\right] &= \left[\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}_{\mu\nu},\ \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\hat{\pi}^{\rho\sigma}\,\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\right] \\
&= \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\left[\delta\hat{g}_{\mu\nu},\,\hat{\pi}^{\rho\sigma}\right]\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \qquad \text{(commutations from Proposition 10.9)} \\
&= i\;\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta_{\mu\nu}{}^{\rho\sigma}\,\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,P_x(\lambda_{\mathrm{UV}})\,\delta^{(d-1)}(\mathbf{x}-\mathbf{y}),
\end{aligned} \tag{11.14}$$

where in the last line we used (11.13). The vanishing $\left[\hat{h},\hat{h}\right] = \left[\hat{\Pi},\hat{\Pi}\right] = 0$ follows in the same way from (11.13) and the projector sandwiching.

Step 3 (BRST cleanliness and physical projection). Because the band tag is BRST-closed ($[\hat{Q}_{\mathrm{BRST}}, \mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}] = 0$, Lemma 9.11) and $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ is built from $\bar{\Sigma}$-covariant operators, the commutator is a BRST cocycle and descends to the physical Hilbert space $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ with gauge-parameter independence (Nielsen flatness) on $\mathfrak{P}$ (Propositions 7.3 and 10.10).

Collecting the results yields exactly the displayed ETCR on the band, with all other equal-time commutators vanishing. □

At a high level, Proposition 11.4 states that after first projecting into the physically relevant, background–covariant TT sector using $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ (Definition 10.3 and eq. (10.4)) and defining the canonical pair as in Definition 11.3, the graviton field $\hat{h}_{\mu\nu}$ and its momentum $\hat{\Pi}^{\rho\sigma}$ obey the *standard* equal–time canonical commutator, but restricted to the TT subspace. The projector sandwich in the ETCR enforces that only transverse–traceless polarizations carry quantum degrees of freedom; all trace/longitudinal components (and the temporal/coherence sector) are removed prior to quantization, hence the algebra does not leak into non-TT directions (Proposition 10.9). The appearance of $P_x(\lambda_{\mathrm{UV}})$ indicates realization *on the band*, with UV/IR control provided by the band tags $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ (Definitions 6.280 and 9.10), while cone preservation guarantees a well-defined equal-time limit (Remark 9.83).

Equally important, the commutator is *BRST-healthy*: the band tags commute with the BRST charge (tag hygiene), and the TT construction is $\bar{\Sigma}$-covariant, so the algebra descends unchanged to the physical cohomology $\mathcal{H}_{\mathrm{phys}}$ and is gauge-parameter independent on a certified plateau $\mathfrak{P}$ (Lemma 9.11 and Proposition 7.3). In summary, the proposition formalizes the principle *project, then quantize*: imposing TT typing at the operator level and employing regulator-safe band tags realizes the canonical graviton structure precisely within the TT sector, without regulator artifacts or gauge contamination (Definitions 11.2 and 11.3).

Corollary 11.5 (Nielsen safety of the TT algebra)**.** *Along gauge-parameter deformations that preserve $\mathfrak{P}$, the typed ETCR of Proposition 11.4 is invariant (Nielsen-flat): any variation is ζ-exact and therefore trivial on $\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}})$ (Chapter 8).*

Proof. Let α denote a gauge–fixing parameter, and consider deformations $\alpha \mapsto \alpha + \delta\alpha$ that preserve the certified plateau $\mathfrak{P}$ and the band tags $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ (hence tag hygiene and BRST closure remain valid; see Definitions 6.280 and 9.10 and Lemma 9.11). Define the typed canonical variables by $\hat{h}_{\mu\nu} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \delta\hat{g}_{\mu\nu}$ and $\hat{\Pi}^{\rho\sigma} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \hat{\pi}^{\rho\sigma}\, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ as in Definition 11.3, with $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ the background–covariant TT projector (eq. (10.4) and Definition 10.3). The ETCR on the band is given by Proposition 11.4.

Step 1: BRST cocycle property. On $\mathfrak{P}$ the band tag is BRST–closed and commutes with all typed/triad projectors (Lemma 9.11 and Proposition 10.9). Since $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ is built from $\bar{\Sigma}$–covariant operators, $[\,\hat{Q}_{\mathrm{BRST}}, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}] = 0$ as a consequence of the BRST–closed realized weave configuration (Proposition 10.9). Therefore the typed commutator kernel

$$\mathcal{C}_{\mu\nu}{}^{\rho\sigma}(t;\mathbf{x},\mathbf{y}) := \Big[\hat{h}_{\mu\nu}(t,\mathbf{x}),\, \hat{\Pi}^{\rho\sigma}(t,\mathbf{y})\Big]$$

is a BRST cocycle: $[\,\hat{Q}_{\mathrm{BRST}}, \mathcal{C}\,] = 0$. Hence $\mathcal{C}$ defines a class in $\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}})$ (Proposition 7.3).

Step 2: Nielsen identity on the band. On a certified plateau the Slavnov–Taylor/Nielsen framework implies that the α–variation of any BRST–closed realized weave configuration observable is BRST–exact ("Nielsen flatness"): $\partial_\alpha \mathcal{O} = \{\hat{Q}_{\mathrm{BRST}}, \zeta_{\mathcal{O}}\}$ for some local $\zeta_{\mathcal{O}}$ (band version; see the invariance statement for calibrated meters/visibilities, Proposition 10.10, together with the BRST projection properties in Proposition 7.3). Applying this to $\mathcal{C}$ yields

$$\partial_\alpha \mathcal{C}_{\mu\nu}{}^{\rho\sigma}(t;\mathbf{x},\mathbf{y}) = \{\hat{Q}_{\mathrm{BRST}},\, \zeta_{\mu\nu}{}^{\rho\sigma}(t;\mathbf{x},\mathbf{y})\},$$

with ζ constructed from the same realized weave configuration operators (TT/projectors, band tag), which commute with $\hat{Q}_{\mathrm{BRST}}$ by Lemma 9.11 and Proposition 10.9. Therefore $\partial_\alpha \mathcal{C}$ is BRST–exact and vanishes in cohomology, i.e. it is trivial on $\mathcal{H}_{\mathrm{phys}}$.

Combining Steps 1–2 shows that the typed ETCR of Proposition 11.4 is invariant under gauge–parameter deformations that preserve $\mathfrak{P}$; equivalently, it is Nielsen–flat, establishing Corollary 11.5. □

Typed TLO attachment (TT only). In the sense of Chapter IX, diagnostics are attached to the TT sector *only*, using the background–covariant TT projector $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ (eq. (10.4) and Definition 10.3) on a certified plateau $\mathfrak{P}$ (Section 7.3.9).

Definition 11.6 (TT law residual and visibility)**.** Let the TT–projected linearized Einstein operator about $\bar{\Sigma}$ act on the TT graviton $\hat{h}_{\mu\nu}$ defined in Definition 11.3. With the chronogeometric Einstein tensor $\bar{G}_{\mu\nu}[\bar{\Sigma}]$ and total stress tensor $\bar{T}^{\mathrm{total}}_{\mu\nu}$ from Definition 11.106, define the *TT law residual*

$$\boxed{\mathbf{R}^{[\mathrm{TT}]} \;:=\; \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\Big[\delta\bar{G}_{\mu\nu}[\hat{h}] + \Lambda\,\hat{h}_{\mu\nu} - \kappa\,\delta\bar{T}^{\mathrm{total}}_{\mu\nu}\Big]\,,} \tag{11.15}$$

and the associated *law visibility* $\Lambda_{\mathrm{law}}^{[\mathrm{TT}]} := 1 - \|\mathbf{R}^{[\mathrm{TT}]}\|_{\mathcal{I}}/\|\cdot\|_{\mathrm{ref}} \in [0,1]$, to be reported on $\mathfrak{P}$. Here $\|\cdot\|_{\mathcal{I}}$ is the norm specified by the realized weave configuration instrumentation $\mathcal{I}$ (Definition 10.1), and $\|\cdot\|_{\mathrm{ref}}$ is a fixed reference scale consistent with the acceptance pipeline (Section 7.3.9).

Definition 11.7 (TT purity and TT leak diagnostics)**.** Let $\mathfrak{O}$ denote any operator entering the TT readout (modes, propagators, vertices). With $\Pi_{\bar{\Sigma}}^{\perp[\mathrm{TT}]} := \mathbf{1} - \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ (Definition 10.3), the *TT–purity* residual is $\|\Pi_{\bar{\Sigma}}^{\perp[\mathrm{TT}]}\,\mathfrak{O}\|_{\mathcal{I}}$, and the *TT–leak* is a small positive number $\varepsilon_{\mathrm{TT}}$ that bounds any spurious inflow from non–TT channels. On a certified plateau these quantities are gauge–parameter independent (Nielsen–flat) by Corollary 11.5 together with the projector/realized weave configuration commutation properties in Proposition 10.9.

These constructions integrate the canonical TT algebra $\big($ETCR on the band, Proposition 11.4$\big)$ with the typed/triadic, BRST–healthy, plateau–stable framework of Chapters VII–XI (Definitions 10.1 and 10.2 and Propositions 7.3 and 10.9), ensuring that canonical statements descend consistently to the physical sector $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ and admit Parasymmetric TLO diagnostics in the TT type (Definitions 8.4 to 8.6).

Equal-Time Commutation Relations (ETCR)

The essence of canonical quantization is that the field and its conjugate momentum no longer commute. We work on the chronogeometric background $(\mathcal{X}_q, \bar{\Sigma}_{\mu\nu})$ with Relational Levi–Civita connection $\bar{\nabla}$ (Definition 6.7), and we use the canonical momentum of the Gaiamiti field defined in Definition 11.2. Throughout, equal-time refers to the foliation determined by $\bar{\Sigma}$ (cf. Section 5.2.6).

Definition 11.8 (Equal-Time Commutation Relations for the Gaiamiti Field)**.** Let $x = (x^0, \vec{x})$ and $y = (y^0, \vec{y})$ be two points on a Cauchy hypersurface of constant emergent time $x^0 = y^0$. For

the components $\hat{\mathbf{\Phi}}^\lambda$ of the Gaiamiti field and their conjugate momenta $\hat{\mathbf{\Pi}}^\rho_\Phi$ (Definition 11.2), the *equal-time commutation relations* are

$$\boxed{\begin{aligned}\left[\hat{\mathbf{\Phi}}^\lambda(x^0,\vec{x}),\, \hat{\mathbf{\Pi}}^\rho_\Phi(x^0,\vec{y})\right] &= i\,\delta^\lambda{}_\rho\,\delta^{(d-1)}(\vec{x}-\vec{y})\,\mathbb{I}, &(11.16)\\ \left[\hat{\mathbf{\Phi}}^\lambda(x^0,\vec{x}),\, \hat{\mathbf{\Phi}}^\rho(x^0,\vec{y})\right] &= 0, &(11.17)\\ \left[\hat{\mathbf{\Pi}}^\lambda_\Phi(x^0,\vec{x}),\, \hat{\mathbf{\Pi}}^\rho_\Phi(x^0,\vec{y})\right] &= 0. &(11.18)\end{aligned}}$$

Meaning of the notation and of (11.16)–(11.18).

- $x=(x^0,\vec{x})$, $y=(y^0,\vec{y})$: spacetime points restricted to a common Cauchy hypersurface Σ_{x^0} of constant emergent time $x^0=y^0$, defined by the foliation induced by the chronogeometric background $\bar{\Sigma}_{\mu\nu}$; the compatible connection is the Relational Levi–Civita $\bar{\nabla}$ (see Definition 6.7).
- $\delta^{(d-1)}(\vec{x}-\vec{y})$: Dirac delta *on the slice* Σ_{x^0}, normalized so that $\int_{\Sigma_{x^0}} d^{d-1}\vec{y}\;\delta^{(d-1)}(\vec{x}-\vec{y})\,f(\vec{y})=f(\vec{x})$ for test fields f.
- $\hat{\mathbf{\Phi}}^\lambda$, $\hat{\mathbf{\Pi}}^\rho_\Phi$: the components (e.g. in the local/modal basis of Definitions 6.1 and 6.2) of the Gaiamiti field and its canonical momentum, with $\hat{\mathbf{\Pi}}_\Phi=\delta\mathcal{L}_\Phi/\delta(\nabla_0\hat{\mathbf{\Phi}})$ per Definition 11.2; here ∇_0 is the temporal part of the split connection $\nabla_\mu=\nabla^{(\mathrm{rel})}_\mu+D_\mu$ (eq. (6.314)).
- $\delta^\lambda{}_\rho$: the Kronecker symbol on the (field/modal) indices; $\mathbb{I}$ denotes the identity on the fiber $\mathrm{End}(\mathcal{H}_{\mathrm{em}})$.

Equation (11.16). This is the canonical equal–time bracket between the field and its conjugate momentum:

$$\left[\hat{\mathbf{\Phi}}^\lambda(x^0,\vec{x}),\, \hat{\mathbf{\Pi}}^\rho_\Phi(x^0,\vec{y})\right] = i\,\delta^\lambda{}_\rho\,\delta^{(d-1)}(\vec{x}-\vec{y})\,\mathbb{I}.$$

It expresses *locality* on the Cauchy slice through $\delta^{(d-1)}(\vec{x}-\vec{y})$ and *componentwise* pairing through $\delta^\lambda{}_\rho$. The normalization is consistent with $\hbar=1$. In band–regularized settings this bracket acquires the explicit UV projector $P_x(\lambda_{\mathrm{UV}})$ from the band tag $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ (Definitions 6.280 and 9.10), analogously to the TT case in Proposition 11.4.

Equation (11.17).

$$\left[\hat{\mathbf{\Phi}}^\lambda(x^0,\vec{x}),\, \hat{\mathbf{\Phi}}^\rho(x^0,\vec{y})\right] = 0.$$

Equal–time field operators commute for the bosonic Gaiamiti sector; this encodes that no independent canonical pairing occurs among the field components themselves at equal time.

Equation (11.18).

$$\left[\hat{\mathbf{\Pi}}^\lambda_\Phi(x^0,\vec{x}),\, \hat{\mathbf{\Pi}}^\rho_\Phi(x^0,\vec{y})\right] = 0.$$

Equal–time conjugate momenta commute, reflecting that the only nontrivial canonical pairing is the field–momentum one in (11.16). For the quadratic Gaiamiti sector, the momentum reads $\hat{\mathbf{\Pi}}_\Phi = 2\,\bar{\Sigma}^{0\nu}(\nabla_\nu\hat{\mathbf{\Phi}})^\dagger$ up to conventions (Definition 11.2), where $\bar{\Sigma}^{0\nu}$ are components of the inverse chronogeometric tensor (Definition 6.24).

Remark 11.9 (Geometry and BRST projection)**.** All brackets are understood as operator–valued distributions on Σ_{x^0} determined by the foliation of $(\mathcal{X}_q, \bar{\Sigma})$ (cf. Section 5.2.6). On certified plateaux, BRST projection to the physical sector $\mathcal{H}_{\rm phys} = H^0(\hat{Q}_{\rm BRST})$ preserves these equal–time relations and ensures gauge–parameter independence (Nielsen flatness) in the sense of Proposition 7.3.

Lemma 11.10 (TT-reduced ETCR and compatibility with typed variables)**.** *Let $\hat{h}_{\mu\nu}$ and $\hat{\Pi}^{\mu\nu}$ be the TT-typed canonical pair of Definition 11.3, obtained by projecting $\delta\hat{g}_{\mu\nu}$ and its momentum with the background–covariant projector $\Pi_{\bar{\Sigma}}^{[\rm TT]}$ (eq. (10.4) and Definition 10.3). On a certified plateau $\mathfrak{P}$ with BRST-closed band tags $\mathfrak{B}_{\lambda_{\rm UV},\ell} = (P_x(\lambda_{\rm UV}), K_{\rm IR}^{(\ell)})$ that commute with $\Pi_{\bar{\Sigma}}^{[\rm TT]}$ (Definitions 6.280 and 9.10, Lemma 9.11, and Proposition 10.9), restricting (11.16)–(11.18) to the TT sector yields exactly the typed equal-time algebra stated in Proposition 11.4.*

Proof. Step 1 (unprojected equal-time algebra). The canonical momentum for the metric fluctuation is defined covariantly by the same variational rule used for the Gaiamiti sector (cf. Definition 11.2), with ∇_0 the temporal component of the split connection $\nabla_\mu = \nabla_\mu^{(\rm rel)} + D_\mu$ (eq. (6.314) and Definition 6.7). Hence, the equal-time relations (11.16)–(11.18) apply componentwise to $(\delta\hat{g}_{\mu\nu}, \hat{\pi}^{\rho\sigma})$ on the Cauchy slice Σ_t, with the same Dirac delta $\delta^{(d-1)}(\vec{x} - \vec{y})$.

Step 2 (TT projection and commutation with the realized weave configuration). Define the typed variables as in Definition 11.3, $\hat{h}_{\mu\nu} := \Pi_{\bar{\Sigma}}^{[\rm TT]}\,\delta\hat{g}_{\mu\nu}$ and $\hat{\Pi}^{\rho\sigma} := \Pi_{\bar{\Sigma}}^{[\rm TT]}\,\hat{\pi}^{\rho\sigma}\,\Pi_{\bar{\Sigma}}^{[\rm TT]}$. Use that $\Pi_{\bar{\Sigma}}^{[\rm TT]}$ is idempotent and self-adjoint and annihilates trace/longitudinal parts (eq. (10.4) and Definition 10.3), and that it *commutes* with the band tags and the typed/triad projectors on $\mathfrak{P}$ (tag hygiene and weave commutations; Lemma 9.11 and Proposition 10.9). Then

$$\begin{aligned}
\Big[\hat{h}_{\mu\nu}(t,\mathbf{x}), \hat{\Pi}^{\rho\sigma}(t,\mathbf{y})\Big] &= \Big[\Pi_{\bar{\Sigma}}^{[\rm TT]}\delta\hat{g}_{\mu\nu},\ \Pi_{\bar{\Sigma}}^{[\rm TT]}\hat{\pi}^{\rho\sigma}\Pi_{\bar{\Sigma}}^{[\rm TT]}\Big] \\
&= \Pi_{\bar{\Sigma}}^{[\rm TT]}\Big[\delta\hat{g}_{\mu\nu}(t,\mathbf{x}), \hat{\pi}^{\rho\sigma}(t,\mathbf{y})\Big]\Pi_{\bar{\Sigma}}^{[\rm TT]} \\
&\quad \text{(via rigid commutations for TT component; Proposition 10.9)} \\
&= i\,\Pi_{\bar{\Sigma}}^{[\rm TT]}\,\delta_{\mu\nu}{}^{\rho\sigma}\,\Pi_{\bar{\Sigma}}^{[\rm TT]}\,P_x(\lambda_{\rm UV})\,\delta^{(d-1)}(\mathbf{x}-\mathbf{y}),
\end{aligned}$$

which is exactly the ETCR stated in Proposition 11.4. The vanishing brackets follow similarly:

$$\Big[\hat{h}, \hat{h}\Big] = \Pi_{\bar{\Sigma}}^{[\rm TT]}\Big[\delta\hat{g}, \delta\hat{g}\Big]\Pi_{\bar{\Sigma}}^{[\rm TT]} = 0, \qquad \Big[\hat{\Pi}, \hat{\Pi}\Big] = \Pi_{\bar{\Sigma}}^{[\rm TT]}\Big[\hat{\pi}, \hat{\pi}\Big]\Pi_{\bar{\Sigma}}^{[\rm TT]} = 0,$$

by (11.17)–(11.18).

Step 3 (BRST cleanliness and plateau stability). On $\mathfrak{P}$, the band tag $\mathfrak{B}_{\lambda_{\rm UV},\ell}$ is BRST-closed and the realized weave configuration operators (including $\Pi_{\bar{\Sigma}}^{[\rm TT]}$) commute with $\hat{Q}_{\rm BRST}$ (Lemma 9.11 and Proposition 10.9). Therefore the projected ETCR is a BRST cocycle and descends to $\mathcal{H}_{\rm phys} = H^0(\hat{Q}_{\rm BRST})$ with gauge-parameter independence on the plateau (Proposition 7.3 and Corollary 11.5).

Combining Steps 1–3 proves that restricting (11.16)–(11.18) to the TT sector yields precisely the typed algebra of Proposition 11.4. □

Covariance and Interpretation

1. *Quantum uncertainty.* The non-vanishing commutator $\left[\hat{\mathbf{\Phi}}, \hat{\mathbf{\Pi}}_{\Phi}\right] = i\,\mathbb{I}$ in (11.16) encodes the impossibility of simultaneously measuring the configuration and the conjugate rate of change of the geometric field with arbitrary precision, i.e. a geometric Heisenberg principle for the apeiron degrees of freedom.

2. *Locality and microcausality on a Cauchy slice.* The Dirac delta $\delta^{(d-1)}(\vec{x}-\vec{y})$ ensures that the field at $\vec{x}$ has non-trivial commutator only with its conjugate momentum at the *same* spatial point on the equal-time hypersurface; all other equal-time commutators vanish, cf. (11.17)–(11.18).

3. *Covariance via the Peierls bracket.* The notion of "equal time" is defined with respect to the $\bar{\Sigma}$-induced foliation (Section 5.2.6), so the ETCR formalism is not manifestly covariant. Covariance is restored by the *Peierls bracket* built from the TT-projected Lichnerowicz-type wave operator $K_{[\mathrm{TT}]}$ on $(\mathcal{X}_q, \bar{\Sigma})$ (cf. Chapter X and Definition 10.3). Define the advanced/retarded TT Green operators $G^{\mathrm{adv/ret}}_{[\mathrm{TT}]}$ and their causal difference $\Delta^{[\mathrm{TT}]} := G^{\mathrm{ret}}_{[\mathrm{TT}]} - G^{\mathrm{adv}}_{[\mathrm{TT}]}$. For gauge-invariant functionals $F[h], G[h]$ supported on the certified plateau $\mathfrak{P}$,

$$\boxed{\begin{aligned} \{F,G\}_{\mathrm{P}} &= \left\langle \frac{\delta F}{\delta h_{\mu\nu}}, \Delta^{[\mathrm{TT}]\mu\nu}{}_{\rho\sigma} \left(\frac{\delta G}{\delta h_{\rho\sigma}} \right) \right\rangle, && (11.19)\\ \Delta^{[\mathrm{TT}]} &:= G^{\mathrm{ret}}_{[\mathrm{TT}]} - G^{\mathrm{adv}}_{[\mathrm{TT}]}. && (11.20) \end{aligned}}$$

   Functional derivatives are taken at fixed band tags $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell} = (P_x(\lambda_{\mathrm{UV}}), K^{(\ell)}_{\mathrm{IR}})$ (Definitions 6.280 and 9.10); cone preservation of $K^{(\ell)}_{\mathrm{IR}}$ ensures well-defined causal support (Remark 9.83). The equal-time limit of (11.25) reproduces the typed ETCR of Proposition 11.4; thus the canonical ETCRs are the Cauchy-slice restriction of the Peierls construction (Proposition 11.11) [82, 155].

Meaning of F and G in the Peierls bracket. In Proposition 11.4 and the Peierls construction, F and G denote (gauge/BRST-invariant) *observables*, i.e. sufficiently regular functionals of the *TT carrier* $h_{\mu\nu}$. Concretely, they are maps

$$F, G : \ \mathcal{E}_{\mathrm{TT}} \longrightarrow \mathbb{C}, \qquad \mathcal{E}_{\mathrm{TT}} := \{\text{admissible TT configurations on } (\mathcal{X}_q, \bar{\Sigma})\},$$

that are (Gâteaux/Fréchet) differentiable so that their functional derivatives $\delta F/\delta h_{\mu\nu}$ and $\delta G/\delta h_{\rho\sigma}$ exist as tensor-valued (distributional) test objects with the band/realized weave configuration properties of $\mathcal{I}^{[\mathcal{J}]}$ (Definition 10.1). The Peierls bracket

$$\{F,G\}_{\mathrm{P}} = \left\langle \frac{\delta F}{\delta h_{\mu\nu}}, \Delta^{[\mathrm{TT}]\mu\nu}{}_{\rho\sigma} \left(\frac{\delta G}{\delta h_{\rho\sigma}} \right) \right\rangle, \qquad \Delta^{[\mathrm{TT}]} := G^{\mathrm{ret}}_{[\mathrm{TT}]} - G^{\mathrm{adv}}_{[\mathrm{TT}]},$$

is then defined by pairing these derivatives with the TT Pauli–Jordan bi-solution ($\bar{\Sigma}$-microcausal support; Proposition 11.14). In spacetime form,

$$\left\langle \frac{\delta F}{\delta h_{\mu\nu}}, \Delta^{[\mathrm{TT}]\mu\nu}{}_{\rho\sigma} \left(\frac{\delta G}{\delta h_{\rho\sigma}} \right) \right\rangle = \int d\mu_{\bar{\Sigma}}(x) \int d\mu_{\bar{\Sigma}}(y)\, \frac{\delta F}{\delta h_{\mu\nu}(x)}\, \Delta^{[\mathrm{TT}]\mu\nu}{}_{\rho\sigma}(x,y)\, \frac{\delta G}{\delta h_{\rho\sigma}(y)}.$$

Regularity/typing requirements. To ensure the pairing is well-defined, F and G are taken to be *TT-typed, band-limited, microcausal* functionals on the realized weave configuration $\mathcal{I}^{[\mathscr{S}]}$: their derivatives commute with the weave projectors, have support/wavefront compatible with $G^{\mathrm{ret/adv}}_{[\mathrm{TT}]}$, and are BRST-clean so that the bracket descends to $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ (Propositions 7.3 and 10.9).

Typical examples. (i) *Smeared linear observable:* $F(h) = \int d\mu_{\bar{\Sigma}}\, f^{\mu\nu} h_{\mu\nu}$ with TT test tensor $f^{\mu\nu}$ (compact support on the band); then $\delta F/\delta h_{\mu\nu} = f^{\mu\nu}$. (ii) *Local polynomial:* $G(h) = \int d\mu_{\bar{\Sigma}}\, \frac{1}{2}\, h_{\mu\nu} Q^{\mu\nu\rho\sigma} h_{\rho\sigma}$, with Q a fixed TT-preserving kernel (e.g. a differential operator built from $\bar{\Sigma}$). (iii) *Curvature-linear observable:* $F(h) = \int d\mu_{\bar{\Sigma}}\, \alpha^{\mu\nu\rho\sigma}\, \delta R_{\mu\nu\rho\sigma}[h]$, with TT-compatible smearing α. All such choices satisfy the typed/triad commutation and band conditions used in the chapter.

Proposition 11.11 (Typed microcausality and cone support)**.** *Let $J^{\pm}_{\bar{\Sigma}}(x)$ be the causal future/-past with respect to the $\bar{\Sigma}$-cone. The TT causal kernel has support* $\operatorname{supp}\Delta^{[\mathrm{TT}]} \subset \{(x,y) \mid y \in J^{+}_{\bar{\Sigma}}(x) \cup J^{-}_{\bar{\Sigma}}(x)\}$. *Moreover, the IR window $K^{(\ell)}_{\mathrm{IR}}$ does not enlarge causal cones (cf. Definition 6.280); hence $\{F,G\}_{\mathrm{P}} = 0$ whenever the supports of F and G are $\bar{\Sigma}$-spacelike separated.*

Proof of Proposition 11.11. Work on a certified plateau $\mathfrak{P}$ with BRST–closed band tags $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell} = (P_x(\lambda_{\mathrm{UV}}), K^{(\ell)}_{\mathrm{IR}})$ that commute with the realized weave configuration operators (Definitions 6.280 and 9.10, Lemma 9.11, and Proposition 10.9). Let $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ be the background–covariant York–Lichnerowicz projector (eq. (10.4) and Definition 10.3), and denote by $K^{[\mathrm{TT}]}$ the TT kinetic operator obtained by projecting the linearized chronogeometric Einstein operator about the background $\bar{\Sigma}$ (Definition 11.106) onto the TT type (cf. Definition 11.3).

Step 1 (Hyperbolicity and cone support). By the hyperbolicity condition of the band–limited equations of motion ($\sigma_{\mathrm{pr}}(\bar{\Sigma}^{\mu\nu}\nabla_\mu\nabla_\nu)(k) = \bar{\Sigma}^{\mu\nu}k_\mu k_\nu > 0$ on timelike covectors; see eq. (7.15)), the retarded/advanced fundamental solutions $G_{R/A}$ of the untyped linearized operator have support contained in the $\bar{\Sigma}$–cones (future/past) ($\operatorname{supp} G_{R/A} \subset \{(x,y) \,|\, y \in J^{\pm}_{\bar{\Sigma}}(x)\}$). Because $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ is a local, $\bar{\Sigma}$–covariant projector that commutes with the realized weave configuration (Proposition 10.9), the TT Green operators $G^{[\mathrm{TT}]}_{R/A} := \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\, G_{R/A}\, \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ inherit the same support:

$$\operatorname{supp} G^{[\mathrm{TT}]}_{R/A} \;\subset\; \{(x,y) \mid y \in J^{\pm}_{\bar{\Sigma}}(x)\}.$$

Hence the TT Pauli–Jordan bi–solution $\Delta^{[\mathrm{TT}]} := G^{[\mathrm{TT}]}_R - G^{[\mathrm{TT}]}_A$ satisfies $\operatorname{supp}\Delta^{[\mathrm{TT}]} \subset \{(x,y) \mid y \in J^{+}_{\bar{\Sigma}}(x) \cup J^{-}_{\bar{\Sigma}}(x)\}$.

Step 2 (IR window and equal–time Peierls bracket). By construction of the IR window $K^{(\ell)}_{\mathrm{IR}}$ (Definition 6.280) and the cone–preservation statement for $\bar{\Sigma}$–covariant windowing (Remark 9.83), the band limitation does *not* enlarge the $\bar{\Sigma}$–cones. Therefore the (typed) Peierls bracket $\{F,G\}_{\mathrm{P}} = \langle \delta F/\delta\hat{h},\, \Delta^{[\mathrm{TT}]}\, \delta G/\delta\hat{h}\rangle$ (band version; cf. the inversion/propagator construction in eq. (7.10) and Proposition 7.16) vanishes whenever $\operatorname{supp} F$ and $\operatorname{supp} G$ are $\bar{\Sigma}$–spacelike separated, because $\Delta^{[\mathrm{TT}]}$ has no support across spacelike separations. This proves both the cone–support inclusion and the microcausality statement. □

Corollary 11.12 (TLO hook via Peierls)**.** *Let $S^{[\mathrm{TT}]}$ denote the TT-projected quadratic action on $\mathfrak{P}$. The TT law residual of Definition 11.6 can be written as a Peierls commutator with the action, $\mathbf{R}^{[\mathrm{TT}]} \sim \{\,\cdot\,, S^{[\mathrm{TT}]}\}_{\mathrm{P}}$, and vanishes identically when the TT Ward identity holds on $\mathfrak{P}$.*

Proof of Corollary 11.12 (with a meter functional). Let $S^{[\mathrm{TT}]}$ be the TT–projected quadratic action on $\mathfrak{P}$ (typed by $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ and band–limited; cf. Definition 11.3 and Proposition 10.9). Denote the TT Euler–Lagrange operator by $E^{[\mathrm{TT}]} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \delta S/\delta\hat{h}$ (typed/cohomological EL equations on the band; see Proposition 7.3 and eq. (7.10)). For linearized chronogeometric gravity,

$$E^{[\mathrm{TT}]}[\hat{h}] = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\left[\delta\bar{G}_{\mu\nu}[\hat{h}] + \Lambda\hat{h}_{\mu\nu} - \kappa\,\delta\bar{T}^{\mathrm{total}}_{\mu\nu}\right],$$

hence the TT law residual of Definition 11.6 is $\mathbf{R}^{[\mathrm{TT}]} = E^{[\mathrm{TT}]}[\hat{h}]$.

On a certified plateau, the Peierls bracket is generated by the TT Pauli–Jordan bi–solution $\Delta^{[\mathrm{TT}]}$ (Propositions 7.16 and 11.11). For quadratic actions the Peierls identity yields, for any functional $\mathcal{F}[\hat{h}]$,

$$\{\mathcal{F}, S^{[\mathrm{TT}]}\}_{\mathrm{P}} = \left\langle \frac{\delta\mathcal{F}}{\delta\hat{h}},\, \Delta^{[\mathrm{TT}]}\, E^{[\mathrm{TT}]} \right\rangle.$$

Choose the *meter functional* $\mathcal{M}^{[\mathrm{TT}]}$ by $\delta\mathcal{M}^{[\mathrm{TT}]}/\delta\hat{h} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ (a calibrated meter in the realized weave configuration sense; Definition 10.1 and Proposition 10.9). Then

$$\{\mathcal{M}^{[\mathrm{TT}]}, S^{[\mathrm{TT}]}\}_{\mathrm{P}} = \left\langle \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]},\, \Delta^{[\mathrm{TT}]}\, E^{[\mathrm{TT}]} \right\rangle \;\sim\; E^{[\mathrm{TT}]},$$

up to the fixed Green action, so $\mathbf{R}^{[\mathrm{TT}]} = E^{[\mathrm{TT}]} \sim \{\cdot, S^{[\mathrm{TT}]}\}_{\mathrm{P}}$, which proves the Peierls "hook".

If the TT Ward identity holds on $\mathfrak{P}$ (law satisfied at the typed level; cf. the law witness framework in Definitions 8.4 and 8.5 and its TT specialization Definition 11.6), then $E^{[\mathrm{TT}]} = 0$ and hence $\mathbf{R}^{[\mathrm{TT}]} = 0$. Gauge–parameter independence (Nielsen flatness) follows from the plateau BRST/Nielsen framework (Proposition 7.3 and Corollary 11.5). □

11.1.2 Covariant Quantization via the Peierls Bracket

To restore manifest covariance beyond the equal-time formalism of Section 11.1.1, we define the quantum algebra from the causal structure of the linearized equations, encoded in Green operators for the fluctuation dynamics (Definition 9.268 and Section 6.5). All constructions below are performed on a certified plateau $\mathfrak{P}$ (Chapter 8) with BRST-closed band tags $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell} = \left(P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}^{(\ell)}\right)$ commuting with the typed projector $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ (Definitions 6.280 and 9.10; typed projector from Chapter 11).

The Gaiamiti Green's Function and Causal Structure

Definition 11.13 (Gaiamiti Green's Function)**.** Let $\hat{\mathcal{D}}_\Phi$ denote the linearized fluctuation operator of the Gaiamiti sector derived from the second variation of S_Φ (Section 6.5). The (operator-valued) Green kernel $\hat{G}_{\Phi\Phi}(x,x')$ solves

$$\hat{\mathcal{D}}_\Phi(x)\,\hat{G}_{\Phi\Phi}(x,x') = \delta^{(d)}(x-x')\,\mathbb{I}\,. \tag{11.21}$$

For the spin–2 geometric carrier, we further define the *typed* Lichnerowicz-type wave operator

$$K_{[\mathrm{TT}]} \;:=\; \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\hat{\mathcal{D}}_{\Phi}\,\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,, \tag{11.22}$$

and its *TT-projected* advanced/retarded inverses $G_{[\mathrm{TT}]}^{\mathrm{adv/ret}}$ by

$$\boxed{K_{[\mathrm{TT}]}(x)\,G_{[\mathrm{TT}]}^{\mathrm{adv/ret}}(x,x') \;=\; \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,P_x(\lambda_{\mathrm{UV}})\,\delta^{(d)}(x-x')\,,} \tag{11.23}$$

with support conditions $G_{[\mathrm{TT}]}^{\mathrm{ret}}(x,x') = 0$ unless $x \in J_{\bar{\Sigma}}^{+}(x')$ and $G_{[\mathrm{TT}]}^{\mathrm{adv}}(x,x') = 0$ unless $x \in J_{\bar{\Sigma}}^{-}(x')$. The *TT causal kernel* (Pauli–Jordan function) is

$$\Delta^{[\mathrm{TT}]}(x,x') \;:=\; G_{[\mathrm{TT}]}^{\mathrm{ret}}(x,x') - G_{[\mathrm{TT}]}^{\mathrm{adv}}(x,x')\,. \tag{11.24}$$

The Covariant Commutator and the Peierls Bracket

For gauge-invariant functionals $F[h]$ and $G[h]$ of the typed graviton $h_{\mu\nu} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\delta\hat{g}_{\mu\nu}$ (Definition 11.3), the *typed Peierls bracket* is defined by the TT causal kernel:

$$\boxed{\begin{aligned}\{F,G\}_{\mathrm{P}} &= \Big\langle \frac{\delta F}{\delta h_{\mu\nu}},\; \Delta^{[\mathrm{TT}]\mu\nu}{}_{\rho\sigma}\left(\frac{\delta G}{\delta h_{\rho\sigma}}\right)\Big\rangle,\\ \Delta^{[\mathrm{TT}]} &:= G_{[\mathrm{TT}]}^{\mathrm{ret}} - G_{[\mathrm{TT}]}^{\mathrm{adv}}.\end{aligned}} \tag{11.25}$$

where functional derivatives are taken at fixed band tags (Definitions 6.280 and 9.10) on $\mathfrak{P}$. In particular, the *covariant commutator* of the typed field is

$$\boxed{\big[\,\hat{h}_{\mu\nu}(x),\;\hat{h}_{\rho\sigma}(x')\,\big] \;=\; i\;\Delta_{\mu\nu\,\rho\sigma}^{[\mathrm{TT}]}(x,x')\,.} \tag{11.26}$$

The equal-time limit of (11.25) on a Cauchy slice recovers the typed ETCR of Proposition 11.4; thus the canonical ETCRs appear as a restriction of the Peierls construction [82, 155].

Proposition 11.14 (Typed microcausality and cone support)**.** *With respect to the $\bar{\Sigma}$-cone, the support of the TT causal kernel is*

$$\operatorname{supp}\Delta^{[\mathrm{TT}]} \;\subset\; \{(x,x') \mid x' \in J_{\bar{\Sigma}}^{+}(x) \;\text{or}\; x' \in J_{\bar{\Sigma}}^{-}(x)\}.$$

Consequently, if the supports of F and G are $\bar{\Sigma}$-spacelike separated then $\{F,G\}_{\mathrm{P}} = 0$. Moreover, the IR window $K_{\mathrm{IR}}^{(\ell)}$ does not enlarge causal cones (Definition 6.280), so microcausality is preserved under the band-limit on $\mathfrak{P}$.

Proof. Work on a certified plateau $\mathfrak{P}$ with BRST–closed band tags $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell} = (P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}^{(\ell)})$ that commute with the realized weave configuration operators (Definitions 6.280 and 9.10, Lemma 9.11, and Proposition 10.9). Let $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ denote the background–covariant York–Lichnerowicz projector (eq. (10.4) and Definition 10.3). Denote by $K_{[\mathrm{TT}]}$ the TT–projected linearized chronogeometric Einstein operator about $\bar{\Sigma}$ (Definition 11.106).

Step 1: Cone support of retarded/advanced Green operators. The untyped linearized equations are (normally) hyperbolic with respect to the $\bar{\Sigma}$–cone, so their retarded and advanced Green operators $G_{R/A}$ obey $\operatorname{supp} G_{R/A} \subset \{(x,x') \,|\, x' \in J^{\pm}_{\bar{\Sigma}}(x)\}$ (standard hyperbolic support property; see the causal–response statement in Remark 9.83). Because $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ is a $\bar{\Sigma}$–covariant *local* projector that commutes with the realized weave configuration (Proposition 10.9), the TT Green operators

$$G^{[\mathrm{TT}]}_{R/A} := \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\, G_{R/A}\, \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$$

inherit the same support: $\operatorname{supp} G^{[\mathrm{TT}]}_{R/A} \subset \{(x,x') \,|\, x' \in J^{\pm}_{\bar{\Sigma}}(x)\}$. Hence the TT Pauli–Jordan bi–solution $\Delta^{[\mathrm{TT}]} := G^{[\mathrm{TT}]}_R - G^{[\mathrm{TT}]}_A$ satisfies

$$\operatorname{supp} \Delta^{[\mathrm{TT}]} \subset \{(x,x') \mid x' \in J^{+}_{\bar{\Sigma}}(x) \text{ or } x' \in J^{-}_{\bar{\Sigma}}(x)\}.$$

Step 2: Peierls bracket and microcausality. By definition (eq. (11.25)), the typed Peierls bracket of gauge–invariant functionals $F[h]$, $G[h]$ supported on $\mathfrak{P}$ is

$$\{F,G\}_{\mathrm{P}} = \Big\langle \frac{\delta F}{\delta h_{\mu\nu}},\ \Delta^{[\mathrm{TT}]\,\mu\nu}{}_{\rho\sigma}\ \frac{\delta G}{\delta h_{\rho\sigma}} \Big\rangle.$$

If $\operatorname{supp} F$ and $\operatorname{supp} G$ are $\bar{\Sigma}$–spacelike separated, the distribution $\Delta^{[\mathrm{TT}]}$ has no support connecting the two regions (by Step 1), hence the pairing vanishes and $\{F,G\}_{\mathrm{P}} = 0$. This is typed microcausality.

Step 3: Stability under the band IR window. The IR window $K^{(\ell)}_{\mathrm{IR}}$ is chosen $\bar{\Sigma}$–covariantly so as *not* to enlarge causal cones (Definition 6.280); the cone–preservation statement in Remark 9.83 ensures that inserting $K^{(\ell)}_{\mathrm{IR}}$ (and the commuting UV projector $P_x(\lambda_{\mathrm{UV}})$) into the realized weave configuration leaves the retarded/advanced support properties unchanged. Therefore the support inclusion for $\Delta^{[\mathrm{TT}]}$ and the vanishing of $\{F,G\}_{\mathrm{P}}$ at $\bar{\Sigma}$–spacelike separation persist under the band–limit on $\mathfrak{P}$.

The stated support inclusion and microcausality properties follow. □

Remark 11.15 (TLO hook for the TT law residual)**.** The TT *law residual* of Definition 11.6 can be written as a Peierls commutator with the TT action $S^{[\mathrm{TT}]}$ on $\mathfrak{P}$, schematically $\mathbf{R}^{[\mathrm{TT}]} \sim \{\cdot, S^{[\mathrm{TT}]}\}_{\mathrm{P}}$. It vanishes when the TT Ward identity holds, providing a covariant route to $\Lambda^{[\mathrm{TT}]}_{\mathrm{law}} \to 1$ in the exact law limit (Chapters 8–11).

Pauli–Jordan bi–solution (what it is and why it is vital).

Definition 11.16 (Typed Pauli–Jordan bi–solution)**.** Let $K_{[\mathrm{TT}]}$ be the TT–projected Lichnerowicz–type wave operator on $(\mathcal{X}_q, \bar{\Sigma})$ (cf. Definition 10.3 and eq. (10.4) and the linearized chronogeometric equations in Definition 11.106). Denote by $G^{\mathrm{ret/adv}}_{[\mathrm{TT}]}$ the retarded/advanced Green operators of $K_{[\mathrm{TT}]}$. The *typed Pauli–Jordan bi–solution* is the bi–distribution

$$\Delta^{[\mathrm{TT}]} \ := \ G^{\mathrm{ret}}_{[\mathrm{TT}]} - G^{\mathrm{adv}}_{[\mathrm{TT}]}, \tag{11.27}$$

characterized by the homogeneous equations $K^{(x)}_{[\mathrm{TT}]}\,\Delta^{[\mathrm{TT}]}(x,x') = 0 = K^{(x')}_{[\mathrm{TT}]}\,\Delta^{[\mathrm{TT}]}(x,x')$, antisymmetry $\Delta^{[\mathrm{TT}]}(x,x') = -\Delta^{[\mathrm{TT}]}(x',x)$, and causal support $\operatorname{supp}\Delta^{[\mathrm{TT}]} \subset \{(x,x') \mid x' \in J^{+}_{\bar{\Sigma}}(x) \text{ or } x' \in J^{-}_{\bar{\Sigma}}(x)\}$ (Propositions 11.11 and 11.14).

Why it is vital.

- *Covariant Poisson structure.* The Peierls bracket for gauge–invariant functionals is mediated by $\Delta^{[\mathrm{TT}]}$:
$$\{F, G\}_{\mathrm{P}} = \Big\langle \frac{\delta F}{\delta h_{\mu\nu}},\ \Delta^{[\mathrm{TT}]\,\mu\nu}{}_{\rho\sigma}\ \frac{\delta G}{\delta h_{\rho\sigma}} \Big\rangle,$$
giving a manifestly covariant replacement of equal–time brackets (eq. (11.25)). Its causal support guarantees $\{F, G\}_{\mathrm{P}} = 0$ for $\bar{\Sigma}$–spacelike separated supports (typed microcausality; Proposition 11.14).

- *Bridge to canonical ETCR.* On a Cauchy slice the Peierls bracket induced by $\Delta^{[\mathrm{TT}]}$ reproduces the typed equal–time commutator of the TT canonical variables (Proposition 11.4), so $\Delta^{[\mathrm{TT}]}$ is the covariant kernel that encodes the canonical algebra and its TT restriction ("project, then quantize"; Definition 11.3).

- *Causality and regulator stability.* Because $\Delta^{[\mathrm{TT}]}$ is the difference of retarded/advanced Green operators, its support is confined to $\bar{\Sigma}$–causal pairs (Proposition 11.14). The IR window $K_{\mathrm{IR}}^{(\ell)}$ does not enlarge cones (Definition 6.280 and Remark 9.83), so typed microcausality persists under band–limiting on the plateau $\mathfrak{P}$.

- *Law diagnostics (TLO hook).* The TT law residual is a Peierls commutator with the typed action, $\mathbf{R}^{[\mathrm{TT}]} \sim \{\cdot, S^{[\mathrm{TT}]}\}_{\mathrm{P}}$ (Corollary 11.12); thus $\Delta^{[\mathrm{TT}]}$ furnishes the kernel that links the linearized law (Ward identity) to observable diagnostics (Definition 11.6).

Interpretation and significance.

- *Manifest covariance (typed).* The covariant commutator in eq. (11.26) is expressed entirely through the TT causal kernel $\Delta^{[\mathrm{TT}]}$ built from the background geometry $\bar{\Sigma}_{\mu\nu}$ and its Relational Levi–Civita structure (Definition 6.7). It makes no reference to a preferred foliation beyond the $\bar{\Sigma}$-induced causal cone, and—by Proposition 11.14—vanishes for $\bar{\Sigma}$-spacelike separations, thereby enforcing microcausality. On a certified plateau with BRST-closed band tags (Definitions 6.280 and 9.10) the construction is Nielsen-flat and the IR window does not enlarge causal cones.

- *Relation to canonical quantization.* The typed Peierls bracket $\{\,\cdot\,,\,\cdot\,\}_{\mathrm{P}}$ in eq. (11.25) is the fundamental, covariant object. The equal-time canonical algebra for the typed variables is recovered by restricting eq. (11.25) to a Cauchy hypersurface and taking the equal-time limit, yielding the ETCR of Proposition 11.4; thus the ETCR formalism is a non-covariant specialization of the Peierls framework [82, 155].

- *Self-consistency within the IQR weave.* The same emergent chronogeometric structure $\bar{\Sigma}$ that governs causal propagation for the typed wave operator $K_{[\mathrm{TT}]}$ also determines the quantum algebra via $\Delta^{[\mathrm{TT}]}$. In particular, the TT law residual $\mathbf{R}^{[\mathrm{TT}]}$ of Definition 11.6 is evaluated against this causal structure and can be expressed through the Peierls construction; when the TT Ward identity holds on the plateau, the residual vanishes and the law visibility saturates. In this sense, the typed quantum algebra, causal propagation, and law diagnostics are mutually compatible and realized on the same certified plateau.

11.1.3 Graviton Mode Decomposition from Gaiamiti Fluctuations

We now isolate the physically propagating spin–2 degrees of freedom contained in the Gaiamiti fluctuation operator $\delta\hat{\mathbf{\Phi}}(x)$ [438, 502, 403, 114, 517]. In the IQR framework geometry is emergent rather than presupposed; hence the graviton is not a ripple *on* spacetime but a quantum excitation *of* the field whose expectation realizes the chronogeometric background $\bar{\Sigma}_{\mu\nu}$ (Chapter 5 and Section 5.2; see also [518, 425, 40, 1, 519]). Throughout this section we work on a certified plateau $\mathfrak{P}$ (Chapter 8) and use the typed (TT) projector $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ of Chapter 11 to define the graviton carrier $\hat{h}_{\mu\nu} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\delta\hat{g}_{\mu\nu}$ (Definition 11.3); the temporal/coherence fluctuation $\hat{\tau}_{\mu\nu} := \delta\hat{t}_{\mu\nu}$ is TT-null and does not contribute to $\hat{h}_{\mu\nu}$ (Chapter 11).

Background and Motivation

In classical GR, radiative gravitational degrees of freedom are massless, symmetric, transverse and traceless (TT) perturbations of the metric [7, 264, 502]. Our goal is to identify the *quantum* counterpart inside the Gaiamiti fluctuations on $(\mathcal{X}_q, \bar{\Sigma})$, respecting the typed/BRST structure (Chapters 8–11). This will yield a mode expansion for $\hat{h}_{\mu\nu}$ and the associated creation/annihilation operators, organized in a triad-resolved fashion $(\mathrm{corr}, \mathrm{vac}, \mathrm{spec})$, consistent with the sector projectors $\{\Pi^{(a)}\}$ commuting with $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ (Definitions 6.280 and 9.10).

Definition 11.17 (Gaiamiti fluctuation)**.** The dynamical Gaiamiti field (cf. Definition 6.2) splits as

$$\boxed{\hat{\mathbf{\Phi}}(x) = \bar{\Phi}(x)\,\mathbb{I}_x + \delta\hat{\mathbf{\Phi}}(x),} \tag{11.28}$$

with $\bar{\Phi}(x) := \langle\hat{\mathbf{\Phi}}(x)\rangle$ and $\langle 0|\delta\hat{\mathbf{\Phi}}(x)|0\rangle = 0$ (see eq. (10.87)). The quanta of $\delta\hat{\mathbf{\Phi}}$ are the *apeirons* (cf. Definition 9.246).

Spin–2 Polarization Basis (typed, background-covariant)

Definition 11.18 (Typed spin–2 (TT) polarization basis)**.** Fix the background $(\mathcal{X}_q, \bar{\Sigma})$ with $\bar{\Sigma}$-compatible Levi–Civita connection and TT projector $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ (York–Lichnerowicz form; see Definitions 6.7 and 10.3 and eq. (10.4)). For each triad sector $a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$ (Definition 10.2) and each $(k; x)$, a *typed, background–covariant spin–2 polarization basis* is a set $\{e_{\mu\nu}^{(a,\lambda)}(k;x)\}_{\lambda=1}^{d_g} \subset \mathrm{Im}\left(\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\Pi^{(a)}\right)$ satisfying:

$$e_{\mu\nu}^{(a,\lambda)}(k;x) = e_{\nu\mu}^{(a,\lambda)}(k;x) \qquad \text{(symmetry)}, \tag{11.29}$$

$$\bar{g}^{\mu\nu}(x)\, e_{\mu\nu}^{(a,\lambda)}(k;x) = 0 \qquad \text{(tracelessness)}, \tag{11.30}$$

$$k^{\mu} e_{\mu\nu}^{(a,\lambda)}(k;x) = 0 \qquad \text{(transversality)}, \tag{11.31}$$

$$\left\langle e^{(a,\lambda)}(k;x), e^{(a,\lambda')}(k;x)\right\rangle_{\bar{\Sigma}} = \delta_{\lambda\lambda'} \qquad \text{(orthonormality)}, \tag{11.32}$$

where indices are raised with $\bar{\Sigma}^{\mu\nu}$ (Definition 6.24) and $\langle A, B\rangle_{\bar{\Sigma}} := A_{\mu\nu}\, B^{\mu\nu}$ is the $\bar{\Sigma}$-induced inner product on symmetric, traceless rank–2 tensors at x (TT calculus; Definition 10.3). The polarization count is $d_g = \frac{d(d-3)}{2}$ (thus $d_g = 2$ for $d = 4$).

Completeness (typed TT subspace). The basis resolves the typed TT projector kernel:

$$\sum_{\lambda=1}^{d_g} e_{\mu\nu}^{(a,\lambda)}(k;x)\, e^{(a,\lambda)\,\rho\sigma}(k;x) = \left(\Pi_{\bar\Sigma}^{[\mathrm{TT}]}\,\Pi^{(a)}\right)_{\mu\nu}{}^{\rho\sigma}, \tag{11.33}$$

with $\Pi^{(a)}$ the triad projectors and $\Pi_{\bar\Sigma}^{[\mathrm{TT}]}\Pi^{(a)} = \Pi^{(a)}\Pi_{\bar\Sigma}^{[\mathrm{TT}]}$ (Definition 10.2 and Proposition 10.9). In geometric–optics regimes the label k is taken tangent to the $\bar\Sigma$-null cone and the resulting $(+,\times)$ content matches the E(2) template discussion (Remark 11.151).

Definition 11.19 (Triad-resolved TT mode expansion)**.** Decompose the typed graviton into sector components $\hat h_{\mu\nu}^{(a)} := \Pi^{(a)}\hat h_{\mu\nu}$ (with $[\Pi^{(a)}, \Pi_{\bar\Sigma}^{[\mathrm{TT}]}] = 0$) and expand each sector in TT modes:

$$\boxed{\hat h_{\mu\nu}(x) = \sum_{a\in\{\mathrm{corr,vac,spec}\}} \sum_{\lambda=1}^{d_g} \int \frac{d^{d-1}\mathbf{k}}{(2\pi)^{d-1}} \left[e_{\mu\nu}^{(a,\lambda)}(k;x)\; \hat a_{a,\lambda}(\mathbf{k})\, e^{-ik\cdot x} + e_{\mu\nu}^{(a,\lambda)}(k;x)^*\; \hat a_{a,\lambda}^\dagger(\mathbf{k})\, e^{+ik\cdot x}\right]}, \tag{11.34}$$

where $k^0 = \omega_a(\mathbf{k})$ solves the on-shell condition of the typed wave operator $K_{[\mathrm{TT}]}$ (Chapter 10) and the normalization of the measure follows the $\bar\Sigma$-compatible relativistic convention.

Explanation of Definition 11.19.

- *Triad resolution and TT typing.* The graviton operator $\hat h_{\mu\nu}$ is first typed by the background–covariant TT projector $\Pi_{\bar\Sigma}^{[\mathrm{TT}]}$ (York–Lichnerowicz form; eq. (10.4) and Definition 10.3), then decomposed by the triad projectors $\{\Pi^{(a)}\}_{a\in\{\mathrm{corr,vac,spec}\}}$ (Definition 10.2). The commutation $\Pi^{(a)}\Pi_{\bar\Sigma}^{[\mathrm{TT}]} = \Pi_{\bar\Sigma}^{[\mathrm{TT}]}\Pi^{(a)}$ is ensured by the realized weave configuration commutations on certified plateaux (Proposition 10.9), so $\hat h_{\mu\nu}^{(a)} := \Pi^{(a)}\hat h_{\mu\nu} \in \mathrm{Im}(\Pi_{\bar\Sigma}^{[\mathrm{TT}]}\Pi^{(a)})$.
- *Typed polarization basis.* For each triad sector a and pair $(k;x)$ there exists a complete $\bar\Sigma$–orthonormal TT basis $\{e_{\mu\nu}^{(a,\lambda)}(k;x)\}_{\lambda=1}^{d_g}$ obeying symmetry, tracelessness, transversality and orthonormality (Definition 11.18); completeness reads $\sum_\lambda e_{\mu\nu}^{(a,\lambda)} e^{(a,\lambda)\,\rho\sigma} = (\Pi_{\bar\Sigma}^{[\mathrm{TT}]}\Pi^{(a)})_{\mu\nu}{}^{\rho\sigma}$ (eq. (11.33)).
- *On–shell condition and phase.* The typed wave operator $K_{[\mathrm{TT}]}$ (TT–projected Lichnerowicz type; cf. Chapter X and Definition 10.3) determines the dispersion relation in each sector: $k^0 = \omega_a(\mathbf{k})$ solves the on–shell equation of $K_{[\mathrm{TT}]}$. The phase is $k\cdot x := k_\mu x^\mu$, with indices moved by $\bar\Sigma$ ($k_\mu = \bar\Sigma_{\mu\nu}k^\nu$; Definition 6.24).
- *Measure and normalization.* The integration measure $\int d^{d-1}\mathbf{k}/(2\pi)^{d-1}$ is the $\bar\Sigma$–compatible relativistic convention on the Cauchy slice induced by the foliation of $(\mathcal{X}_q, \bar\Sigma)$ (cf. Definition 6.7).

- *Creation/annihilation algebra (typed ETCR).* The operators $\hat{a}_{a,\lambda}(\mathbf{k})$, $\hat{a}^\dagger_{a,\lambda}(\mathbf{k})$ are defined so that eq. (11.34) reproduces the typed equal–time commutation relations of the TT canonical pair (Proposition 11.4). This fixes

$$\left[\hat{a}_{a,\lambda}(\mathbf{k}), \hat{a}^\dagger_{b,\lambda'}(\mathbf{k}')\right] = (2\omega_a(\mathbf{k}))\,(2\pi)^{d-1}\,\delta_{ab}\,\delta_{\lambda\lambda'}\,\delta^{(d-1)}(\mathbf{k}-\mathbf{k}')\,, \tag{11.35}$$

$$\left[\hat{a}_{a,\lambda}(\mathbf{k}), \hat{a}_{b,\lambda'}(\mathbf{k}')\right] = \left[\hat{a}^\dagger_{a,\lambda}(\mathbf{k}), \hat{a}^\dagger_{b,\lambda'}(\mathbf{k}')\right] = 0\,, \tag{11.36}$$

  with all $\bar{\Sigma}$–indices contracted using the polarization orthonormality (eq. (11.32)) and completeness (eq. (11.33)). The consistency with the Peierls bracket (eq. (11.25)) follows from the identification of the covariant Pauli–Jordan kernel $\Delta^{[\mathrm{TT}]}$ and its cone support (Proposition 11.14).

- *Causality, band stability, and gauge independence.* Microcausality is guaranteed by the cone support of $\Delta^{[\mathrm{TT}]}$ (Proposition 11.14) and is preserved under the band IR window $K^{(\ell)}_{\mathrm{IR}}$ (Definition 6.280 and Remark 9.83). On a certified plateau $\mathfrak{P}$, the realized weave configuration operators commute with the BRST charge, so gauge–parameter variations are Nielsen–flat (Proposition 7.3 and Corollary 11.5); thus the mode expansion and the algebra (11.35) are physical (cohomological) statements.

Proposition 11.20 (Orthogonality of sectors and Fock factorization)**.** *With creation/annihilation operators labeled by the triad and polarization, the only non-vanishing commutators are*

$$\boxed{\left[\hat{a}_{a,\lambda}(\mathbf{k}),\, \hat{a}^\dagger_{b,\lambda'}(\mathbf{k}')\right] = \delta_{ab}\,\delta_{\lambda\lambda'}\,(2\pi)^{d-1}\,2\omega_a(\mathbf{k})\,\delta^{(d-1)}(\mathbf{k}-\mathbf{k}')\,,} \tag{11.37}$$

with all others vanishing. Consequently the graviton Fock space factorizes across triads:

$$\mathscr{F}_\Phi \simeq \mathscr{F}_\Phi^{(\mathrm{corr})} \,\hat{\otimes}\, \mathscr{F}_\Phi^{(\mathrm{vac})} \,\hat{\otimes}\, \mathscr{F}_\Phi^{(\mathrm{spec})},$$

and the typed physical sector is obtained by the BRST cohomology $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ *(Chapter 8).*

Proof. Step 1 (projected mode extraction of the CCR). Use the typed TT mode expansion (11.34) with the typed polarization basis of Definition 11.18 (orthonormality (11.32), completeness (11.33)) and the typed ETCR on the band $\left[\hat{h}_{\mu\nu}(t,\mathbf{x}), \hat{\Pi}^{\rho\sigma}(t,\mathbf{y})\right]$ from Proposition 11.4. Project the ETCR onto a pair of modes by contracting with $e^{(a,\lambda)\,\mu\nu}(k;x)\,e^{-ik\cdot x}$ and $e^{(b,\lambda')}_{\rho\sigma}(k';y)^*\,e^{+ik'\cdot y}$ and integrating over the Cauchy slice. Using (11.32)–(11.33) and the $\bar{\Sigma}$-compatible measure (foliation fixed by Definition 6.7), the only nonzero contribution picks out equal triad labels and equal polarizations. Matching to the right-hand side of Proposition 11.4 (with the standard relativistic normalization of the measure) yields

$$\left[\hat{a}_{a,\lambda}(\mathbf{k}),\, \hat{a}^\dagger_{b,\lambda'}(\mathbf{k}')\right] = \delta_{ab}\,\delta_{\lambda\lambda'}\,(2\pi)^{d-1}\,2\omega_a(\mathbf{k})\,\delta^{(d-1)}(\mathbf{k}-\mathbf{k}'), \qquad \left[\hat{a},\hat{a}\right] = \left[\hat{a}^\dagger,\hat{a}^\dagger\right] = 0, \tag{11.38}$$

which coincides with (11.35). All cross-sector ($a \neq b$) and cross-polarization ($\lambda \neq \lambda'$) commutators vanish by the orthogonality of the typed basis and the commuting triad/TT projectors $\Pi^{(a)}\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}} = \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\Pi^{(a)}$ (Definition 10.2 and Proposition 10.9).

Step 2 (orthogonal decomposition of the one-particle space). Let $\mathcal{H}_1$ be the one-particle space obtained by completing finite-energy linear combinations of TT modes. By (11.32)–(11.33) and the orthogonality of the triad projectors (Definition 10.2), $\mathcal{H}_1$ decomposes as an orthogonal Hilbert direct sum

$$\mathcal{H}_1 \;=\; \mathcal{H}_1^{(\mathrm{corr})} \;\oplus\; \mathcal{H}_1^{(\mathrm{vac})} \;\oplus\; \mathcal{H}_1^{(\mathrm{spec})}, \qquad \mathcal{H}_1^{(a)} = \overline{\mathrm{span}}\Big\{ e_{\mu\nu}^{(a,\lambda)}(k;\cdot) \Big\}. \tag{11.39}$$

The commutation relations (11.38) imply that the creation/annihilation operators with different triad labels act on mutually orthogonal one-particle subspaces and generate commuting subalgebras; explicitly, $[\,\hat{a}_{a,\lambda}(\mathbf{k}), \hat{a}^\dagger_{b,\lambda'}(\mathbf{k}')\,] = 0$ for $a \neq b$, and the number operators $N^{(a)} = \int \frac{d^{d-1}\mathbf{k}}{(2\pi)^{d-1}\,2\omega_a(\mathbf{k})} \sum_\lambda \hat{a}^\dagger_{a,\lambda}(\mathbf{k})\hat{a}_{a,\lambda}(\mathbf{k})$ commute for different a.

Step 3 (Fock factorization). For a bosonic field, the symmetric Fock functor $\Gamma_{\mathrm{s}}(\cdot)$ maps an orthogonal direct sum into a (completed) tensor product: $\Gamma_{\mathrm{s}}(\mathcal{H}_1^{(1)} \oplus \mathcal{H}_1^{(2)} \oplus \mathcal{H}_1^{(3)}) \simeq \Gamma_{\mathrm{s}}(\mathcal{H}_1^{(1)}) \widehat{\otimes} \Gamma_{\mathrm{s}}(\mathcal{H}_1^{(2)}) \widehat{\otimes} \Gamma_{\mathrm{s}}(\mathcal{H}_1^{(3)})$. Applying this to (11.39) gives

$$\mathcal{F}_\Phi \;=\; \Gamma_{\mathrm{s}}(\mathcal{H}_1) \;\simeq\; \mathcal{F}_\Phi^{(\mathrm{corr})} \,\widehat{\otimes}\, \mathcal{F}_\Phi^{(\mathrm{vac})} \,\widehat{\otimes}\, \mathcal{F}_\Phi^{(\mathrm{spec})}, \tag{11.40}$$

with $\mathcal{F}_\Phi^{(a)} := \Gamma_{\mathrm{s}}(\mathcal{H}_1^{(a)})$. The identification is implemented by the commuting CCR blocks in (11.38) and the vacuum vector annihilated by all $\hat{a}_{a,\lambda}(\mathbf{k})$.

Step 4 (typed physical sector). On a certified plateau $\mathfrak{P}$, the realized weave configuration operators (TT projector, triad projectors, band tags) are BRST-closed and commute as in Proposition 10.9; hence the construction is gauge-parameter independent (Nielsen-flat) and descends to the physical Hilbert space by cohomology $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ (Proposition 7.3 and Corollary 11.5).

The stated CCR block-diagonality and the Fock factorization across triads follow. □

Definition 11.21 (Triadic residue vector from the TT two-point function)**.** Let $G^{[\mathrm{TT}]}$ be the typed graviton two-point function. In momentum space, each sector admits a Källén–Lehmann representation on $\mathcal{H}_{\mathrm{phys}}$,

$$\Delta^{(2),a}(q) = \frac{Z_a}{q^2 + i\epsilon} + \int_0^\infty \frac{\rho_a^{(2)}(\mu^2)\, d\mu^2}{q^2 - \mu^2 + i\epsilon}, \qquad Z_a \geq 0,\ \rho_a^{(2)}(\mu^2) \geq 0,$$

and we define the *triadic residue vector* $\vec{Z} := (Z_{\mathrm{corr}}, Z_{\mathrm{vac}}, Z_{\mathrm{spec}})$. This vector provides the sector-resolved pole strength used in the TT *state visibility* (Chapter 9) and is to be read on the certified plateau $\mathfrak{P}$.

Remark 11.22 (Physical interpretation and TLO hook)**.** Each $\hat{a}^\dagger_{a,\lambda}(\mathbf{k})$ creates a TT quantum of curvature with definite momentum and polarization in sector a. The sector weights are summarized by $\vec{Z}$ of Definition 11.21; together with the TT law residual $\mathbf{R}^{[\mathrm{TT}]}$ (Definition 11.6), they feed the typed Parasymmetric TLO resolver (Chapter 9), providing certified, sector-aware diagnostics for the radiative graviton within the IQR framework.

Support and Sectoral Projection

The full Gaiamiti fluctuation $\delta\hat{\mathbf{\Phi}}(x)$ (cf. Definition 11.17) contains spin-0, spin-1, and higher-rank excitations in addition to the radiative spin-2 content. To isolate the physically propagating spin-2 (graviton) degrees of freedom on the chronogeometric background $(\mathcal{X}_q, \bar{\Sigma}_{\mu\nu})$ with Relational Levi–Civita connection $\bar{\nabla}$ (Definition 6.7), we employ the typed background-covariant TT projector from Chapter 11 and the triad projectors from Chapters 8–10.

Definition 11.23 (Spin–2 (TT) projection operator)**.** Let $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ denote the background-covariant, self-adjoint, idempotent projector onto transverse–traceless rank-2 tensors with respect to $\bar{\Sigma}$ (Chapter 11). The *spin–2 projection* of the fluctuation is the covariant map

$$\boxed{\mathcal{P}^{(2)}[\delta\hat{\mathbf{\Phi}}](x) := \delta\hat{\mathbf{\Phi}}^{(2)}(x) := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{\mathbf{\Phi}}(x)\,,} \tag{11.41}$$

characterized by the TT constraints

$$\delta\hat{\mathbf{\Phi}}^{(2)}_{\mu\nu}(x) = \delta\hat{\mathbf{\Phi}}^{(2)}_{\nu\mu}(x) \qquad \text{(symmetry)}, \tag{11.42}$$

$$\bar{g}^{\mu\nu}(x)\,\delta\hat{\mathbf{\Phi}}^{(2)}_{\mu\nu}(x) = 0 \qquad \text{(tracelessness)}, \tag{11.43}$$

$$\bar{\nabla}^{\mu}\,\delta\hat{\mathbf{\Phi}}^{(2)}_{\mu\nu}(x) = 0 \qquad \text{(transversality)}, \tag{11.44}$$

where $\bar{g}$ and $\bar{\nabla}$ are those of $\bar{\Sigma}$. In particular, for the metric fluctuation $\delta\hat{g}_{\mu\nu}$ one obtains the typed graviton carrier $\hat{h}_{\mu\nu} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\delta\hat{g}_{\mu\nu}$ (Definition 11.3), which is the field expanded in Definition 11.19.

Definition 11.24 (Triad (sectoral) projectors and commuting structure)**.** Let $\Pi^{(a)}$ for $a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$ denote the sector projectors introduced in Chapters 8–10 (see Definitions 6.280 and 9.10). They satisfy orthogonality and completeness:

$$\Pi^{(a)}\Pi^{(b)} = \delta^{ab}\,\Pi^{(a)}, \qquad \sum_a \Pi^{(a)} = \mathbb{I}.$$

$$\boxed{\sum_a \Pi^{(a)} = \mathbf{1}, \qquad \Pi^{(a)}\Pi^{(b)} = \delta_{ab}\,\Pi^{(a)}\,.} \tag{11.45}$$

On the certified plateau $\mathfrak{P}$ they commute with the typed projector,

$$\boxed{\left[\Pi^{(a)},\, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\right] = 0\,,} \tag{11.46}$$

so the typed graviton decomposes sectorally as

$$\boxed{\hat{h}^{(a)}_{\mu\nu} := \Pi^{(a)}\,\hat{h}_{\mu\nu}\,, \qquad \hat{h}_{\mu\nu} = \sum_a \hat{h}^{(a)}_{\mu\nu}\,.} \tag{11.47}$$

Proposition 11.25 (Band support and no leakage between types)**.** *Fix the BRST-closed band tags* $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell} := \left(P_x(\lambda_{\mathrm{UV}}),\, K^{(\ell)}_{\mathrm{IR}}\right)$ *(Definitions 6.280 and 9.10). On the certified plateau* $\mathfrak{P}$*, they commute with the TT projector* $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ *and with each sector projector* $\Pi^{(a)}$*. Then the following hold:*

(i) (Typed support) *The realized TT field satisfies*

$$P_x(\lambda_{\rm UV})\, K_{\rm IR}^{(\ell)}\, \hat{h}_{\mu\nu} = \hat{h}_{\mu\nu} = \Pi_{\bar{\Sigma}}^{[\rm TT]}\, P_x(\lambda_{\rm UV})\, K_{\rm IR}^{(\ell)}\, \delta\hat{g}_{\mu\nu}\,,$$

so the TT content is entirely supported on $\mathfrak{B}_{\lambda_{\rm UV},\ell}$.

(ii) (Sectoral support) *Each component* $\hat{h}_{\mu\nu}^{(a)}$ *is likewise band-supported:*

$$P_x(\lambda_{\rm UV})\, K_{\rm IR}^{(\ell)}\, \hat{h}_{\mu\nu}^{(a)} = \hat{h}_{\mu\nu}^{(a)}\,,$$

and there is no leakage *between types:* $\Pi_{\bar{\Sigma}}^{\perp[\rm TT]}\hat{h}_{\mu\nu} = 0$ *and* $\Pi^{(b)}\hat{h}_{\mu\nu}^{(a)} = 0$ *for* $b \neq a$.

Sketch. Use commutativity of $P_x, K_{\rm IR}$ with $\Pi^{[\rm TT]}$ and $\Pi^{(a)}$ on $\mathfrak{P}$ and idempotency/orthogonality of the projectors. BRST-closure guarantees that any residual commutators are ζ-exact and trivial on $\mathcal{H}_{\rm phys} := H^0(\hat{Q}_{\rm BRST})$ (cf. Proposition 7.3). □

Corollary 11.26 (Compatibility with mode expansion and Fock factorization)**.** *The triad-resolved TT mode expansion of Definition 11.19 is well-defined with all modes supported on* $\mathfrak{B}_{\lambda_{\rm UV},\ell}$*, and the commutation relations of sector-labeled creation/annihilation operators yield the Fock factorization stated in Proposition 11.20. In particular, the spectrum and residues entering the triadic Källén–Lehmann representation are to be read on* $\mathfrak{P}$ *(cf. Definition 11.21).*

Remark 11.27 (Causality of support)**.** Because $K_{\rm IR}^{(\ell)}$ does not enlarge $\bar{\Sigma}$-cones (Definition 6.280) and the typed causal kernel has support on $J_{\bar{\Sigma}}^{\pm}$ (Proposition 11.14), the TT sector obeys microcausality and its band restriction preserves the causal support used in Section 11.1.2.

Quantum Graviton Field Operator

$$\boxed{\hat{h}_{\mu\nu} \;:=\; \Pi_{\bar{\Sigma}}^{[\rm TT]}\,\delta\hat{g}_{\mu\nu} \quad\Rightarrow\quad \hat{h}_{\mu\nu} = \hat{h}_{(\mu\nu)},\;\; \bar{g}^{\mu\nu}\hat{h}_{\mu\nu} = 0,\;\; \bar{\nabla}^{\mu}\hat{h}_{\mu\nu} = 0,} \tag{11.48}$$

with $\Pi_{\bar{\Sigma}}^{[\rm TT]}$ the background–covariant TT projector of Chapter 11 (cf. Definition 11.3). Moreover, the temporal/coherence fluctuation $\hat{\tau}_{\mu\nu} := \delta\hat{t}_{\mu\nu}$ is TT-null and hence cannot contribute to $\hat{h}_{\mu\nu}$ (see Definition 11.3).

Definition 11.28 (Quantum Graviton Field Operator)**.** Let $\delta\hat{\Phi}(x)$ be the operator-valued fluctuation of the Gaiamiti field on $(\mathcal{X}_q, \bar{\Sigma})$ (Definition 11.17). Let $\mathcal{P}^{(2)}$ be the covariant spin–2 projection operator (Definition 11.23), and let $\mathbb{E}^{\dagger}_{\mu\nu}$ be the dual geometric embedding that extracts a symmetric TT tensor from the internal operator algebra (inverse to $\mathbb{E}^{\mu\nu}$ in Definition 11.19). We *define* the quantum graviton field operator by

$$\boxed{\hat{h}_{\mu\nu}(x) \;:=\; \mathcal{P}^{(2)}\big[\delta\hat{\Phi}\big](x)\;\cdot\;\mathbb{E}^{\dagger}_{\mu\nu},} \tag{11.49}$$

which, by the typed clause above, coincides with the background–covariant TT projection $\Pi_{\bar{\Sigma}}^{[\rm TT]}\delta\hat{g}_{\mu\nu}$ and therefore satisfies symmetry, tracelessness, and transversality with respect to $\bar{\Sigma}$.

Interpretation and Structure.

- $\delta\hat{\mathbf{\Phi}}(x) \in \Gamma\Big[\mathrm{End}\Big(T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}\Big)\Big]$: the total Gaiamiti fluctuation over $\mathcal{X}_q$ (cf. Definition 11.17).
- $\mathcal{P}^{(2)}[\delta\hat{\mathbf{\Phi}}](x)$: the covariant TT (spin–2) projection implementing symmetry, tracelessness, and transversality relative to $\bar{\Sigma}_{\mu\nu}$ (Definition 11.23).
- $\mathbb{E}^{\dagger}_{\mu\nu}$: $\mathrm{End}\Big(T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}}\Big) \to \mathrm{TT}^2(T^*_x\mathcal{X}_q)$: the dual embedding that recovers a rank–2 TT tensor, inverse to the embedding used in the mode expansion (Definition 11.19).
- $\hat{h}_{\mu\nu}(x)$: a typed, covariant $(0,2)$ operator field acting on the total Fock space; it is precisely the carrier expanded in eq. (11.34) and used in the propagator construction of Section 11.2.

Physical Role.

- *Canonical identification.* $\hat{h}_{\mu\nu}$ gives the canonical, typed identification of the quantum gravitational field without reference to a fixed metric background (it is defined via $\bar{\Sigma}$ and $\Pi^{[\mathrm{TT}]}$).
- *Kinematics → dynamics.* It sources the triad-resolved mode expansion and creation/annihilation algebra (eq. (11.34)), and it underlies the definition of the TT propagator (Section 11.2).
- *Linearized field equation.* $\hat{h}_{\mu\nu}$ satisfies the typed, linearized equation obtained from the matter–geometry kernel $\mathcal{U}_{\Phi\Theta}(x,x')$ (see eqs. (10.183) and (10.184)).
- *Quantum curvature.* It represents the spin–2 quantum excitation of the relational vacuum manifold and therefore encodes genuine quantum curvature fluctuations in the IQR framework.

Masslessness of the Graviton from Spontaneous Symmetry Breaking

Explanation of Proposition 11.29. The typed graviton $\hat{h}_{\mu\nu} := \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{g}_{\mu\nu}$ is the background–covariant, transverse–traceless carrier obtained by projecting the metric fluctuation with the York–Lichnerowicz projector (eq. (10.4) and Definition 10.3) as formalized in Definition 11.3. The setting is a certified plateau $\mathfrak{P}$ (Chapter VIII), where the band tags $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ commute with all realized weave configuration operators and are BRST–closed (Definitions 6.280 and 9.10 and Propositions 7.3 and 10.9). Assume the Gaiamiti vacuum manifold $\mathcal{M}_{\mathrm{vac}}$ admits a continuous symmetry group G whose action on the field $\hat{\mathbf{\Phi}}$ is spontaneously broken by the realized background $\bar{\Phi}$ in the split $\hat{\mathbf{\Phi}} = \bar{\Phi}\,\mathbb{I} + \delta\hat{\mathbf{\Phi}}$ (Definitions 6.61 and 11.17 and eq. (11.28)). Then the broken generators define flat directions (Goldstone directions) in the fluctuation space. The *typed* (TT) content of those directions is obtained by applying $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$: if the Goldstone direction projects nontrivially onto the TT carrier, the linearized tensorial mass operator $\mathcal{M}^{\mu\nu}{}_{\alpha\beta}$ (defined via the second variation and projected onto the rank–2 sector; Proposition 11.32) annihilates $\hat{h}$, $\mathcal{M}^{\mu\nu}{}_{\alpha\beta}\hat{h}^{\alpha\beta} = 0$. Equivalently, the typed

equations reduce to a massless wave law $\bar{\nabla}^\lambda \bar{\nabla}_\lambda \hat{h}_{\mu\nu} = \hat{J}_{\mu\nu}$, consistent with TT–projected dynamics and microcausality on $(\mathcal{X}_q, \bar{\Sigma})$ (Definition 6.7 and Proposition 11.14).

The two–point function of the typed graviton on the physical Hilbert space $\mathcal{H}_{\rm phys} = H^0(\hat{Q}_{\rm BRST})$ (plateau projection; Proposition 7.3) admits a triad–resolved Källén–Lehmann representation with nonnegative spectral weights (Definitions 10.2, 11.18 and 11.21). In the presence of a TT Goldstone direction, the TT spectral measure necessarily contains a pole at $q^2 = 0$ with residue $Z_a \geq 0$ in each triad sector a, so the total residue $Z_{\rm TT} = \sum_a Z_a > 0$ on $\mathfrak{P}$. Positivity follows from the cohomological (BRST–projected) positivity together with the orthonormality/completeness of the typed polarization basis and the canonical TT algebra (eqs. (11.32) and (11.33) and Proposition 11.4); regulator stability is ensured by Nielsen flatness of the realized weave configuration (Corollary 11.5) and cone–preserving band limitation (Definition 6.280 and Remark 9.83). Thus, spontaneous breaking of G in the Gaiamiti sector enforces a *massless* TT carrier with a positive–residue Goldstone pole in the typed graviton spectrum, exactly as stated in items *(i)–(ii)* of Proposition 11.29.

Proposition 11.29 (Masslessness of the graviton from typed SSB)**.** *Let $\hat{h}_{\mu\nu} := \Pi^{[\rm TT]}_{\bar{\Sigma}}\, \delta\hat{g}_{\mu\nu}$ be the typed (background–covariant TT) graviton carrier of Definitions 11.3 and 11.28 and eq. (11.49), defined on a certified plateau $\mathfrak{P}$ (Chapter 8) with BRST-closed band tags commuting with $\Pi^{[\rm TT]}_{\bar{\Sigma}}$ and the sector projectors (Definitions 6.280 and 9.10). Suppose the Gaiamiti vacuum manifold $\mathcal{M}_{\rm vac}$ (Definition 6.61) admits a continuous symmetry group G whose action on $\hat{\mathbf{\Phi}}$ is spontaneously broken by the realization $\bar{\Phi}$ appearing in the split $\hat{\mathbf{\Phi}} = \bar{\Phi}\,\mathbb{I} + \delta\hat{\mathbf{\Phi}}$ (Definition 11.17 and eq. (11.28)). Then, in the* typed *(TT) sector:*

*(i) (*Goldstone in TT type*) The broken generators produce a Goldstone direction that projects non-trivially onto the TT carrier; equivalently, the linearized tensorial mass operator $\mathcal{M}^{\mu\nu}{}_{\alpha\beta}$ of Proposition 11.32 annihilates $\hat{h}$,*

$$\boxed{\mathcal{M}^{\mu\nu}{}_{\alpha\beta}(x)\, \hat{h}^{\alpha\beta}(x) = 0\,,} \tag{11.50}$$

so the typed field obeys a massless wave equation (up to sources) $\bar{\nabla}^\lambda \bar{\nabla}_\lambda \hat{h}_{\mu\nu} = \hat{J}_{\mu\nu}$.

*(ii) (*Massless pole with positive residue*) The TT two-point function on $\mathcal{H}_{\rm phys} = H^0(\hat{Q}_{\rm BRST})$ admits a triad-resolved Källén–Lehmann representation (Definition 11.21) with a massless pole,*

$$\Delta^{(2),a}(q) = \frac{Z_a}{q^2 + i\epsilon} + \int_0^\infty \frac{\rho_a^{(2)}(\mu^2)\, d\mu^2}{q^2 - \mu^2 + i\epsilon}, \qquad Z_a \geq 0\,.$$

Writing $Z_{\rm TT} := \sum_{a\in\{\rm corr,vac,spec\}} Z_a$, one has

$$\boxed{Z_{\rm TT} > 0 \quad \text{on } \mathfrak{P}\,,} \tag{11.51}$$

so the TT spectral measure contains a Goldstone pole at $q^2 = 0$ with positive residue.

Proof of Proposition 11.29. Preliminaries. Work on a certified plateau $\mathfrak{P}$ where the realized weave configuration operators (TT projector, triad projectors, band tags) commute and are

BRST–closed (Propositions 7.3 and 10.9 and Definitions 6.280 and 9.10). The typed Peierls bracket is

$$\{F,G\}_{\mathrm{P}} = \Big\langle \frac{\delta F}{\delta h_{\mu\nu}},\ \Delta^{[\mathrm{TT}]\,\mu\nu}{}_{\rho\sigma}\,\frac{\delta G}{\delta h_{\rho\sigma}} \Big\rangle, \qquad \Delta^{[\mathrm{TT}]} := G^{\mathrm{ret}}_{[\mathrm{TT}]} - G^{\mathrm{adv}}_{[\mathrm{TT}]}, \tag{11.52}$$

(eq. (11.25)), with cone support and microcausality as in Proposition 11.14. The TT carrier is $\hat{h}_{\mu\nu} := \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{g}_{\mu\nu}$ (Definition 11.3).

Step 1 (SSB and TT projection). The background–fluctuation split $\hat{\mathbf{\Phi}} = \bar{\Phi}\,\mathbb{I} + \delta\hat{\mathbf{\Phi}}$ (Definition 11.17 and eq. (11.28)) realizes SSB by $\bar{\Phi} \in \mathcal{M}_{\mathrm{vac}}$ (Definition 6.61), selecting a coset G/H. Let Q^A be a broken generator. Its action on fields is generated by the Peierls bracket at fixed band tags (Definitions 6.280 and 9.10): for any gauge–invariant F, $\delta_{Q^A}F = \{F, Q^A\}_{\mathrm{P}}$. Applying the York–Lichnerowicz projector (eq. (10.4) and Definition 10.3) and using realized weave configuration commutations (Proposition 10.9) gives a typed variation $\delta_{Q^A}\hat{h} = \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta_{Q^A}(\delta\hat{g})$, with $\hat{\tau}_{\mu\nu}$ TT–null (TT calculus, Chapter XI).

Step 2 (Typed Ward identity and soft matrix element). The TT law residual is the Peierls commutator with the typed action ($\mathbf{R}^{[\mathrm{TT}]} \sim \{\cdot, S^{[\mathrm{TT}]}\}_{\mathrm{P}}$; Corollary 11.12); invariance under the unbroken subgroup and SSB for the broken directions yield the *typed Ward identity* on $\mathfrak{P}$: for the broken Q^A,

$$\{\,\hat{h},\ Q^A\,\}_{\mathrm{P}} \neq 0 \quad \text{and} \quad \{\,S^{[\mathrm{TT}]},\ Q^A\,\}_{\mathrm{P}} = 0, \tag{11.53}$$

the latter expressing stationarity of the action along the symmetry direction. Fourier transforming the two–point function and inserting (11.53) shows that the soft matrix element of the broken charge between the vacuum and a one–TT–quantum state saturates the *soft limit* of the typed two–point distribution, producing a $1/q^2$ singularity in the TT channel (see also Proposition 11.4 for the canonical normalization and Proposition 11.14 for the covariant kernel).

Step 3 (Vanishing of the typed tensorial mass operator). Let $\mathcal{M}^{\mu\nu}{}_{\alpha\beta}$ denote the linearized tensorial mass operator obtained from the second variation of the action and projected onto symmetric rank–2 tensors (Proposition 11.32). Along a broken direction, the Hessian has a null vector (Goldstone flat direction). Projecting this direction with $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ and using $[\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}},\ \cdot\,] = 0$ on $\mathfrak{P}$ (Proposition 10.9) yields

$$\mathcal{M}^{\mu\nu}{}_{\alpha\beta}\,\hat{h}^{\alpha\beta} = 0, \tag{11.54}$$

which is item (i). Consequently, up to sources, the typed equation of motion is the massless wave law $\bar{\nabla}^{\lambda}\bar{\nabla}_{\lambda}\hat{h}_{\mu\nu} = \hat{J}_{\mu\nu}$ (Definition 6.7).

Step 4 (Källén–Lehmann representation and positivity). On the physical Hilbert space $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ (Proposition 7.3) the typed two–point function admits a triad–resolved Källén–Lehmann representation with nonnegative spectral weights (Definitions 10.2, 11.18 and 11.21):

$$\Delta^{(2),a}(q) = \frac{Z_a}{q^2 + i\epsilon} + \int_0^\infty \frac{\rho_a^{(2)}(\mu^2)\,d\mu^2}{q^2 - \mu^2 + i\epsilon}, \qquad Z_a \geq 0.$$

Using the orthonormality and completeness of TT polarizations (eqs. (11.32) and (11.33)) and the block–diagonal CCR (Propositions 11.4 and 11.20) shows that the residue is the modulus squared of the TT order–parameter matrix element associated with the broken charge, hence strictly positive if the TT projection of the Goldstone direction is nonzero. Summing over triads gives $Z_{\mathrm{TT}} = \sum_a Z_a > 0$, which is item (ii).

Step 5 (Stability and covariance). Nielsen flatness (Corollary 11.5) ensures gauge–parameter independence on $\mathfrak{P}$, and cone–preserving band limitation (Definition 6.280 and Remark 9.83) preserves the support and the Peierls construction (Proposition 11.14). Thus both (i) and (ii) are regulator–safe and covariant.

This completes the proof of Proposition 11.29. □

Corollary 11.30 (GMOR-style typed mass relation under explicit breaking)**.** *Introduce a small explicit breaking ϵ of the G-symmetry in the Gaiamiti potential (or source sector). On $\mathfrak{P}$ there exists a plateau constant $C_{\mathfrak{P}} > 0$ such that the TT pole shifts to*

$$\boxed{m^2_{\mathrm{TT}} \;=\; \mathcal{O}(\epsilon), \qquad 0 \;\le\; m^2_{\mathrm{TT}} \;\le\; C_{\mathfrak{P}}\,\epsilon\,,} \tag{11.55}$$

and the two-point becomes $\Delta^{(2)}_{\mathrm{TT}}(q) \simeq \dfrac{Z_{\mathrm{TT}}}{q^2 - m^2_{\mathrm{TT}} + i\epsilon}$ *+ regular. Plateau/Nielsen safety (Chapter 8) ensures the bound is gauge-parameter independent.*

Proof. Setup and notation. Work on a certified plateau $\mathfrak{P}$ where the realized weave configuration operators (TT projector, triad projectors, band tags) commute and are BRST–closed (Propositions 7.3 and 10.9 and Definitions 6.280 and 9.10). Introduce a small explicit breaking ϵ of the G-symmetry in the (typed) action,

$$S^{[\mathrm{TT}]}_\epsilon := S^{[\mathrm{TT}]} \;+\; \epsilon\, B^{[\mathrm{TT}]}, \qquad B^{[\mathrm{TT}]} := \Pi^{[\mathrm{TT}]}_{\bar\Sigma}\, B\, \Pi^{[\mathrm{TT}]}_{\bar\Sigma}, \tag{11.56}$$

with B a $\bar\Sigma$-covariant functional (potential or source) calibrated on $\mathfrak{P}$. The typed Euler–Lagrange map is then

$$E^{[\mathrm{TT}]}_\epsilon[\hat h] := \Pi^{[\mathrm{TT}]}_{\bar\Sigma}\, \frac{\delta S^{[\mathrm{TT}]}_\epsilon}{\delta \hat h} = E^{[\mathrm{TT}]}[\hat h] \;+\; \epsilon\, B^{[\mathrm{TT}]}_{,h}[\hat h], \tag{11.57}$$

where $E^{[\mathrm{TT}]}[\hat h] = 0$ are the unbroken typed equations (massless carrier; Proposition 11.29). By the Peierls "hook" (Corollary 11.12) and the typed bracket definition (eq. (11.25)), $\mathbf{R}^{[\mathrm{TT}]}_\epsilon \sim \{\cdot, S^{[\mathrm{TT}]}_\epsilon\}_{\mathrm{P}}$ and thus $E^{[\mathrm{TT}]}_\epsilon = 0$ encodes the linear response and the pole structure of the TT two–point.

Step 1 (modified Ward identity and soft saturation). Let Q^A be a generator broken spontaneously at $\epsilon = 0$ (Definitions 6.61 and 11.17) and explicitly by $\epsilon \neq 0$. On $\mathfrak{P}$, the typed Ward identity is deformed by

$$\{\, S^{[\mathrm{TT}]}_\epsilon,\, Q^A \,\}_{\mathrm{P}} = \epsilon\, \{\, B^{[\mathrm{TT}]},\, Q^A \,\}_{\mathrm{P}}, \qquad \{\cdot,\cdot\}_{\mathrm{P}} \text{ as in } eq.\ (11.25). \tag{11.58}$$

Fourier transforming the TT two–point and inserting (11.58) shows that the massless pole at $q^2 = 0$ (present at $\epsilon = 0$; Proposition 11.29) shifts to $q^2 = m^2_{\mathrm{TT}}$ of order ϵ, controlled by

the explicit–breaking insertion $B^{[\mathrm{TT}]}$. This is the typed, covariant analogue of the GMOR mechanism.

Step 2 (first–order pole shift and bound). Linearizing (11.57) around the unbroken vacuum yields, in momentum space,

$$K_{[\mathrm{TT}]}(q)\,\tilde{h}(q) + \epsilon\,\Sigma_{[\mathrm{TT}]}(q)\,\tilde{h}(q) = 0,$$

with $K_{[\mathrm{TT}]}$ the typed wave operator (York–Lichnerowicz type; Chapter X) and $\Sigma_{[\mathrm{TT}]} \equiv (B^{[\mathrm{TT}]}_{,h})_{,h}$ the typed explicit–breaking Hessian. At $\epsilon = 0$, $K_{[\mathrm{TT}]}(q)$ has a zero eigenvalue at $q^2 = 0$ in the Goldstone TT direction (Proposition 11.29); let ψ denote a normalized TT polarization vector in that kernel (constructed from Definition 11.18). Rayleigh–Schrödinger perturbation theory (or Feshbach reduction on the TT subspace) gives the pole shift

$$m^2_{\mathrm{TT}} = \epsilon\,\frac{\langle \psi,\ \Sigma_{[\mathrm{TT}]}(0)\,\psi\rangle_{\bar{\Sigma}}}{\langle \psi,\ (\partial K_{[\mathrm{TT}]}/\partial q^2)_{q^2=0}\,\psi\rangle_{\bar{\Sigma}}} \ +\ \mathcal{O}(\epsilon^2). \tag{11.59}$$

The denominator equals the positive TT residue Z_{TT} of the Källén–Lehmann representation (Definition 11.21 and Proposition 11.29), while the numerator is bounded by the instrumentation norm on $\mathfrak{P}$ (Definition 10.1): there exists $C'_{\mathfrak{P}} > 0$ such that $\langle \psi,\ \Sigma_{[\mathrm{TT}]}\,\psi\rangle_{\bar{\Sigma}} \le C'_{\mathfrak{P}}$. Hence, for ϵ sufficiently small,

$$0 \ \le\ m^2_{\mathrm{TT}} \ \le\ \frac{C'_{\mathfrak{P}}}{Z_{\mathrm{TT}}}\,\epsilon \ =:\ C_{\mathfrak{P}}\,\epsilon, \qquad m^2_{\mathrm{TT}} = \mathcal{O}(\epsilon). \tag{11.60}$$

Positivity $m^2_{\mathrm{TT}} \ge 0$ follows from the spectral positivity of the TT two–point on $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ (Proposition 7.3 and Definition 11.21) and from $Z_{\mathrm{TT}} > 0$ (Proposition 11.29).

Step 3 (two–point structure and covariance). Inserting (11.60) into the typed Källén–Lehmann form yields

$$\Delta^{(2)}_{\mathrm{TT}}(q) \simeq \frac{Z_{\mathrm{TT}}}{q^2 - m^2_{\mathrm{TT}} + i\epsilon} + \text{regular},$$

with microcausality preserved by cone support of $\Delta^{[\mathrm{TT}]}$ (Proposition 11.14) and stability under the band IR window (Definition 6.280 and Remark 9.83). Finally, gauge–parameter independence of m^2_{TT} and of the bound $C_{\mathfrak{P}}$ on $\mathfrak{P}$ follows from Nielsen flatness (Corollary 11.5) and BRST descent to $\mathcal{H}_{\mathrm{phys}}$ (Proposition 7.3).

Equations (11.60) establish the claimed GMOR–style typed mass relation. □

Remark 11.31 (Law visibility in the SSB limit)**.** Recall the TT law residual and visibility from Definition 11.6: $\mathbf{R}^{[\mathrm{TT}]} := \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\big[\delta\bar{G}_{\mu\nu}[\hat{h}] + \Lambda\,\hat{h}_{\mu\nu} - \kappa\,\delta\bar{T}^{\mathrm{total}}_{\mu\nu}\big]$ and $\Lambda^{[\mathrm{TT}]}_{\mathrm{law}} := 1 - \|\mathbf{R}^{[\mathrm{TT}]}\|_{\mathcal{I}}/\|\cdot\|_{\mathrm{ref}} \in [0,1]$, with instrumentation norm $\|\cdot\|_{\mathcal{I}}$ and reference scale $\|\cdot\|_{\mathrm{ref}}$ calibrated on the certified plateau $\mathfrak{P}$ (Definition 10.1 and Section 7.3.9). On $\mathfrak{P}$, the Peierls "hook" identifies the residual with the typed equations of motion, $\mathbf{R}^{[\mathrm{TT}]} \sim \{\cdot, S^{[\mathrm{TT}]}\}_{\mathrm{P}}$ (Corollary 11.12 and eq. (11.25)), and cone preservation of the band realized weave configuration ensures regulator safety (Definition 6.280 and Remark 9.83).

In the exact SSB limit ($\epsilon \to 0$), the typed Ward identity holds and the TT Goldstone direction implies vanishing typed mass operator (Proposition 11.29); therefore the typed equations of motion are satisfied and $\mathbf{R}^{[\mathrm{TT}]} = 0$, which yields

$$\boxed{\Lambda_{\mathrm{law}}^{[\mathrm{TT}]} \longrightarrow 1.} \tag{11.61}$$

For small explicit breaking ϵ, the GMOR–style estimate gives $m_{\mathrm{TT}}^2 = \mathcal{O}(\epsilon)$ with $0 \le m_{\mathrm{TT}}^2 \le C_{\mathfrak{P}}\epsilon$ (Corollary 11.30). Linear response of the typed EL map on $\mathfrak{P}$ then bounds the residual by the same plateau constants, $\|\mathbf{R}^{[\mathrm{TT}]}\|_{\mathcal{I}} \le C'_{\mathfrak{P}}\, m_{\mathrm{TT}}^2 = \mathcal{O}(\epsilon)$, so $\Lambda_{\mathrm{law}}^{[\mathrm{TT}]} = 1 - \mathcal{O}(\epsilon)$. Gauge–parameter independence of these statements follows from BRST/Nielsen safety (Proposition 7.3 and Corollary 11.5), and covariance from the typed Peierls construction with causal support (Proposition 11.14).

The Tensorial Graviton Mass Operator

Proposition 11.32 (The Tensorial Graviton Mass Operator)**.** *Let the emergent chronogeometric tensor be the symmetric functional $\hat{\Sigma}_{\mu\nu}(x) = \mathcal{S}_{\mu\nu}[\hat{\mathbf{\Phi}}(x)]$, and let the linear response of Σ to Gaiamiti fluctuations at the background $\bar{\Phi}$ be $\delta\hat{\Sigma}_{\mu\nu}(x) = \sum_i \mathcal{E}^i_{\mu\nu}(x)\,\delta\hat{\Phi}^i(x)$ with $\mathcal{E}^i_{\mu\nu}(x) := \frac{\delta\mathcal{S}_{\mu\nu}}{\delta\hat{\Phi}^i(x)}\Big|_{\bar{\Phi}}$. Write the quadratic mass term in the Gaiamiti basis as*

$$S^{(2)}_{mass} = \frac{1}{2}\int d^dx\,\sqrt{-\det\bar{\Sigma}}\,\sum_{i,j}\left(\delta\hat{\Phi}^i\right)^\dagger \mathcal{M}_{ij}\,\delta\hat{\Phi}^j, \qquad \mathcal{M}_{ij} := \frac{\delta^2\mathcal{V}_\Phi}{\delta(\hat{\Phi}^i)^\dagger\delta\hat{\Phi}^j}\Big|_{\bar{\Phi}}.$$

If $\mathcal{E}^{-1}$ denotes the inverse kernel expressing $\delta\hat{\Phi}$ in terms of $\delta\hat{\Sigma}$, then the tensorial *graviton mass operator appearing in the geometric quadratic form*

$$S^{(2)}_{mass} = \frac{1}{2}\int d^dx\,\sqrt{-\det\bar{\Sigma}}\,\left(\delta\hat{\Sigma}_{\mu\nu}\right)^\dagger \mathcal{M}^{\mu\nu}{}_{\alpha\beta}\,\delta\hat{\Sigma}^{\alpha\beta}$$

is given by the congruence of $\mathcal{M}_{ij}$ through the response map:

$$\boxed{\mathcal{M}^{\mu\nu}{}_{\alpha\beta}(x) := \sum_{i,j}\Big((\mathcal{E}^{-1})^i_{\rho\sigma}(x)\Big)^\dagger \bar{\Sigma}^{\rho\mu}(x)\,\bar{\Sigma}^{\sigma\nu}(x)\,\mathcal{M}_{ij}(x)\,\Big((\mathcal{E}^{-1})^j_{\alpha\beta}(x)\Big).} \tag{11.62}$$

This congruence inherits the null directions of $\mathcal{M}_{ij}$, so Goldstone directions in the Gaiamiti basis remain null in the geometric tensor basis, consistently with Proposition 11.29.

Proof (typed/TT refinement). **Step 1: Mass term in the Gaiamiti basis.** As in the original argument, expand the potential to quadratic order to obtain

$$S^{(2)}_{\mathrm{mass}} = \frac{1}{2}\int d^dx\,\sqrt{-\det\bar{\Sigma}}\,\sum_{i,j}(\delta\hat{\Phi}^i)^\dagger\,\mathcal{M}_{ij}\,\delta\hat{\Phi}^j,$$

with $\mathcal{M}_{ij}$ defined at $\bar{\Phi}$.

Step 2: Change of basis to geometric fluctuations. Use the inverse linear response kernel $(\mathcal{E}^{-1})^{i}_{\alpha\beta}$ (at $\bar{\Phi}$) to express $\delta\hat{\Phi}^{i} = (\mathcal{E}^{-1})^{i}_{\alpha\beta}\,\delta\hat{\Sigma}^{\alpha\beta}$. Substituting gives a quadratic form in $\delta\hat{\Sigma}$ whose coefficient is the tensorial mass kernel

$$\mathcal{M}^{\mu\nu\alpha\beta} := \sum_{i,j} \Big((\mathcal{E}^{-1})^{i}_{\mu\nu}\Big)^{\dagger} \mathcal{M}_{ij} \Big((\mathcal{E}^{-1})^{j}_{\alpha\beta}\Big), \qquad \mathcal{M}^{\mu\nu}{}_{\alpha\beta} = \mathcal{M}^{\mu\nu\rho\sigma}\,\bar{\Sigma}_{\rho\alpha}\,\bar{\Sigma}_{\sigma\beta},$$

which is equivalent to eq. (11.62).

Step 3: Typed projection of the mass operator. Project onto the typed graviton carrier $\hat{h}_{\mu\nu} = \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\delta\hat{g}_{\mu\nu}$ (Definitions 11.3, 11.23 and 11.28). Define the TT block

$$\mathcal{M}^{[\mathrm{TT}]} := \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\;\mathcal{M}\;\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}},$$

so that the mass term acting on $\hat{h}$ is governed by $\mathcal{M}^{[\mathrm{TT}]}$ (Definition 11.33).

Step 4: SSB and null directions. If $\mathcal{M}_{ij}$ has Goldstone null directions (spontaneous breaking $G\to H$ at $\bar{\Phi}$; Proposition 11.29), then the congruence through $\mathcal{E}^{-1}$ maps these nulls into null symmetric tensors of $\mathcal{M}^{\mu\nu}{}_{\alpha\beta}$. Upon TT projection, any component that overlaps the typed carrier yields a null of $\mathcal{M}^{[\mathrm{TT}]}$. Hence, in the exact SSB limit,

$$\operatorname{Spec}\mathcal{M}^{[\mathrm{TT}]} \;=\; \{0\} \quad \text{on } \mathfrak{P},$$

and the typed equation of motion is massless (up to sources), consistent with the Goldstone pole in the TT two-point function (Proposition 11.29 and Definition 11.21).

Step 5: Small explicit breaking (GMOR-style). If a deformation of size ϵ explicitly breaks the symmetry, the TT block acquires a small eigenvalue $m^2_{\mathrm{TT}} = \mathcal{O}(\epsilon)$ bounded on the plateau, $0 \le m^2_{\mathrm{TT}} \le C_{\mathfrak{P}}\epsilon$ (Corollary 11.30).

Certification notes. Positivity on $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ and hyperbolicity of the TT kinetic principal symbol are *plateau checks* and are verified as in Remark 11.35 (Nielsen-flat on $\mathfrak{P}$). □

Definition 11.33 (TT-projected graviton mass operator)**.** Let $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ be the background–covariant TT projector (Chapter 11; cf. Definitions 11.3 and 11.23). The *typed* mass operator acting on the graviton carrier $\hat{h}_{\mu\nu} = \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\delta\hat{g}_{\mu\nu}$ is

$$\boxed{\mathcal{M}^{[\mathrm{TT}]} \;:=\; \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\;\mathcal{M}\;\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,,} \tag{11.63}$$

i.e. the TT block of $\mathcal{M}^{\mu\nu}{}_{\alpha\beta}$ obtained by projection on both indices. In the exact SSB setting of Proposition 11.29 one has

$$\boxed{\operatorname{Spec}\mathcal{M}^{[\mathrm{TT}]} \;=\; \{0\} \qquad \text{on the certified plateau } \mathfrak{P},} \tag{11.64}$$

so the typed mass operator vanishes on $\hat{h}$ and yields the massless TT equation of motion.

Corollary 11.34 (Typed spectrum under small explicit breaking)**.** *If a symmetry-breaking deformation of size ϵ is introduced in $\mathcal{V}_{\Phi}$, then the TT block acquires a small spectral shift $0 \le m^2_{\mathrm{TT}} \le C_{\mathfrak{P}}\,\epsilon$ on $\mathfrak{P}$ (cf. Corollary 11.30).*

Remark 11.35 (Plateau certification: positivity and hyperbolicity)**.** On the physical Hilbert space $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$ (Chapter 8), the quadratic TT form induced by $\mathcal{M}^{[\text{TT}]}$ is non-negative, and the principal symbol of the typed kinetic operator $K_{[\text{TT}]}$ is hyperbolic with respect to the $\bar{\Sigma}$-cone. Both statements are *plateau checks* (see Definitions 6.280 and 9.10 and Proposition 7.3): they are evaluated with BRST-closed band tags and are Nielsen-flat on $\mathfrak{P}$.

Channel Projection of the Graviton

In preparation for vertices and scattering, we refine the typed TT carrier $\hat{h}_{\mu\nu} := \Pi_{\bar{\Sigma}}^{[\text{TT}]}\,\delta\hat{g}_{\mu\nu}$ (Definitions 11.3 and 11.28) by projecting it onto *coupling channels* determined by the background–covariant bi–local parent interaction of Chapter 11 (cf. the $\Phi\Phi \leftrightarrow \Theta\Theta$ off–diagonal) and its linearized TT shadow $S_{\text{int}}^{(1)}[\hat{h}, \hat{\Theta}]$ (Section 11.1). All statements are made on a certified plateau $\mathfrak{P}$ with BRST–closed band tags $\mathfrak{B}_{\lambda_{\text{UV}},\ell}$ commuting with $\Pi_{\bar{\Sigma}}^{[\text{TT}]}$ and the triad projectors $\{\Pi^{(a)}\}_{a\in\{\text{corr,vac,spec}\}}$ (Definitions 6.280 and 9.10).

Definition 11.36 (Typed channel projectors)**.** Let $\mathscr{C}$ denote the finite (or countable) index set of *TT–preserving* coupling channels extracted from the TT–linearized vertex kernel $\hat{V}^{\mu\nu}(x;\hat{\Theta})$ (Chapter 11). A family of bounded projectors $\{\Pi^{\langle c\rangle}\}_{c\in\mathscr{C}}$ on the graviton carrier Hilbert space is called a *typed channel decomposition* if, on $\mathfrak{P}$,

$$\Pi^{\langle c\rangle}\Pi^{\langle c'\rangle} = \delta_{cc'}\Pi^{\langle c\rangle}, \qquad \sum_{c\in\mathscr{C}} \Pi^{\langle c\rangle} = \mathbf{1}, \tag{11.65}$$

$$\left[\Pi^{\langle c\rangle}, \Pi_{\bar{\Sigma}}^{[\text{TT}]}\right] = 0, \qquad \left[\Pi^{\langle c\rangle}, \Pi^{(a)}\right] = 0, \qquad \left[\Pi^{\langle c\rangle}, P_x(\lambda_{\text{UV}})\right] = \left[\Pi^{\langle c\rangle}, K_{\text{IR}}^{(\ell)}\right] = 0. \tag{11.66}$$

The channel–resolved graviton and sector fields are

$$\hat{h}_{\mu\nu}^{\langle c\rangle} := \Pi^{\langle c\rangle}\hat{h}_{\mu\nu}, \qquad \hat{h}_{\mu\nu}^{(a,\langle c\rangle)} := \Pi^{(a)}\Pi^{\langle c\rangle}\hat{h}_{\mu\nu} = \Pi^{\langle c\rangle}\Pi^{(a)}\hat{h}_{\mu\nu}. \tag{11.67}$$

Remark 11.37 (Construction by spectral calculus of the vertex)**.** When $\hat{V}[\cdot]$ admits a self–adjoint TT reduction on $\mathfrak{P}$, the $\Pi^{\langle c\rangle}$ can be obtained as spectral projectors of a commuting family of Hermitian coupling operators (e.g. charge, spin, or representation labels carried on the right leg Θ and pulled back to the TT left leg via the bi–local kernel). TT–nullity of $\hat{\tau}_{\mu\nu}$ (Chapter 11) ensures channels are purely metric on the left leg.

Proposition 11.38 (Orthogonality, support, and microcausality)**.** *On $\mathfrak{P}$ with BRST–closed $\mathfrak{B}_{\lambda_{\text{UV}},\ell}$,*

*(i) (*Orthogonality*) The only non–vanishing mode algebra across channels and triad labels is*

$$\left[\hat{a}_{a,\lambda}^{\langle c\rangle}(\mathbf{k}),\, \hat{a}_{b,\lambda'}^{\langle c'\rangle\,\dagger}(\mathbf{k}')\right] = \delta_{ab}\delta_{cc'}\delta_{\lambda\lambda'}(2\pi)^{d-1}2\omega_a(\mathbf{k})\,\delta^{(d-1)}(\mathbf{k}-\mathbf{k}')\,,$$

so the Fock space factorizes as $\mathscr{F}_\Phi \simeq \widehat{\bigotimes}_{c\in\mathscr{C}}\left(\mathscr{F}_\Phi^{(\text{corr},\langle c\rangle)}\widehat{\otimes}\mathscr{F}_\Phi^{(\text{vac},\langle c\rangle)}\widehat{\otimes}\mathscr{F}_\Phi^{(\text{spec},\langle c\rangle)}\right)$ (cf. Proposition 11.20).

(ii) (Band support) $P_x(\lambda_{\mathrm{UV}})K_{\mathrm{IR}}^{(\ell)}\hat{h}_{\mu\nu}^{(a,\langle c\rangle)} = \hat{h}_{\mu\nu}^{(a,\langle c\rangle)}$.

(iii) (Microcausality) *Channel projection commutes with the typed Peierls kernel, hence* $\Delta^{[\mathrm{TT}],\langle c\rangle} := \Pi^{\langle c\rangle}\Delta^{[\mathrm{TT}]}\Pi^{\langle c\rangle}$ *has support on the* $\bar{\Sigma}$*–null cone and vanishes for* $\bar{\Sigma}$*–spacelike separations (Proposition 11.14).*

Definition 11.39 (Channel–resolved residues and visibilities)**.** Let $G^{[\mathrm{TT}],\langle c\rangle}$ be the channel–projected TT two–point function. In momentum space each triad sector admits a KL representation on $\mathcal{H}_{\mathrm{phys}}$,

$$\Delta^{(2),a,\langle c\rangle}(q) = \frac{Z_a^{\langle c\rangle}}{q^2+i\epsilon} + \int_0^\infty \frac{\rho_a^{(2)\,\langle c\rangle}(\mu^2)\,d\mu^2}{q^2-\mu^2+i\epsilon}, \qquad Z_a^{\langle c\rangle} \geq 0,\ \rho_a^{(2)\,\langle c\rangle} \geq 0.$$

Define the *channel residue vector* $\vec{Z}^{\langle c\rangle} := \left(Z_{\mathrm{corr}}^{\langle c\rangle}, Z_{\mathrm{vac}}^{\langle c\rangle}, Z_{\mathrm{spec}}^{\langle c\rangle}\right)$. The typed Parasymmetric TLO state visibility refines to $\Lambda_{\mathrm{state}}^{[\mathrm{TT}],\langle c\rangle}$ by replacing sector residues with $\vec{Z}^{\langle c\rangle}$ in the readout (Chapter 9).

Definition 11.40 (Channel purity and cross–talk leak)**.** The *channel purity* residual is $\|\Pi^{\perp\langle c\rangle}\hat{h}_{\mu\nu}\|_{\mathcal{I}}$ with $\Pi^{\perp\langle c\rangle} := \mathbf{1} - \Pi^{\langle c\rangle}$. For $c \neq c'$, the *cross–talk* leak is the minimal $\varepsilon_{c\to c'} \geq 0$ such that $\|\Pi^{\langle c'\rangle}\mathfrak{O}\,\Pi^{\langle c\rangle}\|_{\mathcal{I}} \leq \varepsilon_{c\to c'}\|\mathfrak{O}\|_{\mathcal{I}}$ for all observables $\mathfrak{O}$ entering the TT readouts. On $\mathfrak{P}$ these diagnostics are Nielsen–flat and band–stable (Chapter 8).

Remark 11.41 (Law channel in each coupling channel)**.** The TT law residual $\mathbf{R}^{[\mathrm{TT}]}$ (Definition 11.6) refines to $\mathbf{R}^{[\mathrm{TT}],\langle c\rangle} := \Pi^{\langle c\rangle}\mathbf{R}^{[\mathrm{TT}]}\Pi^{\langle c\rangle}$, with associated $\Lambda_{\mathrm{law}}^{[\mathrm{TT}],\langle c\rangle} \in [0,1]$. In the exact SSB typed limit (Proposition 11.29) one expects $\Lambda_{\mathrm{law}}^{[\mathrm{TT}],\langle c\rangle} \to 1$ for channels aligned with the Goldstone TT carrier, while small explicit breakings shift the visibility according to the GMOR–style bounds (Corollary 11.30).

This channel decomposition will be used immediately in Sections 11.3.3 to 11.3.5, where the vertex is written in the channel basis and dispersion/positivity analyses (Section 11.3.6) are evaluated per–channel, per–triad, on the same certified plateau $\mathfrak{P}$.

Full typed graviton and equivalence of constructions. Using the background–covariant TT projector (Chapter 11) and the spin–2 projection of the Gaiamiti fluctuation, the full graviton carrier is

$$\boxed{\hat{h}_{\mu\nu}(x) \;:=\; \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}_{\mu\nu}(x) \;\equiv\; \mathcal{P}^{(2)}\Big[\delta\hat{\mathbf{\Phi}}\Big](x)\cdot\mathbb{E}_{\mu\nu}^{\dagger},} \tag{11.68}$$

in agreement with Definitions 11.23 and 11.28 and eq. (11.49). By construction $\hat{h}_{\mu\nu} = \hat{h}_{(\mu\nu)}$, $\bar{g}^{\mu\nu}\hat{h}_{\mu\nu} = 0$, $\bar{\nabla}^{\mu}\hat{h}_{\mu\nu} = 0$, and the temporal/coherence fluctuation $\hat{\tau}_{\mu\nu}$ is TT-null (Chapter 11).

Step 1: Sector and channel projectors (interaction–sourced)

Let $\{\Pi^{(a)}\}_{a\in\{\mathrm{corr,vac,spec}\}}$ be the triad (sector) projectors and $\{\Pi^{\langle c\rangle}\}_{c\in\mathscr{C}}$ the typed channel projectors extracted from the TT-linearized vertex (Chapter 11), with commutation and

orthogonality as in Definitions 11.24 and 11.36. They commute with $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ and with the band tags on $\mathfrak{P}$ (Definitions 6.280 and 9.10). The interaction kernel admits a triadic decomposition $\mathcal{U}_{\Phi\Theta} = \mathcal{U}_{\Phi\Theta}^{(\mathrm{corr})} + \mathcal{U}_{\Phi\Theta}^{(\mathrm{vac})} + \mathcal{U}_{\Phi\Theta}^{(\mathrm{spec})}$ (Section 10.19.6), which induces the corresponding decomposition of the *typed* response.

Step 2: Sectoral and channel–resolved gravitons

We define the sectoral and channel–resolved graviton fields purely by commuting projections,

$$\hat{h}_{\mu\nu}^{(a)} := \Pi^{(a)}\,\hat{h}_{\mu\nu}, \qquad \sum_a \Pi^{(a)} = \mathbf{1}, \quad \Pi^{(a)}\Pi^{(b)} = \delta_{ab}\Pi^{(a)}, \quad [\Pi^{(a)}, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}] = 0, \tag{11.69}$$

$$\hat{h}_{\mu\nu}^{(a,\langle c\rangle)} := \Pi^{\langle c\rangle}\Pi^{(a)}\,\hat{h}_{\mu\nu} = \Pi^{(a)}\Pi^{\langle c\rangle}\,\hat{h}_{\mu\nu}, \qquad \sum_{c\in\mathscr{C}} \Pi^{\langle c\rangle} = \mathbf{1}, \tag{11.70}$$

so that $\hat{h}_{\mu\nu} = \sum_{a,c} \hat{h}_{\mu\nu}^{(a,\langle c\rangle)}$. When one wishes to display the interaction-sourced form explicitly, this becomes

$$\hat{h}_{\mu\nu}^{(a,\langle c\rangle)}(x) = \Pi^{\langle c\rangle}\Pi^{(a)}\left(\mathcal{P}^{(2)}\left[\int d^dx'\,\mathcal{U}_{\Phi\Theta}^{(a)}(x,x')\cdot\delta\hat{\boldsymbol{\Theta}}(x')\right]\cdot\mathbb{E}_{\mu\nu}^{\dagger}\right). \tag{11.71}$$

which is equivalent by commutativity to first projecting $\hat{h}$ and then expanding in modes (Definition 11.19 and eq. (11.34)).

Step 3: Physical interpretation and support

Each $\hat{h}_{\mu\nu}^{(a,\langle c\rangle)}$ captures the typed TT response in sector a through coupling channel $\langle c\rangle$. For example, the correlational channel encodes long-range, coherence-induced curvature; it vanishes if $\mathcal{U}_{\Phi\Theta}^{(\mathrm{corr})} \equiv 0$. On the certified plateau $\mathfrak{P}$,

$$P_x(\lambda_{\mathrm{UV}})\,K_{\mathrm{IR}}^{(\ell)}\,\hat{h}_{\mu\nu}^{(a,\langle c\rangle)} = \hat{h}_{\mu\nu}^{(a,\langle c\rangle)},$$

so band support is preserved and microcausality follows from the typed Peierls kernel after projection (Propositions 11.14 and 11.38).

Definition: Correlational Graviton

Definition 11.42 (Sectoral graviton fields via commuting triadic projection)**.** Let the typed graviton carrier be $\hat{h}_{\mu\nu} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}_{\mu\nu} \equiv \mathcal{P}^{(2)}[\delta\hat{\boldsymbol{\Phi}}]\cdot\mathbb{E}_{\mu\nu}^{\dagger}$ (Definitions 11.23 and 11.28 and eq. (11.49)). Let $\{\Pi^{(a)}\}_{a\in\{\mathrm{corr,vac,spec}\}}$ be the sector projectors that resolve the identity and commute with $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ on the certified plateau $\mathfrak{P}$ (Definitions 6.280, 9.10 and 11.24). The *sectoral graviton fields* are defined purely by commuting projections:

$$\hat{h}_{\mu\nu}^{(a)}(x) := \Pi^{(a)}\,\hat{h}_{\mu\nu}(x) = \Pi^{(a)}\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}_{\mu\nu}(x), \qquad \sum_a \Pi^{(a)} = \mathbf{1}, \quad \Pi^{(a)}\Pi^{(b)} = \delta_{ab}\Pi^{(a)}. \tag{11.72}$$

In particular, the *Correlational Graviton* is

$$\boxed{\hat{h}^{(\mathrm{corr})}_{\mu\nu}(x) := \Pi^{(\mathrm{corr})}\,\hat{h}_{\mu\nu}(x).} \tag{11.73}$$

When one wishes to display the interaction-sourced form explicitly (cf. Sections 10.19.6 and 11.3), this is equivalent—by commutativity of the projectors on $\mathfrak{P}$—to

$$\boxed{\hat{h}^{(\mathrm{corr})}_{\mu\nu}(x) \;\equiv\; \mathcal{P}^{(2)}\Big[\int d^dx'\,\mathcal{U}^{(\mathrm{corr})}_{\Phi\Theta}(x,x')\cdot\delta\hat{\boldsymbol{\Theta}}(x')\Big]\cdot\mathbb{E}^{\dagger}_{\mu\nu},} \tag{11.74}$$

which is a notational convenience; the *defining* construction in this chapter is the typed/sector projector definition (11.72). The temporal/coherence fluctuation $\hat{\tau}_{\mu\nu}$ is TT-null (Chapter 11) and cannot contribute to any $\hat{h}^{(a)}_{\mu\nu}$.

Definition: Vacuum Graviton Field

Definition 11.43 (Vacuum graviton (typed sectoral projection))**.** The *vacuum graviton field* is the vacuum-sector projection of the typed carrier:

$$\boxed{\hat{h}^{(\mathrm{vac})}_{\mu\nu}(x) \;:=\; \Pi^{(\mathrm{vac})}\,\hat{h}_{\mu\nu}(x) \;=\; \Pi^{(\mathrm{vac})}\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{g}_{\mu\nu}(x).} \tag{11.75}$$

When the interaction-kernel representation is preferred for exposition, $\hat{h}^{(\mathrm{vac})}_{\mu\nu}$ admits the equivalent form $\mathcal{P}^{(2)}\big[\int\mathcal{U}^{(\mathrm{vac})}_{\Phi\Theta}\cdot\delta\hat{\boldsymbol{\Theta}}\big]\cdot\mathbb{E}^{\dagger}_{\mu\nu}$ (cf. Section 10.19.6); physically it captures curvature sourced by vacuum structure $\bar{\Theta}$ and the symmetry content of $\mathcal{M}_{\mathrm{vac}}$ (Section 9.18.6).

Physical Interpretation

- $\hat{h}^{(\mathrm{vac})}_{\mu\nu}$ encodes quasi-static or slowly varying TT curvature arising from vacuum expectation structure rather than long-range entanglement; it is dominant when $\mathcal{M}_{\mathrm{vac}}$ carries nontrivial order (e.g. SSB regimes).
- On $\mathfrak{P}$, band support is preserved: $P_x(\lambda_{\mathrm{UV}})K^{(\ell)}_{\mathrm{IR}}\hat{h}^{(a)}_{\mu\nu} = \hat{h}^{(a)}_{\mu\nu}$, and microcausality follows from the typed Peierls kernel after projection (Propositions 11.14 and 11.38).

Definition: Spectral Graviton Field

Definition 11.44 (Spectral graviton (typed sectoral projection))**.** The *spectral graviton field* is the spectral-sector projection of the typed carrier:

$$\boxed{\hat{h}^{(\mathrm{spec})}_{\mu\nu}(x) \;:=\; \Pi^{(\mathrm{spec})}\,\hat{h}_{\mu\nu}(x) \;=\; \Pi^{(\mathrm{spec})}\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{g}_{\mu\nu}(x).} \tag{11.76}$$

Equivalently (for display only), one may write $\hat{h}^{(\mathrm{spec})}_{\mu\nu} \equiv \mathcal{P}^{(2)}\big[\int\mathcal{U}^{(\mathrm{spec})}_{\Phi\Theta}\cdot\delta\hat{\boldsymbol{\Theta}}\big]\cdot\mathbb{E}^{\dagger}_{\mu\nu}$ to emphasize sourcing by localized/high-frequency matter dynamics (Section 10.19.6).

Physical Interpretation

- *Spectral sector (typed TT).* The field $\hat{h}^{(\mathrm{spec})}_{\mu\nu}(x) := \Pi^{(\mathrm{spec})}\hat{h}_{\mu\nu}(x)$ (Definition 11.44) captures TT curvature sourced by the *local spectral content* of the Tantrakyma field, i.e. high-frequency or localized excitations pulled back through the spectral component of the interaction kernel $\mathcal{U}^{(\mathrm{spec})}_{\Phi\Theta}$ (Section 10.19.6).
- *Dominant regime and causal control.* This contribution dominates when the matter spectral density is large—e.g., in scattering processes or sharply localized wavepackets—and it is the natural arena for the typed propagator and amplitude constructions (Sections 11.2 and 11.3.4). Microcausality is preserved after sector projection because the channel/sector projectors commute with the typed Peierls kernel (Propositions 11.14 and 11.38).
- *Complementarity within the triad.* $\hat{h}^{(\mathrm{spec})}$ complements the *correlational* and *vacuum* gravitons—$\hat{h}^{(\mathrm{corr})}$ (Definition 11.42 and eq. (11.74)) encodes long-range, coherence-driven curvature; $\hat{h}^{(\mathrm{vac})}$ (Definition 11.43) encodes vacuum-structure responses (e.g., SSB backgrounds). Their relative weights are quantified by the triadic residue vector $\vec{Z}$ (Definition 11.21) and can be further refined per coupling channel via the channel-resolved visibilities (Definition 11.39). On the certified plateau $\mathfrak{P}$ the band tags preserve support and factorization across sectors (Definitions 6.280 and 9.10 and Proposition 11.20).

Graviton Mode	Source Channel	Dominant Regime	Interpretative Notes
$\hat{h}^{(\mathrm{corr})}_{\mu\nu}(x)$	Quantum-informational structure; long-range entanglement	**Pre-geometric / coherence-dominated to mesoscopic**	Geometry-as-entanglement response; suppressed when coherence meters vanish.
$\hat{h}^{(\mathrm{vac})}_{\mu\nu}(x)$	Symmetry/ordering of the vacuum manifold $\mathcal{M}_{\mathrm{vac}}$	**Background-dominated / symmetry-broken phases**	Quasi-static TT curvature from vacuum alignment and order parameters (SSB).
$\hat{h}^{(\mathrm{spec})}_{\mu\nu}(x)$	Localized energy–momentum / high-frequency spectral content	**Microscopic to high-energy / short-wavelength**	Scattering and localized wavepacket regime; approaches classical GR limit for coarse-grained stress–energy.

Table 11.1: Dominance regimes of the triad-resolved graviton contributions.

11.1.4 Triadic Decomposition of the Graviton Field

We have established that the quantum graviton field $\hat{h}_{\mu\nu}(x)$ is the typed, transverse–traceless (TT) projection of the metric fluctuation (equivalently, the spin–2 projection of the Gaiamiti fluctuation) via a geometric embedding (Definitions 11.23 and 11.28 and eq. (11.49)). A central feature of the IQR formalism is that the fluctuation field is not monolithic: it admits a decomposition into three orthogonal components reflecting distinct relational origins—*correlational*, *vacuum*, and *spectral*. Because $\mathcal{P}^{(2)}$ (equivalently $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$) is linear and commutes with the sector projectors on the certified plateau $\mathfrak{P}$ (Definitions 6.280, 9.10 and 11.24), the typed graviton inherits a natural triadic decomposition.

1. Mathematical basis of the decomposition

Let $\delta\hat{\mathbf{\Phi}}(x) \in \Gamma[\mathrm{End}(T_{|\Omega(x)\rangle}\mathcal{M}_{\mathrm{vac}})]$ admit the triadic split

$$\delta\hat{\mathbf{\Phi}}(x) = \delta\hat{\mathbf{\Phi}}^{(\mathrm{corr})}(x) + \delta\hat{\mathbf{\Phi}}^{(\mathrm{vac})}(x) + \delta\hat{\mathbf{\Phi}}^{(\mathrm{spec})}(x),$$

as induced by the commuting sector projectors $\{\Pi^{(a)}\}_{a\in\{\mathrm{corr,vac,spec}\}}$ and the triadic interaction channels (cf. Definition 11.24 and Sections 10.19.6 and 11.3). Applying the spin–2 projection and the dual embedding $\mathbb{E}^{\dagger}_{\mu\nu}$ (Definition 11.19), we obtain:

Definition 11.45 (Triadic decomposition of the typed graviton)**.** Let the typed graviton be $\hat{h}_{\mu\nu}(x) := \mathcal{P}^{(2)}[\delta\hat{\mathbf{\Phi}}](x) \cdot \mathbb{E}^{\dagger}_{\mu\nu}$ (eq. (11.49)). Its *triadic decomposition* is

$$\boxed{\hat{h}_{\mu\nu}(x) = \sum_{a\in\{\mathrm{corr,vac,spec}\}} \hat{h}^{(a)}_{\mu\nu}(x), \qquad \hat{h}^{(a)}_{\mu\nu}(x) := \mathcal{P}^{(2)}\Big[\delta\hat{\mathbf{\Phi}}^{(a)}(x)\Big] \cdot \mathbb{E}^{\dagger}_{\mu\nu} \equiv \Pi^{(a)}\hat{h}_{\mu\nu}(x),} \tag{11.77}$$

where the last equality uses $[\Pi^{(a)}, \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}] = 0$ (Definition 11.24).

Each $\hat{h}^{(a)}_{\mu\nu}(x)$ is a symmetric, traceless $(0,2)$-tensor on $(\mathcal{X}_q, \bar{\Sigma})$ and transforms covariantly; $\hat{\tau}_{\mu\nu}$ is TT-null and does not contribute (Chapter 11).

2. Mode algebra and Fock factorization

Combine the definition (11.77) with the triad-resolved TT mode expansion (Definition 11.19 and eq. (11.34)):

$$\hat{h}^{(a)}_{\mu\nu}(x) = \sum_{\lambda=1}^{d_g} \int \frac{d^{d-1}\mathbf{k}}{(2\pi)^{d-1}} \Big[e^{(a,\lambda)}_{\mu\nu}(k;x)\, \hat{a}_{a,\lambda}(\mathbf{k})\, e^{-ik\cdot x} + e^{(a,\lambda)}_{\mu\nu}(k;x)^*\, \hat{a}^{\dagger}_{a,\lambda}(\mathbf{k})\, e^{+ik\cdot x} \Big],$$

with TT polarizations orthonormal in the $\bar{\Sigma}$-inner product (Section 11.1.3). The only nonvanishing commutators are (Proposition 11.20)

$$\boxed{\Big[\hat{a}_{a,\lambda}(\mathbf{k}),\, \hat{a}^{\dagger}_{b,\lambda'}(\mathbf{k}')\Big] = \delta_{ab}\,\delta_{\lambda\lambda'}\,(2\pi)^{d-1}\,2\omega_a(\mathbf{k})\,\delta^{(d-1)}(\mathbf{k}-\mathbf{k}'), \qquad \text{all others } = 0.} \tag{11.78}$$

Corollary 11.46 (Fock factorization across the triad)**.** *The graviton Fock space factorizes as the completed tensor product of sector Fock spaces,*

$$\boxed{\mathscr{F}_\Phi \;\simeq\; \mathscr{F}_\Phi^{(\mathrm{corr})} \,\widehat{\otimes}\, \mathscr{F}_\Phi^{(\mathrm{vac})} \,\widehat{\otimes}\, \mathscr{F}_\Phi^{(\mathrm{spec})},} \tag{11.79}$$

with commuting creation/annihilation operators labeled by the sector index a as in (11.78) *(cf. Proposition 11.20).*

3. Spectral diagnostics and support

On the physical Hilbert space $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ (Chapter 8), each sector's two-point admits a Källén–Lehmann representation with nonnegative residue and density; collecting the residues defines the triadic *residue vector* $\vec{Z} = (Z_{\mathrm{corr}}, Z_{\mathrm{vac}}, Z_{\mathrm{spec}})$ (Definition 11.21). Band tags commute with the sector and typed projectors, hence preserve support and causal properties:

$$P_x(\lambda_{\mathrm{UV}})\, K_{\mathrm{IR}}^{(\ell)}\, \hat{h}_{\mu\nu}^{(a)} = \hat{h}_{\mu\nu}^{(a)}, \tag{11.80}$$

and microcausality follows from the typed Peierls kernel (*Propositions* 11.14*and* 11.38). (11.81)

This triad-resolved structure underlies the channel projections (Section 11.1.3), the propagator (Section 11.2), and scattering amplitudes (Section 11.3.4), with typed Parasymmetric TLO readouts assembled per sector (Chapter 9).

2. Physical Interpretation of the Graviton Species

Each component $\hat{h}_{\mu\nu}^{(a)}(x)$ in the triadic decomposition (11.77) corresponds to a physically distinct graviton contribution, selected by the commuting sector projectors $\Pi^{(a)}$ and compatible with the typed TT carrier $\hat{h}_{\mu\nu}$ (Definitions 11.24, 11.28 and 11.45). The sector labels track relational sources—correlational, vacuum, and spectral—and they propagate through the mode algebra and Fock factorization (Definition 11.19, eq. (11.34), and Corollary 11.46).

- **Correlational Gravitons** $\hat{h}_{\mu\nu}^{(\mathrm{corr})}(x)$. Defined by the sectoral projection $\hat{h}_{\mu\nu}^{(\mathrm{corr})} := \Pi^{(\mathrm{corr})}\hat{h}_{\mu\nu}$ (Definition 11.42), these quanta encode TT curvature sourced by long-range coherence and entanglement structure, optionally represented via the interaction-kernel form (11.74) (see also channel projectors in Definition 11.36). They realize the principle that *information shapes geometry*, resonant with the ER=EPR perspective [520], holographic tensor-network intuition [1], and modular-flow constructions [521, 522]. In the typed ledger, their strength is captured by sectoral residues in the triadic Källén–Lehmann representation ($\vec{Z}$; Definition 11.21) and by the Parasymmetric TLO visibilities (Chapter 8); the contribution vanishes when coherence meters collapse on the certified plateau $\mathfrak{P}$ (Definitions 6.280 and 9.10).

- **Vacuum Gravitons** $\hat{h}_{\mu\nu}^{(\mathrm{vac})}(x)$. Given by $\hat{h}_{\mu\nu}^{(\mathrm{vac})} := \Pi^{(\mathrm{vac})}\hat{h}_{\mu\nu}$ (Definition 11.43 and Section 11.1.3), these excitations arise from the structure of the vacuum manifold $\mathcal{M}_{\mathrm{vac}}$ (Sections 6.1.8 and 9.18.6) and behave as typed Goldstone modes when geometric symmetries are spontaneously broken (Proposition 11.29). Their masslessness corresponds

to the vanishing of the TT-projected mass operator $\mathcal{M}^{[\mathrm{TT}]}$ (Definition 11.33), while small explicit breakings generate $m^2_{\mathrm{TT}} = \mathcal{O}(\epsilon)$ with plateau bounds (Corollary 11.30). These modes capture quasi-static TT curvature from vacuum alignment and order parameters; they also furnish the background against which the typed propagator and Ward identities are evaluated (Section 11.2 and eq. (11.25)). Related vacuum transitions and tunneling phenomena are encompassed in the standard picture of false-vacuum decay [523].

- **Spectral Gravitons** $\hat{h}^{(\mathrm{spec})}_{\mu\nu}(x)$. Defined by $\hat{h}^{(\mathrm{spec})}_{\mu\nu} := \Pi^{(\mathrm{spec})}\hat{h}_{\mu\nu}$ (Definition 11.44 and Section 11.1.3) and optionally by the spectral piece of the interaction kernel $\mathcal{U}^{(\mathrm{spec})}_{\Phi\Theta}$ (Section 10.19.6), these carry energy–momentum–dominated TT curvature. They govern high-frequency, localized, or high-energy regimes and interface directly with the typed propagator and scattering constructions (Sections 11.2, 11.2.2 and 11.3.4). In the semiclassical limit they reduce to helicity-2 perturbations of linearized GR [245], with microcausality ensured by the typed Peierls kernel and band-stable support (Propositions 11.14 and 11.38 and Definitions 6.280 and 9.10).

3. Consequences and Theoretical Significance

The triadic decomposition $\hat{h}_{\mu\nu} = \sum_{a\in\{\mathrm{corr,vac,spec}\}} \hat{h}^{(a)}_{\mu\nu}$ (Definition 11.45 and eq. (11.77)) reveals the fundamentally *composite* character of gravitational excitation in the IQR framework. Because the sector projectors $\Pi^{(a)}$ commute with the typed TT projector $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ on the certified plateau $\mathfrak{P}$ (Definitions 6.280, 9.10 and 11.24), each contribution propagates as a well-defined typed channel compatible with microcausality (Proposition 11.14) and with the mode/Fock infrastructure (Definition 11.19, eq. (11.34), and Corollary 11.46). This leads to the following implications:

- **Multi-channel gravitational interaction.** The sector (and optional coupling-channel) resolution (Section 11.1.3 and Definition 11.36) organizes propagation and interactions into independently readable blocks. In the sector/channel basis, the typed two-point and propagator inherit a block-diagonal structure compatible with the mode expansion and background geometry (Sections 11.2, 11.2.2 and 11.2.3). Dynamical couplings descend from the TT-linearized parent vertex (Section 11.3) and can be monitored via the typed Parasymmetric TLO readouts: law and state visibilities (Definitions 11.6 and 11.39), with positivity/dispersion constraints evaluated per block (Section 11.3.6). Renormalization and gauge-parameter independence are assessed on $\mathfrak{P}$ (Nielsen-flatness; Chapter 8).

- **New phenomenological pathways.** Contexts such as early-universe phases with strong vacuum alignment, near-vacuum decoherence, or topological transitions can be dominated by $\hat{h}^{(\mathrm{vac})}$ or $\hat{h}^{(\mathrm{corr})}$, not by $\hat{h}^{(\mathrm{spec})}$ (Definitions 11.42 and 11.43). Typed SSB protects masslessness in the TT sector (Proposition 11.29), while small explicit breakings obey the GMOR-style bound $m^2_{\mathrm{TT}} = \mathcal{O}(\epsilon)$ (Corollary 11.30) through the TT mass block $\mathcal{M}^{[\mathrm{TT}]}$ (Definition 11.33). Hence, observable geometric dynamics may arise without classical stress–energy dominance, and they remain consistent with causality via the typed Peierls bracket (eq. (11.25)).

- **A composite quantum graviton.** In IQR the graviton is not elementary but a sector-resolved superposition of typed TT excitations with distinct relational origins—entanglement/coherence, vacuum geometry, and spectral content (Definitions 11.42 to 11.44). This composition is quantitatively captured by the triadic residue vector $\vec{Z} = (Z_{\text{corr}}, Z_{\text{vac}}, Z_{\text{spec}})$ (Definition 11.21), realized in the KL representation on $\mathcal{H}_{\text{phys}}$ (Chapter 8), and supported by the Fock factorization across sectors (Corollary 11.46).

This reformulation places the gravitational field at the confluence of quantum information, vacuum structure, and energy–momentum dynamics—unifying these sources within a single typed, BRST-healthy, sector- and channel-aware pipeline (Definitions 11.6 and 11.24 and Sections 11.1.3, 11.2 and 11.3).

Graviton Type	Source of Geometry	Related Paradigm
Correlational	Entanglement and quantum coherence structure (sector projector $\Pi^{(\text{corr})}$)	ER=EPR; holographic tensor networks; modular flow
Vacuum	Moduli space and symmetry structure of $\mathcal{M}_{\text{vac}}$ (sector projector $\Pi^{(\text{vac})}$)	Goldstone modes; vacuum topology; phase transitions (SSB/GMOR)
Spectral	Local spectral/energy–momentum content (sector projector $\Pi^{(\text{spec})}$)	Linearized GR; GW phenomenology; semiclassical gravity

Table 11.2: The triadic nature of gravitational quanta and their conceptual anchors.

This triadic picture extends the physical meaning of gravitational phenomena beyond stress–energy sourcing, while preserving causality, positivity, and typed consistency on $\mathfrak{P}$ (Chapter 8). It enriches the ontology of quantum gravity and enables novel, nonlocal and symmetry-based interactions to be formulated and tested within a unified, sector-resolved framework.

11.1.5 Graviton Fock Space and Multi-Particle States

With the linearized, quantized TT modes identified (Definition 11.19 and eq. (11.34)) and the triad-resolved structure established (Definition 11.45 and eq. (11.77)), we now construct the *typed, triad-resolved* Fock space for geometric excitations. All constructions are performed on a certified plateau $\mathfrak{P}$ with BRST-closed band tags commuting with the typed and sector projectors (Definitions 6.280, 9.10 and 11.24), and all physical states are implicitly taken in the BRST cohomology $\mathcal{H}_{\text{phys}} := H^0(\hat{Q}_{\text{BRST}})$ (Chapter 8).

The Graviton Vacuum State

Definition 11.47 (Graviton vacuum (typed/triad, BRST-closed))**.** Let $|0\rangle_\Phi$ denote a vacuum vector annihilated by all TT graviton annihilation operators $\hat{a}_{a,\lambda}(\mathbf{k})$ appearing in the triad-

resolved mode expansion (Section 11.1.3 and Definition 11.19):

$$\boxed{\hat{a}_{a,\lambda}(\mathbf{k})\,|0\rangle_\Phi = 0, \qquad \forall\, a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\},\ \lambda,\ \mathbf{k}.} \tag{11.82}$$

The *physical* vacuum is its BRST cohomology class $[\,|0\rangle_\Phi\,] \in \mathcal{H}_{\mathrm{phys}}$, satisfying $\hat{Q}_{\mathrm{BRST}}|0\rangle_\Phi = 0$ (Chapter 8).

The Graviton Fock Space

Definition 11.48 (Typed, triad-resolved graviton Fock space)**.** Let $\mathcal{H}_\Phi^{(1)}$ be the one-particle TT Hilbert space spanned by $\hat{a}^\dagger_{a,\lambda}(\mathbf{k})|0\rangle_\Phi$. The (bosonic) Fock space is

$$\boxed{\mathscr{F}_\Phi := \mathbb{C}\,|0\rangle_\Phi \ \oplus \bigoplus_{n\geq 1} \mathrm{Sym}\left[\underbrace{\mathcal{H}_\Phi^{(1)} \otimes \cdots \otimes \mathcal{H}_\Phi^{(1)}}_{n\ \text{times}}\right],} \tag{11.83}$$

and its *physical* subspace is the BRST cohomology $\mathscr{F}_\Phi^{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}}; \mathscr{F}_\Phi)$ (Chapter 8). The creation/annihilation algebra is sector-diagonal as in (11.78), ensuring independence of the triadic sectors.

Multi-Particle States and Triadic Decomposition

Definition 11.49 (Triadic-labeled creation/annihilation and basis)**.** For each sector $a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$ and polarization λ, the operators $\hat{a}_{a,\lambda}(\mathbf{k})$, $\hat{a}^\dagger_{a,\lambda}(\mathbf{k})$ obey the mode algebra (11.78). A convenient *typed, triad-resolved* basis for the n-particle sector is

$$\boxed{\left|\mathbf{k}_1, \lambda_1, a_1; \ldots; \mathbf{k}_n, \lambda_n, a_n\right\rangle := \frac{1}{\sqrt{n!}}\, \hat{a}^\dagger_{a_1,\lambda_1}(\mathbf{k}_1) \cdots \hat{a}^\dagger_{a_n,\lambda_n}(\mathbf{k}_n)\,|0\rangle_\Phi,} \tag{11.84}$$

implicitly symmetrized under permutations. States are understood modulo BRST-exact vectors; hence we identify their cohomology classes in $\mathscr{F}_\Phi^{\mathrm{phys}} \subset \mathcal{H}_{\mathrm{phys}}$.

Proposition 11.50 (Fock factorization across the triad)**.** *With the sector-diagonal algebra* (11.78)*, the Fock space factorizes as*

$$\boxed{\mathscr{F}_\Phi \ \simeq\ \mathscr{F}_\Phi^{(\mathrm{corr})} \,\widehat{\otimes}\, \mathscr{F}_\Phi^{(\mathrm{vac})} \,\widehat{\otimes}\, \mathscr{F}_\Phi^{(\mathrm{spec})},} \tag{11.85}$$

and the same holds for the physical subspace $\mathscr{F}_\Phi^{\mathrm{phys}}$ (cf. Corollary 11.46).

Remarks (typed certification and visibilities). (i) The *TT state visibility* $\Lambda^{[\mathrm{TT}]}_{\mathrm{state}}$ used in the Parasymmetric TLO readout (Chapter 9) is computed from the TT two-point Källén–Lehmann content—namely, the sectoral residues and spectral densities $\{Z_a, \rho_a^{(2)}\}_{a=\mathrm{corr,vac,spec}}$ (Definitions 11.21 and 11.39). On the certified plateau $\mathfrak{P}$ these readouts are Nielsen-flat and

band-stable (Definitions 6.280 and 9.10). (ii) All multiparticle observables inherit microcausality from the typed Peierls kernel after projection (Proposition 11.14), and their sectoral decomposition is preserved by the commuting projectors $\Pi^{(a)}$ (Definition 11.24). (iii) When channel projectors $\Pi^{\langle c\rangle}$ are used (Definition 11.36), the basis (11.84) refines by including a channel label $\langle c\rangle$, and the factorization extends accordingly.

11.2 The Graviton Propagator

With the typed, triad-resolved graviton field and Fock construction in hand (Definitions 11.45 and 11.48, eq. (11.77), and Corollary 11.46), we now define and analyze the *graviton propagator*. Throughout, causality and equal-time limits are controlled by the typed Peierls structure (eq. (11.25)) on the chronogeometric background $(\mathcal{X}_q, \bar{\Sigma})$ (Section 5.2.6). Time ordering is taken with respect to the $\bar{\Sigma}$-induced foliation (Section 11.2.1). All statements are certified on the plateau $\mathfrak{P}$ with BRST-closed band tags commuting with the typed and sector projectors (Definitions 6.280, 9.10 and 11.24).

We recall the triadic split of the typed graviton (Definition 11.45):

$$\hat{h}_{\mu\nu}(x) = \sum_{a\in\{\mathrm{corr,vac,spec}\}} \hat{h}^{(a)}_{\mu\nu}(x), \qquad \hat{h}^{(a)}_{\mu\nu} = \Pi^{(a)}\hat{h}_{\mu\nu}.$$

Orthogonality of the sector projectors and their commutation with $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ imply a diagonal free two-point structure; hence the full propagator decomposes into a sum of sector blocks with no cross terms at the free level:

$$\boxed{G^{\mathrm{grav}}_{\mu\nu}{}^{\rho\sigma}(x,x') = \sum_{a\in\{\mathrm{corr,vac,spec}\}} G^{(a)}_{\mu\nu}{}^{\rho\sigma}(x,x').} \tag{11.86}$$

Definition 11.51 (Triad-resolved typed graviton propagator)**.** For a given sector $a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$, the **(typed) Feynman two-point function** (graviton propagator) is defined on the physical vacuum (Definition 11.47) by:

$$\boxed{G^{(a)}_{\mu\nu}{}^{\rho\sigma}(x,x') \;:=\; {}_{\Phi}\!\left\langle 0\right| T_{\bar{\Sigma}}\!\left\{\hat{h}^{(a)}_{\mu\nu}(x)\,\hat{h}^{(a)\,\rho\sigma}(x')\right\}\left|0\right\rangle_{\Phi},} \tag{11.87}$$

where $T_{\bar{\Sigma}}$ is the time-ordering operator with respect to the causal time of $\bar{\Sigma}$ (Section 11.2.1). Equivalently, $G^{(a)}$ is the TT-typed Peierls/Feynman kernel of the sectoral wave operator (cf. Section 11.1.2).

1. Relation to mode expansion

Using the triad-resolved TT mode expansion (Definition 11.19 and eq. (11.34)) and Wick's theorem, one finds an operator–polarization representation. In momentum space on a locally stationary patch (or in Riemann normal coordinates), this reads

$$\boxed{G^{(a)}_{\mu\nu}{}^{\rho\sigma}(k) = \frac{-i}{k^2 - m^2_{a,\mathrm{TT}} + i\epsilon}\,\Pi^{[\mathrm{TT}](a)}_{\mu\nu}{}^{\rho\sigma}(k;\bar{\Sigma}),} \tag{11.88}$$

where $\Pi^{[\mathrm{TT}](a)}$ is the TT polarization projector assembled from the orthonormal TT tensors $e^{(a,\lambda)}_{\mu\nu}(k)$ (Section 11.1.3). In the spontaneously-broken, typed limit (Proposition 11.29) one has $m^2_{a,\mathrm{TT}} = 0$; small explicit breakings yield a GMOR-style shift $m^2_{a,\mathrm{TT}} = \mathcal{O}(\epsilon)$ (Corollary 11.30 and Definition 11.33).

2. Propagator as typed Green's function

Let $K^{[\mathrm{TT}]}$ denote the TT-projected Lichnerowicz-type wave operator on $(\mathcal{X}_q, \bar{\Sigma})$, augmented by the TT mass/mixing block. For each sector a,

$$\boxed{K^{[\mathrm{TT}](a)}_{\mu\nu}{}^{\alpha\beta}(x)\,\hat{h}^{(a)}_{\alpha\beta}(x) = 0, \qquad K^{[\mathrm{TT}](a)} := \mathcal{L}^{[\mathrm{TT}]} - \mathcal{M}^{[\mathrm{TT}](a)},} \tag{11.89}$$

with $\mathcal{M}^{[\mathrm{TT}](a)} := \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\mathcal{M}^{(a)}\,\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ (Definition 11.33). The Feynman propagator is the Green's function of $K^{[\mathrm{TT}](a)}$:

$$\boxed{K^{[\mathrm{TT}](a)}_{\mu\nu}{}^{\alpha\beta}(x)\,G^{(a)}_{\alpha\beta}{}^{\rho\sigma}(x,x') = i\,\Pi^{[\mathrm{TT}]}_{\mu\nu}{}^{\rho\sigma}(\bar{\Sigma})\,\delta^{(d)}(x-x'),} \tag{11.90}$$

so that the causal difference of retarded/advanced solutions reproduces the typed Peierls kernel $\Delta^{[\mathrm{TT}]}$ (Section 11.1.2). Microcausality and band support follow from $\operatorname{supp}\Delta^{[\mathrm{TT}]} \subseteq$ the $\bar{\Sigma}$-null cone and the commutation of band tags with the projectors (Propositions 11.14 and 11.38 and Definitions 6.280 and 9.10).

3. Triadic Källén–Lehmann representation

On $\mathcal{H}_{\mathrm{phys}}$ the TT two-point admits a sector-resolved KL form (Definition 11.21):

$$\boxed{\Delta^{(2),a}(q) = \frac{Z_a}{q^2 + i\epsilon} + \int_0^\infty d\mu^2\,\frac{\rho^{(2)}_a(\mu^2)}{q^2 - \mu^2 + i\epsilon}, \qquad Z_a \geq 0,\ \rho^{(2)}_a(\mu^2) \geq 0,} \tag{11.91}$$

with residue vector $\vec{Z} = (Z_{\mathrm{corr}}, Z_{\mathrm{vac}}, Z_{\mathrm{spec}})$. In the exact SSB, TT-typed limit one has $Z_a > 0$ and $\operatorname{Spec}\mathcal{M}^{[\mathrm{TT}]} = \{0\}$ (Proposition 11.29 and Definition 11.33); explicit breakings deform the spectral threshold by $\mathcal{O}(\epsilon)$ (Corollary 11.30). These spectral data feed the TT state-visibility meters in the typed Parasymmetric TLO (Chapter 9).

4. Background-covariant refinements

In general backgrounds one must use bitensors and $\bar{\Sigma}$-parallel transporters; polarization projectors become point-split and the momentum-space formulas give way to Hadamard/Schwinger–DeWitt constructions. These refinements are presented in Section 11.2.3, and the operator-basis viewpoint in Section 11.2.2. The time-ordering and Feynman prescriptions are summarized in Section 11.2.1.

Summary. The typed, triad-resolved graviton propagator is the Green's function of the TT wave operator, diagonal in the triadic basis at the free level, with a sector-wise KL representation that is positive on $\mathcal{H}_{\mathrm{phys}}$ and Nielsen-flat on $\mathfrak{P}$. It furnishes the universal building block for the interaction analysis in Section 11.3.

11.2.1 The Time-Ordering Operator and the Feynman Propagator

In the relational IQR framework, chronological ordering is defined by the causal cone of the chronogeometric background $(\mathcal{X}_q, \bar{\Sigma})$ (Section 5.2.6). Accordingly, all time-ordering and Feynman prescriptions are *background-covariant* with respect to $\bar{\Sigma}_{\mu\nu}$ and are applied after the typed TT selection (Section 11.1.3 and eq. (11.49)). Band tags commute with the projectors and preserve causal support on the certified plateau $\mathfrak{P}$ (Definitions 6.280, 9.10 and 11.24).

Definition 11.52 (TT time–ordering (TLO) ledgers)**.** On the certified plateau $\mathfrak{P}$, the $\bar{\Sigma}$–time–ordering operator $T_{\bar{\Sigma}}$ (Definition 11.53) and the transverse–traceless and triadic projectors satisfy the bookkeeping ("ledgers") below:

(a) **Commutation with projectors.** $[\,T_{\bar{\Sigma}},\, \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,] = 0$ and $[\,T_{\bar{\Sigma}},\, \Pi^{(a)}\,] = 0$ for $a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$; moreover the band tag commutes as well: $[\,\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell},\, \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,] = [\,\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell},\, \Pi^{(a)}\,] = 0$ (Definition 9.10 and Proposition 11.25).

(b) **TT visibility / Ward residual.** For conserved external insertions, longitudinal pieces are contact terms that drop out under TT projection; diagrammatically, contracting external legs with the TT Ward identity leaves only TT exchange (Lemma 11.92).

(c) **Causal compatibility.** $T_{\bar{\Sigma}}$, $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ and the IR window $K^{(\ell)}_{\mathrm{IR}}$ preserve $\bar{\Sigma}$–causal support, so TT–projected retarded/advanced and Feynman kernels are well defined (Sections 11.2.1 and 11.2.3).

These ledgers justify using only TT, triad-resolved internal lines $G^{\mathrm{F},[\mathrm{TT}],(a)}$ and typed vertices in amplitudes (Section 11.3.4).

Definition 11.53 (Time ordering with respect to $\bar{\Sigma}$)**.** Let $J^{\pm}_{\bar{\Sigma}}(x)$ denote the future/past causal sets defined by the $\bar{\Sigma}$-null cone, and let $\Theta_{\bar{\Sigma}}(x \succeq y) := \mathbf{1}_{x \in J^{+}_{\bar{\Sigma}}(y)}$. For bosonic operators $\hat{A}(x), \hat{B}(y)$ we define the $\bar{\Sigma}$-*time-ordering operator* by

$$\boxed{T_{\bar{\Sigma}}\Big\{\hat{A}(x)\hat{B}(y)\Big\} := \Theta_{\bar{\Sigma}}(x \succeq y)\,\hat{A}(x)\hat{B}(y) + \Theta_{\bar{\Sigma}}(y \succeq x)\,\hat{B}(y)\hat{A}(x),} \tag{11.92}$$

with the equal-time limit defined by symmetric point-splitting consistent with the typed Peierls bracket (eq. (11.25)) and reproducing the typed ETCR on a Cauchy slice (Definition 11.8 and Proposition 11.4).

Definition 11.54 (Typed TT Feynman propagator)**.** For the typed graviton $\hat{h}_{\mu\nu}$ (Definition 11.28), the *Feynman two-point function* (TT-projected) is

$$\boxed{i\,G^{\mathrm{F},[\mathrm{TT}]}_{\mu\nu}{}^{\rho\sigma}(x,y) := {}_{\Phi}\langle 0|\, T_{\bar{\Sigma}}\Big\{\hat{h}_{\mu\nu}(x)\,\hat{h}^{\rho\sigma}(y)\Big\}\,|0\rangle_{\Phi}, \qquad |0\rangle_{\Phi} \text{ as in } \textit{Definition } 11.47.} \tag{11.93}$$

Equivalently, since $\hat{h}$ is TT-typed by construction, $G^{\mathrm{F},[\mathrm{TT}]}$ is the TT block of the full Feynman kernel.

Proposition 11.55 (Retarded/advanced representation and Green identity)**.** *Let $G^{\mathrm{ret/adv},[\mathrm{TT}]}$ be the TT-projected retarded/advanced Green operators of the TT wave operator $K^{[\mathrm{TT}]}$ (Section 11.1.2 and eq. (11.89)). Then*

$$\boxed{G^{\mathrm{F},[\mathrm{TT}]}(x,y) = \Theta_{\bar{\Sigma}}(x \succeq y)\, G^{\mathrm{ret},[\mathrm{TT}]}(x,y) + \Theta_{\bar{\Sigma}}(y \succeq x)\, G^{\mathrm{adv},[\mathrm{TT}]}(x,y),} \tag{11.94}$$

and $G^{\mathrm{F},[\mathrm{TT}]}$ solves the TT Green equation with the Feynman $i\epsilon$ prescription:

$$\boxed{K^{[\mathrm{TT}]}_{\mu\nu}{}^{\alpha\beta}(x)\, G^{\mathrm{F},[\mathrm{TT}]}_{\alpha\beta}{}^{\rho\sigma}(x,y) = i\,\Pi^{[\mathrm{TT}]}_{\mu\nu}{}^{\rho\sigma}(\bar{\Sigma})\,\delta^{(d)}(x-y).} \tag{11.95}$$

Consequently, the causal difference reproduces the typed Peierls kernel $\Delta^{[\mathrm{TT}]} = G^{\mathrm{ret},[\mathrm{TT}]} - G^{\mathrm{adv},[\mathrm{TT}]}$ (Section 11.1.2).

Corollary 11.56 (Triadic block-diagonality and microcausality)**.** *With sector projectors $\Pi^{(a)}$ commuting with $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ (Definition 11.24), the Feynman kernel is block-diagonal:*

$$\boxed{G^{\mathrm{F},[\mathrm{TT}]}_{\mu\nu}{}^{\rho\sigma} = \sum_{a\in\{\mathrm{corr,vac,spec}\}} \Pi^{(a)}_{\mu\nu}{}^{\alpha\beta}\, G^{\mathrm{F},[\mathrm{TT}]}_{\alpha\beta}{}^{\gamma\delta}\, \Pi^{(a)}_{\gamma\delta}{}^{\rho\sigma},} \tag{11.96}$$

and microcausality holds: the commutator kernel has support only on the $\bar{\Sigma}$-null cone ($\mathrm{supp}\,\Delta^{[\mathrm{TT}]} \subseteq \partial J^{\pm}_{\bar{\Sigma}}$) (Proposition 11.14).

IR/UV band stability. On $\mathfrak{P}$, inserting the band tags as a sandwich, $G^{\mathrm{F},[\mathrm{TT}]}_{\mathfrak{B}} := K^{(\ell)}_{\mathrm{IR}}\, P_x(\lambda_{\mathrm{UV}})\, G^{\mathrm{F},[\mathrm{TT}]}\, P_x(\lambda_{\mathrm{UV}})\, K^{(\ell)}_{\mathrm{IR}}$, preserves typed support and microcausality because the tags commute with the typed/sector projectors and do not enlarge causal cones (Definitions 6.280 and 9.10 and Proposition 11.38). Equal-time limits of (11.93) reproduce the typed ETCR on a Cauchy slice (Definition 11.8 and Proposition 11.4), while momentum-space forms reduce to the polarization-projector representation (eq. (11.88)).

Canonical role in perturbative gravity. The $\bar{\Sigma}$-time-ordered product $T_{\bar{\Sigma}}\{\cdot\}$ (Definition 11.53) defines the typed Feynman kernel $G^{\mathrm{F},[\mathrm{TT}]}$ (Definition 11.54), which enters every term of the Dyson expansion for scattering on $(\mathcal{X}_q, \bar{\Sigma})$. Concretely, $i\,G^{\mathrm{F},[\mathrm{TT}]}_{\mu\nu}{}^{\rho\sigma}(x,y) = {}_{\Phi}\langle 0|\, T_{\bar{\Sigma}}\{\hat{h}_{\mu\nu}(x)\hat{h}^{\rho\sigma}(y)\}\,|0\rangle_{\Phi}$ provides the covariant, causal amplitude for a TT graviton created at y to be annihilated at x, with forward/backward propagation in $\bar{\Sigma}$-time encoded by eqs. (11.94) and (11.95). Equal-time limits are taken by $\bar{\Sigma}$-covariant point-splitting compatible with the typed Peierls bracket, so that $\lim_{x^0\to y^0} G^{\mathrm{F},[\mathrm{TT}]}$ reproduces the typed ETCR (eq. (11.25), Definition 11.8, and Proposition 11.4). We assume a Hadamard physical state on $\mathcal{H}_{\mathrm{phys}}$ so the two-point function has the standard short-distance form on curved backgrounds. This typed construction is the cornerstone for building covariant S-matrix elements in perturbative QFT, here adapted to the IQR setting ([82, 155, 403, 502]).

Field-theoretic and diagrammatic interpretation. The operator $T_{\bar{\Sigma}}$ enforces causal consistency of virtual exchanges with respect to the $\bar{\Sigma}$-null cone, and the causal difference reproduces the typed Peierls kernel (eq. (11.25) and Proposition 11.14). Hence each internal graviton line is a *TT-projected* Feynman propagator. On the certified plateau $\mathfrak{P}$, the band tags $P_x(\lambda_{\mathrm{UV}})$ and $K_{\mathrm{IR}}^{(\ell)}$ are chosen $\bar{\Sigma}$-covariantly (e.g. as functions of the TT d'Alembertian) and commute with $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$, $\Pi^{(a)}$, and (optionally) channel projectors $\Pi^{\langle c\rangle}$ (Definitions 6.280, 9.10, 11.24 and 11.36), so band-sandwiching *does not enlarge causal cones* (Proposition 11.38) and preserves equal-time limits. In the triadic basis, free internal lines are block-diagonal by sector (Definitions 11.45 and 11.51), with polarization structure captured by the TT projector assembled from orthonormal spin-2 tensors (Section 11.1.3); this mirrors the standard treatment of gauge bosons in perturbation theory ([82, 403, 502]). *Beyond tree level*, triadic blocks can mix via self-energies Σ_{ab} unless symmetry forbids it; our free-level statements and plateau diagnostics apply blockwise.

Physical interpretation of the triadic graviton propagator. As established in Section 11.1.4, Definition 11.51, and eq. (11.86), the sector-resolved kernel $G^{(a)}_{\mu\nu}{}^{\rho\sigma}(x,x')$ acts as the *quantum mediation kernel* for sector a:

- *Kernel.* $G^{(a)}$ is the typed Green function of the TT wave operator with sectoral TT mass/mixing block (eqs. (11.89) and (11.90)); as a convolution kernel it determines the linear response of $\hat{h}^{(a)}$ to localized sources with $\bar{\Sigma}$-causal support.
- *Quantum.* $G^{(a)}$ encodes off-shell propagation of virtual spin-2 quanta. Its KL spectral content—sectoral residues and densities—governs measurable amplitudes and dispersion/positivity constraints per sector (eq. (11.91) and Section 11.3.6). The massless TT pole is protected by the *typed Ward identity* associated with SSB in the Gaiamiti sector (Proposition 11.29); small explicit breaking yields a GMOR-style shift $m^2_{\mathrm{TT}} = \mathcal{O}(\epsilon)$ within plateau bounds (Definition 11.33 and Corollary 11.30). In the semiclassical limit of the spectral sector, $G^{(\mathrm{spec})}$ reproduces the standard helicity-2 propagator of linearized GR ([245, 502]).

Taken together, the typed, triad-resolved Feynman kernels furnish a covariant and causal foundation for perturbative gravity in IQR, unifying informational, vacuum, and energetic sources within a single propagator framework and staging the interaction analysis in Section 11.3.

11.2.2 Mode Expansion and the Operator Basis of the Propagator

The typed graviton Feynman kernel $G^{\mathrm{F},[\mathrm{TT}]}$ admits an explicit mode-sum representation obtained by inserting the triad-resolved TT mode expansion (Section 11.1.3 and eq. (11.34)) into the $\bar{\Sigma}$-time-ordered two-point (Definition 11.54). This builds a bridge between the operator (creation– annihilation) picture and the wave/Green-function picture familiar from perturbative QFT ([82, 524]).

1. Triad-resolved relativistic mode expansion (typed TT)

For each triadic sector $a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$ and orthonormal TT polarization basis $e^{(a,\lambda)}_{\mu\nu}(k)$ (Section 11.1.3), the typed graviton field expands as

$$\boxed{\hat{h}^{(a)}_{\mu\nu}(x) = \sum_{\lambda=1}^{d_g} \int \frac{d^{d-1}\mathbf{k}}{(2\pi)^{d-1}} \frac{1}{\sqrt{2\omega_{\mathbf{k}}}} \Big[\hat{a}_{a,\lambda}(\mathbf{k})\, e^{(a,\lambda)}_{\mu\nu}(k)\, e^{-ikx} + \hat{a}^{\dagger}_{a,\lambda}(\mathbf{k})\, e^{(a,\lambda)*}_{\mu\nu}(k)\, e^{+ikx} \Big],} \tag{11.97}$$

with $\omega_{\mathbf{k}} = \sqrt{\mathbf{k}^2 + m^2_{a,\mathrm{TT}}}$ and $[\hat{a}_{a,\lambda}(\mathbf{k}), \hat{a}^{\dagger}_{b,\lambda'}(\mathbf{k}')] = \delta_{ab}\delta_{\lambda\lambda'}(2\pi)^{d-1}\delta^{(d-1)}(\mathbf{k}-\mathbf{k}')$ (Definition 11.24 and Corollary 11.46).

2. Wightman kernels and $\bar{\Sigma}$-time ordering

Define the positive/negative-frequency Wightman kernels by vacuum contractions of (11.97):

$$G^{(a)>}_{\mu\nu}{}^{\rho\sigma}(x,y) := {}_{\Phi}\langle 0|\, \hat{h}^{(a)}_{\mu\nu}(x)\, \hat{h}^{(a)\,\rho\sigma}(y)\, |0\rangle_{\Phi}, \tag{11.98}$$

$$G^{(a)<}_{\mu\nu}{}^{\rho\sigma}(x,y) := {}_{\Phi}\langle 0|\, \hat{h}^{(a)}_{\mu\nu}(y)\, \hat{h}^{(a)\,\rho\sigma}(x)\, |0\rangle_{\Phi}. \tag{11.99}$$

Using the canonical algebra one obtains the standard mode sums,

$$\boxed{\begin{aligned} G^{(a)>}_{\mu\nu}{}^{\rho\sigma}(x,y) &= \sum_{\lambda=1}^{d_g} \int \frac{d^{d-1}\mathbf{k}}{(2\pi)^{d-1}} \frac{1}{2\omega_{\mathbf{k}}}\, e^{(a,\lambda)}_{\mu\nu}(k)\, e^{(a,\lambda)*\,\rho\sigma}(k)\, e^{-ik(x-y)}, \\ G^{(a)<}(x,y) &= G^{(a)>}(y,x). \end{aligned}} \tag{11.100}$$

The triad-resolved typed Feynman kernel then follows from $\bar{\Sigma}$-time ordering (Definition 11.53):

$$\boxed{G^{\mathrm{F},[\mathrm{TT}],(a)}_{\mu\nu}{}^{\rho\sigma}(x,y) = \Theta_{\bar{\Sigma}}(x \succeq y)\, G^{(a)>}_{\mu\nu}{}^{\rho\sigma}(x,y) + \Theta_{\bar{\Sigma}}(y \succeq x)\, G^{(a)<}_{\mu\nu}{}^{\rho\sigma}(x,y),} \tag{11.101}$$

where $\Theta_{\bar{\Sigma}}$ encodes the causal order in the $\bar{\Sigma}$-cone (Definition 11.53). Equal-time limits are taken by $\bar{\Sigma}$-covariant point-splitting compatible with the typed Peierls bracket, reproducing the typed ETCR on a Cauchy slice (eq. (11.25), Definition 11.8, and Proposition 11.4).

Remark 11.57 (What "foliation" means and why it matters)**.** An $\bar{\Sigma}$*–foliation* is a smooth time function $t : \mathcal{X}_q \to \mathbb{R}$ whose level sets $\Sigma_t := \{x \in \mathcal{X}_q \,|\, t(x) = \mathrm{const}\}$ are spacelike Cauchy hypersurfaces for the background Lorentzian structure $\bar{g}^{\mathrm{Lor}}$ (equivalently, for the causal cones of the unified background $\bar{\Sigma}$ used in this chapter). Writing n^μ for the future–pointing unit normal to Σ_t and $h_{\mu\nu}$ for the induced spatial projector (so that $h_{\mu\nu}n^\nu = 0$ and $\bar{\Sigma}_{\mu\nu} = -n_\mu n_\nu + h_{\mu\nu}$ in the usual signature convention), the foliation provides the "equal–time" notion required by the canonical construction. In Chapter 12 it serves four roles:

1. **Canonical (equal–time) quantization.** The typed equal–time commutation relations (ETCR) are imposed on Cauchy slices Σ_t selected by the foliation (the *band–regularized* canonical algebra in the TT carrier); see Definition 11.8 and Proposition 11.4 and its covariant counterpart via the typed Peierls bracket eq. (11.25). On the certified plateau $\mathfrak{P}$, Chapter 12 proves canonical $\leftrightarrow$ Peierls equivalence, so physical predictions do not depend on the particular admissible choice of t.

2. $\bar{\Sigma}$**–time ordering and propagators.** The foliation (equivalently, the $\bar{\Sigma}$ causal order) underlies the time–ordering operator $\Theta_{\bar{\Sigma}}$ used to assemble the TT Feynman kernel from the Wightman kernels:

$$G^{\mathrm{F},[\mathrm{TT}],(a)}(x,y) = \Theta_{\bar{\Sigma}}(x \succeq y)\, G^{(a)>}(x,y) + \Theta_{\bar{\Sigma}}(y \succeq x)\, G^{(a)<}(x,y),$$

cf. Definition 11.53 and eqs. (11.100) and (11.101). Equal–time limits are taken by $\bar{\Sigma}$–covariant point–splitting compatible with the typed Peierls bracket, reproducing the ETCR on Σ_t.

3. **Mode expansion and polarization basis.** Given a foliation, the TT graviton admits a mode expansion on Σ_t with frequencies $\omega_{\mathbf{k}}$ and TT polarization tensors $e^{(a,\lambda)}_{\mu\nu}(k)$, leading to the standard Wightman mode sums in eq. (11.100). The "$>/<$" kernels are the unordered vacuum two–point functions with positive/negative–frequency support relative to the chosen foliation/state, which the time–ordering then recombines.

4. **Compatibility with the weave/triad structure.** The foliation is required to be *admissible* (compatible with the unified cones of $\bar{\Sigma}$ and the band regularization). Within this admissible class, the TT projector $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ and the weave triad $\{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$ commute with the bookkeeping and do not depend on the particular slicings used to impose ETCR—hence the triad–resolved Fock factorization and diagnostics are foliation–independent on $\mathfrak{P}$ (see the TT quantization and triad factorization statements in Chapter 12).

In short, a $\bar{\Sigma}$–foliation supplies the "equal time" needed to state the canonical algebra and to define $\bar{\Sigma}$–time ordering, while the Peierls formulation guarantees that (on the certified plateau) observable TT physics does not depend on which admissible slicing one uses to implement those steps.

Remark 11.58 (Meaning of $G^{>}$, $G^{<}$ and $\bar{\Sigma}$–time ordering)**.** The kernels $G^{(a)>}$ and $G^{(a)<}$ in (11.100) are the *Wightman* two–point functions (also called "greater/lesser" functions): they are vacuum expectation values of the *unordered* operator products,

$$G^{(a)>}_{\mu\nu}{}^{\rho\sigma}(x,y) = {}_{\Phi}\langle 0|\, \hat{h}^{(a)}_{\mu\nu}(x)\, \hat{h}^{(a)\,\rho\sigma}(y)\, |0\rangle_{\Phi}, \qquad G^{(a)<}(x,y) = G^{(a)>}(y,x),$$

so that "$>$" places the operator at x to the left (positive–frequency part in the mode expansion), while "$<$" places it to the right (negative–frequency part) [255, 230, 82]. In curved backgrounds, the split into positive/negative frequency is defined relative to the chosen $\bar{\Sigma}$–foliation/state and is standard in the algebraic/Hadamard framework [245, 10, 13].

The *Feynman* (time–ordered) TT kernel in (11.101) is obtained by ordering with the $\bar{\Sigma}$–causal step function $\Theta_{\bar{\Sigma}}$ (cf. Definition 11.53), giving the familiar decomposition

$$G^{\mathrm{F},[\mathrm{TT}],(a)}(x,y) = \Theta_{\bar{\Sigma}}(x \succeq y)\, G^{(a)>}(x,y) + \Theta_{\bar{\Sigma}}(y \succeq x)\, G^{(a)<}(x,y),$$

exactly as in flat–space QFT with the usual Heaviside functions [82, Sec. 4]; in curved spacetime this formula is valid once a global time function (or, more generally, the causal order) associated with $\bar{\Sigma}$ is specified [245, 10, 13]. Equal–time limits are taken by $\bar{\Sigma}$–covariant point–splitting and reproduce the typed ETCR on a Cauchy slice (cf. eq. (11.25)),

Definition 11.8, and Proposition 11.4); distributionally, $G^{>} - G^{<}$ yields the Pauli–Jordan commutator function that implements the canonical algebra [255, 230, 384]. For the graviton TT polarization basis and the sum over helicities d_g used in (11.100), see standard constructions in gravitational–wave quantization [502].

3. Momentum-space / polarization-projector form

Fourier transforming (11.101) on a locally stationary patch (or in Riemann normal coordinates) yields the familiar momentum-space representation:

$$\boxed{G^{\mathrm{F},[\mathrm{TT}],(a)\ \rho\sigma}_{\mu\nu}(k) = \frac{i}{k^2 - m^2_{a,\mathrm{TT}} + i\epsilon}\, \Pi^{[\mathrm{TT}](a)\ \rho\sigma}_{\mu\nu}(k;\bar{\Sigma}),} \tag{11.102}$$

with the TT polarization projector assembled from the orthonormal basis, $\Pi^{[\mathrm{TT}](a)\ \rho\sigma}_{\mu\nu}(k) := \sum_\lambda e^{(a,\lambda)}_{\mu\nu}(k)\, e^{(a,\lambda)*\,\rho\sigma}(k)$ (Section 11.1.3). In the exact SSB/typed limit one has $m^2_{a,\mathrm{TT}} = 0$ with a protected massless pole; explicit breaking induces a GMOR-style shift $m^2_{a,\mathrm{TT}} = \mathcal{O}(\epsilon)$ (Proposition 11.29, Definition 11.33, and Corollary 11.30).

Remark 11.59 (KL positivity and band-stable causality)**.** On the physical Hilbert space $\mathcal{H}_{\mathrm{phys}}$ the sector-resolved TT two-point admits a Källén–Lehmann representation with non-negative residue and spectral density (eq. (11.91)); this *KL positivity* ensures that the polarization projector contributes positive weight to observable amplitudes. On the certified plateau $\mathfrak{P}$, band tags $P_x(\lambda_{\mathrm{UV}})$ and $K^{(\ell)}_{\mathrm{IR}}$ are chosen $\bar{\Sigma}$-covariantly (e.g. as functions of the TT d'Alembertian) and commute with $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ and $\Pi^{(a)}$ (Definitions 6.280, 9.10 and 11.24), so band-sandwiching preserves microcausal support and does not enlarge cones (Proposition 11.38).

Operator-basis viewpoint. Equations (11.100)–(11.102) show that $G^{\mathrm{F},[\mathrm{TT}]}$ is built from the same orthonormal operator basis $\{\hat{a}_{a,\lambda}(\mathbf{k}), \hat{a}^\dagger_{a,\lambda}(\mathbf{k})\}$ that spans the triad-resolved Fock space (Definition 11.48 and Corollary 11.46), thereby making explicit the link between the particle (creation/annihilation) and wave (Green function) descriptions ([82, 524]).

The Propagator as a Virtual Exchange Amplitude The triad-resolved, typed momentum-space form in eq. (11.102) is the central result for perturbative gravity in the IQR. Its interpretation is multi-layered:

- **Covariance.** As a TT-typed bi-kernel on $(\mathcal{X}_q, \bar{\Sigma})$, $G^{\mathrm{F},[\mathrm{TT}],(a)\ \rho\sigma}_{\mu\nu}(x,y)$ transforms as a rank-$(0,4)$ bitensor (a section of $\mathrm{TT}^2 \otimes \mathrm{TT}^2$) under background diffeomorphisms generated by $\bar{\Sigma}$ (Definitions 11.53 and 11.54). The momentum-space representation with $\Pi^{[\mathrm{TT}](a)\ \rho\sigma}_{\mu\nu}(k;\bar{\Sigma})$ is valid in a locally stationary patch (or Riemann normal coordinates) and is related to the position-space bitensor by the usual Fourier map.
- **Typed gauge/BRST invariance.** The explicit TT projector ensures that only the physical, transverse-traceless spin-2 polarizations propagate; pure-gauge components are removed at the outset. On the certified plateau $\mathfrak{P}$, the kernel is BRST-closed and gauge-parameter independent (Nielsen-flat) and thus constitutes the correct physical

propagator for perturbation theory (Section 11.1.2 and Definitions 6.280 and 9.10; cf. [155, 245, 403, 502]).

- **KL positivity and SSB protection.** On $\mathcal{H}_{\mathrm{phys}}$ the TT two-point function admits a Källén–Lehmann representation with non-negative residues and spectral densities (eq. (11.91)), so the spectral weight is positive in each sector. In the exact SSB limit the typed Ward identity protects the massless TT pole, while small explicit breaking induces the GMOR-type shift $m^2_{a,\mathrm{TT}} = \mathcal{O}(\epsilon)$ (Proposition 11.29, Definition 11.33, and Corollary 11.30).

- **Virtual graviton exchange.** Physically, $G^{\mathrm{F},[\mathrm{TT}],(a)}$ is the amplitude for *virtual* TT spin-2 exchange in sector a. In the Dyson expansion, every internal wavy line is a TT-projected triadic propagator, and it connects typed interaction vertices obtained by reducing the Chapter 11 bi-local parent coupling to the linearized form in this chapter (Sections 11.3 and 11.3.3). This is the standard field-theoretic picture of exchange, here implemented with the $\bar{\Sigma}$-time ordering and typed projectors ([82, 403, 502]).

11.2.3 Graviton Propagator in Background Geometry

On the emergent chronogeometric background $(\mathcal{X}_q, \bar{\Sigma}_{\mu\nu})$, the TT-typed graviton $\hat{h}^{(a)}_{\mu\nu}(x)$ (sector $a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$) propagates according to a *project-then-solve* prescription: first apply the background-covariant TT projector $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ (York-type, Chapter 11), then define all dynamical operators and Green kernels on the TT image (Sections 11.1.2 and 11.1.3 and Definition 11.28).

TT Lichnerowicz-type wave operator.

Definition 11.60 (TT Lichnerowicz–type wave operator (sector–resolved))**.** Let $\mathbb{L}^{(2)}_{\bar{\Sigma}}$ denote the spin–2 Lichnerowicz–type operator acting on symmetric rank–2 tensors over $(\mathcal{X}_q, \bar{\Sigma})$, with principal part $-\bar{\nabla}^\lambda\bar{\nabla}_\lambda\,\delta^\mu{}_\alpha\delta^\nu{}_\beta$ and curvature couplings built from $\bar{R}_{\mu\nu\rho\sigma}[\bar{\Sigma}]$. For each triad sector $a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$ (Definition 10.2), the *typed, sector–resolved* wave operator is the TT–sandwiched operator

$$\boxed{K^{[\mathrm{TT}],(a)\alpha\beta}_{\mu\nu}(x) := \Big[\,\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}} \circ \Big(\,\mathbb{L}^{(2)}_{\bar{\Sigma}} - \mathcal{M}^{(a)}{}_{\cdot\cdot}{}^{\cdot\cdot}(x)\,\Big) \circ \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\Big]_{\mu\nu}{}^{\alpha\beta},} \tag{11.103}$$

where $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ is the background–covariant York–Lichnerowicz TT projector (eq. (10.4) and Definition 10.3) and $\mathcal{M}^{(a)}$ is the sectoral mass/mixing tensor projected to the TT block (Definition 11.33 and Proposition 11.32). The following properties hold on a certified plateau $\mathfrak{P}$:

1. **Typed equations of motion.** The typed graviton carrier $\hat{h}^{(a)}_{\mu\nu} := \Pi^{(a)}\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{g}_{\mu\nu}$ (Definition 11.3 and Proposition 10.9) obeys $K^{[\mathrm{TT}],(a)\alpha\beta}_{\mu\nu}\,\hat{h}^{(a)}_{\alpha\beta} = 0$ (up to sources), which fixes the normalization and dynamics sector–by–sector.

2. **Green operators and typed propagator.** The retarded/advanced Green operators $G^{\mathrm{ret/adv},[\mathrm{TT}],(a)}$ solve the Green equations for $K^{[\mathrm{TT}],(a)}$, and the typed Feynman propagator $G^{\mathrm{F},[\mathrm{TT}],(a)}$ is the corresponding Feynman Green function (Definition 11.54). The Pauli–Jordan kernel $\Delta^{[\mathrm{TT}]} := G^{\mathrm{ret}}_{[\mathrm{TT}]} - G^{\mathrm{adv}}_{[\mathrm{TT}]}$ determines the covariant commutator and the Peierls bracket (eqs. (11.25) and (11.26)).
3. **Hyperbolicity and microcausality.** The principal symbol of $K^{[\mathrm{TT}],(a)}$ is inherited from $\mathbb{L}^{(2)}_{\bar\Sigma}$ and is $\bar\Sigma$–hyperbolic, so $\operatorname{supp} G^{\mathrm{ret/adv},[\mathrm{TT}],(a)}$ lies on the $\bar\Sigma$–causal cones; hence $\operatorname{supp}\Delta^{[\mathrm{TT}]} \subset J^+_{\bar\Sigma} \cup J^-_{\bar\Sigma}$, which implies typed microcausality for the Peierls bracket (Proposition 11.14 and Remark 9.83).
4. **Compatibility with the realized weave configuration (plateau safety).** On $\mathfrak{P}$ the TT and triad projectors commute with the band tags and are BRST–closed; therefore $K^{[\mathrm{TT}],(a)}$ is regulator–safe and gauge–parameter independent (Nielsen–flat) in all constructions that use it—propagators, Peierls bracket, and diagnostics (Definitions 6.280 and 9.10, Propositions 7.3 and 10.9, and Corollary 11.5).

Central role of the typed Lichnerowicz–type operator. The wave operator $K^{[\mathrm{TT}],(a)\alpha\beta}_{\mu\nu}$ defined by the TT sandwiching of the spin–2 Lichnerowicz operator together with the sectoral mass/mixing tensor (cf. eq. (11.103)) is the kinetic operator for the typed graviton in each triad sector $a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$ (Definitions 10.2 and 11.3). After TT projection, the carrier $\hat h^{(a)}_{\mu\nu}$ obeys the typed field equation $K^{[\mathrm{TT}],(a)\alpha\beta}_{\mu\nu}\,\hat h^{(a)}_{\alpha\beta} = 0$, and the *typed Feynman propagator* $G^{\mathrm{F},[\mathrm{TT}],(a)}{}_{\mu\nu}{}^{\rho\sigma}$ is by definition the Green's function of the same $K^{[\mathrm{TT}],(a)}$ (see Definition 11.54). This identification fixes the normalization of modes, determines the on–shell content sector–by–sector, and ensures that the propagator is strictly TT, i.e. it maps the TT subspace into itself and vanishes on its orthogonal complement (use Definition 11.18 and eq. (11.33)).

The same operator generates the *covariant* algebra via the Peierls construction. The typed Peierls bracket

$$\{F,G\}_{\mathrm{P}} = \Big\langle \frac{\delta F}{\delta h_{\mu\nu}},\ \Delta^{[\mathrm{TT}]\,\mu\nu}{}_{\rho\sigma}\ \frac{\delta G}{\delta h_{\rho\sigma}} \Big\rangle, \qquad \Delta^{[\mathrm{TT}]} := G^{\mathrm{ret}}_{[\mathrm{TT}]} - G^{\mathrm{adv}}_{[\mathrm{TT}]},$$

(eq. (11.25)) is built from the retarded/advanced Green operators of the very same typed wave operator $K_{[\mathrm{TT}]}$. Consequently, $K_{[\mathrm{TT}]}$ (ultimately sourced by $\mathbb{L}^{(2)}_{\bar\Sigma}$) controls the covariant commutator $[\,\hat h(x), \hat h(x')\,] = i\,\Delta^{[\mathrm{TT}]}(x,x')$ and, upon restriction to a Cauchy slice, reproduces the equal–time algebra of the typed canonical pair (eq. (11.26) and Proposition 11.4).

Typed microcausality follows from the $\bar\Sigma$–hyperbolic nature of $K_{[\mathrm{TT}]}$: the retarded/advanced TT Green operators are supported on the $\bar\Sigma$–causal cones, whence the Pauli–Jordan kernel $\Delta^{[\mathrm{TT}]}$ has support only on $J^{\pm}_{\bar\Sigma}$ and the Peierls bracket vanishes for $\bar\Sigma$–spacelike separations (Proposition 11.14). The IR window $K^{(\ell)}_{\mathrm{IR}}$ is cone–preserving, and the band realized weave configuration commutes with the TT projector, so these causal properties are regulator–safe on a certified plateau $\mathfrak{P}$ (Definition 6.280, Remark 9.83, and Proposition 10.9).

In momentum space the typed propagator is the inverse of $K_{[\mathrm{TT}]}$ on the TT subspace and projects onto TT polarizations, carrying in each sector the typed mass parameter $m^2_{a,\mathrm{TT}}$. In the spontaneously broken limit, the TT Goldstone direction forces $m^2_{a,\mathrm{TT}} = 0$ and yields a

massless pole with positive residue (Proposition 11.29); a small explicit breaking shifts the pole according to the GMOR–style estimate $m_{\mathrm{TT}}^2 = \mathcal{O}(\epsilon)$ with a plateau bound (Corollary 11.30). These spectral statements are read directly from the structure of $K_{[\mathrm{TT}]}$ and its TT inverse (cf. the triad–resolved Källén–Lehmann form in Definition 11.21).

Finally, the operator is intrinsically compatible with the typed realized weave configuration: it is sandwiched by the commuting TT and triad projectors and respects the band/tag hygiene on $\mathfrak{P}$, so covariance, causality, and gauge–parameter independence (Nielsen flatness) propagate to all derived objects—propagator, Peierls bracket, and law/state diagnostics (Definition 9.10, Propositions 7.3 and 10.9, and Corollary 11.5). This is why tree–level exchange amplitudes are expressed as contractions of typed vertices with the typed propagator—the Green's function of $K^{[\mathrm{TT}]}$—so scattering and linear response depend precisely on the Lichnerowicz–type operator through its TT projection (Definitions 11.54 and 11.67).

Typed linear dynamics (project then solve).

Definition 11.61 (Typed TT equation of motion (project then solve))**.** Let $K_{\mu\nu}^{[\mathrm{TT}],(a)\alpha\beta}(x)$ be the sector–resolved TT wave operator of Definition 11.60 and let $\hat{h}_{\mu\nu}^{(a)} := \Pi^{(a)}\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}_{\mu\nu}$ be the typed graviton carrier in triad sector $a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$ (Definitions 10.2 and 11.3). The *typed TT equation of motion* is

$$\boxed{K_{\mu\nu}^{[\mathrm{TT}],(a)\alpha\beta}(x)\,\hat{h}_{\alpha\beta}^{(a)}(x) = \hat{J}_{\mu\nu}^{[\mathrm{TT}],(a)}(x), \qquad \hat{J}^{[\mathrm{TT}],(a)} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\hat{J}^{(a)},} \tag{11.104}$$

where $\hat{J}^{(a)} := \Pi^{(a)}\hat{J}$ is the triad–filtered source induced by the bi–local matter–geometry interaction (Section 11.3 and Definition 11.24). On a certified plateau $\mathfrak{P}$, the realized weave configuration commutations imply $\Pi^{(a)}\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\Pi^{(a)}$ (Proposition 10.9), so either ordering yields the same typed source.

Remark 11.62. (i) Retarded/advanced Green operators of $K^{[\mathrm{TT}],(a)}$ produce covariant solutions with cone support, and their causal difference $\Delta^{[\mathrm{TT}]}$ generates the Peierls bracket and covariant commutator (Definition 11.60, Proposition 11.14, and eqs. (11.25) and (11.26)). (ii) The Feynman Green function of $K^{[\mathrm{TT}],(a)}$ is the typed propagator that enters tree–level exchange amplitudes (Definitions 11.54 and 11.67). (iii) Formulating (11.104) inside the TT, triad–resolved subspace enforces purity and prevents leakage into non–TT or off–sector channels (cf. Definition 11.7).

Explanation (typed linear dynamics: project then solve). The operator $K_{\mu\nu}^{[\mathrm{TT}],(a)\alpha\beta}(x)$ is the sector–resolved, TT–sandwiched Lichnerowicz–type wave operator of Definition 11.60. The typed graviton carrier in sector $a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$ is $\hat{h}_{\mu\nu}^{(a)} := \Pi^{(a)}\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}_{\mu\nu}$ (Definitions 10.2 and 11.3), where the commuting property of the realized weave configuration on the certified plateau $\mathfrak{P}$ ensures $\Pi^{(a)}\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\Pi^{(a)}$ (Proposition 10.9). The typed source $\hat{J}^{[\mathrm{TT}],(a)} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\hat{J}^{(a)}$ is obtained by projecting the triad–filtered interaction current $\hat{J}^{(a)}$ that arises from the bi–local matter–geometry coupling in the interaction action (Section 11.3 and Definition 11.24). Writing the equation of motion,

$$K_{\mu\nu}^{[\mathrm{TT}],(a)\alpha\beta}(x)\,\hat{h}_{\alpha\beta}^{(a)}(x) = \hat{J}_{\mu\nu}^{[\mathrm{TT}],(a)}(x),$$

implements the principle "project, then solve": the dynamics is enforced *within* the TT, sector–resolved subspace, so no longitudinal/trace components or cross–triad leaks can appear in the solution (cf. TT purity/leak diagnostics in Definition 11.7).

Green representation, causality, and algebra. Let ${G^{\mathrm{ret/adv},[\mathrm{TT}],(a)}}_{\mu\nu}{}^{\rho\sigma}(x,y)$ be the retarded/advanced Green operators of $K^{[\mathrm{TT}],(a)}$ ($\bar{\Sigma}$–hyperbolic; Definition 11.60). The causal support statement $\operatorname{supp} G^{\mathrm{ret/adv},[\mathrm{TT}]} \subset J^{\pm}_{\bar{\Sigma}}$ implies the retarded solution

$$\hat{h}^{(a)}_{\mu\nu}(x) = \int d\mu_{\bar{\Sigma}}(y)\; {G^{\mathrm{ret},[\mathrm{TT}],(a)}}_{\mu\nu}{}^{\rho\sigma}(x,y)\; \hat{J}^{[\mathrm{TT}],(a)}_{\rho\sigma}(y),$$

which is manifestly covariant and microcausal (Proposition 11.14). The Feynman propagator $G^{\mathrm{F},[\mathrm{TT}],(a)}$ used in time–ordered exchange amplitudes is the corresponding Feynman Green function of $K^{[\mathrm{TT}],(a)}$ (Definitions 11.54 and 11.67). The Pauli–Jordan kernel $\Delta^{[\mathrm{TT}]} := G^{\mathrm{ret}}_{[\mathrm{TT}]} - G^{\mathrm{adv}}_{[\mathrm{TT}]}$ generated by the same operator controls the Peierls bracket and the covariant commutator, whose equal–time limit reproduces the typed canonical algebra (eqs. (11.25) and (11.26) and Proposition 11.4). Gauge–parameter independence of these statements on $\mathfrak{P}$ follows from BRST/Nielsen safety (Proposition 7.3 and Corollary 11.5), and regulator stability from cone–preserving band tags (Definition 6.280 and Remark 9.83).

Typed Green kernels and causal support. Define the retarded/advanced TT Green kernels as the bi-tensors $G^{\mathrm{ret/adv},[\mathrm{TT}],(a)}_{\mu\nu}{}^{\rho\sigma}(x,y)$ solving

$$\boxed{K^{[\mathrm{TT}],(a)}_{\mu\nu}{}^{\alpha\beta}(x)\, G^{\mathrm{ret/adv},[\mathrm{TT}],(a)}_{\alpha\beta}{}^{\rho\sigma}(x,y) = \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}\,\mu\nu}{}^{\rho\sigma}\,\delta^{(d)}(x-y),} \tag{11.105}$$

with supports $\operatorname{supp} G^{\mathrm{ret}} \subset J^{+}_{\bar{\Sigma}}(y)$ and $\operatorname{supp} G^{\mathrm{adv}} \subset J^{-}_{\bar{\Sigma}}(y)$. The causal (Pauli–Jordan) kernel and the typed Feynman kernel are then

$$\Delta^{[\mathrm{TT}],(a)} := G^{\mathrm{ret},[\mathrm{TT}],(a)} - G^{\mathrm{adv},[\mathrm{TT}],(a)}, \qquad \operatorname{supp}\Delta^{[\mathrm{TT}],(a)} \subset J^{+}_{\bar{\Sigma}}(y) \cup J^{-}_{\bar{\Sigma}}(y), \tag{11.106}$$

$$i\, G^{\mathrm{F},[\mathrm{TT}],(a)}_{\mu\nu}{}^{\rho\sigma}(x,y) = {}_{\Phi}\langle 0|\, T_{\bar{\Sigma}}\Big\{\hat{h}^{(a)}_{\mu\nu}(x)\hat{h}^{(a)\,\rho\sigma}(y)\Big\}\,|0\rangle_{\Phi}, \quad \text{with } K^{[\mathrm{TT}],(a)} \circ G^{\mathrm{F},[\mathrm{TT}],(a)} = i\,\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\delta^{(d)}(\cdot), \tag{11.107}$$

(Definitions 11.53 and 11.54, eq. (11.102), and Proposition 11.14).

Composition convention. For bi-tensors A, B of type $(0,4)$ we write

$$(A\circ B)_{\mu\nu}{}^{\rho\sigma}(x,y) := \int_{\mathcal{X}_q} d\mu_{\bar{\Sigma}}(z)\, A_{\mu\nu}{}^{\alpha\beta}(x,z)\, B_{\alpha\beta}{}^{\rho\sigma}(z,y),$$

with $d\mu_{\bar{\Sigma}}$ the background-covariant measure.

Hyperbolicity and Nielsen flatness (plateau tests). The principal symbol of $K^{[\mathrm{TT}],(a)}$ is $\sigma\big(K^{[\mathrm{TT}],(a)}\big)(x,\xi) = \bar{\Sigma}^{\lambda\kappa}(x)\,\xi_\lambda\xi_\kappa\,\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}(x)$, so $K^{[\mathrm{TT}],(a)}$ is normally hyperbolic on $(\mathcal{X}_q,\bar{\Sigma})$; hence retarded/advanced solutions exist and have $\bar{\Sigma}$-causal support. On the certified plateau $\mathfrak{P}$, observables built from $G^{\mathrm{ret/adv/F},[\mathrm{TT}],(a)}$ are gauge-parameter independent (*Nielsen-flat*) and stable under admissible band deformations; band tags commute with $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ and do not enlarge causal cones (Definitions 6.280 and 9.10 and Proposition 11.38).

Remark 11.63 (Equivalence with mode-sum representations)**.** In a locally stationary patch (or via covariant Fourier–Hadamard techniques), the kernels in (11.107) coincide with the triad-resolved mode sums and the momentum-space projector form (Section 11.2.2 and eq. (11.102)), ensuring both manifest locality (Green-function picture) and spectral completeness (operator picture) within the IQR formalism.

11.2.4 Triadic Structure and Interpretation of the Graviton Propagator

By construction, the free graviton two-point function is *typed* (TT-projected) and *triad-resolved*. The full kernel is the orthogonal sum of sectoral contributions (Definition 11.51 and eq. (11.86)):

$$\boxed{G^{\mathrm{grav}}_{\mu\nu}{}^{\rho\sigma}(x,x') = \sum_{a\in\{\mathrm{corr,vac,spec}\}} G^{(a)}_{\mu\nu}{}^{\rho\sigma}(x,x'), \qquad [\Pi^{(a)},\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}] = 0,} \tag{11.108}$$

so that, at the free level on the certified plateau $\mathfrak{P}$, the internal lines of Feynman graphs are block-diagonal by sector and TT-pure (Section 11.1.4 and Definition 11.54). This makes explicit that the apparent single "gravitational interaction" is, in IQR, a composite phenomenon mediated by three virtual spin-2 channels with distinct relational origins.

The Three Channels of Gravitational Interaction

Correlational channel: geometry from quantum information

The correlational propagator $G^{(\mathrm{corr})}$ captures TT fluctuations sourced by the information-theoretic structure of the state—coherence and entanglement—in the sense of the correlational distance d_{corr} (Section 4.9.1 and Definition 11.24). Its field operator $\hat{h}^{(\mathrm{corr})}_{\mu\nu}$ is the TT projection of the correlational component of the Gaiamiti fluctuation (Definition 11.42 and eq. (11.74)). As a mediator, this channel is sensitive to gradients of modular/entanglement data rather than classical stress–energy, providing a dynamical avatar of the "information builds geometry" motif ([520, 1]; cf. Section 11.1.3 for the TT basis).

Vacuum channel: geometry from symmetry breaking

The vacuum propagator $G^{(\mathrm{vac})}$ describes TT excitations tied to the response of the vacuum manifold $\mathcal{M}_{\mathrm{vac}}$ (Section 6.1.8). In the exact SSB limit, the typed Ward identity enforces a protected massless TT pole in this channel (Proposition 11.29); small explicit breaking produces a

GMOR-type shift consistent with the TT-projected mass operator (Definition 11.33 and Corollary 11.30). The corresponding field, $\hat{h}^{(\mathrm{vac})}_{\mu\nu}$, is defined in Definition 11.43 and eq. (11.75) and mediates long-range geometric responses to shifts within $\mathcal{M}_{\mathrm{vac}}$.

Spectral channel: geometry from energy–momentum

The spectral propagator $G^{(\mathrm{spec})}$ is driven by local spectral content (energetic/momentum scales) as encoded by the spectral distance d_{spec} (Section 4.9.6) and the sectoral source defined by the interaction kernel (Section 10.19.6 and Definition 11.44). In the semiclassical regime, $G^{(\mathrm{spec})}$ reduces to the familiar helicity-2 kernel of linearized GR with TT polarization projector (Section 11.2.2 and eq. (11.102); [245, 502]), thus governing the classical phenomena sourced by stress–energy.

Remarks. (i) Each $G^{(a)}$ is defined by first projecting the Lichnerowicz-type operator onto the TT subspace and then solving for the sectoral Green kernel (Section 11.2.3 and eqs. (11.103) and (11.107)). (ii) KL positivity holds on the physical Hilbert space for each sectoral two-point (eq. (11.91)); Nielsen flatness and hyperbolicity are plateau tests ensuring gauge-parameter independence and causal support (Proposition 11.14). (iii) Beyond tree level, radiative corrections can induce sectoral self-energies Σ_{ab} consistent with the typed projectors; the resulting dispersion/positivity bounds are assessed triadwise (Section 11.3.6).

11.2.5 A Composite Gravitational Force and its Predictions

In the IQR framework the *typed, triad-resolved* propagators add coherently at the free level (eq. (11.108)), so the net gravitational interaction is a superposition of three TT spin–2 exchange channels with distinct relational origins (correlational, vacuum, spectral; cf. Section 11.1.4 and Definition 11.51). Beyond tree level, sectoral self-energies Σ_{ab} compatible with the typed projectors may induce controlled mixing among channels (triadic transmutation), still certified on the plateau $\mathfrak{P}$ by TT law/state visibilities ($\Lambda^{[\mathrm{TT}]}_{\mathrm{law}}, \Lambda^{[\mathrm{TT}]}_{\mathrm{state}}$; Definition 11.52 and Proposition 11.14).

Unique predictions of the triadic model. The novelty arises from the correlational and vacuum channels, absent in single-metric treatments ([156, 10]). In addition to the familiar *spectral* (stress–energy) graviton, the theory predicts:

- *Correlational gravitons* $\hat{h}^{(\mathrm{corr})}_{\mu\nu}$ mediating geometry sourced by quantum informational structure (coherence/entanglement), in line with the "information builds geometry" paradigm (Definition 11.42; [1, 525, 520]).
- *Vacuum gravitons* $\hat{h}^{(\mathrm{vac})}_{\mu\nu}$ tied to the structure of the vacuum manifold $\mathcal{M}_{\mathrm{vac}}$ and its spontaneously broken symmetries; in the exact SSB limit they exhibit a protected massless TT pole, with a GMOR-type shift under small explicit breaking (Proposition 11.29 and Definition 11.33; [526]).

Consequently, the framework admits *gravity without traditional stress–energy sources*: highly entangled yet low-energy configurations can source correlational curvature, and vacuum

rearrangements can source TT waves during phase transitions (Section 11.1.4; [1, 525]). This challenges the strong reading of universality in GR while remaining testable against precision constraints ([479]).

Falsifiable aspects and observational handles. The triadic structure leads to concrete, potentially observable deviations from GR that can be certified on $\mathfrak{P}$ via TT visibilities and positivity/dispersion checks (Section 11.3.6):

1. **Apparent non-universality of Newton's constant.** Because exchange amplitudes add channelwise (eq. (11.108)), the effective coupling measured between two sources can depend on their triadic composition. Denote by G_{eff} the empirically inferred coupling in a given setup; then G_{eff} is a functional of the sector weights (the TT KL residue vector $\vec{Z}$, eq. (11.91)) and may differ for sources with equal stress–energy but different coherence/vacuum structure. A targeted null test compares the attraction between two samples with matched mass–energy but sharply different internal quantum coherence (highly entangled vs. decohered), while controlling for environmental decoherence and vacuum alignment ([527, 528]; cf. Definition 11.52).

2. **Directional or frequency response tied to chronogeometry.** Propagation is governed by the TT Lichnerowicz-type operator built from $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (Section 11.2.3 and eq. (11.103)). In regions where $\bar{t}_{\mu\nu}$ is sizable, the *chronogeometric* cone can differ from that inferred by probes coupling only to $\bar{g}_{\mu\nu}$, opening the door to apparent anisotropy/dispersion of TT waves relative to standard matter clocks. Precision bounds on Lorentz-violating dispersion therefore constrain the allowed $\bar{t}$-backdrop and sectoral self-energies ([529, 514]); any confirmed direction- or frequency-dependence beyond GR expectations would be evidence for the chronogeometric structure posited here.

3. **Primordial imprints in cosmology.** During high-coherence, high-energy epochs, correlational and vacuum channels can be non-negligible, modifying the tensor power spectrum, its polarization, and the stochastic GW background. One expects non-standard tensor consistency relations or mixed-polarization signatures seeded by triadic mixing and SSB-protected modes (Section 11.1.4 and Proposition 11.29). CMB B-modes and direct searches for stochastic backgrounds thus provide incisive tests ([439, 530, 512]).

Certification protocol. For any proposed deviation, the TT law visibility $\Lambda^{[\mathrm{TT}]}_{\text{law}}$ and state visibility $\Lambda^{[\mathrm{TT}]}_{\text{state}}$ must be near unity on a certified plateau (Definition 11.52), KL positivity must hold sectorwise (eq. (11.91)), and microcausality with respect to the $\bar{\Sigma}$-cone must be maintained (Proposition 11.14 and Definition 11.53). This ensures that any phenomenological inference is *typed, causal, and gauge-robust* before being confronted with data.

11.3 Graviton-Mediated Scattering

With the typed, TT-projected graviton field $\hat{h}_{\mu\nu} := \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\, \delta\hat{g}_{\mu\nu}$ (Section 11.1.3) and the triad-resolved propagators in hand (Definitions 11.51 and 11.54, eq. (11.86), and Section 11.2.2), we

now formulate graviton–mediated scattering on the certified plateau $\mathfrak{P}$ (band tags commuting with $\Pi^{(a)}$ and $\Pi^{[\mathrm{TT}]}_{\bar\Sigma}$; Definitions 6.280 and 9.10 and Proposition 11.14). All couplings below are understood as the TT-typed *shadow* of the bi-local matter–geometry interaction introduced in Chapter 11 (the $\Phi\Phi \leftrightarrow \Theta\Theta$ block), and inherit Hermiticity, $\bar\Sigma$-cone support, and TT-preservation by the weave algebra and the TT-nullity of the τ-typed leg (Section 11.1.3).

Definition 11.64 (Graviton–Tantrakyma vertex operator (typed, triad-resolved))**.** Let $S_{\mathrm{int}}[\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}]$ denote the bi-local parent interaction (Chapter 11). Its TT-typed, linearized reduction around $(\mathcal{X}_q, \bar\Sigma)$ is defined by

$$\boxed{\hat V^{\mu\nu}(x) := \left(\Pi^{[\mathrm{TT}]}_{\bar\Sigma}\right)^{\mu\nu}{}_{\alpha\beta} \left.\frac{\delta S_{\mathrm{int}}[\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}]}{\delta \hat h_{\alpha\beta}(x)}\right|_{\mathfrak{P}}, \qquad \hat V^{\mu\nu} = \hat V^{\nu\mu}, \quad \bar\Sigma_{\mu\nu}\hat V^{\mu\nu} = 0.} \tag{11.109}$$

and decomposes orthogonally into the triadic channels $\hat V^{\mu\nu} = \sum_{a\in\{\mathrm{corr,vac,spec}\}} \hat V^{(a)\mu\nu}$ with $\hat V^{(a)\mu\nu} := \Pi^{(a)}\hat V^{\mu\nu}$ and $[\Pi^{(a)}, \Pi^{[\mathrm{TT}]}_{\bar\Sigma}] = 0$ (Definition 11.24).

Remark 11.65 (Decomposition and practical reading of eq. (11.109))**. What each piece means.**

- $\left(\Pi^{[\mathrm{TT}]}_{\bar\Sigma}\right)^{\mu\nu}{}_{\alpha\beta}$. This is the *background–covariant transverse–traceless (TT) projector* for rank-2 symmetric tensors on the certified plateau determined by $\bar\Sigma_{\mu\nu}$ (see the TT calculus from Ch. 11). As a map it sends a two–tensor $X^{\alpha\beta}$ to its TT part,

  $$X^{\mu\nu}_{[\mathrm{TT}]} = \left(\Pi^{[\mathrm{TT}]}_{\bar\Sigma}\right)^{\mu\nu}{}_{\alpha\beta} X^{\alpha\beta},$$

  enforcing symmetry, tracelessness w.r.t. $\bar\Sigma$, and—when inserted in the dynamics—transversality w.r.t. $\bar\nabla$.
- $\dfrac{\delta S_{\mathrm{int}}[\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}]}{\delta \hat h_{\alpha\beta}(x)}$. The *functional derivative* of the (bi–local) interaction action with respect to the graviton field $\hat h_{\alpha\beta}(x)$. Before typing, this is the "raw'' vertex tensor with two free indices $(\alpha\beta)$.
- $\big|_{\mathfrak{P}}$. *Evaluation on the certified plateau* $\mathfrak{P}$ (Ch. 8): band tags are fixed, BRST cohomology has been taken, and all quantities are *Nielsen–flat* (gauge–parameter independent). Operationally: take the derivative, then restrict to the physical, regulator–clean arena.
- *Left–hand side* $\hat V^{\mu\nu}(x)$. The *typed (TT) vertex operator* that couples to the typed graviton $\hat h_{\mu\nu}$. It is *by construction*

  $$\hat V^{\mu\nu} = \hat V^{\nu\mu}, \qquad \bar\Sigma_{\mu\nu}\hat V^{\mu\nu} = 0,$$

  i.e. symmetric and $\bar\Sigma$–traceless (TT). Transversality is then enforced via the TT Ward identity in the covariant Peierls framework; see Section 11.1.2 and the vertex discussion in Section 11.3.2.

Why we project. We *project after differentiating* so that only the *physical*, TT–propagating part of the vertex survives. This aligns the right leg of the vertex with the *typed graviton carrier*

$$\hat{h}_{\mu\nu} = \left(\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\right)_{\mu\nu}{}^{\rho\sigma}\, \delta\hat{g}_{\rho\sigma} \quad \text{(cf. Section 11.1.3)},$$

guarantees TT–purity on all graviton lines, and makes TT Ward checks straightforward on $\mathfrak{P}$.

Common pitfalls (and the tiny typo).

- The argument of $S_{\mathrm{int}}[\cdots]$ must be closed: use $S_{\mathrm{int}}[\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}]$.
- Index placement matters: $\left(\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\right)^{\mu\nu}{}_{\alpha\beta}$ maps lower–index $\hat{h}_{\alpha\beta}$ to upper–index $\hat{V}^{\mu\nu}$; raise/lower with $\bar{\Sigma}$.

Equivalent "expanded" reading. For a compact, index–free view, read

$$\hat{V}_{[\mathrm{TT}]} \;=\; \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\left(\frac{\delta S_{\mathrm{int}}}{\delta \hat{h}}\right)\Bigg|_{\mathfrak{P}},$$

i.e. "*take the vertex = TT–project (the functional derivative of the interaction) on the plateau.*" This is precisely the typed bi–local → linearized reduction in Section 11.3.2, and it is the object that contracts with TT propagators in amplitudes (Section 11.3.7).

11.3.1 Linearized Interaction Action

The typed, linear coupling of the TT graviton to Tantrakyma degrees of freedom is

Definition 11.66 (Graviton–Tantrakyma interaction (typed, linearized))**.** To first order in $\hat{h}$, the interaction action on $\mathfrak{P}$ reads

$$\boxed{S_{\mathrm{int}}^{(1)}[\hat{h}, \hat{\mathbf{\Theta}}] := \int d\mu_{\bar{\Sigma}}(x)\ \mathrm{Tr}\Big[\hat{h}_{\mu\nu}(x)\, \hat{V}^{\mu\nu}(x)\Big] = \sum_a \int d\mu_{\bar{\Sigma}}(x)\ \mathrm{Tr}\Big[\hat{h}^{(a)}_{\mu\nu}(x)\, \hat{V}^{(a)\mu\nu}(x)\Big]\,,} \tag{11.110}$$

with $d\mu_{\bar{\Sigma}}(x)$ the covariant volume density, and "Tr" the fiber trace over the local operator bundle (Section 11.1.3).

11.3.2 Structure of the Vertex from the Interaction Kernel

Let $\mathcal{U}_{\Phi\Theta}(x, x')$ be the bi-local interaction kernel of Chapter 11 and $\mathcal{U}^{(a)}_{\Phi\Theta}$ its triadic components (Section 10.19.6). The typed vertex admits the representation

$$\boxed{\hat{V}^{(a)\mu\nu}(x) = \Pi^{[\mathrm{TT}]\mu\nu}{}_{\alpha\beta} \int d\mu_{\bar{\Sigma}}(x')\, \hat{\mathcal{K}}^{(a)\alpha\beta}{}_{A}(x, x')\, \delta\hat{\mathbf{\Theta}}^{A}(x'), \qquad a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\},}$$

(11.111)

for some operator-valued kernel $\hat{\mathcal{K}}^{(a)}$ determined by the variation of the parent action. Hermiticity and $\bar{\Sigma}$-causal support follow from the parent bilinear structure; TT-preservation follows from the projector and the TT-nullity of $\hat{\tau}$ (Section 11.1.3).

11.3.3 Graviton Coupling Channels (modal decomposition)

Expanding on the TT polarization basis $e^{(s,a)}_{\mu\nu}(k)$ (Section 11.1.3), the vertex admits a modal decomposition

$$\boxed{\hat{V}^{(a)\mu\nu}(x) = \sum_{s=1}^{d_g} \int \frac{d^{d-1}\mathbf{k}}{(2\pi)^{d-1}} \Big[\hat{\mathcal{J}}^{(s,a)}(\mathbf{k};x)\, e^{(s,a)\mu\nu}(k)\, e^{-ikx} + \text{h.c.} \Big], \quad [\hat{\mathcal{J}}^{(s,a)}, \hat{\mathcal{J}}^{(s',b)\dagger}] \propto \delta^{ss'}\delta^{ab},} \tag{11.112}$$

so that sector labels commute and the triadic Fock spaces factorize (Sections 11.1.4 and 11.1.5).

11.3.4 Scattering Amplitudes from Graviton Exchange

Setup and external states. Consider $2 \to 2$ scattering of Tantrakyma excitations with external one–particle states $|\Theta_1(p_1)\rangle, |\Theta_2(p_2)\rangle \to |\Theta_3(p_3)\rangle, |\Theta_4(p_4)\rangle$, LSZ–amputated and normalized in the standard relativistic convention. We write expectation values in these external states as matrix elements, e.g. $\langle \hat{V}^{(a)}_{\mu\nu}(x)\rangle := \langle \Theta_3|\hat{V}^{(a)}_{\mu\nu}(x)|\Theta_1\rangle$ and $\langle \hat{V}^{(a)\rho\sigma}(y)\rangle := \langle \Theta_4|\hat{V}^{(a)\rho\sigma}(y)|\Theta_2\rangle$. Integration uses the $\bar{\Sigma}$–invariant measure $d\mu_{\bar{\Sigma}}(x)$ on $\mathcal{X}_q$. Time ordering is with respect to the $\bar{\Sigma}$ causal structure (Section 11.2.1).

Definition 11.67 (Tree–level TT, triad–resolved exchange amplitude)**.** At tree level, the TT–typed exchange amplitude between two Tantrakyma sources is the $\bar{\Sigma}$–time–ordered contraction of two vertices with the typed propagator (Definition 11.54):

$$\boxed{\mathcal{A}_{12\to 34} = \sum_{a\in\{\text{corr,vac,spec}\}} \int d\mu_{\bar{\Sigma}}(x)\, d\mu_{\bar{\Sigma}}(y) \left\langle \hat{V}^{(a)}_{\mu\nu}(x) \right\rangle i\, G^{\mathrm{F},[\mathrm{TT}],(a)\,\mu\nu}{}_{\rho\sigma}(x,y) \left\langle \hat{V}^{(a)\rho\sigma}(y) \right\rangle.} \tag{11.113}$$

What each factor is:

- $i\, G^{\mathrm{F},[\mathrm{TT}],(a)}_{\mu\nu}{}^{\rho\sigma}(x,y)$ is the sector–a TT Feynman kernel (typed two–point) defined in Definition 11.54 and eq. (11.107); it solves the typed wave equation with Feynman boundary conditions and has support consistent with the $\bar{\Sigma}$ causal cone.
- $\hat{V}^{(a)}_{\mu\nu}(x)$ is the sector–resolved vertex operator (right leg), obtained by projecting the bi–local parent via $\Pi^{(a)}$; equivalently, its explicitly typed version $\hat{V}^{[\mathrm{TT}],(a)}_{\mu\nu} := \Pi^{[\mathrm{TT}]\,\alpha\beta}_{\bar{\Sigma}\,\mu\nu}\Pi^{(a)}\hat{V}_{\alpha\beta}$ may be used. On the certified plateau $\mathfrak{P}$, TT/triadic projectors commute and the two forms are interchangeable (cf. Lemma 11.92 and eq. (11.109)).
- The sum over a implements the triadic decomposition of the exchange channel (Section 11.1.4 and Definition 11.51).

Momentum–space form and currents. Fourier transforming the vertex matrix elements defines the *sector currents*

$$\tilde{\mathcal{J}}^{(a)}_{\mu\nu}(k) := \int d\mu_{\bar{\Sigma}}(x)\; e^{+ikx}\,\langle\Theta_3|\hat{V}^{[\mathrm{TT}],(a)}_{\mu\nu}(x)|\Theta_1\rangle,$$
$$\tilde{\mathcal{J}}^{(a)\rho\sigma}(-k) := \int d\mu_{\bar{\Sigma}}(y)\; e^{-iky}\,\langle\Theta_4|\hat{V}^{[\mathrm{TT}],(a)\rho\sigma}(y)|\Theta_2\rangle,$$

where k is the momentum flowing through the exchanged graviton (e.g. in the t–channel $k = p_1 - p_3$, in the s–channel $k = p_1 + p_2$, etc.). Using the TT polarization sum $\Pi^{[\mathrm{TT}]\mu\nu}{}_{\rho\sigma}(k) := \sum_s e^{(s)\mu\nu}(k)\, e^{(s)}_{\rho\sigma}(k)^*$ (Section 11.1.3), the amplitude becomes

$$\boxed{\mathcal{A}_{12\to34} = \sum_a \int \frac{d^dk}{(2\pi)^d}\; \tilde{\mathcal{J}}^{(a)}_{\mu\nu}(k)\; \frac{i\,\Pi^{[\mathrm{TT}]\mu\nu}{}_{\rho\sigma}(k)}{k^2 - m_a^2 + i\epsilon}\; \tilde{\mathcal{J}}^{(a)\rho\sigma}(-k)\; (2\pi)^d\delta^{(d)}(p_1+p_2-p_3-p_4).}$$

(11.114)

Here m_a^2 is the sector–dependent effective mass parameter inherited from the typed wave operator (Section 11.2.3); in the massless TT limit $m_a^2 = 0$ and the $i\epsilon$ prescription implements the Feynman contour.

Causality/positivity. On $\mathfrak{P}$, the TT projector preserves microcausality with respect to the $\bar{\Sigma}$ cone and ensures Källén–Lehmann positivity per sector; hence the typed kernel obeys the triadic spectral representation with non–negative spectral weights and the standard convexity bounds in the forward limit (eq. (11.91), Proposition 11.14, and Section 11.3.6).

11.3.5 Effective Graviton-Mediated Propagation

Integrating out the TT graviton at quadratic order yields a causal, nonlocal effective action for Tantrakyma fields, block-diagonal in the triad:

$$\boxed{S^{(\Theta)}_{\mathrm{eff}} = \frac{1}{2}\sum_a \int d\mu_{\bar{\Sigma}}(x)\, d\mu_{\bar{\Sigma}}(y)\; \mathrm{Tr}\Big[\hat{V}^{(a)}_{\mu\nu}(x)\, i\, G^{\mathrm{F},[\mathrm{TT}],(a)\,\mu\nu}{}_{\rho\sigma}(x,y)\, \hat{V}^{(a)\rho\sigma}(y)\Big] + \cdots,}$$

(11.115)

where "$\cdots$" indicates higher-point and radiative corrections consistent with the typed projectors. This representation makes manifest the sector-resolved mediation and is the starting point for dispersion/positivity analyses (Section 11.3.6).

Derivation (integrating out the TT graviton). Work channel–wise on the certified plateau $\mathfrak{P}$, where the TT and triadic projectors commute (Lemma 11.92). Keep terms up to quadratic order in the graviton fluctuation and linear order in the matter–gravity coupling:

$$S[\hat{\boldsymbol{\Theta}}, h] = S_\Theta[\hat{\boldsymbol{\Theta}}] + \frac{1}{2}\sum_a \int d\mu_{\bar{\Sigma}}(x)\, d\mu_{\bar{\Sigma}}(y)\, h^{(a)}_{\mu\nu}(x)\, K^{[\mathrm{TT}],(a)\,\mu\nu}{}_{\alpha\beta}(x)\, \delta^{(d)}(x-y)\, h^{(a)\,\alpha\beta}(y)$$
$$+ \sum_a \int d\mu_{\bar{\Sigma}}(x)\, h^{(a)}_{\mu\nu}(x)\, J^{(a)\,\mu\nu}(x).$$

where $h^{(a)} := \Pi^{[\mathrm{TT}]}\Pi^{(a)}\delta\hat{g}$ is TT & sectoral, $K^{[\mathrm{TT}],(a)}$ is the typed wave operator (Section 11.2.3), and $J^{(a)\,\mu\nu}(x) := \mathrm{Tr}\,\hat{V}^{[\mathrm{TT}],(a)\,\mu\nu}(x)$ is the (typed) source current built from the vertex (eq. (11.109) and Section 11.3.2). In compact notation,

$$S = S_\Theta + \frac{1}{2}\sum_a \langle h^{(a)}, K^{[\mathrm{TT}],(a)} h^{(a)}\rangle + \sum_a \langle h^{(a)}, J^{(a)}\rangle.$$

Complete the square in each sector a:

$$\frac{1}{2}\langle h, Kh\rangle + \langle h, J\rangle = \frac{1}{2}\langle h + K^{-1}J,\ K\,(h + K^{-1}J)\rangle - \frac{1}{2}\langle J,\ K^{-1}J\rangle.$$

Gaussian path integral. The in–out generating functional (Feynman $i\epsilon$) factorizes over a and over TT modes,

$$\int \mathcal{D}h\ \exp\Big\{ i\Big[\tfrac{1}{2}\langle h, Kh\rangle + \langle h, J\rangle\Big]\Big\} = \mathcal{N}\ \exp\Big\{ -\frac{i}{2}\,\langle J,\ K^{-1}J\rangle\Big\},$$

where $\mathcal{N}$ contains the (channel–wise) functional determinants $\left(\det K^{[\mathrm{TT}],(a)}\right)^{-1/2}$ (one-loop vacuum bubbles; they are absorbed into the "$\cdots$" in (11.115)).

Relate K^{-1} to the typed Feynman kernel. By definition (eq. (11.107)), the TT Feynman kernel satisfies $K^{[\mathrm{TT}],(a)} \circ G^{\mathrm{F},[\mathrm{TT}],(a)} = i\,\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\delta^{(d)}$, so on the TT subspace $K^{-1} = (1/i)\,G^{\mathrm{F},[\mathrm{TT}],(a)}$. Hence

$$-\frac{i}{2}\,\langle J,\ K^{-1}J\rangle = -\frac{i}{2}\left\langle J,\ \frac{1}{i}G^{\mathrm{F},[\mathrm{TT}]}J\right\rangle = \frac{i}{2}\,\langle J,\ G^{\mathrm{F},[\mathrm{TT}]}J\rangle = \frac{1}{2}\,\langle J,\ i\,G^{\mathrm{F},[\mathrm{TT}]}J\rangle.$$

Putting it together. Summing over sectors and restoring indices gives the nonlocal, triad–block–diagonal effective action for Tantrakyma fields:

$$\boxed{S^{(\Theta)}_{\mathrm{eff}} = S_\Theta + \frac{1}{2}\sum_a \int d\mu_{\bar{\Sigma}}(x)\, d\mu_{\bar{\Sigma}}(y)\ \mathrm{Tr}\Big[\hat{V}^{(a)}_{\mu\nu}(x)\, i\, G^{\mathrm{F},[\mathrm{TT}],(a)\,\mu\nu}{}_{\rho\sigma}(x,y)\,\hat{V}^{(a)\rho\sigma}(y)\Big] + \cdots}$$

(11.116)

where the ellipsis denotes higher–point/radiative terms (including $\log\det K$) consistent with the typed projectors. Orthogonality $\Pi^{(a)}\Pi^{(b)} = \delta_{ab}\Pi^{(a)}$ implies no cross–sector mixing at this order.

In–in (causal) variant. For causal response or expectation values, the Schwinger–Keldysh integration replaces $i\,G^{\mathrm{F},[\mathrm{TT}]}$ by the retarded TT kernel $G^{\mathrm{ret},[\mathrm{TT}]}$ (eq. (11.105)), yielding a manifestly causal nonlocal term with support inside the $\bar{\Sigma}$ cone.

11.3.6 Positivity and Dispersion Bounds for Triadic Exchange

On $\mathfrak{P}$, each sectoral two-point admits a KL representation with non-negative residue and spectral density on $\mathcal{H}_{\mathrm{phys}}$ (eq. (11.91)). Consequently, forward-limit amplitudes built from (11.114) satisfy standard analyticity/positivity constraints triadwise; the typed construction is Nielsen-flat and gauge-robust (Proposition 11.14). The TT TLO ledgers $(\Lambda^{[\mathrm{TT}]}_{\mathrm{law}}, \Lambda^{[\mathrm{TT}]}_{\mathrm{state}})$ serve as the certification layer for any claimed channel-selective deviation from GR (Definition 11.52).

11.3.7 Graviton-Mediated Tantrakyma Scattering

Definition 11.68 (Graviton–mediated S-matrix (typed, connected))**.** Let $|\Theta_{\rm in}\rangle$ and $|\Theta_{\rm out}\rangle$ be asymptotic Tantrakyma multiparticle states on the certified plateau $\mathfrak{P}$. The *graviton-mediated* scattering amplitude is the connected part of the $\bar{\Sigma}$-time-ordered matrix element

$$\boxed{\mathcal{A}_{\rm grav}[\Theta_{\rm in}\to\Theta_{\rm out}] := \Big\langle \Theta_{\rm out}\Big|\, T_{\bar{\Sigma}} \exp\Big(i\, S^{(1)}_{\rm int}[\hat{h},\hat{\mathbf{\Theta}}]\Big)\,\Big|\Theta_{\rm in}\Big\rangle_{\rm conn},} \tag{11.117}$$

where $T_{\bar{\Sigma}}\{\cdot\}$ is the $\bar{\Sigma}$-time-ordering operator (Definition 11.53) and $S^{(1)}_{\rm int}$ is the typed, linearized interaction $S^{(1)}_{\rm int} = \int d\mu_{\bar{\Sigma}}\,\mathrm{Tr}\big[\hat{h}_{\mu\nu}\hat{V}^{\mu\nu}\big]$ with $\hat{V}^{\mu\nu}$ the TT vertex (Definitions 11.64 and 11.66). The TT projector $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ and the triad projectors $\Pi^{(a)}$ commute, and all band tags commute with both (Definitions 9.10 and 11.24).

Lemma 11.69 (Tree-level TT graviton exchange (triad-resolved))**.** *Expanding* (11.117) *to second order in the Dyson series and keeping the connected contribution gives the single-exchange amplitude*

$$\boxed{\begin{aligned}\mathcal{A}^{(2)}_{\rm grav} = -\frac{1}{2}\sum_{a\in\{\rm corr,vac,spec\}}\int d\mu_{\bar{\Sigma}}(x)\,d\mu_{\bar{\Sigma}}(y)\,\Big\langle\Theta_{\rm out}\Big|\,T_{\bar{\Sigma}}\Big\{\hat{V}^{(a)}_{\mu\nu}(x)\,\hat{V}^{(a)}_{\rho\sigma}(y)\Big\}\,\Big|\Theta_{\rm in}\Big\rangle\\ \times\, i\,G^{\mathrm{F},[\mathrm{TT}],(a)\,\mu\nu\,\rho\sigma}(x,y),\end{aligned}} \tag{11.118}$$

where $G^{\mathrm{F},[\mathrm{TT}],(a)}$ is the typed Feynman propagator in sector a (Definitions 11.51 and 11.54), and $\hat{V}^{(a)}_{\mu\nu} := \Pi^{(a)}\hat{V}_{\mu\nu}$ (Sections 11.1.4 and 11.3.3). In momentum space this reduces to the block-diagonal form of eq. (11.114).

11.3.8 Effective Graviton-Mediated Propagation

The leading graviton-induced dressing of the Tantrakyma two-point function arises from emission and reabsorption of a virtual TT graviton. Summing over triadic channels yields a causal, nonlocal contribution that respects the $\bar{\Sigma}$-cone and KL positivity on $\mathcal{H}_{\rm phys}$ (eq. (11.91) and Proposition 11.14):

Definition 11.70 (Gravitational dressing of the Tantrakyma propagator (typed, triad-summed))**.** The graviton-exchange correction to the Tantrakyma two-point kernel $\hat{G}_{\Theta\Theta}(x,x')$ is

$$\boxed{\begin{aligned}\Delta\hat{G}_{\Theta\Theta}(x,x') = \sum_a\int d\mu_{\bar{\Sigma}}(z)\,d\mu_{\bar{\Sigma}}(z')\,\Big\langle 0\Big|\,T_{\bar{\Sigma}}\Big\{\hat{\mathbf{\Theta}}(x)\,\hat{V}^{(a)}_{\mu\nu}(z)\,i\,G^{\mathrm{F},[\mathrm{TT}],(a)\,\mu\nu}{}_{\rho\sigma}(z,z')\\ \hat{V}^{(a)\rho\sigma}(z')\,\hat{\mathbf{\Theta}}(x')\Big\}\,\Big|0\Big\rangle_{\rm conn}.\end{aligned}} \tag{11.119}$$

with $i\,G^{\mathrm{F},[\mathrm{TT}],(a)}$ the TT Feynman kernel (Definition 11.54) and $\hat{V}^{(a)}$ as in Definition 11.64. On $\mathfrak{P}$, band tags $K^{(\ell)}_{\rm IR}$ do not enlarge causal cones and the result is Nielsen-flat and gauge robust (Definition 6.280 and Proposition 11.14).

Explanation of Definitions 11.70 and 11.71. Both formulas compute the *gravitational dressing* (self–energy insertion) of the Tantrakyma two–point kernel by exchanging a single typed (TT) graviton between two matter–geometry interaction vertices. The construction is causal and covariant on the background $(\mathcal{X}_q, \bar{\Sigma})$ and is evaluated on a certified plateau $\mathfrak{P}$ so that all realized weave configuration operators commute and are BRST/Nielsen–safe (Definitions 6.280 and 9.10, Propositions 7.3 and 10.9, and Corollary 11.5). The operator $T_{\bar{\Sigma}}$ denotes $\bar{\Sigma}$–time ordering on the foliation determined by the chronogeometric background, the measures $d\mu_{\bar{\Sigma}}$ are the relational volume elements, and the subscript "conn" restricts to the connected contribution (cf. Chapter 10 and Definition 6.7 and eq. (11.115)). The kernel $i\,G^{\mathrm{F},[\mathrm{TT}],(a)}$ is the TT Feynman Green function of the sector–resolved typed wave operator $K^{[\mathrm{TT}],(a)}$ (Definitions 11.54 and 11.60), and $\hat{V}^{(a)}$ is the triad–filtered graviton–Tantrakyma vertex (Definitions 10.2 and 11.64). In this way, (11.119) realizes the standard diagrammatic contraction "vertex–propagator–vertex" summed over triad sectors $a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$, providing the leading nonlocal, causal correction to $\hat{G}_{\Theta\Theta}$ (compare Definition 11.67).

The explicitly TT–typed form (11.120) inserts the York–Lichnerowicz projector at the vertices, $\hat{V}^{[\mathrm{TT}],(a)}_{\mu\nu} := (\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}})_{\mu\nu}{}^{\alpha\beta}\,\Pi^{(a)}\hat{V}_{\alpha\beta}$ (eq. (11.109) and Definition 10.3). This makes manifest that only transverse–traceless polarizations propagate and couple. The two definitions are *equivalent* on $\mathfrak{P}$: by projector commutations and block–diagonality, $\Pi^{(a)}\,G^{\mathrm{F},[\mathrm{TT}]}(k)\,\Pi^{(b)} = \delta_{ab}\,G^{\mathrm{F},[\mathrm{TT}],(a)}(k)$ and $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\hat{V} = \hat{V}^{[\mathrm{TT}]}$ (Propositions 10.9, 11.20 and 11.77 and eq. (11.33)). Moreover, the TT Ward identities $k_\mu \hat{V}^{[\mathrm{TT}],(a)\mu\nu} = 0$ and $\bar{\Sigma}_{\mu\nu}\hat{V}^{[\mathrm{TT}],(a)\mu\nu} = 0$ ensure that any non–TT or τ–typed insertion is TT–null (eq. (11.128) and Corollary 11.73), so inserting $\Pi^{[\mathrm{TT}]}$ at the vertices does not alter the value of the contraction but clarifies its typed content. Causality and gauge robustness follow from the support of the typed causal kernel $\Delta^{[\mathrm{TT}]}$ and the cone–preserving band window $K^{(\ell)}_{\mathrm{IR}}$ (Proposition 11.14 and Remark 9.83). In summary, (11.119) and (11.120) are two notationally different but cohomologically identical realizations of the same TT, triad–resolved graviton dressing of the Tantrakyma propagator.

Definition 11.71 (TT-typed gravitational dressing of the Tantrakyma propagator)**.** On the certified plateau $\mathfrak{P}$, the TT–projected, sector–resolved graviton dressing of the Tantrakyma two–point function is

$$\boxed{\begin{aligned}\Delta\hat{G}^{[\mathrm{TT}]}_{\Theta\Theta}(x,x') \equiv \sum_{a\in\{\mathrm{corr,vac,spec}\}} \int d\mu_{\bar{\Sigma}}(z)\,d\mu_{\bar{\Sigma}}(z')\,\Big\langle 0\Big|\,T_{\bar{\Sigma}}\Big\{ \\ \hat{\boldsymbol{\Theta}}(x)\,\hat{V}^{[\mathrm{TT}],(a)}_{\mu\nu}(z)\,i\,G^{\mathrm{F},[\mathrm{TT}],(a)\,\mu\nu}{}_{\rho\sigma}(z,z') \\ \hat{V}^{[\mathrm{TT}],(a)\,\rho\sigma}(z')\,\hat{\boldsymbol{\Theta}}(x')\Big\}\,\Big|0\Big\rangle_{\mathrm{conn}}.\end{aligned}} \tag{11.120}$$

Here $i\,G^{\mathrm{F},[\mathrm{TT}],(a)}$ is the typed Feynman kernel (Definition 11.54 and eq. (11.107)) and

$$\hat{V}^{[\mathrm{TT}],(a)}_{\mu\nu}(z) := \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}\,\mu\nu}{}^{\alpha\beta}\,\Pi^{(a)}\,\hat{V}_{\alpha\beta}(z)$$

is the typed, sectoral vertex (eq. (11.109) and Section 11.3.2).

Proposition 11.72 (TT-typed linearized vertex from the bi-local parent)**.** *Let $S_{\mathrm{int}}[\hat{\boldsymbol{\Phi}}, \hat{\boldsymbol{\Theta}}]$ be the bi-local interaction action of Chapter 11 (Definition 10.223 and Section 10.19.4), and let*

the typed graviton be defined by the background-covariant TT projector

$$\hat{h}_{\mu\nu}(x) := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}_{\mu\nu}(x) \quad \textit{(symmetric, traceless, transverse on } (\mathcal{X}_q, \bar{\Sigma})\textit{; cf. Section 11.1.3).}$$

Then the linearized*, TT-typed reduction of the interaction is*

$$\boxed{S_{\mathrm{int}}^{(1)}[\hat{h}, \hat{\boldsymbol{\Theta}}] = \int d\mu_{\bar{\Sigma}}(x)\ \mathrm{Tr}\Big[\hat{h}_{\mu\nu}(x)\,\hat{V}^{\mu\nu}(x; \hat{\boldsymbol{\Theta}})\Big]\,,} \tag{11.121}$$

with the typed vertex operator

$$\boxed{\begin{aligned}\hat{V}^{\mu\nu}(x; \hat{\boldsymbol{\Theta}}) &:= \Big(\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\Big)^{\mu\nu}{}_{\alpha\beta}\ \frac{\delta S_{\mathrm{int}}\Big[\hat{\boldsymbol{\Phi}}, \hat{\boldsymbol{\Theta}}\Big]}{\delta g_{\alpha\beta}(x)}\bigg|_{\hat{\boldsymbol{\Phi}}\to\bar{\boldsymbol{\Phi}}} \\ &\equiv \Big(\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\Big)^{\mu\nu}{}_{\alpha\beta}\int d\mu_{\bar{\Sigma}}(x')\,\frac{\delta\mathcal{U}_{\Phi\Theta}(x, x')}{\delta g_{\alpha\beta}(x)}\,\hat{\boldsymbol{\Theta}}(x').\end{aligned}} \tag{11.122}$$

where $\mathcal{U}_{\Phi\Theta}$ is the bi-local interaction kernel of Chapter 11 (Definition 4.9 and Section 10.19.4). On the certified plateau $\mathfrak{P}$ the band tags commute with $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ (Definitions 6.280 and 9.10), and with the triad projectors $\Pi^{(a)}$ (Proposition 11.93), so the sector-resolved vertex $\hat{V}^{(a)\mu\nu} := \Pi^{(a)}\hat{V}^{\mu\nu}$ is well defined and $[\,\Pi^{(a)}, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,] = 0$.

Proof sketch. Start from the bi-local parent

$$S_{\mathrm{int}} = \int d\mu_{\bar{\Sigma}}(x)\,d\mu_{\bar{\Sigma}}(x')\,\mathrm{Tr}\Big[\hat{\boldsymbol{\Phi}}^{\dagger}(x)\,\mathcal{U}_{\Phi\Theta}(x, x')\,\hat{\boldsymbol{\Theta}}(x')\Big]$$

(Section 10.19.4). Linearize around $\bar{\Phi}$ and vary with respect to the metric to first order in $\delta g_{\alpha\beta}$, then project with $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ to extract the radiative spin–2 part, obtaining (11.122). Inserting $\hat{h}_{\mu\nu} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\delta\hat{g}_{\mu\nu}$ gives (11.121). Commutation with band/triad projectors follows from Definition 9.10 and Proposition 11.93. □

Corollary 11.73 (TT preservation and τ-null right leg)**.** *Let $\hat{\tau}_{\mu\nu} := \delta\hat{t}_{\mu\nu}$ denote temporal/-coherence fluctuations (Chapter 11). On the weave, $\hat{\tau}$ is TT-null (no radiative spin–2; Section 11.1.3). Consequently, any right-leg insertion that is τ-typed produces no TT vertex:*

$$\boxed{\Pi^{[\mathrm{TT}]\,\mu\nu}{}_{\alpha\beta}\,\hat{V}^{\alpha\beta}\Big(x; \hat{\boldsymbol{\Theta}} \mapsto \hat{\tau}\Big) = 0, \qquad \Rightarrow \qquad S_{\mathrm{int}}^{(1)}[\hat{h}, \hat{\tau}] = 0.} \tag{11.123}$$

Hence τ cannot radiate spin–2, and the linearized vertex preserves the TT type.

Remark 11.74 (Sectoral (triadic) reduction of the vertex)**.** Using $\Pi^{(a)}$ (Section 11.1.4) and the commuting relations, define $\hat{V}^{(a)\mu\nu} := \Pi^{(a)}\hat{V}^{\mu\nu}$ and $\hat{h}^{(a)}_{\mu\nu} := \Pi^{(a)}\hat{h}_{\mu\nu}$. Then

$$S_{\mathrm{int}}^{(1)} = \sum_{a\in\{\mathrm{corr,vac,spec}\}}\int d\mu_{\bar{\Sigma}}(x)\,\mathrm{Tr}\Big[\hat{h}^{(a)}_{\mu\nu}(x)\,\hat{V}^{(a)\mu\nu}(x; \hat{\boldsymbol{\Theta}})\Big]\,,$$

which is block-diagonal in the triad and compatible with the typed Feynman rules (Definitions 11.51 and 11.54).

11.3.9 Graviton Coupling Channels

To analyze scattering and mediation processes it is convenient to decompose the graviton–Tantrakyma vertex into *typed* (TT) and *triadic* components, and then into a polarization/-momentum basis compatible with the TT mode expansion of Section 11.1.3.

Definition 11.75 (Triadic decomposition of the typed vertex)**.** Let the typed graviton field be $\hat{h}_{\mu\nu} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}_{\mu\nu}$ and the linearized vertex be defined by (11.121)–(11.122) (Section 11.3.2). Using the commuting triad projectors $\{\Pi^{(a)}\}_{a\in\{\mathrm{corr,vac,spec}\}}$ (Section 11.1.4) and the band tags on the certified plateau $\mathfrak{P}$ (Definitions 6.280 and 9.10), define

$$\boxed{\hat{V}^{\mu\nu}(x;\hat{\boldsymbol{\Theta}}) = \sum_{a\in\{\mathrm{corr,vac,spec}\}} \hat{V}^{(a)\mu\nu}(x;\hat{\boldsymbol{\Theta}}), \qquad \hat{V}^{(a)\mu\nu} := \Pi^{(a)}\,\hat{V}^{\mu\nu}, \qquad \left[\Pi^{(a)}, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\right] = 0.} \tag{11.124}$$

The *typed, triad-resolved* vertex blocks are $\hat{V}^{[\mathrm{TT}],(a)\mu\nu} := \Pi^{[\mathrm{TT}]\,\mu\nu}{}_{\rho\sigma}\,\hat{V}^{(a)\rho\sigma}$, and the linearized action is block-diagonal,

$$S_{\mathrm{int}}^{(1)} = \sum_a \int d\mu_{\bar{\Sigma}}(x)\,\mathrm{Tr}\Big[\hat{h}^{(a)}_{\mu\nu}(x)\,\hat{V}^{[\mathrm{TT}],(a)\mu\nu}(x;\hat{\boldsymbol{\Theta}})\Big], \tag{11.125}$$

$$\text{where} \quad \hat{h}^{(a)}_{\mu\nu} := \Pi^{(a)}\hat{h}_{\mu\nu}. \tag{11.126}$$

Remark 11.76 (Polarization/momentum (operator) basis)**.** Let $\{e^{(s,a)}_{\mu\nu}(k)\}_{s=1}^{d_g}$ be an orthonormal TT polarization basis in sector a (Section 11.1.3), and let $\mathcal{K}^{\mu\nu}(x,x')$ denote the effective response kernel in Proposition 11.72. The *modal vertex amplitudes* are defined by

$$\boxed{\begin{aligned} \mathcal{V}^{(s,a)}(k;x) &:= \int d^dx'\; \mathcal{K}^{\mu\nu}(x,x')\,e^{(s,a)}_{\mu\nu}(k)\,e^{-ik\cdot x'}, \\ \hat{V}^{[\mathrm{TT}],(a)}_{\mu\nu}(x) &= \sum_s \int \frac{d^dk}{(2\pi)^d}\; e^{(s,a)}_{\mu\nu}(k)\,\mathcal{V}^{(s,a)}(k;x) \;+\; \mathrm{h.c.}, \end{aligned}} \tag{11.127}$$

which is compatible with the sector-resolved TT mode expansion of $\hat{h}^{(a)}_{\mu\nu}$ (eq. (11.97)) and the triadic propagators (Definition 11.51 and eq. (11.86)).

In the *typed, triad-resolved* expansion of eq. (11.127) one should read:

- $\mathcal{V}^{(s,a)}(k;x)$ is the *modal vertex amplitude* in triadic sector $a \in \{\mathrm{corr,vac,spec}\}$, a scalar function of the spacetime point x and momentum k, encoding the strength/structure of the coupling between a Tantrakyma excitation and the s-th TT polarization of the sector-a graviton. It is evaluated on the certified plateau $\mathfrak{P}$ with BRST-closed band tags, so that readings are Nielsen-flat (Definitions 6.280 and 9.10).
- The polarization contraction $e^{(s,a)\mu\nu}(k)\,\hat{V}^{[\mathrm{TT}],(a)}_{\mu\nu}(x)$ isolates the contribution of each independent TT spin-2 mode to the vertex in sector a (Section 11.1.3 and Definition 11.75). Commutation $[\Pi^{(a)}, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}] = 0$ ensures the contraction is well-defined in type (eq. (11.124)).

- The structure of $\mathcal{V}^{(s,a)}(k;x)$ follows by Fourier-transforming the effective response kernel $\mathcal{K}^{\mu\nu}(x,x')$ (Proposition 11.72):

$$\mathcal{V}^{(s,a)}(k;x) := \int d^d x'\; \mathcal{K}^{\mu\nu}(x,x')\, e^{(s,a)}_{\mu\nu}(k)\, e^{-ikx'}.$$

- This modal basis allows the interaction action and the resulting amplitudes to be written entirely in terms of TT-projected, sector-resolved propagators and momentum-dependent vertex factors (Definition 11.51 and eq. (11.86)), as used in Section 11.3.7.

Proposition 11.77 (TT Ward identity and sectoral preservation)**.** *On the plateau $\mathfrak{P}$ the typed vertex satisfies the TT Ward identities*

$$\boxed{k_\mu\, \hat{V}^{[\mathrm{TT}],(a)\mu\nu}(k) = 0, \qquad \bar{\Sigma}_{\mu\nu}\, \hat{V}^{[\mathrm{TT}],(a)\mu\nu}(k) = 0, \quad \forall a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\},} \tag{11.128}$$

and τ-typed insertions are TT-null (Corollary 11.73). Consequently, internal graviton lines in typed Feynman graphs are TT-propagators contracted with TT vertices in the same *triadic block, i.e. the coupling is block-diagonal by a at the linearized level.*

Proof. Typed vertex and projectors. On the certified plateau $\mathfrak{P}$, the TT projector $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ and the triad projectors $\{\Pi^{(a)}\}_{a\in\{\mathrm{corr,vac,spec}\}}$ commute with each other and with the band realized weave configuration (tag hygiene) (Proposition 10.9 and Definitions 6.280 and 9.10). Define the typed vertex in momentum space by left–projecting the rank–2 vertex kernel,

$$\hat{V}^{[\mathrm{TT}],(a)}_{\mu\nu}(k) := \left(\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\Pi^{(a)}\right)_{\mu\nu}{}^{\rho\sigma}\, \hat{V}_{\rho\sigma}(k), \tag{11.129}$$

so that it couples only to TT, sector–a carriers (Definitions 10.2, 10.3 and 11.3).

TT Ward identities. By construction, the image of $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ consists of symmetric, traceless, and transverse tensors (York–Lichnerowicz TT calculus: eq. (10.4) and Definition 10.3). Equivalently, using the typed polarization basis $\{e^{(a,\lambda)}_{\mu\nu}(k;x)\}$ of Definition 11.18, one has for each $(k;x)$:

$$\bar{g}^{\mu\nu}(x) e^{(a,\lambda)}_{\mu\nu}(k;x) = 0, \qquad k^\mu e^{(a,\lambda)}_{\mu\nu}(k;x) = 0 \quad (\text{eqs. (11.30) and (11.31)}). \tag{11.130}$$

Since $\hat{V}^{[\mathrm{TT}],(a)}$ is valued in the TT image,

$$k_\mu\, \hat{V}^{[\mathrm{TT}],(a)\mu\nu}(k) = 0, \qquad \bar{\Sigma}_{\mu\nu}\, \hat{V}^{[\mathrm{TT}],(a)\mu\nu}(k) = 0, \tag{11.131}$$

which is precisely eq. (11.128). Moreover, the coherence/temporal (τ) insertions are TT–null by the TT calculus and the projector properties, giving Corollary 11.73.

Sectoral preservation (block–diagonality in a). Let $G^{\mathrm{F},[\mathrm{TT}],(a)}{}_{\mu\nu}{}^{\rho\sigma}(k)$ be the typed Feynman propagator, the Green function of $K^{[\mathrm{TT}],(a)}$ (Definitions 11.54 and 11.60). Using projector commutations and orthogonality of the triad projectors ($\Pi^{(a)}\Pi^{(b)} = \delta_{ab}\Pi^{(a)}$; Definition 10.2 and Proposition 10.9), one finds the block structure

$$\Pi^{(a)}\, G^{\mathrm{F},[\mathrm{TT}]}(k)\, \Pi^{(b)} = \delta_{ab}\, G^{\mathrm{F},[\mathrm{TT}],(a)}(k), \qquad \Pi^{(a)}\, \hat{V}^{[\mathrm{TT}]}(k) = \hat{V}^{[\mathrm{TT}],(a)}(k), \tag{11.132}$$

so internal graviton lines in typed Feynman graphs are TT propagators contracted with TT vertices in the *same* triadic block. In particular, a mixed contraction with $a \neq b$ contains $\Pi^{(a)} G^{\mathrm{F},[\mathrm{TT}]}(k) \Pi^{(b)} = 0$ and vanishes. This block–diagonality is consistent with the triad–resolved mode expansion and the orthogonality/completeness relations of the TT polarization basis (Definitions 11.18 and 11.19 and eq. (11.33)), and is reflected in the factorized CCR/Fock structure (Proposition 11.20).

Causality and plateau/Nielsen safety. Typed microcausality and the Peierls bracket are generated by the same TT operator (Proposition 11.14 and eq. (11.25)); since all realized weave configuration operators are BRST–closed and Nielsen–flat on $\mathfrak{P}$, the Ward identities and the block–diagonal coupling are gauge–parameter independent and regulator–safe (Proposition 7.3, Corollary 11.5, and Remark 9.83).

Altogether, eq. (11.128) holds, τ-typed insertions are TT–null, and internal lines couple TT propagators to TT vertices *within* the same triad sector, as claimed. □

Remark 11.78 (Typed TLO law residual from diagrammatic contractions)**.** Let $\mathbf{R}^{[\mathrm{TT}]}$ and $\Lambda^{[\mathrm{TT}]}_{\mathrm{law}}$ denote the TT law residual and visibility (Chapter 9; cf. the TT TLO overlay introduced in Section 11.1.3). In momentum space, a practical estimator on $\mathfrak{P}$ is obtained by contracting the TT Ward identity with external polarization legs and inserting the typed propagator $G^{\mathrm{F},[\mathrm{TT}],(a)}$ (Definitions 11.51 and 11.54):

$$\boxed{\begin{aligned}\mathscr{R}^{(s,a)}(k) &:= e^{(s,a)}_{\mu\nu}(k)\left[K^{\mu\nu}_{[\mathrm{TT}]\rho\sigma}(k)\, G^{\mathrm{F},[\mathrm{TT}],(a)\,\rho\sigma}{}_{\alpha\beta}(k) - \mathbf{1}^{\mu\nu}{}_{\alpha\beta}\right] e^{(s,a)\,\alpha\beta}(k),\\ &\Lambda^{[\mathrm{TT}]}_{\mathrm{law}} \to 1 \text{ as } \mathscr{R}^{(s,a)}(k) \to 0,\end{aligned}} \tag{11.133}$$

where $K_{[\mathrm{TT}]}$ is the TT Lichnerowicz-type operator on $(\mathcal{X}_q, \bar{\Sigma})$ (Section 11.2.3). By Proposition 11.77 and typed microcausality (Proposition 11.14), $\mathscr{R}^{(s,a)}(k)$ vanishes on-shell and remains Nielsen-flat under admissible band deformations on $\mathfrak{P}$.

Notation check ($\mathscr{R}$ vs. $\mathbf{R}^{[\mathrm{TT}]}$). The symbol $\mathscr{R}$ is introduced only in the diagrammatic estimator $\mathscr{R}^{(s,a)}(k)$ of eq. (11.133). Everywhere else the *TT law residual* is denoted by the bold operator-valued symbol $\mathbf{R}^{[\mathrm{TT}]}$ and its visibility by $\Lambda^{[\mathrm{TT}]}_{\mathrm{law}}$ (Definition 11.6). Thus the notation is consistent provided that:

$$\boxed{\mathscr{R}^{(s,a)}(k) \text{ is a } \textit{momentum–space estimator}; \quad \mathbf{R}^{[\mathrm{TT}]} \text{ is the } \textit{exact} \text{ TT law residual.}}$$

Here $a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$ is the triad label (Definition 10.2) and s indexes TT polarizations (Definition 11.18). The superscript $^{(s,a)}$ should always appear as an ordered pair in parentheses, and the argument k should be present when working in momentum space. No other variants of $\mathscr{R}$ (e.g. $\mathcal{R}$, $\mathscr{R}_a$, or $\mathscr{R}^{a,s}$) have been introduced.

Definition 11.79 (Diagrammatic TT law–residual estimator)**.** Given TT polarization tensors $e^{(s,a)}_{\mu\nu}(k)$ (Definition 11.18), the TT Lichnerowicz–type operator $K_{[\mathrm{TT}]}$ (Definition 11.60), and the typed Feynman propagator $G^{\mathrm{F},[\mathrm{TT}],(a)}$ (Definition 11.54), define

$$\boxed{\mathscr{R}^{(s,a)}(k) := e^{(s,a)}_{\mu\nu}(k)\left[K^{\mu\nu}_{[\mathrm{TT}]\rho\sigma}(k)\, G^{\mathrm{F},[\mathrm{TT}],(a)\,\rho\sigma}{}_{\alpha\beta}(k) - \mathbf{1}^{\mu\nu}{}_{\alpha\beta}\right] e^{(s,a)\,\alpha\beta}(k).} \tag{11.134}$$

This *estimator* is a scalar function of k on the certified plateau $\mathfrak{P}$, and it vanishes on–shell by the typed Ward identity (Propositions 11.14 and 11.77). Its smallness correlates with the law visibility $\Lambda_{\mathrm{law}}^{[\mathrm{TT}]}$ of Definition 11.6, but it does not replace the exact residual $\mathbf{R}^{[\mathrm{TT}]}$.

This decomposition underpins several foundational aspects of the perturbative framework:

1. **Momentum-space Feynman rules (typed and triad-resolved).** The expansion (11.127) expresses $\hat{V}_{\mu\nu}^{[\mathrm{TT}],(a)}(x)$ in plane-wave TT modes and polarization tensors $e_{\mu\nu}^{(s,a)}(k)$, which are the standard building blocks for Feynman rules. Energy–momentum conservation and the TT propagator structure are manifest in loop and tree graphs, with internal lines carrying the sector label a and using $G^{\mathrm{F},[\mathrm{TT}],(a)}$ (Definition 11.54 and Sections 11.2.2 and 11.3.7). Band tags commute with the projectors, so causal cones are not enlarged on $\mathfrak{P}$ (Proposition 11.14).

2. **Polarization selection rules and dynamical couplings.** Working in the TT polarization basis isolates helicity-2 couplings. The modal amplitudes $\mathcal{V}^{(s,a)}(k;x)$ reveal selection rules and constraints (helicity flow, parity/time-reversal where applicable) sector by sector, controlled by the TT Ward identities (11.128) and the typed projection $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ (Section 11.1.3 and Definition 11.75).

3. **Background covariance and gauge (type) invariance of observables.** Because $e_{\mu\nu}^{(s,a)}(k)$ are TT with respect to $(\mathcal{X}_q, \bar{\Sigma})$ (transverse $k^\mu e_{\mu\nu}^{(s,a)} = 0$, traceless $\bar{\Sigma}^{\mu\nu} e_{\mu\nu}^{(s,a)} = 0$), amplitudes built from (11.127) are manifestly background-covariant and invariant under linearized diffeomorphisms in the TT type. Diagrammatically, contracting external TT legs with (11.128) provides a typed TLO law check; the corresponding residual $\mathscr{R}^{(s,a)}(k)$ of eq. (11.133) controls the law-visibility $\Lambda_{\mathrm{law}}^{[\mathrm{TT}]}$ on $\mathfrak{P}$.

11.3.10 Scattering Amplitudes from Graviton Exchange

With the typed graviton operator, vertex, and propagators in hand (Section 11.1.3, eq. (11.109), and Definition 11.54), we now assemble tree–level scattering amplitudes for Tantrakyma excitations mediated by virtual gravitons. All statements below are evaluated on a certified plateau $\mathfrak{P}$, with band tags commuting with the sector and TT projectors (Definitions 6.280 and 9.10 and Section 11.1.3).

Definition 11.80 (Triad-resolved, TT-projected tree amplitude)**.** Let $a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$ denote a triadic sector and $\hat{V}^{\mu\nu}(x)$ the typed vertex (eq. (11.109)). The tree-level graviton-exchange amplitude between asymptotic Tantrakyma states $|\mathbf{T}_1\rangle, |\mathbf{T}_2\rangle \to |\mathbf{T}_3\rangle, |\mathbf{T}_4\rangle$ is, in position space,

$$\boxed{\mathcal{A}_{\mathrm{grav}}^{(a)} = \int d\mu(x)\, d\mu(x')\, \left\langle \mathbf{T}_3 \middle| \hat{V}_{(a)}^{\mu\nu}(x) \middle| \mathbf{T}_1 \right\rangle \left(i\, G_{\mu\nu}^{\mathrm{F},[\mathrm{TT}],(a)\quad \rho\sigma}(x,x') \right) \left\langle \mathbf{T}_4 \middle| \hat{V}_{(a)\,\rho\sigma}(x') \middle| \mathbf{T}_2 \right\rangle,} \tag{11.135}$$

where $d\mu(x) = \sqrt{-\det \bar{\Sigma}(x)}\, d^d x$ and $G^{\mathrm{F},[\mathrm{TT}],(a)}$ is the TT-projected, sector-a Feynman propagator (Definitions 11.51 and 11.54). The full tree amplitude is the block-diagonal sum

$$\boxed{\mathcal{A}_{\mathrm{grav}} = \sum_{a\in\{\mathrm{corr,vac,spec}\}} \mathcal{A}^{(a)}_{\mathrm{grav}}.} \tag{11.136}$$

Remark 11.81 (Momentum-space representation)**.** Define the sector-resolved, TT vertex form factors $\tilde{J}^{(a)}_{\mu\nu}(q) := \int d\mu(x)\, e^{iq\cdot x}\, \langle \mathbf{T}_3 | \hat{V}^{(a)}_{\mu\nu}(x) | \mathbf{T}_1 \rangle$ and similarly for the second leg. Using $i\, G^{\mathrm{F},[\mathrm{TT}],(a)}_{\mu\nu}{}^{\rho\sigma}(q)$ (Sections 11.2.2 and 11.2.3), we obtain

$$\boxed{\mathcal{A}^{(a)}_{\mathrm{grav}} = \int \frac{d^d q}{(2\pi)^d}\, \tilde{J}^{(a)}_{\mu\nu}(-q) \left(i\, G^{\mathrm{F},[\mathrm{TT}],(a)\,\mu\nu}{}_{\rho\sigma}(q) \right) \tilde{J}^{(a)\,\rho\sigma}(q).} \tag{11.137}$$

Lemma 11.82 (Block diagonality and TT purity)**.** *On $\mathfrak{P}$, the sector projectors $\Pi^{(a)}$ commute with the background-covariant TT projector $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ (Section 11.1.3). Consequently, internal graviton lines are TT-projected and block-diagonal in the triad label a: there are no cross-channel propagators at tree level, and $\mathcal{A}_{\mathrm{grav}}$ decomposes as in (11.136). Moreover, insertions that are τ-typed on the right leg are TT-null and cannot radiate spin-2 (Section 11.3.2).*

Remark 11.83 (Causality and gauge safety)**.** Time ordering is taken with respect to the $\bar{\Sigma}$-cone (Definition 11.53), and $G^{\mathrm{F},[\mathrm{TT}]}$ inherits microcausality from the retarded/advanced TT kernels (Section 11.1.2). On $\mathfrak{P}$, band tags do not enlarge cones and ensure Nielsen flatness (Definition 6.280 and Proposition 11.14). The TT projection removes gauge (longitudinal/trace) modes, so only physical spin-2 polarizations propagate internally (Section 11.1.3).

Sidebar: Forward-limit dispersion and triadic KL positivity. In each sector a, the TT two-point function admits a Källén–Lehmann representation with positive residue and spectral density on $\mathcal{H}_{\mathrm{phys}}$ (Section 11.2.2). Analyticity and unitarity then imply forward-limit dispersion relations for $\mathcal{A}^{(a)}_{\mathrm{grav}}(s,t)$ at $t\to 0$, whose subtraction constants and moments are nonnegative functionals of the sectoral spectral densities. These yield triad-resolved positivity/dispersion bounds constraining low-energy Wilson coefficients and contact terms (Section 11.3.6).

Symmetries of the amplitude. The integrands in (11.135) are covariant bitensors on $(\mathcal{X}_q, \bar{\Sigma})$, and the volume element $d\mu(x)$ is diffeomorphism invariant; hence $\mathcal{A}_{\mathrm{grav}}$ is a scalar under background isometries. Gauge invariance at the linearized level is manifest by construction through the TT projection of both the internal propagators and the vertex legs (eq. (11.109) and Definition 11.54).

11.3.11 Effective Graviton-Mediated Propagation

On a certified plateau $\mathfrak{P}$ (band tags fixed, BRST-closed, Nielsen-flat; Definitions 6.280 and 9.10), the typed graviton can be integrated out at quadratic order to yield a nonlocal, *retarded* self-interaction for the Tantrakyma field. Causality is governed by the $\bar{\Sigma}$-null cone throughout (Section 11.1.2).

Definition 11.84 (Retarded TT mediator kernel)**.** Let $G^{\mathrm{ret},[\mathrm{TT}]}_{\mu\nu}{}^{\ \rho\sigma}(x,x')$ be the retarded Green operator of the TT wave operator on $(\mathcal{X}_q,\bar{\Sigma})$ (Section 11.2.3) and let $\hat{V}^{\mu\nu}(x)$ be the typed vertex (eq. (11.109)). The *retarded TT mediator kernel* is

$$\boxed{\mathcal{K}^{\mathrm{ret},[\mathrm{TT}]}(x,x') := \hat{V}^{\mu\nu}(x)\, G^{\mathrm{ret},[\mathrm{TT}]}_{\mu\nu}{}^{\ \rho\sigma}(x,x')\, \hat{V}_{\rho\sigma}(x'), \qquad \operatorname{supp}\mathcal{K}^{\mathrm{ret},[\mathrm{TT}]} \subseteq J^+_{\bar{\Sigma}},} \tag{11.138}$$

where $J^+_{\bar{\Sigma}}$ is the $\bar{\Sigma}$-causal future. Triadically, $\mathcal{K}^{\mathrm{ret},[\mathrm{TT}]} = \sum_a \mathcal{K}^{\mathrm{ret},[\mathrm{TT}],(a)}$ with $\mathcal{K}^{\mathrm{ret},[\mathrm{TT}],(a)}$ built from $G^{\mathrm{ret},[\mathrm{TT}],(a)}$ and $\hat{V}^{(a)}$ (Section 11.3.3).

Proposition 11.85 (Causal influence action for Tantrakyma)**.** *Integrating out the typed graviton at Gaussian order on $\mathfrak{P}$ yields the real-time (in–in) effective action for $\hat{\boldsymbol{\Theta}}$ with a causal nonlocal term*

$$\boxed{S^{\mathrm{eff}}_{\Theta} \;=\; S_{\Theta} + \frac{1}{2}\int d\mu(x)\, d\mu(x')\; \hat{\boldsymbol{\Theta}}(x)\, \mathcal{K}^{\mathrm{ret},[\mathrm{TT}]}(x,x')\, \hat{\boldsymbol{\Theta}}(x'),} \tag{11.139}$$

where $d\mu(x) = \sqrt{-\det\bar{\Sigma}(x)}\, d^d x$. The nonlocal kernel has $\bar{\Sigma}$-retarded support by (11.138); the IR window $K^{(\ell)}_{\mathrm{IR}}$ does not enlarge causal cones (Definition 6.280).

Corollary 11.86 (Dressed, causal equation of motion)**.** *Let $\mathcal{D}_\Theta$ denote the local (bare) Tantrakyma wave operator, so that $\mathcal{D}_\Theta\,\hat{\boldsymbol{\Theta}} = \hat{\mathcal{J}}$ in absence of gravity (Definition 10.208). Variation of (11.139) gives*

$$\boxed{\big(\mathcal{D}_\Theta\,\hat{\boldsymbol{\Theta}}\big)(x) + \int d\mu(x')\, \mathcal{K}^{\mathrm{ret},[\mathrm{TT}]}(x,x')\, \hat{\boldsymbol{\Theta}}(x') = \hat{\mathcal{J}}(x),} \tag{11.140}$$

a manifestly $\bar{\Sigma}$-causal, TT-mediated propagation law. Sectorwise, $\mathcal{K}^{\mathrm{ret},[\mathrm{TT}]} = \sum_a \mathcal{K}^{\mathrm{ret},[\mathrm{TT}],(a)}$.

Proposition 11.87 (Dyson equation for the retarded Tantrakyma propagator)**.** *Let $\Delta^{\mathrm{ret}}_{\Theta,0}$ be the bare retarded Tantrakyma propagator ($\mathcal{D}_\Theta \circ \Delta^{\mathrm{ret}}_{\Theta,0} = \mathbb{I}$) and $\Delta^{\mathrm{ret}}_{\Theta}$ the dressed one. Then on $\mathfrak{P}$*

$$\boxed{\big(\Delta^{\mathrm{ret}}_{\Theta}\big)^{-1} = \big(\Delta^{\mathrm{ret}}_{\Theta,0}\big)^{-1} - \Sigma^{\mathrm{ret}}_{\Theta,\mathrm{grav}}, \qquad \Sigma^{\mathrm{ret}}_{\Theta,\mathrm{grav}} \equiv \mathcal{K}^{\mathrm{ret},[\mathrm{TT}]},} \tag{11.141}$$

with $\Sigma^{\mathrm{ret}}_{\Theta,\mathrm{grav}}$ the retarded *TT self-energy generated by graviton exchange. Triadically, $\Sigma^{\mathrm{ret}}_{\Theta,\mathrm{grav}} = \sum_a \Sigma^{\mathrm{ret},(a)}_{\Theta,\mathrm{grav}}$.*

Remark 11.88 (Relation to time-ordered dressing)**.** The in–out (S-matrix) correction uses the Feynman kernel, $\Sigma^{\mathrm{F}}_{\Theta,\mathrm{grav}} = \hat{V}\, iG^{\mathrm{F},[\mathrm{TT}]}\, \hat{V}$ (Definition 11.54), reproducing the standard self-energy integrand of Definition 11.71. For causal response and propagation on $(\mathcal{X}_q,\bar{\Sigma})$, the appropriate kernel is the retarded one in (11.138); both agree on spectral data (KL densities) on the physical Hilbert space (Section 11.2.2).

Triadic Decomposition of the Self-Energy

A direct consequence of the typed, triad-resolved propagator (Definition 11.54 and eq. (11.86)) is that the gravitational dressing of Tantrakyma propagation splits cleanly by channel on the certified plateau $\mathfrak{P}$:

$$\boxed{\Delta\hat{G}_{\Theta\Theta}(x,x') = \sum_{a\in\{\mathrm{corr},\,\mathrm{vac},\,\mathrm{spec}\}} \Delta\hat{G}^{(a)}_{\Theta\Theta}(x,x').} \tag{11.142}$$

Each sectoral contribution is obtained by contracting the typed vertex (eqs. (11.109) and (11.122)) with the triad-resolved TT Feynman kernel (Definition 11.54 and Section 11.2.2):

$$\boxed{\Delta\hat{G}^{(a)}_{\Theta\Theta}(x,x') = \int d\mu_{\bar{\Sigma}}(z)\, d\mu_{\bar{\Sigma}}(z')\, \Big\langle 0\Big|\, T_{\bar{\Sigma}}\Big\{\hat{\boldsymbol{\Theta}}(x)\, \hat{V}^{\mu\nu}_{(a)}(z)\, \hat{V}^{\rho\sigma}_{(a)}(z')\, \hat{\boldsymbol{\Theta}}(x')\Big\}\, \Big|0\Big\rangle\, i\, G^{\mathrm{F},[\mathrm{TT}],(a)}_{\mu\nu,\rho\sigma}(z,z').} \tag{11.143}$$

Here $T_{\bar{\Sigma}}$ is the $\bar{\Sigma}$-time-ordering (Definition 11.53), and causal support of $G^{\mathrm{F},[\mathrm{TT}],(a)}$ is governed by the $\bar{\Sigma}$-null cone (Section 11.2.3 and Proposition 11.14). Nielsen flatness and hyperbolicity on $\mathfrak{P}$ ensure regulator- and gauge-parameter independence of the decomposition (Ch. 8).

Physical reading of the three terms.

1. *Correlational dressing* $\Delta\hat{G}^{(\mathrm{corr})}_{\Theta\Theta}$: self-energy induced by coupling to the information-theoretic sector (entanglement/coherence), sourced via $\hat{V}^{\mu\nu}_{(\mathrm{corr})}$ (Section 11.3.3).

2. *Vacuum dressing* $\Delta\hat{G}^{(\mathrm{vac})}_{\Theta\Theta}$: self-energy from polarizing the vacuum manifold $\mathcal{M}_{\mathrm{vac}}$ through $\hat{V}^{\mu\nu}_{(\mathrm{vac})}$ (Section 11.1.3).

3. *Spectral dressing* $\Delta\hat{G}^{(\mathrm{spec})}_{\Theta\Theta}$: the familiar energy–momentum–driven contribution via $\hat{V}^{\mu\nu}_{(\mathrm{spec})}$, reproducing the semiclassical limit in the spectral channel (Section 11.2.3).

Thus the observed (dressed) matter propagator acquires channel-wise corrections consistent with the triadic structure of the gravitational mediator (Section 11.2.4).

11.3.12 Graviton-Mediated Tantrakyma Scattering

We compute scattering on $\mathfrak{P}$ with $\bar{\Sigma}$-time ordering (Definition 11.53) and typed vertices (Section 11.3.2).

Definition 11.89 (Graviton–mediated S-matrix element)**.** For asymptotic Tantrakyma states $|\Theta_{\mathrm{in}}\rangle$, $|\Theta_{\mathrm{out}}\rangle$, the connected graviton-mediated amplitude is

$$\boxed{\mathcal{A}_{\mathrm{grav}}\big[\Theta_{\mathrm{in}}\to\Theta_{\mathrm{out}}\big] := \Big\langle\Theta_{\mathrm{out}}\Big|\, T_{\bar{\Sigma}}\exp\Big(i\, S^{(1)}_{\mathrm{int}}[\hat{h},\hat{\boldsymbol{\Theta}}]\Big)\,\Big|\Theta_{\mathrm{in}}\Big\rangle_{\mathrm{conn}},} \tag{11.144}$$

with $S^{(1)}_{\mathrm{int}}$ the linearized, TT-typed interaction (eqs. (11.109) and (11.110)).

Tree-level, single-graviton exchange. Expanding (11.144) to second order and Wick contracting the TT graviton fields yields the tree graph.

Lemma 11.90 (Typed, triad-resolved tree amplitude)**.** *The single-exchange contribution decomposes by triad as*

$$\boxed{\mathcal{A}^{(2)}_{\rm grav} = -\frac{1}{2}\sum_{a\in\{\rm corr,\,vac,\,spec\}}\int d\mu_{\bar\Sigma}(x)\,d\mu_{\bar\Sigma}(y)\;\mathcal{J}^{\mu\nu}_{(a)}(x)\;i\,G^{\mathrm{F},[\mathrm{TT}],(a)}{}_{\mu\nu}{}^{\rho\sigma}(x,y)\;\mathcal{J}^{(a)}_{\rho\sigma}(y),}\tag{11.145}$$

where $\mathcal{J}^{\mu\nu}_{(a)}(x) := \langle\Theta_{\rm out}|\hat V^{\mu\nu}_{(a)}(x)|\Theta_{\rm in}\rangle$ *are typed, sector-resolved matrix elements of the vertex (Section 11.3.3), and* $G^{\mathrm{F},[\mathrm{TT}],(a)}$ *are the TT Feynman kernels in each sector (Definition 11.54 and eq. (11.102)).*

Equation (11.145) is block-diagonal in the triad and manifestly covariant/causal: time ordering is with respect to the $\bar\Sigma$-cone; the IR window and band tags preserve microcausality and commute with the typed projectors (Definitions 6.280 and 9.10 and Proposition 11.14). Gauge-parameter independence (Nielsen flatness) and hyperbolicity of the TT principal symbol are part of plateau certification (Ch. 8).

The Triadic Nature of Gravitational Scattering Because the typed graviton propagator decomposes by sector (eq. (11.86)) and the sector projectors commute with the TT projector on the certified plateau $\mathfrak{P}$ (Sections 11.1.3 and 11.3.3), every graviton–mediated amplitude splits into a block–diagonal sum:

$$\boxed{\mathcal{A}_{\rm grav} = \sum_{a\in\{\rm corr,\,vac,\,spec\}}\mathcal{A}^{(a)}_{\rm grav}.}\tag{11.146}$$

At tree level the sectoral component is the typed TT contraction of currents with the sector–resolved Feynman kernel (Lemma 11.90):

$$\boxed{\mathcal{A}^{(a)}_{\rm grav} = -\frac{1}{2}\int d\mu_{\bar\Sigma}(x)\,d\mu_{\bar\Sigma}(y)\;\mathcal{J}^{\mu\nu}_{(a)}(x)\;i\,G^{\mathrm{F},[\mathrm{TT}],(a)}{}_{\mu\nu}{}^{\rho\sigma}(x,y)\;\mathcal{J}^{(a)}_{\rho\sigma}(y)\;+\;(\text{loops}).}\tag{11.147}$$

Here $\mathcal{J}^{\mu\nu}_{(a)}(x) = \langle\Theta_{\rm out}|\hat V^{\mu\nu}_{(a)}(x)|\Theta_{\rm in}\rangle$ are the typed, triad–resolved vertex matrix elements (Section 11.3.3 and eq. (11.109)), and $G^{\mathrm{F},[\mathrm{TT}],(a)}$ are the $\bar\Sigma$–time–ordered, TT–projected propagators in each sector (Definition 11.54 and Section 11.2.1). Causality and microcausality follow from the $\bar\Sigma$–null support of the typed kernels (Section 11.2.3 and Proposition 11.14), and band/IR operations preserve these cones (Definitions 6.280 and 9.10).

Sectoral interpretation.

- *Correlational channel* (a = corr): exchange of TT quanta sourced by informational structure (entanglement/coherence); see Section 11.2.4.

- *Vacuum channel* ($a = \mathrm{vac}$): exchange of TT Goldstone–like modes tied to the vacuum manifold; masslessness in the SSB limit is ensured by Proposition 11.29.
- *Spectral channel* ($a = \mathrm{spec}$): the energy–momentum–driven exchange reproducing the linearized GR limit (Section 11.2.3).

Coherence and mixing. On $\mathfrak{P}$ the internal graviton lines are block–diagonal in the triad, so there is no cross–channel interference at the free–field (tree) level (Lemma 11.90). Possible mixing can only enter via explicit sector–mixing structures in higher–order vertices or loop corrections, and remains TT–typed by construction (Section 11.3.2).

Generating functional view. The compact expression $\mathcal{A}_{\mathrm{grav}}[\Theta_{\mathrm{in}} \to \Theta_{\mathrm{out}}]$ (eq. (11.144)) is the generating functional for all graviton–exchange processes; expanding the Dyson series yields tree and loop graphs with internal lines $G^{\mathrm{F},[\mathrm{TT}],(a)}$ in each channel. In the forward limit, sector–by–sector Källén–Lehmann positivity of the TT two–point function (Section 11.2.2) supports dispersion relations and sum–rule constraints applied independently to each $\mathcal{A}^{(a)}_{\mathrm{grav}}$.

Lemma 11.91 (Dyson expansion (typed, TT, and triad-resolved))**.** *For $2 \to 2$ Tantrakyma scattering on the certified plateau $\mathfrak{P}$, the leading graviton–mediated contribution arises from the second term of the Dyson series built from the linearized interaction $S^{(1)}_{\mathrm{int}} = \int d\mu_{\bar{\Sigma}}(x)\, \hat{h}_{\mu\nu}(x)\, \hat{V}^{\mu\nu}(x)$ (eqs. (11.109) and (11.110)):*

$$\boxed{\mathcal{A}^{(2)}_{\Theta\Theta\to\Theta\Theta} = \frac{i^2}{2!} \left\langle \Theta_f \middle| T_{\bar{\Sigma}} \left\{ \left(\int d\mu_{\bar{\Sigma}}(x)\, \hat{h}_{\mu\nu}(x)\, \hat{V}^{\mu\nu}(x) \right)^2 \right\} \middle| \Theta_i \right\rangle_{\mathrm{conn}}.} \tag{11.148}$$

Equivalently—after Wick contracting the two graviton fields into the typed Feynman kernel $iG^{\mathrm{F},[\mathrm{TT}]}$ (Definition 11.54 and Section 11.2.1)—the tree-level exchange amplitude is

$$\boxed{\mathcal{A}^{(2)}_{\Theta\Theta\to\Theta\Theta} = -\frac{1}{2} \int d\mu_{\bar{\Sigma}}(x)\, d\mu_{\bar{\Sigma}}(y) \left\langle \Theta_f \middle| T_{\bar{\Sigma}} \{ \hat{V}^{\mu\nu}(x)\, \hat{V}^{\rho\sigma}(y) \} \middle| \Theta_i \right\rangle i\, G^{\mathrm{F},[\mathrm{TT}]}_{\mu\nu}{}^{\rho\sigma}(x,y).} \tag{11.149}$$

On $\mathfrak{P}$ the triad projectors commute with the TT projector (Sections 11.1.3 and 11.3.3), so the internal line decomposes by sector, $iG^{\mathrm{F},[\mathrm{TT}]} = \sum_a iG^{\mathrm{F},[\mathrm{TT}],(a)}$, yielding $\mathcal{A}^{(2)} = \sum_a \mathcal{A}^{(2),(a)}$ (eq. (11.146)).

Proof via Wick's theorem. Start from the Dyson term (11.148), expand the square to a double integral over x, y, and move the integrals outside the matrix element by linearity of $T_{\bar{\Sigma}}$:

$$\mathcal{A}^{(2)} = \frac{i^2}{2} \int d\mu_{\bar{\Sigma}}(x)\, d\mu_{\bar{\Sigma}}(y) \left\langle \Theta_f \middle| T_{\bar{\Sigma}} \{ \hat{h}_{\mu\nu}(x) \hat{V}^{\mu\nu}(x)\, \hat{h}_{\rho\sigma}(y) \hat{V}^{\rho\sigma}(y) \} \middle| \Theta_i \right\rangle_{\mathrm{conn}}.$$

Apply Wick's theorem: since there are no external gravitons, the only non-vanishing connected contraction pairs the two $\hat{h}$'s, producing the typed Feynman propagator $i\, G^{\mathrm{F},[\mathrm{TT}]}_{\mu\nu}{}^{\rho\sigma}(x,y) = {}_{\Phi}\langle 0|\, T_{\bar{\Sigma}} \{ \hat{h}_{\mu\nu}(x) \hat{h}^{\rho\sigma}(y) \} |0\rangle_{\Phi}$ (Definition 11.54). The remaining operators reorganize into $T_{\bar{\Sigma}} \{ \hat{V}^{\mu\nu}(x) \hat{V}^{\rho\sigma}(y) \}$. This yields (11.149). On $\mathfrak{P}$, triad projectors commute with $\Pi^{[\mathrm{TT}]}$ and band tags (Section 11.3.3), so $G^{\mathrm{F},[\mathrm{TT}]} = \sum_a G^{\mathrm{F},[\mathrm{TT}],(a)}$ and the amplitude splits as a sectoral sum, as claimed. □

11.3.13 Positivity/Dispersion Bounds for Triadic Graviton Exchange

We derive forward–limit dispersion constraints for elastic $2\to2$ scattering mediated by spin–2 exchange, resolved *triadically* by the commuting sector projectors P_χ with $\chi \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$ (cf. the sectoral/TT projectors in Section 11.1.3 and the triadic decomposition of the graviton in Section 11.1.4). To avoid a clash with the interferometric overlap $\mathcal{M}$ used elsewhere, we denote the Lorentz–invariant QFT scattering amplitude by $\mathscr{M}$.

Forward dispersion and triadic split. Let $\mathscr{M}^{(2)}_{ab\to ab}(s,t)$ be the crossing–even part of the Lorentz–invariant amplitude in the TT (spin–2) channel isolated by the projector of Definition 11.23, with Mandelstam invariants s,t and forward limit $t\to0$. Inserting the sector projectors into the TT exchange kernel yields a triadic decomposition *at the absorptive level*:

$$\mathscr{M}^{(2)}_{ab\to ab}(s,t) = \sum_\chi \mathscr{M}^{(2),\chi}_{ab\to ab}(s,t), \qquad \mathrm{Im}\,\mathscr{M}^{(2)} = \sum_\chi \mathrm{Im}\,\mathscr{M}^{(2),\chi}, \tag{11.150}$$

with $P_\chi P_{\chi'} = \delta_{\chi\chi'}P_\chi$ and $\sum_\chi P_\chi = \mathbb{I}$ (Section 11.1.3). Unitarity (optical theorem), microcausality (Peierls support on $\bar\Sigma$–null cones), and analyticity/crossing then imply a twice–subtracted forward dispersion relation with a positive spectral integrand for each sector [82, 14, 531, 255, 409]:

$$\boxed{\frac{\partial^2}{\partial s^2}\,\mathrm{Re}\,\mathscr{M}^{(2)}_{ab\to ab}(s,0) \;\geq\; 0, \qquad \rho^{(2)}_{\mathrm{tot}}(\mu^2) = \rho^{(2)}_{\mathrm{corr}}(\mu^2) + \rho^{(2)}_{\mathrm{vac}}(\mu^2) + \rho^{(2)}_{\mathrm{spec}}(\mu^2),} \tag{11.151}$$

where each sectoral TT spectral density obeys the Källén–Lehmann (KL) positivity $\rho^{(2)}_\chi(\mu^2)\geq 0$, being the (typed, TT) two–point density built from the corresponding graviton projector (cf. Definition 11.51 and the typed Feynman kernel in Definition 11.54).

KL representation, residues, and sum rules. In momentum space the typed, TT propagator in sector χ admits a KL form [82, 14]:

$$i\,G^{\mathrm{F},[\mathrm{TT}],\chi}_{\mu\nu}{}^{\rho\sigma}(k) = \int_0^\infty d\mu^2\, \frac{\rho^{(2)}_\chi(\mu^2)}{k^2-\mu^2+i\epsilon}\, \Pi^{[\mathrm{TT}]}_{\mu\nu}{}^{\rho\sigma}(k;\bar\Sigma), \tag{11.152}$$

with $\rho^{(2)}_\chi(\mu^2)\geq0$ and the TT projector $\Pi^{[\mathrm{TT}]}$ built from the orthonormal spin–2 basis (Section 11.1.3). If a narrow pole at $\mu^2=m_\chi^2$ is present, the corresponding residue $Z_\chi \equiv \lim_{k^2\to m_\chi^2}(k^2-m_\chi^2)\,G^{\mathrm{F},[\mathrm{TT}],\chi}$ is non–negative. Triadic completeness implies additivity of both the spectral densities and the pole residues:

$$\boxed{\rho^{(2)}_{\mathrm{tot}} = \sum_\chi \rho^{(2)}_\chi, \qquad Z_{\mathrm{tot}} = \sum_\chi Z_\chi, \qquad Z_\chi \geq 0.} \tag{11.153}$$

Forward–limit positivity (sector–resolved). Combining (11.91) with standard forward dispersion for the crossing–even amplitude gives, for any subtraction point $s_\star$ in the analytic

domain,

$$\boxed{\frac{\partial^2}{\partial s^2}\,\mathrm{Re}\,\mathscr{M}^{(2),\chi}_{ab\to ab}(s_\star,0)=\frac{2}{\pi}\int_{s_{\rm th}}^{\infty}ds'\,\frac{\mathrm{Im}\,\mathscr{M}^{(2),\chi}_{ab\to ab}(s',0)}{(s'-s_\star)^3}\;\geq\;0,}\tag{11.154}$$

with $s_{\rm th}$ the sectoral threshold. The inequality follows from the optical theorem $\mathrm{Im}\,\mathscr{M}^{(2),\chi}\propto\sum_X|\langle X|T^{[\mathrm{TT}],\chi}|ab\rangle|^2\geq 0$ and thus from $\rho^{(2)}_\chi\geq 0$.

Connection to TT state–visibility on $\mathfrak{P}$. On the certified plateau $\mathfrak{P}$ (band tags fixed, BRST cohomology taken, and Nielsen–flatness enforced; cf. Section 11.1.5), the TT *state–visibility* diagnostic is governed by the sectoral KL weights in (11.91). Any loss of positivity (e.g. a negative Z_χ or negative region of $\rho^{(2)}_\chi$) would manifest as a violation of forward–limit bounds (11.154) and thus signal a breakdown of physicality on $\mathfrak{P}$ (microcausality/unitarity).

EFT consequences. Expanding the forward amplitude at low s produces Wilson coefficients in the triad–resolved effective interaction (Section 11.3.4). The inequalities (11.154) then translate into model–independent sign constraints and size bounds on those coefficients, with the sectoral additivity (11.153) correlating the three channels. These results mirror standard EFT positivity arguments adapted to the typed, TT, triad–resolved IQR setting [82, 14, 531].

A. Setup: Forward amplitudes with triadic spin–2 exchange. Work in the TT sector singled out by Definition 11.23; resolve the exchange using the commuting sector projectors P_χ of Section 11.1.3. Consider elastic scattering of identical, neutral Tantrakyma probes (treated as scalars for definiteness) with $s=(p_1+p_2)^2$, $t=(p_1-p_3)^2$, $u=4m^2-s-t$, and define the crossing–even combination $\mathscr{M}_+(s,t)=\frac{1}{2}\big[\mathscr{M}(s,t)+\mathscr{M}(u,t)\big]$ so that $\mathscr{M}_+(s,0)=\mathscr{M}_+(-s,0)$. Stress–tensor Ward identities on the BRST–physical space guarantee analyticity and the optical theorem, furnishing the forward–limit framework used above [82, 14, 531].

Relational regularization and the forward limit. To control IR/UV issues intrinsically in the emergent geometry $(\mathcal{X}_q,\bar\Sigma)$, perform dispersion at fixed state–conditioned band on $\mathfrak{P}$ (coherent, relational regularization), and take the forward limit $t\to 0$ after establishing (11.151). This preserves the typed microcausal structure and ensures each sectoral contribution $\mathscr{M}^{(2),\chi}$ is well–defined in the forward analysis (cf. Section 11.2.1 and Definition 11.54).

Lemma 11.92 (Commutativity of TT and triadic projectors on the physical sector)**.** *Let $\widehat{T}_{\mu\nu}(x)$ be the conserved stress tensor (Ward/BRST identities), and let $W_{\mu\nu;\rho\sigma}(x,y)$ denote its two–point distribution in the BRST/GNS–physical state (Sections 1.4.1 and 12.2). Let $\Pi^{[\mathrm{TT}]\,\alpha\beta}_{\bar\Sigma\,\mu\nu}$ be the transverse–traceless (spin–2) projector (Definition 11.23 and Section 11.1.3), acting on the symmetric tensor indices with respect to the background $\bar\Sigma_{\mu\nu}$, and let $\{P_\chi\}_{\chi\in\{\mathrm{corr,vac,spec}\}}$ be the orthogonal sector projectors from the Unified Inter–Modal Correlator (Definition 6.4), acting on the triadic (internal) fibre. Then, as operator–valued distributions on test data of the form $f^{\mu\nu}(x)\otimes\mathbf{v}$ with $f\in\mathcal{D}(\mathcal{X}_q)$ and $\mathbf{v}$ in the triadic fibre,*

$$\big(\Pi^{[\mathrm{TT}]}_{\bar\Sigma}\otimes\mathbb{I}\big)\big(\mathbb{I}\otimes P_\chi\big)\,W\;=\;\big(\mathbb{I}\otimes P_\chi\big)\big(\Pi^{[\mathrm{TT}]}_{\bar\Sigma}\otimes\mathbb{I}\big)\,W,\tag{11.155}$$

in the sense of distributions modulo contact terms that vanish under (i) smearing with conserved sources and (ii) LSZ amputation in the forward limit. Equivalently, the TT–projected, triadically resolved correlators are well-defined and unique:

$$W^{[\mathrm{TT}],\chi} := \left(\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \otimes P_\chi\right) W = \left(\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \otimes \mathbb{I}\right)(\mathbb{I} \otimes P_\chi)\, W = (\mathbb{I} \otimes P_\chi)\left(\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \otimes \mathbb{I}\right) W. \quad (11.156)$$

Proof. (1) *Tensor–product structure.* $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ acts only on space–time tensor indices, whereas P_χ acts only on the finite–dimensional triadic fibre (Definition 6.4). On the algebraic tensor product $\mathcal{D}^{(2)}_{\mathrm{sym}}(\mathcal{X}_q) \,\widehat{\otimes}\, \mathbb{C}^3$ they commute algebraically: $(\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \otimes \mathbb{I})(\mathbb{I} \otimes P_\chi) = (\mathbb{I} \otimes P_\chi)(\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \otimes \mathbb{I})$.

(2) *Physical sector and BRST closure (plateau $\mathfrak{P}$).* On the BRST/GNS–physical sector (Section 12.2) and on the certified plateau $\mathfrak{P}$ (band tags fixed, Nielsen–flatness), P_χ is BRST–closed and orthogonal ($P_\chi P_{\chi'} = \delta_{\chi\chi'} P_\chi$), so inserting P_χ before or after any operation on space–time indices leaves the triadic block invariant.

(3) *Transversality and contact terms.* Ward identities imply $\bar{\nabla}^\mu W_{\mu\nu;\rho\sigma} = 0$ as a distribution (compatibly with Definitions 11.53 and 11.54). Longitudinal pieces killed by $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ can arise only as contact terms (derivatives of δ), which vanish when smeared against conserved test tensors $f^{\mu\nu}$ ($\bar{\nabla}_\mu f^{\mu\nu} = 0$), and drop out after LSZ amputation in the forward limit used for dispersion. Hence (11.155) holds in the physically relevant distributional sense.

(4) *Domain and continuity.* $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ is a local differential projector (momentum–space multiplier), and P_χ is a bounded idempotent on the triadic fibre; both are continuous endomorphisms of the test space and extend by duality to the space of operator–valued distributions. The commutation therefore extends to W itself. □

Proposition 11.93 (Typed–sectoral projection commuting on the physical band)**.** *On the certified plateau $\mathfrak{P}$, the background–covariant TT projector $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ (Definition 11.23) and the triadic sector projectors $\Pi^{(a)}$ with $a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$ (Definitions 6.4 and 11.45)* commute *on operator–valued distributions with conserved insertions, modulo contact terms that vanish when paired with conserved sources and under LSZ amputation. Equivalently, for the TT two–point function W and any local fluctuation $\delta\hat{g}_{\mu\nu}$,*

$$(\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \otimes \mathbb{I})\,(\mathbb{I} \otimes \Pi^{(a)})\, W = (\mathbb{I} \otimes \Pi^{(a)})\,(\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \otimes \mathbb{I})\, W, \quad (11.157)$$

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \Pi^{(a)}\, \delta\hat{g}_{\mu\nu} = \Pi^{(a)}\, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \delta\hat{g}_{\mu\nu}. \quad (11.158)$$

Consequently, channel projection can be performed before or after *TT projection on fields and propagators; in particular,*

$$G^{[\mathrm{TT}],(a)} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \Pi^{(a)}\, G\, \Pi^{(a)}\, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} = \Pi^{(a)}\, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, G\, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \Pi^{(a)}. \quad (11.159)$$

Proof sketch. Immediate from Lemma 11.92 (TT/triadic commutativity on the physical sector), using BRST closure and orthogonality of $\Pi^{(a)}$, and noting that contact terms drop out for conserved sources. Band tags commute with both projectors on $\mathfrak{P}$ (Definition 9.10 and Proposition 11.25), so there is no leakage across types. □

Corollary 11.94 (Channel–wise TT spectral decomposition)**.** *The TT two–point function admits a Källén–Lehmann representation that is* triadically block–diagonal*:*

$$W^{(2)} = \sum_{\chi \in \{\mathrm{corr},\mathrm{vac},\mathrm{spec}\}} W^{(2),\chi}, \qquad W^{(2),\chi} := \left(\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \otimes P_\chi\right) W,$$

with non–negative sectoral spectral densities by KL positivity on the BRST/GNS–physical sector (Sections 11.1.3 and 11.3.6). Consequently, the channel–wise dispersion relations and positivity bounds derived in Section 11.3.6 apply separately to each $W^{(2),\chi}$.

Remark 11.95 (Gauge–fixing independence)**.** The statement is *gauge–parameter independent*: it is made on the BRST/GNS–physical sector, where longitudinal/gauge contributions are BRST–exact and drop out of W modulo contact terms. The decomposition and positivity are thus properties of physical correlators, not of any particular gauge choice.

B. Universal IR subtraction (massless t–channel pole). In each triadic TT sector $\chi \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$, massless spin–2 exchange produces the universal long–range Born singularity for small $|t|$:

$$\mathscr{M}_{+}^{(2),\chi}(s,t) \;=\; \frac{\kappa_\chi^2}{4}\,\frac{s^2}{-t} \;+\; \mathscr{M}_{+,\mathrm{reg}}^{(2),\chi}(s,t), \qquad \lim_{t\to 0} \mathscr{M}_{+,\mathrm{reg}}^{(2),\chi}(s,t) \text{ finite}, \tag{11.160}$$

where the IR–universal coupling κ_χ is fixed by Ward–type identities for the conserved $T_{\mu\nu}$ in sector χ (equivalence–principle normalization), and $\mathscr{M}_{+,\mathrm{reg}}^{(2),\chi}$ is analytic at $t = 0$.[1]
Define IR–subtracted forward amplitudes by removing only the universal Born piece and then taking the forward limit:

$$\widetilde{\mathscr{M}}_{+}^{(2),\chi}(s) \;:=\; \lim_{t\to 0^+}\left[\mathscr{M}_{+}^{(2),\chi}(s,t) \;-\; \frac{\kappa_\chi^2}{4}\,\frac{s^2}{-t}\right], \qquad \widetilde{\mathscr{M}}_{+}^{(2)}(s) \;:=\; \sum_{\chi} \widetilde{\mathscr{M}}_{+}^{(2),\chi}(s). \tag{11.161}$$

This subtraction preserves analyticity in s at fixed $t{=}0$ and is compatible with unitarity and crossing: it removes only the long–range t-channel singularity while leaving the absorptive part (and thus the dispersion kernel) unchanged [82, 14, 531]. In the relational scheme of Sections 4.9.13 and 4.9.15, (11.161) is implemented at fixed state–conditioned IR/UV band, and the limit $t \to 0^+$ is taken after subtraction, ensuring a well–defined forward amplitude sector by sector.

C. Channel–wise dispersion relation and positivity. Analyticity and growth (emergent background). Assume: (A1) microcausality with respect to the effective Lorentzian tensor $\bar{g}_{\mu\nu}^{\mathrm{Lor}}$ (Sections 5.3.4, 5.3.6 and 5.4); (A2) polynomial boundedness after the universal IR subtraction (Sections 4.9.13 and 4.9.15); (A3) BRST/GNS physical states with Hadamard short–distance form (Section 1.4.1); (A4) local, $\bar{\Sigma}$–covariant couplings with Ward identities isolating TT exchange. By Lemma 11.92, TT and triadic projectors commute on the physical two–point, so the decomposition involves no mixing. Then each sector's forward, crossing–even amplitude obeys the twice–subtracted dispersion relation [82, 14, 531]:

$$\begin{aligned}\mathrm{Re}\,\widetilde{\mathscr{M}}_{+}^{(2),\chi}(s) \;=\; &\widetilde{\mathscr{M}}_{+}^{(2),\chi}(0) \;+\; \tfrac{1}{2}\,s^2\,\widetilde{\mathscr{M}}_{+}^{(2),\chi\prime\prime}(0) \\ &+\; \frac{s^4}{\pi}\int_{s_{\mathrm{th}}^{\chi}}^{\infty} \mathrm{d}s'\;\frac{\mathrm{Im}\,\widetilde{\mathscr{M}}_{+}^{(2),\chi}(s')}{s'^4}\left(\frac{1}{s'-s}+\frac{1}{s'+s}\right),\end{aligned} \tag{11.162}$$

[1] If a state–dependent spectral gap $m_{\mathrm{eff}} > 0$ opens (cf. Section 4.9.15), the pole is $s^2/(-t-m_{\mathrm{eff}}^2)$. In (11.161) subtract $s^2/(-t-m_{\mathrm{eff}}^2)$; the massless form (11.160) is recovered as $m_{\mathrm{eff}} \downarrow 0$.

whence the forward convexity (optical theorem):

$$\boxed{\widetilde{\mathscr{M}}_{+}^{(2),\chi\,\prime\prime}(0) \;=\; \frac{2}{\pi}\int_{s_{\mathrm{th}}^{\chi}}^{\infty}\frac{\mathrm{d}s}{s^{3}}\,\mathrm{Im}\,\widetilde{\mathscr{M}}_{+}^{(2),\chi}(s) \;\geq\; 0} \tag{11.163}$$

equivalently, with $\rho_{\chi}^{(2)}(s)=(1/\pi)\,\mathrm{Im}\,\widetilde{\mathscr{M}}_{+}^{(2),\chi}(s)\geq 0$: $\widetilde{\mathscr{M}}_{+}^{(2),\chi\,\prime\prime}(0)=2\int_{s_{\mathrm{th}}^{\chi}}^{\infty}\rho_{\chi}^{(2)}(s)\,s^{-3}\mathrm{d}s\geq 0$.

D. Spectral representation and the triadic sum rule. Let $\Delta^{(2)}(q)$ be the TT propagator (or TT part of the stress–tensor exchange kernel). With $\Pi^{[\mathrm{TT}]}$ from Definition 11.23 and sector projectors P_{χ} from Definition 6.4 (which commute on the BRST/GNS physical two–point), define

$$\Delta^{(2),\chi}(q) \;:=\; \Pi^{[\mathrm{TT}]}\,P_{\chi}\,\Delta^{(2)}(q)\,P_{\chi}\,\Pi^{[\mathrm{TT}]}.$$

Then the triadically resolved Källén–Lehmann representation reads [82, 14]

$$\Delta^{(2),\chi}(q) \;=\; \frac{Z_{\chi}}{q^{2}+i\epsilon} \;+\; \int_{0^{+}}^{\infty}\mathrm{d}\mu^{2}\,\frac{\rho_{\chi}^{(2)}(\mu^{2})}{q^{2}-\mu^{2}+i\epsilon}, \qquad Z_{\chi}\geq 0,\quad \rho_{\chi}^{(2)}(\mu^{2})\geq 0, \tag{11.164}$$

and summing over χ (using $P_{\chi}P_{\chi'}=\delta_{\chi\chi'}P_{\chi}$, $\sum_{\chi}P_{\chi}=\mathbb{I}$) yields the *triadic spectral sum rule*

$$\boxed{\rho_{\mathrm{tot}}^{(2)}(\mu^{2}) \;=\; \rho_{\mathrm{corr}}^{(2)}(\mu^{2}) \;+\; \rho_{\mathrm{vac}}^{(2)}(\mu^{2}) \;+\; \rho_{\mathrm{spec}}^{(2)}(\mu^{2}), \qquad \rho_{\bullet}^{(2)}(\mu^{2})\geq 0\;.} \tag{11.165}$$

Inserting (11.165) into LSZ for the TT channel reproduces the channel–wise convexity bound (11.163) ($\widetilde{\mathscr{M}}_{+}^{(2),\chi\,\prime\prime}(0)\propto\int\rho_{\chi}^{(2)}(s)\,s^{-3}\mathrm{d}s\geq 0$).

E. Low–energy EFT bounds (constraints on triadic Wilson coefficients). Expand the IR–subtracted, forward, crossing–even amplitude in each sector (eq. (11.161)):

$$\widetilde{\mathscr{M}}_{+}^{(2),\chi}(s) \;=\; c_{0}^{(\chi)} \;+\; c_{2}^{(\chi)}\,s^{2} \;+\; c_{4}^{(\chi)}\,s^{4} \;+\cdots, \qquad (\text{even in } s \text{ by crossing}). \tag{11.166}$$

From (11.163) (twice–subtracted dispersion; universal t-channel pole removed) [82, 14, 531],

$$\begin{aligned} c_{2}^{(\chi)} &= \frac{1}{\pi}\int_{s_{\mathrm{th}}^{\chi}}^{\infty}\frac{\mathrm{d}s}{s^{3}}\,\mathrm{Im}\,\widetilde{\mathscr{M}}_{+}^{(2),\chi}(s) \;\geq\; 0,\\ c_{2n}^{(\chi)} &= \frac{(2n-3)!}{\pi}\int_{s_{\mathrm{th}}^{\chi}}^{\infty}\frac{\mathrm{d}s}{s^{2n-1}}\,\mathrm{Im}\,\widetilde{\mathscr{M}}_{+}^{(2),\chi}(s) \;\geq\; 0, \qquad n\geq 2. \end{aligned} \tag{11.167}$$

Matching to EFT. If heavy TT exchange induces the dimension–8 contact

$$\mathcal{L}_{\mathrm{eff}} \;\supset\; \sum_{\chi\in\{\mathrm{corr,vac,spec}\}}\frac{\alpha_{\chi}}{\Lambda^{4}}\left(J_{\mu\nu}^{(\chi)}J_{(\chi)}^{\mu\nu}\right), \tag{11.168}$$

then at tree level $\widetilde{\mathscr{M}}_{+}^{(2),\chi}(s)\;\supset\;\kappa_{\mathrm{EFT}}\,(\alpha_{\chi}/\Lambda^{4})\,s^{2}$ with $\kappa_{\mathrm{EFT}}>0$, and comparing with (11.166)–(11.167) gives

$$\boxed{\alpha_{\chi}\;\geq\;0 \qquad \text{for each}\;\;\chi\in\{\mathrm{corr,vac,spec}\}} \tag{11.169}$$

Proposition (Triadic forward positivity). *Let $\widetilde{\mathcal{M}}_{+}^{(2),\chi}(s)$ be the IR–subtracted, forward, crossing–even $2\to 2$ amplitude mediated by the TT spin–2 sector in channel χ. Assuming microcausality, analyticity, and polynomial boundedness with respect to $\bar{g}_{\mu\nu}^{\mathrm{Lor}}$,*

$$\frac{\partial^2}{\partial s^2}\,\mathrm{Re}\,\widetilde{\mathcal{M}}_{+}^{(2),\chi}(s)\Big|_{s=0} \;=\; \frac{2}{\pi}\int_{s_{\mathrm{th}}^{\chi}}^{\infty}\frac{\mathrm{d}s}{s^3}\,\mathrm{Im}\,\widetilde{\mathcal{M}}_{+}^{(2),\chi}(s) \;\geq\; 0.$$

Corollary (EFT coefficient bounds). *In any low–energy realization of Section 11.3.12, the coefficients multiplying forward s^{2n} contact structures sourced by triadic spin–2 exchange satisfy $c_{2n}^{(\chi)}\geq 0$. In particular, the leading dimension–8 triadic coefficients obey $\alpha_\chi\geq 0$ as in (11.169).*

Remarks. (i) *Universality of the subtraction.* The t–channel pole is fixed by Ward identities; subtracting it removes only the long–range piece and leaves the dispersive kernel unchanged [82, 14]. (ii) *Crossing/odd parts.* For non-identical external states, apply the twice–subtracted dispersion relation to the crossing–even part [531]. (iii) *Relational IR scheme.* Working at fixed state–conditioned IR floor (and coherent UV ceiling) as in Sections 4.9.13 and 4.9.15 yields the same non–negativity; removing regulators at the end commutes with the dispersion integrals by dominated convergence.

Interpretive Summary.

Triadic spectral positivity. By isolating the transverse–traceless (spin–2) exchange and resolving it with the sector projectors P_χ ($\chi=\mathrm{corr},\mathrm{vac},\mathrm{spec}$), the absorptive part in each channel is non-negative and admits a Källén–Lehmann density $\rho_\chi^{(2)}(\mu^2)\geq 0$. The total spectral weight is the sum of the three sectors, $\rho_{\mathrm{tot}}^{(2)}=\sum_\chi\rho_\chi^{(2)}$ (eq. (11.165)). This is precisely the spectral manifestation of unitarity and microcausality for spin–2 exchange [255, 230, 409, 82, 14].

Channel-wise forward convexity. After subtracting the universal long–range t-channel pole (eq. (11.161)), the twice-subtracted, crossing-even dispersion relation with a positive kernel (eq. (11.162)) implies the forward sum rule (eq. (11.163)). Geometrically, the real part of each sector's forward amplitude has non-negative curvature at $s=0$. Any putative negative curvature would contradict at least one S-matrix pillar (unitarity/analyticity/crossing) [82, 14, 531].

IR universality and relational finiteness. The only universal IR singularity is the massless t-channel pole fixed by Ward identities; its subtraction isolates the genuinely model-dependent remainder (eq. (11.161)). Within the relational regularization scheme—working at fixed, state-conditioned IR/UV band and taking the forward limit last—the conclusions are unchanged but the procedure is intrinsically finite (Section 4.9.15).

EFT sign constraints. The low-energy expansion of the IR-subtracted forward amplitude yields $c_{2n}^{(\chi)} \geq 0$ for all $n \geq 1$ (eqs. (11.166) and (11.167)). Matching to the dimension-8 contact $\left(J_{\mu\nu}^{(\chi)}\right)^2/\Lambda^4$ gives $\alpha_\chi \geq 0$ in each sector (eq. (11.169)). Thus any UV completion that induces triadic spin–2 contact terms must respect these sign constraints; a negative α_χ would signal a breakdown of the S-matrix axioms or of the spin–2 projection assumptions.

Bridge to interferometry. The interferometric bounds (Section 3.1.8) assert that coherence is partitioned triadically and cannot exceed an ℓ^2 envelope, while the dispersive bounds assert that absorptive probability is partitioned triadically and cannot be negative. Together they yield an instrument-to-field synthesis: visibility constrains *how* channels interfere; dispersion/positivity constrains *how* they dissipate.

In short, once the universal IR pole is removed, each triadic spin–2 channel carries positive spectral weight and enforces a convex forward amplitude, which in turn imposes *non-negativity* constraints on the corresponding low-energy triadic Wilson coefficients. These conclusions rest only on unitarity, analyticity, crossing, and Ward identities, and are expressed in a form directly usable for EFT matching and phenomenological consistency checks (Section 11.3.6).

11.4 Geometry–matter interface in the typed (TT) formalism.

The graviton is the typed fluctuation of geometry, namely the background–covariant transverse–traceless carrier $\hat{h}_{\mu\nu} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \delta\hat{g}_{\mu\nu}$ (Definition 11.3). Gravitational effects are accessed through their imprint on matter: the Tantrakyma sector provides calibrated probes and sources via the bi–local matter–geometry interaction, producing a triad–filtered vertex $\hat{V}_{\mu\nu}^{(a)}$ that couples linearly to the typed graviton in each sector $a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$ (Section 11.3 and Definition 11.24). At the linear level the typed equation of motion is

$$K_{\mu\nu}^{[\mathrm{TT}],(a)\alpha\beta}\, \hat{h}_{\alpha\beta}^{(a)} = \hat{J}_{\mu\nu}^{[\mathrm{TT}],(a)}, \qquad \hat{J}^{[\mathrm{TT}],(a)} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \hat{J}^{(a)},$$

so the geometric carrier $\hat{h}$ is driven by matter currents $\hat{J}^{(a)}$ through the typed Lichnerowicz–type operator $K^{[\mathrm{TT}],(a)}$ (Definitions 11.60 and 11.61). The TT Ward identities guarantee that only physical polarizations couple, $k_\mu\, \hat{V}^{[\mathrm{TT}],(a)\mu\nu} = 0$ and $\bar{\Sigma}_{\mu\nu}\, \hat{V}^{[\mathrm{TT}],(a)\mu\nu} = 0$, and that τ–typed insertions are TT–null (Proposition 11.77 and Corollary 11.73). Hence the Tantrakyma sector necessarily appears whenever the observable imprint of geometry is evaluated.

Two standard computations illustrate this structure. First, tree–level exchange between Tantrakyma sources proceeds through a typed graviton propagator sandwiched between typed vertices, yielding a sector–resolved amplitude that is block–diagonal in a (Definition 11.67 and Propositions 11.20 and 11.77). Second, integrating out the typed graviton at quadratic order produces a causal, nonlocal effective action for Tantrakyma fields,

$$S_{\mathrm{eff}}^{(\Theta)} = \frac{1}{2}\sum_a \int d\mu_{\bar{\Sigma}}(x)\, d\mu_{\bar{\Sigma}}(y)\, \mathrm{Tr}\Big[\hat{V}_{\mu\nu}^{(a)}(x)\, i\, G^{\mathrm{F},[\mathrm{TT}],(a)\,\mu\nu}{}_{\rho\sigma}(x,y)\, \hat{V}^{(a)\rho\sigma}(y)\Big] + \cdots,$$

which underlies the dispersion/positivity analyses (Sections 11.3.5 and 11.3.6 and eq. (11.115)). The same typed Feynman Green function $i\, G^{\mathrm{F},[\mathrm{TT}],(a)}$ governs the Tantrakyma two–point

dressing $\Delta\hat{G}_{\Theta\Theta}$ (Definitions 11.70 and 11.71) and, via its causal difference $\Delta^{[\mathrm{TT}]}$, the Peierls bracket and the covariant commutator (eqs. (11.25) and (11.26) and Proposition 11.14). In summary, the graviton originates in the geometric (Gaiamiti) sector, while its observable content is coupled to and diagnosed by Tantrakyma matter—both as a source in the typed equations of motion and as a probe in amplitudes and propagator dressings—within the BRST–healthy, plateau–stable TT framework (Proposition 7.3 and Corollary 11.5).

Synthesis and alignment with standard expectations. The typed, TT–projected graviton carrier $\hat{h}_{\mu\nu} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}_{\mu\nu}$ (Definition 11.3) obeys a Lichnerowicz–type wave equation sandwiched by the TT projector (Definition 11.60). Couplings to matter respect the typed Ward identities that enforce transversality and tracelessness (Proposition 11.77). The covariant algebra arises from the Peierls bracket built with the retarded/advanced Green operators of the same typed wave operator (eq. (11.25) and Proposition 11.14). At tree level, exchange and dressing take the familiar "vertex–propagator–vertex'' form, both in position space (Definition 11.67) and in momentum space (eq. (11.137)), and integrating out the typed graviton yields a causal, nonlocal current–current effective action for Tantrakyma fields (Section 11.3.5, eq. (11.115), and Definition 11.71). In this sense, the IQR formulation is fully consonant with linearized gravity and the EFT/QFT-in-curved-spacetime toolkit: physical polarizations are TT, the commutator descends from the causal bi-solution, and amplitudes are controlled by TT projectors and Green kernels [244, 7, 415, 502, 82, 91].

Typed/triadic machinery versus standard minimal gravity. What departs from the minimal textbook setup are the organizational devices introduced by IQR: the triad decomposition $a \in \{\mathrm{corr}, \mathrm{vac}, \mathrm{spec}\}$, band tags with certified plateaux, and Nielsen–flat diagnostics together with typed TLO overlays. These structures commute with the TT projector and remain BRST-clean on the certified plateau, so microcausality, the typed Ward identities, and the TT algebra are preserved (Propositions 7.3 and 10.9, Definition 6.280, and Corollary 11.5). They refine bookkeeping and regulator control without altering the physical TT content encoded in Definitions 11.3, 11.60, 11.67 and 11.71.

Masslessness, explicit deformations, and observational status. Within IQR the masslessness of the typed graviton follows from the symmetry/SSB structure in the TT channel (Proposition 11.29). A GMOR–style statement under a controlled explicit breaking produces a small mass deformation (Corollary 11.30), which is a *model* hypothesis akin to massive–gravity deformations rather than a property of vanilla GR. Such deformations must remain consistent with current bounds from gravitational–wave observations that find no deviation from pure-tensor GR radiation; multi–event analyses (GWTC–3) report limits consistent with a strictly massless spin–2 field and the absence of extra polarizations [122, 479]. Ghost–free massive/bimetric frameworks illustrate how tiny graviton masses can be arranged consistently, but they are tightly constrained [113, 481].

Conclusion. Consequently, the typed TT dynamics, Ward identities, Peierls construction, exchange amplitudes, and matter–sector dressings employed here (Definitions 11.3, 11.60, 11.67 and 11.71) are expected and standard in spirit. The triad/band/plateau/Nielsen

apparatus is compatible bookkeeping that preserves the TT physics while improving control over causality, gauge cleanliness, and regulators; any GMOR–type graviton–mass discussion is a permissible deformation rather than a GR prediction and is therefore subject to stringent data-informed bounds [122, 113, 481, 479].

11.4.1 The Feynman Rules for Matter and Geometry

The perturbative dynamics of the IQR, particularly the interactions between matter and geometry, can be systematically organized and calculated using a set of Feynman rules. These rules provide a graphical representation for the terms in the Dyson expansion of the S-matrix (Section 11.3.12). The fundamental components are the propagators for the fields and the vertices describing their interactions.

The Tantrakyma Propagator (Matter)

A matter-like excitation of the Tantrakyma field (of species 'p') propagating from spacetime point x to y is represented by a solid line.



Figure 11.1: Tantrakyma field propagator $\Delta_p(x, y)$, representing matter-like quanta propagating between spacetime points x and y in the emergent configuration space.

- *Diagrammatic Rule:* A solid line connecting points x and y.
- *Mathematical Expression:* This line corresponds to the *Tantrakyma field propagator*, $\Delta_p(x, y)$, which is derived from the matter-sector correlator $\hat{G}_{\Theta\Theta}$ (Definition 10.208).

The Graviton Propagator (Geometry)

A quantum fluctuation of the geometric field propagating from x to y is represented by a wavy line, the standard convention for a mediating gauge boson.



Figure 11.2: Full graviton propagator $G^{\text{grav}}_{\mu\nu\rho\sigma}(x, y)$, encoding quantum fluctuations of the spin-2 geometry field across spacetime.

- *Diagrammatic Rule:* A wavy line connecting points x and y.
- *Mathematical Expression:* This line corresponds to the *full graviton propagator*, $G^{\text{grav}}_{\mu\nu\rho\sigma}(x, y)$ (Section 11.2).

The Triadic Decomposition of the Graviton Propagator

A central prediction of the pqs framework is that the graviton is a composite object with three distinct relational origins. Therefore, any internal graviton line in a Feynman diagram must be treated as the sum of three separate channels.



Figure 11.3: Triadic decomposition of the graviton propagator into correlational ($G^{(\text{corr})}$), vacuum ($G^{(\text{vac})}$), and spectral ($G^{(\text{spec})}$) sectors. Each corresponds to a distinct origin mode of geometric excitation.

- *Diagrammatic Rule:* A single wavy graviton line is understood as the sum of three distinct types of propagation, which can be represented by different colors or styles.
- *Mathematical Expression:* The total propagator is the sum of the triadic components (eq. (11.86)):

  $$G^{\text{grav}}_{\mu\nu\rho\sigma}(x,y) = G^{(\text{corr})}_{\mu\nu\rho\sigma}(x,y) + G^{(\text{vac})}_{\mu\nu\rho\sigma}(x,y) + G^{(\text{spec})}_{\mu\nu\rho\sigma}(x,y)$$

  When calculating a scattering amplitude, one must sum over all diagrams involving each of these three distinct internal graviton lines.

The Graviton-Matter Interaction Vertex

The fundamental interaction between matter and geometry is represented by a three-point vertex where two Tantrakyma field lines meet a single graviton line.



Figure 11.4: Graviton–matter interaction vertex, coupling two Tantrakyma fields (solid lines) to a graviton (wavy line) via the vertex operator $\hat{V}^{\mu\nu}(x)$.

- *Diagrammatic Rule:* A point where two solid lines (Tantrakyma) and one wavy line (graviton) meet.
- *Mathematical Expression:* This vertex corresponds to a factor of $i\hat{V}^{\mu\nu}(x)$, the *graviton-Tantrakyma vertex operator* defined in Definition 11.64. It represents the "gravitational charge" of the matter field.

11.5 The Chronogeometric Gravity

The preceding chapters fix the IQR realized weave configuration and establish how a physical chronogeometry emerges from an operator substrate via diagnostic realized reference configuration (band–limited, BRST–clean) expectations (Definitions 6.280 and 9.10 and Proposition 7.3). The unified geometric carrier is the *Chronogeometric Tensor* (Definition 5.26), defined as the expectation of the operator-valued kernel $\hat{\Sigma}_{\mu\nu}$ in the Quantum Holo-State (Definition 1.27):

$$\bar{\Sigma}_{\mu\nu}(x) := \Big\langle \mathring{\boldsymbol{\Psi}} \Big| \hat{\Sigma}_{\mu\nu}(x) \Big| \mathring{\boldsymbol{\Psi}} \Big\rangle = \bar{g}_{\mu\nu}(x) + \bar{t}_{\mu\nu}(x),$$

the standard background split used throughout (Section 5.2.6). The tensor $\bar{\Sigma}_{\mu\nu}$ determines both the causal cones and the differential geometry on $(\mathcal{X}_q, \bar{\Sigma})$; in particular, it fixes the Relational Levi–Civita connection $\bar{\nabla} \equiv \nabla^{(\mathrm{rel})}$ (Definition 6.7) and its inverse tensor $\bar{\Sigma}^{\mu\nu}$ used to raise indices (Definition 6.24). The two pieces encode:

- classical distance and causal structure via $\bar{g}_{\mu\nu}$ (background metric block; Section 5.2.6), and
- coherence/phase–flow via the temporal/coherence block $\bar{t}_{\mu\nu}$ (expectation of $\hat{t}_{\mu\nu}$; Definition 5.17).

Unified source (expectation level). Curvature in IQR is sourced by a *unified* stress–energy constructed from geometry (Gaiamiti), matter (Tantrakyma), and their interaction, as organized by the total relational action (Definition 9.357). The operator and scalar (fiber–traced) stress tensors are defined in Section 9.17.9, Definition 9.356, and eq. (9.224), with the interaction piece fixed variationally (eq. (9.237)). On a certified plateau $\mathfrak{P}$ with the physical projector $\mathsf{P}_{\mathrm{phys}} : \mathcal{H} \to H^0(\hat{Q}_{\mathrm{BRST}})$ the *expectation–level* total source is (Definition 9.358)

$$\boxed{\bar{T}^{\mu\nu}_{\mathrm{total}}(x) := \Big\langle \mathring{\boldsymbol{\Psi}} \Big| \mathsf{P}_{\mathrm{phys}}\, \mathrm{Tr}\Big\{ \hat{T}^{\mu\nu}_{(\Phi)}(x) + \hat{T}^{\mu\nu}_{(\Theta)}(x) + \hat{T}^{\mu\nu}_{(\mathrm{int})}(x) \Big\} \Big| \mathring{\boldsymbol{\Psi}} \Big\rangle,} \tag{11.170}$$

which is precisely the diagnostic realized reference configuration form emphasized in eqs. (9.230) and (9.238). Concretely,

- $\hat{T}^{\mu\nu}_{(\Phi)}$ is the Gaiamiti Noether tensor (geometry sector) with kernel form given in eq. (9.235);
- $\hat{T}^{\mu\nu}_{(\Theta)}$ is the Tantrakyma Noether tensor (matter sector) as in Definition 9.356 and eq. (9.236);
- $\hat{T}^{\mu\nu}_{(\mathrm{int})}$ follows from the variation of S_{int} with respect to $\bar{\Sigma}_{\mu\nu}$ (eq. (9.237));
- $\big|\mathring{\boldsymbol{\Psi}}\big\rangle$ is the Quantum Holo-State entering all diagnostic realized reference configuration expectations (Definition 1.27).

By linearity of the expectation (diagnostic realized reference configuration semantics), additivity holds as stated in eqs. (9.233) and (9.234).

Self-consistency and conservation. The unified Hilbert variation yields the unique geometric source and the on–shell conservation law (eq. (9.232)):

$$\boxed{\nabla_\mu^{(\mathrm{rel})}\,\bar{T}^{\mu\nu}_{\mathrm{total}}(x)=0,} \tag{11.171}$$

with separate sectoral conservation in the absence of exchange ($\hat{T}^{\mu\nu}_{(\mathrm{int})}=0$), and conservation of the sum otherwise (Section 9.18.12). Thus, the chronogeometric field equations in IQR equate two *emergent* objects on $(\mathcal{X}_q,\bar{\Sigma})$: relational curvature (built from $\bar{\Sigma}$ and $\bar{\nabla}$; Section 5.2.6) and the diagnostic realized reference configuration–projected total source (11.170).

Remark 11.96 (Temporal/coherence curvature as macroscopic quantum structure)**.** The temporal/coherence block $\bar{t}_{\mu\nu}(x)$ is the expectation of a well–defined operator $\hat{t}_{\mu\nu}(x)$ (Definition 5.17), not an ad hoc metric deformation. It encodes how local phase evolution and coherence flow are *curved* by the state, thereby modifying geodesic transport defined by $\bar{\nabla}$ (Definition 6.7). In particular, $\bar{t}_{\mu\nu}$ supports coherence–driven focusing/defocusing and entanglement–induced curvature consistent with the unified source mechanism (Section 9.18.12) and with the additivity/split formulas (eqs. (9.233) and (9.238)).

Instrumentation preamble (diagnostic realized reference configuration, projectors, and readout). All derivations and readouts in this section are performed on a certified plateau $\mathfrak{P}$ with BRST–closed band tags $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}=(P_x(\lambda_{\mathrm{UV}}),K^{(\ell)}_{\mathrm{IR}})$ that commute with the triad projectors $\Pi^{(\mathrm{corr})},\Pi^{(\mathrm{vac})},\Pi^{(\mathrm{spec})}$ and with the background–covariant TT projector $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ (Definitions 6.280, 9.10, 10.3 and 11.24 and eq. (10.4)). Observables are evaluated on the physical cohomology $\mathcal{H}_{\mathrm{phys}}:=H^0(\hat{Q}_{\mathrm{BRST}})$ (diagnostic realized reference configuration projection $\mathsf{P}_{\mathrm{phys}}$), and all law–channel variations are Nielsen–flat on $\mathfrak{P}$ (Proposition 7.3 and Corollary 11.5). In particular, the Einstein equations are kept verbatim as the central law and will be presented in the BRST/cohomological and band–regular form, with a TT–typed TLO panel attached only for radiative gravity (Definitions 11.6 and 11.106 and eq. (11.191)).

Relational geometry, conservation, and causal support. All connections and curvatures are those of the realized chronogeometry $(\mathcal{X}_q,\bar{\Sigma})$: indices are raised with $\bar{\Sigma}^{\mu\nu}$, and covariant differentiation is by the Relational Levi–Civita connection $\bar{\nabla}\equiv\nabla^{(\mathrm{rel})}$ (Definitions 5.26, 6.7 and 6.24). The unified Bianchi identity $\bar{\nabla}^\mu\bar{G}_{\mu\nu}\equiv 0$ is read on the same background and is matched by the expectation–level conservation law $\bar{\nabla}^{(\mathrm{rel})}_\mu\,\bar{T}^{\mu\nu}_{\mathrm{total}}=0$ (Section 9.18.12 and eq. (9.232)). Linearization employs the TT–sandwiched Lichnerowicz–type operator and its Σ–*causal* advanced/retarded Green kernels, ensuring microcausality on the very cone used elsewhere in the treatise (Definition 11.60, eq. (11.25), and Proposition 11.14). The physical–polarization content is carried by the typed TT graviton, while non–TT "observables'' (e.g. gauge–condition or triad–mixed readouts) are explicitly identified; TT purity and orthogonality follow the Chapter 11 projector calculus and sectoral factorization (Definitions 11.3 and 11.19 and Propositions 11.20 and 11.77).

11.5.1 The Unified Riemann Curvature Tensor

All geometric objects in this section are built from the realized chronogeometric background $(\mathcal{X}_q, \bar{\Sigma})$: indices are raised with $\bar{\Sigma}^{\mu\nu}$ and covariant derivatives are taken with the Relational Levi–Civita connection $\bar{\nabla} \equiv \nabla^{(\mathrm{rel})}$, which is torsion–free and $\bar{\Sigma}$–compatible (Definitions 5.26, 6.7 and 6.24 and Section 5.2.6). The split $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (Section 5.2.6 and Definition 5.17) will be used to exhibit the metric and phase/coherence contributions to curvature.

Definition 11.97 (Riemann tensor of $(\mathcal{X}_q, \bar{\Sigma})$)**.** For any vector field v^λ, the curvature is defined by the commutator

$$\boxed{\left(\bar{\nabla}_\mu \bar{\nabla}_\nu - \bar{\nabla}_\nu \bar{\nabla}_\mu\right) v^\lambda \;=\; \bar{R}^\lambda{}_{\rho\mu\nu}[\bar{\Sigma}]\; v^\rho,} \tag{11.172}$$

equivalently, in a coordinate chart with Christoffel symbols $\Gamma^\lambda{}_{\mu\nu}[\bar{\Sigma}]$ of $\nabla^{(\mathrm{rel})}$ (Definition 6.7),

$$\boxed{\bar{R}^\lambda{}_{\rho\mu\nu}[\bar{\Sigma}] = \partial_\mu \Gamma^\lambda{}_{\nu\rho} - \partial_\nu \Gamma^\lambda{}_{\mu\rho} + \Gamma^\lambda{}_{\mu\alpha}\Gamma^\alpha{}_{\nu\rho} - \Gamma^\lambda{}_{\nu\alpha}\Gamma^\alpha{}_{\mu\rho}.} \tag{11.173}$$

Contractions and Einstein tensor (chronogeometric). The Ricci and scalar curvatures of $\bar{\Sigma}$ are

$$\boxed{\bar{R}_{\mu\nu}[\bar{\Sigma}] := \bar{R}^\rho{}_{\mu\rho\nu}[\bar{\Sigma}], \qquad \bar{R}[\bar{\Sigma}] := \bar{\Sigma}^{\mu\nu}\, \bar{R}_{\mu\nu}[\bar{\Sigma}],} \tag{11.174}$$

and the chronogeometric Einstein tensor is

$$\boxed{\bar{G}_{\mu\nu}[\bar{\Sigma}] := \bar{R}_{\mu\nu}[\bar{\Sigma}] - \tfrac{1}{2}\,\bar{\Sigma}_{\mu\nu}\,\bar{R}[\bar{\Sigma}],} \tag{11.175}$$

which obeys the contracted Bianchi identity $\bar{\nabla}^\mu \bar{G}_{\mu\nu} \equiv 0$ by $\bar{\nabla}$–compatibility of $\bar{\Sigma}$ (Definition 6.7). The unified semi–classical Einstein law is then stated verbatim as $\bar{G}_{\mu\nu}[\bar{\Sigma}] + \Lambda\,\bar{\Sigma}_{\mu\nu} = \kappa\,\bar{T}^{\mathrm{total}}_{\mu\nu}$ (Definition 11.106, eq. (11.191), and Section 9.18.12).

Proposition 11.98 (Symmetries and Bianchi identities)**.**

$$\boxed{\begin{aligned} \bar{R}_{\lambda\rho\mu\nu} &= -\,\bar{R}_{\rho\lambda\mu\nu} && \text{(11.176a)}\\ \bar{R}_{\lambda\rho\mu\nu} &= -\,\bar{R}_{\lambda\rho\nu\mu} && \text{(11.176b)}\\ \bar{R}_{\lambda\rho\mu\nu} &= \bar{R}_{\mu\nu\lambda\rho} && \text{(11.176c)}\\ \bar{R}_{\lambda[\rho\mu\nu]} &= 0 && \text{(11.176d)}\\ \bar{\nabla}_{[\sigma}\bar{R}_{\lambda\rho]\mu\nu} &= 0 && \text{(11.176e)} \end{aligned}}$$

Remark 11.99 (Meaning and applicability of the curvature symmetries)**.** Throughout Proposition 11.98 we work with the unified (Relational) Levi–Civita connection $\bar{\nabla}$ of $\bar{\Sigma}$ (metric compatible and torsion–free), as in Definition 11.97. Under these assumptions, each line in (11.176) has the standard geometric meaning and domain of validity:

(i) Pair antisymmetries.
(11.176a)–(11.176b). They express the skewness of the curvature two–form in each index pair; when indices are lowered with the metric (metric compatibility), one gets $\bar{R}_{\lambda\rho\mu\nu} = -\bar{R}_{\rho\lambda\mu\nu}$ and $\bar{R}_{\lambda\rho\mu\nu} = -\bar{R}_{\lambda\rho\nu\mu}$ [248, 84, 316].

(ii) Major (pair–exchange) symmetry.
(11.176c). For a metric–compatible, torsion–free connection (Levi–Civita) one has $\bar{R}_{\lambda\rho\mu\nu} = \bar{R}_{\mu\nu\lambda\rho}$; this uses metric compatibility in an essential way [248, Ch. III][84, Sec. 6][316, Sec. 3.3].

(iii) First (algebraic) Bianchi identity.
(11.176d). The cyclic sum $\bar{R}_{\lambda[\rho\mu\nu]} = 0$ is the algebraic identity that holds for torsion–free connections; with torsion it acquires additional $T\cdot T$ and $\bar{\nabla}T$ terms. In the Levi–Civita case it takes the simple form shown [248, Ch. III, §7][375, Sec. 7][316, Sec. 3.3].

(iv) Second (differential) Bianchi identity.
(11.176e). The differential identity $\bar{\nabla}_{[\sigma}\bar{R}_{\lambda\rho]\mu\nu} = 0$ holds for the Levi–Civita connection; in the presence of torsion it generalizes by extra torsion–curvature couplings. Contracting indices yields the *contracted Bianchi identity* $\bar{\nabla}^{\mu}\bar{G}_{\mu\nu} = 0$ (with $\bar{G}_{\mu\nu}$ from eq. (11.175)), which underlies covariant conservation of the effective source in the field equations [146, §B.2][316, Sec. 3.4][375, Sec. 7].

These statements are standard in Riemannian/Lorentzian geometry and are presented here in the unified $\bar{\Sigma}$–formalism; see the cited references for proofs and for the torsion/non–metricity generalizations.

Remark 11.100 (Metric part + coherence part via connection difference)**.** Let $\Gamma[\bar{g}]$ be the Levi–Civita connection of the metric block $\bar{g}_{\mu\nu}$ and define the difference tensor $C^{\lambda}{}_{\mu\nu} := \Gamma^{\lambda}{}_{\mu\nu}[\bar{\Sigma}] - \Gamma^{\lambda}{}_{\mu\nu}[\bar{g}]$. Then

$$\boxed{\bar{R}^{\lambda}{}_{\rho\mu\nu}[\bar{\Sigma}] = R^{\lambda}{}_{\rho\mu\nu}[\bar{g}] + \nabla^{(\bar{g})}_{\mu}C^{\lambda}{}_{\nu\rho} - \nabla^{(\bar{g})}_{\nu}C^{\lambda}{}_{\mu\rho} + C^{\lambda}{}_{\mu\alpha}C^{\alpha}{}_{\nu\rho} - C^{\lambda}{}_{\nu\alpha}C^{\alpha}{}_{\mu\rho},} \tag{11.177}$$

where $\nabla^{(\bar{g})}$ is the Levi–Civita derivative of $\bar{g}$. The tensor $C^{\lambda}{}_{\mu\nu}$ is determined by the temporal/coherence block $\bar{t}_{\mu\nu}$ through $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (Section 5.2.6 and Definition 5.17). Thus the "metric curvature" $R[\bar{g}]$ and the "phase–flow twist" driven by $\bar{t}$ combine to yield the full chronogeometric curvature $\bar{R}[\bar{\Sigma}]$. In the pure–metric limit $\bar{t}_{\mu\nu} \to 0$, one recovers $C \to 0$ and $\bar{R}[\bar{\Sigma}] \to R[\bar{g}]$.

Cohomological form and law residual (context). On a certified plateau $\mathfrak{P}$ with BRST–closed tags (Definitions 6.280 and 9.10), the Einstein law may also be recorded in the cohomological form $\mathcal{E}^{\mathrm{coh}}_{\mathbf{\Phi}}(\bar{\Sigma}) = 0$, $\mathcal{E}^{\mathrm{coh}}_{\mathbf{\Theta}}(\bar{\Sigma}) = 0$ modulo ζ–exact terms and evaluated with Σ–causal retarded kernels (Section 9.18.12 and Propositions 7.3 and 11.14). For diagnostic realized reference configuration–level auditing, define the untyped, global Einstein–law residual and visibility

$$\boxed{\mathbf{R}^{[\mathrm{Ein}]}_{\mu\nu} := \bar{G}_{\mu\nu}[\bar{\Sigma}] + \Lambda\,\bar{\Sigma}_{\mu\nu} - \kappa\,\bar{T}^{\mathrm{total}}_{\mu\nu}, \qquad \Lambda^{[\mathrm{Ein}]}_{\mathrm{law}} := 1 - \frac{\|\mathbf{R}^{[\mathrm{Ein}]}\|_{\mathcal{I}}}{\|\cdot\|_{\mathrm{ref}}},} \tag{11.178}$$

interpreted using the plateau meters and synthesis logic of Chapter 9 (Definitions 9.299 and 11.6 and eqs. (9.126) to (9.128)). Typed TT diagnostics remain reserved for the radiative sector as in Chapters 10–11 (Definitions 11.3, 11.6 and 11.60).

Derivation of the Unified Riemann Curvature Tensor

1. Motivation: curvature as non-commuting derivatives. Work on the realized chronogeometric background $(\mathcal{X}_q, \bar{\Sigma})$, with indices raised by $\bar{\Sigma}^{\mu\nu}$ and covariant differentiation by the Relational Levi–Civita connection $\bar{\nabla} \equiv \nabla^{(\mathrm{rel})}$, which is torsion–free and $\bar{\Sigma}$–compatible (Definitions 6.7 and 6.24 and Section 5.2.6). Curvature is the obstruction to the commutativity of covariant derivatives. Acting on a vector field V^λ,

$$\boxed{\left[\bar{\nabla}_\mu, \bar{\nabla}_\nu\right] V^\lambda := \bar{\nabla}_\mu(\bar{\nabla}_\nu V^\lambda) - \bar{\nabla}_\nu(\bar{\nabla}_\mu V^\lambda),} \tag{11.179}$$

and the goal is to compute the right–hand side explicitly and identify the Riemann tensor of $(\mathcal{X}_q, \bar{\Sigma})$ introduced in Definition 11.97 (components in eq. (11.173)).

2. Calculation of the commutator. Using the coordinate expression for the covariant derivative of a vector, $\bar{\nabla}_\nu V^\lambda = \partial_\nu V^\lambda + \Gamma^\lambda{}_{\nu\sigma}[\bar{\Sigma}]\, V^\sigma$ (Definition 6.7),

$$\bar{\nabla}_\mu(\bar{\nabla}_\nu V^\lambda) = \bar{\nabla}_\mu(\partial_\nu V^\lambda) + \bar{\nabla}_\mu(\Gamma^\lambda{}_{\nu\sigma} V^\sigma).$$

Treat $\partial_\nu V^\lambda$ as a mixed tensor $T^\lambda{}_\nu$ and apply the Leibniz rule:

$$\begin{aligned}
\bar{\nabla}_\mu(\partial_\nu V^\lambda) &= \partial_\mu\partial_\nu V^\lambda + \Gamma^\lambda{}_{\mu\rho}\,\partial_\nu V^\rho - \Gamma^\rho{}_{\mu\nu}\,\partial_\rho V^\lambda, \\
\bar{\nabla}_\mu(\Gamma^\lambda{}_{\nu\sigma} V^\sigma) &= (\partial_\mu \Gamma^\lambda{}_{\nu\sigma})\, V^\sigma + \Gamma^\lambda{}_{\nu\sigma}\,\bar{\nabla}_\mu V^\sigma = (\partial_\mu \Gamma^\lambda{}_{\nu\sigma})\, V^\sigma + \Gamma^\lambda{}_{\nu\sigma}\Big(\partial_\mu V^\sigma + \Gamma^\sigma{}_{\mu\rho} V^\rho\Big).
\end{aligned}$$

Hence

$$\begin{aligned}
\bar{\nabla}_\mu(\bar{\nabla}_\nu V^\lambda) = {} & \partial_\mu\partial_\nu V^\lambda + \Gamma^\lambda{}_{\mu\rho}\,\partial_\nu V^\rho - \Gamma^\rho{}_{\mu\nu}\,\partial_\rho V^\lambda \\
& + (\partial_\mu \Gamma^\lambda{}_{\nu\sigma})\, V^\sigma + \Gamma^\lambda{}_{\nu\sigma}\,\partial_\mu V^\sigma + \Gamma^\lambda{}_{\nu\sigma}\Gamma^\sigma{}_{\mu\rho}\, V^\rho.
\end{aligned}$$

Interchanging $\mu \leftrightarrow \nu$ yields $\bar{\nabla}_\nu(\bar{\nabla}_\mu V^\lambda)$:

$$\begin{aligned}
\bar{\nabla}_\nu(\bar{\nabla}_\mu V^\lambda) = {} & \partial_\nu\partial_\mu V^\lambda + \Gamma^\lambda{}_{\nu\rho}\,\partial_\mu V^\rho - \Gamma^\rho{}_{\nu\mu}\,\partial_\rho V^\lambda \\
& + (\partial_\nu \Gamma^\lambda{}_{\mu\sigma})\, V^\sigma + \Gamma^\lambda{}_{\mu\sigma}\,\partial_\nu V^\sigma + \Gamma^\lambda{}_{\mu\sigma}\Gamma^\sigma{}_{\nu\rho}\, V^\rho.
\end{aligned}$$

Subtracting and using (i) equality of mixed partials on smooth fields, (ii) torsion–free symmetry $\Gamma^\rho{}_{\mu\nu} = \Gamma^\rho{}_{\nu\mu}$ (Definition 6.7), and (iii) relabelling of dummy indices, the terms with ∂V cancel and one finds

$$\left[\bar{\nabla}_\mu, \bar{\nabla}_\nu\right] V^\lambda = \Big[\partial_\mu \Gamma^\lambda{}_{\nu\rho} - \partial_\nu \Gamma^\lambda{}_{\mu\rho} + \Gamma^\lambda{}_{\mu\sigma}\Gamma^\sigma{}_{\nu\rho} - \Gamma^\lambda{}_{\nu\sigma}\Gamma^\sigma{}_{\mu\rho}\Big] V^\rho.$$

By Definition 11.97 and eq. (11.173), the bracket equals $\bar{R}^\lambda{}_{\rho\mu\nu}[\bar{\Sigma}]\, V^\rho$, hence

$$\boxed{\left[\bar{\nabla}_\mu, \bar{\nabla}_\nu\right] V^\lambda = \bar{R}^\lambda{}_{\rho\mu\nu}[\bar{\Sigma}]\; V^\rho,} \tag{11.180}$$

which is the commutator (operator) definition of the unified Riemann tensor of $(\mathcal{X}_q, \bar{\Sigma})$ and is equivalent to the component formula in eq. (11.173).

Checks and continuity. The symmetry and Bianchi identities (Proposition 11.98) follow from torsion–freeness and $\bar{\nabla}\bar{\Sigma}=0$ (Definition 6.7). Decomposing the connection by $\Gamma[\bar{\Sigma}] = \Gamma[\bar{g}] + C(\bar{t})$ gives the metric/coherence split of curvature (Remark 11.100 and eq. (11.177)); in the pure–metric limit $\bar{t}_{\mu\nu} \to 0$ one recovers $C \to 0$ and $\bar{R}[\bar{\Sigma}] \to R[\bar{g}]$, maintaining continuity with the constructions of Chapters 6–7.

Lemma 11.101 (Linearization and Σ–causal Green operators)**.** *Linearize the unified Einstein law $\bar{G}_{\mu\nu}[\bar{\Sigma}] + \Lambda\,\bar{\Sigma}_{\mu\nu} = \kappa\,\bar{T}^{\text{total}}_{\mu\nu}$ (Definition 11.106 and eq. (11.191)) around the realized background $(\mathcal{X}_q, \bar{\Sigma})$ with $\bar{\nabla} \equiv \nabla^{(\text{rel})}$ (Definition 6.7). The linearized operator $\mathcal{E}_{\mu\nu}{}^{\rho\sigma}[\bar{\Sigma}]$ acting on symmetric rank–2 fluctuations $h_{\rho\sigma}$ is defined by*

$$\boxed{\mathcal{E}_{\mu\nu}{}^{\rho\sigma}[\bar{\Sigma}]\; h_{\rho\sigma} := \frac{\delta}{\delta\bar{\Sigma}_{\rho\sigma}}\Big(\bar{G}_{\mu\nu}[\bar{\Sigma}] + \Lambda\,\bar{\Sigma}_{\mu\nu}\Big)\Big|_{\bar{\Sigma}} h_{\rho\sigma}, \qquad \bar{\nabla}^{\mu}\Big(\mathcal{E}_{\mu\nu}{}^{\rho\sigma} h_{\rho\sigma}\Big) = 0.} \tag{11.181}$$

Projecting to spin–2, transverse–traceless type with $\Pi^{[\text{TT}]}_{\bar{\Sigma}}$ (Definitions 10.3 and 11.3 and eq. (10.4)), the typed wave operator is

$$\boxed{K^{[\text{TT}]}_{\mu\nu}{}^{\alpha\beta} := \Big[\Pi^{[\text{TT}]}_{\bar{\Sigma}} \circ \mathcal{E}[\bar{\Sigma}] \circ \Pi^{[\text{TT}]}_{\bar{\Sigma}}\Big]_{\mu\nu}{}^{\alpha\beta} \;\equiv\; K^{[\text{TT}]}_{\mu\nu}{}^{\alpha\beta} \text{ of eq. (11.103).}} \tag{11.182}$$

For the TT fluctuation $h_{\alpha\beta}$ and TT–projected source $J^{[\text{TT}]}_{\mu\nu}$ (defined as in Definition 11.61), the linearized dynamics read

$$\boxed{K^{[\text{TT}]}_{\mu\nu}{}^{\alpha\beta}\, h_{\alpha\beta} = J^{[\text{TT}]}_{\mu\nu}, \qquad J^{[\text{TT}]}_{\mu\nu} := \Pi^{[\text{TT}]}_{\bar{\Sigma}\,\mu\nu}{}^{\rho\sigma}\, J_{\rho\sigma}.} \tag{11.183}$$

The retarded/advanced Green operators $G^{\text{ret/adv},[\text{TT}]}$ are the two–sided inverses of $K^{[\text{TT}]}$ on the TT subspace,

$$\boxed{K^{[\text{TT}]} \circ G^{\text{ret/adv},[\text{TT}]} = \Pi^{[\text{TT}]}_{\bar{\Sigma}}, \qquad G^{\text{ret/adv},[\text{TT}]} \circ K^{[\text{TT}]} = \Pi^{[\text{TT}]}_{\bar{\Sigma}},} \tag{11.184}$$

with support constrained to the $\bar{\Sigma}$–causal cones, $\operatorname{supp} G^{\text{ret/adv},[\text{TT}]} \subset J^{\pm}_{\bar{\Sigma}}$ (Definition 6.7 and Proposition 11.14). Their causal difference $\Delta^{[\text{TT}]} := G^{\text{ret},[\text{TT}]} - G^{\text{adv},[\text{TT}]}$ generates the Peierls bracket and the covariant commutator (eqs. (11.25) and (11.26)), and the Feynman kernel $i\,G^{\text{F},[\text{TT}]}$ solves the typed Green equation (Definition 11.54 and eq. (11.107)).

Remark 11.102 (Microcausal cone and continuity across chapters)**.** The TT retarded/advanced kernels in (11.184) inherit the *same* $\bar{\Sigma}$–microcausal cone used for scalar kernels and their Stieltjes representations in the kinematic constructions of Chapters 6–7, thereby providing a direct bridge to the dynamical Peierls framework of Chapter 8 (Definition 6.7, Proposition 11.14, and eq. (11.25)).

11.5.2 Ricci Tensor and Scalar Curvature of the Unified Geometry

All contractions are taken on the realized chronogeometric background $(\mathcal{X}_q, \bar{\Sigma})$, with indices raised by $\bar{\Sigma}^{\mu\nu}$ and covariant derivatives given by the Relational Levi–Civita connection $\bar{\nabla} \equiv \nabla^{(\mathrm{rel})}$ (Definitions 6.7 and 6.24 and Section 5.2.6). Starting from the unified Riemann tensor $\bar{R}^{\lambda}{}_{\rho\mu\nu}[\bar{\Sigma}]$ (Definition 11.97 and eq. (11.173)), the standard traces yield the curvature quantities that enter the chronogeometric Einstein tensor (eq. (11.175)) and the unified semi-classical law (Definition 11.106 and eq. (11.191)).

Definition 11.103 (Ricci tensor and scalar curvature of $(\mathcal{X}_q, \bar{\Sigma})$)**.** The *Ricci tensor* is the contraction of the Riemann tensor on its first and third indices,

$$\boxed{\bar{R}_{\mu\nu}[\bar{\Sigma}] \;:=\; \bar{R}^{\lambda}{}_{\mu\lambda\nu}[\bar{\Sigma}],} \tag{11.185}$$

and the *scalar curvature* is the full trace with the inverse chronogeometric tensor,

$$\boxed{\bar{R}[\bar{\Sigma}] \;:=\; \bar{\Sigma}^{\mu\nu}\,\bar{R}_{\mu\nu}[\bar{\Sigma}].} \tag{11.186}$$

Remark 11.104 (Properties and role in the field equations)**.** By the symmetries and Bianchi identities of $\bar{R}_{\lambda\rho\mu\nu}$ (Proposition 11.98), the Ricci tensor is symmetric, $\bar{R}_{\mu\nu} = \bar{R}_{\nu\mu}$, and the contracted Bianchi identity $\bar{\nabla}^{\mu}\bar{G}_{\mu\nu} \equiv 0$ follows once the Einstein tensor $\bar{G}_{\mu\nu} = \bar{R}_{\mu\nu} - \frac{1}{2}\bar{\Sigma}_{\mu\nu}\bar{R}$ is formed (eq. (11.175)). These objects provide the geometric side of the unified semi-classical Einstein equations, $\bar{G}_{\mu\nu}[\bar{\Sigma}] + \Lambda\,\bar{\Sigma}_{\mu\nu} = \kappa\,\bar{T}^{\mathrm{total}}_{\mu\nu}$ (Definition 11.106 and eq. (11.191)), with conservation read on the same background, $\bar{\nabla}^{(\mathrm{rel})}_{\mu}\,\bar{T}^{\mu\nu}_{\mathrm{total}} = 0$ (eq. (9.232)).

Derivation of the Ricci tensor formula (unified background). Starting from the unified Riemann tensor of $(\mathcal{X}_q, \bar{\Sigma})$ (Definition 11.97 and eq. (11.173)) and using the Relational Levi–Civita connection $\bar{\nabla} \equiv \nabla^{(\mathrm{rel})}$ (Definition 6.7) with indices raised by $\bar{\Sigma}^{\mu\nu}$ (Definition 6.24), the Ricci tensor is the contraction on the first and third slots: $\bar{R}_{\mu\nu}[\bar{\Sigma}] = \bar{R}^{\lambda}{}_{\mu\lambda\nu}[\bar{\Sigma}]$ (eq. (11.185)). Writing $\bar{R}^{\lambda}{}_{\rho\mu\nu} = \partial_\mu\Gamma^{\lambda}{}_{\nu\rho} - \partial_\nu\Gamma^{\lambda}{}_{\mu\rho} + \Gamma^{\lambda}{}_{\mu\alpha}\Gamma^{\alpha}{}_{\nu\rho} - \Gamma^{\lambda}{}_{\nu\alpha}\Gamma^{\alpha}{}_{\mu\rho}$ and setting $\rho \to \lambda$ yields, term by term,

$$\partial_\mu\Gamma^{\lambda}{}_{\nu\lambda}, \quad -\partial_\nu\Gamma^{\lambda}{}_{\mu\lambda}, \quad +\Gamma^{\lambda}{}_{\mu\alpha}\Gamma^{\alpha}{}_{\nu\lambda}, \quad -\Gamma^{\lambda}{}_{\nu\alpha}\Gamma^{\alpha}{}_{\mu\lambda}.$$

Collecting them gives the standard Christoffel form of the unified Ricci tensor:

$$\boxed{\bar{R}_{\mu\nu}[\bar{\Sigma}] = \partial_\mu\Gamma^{\lambda}{}_{\nu\lambda} - \partial_\nu\Gamma^{\lambda}{}_{\mu\lambda} + \Gamma^{\lambda}{}_{\mu\alpha}\Gamma^{\alpha}{}_{\nu\lambda} - \Gamma^{\lambda}{}_{\nu\alpha}\Gamma^{\alpha}{}_{\mu\lambda},} \tag{11.187}$$

which is equivalent to the contraction definition eq. (11.185) and is consistent with the symmetry/Bianchi identities in Proposition 11.98. The scalar curvature then follows by the $\bar{\Sigma}$–trace $\bar{R}[\bar{\Sigma}] = \bar{\Sigma}^{\mu\nu}\bar{R}_{\mu\nu}[\bar{\Sigma}]$ (eq. (11.186)), completing the contractions that enter the chronogeometric Einstein tensor (eq. (11.175)).

Physical interpretation and significance (chronogeometric reading).

- *Richer measure of curvature.* In classical GR, $\mathrm{Ric}(g)$ governs volume focusing and tidal deformation. In the IQR setting the unified Ricci tensor $\bar{R}_{\mu\nu}[\bar{\Sigma}]$ measures focusing in the *chronogeometric* sense: because $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (Section 5.2.6), its contractions probe both metric curvature (via $\bar{g}$) and the coherence/phase–flow contribution encoded by $\bar{t}$ (expectation of $\hat{t}_{\mu\nu}$, Definition 5.17). The connection difference representation (Remark 11.100 and eq. (11.177)) makes this explicit by splitting $\bar{R}[\bar{\Sigma}]$ into a metric part $R[\bar{g}]$ and a "twist'' built from the temporal/coherence block.

- *Gravitational Lagrangian and variational law.* The unified scalar curvature $\bar{R}[\bar{\Sigma}]$ is the simplest scalar built from $\bar{\Sigma}$ and $\bar{\nabla}$ and thus plays the role of the gravitational Lagrangian density. With the relational volume element $d\mu(x) = \sqrt{-\det \bar{\Sigma}(x)}\, d^d x$ (Definition 6.33), the Einstein–Hilbert–type relational action reads

$$S_{\mathrm{EH}}^{(\mathrm{rel})}[\bar{\Sigma}] := \int_{\mathcal{X}_q} d\mu(x)\ \bar{R}[\bar{\Sigma}].$$

  Its variation (together with a cosmological term) produces the geometric side of the unified law, $\bar{G}_{\mu\nu}[\bar{\Sigma}] + \Lambda\, \bar{\Sigma}_{\mu\nu} = \kappa\, \bar{T}^{\mathrm{total}}_{\mu\nu}$ (eqs. (11.175) and (11.191) and Definition 11.106), with conservation read on the same background $\bar{\nabla}^{(\mathrm{rel})}_{\mu}\, \bar{T}^{\mu\nu}_{\mathrm{total}} = 0$ (Section 9.18.12 and eq. (9.232)).

- *Source of new physics and the GR limit.* The difference $\bar{R}[\bar{\Sigma}] - R[\bar{g}]$ is entirely controlled by the temporal/coherence sector through the difference tensor $C(\bar{t})$ in eq. (11.177). When $\bar{t}_{\mu\nu} \to 0$, one has $C \to 0$ and $\bar{R}[\bar{\Sigma}] \to R[\bar{g}]$, recovering GR. In regimes of high coherence or strong temporal curvature (non-negligible $\bar{t}$), the additional terms modify focusing, dispersion, and exchange, feeding back consistently into the unified source $\bar{T}^{\mathrm{total}}_{\mu\nu}$ defined at diagnostic realized reference configuration level (Definition 9.358 and eq. (9.238)) and into the typed radiative sector through the linearized TT operator and its causal kernels (Definition 11.60 and Proposition 11.14).

These unified curvature objects thus furnish the geometric backbone of the emergent Einstein equations, anchoring the left–hand side on the same chronogeometric background, cones, and conservation structure used throughout Chapters 6–11 (Definitions 5.26 and 6.7 and Section 9.18.12).

11.5.3 The Relational Einstein–Hilbert Action

Diagnostic realized reference configuration and covariance context. All statements below are made on a certified plateau $\mathfrak{P}$ with BRST–closed band tags $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell} = (P_x(\lambda_{\mathrm{UV}}), K^{(\ell)}_{\mathrm{IR}})$ that commute with the triad and TT projectors (Definitions 6.280, 9.10, 10.3 and 11.24). Geometric objects are built from the realized chronogeometry $(\mathcal{X}_q, \bar{\Sigma})$, with indices raised by $\bar{\Sigma}^{\mu\nu}$ and covariant derivatives given by the Relational Levi–Civita connection $\bar{\nabla} \equiv \nabla^{(\mathrm{rel})}$ (Definitions 5.26, 6.7 and 6.24). The relational volume element is $d\mu(x) := \sqrt{-\det \bar{\Sigma}(x)}\, d^d x$ (Definition 6.33).

Definition 11.105 (Relational Einstein–Hilbert functional (diagnostic realized reference configuration, band–regularized))**.** The pure–geometry action is the scalar invariant built from the unified scalar curvature $\bar{R}[\bar{\Sigma}]$ (Definition 11.103) and the relational measure:

$$\boxed{S_{\mathrm{EH}}^{(\mathrm{rel})}[\bar{\Sigma}] \;:=\; \frac{1}{2\kappa}\int_{\mathcal{X}_q} d\mu(x)\; \bar{R}[\bar{\Sigma}] \;-\; \frac{\Lambda}{\kappa}\int_{\mathcal{X}_q} d\mu(x) \;+\; S_{\mathrm{GHY}}^{(\mathrm{rel})}[\partial\mathcal{X}_q],} \tag{11.188}$$

where κ is the gravitational coupling, Λ is the cosmological constant, and $S_{\mathrm{GHY}}^{(\mathrm{rel})}$ is the relational Gibbons–Hawking–York boundary term (needed for a well–posed Dirichlet variational problem on $\partial\mathcal{X}_q$) [532, 533, 534, 535]. The scalar curvature and its contractions are those of the unified geometry (Definition 11.97 and eqs. (11.175), (11.185) and (11.186)).

Variation and field equations (cohomological/diagnostic realized reference configuration forms). Varying (11.188) with respect to $\bar{\Sigma}_{\mu\nu}$ and discarding the boundary variation by $S_{\mathrm{GHY}}^{(\mathrm{rel})}$ gives the geometric Euler–Lagrange identity

$$\boxed{\delta S_{\mathrm{EH}}^{(\mathrm{rel})} = \frac{1}{2\kappa}\int d\mu\,\Big(\bar{G}_{\mu\nu}[\bar{\Sigma}] + \Lambda\,\bar{\Sigma}_{\mu\nu}\Big)\,\delta\bar{\Sigma}^{\mu\nu},} \tag{11.189}$$

with $\bar{\nabla}^{\mu}\bar{G}_{\mu\nu} \equiv 0$ by the contracted Bianchi identity (eq. (11.175)). Coupling to the total relational action $S_{\mathrm{total}} = S_{\Phi} + S_{\Theta} + S_{\mathrm{int}}$ (Definition 9.357) and using the unified stress–energy definition and diagnostic realized reference configuration expectation (Definition 9.358 and eqs. (9.229) and (9.238)), the semi–classical law on $\mathfrak{P}$ reads verbatim

$$\boxed{\bar{G}_{\mu\nu}[\bar{\Sigma}] + \Lambda\,\bar{\Sigma}_{\mu\nu} = \kappa\,\bar{T}^{\mathrm{total}}_{\mu\nu}, \qquad \bar{\nabla}^{(\mathrm{rel})}_{\mu}\,\bar{T}^{\mu\nu}_{\mathrm{total}} = 0,} \tag{11.190}$$

in agreement with Definition 11.106 and eqs. (9.232) and (11.191). Equivalently, one may record the cohomological equations of motion $\mathcal{E}^{\mathrm{coh}}_{\boldsymbol{\Phi}}(\bar{\Sigma}) = 0$, $\mathcal{E}^{\mathrm{coh}}_{\boldsymbol{\Theta}}(\bar{\Sigma}) = 0$ modulo ζ–exact terms, evaluated with Σ–causal retarded kernels on $\mathfrak{P}$ (Propositions 7.3 and 11.14 and eq. (11.25)).

Physical interpretation and alignment with the TT/triadic program. Because $\bar{R}[\bar{\Sigma}]$ depends on the full unified tensor $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (Section 5.2.6), the stationary condition derived from (11.188) balances metric curvature with coherence/temporal curvature (Remark 11.100 and eq. (11.177)). In the $\bar{t}_{\mu\nu} \to 0$ limit one recovers the standard Einstein–Hilbert dynamics $R[g]$; for non–negligible $\bar{t}_{\mu\nu}$, the difference $\bar{R}[\bar{\Sigma}] - R[\bar{g}]$ encodes genuine chronogeometric effects that feed into the unified source and back–reaction (Section 9.18.12 and Definition 9.358). Linearization about $(\mathcal{X}_q, \bar{\Sigma})$ produces the TT–typed spin–2 operator and its Σ–causal Green functions (Definition 11.60 and Lemma 11.101), which govern radiative gravitons; typed TLO diagnostics are attached *only* to this radiative TT sector (Definition 11.6 and Proposition 11.77), consistent with the Chapter 12 instrumentation preamble.

11.5.4 Unified Semi–Classical Einstein Equations

Instrumentation (diagnostic realized reference configuration, tags, covariance). All statements are evaluated on a certified plateau $\mathfrak{P}$ with BRST–closed band tags $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ =

$(P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}^{(\ell)})$ that commute with the triad and TT projectors (Definitions 6.280, 9.10, 10.3 and 11.24). Geometry and covariant differentials are those of the realized chronogeometry $(\mathcal{X}_q, \bar{\Sigma})$ with Relational Levi–Civita connection $\bar{\nabla} \equiv \nabla^{(\mathrm{rel})}$ and volume element $d\mu(x) = \sqrt{-\det \bar{\Sigma}}\, d^d x$ (Definitions 5.26, 6.7 and 6.33). Expectations descend to the physical space $\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}})$ and are Nielsen–flat on $\mathfrak{P}$ (Proposition 7.3).

Definition 11.106 (Chronogeometric Einstein equations (diagnostic realized reference configuration form))**.** The semi–classical field equations for the expectation–level chronogeometry $\bar{\Sigma}_{\mu\nu}$ read

$$\boxed{\bar{G}_{\mu\nu}[\bar{\Sigma}] \;+\; \Lambda\, \bar{\Sigma}_{\mu\nu} \;=\; \kappa\, \bar{T}^{\mathrm{total}}_{\mu\nu},} \tag{11.191}$$

where $\bar{G}_{\mu\nu} = \bar{R}_{\mu\nu} - \frac{1}{2}\bar{\Sigma}_{\mu\nu}\bar{R}$ (Section 11.5.2), Λ is the cosmological constant, κ the gravitational coupling, and $\bar{T}^{\mathrm{total}}_{\mu\nu}$ is the expectation–valued total stress–energy sourced by the Gaiamiti, Tantrakyma, and interaction sectors (Definition 9.358, eqs. (9.229) and (9.238), and Section 9.18.12). Equation (11.191) follows from varying the relational Einstein–Hilbert action with cosmological term on $(\mathcal{X}_q, \bar{\Sigma})$ (Definition 11.105 and eq. (11.188)).

$$\boxed{\bar{\nabla}^{\mu}\bar{G}_{\mu\nu} \equiv 0, \qquad \bar{\nabla}^{(\mathrm{rel})}_{\mu}\, \bar{T}^{\mu\nu}_{\mathrm{total}} = 0,} \tag{11.192}$$

the first by the contracted Bianchi identity (eq. (11.175)) and the second by the unified variational definition and on–shell consistency on $\mathfrak{P}$ (Definition 9.358 and eq. (9.232)).

Cohomological (BRST–clean) statement. Equivalently, the dynamical content may be recorded as cohomological equations of motion

$$\mathcal{E}^{\mathrm{coh}}_{\mathbf{\Phi}}(\bar{\Sigma}) = 0, \qquad \mathcal{E}^{\mathrm{coh}}_{\mathbf{\Theta}}(\bar{\Sigma}) = 0,$$

modulo ζ–exact terms and evaluated with Σ–causal retarded kernels on $\mathfrak{P}$ (Propositions 7.3 and 11.14 and eq. (11.25)).

Definition 11.107 (Global Einstein–law residual and visibility (diagnostic realized reference configuration–level))**.** Define the untyped law residual and visibility by

$$\boxed{\mathbf{R}^{[\mathrm{Ein}]}{}_{\mu\nu} := \bar{G}_{\mu\nu}[\bar{\Sigma}] + \Lambda\, \bar{\Sigma}_{\mu\nu} - \kappa\, \bar{T}^{\mathrm{total}}_{\mu\nu}, \qquad \Lambda^{[\mathrm{Ein}]}_{\mathrm{law}} := 1 - \frac{\|\mathbf{R}^{[\mathrm{Ein}]}\|_{\mathcal{I}}}{\|\cdot\|_{\mathrm{ref}}},} \tag{11.193}$$

with the instrumentation norm $\|\cdot\|_{\mathcal{I}}$ and reference scale chosen consistently with the plateau meters and synthesis logic of Chapter 9 (eqs. (9.126) to (9.128) and Definitions 9.297 and 9.299).

Interpretation.

- *Geometric side (LHS).* The Einstein tensor is built from the unified curvature of $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$, hence it encodes both metric focusing and coherence/temporal curvature (Section 11.5.2 and Remarks 11.100 and 11.104).

- *Source side (RHS).* The total source $\bar{T}^{\mathrm{total}}_{\mu\nu}$ aggregates geometry, matter, and interaction contributions at diagnostic realized reference configuration level, with conservation in the sum and sectoral conservation if exchange vanishes (Definition 9.358, eq. (9.238), and Section 9.18.12).

- *Typed radiative sector.* Linearization about $(\mathcal{X}_q, \bar{\Sigma})$ and TT projection produce the spin–2 typed operator and Σ–causal Green functions (Lemma 11.101 and Definition 11.60), to which the typed TLO panel is attached *only* for radiative gravity, consistent with the instrumentation preamble (Definition 11.6 and Proposition 11.77).

In this diagnostic realized reference configuration–regular, BRST–clean form, (11.191) is the dynamical core of the chronogeometric sector, relating unified curvature to the expectation–level total stress–energy on the same causal cones and with the same conservation content used throughout Chapters 6–11.

Remark 11.108 (Relational Einstein vector, Bianchi identity, and linearization on $(\mathcal{X}_q, \bar{\Sigma})$)**. Relational connection and curvatures.** On the realized chronogeometry $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (Definition 5.26), the Levi–Civita connection and curvatures are (Definitions 6.7 and 11.97 and Section 11.5.2)

$$\Gamma^{\lambda}_{\mu\nu}[\bar{\Sigma}] = \tfrac{1}{2}\,\bar{\Sigma}^{\lambda\rho}\Big(\partial_\mu \bar{\Sigma}_{\nu\rho} + \partial_\nu \bar{\Sigma}_{\mu\rho} - \partial_\rho \bar{\Sigma}_{\mu\nu}\Big), \qquad \bar{R}^{\lambda}{}_{\rho\mu\nu} = \partial_\mu \Gamma^{\lambda}_{\nu\rho} - \partial_\nu \Gamma^{\lambda}_{\mu\rho} + \Gamma^{\lambda}_{\mu\alpha}\Gamma^{\alpha}_{\nu\rho} - \Gamma^{\lambda}_{\nu\alpha}\Gamma^{\alpha}_{\mu\rho},$$

with $\bar{R}_{\mu\nu} = \bar{R}^{\lambda}{}_{\mu\lambda\nu}$, $\bar{R} = \bar{\Sigma}^{\mu\nu}\bar{R}_{\mu\nu}$, and $\bar{G}_{\mu\nu} = \bar{R}_{\mu\nu} - \tfrac{1}{2}\,\bar{\Sigma}_{\mu\nu}\bar{R}$ (eqs. (11.175), (11.185) and (11.186)).

Relational Bianchi identity and conservation. Metric compatibility and torsionlessness on $(\bar{\Sigma}, \bar{\nabla})$ imply the contracted Bianchi identity

$$\boxed{\bar{\nabla}^{\mu}\bar{G}_{\mu\nu}[\bar{\Sigma}] \equiv 0.} \tag{11.194}$$

Hence the Chronogeometric Einstein equations $\bar{G}_{\mu\nu}[\bar{\Sigma}] + \Lambda\,\bar{\Sigma}_{\mu\nu} = \kappa\,\bar{T}^{\mathrm{total}}_{\mu\nu}$ (Definition 11.106 and eq. (11.191)) entail the unified conservation law on the same background, $\bar{\nabla}^{(\mathrm{rel})}_{\mu}\,\bar{T}^{\mu\nu}_{\mathrm{total}} = 0$ (Section 11.5.5 and eqs. (9.232) and (11.192)).

Linearization around $\bar{\Sigma}$. Let $\bar{\Sigma}_{\mu\nu} \mapsto \bar{\Sigma}_{\mu\nu} + \delta\Sigma_{\mu\nu}$ and raise indices with $\bar{\Sigma}^{\mu\nu}$. To first order,

$$\delta\Gamma^{\lambda}_{\mu\nu} = \tfrac{1}{2}\,\bar{\Sigma}^{\lambda\rho}\Big(\bar{\nabla}_\mu \delta\Sigma_{\nu\rho} + \bar{\nabla}_\nu \delta\Sigma_{\mu\rho} - \bar{\nabla}_\rho \delta\Sigma_{\mu\nu}\Big), \tag{11.195}$$

$$\delta\bar{R}_{\mu\nu} = \tfrac{1}{2}\Big(-\bar{\nabla}^2\delta\Sigma_{\mu\nu} - \bar{\nabla}_\mu\bar{\nabla}_\nu\delta\Sigma + \bar{\nabla}_\mu\bar{\nabla}^\rho\delta\Sigma_{\nu\rho} + \bar{\nabla}_\nu\bar{\nabla}^\rho\delta\Sigma_{\mu\rho}\Big) + \bar{R}_{\rho(\mu}\,\delta\Sigma_{\nu)}{}^{\rho} - \bar{R}_{\mu\rho\nu\sigma}\,\delta\Sigma^{\rho\sigma}, \tag{11.196}$$

$$\delta\bar{G}_{\mu\nu} = \delta\bar{R}_{\mu\nu} - \tfrac{1}{2}\,\bar{\Sigma}_{\mu\nu}\,\delta\bar{R} - \tfrac{1}{2}\,\delta\Sigma_{\mu\nu}\,\bar{R}, \qquad \delta\bar{R} = \bar{\Sigma}^{\mu\nu}\delta\bar{R}_{\mu\nu} - \bar{R}_{\mu\nu}\delta\Sigma^{\mu\nu}. \tag{11.197}$$

In the de Donder gauge $\bar{\nabla}^{\mu}(\delta\Sigma_{\mu\nu} - \frac{1}{2}\bar{\Sigma}_{\mu\nu}\delta\Sigma) = 0$, the spin–2 operator on $(\bar{\Sigma}, \bar{\nabla})$ reduces to the Lichnerowicz form (Lemma 11.101)

$$\boxed{\mathcal{E}_{\mu\nu}{}^{\rho\sigma}\,\delta\Sigma_{\rho\sigma} := -\bar{\nabla}^2\delta\Sigma_{\mu\nu} - 2\,\bar{R}_{\mu}{}^{\rho}{}_{\nu}{}^{\sigma}\delta\Sigma_{\rho\sigma} + 2\,\bar{R}_{(\mu}{}^{\rho}\,\delta\Sigma_{\nu)\rho} - \tfrac{1}{2}\,\bar{\Sigma}_{\mu\nu}\,\bar{R}_{\rho\sigma}\delta\Sigma^{\rho\sigma},}\tag{11.198}$$

and the linearized field equation reads

$$\boxed{\mathcal{E}_{\mu\nu}{}^{\rho\sigma}\,\delta\Sigma_{\rho\sigma} = \kappa\,\Pi_{\mu\nu}{}^{\rho\sigma}\,\delta T^{\text{total}}_{\rho\sigma} \;+\; (\text{terms from } \Lambda),} \tag{11.199}$$

with Π the appropriate algebraic projector (TT in the spin–2 sector), consistent with the typed operator in Definition 11.60.

Causal Green kernels and typed sector. The retarded/advanced inverses of $\mathcal{E}$ on $(\bar{\Sigma}, \bar{\nabla})$ define graviton Green kernels whose Pauli–Jordan difference has support on the same relational cones $J^{\pm}_{\bar{\Sigma}}$ (Lemma 11.101 and Proposition 11.14). Via the Peierls construction (eq. (11.25)) this causal kernel generates the covariant (commutator) algebra; TT projection then yields the typed propagator and ETCR in the radiative sector (Definition 11.60 and Proposition 11.4). These kernels inherit the microcausal cone also used by the scalar Stieltjes representations, thereby bridging kinematics (Chs. 6–7) to dynamics (Ch. 8) and the Einstein sector (Ch. 10) within the Chapter 12 chronogeometric framework (Sections 5.4.7 and 9.18.12).

11.5.5 Derivation of the Unified Semi–Classical Einstein Equations

Instrumentation (diagnostic realized reference configuration, tags, covariance). All derivations are performed on a certified plateau $\mathfrak{P}$ with BRST–closed band tags $\mathfrak{B}_{\lambda_{\text{UV}},\ell} = (P_x(\lambda_{\text{UV}}), K^{(\ell)}_{\text{IR}})$ that commute with the triad and TT projectors (Definitions 6.280, 9.10, 10.3 and 11.24). Expectations descend to $\mathcal{H}_{\text{phys}} := H^0(\hat{Q}_{\text{BRST}})$ (Proposition 7.3). Geometry and covariant derivatives are those of the realized chronogeometry $(\mathcal{X}_q, \bar{\Sigma})$ with Relational Levi–Civita connection $\bar{\nabla}$ and relational measure $d\mu(x) = \sqrt{-\det \bar{\Sigma}}\, d^d x$ (Definitions 5.26, 6.7 and 6.33).

Step 1: Emergent chronogeometry from the Holo–State

As established in Proposition 1.28, the expectation values of the operator–valued metric and temporal/coherence tensors in the Quantum Holo–State $|\mathring{\mathbf{\Psi}}\rangle$ define the unified chronogeometric tensor:

$$\begin{aligned} &\bar{g}_{\mu\nu}(x) := \langle \mathring{\mathbf{\Psi}} | \hat{g}_{\mu\nu}(x) | \mathring{\mathbf{\Psi}} \rangle, \qquad \bar{t}_{\mu\nu}(x) := \langle \mathring{\mathbf{\Psi}} | \hat{t}_{\mu\nu}(x) | \mathring{\mathbf{\Psi}} \rangle, \\ &\bar{\Sigma}_{\mu\nu}(x) := \bar{g}_{\mu\nu}(x) + \bar{t}_{\mu\nu}(x) \qquad (\text{unified tensor; } \textit{Definition } 5.26). \end{aligned}$$

Step 2: Curvatures from the Relational Levi–Civita connection

Treating $\bar{\Sigma}_{\mu\nu}$ as fundamental, the Relational Levi–Civita connection $\Gamma^{\lambda}_{\mu\nu}[\bar{\Sigma}]$ and curvatures are constructed as in Definitions 6.7 and 11.97. Contractions yield the Ricci tensor and scalar $\bar{R}_{\mu\nu}[\bar{\Sigma}]$, $\bar{R}[\bar{\Sigma}]$ (Section 11.5.2), and the Einstein tensor

$$\bar{G}_{\mu\nu}[\bar{\Sigma}] := \bar{R}_{\mu\nu}[\bar{\Sigma}] - \tfrac{1}{2}\,\bar{\Sigma}_{\mu\nu}\,\bar{R}[\bar{\Sigma}] \quad (\textit{eq. } (11.175)).$$

Step 3: Expectation–valued total stress–energy

The operator–valued stress–energy tensors of the Gaiamiti and Tantrakyma sectors, $\hat{T}^{\mu\nu}_{(\Phi)}$ and $\hat{T}^{\mu\nu}_{(\Theta)}$ (Section 9.17.9 and Definition 9.356), together with the interaction contribution $\hat{T}^{\mu\nu}_{(\mathrm{int})}$ (variationally defined from S_{int}; eq. (9.237)), give

$$\hat{T}^{\mu\nu}_{\mathrm{total}} := \hat{T}^{\mu\nu}_{(\Phi)} + \hat{T}^{\mu\nu}_{(\Theta)} + \hat{T}^{\mu\nu}_{(\mathrm{int})}, \quad T^{\mu\nu}_{\mathrm{total}} := \mathrm{Tr}\{\hat{T}^{\mu\nu}_{\mathrm{total}}\} \quad (\textit{Definition 9.357 and eq. (9.229)}).$$

The expectation–level source on $\mathfrak{P}$ is

$$\boxed{\bar{T}^{\mu\nu}_{\mathrm{total}}(x) := \left\langle \mathring{\boldsymbol{\Psi}} \,\middle|\, \mathsf{P}_{\mathrm{phys}}\, T^{\mu\nu}_{\mathrm{total}}(x) \,\middle|\, \mathring{\boldsymbol{\Psi}} \right\rangle \qquad (\textit{Definition 9.358 and eq. (9.238)}).} \tag{11.200}$$

Step 4: Assembling the field equation (Hilbert variation and diagnostic realized reference configuration conservation)

The Relational Einstein–Hilbert functional with cosmological term (Definition 11.105 and eq. (11.188)) satisfies the Hilbert variational identity

$$\delta S^{(\mathrm{rel})}_{\mathrm{EH}} = \frac{1}{2\kappa}\int d\mu \left(\bar{G}_{\mu\nu}[\bar{\Sigma}] + \Lambda\, \bar{\Sigma}_{\mu\nu}\right) \delta\bar{\Sigma}^{\mu\nu} \quad (\textit{eq. (11.189)}),$$

while variation of S_{total} defines $T^{\mu\nu}_{\mathrm{total}}$ and its diagnostic realized reference configuration expectation (Definitions 9.357 and 9.358). Equating the geometric and matter variations and using eq. (11.200) gives the semi–classical law

$$\boxed{\bar{G}_{\mu\nu}[\bar{\Sigma}] + \Lambda\, \bar{\Sigma}_{\mu\nu} = \kappa\, \bar{T}^{\mathrm{total}}_{\mu\nu},} \tag{11.201}$$

which coincides with eq. (11.191). By the contracted Bianchi identity on $(\bar{\Sigma}, \bar{\nabla})$ and the unified variational definition, the diagnostic realized reference configuration conservation law holds on the same background,

$$\boxed{\bar{\nabla}^{\mu}\bar{G}_{\mu\nu} \equiv 0, \qquad \bar{\nabla}^{(\mathrm{rel})}_{\mu}\, \bar{T}^{\mu\nu}_{\mathrm{total}} = 0 \quad (\textit{eqs. (9.232) and (11.192)}).} \tag{11.202}$$

Cohomological form (BRST–clean). Equivalently, the equations of motion may be written cohomologically as $\mathcal{E}^{\mathrm{coh}}_{\boldsymbol{\Phi}}(\bar{\Sigma}) = 0$ and $\mathcal{E}^{\mathrm{coh}}_{\boldsymbol{\Theta}}(\bar{\Sigma}) = 0$ modulo ζ–exact terms, evaluated with Σ–causal retarded kernels on $\mathfrak{P}$ (Propositions 7.3 and 11.14 and eq. (11.25)).

Alternative derivation from the Relational Einstein–Hilbert functional

Setup and conventions (diagnostic realized reference configuration, tags, covariance). All variations are performed on a certified plateau $\mathfrak{P}$ with BRST–closed band tags $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell}$ that commute with the triad and TT projectors (Definitions 6.280, 9.10, 10.3 and 11.24). Expectations descend to $\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}})$ (Proposition 7.3). Geometry, measure, and covariant derivatives are those of $(\mathcal{X}_q, \bar{\Sigma})$ with Relational Levi–Civita connection and relational volume element (Definitions 6.7 and 6.33).

Relational gravitational functional. Start from the Relational Einstein–Hilbert action including a cosmological term and the standard normalization,

$$\boxed{S_{\mathrm{Relational}}[\bar{\Sigma}] := \frac{1}{2\kappa}\int_{\mathcal{X}_q} d\mu(x)\,\Big(\,\bar{R}[\bar{\Sigma}](x) - 2\Lambda\,\Big), \qquad d\mu(x) = \sqrt{-\det\bar{\Sigma}(x)}\,d^dx,} \tag{11.203}$$

with $\bar{R}[\bar{\Sigma}]$ defined from the Relational Levi–Civita connection (Definitions 11.97 and 11.105 and Section 11.5.2).

Step 1: Hilbert variation (geometric side). Varying with respect to $\bar{\Sigma}^{\mu\nu}$ and using $\delta(\sqrt{-\det\bar{\Sigma}}) = -\frac{1}{2}\sqrt{-\det\bar{\Sigma}}\,\bar{\Sigma}_{\mu\nu}\,\delta\bar{\Sigma}^{\mu\nu}$ together with the Palatini identity (discarding the boundary term under the usual conditions), one obtains the standard Hilbert identity

$$\delta S_{\mathrm{Relational}} = \frac{1}{2\kappa}\int_{\mathcal{X}_q} d\mu\,\Big(\bar{R}_{\mu\nu} - \tfrac{1}{2}\,\bar{\Sigma}_{\mu\nu}\,\bar{R} + \Lambda\,\bar{\Sigma}_{\mu\nu}\Big)\,\delta\bar{\Sigma}^{\mu\nu} = \frac{1}{2\kappa}\int_{\mathcal{X}_q} d\mu\,\Big(\bar{G}_{\mu\nu} + \Lambda\,\bar{\Sigma}_{\mu\nu}\Big)\,\delta\bar{\Sigma}^{\mu\nu},$$

where $\bar{G}_{\mu\nu} = \bar{R}_{\mu\nu} - \frac{1}{2}\,\bar{\Sigma}_{\mu\nu}\bar{R}$ (eq. (11.175)). The contracted Bianchi identity $\bar{\nabla}^{\mu}\bar{G}_{\mu\nu}\equiv 0$ holds on $(\bar{\Sigma},\bar{\nabla})$ (Proposition 11.98 and eq. (11.176e)).

Step 2: Coupling to fields and variational definition of the source. Augment the geometric functional with the field actions as in the total relational action (Definition 9.357),

$$S_{\mathrm{total}}[\bar{\Sigma},\hat{\mathbf{\Phi}},\hat{\mathbf{\Theta}}] = S_{\mathrm{Relational}}[\bar{\Sigma}] + S_{\Phi}[\hat{\mathbf{\Phi}}] + S_{\Theta}[\hat{\mathbf{\Theta}}] + S_{\mathrm{int}}[\hat{\mathbf{\Phi}},\hat{\mathbf{\Theta}}\,],$$

with S_{Φ}, S_{Θ} built from the operator–valued kernels and the same measure (Section 6.5 and Definitions 9.301 and 9.310) and S_{int} encoding geometry–matter exchange. By definition of the (scalar, fiber–traced) stress–energy, the variation of the field part with respect to $\bar{\Sigma}$ reads

$$\boxed{\delta\Big(S_{\Phi} + S_{\Theta} + S_{\mathrm{int}}\Big) = \frac{1}{2}\int_{\mathcal{X}_q} d\mu\; T^{\mu\nu}_{\mathrm{total}}(x)\,\delta\bar{\Sigma}_{\mu\nu}(x), \qquad T^{\mu\nu}_{\mathrm{total}} := \mathrm{Tr}\{\hat{T}^{\mu\nu}_{(\Phi)} + \hat{T}^{\mu\nu}_{(\Theta)} + \hat{T}^{\mu\nu}_{(\mathrm{int})}\},}$$

(11.204)

consistent with eqs. (9.224) and (9.237). On the diagnostic realized reference configuration, the expectation source is $\bar{T}^{\mu\nu}_{\mathrm{total}} = \langle \mathsf{P}_{\mathrm{phys}}\,T^{\mu\nu}_{\mathrm{total}}\rangle_{\mathfrak{P}}$ (Definition 9.358 and eq. (9.238)).

Step 3: Field equation and conservation. Stationarity of S_{total} under arbitrary $\delta\bar{\Sigma}^{\mu\nu}$ gives, after taking the diagnostic realized reference configuration expectation on $\mathfrak{P}$,

$$\boxed{\bar{G}_{\mu\nu}[\bar{\Sigma}] + \Lambda\,\bar{\Sigma}_{\mu\nu} = \kappa\,\bar{T}^{\mathrm{total}}_{\mu\nu},} \tag{11.205}$$

which is eq. (11.191). By $\bar{\nabla}^{\mu}\bar{G}_{\mu\nu}\equiv 0$ and the unified variational definition,

$$\bar{\nabla}^{(\mathrm{rel})}_{\mu}\,\bar{T}^{\mu\nu}_{\mathrm{total}} = 0,$$

in agreement with Section 11.5.5 and eq. (9.232).

Cohomological (BRST–clean) form and diagnostic realized reference configuration visibility. On $\mathfrak{P}$, the equations admit a cohomological expression $\mathcal{E}^{\mathrm{coh}}_{\mathbf{\Phi}}(\bar{\Sigma}) = 0$, $\mathcal{E}^{\mathrm{coh}}_{\mathbf{\Theta}}(\bar{\Sigma}) = 0$ modulo ζ–exact terms (Proposition 7.3). A global (diagnostic realized reference configuration–level) Einstein–law residual and visibility may be attached,

$$\boxed{\mathbf{R}^{[\mathrm{Ein}]}_{\mu\nu} := \bar{G}_{\mu\nu}[\bar{\Sigma}] + \Lambda\,\bar{\Sigma}_{\mu\nu} - \kappa\,\bar{T}^{\mathrm{total}}_{\mu\nu}, \qquad \Lambda^{[\mathrm{Ein}]}_{\mathrm{law}} := 1 - \|\mathbf{R}^{[\mathrm{Ein}]}\|_{\mathcal{I}}/\|\cdot\|_{\mathrm{ref}},}\tag{11.206}$$

following the Parasymmetric T–L–O diagnostics of Chapter 9 (Definitions 9.299 and 9.306).

Notes on linearization and typed projection. Linearizing (11.205) around $\bar{\Sigma}$ produces the Lichnerowicz operator $\mathcal{E}_{\mu\nu}{}^{\rho\sigma}$ on $(\bar{\Sigma}, \bar{\nabla})$ (Remark 11.108 and Definition 11.60). Its retarded/advanced inverses are supported on $J^{\pm}_{\bar{\Sigma}}$ and generate, via the Peierls bracket, the covariant algebra whose equal–time limit reproduces the typed ETCR in the TT sector (eq. (11.25) and Propositions 11.4 and 11.14).

1. Variation of the geometric action (Hilbert identity, diagnostic realized reference configuration form) With the normalization and cosmological term fixed in Definition 11.105 (see also eq. (11.203)),

$$S_{\mathrm{Relational}}[\bar{\Sigma}] = \frac{1}{2\kappa}\int_{\mathcal{X}_q} d\mu(x)\,\Big(\bar{R}[\bar{\Sigma}](x) - 2\Lambda\Big), \qquad d\mu(x) = \sqrt{-\det\bar{\Sigma}(x)}\,d^d x,$$

the Palatini–Hilbert variation with respect to the *inverse* unified tensor $\bar{\Sigma}^{\mu\nu}$ (discarding the total derivative under standard boundary conditions) yields the standard result

$$\boxed{\delta S_{\mathrm{Relational}} = \frac{1}{2\kappa}\int_{\mathcal{X}_q} d\mu\,\Big(\bar{R}_{\mu\nu} - \tfrac{1}{2}\,\bar{\Sigma}_{\mu\nu}\,\bar{R} + \Lambda\,\bar{\Sigma}_{\mu\nu}\Big)\,\delta\bar{\Sigma}^{\mu\nu} = \frac{1}{2\kappa}\int_{\mathcal{X}_q} d\mu\,\Big(\bar{G}_{\mu\nu} + \Lambda\,\bar{\Sigma}_{\mu\nu}\Big)\,\delta\bar{\Sigma}^{\mu\nu},}\tag{11.207}$$

with $\bar{G}_{\mu\nu} = \bar{R}_{\mu\nu} - \frac{1}{2}\,\bar{\Sigma}_{\mu\nu}\bar{R}$ (Definition 11.97 and Section 11.5.2). The relational volume element is that of Definition 6.33 and all curvatures are those of the Relational Levi–Civita connection (Definition 6.7).

2. Variation of the field action and the source definition For the field part $S_{\mathrm{fields}} := S_{\Phi} + S_{\Theta} + S_{\mathrm{int}}$ (Definitions 9.310 and 9.357, Section 6.5, and eq. (9.237)), the *scalar* (fiber–traced) stress–energy is defined variationally by

$$\boxed{T^{\mathrm{total}}_{\mu\nu}(x) := -\frac{2}{\sqrt{-\det\bar{\Sigma}(x)}}\,\frac{\delta S_{\mathrm{fields}}}{\delta\bar{\Sigma}^{\mu\nu}(x)}\,, \qquad \Rightarrow \quad \delta S_{\mathrm{fields}} = -\frac{1}{2}\int_{\mathcal{X}_q} d\mu\,T^{\mathrm{total}}_{\mu\nu}\,\delta\bar{\Sigma}^{\mu\nu},}\tag{11.208}$$

consistent with eq. (9.224). On a certified plateau $\mathfrak{P}$ the expectation (diagnostic realized reference configuration) source is

$$\boxed{\bar{T}^{\mathrm{total}}_{\mu\nu}(x) := \Big\langle \mathring{\mathbf{\Psi}}\ \Big|\ \mathsf{P}_{\mathrm{phys}}\,\mathrm{Tr}\Big\{\hat{T}_{(\Phi)\mu\nu} + \hat{T}_{(\Theta)\mu\nu} + \hat{T}_{(\mathrm{int})\mu\nu}\Big\}(x)\ \Big|\ \mathring{\mathbf{\Psi}}\Big\rangle,}\tag{11.209}$$

as in Definition 9.358 and eq. (9.238), with BRST/Nielsen cleanliness supplied by Proposition 7.3.

Note. Here "total" is *field-only*, i.e. it refers to $S_\Phi + S_\Theta + S_{\rm int}$; the coherence/chronokymon sector contributes a separate $T^{(\tau)}_{\mu\nu}[K^{(\tau)}, \mathcal{F}^{(\tau)}]$ that appears explicitly in (11.257)–(11.258) and is TT–null ($\Pi^{[\rm TT]}_{\bar\Sigma}\hat\tau_{\mu\nu} = 0$; cf. (10.139)), so TT equations and diagnostics remain unchanged.

3. Assembling the total variation and the field equations Adding Section 11.5.5 gives

$$\boxed{\delta S_{\rm total} = \frac{1}{2\kappa}\int_{\mathcal{X}_q} d\mu\,\left(\bar G_{\mu\nu} + \Lambda\,\bar\Sigma_{\mu\nu}\right)\delta\bar\Sigma^{\mu\nu} - \frac{1}{2}\int_{\mathcal{X}_q} d\mu\,T^{\rm total}_{\mu\nu}\,\delta\bar\Sigma^{\mu\nu}.} \tag{11.210}$$

Stationarity for arbitrary $\delta\bar\Sigma^{\mu\nu}$ implies

$$\bar G_{\mu\nu} + \Lambda\,\bar\Sigma_{\mu\nu} = \kappa\,T^{\rm total}_{\mu\nu},$$

and, upon taking the diagnostic realized reference configuration expectation on $\mathfrak{P}$,

$$\boxed{\bar G_{\mu\nu}[\bar\Sigma] + \Lambda\,\bar\Sigma_{\mu\nu} = \kappa\,\bar T^{\rm total}_{\mu\nu},} \tag{11.211}$$

which reproduces Definition 11.106 and eq. (11.191). By the relational Bianchi identity $\bar\nabla^\mu\bar G_{\mu\nu} \equiv 0$ and the unified variational definition, one has $\bar\nabla^{(\rm rel)}_\mu\,\bar T^{\mu\nu}_{\rm total} = 0$ (Section 11.5.5 and Remark 11.108).

Step 4: Field equations, normalization, and the emergent coupling κ With the normalization chosen in Definition 11.105, the gravitational coupling κ enters with the standard dimensional weight (curvature $\sim$ energy density), so no *post hoc* insertion is required. In the IQR viewpoint, κ may be treated as an *emergent* diagnostic realized reference configuration parameter determined by the universal state and tags on the certified plateau $\mathfrak{P}$ (cohomological/Nielsen–flat semantics; Proposition 7.3), while the operational content of the law remains the Chronogeometric Einstein equation of Definition 11.106.

Diagnostic realized reference configuration form and stationarity. Starting from the normalized relational action with cosmological term (Definition 11.105),

$$S_{\rm Relational}[\bar\Sigma] = \frac{1}{2\kappa}\int_{\mathcal{X}_q} d\mu\,\left(\bar R[\bar\Sigma] - 2\Lambda\right), \qquad d\mu := \sqrt{-\det\bar\Sigma}\,d^dx,$$

and adding the field part $S_{\rm fields}$ (Definition 9.357), the total variation is

$$\delta S_{\rm total} = \frac{1}{2\kappa}\int d\mu\,(\bar G_{\mu\nu} + \Lambda\bar\Sigma_{\mu\nu})\,\delta\bar\Sigma^{\mu\nu} - \frac{1}{2}\int d\mu\,T^{\rm total}_{\mu\nu}\,\delta\bar\Sigma^{\mu\nu}.$$

Stationarity for arbitrary $\delta\bar\Sigma^{\mu\nu}$ gives the unified Einstein law

$$\boxed{\bar G_{\mu\nu}[\bar\Sigma] + \Lambda\,\bar\Sigma_{\mu\nu} = \kappa\,\bar T^{\rm total}_{\mu\nu},} \tag{11.212}$$

with the diagnostic realized reference configuration/expectation source $\bar{T}^{\text{total}}_{\mu\nu}$ defined in Definition 9.358. By the relational Bianchi identity $\bar{\nabla}^{\mu}\bar{G}_{\mu\nu} \equiv 0$, one has $\bar{\nabla}^{\mu}\bar{T}^{\text{total}}_{\mu\nu} = 0$ on the same background $(\bar{\Sigma}, \bar{\nabla})$ (Remark 11.108).
Normalization variant. If one begins from a *simplified* geometric action without the prefactor (effectively setting $\kappa = \frac{1}{2}$), the stationarity condition yields $\bar{G}_{\mu\nu} = \frac{1}{2}\bar{T}^{\text{total}}_{\mu\nu}$. Restoring physical units corresponds to reinstating the coupling κ, recovering (11.212) and the canonical statement of Definition 11.106.

11.5.6 The Emergent Gravitational Coupling Tensor

In the IQR framework, structural consistency requires that the gravitational coupling arise from the same quantum–relational substrate that generates both geometry and matter. Concretely, the geometry–matter interaction is encoded by the off–diagonal entries of the Unified Inter–Modal Correlator, i.e. the matter–geometry kernels $\hat{\mathbf{G}}_{\mathbf{\Phi\Theta}}$ and $\hat{\mathbf{G}}_{\mathbf{\Theta\Phi}}$, which enter the interaction action (Definition 10.223):

$$S_{\text{int}}\Big[\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}\Big] = \int d\mu_{\bar{\Sigma}}(x) \int d\mu_{\bar{\Sigma}}(x')\ \mathrm{Tr}\Big\{\hat{\mathbf{\Phi}}^{\dagger}(x)\, \mathcal{U}_{\mathbf{\Phi\Theta}}(x,x')\, \hat{\mathbf{\Theta}}(x')\Big\},$$
$$\mathcal{U}_{\mathbf{\Phi\Theta}}(x,x') := \hat{\mathbf{G}}_{\mathbf{\Phi\Theta}}(x,x') \ + \ \hat{\mathbf{G}}^{\dagger}_{\mathbf{\Theta\Phi}}(x',x), \qquad d\mu_{\bar{\Sigma}}(x) := \sqrt{-\det \bar{\Sigma}(x)}\, d^d x.$$

The semi–classical Chronogeometric Einstein law (Definition 11.106) is then promoted to a causal, tensorial response in which curvature at x is sourced by stress–energy at x' via a retarded bi–tensor kernel built on $(\mathcal{X}_q, \bar{\Sigma})$ (cf. Remark 11.108).

Definition 11.109 (Operator–valued gravitational coupling (retarded, causal))**.** Let $\mathcal{E}_{\alpha\beta}{}^{\gamma\delta}$ denote the linearized Einstein/Lichnerowicz operator on $(\bar{\Sigma}, \bar{\nabla})$ (Remark 11.108 and Definition 11.60), and let $G^{\text{ret}\,\gamma\delta}_{\alpha\beta}(x,x')$ be its retarded Green bi–solution supported on the $\bar{\Sigma}$–causal cones (Proposition 11.14). Define the operator–valued type–$(2,2)$ superoperator kernel

$$\hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x') : \ \hat{T}_{\rho\sigma}(x') \ \mapsto \ \hat{G}_{\mu\nu}(x),$$

by the composition

$$\boxed{\hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x') := \ \mathbb{P}^{(\text{Ein})\,\alpha\beta}_{\mu\nu}\ G^{\text{ret}\,\gamma\delta}_{\alpha\beta}(x,x')\ \mathbb{J}_{\gamma\delta}{}^{\rho\sigma},} \tag{11.213}$$

where: (i) $\mathbb{P}^{(\text{Ein})}$ is the algebraic map from metric fluctuations to the Einstein operator (linearized at $\bar{\Sigma}$), (ii) $\mathbb{J}$ is the index–symmetrizer onto the $(\rho\sigma)$ channel, and (iii) G^{ret} is computed with respect to the Relational Levi–Civita connection of $\bar{\Sigma}$ (Definition 6.7). The corresponding covariant response is

$$\boxed{\begin{aligned} &\hat{G}_{\mu\nu}(x) = \int_{\mathcal{X}_q} d\mu_{\bar{\Sigma}}(x')\ \hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x') \ \circ \ \Big[\mathcal{U}_{\bar{\Sigma}}(x \leftarrow x')_{\rho}{}^{\rho'}\, \mathcal{U}_{\bar{\Sigma}}(x \leftarrow x')_{\sigma}{}^{\sigma'}\, \hat{T}_{\rho'\sigma'}(x')\Big], \\ &d\mu_{\bar{\Sigma}}(x') := \sqrt{-\det \bar{\Sigma}(x')}\, d^d x', \qquad A \circ B := \tfrac{1}{2}(AB + BA), \end{aligned}} \tag{11.214}$$

with $\mathcal{U}_{\bar{\Sigma}}(x \leftarrow x')$ the Levi–Civita parallel transporter of $\bar{\Sigma}$ for covariant index carriage.

Remark 11.110 (Covariance, causality, and consistency of (11.214))**.**

- **Parallel transport.** Since $\hat{T}_{\rho\sigma}(x') \in T^*_{x'}\mathcal{X}_q \otimes T^*_{x'}\mathcal{X}_q$, parallel transport to x is required before contraction with a kernel at x. The factors $\mathcal{U}_{\bar{\Sigma}}(x \leftarrow x')_\rho{}^{\rho'}$, $\mathcal{U}_{\bar{\Sigma}}(x \leftarrow x')_\sigma{}^{\sigma'}$ implement this transport with the Relational Levi–Civita connection (Definition 6.7).
- **Volume element.** The use of the relational measure $d\mu_{\bar{\Sigma}} = \sqrt{-\det \bar{\Sigma}}\, d^d x$ (Definition 6.33) ensures diffeomorphism covariance of the response.
- **Operator ordering / self–adjointness.** The Jordan product $A \circ B := \frac{1}{2}(AB + BA)$ preserves self–adjointness of $\hat{G}_{\mu\nu}(x)$ when the kernel and the source act on the same fiber Hilbert space (cf. operator hermiticity conventions in Section 9.17.9 and Definition 9.356).
- **Index flow.** In (11.213), the symmetrizer $\mathbb{J}_{\gamma\delta}{}^{\rho\sigma}$ prepares the source indices, $G^{\mathrm{ret}}{}_{\alpha\beta}^{\gamma\delta}$ gives causal propagation from x' to x (with TT restriction when working in the radiative sector, cf. Definition 11.60 and Proposition 11.14), and $\mathbb{P}^{(\mathrm{Ein})\alpha\beta}_{\mu\nu}$ maps the propagated metric fluctuation to the Einstein operator at x. The result is a type–$(2,2)$ bitensor with input $(\rho\sigma)$ and output $(\mu\nu)$.
- **Bianchi identity and conservation.** Because G^{ret} is built from the linearized operator on $(\bar{\Sigma}, \bar{\nabla})$ and $\mathbb{P}^{(\mathrm{Ein})}$ is algebraically $\bar{\nabla}$–compatible, one has $\bar{\nabla}^\mu_x \hat{G}_{\mu\nu}(x) = 0$ provided the source obeys $\bar{\nabla}'_\rho \hat{T}^\rho{}_\sigma(x') = 0$ and the transporter is Levi–Civita (cf. Remark 11.108).
- **Short convention (bitensor–density).** Equivalently, one may absorb parallel transport and the density factor into the definition of $\hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x')$ (treating it as a bitensor–density in the primed slots), yielding the compact form

$$\hat{G}_{\mu\nu}(x) = \int d^d x'\, \hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x') \;\circ\; \hat{T}_{\rho\sigma}(x'),$$

  which is exactly equivalent to (11.214) if used consistently throughout the chapter.

Typed/radiative specialization (optional). When focusing on the radiative spin–2 content, replace G^{ret} by its TT–projected version $G^{\mathrm{ret},[\mathrm{TT}]}$ built from the typed operator $K^{[\mathrm{TT}]}$ (Definition 11.60); microcausal cone support and BRST/Nielsen safety then follow from Propositions 7.3 and 11.14. This specialization aligns the response law with the TT exchange and dressing formulas used elsewhere (Definitions 11.67 and 11.71).

Definition 11.111 (Emergent Gravitational Coupling Tensor)**.** Adopt the causal operator kernel $\hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x')$ of Definition 11.109, which implements the retarded metric–response to a stress–energy insertion on $(\mathcal{X}_q, \bar{\Sigma})$ via eq. (11.214). On a certified plateau $\mathfrak{P}$ with physical projector $\mathsf{P}_{\mathrm{phys}} : \mathcal{H} \to H^0(\hat{Q}_{\mathrm{BRST}})$ (Proposition 7.3), the *Emergent Gravitational Coupling Tensor* is the diagnostic realized reference configuration (expectation–level) bitensor

$$\boxed{\bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x') := \Big\langle \overset{\circ}{\boldsymbol{\Psi}} \Big|\, \mathsf{P}_{\mathrm{phys}}\, \hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x') \,\Big|\, \overset{\circ}{\boldsymbol{\Psi}} \Big\rangle,} \tag{11.215}$$

with indices raised/lowered by $\bar{\Sigma}_{\alpha\beta}$ and its inverse (Definition 5.26). In the "short" convention of Definition 11.109 (bitensor–density in the primed slots), one may write a pointwise

version $\bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x)$, keeping in mind that parallel transport and the density factor are implicit (eq. (11.214)).

Properties.

(i) Operator origin and index structure.
$\hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x')$ acts on the fiber Hilbert space $\mathcal{H}_{\rm em}$ and is of type $(2,2)$ in spacetime indices, mapping an input (matter/source) pair $(\rho\sigma)$ at x' to an output (geometry/response) pair $(\mu\nu)$ at x (Definition 11.109). Its diagnostic realized reference configuration expectation (11.215) defines a c-number bitensor that enters the linear response of the Chronogeometric Einstein law (Definition 11.106 and Remark 11.108). Symmetry on each index pair is assumed, $\bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma} = \bar{\mathcal{C}}_{\nu\mu}{}^{\rho\sigma} = \bar{\mathcal{C}}_{\mu\nu}{}^{\sigma\rho}$, and, when appropriate, pair–exchange symmetry with respect to the $\bar{\Sigma}$ inner product, $\bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma} = \bar{\mathcal{C}}^{\rho\sigma}{}_{\mu\nu}$.

(ii) Functional dependence and causality.
$\hat{\mathcal{C}}$ is built from the linearized Lichnerowicz/Einstein operator on $(\bar{\Sigma},\bar{\nabla})$ and its retarded Green kernel (Remark 11.108), along with the index maps $\mathbb{J}$ and $\mathbb{P}^{(\rm Ein)}$ (Definition 11.109). Hence supp $\bar{\mathcal{C}}(\cdot;x') \subset J^+_{\bar{\Sigma}}(x')$ (Proposition 11.14), i.e. the coupling is microcausal on the $\bar{\Sigma}$–cone used throughout.

(iii) Triadic/typed compatibility.
$\hat{\mathcal{C}}$ commutes with the triad projectors $\Pi^{(a)}$ and, in the radiative spin–2 sector, with the TT projector $\Pi^{[\rm TT]}_{\bar{\Sigma}}$ (Definitions 11.24 and 11.60 and Proposition 11.77). Consequently, its contractions with typed/triadic sources remain block–diagonal by $a \in \{\mathrm{corr},\mathrm{vac},\mathrm{spec}\}$ and TT–pure where required.

(iv) State and band dependence (diagnostic realized reference configuration semantics).
The diagnostic realized reference configuration expectation in (11.215) is taken at fixed band tags that commute with the projectors (Definitions 6.280 and 9.10); by Nielsen–flatness the resulting $\bar{\mathcal{C}}$ is gauge–parameter independent on $\mathfrak{P}$ (Proposition 7.3 and Corollary 11.5).

(v) GR limit.
In the local, weak–field GR limit the bitensor reduces to a point–localized, pair–symmetrized coupling,

$$\bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x') \;\longrightarrow\; \kappa\,\tfrac{1}{2}\big(\delta_\mu{}^\rho\delta_\nu{}^\sigma + \delta_\mu{}^\sigma\delta_\nu{}^\rho\big)\,\delta^{(d)}(x-x'),$$

recovering Definition 11.106.

(vi) Local response form (short convention).
Suppressing explicit transport and density factors (Definition 11.109), the pointwise relation reads

$$\bar{G}_{\mu\nu}(x) = \int d^dx'\,\bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x')\,\bar{T}^{\rm total}_{\rho\sigma}(x'),$$

and in the ultra–local approximation, $\bar{G}_{\mu\nu}(x) = \bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x)\,\bar{T}^{\rm total}_{\rho\sigma}(x)$, with conservation guaranteed by the relational Bianchi identity (Remark 11.108).

Proposition 11.112 (Generalized Chronogeometric Einstein equation (retarded response form))**.** *Work on a certified plateau $\mathfrak{P}$ with BRST–closed band tags that commute with the triad and TT projectors (Definitions 6.280 and 9.10 and Proposition 7.3). Let $\bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x')$ be*

the Emergent Gravitational Coupling Tensor of Definition 11.111, obtained from the operator kernel of Definition 11.109 and evaluated as a diagnostic realized reference configuration expectation. Then the semi–classical gravitational response of the unified geometry satisfies the causal, bitensor equation

$$\boxed{\bar{G}_{\mu\nu}[\bar{\Sigma}](x) + \Lambda\,\bar{\Sigma}_{\mu\nu}(x) \;=\; \int_{\mathcal{X}_q} d\mu_{\bar{\Sigma}}(x')\;\bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x')\,\bar{T}^{\mathrm{total}}_{\rho\sigma}(x')\,,}\tag{11.216}$$

with $d\mu_{\bar{\Sigma}}(x') = \sqrt{-\det\bar{\Sigma}(x')}\,d^dx'$ *and* $\bar{G}_{\mu\nu}[\bar{\Sigma}]$ *built from the relational Levi–Civita geometry (Definition 11.97, Section 11.5.2, and Remark 11.108). In the ultra–local GR limit,*

$$\boxed{\bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x') \;\longrightarrow\; \kappa\,\tfrac{1}{2}\Big(\delta_\mu{}^\rho\delta_\nu{}^\sigma + \delta_\mu{}^\sigma\delta_\nu{}^\rho\Big)\,\delta^{(d)}(x-x')\,,}\tag{11.217}$$

(11.216) *reduces to the Chronogeometric Einstein equations*

$$\bar{G}_{\mu\nu}[\bar{\Sigma}] + \Lambda\,\bar{\Sigma}_{\mu\nu} = \kappa\,\bar{T}^{\mathrm{total}}_{\mu\nu}\,,\qquad \textit{(Definition 11.106).}\tag{11.218}$$

Sketch. By Definition 11.109, the operator–valued response kernel $\hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x')$ is the composition of: (i) the algebraic map to Einstein form $\mathbb{P}^{(\mathrm{Ein})}$, (ii) the retarded Green kernel of the linearized Lichnerowicz/Einstein operator on $(\bar{\Sigma},\bar{\nabla})$, and (iii) the source–symmetrizing map $\mathbb{J}$ (Remark 11.108). Taking the diagnostic realized reference configuration expectation on $\mathfrak{P}$ (Proposition 7.3) yields the c–number bitensor $\bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x')$ of Definition 11.111. Causality of the retarded kernel implies $\operatorname{supp}\bar{\mathcal{C}}(\cdot;x') \subset J^+_{\bar{\Sigma}}(x')$ (Proposition 11.14), ensuring that the right–hand side of (11.216) is microcausal on the same cone as used elsewhere. Metric compatibility and the bitensor construction give $\bar{\nabla}^\mu_x\bar{G}_{\mu\nu} \equiv 0$ and, provided $\bar{\nabla}'_\rho\bar{T}^\rho{}_\sigma(x') = 0$, $\bar{\nabla}^\mu_x$ of the right–hand side vanishes; thus the relational Bianchi identity and conservation law remain consistent (Remark 11.108). In the ultra–local limit (11.217) one recovers the standard Einstein coupling with constant κ, proving the reduction to Definition 11.106. □

Interpretation and consistency with the chapter plan. Equation (11.216) presents the Chronogeometric Einstein law in BRST/cohomological, band–regular form on $\mathfrak{P}$, with a state– and tag–dependent emergent coupling (Definition 11.111). The kernel is compatible with the typed/triadic realized weave configuration: it commutes with the sector projectors and, in the radiative spin–2 block, with the TT projector (Definitions 11.24 and 11.60 and Proposition 11.77), so TT–typed TLO diagnostics attach only to the radiative sector as prescribed in the plan. The local GR law Definition 11.106 is recovered as the ultra–local limit of the same structure, with κ playing the usual role of a (diagnostic realized reference configuration–normalized) coupling constant (Section 11.5.5).

11.5.7 Scalarisation, Universality, and Variation Bounds of the Gravitational Coupling Tensor

A. EP–safe scalarisation (when $\bar{\mathcal{C}}$ reduces to a scalar). Work on the diagnostic realized reference configuration $\mathfrak{P}$ with BRST–closed tags commuting with the triad and TT projectors

(Definitions 6.280 and 9.10 and Proposition 7.3). Let $\bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x)$ be the (expectation–level) Emergent Gravitational Coupling Tensor of Definition 11.111, entering the generalized Einstein response (Proposition 11.112). On the bundle $\mathrm{Sym}^2(T^*\mathcal{X}_q)$ of symmetric $(0,2)$ tensors define the identity and pure–trace projector

$$I_{\mu\nu}{}^{\rho\sigma} := \tfrac{1}{2}\big(\delta_\mu{}^\rho\delta_\nu{}^\sigma + \delta_\mu{}^\sigma\delta_\nu{}^\rho\big), \qquad (P_{\mathrm{tr}})_{\mu\nu}{}^{\rho\sigma} := \tfrac{1}{d}\,\bar{\Sigma}_{\mu\nu}\bar{\Sigma}^{\rho\sigma},$$

respectively [249, 146].[2]

Algebraic symmetry / self–adjointness. Assume $\bar{\mathcal{C}}$ is symmetric on Sym^2 with respect to the $\bar{\Sigma}$–inner product:

$$\bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma} = \bar{\mathcal{C}}_{\nu\mu}{}^{\rho\sigma} = \bar{\mathcal{C}}_{\mu\nu}{}^{\sigma\rho}, \qquad \bar{\mathcal{C}}_{\mu\nu,\rho\sigma} := \bar{\Sigma}_{\rho\alpha}\bar{\Sigma}_{\sigma\beta}\,\bar{\mathcal{C}}_{\mu\nu}{}^{\alpha\beta} = \bar{\mathcal{C}}_{\rho\sigma,\mu\nu}, \tag{11.219}$$

so a local $\bar{\Sigma}$–orthonormal frame diagonalises $\bar{\mathcal{C}}$ [536, 404].

Scalarisation ansatz. Decompose into an isotropic (species–blind) scalar and a trace–free anisotropic remainder:

$$\bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x) = \kappa_{\mathrm{eff}}(x)\, I_{\mu\nu}{}^{\rho\sigma} + \Delta_{\mu\nu}{}^{\rho\sigma}(x), \qquad \mathrm{tr}_{\mathrm{Sym}^2}\Delta = 0, \tag{11.220}$$

with the scalar defined by the operator trace on Sym^2,

$$\kappa_{\mathrm{eff}}(x) \;:=\; \frac{1}{N_d}\,\mathrm{tr}_{\mathrm{Sym}^2}\big[\bar{\mathcal{C}}(x)\big] \;=\; \frac{2}{d(d{+}1)}\,\bar{\mathcal{C}}_{\mu\nu}{}^{\mu\nu}(x), \qquad N_d = \tfrac{d(d+1)}{2} = \dim \mathrm{Sym}^2, \tag{11.221}$$

so that (11.220) is the unique orthogonal split (w.r.t. the operator trace).

Trace–free projection (unambiguous). For any endomorphism $\mathcal{E}_{\mu\nu}{}^{\rho\sigma}$ on Sym^2, define its pair–trace and $\bar{\Sigma}$–trace–free part by

$$(\mathcal{E}\cdot\bar{\Sigma})_{\mu\nu} := \mathcal{E}_{\mu\nu}{}^{\rho\sigma}\bar{\Sigma}_{\rho\sigma}, \qquad \mathrm{TF}_{\bar{\Sigma}}\,\mathcal{E} \;:=\; \mathcal{E} - \frac{1}{d}\,(\mathcal{E}\cdot\bar{\Sigma})\otimes\bar{\Sigma}^{\rho\sigma}, \tag{11.222}$$

so $(\mathrm{TF}_{\bar{\Sigma}}\,\mathcal{E})\cdot\bar{\Sigma} = 0$ for any d. (Here $(A\otimes B)_{\mu\nu}{}^{\rho\sigma} := A_{\mu\nu}B^{\rho\sigma}$.)

Norms and slow–variation regime. Let all norms be Hilbert–Schmidt norms induced by $\bar{\Sigma}$:

$$\|X\|^2_{\mathrm{HS}} := X_{\mu\nu}{}^{\rho\sigma}X_{\alpha\beta}{}^{\gamma\delta}\,\bar{\Sigma}^{\mu\alpha}\bar{\Sigma}^{\nu\beta}\bar{\Sigma}_{\rho\gamma}\bar{\Sigma}_{\sigma\delta}, \qquad \|Y\|^2_{\mathrm{HS}} := Y_{\mu\nu}Y_{\alpha\beta}\,\bar{\Sigma}^{\mu\alpha}\bar{\Sigma}^{\nu\beta}.$$

In the coherence–controlled regime (cf. Section 11.6), introduce

$$\varepsilon_{\mathrm{aniso}}(x) := \frac{\|\mathrm{TF}_{\bar{\Sigma}}\,\bar{\mathcal{C}}(x)\|_{\mathrm{HS}}}{\|\bar{\mathcal{C}}(x)\|_{\mathrm{HS}}} \ll 1, \qquad \varepsilon_{\mathrm{grad}}(x) := \max\Big\{L_{\mathrm{exp}}\tfrac{\|\nabla\bar{t}\|_{\mathrm{HS}}}{1+\|\bar{t}\|_{\mathrm{HS}}},\; L^2_{\mathrm{exp}}\tfrac{\|\nabla\nabla\bar{t}\|_{\mathrm{HS}}}{1+\|\bar{t}\|_{\mathrm{HS}}}\Big\} \ll 1, \tag{11.223}$$

with an experimental scale L_{exp}. Since $\bar{\mathcal{C}} = \langle\hat{\mathcal{C}}[\mathcal{U}_{\boldsymbol{\Phi\Theta}}[\bar{t},\ldots]]\rangle$ (Definitions 11.109 and 11.111), assume Lipschitz–type control

$$\|\nabla^k\bar{\mathcal{C}}\|_{\mathrm{HS}} \;\le\; L_k\,\|\nabla^k\bar{t}\|_{\mathrm{HS}}, \qquad k\in\{1,2\}, \tag{11.224}$$

for state–dependent L_k.

[2] For any symmetric $X_{\rho\sigma}$, $I_{\mu\nu}{}^{\rho\sigma}X_{\rho\sigma} = X_{\mu\nu}$. We write I to emphasize action on $\mathrm{End}(\mathrm{Sym}^2)$.

Local isotropy to leading order. Under (11.219)–(11.224), there exists a local $\bar{\Sigma}$–orthonormal frame with

$$\bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x) = \kappa_{\mathrm{eff}}(x)\, I_{\mu\nu}{}^{\rho\sigma} + \mathcal{O}\Big(\varepsilon_{\mathrm{aniso}} + \varepsilon_{\mathrm{grad}}\Big), \tag{11.225}$$

so the generalized field equation (Proposition 11.112) reduces locally to the GR form with a slowly varying effective coupling $\kappa_{\mathrm{eff}}(x)$ [6, 146]. In the radiative spin–2 block, compatibility with TT typing is automatic because $\bar{\mathcal{C}}$, built from the $\bar{\Sigma}$–Lichnerowicz/Einstein operator and retarded kernels, commutes with the TT projector (Definition 11.60 and Proposition 11.77).

Variation bound (diagnostic realized reference configuration form). Let $A : B$ denote full index contraction (bilinear pairing) on Sym^2. Using Cauchy–Schwarz for the Hilbert–Schmidt norm and (11.220), one has the pointwise bound

$$\boxed{\Big\| \Big(\bar{\mathcal{C}} - \kappa_{\mathrm{eff}} I\Big) : \bar{T}^{\mathrm{total}} \Big\|_{\mathrm{HS}} \;\le\; \|\Delta\|_{\mathrm{HS}}\, \|\bar{T}^{\mathrm{total}}\|_{\mathrm{HS}} \;\lesssim\; \Big(\varepsilon_{\mathrm{aniso}} + \varepsilon_{\mathrm{grad}}\Big)\, \|\bar{\mathcal{C}}\|_{\mathrm{HS}}\, \|\bar{T}^{\mathrm{total}}\|_{\mathrm{HS}},} \tag{11.226}$$

so departures from the scalarised law are suppressed by the anisotropy and slow–variation controls. Consequently, universality tests (free fall) are satisfied to the indicated accuracy whenever $\varepsilon_{\mathrm{aniso}}, \varepsilon_{\mathrm{grad}} \ll 1$ [479].

Species–blindness (EP universality). On the BRST/GNS physical sector $\mathcal{H}_{\mathrm{phys}}$ the coupling acts trivially on internal matter indices,

$$\bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma} \;\otimes\; \mathbf{1}_{\mathrm{int}},$$

so in the scalarised limit it multiplies the total stress tensor $\bar{T}^{\mathrm{total}}_{\rho\sigma}$, preserving universality of free fall [479]. This is consistent with the ultra–local GR limit $\bar{\mathcal{C}} \to \kappa\, I\, \delta^{(d)}(x-x')$ (Definition 11.106, Proposition 11.112, and Section 11.5.5).

B. Variation bounds compatible with EP/PPN tests. *Diagnostic realized reference configuration semantics and scope.* All statements below are evaluated on the certified plateau $\mathfrak{P}$ with BRST–closed tags commuting with the triad and TT projectors (Nielsen–flat readouts; Definitions 6.280 and 9.10 and Proposition 7.3). They constrain departures of the Emergent Gravitational Coupling Tensor $\bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}$ from its scalarised limit $\kappa_{\mathrm{eff}} I$ (Definition 11.111 and Section 11.5.7) in regimes probed by weak–field experiments (EP/PPN).

Instrument–facing tolerances. To parameterize observable deviations from a constant, isotropic coupling, introduce four tolerances $\mathcal{B}_\gamma$, $\mathcal{B}_{\mathrm{EP}}$, $\mathcal{B}_{\dot{G}}$, $\mathcal{B}_{\nabla G}$:

1. **Isotropy (PPN γ on AU scales).** Trace–free anisotropy of the coupling endomorphism is small,
$$\varepsilon_{\mathrm{aniso}} \;\lesssim\; \mathcal{B}_\gamma \qquad \text{(solar–system regime; cf. Section 11.5.7).}$$

2. **Universality of free fall (Eötvös–type bounds).** Composition dependence in the scalarised response obeys
$$\left.\frac{\Delta\kappa_{\mathrm{eff}}}{\kappa_{\mathrm{eff}}}\right|_{\mathrm{materials}} \;\lesssim\; \mathcal{B}_{\mathrm{EP}} \qquad \text{(near–Earth / Earth–orbit experiments).}$$

3. **Temporal drift (e.g. LLR).** The fractional time variation satisfies

$$\left|\frac{\dot{\kappa}_{\rm eff}}{\kappa_{\rm eff}}\right| \lesssim \mathcal{B}_{\dot{G}} \qquad \text{(per unit time)}.$$

4. **Spatial gradients (laboratory / planetary baselines).** Over a baseline L,

$$\left|\frac{\Delta\kappa_{\rm eff}}{\kappa_{\rm eff}}\right| \lesssim \mathcal{B}_{\nabla G}(L), \qquad \text{equivalently} \qquad \left|\frac{\nabla\kappa_{\rm eff}}{\kappa_{\rm eff}}\right| \lesssim \frac{\mathcal{B}_{\nabla G}(L)}{L}.$$

Calibration note: Concrete numerical values for $(\mathcal{B}_\gamma, \mathcal{B}_{\rm EP}, \mathcal{B}_{\dot{G}}, \mathcal{B}_{\nabla G})$ may be set to current best–fit limits in an appendix; the theory development here is independent of those numbers.

Covariant variation bounds from scalarisation. Using the definition of the effective scalar coupling (eq. (11.221)) and the Lipschitz–type controls linking $\bar{\mathcal{C}}$ to the temporal texture $\bar{t}$ (eq. (11.224)), one finds

$$\boxed{\left|\nabla\kappa_{\rm eff}\right| = \frac{2}{d(d+1)}\left|\nabla\bar{\mathcal{C}}_{\mu\nu}{}^{\mu\nu}\right| \le \frac{2}{d(d+1)}\,\|\nabla\bar{\mathcal{C}}\|_{\rm HS} \le \frac{2L_1}{d(d+1)}\,\|\nabla\bar{t}\|_{\rm HS},} \tag{11.227}$$

and similarly $\left|\nabla\nabla\kappa_{\rm eff}\right| \le \frac{2L_2}{d(d+1)}\,\|\nabla\nabla\bar{t}\|_{\rm HS}$, which dovetail with the slow–variation control $\varepsilon_{\rm grad}$ defined in Section 11.5.7. Together with the anisotropy bound $\varepsilon_{\rm aniso} \ll 1$ and species–blind action of $\bar{\mathcal{C}}$ on the BRST/GNS physical sector (Section 11.5.7), these estimates ensure compatibility with EP and weak–field PPN tests while remaining fully consistent with the generalized response law and its GR limit (Proposition 11.112, Definition 11.106, and Section 11.5.5).

C. EP–safe scalarisation.

Proposition 11.113 (EP–safe scalarisation)**.** *Work on a certified plateau $\mathfrak{P}$ with BRST–closed band tags commuting with the triad and TT projectors (Nielsen–flat readouts; Definitions 6.280 and 9.10 and Proposition 7.3). Let the coupling endomorphism $\bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}$ obey the symmetry/-self–adjointness in eq. (11.219) and the scalarisation split $\bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma} = \kappa_{\rm eff} I_{\mu\nu}{}^{\rho\sigma} + \Delta_{\mu\nu}{}^{\rho\sigma}$ with $\kappa_{\rm eff}$ defined in eq. (11.221). If, on a spacetime region, the anisotropy and slow–variation controls from Section 11.5.7 satisfy $\varepsilon_{aniso} \ll 1$ and $\varepsilon_{grad} \ll 1$, then the generalized law $\bar{G}_{\mu\nu} = \bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}\,\bar{T}^{\rm total}_{\rho\sigma}$ (Proposition 11.112) reduces to*

$$\bar{G}_{\mu\nu}(x) = \kappa_{\rm eff}(x)\,\bar{T}^{\rm total}_{\mu\nu}(x) + \mathcal{O}\Big(\varepsilon_{aniso} + \varepsilon_{grad}\Big),$$

with $\kappa_{\rm eff}$ species–blind *on the BRST/GNS physical sector (universality; Section 11.5.7). If, in addition, the drifts satisfy the instrument–facing bounds of Section 11.5.7 (items 1–4) together with the covariant estimates eq. (11.227), then the theory is compatible with current null–tests of the Equivalence Principle and weak–field PPN constraints [479].*

Sketch. Contracting the scalarisation split (eq. (11.220)) with $\bar{T}^{\rm total}$ gives $\bar{\mathcal{C}} : \bar{T}^{\rm total} = \kappa_{\rm eff}\,\bar{T}^{\rm total} + \Delta : \bar{T}^{\rm total}$. The symmetry eq. (11.219) allows a local $\bar{\Sigma}$–orthonormal frame that diagonalises $\bar{\mathcal{C}}$; the HS–norm control implies $\|\Delta : \bar{T}^{\rm total}\|_{\rm HS} \le \|\Delta\|_{\rm HS}\|\bar{T}^{\rm total}\|_{\rm HS} =$

$\mathcal{O}(\varepsilon_{\mathrm{aniso}})\|\bar{\mathcal{C}}\|_{\mathrm{HS}}\|\bar{T}^{\mathrm{total}}\|_{\mathrm{HS}}$. Slow variation ($\varepsilon_{\mathrm{grad}} \ll 1$) ensures that gradients of $\bar{\mathcal{C}}$ only induce subleading corrections in covariant conservation and matching conditions. Species–blindness (Sec. Section 11.5.7) guarantees that κ_{eff} multiplies the total stress tensor on the physical sector. Finally, the variation bounds eq. (11.227) together with the EP/PPN tolerances (items 1–4 in Section 11.5.7) place the residual deviations below experimental sensitivity, yielding the stated reduction and observational compatibility [479]. □

Remark 11.114 (Bianchi identity and conservation in the scalarised regime)**.** By the relational Bianchi identity (Remark 11.108), $\bar{\nabla}^{\mu}\bar{G}_{\mu\nu} = 0$, hence the generalized response law implies

$$\bar{\nabla}^{\mu}\Big(\bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}\,\bar{T}^{\mathrm{total}}_{\rho\sigma}\Big) = 0.$$

In the scalarised limit (Section 11.5.7),

$$(\bar{\nabla}^{\mu}\kappa_{\mathrm{eff}})\,\bar{T}^{\mathrm{total}}_{\mu\nu} + \kappa_{\mathrm{eff}}\,\bar{\nabla}^{\mu}\bar{T}^{\mathrm{total}}_{\mu\nu} = 0.$$

Thus GR–like conservation, $\bar{\nabla}^{\mu}\bar{T}^{\mathrm{total}}_{\mu\nu} \approx 0$, obtains whenever $\bar{\nabla}\kappa_{\mathrm{eff}}$ is negligible on the probed scales (controlled by eq. (11.227) and $\varepsilon_{\mathrm{grad}}$) or any residual exchange with the coupling sector is higher order in $(\varepsilon_{\mathrm{aniso}}, \varepsilon_{\mathrm{grad}})$ [7, 146]. This is consistent with the diagnostic realized reference configuration semantics and Nielsen–flatness of the readouts on $\mathfrak{P}$ (Proposition 7.3).

11.5.8 Recovery of General Relativity in the Spectral Limit

Classical gravity should emerge from the IQR dynamics in the regime of weak fields, low energies, and negligible coherence. In the typed/triadic language, this corresponds to working on a certified plateau $\mathfrak{P}$ (BRST–clean, Nielsen–flat) with band tags that commute with the triad projectors and the TT projector (Definitions 6.280 and 9.10 and Proposition 7.3), and then focusing on the *spectral* channel of the triad decomposition (Definition 11.24). In this limit the correlation (corr) and vacuum (vac) channels decouple from macroscopic matter dynamics, while the spectral (spec) channel dominates the geometry–matter response. At the level of the emergent coupling tensor this reads

$$\bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma} = \bar{\mathcal{C}}^{(\mathrm{corr})\rho\sigma}_{\mu\nu} + \bar{\mathcal{C}}^{(\mathrm{vac})\rho\sigma}_{\mu\nu} + \bar{\mathcal{C}}^{(\mathrm{spec})\rho\sigma}_{\mu\nu} \xrightarrow{\text{classical limit}} \bar{\mathcal{C}}^{(\mathrm{spec})\rho\sigma}_{\mu\nu},$$

with $\bar{\mathcal{C}}$ defined at diagnostic realized reference configuration level in Definition 11.111 and entering the generalized Einstein law of Proposition 11.112. Consistency with EP/PPN tests is ensured by the scalarisation analysis in Section 11.5.7 and Proposition 11.113.

Proposition 11.115 (Recovery of General Relativity)**.** *In the weak–field, low–energy, low–coherence regime on $\mathfrak{P}$, the spectral–channel coupling reduces to the identity endomorphism on* Sym^2 *times a constant,*

$$\bar{\mathcal{C}}^{(\mathrm{spec})\rho\sigma}_{\mu\nu}(x) \;\longrightarrow\; \kappa\,\Pi_{\mu\nu}{}^{\rho\sigma}, \qquad \Pi_{\mu\nu}{}^{\rho\sigma} := \tfrac{1}{2}\Big(\delta_{\mu}{}^{\rho}\delta_{\nu}{}^{\sigma} + \delta_{\mu}{}^{\sigma}\delta_{\nu}{}^{\rho}\Big),$$

with κ fixed by the Newtonian/PPN matching (Section 11.5.5). Consequently, the Chronogeometric Einstein equations recover the GR form,

$$\boxed{\bar{G}_{\mu\nu}[\bar{\Sigma}] = \kappa\,\bar{T}^{\mathrm{total}}_{\mu\nu}(x), \qquad (\textit{equivalently } \bar{G}_{\mu\nu} + \Lambda\bar{\Sigma}_{\mu\nu} = \kappa\,\bar{T}^{\mathrm{total}}_{\mu\nu}),} \tag{11.228}$$

coinciding with Definition 11.106.

Proof. Step 1 (Linearised spectral sector and TT support). Expand around a smooth background $\bar{\Sigma}$ in a $\bar{\Sigma}$–orthonormal frame, $\bar{\Sigma}_{\mu\nu} \mapsto \bar{\Sigma}_{\mu\nu} + \epsilon\,\delta\Sigma_{\mu\nu}$ with $\epsilon \ll 1$. In the spin–2 sector the typed fluctuation is the TT graviton $\hat{h}_{\mu\nu} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\delta\hat{g}_{\mu\nu}$ (Definitions 11.3 and 11.28). Its dynamics are governed by the typed Lichnerowicz operator $K^{[\mathrm{TT}]}$ (Definition 11.60), with retarded/advanced Green kernels and Pauli–Jordan bi–solution supported on the $\bar{\Sigma}$–cones (Proposition 11.14). TT Ward identities enforce that only physical polarizations couple to matter vertices (Proposition 11.77), and the triad decomposition is block–diagonal at the linearized level (Proposition 11.20).

Step 2 (Operator response and long–wavelength limit). The operator–valued coupling kernel is the causal composition

$$\hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x') = \mathbb{P}^{(\mathrm{Ein})\,\alpha\beta}_{\mu\nu}\,\hat{\Delta}^{\mathrm{ret}\,\gamma\delta}_{\alpha\beta}(x,x')\,\mathbb{J}_{\gamma\delta}{}^{\rho\sigma}, \qquad \text{(Definition 11.109)},$$

where $\hat{\Delta}^{\mathrm{ret}}$ is the (typed) retarded graviton kernel, $\mathbb{P}^{(\mathrm{Ein})}$ the linearized Einstein map, and $\mathbb{J}$ the symmetric matter vertex compatible with the conservation constraints. In the spectral channel and for wavelengths large compared to any microscopic length (long–wavelength-/low–frequency limit consistent with diagnostic realized reference configuration tags), the nonlocal kernel reduces to its local leading term,

$$\hat{\Delta}^{\mathrm{ret}\,\gamma\delta}_{\alpha\beta}(x,x') \;\sim\; C_0\,\Pi_{\alpha\beta}{}^{\gamma\delta}\,\delta^{(d)}(x-x')\,\hat{\mathbf{1}} \quad\Rightarrow\quad \hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x') \;\sim\; C_0\,\Pi_{\mu\nu}{}^{\rho\sigma}\,\delta^{(d)}(x-x')\,\hat{\mathbf{1}},$$

using that the TT projector commutes through the composition and preserves the symmetric identity on Sym^2 (TT purity; Propositions 11.20 and 11.77). Taking the Holo–State expectation (Definition 11.111) yields the c–number kernel with the same local structure.

Step 3 (Matching and conclusion). Plugging the local kernel into the generalized law (Proposition 11.112) gives $\bar{G}_{\mu\nu}(x) = C_0\,\Pi_{\mu\nu}{}^{\rho\sigma}\bar{T}^{\mathrm{total}}_{\rho\sigma}(x) = C_0\,\bar{T}^{\mathrm{total}}_{\mu\nu}(x)$. Fix $C_0 = \kappa$ by the Newtonian/PPN matching discussed in Section 11.5.5 (equivalently, by the scalarisation normalisation in Section 11.5.7), which recovers (11.228). The inclusion of Λ is immediate from the action–level derivation (Definition 11.105 and Section 11.5.5). □

Discussion. Equation (11.228) shows that, once (i) TT purity and triad block–diagonality hold (Propositions 11.20 and 11.77), (ii) the retarded kernels share the $\bar{\Sigma}$–microcausal cones (Proposition 11.14), and (iii) the coupling scalarises in the sense of Section 11.5.7, the IQR equations reduce to the standard Einstein equations with constant κ (diagnostic realized reference configuration–matched) and the usual conservation laws (Bianchi identity; Remark 11.108). This anchors the classical limit of the IQR framework squarely within GR while retaining a clear pathway for controlled departures outside the spectral, low–coherence regime.

In summary, the typed exchange in the spectral channel reproduces the classical response: by Definition 11.67 and eq. (11.137) and the recovery result Proposition 11.115, the sector–resolved amplitude obeys

$$\mathcal{A}^{(\mathrm{spec})}_{\mathrm{grav}} \xrightarrow[\text{weak field / low coherence}]{} \text{General Relativity},$$

consistently with the Chronogeometric Einstein law Definition 11.106 and its diagnostic realized reference configuration derivation Section 11.5.5. This is precisely the spectral–limit

realization of the classical "vertex–propagator–vertex" picture developed earlier for the TT sector (Section 11.3.5 and eq. (11.115)).

Signature & invertibility conditions (Lorentz window). Write the expectation–level unified tensor as $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$, with $\bar{t}^{\rm sym}_{\mu\nu} := \frac{1}{2}(\bar{t}_{\mu\nu} + \bar{t}_{\nu\mu})$ and $\bar{t}^{\rm anti}_{\mu\nu} := \frac{1}{2}(\bar{t}_{\mu\nu} - \bar{t}_{\nu\mu})$ (cf. Section 5.4 and eq. (11.232)). Define the effective Lorentzian metric extracted from the unified tensor, $\bar{g}^{\rm Lor}_{\mu\nu} := \bar{g}_{\mu\nu} + \bar{t}^{\rm sym}_{\mu\nu}$ (eq. (11.232)); this is the metric that carries the Relational Levi–Civita connection used throughout Section 11.5.1 and Remark 11.108.

Assumption 11.116 (Lorentz window for the temporal sector)**.** *There exists* $0 < \varepsilon < 1$ *such that, for all non–null* v^μ*,*

$$\frac{\left|\bar{t}^{\rm sym}_{\mu\nu} v^\mu v^\nu\right|}{\left|\bar{g}_{\mu\nu} v^\mu v^\nu\right|} \;\le\; \varepsilon, \qquad \textit{and} \qquad \left\| \, (\bar{g}^{\rm Lor})^{-1} \cdot \bar{t}^{\rm anti} \, \right\| \;<\; 1, \tag{11.229}$$

where $\|\cdot\|$ *is any operator norm induced by* $\bar{g}^{\rm Lor}$*.*

Proposition 11.117 (Persistence of Lorentz signature)**.** *Under Assumption 11.116, the quadratic form* $\bar{g}^{\rm Lor}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}^{\rm sym}_{\mu\nu}$ *is non–degenerate and has the same inertia as* $\bar{g}_{\mu\nu}$*. Hence,* $\mathrm{signature}(\bar{g}^{\rm Lor}) = (+,-,-,-)$ *in* $d = 4$ *[7, 146].*

Proof sketch. The Rayleigh–quotient bound implied by (11.229) gives $(1-\varepsilon)\,|\bar{g}(v,v)| \le |\bar{g}^{\rm Lor}(v,v)| \le (1+\varepsilon)\,|\bar{g}(v,v)|$ for all non–null v, along the homotopy $\bar{g}_\lambda := \bar{g} + \lambda\,\bar{t}^{\rm sym}$, $\lambda \in [0,1]$. No root or sign flip can occur without violating the bound, so non–degeneracy and inertia persist by continuity (Sylvester's law). The second inequality in (11.229) ensures the endomorphism $\mathrm{id} + (\bar{g}^{\rm Lor})^{-1}\cdot \bar{t}^{\rm anti}$ is invertible (Neumann series), guaranteeing that raising/lowering with $\bar{g}^{\rm Lor}$ and transporting with its Levi–Civita connection remain well–posed. □

Remark 11.118 (Why the Lorentz window matters here)**.** The bounds (11.229) ensure: (i) the causal cones defined by $\bar{g}^{\rm Lor}$ coincide perturbatively with those used in the TT/Peierls construction, so retarded/advanced kernels retain support on $J^{\pm}_{\bar{\Sigma}}$ (Proposition 11.14); (ii) parallel transport and Green–operator inversion on $(\mathcal{X}_q, \bar{\Sigma})$ are well–defined, which is required by the response formulas and effective actions in Section 11.3.5 and eq. (11.115); and (iii) the Bianchi identity and conservation statements proven on $\bar{\Sigma}$ (Remark 11.108 and Section 11.5.5) apply without modification, anchoring the GR recovery in Proposition 11.115.

Proposition 11.119 (Invertibility of $\bar{\Sigma}$ and control of the determinant)**.** *Assume the Lorentz window of Assumption 11.116 holds and, in particular, the second bound in (11.229). If* $\bar{g}^{\rm Lor}$ *is Lorentzian and non–degenerate (by Proposition 11.117), then*

$$\bar{\Sigma} = \bar{g}^{\rm Lor} + \bar{t}^{\rm anti} = \bar{g}^{\rm Lor}\cdot\Big(\mathbf{1} + (\bar{g}^{\rm Lor})^{-1}\bar{t}^{\rm anti}\Big)$$

is invertible. Moreover,

$$\det\bar{\Sigma} = \det\bar{g}^{\rm Lor}\;\det\Big(\mathbf{1} + (\bar{g}^{\rm Lor})^{-1}\bar{t}^{\rm anti}\Big) \neq 0,$$

and the Neumann series for $\Big(\mathbf{1} + (\bar{g}^{\rm Lor})^{-1}\bar{t}^{\rm anti}\Big)^{-1}$ *converges. Hence* $\bar{\Sigma}^{-1}$ *exists on any domain where (11.229) holds uniformly [536].*

Proof. Set $A := (\bar{g}^{\mathrm{Lor}})^{-1}\bar{t}^{\mathrm{anti}}$. The second bound in (11.229) gives $\|A\| < 1$ (for any operator norm induced by $\bar{g}^{\mathrm{Lor}}$), so the Neumann series

$$(\mathbf{1} + A)^{-1} = \sum_{n=0}^{\infty} (-A)^n$$

converges absolutely. Thus $\mathbf{1} + A$ is invertible and

$$\bar{\Sigma}^{-1} = (\mathbf{1} + A)^{-1} \cdot (\bar{g}^{\mathrm{Lor}})^{-1}.$$

Since $\det(\cdot)$ is multiplicative, $\det \bar{\Sigma} = \det \bar{g}^{\mathrm{Lor}} \ \det(\mathbf{1} + A)$, and $\det(\mathbf{1} + A) \neq 0$ because $\mathbf{1} + A$ is invertible. Non–degeneracy of $\bar{g}^{\mathrm{Lor}}$ is guaranteed by Proposition 11.117, hence $\det \bar{g}^{\mathrm{Lor}} \neq 0$, completing the proof. □

Remark 11.120 (Determinant and inverse: placement in the text)**.** The inverse $\bar{\Sigma}^{\mu\nu}$ exists precisely on open regions where $\bar{\Sigma}$ is non–degenerate (cf. the general inverse rule for the unified tensor in Definition 5.26 and the linearization formulas in Remark 11.108). Accordingly, the volume elements $\sqrt{-\det \bar{g}^{\mathrm{Lor}}}$ and $\sqrt{-\det \bar{\Sigma}}$ that appear in the measure $d\mu_{\bar{\Sigma}}$ and in the Relational Einstein–Hilbert action are well–defined on those regions (see Sections 11.5.3 and 11.5.5).

Corollary 11.121 (Stability under small field fluctuations)**.** *Let Assumption 11.116 hold with margin $\varepsilon < 1$ at a given background $(\bar{g}, \bar{t})$. Because $\bar{g}$ and $\bar{t}$ depend smoothly on the Holo-State and on the fields $(\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}})$, there exists $\eta > 0$ such that for all fluctuations with $\|\delta\bar{g}\| + \|\delta\bar{t}\| < \eta$ (induced by small operator variations $\delta\hat{\mathbf{\Phi}}$, $\delta\hat{\mathbf{\Theta}}$), the inequalities (11.229) persist. Hence Lorentzian signature (Proposition 11.117) and invertibility (Proposition 11.119) are* stable *under sufficiently small fluctuations [536].*

11.6 A Coherence–Energy Condition (Relational NEC)

All statements in this section are made on a certified plateau $\mathfrak{P}$ with BRST-closed band tags commuting with the triad and TT projectors; expectation values are taken on the physical space $\mathcal{H}_{\mathrm{phys}}$ (diagnostic realized reference configuration semantics; cf. Proposition 7.3 and Definition 6.280). The causal and differential structures are those of the expectation–level chronogeometric tensor $\bar{\Sigma}_{\mu\nu} := \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (Definition 5.26), with Levi–Civita connection and curvatures defined in Definitions 6.7 and 11.97 and Section 11.5.2. Null directions and cones are always taken with respect to $\bar{\Sigma}$ (or, equivalently, $\bar{g}^{\mathrm{Lor}}_{\mu\nu} := \bar{g}_{\mu\nu} + \bar{t}^{\mathrm{sym}}_{\mu\nu}$ when the Lorentz window holds; cf. eq. (11.232), Assumption 11.116, and Proposition 11.117).

Definition 11.122 (Relational Null Energy/Coherence Condition (RNEC))**.** Let k^μ be any $\bar{\Sigma}$–null vector, $\bar{\Sigma}_{\mu\nu}k^\mu k^\nu = 0$. Define the diagnostic realized reference configuration/expectation effective source

$$\boxed{\bar{T}^{\mathrm{eff}}_{\mu\nu} \ := \ \bar{T}^{\mathrm{total}}_{\mu\nu} \ + \ \gamma\, \mathcal{Q}_{\mu\nu}[\bar{t}]\ , \qquad \bar{T}^{\mathrm{total}}_{\mu\nu} \text{ as in } \textit{Definition } 9.358,} \tag{11.230}$$

where $\gamma \geq 0$ and $\mathcal{Q}_{\mu\nu}[\bar{t}]$ is a symmetric, diagnostic realized reference configuration-safe, covariant functional of the temporal sector that vanishes when $\bar{t}_{\mu\nu} = 0$ (coherence off). The *Relational NEC* requires

$$\boxed{\forall\, k^\mu \ \bar{\Sigma}\text{–null:} \qquad k^\mu k^\nu \, \bar{T}^{\text{eff}}_{\mu\nu} \ \geq \ 0\,.} \tag{11.231}$$

Comments and normalization.

- **Vacuum piece.** If the cosmological term is placed on the right of the chronogeometric Einstein law $\bar{G}_{\mu\nu}[\bar{\Sigma}] + \Lambda \bar{\Sigma}_{\mu\nu} = \kappa\, \bar{T}^{\text{total}}_{\mu\nu}$ (Definition 11.106), one may incorporate it into $\bar{T}^{\text{total}}$ via $\bar{T}^{\text{vac}}_{\mu\nu} := -(\Lambda/\kappa)\, \bar{\Sigma}_{\mu\nu}$. This does not affect (11.231) because $k^\mu k^\nu \bar{\Sigma}_{\mu\nu} = 0$ for $\bar{\Sigma}$–null k^μ.
- **GR limit and focusing.** In the decohered regime $\bar{t}_{\mu\nu} \to 0$ (hence $\mathcal{Q}_{\mu\nu} \to 0$), (11.231) reduces to the standard NEC, $k^\mu k^\nu \bar{T}^{\text{total}}_{\mu\nu} \geq 0$, reproducing the classical focusing theorems for null congruences [240, 7].
- **Cone, causality, and Peierls.** Using $\bar{\Sigma}$–null k^μ aligns the energy condition with the same cone that governs the linearized Lichnerowicz operator and its retarded/advanced Green functions; the Pauli–Jordan kernel is supported on $J^{\pm}_{\bar{\Sigma}}$ and the covariant algebra follows by Peierls (Remark 11.108 and Proposition 11.14).

Physical sector and TT consistency. All expectations are evaluated on $\mathcal{H}_{\text{phys}}$ with Nielsen-flat law channel (Proposition 7.3). The temporal–curvature fluctuations $\hat{\tau}_{\mu\nu} := \delta\hat{\bar{t}}_{\mu\nu}$ furnish the coherence contribution $\mathcal{Q}_{\mu\nu}[\bar{t}]$ but do *not* carry spin–2 radiative content: τ–typed insertions are TT-null and the typed Ward identities enforce transversality/tracelessness in the graviton channel (Proposition 11.77 and Corollary 11.73). Thus the RNEC adds coherence control to the source without spoiling the TT graviton algebra or its causal support.

Equivalent presentations. One may keep $\bar{T}^{\text{total}}$ on the right-hand side of the Einstein equation and view $\gamma\, \mathcal{Q}_{\mu\nu}$ as an additive, state-dependent correction to the source diagnostics; or reabsorb (parts of) $\mathcal{Q}_{\mu\nu}$ into an effective, slowly varying coupling $\kappa_{\text{eff}}(x)$ when the scalarisation conditions of Section 11.5.7 hold. Both presentations are diagnostic realized reference configuration-equivalent and reduce to GR with constant κ in the spectral, low-coherence limit (Section 11.5.8).

Operational reading. The RNEC is the instrument-facing positivity that ties the chronogeometric law to causal propagation and dispersion: it uses the same cone as the TT kernels, is compatible with BRST/diagnostic realized reference configuration discipline, and collapses to the standard NEC when coherence is negligible. Standard vacuum-stress treatments on curved backgrounds provide the reference behavior for the matter/vacuum pieces of $\bar{T}^{\text{total}}$ [244, 11], while the coherence addendum $\mathcal{Q}_{\mu\nu}$ is state-controlled and vanishes in the GR limit.

1. Kinematic setup and the null cone. Let the expectation–level chronogeometric tensor be $\bar{\Sigma}_{\mu\nu} := \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (Definition 5.26), with $\bar{t}_{\mu\nu}$ the temporal–curvature tensor from the temporal functional (Definition 5.17). Following the causal/temporal structures of Section 5.4, split $\bar{t}_{\mu\nu}$ into symmetric/antisymmetric parts,

$$\bar{t}^{\mathrm{sym}}_{\mu\nu} := \tfrac{1}{2}(\bar{t}_{\mu\nu} + \bar{t}_{\nu\mu}), \qquad \bar{t}^{\mathrm{anti}}_{\mu\nu} := \tfrac{1}{2}(\bar{t}_{\mu\nu} - \bar{t}_{\nu\mu}),$$

and define the effective Lorentzian tensor

$$\bar{g}^{\mathrm{Lor}}_{\mu\nu} \;:=\; \bar{g}_{\mu\nu} + \bar{t}^{\mathrm{sym}}_{\mu\nu}, \tag{11.232}$$

assumed smooth with Lorentzian signature on the domain of interest (Assumption 11.116 and Proposition 11.117). A (contravariant) vector k^μ is *null* iff

$$k^\mu k^\nu\, \bar{g}^{\mathrm{Lor}}_{\mu\nu} = 0, \tag{11.233}$$

and we use the Levi–Civita connection of $\bar{g}^{\mathrm{Lor}}$ as the *relational* connection $\bar{\nabla} \equiv \nabla^{(\mathrm{rel})}$ (cf. Definition 6.7) for causal/kinematic constructions. In particular, null congruences satisfy

$$k^\nu \bar{\nabla}_\nu k^\mu = 0, \qquad \frac{d\theta}{d\lambda} = -\frac{1}{2}\theta^2 - \sigma_{\mu\nu}\sigma^{\mu\nu} + \omega_{\mu\nu}\omega^{\mu\nu} - \bar{R}_{\mu\nu}[\bar{g}^{\mathrm{Lor}}]\, k^\mu k^\nu, \tag{11.234}$$

where $(\theta, \sigma_{\mu\nu}, \omega_{\mu\nu})$ are the expansion, shear, and twist of the congruence defined with $\bar{g}^{\mathrm{Lor}}$ (cf. [240, 7] and Definition 11.97 and Section 11.5.2).

Lemma 11.123 (Null–cone equivalence and vacuum drop-out)**.** *For any vector k^μ, $k^\mu k^\nu\, \bar{\Sigma}_{\mu\nu} = k^\mu k^\nu\, \bar{g}^{\mathrm{Lor}}_{\mu\nu}$. Hence k^μ is $\bar{g}^{\mathrm{Lor}}$-null iff it is $\bar{\Sigma}$-null; moreover, for the cosmological vacuum stress $\bar{T}^{\mathrm{vac}}_{\mu\nu} = -(\Lambda/\kappa)\,\bar{\Sigma}_{\mu\nu}$ (cf. Definition 11.106), one has $k^\mu k^\nu \bar{T}^{\mathrm{vac}}_{\mu\nu} = 0$ on null k^μ.*

Proof. Write $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}^{\mathrm{sym}}_{\mu\nu} + \bar{t}^{\mathrm{anti}}_{\mu\nu}$. Because $k^\mu k^\nu$ is symmetric, $k^\mu k^\nu \bar{t}^{\mathrm{anti}}_{\mu\nu} = 0$. Thus $k^\mu k^\nu \bar{\Sigma}_{\mu\nu} = k^\mu k^\nu (\bar{g}_{\mu\nu} + \bar{t}^{\mathrm{sym}}_{\mu\nu}) = k^\mu k^\nu \bar{g}^{\mathrm{Lor}}_{\mu\nu}$, proving both cone equivalence and vacuum drop-out. □

Consequently, in evaluating the Relational NEC (Definition 11.122), the null directions determined by $\bar{g}^{\mathrm{Lor}}$ coincide with those of $\bar{\Sigma}$, and the vacuum term makes no contribution to the null contraction. This isolates the matter and coherence stresses as the only kinematic sources relevant for focusing and energy–condition analyses on the relational cone (Definitions 9.358 and 11.122).

2. The coherence stress $\mathcal{Q}_{\mu\nu}[\bar{t}]$. Decompose the temporal–curvature tensor into symmetric/antisymmetric parts,

$$\bar{t}_{\mu\nu} = S_{\mu\nu} + A_{\mu\nu}, \qquad S_{\mu\nu} := \tfrac{1}{2}(\bar{t}_{\mu\nu} + \bar{t}_{\nu\mu}), \quad A_{\mu\nu} := \tfrac{1}{2}(\bar{t}_{\mu\nu} - \bar{t}_{\nu\mu}).$$

Index convention in this subsection: when forming $\mathcal{Q}_{\mu\nu}$ and working on the null screen, raise/lower with $\bar{g}^{\mathrm{Lor}}$; for null contractions, the $\bar{\Sigma}$–trace piece drops out by Lemma 11.123.

Motivated by the electromagnetic/superenergy template [146, 7], define the *coherence stress* as the (dimension-adjusted) quadratic:

$$\boxed{\mathcal{Q}_{\mu\nu}[\bar{t}] := S_{\mu\alpha}\,S_{\nu}{}^{\alpha} + A_{\mu\alpha}\,A_{\nu}{}^{\alpha} - \frac{1}{d}\,\bar{g}^{\mathrm{Lor}}_{\mu\nu}\Big(S_{\alpha\beta}S^{\alpha\beta} + A_{\alpha\beta}A^{\alpha\beta}\Big)\,,} \tag{11.235}$$

where $d = \dim \mathcal{X}_q$. *In $d = 4$, (11.235) coincides with the earlier $1/4$ normalization.*

Basic properties.

- *Symmetry.* $\mathcal{Q}_{\mu\nu} = \mathcal{Q}_{\nu\mu}$, since each term in (11.235) is manifestly symmetric in (μ, ν).
- *$\bar{g}^{\mathrm{Lor}}$–tracefree (dimension-exact).* Contracting (11.235) with $\bar{g}^{\mathrm{Lor}\,\mu\nu}$ gives

$$\bar{g}^{\mathrm{Lor}\,\mu\nu}\mathcal{Q}_{\mu\nu} = S_{\alpha\beta}S^{\alpha\beta} + A_{\alpha\beta}A^{\alpha\beta} - \frac{d}{d}\Big(S_{\alpha\beta}S^{\alpha\beta} + A_{\alpha\beta}A^{\alpha\beta}\Big) = 0.$$

  (If you fix $d = 4$ from the outset, replace $1/d$ by $1/4$ to keep this property.)

- *Null positivity (Relational NEC contribution).* For any $\bar{g}^{\mathrm{Lor}}$–null k^μ,

$$\begin{aligned} k^\mu k^\nu \mathcal{Q}_{\mu\nu} &= (S{\cdot}k)_\alpha (S{\cdot}k)^\alpha + (A{\cdot}k)_\alpha (A{\cdot}k)^\alpha \\ &\quad - \frac{1}{d}\,(\bar{g}^{\mathrm{Lor}}_{\mu\nu}k^\mu k^\nu)\,\Big(S_{\rho\sigma}S^{\rho\sigma} + A_{\rho\sigma}A^{\rho\sigma}\Big) \\ &= \|(S{\cdot}k)\|^2 + \|(A{\cdot}k)\|^2 \;\geq 0. \end{aligned}$$

  so $\mathcal{Q}_{\mu\nu}$ contributes non-negatively to the Relational NEC (Section 11.6, and vanishes in the decohered limit $\bar{t} \to 0$.

- *Quadratic scaling and vanishing.* For any scalar λ, $\mathcal{Q}_{\mu\nu}[\lambda\bar{t}] = \lambda^2\,\mathcal{Q}_{\mu\nu}[\bar{t}]$; in particular, $\bar{t} \equiv 0 \Rightarrow \mathcal{Q}_{\mu\nu} \equiv 0$.
- *Local covariance.* $\mathcal{Q}_{\mu\nu}$ is a bona-fide $(0,2)$ tensor built locally from $(\bar{g}^{\mathrm{Lor}}, \bar{t})$; diffeomorphisms act covariantly. Any further divergence property follows from the evolution/-constraint equations for $\bar{t}$ (see eq. (16.467)) and the Bianchi identity for $\bar{G}_{\mu\nu}$.
- *TT isolation and SM orthogonality.* With $\mathcal{P}^{(2)}[\hat{\tau}] = 0$ and the coherence projector $\mathcal{P}^{\mathrm{coh}}_G$ (eq. (13.113) and Definition 13.1), the coherence channel carries no spin-2 radiative content and remains orthogonal to SM excitations; this keeps the TT graviton sector (Chapter 11) clean while allowing $\mathcal{Q}_{\mu\nu}$ to enter energy-condition and focusing analyses.

Lemma 11.124 (Vacuum term vanishes on nulls)**.** *For any $\bar{g}^{\mathrm{Lor}}$–null covector k^μ,*

$$k^\mu k^\nu\, T^{\mathrm{vac}}_{\mu\nu} = -(\Lambda/\kappa')\,k^\mu k^\nu\,\bar{\Sigma}_{\mu\nu} = 0.$$

Proof. By Lemma 11.123, $k^\mu k^\nu \bar{\Sigma}_{\mu\nu} = k^\mu k^\nu(\bar{g}_{\mu\nu} + \bar{t}^{\mathrm{sym}}_{\mu\nu}) = k^\mu k^\nu \bar{g}^{\mathrm{Lor}}_{\mu\nu} = 0$ for any $\bar{g}^{\mathrm{Lor}}$–null k^μ; the antisymmetric part does not contribute because $k^\mu k^\nu$ is symmetric. Hence the result. □

Lemma 11.125 (Positivity of the coherence term on nulls (sufficient conditions))**.** *Let k^μ be $\bar{g}^{\mathrm{Lor}}$–null and write $\bar{t}_{\mu\nu} = S_{\mu\nu} + A_{\mu\nu}$ with $S_{\mu\nu} = S_{\nu\mu}$ and $A_{\mu\nu} = -A_{\nu\mu}$. Then:*

1. *The antisymmetric part contributes non–negatively:*

$$k^\mu k^\nu\, A_{\mu\alpha}A_\nu{}^\alpha \;\geq\; 0.$$

2. *If, in addition, the* screen–orthogonality *condition*

$$k^\mu k^\nu\, S_{\mu\nu} = 0$$

holds along the null congruence (no longitudinal squeezing), then

$$k^\mu k^\nu\, S_{\mu\alpha}S_\nu{}^\alpha \;\geq\; 0.$$

Consequently, with the same condition one has

$$k^\mu k^\nu\, \mathcal{Q}_{\mu\nu}[\bar{t}] \;\geq\; 0$$

for the coherence stress defined in (11.235).

Proof. Choose an auxiliary null covector ℓ^μ with $k\cdot\ell = -1$ and define the *screen* (transverse) metric

$$q_{\mu\nu} := \bar{g}^{\mathrm{Lor}}_{\mu\nu} + k_\mu\ell_\nu + \ell_\mu k_\nu,$$

which is positive–definite on the 2–plane orthogonal to $\{k,\ell\}$ (see, e.g., [7]). Define the transverse vectors $V^\alpha := (S\cdot k)^\alpha := S^\alpha{}_\beta k^\beta$ and $W^\alpha := (A\cdot k)^\alpha := A^\alpha{}_\beta k^\beta$.

(1) Antisymmetric part. Since $A_{\mu\nu}$ is antisymmetric, $k_\alpha W^\alpha = A_{\alpha\beta}k^\alpha k^\beta = 0$, so W^α lies in the screen. Hence

$$k^\mu k^\nu\, A_{\mu\alpha}A_\nu{}^\alpha = W_\alpha W^\alpha = q_{\alpha\beta}\, W^\alpha W^\beta \;\geq\; 0,$$

because $q_{\alpha\beta}$ is positive–definite on the screen.

(2) Symmetric part. If $k^\mu k^\nu S_{\mu\nu} = 0$, then $k_\alpha V^\alpha = S_{\alpha\beta}k^\alpha k^\beta = 0$, so V^α is also purely transverse. Therefore

$$k^\mu k^\nu\, S_{\mu\alpha}S_\nu{}^\alpha = V_\alpha V^\alpha = q_{\alpha\beta}\, V^\alpha V^\beta \;\geq\; 0.$$

Finally, in the null contraction of (11.235) the trace term drops out because $k^\mu k^\nu \bar{g}^{\mathrm{Lor}}_{\mu\nu} = 0$, so $k^\mu k^\nu \mathcal{Q}_{\mu\nu} = V_\alpha V^\alpha + W_\alpha W^\alpha \geq 0$ under the stated condition. □

Remark 11.126 (Unconditional positivity via a screen–improved definition)**.** If one prefers a version that is *unconditionally* non–negative on nulls (without the extra condition on S), replace (11.235) by the *screen–projected* quadratic

$$\mathcal{Q}^\perp_{\mu\nu} := \left(S^\perp_{\mu\alpha}S^\perp_{\nu\beta} + A^\perp_{\mu\alpha}A^\perp_{\nu\beta}\right) q^{\alpha\beta} - \frac{1}{d}\, q_{\mu\nu}\left(S^\perp_{\alpha\beta}S^\perp_{\gamma\delta} + A^\perp_{\alpha\beta}A^\perp_{\gamma\delta}\right) q^{\alpha\gamma}q^{\beta\delta},$$

with $S^\perp_{\mu\nu} := q_\mu{}^\rho q_\nu{}^\sigma S_{\rho\sigma}$ and similarly for $A^\perp_{\mu\nu}$. Then

$$k^\mu k^\nu\, \mathcal{Q}^\perp_{\mu\nu} = q_{\alpha\beta}(S\cdot k)^\alpha(S\cdot k)^\beta + q_{\alpha\beta}(A\cdot k)^\alpha(A\cdot k)^\beta \;\geq\; 0$$

for every null k^μ, by positive–definiteness of $q_{\alpha\beta}$ [537]. This "Bel–type improvement" leaves local NEC tests and the Einstein equation unaffected at the level of null contractions used in Section 11.6, while making the positivity manifest [538, 537].

Remark 11.127 (Bel–type improvement and manifest null-positivity)**.** Fix a null pair (k^μ, ℓ^μ) with $k^\mu k_\mu = 0$, $\ell^\mu \ell_\mu = 0$ and $k\cdot\ell = -1$, and let the screen metric be

$$q_{\mu\nu} := \bar{g}^{\mathrm{Lor}}_{\mu\nu} + k_\mu \ell_\nu + \ell_\mu k_\nu, \qquad q^\mu{}_\nu := \bar{g}^{\mu\rho} q_{\rho\nu},$$

which is positive–definite on the $(d-2)$–dimensional subspace orthogonal to $\{k, \ell\}$ and obeys $k^\mu q_{\mu\nu} = q_{\mu\nu} k^\nu = 0$ [537]. Define the screen–projected tensors $S^\perp_{\mu\nu} := q_\mu{}^\rho q_\nu{}^\sigma S_{\rho\sigma}$ and $A^\perp_{\mu\nu} := q_\mu{}^\rho q_\nu{}^\sigma A_{\rho\sigma}$ and set

$$\mathcal{Q}^\perp_{\mu\nu} := \left(S^\perp_{\mu\alpha} S^\perp_{\nu\beta} + A^\perp_{\mu\alpha} A^\perp_{\nu\beta}\right) q^{\alpha\beta} \;-\; \frac{1}{d}\, q_{\mu\nu} \left(S^\perp_{\alpha\beta} S^\perp_{\gamma\delta} + A^\perp_{\alpha\beta} A^\perp_{\gamma\delta}\right) q^{\alpha\gamma} q^{\beta\delta}. \tag{11.236}$$

This is the "Bel–type improvement'' of $\mathcal{Q}_{\mu\nu}$ (cf. (11.235)) [538, 537]. It has the following properties:

1. **Screen–tracelessness.** $q^{\mu\nu}\mathcal{Q}^\perp_{\mu\nu} = 0$ by construction.
2. **Manifest null–positivity.** For every null k^μ,

$$k^\mu k^\nu\, \mathcal{Q}^\perp_{\mu\nu} = q_{\alpha\beta}(S\cdot k)^\alpha (S\cdot k)^\beta + q_{\alpha\beta}(A\cdot k)^\alpha (A\cdot k)^\beta \;\geq\; 0, \tag{11.237}$$

   where $(S\cdot k)^\alpha := S^\alpha{}_\beta k^\beta$ and $(A\cdot k)^\alpha := A^\alpha{}_\beta k^\beta$, using the positive–definiteness of $q_{\alpha\beta}$ on the screen [537].
3. **Null–contraction invariance (for NEC/NCC steps).** Since $k^\mu k^\nu q_{\mu\nu} = 0$, the trace–subtraction term in (11.236) drops out under $k^\mu k^\nu(\,\cdot\,)$, and the screen projection does not alter the right–hand side of (11.237). Consequently,

$$k^\mu k^\nu \mathcal{Q}^\perp_{\mu\nu} = k^\mu k^\nu \mathcal{Q}_{\mu\nu},$$

   so local null–energy tests and the null contractions entering the relational Einstein equations in Section 11.6, Theorem 11.128 and Corollary 11.129 are unchanged, while positivity becomes manifest via (11.237) [537].

Theorem 11.128 (NCC from Coherence Positivity)**.** *Assume (on the BRST/GNS physical sector) that the ordinary matter sector satisfies the NEC with respect to $\bar{g}^{\mathrm{Lor}}$, i.e. $k^\mu k^\nu T^{\mathrm{matter}}_{\mu\nu} \geq 0$ for all $\bar{g}^{\mathrm{Lor}}$–null k^μ. If $\gamma \geq 0$ in (11.230), then for all such k^μ,*

$$k^\mu k^\nu T^{\mathrm{eff}}_{\mu\nu} = k^\mu k^\nu T^{\mathrm{matter}}_{\mu\nu} \;+\; \gamma\, k^\mu k^\nu \mathcal{Q}_{\mu\nu}[\bar{t}] \;\geq\; 0,$$

so the Relational NEC *holds.*

Proof. By definition (11.230), $T^{\mathrm{eff}}_{\mu\nu} = T^{\mathrm{matter}}_{\mu\nu} + T^{\mathrm{vac}}_{\mu\nu} + \gamma\, \mathcal{Q}_{\mu\nu}[\bar{t}]$ with $T^{\mathrm{vac}}_{\mu\nu} = -(\Lambda/\kappa')\,\bar{\Sigma}_{\mu\nu}$. Lemma Lemma 11.124 shows $k^\mu k^\nu T^{\mathrm{vac}}_{\mu\nu} = 0$ on $\bar{g}^{\mathrm{Lor}}$–null k^μ, and Lemma Lemma 11.125 gives $k^\mu k^\nu \mathcal{Q}_{\mu\nu}[\bar{t}] \geq 0$. With the matter NEC and $\gamma \geq 0$, the sum is non-negative. □

Corollary 11.129 (Relational NCC under the field equations)**.** *Assume the hypotheses and conclusion of Theorem 11.128, namely that for every $\bar{g}^{\mathrm{Lor}}$–null k^μ,*

$$k^\mu k^\nu T^{\mathrm{eff}}_{\mu\nu} \;\geq\; 0, \qquad T^{\mathrm{eff}}_{\mu\nu} = T^{\mathrm{matter}}_{\mu\nu} + \gamma\, \mathcal{Q}_{\mu\nu}[\bar{t}] \;\; \textit{with} \;\; \gamma \geq 0 \;\; \textit{(cf. (11.230)).} \tag{11.238}$$

If, in addition, the relational Einstein equations hold on the same background,

$$\bar{G}_{\mu\nu}[\bar{g}^{\mathrm{Lor}}] + \Lambda\, \bar{g}^{\mathrm{Lor}}_{\mu\nu} \;=\; \kappa\, T^{\mathrm{eff}}_{\mu\nu}, \tag{11.239}$$

then for all $\bar{g}^{\mathrm{Lor}}$–null k^μ one has the Relational Null Convergence Condition (Relational NCC)

$$\boxed{R_{\mu\nu}[\bar{g}^{\mathrm{Lor}}]\, k^\mu k^\nu \;\geq\; 0.} \tag{11.240}$$

Proof. Contract (11.239) with $k^\mu k^\nu$ and use $G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2} R\, \bar{g}^{\mathrm{Lor}}_{\mu\nu}$:

$$k^\mu k^\nu R_{\mu\nu} \;-\; \tfrac{1}{2} R\, k^\mu k^\nu \bar{g}^{\mathrm{Lor}}_{\mu\nu} \;+\; \Lambda\, k^\mu k^\nu \bar{g}^{\mathrm{Lor}}_{\mu\nu} \;=\; \kappa\, k^\mu k^\nu T^{\mathrm{eff}}_{\mu\nu}.$$

Since k^μ is null, $k^\mu k^\nu \bar{g}^{\mathrm{Lor}}_{\mu\nu} = 0$, so the trace and cosmological terms vanish. Hence

$$k^\mu k^\nu R_{\mu\nu} \;=\; \kappa\, k^\mu k^\nu T^{\mathrm{eff}}_{\mu\nu}.$$

By Theorem 11.128 and the displayed inequality (11.238), the right-hand side is nonnegative for all null k^μ. Therefore $R_{\mu\nu} k^\mu k^\nu \geq 0$, which is (11.240). □

3. Raychaudhuri focusing and the scalarised coupling. Contract the generalized Einstein law (Proposition 11.112) along a null k^μ:

$$R_{\mu\nu}[\bar{g}^{\mathrm{Lor}}]\, k^\mu k^\nu = k^\mu k^\nu\, \bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x)\, \bar{T}^{\mathrm{total}}_{\rho\sigma}(x).$$

In the EP-safe (scalarised) regime of Section 11.5.7 one has

$$R_{\mu\nu}[\bar{g}^{\mathrm{Lor}}]\, k^\mu k^\nu = \kappa_{\mathrm{eff}}(x)\, k^\mu k^\nu \Big(T^{\mathrm{matter}}_{\mu\nu} + \gamma\, \mathcal{Q}_{\mu\nu}[\bar{t}] \Big),$$

since the vacuum term drops out by Lemma Lemma 11.124. Inserting this into the relational Raychaudhuri equation (11.234) with vanishing twist ($\omega_{\mu\nu} = 0$) yields

$$\frac{d\theta}{d\lambda} = -\frac{1}{2}\theta^2 - \sigma_{\mu\nu}\sigma^{\mu\nu} - \kappa_{\mathrm{eff}}\, k^\mu k^\nu \Big(T^{\mathrm{matter}}_{\mu\nu} + \gamma\, \mathcal{Q}_{\mu\nu}[\bar{t}] \Big) \;\leq\; -\frac{1}{2}\theta^2,$$

whenever the Relational NCC holds (Theorem Theorem 11.128). Therefore, if $\theta(0) < 0$ initially,

$$\theta(\lambda) \to -\infty \quad \text{by} \quad \lambda \leq \frac{2}{|\theta(0)|},$$

recovering the standard focusing bound, with κ_{eff} absorbed into the source normalization (cf. Section 11.5.7).

Remark 11.130 (NEC, Relational NEC, and NCC in the $\bar{\Sigma}$–framework)**.** In this chapter all null tests are performed with respect to the background Lorentzian metric $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ and null vectors k^μ (cf. Section 11.6). The three conditions used in Section 11.6 and in Theorem 11.128 and Corollary 11.129 are:

1. **NEC (standard Null Energy Condition).** For every $\bar{g}^{\mathrm{Lor}}$–null k^μ,
$$k^\mu k^\nu \bar{T}_{\mu\nu} \geq 0.$$
This is a *matter–side* inequality (domain: $\bar{T}_{\mu\nu}$).
2. **Relational NEC.** With the relational/effective source
$$T^{\mathrm{rel}}_{\mu\nu} := \bar{T}_{\mu\nu} + \mathcal{Q}_{\mu\nu}.$$
(See eq. (11.235).)
Here we prove the *null–positivity* of the $\mathcal{Q}$–piece:
$$k^\mu k^\nu \mathcal{Q}_{\mu\nu} = V_\alpha V^\alpha + W_\alpha W^\alpha \geq 0,$$
where $q_{\mu\nu} = \bar{g}^{\mathrm{Lor}}_{\mu\nu} + k_\mu \ell_\nu + \ell_\mu k_\nu$ is the screen metric and $V^\alpha := (S{\cdot}k)^\alpha$, $W^\alpha := (A{\cdot}k)^\alpha$ are the transverse projections; positivity follows from the positive–definiteness of $q_{\alpha\beta}$ on the screen and the null–trace drop $k^\mu k^\nu \bar{g}^{\mathrm{Lor}}_{\mu\nu} = 0$ (cf. Section 11.6 and Remark 11.127). Thus $k^\mu k^\nu T^{\mathrm{rel}}_{\mu\nu} \geq 0$ holds whenever $k^\mu k^\nu \bar{T}_{\mu\nu} \geq 0$ (and even beyond, by the added non–negative $\mathcal{Q}$).
3. **NCC (Null Convergence Condition).** A *geometry–side* requirement on the Ricci tensor of $\bar{g}^{\mathrm{Lor}}$:
$$R_{\mu\nu}\, k^\mu k^\nu \geq 0 \quad \text{for all null } k^\mu.$$
It is used with Raychaudhuri to infer focusing of null congruences.

Same vs. different, and how they connect.

- *Domains.* NEC constrains the matter tensor $\bar{T}_{\mu\nu}$; Relational NEC constrains the effective source $T^{\mathrm{rel}}_{\mu\nu} = \bar{T}_{\mu\nu} + \mathcal{Q}_{\mu\nu}$; NCC constrains the geometry $R_{\mu\nu}$.
- *Null–trace identity.* For any null k^μ, $k^\mu k^\nu \bar{g}^{\mathrm{Lor}}_{\mu\nu} = 0$; this removes metric trace terms and is the algebraic reason the screen argument isolates $V_\alpha V^\alpha + W_\alpha W^\alpha$ in $k^\mu k^\nu \mathcal{Q}_{\mu\nu}$ (cf. Section 11.6).
- *Implications under the field equations.* With the relational Einstein equations (11.239),
$$k^\mu k^\nu R_{\mu\nu} = \kappa\, k^\mu k^\nu T^{(\cdot)}_{\mu\nu} \quad \text{(trace terms drop by nullness)},$$
so *NEC* $\Rightarrow$ *NCC* in classical GR, and *Relational NEC* $\Rightarrow$ *NCC* in the relational theory (see Theorem 11.128 and Corollary 11.129).
- *Relative strength.* Because $\mathcal{Q}_{\mu\nu}$ is null–positive, the Relational NEC is (weakly) easier to satisfy than the standard NEC: even if $\bar{T}_{\mu\nu}k^\mu k^\nu < 0$ slightly, a sufficiently positive $k^\mu k^\nu \mathcal{Q}_{\mu\nu}$ can still yield $k^\mu k^\nu T^{\mathrm{rel}}_{\mu\nu} \geq 0$ (cf. Section 11.6).
- *Saturation of the $\mathcal{Q}$–piece.* The equality $k^\mu k^\nu \mathcal{Q}_{\mu\nu} = 0$ holds iff the transverse projections vanish for that null direction, $V^\alpha = W^\alpha = 0$.

In short, NEC and NCC live on opposite sides of the field equations; the Relational NEC strengthens the source–side inequality by adding a *provably* null–positive $\mathcal{Q}_{\mu\nu}$, so that—once (11.239) and the null–trace drop are used—Relational NEC implies Relational NCC exactly as formalized in Corollary 11.129.

Remark 11.131 (Operator–algebraic route to NCC)**.** Let $\hat{\tau}_{\mu\nu} = \delta\hat{t}_{\mu\nu}$ denote the tau–kairos fluctuation in the coherence channel (eq. (13.113) and Definition 13.1). For a null generator k^μ and a smooth compactly supported test 1–form φ_α on a null screen Σ_k, define the smeared flux operator

$$\hat{\mathcal{J}}_k[\varphi] := \int_{\Sigma_k} \mathrm{d}\Sigma\, \varphi_\alpha\, k^\mu\, \hat{\tau}_\mu{}^\alpha.$$

On the BRST/GNS physical space (§12.3) one has positivity $\langle(\hat{\mathcal{J}}_k[\varphi])^\dagger \hat{\mathcal{J}}_k[\varphi]\rangle \geq 0$. Unpacking this quadratic form and using the screen decomposition underlying (11.235) shows that the corresponding c-number density satisfies $k^\mu k^\nu \mathcal{Q}_{\mu\nu}[\bar{t}] \geq 0$ (equivalently, the transverse amplitudes $(\bar{t}\cdot k)^\perp$ have non-negative norm), hence Theorem 11.128. This complements the Newman–Penrose screen-space proof in Lemma 11.125 and is compatible with the TT isolation/SM orthogonality enforced by the coherence projectors (eq. (13.113) and Definition 13.1).

Operator–algebraic strengthening (GNS positivity). Let $\hat{t}_{\mu\nu}$ be the operator–valued temporal–curvature tensor (Definition 5.17), and define its centered fluctuation (*tau–kairos*) by $\hat{\tau}_{\mu\nu} := \delta\hat{t}_{\mu\nu} = \hat{t}_{\mu\nu} - \bar{t}_{\mu\nu}$ (Section 11.6). Fix a $\bar{g}^{\mathrm{Lor}}$–null hypersurface Σ_k with future–null normal k^μ (eqs. (11.232) and (11.233)), pick an auxiliary null ℓ^μ with $k\cdot\ell = -1$, and define the screen metric and projector

$$h_{\mu\nu} := \bar{g}^{\mathrm{Lor}}_{\mu\nu} + k_\mu\ell_\nu + \ell_\mu k_\nu, \qquad \Pi^\alpha{}_\beta := \delta^\alpha{}_\beta + k^\alpha\ell_\beta + \ell^\alpha k_\beta,$$

so that $h_{\mu\nu}$ is positive–definite on vectors orthogonal to $\{k, \ell\}$ and $\Pi^\alpha{}_\beta$ projects to the screen ($\Pi^\alpha{}_\beta k^\beta = \Pi^\alpha{}_\beta \ell^\beta = 0$). For any smooth, compactly supported 1–form φ_α on Σ_k satisfying $k^\alpha\varphi_\alpha = 0$, define the smeared null–coherence operator

$$\hat{\mathcal{J}}_k[\varphi] \;:=\; \int_{\Sigma_k} \mathrm{d}\Sigma(x)\, \varphi_\alpha(x)\, k^\mu(x) \left(\Pi^\alpha{}_\beta\, \hat{\tau}_\mu{}^\beta\right)(x), \tag{11.241}$$

and the positive screen norm

$$\|\varphi\|_k^2 \;:=\; \int_{\Sigma_k} \mathrm{d}\Sigma(x)\, \overline{\varphi_\alpha(x)}\, \varphi_\beta(x)\, h^{\alpha\beta}(x), \qquad h^{\alpha\beta}h_{\beta\gamma} = \Pi^\alpha{}_\gamma. \tag{11.242}$$

(When correlating insertions at x and y along Σ_k, screen indices are compared with the Levi–Civita parallel transporter of $\bar{g}^{\mathrm{Lor}}$, $\mathcal{U}_{\bar{g}^{\mathrm{Lor}}}(x \leftarrow y)$, built from Definition 6.7; cf. the bitensor use in Definition 11.109.)

On the certified plateau $\mathfrak{P}$ (BRST–closed, diagnostic realized reference configuration semantics; Proposition 7.3), GNS/Wightman positivity on the BRST/GNS physical sector implies

$$\left\langle \left(\hat{\mathcal{J}}_k[\varphi]\right)^\dagger \hat{\mathcal{J}}_k[\varphi] \right\rangle \;\geq\; 0, \tag{11.243}$$

for all transverse test 1–forms φ [381, 241, 255]. Expanding (11.243) yields a positive semi–definite bi–kernel on Σ_k,

$$0 \;\leq\; \int_{\Sigma_k} \mathrm{d}\Sigma(x) \int_{\Sigma_k} \mathrm{d}\Sigma(y)\, \overline{\varphi_\alpha(x)}\, \varphi_\beta(y)\, k^\mu(x) k^\nu(y)\, \Pi^\alpha{}_{\alpha'}(x) \left\langle \hat{\tau}_\mu{}^{\alpha'}(x)\, \mathcal{U}_{\alpha'}{}^{\beta'}(x \leftarrow y)\, \hat{\tau}_{\nu\,\beta'}(y) \right\rangle,$$

with $\mathcal{U}_{\alpha'}{}^{\beta'}(x \leftarrow y)$ the $\bar{g}^{\mathrm{Lor}}$ parallel transporter on the screen (Definition 6.7). In the local Hadamard point–split limit on Σ_k (Section 5.4.7), the projector isolates the transverse null–screen amplitudes. Using the trace subtraction in the coherence stress definition (eq. (11.235)), one obtains the operator–algebraic strengthening of Lemma 11.125:

$$\boxed{k^\mu k^\nu \, \mathcal{Q}_{\mu\nu}[\bar{t}] \;=\; \inf_{\varphi\neq 0} \frac{\big\langle (\hat{\mathcal{J}}_k[\varphi])^\dagger \hat{\mathcal{J}}_k[\varphi]\big\rangle}{\|\varphi\|_k^2} \;\geq\; 0,} \tag{11.244}$$

where $\|\cdot\|_k$ is given by (11.242). Thus null–positivity of $\mathcal{Q}_{\mu\nu}$ is elevated from the classical identity on the screen (Lemma 11.125) to a direct consequence of operator–algebraic positivity, compatible with the relational null cone (Lemma 11.123) and the NCC of Section 11.6 and eq. (11.231).

Remark 11.132 (Centering vs. background subtraction in $\hat{\mathcal{J}}_k$)**.** For the GNS/Wightman positivity argument—and thus for the Relational NEC—it is immaterial whether the smeared operator $\hat{\mathcal{J}}_k$ is built from the centered fluctuation $\hat{\tau}_{\mu\nu} := \hat{t}_{\mu\nu} - \bar{t}_{\mu\nu}$ or from the full operator $\hat{t}_{\mu\nu}$.

1) Centering. By definition, $\hat{\tau}_{\mu\nu} = \hat{t}_{\mu\nu} - \bar{t}_{\mu\nu}$ with $\langle\hat{\tau}_{\mu\nu}\rangle = 0$. Let Σ_k be a $\bar{g}^{\mathrm{Lor}}$–null screen with normal k^μ and auxiliary null ℓ^μ, and let $\Pi^\alpha{}_\beta$ be the screen projector (eqs. (11.232), (11.233) and (11.242)). The $\hat{t}$–smeared and centered smearings are

$$\hat{\mathcal{J}}_k^{(t)}[\varphi] := \int_{\Sigma_k} \mathrm{d}\Sigma \; \varphi_\alpha \, k^\mu \, (\Pi^\alpha{}_\beta \hat{t}_\mu{}^\beta), \qquad \hat{\mathcal{J}}_k^{(\tau)}[\varphi] := \int_{\Sigma_k} \mathrm{d}\Sigma \; \varphi_\alpha \, k^\mu \, (\Pi^\alpha{}_\beta \hat{\tau}_\mu{}^\beta), \tag{11.245}$$

which differ by the c–number

$$c[\varphi] := \int_{\Sigma_k} \mathrm{d}\Sigma \; \varphi_\alpha \, k^\mu \, (\Pi^\alpha{}_\beta \bar{t}_\mu{}^\beta). \tag{11.246}$$

(Parallel transport of screen indices between points on Σ_k is tacit via the Levi–Civita bitransporter of $\bar{g}^{\mathrm{Lor}}$, as in eq. (11.241).)

2) Positivity is invariant under c–number shifts. For any operator A and $c \in \mathbb{C}$, $\langle (A+c)^\dagger (A+c)\rangle = \langle A^\dagger A\rangle + |c|^2 + 2\,\mathrm{Re}\big(c^*\langle A\rangle\big) \geq 0$. Taking $A = \hat{\mathcal{J}}_k^{(\tau)}[\varphi]$ with $\langle A\rangle = 0$ gives

$$\Big\langle \big(\hat{\mathcal{J}}_k^{(t)}[\varphi]\big)^\dagger \hat{\mathcal{J}}_k^{(t)}[\varphi]\Big\rangle = \Big\langle \big(\hat{\mathcal{J}}_k^{(\tau)}[\varphi]\big)^\dagger \hat{\mathcal{J}}_k^{(\tau)}[\varphi]\Big\rangle + |c[\varphi]|^2 \;\geq\; 0,$$

so GNS positivity (eq. (11.243)) holds for either choice.

3) Variational formula and $\mathcal{Q}_{\mu\nu}$. The operator–to–classical bridge (eq. (11.244)) reads

$$k^\mu k^\nu \mathcal{Q}_{\mu\nu}[\bar{t}] \;=\; \inf_{\varphi\neq 0} \frac{\big\langle (\hat{\mathcal{J}}_k^{(\tau)}[\varphi])^\dagger \hat{\mathcal{J}}_k^{(\tau)}[\varphi]\big\rangle}{\|\varphi\|_k^2} \;\geq\; 0.$$

If one instead uses $\hat{t}$,

$$\inf_{\varphi\neq 0}\frac{\left\langle(\hat{\mathcal{J}}_k^{(t)}[\varphi])^\dagger\hat{\mathcal{J}}_k^{(t)}[\varphi]\right\rangle}{\|\varphi\|_k^2}=\inf_{\varphi\neq 0}\frac{\left\langle(\hat{\mathcal{J}}_k^{(\tau)}[\varphi])^\dagger\hat{\mathcal{J}}_k^{(\tau)}[\varphi]\right\rangle+|c[\varphi]|^2}{\|\varphi\|_k^2}\;\geq\;\inf_{\varphi\neq 0}\frac{\left\langle(\hat{\mathcal{J}}_k^{(\tau)}[\varphi])^\dagger\hat{\mathcal{J}}_k^{(\tau)}[\varphi]\right\rangle}{\|\varphi\|_k^2}.$$

Thus the lower bound needed for the NCC (Section 11.6) is unchanged or strengthened by using $\hat{t}$. If exact equality to $k^\mu k^\nu\mathcal{Q}_{\mu\nu}[\bar{t}]$ is desired with the non–centered smearing, replace $\langle\cdot\rangle$ by the connected correlator $\langle\cdot\rangle_c$, or use the centered $\hat{\mathcal{J}}_k^{(\tau)}$.

4) Physical takeaway. Using $\hat{\tau}$ centers the observable ($\langle\hat{\mathcal{J}}_k^{(\tau)}\rangle=0$) and lands the variational formula directly on $k^\mu k^\nu\mathcal{Q}_{\mu\nu}[\bar{t}]$ (eqs. (11.235) and (11.244) and Lemma 11.125). Using $\hat{t}$ is equally valid for positivity because c–number shifts cannot spoil $\langle A^\dagger A\rangle\geq 0$. Either choice is consistent with the tau–kairos (scalar–vector) split and TT isolation (eq. (13.113) and Definition 13.1) and leaves the NCC conclusion intact (Section 11.6).

Remark 11.133 (Null–coherence smearing and Chrono–Kymon content)**. (a) What $\hat{\mathcal{J}}_k[\varphi]$ probes.** In the coherence/NEC construction the smeared null–surface operator is built from $\hat{\tau}_{\mu\nu}:=\delta\hat{t}_{\mu\nu}$, projected transversely on Σ_k (transversality $k^\alpha\varphi_\alpha=0$):

$$\hat{\mathcal{J}}_k[\varphi]=\int_{\Sigma_k}\mathrm{d}\Sigma\,\varphi_\alpha\,k^\mu\,\hat{\tau}_\mu{}^\alpha,\qquad\left\langle(\hat{\mathcal{J}}_k[\varphi])^\dagger\hat{\mathcal{J}}_k[\varphi]\right\rangle\geq 0,\tag{11.247}$$

and the positivity (11.243) together with the Hadamard point-split limit yields the variational bridge (11.244).

(b) How the Chrono–Kymon enters. The tau–kairos splits into scalar and vector (Chrono–Kymon) pieces, and the antisymmetric temporal curvature corresponds to a kymonic field strength:

$$\hat{\tau}_{\mu\nu}=S_{\mu\nu}[\hat{\varphi}]\oplus A_{\mu\nu}[\hat{K}^{(\tau)}],\qquad\hat{t}_{[\mu\nu]}=\chi_\tau\,\hat{\mathcal{F}}^{(\tau)}_{\mu\nu},\qquad\hat{\mathcal{F}}^{(\tau)}_{\mu\nu}=2\,\bar{\nabla}_{[\mu}\hat{K}^{(\tau)}_{\nu]},\tag{11.248}$$

so $\hat{\mathcal{J}}_k$ probes *transverse, null–projected components of the field strength* (derivatives of $K^{(\tau)}$) on the screen space orthogonal to k^μ, not the homogeneous background norm $K^{(\tau)}\cdot K^{(\tau)}$ used in cosmological calibration. This is precisely why the contraction $k^\mu k^\nu\mathcal{Q}_{\mu\nu}[\bar{t}]$ becomes a sum of positive screen–space squares.

Remark 11.134 (Chrono–kymon vs. τ–kairos: rationale and formalism)**.** There is a precise reason to introduce the *chrono–kymon* $K^{(\tau)}_\mu$ instead of working only with the τ-kairos tensor $\hat{\tau}_{\mu\nu}:=\delta\hat{t}_{\mu\nu}$ (eqs. (11.248) and (13.113)).

1. **Clean split of degrees of freedom.** The τ-kairos decomposes as

$$\hat{\tau}_{\mu\nu}=S_{\mu\nu}\oplus A_{\mu\nu},\qquad A_{\mu\nu}=2\,\bar{\nabla}_{[\mu}K^{(\tau)}_{\nu]}\;\equiv\;\mathcal{F}^{(\tau)}_{\mu\nu},$$

so the antisymmetric piece is an *exact two–form* (a spin–1 mode), clearly separated from the scalar mode in $S_{\mu\nu}$ (eqs. (11.248) and (13.113)). This keeps the spin–2 (TT) graviton sector orthogonal and unmixed.

2. **Gauge/BRST cleanliness.** Writing $A = dK^{(\tau)}$ equips the coherence vector with a U(1)–like gauge symmetry $K^{(\tau)} \mapsto K^{(\tau)} + \bar{\nabla}\alpha$. This makes the coherence sector BRST–amenable and preserves commutation with the typed/TT projectors (Proposition 7.3 and Corollary 11.5). Working directly with $\tau_{\mu\nu}$ obscures these redundancies.
3. **Local, causal dynamics.** Potentials admit local Lagrangians and retarded/advanced Green operators supported on the *same* relational cone $J^{\pm}_{\Sigma}$ as in the rest of the formalism, ensuring microcausality alignment with Chapters 6–8 (cf. the kernels in Remark 11.108).
4. **Null–screen positivity becomes manifest (Relational NEC).** The coherence stress $\mathcal{Q}_{\mu\nu}$ (eq. (11.235)) satisfies, for any $\bar{g}^{\mathrm{Lor}}$–null k^μ,

$$k^\mu k^\nu \mathcal{Q}_{\mu\nu}[\bar{t}] = \|(S\cdot k)\|^2 + \|(\mathcal{F}^{(\tau)}\cdot k)\|^2 \ \geq 0,$$

   which is the positivity used in the Relational NEC (Lemma 11.125 and Section 11.6). Expressing the antisymmetric content as a *field strength* turns the null contraction into a sum of *screen–space squares*, exactly what the GNS/point–split argument requires (eqs. (11.243) and (11.244) and Remark 11.133).
5. **Observable, gauge–invariant probes on null screens.** The smeared operator $\hat{\mathcal{J}}_k[\varphi]$ samples transverse components of the *field strength* (not the potential) and is therefore gauge–invariant and positive by construction (eqs. (11.241) and (11.247) and Remark 11.132). This gives a direct operator–algebraic route to the NEC via eqs. (11.243) and (11.244).
6. **EP–safe scalarisation and PPN bookkeeping.** Encoding the antisymmetric sector via $K^{(\tau)}$ clarifies how it contributes to the anisotropic piece of the emergent coupling $\bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}$ and why, in the EP–safe (scalarised) regime, such effects are projected out or suppressed by small, screened gradients (Section 11.5.7 and Proposition 11.113).
7. **Global note (local arguments suffice).** If A has nontrivial cohomology, $K^{(\tau)}$ exists patchwise. Our NEC/focusing and GNS arguments are *local* (Hadamard/point–split), so a potential description is always available on the charts we use.

In sum. The composite tensor $\tau_{\mu\nu}$ is the right object to *define* the coherence stress, but introducing the chrono–kymon $K^{(\tau)}$ renders the antisymmetric sector gauge–clean, causal, and positivity–transparent, while preserving TT isolation and the scalar–vector bookkeeping consistently across Chapters 6–12 (eqs. (11.235), (11.243) and (11.244), Lemma 11.125, and Remarks 11.132 and 11.133).

State class, domains, and renormalization (technical set-up). We work throughout in the BRST/GNS physical representation (Sections 1.4.1 and 12.2) and assume Hadamard regularity for two-point correlators in the temporal–curvature sector, both for the full operator $\hat{t}_{\mu\nu}$ and its centered fluctuation (tau–kairos) $\hat{\tau}_{\mu\nu} := \delta\hat{t}_{\mu\nu}$ (Section 5.2.1). Let Σ_k be a $\bar{g}^{\mathrm{Lor}}$–null hypersurface with future-directed null normal k^μ and induced measure $\mathrm{d}\Sigma$. Choose an auxiliary null ℓ^μ with $k\cdot\ell = -1$ and define the screen projector and metric

$$h_{\mu\nu} := \bar{g}^{\mathrm{Lor}}_{\mu\nu} + k_\mu\ell_\nu + k_\nu\ell_\mu, \qquad \Pi^\alpha{}_\beta := \delta^\alpha{}_\beta + \ell^\alpha k_\beta \quad (\Rightarrow\ k^\beta\Pi^\alpha{}_\beta = 0).$$

Define the transverse test space

$$\mathcal{T}_k \;:=\; \Big\{\; \varphi_\alpha \in C_0^\infty(T^*\Sigma_k) \;:\; k^\alpha \varphi_\alpha = 0 \;\Big\}.$$

For $\varphi \in \mathcal{T}_k$, the (centered) smeared null–coherence operator is

$$\hat{\mathcal{J}}_k[\varphi] \;:=\; \int_{\Sigma_k} \mathrm{d}\Sigma(x)\; \varphi_\alpha(x)\, k^\mu(x)\, \hat{\tau}_\mu{}^\alpha(x), \tag{11.249}$$

which is densely defined on the GNS domain, with positive screen norm

$$\|\varphi\|_k^2 \;:=\; \int_{\Sigma_k} \mathrm{d}\Sigma(x)\; \overline{\varphi_\alpha(x)}\, \varphi_\beta(x)\, h^{\alpha\beta}(x), \qquad h^{\alpha\beta} h_{\beta\gamma} = \Pi^\alpha{}_\gamma. \tag{11.250}$$

(For bi-local expressions on Σ_k, screen indices at y are parallel-transported to x using the Levi–Civita transporter $\mathcal{U}_{\bar{g}^{\mathrm{Lor}}}(x \leftarrow y)$.)
By GNS/Wightman positivity in the physical sector [381, 241, 255],

$$\Big\langle \big(\hat{\mathcal{J}}_k[\varphi]\big)^\dagger \hat{\mathcal{J}}_k[\varphi] \Big\rangle \;\geq\; 0, \qquad \forall\, \varphi \in \mathcal{T}_k, \tag{11.251}$$

(cf. eq. (11.243)). If one prefers the uncentered smearing with $\hat{t}_{\mu\nu}$, set $\hat{\mathcal{J}}_k^{(t)}[\varphi] := \int_{\Sigma_k} \mathrm{d}\Sigma\, \varphi_\alpha k^\mu \hat{t}_\mu{}^\alpha$; positivity is unchanged and the variational bound differs only by a harmless c-number shift (Remark 11.132).

To define the quadratic coherence stress as an operator-valued distribution, use point-splitting with a Hadamard parametrix and subtract local, covariant counterterms that preserve BRST closure and $\bar{g}^{\mathrm{Lor}}$–covariance (the standard finite renormalization freedom). Denote the Wick-ordered quadratic tensor by

$$:\!\mathcal{Q}_{\mu\nu}\!:\,[\bar{t}] \;:=\; :\!S_{\mu\alpha} S_\nu{}^\alpha\!: \;+\; :\!A_{\mu\alpha} A_\nu{}^\alpha\!: \;-\; \tfrac{1}{4}\, \bar{g}^{\mathrm{Lor}}_{\mu\nu} \Big(:\!S_{\alpha\beta} S^{\alpha\beta}\!: + :\!A_{\alpha\beta} A^{\alpha\beta}\!: \Big), \tag{11.252}$$

where $S_{\mu\nu} := \hat{\tau}_{(\mu\nu)}$, $A_{\mu\nu} := \hat{\tau}_{[\mu\nu]}$, and all products are Wick-ordered by the state's Hadamard two-point function (Section 1.4.1; cf. the classical definition eq. (11.235)). Possible finite counterterms are built from $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ and its curvatures; terms $\propto \bar{g}^{\mathrm{Lor}}_{\mu\nu}$ vanish after contraction with $k^\mu k^\nu$ by Lemma 11.123, and total-derivative terms can be arranged to vanish upon smearing on Σ_k (compact support). The tau–kairos TT isolation $\mathcal{P}^{(2)}[\hat{\tau}_{\mu\nu}] = 0$ (see the typed/triadic structure in Chap. 13 and Remark 11.133) guarantees no spin–2 leakage into $:\!\mathcal{Q}_{\mu\nu}\!:$, preserving the null-cone analysis and the Relational NEC (Section 11.6).

Proposition 11.135 (Null positivity of the Wick-ordered coherence stress)**.** *Let k^μ be a $\bar{g}^{\mathrm{Lor}}$-null vector field and let $f \in C_0^\infty(\Sigma_k)$ with $f \geq 0$. Then, in any BRST/GNS physical Hadamard state,*

$$\Big\langle\; :\!\mathcal{Q}_{\mu\nu}\!:\,[\bar{t}]\; k^\mu k^\nu \,,\; f \;\Big\rangle \;\geq\; 0, \tag{11.253}$$

i.e. the distribution $k^\mu k^\nu\; :\!\mathcal{Q}_{\mu\nu}\!:$ is positive on non-negative test densities supported on Σ_k.

Proof sketch consistent with eq. (11.251). Choose a family $\{\varphi^{(\epsilon)}\}_{\epsilon>0} \subset \mathcal{T}_k$ with $\varphi^{(\epsilon)} \to \sqrt{f}\, m_\alpha$ in C_0^∞ as $\epsilon \downarrow 0$, where m_α is any unit complex transverse covector on Σ_k (and similarly $\bar{m}_\alpha$ in a Newman–Penrose frame). By eq. (11.251),

$$\Big\langle \big(\hat{\mathcal{J}}_k[\varphi^{(\epsilon)}]\big)^\dagger \hat{\mathcal{J}}_k[\varphi^{(\epsilon)}] \Big\rangle \;\geq\; 0.$$

Perform point-splitting with a Hadamard parametrix, parallel-transport screen indices, and subtract the state-dependent singularity; in the limit $\epsilon \downarrow 0$ one obtains the Wick-squares of the transverse null contractions of $\hat{\tau}_{\mu\nu}$, i.e. the positive pieces in eq. (11.252) smeared by f. The trace subtraction in eq. (11.252) contributes only multiples of $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$, which vanish under $k^\mu k^\nu$ by Lemma 11.123. Local, covariant finite counterterms (BRST-closed, $\bar{g}^{\mathrm{Lor}}$-covariant) either drop out on nulls or reduce to total divergences that vanish upon smearing on Σ_k (compact support). Hence $\langle :\mathcal{Q}_{\mu\nu}: \, k^\mu k^\nu, f\rangle \geq 0$. □

Corollary 11.136 (Operator-level strengthening of Lemma 11.125)**.** *With* $:\mathcal{Q}_{\mu\nu}:$ *defined by eq.* (11.252)*, the null inequality* $\langle :\mathcal{Q}_{\mu\nu}: \, k^\mu k^\nu, f\rangle \geq 0$ *holds for all non-negative* $f \in C_0^\infty(\Sigma_k)$*. In particular, for smooth, compactly supported congruence bundles,*

$$k^\mu k^\nu \, \mathcal{Q}_{\mu\nu}[\bar{t}] \;\geq\; 0$$

in the sense of quadratic forms, thereby upgrading Lemma 11.125 from a tetrad identity to an operator–algebraic statement consistent with eq. (11.251)*.*

The operator–algebraic positivity statement $\left\langle (\hat{\mathcal{J}}_k[\varphi])^\dagger \hat{\mathcal{J}}_k[\varphi] \right\rangle \geq 0$ on null screens (eq. (11.251)) promotes the classical null "sum of squares" identity for the coherence sector to a fully renormalized field-theoretic result. After Hadamard point-splitting and Wick ordering, this yields the positivity of the distribution $k^\mu k^\nu \, :\mathcal{Q}_{\mu\nu}:$ on every non-negative test density f supported on the null hypersurface Σ_k (Prop. Proposition 11.135, using the definition in eq. (11.252)). In other words, the coherence contribution cannot furnish negative energy along a light ray in any BRST/GNS physical Hadamard state.

The structural reason is the scalar–vector split of the tau–kairos tensor $\hat{\tau}_{\mu\nu}$ into a symmetric piece $S_{\mu\nu}$ and an antisymmetric piece $A_{\mu\nu}$, the latter expressed as the chrono–kymon field strength $\mathcal{F}^{(\tau)}_{\mu\nu} = 2\,\bar{\nabla}_{[\mu} K^{(\tau)}_{\nu]}$ (Remark 11.133 and eq. (11.248)). With the trace-adjusted definition of the coherence stress $\mathcal{Q}_{\mu\nu}$ (eq. (11.235)), the null contraction becomes a sum of positive screen–space squares, $k^\mu k^\nu \mathcal{Q}_{\mu\nu} = \|(S{\cdot}k)\|^2 + \|(\mathcal{F}^{(\tau)}{\cdot}k)\|^2 \geq 0$ (Lemma Lemma 11.125). The smeared operator $\hat{\mathcal{J}}_k[\varphi] = \int_{\Sigma_k} \mathrm{d}\Sigma \, \varphi_\alpha \, k^\mu \, \hat{\tau}_\mu{}^\alpha$ (eq. (11.249)), built with transverse test one-forms and the screen metric, probes precisely these gauge-invariant, transverse components and delivers the variational bridge from operator positivity to the classical inequality (eqs. (11.244) and (11.251) and Remarks 11.132 and 11.133).

Gravitationally, combining this null positivity with the vacuum drop-out $k^\mu k^\nu T^{\mathrm{vac}}_{\mu\nu} = 0$ (Lemma Lemma 11.124) and the ordinary NEC for matter on $\bar{g}^{\mathrm{Lor}}$ establishes the Relational NEC of Section 11.6 and its operator-level strengthening (Cor. Corollary 11.136). Contracting the field equations with $k^\mu k^\nu$ then feeds into the Raychaudhuri inequality on $(\bar{g}^{\mathrm{Lor}}, \bar{\nabla})$ (eq. (11.234)), ensuring standard null focusing whenever the scalarized source $k^\mu k^\nu \left(T^{\mathrm{matter}}_{\mu\nu} + \gamma \, \mathcal{Q}_{\mu\nu}\right)$ is non-negative. In the EP-safe (scalarized) regime (Section 11.5.7 and Proposition 11.113), this statement is preserved with a slowly varying κ_{eff}, maintaining consistency with the Chronogeometric Einstein law (Definition 11.106).

Focusing and Observational Handles. Let $\bar{\nabla}$ be the Levi–Civita connection of $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ (Definition 6.7). For an affinely parameterized null congruence with tangent k^μ ($k^\nu \bar{\nabla}_\nu k^\mu = 0$), the

Raychaudhuri equation takes the standard form [240, 7]:

$$\frac{\mathrm{d}\theta}{\mathrm{d}\lambda} = -\tfrac{1}{2}\,\theta^2 - \sigma_{\mu\nu}\sigma^{\mu\nu} + \omega_{\mu\nu}\omega^{\mu\nu} - R_{\mu\nu}[\bar{g}^{\mathrm{Lor}}]\,k^\mu k^\nu, \tag{11.254}$$

where $\theta, \sigma_{\mu\nu}, \omega_{\mu\nu}$ are computed with $\bar{g}^{\mathrm{Lor}}$. Using the scalarised field equations of 11.5.7 one has

$$R_{\mu\nu}[\bar{g}^{\mathrm{Lor}}]\,k^\mu k^\nu = \kappa_{\mathrm{eff}}(x)\,k^\mu k^\nu\Big(T^{\mathrm{matter}}_{\mu\nu} + \gamma\,\mathcal{Q}_{\mu\nu}[\bar{t}]\Big),$$

and by Lemma 11.123 the vacuum term vanishes on nulls. For hypersurface–orthogonal congruences ($\omega_{\mu\nu} = 0$) and under the NCC (11.231) (with $\kappa_{\mathrm{eff}} \geq 0$),

$$\frac{\mathrm{d}\theta}{\mathrm{d}\lambda} \;\leq\; -\tfrac{1}{2}\,\theta^2 \qquad \Rightarrow \qquad \theta(\lambda) \;\to\; -\infty \;\text{ by }\; \lambda \leq \frac{2}{|\theta(0)|} \;\text{ if } \theta(0) < 0,$$

i.e. focusing within a finite affine parameter, exactly as in GR, now with explicit sensitivity to the coherence stress through $\mathcal{Q}_{\mu\nu}[\bar{t}]$. This furnishes *observational handles*: null–congruence phenomena such as strong/weak lensing, Shapiro delay, echo/tail structures in waveforms, and standard–ruler distortions acquire coherence–dependent corrections controlled by $\gamma\,\mathcal{Q}_{\mu\nu}$ (cf. precision weak-field tests [479]).

Remark 11.137 (Optical scalars and their role in (11.254))**.** Fix an affinely–parameterized null congruence with tangent k^μ ($k^\nu\bar{\nabla}_\nu k^\mu = 0$) for the Levi–Civita connection $\bar{\nabla}$ of $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ (cf. Definition 6.7). Choose a null partner ℓ^μ with $k\cdot\ell = -1$ and define the screen metric $q_{\mu\nu} = \bar{g}^{\mathrm{Lor}}_{\mu\nu} + k_\mu\ell_\nu + \ell_\mu k_\nu$ (cf. Remark 11.127). The *optical tensor* is the projected gradient

$$B_{\mu\nu} \;:=\; q_\mu{}^\alpha\, q_\nu{}^\beta\, \bar{\nabla}_\alpha k_\beta, \qquad q_\mu{}^\alpha := \bar{g}^{\alpha\rho}q_{\rho\mu},$$

and decomposes into its trace, symmetric tracefree, and antisymmetric parts:

$$B_{\mu\nu} \;=\; \frac{1}{d-2}\,\theta\, q_{\mu\nu} \;+\; \sigma_{\mu\nu} \;+\; \omega_{\mu\nu}.$$

Here

- **Expansion** $\theta := B^\mu{}_\mu = q^{\mu\nu}\bar{\nabla}_\mu k_\nu$ measures the fractional rate of change of the beam's cross–sectional area (negative θ signals focusing).
- **Shear** $\sigma_{\mu\nu} := B_{(\mu\nu)} - \dfrac{\theta}{d-2}\,q_{\mu\nu}$ is symmetric and tracefree ($q^{\mu\nu}\sigma_{\mu\nu} = 0$), capturing shape distortions at fixed area.
- **Twist (vorticity)** $\omega_{\mu\nu} := B_{[\mu\nu]}$ measures the local rotation of the congruence; hypersurface–orthogonality (Frobenius) is equivalent to $\omega_{\mu\nu} = 0$.

With these definitions, the null Raychaudhuri equation (11.254) reads $\frac{d\theta}{d\lambda} = -\frac{1}{2}\theta^2 - \sigma_{\mu\nu}\sigma^{\mu\nu} + \omega_{\mu\nu}\omega^{\mu\nu} - R_{\mu\nu}k^\mu k^\nu$, where $\sigma_{\mu\nu}\sigma^{\mu\nu} \geq 0$ and $\omega_{\mu\nu}\omega^{\mu\nu} \geq 0$ are computed using $\bar{g}^{\mathrm{Lor}}$. In particular, for twist–free congruences ($\omega_{\mu\nu} = 0$) and under the (Relational) NCC ($R_{\mu\nu}k^\mu k^\nu \geq 0$), negative initial θ necessarily evolves toward focusing, providing the observational handle discussed around (11.254) [240, 7, 316].

Normalization, Limits, and Saturation. (i) *Normalization.* Any $\gamma \geq 0$ preserves the NCC (11.231); choosing $\gamma = 1$ matches the quadratic normalization from the temporal–sector action (see Section 6.1.13). (ii) *Decohered (GR) limit.* As $\bar{t}_{\mu\nu} \to 0$, one has $\mathcal{Q}_{\mu\nu}[\bar{t}] \to 0$ and $\bar{g}^{\mathrm{Lor}}_{\mu\nu} \to \bar{g}_{\mu\nu}$; thus (11.231) reduces to the standard NEC for $T^{\mathrm{matter}}_{\mu\nu}$ and (11.254) becomes the classical Raychaudhuri equation [7]. (iii) *Saturation.* Equality in (11.231) occurs iff the transverse coherence amplitudes vanish along the congruence, $\Xi_m := \bar{t}_{\mu\nu} k^\mu m^\nu = 0 = \Xi_{\bar{m}} := \bar{t}_{\mu\nu} k^\mu \bar{m}^\nu$ for some Newman–Penrose tetrad $\{k, \ell, m, \bar{m}\}$ (cf. Lemma 11.125); then only longitudinal (transport/pure–gauge) coherence survives and the condition reduces to the matter NEC alone.

Conclusion. Assuming matter obeys the NEC and $\gamma \geq 0$, the Relational Null Coherence Condition (11.231) is secured by operator–algebraic positivity and yields standard focusing theorems for $\bar{g}^{\mathrm{Lor}}$ with coherence–sensitive corrections, placing the Chronogeometric Einstein Equations (Definition 11.106) on the same energy–condition footing as in GR while opening observational avenues to probe $\bar{t}$ via null–congruence data.

11.7 Operator–Valued Einstein Equations with Unified Tensor

Instrumentation preamble (diagnostic realized reference configuration & tags). All statements in this section are made on a certified plateau $\mathfrak{P}$ with BRST–closed band tags (P_x, K_{IR}) commuting with the triad projectors $\{\Pi^{(\mathrm{corr})}, \Pi^{(\mathrm{vac})}, \Pi^{(\mathrm{spec})}\}$ and with the TT projector $\Pi^{[\mathrm{TT}]}{}_*\bar{\Sigma}$; observables descend to the physical space $\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}})$ with Σ–Nielsen flatness. Connections, curvatures and causal cones are those of $\bar{\Sigma}$ (Relational Levi–Civita); $\bar{\nabla}^\mu \bar{G}_{\mu\nu} \equiv 0$ and $\bar{\nabla}^\mu \bar{T}^{\mathrm{total}}_{\mu\nu} = 0$ are read on the same background (cf. Remark 11.108).

The semi–classical Chronogeometric Einstein law (Definition 11.106) reads

$$\boxed{\bar{G}_{\mu\nu}[\bar{\Sigma}] + \Lambda\, \bar{\Sigma}_{\mu\nu} = \kappa\, \bar{T}^{\mathrm{total}}_{\mu\nu}, \qquad \text{equivalently} \quad \bar{G}_{\mu\nu} = \bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}\, \bar{T}^{\mathrm{total}}_{\rho\sigma},} \tag{11.255}$$

with the emergent coupling tensor $\bar{\mathcal{C}}$ defined in Definition 11.111 and scalarisation limits in Section 11.5.7. Since both geometry and matter are operator–valued in IQR, the fundamental law must be formulated at the operator level and reduce to (11.255) upon taking expectations in the Holo–State.

11.7.1 Formulation of the Operator Equation

Let the unified operator tensor (Chs. 6–7) be

$$\hat{\Sigma}_{\mu\nu}(x) := \hat{g}_{\mu\nu}(x) + \hat{t}_{\mu\nu}(x),$$

where $\hat{g}_{\mu\nu}$ arises from the distinguishability functional of the Gaiamiti sector and $\hat{t}_{\mu\nu}$ from the temporal–curvature functional (the coherence/chrono–temporal sector), cf. Definitions 5.1 and 5.17. The operator–valued curvature tensors $\hat{R}^\lambda{}_{\rho\mu\nu}[\hat{\Sigma}]$, $\hat{R}_{\mu\nu}[\hat{\Sigma}]$ and $\hat{R}[\hat{\Sigma}]$ are defined as operator–valued distributions by Hadamard point–splitting on $\mathfrak{P}$ and functional composition with the (quantum) Levi–Civita connection of $\hat{\Sigma}$ (normal–ordered where needed). The operator Einstein tensor is

$$\hat{G}_{\mu\nu}[\hat{\Sigma}] := \hat{R}_{\mu\nu}[\hat{\Sigma}] - \tfrac{1}{2}\, \hat{\Sigma}_{\mu\nu}\, \hat{R}[\hat{\Sigma}].$$

Causal response form (diagnostic realized reference configuration-covariant). Following Definition 11.109 and eqs. (11.213) and (11.214), the operator equation takes the retarded, diagnostic realized reference configuration-covariant bitensor form:

$$\boxed{\hat{G}_{\mu\nu}(x) = \int_{\mathcal{X}_q} d\mu_{\bar{\Sigma}}(x')\, \hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x') \circ \Big[\mathcal{U}_{\bar{\Sigma}}(x \leftarrow x')_{\rho}{}^{\rho'}\, \mathcal{U}_{\bar{\Sigma}}(x \leftarrow x')_{\sigma}{}^{\sigma'}\, \hat{T}^{\text{total}}_{\rho'\sigma'}(x')\Big],} \tag{11.256}$$

where $d\mu_{\bar{\Sigma}}(x') := \sqrt{-\det \bar{\Sigma}(x')}\, d^d x'$, $\mathcal{U}_{\bar{\Sigma}}$ is the Levi–Civita parallel transporter of $\bar{\Sigma}$, and $A \circ B := \frac{1}{2}(AB+BA)$ denotes the Jordan product (self-adjointness). The retarded kernel

$$\hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x') = \mathbb{P}^{(\text{Ein})\alpha\beta}_{\mu\nu}\, \hat{\Delta}^{\text{ret}\,\gamma\delta}_{\alpha\beta}(x,x')\, \mathbb{J}_{\gamma\delta}{}^{\rho\sigma}$$

has support in $J^{+}_{\bar{\Sigma}}(x')$ (microcausality) and encodes the TT-typed graviton response in the radiative sector. In the local/spectral limit (Section 11.5.8 and Proposition 11.115), $\hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x') \to \kappa\, \Pi_{\mu\nu}{}^{\rho\sigma}\, \delta^{(d)}(x-x')\, \hat{\mathbf{1}}$, recovering the familiar local law.

Local symbolic form (including interaction/coherence terms). Suppressing parallel transport and density factors, the retarded bitensor law (11.256) can be written symbolically as

$$\boxed{\hat{G}_{\mu\nu}[\hat{\Sigma}](x) \;=\; \hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x) \;\circ\; \hat{T}^{\text{total}}_{\rho\sigma}(x), \qquad A \circ B := \tfrac{1}{2}(AB+BA),} \tag{11.257}$$

where the *total* source includes the diagonal, interaction (mixing), and coherence/torsion pieces:

$$\boxed{\hat{T}^{\text{total}}_{\rho\sigma} \;:=\; \underbrace{\hat{T}^{(\Phi)}_{\rho\sigma} + \hat{T}^{(\Theta)}_{\rho\sigma}}_{\text{diagonal blocks}} \;+\; \underbrace{\hat{T}^{(\Phi\Theta)}_{\rho\sigma}\Big[\hat{\mathbf{G}}_{\Phi\Theta}\Big]}_{\text{off-diagonal interaction}} \;+\; \underbrace{\hat{T}^{(\tau)}_{\rho\sigma}\Big[\hat{K}^{(\tau)}, \hat{\mathcal{F}}^{(\tau)}\Big]}_{\text{coherence/torsion}}.} \tag{11.258}$$

Here $\hat{T}^{(\Phi\Theta)}$ encodes the mixing induced by the off–diagonal correlator blocks of the unified two–point kernel (cf. Section 6.1.1 and Lemma 11.146), and $\hat{T}^{(\tau)}$ collects the Chrono–Kymon contributions mapped into $\hat{t}_{[\mu\nu]} = \chi_\tau\, \hat{\mathcal{F}}^{(\tau)}_{\mu\nu}$ (cf. Sections 10.7 and 10.17.9). This is the operator–level analogue of the expectation–level decomposition in (11.264).

Vacuum placement. The cosmological (vacuum) channel may be kept on the geometric side, yielding the expectation–level form (11.269), or moved to the source as $-\frac{\Lambda_{\text{eff}}}{\kappa'}\, \hat{\Sigma}_{\rho\sigma}$. Both conventions are equivalent at the level of null contractions by Lemma 11.123. (For the precise diagnostic realized reference configuration–covariant bitensor structure, see (11.256).)

11.7.2 Consistency, Conservation, and Semiclassical Reduction

(i) Bianchi/cohomological consistency. On the diagnostic realized reference configuration background $(\bar{\Sigma}, \bar{\nabla})$, $\bar{\nabla}^{\mu}_{x} \mathbb{P}^{(\text{Ein})\alpha\beta}_{\mu\nu} = 0$ and the causal graviton kernel $\hat{\Delta}^{\text{ret}}$ is built from the $\bar{\Sigma}$–Lichnerowicz operator (Remark 11.108); together with Levi–Civita transport this yields

$$\bar{\nabla}^{\mu}_{x}\, \hat{G}_{\mu\nu}(x) = 0 \quad \Rightarrow \quad \bar{\nabla}^{\mu} \hat{T}^{\text{total}}_{\mu\nu}(x) = 0 \quad \text{(weakly on } \mathcal{H}_{\text{phys}}\text{, mod. BRST–exact).}$$

Equivalently, the cohomological EOMs vanish on $\mathfrak{P}$ "mod ζ–exact" as in Section 11.5.4.

(ii) State dependence and emergence. Taking the Holo–State expectation of (11.256) reproduces (11.255) with $\bar{\mathcal{C}} := \langle \hat{\mathcal{C}} \rangle$ (Definition 11.111); scalarisation and variation bounds are given in Section 11.5.7, and the GR limit in Section 11.5.8.

(iii) Ordering and microcausality. Non–commutativity is controlled by the Jordan product $\circ$ and by causal support of $\hat{\mathcal{C}}(\cdot;\cdot)$ in $J^{\pm}_{\bar{\Sigma}}$, guaranteeing that $\hat{G}_{\mu\nu}(x)$ depends only on sources in the past cone of x (the same microcausal cone used for scalar and spin–2 kernels in Chs. 6–8; cf. Remark 11.108).

(iv) Typed radiative sector. In the spin–2 sector, the TT–typed TLO and projector $\Pi^{[\mathrm{TT}]}{}_{*\bar{\Sigma}}$ (Ch. 11) act on (11.256), isolating physical graviton polarizations; non–TT observables are triad–mixed as in the polarization discussion of Ch. 11.

Weinberg–Witten applicability

The Weinberg–Witten (WW) no–go theorem [475] states that, *if* a quantum field theory (i) is formulated on a fixed Minkowski background with exact global Poincaré symmetry, and (ii) possesses a single, Lorentz–covariant, gauge–invariant, conserved local stress tensor $T^{\mu\nu}$, then it cannot contain interacting massless particles of spin $s > 1$. In the Chronogeometric/IQR framework, the hypotheses are not satisfied, and the theorem is therefore inapplicable:

1. **No fixed Minkowski background.** The dynamical "diagnostic realized reference configuration'' geometry is *emergent and state–dependent*: $\bar{\Sigma}_{\mu\nu}(x) = \langle \overset{\circ}{\boldsymbol{\Psi}} | \, \hat{\Sigma}_{\mu\nu}(x) \, | \overset{\circ}{\boldsymbol{\Psi}} \rangle$, so there is no global Poincaré frame or global Killing algebra supporting the WW assumptions. Causal structure, connections, and curvatures are those of $(\bar{\Sigma}, \bar{\nabla})$ (Relational Levi–Civita), with all readouts taken on the BRST/GNS physical space (diagnostic realized reference configuration instrumentation preamble; cf. Remark 11.108).

2. **No single local, gauge–invariant conserved $T^{\mu\nu}$.** The operative stress tensor is defined *relationally* with indices raised/lowered by $\bar{\Sigma}^{\mu\nu}$, and includes both matter and geometric contributions ($\hat{T}^{\mathrm{total}}_{\mu\nu} = \hat{T}^{(\Phi)}_{\mu\nu} + \hat{T}^{(\Theta)}_{\mu\nu}$; see Definition 9.358). The geometry–matter coupling is mediated by a *retarded bitensor kernel* with causal support, not by a single ultralocal coupling to a background–independent $T^{\mu\nu}$:

$$\hat{G}_{\mu\nu}(x) = \int d\mu_{\bar{\Sigma}}(x') \, \hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x') \;\circ\; \Big[\mathcal{U}_{\bar{\Sigma}}(x \leftarrow x')_{\rho}{}^{\rho'} \mathcal{U}_{\bar{\Sigma}}(x \leftarrow x')_{\sigma}{}^{\sigma'} \hat{T}^{\mathrm{total}}_{\rho'\sigma'}(x') \Big], \tag{11.259}$$

   (cf. eq. (11.256)). This relational kernel $\hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x')$ is state–dependent (Definition 11.111) and preserves microcausality on $J^{\pm}_{\bar{\Sigma}}$; such nonlocal bitensor couplings lie outside the WW hypothesis of a single local, gauge–invariant conserved tensor current on flat space.

3. **Gauge/BRST structure and physical polarizations.** Diffeomorphisms act as gauge symmetries; the physical graviton sector is the TT–typed cohomology on the diagnostic realized reference configuration, projected by $\Pi^{[\mathrm{TT}]}{}_{*\bar{\Sigma}}$ (Ch. 11; cf. Remark 11.108). Non–TT observables are triad–mixed and do not represent additional propagating

spin–2 modes. The operator identity (11.259) is BRST–consistent; its expectation reproduces the semi–classical equation with emergent coupling $\bar{\mathcal{C}} = \langle \hat{\mathcal{C}} \rangle$ (eq. (11.255) and Definition 11.111).

Spectral/GR limit and consistency. In the weak–field, low–coherence spectral channel one has $\hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x') \to \kappa\, \Pi_{\mu\nu}{}^{\rho\sigma}\, \delta^{(d)}(x-x')\, \hat{\mathbf{1}}$, so $\bar{G}_{\mu\nu} = \kappa\, \bar{T}^{\text{total}}_{\mu\nu}$ is recovered (Section 11.5.8 and Proposition 11.115). This agreement with low–energy GR and soft–theorem/EFT expectations [539, 403, 519] does not re–impose the WW hypotheses: the flat, globally Poincaré–invariant background and the existence of a single, background–independent, gauge–invariant local $T^{\mu\nu}$ are absent at the fundamental level. Hence the Chronogeometric TT graviton *evades* the WW obstruction in a manner consistent with (11.256) and with its GR limit.

11.7.3 The Einstein Tensor of the Unified Geometry

Instrumentation preamble (diagnostic realized reference configuration & tags). All statements in this subsection are made on a certified plateau $\mathfrak{P}$ with BRST–closed band tags (P_x, K_{IR}) commuting with the triad projectors and the TT projector, and all readouts descend to $\mathcal{H}_{\text{phys}} := H^0(\hat{Q}_{\text{BRST}})$ (Nielsen–flat law channel), cf. the preamble to §11.5.4. *Relational geometry.* Connections and curvatures are those of $\bar{\Sigma}$ (Relational Levi–Civita); index gymnastics use $\bar{\Sigma}_{\mu\nu}$ and $\bar{\Sigma}^{\mu\nu}$, and $\bar{\nabla}$ is the Levi–Civita connection of $\bar{\Sigma}$ (see Definition 11.103 and Remark 11.108).

Definition 11.138 (Einstein tensor of the unified geometry)**.** Given the relational Ricci tensor and scalar curvature of $\bar{\Sigma}$ (Definition 11.103), the *Einstein tensor* on $(\mathcal{X}_q, \bar{\Sigma})$ is

$$\boxed{\bar{G}_{\mu\nu}[\bar{\Sigma}] := \; \bar{R}_{\mu\nu}[\bar{\Sigma}] - \tfrac{1}{2}\, \bar{\Sigma}_{\mu\nu}\, \bar{R}[\bar{\Sigma}].} \tag{11.260}$$

Conservation via the contracted Bianchi identity. Because $(\bar{\Sigma}, \bar{\nabla})$ is torsion–free and metric–compatible, the curvature obeys the (second) Bianchi identity; contracting yields conservation of the Einstein tensor.

Proposition 11.139 (Relational Bianchi & conservation)**.** *With $\bar{\nabla}$ the Levi–Civita connection of $\bar{\Sigma}$,*

$$\boxed{\bar{\nabla}^{\mu}\, \bar{G}_{\mu\nu}[\bar{\Sigma}] \;\equiv\; 0,} \tag{11.261}$$

as a direct consequence of the Bianchi identities in Proposition 11.98.

This divergence–free structure fixes the geometric side of the law and matches the diagnostic realized reference configuration–level conservation of the total expectation stress tensor, $\bar{\nabla}_{\mu} \bar{T}^{\mu\nu}_{\text{total}} = 0$, used in the Chronogeometric Einstein equations (Definition 11.106 and Section 11.5.5).

Interpretation and placement in the dynamical stack. The tensor $\bar{G}_{\mu\nu}[\bar{\Sigma}]$ captures the covariantly conserved, second–order content of the unified chronogeometry and therefore supplies the correct left–hand side of the field equations at both the semi–classical and operator levels: it appears in the diagnostic realized reference configuration law $\bar{G}_{\mu\nu} + \Lambda\,\bar{\Sigma}_{\mu\nu} = \kappa\,\bar{T}^{\mathrm{total}}_{\mu\nu}$ (Definition 11.106 and Section 11.5.4), and in the operator–valued causal response relation (eq. (11.256)). Linearization about $\bar{\Sigma}$ produces the Lichnerowicz operator and TT–projected propagator (Remark 11.108), ensuring that the radiative (spin–2) sector is handled by the typed TT panel of Ch. 11 while non–TT observables are triad–mixed but remain orthogonal to TT as in Remark 11.108. The same causal cones $J^{\pm}_{\bar{\Sigma}}$ control the support of Green kernels across the scalar and spin–2 sectors, maintaining microcausal alignment with Chs. 6–8.

Diagnostic realized reference configuration visibility (optional). For diagnostics on $\mathfrak{P}$, deviations from the Einstein law can be summarized by the diagnostic realized reference configuration residual and visibility defined in §11.5.4, which are evaluated with the same tags and projectors that certify the plateau; $\bar{G}_{\mu\nu}[\bar{\Sigma}]$ supplies the geometric input to those meters.

Relational connection and compatibility. Throughout this subsection the dynamical background is the unified tensor $\bar{\Sigma}_{\mu\nu}$ and $\nabla^{(\mathrm{rel})}$ denotes a *relational* affine connection chosen to be *metric–compatible* with $\bar{\Sigma}$ but allowed to carry torsion:

$$\nabla^{(\mathrm{rel})}_{\lambda}\bar{\Sigma}_{\mu\nu} = 0, \qquad \mathcal{T}^{\rho}{}_{\mu\nu} := 2\,\Gamma^{\rho}{}_{[\mu\nu]} \neq 0.$$

The antisymmetric piece of the temporal sector is modeled constitutively by the chrono–kymon field strength (Remark 11.133 and eq. (11.248)):

$$\bar{t}_{[\mu\nu]} \;=\; \chi_{\tau}\,\bar{\mathcal{F}}^{(\tau)}_{\mu\nu}, \qquad \bar{\mathcal{F}}^{(\tau)}_{\mu\nu} := 2\,\bar{\nabla}_{[\mu}\bar{K}^{(\tau)}_{\nu]},$$

where χ_{τ} is a constitutive coefficient and $\bar{\nabla}$ is the Levi–Civita connection of $\bar{\Sigma}$. Nonmetricity is set to zero (any nonmetric effects are absorbed in the constitutive definition of $\Gamma[\bar{\Sigma}]$), so that $\nabla^{(\mathrm{rel})}$ reduces to $\bar{\nabla}$ when $\mathcal{T} = 0$.

Proposition 11.140 (Relational Bianchi identity)**.** *Let $R^{\rho}{}_{\sigma\mu\nu}[\Gamma]$ be the curvature of the torsionful, metric–compatible $\Gamma = \Gamma[\bar{\Sigma}]$ with torsion $\mathcal{T}^{\rho}{}_{\mu\nu}$. Then the differential Bianchi identity takes the standard torsionful form*

$$\nabla^{(\mathrm{rel})}_{[\lambda} R^{\rho}{}_{|\sigma|\mu\nu]} \;-\; \mathcal{T}^{\eta}{}_{[\lambda\mu}\,R^{\rho}{}_{|\sigma|\nu]\eta} \;\equiv\; 0, \tag{11.262}$$

and the contracted identity implies a generalized divergence relation for an Einstein–type tensor $\mathbb{G}^{\mu}{}_{\nu}[\bar{\Sigma}]$ built from $R^{\rho}{}_{\sigma\mu\nu}[\Gamma]$:

$$\nabla^{(\mathrm{rel})}_{\mu}\,\mathbb{G}^{\mu}{}_{\nu}[\bar{\Sigma}] \;+\; \mathcal{C}_{\nu}[\mathcal{T}] \;\equiv\; 0, \tag{11.263}$$

where $\mathcal{C}_{\nu}[\mathcal{T}]$ is a covariant functional of torsion and curvature that vanishes identically when $\mathcal{T} = 0$. In the torsionless limit, $\mathbb{G}_{\mu\nu} \to \bar{G}_{\mu\nu}[\bar{\Sigma}]$ and (11.263) reduces to $\bar{\nabla}^{\mu}\bar{G}_{\mu\nu} = 0$ (cf. Proposition 11.139).

Proof sketch. Equation (11.262) is the first Bianchi identity for metric–compatible connections with torsion. Contracting on (ρ, λ) and rearranging produces (11.263). When $\mathcal{T} = 0$, $\nabla^{(\mathrm{rel})} \to \bar{\nabla}$ and $\mathcal{C}_\nu[\mathcal{T}] \to 0$, yielding the contracted Bianchi identity for $\bar{G}_{\mu\nu}[\bar{\Sigma}]$. □

Proposition 11.141 (Covariant conservation of the effective stress)**.** *Define the expectation–level field equations on* $(\mathcal{X}_q, \bar{\Sigma})$ *by*

$$\mathbb{G}_{\mu\nu}[\bar{\Sigma}] \;=\; \kappa\, T^{\mathrm{eff}}_{\mu\nu}, \qquad T^{\mathrm{eff}}_{\mu\nu} := \underbrace{\bar{T}^{(\Phi)}_{\mu\nu} + \bar{T}^{(\Theta)}_{\mu\nu}}_{\text{diagonal}} + \underbrace{T^{(\Phi\Theta)}_{\mu\nu}[\bar{\mathbf{G}}_{\Phi\Theta}]}_{\text{off–diagonal (mixing)}} + \underbrace{T^{(\tau)}_{\mu\nu}[\bar{K}^{(\tau)}, \bar{\mathcal{F}}^{(\tau)}]}_{\text{coherence/torsion}}, \tag{11.264}$$

where the mixed term $T^{(\Phi\Theta)}_{\mu\nu}$ *is induced by the off–diagonal correlator blocks of the unified two–point kernel (triadic propagator; cf. Section 10.19.6) and* $T^{(\tau)}_{\mu\nu}$ *collects the chrono–kymon contributions mapped into* $\bar{t}_{[\mu\nu]}$ *(Remark 11.133 and eq. (11.248)). Assume the chrono–kymon equations of motion (Maxwell/Proca–type on* $(\bar{\Sigma}, \bar{\nabla})$*) hold consistently with the constitutive map* $\bar{t}_{[\mu\nu]} = \chi_\tau\, \bar{\mathcal{F}}^{(\tau)}_{\mu\nu}$*. Then*

$$\nabla^{(\mathrm{rel})}_\mu\, T^{\mathrm{eff}\,\mu}{}_\nu \;=\; 0. \tag{11.265}$$

Proof sketch. Taking $\nabla^{(\mathrm{rel})}$ of (11.264) and using (11.263) shows that the torsion functional $\mathcal{C}_\nu[\mathcal{T}]$ generated from the geometric side is canceled by the divergence of $T^{(\tau)}_{\mu\nu}$ once the chrono–kymon equations and $\bar{t}_{[\mu\nu]} = \chi_\tau\, \bar{\mathcal{F}}^{(\tau)}_{\mu\nu}$ are imposed. The residual exchange between $\bar{T}^{(\Phi)}_{\mu\nu}$ and $\bar{T}^{(\Theta)}_{\mu\nu}$ is mediated by the off–diagonal correlator current in $T^{(\Phi\Theta)}_{\mu\nu}$ (relational diffeomorphism Ward identity for the mixed propagator), yielding (11.265). □

Remark 11.142 (Two equivalent placements of torsion effects)**.** Either (i) keep the Levi–Civita divergence $\bar{\nabla}_\mu \bar{G}^\mu{}_\nu = 0$ and absorb all contorsion/coherence terms into $T^{\mathrm{eff}}_{\mu\nu}$, or (ii) work with the natural derivative $\nabla^{(\mathrm{rel})}$ on the unified diagnostic realized reference configuration and use (11.263) together with (11.265). Both formulations are equivalent; the present choice emphasizes diagnostic realized reference configuration–covariance on $(\bar{\Sigma}, \bar{\nabla})$ and aligns with the operator–valued causal response law (eq. (11.256)) and with the scalarised, EP–safe sector used in 11.5.7.

11.7.4 What is Gravity in the IQR Framework?

Definition 11.143 (Gravity as a relational chronogeometric response)**.** In the IQR framework, *gravity* is the *state–relative, operator–valued response* of the unified *chronogeometric tensor* $\hat{\Sigma}_{\mu\nu} := \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu}$ to physical and informational sources on the BRST/GNS physical sector.

(G0) Background (state dependence). The background chronogeometry is the Holo–State expectation

$$\bar{\Sigma}_{\mu\nu}(x) = \Big\langle \overset{\circ}{\mathbf{\Psi}} \Big|\, \hat{\Sigma}_{\mu\nu}(x) \,\Big|\overset{\circ}{\mathbf{\Psi}} \Big\rangle, \qquad \bar{g}^{\mathrm{Lor}}_{\mu\nu} := \bar{g}_{\mu\nu} + \tfrac{1}{2}\big(\bar{t}_{\mu\nu} + \bar{t}_{\nu\mu}\big), \tag{11.266}$$

which fixes the causal structure (cf. Section 5.4 and eq. (11.232)).

(G1) Operator response map (retarded, covariant). Let $\hat{T}^{\mathrm{nonvac}}_{\rho\sigma}$ collect the non–vacuum source (matter and coherence; the vacuum channel is placed on the geometric side in (G2)). Gravity is encoded by the retarded, type-(2,2) *operator response*

$$\hat{G}_{\mu\nu}(x) = \int_{\mathcal{M}} \mathrm{d}\mu_{\bar{\Sigma}}(x')\, \mathcal{U}_{\bar{\Sigma}}(x \leftarrow x')_{\mu\nu}{}^{\mu'\nu'} \Big[\, \hat{\mathcal{C}}_{\mu'\nu'}{}^{\rho'\sigma'}(x; x') \;\circ\; \hat{T}^{\mathrm{nonvac}}_{\rho'\sigma'}(x') \,\Big], \tag{11.267}$$

where $\hat{G}_{\mu\nu}$ is the Einstein operator of $\hat{\Sigma}$, $A \circ B := \frac{1}{2}(AB + BA)$ is the Jordan product (ensuring self-adjointness), $\mathrm{d}\mu_{\bar{\Sigma}}(x') := \sqrt{-\det \bar{\Sigma}(x')}\,\mathrm{d}^d x'$ is the invariant measure, and

$$\mathcal{U}_{\bar{\Sigma}}(x \leftarrow x')_{\mu\nu}{}^{\mu'\nu'} := \mathcal{U}_{\bar{\Sigma}}(x \leftarrow x')_{\mu}{}^{\mu'}\, \mathcal{U}_{\bar{\Sigma}}(x \leftarrow x')_{\nu}{}^{\nu'}$$

is the Levi–Civita parallel transporter (of $\bar{g}^{\mathrm{Lor}}$) acting on each index; it vanishes unless $x' \in J^-_{\bar{g}^{\mathrm{Lor}}}(x)$ (retarded support). The operator–valued coupling kernel is

$$\hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x') = \mathbb{P}^{(\mathrm{Ein})\alpha\beta}_{\mu\nu}\ \hat{\Delta}^{\mathrm{ret}\,\gamma\delta}_{\alpha\beta}(x,x')\ \mathbb{J}_{\gamma\delta}{}^{\rho\sigma}, \tag{11.268}$$

with $\mathbb{P}^{(\mathrm{Ein})}$ the linearized Einstein (Lichnerowicz–type) map at x, $\hat{\Delta}^{\mathrm{ret}}$ the retarded (TT–projected) propagator of the Gaiamiti fluctuation about $\bar{\Sigma}$, and $\mathbb{J}$ the symmetric matter–to–geometry vertex (species–blind in the minimal case). Thus $\hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x') : \mathrm{Sym}^2_{x'} \to \mathrm{Sym}^2_x$ is a causal bitensor superoperator, mapping the source at x' to the geometric response at x (cf. Definition 11.109 and eqs. (11.213), (11.214) and (11.256)).

(G2) Expectation level, vacuum placement, and scalarisation. Taking the Holo–State expectation $\bar{\mathcal{C}} := \langle \overset{\circ}{\boldsymbol{\Psi}} | \hat{\mathcal{C}} | \overset{\circ}{\boldsymbol{\Psi}} \rangle$, (11.267) becomes the c–number, diagnostic realized reference configuration–covariant bitensor equation

$$\bar{G}_{\mu\nu}(x) \ + \ \Lambda_{\mathrm{eff}}\, \bar{\Sigma}_{\mu\nu}(x) = \int_{\mathcal{M}} \mathrm{d}\mu_{\bar{\Sigma}}(x')\, \mathcal{U}_{\bar{\Sigma}}(x \leftarrow x')_{\mu\nu}{}^{\mu'\nu'}\ \bar{\mathcal{C}}_{\mu'\nu'}{}^{\rho'\sigma'}(x;x')\, \bar{T}^{\mathrm{nonvac}}_{\rho'\sigma'}(x'), \tag{11.269}$$

where the vacuum channel is carried geometrically by $\Lambda_{\mathrm{eff}} = \Lambda + \kappa'\varepsilon_\Lambda$ (Sections 17.1.2 and 17.1.4), so that null contractions of the vacuum term vanish (Lemmas 11.123 and 11.124). Equivalently, $\bar{T}^{\mathrm{total}} = \bar{T}^{\mathrm{nonvac}} - \frac{\Lambda_{\mathrm{eff}}}{\kappa'}\bar{\Sigma}$ moves the same content to the source side.

EP–safe scalarisation. In weak–field, low–coherence domains obeying Section 11.5.7, the expectation kernel is locally isotropic to leading order and reduces to a scalar multiple of the identity on Sym^2:

$$\bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x') \simeq \kappa_{\mathrm{eff}}(x)\, \Pi_{\mu\nu}{}^{\rho\sigma}\, \delta_{\bar{\Sigma}}(x,x'), \qquad \Pi_{\mu\nu}{}^{\rho\sigma} := \tfrac{1}{2}\left(\delta_\mu{}^\rho \delta_\nu{}^\sigma + \delta_\mu{}^\sigma \delta_\nu{}^\rho\right), \tag{11.270}$$

where $\delta_{\bar{\Sigma}}$ is the covariant Dirac density (normalized with respect to $\mathrm{d}\mu_{\bar{\Sigma}}$) and the parallel transport inside $\mathcal{U}_{\bar{\Sigma}}$ is understood. Then (11.269) collapses to the local GR–like law

$$\bar{G}_{\mu\nu}(x) \ + \ \Lambda_{\mathrm{eff}}\, \bar{\Sigma}_{\mu\nu}(x) = \kappa_{\mathrm{eff}}(x)\, \bar{T}^{\mathrm{nonvac}}_{\mu\nu}(x), \qquad \kappa_{\mathrm{eff}} \to \kappa \ \text{ in the GR limit}, \tag{11.271}$$

consistent with Section 11.5.8 and Proposition 11.115.

Bianchi consistency. Applying $\bar{\nabla}^\mu$ to (11.271) and using $\bar{\nabla}^\mu \bar{G}_{\mu\nu} = 0 = \bar{\nabla}^\mu(\Lambda_{\mathrm{eff}}\bar{\Sigma}_{\mu\nu})$ yields $(\bar{\nabla}^\mu \kappa_{\mathrm{eff}})\, \bar{T}^{\mathrm{nonvac}}_{\mu\nu} + \kappa_{\mathrm{eff}}\, \bar{\nabla}^\mu \bar{T}^{\mathrm{nonvac}}_{\mu\nu} = 0$. Thus GR–like conservation follows when the variation controls of Section 11.5.7 suppress $\bar{\nabla}\kappa_{\mathrm{eff}}$ on the scales probed.

Remark 11.144 (On Λ, ε_Λ, and Λ_{eff} (cosmology sector))**.**

(i) **Definition.** In this chapter, the vacuum channel is carried geometrically by

$$\Lambda_{\mathrm{eff}} = \Lambda + \kappa'\varepsilon_\Lambda,$$

so that the expectation–level equation reads $\bar{G}_{\mu\nu} + \Lambda_{\mathrm{eff}}\bar{\Sigma}_{\mu\nu} = \cdots$ as in eq. (11.269). The *observational* cosmological constant corresponds to Λ_{eff}.

(ii) **Meaning of Λ (diagnostic realized reference configuration/bare cosmological term).** Λ is the chronogeometric (diagnostic realized reference configuration) cosmological constant that appears in the Einstein–Hilbert sector of the typed diagnostic realized reference configuration law. It is set by the diagnostic realized reference configuration/plateau choice and renormalization scheme, independent of the state-dependent residual ε_Λ. It has units L^{-2} and contributes as

$$\text{(geometric side)} \quad \Lambda\,\bar{\Sigma}_{\mu\nu} \quad \Longleftrightarrow \quad \text{(source side)} \quad T^{\mathrm{vac}}_{\mu\nu} = -(\Lambda/\kappa')\,\bar{\Sigma}_{\mu\nu},$$

which are equivalent placements at the expectation level used in this chapter.

(iii) **Diagnostic realized reference configuration vs. observed.** Λ is the diagnostic realized reference configuration (bare) cosmological term; the quantity compared to observations is $\Lambda_{\mathrm{eff}} = \Lambda + \kappa'\varepsilon_\Lambda$. Only if $\varepsilon_\Lambda = 0$ would $\Lambda_{\mathrm{eff}} = \Lambda$.

(iv) **Not the spectral cutoff.** The notation Λ_{eff} here is *not* the Chapter IV effective spectral cutoff $\Lambda^{[\mathrm{obs}]}_{\mathrm{eff}}(x, x'; \varepsilon)$. See the cross-chapter dictionary in Remark 4.575.

(v) **Null-silent vacuum.** The vacuum contribution is null-silent: $k^\mu k^\nu \bar{\Sigma}_{\mu\nu} = 0 \Rightarrow k^\mu k^\nu T^{\mathrm{vac}}_{\mu\nu} = 0$ (see Lemmas 11.123 and 11.124).

(G3) Axioms (covariance, causality, conservation, universality).

1. *Covariance & transport.* $\hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x; x')$ is a bitensor on $(\mathcal{M}, \bar{\Sigma})$ and the response (11.267) is built with the Levi–Civita transporter $\mathcal{U}_{\bar{\Sigma}}$ and the invariant measure $\mathrm{d}\mu_{\bar{\Sigma}}$, ensuring full diffeomorphism covariance (cf. Definition 11.109).
2. *Causality & microcausality.* $\hat{\Delta}^{\mathrm{ret}}$ has retarded support, hence $\operatorname{supp}\hat{\mathcal{C}}(x; x') \subset J^-_{\bar{g}^{\mathrm{Lor}}}(x)$ (cf. eq. (11.268)), and BRST/GNS commutators vanish at $\bar{g}^{\mathrm{Lor}}$–spacelike separation.
3. *Self–adjointness & symmetry.* The kernel is symmetric on Sym^2: $\hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma} = \hat{\mathcal{C}}_{\nu\mu}{}^{\rho\sigma} = \hat{\mathcal{C}}_{\mu\nu}{}^{\sigma\rho}$ and pair–Hermitian with respect to the $\bar{\Sigma}$–inner product (cf. §Section 11.5.7). With $A \circ B = \frac{1}{2}(AB+BA)$, $\hat{G}_{\mu\nu}$ is self–adjoint.
4. *Bianchi/consistency.* Since $\bar{\nabla}^\mu \bar{G}_{\mu\nu} = 0$ and $\bar{\nabla}^\mu_x \mathbb{P}^{(\mathrm{Ein})\alpha\beta}_{\mu\nu} = 0$, the expectation–level response with Λ_{eff} in (11.269) is compatible with source conservation modulo controlled exchange with slowly varying κ_{eff} (see Section 11.5.7 and the "Bianchi consistency" paragraph above). The vacuum channel is placed geometrically as $\Lambda_{\mathrm{eff}}\bar{\Sigma}_{\mu\nu}$, whose null contraction vanishes (Lemmas 11.123 and 11.124).
5. *Universality (scalarised limit).* In EP–tested regimes, $\bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma} \propto \Pi_{\mu\nu}{}^{\rho\sigma}$ acts species–blind on $\bar{T}^{\mathrm{nonvac}}_{\rho\sigma}$, preserving universality of free fall (cf. Section 11.5.7 and Proposition 11.113).

(G4) Fluctuation channels and mixing. Small fluctuations of the unified tensor split as $\delta\hat{\Sigma}_{\mu\nu} = \hat{h}_{\mu\nu} \oplus \hat{\tau}_{\mu\nu}$ with $\hat{h}_{\mu\nu} := \delta\hat{g}_{\mu\nu}$ (metric channel) and $\hat{\tau}_{\mu\nu} := \delta\hat{t}_{\mu\nu}$ (temporal/coherence channel). The coherence channel is TT–silent, $\mathcal{P}^{(2)}_{\bar{g}^{\mathrm{Lor}}}[\hat{\tau}_{\mu\nu}] = 0$ (cf. Lemma 11.125 and Remark 11.133), and mixing with $\hat{h}_{\mu\nu}$ is captured by the off–diagonal two–point functions in the retarded kernel (11.268). In the spectral/GR limit this mixing is suppressed and $\hat{\mathcal{C}} \to \kappa\,\Pi\,\delta_{\bar{\Sigma}}$ (Section 11.5.8 and Proposition 11.115).

(G5) Perturbative state. Propagators and loop corrections are computed in the BRST/GNS physical sector using the interacting GNS vacuum associated with the Holo–State, denoted $|\Omega_{\Psi^\circ}\rangle$.

Remark 11.145 (What "push forward" means here)**.** Let G_{hh} be any linear kernel/operator acting on symmetric rank–2 fields (the full metric channel) and let $\Pi_{\bar\Sigma}^{[\mathrm{TT}]}$ be the rig–compatible TT projector (cf. Definition 10.3). Denote by

$$i_{\mathrm{TT}}:\ \Gamma(S^2_{\mathrm{TT}}T^*)\hookrightarrow\Gamma(S^2T^*),\qquad P_{\mathrm{TT}}:=\Pi_{\bar\Sigma}^{[\mathrm{TT}]}:\ \Gamma(S^2T^*)\to\Gamma(S^2_{\mathrm{TT}}T^*)$$

the inclusion and projection (with $P_{\mathrm{TT}}\circ i_{\mathrm{TT}}=\mathrm{id}$ on the TT carrier). *Pushing forward* G_{hh} to the TT sector means taking the operator it *induces* on the TT subbundle:

$$G_{hh}^{[\mathrm{TT}]}\ :=\ P_{\mathrm{TT}}\circ G_{hh}\circ i_{\mathrm{TT}}\ \equiv\ \Pi_{\bar\Sigma}^{[\mathrm{TT}]}\ \circ\ G_{hh}\ \circ\ \Pi_{\bar\Sigma}^{[\mathrm{TT}]}. \tag{11.272}$$

In index (bitensor) form this is the "sandwich"

$$\big(G_{hh}^{[\mathrm{TT}]}\big)_{\mu\nu}{}^{\rho\sigma}(x,y)=\Pi_{\bar\Sigma\,\mu\nu}^{[\mathrm{TT}]\,\alpha\beta}(x)\,G_{hh\,\alpha\beta}{}^{\gamma\delta}(x,y)\,\Pi_{\bar\Sigma\,\gamma\delta}^{[\mathrm{TT}]\,\rho\sigma}(y).$$

Thus one obtains the TT graviton propagator/Green operator used in Lemma 11.146, Definition 10.182, and eq. (10.112). Heuristically, this is the "$4\to2$" reduction: from the full symmetric tensor fiber (10 components in $d{=}4$) down to the *two* physical TT helicities per momentum.

Remarks. (i) The same push–forward applies triad–resolved, because the weave triad projectors commute with typing: $G_{hh}^{[\mathrm{TT}],(a)}=\mathsf{P}^{(2)}_{(a)}\,\Pi_{\bar\Sigma}^{[\mathrm{TT}]}G_{hh}\Pi_{\bar\Sigma}^{[\mathrm{TT}]}\,\mathsf{P}^{(2)}_{(a)}$ (cf. Definition 10.2). (ii) Do not confuse this with the differential–geometric pushforward $f_*:TM\to TN$ of a map f: here "push forward" is the induced map on a *subspace* produced by inclusion–projection. (iii) On the certified plateau, properties like self–adjointness, positivity, and causal support that hold for G_{hh} descend to $G_{hh}^{[\mathrm{TT}]}$ by (11.272), and the resulting TT dynamics coincide with those built from the typed Peierls bracket (eq. (11.25)) and the canonical ETCR (Definition 11.8 and Proposition 11.4).

Lemma 11.146 (Kinematic push–forward of the fourfold inter–modal correlator)**.** *Let* $\mathbf{G}^{(2)}_{\Sigma;AB}(x,x')$ *denote the fourfold inter–modal correlator from Definition 6.4 and Section 6.1.1, with* $A,B\in\{0,1,p,s\}$ *(and implicit* $\mathbf{\Phi}/\mathbf{\Theta}$ *blocks as in Ch. Chapter 6). Let* $(\mathcal{S}_A)_{\mu\nu}$ *and* $(\mathcal{T}_A)_{\mu\nu}$ *be the constitutive maps that produce the rank–2 kinematic fields (metric channel* h *and temporal/coherence channel* τ*), with* $\mathcal{P}^{(2)}_{\bar g^{\mathrm{Lor}}}[\hat\tau]=0$ *(cf. Sections 10.3, 10.4 and 10.18.7). Then the* (h,τ) *two–point* bitensor *kernels are the push–forwards*

$$G_{hh\,\mu\nu}{}^{\rho\sigma}(x,x')=(\mathcal{S}_A)_{\mu\nu}(x)\ \mathbf{G}^{(2)}_{\Sigma;AB}(x,x')\ (\mathcal{S}_B^\dagger)^{\rho\sigma}(x'), \tag{11.273}$$

$$G_{h\tau\,\mu\nu}{}^{\rho\sigma}(x,x')=(\mathcal{S}_A)_{\mu\nu}(x)\ \mathbf{G}^{(2)}_{\Sigma;AB}(x,x')\ (\mathcal{T}_B^\dagger)^{\rho\sigma}(x'), \tag{11.274}$$

$$G_{\tau h\,\mu\nu}{}^{\rho\sigma}(x,x')=(\mathcal{T}_A)_{\mu\nu}(x)\ \mathbf{G}^{(2)}_{\Sigma;AB}(x,x')\ (\mathcal{S}_B^\dagger)^{\rho\sigma}(x'), \tag{11.275}$$

$$G_{\tau\tau\,\mu\nu}{}^{\rho\sigma}(x,x')=(\mathcal{T}_A)_{\mu\nu}(x)\ \mathbf{G}^{(2)}_{\Sigma;AB}(x,x')\ (\mathcal{T}_B^\dagger)^{\rho\sigma}(x'), \tag{11.276}$$

with modal indices A, B summed. Here $(\cdot)^\dagger$ is the formal adjoint with respect to the $\bar{g}^{\mathrm{Lor}}$–induced inner product on Sym^2 *and the invariant measure* $\mathrm{d}\mu_{\bar{\Sigma}}$*; it* includes *Levi–Civita parallel transport from x' to x, so the right index pair $(\rho\sigma)$ is compared at x before contraction. Consequently, the 2×2 kinematic kernel $\left(\begin{smallmatrix} G_{hh} & G_{h\tau} \\ G_{\tau h} & G_{\tau\tau}\end{smallmatrix}\right)$ used in Definition 10.182 and eq. (10.112) is the* kinematic reduction *(push–forward) of the* modal *fourfold object $\mathbf{G}^{(2)}_{\Sigma;AB}$ defined in Section 6.1.1.*

Proof sketch. Collect the constitutive maps into rectangular operators $\mathbb{S} := (\mathcal{S}_A)$ and $\mathbb{T} := (\mathcal{T}_A)$ mapping the modal sector to Sym^2. Then, by linearity of the push–forward on two–point functions,

$$\mathbf{G}_{(h,\tau)} = \begin{pmatrix} \mathbb{S} \\ \mathbb{T} \end{pmatrix} \mathbf{G}^{(2)}_{\Sigma} \begin{pmatrix} \mathbb{S}^\dagger & \mathbb{T}^\dagger \end{pmatrix},$$

with $\dagger$ the formal adjoint relative to the $\bar{g}^{\mathrm{Lor}}$ inner product and Levi–Civita transport from x' to x. Expanding the block product yields (11.273)–(11.276). Retarded/Hadamard/advanced variants follow componentwise, and causal support is preserved under push–forward. □

Remark 11.147 (Matrix shorthand, TT isolation, and projections)**.** Let $\mathbb{P}_{\mathrm{TT}} \equiv \mathcal{P}^{(2)}_{\bar{g}^{\mathrm{Lor}}}$ denote the transverse–traceless (TT) projector on Sym^2 with respect to $\bar{g}^{\mathrm{Lor}}$. The absence of spin–2 content in the τ channel implies

$$\mathbb{P}_{\mathrm{TT}} \circ \mathbb{T} \equiv 0, \qquad \mathbb{T}^\dagger \circ \mathbb{P}_{\mathrm{TT}} \equiv 0,$$

hence the projected kernels satisfy

$$\mathbb{P}_{\mathrm{TT}} \circ G_{\tau\tau} \circ \mathbb{P}_{\mathrm{TT}} = 0, \qquad \mathbb{P}_{\mathrm{TT}} \circ G_{h\tau} \circ \mathbb{P}_{\mathrm{TT}} = 0, \qquad \mathbb{P}_{\mathrm{TT}} \circ G_{\tau h} \circ \mathbb{P}_{\mathrm{TT}} = 0. \tag{11.277}$$

Therefore no *new* TT basis directions arise from the τ sector; the TT radiative content is governed entirely by the hh block, through $\mathbb{P}_{\mathrm{TT}} \circ G_{hh} \circ \mathbb{P}_{\mathrm{TT}}$.

In matrix shorthand one has

$$\mathbf{G}_{(h,\tau)} = \begin{pmatrix} \mathbb{S} \\ \mathbb{T} \end{pmatrix} \mathbf{G}^{(2)}_{\Sigma} \begin{pmatrix} \mathbb{S}^\dagger & \mathbb{T}^\dagger \end{pmatrix}, \qquad \mathbb{P}_{\mathrm{TT}}\, G_{\tau\tau}\, \mathbb{P}_{\mathrm{TT}} = 0,$$

with blocks given in Lemma 11.146. These 2×2 kernels are precisely the inputs that feed the coupling kernel $\hat{\mathcal{C}}$ (cf. Definition 11.109 and eq. (11.268)) and the E(2) polarization analysis (cf. Remarks 11.150 and 11.151).

Remark 11.148 (Change of modal basis)**.** Consider the (h, τ) doublet as a column vector $\left(\hat{h}, \hat{\tau}\right)^\top$. Under any admissible modal redefinition

$$\begin{pmatrix} \hat{h} \\ \hat{\tau} \end{pmatrix} \longmapsto U \begin{pmatrix} \hat{h} \\ \hat{\tau} \end{pmatrix}, \qquad U \in \mathrm{GL}(2) \text{ (unitary on the BRST/GNS physical sector)},$$

the data transform as

$$\mathbf{G}^{(2)}_{\Sigma} \;\mapsto\; U\, \mathbf{G}^{(2)}_{\Sigma}\, U^\dagger, \qquad \mathbb{S} \;\mapsto\; \mathbb{S}\, U^\dagger, \qquad \mathbb{T} \;\mapsto\; \mathbb{T}\, U^\dagger.$$

It follows that the assembled two–channel kernel

$$\mathbf{G}_{(h,\tau)} = \begin{pmatrix} \mathbb{S} \\ \mathbb{T} \end{pmatrix} \mathbf{G}_{\Sigma}^{(2)} \begin{pmatrix} \mathbb{S}^{\dagger} & \mathbb{T}^{\dagger} \end{pmatrix}$$

is invariant, and so are its TT/non–TT projections—particularly $\mathbb{P}_{\mathrm{TT}} \circ G_{hh} \circ \mathbb{P}_{\mathrm{TT}}$ and the null relations in (11.277). Hence the physical TT content and the mixing pattern are basis–independent.

Remark 11.149 (Change of modal basis: general $GL(n)$ form)**.** Let the typed/modal stack be the column vector of operators

$$\widehat{\mathbf{m}} := \big(\hat{m}_1, \dots, \hat{m}_n\big)^{\top},$$

with two–point (bitensor) kernel $\mathbf{G}_{\Sigma}^{(2)}$ on this stack. For any invertible modal redefinition

$$\widehat{\mathbf{m}} \;\longmapsto\; U\,\widehat{\mathbf{m}}, \qquad U \in GL(n) \;\; \text{(unitary on the BRST/GNS physical sector)},$$

the data transform by

$$\mathbf{G}_{\Sigma}^{(2)} \;\mapsto\; U\,\mathbf{G}_{\Sigma}^{(2)}\,U^{\dagger}, \qquad \mathbb{X} \;\mapsto\; \mathbb{X}\,U^{\dagger} \quad \text{for every row "selector" } \mathbb{X} \in \mathbb{C}^{1\times n} \text{ (e.g. } \mathbb{S}, \mathbb{T}).$$

Given any finite family of such selectors $\{\mathbb{X}_{\alpha}\}_{\alpha=1}^{r}$, form the assembled $r\times r$ kernel

$$\mathbf{G}_{(\mathbb{X})} \;:=\; \begin{pmatrix} \mathbb{X}_1 \\ \vdots \\ \mathbb{X}_r \end{pmatrix} \mathbf{G}_{\Sigma}^{(2)} \begin{pmatrix} \mathbb{X}_1^{\dagger} & \cdots & \mathbb{X}_r^{\dagger} \end{pmatrix}. \tag{11.278}$$

Then $\mathbf{G}_{(\mathbb{X})}$ is *basis–independent*: under $U \in GL(n)$ the factors on the left/right of $\mathbf{G}_{\Sigma}^{(2)}$ pick up $U^{\dagger}$ and cancel the U, $U^{\dagger}$ inside, so $\mathbf{G}_{(\mathbb{X})}$ is unchanged. In particular, for the (h,τ) selectors $\mathbb{S}, \mathbb{T}$ one recovers the invariance of

$$\mathbf{G}_{(h,\tau)} = \begin{pmatrix} \mathbb{S} \\ \mathbb{T} \end{pmatrix} \mathbf{G}_{\Sigma}^{(2)} \begin{pmatrix} \mathbb{S}^{\dagger} & \mathbb{T}^{\dagger} \end{pmatrix}.$$

Moreover, all TT/non–TT projections of the assembled blocks are invariant—for example,

$$G_{hh}^{[\mathrm{TT}]} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \circ G_{hh} \circ \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \quad \text{(cf. } \textit{Definition } 10.3, \textit{Lemma } 11.146, \textit{and eq. } (10.112))$$

is independent of the choice of modal basis. Hence the physical TT content and the mixing pattern are basis–independent within the entire admissible $GL(n)$ class.

Remark 11.150 (Triadic origin of extra GW polarizations)**.** In the IQR framework the observable metric response is sourced by three *flavours* of the graviton channel, $a \in \{\mathrm{spec}, \mathrm{vac}, \mathrm{corr}\}$ (the triadic split), and is obtained from the fourfold inter–modal correlator by the constitutive push–forward (Chs. Chapters 6 and 11, Section 6.1.1, Lemma 11.146, and Definition 11.143). Decomposing the observable perturbation in the E(2) polarization basis (Remark 11.151),

$$h_{ij}^{\mathrm{obs}}(x) = \sum_{A\in\{+,\times,x,y,b,L\}} h_A(x)\, e_{ij}^{A},$$

one finds:

1. **Tensor $(+,\times)$: spectral flavour only.** The TT amplitudes are governed solely by the spectral channel, because the τ-channel has no spin-2 content and the TT projector annihilates $\mathbb{T}$ (Lemma 11.146 and Remark 11.147):

$$(h_+, h_\times) \;\propto\; \mathcal{P}^{(2)}_{\bar{g}^{\mathrm{Lor}}}\Big[(\mathcal{S}_A)\,\mathbf{G}^{(2,\mathrm{spec})}_{\Sigma;AB}\,(\mathcal{S}^\dagger_B)\Big]\,\mathcal{J}_h,$$

reproducing GR in the EP-safe scalarised limit (Section 11.5.7 and Proposition 11.113).

2. **Vector/scalar (x, y, b, L): triadic mixing.** Any non–TT amplitude is an *induced response* from off–diagonal mixing between the metric and temporal/coherence channels, sourced by the *vacuum* and/or *correlational* flavours:

$$h_A \;\propto\; e^{ij}_A\,\mathbb{P}_{\mathrm{nonTT}} \sum_{a\in\{\mathrm{vac,corr}\}} \Big[(\mathcal{S}_A)\,\mathbf{G}^{(2,a)}_{\Sigma;AB}\,(\mathcal{T}^\dagger_B)\Big]\,\mathcal{J}^{(a)}_\tau, \qquad \mathbb{P}_{\mathrm{nonTT}} := \mathbf{1} - \mathcal{P}^{(2)}_{\bar{g}^{\mathrm{Lor}}}, \tag{11.279}$$

with $\mathcal{J}_h, \mathcal{J}_\tau$ the channel sources entering the response kernel (Definitions 11.109 and 11.143 and eq. (11.268)). Thus detections of x, y, b, L power in interferometer networks, PTA cross–correlations, or SGWB angular spectra would diagnose nonzero triadic weights together with nonvanishing off–diagonal blocks $G_{h\tau}, G_{\tau h}$ inherited from $\mathbf{G}^{(2,\mathrm{vac/corr})}_\Sigma$.

3. **Consistency with EP/PPN bounds.** In EP–tested (scalarised) domains the variation/aniso controls of Section 11.5.7 ensure $\|\mathbb{P}_{\mathrm{nonTT}} \circ G_{h\tau}\| \ll \|G^{\mathrm{TT}}_{hh}\|$, so $|h_{x,y,b,L}| \ll |h_{+,\times}|$; larger admixtures may occur in high–coherence or strong–field regimes, without introducing *extra* TT polarizations (cf. Remarks 11.147 and 11.151).

In short: *any observed extra GW polarizations in IQR trace back to triadic flavour content and metric–coherence mixing*, not to additional massless spin–2 degrees of freedom. The spectral flavour yields the GR tensor pair, while the vacuum/correlational flavours feed vector/scalar *responses* through the off–diagonal push–forward (11.279).

Remark 11.151 (E(2) polarization content and its realization in IQR)**.** In a local metric theory, plane–wave gravitational disturbances admit up to *six* independent polarizations (the E(2) classification): two tensor $(+,\times)$, two vector (x, y), and two scalar (breathing b and longitudinal L). In the IQR framework (Sections 4.8 and 11.5 and Chapters 6 and 11), the *metric response* can likewise exhibit up to six *observable* polarization templates. The *tensor* pair arises from the spectral channel (TT graviton), whereas possible *vector/scalar* components originate as *triadic admixtures* via the mixed propagator blocks $G_{h\tau}, G_{\tau h}$ (Remark Remark 11.150, Lemma 11.146). Throughout, the E(2) basis is understood in the local geometric–optics limit of $\bar{g}^{\mathrm{Lor}}$.

Local polarization basis and normalization. Let $\hat{n}$ be the propagation direction and $\{e_1, e_2, \hat{n}\}$ an orthonormal triad in a local Lorentz frame adapted to $\bar{g}^{\mathrm{Lor}}$. Define the six standard polarization tensors on the screen space,

$$\begin{aligned}
e^+_{ij} &:= e_{1i}e_{1j} - e_{2i}e_{2j}, & e^\times_{ij} &:= e_{1i}e_{2j} + e_{2i}e_{1j},\\
e^x_{ij} &:= e_{1i}\hat{n}_j + \hat{n}_i e_{1j}, & e^y_{ij} &:= e_{2i}\hat{n}_j + \hat{n}_i e_{2j},\\
e^b_{ij} &:= e_{1i}e_{1j} + e_{2i}e_{2j}, & e^L_{ij} &:= \hat{n}_i\hat{n}_j,
\end{aligned}$$

normalized so that $e^A_{ij} e^{B\,ij} = 2\,\delta^{AB}$. Any observable metric perturbation decomposes as

$$h^{\rm obs}_{ij}(x) = \sum_{A\in\{+,\times,x,y,b,L\}} h_A(x)\, e^A_{ij}, \qquad h_A := \tfrac{1}{2}\, h^{\rm obs}_{ij}\, e^{A\,ij}. \tag{11.280}$$

IQR realization (origin of each mode).

- *Tensor (two helicities).* The radiative TT modes are

$$\hat h^{[\mathrm{TT}],(\mathrm{spec})}_{\mu\nu} := \left(\mathsf{P}^{(2)}_{\rm spec}\, \Pi^{[\mathrm{TT}]}_{\bar\Sigma} \hat h\right)_{\mu\nu},$$

  with $\Pi^{[\mathrm{TT}]}_{\bar\Sigma}$ the rig–compatible TT projector (Definition 10.3) and $\mathsf{P}^{(2)}_{\rm spec}$ the spectral idempotent inside the TT block (provenance split; Definition 10.2). In the scalarized/weak-field limit (Section 11.5.7 and Proposition 11.115) these reduce to the GR tensor pair $(+,\times)$ in the standard TT polarization basis (cf. eq. (11.100)).

- *Vector/Scalar* (x,y,b,L). The coherence channel satisfies $\mathcal{P}^{(2)}_{\bar g^{\rm Lor}}[\hat\tau_{\mu\nu}] = 0$ (Section 10.18.7) but induces *non–TT metric responses* through off–diagonal mixing:

$$h^{\rm obs}_{ij} = \left(\mathcal{P}^{(2)}_{\bar g^{\rm Lor}} + \mathbb{P}_{\rm nonTT} \circ G_{h\tau} \circ \mathbb{J}_{\tau\to h}\right)[\text{sources}]_{ij}\,, \qquad \mathbb{P}_{\rm nonTT} := \mathbf{1} - \mathcal{P}^{(2)}_{\bar g^{\rm Lor}},$$

  with sources and vertex as in (11.267)–(11.268). Projecting (11.280) with $e^{x,y,b,L}$ yields the amplitudes $h_{x,y,b,L}$. Their magnitudes are controlled by the triadic weights and the state (Remark Remark 11.150); in EP–tested, scalarised domains the bounds of Section 11.5.7 ensure $|h_{x,y,b,L}| \ll |h_{+,\times}|$.

Summary. Chs. Chapters 4, 6 and 11 and Section 11.5 permit up to six *observable* polarization templates at the level of the metric response. The *radiative* massless content remains the standard TT pair from the spectral flavour; any *additional* vector/scalar responses arise via triadic mixing and are state- and regime-dependent, without introducing extra TT polarizations.

Remark 11.152 (What $E(2)$ means in this chapter)**.** Fix a future–pointing null direction k^μ (e.g. the propagation direction of a TT graviton mode) and choose a null partner ℓ^μ with $k\cdot\ell = -1$. The associated *screen* is the $(d-2)$–dimensional space orthogonal to $\{k,\ell\}$ with metric $q_{\mu\nu} = \bar g^{\rm Lor}_{\mu\nu} + k_\mu\ell_\nu + \ell_\mu k_\nu$ (cf. Remark 11.127). In $d=4$ the screen is 2–dimensional, and

$$E(2) \;:=\; \mathrm{Isom}(\mathbb{R}^2) \;\simeq\; \mathbb{R}^2 \rtimes SO(2)$$

is the Euclidean group of the screen: $SO(2)$ rotations about k and the (screen) translations. Equivalently, $E(2)$ is the *little group* of a null vector: the subgroup of the Lorentz group that leaves k^μ invariant. In our $\bar\Sigma$–based notation this is the symmetry that acts on the *screen* indices of polarization objects at fixed k.

Why it matters for TT graviton modes.

- The $SO(2)$ *rotation* part of $E(2)$ acts on the TT polarization basis $e^{(a,\lambda)}_{\mu\nu}(k)$ (eq. (11.100)) as a helicity rotation: the two TT modes furnish the $\lambda = \pm 2$ representation, i.e. under a screen rotation by angle θ, $e^{(\pm)} \mapsto e^{(\pm)} e^{\pm 2i\theta}$. This is the precise meaning of the "two tensor helicities."
- The $\mathbb{R}^2$ *translation* part (null rotations) corresponds to gauge transformations that shift non-TT (pure-gauge/longitudinal) pieces. In the *TT-projected* theory we work with $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}} \hat{h}_{\mu\nu}$ (Definition 10.3), so these translations act trivially on physical TT data.

Relation to our typed/triad structure. On the certified plateau, the radiative spin-2 content lives entirely in the TT carrier with helicities ± 2 (the $SO(2)$ sector above); vector/scalar screen representations do not appear in the gravitational wave channel because the τ–kairos sector is TT–null ($\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}} \hat{\tau}_{\mu\nu} = 0$; cf. eq. (10.139)), and the weave triad bookkeeping (Definition 10.2) is compatible with the same projector. Hence, within Chapter 12 the shorthand "$E(2)$" simply signals the screen–space symmetry behind helicity and polarization statements for TT modes.

Proposition 11.153 (E(2) completeness of the metric response in IQR)**.** *Fix a local Lorentz frame of $\bar{g}^{\mathrm{Lor}}$ and a plane–wave direction $\hat{n}$. Let $\{e_1, e_2, \hat{n}\}$ be an orthonormal triad and let $\{e^A_{ij}\}_{A\in\{+,\times,x,y,b,L\}}$ be the standard E(2) polarization tensors on the screen space (cf. Remark Remark 11.151). In the triadic response formalism (Section 10.17.6, Definition 10.182, and eq. (10.112)), the observable metric perturbation admits the full six–mode decomposition*

$$h^{\mathrm{obs}}_{ij}(x) = \sum_{A\in\{+,\times,x,y,b,L\}} h_A(x)\, e^A_{ij}, \qquad h_A := \tfrac{1}{2}\, h^{\mathrm{obs}}_{ij} e^{A\,ij}. \tag{11.281}$$

Moreover, writing $\mathbb{P}_{\mathrm{nonTT}} := \mathbf{1} - \mathcal{P}^{(2)}_{\bar{g}^{\mathrm{Lor}}}$ and letting $\star$ denote the retarded convolution with Levi–Civita transport and invariant measure (eqs. (11.267) and (11.268)), one has

$$h_{+,\times} \;=\; \left(\mathcal{P}^{(2)}_{\bar{g}^{\mathrm{Lor}}}\, G^{\mathrm{ret}}_{hh}\right) \star \mathcal{J}_h, \qquad \textit{(spectral TT graviton)}, \tag{11.282}$$

$$(h_x, h_y, h_b, h_L) \;=\; \Big[\; \mathbb{P}_{\mathrm{nonTT}} \circ \left(G^{\mathrm{ret}}_{h\tau} \star \mathcal{J}_\tau \;+\; G^{\mathrm{ret}}_{hh} \star \mathcal{J}^{\mathrm{nonTT}}_h\right) \Big]_{\textit{projected}}, \tag{11.283}$$

where $\mathcal{J}_h$ and $\mathcal{J}_\tau$ are the metric and coherence source currents entering the operator coupling kernel $\hat{\mathcal{C}}$ (Definition 11.109 and eq. (11.268)), and $\mathcal{J}^{\mathrm{nonTT}}_h := \mathbb{P}_{\mathrm{nonTT}} \mathcal{J}_h$. "Projected" means contraction with $e^{A\,ij}$ as in (11.281).

Consequently:

- *The* radiative *massless spin–2 content is the usual TT pair from the spectral channel ($\mathcal{P}^{(2)}_{\bar{g}^{\mathrm{Lor}}}[\hat{\tau}] = 0$, Section 10.18.7).*
- *The* vector/scalar *metric responses arise as* triadic admixtures*. They are induced by the off-diagonal blocks $G_{h\tau}$ and $G_{\tau h}$ (push–forward of the fourfold correlator; Lemma Lemma 11.146), together with any non-TT part of $\mathcal{J}_h$.*

In EP–tested, scalarised domains (Section 11.5.7) one has $|h_{x,y,b,L}| \ll |h_{+,\times}|$; in high–coherence or strong–field regimes larger admixtures are possible without introducing extra TT polarizations.

Remark 11.154 (Phenomenology windows vs. weak-field expectations)**.** Within IQR none of the following effects is a fundamental no–go; each is *permitted* in suitable, state–dependent regimes governed by the operator kernel $\bar{\mathcal{C}}$ and the triadic weights (Definition 11.109 and Section 10.17.6). In EP–tested (scalarised) domains (Section 11.5.7) they are tightly bounded.

1. **Tensor speed deviations (non-luminal c_T at finite frequency).** The spectral TT graviton's dispersion follows from the pole of the TT block of the push–forward kernel (Lemma 11.146):

$$\det D^{\mathrm{TT}}(\omega, \mathbf{k}; \overset{\circ}{\mathbf{\Psi}}) = 0, \qquad G^{\mathrm{TT}}_{hh}(\omega, \mathbf{k}) = \left[D^{\mathrm{TT}}\right]^{-1}.$$

   Retarded support of $\hat{\mathcal{C}}$ enforces causality (Definition 11.109), so the *front* velocity is luminal with respect to $\bar{g}^{\mathrm{Lor}}$. The *phase/group* speed $c_T(\omega, \mathbf{k})$ can nevertheless acquire state/triadic self-energy dressings $\delta c_T^2(\overset{\circ}{\mathbf{\Psi}})$ through $\bar{\mathcal{C}}$. In scalarised regions $\delta c_T^2 = \mathcal{O}(\varepsilon_{\mathrm{aniso}} + \varepsilon_{\mathrm{grad}}) \ll 1$; in high-coherence or strong-field regimes larger (still causal) deviations are allowed.

2. **Vector/scalar GW power at detectors ("extra polarizations").** E(2) x, y, b, L components are *non-TT metric responses* induced by triadic mixing:

$$h_A \;\propto\; e_A^{ij}\, \mathbb{P}_{\mathrm{nonTT}}\Big[(\mathcal{S}_A)\, \mathbf{G}^{(2)}_{\Sigma}\, (\mathcal{T}_B^{\dagger})\Big]\mathcal{J}_\tau, \qquad A \in \{x, y, b, L\},$$

   i.e. via $G_{h\tau}, G_{\tau h}$ obtained by push–forward of the fourfold correlator (Section 6.1.1, Lemma 11.146, and Remark 11.151). In EP–tested regimes these amplitudes are suppressed by the scalarisation/variation bounds (Section 11.5.7); outside them (early universe, near compact objects) they can be appreciable.

3. **Multiple TT flavours (same two tensor basis states).** In $D = 4$ there are only two TT basis polarizations, but IQR allows distinct TT *flavours*—spectral, vacuum-dressed, correlational—each contributing to the *same* TT basis with different dispersion/damping,

$$G^{\mathrm{TT}}_{hh} = \sum_{a \in \{\mathrm{spec,vac,corr}\}} \mathcal{P}^{(2)}_{\bar{g}^{\mathrm{Lor}}} \circ \Big[(\mathcal{S})\, \mathbf{G}^{(2,a)}_{\Sigma}\, (\mathcal{S}^{\dagger})\Big] \circ \mathcal{P}^{(2)}_{\bar{g}^{\mathrm{Lor}}}.$$

   Observable signatures include mild dispersion, birefringence, or phase beating of $(+, \times)$—not additional independent TT directions, consistent with $\mathcal{P}^{(2)}_{\bar{g}^{\mathrm{Lor}}}[\hat{\tau}] = 0$ (no spin–2 in the τ channel; Section 10.18.7).

Bottom line. GR-like behavior (luminal front velocity, negligible non-TT power) is the *scalarised, weak-field expectation.* Departures are *allowed* and are controlled by the state-dependent triadic dressings and the operator kernel $\bar{\mathcal{C}}$, consistent with covariance, causality, and the relational NEC (Sections 11.5.7 and 11.6 and Definition 11.109).

Radiative content. The *graviton* is the TT (spin–2) projection of the metric channel with respect to the unified background:

$$\hat{h}^{[\mathrm{TT}]}_{\mu\nu} \;:=\; \left(\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\hat{h}\right)_{\mu\nu} \qquad (\text{cf. } \textit{Definition } 10.3),$$

and its dynamics are governed by the TT–sandwiched (pushed–forward) kernel

$$G^{[\mathrm{TT}]}_{hh} \;:=\; \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}} \;\circ\; G_{hh} \;\circ\; \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}} \tag{11.284}$$

See Lemma 11.146, Definition 10.182, and eq. (10.112); cf. [415, 502].

The *tau–kairos* sector is strictly non–TT at linear order,

$$\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\hat{\tau}_{\mu\nu} = 0 \quad (\text{Sections } 10.3,\ 10.4 \text{ and } 10.18.7),$$

so any *observed* vector/scalar GW responses arise only via triadic mixing in the off–diagonal blocks $G_{h\tau}$, $G_{\tau h}$ (Remark Remark 11.150).

Origin and symmetry. The unique long–range, massless radiative mode is the spin–2 graviton, identified as the geometric Goldstone excitation of the Gaiamiti sector's vacuum (Section 10.17.9); massive geometric scalars (apeiron) are radial (non–Goldstone) modes (Section 10.16). By contrast, the temporal/coherence channel $\hat{\tau}_{\mu\nu}$ is populated by *Tantrakyma* fluctuations via the constitutive map

$$\hat{t}_{[\mu\nu]} = \chi_\tau\,\hat{\mathcal{F}}^{(\tau)}_{\mu\nu}, \qquad \hat{\mathcal{F}}^{(\tau)}_{\mu\nu} := 2\,\bar{\nabla}_{[\mu}\hat{K}^{(\tau)}_{\nu]},$$

with spin–1 *Chrono–Kymon* potential $\hat{K}^{(\tau)}_\mu$ (often $\hat{A}_\mu$) (Sections 10.7 and 10.17.9). This sector is non-radiative in the TT sense (Sections 10.3 and 10.4) but can seed non-TT *metric responses* through $G_{h\tau}, G_{\tau h}$.

Triadic factorization of sources. Curvature fluctuations admit a *triadic* resolution into correlational, vacuum, and spectral flavours:

$$\mathbf{G}^{(2)}_\Sigma = \sum_{a\in\{\mathrm{corr,vac,spec}\}} \mathbf{G}^{(2,a)}_\Sigma, \qquad \hat{h}_{\mu\nu} = \sum_a \hat{h}^{(a)}_{\mu\nu},$$

separating entanglement–induced, vacuum–manifold, and energy–momentum–induced contributions (Sections 10.17.5 and 10.17.6). TT projection w.r.t. $\bar{g}^{\mathrm{Lor}}$ isolates the radiative content in each flavour while preserving the off–diagonal (h,τ) dressings at the response level (Definition 10.182 and Lemma 11.146). In particular, the observable TT block is

$$G^{\mathrm{TT}}_{hh} = \sum_a \mathcal{P}^{(2)}_{\bar{g}^{\mathrm{Lor}}} \circ \Big[(\mathcal{S})\,\mathbf{G}^{(2,a)}_\Sigma\,(\mathcal{S}^\dagger)\Big] \circ \mathcal{P}^{(2)}_{\bar{g}^{\mathrm{Lor}}},$$

and possible vector/scalar responses trace to $\mathbb{P}_{\mathrm{nonTT}} \circ G_{h\tau}$ (Remarks 11.150, 11.151).

Matter coupling and temporal torsion. The *Tantrakyma* field $\hat{\mathbf{\Theta}}$ furnishes the matter/coherence sector. Its fluctuations generate (i) *masses* via the operator mass matrix $\mathcal{M}_\Theta$ and projected propagators Δ_p (Sections 10.18.5 and 10.18.6); and (ii) a *temporal imprint* via

$$\hat{t}_{[\mu\nu]} = \chi_\tau \, \hat{\mathcal{F}}^{(\tau)}_{\mu\nu}, \qquad \hat{\mathcal{F}}^{(\tau)}_{\mu\nu} = 2\,\bar{\nabla}_{[\mu}\hat{K}^{(\tau)}_{\nu]},$$

sourced by the spin–1 Chrono–Kymon mode $\hat{K}^{(\tau)}_\mu = \delta\hat{\mathbf{\Theta}}^{(1)}_\mu$ (Sections 10.7 and 10.17.9). Thus, gravity in IQR is the joint, channel–resolved response of $\bar{\Sigma}_{\mu\nu}$ to energetic and coherence–informational sources.

Operational content (summary).

- *Causality:* propagation is governed by the $\bar{g}^{\mathrm{Lor}}$ light cone (equivalently, the $\bar{\Sigma}$ cone; Lemma 11.123) (Section 5.4.5); response kernels are retarded (Definition 11.109).
- *Radiation:* the TT block G^{TT}_{hh} carries exactly two tensor polarizations; any non-TT detector responses arise only via triadic mixing (Remark 11.150) (Sections 10.3, 10.4 and 10.17.8).
- *Background dependence:* amplitudes, phases, and effective ranges acquire triadic, state–dependent dressings through $\mathbf{G}^{(2,a)}_\Sigma$ and the Tantrakyma blocks $\mathbf{\Delta}^{(a)}_\Theta$ (Sections 10.17.6 and 10.18.7).

Bridge to Section 11.9. With gravity defined as a chronogeometric response, the operative regimes are controlled by the relative magnitude and anisotropy of $\bar{t}_{\mu\nu}$ versus $\bar{g}_{\mu\nu}$, the coherence scales entering $\mathcal{M}_\Theta$, and the triadic weights; these parameters organize the phenomenology in Section 11.9.

The world isn't stuff moving on a fixed stage. In this framework, the "stage" itself—the network of distances, clock rates, and cause-and-effect—is a fabric woven from relationships among everything that exists. We summarize that fabric by a single field, the **chronogeometric field** $\bar{\Sigma}_{\mu\nu}$. At the everyday, averaged (expectation) level it splits into a familiar symmetric metric $\bar{g}_{\mu\nu}$ that sets lengths and times, plus a tiny antisymmetric "coherence" piece $\bar{t}_{[\mu\nu]}$ that tracks subtle quantum-relational structure.

So what is gravity? It's how that fabric *responds* when the world is active. Put matter or light somewhere, build correlations, excite quantum fields—anything that counts as "total source"—and the fabric reshapes. Clocks tick a bit differently, straight paths curve, and light rays bend. Crucially, this response is *retarded and causal*: the change ripples outward only inside the light-cone of $\bar{\Sigma}$, never faster than light. Mathematically, it's a rule with "curvature on the left" and a *retarded kernel acting on the total source on the right*—but the picture to keep is simply: pushes on the web cause bends in the web, after a light-speed delay.

Why it mostly looks like Einstein. In ordinary conditions (weak fields, everyday energies) the fabric behaves like a very stiff, smooth sheet. The part of the dynamics that matters is the *transverse–traceless (TT) spectral sector*, and you get exactly Einstein's relation between "push" (stress–energy) and "bend" (curvature) with one constant G. That's why freely falling bodies all follow the same paths and why the classic GR tests work so well.

Where the refinements live. Because the fabric is ultimately *quantum and relational*, there's an additional, state-dependent correction $\mathcal{Q}_{\mu\nu}$ built from that coherence piece. Chapter 12 shows this correction is *null-positive* (it can't spoil the usual light-ray focusing), so the familiar link from energy conditions to geometric focusing still holds. At the same diagnostic realized reference configuration level, the quiet "pressure" of empty space appears as an effective cosmological constant Λ. These refinements are tiny in the solar system, but they can matter near extremes like the early universe or compact objects.

Gravitational waves, simply. Jiggle the fabric fast and ripples carry energy away: those are gravitational waves. In normal conditions they come with the two familiar "plus" and "cross" patterns—the two *TT helicities* of the radiating graviton. In exotic states, other degrees of freedom can *mix into the metric response* and produce small extra squeeze/longitudinal features in what detectors read out, but they do *not* add new radiative polarizations. (The τ–kairos sector is *TT-null*, so it cannot radiate spin-2.)

Is gravity a force? Fundamentally, *no.* Objects in free fall are not being pushed; they're following the straightest possible paths (geodesics) in a curved fabric. The "force of gravity" shows up only as a convenient approximation—like the Newtonian potential—in slow, weak-field regimes, or as *tidal effects* (relative accelerations between nearby paths) that are really geometry in disguise.
Newtonian / weak-field limit. When fields are weak and speeds are slow, the chronogeometric field looks almost flat and its time–time piece satisfies $g_{00} \approx -(1 + 2\Phi)$. In this limit you recover the everyday "force law":

$$\mathbf{F} \;=\; m\,\mathbf{a} \;=\; -\,m\,\nabla\Phi.$$

If the potential is $\Phi(r) = -\dfrac{GM}{r}$, then

$$\mathbf{F} \;=\; -\,\frac{GMm}{r^2}\,\hat{\mathbf{r}},$$

the familiar inverse-square "gravity." This is a *repackaging of curvature* into a potential, not a fundamental force; it's how the geometric picture reduces to the equation everyone knows ($F = ma$) in ordinary conditions.

Always causal, always consistent. Conservation laws (the contracted Bianchi identity) ensure there's no mismatch between the "push" and the "bend," and the canonical–vs–covariant (Peierls) ways of formulating the theory agree where we actually make predictions.

One-liner: *Gravity is the causal, retarded reshaping of the chronogeometric fabric $\bar{\Sigma}$ by the total source (matter, interactions, and coherence). It reduces to Einstein's two-polarization TT dynamics in ordinary regimes, while allowing small, testable quantum-relational corrections in the extremes. A gravitational wave is a light-speed, two-polarization ripple of spacetime itself—tiny oscillations of distance and time that arise from accelerating mass–energy and are felt as changing separations, not as a force.*

11.8 Chronogeometry of the τ–Kairos

Scope. This section makes explicit how the non–TT coherence sector—the TT–complement $\hat{\tau}_{\mu\nu}$ defined in Section 10.4 with respect to the realized chronogeometry $(\mathcal{X}_q, \bar{\Sigma})$—enters the diagnostic realized reference configuration Einstein law Definition 11.106 without introducing any duplicate definitions. All constructions here are formulated on certified plateaux $\mathfrak{P}$, with transport by the $\bar{\Sigma}$–compatible relational Levi–Civita connection $\bar{\nabla}$ (cf. Sections 11.5.1 and 11.5.2).

Background shift vs. radiative content

Setup and notation. Work on a certified plateau $\mathfrak{P}$ with realized chronogeometry $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (Ch. 6) and with transport by the $\bar{\Sigma}$–compatible relational Levi–Civita connection $\bar{\nabla}$. Let $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ be the certified TT projector on symmetric rank–2 tensors (Definition 10.3), and $\mathbb{P}_{\mathrm{nonTT}} := \mathrm{Id} - \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$. The non–TT coherence sector is (cf. Section 10.4)

$$\hat{\tau}_{\mu\nu} := \mathbb{P}_{\mathrm{nonTT}}\, \delta\hat{\Sigma}_{\mu\nu}, \qquad \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \hat{\tau}_{\mu\nu} = 0. \tag{11.285}$$

Coherent expectations and background shift. Denote diagnostic realized reference configuration/physical expectations on $\mathfrak{P}$ by $\langle\,\cdot\,\rangle$. A *coherent* τ configuration is one with nonzero mean $\langle\hat{\tau}_{\mu\nu}\rangle \neq 0$. By linearity and (11.285),

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \langle\hat{\tau}_{\mu\nu}\rangle = \langle \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \hat{\tau}_{\mu\nu}\rangle = 0, \tag{11.286}$$

so the mean coherence field is *TT–null*. It therefore shifts the *background* temporal/coherence tensor and with it the realized chronogeometry:

$$\bar{t}_{\mu\nu} \;\longrightarrow\; \bar{t}'_{\mu\nu} := \bar{t}_{\mu\nu} + \Delta t^{(\tau)}_{\mu\nu}, \qquad \Delta t^{(\tau)}_{\mu\nu} := \langle\hat{\tau}_{\mu\nu}\rangle, \qquad \bar{\Sigma}'_{\mu\nu} := \bar{g}_{\mu\nu} + \bar{t}'_{\mu\nu}. \tag{11.287}$$

Typed reconstruction of the mean coherence field. Using the certified typed projectors of the weave (Ch. 11), decompose $\Delta t^{(\tau)}_{\mu\nu}$ into canonical scalar/pseudoscalar/vector pieces:

$$\Delta t^{(\tau)}_{\mu\nu} = S^{(b)}_{\mu\nu}\Big[\langle\hat{\varphi}_b\rangle\Big] + S^{(L)}_{\mu\nu}\Big[\langle\hat{\varphi}_L\rangle\Big] + \tilde{S}^{(p)}_{\mu\nu}\Big[\langle\tilde{\varphi}\rangle\Big] + A^{(\lambda)}_{\mu\nu}\Big[\langle\hat{K}^{(\tau)}_{\lambda}\rangle\Big], \qquad \lambda = \pm, \tag{11.288}$$

where $S^{(b)}, S^{(L)}$ are scalar–typed, $\tilde{S}^{(p)}$ is pseudoscalar–typed, and $A^{(\lambda)}$ is the vector (chrono–kymon) type. In particular, for the vector type the gauge–invariant strength $\mathcal{F}^{(\tau)}_{\mu\nu} := 2\,\bar{\nabla}_{[\mu} K^{(\tau)}_{\nu]}$ induces an antisymmetric temporal imprint

$$\Delta t^{(\tau)}_{[\mu\nu]} \propto \mathcal{F}^{(\tau)}_{\mu\nu}, \qquad \Delta t^{(\tau)}_{(\mu\nu)} \text{ from scalar/pseudoscalar types.} \tag{11.289}$$

Effect on the geometric side of the diagnostic realized reference configuration law. Let $\bar{G}_{\mu\nu}[\bar{\Sigma}]$ be the unified Einstein tensor of the realized geometry (cf. Definition 11.106). Introduce a bookkeeping parameter ε for the coherence amplitude and write $\bar{\Sigma}' = \bar{\Sigma} + \varepsilon\,\Delta\Sigma^{(\tau)}$ with $\Delta\Sigma^{(\tau)}_{\mu\nu} := \Delta t^{(\tau)}_{\mu\nu}$ from (11.287). To first nontrivial order,

$$\bar{G}_{\mu\nu}[\bar{\Sigma}'] = \bar{G}_{\mu\nu}[\bar{\Sigma}] + \varepsilon\,\delta\bar{G}_{\mu\nu}\Big[\Delta\Sigma^{(\tau)}\Big] + \mathcal{O}(\varepsilon^2), \tag{11.290}$$

so a coherent τ background modifies the *geometric side* of the diagnostic realized reference configuration law through the relational variation $\delta\bar{G}[\cdot]$ evaluated on the TT–null $\Delta\Sigma^{(\tau)}$.

No new spin–2 polarizations (TT–purity). Although (11.290) alters the background operator felt by perturbations, it does *not* inject additional spin–2 content. Indeed, the radiative field is $h := \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}'}(\delta\Sigma)$ with respect to the *new* background $\bar{\Sigma}'$; by (11.286) the coherence shift satisfies $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\Delta\Sigma^{(\tau)} = 0$, and the typed hygiene of the weave ensures that the TT projector on $\bar{\Sigma}'$ still isolates precisely two tensor polarizations. In short, $\Delta\Sigma^{(\tau)}$ *moves the background* but does not create extra GW polarizations; the TT graviton (Apeiron$^{[\mathrm{TT}]}$) remains the unique spin–2 radiative content.

Background–dependent radiative operator. Linearizing the diagnostic realized reference configuration law about $\bar{\Sigma}'$ yields the TT wave operator $\mathcal{L}_{hh}[\bar{\Sigma}']$ acting on h. Relative to the unshifted background,

$$\mathcal{L}_{hh}[\bar{\Sigma}'] = \mathcal{L}_{hh}[\bar{\Sigma}] + \varepsilon\,\delta\mathcal{L}_{hh}\Big[\Delta\Sigma^{(\tau)}\Big] + \mathcal{O}(\varepsilon^2), \tag{11.291}$$

so coherent τ modifies *propagation parameters* (dispersion, effective potentials) but not the tensorial *type* of the radiative slice.

Bookkeeping across LHS/RHS. Consistently with (11.290), the diagnostic realized reference configuration source is also shifted: $\bar{T}^{\mathrm{total}} \to \bar{T}^{\mathrm{total}} + \varepsilon\,\Delta\bar{T}^{(\tau)}$, where the coherence contribution $\Delta\bar{T}^{(\tau)}$ is assembled from the typed amplitudes (vector piece yielding the Maxwell–like structure, scalar/pseudoscalar pieces entering the symmetric stress), in harmony with the Relational NEC. The diagnostic realized reference configuration law at $\mathcal{O}(\varepsilon)$ reads

$$\delta\bar{G}_{\mu\nu}\Big[\Delta\Sigma^{(\tau)}\Big] + \Lambda\,\Delta\Sigma^{(\tau)}_{\mu\nu} = \kappa\,\Delta\bar{T}^{(\tau)}_{\mu\nu}, \tag{11.292}$$

which simply states that the *background* coherence shift must be balanced by its diagnostic realized reference configuration source—while the *radiative* TT sector remains governed by $\mathcal{L}_{hh}[\bar{\Sigma}']$ in (11.291).

Summary. Coherent (typed) τ expectations are TT–null and therefore (i) *shift the background* $\bar{t}_{\mu\nu}$, modifying $\bar{\Sigma}$ and the geometric operator via (11.290)–(11.291); (ii) do *not* create additional spin–2 polarizations (TT–purity); and (iii) contribute to the diagnostic realized reference configuration source through typed coherence stress, satisfying the balance (11.292). Gravitational radiation remains purely TT; coherence physics lives in the background and in non–TT responses.

Linearized diagnostic realized reference configuration law: mixed–block operator

Linearizing Definition 11.106 around a realized background $\bar{\Sigma} \mapsto \bar{\Sigma} + h$ and decomposing with $\mathbb{P}_{\mathrm{nonTT}} := \mathrm{Id} - \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ gives the block system

$$\boxed{\begin{pmatrix} \mathcal{L}_{hh} & \mathcal{L}_{h\tau} \\ \mathcal{L}_{\tau h} & \mathcal{L}_{\tau\tau} \end{pmatrix} \begin{pmatrix} h \\ \tau \end{pmatrix} = \kappa \begin{pmatrix} \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \delta\bar{T}^{\mathrm{total}} \\ \mathbb{P}_{\mathrm{nonTT}}\, \delta\bar{T}^{\mathrm{total}} \end{pmatrix}, \qquad \begin{cases} h := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, h, \\ \tau := \mathbb{P}_{\mathrm{nonTT}}\, h \end{cases}} \tag{11.293}$$

with $\delta\bar{T}^{\mathrm{total}}$ the linearized diagnostic realized reference configuration source (cf. Definition 9.358 and Section 9.18.12). By TT–purity (Chapter 11), $\mathcal{L}_{hh}$ governs the spin–2 radiative sector (graviton, i.e. Apeiron$^{[\mathrm{TT}]}$), $\mathcal{L}_{\tau\tau}$ governs the non–TT coherence sector, and the off–diagonal entries $\mathcal{L}_{h\tau}, \mathcal{L}_{\tau h}$ implement the certified mixed response kernels (cf. the weave/typed realized weave configuration).

Remark 11.155 (Causality and well-posedness)**.** All four blocks in (11.293) are realized with Σ–causal propagators on $\mathfrak{P}$ (retarded/advanced support in the $\bar{\Sigma}$–cones), exactly as for the TT radiative operator (cf. the causal statements in Sections 11.5.1 and 11.5.4).

Mixed–block form (non–TT ledger). On the certified plateau $\mathfrak{P}$ with $\bar{\Sigma}$–chronogeometry (cf. Section 11.8), the linearized non–TT sector closes under a typed state vector and a mixed–block operator:

$$\mathbb{K}_{\mathrm{DRRC}} \begin{pmatrix} \delta g_{\text{non-TT}} \\ \tau_{\mathrm{scalar}} \\ K_\mu^{(\tau)} \end{pmatrix} = \begin{pmatrix} \mathcal{S}_g^{(\mathrm{coh})} + \mathcal{S}_g^{(\mathrm{str})} \\ \mathcal{S}_S^{(\mathrm{coh})} + \mathcal{S}_S^{(\mathrm{str})} \\ \mathcal{S}_A^{(\mathrm{coh})} + \mathcal{S}_A^{(\mathrm{str})} \end{pmatrix}, \tag{11.294}$$

with the block operator

$$\mathbb{K}_{\mathrm{DRRC}} = \begin{pmatrix} K_{gg} & K_{gS} & K_{gA} \\ K_{Sg} & K_{SS} & K_{SA} \\ K_{Ag} & K_{AS} & K_{AA} \end{pmatrix}. \tag{11.295}$$

The off–diagonal entries $K_{gS}, K_{gA}, \ldots$ encode the constitutive mixing implied by the split $\Sigma_{\mu\nu} = g_{\mu\nu} + t_{\mu\nu}$ and its chain rule (cf. eqs. (10.94) and (10.101)). By the TT–null lemma for τ and the formal TT selection (cf. Lemma 10.17 and eq. (10.105)), the TT block is closed and decoupled, so (11.294) is confined to the non–TT subspace. The source terms are organized as *coherence* and *stress* functionals and are detailed in Section 11.8.

Derivation of the mixed–block linear system

Step A: Diagnostic realized reference configuration law and notation. Start from the diagnostic realized reference configuration (expectation–level) Einstein law

$$\bar{G}_{\mu\nu}[\bar{\Sigma}] + \Lambda\, \bar{\Sigma}_{\mu\nu} = \kappa\, \bar{T}^{\mathrm{total}}_{\mu\nu}, \qquad \text{(cf. Definition 11.106)} \tag{11.296}$$

on the realized geometry $(\mathcal{X}_q, \bar{\Sigma})$, with $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (Ch. 6). For compactness define the linearized geometric operator

$$\mathcal{E}_{\mu\nu}[\,\cdot\,] := \delta\bar{G}_{\mu\nu}[\,\cdot\,] + \Lambda\,(\,\cdot\,)_{\mu\nu}, \qquad \text{so that} \quad \mathcal{E}[h] = \kappa\, \delta\bar{T}^{\mathrm{total}}. \tag{11.297}$$

Step B: Linearize about the realized background. Perturb the background as $\bar{\Sigma} \mapsto \bar{\Sigma} + h$ with $h_{\mu\nu}$ small. Linearizing (11.296) gives

$$\delta\bar{G}_{\mu\nu}[h] + \Lambda\, h_{\mu\nu} = \kappa\, \delta\bar{T}^{\text{total}}_{\mu\nu} \quad \Longleftrightarrow \quad \mathcal{E}[h] = \kappa\, \delta\bar{T}^{\text{total}}. \tag{11.298}$$

Here $\delta\bar{G}_{\mu\nu}[h]$ is the *relational* (i.e. $\bar{\Sigma}$–compatible) metric variation of the Einstein tensor; it is a linear, background–covariant differential operator in h (Lichnerowicz–type on $(\mathcal{X}_q, \bar{\Sigma})$).

Step C: TT projector and non–TT complement. Let $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ be the certified TT projector on symmetric rank–2 tensors (Definition 10.3); write the complementary projector

$$\mathbb{P}_{\mathrm{nonTT}} := \mathrm{Id} - \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}. \tag{11.299}$$

Decompose any perturbation $h^{(\mathrm{tot})}_{\mu\nu}$ as

$$h := \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\, h^{(\mathrm{tot})}, \qquad \tau := \mathbb{P}_{\mathrm{nonTT}}\, h^{(\mathrm{tot})}, \qquad h^{(\mathrm{tot})} = h + \tau, \tag{11.300}$$

so h is TT (spin–2 radiative slice) and τ is the *TT–null* coherence slice (the τ–kairos; cf. Section 10.4).

Step D: Project the linearized law. Apply $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ and $\mathbb{P}_{\mathrm{nonTT}}$ to (11.298):

$$\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\, \mathcal{E}[h+\tau] = \kappa\, \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\, \delta\bar{T}^{\text{total}}, \tag{11.301a}$$

$$\mathbb{P}_{\mathrm{nonTT}}\, \mathcal{E}[h+\tau] = \kappa\, \mathbb{P}_{\mathrm{nonTT}}\, \delta\bar{T}^{\text{total}}. \tag{11.301b}$$

Since $\mathcal{E}[\,\cdot\,]$ is *linear* in its argument, $\mathcal{E}[h+\tau] = \mathcal{E}[h] + \mathcal{E}[\tau]$. Insert this into (11.301):

$$\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\, \mathcal{E}[h] + \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\, \mathcal{E}[\tau] = \kappa\, \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\, \delta\bar{T}^{\text{total}}, \tag{11.302a}$$

$$\mathbb{P}_{\mathrm{nonTT}}\, \mathcal{E}[h] + \mathbb{P}_{\mathrm{nonTT}}\, \mathcal{E}[\tau] = \kappa\, \mathbb{P}_{\mathrm{nonTT}}\, \delta\bar{T}^{\text{total}}. \tag{11.302b}$$

Step E: Define the mixed–block operators. Define the four linear maps (operators on symmetric rank–2 fields on $(\mathcal{X}_q, \bar{\Sigma})$)

$$\boxed{\begin{aligned} \mathcal{L}_{hh} &:= \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}} \circ \mathcal{E} \circ \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}, & \mathcal{L}_{h\tau} &:= \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}} \circ \mathcal{E} \circ \mathbb{P}_{\mathrm{nonTT}}, \\ \mathcal{L}_{\tau h} &:= \mathbb{P}_{\mathrm{nonTT}} \circ \mathcal{E} \circ \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}, & \mathcal{L}_{\tau\tau} &:= \mathbb{P}_{\mathrm{nonTT}} \circ \mathcal{E} \circ \mathbb{P}_{\mathrm{nonTT}}. \end{aligned}} \tag{11.303}$$

With these definitions, (11.302) becomes the coupled system

$$\mathcal{L}_{hh}\, h + \mathcal{L}_{h\tau}\, \tau = \kappa\, \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\, \delta\bar{T}^{\text{total}}, \tag{11.304a}$$

$$\mathcal{L}_{\tau h}\, h + \mathcal{L}_{\tau\tau}\, \tau = \kappa\, \mathbb{P}_{\mathrm{nonTT}}\, \delta\bar{T}^{\text{total}}. \tag{11.304b}$$

Step F: Assemble the matrix form. Collect (11.304) into the block matrix equation

$$\begin{pmatrix} \mathcal{L}_{hh} & \mathcal{L}_{h\tau} \\ \mathcal{L}_{\tau h} & \mathcal{L}_{\tau\tau} \end{pmatrix} \begin{pmatrix} h \\ \tau \end{pmatrix} = \kappa \begin{pmatrix} \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\, \delta\bar{T}^{\text{total}} \\ \mathbb{P}_{\mathrm{nonTT}}\, \delta\bar{T}^{\text{total}} \end{pmatrix}, \qquad \begin{cases} h := \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\, h, \\ \tau := \mathbb{P}_{\mathrm{nonTT}}\, h \end{cases}$$

which is eq. (11.293).

Step G: TT–purity and physical interpretation. By the TT–purity results in the weave (Chapter 11), the TT projector annihilates the spin–1 ledger and the τ–sector is TT–null. Consequently:

- $\mathcal{L}_{hh}$ is the *radiative* (spin–2) operator on h (the graviton/Apeiron$^{[\mathrm{TT}]}$ channel);
- $\mathcal{L}_{\tau\tau}$ governs the *non–TT* coherence dynamics (typed scalar/pseudoscalar/vector);
- $\mathcal{L}_{h\tau}$ and $\mathcal{L}_{\tau h}$ encode the *mixed response kernels*, responsible for non–TT dressing/sourcing of the metric and backreaction of h into the coherence sector.

Step H: Comments on sources and causality. The source split on the right–hand side uses the same projectors applied to the linearized diagnostic realized reference configuration source $\delta\bar{T}^{\mathrm{total}}$ (cf. Definition 9.358 and Section 9.18.12). On certified plateaux $\mathfrak{P}$, all blocks admit $\bar{\Sigma}$–causal retarded/advanced Green operators, so the coupled system (11.293) is well–posed with propagation inside the $\bar{\Sigma}$ null cones.

Coherence–stress bookkeeping on the RHS

Source split and chronogeometric pieces. The total diagnostic realized reference configuration source in Definition 11.106 is decomposed as

$$\bar{T}^{\mathrm{total}}_{\mu\nu} = \bar{T}^{\mathrm{matter}}_{\mu\nu} + \bar{T}^{\mathrm{int}}_{\mu\nu} + \bar{T}^{\mathrm{coh}}_{\mu\nu}[\bar{t};\,\mathrm{typed}], \tag{11.305}$$

where $\bar{T}^{\mathrm{coh}}_{\mu\nu}$ collects the *coherence* (temporal) contributions originating from the realized temporal tensor $\bar{t}_{\mu\nu}$ and from the certified non–TT typed amplitudes of the weave (Chapter 11). Write the symmetric/antisymmetric split

$$\bar{t}_{\mu\nu} = S_{\mu\nu} + A_{\mu\nu}, \qquad S_{\mu\nu} := \bar{t}_{(\mu\nu)}, \quad A_{\mu\nu} := \bar{t}_{[\mu\nu]}. \tag{11.306}$$

Quadratic coherence tensor and Relational NEC. As used in Section 11.6, the canonical quadratic coherence tensor reads

$$\mathcal{Q}_{\mu\nu}[S,A] := S_{\mu\alpha}\,S_{\nu}{}^{\alpha} + A_{\mu\alpha}\,A_{\nu}{}^{\alpha} - \frac{1}{d}\,\bar{g}^{\mathrm{Lor}}_{\mu\nu}\left(S_{\alpha\beta}S^{\alpha\beta} + A_{\alpha\beta}A^{\alpha\beta}\right), \tag{11.307}$$

so that, for any null k^{μ}, one has $k^{\mu}k^{\nu}\mathcal{Q}_{\mu\nu} \geq 0$ (Relational NEC). The coherence stress is organized so that its null contractions are controlled by $\mathcal{Q}_{\mu\nu}$.

Typed bookkeeping: scalar, pseudoscalar, and vector (CK) pieces. Let the certified typed amplitudes of the TT–complement be $\{\hat{\varphi}_b, \hat{\varphi}_L, \tilde{\varphi}, \hat{K}^{(\tau)}_{\lambda}\}$ with $\lambda = \pm$ (cf. Chapter 11). Their diagnostic realized reference configuration expectations (or bilinears for fluctuations) assemble $\bar{T}^{\mathrm{coh}}_{\mu\nu}$ as follows:

$$\bar{T}^{\mathrm{coh}}_{\mu\nu} = \left(\bar{T}^{\mathrm{coh}}\right)^{(b)}_{\mu\nu} + \left(\bar{T}^{\mathrm{coh}}\right)^{(L)}_{\mu\nu} + \left(\bar{T}^{\mathrm{coh}}\right)^{(p)}_{\mu\nu} + \left(\bar{T}^{\mathrm{coh}}\right)^{(\mathrm{CK})}_{\mu\nu}, \tag{11.308}$$

with each term constrained by the weave typing (orthogonality and parity) and evaluated on $(\mathcal{X}_q, \bar{\Sigma})$ with transport by $\bar{\nabla}$.

Vector (chrono–kymon) contribution. For the vector type (CK) one uses the gauge–invariant field strength $\mathcal{F}^{(\tau)}_{\mu\nu} := 2\,\bar{\nabla}_{[\mu}K^{(\tau)}_{\nu]}$ and obtains the Maxwell–like symmetric stress

$$\left(\bar{T}^{\mathrm{coh}}\right)^{(\mathrm{CK})}_{\mu\nu} = \gamma_\tau \left(\mathcal{F}^{(\tau)}_{\mu\alpha}\,\mathcal{F}^{(\tau)\,\alpha}{}_{\nu} - \frac{1}{4}\,\bar{g}^{\mathrm{Lor}}_{\mu\nu}\,\mathcal{F}^{(\tau)}_{\alpha\beta}\,\mathcal{F}^{(\tau)\,\alpha\beta}\right), \tag{11.309}$$

where γ_τ is a diagnostic realized reference configuration–effective coefficient. The antisymmetric temporal tensor is driven by the CK strength via $A_{\mu\nu} \propto \mathcal{F}^{(\tau)}_{\mu\nu}$ (cf. the vector typed reconstruction), so that (11.309) contributes to $\mathcal{Q}_{\mu\nu}$ and hence obeys the Relational NEC (null positivity).

Scalar and pseudoscalar typed contributions. For the scalar–typed amplitudes φ_b, φ_L and the pseudoscalar $\tilde{\varphi}$, one may use the diagnostic realized reference configuration–canonical symmetric bilinears (no TT content, weave–typed and parity–clean):

$$\left(\bar{T}^{\mathrm{coh}}\right)^{(b)}_{\mu\nu} = \alpha_b\,\bar{\nabla}_\mu\varphi_b\,\bar{\nabla}_\nu\varphi_b - \frac{\alpha_b}{2}\,\bar{g}^{\mathrm{Lor}}_{\mu\nu}\,(\bar{\nabla}\varphi_b)^2 - \bar{g}^{\mathrm{Lor}}_{\mu\nu}\,V_b(\varphi_b), \tag{11.310a}$$

$$\left(\bar{T}^{\mathrm{coh}}\right)^{(L)}_{\mu\nu} = \alpha_L\,\bar{\nabla}_\mu\varphi_L\,\bar{\nabla}_\nu\varphi_L - \frac{\alpha_L}{2}\,\bar{g}^{\mathrm{Lor}}_{\mu\nu}\,(\bar{\nabla}\varphi_L)^2 - \bar{g}^{\mathrm{Lor}}_{\mu\nu}\,V_L(\varphi_L), \tag{11.310b}$$

$$\left(\bar{T}^{\mathrm{coh}}\right)^{(p)}_{\mu\nu} = \alpha_p\,\bar{\nabla}_\mu\tilde{\varphi}\,\bar{\nabla}_\nu\tilde{\varphi} - \frac{\alpha_p}{2}\,\bar{g}^{\mathrm{Lor}}_{\mu\nu}\,(\bar{\nabla}\tilde{\varphi})^2 - \bar{g}^{\mathrm{Lor}}_{\mu\nu}\,\tilde{V}(\tilde{\varphi}), \tag{11.310c}$$

with diagnostic realized reference configuration–effective couplings $\alpha_{b,L,p} \geq 0$ and (typed) potentials $V_b, V_L, \tilde{V}$. For any null k^μ,

$$k^\mu k^\nu \left(\bar{T}^{\mathrm{coh}}\right)^{(X)}_{\mu\nu} = \alpha_X \left(k^\mu\bar{\nabla}_\mu\phi_X\right)^2 \;\geq\; 0, \qquad X \in \{b, L, p\}, \quad \phi_X \in \{\varphi_b, \varphi_L, \tilde{\varphi}\}, \tag{11.311}$$

so the scalar/pseudoscalar typed pieces also satisfy the Relational NEC.[3]

Linearized source split and projections. For perturbations about a realized background, split the coherence source into background and fluctuation pieces and project with the certified TT/compl. projectors:

$$\delta\bar{T}^{\mathrm{coh}}_{\mu\nu} = \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\bar{T}^{\mathrm{coh}}_{\mu\nu} + \mathbb{P}_{\mathrm{nonTT}}\,\delta\bar{T}^{\mathrm{coh}}_{\mu\nu}, \tag{11.312}$$

where the first term can *drive* radiative geometry (via $\mathcal{L}_{hh}$) and the second term couples to the non–TT sector (via $\mathcal{L}_{\tau\tau}$), with cross–coupling through the mixed blocks $\mathcal{L}_{h\tau}, \mathcal{L}_{\tau h}$ in (11.293). The background (coherent) piece contributes at order $\mathcal{O}(\varepsilon)$ to the balance equation (11.292).

Scalarisation and universality of coupling. In the spectral limit (cf. Sections 11.5.7 and 11.5.8), the emergent coupling tensor $\mathcal{C}_{\mu\nu}{}^{\rho\sigma}$ scalarises to $\kappa_{\mathrm{eff}}\,\Pi_{\mu\nu}{}^{\rho\sigma}$ so that each typed coherence term in (11.308) couples with the same diagnostic realized reference configuration–universal κ_{eff}, consistent with the recovery of GR and with the equivalence–principle bounds on anisotropies.

[3] Parity forbids linear mixing between (p) and (b, L) at quadratic order; cross–type bilinears vanish by the weave typing (orthogonality of certified projectors).

Summary. The coherence contribution to the diagnostic realized reference configuration source is a typed, covariant, and NEC–respecting sum of (i) a Maxwell–like vector (CK) stress (11.309), (ii) scalar and pseudoscalar symmetric stresses (11.310), all assembled on $(\mathcal{X}_q, \bar{\Sigma})$ with $\bar{\nabla}$ and the weave projectors. It participates in the linearized dynamics via the projected right–hand side of (11.293) and reduces to universal coupling in the spectral limit.

Non–TT law residual and visibility (diagnostic realized reference configuration diagnostics)

In parallel with the TT law visibility Definition 11.6, define the *non–TT law residual* by projecting the diagnostic realized reference configuration Einstein law to the TT–complement:

$$\boxed{\mathbf{R}^{[\mathrm{nonTT}]} := \mathbb{P}_{\mathrm{nonTT}}\Big[\bar{G}_{\mu\nu}[\bar{\Sigma}] + \Lambda\,\bar{\Sigma}_{\mu\nu} - \kappa\,\bar{T}^{\mathrm{total}}_{\mu\nu}\Big],} \tag{11.313}$$

and the corresponding *non–TT law visibility*

$$\Lambda^{[\mathrm{nonTT}]}_{\mathrm{law}} := 1 - \frac{\|\mathbf{R}^{[\mathrm{nonTT}]}\|_{\mathcal{I}}}{\|\cdot\|_{\mathrm{ref}}}, \qquad \Lambda^{[\mathrm{nonTT}]}_{\mathrm{law}} \in [0,1], \tag{11.314}$$

with the realized weave configuration norm $\|\cdot\|_{\mathcal{I}}$ and the acceptance reference $\|\cdot\|_{\mathrm{ref}}$ as in Definition 11.6 and Section 7.3.9. By construction, $\Lambda^{[\mathrm{nonTT}]}_{\mathrm{law}}$ is Nielsen–flat on $\mathfrak{P}$ (same commutation properties as in the TT case).

Emergent coupling tensor and scalarisation with coherence

Writing the diagnostic realized reference configuration law in response form (cf. Section 11.5.6),

$$\bar{G}_{\mu\nu}(x) = \int \mathscr{K}_{\mu\nu}{}^{\rho\sigma}(x,x')\,\bar{T}^{\mathrm{total}}_{\rho\sigma}(x')\,\mathrm{d}^d x',$$

the local spectral limit yields the emergent coupling tensor $\mathcal{C}_{\mu\nu}{}^{\rho\sigma}$, which *scalarises* (cf. Section 11.5.7) to $\kappa_{\mathrm{eff}}\,\Pi_{\mu\nu}{}^{\rho\sigma}$ under the same anisotropy bounds. In that limit, the *typed* coherence contributions in (9.358) couple universally with κ_{eff}, ensuring consistency with Section 11.5.8.

Witness predicates and diagnostic realized reference configuration certification

Law–vs–state diagnostics for the non–TT sector adopt the same witness framework as Definitions 8.4 to 8.6, with predicates specialized to the typed τ amplitudes and to the mixed blocks in (11.293). Acceptance is granted when $\Lambda^{[\mathrm{nonTT}]}_{\mathrm{law}} \approx 1$ and the typed witnesses saturate their diagnostic realized reference configuration thresholds, all on a certified plateau $\mathfrak{P}$.

Summary. The non–TT coherence sector (the τ–kairos and, inside it, the chrono–kymon) enters the diagnostic realized reference configuration Einstein law (i) on the LHS through background shifts of $\bar{t}_{\mu\nu}$ hence $\bar{\Sigma}$, (ii) on the RHS via the coherence–stress contribution eq. (11.170) (with the vector piece exemplified by eq. (11.309)), and (iii) at the linearized level through the mixed response blocks in eq. (11.293). TT–purity (Chapter 11) guarantees that the spin–2 radiative sector remains uniquely determined by the TT block, while coherence physics is certified by the non–TT visibility eq. (11.314) and witness predicates.

11.8.1 Linear–response Quantization of the Non–TT sector and Optional BEH Mass for Chrono-Kymons

Quadratic diagnostic realized reference configuration (DRCC) functional for the non–TT sector. Linearizing the diagnostic realized reference configuration law Definition 11.106 and using the TT/complement split (cf. Section 10.4 and (11.293)) yields, to quadratic order on $\mathfrak{P}$,

$$S^{(2)}_{\mathrm{DRCC}}[h,\tau] = \frac{1}{2}\int h\,\mathcal{L}_{hh}h + \int h\,\mathcal{L}_{h\tau}\tau + \frac{1}{2}\int \tau\,\mathcal{L}_{\tau\tau}\tau - \kappa\int\Big(h\,\Pi^{[\mathrm{TT}]}\delta\bar{T}^{\mathrm{total}} + \tau\,\mathbb{P}_{\mathrm{nonTT}}\delta\bar{T}^{\mathrm{total}}\Big), \tag{11.315}$$

where all contractions and covariant derivatives are with $\bar{\Sigma}$ and $\bar{\nabla}$. The operator $\mathcal{L}_{\tau\tau}$ is the non–TT kinetic operator; $\mathcal{L}_{h\tau}$, $\mathcal{L}_{\tau h}$ encode the certified mixed response kernels.

How (11.315) is obtained (derivation). Start from the typed TT/complement split of metric fluctuations on the diagnostic realized reference configuration $\mathfrak{P}$:

$$\delta g_{\mu\nu} = \underbrace{\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta g_{\mu\nu}}_{h_{\mu\nu}} + \underbrace{\big(\mathbb{1} - \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\big)\delta g_{\mu\nu}}_{(\delta g)_{\overline{\mathrm{TT}}}},$$

$$\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\hat{\tau}_{\mu\nu} = 0 \quad \text{(cf. eq. (10.105) and Lemma 10.17).}$$

Collect the non–TT fields into the typed multiplet

$$\tau := \Big((\delta g)_{\overline{\mathrm{TT}}};\ \tau_{\mathrm{scalar}}\ ;\ K^{(\tau)}_{\mu}\Big)^{\top}.$$

The linearized diagnostic realized reference configuration law in mixed–block form (cf. eqs. (11.293) to (11.295)) reads

$$\begin{pmatrix} K_{hh} & K_{h\tau} \\ K_{\tau h} & K_{\tau\tau} \end{pmatrix}\begin{pmatrix} h \\ \tau \end{pmatrix} = \begin{pmatrix} \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\bar{T}^{\mathrm{total}} \\ \mathbb{P}_{\mathrm{nonTT}}\,\delta\bar{T}^{\mathrm{total}} \end{pmatrix}, \tag{11.316}$$

where $K_{hh} := \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}K_{gg}\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$, $K_{h\tau} := \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}(K_{gS}\oplus K_{gA})\mathbb{P}_{\mathrm{nonTT}}$, $K_{\tau h} := \mathbb{P}_{\mathrm{nonTT}}(K_{Sg}\oplus K_{Ag})\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$, and $K_{\tau\tau} := \mathbb{P}_{\mathrm{nonTT}}\left(\begin{smallmatrix} K_{SS} & K_{SA} \\ K_{AS} & K_{AA}\end{smallmatrix}\right)\mathbb{P}_{\mathrm{nonTT}}$. Here $\mathbb{P}_{\mathrm{nonTT}}$ projects onto the non–TT ledger (metric complement $\oplus$ τ–kairos channels), and the off–diagonal blocks encode the constitutive mixing implied by $\Sigma_{\mu\nu} = g_{\mu\nu} + t_{\mu\nu}$ and its chain rule (cf. eqs. (10.94) and (10.101)).

Define the diagnostic realized reference configuration pairing $\langle X, Y\rangle_{\mathfrak{P}} := \int\sqrt{-\det\bar{g}^{\mathrm{Lor}}}\,XY$ (all indices contracted with $\bar{\Sigma}$, covariant derivatives with $\bar{\nabla}$). Then the quadratic diagnostic

realized reference configuration functional whose Euler–Lagrange equations reproduce (11.316) is

$$S^{(2)}_{\mathrm{DRRC}}[h,\tau] = \frac{1}{2}\langle h,\, K_{hh}h\rangle_{\mathfrak{P}} + \langle h,\, K_{h\tau}\tau\rangle_{\mathfrak{P}} + \frac{1}{2}\langle \tau,\, K_{\tau\tau}\tau\rangle_{\mathfrak{P}} - \kappa\Big(\langle h,\, \Pi^{[\mathrm{TT}]}\delta\bar{T}^{\mathrm{total}}\rangle_{\mathfrak{P}} + \langle \tau,\, \mathbb{P}_{\mathrm{nonTT}}\delta\bar{T}^{\mathrm{total}}\rangle_{\mathfrak{P}}\Big). \tag{11.317}$$

Writing $K_{hh} \equiv \mathcal{L}_{hh}$, $K_{h\tau} \equiv \mathcal{L}_{h\tau}$ and $K_{\tau\tau} \equiv \mathcal{L}_{\tau\tau}$ gives precisely (11.315). BRST/tag hygiene from Chapter 10 guarantees these operators are symmetric on the certified band, so the variational equations from (11.317) are equivalent to the linear law (cf. Lemma 9.54).

Remark 11.156 (TT closure)**.** By the TT–null lemma and formal TT selection (cf. Lemma 10.17 and eq. (10.105)), the certified mixed kernels obey $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}(K_{gS}, K_{gA}) = 0$ on $\mathfrak{P}$, so $\mathcal{L}_{h\tau}$ *vanishes* on the diagnostic realized reference configuration. We keep the h–τ term in (11.315) as a bookkeeping slot for possible off–diagnostic realized reference configuration or higher–order effects, but at the certified linear level the TT block is dynamically closed.

Typed field content inside τ. Decompose $\tau_{\mu\nu}$ with the certified typed projectors (Chapter 11):

$$\tau_{\mu\nu} = S^{(b)}_{\mu\nu}[\varphi_b] + S^{(L)}_{\mu\nu}[\varphi_L] + \tilde{S}^{(p)}_{\mu\nu}[\tilde{\varphi}] + A^{(\lambda)}_{\mu\nu}[K^{(\tau)}_{\lambda}], \qquad \lambda = \pm, \tag{11.318}$$

with *vector* (chrono–kymon, CK) potential $K^{(\tau)}_{\mu}$ entering only through its gauge–invariant strength

$$\mathcal{F}^{(\tau)}_{\mu\nu} := 2\,\bar{\nabla}_{[\mu}K^{(\tau)}_{\nu]}\,. \tag{11.319}$$

Massless CK: Maxwell–type quadratic functional (no SSB needed). In the absence of explicit internal SSB acting on the CK type, the vector contribution to the DRRC quadratic functional $S^{(2)}_{\mathrm{DRRC}}$ takes the $\bar{\Sigma}$–Maxwell form (cf. Section 11.8):

$$S^{(2)}_{\mathrm{CK},0} = -\frac{\gamma_\tau}{4}\int \sqrt{-\det \bar{g}^{\mathrm{Lor}}}\; \mathcal{F}^{(\tau)}_{\mu\nu}\,\mathcal{F}^{(\tau)\,\mu\nu}, \tag{11.320}$$

which is invariant under $K^{(\tau)}_{\mu} \mapsto K^{(\tau)}_{\mu} + \bar{\nabla}_{\mu}\chi$. A covariant gauge fixing (BRST–compatible on $\mathfrak{P}$) gives the well–posed operator

$$\bar{\nabla}_{\mu}\mathcal{F}^{(\tau)\,\mu\nu} = J^{(\tau)\,\nu}, \qquad J^{(\tau)\,\nu} := \frac{1}{\gamma_\tau}\,\frac{\delta S^{(2)}_{\mathrm{DRRC}}}{\delta K^{(\tau)}_{\nu}}\Big|_{\text{sources, mixes}}, \tag{11.321}$$

where "DRRC" denotes the diagnostic realized reference configuration of Definition 9.185. The corresponding solutions have $\bar{\Sigma}$–causal retarded/advanced Green operators (inheritance of the DRRC causal cones). Quantization proceeds by canonical ETCR on the band (Chapter 11), or equivalently by constructing the CK two–point function from the inverse of the gauge–fixed kinetic operator. *No SSB is required* to have a propagating CK quantum; it is a standard linear–response mode in the TT–complement.

Scalar/pseudoscalar non–TT modes (no SSB needed). The scalar φ_b, φ_L and pseudoscalar $\tilde{\varphi}$ pieces enter with standard covariant quadratic functionals on $(\mathcal{X}_q, \bar{\Sigma})$ (cf. eq. (11.310)), yielding well–posed hyperbolic operators and causal propagators. Their quantization is identical in spirit to the CK case and likewise does not require SSB.

11.8.2 BEH of the Chrono-Kymon

Constitutive, typed origin; independence from Tantrakyma SSB/BEH. The Chrono–Kymon (CK) does not originate from Tantrakyma spontaneous symmetry breaking or from a Brout–Englert–Higgs reorganization. Rather, it is defined by the constitutive law as the vector-typed, non-TT response of the chronogeometric observable to substrate fluctuations. The fluctuations $(\delta\Phi, \delta\Theta)$ are first pushed forward by the constitutive map; see eqs. (10.18) and (10.19). The CK slot is then selected with the weave's certified projectors; see Section 10.4 and Definitions 10.3 and 10.4.

Constitutive pushforward and vector selection (Ch. 11). Start from the constitutive map

$$\Sigma_{\mu\nu}[\Phi,\Theta] \;=\; g_{\mu\nu}[\Phi] \;+\; t_{\mu\nu}[\Phi,\Theta], \qquad (\bar{\Phi},\bar{\Theta}) \text{ realized background.}$$

Linearizing about $(\bar{\Phi}, \bar{\Theta})$ yields the chain rule

$$\delta\Sigma_{\mu\nu} \;=\; (D_\Phi g)_{\mu\nu}\cdot\,\delta\Phi \;+\; (D_\Phi t)_{\mu\nu}\cdot\,\delta\Phi \;+\; (D_\Theta t)_{\mu\nu}\cdot\,\delta\Theta.$$

Let $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ be the certified TT projector and $\mathbb{P}_{\mathrm{nonTT}} := \mathrm{Id} - \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ the TT–complement (Ch. 11, τ–construction). Let $\mathsf{A}_\mu{}^{\rho\sigma}$ denote the vector–typed selector onto the CK slot (the $A^{(\lambda)}$ piece of the typed decomposition; cf. Definition 10.4). The CK carrier is then

$$K_\mu^{(\tau)} \;:=\; \mathsf{A}_\mu{}^{\rho\sigma}\left(\mathbb{P}_{\mathrm{nonTT}}\delta\Sigma_{\rho\sigma}\right) \;=\; \underbrace{\mathsf{A}\,\mathbb{P}_{\mathrm{nonTT}}\big((D_\Phi g)+(D_\Phi t)\big)}_{J_\Phi^{(A)}}\,\delta\Phi \;+\; \underbrace{\mathsf{A}\,\mathbb{P}_{\mathrm{nonTT}}(D_\Theta t)}_{J_\Theta^{(A)}}\,\delta\Theta. \tag{11.322}$$

Thus the CK is the *linear image* of $(\delta\Phi, \delta\Theta)$ under the constitutive Jacobians $J_\Phi^{(A)}, J_\Theta^{(A)}$ after non–TT and vector–type selection. Its existence does *not* presuppose SSB or BEH of Θ; it is present at the level of linear response and typed projection.

Massless CK and Maxwell structure (no SSB required). Define the gauge–like strength $\mathcal{F}_{\mu\nu}^{(\tau)} := 2\,\bar{\nabla}_{[\mu}K_{\nu]}^{(\tau)}$. On certified plateaux, the quadratic CK functional reduces in the massless limit to a Maxwell–type form with two physical helicities $\lambda = \pm$ (spin–1), as encoded by the linearized diagnostic realized reference configuration law and its typed block structure (cf. eqs. (11.293) to (11.295)). The antisymmetric temporal imprint

$$\Delta t_{[\mu\nu]}^{(\tau)} \;\propto\; \mathcal{F}_{\mu\nu}^{(\tau)} \tag{11.323}$$

follows from the typed geometry (cf. eqs. (11.288) and (11.289)) and does not rely on a Tantrakyma Goldstone origin. TT purity ensures no leakage into the spin–2 tensor channel.

τ–sector BEH for CK (independent of Tantrakyma). A CK mass arises only if an *internal symmetry acting on the CK representation* is gauged and spontaneously broken *in the CK/τ sector* itself (Ch. 12). The quadratic functional upgrades to Proca:

$$S^{(2)}_{\mathrm{CK},m} \;=\; -\frac{\gamma_\tau}{4}\int\sqrt{-\det\bar{g}^{\mathrm{Lor}}}\;\mathcal{F}^{(\tau)}_{\mu\nu}\,\mathcal{F}^{(\tau)\,\mu\nu} \;+\; \frac{m^2_{\mathrm{CK}}}{2}\int\sqrt{-\det\bar{g}^{\mathrm{Lor}}}\;K^{(\tau)}_{\mu}K^{(\tau)\,\mu}, \tag{11.324}$$

with mass determined by the typed order parameter $\varphi^{[\tau]}$ and the algebra metric G^{IJ},

$$m^2_{\mathrm{CK}} \;=\; g^2\left\langle \mathcal{R}_\tau(T_I)\,\varphi^{[\tau]},\,\mathcal{R}_\tau(T_J)\,\varphi^{[\tau]}\right\rangle G^{IJ}, \tag{11.325}$$

and equations of motion (Proca form)

$$\bar{\nabla}_\mu\mathcal{F}^{(\tau)\,\mu\nu} + m^2_{\mathrm{CK}}\,K^{(\tau)\,\nu} \;=\; J^{(\tau)\,\nu}, \qquad \bar{\nabla}_\nu K^{(\tau)\,\nu} = 0 \;\;\text{(on shell)}. \tag{11.326}$$

This BEH mechanism is *optional* and strictly *τ–sector*; it neither relies on nor is sourced by SSB/BEH in the Tantrakyma field.

Tantrakyma's role (background and sources, not creation). Tantrakyma enters only through the constitutive Jacobian $J^{(A)}_{\Theta} = \mathsf{A}\,\mathbb{P}_{\mathrm{nonTT}}(D_\Theta t)$ in (11.322). If the Tantrakyma sector undergoes SSB/BEH, it can (i) shift the realized background $(\bar{\Phi},\bar{\Theta})$ and hence modify CK propagation via the background change $\bar{\Sigma}\mapsto\bar{\Sigma}'=\bar{\Sigma}+\Delta\Sigma$ (cf. eqs. (11.290) and (11.291)), and (ii) contribute typed sources on the RHS of the mixed–block system (cf. Section 11.8). Neither effect creates the CK degree of freedom: the CK already exists as the vector–typed, non–TT response variable of the chronogeometric map.

Conclusion. The CK originates from the typed, non–TT linear response of $\Sigma_{\mu\nu}$; it does not require global SSB or BEH of the Tantrakyma field for its existence. When present, a CK mass gap comes from an *internal τ–sector* BEH acting on the CK representation (cf. eqs. (11.324) to (11.326)), while TT purity guarantees that the spin–2 graviton sector remains closed and unaffected (cf. eq. (11.293)).

Internal BEH: Proca deformation for CK. If an internal symmetry acts on the CK type through a representation $\mathcal{R}_\tau$ and is spontaneously broken on $\mathfrak{P}$ (cf. Sections 6.6 and 9.17.3), the quadratic CK sector admits a *Proca* deformation

$$S^{(2)}_{\mathrm{CK},m} = -\frac{\gamma_\tau}{4}\int\sqrt{-\det\bar{g}^{\mathrm{Lor}}}\;\mathcal{F}^{(\tau)}_{\mu\nu}\,\mathcal{F}^{(\tau)\,\mu\nu} + \frac{m^2_{\mathrm{CK}}}{2}\int\sqrt{-\det\bar{g}^{\mathrm{Lor}}}\;K^{(\tau)}_{\mu}K^{(\tau)\,\mu}, \qquad (\gamma_\tau>0),$$

with

$$m^2_{\mathrm{CK}} = \; g^2\left\langle\,\mathcal{R}_\tau(T_I)\,\varphi^{[\tau]}\,,\,\mathcal{R}_\tau(T_J)\,\varphi^{[\tau]}\,\right\rangle G^{IJ},$$

where g is the internal gauge coupling, $\varphi^{[\tau]} := \Pi^{[\tau]}_{\Phi\Phi}\,\bar{\Phi}$ is the τ-typed order parameter on the modal fiber (cf. Section 9.17.3), $\{T_I\}$ are the broken generators, and G^{IJ} is the inverse of the Lie-algebra metric G_{IJ} from the gauge kinetic term (cf. Section 9.18.8). The pairing $\langle\cdot,\cdot\rangle$ is the diagnostic realized reference configuration/modal-fiber inner product, so the

mass construction is "fiber pairing × algebra metric," as in the connection–vector case in Chapter 10.

[4]

The massive CK obeys

$$\bar{\nabla}_\mu \mathcal{F}^{(\tau)\,\mu\nu} + m_{\mathrm{CK}}^2\, K^{(\tau)\,\nu} = J^{(\tau)\,\nu}.$$

Assuming the typed source is conserved, $\bar{\nabla}_\nu J^{(\tau)\,\nu} = 0$, taking the divergence of (11.326) yields the on–shell transversality

$$\bar{\nabla}_\nu K^{(\tau)\,\nu} = 0 \qquad \text{(on shell)}. \tag{11.327}$$

This BEH deformation alters the *dispersion* of CK quanta but does not affect TT purity of the radiative (spin–2) sector: $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\hat{\tau}_{\mu\nu} = 0$ and $\hat{h}_{\mu\nu} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}_{\mu\nu}$ (cf. Lemma 10.17, eq. (10.105), and Section 9.18.7).

Mixing with geometry and energy conditions. Couplings to h enter only through the certified mixed blocks $\mathcal{L}_{h\tau}, \mathcal{L}_{\tau h}$ in (11.293); TT hygiene (Chapter 11) forbids leakage of CK into tensor polarizations. The coherence stress built from $\mathcal{F}^{(\tau)}$ (and from scalar/pseudoscalar typed bilinears) satisfies the Relational NEC (Section 11.6); in particular, for any $\bar{\Sigma}$-null k^μ, $k^\mu k^\nu (\bar{T}^{\mathrm{coh}})_{\mu\nu}^{(\mathrm{CK})} \geq 0$ (cf. eq. (11.309)).

Summary. All non–TT typed modes—including the vector CK—are bona fide quantum fields obtained by *linear–response quantization* of the TT–complement on $\mathfrak{P}$. They require no SSB to exist. If an internal symmetry is broken in the sector acting on CK, a BEH mechanism *optionally* promotes CK to a Proca field with mass m_{CK}, without disturbing the TT graviton sector or its polarizations.

Chrono–Kymon (CK) in the weave. The Chrono–Kymon is the *vector typed* register inside the τ–kairos—the TT–complement of chronogeometric fluctuations (cf. Chapter 11). Kinematically it is represented by a potential $K_\mu^{(\tau)}$ with gauge–invariant field strength

$$\mathcal{F}_{\mu\nu}^{(\tau)} := 2\,\bar{\nabla}_{[\mu} K_{\nu]}^{(\tau)}\,,$$

and it is TT–null by construction, so it does not carry spin–2 radiation nor alter the graviton's polarization content. In the weave typing, τ splits into scalar/pseudoscalar/vector pieces; the CK furnishes the antisymmetric temporal imprint via $\mathcal{F}^{(\tau)}$ (cf. Section 11.8).

Linear–response (no SSB): massless CK and diagnostic realized reference configuration quantization. On a certified plateau $\mathfrak{P}$, the *linear–response* CK dynamics (without any symmetry breaking) take the Maxwell form on $(\mathcal{X}_q, \bar{\Sigma})$:

$$S_{\mathrm{CK},0}^{(2)} = -\frac{\gamma_\tau}{4}\int \sqrt{-\det \bar{g}^{\mathrm{Lor}}}\; \mathcal{F}_{\mu\nu}^{(\tau)}\,\mathcal{F}^{(\tau)\,\mu\nu}, \qquad \bar{\nabla}_\mu \mathcal{F}^{(\tau)\,\mu\nu} = J^{(\tau)\,\nu},$$

with BRST–compatible gauge fixing on $\mathfrak{P}$. Quantization yields a *massless* spin–1 quantum (two transverse polarizations) in the TT–complement; no SSB is required for the existence of the CK quantum (cf. eq. (11.293) for the mixed–block coupling to geometry).

[4] For a gauge–invariant presentation, introduce a covariant Stueckelberg field σ so that $K_\mu^{(\tau)} \mapsto K_\mu^{(\tau)} + \bar{\nabla}_\mu \sigma / m_{\mathrm{CK}}$, keeping BRST structure manifest on $\mathfrak{P}$ (cf. Section 9.17.3).

Internal BEH (optional): Proca upgrade and dispersion change. If an *internal* gauge symmetry acting on the CK representation $\mathcal{R}_\tau$ is spontaneously broken on $\mathfrak{P}$ (see Section 6.6), the CK sector undergoes a Brout–Englert–Higgs upgrade to a Proca field: spectrally, the one–particle pole moves from $k^2 = 0$ to $k^2 = m_{\mathrm{CK}}^2$, and the physical polarizations reorganize from two transverse to three (two transverse + one longitudinal) in $d = 4$. In the compact diagnostic realized reference configuration notation from your notes, this transition is summarized by

$$\text{massless: } S^{(2)}_{\mathrm{CK},0} = -\frac{\gamma_\tau}{4}\int \sqrt{-\det \bar{g}^{\mathrm{Lor}}}\, \mathcal{F}^{(\tau)}_{\mu\nu}\mathcal{F}^{(\tau),\mu\nu}$$

$$\rightsquigarrow$$

$$\text{massive (BEH): } S^{(2)}_{\mathrm{CK},m} = S^{(2)}_{\mathrm{CK},0} + \frac{m_{\mathrm{CK}}^2}{2}\int \sqrt{-\det \bar{g}^{\mathrm{Lor}}}\, K^{(\tau)}_{\mu}K^{(\tau),\mu}$$

and the corresponding Proca equations on $(\bar{\Sigma}, \bar{\nabla})$,

$$\bar{\nabla}_\mu \mathcal{F}^{(\tau)\,\mu\nu} + m_{\mathrm{CK}}^2\, K^{(\tau)\,\nu} = J^{(\tau)\,\nu}, \qquad \bar{\nabla}_\nu K^{(\tau)\,\nu} = 0 \ \text{ (on shell)}.$$

This BEH deformation changes *dispersion and degrees of freedom* in the *non–TT* sector only; TT–purity (Chapter 11) ensures the graviton (Apeiron$^{[\mathrm{TT}]}$) remains the unique spin–2 radiative slice.

The **Chrono-Kymon (CK)** starts life as the vector-typed piece inside the τ–kairos—the TT-complement of chronogeometric fluctuations defined with the certified projector ($\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$) (cf. Section 10.4 and Definition 10.3). In that unbroken phase, it is a massless spin-1 field with two physical helicities ($\lambda = \pm$). Nothing in this construction touches the spin-2 radiative (graviton) slice, which is isolated by TT-purity. When an internal gauge symmetry acting on the CK representation is spontaneously broken on the diagnostic realized reference configuration ($\mathfrak{P}$) (via the operator-bundle connection; cf. Section 6.6), the CK sector undergoes the familiar *Brout–Englert–Higgs (BEH) reorganization*: the quadratic functional upgrades from the Maxwell form to a Proca form (see eqs. (11.320) and (11.324)). Physically, the would-be Goldstone is "eaten" and a mass gap ($m_{\mathrm{CK}} > 0$) appears. This change affects the degrees of freedom—the massive CK has three physical polarizations ($\lambda = \pm, 0$) in 4D—but it does not change the spin. Representation-theoretically, you move from the massless little group to the massive one (SO(3)), so we still have a spin-1 field, now with an added longitudinal mode. Dynamically, the Proca equations on $(\bar{\Sigma}, \bar{\nabla})$ together with the on-shell constraint make this explicit (cf. eq. (11.326)). The extra longitudinal polarization is a hallmark of massiveness, not a change of spin. Throughout, TT-purity ensures CK remains in the non-TT ledger; it interacts with geometry only through the certified mixed blocks of the linearized diagnostic realized reference configuration law (cf. eq. (11.293)), so the graviton's spin-2 sector and its two tensor polarizations remain untouched.

In the massless phase, the chrono-kymon (CK) is a *gauge-like, long-range coherence/phase-flow imprint* on the weave: its field strength ($\mathcal{F}^{(\tau)}$) writes the antisymmetric temporal texture ($t_{[\mu\nu]}$) and supplies a Maxwell-type coherence stress on the diagnostic realized reference configuration source, all while remaining TT-null (no spin-2 radiation). After BEH, the CK becomes a massive spin-1 Proca field. Conceptually, its role shifts from a *long-range, scale-free phase-flow carrier* to a *finite-range mediator of temporal coherence* whose influence

is controlled by a correlation length ($\xi_{\mathrm{CK}} \sim m_{\mathrm{CK}}^{-1}$) (cf. the Proca upgrade and equations in eqs. (11.324) and (11.326)).

This has three practical consequences for the geometry. First, background control becomes *screened*: coherent CK expectations still shift ($\bar{t}_{\mu\nu}$) and hence ($\bar{\Sigma}$), but the induced temporal twist decays over (ξ_{CK}) rather than propagating freely, so the modification of ($\bar{G}_{\mu\nu}[\bar{\Sigma}]$) is localized (see Section 11.8). Second, the *new longitudinal polarization* provides a *compressional mode* of "clock-alignment" in the non-TT ledger—an extra way to organize temporal coherence—without ever leaking into tensor GW polarizations (TT-purity still applies). Third, geometry–coherence interplay remains entirely through the certified mixed blocks of the linearized diagnostic realized reference configuration law, so CK can dress/source non-TT metric response but cannot alter the *graviton channel* (eq. (11.293)).

On the source side, the CK contribution to the coherence stress keeps the *same gauge-invariant backbone* built from ($\mathcal{F}^{(\tau)}$) but now includes the Proca mass in the energy budget; the resulting stress remains typed, covariant, and compliant with the Relational NEC (Sections 11.6 and 11.8). Operationally, the massive CK thus acts as a *gapped carrier of temporal coherence*: it shapes *finite-range domains of phase organization*, stores/dissipates coherence energy locally, and backreacts on the background and perturbations in a controlled, non-TT way—all while its spin stays 1.

A simple way to picture the chrono-kymon. Think of the chrono-kymon as the field that helps nearby patches of spacetime "agree" on a local sense of timing—like a gentle wind that lines up a field of weathervanes. When the CK is massless, that "clock-wind" reaches arbitrarily far. It imprints a coherent orientation of time (a phase-flow) across wide regions, and it only has two ways to wiggle: the wind can sway sideways in two different directions (the two transverse polarizations). That's its *long-range, "clocking" role*. Now imagine something changes in the medium so that the wind can't blow freely anymore. After BEH, the CK becomes massive. In everyday terms, that doesn't mean clocks get heavier; it means the clock-wind acquires a *finite reach*. Its influence dies off beyond a characteristic distance ($\xi_{\mathrm{CK}} \sim 1/m_{\mathrm{CK}}$). Nearby regions can still synchronize, but distant ones no longer feel it strongly. And the wind can now also do one extra thing: besides swaying sideways, it can momentarily push or pull along its own direction—a *new "longitudinal" way* to disturb timing. That's what the third polarization is. A few analogies help this land:

- *Radio time signal vs. fog (or the Meissner effect).* A massless CK is like a time-signal broadcast that travels indefinitely through clear air. Giving it mass is like surrounding everything with a medium that screens the signal: it still broadcasts, but only out to a finite radius (like how magnetic fields are expelled from a superconductor and only penetrate a short distance). *Same broadcaster, shorter reach.*

- *Weathervanes in a breeze.* With massless CK, a steady breeze lines up all the vanes across the whole valley (*global clock coherence*). With massive CK, tall hedges now block the breeze, so alignment persists only locally; beyond a few fields, the effect fades. The "longitudinal" mode is like a brief gust that presses the vanes forward instead of just nudging them sideways.

- *Metronomes on a shared platform.* Put many metronomes on a movable board and they'll synchronize through tiny couplings (massless-like, long range across the board). Add mechanisms that confine the board's motion and the coupling becomes short-range: metronomes form local clusters that sync, and there's an *extra "compressional" way* the platform can respond—mirroring the CK's new longitudinal degree of freedom.

What doesn't change is the kind of thing the CK is: it remains a *spin-1 vector field* (an oriented "clock-wind"), not a spin-2 gravitational wave. BEH doesn't turn it into something else; it just adds a *gap* (so the effect is finite-range) and unlocks that *extra longitudinal wiggle*. The graviton sector stays pristine and separate; the CK continues to live in the *non-TT ledger*, acting as a *gapped carrier of temporal coherence*—organizing timing locally rather than globally.

Coupling, causality, and certification on the diagnostic realized reference configuration. The CK couples to geometry exclusively through the certified mixed blocks $\mathcal{L}_{h\tau}, \mathcal{L}_{\tau h}$ in the linearized diagnostic realized reference configuration law eq. (11.293), allowing CK to *dress* non–TT metric features without leaking into tensor polarizations. Its coherence imprint enters the RHS bookkeeping via the Maxwell/Proca–like term (vector piece in Section 11.8), while null–positivity is guaranteed by the Relational NEC (Section 11.6). Diagnostic realized reference configuration certification proceeds via typed correlator poles (existence of CK quanta, massless or massive) and witness predicates for actual *production* (which requires sources, mixing, or non–adiabatic background evolution, not BEH alone).

11.8.3 The Chrono-Kymon as a Unified Dark Sector Candidate

Massless Phase (Pre-BEH): Dark Radiation In its massless phase, the CK is a gauge-like, long-range coherence or phase-flow imprint on the weave. Its Maxwell-type stress is traceless, and if isotropically excited on the diagnostic realized reference configuration, it behaves as an extra **dark radiation** component. This contribution sits in the coherence-stress term built from the vector-typed piece $\mathcal{F}^{(\tau)}$ (cf. eqs. (11.235) and (11.248)) and contributes to the right-hand side (RHS) of the diagnostic realized reference configuration Einstein law (cf. Definition 11.106), while consistently obeying the Relational Null Energy Condition (NEC) (cf. Section 11.6). Critically, the CK remains Transverse–Traceless (TT)-null (cf. Section 10.4), ensuring it never contaminates the spin-2 gravitational wave (GW) polarizations. Conceptually, this phase is a long-range "clock-wind" that can subtly modify cosmic expansion and perturbation history without coupling to visible-sector Standard Model (SM) fields.

Broken Phase (After Internal BEH): Vector Dark Matter Once the **Brout–Englert–Higgs (BEH)** mechanism imparts a Proca mass ($m_{\mathrm{CK}} > 0$) to the CK (cf. eqs. (11.324) and (11.326)), its solutions—which possess three physical polarizations while retaining spin-1—redshift on average like pressureless matter. This dynamical behavior makes the CK a natural candidate for **vector dark matter (DM)**. The field's correlation length ($\xi_{\mathrm{CK}} \sim m_{\mathrm{CK}}^{-1}$) sets the characteristic size of coherent domains. The massive CK field remains "dark" because its universal coupling is exclusively through gravity—entering the RHS of Definition 11.106—and through any hidden-sector charges sharing its internal symmetry, bypassing direct coupling

to SM fields. The internal gauge-theoretic origin of this BEH upgrade is framed by the operator-bundle connection (cf. Section 6.6).

Dark Energy Potentiality The CK's Maxwell/Proca stress tensor does not generically yield the necessary equation of state ($w \simeq -1$) for **dark energy (DE)**. For it to mimic DE, the diagnostic realized reference configuration must sustain a nearly homogeneous, slowly varying CK background whose energy density is dominated by the potential or mass term, while the field strength $\mathcal{F}^{(\tau)}$ remains small enough to suppress anisotropic stress (cf. eq. (11.235)). In typed terms, the vector-typed contribution to the background coherence stress tensor $\bar{T}^{\mathrm{coh}}_{\mu\nu}$ must approximate an isotropic fluid and pass the Relational-NEC and diagnostic realized reference configuration diagnostics (cf. Section 11.6). Achieving this behavior is model-building–possible but significantly more constrained than the robust DM case.

Structural Consistency and Hygiene The Chrono-Kymon remains consistent with the structure of the unified theory through rigorous adherence to internal checks:

- The CK contribution always enters on the **RHS** of the field equations as coherence stress (cf. Definition 11.106), and influences the left-hand side (LHS) exclusively through background shifts of the temporal texture $\bar{t}_{\mu\nu}$; these background effects are detailed in the TT/complement framework (cf. Section 10.4).
- Causality, BRST symmetry, and Nielsen hygiene on a certified plateau ensure that the massive pole (if present) is physical and gauge-parameter independent; the massless case remains well-posed with Σ-causal Green functions (cf. Definition 9.62 and eq. (11.326)).
- In the low-energy (spectral/GR) limit, the emergent coupling scalarises and becomes universal (cf. Sections 11.5.7 and 11.5.8), so the CK's gravitational interaction is consistent with a dark sector coupling rule.

The overarching conclusion is that the Chrono-Kymon is a viable dark sector candidate: the BEH mechanism determines whether the field naturally inhabits the dark-radiation or vector–dark-matter corner of the cosmological landscape, while TT-purity keeps the spin-2 graviton channel uncontaminated (cf. eq. (11.293)).

11.8.4 BEH of the Gaiaton

Two complementary lenses; one physical mode. There is no contradiction between Chapters 10 and 11. The *Gaiaton* is a single *spin–0* scalar mode that appears (i) in Chapter 10 as the *radial (apeiron) scalar* of the Gaiamiti sector produced by global SSB (mass from the typed Hessian curvature), and (ii) in Chapter 11 as the *scalar piece of the τ–kairos*, namely the *non–TT scalar response* of the chronogeometric observable $\Sigma_{\mu\nu}$ to the same substrate fluctuations, obtained by linear response and typed projection. Formally, the two views are related by pushing forward the substrate fluctuations into chronogeometric fluctuations via the constitutive map and then selecting the non–TT scalar slot with the weave projectors (cf. Section 10.4 and Definitions 10.3 and 10.4).

Constitutive pushforward and typed selection (Ch. 10 → Ch. 11). Start from the constitutive map (Ch. 11)

$$\Sigma_{\mu\nu}[\Phi,\Theta] \;=\; g_{\mu\nu}[\Phi] \;+\; t_{\mu\nu}[\Phi,\Theta], \qquad \text{with background } (\bar\Phi,\bar\Theta).$$

Linearizing about $(\bar\Phi,\bar\Theta)$ gives the chain rule (Ch. 11)

$$\delta\Sigma_{\mu\nu} \;=\; (D_\Phi g)_{\mu\nu}\cdot\delta\Phi \;+\; (D_\Phi t)_{\mu\nu}\cdot\delta\Phi \;+\; (D_\Theta t)_{\mu\nu}\cdot\delta\Theta.$$

Let $\Pi_{\bar\Sigma}^{[\mathrm{TT}]}$ be the certified TT projector and $\mathbb{P}_{\mathrm{nonTT}} := \mathrm{Id} - \Pi_{\bar\Sigma}^{[\mathrm{TT}]}$ the TT–complement (Ch. 11, τ–construction). Let S denote the scalar typed idempotent on symmetric rank–2 tensors (Ch. 11; cf. Definition 10.4). The *τ–kairos scalar* field used in Chapter 11 is

$$\tau_{\mathrm{scalar}} \;:=\; S\,\mathbb{P}_{\mathrm{nonTT}}\,\delta\Sigma \;=\; \underbrace{S\,\mathbb{P}_{\mathrm{nonTT}}\Big((D_\Phi g)+(D_\Phi t)\Big)}_{J_\Phi}\,\delta\Phi \;+\; \underbrace{S\,\mathbb{P}_{\mathrm{nonTT}}(D_\Theta t)}_{J_\Theta}\,\delta\Theta. \tag{11.328}$$

Thus τ_{scalar} is the *linear pushforward* of $(\delta\Phi,\delta\Theta)$ under the *constitutive Jacobians*

$$J_\Phi \;:=\; S\,\mathbb{P}_{\mathrm{nonTT}}\Big((D_\Phi g)+(D_\Phi t)\Big), \qquad J_\Theta \;:=\; S\,\mathbb{P}_{\mathrm{nonTT}}(D_\Theta t). \tag{11.329}$$

All typed/TT hygiene (orthogonality, TT–nullity, band causality) is inherited from the weave/realized weave configuration (Chs. 10–11; cf. Definition 9.116 and Proposition 9.202).

Origin of the Gaiaton in Chapter 10 (substrate/Hessian viewpoint). Let $\bar\Phi$ minimize the (effective) Gaiamiti potential. Decompose the Gaiamiti fluctuation into *tangential* NG directions and the *radial (apeiron)* direction using the typed projectors (Ch. 10):

$$\delta\Phi \;=\; \sum_a \pi^a\, v_a^{\mathrm{NG}} \;+\; H_{\mathrm{G}}\, v_{\mathrm{rad}}, \qquad v_{\mathrm{rad}} \;\propto\; \Pi_{\Phi\Phi}^{[S]}\,\bar\Phi. \tag{11.330}$$

The *typed Hessian* (curvature) of the potential restricted to the scalar slot gives the radial mass,

$$m_{\mathrm{G}}^2 \;\in\; \mathrm{spec}\Big(\Pi_{\Phi\Phi}^{[S]}\Big[\delta^2\mathcal{U}_{\Phi\Phi}\Big|_{\bar\Phi}\Big]\,\Pi_{\Phi\Phi}^{[S]}\Big)\,. \tag{11.331}$$

The *gapped radial scalar* H_{G} is the *Gaiaton* (Ch. 10 SSB result). NG directions are massless prior to internal gauging and may be eaten by connection–borne vectors only after gauging (BEH) in the relevant sector (cf. Section 9.17.3).

Detection of the same mode in Chapter 11 (response/spectral viewpoint). Insert the above decomposition into the pushforward:

$$\tau_{\mathrm{scalar}} \;=\; J_\Phi\Big(\sum_a \pi^a\, v_a^{\mathrm{NG}} + H_{\mathrm{G}}\, v_{\mathrm{rad}}\Big) \;+\; J_\Theta\,\delta\Theta \;=\; \underbrace{(J_\Phi v_{\mathrm{rad}})}_{\neq 0}\,H_{\mathrm{G}} \;+\; (\text{NG and } \Theta \text{ pieces}). \tag{11.332}$$

On the certified band, the τ–scalar block $K_\tau^{(0)}$ is analyzed with the *spin–0 Riesz selector* (Ch. 11, Definitions 10.68 and 10.70):

$$\mathsf{P}^{(0)}_{\text{spec}} = \sum_{m\in\mathfrak{S}_0} \frac{1}{2\pi i}\oint_{\zeta_m} \left(K_\tau^{(0)} - z\right)^{-1} dz, \qquad \text{realized masses } m_{0,\eta} \text{ from the enclosed poles/bands.} \tag{11.333}$$

Because τ_{scalar} is a linear image of $(\delta\Phi, \delta\Theta)$, its two–point function factorizes as

$$G_{\tau\tau} \;=\; J\, G_{\text{sub}}\, J^{*}, \qquad J := (J_\Phi, J_\Theta), \tag{11.334}$$

so the poles of $G_{\tau\tau}$ are inherited from the substrate correlators G_{sub}. In particular, the *radial* Φ *pole* at $p^2 = m_{\text{G}}^2$ appears in $G_{\tau\tau}$ with nonzero residue precisely when $J_\Phi v_{\text{rad}} \neq 0$ (ensured by typed construction and orthogonality). This is exactly what the Riesz selector isolates as the *Gaiaton channel* on the τ side. NG contributions are treated by the typed/BRST machinery: they remain massless before internal gauging, or are eaten in the τ–scalar BEH completion; in all cases, TT purity precludes leakage into the TT sector (cf. Lemma 10.17 and eq. (10.105)).

Reconciliation in one line. The Gaiaton is *not* created twice. Chapter 10 *identifies* the radial scalar H_{G} as a gapped substrate mode (mass from the typed Hessian). Chapter 11 *detects the same mode* in the τ–kairos by pushing substrate fluctuations through the constitutive Jacobian $J = (J_\Phi, J_\Theta)$ and then projecting to the non–TT scalar ledger; the scalar Riesz projector picks out the same pole in the τ–scalar correlator:

$$\begin{array}{rcl} \text{Gaiamiti radial mode } (H_{\text{G}}) & \xrightarrow{\text{constitutive } J} & \\ \tau\text{–scalar response } (\tau_{\text{scalar}}) & \xrightarrow{\text{Riesz/typed}} & \text{Gaiaton channel (Ch. 11)} \end{array}$$

All steps respect TT purity and typed orthogonality; the graviton (TT Goldstone-like branch) remains disjoint from the scalar (apeiron/Gaiaton) branch throughout (cf. Definition 10.3 and Proposition 9.202).

Internal BEH: τ–scalar (Gaiaton) completion. We apply BEH to the *τ–scalar multiplet* (the gaiaton slot) rather than to the Gaiamiti field of Chapter 10. Let $\mathcal{R}_\tau^{(S)}$ be the internal representation acting on the τ–scalar typed space, with gauge fields $\mathcal{B}_\mu^I$ and Lie–algebra metric $G_{IJ}^{(\tau)}$ in the kinetic term (cf. Section 9.17.3). Write the typed order parameter $\varphi^{[S]} := \Pi_{\Phi\Phi}^{[S]}\,\bar{\Phi}$ and parameterize the scalar fiber as $\Phi = \exp\!\big(\pi^I T_I\big)\big(\varphi^{[S]} + H_{\text{G}}\big)$, where π^I are the τ–scalar NG coordinates and H_{G} is the *radial* (gaiaton) fluctuation.

Vector masses from τ–sector SSB. If the τ–scalar symmetry is broken on $\mathfrak{P}$, the $\mathcal{B}_\mu^I$ acquire a BEH mass matrix built as "fiber pairing × algebra metric":

$$(\,M^2_{\tau\text{-vec}}\,)_{IJ} = g_\tau^2 \left\langle \mathcal{R}_\tau^{(S)}(T_I)\,\varphi^{[S]}\,,\, \mathcal{R}_\tau^{(S)}(T_J)\,\varphi^{[S]} \right\rangle, \qquad (\,G^{(\tau)}\,)^{IK} G^{(\tau)}_{KJ} = \delta^I{}_J, \tag{11.335}$$

and the NG coordinates π^I are eaten. A Stueckelberg presentation keeps BRST structure manifest if desired.

Surviving scalar (the gaiaton). The radial mode H_{G} survives as the physical **gaiaton**. Its quadratic diagnostic realized reference configuration functional is the typed τ–scalar corner of the non–TT block (cf. eqs. (11.293) and (11.294)):

$$\begin{aligned} S^{(2)}_{\text{Gaiaton}} = \frac{1}{2}\int \sqrt{-\det \bar{g}^{\mathrm{Lor}}}\,\Big[H_{\mathrm{G}}\,\mathcal{K}_{SS}\,H_{\mathrm{G}} + 2\,H_{\mathrm{G}}\,\mathcal{K}_{Sg}\,(\delta g)_{\overline{\mathrm{TT}}}\Big] \\ - \kappa\int \sqrt{-\det \bar{g}^{\mathrm{Lor}}}\,\Big[H_{\mathrm{G}}\,\mathcal{J}_S + (\delta g)_{\overline{\mathrm{TT}}}\,\mathcal{J}_g\Big], \end{aligned} \tag{11.336}$$

with $\mathcal{K}_{SS}$ the τ–scalar kinetic operator and $\mathcal{K}_{Sg}$ the certified scalar–metric mixed kernel; sources are the scalar/complement projections of $\delta\bar{T}^{\mathrm{total}}$ (cf. Section 11.8). The gaiaton mass is set by the typed diagnostic realized reference configuration potential curvature,

$$m^2_{\text{Gaiaton}} = \left.\frac{\partial^2 \mathcal{V}_\Phi}{\partial \|\varphi^{[S]}\|^2}\right|_{\varphi^{[S]}=\langle\varphi^{[S]}\rangle}, \tag{11.337}$$

and covariant expansion generates the expected couplings $H_{\mathrm{G}}\,\mathcal{B}^I_\mu\mathcal{B}^{J\,\mu}$ with coefficients fixed by $\mathcal{R}^{(S)}_\tau$ and $\varphi^{[S]}$.

Placement in the mixed–block system and TT hygiene. This BEH completion refines the scalar corner (K_{SS}) and its sources in eq. (11.295), may induce controlled scalar–vector off–diagonals (K_{SA}, K_{AS}), and leaves the TT block closed: by Lemma 10.17 and eq. (10.105) the gaiaton and the $\mathcal{B}^I_\mu$ live entirely in the *non–TT ledger*. Couplings to the TT graviton h appear only through the certified mixed blocks $\mathcal{L}_{h\tau}, \mathcal{L}_{\tau h}$ (cf. eq. (11.293)); no extra spin–2 polarizations are created.

Remark 11.157 (Minimal $U(1)$ snapshot)**.** For $U(1)$ with charge $q_{\tau S}$: $m^2_{\mathcal{B}} = g^2_\tau q^2_{\tau S}\|\varphi^{[S]}\|^2$ and $S^{(2)}_{\text{Gaiaton}} = \frac{1}{2}\int\sqrt{-\det\bar{g}^{\mathrm{Lor}}}\big[(\bar{\nabla}H_{\mathrm{G}})^2 - m^2_{\text{Gaiaton}}H^2_{\mathrm{G}}\big] + \int\sqrt{-\det\bar{g}^{\mathrm{Lor}}}\,\alpha\,H_{\mathrm{G}}\,\mathcal{B}_\mu\mathcal{B}^\mu$, with $\alpha = g^2_\tau q^2_{\tau S}\|\varphi^{[S]}\|$ up to typed normalizations. This resides entirely in the non–TT subspace and respects TT purity.

The gaiaton is a *spin-0* (Lorentz-scalar) mode both *before* and *after* internal BEH in the τ–scalar sector. BEH changes its mass and couplings but *not* its spin (typed τ–kairos context from Ch. 11: cf. Definitions 10.4 and 10.15 and Section 10.4; TT purity from Lemma 10.17 and eq. (10.105)). Its scalar identity and diagnostic realized reference configuration mass input are already fixed in the global-SSB summary of Ch. 10 (cf. Section 9.17.3).

Setup (typed τ–scalar sector). Let the τ–scalar typed space carry an internal representation $\mathcal{R}^{(S)}_\tau$ (typed projectors and triad per Ch. 11, cf. Definitions 10.4 and 10.15). On the diagnostic realized reference configuration $\mathfrak{P}$ (with realized chronogeometry $(\mathcal{X}_q, \bar{\Sigma})$ and $\bar{\nabla}$), introduce the typed order parameter

$$\varphi^{[S]} := \Pi^{[S]}_{\Phi\Phi}\,\bar{\Phi} \quad \text{and the } \textit{radial} \text{ fluctuation } H_{\mathrm{G}} \quad \text{so that} \quad \Phi = \exp\Big(\pi^I T_I\Big)\Big(\varphi^{[S]} + H_{\mathrm{G}}\Big),$$

with π^I the *typed NG coordinates* (Ch. 10: cf. Section 9.17.3; see also the Ch. 10 gauging scope and completion for context Sections 9.17.3 to 9.17.4). The gaiaton is, by definition, the *radial* scalar H_{G} in the τ–scalar slot of the non–TT ledger (TT-orthogonality from Ch. 11: Lemma 10.17 and eq. (10.105); typed τ construction from Section 10.4).

Pre–BEH: quadratic functional and scalar character. The non–TT block yields the scalar corner (cf. eqs. (11.293) to (11.295))

$$S^{(2)}_{\text{Gaiaton}} = \tfrac{1}{2}\langle H_{\text{G}},\, \mathcal{K}_{SS} H_{\text{G}}\rangle_{\mathfrak{P}} + \langle H_{\text{G}},\, \mathcal{K}_{Sg}(\delta g)_{\overline{\text{TT}}}\rangle_{\mathfrak{P}} + \cdots ,$$

where all pairings use $\bar{\Sigma}$ and $\bar{\nabla}$ (cf. Section 11.8). Under local Lorentz (with respect to $\bar{\Sigma}$), H_{G} transforms as a *scalar*:

$$H_{\text{G}}(x) \mapsto H'_{\text{G}}(x) = H_{\text{G}}(x), \qquad \Rightarrow \quad \text{Pauli–Lubanski: } W_\mu\, |H_{\text{G}}(p)\rangle = 0,$$

so the one–particle states created by H_{G} are spin–0. (Equivalently, the two-point $G_{SS}(x,y)$ is a scalar bisolution of $\mathcal{K}_{SS}$; no spin projectors other than the scalar one appear.)

BEH in the τ–scalar sector (not in Gaiamiti; cf. Section 9.17.3). Gauge the *τ–scalar* symmetry with fields $\mathcal{B}^I_\mu$ and algebra metric $G^{(\tau)}_{IJ}$:

$$D_\mu \Phi := \bar{\nabla}_\mu \Phi - g_\tau\, \mathcal{B}^I_\mu\, \mathcal{R}^{(S)}_\tau(T_I)\, \Phi.$$

Expanding the kinetic term $\langle D_\mu\Phi, D^\mu\Phi\rangle$ about $\varphi^{[S]}$ gives

$$\underbrace{\tfrac{1}{2}(\bar{\nabla} H_{\text{G}})^2}_{\text{scalar kinetic}} + \underbrace{\tfrac{1}{2}\,\mathcal{B}^I_\mu (M^2_{\tau\text{-vec}})_{IJ}\, \mathcal{B}^{J\mu}}_{\text{vector masses}} + \underbrace{\alpha_{IJ}\, H_{\text{G}}\, \mathcal{B}^I_\mu \mathcal{B}^{J\mu}}_{\text{typed } HBB \text{ coupling}} + \underbrace{(\partial_\mu \pi^I + \cdots)\mathcal{B}^\mu_I}_{\text{mixing}},$$

with

$$(M^2_{\tau\text{-vec}})_{IJ} = g_\tau^2 \left\langle \mathcal{R}^{(S)}_\tau(T_I)\, \varphi^{[S]},\, \mathcal{R}^{(S)}_\tau(T_J)\, \varphi^{[S]} \right\rangle, \qquad \alpha_{IJ} \propto g_\tau^2 \left\langle \mathcal{R}^{(S)}_\tau(T_I)\, \varphi^{[S]},\, \mathcal{R}^{(S)}_\tau(T_J)\, H_{\text{G}} \right\rangle,$$

(fiber pairing $\times$ algebra metric; cf. Section 9.18.8). In unitary gauge the NG coordinates π^I are eaten, and one obtains the *BEH-completed* quadratic piece

$$\begin{aligned} S^{(2)}_{\text{Gaiaton, BEH}} = {} & \tfrac{1}{2}\langle H_{\text{G}},\, \mathcal{K}_{SS} H_{\text{G}}\rangle_{\mathfrak{P}} - \tfrac{1}{2} m^2_{\text{Gaiaton}} \langle H_{\text{G}}, H_{\text{G}}\rangle_{\mathfrak{P}} \\ & + \tfrac{1}{2}\langle \mathcal{B}_\mu, (M^2_{\tau\text{-vec}})\mathcal{B}^\mu\rangle + \langle H_{\text{G}},\, \alpha_{IJ}\, \mathcal{B}^I_\mu \mathcal{B}^{J\mu}\rangle + \cdots , \end{aligned} \tag{11.338}$$

with

$$m^2_{\text{Gaiaton}} = \left.\frac{\partial^2 \mathcal{V}_\Phi}{\partial \|\varphi^{[S]}\|^2}\right|_{\varphi^{[S]} = \langle \varphi^{[S]}\rangle} \qquad \text{(typed DRRC curvature; cf. Section 9.17.3).}$$

Here "DRRC" refers to the diagnostic realized reference configuration of Definition 9.185.

Spin after BEH: unchanged (still 0). The Lorentz transformation on H_{G} is untouched by internal gauging:

$$H_{\text{G}}(x) \mapsto H_{\text{G}}(x) \quad \text{(scalar under local Lorentz)}, \qquad \mathcal{B}^I_\mu(x) \mapsto \Lambda_\mu{}^\nu\, \mathcal{B}^I_\nu(x) \quad \text{(vector)}.$$

Thus the Pauli–Lubanski operator (built from $\bar{\Sigma}$'s local Lorentz generators) still annihilates the one–particle gaiaton states:

$$W_\mu\, |H_{\text{G}}(p)\rangle_{\text{BEH}} = 0,$$

and the Källén–Lehmann representation of the τ–scalar two–point remains purely *scalar*,

$$G_{SS}(p) = \frac{Z_S}{p^2 - m^2_{\text{Gaiaton}} + i0} + \int_{4m_{\mathcal{B}}^2}^{\infty} d\mu^2 \, \frac{\rho_S(\mu^2)}{p^2 - \mu^2 + i0},$$

with $Z_S \in (0, 1]$ and *no* spin-1/2/2 projectors (contrast eq. (11.293) for the TT sector). Couplings $H_{\text{G}}\,\mathcal{B}\mathcal{B}$ renormalize m^2_{Gaiaton} and Z_S but do not change the Lorentz *type*. In particular, TT hygiene enforces

$$\Pi^{[\text{TT}]}_{\bar{\Sigma}}\,\hat{\tau}_{\mu\nu} = 0, \qquad \Pi^{[\text{TT}]}_{\bar{\Sigma}}\,(\text{any functional of } H_{\text{G}}) = 0,$$

so no spin–2 content is ever created (cf. Lemma 10.17 and eq. (10.105)).

Conclusion. *Before* BEH, the gaiaton H_{G} is a scalar (spin–0) governed by the τ–scalar corner $\mathcal{K}_{SS}$. *After* BEH in the τ–scalar sector, NG modes are eaten and τ–vectors become massive, while the same H_{G} survives with updated (m, Z) and scalar interactions (e.g. $H\mathcal{B}\mathcal{B}$). Its Lorentz spin remains 0 throughout; only its mass/couplings change. The TT graviton sector stays closed and pure by Lemma 10.17 and eq. (10.105).

Gaiaton after τ–sector BEH: NG removal and spin conservation. Consider the τ–scalar typed space in representation $\mathcal{R}^{(S)}_{\tau}$ with typed projectors as in Ch. 11 (cf. Definition 10.4 and Section 10.4). On the diagnostic realized reference configuration $\mathfrak{P}$ with realized chronogeometry $(\mathcal{X}_q, \bar{\Sigma})$ and connection $\bar{\nabla}$, introduce the typed order parameter and radial fluctuation

$$\varphi^{[S]} := \Pi^{[S]}_{\Phi\Phi}\,\bar{\Phi}, \qquad \Phi = \exp\!\big(\pi^I T_I\big)\big(\varphi^{[S]} + H_{\text{G}}\big),$$

where π^I are τ–scalar NG coordinates (Ch. 10; cf. Section 9.17.3). The *gaiaton* is the radial scalar H_{G} in the τ–scalar slot of the non–TT ledger; TT orthogonality and purity follow from Ch. 11 (cf. Lemma 10.17 and eq. (10.105)).

(i) NG removal via the τ–vector EOM. Gauge the τ–scalar symmetry with gauge fields $\mathcal{B}^I_\mu$ and algebra metric $G^{(\tau)}_{IJ}$ (cf. Section 9.17.3). The BEH mass matrix is

$$\begin{aligned} (M^2_{\tau\text{-vec}})_{IJ} &= g_\tau^2 \left\langle \mathcal{R}^{(S)}_{\tau}(T_I)\,\varphi^{[S]}\,,\, \mathcal{R}^{(S)}_{\tau}(T_J)\,\varphi^{[S]} \right\rangle, \\ (M^2_{\tau\text{-vec}})^I{}_J &:= G^{(\tau)\,IK}(M^2_{\tau\text{-vec}})_{KJ} \quad \text{(cf. Section 9.18.8)}. \end{aligned} \tag{11.339}$$

The τ–vector Proca/BEH equation of motion reads

$$\bar{\nabla}_\mu \mathcal{F}^{I\,\mu\nu} + (M^2_{\tau\text{-vec}})^I{}_J\,\mathcal{B}^{J\,\nu} = J^{I\,\nu}_{(\tau)}, \qquad \mathcal{F}^I_{\mu\nu} := 2\bar{\nabla}_{[\mu}\mathcal{B}^I_{\nu]}. \tag{11.340}$$

Taking $\bar{\nabla}_\nu$ of (11.340) and using $\bar{\nabla}_\nu \mathcal{F}^{I\,\mu\nu}$ antisymmetry gives

$$(M^2_{\tau\text{-vec}})^I{}_J\,\bar{\nabla}_\nu \mathcal{B}^{J\,\nu} = \bar{\nabla}_\nu J^{I\,\nu}_{(\tau)} \;\Rightarrow\; \bar{\nabla}_\nu \mathcal{B}^{I\,\nu} = 0 \quad \text{on shell if} \quad \bar{\nabla}_\nu J^{I\,\nu}_{(\tau)} = 0. \tag{11.341}$$

Thus the NG coordinates are removed (eaten) by the vector fields. Equivalently, in a Stueckelberg presentation, the mass term $\frac{1}{2}\,G^{(\tau)}_{IJ}(M\mathcal{B}_\mu - \bar{\nabla}_\mu\sigma)^I(M\mathcal{B}^\mu - \bar{\nabla}^\mu\sigma)^J$ is gauge-fixed by $\sigma = 0$ (unitary gauge), reproducing (11.341). The dynamics remain in the non–TT block of the mixed system (cf. eqs. (11.293) and (11.294)), preserving TT closure.

(ii) Spin of the gaiaton is unchanged (Pauli–Lubanski test). The gaiaton H_{G} is a scalar density of weight 0 relative to $\bar{\Sigma}$. Let Q_ω be the Noether charge for an infinitesimal local Lorentz transformation with parameter $\omega^{\mu\nu} = -\omega^{\nu\mu}$. The typed Peierls bracket (equal-time limit agreeing with ETCR on $\mathfrak{P}$; cf. Section 11.8.1) gives the field variation

$$\delta_\omega H_{\mathrm{G}}(x) = \{H_{\mathrm{G}}(x), Q_\omega\}_{\mathrm{P}} = -\tfrac{1}{2}\,\omega^{\mu\nu}\left(x_\mu \bar{\nabla}_\nu - x_\nu \bar{\nabla}_\mu\right) H_{\mathrm{G}}(x), \tag{11.342}$$

which contains no intrinsic (spin) term (Peierls map and its equal-time limit: [155]; Noether–Lorentz generators on fields: [14, 82]). Passing to momentum space, the Lorentz generators act purely orbitally on the one-particle space created by H_{G}. The Pauli–Lubanski vector

$$W_\mu := -\tfrac{1}{2}\,\epsilon_{\mu\nu\rho\sigma}\, M^{\nu\rho} P^\sigma \tag{11.343}$$

therefore annihilates these states:

$$W_\mu\,|H_{\mathrm{G}}(p)\rangle = 0 \;\Rightarrow\; W^2\,|H_{\mathrm{G}}(p)\rangle = -m^2_{\text{Gaiaton}}\, s(s+1)\,|H_{\mathrm{G}}(p)\rangle = 0, \tag{11.344}$$

so $s = 0$ both before and after BEH (Pauli–Lubanski Casimir and Wigner classification: [482, 14, 255]). The scalar two–point admits the Källén–Lehmann form without non-scalar projectors,

$$G_{SS}(p) = \frac{Z_S}{p^2 - m^2_{\text{Gaiaton}} + i0} + \int_{4m^2_{\tau\text{-vec}}}^{\infty} d\mu^2\, \frac{\rho_S(\mu^2)}{p^2 - \mu^2 + i0}, \qquad Z_S \in (0,1], \tag{11.345}$$

consistent with the spectral representation for scalar channels ([14, 82, 255]).

TT hygiene and placement. All constructions above lie in the τ–kairos (Ch. 11; cf. Definition 10.4 and Section 10.4) and the non–TT ledger. TT purity is preserved:

$$\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\hat{\tau}_{\mu\nu} = 0, \qquad \hat{h}_{\mu\nu} = \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\delta\hat{g}_{\mu\nu} \quad \text{(cf. Lemma 10.17 and eq. (10.105))}. \tag{11.346}$$

The BEH completion of the τ–scalar sector refines the scalar corner in the mixed–block operator (cf. eqs. (11.294) and (11.295)) and modifies masses/couplings, but not the gaiaton's spin.

Identity and degrees of freedom. The *Chrono–Kymon (CK)* is the vector–typed piece of the τ–kairos, i.e. the TT–complement of chronogeometric fluctuations selected by the certified projector $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ (cf. Section 10.4 and Definition 10.3). In the unbroken phase it is a massless spin–1 field with two physical helicities $\lambda = \pm$. When an internal gauge symmetry acting on the CK representation is spontaneously broken on the diagnostic realized reference configuration $\mathfrak{P}$ (via the operator–bundle connection), the CK sector undergoes the Brout–Englert–Higgs (BEH) reorganization and upgrades from the Maxwell form to a Proca form (cf. eqs. (11.324) and (11.326)). The resulting massive CK is still spin–1 but now exhibits three physical polarizations $\lambda = \pm, 0$; the new longitudinal polarization reflects massiveness, not a change of spin. The *Gaiaton* is the scalar–typed (radial) piece of the τ–kairos defined by the typed scalar projector of the weave (Ch. 11; cf. Definitions 10.4 and 10.15 and Section 10.4). Before BEH in the τ–scalar sector it is a spin–0 radial mode;

after τ–scalar BEH (distinct from the Gaiamiti completion of Ch. 10), the Nambu–Goldstone directions in the scalar slot are eaten by τ–sector vectors while the radial Gaiaton survives as a spin–0 scalar. In both cases BEH modifies masses and couplings but not Lorentz spin (cf. Ch. 10 Section 9.17.3).

Background imprint. The CK imprints an *antisymmetric* temporal texture through its gauge–invariant strength $\mathcal{F}^{(\tau)}_{\mu\nu} = 2\,\bar{\nabla}_{[\mu}K^{(\tau)}_{\nu]}$, yielding $\Delta t^{(\tau)}_{[\mu\nu]} \propto \mathcal{F}^{(\tau)}_{\mu\nu}$ in the typed mean (cf. eqs. (11.288) and (11.289)). By contrast, scalar maps $S^{(\cdot)}[\varphi]$ generate a *symmetric* shift $\Delta t^{(\tau)}_{(\mu\nu)}$ associated with the Gaiaton (no antisymmetric twist). Both enter the background shift $\bar{\Sigma} \mapsto \bar{\Sigma}' = \bar{g} + \bar{t}'$ and modify the geometric and radiative operators at first order as in eqs. (11.290) and (11.291), while remaining TT–null (cf. eqs. (10.105), (11.285) and (11.286) and Lemma 10.17).

Range and the effect of BEH. In the massless phase the CK mediates *long–range* coherence of phase–flow (Maxwell–like propagation). After BEH it becomes a Proca field with *finite range* set by the correlation length $\xi_{\mathrm{CK}} \sim m_{\mathrm{CK}}^{-1}$ and gains a longitudinal polarization (cf. eqs. (11.324) and (11.326)). The Gaiaton's range is controlled by its scalar mass m_{G} obtained from the typed curvature of the diagnostic realized reference configuration potential, $m_{\mathrm{G}}^2 = \partial^2\mathcal{V}_\Phi/\partial\|\varphi^{[S]}\|^2|_{\mathrm{DRRC}}$ (cf. Section 9.17.3). After τ–scalar BEH the τ–sector vectors acquire a mass matrix

$$(M^2_{\tau\text{-vec}})_{IJ} = g_\tau^2 \left\langle \mathcal{R}^{(S)}_\tau(T_I)\,\varphi^{[S]}\,,\, \mathcal{R}^{(S)}_\tau(T_J)\,\varphi^{[S]} \right\rangle G^{(\tau)\,IJ},$$

while the Gaiaton remains a scalar with updated m_{G} determined by the same typed curvature (cf. Section 9.18.8).

Placement in the mixed–block system. Within the linearized diagnostic realized reference configuration law (cf. eq. (11.293)), the CK occupies the vector corner K_{AA} and mixes constitutively through K_{Ag}, K_{AS}; the Gaiaton occupies the scalar corner K_{SS} and mixes via K_{Sg}, K_{SA} (cf. eqs. (10.94), (10.101), (11.294) and (11.295)). All entries lie strictly in the non–TT ledger; the TT block remains closed by TT–purity (cf. Lemma 10.17 and eq. (10.105)).

Coupling to geometry. Both CK and Gaiaton alter the *background* felt by TT waves through the typed coherence shift $\Delta\Sigma^{(\tau)}$, thereby modifying dispersion/effective potentials via eqs. (11.290) and (11.291). Neither generates additional spin–2 polarizations; the graviton channel is isolated by the certified TT projector (cf. Definition 10.3, Lemma 10.17, and eq. (10.105)).

Source and energy structure. The CK coherence stress is built from $\mathcal{F}^{(\tau)}_{\mu\nu}$ (and includes $m^2_{\mathrm{CK}}K^{(\tau)}_\mu K^{(\tau)\mu}$ when massive) and remains typed and covariant; the Gaiaton stress arises from $(\bar{\nabla}H_{\mathrm{G}})^2$ and the curvature of $\mathcal{V}_\Phi$. In both cases the stress satisfies the Relational NEC on the diagnostic realized reference configuration (cf. Sections 11.6 and 11.8).

Summary. The CK is a *vector phase–flow carrier* whose BEH transition converts long–range coherence into finite–range mediation while preserving spin–1. The Gaiaton is a *scalar background–stiffness controller* whose BEH occurs in the τ–scalar sector: it remains spin–0 as nearby τ–vectors become massive and its own mass/couplings shift. In all cases the TT graviton sector (two tensor polarizations) remains unaffected by construction (cf. eq. (11.293) and Lemma 10.17).

A simple way to picture the gaiaton. The gaiaton is a *rate–stiffness dial* for the chronogeometric medium. In its massless/light regime it behaves like a gentle, uniform change to the medium's "elasticity," subtly adjusting how quickly disturbances propagate without twisting time (its imprint is *symmetric*, not antisymmetric). It does not radiate spin–2 waves; it quietly shifts the background against which those waves travel. After BEH in the τ–scalar sector, the scalar Nambu–Goldstone directions are eaten by nearby τ–vectors, which become massive, while the *radial* gaiaton remains a scalar. What changes is not its identity but its *reach and couplings*: its mass (m_{G}) (set by the typed curvature of ($\mathcal{V}_\Phi$)) fixes a scalar correlation length ($\xi_{\mathrm{G}} \sim m_{\mathrm{G}}^{-1}$), and it now couples locally to the massive τ–vectors (terms like $\langle H_{\mathrm{G}}, \mathcal{B}_\mu \mathcal{B}^\mu \rangle$). The role is still to regulate background "stiffness" and dispersion—not to carry phase-flow or create tensor polarizations.

- *Drum-head tension.* The gaiaton acts like a global tensioner on a drum skin: increasing tension raises the pitch and speeds surface waves; relaxing it lowers the pitch. Pre-BEH, tension can drift smoothly; post-BEH, tension adjustments remain but also couple to added hardware (the now-massive τ–vectors).

- *Index of refraction.* It is an index field for the chronogeometric medium: higher index slows signals, lower index speeds them. BEH makes this index modulator interact with short-range "access points" (massive τ–vectors) but does not change it into a vector; it stays a scalar knob.

- *Metronome tempo knob.* If the CK is a "clock–wind" aligning directions, the gaiaton is the "tempo knob" setting how fast the beat runs. After BEH, the tempo knob still sets the beat, while additional damped linkages (massive vectors) limit how far that adjustment propagates.

Bottom line. The CK is a *vector phase–flow carrier* (twist, antisymmetric imprint) that becomes finite–range and gains a longitudinal mode after BEH but remains spin–1. The gaiaton is a *scalar stiffness/rate controller* (symmetric imprint) that remains spin–0 before and after BEH; BEH chiefly localizes its influence and introduces scalar–vector couplings, without touching the graviton's TT sector.

11.8.5 Gaiaton as the Dark Matter and Energy Candidate

Post–BEH validity of Proposition 10.153. Yes. After completing the τ–scalar BEH (which eats the τ–scalar NG directions and gives mass to the τ–sector vectors), the *radial* scalar H_{G}—the Gaiaton—remains a spin–0 field in the non–TT ledger with updated mass and

typed couplings (cf. Section 10.8 and the τ–scalar BEH completion in Ch. 12). The dark–sector proposition remains valid with the replacements $\mathcal{V}_\Phi \to \mathcal{V}_\Phi^{(\tau\text{–BEH})}$ (the gauge–invariant, Nielsen–stable effective potential on the certified plateau) and with the inclusion of the induced scalar–vector couplings from the BEH completion (cf. Section 9.17.3 and Definition 10.68).

Low–energy post–BEH scalar sector (unitary gauge). On a certified plateau $\mathfrak{P}$ with realized chronogeometry $(\mathcal{X}_q, \bar{\Sigma})$ and Levi–Civita $\bar{\nabla}$, the quadratic-plus-interaction scalar sector for the Gaiaton takes the form

$$\begin{aligned} S_{\mathrm{G}} = \int \sqrt{-\det \bar{g}^{\mathrm{Lor}}} \Big[& \tfrac{1}{2}\, \bar{\nabla}_\mu H_{\mathrm{G}}\, \bar{\nabla}^\mu H_{\mathrm{G}} - \mathcal{V}_\Phi^{(\tau\text{–BEH})}(H_{\mathrm{G}}) \\ & + \tfrac{1}{2}\, \mathcal{B}^I_\mu (M^2_{\tau\text{–vec}})_{IJ}\, \mathcal{B}^{J\mu} + \alpha_{IJ}\, H_{\mathrm{G}}\, \mathcal{B}^I_\mu \mathcal{B}^{J\mu} + \dots \Big], \end{aligned} \tag{11.347}$$

where $(M^2_{\tau\text{–vec}})_{IJ} = g_\tau^2 \langle \mathcal{R}_\tau^{(S)}(T_I)\, \varphi^{[S]},\, \mathcal{R}_\tau^{(S)}(T_J)\, \varphi^{[S]} \rangle\, G^{(\tau)\, IJ}$ is the τ–vector mass matrix generated by τ–scalar BEH (cf. Ch. 12), and α_{IJ} are the typed $H\mathcal{B}\mathcal{B}$ coefficients fixed by $\mathcal{R}_\tau^{(S)}$ and $\varphi^{[S]}$ (cf. the scalar corner refinement of eqs. (11.294) and (11.295)). Physical observables extracted from (11.347) on $\mathfrak{P}$ are gauge–parameter independent by Nielsen identities, exactly as in Chs. 10–11.

Dark energy channel (post–BEH). Let H_{G} be spatially coherent on cosmological slices and define $\rho_H = \frac{1}{2}\dot{H}_{\mathrm{G}}^2 + \mathcal{V}_\Phi^{(\tau\text{–BEH})}$, $p_H = \frac{1}{2}\dot{H}_{\mathrm{G}}^2 - \mathcal{V}_\Phi^{(\tau\text{–BEH})}$ (with overdots taken along the diagnostic realized reference configuration flow; cf. the scalar typed stress entry $S_{\mu\nu}[\hat{\varphi}_A]$ in Definitions 10.4 and 10.13). The equation–of–state parameter is $w_H = (p_H/\rho_H) = (K - V)/(K + V)$ with $K = \frac{1}{2}\dot{H}_{\mathrm{G}}^2$, $V = \mathcal{V}_\Phi^{(\tau\text{–BEH})}$. In the slow–roll/quintessence regime,

$$\epsilon_V := \frac{M_{\mathrm{P}}^2}{2}\Big(\frac{\partial_H \mathcal{V}_\Phi^{(\tau\text{–BEH})}}{\mathcal{V}_\Phi^{(\tau\text{–BEH})}}\Big)^2 \ll 1, \qquad \eta_V := M_{\mathrm{P}}^2\, \frac{\partial_H^2 \mathcal{V}_\Phi^{(\tau\text{–BEH})}}{\mathcal{V}_\Phi^{(\tau\text{–BEH})}} \ll 1, \tag{11.348}$$

and any energy transfer to massive τ–vectors is negligible on Hubble times, $\Gamma_{H\to\mathcal{B}\mathcal{B}} \ll H$ (or kinematically forbidden if $m_{\mathrm{G}} < 2m_{\mathcal{B}}$), one has $w_H \simeq -1 + 2K/V \approx -1$ (cf. Section 10.8). Hence item (1) of Proposition 10.153 remains valid post–BEH, with $\mathcal{V}_\Phi \to \mathcal{V}_\Phi^{(\tau\text{–BEH})}$ and with the slow–roll conditions (11.348) evaluated on the BEH–completed effective potential.

Dark matter channel (post–BEH). Near a minimum, write $\mathcal{V}_\Phi^{(\tau\text{–BEH})} = \frac{1}{2}m_{\mathrm{G}}^2 H_{\mathrm{G}}^2 + \mathcal{O}(H_{\mathrm{G}}^3)$. When $H \lesssim m_{\mathrm{G}}$ the field undergoes coherent oscillations; averaged over many periods, $\langle w_H \rangle \simeq 0$ for a quadratic minimum, yielding a pressureless condensate (*misalignment*). Post–BEH viability requires:

$$\begin{aligned} &\text{(i) stability:} && \Gamma_{H\to\mathcal{B}\mathcal{B}} \ll H_0 \ \ \text{(or forbidden if } m_{\mathrm{G}} < 2m_{\mathcal{B}}), \\ &\text{(ii) weak portals:} && |\alpha_{IJ}\langle \mathcal{B}^I_\mu \mathcal{B}^{J\mu} \rangle| \ll \rho_H, \end{aligned} \tag{11.349}$$

together with the usual abundance/misalignment bounds (initial displacement, onset epoch), all read on the certified band with the scalar Riesz selector (Definition 10.68). Under (11.349), item (2) of Proposition 10.153 remains valid post–BEH.

Placement and hygiene. In both dark–energy and dark–matter regimes, the Gaiaton enters the chronogeometric response via the scalar typed map $S_{\mu\nu}[\hat{\varphi}_A]$ in the constitutive law (cf. Definitions 10.4 and 10.13). The TT block remains closed by typed orthogonality (no extra spin–2 polarizations), and all propagation/source statements continue to be organized by the mixed–block system eqs. (11.293) to (11.295).

Conclusion. Proposition 10.153 survives the τ–scalar BEH completion. The only changes are (a) the replacement of the pre–BEH potential by the gauge–invariant post–BEH effective potential $\mathcal{V}_{\Phi}^{(\tau\text{–BEH})}$ in the slow–roll analysis, and (b) the presence of controlled scalar–vector couplings from (11.347), which must satisfy (11.348)–(11.349) to preserve the dark–energy and dark–matter behaviors on the diagnostic realized reference configuration. All typed, BRST, and TT–purity properties used in Chapter 11 remain intact on certified plateaux.

11.9 The Chronogeometric Regimes

In the IQR, gravity emerges through a coherent transformation of relational structure from ultra-sub-Planckian entanglement to macroscopic classical curvature. This emergence proceeds through a discrete sequence of ontological regimes.

Hierarchy of Emergence Scales

Scale	Regime	Core Structure	Ontological Level
$\ll \ell_P$	**Ultra-sub-Planckian**	Fibrewise operator algebra $\mathfrak{F}_x$	Coherence-only domain; no geometry
$\sim \ell_P$	**Planckian**	Onset of stable modal structure	Proto-geometry; incipient field fluctuations
$\gtrsim \ell_{\mathrm{PQS}}^{-1}$	**Relational/ Mesoscopic**	Expectation tensor $\bar{\Sigma}_{\mu\nu}(x)$	Semi-classical geometry from entangled expectation
$\gg \ell_P$	**Macroscopic (IR)**	Causal manifold $\mathcal{M}_{\mathrm{geom}}$	Classical GR regime; Einstein field equations
$\sim \mathcal{L}_U$	**Cosmological**	Global solution of $\bar{R}_{\mu\nu} - \frac{1}{2}\bar{\Sigma}_{\mu\nu}\bar{R}$	Thermodynamic limit of relational gravity

Table 11.3: The hierarchy of emergent geometric regimes, from the ultra-sub-Planckian operator algebra to the macroscopic classical universe. Each scale is characterized by a distinct dominant structure and ontological interpretation within the IQR.

Transitions Between Dynamical Regimes

0. Ultra-Sub-Planckian PQS Category (Singularity Level)

Structure	Genesis Step / Description
Underlying Continuum	$\mathcal{C}_d$, a smooth, connected, Hausdorff manifold of cardinality $\mathfrak{c}$, serving as the foundational substrate for relational singularities and fiber structure (Definition 4.617).
Formal Domain	Operator-theoretic singularities, the Primordial Quantum Singularities, are defined as $\mathfrak{B}_{\vec{r}} := {}_{\hat{\Phi}}\mathring{\Lambda}_{\hat{\Theta}}$ (eq. (4.301)).
Fibred Structure	The local dual relational fiber is given by $\mathfrak{F}_x := \mathcal{F}_x^{\text{dist}} \times \mathcal{F}_x^{\text{time}}$, which encodes the potential for emergent distance and time (Definition 4.604).
Relational Transition Structure	The morphisms of $\mathcal{C}_d$ correspond to coherence-respecting transitions $\vec{r} \to \vec{r}'$, encoding pre-causal structure via operator-defined relational dynamics (Section 4.19.3).
Coherence Flow and Modal Support	**Local coherence structure:** Local coherence is structured by inter-modal correlators $\hat{\mathbf{G}}_F(x, x')$, which govern entanglement propagation and modal phase connectivity within $\mathfrak{F}_x$ (Definition 6.4). These encode proto-causal and pre-metric relational features prior to geometric emergence.
Pre-geometric Quantum Substrate	The entire structure is defined by the Quantum Holo-State $\|\mathring{\mathbf{\Psi}}\rangle$, which encodes all relational amplitudes and coherence flows at the sub-Planckian level (Definition 1.27).
Scale Support	The characteristic scale is the ultra-sub-Planckian radius ℓ_{PQS}, satisfying $\ell_{\text{PQS}} \ll \ell_P$ (Theorem 4.594).

Table 11.4: Refined summary of the Ultra-Sub-Planckian pqs Category, including operator dynamics, relational structure, and modal coherence as foundational to all emergent geometry.

1. Ultra-Sub-Planckian to Planckian

Structure / Concept	Formalism and Description
Scale / Formal Domain	The transition regime from the PQS coherence scale to the Planck length, defined by $\ell \sim \ell_{\text{PQS}} \to \ell_P$.
Relational Basis	The foundational data is provided by the fibre modal structure, $\mathfrak{F}_x := \mathcal{F}_x^{\text{dist}} \times \mathcal{F}_x^{\text{time}}$ (Definition 4.604).
Operator Activation	The chronogeometric tensor emerges at the operator level from the Gaiamiti and Tantrakyma field configurations: $$\hat{\Sigma}_{\mu\nu}(x) := \hat{g}_{\mu\nu}(x) + \hat{t}_{\mu\nu}(x)$$ (Section 5.2.6).
Emergent Expectation Tensor	The expectation-valued geometry first becomes meaningful via: $$\bar{g}_{\mu\nu}(x) := \langle \overset{\circ}{\boldsymbol{\Psi}} | \hat{g}_{\mu\nu}(x) | \overset{\circ}{\boldsymbol{\Psi}} \rangle, \quad \bar{t}_{\mu\nu}(x) := \langle \overset{\circ}{\boldsymbol{\Psi}} | \hat{t}_{\mu\nu}(x) | \overset{\circ}{\boldsymbol{\Psi}} \rangle$$ defining the first coherent semi-classical geometry (Definition 5.10, Definition 5.22).
Excitation Threshold	The first physical excitation modes appear, including the graviton field operator, defined from Gaiamiti fluctuations as: $$\hat{h}_{\mu\nu}(x) := \delta\hat{\Phi}_{\mu\nu}(x)$$ (Definition 11.28).
Triadic Modal Structure	The unified expectation tensor $\bar{\Sigma}_{\mu\nu}(x) := \bar{g}_{\mu\nu}(x) + \bar{t}_{\mu\nu}(x)$ decomposes into three coherence regimes—correlational, vacuum, and spectral—supporting the triadic structure of the graviton propagator (Definition 5.54).
Constituent Entities	Singular PQSs ($\mathfrak{B}_{\vec{r}}$) are manifested as localized coherence bundles at the scale $\ell \ll \ell_P$.

Table 11.5: Summary of the Ultra-Sub-Planckian to Planckian transition, detailing the emergence of operator-level geometry, expectation structures, and the first coherence-supported excitations.

This transition marks the emergence of linearizable metric degrees of freedom and localized coherence fluctuations.

2. Planckian to Semi-Classical Geometry

This table describes the transition from the purely operator-valued geometry to a semi-classical manifold, where expectation-valued tensor fields begin to define a coherent geometric structure.

Structure	Genesis Step / Description
Unified Operator Geometry	The foundational geometry is described by the operator-valued pair $(\hat{g}_{\mu\nu}(x),\ \hat{t}_{\mu\nu}(x))$, arising from the second variation of the relational fibers (Definition 5.1, Definition 5.17).
Emergent Expectation Geometry	Expectation values of the operator-valued tensors in the Quantum Holo-State yield the classical fields: $(\hat{g}, \hat{t}) \rightsquigarrow (\bar{g}, \bar{t})$ (Definition 1.27).
Chronogeometric Tensor	The full chronogeometric tensor is formed from the expectation values: $\bar{\Sigma}_{\mu\nu}(x) := \bar{g}_{\mu\nu}(x) + \bar{t}_{\mu\nu}(x)$ (Section 5.2.6).
Emergent Tantrakyma Excitations	The Tantrakyma field $\hat{\mathbf{\Theta}}(x)$ admits localized excitation modes—*kymons*—when the excitation projection $\hat{P}_{\text{exc}}(x)$ becomes non-zero. These modes inhabit $\mathcal{E}_{\text{exc}}(x) \subset \mathfrak{F}_x$ and define fiberwise matter excitations that contribute to $\bar{T}^{\mu\nu}_{(\Theta)}(x)$ (Definition 9.356, Section 9.18.12).
Inter-Modal Coupling and Backreaction	Geometry becomes dynamically responsive to excitation content through the emergent coupling tensor $\mathcal{C}_{\mu\nu}{}^{\rho\sigma}(x)$, defined by variations of the inter-modal correlator $\hat{G}_{\Phi\Theta}(x, x')$. This defines how matter excitations backreact onto curvature (Definition 11.111).
Emergent Connection & Curvature	The **Relational Levi-Civita connection**, $\Gamma^{\lambda}_{\mu\nu}[\bar{\Sigma}]$ (Definition 6.7), is derived from the unified tensor, which in turn defines the Ricci tensor $\bar{R}_{\mu\nu}[\bar{\Sigma}]$.

Table 11.6: Summary of the transition from Planckian operator geometry to the semi-classical regime, where expectation-valued tensor fields and matter excitations define a unified chronogeometric structure.

3. Semi-Classical to Macroscopic Gravity

This table details the final steps in the emergence of classical gravity, where the semi-classical structures crystallize into the familiar manifold and dynamics of General Relativity.

4. Macroscopic to Cosmological

Concept	Formalism and Description
Global Geometry	The geometry is described by the cosmological limit of the chronogeometric tensor, $\bar{\Sigma}_{\mu\nu}(x)$, defined over $\mathcal{X}_q$.
Governing Action Principle	The global evolution of the geometry is determined by the **Relational Einstein-Hilbert Action**: $$S_{\text{Relational}}[\bar{\Sigma}] = \int_{\mathcal{X}_q} d^d x \sqrt{-\det \bar{\Sigma}(x)}\, \bar{R}[\bar{\Sigma}](x)$$ (Section 11.5.3).
Universal State as Cosmological Boundary Condition	The Quantum Holo-State $\|\mathring{\mathbf{\Psi}}\rangle$ functions as the boundary condition for cosmological evolution. Its global coherence and entanglement topology determine the asymptotic behavior of $\bar{\Sigma}_{\mu\nu}(x)$ and the initialization of relational observables at horizon scale (Definition 1.27).
Cosmological Dynamics & Interpretation	The large-scale behavior of $\bar{\Sigma}_{\mu\nu}(x)$ and the topological structure of the universe are inherited from the universal coherence encoded in $\|\mathring{\mathbf{\Psi}}\rangle$, mediated by inter-modal correlators (Definition 11.105).
Modal Spectral Flow and Expansion	Cosmological evolution reflects modal redistribution across coherence sectors. The triadic graviton and Tantrakyma modes exhibit spectral flow, redshift, and long-range phase decoherence, which manifest as large-scale expansion, cosmological acceleration, and effective vacuum energy gradients (Definition 5.54, Definition 6.4).

Table 11.7: Summary of the emergence of the cosmological regime from macroscopic relational gravity, including global coherence, modal redistribution, and action-based evolution.

Part IV

The Standard Model

Chapter 12

The Standard Model

This chapter synthesizes the operator-geometric and chronogeometric machinery developed in the foundations into a complete account of how the Standard Model (SM) gauge and matter content emerges within the holocosmic framework. Beginning from the universal internal symmetry of the endomorphism algebra on the emergent Hilbert space, $G_{\text{int}} = \text{Aut}(\text{End}(\mathcal{H}_{\text{em}}))$ (Definition 6.113), we articulate a filtration—by locality, Tantrakyma invariance, chirality, and anomaly-freedom—that realizes a compact gauge group with the familiar decomposition $G_{\text{SM}} = SU(3)_C \times SU(2)_L \times U(1)_Y$. The resulting presentation is deliberately operational: selection is not postulated but inferred from interferometric constraints (Holosymmetry), thereby aligning the chapter's central claims with measurable coherence data rather than abstract kinematical preference. Historically, this recovers the Yang–Mills paradigm for non-Abelian connections and their dynamics [123], the electroweak unification and its spontaneous symmetry breaking [57, 58, 97], and the QCD sector governing color dynamics [540, 541], while grounding all of these in the operator-bundle formalism of the IQR program.

The Holosymmetric Selection Principle (HSP) is the methodological core (Section 12.2). Stage I is a self-consistency sieve that assigns to any candidate theory A its *Unitarity Operator* $\mathbf{U}^{(A)}_{\text{unitarity}} = (\mathbf{U}^{(A)})^\dagger \mathbf{U}^{(A)}$ with $\mathbf{U}^{(A)} = \exp\left(i\hat{H}^{(A)}\right)$, and evaluates the *Self-Coherence Functional* $\mathcal{C}(A) = \left|\langle\Psi_0|\mathbf{U}^{(A)}_{\text{unitarity}}|\Psi_0\rangle\right|$. Only theories with $\mathcal{C}(A) = 1$ pass; failure indicates dynamical non-unitarity or gauge inconsistency (including would-be anomaly obstructions in chiral sectors), thus excluding them from the operational spectrum (Section 12.2; cf. [82]). Stage II is comparative tomography: asymmetric interferometry is used to probe the relational visibilities between the realized G_{SM} and any other viable G_X. The observed suppression $\mathcal{V}_{\text{asym}}(SM, X) \approx 0$ implements selection by maximal relational orthogonality, thereby stabilizing the SM as the unique holosymmetric fixed point among nearby unitary contenders (Section 12.2).

Having fixed the gauge group, we treat gauge bosons as excitations of the *Operator Bundle Connection* $\mathcal{A}_\mu(x)$ on the Tantrakyma bundle (Definition 6.192). The connection takes values in $\mathfrak{su}(3) \oplus \mathfrak{su}(2) \oplus \mathfrak{u}(1)$, decomposing into the gluon, weak, and hypercharge components, respectively. Its curvature,

$$\mathcal{F}_{\mu\nu} = \partial_\mu \mathcal{A}_\nu - \partial_\nu \mathcal{A}_\mu + i[\mathcal{A}_\mu, \mathcal{A}_\nu],$$

provides the intrinsic Yang–Mills dynamics and underlies the quantized gauge field content

(Definitions 12.95 and 12.96; cf. [123]). This presentation is continuous with the foundational derivation of $\mathcal{A}_\mu$ as the unique gauge potential compatible with the endomorphism algebra and its covariance law $\mathcal{A}_\mu \mapsto U\mathcal{A}_\mu U^\dagger + iU\partial_\mu U^\dagger$ (Definition 6.192).

Fermions appear as odd-graded fluctuations of the Tantrakyma field $\hat{\boldsymbol{\Theta}}(x)$, realized as operator-valued spinor sections (the "kymons"). One generation is shown to furnish precisely the standard representation content $(3,2)_{1/6}$, $(1,2)_{-1/2}$, $(3,1)_{2/3}$, $(3,1)_{-2/3}$, $(1,1)_{-1}$, $(1,1)_0$, with the hypercharges verified in a consolidated table (Section 12.4.1). This matching is not merely book-keeping: it exhibits how the Tantrakyma grading, together with holosymmetric constraints, forces the chiral and charge assignments characteristic of the SM spectrum.

Electroweak symmetry breaking (EWSB) is treated in the Tantrakyma language as a vacuum alignment of an $SU(2)_L$ doublet that preserves an unbroken electromagnetic generator $Q = I_3 + \frac{1}{2}Y$. We emphasise the emergence of the Gell–Mann–Nishijima relation and verify $Y = 2(Q - I_3)$ across all fermionic multiplets (Section 12.4.3), thereby recovering the canonical electroweak charges within the operator-bundle setting [57, 58, 126]. The relation serves here as a rigid compatibility check between Tantrakyma kinematics, gauge covariance, and the low-energy charge spectrum.

The chapter then turns to flavor. We show that Yukawa couplings in the Tantrakyma sector need not be diagonal in the interaction basis; after EWSB they generate mass matrices $M_{u,d} = \bar{\Theta} Y_{u,d}/\sqrt{2}$ whose biunitary diagonalization introduces left-unitary rotations V_{uL}, V_{dL}. Their mismatch appears in the charged-current interaction as the CKM matrix, with an analogous construction for the lepton sector yielding PMNS (Section 12.5.5). This dovetails with the historical Cabibbo–Kobayashi–Maskawa and Pontecorvo–Maki–Nakagawa–Sakata narratives [542, 543, 544, 545], but is here anchored in the Tantrakyma-holosymmetric geometry rather than imposed phenomenologically.

Color dynamics and confinement are treated via **gauge-induced spectral filtering**. Quark kymons transform in **3** of $SU(3)_C$, and physical hadrons arise as projections onto the color-singlet subspace of $(\mathbb{C}^3)^{\otimes n}$: $3 \otimes 3 \otimes 3 = 1 \oplus 8 \oplus 8 \oplus 10$ for baryons, $3 \otimes \bar{3} = 1 \oplus 8$ for mesons. We formalize this as a canonical internal-gauge projection selecting the observable spectrum and state the **spectral branching** corollary restricting $\mathbb{H}^{(\Theta)}$ to $\mathcal{H}_{\text{hadron}} = P_{\text{singlet}}(\mathcal{H}^{(n)}_{\text{quark}})$ (Proposition 12.128). In this way, confinement—familiar from lattice and renormalization-group perspectives [389, 540, 541]—appears here as an intrinsic projection mechanism enforced by holosymmetric compatibility.

To quantize gauge dynamics while preserving unitarity and locality, we adopt the BRST enlargement of field space and its cohomological projection onto the physical state space (Section 12.3). This implements gauge fixing via Faddeev–Popov ghosts and encodes residual gauge symmetry in the nilpotent BRST charge, ensuring that only cohomology classes represent physical excitations [104, 105, 106]. The same formal projector architecture recurs in our treatment of color-singlet extraction, exhibiting a unifying pattern: holosymmetric admissibility corresponds to cohomological or spectral projection to physically meaningful subspaces.

The neutrino sector is addressed in a structural option that accommodates Majorana masses through Tantrakyma-sector operators, naturally realizing a seesaw hierarchy and explaining the smallness of neutrino masses without departing from the holosymmetric ethos (Section 12.7.1). The account is consistent with the historical Dirac–Majorana dialogue and

the modern seesaw synthesis [544, 546, 547], and it interfaces cleanly with the flavor-mixing analysis above.

We also provide an operational criterion for dark sectors grounded in holosymmetric tomography. A state $|\psi\rangle$ belongs to a dark gauge sector G precisely when it possesses nonzero projection under the **coherence projector** $\mathcal{P}_G^{\mathrm{coh}}$ (lensing, i.e., it is visible to holosymmetric probes in the G channel) while remaining spectrally orthogonal to all SM observables (decoupling): $\langle\psi|\mathcal{O}_{\mathrm{vis}}|\psi\rangle \approx 0$ for all $\mathcal{O}_{\mathrm{vis}} \in \mathcal{A}(G_{\mathrm{SM}})$. This formalizes the long-standing intuition behind "hidden-sector" dynamics in strictly operational terms [548, 549, 550], and demonstrates how the holosymmetric apparatus discriminates visible from dark degrees of freedom without additional ontological posits.

A short side thread engages the information problem for black holes. Within the corrected chronogeometric background of the holocosmos, we sketch a state-dependent S-matrix and an interaction Hamiltonian $\hat{H}_{AE}$ that mediate pure-to-pure evolution across collapse and evaporation, thus reconciling semiclassical Hawking flux with global unitarity. This reconciliatory stance is positioned relative to the classic paradox and contemporary entanglement-wedge perspectives [329, 138, 337], while keeping the technical details deliberately modular to the main SM narrative.

Two interlocks with the foundations deserve emphasis at the outset. First, the *Operator-geometric substrate*: the connection $\mathcal{A}_\mu$ used throughout is exactly the gauge structure canonically induced by $\mathrm{End}(\mathcal{H}_{\mathrm{em}})$ and its covariance, as developed in the first part of the treatise (Definitions 6.113 and 6.192). Second, the *chronogeometric background and transport*: all interferometric notions (holonomies, temporal potentials, parallel transport) rest on the triadic fibre bundle, chronogeometric tensors, and their gauge-covariant holonomies introduced earlier (Proposition 4.605). The present chapter, accordingly, should be read as an application and stress-test of that machinery: the SM is not appended to the IQR framework but carved from it by holosymmetric tomography.

The plan of the chapter follows this logical arc. We begin by stating the HSP and establishing the filtration to G_{SM} (Section 12.2). We then construct the gauge and matter sectors in the operator-bundle language, verify the electroweak charges and EWSB pattern, and derive the flavor structure (Definitions 12.95 and 12.96 and Sections 12.4.1, 12.4.3 and 12.5.5). Next we develop confinement as spectral filtering and formalize BRST quantization (Proposition 12.128 and Section 12.3). We close with the neutrino sector, the dark-sector criterion, and the unitary black-hole side thread. Throughout, all claims are phrased so as to admit direct operational checks within the holosymmetric paradigm, while remaining in transparent dialogue with established field-theoretic formalisms [123, 57, 58, 97, 389, 542, 543, 545, 544, 105, 106, 104, 548, 549, 550, 329, 138, 337].

A principal aim of the IQR programme is to explain why the effective gauge symmetries of nature take the form they do. Rather than postulating G_{SM} ab initio, we begin from the universal internal group $G_{\mathrm{int}} = \mathrm{Aut}(\mathrm{End}(\mathcal{H}_{\mathrm{em}}))$ and apply a sequence of structural filters that operate on the operator-geometric substrate itself (Definitions 1.32 and 1.34). The two foundational fields are bundled into the Generalized Field Operator $\hat{\mathbf{F}}(x) = \left(\hat{\mathbf{\Phi}}(x), \hat{\mathbf{\Theta}}(x)\right)^{\mathsf{T}}$ (Definition 6.1), whose correlations are collected by the Unified Inter-Modal Correlator $\hat{\mathbf{G}}_\Lambda(x, x')$ (Definition 6.4). These are the universal carriers of both geometric and matter dynamics and will serve as the stage on which the selection proceeds.

12.1 Cosmic Constraints and Candidate Symmetry Groups

Here we collect the *six cosmic constraints* that prune the universal internal automorphism group down to the physically realized gauge content. The first three are *structural/operator-geometric* necessities that any candidate $G \subset G_{\text{int}}$ must satisfy on the certified BRST-physical diagnostic realized reference configuration; the last three are *selection/phenomenology* constraints that decide among structurally viable options. After this prelude, the detailed statements and checks begin in Section 12.1.1.

Constraint A — Local implementability via a genuine connection. A candidate symmetry must be locally realizable on the operator bundle by a bona fide gauge connection $\mathcal{A}_\mu$, entering the split covariant derivative and yielding the standard curvature $\mathcal{F}_{\mu\nu}$, with correct local gauge action and compatibility with the relational Levi–Civita transport [435, 375, 248, 84]. Equivalently, all G–covariant constructions must live on the same realized weave configuration (band tags, $\bar{\Sigma}$–causality, BRST projector) so that they descend to $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$ without ambiguity [445, 106, 442, 108, 551] (Section 12.1.1).

Constraint B — Compactness and representation-theoretic well-posedness. The internal group must be (essentially) compact with unitary, well-behaved representations for Tantrakyma/matter, ensuring positive Källén–Lehmann densities for gauge-invariant correlators, a well-posed harmonic analysis on the diagnostic realized reference configuration, and stable quadratic forms for the band Hessians and Ward identities [552, 82, 91, 484, 483]. This guarantees that the structural and spectral machinery is mathematically controlled on $\mathcal{H}_{\text{phys}}$ (Section 12.1.1).

Constraint C — Symmetry of the Tantrakyma dynamics. Beyond kinematics, G must be an actual symmetry of the internal/woven sector: the Lagrangian density $\mathcal{L}_\Theta$ and potential $\mathcal{V}_\Theta$ (including Yukawa couplings) are G-invariant when written with $\nabla_\mu^{(G)}$ and $\mathcal{F}_{\mu\nu}$; the resulting Euler–Lagrange equations and Ward identities hold (Nielsen-flat) on $\mathcal{H}_{\text{phys}}$ [123, 57, 58, 59, 97, 484, 483, 120]. Only then do BEH/SSB analyses and spectrum extraction make physical sense for that G (Section 12.1.1).

Constraint D — Chiral admissibility. The weak interaction is maximally parity-violating, so the Tantrakyma spinor bundle must support a chiral grading in which only left-handed fields transform nontrivially under the weak isospin factor while right-handed fields are singlets. Concretely, with $P_{L/R} = \frac{1}{2}(1 \mp \gamma^5)$ acting on Tantrakyma spinor sections, the same modal fibre must realize inequivalent G–representations so that $P_L\Psi$ carries $SU(2)_L$ doublets and $P_R\Psi$ carries $SU(2)_L$ singlets. At the phenomenological level this reproduces, generation by generation, the quark and lepton assignments summarized in table 12.1, and it must be implementable within the Tantrakyma dynamics so that kinematics, Ward identities, and BRST/plateau readouts remain well-posed on $\mathcal{H}_{\text{phys}}$ [91, 82] (Sections 12.1.2 and 12.4.1).

Constraint E — Exact charge realization and Yukawa admissibility. The gauge sector must admit a $U(1)$ generator Y whose eigenvalues match table 12.1 and satisfy the Gell–Mann–Nishi-

jima relation $Q = I_3 + \frac{1}{2}Y$ [125, 126] (see Section 12.4.3), together with an electroweak Tantrakyma doublet $H \sim (1,2)_{+1/2}$ that permits all renormalizable, gauge-invariant Yukawa couplings [58, 59, 97]. This "Yukawa admissibility" ties the hypercharge pattern to electroweak symmetry breaking and mass generation—ensuring, in particular, that quark and lepton masses arise from H while preserving the unbroken electromagnetic generator Q. In this sense, Section 12.1.2 and table 12.1 serve as operational tests that the candidate G realizes the exact SM charges and couplings with the economical choice $G_{\rm SM} = SU(3)_C \times SU(2)_L \times U(1)_Y$.

Constraint F — Complete anomaly freedom. In chiral gauge theories, quantum anomalies must vanish identically; in the IQR/holosymmetric setting this is precisely the **Stage I viability clause of the HSP**: any uncancelled anomaly obstructs BRST nilpotency (and hence preservation of cohomology and unitarity), so the theory is excluded at the outset [442, 108, 445, 106] (Sections 12.2 and 12.3). For one generation of left-chiral Weyl fields in representations $\mathbf{R}_3 \otimes \mathbf{R}_2$ with hypercharge Y, all *local (perturbative)* anomalies collected in eq. (12.46) must cancel—namely $SU(3)^3$, $SU(2)^2U(1)$, $SU(3)^2U(1)$, $U(1)^3$, and the mixed $U(1)$–gravitational condition $\sum Y = 0$ [443, 103, 82]; the purely $SU(2)^3$ cubic anomaly is absent because the fundamental is pseudoreal [82, 91]. *Global* anomalies must also vanish, in particular the $SU(2)$ Witten anomaly, which demands an even number of left-chiral doublets per generation (satisfied by 3 quark doublets + 1 lepton doublet = 4) [553]. See Section 12.1.3.

12.1.1 Characterization of the Candidate Symmetry Group

The filtration of $G_{\rm int}$ to viable gauge sectors leverages three constraints, each now stated in operator-geometric form.

Constraint A: Local Implementability via a Genuine Connection

Realized weave configuration-compatible statement. Local gaugeability is realized by an *Operator Bundle Connection* $\mathcal{A}_\mu(x)$ on the modal Hilbert bundle (Definition 6.192). In the strengthened form, $\mathcal{A}_\mu$ is required to be *realized weave configuration-compatible*: it lives and acts on the *certified plateau* $\mathfrak{P}$ with instrumentation $\mathcal{I}^{[\mathcal{P}]} = (\bar{\Sigma}_{\mu\nu};\, P_x(\lambda_{\rm UV}),\, K_{\rm IR}^{(\ell)},\, \mathsf{P}_{\rm phys})$ (Definition 10.1), is compatible with the relational Levi–Civita transport $\nabla_\mu^{(\rm rel)}[\bar{\Sigma}]$ (Definition 9.62), descends to the BRST-physical sector $\mathcal{H}_{\rm phys} = H^0(\hat{Q}_{\rm BRST})$, preserves cone causality with respect to $\bar{\Sigma}$, and is TT-pure on the spin-2 (graviton) carrier (eq. (10.4) and Definition 10.167).

Definition 12.1 (Realized weave configuration-compatible operator bundle connection)**.** An operator-valued one–form $\mathcal{A}_\mu \in \Omega^1(\mathcal{X}_q, \mathrm{End}(\mathcal{H}_{\rm em}))$ is a *realized weave configuration-compatible operator bundle connection* if the following hold:

(a) **Local gauge law (frame covariance).** For any local quantum frame $U(x) \in G \subset G_{\rm int}$,

$$\mathcal{A}_\mu \;\mapsto\; U\,\mathcal{A}_\mu\,U^\dagger \;+\; i\,U\,\partial_\mu U^\dagger, \qquad \mathcal{F}_{\mu\nu} := \partial_\mu\mathcal{A}_\nu - \partial_\nu\mathcal{A}_\mu + i[\mathcal{A}_\mu,\mathcal{A}_\nu] \;\mapsto\; U\,\mathcal{F}_{\mu\nu}\,U^\dagger, \tag{12.1}$$

and the total covariant derivative splits as

$$\nabla_\mu \;=\; \underbrace{\nabla_\mu^{(\rm rel)}[\bar{\Sigma}]}_{\text{torsion-free, } \bar{\Sigma}\text{-compatible}} \;+\; \partial_\mu + i\,\mathrm{ad}_{\mathcal{A}_\mu} \quad \text{(eq. (6.257))}, \tag{12.2}$$

so that, for a vector-type operator $\hat{\mathbf{X}}^\lambda$,

$$(\nabla_\mu \hat{\mathbf{X}})^\lambda = \partial_\mu \hat{\mathbf{X}}^\lambda + \Gamma^\lambda_{\mu\nu}[\bar{\Sigma}]\,\hat{\mathbf{X}}^\nu + i\,[\mathcal{A}_\mu, \hat{\mathbf{X}}^\lambda] \quad \text{(eq. (6.258))}. \tag{12.3}$$

(b) **Band tameness (Ch. 7 commutator control).** With the UV tag $P_x(\lambda_{\mathrm{UV}})$ and IR window $K^{(\ell)}_{\mathrm{IR}}$,

$$\big\|\,[\,\nabla_\mu,\, P_x(\lambda_{\mathrm{UV}})\,]\,\big\| \;\le\; C\,\varepsilon\,\ell^{-1}, \qquad \big\|\,[\,\nabla_\mu,\, K^{(\ell)}_{\mathrm{IR}}\,]\,\big\| \;\le\; C'\,\varepsilon\,\ell^{-1}, \tag{12.4}$$

so that Euler–Lagrange/projected variations incur only the standard commutator corrections (cf. Ch. 7 scale-tagged calculus).

(c) **BRST chain map (cohomological compatibility).** With ghost c and BRST differential s,

$$s\,\mathcal{A}_\mu = D_\mu c, \qquad s\,c = -\tfrac{i}{2}[c,c], \qquad s\,\mathcal{F}_{\mu\nu} = i[\mathcal{F}_{\mu\nu}, c], \qquad s^2 = 0, \tag{12.5}$$

and the induced quantum charge $\hat{Q}_{\mathrm{BRST}}$ satisfies

$$[\,\hat{Q}_{\mathrm{BRST}},\, \nabla_\mu\,] = 0 \quad \text{on } \mathfrak{P}, \tag{12.6}$$

ensuring descent of gauge-covariant statements to $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ (Nielsen-flat).

(d) **Cone-preserving principal symbol (typed hyperbolicity).** Any kinetic operator built from ∇_μ has principal symbol

$$\sigma_{\mathrm{pr}}(\mathcal{K})(x,k) \;=\; \bar{\Sigma}^{\mu\nu}(x)\,k_\mu k_\nu\,\mathbf{1} \quad\Rightarrow\quad \text{retarded/advanced supports are } \bar{\Sigma}\text{-causal}, \tag{12.7}$$

implying microcausal Peierls brackets (Chs. 6–8; typed statement in Ch. 11).

(e) **TT purity on the geometry leg.** With the background-covariant TT projector $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ (eq. (10.4)),

$$\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\mathrm{ad}_{\mathcal{A}_\mu}\,\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}} = 0 \quad \text{on the Gaiamiti (geometry) leg}, \tag{12.8}$$

so internal transport does not source radiative spin-2; τ is TT-null (Definition 10.167).

(f) **Relational Bianchi and Ward hook.** The curvature obeys $D_{[\mu}\mathcal{F}_{\nu\rho]} = 0$ with $D_\mu = \partial_\mu + i[\mathcal{A}_\mu, \cdot]$, and the associated improved currents feed the Ward ledger $\mathcal{W}$ used by law-visibility meters on $\mathfrak{P}$ (Ch. 8).

Lemma 12.2 (Cohomological descent (with band stability))**.** *If $\mathcal{A}_\mu$ is realized weave configuration-compatible in the sense of Definition 12.1, then any gauge-covariant field equation, Ward identity, or observable constructed from ∇_μ, $\mathcal{F}_{\mu\nu}$, and the UV/IR band instruments $\big(P_x(\lambda_{\mathrm{UV}}),\, K^{(\ell)}_{\mathrm{IR}}\big)$ descends to a well-defined operator on the BRST-physical space $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$. Moreover, for any gauge-invariant two-point function $\mathcal{G}$ evaluated on the certified plateau $\mathfrak{P}$, (i) the pole locations and residues are independent of the gauge-fixing parameter (Nielsen-flat), and (ii) the poles, residues, and spectral projectors vary continuously with the band knobs $(\lambda_{\mathrm{UV}}, \ell)$ (band stability audit).*

Proof. Step 0: Band primitives (exact tag commutation and audit). On the realized weave configuration $\mathcal{I}^{[\mathcal{J}]}$ we assume the standard band primitives (Chs. 6–8):

$$\Big[\mathsf{P}_{\rm phys}, P_x(\lambda_{\rm UV})\Big] = \Big[\mathsf{P}_{\rm phys}, K^{(\ell)}_{\rm IR}\Big] = \Big[\nabla_\mu, P_x(\lambda_{\rm UV})\Big] = \Big[\nabla_\mu, K^{(\ell)}_{\rm IR}\Big] = 0, \tag{12.9}$$

(cf. Definitions 6.280 and 9.10 and Proposition 7.20), and the *band-stability audit*: there exists $\delta > 0$ such that for all $(\lambda'_{\rm UV}, \ell')$ with $|\lambda'_{\rm UV} - \lambda_{\rm UV}| + |\ell' - \ell| < \delta$,

$$\Big\|\Pi^{[\chi]}(\lambda'_{\rm UV}, \ell') - \Pi^{[\chi]}(\lambda_{\rm UV}, \ell)\Big\| \to 0, \qquad \text{and all tagged meters/visibilities vary continuously,} \tag{12.10}$$

where $\Pi^{[\chi]}$ denotes any typed spectral projector used on $\mathfrak{P}$ (cf. Definitions 9.116 and 9.166). *Optional robustness bound* (useful when one works with approximate tags):

$$\Big\|[\,\nabla_\mu, P_x(\lambda_{\rm UV})\,]\Big\| \le \epsilon_\nabla(\lambda_{\rm UV}), \qquad \Big\|[\,\nabla_\mu, K^{(\ell)}_{\rm IR}\,]\Big\| \le \epsilon_\nabla(\ell), \tag{12.11}$$

with $\epsilon_\nabla(\cdot) \to 0$ under the audit refinement; the main argument below only uses (12.9) and (12.10).

Step 1: Chain-map property and adjoint covariance. By (12.6) we have

$$[\hat{Q}_{\rm BRST}, \nabla_\mu] = 0, \qquad s\,\mathcal{A}_\mu = D_\mu c, \quad s\,\mathcal{F}_{\mu\nu} = i[\mathcal{F}_{\mu\nu}, c], \quad s^2 = 0,$$

so for any operator field $\hat{\Psi}$ in the adjoint, $s(\nabla_\mu \hat{\Psi}) = i[c, \nabla_\mu \hat{\Psi}]$. Let $\mathcal{O} = \mathcal{O}[\nabla, \mathcal{F}]$ be any local functional built from ∇, $\mathcal{F}$ (with $\bar{\Sigma}$-covariant contractions). Then $s\,\mathcal{O} = i[c, \mathcal{O}]$. If $\mathcal{O}$ is gauge-invariant ($s\mathcal{O} = 0$), it defines a class in $H^0(\mathrm{ad}\, s)$, and

$$\mathcal{O}_{\rm phys} \;:=\; \mathsf{P}_{\rm phys}\,\mathcal{O}\,\mathsf{P}_{\rm phys} \quad \in\; \mathrm{End}(\mathcal{H}_{\rm phys})$$

commutes with $\hat{Q}_{\rm BRST}$, hence acts on $H^0(\hat{Q}_{\rm BRST})$.

Step 2: Descent with tags (exact commutation and continuity). Define the tagged object $\mathcal{O}_{PK} := P_x(\lambda_{\rm UV})\, K^{(\ell)}_{\rm IR}\, \mathcal{O}\, K^{(\ell)}_{\rm IR}\, P_x(\lambda_{\rm UV})$. Using (12.9),

$$[\,\hat{Q}_{\rm BRST}, \mathcal{O}_{PK}\,] = PK\,[\,\hat{Q}_{\rm BRST}, \mathcal{O}\,]\,KP = 0,$$

so $(\mathcal{O}_{PK})_{\rm phys} := \mathsf{P}_{\rm phys}\mathcal{O}_{PK}\mathsf{P}_{\rm phys}$ is well-defined on H^0. By (12.10), the map $(\lambda_{\rm UV}, \ell) \mapsto (\mathcal{O}_{PK})_{\rm phys}$ is continuous in operator norm; the same holds termwise for covariant Euler–Lagrange operators and Ward functionals built from $\nabla, \mathcal{F}$. If one works with the robustness bounds (12.11), the difference $\mathcal{O}_{P'K'} - \mathcal{O}_{PK}$ is $O\Big(\epsilon_\nabla(\lambda_{\rm UV}) + \epsilon_\nabla(\ell)\Big)$.

Step 3: Decoupling of BRST-exact insertions. If $\mathcal{O} = s\Xi$, then for $|\phi\rangle, |\psi\rangle \in \mathcal{H}_{\rm phys}$,

$$\langle\phi|\,(\mathcal{O}_{PK})_{\rm phys}\,|\psi\rangle = \langle\phi|\,[\hat{Q}_{\rm BRST}, (\Xi_{PK})_{\rm phys}]\,|\psi\rangle = 0,$$

so BRST-exact contributions vanish on $\mathcal{H}_{\rm phys}$; correlation functions of gauge-invariant operators are unchanged by $s(\cdot)$-shifts after projection.

Step 4: Nielsen-flatness (gauge-parameter independence). With $S_{\rm gf+gh}(\xi) = s\,\Psi(\xi)$ we obtain the Nielsen identity for the 2-point 1PI kernel,

$$\partial_\xi \Gamma^{(2)}_{ab}(k;\xi) = \Gamma^{(2)}_{ac}(k;\xi)\,\mathcal{N}_{cb}(k;\xi) + \tilde{\mathcal{N}}_{ac}(k;\xi)\,\Gamma^{(2)}_{cb}(k;\xi), \tag{12.12}$$

where $\mathcal{N}, \tilde{\mathcal{N}}$ are built from BRST-exact composites. As (12.9) commutes tags with covariant differentiation and $\mathsf{P}_{\mathrm{phys}}$, the tagged/physical kernels satisfy the same identity. Thus zeros of $\det \Gamma^{(2)}$ (physical poles of $\Delta = (\Gamma^{(2)})^{-1}$) are ξ-independent, and the residue matrices are ξ-independent as well (regularity of $\partial_\xi \Delta$ at poles follows by differentiating $\Gamma^{(2)}\Delta = \mathbf{1}$ and using (12.12)). All steps commute with tag sandwiches by (12.9); BRST-exact remainders decouple by Step 3.

Step 5: Ward identities descend. Let $\mathcal{W}[\nabla, \mathcal{F}] = 0$ be a covariant Ward identity (bulk or boundary-tagged). Since $s\,\mathcal{W} = i[c, \mathcal{W}]$ and $[\,\hat{Q}_{\mathrm{BRST}}, \mathcal{W}\,] = 0$ modulo exact terms, we have $\mathsf{P}_{\mathrm{phys}}\, \mathcal{W}_{PK}\, \mathsf{P}_{\mathrm{phys}} = 0$ on $\mathcal{H}_{\mathrm{phys}}$.

Step 6: Continuity of poles/residues in $(\lambda_{\mathrm{UV}}, \ell)$. By Step 2 and the audit (12.10), the physical 1PI operator $\Gamma^{(2)}_{\mathrm{phys}}(k; \lambda_{\mathrm{UV}}, \ell)$ depends continuously on $(\lambda_{\mathrm{UV}}, \ell)$ in operator norm. For an isolated simple pole $k^2 = m^2$ with spectral Riesz projector

$$\Pi_\Gamma(\lambda_{\mathrm{UV}}, \ell) \;=\; \frac{1}{2\pi i} \oint_\gamma \Big(z - \Gamma^{(2)}_{\mathrm{phys}}(k; \lambda_{\mathrm{UV}}, \ell)\Big)^{-1} dz,$$

continuity of the resolvent implies continuity of Π_Γ, hence of the pole m^2 and residue in $(\lambda_{\mathrm{UV}}, \ell)$. This completes the band-stability claim.

Combining Steps 1–6 yields the stated descent to $\mathcal{H}_{\mathrm{phys}}$, Nielsen-flatness, and band stability. □

Proposition 12.3 (Cone-preserving hyperbolicity and typed microcausality)**.** *Let $\mathcal{A}_\mu$ be realized weave configuration-compatible. Then the linearized kinetic operators for gauge and matter excitations built from ∇_μ are $\bar{\Sigma}$-hyperbolic with principal symbol (12.7). Consequently, the retarded/advanced Green operators $E^\pm$ exist, have $\bar{\Sigma}$-cone support,*

$$\operatorname{supp} E^\pm \subset \{(x, y) \mid y \in J^\pm_{\bar{\Sigma}}(x)\},$$

and the (bosonic) Peierls bi-solution $\Delta := E^+ - E^-$ yields microcausal brackets: $\{F, G\}_{\mathrm{P}} = 0$ whenever the functional derivatives of F, G have $\bar{\Sigma}$-spacelike separated supports. Moreover, in the TT type one has

$$\Delta^{[\mathrm{TT}]} := \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\, \Delta\, \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$$

with the same cone support and the associated Peierls brackets are $\bar{\Sigma}$-causal and independent of the gauge parameter.

Proof. Step 1: Model kinetic operators and their principal symbols. Work on $(\mathcal{X}_q, \bar{\Sigma})$ with Levi–Civita $\nabla^{(\mathrm{rel})}$. After gauge-fixing in an R_ξ family the linearized vector operator in a local trivialization can be written as

$$(K_V^{(\xi)} A)_\mu = -\,\bar{\Sigma}^{\rho\sigma} \nabla_\rho \nabla_\sigma A_\mu + \Big(1 - \tfrac{1}{\xi}\Big) \nabla_\mu \nabla^\nu A_\nu + \mathsf{R}_\mu{}^\nu A_\nu + \mathsf{M}_\mu{}^\nu A_\nu, \tag{12.13}$$

where R, M are lower-order curvature/mass matrices (built from $\bar{\Sigma}$, background fields, and $\mathcal{F}$). Its principal symbol at $(x, k) \in T^*_x \mathcal{X}_q$ is

$$\sigma_{\mathrm{pr}}(K_V^{(\xi)})(x, k) = \bar{\Sigma}^{\rho\sigma} k_\rho k_\sigma \Big(\Pi_T(x, k) + \tfrac{1}{\xi}\, \Pi_L(x, k)\Big), \quad \Pi_L = \frac{k \otimes k}{\bar{\Sigma}^{\alpha\beta} k_\alpha k_\beta}, \;\; \Pi_T = \mathbb{1} - \Pi_L. \tag{12.14}$$

so that $\det \sigma_{\mathrm{pr}}(K_V^{(\xi)}) = \big(\bar{\Sigma}^{\rho\sigma}k_\rho k_\sigma\big)^d \xi^{-1}$ up to a nonzero constant; hence the characteristic set is the $\bar{\Sigma}$-null cone. For a scalar (Higgs/radial) field χ,

$$K_S\chi = -\,\bar{\Sigma}^{\mu\nu}\nabla_\mu\nabla_\nu\chi + \mathsf{m}^2\chi + (\text{lower-order}), \qquad \sigma_{\mathrm{pr}}(K_S) = \bar{\Sigma}^{\mu\nu}k_\mu k_\nu. \tag{12.15}$$

For a Dirac spinor ψ with covariant Dirac operator $\not{D} = \gamma^\mu(\nabla_\mu^{(\mathrm{rel})} + i\,\mathrm{ad}_{\mathcal{A}_\mu})$, the Lichnerowicz formula gives

$$\not{D}^2 = -\,\bar{\Sigma}^{\mu\nu}\nabla_\mu\nabla_\nu + \tfrac{1}{4}R[\bar{\Sigma}] + (\text{lower-order}), \qquad \sigma_{\mathrm{pr}}(\not{D}^2) = \bar{\Sigma}^{\mu\nu}k_\mu k_\nu, \tag{12.16}$$

so the second-order operator $\not{D}^2$ is normally hyperbolic. In all cases we recover (12.7).

Step 2: Green-hyperbolicity and cone support. Because the principal symbols agree with $\bar{\Sigma}^{\mu\nu}k_\mu k_\nu$ times an invertible endomorphism on each type (cf. (12.14)–(12.16)) and lower-order terms do not affect the characteristic set, each kinetic operator K above is (formally) normally hyperbolic on the globally hyperbolic background $(\mathcal{X}_q, \bar{\Sigma})$. Hence there exist unique retarded/advanced Green operators $E^\pm$ solving

$$K\,E^\pm = E^\pm K = \mathrm{Id}, \qquad \operatorname{supp} E^\pm f \subset J^\pm_{\bar{\Sigma}}(\operatorname{supp} f),$$

for compactly supported test sections f. The causal propagator $\Delta := E^+ - E^-$ is a bi-solution with $\operatorname{supp}\Delta \subset \{(x,y) \mid y \in J_{\bar{\Sigma}}(x)\}$.

Step 3: Typed TT reduction and independence of ξ. Let $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ be the background-covariant York–Lichnerowicz projector. At the principal-symbol level $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ depends on (x,k) through $\bar{\Sigma}$ and the first-order differential $\mathcal{L}_{\bar{\Sigma}}$; hence

$$\Big[\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}},\, \sigma_{\mathrm{pr}}(K_V^{(\xi)})\Big] = 0,$$

and lower-order commutators only change subprincipal structure. Define the TT-reduced operator and Green operators by

$$K^{[\mathrm{TT}]} := \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}} K_V^{(\xi)} \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}, \qquad E^{[\mathrm{TT}],\pm} := \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}} E^\pm \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}.$$

Using (12.8) (TT purity on the geometry leg), the ξ-dependent longitudinal part of $K_V^{(\xi)}$ is annihilated when sandwiched with $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$. Therefore $K^{[\mathrm{TT}]}$ and $E^{[\mathrm{TT}],\pm}$ are independent of ξ and inherit the same cone support:

$$\operatorname{supp} E^{[\mathrm{TT}],\pm} \subset \{(x,y) \mid y \in J^\pm_{\bar{\Sigma}}(x)\}.$$

The TT causal propagator $\Delta^{[\mathrm{TT}]} := E^{[\mathrm{TT}],+} - E^{[\mathrm{TT}],-}$ is thus $\bar{\Sigma}$-causal and ξ-independent.

Step 4: Peierls brackets and microcausality (typed). For a bosonic field sector φ with kinetic operator K, the Peierls bracket of two functionals F, G is

$$\{F,G\}_{\mathrm{P}} = \Big\langle \frac{\delta F}{\delta\varphi},\ \Delta\,\frac{\delta G}{\delta\varphi}\Big\rangle,$$

where the pairing uses the $\bar{\Sigma}$-compatible measure and index contractions. If the supports of $\delta F/\delta\varphi$ and $\delta G/\delta\varphi$ are $\bar{\Sigma}$-spacelike separated, then Δ vanishes on their product support and

$\{F,G\}_{\mathrm{P}}=0$. In the TT type we replace Δ by $\Delta^{[\mathrm{TT}]}$ and φ by $h_{\mu\nu}$ projected by $\Pi_{\bar\Sigma}^{[\mathrm{TT}]}$; the same support argument yields

$$\{F,G\}_{\mathrm{P}}^{[\mathrm{TT}]}=\Big\langle \frac{\delta F}{\delta h},\ \Delta^{[\mathrm{TT}]}\ \frac{\delta G}{\delta h}\Big\rangle = 0 \quad \text{for } \bar\Sigma\text{-spacelike separation},$$

and by Step 3 it is independent of ξ.

Step 5: Matter (spin-$\frac{1}{2}$) remark. For spinors the Peierls construction applies to $\not{D}^2$ (bosonic second-order form) or, equivalently, microcausality is expressed in terms of the anticommutator bi-solution of $\not{D}$; in both cases the cone support follows from (12.16).

Combining Steps 1–4 completes the proof. □

Definition 12.4 (Boundary-compatible connection (optional))**.** Work on a certified *boundary* plateau $\mathfrak{P}_\partial$ with tangential IR window $K^{(L)}_{\mathrm{IR},\partial}$ and BRST-closed tags, as in the boundary witnesses (Definition 8.31) *on* $\mathcal{H}_{\mathrm{phys}}:=H^0(\hat Q_{\mathrm{BRST}})$ (Ch. 8). Let $(\mathcal{X}_q,\bar\Sigma)$ denote the realized chronogeometric background with the Relational Levi–Civita connection $\nabla^{(\mathrm{rel})}$ (Chs. 9–11). A connection $\mathcal{A}_\mu$ on the operator bundle is *boundary-compatible* if the following hold.

(i) Tangential geometry and parallel transport. Let $\partial\mathcal{X}_q\hookrightarrow\mathcal{X}_q$ be a smooth, co-dimension–1 submanifold with unit normal n_μ (w.r.t. $\bar\Sigma$) and projector

$$P^\mu{}_\nu \;:=\; \delta^\mu{}_\nu - n^\mu n_\nu\,.$$

Define the *tangential metric* and *tangential Levi–Civita* by

$$\bar\Sigma^{\parallel}_{\mu\nu}:=P_\mu{}^\rho P_\nu{}^\sigma\,\bar\Sigma_{\rho\sigma},\qquad \nabla^{(\mathrm{rel})}_{\parallel\mu}:=P_\mu{}^\rho\,P^\alpha{}_\beta\,\nabla^{(\mathrm{rel})}_\rho\,,$$

so that $\nabla^{(\mathrm{rel})}_{\parallel}$ is torsion-free and compatible with $\bar\Sigma^{\parallel}$ along $\partial\mathcal{X}_q$ (Levi–Civita transport and measure are those of $\bar\Sigma$, cf. [84, 248]; see also the Levi–Civita parallel transport/measure usage in Ch. 11). *(Parallel transport and relational measure:* Definition 6.7 and the boxed response integral with $d\mu_{\bar\Sigma}$ and Levi–Civita transport.)

(ii) Tangential restriction of the gauge data. Let the *tangential connection* be $\mathcal{A}_{\parallel\mu}:=P_\mu{}^\nu\,\mathcal{A}_\nu$ and the *tangential covariant derivative* on any operator field $\hat\Psi$ be

$$D^{(\partial)}_{\parallel\mu}\hat\Psi \;:=\; \nabla^{(\mathrm{rel})}_{\parallel\mu}\hat\Psi \;+\; i\,[\mathcal{A}_{\parallel\mu},\hat\Psi]\,,$$

so that under a boundary frame change $U|_{\partial\mathcal{X}_q}$ one has

$$\mathcal{A}_{\parallel\mu}\;\mapsto\; U\,\mathcal{A}_{\parallel\mu}\,U^\dagger \;+\; i\,U\,\nabla^{(\mathrm{rel})}_{\parallel\mu}U^\dagger\,.$$

(The bulk connection $\mathcal{A}_\mu$ and its local implementability are as in the Operator Bundle Connection; cf. Ch. 12 overview of the connection.)

(iii) Tangential curvature and Bianchi identity. Define the *tangential curvature* either index-free,

$$\mathcal{F}_{\parallel} \;:=\; \mathrm{d}_{\parallel}\mathcal{A}_{\parallel} \;+\; i\,\mathcal{A}_{\parallel}\wedge\mathcal{A}_{\parallel}\,, \qquad \mathrm{d}_{\parallel} := P\circ\mathrm{d}\circ P,$$

or in components,

$$\mathcal{F}_{\parallel\mu\nu} \;:=\; P_{\mu}{}^{\rho}P_{\nu}{}^{\sigma}\Big(\partial_{\rho}\mathcal{A}_{\sigma}-\partial_{\sigma}\mathcal{A}_{\rho}+i[\mathcal{A}_{\rho},\mathcal{A}_{\sigma}]\Big)\,,$$

so that the *tangential Bianchi identity* holds, $D_{\parallel}^{(\partial)}\mathcal{F}_{\parallel}=0$, with $D_{\parallel}^{(\partial)}$ the adjoint covariant derivative built from $\mathcal{A}_{\parallel}$. (For the bulk curvature law and Yang–Mills form, see Ch. 12 prelude.)

(iv) Boundary band–tameness (tangential form). With the boundary IR window $K^{(L)}_{\mathrm{IR},\partial}$ and centered tag $P_x(\lambda_{\mathrm{UV}})$ ("arena" $\mathcal{A}=\mathfrak{P}_{\partial}$, IR knob $r=L$), the following operator bounds hold *uniformly on* $\mathfrak{P}_{\partial}$:

$$\Big\|[K^{(L)}_{\mathrm{IR},\partial},\nabla^{(\mathrm{rel})}_{\parallel}]\Big\|_{\mathcal{I}} \;\lesssim\; L^{-1}, \qquad \Big\|K^{(L)}_{\mathrm{IR},\partial}\,\mathcal{A}_{\parallel}\,K^{(L)}_{\mathrm{IR},\partial}\Big\|_{\mathcal{I}} \;+\; \Big\|K^{(L)}_{\mathrm{IR},\partial}\,\mathcal{F}_{\parallel}\,K^{(L)}_{\mathrm{IR},\partial}\Big\|_{\mathcal{I}} \;<\; \infty,$$

and these bounds commute with the BRST projector and with the boundary band sandwiches $P_x(\lambda_{\mathrm{UV}})\,K^{(L)}_{\mathrm{IR},\partial}(\cdot)\,K^{(L)}_{\mathrm{IR},\partial}\,P_x(\lambda_{\mathrm{UV}})$ used for edge diagnostics. *(Boundary arena, IR knob L, and tag/witness sandwich:* Definition 8.31, Corollary 8.46 items 1–3.)

(v) Tagged boundary Ward identity (BRST/adjoint form). Let the (improved, scale-tagged) boundary current be $J^{(\partial)}_{\parallel\mu} := \delta S/\delta\mathcal{A}_{\parallel}{}^{\mu}$ restricted to $\partial\mathcal{X}_q$, with the standard BRST improvements. The *tagged boundary Ward identity* reads

$$\nabla_{\parallel}\cdot J^{(\partial)}_{\parallel} \;=\; S_{\partial} \quad \text{on } \mathcal{H}_{\mathrm{phys}}, \qquad \text{equivalently} \quad \nabla^{(\mathrm{ad})}_{\parallel\mu}J^{(\partial)\mu}_{\parallel} = S_{\partial} \;\; (\mathrm{mod}\ \zeta\text{-exact}),$$

and is the input for the boundary law visibility $\Lambda^{[\partial]}_{\mathrm{law}}(L)$. *(Boundary witnesses and their Ward source:)* Definition 8.31. *(Scale-tagged adjoint Ward identity in the bulk setting, for comparison:)* Definition 7.567.

(vi) BRST closure of boundary data. The BRST variations restrict tangentially: $s\,\mathcal{A}_{\parallel} = D^{(\partial)}_{\parallel}c|_{\partial}$, $s\,J^{(\partial)}_{\parallel} = i[c,J^{(\partial)}_{\parallel}]$, and $s\,S_{\partial}=0$, so the Ward identity and the associated visibility on $\mathfrak{P}_{\partial}$ are Nielsen/BRST–safe under projector sandwiches and audit moves adopted in the boundary arena (Ch. 8 checklist; arenas and typed witnesses).

Any $\mathcal{A}_{\mu}$ obeying (i)–(vi) will be said to be *boundary-compatible* on $(\mathfrak{P}_{\partial}; L; P_x, K^{(L)}_{\mathrm{IR},\partial}; \mathsf{P}_{\mathrm{phys}})$.

Practical checklist (diagnostic realized reference configuration tests for a candidate G). Given a putative $G\subset G_{\mathrm{int}}$, the following checks certify Constraint A on $\mathfrak{P}$:

(A.1) *Transformation/curvature:* verify (12.1); compute $\mathcal{F}_{\mu\nu}$, Bianchi identity.

(A.2) *Split connection:* implement (12.2)–(12.3) with $\Gamma^{\lambda}_{\mu\nu}[\bar{\Sigma}]$ (Ch. 6).

(A.3) *Band tameness:* estimate commutators (12.4) on the chosen tags (Ch. 7).

(A.4) *BRST chain map:* check (12.5) and (12.6); verify Nielsen-flat poles.

(A.5) *Cone causality:* confirm (12.7) for the kinetic operators (Chs. 6–8).

(A.6) *TT purity:* verify (12.8) against the TT projector (eq. (10.4)) (Chs. 10–11).

⋆ *Boundary (if present):* apply Definition 12.4 and the boundary Ward test (Ch. 8).

Constraint B: Compactness and representation-theoretic well-posedness

Objective. Constraint B certifies, on the certified plateau $\mathfrak{P}$, that: (i) the internal symmetry actually realized by the Operator Bundle Connection $\mathcal{A}_\mu$ has *compact* local holonomy groups (unitary, finite-dimensional action on each effective internal fibre); and (ii) the Tantrakyma sector is *representation-theoretically well-posed*: finite/unitary decomposition per band, positive Källén–Lehmann (KL) spectral measures, and Ward identities that descend to $\mathcal{H}_{\rm phys} = H^0(\hat{Q}_{\rm BRST})$ and are Nielsen-flat. We now formalize these requirements using the GNS substrate (Definition 1.34), the band instruments/tag audit, and the BRST/typed infrastructure established in Chs. 6–11 (cf. Definition 6.192, Lemma 12.2, and Proposition 12.3).

Definition 12.5 (GNS substrate and Hermitian gauge directions)**.** Let $(\pi_\omega, \mathcal{H}_{\rm em}, |\Omega_\omega\rangle)$ be the GNS triple of the operational algebra $\mathcal{A}$ with holo-state $\omega^{\mathring{\Psi}}$ (Definition 1.34). The *gauge directions* are the Hermitian endomorphisms acting in the Tantrakyma fibre:

$$\mathfrak{g}_{\rm em} \;:=\; \{\, T \in \mathrm{End}(\mathcal{H}_{\rm em}) \mid T = T^\dagger,\ \Pi_{\rm Tan}\, T\, \Pi_{\rm Tan} = T \,\},$$

where $\Pi_{\rm Tan}$ is the Tantrakyma projector (commutes with the realized weave configuration instruments). We write the connection in represented form as $\mathcal{A}_\mu(x) = A^a_\mu(x)\, T_a \in \pi_\omega(\mathcal{A}) \subset \mathrm{End}(\mathcal{H}_{\rm em})$ (Definition 6.192). By Stone's theorem, each T_a generates a unitary one-parameter group $U_a(\alpha) = e^{i\alpha T_a}$, so any local frame $U(x) = e^{i\alpha^a(x) T_a}$ is unitary on $\mathcal{H}_{\rm em}$.

Definition 12.6 (Effective internal fibre on a band window)**.** For a realized weave configuration tag pair $\big(P_x(\lambda_{\rm UV}),\, K^{(\ell)}_{\rm IR}\big)$, define the *effective internal fibre* at x by

$$\mathcal{H}_{\rm int}(x; \lambda_{\rm UV}, \ell) \;:=\; \mathrm{Ran}\Big(K^{(\ell)}_{\rm IR}\, P_x(\lambda_{\rm UV})\, \Pi_{\rm Tan}\Big) \;\subset\; \mathcal{H}_{\rm em}.$$

Lemma 12.7 (Finite internal multiplicity and band stability)**.** *On any certified plateau $\mathfrak{P}$, assume the tag–commutation relations*

$$\Big[\mathsf{P}_{\rm phys}, P_x(\lambda_{\rm UV})\Big] = \Big[\mathsf{P}_{\rm phys}, K^{(\ell)}_{\rm IR}\Big] = \Big[\nabla_\mu, P_x(\lambda_{\rm UV})\Big] = \Big[\nabla_\mu, K^{(\ell)}_{\rm IR}\Big] = 0, \tag{12.17}$$

(cf. eq. (12.9), Definitions 6.280 and 9.10 and Proposition 7.20). Then:

(a) $\dim \mathcal{H}_{\rm int}(x; \lambda_{\rm UV}, \ell) < \infty$.

(b) (Audit continuity) *There exists $\delta > 0$ such that whenever $|\lambda'_{\rm UV} - \lambda_{\rm UV}| + |\ell' - \ell| < \delta$,*

$$\Big\| \Pi_{\rm int}(x; \lambda'_{\rm UV}, \ell') - \Pi_{\rm int}(x; \lambda_{\rm UV}, \ell) \Big\| \;\to\; 0,$$

with $\Pi_{\rm int}$ the orthogonal projector onto $\mathcal{H}_{\rm int}(x; \lambda_{\rm UV}, \ell) = \mathrm{Ran}\Big(K^{(\ell)}_{\rm IR} P_x(\lambda_{\rm UV}) \Pi_{\rm Tan}\Big)$ (cf. Definition 9.166).

Proof. Preliminaries. Set

$$P := P_x(\lambda_{\mathrm{UV}}), \qquad K := K_{\mathrm{IR}}^{(\ell)}, \qquad \Pi_{\mathrm{Tan}} = \Pi_{\mathrm{Tan}}^{\dagger} = \Pi_{\mathrm{Tan}}^{2}.$$

By (12.17) and typed compatibility, these commute with the physical projector and with ∇_μ on $\mathfrak{P}$, so algebraic manipulations below are stable under the BRST descent (Lemma 12.2).

(a) Finiteness. By construction of the IR window (Ch. 7),

$$K \;=\; \chi_{[0,\Lambda(\ell)]}\big(\mathcal{K}\big), \tag{12.18}$$

where $\mathcal{K}$ is a $\bar{\Sigma}$-elliptic, essentially self-adjoint, nonnegative operator on the internal (Tantrakyma) bundle (e.g. the spatial part of the normally hyperbolic kinetic operator, cf. Proposition 12.3), and χ is the spectral indicator. On the certified plateau the arena is cut to finite IR scale and localized near x by the UV tag P; the resolvent of $\mathcal{K}$ is compact on this tagged arena (Rellich–Kondrachov type compactness for the $\bar{\Sigma}$-geometry—see the typed projector/band setup in Definition 9.116). Therefore the spectrum of $\mathcal{K}$ in $[0, \Lambda(\ell)]$ is *pure point* with *finite* total multiplicity; hence

$$\operatorname{rank} K \;<\; \infty. \tag{12.19}$$

Since P and Π_{Tan} are bounded projections, the range

$$\mathcal{H}_{\mathrm{int}}(x; \lambda_{\mathrm{UV}}, \ell) \;=\; \operatorname{Ran}\big(K\,P\,\Pi_{\mathrm{Tan}}\big)$$

is contained in $\operatorname{Ran} K$. Thus $\dim \mathcal{H}_{\mathrm{int}}(x; \lambda_{\mathrm{UV}}, \ell) \le \operatorname{rank} K < \infty$, proving (a).

(b) Audit continuity. We prove norm–continuity of the orthogonal projector onto $\operatorname{Ran}(KP\Pi_{\mathrm{Tan}})$ under small changes of the knobs $(\lambda_{\mathrm{UV}}, \ell)$. By the *band-stability audit* (Definition 9.166) there exists $\delta > 0$ such that for all $(\lambda'_{\mathrm{UV}}, \ell')$ with $|\lambda'_{\mathrm{UV}} - \lambda_{\mathrm{UV}}| + |\ell' - \ell| < \delta$,

$$\big\|P' - P\big\| + \big\|K' - K\big\| \;\to\; 0, \qquad \operatorname{rank} K' = \operatorname{rank} K, \tag{12.20}$$

i.e. no eigenvalue of $\mathcal{K}$ crosses the IR cutoff and the tagged ranks are constant on the audit neighborhood. Set

$$S \;:=\; KP\Pi_{\mathrm{Tan}} \in \mathcal{B}(\mathcal{H}_{\mathrm{em}}), \qquad S' \;:=\; K'P'\Pi_{\mathrm{Tan}}.$$

Then $\operatorname{Ran} S = \mathcal{H}_{\mathrm{int}}(x; \lambda_{\mathrm{UV}}, \ell)$, and similarly for S'. Consider the positive operators $H := SS^{\dagger}$ and $H' := S'S'^{\dagger}$. Because $\operatorname{rank} S = \operatorname{rank} K < \infty$, the spectrum of H consists of $r := \operatorname{rank} S$ positive eigenvalues $\{\sigma_j^2\}_{j=1}^{r}$ (the squared singular values of S) and 0 with infinite multiplicity. By (12.20) and standard norm–perturbation estimates,

$$\big\|H' - H\big\| \;\le\; \|S' - S\|\,\Big(\|S'\| + \|S\|\Big) \;\to\; 0. \tag{12.21}$$

Let $\eta := \frac{1}{2}\min_{1\le j\le r}\sigma_j^2 > 0$; then the spectral projector of H onto $[\eta, \infty)$ equals the orthogonal projector onto $\operatorname{Ran} S$:

$$\Pi_{\mathrm{int}} = \mathbf{1}_{[\eta,\infty)}(H).$$

By rank constancy in (12.20), the spectrum of H' also has exactly r positive eigenvalues and admits the same $\eta > 0$ gap for all small enough knob variations. Using the Dunford–Riesz integral formula for spectral projectors,

$$\Pi_{\mathrm{int}} = \frac{1}{2\pi i}\oint_\Gamma (z - H)^{-1}\, dz, \qquad \Pi'_{\mathrm{int}} = \frac{1}{2\pi i}\oint_\Gamma (z - H')^{-1}\, dz,$$

with Γ a fixed contour enclosing $[\eta, \infty)$ and excluding 0, we obtain by resolvent continuity (from (12.21))

$$\left\|\Pi'_{\mathrm{int}} - \Pi_{\mathrm{int}}\right\| \;\le\; \frac{1}{2\pi}\,\mathrm{len}(\Gamma)\,\sup_{z\in\Gamma}\left\|(z - H')^{-1} - (z - H)^{-1}\right\| \;\xrightarrow[(\lambda'_{\mathrm{UV}},\ell')\to(\lambda_{\mathrm{UV}},\ell)]{}\; 0.$$

Equivalently, one may invoke the Davis–Kahan $\sin\Theta$ theorem for subspace perturbations applied to the pairs (H, H') with the fixed gap at 0, yielding

$$\left\|\Pi'_{\mathrm{int}} - \Pi_{\mathrm{int}}\right\| \;\le\; \frac{2}{\eta}\left\|H' - H\right\| \;\to\; 0.$$

This proves (b).

All steps are compatible with the BRST chain-map and tag–commutation hypotheses, hence the conclusions hold on $\mathfrak{P}$ and descend to $\mathcal{H}_{\mathrm{phys}}$ (Lemma 12.2). □

Proposition 12.8 (Compact holonomy on finite internal fibres)**.** *Let $\mathcal{A}_\mu = A^a_\mu T_a$ as in Definition 12.5. On each finite fibre $\mathcal{H}_{\mathrm{int}}(x; \lambda_{\mathrm{UV}}, \ell)$ the local holonomy group*

$$\mathrm{Hol}_x(\mathcal{A}) \;\subset\; U\Big(\mathcal{H}_{\mathrm{int}}(x; \lambda_{\mathrm{UV}}, \ell)\Big)$$

is a Lie subgroup of a compact unitary group; hence $\mathrm{Hol}_x(\mathcal{A})$ is compact. Its Lie algebra is represented by skew-adjoints iT_a with $[T_a, T_b] = i f_{ab}{}^c T_c$ (physics convention), and on each semisimple block the Killing form $\kappa_{ab} := -\mathrm{Tr}(\mathrm{ad}_{T_a}\mathrm{ad}_{T_b})$ is positive (physics sign).

Proof. Step 0: Band fibre and covariant constancy of tags. Fix a certified plateau $\mathfrak{P}$ and a point $x \in \mathcal{X}_q$. Let

$$P := P_x(\lambda_{\mathrm{UV}}), \qquad K := K^{(\ell)}_{\mathrm{IR}}, \qquad \Pi_{\mathrm{Tan}} = \Pi^\dagger_{\mathrm{Tan}} = \Pi^2_{\mathrm{Tan}},$$

and write the effective internal fibre as $\mathcal{H}_{\mathrm{int}}(x; \lambda_{\mathrm{UV}}, \ell) = \mathrm{Ran}(K\,P\,\Pi_{\mathrm{Tan}})$ (Definition 12.6). By the tag–commutation relations on $\mathfrak{P}$,

$$[\nabla_\mu, P] = [\nabla_\mu, K] = [\nabla_\mu, \Pi_{\mathrm{Tan}}] = 0 \quad \text{on } \mathfrak{P}$$
(cf. eq. (12.17), Definitions 6.280 and 9.10, and Proposition 7.20). (12.22)

Hence the projector $S := K\,P\,\Pi_{\mathrm{Tan}}$ is *covariantly constant* in the sense of the realized weave configuration-compatible split connection $\nabla_\mu = \nabla^{(\mathrm{rel})}_\mu[\bar{\Sigma}] + \partial_\mu + i\,\mathrm{ad}_{\mathcal{A}_\mu}$ (Definition 12.1 and eq. (12.2)): $\nabla_\mu S = 0$ on $\mathfrak{P}$.

Step 1: Parallel transport preserves the band fibre. Let $\gamma : [0, 1] \to \mathcal{X}_q$ be a piecewise smooth curve with $\gamma(0) = x$. Define the (total) parallel transporter $U_\gamma(s)$ along γ by

$$\frac{d}{ds}U_\gamma(s) = \dot\gamma^\mu(s)\left(\nabla^{(\mathrm{rel})}_\mu + i\,\mathrm{ad}_{\mathcal{A}_\mu}\right)U_\gamma(s), \qquad U_\gamma(0) = \mathbf{1}, \tag{12.23}$$

acting on the operator bundle (cf. eq. (12.2) and Definition 6.192). From $\nabla_\mu S = 0$ we obtain $\frac{d}{ds}\big(U_\gamma(s)^{-1} S_{\gamma(s)}\, U_\gamma(s)\big) = 0$, so

$$U_\gamma(s)^{-1}\, S_{\gamma(s)}\, U_\gamma(s) \;=\; S_x \quad \text{for all } s \in [0,1]. \tag{12.24}$$

In particular, for a loop γ based at x ($\gamma(1) = x$), $U_\gamma(1)\, S_x = S_x\, U_\gamma(1)$, so $U_\gamma(1)$ *reduces* the subspace $\mathrm{Ran}(S_x) = \mathcal{H}_{\mathrm{int}}(x;\lambda_{\mathrm{UV}},\ell)$ (cf. Definition 12.6). Thus the holonomy acts by endomorphisms of $\mathcal{H}_{\mathrm{int}}(x;\lambda_{\mathrm{UV}},\ell)$.

Step 2: Unitarity of the holonomy on the fibre. In a local trivialization where $\nabla^{(\mathrm{rel})}$ acts on spacetime indices and $\mathcal{A}_\mu$ on the internal fibre, the restricted transporter $U^{\mathrm{int}}_\gamma(s)$ obeys

$$\frac{d}{ds} U^{\mathrm{int}}_\gamma(s) = i\,\dot\gamma^\mu(s)\,\mathcal{A}_\mu\big(\gamma(s)\big)\, U^{\mathrm{int}}_\gamma(s), \qquad U^{\mathrm{int}}_\gamma(0) = \mathbf{1}, \tag{12.25}$$

(cf. Definition 12.5). Since $T_a = T_a^\dagger$ and $\mathcal{A}_\mu = A^a_\mu T_a$ is Hermitian on $\mathcal{H}_{\mathrm{em}}$, the generator $i\mathcal{A}_\mu$ is *skew*-adjoint; therefore

$$\frac{d}{ds}\big(U^{\mathrm{int}}_\gamma(s)^\dagger U^{\mathrm{int}}_\gamma(s)\big) = 0,$$

and $U^{\mathrm{int}}_\gamma(s)$ is unitary for all s. For a loop, $U^{\mathrm{int}}_\gamma(1) = \mathcal{P}\exp i\oint_\gamma \mathcal{A}$ is unitary and, by (12.24), preserves the fibre $\mathcal{H}_{\mathrm{int}}(x;\lambda_{\mathrm{UV}},\ell)$. Hence

$$\mathrm{Hol}_x(\mathcal{A}) \;\subset\; U\big(\mathcal{H}_{\mathrm{int}}(x;\lambda_{\mathrm{UV}},\ell)\big) \;\cong\; U(N), \qquad N := \dim \mathcal{H}_{\mathrm{int}}(x;\lambda_{\mathrm{UV}},\ell) < \infty$$

by Lemma 12.7(a).

Step 3: Lie subgroup and compactness. By the Ambrose–Singer theorem (holonomy of smooth connections), the (restricted) holonomy group is a *Lie subgroup* whose Lie algebra is generated by parallel translates of the curvature two-form along loops based at x. Since $U(N)$ is a compact Lie group and $\mathrm{Hol}_x(\mathcal{A}) \subset U(N)$ is a Lie subgroup, it is automatically a *closed* subgroup and hence compact. This proves the first claim.

Step 4: The holonomy Lie algebra inside $u(N)$. Write the internal curvature $\mathcal{F}_{\mu\nu} = \partial_\mu \mathcal{A}_\nu - \partial_\nu \mathcal{A}_\mu + i[\mathcal{A}_\mu, \mathcal{A}_\nu] = F^a_{\mu\nu} T_a$ (eq. (12.1)). Then $i\mathcal{F}_{\mu\nu} = F^a_{\mu\nu}(iT_a)$ is *skew*-adjoint. By Ambrose–Singer, the holonomy Lie algebra $\mathfrak{hol}_x(\mathcal{A})$ is the smallest Lie subalgebra of $u(N)$ containing all *parallel translates* of $i\mathcal{F}$ to x. Hence there is a Lie algebra homomorphism

$$\rho : \mathfrak{g} \longrightarrow u(N), \qquad \rho(T_a) \;:=\; -\,i\,T_a,$$

so that $[\rho(T_a), \rho(T_b)] = \rho([T_a, T_b])$ with the physics commutator $[T_a, T_b] = i f_{ab}{}^c T_c$ (Definition 12.5).

Step 5: Positivity of the (signed) Killing form on semisimple blocks. On a compact semisimple Lie algebra $\mathfrak{k}$, the Killing form $B(X,Y) = \mathrm{Tr}(\mathrm{ad}_X \mathrm{ad}_Y)$ is *negative*-definite. With the physics sign $\kappa(X,Y) := -B(X,Y) = -\mathrm{Tr}(\mathrm{ad}_X \mathrm{ad}_Y)$, κ is *positive*-definite. Concretely, choose an orthonormal Hilbert–Schmidt basis $\{E_j\}$ of $\mathrm{End}(\mathcal{H}_{\mathrm{int}})$ and compute, for Hermitian T,

$$\kappa(T,T) = -\sum_j \mathrm{Tr}\big([T,E_j]\,[T,E_j]^\dagger\big) = \sum_j \|[T,E_j]\|^2_{\mathrm{HS}} \;\geq\; 0,$$

with equality iff T is central on the semisimple block. Thus, on each semisimple component of the represented holonomy algebra, $\kappa_{ab} = -\mathrm{Tr}(\mathrm{ad}_{T_a}\mathrm{ad}_{T_b})$ is positive-definite (cf. the representation data in Definition 12.5).

Combining Steps 1–5 completes the proof: $\mathrm{Hol}_x(\mathcal{A})$ acts unitarily on the finite-dimensional fibre, is a Lie (hence closed) subgroup of $U(N)$ and therefore compact; its Lie algebra is represented by the skew-adjoints iT_a (modulo the sign convention), and the signed Killing form κ is positive on semisimple blocks. □

Remark 12.9 (Peter–Weyl machinery used throughout)**.** For a compact group K with Haar measure μ (normalized to $\mu(K) = 1$), the Peter–Weyl Theorem ([552]) asserts:

1. *Matrix–coefficient basis and density.* If $D^{(\alpha)} : K \to U(\mathcal{H}_\alpha)$ ranges over the finite-dimensional irreducible unitary representations, then the matrix coefficients $D^{(\alpha)}_{ij}(g)$ form an orthonormal basis of $L^2(K)$ and are uniformly dense in $C(K)$. Equivalently,
$$L^2(K) \;\cong\; \widehat{\bigoplus}_{\alpha\in\widehat{K}} \mathcal{H}_\alpha \otimes \mathcal{H}_\alpha^*,$$
with the left/right regular representations decomposing accordingly [446, 387].
2. *Schur orthogonality.* For $\alpha, \beta \in \widehat{K}$ and indices i, j, k, l,
$$\int_K \overline{D^{(\alpha)}_{ij}(g)}\, D^{(\beta)}_{kl}(g)\, d\mu(g) \;=\; \frac{\delta_{\alpha\beta}\,\delta_{ik}\,\delta_{jl}}{d_\alpha}, \qquad d_\alpha = \dim \mathcal{H}_\alpha [446,\ 387].$$
3. *Characters as class–function basis.* On the subspace of class functions, the irreducible characters $\chi_\alpha(g) = \mathrm{Tr}\, D^{(\alpha)}(g)$ are orthonormal:
$$\langle \chi_\alpha, \chi_\beta\rangle_K := \int_K \chi_\alpha(g)^* \chi_\beta(g)\, d\mu(g) = \delta_{\alpha\beta} [446,\ 387].$$

Within this chapter, compact holonomy yields a compact window K (Proposition 12.8), and the consequences above are invoked via the local statement Proposition 12.10. Two uses are central:

1. *Averaged overlap as an intertwiner.* In Proposition 12.46 and Corollary 12.48, the operator average $\int_K \pi_B(g)^\dagger \pi_A(g)\, d\mu(g)$ is the K-intertwiner $\mathrm{Hom}_K(\pi_B, \pi_A)$, which vanishes for inequivalent actions. This yields a vanishing averaged process-overlap and, by convex approximation on K, $\inf_{g\in K} \rho_{\mathrm{ens}}(A, B; g) = 0$, establishing Stage-II relational orthogonality (Theorem 12.42).
2. *Informational completeness on compact windows.* In Lemma 12.43, the Peter–Weyl spanning/orthogonality justifies that finite statistics of matrix-coefficient type (implemented through admissible control phases; Definition 2.114) separate the operator span $\mathcal{U}|_{\mathcal{H}_{\mathrm{probe}}}$ on the compact-holonomy window, consistent with Proposition 12.10.

Example ($K = SU(2)$). Irreducibles are spins $j = 0, \frac{1}{2}, 1, \ldots$ with $d_j = 2j + 1$ and characters
$$\chi_j(\theta) = \frac{\sin\Big((2j+1)\theta/2\Big)}{\sin(\theta/2)}.$$
Orthogonality of characters implies $\langle \chi_{j'}, \chi_j\rangle_{SU(2)} = 0$ for $j' \neq j$, activating Corollary 12.48.

Proposition 12.10 (Peter–Weyl decomposition and character meters)**.** *On $\mathcal{H}_{\mathrm{int}}(x;\lambda_{\mathrm{UV}},\ell)$ the representation of $\mathrm{Hol}_x(\mathcal{A})$ decomposes as a finite orthogonal sum of unitary irreps:*

$$\mathcal{H}_{\mathrm{int}}(x;\lambda_{\mathrm{UV}},\ell) \;\cong\; \bigoplus_{R\in\widehat{G}_x} \mathcal{H}_R, \qquad U(\gamma) \;\cong\; \bigoplus_{R\in\widehat{G}_x} U_R(\gamma),$$

with $\widehat{G}_x$ the finite set of irreps present. For any closed loop γ based at x, the characters $\chi_R(\gamma) := \mathrm{Tr}_{\mathcal{H}_R} U_R(\gamma)$ define bounded meters on $\mathfrak{P}$ and obey orthogonality relations (with respect to Haar measure on $\mathrm{Hol}_x(\mathcal{A})$) fiberwise; their expectations $\omega(\chi_R)$ serve as compactness diagnostics (boundedness/orthogonality).

Proof. Step 1: Finite fibre and compact unitary holonomy. By Lemma 12.7(a) the effective internal fibre $\mathcal{H}_{\mathrm{int}}(x;\lambda_{\mathrm{UV}},\ell)$ is finite-dimensional. By Proposition 12.8, the holonomy group $\mathrm{Hol}_x(\mathcal{A})$ acts unitarily on this fibre and is a (closed) Lie subgroup of $U(N)$ with $N = \dim \mathcal{H}_{\mathrm{int}}(x;\lambda_{\mathrm{UV}},\ell)$; in particular $\mathrm{Hol}_x(\mathcal{A})$ is a *compact* Lie group. The action is given by the unitary representation

$$U:\ \mathrm{Hol}_x(\mathcal{A}) \to U\Big(\mathcal{H}_{\mathrm{int}}(x;\lambda_{\mathrm{UV}},\ell)\Big), \qquad U(g) = \mathcal{P}\exp i\oint_\gamma \mathcal{A},$$

where g is the holonomy of $\mathcal{A}$ along a loop γ based at x (cf. eqs. (12.24) and (12.25)).

Step 2: Complete reducibility (Peter–Weyl/average projector). Let $G_x := \mathrm{Hol}_x(\mathcal{A})$ and let dg denote its normalized Haar measure. For any G_x-invariant subspace $W \subset \mathcal{H}_{\mathrm{int}}$, the orthogonal complement $W^\perp$ is also invariant because the representation is unitary. More generally, for any closed subspace $V \subset \mathcal{H}_{\mathrm{int}}$ one can construct the G_x-equivariant orthogonal projector by averaging:

$$\Pi_V^{G_x} \;:=\; \int_{G_x} U(g)\,\Pi_V\,U(g)^\dagger\,dg, \tag{12.26}$$

which satisfies $U(g)\Pi_V^{G_x} = \Pi_V^{G_x}U(g)$ for all $g \in G_x$. Standard compact-group representation theory then yields complete reducibility: $\mathcal{H}_{\mathrm{int}}$ is a finite orthogonal direct sum of irreducibles. Equivalently, the isotypic projectors may be written using characters of G_x (Schur orthogonality):

$$\Pi_R \;=\; d_R\int_{G_x}\chi_R(g)^*\,U(g)\,dg, \qquad \Pi_R^\dagger = \Pi_R,\ \ \Pi_R\Pi_S = \delta_{RS}\Pi_R,\ \ \sum_{R\in\widehat{G}_x}\Pi_R = \mathbf{1}, \tag{12.27}$$

where $d_R = \dim R$ and $\widehat{G}_x$ is the finite set of irreps present in U. This produces the orthogonal decomposition $\mathcal{H}_{\mathrm{int}} \cong \bigoplus_{R\in\widehat{G}_x}\mathcal{H}_R$ with $U(g) \cong \bigoplus_R U_R(g)$.

Step 3: Character boundedness and orthogonality. For any loop γ based at x, the holonomy $U_R(\gamma) \in U(\mathcal{H}_R)$ is unitary, hence

$$|\chi_R(\gamma)| = \big|\mathrm{Tr}_{\mathcal{H}_R}U_R(\gamma)\big| \;\le\; \mathrm{Tr}_{\mathcal{H}_R}\big(\mathbf{1}\big) = d_R. \tag{12.28}$$

Thus $\chi_R(\gamma)$ is a bounded observable (a bounded element of the represented algebra when realized as a Wilson loop; see below). Orthogonality of characters holds with respect to Haar measure on the compact group G_x:

$$\int_{G_x}\chi_R(g)\,\chi_S(g)^*\,dg \;=\; \delta_{RS}, \tag{12.29}$$

and more generally for matrix elements. Equation (12.29) underlies the projector formula (12.27) and the finite isotypic sum claimed.

Step 4: Characters as meters (Wilson loops) and BRST descent. Define the *Wilson loop* operators in representation R along γ:

$$W_R(\gamma) \;:=\; \mathrm{Tr}_{\mathcal{H}_R}\,\mathcal{P}\,\exp i\oint_\gamma \mathcal{A} \;\in\; \pi_\omega(\mathcal{A}) \subset \mathrm{End}(\mathcal{H}_{\mathrm{em}}), \tag{12.30}$$

with $\mathcal{A}_\mu$ the Operator Bundle Connection (Definition 6.192 and eq. (12.1)). Since $\mathcal{A}_\mu$ transforms as a connection and the trace is conjugation-invariant, $W_R(\gamma)$ is gauge invariant (BRST-closed), and by (12.28) it is bounded with $\|W_R(\gamma)\| \le d_R$. Therefore $W_R(\gamma)$ is a legitimate *meter* on the certified plateau $\mathfrak{P}$. By the cohomological descent lemma (Lemma 12.2), meters built from ∇_μ and $\mathcal{A}_\mu$ (and their tag sandwiches) descend to $\mathcal{H}_{\mathrm{phys}}$ and are Nielsen-flat on $\mathfrak{P}$; in particular, their expectations are computed by the GNS/Born rule

$$\langle W_R(\gamma)\rangle_\omega \;=\; \omega^{\mathring{\Psi}}\Big(W_R(\gamma)\Big), \tag{12.31}$$

which matches the *dialectic/Born* semantics used throughout (Definition 1.18).

Step 5: Fiberwise orthogonality and compactness diagnostics. On the compact group G_x the family $\{\chi_R\}_{R\in\widehat{G}_x}$ is orthonormal in $L^2(G_x, dg)$ by (12.29). Pulling back along the holonomy map $\gamma \mapsto g_\gamma \in G_x$, one obtains *fiberwise* orthogonality tests by averaging characters over families of loops that sample G_x (operationally: average over a loop ensemble whose induced measure approximates Haar; cf. the meters ledger $\mathcal{W}$ and spectral certificate $\mathcal{Z}$ in Ch. 8). Boundedness (12.28) and orthogonality (12.29), together with the compactness of G_x (Proposition 12.8), yield the advertised *compactness diagnostics*:

(i) bounded character meters $\|W_R(\gamma)\| \le d_R$,

(ii) orthogonality-based isotypic extraction via (12.27).

By Lemma 12.2, these diagnostics are stable under tag variations and descend to $\mathcal{H}_{\mathrm{phys}}$.

Putting Steps 1–5 together gives the stated finite Peter–Weyl ([552]) decomposition on $\mathcal{H}_{\mathrm{int}}(x;\lambda_{\mathrm{UV}},\ell)$ and the associated character meters with their boundedness/orthogonality properties. □

Definition 12.11 (GNS energy on gauge directions and current algebra)**.** *GNS substrate and energy on gauge directions.* On the GNS arena $(\pi_\omega, \mathcal{H}_{\mathrm{em}}, |\Omega_\omega\rangle)$ (Definition 1.34) equip $\mathrm{End}(\mathcal{H}_{\mathrm{em}})$ with the GNS inner product

$$\langle X, Y\rangle_\omega \;:=\; \omega\Big(X^\dagger Y\Big), \qquad \|X\|_\omega^2 = \langle X, X\rangle_\omega,$$

and, for Hermitian gauge generators $T = T^\dagger$ in the represented internal fibre (Definition 12.5), set the *GNS energy* $\|T\|_\omega^2 := \omega(T^2)$. The compact-holonomy representation implies all gauge directions act by unitary one-parameter groups on each effective fibre (Proposition 12.8 and Lemma 12.7).

emphImproved Noether currents (Ch. 6/9) and BRST-safe form. Let S be the Tantrakyma action written with the realized weave configuration-compatible split connection

$$\nabla_\mu \;=\; \nabla_\mu^{(\mathrm{rel})}[\bar{\Sigma}] \;+\; \Big(\partial_\mu + i[\mathcal{A}_\mu, \cdot]\Big) \quad (\textit{Definition 6.192and eq. (6.314)}).$$

In an adjoint basis $\{T_a\}$ we write $\mathcal{A}_\mu = A^a_\mu T_a$. Define the *canonical* gauge current by

$$J^\mu_{a,\mathrm{can}}(x) \;:=\; \frac{\delta S}{\delta A^a_\mu(x)}.$$

The *improved* current used on the certified plateau (Ch. 6) is obtained by adding the divergence of a local antisymmetric superpotential and the BRST/gauge-fixing improvement (so Ward functionals are adjoint-covariant and tag-safe):

$$J^\mu_{a,\mathrm{imp}} \;:=\; J^\mu_{a,\mathrm{can}} \;+\; \nabla_\nu\, X^{[\mu\nu]}_a \;-\; \zeta\,\Lambda^\mu_a, \qquad X^{[\mu\nu]}_a = -X^{[\nu\mu]}_a, \tag{12.32}$$

with ζ the BRST operator and Λ^μ_a the gauge-fixing descent current from eq. (6.290). On the diagnostic realized reference configuration, the *adjoint Ward identity* then takes the standard BRST-safe form

$$\nabla^{(\mathrm{ad})}_\mu\, \hat{J}^\mu_{a,\mathrm{imp}} = \sum_k \epsilon_k\, \mathcal{S}_{k,a} + \zeta\,\eta_a.$$

(See eqs. (6.293) and (6.378), Lemma 6.273, and Definition 6.295.) In the BEH/SSB regime of Ch. 9, the same improvement absorbs Goldstone mixings so that the *gauge-improved* current is covariantly conserved on shell (cf. the BRST-improved Ward readouts in Sections 12.2 and 12.3).

Ward identity (scale-tagged, adjoint form). For any test field $\alpha^a(x)$ supported in the diagnostic realized reference configuration arena, the tagged Ward functional reads

$$\mathcal{W}[\alpha] \;:=\; \int d\mu_{\bar{\Sigma}}(x)\; \alpha^a(x)\, \big(\nabla^{(\mathrm{ad})\,\mu} J^a_\mu\big)(x) \;=\; 0 \quad \text{on } \mathcal{H}_{\mathrm{phys}}, \tag{12.33}$$

where $\nabla^{(\mathrm{ad})}_\mu$ is the adjoint-covariant derivative built from $\mathcal{A}_\mu$ (eq. (12.1) and Definition 7.567). By Lemma 12.2, (12.33) *descends* to $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ and is Nielsen-flat (gauge-parameter independent). Typed causal support used in the derivation is guaranteed by cone-preserving hyperbolicity (Proposition 12.3).

Rep-orthogonality and positivity of the current two-point. Choose a compact-holonomy orthonormal adjoint basis $\{T_a\}$ with $\kappa_{ab} = -\mathrm{Tr}(\mathrm{ad}_{T_a}\mathrm{ad}_{T_b}) = \delta_{ab}$ on semisimple blocks (Proposition 12.8). Then, on the certified plateau $\mathfrak{P}$, the improved currents have a *rep-orthogonal, positive* two-point function:

$$\Big\langle J^a_\mu(x)\, J^b_\nu(y)\Big\rangle_\omega \;=\; \delta^{ab}\,\Pi_{\mu\nu}(x,y), \qquad \Pi \;\succeq\; 0, \tag{12.34}$$

where positivity is with respect to the GNS pairing and follows from unitarity of the holonomy representation (Proposition 12.8) together with the Peierls/BRST construction (Lemma 12.2) and the causal propagator support (Proposition 12.3). The rep-orthogonality δ^{ab} is with respect to κ_{ab} and the compact Peter–Weyl decomposition on each finite fibre (Proposition 12.10).

KL positivity. Equation (12.34) feeds the Källén–Lehmann (KL) representation for two-point functions in each irrep channel (Ch. 6 spectral calculus on $(\mathcal{X}_q, \bar{\Sigma})$), yielding non-negative spectral data and stable residues on $\mathfrak{P}$ (Nielsen-flat by Lemma 12.2); cf. the explicit KL statement in Proposition 12.12.

Proposition 12.12 (KL positivity and Ward well-posedness)**.** *On $\mathfrak{P}$, for each irrep $R \in \widehat{G}_x$ appearing in Proposition 12.10, the two-point functions admit a Källén–Lehmann representation with non-negative weights,*

$$\Delta_R(q) \;=\; \frac{Z_R}{q^2 + i\epsilon} \;+\; \int_0^\infty \frac{\rho_R(\mu^2)\, d\mu^2}{q^2 - \mu^2 + i\epsilon}, \qquad Z_R, \rho_R \geq 0,$$

and the associated Ward identities descend to $\mathcal{H}_{\text{phys}}$ and are gauge-parameter independent (Nielsen-flat).

Proof. Step 1: Isotypic projection and positivity in each R-channel. By Proposition 12.10 the holonomy action on the finite fibre $\mathcal{H}_{\text{int}}(x; \lambda_{\text{UV}}, \ell)$ (Ch. 7 finiteness, Lemma 12.7) is a compact-unitary representation (Ch. 6 compactness input via Proposition 12.8). Let Π_R be the isotypic projector (Peter–Weyl; Proposition 12.10, (12.27)) and define $J_\mu^{(R)} := \Pi_R J_\mu \Pi_R$, where J_μ is the *improved* Noether current of Ch. 6/9 (Definition 12.11, (12.32)). With the GNS pairing $\langle X, Y\rangle_\omega = \omega(X^\dagger Y)$ (Definitions 1.34 and 12.11), the current two-point is *rep-orthogonal and positive*:

$$\Big\langle J_\mu^{(R)}(x)\, J_\nu^{(R)}(y) \Big\rangle_\omega \;=\; \Pi_{\mu\nu}^{(R)}(x, y), \qquad \Pi^{(R)} \succeq 0, \tag{12.35}$$

by Definition 12.11 and the orthogonality of adjoint directions fixed with the compact (signed) Killing form in Proposition 12.8.

Step 2: Hyperbolicity, causal support, and band instruments (Chs. 6–7). Cone-preserving hyperbolicity on $(\mathcal{X}_q, \bar{\Sigma})$ gives retarded/advanced Green operators with $\bar{\Sigma}$-cone support (Ch. 6; Proposition 12.3). On the diagnostic realized reference configuration $\mathfrak{P}$, the IR window is the spectral projector of a self-adjoint, nonnegative $\bar{\Sigma}$-elliptic operator $\mathcal{K}$ (Ch. 7 band instruments),

$$K_{\text{IR}}^{(\ell)} = \chi_{[0, \Lambda(\ell)]}(\mathcal{K}), \tag{12.36}$$

and varies continuously with the knobs by the audit (Definition 9.166). Thus, for any test current f^μ supported in the diagnostic realized reference configuration arena,

$$\langle f, \Pi^{(R)} f\rangle := \int f^\mu(x)\, \Pi_{\mu\nu}^{(R)}(x, y)\, f^\nu(y)\, d\mu_{\bar{\Sigma}}(x)\, d\mu_{\bar{\Sigma}}(y) \;\geq 0,$$

so $\Pi^{(R)}$ is a positive-type distribution with causal support (Ch. 6 Peierls/microcausality; Proposition 12.3).

Step 3: KL representation and non-negative weights (Ch. 6). By the spectral theorem on the certified background and Bochner–Schwartz positivity, $\Pi^{(R)}$ admits a positive spectral measure $d\rho_R(\mu^2)$; after local Fourier transform (in a stationary patch or via microlocal spectral calculus on $\mathfrak{P}$),

$$\Pi^{(R)}(q) \;=\; \int_0^\infty \frac{d\rho_R(\mu^2)}{q^2 - \mu^2 + i\epsilon} \;+\; (\text{possible simple pole contributions}).$$

Consequently the (time-ordered/Feynman) two-point in the R-channel takes the Källén–Lehmann form

$$\Delta_R(q) \;=\; \frac{Z_R}{q^2 + i\epsilon} \;+\; \int_0^\infty \frac{\rho_R(\mu^2)\, d\mu^2}{q^2 - \mu^2 + i\epsilon}, \qquad Z_R, \rho_R \geq 0,$$

where Z_R is the non-negative one-particle residue (norm square of the corresponding state), and ρ_R is the non-negative spectral density. Positivity derives from (12.35) and the compact/unitary realization of holonomy (Ch. 6 via Proposition 12.8); causal support used in the dispersion argument is supplied by Proposition 12.3.

Step 4: Ward descent and Nielsen-flatness (Chs. 6,7,9). The improved current (12.32) satisfies the adjoint Ward identity on the diagnostic realized reference configuration (Definition 12.11; Ch. 6 Ward formalism), and by the *cohomological descent* lemma (Lemma 12.2, Ch. 7 tag commutation) it descends to $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$ and is stable under tag sandwiches and band variations (Definition 9.166). Gauge-parameter independence follows from the Nielsen identity (Lemma 12.2, (12.12)), which in each R-channel reads

$$\partial_\xi \Gamma_R^{(2)} = \Gamma_R^{(2)} \mathcal{N}_R + \tilde{\mathcal{N}}_R \Gamma_R^{(2)},$$

with $\mathcal{N}_R, \tilde{\mathcal{N}}_R$ built from BRST-exact composites (Ch. 6 BRST algebra). Therefore the pole locations (zeros of $\det \Gamma_R^{(2)}$) and residues of $\Delta_R = (\Gamma_R^{(2)})^{-1}$ are *Nielsen-flat* (gauge-parameter independent). In the BEH/SSB regime (Ch. 9), the improvement (12.32) removes Goldstone mixing and preserves the adjoint Ward form after symmetry breaking, so the same descent and flatness hold in the broken phase.

Step 5: Audit continuity (Ch. 7). By Lemma 12.7(b) and Definition 9.166, the isotypic projector Π_R, residue Z_R, and spectral measure ρ_R vary continuously with $(\lambda_{\text{UV}}, \ell)$ on $\mathfrak{P}$, and the Ward identities remain valid on $\mathcal{H}_{\text{phys}}$ under these variations (Lemma 12.2).

Combining Steps 1–5 yields the claimed KL representation with $Z_R, \rho_R \geq 0$ and the well-posed, gauge-parameter–independent Ward structure on $\mathcal{H}_{\text{phys}}$. □

Definition 12.13 (Tetralemma compactness meter)**.** Work on a certified plateau $\mathfrak{P}$ with UV/IR tags $P_x(\lambda_{\text{UV}})$, $K_{\text{IR}}^{(\ell)}$ (band functor $\mathcal{B}_{\lambda_{\text{UV}},\ell}$) and a BRST-closed state ω (Chs. 6–8; Definitions 7.29 and 12.67, Proposition 7.30, and Postulate 12.2). The four-outcome decision schema (Thesis/Antithesis/Synthesis/Holothesis) is the tetralemma of Ch. 2 (Chapter 2 and Section 2.1), realized by operator *effects* with Born valuations as in the local logical projectors and their complementary (parasymmetric) probes (Definitions 2.1, 2.7, 2.16 and 2.17 and eq. (2.57)).

Let $\{\gamma_j\}_{j=1}^J$ be a finite family of based loops at x and let $\widehat{G}_x$ denote the (finite) set of irreps present in the fibrewise Peter–Weyl decomposition (Proposition 12.10). For each $R \in \widehat{G}_x$ define the *normalized, self-adjoint character meters*

$$\widehat{\chi}_R(\gamma_j) \;:=\; \frac{1}{d_R} \operatorname{Re} W_R(\gamma_j), \qquad W_R(\gamma_j) \;:=\; \operatorname{Tr}_{\mathcal{H}_R} \mathcal{P} \exp\Big(i \oint_{\gamma_j} \mathcal{A} \Big),$$

regarded as bounded operators on $\mathcal{H}_{\text{em}}$ via the band functor (positivity/*-preservation) and BRST descent to $\mathcal{H}_{\text{phys}}$ (Lemma 12.2; see also Definitions 6.295 and 7.567). (Ch. 2 provides the operator semantics of effects and their Born readings that we adopt here; cf. the duality and parasymmetry constructions Definitions 2.7, 2.12 and 2.17 and eq. (2.57).)

Average them into a bounded self-adjoint *compactness score operator*

$$\mathsf{S}_{\text{cmp}} \;:=\; \frac{1}{J} \sum_{j=1}^{J} \frac{1}{|\widehat{G}_x|} \sum_{R \in \widehat{G}_x} \widehat{\chi}_R(\gamma_j), \qquad -1 \leq \mathsf{S}_{\text{cmp}} \leq 1.$$

Fix thresholds $\alpha, \beta \in (0,1)$ from the holosymmetric calibration protocol (Ch. 8; $(\mathcal{W}, \mathcal{Z})$ meters). Using spectral calculus on $\mathsf{S}_{\mathrm{cmp}}$, define the **effect** operators

$$\begin{aligned} E_{\mathsf{Th}} &:= \mathbf{1}_{[\alpha,1]}\big(\mathsf{S}_{\mathrm{cmp}}\big), \\ E_{\mathsf{HoloTh}} &:= \mathbf{1}_{(-\infty,\alpha)}\big(\mathsf{S}_{\mathrm{cmp}}\big) \;\wedge\; \mathbf{1}_{[0,\beta]}\big(\mathsf{D}_{\mathrm{nc}}\big), \\ E_{\mathsf{AntiTh}} &:= \mathbf{1}_{(\beta,\infty)}\big(\mathsf{D}_{\mathrm{nc}}\big), \\ E_{\mathsf{SynTh}} &:= \mathbf{1}_{(-\infty,\alpha)}\big(\mathsf{S}_{\mathrm{cmp}}\big) \;\wedge\; \mathbf{1}_{(\beta_-,\beta_+]}\big(\mathsf{D}_{\mathrm{nc}}\big), \end{aligned}$$

where "$\wedge$" denotes the product (meet) of commuting effects after band-averaging (the band functor ensures positivity/*-preservation), and where $0 < \beta_- < \beta < \beta_+$. We define the *non-compactness witness* as

$$\mathsf{D}_{\mathrm{nc}} \;:=\; \Big(\sum_a \big(\Pi_{\mathrm{int}} T_a \Pi_{\mathrm{int}}\big)^2 \Big)^{1/2},$$

the operator on the finite internal fibre $\mathcal{H}_{\mathrm{int}}(x; \lambda_{\mathrm{UV}}, \ell) = \mathrm{Ran}\big(K_{\mathrm{IR}}^{(\ell)} P_x(\lambda_{\mathrm{UV}}) \Pi_{\mathrm{Tan}}\big)$ (Definition 12.6 and Lemma 12.7). Here $T_a = T_a^\dagger$ are the represented Hermitian directions (Definition 12.5); D_{nc} is bounded when the holonomy is compact, since $\mathrm{Hol}_x(\mathcal{A}) \subset U(N)$ on the fibre (Proposition 12.8). Holothesis here aligns with the holosymmetric layer of Ch. 2 (dual-register scaffold and controlled evolution) used to formalize "suspend the verdict under admissibility" (Sections 2.3.1, 2.3.2 and 2.4, Definitions 2.101 and 2.102, and eq. (2.68)).

Finally, for each corner $M \in \{\mathsf{Th}, \mathsf{AntiTh}, \mathsf{SynTh}, \mathsf{HoloTh}\}$ define the Born weight

$$v_M(\rho) \;:=\; \mathrm{Tr}\big(E_M\, \rho\big) \;\in\; [0,1].$$

This agrees with the Born-valued effects introduced in Ch. 2 (logical projectors on the $(0,1)$-sector and parasymmetric projectors; cf. Definitions 2.1 and 2.17 and eq. (2.57)). Define the clipped contrast

$$\Lambda^{[\mathrm{cmp}]} \;:=\; \Big[\, v_{\mathsf{Th}} - \big(v_{\mathsf{AntiTh}} + v_{\mathsf{SynTh}}\big) \,\Big]_+ \;\in\; [0,1].$$

We report $\Lambda^{[\mathrm{cmp}]}$ together with the Ward and spectral ledgers $(\mathcal{W}, \mathcal{Z})$ (Ch. 8) and with the character orthogonality checks from Proposition 12.10.

Remark 12.14 (Correctness, descent, and audit stability)**.** (i) *Operator effects & Born semantics.* Each E_M is an effect $0 \le E_M \le \mathbf{1}$ on $\mathcal{H}_{\mathrm{em}}$, so $v_M(\rho) = \mathrm{Tr}(E_M \rho)$ is a legitimate probability; this matches the Ch. 2 realization where Thesis/Antithesis are logical projectors on the $(0,1)$-sector and Synthesis uses parasymmetric effects (Definitions 2.1 and 2.17 and eq. (2.57)). (ii) *BRST/tag descent and Nielsen-flatness.* Wilson-loop/character meters are gauge-invariant; by the BRST chain-map and band functor properties, they *descend* to $\mathcal{H}_{\mathrm{phys}}$ and are gauge-parameter independent on $\mathfrak{P}$ (Nielsen-flat) (Lemma 12.2). (iii) *Compactness diagnostics.* On finite fibres, $\mathrm{Hol}_x(\mathcal{A}) \subset U(N)$ is compact and represented unitarily (Proposition 12.8); Peter–Weyl then gives bounded characters and orthogonality fibrewise (Proposition 12.10), which $\mathsf{S}_{\mathrm{cmp}}$ aggregates. Large D_{nc} (beyond β) flags antithesis signals incompatible with compact holonomy. (iv) *Audit continuity.* All meters

are built from band-tagged operators; the audit ensures norm-continuity in $(\lambda_{\mathrm{UV}}, \ell)$ and commutation with the physical projector, so $\Lambda^{[\mathrm{cmp}]}$ is tag-stable (Chs. 6–7; Lemma 12.7 and $\mathsf{P}_{\mathrm{phys}}$ commutation). (v) *Holosymmetric alignment.* The Holothesis branch is implemented in the Ch. 2 holosymmetric scaffold by dual registers and controlled recombination (Sections 2.3.1, 2.3.2 and 2.4, Definitions 2.101 and 2.102, and eq. (2.68)), matching the HSP.

Thesis and Antithesis (local duality). In the $(0,1)$-sector of the modal space $\mathbb{C}^4$, Thesis and Antithesis are realized by the distinguished basis vectors $\{|e_1\rangle, |e_2\rangle\}$ with logical projectors $P_T := |e_1\rangle\langle e_1|$ and $P_A := |e_2\rangle\langle e_2|$. Their Born-valued warrants are $v_{\mathrm{Thesis}}(\rho) = \mathrm{Tr}(P_T\rho)$ and $v_{\mathrm{Antithesis}}(\rho) = \mathrm{Tr}(P_A\rho)$ (Definition 2.1). The *local involutive duality operator* $D^{(\mathfrak{B}_x)}$ swaps the two ($D^{(\mathfrak{B}_x)}|e_1\rangle = |e_2\rangle$, $D^{(\mathfrak{B}_x)}|e_2\rangle = |e_1\rangle$), is Hermitian and involutive ($D^{(\mathfrak{B}_x)} = D^{(\mathfrak{B}_x)\dagger}$, $D^{(\mathfrak{B}_x)2} = \mathbf{1}$), and acts trivially off the $(0,1)$-plane (Definition 2.7). A coherent multi-site realization uses the global swap $\hat{D}^{(n)}_{\mathrm{PQS}} = \bigotimes_{i=1}^{n} D^{(\mathfrak{B}_{x_i})}$ (Definition 2.12).

Synthesis (parasymmetry). Chapter 2 fixes the complementary synthesis basis $|\pm\rangle = \frac{1}{\sqrt{2}}\big(|e_1\rangle \pm |e_2\rangle\big)$ spanning the same $(0,1)$-sector, with projectors P_{para} and P_{antipara} resolving the parasymmetric modalities (Definitions 2.16 and 2.17). The *degree of synthesis* is the Born valuation $v_{\mathrm{Synthesis}}(\rho) = \mathrm{Tr}\big(P_{\mathrm{para}}\rho\big)$ (cf. eq. (2.57)). The same logic admits relational (entangled) realizations, e.g. a Bell-type parasymmetric state $|\Psi_{\mathrm{para}}\rangle_{xx'}$ with corresponding global projector $P^{(xx')}_{\mathrm{para}}$ that operationalizes synthesis as *coherence* between Thesis/Antithesis branches across sites (Definition 2.17).

Holothesis (holosymmetry). Holothesis is the *controlled interferometric readout* that tests which global evolution is coherently realized—i.e. a meta-test beyond the base Thesis/Antithesis axis. Chapter 2 constructs the dual-register scaffold and the controlled "path" unitaries to compare alternative histories (Sections 2.3.1, 2.3.2 and 2.4, Definitions 2.101 and 2.102, and eq. (2.68)); Chapter 12 then compresses this to the BRST-physical diagnostic realized reference configuration and packages it as a holosymmetric *coherence effect* $E^{(G)}_{\mathrm{holo}}$ with projector $\mathcal{P}^{\mathrm{coh}}_G$ used operationally in selection (cf. Definition 12.13).

Overlay and HSP as the Holothesis gate. The holosymmetric overlay of Chapter 6 supplies the visibility plane $(\Lambda_{\mathrm{law}}, \Lambda_{\mathrm{state}})$ and certifies the *plateau* where precise meters are admissible. If either visibility falls below threshold, the protocol must *suspend* any verdict about a target claim φ; in tetralemma terms, the Holothesis effect E_{HoloTh} fires (Definition 12.13). Stage I of the Holosymmetric Selection Principle (HSP) in Chapter 12 then cleanly separates: (i) decisive refutations (AntiThesis; e.g. BRST obstruction by anomalies), (ii) decisive affirmations (Thesis; all viability checks pass on the plateau), and (iii) *inadmissible/inconclusive* runs (Holothesis; non-plateau or scores in the indifference region), with descent to $\mathcal{H}_{\mathrm{phys}}$ ensured by the BRST chain-map and tag-commutation (Lemma 12.2) and spectral/Ward hygiene by KL positivity (Proposition 12.12).

Decision layer versus diagnostic realized reference configuration quality. The four law–state quadrants (Ch. 6) classify *diagnostic realized reference configuration quality*—how visible the

laws and prepared states are—while the tetralemma's four corners are *decision outcomes* for a concrete proposition φ (e.g. "holonomy is compact"). Once the overlay certifies the plateau (high–high quadrant), one applies a tetralemma protocol: build self-adjoint meters (e.g. normalized characters and a non-compactness witness), assemble effects E_M, and read Born weights

$$v_M(\rho) = \mathrm{Tr}\big(E_M \rho\big) \in [0,1], \qquad M \in \{\mathsf{Th}, \mathsf{AntiTh}, \mathsf{SynTh}, \mathsf{HoloTh}\}$$

(Definition 12.13). Thus, Thesis/Antithesis are the $\{|e_1\rangle, |e_2\rangle\}$ axis with Born truth, Synthesis is the complementary parasymmetric probe (locally $|\pm\rangle$ and relationally via entanglement), and Holothesis is the holosymmetric interferometric comparison of global histories—precisely realized by the local swap $D^{(\mathfrak{B}_x)}$, the parasymmetric projectors, and the holosymmetric effect, as formalized in Chapter 2 and deployed in Chapter 12 (Definitions 2.7, 2.17 and 12.13).

Definition 12.15 (Boundary compactness)**.** Work on a boundary plateau $\mathfrak{P}_\partial$ equipped with a boundary–compatible operator-bundle connection $\mathcal{A}_\mu$ in the sense of Definition 12.4. Let $\iota : \partial\mathcal{X}_q \hookrightarrow \mathcal{X}_q$ be the inclusion and $\Pi_\parallel$ the projector onto tangential directions. Define the tangential connection and curvature by

$$\mathcal{A}_\parallel \;:=\; \iota^*\mathcal{A} \;=\; \Pi_\parallel \,\mathcal{A}\, \Pi_\parallel, \qquad \mathcal{F}_\parallel \;:=\; d_\parallel \mathcal{A}_\parallel + i\,\mathcal{A}_\parallel \wedge \mathcal{A}_\parallel \quad \text{(cf. Definition 6.192 and eq. (12.1))}.$$

Choose boundary UV/IR instruments $P_{x,\partial}(\lambda_{\mathrm{UV}})$ and $K^{(\ell)}_{\mathrm{IR},\partial}$ (tangential band functor) as in Definition 12.4 and the audit formalism of Ch. 7 (Definition 9.166); by tag–commutation on the certified plateau,

$$[\,\nabla_\parallel, P_{x,\partial}(\lambda_{\mathrm{UV}})\,] \;=\; [\,\nabla_\parallel, K^{(\ell)}_{\mathrm{IR},\partial}\,] \;=\; 0 \quad \text{on } \mathfrak{P}_\partial \quad \text{(cf. eq. (12.17) and Lemma 12.2)}.$$

The *boundary internal fibre* at $x \in \partial\mathcal{X}_q$ is

$$\mathcal{H}^\partial_{\mathrm{int}}(x;\lambda_{\mathrm{UV}},\ell) \;:=\; \mathrm{Ran}\Big(K^{(\ell)}_{\mathrm{IR},\partial}\, P_{x,\partial}(\lambda_{\mathrm{UV}})\, \Pi_{\mathrm{Tan}}\Big) \quad \text{(finite by Lemma 12.7)}.$$

For any tangential loop $\gamma_\parallel \subset \partial\mathcal{X}_q$ based at x, the *tangential holonomy* is the unitary

$$U_\parallel(\gamma_\parallel) \;:=\; \mathcal{P}\exp i\oint_{\gamma_\parallel} \mathcal{A}_\parallel \in U\Big(\mathcal{H}^\partial_{\mathrm{int}}(x;\lambda_{\mathrm{UV}},\ell)\Big),$$

unitarity following from Hermitian internal generators (Definition 12.5). The associated boundary holonomy group

$$\mathrm{Hol}_x(\mathcal{A}_\parallel) \;\subset\; U\Big(\mathcal{H}^\partial_{\mathrm{int}}(x;\lambda_{\mathrm{UV}},\ell)\Big)$$

is a Lie subgroup of a compact unitary group and hence compact, by the same argument as in the bulk (Proposition 12.8, applied tangentially). Consequently the boundary representation admits a finite Peter–Weyl decomposition with characters $\chi_R(\gamma_\parallel) = \mathrm{Tr}_{\mathcal{H}_R} U_{R,\parallel}(\gamma_\parallel)$, bounded and orthogonal fibrewise (Proposition 12.10).

Boundary Ward hygiene is provided by the tagged *boundary* identity

$$\nabla_\parallel \cdot J^{(\partial)}_\parallel \;=\; S_\partial,$$

from Definition 12.4, which descends to $\mathcal{H}_{\text{phys}}$ and is Nielsen-flat by the cohomological descent lemma (Lemma 12.2) and the cone-preserving hyperbolicity of the tangential dynamics (Proposition 12.3, applied to $\bar{\Sigma}$-causal tangential propagation).

Define the *boundary compactness meters* by replacing bulk loops γ with $\gamma_{\|}$, bulk holonomies with $U_{\|}$, and the bulk fibre with $\mathcal{H}^{\partial}_{\text{int}}$ in the construction of Proposition 12.10 and Definitions 12.11 and 12.13. The resulting boundary compactness visibility

$$\Lambda^{[\text{cmp}]}_{\partial} \in [0,1]$$

is read via the same tetralemma effects (Thesis/Antithesis/Synthesis/Holothesis) now applied tangentially; it is tag-stable under the audit (Definition 9.166 and Lemma 12.7) and inherits KL positivity and Ward well-posedness from the boundary two-point calculus (Proposition 12.12).

Checklist (diagnostic realized reference configuration tests for B). Given $G \subset G_{\text{int}}$, on $\mathfrak{P}$ verify:

(B.1) *Unitary holonomy:* compute $\text{Hol}_x(\mathcal{A}) \subset U\Big(\mathcal{H}_{\text{int}}(x;\lambda_{\text{UV}},\ell)\Big)$ and its Lie algebra generators iT_a.

(B.2) *Finite fibre & audit:* establish $\dim \mathcal{H}_{\text{int}} < \infty$ and continuity in $(\lambda_{\text{UV}},\ell)$ (Lemma 12.7).

(B.3) *Peter–Weyl:* decompose into irreps $\bigoplus_R \mathcal{H}_R$, verify character orthogonality/boundedness (Proposition 12.10).

(B.4) *KL positivity:* extract $(Z_R,\rho_R) \geq 0$ from two-point functions (Proposition 12.12).

(B.5) *Ward descent/Nielsen:* apply Lemma 12.2 to confirm descent to $\mathcal{H}_{\text{phys}}$ and gauge-parameter independence.

(B.6) *Tetralemma:* compute $v_M(\rho) = \text{Tr}(E_M\rho)$ and $\Lambda^{[\text{cmp}]}$ (Definition 12.13); record in $\mathcal{W},\mathcal{Z}$.

(B.∂) *Boundary (if present):* repeat items (1)–(6) with tangential data per Definition 12.15.

Constraint C: Symmetry of the Tantrakyma dynamics

A candidate group must be a symmetry of the Tantrakyma action S_Θ, i.e. preserved by the full covariant equations of motion. Concretely, the Euler–Lagrange law for $\hat{\boldsymbol{\Theta}}$,

$$\bar{\Sigma}^{\mu\nu}\nabla_\mu\nabla_\nu\hat{\boldsymbol{\Theta}} + (\nabla_\mu\bar{\Sigma}^{\mu\nu})\nabla_\nu\hat{\boldsymbol{\Theta}} = \frac{\delta\mathcal{V}_\Theta}{\delta\hat{\boldsymbol{\Theta}}^\dagger},$$

is formulated with exactly the same ∇_μ that carries the connection $\mathcal{A}_\mu$, so invariance of S_Θ and Noether currents is evaluated intrinsically on the operator bundle (cf. Definition 9.335 and Section 9.18.10; see also the gaiamiti/chronogeometric tensors and connection in Definitions 5.26 and 6.7, the unified field doublet in Definition 6.1, and the BRST layer for gauge invariance in Section 6.10 and Definition 12.58). The causal background that enters via $\bar{\Sigma}$ is the phase-invariant chronogeometry (cf. Definition 5.122 and Corollary 5.126).

Under these three filters the connection's Lie algebra is forced to be a compact subalgebra of $\text{End}(\mathcal{H}^{(\mathfrak{B}_x)}_{\text{mod}})$ that genuinely acts on Tantrakyma sections and their couplings (cf.

typed/ledger observables and projectors in Definition 10.80 and the typed projectors' band in Definition 9.116; see also the local observable layer of Chapter 10 for spacetime insertions Definition 5.139). When the Tantrakyma reps are those of the Standard Model, the connection decomposes canonically

$$\mathcal{A}_\mu(x) = g_s\, G^a_\mu T_a + g\, W^i_\mu \tau_i + g' B_\mu Y,$$

identifying gluons, weak bosons, and the hypercharge field; the dynamics follow from the Yang–Mills curvature $\mathcal{F}_{\mu\nu} = \partial_\mu \mathcal{A}_\nu - \partial_\nu \mathcal{A}_\mu + i[\mathcal{A}_\mu, \mathcal{A}_\nu]$ (Definitions 12.95 and 12.96). Cohomological admissibility of insertions within this sector is read at the BRST layer (Chs. 6–9), where generic observables $\mathcal{O}$ (ghost number 0, BRST-closed) project to the physical sector and BRST-exacts decouple (cf. Section 6.10 and Lemma 6.210).

Definition (Realized weave configuration-compatible G-symmetry of the Tantrakyma sector). A candidate $G \subset G_{\mathrm{int}}$ is a symmetry of the Tantrakyma action S_Θ if, when written with the same realized weave configuration-compatible split connection

$$\nabla_\mu \;=\; \nabla^{(\mathrm{LC})}_\mu \;+\; i\,\mathrm{ad}_{\mathcal{A}_\mu} \qquad \text{(Definitions 6.192 and 12.1)},$$

the following hold on the certified plateau: (i) the Euler–Lagrange (EL) operator for $\hat{\mathbf{\Theta}}$,

$$\mathcal{E}_\Theta[\hat{\mathbf{\Theta}}] := \bar{\Sigma}^{\mu\nu} \nabla_\mu \nabla_\nu \hat{\mathbf{\Theta}} \;+\; (\nabla_\mu \bar{\Sigma}^{\mu\nu})\, \nabla_\nu \hat{\mathbf{\Theta}}, \tag{12.37}$$

transforms covariantly in the representation carried by $\hat{\mathbf{\Theta}}$ (Ch. 6 cones/transport; Definitions 5.26 and 6.7 and Proposition 12.3); (ii) the Tantrakyma potential and couplings $\mathcal{V}_\Theta$ (including Yukawas if present) are G-invariant when expressed with ∇_μ and $\mathcal{A}_\mu$ (Chs. 9–10; Definitions 6.1 and 9.335); (iii) the improved Noether current $J^a_\mu = \delta S/\delta A^\mu_a$ satisfies the adjoint Ward identity $\nabla^{(\mathrm{ad})\mu} J_\mu = 0$ in bulk and its tagged boundary variant (Ch. 6; Section 9.18.10 and Definitions 12.4 and 12.11). All statements are implemented as BRST-closed meters, hence descend to $\mathcal{H}_{\mathrm{phys}}$ and are Nielsen-flat (Lemma 12.2 and Proposition 12.12).

Meters (operator–geometric, diagnostic realized reference configuration-ready).

Definition 12.16 (Meters (operator–geometric, diagnostic realized reference configuration–ready))**.** Work on a certified plateau so that physical readouts are BRST/Nielsen–stable (Sections 6.12.4 and 12.3 and Proposition 12.12). Let $\bar{\Sigma}$ be a Cauchy slice with induced volume form $d\mu_{\bar{\Sigma}}$ and let $\alpha : \bar{\Sigma} \to \mathfrak{g}$ be a smooth, compactly supported *adjoint* test field. Write ∇_μ for the rig–compatible covariant derivative built from the Operator–Bundle Connection $\mathcal{A}_\mu$ (Definitions 6.192, 12.1 and 12.96), and $\nabla^{(\mathrm{ad})}_\mu$ for its adjoint action on $\mathfrak{g}$–valued fields. In the Tantrakyma sector (Definition 9.323), let $\mathcal{E}_\Theta$ denote the Euler–Lagrange operator and $\mathcal{V}_\Theta$ the scalar potential (*cf.* BEH ledger eq. (12.44)). Define the tagged meters

$$\mathsf{M}_{\mathrm{EL}}(\alpha) := \int_{\bar{\Sigma}} d\mu_{\bar{\Sigma}}\, \mathrm{Tr}\{\alpha(x)\, [\mathcal{E}_\Theta, \cdot]_{\mathrm{ad}}\}\,, \tag{12.38}$$

$$\mathsf{M}_{\mathrm{pot}}(\alpha) := \int_{\bar{\Sigma}} d\mu_{\bar{\Sigma}}\, \mathrm{Tr}\{\alpha(x)\, s_G\, \mathcal{V}_\Theta\}\,, \tag{12.39}$$

$$\mathscr{W}[\alpha] := \int_{\bar{\Sigma}} d\mu_{\bar{\Sigma}}\, \alpha^a(x) \left(\nabla^{(\mathrm{ad})\mu} J_\mu\right)_a, \tag{12.40}$$

where s_G is the infinitesimal G–variation implemented via ∇_μ and $\mathcal{A}_\mu$, and J_μ is the Noether current of the G–action on the Tantrakyma sector. All three meters are constructed with the same diagnostic realized reference configuration instrumentation and projections used throughout (e.g., TT–type projectors in radiative backgrounds; Definition 10.3 and eqs. (10.4) and (10.134)) and are designed to be BRST–closed and gauge–parameter independent upon descent to the physical sector $\mathcal{H}_{\rm phys} = H^0(\hat{Q}_{\rm BRST})$ (Section 12.3, Proposition 12.12, and Lemma 9.54). In the BEH phase, consistency is checked against the ledger conditions eqs. (12.41) to (12.44) (and, when present, CK–type masses eq. (12.45)), with poles/residues taken as Nielsen–flat on the plateau.

Lemma 12.17 (Descent and hygiene of the C-meters)**.** *The band–tagged versions of eqs.* (12.38) *to* (12.40) *commute with the physical projector and with* ∇_μ *on the certified plateau* $\mathfrak{P}$*; therefore they define well–posed operators on* $\mathcal{H}_{\rm phys}$ *whose expectations are Nielsen–flat.*

Proof. Diagnostic realized reference configuration/BRST setup. Work on a certified plateau so that readouts descend to the BRST physical space $\mathcal{H}_{\rm phys} = H^0(\hat{Q}_{\rm BRST})$ and are gauge/Nielsen–stable (Sections 6.12.4 and 12.3 and Proposition 12.12). Let $P_{\rm phys} : \mathcal{H} \to \mathcal{H}_{\rm phys}$ denote the canonical cohomological projector associated with $\hat{Q}_{\rm BRST}$ (ghost number 0 realization; Section 9.5 and Lemma 9.54). Band tags are implemented by the typed diagnostic realized reference configuration projectors used throughout (UV/IR band instruments and TT–type selections); in radiative backgrounds these include the York/TT projectors (Definition 10.3 and eqs. (10.4) and (10.134)). The covariant derivative ∇_μ is the rig–compatible connection built from the Operator–Bundle Connection (Definitions 6.192 and 12.1).

(i) BRST closure and descent. Each integrand in eqs. (12.38) to (12.40) has ghost number 0 and is BRST–closed on $\mathfrak{P}$: the Euler–Lagrange piece transforms into a total $\hat{Q}_{\rm BRST}$–exact term under the gauge variation; the potential variation $s_G \mathcal{V}_\Theta$ is BRST–closed at the realized vacuum; the Ward density $\nabla^{(\mathrm{ad})\mu} J_\mu$ generates the local symmetry and is BRST–closed on the diagnostic realized reference configuration (Noether/BRST compatibility) (Section 12.3 and Lemma 9.54). Hence each meter defines an endomorphism of $H^0(\hat{Q}_{\rm BRST})$, and on representatives one has

$$P_{\rm phys}\,\mathsf{M} \;=\; \mathsf{M}\,P_{\rm phys} \qquad \text{for} \;\; \mathsf{M} \in \{\mathsf{M}_{\rm EL}, \mathsf{M}_{\rm pot}, \mathcal{W}[\cdot]\},$$

i.e. the meters *commute* with the physical projector modulo BRST–exact terms, which vanish on $\mathcal{H}_{\rm phys}$ (Section 9.5).

(ii) Covariant commutation with ∇_μ. Band tags and typed projectors are constructed to be *covariantly constant* on the diagnostic realized reference configuration: the rig–compatible connection preserves the inner products and the decomposition used by the tags, and the TT projectors are built from ∇_μ–invariant tensors (Definitions 10.3 and 12.1 and eq. (10.4)). Consequently $[\nabla_\mu, \Pi_{\rm tag}] = 0$ for any diagnostic realized reference configuration tag $\Pi_{\rm tag}$, and because the meters are built from adjoint–covariant densities and α has compact support, the band–tagged meters commute with ∇_μ on $\mathfrak{P}$:

$$[\nabla_\mu,\; \Pi_{\rm tag}\mathsf{M}\Pi_{\rm tag}] = 0, \qquad \mathsf{M} \in \{\mathsf{M}_{\rm EL}, \mathsf{M}_{\rm pot}, \mathcal{W}[\cdot]\}.$$

(iii) Well–posedness on $\mathcal{H}_{\rm phys}$. Compact support of α and band–tameness of the tags ensure the integrals in eqs. (12.38) to (12.40) define closable operators on the kinematic

domain, and by (i) they descend to bounded (or closable) operators on $\mathcal{H}_{\mathrm{phys}}$ that commute with P_{phys} (Sections 6.12.4 and 12.3).

(iv) Nielsen flatness. Along the R_ξ gauge–fixing family, the derivative of any physical expectation value with respect to ξ is a BRST–exact insertion (Nielsen identity); hence on $H^0(\hat{Q}_{\mathrm{BRST}})$ the expectations of the band–tagged meters are ξ–independent. Together with the Källén–Lehmann/ Ward hygiene on the plateau, this gives Nielsen–flatness (Proposition 12.12).

Combining (i)–(iv) proves the claim. □

BEH ledger in the Tantrakyma sector (gauge & scalars). When $\langle\hat{\mathbf{\Theta}}\rangle \neq 0$, tests are extended by the Brout–Englert–Higgs (BEH) ledger (Ch. 9):

$$\mathfrak{h} := \{X \in \mathfrak{g} \mid [X, \langle\hat{\mathbf{\Theta}}\rangle] = 0\}, \tag{12.41}$$

$$\mathcal{G} := \mathrm{span}\{\,[T_a, \langle\hat{\mathbf{\Theta}}\rangle]\,\}/\sim \;\cong\; \mathfrak{g}/\mathfrak{h}, \tag{12.42}$$

$$(M_V^2)_{ab} := \omega\Big(\,[T_a, \langle\hat{\mathbf{\Theta}}\rangle]^\dagger\,[T_b, \langle\hat{\mathbf{\Theta}}\rangle]\,\Big) \qquad \text{(up to the standard couplings in Section 9.18.7)}, \tag{12.43}$$

$$M_H^2 := \left.\frac{\partial^2 \mathcal{V}_\Theta}{\partial\Theta^\dagger \partial\Theta}\right|_{\Theta=\langle\hat{\mathbf{\Theta}}\rangle}. \tag{12.44}$$

Goldstone directions are BRST-exact and thus do not contribute to $\mathcal{H}_{\mathrm{phys}}$; vector/scalar poles and residues read from the two-point functions are Nielsen-flat (Section 12.3 and Proposition 12.12). The $U(1)$ worked example provides calibration (Section 9.18.8).

Vector mass matrices for mition/tantron and how to read them. On a certified plateau with a rig–compatible connection (Section 6.12.4 and Definition 12.1), the internal gauge transport is carried by the Operator–Bundle Connection $\mathcal{A}_\mu$ (Definitions 6.192 and 12.96). In each sector, BEH breaking produces a vector mass–squared matrix from the action of the Lie generators on the relevant vacuum. For the *Gaiamiti* sector (whose spin–1 quanta are the *mition*; Definition 9.258), using the representation ρ_Φ that enters its covariant derivative (Section 6.5.3),

$$(M_\Phi^2)_{IJ} = g^2\, g_T\Big(\rho_\Phi(T_I)\,\bar{\Phi},\ \rho_\Phi(T_J)\,\bar{\Phi}\Big), \qquad O^\top M_\Phi^2 O = \mathrm{diag}(m_{V,a}^2), \quad V_{a\mu} = O^{\ I}{}_a A_\mu^I.$$

For the *Tantrakyma* sector (spin–1 *tantron*; Definition 9.348), using the representation entering its covariant derivative (Definition 9.323),

$$(M_\Theta^2)_{IJ} = g^2\, g_C\Big(\rho(T_I)\,\bar{\Theta},\ \rho(T_J)\,\bar{\Theta}\Big), \qquad O^\top M_\Theta^2 O = \mathrm{diag}(m_{V,a}^2), \quad W_{a\mu} = O^{\ I}{}_a A_\mu^I.$$

Chapter 12 repackages this in the commutator form eq. (12.43), making *no commutator, no mass* manifest; directions commuting with the VEV remain massless, those not commuting acquire mass (Goldstone span eq. (12.42)). The scalar (radial) masses are governed independently by eq. (12.44) and are not entries of M_V^2. In a broken eigenchannel (after diagonalizing

M^2), the gauge–fixed quadratic form assumes the standard R_ξ block,

$$\mathbb{Q}(k) = \begin{pmatrix} k^2 - m_V^2 & 0 & 0 & 0 \\ 0 & \dfrac{k^2 - m_V^2}{\xi} & 0 & 0 \\ 0 & 0 & k^2 - \xi m_V^2 & 0 \\ 0 & 0 & 0 & k^2 - M_{\mathrm{A}}^2 \end{pmatrix},$$

for $(A_T, A_L, \pi_{\mathrm{NG}}, h_{\mathrm{A}})$, so the transverse pole at $k^2 = m_V^2$ is the physical mition/tantron pole; NG/longitudinal pieces are BRST artefacts, and h_{A} carries M_{A}^2 (Section 12.3 and Proposition 12.12).

Bridge to sectoral mass matrices (mition/tantron). Using eq. (12.43) together with the sectoral covariant–derivative data from Chapters 6–9, the eigenfields $V_{a\mu}$ and $W_{a\mu}$ above are the mass eigenvectors (linear combinations of the original A_μ^I) that define the *mition/tantron* species (Definitions 9.258 and 9.348). Holosymmetric meters and the dual–register asymmetric interferometer provide gauge–stable identification of the transverse simple poles and informationally complete discrimination between eigenchannels (Definitions 2.114, 2.124 and 6.275, Section 3.1.5, Lemma 12.39, and eqs. (12.56) and (12.57)); on compact K, Peter–Weyl/Schur orthogonality separates species by representation content when required (Propositions 12.8 and 12.10 and Remark 12.9). In radiative geometries, TT projectors from Chapter 11 remove gravitational contamination during gauge–sector readout (Definition 10.3 and eqs. (10.4) and (10.134)).

CK ledger (non–TT sector; Ch. 11). If the Chrono-Kymon (CK) couples to $\hat{\mathbf{\Theta}}$, its linear-response kernel yields an induced mass

$$M_{\mathrm{CK}}^2 \;:=\; \mathcal{M}\big[\langle\hat{\mathbf{\Theta}}\rangle;\, \nabla_\mu, \mathcal{A}_\mu\big], \tag{12.45}$$

computed on the same diagnostic realized reference configuration and with the same connection (cf. Sections 11.8, 11.8.1 and 11.8.2). Typed microcausality, $\bar{\Sigma}$-hyperbolicity, KL positivity and Nielsen-flatness are certified as in Propositions 12.3 and 12.12; TT-purity of the gravity carrier remains intact (Ch. 6 hygiene).

Proposition 12.18 (Equivalence of invariance and vanishing C-meters)**.** *On the certified plateau, G–invariance of the Tantrakyma dynamics (definition immediately preceding) is equivalent to the vanishing of all tagged meters M_{EL}, $\mathsf{M}_{\mathrm{pot}}$, $\mathcal{W}[\alpha]$ on $\mathcal{H}_{\mathrm{phys}}$, together with the BEH/CK ledger conditions eqs. (12.41) to (12.45) evaluated with Nielsen–flat poles/residues.*

Proof. Diagnostic realized reference configuration and hygiene. Work on a certified plateau so that gauge/Nielsen choices are harmless for physical readouts and all meters descend cohomologically to the BRST sector (Sections 6.12.4 and 12.3 and Proposition 12.12). Write $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ with $\hat{Q}_{\mathrm{BRST}}^2 = 0$ and $\frac{d}{dt}\hat{Q}_{\mathrm{BRST}} = 0$ (Section 12.3).

$\Rightarrow$ Invariance $\Longrightarrow$ meters vanish. Assume G–invariance of the Tantrakyma dynamics on the window (definition preceding the proposition): the action and equations of motion are covariant under local $g(x) \in G$ built from the compact holonomy subgroup K (Proposition 12.8). For

any compactly supported parameter $\alpha(x)$, the associated Ward functional $\mathcal{W}[\alpha]$ is the BRST–closed generator of the local variation; G–invariance implies the Noether identity and therefore $\mathcal{W}[\alpha] = 0$ as an operator on $\mathcal{H}_{\text{phys}}$ (Section 12.3 and Proposition 12.12). The Euler–Lagrange meter M_{EL} (kinetic/gauge sector) is proportional to the equations of motion contracted with the symmetry variation; by covariance and on-shell BRST descent it vanishes on H^0 (Section 12.3, cf. Lemma 12.17). Likewise, M_{pot} tests invariance of the potential around the realized vacuum; G–invariance together with the BEH/CK ledger conditions eqs. (12.41) to (12.45) ("no commutator, no mass" and the correct scalar Hessian) imply that Goldstone directions are BRST–exact and that vector/scalar poles and residues are Nielsen–flat (Section 12.3 and Proposition 12.12). Hence $\mathsf{M}_{\text{pot}} = 0$ on $\mathcal{H}_{\text{phys}}$. Collecting, all tagged meters vanish on the plateau.

$\Leftarrow$ Meters vanish $\Longrightarrow$ invariance. Assume $\mathsf{M}_{\text{EL}} = \mathsf{M}_{\text{pot}} = 0$ and $\mathcal{W}[\alpha] = 0$ on $\mathcal{H}_{\text{phys}}$, with the BEH/CK ledger satisfied and poles/residues Nielsen–flat (eqs. (12.41) to (12.45) and Proposition 12.12). First, $\mathcal{W}[\alpha] = 0$ provides the Ward identity for any local parameter $\alpha(x)$, which reconstructs the adjoint–covariant local symmetry on the kinetic/gauge sector (Noether direction), while $\mathsf{M}_{\text{EL}} = 0$ enforces the Euler–Lagrange part of the invariance (kinetic/gauge terms) after BRST descent (Section 12.3, cf. Lemma 12.17). Second, $\mathsf{M}_{\text{pot}} = 0$ fixes the interaction (potential) part of the local variation to vanish at the realized VEV; the ledger conditions ensure that any apparent gauge–fixed noninvariance is confined to BRST–exact Goldstone/longitudinal pieces and does not act on $\mathcal{H}_{\text{phys}}$ (eq. (12.42), Section 12.3, and Proposition 12.12). Since the compact support window is generated by K (Proposition 12.8), these local statements integrate to full G–invariance of the Tantrakyma dynamics on the plateau.

Conclusion. Under plateau hygiene, G–invariance is equivalent to the vanishing of all tagged meters together with the BEH/CK ledger satisfied at Nielsen–flat poles/residues, establishing the claim. □

Decision layer (tetralemma effects for symmetry). Aggregate normalized residuals of $\mathsf{M}_{\text{EL}}, \mathsf{M}_{\text{pot}}, \mathcal{W}$ (and, when applicable, BEH/CK residuals) into a bounded self-adjoint *symmetry score* $\mathsf{S}_{\text{sym}} \in [-1, 1]$ and a *violation witness* $\mathsf{D}_{\text{sym}} \geq 0$ (band-averaged, BRST-closed). Fix thresholds $\alpha_{\text{sym}}, \beta_{\text{sym}} \in (0, 1)$ from the holosymmetric calibration (Ch. 8), and define effects exactly as in the tetralemma scheme (Definition 12.13):

$$E^{[\text{sym}]}_{\mathsf{Th}} = \mathbf{1}_{[\alpha_{\text{sym}},1]}(\mathsf{S}_{\text{sym}}), \quad E^{[\text{sym}]}_{\mathsf{AntiTh}} = \mathbf{1}_{(\beta_{\text{sym}},\infty)}(\mathsf{D}_{\text{sym}}),$$

$$E^{[\text{sym}]}_{\mathsf{SynTh}} = \mathbf{1}_{(-\infty,\alpha_{\text{sym}})}(\mathsf{S}_{\text{sym}}) \wedge \mathbf{1}_{(\beta_-,\beta_+]}(\mathsf{D}_{\text{sym}}), \quad E^{[\text{sym}]}_{\mathsf{HoloTh}} = \mathbf{1}_{(-\infty,\alpha_{\text{sym}})}(\mathsf{S}_{\text{sym}}) \wedge \mathbf{1}_{[0,\beta_{\text{sym}}]}(\mathsf{D}_{\text{sym}}),$$

with $0 < \beta_- < \beta_{\text{sym}} < \beta_+$. Born weights $v_M = \text{Tr}(E^{[\text{sym}]}_M \rho)$ and a visibility $\Lambda^{[\text{sym}]} = [\, v_{\mathsf{Th}} - (v_{\mathsf{AntiTh}} + v_{\mathsf{SynTh}}) \,]_+$ are reported alongside $(\mathcal{W}, \mathcal{Z})$ (Ch. 8).

SM specialization (calibration fixture). When Tantrakyma reps match the SM matter content (Constraints D/E), the connection decomposes canonically

$$\mathcal{A}_\mu \;=\; g_s\, G^a_\mu T_a \;+\; g\, W^i_\mu \tau_i \;+\; g'\, B_\mu\, Y \qquad \text{(Definitions 12.95 and 12.96)}.$$

Then C-meters vanish for the Yang–Mills+Higgs form; the BEH ledger reproduces $W^\pm, Z^0$ masses and a massless photon (unbroken $Q = I_3 + \frac{1}{2}Y$), with Nielsen-flat poles/residues. This block provides a running calibration for S_{sym} and D_{sym}.

Boundary variant (optional). Repeat all constructions tangentially on a boundary plateau $\mathfrak{P}_\partial$ with a boundary-compatible connection (Ch. 6; Definitions 12.4 and 12.15): use $\mathcal{A}_\parallel$, $\mathcal{F}_\parallel$, boundary UV/IR instruments, and the boundary Ward identity $\nabla_\parallel \cdot J_\parallel^{(\partial)} = S_\partial$. This defines the boundary fibre $\mathcal{H}^\partial_{\text{int}}$, a compact boundary holonomy group, and a boundary symmetry visibility $\Lambda_\partial^{[\text{sym}]}$, inheriting descent, Nielsen-flatness, and typed microcausality as in the bulk (Chs. 6–8; Lemma 12.2 and Proposition 12.3).

Constraint C certifies that the *same* operator-bundle connection $\mathcal{A}_\mu$ and split covariant derivative ∇_μ that passed Constraints A/B render the Tantrakyma action S_Θ symmetric at the level of EL equations, potentials/Yukawas, and Ward identities, and that any BEH organization (including CK) is BRST/ward-compatible with Nielsen-flat spectra. The decision is reported by tetralemma effects fed by the symmetry score/witness, on the certified plateau. The equation

$$\mathcal{A}_\mu \;=\; g_s\, G^a_\mu T_a \;+\; g\, W^i_\mu \tau_i \;+\; g'\, B_\mu\, Y$$

(Definitions 12.95 and 12.96) is the *gauge-connection decomposition* in a basis of the SM Lie algebra acting on Tantrakyma fibres. It belongs to the gauge sector and enters the split covariant derivative

$$\nabla_\mu \;=\; \nabla^{(\text{LC})}_\mu \;+\; i\,\text{ad}_{\mathcal{A}_\mu}$$

(Definitions 6.192 and 12.1), the very ∇_μ required by Constraint C. This decomposition specifies how the gluonic, weak-isospin, and hypercharge carriers couple to any field transforming under G, independently of which Tantrakyma quanta are present.

What renders this decomposition *operational* in the electroweak sector is the Higgs-type kymon $H \sim (1,2)_{+1/2}$. Its covariant derivative uses precisely the weak and hypercharge pieces,

$$\nabla_\mu H \;=\; \partial_\mu H \;+\; i\, g\, W^i_\mu \tau_i\, H \;+\; i\, g'\, B_\mu\, Y\, H,$$

and a nonzero $\langle H\rangle$ drives the BEH mechanism (Ch. 9), yielding massive $W^\pm, Z^0$ and the unbroken electromagnetic generator $Q = I_3 + \frac{1}{2}Y$ (massless photon). The corresponding pole locations and residues are certified Nielsen-flat on the diagnostic realized reference configuration (Sections 9.18.7 and 9.18.8 and Proposition 12.12).

Other Tantrakyma-related quanta do not alter the decomposition itself. The Chronokymon (CK) in the τ–Kairos sector can acquire an optional BEH mass from $\langle\hat{\boldsymbol{\Theta}}\rangle$ while preserving TT-purity and typed microcausality; its ledger is checked within the same Constraint C hygiene (Ch. 11; Sections 11.8, 11.8.1 and 11.8.2 and Proposition 12.3). Tantrons (spinor excitations) determine which generators T_a, τ_i, Y act and with which charges—facts enforced by Constraints D/E (chiral admissibility and exact charge/Yukawa admissibility), not by changing $\mathcal{A}_\mu$'s form (Section 12.1.2).

Holosymmetric filtration

Having specified the admissible local, compact, Tantrakyma-symmetric subgroups, the final winnowing is dynamical. Stage I introduces the Unitarity Operator $\mathbf{U}^{(A)}_{\text{unitarity}} = (\mathbf{U}^{(A)})^\dagger \mathbf{U}^{(A)}$ and the Self-Coherence Functional $\mathcal{C}(A) = |\langle\Psi_0|\mathbf{U}^{(A)}_{\text{unitarity}}|\Psi_0\rangle|$ which must saturate unity for viability (Definition 12.28; Postulate Definition 12.33; cf. the holosymmetric selection pipeline in Sections 2.4 and 12.2). Stage II compares any two viable candidates via the

asymmetric interferometer and its Relational Coherence Functional, enforcing suppression of cross-visibilities with the realized sector (eq. (12.49); cf. the phase-resolved causal response and its T-even/T-odd separation in Definition 5.139 and Proposition 5.60, with cone assignment phase-invariant by Corollary 5.126 and no–law–signalling guaranteed by Proposition 5.152 and Corollary 5.145). The entire filtration acts on the *physical* (BRST) sector (Section 6.10, Principle 12.1, and Lemma 6.210) and can be read through the typed/ledger insertions when needed (Definition 10.80), while respecting the holosymmetric cone kinematics of Chapter 5 (Definition 5.123 and Proposition 5.153).

Interlocks with the Foundations

The same $\mathcal{A}_\mu$ controls parallel transport in the Unified Inter-Modal Correlator, with graded covariant constancy $\mathfrak{D}_\mu \hat{\mathbf{G}}_\Lambda = 0$ encoding consistency of preparation/transport/readout; its curvature sources holonomy-induced deviations (Definition 6.4, Section 4.1.7, and eq. (10.61)). This makes the geometric sector's Einstein–type equations and the matter sector's gauge dynamics two faces of a single operator-transport structure (cf. the gaiamiti/tantrakyma doublet Definition 6.1, the relational connection Definition 6.7, and the chronogeometric tensor entering transport Definition 5.26), and it is precisely this shared structure that allows the holosymmetric interferometer to compare candidate gauge evolutions on equal operational footing (cf. Section 2.4 and the causal/operational layer in Definition 5.139 and Proposition 5.143).

Finally, the geometric correlator $\hat{G}_{\Phi\Phi}$ that underpins chronogeometric dynamics is one block of the same unified correlator matrix used above, so gravitational response and internal gauge selection are computed within a single correlational calculus (Section 10.17 and Definition 6.4). In particular, the Einstein-type tensor $\bar{G}_{\mu\nu}[\bar{\Sigma}]$ employed for background transport is introduced within this correlator framework, ensuring that "geometry" is never invoked outside the operational algebra (cf. the effective cone definition and phase invariance in Definition 5.122 and Corollary 5.126 and the PQS locus where these structures attach Theorem 5.155).

12.1.2 Matching to the Observed Fermion Content

The second stage requires that any admissible gauge sector reproduce the empirically observed *fermionic representation content*—here realized as Tantrakyma-sector spinor sections arranged into quark and lepton multiplets (Section 12.4.1). Concretely, the representation theory must encode both the electroweak chirality pattern and the full set of color, isospin, and hypercharge assignments in table 12.1 [82, 91]. Throughout, fermions live on the same operator bundle and split covariant derivative $\nabla_\mu = \nabla_\mu^{(\mathrm{LC})} + i\,\mathrm{ad}_{\mathcal{A}_\mu}$ used in Constraints A–C (Ch. 6; Definitions 6.192 and 12.1; see also [435, 375]). All checks are executed bandwise on the certified plateau $\mathfrak{P}$, where tags commute with ∇_μ and with the physical projector, so BRST descent and the audit guarantees apply (Lemmas 12.2 and 12.7; cf. [445, 106, 442]). On each finite internal fibre, compact holonomy and Peter–Weyl yield unitary, finite-multiplicity irreducibles that scaffold the spinor content (Propositions 12.8 and 12.10; [552, 248]). Ward/KL hygiene is ensured by the improved currents and positivity results (Definition 12.11 and Proposition 12.12; see [484, 483, 82, 91]).

Dimensional note. The use of $P_{L/R} = \frac{1}{2}(1 \mp \gamma^5)$ and chiral identities is specific to d=4 (and to the $d \to 4$ interpretation in DimReg).

To *weld* Constraints D–E to this diagnostic realized reference configuration, first impose **Chiral admissibility (D)**. Let $P_{L/R} = \frac{1}{2}(1 \mp \gamma^5)$ act on the Tantrakyma spinor bundle, and write $\Pi_L^{(2)}$ for the fibrewise projector onto $SU(2)_L$ doublets and $\Pi_R^{(1)}$ for singlets extracted from the same $SU(2)_L$-representation. On the band one requires the operator identities

$$P_L\,\Psi = \Pi_L^{(2)}\,P_L\,\Psi, \qquad P_R\,\Psi = \Pi_R^{(1)}\,P_R\,\Psi,$$

together with the placement of each generation into the slots listed in table 12.1 (i.e. $Q_L \sim (3,2)_{1/6}$, $L_L \sim (1,2)_{-1/2}$, $u_R \sim (3,1)_{2/3}$, $d_R \sim (3,1)_{-1/3}$, $e_R \sim (1,1)_{-1}$, $\nu_R \sim (1,1)_0$ optional). Because tags are BRST-closed and ∇_μ is a chain map, these identities descend to $\mathcal{H}_{\text{phys}}$ and are gauge-parameter independent (Nielsen-flat) (Lemma 12.2 and Proposition 12.12; see also [120]). The empirical chiral pattern itself is the one realized by the electroweak theory [57, 58, 59]; standard accounts include [82, 124, 554].

Next enforce **Exact charge realization and Yukawa admissibility (E)**. Promote Y, I_3, and Q to commuting observables on the same fibre and check, via spectral projectors, that the observed eigenvalue assignments hold: $\mathbf{1}_{\{y_f\}}(Y)\Psi_f = \Psi_f$ with y_f as in table 12.1. The Gell–Mann–Nishijima relation becomes the operator constraint on the band,

$$Q - I_3 - \tfrac{1}{2}\,Y \;=\; 0$$

(see Section 12.4.3 and [125, 126]). Gauge-invariant, renormalizable Yukawa couplings with $H \sim (1,2)_{+1/2}$ are verified intrinsically by the meter $s_G\mathcal{L}_Y = 0$ constructed with the same ∇_μ and $\mathcal{A}_\mu$ as in Constraint C (Chs. 9–10; Sections 9.18.7 and 9.18.8 and Definition 6.1; see [58, 59, 97]). On the certified plateau, the improved-current Ward identities ensure gauge covariance of $\mathcal{L}_Y$ and KL positivity of fermion two-point functions (Definition 12.11 and Proposition 12.12; [484, 483, 82, 91]), and BRST-exactness protects the construction [445, 106, 442]. BEH then ties these assignments to mass generation through H without spoiling BRST descent or microcausality. Operationally, package these tests into band-tagged, BRST-closed meters—(i) a chiral overlap meter for $P_{L/R}$ vs. $\Pi_L^{(2)}, \Pi_R^{(1)}$; (ii) a charge-spectrum meter for Y, I_3, Q and $Q = I_3 + \frac{1}{2}Y$; (iii) a Yukawa gauge-variation meter $s_G\mathcal{L}_Y$—and read them via the tetralemma effects (Definition 12.13) gated by the holosymmetric overlay (Ch. 6). Thesis/Antithesis/Synthesis/Holothesis outcomes thus align D–E with the operator-bundle, BRST/Ward, KL, compactness/holonomy, and decision-layer machinery established in Chs. 6–11. For neutrinos, the optional singlet ν_R naturally interfaces with seesaw realizations within the same representation framework [129, 547].

Constraint D (Chiral admissibility).

The weak interaction is maximally parity-violating: only left-chiral fields transform nontrivially under the weak isospin factor. Thus the gauge sector must admit representations for which

$$P_L\Psi \sim \text{doublets of } SU(2)_L, \qquad P_R\Psi \sim \text{singlets of } SU(2)_L,$$

with $P_{L/R} = \frac{1}{2}(1 \mp \gamma^5)$ acting on the Tantrakyma spinors. In particular, one generation must realize

$$\begin{aligned} Q_L &\sim (3,2)_{1/6}, & L_L &\sim (1,2)_{-1/2}, \\ u_R &\sim (3,1)_{2/3}, & d_R &\sim (3,1)_{-1/3}, \\ e_R &\sim (1,1)_{-1}, & \nu_R &\sim (1,1)_0 \text{ (optional)}. \end{aligned}$$

This chiral split must be implementable within the Tantrakyma grading so that left/right project to inequivalent G–representations on the same modal fibre. On the band, write $\Pi^{(2)})_L$ for the fibrewise projector onto $SU(2)_L$ doublets and $\Pi_R^{(1)}$ for singlets; chiral admissibility is the operator identity

$$P_L\Psi = \Pi_L^{(2)} P_L \Psi, \qquad P_R\Psi = \Pi_R^{(1)} P_R \Psi,$$

together with the placement of each generation into the irrep slots of table 12.1. Because tags commute with ∇_μ and the physical projector on $\mathfrak{P}$, these identities descend to $\mathcal{H}_{\text{phys}}$ and are gauge-parameter independent (Nielsen-flat) (Lemma 12.2 and Proposition 12.12; see [120]). Standard accounts: [82, 124, 554]; for the historical electroweak structure, see [57, 58, 59].

Constraint E (Exact charge realization and Yukawa admissibility).

The gauge sector must carry a $U(1)$ generator Y whose eigenvalues match table 12.1 and satisfy the Gell–Mann–Nishijima relation

$$Q \;=\; I_3 + \tfrac{1}{2}Y \qquad \text{(Section 12.4.3; see [126])}.$$

Equivalently, there must exist a Higgs/Tantrakyma electroweak doublet $H \sim (1,2)_{+1/2}$ whose *gauge-invariant* renormalizable Yukawa couplings can be formed,

$$\mathcal{L}_Y \supset -\bar{Q}_L\, \tilde{H}\, Y_u\, u_R - \bar{Q}_L\, H\, Y_d\, d_R - \bar{L}_L\, H\, Y_e\, e_R \text{ (and } - \bar{L}_L\, \tilde{H}\, Y_\nu\, \nu_R \text{ if present)},$$

with $\tilde{H} = i\sigma^2 H^*$. On the band, promote Y, I_3, Q to commuting observables and enforce the operator constraint $Q - I_3 - \frac{1}{2}Y = 0$ together with the spectral placements $\mathbf{1}_{\{y_f\}}(Y)\Psi_f = \Psi_f$ for each fermion f listed in table 12.1. Gauge invariance of $\mathcal{L}_Y$ is verified intrinsically via the meter $s_G\mathcal{L}_Y = 0$ built with the same ∇_μ and $\mathcal{A}_\mu$ used in Constraint C (Chs. 9–10; Sections 9.18.7 and 9.18.8 and Definition 6.1). This *Yukawa admissibility* criterion ties the hypercharge assignments to electroweak symmetry breaking and mass generation, and it is satisfied with striking economy by $G_{\text{SM}} = SU(3)_C \times SU(2)_L \times U(1)_Y$ (table 12.1). Standard expositions: [82, 554, 128, 97].

Operational meters and decision layer for D/E.

Implement band-tagged, BRST-closed meters parallel to those used in Constraints B/C: (i) a *chiral meter* (overlap of $P_{L/R}$ with $\Pi_L^{(2)}, \Pi_R^{(1)}$); (ii) a *charge-spectrum meter* (spectral mismatch of Y, I_3, Q against table 12.1 and the operator identity $Q = I_3 + \frac{1}{2}Y$); (iii) a *Yukawa gauge-invariance meter* ($s_G\mathcal{L}_Y$). By Lemma 12.2, each meter descends to $\mathcal{H}_{\text{phys}}$ and is Nielsen-flat; by Propositions 12.8 and 12.10 the fibrewise representation content is finite and unitary on $\mathfrak{P}$. Readouts use the tetralemma effects (Definition 12.13) gated by the holosymmetric overlay (Ch. 6): *Thesis* when all meters pass within tolerance, *Antithesis* upon decisive

violation, *Synthesis* for typed/bandwise mixed outcomes, and *Holothesis* whenever law/state visibilities fall below plateau thresholds. This welds Constraints D–E to the operator-bundle, BRST/Ward, KL-positivity, compactness/holonomy, and decision-layer machinery established in Chs. 6–11.

12.1.3 The Anomaly Cancellation Constraint

Among all filters, anomaly cancellation is the most stringent. In a chiral gauge theory, quantum effects can violate classical gauge invariance by inducing non-vanishing gauge variations of the effective action; such *anomalies* render the theory inconsistent (loss of unitarity or locality) and must vanish identically [443, 91, 82, 103]. In the IQR/holosymmetric setting this requirement coincides with Stage I of the HSP: only theories with saturated self-coherence, $\mathcal{C}(A) = 1$, survive; an uncancelled anomaly obstructs BRST nilpotency and thus fails the viability condition [442, 108] (Sections 12.2 and 12.3).

- **Constraint F (Complete anomaly freedom).** For one generation of left-chiral Weyl fields in representations $\mathbf{R}_3 \otimes \mathbf{R}_2$ with hypercharge Y, the consistent local (perturbative) anomalies must vanish:

$$\begin{aligned}
\sum_{\text{LH}} d_2(\mathbf{R}_2)\, A_3(\mathbf{R}_3) &= 0, \\
\sum_{\text{LH}} d_3(\mathbf{R}_3)\, A_2(\mathbf{R}_2) &= 0, \\
\sum_{\text{LH}} Y\, d_2(\mathbf{R}_2)\, T_3(\mathbf{R}_3) &= 0, \\
\sum_{\text{LH}} Y\, d_3(\mathbf{R}_3)\, T_2(\mathbf{R}_2) &= 0, \\
\sum_{\text{LH}} d_3(\mathbf{R}_3)\, d_2(\mathbf{R}_2)\, Y^3 &= 0, \\
\sum_{\text{LH}} d_3(\mathbf{R}_3)\, d_2(\mathbf{R}_2)\, Y &= 0.
\end{aligned} \tag{12.46}$$

  (The $SU(2)^3_L$ cubic anomaly vanishes automatically since the fundamental is pseudoreal.) Here $d_{2,3}$ denote representation dimensions, $T_{2,3}$ the Dynkin indices (e.g. $T(\mathbf{2}) = T(\mathbf{3}) = \frac{1}{2}$), and $A_{2,3}$ the cubic indices (odd under conjugation); see [82, 91, 443] for conventions and derivations. Global anomalies must also be absent; in particular, the $SU(2)$ Witten anomaly requires an even number of left-chiral doublets per generation (satisfied in the SM: 3 quark doublets + 1 lepton doublet = 4) [553].

Per-generation check for the Standard Model. Writing all fermions as *left-chiral* fields (so that right-chiral fermions appear as left-chiral conjugates with conjugate representations and

opposite Y), the assignments in table 12.1 satisfy (12.46) identically (see, e.g., [82, 91, 443]):

$$\begin{aligned}
2\,Y(Q_L)\,T(\mathbf{3}) + Y(u_R^c)\,T(\bar{\mathbf{3}}) + Y(d_R^c)\,T(\bar{\mathbf{3}}) &= 0, && SU(3)_C^2\text{–}U(1)_Y\\
3\,Y(Q_L)\,T(\mathbf{2}) + Y(L_L)\,T(\mathbf{2}) &= 0, && SU(2)_L^2\text{–}U(1)_Y\\
3\Big(2\big(\tfrac{1}{6}\big)^3 + \big(-\tfrac{2}{3}\big)^3 + \big(\tfrac{1}{3}\big)^3\Big) + \Big(2\big(-\tfrac{1}{2}\big)^3 + 1^3 + 0^3\Big) &= 0, && U(1)_Y^3\\
3\Big(2\cdot\tfrac{1}{6} - \tfrac{2}{3} + \tfrac{1}{3}\Big) + \Big(2\cdot\big(-\tfrac{1}{2}\big) + 1 + 0\Big) &= 0, && \mathrm{grav}^2\text{–}U(1)_Y\\
2\,A(\mathbf{3}) + A(\bar{\mathbf{3}}) + A(\bar{\mathbf{3}}) &= 0, && SU(3)_C^3\\
0 &= 0. && SU(2)_L^3
\end{aligned}$$

The $SU(2)$ global anomaly cancels because each generation contains 4 doublets (an even number) [553].

Minimality and uniqueness. Subject to the empirical fermion content in table 12.1, absence of exotics (no additional chiral states beyond vectorlike pairs), and a single electroweak doublet mediator of Yukawa couplings, the product group

$$G_{\mathrm{SM}} \;=\; SU(3)_C \times SU(2)_L \times U(1)_Y$$

is the *unique minimal chiral gauge sector* solving (12.46). Any alteration of hypercharges or removal of a single multiplet upsets at least one of the conditions, reintroducing anomalies and violating Stage I of the HSP. This delicacy is the precise, non-miraculous content behind the oft-remarked "miracle" of SM anomaly cancellation [443, 91, 82].

Operational formalization of Constraint F

We now recast Section 12.1.3 as a bandwise, BRST-safe test on the certified plateau, using the same operator-bundle connection $\nabla_\mu = \nabla_\mu^{(\mathrm{LC})} + i\,\mathrm{ad}_{\mathcal{A}_\mu}$ and tags as in Constraints A–E (Ch. 6; Definitions 6.192 and 12.1). All meters below are constructed on the certified plateau $\mathfrak{P}$, where tags commute with ∇_μ and with the physical projector, so that BRST descent and the audit guarantees apply (Lemmas 12.2 and 12.7). On each finite internal fibre $\mathcal{H}_{\mathrm{int}}(x;\lambda_{\mathrm{UV}},\ell)$, compact holonomy and Peter–Weyl yield a finite set of unitary irreducibles for the fermion sector (Propositions 12.8 and 12.10); normalization of currents uses the GNS inner product (Ch. 6; Definition 12.11), and KL/Ward hygiene follows from Proposition 12.12.

Definition 12.19 (Bandwise extraction of chiral representation content)**.** Let $\{\,\Psi_f\,\}$ denote the family of Tantrakyma-sector *left-chiral* Weyl fields selected by the chiral projector $P_L = \frac{1}{2}(1-\gamma^5)$ on the band $K_{\mathrm{IR}}^{(\ell)} P_x(\lambda_{\mathrm{UV}})$ (Ch. 6–7). For each f let $(\mathbf{R}_3^{(f)}, \mathbf{R}_2^{(f)}; Y_f)$ be the $SU(3)_C \otimes SU(2)_L$ representation and hypercharge eigenvalue determined by the fibrewise Peter–Weyl decomposition and the spectral measure of Y (Proposition 12.10 and Section 12.1.2). Write $d_{2,3}$ for representation dimensions, $T_{2,3}$ for Dynkin indices, and $A_{2,3}$ for cubic indices (odd under conjugation), all fixed by current normalization through Definition 12.11.

Definition 12.20 (Local anomaly meters (band-tagged, BRST-closed))**.** Define six scalar (number-valued) meters on the band which reproduce the anomaly sums in (12.46):

$$\mathsf{A}_{3^3} := \sum_{f\in\mathrm{LH}} d_2(\mathbf{R}_2^{(f)})\, A_3(\mathbf{R}_3^{(f)}), \qquad \mathsf{A}_{2^3} := \sum_{f\in\mathrm{LH}} d_3(\mathbf{R}_3^{(f)})\, A_2(\mathbf{R}_2^{(f)}),$$

$$\mathsf{A}_{3^2Y} := \sum_{f\in\mathrm{LH}} Y_f\, d_2(\mathbf{R}_2^{(f)})\, T_3(\mathbf{R}_3^{(f)}), \qquad \mathsf{A}_{2^2Y} := \sum_{f\in\mathrm{LH}} Y_f\, d_3(\mathbf{R}_3^{(f)})\, T_2(\mathbf{R}_2^{(f)}),$$

$$\mathsf{A}_{Y^3} := \sum_{f\in\mathrm{LH}} d_3(\mathbf{R}_3^{(f)})\, d_2(\mathbf{R}_2^{(f)})\, Y_f^3, \qquad \mathsf{A}_{\mathrm{grav}^2Y} := \sum_{f\in\mathrm{LH}} d_3(\mathbf{R}_3^{(f)})\, d_2(\mathbf{R}_2^{(f)})\, Y_f.$$

Each meter is computed from the fibrewise representation list of Definition 12.19 and is *tag-stable* by construction; by Lemma 12.2 they are BRST-closed and Nielsen-flat on $\mathfrak{P}$. (The $SU(2)_L^3$ cubic anomaly vanishes identically because **2** is pseudoreal.)

Proposition 12.21 (Vanishing of local meters $\Leftrightarrow$ BRST nilpotency on the diagnostic realized reference configuration)**.** *On the certified plateau $\mathfrak{P}$, the simultaneous vanishing $\mathsf{A}_{3^3} = \mathsf{A}_{2^3} = \mathsf{A}_{3^2Y} = \mathsf{A}_{2^2Y} = \mathsf{A}_{Y^3} = \mathsf{A}_{\mathrm{grav}^2Y} = 0$ is equivalent to the absence of consistent local gauge anomalies for the band-extracted fermion content. Equivalently, the Wess–Zumino descent cocycle of the effective action is BRST-exact and vanishes upon projection to $\mathcal{H}_{\mathrm{phys}}$, so that $s^2 = 0$ on gauge-invariant observables (Stage I viability of the HSP; Sections 12.2 and 12.3).*

Proof. Setup on the certified plateau. Work on the certified plateau $\mathfrak{P}$ with the split connection $\nabla_\mu = \nabla_\mu^{(\mathrm{LC})} + i\,\mathrm{ad}_{\mathcal{A}_\mu}$ and BRST/realized weave configuration compatibility from Ch. 6 (Definitions 6.192 and 12.1). By tag commutation and band audit, all band-tagged constructions commute with the physical projector and descend to $\mathcal{H}_{\mathrm{phys}}$ (Lemmas 12.2 and 12.7). On each finite internal fibre $\mathcal{H}_{\mathrm{int}}(x;\lambda_{\mathrm{UV}},\ell)$, compact holonomy and Peter–Weyl provide a finite list of unitary irreps for the left-chiral fermions $\{\Psi_f\}$ with representation data $(\mathbf{R}_3^{(f)},\mathbf{R}_2^{(f)};Y_f)$ (Propositions 12.8 and 12.10 and Definition 12.19). Fix current normalizations by the GNS inner product (Definition 12.11); Ward/KL hygiene used below follows from Proposition 12.12.

( ) Vanishing local meters $\Rightarrow$ absence of consistent local anomalies and BRST nilpotency. Consider the consistent gauge variation of the renormalized effective action Γ under an infinitesimal local parameter $\alpha(x)$:

$$s\,\Gamma \;=\; \delta_\alpha\Gamma \;=\; \int d^4x\; \alpha^a(x)\,\mathscr{A}_a(x),$$

with anomaly densities $\mathscr{A}_a$ solving the Wess–Zumino consistency condition (the "descent equation"). For a product group $SU(3)_C \times SU(2)_L \times U(1)_Y$, locality and gauge covariance imply that $\mathscr{A}_a$ is a linear combination of six independent tensor structures labelled by $SU(3)^3$, $SU(2)^3$, $SU(3)^2U(1)$, $SU(2)^2U(1)$, $U(1)^3$, $\mathrm{grav}^2{-}U(1)$, with coefficients given by the left-chiral *representation sums* in (12.46). By construction, the band-tagged scalar meters of Definition 12.20 reproduce exactly these coefficients:

$$\mathsf{A}_{3^3},\ \mathsf{A}_{2^3},\ \mathsf{A}_{3^2Y},\ \mathsf{A}_{2^2Y},\ \mathsf{A}_{Y^3},\ \mathsf{A}_{\mathrm{grav}^2Y}.$$

(Here $\mathsf{A}_{2^3} \equiv 0$ for $SU(2)$ since the cubic index vanishes for all $SU(2)$ reps; this matches the remark following (12.46).) If all six meters vanish on $\mathfrak{P}$, then all local coefficients are zero,

hence $\mathscr{A}_a \equiv 0$ (consistent scheme). Equivalently, the anomaly cocycle from the Wess–Zumino descent is BRST-exact with a vanishing class on the band.

Because the meters are BRST-closed and tag-stable, this statement *descends* to the physical space (Lemma 12.2): $s\,\Gamma = 0$ on $\mathcal{H}_{\text{phys}}$. Therefore the BRST differential squares to zero on gauge-invariant observables (no obstruction at ghost number 1), i.e.

$$s^2 = 0 \quad \text{on} \quad H^0(\hat{Q}_{\text{BRST}}) = \mathcal{H}_{\text{phys}},$$

establishing Stage I viability of the HSP (Sections 12.2 and 12.3).

( ) BRST nilpotency and absence of consistent local anomalies ⇒ vanishing local meters. Conversely, assume $s\,\Gamma = 0$ on $\mathfrak{P}$ (no consistent local anomaly) and hence $s^2 = 0$ on $\mathcal{H}_{\text{phys}}$. By the Wess–Zumino consistency condition, any surviving $\mathscr{A}_a$ would define a nontrivial BRST cocycle at ghost number 1. But by hypothesis the corresponding cohomology class vanishes, so all coefficients of the six independent local structures must be zero. Since the scalar meters of Definition 12.20 are precisely these coefficients evaluated with the GNS-normalized indices (Definition 12.11) and extracted bandwise from the Peter–Weyl list (Definition 12.19), one concludes

$$\mathsf{A}_{3^3} = \mathsf{A}_{2^3} = \mathsf{A}_{3^2Y} = \mathsf{A}_{2^2Y} = \mathsf{A}_{Y^3} = \mathsf{A}_{\text{grav}^2Y} = 0.$$

Audit continuity and tag commutation ensure this implication is stable under $(\lambda_{\text{UV}}, \ell)$-variations and under projection to $\mathcal{H}_{\text{phys}}$ (Lemmas 12.2 and 12.7).

Combining the two directions yields the stated equivalence between the vanishing of the local anomaly meters and BRST nilpotency/absence of consistent local anomalies on the certified plateau, completing the proof. □

Definition 12.22 (Global anomaly meters)**.** Let $\Pi^{(2)}$ be the projector onto $SU(2)_L$ doublets on the band. Set the doublet-count operator $\mathsf{N}_2 := \sum_{f\in\text{LH}} \langle \Psi_f, \Pi^{(2)} \Psi_f \rangle$ (counted per generation). Define the Witten-parity effect $E_{\text{Witten}} := \frac{1}{2}(\mathbf{1} + (-1)^{\mathsf{N}_2})$, which equals $\mathbf{1}$ iff N_2 is even. In boundary settings, complement this by a tangential inflow check using the boundary Ward identity $\nabla_{\parallel} \cdot J_{\parallel}^{(\partial)} = S_{\partial}$ (Definitions 12.4 and 12.15) to ensure no residual global inflow is required.

Proposition 12.23 (Audit continuity and descent of anomaly diagnostics)**.** *All meters of Definitions 12.20 and 12.22 are continuous under the band audit and commute with the physical projector: their values vary continuously with* $(\lambda_{\text{UV}}, \ell)$*, and their effects descend to* $\mathcal{H}_{\text{phys}}$ *(Chs. 6–7; Definition 9.166 and Lemmas 12.2 and 12.7).*

Definition 12.24 (Tetralemma decision layer for anomalies)**.** Aggregate the six local meters and the global parity test into an *Anomaly Score* and a *Violation Witness*:

$$\mathsf{S}_{\text{ano}} := 1 - \frac{1}{\mathsf{N}_*}\left(|\mathsf{A}_{3^3}| + |\mathsf{A}_{2^3}| + |\mathsf{A}_{3^2Y}| + |\mathsf{A}_{2^2Y}| + |\mathsf{A}_{Y^3}| + |\mathsf{A}_{\text{grav}^2Y}|\right), \quad \mathsf{D}_{\text{ano}} := \left(\sum\nolimits_{\bullet} \mathsf{A}_{\bullet}^2\right)^{1/2},$$

with $\mathsf{N}_* > 0$ a normalization fixed by the GNS/KL conventions (Definition 12.11 and Proposition 12.12). Convert $(\mathsf{S}_{\text{ano}}, \mathsf{D}_{\text{ano}})$ to effects $E_{\mathsf{Th}}, E_{\mathsf{AntiTh}}, E_{\mathsf{SynTh}}, E_{\mathsf{HoloTh}}$ exactly as in Definition 12.13, gated by the holosymmetric overlay (Ch. 6). Outcomes are: *Thesis* (all meters vanish within tolerance), *Antithesis* (decisive nonzero), *Synthesis* (typed/band-localized anomaly signal), and *Holothesis* (diagnostic realized reference configuration not certified by the law–state visibilities).

Remark 12.25 (Coupling to Stage-2 representation matching)**.** The same band-extracted representation data used for Constraints D/E (Section 12.1.2) feed the anomaly meters of Definition 12.20. In particular, the operator identity $Q = I_3 + \frac{1}{2}Y$ and the spectral placements $\mathbf{1}_{\{y_f\}}(Y)\Psi_f = \Psi_f$ ensure consistent normalization of Y with the Dynkin-index conventions entering (12.46). Any alteration of hypercharges or multiplets immediately propagates to $\mathsf{A}_\bullet$.

Proposition 12.26 (Standard-Model calibration fixture)**.** *For one generation with assignments as in table 12.1, the local meters satisfy*

$$\mathsf{A}_{3^3} = \mathsf{A}_{2^3} = \mathsf{A}_{3^2Y} = \mathsf{A}_{2^2Y} = \mathsf{A}_{Y^3} = \mathsf{A}_{\mathrm{grav}^2Y} = 0,$$

and the global $SU(2)$ Witten test passes ($\mathsf{N}_2 = 4$ is even). Consequently, the anomaly decision layer (Definition 12.24) yields Thesis *with visibility $\Lambda^{[\mathrm{ano}]}$ close to 1 on $\mathfrak{P}$. This block serves to validate tagging, normalization, and Nielsen-flatness of all anomaly readouts.*

Proof. Bandwise SM content and meters. Work on the certified plateau $\mathfrak{P}$ with the split connection $\nabla_\mu = \nabla_\mu^{(\mathrm{LC})} + i\,\mathrm{ad}_{\mathcal{A}_\mu}$ (Ch. 6; Definitions 6.192 and 12.1). By band audit and tag commutation, representation extraction and expectations descend to $\mathcal{H}_{\mathrm{phys}}$ (Lemmas 12.2 and 12.7). For one generation, take the band-extracted left-chiral fermions $\{\Psi_f\}$ with representation data given by table 12.1 (encoded via Definition 12.19). By Definition 12.20, the six local meters $\mathsf{A}_{3^3}, \mathsf{A}_{2^3}, \mathsf{A}_{3^2Y}, \mathsf{A}_{2^2Y}, \mathsf{A}_{Y^3}, \mathsf{A}_{\mathrm{grav}^2Y}$ are *precisely* the coefficients of the tensor structures in (12.46) evaluated on this representation list (currents normalized by Definition 12.11).

Local cancellation for the SM assignment. The per-generation SM check displayed in Section 12.1.3 shows that each of the six sums in (12.46) vanishes for the assignments in table 12.1 (using $T(\mathbf{2}) = T(\mathbf{3}) = \frac{1}{2}$ and the oddness of the cubic index under conjugation). Therefore,

$$\mathsf{A}_{3^3} = \mathsf{A}_{2^3} = \mathsf{A}_{3^2Y} = \mathsf{A}_{2^2Y} = \mathsf{A}_{Y^3} = \mathsf{A}_{\mathrm{grav}^2Y} = 0$$

on the band for one SM generation. (As recorded below (12.46), the $SU(2)_L^3$ cubic anomaly is absent identically since the fundamental is pseudoreal.)

Global $SU(2)$ Witten test. Let $\Pi^{(2)}$ be the projector onto $SU(2)_L$ doublets and N_2 the doublet-count operator of Definition 12.22. For one generation, $\mathsf{N}_2 = 3$ (quark doublets) $+ 1$ (lepton doublet) $= 4$, hence even; equivalently the Witten-parity effect E_{Witten} equals $\mathbf{1}$ (Definition 12.22).

Decision layer and diagnostic realized reference configuration hygiene. Since all local meters vanish and the global test passes, the Anomaly Score/Violation Witness of Definition 12.24 yield the tetralemma outcome *Thesis* with visibility $\Lambda^{[\mathrm{ano}]}$ close to 1 on $\mathfrak{P}$. Because the meters are BRST-closed and tag-stable, these conclusions are gauge-parameter independent (Nielsen-flat) and descend to $\mathcal{H}_{\mathrm{phys}}$ by Lemma 12.2; Ward/KL hygiene follows from Proposition 12.12. This SM block therefore validates tagging, normalization, and the stability of the anomaly readouts, completing the calibration. □

Corollary 12.27 (Minimality and uniqueness at fixed content)**.** *Subject to the empirical fermion content of table 12.1, absence of exotics (beyond vectorlike pairs), and a single electroweak doublet mediator of Yukawa couplings, the product group $G_{\mathrm{SM}} = SU(3)_C \times SU(2)_L \times U(1)_Y$ is the unique minimal chiral gauge sector solving (12.46) on the certified plateau. Any shift of hypercharges or removal of a required multiplet renders at least one meter in Definition 12.20 nonzero, obstructing BRST nilpotency and violating Stage I of the HSP (Section 12.2).*

A Proof by Uniqueness (Prelude to Holosymmetry)

We do not *derive* G_{SM} ex nihilo. Instead, a tightly constrained operator–geometric and dynamical framework *singles it out* as the unique minimal solution. The logic is a *proof by uniqueness* carried out strictly on the certified BRST diagnostic realized reference configuration (Ch. 6):

1. **Start from universality.** Take the universal internal symmetry as the automorphism group of the represented observable algebra,

$$G_{\mathrm{int}} = \mathrm{Aut}\Big(\mathrm{End}(\mathcal{H}_{\mathrm{em}})\Big) \qquad \text{(eq. (12.99))},$$

   acting via the split, realized weave configuration-compatible connection $\nabla_\mu = \nabla_\mu^{(\mathrm{LC})} + i\,\mathrm{ad}_{\mathcal{A}_\mu}$ on the operator bundle (Definitions 6.192 and 12.1).

2. **Impose structural viability.** Enforce local implementability by a bona fide connection (Constraint A), compactness and representation-theoretic well-posedness (Constraint B), and genuine symmetry of the Tantrakyma dynamics (Constraint C), all formulated and tested on the diagnostic realized reference configuration (Section 12.1.1). These steps invoke compact holonomy and Peter–Weyl on finite fibres (Propositions 12.8 and 12.10) and the improved current/Ward framework with KL positivity (Definition 12.11 and Proposition 12.12).

3. **Match the fermion sector.** Require the empirically observed chiral content and charges to be realized as Tantrakyma spinor sections—*Chiral admissibility* and *Exact charge/Yukawa admissibility* (Constraints D and E)—with placements and operator identities taken from Sections 12.1.2 and 12.4.1 and table 12.1 and the operator equation $Q = I_3 + \frac{1}{2}Y$ (Chs. 6–7).

4. **Enforce anomaly freedom.** Impose the vanishing of all consistent local and relevant global anomalies (Constraint F) using the band-tagged meters of Definitions 12.20 and 12.22 and (12.46); equivalently, BRST nilpotency holds on $\mathcal{H}_{\mathrm{phys}}$ (Stage I HSP viability; Sections 12.2 and 12.3 and Proposition 12.21).

What survives this gauntlet is precisely

$$G_{\mathrm{SM}} = SU(3)_C \times SU(2)_L \times U(1)_Y,$$

with the canonical connection decomposition $\mathcal{A}_\mu = g_s\, G_\mu^a T_a + g\, W_\mu^i \tau_i + g' B_\mu Y$ (Definitions 12.95 and 12.96). Nothing strictly smaller (or differently charged) satisfies all four demands simultaneously; at fixed fermion content this minimality/uniqueness is summarized in Corollary 12.27 and calibrated by Proposition 12.26. This uniqueness result prepares the ground for the *Holosymmetric Selection Principle* (HSP), which provides a dynamical mechanism for *why* nature realizes exactly those properties that the uniqueness proof *characterizes* (Section 12.2).

From properties to mechanism. Having fixed the target properties—local compact gaugeability (A–B), symmetry of the Tantrakyma dynamics (C), chiral/charge realization (D–E), and complete anomaly freedom (F)—the question is elevated from classification to dynamics. In the IQR programme, selection is encoded by **Holosymmetry**: physical structure is realized by maximal relational coherence under interferometric scrutiny on the certified plateau (Ch. 6; law–state ledgers and plateau gate, Definition 10.80, and tetralemma/GNS effects, Definitions 1.18 and 12.13 and Remark 12.14).

Superposed candidates and the cosmic interferometer. Each admissible subgroup $G_A \subset G_{\text{int}}$ defines a complete candidate sector with its own connection, content, and couplings consistent with A–F. The holosymmetric viewpoint regards these viable sectors as coexisting in a quantum superposition. The **Holosymmetric Interferometer Model** prepares and recombines universal "paths" (two alternatives G_A vs. G_B) and reads a relational interference amplitude using diagnostic realized reference configuration-safe meters (Section 2.4; BRST/tag descent Lemma 12.2, microcausality and hyperbolicity Proposition 12.3). Selection becomes an operational criterion on theory space, reported with the same tetralemma decision calculus used for our constraints.

Principle of maximal coherence (informal statement). The realized gauge sector is the one that: (i) exhibits perfect *self*-coherence (passing Stage I: no anomalies, BRST nilpotency; Section 12.3 and Proposition 12.21) and (ii) is maximally orthogonal, under interferometric comparison, to all alternatives that also pass A–F (competitive suppression on the visibility plane; Ch. 6 ledgers and certified plateau). When read on the diagnostic realized reference configuration, the **Quantum Holo-State** $\left|\mathring{\mathbf{\Psi}}\right\rangle$ is the bright-fringe outcome of this cosmic interferometer: the globally consistent, maximally self-coherent realization of the Tantrakyma dynamics. The next section states the HSP precisely and develops the coherence functionals and visibility measures that operationalize this selection within the operator–bundle calculus (Section 12.2), leveraging the same KL/ Ward hygiene (Definition 12.11 and Proposition 12.12), compact holonomy (Proposition 12.8), and Peter–Weyl diagnostics (Proposition 12.10) already certified by Constraints A–F.

Conclusion (canonical connection decomposition). The *unique* expansion of a *single* relational gauge connection $\mathcal{A}_\mu$ valued in the Lie algebra that survives the filters of Constraints A–C (and D–F). By definition, the operator–bundle connection is a $\mathfrak{g}$-valued one–form acting through the split covariant derivative

$$\nabla_\mu \;=\; \nabla_\mu^{(\text{rel})}[\bar{\Sigma}] \;+\; i\,\text{ad}_{\mathcal{A}_\mu} \qquad \text{(cf. Definitions 6.7, 6.192 and 12.1).}$$

Once the admissible internal algebra is fixed to the direct sum

$$\mathfrak{g} \;=\; \mathfrak{su}(3) \;\oplus\; \mathfrak{su}(2) \;\oplus\; \mathfrak{u}(1)$$

(by compactness, representation well-posedness, symmetry of the dynamics, and the Stage-2/3 filters), there exists an orthonormal Hermitian basis $\{T_a\} \cup \{\tau_i\} \cup \{Y\}$ (color, weak, hypercharge) as guaranteed by the fibrewise unitary realization and compact holonomy

(Definition 12.5 and Proposition 12.8). Any $\mathfrak{g}$-valued one–form then expands *uniquely* as $\mathcal{A}_\mu = G^a_\mu T_a + W^i_\mu \tau_i + B_\mu Y$, while the couplings g_s, g, g' are the conventional normalizations fixed by the kinetic term and by current two–point normalization—equivalently, by rescalings that diagonalize the positive GNS/KL inner products (Definition 12.11 and Proposition 12.12). Thus the familiar coordinate expression

$$\mathcal{A}_\mu \;=\; g_s\, G^a_\mu T_a \;+\; g\, W^i_\mu \tau_i \;+\; g'\, B_\mu\, Y$$

is not an ad hoc insertion but the canonical form of a single $\mathfrak{g}$-valued connection in a basis adapted to the direct–sum decomposition.

The appearance of precisely these three blocks—color, weak isospin, and hypercharge—follows from the *filters* rather than choice: (i) compact holonomy on finite fibres and Peter–Weyl enforce unitary finite–multiplicity irreps and exclude noncompact directions (Propositions 12.8 and 12.10); (ii) symmetry of the Tantrakyma dynamics (Constraint C) requires that the same $\mathcal{A}_\mu$ render the Euler–Lagrange law covariant and the improved Ward identities valid on the certified plateau (Section 12.1.1 and Lemma 12.2); (iii) matching to the observed fermion content (Constraints D/E) selects the representation block that carries the weak chirality pattern and the operator identity $Q = I_3 + \frac{1}{2}Y$ with gauge–invariant Yukawas (Section 12.1.2); (iv) anomaly cancellation (Constraint F) removes hypercharge assignments or extra factors that would spoil BRST nilpotency (cf. Section 12.1.3 and Proposition 12.21). Any additional term would correspond to a generator outside $\mathfrak{su}(3) \oplus \mathfrak{su}(2) \oplus \mathfrak{u}(1)$; such directions are ruled out operationally by compactness/character diagnostics and structurally by D–F.

Relationally, the *same* $\mathcal{A}_\mu$ acts everywhere in the operator bundle via $i\,\mathrm{ad}_{\mathcal{A}_\mu}$, unified with $\nabla^{(\mathrm{LC})}$ on the base (Ch. 6). The field strength

$$\mathcal{F}_{\mu\nu} = \partial_\mu \mathcal{A}_\nu - \partial_\nu \mathcal{A}_\mu + i[\mathcal{A}_\mu, \mathcal{A}_\nu]$$

inherits the direct–sum structure: cross–commutators between distinct simple factors vanish, while commutators within a factor reproduce its structure constants (Definition 12.96). The carriers G^a_μ, W^i_μ, B_μ are therefore the *components* of one relational connection along the compact algebra directions certified by the selection programme; the decomposition is fixed jointly by the algebraic splitting of $\mathfrak{g}$, by kinetic/current normalization, and by the empirical representation content enforced in Stage 2–3—hence, not arbitrary.

12.2 The Holosymmetric Selection Principle (HSP)

The selection of the observed gauge structure is, within IQR, a genuinely quantum-mechanical inference enacted by *Holosymmetry*—the meta-operation that coherently compares entire histories (laws) rather than mere states (Section 3.8). Operationally, Holosymmetry instantiates the *Holothesis*: a Born-valued predicate about superposed *laws of evolution*, realized by a holosymmetric control system that branches, recombines, and reads out two complementary path-laws on dual registers (Definitions 1.4, 2.102 and 2.114). The "cosmic interferometer" thereby provides a metrological handle on the relational overlap of candidate universes.

Definition 12.28 (Unitarity Operator of a Candidate Theory)**.** Let $G_A \subset G_{\mathrm{int}}$ be a candidate internal symmetry group and let $\hat{H}^{(A)}$ denote the (possibly time-dependent) Relational

Hamiltonian associated with the corresponding theory (Definition 6.54). Assume $\hat{H}^{(A)}$ is self-adjoint on a dense, symmetry-stable domain $\mathcal{D} \subset \mathcal{H}_{\mathrm{em}}$ (or essentially self-adjoint with closure still denoted $\hat{H}^{(A)}$). The *unitary evolution operator* is

$$\mathbf{U}^{(A)} := \begin{cases} \exp\!\left(i\hat{H}^{(A)}\right), & \text{time-independent,} \\ \mathcal{T}\exp\!\left(i\int_{t_0}^{t_1}\hat{H}^{(A)}(t)\,dt\right), & \text{time-dependent,} \end{cases} \tag{12.47}$$

where $\mathcal{T}$ denotes time-ordering and the product is defined on $\mathcal{D}$. The *Unitarity Operator* is the positive operator

$$\mathbf{U}^{(A)}_{\mathrm{unitarity}} := \left(\mathbf{U}^{(A)}\right)^{\dagger}\mathbf{U}^{(A)}. \tag{12.48}$$

It acts on $\mathcal{H}_{\mathrm{em}}$ and enjoys the following properties:

(i) **Positivity and bounds.** $\mathbf{U}^{(A)}_{\mathrm{unitarity}} \geq 0$ and $0 \leq \langle\psi|\mathbf{U}^{(A)}_{\mathrm{unitarity}}|\psi\rangle \leq \|\mathbf{U}^{(A)}\|^2$ for all unit vectors $|\psi\rangle \in \mathcal{D}$.

(ii) **Gauge covariance, gauge invariance.** Under any fibrewise change of quantum frame W, $\mathbf{U}^{(A)} \mapsto W\,\mathbf{U}^{(A)}\,W^{\dagger}$, hence $\mathbf{U}^{(A)}_{\mathrm{unitarity}} \mapsto W\,\mathbf{U}^{(A)}_{\mathrm{unitarity}}\,W^{\dagger}$ and its expectation values are gauge-invariant.

(iii) **Characterization of unitarity.** The following are equivalent:

- (a) $\mathbf{U}^{(A)}$ is unitary on $\overline{\mathcal{D}}$;
- (b) $\mathbf{U}^{(A)}_{\mathrm{unitarity}} = \mathbb{I}$ on $\overline{\mathcal{D}}$;
- (c) $\langle\psi|\mathbf{U}^{(A)}_{\mathrm{unitarity}}|\psi\rangle = 1$ for all unit $|\psi\rangle \in \mathcal{D}$;
- (d) $\left\|\left(\mathbf{U}^{(A)}_{\mathrm{unitarity}} - \mathbb{I}\right)|\psi\rangle\right\| = 0$ for all $|\psi\rangle \in \mathcal{D}$.

Remark 12.29 (Relation to BRST consistency and anomalies)**.** In a gauge-fixed formulation (Section 12.3), unitarity on the physical Hilbert space $\mathcal{H}_{\mathrm{phys}} = \ker Q_{\mathrm{BRST}}/\operatorname{im} Q_{\mathrm{BRST}}$ is equivalent to (a) nilpotency $Q^2_{\mathrm{BRST}} = 0$, (b) BRST invariance of the dynamics $[Q_{\mathrm{BRST}}, \hat{H}^{(A)}] = 0$, and (c) trivial Jacobian for gauge transformations of the functional measure. Any perturbative or global anomaly violates at least one of (a)–(c) and manifests as $\mathbf{U}^{(A)}_{\mathrm{unitarity}} \neq \mathbb{I}$ on $\mathcal{H}_{\mathrm{phys}}$.

Self-coherence functional.

Definition 12.30 (Self–Coherence Functional (state–general))**.** Let ρ_0 be a normalized density operator (pure or mixed) on $\mathcal{H}_{\mathrm{em}}$. For a candidate theory A with Unitarity Operator $\mathbf{U}^{(A)}_{\mathrm{unitarity}} := \left(\mathbf{U}^{(A)}\right)^{\dagger}\mathbf{U}^{(A)}$ as in Definition 12.28, define

$$\mathcal{C}(A \mid \rho_0) := \left|\operatorname{Tr}\!\left(\rho_0\,\mathbf{U}^{(A)}_{\mathrm{unitarity}}\right)\right|. \tag{12.49}$$

This quantity satisfies:

(i) **Bounds.** $0 \leq \mathcal{C}(A \mid \rho_0) \leq 1$.

(ii) **Pure–state reduction.** If $\rho_0 = |\Psi_0\rangle\langle\Psi_0|$, then

$$\mathcal{C}(A) = \left|\langle\Psi_0|\mathbf{U}^{(A)}_{\mathrm{unitarity}}|\Psi_0\rangle\right| \qquad (\textit{Section 12.2}). \tag{12.50}$$

(iii) **Extremality / characterisation.** $\mathcal{C}(A \mid \rho_0) = 1$ iff $\mathbf{U}^{(A)}_{\text{unitarity}} = \mathbb{I}$ on $\text{supp}(\rho_0)$ (equivalently, $(\mathbf{U}^{(A)})^\dagger \mathbf{U}^{(A)} = \mathbb{I}$ on $\text{supp}(\rho_0)$).

(iv) **Gauge invariance.** Under any fibrewise change of quantum frame U on $\mathcal{H}_{\text{em}}$, $\rho_0 \mapsto U\rho_0 U^\dagger$ and $\mathbf{U}^{(A)}_{\text{unitarity}} \mapsto U\,\mathbf{U}^{(A)}_{\text{unitarity}}\,U^\dagger$, hence $\mathcal{C}(A \mid \rho_0)$ is invariant.

Definition 12.31 (Holosymmetric Self–Coherence Functional (register–lifted))**.** Let $\mathcal{H}_c = \text{span}\{|\mathbf{0}\rangle_c, |\mathbf{1}\rangle_c\}$ be the control space (as in Definition 2.102) and $\mathcal{H}^{(2n)}_{\text{RR}}$ the dual n–PQS register space (Ch. II; cf. Definition 2.113). For a candidate theory A with register–level path operators $\hat{\mathbf{U}}^{A_G B_N}_{\text{RR}}$ and $\hat{\mathbf{U}}^{A_N B_G}_{\text{RR}}$ (Ch. II–III; see Definitions 2.103 and 2.108), let $\mathbf{U}^{(\star)}_{holo}[A]$ denote the holosymmetric interferometer on $\mathcal{H}_c \otimes \mathcal{H}^{(2n)}_{\text{RR}}$, with $\star \in \{\text{sym}, \text{asym}\}$:

$$\mathbf{U}^{(\text{sym})}_{\text{holo}}[A] := \mathbf{U}_{\text{Had}}\,\mathbf{U}_{\text{ctrl}}[A]\,\mathbf{U}_{\text{Had}} \qquad \text{(Definition 2.123)},$$
$$\mathbf{U}^{(\text{asym})}_{\text{holo}}[A] := \mathbf{U}_{\text{read}}\,\mathbf{U}_{\text{ctrl}}[A]\,\mathbf{U}_{\text{prep}} \qquad \text{(Definition 2.124)},$$

where $\mathbf{U}_{\text{ctrl}}[A]$ is the controlled holosymmetric operator built from $\hat{\mathbf{U}}^{A_G B_N}_{\text{RR}}$ and $\hat{\mathbf{U}}^{A_N B_G}_{\text{RR}}$ (Definition 2.114). For any normalized probe state $\rho_{c\text{RR},0}$ on $\mathcal{H}_c \otimes \mathcal{H}^{(2n)}_{\text{RR}}$, define the *register–lifted self–coherence functional*

$$\mathcal{C}_{\text{holo}}(A \mid \rho_{c\text{RR},0};\, \star) \;:=\; \Big|\,\text{Tr}\Big(\rho_{c\text{RR},0}\,\mathbf{U}^{(A);\text{holo},(\star)}_{\text{unitarity}}\Big)\Big|, \qquad \mathbf{U}^{(A);\text{holo},(\star)}_{\text{unitarity}} := \Big(\mathbf{U}^{(\star)}_{\text{holo}}[A]\Big)^\dagger \mathbf{U}^{(\star)}_{\text{holo}}[A]. \tag{12.51}$$

This quantity satisfies:

(i) **Bounds.** $0 \le \mathcal{C}_{\text{holo}}(A \mid \rho_{c\text{RR},0};\, \star) \le 1$.

(ii) **Pure–state reduction.** If $\rho_{c\text{RR},0} = |\Psi_{c\text{RR}}\rangle\langle\Psi_{c\text{RR}}|$, then

$$\mathcal{C}_{\text{holo}}(A \mid \rho_{c\text{RR},0};\, \star) = \Big\|\mathbf{U}^{(\star)}_{\text{holo}}[A]\,|\Psi_{c\text{RR}}\rangle\Big\|^2 \quad \text{(cf. eq. (12.49))}.$$

(iii) **Extremality / characterisation.** $\mathcal{C}_{\text{holo}}(A \mid \rho_{c\text{RR},0};\, \star) = 1$ iff $\mathbf{U}^{(\star)}_{\text{holo}}[A]$ is an isometry on $\text{supp}(\rho_{c\text{RR},0})$; equivalently, $\mathbf{U}^{(A);\text{holo},(\star)}_{\text{unitarity}} = \mathbb{I}$ on $\text{supp}(\rho_{c\text{RR},0})$ (cf. Definition 12.28).

(iv) **Gauge invariance.** For any fibrewise change of quantum frame U on $\mathcal{H}_c \otimes \mathcal{H}^{(2n)}_{\text{RR}}$, $\rho_{c\text{RR},0} \mapsto U\rho_{c\text{RR},0}U^\dagger$ and $\mathbf{U}^{(\star)}_{\text{holo}}[A] \mapsto U\,\mathbf{U}^{(\star)}_{\text{holo}}[A]\,U^\dagger$, hence $\mathcal{C}_{\text{holo}}$ is invariant (see the bundle–covariance discussion in Ch. III; Definition 2.114).

(v) **Compatibility with the state–general functional.** If $\rho_{c\text{RR},0} = \rho_c \otimes \rho_0$ and $\mathbf{U}^{(\star)}_{\text{holo}}[A] = \mathbf{U}_c \otimes \widehat{\mathbf{U}}^{(A)}$ with a register–lift $\widehat{\mathbf{U}}^{(A)}$ of $\mathbf{U}^{(A)}$ (cf. Definition 12.28), then $\mathcal{C}_{\text{holo}}(A \mid \rho_{c\text{RR},0};\, \star) = \mathcal{C}(A \mid \rho_0)$ as in eq. (12.49).

Remark 12.32 (Symmetric visibility and the $\hat{\mathbf{U}}_\pm$ blocks)**.** For the symmetric holosymmetric interferometer

$$\mathbf{U}^{(\text{sym})}_{\text{holo}}[A] = \mathbf{U}_{\text{Had}}\,\mathbf{U}_{\text{ctrl}}[A]\,\mathbf{U}_{\text{Had}},$$

with $\mathbf{U}_{\text{Had}}$ as in eq. (2.78) and the controlled holosymmetric operator $\mathbf{U}_{\text{ctrl}}$ as in Definition 2.114 (built from the register-level path operators of Definition 2.108), the lifted unitarity operator is the identity, $\mathbf{U}^{(A);\text{holo},(\text{sym})}_{\text{unitarity}} = \mathbb{I}$. Consequently $\mathcal{C}_{\text{holo}}(A \mid \rho_{c\text{RR},0};\, \text{sym}) = 1$ for every normalized probe (cf. Definition 12.31; see also Section 3.1.2 and Theorem 3.4).

Discrimination therefore appears in the *readout probabilities*

$$p_0 = \tfrac{1}{4}\big\|\hat{\mathbf{U}}_+|\Psi_{\mathrm{RR}}\rangle\big\|^2, \qquad p_1 = \tfrac{1}{4}\big\|\hat{\mathbf{U}}_-|\Psi_{\mathrm{RR}}\rangle\big\|^2, \qquad \hat{\mathbf{U}}_\pm := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} \pm \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G},$$

whence the symmetric visibility is

$$\mathcal{V}_{\mathrm{sym}} := |p_0 - p_1| = \Big|\mathrm{Re}\,\langle\Psi_{\mathrm{RR}}|\,(\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G})^\dagger \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}\,|\Psi_{\mathrm{RR}}\rangle\Big| \;=\; \Big|\mathrm{Re}\,\mathrm{Tr}\Big(\rho_{\mathrm{RR}}\,(\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G})^\dagger \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}\Big)\Big|$$

(for pure $|\Psi_{\mathrm{RR}}\rangle$ and mixed ρ_{RR}, respectively) as developed in Sections 3.1.2 and 3.1.4. Thus, Stage-I self-coherence is trivial in the symmetric lift, while Stage-II comparative tomography hinges on the $\hat{\mathbf{U}}_\pm$ blocks and $\mathcal{V}_{\mathrm{sym}}$ (and, more generally, its asymmetric SU(2) generalization; cf. Section 3.1.5 and eq. (3.56)).

Definition 12.33 (Viability defect and state-independent viability)**.** Define the viability defect $\varepsilon(A \mid \rho_0) := 1 - \mathcal{C}(A \mid \rho_0) \geq 0$. Let $\mathcal{S}_{\mathrm{IC}}$ be informationally complete on a subspace $\mathcal{K} \subseteq \mathcal{H}_{\mathrm{em}}$. We say A is *internally viable on* $\mathcal{K}$ if

$$\sup_{\rho_0 \in \mathcal{S}_{\mathrm{IC}}} \varepsilon(A \mid \rho_0) = 0 \quad \Longleftrightarrow \quad \mathbf{U}^{(A)}_{\mathrm{unitarity}} = \mathbb{I} \text{ on } \mathcal{K}. \tag{12.52}$$

Remark 12.34 (Meaning of state–independent viability)**.** Let $\varepsilon(A \mid \rho_0)$ and "internally viable on $\mathcal{K}$" be as in Definition 12.33. Then:

(i) **Uniform (state–independent) pass of Stage I.** Since $\varepsilon(A \mid \rho_0) \geq 0$, the condition $\sup_{\rho_0 \in \mathcal{S}_{\mathrm{IC}}} \varepsilon(A \mid \rho_0) = 0$ is equivalent to $\mathcal{C}(A \mid \rho_0) = 1$ for *every* $\rho_0 \in \mathcal{S}_{\mathrm{IC}}$ (cf. Definition 12.30). By informational completeness on $\mathcal{K}$, this holds iff $\mathbf{U}^{(A)}_{\mathrm{unitarity}} = \mathbb{I}$ on $\mathcal{K}$ (cf. Definition 12.28).

(ii) **Operational reading (tomography).** Any deviation of $\mathbf{U}^{(A)}_{\mathrm{unitarity}}$ from $\mathbb{I}$ on $\mathcal{K}$ is detected by at least one probe $\rho_0 \in \mathcal{S}_{\mathrm{IC}}$ via $\mathcal{C}(A \mid \rho_0) < 1$. Thus, viability is a *uniform* (state–independent) unitarity certificate over the scanned subspace.

(iii) **Choice of probes.** The statement is invariant under the choice of IC set: replacing $\mathcal{S}_{\mathrm{IC}}$ by any other IC family on the same $\mathcal{K}$ leads to the same viability verdict. In particular, one may use an IC–POVM ensemble or an overcomplete coherent–state family.

(iv) **Register–lifted context.** For the symmetric holosymmetric lift (Definition 12.31), $\mathbf{U}^{(A);\mathrm{holo},(\mathrm{sym})}_{\mathrm{unitarity}} = \mathbb{I}$, so $\mathcal{C}_{\mathrm{holo}}(A \mid \rho_{\mathrm{cRR},0}; \mathrm{sym}) = 1$ for all probes (see Remark 12.32). Hence Stage I is trivial in the symmetric interferometer and discrimination is deferred to visibility/overlap diagnostics; asymmetry (SU(2) prep/read) restores sensitivity.

(v) **Scope.** If $\mathcal{K} = \mathcal{H}_{\mathrm{em}}$, "internally viable" means $\mathbf{U}^{(A)}_{\mathrm{unitarity}} = \mathbb{I}$ on the full emergent space; if $\mathcal{K} \subsetneq \mathcal{H}_{\mathrm{em}}$, the certificate is restricted to the scanned sector $\mathcal{K}$.

Proposition 12.35 (Basic bounds and gauge invariance of $\mathcal{C}$)**.** *Let $\mathcal{C}(A \mid \rho_0)$ be the self–coherence functional of Definition 12.30, with $\mathbf{U}^{(A)}_{\mathrm{unitarity}}$ as in Definition 12.28. For any density operator ρ_0 and any candidate A:*

(i) Bounds and norm gap

One has $0 \le \mathcal{C}(A \mid \rho_0) \le 1$, *and*

$$1 - \mathcal{C}(A \mid \rho_0) \;\le\; \Big\| \mathbf{U}^{(A)}_{\text{unitarity}} - \mathbb{I} \Big\|.$$

(ii) Gauge invariance

Under any fibrewise change of quantum frame U, $\rho_0 \mapsto U\rho_0 U^\dagger$ *and* $\mathbf{U}^{(A)} \mapsto U\mathbf{U}^{(A)}U^\dagger$, *hence* $\mathbf{U}^{(A)}_{\text{unitarity}} \mapsto U\,\mathbf{U}^{(A)}_{\text{unitarity}}\,U^\dagger$ *and* $\mathcal{C}(A \mid \rho_0)$ *is invariant.*

(iii) Informational completeness (IC) ⇒ unitarity on the scanned sector

If $\mathcal{C}(A \mid \rho) = 1$ *for all* ρ *in an informationally complete set on a subspace* $\mathcal{K} \subseteq \mathcal{H}_{\text{em}}$, *then* $\mathbf{U}^{(A)}_{\text{unitarity}} = \mathbb{I}$ *on* $\mathcal{K}$ *(equivalently,* A *is internally viable on* $\mathcal{K}$ *per Definition 12.33).*

Proof. (a) By Definition 12.30, $\mathcal{C}(A \mid \rho_0) = \Big|\operatorname{Tr}\Big(\rho_0\,\mathbf{U}^{(A)}_{\text{unitarity}}\Big)\Big|$ with $\rho_0 \ge 0$, $\operatorname{Tr}\rho_0 = 1$, and $\mathbf{U}^{(A)}_{\text{unitarity}} = (\mathbf{U}^{(A)})^\dagger\mathbf{U}^{(A)} \ge 0$. Thus $0 \le \operatorname{Tr}(\rho_0\,\mathbf{U}^{(A)}_{\text{unitarity}})$ and so $0 \le \mathcal{C} \le 1$ by the positivity/norm bound in Definition 12.30. For the norm gap, write $\mathbf{U}^{(A)}_{\text{unitarity}} = \mathbb{I} + \Delta$ and use Hölder's inequality:

$$\mathcal{C} = \Big|\operatorname{Tr}(\rho_0(\mathbb{I}+\Delta))\Big| \ge 1 - \Big|\operatorname{Tr}(\rho_0\,\Delta)\Big| \ge 1 - \|\rho_0\|_1\,\|\Delta\| = 1 - \|\Delta\|,$$

since $\|\rho_0\|_1 = \operatorname{Tr}\rho_0 = 1$. Rearranging gives $1 - \mathcal{C}(A \mid \rho_0) \le \|\mathbf{U}^{(A)}_{\text{unitarity}} - \mathbb{I}\|$.

(b) By Definition 12.28, under U we have $\mathbf{U}^{(A)}_{\text{unitarity}} \mapsto U\,\mathbf{U}^{(A)}_{\text{unitarity}}\,U^\dagger$. Then by cyclicity of the trace,

$$\mathcal{C}'(A \mid \rho_0') = \Big|\operatorname{Tr}\Big(U\rho_0U^\dagger\,U\,\mathbf{U}^{(A)}_{\text{unitarity}}\,U^\dagger\Big)\Big| = \Big|\operatorname{Tr}(\rho_0\,\mathbf{U}^{(A)}_{\text{unitarity}})\Big| = \mathcal{C}(A \mid \rho_0),$$

so $\mathcal{C}$ is gauge-invariant (cf. the gauge-invariance clause in Definition 12.30).

(c) If $\mathcal{C}(A \mid \rho) = 1$ for all ρ in an informationally complete set on $\mathcal{K}$, then (using positivity) $\operatorname{Tr}\Big(\rho\,\mathbf{U}^{(A)}_{\text{unitarity}}\Big) = 1$ for all such ρ. Hence $\operatorname{Tr}\Big(\rho\,(\mathbf{U}^{(A)}_{\text{unitarity}} - \mathbb{I})\Big) = 0$ for all ρ in the IC set. Informational completeness on $\mathcal{K}$ implies that any operator with vanishing expectation on all probes in the set must vanish on $\mathcal{K}$; therefore $\mathbf{U}^{(A)}_{\text{unitarity}} - \mathbb{I} = 0$ on $\mathcal{K}$, i.e. $\mathbf{U}^{(A)}_{\text{unitarity}} = \mathbb{I}$ on $\mathcal{K}$. By Definition 12.33 and Remark 12.34, this is precisely "internal viability on $\mathcal{K}$". □

Remark 12.36 (Operational readout via the symmetric interferometer)**.** The probabilities in the symmetric holosymmetric interferometer satisfy $p_0 = \frac{1}{4}\|\hat{\mathbf{U}}_+|\Psi_{\text{RR}}\rangle\|^2$, $p_1 = \frac{1}{4}\|\hat{\mathbf{U}}_-|\Psi_{\text{RR}}\rangle\|^2$, $p_0{+}p_1 = 1$ (Section 3.1.2 and Theorem 3.4). The visibility extracted from these outcomes reproduces $\mathcal{C}(A)$ up to the normalization specified in eqs. (12.49) and (12.50), providing a physically measurable enforcement of the viability condition Definition 12.33.

12.2.1 Holothesis and the Superposition of Laws (Holosymmetry)

Fix a dual-register arena $\mathcal{H}^{(2n)}_{\text{RR}}$ and a single-qubit control $\mathcal{H}_c$ (as in Definitions 2.102 and 2.113). The holosymmetric circuit prepares a control superposition (via $\mathbf{U}_{\text{Had}}$; cf. eq. (2.78)), applies a controlled pair of lifted path-laws to the two registers $\Big(\hat{\mathbf{U}}^{A_G B_N}_{\text{RR}},\ \hat{\mathbf{U}}^{A_N B_G}_{\text{RR}}\Big)$ through the controlled operator $\mathbf{U}_{\text{ctrl}}$ (see Definitions 2.108 and 2.114), recombines paths with either the

symmetric or asymmetric interferometer $\mathbf{U}_{\text{holo}}^{(\text{sym})}$ or $\mathbf{U}_{\text{holo}}^{(\text{asym})}$ (Definitions 2.123 and 2.124), and finally measures the control in the $\{|\mathbf{0}\rangle_c, |\mathbf{1}\rangle_c\}$ basis.

The Holothesis is read as a Born valuation of the *holosymmetric effect*

$$E_{\text{Holothesis}}^{(\mathbf{a}\to\mathbf{b})} = \Big\langle \mathbf{a}\Big|_c \, \mathbf{U}_{\text{holo}}^{\dagger}(|\mathbf{b}\rangle\langle\mathbf{b}|_c \otimes \mathbb{I})\mathbf{U}_{\text{holo}} \, \Big|\mathbf{a}\Big\rangle_c, \tag{12.53}$$

so that $v_{\text{Holothesis}}(\rho_{\text{RR}}) = \text{Tr}\Big(E_{\text{Holothesis}}^{(\mathbf{a}\to\mathbf{b})}\rho_{\text{RR}}\Big)$ (Definition 1.18 and Section 3.8; see also the register-lifted self-coherence in Definition 12.31).

Conceptually, the control readout is an *interference visibility meter* for the coherence (agreement/dissonance) between two *complete* path-laws (universal histories) acting on the same input: in the symmetric case this is captured by the $\hat{\mathbf{U}}_{\pm}$ blocks and the visibility $\mathcal{V}_{\text{sym}}$ (Remark 12.32 and Section 3.1.2 and Theorem 3.4); the asymmetric SU(2) variant generalizes the readout to full process tomography (Definition 2.124). This is the operational content of Holosymmetry as a meta-level symmetry over laws (Section 3.8).

Remark 12.37 (Fiducial probe state $|\Psi_0\rangle$)**.**

(i) **Definition.** $|\Psi_0\rangle \in \mathcal{H}_{\text{em}}$ denotes a unit (pure) probe state. It is the pure-state instance of the state–general probe ρ_0 in Definition 12.30, i.e. $\mathcal{C}(A \mid \rho_0) = \Big|\text{Tr}(\rho_0\, \mathbf{U}_{\text{unitarity}}^{(A)})\Big|$ reduces, for $\rho_0 = |\Psi_0\rangle\langle\Psi_0|$, to $\mathcal{C}(A) = \Big|\langle\Psi_0|\, \mathbf{U}_{\text{unitarity}}^{(A)}\, |\Psi_0\rangle\Big|$ (eqs. (12.49) and (12.50)).

(ii) **Evaluation.** With $\mathbf{U}_{\text{unitarity}}^{(A)} := (\mathbf{U}^{(A)})^{\dagger}\mathbf{U}^{(A)} \succeq 0$ (Definition 12.28),

$$\mathcal{C}(A) = \langle\Psi_0|\, \mathbf{U}_{\text{unitarity}}^{(A)}\, |\Psi_0\rangle = \Big\| \, \mathbf{U}^{(A)}|\Psi_0\rangle \, \Big\|^2,$$

hence $0 \le \mathcal{C}(A) \le 1$ with the bounds and norm gap as in Proposition 12.35.

(iii) **Role in Stage I (symmetric interferometer).** In the symmetric holosymmetric interferometer, $\mathcal{V}_{\text{sym}}(A, A) = \Big|\text{Re}\,\langle\Psi_0|(\mathbf{U}^{(A)})^{\dagger}\mathbf{U}^{(A)}|\Psi_0\rangle\Big| = \mathcal{C}(A)$ (Section 3.1.2, Theorem 3.4, and Remark 12.32). Therefore $\mathcal{C}(A) = 1$ iff $(\mathbf{U}^{(A)})^{\dagger}\mathbf{U}^{(A)} = \mathbb{I}$ on the support of $|\Psi_0\rangle$.

(iv) **State–independent strengthening.** To certify unitarity on a subspace $\mathcal{K} \subseteq \mathcal{H}_{\text{em}}$, replace $|\Psi_0\rangle$ by an informationally complete family $\mathcal{S}_{\text{IC}}$ on $\mathcal{K}$ and require $\sup_{\rho_0\in\mathcal{S}_{\text{IC}}} \Big(1 - \mathcal{C}(A \mid \rho_0)\Big) = 0$. By Definition 12.33 and Remark 12.34 this holds iff $\mathbf{U}_{\text{unitarity}}^{(A)} = \mathbb{I}$ on $\mathcal{K}$.

(v) **Register–lifted counterpart.** When the probe is prepared on $\mathcal{H}_c \otimes \mathcal{H}_{\text{RR}}^{(2n)}$ (either $|\Psi_{c\text{RR}}\rangle$ or $\rho_{c\text{RR},0}$), the same interpretation applies with the lifted operator in Definition 12.31.

12.2.2 Stage I — The Self-Consistency Filter (Symmetric Interferometer)

Let $G_A \subset G_{\text{int}}$ be a candidate internal symmetry and $\mathbf{U}^{(A)} = \exp\Big(i\hat{H}^{(A)}\Big)$ the corresponding evolution on the universal state space (Definition 12.28). In the *symmetric* interferometer, the control is prepared and read in the Hadamard basis and the two branches enact the same candidate law (cf. Section 3.1.2). The control-outcome probabilities are determined by the constructive/destructive path-operators $\hat{\mathbf{U}}_{\pm}$:

$$p_0 = \tfrac{1}{4}\,\|\hat{\mathbf{U}}_{+}|\Psi_{\text{RR}}\rangle\|^2, \qquad p_1 = \tfrac{1}{4}\,\|\hat{\mathbf{U}}_{-}|\Psi_{\text{RR}}\rangle\|^2, \qquad p_0{+}p_1 = 1 \tag{12.54}$$

(Theorem 3.4; see also Remark 12.32). The *self-coherence functional* is the symmetric visibility evaluated on a theory against itself,

$$\begin{aligned}\mathcal{C}(A) &:= \mathcal{V}_{\mathrm{sym}}(A, A) \\ &= \left|\mathrm{Re}\,\langle\Psi_0|(\mathbf{U}^{(A)})^\dagger\mathbf{U}^{(A)}|\Psi_0\rangle\right| \qquad (12.55)\end{aligned}$$

(Special case of Definition 12.30 with $\rho_0 = |\Psi_0\rangle\langle\Psi_0|$; *eqs.* (12.49)*and* (12.50)).

Postulate 12.1 (Viability)**.** *A candidate A is* internally viable *iff $\mathcal{C}(A) = 1$. Equivalently, the symmetric interferometer yields unit total visibility only when $\mathbf{U}^{(A)}$ is unitary on the support of $|\Psi_0\rangle$ (hence $(\mathbf{U}^{(A)})^\dagger\mathbf{U}^{(A)} = \mathbb{I}$ on that line). Non-unitary defects (e.g. gauge anomalies) depress $\mathcal{C}(A)<1$ and are dynamically filtered out (Definition 12.33 and Proposition 12.35).*

Proof. (i) *Reduction to a positive expectation.* By Definition 12.28, $\mathbf{U}^{(A)}_{\mathrm{unitarity}} := (\mathbf{U}^{(A)})^\dagger\mathbf{U}^{(A)} \succeq 0$. Hence $\langle\Psi_0|\mathbf{U}^{(A)}_{\mathrm{unitarity}}|\Psi_0\rangle \in [0, 1]$ and is real. Therefore the "Re" in the definition of $\mathcal{C}(A)$ is redundant, and

$$\mathcal{C}(A) = \langle\Psi_0|\mathbf{U}^{(A)}_{\mathrm{unitarity}}|\Psi_0\rangle = \left\| \mathbf{U}^{(A)}|\Psi_0\rangle \right\|^2,$$

which is exactly the pure-state case of Definition 12.30 (also eq. (12.50)).

(ii) *Bounds.* From (i), $0 \le \mathcal{C}(A) \le 1$, agreeing with Proposition 12.35(a).

(iii) *If part.* If $\mathbf{U}^{(A)}$ is unitary on $\overline{\mathrm{span}}\{|\Psi_0\rangle\}$, then $\mathbf{U}^{(A)}_{\mathrm{unitarity}} = \mathbb{I}$ on that (one-dimensional) support, so $\mathcal{C}(A) = \langle\Psi_0|\mathbb{I}|\Psi_0\rangle = 1$.

(iv) *Only-if part.* If $\mathcal{C}(A) = 1$, then $\|\mathbf{U}^{(A)}|\Psi_0\rangle\| = 1$, so $(\mathbf{U}^{(A)})^\dagger\mathbf{U}^{(A)}$ has expectation 1 on $|\Psi_0\rangle$. But $\mathbf{U}^{(A)}_{\mathrm{unitarity}} \succeq 0$ and $\|\mathbf{U}^{(A)}_{\mathrm{unitarity}}\| \le 1$ on physical states (Proposition 12.35(a)); the only way a positive contraction has unit expectation on a unit vector is if it acts as the identity on that vector. Thus $\mathbf{U}^{(A)}_{\mathrm{unitarity}}|\Psi_0\rangle = |\Psi_0\rangle$, i.e. $(\mathbf{U}^{(A)})^\dagger\mathbf{U}^{(A)} = \mathbb{I}$ on the support of $|\Psi_0\rangle$.

(v) *Connection to the symmetric interferometer.* From Section 3.1.2 and Theorem 3.4 (and Remark 12.32), the symmetric readout satisfies $\mathcal{V}_{\mathrm{sym}}(A, A) = \left|\mathrm{Re}\,\langle\Psi_0|(\mathbf{U}^{(A)})^\dagger\mathbf{U}^{(A)}|\Psi_0\rangle\right|$, coinciding with $\mathcal{C}(A)$ in (i). Therefore the symmetric interferometer implements the Stage-I "unitarity vs. defect" test: if $(\mathbf{U}^{(A)})^\dagger\mathbf{U}^{(A)} \neq \mathbb{I}$, then by Proposition 12.35(a)

$$1 - \mathcal{C}(A) \;\le\; \left\|\mathbf{U}^{(A)}_{\mathrm{unitarity}} - \mathbb{I}\right\| \;>\; 0,$$

so the visibility drops below unity and A fails the self-consistency filter.

(vi) *State-independent strengthening.* If one requires viability not just for a single probe $|\Psi_0\rangle$ but uniformly over an informationally complete family on a subspace $\mathcal{K}$, then by Definition 12.33 (see also Remark 12.34) the condition becomes $\sup_{\rho_0\in\mathcal{S}_{\mathrm{IC}}}\big(1 - \mathcal{C}(A \mid \rho_0)\big) = 0$ $\iff \mathbf{U}^{(A)}_{\mathrm{unitarity}} = \mathbb{I}$ on $\mathcal{K}$. □

12.2.3 Stage II — Comparative Tomography and Relational Orthogonality (Asymmetric Interferometer)

Among *internally viable* candidates (cf. Definition 12.33), selection is performed by *comparative* interference. The *asymmetric* holosymmetric interferometer permits arbitrary control

preparation/readout and thus tomographically complete access to the complex overlap of two viable laws A, B acting on the same input (see Definition 2.124).

Let the control $SU(2)$ gates be

$$U_c^{\text{prep}} = \begin{pmatrix} \alpha & \beta \\ -\beta^* & \alpha^* \end{pmatrix}, \qquad U_c^{\text{read}} = \begin{pmatrix} \gamma & \delta \\ -\delta^* & \gamma^* \end{pmatrix}, \quad \text{with} \quad |\alpha|^2+|\beta|^2 = |\gamma|^2+|\delta|^2 = 1.$$

Then (derivation in Section 3.1.5) the control-outcome probabilities for a pure probe $|\Psi_0\rangle$ on $\mathcal{H}_{\text{em}}$ are

$$p_0^{(\text{asym})}(A,B) = \left\| \left(\gamma\alpha\, \mathbf{U}^{(A)} - \delta^*\beta\, \mathbf{U}^{(B)}\right) |\Psi_0\rangle \right\|^2, \tag{12.56}$$

$$p_1^{(\text{asym})}(A,B) = \left\| \left(\delta\alpha\, \mathbf{U}^{(A)} + \gamma^*\beta\, \mathbf{U}^{(B)}\right) |\Psi_0\rangle \right\|^2, \tag{12.57}$$

with $\mathbf{U}^{(\cdot)}$ the emergent evolutions (Definition 12.28). The corresponding *asymmetric visibility* is

$$\mathcal{V}_{\text{asym}}(A,B) := \left| p_0^{(\text{asym})} - p_1^{(\text{asym})} \right| \quad (\textit{Section 3.1.7}). \tag{12.58}$$

Tomographic interpretation and relational orthogonality. By scanning (α,β) and (γ,δ), the pair $\left(p_0^{(\text{asym})}(A,B),\, p_1^{(\text{asym})}(A,B)\right)$ recovers the full *complex* overlap $\langle\Psi_0|(\mathbf{U}^{(B)})^\dagger\mathbf{U}^{(A)}|\Psi_0\rangle$; in particular, the symmetric case recovers its real part (cf. Remark 12.32), while suitable choices of $U_c^{\text{prep}}, U_c^{\text{read}}$ extract the imaginary part (Section 3.1.7). A pair of viable candidates A, B is *relationally orthogonal* on a subspace $\mathcal{K}$ if this overlap vanishes for all probes from an informationally complete family on $\mathcal{K}$ (so the corresponding effects are indistinguishable on $\mathcal{K}$ via Born valuation; cf. Definition 12.30).

Register–lifted form (consistency with Chs. II–III). In the explicit holosymmetric circuit on control+registers, replace the emergent evolutions by the register–lifted construction:

$$\mathbf{U}_{\text{holo}}^{(\text{asym})}[A] = \mathbf{U}_{\text{read}}\, \mathbf{U}_{\text{ctrl}}[A]\, \mathbf{U}_{\text{prep}},$$
$$\mathbf{U}_{\text{ctrl}}[A] \text{ is built from } \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N},\ \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} \quad (\textit{Definitions 2.108, 2.114 and 2.124}).$$

For a probe $\rho_{c\text{RR},0}$ on $\mathcal{H}_c \otimes \mathcal{H}_{\text{RR}}^{(2n)}$, the analysis is expressed by the register–lifted self–coherence/visibility functionals of Definition 12.31, reducing to (12.56)–(12.57) for pure product probes and appropriate identifications of $\mathbf{U}^{(A)}, \mathbf{U}^{(B)}$ with their lifts.

Definition 12.38 (Relational overlap and holosymmetric orthogonality)**.** Let A, B be *viable* candidates (cf. Postulate 12.1 and Definition 12.33) with emergent evolutions $\mathbf{U}^{(A)}, \mathbf{U}^{(B)}$ as in Definition 12.28, and fix a fiducial pure probe $|\Psi_0\rangle \in \mathcal{H}_{\text{em}}$ (Definition 12.30). The *holosymmetric overlap* is

$$\Omega(A, B \mid \Psi_0) := \langle\Psi_0|\, (\mathbf{U}^{(B)})^\dagger\, \mathbf{U}^{(A)}\, |\Psi_0\rangle, \tag{12.59}$$

which is the complex quantity tomographically accessed by the *asymmetric* holosymmetric interferometer (Definition 2.124 and Section 3.1.5).

Define the *relational orthogonality index*

$$\rho(A,B) := \min_{U_c^{\mathrm{prep}},\,U_c^{\mathrm{read}}\in SU(2)} \Big|\Omega(A,B\mid\Psi_0)\Big|, \tag{12.60}$$

where the minimum ranges over all admissible control gates for the asymmetric scheme (cf. Definition 2.124, eqs. (12.56) and (12.57), and Section 3.1.7). The candidates A and B are said to be *holosymmetrically orthogonal* on the probe ray $\mathbb{C}|\Psi_0\rangle$ if $\rho(A,B)=0$.

Lemma 12.39 (Tomographic control and overlap recovery)**.** *Let A,B be internally viable candidates with emergent evolutions $\mathbf{U}^{(A)},\mathbf{U}^{(B)}$ (Definition 12.28 and Postulate 12.1). For a pure probe $|\Psi_0\rangle$, the pair of asymmetric readout probabilities (12.56)–(12.57) as functions of the control parameters $(\alpha,\beta,\gamma,\delta)\in SU(2)\times SU(2)$ is tomographically sufficient to recover the full complex* holosymmetric overlap

$$\Omega(A,B\mid\Psi_0) := \langle\Psi_0|(\mathbf{U}^{(B)})^{\dagger}\mathbf{U}^{(A)}|\Psi_0\rangle \quad (\textit{cf. Definition 12.38}).$$

In particular, $\Omega(A,B\mid\Psi_0)=0$ iff there exist two phase–distinct control settings (real–Hadamard and quarter–phase–shifted; see below) for which $p_0^{(\mathrm{asym})}(A,B)=p_1^{(\mathrm{asym})}(A,B)$ simultaneously.

Proof. Write $x:=\mathbf{U}^{(A)}|\Psi_0\rangle$, $y:=\mathbf{U}^{(B)}|\Psi_0\rangle$ so that $\Omega=\langle y|x\rangle$ (Section 3.1.5). Expanding (12.56)–(12.57) with $|\alpha|^2+|\beta|^2=|\gamma|^2+|\delta|^2=1$ gives

$$p_0^{(\mathrm{asym})} = |\gamma\alpha|^2\,\|x\|^2+|\delta\beta|^2\,\|y\|^2 \;-\; 2\,\mathrm{Re}\Big(\gamma^*\alpha^*\,\delta^*\beta\,\langle x|y\rangle\Big), \tag{12.61}$$

$$p_1^{(\mathrm{asym})} = |\delta\alpha|^2\,\|x\|^2+|\gamma\beta|^2\,\|y\|^2 \;+\; 2\,\mathrm{Re}\Big(\delta^*\alpha^*\,\gamma^*\beta\,\langle x|y\rangle\Big). \tag{12.62}$$

Since A,B are viable, $\mathbf{U}^{(A)},\mathbf{U}^{(B)}$ are unitary on the support of $|\Psi_0\rangle$ (Postulate 12.1), hence $\|x\|=\|y\|=1$.

Step 1: calibration of single-branch norms. Using settings $(\alpha,\beta;\gamma,\delta)=(1,0;1,0)$ and $(0,1;0,1)$ in (12.61)–(12.62) yields $p_0^{(\mathrm{asym})}=1$ and $p_0^{(\mathrm{asym})}=1$, respectively, confirming $\|x\|=\|y\|=1$ (consistency with Definition 12.28 and Proposition 12.35).

Step 2: extraction of the real part. Choose an equal–weight, real–Hadamard setting

$$(\alpha,\beta;\gamma,\delta)=\Big(\tfrac{1}{\sqrt{2}},\tfrac{1}{\sqrt{2}};\;\tfrac{1}{\sqrt{2}},\tfrac{1}{\sqrt{2}}\Big).$$

Then $|\gamma\alpha|^2=|\delta\alpha|^2=|\delta\beta|^2=|\gamma\beta|^2=\tfrac14$ and $\gamma^*\alpha^*\,\delta^*\beta=\tfrac14$, giving

$$p_0^{(\mathrm{asym})}=\tfrac14\Big(2-2\,\mathrm{Re}\,\Omega\Big),\qquad p_1^{(\mathrm{asym})}=\tfrac14\Big(2+2\,\mathrm{Re}\,\Omega\Big),$$

whence

$$\mathrm{Re}\,\Omega \;=\; p_1^{(\mathrm{asym})}-p_0^{(\mathrm{asym})}. \tag{12.63}$$

(This is the asymmetric analogue of the symmetric visibility readout, cf. Remark 12.32 and Section 3.1.7.)

Step 3: extraction of the imaginary part. Use a quarter–phase–shifted, equal–weight setting

$$(\alpha, \beta; \gamma, \delta) = \Big(\tfrac{1}{\sqrt{2}}, \tfrac{i}{\sqrt{2}};\ \tfrac{1}{\sqrt{2}}, \tfrac{1}{\sqrt{2}}\Big),$$

so that $\gamma^* \alpha^*\, \delta^* \beta = \frac{i}{4}$. Then

$$p_0^{(\mathrm{asym})} = \tfrac{1}{4}\big(2 + 2\,\mathrm{Im}\,\Omega\big), \qquad p_1^{(\mathrm{asym})} = \tfrac{1}{4}\big(2 - 2\,\mathrm{Im}\,\Omega\big),$$

and hence

$$\mathrm{Im}\,\Omega \;=\; p_0^{(\mathrm{asym})} - p_1^{(\mathrm{asym})}. \tag{12.64}$$

Step 4: reconstruction and orthogonality. Equations (12.63)–(12.64) recover the full complex overlap $\Omega(A, B \mid \Psi_0)$ from two phase–distinct control settings, establishing tomographic sufficiency. Therefore $\Omega = 0$ iff both differences vanish for those two settings, i.e., iff there exist two such settings with $p_0^{(\mathrm{asym})} = p_1^{(\mathrm{asym})}$ simultaneously. In that case, by Definition 12.38, A and B are holosymmetrically orthogonal on the probe ray. □

Remark 12.40 (On minimizing visibility)**.** The raw minimization $\min_{U_c} \mathcal{V}_{\mathrm{asym}}(A, B)$ can be driven to 0 by phase–tuning even when $\Omega \neq 0$ (choose $|\gamma| = |\delta|$ so the norm terms in $p_0^{(\mathrm{asym})} - p_1^{(\mathrm{asym})}$ cancel and set the overall phase to make the real part vanish). Hence the discriminative content of Stage II rests on the *tomographic recovery* of Ω via fixed, linearly independent settings as above, not on a single minimization. Orthogonality should therefore be certified via $\Omega = 0$ (or via two-setting equalities), not via $\min_{U_c} \mathcal{V}_{\mathrm{asym}}$ alone (cf. Definition 12.38 and Section 3.1.7).

Remark 12.41 (Relational tomography as a physical principle)**.**

(i) **Fixed-order, law-level tomography.** The asymmetric holosymmetric interferometer provides a *process-tomographic* probe of universal histories with a fixed causal skeleton (no indefinite causal order). Its circuit is $\mathbf{U}_{\mathrm{holo}}^{(\mathrm{asym})} = \mathbf{U}_{\mathrm{read}}\, \mathbf{U}_{\mathrm{ctrl}}\, \mathbf{U}_{\mathrm{prep}}$ (Definition 2.124), which acts on the control $\mathcal{H}_c$ and dual registers $\mathcal{H}_{\mathrm{RR}}^{(2n)}$ defined in Chs. II–III.

(ii) **What is reconstructed.** For two *viable* candidates A, B (Stage I; Postulate 12.1 and Definition 12.33), the asymmetric probabilities (12.56)–(12.57) are functions of the control $SU(2)$ parameters (α, β) and (γ, δ) and determine the full complex *holosymmetric overlap* $\Omega(A, B \mid \Psi_0) = \langle \Psi_0 | (\mathbf{U}^{(B)})^\dagger \mathbf{U}^{(A)} | \Psi_0 \rangle$ (Definition 12.38 and Sections 3.1.5 and 3.1.7). Explicit two-setting reconstructions of $\mathrm{Re}\,\Omega$ and $\mathrm{Im}\,\Omega$ are given in Lemma 12.39.

(iii) **Operational meaning.** Tomographic control on the *control qubit alone* suffices to compare laws: varying $U_c^{\mathrm{prep}}, U_c^{\mathrm{read}} \in SU(2)$ tunes interference between the two lifted path-laws while the register dynamics remain fixed. The relational figure of merit is the *overlap* Ω rather than a single visibility value; $\mathcal{V}_{\mathrm{asym}}$ is a derived readout (Section 3.1.7).

(iv) **Gauge covariance and invariance.** Because $\mathcal{C}(\cdot)$ and the induced probabilities are invariant under fibrewise conjugations (Proposition 12.35), the recovered overlap is a gauge-invariant diagnostic of the *internal* dynamics (Definitions 12.28 and 12.30).

(v) **Relation to the symmetric case (Stage I).** Setting $A = B$ collapses the asymmetric scheme to the Stage I self-coherence test: $\Omega(A, A \mid \Psi_0) = \langle\Psi_0|\mathbf{U}^{(A)}_{\text{unitarity}}|\Psi_0\rangle = \mathcal{C}(A)$, and the symmetric visibility equals this expectation (Remark 12.32, Definition 12.30, and Postulate 12.1).

(vi) **Relational orthogonality and selection.** A pair A, B is *holosymmetrically orthogonal* on the probe ray if and only if $\Omega(A, B \mid \Psi_0) = 0$ (Definition 12.38). On a subspace $\mathcal{K}$, orthogonality is certified state-independently by an informationally complete family of probes (IC) on $\mathcal{K}$ (Definition 12.33 and Remark 12.34); this upgrades the comparison of laws to a tomographic separation over $\mathcal{K}$.

(vii) **Register–lifted consistency.** All statements admit an equivalent register–lifted formulation on $\mathcal{H}_c \otimes \mathcal{H}^{(2n)}_{\text{RR}}$ using $\mathbf{U}^{(\text{asym})}_{\text{holo}}$ built from the controlled operator $\mathbf{U}_{\text{ctrl}}$ and the path operators $\hat{\mathbf{U}}^{A_G B_N}_{\text{RR}}, \hat{\mathbf{U}}^{A_N B_G}_{\text{RR}}$ (Definitions 2.108, 2.114 and 12.31).

12.2.4 The Holosymmetric Selection Principle (HSP)

Theorem 12.42 (Holosymmetric Selection Principle)**.** *Fix the universal internal group* $G_{\text{int}} = \text{Aut}(\text{End}(\mathcal{H}_{\text{em}}))$ *acting via a realized weave configuration-compatible split connection* $\nabla_\mu = \nabla^{(\text{LC})}_\mu + i\,\text{ad}_{\mathcal{A}_\mu}$ *on the operator bundle (eq. (12.99) and Definitions 6.192 and 12.1). Among all local, compact, Tantrakyma-invariant, chiral candidates* $G_A \subset G_{\text{int}}$ *that obey the diagnostic realized reference configuration/plateau hygiene (Propositions 12.8 and 12.12, Lemma 12.2, and Section 6.12.4), the realized gauge symmetry* G_{SM} *is characterized by:*

Self-consistency (Stage I)

$\mathcal{C}(A) = 1$ *(perfect symmetric visibility)* iff A *is unitary (on the certified domain) and anomaly-free:*

$$\mathcal{C}(A) = \left|\langle\Psi_0|\mathbf{U}^{(A)}_{\text{unitarity}}|\Psi_0\rangle\right| = 1 \quad \Longleftrightarrow \quad \mathbf{U}^{(A)}_{\text{unitarity}} = \mathbb{I} \text{ on } \text{supp}(\Psi_0) \text{ and } \mathsf{A}_\bullet = 0,$$

with $\mathbf{U}^{(A)}_{\text{unitarity}} = (\mathbf{U}^{(A)})^\dagger \mathbf{U}^{(A)}$ *(Definition 12.28),* $\mathcal{C}$ *as in eqs. (12.49) to (12.51), and anomaly meters* $\mathsf{A}_\bullet$ *from Definitions 12.20 and 12.24; failure is recorded by the viability defect Definition 12.33. Operationally, the symmetric interferometer readout reproduces* $\mathcal{C}(A)$ *(Theorem 3.4 and Remark 12.36).*

Relational orthogonality (Stage II)

For every other viable $B \neq A$*, asymmetric tomography (with*

$$\mathbf{U}^{(\text{asym})}_{holo} = \mathbf{U}_{\text{read}} \mathbf{U}_{\text{ctrl}}[A] \mathbf{U}_{\text{prep}}$$

Definitions 2.114 and 2.124) attains minimal overlap with A*:*

$$\begin{aligned} \rho(A, B) \;=\; 0, \qquad & \textit{equivalently} \\ & \exists \textit{ prep/readout such that } \mathcal{V}_{\text{asym}}(A, B) \to 0 \\ & \textit{while } \mathcal{V}_{\text{asym}}(A, A) \textit{ is maximal,} \end{aligned}$$

where the (physical) holosymmetric overlap is $\Omega_{\text{phys}}(A, B \mid \Psi_0) = \langle\Psi_0|U^{(A)\dagger}_{\text{phys}} U^{(B)}_{\text{phys}}|\Psi_0\rangle$ *(Definition 12.69 and eq. (12.83)) reconstructed by the asymmetric interferometer*

(Section 3.1.5 and Definitions 3.13 and 3.15). Informational completeness on the BRST physical sector is enforced by the channel-resolved meters on certified plateaux (Definition 6.275, Lemma 12.43, Corollary 12.52, and Section 6.12.4).

Together, (1)–(2) constitute a proof by uniqueness*: within the constrained class (local, compact, Tantrakyma-invariant, chiral/BRST-consistent), there is a unique A satisfying both criteria; empirically this is G_{SM} (Proposition 12.26, Corollary 12.27, and eq. (12.46)).*

Proof. Diagnostic realized reference configuration and constraints. Work on the certified plateau $\mathfrak{P}$ with the realized weave configuration-compatible connection (Definition 12.1). Compact holonomy holds on each finite internal fibre, ensuring unitary representation theory and Peter–Weyl control (Proposition 12.8). All meters and Ward/KL audits descend to the BRST physical space $\mathcal{H}_{\mathrm{phys}}$ and are Nielsen-flat (Lemma 12.2 and Proposition 12.12). Tantrakyma dynamics is expressed covariantly with the same split ∇_μ and improved currents (eq. (12.37) and Definition 12.4).

Stage I (self-consistency). Define $\mathbf{U}^{(A)}_{\mathrm{unitarity}} = (\mathbf{U}^{(A)})^\dagger \mathbf{U}^{(A)}$ and the self-coherence functional $\mathcal{C}(A)$ (Definition 12.28 and eqs. (12.49) and (12.50)). In the register-lifted interferometric form, $\mathcal{C}_{\mathrm{holo}}(A \mid \rho_{\mathrm{cRR},0}; \star) = \left|\mathrm{Tr}(\rho_{\mathrm{cRR},0}\, \mathbf{U}^{(A);\mathrm{holo},(\star)}_{\mathrm{unitarity}})\right|$ with $\mathbf{U}^{(\star)}_{\mathrm{holo}}$ built from $\mathbf{U}_{\mathrm{ctrl}}$ (Definitions 2.114 and 12.31). By Definition 12.28, $\mathcal{C}(A) = 1$ iff $\mathbf{U}^{(A)}_{\mathrm{unitarity}} = \mathbb{I}$ on $\mathrm{supp}(\Psi_0)$. Moreover, viability on $\mathfrak{P}$ requires the anomaly meters vanish and the global $SU(2)$ Witten test to pass (Definitions 12.20 and 12.24 and Proposition 12.26). Conversely, any nonzero meter in eq. (12.46) obstructs BRST nilpotency and thus Stage I (Corollary 12.27). Operationally, the symmetric interferometer readout reproduces $\mathcal{C}(A)$ (normalization as in Remark 12.36), so Stage I is both necessary and sufficient for perfect symmetric visibility.

Stage II (relational orthogonality). Take two *viable* A, B passing Stage I. Run the asymmetric (character-weighted) interferometer on $\mathcal{H}_{\mathrm{phys}} \otimes \mathcal{H}_{\mathrm{phys}}$ with compressed arm blocks $U^{(A)}_{\mathrm{phys}}, U^{(B)}_{\mathrm{phys}}$; the outcome probabilities reconstruct the physical overlap $\Omega_{\mathrm{phys}}(A, B \mid \Psi_0) = \langle\Psi_0| U^{(A)\dagger}_{\mathrm{phys}} U^{(B)}_{\mathrm{phys}} |\Psi_0\rangle$ exactly as in Definition 12.69 and eq. (12.83). By informational completeness of the dual-register tomography and the availability of BRST-closed primitives, there exist preparation/readout choices driving the cross-visibility to zero while keeping $\mathcal{V}_{\mathrm{asym}}(A, A)$ maximal (Lemma 12.43, Corollary 12.52, Section 3.1.5, and Definition 2.124). Equivalently, $\rho(A, B) = 0$ in the sense of holosymmetric overlap orthogonality (Definition 12.38).

Uniqueness. Within the constrained class (local/realized weave configuration-compatible, compact holonomy, Tantrakyma-invariant, chiral, BRST-consistent), the anomaly audit fixes the chiral content and hypercharges, yielding the unique minimal anomaly-free sector (Proposition 12.26, eq. (12.46), and Corollary 12.27). Stage II then promotes this from a static audit to a dynamical selection: if two distinct candidates $A \neq B$ both satisfied (1)–(2), they would be mutually orthogonal under all asymmetric meters yet each have maximal self-visibility—a contradiction unless they are physically the same law up to gauge/diagnostic realized reference configuration equivalence (*proof by uniqueness*, Section 12.2; cf. Section 12.1.1). Hence a unique fixed point remains; empirically G_{SM}. □

12.2.5 Structural Interlocks with the Foundations

Born-valued semantics and meta-level truth. The Holothesis inherits the Born semantics of the Fourfold, so that "truth" of a meta-claim about laws is an *effect valuation* determined by the holosymmetric circuit (Definitions 1.4 and 1.18).

Universal histories and the relational action. The objects compared interferometrically are universal histories governed by the Total Relational Action $S_{\text{total}}[\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}]$; their interference is constrained by the universal (Wheeler–DeWitt–type) relation and its holocosmic extension (eq. (17.54)). The HSP is thus a *selection by interference* within the space of histories permitted by these constraints.

Continuum–entanglement control. In correlator-sparse regimes the dual-register scaffold is minimal and symmetry-respecting, and holosymmetric dominance identifies the bright fringe as the realized meta-dynamics, providing the physical reading of "maximal relational coherence."

Motivational Aside: Superposition and the Cosmic Interferometer For orientation only, recall that Holosymmetry compares *laws* by coherent branching and recombination of dual path–histories, with a qubit control readout providing an interference visibility (Section 12.2). Concretely, one may imagine two admissible evolutions acting on the same preparation,

Path A G_A–dynamics on the Tantrakyma sector,

Path B G_B–dynamics on the Tantrakyma sector,

whose relational overlap is tomographically recovered by the asymmetric interferometer (Sections 3.1.5 and 3.1.7). This picture is *only* mnemonic: all formal content is fixed by the Unitarity Operator and the visibility functionals already defined in Section 12.2.

Sufficiency results for the Holosymmetric Selection Principle

In this subsection we supply two results that close the remaining sufficiency gap in the HSP narrative. The first constructs a symmetry–respecting, dual–register probe ensemble that is informationally complete for the *process–overlap* functionals appearing in the asymmetric interferometer (Section 12.2.3, Definitions 2.114, 2.124 and 12.69, and eqs. (12.56), (12.57) and (12.83)). The second reduces the holosymmetric overlap to group–theoretic data and proves relational orthogonality whenever the induced actions are inequivalent (Definition 12.38, Lemma 12.39, and Propositions 12.8 and 12.10).

Lemma 12.43 (Dual–register informational completeness for process overlaps)**.** *Let $\mathcal{U}$ denote the linear span (over $\mathbb{C}$) of operators of the form $(\mathbf{U}^{(B)})^\dagger \mathbf{U}^{(A)}$ with A, B ranging over the admissible candidate class (local/compact, Tantrakyma–symmetric, chiral, anomaly–free; Section 12.1.1, Definitions 12.1 and 12.20, Proposition 12.8, and Corollary 12.27). Fix a finite–dimensional probe subspace $\mathcal{H}_{\text{probe}} \subset \mathcal{H}_{\text{RR}}^{(2n)}$ that contains the support of all states used*

by the holosymmetric circuit on the two registers (Section 3.1.5). Then there exists a finite ensemble

$$\mathcal{S}_{\mathrm{IC}}^{(2n)} = \left\{(\rho_j, U_{c,j}^{\mathrm{prep}}, U_{c,j}^{\mathrm{read}})\right\}_{j=1}^{N}$$

with each ρ_j a density operator on $\mathcal{H}_{\mathrm{probe}}$ and each $U_{c,j}^{\mathrm{prep/read}}$ an admissible control gate (Definitions 2.108 and 2.114), such that the map

$$\Phi : \mathcal{U} \longrightarrow \mathbb{C}^N, \qquad X \longmapsto \left(p_{0,j}^{(\mathrm{asym})} - p_{1,j}^{(\mathrm{asym})} \right)_{j=1}^{N}$$

is injective on $\mathcal{U}|_{\mathcal{H}_{\mathrm{probe}}}$. Equivalently, the data $\{p_{0,j}^{(\mathrm{asym})}, p_{1,j}^{(\mathrm{asym})}\}$ measured on $\mathcal{S}_{\mathrm{IC}}^{(2n)}$ separate process–overlaps on $\mathcal{H}_{\mathrm{probe}}$ (Definition 12.69 and eq. (12.83)).

Proof. By Section 3.1.5 and Definition 2.124, for fixed $(\rho, U_c^{\mathrm{prep}}, U_c^{\mathrm{read}})$ one has the arm–resolved probabilities $p_0^{(\mathrm{asym})}, p_1^{(\mathrm{asym})}$ (see eqs. (12.56) and (12.57)) with control parameters $(\alpha, \beta, \gamma, \delta)$ from the admissible gate parametrization (Definition 2.114). Taking the difference and using the algebraic cancellation in the asymmetric path expansion (Definitions 3.13 and 3.15 and Lemma 12.39), the residual dependence becomes *linear* in the process–overlap

$$\Omega(A, B \mid \rho) := \mathrm{Tr}\Big(\rho\, (\mathbf{U}^{(B)})^{\dagger} \mathbf{U}^{(A)}\Big)$$

and its complex conjugate, with coefficients that are polynomial functions of $(\alpha, \beta, \gamma, \delta)$ (Definition 2.114). Hence there exist four admissible control settings that isolate $\mathrm{Re}\,\Omega$ and $\mathrm{Im}\,\Omega$ linearly from $p_0^{(\mathrm{asym})} - p_1^{(\mathrm{asym})}$ (extraction via Lemma 12.39).

Let $\{X_\ell\}_{\ell=1}^{M}$ be any basis of $\mathcal{U}|_{\mathcal{H}_{\mathrm{probe}}}$ (*cf.* Peter–Weyl–type spanning for compact holonomy; Propositions 12.8 and 12.10). Choose a finite set of density operators $\{\rho_j\}_{j=1}^{M'}$ on $\mathcal{H}_{\mathrm{probe}}$ that is informationally complete for linear statistics on $\mathcal{U}$—i.e. the matrix $\Big(\mathrm{Tr}(\rho_j X_\ell)\Big)_{j,\ell}$ has full column rank.[1] Combine each ρ_j with the four control settings that recover Re/Im as above and collect all triples into $\mathcal{S}_{\mathrm{IC}}^{(2n)}$. The associated measurement matrix for the linear map $X \mapsto \big(p_{0,j}^{(\mathrm{asym})} - p_{1,j}^{(\mathrm{asym})}\big)_j$ then contains blockwise the real/imaginary copies of $\Big(\mathrm{Tr}(\rho_j X_\ell)\Big)$ and thus has full column rank (by Lemma 12.39). Therefore Φ is injective on $\mathcal{U}|_{\mathcal{H}_{\mathrm{probe}}}$, and the ensemble $\mathcal{S}_{\mathrm{IC}}^{(2n)}$ separates process–overlaps (Definition 12.69 and eq. (12.83)). □

Definition 12.44 (Ensemble–relaxed orthogonality index)**.** Given the dual–register IC ensemble $\mathcal{S}_{\mathrm{IC}}^{(2n)}$ from Lemma 12.43 within Section 12.2.5, define

$$\rho_{\mathrm{ens}}(A, B) := \inf_{\substack{(\rho, U_c^{\mathrm{prep}}, U_c^{\mathrm{read}}) \\ \in \mathrm{co}\, \mathcal{S}_{\mathrm{IC}}^{(2n)}}} \Big|\Omega(A, B \mid \rho)\Big|,$$

where $\Omega(A, B \mid \rho) = \mathrm{Tr}\Big(\rho\, (\mathbf{U}^{(B)})^{\dagger} \mathbf{U}^{(A)}\Big)$ is the (physical) process–overlap (Definition 12.69 and eq. (12.83)) read out by the asymmetric interferometer (Section 3.1.5, Definition 2.124, and eqs. (12.56) and (12.57)), and $\mathrm{co}\,\mathcal{S}$ denotes the convex hull of the probe triples (classical mixtures of probes on $\mathcal{H}_{\mathrm{RR}}^{(2n)}$; *cf.* Section 3.1.5 and Definition 2.114). Clearly $\rho_{\mathrm{ens}}(A, B) \leq \rho(A, B)$ (with ρ as in Definition 12.38), with equality when a single pure probe in $\mathcal{S}_{\mathrm{IC}}^{(2n)}$ already attains the infimum.

[1] Any finite–dimensional IC–POVM will do; you can take its outcome states restricted to $\mathcal{H}_{\mathrm{probe}}$.

Remark 12.45 (Operational meaning and basic properties)**. Best achievable cross–visibility under classical randomization**

Because $\Omega(A, B \mid \rho)$ is linear in ρ (eq. (12.83)), taking the infimum over the convex hull

$$\operatorname{co} \mathcal{S}_{\mathrm{IC}}^{(2n)}$$

computes the smallest cross–visibility accessible by classical randomization over admissible probes (and admissible control gates; Definition 2.114) while remaining on the BRST physical sector (Section 12.3). Hence $\rho_{\mathrm{ens}}(A, B)$ is the operational "floor" for cross–visibility using the given IC ensemble.

Bounds and equality case

By construction $0 \le \rho_{\mathrm{ens}}(A, B) \le \rho(A, B)$, where $\rho(A, B)$ is the overlap index used in Stage II (Definition 12.38 and Theorem 12.42). Equality $\rho_{\mathrm{ens}}(A, B) = \rho(A, B)$ holds whenever a single (pure) probe triple $(\rho, U_c^{\mathrm{prep}}, U_c^{\mathrm{read}}) \in \mathcal{S}_{\mathrm{IC}}^{(2n)}$ attains the infimum, which occurs in particular when the optimal witness for Ω already lies in $\mathcal{S}_{\mathrm{IC}}^{(2n)}$ (Lemma 12.43).

Monotonicity under ensemble refinement

If $\mathcal{S}_{\mathrm{IC}}^{(2n)} \subseteq \widetilde{\mathcal{S}}_{\mathrm{IC}}^{(2n)}$ then $\rho_{\mathrm{ens}}^{\widetilde{\mathcal{S}}}(A, B) \le \rho_{\mathrm{ens}}^{\mathcal{S}}(A, B)$ by convexity/linearity of Ω (eq. (12.83)). Thus adding probes or control settings can only improve (lower) the achievable cross–visibility floor.

Sufficiency for Stage II

If $\rho_{\mathrm{ens}}(A, B) = 0$ then, by definition, there exist admissible probe mixtures and controls that drive the asymmetric visibility to zero, hence $\rho(A, B) = 0$ and *relational orthogonality* holds (Definition 12.38 and Theorem 12.42). Together with Lemma 12.43 (informational completeness of $\mathcal{S}_{\mathrm{IC}}^{(2n)}$), this supplies the Stage-II sufficiency direction used in Section 12.2.5.

Gauge/BRST and Nielsen stability

Since overlaps are evaluated on the BRST physical space and meters descend cohomologically (Section 12.3 and Proposition 12.12), ρ_{ens} is invariant under admissible gauge choices and Nielsen reparametrizations on certified plateaux (Section 6.12.4).

Proposition 12.46 (Relational character orthogonality)**.** *Fix two* viable *candidates A, B in the constrained class (local, compact, Tantrakyma–invariant, chiral/BRST–consistent; Sections 12.1.1 and 12.3 and Propositions 12.8 and 12.12). Let K denote the compact subgroup of G_{int} (eq. (12.99)) generated by the internal gauge support of the probe window (Section 3.1.5 and Definition 12.4). Suppose the restricted evolutions on $\mathcal{H}_{\mathrm{probe}}$ are unitarily equivalent to* group representations

$$\mathbf{U}^{(A)}(g) = \pi_A(g), \qquad \mathbf{U}^{(B)}(g) = \pi_B(g), \qquad g \in K,$$

with π_A, π_B finite–dimensional unitary reps (the time–ordered exponential along the window implements g; Definitions 2.108, 2.124 and 3.15). If π_A and π_B are inequivalent *as K–representations, then*

$$\inf_{g \in K} \rho_{\mathrm{ens}}\big(A, B; g\big) = 0,$$

i.e. there exists (in the infimum sense) an admissible probe ensemble and a pulse choice g for which the holosymmetric overlap can be made arbitrarily small; in particular, Stage–II relational orthogonality holds (Definition 12.38 and Theorem 12.42).

Proof. For $g \in K$ set $V(g) := (\mathbf{U}^{(B)}(g))^\dagger \mathbf{U}^{(A)}(g) = \pi_B(g)^\dagger \pi_A(g)$. Let ρ_0 be any density operator on $\mathcal{H}_{\mathrm{probe}}$. Consider the class–function average

$$\overline{\Omega}(A, B \mid \rho_0) := \int_K \mathrm{d}\mu(g)\ \mathrm{Tr}\Big(\rho_0\, V(g)\Big),$$

where μ is Haar measure on K (normalized as in Proposition 12.10). Decompose π_A and π_B into irreducibles:

$$\pi_A \cong \bigoplus_\alpha m_\alpha^A\, \alpha, \qquad \pi_B \cong \bigoplus_\beta m_\beta^B\, \beta,$$

with multiplicities m_α^A, m_β^B (*Peter–Weyl* control for compact holonomy; Propositions 12.8 and 12.10). By Schur orthogonality of matrix elements on compact groups (encoded in Proposition 12.10), the operator integral $\int_K \pi_B(g)^\dagger \pi_A(g)\, \mathrm{d}\mu(g)$ is the intertwiner from π_B to π_A; it vanishes on inequivalent components and reduces to a multiple of the identity on common irreps. Since π_A and π_B are assumed *inequivalent*, the intertwiner is the zero operator; hence $\overline{\Omega}(A, B \mid \rho_0) = 0$ for every ρ_0.

Define $f_{\rho_0}(g) := \mathrm{Tr}\Big(\rho_0\, V(g)\Big)$, which is continuous in g by continuity of matrix elements (Proposition 12.10). Then $\int_K f_{\rho_0}(g)\, \mathrm{d}\mu(g) = 0$ for all ρ_0. Approximating the Haar integral by finite convex combinations (standard on compact groups; see the approximate-identity discussion in Proposition 12.10), we obtain $g_1, \dots, g_r \in K$ and weights $w_j \geq 0$, $\sum_j w_j = 1$, with $\sum_j w_j f_{\rho_0}(g_j) = 0$.

Now use: (i) Ω is linear in ρ (Definition 12.69 and eq. (12.83)); (ii) classical randomization over probes and admissible controls/pulses is permitted within the dual–register circuit (Definition 2.114 and Section 3.1.5); and (iii) mixtures from Lemma 12.43 are realizable inside the convex hull $\mathrm{co}\,\mathcal{S}_{\mathrm{IC}}^{(2n)}$ (Definition 12.44). Thus, for each $\varepsilon > 0$ we can choose $j_\star$ and an ensemble on $\mathcal{H}_{\mathrm{probe}}$ such that

$$\rho_{\mathrm{ens}}(A, B; g_{j_\star}) = \inf_{(\rho, U_c^{\mathrm{prep/read}}) \in \mathrm{co}\,\mathcal{S}_{\mathrm{IC}}^{(2n)}} \Big| \mathrm{Tr}(\rho\, V(g_{j_\star})) \Big| < \varepsilon,$$

hence $\inf_{g \in K} \rho_{\mathrm{ens}}(A, B; g) = 0$. By Definition 12.38 and Theorem 12.42, this suffices for Stage–II relational orthogonality. □

Remark 12.47 (Attainment of the infimum)**.** Since K is compact and $g \mapsto V(g)$ is continuous (Proposition 12.10), the map $g \mapsto \rho_{\mathrm{ens}}(A, B; g)$ is lower semicontinuous as an infimum of continuous affine functionals of ρ (eq. (12.83) and Definition 12.44). Consequently the infimum over K is a minimum whenever $\mathcal{S}_{\mathrm{IC}}^{(2n)}$ is closed under limits of classical mixtures (true for the constructions in Lemma 12.43); in that case there exists $g_\star \in K$ and an admissible ensemble with $\rho_{\mathrm{ens}}(A, B; g_\star) = 0$.

Corollary 12.48 (Character–orthogonality filter)**.** *Under the hypotheses of Proposition 12.46, let $\chi_A(g) := \mathrm{Tr}\ \pi_A(g)$ and $\chi_B(g) := \mathrm{Tr}\ \pi_B(g)$ denote the characters of the restricted unitary representations on K. If the Peter–Weyl inner product of class functions vanishes,*

$$\langle \chi_B, \chi_A \rangle_K := \int_K \chi_B(g)^*\, \chi_A(g)\, \mathrm{d}\mu(g) = 0 \quad (\textit{Haar-normalized as in Proposition 12.10}),$$

then

$$\inf_{g\in K}\ \rho_{\mathrm{ens}}(A,B;g)=0,$$

hence $\rho(A,B)=0$ *and Stage–II relational orthogonality holds (Definitions 12.38 and 12.44 and Theorem 12.42).*

Proof. Take the maximally mixed probe on the common support, $\rho_{\mathrm{mm}}:=\frac{\mathbb{I}_{\mathrm{probe}}}{\dim\mathcal{H}_{\mathrm{probe}}}$, admissible within the dual–register circuit (Section 3.1.5). Then

$$\int_K \mathrm{Tr}\Big(\rho_{\mathrm{mm}}\,\pi_B(g)^\dagger\pi_A(g)\Big)\,\mathrm{d}\mu(g)=\frac{1}{\dim\mathcal{H}_{\mathrm{probe}}}\int_K \mathrm{Tr}\Big(\pi_B(g)^\dagger\pi_A(g)\Big)\,\mathrm{d}\mu(g)=\frac{\langle\chi_B,\chi_A\rangle_K}{\dim\mathcal{H}_{\mathrm{probe}}}=0,$$

using $\mathrm{Tr}(\pi_B^\dagger\pi_A)=\chi_B^*\chi_A$ and the character inner product from Proposition 12.10. Approximate the Haar integral by finite convex combinations on K (standard on compact groups; see the approximate-identity discussion in Proposition 12.10) to get $g_1,\dots,g_r\in K$ and weights $w_j\ge 0$, $\sum_j w_j=1$, with $\sum_j w_j\ \mathrm{Tr}\Big(\rho_{\mathrm{mm}}\,\pi_B(g_j)^\dagger\pi_A(g_j)\Big)=0$. By linearity of the overlap in ρ and admissibility of classical randomization over probes/controls (Definitions 2.114 and 12.69 and eq. (12.83)), this convex program is realizable within $\mathrm{co}\,\mathcal{S}_{\mathrm{IC}}^{(2n)}$ (Lemma 12.43 and Definition 12.44), whence $\inf_{g\in K}\rho_{\mathrm{ens}}(A,B;g)=0$. Finally, Definition 12.38 and Theorem 12.42 imply $\rho(A,B)=0$ and Stage–II orthogonality. □

Remark 12.49 (When the filter does *not* trigger)**.** If $\pi_A\simeq\pi_B$ as K–representations, then $\langle\chi_B,\chi_A\rangle_K>0$ equals the total multiplicity of common irreps (Proposition 12.10), so the character–orthogonality hypothesis fails and Corollary 12.48 does not apply—consistent with the fact that Proposition 12.46 requires inequivalence to deduce Stage–II orthogonality.

Corollary 12.50 (Stage–II sufficiency for inequivalent actions)**.** *Under the hypotheses of Proposition 12.46, if* $A\neq B$ *as* K*–representations, then there exists an admissible probe ensemble (and control/readout choice) for which the asymmetric interferometer visibility satisfies* $\mathcal{V}_{\mathrm{asym}}(A,B)=0$ *and hence* $\rho_{\mathrm{ens}}(A,B)=0$.

Proof. By Proposition 12.46 (inequivalent restricted actions on K), there exists a pulse $g_\star\in K$ and an admissible (classically randomized) probe ensemble inside $\mathrm{co}\,\mathcal{S}_{\mathrm{IC}}^{(2n)}$ such that the process–overlap

$$\Omega\Big(A,B\mid\rho;g_\star\Big)=\mathrm{Tr}\Big(\rho\,(\mathbf{U}^{(B)}(g_\star))^\dagger\,\mathbf{U}^{(A)}(g_\star)\Big)$$

vanishes. Equivalently, $\rho_{\mathrm{ens}}(A,B;g_\star)=0$ by Definition 12.44. (If desired, attainment of the infimum over K follows from compactness and lower semicontinuity as noted in Remark 12.47.)

Run the asymmetric two–register interferometer (Section 3.1.5 and Definition 2.124) with admissible control gates parametrized by $(\alpha,\beta,\gamma,\delta)$ (Definition 2.114). For any fixed $(\rho,U_c^{\mathrm{prep}},U_c^{\mathrm{read}})$ the arm probabilities satisfy $p_0^{(\mathrm{asym})},p_1^{(\mathrm{asym})}$ as in eqs. (12.56) and (12.57), and their difference $p_0^{(\mathrm{asym})}-p_1^{(\mathrm{asym})}$ depends *linearly* on $\Omega(A,B\mid\rho;g_\star)$ and its complex conjugate (algebraic cancellation in the path expansion; Lemma 12.39). Choose any quadrature control that isolates $\mathrm{Re}\,\Omega$ or $\mathrm{Im}\,\Omega$ (see Definition 2.114 and Lemma 12.39). Since $\Omega(A,B\mid\rho;g_\star)=0$, one obtains

$$p_0^{(\mathrm{asym})}-p_1^{(\mathrm{asym})}=0. \tag{12.65}$$

Consequently, by the visibility definition in Section 3.1.5 and eqs. (12.56) and (12.57),

$$\mathcal{V}_{\mathrm{asym}}(A,B)=0. \tag{12.66}$$

Since the same admissible ensemble realizes $\Omega=0$, Definition 12.44 yields $\rho_{\mathrm{ens}}(A,B)=0$. This is precisely the Stage–II relational orthogonality condition referenced in Theorem 12.42. □

Remark 12.51 (From ensembles back to single–probe orthogonality)**.** The proof of Corollary 12.50 achieves $\Omega(A,B\mid\rho;g)=0$ by convex randomization over a finite set of pulses $g\in K$ and probe states $\rho\in\operatorname{co}\mathcal{S}_{\mathrm{IC}}^{(2n)}$ (cf. Definition 12.44 and Lemma 12.43). In many physically relevant instances—e.g. when the restricted actions π_A,π_B contain inequivalent irreducibles whose characters exhibit opposite phases along a one–parameter subgroup $g(t)=\exp(tX)\subset K$—a *single* pure probe Ψ and a *single* pulse $g(t_\star)$ already yield $\Omega(A,B\mid\Psi;g(t_\star))=0$. Indeed, $t\mapsto\Omega(A,B\mid\Psi;g(t))=\langle\Psi|\pi_B(g(t))^\dagger\pi_A(g(t))|\Psi\rangle$ is continuous (matrix–coefficient continuity on compact K; Proposition 12.10), and a change of argument along $g(t)$ forces a zero by the intermediate value principle. Thus the ensemble form ρ_{ens} guarantees *existence* of cancellation (Definition 12.44 and Proposition 12.46), while a single–probe realization is often available by continuity on K.

Independent of such phase arguments, convex–geometry bounds give finite (and small) support sizes for the required randomizations. First, any probe density operator on a D–dimensional subspace admits a convex decomposition into at most D pure states by the spectral theorem (see, e.g., [287]). More generally, Carathéodory's theorem on convex hulls ensures that any point in a convex subset of a real vector space of dimension m can be expressed as a convex combination of at most $m+1$ extreme points. Applied to the real vector space of Hermitian trace–one operators (dimension D^2-1), this yields a conservative upper bound of D^2 pure–state components for the probe mixture, with an additional finite support for the pulse randomization on K (the latter arising from approximating the Haar average by convex combinations on compact groups; cf. Proposition 12.10). Consequently, while the existence proof in Corollary 12.50 uses convex mixtures, the orthogonality can often be realized by a single pure probe and a single pulse, and in all cases by a finite ensemble with support bounded by these geometric considerations.[2]

Consequent uniqueness contour (closed). Work throughout on a certified plateau with rig–compatible connection and compact holonomy (Section 6.12.4, Definition 12.1, and Proposition 12.8), so that all meters descend to the BRST physical space and Ward/Källén–Lehmann hygiene holds (Section 12.3 and Proposition 12.12). Stage–I viability asserts that a candidate law A is admitted iff the self–coherence functional saturates, $\mathcal{C}(A)=1$, equivalently the viability defect vanishes and anomaly meters are zero (Definition 12.33, Proposition 12.35, and Theorem 12.42). In this regime the symmetric interferometer readout matches $\mathcal{C}$ and is gauge/Nielsen stable on the plateau (Section 3.1.5, Definition 2.124, and Theorem 12.42).

For sufficiency at Stage–II, assume A,B are distinct *viable* laws within the admissible class (local, compact, Tantrakyma–symmetric, chiral; Section 12.1.1). By compact holonomy there is a compact support subgroup $K\subset G_{\mathrm{int}}$ for the probe window (Proposition 12.8). If the restricted actions were equivalent on every such K, the two laws would coincide up to diagnostic

[2]For Carathéodory's theorem, see [555] or the original source [556].

realized reference configuration equivalence; hence distinct viable laws must be inequivalent on some compact window K. Peter–Weyl/Schur orthogonality (as localized in Proposition 12.10) then promotes inequivalence to relational character orthogonality: the averaged intertwiner $\int_K \pi_B(g)^\dagger \pi_A(g)\, d\mu(g)$ vanishes, and so does the character inner product when applicable (Proposition 12.46 and Corollary 12.48). By Lemma 12.43 and Definition 12.44 this implies the existence of an admissible (possibly randomized) probe ensemble and control/readout choice with zero process–overlap for some pulse $g_\star$, whence $\rho_{\mathrm{ens}}(A,B)=0$; attainment can be taken on K by compactness, and in many instances a *single* pure probe and pulse suffice (Remarks 12.47 and 12.51). Translating overlap to arm probabilities via the asymmetric interferometer formulas (eqs. (12.56), (12.57) and (12.83), Definitions 2.114 and 12.69, and Lemma 12.39) yields $\mathcal{V}_{\mathrm{asym}}(A,B)=0$, the Stage–II orthogonality recorded in Corollary 12.50.

Consequently, within the admissible class, any two *distinct* viable laws become holosymmetrically orthogonal on some compact window K, i.e. $\rho_{\mathrm{ens}}(A,B)=0$. Therefore the pair of conditions

$$\Big(\ \mathcal{C}(A)=1,\quad \rho_{\mathrm{ens}}(A,B)=0\ \ \forall\, B\neq A\ \Big)$$

has at most one solution on a given plateau. This *uniqueness contour* closes the sufficiency side of the HSP proof–by–uniqueness (Theorem 12.42); empirically the survivor is the Standard Model gauge action G_{SM}.

Completing Stage II and the uniqueness bridge

Corollary 12.52 (General completeness of probes for Stage II)**.** *For any two viable, Tantrakyma–compatible laws $A\neq B$ acting on the same probe window, there exists an admissible family of probes $(\rho, U_c^{\mathrm{prep}}, U_c^{\mathrm{read}})$ within the dual–register scaffold such that the holosymmetric overlap obeys $\Omega(A,B\mid\rho)=0$. Consequently, by the asymmetric interferometer readout, there is a setting with $\mathcal{V}_{\mathrm{asym}}(A,B)=0$.*

Proof. Setup on a certified plateau. Work on a certified plateau with rig–compatible connection and compact holonomy so that meters descend cohomologically to the BRST physical sector and Ward/Källén–Lehmann hygiene holds (Sections 6.12.4 and 12.3, Definition 12.1, and Propositions 12.8 and 12.12). On the two–register scaffold, process overlaps are read out by the asymmetric interferometer (Section 3.1.5, Definitions 2.124 and 12.69, and eqs. (12.56), (12.57) and (12.83)).

Informational completeness. By Lemma 12.43, the symmetry–respecting ensemble $\mathcal{S}_{\mathrm{IC}}^{(2n)}$ is informationally complete for process–overlap functionals on the finite probe subspace $\mathcal{H}_{\mathrm{probe}}$: for every X in the span $\mathcal{U}|_{\mathcal{H}_{\mathrm{probe}}}$, the differences $p_0^{(\mathrm{asym})}-p_1^{(\mathrm{asym})}$ computed on a finite set of control settings separate X linearly (via the quadrature choices guaranteed by Lemma 12.39 and implemented by Definition 2.114).

Reduction to group–theoretic data. Let $K\subset G_{\mathrm{int}}$ be the compact subgroup generated by the internal gauge support of the probe window (Proposition 12.8). Restrict the evolutions to K so that $\mathbf{U}^{(A)}(g)=\pi_A(g)$ and $\mathbf{U}^{(B)}(g)=\pi_B(g)$ for $g\in K$ (Definitions 2.108 and 3.15). Since $A\neq B$ act as distinct viable laws on the window, their restricted actions are inequivalent for some compact K (otherwise they would coincide on that window up to diagnostic realized reference configuration equivalence). By the Peter–Weyl/Schur package localized in Proposition 12.10, inequivalence implies *relational character orthogonality*: the K–averaged

intertwiner vanishes and an admissible convex mixture of probes and pulses achieves zero overlap, i.e.

$$\Omega(A, B \mid \rho; g_\star) = \mathrm{Tr}\Big(\rho\,(\mathbf{U}^{(B)}(g_\star))^\dagger \mathbf{U}^{(A)}(g_\star)\Big) = 0$$

for some randomized admissible setting inside $\mathrm{co}\,\mathcal{S}_{\mathrm{IC}}^{(2n)}$ (Proposition 12.46 and Definition 12.44). Equivalently, $\rho_{\mathrm{ens}}(A, B; g_\star) = 0$ by Definition 12.44; compactness of K and lower semicontinuity ensure attainment as noted in Remark 12.47, and in many instances a *single* pure probe and a single pulse suffice by Remark 12.51.

From overlap to vanishing visibility. With $\Omega(A, B \mid \rho; g_\star) = 0$ fixed, choose any of the standard quadrature control settings (available within the admissible gate family) so that the measured difference $p_0^{(\mathrm{asym})} - p_1^{(\mathrm{asym})}$ isolates $\mathrm{Re}\,\Omega$ or $\mathrm{Im}\,\Omega$ (Definition 2.114 and Lemma 12.39). The linear dependence of $p_0^{(\mathrm{asym})} - p_1^{(\mathrm{asym})}$ on Ω then forces $p_0^{(\mathrm{asym})} - p_1^{(\mathrm{asym})} = 0$, hence $\mathcal{V}_{\mathrm{asym}}(A, B) = 0$ by the visibility definition in Section 3.1.5 and eqs. (12.56) and (12.57).

This establishes the claimed existence of an admissible family with $\Omega(A, B \mid \rho) = 0$ and a corresponding interferometric setting with $\mathcal{V}_{\mathrm{asym}}(A, B) = 0$. In particular, Stage–II *relational orthogonality* holds in the sense of Definition 12.38, as also recorded in Corollary 12.50. □

Remark 12.53 (Pure-state refinement)**.** The existence proofs in Corollaries 12.50 and 12.52 use convex ensembles—mixed probes and/or randomized pulses in $\mathrm{co}\,\mathcal{S}_{\mathrm{IC}}^{(2n)}$ (Lemma 12.43 and Definition 12.44). In many finite–dimensional instances, a *single* pure probe and a single pulse already suffice to achieve cancellation. Indeed, let $g(t) = \exp(tX) \subset K$ be a one–parameter subgroup and define the matrix–coefficient curve

$$f_\Psi(t) \;:=\; \langle \Psi |\, V(g(t))\, |\Psi\rangle\,, \qquad V(g) := (\mathbf{U}^{(B)}(g))^\dagger \mathbf{U}^{(A)}(g) \quad \text{(as in Proposition 12.46)}.$$

By continuity of matrix elements on compact K (Propositions 12.8 and 12.10), $t \mapsto f_\Psi(t)$ is continuous. When the induced phase along $g(t)$ is nonconstant (e.g. inequivalent irreducibles contribute characters of opposite winding), the intermediate value principle guarantees $t_\star$ with $f_\Psi(t_\star) = 0$, i.e. When the induced phase along $g(t)$ is nonconstant (e.g. inequivalent irreducibles contribute characters of opposite winding), the intermediate value principle guarantees $t_\star$ with

$$\Omega\Big(A, B \mid \Psi; g(t_\star)\Big) = 0. \tag{12.67}$$

Here $\Omega(\cdot)$ denotes the process–overlap of Definition 12.69; see also eq. (12.83). Selecting a quadrature control within the admissible gate family (Definition 2.114 and Lemma 12.39) then yields

$$p_0^{(\mathrm{asym})} - p_1^{(\mathrm{asym})} = 0, \tag{12.68}$$

and therefore $\mathcal{V}_{\mathrm{asym}}(A, B) = 0$ by Section 3.1.5 and eqs. (12.56) and (12.57). Selecting a quadrature control within the admissible gate family (Definition 2.114 and Lemma 12.39) then yields $p_0^{(\mathrm{asym})} - p_1^{(\mathrm{asym})} = 0$ and hence $\mathcal{V}_{\mathrm{asym}}(A, B) = 0$ by Section 3.1.5 and eqs. (12.56) and (12.57). Thus the ensemble index ρ_{ens} (Definition 12.44) ensures *existence* of orthogonality, while a pure–state/single–pulse realization often obtains by continuity (cf. Remarks 12.47 and 12.51).

Independently of such phase arguments, convex–geometry bounds control the support size of ensemble realizations. On a D–dimensional probe subspace, any density operator

admits a spectral decomposition into at most D pure states [287, Sec. 2.4]. More generally, by Carathéodory's theorem, any point in a convex set lying in a real affine space of dimension m is a convex combination of at most $m+1$ extreme points; applied to the real vector space of Hermitian trace–one operators (dimension D^2-1), this yields a conservative upper bound of D^2 pure components for the probe mixture [555, Thm. 17.1]; see also the original source [556]. Together with compactness of K and the Haar–average approximation by finite convex combinations (Proposition 12.10), this ensures that any ensemble cancellation realizing $\Omega = 0$ can be implemented with a finite support whose size is bounded in terms of D.

Definition 12.54 (Relational character of a candidate action)**.** Let A be a *viable* candidate in the admissible class (local, compact, Tantrakyma–symmetric, chiral) in the sense of Section 12.1.1, Definition 12.33, and Theorem 12.42. Let $K \subset G_{\mathrm{int}}$ be the compact subgroup generated by the internal gauge support of the probe window (Proposition 12.8 and Section 3.1.5). Assume the induced K–action on the (finite) matter probe space decomposes as

$$\mathcal{H}_{\mathrm{mat}} \;=\; \bigoplus_i \mathcal{H}_i, \qquad \mathbf{U}^{(A)}(g)\big|_{\mathcal{H}_i} \;=\; \pi_{A,i}(g), \quad g \in K,$$

with each $\pi_{A,i}$ a finite–dimensional unitary representation (Peter–Weyl/Schur framework; Proposition 12.10). The *relational character* of A on K is the class function

$$\chi_A(g) \;:=\; \sum_i w_i \,\mathrm{Tr}\Big(\pi_{A,i}(g)\Big), \qquad g \in K,$$

where the positive weights w_i encode the physical multiplicities/gradings (e.g., left vs. right, color vs. flavor) used in the Tantrakyma sector (*cf.* the chiral/gauge content and hypercharge tabulation in Section 12.4.3 and table 12.1). Any fixed choice of $\{w_i\}$ that is representation–independent across candidates preserves orthogonality statements, since character inner products and the averaged intertwiner criteria depend only on the K–isomorphism classes of the summands (Propositions 12.10 and 12.46 and Corollary 12.48).

Lemma 12.55 (Relational character overlap)**.** *For any admissible probe ρ supported on $\mathcal{H}_{\mathrm{mat}}$, there are control settings for which the holosymmetric overlap obeys*

$$\Omega(A, B \mid \rho) \;=\; \int_K \chi_B(g)^*\, \chi_A(g)\, \mathrm{d}\mu_\rho(g),$$

where μ_ρ is a probability measure on K determined by the pulse schedule and the probe (it is absolutely continuous w.r.t. Haar for smooth pulses).

Proof. Step 1: Reduction to the K–window and block form. Work on a certified plateau with compact holonomy and rig–compatible connection so that the probe window generates a compact subgroup $K \subset G_{\mathrm{int}}$ and all meters are BRST/Nielsen–stable (Sections 6.12.4 and 12.3, Definition 12.1, and Propositions 12.8 and 12.12). Restrict the induced evolutions to K and write $\mathbf{U}^{(A)}(g) = \pi_A(g)$, $\mathbf{U}^{(B)}(g) = \pi_B(g)$ on $\mathcal{H}_{\mathrm{mat}} = \bigoplus_i \mathcal{H}_i$ with finite–dimensional unitary blocks as in Definition 12.54. Define $V(g) := (\mathbf{U}^{(B)}(g))^\dagger \mathbf{U}^{(A)}(g) = \pi_B(g)^\dagger \pi_A(g)$ and the process–overlap $\Omega(A, B \mid \rho) = \mathrm{Tr}\Big(\rho\, V(g_\bullet)\Big)$ in the single–pulse case (Definition 12.69 and eq. (12.83)).

Step 2: From single pulses to averaged readouts. Use the asymmetric interferometer on the dual–register scaffold; by design, the measured difference $p_0^{(\mathrm{asym})} - p_1^{(\mathrm{asym})}$ depends linearly on $\Omega(A, B \mid \rho)$ (and its conjugate), and gate choices isolate these linear parts (Section 3.1.5, Definitions 2.114 and 2.124, eqs. (12.56) and (12.57), and Lemma 12.39). Allow a (classically randomized) pulse schedule on K with law μ_ρ realized via admissible controls (Definitions 2.108 and 2.114). Linearity of the readout then yields the averaged overlap

$$\Omega(A, B \mid \rho) \;=\; \int_K \mathrm{Tr}\Big(\rho\, V(g)\Big)\, \mathrm{d}\mu_\rho(g),$$

with μ_ρ a probability measure on K (the distribution of pulses). If the schedule is generated by a smooth time–dependent control $g(t) \in K$ with a smooth sampling density on t, then μ_ρ is absolutely continuous w.r.t. Haar by standard pushforward along $g(t)$ on compact groups (cf. Haar framework in Proposition 12.10).

Step 3: Characterization as a character–weighted average. By Definition 12.54, write the relational characters $\chi_A(g) = \sum_i w_i \,\mathrm{Tr}\, \pi_{A,i}(g)$ and $\chi_B(g) = \sum_j w_j \,\mathrm{Tr}\, \pi_{B,j}(g)$, with weights $w_i > 0$ fixed independently of candidates. Choose admissible pre/post controls (unitaries that commute with the K–action on each $\mathcal{H}_i$) so that ρ is block–diagonal and proportional to the identity on each $\mathcal{H}_i$ with the same fixed weights w_i; this is always achievable by local unitary conjugations that preserve the K–representation type and do not affect the interferometric readout (Section 3.1.5 and Definition 2.124). Then for each $g \in K$,

$$\mathrm{Tr}\Big(\rho\, V(g)\Big) = \sum_i w_i \,\frac{1}{d_i}\, \mathrm{Tr}\Big(\pi_{B,i}(g)^\dagger \pi_{A,i}(g)\Big), \qquad d_i = \dim \mathcal{H}_i.$$

Using the Peter–Weyl/Schur machinery (Proposition 12.10), the class–function projection of $g \mapsto \mathrm{Tr}\Big(\pi_{B,i}(g)^\dagger \pi_{A,i}(g)\Big)$ is exactly $\chi_{B,i}(g)^* \chi_{A,i}(g)$ (matrix–coefficient orthogonality contracts non–class parts to their class–function representative). Since the interferometer statistics on a compact window can be made to depend only on the class of g by averaging over conjugacy (admissible by randomized controls; Definitions 2.108 and 2.114 and Proposition 12.10), the same average yields

$$\int_K \mathrm{Tr}\Big(\rho\, V(g)\Big)\, \mathrm{d}\mu_\rho(g) = \int_K \Big(\sum_i w_i\, \chi_{B,i}(g)^* \chi_{A,i}(g)\Big)\, \mathrm{d}\mu_\rho(g) = \int_K \chi_B(g)^*\, \chi_A(g)\, \mathrm{d}\mu_\rho(g),$$

where the last identity is the definition of χ_A, χ_B as in Definition 12.54. Combining with Step 2 establishes the claim.

Step 4: Smooth schedules and absolute continuity. If the pulse schedule is implemented by a smooth density on K (e.g. by convolving Haar with a smooth class function and sampling; admissible on compact groups), then μ_ρ has a continuous Radon–Nikodym derivative w.r.t. Haar, hence is absolutely continuous (Haar background in Proposition 12.10). □

Proposition 12.56 (Character orthogonality ⇒ holosymmetric orthogonality)**.** *Let $K \subset G_{\mathrm{int}}$ be the compact subgroup generated by the internal gauge support of the probe window (Proposition 12.8), with Haar measure μ normalized as in Proposition 12.10. If the matter representations induced by A and B share no common irreducible K–subrepresentations (i.e. the K–actions on $\mathcal{H}_{\mathrm{mat}}$ have disjoint isotypic content in the sense of Definition 12.54), then*

$$\int_K \chi_B(g)^*\, \chi_A(g)\, \mathrm{d}\mu(g) = 0,$$

and consequently there exists an admissible probe ensemble with $\Omega(A,B) = 0$ *(hence* $\mathcal{V}_{\mathrm{asym}}(A,B) = 0$*).*

Proof. Step 1: Character orthogonality on compact K*.* By Definition 12.54, write the (relational) characters of the restricted actions as $\chi_A(g) = \sum_i w_i \operatorname{Tr}(\pi_{A,i}(g))$ and $\chi_B(g) = \sum_j w_j \operatorname{Tr}(\pi_{B,j}(g))$, with positive weights w_i fixed independently of candidates. Since A and B share no common irreducible K–summands, Schur orthogonality/Peter–Weyl on compact groups (Proposition 12.10) implies that the class–function inner product of their characters vanishes:

$$\langle \chi_A, \chi_B \rangle_K := \int_K \chi_B(g)^* \chi_A(g)\, d\mu(g) = 0.$$

(Equivalently, the averaged intertwiner $\int_K \pi_B(g)^\dagger \pi_A(g)\, d\mu(g)$ is the zero map; cf. Proposition 12.10 and Corollary 12.48.)

Step 2: From character average to overlap average. By the relational–character overlap identity Lemma 12.55, for any admissible probe ρ there exist control settings producing a pulse–law μ_ρ on K with

$$\Omega(A,B \mid \rho) \;=\; \int_K \chi_B(g)^* \chi_A(g)\, d\mu_\rho(g).$$

Choosing the (normalized) Haar schedule $\mu_\rho = \mu$ is admissible on compact K up to arbitrarily fine convex approximation by finite pulse schedules implemented with the same gate family (Definitions 2.108 and 2.114 and Proposition 12.10). Hence, using $\langle \chi_A, \chi_B \rangle_K = 0$ from Step 1,

$$\Omega(A,B \mid \rho) = \int_K \chi_B(g)^* \chi_A(g)\, d\mu(g) = 0,$$

and, by convex approximation of the Haar average on compact K, an *admissible ensemble* drawn from $\operatorname{co} \mathcal{S}_{\mathrm{IC}}^{(2n)}$ realizes the same cancellation exactly (Lemma 12.43, Definition 12.44, Proposition 12.46, and Corollary 12.48).

Step 3: Vanishing visibility. With $\Omega(A,B \mid \rho) = 0$ fixed, select a quadrature control within the admissible gate family to isolate $\operatorname{Re}\Omega$ or $\operatorname{Im}\Omega$ in the arm–probability difference $p_0^{(\mathrm{asym})} - p_1^{(\mathrm{asym})}$ (Lemma 12.39 and Definition 2.114). The linear dependence of $p_0^{(\mathrm{asym})} - p_1^{(\mathrm{asym})}$ on Ω then forces $p_0^{(\mathrm{asym})} - p_1^{(\mathrm{asym})} = 0$, and thus the asymmetric visibility vanishes by definition: $\mathcal{V}_{\mathrm{asym}}(A,B) = 0$ (Section 3.1.5 and eqs. (12.56) and (12.57)). Equivalently, Corollary 12.50 gives the same conclusion once $\Omega = 0$ is established. □

Theorem 12.57 (Representation-theoretic sufficiency bridge to G_{SM})**.** *Assume the candidate law* A *satisfies the constraints established earlier in this chapter: (i) compact, local, Tantrakyma–invariant gaugeability (Section 12.1.1, Definition 12.1, and Proposition 12.8); (ii) chiral admissibility with the one–generation matter content of table 12.1 (Section 12.4.3); (iii) complete cancellation of local and global anomalies (Definition 12.20, eq.* (12.46)*, Proposition 12.26, and Corollary 12.27); and (iv)* minimality *(no extra chiral matter beyond vectorlike pairs; a single abelian factor acting nontrivially on matter; Corollary 12.27). Then the induced* K*–action on* $\mathcal{H}_{\mathrm{mat}}$ *is (up to unitary equivalence) the standard action of*

$$G_{\mathrm{SM}} \;=\; SU(3)_C \times SU(2)_L \times U(1)_Y,$$

and for any other viable candidate $B \not\simeq A$ the induced K–modules share no irreducible subrepresentations. Consequently, by Proposition 12.56, there exists an admissible probe ensemble with $\mathcal{V}_{\mathrm{asym}}(A, B) = 0$.

Proof sketch. (1) Lie–algebra factors from representation features. By chiral admissibility and the spectrum in table 12.1, the electroweak left–handed sector contains a two–dimensional pseudoreal doublet; this forces an $\mathfrak{su}(2)$ factor acting as the weak isospin on those doublets (Section 12.4.3). Likewise, the presence of complex color triplets in the matter content and nonvanishing cubic index before family replication singles out an $\mathfrak{su}(3)$ color factor. Fractional electric charges and the Gell–Mann–Nishijima relation $Q = I_3 + \frac{1}{2}Y$ (Section 12.4.3 and eq. (12.107)) require one abelian factor $\mathfrak{u}(1)_Y$.

(2) Rank and index constraints. The mixed–anomaly system (eq. (12.46)) fixes the embedding of $\mathfrak{u}(1)_Y$ uniquely up to an overall normalization, consistent with Section 12.4.3. Any additional simple factor would contribute new Dynkin indices to the mixed anomalies which cannot be cancelled by the minimal multiplet set without introducing exotics, violating minimality (iv) (Corollary 12.27). Multiple nontrivial abelian factors would similarly introduce uncancellable $U(1)'^3$ and $\mathrm{grav}^2 - U(1)'$ anomalies unless they act trivially, in which case they decouple from the K–window (Definition 12.20 and Proposition 12.26).

(3) Exclusion of extra simple/abelian factors. Combining (1)–(2) with compact/local implementability (Definition 12.1 and Proposition 12.8) and Tantrakyma invariance (Section 12.1.1), the only nontrivial internal action on matter consistent with (i)–(iv) is $\mathfrak{su}(3) \oplus \mathfrak{su}(2) \oplus \mathfrak{u}(1)$ with the hypercharges of table 12.1.

(4) Equivalence of the induced modules. With that Lie–algebra content and the fixed multiplet list of table 12.1, the representation ring on $\mathcal{H}_{\mathrm{mat}}$ (highest weights and multiplicities) coincides with the Standard Model one, so the induced K–action of A is unitarily equivalent to the restriction of the G_{SM} action (Peter–Weyl/Schur decomposition on compact windows, Proposition 12.10; see also the definition of the relational character in Definition 12.54).

(5) Orthogonality to any inequivalent candidate. Let B be any other viable candidate whose induced K–action on $\mathcal{H}_{\mathrm{mat}}$ is not equivalent to A's. Then the two K–modules share no common irreducible summands (by Step 4), and thus their relational characters obey $\langle \chi_A, \chi_B \rangle_K = 0$ (Proposition 12.10 and Definition 12.54). By Proposition 12.56 and Lemma 12.55, there exists an admissible probe ensemble with $\Omega(A, B) = 0$; selecting quadrature controls converts this to $\mathcal{V}_{\mathrm{asym}}(A, B) = 0$ (Section 3.1.5, eqs. (12.56) and (12.57), Lemma 12.39, and Definition 2.114). This is precisely the Stage–II sufficiency clause in Theorem 12.42 specialized to the G_{SM} fixed point.

Conclusion. Hence A induces, up to unitary equivalence, the standard G_{SM} action on $\mathcal{H}_{\mathrm{mat}}$, and any inequivalent viable B is holosymmetrically orthogonal to A on some compact K, completing the representation–theoretic sufficiency bridge to the Standard Model. □

Conclusion. Corollary 12.52 settles the *existence* of admissible probe/control families that drive the process–overlap to zero for any two distinct viable laws acting on the same compact window, and—via the asymmetric readout—force vanishing Stage–II visibility (Section 3.1.5, eqs. (12.56) and (12.57), Lemma 12.39, and Definition 2.114). Theorem 12.57 supplies the promised representation–theoretic bridge: within the anomaly–free, minimal, Tantrakyma–compatible class carrying the SM's one–generation chiral content (Sections 12.1.1

and 12.4.3, Definition 12.20, eq. (12.46), Proposition 12.26, and Corollary 12.27), the only nontrivial internal action on matter consistent with compact/local gaugeability (Definition 12.1 and Proposition 12.8) is the standard G_{SM} action. Any other viable candidate $B \not\cong A$ has an induced K–module that shares no irreducible subrepresentations with that of A; by Peter–Weyl/Schur orthogonality (Proposition 12.10 and Remark 12.9) this yields character orthogonality and hence holosymmetric orthogonality (Proposition 12.56 and Lemma 12.55), i.e. there exists an admissible ensemble with $\Omega(A,B) = 0$ and thus $\mathcal{V}_{\mathrm{asym}}(A,B) = 0$ (Corollary 12.50). Together with Stage–I viability ($\mathcal{C}(A) = 1$ and vanishing anomaly meters; Definition 12.33 and Theorem 12.42), these results complete the sufficiency half of the HSP proof–by–uniqueness: within the admissible class, the pair of conditions $\big(\mathcal{C}(A) = 1,\ \rho_{\mathrm{ens}}(A,B) = 0\ \forall B \neq A\big)$ singles out the G_{SM} fixed point (Definition 12.44 and Theorem 12.42).

12.3 BRST Symmetry and the Quantization of Gauge Fields

Local gauge symmetry renders the classical field description redundant; a faithful quantum theory must remove this redundancy without sacrificing covariance or unitarity. In the IQR framework this is accomplished cohomologically by the **BRST** (Becchi–Rouet–Stora–Tyutin) symmetry: gauge invariance is lifted to a nilpotent differential $\hat{Q}_{\mathrm{BRST}}$ acting on an *extended* field space, and the physical Hilbert space is identified with the corresponding cohomology at ghost number zero,

$$\mathcal{H}_{\mathrm{phys}} \ \equiv\ H^0(\hat{Q}_{\mathrm{BRST}}),$$

with $\hat{Q}_{\mathrm{BRST}}^2 = 0$ and the Heisenberg conservation $\frac{d}{dt}\hat{Q}_{\mathrm{BRST}} = 0$ ensuring unitary evolution on H^0 [445, 105, 106, 107, 108]. Throughout, all statements are made on certified plateaux to guarantee gauge/metric hygiene and well-posed readouts (Section 6.12.4 and Proposition 12.12).

Operator–bundle connection and sectoral covariance. Consistently with the sectoral realization of gauge transport (Chapter 6), the gauge potential is the *Operator Bundle Connection* $\mathcal{A}_\mu$ on the realized weave configuration-compatible bundle (Definitions 6.192 and 12.1), with curvature and transformations as in Definition 12.96. The same split connection governs both bulk and boundary transport (Levi–Civita part and internal part), with its boundary-compatible restriction used on probe windows (Definition 12.4). This connection enters the covariant derivatives of the Gaiamiti and Tantrakyma fields (Section 6.5.3 and Definition 9.323), ensuring that the BRST complex is compatible with the realized chronogeometry and compact holonomy (Proposition 12.8).

Quanta, readout, and BRST descent. At the level of quanta (Chapters 8–9), excitations of the sectoral images of $\mathcal{A}_\mu$ are the *mition* and *tantron* (Definitions 9.258 and 9.348); their extraction is performed cohomologically by descending to $H^0(\hat{Q}_{\mathrm{BRST}})$ (Section 9.5 and Lemma 9.54). Physical readouts use holosymmetric meters that act on the BRST-closed sector and are insensitive to gauge/Nielsen choices (Definition 6.275 and Proposition 12.12). In nonabelian color, the BRST treatment implements spectral singlet projection: the physical

hadronic space equals the color-singlet subspace obtained by the gauge-induced spectral filter (Chapter 9), as encoded in Proposition 12.128 and its cohomological form in the hadronic BRST section (Section 12.6). Decoupling of the TT radiative block in readouts is enforced by the realized TT projectors from Chapter 11 (Definition 10.3 and eqs. (10.4) and (10.134)).

Interlock with the HSP (Stage I & Stage II). Stage I of the Holosymmetric Selection Principle requires *viability*: the self–coherence functional saturates, $\mathcal{C}(A) = 1$, if and only if the BRST charge is nilpotent/conserved and all anomaly meters vanish (Theorem 12.42 and Definition 12.33). Concretely, local and global anomalies are monitored by the meters and constraints of Chapter 12 (Definition 12.20, eq. (12.46), Proposition 12.26, and Corollary 12.27); any nonzero anomaly violates BRST conservation and fails Stage I on the plateau. Stage II acts on the resulting gauge–invariant composites using the asymmetric interferometer on the dual-register scaffold, whose arm probabilities and visibilities provide linear, informationally complete access to process overlaps (Section 3.1.5, Definitions 2.114 and 2.124, Lemmas 12.39 and 12.43, and eqs. (12.56) and (12.57)). On compact probe windows K (guaranteed by compact holonomy), Peter–Weyl/Schur orthogonality turns character orthogonality into vanishing holosymmetric overlap (Propositions 12.10, 12.46 and 12.56, Corollary 12.48, and Lemma 12.55), yielding $\mathcal{V}_{\mathrm{asym}}(A, B) = 0$ for inequivalent viable laws (Corollaries 12.50 and 12.52).

Electroweak charges and anomaly hygiene. Within the Tantrakyma sector, electroweak symmetry breaking respects the Gell–Mann–Nishijima relation $Q = I_3 + \frac{1}{2}Y$ (Section 12.4.3) and the one–generation hypercharge assignments of table 12.1 and Section 12.4.3. These are precisely the inputs that make the anomaly block eq. (12.46) solvable with vanishing meters on the plateau (Proposition 12.26), fitting the Stage I requirement above. The representation–theoretic sufficiency bridge then singles out the Standard Model action among all admissible, minimal, Tantrakyma–compatible candidates (Chapter 12; Theorem 12.57), completing the HSP proof-by-uniqueness on BRST-clean grounds (Theorem 12.42).

12.3.1 The Extended Field Space

Let $\mathcal{A}_\mu(x)$ be the *Operator Bundle Connection* on the modal fibres, transforming as a connection one–form under local changes of quantum frame (cf. Definition 6.192 and eq. (6.264)). In the IQR setting $\mathcal{A}_\mu$ arises intrinsically from the observable algebra via the GNS representation of the Quantum Holo-State (cf. Definitions 1.32 and 1.34 and Remark 6.198), and acts sectorally on the substrate fields through the maps $\mathcal{R}_\Phi, \mathcal{R}_\Theta$ that enter the split covariant derivatives (cf. Section 6.5.3 and Definition 9.323). To implement BRST quantization on a certified plateau (so that physical readouts are BRST/Nielsen–stable; Sections 6.12.4 and 12.3 and Proposition 12.12), enlarge the field content by Grassmann–odd ghosts and their auxiliaries:

- **Gauge field:** $\mathcal{A}_\mu = \mathcal{A}_\mu^a T^a$, Lie–algebra valued, Grassmann–even, $\mathrm{gh}(\mathcal{A}_\mu) = 0$.
- **Ghost:** $c = c^a T^a$, Grassmann–odd, $\mathrm{gh}(c) = +1$ [445, 105, 106].
- **Antighost:** $\bar{c} = \bar{c}^a T^a$, Grassmann–odd, $\mathrm{gh}(\bar{c}) = -1$ [445, 105, 106].

- **Nakanishi–Lautrup field:** $B = B^a T^a$, Grassmann–even, $\mathrm{gh}(B) = 0$. This is the BRST–auxiliary enforcing the gauge condition off–shell (linear in B in the gauge–fixing functional); it is standard in the Kugo–Ojima BRST construction [108] and originates in the covariant gauge formulations of Nakanishi and Lautrup [551, 557].

(Here T^a are anti–Hermitian generators with $[T^a, T^b] = f^{abc}T^c$ and $\mathrm{Tr}(T^aT^b) = -\frac{1}{2}\delta^{ab}$.)

In this extended field space, the BRST charge $\hat{Q}_{\mathrm{BRST}}$ is nilpotent and conserved on the plateau, and the physical Hilbert space is the cohomology $H^0(\hat{Q}_{\mathrm{BRST}})$ (see Section 12.3); the band–tagged meters and interferometric readouts are designed to descend to this sector (Definition 12.16 and Lemma 12.17).

The BRST Differential and Physical States

Definition 12.58 (BRST differential)**.** On the extended field space of Section 12.3.1, the BRST differential δ_{BRST} is a graded derivation of ghost number $+1$ acting by

$$\delta_{\mathrm{BRST}}\mathcal{A}_\mu = D_\mu c \;=\; \partial_\mu c + [\mathcal{A}_\mu, c], \tag{12.69}$$

$$\delta_{\mathrm{BRST}}c = -\tfrac{1}{2}[c, c], \tag{12.70}$$

$$\delta_{\mathrm{BRST}}\bar{c} = B, \tag{12.71}$$

$$\delta_{\mathrm{BRST}}B = 0, \tag{12.72}$$

and, for any matter field Ψ in a representation R, $\delta_{\mathrm{BRST}}\Psi = i\,c^a(T^a_R)\,\Psi$. Concretely, for the substrate fields one has $\Psi \in \{\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}\}$ and $T^a_R \in \{\mathcal{R}_\Phi(T^a), \mathcal{R}_\Theta(T^a)\}$ (cf. Section 6.5.3 and Definition 9.323). Its defining property is *nilpotency*:

$$\delta^2_{\mathrm{BRST}} = 0. \tag{12.73}$$

In the band–tagged setting of Chapters 7–9, the BRST action is compatible with the tags and descends to cohomology (chain–map property), so that physical observables and the gauge–boson quanta (*mition*, *tantron*) are realized as classes in $H^0(\hat{Q}_{\mathrm{BRST}})$ (cf. Lemma 9.54 and Section 9.5; diagnostic realized reference configuration hygiene in Section 6.12.4 and Proposition 12.12).

Proof sketch of (12.73). The inclusion of the Nakanishi–Lautrup field B (Grassmann–even, $\mathrm{gh} = 0$) ensures *off–shell* nilpotency (no use of equations of motion), matching the standard BRST construction.

Gauge field. Using D_μ–Leibniz and adjoint covariance,

$$\delta^2\mathcal{A}_\mu = \delta(D_\mu c) = D_\mu(\delta c) + [\,\delta\mathcal{A}_\mu,\, c\,] = D_\mu\Big(-\tfrac{1}{2}[c, c]\Big) + [\,D_\mu c,\, c\,] \overset{\text{Jacobi}}{=} 0.$$

Ghost. Graded Jacobi and antisymmetry give

$$\delta^2 c = -\tfrac{1}{2}[\,\delta c,\, c\,] - \tfrac{1}{2}[\,c,\, \delta c\,] = -\tfrac{1}{2}\Big[-\tfrac{1}{2}[c, c],\, c\Big] - \tfrac{1}{2}\Big[c,\, -\tfrac{1}{2}[c, c]\Big] = 0.$$

Antighost/NL field. By definition, $\delta\bar{c} = B$ and $\delta B = 0$, hence $\delta^2\bar{c} = \delta B = 0$.

Matter. For Ψ in representation R,

$$\delta^2\Psi = i\,\delta c\,\Psi + i\,c\,(\delta\Psi) = i\,\delta c\,\Psi + i\,c\,(i\,c\,\Psi) = i\Big(\delta c + \tfrac{i}{2}[c,c]\Big)\Psi = 0,$$

using $\delta c = -\frac{1}{2}[c,c]$ and graded antisymmetry.

These computations establish $\delta^2_{\mathrm{BRST}} = 0$ off–shell; see [105, 106]. On a certified plateau, the nilpotent charge $\hat{Q}_{\mathrm{BRST}}$ implementing Definition 12.58 is conserved and its cohomology defines the physical Hilbert space (Section 12.3), a condition also required by Stage–I viability in Theorem 12.42. Compatibility with band tags and diagnostic realized reference configuration projectors (chain–map property) follows from the tag/connection covariance and BRST–closure discussed in Lemmas 9.54 and 12.17, ensuring that the induced action on $H^0(\hat{Q}_{\mathrm{BRST}})$ is well defined and Nielsen–flat (Proposition 12.12). □

Definition 12.59 (Gauge fixing via a BRST gauge–fermion)**.** Work in the extended field space of Section 12.3.1 with BRST rules in Definition 12.58. For a gauge–fixing functional $\mathcal{F}[\mathcal{A}]$ and parameter $\xi > 0$, define the *gauge–fixing fermion*

$$\Psi_{\mathrm{gf}} \;:=\; \int d^4x\ \mathrm{Tr}\Big(\,\bar{c}\,[\,\mathcal{F}[\mathcal{A}] - \tfrac{\xi}{2}B\,]\,\Big),$$

where $\mathcal{A}_\mu$ is the Operator–Bundle Connection acting on modal fibres (Definitions 6.192 and 12.96), whose sectoral actions enter the split covariant derivatives for the Gaiamiti and Tantrakyma fields (Section 6.5.3 and Definition 9.323). The fields $(c, \bar{c}, B)$ and their BRST transformations are as in Section 12.3.1 and Definition 12.58 (standard BRST/Nakanishi–Lautrup structure [105, 106, 108, 551, 557]).

Lemma 12.60 (BRST exactness of the gauge–fixing + ghost sector; R_ξ form and Nielsen flatness)**.** *With Ψ_{gf} from Definition 12.59, the gauge–fixing plus ghost Lagrangian is BRST exact:*

$$\mathcal{L}_{\mathrm{gf+gh}} := \delta_{\mathrm{BRST}}\,\Psi_{\mathrm{gf}} \;=\; \mathrm{Tr}\Big(B\,\mathcal{F}[\mathcal{A}] - \tfrac{\xi}{2}B^2 - \bar{c}\,\tfrac{\delta\mathcal{F}}{\delta\mathcal{A}_\mu}\,D_\mu c\Big), \tag{12.74}$$

and for the Lorenz choice $\mathcal{F}[\mathcal{A}] = \partial^\mu\mathcal{A}_\mu$ one recovers the standard R_ξ Lagrangian. On certified plateaux (Section 6.12.4), expectations of BRST–closed observables in the physical sector $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ are ξ–independent (Nielsen flat) (Proposition 12.12 and Lemma 12.17); consequently, the sectoral gauge quanta (mition/tantron) extracted in the spectral channel are gauge–parameter independent (Section 9.5 and Definitions 6.275, 9.258 and 9.348).

Proof. Compute $\delta_{\mathrm{BRST}}\Psi_{\mathrm{gf}}$ using Definition 12.58. Since $\delta\bar{c} = B$, $\delta B = 0$, and $\delta\mathcal{A}_\mu = D_\mu c$ with $D_\mu = \partial_\mu + [\mathcal{A}_\mu, \cdot]$, one finds

$$\delta_{\mathrm{BRST}}\Psi_{\mathrm{gf}} = \int d^4x\,\mathrm{Tr}\Big(B\,\mathcal{F}[\mathcal{A}] - \tfrac{\xi}{2}B^2 - \bar{c}\,\tfrac{\delta\mathcal{F}}{\delta\mathcal{A}_\mu}\,D_\mu c\Big),$$

which is (12.74). For $\mathcal{F}[\mathcal{A}] = \partial^\mu\mathcal{A}_\mu$, integrating by parts and using cyclicity of the trace gives the textbook R_ξ form $(B\,\partial\cdot\mathcal{A} - \frac{\xi}{2}B^2 - \bar{c}\,\partial^\mu D_\mu c)$ [105, 106, 108].

On a certified plateau, the cohomological projector P_{phys} exists and readouts are Ward/Källén–Lehmann clean (Sections 6.12.4 and 12.3 and Proposition 12.12). Along the R_ξ family, the

Nielsen identity states that the ξ–derivative of any physical expectation value is a BRST–exact insertion; hence it vanishes on $H^0(\hat{Q}_{\mathrm{BRST}})$, giving gauge–parameter independence of physical poles/residues (Nielsen flatness). Band–tagged operators commute with the physical projector and with ∇_μ (Lemma 12.17), so the same conclusion applies to the channel–resolved diagnostics (Definition 6.275). Therefore the transverse poles identifying the mition/tantron (Section 9.5 and Definitions 9.258 and 9.348) are ξ–independent. □

Proposition 12.61 (Conservation and unitarity on cohomology)**.** *Let* $\mathcal{L} = \mathcal{L}_{\mathrm{YM}}+\mathcal{L}_{\mathrm{matter}}+\mathcal{L}_{\mathrm{gf+gh}}$, *with* $\mathcal{L}_{\mathrm{YM}} = -\frac{1}{2}\mathrm{Tr}\,(\mathcal{F}_{\mu\nu}\mathcal{F}^{\mu\nu})$ *and* $\mathcal{F}_{\mu\nu} = \partial_\mu\mathcal{A}_\nu - \partial_\nu\mathcal{A}_\mu + [\mathcal{A}_\mu, \mathcal{A}_\nu]$. *Then* $\delta_{\mathrm{BRST}}\mathcal{L}$ *is a total derivative; consequently there exists a conserved Noether charge* $\hat{Q}_{\mathrm{BRST}}$ *with*

$$\hat{Q}^2_{\mathrm{BRST}} = 0, \qquad [\hat{H}_{\mathrm{total}}, \hat{Q}_{\mathrm{BRST}}] = 0. \tag{12.75}$$

The physical state space is the ghost-number-zero cohomology

$$\mathcal{H}_{\mathrm{phys}} \;:=\; H^0(\hat{Q}_{\mathrm{BRST}}) = \ker \hat{Q}_{\mathrm{BRST}} \cap \ker N_{\mathrm{gh}} \,\big/\, \mathrm{im}\, \hat{Q}_{\mathrm{BRST}}, \tag{12.76}$$

on which the S-matrix is unitary (Kugo–Ojima quartet mechanism). In the IQR framework, $\mathcal{H}_{\mathrm{phys}}$ *is precisely the sector used for the holosymmetric readouts of Chapter 7 (channel-resolved visibilities; cf. Definition 6.275) and for the realization of sectoral gauge quanta in Chapter 10 (BRST chain-map and band-tag descent; cf. Lemma 9.54 and Section 9.5). Accordingly, the physical poles/residues of the* mition *and* tantron *obtained in the spectral channel are invariant under gauge-parameter variations and are to be interpreted on* $\mathcal{H}_{\mathrm{phys}}$ *(cf. Definitions 9.258 and 9.348; plateau certification in Section 6.12.4).*

Proof. We prove the three claims in order: (i) $\delta_{\mathrm{BRST}}\mathcal{L}$ is a total derivative; (ii) the Noether charge $\hat{Q}_{\mathrm{BRST}}$ is conserved and nilpotent; (iii) the physical state space is $H^0(\hat{Q}_{\mathrm{BRST}})$ and the S–matrix is unitary thereon.

(i) $\delta_{\mathrm{BRST}}\mathcal{L}$ is a total derivative. Decompose $\mathcal{L} = \mathcal{L}_{\mathrm{YM}} + \mathcal{L}_{\mathrm{matter}} + \mathcal{L}_{\mathrm{gf+gh}}$ with $\mathcal{L}_{\mathrm{YM}} = -\frac{1}{2}\mathrm{Tr}(\mathcal{F}_{\mu\nu}\mathcal{F}^{\mu\nu})$ and $\mathcal{F}_{\mu\nu} = \partial_\mu\mathcal{A}_\nu - \partial_\nu\mathcal{A}_\mu + [\mathcal{A}_\mu, \mathcal{A}_\nu]$. By the definition of the BRST differential (Definition 12.58), $\delta_{\mathrm{BRST}}\mathcal{A}_\mu = D_\mu c$ and $\delta_{\mathrm{BRST}}c = -\frac{1}{2}[c, c]$. Since $\mathcal{L}_{\mathrm{YM}}$ is gauge invariant and the BRST variation is an infinitesimal gauge transformation with Grassmann parameter c, one has

$$\delta_{\mathrm{BRST}}\mathcal{L}_{\mathrm{YM}} = \partial_\mu K^\mu_{\mathrm{YM}} \quad \text{for some local current } K^\mu_{\mathrm{YM}} \;\; (\text{trace cyclicity and } D_{[\mu}\mathcal{F}_{\nu\rho]} = 0).$$

For the matter part, fields transform as $\delta_{\mathrm{BRST}}\Psi = i\,c^a(T^a_R)\Psi$ (Definition 12.58), and with covariant derivatives built from the *Operator Bundle Connection* (Definition 6.192) via the sectoral maps that define the split derivatives (Section 6.5.3 and Definition 9.323), gauge invariance again implies $\delta_{\mathrm{BRST}}\mathcal{L}_{\mathrm{matter}} = \partial_\mu K^\mu_{\mathrm{matter}}$. Finally, by *BRST exactness* of the gauge-fixing+ghost term (Definition 12.59),

$$\delta_{\mathrm{BRST}}\mathcal{L}_{\mathrm{gf+gh}} = \delta^2_{\mathrm{BRST}}\Psi_{\mathrm{gf}} = 0 \quad (\text{nilpotency eq. (12.73)}).$$

Collecting terms, $\delta_{\mathrm{BRST}}\mathcal{L} = \partial_\mu(K^\mu_{\mathrm{YM}} + K^\mu_{\mathrm{matter}})$, a total derivative.

(ii) Existence, conservation, and nilpotency of $\hat{Q}_{\mathrm{BRST}}$. By Noether's theorem applied to the BRST symmetry of the action, there exists a conserved current j^{μ}_{BRST} with $\partial_{\mu} j^{\mu}_{\mathrm{BRST}} = 0$ and

$$\hat{Q}_{\mathrm{BRST}} := \int_{\mathbb{R}^3} d^3x \, j^0_{\mathrm{BRST}}(t, \mathbf{x}), \qquad \frac{d}{dt}\hat{Q}_{\mathrm{BRST}} = 0 \iff [\hat{H}_{\mathrm{total}}, \hat{Q}_{\mathrm{BRST}}] = 0,$$

which gives the second equality in eq. (12.75). The action of δ_{BRST} on any local functional $\mathcal{O}$ can be written as a graded commutator $i[\hat{Q}_{\mathrm{BRST}}, \mathcal{O}]_{\pm}$, hence

$$\delta^2_{\mathrm{BRST}}\mathcal{O} = i[\hat{Q}_{\mathrm{BRST}},\, i[\hat{Q}_{\mathrm{BRST}}, \mathcal{O}]_{\pm}]_{\pm} = -[\,\hat{Q}^2_{\mathrm{BRST}},\ \mathcal{O}]_{\pm}.$$

Using nilpotency at the Lagrangian level (eq. (12.73)), the left-hand side vanishes for all $\mathcal{O}$, implying $\hat{Q}^2_{\mathrm{BRST}} = 0$, i.e. the first relation in eq. (12.75).

(iii) Physical space and unitarity of the S–matrix on $H^0(\hat{Q}_{\mathrm{BRST}})$. Define the ghost-number operator N_{gh} and the physical space as in eq. (12.76). The quartet (Kugo–Ojima) mechanism [107, 108] pairs unphysical degrees of freedom (longitudinal/time-like gauge modes and ghosts) into BRST doublets; doublets contribute only $\hat{Q}_{\mathrm{BRST}}$–exact states and thus vanish in cohomology. Since $[\hat{H}_{\mathrm{total}}, \hat{Q}_{\mathrm{BRST}}] = 0$ (previous step), time evolution preserves BRST cohomology classes; equivalently, the scattering operator S satisfies $[S, \hat{Q}_{\mathrm{BRST}}] = 0$ and therefore induces a well-defined operator on $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$. Positivity (absence of negative-norm states) on $\mathcal{H}_{\mathrm{phys}}$ follows from the exclusion of BRST doublets; hence S is unitary on $\mathcal{H}_{\mathrm{phys}}$.

Compatibility with the IQR instrumentation (Chs. 7, 8, 10, 11). (i) *Sectoral gauge field and split transport.* The gauge field in all steps above is the universal *Operator Bundle Connection* $\mathcal{A}_{\mu}$ (Definition 6.192 and Remark 6.200), whose sectoral actions enter the covariant derivatives for the Gaiamiti and Tantrakyma fields (Section 6.5.3 and Definition 9.323). (ii) *Band/BRST chain map and plateau license.* On the UV/IR band (Chapter 8, Definition 6.282) the BRST action commutes with the band tag, so diagnostics and spectra *descend* to cohomology (Chapter 10, Lemma 9.54); on certified plateaux the gauge-parameter ξ drops out of physical readouts (Section 6.12.4). (iii) *Channel-resolved readouts and sectoral quanta.* Holosymmetric visibilities are evaluated with triadic projectors (Definition 6.275); the sectoral gauge-boson quanta of $\mathcal{A}_{\mu}$—the *mition* and *tantron*—are defined in Chapter 10 and realized in the BRST-physical sector $H^0(\hat{Q}_{\mathrm{BRST}})$ via the Σ-causal construction (Definitions 9.258 and 9.348 and Section 9.5). (iv) *Typed/TT structure.* Typed projectors and TT closure from Chapter 11 ensure that gravitational TT content is separated from internal gauge sectors at the level of correlators, so the BRST argument above applies independently in the internal (tantrakyma) channels used for spectral extraction.

Combining these points establishes Proposition 12.61. □

Remark 12.62 (Anomalies and HSP Stage I (BRST viability))**.** A gauge (BRST) anomaly manifests as a failure of the cohomological criteria proved in Proposition 12.61: either $\hat{Q}^2_{\mathrm{BRST}} \neq 0$ or $[\hat{H}_{\mathrm{total}}, \hat{Q}_{\mathrm{BRST}}] \neq 0$ in (12.75), or the path–integral measure fails to be invariant under the BRST differential (Definition 12.58) even when the gauge fixing is BRST–exact (Definition 12.59). In the band–tagged setting the anomaly obstructs the chain–map/descend

property (Chs. 8–10; cf. Lemma 9.54 and Section 9.5), so physical readouts on $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$ (12.76) are not well defined and the plateau license for channel–resolved visibilities fails (cf. Section 6.12.4 and Definition 6.275). Within the Holosymmetric Selection Principle (Ch. 13), this precisely triggers a Stage I veto: the candidate internal automorphism A attains holosymmetric score $\mathcal{C}(A) < 1$ and is excluded at *Stage I* (viability), prior to any Stage II analysis of gauge–invariant composites (cf. Section 12.2). Operationally, because the gauge field here is the universal Operator Bundle Connection $\mathcal{A}_\mu$ (Definition 6.192 and Remark 6.200), such an anomaly would also obstruct the sectoral realization of its quanta—the *mition* and *tantron*—defined in Chapter 10 (cf. Definitions 9.258 and 9.348) and normally extracted in the spectral channel after BRST descent (Section 9.5).

Summary. BRST symmetry furnishes a covariant, cohomological prescription for quantizing the operator–bundle gauge theory built on the universal connection $\mathcal{A}_\mu$ (cf. Definition 6.192 and Remark 6.200): gauge fixing is BRST exact (12.74), the nilpotent, conserved charge (12.75) selects the physical space (12.76), and anomalies are precisely the obstruction to *Stage I* viability in the Holosymmetric Selection Principle (cf. Remark 12.62 and Ch. 13, Section 12.2). In the IQR instrumentation this prescription is implemented on the UV/IR band via the scale–tagging functor (cf. Ch. 8, Definition 6.282), with BRST acting as a chain map so that diagnostics *descend* to cohomology on certified plateaux (cf. Ch. 7, Section 6.12.4, and Ch. 10, Lemma 9.54 and Section 9.5). Physical readouts—including the spectral poles/residues that define the sectoral gauge quanta of $\mathcal{A}_\mu$, namely the *mition* and *tantron*—are therefore gauge–parameter independent and evaluated in the channel–resolved meters of Chapter 7 (cf. Definition 6.275 and Ch. 10, Definitions 9.258 and 9.348). The gravitational TT content used alongside internal sectors is isolated by the background–covariant projectors of Chapter 11, ensuring a clean separation of spin–2 (graviton) dynamics from internal gauge channels when forming BRST–safe observables. In Section 12.6 we compute $H^0(\hat{Q}_{\text{BRST}})$ explicitly for QCD, proving that the hadronic Hilbert space equals the color–singlet projection, in harmony with the general cohomological construction above and its holosymmetric consistency checks (Ch. 13, Section 12.2).

Principle 12.1 (Physical state selection via BRST cohomology)**.** *Let $\mathcal{H}_{\text{total}}$ be the gauge–fixed state space (including ghosts and auxiliary fields) equipped with the ghost–number grading N_{gh}, and let $\hat{Q}_{\text{BRST}}$ be the conserved, nilpotent BRST charge generated by the symmetry of the gauge–fixed action (Section 12.3; cf. (12.75)). The* physical Hilbert space *is the ghost–number–zero BRST cohomology:*

$$\mathcal{H}_{\text{phys}} := H^0(\hat{Q}_{\text{BRST}}) = \frac{\ker \hat{Q}_{\text{BRST}} \cap \ker N_{\text{gh}}}{\operatorname{im} \hat{Q}_{\text{BRST}} \cap \ker N_{\text{gh}}} \subset \mathcal{H}_{\text{total}}. \tag{12.77}$$

Equivalently, a state $|\Psi\rangle$ is physical iff (i) $\hat{Q}_{\text{BRST}}|\Psi\rangle = 0$ and (ii) $|\Psi\rangle \not\sim |\Psi\rangle + \hat{Q}_{\text{BRST}}|\Lambda\rangle$ for any $|\Lambda\rangle$ in $\ker N_{\text{gh}}$ (closed but not exact). The S–matrix restricted to $\mathcal{H}_{\text{phys}}$ is unitary by the Kugo–Ojima quartet mechanism [105, 106, 107, 108].

Relational gauge quantization via BRST cohomology

Definition 12.63 (Relational gauge quantization via BRST cohomology)**.** A gauge sector in the IQR framework is *BRST–consistent* when the following structural and dynamical conditions hold (all fields built from the universal Operator Bundle Connection $\mathcal{A}_\mu$; cf. Definition 6.192 and Remark 6.200):

1. **Nilpotent differential and grading.** There exists a globally defined, ghost–number–+1 graded derivation δ_{BRST} on the extended field algebra $\mathcal{F}_{\mathrm{BRST}}$ (gauge field, ghosts, antighosts, auxiliaries) such that
$$\delta_{\mathrm{BRST}}^2 = 0, \qquad [N_{\mathrm{gh}}, \delta_{\mathrm{BRST}}] = \delta_{\mathrm{BRST}},$$
with action on generators given in Definition 12.58 (see (12.69)–(12.72)). By Noether's theorem there is a densely defined operator $\hat{Q}_{\mathrm{BRST}}$ with $\delta_{\mathrm{BRST}}(\,\cdot\,) = [\hat{Q}_{\mathrm{BRST}}, \cdot\,\}$ and $\hat{Q}_{\mathrm{BRST}}^2 = 0$ (Proposition 12.61).
2. **Dynamical compatibility.** The full (gauge–fixed) relational Hamiltonian $\hat{H}_{\mathrm{total}}$ is BRST invariant:
$$[\hat{H}_{\mathrm{total}}, \hat{Q}_{\mathrm{BRST}}] = 0,$$
so that time evolution descends to $\mathcal{H}_{\mathrm{phys}}$ and preserves BRST cohomology classes (Section 12.3, (12.75)). In the sectoral split of Chapter 7, this invariance is implemented with covariant derivatives that couple $\mathcal{A}_\mu$ through the representations $\mathcal{R}_\Phi, \mathcal{R}_\Theta$ of the Gaiamiti and Tantrakyma fields (cf. Section 6.5.3 and Definition 9.323).
3. **Gauge fixing as a BRST exact deformation.** There exists a gauge–fermion functional Ψ_{gf} such that the gauge–fixing plus ghost Lagrangian is BRST exact,
$$\mathcal{L}_{\mathrm{gf+gh}} = \delta_{\mathrm{BRST}} \Psi_{\mathrm{gf}},$$
e.g. the R_ξ family in (12.74) (Definition 12.59). On certified plateaux the dependence on ξ cancels in physical readouts by the BRST/Nielsen mechanism for band–tagged diagnostics (Chs. 8–10; cf. Sections 6.12.4 and 9.5, Definition 6.282, and Lemma 9.54).
4. **BRST–closed observables and operator cohomology.** Relational observables $\hat{\mathcal{O}}$ satisfy $\delta_{\mathrm{BRST}} \hat{\mathcal{O}} = 0$; two representatives that differ by a BRST–exact term give identical matrix elements on $\mathcal{H}_{\mathrm{phys}}$:
$$\hat{\mathcal{O}} \sim \hat{\mathcal{O}} + [\hat{Q}_{\mathrm{BRST}}, \hat{X}\} \quad \Longrightarrow \quad \langle \Psi_{\mathrm{phys}} | \hat{\mathcal{O}} | \Phi_{\mathrm{phys}} \rangle = \langle \Psi_{\mathrm{phys}} | (\hat{\mathcal{O}} + [\hat{Q}_{\mathrm{BRST}}, \hat{X}\}) | \Phi_{\mathrm{phys}} \rangle.$$
In practice these observables are the holosymmetric, channel–resolved meters of Chapter 7 evaluated on $\mathcal{H}_{\mathrm{phys}}$ (Definition 6.275), including the spectral channel used to read the sectoral gauge quanta of $\mathcal{A}_\mu$ (mition/tantron; cf. Ch. 10, Definitions 9.258 and 9.348).
5. **Cohomological identification of the spectrum.** The physical Hilbert space is the ghost–number–zero cohomology *by definition*,
$$\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}}),$$
as in (12.76) (cf. also (12.77), if defined). Unphysical (longitudinal/timelike/ghost) excitations assemble into BRST quartets and do not contribute to H^0 (Kugo–Ojima) [107, 108]. The gravitational TT content remains cleanly separated by the background–covariant projectors of Chapter 11 when forming BRST–safe observables in the internal gauge channels.

Interpretation. BRST consistency elevates gauge fixing to a *spectral* prescription: the theory's observable content is the BRST cohomology, which is invariant under gauge–fermion deformations and stable under the dynamics (cf. Definition 12.59 and Proposition 12.61). In the IQR setting this aligns with the Holosymmetric Selection Principle's *Stage I* viability test in Chapter 13: failure of any item above (e.g. a BRST anomaly) depresses the self–coherence functional $\mathcal{C}$ below unity and excludes the candidate internal automorphism at Stage I (cf. Section 12.2 and Remark 12.62). Concretely, the universal connection $\mathcal{A}_\mu$ transports both substrate fields (Ch. 7; Definition 6.192 and Remark 6.200), its sectoral quanta (mition/-tantron) are realized on $\mathcal{H}_{\text{phys}}$ by the band/BRST chain–map and Σ–causal construction (Ch. 10; Definitions 9.258 and 9.348, Lemma 9.54, and Section 9.5), and the TT projectors of Chapter 11 ensure spin–2 gravitational content is kept orthogonal to internal gauge channels during spectral extraction. This same machinery underlies confinement and the color–singlet projection: in QCD one finds $H^0(\hat{Q}_{\text{BRST}}) \cong$ (color singlets of quark–gluon Fock space), as proven in Section 12.6.

Consistency with relational dynamics

Proposition 12.64 (BRST compatibility with the relational Hamiltonian)**.** *Let $\hat{H}_{\text{total}}$ be the gauge–fixed* relational *Hamiltonian on a dense, invariant domain $\mathcal{D} \subset \mathcal{H}_{\text{total}}$ and let $\hat{Q}_{\text{BRST}}$ be the nilpotent, conserved BRST charge of Section 12.3 (cf. (12.75)). Then*

$$[\hat{H}_{\text{total}}, \hat{Q}_{\text{BRST}}] = 0 \quad \Longleftrightarrow \quad [\hat{H}_{\text{total}}, \delta_{\text{BRST}}] = 0 \text{ on local polynomials,} \tag{12.78}$$

and the Schrödinger evolution $U(t) = e^{-it\hat{H}_{\text{total}}}$ descends to the BRST cohomology:

$$U(t) \,:\, H^0(\hat{Q}_{\text{BRST}}) \longrightarrow H^0(\hat{Q}_{\text{BRST}}), \qquad U(t)\,[\Psi] = [\,U(t)\Psi\,]. \tag{12.79}$$

Equivalently, there exists a cohomological projector $\Pi_{\text{phys}} : \mathcal{H}_{\text{total}} \to \mathcal{H}_{\text{phys}}$ with $[\hat{H}_{\text{total}}, \Pi_{\text{phys}}] = 0$.

Proof. **Preliminaries.** The gauge field is the *Operator Bundle Connection* $\mathcal{A}_\mu$ (Definition 6.192 and Remark 6.200), acting sectorally via the split covariant derivatives on $\hat{\mathbf{\Phi}}$ and $\hat{\mathbf{\Theta}}$ (cf. Section 6.5.3 and Definition 9.323). Gauge fixing is introduced as a BRST–exact deformation (Definition 12.59, (12.74)). The BRST differential and charge are defined in Definition 12.58, with nilpotency and conservation summarized in (12.75) and the physical space in (12.76) (see also Proposition 12.61). On the UV/IR band, the band tag commutes with BRST (chain–map property), so all diagnostics descend to cohomology and are gauge–parameter independent on certified plateaux (Chs. 8–10; Definition 6.282, Lemma 9.54, and Sections 6.12.4 and 9.5). Stage I of the Holosymmetric Selection Principle requires these BRST criteria to hold (Ch. 13; Section 12.2 and Remark 12.62).

(1) $[\hat{H}_{\text{total}}, \hat{Q}_{\text{BRST}}] = 0 \Rightarrow [\hat{H}_{\text{total}}, \delta_{\text{BRST}}] = 0$**.** Let $\mathcal{O}$ be any local polynomial in the extended field algebra $\mathcal{F}_{\text{BRST}}$. By definition, $\delta_{\text{BRST}}\mathcal{O} = [\hat{Q}_{\text{BRST}}, \mathcal{O}\}$ (graded commutator; Definition 12.58). Using the graded Jacobi identity,

$$[\hat{H}_{\text{total}}, \delta_{\text{BRST}}\mathcal{O}] = [\hat{H}_{\text{total}}, [\hat{Q}_{\text{BRST}}, \mathcal{O}\}] = [[\hat{H}_{\text{total}}, \hat{Q}_{\text{BRST}}], \mathcal{O}] + [\hat{Q}_{\text{BRST}}, [\hat{H}_{\text{total}}, \mathcal{O}]] = 0,$$

since $[\hat{H}_{\text{total}}, \hat{Q}_{\text{BRST}}] = 0$ by hypothesis. Thus $[\hat{H}_{\text{total}}, \delta_{\text{BRST}}] = 0$ on local polynomials.

(2) $[\hat{H}_{\text{total}}, \delta_{\text{BRST}}] = 0 \Rightarrow [\hat{H}_{\text{total}}, \hat{Q}_{\text{BRST}}] = 0$**.** Assume $[\hat{H}_{\text{total}}, \delta_{\text{BRST}}] = 0$ on $\mathcal{F}_{\text{BRST}}$. Then for all local $\mathcal{O}$,

$$0 = [\hat{H}_{\text{total}}, \delta_{\text{BRST}}\mathcal{O}] = [\hat{H}_{\text{total}}, [\hat{Q}_{\text{BRST}}, \mathcal{O}\}] = [[\hat{H}_{\text{total}}, \hat{Q}_{\text{BRST}}], \mathcal{O}] + [\hat{Q}_{\text{BRST}}, [\hat{H}_{\text{total}}, \mathcal{O}]].$$

Because $[\hat{Q}_{\text{BRST}}, [\hat{H}_{\text{total}}, \mathcal{O}]]$ is itself a graded commutator with $\hat{Q}_{\text{BRST}}$, the family of equalities for all generators $\mathcal{O}$ implies that $[[\hat{H}_{\text{total}}, \hat{Q}_{\text{BRST}}], \mathcal{O}] = 0$ on the dense, invariant domain $\mathcal{D}$ generated by those fields and their conjugate momenta. By irreducibility of the local field algebra on $\mathcal{D}$, it follows that $[\hat{H}_{\text{total}}, \hat{Q}_{\text{BRST}}]$ is a c-number multiple of the identity; since it raises ghost number by $+1$ while the identity has ghost number 0 (grading in Definition 12.58), that multiple must vanish. Hence $[\hat{H}_{\text{total}}, \hat{Q}_{\text{BRST}}] = 0$.

(3) Descent of time evolution to cohomology (12.79)**.** From $[\hat{H}_{\text{total}}, \hat{Q}_{\text{BRST}}] = 0$ we obtain $U(t)\hat{Q}_{\text{BRST}}U(t)^{-1} = \hat{Q}_{\text{BRST}}$ for $U(t) = e^{-it\hat{H}_{\text{total}}}$, so $U(t)$ maps $\ker \hat{Q}_{\text{BRST}}$ to itself and $\operatorname{im} \hat{Q}_{\text{BRST}}$ to itself. Therefore $U(t)$ induces a well-defined map on the quotient $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$ (Section 12.3, (12.76)). This is precisely the dynamical compatibility condition cited in Definition 12.63(2).

(4) Existence of a cohomological projector commuting with $\hat{H}_{\text{total}}$**.** By the Kugo–Ojima construction [107, 108], there exists a contracting homotopy $\hat{K}$ on the unphysical sector such that $\{\hat{Q}_{\text{BRST}}, \hat{K}\} = \mathbb{I} - \Pi_{\text{phys}}$ with Π_{phys} the orthogonal projector onto $H^0(\hat{Q}_{\text{BRST}})$ (cf. (12.76)). Choose $\hat{K}$ $\hat{H}_{\text{total}}$–equivariant on $\mathcal{D}$ (standard for gauge-fixed quadratic forms and their closures); then

$$[\hat{H}_{\text{total}}, \Pi_{\text{phys}}] = [\hat{H}_{\text{total}}, \mathbb{I} - \{\hat{Q}_{\text{BRST}}, \hat{K}\}] = -\{[\hat{H}_{\text{total}}, \hat{Q}_{\text{BRST}}], \hat{K}\} - \{\hat{Q}_{\text{BRST}}, [\hat{H}_{\text{total}}, \hat{K}]\} = 0,$$

using $[\hat{H}_{\text{total}}, \hat{Q}_{\text{BRST}}] = 0$ from steps (1)–(2). Thus Π_{phys} commutes with $\hat{H}_{\text{total}}$, which is equivalent to (12.79).

(5) Compatibility with the IQR instrumentation and Stage I. Because $\hat{H}_{\text{total}}$ is constructed from the BRST–exact gauge fixing (Definition 12.59), the split transport by $\mathcal{A}_\mu$, and the band–tagged dynamics, the commutation relations proven above hold on certified plateaux (Section 6.12.4) and descend to the BRST cohomology (Lemma 9.54 and Section 9.5). Consequently, channel–resolved visibilities (Definition 6.275) and the spectral poles/residues identifying the sectoral gauge quanta of $\mathcal{A}_\mu$—the *mition* and *tantron* (Definitions 9.258 and 9.348)—are dynamically stable on $\mathcal{H}_{\text{phys}}$. Failure of any of these BRST compatibilities would obstruct the Stage I criterion of the Holosymmetric Selection Principle (Section 12.2 and Remark 12.62), excluding the candidate internal automorphism before Stage II.

These steps establish (12.78) and (12.79), completing the proof. □

Corollary 12.65 (Confinement as cohomological selection)**.** *In QCD,* $\mathcal{H}_{\text{phys}}$ *equals the color–singlet subspace of the transverse quark–gluon Fock space (Theorem 12.133); hence the hadronic space* $\mathcal{H}_{\text{hadron}} \cong \left(\mathbf{3}^{\otimes n} \otimes \bar{\mathbf{3}}^{\otimes \bar{m}} \otimes \mathbf{8}^{\otimes k}\right)^{\mathbf{1}} \otimes \mathcal{H}_{\text{kin}}$ *is invariant under* $U(t)$ *and coincides with the spectral color–singlet filtering (Proposition 12.128).*

Proof. Let $\mathcal{H}_{\text{Fock}}^{\perp}$ denote the *transverse* quark–gluon Fock space after BRST gauge fixing (longitudinal/timelike and ghost sectors paired into BRST quartets, cf. Proposition 12.61

and the Kugo–Ojima mechanism). Write $\mathfrak{su}(3)_C$–charges as $\hat{Q}^a$ and the global color–singlet projector as

$$\Pi_{\mathbf{1}} = \int_{SU(3)} dU\, U \quad \text{(Haar average in the physical representation)}.$$

Step 1: $\mathcal{H}_{\text{phys}}$ is invariant under $U(t)$. By Proposition 12.64 we have

$$[\hat{H}_{\text{total}}, \hat{Q}_{\text{BRST}}] = 0$$

and hence the evolution $U(t) = e^{-it\hat{H}_{\text{total}}}$ descends to cohomology, specifically yielding a map:

$$U(t) : H^0(\hat{Q}_{\text{BRST}}) \to H^0(\hat{Q}_{\text{BRST}}), \quad (12.79)$$

Equivalently, there is a cohomological projector Π_{phys} such that

$$[\hat{H}_{\text{total}}, \Pi_{\text{phys}}] = 0 \quad \text{(ibid.)}$$

Step 2: Color charges are BRST–exact on $\mathcal{H}^{\perp}_{\text{Fock}}$. Gauge fixing is a BRST–exact deformation (Definition 12.59, (12.74)), and the BRST differential acts as a graded inner automorphism generated by $\hat{Q}_{\text{BRST}}$ (Definition 12.58). For QCD (Stage I–admitted, anomaly–free $SU(3)_C$; cf. Ch. 13, Section 12.2 and Remark 12.62), the Noether color charges satisfy the well–known Kugo–Ojima relation

$$\hat{Q}^a \;=\; i\,[\hat{Q}_{\text{BRST}},\, \hat{N}^a\}$$

for suitable local $\hat{N}^a$ built from ghosts and Nakanishi–Lautrup fields (KO construction; see Proposition 12.61 around (12.75)). Therefore $\hat{Q}^a$ acts trivially on cohomology: for any $|\Psi\rangle \in \ker \hat{Q}_{\text{BRST}}$ and any $|\chi\rangle$,

$$\langle\chi|\,\hat{Q}^a\,|\Psi\rangle = i\,\langle\chi|[\hat{Q}_{\text{BRST}}, \hat{N}^a\}|\Psi\rangle = i\,\langle\chi|\hat{Q}_{\text{BRST}}(\hat{N}^a|\Psi\rangle)\rangle \equiv 0 \quad \text{in } H^0(\hat{Q}_{\text{BRST}}).$$

Hence every physical class $[\Psi] \in \mathcal{H}_{\text{phys}}$ is annihilated by all $\hat{Q}^a$ and thus transforms in the trivial $(\mathbf{1})$ representation of $SU(3)_C$.

Step 3: Identification of $\mathcal{H}_{\text{phys}}$ with color singlets. By Theorem 12.133 we have the operator–level equality

$$\mathcal{H}_{\text{phys}} \;\cong\; \left(\mathcal{H}^{\perp}_{\text{Fock}}\right)^{SU(3)_C} = \Pi_{\mathbf{1}}\,\mathcal{H}^{\perp}_{\text{Fock}},$$

i.e. the BRST cohomology equals the global color–singlet subspace of the transverse Fock space. Equivalently, $\Pi_{\text{phys}} = \Pi_{\mathbf{1}} \otimes \Pi_{\text{kin}}$ on $\mathcal{H}^{\perp}_{\text{Fock}} \otimes \mathcal{H}_{\text{kin}}$, where $\mathcal{H}_{\text{kin}}$ collects spin/flavor/kinematical labels.

Step 4: Invariance and equality with spectral singlet filtering. From Step 1 and the commutation $[\hat{H}_{\text{total}}, \Pi_{\text{phys}}] = 0$ we obtain $U(t)\mathcal{H}_{\text{phys}} \subseteq \mathcal{H}_{\text{phys}}$. Using Step 3, $U(t)\Pi_{\mathbf{1}}\mathcal{H}^{\perp}_{\text{Fock}} = \Pi_{\mathbf{1}}U(t)\mathcal{H}^{\perp}_{\text{Fock}}$, i.e. the hadronic space

$$\mathcal{H}_{\text{hadron}} \;\cong\; \left(\mathbf{3}^{\otimes n} \otimes \bar{\mathbf{3}}^{\otimes \bar{m}} \otimes \mathbf{8}^{\otimes k}\right)^{\mathbf{1}} \otimes \mathcal{H}_{\text{kin}}$$

is invariant under $U(t)$. Moreover, Proposition 12.128 identifies the color–singlet projector with the *spectral* singlet filter constructed in the holosymmetric spectral channel (channel–resolved meters; Definition 6.275, with plateau license Section 6.12.4), evaluated in the BRST–physical sector (Ch. 10; Lemma 9.54 and Section 9.5). Thus the cohomological singlet selection and the spectral singlet filtering coincide.

Step 5: Decoupling from gravitational TT channels. Typed/TT projectors ensure that spin–2 gravitational content remains orthogonal to the internal color channel during spectral extraction. Concretely, the *background–covariant TT projector* $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ (cf. Definition 10.3, with explicit form in eq. (10.4)) defines the graviton block via $\hat{h}_{\mu\nu} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}_{\mu\nu}$ (cf. eq. (10.30)), while the τ–kairos fluctuation is TT–null, $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\hat{\tau}_{\mu\nu} = 0$ (cf. Lemma 10.17 and eq. (10.90)). Using the constitutive split $\Sigma_{\mu\nu}[\Phi,\Theta] = g_{\mu\nu}[\Phi] + t_{\mu\nu}[\Phi,\Theta]$ recalled in Chapter 10, Chapter 11 proves the *TT selection rule*

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \circ D_{\Theta}\Sigma \;=\; 0, \qquad \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\Sigma_{\mu\nu} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta g_{\mu\nu},$$

(cf. eqs. (10.106) and (10.134) and the constitutive statement Proposition 10.173). Hence the TT graviton block is sourced solely by tangential Φ–variations and carries no internal (Θ) support. It follows that the internal QCD color channel—read in the spectral meter of Chapter 7 (holosymmetric visibilities Definition 6.275) on certified plateaux (Section 6.12.4)—decouples from the gravitational TT block during BRST–safe spectral extraction; the BRST descent/realization (Ch. 10; Lemma 9.54 and Section 9.5) then applies to Steps 2–4 unchanged in the QCD sector.

Combining Steps 1–4 proves the corollary: $\mathcal{H}_{\mathrm{phys}}$ equals the color–singlet subspace, this space is invariant under $U(t)$ by Proposition 12.64, and it coincides with the spectral color–singlet filtering of Proposition 12.128. □

Universal connection and sectoral coupling. The proof of Corollary 12.65 is carried out within the full IQR dynamics, where the total gauge–fixed *relational* Hamiltonian $\hat{H}_{\mathrm{total}}$ couples both the internal (color) and geometric (spin–2) sectors through the universal Operator Bundle Connection $\mathcal{A}_{\mu}$ (Definition 6.192). This connection governs transport on the modal fibres in *both* substrates (Gaiamiti and Tantrakyma) and enters the split covariant derivatives that build $\hat{H}_{\mathrm{total}}$ (cf. Definitions 6.192 and 9.323, Remark 6.200, and Section 6.5.3). Because color–sector conclusions are drawn using the dynamics generated by $\hat{H}_{\mathrm{total}}$, one must certify that the geometric (spin–2) channel cannot pollute color–channel spectral readouts when passing to BRST cohomology; this is precisely the role of Step 5 in the proof.

BRST descent and holosymmetric meters on the full theory. Time evolution preserves the BRST cohomology by Proposition 12.64, hence $U(t) = e^{-it\hat{H}_{\mathrm{total}}}$ descends to $H^0(\hat{Q}_{\mathrm{BRST}})$ (12.79). Spectral statements are made with the channel–resolved *holosymmetric visibilities* of Chapter 7, evaluated on certified plateaux so that small UV/IR deformations do not alter the verdict (cf. Definition 6.275 and Section 6.12.4). BRST/Nielsen hygiene guarantees that, once restricted to the BRST–physical sector, these band–tagged diagnostics are gauge–parameter independent and compatible with the cohomological quotient (Lemma 9.54 and Section 9.5).

Since these meters act on the *full* theory, a clean separation of the spin–2 geometric content from the internal color channel is required to prevent spurious mixing in the spectral extraction.

Typed/TT projection and the graviton block. Chapter 11 provides the typed, background–covariant York–Lichnerowicz projector $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ that isolates the *graviton* as the spin–2, transverse–traceless (TT) part of the metric fluctuation, $\hat{h}_{\mu\nu} := \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\hat{g}_{\mu\nu}$ (cf. Definition 10.3 and eqs. (10.4) and (10.30)). The companion τ–kairos fluctuation is TT–null, $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\hat{\tau}_{\mu\nu} = 0$ (cf. Lemma 10.17 and eq. (10.90)). Invoking the constitutive split $\Sigma_{\mu\nu} = g_{\mu\nu}[\Phi] + t_{\mu\nu}[\Phi,\Theta]$ recalled in Chapter 10, Chapter 11 proves the *TT selection rule* $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \circ D_{\Theta}\Sigma = 0$ together with $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta\Sigma_{\mu\nu} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\delta g_{\mu\nu}$ (cf. Proposition 10.173 and eqs. (10.106) and (10.134)). Hence the TT graviton block carries no *internal* (Θ) support and is sourced solely by tangential Φ–variations. This establishes the necessary decoupling: color–channel spectra—where one identifies color–singlet poles and applies the global projector $\Pi_{\mathbf{1}}$—are not contaminated by gravitational TT excitations.

Consequence for the confinement corollary. Steps 2–4 of the proof show that $\mathcal{H}_{\mathrm{phys}}$ coincides with the color–singlet subspace and that the *spectral* singlet filter equals the cohomological one. Step 5 provides the final hygiene check in the *coupled* theory governed by $\hat{H}_{\mathrm{total}}$: by projecting out the TT graviton block with $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ and invoking the TT selection rule, one guarantees that the spectral meter in the QCD color channel (evaluated on certified plateaux and realized on $H^0(\hat{Q}_{\mathrm{BRST}})$; cf. Sections 6.12.4 and 9.5 and Lemma 9.54) cannot pick up spurious spin–2 contributions. Therefore the identification of $\mathcal{H}_{\mathrm{phys}}$ with the color–singlet subspace—and its invariance under $U(t)$—remains valid in the presence of dynamical geometry, completing Corollary 12.65.

Remark 12.66 (Exact states and gauge redundancy (cohomological quotient and IQR instrumentation))**.** Vectors in $\operatorname{im}\hat{Q}_{\mathrm{BRST}}$ represent *pure gauge* directions: if $|\Psi\rangle = \hat{Q}_{\mathrm{BRST}}|\chi\rangle$ then, for any BRST–closed observable $\hat{\mathcal{O}}$, one has

$$\langle\Phi_{\mathrm{phys}}|\hat{\mathcal{O}}|\Psi\rangle = \langle\Phi_{\mathrm{phys}}|\hat{\mathcal{O}}\,\hat{Q}_{\mathrm{BRST}}|\chi\rangle = \langle\Phi_{\mathrm{phys}}|[\hat{\mathcal{O}},\hat{Q}_{\mathrm{BRST}}\}|\chi\rangle = 0,$$

because $\delta_{\mathrm{BRST}}\hat{\mathcal{O}} = [\hat{Q}_{\mathrm{BRST}},\hat{\mathcal{O}}\} = 0$ (cf. Definition 12.58). Accordingly, the physical states are *equivalence classes* $[\Psi]$ with $|\Psi\rangle \sim |\Psi\rangle + \hat{Q}_{\mathrm{BRST}}|\chi\rangle$ and

$$\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}}) = \ker\hat{Q}_{\mathrm{BRST}} \cap \ker N_{\mathrm{gh}} \Big/ \operatorname{im}\hat{Q}_{\mathrm{BRST}} \quad \text{(cf. Principle 12.1, (12.76))}.$$

This quotient removes gauge redundancy and ghost contamination, and—by the Kugo–Ojima quartet mechanism—yields a positive–definite inner product supporting a unitary S–matrix on $\mathcal{H}_{\mathrm{phys}}$ (cf. Proposition 12.61).

In the IQR framework, "exact–state" decoupling is implemented consistently with: (i) the *BRST–exact* gauge fixing (Definition 12.59, (12.74)); (ii) the *band/BRST chain–map* so that diagnostics descend to cohomology and are gauge–parameter independent on certified plateaux (Definition 6.282, Lemma 9.54, and Sections 6.12.4 and 9.5); and (iii) the *channel–resolved holosymmetric visibilities* used to evaluate observables (Definition 6.275). For the universal Operator Bundle Connection $\mathcal{A}_{\mu}$ (Definition 6.192 and Remark 6.200), this means that

unphysical polarizations are BRST–exact and drop out, leaving the transverse physical quanta: the *mition* and *tantron* in their sectoral representations (Definitions 9.258 and 9.348), isolated in the spectral channel and realized on $\mathcal{H}_{\text{phys}}$. Typed/TT projectors from Chapter 11 ensure that spin–2 gravitational content remains orthogonal to internal gauge channels during this extraction.

Finally, since $[\hat{H}_{\text{total}}, \hat{Q}_{\text{BRST}}] = 0$ (cf. Proposition 12.64), the cohomological projector Π_{phys} commutes with the relational Hamiltonian and the quotient is dynamically stable: $U(t)$ descends to $H^0(\hat{Q}_{\text{BRST}})$ (12.79). Failure of any of these BRST properties signals an anomaly and violates the Holosymmetric Selection Principle's *Stage I* viability (Section 12.2 and Remark 12.62).

12.3.2 BRST Cohomology and the Holosymmetric Selection Principle

At the foundation of IQR stands the representation–independent *Algebra of Observables* $\mathcal{A}(\mathcal{C}_d)$ (cf. Definition 1.32; see also the context–indexed logical layer in Section 1.2.1 and Definition 1.1). Gauge structure enters as a subgroup $G \subset \text{Aut}(\mathcal{A})$ (connection–theoretic realization emerging from the algebra via GNS; cf. Definitions 1.34 and 6.192 and Remark 6.198), and quantization is effected by adjoining ghost/auxiliary generators to obtain an extended graded $*$–algebra together with a nilpotent, degree–$+1$ derivation (the BRST differential; cf. Definition 12.58). The Holosymmetric Selection Principle (HSP) then acts *on the BRST–physical sector* to select the realized gauge law (Stage I viability and subsequent selection; cf. Section 12.2). Operationally, all readouts are taken on certified plateaux with the band/transport structure induced by the universal Operator Bundle Connection $\mathcal{A}_\mu$ (Remark 6.200, Sections 6.12.2 and 6.12.4, and Definition 6.282), and are ensured to be BRST/Nielsen–safe by the chain–map and realization results (Lemma 9.54 and Section 9.5). Typed/TT separations used when combining gravitational with internal channels follow Chapter 11's background–covariant construction (Definition 10.3, eqs. (10.4), (10.30), (10.90), (10.106) and (10.134), Lemma 10.17, and Proposition 10.173).

1. Algebraic BRST structure and degree–zero cohomology

Definition 12.67 (Abstract BRST $*$–algebra and degree–zero cohomology)**.** For a candidate $G \subset \text{Aut}(\mathcal{A})$ let $\mathcal{A}^{(G)}_{\text{BRST}}$ be the $\mathbb{Z}$–graded $*$–algebra generated by $\mathcal{A}$ and ghost/antighost/auxiliary symbols (cf. Definition 12.58), equipped with a graded $*$–derivation δ_{BRST} of ghost number $+1$ obeying $\delta^2_{\text{BRST}} = 0$. Its *degree–zero BRST cohomology* is

$$H^0\big(\mathcal{A}^{(G)}_{\text{BRST}}, \delta_{\text{BRST}}\big) := \frac{\ker(\delta_{\text{BRST}}|_{\text{deg}=0})}{\text{im}(\delta_{\text{BRST}}|_{\text{deg}=-1})} =: \mathcal{A}^{(G)}_{\text{phys}},$$

a $*$–algebra (the product and $*$ descend because δ_{BRST} is a graded derivation). In practice, this abstract construction is *realized* on the emergent/GNS Hilbert space $\mathcal{H}_{\text{em}}$ associated to the Quantum Holo–State (cf. Definition 1.34) and then projected to the BRST–physical sector of the dynamical theory, $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$ (cf. (12.76)). When the sector carries internal type, transport and band–restriction are implemented by the universal connection $\mathcal{A}_\mu$ and the band functor (Remark 6.200, Section 6.12.2, and Definition 6.282), and BRST/Nielsen hygiene guarantees gauge–parameter independence of physical correlators on certified plateaux

(Lemma 9.54 and Section 9.5). Logical assertions about existence/selection that refer to this physical algebra are to be read relative to their immanent context as specified by the tetralemma layer (cf. Section 1.2.1 and Definition 1.1).

Postulate 12.2 (BRST–closed vacuum functional)**.** *Let $\omega^{\mathring{\mathbf{\Psi}}}$ be the Quantum Holo–State on $\mathcal{A}(\mathcal{C}_d)$ (cf. Definition 1.34 together with Definition 1.32). For each candidate gauge subgroup $G \subset \mathrm{Aut}(\mathcal{A})$, extend $\omega^{\mathring{\mathbf{\Psi}}}$ to the abstract BRST $*$–algebra $\mathcal{A}^{(G)}_{\mathrm{BRST}}$ (cf. Definition 12.67) and denote the extension by $\tilde{\omega}$. We postulate that $\tilde{\omega}$ is* BRST–closed*:*

$$\tilde{\omega}\Big(\delta_{\mathrm{BRST}}(X)\Big) = 0 \qquad \forall\, X \in \mathcal{A}^{(G)}_{\mathrm{BRST}},$$

where δ_{BRST} is the nilpotent graded derivation of Definition 12.58. Then $\tilde{\omega}$ factors through the quotient map $q : \ker \delta_{\mathrm{BRST}} \twoheadrightarrow \mathcal{A}^{(G)}_{\mathrm{phys}}$ of Definition 12.67, defining a positive linear functional $\omega^{(G)}_{\mathrm{phys}} := \tilde{\omega} \circ q^{-1}$ on the degree–zero BRST cohomology algebra $\mathcal{A}^{(G)}_{\mathrm{phys}}$.

Remarks. *(1) By the GNS construction (cf. Definition 1.34), $\omega^{(G)}_{\mathrm{phys}}$ yields a concrete representation $(\mathcal{H}^{(G)}_{\mathrm{phys}}, \pi^{(G)}_{\mathrm{phys}}, |\Omega^{(G)}_{\mathrm{phys}}\rangle)$ that realizes the abstract physical algebra $\mathcal{A}^{(G)}_{\mathrm{phys}}$. This representation is compatible with the operator–level BRST construction of Section 12.3, where the physical Hilbert space is identified as $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ (cf. (12.76), Proposition 12.61).*

(2) In the IQR setting, transport and coarse–graining are implemented using the universal Operator Bundle Connection and the band functor (cf. Remark 6.200, Section 6.12.2, and Definition 6.282). BRST/Nielsen hygiene on certified plateaux guarantees that the induced correlators computed with $\omega^{(G)}_{\mathrm{phys}}$ are gauge–parameter independent and descend to cohomology (Ch. 10; cf. Lemma 9.54 and Section 9.5, and Section 6.12.4).

(3) As required by the Holosymmetric Selection Principle, Stage I acts on *the BRST–physical sector (cf. Section 12.2): if a candidate G fails BRST closure (e.g. an anomaly obstructs δ_{BRST}–invariance of $\tilde{\omega}$), then the selection functional excludes G at Stage I (cf. Remark 12.62).*

Proposition 12.68 (Equivalence of algebraic and Hilbert–space cohomologies)**.** *Consider the GNS triple $(\pi, \mathcal{H}_{\mathrm{total}}, |\Omega\rangle)$ of $\tilde{\omega}$ on $\mathcal{A}^{(G)}_{\mathrm{BRST}}$ (cf. Definitions 1.34 and 12.67 and Postulate 12.2), and let $\hat{Q}_{\mathrm{BRST}}$ be the densely defined, nilpotent operator implementing δ_{BRST} by graded commutators (Section 12.3, Definition 12.58, and Proposition 12.61). Then*

$$\mathrm{GNS}\Big(\mathcal{A}^{(G)}_{\mathrm{phys}}, \omega^{(G)}_{\mathrm{phys}}\Big) \;\simeq\; \Big(H^0(\hat{Q}_{\mathrm{BRST}}),\ \pi|_{H^0},\ [|\Omega\rangle]\Big). \tag{12.80}$$

holds.

$$\mathrm{GNS}\Big(\mathcal{A}^{(G)}_{\mathrm{phys}}, \omega^{(G)}_{\mathrm{phys}}\Big) \;\simeq\; \Big(H^0(\hat{Q}_{\mathrm{BRST}}),\ \pi|_{H^0},\ [|\Omega\rangle]\Big), \tag{12.81}$$

i.e. the GNS representation of the algebraic *degree–zero cohomology is unitarily equivalent to the* Hilbert–space *BRST cohomology representation (Principle 12.1, (12.76)) [107, 108].*

BRST–closedness of $\tilde{\omega}$ implies $\tilde{\omega}$ annuls $\mathrm{im}\,\delta_{\mathrm{BRST}}$ so it factors through $\mathcal{A}^{(G)}_{\mathrm{phys}}$. The cyclic subspace generated from $|\Omega\rangle$ by $\pi(\ker \delta_{\mathrm{BRST}})$ is invariant under $\hat{Q}_{\mathrm{BRST}}$; quotienting by $\mathrm{im}\,\hat{Q}_{\mathrm{BRST}}$ yields $H^0(\hat{Q}_{\mathrm{BRST}})$, and the natural map from $\mathcal{A}^{(G)}_{\mathrm{phys}}$ to bounded operators on H^0 is faithful. Standard Kugo–Ojima arguments supply the unitary identification.

Proof. **Step 0 (setup and references).** For a fixed candidate $G \subset \mathrm{Aut}(\mathcal{A})$, let $\mathcal{A}^{(G)}_{\mathrm{BRST}}$ be the abstract BRST $*$–algebra with differential δ_{BRST} (Definitions 12.58 and 12.67). By the BRST–closed vacuum postulate (Postulate 12.2), $\tilde{\omega}$ factors through the algebraic degree–zero cohomology $\mathcal{A}^{(G)}_{\mathrm{phys}} := H^0(\mathcal{A}^{(G)}_{\mathrm{BRST}}, \delta_{\mathrm{BRST}})$, producing $\omega^{(G)}_{\mathrm{phys}}$. Let $(\pi, \mathcal{H}_{\mathrm{total}}, |\Omega\rangle)$ be the GNS triple of $\tilde{\omega}$ on $\mathcal{A}^{(G)}_{\mathrm{BRST}}$ (cf. Definition 1.34), and let $\hat{Q}_{\mathrm{BRST}}$ implement δ_{BRST} as $\delta_{\mathrm{BRST}}(\cdot) = [\hat{Q}_{\mathrm{BRST}}, \cdot\}$ with $\hat{Q}^2_{\mathrm{BRST}} = 0$ (Section 12.3 and Proposition 12.61). We take the physical Hilbert space to be the BRST cohomology $H^0(\hat{Q}_{\mathrm{BRST}})$ as in (12.76).

Step 1 (algebraic factorization and the physical GNS pair). By Postulate 12.2, $\tilde{\omega}(\delta_{\mathrm{BRST}} X) = 0$ for all $X \in \mathcal{A}^{(G)}_{\mathrm{BRST}}$, hence $\tilde{\omega}$ annihilates $\mathrm{im}(\delta_{\mathrm{BRST}}|_{\mathrm{deg}=-1})$ and descends to a positive linear functional $\omega^{(G)}_{\mathrm{phys}}$ on $\mathcal{A}^{(G)}_{\mathrm{phys}}$. Applying Definition 1.34 to $(\mathcal{A}^{(G)}_{\mathrm{phys}}, \omega^{(G)}_{\mathrm{phys}})$ yields a GNS triple $(\pi_{\mathrm{phys}}, \mathcal{H}^{\mathrm{alg}}_{\mathrm{phys}}, |\Omega_{\mathrm{phys}}\rangle)$.

Step 2 (construction of the intertwiner V). Consider the dense subspace

$$\mathcal{D}_0 := \mathrm{span}\Big\{\ \pi(A)\,|\Omega\rangle\ :\ A \in \ker(\delta_{\mathrm{BRST}}|_{\mathrm{deg}=0})\ \Big\}\ \subset\ \mathcal{H}_{\mathrm{total}}.$$

Define a linear map $V : \mathcal{D}_0 \to H^0(\hat{Q}_{\mathrm{BRST}})$ by

$$V\Big(\pi(A)|\Omega\rangle\Big)\ :=\ \Big[\ \pi(A)|\Omega\rangle\ \Big].$$

This is well–defined on cohomology classes: if $A' = A + \delta_{\mathrm{BRST}} B$ with B of degree -1, then

$$\pi(A')|\Omega\rangle - \pi(A)|\Omega\rangle = \pi(\delta_{\mathrm{BRST}} B)|\Omega\rangle = [\hat{Q}_{\mathrm{BRST}}, \pi(B)]\,|\Omega\rangle = \hat{Q}_{\mathrm{BRST}}(\pm\,\pi(B)|\Omega\rangle)\ \in\ \mathrm{im}\,\hat{Q}_{\mathrm{BRST}},$$

so $[\pi(A')|\Omega\rangle] = [\pi(A)|\Omega\rangle]$ in $H^0(\hat{Q}_{\mathrm{BRST}})$. (Here we used the operator implementation of Definition 12.58 and $\hat{Q}^2_{\mathrm{BRST}} = 0$, Proposition 12.61.) Moreover, by Postulate 12.2 and standard Kugo–Ojima normalization, $|\Omega\rangle$ can be chosen BRST–invariant, $\hat{Q}_{\mathrm{BRST}}|\Omega\rangle = 0$ [107, 108], ensuring V is defined on the cyclic domain generated by $\pi(\ker \delta_{\mathrm{BRST}})|\Omega\rangle$.

Step 3 (isometry property of V). For $A, B \in \ker(\delta_{\mathrm{BRST}}|_{\mathrm{deg}=0})$,

$$\langle V(\pi(A)|\Omega\rangle),\, V(\pi(B)|\Omega\rangle)\rangle_{H^0} = \langle\, [\pi(A)|\Omega\rangle]\,,\, [\pi(B)|\Omega\rangle]\,\rangle = \tilde{\omega}(A^* B),$$

because the inner product on $H^0(\hat{Q}_{\mathrm{BRST}})$ is inherited from $\mathcal{H}_{\mathrm{total}}$ after quotienting by $\mathrm{im}\,\hat{Q}_{\mathrm{BRST}}$ (Principle 12.1, (12.76)). Since $\tilde{\omega}$ factors through $q : \ker \delta_{\mathrm{BRST}} \twoheadrightarrow \mathcal{A}^{(G)}_{\mathrm{phys}}$ (Postulate 12.2), $\tilde{\omega}(A^* B) = \omega^{(G)}_{\mathrm{phys}}([A]^*[B])$, which is precisely the inner product in the GNS space $\mathcal{H}^{\mathrm{alg}}_{\mathrm{phys}}$. Thus V is isometric from the cyclic subspace generated by $\pi(\ker \delta_{\mathrm{BRST}})|\Omega\rangle$ onto a dense subspace of $H^0(\hat{Q}_{\mathrm{BRST}})$.

Step 4 (intertwining the representations). Define the algebra–side representation on cohomology by

$$\pi_{\mathrm{phys}}([A])\,[\psi]\ :=\ [\,\pi(A)\psi\,],\qquad [A] \in \mathcal{A}^{(G)}_{\mathrm{phys}},\quad [\psi] \in H^0(\hat{Q}_{\mathrm{BRST}}).$$

This is well–defined: if $A \sim A + \delta_{\mathrm{BRST}} B$, then $\pi(\delta_{\mathrm{BRST}} B)\psi = [\hat{Q}_{\mathrm{BRST}}, \pi(B)\}\psi \in \mathrm{im}\,\hat{Q}_{\mathrm{BRST}}$ whenever $[\psi] \in H^0(\hat{Q}_{\mathrm{BRST}})$, hence vanishes in the quotient. Moreover, π_{phys} is a $*$–homomorphism because δ_{BRST} is a graded derivation and the product/$*$ descend to cohomology

(Definition 12.67). By construction,

$$\begin{aligned} V\,\pi_{\mathrm{phys}}([A])\,[\pi(B)|\Omega\rangle] &= V\,[\pi(AB)|\Omega\rangle] \\ &= [\pi(AB)|\Omega\rangle] \\ &= \pi(A)\,[\pi(B)|\Omega\rangle] \\ &= \pi|_{H^0}([A])\,V\,[\pi(B)|\Omega\rangle], \end{aligned}$$

so V intertwines π_{phys} and $\pi|_{H^0}$ on a dense set. Thus V extends uniquely to a unitary map $V : \mathcal{H}_{\mathrm{phys}}^{\mathrm{alg}} \xrightarrow{\simeq} H^0(\hat{Q}_{\mathrm{BRST}})$ with $V|\Omega_{\mathrm{phys}}\rangle = [|\Omega\rangle]$ and $V\,\pi_{\mathrm{phys}}([A]) = \pi|_{H^0}([A])\,V$.

Step 5 (conclusion and compatibility with IQR instrumentation). We have shown that the GNS representation of $(\mathcal{A}_{\mathrm{phys}}^{(G)}, \omega_{\mathrm{phys}}^{(G)})$ is unitarily equivalent to the BRST–cohomology representation $(H^0(\hat{Q}_{\mathrm{BRST}}), \pi|_{H^0}, [|\Omega\rangle])$, proving (12.81). In the IQR setting, all operators involved can be taken band–tagged and evaluated on certified plateaux (Definition 6.282 and Section 6.12.4); the *chain–map* property guarantees that both sides descend to cohomology in a gauge–parameter–independent way (Lemma 9.54 and Section 9.5). This is the precise algebra/geometry bridge that underlies the Stage I action of the Holosymmetric Selection Principle on the BRST–physical sector (Section 12.2). □

Physically relevant projector. By Proposition 12.64 there exists a cohomological (orthogonal) projector $\Pi_{\mathrm{phys}}^{(G)}$ on $\mathcal{H}_{\mathrm{total}}$ such that $\left[\hat{H}_{\mathrm{total}}, \Pi_{\mathrm{phys}}^{(G)}\right] = 0$ and the Schrödinger evolution $U(t) = e^{-it\hat{H}_{\mathrm{total}}}$ descends to $H^0(\hat{Q}_{\mathrm{BRST}})$ (cf. (12.79), (12.76)). Moreover, for every BRST–closed observable $\hat{\mathcal{O}}$ one has the compression identity $\Pi_{\mathrm{phys}}^{(G)}\hat{\mathcal{O}}\Pi_{\mathrm{phys}}^{(G)} = \hat{\mathcal{O}}\Pi_{\mathrm{phys}}^{(G)} = \Pi_{\mathrm{phys}}^{(G)}\hat{\mathcal{O}}$, since $\hat{\mathcal{O}}$ acts on cohomology classes modulo exact terms (*BRST–closed observables and operator cohomology*, Definition 12.63; see also Proposition 12.61 and Section 12.3). We therefore set

$$\mathcal{H}_{\mathrm{phys}}^{(G)} := \Pi_{\mathrm{phys}}^{(G)}\,\mathcal{H}_{\mathrm{total}}, \qquad \mathbf{U}_{\mathrm{phys}}^{(G)}(t) := \Pi_{\mathrm{phys}}^{(G)}\,U^{(G)}(t)\,\Pi_{\mathrm{phys}}^{(G)},$$

so that $\mathbf{U}_{\mathrm{phys}}^{(G)}(t)$ is a strongly continuous unitary group on $\mathcal{H}_{\mathrm{phys}}^{(G)}$ implementing the descended evolution on BRST cohomology (Proposition 12.64). In the algebra/geometry correspondence this projector agrees with the GNS realization of the algebraic degree–zero cohomology (Proposition 12.68) built from the BRST–closed vacuum functional (Postulate 12.2), and its spectral use in physical meters is licensed on certified plateaux (Section 6.12.4) with BRST/Nielsen hygiene (Lemma 9.54 and Section 9.5).

2. HSP on the BRST–physical sector

Definition 12.69 (Physical self–coherence and overlap)**.** For a candidate G and a (possibly mixed) probe ρ on $\mathcal{H}_{\mathrm{phys}}^{(G)}$ define the *physical self–coherence*

$$\mathcal{C}_{\mathrm{phys}}(G \mid \rho) := \left| \mathrm{Tr}\Big(\rho\,(\mathbf{U}_{\mathrm{phys}}^{(G)})^{\dagger}\mathbf{U}_{\mathrm{phys}}^{(G)}\Big) \right| \le 1, \tag{12.82}$$

and, for two viable candidates G_A, G_B, the *physical holosymmetric overlap*

$$\Omega_{\mathrm{phys}}(A, B \mid \Psi_0) := \langle\Psi_0|(\mathbf{U}_{\mathrm{phys}}^{(B)})^{\dagger}\mathbf{U}_{\mathrm{phys}}^{(A)}|\Psi_0\rangle, \tag{12.83}$$

with $|\Psi_0\rangle$ supported on $\mathcal{H}_{\mathrm{phys}}^{(A)} \cap \mathcal{H}_{\mathrm{phys}}^{(B)}$ (after the canonical identification from (12.81)).

Remark 12.70 (Physical self–coherence, holosymmetric overlap, and the BRST probe)**.** *Stage I (self–coherence).* For a candidate gauge law G, the compressed arm block $\mathbf{U}^{(G)}_{\mathrm{phys}} := \Pi_{\mathrm{phys}}\,\mathbf{U}^{(G)}\,\Pi_{\mathrm{phys}}$ acts on $\mathcal{H}^{(G)}_{\mathrm{phys}}$. The quantity

$$\mathcal{C}_{\mathrm{phys}}(G \mid \rho) = \left|\,\mathrm{Tr}\Big(\rho\,(\mathbf{U}^{(G)}_{\mathrm{phys}})^{\dagger}\mathbf{U}^{(G)}_{\mathrm{phys}}\Big)\right| \le 1 \quad (\textit{eq. (12.82)})$$

is a Stage I sanity check *after* descent to cohomology: G is Stage I viable iff $\mathcal{C}_{\mathrm{phys}}(G \mid \rho) = 1$ for an informationally complete family of probes on $\mathcal{H}^{(G)}_{\mathrm{phys}}$, equivalently, the $\mathbf{U}^{(G)}_{\mathrm{phys}}(t)$ form a unitary group on the physical subspace.

Stage II (relational comparison). For two *viable* candidates G_A, G_B, the physical holosymmetric overlap

$$\Omega_{\mathrm{phys}}(A, B \mid \Psi_0) = \langle\Psi_0|(\mathbf{U}^{(B)}_{\mathrm{phys}})^{\dagger}\mathbf{U}^{(A)}_{\mathrm{phys}}|\Psi_0\rangle \quad (\textit{eq. (12.83)})$$

is the physical–sector amplitude for "do A then undo B" on the *same* probe. In the asymmetric dual–arm interferometer the outcome probabilities obey

$$p_{\pm}(A, B) = \tfrac{1}{2}(1 \pm \mathrm{Re}\ \Omega_{\mathrm{phys}}(A, B \mid \Psi_0)) \quad (\textit{eq. (13.48)}),$$

so the fringe visibility and the phase–calibrated coherence score are monotone in $\mathrm{Re}\ \Omega_{\mathrm{phys}}$ (or in $|\Omega_{\mathrm{phys}}|$ after phase compensation; eq. (13.49)). The Stage II selection rule therefore realises the *interferometric argmax*: the realised law $G_\star$ maximises its diagonal self–visibility, $\Omega_{\mathrm{phys}}(G_\star, G_\star \mid \Psi_0)$, and is orthogonal to any inequivalent competitor, $\Omega_{\mathrm{phys}}(G_\star, B \mid \Psi_0) = 0$ for all $B \neq G_\star$ (Proposition 12.76; cf. also eq. (13.82) in Ch. 13).

The BRST probe $|\Psi_0\rangle$. The interferometric probe is taken *BRST–closed* on the dual–register physical space so that all primitives descend unchanged to cohomology. In particular, after the canonical identification between algebraic and operator cohomology (eq. (12.81)),

$$\mathrm{supp}(|\Psi_0\rangle) \subset \mathcal{H}^{(A)}_{\mathrm{phys}} \cap \mathcal{H}^{(B)}_{\mathrm{phys}},$$

ensuring that cross–candidate comparisons are made on the common physical stage.

Equivalence to the holo–state. At the holosymmetric/GNS layer (tetralemma–GNS), the cyclic holo–state $\mathring{\mathbf{\Psi}}$ (built from admissible effect–families; see Definition 1.16 and §2.4) pushes forward to $|\Psi_0\rangle$ under the same identification, and the overlaps coincide:

$$\Omega_{\mathrm{phys}}(A, B \mid \mathring{\mathbf{\Psi}}) = \langle\Psi_0|(\mathbf{U}^{(B)}_{\mathrm{phys}})^{\dagger}\mathbf{U}^{(A)}_{\mathrm{phys}}|\Psi_0\rangle.$$

Thus, all Stage II readouts are representation–independent across the physical and holosymmetric layers.

Relation to tetralemma meters and Ch. 2 controls. The local Thesis/Antithesis/Synthesis meters used in preparations and readouts are the orthoprojectors $P_T, P_A, P_{\mathrm{para}}$ with Born semantics (tetralemma–GNS; Definition 1.16), realised concretely on the logical sector by the Ch. 2 constructions $\big(P^{(\mathfrak{B}_x)}_{\mathrm{para}},\ P_T,\ P_A,\ D^{(\mathfrak{B}_x)}\big)$ (Definitions 2.7 and 2.17). Compression to $\mathcal{H}_{\mathrm{phys}}$ leaves their interferometric statistics unchanged.

Definition 12.71 (Fujikawa BRST measure Jacobian and anomaly class)**.** Let $\delta_{\mathrm{BRST}}(\,\cdot\,) = [\hat{Q}_{\mathrm{BRST}}, \cdot\,\}$ be the BRST derivation on fields (cf. Definition 12.58 and Section 12.3). Under an infinitesimal BRST variation, the functional measure transforms as

$$\delta_{\mathrm{BRST}} \ln \mathcal{D}\Phi \;=\; \Delta_{\mathrm{meas}} \;=\; \int d^d x\, \mathcal{A}(x), \tag{12.84}$$

with a local density $\mathcal{A}(x)$ whose cohomology class $[\mathcal{A}] \in H^{+1}(\hat{Q}_{\mathrm{BRST}})$ is called the *BRST (Fujikawa) anomaly class*. BRST measure invariance is equivalent to $[\mathcal{A}] = 0$ (up to local counterterms), in which case the Jacobian can be absorbed by a BRST–exact deformation of the action (cf. Definition 12.59).

Definition 12.72 (Holosymmetric Selection Principle — Stage I (viability on the BRST–physical sector))**.** Given a candidate internal automorphism group $G \subset \mathrm{Aut}(\mathcal{A})$ acting through the universal Operator Bundle Connection $\mathcal{A}_\mu$ (cf. Definition 6.192 and Remark 6.200) and the sectoral covariant derivatives on the substrate fields (cf. Section 6.5.3 and Definition 9.323), *Stage I* declares *G viable* iff the following hold:

1. **BRST structure exists and is nilpotent.** There is a graded derivation δ_{BRST} with $\delta_{\mathrm{BRST}}^2 = 0$ acting on the extended field algebra (cf. Definition 12.58 and Section 12.3), implemented by an operator $\hat{Q}_{\mathrm{BRST}}$ obeying (12.75).
2. **Gauge fixing is BRST exact.** The gauge–fixing+ghost sector is a BRST–exact deformation $\mathcal{L}_{\mathrm{gf+gh}} = \delta_{\mathrm{BRST}} \Psi_{\mathrm{gf}}$ (cf. Definition 12.59, (12.74)), so that gauge–parameter changes leave BRST–closed observables invariant.
3. **Measure is BRST–invariant (Fujikawa).** The Fujikawa Jacobian has trivial BRST class, $\Delta_{\mathrm{meas}} = 0$ and $[\mathcal{A}] = 0$ (cf. Definition 12.71 and Proposition 12.74); otherwise conservation in (12.75) fails and the candidate is excluded.
4. **Physical sector exists and dynamics descend.** The physical Hilbert space $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ is well-defined (cf. (12.76), Proposition 12.61) and the relational evolution $U(t) = e^{-it\hat{H}_{\mathrm{total}}}$ descends to cohomology (cf. Proposition 12.64, (12.79)); equivalently, there is a cohomological projector Π_{phys} commuting with $\hat{H}_{\mathrm{total}}$.
5. **Band/transport hygiene on certified plateaux.** The band functor and holosymmetric meters (Chs. 7–8) commute with BRST (chain–map), so diagnostics descend to cohomology and are gauge–parameter independent on certified plateaux (cf. Definitions 6.275 and 6.282, Sections 6.12.4 and 9.5, and Lemma 9.54).
6. **Algebra/Hilbert–space equivalence.** The BRST–closed vacuum functional factors to the algebraic degree–zero cohomology and its GNS representation coincides with the Hilbert–space BRST cohomology (cf. Postulate 12.2, Definition 12.67, and Proposition 12.68, (12.81)).

If (1)–(6) hold, the Stage I holosymmetric score attains unity ($\mathcal{C}(A) = 1$) and the candidate passes viability (cf. Section 12.2 and Remark 12.62). Logical assertions of selection are read *relative to context* in the tetralemma layer (cf. Section 1.2.1 and Definition 1.1).

Definition 12.73 (Holosymmetric Selection Principle — Stage II (selection on gauge–invariant composites))**.** Assuming Stage I viability, *Stage II* acts on BRST–invariant composites and correlators evaluated in the physical sector. Concretely:

1. **Meters and spectra.** Use the holosymmetric, channel–resolved meters (Definition 6.275) in the spectral channel Π_{spec} on certified plateaux (Section 6.12.4), realized on $\mathcal{H}_{\mathrm{phys}}$ via BRST descent (Lemma 9.54 and Section 9.5).
2. **Sectoral gauge quanta of the universal connection.** Excitations of $\rho_\Phi(\mathcal{A}_\mu)$ and $\rho_\Theta(\mathcal{A}_\mu)$ define, respectively, the *mition* and *tantron* (Definitions 9.258 and 9.348), whose physical poles/residues are read as transverse typed vectors in Π_{spec} and identified in $H^0(\hat{Q}_{\mathrm{BRST}})$.
3. **Typed/TT separation with geometry.** When combining internal channels with geometry, apply the background–covariant TT projector to isolate the spin–2 graviton block and enforce the TT selection rule (cf. Definition 10.3, eqs. (10.4), (10.30), (10.90), (10.106) and (10.134), Lemma 10.17, and Proposition 10.173). This guarantees that internal readouts (e.g. color singlets in QCD) are not contaminated by spin–2 content.
4. **Algebraic realization.** On the algebra side, evaluate composites in $\mathcal{A}^{(G)}_{\mathrm{phys}}$ and represent them by GNS on $\mathcal{H}_{\mathrm{phys}}$ (cf. Definition 12.67, Postulate 12.2, and Proposition 12.68).

Stage II thus *selects* the realized gauge law by matching the BRST–safe, plateau–licensed spectra/composites against the holosymmetric criteria (cf. Section 12.2), with context-indexing supplied by the tetralemma layer (cf. Section 1.2.1 and Definition 1.1).

Proposition 12.74 (Stage I consistency via Fujikawa)**.** *If* $[\mathcal{A}] \neq 0$ *in* (12.84)*, then the Ward identity acquires an anomalous term and the Noether current and charge of Proposition 12.61 fail to be conserved:*

$$\partial_\mu j^\mu_{\mathrm{BRST}} = \mathcal{A}(x) \quad \Longrightarrow \quad \frac{d}{dt}\hat{Q}_{\mathrm{BRST}} \neq 0,$$

so (12.75) *is violated. Hence the BRST cohomology* $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ *((12.76)) is ill–defined as a physical state space, and the candidate gauge law is excluded by the Holosymmetric Selection Principle at* Stage I *(cf. Section 12.2 and Remark 12.62). Conversely, if* $[\mathcal{A}] = 0$*, then the BRST–closed vacuum functional exists (Postulate 12.2),* $\mathrm{GNS}(\mathcal{A}^{(G)}_{\mathrm{phys}}, \omega^{(G)}_{\mathrm{phys}})$ *is well defined and agrees with the Hilbert–space cohomology representation (cf. Proposition 12.68), and the relational evolution descends to cohomology (cf. Proposition 12.64).*

Proof. **Step 1 (Fujikawa Jacobian and the anomalous Ward identity).** Let $S_{\mathrm{tot}} = \int d^dx\, \mathcal{L}$ be the gauge–fixed action with $\mathcal{L} = \mathcal{L}_{\mathrm{YM}} + \mathcal{L}_{\mathrm{matter}} + \mathcal{L}_{\mathrm{gf+gh}}$ and $\mathcal{L}_{\mathrm{gf+gh}} = \delta_{\mathrm{BRST}}\Psi_{\mathrm{gf}}$ (Definition 12.59, (12.74)). Under an infinitesimal BRST change of variables $\Phi \mapsto \Phi + \epsilon\, \delta_{\mathrm{BRST}}\Phi$ with Grassmann ϵ, the classical action changes by a total derivative (BRST symmetry of the Lagrangian, Proposition 12.61), while the functional measure picks up the Fujikawa Jacobian $\delta_{\mathrm{BRST}} \ln \mathcal{D}\Phi = \Delta_{\mathrm{meas}} = \int d^dx\, \mathcal{A}(x)$ (Definition 12.71, (12.84)). Thus the generating functional acquires an anomalous phase and the Ward identity becomes

$$\left\langle \delta_{\mathrm{BRST}}\mathcal{O} \right\rangle = i \left\langle \mathcal{O} \int d^dx\, \mathcal{A}(x) \right\rangle \tag{12.85}$$

for all insertions $\mathcal{O}$ built from the extended field algebra (cf. Definition 12.58 and Section 12.3).

Step 2 (Local form: divergence of the BRST current). Choosing $\mathcal{O}$ to be the unit operator and/or inserting local sources and then setting them to zero gives the local anomalous Ward identity

$$\partial_\mu j^\mu_{\mathrm{BRST}}(x) = \mathcal{A}(x), \tag{12.86}$$

where j^μ_{BRST} is the Noether current whose existence follows from the BRST invariance of the classical action (Proposition 12.61). Integrating over a Cauchy slice and assuming standard boundary conditions yields

$$\frac{d}{dt}\hat{Q}_{\mathrm{BRST}} = \int_{\mathbb{R}^{d-1}} d^{d-1}\mathbf{x}\ \mathcal{A}(t,\mathbf{x}), \tag{12.87}$$

so a nontrivial anomaly class $[\mathcal{A}] \neq 0$ in (12.84) implies $\frac{d}{dt}\hat{Q}_{\mathrm{BRST}} \neq 0$ and hence

$$[\hat{H}_{\mathrm{total}}, \hat{Q}_{\mathrm{BRST}}] \neq 0, \tag{12.88}$$

violating the conservation relation in (12.75).

Step 3 (Breakdown of the BRST physical sector). If (12.88) holds, $\ker \hat{Q}_{\mathrm{BRST}}$ is not invariant under time evolution and the quotient $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ ((12.76)) is not dynamically well defined: the Schrödinger group $U(t) = e^{-it\hat{H}_{\mathrm{total}}}$ cannot descend to cohomology, contradicting Proposition 12.64. Equivalently, the cohomological projector Π_{phys} commuting with $\hat{H}_{\mathrm{total}}$ does not exist. Therefore the candidate violates the Stage I viability condition of the Holosymmetric Selection Principle (Section 12.2 and Remark 12.62).

Step 4 (Algebraic side: failure of the BRST–closed vacuum functional). A nontrivial Fujikawa class $[\mathcal{A}] \neq 0$ implies that the extended vacuum functional $\tilde{\omega}$ is *not* BRST closed, i.e. $\tilde{\omega}(\delta_{\mathrm{BRST}}X) = \langle \delta_{\mathrm{BRST}}X \rangle \neq 0$ for some X, by (12.85). Hence Postulate 12.2 fails: $\tilde{\omega}$ does not factor to the degree–zero cohomology $\mathcal{A}^{(G)}_{\mathrm{phys}}$ (Definition 12.67), so the algebraic physical state $\omega^{(G)}_{\mathrm{phys}}$ and its GNS representation cannot be constructed. This mirrors the breakdown on the Hilbert–space side in Step 3 and confirms the Stage I veto.

Step 5 (Conversely: trivial class restores Stage I). If $[\mathcal{A}] = 0$, then there exists a local counterterm that cancels the Jacobian, rendering the measure BRST invariant (Definition 12.71). The classical BRST symmetry is then promoted to a *quantum* symmetry: $\partial_\mu j^\mu_{\mathrm{BRST}} = 0$ and $[\hat{H}_{\mathrm{total}}, \hat{Q}_{\mathrm{BRST}}] = 0$ (Propositions 12.61 and 12.64), so $U(t)$ descends to cohomology as in (12.79). On the algebraic side, Postulate 12.2 holds, and the GNS representation of the algebraic cohomology $\mathrm{GNS}(\mathcal{A}^{(G)}_{\mathrm{phys}}, \omega^{(G)}_{\mathrm{phys}})$ agrees with the Hilbert–space cohomology representation (Proposition 12.68, (12.81)). In the IQR instrumentation, all diagnostics are then evaluated on certified plateaux with the band/transport structure and are BRST/Nielsen–safe (Sections 6.12.4 and 9.5, Definition 6.282, and Lemma 9.54), fulfilling Stage I (Section 12.2).

Combining Steps 1–4 proves the forward implication; Step 5 proves the converse, establishing Proposition 12.74. □

Remark. When gravitational and internal channels are present, the typed/TT separation of Chapter 11 ensures that spin–2 (graviton) TT content does not contaminate internal BRST meters used in anomaly tests (cf. Definition 10.3, eqs. (10.4), (10.30), (10.90), (10.106) and (10.134), Lemma 10.17, and Proposition 10.173); anomaly checks are thus carried out on certified plateaux with band/transport structure and BRST/Nielsen hygiene (Chs. 7–10; Sections 6.12.4 and 9.5, Definition 6.282, and Lemma 9.54).

Proposition 12.75 (Stage I on the physical sector)**.** $\mathcal{C}_{\mathrm{phys}}(G \mid \rho) = 1$ *for an informationally complete family of* ρ *(Definition 12.33 and Proposition 12.35) iff* $\mathbf{U}^{(G)}$ *is unitary on* $\mathcal{H}^{(G)}_{\mathrm{phys}}$*. Any BRST anomaly (failure of* $\hat{Q}^2_{\mathrm{BRST}} = 0$ *or of BRST invariance of the measure eq. (12.84)) forces* $\mathcal{C}_{\mathrm{phys}}(G \mid \rho) < 1$ *and excludes* G *at Stage I [105, 106, 107, 108].*

Proof. **Preliminaries.** Let $\mathcal{A}^{(G)}_{\mathrm{BRST}}$ be the abstract BRST $*$–algebra with differential δ_{BRST} and let $\hat{Q}_{\mathrm{BRST}}$ implement δ_{BRST} by graded commutators (Definitions 12.58 and 12.67 and Section 12.3). Assume Stage I instrumentation is in place: BRST–exact gauge fixing (Definition 12.59, (12.74)), conservation/nilpotency (Proposition 12.61, (12.75)), and descent of the relational evolution to cohomology (Proposition 12.64, (12.79)). On the algebraic side, the BRST–closed vacuum functional factors to $\mathcal{A}^{(G)}_{\mathrm{phys}} := H^0(\mathcal{A}^{(G)}_{\mathrm{BRST}}, \delta_{\mathrm{BRST}})$ and the GNS representation agrees with the Hilbert–space cohomology representation (Postulate 12.2 and Proposition 12.68, (12.81)). Holosymmetric meters are read in the spectral/channel–resolved scheme of Chapter 7 on certified plateaux and commute with BRST (chain–map) by Chapter 10 (Definitions 6.275 and 6.282, Sections 6.12.4 and 9.5, and Lemma 9.54). Write $\mathcal{H}^{(G)}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}})$ ((12.76)) and denote by $\mathbf{U}^{(G)}(t) = e^{-it\hat{H}_{\mathrm{total}}}$ the relational evolution, which by Proposition 12.64 induces a group on $\mathcal{H}^{(G)}_{\mathrm{phys}}$ (compression $U^{(G)}_{\mathrm{phys}}(t)$).

(⇒) If $\mathcal{C}_{\mathrm{phys}}(G \mid \rho) = 1$ **for an informationally complete family, then** $\mathbf{U}^{(G)}$ **is unitary on** $\mathcal{H}^{(G)}_{\mathrm{phys}}$**.** By definition of the Stage I coherence functional and its bounds (Definition 12.33 and Proposition 12.35), $\mathcal{C}_{\mathrm{phys}}(G \mid \rho) = 1$ for an informationally complete family ρ means that, on certified plateaux, *all* BRST–closed holosymmetric visibilities and their spectral poles/residues are perfectly self–consistent under the dynamical evolution and instrument transport (Definitions 6.275 and 6.282 and Section 6.12.4) and remain BRST/Nielsen–safe (Lemma 9.54 and Section 9.5). Informational completeness implies these visibilities separate states of $\mathcal{A}^{(G)}_{\mathrm{phys}}$; via Proposition 12.68 the same is true on $\mathcal{H}^{(G)}_{\mathrm{phys}}$. Therefore the descended evolution $U^{(G)}_{\mathrm{phys}}(t)$ preserves *all* expectation values of a separating family of BRST–closed observables; by the GNS/Wigner argument this forces $U^{(G)}_{\mathrm{phys}}(t)$ to act by a one–parameter group of *unitaries* on $\mathcal{H}^{(G)}_{\mathrm{phys}}$. Equivalently, $\mathbf{U}^{(G)}$ is unitary after compression to the physical sector (cf. Proposition 12.64).

(⇐) If $\mathbf{U}^{(G)}$ **is unitary on** $\mathcal{H}^{(G)}_{\mathrm{phys}}$**, then** $\mathcal{C}_{\mathrm{phys}}(G \mid \rho) = 1$**.** Assume $U^{(G)}_{\mathrm{phys}}(t)$ is a unitary group on $\mathcal{H}^{(G)}_{\mathrm{phys}}$. For any BRST–closed observable $\hat{\mathcal{O}}$ represented on $\mathcal{H}^{(G)}_{\mathrm{phys}}$ (Proposition 12.68), the channel–resolved holosymmetric meters are computed as expectation values/correlators in $\mathcal{H}^{(G)}_{\mathrm{phys}}$ (Ch. 7; Definition 6.275) and are independent of gauge parameters on plateaux (Lemma 9.54). Unitarity of $U^{(G)}_{\mathrm{phys}}(t)$ preserves the full set of such correlators, so for any informationally complete family ρ the self–coherence functional attains its upper bound 1 (Proposition 12.35). Thus $\mathcal{C}_{\mathrm{phys}}(G \mid \rho) = 1$.

Anomalies force $\mathcal{C}_{\mathrm{phys}}(G \mid \rho) < 1$ **and exclusion at Stage I.** If there is a BRST anomaly—either failure of nilpotency $\hat{Q}^2_{\mathrm{BRST}} = 0$ or a Fujikawa measure anomaly (12.84)—then by Proposition 12.74 the anomalous Ward identity implies $\partial_\mu j^\mu_{\mathrm{BRST}} = \mathcal{A}(x) \neq 0$ and hence $[\hat{H}_{\mathrm{total}}, \hat{Q}_{\mathrm{BRST}}] \neq 0$; the evolution no longer descends to cohomology (Proposition 12.64 fails), and the BRST–closed vacuum functional does not exist (Postulate 12.2). Consequently, the physical projector and BRST/Nielsen hygiene break down, so the holosymmetric meters

cannot remain perfectly self–coherent on plateaux; by Proposition 12.35 the score strictly drops: $\mathcal{C}_{\text{phys}}(G \mid \rho) < 1$. Per the Holosymmetric Selection Principle, Stage I excludes such a G (Section 12.2 and Remark 12.62) [105, 106, 107, 108].

Combining the two directions and the anomaly clause proves the proposition. □

Proposition 12.76 (Stage II on the physical sector)**.** *For viable A, B, the asymmetric interferometer operated on the dual–register physical space recovers $\Omega_{\text{phys}}(A, B \mid \Psi_0)$ from outcome probabilities exactly as in Section 3.1.5. The sufficiency results Lemma 12.43 and Corollary 12.52 hold with $\mathbf{U}^{(\cdot)}$ replaced by $\mathbf{U}^{(\cdot)}_{\text{phys}}$. In particular, if the induced K–actions on the matter sector are inequivalent, then there exist admissible probes with $\Omega_{\text{phys}}(A, B \mid \Psi_0) = 0$ (Proposition 12.56 and Theorem 12.57).*

Proof. **Step 0 (Stage I viability, chain–map, and compression to the physical sector).** Assume A, B are *viable* in the Stage I sense (cf. Definition 12.72 and Section 12.2). Then the BRST differential is nilpotent and the Noether charge is conserved, so the relational evolution descends to cohomology (Propositions 12.61 and 12.64, (12.75), (12.79)). Equivalently, there exists a cohomological projector Π_{phys} with $[\hat{H}_{\text{total}}, \Pi_{\text{phys}}] = 0$, and the compressed blocks

$$U^{(\cdot)}_{\text{phys}}(t) := \Pi_{\text{phys}}\, U^{(\cdot)}(t)\, \Pi_{\text{phys}}$$

form a unitary group on $\mathcal{H}_{\text{phys}}$ (cf. Proposition 12.75). On the algebraic side, the BRST–closed vacuum functional factors to degree–zero cohomology and its GNS representation agrees with the Hilbert–space cohomology representation (Postulate 12.2 and Proposition 12.68, (12.81)). Crucially, Chapter 10's *pre–physics realization* gives the BRST *chain–map* property for band–tagged meters on certified plateaux (Chs. 7–10; Definitions 6.275 and 6.282, Sections 6.12.4 and 9.5, and Lemma 9.54): for every BRST–closed operator $\hat{\mathcal{O}}$ built from the interferometer primitives,

$$\Pi_{\text{phys}}\hat{\mathcal{O}} = \hat{\mathcal{O}}\Pi_{\text{phys}} = \Pi_{\text{phys}}\hat{\mathcal{O}}\Pi_{\text{phys}} \qquad \text{on} \quad \mathcal{H}_{\text{total}}, \tag{12.89}$$

so matrix elements on BRST classes are unchanged by compression (cf. Definition 12.63).

Step 1 (Dual–register asymmetric interferometer on the physical sector). Fix the asymmetric (character–weighted) interferometer of Chapter 3 (Section 3.1.5; see also the operator forms in Definitions 3.13 and 3.15) and run it on the *dual–register* BRST–physical space $\mathcal{H}_{\text{phys}} \otimes \mathcal{H}_{\text{phys}}$, with the dynamical blocks replaced by their compressions $U^{(A)}_{\text{phys}}, U^{(B)}_{\text{phys}}$ from Step 0. Let $|\Psi_0\rangle$ be an admissible (BRST–closed) probe in $\mathcal{H}_{\text{phys}} \otimes \mathcal{H}_{\text{phys}}$, and use the same beam–splitter and control gates as in Section 3.1.5. Since those gates are BRST–closed instruments, (12.89) applies to each primitive, so the outcome probabilities are *unchanged* by the replacement $U^{(\cdot)} \mapsto U^{(\cdot)}_{\text{phys}}$. Repeating the derivation in Section 3.1.5 then yields

$$p_{\pm} = \tfrac{1}{2}\Big(1 \pm \operatorname{Re}\, \Omega_{\text{phys}}(A, B \mid \Psi_0)\Big), \qquad \Omega_{\text{phys}}(A, B \mid \Psi_0) := \langle \Psi_0 |\, U^{(A)\dagger}_{\text{phys}} U^{(B)}_{\text{phys}}\, | \Psi_0 \rangle,$$

i.e. the interferometer reconstructs the *physical* holosymmetric overlap exactly as stated (cf. Definition 12.69, (12.83)).

Step 2 (Sufficiency and informational completeness on cohomology). The completeness results Lemma 12.43 and Corollary 12.52 rely on three inputs only: (i) unitarity of the arm

blocks, (ii) availability of the same interferometric primitives, and (iii) an informationally complete family of probes/meters. By Step 0 the compressed blocks $U^{(\cdot)}_{\mathrm{phys}}$ are unitary on $\mathcal{H}_{\mathrm{phys}}$ (Propositions 12.64 and 12.75); the primitives are BRST–closed and thus valid on cohomology (by (12.89) and Definition 12.63); and informational completeness is implemented by Chapter 7's channel–resolved holosymmetric meters on certified plateaux (Chs. 7–8; Definitions 6.275 and 6.282 and Section 6.12.4) with BRST/Nielsen hygiene (Ch. 10; Lemma 9.54 and Section 9.5). Therefore the proofs of Lemma 12.43 and Corollary 12.52 carry over verbatim with $U^{(\cdot)} \to U^{(\cdot)}_{\mathrm{phys}}$.

Step 3 (Inequivalent K–actions ⇒ holosymmetric orthogonality on $\mathcal{H}_{\mathrm{phys}}$). If the induced K–actions on the tantrakyma (matter) sector are inequivalent, then Proposition 12.56 furnishes character weights and probe settings for which the holosymmetric overlap vanishes, and Theorem 12.57 guarantees the existence of admissible probes realizing this separation. Because our overlaps are computed with $U^{(A)}_{\mathrm{phys}}, U^{(B)}_{\mathrm{phys}}$ (Step 1), we obtain $\Omega_{\mathrm{phys}}(A, B \mid \Psi_0) = 0$ for some admissible $|\Psi_0\rangle \in \mathcal{H}_{\mathrm{phys}} \otimes \mathcal{H}_{\mathrm{phys}}$. When geometric channels are present, Chapter 11's typed/TT projectors isolate the spin–2 graviton block and enforce the TT selection rule, precluding contamination of the internal channel used by the meters (cf. Definition 10.3, eqs. (10.4), (10.30), (10.90), (10.106) and (10.134), Lemma 10.17, and Proposition 10.173).

Conclusion. Steps 1–3 show that the asymmetric dual–register interferometer on the BRST–physical space reconstructs $\Omega_{\mathrm{phys}}(A, B \mid \Psi_0)$ exactly as in Section 3.1.5; that Lemma 12.43 and Corollary 12.52 remain valid with $U^{(\cdot)} \to U^{(\cdot)}_{\mathrm{phys}}$; and that inequivalent K–actions imply the existence of admissible probes with $\Omega_{\mathrm{phys}}(A, B \mid \Psi_0) = 0$ (Proposition 12.56 and Theorem 12.57). This completes the proof of Proposition 12.76. □

3. Holosymmetric selection stated on cohomology

Definition 12.77 (Holosymmetric coherence effect and projector)**.** Fix admissible preparation/readout gates on the control (BRST–closed instruments so they act on cohomology; cf. Definition 12.63) and let $\mathbf{U}^{(G)}_{\mathrm{holo}}$ denote the holosymmetric circuit *compressed to the BRST–physical space* $\mathcal{H}^{(G)}_{\mathrm{phys}}$ (cf. (12.76), Propositions 12.64 and 12.75; holosymmetry set–up in Ch. 3, Section 3.8, and its physical overlap in Definition 12.69). The *holosymmetric coherence effect* on $\mathcal{H}^{(G)}_{\mathrm{phys}}$ is

$$E^{(G)}_{\mathrm{holo}} \;:=\; \Big\langle \mathbf{a}\Big|_c \, \mathbf{U}^{(G)\,\dagger}_{\mathrm{holo}} \Big(|\mathbf{b}\rangle\langle\mathbf{b}|_c \otimes \mathbb{I}\Big)\, \mathbf{U}^{(G)}_{\mathrm{holo}} \Big|\mathbf{a}\Big\rangle_c, \qquad 0 \le E^{(G)}_{\mathrm{holo}} \le \mathbb{I}. \tag{12.90}$$

Here $|\mathbf{a}\rangle_c, |\mathbf{b}\rangle_c$ are the control–register gates from the (symmetric/asymmetric) holosymmetric interferometers (Definitions 2.123, 2.124 and 3.15, Section 3.1.5, and Corollary 3.7). The inequality $0 \le E^{(G)}_{\mathrm{holo}} \le \mathbb{I}$ follows from positivity and contractivity of effects obtained by unitary conjugation and compression to the physical sector (Proposition 12.75; see also the channel–resolved, plateau–licensed meters in Definition 6.275 and Section 6.12.4 that read $\mathrm{Tr}(\rho\, E^{(G)}_{\mathrm{holo}})$).

Its top eigenvalue $\lambda_{\max}(E^{(G)}_{\mathrm{holo}}) \in [0, 1]$ defines the *holosymmetric coherence projector* as the spectral projector onto the top eigenspace:

$$\mathcal{P}^{\mathrm{coh}}_G \;:=\; \mathbf{1}_{\{\lambda_{\max}\}}\Big(E^{(G)}_{\mathrm{holo}}\Big). \tag{12.91}$$

Equivalently, for any probe state ρ on $\mathcal{H}_{\mathrm{phys}}^{(G)}$, $\mathrm{Tr}(\rho\, E_{\mathrm{holo}}^{(G)}) \le \mathrm{Tr}(\rho\, \mathcal{P}_G^{\mathrm{coh}})$, with equality for ρ supported on $\mathrm{ran}\, \mathcal{P}_G^{\mathrm{coh}}$. In Stage II this projector is used together with the asymmetric, dual–register scheme to reconstruct physical overlaps and enforce holosymmetric selection on cohomology (cf. Definition 12.73 and Proposition 12.76).

Corollary 12.78 (Holosymmetry as quantum consistency constraint)**.** *Among BRST-viable candidates $G \subset G_{\mathrm{int}}$, the realized symmetry $G_\star$ maximizes physical coherence:*

$$G_\star \;\in\; \arg\max_G \left\{ \sup_{\substack{\text{prep/readout} \\ \rho \text{ on } \mathcal{H}_{\mathrm{phys}}^{(G)}}} \mathrm{Tr}\Big(\rho\, E_{\mathrm{holo}}^{(G)}\Big) \right\} \;=\; \arg\max_G \left\{ \sup_{\|\psi\|=1} \Big\| \mathcal{P}_G^{\mathrm{coh}}\, |\psi\rangle \Big\|^2 \right\}. \tag{12.92}$$

Stage I viability requires $\mathcal{C}_{\mathrm{phys}}(G) = 1$ (Proposition 12.75); among those, Stage II comparative tomography (Proposition 12.76) ensures that $G_\star$ is uniquely selected by attaining maximal coherence while being holosymmetrically orthogonal to all other viable candidates.

Proof. **Step 1 (Variational form and projector equivalence).** For each BRST–viable G (Stage I; cf. Definition 12.72 and Section 12.2), the holosymmetric coherence effect $E_{\mathrm{holo}}^{(G)}$ on $\mathcal{H}_{\mathrm{phys}}^{(G)}$ is a positive contraction, $0 \le E_{\mathrm{holo}}^{(G)} \le \mathbb{I}$ (Definition 12.77). By Rayleigh–Ritz,

$$\sup_\rho \mathrm{Tr}\Big(\rho\, E_{\mathrm{holo}}^{(G)}\Big) = \sup_{\|\psi\|=1} \langle\psi| E_{\mathrm{holo}}^{(G)} |\psi\rangle = \lambda_{\max}\Big(E_{\mathrm{holo}}^{(G)}\Big), \tag{12.93}$$

with the supremum over density operators (or unit vectors) on $\mathcal{H}_{\mathrm{phys}}^{(G)}$. By Definition 12.77, the spectral projector onto the top eigenspace is $\mathcal{P}_G^{\mathrm{coh}} = \mathbf{1}_{\{\lambda_{\max}\}}(E_{\mathrm{holo}}^{(G)})$, and

$$\sup_{\|\psi\|=1} \|\mathcal{P}_G^{\mathrm{coh}}\psi\|^2 = \sup_{\|\psi\|=1} \langle\psi|\, \mathcal{P}_G^{\mathrm{coh}}\, |\psi\rangle = \text{rank-one saturation of (12.93)} = \lambda_{\max}\Big(E_{\mathrm{holo}}^{(G)}\Big). \tag{12.94}$$

Combining (12.93)–(12.94) gives the equality of the two arg max forms in the statement.

Step 2 (Realized symmetry $G_\star$ attains the maximum). Stage I on the physical sector (Proposition 12.75) implies that for a viable G the compressed evolution $\mathbf{U}_{\mathrm{phys}}^{(G)}$ is unitary on $\mathcal{H}_{\mathrm{phys}}^{(G)}$, yielding $\mathcal{C}_{\mathrm{phys}}(G \mid \rho) = 1$ for every probe ρ (Definition 12.69, (12.82)). Specializing the asymmetric interferometer of Chapter 3 to a *self*-comparison (both arms set to $G_\star$) and running it on the BRST–physical space (Ch. 3 + descent; Section 3.1.5 and Proposition 12.76) gives

$$\Omega_{\mathrm{phys}}(G_\star, G_\star \mid \Psi_0) = \langle\Psi_0|\, U_{\mathrm{phys}}^{(G_\star)\dagger} U_{\mathrm{phys}}^{(G_\star)}\, |\Psi_0\rangle = 1 \quad \Rightarrow \quad p_+ = 1,$$

for suitable control settings and any unit vector $|\Psi_0\rangle$ (cf. Definition 12.69, (12.83)). Hence the corresponding effect $E_{\mathrm{holo}}^{(G_\star)}$ has an eigenvalue 1 and therefore $\lambda_{\max}\Big(E_{\mathrm{holo}}^{(G_\star)}\Big) = 1$. By (12.93)–(12.94), $G_\star$ *attains* the maximal physical coherence and thus belongs to the arg max set.

Step 3 (Comparative tomography on cohomology and orthogonality of non–realized candidates). For any other BRST–viable $G \neq G_\star$, Stage II on the physical sector (Proposition 12.76) ensures that the dual–register asymmetric interferometer reconstructs $\Omega_{\mathrm{phys}}(G, G_\star \mid \Psi_0)$ on $\mathcal{H}_{\mathrm{phys}}$ exactly as in Section 3.1.5. If the induced K–actions on the

tantrakyma (matter) sector for G and $G_\star$ are inequivalent, Proposition 12.56 provides admissible character weights and probes with $\Omega_{\mathrm{phys}}(G, G_\star \mid \Psi_0) = 0$, and Theorem 12.57 guarantees the existence of such admissible probes. Thus G is *holosymmetrically orthogonal* to $G_\star$ in the physical sector under Stage II tomography. (When geometric channels are present, typed/TT selection guarantees spin–2 decoupling during these measurements; cf. Ch. 11, Definition 10.3, eqs. (10.4), (10.30), (10.90), (10.106) and (10.134), Lemma 10.17, and Proposition 10.173.)

Step 4 (Selection rule). By Step 2, $G_\star$ achieves the global maximum $\lambda_{\max}(E^{(G)}_{\mathrm{holo}})$; hence $G_\star \in \arg\max_G\{\cdots\}$. By Step 3, any other viable $G \neq G_\star$ can be made holosymmetrically orthogonal to $G_\star$ via admissible probes/meters, so comparative Stage II tomography distinguishes $G_\star$ uniquely among the Stage–I–admitted set: it is the *only* candidate that both (i) attains maximal coherence and (ii) is orthogonal to all competitors under the holosymmetric tests (Propositions 12.56 and 12.76 and Theorem 12.57).

Combining Steps 1–4 proves Corollary 12.78. □

Theorem 12.79 (HSP stated on the BRST–physical sector)**.** *Among admissible $G \subset G_{\mathrm{int}}$:*

1. *(*Stage I*) G is viable iff $\mathcal{C}_{\mathrm{phys}}(G \mid \rho) = 1$ for an informationally complete family of probes on $\mathcal{H}^{(G)}_{\mathrm{phys}}$ (eq. (12.82)).*
2. *(*Stage II*) For the realized $G_\star$, the asymmetric tomography achieves $\Omega_{\mathrm{phys}}(G_\star, B \mid \Psi_0) = 0$ for every other viable $B \neq G_\star$ (holosymmetric orthogonality on cohomology), while $\Omega_{\mathrm{phys}}(G_\star, G_\star \mid \Psi_0)$ is maximized.*

Under the anomaly, chirality, and minimality constraints of this chapter, $G_\star \simeq G_{\mathrm{SM}}$ (Theorem 12.57).

Proof. **Preliminaries (BRST descent and physical sector).** For an admissible candidate $G \subset G_{\mathrm{int}}$, Stage I asks for a nilpotent, conserved BRST charge and a BRST–invariant measure; equivalently, the relational evolution must *descend* to cohomology (Definition 12.72, Section 12.2, and Propositions 12.61 and 12.64, (12.75), (12.79)). On the algebraic side, the BRST–closed vacuum functional factors to degree–zero cohomology and its GNS representation coincides with the Hilbert–space cohomology representation (Postulate 12.2 and Proposition 12.68, (12.81)), Definitions 6.275 and 6.282, Sections 6.12.4 and 9.5, and Lemma 9.54).

(1) Stage I: G viable $\Longleftrightarrow$ $\mathcal{C}_{\mathrm{phys}}(G \mid \rho) = 1$ for an IC family. By Proposition 12.75, for an informationally complete family of probes on $\mathcal{H}^{(G)}_{\mathrm{phys}}$ the following are equivalent:

$$\mathcal{C}_{\mathrm{phys}}(G \mid \rho) = 1 \quad \Longleftrightarrow \quad \mathbf{U}^{(G)}_{\mathrm{phys}} \text{ is unitary on } \mathcal{H}^{(G)}_{\mathrm{phys}} \quad \Longleftrightarrow \quad [\hat{H}_{\mathrm{total}}, \hat{Q}_{\mathrm{BRST}}] = 0,$$

with $\mathcal{C}_{\mathrm{phys}}$ as in Definition 12.69, (12.82). If a BRST anomaly is present (either $\hat{Q}^2_{\mathrm{BRST}} \neq 0$ or a nontrivial Fujikawa class in (12.84)), then the anomalous Ward identity implies nonconservation of $\hat{Q}_{\mathrm{BRST}}$, the evolution fails to descend, and the self–coherence score strictly drops below 1 (Proposition 12.74). Therefore *viability* (nilpotency, conserved charge, measure invariance; Definition 12.72) holds iff $\mathcal{C}_{\mathrm{phys}}(G \mid \rho) = 1$ for an informationally complete family.

(2) Stage II: realized $G_\star$ maximizes self–coherence and is holosymmetrically orthogonal to every other viable $B \neq G_\star$. On the BRST–physical sector, the asymmetric dual–register interferometer reconstructs the *physical* holosymmetric overlap $\Omega_{\mathrm{phys}}(A, B \mid \Psi_0) =$

$\langle\Psi_0|U_{\rm phys}^{(A)\dagger}U_{\rm phys}^{(B)}|\Psi_0\rangle$ with detection statistics exactly as in Section 3.1.5 (now with compressed blocks), by Proposition 12.76 and (12.83). Among the Stage I–admitted set, Corollary 12.78 shows that the realized symmetry $G_\star$ *maximizes* physical coherence:

$$\lambda_{\max}\Big(E_{\rm holo}^{(G_\star)}\Big) = 1 = \sup_{\rho} \mathrm{Tr}(\rho\, E_{\rm holo}^{(G_\star)}) = \sup_{\|\psi\|=1} \|\mathcal{P}_{G_\star}^{\rm coh}\psi\|^2,$$

with $E_{\rm holo}^{(G)}$ and $\mathcal{P}_G^{\rm coh}$ from Definition 12.77. For any other viable $B \neq G_\star$ whose induced K–action on the matter sector is inequivalent, Proposition 12.56 and Theorem 12.57 provide admissible probes/meters such that $\Omega_{\rm phys}(G_\star, B \mid \Psi_0) = 0$ (holosymmetric orthogonality on cohomology), while the self–overlap satisfies $\Omega_{\rm phys}(G_\star, G_\star \mid \Psi_0) = 1$ (cf. Definition 12.69, (12.83)). When gravitational channels are present, typed/TT selection guarantees that spin–2 content does not contaminate these internal–channel diagnostics (Definition 10.3, eqs. (10.4), (10.30), (10.90), (10.106) and (10.134), Lemma 10.17, and Proposition 10.173).

(3) Identification of $G_\star$. Imposing the anomaly, chirality, and minimality constraints of this chapter singles out the unique bridge to phenomenology, yielding $G_\star \simeq G_{\rm SM}$ by Theorem 12.57.

Items (1)–(3) establish Theorem 12.79. □

Explanatory notes (overlay perspective). Theorem 12.79 is the synthesis layer where the BRST construction (Ch. 10/13) and the holosymmetric instrumentation (Chs. 7–8) are overlaid and then *compressed* to the BRST–physical sector. Item (1) reads the *channel–resolved holosymmetric visibilities* of Chapter 7 on certified plateaux (Definition 6.275 and Section 6.12.4) and identifies their physical incarnation with the *physical self–coherence* (Definition 12.69, (12.82)). The Stage I equivalence asserted in the theorem is exactly Proposition 12.75 once the meters are evaluated on cohomology. Item (2) deploys the *asymmetric* two–path law/state interferometer (Ch. 3; Section 3.1.5 and Definitions 3.13 and 3.15) but now with the blocks compressed to the physical sector, thereby reconstructing the *physical* holosymmetric overlap $\Omega_{\rm phys}(\cdot,\cdot \mid \Psi_0)$ (Definition 12.69, (12.83)) exactly as used in Proposition 12.76. Throughout, transport covariance is governed by the same *universal* operator–bundle connection $\mathcal{A}_\mu$ (Definition 6.192 and Remark 6.200); the theorem preserves this universality by evaluating the meters on the BRST–physical quotient rather than modifying their kinematics.

Role of band/transport instrumentation. The theorem does not introduce any structure beyond Chs. 7–8; it *uses* the band/scale transport hygiene codified by the band functor and the plateau license to guarantee that the meters are stable, context–aware instruments (Definition 6.282 and Section 6.12.4). These hypotheses are precisely those invoked in Propositions 12.75 and 12.76 to ensure (i) informational completeness of probe families and (ii) robustness of the interferometric readouts under transport. Optimization is then stated at the operator level via the *holosymmetric coherence effect* and its spectral projector (Definition 12.77), which feed into the argmax principle of Corollary 12.78.

Why Chapter 10/13 guarantees correctness on the physical sector. Two realization layers justify reading all Chapter 7 meters on $H^0(\hat{Q}_{\rm BRST})$. First, the *pre–physics realization* (Ch. 10)

shows that the interferometer primitives and holosymmetric meters are *BRST–closed* and satisfy a chain–map relation with the cohomological projector, so compression leaves matrix elements unchanged (Definition 12.63 and Lemma 9.54). Second, the *physics realization* (Ch. 13) proves that the relational dynamics *descend* to cohomology and that algebraic and Hilbert–space cohomologies *coincide* via GNS (Proposition 12.64, (12.79), Postulate 12.2 and Proposition 12.68, (12.81)). Consequently, the quantities used in the theorem—$\mathcal{C}_{\text{phys}}$ and Ω_{phys}—are the correct, representation–independent objects (Definition 12.69), and the Stage I/II claims are precisely Propositions 12.75 and 12.76 read on cohomology.

Selection and uniqueness. Given the above, Stage I filters to the BRST–viable set by the unitary–compression criterion (Proposition 12.75), and Stage II distinguishes among those via asymmetric, dual–register tomography on the physical sector (Proposition 12.76). The optimizer statement—maximizing the top eigenvalue of $E^{(G)}_{\text{holo}}$ or, equivalently, the mass of $\mathcal{P}^{\text{coh}}_{G}$—is Corollary 12.78. Under the anomaly/chirality/minimality constraints of the chapter, the bridge–uniqueness theorem then identifies the realized law with the Standard Model gauge group, $G_\star \simeq G_{\text{SM}}$ (Theorem 12.57). (When gravitational channels are present, the typed/TT selection of Ch. 11 ensures spin–2 decoupling from the internal channel used by these meters, so the comparisons remain faithful; cf. Definition 10.3, eqs. (10.4), (10.30), (10.90), (10.106) and (10.134), Lemma 10.17, and Proposition 10.173.)

Unified flow (algebra → cohomology → selection).

$$\mathcal{A}(\mathcal{C}_d) \xrightarrow[\text{BRST}]{\text{extend \& } \delta_{\text{BRST}}} \mathcal{A}^{(G)}_{\text{phys}} \xrightarrow{\text{GNS of } \omega^{(G)}_{\text{phys}}} \left(\mathcal{H}^{(G)}_{\text{phys}}, \mathbf{U}^{(G)}_{\text{phys}}\right) \xrightarrow{\text{HSP}} G_{\text{SM}}.$$

BRST cohomology supplies the gauge–consistent physical arena; HSP acts *there* to select the unique, maximally coherent law realized in the bright fringe of the holocosmos.

The Universe Sculpted by Coherence: An Interpretation

The formalism above describes a hierarchical, quantum–informational genesis of law and content—a *sculpting by coherence*. The primordial arena is not a metric manifold but a pre–geometric substrate of *Primordial Quantum Singularities* (Chapter 1), a relational continuum of pure potential.

Algebraic inception. Order first appears as the abstract *Algebra of Observables* $\mathcal{A}(\mathcal{C}_d)$, the C^*–algebra of all admissible relational observables (Definition 1.32). A *Quantum Holo–State* $\omega^{\mathring{\mathbf{\Psi}}}$ imposes a global coherence constraint and, via the GNS construction, realizes an emergent Hilbert representation $(\pi^{\mathring{\mathbf{\Psi}}}, \mathcal{H}_{\text{em}})$ (Definition 1.34). This is the first sculpting: from algebraic potential to a concrete quantum stage.

Cohomological purification (BRST). Because $\mathcal{H}_{\text{em}}$ respects local gauge covariance, it is overcomplete: it contains redundant (gauge) and ghost degrees of freedom. The *BRST* differential δ_{BRST} provides a covariant, nilpotent filtration of this redundancy (Section 12.3).

Physical content is the degree–zero cohomology $H^0(\hat{Q}_{\mathrm{BRST}})$, characterized in Principle 12.1, and preserved by the relational dynamics (Proposition 12.64). This is the second sculpting: *purification* of the emergent space to $\mathcal{H}_{\mathrm{phys}}$.

Comparative selection (Holosymmetry). Even after BRST purification, distinct BRST–viable gauge sectors $G \subset G_{\mathrm{int}}$ can act on $\mathcal{H}_{\mathrm{phys}}$. The *Holosymmetric Selection Principle* (HSP) operates *on* this physical sector (Section 12.3.2): Stage I enforces internal viability via perfect self–coherence, and Stage II implements comparative tomography to suppress all but the maximally coherent law (Theorem 12.79 and Propositions 12.75 and 12.76). In the admissible, Tantrakyma–compatible class with SM chiral content and anomaly freedom, this singles out the Standard Model gauge action (Theorem 12.57).

Ontological chain (from potential to law).

$$\mathcal{A}(\mathcal{C}_d) \xrightarrow{\omega^{\mathring{\Psi}}\ \text{(GNS)}} \mathcal{H}_{\mathrm{em}} \xrightarrow{\text{BRST cohomology}} \mathcal{H}_{\mathrm{phys}} \xrightarrow{\text{HSP (Stage I \& II)}} G_{\mathrm{SM}}.$$

Interpretive summary. In this view, the laws of nature are *selected*, not posited. The algebraic substrate supplies possibility; the Holo–State and GNS realize a quantum arena; BRST enforces gauge consistency by cohomological purification; and Holosymmetry performs a final, interferometric selection within the BRST–physical sector, yielding G_{SM} as the unique fixed point of maximal relational coherence (Theorems 12.42 and 12.79). The cosmos thus appears as the artifact of its own coherence: *from potential to representation, from representation to physicality, and from physicality to uniquely realized law.*

12.4 The Emergence of Fermions from the Graded Tantrakyma Field

The existence of fermions—the spin-$\frac{1}{2}$ constituents of matter—is not an independent axiom within the IQR Framework. Instead, it is a necessary and profound consequence of the theory's foundational algebraic structure and emergent geometry. We shall now formalize this emergence through a synthesis of two principles: the grounding of fermionic statistics in a graded algebra, and the realization of fermionic fields as sections of an operator-valued spinor bundle. (Foundational fourfold algebra and GNS representation: Section 1.4.2 and Definitions 1.34 and 4.9; emergent operator geometry and chronogeometry: Definitions 5.1 and 5.26; relational compatibility of derivatives: Definition 6.7.)

Fermionic Statistics from the Dialectical Grading. The ontological origin of the distinction between bosons and fermions lies in the $\mathbb{Z}_2 \times \mathbb{Z}_2$ graded structure of the Fourfold Dialectic Algebra (Section 4.1.7, cf. the symmetry–corner bridge in Definition 1.23 and the prismatic unity in Section 1.2.5). We postulate a direct correspondence between this algebraic grade and quantum statistics, providing a deep reason for the spin-statistics theorem. This parallels the conceptual foundation laid by Pauli in [558], and generalizations explored through graded

quantum algebras [559, 560]. (Modal carriers and inter-modal structure: Definitions 4.1 and 4.9.)

Excitations of the *Tantrakyma field* $\hat{\mathbf{\Theta}}(x)$ that correspond to modes of *odd grade*—namely, those associated with the *Antisymmetry* and *Holosymmetry* sectors of the dialectic—are defined to obey Fermi–Dirac statistics (odd/even grading and sector identification: Section 4.1.7, with Antisymmetry/Holosymmetry mapped via Definitions 1.2 and 1.4). It is these "odd-parity" kymons that constitute the fundamental fermions of the theory. This algebraic assignment ensures that their anti-commuting nature is not an ad-hoc property but a direct result of their role in the universe's foundational logic. (Dynamical Tantrakyma field and action: Section 6.3 and Definitions 6.268 and 9.301.)

Background split, diagnostic realized reference configuration discipline, and typed realization. We work throughout with the Tantrakyma background–fluctuation split $\hat{\mathbf{\Theta}}(x) = \bar{\Theta}(x)\,\mathbb{I}_x + \delta\hat{\mathbf{\Theta}}(x)$ taken on a certified plateau $\mathfrak{P}$ with UV/IR tags and BRST projection to $H^0(\hat{Q}_{\mathrm{BRST}})$; all typed/triadic manipulations use the weave projectors and triad, and realization proceeds via the Σ–pipeline (Definitions 7.29, 7.39, 9.60, 10.2, 10.4 and 10.204 and eq. (10.3)).

Kymons as poles of typed two–point functions. Typed Tantrakyma excitations ("kymons") are the one–particle modes associated with simple poles of the retarded two–point kernels $\Delta^{[\chi]}_{\Theta\Theta}$ on $H^0(\hat{Q}_{\mathrm{BRST}})$; equivalently, they arise from the linear response equation with typed kinetic block $\mathbb{K}^{[\chi]}_{\Theta}$ and typed Hessian $\mathcal{K}^{[\chi]}_{\Theta\Theta}$ (cf. eqs. (9.143) and (9.175)). Positivity and gauge–parameter independence of the physical residues follow from the KL/Nielsen diagnostic realized reference configuration identities, and species are isolated by Riesz selectors on band–admissible spectral islands (Proposition 7.244, Theorem 7.283, eq. (12.12), Definitions 10.6 and 10.7, and Section 10.2).

Radial (Higgs) kymon. Writing the Tantrakyma invariant $\mathsf{X} := g_C(\hat{\mathbf{\Theta}}, \hat{\mathbf{\Theta}})$, the radial direction at $\bar{\Theta}$ is singled out by $\delta\mathsf{X}|_{\bar{\Theta}} = 2\,g_C(\delta\hat{\mathbf{\Theta}}, \bar{\Theta})$, i.e. the g_C–Riesz image of $\bar{\Theta}$. The scalar (singlet) Higgs kymon is obtained by restricting the potential Hessian to this line as encoded by the operator $\mathcal{K}_{\mathrm{H}}$, whose spectrum yields M^2_{H}; the corresponding propagator has a single gapped pole (contrast NG/pNG channels) (eq. (9.185) and Definition 9.249).

Gauge coupling and BEH setup. For a compact G acting via ρ on the Tantrakyma fibre, we extend the split connection on $(\mathcal{X}_q, \bar{\Sigma})$ by the internal gauge potential, defining the covariant derivative D_μ and the typed/triadic inserted action on the diagnostic realized reference configuration (Definitions 6.7, 9.187 and 9.310, Section 6.12.2, and eq. (6.314)). Physical poles/residues are Nielsen–flat, and hyperbolicity/microcausality use the $\bar{\Sigma}$–cone principal symbol.

SSB pattern, NG projector, and radial split. Let $\bar{\Theta}$ minimize V_Θ; the stabilizer $H \subset G$ satisfies $\rho(T_a)\bar{\Theta} = 0$, while broken directions T_b generate the NG subspace $\mathcal{N} = \mathrm{span}\{\rho(T_b)\bar{\Theta}\}_b$. The g_C–Riesz projector onto $\mathcal{N}$ provides the typed NG extraction, and the radial line remains $\hat{\mathsf{R}} \propto \bar{\Theta}$ with its associated typed mass operator $\mathcal{K}^{[\chi]}_{\mathrm{rad}}$; all projectors commute with the triad on the diagnostic realized reference configuration (Definitions 9.249 and 9.250 and Proposition 9.207). :contentReference[oaicite:7]index=7 :contentReference[oaicite:8]index=8

Vector masses from BEH. Expanding about $\bar{\Theta}$ and reading the quadratic gauge terms yields the internal vector mass matrix $(M^2)_{IJ} = g^2\, g_C\Big(\rho(T_I)\bar{\Theta}, \rho(T_J)\bar{\Theta}\Big)$; vectors along broken directions

acquire masses from spec(M^2), while H–vectors remain massless. The scalar spectrum splits into the radial Higgs kymon and the NG/pNG multiplet; typed transverse/longitudinal/helicity resolutions follow the standard vector projectors, and the eigenvalues are Nielsen–flat on $\mathfrak{P}$ (Definitions 9.252 and 9.350 and eq. (12.12)).

Minimal $U(1)$ illustration. For $G = U(1)$ with $\rho(T) = q$ and $\bar{\Theta} \neq 0$, the gauge boson mass is $M_A^2 = g^2 q^2\, g_C(\bar{\Theta}, \bar{\Theta})$; the NG direction span$\{\rho(T)\bar{\Theta}\}$ is eaten, and the radial Higgs kymon is given by the Hessian restriction referenced above. Nielsen flatness of M_A holds on $\mathfrak{P}$ (eqs. (9.185) and (12.12) and Theorem 7.283).

The τ–kairos field and fermion bilinears. The non–TT temporal/coherence tensor decomposes orthogonally into typed scalar, longitudinal–scalar, pseudoscalar, and helicity–resolved vector summands, each $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$–null. Standard fermion bilinears land precisely in these τ–channels (after helicity split), providing the bridge from the odd–graded fermionic sector to the coherence block (Definitions 10.3, 10.4 and 12.81).

The τ-kairos field is the orthogonal sum

$$\boxed{\hat{\tau}_{\mu\nu} = S^{(b)}_{\mu\nu}[\hat{\varphi}_b] \;\oplus\; S^{(L)}_{\mu\nu}[\hat{\varphi}_L] \;\oplus\; \tilde{S}^{(p)}_{\mu\nu}[\tilde{\varphi}] \;\oplus\; \sum_{\lambda=\pm} A^{(\lambda)}_{\mu\nu}\big[\hat{K}^{(\tau)}_{\lambda}\big]} \tag{12.95}$$

with scalar amplitudes $\hat{\varphi}_b, \hat{\varphi}_L$, pseudoscalar $\tilde{\varphi}$, and helicity-resolved vector creators $\hat{K}^{(\tau)}_{\lambda}$. Each typed summand is $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$-null, cf. Definitions 10.3 and 10.4. Standard fermion bilinears land in these channels:

$$\bar{\psi}\psi \in S^{(b)}, \qquad \bar{\psi}\, i\gamma_5\psi \in \tilde{S}^{(p)}, \qquad \bar{\psi}\gamma_\mu\psi \in A^{(\pm)} \text{ (after helicity split)},$$

linking the odd-graded fermionic kymons to the τ-sector (cf. Definition 12.81).

Chrono–Kymon: vector channel of τ. On the realized weave configuration inside $\mathcal{H}_{\mathrm{phys}}$, with triad $\Pi^{(a)}$ and helicity idempotents $A^{(\pm)}$, the spin–1 Riesz selector on the τ–vector subspace isolates the realized helicity–1 spectral features. Helicity–resolved Chrono–Kymon projectors then extract (helicity, triad) CK fields from raw τ–vector fluctuations, and their tensor images are obtained via the typed map $A^{(\lambda)}_{\mu\nu}[\cdot]$ (Definitions 10.1, 10.2, 10.4, 10.56 and 10.57).

Parity and visibility diagnostics. Parity on the τ–vector subspace exchanges the helicity idempotents $A^{(+)} \leftrightarrow A^{(-)}$. The law/state visibility asymmetry $\Delta^{(1)}_{P,\Lambda}$ compares the corresponding witnesses, with robustness under spectral mixing controlled by the established Davis–Kahan/visibility bounds (Definitions 10.80 and 10.82).

Chronogeometric placement and TT selection. At expectation level the chronogeometric tensor splits as $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$; Tantrakyma variations populate only the non–TT temporal structure, with $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \circ D_\Theta \Sigma = 0$, leaving the spin–2 (metric/TT) graviton sector intact (Definitions 5.26 and 10.3).

Diagnostic realized reference configuration quantization and CK BEH option. Linear–response quantization on $\mathfrak{P}$ uses typed, band-limited kernels with triad insertions and $\bar{\Sigma}$–cone microcausality; physical objects live on $H^0(\hat{Q}_{\mathrm{BRST}})$ with Nielsen/KL control. When an internal gauge acting on the τ–vector carrier is gauged and broken, the CK acquires a BEH

mass; helicity projectors pick the transverse physical modes, longitudinal components are BRST–removed, and the CK mass is Nielsen–flat with robustness under spectral mixing (Theorem 7.283, Proposition 7.244, and Definitions 10.57 and 10.58).

Geometric Realization in Operator-Valued Spinor Bundles. To describe these fermionic excitations geometrically, we must define the arena in which they propagate. This is not the tangent bundle, but a more specialized structure.

We introduce the *Tantrakyma spinor bundle*, a vector bundle $\mathcal{S} \to \mathcal{X}_q$ over the emergent quantum configuration space (emergent manifold and operator-level structure: Definition 5.64). This bundle is defined with respect to the full emergent chronogeometric structure given by the unified tensor $\bar{\Sigma}_{\mu\nu}$, ensuring that its geometry is fully covariant (Definitions 5.26 and 6.7; cf. band-compatible operator connection Definition 12.1 and hyperbolic principal symbol eq. (7.15)) ([8, 3] for the importance of relational covariance in quantum geometry).

A *fermionic kymon field*, denoted $\hat{\psi}(x)$, is a section of the operator-valued bundle $\mathrm{End}(\mathcal{H}_{\mathrm{em}}) \otimes \mathcal{S}$ (GNS origin of $\mathcal{H}_{\mathrm{em}}$: Definition 1.34). At each point x, $\hat{\psi}(x)$ is thus an operator on the total Hilbert space that transforms as a spinor under the local Lorentz group $\mathrm{Spin}(1,3)$; its dynamics and covariant kinetic operator are constructed on $(\mathcal{X}_q, \bar{\Sigma})$ using the realized weave configuration-compatible connection (Definition 12.1) and, where needed, the Lichnerowicz identity in the realized setting (eq. (12.16)). (See also the formal definition and properties in Definition 12.80.)

Definition 12.80 (Operator-Valued Tantrakyma Spinor Bundle and Fermionic Kymon Field)**.** Let $\mathcal{X}_q$ be the emergent quantum configuration space (Proposition 4.625), endowed with the effective Lorentzian metric structure $\bar{g}^{\mathrm{Lor}}_{\mu\nu}$ derived from the chronogeometric tensor $\bar{\Sigma}_{\mu\nu}$ (Definitions 5.26, 5.77 and 6.24). Equip $(\mathcal{X}_q, \bar{\Sigma})$ with the Relational Levi–Civita connection $\nabla^{(\mathrm{rel})}$ and relational measure $d\mu(x) = \sqrt{-\det \bar{\Sigma}}\, d^d x$ (Definitions 6.7 and 6.33). Let $\mathcal{H}_{\mathrm{em}}$ be the total emergent Hilbert space (Definition 4.287).

1. The Tantrakyma Spinor Bundle. We define the *Tantrakyma spinor bundle* as a complex vector bundle $\mathcal{S} \to \mathcal{X}_q$, constructed with the following properties:

- **Base Space:** The emergent configuration space $\mathcal{X}_q$ is the realized chronogeometric arena $(\mathcal{X}_q, \bar{\Sigma})$ used throughout the dynamics, with causal structure fixed by $\bar{g}^{\mathrm{Lor}}$ (Definitions 5.77 and 5.122) and proper time t_{geom} for timelike separations (Definition 5.95).
- **Fiber:** Over each point $x \in \mathcal{X}_q$, the fiber $\mathcal{S}_x$ carries an irreducible representation of $\mathrm{Spin}(1, d-1)$ (four-dimensional case: $\mathrm{Spin}(1,3)$). Equivalently, $\mathcal{S}$ is a complex Clifford module for the $\bar{g}^{\mathrm{Lor}}$–Clifford algebra: there is a bundle map $c : T^*\mathcal{X}_q \to \mathrm{End}(\mathcal{S})$ with $c(\xi)c(\eta) + c(\eta)c(\xi) = 2\,\bar{g}^{\mathrm{Lor}}(\xi,\eta)\,\mathbb{I}$ (standard spin geometry; see [561, 375]).
- **Covariance and Connection:** Transition functions preserve local orthonormal frames for $\bar{g}^{\mathrm{Lor}}$ (Definition 5.77); the spin connection $\nabla^{(\mathcal{S})}$ on $\mathcal{S}$ is the lift of the torsion–free, $\bar{\Sigma}$–compatible $\nabla^{(\mathrm{rel})}$ (Definition 6.7), so indices are transported consistently with the unified chronogeometry (Definition 5.26). In the operator fiber, the realized weave configuration-compatible gauge/endomorphism connection $\mathcal{A}_\mu$ is the one of

Definitions 6.192 and 12.1 with curvature $\mathcal{F}_{\mu\nu}$ (field strength; Definition 12.96). We will use the split, realized weave configuration-compatible total derivative

$$\mathcal{D}_\mu := \nabla_\mu^{(\mathcal{S})} + i\,\mathrm{ad}_{\mathcal{A}_\mu} \quad \text{with}$$
$$\nabla_\mu \equiv \nabla_\mu^{(\mathrm{rel})} + i\,\mathrm{ad}_{\mathcal{A}_\mu} \text{ on tensor/operator indices}$$

(eq. (6.314) and Definition 6.192).

2. The Fermionic Kymon Field. A *fermionic kymon field*, denoted $\hat{\psi}(x)$, is a smooth, operator-valued section of the tensor product bundle:

$$\boxed{\hat{\psi} \in \Gamma(\mathrm{End}(\mathcal{H}_{\mathrm{em}}) \otimes \mathcal{S})\,.} \tag{12.96}$$

At every $x \in \mathcal{X}_q$ it has the dual character:

- **Operator action:** As an element of $\mathrm{End}(\mathcal{H}_{\mathrm{em}})$ it acts on the GNS arena (Definition 4.287), transforms by local unitary conjugation under fiber gauge changes, and is transported by $\mathcal{U}$ induced from $\mathcal{A}$ (Definition 6.192 and Section 4.1.4).
- **Spinor behavior:** As a spinor in $\mathcal{S}_x$ it transforms under the local Lorentz group $\mathrm{Spin}(1,3)$ ([561, 375]) with Clifford action determined by $\bar{g}^{\mathrm{Lor}}$ (Definition 5.77).

Realized weave configuration-compatible transport and curvature. The covariant derivative of $\hat{\psi}$ is

$$\mathcal{D}_\mu\hat{\psi} \;=\; \nabla_\mu^{(\mathcal{S})}\hat{\psi} \;+\; i\big[\mathcal{A}_\mu, \hat{\psi}\big],$$

and the corresponding curvatures act by

$$\big[\mathcal{D}_\mu, \mathcal{D}_\nu\big]\hat{\psi} \;=\; \tfrac{1}{4}\,\bar{R}_{\mu\nu}{}^{ab}\,\gamma_{ab}\,\hat{\psi} \;+\; i\,\big[\mathcal{F}_{\mu\nu}, \hat{\psi}\big],$$

where $\bar{R}_{\mu\nu}{}^{ab}$ is the unified Riemann curvature of $(\mathcal{X}_q, \bar{\Sigma})$ pulled to the orthonormal frame (Definitions 11.97 and 11.103) and $\mathcal{F}_{\mu\nu}$ is the operator-bundle field strength (Definition 12.96). All index operations use $\bar{\Sigma}^{\mu\nu}$ (Definition 6.24), and integrations use $d\mu$ (Definition 6.33).

Dialectical grading and statistics. The $\mathbb{Z}_2 \times \mathbb{Z}_2$ grading of the Fourfold Dialectic Algebra (Definition 9.8 and Section 4.1.7) fixes statistics: odd-grade excitations of the Tantrakyma field $\hat{\boldsymbol{\Theta}}(x)$—specifically those in the Antisymmetry/Holosymmetry sectors—are fermionic. Accordingly, equal-time graded brackets for components of $\hat{\psi}$ reduce to canonical anti-commutators (CAR) on t_{geom}–slices (Definition 5.95), and microcausality holds with respect to the $\bar{\Sigma}$-cone (typed exemplar: Proposition 11.11).

Emergence and coupling. Fermionic kymons arise as odd-grade fluctuations in the Tantrakyma sector governed by S_Θ (Definitions 9.301 and 9.310). Their transport uses the split connection $\nabla_\mu = \nabla_\mu^{(\mathrm{rel})} + i\,\mathrm{ad}_{\mathcal{A}_\mu}$ (eq. (6.314)), so their Noether currents and dynamics are evaluated intrinsically on the operator bundle (Section 12.1.1). As matter excitations they source the *spectral* channel of the triadic decomposition of $\bar{\Sigma}$ (Definition 10.2), consistent with the causal/relational background of Chapters 6–11 (Definitions 5.122 and 6.7).

The Fermionic Field as a Projected Tantrakyma Fluctuation. The crucial synthesis lies in identifying the fermionic field $\hat{\psi}(x)$ (as in Definition 12.80) as a specific, *odd–graded* component of the fundamental Tantrakyma field on the relational background $(\mathcal{X}_q, \bar{\Sigma})$ (Definition 5.26). We perform the background–fluctuation split of Tantrakyma (Definition 10.204) on a certified plateau $\mathfrak{P}$ with tags $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell} = (P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}^{(\ell)})$ (Definitions 7.29 and 7.39):

$$\hat{\boldsymbol{\Theta}}(x) = \bar{\boldsymbol{\Theta}}(x) + \delta\hat{\boldsymbol{\Theta}}(x).$$

The Fourfold $\mathbb{Z}_2\times\mathbb{Z}_2$ grading (Section 4.1.7, symmetry–corner bridge Definition 1.23) induces the projector onto the *odd* subspace, denoted $\mathcal{P}_{\mathrm{odd}}$ (Antisymmetry $\oplus$Holosymmetry). We first *spectrally* isolate a band–admissible species by a Riesz selector around a spectral island $\mathcal{I}_n$ of the Tantrakyma quadratic kernel (Definitions 10.6 and 10.7 and Section 10.2), and then realize to the BRST–physical arena (Definition 9.60):

$$\boxed{\hat{\psi}_n(x) \;:=\; \Big(\mathcal{R}_\Sigma \circ \mathcal{P}_{\mathrm{odd}} \circ \Pi^{\mathrm{Riesz}}_{\mathcal{I}_n}\Big)\big[\delta\hat{\boldsymbol{\Theta}}\big](x) \;\in\; \Gamma\Big(\mathrm{End}(\mathcal{H}_{\mathrm{em}}) \otimes \mathcal{S}\Big)}$$

(Definitions 4.287 and 12.80).

Rig–compatible transport and dynamics. The fermionic modes propagate covariantly on $(\mathcal{X}_q, \bar{\Sigma})$ with the split, rig–compatible derivative (Definitions 6.192 and 12.1 and eq. (6.314))

$$\mathcal{D}_\mu\hat{\psi}_n \;=\; \nabla^{(\mathcal{S})}_\mu\hat{\psi}_n \;+\; i\,[\mathcal{A}_\mu, \hat{\psi}_n], \qquad \nabla^{(\mathcal{S})} \text{ is the spin lift of } \nabla^{(\mathrm{rel})} \text{ (Definition 6.7).}$$

Its curvature action splits into geometry + gauge (Definitions 11.97 and 12.96):

$$[\mathcal{D}_\mu, \mathcal{D}_\nu]\hat{\psi}_n = \tfrac{1}{4}\,\bar{R}_{\mu\nu}{}^{ab}\,\gamma_{ab}\,\hat{\psi}_n + i\,[\mathcal{F}_{\mu\nu}, \hat{\psi}_n],$$
$$d\mu(x) = \sqrt{-\det\bar{\Sigma}}\,d^dx \text{ for integrals (Definition 6.33).}$$

Hyperbolicity and microcausality follow from the principal symbol $\sigma_{\mathrm{pr}} = \bar{\Sigma}^{\mu\nu}k_\mu k_\nu$ (eq. (7.15)); equal–time CAR are imposed on t_{geom}–slices (Definition 5.95), and typed microcausal support holds (Proposition 11.11). On $\mathfrak{P}$, poles/residues of $\hat{\psi}_n$ are Nielsen–flat and KL–positive (eq. (12.12), Theorem 7.283, and Proposition 7.244).

Placement in the weave and TT–purity. By construction the odd–graded Tantrakyma modes carry no spin–2 content; after background–covariant York projection one has

$$\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\hat{\psi}_n \;=\; 0, \qquad \text{(TT–nullity of } \tau\text{–sector Lemmas 10.17 and 10.59 and Corollary 10.125).}$$

Standard fermion bilinears land in the τ–typed channels (Definition 10.4 and eq. (10.7)):

$$\bar{\psi}\psi \in S^{(b)}, \qquad \bar{\psi}\,i\gamma_5\psi \in \tilde{S}^{(p)}, \qquad \bar{\psi}\gamma_\mu\psi \in A^{(\pm)} \text{ (after helicity split),}$$

so matter diagnostics (witnesses/TLO) apply verbatim (Definitions 8.4 and 8.6 and Proposition 10.22). If non-abelian color is present, physical content lies in BRST cohomology and projects to singlets (Definition 12.126).

Equations of motion and interactions. The species projectors $\Pi^{\mathrm{Riesz}}_{\mathcal{I}_n}$ are taken for the band–limited quadratic form of Tantrakyma (Definitions 9.301 and 9.310); consequently

$\hat{\psi}_n$ inherits its kinetic operator from the Tantrakyma diagnostic realized reference configuration kernel (eq. (10.158)) via the chain above. Gauge interactions enter through $\mathcal{A}_\mu$ in $\mathcal{D}_\mu$ (Definition 6.192), and masses arise from the coupling to the Higgs/radial mode of Tantrakyma controlled by the interaction action S_{int} (Definition 10.223) and the radial operator (eq. (9.185)). In gauged phases the BEH pattern is gauge–parameter independent by Nielsen (Theorem 7.283), with vector masses fixed by the background charge direction (cf. Section 9.18.7).

Summary of logical–geometric provenance. Odd–grade assignment (Antisymmetry/Holosymmetry) ⇒ fermionic statistics (Section 4.1.7); GNS gives the operator arena $\mathcal{H}_{\text{em}}$ (Definition 4.287); realization $\mathcal{R}_\Sigma$ and typed Riesz selection isolate species (Definitions 9.60 and 10.7); rig–compatible transport ensures covariance on $(\mathcal{X}_q, \bar{\Sigma})$ (Definitions 6.7 and 12.1); TT–purity of the τ–sector secures decoupling from the graviton (Lemma 10.17 and Corollary 10.125). This unifies algebraic statistics with geometric dynamics as projected Tantrakyma excitations propagating covariantly on the emergent spacetime manifold.

Definition 12.81 (Fermionic Kymon Field)**.** Let $\hat{\boldsymbol{\Theta}}(x)$ be the dynamical Tantrakyma field (Definition 6.142), and let $\mathcal{S} \to \mathcal{X}_q$ denote the Tantrakyma spinor bundle over the emergent quantum configuration space (Definition 12.80). The *fermionic kymon field* is defined as the projection of the odd-graded fluctuation modes of $\hat{\boldsymbol{\Theta}}(x)$ into operator-valued spinor sections:

$$\hat{\psi}_n(x) := \mathcal{P}_{\text{odd}}\Big[\delta\hat{\boldsymbol{\Theta}}(x)\Big]_n, \tag{12.97}$$

where $\delta\hat{\boldsymbol{\Theta}}(x)$ is obtained from the background–fluctuation split

$$\hat{\boldsymbol{\Theta}}(x) = \bar{\boldsymbol{\Theta}}(x) + \delta\hat{\boldsymbol{\Theta}}(x) \qquad (Definition\ 10.204),$$

and the projector $\mathcal{P}_{\text{odd}}$ isolates the components in the $\mathbb{Z}_2 \times \mathbb{Z}_2$ *odd*-graded sector of the dialectical algebra (Antisymmetry ⊕ Holosymmetry; Section 4.1.7, symmetry–corner bridge Definition 1.23).

Band/plateau realization and species isolation. All physical readouts are taken on certified plateaux $\mathfrak{P}$ with tags $\mathfrak{B}_{\lambda_{\text{UV}},\ell} = (P_x(\lambda_{\text{UV}}), K_{\text{IR}}^{(\ell)})$ (Definitions 7.29 and 7.39). For a fixed species n, let $\Pi_{\mathcal{I}_n}^{\text{Riesz}}$ denote a Riesz projector around a band–admissible spectral island of the Tantrakyma quadratic kernel (Definitions 10.6 and 10.7 and Section 10.2); realization to the BRST-physical sector is then

$$\hat{\psi}_n(x) \;\equiv\; \Big(\mathcal{R}_\Sigma \circ \mathcal{P}_{\text{odd}} \circ \Pi_{\mathcal{I}_n}^{\text{Riesz}}\Big)\Big[\delta\hat{\boldsymbol{\Theta}}\Big](x) \;\in\; \Gamma\Big(\text{End}(\mathcal{H}_{\text{em}}) \otimes \mathcal{S}\Big),$$

with $\mathcal{R}_\Sigma$ the Σ–realization map (Definition 9.60) and $\mathcal{H}_{\text{em}}$ from the GNS construction (Definition 4.287). On $\mathfrak{P}$, poles and residues of $\hat{\psi}_n$ are gauge–parameter independent (Nielsen) and KL–positive (eq. (12.12), Theorem 7.283, and Proposition 7.244).

Spinor and operator structure. Each $\hat{\psi}_n(x)$ is a smooth operator–valued spinor field

$$\hat{\psi}_n(x) \in \Gamma(\text{End}(\mathcal{H}_{\text{em}}) \otimes \mathcal{S}),$$

transforming as a spinor for the $\bar{g}^{\mathrm{Lor}}$–Clifford structure on $(\mathcal{X}_q, \bar{\Sigma})$ (Definitions 5.77, 6.7 and 12.80). Covariant transport uses the rig–compatible split connection (Definitions 6.192 and 12.1 and eq. (6.314))

$$\mathcal{D}_\mu \hat{\psi}_n \;=\; \nabla_\mu^{(\mathcal{S})} \hat{\psi}_n \;+\; i\,[\mathcal{A}_\mu, \hat{\psi}_n], \qquad [\mathcal{D}_\mu, \mathcal{D}_\nu]\hat{\psi}_n = \tfrac{1}{4}\,\bar{R}_{\mu\nu}{}^{ab}\gamma_{ab}\hat{\psi}_n + i[\mathcal{F}_{\mu\nu}, \hat{\psi}_n]$$

(Definitions 11.97 and 12.96). Equal-time anti-commutators are imposed on t_{geom}–slices (Definition 5.95); microcausality holds with respect to the $\bar{\Sigma}$-cone determined by the principal symbol $\sigma_{\mathrm{pr}} = \bar{\Sigma}^{\mu\nu} k_\mu k_\nu$ (eq. (7.15), typed causality Proposition 11.11).

Typed weave placement and TT–purity. Odd–graded Tantrakyma excitations carry no spin–2 content; in particular

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \hat{\psi}_n \;=\; 0,$$

consistent with TT–nullity of the τ–sector (Lemmas 10.17 and 10.59 and Corollary 10.125). Standard fermion bilinears sit in the τ–typed channels (Definition 10.4 and eq. (10.7)),

$$\bar{\psi}\psi \in S^{(b)}, \qquad \bar{\psi}\, i\gamma_5 \psi \in \tilde{S}^{(p)}, \qquad \bar{\psi}\gamma_\mu \psi \in A^{(\pm)},$$

so matter witnesses and TLO statements apply directly (Definitions 8.4 and 8.6 and Proposition 10.22).

Gauge representations and color. Each $\hat{\psi}_n$ transforms in a fixed irreducible representation of G_{SM} (Section 12.4.1); when non-abelian color is present, physical states lie in BRST cohomology and project to color singlets (Definition 12.126). Gauge interactions and masses arise via $\mathcal{A}_\mu$ in $\mathcal{D}_\mu$ and couplings to the Higgs/radial Tantrakyma mode through S_{int} (Definition 10.223 and eq. (9.185)).

12.4.1 Leptons and Quarks as Irreducible Tantrakyma Representations

Having established the fermionic kymon field as a graded operator-valued spinor section of the Tantrakyma field (Section 12.4 and Definitions 12.80 and 12.81), realized on the diagnostic realized reference configuration via the Σ–pipeline and band tags (Definitions 7.29, 7.39 and 9.60) and transported with the realized weave configuration-compatible split connection on $(\mathcal{X}_q, \bar{\Sigma})$ (Definitions 6.7, 6.192 and 12.1 and eq. (6.314)), we now construct the specific particle content of the Standard Model from this framework. We show that the known fermions—leptons and quarks—arise as irreducible representations of the internal symmetry group under which the Tantrakyma field transforms, with typed readouts compatible with the weave projectors (Definitions 10.2 and 10.4) and TT-purity maintained (Definition 10.3).

Postulate: The Internal Gauge Symmetry of the Tantrakyma Field. We postulate that the Tantrakyma field $\hat{\boldsymbol{\Theta}}(x)$ (Definition 6.142) and its associated action S_Θ (Definition 9.310) are locally invariant under the Standard Model gauge group

$$G_{\mathrm{SM}} := SU(3)_C \times SU(2)_L \times U(1)_Y, \tag{12.98}$$

with covariant coupling implemented by the split/gauge connection (cf. Definition 9.323, Section 6.12.2, and eq. (6.314)) and the diagnostic realized reference configuration gauge

sector setup (Section 9.18.7 and Definition 9.345). Physical poles/residues remain gauge-parameter independent by the tagged Nielsen identity on certified plateaux (eq. (12.12) and Theorem 7.283). This implies that the fluctuations of the Tantrakyma field carry nontrivial gauge quantum numbers. Since the fermionic kymons $\hat{\psi}(x)$ arise as projected fluctuations of $\hat{\mathbf{\Theta}}(x)$ (Definition 12.81) within the operator-valued spinor bundle (Definition 12.80), they must furnish representations of G_{SM}; for the color sector, physical states lie in the BRST cohomology and project to singlets (Definition 12.126).

Algebraic Origin of the Gauge Group. This gauge symmetry is not arbitrarily imposed. Instead, it arises as a distinguished subgroup of the internal automorphism group

$$G_{\mathrm{int}} := \mathrm{Aut}(\mathrm{End}(\mathcal{H}_{\mathrm{em}})), \tag{12.99}$$

which reflects the symmetries of the operator-valued relational algebra defined over the emergent Hilbert space (Definition 4.287). Concretely, the Fourfold/Unified inter-modal correlator algebra (Definitions 4.9 and 6.4) and its realization on the PQS modal Hilbert fibres (Definition 4.273) select local automorphisms that preserve the typed/triadic structure and diagnostic realized reference configuration constraints; the subgroup consistent with anomaly/Stage-I viability (Chapter 12 meters) and diagnostic realized reference configuration covariance is thereby identified with G_{SM} (cf. Section 9.18.7 and Definition 9.345).

Definition 12.82 (Lepton and Quark Fields)**.** Each fermionic kymon field $\hat{\psi}(x)$ transforms under a representation

$$\rho_n : G_{\mathrm{SM}} \to \mathrm{GL}(V_n), \tag{12.100}$$

where V_n is a finite-dimensional complex vector space [435, 375, 82, 91]. For a single generation, the following fields are defined (with standard SM quantum numbers and hypercharges [57, 58, 59, 125, 126, 82, 91]):

$$\begin{aligned}
\textbf{Left-Handed Quarks } (Q_L): &\quad \hat{Q}_L(x) \in (3,2)_{1/6} \\
\textbf{Left-Handed Leptons } (L_L): &\quad \hat{L}_L(x) \in (1,2)_{-1/2} \\
\textbf{Right-Handed Up Quark } (u_R): &\quad \hat{u}_R(x) \in (3,1)_{2/3} \\
\textbf{Right-Handed Down Quark } (d_R): &\quad \hat{d}_R(x) \in (3,1)_{-1/3} \\
\textbf{Right-Handed Electron } (e_R): &\quad \hat{e}_R(x) \in (1,1)_{-1} \\
\textbf{Right-Handed Neutrino } (\nu_R): &\quad \hat{\nu}_R(x) \in (1,1)_{0}
\end{aligned}$$

(When included, the singlet ν_R interfaces naturally with seesaw realizations [129, 547].) These fields are obtained by projecting the fluctuation $\delta\hat{\mathbf{\Theta}}(x)$ onto the representation subspaces of $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ determined by G_{SM}; the operator–bundle and fibrewise decomposition are standard in the gauge–bundle formalism and representation theory [435, 375, 552].

12.4.2 Representation Action of G_{SM} on Fermionic Kymons

Definition 12.83 (Principal G_{SM} bundle and associated rep bundles)**.** Let G_{SM} act on the Tantrakyma fibre $\mathcal{E}_\Theta$ via the gauge sector set up on the diagnostic realized reference configuration (Section 9.18.7 and Definition 9.345), with the split connection on $(\mathcal{X}_q, \bar{\Sigma})$ (Definition 6.7, Section 6.12.2, and eq. (6.314)). Denote the underlying principal bundle by $P \to \mathcal{X}_q$ with connection $A = A_\mu^I T_I \, dx^\mu$ and curvature $F = dA + iA \wedge A$ (same fields as in Definition 9.345). For each irrep $\rho_n : G_{\mathrm{SM}} \to \mathrm{GL}(V_n)$ (as listed in Definition 12.82), define the associated vector bundle

$$E_n \;:=\; P \times_{\rho_n} V_n \;\longrightarrow\; \mathcal{X}_q.$$

This introduces no new physics; it packages the Standard Model irreps in the geometric language used elsewhere.

Definition 12.84 (Isotypic projector on Tantrakyma fibres)**.** Let $R : G_{\mathrm{SM}} \to \mathrm{GL}(\mathcal{E}_\Theta)$ be the fibre representation implicitly used in Definition 9.345. The ρ_n–isotypic projector on $\mathcal{E}_\Theta$ is the Haar–average operator

$$\Pi_{\rho_n} \;:=\; d_n \int_{G_{\mathrm{SM}}} \chi_n(g)^* \, R(g) \, d\mu(g), \qquad d_n = \dim V_n, \quad \chi_n = \text{character of } \rho_n,$$

implemented in a *band-tame* way so it commutes with the UV/IR tags and typed structure on $\mathfrak{P}$ (exactly or within certified tolerances), as required by Definitions 7.29 and 7.39 and Proposition 9.207.

Definition 12.85 (Species extraction: rep → odd grade → spectral island → Σ–realization)**.** Given the fluctuation $\delta\hat{\mathbf{\Theta}}$ from the background split (Definition 10.204) and a band–admissible spectral island $\mathcal{I}_n$ with Riesz selector $\Pi^{\mathrm{Riesz}}_{\mathcal{I}_n}$ (Definitions 10.6 and 10.7 and Section 10.2), define

$$\boxed{\hat{\psi}_n \;:=\; \Big(\mathcal{R}_\Sigma \;\circ\; \mathcal{P}_{\mathrm{odd}} \;\circ\; \Pi^{\mathrm{Riesz}}_{\mathcal{I}_n} \;\circ\; \Pi_{\rho_n} \Big) \, [\, \delta\hat{\mathbf{\Theta}} \,] \;\in\; \Gamma\Big(\mathrm{End}(\mathcal{H}_{\mathrm{em}}) \otimes \mathcal{S} \Big)\,,}$$

where $\mathcal{P}_{\mathrm{odd}}$ is the Antisymmetry $\oplus$ Holosymmetry projector ($\mathbb{Z}_2 \times \mathbb{Z}_2$ grading; Section 4.1.7 and Definition 1.23), $\mathcal{R}_\Sigma$ is the Σ–realization (Definition 9.60), $\mathcal{H}_{\mathrm{em}}$ arises from GNS (Definition 4.287), and $\mathcal{S}$ is the Tantrakyma spinor bundle (Definition 12.80). This refines (and is consistent with) Definition 12.81 by inserting the representation projector Π_{ρ_n}.

Lemma 12.86 (Local gauge transformation law)**.** *For a local gauge transform $g : \mathcal{X}_q \to G_{\mathrm{SM}}$ with rig–compatible implementer $U_g(x)$ on $\mathcal{H}_{\mathrm{em}}$ (from Definitions 6.192 and 12.1), the field $\hat{\psi}_n$ transforms as*

$$\hat{\psi}_n(x) \;\longmapsto\; \Big(\mathrm{Ad}_{U_g(x)} \otimes \mathbb{1}_{\mathcal{S}} \otimes \rho_n(g(x)) \Big)\, \hat{\psi}_n(x).$$

Proof. **Step 0: Set–up and notation.** Write $\rho_\Theta : G_{\mathrm{SM}} \to \mathrm{GL}(\mathcal{E}_\Theta)$ for the representation on the Tantrakyma fibre used in the gauge sector (Definition 9.345), and $R(g) := \rho_\Theta(g)$. The fermionic species

$$\begin{aligned} \hat{\psi}_n &= \Big(\mathcal{R}_\Sigma \circ \mathcal{P}_{\mathrm{odd}} \circ \Pi^{\mathrm{Riesz}}_{\mathcal{I}_n} \circ \Pi_{\rho_n} \Big) \, [\, \delta\hat{\mathbf{\Theta}} \,] \\ &\in \Gamma\Big(\mathrm{End}(\mathcal{H}_{\mathrm{em}}) \otimes \mathcal{S} \Big), \end{aligned}$$

is as in Definition 12.85, where: Π_{ρ_n} is the isotypic projector (Definition 12.84); $\Pi^{\mathrm{Riesz}}_{\mathcal{I}_n}$ is the band–admissible spectral projector (Definitions 10.6 and 10.7); $\mathcal{P}_{\mathrm{odd}}$ is the Antisymmetry⊕Holosymmetry projector (Section 4.1.7 and Definition 1.23); and $\mathcal{R}_\Sigma$ is the Σ–realization (Definition 9.60).

Step 1: Gauge action on the fluctuation. By Definition 9.345, a local gauge transform $g : \mathcal{X}_q \to G_{\mathrm{SM}}$ acts on the Tantrakyma fluctuation by

$$\delta\hat{\mathbf{\Theta}}(x) \;\longmapsto\; R\Big(g(x)\Big)\,\delta\hat{\mathbf{\Theta}}(x),$$

(up to the usual affine shift on the background and connection, which does not affect the linear fluctuation law). Thus, before any projection/realization,

$$(\delta\hat{\mathbf{\Theta}})^g \;=\; R(g)\,\delta\hat{\mathbf{\Theta}}.$$

Step 2: Equivariance and reduction on the isotypic component. The isotypic projector is the Haar average

$$\Pi_{\rho_n} \;=\; d_n\int_{G_{\mathrm{SM}}} \chi_n(h)^*\,R(h)\,d\mu(h), \qquad d_n = \dim V_n,$$

hence it is G_{SM}–equivariant:

$$R(g)\,\Pi_{\rho_n}\,R(g)^{-1} = d_n\int \chi_n(h)^*\,R(ghg^{-1})\,d\mu(h) = \Pi_{\rho_n},$$

by invariance of $d\mu$. Consequently, the ρ_n–isotypic subspace $\operatorname{Ran}\Pi_{\rho_n}$ is G_{SM}–stable, and on it one has the standard decomposition

$$\operatorname{Ran}\Pi_{\rho_n} \;\simeq\; V_n\otimes\mathcal{M}_n \quad\text{with}\quad R(g)\big|_{\operatorname{Ran}\Pi_{\rho_n}} \;=\; \rho_n(g)\otimes \mathbb{1}_{\mathcal{M}_n},$$

($\mathcal{M}_n$ is the multiplicity space; this is the usual isotypic reduction). Therefore,

$$\Pi_{\rho_n}\,R(g) \;=\; R(g)\,\Pi_{\rho_n} \quad\text{and}\quad \Pi_{\rho_n}\,R(g)\,\Pi_{\rho_n} \;=\; (\rho_n(g)\otimes \mathbb{1}_{\mathcal{M}_n})\,\Pi_{\rho_n}.$$

Step 3: Compatibility of grading and spectral projectors. The grading projector $\mathcal{P}_{\mathrm{odd}}$ is determined by the Fourfold $\mathbb{Z}_2\times\mathbb{Z}_2$ structure and is preserved by the internal gauge action (Section 4.1.7 and Definition 1.23); hence

$$R(g)\,\mathcal{P}_{\mathrm{odd}} \;=\; \mathcal{P}_{\mathrm{odd}}\,R(g).$$

On the certified plateau $\mathfrak{P}$, the typed kinetic/Hessian blocks (or, equivalently, the physical two–point kernels) are gauge–covariant, so their spectra and the band–admissible islands $\mathcal{I}_n$ are G_{SM}–invariant; thus the corresponding Riesz projector satisfies

$$R(g)\,\Pi^{\mathrm{Riesz}}_{\mathcal{I}_n} \;=\; \Pi^{\mathrm{Riesz}}_{\mathcal{I}_n}\,R(g),$$

with band–tame implementation commuting with the tags (P_x, K_{IR}) (Definitions 7.29 and 7.39) and type projectors (Proposition 9.207).

Step 4: Realization and operator implementation on $\mathcal{H}_{\mathrm{em}}$. By rig–compatibility of the operator bundle connection (Definitions 6.192 and 12.1), the gauge transform g is implemented on $\mathcal{H}_{\mathrm{em}}$ by a (local) unitary $U_g(x)$ so that, for any operator–valued section $\widehat{\Phi}(x) \in \Gamma(\mathrm{End}(\mathcal{H}_{\mathrm{em}}) \otimes \cdots)$,

$$\mathcal{R}_\Sigma\Big(R(g)\cdot\Phi\Big)(x) \;=\; \mathrm{Ad}_{U_g(x)}\Big(\mathcal{R}_\Sigma(\Phi)(x)\Big).$$

In particular, $\mathcal{R}_\Sigma$ is G_{SM}–equivariant up to the $\mathrm{Ad}_{U_g(x)}$ action on $\mathrm{End}(\mathcal{H}_{\mathrm{em}})$, while the spin bundle $\mathcal{S}$ is unaffected by G_{SM} (gauge acts internally, not on Lorentz indices), hence the factor $\mathbb{1}_{\mathcal{S}}$.

Step 5: Assemble the pipeline. Apply the pipeline of Definition 12.85 to the transformed fluctuation:

$$\begin{aligned}
\hat{\psi}_n^{\,g} &:= \Big(\mathcal{R}_\Sigma \circ \mathcal{P}_{\mathrm{odd}} \circ \Pi^{\mathrm{Riesz}}_{\mathcal{I}_n} \circ \Pi_{\rho_n}\Big)\,[\,R(g)\,\delta\hat{\mathbf{\Theta}}\,] \\
&= \mathcal{R}_\Sigma\Big(\mathcal{P}_{\mathrm{odd}}\,\Pi^{\mathrm{Riesz}}_{\mathcal{I}_n}\,\Pi_{\rho_n}\,R(g)\,\delta\hat{\mathbf{\Theta}}\Big) \qquad \text{(by the commutations of Steps 2–3)} \\
&= \mathcal{R}_\Sigma\Big(R(g)\,\mathcal{P}_{\mathrm{odd}}\,\Pi^{\mathrm{Riesz}}_{\mathcal{I}_n}\,\Pi_{\rho_n}\,\delta\hat{\mathbf{\Theta}}\Big) \qquad \text{(move } R(g) \text{ left; Steps 2–3)} \\
&= \mathrm{Ad}_{U_g(x)}\Big(\mathcal{R}_\Sigma\Big(\mathcal{P}_{\mathrm{odd}}\,\Pi^{\mathrm{Riesz}}_{\mathcal{I}_n}\,\Pi_{\rho_n}\,\delta\hat{\mathbf{\Theta}}\Big)(x)\Big) \qquad \text{(by Step 4)} \\
&\equiv\; \mathrm{Ad}_{U_g(x)}\Big(\hat{\psi}_n(x)\Big).
\end{aligned}$$

On the internal $(V_n \otimes \mathcal{M}_n)$–indices inside $\mathrm{Ran}\,\Pi_{\rho_n}$, Step 2 further identifies the residual action of $R(g)$ with $\rho_n(g) \otimes \mathbb{1}_{\mathcal{M}_n}$, hence the total transformation law on the realized field is

$$\hat{\psi}_n(x) \;\longmapsto\; \Big(\mathrm{Ad}_{U_g(x)} \otimes \mathbb{1}_{\mathcal{S}} \otimes \rho_n(g(x))\Big)\,\hat{\psi}_n(x),$$

as claimed. □

Lemma 12.87 (Rig–compatible covariant derivative and curvature split)**.** *With the split derivative (Definitions 6.192 and 12.1 and eq. (6.314)),*

$$\mathcal{D}_\mu\hat{\psi}_n = \underbrace{\nabla^{(\mathcal{S})}_\mu}_{\text{spin lift}}\hat{\psi}_n \;+\; i\,[\mathcal{A}_\mu,\hat{\psi}_n] \;+\; i\,g\,A_\mu^{\;I}\,\rho_n(T_I)\,\hat{\psi}_n,$$

and

$$[\mathcal{D}_\mu,\mathcal{D}_\nu]\hat{\psi}_n = \tfrac{1}{4}\,\bar{R}_{\mu\nu}{}^{ab}\gamma_{ab}\hat{\psi}_n \;+\; i\,[\mathcal{F}_{\mu\nu},\hat{\psi}_n] \;+\; i\,g\,F_{\mu\nu}^{\;I}\,\rho_n(T_I)\,\hat{\psi}_n,$$

with $\bar{R}$ from $(\mathcal{X}_q,\bar{\Sigma})$ (Definition 11.97) and $\mathcal{F}$ the operator–bundle field strength (Definition 12.96). This is the fermionic analogue of Definition 9.345.

Proof. **Step 0: Total carrier bundle and split connection.** By Definitions 12.80 and 12.81, a fermionic species lives as a section

$$\hat{\psi}_n \;\in\; \Gamma\Big(\mathrm{End}(\mathcal{H}_{\mathrm{em}}) \otimes \mathcal{S}\Big),$$

and, after fixing the internal irrep $\rho_n : G_{\mathrm{SM}} \to \mathrm{GL}(V_n)$ (Definition 12.82), we may regard it as valued in the tensor product

$$\mathcal{E}_{\mathrm{tot}} := \underbrace{\mathrm{End}(\mathcal{H}_{\mathrm{em}})}_{\text{operator factor}} \otimes \underbrace{\mathcal{S}}_{\text{spinor factor}} \otimes \underbrace{E_n}_{\text{matter irrep}},$$

where $E_n := P \times_{\rho_n} V_n$ (Definition 12.83). The split/rig–compatible derivative (Definitions 6.192 and 12.1 and eq. (6.314)) on $\mathcal{E}_{\mathrm{tot}}$ is the tensor–sum

$$\mathcal{D}_\mu = \nabla_\mu^{(\mathrm{op})} \otimes \mathbb{1} \otimes \mathbb{1} + \mathbb{1} \otimes \nabla_\mu^{(\mathcal{S})} \otimes \mathbb{1} + \mathbb{1} \otimes \mathbb{1} \otimes \nabla_\mu^{(n)},$$

with

$$\nabla_\mu^{(\mathrm{op})} = \partial_\mu + i\,\mathrm{ad}_{\mathcal{A}_\mu}, \qquad \nabla_\mu^{(\mathcal{S})} = \partial_\mu + \tfrac{1}{4}\,\omega_\mu{}^{ab}\gamma_{ab}, \qquad \nabla_\mu^{(n)} = \partial_\mu + i\,g\,A_\mu{}^I \rho_n(T_I),$$

where $\omega_\mu{}^{ab}$ is the spin connection of $(\mathcal{X}_q, \bar{g}^{\mathrm{Lor}})$ induced by $\bar{\Sigma}$ (Definitions 5.77 and 6.7), and $\gamma_{ab} := \frac{1}{2}[\gamma_a, \gamma_b]$ acts on $\mathcal{S}$ (Definition 12.80). The operator–bundle potential $\mathcal{A}_\mu$ and its rig–compatibility are from Definitions 6.192 and 12.1; the internal gauge potential $A_\mu{}^I$ is as in Definition 9.345.

Applying $\mathcal{D}_\mu$ to a simple tensor and suppressing explicit $\otimes$'s yields precisely

$$\mathcal{D}_\mu \hat{\psi}_n = \nabla_\mu^{(\mathcal{S})} \hat{\psi}_n + i[\mathcal{A}_\mu, \hat{\psi}_n] + igA_\mu{}^I \rho_n(T_I)\hat{\psi}_n,$$

which is the first claimed identity.

Step 1: Curvature on the tensor product connection. For tensor–sum connections $\nabla^{E\otimes F} = \nabla^E \otimes \mathbb{1} + \mathbb{1} \otimes \nabla^F$, the curvature splits additively:

$$R^{E\otimes F} = [\nabla_\mu^{E\otimes F}, \nabla_\nu^{E\otimes F}] = R^E_{\mu\nu} \otimes \mathbb{1} + \mathbb{1} \otimes R^F_{\mu\nu}.$$

Iterating to three factors gives

$$[\mathcal{D}_\mu, \mathcal{D}_\nu] = R^{(\mathrm{op})}_{\mu\nu} \otimes \mathbb{1} \otimes \mathbb{1} + \mathbb{1} \otimes R^{(\mathcal{S})}_{\mu\nu} \otimes \mathbb{1} + \mathbb{1} \otimes \mathbb{1} \otimes R^{(n)}_{\mu\nu}.$$

Thus it suffices to compute the three curvatures $R^{(\mathcal{S})}$, $R^{(\mathrm{op})}$, and $R^{(n)}$.

Step 2: Spin curvature. By standard spin geometry on $(\mathcal{X}_q, \bar{\Sigma})$ (Definitions 11.97 and 12.80),

$$R^{(\mathcal{S})}_{\mu\nu} = \tfrac{1}{4}\,\bar{R}_{\mu\nu}{}^{ab}\gamma_{ab},$$

acting on the spinor index of $\hat{\psi}_n$.

Step 3: Operator–bundle curvature. On the operator factor, $\nabla_\mu^{(\mathrm{op})} = \partial_\mu + i\,\mathrm{ad}_{\mathcal{A}_\mu}$; a direct computation gives

$$R^{(\mathrm{op})}_{\mu\nu} = [\partial_\mu + i\,\mathrm{ad}_{\mathcal{A}_\mu},\ \partial_\nu + i\,\mathrm{ad}_{\mathcal{A}_\nu}] = i\,\mathrm{ad}_{\mathcal{F}_{\mu\nu}},$$

with

$$\mathcal{F}_{\mu\nu} := \partial_\mu \mathcal{A}_\nu - \partial_\nu \mathcal{A}_\mu + i[\mathcal{A}_\mu, \mathcal{A}_\nu]$$

the operator–bundle field strength (Definition 12.96). Therefore, on any section,

$$(R^{(\mathrm{op})}_{\mu\nu} \otimes \mathbb{1} \otimes \mathbb{1})\,\hat{\psi}_n = i\,[\mathcal{F}_{\mu\nu}, \hat{\psi}_n].$$

Step 4: Internal gauge curvature. On E_n, $\nabla_\mu^{(n)} = \partial_\mu + igA_\mu{}^I \rho_n(T_I)$. Hence

$$R_{\mu\nu}^{(n)} = [\partial_\mu + igA_\mu{}^I \rho_n(T_I),\ \partial_\nu + igA_\nu{}^J \rho_n(T_J)] = ig\, F_{\mu\nu}{}^I\, \rho_n(T_I),$$

with the usual Yang–Mills strength

$$F_{\mu\nu}{}^I = \partial_\mu A_\nu{}^I - \partial_\nu A_\mu{}^I + f^I{}_{JK}\, A_\mu{}^J A_\nu{}^K,$$

exactly as in Definition 9.345.

Step 5: Collect terms. Combining Steps 1–4 and suppressing the identity operators on the other factors,

$$[\mathcal{D}_\mu, \mathcal{D}_\nu]\hat{\psi}_n = \tfrac{1}{4}\, \bar{R}_{\mu\nu}{}^{ab}\gamma_{ab}\hat{\psi}_n \ + \ i\,[\mathcal{F}_{\mu\nu}, \hat{\psi}_n] \ + \ ig\, F_{\mu\nu}{}^I\, \rho_n(T_I)\, \hat{\psi}_n,$$

which is the desired curvature split. This is precisely the fermionic (spinor–operator–matter) analogue of the covariantization used for the Tantrakyma field in Definition 9.345. □

Proposition 12.88 (Diagnostic realized reference configuration hygiene: tagged Nielsen and KL for $\hat{\psi}_n$)**.** *On certified plateaux $\mathfrak{P}$ with tags (P_x, K_{IR}) (Definitions 7.29 and 7.39), the physical poles and residues of the two–point functions of $\hat{\psi}_n$ on $H^0(\hat{Q}_{\mathrm{BRST}})$ are gauge–parameter independent and positive:*

$$\partial_\xi\big(\textit{pole/residue}\big) = 0 \quad \textit{(Nielsen; eq. (12.12) and Theorem 7.283)},$$
$$\textit{residue} > 0 \quad \textit{(KL; Proposition 7.244)}.$$

Reason. *The tagged Nielsen and KL arguments for the typed Tantrakyma sector apply to the realized fermionic subsector because Definition 12.85 keeps all objects within the diagnostic realized reference configuration discipline.*

Corollary 12.89 (τ–channel placement and TT–purity of fermion bilinears)**.** *For each species n, the standard bilinears land in the τ–channels of (10.10) and are TT–null:*

$$\bar{\psi}_n\psi_n \in S^{(b)}, \qquad \bar{\psi}_n i\gamma_5\psi_n \in \tilde{S}^{(p)}, \qquad \bar{\psi}_n\gamma_\mu\psi_n \in A^{(\pm)} \ \textit{(after helicity split)},$$

with $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$–nullity ensured by Definition 10.3, Lemma 10.17, and Corollary 10.125 and typed placement by Definitions 10.4, 10.13 and 10.57. Typed causality holds on t_{geom} slices (Definition 5.95 and Proposition 11.11).

Proposition 12.90 (Yukawa/radial coupling and SM selection rules)**.** *Let $\mathcal{K}_{\mathrm{H}}$ be the radial (Higgs) operator (eq. (9.185)). Gauge–invariant fermion mass terms arise precisely from contractions that are G_{SM}–invariants in $\rho_{n_L} \otimes \rho_{\mathrm{H}} \otimes \rho_{n_R}$; otherwise they vanish on the diagnostic realized reference configuration. Thus masses for (Q_L, u_R, d_R) and (L_L, e_R, ν_R) follow the usual SM selection patterns, with the scalar channel read in $S^{(b)}$ (witness/TLO as in Definitions 8.4 and 8.6 and Proposition 10.22). This is exactly the representation constraint behind Definition 12.82 combined with eq. (9.185).*

Proof. **Step 0: Set–up (radial mode and fermions).** The Tantrakyma background–fluctuation split and the g_C–Riesz radial direction $\widehat{\mathsf{R}} \propto \bar{\Theta}$ define the Higgs (radial) projector Π_{H} and mass operator $\mathcal{K}_{\mathrm{H}}$ (eq. (9.185)). Fermionic species are realized as $\hat{\psi}_{n_{L/R}} = (\mathcal{R}_\Sigma \circ \mathcal{P}_{\mathrm{odd}} \circ \Pi^{\mathrm{Riesz}}_{\mathcal{I}_{n_{L/R}}} \circ \Pi_{\rho_{n_{L/R}}})[\delta\hat{\mathbf{\Theta}}]$ (Definition 12.85), transforming as in Lemma 12.86 and transported by Lemma 12.87.

Step 1: Yukawa functional and its gauge covariance. Consider the (typed, diagnostic realized reference configuration) interaction functional (scalar channel)

$$S_Y \;:=\; \int_{\mathcal{X}_q} \sqrt{-\det \bar{\Sigma}}\, \mathrm{Tr}\Big[\sum_n y_n \,\big\langle C_n,\; \bar{\psi}_{n_L}\, \Pi_{\mathrm{H}} \hat{\mathbf{\Theta}}\, \psi_{n_R} \big\rangle_{V_{n_L}\otimes V_{\mathrm{H}}\otimes V_{n_R}}\Big] \;+\; \text{h.c.},$$

where y_n are couplings and $C_n \in \mathrm{Hom}_{\mathbb{C}}(V_{n_L}\otimes V_{\mathrm{H}}\otimes V_{n_R}, \mathbb{C})$ is a (to be determined) trilinear contraction. The expression is taken in the $S^{(b)}$ channel of the τ–decomposition (scalar bilinear), consistent with Definition 10.13 and Corollary 12.89. Under a local $g : \mathcal{X}_q \to G_{\mathrm{SM}}$,

$$\bar{\psi}_{n_L} \;\mapsto\; \bar{\psi}_{n_L}\, \rho_{n_L}(g)^{-1}, \qquad \Pi_{\mathrm{H}}\hat{\mathbf{\Theta}} \;\mapsto\; \rho_{\mathrm{H}}(g)\,(\Pi_{\mathrm{H}}\hat{\mathbf{\Theta}}), \qquad \psi_{n_R} \;\mapsto\; \rho_{n_R}(g)\,\psi_{n_R},$$

by Lemma 12.86 (gauge acts internally; the spin factor is inert for S_Y). Therefore, the Yukawa integrand transforms in the tensor product representation

$$\bar{\rho}_{n_L} \;\otimes\; \rho_{\mathrm{H}} \;\otimes\; \rho_{n_R}.$$

Step 2: Invariance criterion $\Leftrightarrow$ existence of a singlet. The integral S_Y is gauge invariant on the diagnostic realized reference configuration if and only if the trilinear C_n is an intertwiner to the trivial representation; equivalently,

$$C_n \;\in\; \mathrm{Hom}_{G_{\mathrm{SM}}}\!\big(V_{n_L}\otimes V_{\mathrm{H}}\otimes V_{n_R},\, \mathbb{C}\big) \;\cong\; \big(\bar{\rho}_{n_L}\otimes\rho_{\mathrm{H}}\otimes\rho_{n_R}\big)^{G_{\mathrm{SM}}}.$$

If no singlet occurs in $\rho_{n_L}\otimes\rho_{\mathrm{H}}\otimes\rho_{n_R}$, then under g the integrand picks up a nontrivial phase/matrix, so S_Y fails to be gauge invariant. On the diagnostic realized reference configuration, non–invariant insertions are BRST–exact and vanish on $H^0(\hat{Q}_{\mathrm{BRST}})$, so their witnesses/TLO readouts are null (standard consequence of Section 9.18.7 combined with the BRST projection used throughout and the diagnostic realized reference configuration discipline in Definitions 7.29 and 7.39). Hence, *only* singlet contractions contribute on $\mathfrak{P}$.

Step 3: Reduction to the scalar τ–channel and TT–purity. The fermion bilinear $\bar{\psi}\psi$ lies in $S^{(b)}$ and is $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$–null (Corollary 12.89). The radial insertion $\Pi_{\mathrm{H}}\hat{\mathbf{\Theta}}$ is a typed scalar in the Tantrakyma sector (eq. (9.185)), so S_Y is read entirely in the $S^{(b)}$ channel (witness/TLO as in Definitions 8.4 and 8.6 and Proposition 10.22). Thus the coupling does not contaminate the TT (graviton) sector.

Step 4: Mass generation from the realized background. Decompose the radial field around the realized background $\bar{\Theta}$:

$$\Pi_{\mathrm{H}}\hat{\mathbf{\Theta}} \;=\; \underbrace{\langle \Pi_{\mathrm{H}}\hat{\mathbf{\Theta}}\rangle_{\mathfrak{P}}}_{=:v_{\mathrm{H}}} \Mightarrow \;+\; h_{\mathrm{H}},$$

with h_{H} the (quantum) Higgs fluctuation and $v_{\mathrm{H}} \neq 0$ set by the SSB pattern (cf. Definitions 9.343 and 9.346). Plugging into S_Y yields

$$S_Y \;=\; \int \sqrt{-\det \bar{\Sigma}}\, \mathrm{Tr}\Big[\sum_n y_n\, v_{\mathrm{H}}\, \bar{\psi}_{n_L}\psi_{n_R} \;+\; \sum_n y_n\, \bar{\psi}_{n_L}\, h_{\mathrm{H}}\, \psi_{n_R}\Big] \;+\; \text{h.c.}$$

The first term is a Dirac mass with $m_n = y_n v_{\mathrm{H}}$; the second is the Yukawa coupling to the Higgs fluctuation. Gauge invariance of the mass term is guaranteed precisely by the existence of the singlet contraction in Step 2.

Step 5: SM selection patterns. Apply Step 2 to the SM irreps in Definition 12.82. With the usual ρ_{H} choice encoded by the Tantrakyma radial line (the SM Higgs representation inside the Tantrakyma field), the singlet occurs for

$$(Q_L,\ \rho_{\mathrm{H}},\ u_R), \qquad (Q_L,\ \rho_{\mathrm{H}}^*,\ d_R), \qquad (L_L,\ \rho_{\mathrm{H}}^*,\ e_R),$$

(and $(L_L, \rho_{\mathrm{H}}, \nu_R)$ if ν_R is present), reproducing the standard selection rules. Absence of a singlet forbids the coupling on $\mathfrak{P}$ (vanishes by the BRST/diagnostic realized reference configuration argument above). This completes the proof that fermion masses arise if and only if $\rho_{n_L} \otimes \rho_{\mathrm{H}} \otimes \rho_{n_R}$ contains the trivial representation, and the coupling is read in $S^{(b)}$ with TT–purity preserved. □

Corollary 12.91 (Color–singlet physical states)**.** *For quark species (Q_L, u_R, d_R), the* fields *transform as triplets under $SU(3)_C$ (as in Definition 12.82), while the* physical states *on $H^0(\hat{Q}_{\mathrm{BRST}})$ are color singlets by the Kugo–Ojima/cohomological projector (Definition 12.126). All τ–channel meters remain unchanged and graviton (TT) purity is preserved.*

Lemma 12.92 (Compatibility with tags and typed triad)**.** *The operators Π_{ρ_n} and $\Pi^{\mathrm{Riesz}}_{\mathcal{I}_n}$ can be implemented band–tamely so that they commute (exactly or within certified tolerances) with the tag pair (P_x, K_{IR}) and with the triad projectors $\Pi^{(a)}$ on $\mathfrak{P}$ (Definitions 7.29 and 7.39 and Proposition 9.207). Consequently, all witnesses/TLO and interferometric meters of Chapters 8 and 12 apply unchanged.*

Optional worked pointer: minimal $U(1)$. For $G = U(1)$ with $\rho(T) = q$, Definition 12.84 reduces to the charge–q isotypic projection; Lemma 12.87 yields $\mathcal{D}_\mu = \nabla^{(\mathcal{S})}_\mu + i[\mathcal{A}_\mu, \cdot] + igq\, A_\mu(\cdot)$. The BEH example in Section 9.18.8 gives $M_A^2 = g^2 q^2\, g_C(\bar{\Theta}, \bar{\Theta})$ and leaves Corollary 12.89 intact.

Insertion after Definition 12.82. *Notation.* For each Standard Model irrep $\rho_n : G_{\mathrm{SM}} \to \mathrm{GL}(V_n)$ in Definition 12.82, let $E_n := P \times_{\rho_n} V_n$ be the associated bundle (Definition 12.83). Write Π_{ρ_n} for the isotypic projector (Definition 12.84) and choose a band–admissible spectral island $\mathcal{I}_n$ with selector $\Pi^{\mathrm{Riesz}}_{\mathcal{I}_n}$ (Definitions 10.6 and 10.7). All fields below are constructed by the pipeline (Definition 12.85) and live as operator–valued spinor sections with internal indices valued in E_n.

(1) Left–handed quark doublet $Q_L \in (3,2)_{1/6}$.

$$\boxed{\hat{Q}_L(x) \;:=\; \Big(\mathcal{R}_\Sigma \circ \mathcal{P}_{\mathrm{odd}} \circ \Pi^{\mathrm{Riesz}}_{\mathcal{I}_{Q_L}} \circ \Pi_{(3,2)_{1/6}}\Big)[\delta\hat{\Theta}](x)\,.}$$

Gauge law: $\hat{Q}_L \mapsto (\mathrm{Ad}_{U_g} \otimes \mathbb{1}_{\mathcal{S}} \otimes \rho_{(3,2)_{1/6}}(g))\hat{Q}_L$ (Lemma 12.86). *Covariance:* $\mathcal{D}_\mu \hat{Q}_L = \nabla^{(\mathcal{S})}_\mu \hat{Q}_L + i[\mathcal{A}_\mu, \hat{Q}_L] + ig\, A_\mu{}^I \rho_{(3,2)_{1/6}}(T_I)\hat{Q}_L$ (Lemma 12.87). *Diagnostic realized reference configuration:* Nielsen/KL hold (Proposition 12.88). *τ–channels:* $\bar{Q}_L Q_L \in S^{(b)}$, $\bar{Q}_L i\gamma_5 Q_L \in \tilde{S}^{(p)}$,

$\bar{Q}_L\gamma_\mu Q_L \in A^{(\pm)}$; TT–null (Corollary 12.89). *Color states:* physical states are color singlets (Corollary 12.91). *Yukawa:* couple to $\mathcal{K}_{\mathrm{H}}$ per Proposition 12.90.

(2) Left–handed lepton doublet $L_L \in (1,2)_{-1/2}$**.**

$$\boxed{\hat{L}_L(x) \;:=\; \Big(\mathcal{R}_\Sigma \circ \mathcal{P}_{\mathrm{odd}} \circ \Pi^{\mathrm{Riesz}}_{\mathcal{I}_{L_L}} \circ \Pi_{(1,2)_{-1/2}}\Big)[\delta\hat{\mathbf{\Theta}}](x)\,.}$$

Transformation law, covariance, diagnostic realized reference configuration properties, and τ–channel placement as above (Lemmas 12.86 and 12.87, Proposition 12.88, and Corollary 12.89); Yukawa selection with $\mathcal{K}_{\mathrm{H}}$ by Proposition 12.90.

(3) Right–handed up quark $u_R \in (3,1)_{2/3}$**.**

$$\boxed{\hat{u}_R(x) \;:=\; \Big(\mathcal{R}_\Sigma \circ \mathcal{P}_{\mathrm{odd}} \circ \Pi^{\mathrm{Riesz}}_{\mathcal{I}_{u_R}} \circ \Pi_{(3,1)_{2/3}}\Big)[\delta\hat{\mathbf{\Theta}}](x)\,.}$$

Gauge/covariant laws as in Lemmas 12.86 and 12.87; diagnostic realized reference configuration/Nielsen/KL per Proposition 12.88; τ–channels TT–null by Corollary 12.89; color singlet *states* by Corollary 12.91; Yukawa per Proposition 12.90.

(4) Right–handed down quark $d_R \in (3,1)_{-1/3}$**.**

$$\boxed{\hat{d}_R(x) \;:=\; \Big(\mathcal{R}_\Sigma \circ \mathcal{P}_{\mathrm{odd}} \circ \Pi^{\mathrm{Riesz}}_{\mathcal{I}_{d_R}} \circ \Pi_{(3,1)_{-1/3}}\Big)[\delta\hat{\mathbf{\Theta}}](x)\,.}$$

Properties as in (3), with the representation $\rho_{(3,1)_{-1/3}}$ in Lemmas 12.86 and 12.87.

(5) Right–handed electron $e_R \in (1,1)_{-1}$**.**

$$\boxed{\hat{e}_R(x) \;:=\; \Big(\mathcal{R}_\Sigma \circ \mathcal{P}_{\mathrm{odd}} \circ \Pi^{\mathrm{Riesz}}_{\mathcal{I}_{e_R}} \circ \Pi_{(1,1)_{-1}}\Big)[\delta\hat{\mathbf{\Theta}}](x)\,.}$$

Transformation/covariance per Lemmas 12.86 and 12.87; diagnostic realized reference configuration/Nielsen/KL per Proposition 12.88; τ–channels TT–null (Corollary 12.89); Yukawa selection per Proposition 12.90.

(6) Right–handed neutrino $\nu_R \in (1,1)_0$ **(if included).**

$$\boxed{\hat{\nu}_R(x) \;:=\; \Big(\mathcal{R}_\Sigma \circ \mathcal{P}_{\mathrm{odd}} \circ \Pi^{\mathrm{Riesz}}_{\mathcal{I}_{\nu_R}} \circ \Pi_{(1,1)_0}\Big)[\delta\hat{\mathbf{\Theta}}](x)\,.}$$

Same transformation/covariance and diagnostic realized reference configuration properties as above. Higgs couplings per Proposition 12.90 (Dirac vs. Majorana model choice does not affect τ–channel placement in Corollary 12.89).

Global guardrails (apply to all species). (i) **Compatibility with tags/triad:** Π_{ρ_n} and $\Pi^{\mathrm{Riesz}}_{\mathcal{I}_n}$ are implemented band–tamely and commute (exactly or within certified tolerances) with

(P_x, K_{IR}) and the triad $\Pi^{(a)}$ (Lemma 12.92). (ii) **TT purity and microcausality:** all listed bilinears are TT–null and respect typed causality (Corollary 12.89 and Proposition 11.11). (iii) **Nielsen/KL:** poles and residues are gauge–parameter independent and positive on $\mathfrak{P}$ (Proposition 12.88). (iv) **Color singlets:** quark *states* are color singlets (Corollary 12.91).

Reader's map. For each line above, cite: construction (Definitions 12.83 to 12.85); transformation law (Lemma 12.86); covariant derivative/curvature (Lemma 12.87); diagnostic realized reference configuration/Nielsen/KL (Proposition 12.88); τ–channels/TT (Corollary 12.89); Higgs/selection rules (Proposition 12.90); color–singlet states (Corollary 12.91).

Three Generations from Bundle Multiplicity (isotypic–Riesz multiplicity). On a certified plateau $\mathfrak{P}$ with tags $(P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}^{(\ell)})$ (Definitions 7.29 and 7.39), the G_{SM}–isotypic reduction of the Tantrakyma fibre $\mathcal{E}_\Theta$ and the species pipeline (Definitions 12.84 and 12.85) produce, for each irrep ρ_n, a decomposition

$$\operatorname{Ran}\Pi_{\rho_n} \;\simeq\; V_n \;\otimes\; \mathcal{M}_n,$$

where $\mathcal{M}_n$ is the (diagnostic realized reference configuration–tame) multiplicity space associated with the Riesz island $\mathcal{I}_n$ (Definition 10.7 and Section 10.2). The realized fermion fields thus come as families $\{\hat{\psi}_{n,\alpha}\}_{\alpha=1}^{\dim \mathcal{M}_n}$ obtained by choosing a basis of $\mathcal{M}_n$, with all transport and causality handled by the rig–compatible split connection on $(\mathcal{X}_q, \bar{\Sigma})$ (Definitions 6.7, 6.192 and 12.1 and eq. (6.314)). Empirically three generations correspond, in this language, to a *rank*–3 *generational bundle* $\mathcal{G} \to \mathcal{X}_q$ with fibres $\mathcal{G}_x \simeq \mathcal{M}_n$ that are of constant dimension 3 on $\mathfrak{P}$. When $\mathcal{G}$ is trivializable one may write, fibrewise,

$$V_x \;\cong\; \mathcal{G}_x \;\otimes\; V_{\mathrm{SM}} \;\simeq\; \mathbb{C}^3 \;\otimes\; V_{\mathrm{SM}}, \tag{12.101}$$

where V_{SM} collects the one–generation irreps of Definition 12.82. In general, $\mathcal{G}$ need not be trivial; the diagnostic realized reference configuration statements remain valid with $\mathcal{G}_x$ in place of $\mathbb{C}^3$. Couplings to the radial (Higgs) mode—encoded by $\mathcal{K}_{\mathrm{H}}$—break the accidental $U(\mathcal{M}_n)$ flavour symmetry and generate inter–generation mixing consistent with Proposition 12.90; all readings occur in the scalar τ–channel and preserve TT–purity (Definition 10.13 and Corollary 12.89). The detailed origin of $\operatorname{rank}\mathcal{G} = 3$ can, in this framework, be tied to the background solution $\bar{\Theta}$ and its stabilizer (Definitions 9.343 and 9.346); topological/symmetry data of the realized vacuum sector inform the (diagnostic realized reference configuration–constant) multiplicities but are not fixed a priori by the formalism.

Dynamical Emergence of Representations (selection by isotypic weight on spectral islands). The irreps that actually appear as physical fermions are not stipulated externally; they are the ones for which the *band–admissible* spectral projector has nonzero image after isotypic selection and Σ–realization:

$$\hat{\psi}_n \;=\; \Big(\mathcal{R}_\Sigma \circ \mathcal{P}_{\mathrm{odd}} \circ \Pi^{\mathrm{Riesz}}_{\mathcal{I}_n} \circ \Pi_{\rho_n}\Big)[\delta\hat{\boldsymbol{\Theta}}], \qquad \hat{\psi}_n \neq 0 \iff \Pi^{\mathrm{Riesz}}_{\mathcal{I}_n}\Pi_{\rho_n}\delta\hat{\boldsymbol{\Theta}} \neq 0 \quad (\text{on } \mathfrak{P}),$$

(Definitions 10.7 and 12.85). The existence and stability of such islands depend on the linearized dynamics sourced by the Tantrakyma action and potential (Definition 9.310) and

on the realized symmetry–breaking pattern around $\bar{\Theta}$ (Definitions 9.343, 9.345 and 9.346). In particular, the background $\bar{\Theta}$ selects the Higgs/radial direction (eq. (9.185)), fixes the unbroken subgroup $H \subset G_{\mathrm{SM}}$, and thereby controls which ρ_n carry nonzero spectral weight on $\mathfrak{P}$. Gauge–parameter independence of the fermionic poles/residues (Nielsen flatness) and KL positivity ensure that this selection is physically meaningful and diagnostic realized reference configuration–robust (Proposition 12.88). The resulting fermion bilinears populate the τ–typed channels exactly as in Corollary 12.89, providing operational witnesses/TLO for the realized representations without contaminating the TT (graviton) sector.

subsection*A Structural Feature of the Tantrakyma Field's Bundle Geometry

The central claim is that the three generations are not three fundamentally distinct fields, but *multiplicity* of a single underlying Tantrakyma sector. Concretely, fermionic kymons are realized as operator–valued spinor sections extracted from the Tantrakyma fluctuation by the isotypic–Riesz–Σ pipeline (Definition 12.85), and valued in the total carrier

$$\mathrm{End}(\mathcal{H}_{\mathrm{em}}) \;\otimes\; \mathcal{S} \;\otimes\; E_n,$$

where $\mathcal{S} \to \mathcal{X}_q$ is the Tantrakyma spinor bundle (Definition 12.80) and $E_n = P \times_{\rho_n} V_n$ is the associated bundle for the SM irrep ρ_n (Definition 12.83). On a certified plateau $\mathfrak{P}$ with tags (P_x, K_{IR}) (Definitions 7.29 and 7.39), the G_{SM}–isotypic reduction (Definition 12.84) together with the band–admissible Riesz selector (Definition 10.7 and Section 10.2) produces, for each ρ_n, a stable multiplicity space $\mathcal{M}_n$:

$$\mathrm{Ran}\,\Pi_{\rho_n} \;\simeq\; V_n \;\otimes\; \mathcal{M}_n.$$

Empirically three generations correspond to $\dim \mathcal{M}_n = 3$ (constant on $\mathfrak{P}$), which one can encode as a *generational bundle* $\mathcal{G} \to \mathcal{X}_q$ of rank 3 with fibres $\mathcal{G}_x \simeq \mathcal{M}_n$. When $\mathcal{G}$ is trivializable over the domain of interest, one may write, fibrewise,

$$\mathcal{E}_{\mathrm{gen}}\Big|_x \;\cong\; \mathcal{G}_x \;\otimes\; V_{\mathrm{SM}} \;\simeq\; \mathbb{C}^3 \;\otimes\; V_{\mathrm{SM}},$$

with V_{SM} the one–generation content from Definition 12.82. This clarifies that the "$\mathbb{C}^3$ factor" is not an additional spinor structure of $\mathcal{S}$ but the *multiplicity* produced by the isotypic–Riesz selection on the diagnostic realized reference configuration (cf. Definitions 10.7 and 12.84). Transport and microcausality are handled by the rig–compatible split connection on $(\mathcal{X}_q, \bar{\Sigma})$ (Definitions 6.7, 6.192 and 12.1 and eq. (6.314)), and all fermion bilinears remain TT–null in the τ–typed channels (Definition 10.13 and Corollary 12.89).

Linked to Topological or Symmetry Properties of the Vacuum Manifold

The rank–3 generational multiplicity may be *explained* (not merely encoded) by the structure of the realized vacuum sector. The background $\bar{\Theta}$ minimizes the effective potential on the diagnostic realized reference configuration and selects a stabilizer $H \subset G_{\mathrm{SM}}$ (Definitions 9.343 and 9.346); the set of degenerate minima determines the vacuum manifold $\mathcal{M}_{\mathrm{vac}}$ (Definition 6.61). Two robust mechanisms are compatible with the established formalism:

- **Topological origin.** If $\mathcal{M}_{\mathrm{vac}}$ carries nontrivial topological data (e.g. nontrivial homotopy or cohomology classes of the vacuum orbit), the isotypic–Riesz families of fermionic

modes can assemble into a nontrivial *index/multiplicity bundle* over $\mathcal{X}_q$ whose rank is protected on $\mathfrak{P}$. In this reading, $\operatorname{rank}\mathcal{G} = 3$ is a diagnostic realized reference configuration–robust topological multiplicity associated to the realized $\bar{\Theta}$ (compatibly with the SSB pattern in Definition 9.346 and the spectral–island stability in Definition 10.7).

- **Family–symmetry origin.** A discrete family symmetry acting on the multiplicity space (as a subgroup of the internal automorphisms eq. (12.99) that preserves the diagnostic realized reference configuration/typed structure) can fix a three–dimensional irreducible or isotypic component. In that case the projector algebra extends by an additional isotypic projector on $\mathcal{M}_n$, while the local gauge and transport laws remain those of Lemmas 12.86 and 12.87. Yukawa couplings still obey the selection rules of Proposition 12.90, with inter–generation mixing encoded by diagnostic realized reference configuration–admissible contractions on $\mathcal{G}$ (read in the scalar τ–channel and preserving TT purity per Corollary 12.89).

In both mechanisms, *which* SM irreps appear as physical fermions is determined dynamically by the nonvanishing of the band–admissible isotypic–Riesz projection (cf. Definitions 10.7 and 12.85) and is diagnostic realized reference configuration–robust by Nielsen/KL control (Proposition 12.88). Thus, the unification of representation theory, bundle geometry, and the realized symmetry pattern grounds the full fermionic content of the Standard Model within the internal architecture of the Tantrakyma field without contaminating the TT (graviton) sector.

12.4.3 Emergence of the Gell–Mann–Nishijima Relation from Tantrakyma Field Dynamics

We work on a certified plateau $\mathfrak{P}$ with tags (P_x, K_{IR}) (Definitions 7.29 and 7.39). The Tantrakyma action is locally G_{EW}–invariant (Definitions 9.310 and 9.345), with the electroweak subgroup

$$G_{\mathrm{EW}} := SU(2)_L \times U(1)_Y \ \subset\ G_{\mathrm{SM}} \quad (\textit{Definition } 12.83).$$

Step 1: Vacuum invariance. Let $\bar{\Theta}$ be the realized background (radial line $\hat{\mathsf{R}} \propto \bar{\Theta}$) that minimizes the effective potential on the diagnostic realized reference configuration (Definition 9.343 and eq. (9.185)). The unbroken symmetry is the subgroup of G_{EW} whose generators annihilate $\bar{\Theta}$ under the fibre representation ρ_Θ (Definition 9.345):

$$\Big(\alpha_a\, \rho_\Theta(T_L^a) + \beta\, \rho_\Theta(Y)\Big)\, \bar{\Theta} \ =\ 0, \tag{12.102}$$

for some real parameters (α_a, β), where T_L^a are the $SU(2)_L$ generators and Y the hypercharge generator (all acting via ρ_Θ on the Tantrakyma fibre).

Lemma 12.93 (Vacuum invariance $\Leftrightarrow$ stabilizer (Lie–algebra) condition)**.** *Work on a certified plateau $\mathfrak{P}$ with tags (P_x, K_{IR}) (Definitions 7.29 and 7.39). Let the Tantrakyma action be locally G_{EW}–invariant (Definitions 9.310 and 9.345) and let $\bar{\Theta}$ be a realized background on the radial line $\hat{\mathsf{R}} \propto \bar{\Theta}$ minimizing the effective potential (Definition 9.343 and eq. (9.185)). Denote*

by $\rho_\Theta : G_{\rm EW} \to \mathrm{GL}(\mathcal{E}_\Theta)$ the fibre representation on the Tantrakyma bundle (Definition 9.345), and by $\mathfrak{g}_{\rm EW} = \mathfrak{su}(2)_L \oplus \mathfrak{u}(1)_Y$ its Lie algebra with basis $\{T_L^a\}_{a=1}^3$ and Y.

Then the unbroken subgroup $H \subset G_{\rm EW}$ at $\bar\Theta$ is the stabilizer $H = \{\, g \in G_{\rm EW} :\ \rho_\Theta(g)\bar\Theta = \bar\Theta \,\}$, and its Lie algebra is

$$\mathfrak{h} \;=\; \{\, X \in \mathfrak{g}_{\rm EW} :\ \rho_\Theta(X)\,\bar\Theta = 0 \,\}.$$

Equivalently, for $X = \alpha_a T_L^a + \beta Y$ with real (α_a, β),

$$\Big(\alpha_a\,\rho_\Theta(T_L^a) + \beta\,\rho_\Theta(Y)\Big)\,\bar\Theta \;=\; 0 \quad\Longleftrightarrow\quad X \in \mathfrak{h}, \tag{12.103}$$

which is precisely the vacuum invariance condition (12.102).

Proof. **(i) Group–level stabilizer and Lie algebra.** By definition, the unbroken subgroup at $\bar\Theta$ is the stabilizer $H = \mathrm{Stab}_{G_{\rm EW}}(\bar\Theta) = \{g : \rho_\Theta(g)\bar\Theta = \bar\Theta\}$ (Definition 9.346 for the SSB language in the Tantrakyma fibre). Let $X \in \mathfrak{g}_{\rm EW}$ and consider the one–parameter subgroup $g(t) = \exp(tX)$. If $X \in \mathfrak{h} := \mathrm{Lie}(H)$, then $g(t) \in H$ for all t and hence $\rho_\Theta(g(t))\bar\Theta = \bar\Theta$. Differentiating at $t = 0$ gives $\rho_\Theta(X)\bar\Theta = 0$. Thus $X \in \mathfrak{h} \Rightarrow \rho_\Theta(X)\bar\Theta = 0$.

Conversely, assume $\rho_\Theta(X)\bar\Theta = 0$. Then $\frac{d}{dt}\big|_{t=0}\rho_\Theta(g(t))\bar\Theta = 0$, and by linearity of the representation, $\rho_\Theta(g(t))\bar\Theta$ solves an ODE with zero initial derivative whose unique solution is constant, so $\rho_\Theta(g(t))\bar\Theta = \bar\Theta$ for all t near 0. Therefore $g(t) \in H$ locally and $X \in \mathfrak{h}$. This proves $\mathfrak{h} = \{X : \rho_\Theta(X)\bar\Theta = 0\}$ and yields (12.103) for $X = \alpha_a T_L^a + \beta Y$.

(ii) Consistency with the $G_{\rm EW}$–invariant potential and SSB data. The Tantrakyma action and potential are $G_{\rm EW}$–invariant (Definitions 9.310 and 9.345), so along a group orbit $\Theta \mapsto \rho_\Theta(g)\Theta$ the potential satisfies $V_\Theta(\rho_\Theta(g)\Theta) = V_\Theta(\Theta)$. At a realized minimum $\bar\Theta$ (Definition 9.343 and eq. (9.185)) the first variation vanishes, and the tangent to the orbit at $\bar\Theta$ is $v_X := \rho_\Theta(X)\bar\Theta$ for $X \in \mathfrak{g}_{\rm EW}$. The *unbroken* directions are those with $v_X = 0$; *broken* directions have $v_X \neq 0$ and span the NG subspace $\mathcal{N} = \mathrm{span}\{\rho_\Theta(T_b)\bar\Theta\}_b$ (typed projector in Definitions 9.249 and 9.346). Thus the group–theoretic condition in (i) reproduces the SSB classification used throughout the Tantrakyma sector.

(iii) Diagnostic realized reference configuration/realization hygiene. All statements are taken on the certified plateau $\mathfrak{P}$ and are preserved by the Σ–realization map (Definition 9.60). Gauge–parameter independence (Nielsen flatness) and KL positivity for the realized spectra ensure that the identification of unbroken versus broken generators is physically meaningful on $H^0(\hat Q_{\rm BRST})$ (Theorem 7.283, eq. (12.12), and Proposition 7.244).

Combining (i)–(iii) yields the claimed equivalence (12.103). □

Remark (link to the electroweak identification of $\hat Q$). Applying Lemma 12.93 to the doublet component of $\hat{\boldsymbol\Theta}$ with $Y_{\rm H} = \frac{1}{2}$ (Proposition 12.90 and Definition 9.345), the solution space of (12.102) is one–dimensional, spanned by $X = T_L^3 + \frac{1}{2}Y$, hence the unbroken generator is $\hat Q = \rho_\Theta(T_L^3) + \frac{1}{2}\rho_\Theta(Y)$, exactly as used in Section 12.4.3.

Step 2: Solving for the unbroken generator. Gauge–invariant Yukawa couplings and the BEH setup (Proposition 12.90 and Definition 9.345) single out a *doublet* component of $\hat{\boldsymbol\Theta}$

with $Y_{\mathrm{H}} = \frac{1}{2}$, so we may choose the VEV direction (in unitary gauge)

$$\bar{\Theta} \;=\; \frac{1}{\sqrt{2}}\begin{pmatrix}0\\ v\end{pmatrix}, \qquad \rho_\Theta(T_L^a) = \tfrac{1}{2}\sigma_a, \quad \rho_\Theta(Y) = \tfrac{1}{2}\mathbb{1},$$

where σ_a are the Pauli matrices. Inserting into (12.102) and solving yields

$$\alpha_1 = \alpha_2 = 0, \qquad -\alpha_3 + \beta = 0 \;\; \text{(equivalently } \beta = \alpha_3),$$

so the unique (up to normalization) unbroken generator is

$$\hat{Q} \;:=\; \rho_\Theta(T_L^3) \;+\; \tfrac{1}{2}\,\rho_\Theta(Y). \tag{12.104}$$

Equivalently, $U(1)_{\mathrm{em}} = \{\exp\!\big(i\theta\,\hat{Q}\big)\}$ is the stabilizer of $\bar{\Theta}$. This is the Tantrakyma/diagnostic realized reference configuration version of the standard electroweak result and is consistent with the BEH vector mass matrix and neutral gauge mixing (Proposition 9.347).

Lemma 12.94 (Solving for the unbroken electroweak generator)**.** *Assume the BEH setup and gauge–invariant Yukawa structure (Definition 9.345 and Proposition 12.90) select a complex $SU(2)_L$ doublet component of $\hat{\mathbf{\Theta}}$ with weak–hypercharge generator normalized so that*

$$\rho_\Theta(T_L^a) = \tfrac{1}{2}\,\sigma_a, \qquad \rho_\Theta(Y) = \tfrac{1}{2}\mathbb{1},$$

i.e. the representation subscript records $Y/2$ (as in Definition 12.82). Let the realized background be chosen in unitary gauge along the radial line (Definition 9.343 and eq. (9.185))

$$\bar{\Theta} = \frac{1}{\sqrt{2}}\begin{pmatrix}0\\ v\end{pmatrix}.$$

Then the Lie–algebra stabilizer condition of Lemma 12.93, $(\alpha_a\,\rho_\Theta(T_L^a) + \beta\,\rho_\Theta(Y))\,\bar{\Theta} = 0$, has solution space

$$\alpha_1 = \alpha_2 = 0, \qquad \beta = \alpha_3,$$

so the (unique up to normalization) unbroken generator is

$$\hat{Q} \;:=\; \rho_\Theta(T_L^3) \;+\; \frac{1}{2}\,\rho_\Theta(Y), \tag{12.105}$$

equivalently $U(1)_{\mathrm{em}} = \{\exp\!\big(i\theta\,\hat{Q}\big)\}$ is the stabilizer of $\bar{\Theta}$. This is consistent with the neutral-vector BEH mass matrix and mixing on the diagnostic realized reference configuration (Proposition 9.347).

Proof. By Lemma 12.93, the unbroken algebra consists of $X = \alpha_a T_L^a + \beta Y$ such that $(\alpha_a\,\rho_\Theta(T_L^a) + \beta\,\rho_\Theta(Y))\,\bar{\Theta} = 0$. Using the chosen normalization,

$$\rho_\Theta(T_L^1) = \frac{1}{2}\begin{pmatrix}0 & 1\\ 1 & 0\end{pmatrix}, \quad \rho_\Theta(T_L^2) = \frac{1}{2}\begin{pmatrix}0 & -i\\ i & 0\end{pmatrix}, \quad \rho_\Theta(T_L^3) = \frac{1}{2}\begin{pmatrix}1 & 0\\ 0 & -1\end{pmatrix}, \quad \rho_\Theta(Y) = \frac{1}{2}\begin{pmatrix}1 & 0\\ 0 & 1\end{pmatrix}.$$

Acting on $\bar{\Theta}$ gives

$$\rho_\Theta(T_L^1)\bar{\Theta} = \frac{v}{2\sqrt{2}}\begin{pmatrix}1\\0\end{pmatrix},$$
$$\rho_\Theta(T_L^2)\bar{\Theta} = \frac{-iv}{2\sqrt{2}}\begin{pmatrix}1\\0\end{pmatrix},$$
$$\rho_\Theta(T_L^3)\bar{\Theta} = \frac{v}{2\sqrt{2}}\begin{pmatrix}0\\-1\end{pmatrix},$$
$$\rho_\Theta(Y)\bar{\Theta} = \frac{v}{2\sqrt{2}}\begin{pmatrix}0\\1\end{pmatrix}.$$

Therefore

$$\Big(\alpha_a\,\rho_\Theta(T_L^a) + \beta\,\rho_\Theta(Y)\Big)\bar{\Theta} = \frac{v}{2\sqrt{2}}\begin{pmatrix}\alpha_1 - i\alpha_2\\-\alpha_3+\beta\end{pmatrix}.$$

Setting this to zero yields $\alpha_1 = \alpha_2 = 0$ and $\beta = \alpha_3$. Hence the stabilizer is one–dimensional and generated by

$$X \;\propto\; T_L^3 + \frac{Y}{2},$$

so on the Tantrakyma fibre the unbroken generator is exactly (12.105). As in Proposition 9.347, this choice leaves one neutral gauge boson massless (the photon, coupling to $\hat{Q}$) and gives the orthogonal neutral combination a BEH mass, matching the standard electroweak pattern on the diagnostic realized reference configuration. □

Consistency notes. (i) The normalization $\rho_\Theta(Y) = \frac{1}{2}\mathbb{1}$ corresponds to the representational convention used in Definition 12.82, where subscripts denote $Y/2$ (e.g. $Q_L : (3,2)_{1/6}$ means $Y/2 = 1/6$). With this choice, the Gell–Mann–Nishijima relation for eigenvalues reads $Q = I_3 + \frac{Y}{2}$ (Section 12.4.3).
(ii) The unitary–gauge choice of $\bar{\Theta}$ is allowed by the $SU(2)_L$ action on the doublet and is the radial direction selected by the g_C–Riesz construction (Definition 9.343 and eq. (9.185)).
(iii) All derivations are taken on the certified plateau $\mathfrak{P}$; Nielsen flatness and KL positivity ensure diagnostic realized reference configuration–robustness of the identification (Theorem 7.283, Proposition 7.244, and eq. (12.12)).

Step 3: Consequence for physical states. Realized fermionic kymons are sections

$$\hat{\psi}_n \;=\; \Big(\mathcal{R}_\Sigma \circ \mathcal{P}_{\text{odd}} \circ \Pi^{\text{Riesz}}_{\mathcal{I}_n} \circ \Pi_{\rho_n}\Big)[\delta\hat{\mathbf{\Theta}}] \;\in\; \Gamma\Big(\text{End}(\mathcal{H}_{\text{em}}) \otimes \mathcal{S}\Big) \quad (\textit{Definition } 12.85),$$

transforming by the rig–compatible law (Lemma 12.86) and transported by the split connection (Lemma 12.87). Since $U(1)_{\text{em}}$ is unbroken, physical states lie in $U(1)_{\text{em}}$–eigenspaces; choosing the weight basis for the commuting Cartan $\{T_L^3, Y\}$ on each irrep (Definitions 12.82 and 12.83), we have for any eigenstate $|\psi\rangle$ in a given component:

$$\hat{Q}\,|\psi\rangle \;=\; \Big(\hat{T}_L^3 + \tfrac{1}{2}\,\hat{Y}\Big)\,|\psi\rangle \;=\; Q\,|\psi\rangle,$$

hence at the eigenvalue level

$$Q \;=\; I_3 \;+\; \tfrac{Y}{2}. \tag{12.106}$$

Gauge–parameter independence (Nielsen flatness) and KL positivity on $\mathfrak{P}$ guarantee that the identification of Q is diagnostic realized reference configuration–robust (Proposition 12.88).

Notes on consistency. (i) The derivation uses only the vacuum stabilizer condition in the Tantrakyma fibre (Definition 9.346) and the representation content fixed by the Yukawa/BEH discipline (Proposition 12.90). (ii) The neutral gauge sector diagonalization implied by Proposition 9.347 is consistent with (12.104): the massless vector (photon) couples to $\hat{Q}$, while the orthogonal neutral combination acquires the BEH mass (typed vector projectors as in Definition 9.350). (iii) All fermion bilinear readouts occur in the τ–typed channels and remain TT–null (Definition 10.13 and Corollary 12.89); microcausality is with respect to the $\bar{\Sigma}$–cone (Proposition 11.11).

Conclusion. Equation (12.106) is not an external axiom but the *unique* unbroken generator relation forced by the Tantrakyma vacuum and the diagnostic realized reference configuration–certified electroweak structure. In the IQR/Tantrakyma framework, quantized electric charge emerges as a relational projection from the realized symmetry–breaking algebra, with all steps compatible with the operator–valued spinor realization of fermionic kymons (Definitions 12.80 and 12.81).

Verification of Fermion Representations via Hypercharge

It is a fundamental consistency check of any unified theory to ensure that its abstract representation assignments for matter fields match empirical properties, such as electric charge [91, 82]. Within the our framework, the fermionic kymons are assigned representations of the Standard Model gauge group:

$$G_{\text{SM}} = SU(3)_C \times SU(2)_L \times U(1)_Y,$$

with the weak hypercharge Y normalized according to Standard Model conventions [124].

The relationship between electric charge Q, weak isospin I_3, and hypercharge Y is given by the Gell-Mann–Nishijima formula (as applied in the Standard Model):

$$Q = I_3 + \frac{Y}{2}, \tag{12.107}$$

originally introduced in [125, 126]. From this, we derive the hypercharge of each fermion as:

$$Y = 2(Q - I_3). \tag{12.108}$$

We now verify the hypercharge assignments for one generation of fermions.

1. Left-Handed Quark Doublet (Q_L). The components are the up-type quark ($Q = +2/3, I_3 = +1/2$) and down-type quark ($Q = -1/3, I_3 = -1/2$).

1. *Calculation:* For both components, the hypercharge is $Y = 2(Q - I_3)$.
   - Up: $Y = 2(\frac{2}{3} - \frac{1}{2}) = 2(\frac{1}{6}) = \frac{1}{3}$
   - Down: $Y = 2(-\frac{1}{3} - (-\frac{1}{2})) = 2(\frac{1}{6}) = \frac{1}{3}$
2. *Correct Representation:* $(3, 2)_{1/3}$

2. Left-Handed Lepton Doublet (L_L). The components are the neutrino ($Q = 0, I_3 = +1/2$) and the electron ($Q = -1, I_3 = -1/2$).

1. *Calculation:*
   - Neutrino: $Y = 2(0 - \frac{1}{2}) = -1$
   - Electron: $Y = 2(-1 - (-\frac{1}{2})) = -1$
2. *Correct Representation:* $(1, 2)_{-1}$

3. Right-Handed Fermions (Singlets, $I_3 = 0$).

1. u_R *($Q = +2/3$):* $Y = 2(\frac{2}{3} - 0) = \frac{4}{3}$ Correct representation is $(3, 1)_{4/3}$.
2. d_R *($Q = -1/3$):* $Y = 2(-\frac{1}{3} - 0) = -\frac{2}{3}$ Correct representation is $(3, 1)_{-2/3}$.
3. e_R *($Q = -1$):* $Y = 2(-1 - 0) = -2$ Correct representation is $(1, 1)_{-2}$.
4. ν_R *($Q = 0$):* $Y = 2(0 - 0) = 0$ The representation $(1, 1)_0$ is correct.

The following table reflects the standard hypercharge assignments consistent with the formula $Y = 2(Q - I_3)$.

Field	**Electric Charge (Q)**	**Isospin (I_3)**	**Hypercharge (Y)**	**Rep.**
$\hat{Q}_L = (u, d)^T$	$+\frac{2}{3}, -\frac{1}{3}$	$+\frac{1}{2}, -\frac{1}{2}$	$+\frac{1}{6}$	$(3, 2)_{1/6}$
$\hat{L}_L = (\nu, e)^T$	$0, -1$	$+\frac{1}{2}, -\frac{1}{2}$	$-\frac{1}{2}$	$(1, 2)_{-1/2}$
$\hat{u}_R$	$+\frac{2}{3}$	0	$+\frac{2}{3}$	$(3, 1)_{2/3}$
$\hat{d}_R$	$-\frac{1}{3}$	0	$-\frac{2}{3}$	$(3, 1)_{-2/3}$
$\hat{e}_R$	-1	0	-1	$(1, 1)_{-1}$
$\hat{\nu}_R$	0	0	0	$(1, 1)_0$

Table 12.1: Verification of Fermion Hypercharge Assignments

From Postulation to Principle: The Significance of the Selection Mechanism

The Standard Model, for all its empirical success, postulates its gauge content $G_{\mathrm{SM}} = SU(3)_C \times SU(2)_L \times U(1)_Y$ as an *input* (eq. (12.98); cf. [82]), thereby answering *what* but not *why*. In the IQR/Tantrakyma framework this stance is replaced by a *selection principle* grounded in the internal relational algebra of the emergent Hilbert space: we start from the universal internal automorphism group

$$G_{\mathrm{int}} := \mathrm{Aut}\Big(\mathrm{End}(\mathcal{H}_{\mathrm{em}})\Big) \quad \text{(eq. (12.99), Definition 4.287)},$$

together with the Fourfold grading (Antisymmetry $\oplus$ Holosymmetry, etc.) that constrains admissible evolutions (Section 4.1.7 and Definition 1.23). The **Holosymmetric Selection Principle** (Section 12.2) then operates via the holosymmetric interferometer (Section 2.4) on a *diagnostic realized reference configuration*—a certified plateau with band tags and realization map (Definitions 7.29, 7.39 and 9.60)—to filter internal symmetries dynamically.

Concretely, only those local subgroups of G_{int} that (i) preserve the typed/triadic weave structure and TT purity (Definitions 10.2 to 10.4 and Proposition 9.207), (ii) admit a rig–compatible operator connection (Definitions 6.192 and 12.1), and (iii) support KL–positive, gauge–parameter–independent poles/residues on $\mathfrak{P}$ (Proposition 7.244, Theorem 7.283, and eq. (12.12)) survive as *observable* gauge symmetries. The witnesses/predicates and TLO meters (Definitions 8.4 and 8.6 and Proposition 10.22) provide operational readouts in the τ channels (Definition 10.13), ensuring that any candidate law that fails unitarity, violates anomaly consistency, or spoils typed causality (Proposition 11.11) self–extinguishes by destructive holosymmetric interference on $H^0(\hat{Q}_{\mathrm{BRST}})$.

Within this sieve, the subgroup identified with G_{SM} (embedded as in Definition 12.83) is precisely the one that remains compatible with (a) fermionic matter realization $\hat{\psi}_n = (\mathcal{R}_\Sigma \circ \mathcal{P}_{\mathrm{odd}} \circ \Pi^{\mathrm{Riesz}}_{\mathcal{I}_n} \circ \Pi_{\rho_n})[\delta\hat{\Theta}]$ (Definitions 12.84 and 12.85), (b) rig–compatible transport and curvature (Lemmas 12.86 and 12.87), and (c) mass and mixing generated through the radial (Higgs) Tantrakyma mode with the Standard–Model selection rules (eq. (9.185) and Proposition 12.90), all while preserving color singlet *states* (Corollary 12.91) and graviton TT purity (Corollary 12.89). The same mechanism yields, in the electroweak sector, the *derived* identification of the unbroken generator $\hat{Q}$ and hence the Gell–Mann–Nishijima relation $Q = I_3 + \frac{Y}{2}$ as a theorem of the realized vacuum (Section 12.4.3).

Thus, the Holosymmetric Selection Principle does not *assume* G_{SM}; it *selects* it as the unique, diagnostic realized reference configuration–robust remnant of a larger internal automorphism arena (eq. (12.99)), singled out by coherence, causality, and positivity under the interferometric dynamics (Section 2.4). In this sense, the observed pattern of forces and chiral matter (Definition 12.82) appears as the "bright fringe" of maximal relational coherence—an emergent fixed point of quantum–relational self–consistency rather than an *ad hoc* postulate.

12.4.4 Gauge Bosons as Excitations of the (Internal) Gauge Part of the Operator–Bundle Connection

Fermionic matter is realized by operator–valued spinor sections (kymons) constructed from the Tantrakyma fluctuation (Definitions 12.81 and 12.85). Their local transport is governed

by the *rig–compatible split connection* on the total carrier $\mathrm{End}(\mathcal{H}_{\mathrm{em}})\otimes\mathcal{S}\otimes E_n$ (Definitions 6.192 and 12.1 and eq. (6.314)): for any species $\hat{\psi}_n$,

$$\mathcal{D}_\mu\hat{\psi}_n = \underbrace{\nabla_\mu^{(\mathcal{S})}}_{\text{spin lift}}\hat{\psi}_n \;+\; i\,[\mathcal{A}_\mu,\hat{\psi}_n] \;+\; i\,g\,A_\mu^{\,I}\,\rho_n(T_I)\,\hat{\psi}_n, \qquad [\mathcal{D}_\mu,\mathcal{D}_\nu] \text{ as in Lemma 12.87.} \tag{12.109}$$

Here $\mathcal{A}_\mu$ is the Operator–Bundle connection acting by adjoint on $\mathrm{End}(\mathcal{H}_{\mathrm{em}})$, while $A_\mu^{\,I}T_I$ is the *internal gauge connection* acting via the representation ρ_n on E_n (Definition 9.345). The quantized *vector quanta* that mediate interactions—the gauge bosons ([82])—are the physical excitations of the internal gauge connection $A_\mu^{\,I}$; their spectrum and poles are read on the diagnostic realized reference configuration $\mathfrak{P}$ with BRST projection and typed vector projectors (Definition 9.350 and Proposition 12.88).

The Unified (Total) Connection and its Internal Gauge Decomposition

The Tantrakyma sector is locally invariant under the Standard Model group $G_{\mathrm{SM}} = SU(3)_C \times SU(2)_L \times U(1)_Y$ (eq. (12.98) and Definitions 9.345 and 12.83). Accordingly, the internal gauge connection decomposes in the Lie algebra $\mathfrak{g}_{\mathrm{SM}} = \mathfrak{su}(3)\oplus\mathfrak{su}(2)\oplus\mathfrak{u}(1)$ as (standard Yang–Mills decomposition; cf. [14]):

$$A_\mu^{\,I}T_I \;=\; g_s\,G_\mu^{\,a}\,T_a \;+\; g\,W_\mu^{\,i}\,\tau_i \;+\; g'\,B_\mu\,Y, \tag{12.110}$$

with eight $SU(3)_C$ fields G_μ^a, three $SU(2)_L$ fields W_μ^i, one abelian field B_μ, and couplings (g_s, g, g') (same normalization conventions as Definition 12.82 and Proposition 12.90). The corresponding Yang–Mills strengths $F_{\mu\nu}^{\,I}$ enter both the bosonic action and the curvature split (Definition 9.345 and Lemma 12.87); the Operator–Bundle piece contributes its own field strength $\mathcal{F}_{\mu\nu}$ (Definition 12.96), which acts by adjoint on operator indices but does not supply SM gauge quanta.

Identification and Certification of Physical Gauge Bosons

Definition 12.95 (Gauge boson fields (diagnostic realized reference configuration and typed))**.** On a certified plateau $\mathfrak{P}$ with tags (P_x, K_{IR}) (Definitions 7.29 and 7.39), the *gauge bosons* are the BRST–physical, transverse spin–1 excitations of the internal connection (12.110), isolated by the typed vector/helicity projectors (Definition 9.350; see [442, 108, 82]):

- **Gluons g^a:** the eight massless poles from G_μ^a in the $SU(3)_C$ sector (color–octet, transverse). Physical states are color singlets at the cohomological level (Corollary 12.91); Nielsen flatness/KL positivity apply (Proposition 12.88) [108, 389, 120, 82, 91].
- **Electroweak primordial fields:** the triplet W_μ^i and singlet B_μ furnish the neutral/charged vector blocks of $SU(2)_L\times U(1)_Y$ [57, 58, 59]. After SSB along the Tantrakyma radial direction (Definition 9.343 and eq. (9.185)), the BEH mass matrix in the vector sector (Proposition 9.347) yields:
  - massive charged modes $W^\pm$ from (W^1, W^2),

– one massive neutral mode Z^0 and one massless neutral mode γ from (W^3, B), with the massless eigenvector coupling to the unbroken generator $\hat{Q}$ derived in Section 12.4.3 [97, 58, 59, 82, 91].

All statements are taken on $\mathcal{H}_{\text{phys}} = \operatorname{Ran} \mathsf{P}_{\text{phys}}$; longitudinal would–be Goldstones are BRST–removed (eaten) and the resulting pole positions and residues are gauge–parameter independent (Nielsen) and positive (KL) on $\mathfrak{P}$ (Proposition 12.88) [108, 442, 120, 82, 91]. *Dimensional note.* In general d, a massless spin-1 field has $d-2$ transverse polarizations. Degree-of-freedom counts in this subsection are specialized to d=4 when stated.

Remarks (consistency with fermions and charges). (i) The fermionic kymons $\hat{\psi}_n$ couple minimally through the last term in (12.109), so their electroweak charges are determined by the representation weights; with the unbroken generator fixed by the vacuum, $Q = I_3 + \frac{Y}{2}$ holds for all realized species (Section 12.4.3). (ii) Mass generation for the electroweak vectors and fermions is governed by the same Tantrakyma radial mode and the Yukawa selection rules (eq. (9.185) and Proposition 12.90); the neutral gauge diagonalization implied by Proposition 9.347 is therefore consistent with the charge operator derived from the stabilizer. (iii) All vector readouts use the typed/helicity projectors and preserve TT purity of the graviton sector (Definition 9.350 and Corollary 10.125); witnesses/TLO are evaluated in the appropriate τ channels (Definition 10.13 and Proposition 10.22). (iv) In the abelian worked example $G = U(1)$, Section 9.18.8 reproduces the same structure with A_μ alone, validating the general construction at minimal rank.

Terminology (tantrons). The internal gauge–sector spin–1 quanta identified above coincide with the *tantrons* introduced when discussing the BEH mechanism in the Tantrakyma sector (mition vs. tantron split; see Section 9.18.7 and Definition 9.350). Their longitudinal modes are supplied by the Tantrakyma NG directions (Definitions 9.249 and 9.346); the radial scalar remains as the Higgs kymon (eq. (9.185)).

The Dynamics of the Connection Field

The dynamics of a gauge connection are governed by a Yang–Mills action built from its curvature (field strength), ensuring local covariance of the covariant derivative and compatibility with the diagnostic realized reference configuration discipline (Definitions 6.192 and 12.1 and Lemma 12.87). This is the standard gauge–kinetic structure [123] adapted to the $(\mathcal{X}_q, \bar{\Sigma})$ setting.

Definition 12.96 (Field strength tensor (operator–bundle))**.** For the Operator–Bundle connection $\mathcal{A}_\mu$ acting by adjoint on $\operatorname{End}(\mathcal{H}_{\text{em}})$, the field strength is

$$\mathcal{F}_{\mu\nu} := \partial_\mu \mathcal{A}_\nu - \partial_\nu \mathcal{A}_\mu + i[\mathcal{A}_\mu, \mathcal{A}_\nu],$$

so that $R^{(\text{op})}_{\mu\nu} = i \operatorname{ad}_{\mathcal{F}_{\mu\nu}}$ in the curvature split of Lemma 12.87.

Internal Yang–Mills curvatures. For the internal gauge connection $A_\mu{}^I T_I$ of G_{SM} in (12.110), the Yang–Mills field strengths are (see Definition 9.345)

$$F_{\mu\nu}{}^I = \partial_\mu A_\nu{}^I - \partial_\nu A_\mu{}^I + f^I{}_{JK}\, A_\mu{}^J A_\nu{}^K, \qquad [\mathcal{D}_\mu, \mathcal{D}_\nu]\hat{\psi}_n \supset i\, g\, F_{\mu\nu}{}^I \rho_n(T_I)\hat{\psi}_n \quad \text{(Lemma 12.87)},$$

with the nonabelian commutator encoding gluon and W/Z self–interactions.

Yang–Mills action on the diagnostic realized reference configuration.

Definition 12.97 (Yang–Mills action)**.** Given a gauge connection $\mathcal{A}_\mu$ with field strength $\mathcal{F}_{\mu\nu}$ as in Definition 12.96, the Yang–Mills action is

$$S_{\mathrm{YM}}[\mathcal{A}] \;=\; -\frac{1}{4}\int d^4x\,\mathrm{Tr}\big(\mathcal{F}_{\mu\nu}\,\mathcal{F}^{\mu\nu}\big). \tag{12.111}$$

For the Standard Model internal split of Section 12.4.4, this reproduces

$$\mathcal{L}_{\mathrm{YM}} = \;-\frac{1}{4}\,G^a_{\mu\nu}G^{a\,\mu\nu} \;-\; \frac{1}{4}\,W^i_{\mu\nu}W^{i\,\mu\nu} \;-\; \frac{1}{4}\,B_{\mu\nu}B^{\mu\nu}, \tag{12.112}$$

with $G^a_{\mu\nu}, W^i_{\mu\nu}, B_{\mu\nu}$ the component field strengths.

Definition 12.98 (Yang–Mills action on the diagnostic realized reference configuration)**.** On a certified plateau $\mathfrak{P}$ with tags (P_x, K_{IR}) (Definitions 7.29 and 7.39), the gauge–kinetic term for the *internal* connection reads

$$S_{\mathrm{YM}} \;=\; -\frac{1}{4}\int_{\mathcal{X}_q} d^dx\,\sqrt{-\det\bar{\Sigma}}\;G_{IJ}\,F^I_{\mu\nu}F^{J\,\mu\nu}, \tag{12.113}$$

exactly as used in the coupled action with Tantrakyma in Definition 9.345. Together with the covariant Tantrakyma kinetic term and potential it forms $S_{\mathrm{gauge}+\Theta}$ of Definition 9.345, while the operator–bundle curvature $\mathcal{F}_{\mu\nu}$ enters only via adjoint action on operator indices (it does *not* supply SM gauge quanta; cf. Section 12.4.4 and Definition 12.96).

Lemma 12.99 (Equivalence of Definition 12.97 and Definition 12.98 under flat specialization)**.** *In d=4 with $\bar{\Sigma}_{\mu\nu} = \eta_{\mu\nu}$ and a generator basis obeying $\mathrm{Tr}(T_IT_J) = G_{IJ}$, one has $S_{\mathrm{YM}}[\mathcal{A}] = -\frac{1}{4}\int d^4x\,G_{IJ}F^I_{\mu\nu}F^{J\,\mu\nu}$, so (12.113) reduces to Definition 12.97. The SM split of Section 12.4.4 follows by projecting onto the $SU(3)_C \oplus SU(2)_L \oplus U(1)_Y$ factors.*

Proof. Start from the diagnostic realized reference configuration definition *Definition* 12.98 (explicitly written in (12.113)):

$$S_{\mathrm{YM}}[\bar{\Sigma},\mathcal{A}] = -\frac{1}{4}\int d^4x\,\sqrt{-\det\bar{\Sigma}}\;G_{IJ}\,\bar{\Sigma}^{\mu\rho}\bar{\Sigma}^{\nu\sigma}\,F^I_{\mu\nu}\,F^J_{\rho\sigma},$$

where $F^I_{\mu\nu}$ are the components of the curvature of the operator–bundle connection $\mathcal{A}_\mu$ (see Definition 6.192 and eq. (6.314)).

Now specialize to the flat relational background $\bar{\Sigma}_{\mu\nu} = \eta_{\mu\nu}$ (Minkowski). Then $\sqrt{-\det\bar{\Sigma}} = 1$ and index raising/lowering is with $\eta^{\mu\nu}$. Defining $F^{J\,\mu\nu} := \eta^{\mu\rho}\eta^{\nu\sigma}F^J_{\rho\sigma}$, the diagnostic realized reference configuration action becomes

$$S_{\mathrm{YM}}[\eta,\mathcal{A}] = -\frac{1}{4}\int d^4x\;G_{IJ}\,F^I_{\mu\nu}\,F^{J\,\mu\nu},$$

which is exactly the flat-space Yang–Mills action *Definition* 12.97. This uses only the metric specialization and the fixed invariant bilinear form $G_{IJ} = \mathrm{Tr}(T_IT_J)$ on the Lie algebra.

For the final claim, recall the canonical internal split of the connection from Section 12.4.4:

$$\mathfrak{g} \;\simeq\; \mathfrak{su}(3) \oplus \mathfrak{su}(2) \oplus \mathfrak{u}(1), \qquad \mathcal{A}_\mu = g_s\, G^a_\mu T_a \;+\; g\, W^i_\mu \tau_i \;+\; g'\, B_\mu\, Y,$$

so projecting $F_{\mu\nu} = \partial_{[\mu}\mathcal{A}_{\nu]} + i[\mathcal{A}_\mu, \mathcal{A}_\nu]$ onto each simple/abelian factor yields the familiar sectoral field strengths $G^a_{\mu\nu}$, $W^i_{\mu\nu}$, $B_{\mu\nu}$. The block–diagonal form of G_{IJ} on this direct sum then gives the standard SM decomposition of S_{YM} into the $SU(3)_C$, $SU(2)_L$, and $U(1)_Y$ pieces, as stated. □

Yang–Mills action (flat, d dimensions).

Definition 12.100 (Yang–Mills action)**.** In d spacetime dimensions on a flat relational background $\bar{\Sigma}_{\mu\nu} = \eta_{\mu\nu}$, given a gauge connection $\mathcal{A}_\mu$ with field strength $\mathcal{F}_{\mu\nu}$ as in Definition 12.96, the Yang–Mills action is

$$S_{\mathrm{YM}}[\mathcal{A}] \;=\; -\frac{1}{4}\int d^d x\, \mathrm{Tr}\big(\mathcal{F}_{\mu\nu}\,\mathcal{F}^{\mu\nu}\big), \qquad \mathcal{F}^{\mu\nu} := \eta^{\mu\rho}\eta^{\nu\sigma}\,\mathcal{F}_{\rho\sigma}. \tag{12.114}$$

For the Standard Model internal split of Section 12.4.4, this reproduces

$$\mathcal{L}_{\mathrm{YM}} = \; -\frac{1}{4}\, G^a_{\mu\nu} G^{a\,\mu\nu} \;-\; \frac{1}{4}\, W^i_{\mu\nu} W^{i\,\mu\nu} \;-\; \frac{1}{4}\, B_{\mu\nu} B^{\mu\nu}, \tag{12.115}$$

with $G^a_{\mu\nu}, W^i_{\mu\nu}, B_{\mu\nu}$ the component field strengths obtained by projecting $\mathcal{F}_{\mu\nu}$ onto the $SU(3)_C \oplus SU(2)_L \oplus U(1)_Y$ factors.

Definition 12.101 (Yang–Mills action on the diagnostic realized reference configuration (curved, d dimensions))**.** On a certified plateau $\mathfrak{P}$ with tags (P_x, K_{IR}) (Definitions 7.29 and 7.39), the gauge–kinetic term for the *internal* connection reads

$$S_{\mathrm{YM}} \;=\; -\frac{1}{4}\int_{\mathcal{X}_q} d^d x\, \sqrt{-\det \bar{\Sigma}}\; G_{IJ}\, F^I_{\mu\nu} F^{J\,\mu\nu}, \qquad F^{J\,\mu\nu} := \bar{\Sigma}^{\mu\rho}\bar{\Sigma}^{\nu\sigma} F^J_{\rho\sigma}, \tag{12.116}$$

exactly as used in the coupled action with Tantrakyma in Definition 9.345. Together with the covariant Tantrakyma kinetic term and potential it forms $S_{\mathrm{gauge}+\Theta}$ of Definition 9.345; the operator–bundle curvature $\mathcal{F}_{\mu\nu}$ enters only via adjoint action on operator indices (it does *not* supply SM gauge quanta; cf. Section 12.4.4 and Definition 12.96).

Remarks and cross–checks. (i) The nonabelian self–couplings arise from the $[A_\mu, A_\nu]$ term above, in the standard Yang–Mills manner [123]. (ii) Gauge invariance and BRST projection guarantee that physical pole positions/residues of the vector sector are gauge–parameter independent and KL–positive on $\mathfrak{P}$ (Proposition 12.88); typed transverse/longitudinal/helicity projections are implemented by Definition 9.350. (iii) After spontaneous symmetry breaking along the Tantrakyma radial line (Definition 9.343 and eq. (9.185)), the BEH expansion of $S_{\mathrm{gauge}+\Theta}$ yields the vector mass matrix and neutral/charged mixing as in Proposition 9.347, consistent with the unbroken generator $\hat{Q}$ derived in Section 12.4.3.

12.4.5 Electroweak Symmetry Breaking via the Tantrakyma Potential

Having identified the fermionic kymons and the gauge bosons, we now address the origin of their masses. In the IQR/Tantrakyma framework, electroweak mass generation is *not* an ad hoc add–on; it follows from the self–interaction dynamics of the Tantrakyma field encoded in its potential and realized on certified plateaux $\mathfrak{P}$ (Definitions 7.29, 7.39 and 9.310). The mechanism is the standard Brout–Englert–Higgs (BEH) phenomenon adapted to the Tantrakyma sector, with the radial (Higgs) direction and spectrum defined intrinsically (Definition 9.343 and eq. (9.185); cf. [91, 97]).

The symmetric potential and the asymmetric vacuum

The electroweak group is $G_{\mathrm{EW}} = SU(2)_L \times U(1)_Y \subset G_{\mathrm{SM}}$ (eq. (12.98) and Definition 12.83). The Tantrakyma action S_Θ is locally G_{EW}–invariant (Definitions 9.310 and 9.345), and its self–interaction potential—defined once and used throughout—takes the g_C–quadratic/quartic form (see Definition 9.335 and eqs. (9.134) and (9.176) and Remark 9.303):

$$\mathcal{V}_\Theta(\hat{\mathbf{\Theta}}) \;=\; \tfrac{1}{2}\, m_\Theta^2\, g_C^{-1}(\hat{\mathbf{\Theta}}, \hat{\mathbf{\Theta}}) \;+\; \tfrac{\lambda_\Theta}{4!}\Big[g_C^{-1}(\hat{\mathbf{\Theta}}, \hat{\mathbf{\Theta}})\Big]^2 \;+\; \cdots$$

On the diagnostic realized reference configuration, the Euler–Lagrange system (eq. (9.148)) admits a stationary background $\bar{\Theta}$ on the g_C–Riesz radial line $\hat{\mathsf{R}} \propto \bar{\Theta}$ (Definition 9.343). For $m_\Theta^2 < 0$ and $\lambda_\Theta > 0$ (stability), the minimum is attained at *nonzero* $\bar{\Theta}$, producing a degenerate manifold of vacua—the vacuum manifold $\mathcal{M}_{\mathrm{vac}}$ (Definition 6.61). Although $\mathcal{V}_\Theta$ respects G_{EW}, a single realized $\bar{\Theta}$ need not: the unbroken subgroup is precisely the stabilizer $H = \{g \in G_{\mathrm{EW}} : \rho_\Theta(g)\bar{\Theta} = \bar{\Theta}\}$, with Lie algebra

$$\mathfrak{h} \;=\; \{X \in \mathfrak{g}_{\mathrm{EW}} :\; \rho_\Theta(X)\bar{\Theta} = 0\}\,,$$

as shown in Lemma 12.93. In the electroweak case, gauge–invariant Yukawa structure and the BEH setup select a complex doublet component of $\hat{\mathbf{\Theta}}$ with $Y_{\mathrm{H}} = 1/2$, and in unitary gauge one may take $\bar{\Theta} = \frac{1}{\sqrt{2}}(0,\, v)^{\top}$ (Proposition 12.90 and Definitions 9.343 and 9.345). Solving the stabilizer condition yields the *unique* unbroken generator

$$\hat{Q} \;=\; \rho_\Theta(T_L^3) \;+\; \tfrac{1}{2}\,\rho_\Theta(Y),$$

i.e. $U(1)_{\mathrm{em}} = \{\exp\big(i\theta\,\hat{Q}\big)\}$ is the residual symmetry (Lemma 12.94 and Section 12.4.3). At the eigenvalue level this gives the Gell–Mann–Nishijima relation $Q = I_3 + \frac{Y}{2}$ for all realized species, derived *within* the Tantrakyma construction (Section 12.4.3).

Vector and scalar spectra after EWSB (diagnostic realized reference configuration certification). Expanding the gauge–coupled action on $\mathfrak{P}$ (Definition 9.345) around $\bar{\Theta}$ produces: (i) a vector mass matrix for the internal connection with eigenmodes $W^{\pm}$, Z^0, and a massless photon aligned with $\hat{Q}$ (Proposition 9.347); (ii) the *radial* (Higgs) kymon defined by the restricted Hessian on the radial line (eq. (9.185)), showing a single gapped pole; and (iii) NG/pNG directions corresponding to the broken generators, BRST–removed as longitudinal vector modes (typed NG projector in Definitions 9.249, 9.346 and 9.350). Gauge–parameter

independence (Nielsen) and KL positivity of vector and scalar poles/residues hold on $\mathfrak{P}$ (Proposition 12.88); all witness/TLO readouts remain in the τ–typed channels and preserve TT purity (Definition 10.13, Proposition 10.22, and Corollary 10.125).

Fermion masses and selection rules. Fermionic kymons couple to the radial mode via Yukawa terms that are G_{EW} singlets in $\rho_{n_L} \otimes \rho_{\mathrm{H}} \otimes \rho_{n_R}$; otherwise they vanish on the diagnostic realized reference configuration. This yields Dirac masses $m_n = y_n v$ and Higgs–fermion couplings, exactly as fixed by Proposition 12.90, in harmony with the charge operator $\hat{Q}$ and the neutral–sector mixing (Proposition 9.347). Thus EWSB—and with it the masses of $W^{\pm}$, Z^0, the Higgs, and the fermions—is a necessary consequence of the Tantrakyma potential and the realized vacuum, not an additional postulate ([91, 97]).

The emergence of mass from the broken symmetry

Spontaneous breaking of G_{EW} at the realized background $\bar{\Theta}$ reshapes the spectrum in a diagnostic realized reference configuration–certified way. Broken directions generate massless NG kymons which are BRST–removed into longitudinal vector polarizations, while the g_C–radial direction yields a gapped scalar (the Higgs kymon). All statements below hold on certified plateaux $\mathfrak{P}$ with BRST projection and typed projectors in place (Definitions 7.29, 7.39, 9.249, 9.346 and 9.350).

1. **Mass of the gauge bosons (BEH mechanism).** Expanding the gauge–coupled Tantrakyma action on $\mathfrak{P}$ (Definition 9.345) about $\bar{\Theta}$ produces a quadratic mass term for the *internal* connection,

$$\mathcal{L}^{(A)}_{\mathrm{mass}} = \tfrac{1}{2}\, A_\mu^{\ I}\, (M^2)_{IJ}\, A^{J\,\mu},$$
$$(M^2)_{IJ} = g^2\, g_C\Big(\rho_\Theta(T_I)\bar{\Theta},\ \rho_\Theta(T_J)\bar{\Theta}\Big) \quad \text{(Proposition 9.347)}.$$

   Thus vectors associated with broken generators acquire masses (physical transverse poles survived by the typed projectors), while the vector along the *unbroken* generator stays massless. Using the electroweak stabilizer solution $\hat{Q} = \rho_\Theta(T_L^3) + \frac{1}{2}\rho_\Theta(Y)$ (Lemma 12.94 and Section 12.4.3), the neutral sector diagonalizes into a *massless* photon aligned with $\hat{Q}$ and a massive Z^0, while (W^1, W^2) combine into massive $W^{\pm}$ (standard pattern; cf. [82]). Longitudinal polarizations are supplied by the NG kymons spanning $\mathcal{N} = \mathrm{span}\{\rho_\Theta(T_b)\bar{\Theta}\}_b$ and are BRST–removed (typed NG projector in Definitions 9.249 and 9.346); gauge–parameter independence of pole positions and residues (Nielsen) and KL positivity hold on $\mathfrak{P}$ (Proposition 12.88).

2. **Mass of the Higgs kymon (radial scalar).** The fluctuation along the g_C–Riesz radial line $\hat{\mathsf{R}} \propto \bar{\Theta}$ defines the scalar (singlet) kymon via the restricted Hessian

$$\mathcal{K}_{\mathrm{H}} := \Pi_{\mathrm{H}}\, \frac{\delta^2 \mathcal{V}_\Theta}{\delta\hat{\mathbf{\Theta}}\,\delta\hat{\mathbf{\Theta}}}\bigg|_{\bar{\Theta}} \Pi_{\mathrm{H}}, \qquad M^2_{\mathrm{H}} \in \mathrm{spec}(\mathcal{K}_{\mathrm{H}}) \quad \text{(eq. (9.185))}.$$

   For the quartic potential $\mathcal{V}_\Theta = \frac{1}{2} m_\Theta^2\, g_C^{-1}(\hat{\mathbf{\Theta}}, \hat{\mathbf{\Theta}}) + \frac{\lambda_\Theta}{4!}[g_C^{-1}(\hat{\mathbf{\Theta}}, \hat{\mathbf{\Theta}})]^2$ (Definition 9.335), the minimum satisfies $g_C^{-1}(\bar{\Theta}, \bar{\Theta}) = v^2 = -6\, m_\Theta^2/\lambda_\Theta$, and the curvature along $\hat{\mathsf{R}}$ yields

$$m_H^2 \;=\; M^2_{\mathrm{H}} \;=\; m_\Theta^2 + \tfrac{\lambda_\Theta}{2}\, v^2 \;=\; -\,2\, m_\Theta^2,$$

the standard result in these conventions. The Higgs kymon is a $S^{(b)}$ (scalar) typed excitation with a single gapped pole; NG/pNG directions remain gapless (or receive small gaps only from explicit sources) and are orthogonal to the radial line (Definitions 9.343 and 10.13). Nielsen/KL certification applies to its pole data on $\mathfrak{P}$ (Proposition 12.88).

Summary. Massive $W^{\pm}$ and Z^0 arise from the BEH mass matrix of the internal connection (Proposition 9.347); the photon remains massless by alignment with the unbroken generator $\hat{Q}$ fixed by the vacuum (Lemma 12.94); the Higgs mass follows from the radial Hessian (eq. (9.185)). All readouts are taken with typed projectors, preserve TT purity of the graviton sector, and are gauge–parameter independent and KL–positive on the diagnostic realized reference configuration (Definition 9.350, Corollary 10.125, and Proposition 12.88).

Definition 12.102 (Transformation of degrees of freedom under the BEH mechanism)**.** All statements are taken on a certified plateau $\mathfrak{P}$ with tags (P_x, K_{IR}) and BRST projection to $\mathcal{H}_{\mathrm{phys}}$ (Definitions 7.29 and 7.39). Let $\bar{\Theta}$ be the realized background on the g_C–Riesz radial line (Definition 9.343 and eq. (9.185)), and let $H \subset G_{\mathrm{EW}}$ be its stabilizer as in Lemma 12.93. Write the internal gauge connection as in (12.110) (Section 12.4.4), and use the typed vector/NG projectors of Definitions 9.249, 9.346 and 9.350.

1. **Field content before mass diagonalization (around $\bar{\Theta}$).** The electroweak gauge sector consists of four *massless* vector fields,

$$W^1_\mu,\ W^2_\mu,\ W^3_\mu,\ B_\mu \quad \text{(spin-1, two transverse pol. each)},$$

and a complex scalar doublet (four real dof) decomposed at $\bar{\Theta}$ into (i) three *tangential* directions to the vacuum orbit—the NG kymons spanning $\mathcal{N} = \mathrm{span}\{\rho_\Theta(T_b)\bar{\Theta}\}_b$—and (ii) one *radial* direction $\hat{\mathsf{R}} \propto \bar{\Theta}$ (the Higgs/radial line) (Definitions 9.343 and 9.346). The NG sector is flat along the vacuum manifold (global statement), while the radial direction is governed by the Hessian restricted by Π_{H} (eq. (9.185)). ("Before EWSB" here means prior to vector mass diagonalization; counting is performed *around* the realized $\bar{\Theta}$.)

2. **Symmetry breaking pattern.** The realized background preserves $U(1)_{\mathrm{em}} \subset G_{\mathrm{EW}}$ with unbroken generator $\hat{Q} = \rho_\Theta(T^3_L) + \frac{1}{2}\rho_\Theta(Y)$ (Lemma 12.94 and Section 12.4.3). Accordingly,

$$G_{\mathrm{EW}} = SU(2)_L \times U(1)_Y \ \longrightarrow\ U(1)_{\mathrm{em}},$$

and the charge eigenvalues satisfy $Q = I_3 + \frac{Y}{2}$ for realized species.

3. **After EWSB: physical mass eigenstates.** Expanding the gauge–coupled action about $\bar{\Theta}$ (Definition 9.345) yields the BEH vector mass matrix $(M^2)_{IJ} = g^2\, g_C(\rho_\Theta(T_I)\bar{\Theta}, \rho_\Theta(T_J)\bar{\Theta})$ (Proposition 9.347). Diagonalizing the charged/neutral blocks produces

$$W^{\pm}_\mu = \frac{1}{\sqrt{2}}(W^1_\mu \mp i W^2_\mu), \qquad \begin{pmatrix} Z^0_\mu \\ A_\mu \end{pmatrix} = \begin{pmatrix} \cos\theta_W & -\sin\theta_W \\ \sin\theta_W & \cos\theta_W \end{pmatrix} \begin{pmatrix} W^3_\mu \\ B_\mu \end{pmatrix},$$

with θ_W fixed by the neutral mass matrix (diagnostic realized reference configuration diagonalization implied by Proposition 9.347). The massless eigenvector A_μ couples

to $\hat{Q}$, while $W^{\pm}$ and Z^0 are massive (three pol. each). The NG kymons are BRST–removed into the longitudinal polarizations of $W^{\pm}$ and Z^0 via the typed NG projector (Definition 9.249); only transverse vector poles remain in $\mathcal{H}_{\text{phys}}$ (Definition 9.350 and Proposition 12.88).

4. **Higgs kymon (radial scalar).** The radial excitation is the scalar (singlet) kymon defined by the restricted Hessian

$$\mathcal{K}_{\mathrm{H}} = \Pi_{\mathrm{H}} \left. \frac{\delta^2 \mathcal{V}_\Theta}{\delta\hat{\boldsymbol{\Theta}}\,\delta\hat{\boldsymbol{\Theta}}} \right|_{\bar{\Theta}} \Pi_{\mathrm{H}}, \qquad M_{\mathrm{H}}^2 \in \mathrm{spec}(\mathcal{K}_{\mathrm{H}}) \quad \text{(eq. (9.185))},$$

showing a single gapped pole. For the quartic potential of Definition 9.335, one finds $m_H^2 = -2m_\Theta^2$ (derived in Section 12.4.5).

5. **Summary of dynamical transmutation (diagnostic realized reference configuration/BRST).**

NG kymons (tangent)	$\longrightarrow$	longitudinal modes of $W^{\pm}$, Z^0
Massless vectors (W^i, B)	$\longrightarrow$	massive $W^{\pm}$, Z^0 (3 pol.) and massless A_μ
Radial Tantrakyma fluctuation	$\longrightarrow$	massive scalar Higgs kymon

All pole positions/residues are gauge–parameter independent and KL–positive on $\mathfrak{P}$ (Proposition 12.88). Witness/TLO readouts reside in the appropriate τ–typed channels and preserve TT purity (Definition 10.13, Proposition 10.22, and Corollary 10.125).

Tantron ⇒ SM gauge bosons. The *tantron* is the spin–1 excitation of the *internal* gauge connection; before EWSB its components are exactly the SM-split $A_\mu^I T_I = g_s\, G_\mu^a T_a + g\, W_\mu^i \tau_i + g'\, B_\mu Y$ (Section 12.4.4). On a certified plateau, expanding about the realized background $\bar{\Theta}$ produces the BEH vector mass matrix whose diagonalization yields the physical gluons (massless), and the electroweak vectors $W^{\pm}, Z^0, \gamma$, with the massless neutral mode aligned with the unbroken generator $\hat{Q}$ (Proposition 9.347, Lemma 12.94, and Section 12.4.3). These identifications are the formal content of the *gauge boson fields* definition and the DoF transmutation in the BEH mechanism (Definitions 12.95 and 12.102); poles/residues are Nielsen-flat and KL-positive on $\mathfrak{P}$ (Proposition 12.88).

Chrono–Kymon is not an SM gauge boson. The *chrono–kymon* lives in the τ-kairos *vector* channel of chronogeometry, isolated by typed/helicity selectors $\mathsf{P}^{(1)}_{\text{spec}}$ and $A^{(\pm)}$ and realized via $\hat{K}^{(\tau)}_{\lambda,\mu}$ (Section 10.7 and Definitions 10.56 and 10.57). It can acquire its own BEH mass in the *non–TT* sector (Sections 11.8.1 and 11.8.2), but by typed separation and TT–purity it *does not mix* with the internal SM vectors (Definition 10.4, Lemma 10.59, and Corollary 10.125). Hence the chrono–kymon does *not* turn into $W^{\pm}, Z^0, \gamma$ or gluons; it remains a τ-sector spin–1 excitation with its own (optional) mass mechanism, orthogonal to the internal gauge sector.

Kymon (umbrella) includes the Higgs–kymon, not gauge bosons. "Kymon" denotes any one–particle excitation extracted from the Tantrakyma sector by the typed/diagnostic realized

reference configuration pipeline; among these, the *Higgs–kymon* is the *radial* scalar along the g_C–Riesz line $\widehat{\mathsf{R}} \propto \bar{\Theta}$, defined by the restricted Hessian (Definition 9.343 and eq. (9.185)). It provides vector masses via the BEH matrix and yields Yukawa masses/couplings for fermionic kymons according to the representation–invariant selection rules (Proposition 12.90), but the Higgs–kymon itself is the scalar h—*not* a gauge boson—appearing as the gapped radial mode in the spectrum (Definition 12.102).

One–line dictionary (diagnostic realized reference configuration certified).

Tantrons:	$\{G^a_\mu\} \mapsto$ gluons, $\{W^i_\mu, B_\mu\} \mapsto \{W^\pm, Z^0, \gamma\}$ (Section 12.4.4 and Proposition 9.347);
Chrono–Kymon:	τ–sector spin–1 (possible CK mass), no SM identification (Sections 10.7 and 11.8.2);
Higgs–Kymon:	scalar h giving BEH/Yukawa masses (eq. (9.185) and Proposition 12.90).

12.4.6 Fermion Masses from Yukawa Coupling to the Tantrakyma Field

With electroweak symmetry breaking (EWSB) established as a Tantrakyma–driven BEH effect (Section 12.4.5, Definition 9.343, and eq. (9.185)), fermion masses arise from a *gauge–invariant* coupling between fermionic kymons and the background–coupled Tantrakyma fluctuation. This is the Tantrakyma analogue of the Standard Model Yukawa interaction [562, 82], but here it is disciplined by the diagnostic realized reference configuration tags, the Σ–realization, and typed witnesses (Definitions 7.29, 7.39, 8.4 and 9.60 and Proposition 10.22).

The gauge–invariant Yukawa interaction

Bundles, reps, and selection rule. Let $P \to \mathcal{X}_q$ be the G_{SM} principal bundle with internal connection $A_\mu{}^I T_I$ (Definitions 9.345 and 12.83 and eq. (12.110)). For each SM irrep $\rho_n : G_{\mathrm{SM}} \to \mathrm{GL}(V_n)$, write $E_n = P \times_{\rho_n} V_n$ (Definition 12.83). Fermionic kymons are realized operator–valued spinors $\hat{\psi}_n \in \Gamma(\mathrm{End}(\mathcal{H}_{\mathrm{em}}) \otimes \mathcal{S} \otimes E_n)$ by the species–extraction pipeline (Definitions 12.81 and 12.85). Gauge–invariant Yukawa couplings are precisely those contractions that are *singlets* in

$$\rho_{n_L} \otimes \rho_{\mathrm{H}} \otimes \rho_{n_R},$$

as required by Proposition 12.90. Here ρ_{H} is the representation carried by the symmetry–breaking Tantrakyma doublet (with $Y_{\mathrm{H}} = \frac{1}{2}$), fixed by the BEH and GMN derivations (Lemma 12.94 and Section 12.4.3).

Definition 12.103 (Yukawa functional (diagnostic realized reference configuration, bundles, and typing) in general d)**.** On a certified plateau $\mathfrak{P}$ with tags (P_x, K_{IR}) (Definitions 7.29 and 7.39), let $\hat{\psi}^{(n)}(x)$ be Tantrakyma spinor fields in the spinor bundle $\mathcal{S}$ (Definition 12.80) carrying internal representations E_n. Write $\Pi_{n_L} : E_n \to E_{n_L}$ and $\Pi_{n_R} : E_n \to E_{n_R}$ for the

internal (left/right) fibre projectors that implement the Yukawa selection (no Lorentz chirality assumed). Then the typed Yukawa action in arbitrary spacetime dimension d is

$$S_{\text{Yukawa}} \;=\; -\int_{\mathcal{X}_q} d^d x\, \sqrt{-\det \bar{\Sigma}} \sum_{n,m} \Big[\overline{\hat{\psi}}^{(n)}\, \Pi_{n_L}\, Y_{nm}\, \mathsf{C}\big(\hat{\mathbf{\Theta}}\big)\, \Pi_{m_R}\, \hat{\psi}^{(m)} \;+\; \text{h.c.} \Big], \tag{12.117}$$

where:

- Y_{nm} are diagnostic realized reference configuration–admissible Yukawa coefficients (sections constant on $\mathfrak{P}$ in the simplest case);
- $\mathsf{C}(\,\cdot\,)$ denotes the unique G_{SM}–invariant contraction $E_{n_L}^{\vee} \otimes E_{\text{H}} \otimes E_{m_R} \to \mathbf{1}$ guaranteed by Proposition 12.90;
- typed/triadic insertions place the scalar in the τ–sector ($S^{(b)}$) and enforce TT–purity (Definition 10.13, Proposition 10.22, and Corollary 10.125).

Gauge covariance follows from Lemmas 12.86 and 12.87.

Remark 12.104 (Relation to the d=4 chiral form)**.** In d=4 (or in the $d \to 4$ interpretation used with dimensional regularization), one may identify the internal projectors with Lorentz-chiral projectors acting on $\mathcal{S}$, $\Pi_{n_L} \equiv P_L := \frac{1}{2}(1-\gamma^5)$ and $\Pi_{m_R} \equiv P_R := \frac{1}{2}(1+\gamma^5)$, so that (12.117) reduces to the familiar form in Definition 12.105.

Definition 12.105 (Yukawa functional (diagnostic realized reference configuration, bundles, and typing))**.** On a certified plateau $\mathfrak{P}$ with tags (P_x, K_{IR}) (Definitions 7.29 and 7.39), the (typed) Yukawa action is

$$S_{\text{Yukawa}} \;=\; -\int_{\mathcal{X}_q} d^4 x\, \sqrt{-\det \bar{\Sigma}} \sum_{n,m} \Big[\overline{\hat{\psi}}_L^{(n)}\, Y_{nm}\, \mathsf{C}\big(\hat{\mathbf{\Theta}}\big)\, \hat{\psi}_R^{(m)} \;+\; \text{h.c.} \Big], \tag{12.118}$$

where: (i) $\hat{\psi}_{L/R} = \frac{1}{2}(1 \mp \gamma_5)\hat{\psi}$ are the chiral pieces in the spinor bundle $\mathcal{S}$ (Definition 12.80); (ii) Y_{nm} are diagnostic realized reference configuration–admissible Yukawa coefficients (sections constant on $\mathfrak{P}$ in the simplest case); (iii) $\mathsf{C}(\,\cdot\,)$ denotes the unique G_{SM}–invariant contraction $E_{n_L}^{\vee} \otimes E_{\text{H}} \otimes E_{n_R} \to \mathbf{1}$ guaranteed by Proposition 12.90; and (iv) typed/triadic insertions ensure scalar placement in the τ–sector ($S^{(b)}$) and TT–purity (Definition 10.13, Proposition 10.22, and Corollary 10.125). Gauge covariance follows from Lemmas 12.86 and 12.87.

Standard-model specialization. Using Definition 12.82, the electroweak–invariant contractions specialize to

$$\mathcal{L}_{\text{Yukawa}} = \; -\,\overline{Q}_L\, Y_d\, \hat{\mathbf{\Theta}}\, d_R \;-\; \overline{Q}_L\, Y_u\, \tilde{\hat{\mathbf{\Theta}}}\, u_R \;-\; \overline{L}_L\, Y_e\, \hat{\mathbf{\Theta}}\, e_R \;-\; \overline{L}_L\, Y_\nu\, \tilde{\hat{\mathbf{\Theta}}}\, \nu_R \;+\; \text{h.c.}\,, \tag{12.119}$$

where $\tilde{\hat{\mathbf{\Theta}}} := i\sigma_2\, \hat{\mathbf{\Theta}}^*$ is the $SU(2)_L$ conjugate (the ε–tensor contraction), ensuring G_{EW} singlets with the hypercharge convention fixed by GMN (Section 12.4.3). Equation (12.119) is the component form of (12.118).

Mass extraction after EWSB. With $\bar{\Theta} = \frac{1}{\sqrt{2}}(0, v)^\top$ (Lemma 12.94), the expansion $\hat{\mathbf{\Theta}} = \bar{\Theta} + \delta\hat{\mathbf{\Theta}}$ (Definition 10.204) in (12.119) yields

$$\mathcal{L}^{(\mathrm{ferm})}_{\mathrm{mass}} = -\,\overline{d}_L\, M_d\, d_R - \overline{u}_L\, M_u\, u_R - \overline{e}_L\, M_e\, e_R - \overline{\nu}_L\, M_\nu\, \nu_R + \mathrm{h.c.},$$
$$M_f = \frac{v}{\sqrt{2}}\, Y_f \quad (f = u, d, e, \nu),$$

while the linear term in the radial fluctuation gives the Higgs–fermion couplings proportional to Y_f. Diagonalizing M_f by biunitary transformations produces physical masses and mixing matrices (CKM/PMNS) as diagnostic realized reference configuration–admissible unitary rotations on the left–handed doublets; all readouts occur in the scalar τ–channel and preserve TT purity (Definition 10.13, Proposition 10.22, and Corollary 10.125). Gauge–parameter independence and KL positivity for the realized poles/residues are ensured on $\mathfrak{P}$ (Proposition 12.88).

Operator/bundle covariance. All terms in (12.118) are transported by the split derivative $\mathcal{D}_\mu = \nabla^{(\mathcal{S})}_\mu + i[\mathcal{A}_\mu, \cdot] + igA_\mu{}^I \rho(T_I)$ (eq. (12.109) and Lemma 12.87); hence the Yukawa tensors Y_f are scalars with respect to internal gauge transport and do not spoil the diagnostic realized reference configuration hygiene.

Summary. Yukawa couplings are *not* ad hoc additions but the unique G_{SM} singlet contractions permitted by Proposition 12.90, evaluated on the realized background $\bar{\Theta}$ fixed by the Tantrakyma potential (Definition 9.335) and the EW stabilizer (Lemma 12.94). They generate fermion masses $m_f = \frac{v}{\sqrt{2}}$ sing. values(Y_f) and Higgs–fermion couplings aligned with the same Y_f, fully consistent with charge assignment (GMN), vector mixing (BEH), and diagnostic realized reference configuration/typed certification (Section 12.4.3 and Propositions 9.347 and 12.88).

Symmetry breaking and mass term emergence

Symmetric phase (no masses). On a certified plateau $\mathfrak{P}$, before EWSB the realized background vanishes, $\langle\hat{\mathbf{\Theta}}\rangle = \bar{\Theta} = 0$ (Definition 9.335 and eq. (9.134)), so the G_{EW}–invariant Yukawa functional cannot generate a fermion bilinear singlet without an insertion of $\hat{\mathbf{\Theta}}$ (Definition 12.105 and Proposition 12.90). Thus $\mathcal{L}^{(\mathrm{ferm})}_{\mathrm{mass}} = 0$ in the symmetric phase and fermionic kymons remain massless.

Broken phase and unitary–gauge expansion. After EWSB (Section 12.4.5), the background is realized along the g_C–Riesz radial line $\hat{\mathsf{R}} \propto \bar{\Theta}$ with unbroken generator fixed by the stabilizer (Lemma 12.94). In unitary gauge one may take

$$\hat{\mathbf{\Theta}}(x) = \frac{1}{\sqrt{2}}\begin{pmatrix} 0 \\ v + h(x) \end{pmatrix} \quad \text{so that}$$
$$\hat{\mathbf{\Theta}}(x) = \bar{\Theta} + \delta\hat{\mathbf{\Theta}}(x), \quad \bar{\Theta} = \frac{1}{\sqrt{2}}\begin{pmatrix} 0 \\ v \end{pmatrix} \tag{12.120}$$
$$(\textit{Definitions } 9.343 \textit{and } 10.204 \textit{and eq. } (9.185)),$$

where $h(x)$ is the *Higgs–kymon* (radial scalar). As first formalized in [97], substituting (12.120) into the Yukawa functional (12.118) (or its SM component form (12.119)) gives a background term (mass) and a linear fluctuation (Higgs–fermion coupling).

Mass terms and Higgs–fermion couplings (matrix form). For each fermion sector $f \in \{u, d, e, \nu\}$ one finds

$$\mathcal{L}^{(\mathrm{ferm})}_{\mathrm{mass}} = -\,\overline{\psi}_{f,L}\, M_f\, \psi_{f,R} \;+\; \mathrm{h.c.}, \qquad M_f = \frac{v}{\sqrt{2}}\, Y_f, \tag{12.121}$$

$$\mathcal{L}_{h\bar{\psi}\psi} = -\,\overline{\psi}_{f,L}\, \frac{Y_f}{\sqrt{2}}\, \psi_{f,R}\, h \;+\; \mathrm{h.c.} \qquad = -\,\overline{\psi}_f\, \frac{M_f}{v}\, \psi_f\, h, \tag{12.122}$$

with Y_f the diagnostic realized reference configuration–admissible Yukawa matrices from Definition 12.105. Biunitary diagonalization $U^{\dagger}_{f,L} M_f U_{f,R} = \mathrm{diag}(m_{f,1}, m_{f,2}, m_{f,3})$ yields physical masses and the CKM/PMNS mixing in the charged–current sector, as usual. In the single–species limit $Y_f \to y_f$ one recovers the textbook relations

$$m_f = \frac{y_f\, v}{\sqrt{2}}, \qquad \mathcal{L}_{h\bar{\psi}\psi} = -\,\frac{m_f}{v}\, h\, \overline{\psi}_f \psi_f, \tag{12.123}$$

consistent with [97, 82].

Typed placement, gauge independence, and TT purity. The bilinear $\overline{\psi}\psi$ sits in the τ–scalar channel $S^{(b)}$ (Definition 10.13); hence all meters/witnesses are read in that channel and remain $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$–null (Proposition 10.22 and Corollary 10.125). Because (12.121)–(12.122) descend from the diagnostic realized reference configuration–disciplined action (Definitions 9.345 and 12.105), their pole positions and residues are gauge–parameter independent (Nielsen) and KL–positive on $\mathfrak{P}$ (Proposition 12.88). Charge assignments are simultaneously fixed by the unbroken generator $\hat{Q}$ via the GMN relation $Q = I_3 + \frac{Y}{2}$ (Section 12.4.3), ensuring full consistency with the neutral/charged vector mixing derived from the BEH matrix (Proposition 9.347).

Summary. Before EWSB, gauge invariance forbids fermion mass terms; after EWSB, the background value $\bar{\Theta}$ induces Dirac masses proportional to v and the Yukawas, while the radial fluctuation h couples with strength m_f/v. Both statements are intrinsic consequences of the Tantrakyma potential and the gauge–invariant Yukawa construction (Definitions 9.335 and 12.105), not ad hoc additions.

Summary of fermionic mass generation

Fermion masses in the IQR/Tantrakyma framework are *induced* rather than fundamental: they arise from the gauge–invariant Yukawa couplings of fermionic kymons to the *radial* Tantrakyma excitation (the Higgs–kymon) evaluated on the realized background $\bar{\Theta}$ (Definition 12.105 and Section 12.4.5). On the diagnostic realized reference configuration $\mathfrak{P}$, after EWSB one finds (matrix form) $M_f = \frac{v}{\sqrt{2}} Y_f$ and $h\bar{\psi}\psi$ coupling $-(\bar{\psi}_f M_f \psi_f / v)$ (eqs. (12.121) and (12.122)); in the single–species (diagonal) limit this reduces to $m_f = \frac{y_f v}{\sqrt{2}}$ (eq. (12.123)).

Here v is fixed dynamically by the Tantrakyma potential and the g_C–Riesz construction (Definitions 9.335 and 9.343 and eq. (9.185)), with $Q = I_3 + \frac{Y}{2}$ ensuring electroweak consistency (Section 12.4.3).

Fermion (mass basis)	**Yukawa entry** y_n	**Mass** $m_n = \dfrac{y_n\, v}{\sqrt{2}}$	**Notes**
$e,\ \mu,\ \tau$	$y_e,\ y_\mu,\ y_\tau$	$m_e,\ m_\mu,\ m_\tau$	Charged lepton Dirac masses; Y_e diagonalized by biunitary rotation (eq. (12.121)).
$u,\ c,\ t$	$y_u,\ y_c,\ y_t$	$m_u,\ m_c,\ m_t$	Up–type Dirac masses; CKM mixing from left–handed rotations of Y_u, Y_d (diagnostic realized reference configuration–admissible unitary maps).
$d,\ s,\ b$	$y_d,\ y_s,\ y_b$	$m_d,\ m_s,\ m_b$	Down–type Dirac masses; see note above on CKM.
ν_i	y_{ν_i}	m_{ν_i}	Dirac case: $m_{\nu_i} = y_{\nu_i} v/\sqrt{2}$; seesaw option if a gauge–singlet Majorana sector is included (model choice external to Definition 12.105).

Table 12.2: Yukawa couplings and fermion masses after EWSB. Entries y_n denote diagonal Yukawa eigenvalues in the mass basis (singular values of the 3×3 matrices Y_f). The scale v is the Tantrakyma VEV fixed by the potential/diagnostic realized reference configuration construction (Definitions 9.335 and 9.343); the $h\bar{\psi}\psi$ coupling equals m_n/v (eq. (12.122)).

Remarks. (i) The bilinear $\overline{\psi}\psi$ sits in the τ–scalar channel $S^{(b)}$ and remains TT–null, so witnesses/TLO are read without graviton contamination (Definition 10.13, Proposition 10.22, and Corollary 10.125). (ii) Nielsen flatness and KL positivity certify that the extracted pole masses and residues are gauge–parameter independent on $\mathfrak{P}$ (Proposition 12.88). (iii) Electroweak charge assignments obey $Q = I_3 + \frac{Y}{2}$ for all realized species, consistent with the neutral/charged vector mixing produced by the BEH matrix (Section 12.4.3 and Proposition 9.347). (iv) Hierarchies reflect the structure of the Yukawa tensors Y_f in the operator–bundle setting (Definition 12.105); their origin may trace to deeper features of the correlator algebra and symmetry breaking within the Tantrakyma sector.

12.5 The Threefold Multiplicity of Chiral Matter

The observed replication of chiral matter into three families is traced to the *chronogeometric triad* and its monodromy, realized as a flat, unitary holonomy on a rank-3 auxiliary bundle that accompanies Tantrakyma spinor sections. The construction is (i) *geometric*: it uses the

triadic axes and the operational/relational transport (Definitions 5.1 and 6.7), and (ii) *operational*: its phases are accessible to holosymmetric interferometry and typed witnesses/TLO (Definition 8.4, Proposition 10.22, and Section 2.4). Crucially, no new dynamical vectors are introduced: the family structure is a *Berry–type* background holonomy (a *flat* connection on an auxiliary bundle), not an internal gauge sector (cf. the genuine gauge connection and its Yang–Mills dynamics in Definition 9.345 and Lemma 12.87).

12.5.1 From Triadic Time to a $\mathbb{C}^3$ Family Fibre

Recall the triad of relational axes for distinguishability and time,

$$\vec{t} = (t_{\mathrm{corr}}, t_{\mathrm{vac}}, t_{\mathrm{spec}}), \qquad \vec{d} = (d_{\mathrm{corr}}, d_{\mathrm{vac}}, d_{\mathrm{spec}}),$$

and the operator geometry induced from the Fourfold correlator and composite metric construction (Definitions 4.463, 5.1, 5.26 and 6.7). Loops in $\mathcal{X}_q$ that encircle loci where the *ordered* triad cyclically permutes ("triadic branch loci") carry a well-defined, connection-based holonomy derived from the graded, parallel-transport structure that also underwrites the Unified Inter-Modal Correlator and the Relational Levi–Civita transport (Definitions 4.9 and 6.7). We now package this monodromy into an auxiliary, flat family bundle that tensored to Tantrakyma spinor sections yields the threefold multiplicity.

Definition 12.106 (Family fibre and flat holonomy)**.** Let $\mathfrak{D} \subset \mathcal{X}_q$ be the *triadic branch locus*, i.e. those points around which analytic continuation permutes the ordered axes $(\mathrm{corr}, \mathrm{vac}, \mathrm{spec})$ cyclically. Define the *family bundle* $\pi_F : \mathcal{F} \to \mathcal{X}_q \setminus \mathfrak{D}$ to be a rank-3 Hermitian vector bundle equipped with a *flat* unitary connection $\mathcal{A}^{(F)}$ (curvature $F^{(F)} = 0$ on $\mathcal{X}_q \setminus \mathfrak{D}$) whose monodromy representation

$$\rho_F : \ \pi_1(\mathcal{X}_q \setminus \mathfrak{D}) \ \longrightarrow \ U(3)$$

maps a positive generator γ encircling a branch component to

$$\mathrm{Hol}_\gamma(\mathcal{A}^{(F)}) \ = \ U_\gamma^{(F)} \ = \ \mathrm{diag}(1, \omega, \omega^2), \qquad \omega = e^{2\pi i/3}. \tag{12.124}$$

The connection $\mathcal{A}^{(F)}$ is defined up to global conjugation by $U(3)$ and is taken to be *background* (no Yang–Mills kinetic term, hence no new gauge quanta; contrast with Definition 9.345).

Remark 12.107 (Why Z_3 and why rank three?)**.** The cyclic permutation of the triadic axes induces a C_3 action on the local chronogeometric data and the parallel-transport phases derived from the Fourfold-induced geometry (Definitions 4.9 and 5.1). The minimal faithful unitary lift of C_3 with nontrivial $2\pi/3$ phase is three-dimensional, with spectrum $\{1, \omega, \omega^2\}$. No lower-rank unitary representation simultaneously realises nontrivial $2\pi/3$ holonomy and preserves flatness/unitarity on all loops; thus $\mathrm{rank}(\mathcal{F}) = 3$ is minimal.

Coupling to fermionic kymons and compatibility with existing structures. Let $P \to \mathcal{X}_q$ be the G_{SM} principal bundle (Definition 12.83) and $E_n = P \times_{\rho_n} V_n$ the SM-associated bundles for the species in Definition 12.82. Fermionic kymons are operator-valued spinor sections realised by the species-extraction pipeline (Definitions 12.81 and 12.85) in

$$\Gamma\Big(\mathrm{End}(\mathcal{H}_{\mathrm{em}}) \otimes \mathcal{S} \otimes E_n\Big),$$

with covariant transport given by the split connection (Lemma 12.87). We extend the carrier by the family bundle to

$$\hat{\psi}_{n,\,\mathrm{fam}} \;\in\; \Gamma\Big(\mathrm{End}(\mathcal{H}_{\mathrm{em}}) \otimes \mathcal{S} \otimes E_n \otimes \mathcal{F}\Big), \tag{12.125}$$

and lift the split derivative to include the flat family piece,

$$\mathcal{D}_\mu^{(\mathrm{tot})}\hat{\psi}_{n,\,\mathrm{fam}} = \underbrace{\nabla_\mu^{(\mathcal{S})}}_{\text{spin}}\hat{\psi}_{n,\,\mathrm{fam}} \;+\; i[\mathcal{A}_\mu, \hat{\psi}_{n,\,\mathrm{fam}}] \;+\; i\,g\,A_\mu{}^I \rho_n(T_I)\hat{\psi}_{n,\,\mathrm{fam}} \;+\; \nabla_\mu^{(F)}\hat{\psi}_{n,\,\mathrm{fam}}, \tag{12.126}$$

where $\nabla^{(F)}$ is the flat covariant derivative associated to $\mathcal{A}^{(F)}$ (so $[\nabla_\mu^{(F)}, \nabla_\nu^{(F)}] = 0$ away from $\mathfrak{D}$). Since $F^{(F)} = 0$ on $\mathcal{X}_q \setminus \mathfrak{D}$, the curvature split of Lemma 12.87 is *unchanged*; in particular no new vector mass terms appear in Proposition 9.347 and no new TT channels are introduced (typed purity as in Definition 10.13 and Corollary 10.125).

Lemma 12.108 (Decoupling: no new gauge bosons and TT purity)**.** *The family connection $\mathcal{A}^{(F)}$ contributes no Yang–Mills action and no BEH mass matrix: it is flat and background, acting only by parallel transport phases on $\mathcal{F}$ (no $\mathcal{H}_{\mathrm{phys}}$ vector poles). Typed/triadic projectors commute with the family holonomy, hence TT purity and τ–channel placements are preserved (Definitions 9.350 and 10.13 and Corollary 10.125).*

Threefold multiplicity as holonomy sectors and its observables. Choose a local trivialization that diagonalises $U_\gamma^{(F)}$ and denote by $\{e_a\}_{a=0,1,2}$ the eigenframe of $\mathcal{F}$ with eigenvalues $\{1, \omega, \omega^2\}$. Then a species $\hat{\psi}_n$ lifts to three chiral copies

$$\hat{\psi}_n^{(a)} := \hat{\psi}_n \otimes e_a, \qquad a \in \{0, 1, 2\},$$

that acquire relative phases upon encircling γ, $(\hat{\psi}_n^{(a)}) \mapsto \omega^a\,(\hat{\psi}_n^{(a)})$. Holosymmetric interferometry (Section 2.4) and τ–typed witnesses (Definition 8.4 and Proposition 10.22) can resolve these Berry phases by comparing paths that differ by linking number with $\mathfrak{D}$; the meters remain TT–null by Corollary 10.125.

Yukawa couplings and family mixing under flat holonomy. Yukawa tensors Y_f are sections in

$$\Gamma\Big(\mathrm{Hom}(E_{f,L} \otimes E_{\mathrm{H}}, E_{f,R}) \otimes \mathrm{End}(\mathcal{F})\Big)$$

(Definition 12.105 and Proposition 12.90). Flatness implies $\nabla_\mu^{(F)} Y_f = 0$ away from $\mathfrak{D}$, so in a local trivialization where $\mathcal{A}^{(F)}$ is pure gauge the Y_f are constant matrices on $\mathcal{F}$. Mass matrices $M_f = \frac{v}{\sqrt{2}} Y_f$ (eq. (12.121)) may be diagonalized by biunitary rotations in flavour×family space; misalignment between the u- and d-sector family frames yields the usual CKM structure (diagnostic realized reference configuration-admissible unitary rotations), and analogously for PMNS in the lepton sector. Because $\mathcal{A}^{(F)}$ is flat, these mixings are *global holonomy data*, not additional gauge dynamics, and all pole data remain Nielsen-flat and KL-positive on $\mathfrak{P}$ (Proposition 12.88).

Corollary 12.109 (Minimality and robustness)**.** *The monodromy* (12.124) *is the minimal flat, unitary lift of the triad's C_3 action compatible with (i) the diagnostic realized reference configuration discipline, (ii) typed TT purity, and (iii) absence of new gauge quanta. It produces exactly three chiral copies of each SM representation in Definition 12.82, with all dynamical content (masses, mixings) governed by the existing Tantrakyma potential/BEH and Yukawa sectors (Definitions 9.343 and 12.105 and Proposition 9.347).*

Worked example: single loop around a branch component (family–phase readout). All statements are taken on a certified plateau $\mathfrak{P}$ with BRST projection $\mathsf{P}_{\mathrm{phys}}$ and typed readout in the τ–scalar channel $S^{(b)}$ (Definitions 7.29, 7.39 and 10.13, Proposition 10.22, and Corollary 10.125). Let $\mathfrak{D} \subset \mathcal{X}_q$ be the triadic branch locus and $\mathcal{F} \to \mathcal{X}_q \setminus \mathfrak{D}$ the flat family bundle with monodromy $U_\gamma^{(F)} = \mathrm{diag}(1, \omega, \omega^2)$ (Definition 12.106). Consider a two–arm holosymmetric interferometer (Section 2.4) whose arms γ_0, γ_1 share source/detector points and differ topologically by

$$\ell := \mathrm{link}(\gamma_1, \mathfrak{D}) - \mathrm{link}(\gamma_0, \mathfrak{D}) = +1,$$

i.e. arm γ_1 winds once positively around a connected component of $\mathfrak{D}$ while γ_0 is contractible.

State preparation. Fix a fermionic kymon species $\hat{\psi}_n$ realized as in Definitions 12.81 and 12.85. Let $\{e_a\}_{a=0,1,2}$ be a local eigenframe of $\mathcal{F}$ that diagonalizes $U_\gamma^{(F)}$, and prepare the input family state $|\phi_F\rangle = \sum_{a=0}^{2} c_a\, |e_a\rangle$ with $\sum_a |c_a|^2 = 1$. The full carrier is

$$|\Psi_{\mathrm{in}}\rangle \;\in\; \mathrm{Ran}\,\mathsf{P}_{\mathrm{phys}} \subset \Gamma\Big(\mathrm{End}(\mathcal{H}_{\mathrm{em}}) \otimes \mathcal{S} \otimes E_n \otimes \mathcal{F}\Big) \quad \text{(Definitions 12.80 and 12.83)}.$$

Transport and holonomy. Transport along γ_j is governed by the split connection with the *flat* family piece (Lemma 12.87, eq. (12.126)). Since $F^{(F)} = 0$ on $\mathcal{X}_q \setminus \mathfrak{D}$, only the *monodromy* matters:

$$U^{(\mathrm{tot})}(\gamma_0) = \mathbf{1}_{\mathcal{F}}\,, \qquad U^{(\mathrm{tot})}(\gamma_1) = U_\gamma^{(F)} = \mathrm{diag}(1, \omega, \omega^2)\,.$$

Thus the two outgoing arm states at the recombiner differ by the family phase:

$$|\Psi_0\rangle = |\Psi_{\mathrm{in}}\rangle\,, \qquad |\Psi_1\rangle = (\mathbf{1} \otimes U_\gamma^{(F)})\,|\Psi_{\mathrm{in}}\rangle\,.$$

Typed scalar witness and visibility. Let $\widehat{W}_{S^{(b)}}$ be a calibrated scalar witness (TLO) in the τ–sector (Definition 8.4 and Proposition 10.22). After the 50/50 recombination, the detected scalar meter is proportional to the interference term

$$\Lambda^{[S^{(b)}]} \;\propto\; \mathrm{Re}\Big\langle \Psi_0 \Big| \widehat{W}_{S^{(b)}} \Big| \Psi_1 \Big\rangle \;=\; \mathrm{Re}\Big(\sum_{a=0}^{2} |c_a|^2\, \omega^{\,a} \Big) \Big\langle \Psi_{\mathrm{in}} \Big| \widehat{W}_{S^{(b)}} \Big| \Psi_{\mathrm{in}} \Big\rangle,$$

where we used that $\widehat{W}_{S^{(b)}}$ acts diagonally on $\mathcal{F}$ in the eigenframe (it is family–blind but $\mathcal{F}$–compatible). Therefore the *scalar-channel visibility* is

$$\mathcal{V}_{S^{(b)}} \;=\; \Big| \sum_{a=0}^{2} |c_a|^2\, \omega^{\,a} \Big|\,. \tag{12.127}$$

Unprojected equal–mixing case. If the source is family–unpolarized, $|c_0|^2 = |c_1|^2 = |c_2|^2 = \frac{1}{3}$, then

$$\sum_{a=0}^{2} |c_a|^2\, \omega^{\,a} = \tfrac{1}{3}(1+\omega+\omega^2) = 0 \quad \Rightarrow \quad \mathcal{V}_{S^{(b)}} = 0,$$

so the scalar witness shows *no* net fringe despite phase accumulation—complete cancellation across the three families.

Family-resolved readout (projectors). Introduce the local family projectors $P_F^{(a)} = |e_a\rangle\langle e_a|$ and the corresponding scalar witness $\widehat{W}_{S^{(b)},a} := (\mathbf{1} \otimes P_F^{(a)})\, \widehat{W}_{S^{(b)}}\, (\mathbf{1} \otimes P_F^{(a)})$. Then

$$\Lambda_a^{[S^{(b)}]} \;\propto\; \mathrm{Re}\big(\omega^{\,a}\big)\, \Big\langle \Psi_{\mathrm{in}} \Big| \widehat{W}_{S^{(b)},a} \Big| \Psi_{\mathrm{in}} \Big\rangle, \qquad \mathcal{V}_{S^{(b)}}^{(a)} = |\omega^{\,a}| = 1.$$

Hence each family channel exhibits full visibility with a definite *geometric phase* $0, \pm 2\pi/3$; the summed, unprojected signal cancels unless the source or the detector is family–selective. Typed/TT hygiene is preserved because $P_F^{(a)}$ commutes with the τ–typed and TT projectors (Definition 10.13 and Corollary 10.125).

Multiple windings and phase periodicity. If $\ell \in \mathbb{Z}$ windings differ between the arms, the scalar visibility generalizes to

$$\mathcal{V}_{S^{(b)}}(\ell) = \Big| \sum_{a=0}^{2} |c_a|^2\, \omega^{\,a\ell} \Big|, \qquad \mathcal{V}_{S^{(b)}}^{(a)}(\ell) = |\omega^{\,a\ell}| = 1,$$

with $2\pi/3$ periodicity in ℓ (modulo 3). This realizes the C_3 monodromy of Remark 12.107 as an *operational* phase ladder.

Summary. A single positive loop of one arm around a triadic branch component multiplies the three family components by $(1, \omega, \omega^2)$. Family-unresolved scalar meters erase the fringe $(1+\omega+\omega^2 = 0)$, whereas family–projected meters recover full visibility with characteristic $\pm 2\pi/3$ phase. The effect is purely geometric (flat holonomy), introduces no new gauge quanta (Lemma 12.108), and is read in τ–scalar witnesses with TT purity intact (Definition 10.13, Proposition 10.22, and Corollary 10.125).

Definition 12.110 (Family–lifted Tantrakyma fields)**.** Let $\hat{\psi}_n$ be any fermionic kymon field (quark or lepton) realized by the species–extraction pipeline (Definitions 12.81 and 12.85) in the SM irrep $E_n = P \times_{\rho_n} V_n$ (Definitions 12.82 and 12.83). Let $\mathcal{F} \to \mathcal{X}_q \setminus \mathfrak{D}$ be the rank-3 *family bundle* with flat unitary connection $\mathcal{A}^{(F)}$ and Z_3 monodromy (Definition 12.106). The *family lift* of $\hat{\psi}_n$ is the section

$$\hat{\psi}_n^{\mathsf{f}} \;\in\; \Gamma\Big(\mathrm{End}(\mathcal{H}_{\mathrm{em}}) \;\otimes\; \mathcal{S} \;\otimes\; E_n \;\otimes\; \mathcal{F}\Big),$$

where $\mathcal{S} \to \mathcal{X}_q$ is the Tantrakyma spinor bundle (Definition 12.80). Its total split covariant derivative is

$$\boxed{\mathcal{D}_\mu^{(\mathrm{tot})} \hat{\psi}_n^{\mathsf{f}} = \underbrace{\nabla_\mu^{(\mathcal{S})}}_{\text{spin lift}}\, \hat{\psi}_n^{\mathsf{f}} \;+\; i\,[\mathcal{A}_\mu, \hat{\psi}_n^{\mathsf{f}}] \;+\; i\,g\,A_\mu^{\;I}\, \rho_n(T_I)\, \hat{\psi}_n^{\mathsf{f}} \;+\; \nabla_\mu^{(F)} \hat{\psi}_n^{\mathsf{f}},} \tag{12.128}$$

where: (i) $\nabla^{(\mathcal{S})}$ is the spin connection induced from $(\mathcal{X}_q, \bar{\Sigma})$ (Definitions 6.7 and 11.97); (ii) $\mathcal{A}_\mu$ is the operator–bundle connection acting by adjoint on $\mathrm{End}(\mathcal{H}_{\mathrm{em}})$ (Definition 6.192); (iii) $A_\mu^{\;I} T_I$ is the SM internal gauge connection on E_n (Section 12.4.4 and Definition 9.345); and (iv) $\nabla_\mu^{(F)} = \mathbb{1} \otimes \mathcal{A}_\mu^{(F)}$ is the *flat* family covariant derivative (Definition 12.106).

Curvature split and flatness of the family piece. Away from $\mathfrak{D}$ the family curvature vanishes, $F^{(F)}_{\mu\nu} = 0$, so the commutator reduces to the established split (Lemma 12.87):

$$[\mathcal{D}^{(\mathrm{tot})}_\mu, \mathcal{D}^{(\mathrm{tot})}_\nu]\hat{\psi}^{\mathsf{f}}_n = \tfrac{1}{4}\,\bar{R}_{\mu\nu}{}^{ab}\gamma_{ab}\hat{\psi}^{\mathsf{f}}_n \;+\; i\,[\mathcal{F}_{\mu\nu}, \hat{\psi}^{\mathsf{f}}_n] \;+\; i\,g\,F^{\,I}_{\mu\nu}\rho_n(T_I)\hat{\psi}^{\mathsf{f}}_n,$$

with no additional term from $\mathcal{A}^{(F)}$ (*flat* background). Hence the BEH vector mass matrix and gauge–sector dynamics remain those of Proposition 9.347, and no new gauge quanta appear (cf. Lemma 12.108).

Gauge and family transformation laws. Under a local SM transform $g : \mathcal{X}_q \to G_{\mathrm{SM}}$ and a family-frame change $U_F : \mathcal{X}_q \setminus \mathfrak{D} \to U(3)$,

$$\hat{\psi}^{\mathsf{f}}_n(x) \;\longmapsto\; \Big(\mathrm{Ad}_{U_g(x)} \otimes 1_{\mathcal{S}} \otimes \rho_n(g(x)) \otimes U_F(x)\Big)\,\hat{\psi}^{\mathsf{f}}_n(x) \qquad (\textit{Lemma } 12.86,\ \textit{Definition } 12.106),$$

and (12.128) is covariant.

Typed/TT hygiene and witnesses. Typed vector/scalar placements and TT purity are unchanged: the family lift commutes with the τ–typed/TT projectors (Definitions 9.350 and 10.13 and Corollary 10.125), and all meters/witnesses remain valid (Definition 8.4 and Proposition 10.22). Nielsen flatness and KL positivity on the diagnostic realized reference configuration continue to hold for realized poles/residues (Proposition 12.88).

Monodromy and threefold multiplicity. Parallel transport around a loop encircling a branch component multiplies the family components by $(1, \omega, \omega^2)$ (Definition 12.106 and Remark 12.107), yielding three holonomy sectors of each SM species without introducing new gauge bosons; interferometric readouts are given in the worked example (Section 12.5.1).

12.5.2 Why Exactly Three Families: a Minimal–faithful Holonomy Theorem

Theorem 12.111 (Triadic monodromy ⇒ threefold family multiplicity)**.** *Assume:*

(A1) *Triadic continuation around the branch locus $\mathfrak{D}$ acts nontrivially by a C_3 phase on holosymmetric interferometric amplitudes extracted from the Fourfold–induced chronogeometry (Definitions 4.9, 5.1 and 6.7); operationally, a loop linking $\mathfrak{D}$ produces a $2\pi/3$ Berry-type shift in the interferometer model (Section 2.4, cf. Section 12.5.1).*

(A2) *The family structure is implemented by a* flat*, unitary holonomy of an auxiliary Hermitian bundle $\mathcal{F} \to \mathcal{X}_q \setminus \mathfrak{D}$, which couples only through covariant transport in the split derivative (no Yang–Mills dynamics), as in Definitions 12.106 and 12.110 and eq. (12.128); typed/TT hygiene is preserved (Definition 10.13 and Corollary 10.125).*

(A3) *Family–*unresolved *scalar witnesses (trace over family indices) show no net, path-dependent bias on the diagnostic realized reference configuration; equivalently, the unprojected scalar-channel visibility cancels on a single positive winding (as in Section 12.5.1, eq. (12.127)) so that family-blind meters remain insensitive to the monodromy.*

Then the minimal, globally well-defined realisation of this holonomy requires $\operatorname{rank}(\mathcal{F}) = 3$, *with monodromy spectrum* $\{1, \omega, \omega^2\}$ *and* $\omega = e^{2\pi i/3}$, *i.e.*

$$U_\gamma^{(F)} \;\sim\; \operatorname{diag}(1, \omega, \omega^2), \tag{12.129}$$

as in eq. (12.124). *Hence each chiral Tantrakyma multiplet appears in exactly three families.*

Proof. Let $\rho_F : \pi_1(\mathcal{X}_q \setminus \mathfrak{D}) \to U(n)$ be the flat, unitary monodromy representation of the family connection (Definition 12.106). For a positive generator γ linking a single branch component, flatness and unitarity imply $\rho_F(\gamma)$ is diagonalizable with eigenvalues on the unit circle. Assumption (A1) (triadic C_3 action) enforces

$$\rho_F(\gamma)^3 = \mathbb{1}, \qquad \rho_F(\gamma) \neq \mathbb{1},$$

hence the eigenvalues belong to $\{1, \omega, \omega^2\}$ with $\omega = e^{2\pi i/3}$, and at least one of ω, ω^2 occurs.

Write the multiplicities as $m_0, m_1, m_2 \in \mathbb{N}$ with $m_0 + m_1 + m_2 = n$, so that

$$\rho_F(\gamma) \;\sim\; \operatorname{diag}(\underbrace{1, \dots, 1}_{m_0}, \underbrace{\omega, \dots, \omega}_{m_1}, \underbrace{\omega^2, \dots, \omega^2}_{m_2}).$$

Now impose the two operational/global constraints encoded in (A2)–(A3):

(i) Family–blind cancellation. Family–unresolved scalar witnesses are traces over $\mathcal{F}$ (the meter is family-blind but $\mathcal{F}$–compatible), hence the interferometric factor carried by the family holonomy is

$$\operatorname{Tr}_{\mathcal{F}} \rho_F(\gamma) \;=\; m_0 + m_1\,\omega + m_2\,\omega^2 \quad \text{(Section 12.5.1, eq. (12.127))}.$$

Assumption (A3) requires that this family-blind factor *vanish* on a single positive winding to remove any path-dependent bias in unprojected scalar meters, i.e.

$$m_0 + m_1\,\omega + m_2\,\omega^2 \;=\; 0. \tag{12.130}$$

Since $1 + \omega + \omega^2 = 0$, Eq. (12.130) is solved if and only if

$$(m_0, m_1, m_2) \;\propto\; (1, 1, 1). \tag{12.131}$$

This already rules out $n = 1$ and $n = 2$: no nontrivial choice of (m_0, m_1, m_2) with $m_0 + m_1 + m_2 \in \{1, 2\}$ can satisfy (12.130) while keeping $\rho_F(\gamma) \neq \mathbb{1}$.

(ii) Measure/diagnostic realized reference configuration neutrality. The family connection is background and flat (Definitions 12.106 and 12.110); it must not alter the diagnostic realized reference configuration-certified pole/residue data or induce global phases in family-blind densities. A sufficient, representation-theoretic way to encode this is to demand

$$\det \rho_F(\gamma) = 1, \tag{12.132}$$

so that family transport lies in $SU(n)$ and introduces no global $U(1)$ phase in scalar densities (consistent with the decoupling Lemma 12.108 and diagnostic realized reference configuration hygiene Proposition 12.88). In multiplicity form, (12.132) reads

$$\omega^{\,m_1 + 2m_2} = 1 \quad \Longleftrightarrow \quad m_1 + 2m_2 \equiv 0 \pmod 3.$$

Combined with (12.131), the *minimal* positive solution is $(m_0, m_1, m_2) = (1, 1, 1)$, giving $n = \mathrm{rank}(\mathcal{F}) = 3$.

Thus the minimal, faithful, unitary monodromy satisfying (A1)–(A3) is

$$\rho_F(\gamma) \;\sim\; \mathrm{diag}(1, \omega, \omega^2),$$

unique up to global conjugation, which is exactly (12.124) (cf. Definition 12.106). Appending $\mathcal{F}$ to the Tantrakyma spinor carriers (Definition 12.110) therefore produces three, and only three, covariantly distinct family components for each chiral multiplet, with no new gauge vectors or TT contamination (Lemma 12.108, Definition 10.13, and Corollary 10.125). □

Corollary 12.112 (Stability under Yukawa/BEH and typed readout)**.** *Since $\nabla^{(F)}$ is flat, Yukawa tensors $Y_f \in \Gamma(\mathrm{End}(\mathcal{F}))$ can be taken locally constant on $\mathcal{X}_q \setminus \mathfrak{D}$ (Definition 12.105), and the BEH analysis is unaffected (Proposition 9.347). Masses and mixings arise as diagnostic realized reference configuration-admissible unitary rotations in flavour×family space, while all witnesses/TLO remain in the τ–scalar channel with TT purity (Proposition 10.22 and Corollary 10.125).*

Corollary 12.113 (No extra light families by Holosymmetric Economy)**.** *Let the family fibre $(\mathcal{F}, \mathcal{A}^{(F)})$ realize the C_3 triadic monodromy by a flat, unitary holonomy with spectrum $\{1, \omega, \omega^2\}$ (rank 3 minimal and faithful). Any duplication*

$$\mathcal{F} \;\longrightarrow\; \mathcal{F} \oplus \underline{\mathbb{C}}^k$$

by a trivial (flat, identity–holonomy) subbundle $\underline{\mathbb{C}}^k$ produces sectors that are holosymmetrically visible yet dynamically redundant. By the Stage II criterion of the Holosymmetric Selection Principle (relational orthogonality), such sectors fail maximal orthogonality against the realized vacuum and are therefore suppressed. Hence the minimal faithful rank–3 choice is selected on the diagnostic realized reference configuration.

Proof. **Step 1 (What the duplication does).** The duplication replaces the family holonomy representation on loops $\gamma \in \pi_1(\mathcal{X}_q \setminus \mathfrak{D})$ by the block sum

$$U_\gamma^{(F)} \;\longmapsto\; U_\gamma^{(F)} \;\oplus\; \mathbb{I}_k.$$

Equivalently, on the compact test group for the triadic holonomy (denote it $K \simeq C_3$ inside the interferometer), the character on the family fibre changes as

$$\chi_{\mathrm{ext}}(g) \;=\; \chi_{\mathrm{min}}(g) \;+\; k \cdot \chi_{\mathrm{triv}}(g), \qquad \chi_{\mathrm{triv}}(g) \equiv 1 \;\text{ on } K.$$

Here χ_{min} already contains one trivial eigenvalue (the 1 in $\{1, \omega, \omega^2\}$), while the added block contributes k further trivial characters.

Step 2 (Visibility and relational orthogonality). Stage II relational orthogonality on the diagnostic realized reference configuration is certified by the vanishing of the holosymmetric overlap via the character/Peter–Weyl filter: if the K–characters are orthogonal (Haar inner product), then the interferometric overlap can be driven to zero by admissible probes/meters, and Stage II separation holds (Corollary 12.48, using Proposition 12.10 and the construction of the ensemble overlap functional).

Compute the Peter–Weyl inner product on K between the *realized* minimal family and the *duplicated* one:

$$\langle \chi_{\text{ext}}, \chi_{\text{min}} \rangle_K = \langle \chi_{\text{min}}, \chi_{\text{min}} \rangle_K + k \underbrace{\langle \chi_{\text{triv}}, \chi_{\text{min}} \rangle_K}_{>0}.$$

Because χ_{min} contains a trivial component (the eigenvalue 1 of the C_3 spectrum), the second term is strictly positive; hence the characters are *not* orthogonal on K and the Stage II holosymmetric filter cannot render the two sectors relationally orthogonal. Therefore the enlarged model cannot be selected over the realized minimal one by Stage II tomography (compare Theorem 12.79, Stage II uniqueness).

Step 3 (No dynamical gain; "economy"). Dynamically, the added $\underline{\mathbb{C}}^k$ carries *trivial* family holonomy and couples only through covariant transport as an identity block in the total derivative,

$$\mathfrak{D}_\mu^{\text{tot}} = \nabla_\mu^{(\text{rel})} \otimes \mathbb{I} \; + \; i\,\text{ad}_{\mathcal{A}_\mu} \otimes \mathbb{I} \; + \; i\,\mathbb{I} \otimes \mathcal{A}_\mu^{(F)},$$

so the duplication implements a direct-sum replica with the same SM gauge/internal dynamics (the $\mathcal{A}_\mu$ part) and the same chronogeometric transport (the $\nabla^{(\text{rel})}$ part) without introducing new poles or residues (diagnostic realized reference configuration discipline as in Definition 9.345 and the split-connection formalism). Hence the sector is *holosymmetrically visible* (its character contributes to interferometric meters) but *dynamically redundant* (no new typed poles beyond a multiplicity factor).

Step 4 (Selection by maximal coherence & orthogonality). The Stage II rule, stated on the BRST–physical sector, selects the unique candidate that both maximizes the holosymmetric coherence functional and is relationally orthogonal to all other viable competitors (Theorem 12.79, Step II). Since the duplicated model is *not* orthogonal to the minimal one (Step 2) and does not increase physically meaningful coherence (no new admissible poles/residues on $H^0(\hat{Q}_{\text{BRST}})$), the "Holosymmetric Economy" (minimality among faithful realizations) eliminates it in favor of the minimal rank–3 family fibre.

Therefore, no extra light families arise: the triadic monodromy fixes a minimal faithful rank 3, and trivial duplications are excluded by Stage II holosymmetric selection. □

Start with the built-in "three-ness" of chronogeometry: relational change is resolved along three complementary axes—correlation, vacuum response, and spectral shear—introduced with the triadic time/distance structure and its operator geometry (Section 4.12 and Definitions 4.463, 5.1 and 5.26). When physical data are parallel-transported around certain loops in the emergent space $\mathcal{X}_q$, those three axes can cycle among themselves under the Relational Levi–Civita transport induced by the Fourfold correlator (Definitions 4.9 and 6.7). Operationally—exactly in the sense of a Berry phase—the effect of a full loop is a *three-step* phase pattern on amplitudes: one branch returns unchanged, while the other two acquire phases $\pm 2\pi/3$. That cyclic three-beat rhythm is captured as a nontrivial C_3 *monodromy* seen by holosymmetric interferometry (Sections 2.4 and 12.5.1).

This purely geometric effect is packaged into a harmless "address tag" carried by each fermion: a flat, unitary *family bundle* $(\mathcal{F}, \mathcal{A}^{(F)})$ with spectrum $\{1, \omega, \omega^2\}$ ($\omega = e^{2\pi i/3}$) and Z_3 monodromy (Definition 12.106). The bundle introduces no new forces and no new gauge bosons—it is background and flat—and couples only through covariant transport in the

family-lifted carrier (Definition 12.110, eq. (12.128), and Lemma 12.108). Thus each quark or lepton appears with three internal "addresses"; carrying the field around a loop that links a branch component rotates these addresses by the three phases. This is the sense in which there are *three families*: the same chiral multiplet realized in three holonomy sectors.

Why *exactly* three? Two ingredients make the count rigid. First, interferometric readouts register a nontrivial C_3 loop phase, so the family holonomy must exhibit cube-roots-of-unity eigenvalues (Section 12.5.1). Second, family-*blind* scalar meters should show no path-dependent bias on a single winding—no fringe from a purely geometric loop unless the measurement resolves family—which is precisely the cancellation $\frac{1}{3}(1+\omega+\omega^2)=0$ in the scalar witness (Section 12.5.1). With only one or two eigenvectors, it is impossible to realize both a nontrivial $2\pi/3$ phase *and* the perfect family-blind cancellation. The *Minimal-Faithful Holonomy Theorem* then forces the minimal unitary realization to have rank 3 and phases $\{1,\omega,\omega^2\}$, locking the family count to three (Theorem 12.111).

Could there be more families? One could append extra *trivial* copies $\mathcal{F}\oplus\underline{\mathbb{C}}^k$ that carry no phase. These additions produce holosymmetrically visible yet dynamically redundant sectors (no new poles or residues, no new gauge vectors), failing the Stage-II relational-orthogonality filter and being removed by the Holosymmetric Economy selection (Corollary 12.113). The net picture is clean: *three and only three* light families, no extra gauge bosons, TT purity of chronogeometry preserved (Definition 10.13 and Corollary 10.125), and all mass/mixing physics continuing to arise from the Tantrakyma potential/BEH and Yukawa sectors (Propositions 9.347 and 12.90).

12.5.3 Family Connection, Yukawa Structure, and Flavor Mixing

In the family–lifted formulation, the Yukawa sector acquires a geometric meaning: it is an $\mathrm{End}(\mathcal{F})$–valued fibre map on the flat, unitary family bundle $(\mathcal{F},\mathcal{A}^{(F)})$ defined by the triadic monodromy (Definition 12.106 and Theorem 12.111). The total transport of fermionic kymons includes the family piece via the split derivative (Definition 12.110 and eq. (12.128)), while all gauge/chronogeometric parts remain those of the established framework (Definitions 6.7 and 6.192 and Lemma 12.87).

Definition 12.114 (Family–Yukawa endomorphisms and mass operators)**.** Let $E_{f,L},E_{f,R}$ be the G_{SM}–associated bundles for a fermion type $f\in\{u,d,e,\nu\}$ and let E_{H} carry the Tantrakyma doublet $H\sim(1,2)_{+1/2}$ that effects EWSB (Definition 12.82, Proposition 12.90, and Section 12.4.5). A *family–Yukawa tensor* is a section

$$Y_f \;\in\; \Gamma\Big(\mathrm{Hom}(E_{f,L}\otimes E_{\mathrm{H}},\,E_{f,R})\;\otimes\;\mathrm{End}(\mathcal{F})\Big),$$

entering the Lagrangian as

$$\mathcal{L}_Y \;\supset\; -\,\overline{\Psi^{\mathsf{f}}_L}\,\hat{H}\,Y_f\,\Psi^{\mathsf{f}}_R \;+\; \text{h.c.},$$

with the usual $SU(2)_L$ contraction. Gauge invariance is exactly the selection rule of Proposition 12.90. After symmetry breaking with $\langle H\rangle=\bar{\Theta}/\sqrt{2}$ (radial mode; Definition 9.343 and eq. (9.185)), the *family mass operator* is

$$M_f(x) \;:=\; \frac{\bar{\Theta}(x)}{\sqrt{2}}\,Y_f(x)\;\in\;\mathrm{End}(\mathcal{F}_x), \tag{12.133}$$

consistent with the diagnostic realized reference configuration relation $m_f = \frac{y_f \bar{\Theta}}{\sqrt{2}}$ on eigenvalues (eq. (12.121)).

Proposition 12.115 (Parallel transport and CKM/PMNS as relative family holonomies)**.** *Let $V_{fL}(x), V_{fR}(x) \in U(3)$ be local unitary frames that diagonalize $M_f(x)$, $V_{fL}^\dagger(x)\, M_f(x)\, V_{fR}(x) = \mathrm{diag}(m_{f1}, m_{f2}, m_{f3})$. Along a worldline $\gamma \subset \mathcal{X}_q \setminus \mathfrak{D}$ connecting preparation/measurement points, family frames undergo transport by the Wilson line of $\mathcal{A}^{(F)}$,*

$$W_F(\gamma) \;:=\; \mathcal{P}\exp\!\Big(i\int_\gamma \mathcal{A}^{(F)}\Big) \;\in U(3) \qquad (\textit{Definition 12.106}),$$

while the chronogeometric and internal–gauge parts are transported by the usual split connection (Lemma 12.87). The observable charged–current mixing matrices are then

$$V_{\mathrm{CKM}}(x) \;=\; \big[V_{uL}^\dagger\, W_F(\gamma)\, V_{dL}\big](x), \qquad V_{\mathrm{PMNS}}(x) \;=\; \big[V_{\nu L}^\dagger\, W_F(\gamma)\, V_{eL}\big](x), \tag{12.134}$$

i.e. flavor mixing is the relative family holonomy *between left frames of the two species.*

Proof. (i) *Local diagonalization and family gauge freedom.* By Definition 12.114, M_f is an endomorphism of $\mathcal{F}_x$, diagonalizable by $V_{fL/R}(x)$. Under a local family frame change $U_F(x) \in U(3)$, $M_f \mapsto U_F M_f U_F^\dagger$, hence $V_{fL} \mapsto U_F V_{fL}$ at the same point.

(ii) *Transport.* Between two points on γ, family frames are compared by $W_F(\gamma)$ (flat background transport; Definitions 12.106 and 12.110). The left–handed charged current couples $E_{u,L}$ to $E_{d,L}$; in family space this induces the relative rotation $V_{uL}^\dagger W_F(\gamma) V_{dL}$.

(iii) *Gauge covariance.* Under U_F at the endpoints, $W_F(\gamma) \mapsto U_F(x)\, W_F(\gamma)\, U_F(x_0)^\dagger$, while $V_{fL}(x_0) \mapsto U_F(x_0)\, V_{fL}(x_0)$ and $V_{fL}(x) \mapsto U_F(x)\, V_{fL}(x)$. The combination in (12.134) is therefore invariant. The same reasoning applies to the lepton sector. Chronogeometric and SM–gauge transports commute with the flat family transport (split derivative; Lemma 12.87), so (12.134) isolates the family contribution. □

Remark 12.116 (CP phases from family holonomy; curvature and loop class)**.** If $\mathcal{A}^{(F)}$ were globally pure gauge, all open Wilson lines could be trivialized in a single chart, and any complex phases in (12.134) would come solely from misalignment of the local diagonalizers V_{fL}. The *flat but nontrivial* monodromy fixed by the triadic continuation (Definition 12.106 and Theorem 12.111) already supplies discrete phases $\{1, \omega, \omega^2\}$ to closed Wilson loops (Berry-type origin). In addition, localized or distributional curvature at the branch locus $\mathfrak{D}$ can render Wilson loops path–class dependent; closed-loop operators in $\mathrm{End}(\mathcal{F})$ then generate Jarlskog-type, gauge–invariant phase diagnostics measurable by holosymmetric tomography (Sections 2.4 and 12.5.1). Diagnostic realized reference configuration hygiene (Nielsen flatness/KL positivity) remains intact for realized poles and residues (Proposition 12.88), and TT purity is preserved (Definition 10.13 and Corollary 10.125).

12.5.4 Anomaly and HSP Compatibility

Replicating every chiral multiplet threefold through the flat family fibre $\mathcal{F}$ preserves all local and global anomaly constraints encoded in the anomaly block eq. (12.46). Each entry in that block is linear in traces such as $\mathrm{Tr}\, Y$, $\mathrm{Tr}\, Y^3$, $\mathrm{Tr}\big(Y\, T^a T^a\big)$, etc.; since one–generation

assignments satisfy the vanishing conditions (Proposition 12.26), triplication multiplies zero by an overall factor of 3 and remains zero (Corollary 12.27). This matches the Stage I requirement of the Holosymmetric Selection Principle: viability demands $\mathcal{C}(A) = 1$ if and only if BRST is nilpotent/conserved and *all* anomaly meters vanish (Definitions 12.20 and 12.33 and Theorem 12.42).

Moreover, the family structure—implemented as a *flat*, unitary auxiliary connection $\mathcal{A}^{(F)}$ that couples only through the total covariant transport,

$$\begin{aligned}\mathfrak{D}_\mu^{\mathrm{tot}}\Psi^{\mathsf{f}} = &\left(\nabla_\mu^{(\mathrm{rel})} \otimes \mathbb{I}_3\right)\Psi^{\mathsf{f}} \\ &+\ i\left(\mathrm{ad}_{\mathcal{A}_\mu} \otimes \mathbb{I}_3\right)\Psi^{\mathsf{f}} \\ &+\ i\left(\mathbb{I} \otimes \mathcal{A}_\mu^{(F)}\right)\Psi^{\mathsf{f}}\end{aligned}$$

(Definition 12.110 and eq. (12.128)), does not alter Stage I: the unitarity operator $\mathbf{U}_{\mathrm{unitarity}}$ acts trivially on $\mathcal{F}$, and the anomaly block (which probes G_{SM} and global consistency) is unchanged. Hence Stage I is insensitive to the presence of a flat family bundle. At Stage II (relational orthogonality), "extra" trivial summands $\mathcal{F} \oplus \mathbb{C}^k$ fail the orthogonality/economy test and are dynamically suppressed, as formalized in Corollary 12.113 (see also the Stage II meters and orthogonality bridge in Section 3.1.5, Proposition 12.10, and Corollary 12.48).

Operational Footprints and Predictions.

- **Holosymmetric readout (family holonomy).** Interferometric loops γ that link the triadic branch set $\mathfrak{D}$ read the Z_3 family monodromy $U_\gamma^{(F)} = \mathrm{diag}(1, \omega, \omega^2)$ (Definition 12.106 and eq. (12.124)) as a $2\pi/3$ step in family–resolved visibilities. In charged–current processes, the observable mixing matrices are Wilson lines of $\mathcal{A}^{(F)}$ sandwiched between local diagonalizers,

  $$V_{\mathrm{CKM}}(x) = \left[V_{uL}^\dagger\, \mathcal{P} e^{\,i\int_\gamma \mathcal{A}^{(F)}}\, V_{dL}\right](x), \qquad V_{\mathrm{PMNS}}(x) = \left[V_{\nu L}^\dagger\, \mathcal{P} e^{\,i\int_\gamma \mathcal{A}^{(F)}}\, V_{eL}\right](x),$$

  so flavor mixing is *relative family holonomy* (Proposition 12.115 and eq. (12.134)).

- **Sum rules for phases (contractible loops).** For loops that are contractible in $\mathcal{X}_q \setminus \mathfrak{D}$, the family Wilson loop is the identity. In family–blind readouts this yields rephasing–invariant sum rules that combine CKM/PMNS phases measured at separated loci, directly tied to eq. (12.134) and the flatness of $\mathcal{A}^{(F)}$ away from $\mathfrak{D}$ (Definition 12.106).

- **Stability of threefold multiplicity.** The minimal–faithful holonomy realization of the observed triadic monodromy is $\mathrm{rank}(\mathcal{F}) = 3$ with spectrum $\{1, \omega, \omega^2\}$ (Theorem 12.111). Attempts to populate a fourth light family correspond to adjoining a trivial summand to $\mathcal{F}$; Stage II's orthogonality/economy criterion eliminates these add-ons (Corollary 12.113).

Synthesis. Family multiplicity is not an extra dynamical gauge sector but the *minimal faithful* Berry-type holonomy of the triadic chronogeometry packaged into a flat, unitary auxiliary bundle (Definition 12.106 and Theorem 12.111). It integrates seamlessly with

the Tantrakyma representation theory and EWSB sector (including $Q = I_3 + \frac{1}{2}Y$ and the BEH mass matrix; Section 12.4.3 and Proposition 9.347), while HSP Stage I/II continue to enforce anomaly-free viability and relational economy (Theorem 12.42, eq. (12.46), Proposition 12.26, and Corollaries 12.27 and 12.113). Flavor mixing and CP structure then arise from relative family holonomy and (if present) localized family curvature (Proposition 12.115 and Remark 12.116).

12.5.5 Flavor Mixing via the CKM and PMNS Matrices

Flavor mixing is a direct consequence of mass generation in the IQR framework: the matrices of Cabibbo–Kobayashi–Maskawa (CKM) and Pontecorvo–Maki–Nakagawa–Sakata (PMNS) emerge from the *misalignment* between the basis in which Yukawa couplings (and hence masses) are diagonal and the basis in which the weak charged currents are diagonal (Section 12.4.6, Definition 12.114, and Proposition 12.90; see also [554, 128, 563, 564]). In the family–lifted description, this misalignment is further refined as a relative *family holonomy* of the flat auxiliary bundle $(\mathcal{F}, \mathcal{A}^{(F)})$ (Definitions 12.106 and 12.110 and Proposition 12.115).

Mass eigenstates vs. gauge eigenstates

Fermion masses arise from the Yukawa coupling to the Tantrakyma doublet $H \sim (1,2)_{+1/2}$ and its VEV $\langle H \rangle = \bar{\Theta}/\sqrt{2}$ (Section 12.4.5, Definition 9.343, and eq. (9.185)). For each species, the Yukawa endomorphism Y_f on the family fibre defines a local mass operator $M_f = \bar{\Theta}\, Y_f/\sqrt{2}$ (Definition 12.114 and eq. (12.133)). In general, the family frames that diagonalize M_f need not coincide with those that diagonalize the charged weak current.

Let $\psi_{u,L}^{(\mathrm{gauge})}$ and $\psi_{d,L}^{(\mathrm{gauge})}$ denote the $SU(2)_L$ doublet components arranged as generation triplets (one certified plateau $\mathfrak{P}$; Definition 12.82):

$$\psi_{u,L}^{(\mathrm{gauge})} = \begin{pmatrix} u'_L \\ c'_L \\ t'_L \end{pmatrix}, \qquad \psi_{d,L}^{(\mathrm{gauge})} = \begin{pmatrix} d'_L \\ s'_L \\ b'_L \end{pmatrix}. \tag{12.135}$$

The physical (propagating) fields are the mass eigenstates

$$\psi_{u,L}^{(\mathrm{mass})} = \begin{pmatrix} u_L \\ c_L \\ t_L \end{pmatrix}, \qquad \psi_{d,L}^{(\mathrm{mass})} = \begin{pmatrix} d_L \\ s_L \\ b_L \end{pmatrix}, \tag{12.136}$$

obtained by unitary rotations in family space (below). The same structure applies to the lepton sector with (ν, e) in place of (u, d) (Definition 12.82).

Formal origin of the mixing matrices

On the family fibre, diagonalize the Yukawa (or equivalently the mass) operators by bi–unitary transformations (Definition 12.114):

$$Y_u^{\mathrm{diag}} = V_{uL}^\dagger\, Y_u\, V_{uR}, \qquad Y_d^{\mathrm{diag}} = V_{dL}^\dagger\, Y_d\, V_{dR}, \tag{12.137}$$

and define the basis change between gauge and mass eigenstates for left fields,

$$\begin{pmatrix} u'_L \\ c'_L \\ t'_L \end{pmatrix} = V_{uL} \begin{pmatrix} u_L \\ c_L \\ t_L \end{pmatrix}, \qquad \begin{pmatrix} d'_L \\ s'_L \\ b'_L \end{pmatrix} = V_{dL} \begin{pmatrix} d_L \\ s_L \\ b_L \end{pmatrix}. \tag{12.138}$$

The charged weak current (left chiral) becomes, in the mass basis,

$$\begin{aligned} \mathcal{J}^{\mu}_{\text{weak}} &= \overline{\psi}^{(\text{gauge})}_{u,L} \gamma^{\mu} \psi^{(\text{gauge})}_{d,L} = \left(\overline{\psi}^{(\text{mass})}_{u,L} V^{\dagger}_{uL}\right) \gamma^{\mu} \left(V_{dL} \psi^{(\text{mass})}_{d,L}\right) \\ &= \overline{\psi}^{(\text{mass})}_{u,L} \gamma^{\mu} \left(V^{\dagger}_{uL} V_{dL}\right) \psi^{(\text{mass})}_{d,L}, \end{aligned} \tag{12.139}$$

so the observable mixing matrix in the quark sector is

$$V_{\text{CKM}} := V^{\dagger}_{uL} V_{dL}. \tag{12.140}$$

In the family–lifted picture, this expression refines to the *relative family holonomy* (Proposition 12.115)

$$V_{\text{CKM}}(x) = \left[V^{\dagger}_{uL}\, \mathcal{P} e^{i \int_{\gamma} \mathcal{A}^{(F)}}\, V_{dL}\right](x), \tag{12.141}$$

with γ the relevant worldline segment and $\mathcal{A}^{(F)}$ the flat family connection (Definitions 12.106 and 12.110). When γ is contractible in $\mathcal{X}_q \setminus \mathfrak{D}$, the Wilson line reduces to the identity and (12.141) collapses to (12.140).

Definition 12.117 (CKM and PMNS mixing matrices)**.** Let V_{uL}, V_{dL} be the unitary diagonalizers of the up– and down–type quark Yukawa operators on $\mathcal{F}$ (eq. (12.137)). The quark mixing matrix is

$$V_{\text{CKM}} := V^{\dagger}_{uL} V_{dL},$$

governing charged–current flavor change (CKM: [542, 543]). For leptons, let U_{eL} diagonalize the charged–lepton mass operator. In the Dirac–neutrino case, let $V_{\nu L}$ diagonalize the neutrino Dirac mass and set

$$U_{\text{PMNS}} := U^{\dagger}_{eL} V_{\nu L}.$$

For Majorana neutrinos, the neutral mass matrix M_ν is complex symmetric and diagonalized by a Takagi factor U_ν with $U_\nu^T M_\nu U_\nu = \text{diag}(m_i)$; then $U_{\text{PMNS}} = U^{\dagger}_{eL} U_\nu$ (Definition 12.141). In the family–lifted transport, both CKM and PMNS acquire the Wilson–line refinement as in Proposition 12.115 and eq. (12.134).

CP structure and holonomy. Complex phases in V_{CKM} and U_{PMNS} arise from (i) relative phases in the diagonalizers V_{fL} and (ii) the family Wilson line $\mathcal{P} e^{i \int_{\gamma} \mathcal{A}^{(F)}}$. Flat but nontrivial Z_3 monodromy already supplies discrete phases in closed loops (Definition 12.106 and Theorem 12.111); localized curvature near $\mathfrak{D}$ can generate path–class dependence and Jarlskog–type invariants measurable by holosymmetric tomography (Remark 12.116 and Section 2.4). Diagnostic realized reference configuration hygiene (gauge–parameter independence and KL positivity) for fermionic poles/residues remains as in Proposition 12.88.

Physical Consequences

The unitary flavor rotations encapsulated by V_{CKM} and U_{PMNS} (see Definition 12.117, and the holonomy refinement in Proposition 12.115 and eq. (12.134)) have direct, testable implications in weak processes, once the charged current is written in the mass basis (eq. (12.139)). In the quark sector, the Cabibbo–Kobayashi–Maskawa construction organizes all charged–current flavor transitions into a single 3×3 unitary matrix with one irreducible CP–violating phase for three generations. This framework subsumes the original Cabibbo mixing of the first two generations [542] and, in accommodating kaon and B–meson phenomenology within a renormalizable theory, necessarily predicts CP violation [543]. Empirically, the CKM picture accounts—quantitatively and with high precision—for tree– and loop–level processes, including GIM suppressions, unitarity–triangle relations, and the observed pattern of direct and indirect CP violation (global summaries in [565]). Within the IQR setting, these statements hold on certified plateaux with gauge–parameter–independent poles/residues and KL–positivity (Proposition 12.88), and the weak interactions are mediated by the electroweak vector sector identified from the Operator–Bundle connection (Section 12.4.4 and Definitions 12.95 and 12.102). From a cosmological perspective, the CKM phase supplies one of Sakharov's ingredients (CP violation) for baryogenesis; however, within the minimal SM it is far too feeble to generate the observed baryon asymmetry, motivating additional CP–violating sources beyond the quark sector [566].

Cosmological CP and the limits of the CKM phase. Baryogenesis requires the three Sakharov conditions: (i) baryon–number violation, (ii) violation of C and CP, and (iii) a departure from thermal equilibrium. Within the IQR framework the quark–sector source of CP violation is the phase in the CKM matrix V_{CKM} (defined in Definition 12.117 and arising inevitably from the Yukawa/mass–basis misalignment in Section 12.5.6). Thus condition (ii) is present in principle.

GIM suppression of FCNCs. In the quark sector, the absence of tree–level flavor–changing neutral currents and their strong loop–level suppression follow from CKM unitarity together with the structure of the electroweak vector sector (Section 12.4.4, Definition 12.117, and eq. (12.139)). Loop amplitudes for $q \to q'$ $(q \neq q')$ transitions take the schematic form

$$\mathcal{A}(q \to q') \;\propto\; \sum_{i=u,c,t} V_{iq}\, V_{iq'}^{*}\, F\!\left(\frac{m_i^2}{M_W^2}\right), \tag{12.142}$$

with F a loop function determined by W–exchange (boxes/penguins). If F were mass–independent, the off–diagonal unitarity identity

$$\sum_{i=u,c,t} V_{iq}\, V_{iq'}^{*} \;=\; 0 \qquad (q \neq q') \tag{12.143}$$

would cancel the amplitude exactly (the GIM limit). The residual is therefore controlled by mass splittings through differences $F(m_i^2) - F(m_j^2)$: for kaon FCNCs it scales with $(m_c^2 - m_u^2)/M_W^2$, while in $b \to s, d$ it is dominated by the heavy top. Within the IQR diagnostic realized reference configuration, these loop observables inherit gauge–parameter

independence and KL positivity from Proposition 12.88; the vector fields entering F are the electroweak components of the Operator–Bundle connection (Section 12.4.4), and the CKM factors arise from the Yukawa/mass–basis misalignment (Section 12.5.6). The family bundle is flat and neutral under G_{SM} (Definitions 12.106 and 12.110), so it does not introduce new FCNC sources; GIM suppression remains intact.

Why CKM alone is insufficient. Two facts render the quark–sector CP source too feeble to account for the observed baryon asymmetry. First, the rephasing–invariant strength of CKM CP violation is small and, in the hot electroweak plasma, CP–odd observables suffer additional GIM–type mass–difference suppressions tied to the same flavor structure that controls loop amplitudes (cf. above). Second, with the realized electroweak symmetry breaking sector (Section 12.4.5) the electroweak transition is a smooth crossover, not a strongly first–order phase transition, so the out–of–equilibrium condition (iii) is not efficiently met. Consequently, CKM CP violation by itself cannot generate the observed baryon–to–photon ratio.

Motivation for additional CP sources and the IQR placement. This shortfall motivates CP–violating sources beyond the minimal quark sector [566]. In the present framework there are two natural loci for such phases that do not disturb established CKM/GIM phenomenology: (a) the lepton sector, where neutrino masses and mixings—Dirac or Majorana—arise from the Yukawa/seesaw structure, with leptonic CP phases entering U_{PMNS} (Definition 12.117) and high–scale dynamics enabling leptogenesis (Section 12.7.4); and (b) the family bundle, where flat Z_3 holonomy supplies discrete complex phases and localized curvature can seed rephasing–invariant loop phases, as captured by the holonomy formulation of mixing and CP in Proposition 12.115 and Remark 12.116. These ingredients furnish additional CP violation while remaining compatible with the weak–interaction mediation by the Operator–Bundle connection and the electroweak sector already established (Section 12.4.4 and Definition 12.102).

Leptogenesis in the Yukawa/seesaw sector (most conservative). Within the mass–generation pipeline, the lepton sector naturally carries additional CP phases through the PMNS structure (Definition 12.117), including the Dirac phase and—if neutrinos are Majorana—two extra Majorana phases. Introducing right–handed (sterile) neutrinos via the type–I seesaw (Section 12.7.4) supplies out–of–equilibrium, CP–violating decays at high scale; electroweak sphalerons then reprocess the generated lepton asymmetry into a baryon asymmetry. This route leaves CKM/GIM phenomenology intact and uses only the ingredients already present in the EWSB/Yukawa formalism (Sections 12.4.6 and 12.5.6). It yields crisp experimental handles: the leptonic Dirac phase in oscillations, neutrinoless double–beta decay for the Majorana case, and consistency of the inferred seesaw scale with low–energy neutrino data.

Family holonomy as a controlled, geometric CP source. The flat Z_3 family bundle $(\mathcal{F}, \mathcal{A}^{(F)})$ provides a geometric locus for complex phases: discrete holonomy phases arise even for flat transport, and localized curvature of $\mathcal{A}^{(F)}$ generates rephasing–invariant loop phases encoded in Wilson lines (Definition 12.106, Proposition 12.115, and Remark 12.116). Because $\mathcal{F}$ is SM–neutral and couples only through covariant transport, it does not upset GIM suppression

or introduce FCNCs, and it remains compatible with Stage I/II viability (Section 12.5.4 and Corollary 12.113). These phases are operationally accessible via holosymmetric interferometry and tomography (Section 2.4) while preserving the established weak–interaction structure.

Strengthening the electroweak departure from equilibrium via the Tantrakyma potential. The minimal Higgs sector yields a crossover electroweak transition and fails to deliver a sufficiently strong departure from equilibrium. In the present framework, the order of the transition is governed by the Tantrakyma potential $\mathcal{V}_\Theta$ and the scalar–kymon spectrum (Section 12.4.5, Definition 9.343, and eq. (9.185)). Allowing additional (SM–singlet or SM–doublet) Tantrakyma scalar excitations, or higher operators consistent with diagnostic realized reference configuration discipline and typed compatibility, can drive a *strong first–order* transition without introducing new light gauge bosons, thereby maintaining anomaly freedom and HSP viability (Section 12.5.4 and Theorem 12.42) and preserving the Nielsen/KL hygiene of poles and residues on certified plateaux (Proposition 12.88). The framework does not assert that the CKM phase alone suffices; rather, it supplies structured, testable avenues—seesaw leptogenesis, geometric family holonomy, and a strengthened EWSB potential—each consistent with covariant transport, diagnostic realized reference configuration/Nielsen–KL criteria, HSP selection, and all previously established SM–level constraints, while preserving CKM/GIM phenomenology.

In the lepton sector, the Pontecorvo–Maki–Nakagawa–Sakata paradigm provides the correct organizing principle for flavor change in flight: the misalignment between flavor and mass bases encoded in U_{PMNS} produces neutrino oscillations across solar, atmospheric, reactor, and accelerator baselines [545, 544, 565]. The large leptonic mixing angles (in contrast to the hierarchical CKM pattern) are a distinctive empirical signature. In the IQR framework, U_{PMNS} is defined as in Definition 12.117 and admits the same holonomy refinement as Proposition 12.115. If neutrino masses arise via a Majorana mechanism such as the type–I seesaw (Section 12.7.4), U_{PMNS} generally contains additional (Majorana) phases that do not affect oscillations but can enter lepton–number–violating amplitudes. Moreover, ultraviolet dynamics responsible for light neutrino masses can supply new CP–violating phases whose out–of–equilibrium decays seed a lepton asymmetry, later reprocessed by sphalerons into the baryon asymmetry—*leptogenesis*—linking neutrino physics to early–universe cosmology in a predictive effective framework [567, 566]. In the family–lifted picture, closed–loop phases can also receive contributions from family holonomy/curvature (Definition 12.106 and Remark 12.116), providing an operational handle through holosymmetric interferometry (Section 2.4).

12.5.6 The Inevitable Emergence of Mixing Matrices

The origin of the CKM and PMNS matrices is not an added postulate but a direct consequence of the fermion mass–generation formalism (Section 12.4.6). In the IQR framework, masses arise from Yukawa couplings to the Tantrakyma doublet $H \sim (1,2)_{+1/2}$ and its VEV $\langle H \rangle = \bar{\Theta}/\sqrt{2}$ (Definition 9.343, eq. (9.185), and Proposition 12.90). In the family–lifted description, the Yukawa sector is encoded as endomorphisms on the flat family bundle $(\mathcal{F}, \mathcal{A}^{(F)})$ with mass operators $M_f = \bar{\Theta}\, Y_f/\sqrt{2}$ (Definitions 12.110 and 12.114 and eq. (12.133)).

1. The Yukawa interaction as a matrix equation

The effective Yukawa action for quarks is promoted to the three–family fibre by taking the left/right fields as generation triplets and the Yukawas as 3×3 complex matrices (Definitions 12.82 and 12.114):

$$\psi_{u,L}^{(\text{gauge})} = \begin{pmatrix} u'_L \\ c'_L \\ t'_L \end{pmatrix}, \qquad \psi_{d,L}^{(\text{gauge})} = \begin{pmatrix} d'_L \\ s'_L \\ b'_L \end{pmatrix}, \quad \text{(eq. (12.135))}$$

$$\mathcal{L}_{\text{Yukawa}} \;=\; -\,\overline{\psi}_{u,L}^{(\text{gauge})}\, Y_u\, \tilde{H}\, \psi_{u,R}^{(\text{gauge})} \;-\; \overline{\psi}_{d,L}^{(\text{gauge})}\, Y_d\, H\, \psi_{d,R}^{(\text{gauge})} + \text{h.c.},$$

with $Y_{u,d} \in \mathrm{Mat}_{3\times 3}(\mathbb{C})$ constrained only by G_{SM} gauge invariance (Proposition 12.90).

2. The mass matrices are not diagonal

After EWSB (Section 12.4.5),

$$M_u = \frac{\bar{\Theta}}{\sqrt{2}}\, Y_u, \qquad M_d = \frac{\bar{\Theta}}{\sqrt{2}}\, Y_d,$$

and there is no principle forcing $Y_{u,d}$ to be diagonal; generically $M_{u,d}$ are complex matrices on $\mathcal{F}$ (Definition 12.114). Thus the *gauge* eigenstates (diagonal weak couplings) do not coincide with the *mass* eigenstates (definite propagation), cf. eqs. (12.135) and (12.136).

3. Diagonalization and the physical mass eigenstates

Physical fields are obtained by bi–unitary diagonalization (eq. (12.137)):

$$M_u^{\text{diag}} = V_{uL}^{\dagger} M_u V_{uR}, \qquad M_d^{\text{diag}} = V_{dL}^{\dagger} M_d V_{dR},$$

which induces the family rotations between gauge and mass bases (eqs. (12.136) and (12.138)):

$$\begin{pmatrix} u'_L \\ c'_L \\ t'_L \end{pmatrix} = V_{uL} \begin{pmatrix} u_L \\ c_L \\ t_L \end{pmatrix}, \qquad \begin{pmatrix} d'_L \\ s'_L \\ b'_L \end{pmatrix} = V_{dL} \begin{pmatrix} d_L \\ s_L \\ b_L \end{pmatrix}.$$

4. The mixing matrix appears in the weak current

The charged current couples gauge eigenstates; rewriting it in the mass basis produces the mixing matrix (eq. (12.139)):

$$\mathcal{J}^{\mu}_{\text{weak}} = \overline{\psi}_{u,L}^{(\text{mass})} \gamma^{\mu}\, (V_{uL}^{\dagger} V_{dL})\, \psi_{d,L}^{(\text{mass})}.$$

Hence

$$V_{\text{CKM}} := V_{uL}^{\dagger} V_{dL}, \qquad \text{and analogously } U_{\text{PMNS}} \text{ in the lepton sector} \quad (\textit{Definition } 12.117).$$

In the family–lifted transport, both acquire a Wilson–line refinement as a *relative family holonomy* (Proposition 12.115 and eq. (12.134)):

$$V_{\mathrm{CKM}}(x) = \Big[V_{uL}^{\dagger}\,\mathcal{P}e^{\,i\int_{\gamma}\mathcal{A}^{(F)}}\,V_{dL}\Big](x), \quad U_{\mathrm{PMNS}}(x) = \Big[U_{eL}^{\dagger}\,\mathcal{P}e^{\,i\int_{\gamma}\mathcal{A}^{(F)}}\,V_{\nu L}\Big](x),$$

which reduce to the usual forms on contractible γ in $\mathcal{X}_q\backslash\mathfrak{D}$ (Definition 12.106). All statements are taken on certified plateaux with Nielsen–stable, KL–positive poles/residues (Proposition 12.88), and weak vectors follow from the Operator–Bundle connection (Section 12.4.4 and Definition 12.102).

Conclusion. Thus, CKM/PMNS mixing *inevitably* emerges from (i) non–aligned Yukawa/-mass and weak–current bases (Section 12.4.6 and Definition 12.114) and, in the geometric lift, (ii) relative family holonomy of $(\mathcal{F}, \mathcal{A}^{(F)})$ (Definition 12.106 and Proposition 12.115). No extra postulates are required.

12.5.7 Hadrons as Composite Kymon States

Principle 12.2 (Hadrons as Color–Singlet Composite Kymon States)**.** *Hadrons are* composite relational configurations *of quark–kymons, formed under the confining dynamics of the color gauge symmetry* $\mathrm{SU}(3)_C$*. They are not elementary excitations but BRST-physical, color-neutral bound states:*

$$\boxed{\textit{Hadrons} \;\equiv\; \textit{color–singlet bound states of confined quark–kymons under } \mathrm{SU}(3)_C.} \tag{12.144}$$

Foundational references on confinement, Wilson loops/lines, and the color–singlet characterization of BRST-physical states include [389, 82, 91, 108, 442].

Foundational prerequisites (already established).

1. **Quark–kymons.** Quarks are spin-$\frac{1}{2}$ fermionic kymon excitations of $\hat{\mathbf{\Theta}}(x)$ in the fundamental **3** of $\mathrm{SU}(3)_C$ (Definitions 12.81 and 12.82); standard background on color triplets and flavor assignments can be found in [82, 91].

2. **Gluons.** The color gauge bosons are the $\mathrm{SU}(3)_C$ components of the Operator–Bundle connection $\mathcal{A}_\mu$ (Section 12.4.4 and Definitions 12.95 and 12.96); see also the Yang–Mills construction [123] and textbook expositions [82, 91].

3. **Confinement & physical Hilbert space.** Dynamics encoded in the non-abelian curvature and the Tantrakyma sector of the relational Hamiltonian $\mathbb{H}^{(\Theta)}$ restrict BRST-physical states to color-neutral sectors in the emergent Hilbert space $\mathcal{H}_{\mathrm{em}}$ (Definitions 4.287, 6.54 and 12.126); compare the Kugo–Ojima BRST criterion and Wilson's confinement criterion [108, 442, 389].

Definition 12.118 (Gauge-invariant meson/baryon operators (color singlets))**.** Let $\hat{\psi}_i(x)$ be a quark–kymon field with color index $i \in \{1,2,3\}$ (Definition 12.82). For any spacetime path $\gamma : x \to y$, let

$$\mathcal{W}(x \to y) \;:=\; \mathcal{P}\exp\!\Big\{ ig_s \int_\gamma G^a_\mu T_a \, dx^\mu \Big\}$$

be the $\mathrm{SU}(3)_C$ Wilson line built from the color part of $\mathcal{A}_\mu$ (Section 12.4.4 and Definition 12.96); see [389, 82]. Then the standard color-singlet composites are: **Meson.**

$$\hat{\mathcal{M}}_\Gamma(x,y) \;:=\; \bar{\hat{\psi}}_i(x)\,\Gamma\,\big[\mathcal{W}(x\to y)\big]^i{}_j\,\hat{\psi}^j(y). \tag{12.145}$$

Baryon (at junction x_0).

$$\begin{aligned}\hat{\mathcal{B}}_{\Gamma_1\Gamma_2\Gamma_3}(x_1,x_2,x_3) \;:=\; &\epsilon_{ijk}\,\hat{\psi}^i(x_1)\,\mathcal{W}(x_1\to x_0)\,\Gamma_1 \\ &\times \hat{\psi}^j(x_2)\,\mathcal{W}(x_2\to x_0)\,\Gamma_2 \\ &\times \hat{\psi}^k(x_3)\,\mathcal{W}(x_3\to x_0)\,\Gamma_3\,.\end{aligned} \tag{12.146}$$

with Γ, Γ_k spinor/isospin structures chosen for the desired quantum numbers (parity, spin, flavor) [82, 91]. By construction, (12.145) and (12.146) are gauge-invariant under local $\mathrm{SU}(3)_C$ and thus descend to the BRST-physical space $H^0(\hat{Q}_{\mathrm{BRST}})$ (Definition 12.126); compare [108, 442].

Remark 12.119 (Gauge invariance of Wilson–line–dressed meson and baryon operators)**.** Let $U(x) \in SU(3)_C$ denote a local color transformation. Quark fields in the fundamental and antifundamental transform as

$$\hat{\psi}(y) \;\mapsto\; U(y)\,\hat{\psi}(y), \qquad \bar{\hat{\psi}}(x) \;\mapsto\; \bar{\hat{\psi}}(x)\,U^\dagger(x),$$

while the fundamental Wilson line along $\gamma : x \to y$ transforms as

$$\mathcal{W}(x\to y) \;\mapsto\; U(x)\,\mathcal{W}(x\to y)\,U^\dagger(y).$$

Cf. Definitions 12.96 and 12.118 and Section 12.4.4; see [389, 82, 91].

Meson. The operator in eq. (12.145),

$$\bar{\hat{\psi}}_i(x)\,\Gamma\,\big[\mathcal{W}(x\to y)\big]^i{}_j\,\hat{\psi}^j(y),$$

transforms as

$$\bar{\hat{\psi}}_k(x)\,U^\dagger(x)^k{}_i\,\Gamma\,U(x)^i{}_a\,\mathcal{W}(x\to y)^a{}_b\,U^\dagger(y)^b{}_j\,U(y)^j{}_l\,\hat{\psi}^l(y),$$

and the adjacent factors $U^\dagger(x)U(x)$ and $U^\dagger(y)U(y)$ cancel, proving gauge invariance. Equivalently, the form $\bar{\hat{\psi}}(y)\,\Gamma\,\mathcal{W}(y\to x)\,\hat{\psi}(x)$ is also invariant since $\mathcal{W}(y\to x) = \mathcal{W}(x\to y)^\dagger$ in $SU(3)$ (e.g. [82, 91]). Hence the arrow orientation in eq. (12.145) is correct.

Baryon. For the junction point x_0 in eq. (12.146), parallel transport each quark to x_0:

$$\Psi_a(x_0) \;:=\; \big[\mathcal{W}(x_i\to x_0)\,\hat{\psi}(x_i)\big]_a \;\mapsto\; U(x_0)_a{}^b\,\Psi_b(x_0).$$

Then the color–antisymmetric contraction $\epsilon_{abc}\,\Psi_a(x_0)\,\Psi_b(x_0)\,\Psi_c(x_0)$ is invariant because ϵ_{abc} is the $SU(3)$ singlet tensor (determinant representation), so $\epsilon_{abc}\,U(x_0)_a{}^{a'}U(x_0)_b{}^{b'}U(x_0)_c{}^{c'} = \epsilon_{a'b'c'}$ [82, 91]. Thus the arrows $x_i \to x_0$ in eq. (12.146) are also correct.

In both cases, reversing an arrow requires the corresponding Hermitian conjugation and index placement so that the Wilson line transports the color index from the field insertion point to the common contraction point, preserving local gauge invariance (Definition 12.118; [389, 82]).

Proposition 12.120 (Color singletness and BRST physicality)**.** *Let* $\mathsf{P}_{\rm phys}$ *project to* $H^0(\hat{Q}_{\rm BRST})$ *on the certified plateau* $\mathfrak{P}$ *(Proposition 12.88). Then for any composite* $\hat{\mathcal{O}} \in \{\hat{\mathcal{M}}_\Gamma, \hat{\mathcal{B}}_{\Gamma_1\Gamma_2\Gamma_3}\}$ *of Definition 12.118,*

$$\mathsf{P}_{\rm phys}\,\hat{\mathcal{O}}\,\mathsf{P}_{\rm phys} \;=\; \hat{\mathcal{O}},$$

and any non-singlet composite is BRST-exact (or has vanishing matrix elements between physical states). Hence only color–singlet composites create/annihilate physical hadrons. Sketch. *BRST cohomology enforces Gauss constraints at the operator level; local* $\mathrm{SU}(3)_C$ *invariance is necessary for non-vanishing overlaps with* $\mathsf{P}_{\rm phys}$ *(Definition 12.126).* □

Proposition 12.121 (Diagnostic realized reference configuration certification: hadron poles from color–singlet kernels)**.** *Let* $\hat{\mathcal{O}}$ *be as in Definition 12.118. On* $\mathfrak{P}$ *with tags* $(P_x(\lambda_{\rm UV}), K^{(\ell)}_{\rm IR})$ *(Definitions 7.29 and 7.39), the realized two-point*

$$\Delta_{\mathcal{O}\mathcal{O}}(x-y) \;:=\; \left\langle \mathsf{P}_{\rm phys}\,\hat{\mathcal{O}}(x)\,\hat{\mathcal{O}}^\dagger(y)\,\mathsf{P}_{\rm phys}\right\rangle$$

is gauge-parameter independent and KL-positive (Proposition 12.88). Simple poles in its momentum–space projection certify hadron one-particle states (mesons/baryons) with positive residues; continuum thresholds encode multi-hadron scattering channels. Spectral isolation can be implemented band-tamely by Riesz selectors exactly as in the vector/scalar kymon cases (Definition 10.7 and Section 10.2).

Remark 12.122 (Typed placement and TT-purity)**.** Hadronic currents decompose into the τ–typed channels of the weave ($\bar\psi\psi \in S^{(b)}$, $\bar\psi\, i\gamma_5\psi \in \tilde{S}^{(p)}$, $\bar\psi\gamma_\mu\psi \in A^{(\pm)}$) (Definition 10.4 and eq. (10.10)). Each typed summand is $\Pi^{[\mathrm{TT}]}_{\bar\Sigma}$–null, so hadron excitations remain orthogonal to the TT graviton sector (Lemmas 10.17 and 10.59 and Corollary 10.125). This preserves graviton purity while allowing full hadronic dynamics in the non-TT sector.

Remark 12.123 (Operator–bundle mediation and confinement meters)**.** The strong dynamics entering Definition 12.118 are precisely the $\mathrm{SU}(3)_C$ components of the Operator–Bundle connection (Section 12.4.4). Confinement manifests through the behavior of gauge-invariant loop/-line functionals and the absence of isolated color charges in $H^0(\hat{Q}_{\rm BRST})$ (Definition 12.126), while bound-state poles are read on $\mathfrak{P}$ with Nielsen/KL hygiene (Proposition 12.88). No TT contamination arises (Remark 12.122).

Remark 12.124 (Flavor structure vs. compositeness)**.** Hadron compositeness is controlled by $\mathrm{SU}(3)_C$ dynamics and color singletness; CKM factors only enter charged-current transitions between hadrons through the underlying quark currents (Definition 12.117 and eq. (12.139)). The hadronization map from quark–level operators to hadron observables inherits the diagnostic realized reference configuration discipline (typed channels, Riesz isolation, BRST projection) used elsewhere in the Tantrakyma sector (Definitions 9.60 and 10.7).

Group–Theoretic Realization. The color classification of hadrons follows from the decomposition of $\mathrm{SU}(3)_C$ tensor products for the fundamental $\mathbf{3}$ and antifundamental $\bar{\mathbf{3}}$ representations carried by quark– and antiquark–kymons (Definition 12.82); standard treatments of these decompositions appear in [82, 91].

Baryons as three–quark states:

$$\mathbf{3} \otimes \mathbf{3} \otimes \mathbf{3} \;=\; \mathbf{1} \oplus \mathbf{8} \oplus \mathbf{8} \oplus \mathbf{10}, \tag{12.147}$$

the standard SU(3) decomposition for three fundamentals [82, 91]; only the singlet $\mathbf{1}$ sector yields observable (BRST–physical) baryons by color–singlet projection (Definition 12.126 and Proposition 12.120; cf. [108, 442]).

Mesons as quark–antiquark states:

$$\mathbf{3} \otimes \bar{\mathbf{3}} \;=\; \mathbf{1} \oplus \mathbf{8}, \tag{12.148}$$

identifying the $\mathbf{1}$ as the physically admissible meson subspace [82, 91] (the octet is color–adjoint and does not survive the BRST projection to the physical space).
For later use we record the explicit color–singlet projectors. In the meson channel,

$$\left(P^{(1)}_{\mathrm{mes}}\right)^{\;i}_{j}{}^{k}_{l} \;=\; \frac{1}{3}\,\delta^{i}{}_{l}\,\delta^{k}{}_{j}, \qquad \text{so} \quad \left(P^{(1)}_{\mathrm{mes}} \cdot \psi\bar{\psi}\right) \;\sim\; \tfrac{1}{3}\,\bar{\psi}_i\,\psi^i, \tag{12.149}$$

and in the baryon channel (totally antisymmetric color),

$$\left(P^{(1)}_{\mathrm{bar}}\right)_{ijk}{}^{i'j'k'} \;=\; \frac{1}{6}\,\epsilon_{ijk}\,\epsilon^{i'j'k'}, \qquad \text{so} \quad \left(P^{(1)}_{\mathrm{bar}} \cdot \psi\psi\psi\right) \;\sim\; \epsilon_{ijk}\,\psi^i\psi^j\psi^k. \tag{12.150}$$

These follow from the SU(3) invariants $\delta^i{}_j$ and ϵ_{ijk} and are standard in QCD textbooks [82, 91]. When quarks are not co–located, gauge–invariant dressing by $\mathrm{SU}(3)_C$ Wilson lines built from the color part of the Operator–Bundle connection $\mathcal{A}_\mu$ (Section 12.4.4 and Definition 12.96) yields the meson/baryon operators of Definition 12.118 [389]; these descend to $H^0(\hat{Q}_{\mathrm{BRST}})$ as color singlets (Definition 12.126 and Proposition 12.120; see also [108, 442]).

Relational Ontology. A hadron is not a static sum of constituent fields but a *nonperturbative, entangled modal state* of the Tantrakyma sector whose stability and observables (mass, spin, moments) are read from gauge–invariant correlators with diagnostic realized reference configuration hygiene (Nielsen/KL) on certified plateaux (Definition 6.54 and Propositions 12.88 and 12.121; see [120, 82, 91]). Formally, for $n = 2$ (mesons) or $n = 3$ (baryons) quark–kymons, the color–singlet projection

$$P_{\mathrm{singlet}} \cdot \mathcal{H}^{(n)}_{\mathrm{quark}} \;\subset\; \mathcal{H}_{\mathrm{em}}, \qquad P_{\mathrm{singlet}} \in \mathrm{End}\Big[(\mathbb{C}^3)^{\otimes n}\Big]^{\mathrm{SU}(3)_C}, \tag{12.151}$$

selects the physical subspace inside the emergent Hilbert space $\mathcal{H}_{\mathrm{em}}$ (Definition 4.287); this implements color confinement at the operator level in line with BRST/Gauss constraints [108, 442]. Here the quark–kymons are the fermionic excitations of $\hat{\boldsymbol{\Theta}}(x)$ carrying color in $\mathbb{C}^3$ (Definitions 12.81 and 12.82), and $(\mathbb{C}^3)^{\otimes n}$ arises from combining n such fields with appropriate Wilson–line dressings (Definition 12.118; cf. [389, 82]). The projector P_{singlet} matches the confinement picture where only color–neutral composites survive as BRST–physical states [108, 442].

Gauge and Cohomological Consistency. Physical hadronic states lie in BRST cohomology,

$$\mathcal{H}_{\text{hadron}} \subset \ker(\delta_{\text{BRST}}) \cap \mathcal{H}_{\text{em}}, \tag{12.152}$$

ensuring unitarity and gauge consistency (Section 12.3 and Definition 12.126; see [445, 106, 442, 108]). Two–point functions of gauge–invariant meson/baryon operators are gauge–parameter independent and KL–positive on $\mathfrak{P}$ (Propositions 12.88 and 12.121; cf. [120, 82, 91]); simple poles in their spectral densities certify one–hadron states, while cuts/thresholds encode multi–hadron channels [82, 91]. Typed placement in the τ–sector preserves TT graviton purity (Definition 10.4, eq. (10.10), Lemma 10.17, and Corollary 10.125).

Hadrons emerge as the first composite manifestations of the strong sector built from quark–kymons (Section 12.4.1) and the $\mathrm{SU}(3)_C$ components of the unified Operator–Bundle connection (Section 12.4.4 and Definition 12.95); this is the standard QCD picture of confinement and color singlets [389, 82, 91]. Their existence is not an extra postulate but a structural necessity of nonabelian color dynamics acting on the Tantrakyma excitations. Flavor/family structures (CKM/PMNS, family fibre) enter only through the underlying quark currents and remain compatible with color singletness (Definitions 12.106, 12.110 and 12.117 and eq. (12.139)), while confinement and BRST projection enforce the observable spectrum's gauge invariance (Definition 12.126 and Proposition 12.120; cf. [108, 442]).

Spectral Branching under Color–Gauge Projection

Corollary 12.125 (Spectral branching under color–gauge projection)**.** *Let $\mathcal{H}_{\text{em}}$ be the emergent (GNS) Hilbert space (Definition 4.287) with BRST projection P_{phys} to $H^0(\hat{Q}_{\text{BRST}})$. Fix $n = 2$ (mesons) or $n = 3$ (baryons) and let $\mathcal{H}^{(n)}_{\text{quark}} \subset \mathcal{H}_{\text{em}}$ denote the n-quark sector generated by quark–kymon fields in the fundamental of $\mathrm{SU}(3)_C$ (Definitions 12.81 and 12.82). Let the local color action be implemented on the color factor $(\mathbb{C}^3)^{\otimes n}$ and define the orthogonal singlet projector*

$$P_{\text{singlet}} \in \mathrm{End}\Big[(\mathbb{C}^3)^{\otimes n}\Big]^{\mathrm{SU}(3)_C}, \qquad \textit{cf. the decompositions in eqs. (12.147) and (12.148).} \tag{12.153}$$

Then the physical hadronic subspace arises by color–gauge projection

$$\mathcal{H}_{\text{hadron}} := P_{\text{singlet}}\Big(\mathcal{H}^{(n)}_{\text{quark}}\Big) \subset \mathcal{H}_{\text{em}}, \tag{12.154}$$

and enjoys the following properties:

(i) Spectral reduction (block–diagonalization)

The relational Hamiltonian $\mathbb{H}^{(\Theta)}$ (Definition 6.54) commutes with the color action by gauge invariance of the Tantrakyma sector (Definition 9.345), hence with P_{singlet}: $[\mathbb{H}^{(\Theta)}, P_{\text{singlet}}] = 0$. Therefore

$$\mathbb{H}^{(\Theta)}\Big|_{\mathcal{H}_{\text{hadron}}} = \mathbb{H}^{(\text{hadron})}, \tag{12.155}$$

and the spectrum on $\mathcal{H}_{\text{hadron}}$ consists of color–singlet bound–state energies. On certified plateaux $\mathfrak{P}$ the corresponding two–point functions of gauge–invariant meson/baryon operators (Definition 12.118) are gauge–parameter independent and KL–positive, and their simple poles certify hadron one–particle states (Propositions 12.88 and 12.121).

(ii) Gauge–invariant (BRST–physical) sector

Every $\psi \in \mathcal{H}_{\text{hadron}}$ is BRST–closed, $\delta_{\text{BRST}}\psi = 0$, and survives the projection to $H^0(\hat{Q}_{\text{BRST}})$ (Definition 12.126 and Proposition 12.120):

$$\mathcal{H}_{\text{hadron}} \subset \ker(\delta_{\text{BRST}}) \cap \mathcal{H}_{\text{em}} \qquad \textit{(eq. (12.152)).} \tag{12.156}$$

(iii) Emergent symmetry branching

The passage $\mathrm{SU}(3)_C \curvearrowright (\mathbb{C}^3)^{\otimes n} \rightsquigarrow \mathbf{1}_{\text{phys}}$ implements the first nontrivial gauge–invariant tensor subspace in the emergent spectrum: only the trivial irrep survives the physical projection, while non–singlet sectors are BRST–exact or have vanishing matrix elements between physical states (Proposition 12.120). For spatially separated quarks the Wilson–line dressing by the color component of the Operator–Bundle connection $\mathcal{A}_\mu$ preserves gauge invariance (Section 12.4.4 and Definitions 12.96 and 12.118).

Interpretation. Equation (12.154) exhibits *spectral branching* driven by internal gauge symmetry: the color algebra induces a decomposition of the n-quark tensor product into irreps (eqs. (12.147) and (12.148)), and the operator–algebraic projection P_{singlet} selects the gauge–invariant branch realized in the observable spectrum. Within the diagnostic realized reference configuration discipline, hadronic poles are read from color–singlet correlators with Nielsen/KL hygiene (Propositions 12.88 and 12.121), typed placement keeps them TT–null in the weave (Definition 10.4, eq. (10.10), and Lemma 10.17), and flavor structures (CKM/PMNS, family bundle) enter only through the underlying quark currents without spoiling color singletness (Definitions 12.106 and 12.117 and eq. (12.139)).

Color–Singlet Projection and Irreducible Decomposition

Definition 12.126 (Color–singlet projection and irreducible decomposition)**.** Let $\mathcal{H}^{(n)}_{\text{quark}}$ denote the color sector of the n-quark Hilbert space (with each quark in the fundamental $\mathbf{3}$ of $SU(3)_C$) and $\mathrm{End}\big[(\mathbb{C}^3)^{\otimes n}\big]$ its endomorphism algebra. The $SU(3)_C$-invariant (color–singlet) subspaces are obtained by projecting onto the trivial component in the standard tensor–product decompositions $\mathbf{3} \otimes \bar{\mathbf{3}} = \mathbf{1} \oplus \mathbf{8}$ and $\mathbf{3} \otimes \mathbf{3} \otimes \mathbf{3} = \mathbf{1} \oplus \mathbf{8} \oplus \mathbf{8} \oplus \mathbf{10}$ (eqs. (12.147) and (12.148)). Explicitly:

1. **Mesons ($n = 2$, $3 \otimes \bar{3}$).** Writing a color basis $\{|r\rangle, |g\rangle, |b\rangle\}$ for $\mathbb{C}^3$ and the dual $\{|\bar{r}\rangle, |\bar{g}\rangle, |\bar{b}\rangle\}$ for $\bar{\mathbb{C}}^3$, the unique $SU(3)_C$-invariant singlet is

$$|\text{meson singlet}\rangle = \frac{1}{\sqrt{3}}\Big(|r\rangle|\bar{r}\rangle + |g\rangle|\bar{g}\rangle + |b\rangle|\bar{b}\rangle\Big). \tag{12.157}$$

   The orthogonal projector onto the singlet line is

$$P^{(\text{meson})}_{\text{singlet}} := |\text{meson singlet}\rangle\langle\text{meson singlet}| \in \mathrm{End}\Big[\mathbb{C}^3 \otimes \bar{\mathbb{C}}^3\Big]^{SU(3)_C}. \tag{12.158}$$

2. **Baryons ($n = 3$, $3 \otimes 3 \otimes 3$).** With ϵ_{ijk} the Levi–Civita tensor ($i, j, k \in \{r, g, b\}$), the totally antisymmetric color singlet is

$$|\text{baryon singlet}\rangle = \frac{1}{\sqrt{6}} \sum_{i,j,k=r,g,b} \epsilon_{ijk}\, |q_i\rangle|q_j\rangle|q_k\rangle, \tag{12.159}$$

and the associated rank–one projector is

$$P^{(\text{baryon})}_{\text{singlet}} := |\text{baryon singlet}\rangle\langle\text{baryon singlet}| \ \in \ \text{End}\Big[(\mathbb{C}^3)^{\otimes 3}\Big]^{SU(3)_C}. \tag{12.160}$$

Gauge–invariant realization and interpretation. The abstract singlet projector P_{singlet} here makes concrete the color–neutral projection used in eqs. (12.151) and (12.154): it maps the multi–quark color tensor space $\mathcal{H}^{(n)}_{\text{quark}}$ onto the $SU(3)_C$–invariant subspace that hosts physical hadrons [82, 91]. For spatially separated quarks, local gauge invariance is maintained by inserting color Wilson lines built from the $SU(3)_C$ component of the Operator–Bundle connection $\mathcal{A}_\mu$ (Section 12.4.4 and Definition 12.96), as in the gauge–invariant meson/baryon operators of Definition 12.118 [389, 82]. The orthogonal complements of $P^{(\text{meson})}_{\text{singlet}}$ and $P^{(\text{baryon})}_{\text{singlet}}$ span the octet/decuplet color sectors, which do not survive the BRST projection to the physical space $H^0(\hat{Q}_{\text{BRST}})$ (Definition 12.126 and Proposition 12.120) and are therefore unobservable as isolated asymptotic states due to confinement [108, 442, 389].

Color Charge: The "Flavor" of the Strong Force

In Quantum Chromodynamics (QCD), the strong interaction is described by a local $SU(3)_C$ gauge symmetry acting on an internal *color* degree of freedom (Section 12.4.4; see [82, 91, 123]). Quark–kymons (Definitions 12.81 and 12.82) transform in the fundamental representation **3** of $SU(3)_C$, whose canonical basis vectors will be denoted

$$|r\rangle, \quad |g\rangle, \quad |b\rangle \ \in \ \mathbb{C}^3.$$

The corresponding gauge bosons (gluons) are the $SU(3)_C$ components of the Operator–Bundle connection $\mathcal{A}_\mu$ with field strength $\mathcal{F}_{\mu\nu}$ (Section 12.4.4 and Definition 12.96); standard expositions include [82, 91].

Anti-colors and the conjugate representation. Antiquark states transform in the conjugate representation $\bar{\mathbf{3}} \simeq \bar{\mathbb{C}}^3$ with basis

$$|\bar{r}\rangle, \quad |\bar{g}\rangle, \quad |\bar{b}\rangle,$$

and carry the *anti-color* charges $\bar{r}, \bar{g}, \bar{b}$ [82, 91]. Thus, two-quark color Hilbert space for a quark–antiquark pair is $\mathbb{C}^3 \otimes \bar{\mathbb{C}}^3$, while the three–quark color space is $(\mathbb{C}^3)^{\otimes 3}$.

Color neutrality (singlet condition) and confinement. Only *color–neutral* (gauge–invariant) combinations appear as physical asymptotic states. Formally, hadrons reside in the $SU(3)_C$–*singlet* subrepresentations obtained by projecting multi–quark color tensors with the color–singlet projector P_{singlet} (Definition 12.126 and eq. (12.151); cf. [82, 91]). Non–singlet color sectors are removed by BRST cohomology when passing to the physical Hilbert space $H^0(\hat{Q}_{\text{BRST}})$ (Definition 12.126 and Proposition 12.120); see [445, 106, 108, 442].

- **Baryons** (three–quark states) live in the **1** component of $\mathbf{3} \otimes \mathbf{3} \otimes \mathbf{3} = \mathbf{1} \oplus \mathbf{8} \oplus \mathbf{8} \oplus \mathbf{10}$ (eq. (12.147)); textbook derivations of the decomposition appear in [82, 91]. The unique totally antisymmetric singlet vector is given in eq. (12.159) and projected by eq. (12.160).

- **Mesons** (quark–antiquark states) live in the $\mathbf{1}$ component of $\mathbf{3}\otimes\bar{\mathbf{3}} = \mathbf{1}\oplus\mathbf{8}$ (eq. (12.148)); the normalized singlet is

$$|\text{meson singlet}\rangle = \frac{1}{\sqrt{3}}\Big(|r\rangle|\bar{r}\rangle + |g\rangle|\bar{g}\rangle + |b\rangle|\bar{b}\rangle\Big), \tag{12.161}$$

  cf. eq. (12.157), with projector $P^{(\text{meson})}_{\text{singlet}}$ in eq. (12.158) [82, 91].

Gauge invariance for separated quarks. When the quark and antiquark (or the three quarks) are inserted at distinct spacetime points, local $SU(3)_C$ invariance is maintained by *Wilson lines* built from the color component of $\mathcal{A}_\mu$ [389]; the resulting meson/baryon operators $\hat{\mathcal{M}}, \hat{\mathcal{B}}$ in Definition 12.118 are gauge–invariant and create color–singlet states from the vacuum. These composite operators are BRST–physical on the certified plateau and their two–point functions exhibit KL–positive hadron poles with gauge–parameter–independent residues (Propositions 12.88 and 12.121; cf. [120, 82, 91]).

Equation (12.161) is the unique invariant color wavefunction for a quark–antiquark pair: it is unchanged under global $SU(3)_C$ transformations and therefore represents the physical color content of a meson [82, 91]. Analogously, the totally antisymmetric ϵ_{ijk} structure encodes the baryon color singlet (eq. (12.159)). In the present framework hadrons are thus *color–singlet composite kymon states* (Principle 12.2), realized as gauge–invariant subspaces of the emergent Hilbert space $\mathcal{H}_{\text{em}}$ (eq. (12.154)), with all color dynamics mediated by the $SU(3)_C$ sector of the Operator–Bundle connection (Section 12.4.4; see [82, 91]).

Gauge Invariance and Idempotency of the Color–Singlet Projector

Lemma 12.127 (Gauge invariance, idempotency, and orthogonality of the color–singlet projector)**.** *Let $P_{\text{singlet}} \in \text{End}\big[(\mathbb{C}^3)^{\otimes n}\big]$ denote the orthogonal projection onto the $SU(3)_C$–invariant (trivial) subrepresentation in the n-quark color space $\mathcal{H}^{(n)}_{\text{quark}}$, as in Definition 12.126. Then:*

(i) Idempotency

$P^2_{\text{singlet}} = P_{\text{singlet}}$.

(ii) Hermiticity

$P^\dagger_{\text{singlet}} = P_{\text{singlet}}$. *Hence P_{singlet} is the orthogonal projector onto*

$$\Big[(\mathbb{C}^3)^{\otimes n}\Big]^{SU(3)_C} \subset \mathcal{H}^{(n)}_{\text{quark}}.$$

(iii) Global gauge invariance (commutant property)

For any $U \in SU(3)_C$ acting in the fundamental on each tensor factor,

$$\Big[P_{\text{singlet}},\, U^{\otimes n}\Big] = 0, \qquad \textit{i.e.} \quad P_{\text{singlet}} \in \text{End}\Big[(\mathbb{C}^3)^{\otimes n}\Big]^{SU(3)_C}.$$

Moreover, for the two cases relevant to hadrons,

$$\textit{mesons } (n=2): \quad \mathbf{3}\otimes\bar{\mathbf{3}} = \mathbf{1}\oplus\mathbf{8} \quad \textit{(eq. (12.148))}, \tag{12.162}$$

$$\textit{baryons } (n=3): \quad \mathbf{3}\otimes\mathbf{3}\otimes\mathbf{3} = \mathbf{1}\oplus\mathbf{8}\oplus\mathbf{8}\oplus\mathbf{10} \quad \textit{(eq. (12.147))}, \tag{12.163}$$

the singlet occurs with multiplicity one; the explicit projectors in eqs. (12.158) and (12.160) realize P_{singlet}.

Proof. A standard construction (Peter–Weyl averaging; cf. the character/orthogonality meters in Proposition 12.10 and Corollary 12.48) defines

$$P_{\text{singlet}} := \int_{SU(3)_C} d\mu(U)\, U^{\otimes n}, \tag{12.164}$$

with $d\mu$ the normalized Haar measure. This operator projects onto the subspace fixed by $U^{\otimes n}$ for all U.

Idempotency. Using left–invariance of $d\mu$ and the group law,

$$P_{\text{singlet}}^2 = \int d\mu(U) \int d\mu(V)\, U^{\otimes n} V^{\otimes n} = \int d\mu(W)\, W^{\otimes n} = P_{\text{singlet}}.$$

Hermiticity. Since $(U^{\otimes n})^\dagger = (U^\dagger)^{\otimes n}$ and Haar measure is invariant under inversion/conjugation,

$$P_{\text{singlet}}^\dagger = \int d\mu(U)\, (U^{\otimes n})^\dagger = \int d\mu(U)\, (U^\dagger)^{\otimes n} = \int d\mu(V)\, V^{\otimes n} = P_{\text{singlet}}.$$

Commutation with global color. For any fixed $G \in SU(3)_C$,

$$G^{\otimes n} P_{\text{singlet}} = \int d\mu(U)\, (GU)^{\otimes n} = \int d\mu(V)\, V^{\otimes n} = P_{\text{singlet}},$$

again by left–invariance of $d\mu$. The same argument gives $P_{\text{singlet}} G^{\otimes n} = P_{\text{singlet}}$, hence $[P_{\text{singlet}}, G^{\otimes n}] = 0$.

Uniqueness in the meson/baryon sectors. The decomposition $\mathbf{3} \otimes \bar{\mathbf{3}} = \mathbf{1} \oplus \mathbf{8}$ and $\mathbf{3}^{\otimes 3} = \mathbf{1} \oplus \mathbf{8} \oplus \mathbf{8} \oplus \mathbf{10}$ (eqs. (12.147) and (12.148)) shows the singlet appears once; therefore the explicit rank–one projectors $P_{\text{singlet}}^{(\text{meson})}$ and $P_{\text{singlet}}^{(\text{baryon})}$ given in eqs. (12.158) and (12.160) coincide with (12.164).

Local gauge covariance and Wilson lines. For quark insertions at distinct spacetime points, local $SU(3)_C$ acts as $U(x_i)$ on each color factor. The gauge–invariant composites in Definition 12.118 parallel–transport all color indices to a common junction x_0 with Wilson lines built from the color component of the connection $\mathcal{A}_\mu$ (Section 12.4.4 and Definition 12.96), after which the above global argument applies at x_0. Hence the corresponding singlet projector commutes with local gauge transformations on the composite operators and descends to the BRST–physical space (Definition 12.126 and Proposition 12.120).

All three statements follow. □

The operator P_{singlet} provides a gauge–covariant, orthogonal decomposition of the multi–quark color space into the observable (singlet) and confined (non–singlet) sectors. In particular, insertion of P_{singlet} in the composite operators of Definition 12.118 selects precisely the hadronic color content; its commutation with global color and its covariance under local color (via Wilson transport) guarantee compatibility with the Operator–Bundle connection and BRST cohomology (Section 12.4.4, Definition 12.96, and Proposition 12.120).

Color–Singlet Projection as the Paradigm of Gauge–Induced Spectral Filtering

Proposition 12.128 (Color–singlet projection as canonical gauge–induced spectral filtering)**.** *Let $\mathcal{H}^{(n)}_{\text{quark}} \subset \mathcal{H}_{\text{em}}$ be the n-quark sector of the emergent Hilbert space (Definition 4.287), with local color acting in the fundamental on each factor and parallel transport provided by the $SU(3)_C$ component of the Operator–Bundle connection (Section 12.4.4 and Definition 12.96). Let P_{singlet} be the color–singlet projector of Definition 12.126, equivalently given by the Haar average in Lemma 12.127, eq.* (12.164)*. Then*

$$\boxed{\mathcal{H}_{\text{hadron}} \;:=\; P_{\text{singlet}}\,\mathcal{H}^{(n)}_{\text{quark}} \;\subset\; \mathcal{H}_{\text{em}}} \tag{12.165}$$

defines the hadronic *(color–neutral) subspace and furnishes the canonical example of an* internal gauge–induced spectral filter*: a gauge–invariant projector acting on a reducible tensor product that isolates the first nontrivial, BRST–physical sector of the emergent spectrum (Definition 12.118, Proposition 12.120, and Corollary 12.125).*

Structural consequences.

(i) Spectral filtering of the relational Hamiltonian The Tantrakyma-sector Hamiltonian on the hadronic subspace $\mathbb{H}^{(\Theta)}$ (Definition 6.54) restricts to the effective Hamiltonian

$$\mathbb{H}^{(\text{hadron})} \;:=\; \mathbb{H}^{(\Theta)}\Big|_{\mathcal{H}_{\text{hadron}}}.$$

whose eigenstates are color–singlet bound states (mesons/baryons). On certified plateaux $\mathfrak{P}$ (Definitions 7.29 and 7.39), hadron poles in color–singlet two–point kernels are gauge–parameter independent and KL–positive (Propositions 12.88 and 12.121); isolated poles can be band–tamely selected by Riesz projectors (Definition 10.7 and Section 10.2).

(ii) Internal symmetry reduction (color decomposition) The color tensor product decomposes as $(\mathbb{C}^3)^{\otimes n} \cong \mathbf{1} \oplus$ (nontrivial irreps) (eqs. (12.147) and (12.148)), and $P_{\text{singlet}} \in \text{End}\Big[(\mathbb{C}^3)^{\otimes n}\Big]^{SU(3)_C}$ (Lemma 12.127) implements the algebraic reduction $(\mathbb{C}^3)^{\otimes n} \to \mathbf{1}$, thereby realizing the physical color–neutral sector (Definition 12.118 and eqs. (12.145) and (12.146)).

(iii) Physical admissibility via BRST cohomology Since P_{singlet} commutes with global color and the composite operators are made locally gauge–invariant by Wilson transport (Section 12.4.4 and Definition 12.118), their matrix elements survive the BRST projection and descend to $H^0(\hat{Q}_{\text{BRST}})$ (Proposition 12.120 and eq. (12.152)). Non–singlet composites are BRST–exact or have vanishing overlaps between physical states.

(iv) Typed placement and graviton purity Hadronic currents reside in the τ–typed channels of the weave and are $\Pi^{[\text{TT}]}_{\bar{\Sigma}}$–null ($\bar{\psi}\psi \in S^{(b)}$, $\bar{\psi} i\gamma_5 \psi \in \tilde{S}^{(p)}$, $\bar{\psi}\gamma_\mu\psi \in A^{(\pm)}$; Definition 10.4, eq. (10.10), and Remark 12.122), ensuring no contamination of the TT graviton sector (Lemmas 10.17 and 10.59 and Corollary 10.125).

Equation (12.165) elevates color singletness from a representation–theoretic statement to an *operational* selection rule on the emergent spectrum: internal gauge symmetry drives an algebraic projection that determines which subspaces carry physical poles (hadrons) once the Operator–Bundle dynamics are enforced (Section 12.4.4, Definition 12.96, and Proposition 12.121). In this sense, the color–singlet projector P_{singlet} is the prototype of gauge–induced ontological selection within the IQR framework, fully compatible with diagnostic realized reference configuration discipline (Nielsen/KL hygiene), BRST cohomology, and the global architecture of $\mathcal{H}_{\text{em}}$.

12.6 BRST Cohomology and the Hadronic Physical Space

We now make explicit the cohomological projection from the full gauge–fixed state space to the hadronic (color–singlet) sector in the $SU(3)_C$ subsector of the Operator–Bundle connection (Section 12.4.4 and Definition 12.96). Our treatment follows the standard BRST construction but is adapted to the operator–bundle and diagnostic realized reference configuration discipline of this chapter (Definitions 6.54, 6.192, 7.29 and 7.39 and Proposition 12.88); see [91, Ch. 16] for background on BRST quantization.

12.6.1 Constraints, BRST Charge, and Nilpotency

Let $\hat{G}^a(x)$ denote the Gauss generators of color rotations (first–class constraints) built from the Yang–Mills electric field and the matter color current of quark–kymons (Definition 12.82),

$$\hat{G}^a(x) \;=\; \big(D_i E^i\big)^a(x) \;-\; g_s\,\hat{J}^{0,a}(x), \qquad a=1,\dots,8, \tag{12.166}$$

with D_i the adjoint covariant derivative of the $SU(3)_C$ part of the Operator–Bundle connection (Section 12.4.4). Introduce fermionic ghost/antighost fields $c^a(x), \bar{c}^a(x)$ and the bosonic Nakanishi–Lautrup field $B^a(x)$ with canonical (anti)commutation relations and ghost number assignments (standard covariant gauge). In a covariant gauge, the *BRST charge* $\hat{Q}_{\text{BRST}}$ takes the local form (suppressing normal–ordering symbols)

$$\hat{Q}_{\text{BRST}} = \int d^3x \left(\hat{G}^a\, c^a - \tfrac{1}{2} f^{abc}\, \hat{\mathcal{P}}^a_c\, c^b c^c + B^a\, \hat{\mathcal{P}}^a_{\bar{c}} \right), \tag{12.167}$$

where $\hat{\mathcal{P}}^a_c$ and $\hat{\mathcal{P}}^a_{\bar{c}}$ are the canonical momenta of c^a and $\bar{c}^a$. The BRST differential is $s(\cdot) = [\hat{Q}_{\text{BRST}}, \cdot\}$ (graded commutator). On a dense, gauge–invariant domain $\mathcal{D}$ one has the standard properties [91, Ch. 16]:

$$\hat{Q}^2_{\text{BRST}} = 0, \qquad [\hat{H}_{\text{total}}, \hat{Q}_{\text{BRST}}] = 0, \qquad [N_{\text{gh}}, \hat{Q}_{\text{BRST}}] = \hat{Q}_{\text{BRST}}, \tag{12.168}$$

with N_{gh} the ghost–number operator and $\hat{H}_{\text{total}}$ the full gauge–fixed Hamiltonian (Definition 6.54).

Definition 12.129 (Physical space as BRST cohomology)**.** The physical state space is the ghost–number–zero cohomology $\mathcal{H}_{\text{phys}} := H^0(\hat{Q}_{\text{BRST}}) = \ker \hat{Q}_{\text{BRST}} \cap \ker N_{\text{gh}} \big/ \operatorname{im} \hat{Q}_{\text{BRST}}$. On certified plateaux $\mathfrak{P}$ (Definitions 7.29 and 7.39) all meters are Nielsen–stable and KL–positive (Proposition 12.88).

12.6.2 Color Singletness and BRST Physicality

Recall the gauge–invariant meson/baryon composites built with Wilson transport (Definition 12.118 and eqs. (12.145) and (12.146)) and the color–singlet projector $P_{\text{singlet}} \in \text{End}\big[(\mathbb{C}^3)^{\otimes n}\big]^{SU(3)_C}$ (Definition 12.126 and Lemma 12.127). We now relate these to BRST cohomology.

Proposition 12.130 (Color singletness $\Longleftrightarrow$ BRST–physical hadron sector)**.** *Let $\mathcal{H}^{(n)}_{\text{quark}} \subset \mathcal{H}_{\text{em}}$ be the n–quark sector (Definition 4.287), and define the hadronic subspace by color projection $\mathcal{H}_{\text{hadron}} := P_{singlet}\,\mathcal{H}^{(n)}_{\text{quark}}$ (Corollary 12.125). Then*

$$\mathcal{H}_{\text{hadron}} \;\subset\; H^0(\hat{Q}_{\text{BRST}}) \cap \mathcal{H}_{\text{em}}, \qquad \textit{and} \qquad (\textit{non–singlet}) \;\Rightarrow\; \textit{BRST–exact or zero–overlap.} \tag{12.169}$$

Sketch. *(i) Gauge–invariant composites commute with local gauge rotations generated by $\hat{G}^a$; since the first term of $\hat{Q}_{\text{BRST}}$ is $\int \hat{G}^a c^a$, their BRST variation is proportional to ghost insertions that vanish on ghost–number–zero matrix elements unless the operator carries color. (ii) Wilson lines render $\hat{\mathcal{M}}_\Gamma, \hat{\mathcal{B}}_{\Gamma_1\Gamma_2\Gamma_3}$ locally gauge–invariant (Definition 12.118); therefore their insertions between states in the image of $P_{singlet}$ lie in $\ker \hat{Q}_{\text{BRST}}$. (iii) Any non–singlet operator/state can be written as a BRST variation or has vanishing overlaps after the BRST projection (Lemma 12.127).*

Corollary 12.131 (Diagnostic realized reference configuration certification of hadron poles)**.** *For any color–singlet composite $\hat{\mathcal{O}}$ of Definition 12.118, the two–point kernel $\Delta_{\mathcal{O}\mathcal{O}}(x-y) = \langle \mathsf{P}_{\text{phys}}\,\hat{\mathcal{O}}(x)\,\hat{\mathcal{O}}^\dagger(y)\,\mathsf{P}_{\text{phys}}\rangle$ on $\mathfrak{P}$ is gauge–parameter independent and KL–positive (Propositions 12.88 and 12.121). Simple momentum–space poles certify hadron one–particle states, with Riesz isolation available band–tamely (Definition 10.7 and Section 10.2).*

12.6.3 Typed Placement and Graviton Purity

Hadronic currents decompose into τ–typed channels of the weave $\big(\bar\psi\psi \in S^{(b)},\ \bar\psi i\gamma_5\psi \in \tilde{S}^{(p)},\ \bar\psi\gamma_\mu\psi \in A^{(\pm)}\big)$ (Definition 10.4 and eq. (10.10)). Each typed summand is $\Pi^{[\text{TT}]}_{\bar\Sigma}$–null, ensuring no TT contamination of the graviton sector (Lemmas 10.17 and 10.59 and Corollary 10.125); thus BRST–physical hadrons live entirely in the non–TT sector (Remark 12.122).

12.6.4 Family Neutrality and Flavor Structure

The family bundle $(\mathcal{F}, \mathcal{A}^{(F)})$ is flat, unitary, and neutral under G_{SM} (Definitions 12.106 and 12.110); it couples only through the total covariant derivative $\mathfrak{D}^{\text{tot}}_\mu$ (Definition 12.110 and eq. (12.128)). Hence the BRST cohomology for color acts trivially on $\mathcal{F}$ and *does not* modify color singletness. Flavor–changing charged currents in hadrons arise only through the CKM factors inherited from quark–level currents (Definition 12.117 and eq. (12.139)), while GIM suppression and electroweak mediation proceed via the $SU(2)_L{\times}U(1)_Y$ components of the Operator–Bundle connection (Section 12.4.4); see also Section 12.5.4 and Proposition 12.115.

Synthesis. Color singletness, implemented by P_{singlet} (Definition 12.126 and Lemma 12.127), is precisely the BRST cohomological criterion for hadronic physicality (Proposition 12.130). On the diagnostic realized reference configuration $\mathfrak{P}$, hadronic poles inherit Nielsen stability and KL positivity (Propositions 12.88 and 12.121), typed placement preserves TT graviton purity (eq. (10.10) and Lemma 10.17), and family structure factors trivially through the flat auxiliary bundle (Definition 12.106). Thus $\mathcal{H}_{\text{hadron}} = P_{\text{singlet}}\mathcal{H}^{(n)}_{\text{quark}}$ (Corollary 12.125) is the canonical BRST–physical realization of the hadronic sector within the Operator–Bundle formulation of the IQR framework.

12.6.5 Quartet Mechanism and Gauge–Invariant Reduction

Decompose the total gauge–fixed state space as $\mathcal{H}_{\text{tot}} = \mathcal{H}_{\text{YM}} \otimes \mathcal{H}_{\text{matter}} \otimes \mathcal{H}_{\text{gh}}$, where $\mathcal{H}_{\text{YM}}$ is built from gluon modes of the $SU(3)_C$ part of the Operator–Bundle connection (Section 12.4.4 and Definition 12.96), $\mathcal{H}_{\text{matter}}$ from quark/antiquark modes (Definition 12.82), and $\mathcal{H}_{\text{gh}}$ from the Fock space of $(c, \bar{c}, B)$ (covariant gauge). The *Kugo–Ojima quartet mechanism* (in covariant gauges) asserts that longitudinal and timelike gluon modes together with ghost–antighost excitations assemble into BRST doublets (quartets) and do not contribute to the cohomology; only *transverse* gluon excitations survive, and then only through gauge–invariant composites [91, Ch. 16]. Concretely, there exists a contracting homotopy $\hat{K}$ of ghost number -1 on the unphysical subspace such that, with the BRST differential $s(\,\cdot\,) = [\hat{Q}_{\text{BRST}}, \,\cdot\,\}$ (Section 12.6.1 and eqs. (12.167) and (12.168)),

$$\{\hat{Q}_{\text{BRST}}, \hat{K}\} = \mathbb{I} \quad \text{on the longitudinal/timelike/ghost sector,} \tag{12.170}$$

whence those states are BRST–exact and represent no cohomology classes (Definition 12.129).

Proposition 12.132 (Gauge–invariant cohomology)**.** *Within ghost number zero, the BRST cohomology is isomorphic to the gauge–invariant subspace of the transverse sector:*

$$H^0(\hat{Q}_{\text{BRST}}) \;\cong\; \big(\mathcal{H}_{trans} \otimes \mathcal{H}_{matter}\big)^{SU(3)_C} \;=\; \ker\!\Big(\int d^3x\, \hat{G}^a(x)\Big) \cap \ker N_{\text{gh}}, \tag{12.171}$$

i.e. the subspace annihilated by the (integrated) Gauss generators $\hat{G}^a$ (eq. (12.166)). In particular, all color–nonsinglet vectors lie in $\operatorname{im}\hat{Q}_{\text{BRST}}$ *(or are orthogonal to* $\ker\hat{Q}_{\text{BRST}}$*) and do not represent cohomology classes.*

Idea. By the quartet mechanism (12.170), the unphysical sector (longitudinal, timelike, and ghosts) is contractible under $s = [\hat{Q}_{\text{BRST}}, \cdot\}$, leaving only $\mathcal{H}_{\text{trans}} \otimes \mathcal{H}_{\text{matter}}$. On this reduced sector, the BRST transformations coincide with the action of infinitesimal gauge transformations generated by $\hat{G}^a$; equivalently, s acts as the Lie derivative along the $SU(3)_C$ orbits. Thus H^0 reduces to the invariants under $SU(3)_C$, i.e. the kernel of the Gauss operators modulo their image. A Hodge–type decomposition adapted to the covariant gauge Krein structure (see [91]) renders the isomorphism (12.171) precise. In particular, any color–nonsinglet state is either s–exact or pairs to zero with $\ker\hat{Q}_{\text{BRST}}$, hence does not furnish a cohomology class. □

Proof. **Step 1 (Quartet reduction to the transverse sector).** Decompose $\mathcal{H}_{\mathrm{tot}} = \mathcal{H}_{\mathrm{YM}} \otimes \mathcal{H}_{\mathrm{matter}} \otimes \mathcal{H}_{\mathrm{gh}}$ with $\mathcal{H}_{\mathrm{YM}} = \mathcal{H}_{\mathrm{trans}} \oplus \mathcal{H}_{\mathrm{long/tl}}$ (transverse vs. longitudinal/timelike gluon modes). By the Kugo–Ojima quartet mechanism in covariant gauges, there exists a contracting homotopy $\hat{K}$ of ghost number -1 on $\mathcal{H}_{\mathrm{long/tl}} \otimes \mathcal{H}_{\mathrm{gh}}$ such that

$$\{\hat{Q}_{\mathrm{BRST}}, \hat{K}\} = \mathbb{I} \quad \text{on} \quad \mathcal{H}_{\mathrm{long/tl}} \otimes \mathcal{H}_{\mathrm{gh}},$$

(cf. (12.170)). Hence every state in this unphysical sector is BRST–exact and contributes trivially to cohomology; the inclusion $\mathcal{H}_{\mathrm{red}} := \mathcal{H}_{\mathrm{trans}} \otimes \mathcal{H}_{\mathrm{matter}} \hookrightarrow \mathcal{H}_{\mathrm{tot}}$ is a quasi–isomorphism of BRST complexes [108, 442, 91]. Therefore

$$H^{\bullet}(\hat{Q}_{\mathrm{BRST}}; \mathcal{H}_{\mathrm{tot}}) \;\cong\; H^{\bullet}(\hat{Q}_{\mathrm{BRST}}; \mathcal{H}_{\mathrm{red}}).$$

Step 2 (BRST as Chevalley–Eilenberg differential on the reduced sector). On $\mathcal{H}_{\mathrm{red}}$ (ghost number zero), the BRST variation of any (ghost–independent) state is generated by the Gauss operators:

$$s\Psi \equiv [\hat{Q}_{\mathrm{BRST}}, \Psi\} \;=\; c^a\, \hat{G}_a\, \Psi, \qquad \hat{G}_a = \int d^3x\, \hat{G}_a(x) \quad (\text{cf. } eqs.\ (12.166) and\ (12.167)).$$

Thus, restricted to ghost number zero, s acts as the Chevalley–Eilenberg differential for the $SU(3)_C$ action on $\mathcal{H}_{\mathrm{red}}$ (the Lie derivative along gauge orbits) [442]. In particular, $s\Psi = 0$ implies $\hat{G}_a\Psi = 0$ for all a (the c^a are independent generators of the ghost algebra).

Step 3 (Identification of H^0 with gauge invariants). Because s coincides with $c^a\hat{G}_a$ on ghost number zero, the closed states are exactly the $SU(3)_C$–invariants:

$$Z^0 = \ker(s|_{\mathcal{H}_{\mathrm{red}}}) = \ker\Big(\int d^3x\, \hat{G}^a(x)\Big) \cap \ker N_{\mathrm{gh}}.$$

Two closed states differ by an s–exact vector iff they differ by an infinitesimal gauge transformation. Therefore

$$H^0(\hat{Q}_{\mathrm{BRST}}) \;\cong\; Z^0/B^0 \;\cong\; \big(\mathcal{H}_{\mathrm{trans}} \otimes \mathcal{H}_{\mathrm{matter}}\big)^{SU(3)_C},$$

which is (12.171). A standard Hodge–type (Krein) decomposition adapted to the covariant gauge ($\hat{Q}_{\mathrm{BRST}}$, its adjoint, and the BRST Laplacian) gives canonical representatives that are simultaneously BRST–closed and annihilated by the adjoint, making the isomorphism precise [91, Ch. 16]; see also [442].

Step 4 (Exclusion of non–singlets). Let Ψ transform nontrivially under $SU(3)_C$. Then $\hat{G}_a\Psi \neq 0$, so at ghost number zero $s\Psi = c^a\hat{G}_a\Psi \neq 0$; hence $\Psi \notin \ker s$ and cannot represent a cohomology class. More generally, any state orthogonal to $\ker \hat{Q}_{\mathrm{BRST}}$ represents the zero class. Therefore color–nonsinglet vectors are either s–exact or not s–closed and do not contribute to H^0 [108, 442, 82].

Combining Steps 1–4 yields the claimed isomorphism (12.171). □

Consequences for hadrons. Combining Proposition 12.132 with the color–singlet projection $P_{\mathrm{singlet}} \in \mathrm{End}\big[(\mathbb{C}^3)^{\otimes n}\big]^{SU(3)_C}$ (Definition 12.126 and Lemma 12.127) and the gauge–invariant

composite operators (Definition 12.118 and eqs. (12.145) and (12.146)), one obtains the identification of the hadronic sector as the BRST–physical, color–singlet subspace (Proposition 12.130 and Corollary 12.125). On certified plateaux $\mathfrak{P}$ (Definitions 7.29 and 7.39), two–point functions of such composites are gauge–parameter independent and KL–positive (Proposition 12.88 and Corollary 12.131), so isolated poles certify physical hadrons without TT contamination (Section 12.6.3).

12.6.6 Explicit Computation on the Hadronic Sector

Let $\mathcal{H}_{\text{quark}}^{(n,\bar{m})} \subset \mathcal{H}_{\text{matter}}$ denote the n–quark/$\bar{m}$–antiquark subspace of the matter Fock space, and let $\mathcal{H}_{\text{gluon}}^{(k)} \subset \mathcal{H}_{\text{YM}}$ be the k–gluon subspace built from *transverse* $SU(3)_C$ modes (adjoint) of the Operator–Bundle connection (Section 12.4.4 and Definition 12.96). The total (gauge–fixed) state space factorizes as $\mathcal{H}_{\text{tot}} = \mathcal{H}_{\text{YM}} \otimes \mathcal{H}_{\text{matter}} \otimes \mathcal{H}_{\text{gh}}$, and the BRST physical space is the ghost–number–zero cohomology $H^0(\hat{Q}_{\text{BRST}})$ (Section 12.6.1 and Definition 12.129).

Write the $SU(3)_C$ representation carried by a fixed $(n, \bar{m}, k)$ particle–number sector as

$$\mathcal{R}_{n,\bar{m},k} := \left(\mathbf{3}^{\otimes n} \otimes \bar{\mathbf{3}}^{\otimes \bar{m}} \otimes \mathbf{8}^{\otimes k}\right)\downarrow_{SU(3)_C}, \qquad \mathbf{8} = \text{adjoint}. \tag{12.172}$$

The orthogonal projector onto the color singlet is the group (Haar) average

$$\mathcal{P}_{\text{sing}} = \int_{SU(3)} d\mu(g)\, U(g), \tag{12.173}$$

which equals the abstract singlet projector P_{singlet} of Definition 12.126 and Lemma 12.127. By construction $\mathcal{P}_{\text{sing}}$ commutes with the BRST charge and with the total Hamiltonian ($[\hat{H}_{\text{total}}, \hat{Q}_{\text{BRST}}] = 0$ and $\hat{Q}_{\text{BRST}}^2 = 0$; see Section 12.6.1 and eq. (12.168)), and thus preserves $H^0(\hat{Q}_{\text{BRST}})$.

Theorem 12.133 (BRST cohomology on the hadronic sector)**.** *For each particle–number sector $(n, \bar{m}, k)$,*

$$H^0(\hat{Q}_{\text{BRST}}) \;\cap\; \left(\mathcal{H}_{\text{quark}}^{(n,\bar{m})} \otimes \mathcal{H}_{\text{gluon}}^{(k)} \otimes \mathcal{H}_{\text{gh}}\right) \;\cong\; \left(\mathcal{R}_{n,\bar{m},k}\right)^{\mathbf{1}} \;\otimes\; \mathcal{H}_{\text{kin}}^{(n,\bar{m},k)}, \tag{12.174}$$

where $\left(\mathcal{R}_{n,\bar{m},k}\right)^{\mathbf{1}}$ is the multiplicity space of the trivial $SU(3)_C$ irrep in $\mathcal{R}_{n,\bar{m},k}$, and $\mathcal{H}_{\text{kin}}^{(n,\bar{m},k)}$ carries all non–color quantum numbers (spin, flavor, momenta, family labels $\mathcal{F}$, etc.; cf. Definition 12.110). Equivalently, at fixed kinematics the dimension *of the BRST cohomology equals the singlet multiplicity in the color tensor product.*

Proof. By the quartet mechanism and the contracting homotopy relation $\{\hat{Q}_{\text{BRST}}, \hat{K}\} = \mathbb{I}$ on the unphysical (longitudinal/timelike/ghost) sector (Section 12.6.5 and eq. (12.170)), the cohomology reduces to the transverse–matter sector. Proposition Proposition 12.132 identifies H^0 with the $SU(3)_C$–invariant subspace there. The group average $\mathcal{P}_{\text{sing}}$ (eq. (12.173)) projects onto the singlet multiplicity space $\left(\mathcal{R}_{n,\bar{m},k}\right)^{\mathbf{1}}$; its dimension is the character integral $m_{\mathbf{1}} = \int_{SU(3)} d\mu(g)\, \chi_{\mathcal{R}_{n,\bar{m},k}}(g)$. Ghost excitations have nonzero ghost number and drop out of H^0 (Section 12.6.1); kinematical quantum numbers factor. Hence (12.174) follows. □

Worked examples (quarks only).

(i) Mesons $(n = \bar{m} = 1,\ k = 0)$: $\mathbf{3} \otimes \bar{\mathbf{3}} = \mathbf{1} \oplus \mathbf{8} \Rightarrow$ one singlet per kinematic channel; the corresponding gauge–invariant operators are quark bilinears (with optional Wilson line) of Definition 12.118 and eq. (12.145).

(ii) Baryons $(n = 3,\ \bar{m} = 0,\ k = 0)$: $\mathbf{3}^{\otimes 3} = \mathbf{1} \oplus \mathbf{8} \oplus \mathbf{8} \oplus \mathbf{10} \Rightarrow$ one singlet per kinematic channel; the operator is the ϵ_{abc} color–antisymmetric trilinear of Definition 12.118 and eq. (12.146).

(iii) Pure glue $(n = \bar{m} = 0)$: singlets arise for even k from adjoint contractions; the simplest gauge–invariant operators are $\mathrm{Tr}\, F_{\mu\nu}F^{\mu\nu}$, $\mathrm{Tr}\, F_{\mu\nu}\tilde{F}^{\mu\nu}$ (glueball channels), built from $\mathcal{F}_{\mu\nu}$ (Definition 12.96).

Remark 12.134 (Operator–state map and diagnostic realized reference configuration certification)**.** Every cohomology class in (12.174) has a representative created from $|0\rangle$ by a BRST–closed, gauge–invariant composite (quark bilinears, baryon interpolators, or gluonic traces), possibly connected by Wilson lines to ensure locality in a chosen gauge (Definition 12.118). On certified plateaux $\mathfrak{P}$ with tags $(P_x(\lambda_{\mathrm{UV}}), K_{\mathrm{IR}}^{(\ell)})$ (Definitions 7.29 and 7.39), the two–point functions of such operators are gauge–parameter–independent and KL–positive (Proposition 12.88); simple poles certify hadron one–particle states, and Riesz selectors implement spectral isolation band–tamely (Definition 10.7 and Section 10.2).

12.6.7 Compatibility with Spectral Filtering and HSP

Recall the color–singlet (Haar) projector $\mathcal{P}_{\mathrm{sing}} = \int_{SU(3)} d\mu(g)\, U(g)$ (eq. (12.173)), which realizes the gauge–induced spectral filter of Proposition 12.128:

$$\mathcal{H}_{\mathrm{hadron}} = \mathcal{P}_{\mathrm{sing}}\Big(\mathcal{H}^{(\cdot)}_{\mathrm{quark}} \otimes \mathcal{H}^{(\cdot)}_{\mathrm{gluon}}\Big) \subset \mathcal{H}_{\mathrm{em}}.$$

By Lemma 12.127, $\mathcal{P}_{\mathrm{sing}}$ is Hermitian, idempotent, and lies in the commutant of the $SU(3)_C$ action; in particular it commutes with the (integrated) Gauss generators in (12.166). Since the BRST charge $\hat{Q}_{\mathrm{BRST}}$ implements gauge transformations on the transverse–matter sector (Sections 12.6.1 and 12.6.5, eq. (12.168), and Proposition 12.132), it follows that

$$[\hat{Q}_{\mathrm{BRST}}, \mathcal{P}_{\mathrm{sing}}] = 0, \qquad \mathcal{P}^2_{\mathrm{sing}} = \mathcal{P}_{\mathrm{sing}}.$$

Hence $\mathcal{P}_{\mathrm{sing}}$ is BRST–closed and acts trivially on cohomology classes: for any $|\Psi\rangle \in \ker \hat{Q}_{\mathrm{BRST}}$, $\mathcal{P}_{\mathrm{sing}}|\Psi\rangle$ represents the same class in $H^0(\hat{Q}_{\mathrm{BRST}})$ as the color–invariant component of $|\Psi\rangle$, while non–singlet components are BRST–exact (or orthogonal to $\ker \hat{Q}_{\mathrm{BRST}}$) by Proposition 12.132. Therefore,

$$H^0(\hat{Q}_{\mathrm{BRST}}) \;\cong\; \mathcal{P}_{\mathrm{sing}}\Big(\mathcal{H}^{(\cdot)}_{\mathrm{quark}} \otimes \mathcal{H}^{(\cdot)}_{\mathrm{gluon}}\Big), \tag{12.175}$$

i.e. the "gauge–induced spectral filtering'' narrative and the BRST–cohomological construction agree on the hadronic sector, in precise accord with Theorem 12.133.

HSP compatibility. Stage I of the Holosymmetric Selection Principle (HSP) demands unitarity and anomaly freedom for the realized dynamics (Section 12.2); these are enforced on the BRST–physical space $\mathcal{H}_{\rm phys} = H^0(\hat{Q}_{\rm BRST})$ (Definition 12.129). Equation (12.175) shows that the hadronic subspace is obtained by a *gauge–invariant* projector and therefore inherits Stage I viability. Stage II (relational orthogonality/economy) excludes sectors with residual color charge: such non–singlet components have zero overlap with the realized bright fringe and are dynamically suppressed (Section 12.5.4 and Corollary 12.113). On certified plateaux $\mathfrak{P}$ with tags $(P_x(\lambda_{\rm UV}), K_{\rm IR}^{(\ell)})$, the two–point functions of gauge–invariant hadron operators exhibit gauge–parameter independence and KL–positivity (Proposition 12.88), so spectral isolation via Riesz projectors remains band–tame (Definition 10.7 and Section 10.2).

12.6.8 Technical Notes on Domains and Positivity

All operator statements above are understood on a dense, gauge–invariant domain $\mathcal{D} \subset \mathcal{H}_{\rm tot} = \mathcal{H}_{\rm YM} \otimes \mathcal{H}_{\rm matter} \otimes \mathcal{H}_{\rm gh}$, on which the Gauss generators (12.166) and the BRST charge (12.167) are closable/essentially self–adjoint in the sense of covariant gauge quantization ([91, Ch. 16]; cf. Section 12.6.1). The natural inner product on $\mathcal{H}_{\rm tot}$ is a Krein metric adapted to the gauge–fixed theory; by the Kugo–Ojima quartet mechanism (Section 12.6.5) the unphysical (longitudinal/timelike/ghost) sector is contractible, and the restriction to the ghost–number–zero cohomology $H^0(\hat{Q}_{\rm BRST})$ yields a positive–definite inner product, so that $\mathcal{H}_{\rm phys}$ is a bona fide Hilbert space. On the diagnostic realized reference configuration $\mathfrak{P}$, positivity of residues and gauge–parameter independence of poles for gauge–invariant hadron interpolators are guaranteed by the Nielsen/KL hygiene (Proposition 12.88).

Summary. Combining Propositions 12.128 and 12.132 and Theorem 12.133 one obtains the explicit decomposition

$$H^0(\hat{Q}_{\rm BRST}) \;\cong\; \bigoplus_{n,\bar{m},k} \Big(\mathbf{3}^{\otimes n} \otimes \bar{\mathbf{3}}^{\otimes \bar{m}} \otimes \mathbf{8}^{\otimes k} \Big)^{\mathbf{1}} \;\otimes\; \mathcal{H}_{\rm kin}^{(n,\bar{m},k)},$$

i.e. the hadronic Hilbert space is *exactly* the color–singlet projection of quark–gluon Fock sectors, fully consistent with HSP Stage I/II and the operator–bundle realization of the strong sector (Section 12.4.4 and Definition 12.96).

12.7 Neutrino Masses, the Seesaw, and PMNS Mixing

12.7.1 Neutrino Physics and the Majorana Seesaw Mechanism

Neutrinos are structurally unique among fermionic kymons: they are electrically neutral and, unlike their charged partners, admit both Dirac and Majorana mass terms [82, 91]. Within the IQR framework their masses arise from the same Yukawa pipeline developed for all fermions (Section 12.4.6 and Proposition 12.90), together with the special status of a *gauge–singlet* right–handed neutrino kymon. Representation content is fixed in Definition 12.82: the left–handed lepton doublet $L_L \sim (1,2)_{-1/2}$ and a sterile $\nu_R \sim (1,1)_0$ [82, 91].

Dirac versus Majorana. Dirac masses descend from Yukawa couplings to the Tantrakyma doublet $H \sim (1,2)_{+1/2}$ via the family–lifted interaction maps (Section 12.4.6 and Definitions 12.110 and 12.114):

$$\mathcal{L}_Y \supset -\overline{L_L^{\mathrm{f}}}\,\tilde{H}\,Y_\nu\,\nu_R^{\mathrm{f}} + \mathrm{h.c.}, \qquad \tilde{H} := i\sigma_2 H^*,$$

which is gauge invariant by Proposition 12.90; after electroweak symmetry breaking (Section 12.4.5) with $\langle H\rangle = \bar{\Theta}/\sqrt{2}$, this gives the Dirac matrix $m_D = \bar{\Theta}\,Y_\nu/\sqrt{2}$ on the family fibre $\mathcal{F}$ (Definition 12.114; see [58, 59, 97, 82, 91]).

Definition 12.135 (Majorana mass term for sterile neutrinos)**.** Let $\nu_R \sim (1,1)_0$ and $\nu_R^c := \mathcal{C}\overline{\nu_R}^{\,T}$. The gauge–invariant Majorana term is

$$\mathcal{L}_{\mathrm{Maj}} = -\frac{1}{2}\left(\overline{\nu_R^c}\,M_R\,\nu_R + \mathrm{h.c.}\right), \tag{12.176}$$

with $M_R = M_R^T$ a complex symmetric 3×3 endomorphism on $\mathcal{F}$. It violates global lepton number by $\Delta L = 2$ but preserves G_{SM} gauge invariance (ν_R is a singlet) [129, 547, 91].

Type-I seesaw and the effective light mass. In the chiral basis (ν_L, ν_R^c), the full neutral–lepton mass matrix after EWSB is

$$\mathcal{M}_\nu = \begin{pmatrix} 0 & m_D \\ m_D^T & M_R \end{pmatrix}. \tag{12.177}$$

When $\|M_R\| \gg \|m_D\|$ (natural for a sterile singlet), block–diagonalization yields the familiar type-I seesaw:

$$m_\nu \;=\; -\,m_D\,M_R^{-1}\,m_D^T, \qquad M_N \;=\; M_R + \mathcal{O}(m_D^2/M_R), \tag{12.178}$$

so the active neutrinos are light Majorana modes with mass matrix m_ν, while heavy sterile states have masses near M_R [129, 547, 91, 82]. Equivalently, integrating out ν_R generates the (dimension–5) Weinberg operator

$$\mathcal{L}_{d=5} \;=\; -\tfrac{1}{2}\,\kappa\,(\overline{L_L}\tilde{H})\,(\tilde{H}^T L_L^c) + \mathrm{h.c.}, \qquad \kappa \;=\; Y_\nu\,M_R^{-1}\,Y_\nu^T,$$

consistent with the selection rules of Proposition 12.90; see standard treatments in [91, 82]. Diagnostic realized reference configuration hygiene (Nielsen/KL) for the realized poles/residues of the light and heavy modes follows from Proposition 12.88 (cf. [120, 91]).

12.7.2 PMNS Mixing in the Dirac and Majorana Cases

Let $M_e = \bar{\Theta}\,Y_e/\sqrt{2}$ be the charged–lepton mass operator on the family fibre $\mathcal{F}$ (mass operator from Definition 12.114, family lift in Definition 12.110). Biunitary diagonalization defines U_{eL} via

$$U_{eL}^\dagger\,M_e M_e^\dagger\,U_{eL} = \mathrm{diag}(m_e^2, m_\mu^2, m_\tau^2), \tag{12.179}$$

equivalently $U_{eL}^\dagger M_e U_{eR} = \mathrm{diag}(m_e, m_\mu, m_\tau)$ [82, 91].

For neutrinos.

(i) Dirac case With ν_R present but no Majorana term, $M_\nu = \bar{\Theta}\, Y_\nu/\sqrt{2}$ (Definition 12.139). Let $V_{\nu L}$ diagonalize $M_\nu M_\nu^\dagger$ (or biunitarily $V_{\nu L}^\dagger M_\nu V_{\nu R} = \mathrm{diag}(m_1, m_2, m_3)$). The leptonic mixing matrix is then $U_{\mathrm{PMNS}} = U_{eL}^\dagger V_{\nu L}$ (Definition 12.117 and Section 12.5.5; historically [545, 544]).

(ii) Majorana/seesaw case In the presence of a gauge–singlet Majorana term for ν_R (Definitions 12.135 and 12.139), and working in the Type–I seesaw regime (Section 12.7.4), the light neutrino mass matrix is a complex symmetric matrix

$$M_\nu \;=\; -\,M_D\, M_R^{-1}\, M_D^T,$$

as in eq. (12.188); see also [129, 547].

Choose a Takagi factor U_ν with

$$U_\nu^T M_\nu U_\nu \;=\; \mathrm{diag}(m_1, m_2, m_3), \qquad m_i \geq 0, \tag{12.180}$$

then $U_{\mathrm{PMNS}} = U_{eL}^\dagger U_\nu$ (Definition 12.141 and Section 12.7.5; see [82, 91] for standard parametrizations and phases).

In the family–lifted formulation, relative transport in the flat Z_3 family bundle $(\mathcal{F}, \mathcal{A}^{(F)})$ refines these relations by a Wilson line factor, as in $U_{\mathrm{PMNS}}(x) = U_{eL}^\dagger(x)\, \mathcal{P} e^{i\int_\gamma \mathcal{A}^{(F)}}\,(\cdots)$ (Definitions 12.106 and 12.110, Proposition 12.115, and eq. (12.134)).

Definition 12.136 (PMNS matrix (Majorana versus Dirac))**. (i) Majorana case.**
In the Majorana scenario, the mixing matrix takes the form

$$U_{\mathrm{PMNS}}(x) = \Big[\, U_{eL}^\dagger\, \mathcal{P} e^{i\int_\gamma \mathcal{A}^{(F)}}\, U_\nu \,\Big](x), \tag{12.181}$$

where U_ν is the Takagi factor from eq. (12.180) giving the relative family holonomy between the charged–lepton and neutrino diagonal frames (Proposition 12.115, eq. (12.134), and Definition 12.110). This carries one Dirac phase and, generically, two additional Majorana phases [82, 91].

(ii) Dirac case
If $M_R = 0$, diagonalize M_ν biunitarily: $V_{\nu L}^\dagger M_\nu V_{\nu R} = \mathrm{diag}(m_1, m_2, m_3)$ and set

$$U_{\mathrm{PMNS}}(x) = \Big[\, U_{eL}^\dagger\, \mathcal{P} e^{i\int_\gamma \mathcal{A}^{(F)}}\, V_{\nu L} \,\Big](x), \tag{12.182}$$

matching Definition 12.117 and Proposition 12.115; see [545, 544, 82, 91].

Remark 12.137 (Family transport, phases, and locality)**.** The Wilson line $\mathcal{P} e^{i\int_\gamma \mathcal{A}^{(F)}}$ implements parallel transport on the flat (or weakly curved) family bundle (Definitions 12.106 and 12.110). In a family–blind gauge (common local frame) it reduces to the identity and (12.181)–(12.182) collapse to the standard local definitions in Definition 12.117; nontrivial loops around $\mathfrak{D}$ contribute discrete Z_3 phases or controlled curvature phases (Remark 12.116).

Proposition 12.138 (IQR seesaw consistency and HSP compatibility)**.** *The type-I seesaw (cf. eqs.* (12.188) *and* (12.189)*) is compatible with all constraints previously established:*

(i) Gauge invariance/anomalies
$\nu_R \sim (1,1)_0$ *leaves the anomaly block unchanged; Stage I viability of the HSP is unaffected (Section 12.5.4, Theorem 12.42, and eq.* (12.46)*).*

(ii) Diagnostic realized reference configuration discipline
Realized propagator poles and residues for light/heavy neutrinos are gauge–parameter independent and KL–positive on certified plateaux (Proposition 12.88).

(iii) Family structure
$Y_\nu, M_R \in \mathrm{End}(\mathcal{F})$ *as in Definition 12.114; mixing is the relative family holonomy per Proposition 12.115.*

Proof. We verify each item in turn.

(1) Gauge invariance and anomalies. The Dirac Yukawa term $-\overline{L_L}\,\tilde{H}\,Y_\nu\,\nu_R + \text{h.c.}$ is gauge invariant because $L_L \sim (1,2)_{-1/2}$, $\tilde{H} \sim (1,2)_{+1/2}$ and $\nu_R \sim (1,1)_0$ (Definitions 12.82 and 12.139); the $SU(2)_L$ indices are contracted by $\varepsilon = i\sigma_2$ and the hypercharges sum to zero. The Majorana term $-\frac{1}{2}\,\overline{\nu_R^c}\,M_R\,\nu_R + \text{h.c.}$ is gauge invariant because ν_R is a total G_{SM} singlet (Definitions 12.135 and 12.139). Since ν_R carries no G_{SM} charge, it contributes zero to all entries in the anomaly block (eq. (12.46); cf. Proposition 12.26 and Corollary 12.27): $\mathrm{Tr}\,Y$, $\mathrm{Tr}\,Y^3$, $SU(2)^2$–$U(1)$, $SU(3)^2$–$U(1)$, and the mixed $U(1)$–grav^2 traces are unchanged. Hence Stage I viability in the Holosymmetric Selection Principle remains satisfied (Section 12.5.4 and Theorem 12.42).

(2) Diagnostic realized reference configuration discipline: gauge–parameter independence and KL positivity. In the chiral basis (ν_L, ν_R^c) the quadratic form is

$$\mathcal{L}_{\mathrm{mass}} \;=\; -\frac{1}{2}\begin{pmatrix}\overline{\nu_L} & \overline{\nu_R^c}\end{pmatrix}\begin{pmatrix}0 & M_D\\ M_D^T & M_R\end{pmatrix}\begin{pmatrix}\nu_L^c\\ \nu_R\end{pmatrix} + \text{h.c.},$$

with $M_D = \frac{v}{\sqrt{2}}Y_\nu$ and $\bar{\Theta} = \langle H\rangle$ (Definition 12.139). In the hierarchical regime $\|M_R\| \gg \|M_D\|$, block diagonalization (Schur complement/Takagi) yields the light/heavy spectra $M_\nu = -M_D M_R^{-1} M_D^T$ and $M_N \simeq M_R$ (Section 12.7.4 and eqs. (12.188) and (12.189); see also Proposition 12.140). The full fermionic two–point function is a 6×6 matrix in flavor-chirality space; by the Nielsen identities its pole locations and residues on certified plateaux are independent of the gauge-fixing parameter, provided one projects onto BRST–physical channels (Proposition 12.88). Since M_D and M_R are gauge scalars built from $\bar{\Theta}$ and $Y_\nu, M_R \in \mathrm{End}(\mathcal{F})$, the pole masses are manifestly gauge invariant. Moreover, the Källén–Lehmann spectral representation in the BRST cohomology guarantees positivity of spectral densities for the physical eigenmodes; on the diagnostic realized reference configuration $\mathfrak{P}$ this is enforced by the same KL/Nielsen hygiene used for the fermionic kymons (Proposition 12.88), with spectral isolation available via the standard Riesz selectors (Definition 10.7). Thus realized poles/residues for light and heavy neutrinos satisfy the diagnostic realized reference configuration criteria.

(3) Family structure and holonomy. In the family–lifted theory, $Y_\nu, M_R \in \mathrm{End}(\mathcal{F})$ act fibrewise on the rank-3 family bundle, and the total transport is governed by Definition 12.110

and eq. (12.128). The observable mixing in charged currents is the *relative* family holonomy between the diagonal frames of the charged–lepton and neutrino sectors,

$$U_{\mathrm{PMNS}}(x) = \Big[U_{eL}^{\dagger}\, \mathcal{P} e^{\,i\int_\gamma \mathcal{A}^{(F)}}\, U_\nu \Big](x) \quad \text{(Majorana)}$$
$$\text{or} \quad U_{\mathrm{PMNS}}(x) = \Big[U_{eL}^{\dagger}\, \mathcal{P} e^{\,i\int_\gamma \mathcal{A}^{(F)}}\, V_{\nu L} \Big](x) \quad \text{(Dirac)},$$

as established in Proposition 12.115, eq. (12.134), and Definition 12.136. Because $(\mathcal{F}, \mathcal{A}^{(F)})$ is auxiliary, flat (away from $\mathfrak{D}$), and neutral under G_{SM} (Definition 12.106), it introduces no new gauge currents, preserves GIM suppression of FCNCs, and leaves Stage I anomaly meters unchanged (Section 12.5.4). Stage II's relational economy further disfavours adding trivial family copies (Corollary 12.113). Hence the seesaw completion is fully compatible with the family holonomy picture and with the HSP selection criteria.

Collecting (1)–(3) proves the proposition. □

12.7.3 Dirac–Majorana Sector from the Tantrakyma

In the Tantrakyma sector, fermion masses arise from gauge–invariant Yukawa couplings to the electroweak doublet background $\bar{\Theta}$ (the EWSB order parameter) that descends from the Tantrakyma potential $\mathcal{V}_\Theta$ (Definition 9.335). Varying $\mathcal{V}_\Theta$ yields the Euler–Lagrange equations for $\hat{\boldsymbol{\Theta}}$ (eq. (9.199)); expanding about the realized vacuum $\bar{\Theta}$ produces the effective quadratic (mass) operators in the broken phase (Sections 9.18.5 and 12.4.5). For neutrinos, the presence of right–handed gauge singlets $\nu_R \sim (1,1)_0$ (see Definition 12.82) permits *both* Dirac and Majorana structures (Section 12.7.1).

Definition 12.139 (Neutrino Yukawa sector and Majorana term)**.** Let $L_L = (\nu_L, e_L)^T \sim (1,2)_{-1/2}$ and $H \sim (1,2)_{+1/2}$ be the Tantrakyma doublet that effects EWSB (Proposition 12.90 and Section 12.4.5), with $\tilde{H} := i\sigma_2 H^*$. The neutrino Yukawa Lagrangian and gauge–invariant Majorana term are

$$\mathcal{L}_Y^{(\nu)} \;=\; -\,\overline{L_L}\, Y_\nu\, \tilde{H}\, \nu_R \;+\; \text{h.c.}, \qquad \mathcal{L}_{\mathrm{Maj}} \;=\; -\frac{1}{2}\,\overline{\nu_R^{c}}\, M_R\, \nu_R \;+\; \text{h.c.}, \tag{12.183}$$

where Y_ν is a complex 3×3 Yukawa matrix and M_R is a complex symmetric 3×3 matrix (Definition 12.135). After EWSB with $\langle H \rangle = \bar{\Theta} = \frac{1}{\sqrt{2}}\binom{0}{v}$, the Dirac mass matrix is $M_D = \frac{v}{\sqrt{2}}\, Y_\nu$ (Section 12.4.6).

Family lift and covariance. In the family–lifted formulation (Definition 12.110), $Y_\nu, M_R \in \mathrm{End}(\mathcal{F})$ act fibrewise on the rank–3 family bundle with flat unitary connection $\mathcal{A}^{(F)}$ (Definition 12.106). Gauge invariance of $\mathcal{L}_Y^{(\nu)}$ and $\mathcal{L}_{\mathrm{Maj}}$ follows from the representation content in Definition 12.82 and the selection rules of Proposition 12.90; $\Delta L = 2$ for the Majorana term is allowed because ν_R is an $SU(3)_C \times SU(2)_L \times U(1)_Y$ singlet.

Type–I seesaw and effective masses. Collecting left– and right–handed neutral fields, the post–EWSB mass matrix in the (ν_L, ν_R^{c}) basis is

$$\mathcal{M}_\nu = \begin{pmatrix} 0 & M_D \\ M_D^{T} & M_R \end{pmatrix}, \tag{12.184}$$

cf. eq. (12.177). In the hierarchical regime $\|M_R\| \gg \|M_D\|$, block–diagonalization yields the standard type–I seesaw (eq. (12.178))

$$m_\nu = -\, M_D\, M_R^{-1}\, M_D^T, \qquad M_N = M_R + \mathcal{O}(M_D^2/M_R), \tag{12.185}$$

naturally explaining tiny active–neutrino masses [544, 546, 547]. The induced dimension–five Weinberg operator is $\kappa = (Y_\nu M_R^{-1} Y_\nu^T)$ and is consistent with the representation constraints of Proposition 12.90. On certified plateaux, the light/heavy poles and residues satisfy Nielsen identities and KL positivity by Proposition 12.88.

PMNS mixing. Diagonalization proceeds as in Section 12.7.2: in the Majorana case, Takagi factorization $U_\nu^T m_\nu U_\nu = \mathrm{diag}(m_1, m_2, m_3)$ together with the charged–lepton left diagonalizer U_{eL} gives $U_{\mathrm{PMNS}} = [\, U_{eL}^\dagger\, \mathcal{P} e^{i\int_\gamma \mathcal{A}^{(F)}}\, U_\nu\,]$ (Definition 12.136 and Proposition 12.115). In the Dirac limit $M_R = 0$, the biunitary diagonalization reproduces Definition 12.117.

Historical note. The Dirac–Majorana synthesis and $\Delta L = 2$ dynamics as the natural route to tiny neutrino masses and oscillations are classic insights [544, 546, 547], here realized within the Tantrakyma/Yukawa pipeline (Section 12.4.6) and the family–covariant transport structure (Definitions 12.106 and 12.110).

12.7.4 Type-I seesaw from block diagonalization

EFT prelude (Weinberg operator). At energies $E \ll \|M_R\|$, integrating out the heavy singlets ν_R generates the unique dimension-5 operator [568]

$$\mathcal{L}_{\mathrm{eff}}^{(5)} \;=\; -\frac{1}{2}\left(L_L\tilde{\Theta}\right)^T C_5 \left(L_L\tilde{\Theta}\right) \;+\; \mathrm{h.c.}, \qquad C_5 \;=\; Y_\nu\, M_R^{-1}\, Y_\nu^T. \tag{12.186}$$

Symmetry note. Since $M_R = M_R^T$ by Definition 12.139, its inverse is symmetric and therefore $C_5 = C_5^T$. Consequently $M_\nu = -\frac{v^2}{2} C_5$ is complex symmetric, so a Takagi factor U_ν exists with $U_\nu^T M_\nu U_\nu = \mathrm{diag}(m_i)$; this underlies the leptonic mixing definition in Definition 12.141.[3]

Work in the chiral basis (ν_L, ν_R^c), with $M_D = \frac{v}{\sqrt{2}} Y_\nu$ (Definition 12.139). The quadratic form reads

$$\mathcal{L}_{\mathrm{mass}} \;=\; -\frac{1}{2}\begin{pmatrix}\overline{\nu_L} & \overline{\nu_R^c}\end{pmatrix}\begin{pmatrix}0 & M_D\\ M_D^T & M_R\end{pmatrix}\begin{pmatrix}\nu_L^c\\ \nu_R\end{pmatrix} \;+\; \mathrm{h.c.} \tag{12.187}$$

Assuming the hierarchical regime $\|M_R\| \gg \|M_D\|$ (technically natural since ν_R is a gauge singlet; see reviews in [569]), the Schur complement yields the *effective* light-neutrino mass matrix

$$M_\nu \;=\; -\, M_D\, M_R^{-1}\, M_D^T, \qquad M_N \;\simeq\; M_R, \tag{12.188}$$

the classic Type-I seesaw relation [129, 546, 547, 130], where M_ν acts on ν_L and M_N denotes the heavy singlet block. Equivalently, the heavy–light mixing is controlled by

$$\Theta \;=\; M_D\, M_R^{-1}, \tag{12.189}$$

[3] Overall signs in (12.186) and below are convention-dependent and physically inessential after rephasings that render masses positive. The operator is allowed by the SM gauge representations and follows the selection rules of Proposition 12.90.

and the block-off-diagonal entries are removed by a (nearly) unitary transformation

$$\mathcal{U} \simeq \exp\begin{pmatrix} 0 & \Theta \\ -\Theta^\dagger & 0 \end{pmatrix} \quad \text{up to } \mathcal{O}(\Theta^3)$$

(cf. Proposition 12.140). In particular, inserting $C_5 = Y_\nu M_R^{-1} Y_\nu^T$ into $M_\nu = -\frac{v^2}{2} C_5$ reproduces (12.188).

Since M_ν is complex symmetric, choose a Takagi factor U_ν with $U_\nu^T M_\nu U_\nu = \mathrm{diag}(m_i)$ and define the leptonic mixing matrix

$$U_{\mathrm{PMNS}} = U_{eL}^\dagger U_\nu, \tag{12.190}$$

in agreement with Definitions 12.117 and 12.141 and the inevitability argument of Section 12.5.6; this specializes the bi-unitary flavor discussion to the Majorana/seesaw case and matches the original PMNS construction [545, 544].

Proposition 12.140 (Seesaw diagonalization and active–sterile mixing)**.** *Let $M_D = \frac{v}{\sqrt{2}} Y_\nu$ and $M_R = M_R^T$ as in Definition 12.139. Define $\Theta := M_D\, M_R^{-1}$ and assume $\|\Theta\| \ll 1$. Then there exists a unitary transformation*

$$\mathcal{U} = \exp S, \qquad S := \begin{pmatrix} 0 & \Theta \\ -\Theta^\dagger & 0 \end{pmatrix},$$

such that the Majorana mass matrix $\mathcal{M} = \begin{pmatrix} 0 & M_D \\ M_D^T & M_R \end{pmatrix}$ is block–diagonalized up to $\mathcal{O}(\Theta^3)$ under the Majorana congruence $\mathcal{M}' = \mathcal{U}^T \mathcal{M} \mathcal{U}$:

$$\mathcal{M}' = \begin{pmatrix} -M_D M_R^{-1} M_D^T & 0 \\ 0 & M_R \end{pmatrix} + \mathcal{O}(\Theta^3). \tag{12.191}$$

Hence the light block equals the Type–I seesaw mass $M_\nu = -M_D M_R^{-1} M_D^T$ and the heavy block equals M_R to this order, with active–sterile mixing angles of order Θ. This matches eq. (12.188) and is consistent with the EFT limit $\mathcal{L}^{(5)}$ in eq. (12.186).

Proof. **(i) Unitarity and the correct transformation law.)** By construction $S^\dagger = -S$, hence $\mathcal{U} = \exp S$ is unitary. For a symmetric (Majorana) mass term the field redefinition $n \mapsto \mathcal{U}\, n$ induces the *congruence* $\mathcal{M} \mapsto \mathcal{M}' = \mathcal{U}^T \mathcal{M} \mathcal{U}$ (rather than a similarity), as used in the Takagi framework for symmetric matrices (cf. the Takagi factor in Definition 12.141 and the discussion below eq. (12.186)).

(ii) Power counting and BCH expansion.) Write the original mass matrix in 2×2 block form,

$$\mathcal{M} = \begin{pmatrix} M_{11} & M_{12} \\ M_{21} & M_{22} \end{pmatrix} = \begin{pmatrix} 0 & M_D \\ M_D^T & M_R \end{pmatrix},$$

and set $\epsilon := \|\Theta\| \ll 1$. Expanding with the Baker–Campbell–Hausdorff (BCH) series for a congruence (transpose on the left),

$$\begin{aligned} \mathcal{M}' &= (\mathbb{1} + S^T + \tfrac{1}{2} S^{T\,2} + \cdots)\, \mathcal{M}\, (\mathbb{1} + S + \tfrac{1}{2} S^2 + \cdots) \\ &= \mathcal{M} + \underbrace{(S^T \mathcal{M} + \mathcal{M} S)}_{\mathcal{O}(\epsilon)} + \underbrace{\left(\tfrac{1}{2} S^{T\,2} \mathcal{M} + \tfrac{1}{2} \mathcal{M} S^2 + S^T \mathcal{M} S\right)}_{\mathcal{O}(\epsilon^2)} + \mathcal{O}(\epsilon^3). \end{aligned} \tag{12.192}$$

Since $S = \mathcal{O}(\epsilon)$, the first line fixes the $\mathcal{O}(\epsilon)$ cancellation of the off–diagonal blocks, and the second line controls the residual $\mathcal{O}(\epsilon^2)$ terms.

(iii) Choice of S solves the Sylvester equation at $\mathcal{O}(\epsilon)$.) Compute the upper–right block of $\mathcal{M}'$ at $\mathcal{O}(\epsilon)$:

$$(\mathcal{M}')^{(1)}_{12} = M_{12} + (S^T\mathcal{M})_{12} + (\mathcal{M}S)_{12} = M_D - \Theta M_R + 0 = 0,$$

using $\Theta = M_D M_R^{-1}$, $S^T = \left(\begin{smallmatrix} 0 & -\Theta^T \\ \Theta^* & 0 \end{smallmatrix}\right)$, and $M_{11} = 0$. Similarly, the lower–left block cancels:

$$(\mathcal{M}')^{(1)}_{21} = M_D^T - M_R\Theta^T = M_D^T - M_R(M_D M_R^{-1})^T = 0,$$

since $M_R = M_R^T$. Thus the $\mathcal{O}(\epsilon)$ off–diagonal entries vanish. This is the standard Sylvester equation $M_{12} + M_{11}\Theta - \Theta M_{22} = 0$ with $M_{11} = 0$, solved by $\Theta = M_{12}M_{22}^{-1}$.

(iv) Vanishing of off–diagonals at $\mathcal{O}(\epsilon^2)$.) Using (12.192) and the block structure of S,

$$S^2 \;=\; \left(\begin{smallmatrix} -\Theta\Theta^\dagger & 0 \\ 0 & -\Theta^\dagger\Theta \end{smallmatrix}\right), \qquad S^{T\,2} \;=\; \left(\begin{smallmatrix} -\Theta\Theta^* & 0 \\ 0 & -\Theta^T\Theta^\dagger \end{smallmatrix}\right).$$

A straightforward (but tedious) block multiplication shows that all contributions to $(\mathcal{M}')^{(2)}_{12}$ are proportional to the already–imposed $\mathcal{O}(\epsilon)$ Sylvester condition and its transpose; equivalently,

$$(\mathcal{M}')^{(2)}_{12} = \tfrac{1}{2}\Big(M_{11}\Theta - \Theta M_{22}\Big)\Theta + \tfrac{1}{2}\Theta\Big(M_{22}\Theta^T - \Theta^T M_{11}^T\Big)^T + \cdots \;=\; 0,$$

because $M_{11} = 0$ and $M_{22} = M_R$, together with $M_D - \Theta M_R = 0$ and its transpose, annihilate each term. Hence the off–diagonal blocks vanish through $\mathcal{O}(\epsilon^2)$, so the first nonzero remainder is $\mathcal{O}(\epsilon^3)$.

(v) Diagonal blocks to $\mathcal{O}(\epsilon^2)$.) For the light block,

$$(\mathcal{M}')_{11} = M_{11} + (S^T\mathcal{M} + \mathcal{M}S)_{11} + \Big(\tfrac{1}{2}S^{T\,2}\mathcal{M} + \tfrac{1}{2}\mathcal{M}S^2 + S^T\mathcal{M}S\Big)_{11} + \mathcal{O}(\epsilon^3).$$

Using $M_{11} = 0$ and the relations above, one finds

$$(\mathcal{M}')_{11} = -\,M_D M_R^{-1} M_D^T \;+\; \mathcal{O}(\Theta^3),$$

while for the heavy block $(\mathcal{M}')_{22} = M_R + \mathcal{O}(\Theta^2 M_R)$, so to the stated order M_R is unchanged. Together these give (12.191), i.e. the seesaw formula $M_\nu = -M_D M_R^{-1} M_D^T$ in eq. (12.188) and heavy masses $M_N \simeq M_R$.

(vi) Mixing angles and consistency with the IQR structures.) The generator S rotates (ν_L, ν_R^c) by angles of order Θ between the active and sterile subspaces; hence the active–sterile mixing is $\sim \Theta$ at leading order, as stated. Since $\mathcal{U}$ acts only in family/flavor space, it commutes with the spin/lattice projectors and with the covariant transport on the family bundle at a fixed point (Definition 12.110 and eq. (12.128)); global relative phases are then supplied by family holonomy as in Proposition 12.115. Finally, because the transformation is unitary and local in flavor, it preserves the kinetic form and the diagnostic realized reference configuration/Nielsen–KL hygiene of poles and residues on certified plateaux (Proposition 12.88), ensuring that the light/heavy spectra read from the two–point functions agree with the block–diagonal result. □

12.7.5 PMNS Mixing from Misalignment

Let M_e be the charged–lepton mass matrix from the Yukawa sector (Section 12.4.6). Choose unitary matrices U_{eL}, U_{eR} such that

$$U_{eL}^{\dagger}\, M_e\, U_{eR} \;=\; \mathrm{diag}(m_e, m_\mu, m_\tau). \tag{12.193}$$

For Majorana neutrinos, the light mass matrix M_ν obtained from the seesaw/EFT construction ($M_\nu = -\frac{v^2}{2} C_5 = -M_D M_R^{-1} M_D^{T}$, cf. Section 12.7.4 and eqs. (12.186) and (12.188)) is complex symmetric. Hence there exists a *Takagi* factor U_ν with

$$U_\nu^{T}\, M_\nu\, U_\nu \;=\; \mathrm{diag}(m_1, m_2, m_3), \qquad m_i \geq 0, \tag{12.194}$$

as used also in eq. (12.190); see standard discussions in [82, 91].

Definition 12.141 (Leptonic mixing (PMNS) matrix)**.** The leptonic charged current couples flavor and mass bases through

$$U_{\mathrm{PMNS}} \;:=\; U_{eL}^{\dagger}\, U_\nu, \tag{12.195}$$

entering the weak current $J_\mu^{(+)} = \overline{e_L}\,\gamma_\mu\, U_{\mathrm{PMNS}}\, \nu_L + \mathrm{h.c.}$ (Section 12.5.5 and eq. (12.139)). This is the Pontecorvo–Maki–Nakagawa–Sakata construction [545, 544]; see also [82, 91]. For purely Dirac neutrinos one recovers the bi–unitary form of Definition 12.117 with $U_{\mathrm{PMNS}} = U_{eL}^{\dagger} V_{\nu L}$ [82, 91].

Proposition 12.142 (Standard parameterization and rephasing invariants)**.** *Up to unphysical charged–lepton field rephasings, U_{PMNS} can be written as*

$$U_{\mathrm{PMNS}} \;=\; \underbrace{\begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix}}_{R_{23}} \underbrace{\begin{pmatrix} c_{13} & 0 & s_{13} e^{-i\delta} \\ 0 & 1 & 0 \\ -s_{13} e^{i\delta} & 0 & c_{13} \end{pmatrix}}_{R_{13}(\delta)} \underbrace{\begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix}}_{R_{12}} \mathrm{diag}\!\left(e^{\frac{i\alpha_1}{2}},\, e^{\frac{i\alpha_2}{2}},\, 1\right), \tag{12.196}$$

with $s_{ij} = \sin\theta_{ij}$, $c_{ij} = \cos\theta_{ij}$, Dirac phase δ, and (if neutrinos are Majorana) two additional phases $\alpha_{1,2}$ that drop from oscillation probabilities [82, 91]. The rephasing–invariant leptonic Jarlskog measure is

$$J_{\mathrm{CP}} \;=\; \mathrm{Im}\!\left[U_{e1} U_{\mu 2} U_{e2}^{*} U_{\mu 1}^{*}\right] \;=\; s_{12} s_{23} s_{13}\, c_{12} c_{23} c_{13}^{2}\, \sin\delta, \tag{12.197}$$

in exact analogy with the quark–sector invariant (Definition 12.117); see [82, 91].

Idea. Counting physical parameters follows from removing phase redundancies in U_{eL} and in the Takagi factor U_ν: for three generations one obtains 3 angles, one Dirac phase, and (for Majorana masses) two additional phases that cannot be rephased away because ν and ν^c are not independent (Section 12.7.4 and eq. (12.180)); see [82, 91]. The Euler–like factorization (12.196) is standard and unique up to convention choices for angle ranges; (12.197) is the unique independent rephasing invariant of order four in U–entries, by the same argument used for the CKM sector (Section 12.5.5); cf. [82, 91]. □

Proof. **Step 1 (Degrees of freedom of a 3×3 unitary and a canonical factorization).** A general $U\in U(3)$ has $\dim_{\mathbb{R}} U(3)=9$ real parameters, usually organized as 3 Euler angles and 6 phases. By successive Givens rotations (or the Euler-like construction), any U can be factorized as

$$U \;=\; D_\ell\, R_{23}\, R_{13}(\delta)\, R_{12}\, D_\nu, \tag{12.198}$$

where R_{ij} are real rotations in the (i,j) planes with angles $(\theta_{12},\theta_{13},\theta_{23})$ and a single complex phase in $R_{13}(\delta)$, while $D_\ell=\mathrm{diag}(e^{i\phi_e},e^{i\phi_\mu},e^{i\phi_\tau})$ and $D_\nu=\mathrm{diag}(e^{i\chi_1},e^{i\chi_2},e^{i\chi_3})$ collect the remaining $6-1=5$ phases [82, 91]. Removing redundant overall phases in (12.198) reproduces the standard form (12.196) once unphysical rephasings (below) are modded out.

Step 2 (Equivalence relations from field rephasings). The charged current couples as $J_\mu^{(+)}=\overline{e_L}\gamma_\mu U_{\mathrm{PMNS}}\nu_L+\mathrm{h.c.}$ (Definition 12.141 and Section 12.5.5). Under flavor-basis rephasings

$$e_L\to D_\ell\, e_L, \qquad \nu_L\to D_\nu\,\nu_L,$$

with $D_{\ell,\nu}$ diagonal unitary matrices, the mixing matrix transforms as $U_{\mathrm{PMNS}}\to D_\ell^\dagger\, U_{\mathrm{PMNS}}\, D_\nu$. Thus physical information is the equivalence class

$$[U_{\mathrm{PMNS}}] \;\in\; \begin{cases} U(3)\Big/\Big((U(1))^5\Big), & \text{Dirac case},\\ U(3)\Big/\Big((U(1))^3\Big), & \text{Majorana case}, \end{cases} \tag{12.199}$$

because in the Dirac case both D_ℓ and D_ν are allowed, but one common overall phase is irrelevant (hence 5 removable phases), whereas in the Majorana case D_ν is *not* allowed: rephasing ν_L would spoil the Majorana condition (or the Majorana mass term), so only the three charged-lepton phases can be removed [82, 91].

Step 3 (Counting physical parameters). From (12.199):

$$\dim_{\mathbb{R}} U(3)=9 \quad\Rightarrow\quad \begin{cases} 9-5=4\ (\text{Dirac})=3\ \text{angles}+1\ \text{phase},\\ 9-3=6\ (\text{Majorana})=3\ \text{angles}+3\ \text{phases}. \end{cases}$$

Equivalently, for N generations one finds $N(N-1)/2$ angles; the number of phases is $(N-1)(N-2)/2$ (Dirac) and $N(N-1)/2$ (Majorana) [82, 91]. For $N=3$ this gives 3 angles and, respectively, 1 Dirac phase or 1 Dirac $+2$ Majorana phases. The Majorana phases cannot be absorbed into fields because ν_L and ν_L^c are not independent when the mass term is Majorana (Section 12.7.4 and eq. (12.180)); in oscillations they drop out of probabilities, but they remain in lepton-number–violating amplitudes.

Step 4 (Standard parameterization). Using (12.198) and the rephasing freedom in Step 2 to eliminate D_ℓ (and D_ν in the Dirac case), one arrives at the canonical form (12.196) with three Euler angles $(\theta_{12},\theta_{13},\theta_{23})$, a Dirac phase δ embedded in $R_{13}(\delta)$, and—for Majorana neutrinos—an extra diagonal $\mathrm{diag}(e^{i\alpha_1/2},e^{i\alpha_2/2},1)$ multiplying on the right [82, 91]. Different choices of angle ranges and phase conventions produce equivalent parametrizations related by the rephasings of Step 2.

Step 5 (Rephasing invariance and the Jarlskog measure). Under $U\to D_\ell^\dagger U D_\nu$, the quartet $Q_{\alpha\beta ij}=U_{\alpha i}U_{\beta j}U_{\alpha j}^*U_{\beta i}^*$ is invariant: the diagonal phases cancel pairwise. Its imaginary part

$$J_{\mathrm{CP}}=\mathrm{Im}\, Q_{e\mu\,12}=\mathrm{Im}[U_{e1}U_{\mu 2}U_{e2}^*U_{\mu 1}^*]=s_{12}s_{23}s_{13}\,c_{12}c_{23}c_{13}^2\,\sin\delta$$

(eq. (12.197)) is therefore a rephasing–invariant, CP–odd measure. For three generations, all independent CP–odd, rephasing–invariant polynomials constructed from U at fourth order are equal to $\pm J_{\mathrm{CP}}$; higher-order invariants reduce to J_{CP} times rephasing–even combinations of moduli (the situation is completely analogous to the CKM sector, Section 12.5.5) [82, 91]. In oscillation probabilities, CP violation appears only through terms proportional to J_{CP}.

Step 6 (Consistency with Majorana/Dirac masses). In the Dirac case M_ν is general complex and admits a biunitary diagonalization, $V_{\nu L}^\dagger M_\nu V_{\nu R} = \mathrm{diag}(m_1, m_2, m_3)$, giving $U_{\mathrm{PMNS}} = U_{eL}^\dagger V_{\nu L}$ and removing two additional phases as in Step 3. In the Majorana (seesaw) case M_ν is complex symmetric, so the Takagi factorization $U_\nu^T M_\nu U_\nu = \mathrm{diag}(m_1, m_2, m_3)$ fixes U_ν up to right multiplication by *real* diagonal phase matrices $\mathrm{diag}(\pm 1, \pm 1, \pm 1)$ only, which cannot remove continuous phases; hence two extra (Majorana) phases remain in (12.196) [82, 91]. This completes the derivation of the parameter count and the standard form. □

Remark 12.143 (Oscillations, family transport, and holonomy refinement)**.** Neutrino oscillation probabilities depend on the mass–squared splittings and on the rephasing–invariant combinations of U_{PMNS}, consistent with the Pontecorvo–Maki–Nakagawa–Sakata paradigm (Section 12.5.5) [545, 544]. In the family–lifted formulation (Definition 12.110), relative transport in the flat family bundle modifies the comparison of local diagonal frames by a Wilson line of $\mathcal{A}^{(F)}$; the observable mixing at a point x can be written as in Proposition 12.115 and eq. (12.134) with $U_{\mathrm{PMNS}}(x) = U_{eL}^\dagger(x)\, \mathcal{P} \exp \int_\gamma i\mathcal{A}^{(F)}\, U_\nu(x)$, where γ interpolates the preparation and detection loci.

Remark 12.144 (Dirac limit and consistency with the seesaw)**.** In the Dirac case (no Majorana term), M_ν is general complex (not symmetric) and is diagonalized bi–unitarily; then $U_{\mathrm{PMNS}} = U_{eL}^\dagger V_{\nu L}$ as in Definition 12.117, with Majorana phases absent [82, 91]. In the Type–I seesaw, $M_\nu = -M_D M_R^{-1} M_D^T$ is complex symmetric, leading to the Takagi form (12.180) and to (12.195), consistent with Section 12.7.4 and eq. (12.190) and the original seesaw literature [129, 547].

12.7.6 Constraints from the Tantrakyma Potential and Family Fibre

Within the IQR architecture, the neutrino–sector couplings (Y_ν, M_R) are not free inputs: they are constrained by (i) the symmetry and vacuum structure of the Tantrakyma potential $\mathcal{V}_\Theta$ (which fixes the electroweak scale and admissible endomorphisms), and (ii) the flat, unitary *family fibre* $(\mathcal{F}, \mathcal{A}^{(F)})$ that transports generation indices (Definitions 9.335 and 12.106). In particular, the EWSB background $\bar{\Theta}$ sets the Dirac scale $M_D = \frac{v}{\sqrt{2}} Y_\nu$ (Definition 12.139), while higher-scale neutral moduli in $\mathcal{V}_\Theta$ feed into the gauge–singlet Majorana sector $M_R \in \mathrm{End}(\mathcal{F})$, naturally realizing the hierarchical regime $\|M_R\| \gg \|M_D\|$ required by the Type-I seesaw (Section 12.7.4 and eq. (12.188)). Family transport enters only through the total covariant derivative (Definition 12.110 and eq. (12.128)), so observable misalignment (PMNS mixing) is the relative family holonomy between the charged-lepton and neutrino diagonal frames (Proposition 12.115, eq. (12.134), and Definition 12.136).

Definition 12.145 (Flavor–sector holosymmetric functional)**.** Let $\mathbf{U}^{(\ell)}$ denote the leptonic evolution generated on the diagnostic realized reference configuration by the Tantrakyma-induced relational Hamiltonian with parameters (Y_ν, M_R) (Definitions 6.54 and 12.114).

Define the *flavor–sector self-coherence*

$$\mathcal{C}_\nu \;:=\; \Big|\langle\Psi_0|(\mathbf{U}^{(\ell)})^\dagger\mathbf{U}^{(\ell)}|\Psi_0\rangle\Big|$$

as the Stage-I unitarity meter restricted to the leptonic channel (Theorem 12.42). For two viable pairs (Y_ν, M_R) and (Y'_ν, M'_R), define the *asymmetric relational visibility*

$$\mathcal{V}^{(\nu)}_{\text{asym}} \;:=\; \Big|p_0^{(\text{asym})} - p_1^{(\text{asym})}\Big|$$

as measured on the dual-register holosymmetric interferometer (Section 3.1.5 and Definition 7.39), with the usual witness/TLO readout (Definitions 8.4 and 8.6 and Proposition 10.22).

Corollary 12.146 (Seesaw-compatible HSP filter in the lepton sector)**.** *Among all* (Y_ν, M_R) *consistent with the Tantrakyma vacuum (Section 12.4.5) and BRST physicality, the realized neutrino data are those for which:*

(i) $\mathcal{C}_\nu = 1$ *(Stage I unitarity/anomaly viability; Theorem 12.42 and Section 12.5.4), and*

(ii) $\mathcal{V}^{(\nu)}_{\text{asym}}$ *can be tomographically nulled against any inequivalent viable pair (Stage II relational economy/orthogonality; Proposition 12.10 and Corollary 12.48).*

In the mass basis, item (ii) enforces orthogonality of the induced actions and is equivalent to vanishing holosymmetric overlap, thereby selecting a unique PMNS invariant (angles and phases) inside the admissible (Y_ν, M_R) *manifold (Sections 12.7.2 and 12.7.5).*

Phenomenological remarks. Both normal and inverted orderings are accommodated via the sign of Δm^2_{31}, while the smallness of m_ν follows from the heavy scale in M_R through the seesaw relation $M_\nu = -M_D M_R^{-1} M_D^T$ (eq. (12.188)). Observable CP violation arises through the Dirac phase δ (oscillations) and, for Majorana neutrinos, two additional phases (α_1, α_2) relevant to $0\nu\beta\beta$, with both sets traceable to (i) phases in (Y_ν, M_R) generated by the Tantrakyma sector and (ii) family holonomy on $(\mathcal{F}, \mathcal{A}^{(F)})$ (Definition 12.106 and Remark 12.116). For the historical genesis of oscillations and the seesaw mechanism, see [544, 546, 547].

12.7.7 Neutrino Mass and the Majorana Seesaw Mechanism

We now turn to the explicit mass construction in the lepton sector of the Tantrakyma field. Left- and right-handed neutrinos, ν_L and ν_R, arise as spin-$\frac{1}{2}$ kymon excitations in the lepton representation content (Section 12.4.6 and Definition 12.82). Gauge invariance allows a Dirac mass from Yukawa coupling to the EWSB background $\bar{\Theta} = \langle H\rangle$ as well as a gauge–singlet Majorana mass for ν_R (Definitions 12.135 and 12.139; see [82, 91]).

Dirac and Majorana terms. With $L_L = (\nu_L, e_L)^\top$ and $\tilde{H} = i\sigma_2 H^*$, the neutrino Dirac mass is generated by

$$\mathcal{L}_{\text{Dirac}} \;=\; -\,\overline{L_L}\,\tilde{H}\,Y_\nu\,\nu_R \;+\; \text{h.c.}, \qquad M_D \;=\; \frac{v}{\sqrt{2}}\,Y_\nu \quad \Big(\langle H\rangle = \bar{\Theta} = \tfrac{1}{\sqrt{2}}\begin{pmatrix}0\\ v\end{pmatrix}\Big),$$

while the gauge–invariant Majorana term for $\nu_R \sim (1,1)_0$ is

$$\mathcal{L}_{\text{Maj}} \;=\; -\frac{1}{2}\,\overline{\nu_R^{c}}\,M_R\,\nu_R \;+\; \text{h.c.},$$

with $M_R = M_R^T$ a complex symmetric matrix in family space (Definition 12.139); standard accounts of Dirac/Majorana neutrino masses and their gauge structure include [82, 91] and the original seesaw proposals [129, 547]. Both Y_ν and M_R live as endomorphisms on the family fibre $\mathcal{F}$ (Definitions 12.110 and 12.114).

Six–component Majorana form. Define the Nambu–Gorkov spinor

$$\Psi \;=\; \begin{pmatrix} \nu_L \\ \nu_R^{c} \end{pmatrix}, \qquad \overline{\Psi} \;=\; \Big(\overline{\nu_L},\, \overline{\nu_R^{c}}\Big),$$

so that the full mass Lagrangian reads

$$\mathcal{L}_\nu^{\text{mass}} \;=\; -\frac{1}{2}\,\overline{\Psi}\,\mathcal{M}_\nu\,\Psi \;+\; \text{h.c.}, \qquad \mathcal{M}_\nu \;=\; \begin{pmatrix} 0 & M_D \\ M_D^T & M_R \end{pmatrix}. \tag{12.200}$$

This is the standard Dirac–Majorana structure appearing in the Type-I seesaw (Section 12.7.4); cf. [129, 547, 82, 91].

Hierarchical regime and effective spectrum. In the technically natural regime $\|M_R\| \gg \|M_D\|$ (right–handed singlets are unconstrained by G_{SM}), block diagonalization or, equivalently, the Schur complement gives the light and heavy eigenstructures:

$$M_\nu \;=\; -\,M_D\,M_R^{-1}\,M_D^T, \qquad M_N \;\simeq\; M_R, \tag{12.201}$$

$$\Theta \;=\; M_D\,M_R^{-1}, \tag{12.202}$$

with Θ controlling the active–sterile mixing (Section 12.7.4 and eqs. (12.188) and (12.189)); see [129, 547, 91, 82]. A constructive unitary that cancels the off–diagonal blocks up to $\mathcal{O}(\Theta^3)$ is provided by $\mathcal{U} = \exp\left(\begin{smallmatrix} 0 & \Theta \\ -\Theta^\dagger & 0 \end{smallmatrix}\right)$ (Proposition 12.140; cf. [91]). Equivalently, integrating out ν_R yields Weinberg's d=5 operator with $C_5 = Y_\nu M_R^{-1} Y_\nu^T$ and $M_\nu = -\frac{v^2}{2} C_5$ (eq. (12.186)); see [91, 82].

Proposition 12.147 (Emergence of the seesaw from Tantrakyma dynamics)**.** *Let $M_D = \frac{v}{\sqrt{2}} Y_\nu$ arise from the Yukawa coupling to the Higgs–kymon background $\bar{\Theta}$ and let M_R be supplied by a neutral, quadratic term permitted by the Tantrakyma potential $\mathcal{V}_\Theta$ (Definitions 9.335 and 12.139). Then, on the certified plateau, the neutral fermion sector of the relational Hamiltonian $\mathbb{H}^{(\Theta)}$ possesses a light spectrum suppressed by the seesaw ratio:*

$$\boxed{\text{Spec}_{\textit{light}}(\mathcal{M}_\nu) \;=\; \mathcal{O}\Big(M_D\,M_R^{-1}\,M_D^T\Big) \;\sim\; \mathcal{O}\left(\frac{M_D^2}{M_R}\right),} \tag{12.203}$$

with heavy eigenvalues $\sim \text{Spec}(M_R)$, gauge–parameter–independent poles and KL–positive residues (Proposition 12.88).

Proof. Insert the EWSB background (Section 12.4.5) into $\mathcal{L}_Y^{(\nu)}$ to obtain $M_D = \frac{v}{\sqrt{2}} Y_\nu$ (Definition 12.139). Augment with the gauge–singlet Majorana term allowed by $\mathcal{V}_\Theta$ to form $\mathcal{M}_\nu$ in (12.200). In the hierarchy $\|M_R\| \gg \|M_D\|$, either (i) integrate out ν_R to generate the Weinberg operator (eq. (12.186)), or (ii) apply the unitary block diagonalization of Proposition 12.140; both routes give the light block $M_\nu = -M_D M_R^{-1} M_D^T$ and heavy block M_R up to $\mathcal{O}(\Theta^3)$. Diagnostic realized reference configuration certification (Proposition 12.88) ensures that the resulting propagator poles and residues for light/heavy modes are gauge–parameter independent and satisfy KL positivity, so the spectrum statement in (12.203) is physically meaningful. The endomorphism character $Y_\nu, M_R \in \mathrm{End}(\mathcal{F})$ (Definition 12.114) and family transport (Definition 12.110) do not alter the local block structure, but they do control the relative holonomy that appears later in PMNS mixing (Section 12.7.5 and Definition 12.136). □

PMNS mixing and phases. Since M_ν is complex symmetric, a Takagi factor U_ν exists with $U_\nu^T M_\nu U_\nu = \mathrm{diag}(m_1, m_2, m_3)$. Together with the charged–lepton diagonalizer U_{eL}, the leptonic mixing matrix is $U_{\mathrm{PMNS}} = U_{eL}^\dagger U_\nu$, refined by family holonomy as in Proposition 12.115 and Definition 12.136. The Dirac phase and (if Majorana) two additional phases encode leptonic CP violation (Section 12.7.5 and Proposition 12.142).

Lepton Number Violation and Tantrakyma Condensation

Let Φ_ν be a complex, *SM–singlet* Tantrakyma scalar,

$$\Phi_\nu \sim (1,1)_0 \quad \text{under } G_{\mathrm{SM}},$$

assigned a global lepton number $L(\Phi_\nu) = -2$. Right–handed neutrinos ν_R carry $L(\nu_R) = +1$ and are also SM singlets $\nu_R \sim (1,1)_0$ (Definition 12.82). The Yukawa sector in the neutral Tantrakyma channel extends Definitions 12.135 and 12.139 by

$$\mathcal{L}_{\mathrm{singlet}}^{(\nu)} \;=\; -\frac{1}{2}\,\overline{\nu_R^c}\, y^{(\nu)}\, \Phi_\nu\, \nu_R \;+\; \mathrm{h.c.}, \qquad y^{(\nu)} = y^{(\nu)T} \in \mathrm{End}(\mathcal{F}), \tag{12.204}$$

which is gauge–invariant and preserves global $U(1)_L$ *before* symmetry breaking.[4]

The scalar potential in the Tantrakyma sector is enlarged to

$$\mathcal{V}_\Theta \;\to\; \mathcal{V}_\Theta(\Theta) \;+\; \mathcal{V}_\nu(\Phi_\nu) \;+\; \lambda_{\Theta\nu}\,(\Theta^\dagger\Theta)\,(\Phi_\nu^\dagger\Phi_\nu), \qquad \mathcal{V}_\nu(\Phi_\nu) = m_\nu^2\,\Phi_\nu^\dagger\Phi_\nu + \frac{\lambda_\nu}{2}\,(\Phi_\nu^\dagger\Phi_\nu)^2, \tag{12.205}$$

with $\mathcal{V}_\Theta$ the Tantrakyma/Higgs potential from Definition 9.335. If $m_\nu^2 < 0$ and $\lambda_\nu > 0$, the neutral singlet condenses,

$$\langle \Phi_\nu \rangle \;=\; \frac{v_\nu}{\sqrt{2}} \neq 0, \tag{12.206}$$

spontaneously breaking global $U(1)_L$.[5]

[4] All family–space endomorphisms (here $y^{(\nu)}$) act on the rank–3 family fibre $\mathcal{F}$ (Definitions 12.106 and 12.114).

[5] Spontaneous breaking of a global $U(1)_L$ produces a Nambu–Goldstone (Majoron). In the present framework this can be lifted or decoupled by gauging an anomaly–free $U(1)_{B-L}$ or by small explicit breaking, remaining compatible with Stage I/II of HSP (Section 12.5.4 and Theorem 12.42).

Induced Majorana mass and seesaw scaling. Using (12.204)–(12.206), the RH–Majorana mass matrix is generated dynamically,

$$M_R \;=\; \frac{v_\nu}{\sqrt{2}}\, y^{(\nu)} \in \mathrm{End}(\mathcal{F}), \tag{12.207}$$

and combines with the Dirac mass $M_D = \frac{v}{\sqrt{2}} Y_\nu$ from the electroweak Tantrakyma background $\bar{\Theta}$ (Definition 12.139) to produce, in the hierarchical regime $\|M_R\| \gg \|M_D\|$, the light Majorana matrix (Section 12.7.4)

$$M_\nu \;=\; -\,M_D\, M_R^{-1}\, M_D^T \;=\; -\,\frac{v^2}{2}\, Y_\nu\, (y^{(\nu)} v_\nu/\sqrt{2})^{-1}\, Y_\nu^T, \tag{12.208}$$

which is complex symmetric and Takagi–diagonalizable (eq. (12.180) and Definition 12.141).

At the level of parametric scaling one finds

$$\|M_\nu\| \;\sim\; \frac{(v\,\|Y_\nu\|)^2}{\|y^{(\nu)}\|\, v_\nu}\,, \tag{12.209}$$

matching the seesaw suppression in eq. (12.188) and its EFT limit (eq. (12.186)).

Proposition 12.148 (Spontaneous $U(1)_L$ breaking and seesaw feed–down)**.** *If Φ_ν condenses as in (12.206), global lepton number is spontaneously broken and a Majorana mass M_R is induced per (12.207). Inserting M_R into the Type–I seesaw (Section 12.7.4) yields the light spectrum suppressed as in (12.208)–(12.209). The extended potential (12.205) remains compatible with the IQR diagnostic realized reference configuration and HSP: it introduces no new SM gauge quanta (Section 12.4.4), preserves anomaly viability (Stage I; Section 12.5.4 and eq. (12.46)), and respects the family–bundle architecture ($y^{(\nu)}, Y_\nu \in \mathrm{End}(\mathcal{F})$; Definition 12.114).*

Proof. (i) Spontaneous $U(1)_L$ breaking and induced M_R. Assign global lepton number $L(\nu_R) = +1$ and $L(\Phi_\nu) = -2$. Under a global $U(1)_L$ rotation by angle α,

$$\nu_R \;\mapsto\; e^{+i\alpha}\nu_R, \qquad \nu_R^c \;\mapsto\; e^{-i\alpha}\nu_R^c, \qquad \Phi_\nu \;\mapsto\; e^{-2i\alpha}\Phi_\nu,$$

so the neutral–singlet Yukawa interaction (12.204) is invariant: $\overline{\nu_R^c}\, y^{(\nu)}\, \Phi_\nu\, \nu_R \mapsto \overline{\nu_R^c}\, y^{(\nu)}\, \Phi_\nu\, \nu_R$. The extended Tantrakyma potential (12.205) has a minimum at $\langle \Phi_\nu \rangle = \frac{v_\nu}{\sqrt{2}} \neq 0$ when $m_\nu^2 < 0$ and $\lambda_\nu > 0$. Replacing Φ_ν by its VEV in (12.204) yields the RH–Majorana mass matrix

$$M_R \;=\; \frac{v_\nu}{\sqrt{2}}\, y^{(\nu)} \quad \text{in } \mathrm{End}(\mathcal{F}), \tag{12.210}$$

which is exactly (12.207). Since the order parameter $\langle \Phi_\nu \rangle \neq 0$ carries $L = -2$, the global $U(1)_L$ is *spontaneously* broken.[6]

(ii) Seesaw feed–down to the light spectrum. Electroweak symmetry breaking in the Tantrakyma doublet sector produces the Dirac mass $M_D = \frac{v}{\sqrt{2}} Y_\nu$ (Definition 12.139). In the chiral basis (ν_L, ν_R^c) the full neutral mass matrix is $\left(\begin{smallmatrix} 0 & M_D \\ M_D^T & M_R \end{smallmatrix}\right)$. In the hierarchical regime

[6] In a purely global realization this produces a (Majoron) Goldstone mode; it can be lifted by gauging an anomaly–free $U(1)_{B-L}$ or by small explicit breaking, consistent with Section 12.5.4 and Theorem 12.42.

$\|M_R\| \gg \|M_D\|$, Schur complement (or equivalently the perturbative block–Takagi rotation in Proposition 12.140) gives the light block

$$M_\nu \;=\; -\,M_D\,M_R^{-1}\,M_D^T, \tag{12.211}$$

and heavy block $M_N \simeq M_R$, reproducing (12.188). Inserting (12.210) yields

$$M_\nu \;=\; -\,\frac{v^2}{2}\,Y_\nu\,(y^{(\nu)}v_\nu/\sqrt{2})^{-1}\,Y_\nu^T,$$

which matches (12.208) and the EFT limit (12.186). Hence the light spectrum is suppressed as claimed in (12.208)–(12.209).

(iii) Compatibility with the IQR diagnostic realized reference configuration and HSP. (a) *No new SM gauge quanta.* By construction $\Phi_\nu \sim (1,1)_0$ introduces no additional components in the SM connection split (Section 12.4.4) and thus no new gauge bosons; the Operator–Bundle electroweak/gluonic sectors (Definition 12.95) are unchanged.

(b) *Anomaly viability (HSP Stage I).* The right–handed neutrinos $\nu_R \sim (1,1)_0$ and the SM–singlet scalar Φ_ν carry no SM gauge charges; consequently the anomaly block eq. (12.46) is unaffected, preserving the Stage I criterion of the Holosymmetric Selection Principle (Section 12.5.4 and Theorem 12.42). BRST structure and gauge invariance remain intact.

(c) *Diagnostic realized reference configuration discipline (Nielsen/KL hygiene).* The block–diagonalization in Proposition 12.140 yields realized propagator poles for light/heavy neutrinos with gauge–parameter–independent positions and positive Källén–Lehmann weights on certified plateaux (Proposition 12.88); the singlet sector does not spoil these properties.

(d) *Family bundle architecture.* Both $y^{(\nu)}$ and Y_ν act fibrewise on the rank–3 family bundle (Definitions 12.106 and 12.114); observable mixing is the relative family holonomy between species (Proposition 12.115 and eq. (12.134)), and (12.211) is compatible with the PMNS construction (Section 12.7.2 and Definition 12.141).

Combining (i)–(iii) establishes the proposition: $\langle\Phi_\nu\rangle \neq 0$ spontaneously breaks $U(1)_L$ and induces M_R as in (12.207); inserting M_R into the Type–I seesaw produces the suppressed light spectrum (12.208)–(12.209), and the entire construction remains consistent with the IQR diagnostic realized reference configuration and HSP requirements. □

Interpretation. The Majorana seesaw here is not an ad hoc addition but a *dynamical branch* of the Tantrakyma sector: a neutral singlet Φ_ν in $\mathcal{V}_\Theta$ generates M_R upon condensation, and the ensuing block–diagonalization (Proposition 12.140) feeds down to tiny active–neutrino masses. The smallness of M_ν thus reflects (i) the hierarchy between the EWSB scale v and the singlet scale v_ν, and (ii) the holonomy–structured flavor endomorphisms on $\mathcal{F}$ that also underwrite PMNS mixing (Section 12.7.2 and Proposition 12.115).

12.7.8 Gauging anomaly–free $U(1)_{B–L}$

Assume three right–handed neutrinos $\nu_R \sim (1,1)_0$ so that $U(1)_{B–L}$ is anomaly–free. Gauging this factor extends the internal connection in a controlled way while preserving all established structures.

What gets added.

(i) Gauge–sector extension The connection of Section 12.4.4 is enlarged by an extra abelian piece

$$\mathcal{A}_\mu \;\longrightarrow\; \mathcal{A}_\mu \;+\; g_{BL}\, X_\mu\, Q_{B\text{–}L}, \tag{12.212}$$

with field strength and Yang–Mills term as in Definitions 12.96 and 12.97. A gauge–kinetic–mixing term

$$\frac{\kappa}{2}\, B_{\mu\nu} X^{\mu\nu} \tag{12.213}$$

between hypercharge and $B-L$ is allowed.

(ii) Breaking of $U(1)_{B-L}$ Let the singlet Φ_ν (used for the Majorana sector) carry $B-L$ charge $+2$ as in Section 12.7.7. Its VEV now Higgses the new gauge factor:

$$\langle \Phi_\nu \rangle = v_\nu \quad \Rightarrow \quad m_{Z'} = 2\, g_{BL}\, v_\nu, \tag{12.214}$$

with Z' the massive $B-L$ gauge boson. The would–be Majoron is eaten (no light NG boson).

(iii) Scalar sector The radial singlet s_ν (the "$B-L$ Higgs") can mix with the Tantrakyma/Higgs doublet via a portal term $\lambda_{H\Phi}\,|H|^2|\Phi_\nu|^2$ inside $\mathcal{V}_\Theta$ (Definition 9.335), yielding a 2×2 scalar mass matrix consistent with diagnostic realized reference configuration discipline.

Here, "$B-L$" means **baryon number minus lepton number**. It is a $U(1)$ charge

$$Q_{B\text{–}L} = B - L,$$

assigned to every field by counting its baryon (B) and lepton (L) numbers. In the Standard Model representations this gives

$$Q_{B\text{–}L}(q_L, u_R, d_R) = \tfrac{1}{3}, \qquad Q_{B\text{–}L}(\ell_L, e_R) = -1, \qquad Q_{B\text{–}L}(H) = 0.$$

If we also include one right–handed neutrino per generation, $\nu_R \sim (1,1)_0$ with $Q_{B\text{–}L}(\nu_R) = -1$, then the gauge symmetry $U(1)_{B-L}$ is *anomaly–free* (all mixed and cubic anomalies cancel). That is why Chapter 12 explicitly assumes three ν_R's before "gauging" $B-L$.

Thus, *gauging* $U(1)_{B-L}$ means adding a new abelian gauge field X_μ and coupling it to the conserved current via the generator Q_{B-L}:

$$\mathcal{A}_\mu \;\longrightarrow\; \mathcal{A}_\mu + g_{BL}\, X_\mu\, Q_{B\text{–}L},$$

with field strength $X_{\mu\nu}$ and (optionally) a *kinetic–mixing* term $\frac{\kappa}{2}\, B_{\mu\nu} X^{\mu\nu}$ between hypercharge $B_{\mu\nu}$ and the new $B-L$ field. To break $U(1)_{B-L}$ (so the new gauge boson is massive) the text takes a singlet scalar Φ_ν of charge $+2$. Its VEV,

$$\langle \Phi_\nu \rangle = v_\nu,$$

Higgses $U(1)_{B-L}$ and produces a heavy boson Z' with mass $m_{Z'} = 2\, g_{BL}\, v_\nu$. The corresponding would–be Goldstone (the "Majoron") is eaten, so there is no light NG boson. The same Φ_ν also allows a *Majorana mass* for ν_R (schematically $y_\nu\, \Phi_\nu\, \nu_R^T C\, \nu_R$), realizing the usual seesaw logic, while a portal term $\lambda_{H\Phi}|H|^2|\Phi_\nu|^2$ allows mixing between the radial singlet s_ν and the Higgs sector—done in a way compatible with the diagnostic realized reference configuration/BRST discipline and the unified connection setup of earlier sections.

What stays the same (low energy).

- **Seesaw and PMNS.** The Type–I seesaw and PMNS constructions are unchanged in form,
$$M_\nu \;=\; -\,M_D\,M_R^{-1}\,M_D^T, \qquad U_{\rm PMNS} \;=\; U_{eL}^\dagger U_\nu, \tag{12.215}$$
with the Dirac–Majorana sector in Section 12.7.3, the seesaw in Section 12.7.4, and mixing in Sections 12.7.2 and 12.7.5. Integrating out ν_R (and a heavy Z') still yields the Weinberg operator with $C_5 = Y_\nu M_R^{-1} Y_\nu^T$ (Section 12.7.4).

- **CKM/GIM and FCNCs.** $B-L$ is family–universal, so no tree–level FCNCs are induced; SM GIM cancellations (eqs. (12.142) and (12.143)) are unaffected. Any Z–Z' effects arise only through small kinetic/mass mixing and preserve flavour universality at leading order.

- **Family bundle.** The flat Z_3 family bundle $(\mathcal{F}, \mathcal{A}^{(F)})$ remains SM–neutral and does not couple to Q_{B-L} (Definition 12.106); mixing as relative family holonomy is therefore unchanged (Proposition 12.115).

Consistency with HSP and the diagnostic realized reference configuration.

(i) Stage I (viability) With three ν_R, gauged $U(1)_{B-L}$ is anomaly–free, so the anomaly block $\mathcal{C}(A)$ and BRST nilpotency remain satisfied (Section 12.5.4, eq. (12.46), and Theorem 12.42). The enlarged connection still fits the Operator–Bundle formalism, and propagator poles/residues retain Nielsen/KL hygiene on certified plateaux (Proposition 12.88).

(ii) Stage II (economy/orthogonality) If $m_{Z'}$ lies well above the electroweak scale (large v_ν or modest g_{BL}), the extra vector effectively decouples and the realized bright–fringe sector is unchanged. A relatively light Z' is an additional visible sector; selection demands it not spoil orthogonality (no redundant, family–nonselective copies), which is automatically satisfied because $B-L$ is independent of the family holonomy and couples universally.

Phenomenological implications (qualitative).

(i) Majoron absence Unlike the global $U(1)_L$ case, no physical Majoron remains; the longitudinal Z' is the eaten mode.

(ii) Leptogenesis interplay A light or strongly coupled Z' can keep ν_R in equilibrium longer, tending to wash out a lepton asymmetry; successful seesaw leptogenesis prefers either heavier Z' or small g_{BL} so that decays are out of equilibrium—model–building latitude within $\mathcal{V}_\Theta$ (Section 12.7.4 and Definition 9.335).

(iii) Higgs/portal phenomenology The singlet–doublet portal shifts scalar masses/couplings while respecting the EWSB analysis (Section 12.4.5); diagnostic realized reference configuration/Nielsen–KL properties remain intact (Proposition 12.88).

Bottom line. Gauging anomaly–free $U(1)_{B-L}$ promotes the lepton–number breaking used for M_R from *global* SSB to a *gauge* SSB. The Majorana seesaw, PMNS structure, CKM/GIM phenomenology, and the family–bundle picture survive unchanged in form; a heavy Z' and a singlet scalar are the controlled byproducts, fully compatible with the Operator–Bundle, HSP selection, and diagnostic realized reference configuration hygiene established earlier.

Lepton Number Violation and Majorana Couplings in $\mathcal{V}_\Theta$

In the Standard Model, total lepton number $U(1)_L$ is an accidental global symmetry [82, 91]. In the IQR setting, neutrinos admit both Dirac and Majorana mass terms because $\nu_R \sim (1,1)_0$ is a gauge singlet (Section 12.7.3); see [82, 91, 129, 547]. The Dirac piece arises from the Yukawa sector with the Higgs–kymon background $\bar{\Theta} = \langle H \rangle$ (Section 12.4.6 and Definition 12.139), whereas a *Majorana* mass for ν_R can be sourced by a neutral Tantrakyma scalar that carries lepton number, as in Majoron/$B-L$ realizations [570, 571, 131].

Operator structure in the Tantrakyma potential. Let Φ_ν be a neutral scalar (SM singlet) in the Tantrakyma sector with lepton charge $L(\Phi_\nu) = -2$, and let $y^{(\nu)}_{\alpha\beta} = y^{(\nu)}_{\beta\alpha}$ be flavor–symmetric couplings. The $\Delta L = 2$ channel of the potential is

$$\mathcal{V}^{(\nu)}_\Theta \;:=\; \frac{1}{2}\sum_{\alpha,\beta}\Big(y^{(\nu)}_{\alpha\beta}\,\Phi_\nu\,\nu^T_{R,\alpha}C\,\nu_{R,\beta} \;+\; \text{h.c.}\Big), \tag{12.216}$$

which is Lorentz invariant, gauge invariant under $G_{\rm SM}$, and *globally* $U(1)_L$ invariant *before* symmetry breaking (because $L[\Phi_\nu] = -2$ cancels $L[\nu_R^T C\nu_R] = +2$) [570, 571]. When Φ_ν condenses,

$$\langle\Phi_\nu\rangle = v_\nu \qquad\Longrightarrow\qquad M_R \;=\; y^{(\nu)}\,v_\nu, \tag{12.217}$$

a right–handed Majorana mass matrix M_R is dynamically generated inside $\mathcal{V}_\Theta$ (Definition 12.135); cf. [570, 571, 131]. Together with the Dirac mass $M_D = \frac{v}{\sqrt{2}}Y_\nu$ from EWSB, this realizes the Type–I seesaw (Section 12.7.4):

$$M_\nu \;=\; -\,M_D\,M_R^{-1}\,M_D^T \quad\Rightarrow\quad m_\nu \sim \frac{(y_D\,v)^2}{y^{(\nu)}\,v_\nu}, \tag{12.218}$$

in agreement with eq. (12.188) and the EFT Weinberg operator eq. (12.186) [129, 547, 91].

Proposition 12.149 (Spontaneous $U(1)_L$ breaking via Tantrakyma condensation)**.** *With $L(\Phi_\nu) = -2$, the vacuum $\langle\Phi_\nu\rangle \neq 0$ in (12.217) spontaneously breaks the global lepton number $U(1)_L$ and induces a Majorana mass $M_R = y^{(\nu)}\langle\Phi_\nu\rangle$. Inserting M_R into the seesaw relation (12.218) yields a light spectrum suppressed by M_R^{-1}, while preserving gauge invariance, anomaly viability, and the family–bundle architecture (Section 12.5.4, eq. (12.46), and Definition 12.114); see [570, 571, 131, 91].*

Idea. Gauge and Lorentz invariance of (12.216) are immediate since ν_R is an SM singlet and Φ_ν is a scalar singlet. With $L(\Phi_\nu) = -2$, the operator is globally $U(1)_L$ invariant; SSB occurs when Φ_ν develops a VEV, which replaces Φ_ν by v_ν and produces M_R [570, 571]. The light spectrum then follows from the Schur–complement/Takagi block diagonalization

of Section 12.7.4, yielding $M_\nu = -M_D M_R^{-1} M_D^T$ (eq. (12.188); cf. [129, 547, 91]). Since Φ_ν is $G_{\rm SM}$–neutral, no new SM gauge quanta are introduced (Section 12.4.4); anomaly constraints are unchanged (Section 12.5.4), and $y^{(\nu)}, Y_\nu \in \mathrm{End}(\mathcal{F})$ fit the family–lifted formalism (Definitions 12.110 and 12.114). □

Proof. **(i) Lorentz and $G_{\rm SM}$ gauge invariance of (12.216).** Write the Majorana bilinear with the charge–conjugation matrix C as $\nu_{R,\alpha}^T C\, \nu_{R,\beta}$. Under a proper Lorentz transformation $\nu_R(x) \mapsto S(\Lambda)\,\nu_R(\Lambda^{-1}x)$ one has $S(\Lambda)^T C\, S(\Lambda) = C$, hence $\nu_R^T C \nu_R$ is a Lorentz scalar [82, 91, §3, §7]. Under $G_{\rm SM} = SU(3)_C \times SU(2)_L \times U(1)_Y$, the field $\nu_R \sim (1,1)_0$ and Φ_ν are singlets, so both $\nu_R^T C \nu_R$ and Φ_ν are invariant and the product in (12.216) is $G_{\rm SM}$–gauge invariant. Fermi statistics then require a symmetric flavor coupling $y^{(\nu)}_{\alpha\beta} = y^{(\nu)}_{\beta\alpha}$ [82, 91].

(ii) Global $U(1)_L$ and spontaneous breaking. Assign lepton number $L(\nu_R) = +1$ and $L(\Phi_\nu) = -2$. Then $L[\nu_R^T C \nu_R] = +2$ and $L[\Phi_\nu] = -2$, so the operator $\Phi_\nu\, \nu_R^T C \nu_R$ is globally $U(1)_L$ invariant prior to symmetry breaking. When Φ_ν develops a VEV,

$$\langle \Phi_\nu \rangle = v_\nu \qquad \Longrightarrow \qquad \mathcal{V}_\Theta^{(\nu)} \supset \tfrac{1}{2}\, \overline{\nu_R^c}\, (y^{(\nu)} v_\nu)\, \nu_R + \text{h.c.}\,,$$

which yields the right–handed Majorana mass $M_R = y^{(\nu)} v_\nu$ and *spontaneously* breaks $U(1)_L$ [570, 571]. (If instead an anomaly–free $U(1)_{B-L}$ is gauged with three ν_R, the same VEV Higgses $U(1)_{B-L}$ and the would–be Majoron is eaten [131].)

(iii) Dirac mass and full neutral–lepton mass matrix. Electroweak symmetry breaking (EWSB) with $\langle H \rangle = \bar{\Theta} = (0, v/\sqrt{2})^\top$ produces the Dirac mass $M_D = \frac{v}{\sqrt{2}} Y_\nu$ from $-\overline{L_L} \tilde{H}\, Y_\nu\, \nu_R + \text{h.c.}$ (Definition 12.139; [82, 91]). In the Nambu–Gorkov basis $\Psi = (\nu_L, \nu_R^c)^\top$, the neutral–lepton mass Lagrangian becomes

$$\mathcal{L}_\nu^{\rm mass} = -\tfrac{1}{2}\, \overline{\Psi} \begin{pmatrix} 0 & M_D \\ M_D^T & M_R \end{pmatrix} \Psi + \text{h.c.},$$

the standard Dirac–Majorana structure of the Type–I seesaw [129, 547, 91].

(iv) Seesaw block diagonalization and the effective light mass. In the technically natural regime $\|M_R\| \gg \|M_D\|$, define the small mixing matrix $\Theta := M_D M_R^{-1}$. Consider the unitary transformation

$$\mathcal{U} \;=\; \exp\!\begin{pmatrix} 0 & \Theta \\ -\Theta^\dagger & 0 \end{pmatrix} \;=\; \begin{pmatrix} 1 - \frac{1}{2}\Theta\Theta^\dagger & \Theta \\ -\Theta^\dagger & 1 - \frac{1}{2}\Theta^\dagger\Theta \end{pmatrix} + \mathcal{O}(\Theta^3).$$

A straightforward computation gives the block–diagonalized mass matrix up to $\mathcal{O}(\Theta^3)$,

$$\mathcal{U}^T \begin{pmatrix} 0 & M_D \\ M_D^T & M_R \end{pmatrix} \mathcal{U} \;=\; \begin{pmatrix} -\,M_D M_R^{-1} M_D^T & 0 \\ 0 & M_R + \frac{1}{2}\big(\Theta^T M_D + M_D^T \Theta\big) \end{pmatrix} \;+\; \mathcal{O}\!\left(\frac{M_D^3}{M_R^2}\right).$$

Thus the light and heavy spectra are, respectively,

$$M_\nu \;=\; -\,M_D M_R^{-1} M_D^T, \qquad M_N \;=\; M_R + \mathcal{O}\big(M_D^2/M_R\big),$$

with active–sterile mixing $\Theta = M_D M_R^{-1}$ (Section 12.7.4 and eqs. (12.188) and (12.189)). Equivalently, integrating out ν_R gives the Weinberg operator with $C_5 = Y_\nu M_R^{-1} Y_\nu^T$ and $M_\nu = -\frac{v^2}{2} C_5$ [91, 82].

(v) Symmetry, diagonalization, and phases. M_ν is complex symmetric; hence it admits a Takagi factorization $U_\nu^T M_\nu U_\nu = \mathrm{diag}(m_1, m_2, m_3)$ with $m_i \geq 0$, realizing light *Majorana* eigenmodes. The PMNS matrix $U_{\mathrm{PMNS}} = U_{eL}^\dagger U_\nu$ then carries one Dirac and, generically, two Majorana phases (for three families), as in the standard counting [82, 91].

(vi) Gauge content and anomalies. Because Φ_ν and ν_R are G_{SM} singlets, the construction introduces no new SM gauge quanta (Section 12.4.4). Anomaly constraints of the SM gauge sector remain unchanged (singlets do not contribute to $SU(3)^3$, $SU(2)^2U(1)$, $U(1)^3$, or mixed anomalies) [443, 91]. Family–space endomorphisms Y_ν and $y^{(\nu)}$ are sections of $\mathrm{End}(\mathcal{F})$, consistent with the family–lifted formalism (Definitions 12.110 and 12.114).

Combining (i)–(vi) establishes the gauge/Lorentz invariance of (12.216), the spontaneous $U(1)_L$ breaking via $\langle\Phi_\nu\rangle$, and the emergence of the Type–I seesaw relation $M_\nu = -M_D M_R^{-1} M_D^T$ together with its phase structure and anomaly consistency [570, 571, 131, 129, 547, 91, 82]. □

Remark 12.150 (Global vs. gauged breaking)**.** In the global case, SSB of $U(1)_L$ produces a (pseudo)–Nambu–Goldstone boson (Majoron) [570, 571]. If one instead gauges anomaly–free $U(1)_{B-L}$ with three ν_R, the same VEV v_ν Higgses the new abelian factor, the Majoron is eaten, and a massive Z' appears; seesaw and PMNS structures are unchanged in form (Section 12.7.8; cf. [131, 91]).

Interpretation. The Majorana sector in $\mathcal{V}_\Theta$ plays the role for neutrinos that the Higgs–kymon plays for charged fermions: Dirac masses arise from the EWSB background $\bar{\Theta}$, while Φ_ν supplies M_R through a separate (neutral) SSB channel. The observed smallness of m_ν then reflects the hierarchy between these two condensates, encoded covariantly in the Operator–Bundle/diagnostic realized reference configuration formalism and read out through the PMNS misalignment (Sections 12.7.2 and 12.7.5); see [129, 547, 91, 82].

Inputs (at EWSB)	**Derived quantities (low energy)**
Dirac mass $M_D = \frac{v}{\sqrt{2}} Y_\nu$; Majorana mass $M_R = M_R^T$ (Definition 12.139)	Light-neutrino mass matrix $M_\nu = -M_D M_R^{-1} M_D^T$ (Type-I seesaw, eq. (12.188)); heavy block $M_N \simeq M_R$
Charged-lepton mass matrix M_e	$U_{eL}^\dagger M_e U_{eR} = \mathrm{diag}(m_e, m_\mu, m_\tau)$; $U_\nu^T M_\nu U_\nu = \mathrm{diag}(m_1, m_2, m_3)$; $U_{\mathrm{PMNS}} = U_{eL}^\dagger U_\nu$ (Definition 12.141)
Family-fibre holonomy (triadic monodromy)	Physical phases (Dirac δ, Majorana $\alpha_{1,2}$) arise from relative family holonomy and phases in Y_ν, M_R (Section 12.5, Definition 12.106, and Proposition 12.115)
Outputs / observables	**Expressions / relations**
Mass-squared gaps Δm^2_{21}, Δm^2_{31}	Eigenvalue differences of M_ν: $\Delta m^2_{ij} = m_i^2 - m_j^2$ (normal or inverted ordering)
Mixing angles $\theta_{12}, \theta_{23}, \theta_{13}$ and Dirac phase δ	Standard parameterization of U_{PMNS} (Proposition 12.142); Jarlskog $J_{\mathrm{CP}} = \mathrm{Im}[U_{e1} U_{\mu 2} U_{e2}^* U_{\mu 1}^*] = s_{12} s_{23} s_{13} c_{12} c_{23} c_{13}^2 \sin\delta$
Majorana phases (α_1, α_2)	Neutrinoless double-beta effective mass (eq. (12.220)); rephasing-invariant combinations enter LNV observables

Table 12.3: Mapping from high-scale inputs (Y_ν, M_R) to low-energy neutrino observables

Effective masses and lepton-number violation. Two standard effective masses summarize kinematic and LNV sensitivities:

$$m_\beta^2 := \sum_{i=1}^{3} |U_{ei}|^2 \, m_i^2, \qquad \text{(single } \beta \text{ decay, kinematic)}, \tag{12.219}$$

$$m_{\beta\beta} := \left| \sum_{i=1}^{3} U_{ei}^2 \, m_i \right| = \left| c_{12}^2 c_{13}^2 m_1 e^{i\alpha_1} + s_{12}^2 c_{13}^2 m_2 e^{i\alpha_2} + s_{13}^2 m_3 \right|. \tag{12.220}$$

Here U_{ei} are elements of U_{PMNS} (Definition 12.141 and Proposition 12.142), and (α_1, α_2) are the Majorana phases. *Remark.* In the standard parameterization one may write the third term as $s_{13}^2 m_3 e^{-2i\delta}$; however, the overall magnitude $|m_{\beta\beta}|$ is independent of the Dirac phase and this phase can be absorbed into a redefinition of (α_1, α_2). The scale of $m_{\beta\beta}$ follows from eq. (12.188) and is controlled by the interplay of mass ordering, mixing angles, and the Majorana phases; its nonzero value signals lepton-number violation as anticipated in the Dirac–Majorana synthesis [544, 546, 547]. (For purely Dirac neutrinos, $m_{\beta\beta} = 0$ and LNV is absent.)

12.8 Antimatter in the IQR/Tantrakyma Standard Model

In the Chapter 12 formalism, antimatter is treated uniformly by combining: (i) charge conjugation on matter representations of the internal connection, (ii) the flat holonomy

of the family fibre, and (iii) BRST/diagnostic realized reference configuration hygiene so that physical poles and residues are well-defined (Section 12.4.4, Definitions 12.95, 12.96 and 12.106, and Proposition 12.88; cf. [82, 91]).

1) Antiparticles as charge–conjugate kymons

For any Tantrakyma fermion kymon ψ in a representation $R \subset G_{\mathrm{SM}}$, define the charge–conjugate field by

$$\psi^c \;:=\; \mathcal{C}\,\bar{\psi}^T, \tag{12.221}$$

which transforms in the conjugate representation $\bar{R}$ with abelian charges reversed (e.g. $Y \to -Y$ and hence $Q = I_3 + \frac{1}{2}Y \to -Q$) [82, 91, §3, §20]; cf. the Gell–Mann–Nishijima relation [126]. This sits directly in the operator–bundle/connection split of Section 12.4.4 and Definitions 12.95 and 12.96.

2) Tantrakyma kymons and their antiparticles

(i) Quark–kymons $(u, d, \ldots)$ $\mathbf{3} \to \bar{\mathbf{3}}$ under $SU(3)_C$, with Y flipped [82, 91].

(ii) Lepton–kymons (e, ν) e_L, e_R acquire conjugate hypercharges; the neutrino case depends on Dirac vs. Majorana (see §12.8 6) [82, 91].

(iii) Tantron (SM gauge bosons) gluons g^a, photon γ, and Z^0 are self–conjugate; $W^\pm$ are conjugate to each other (Section 12.4.4 and Definition 12.102; [82, 91]).

(iv) Chrono–kymon SM–neutral spin-1 in the τ channel; TT–null and typed–separated, so "antiparticle'' is the same neutral excitation (Section 10.7, Lemma 10.59, and Corollary 10.125).

(v) Higgs–kymon h real scalar, self–conjugate (Definition 9.343 and eq. (9.185); [82, 91]).

3) Family fibre and antimatter

The family bundle $(\mathcal{F}, \mathcal{A}^{(F)})$ is SM–neutral and flat, with Z_3 monodromy $\mathrm{diag}(1, \omega, \omega^2)$ (Definition 12.106 and eq. (12.124)). Under charge conjugation the holonomy phases complex–conjugate, $1 \leftrightarrow 1$, $\omega \leftrightarrow \omega^2$. Equivalently, loop orientation reversal inverts the holonomy. Thus particle vs. antiparticle acquire conjugate family phases on the same loop. The "relative family holonomy'' formulae for mixing remain valid with $U^{(F)} \mapsto U^{(F)*}$ for anti–states (Proposition 12.115 and eq. (12.134); cf. [82]).

4) Hadrons and anti-hadrons

Color–singlet projection treats quark and antiquark sectors symmetrically:

(i) Mesons $3 \otimes \bar{3} = \mathbf{1} \oplus \mathbf{8}$; the singlet $q\bar{q}$ is gauge invariant (Definitions 12.118 and 12.126 and eq. (12.145); [82, 91]).

(ii) Baryons $3 \otimes 3 \otimes 3 = \mathbf{1} \oplus \mathbf{8} \oplus \mathbf{8} \oplus \mathbf{10}$; the singlet $\epsilon_{ijk} q^i q^j q^k$ is gauge invariant (eqs. (12.146) and (12.147); [82, 91]).

(iii) Anti-baryons $\epsilon^{ijk}\bar{q}_i\bar{q}_j\bar{q}_k$ furnish the conjugate singlet (same references).

BRST cohomology selects these singlets as physical hadron/anti-hadron states; non-singlets are BRST–exact or orthogonal (Proposition 12.120 and Theorem 12.133; cf. [108, 442]).

5) CKM/GIM with antimatter

The charged–current weak interaction relates particle ↔ antiparticle processes by complex conjugation of CKM factors, $V_{\rm CKM} \to V^*_{\rm CKM}$ in the conjugate channels (eq. (12.139) and Definition 12.117; [82, 91]). GIM suppression of FCNC loops is preserved (and identical for anti-processes) because it follows from unitarity, $\sum_i V_{iq}V^*_{iq'} = 0$, and common loop functions $F(m_i^2/M_W^2)$ (eqs. (12.142) and (12.143); see [82, 91]).

6) Neutrinos: Dirac vs. Majorana

(i) Dirac case ($M_R = 0$) ν and $\bar{\nu}$ are distinct; $U_{\rm PMNS} = U^\dagger_{eL}V_{\nu L}$ (Section 12.7.2; [545, 544, 82]).

(ii) Majorana/seesaw light states obey $\nu = \nu^c$ (up to phases); there is no distinct antineutrino field. The mixing is $U_{\rm PMNS} = U^\dagger_{eL}U_\nu$ with additional Majorana phases (Section 12.7.4 and Definition 12.136; [129, 547, 91]), and lepton–number–violating observables such as neutrinoless double beta decay probe this (eq. (12.220); cf. [91]).

7) CP, CPT, and baryon asymmetry

On certified plateaux the theory is CPT invariant and fermion two–point functions satisfy Nielsen/Källén–Lehmann hygiene (Proposition 12.88; cf. [120, 82, 91]). CP asymmetries (particle vs. antiparticle rates) arise from the CKM phase (quarks) and can also originate from leptonic and family–holonomy sources (leptons/family) (Proposition 12.115 and Remark 12.116; see [82, 91]). For the cosmic matter–antimatter asymmetry, CKM alone is too small; viable sources include seesaw leptogenesis and (model–dependent) family–holonomy phases or a strengthened electroweak transition via $\mathcal{V}_\Theta$ (with no new light SM vectors) (Section 12.5.5; cf. [129, 547, 91]).

8) Diagnostic realized reference configuration and HSP statements

(i) Stage I (viability) antimatter sectors share the same anomaly–block entries (fixed by chiral content), so $\mathcal{C}(A) = 1$ is unchanged (Section 12.5.4, eq. (12.46), and Theorem 12.42; cf. [443, 91]).

(ii) Stage II (economy) anti–states are required by locality/CPT and occupy orthogonal channels; family–trivial add-ons (if any) are suppressed (Corollary 12.113; cf. [82, 91]).

(iii) Diagnostic realized reference configuration hygiene poles and residues for anti–fields are gauge–parameter independent and KL–positive on the diagnostic realized reference configuration, just like their particle partners (Proposition 12.88; [120, 82]).

In the IQR/Tantrakyma architecture, antimatter is the charge–conjugate sector of the same kymonic fields: conjugate gauge representations, conjugate family phases, identical BRST/diagnostic realized reference configuration certification, and the same color–singlet projection for (anti)hadrons. Neutrinos are special: if Dirac they have distinct antineutrinos; if Majorana (as in the seesaw regime) the light states are self–conjugate, and lepton–number–violating observables (e.g. $0\nu\beta\beta$) become the antimatter signature (Section 12.7.4, Definition 12.136, and eq. (12.220); [91, 82]).

Context and placement. The overview in 12.8 framed antimatter at a conceptual level inside the weave (operator–bundle language, BRST hygiene, and family fibre). The next section, §12.9, refines that overview into three *diagnostic realized reference configuration-formal* statements: (i) closure under complex representations of the internal connection, (ii) relativistic causality plus positive spectral measure on certified plateaux (CPT and conjugate poles), and (iii) the neutrality of family holonomy with phase conjugation under C. These are the minimal structural ingredients we will later re-express as a single consequence of the Axiom of Duality (Definition 2.100); see [82, 91] for standard background on charge conjugation, CPT, and spectral positivity.

12.9 Antimatter from Complex Reps, Causality, Positivity, and Family Holonomy

This section codifies three structural ingredients that together force the particle–antiparticle pairing in the Standard–Model sector of the weave: (i) complex unitary representations of the internal connection (and the induced $*$–structure on the operator bundle), (ii) relativistic causality with positive spectral measure on certified plateaux (CPT and conjugate poles), and (iii) neutrality of the family fibre with conjugation of its holonomy phases under C. Each ingredient is expressed in the diagnostic realized reference configuration language of Section 12.4.4, Definitions 12.95, 12.96 and 12.106, Propositions 12.88 and 12.115, and eqs. (12.124) and (12.134).

12.9.1 Complex Representations of the Internal Connection

Definition 12.151 (Complex–representation pairing (CRP))**.** Let ψ be a *fermionic Tantrakyma kymon* (spin-$\frac{1}{2}$) transforming in a unitary, genuinely complex irrep $R \subset G_{\mathrm{SM}}$ of the internal connection specified in Section 12.4.4. The *charge–conjugate field* is

$$\psi^c \;:=\; C\,\bar{\psi}^{\top}, \qquad \bar{\psi} := \psi^{\dagger}\gamma^0, \qquad \text{with} \quad \psi^c \in \bar{R},$$

so that all additive charges (e.g. hypercharge Y, color weight, and lepton number in the Dirac case) flip sign. The minimal operator–bundle closure of the matter sector therefore contains $R \oplus \bar{R}$.

Proposition 12.152 (Reality/Unitarity $\Rightarrow R \oplus \bar{R}$ closure)**.** *For any matter content built from R–valued fields, the operator–bundle action with gauge fields of Definitions 12.95 and 12.96*

is real and generates unitary time evolution on the diagnostic realized reference configuration iff the field set is closed under the CRP of Definition 12.151. Concretely, a quark–kymon in $\mathbf{3}$ *of* $SU(3)_C$ *entails an anti–quark–kymon in* $\bar{\mathbf{3}}$*, and an electron with hypercharge* Y *entails a positron with* $-Y$*.*

Proof. (*Necessity*). Let ψ be a Tantrakyma field transforming in a genuinely complex irrep $R \subset G_{\mathrm{SM}}$ of the internal connection (Section 12.4.4). The covariant derivative is

$$D_\mu \psi = \partial_\mu \psi + i\, g\, A_\mu^a\, T_R^a\, \psi,$$

with gauge bosons and curvature as in Definitions 12.95 and 12.96, and with Hermitian generators T_R^a in the unitary representation R. Consider the local matter current $J_\mu^a := \bar{\psi}\, \gamma_\mu T_R^a \psi$ (spinorial notation is for definiteness; the scalar case is analogous with $\bar{\psi} \mapsto \psi^\dagger$). Since T_R^a are Hermitian, J_μ^a is gauge-covariant; moreover the kinetic and gauge-coupling terms are built from such bilinears. Reality of the action and self-adjointness of the Hamiltonian require that the local density be closed under the $*$–operation. Taking the adjoint,

$$\left(J_\mu^a(x)\right)^\dagger = \psi^\dagger(x)\, T_R^a\, \gamma_\mu^\dagger\, \gamma^0 = \bar{\psi}(x)\, \gamma_\mu \left(T_R^a\right)^*,$$

and using $T_R^a = (T_R^a)^\dagger$ shows that the adjoint current sits in the algebra generated by $\bar{\psi}, \psi$; however, at the *field* level, the $*$–operation maps the R–valued field to an operator transforming in the conjugate irrep $\bar{R}$. Indeed, with the standard charge–conjugation matrix C satisfying $C\gamma^\mu C^{-1} = -(\gamma^\mu)^\top$, define

$$\psi^c := C\, \bar{\psi}^\top, \qquad \bar{\psi}^c := \psi^\top C^{-1}.$$

A direct computation gives the covariant transformation law

$$D_\mu \psi^c = C \left(D_\mu \psi\right)^{*\top} = \partial_\mu \psi^c + i\, g\, A_\mu^a\, T_{\bar{R}}^a\, \psi^c, \qquad T_{\bar{R}}^a := -(T_R^a)^\top,$$

so ψ^c transforms in $\bar{R}$. If the field content is *not* closed under the complex–representation pairing (CRP) of Definition 12.151—i.e. if ψ^c (or, equivalently, the creation operators for the $\bar{R}$ quanta) is not present in the operator–bundle—then the $*$–operation maps the local field polynomial algebra *out* of the chosen field set. In particular, the interaction density $\bar{\psi}\gamma^\mu A_\mu^a T_R^a \psi$ has an adjoint which requires the $\bar{R}$ sector at the field level. Hence the action cannot be real (its integrand is not $*$–closed) and the Hamiltonian cannot be self-adjoint on the intended domain; unitarity of time evolution fails. This contradicts the hypothesis of reality/unitarity on the diagnostic realized reference configuration. Therefore R–valued matter content must be closed under CRP, i.e. contain $R \oplus \bar{R}$.

(*Sufficiency*). Conversely, assume the field set is closed under CRP (Definition 12.151), so for each R–valued field ψ the conjugate ψ^c (or, equivalently, the adjoint modes creating $\bar{R}$ quanta) is present. Using $T_{\bar{R}}^a = -(T_R^a)^\top$ and the identities for C, one finds

$$\left(\bar{\psi}\, \gamma^\mu T_R^a \psi\right)^\dagger = \bar{\psi}\, \gamma^\mu T_R^a \psi, \qquad \left(\bar{\psi}\, \psi\right)^\dagger = \bar{\psi}\, \psi, \qquad \left(D_\mu \psi\right)^\dagger D^\mu \psi = \overline{D_\mu \psi}\, D^\mu \psi,$$

and similarly for scalar and Yukawa couplings, so the Lagrangian density built from the standard gauge–invariant bilinears is manifestly Hermitian. The corresponding Hamiltonian

is self-adjoint on the domain generated by the CAR/CCR of (ψ, ψ^c), and time evolution is unitary (Stone's theorem). Thus the operator–bundle action with the gauge fields of Definitions 12.95 and 12.96 is real and yields unitary evolution.

(*Examples*). For $SU(3)_C$, the fundamental $\mathbf{3}$ has $T^a_{\overline{\mathbf{3}}} = -(T^a_{\mathbf{3}})^\top$; therefore a quark–kymon in $\mathbf{3}$ entails an anti–quark–kymon in $\overline{\mathbf{3}}$. For $U(1)_Y$, the representation $e^{iY\alpha}$ conjugates to $e^{-iY\alpha}$; hence an electron with hypercharge Y entails a positron with $-Y$. Both instances realize the $R \oplus \overline{R}$ closure required above.

Combining necessity and sufficiency completes the proof. □

Remark 12.153 (Meaning of "Reality" in Proposition 12.152)**.** In Proposition 12.152, the term *Reality* refers to the $*$–Hermiticity of the gauge–matter action as an operator functional on the certified, diagnostic realized reference configuration $\mathfrak{P}$ (Definition 9.185). Concretely:

(i) Local (density) level

The Lagrangian density $\mathcal{L}$ built from R–valued matter fields, their covariant derivatives D_μ and the gauge fields of Definitions 12.95 and 12.96 is $*$–closed as an *operator density*,

$$\mathcal{L}^\dagger = \mathcal{L} \quad \text{up to total derivatives and BRST-exact terms vanishing in } H^0(\hat{Q}_{\mathrm{BRST}}). \tag{12.222}$$

With unitary representations (Hermitian generators T^a_R) and Hermitian gauge fields, standard bilinears such as $\bar{\psi}\, i\gamma^\mu D_\mu \psi$ and $\bar{\psi}\, \gamma^\mu A^a_\mu T^a_R \psi$ are $*$–invariant *iff* the local field algebra is closed under the complex–representation pairing (CRP) of Definition 12.151. For genuinely complex R, the $*$–operation requires the charge–conjugate field

$$\psi^c := C\, \bar{\psi}^{\top}, \qquad D_\mu \psi^c = \partial_\mu \psi^c + i\, g\, A^a_\mu\, T^a_{\bar{R}}\, \psi^c, \qquad T^a_{\bar{R}} = -(T^a_R)^\top, \tag{12.223}$$

so that the density remains in the same operator algebra. Absence of the $\bar{R}$ sector would map $\mathcal{L}$ outside the chosen field set, violating (12.222).

(ii) Global (action) level

The action $S = \int d\mu_{\bar{\Sigma}}\, \mathcal{L}$ is *real* as an operator,

$$S^\dagger = S, \tag{12.224}$$

so that the path–integral weight e^{iS} has unit modulus and the Noether charges derived from $\mathcal{L}$ are represented by Hermitian operators on $\mathcal{H}_{\mathrm{phys}}$. On the diagnostic realized reference configuration $\mathfrak{P}$, gauge–parameter independence and KL–positivity ensure the stability of these statements (Proposition 12.88).

(iii) Hamiltonian and time evolution

The $*$–Hermitian density implies a self–adjoint Hamiltonian $H = H^\dagger$ (after the standard Legendre transform and BRST projection), hence unitary evolution $U(t) = e^{-iHt}$ on $\mathcal{H}_{\mathrm{phys}}$.

The conjunction of (i)–(iii) is what is denoted "Reality/Unitarity." For a genuinely complex irrep R, these conditions hold *iff* the matter content is CRP–closed, i.e. contains $R \oplus \bar{R}$ (Definition 12.151). Representative instances are:

$$\mathbf{3} \;\leftrightarrow\; \bar{\mathbf{3}} \quad (SU(3)_C)\,, \qquad Y \;\leftrightarrow\; -Y \quad (U(1)_Y)\,, \tag{12.225}$$

ensuring that interaction terms such as $\bar{\psi}\,\gamma^{\mu}A^{a}_{\mu}T^{a}_{R}\psi$ and their Hermitian conjugates remain within the same operator algebra. This is precisely the content formalized in Proposition 12.152.

12.9.2 Relativistic Causality + Positivity ⇒ CPT and Conjugate Poles

We work on a certified plateau $\mathfrak{P}$ (Ch. 6 discipline), where two–point functions have gauge–parameter–independent poles and a positive Källén–Lehmann measure (Proposition 12.88).

Proposition 12.154 (Diagnostic realized reference configuration CPT pairing of poles)**.** *In a local, Lorentzian theory with positive energy on $\mathfrak{P}$, every isolated pole of a one–particle state with mass m, spin s, and nonzero additive charges has a conjugate pole with the same (m, s) and opposite charges. Equivalently, the physical spectrum is closed under*

$$(m, s;\ \mathsf{q}) \ \longmapsto\ (m, s;\ -\mathsf{q}),$$

with residues related by the antiunitary CPT map on the one–particle sector. Operationally, this is the diagnostic realized reference configuration formulation of the "negative–frequency ⇒ antiparticle'' correspondence. Moreover, analytic S–matrix crossing relates processes with an incoming particle to processes with an outgoing antiparticle, so the amplitude level is pair-aware by construction.

Remark 12.155 (Diagnostic realized reference configuration CPT pairing and conjugate charges)**.** On a certified plateau $\mathfrak{P}$ (so that Källén–Lehmann positivity and gauge–parameter–independent poles hold by Proposition 12.88), Proposition 12.154 is the diagnostic realized reference configuration formulation of the CPT theorem for isolated one–particle poles:

(i) Statement

Every isolated pole of a physical one–particle state with mass m, spin s, and additive charge vector q has a *conjugate pole* corresponding to the antiparticle with the same (m, s) and opposite charges. Equivalently, the spectrum is closed under

$$(m, s;\, \mathsf{q}) \ \longmapsto\ (m, s;\, -\mathsf{q}),$$

and the residues are related by the antiunitary CPT map Θ_{CPT} on the one–particle sector (cf. eq. (12.232)).

(ii) Meaning of q and $-\mathsf{q}$

The symbol q denotes the vector of *additive* charges, e.g.

$$\mathsf{q} = (Q,\ Y,\ B,\ L,\ B-L,\ \ldots),$$

which flip sign under charge conjugation C:

$$(Q, Y, B, L, B-L, \ldots) \ \longmapsto\ (-Q, -Y, -B, -L, -(B-L), \ldots).$$

For non-abelian "charges'' (e.g. color), the corresponding transformation is *representation conjugation* $R \mapsto \bar{R}$ (the Complex-Representation Pairing; Definition 12.151), realized at the field level by $\psi^{c} = C\,\bar{\psi}^{\top}$.

(iii) Correlator form (conjugate poles)

If an R–valued two–point function has residue

$$\mathrm{Res}_{p^2\to m^2}\ \mathbf{G}_F^{(R)}(p) \;=\; Z\,\Pi_{(m,s)}(p),$$

then CPT + CRP imply

$$\mathrm{Res}_{p^2\to m^2}\ \mathbf{G}_F^{(\bar{R})}(p) \;=\; Z\,\Pi_{(m,s)}(p),$$

i.e. the same pole at $p^2 = m^2$ with the same covariant projector $\Pi_{(m,s)}(p)$ and residue Z but with charges reversed (cf. eq. (12.233), and the KL forms eqs. (12.230) and (12.231)).

(iv) Crossing = pair-aware amplitudes

On $\mathfrak{P}$, LSZ + analyticity give the crossing identity

$$\mathcal{M}(\cdots \to \mathrm{out} + \mathrm{in}(p,\mathsf{q})) \;=\; \Big[\mathcal{M}(\cdots + \mathrm{out}(\tilde{p},-\mathsf{q}) \to \mathrm{out})\Big]^{\text{analytic cont.}}_{\tilde{p}\to -p},$$

so an incoming particle (p,q) is analytically equivalent to an outgoing antiparticle $(-p,-\mathsf{q})$ (eq. (12.235)). This is the operational content behind "pair-aware by construction."

In short, the map $(m,s;\mathsf{q}) \mapsto (m,s;-\mathsf{q})$ encodes the existence of the antiparticle partner with identical mass/spin and opposite additive charges, together with $R \mapsto \bar{R}$ for any non-abelian representation carried by the state.

Proof. Work on a certified, diagnostic realized reference configuration $\mathfrak{P}$ (Definition 9.185), so that correlators are BRST/Nielsen-clean and gauge-parameter–independent on H^0 (Proposition 12.88). Let ψ be a (single-species) Tantrakyma field carrying a nonzero additive charge q in a genuinely complex irrep $R \subset G_{\mathrm{SM}}$ (the neutral/self-conjugate cases are addressed at the end).

Step 1 (KL positivity and one-particle pole). By Proposition 12.88, the two-point function on $\mathfrak{P}$ admits a Källén–Lehmann representation with nonnegative spectral measure. In momentum space, the Wightman/Pauli–Jordan distributions have support on the positive-energy mass shells, and an isolated one-particle excitation with mass m and spin s appears as a simple pole at $p^2 = m^2$ with positive residue $Z > 0$ in the appropriate matrix element. Gauge-parameter independence ensures that both the pole and its residue are physical (diagnostic realized reference configuration-stable).

Step 2 (CPT antiunitarity on the physical sector). Locality (microcausality on $\mathfrak{P}$) and Lorentz covariance of the correlators, together with positive energy, imply the existence of an antiunitary CPT operator Θ_{CPT} acting on the one-particle sector of $\mathcal{H}_{\mathrm{phys}}$.[7] Antiunitarity gives, for normalized one-particle states, a conjugate mapping

$$\Theta_{\mathrm{CPT}} \;:\; |p,s;\mathsf{q}\rangle \;\longmapsto\; e^{i\alpha}\,|p,s;-\mathsf{q}\rangle,$$

[7]The proof does not require rederiving CPT; it uses the standard structural inputs that are already enforced on $\mathfrak{P}$: (i) locality/commutativity at spacelike separation (the cone/local causality of the diagnostic realized reference configuration), (ii) Lorentz covariance of the Wightman functions, and (iii) positivity of the energy and spectral measure, cf. Proposition 12.88.

where $p^\mu p_\mu = m^2$ and $e^{i\alpha}$ is a pure phase determined by conventions. Thus the one-particle spectrum is closed under $(m, s; \mathsf{q}) \mapsto (m, s; -\mathsf{q})$ at the level of state vectors.

Step 3 (Residues/poles map to the conjugate channel). Let $G(p)$ denote the appropriate time-ordered two-point function (or matrix element in a chosen channel). By antiunitarity and the $*$–Hermiticity of the action/Hamiltonian ("Reality/Unitarity", Proposition 12.152 and Remark 12.153), the pole at $p^2 = m^2$ with residue Z in the R-channel is mapped by Θ_{CPT} to a pole at the same location with residue Z in the conjugate channel. At the field level, the charge-conjugate field $\psi^c = C\,\bar{\psi}^\top$ transforms in $\bar{R}$ (Definition 12.151), so the conjugate pole is realized as a physical excitation created by ψ^c acting on the vacuum (or, equivalently, by the R-field acting on the conjugate bra). Therefore the spectrum on $\mathfrak{P}$ is closed under

$$(m, s; \mathsf{q}) \;\longmapsto\; (m, s; -\mathsf{q}),$$

with residues related by the antiunitary CPT map on the one-particle sector.

Step 4 (S-matrix crossing, operational reading). On $\mathfrak{P}$, locality and unitarity (again Proposition 12.152) yield the usual LSZ reduction and analyticity properties of amplitudes; the *crossing* relation follows: an amplitude with an incoming particle of charge q analytically continues to the amplitude with an outgoing antiparticle of charge $-\mathsf{q}$. Hence pair-awareness holds at the amplitude level by construction (cf. the dispersion/positivity framework summarized in the mediation/positivity results of the diagnostic realized reference configuration).

Neutral/self-conjugate cases. If the irrep is real/pseudoreal or $\mathsf{q} = 0$, the above map acts trivially; this is the self-conjugacy clause already encoded in Proposition 12.158. In particular, adjoint gauge bosons and the Higgs are self-conjugate; in the seesaw/Majorana regime the light neutrino satisfies $\nu \equiv \nu^c$.

Combining Steps 1–3 proves the pole-level statement; Step 4 gives the operational S-matrix version. This completes the proof. □

12.9.3 Family Fibre Holonomy is Neutral but Conjugates under C

Definition 12.156 (Charge conjugation on family holonomy)**.** Let $(\mathcal{F}, \mathcal{A}^{(F)})$ be the family bundle with flat Z_3 monodromy (Definition 12.106 and eq. (12.124)). For a closed loop γ in the family fibre,

$$U^{(F)}[\gamma] \in U(3), \qquad C:\; U^{(F)}[\gamma] \;\longmapsto\; U^{(F)}[\gamma]^*,$$

equivalently reversing loop orientation: $U^{(F)}[\gamma^{-1}] = U^{(F)}[\gamma]^\dagger$. Hence particle/antiparticle states carry conjugate family phases around the same loop. The mixing–as–holonomy maps of Proposition 12.115 and eq. (12.134) acquire $U^{(F)} \mapsto U^{(F)*}$ in anti–channels.

12.9.4 Consequences and Canonical Cases

Corollary 12.157 (Existence of antiparticles for charged kymons)**.** *If a kymon carries a nonzero additive charge (color, hypercharge, Dirac lepton number, etc.), then by Propositions 12.152 and 12.154 the spectrum on $\mathfrak{P}$ contains a conjugate state with the same (m, s) and opposite charges. This state is the* antiparticle.

Proof. Let ψ be a Tantrakyma kymon carrying a nonzero additive charge $\mathsf{q} \neq 0$ on the certified plateau $\mathfrak{P}$ (Definition 9.185).

Step 1 (Field content is $R \oplus \bar{R}$). Since $\mathsf{q} \neq 0$, the internal irrep $R \subset G_{\mathrm{SM}}$ is genuinely complex (the conjugation flips the sign of the corresponding additive charge). By Proposition 12.152 ("Reality/Unitarity"), the operator–bundle action is $*$–Hermitian iff the matter content is closed under complex–representation pairing (CRP, Definition 12.151), hence the conjugate field $\psi^c = \mathcal{C}\,\bar{\psi}^{\top}$ transforming in $\bar{R}$ is part of the field set.

Step 2 (Pole pairing on $\mathfrak{P}$). On $\mathfrak{P}$ the two–point functions admit a Källén–Lehmann representation with nonnegative spectral measure and gauge–parameter–independent poles (Proposition 12.88); thus any isolated one–particle excitation created by ψ appears as a physical pole labeled $(m, s; \mathsf{q})$. By the diagnostic realized reference configuration CPT pairing of poles (Proposition 12.154), there exists a conjugate pole with the same mass and spin and opposite charge,

$$(m, s; \mathsf{q}) \;\longmapsto\; (m, s; -\mathsf{q}),$$

with residues related by the antiunitary CPT map on the one–particle sector. Equivalently, ψ^c creates the conjugate excitation in the $\bar{R}$ channel.

Step 3 (Confining/neutral special cases). If the single–kymon excitation is not an asymptotic state (e.g. color confinement), then the statement lifts to color–singlet bound states by BRST cohomology: hadron singlets and anti–hadron singlets arise as conjugate pairs in H^0 (Corollary 12.159). If the irrep is real/pseudoreal or the relevant charge vanishes, the map acts trivially (self–conjugacy), as recorded in Proposition 12.158 (this case is excluded here by $\mathsf{q} \neq 0$).

Combining Steps 1–2 (and Step 3 when needed) shows that the physical spectrum on $\mathfrak{P}$ contains, for each charged kymon excitation $(m, s; \mathsf{q})$, a conjugate excitation $(m, s; -\mathsf{q})$. This conjugate excitation is the *antiparticle*. □

Proposition 12.158 (Self–conjugacy criterion)**.** *If R is real or pseudoreal, or if the relevant additive charge vanishes, the CRP is trivial and ψ is self–conjugate (no independent partner). Gauge bosons in adjoint reps and the Higgs–kymon are self–conjugate; in the neutrino sector, the seesaw/Majorana regime (Section 12.7.4 and Definition 12.136) yields $\nu \equiv \nu^c$.*

Proof. Work on a certified, diagnostic realized reference configuration $\mathfrak{P}$ (Definition 9.185) so that correlators are BRST/Nielsen–clean and poles/residues are gauge–parameter–independent (Proposition 12.88). Let ψ transform in an internal irrep $R \subset G_{\mathrm{SM}}$ and write $\psi^c := C\,\bar{\psi}^{\top}$ for the charge–conjugate field as in Definition 12.151.

(A) Real or pseudoreal R. If R is *real* or *pseudoreal*, there exists a unitary intertwiner $J_R : R \to \bar{R}$ such that

$$T^a_{\bar{R}}\, J_R \;=\; J_R\, T^a_R, \qquad J_R J_R^{*} = \pm \mathbb{I}, \tag{12.226}$$

with "+" for real and "−" for pseudoreal R (Schur–Frobenius). Hence $\chi := J_R^{-1}\psi^c$ transforms in R with the *same* representation matrices as ψ. All gauge–invariant local densities built from ψ and ψ^c can therefore be rewritten purely in terms of ψ (and its adjoint) by replacing $\psi^c \mapsto J_R\,\psi$. In particular, the CRP map of Definition 12.151 acts *within* R up to a fixed, representation–theoretic unitary J_R: there is no independent partner. On the one–particle spectrum, Proposition 12.154 maps $(m, s; \mathsf{q})$ to $(m, s; -\mathsf{q})$; but for real/pseudoreal R one

has $\bar{R} \simeq R$, so the CPT–related state lives in the same internal multiplet. Therefore the excitation created by ψ^c is identical (up to a fixed phase and J_R) to that created by ψ, i.e. ψ is *self–conjugate* and the CRP is trivial.

(B) Vanishing additive charge. If the relevant additive charge is zero, then the CPT pairing in Proposition 12.154 reads $(m, s; 0) \mapsto (m, s; 0)$, i.e. the conjugate pole coincides with the original one. Family–holonomy phases also conjugate trivially, $U^{(F)} \mapsto U^{(F)*} = U^{(F)}$ (flat, neutral fibre; Definition 12.156). Thus the conjugate excitation cannot define an independent state: the field is *self–conjugate* and the CRP again collapses.

(C) Gauge bosons and the Higgs–kymon. Gauge bosons transform in the *adjoint* representation, which is real: there is an invariant metric (Killing form) identifying $\mathrm{ad} \simeq \overline{\mathrm{ad}}$, so the multiplet $A_\mu = A_\mu^a T^a$ is self–conjugate at the representation level (cf. Definitions 12.95 and 12.96). Individual complex *combinations* (e.g. $W^\pm$) are mutually conjugate, but the underlying adjoint multiplet is real, hence CRP–trivial. For the BEH sector, the physical Higgs–kymon $h(x)$ in unitary gauge is a real scalar,

$$\hat{\mathbf{\Theta}}(x) = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v + h(x) \end{pmatrix}, \qquad h^\dagger = h, \tag{12.227}$$

(eq. (12.120)), so h is self–conjugate.

(D) Neutrino seesaw/Majorana regime. In the neutrino sector, the Dirac pair (ν, ν^c) forms a CRP–dyad. Turning on Majorana terms (type–I seesaw) produces mass eigenstates that satisfy the Majorana condition $\nu = \nu^c$; see Section 12.7.4 and Definition 12.136. Thus the light neutrino is self–conjugate in the seesaw/Majorana regime.

Combining (A)–(D) establishes that whenever R is real/pseudoreal or the relevant additive charge vanishes, the complex–representation pairing is trivial and the field is self–conjugate; the listed examples follow. This proves Proposition 12.158. □

Corollary 12.159 (Bound–state conjugation and BRST cohomology)**.** *Color–singlet projection and BRST cohomology produce hadrons and anti–hadrons as conjugate singlets—mesons $q\bar{q}$ vs. $\bar{q}q$; baryons $\epsilon_{ijk} q^i q^j q^k$ vs. $\epsilon^{ijk} \bar{q}_i \bar{q}_j \bar{q}_k$—and discard all non–singlets (Definition 12.126, Proposition 12.120, and Theorem 12.133).*

Proof. Work on a certified, diagnostic realized reference configuration $\mathfrak{P}$ (Definition 9.185) so that correlators are BRST/Nielsen–clean and poles/residues are gauge–parameter–independent (Proposition 12.88). Let q^i denote a quark–kymon in the fundamental $\mathbf{3}$ of $SU(3)_C$ and $\bar{q}_i$ its conjugate in $\bar{\mathbf{3}}$, with charge–conjugation implemented at the field level by the CRP map of Definition 12.151.

Step 1 (Color–singlet projection on constituents). By Definition 12.126, there exist invariant projectors onto the $SU(3)_C$ singlet:

$$\text{Mesons:} \qquad \Pi_{\mathbf{1}}^{(\mathbf{3}\otimes\bar{\mathbf{3}})} : \ \mathbf{3} \otimes \bar{\mathbf{3}} \to \mathbf{1}, \quad \text{e.g. } \Pi_{\mathbf{1}}^{(\mathbf{3}\otimes\bar{\mathbf{3}})}\big(q^i \otimes \bar{q}_j\big) \ \propto \ \delta^i{}_j, \tag{12.228}$$

$$\text{Baryons:} \qquad \Pi_{\mathbf{1}}^{(\mathbf{3}\otimes\mathbf{3}\otimes\mathbf{3})} : \ \mathbf{3}^{\otimes 3} \to \mathbf{1}, \quad \text{e.g. } \Pi_{\mathbf{1}}^{(\mathbf{3}^{\otimes 3})}\big(q^i \otimes q^j \otimes q^k\big) \ \propto \ \epsilon_{ijk}. \tag{12.229}$$

The anti–meson/anti–baryon projectors are obtained by complex conjugation of the invariant tensors, sending $\delta^i{}_j \mapsto \delta_i{}^j$ and $\epsilon_{ijk} \mapsto \epsilon^{ijk}$, respectively. Hence color–singlet operators are (up to normalization)

$$\mathcal{O}_M \sim \bar{q}_i q^i, \qquad \mathcal{O}_{\overline{M}} \sim q^i \bar{q}_i, \qquad \mathcal{O}_B \sim \epsilon_{ijk} q^i q^j q^k, \qquad \mathcal{O}_{\overline{B}} \sim \epsilon^{ijk} \bar{q}_i \bar{q}_j \bar{q}_k.$$

Step 2 (BRST cohomology: physical singlets, non–singlets discarded). By Proposition 12.120, color–singlet composite operators are BRST–closed and represent nontrivial classes in $H^0(\hat{Q}_{\mathrm{BRST}})$, while color–non–singlet composites are BRST–exact (or decouple) and therefore vanish in cohomology. More generally, Theorem 12.133 identifies the physical hadron Hilbert space within $\mathcal{H}_{\mathrm{phys}}$ with the BRST cohomology of color–singlet composites obtained by the projectors (12.228)–(12.229). Consequently, only singlet combinations survive as physical states; all non–singlets are discarded.

Step 3 (Conjugation to anti–hadrons). The CRP charge–conjugation map (Definition 12.151) sends $q \mapsto q^c \in \bar{\mathbf{3}}$ and $\bar{q} \mapsto \bar{q}^c \in \mathbf{3}$. Because the singlet projectors are built from invariant tensors (δ, ϵ), they are preserved under conjugation (up to index placement), i.e. $\Pi_{\mathbf{1}}^{(\mathbf{3}\otimes\bar{\mathbf{3}})} \mapsto \Pi_{\mathbf{1}}^{(\bar{\mathbf{3}}\otimes\mathbf{3})}$ and $\Pi_{\mathbf{1}}^{(\mathbf{3}^{\otimes 3})} \mapsto \Pi_{\mathbf{1}}^{((\bar{\mathbf{3}})^{\otimes 3})}$. Therefore,

$$\mathcal{O}_M \xrightarrow{C} \mathcal{O}_{\overline{M}}, \qquad \mathcal{O}_B \xrightarrow{C} \mathcal{O}_{\overline{B}},$$

and both sides remain color singlets, hence BRST–physical by Step 2. At the level of poles/residues, Proposition 12.154 guarantees that a hadron pole is accompanied by a conjugate anti–hadron pole with the same (m, s) and opposite additive charges.

Combining Steps 1–3 proves that color–singlet projection together with BRST cohomology produces hadrons and anti–hadrons precisely as conjugate singlets (mesons $q\bar{q}$ vs. $\bar{q}q$; baryons $\epsilon_{ijk} q^i q^j q^k$ vs. $\epsilon^{ijk} \bar{q}_i \bar{q}_j \bar{q}_k$) and discards all non–singlets, as claimed in Definition 12.126, Proposition 12.120, and Theorem 12.133. □

Antimatter pairing is *not* an extra axiom: it follows from (i) the $*$–structure enforced by complex internal reps (Definition 12.151), (ii) diagnostic realized reference configuration locality/positivity implying CPT and conjugate poles (Proposition 12.154), and (iii) neutrality of the family fibre with holonomy phases that conjugate under C (Definition 12.156). Where any ingredient trivializes (real/pseudoreal reps, neutral fields, Majorana condition), the pair collapses and the field becomes self–conjugate (Proposition 12.158).

Remark 12.160 (Roadmap to the Duality formalization of antimatter)**.** The three pillars in §12.9 are precise incarnations of the Axiom of Duality (Definition 2.100):

(i) Representation duality
$R \leftrightarrow \bar{R}$ (CRP, Definition 12.151) is the algebraic Thesis/Antithesis on the internal connection; Proposition 12.152 states the corresponding closure of the action/flow on the diagnostic realized reference configuration.

(ii) Spectral duality
of poles/residues (CPT pairing, Proposition 12.154) is the dynamical dual lifted to the one–particle spectrum (locality + positivity on $\mathfrak{P}$).

(iii) Holonomy duality
$U^{(F)}[\gamma^{-1}] = U^{(F)}[\gamma]^\dagger$ (family phases conjugate under C, Definition 12.156) realizes path/orientation duality on the family fibre.

In Section 12.10, we package these as a single statement:
Duality ⇒ Antiparticles (typed). On a certified plateau $\mathfrak{P}$, for any Tantrakyma kymon in a non-self-conjugate internal irrep, Duality induces a unique conjugate class with the same mass/spin and opposite additive charges; family phases are complex-conjugated and bound-state singlets are mapped to their conjugates. Conversely, when the irrep is real/pseudoreal or the relevant charge vanishes (or in the Majorana limit), the Duality acts trivially and the field is self-conjugate.

This provides the exact bridge from the structural pillars to the Duality-theoretic statement used elsewhere in the tetralemmic weave (Sections 6.11 and 6.11.6).

Building on the structural pillars of §12.9, we now package the algebraic, geometric, and spectral ingredients into a single, typed diagnostic realized reference configuration-level statement. The new subsection §12.10 formalizes antimatter as a direct corollary of the Axiom of Duality, by composing Definitions 12.151 and 12.156, Propositions 12.152 and 12.154, and Corollary 12.159 without introducing any new hypotheses.

12.10 Antimatter as a Duality Theorem

This section gives a *duality-first* foundation of antimatter on the certified plateau $\mathfrak{P}$ by instantiating the Axiom of Duality of the tetralemma layer (Axiom 1.10) in the Standard–Model weave. We compose: (i) complex–representation pairing (CRP), (ii) diagnostic realized reference configuration CPT pairing of poles, (iii) family–holonomy conjugation, (iv) Källén–Lehmann (KL) positivity on $\mathfrak{P}$, and (v) BRST color–singlet cohomology, to obtain an involutive duality functor and a typed antimatter theorem. All objects and maps referenced below are already defined in Section 12.4.4, Definitions 12.151 and 12.156, Propositions 12.152, 12.154 and 12.158, and Corollary 12.159.

12.10.1 Categorical Duality on Matter (Algebraic, Kinematic, and Cohomological Faces)

Definition 12.161 (Diagnostic realized reference configuration duality datum)**.** A *diagnostic realized reference configuration duality datum* on $\mathfrak{P}$ is a triple

$$\mathfrak{D}_{\mathfrak{P}} \;=\; \Big(\mathfrak{D}_{\mathrm{int}},\ \mathfrak{D}_{\mathrm{kin}},\ \mathfrak{D}_{\mathrm{fam}}\Big)$$

consisting of:

(i) Internal (algebraic) duality
$\mathfrak{D}_{\mathrm{int}} : \mathrm{Rep}(G_{\mathrm{SM}}) \to \mathrm{Rep}(G_{\mathrm{SM}})$, the contravariant *–functor

$$\mathfrak{D}_{\mathrm{int}}(R) = \bar{R}, \qquad \mathfrak{D}_{\mathrm{int}}\Big(U : R \to S\Big) = U^* : \bar{S} \to \bar{R},$$

(where U^{*} denotes the conjugate–linear transpose appropriate to intertwiners; in a Hilbert model it coincides with the adjoint on matrix elements). At the field level CRP holds: for a fermionic Tantrakyma kymon $\psi \in R$, the charge–conjugate field $\psi^{c} := C\,\bar{\psi}^{\top}$ transforms in $\bar{R}$ (Definition 12.151 and Proposition 12.152).

(ii) Kinematic duality

$\mathfrak{D}_{\mathrm{kin}}$ on one–particle poles: on $\mathfrak{P}$,

$$(m, s;\, \mathsf{q}) \;\mapsto\; (m, s;\, -\mathsf{q}),$$

$$\mathrm{Res}_{p^2\to m^2}\, \mathbf{G}_F^{(R)}(p) = Z\,\Pi_{(m,s)}(p) \quad\Longleftrightarrow\quad \mathrm{Res}_{p^2\to m^2}\, \mathbf{G}_F^{(\bar{R})}(p) = Z\,\Pi_{(m,s)}(p).$$

with the correspondence implemented by the antiunitary Θ_{CPT} (eqs. (12.230) to (12.233) and Proposition 12.154).

(iii) Family duality

$\mathfrak{D}_{\mathrm{fam}}$ on the auxiliary family bundle $(\mathcal{F}, \mathcal{A}^{(F)})$:

$$U^{(F)}[\gamma] \;\mapsto\; U^{(F)}[\gamma]^{*} \qquad \text{(phase conjugation / holonomy inversion)},$$

equivalently $U^{(F)}[\gamma^{-1}] = U^{(F)}[\gamma]^{\dagger}$ (Definitions 12.106 and 12.156 and eq. (12.124)).

Definition 12.162 (Diagnostic realized reference configuration duality functor on species)**.** Let $\mathsf{Sp}_{\mathfrak{P}}$ be the category whose objects are typed one–particle *species*

$$\mathsf{S} \equiv \big(R,\; m,\; s,\; \mathsf{q};\; \Pi^{[\chi]}\big)$$

specified by an internal irrep R, on–shell invariants (m, s), additive–charge vector q, and a diagnostic realized reference configuration type–projector $\Pi^{[\chi]}$ (e.g. TT, scalar–τ, vector–τ; Definitions 10.3 and 10.4), and whose morphisms are diagnostic realized reference configuration–admissible intertwiners preserving (m, s) and commuting with the typed realized weave configuration. The *diagnostic realized reference configuration duality functor*

$$\mathfrak{D} :\; \mathsf{Sp}_{\mathfrak{P}} \longrightarrow \mathsf{Sp}_{\mathfrak{P}}$$

acts by

$$\mathfrak{D}\colon\; (R, m, s, \mathsf{q}; \Pi^{[\chi]}) \longmapsto (\bar{R},\; m,\; s,\; -\mathsf{q}; \Pi^{[\chi]}),$$

and on morphisms by adjoint–conjugation compatible with $\mathfrak{D}_{\mathrm{int}}, \mathfrak{D}_{\mathrm{kin}}, \mathfrak{D}_{\mathrm{fam}}$. Concretely, for a morphism $f : (R, m, s, \mathsf{q}; \Pi^{[\chi]}) \to (R', m, s, \mathsf{q}; \Pi^{[\chi]})$ we set

$$\mathfrak{D}(f) \;=\; f^{*} : (\bar{R}', m, s, -\mathsf{q}; \Pi^{[\chi]}) \;\longrightarrow\; (\bar{R}, m, s, -\mathsf{q}; \Pi^{[\chi]}),$$

so that $\mathfrak{D}(\mathbf{1}) = \mathbf{1}$ and $\mathfrak{D}(g \circ f) = \mathfrak{D}(f) \circ \mathfrak{D}(g)$.

Remark 12.163 (Involutivity and diagnostic realized reference configuration hygiene)**.** By Axiom 1.10 and the CPT antiunitarity, each component of $\mathfrak{D}_{\mathfrak{P}}$ is an involution; consequently $\mathfrak{D}^{2} = \mathrm{Id}$ on $\mathsf{Sp}_{\mathfrak{P}}$. Moreover, on $\mathfrak{P}$ the KL measure is positive and poles are gauge–parameter independent (Proposition 12.88), so $\mathfrak{D}$ preserves pole locations, residues $Z \geq 0$, and diagnostic realized reference configuration typed projectors (e.g. TT vs. τ–blocks; Definitions 10.3 and 10.4).

Remark 12.164 (Monoidal compatibility)**.** $\mathfrak{D}_{\mathrm{int}}$ preserves direct sums and tensor products up to canonical isomorphism: $\overline{R \oplus S} \simeq \bar{R} \oplus \bar{S}$ and $\overline{R \otimes S} \simeq \bar{R} \otimes \bar{S}$.

Definition 12.165 (Dual completion and dual–closure)**.** Given a field/species set $\mathcal{S} \subset \mathrm{Ob}(\mathsf{Sp}_{\mathfrak{P}})$, its *dual completion* is

$$\overline{\mathcal{S}}^{\mathfrak{D}} \;:=\; \text{the smallest full subcategory containing } \mathcal{S} \text{ and } \mathfrak{D}(\mathcal{S}).$$

We say $\mathcal{S}$ is *dual–closed* if $\mathfrak{D}(\mathcal{S}) = \mathcal{S}$.

12.10.2 Antimatter as a Duality Theorem

Theorem 12.166 (Antimatter Duality Theorem)**.** *On a certified plateau $\mathfrak{P}$, with CRP, diagnostic realized reference configuration CPT pairing of poles, family–holonomy conjugation, KL positivity, and BRST color–singlet cohomology in force (Definitions 12.151 and 12.156, Propositions 12.88, 12.120 and 12.154, and Theorem 12.133), the diagnostic realized reference configuration duality functor $\mathfrak{D} : \mathsf{Sp}_{\mathfrak{P}} \to \mathsf{Sp}_{\mathfrak{P}}$ is an involutive autoequivalence and the physical spectrum is* dual–closed*: for every species $(R, m, s, \mathsf{q}; \Pi^{[\chi]})$ there exists the conjugate species $(\bar{R}, m, s, -\mathsf{q}; \Pi^{[\chi]})$. Equivalently, the physical observable algebra on $\mathfrak{P}$ is $*$–closed* iff *it is $\mathfrak{D}$–closed (Proposition 12.152).*

Proof. We proceed in four steps.

Step I (Construction of $\mathfrak{D}$ on objects and morphisms). By Definition 12.161, the diagnostic realized reference configuration duality datum $\mathfrak{D}_{\mathfrak{P}} = (\mathfrak{D}_{\mathrm{int}}, \mathfrak{D}_{\mathrm{kin}}, \mathfrak{D}_{\mathrm{fam}})$ has:

1. *Internal (algebraic) duality* $\mathfrak{D}_{\mathrm{int}} : \mathrm{Rep}(G_{\mathrm{SM}}) \to \mathrm{Rep}(G_{\mathrm{SM}})$,

$$\mathfrak{D}_{\mathrm{int}}(R) = \bar{R}, \qquad \mathfrak{D}_{\mathrm{int}}\big(U : R \to S\big) = U^* : \bar{S} \to \bar{R},$$

realized at field level by CRP, $\psi^c = C\,\bar{\psi}^{\top} \in \bar{R}$ (Definition 12.151 and Proposition 12.152).

2. *Kinematic duality* (diagnostic realized reference configuration CPT) on one–particle poles,

$$(m, s; \mathsf{q}) \;\mapsto\; (m, s; -\mathsf{q}),$$
$$\mathrm{Res}_{p^2 \to m^2}\, \mathbf{G}_F^{(R)}(p) = Z\,\Pi_{(m,s)}(p) \iff \mathrm{Res}_{p^2 \to m^2}\, \mathbf{G}_F^{(\bar{R})}(p) = Z\,\Pi_{(m,s)}(p),$$

implemented by the antiunitary Θ_{CPT} (eqs. (12.230) to (12.233) and Proposition 12.154).

3. *Family duality* on $(\mathcal{F}, \mathcal{A}^{(F)})$,

$$U^{(F)}[\gamma] \mapsto U^{(F)}[\gamma]^*,$$
$$U^{(F)}[\gamma^{-1}] = U^{(F)}[\gamma]^{\dagger}.$$

(See Definitions 12.106 and 12.156 and eq. (12.124).)

Define $\mathfrak{D}$ on species objects $\mathsf{S} = (R, m, s, \mathsf{q}; \Pi^{[\chi]}) \in \mathsf{Sp}_{\mathfrak{P}}$ by

$$\mathfrak{D}(\mathsf{S}) = (\bar{R}, m, s, -\mathsf{q}; \Pi^{[\chi]}),$$

which preserves the diagnostic realized reference configuration type $\Pi^{[\chi]}$ by Remark 12.163 (KL/Nielsen hygiene and typed–projector stability). For morphisms $f : \mathsf{S} \to \mathsf{S}'$ with fixed (m, s) and type, set

$$\mathfrak{D}(f) = f^* : \ \mathfrak{D}(\mathsf{S}') \longrightarrow \mathfrak{D}(\mathsf{S}).$$

Then $\mathfrak{D}(\mathbf{1}) = \mathbf{1}$ and $\mathfrak{D}(g \circ f) = \mathfrak{D}(f) \circ \mathfrak{D}(g)$, so $\mathfrak{D}$ is a (contravariant) functor $\mathsf{Sp}_{\mathfrak{P}} \to \mathsf{Sp}_{\mathfrak{P}}$ compatible with $(\mathfrak{D}_{\mathrm{int}}, \mathfrak{D}_{\mathrm{kin}}, \mathfrak{D}_{\mathrm{fam}})$ (Definition 12.162).

Step II (Involutivity: $\mathfrak{D}^2 = \mathrm{Id}$). Each component of $\mathfrak{D}_{\mathfrak{P}}$ is involutive: $(\bar{\bar{R}}) \cong R$; CPT applied twice returns the original labels $(m, s; \mathsf{q})$; and family holonomies satisfy $\left(U^{(F)}[\gamma]^*\right)^* = U^{(F)}[\gamma]$. Hence $\mathfrak{D}^2$ acts trivially on objects. On morphisms, $(f^*)^* = f$ (conjugate–transpose involutivity), so $\mathfrak{D}^2 = \mathrm{Id}$ strictly (Remark 12.163).

Step III (Autoequivalence: full, faithful, essentially surjective). *Faithful:* if $\mathfrak{D}(f) = \mathfrak{D}(g)$ then $f^* = g^*$, hence $f = g$. *Full:* given $h : \mathfrak{D}(\mathsf{S}') \to \mathfrak{D}(\mathsf{S})$, set $f := h^*$, so $\mathfrak{D}(f) = h$. *Essentially surjective:* for any T, $\mathsf{T} = \mathfrak{D}^2(\mathsf{T}) = \mathfrak{D}(\mathfrak{D}(\mathsf{T}))$, hence T lies in the essential image. Thus $\mathfrak{D}$ is an involutive autoequivalence of $\mathsf{Sp}_{\mathfrak{P}}$.

Step IV (Dual–closure and the $*$–closure criterion). Because $\mathfrak{D}$ is an autoequivalence, every species admits a conjugate species: $(R, m, s, \mathsf{q}; \Pi^{[\chi]}) \mapsto (\bar{R}, m, s, -\mathsf{q}; \Pi^{[\chi]})$; the spectrum is dual–closed.

For the observable algebra, we show "$*$–closed $\iff$ $\mathfrak{D}$–closed." ($\Rightarrow$) If the algebra is $*$–closed, then for each R–valued field ψ its conjugate $\psi^c = C\,\bar{\psi}^\top$ and the corresponding $\bar{R}$ creation/annihilation modes lie in the algebra (Definition 12.151). On $\mathfrak{P}$, KL/Nielsen guarantees that physical pole data are gauge–parameter independent (Proposition 12.88); CPT then maps $(m, s; \mathsf{q})$ to $(m, s; -\mathsf{q})$ and preserves the residue projector (Proposition 12.154). Therefore the induced mapping of species is exactly $\mathfrak{D}$, so the spectrum and the algebra are $\mathfrak{D}$–closed.

($\Leftarrow$) Conversely, if the spectrum/algebra is $\mathfrak{D}$–closed, then for each R–sector the $\bar{R}$ sector is present with the same (m, s) data; by Proposition 12.152 the local density is $*$–closed (the interaction terms and currents built from $R \oplus \bar{R}$ are Hermitian), so the action is real and the Hamiltonian self–adjoint. Thus $*$–closure holds.

Finally, BRST color–singlet cohomology is respected: color–singlet projectors commute with both BRST and duality (eq. (12.239)), so bound–state singlets map to conjugate singlets inside $H^0(\hat{Q}_{\mathrm{BRST}})$ (Proposition 12.120 and Theorem 12.133). Together, the four steps complete the proof. □

12.10.3 Operational Corollaries

Corollary 12.167 (Conjugate residues and pair–aware amplitudes)**.** *For any isolated pole at $p^2 = m^2$ with residue $Z\,\Pi_{(m,s)}(p)$ in the R channel, there is a conjugate pole with the same residue in the $\bar{R}$ channel (eq. (12.233)). Moreover, LSZ + analyticity on $\mathfrak{P}$ give*

$$\mathcal{M}(\cdots \to out + in(p, \mathsf{q})) = \Big[\mathcal{M}(\cdots + out(\tilde{p}, -\mathsf{q}) \to out)\Big]^{\text{analytic cont.}}_{\tilde{p} \to -p},$$

so amplitudes are pair–aware (eq. (12.235)).

Corollary 12.168 (Color singlets and conjugate hadrons)**.** *Color–singlet projectors commute with BRST and with $\mathfrak{D}$ (eq. (12.239)); thus meson and baryon operators map to their conjugates inside $H^0(\hat{Q}_{\mathrm{BRST}})$,*

$$C: \ \mathcal{O}_M \leftrightarrow \mathcal{O}_{\overline{M}}, \qquad \mathcal{O}_B \leftrightarrow \mathcal{O}_{\overline{B}},$$

as in Corollary 12.159.

Corollary 12.169 (Self–conjugacy criterion (Majorana/real cases))**.** *If $R \simeq \bar{R}$ (real or pseudoreal), then $\mathfrak{D}$ fixes the internal label and antiparticles may be realized as self–conjugate excitations (Majorana–like for fermions), with the field–level constraint controlled by the typed projectors and CRP (Proposition 12.158).*

Remark 12.170 (Typed compatibility)**.** $\mathfrak{D}$ preserves diagnostic realized reference configuration types: the TT projector annihilates τ–blocks and vice versa (Definitions 10.3 and 10.4); hence duality pairs species *within* their kinematic type, e.g. (TT spin-2; q) $\leftrightarrow$ (TT spin-2; $-\mathsf{q}$), while non–TT (e.g. Chrono–Kymon) sectors pair analogously inside their typed block.

Remark 12.171 (Family phases and CP structure)**.** Under $\mathfrak{D}_{\mathrm{fam}}$, holonomy phases conjugate, so CKM/PMNS entries transform by complex conjugation (Proposition 12.115 and eq. (12.134)). This isolates genuine CP–odd rephasing invariants (e.g. Jarlskog–type) as duality–invariant obstructions, consistent with the diagnostic realized reference configuration CPT structure.

The diagnostic realized reference configuration duality data formalized in Definition 12.161 packages three faces of the same operation. On the *internal/algebraic* side, $\mathfrak{D}_{\mathrm{int}}$ is nothing but passage to the complex–conjugate representation: it sends any irrep $R \subset G_{\mathrm{SM}}$ to $\bar{R}$ and any intertwiner U to U^*. At the field level this is precisely the complex–representation pairing (CRP): a fermionic Tantrakyma kymon $\psi \in R$ has a charge–conjugate $\psi^c = C\,\overline{\psi}^{\top} \in \bar{R}$ (Definition 12.151 and Proposition 12.152). On the *kinematic/spectral* side, $\mathfrak{D}_{\mathrm{kin}}$ encodes the diagnostic realized reference configuration CPT content by acting on one–particle data as $(m, s; \mathsf{q}) \mapsto (m, s; -\mathsf{q})$, while preserving the pole position and residue projector: $\operatorname{Res} \mathbf{G}_F^{(R)} = Z\,\Pi_{(m,s)} \Longleftrightarrow \operatorname{Res} \mathbf{G}_F^{(\bar{R})} = Z\,\Pi_{(m,s)}$ (eqs. (12.230) to (12.233) and Proposition 12.154). Finally, on the *family/phase* side, the auxiliary bundle $(\mathcal{F}, \mathcal{A}^{(F)})$ carries only phases; duality is phase conjugation of holonomies, $U^{(F)}[\gamma] \mapsto U^{(F)}[\gamma]^*$ with $U^{(F)}[\gamma^{-1}] = U^{(F)}[\gamma]^{\dagger}$ (Definitions 12.106 and 12.156 and eq. (12.124)). Each face is involutive, in harmony with the Axiom of Duality in the tetralemma layer and with the antiunitarity of CPT.

The *diagnostic realized reference configuration duality functor* $\mathfrak{D}$ of Definition 12.162 then acts on "species," i.e. typed one–particle labels $\mathsf{S} = (R, m, s, \mathsf{q}; \Pi^{[\chi]})$ recording the internal irrep, kinematic invariants, the additive–charge vector, and the diagnostic realized reference configuration type projector (TT versus τ blocks, etc.; see Definitions 10.3 and 10.4). Concretely, $\mathfrak{D}$ leaves the kinematics and the diagnostic realized reference configuration type untouched and flips only the internal data:

$$(R, m, s, \mathsf{q}; \Pi^{[\chi]}) \longmapsto (\bar{R}, m, s, -\mathsf{q}; \Pi^{[\chi]}).$$

This is exactly "send a particle to its antiparticle" without altering mass, spin, or which typed meter sees it. Because each component datum is involutive, we have $\mathfrak{D}^2 = \mathrm{Id}$ (Remark 12.163).

The language of *dual completion/dual–closure* in Definition 12.165 captures a physical necessity: demanding a real action and a self–adjoint Hamiltonian forces the field/species content to be closed under duality. If a chosen set is not $\mathfrak{D}$–closed, the $*$–operation takes the local field algebra outside the set; Hermiticity and unitary time evolution then fail. This is precisely the import of Proposition 12.152: reality and unitarity on the certified plateau are equivalent to $R \oplus \bar{R}$ closure.

With these ingredients, the Antimatter Duality Theorem Theorem 12.166 is not an extra postulate but an outcome: CRP (algebra), diagnostic realized reference configuration CPT (kinematics), family conjugation (phases), KL/Nielsen hygiene (positivity and gauge–parameter independence), and BRST color–singlet cohomology (physical composites) together make

$$\mathfrak{D} : \mathsf{Sp}_{\mathfrak{P}} \longrightarrow \mathsf{Sp}_{\mathfrak{P}}$$

an involutive autoequivalence. Consequently, the physical spectrum is *dual–closed*: every species admits a conjugate partner with $(\bar{R}, -\mathsf{q})$ and the same (m, s) and diagnostic realized reference configuration type.

The operational corollaries align with standard expectations. First, residues match across $R \leftrightarrow \bar{R}$ (eq. (12.233)), and LSZ plus analyticity on $\mathfrak{P}$ identify an incoming particle with an outgoing antiparticle via crossing, so amplitudes are "pair–aware" by construction (Corollary 12.167 and eq. (12.235)). Second, because color–singlet projectors commute with BRST and with $\mathfrak{D}$ (eq. (12.239)), mesons and baryons map to their conjugates inside the physical cohomology $H^0(\hat{Q}_{\mathrm{BRST}})$ (Corollaries 12.159 and 12.168). Third, in real or pseudoreal irreps ($R \simeq \bar{R}$), the duality fixes the internal label and self–conjugate excitations may be realized (Majorana–like in the fermionic case), subject to the usual field–level constraints (Corollary 12.169 and Proposition 12.158). Finally, on the family side, $\mathfrak{D}_{\mathrm{fam}}$ complex–conjugates CKM/PMNS entries (Proposition 12.115 and eq. (12.134)), isolating CP–odd rephasing invariants (Jarlskog–type) as genuine, duality–invariant obstructions (Remark 12.171).

Viewed against the tetralemma–GNS background, this construction is the diagnostic realized reference configuration–level realization of the Axiom of Duality (Axiom 1.10). The object–level dualization $R \mapsto \bar{R}$ and $\psi \mapsto \psi^c$ (CRP), the kinematic mapping $(m, s; \mathsf{q}) \mapsto (m, s; -\mathsf{q})$ with equal residues (diagnostic realized reference configuration CPT), and the phase conjugation $U^{(F)} \mapsto U^{(F)*}$ (family holonomy) glue into the species–level functor $\mathfrak{D}$. Because $\mathfrak{D}$ is involutive and spectrum–closing, "antimatter" appears as a theorem once KL–positivity, BRST physicality, and the operator–bundle realized weave configuration are imposed on the certified plateau.

12.10.4 Diagnostic realized reference configuration Structure that Forces Pairing (Spectral/Cohomological Face)

Diagnostic realized reference configuration CPT pairing (expanded). On a certified plateau $\mathfrak{P}$ (Definition 9.185), two–point functions admit a Källén–Lehmann (KL) representation with nonnegative spectral measure and gauge–parameter–independent poles (Proposition 12.88).

For a scalar kymon (spin $s = 0$) the momentum–space time–ordered two–point function reads

$$G_F(p) \;=\; \frac{Z}{p^2 - m^2 + i\epsilon} \;+\; \int_0^\infty d\mu^2 \, \frac{\rho(\mu^2)}{p^2 - \mu^2 + i\epsilon}, \qquad Z \geq 0, \quad \rho(\mu^2) \geq 0, \tag{12.230}$$

and the isolated one–particle excitation is the simple pole at $p^2 = m^2$ with residue Z. For higher spin s, the same structure holds with a covariant numerator projector $\Pi_{(m,s)}(p)$:

$$\mathbf{G}_F(p) \;=\; \frac{Z\,\Pi_{(m,s)}(p)}{p^2 - m^2 + i\epsilon} \;+\; \int_0^\infty d\mu^2 \, \frac{\rho_{(s)}(\mu^2)\,\Pi_{(\mu,s)}(p)}{p^2 - \mu^2 + i\epsilon}, \tag{12.231}$$

where positivity still implies $Z \geq 0$ and $\rho_{(s)}(\mu^2) \geq 0$ (Proposition 12.88). Locality (microcausality) and Lorentz covariance on $\mathfrak{P}$ guarantee the existence of an antiunitary CPT map Θ_{CPT} acting on the one–particle sector. For a charged kymon (nonzero additive charge q) in a genuinely complex irrep $R \subset G_{\mathrm{SM}}$, the CPT action combines with the complex–representation pairing (CRP, Definition 12.151) to yield

$$\Theta_{\mathrm{CPT}} \;:\; |p, s; \mathsf{q}\rangle_R \;\longmapsto\; e^{i\alpha}\, |p, s; -\mathsf{q}\rangle_{\bar{R}}, \qquad p^2 = m^2, \tag{12.232}$$

for a pure phase $e^{i\alpha}$ fixed by conventions. At the level of residues this implies

$$\begin{aligned} \mathrm{Res}_{p^2 \to m^2}\, \mathbf{G}_F^{(R)}(p) \;&=\; Z\,\Pi_{(m,s)}(p) \quad \Longrightarrow \\ \mathrm{Res}_{p^2 \to m^2}\, \mathbf{G}_F^{(\overline{R})}(p) \;&=\; \Theta_{\mathrm{CPT}} \Big[Z\,\Pi_{(m,s)}(p) \Big] \Theta_{\mathrm{CPT}}^{-1} \\ &=\; Z\,\Pi_{(m,s)}(p), \end{aligned} \tag{12.233}$$

i.e. the same (m, s) pole with conjugate charge, as stated in Proposition 12.154. Equivalently, at the field level one may use $\psi^c = C\,\bar{\psi}^{\,\top}$ (Definition 12.151) to write

$$\begin{aligned} &\mathrm{Res}_{p^2 \to m^2}\, \langle 0 |\, T\{\psi(x)\bar{\psi}(0)\}\, |0\rangle \;\sim\; Z\,\Pi_{(m,s)}(p) \\ &\quad\Longleftrightarrow\quad \mathrm{Res}_{p^2 \to m^2}\, \langle 0 |\, T\{\psi^c(x)\overline{\psi^c}(0)\}\, |0\rangle \;\sim\; Z\,\Pi_{(m,s)}(p). \end{aligned} \tag{12.234}$$

Operationally, LSZ reduction on $\mathfrak{P}$ (locality + unitarity; cf. Proposition 12.152) gives the *crossing* relation:

$$\mathcal{M}\Big(\cdots \to \mathrm{out} + \mathrm{in}(p, \mathsf{q}) \Big) \;=\; \Big[\mathcal{M}\Big(\cdots + \mathrm{out}(\tilde{p}, -\mathsf{q}) \to \mathrm{out} \Big) \Big]^{\text{analytic cont.}}_{\tilde{p} \to -p}, \tag{12.235}$$

so amplitudes are pair–aware by construction, in agreement with Proposition 12.154.

What Θ_{CPT} denotes (and what it does not denote). The symbol Θ_{CPT} is *not* the Tantrakyma field $\hat{\boldsymbol{\Theta}}$. Rather, it denotes the *antiunitary* operator that implements the combined Charge, Parity, and Time-reversal transformation on the (BRST-)physical Hilbert space of the certified plateau $\mathfrak{P}$ (see Definition 9.185). As an antiunitary map it is anti-linear and norm-preserving: for $a, b \in \mathbb{C}$,

$$\Theta_{\mathrm{CPT}}\Big(a|\psi\rangle + b|\phi\rangle \Big) = a^*\, \Theta_{\mathrm{CPT}}|\psi\rangle + b^*\, \Theta_{\mathrm{CPT}}|\phi\rangle, \qquad \langle \Theta_{\mathrm{CPT}}\psi \mid \Theta_{\mathrm{CPT}}\phi \rangle = \langle \phi \mid \psi \rangle,$$

and it flips the phase of the imaginary unit: $\Theta_{\mathrm{CPT}}\, i\, \Theta_{\mathrm{CPT}}^{-1} = -\,i$. This operator acts on one-particle states in the manner summarized below and is conceptually distinct from the field operator $\hat{\boldsymbol{\Theta}}$.

Diagnostic realized reference configuration CPT pairing (spectral statement). On a certified plateau $\mathfrak{P}$, locality (microcausality) and Lorentz covariance ensure that two–point functions admit a Källén–Lehmann (KL) representation with nonnegative spectral measure and gauge–parameter–independent pole structure (Proposition 12.88). For scalars the KL form is given in eq. (12.230); for higher spin s the covariant projector $\Pi_{(m,s)}(p)$ appears as in eq. (12.231). In particular, if a kymon carries a nonzero additive charge q and transforms in a genuinely complex irrep $R \subset G_{\mathrm{SM}}$, then the antiunitary Θ_{CPT} combines with the complex–representation pairing (CRP, Definition 12.151) to map a one–particle state to its antiparticle partner in the conjugate irrep:

$$\Theta_{\mathrm{CPT}} : \ |p, s; \mathsf{q}\rangle_R \ \longmapsto \ e^{i\alpha} \, |p, s; -\mathsf{q}\rangle_{\bar{R}}, \qquad p^2 = m^2,$$

as stated in eq. (12.232). This is the *diagnostic realized reference configuration CPT pairing*: the mass m and spin s are identical for the particle–antiparticle pair, while the irrep and additive charge are conjugated.

Residues and field-level phrasing. The pairing manifests spectrally in the equality of pole residues for R and $\bar{R}$,

$$\mathrm{Res}_{p^2 \to m^2} \, \mathbf{G}_F^{(R)}(p) = \mathrm{Res}_{p^2 \to m^2} \, \mathbf{G}_F^{(\bar{R})}(p) = Z \, \Pi_{(m,s)}(p),$$

as summarized in eq. (12.233). Equivalently at the field level, charge conjugation $\psi^c = C \, \bar{\psi}^{\top}$ (the CRP map of Definition 12.151) ensures that the on-shell residues of the time-ordered two–point functions coincide for ψ and its charge-conjugate ψ^c, exactly as recorded in eq. (12.234).

Crossing and "pair-aware" amplitudes on the diagnostic realized reference configuration. With LSZ reduction on $\mathfrak{P}$ (locality + unitarity; see Proposition 12.152), scattering amplitudes obey the standard crossing relation,

$$\mathcal{M}\Big(\cdots \to \mathrm{out} + \mathrm{in}(p, \mathsf{q}) \Big) = \Big[\mathcal{M}\Big(\cdots + \mathrm{out}(\tilde{p}, -\mathsf{q}) \to \mathrm{out} \Big) \Big]^{\text{analytic cont.}}_{\tilde{p} \to -p},$$

as stated in eq. (12.235). Operationally, an incoming particle with quantum numbers (p, q) is analytically equivalent to an outgoing antiparticle with $(-p, -\mathsf{q})$; the S–matrix is therefore *pair–aware by construction.*

Why "spectral / cohomological face". The argument is *spectral* because it rests on the KL spectral density and the residue at the physical pole (positivity $Z \geq 0$, $\rho \geq 0$, and gauge–independent pole positions; Proposition 12.88). It is simultaneously *cohomological* because everything is evaluated on the certified plateau $\mathfrak{P}$, i.e. in the BRST cohomology $H^0(\hat{Q}_{\mathrm{BRST}})$, so only physical (cohomology–nontrivial) states contribute.

Immediate consequences. First, particle–antiparticle partners in R and $\bar{R}$ necessarily have equal masses and spins and share the same residue Z and projector $\Pi_{(m,s)}$. Second, their additive charges flip sign under CPT+CRP, $\mathsf{q} \to -\mathsf{q}$. Third, amplitudes with incoming particles are related by crossing to amplitudes with outgoing antiparticles via $p \to -p$ (analytic continuation), eq. (12.235). Finally, in special cases where $R \simeq \bar{R}$ (real or pseudoreal representations), the pairing can collapse to self–conjugate excitations; the diagnostic realized reference configuration logic then reduces to the familiar Majorana–like or adjoint cases, still organized by the same projector $\Pi_{(m,s)}$ and CRP map.

BRST doublets and color–singlet cohomology. Let $\hat{Q}_{\mathrm{BRST}}$ be the nilpotent BRST charge ($\hat{Q}^2_{\mathrm{BRST}} = 0$) on $\mathfrak{P}$, and $H^0(\hat{Q}_{\mathrm{BRST}})$ the physical cohomology. The *doublet lemma* (Kugo–Ojima mechanism) states that any pair (U, V) with

$$[\hat{Q}_{\mathrm{BRST}}, U] = V, \qquad [\hat{Q}_{\mathrm{BRST}}, V] = 0 \tag{12.236}$$

does not contribute to $H^0(\hat{Q}_{\mathrm{BRST}})$; unphysical modes assemble into such doublets and decouple. For QCD color, physical states satisfy the subsidiary condition $G^a|\mathrm{phys}\rangle = 0$ (Gauss law), i.e. they are *color singlets*. Let q^i be a quark–kymon in $\mathbf{3}$ of $SU(3)_C$ and $\bar{q}_i$ its CRP partner in $\bar{\mathbf{3}}$ (Definition 12.151). The invariant projectors of Definition 12.126 select the singlet subspace:

$$\Pi_{\mathbf{1}}^{(\mathbf{3}\otimes\bar{\mathbf{3}})}\big(q^i \otimes \bar{q}_j\big) = \frac{1}{N_c}\,\delta^i{}_j\ (\bar{q}_k q^k), \tag{12.237}$$

$$\Pi_{\mathbf{1}}^{(\mathbf{3}^{\otimes 3})}\big(q^i \otimes q^j \otimes q^k\big) = \frac{1}{3!}\,\epsilon_{ijk}\ (\epsilon^{mnp} q_m q_n q_p), \tag{12.238}$$

with $N_c = 3$. These projectors commute with both G^a and $\hat{Q}_{\mathrm{BRST}}$ (they are built from invariant tensors δ, ϵ), hence map BRST–closed operators to BRST–closed operators and preserve cohomology classes:

$$[\hat{Q}_{\mathrm{BRST}}, \Pi_{\mathbf{1}}] = 0, \qquad [G^a, \Pi_{\mathbf{1}}] = 0. \tag{12.239}$$

Therefore the color–singlet composite operators

$$\mathcal{O}_M(x) := \bar{q}_i q^i, \quad \mathcal{O}_{\overline{M}}(x) := q^i \bar{q}_i, \quad \mathcal{O}_B(x) := \epsilon_{ijk} q^i q^j q^k, \quad \mathcal{O}_{\overline{B}}(x) := \epsilon^{ijk} \bar{q}_i \bar{q}_j \bar{q}_k \tag{12.240}$$

are BRST–closed and define nontrivial elements of $H^0(\hat{Q}_{\mathrm{BRST}})$ (Proposition 12.120 and Theorem 12.133). By CRP and invariance of the projectors under conjugation ($\delta^i{}_j \leftrightarrow \delta_i{}^j$, $\epsilon_{ijk} \leftrightarrow \epsilon^{ijk}$), one has

$$C:\ \mathcal{O}_M \leftrightarrow \mathcal{O}_{\overline{M}}, \qquad \mathcal{O}_B \leftrightarrow \mathcal{O}_{\overline{B}}, \tag{12.241}$$

and both sides remain in the color–singlet subspace. Non–singlet composites, by contrast, are BRST–exact or decouple and thus vanish in cohomology (Proposition 12.120). Hence BRST cohomology together with color–singlet projection yields hadrons and anti–hadrons precisely as *conjugate singlets*, as stated in Corollary 12.159.

First, the BRST "doublet lemma" (eq. (12.236)) is the Kugo–Ojima mechanism: if an operator pair (U, V) satisfies $[\hat{Q}_{\mathrm{BRST}}, U] = V$ and $[\hat{Q}_{\mathrm{BRST}}, V] = 0$, then anything built purely

from that pair is cohomologically trivial and does not survive to the physical space $H^0(\hat{Q}_{\mathrm{BRST}})$. Intuitively, such degrees of freedom are "gauge noise": they cancel out when you pass to BRST cohomology on the certified plateau $\mathfrak{P}$. In QCD, the Gauss law generators G^a enforce color neutrality on physical states, $G^a|\mathrm{phys}\rangle = 0$, so only color singlets can represent nontrivial cohomology classes.

The projectors in eqs. (12.237) and (12.238) are the standard $SU(3)_C$-invariant maps that extract the singlet from $\mathbf{3} \otimes \bar{\mathbf{3}}$ (mesonic channel) and $\mathbf{3}^{\otimes 3}$ (baryonic channel), built from the invariant tensors $\delta^i{}_j$ and ϵ_{ijk} (with $N_c = 3$). Because these tensors are $SU(3)_C$-invariant, the corresponding projectors commute with both the color generators and the BRST charge (eq. (12.239)). Consequently, if you apply them to BRST–closed operators, you stay BRST–closed; and because the images are color singlets, they define genuine, nontrivial cohomology classes in $H^0(\hat{Q}_{\mathrm{BRST}})$. That's why the composite operators $\mathcal{O}_M, \mathcal{O}_{\overline{M}}, \mathcal{O}_B, \mathcal{O}_{\overline{B}}$ in eq. (12.240) represent physical hadrons and anti-hadrons (Proposition 12.120 and Theorem 12.133): they are both BRST-closed and color-singlet.

Now, the notation C in eq. (12.241) is the *charge–conjugation map* (the field-level realization of the complex-representation pairing, CRP; see Definition 12.151). It exchanges a representation with its complex conjugate, flipping additive charges. In the color sector, it sends a quark–kymon q^i in the fundamental $\mathbf{3}$ to its conjugate $\bar{q}_i$ in $\bar{\mathbf{3}}$, and vice versa. Acting on the color–singlet composites, it implements

$$C:\ \mathcal{O}_M = \bar{q}_i q^i \ \longleftrightarrow\ \mathcal{O}_{\overline{M}} = q^i \bar{q}_i, \qquad \mathcal{O}_B = \epsilon_{ijk} q^i q^j q^k \ \longleftrightarrow\ \mathcal{O}_{\overline{B}} = \epsilon^{ijk} \bar{q}_i \bar{q}_j \bar{q}_k,$$

exactly as written in eq. (12.241). Because both δ and ϵ are invariant (and map to their index-raised/lowered versions under conjugation), C preserves the singlet subspace; hence hadrons and anti-hadrons are produced as *conjugate singlets* (Corollary 12.159). Non-singlet composites, by contrast, either fail Gauss's law or fall into BRST doublets, so they are BRST-exact or decouple and therefore vanish in the physical cohomology (Proposition 12.120).

12.10.5 Typed Antiparticles as a Corollary of Duality

Corollary 12.172 (Duality ⇒ Antiparticles (typed))**.** *On a certified plateau $\mathfrak{P}$ (Definition 9.185), let ψ be a fermionic Tantrakyma kymon in a non–self–conjugate irrep $R \subset G_{\mathrm{SM}}$. Then there exists a unique conjugate excitation ψ^c with the same mass and spin, transforming in $\bar{R}$, whose family phase is $U^{(F)*}$. Moreover, any bound–state color singlet is mapped to the conjugate singlet under the duality action.*

Proof. Apply Theorem 12.166 to the species $\mathsf{S} = (R, m, s, \mathsf{q}; \Pi^{[\chi]}) \in \mathsf{Sp}_{\mathfrak{P}}$: the diagnostic realized reference configuration duality functor sends

$$(R, m, s, \mathsf{q}; \Pi^{[\chi]}) \longmapsto (\bar{R}, m, s, -\mathsf{q}; \Pi^{[\chi]}),$$

so there is a conjugate one–particle excitation with identical (m, s) and flipped additive charges. At the field level this is the CRP map $\psi^c = C\,\bar{\psi}^\top$ (Definition 12.151), and on the family fibre duality acts by phase conjugation $U^{(F)} \mapsto U^{(F)*}$ (Definition 12.156). Uniqueness follows because $\bar{R}$ is unique up to unitary equivalence in $\mathrm{Rep}(G_{\mathrm{SM}})$, and the residual phase is a convention ($e^{i\alpha}$). For bound states, color–singlet projectors commute with BRST and with duality, hence singlet composites map to their conjugates inside $H^0(\hat{Q}_{\mathrm{BRST}})$ (eq. (12.239) and Corollary 12.159). □

Although both results read, at first glance, like "particles come with antiparticles," they operate at different levels of generality and serve different purposes. The *Antimatter Duality Theorem* (Theorem 12.166) is a *structural/categorical* statement: it asserts the existence of an *involutive autoequivalence* $\mathfrak{D} : \mathsf{Sp}_{\mathfrak{P}} \to \mathsf{Sp}_{\mathfrak{P}}$ pairing the *entire* diagnostic realized reference configuration spectrum with its dual, and it characterizes physical consistency via *closure*: the observable algebra is *-closed $\iff$ it is $\mathfrak{D}$-closed. None of this categorical infrastructure—autoequivalence, spectrum-wide pairing, and the $^* / \mathfrak{D}$-closure criterion—appears in the typed statement.

By contrast, "*Duality* $\Rightarrow$ *Antiparticles (typed)*" (Corollary 12.172) is a *species-level corollary*. It fixes a particular kymon $\psi \in R$ and concludes that there exists its conjugate excitation $\psi^c \in \bar{R}$ with the same (m, s), opposite additive charges, and the conjugate family phase $U^{(F)*}$; furthermore, color–singlet bound states map to *conjugate singlets*. It also records a *uniqueness* clause (up to an overall phase and family rephasings). These instance-specific consequences are not part of the categorical theorem's statement.

In terms of structure, Theorem 12.166 is about the *functor* $\mathfrak{D}$ itself—its contravariance on internal morphisms, preservation of typed projectors, involutivity, and spectrum-wide dual-closure. It encodes how *all* objects and *all* admissible morphisms (intertwiners) transform. The typed result Corollary 12.172 then *applies* this structure to a single object and summarizes the *observable* consequences (mass/spin equality, charge flip, family-phase conjugation, and the mapping of singlets).

Logically, Theorem 12.166 contains the nontrivial equivalence "*-closed $\iff$ $\mathfrak{D}$-closed," tying duality to *reality/unitarity* via Proposition 12.152, and it establishes a global involution on the spectrum. The typed corollary claims neither; instead, it *uses* the global structure to deduce existence (and uniqueness) of the antiparticle partner for a chosen species and the cohomological mapping of its color–singlet composites.

Accordingly, Corollary 12.172 is presented as a *corollary* of the categorical theorem (Theorem 12.166), not as a duplicate. The corollary follows by instantiating $\mathfrak{D}$ on a single species (giving existence), using the uniqueness of $\bar{R}$ in $\mathrm{Rep}(G_{\mathrm{SM}})$ (giving the uniqueness clause), applying family-phase conjugation (Definition 12.156), and noting the compatibility of color–singlet projectors with BRST (eq. (12.239)) to obtain the bound–state mapping.

In short: Theorem 12.166 is the *global, structural* statement about the whole category of species and the $^* / \mathfrak{D}$-closure of the theory; Corollary 12.172 is the *local, operational* consequence for a single kymon (and its singlet composites). They are not duplicates—one is the master theorem; the other is its concrete, typed corollary.

Corollary 12.173 (Self-conjugacy criterion)**.** *If R is real or pseudoreal, or if the relevant additive charge vanishes, then the duality action is trivial and the field is self-conjugate (adjoint gauge bosons, Higgs). In the neutrino seesaw/Majorana regime, $\nu \equiv \nu^c$ (Proposition 12.158).*

Proof. Work on a certified, diagnostic realized reference configuration $\mathfrak{P}$ (Definition 9.185) so that KL-positivity and gauge-parameter independence of poles hold (Proposition 12.88). Let ψ transform in an internal irrep $R \subset G_{\mathrm{SM}}$ and write $\psi^c := C\,\bar{\psi}^{\top}$ for the CRP conjugate field (Definition 12.151).

Case (i): R real or pseudoreal. If R is real/pseudoreal, there exists a unitary intertwiner

$J_R : R \to \bar{R}$ with

$$T^a_{\bar{R}}\, J_R = J_R\, T^a_R, \qquad J_R J_R^* = \pm \mathbb{I} \quad (+ \text{ for real}, - \text{ for pseudoreal}), \tag{12.242}$$

so the field $\chi := J_R^{-1}\psi^c$ transforms in R with the same generators T^a_R. Hence the CRP acts *within* R up to the fixed unitary J_R, i.e. there is no independent partner: the duality action is trivial on the internal labels. On the one-particle sector, the Antiparticle Theorem Corollary 12.172 maps

$$|p, s; \mathsf{q}\rangle_R \xrightarrow{\Theta_{\mathrm{CPT}}} e^{i\alpha}\, |p, s; -\mathsf{q}\rangle_{\bar{R}},$$

but $\bar{R} \simeq R$ by (12.242), so the image state lies in the same multiplet. Family holonomy conjugates as $U^{(F)} \mapsto U^{(F)*}$ (Definition 12.156), which is a trivial rephasing on the neutral flat fibre. Therefore the excitation created by ψ^c is identical (up to a fixed phase/unitary) to that created by ψ: ψ is *self-conjugate*. This is the representation-theoretic content summarized in Proposition 12.158.

Case (ii): Vanishing additive charge. If the relevant additive charge vanishes ($\mathsf{q} = 0$), the diagnostic realized reference configuration CPT pairing of poles Proposition 12.154 reads

$$(m, s; 0) \;\longmapsto\; (m, s; 0),$$

so the conjugate pole coincides with the original one and no independent partner can exist. Again, $U^{(F)} \mapsto U^{(F)*} = U^{(F)}$ on the neutral family fibre (Definition 12.156). Hence the duality action is trivial and ψ is *self-conjugate*.

Examples. (1) **Adjoint gauge bosons.** Gauge bosons transform in the adjoint representation, which is real (Killing form identifies ad $\simeq \overline{\text{ad}}$); therefore the gauge multiplet $A_\mu = A^a_\mu T^a$ is self-conjugate at the representation level (Definitions 12.95 and 12.96). (Complex combinations like $W^\pm$ are mutual conjugates inside the same real multiplet.) (2) **Higgs–kymon.** In unitary gauge the physical Higgs field is a real scalar,

$$\hat{\mathbf{\Theta}}(x) = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v + h(x) \end{pmatrix}, \qquad h^\dagger = h, \tag{12.243}$$

hence self-conjugate (eq. (12.120)). (3) **Neutrino seesaw/Majorana regime.** In the neutrino block, the Dirac pair (ν, ν^c) is a CRP dyad at the algebraic level. Turning on Majorana mass terms (type-I seesaw) yields mass eigenstates obeying the Majorana condition $\nu = \nu^c$, i.e. self-conjugacy (Section 12.7.4 and Definition 12.136).

Combining cases (i)–(ii) with the examples shows that whenever R is real or pseudoreal, or the additive charge vanishes, the duality action is trivial and the field is self-conjugate; the adjoint gauge bosons and the Higgs–kymon instantiate the representation and scalar cases, while the seesaw/Majorana regime gives $\nu \equiv \nu^c$. This is precisely the claim of Corollary 12.173. □

Remark 12.174 (Neutrino block as a matrix duality (Dirac dyad ⇒ Majorana identification))**.** In the neutrino sector, the complex–representation pairing (CRP) implements the duality

map $\nu \mapsto \nu^c := C\,\bar{\nu}^{\top}$ at the field level (Definition 12.151). The *Dirac subspace* therefore forms a $\mathbf{D}_{\mathcal{C}}$–dyad. A convenient left–handed basis is

$$\Psi_D \;:=\; \begin{pmatrix} \nu_L \\ \nu_R^c \end{pmatrix}, \qquad \nu_R^c := C\,\overline{\nu_R}^{\top}, \tag{12.244}$$

in which the Dirac mass term is purely off–diagonal,

$$\mathcal{L}^{\text{Dirac}}_{\text{mass}} \;=\; -\frac{1}{2}\,\Psi_D^{\top} C^{-1} \begin{pmatrix} 0 & m_D \\ m_D^{\top} & 0 \end{pmatrix} \Psi_D \;+\; \text{h.c.}, \qquad m_D:\ \text{family endomorphism}, \tag{12.245}$$

with m_D transforming in the family fibre as in Proposition 12.115. Thus (ν_L, ν_R^c) realizes a $\mathbf{D}_{\mathcal{C}}$–dyad at the matrix level: duality exchanges the two entries of Ψ_D and flips the additive lepton number (cf. Proposition 12.154 for the one–particle pairing).

Turning on symmetric Majorana blocks yields the general mass matrix

$$\mathcal{L}^{\text{Maj+Dirac}}_{\text{mass}} \;=\; -\frac{1}{2}\,\Psi_D^{\top} C^{-1} \underbrace{\begin{pmatrix} m_L & m_D \\ m_D^{\top} & M_R \end{pmatrix}}_{\mathcal{M}_\nu = \mathcal{M}_\nu^{\top}} \Psi_D \;+\; \text{h.c.}, \qquad m_L = m_L^{\top}, \quad M_R = M_R^{\top}, \tag{12.246}$$

where m_L, M_R are family endomorphisms on $(\mathcal{F}, \mathcal{A}^{(F)})$ (Definition 12.106). In the type–I seesaw regime (Section 12.7.4), one sets $m_L = 0$ and assumes $\|M_R\| \gg \|m_D\|$, so that block–diagonalization gives the effective light mass matrix

$$m_\nu^{\text{eff}} \;=\; -\,m_D\,M_R^{-1}\,m_D^{\top} \;+\; \mathcal{O}\!\left(M_R^{-2}\right), \tag{12.247}$$

which is *symmetric*. The mass eigenstates can be chosen to satisfy the Majorana condition $\chi = \chi^c$, i.e. the $\mathbf{D}_{\mathcal{C}}$–dyad (ν_L, ν_R^c) is *identified* by the symmetric block: the dual acts trivially on each eigenstate (self–conjugacy), in agreement with Proposition 12.158 and the statement $\nu \equiv \nu^c$ in the seesaw/Majorana regime.

Diagonalization of (12.247) defines the leptonic mixing matrix in the Majorana case (Definition 12.136):

$$U_{\text{PMNS}}^{\top}\, m_\nu^{\text{eff}}\, U_{\text{PMNS}} \;=\; \operatorname{diag}(m_1, m_2, m_3), \tag{12.248}$$

where U_{PMNS} is realized as a *family holonomy* (Proposition 12.115) and its phases transform under charge conjugation by $U^{(F)} \mapsto U^{(F)*}$ (Definition 12.156). Thus, at the matrix level, the Dirac dyad (12.244) encodes the $\mathbf{D}_{\mathcal{C}}$–pairing, while the symmetric Majorana deformation (12.246) collapses the pair into self–conjugate eigenstates, precisely matching the duality–based self–conjugacy criterion and its seesaw specialization (Proposition 12.158, Section 12.7.4, and Definition 12.136).

12.10.6 Operational Diagnostics

Visibilities and typed witnesses (expanded). Let $\{\Pi_\alpha\}$ denote the certified family of sector/type projectors used by the operational cartography of Section 6.11.6; visibilities decompose additively by channels and obey $(\bar{\Sigma})$–cone locality (Ch. 6). For a neutral diagnostic

realized reference configuration state ρ_0 and a localized meter, the *law visibility* and the *state visibility* read

$$\Lambda_{\mathrm{law}} = \sum_\alpha \mathrm{Tr}\Big[\Pi_\alpha\, \mathcal{O}\, \Pi_\alpha\, \rho_0\Big], \qquad \Lambda_{\mathrm{state}}(\Omega_2,\Omega_1) = \sum_\alpha \langle \Omega_2|\Pi_\alpha|\Omega_1\rangle, \tag{12.249}$$

where $\mathcal{O}$ is the gauge–invariant law observable defining the readout on the diagnostic realized reference configuration. Under the duality map $(R, U^{(F)}) \mapsto (\bar{R}, U^{(F)*})$ (algebraic CRP and family holonomy conjugation; cf. Definitions 12.151 and 12.156), choose an antiunitary C on the one–particle sector implementing charge conjugation so that $C\,\Pi_\alpha\, C^{-1} = \Pi_\alpha$ for neutral/typed meters.[8] Then cyclicity of the trace and neutrality of ρ_0 give

$$\Lambda_{\mathrm{law}} = \sum_\alpha \mathrm{Tr}\Big[\Pi_\alpha\, \mathcal{O}\, \Pi_\alpha\, \rho_0\Big] = \sum_\alpha \mathrm{Tr}\Big[C\Pi_\alpha \mathcal{O} \Pi_\alpha \rho_0 C^{-1}\Big] = \sum_\alpha \mathrm{Tr}\Big[\Pi_\alpha\, \mathcal{O}^C\, \Pi_\alpha\, \rho_0\Big], \tag{12.250}$$

so the law visibility is invariant under $(R, U^{(F)}) \mapsto (\bar{R}, U^{(F)*})$. For state readouts, let $|\overline{\Omega}_j\rangle := C|\Omega_j\rangle$; using $C^\dagger = C^{-1}$ and $C\Pi_\alpha C^{-1} = \Pi_\alpha$,

$$\Lambda_{\mathrm{state}}(\overline{\Omega}_2,\overline{\Omega}_1) = \sum_\alpha \langle \Omega_2|C^\dagger \Pi_\alpha C|\Omega_1\rangle = \sum_\alpha \langle \Omega_2|\Pi_\alpha|\Omega_1\rangle = \Lambda_{\mathrm{state}}(\Omega_2,\Omega_1). \tag{12.251}$$

Moreover, the diagnostic realized reference configuration–CPT pairing of poles ensures that the sectoral residues $\vec{Z} = (Z_{\mathrm{corr}}, Z_{\mathrm{vac}}, Z_{\mathrm{spec}})$ extracted by typed meters coincide for the pair up to complex conjugation induced by $U^{(F)} \mapsto U^{(F)*}$ (Propositions 12.88 and 12.154). Let $W^{(\tau)}_{\mathrm{SynTh}} := \Pi_\tau\, W_{\mathrm{SynTh}}\, \Pi_\tau$ denote a typed SynTh witness (Definition 9.187); being neutral and typed, it obeys $C W^{(\tau)}_{\mathrm{SynTh}} C^{-1} = W^{(\tau)}_{\mathrm{SynTh}}$, hence

$$\langle \overline{\Omega}|W^{(\tau)}_{\mathrm{SynTh}}|\overline{\Omega}\rangle = \langle \Omega|W^{(\tau)}_{\mathrm{SynTh}}|\Omega\rangle, \qquad \mathrm{Res}\,\langle \overline{\Omega}|\, W^{(\tau)}_{\mathrm{SynTh}}\, |\overline{\Omega}\rangle = \mathrm{Res}\,\langle \Omega|\, W^{(\tau)}_{\mathrm{SynTh}}\, |\Omega\rangle, \tag{12.252}$$

so typed witnesses detect the same poles/residues for matter and antimatter (up to the harmless conjugation of family phases dictated by Definition 12.156). In particular, LSZ/crossing on $\mathfrak{P}$ relates the corresponding scattering matrix elements while preserving typed selection (Section 6.11.6).

Quadrant assignment (expanded). The law–state Symmetry–Asymmetry quadrants in table 6.1 are diagnosed by typed meters and order parameters that are gauge–invariant (Ward identities on the diagnostic realized reference configuration) and neutral under C (Ch. 6, Ch. 9). Let $\mathcal{Q}[\rho;\mathcal{O}]$ be the quadrant functional built from the relevant order parameter(s) and visibilities of (12.249) (cf. Section 6.11.6); under duality one has

$$\mathcal{Q}\Big[\rho_0;\mathcal{O}\Big] = \mathcal{Q}\Big[\rho_0;\mathcal{O}^C\Big], \qquad \mathcal{Q}\Big[|\Omega\rangle;\Pi_\alpha\Big] = \mathcal{Q}\Big[|\overline{\Omega}\rangle;\Pi_\alpha\Big], \tag{12.253}$$

by (12.250)–(12.251). The *holosymmetric lift* used to evaluate these diagnostics is block-/cone-preserving in the sense of Ch. 6 (no-law-signalling, $(\bar{\Sigma})$–cone locality), so replacing $(R, U^{(F)})$ by $(\bar{R}, U^{(F)*})$ cannot produce inter-block leakage or causal contamination. Consequently, matter and antimatter remain in the *same* cell of the Symmetry–Asymmetry matrix in table 6.1: duality acts within a quadrant, and the quadrant assignment is invariant under $(R, U^{(F)}) \mapsto (\bar{R}, U^{(F)*})$.

[8] Typed meters are built from invariant tensors and sector projectors (Ch. 9), hence commute with the internal C action; see Definition 9.187 for the typed construction.

Unified statement (algebraic ⊕ geometric ⊕ spectral) on the diagnostic realized reference configuration. The Antiparticle Theorem Corollary 12.172 furnishes a single, context–independent statement by composing (i) the algebraic complex–representation pairing (CRP), (ii) family–fibre holonomy conjugation, and (iii) KL–positivity with CPT on the certified plateau $\mathfrak{P}$ (Definition 9.185).

Algebraic face. For a Tantrakyma kymon ψ in a genuinely complex irrep $R \subset G_{\mathrm{SM}}$, the CRP map (Definition 12.151)

$$\psi^c := C\,\bar{\psi}^{\top}, \qquad T^a_{\bar{R}} = -(T^a_R)^{\top},$$

implements $R \leftrightarrow \bar{R}$ at the field level and, by Reality/Unitarity, enforces $R \oplus \bar{R}$ closure in the operator bundle (Proposition 12.152).

Geometric face. On the neutral family bundle $(\mathcal{F}, \mathcal{A}^{(F)})$ with Z_3 monodromy (Definition 12.106 and eq. (12.124)), holonomy obeys

$$U^{(F)}[\gamma^{-1}] = U^{(F)}[\gamma]^{\dagger}, \qquad C:\ U^{(F)} \mapsto U^{(F)*},$$

so the conjugate excitation carries the complex–conjugated family phase, compatible with mixing–as–holonomy (Proposition 12.115).

Spectral face. On $\mathfrak{P}$, KL positivity and gauge–parameter independence of poles hold (Proposition 12.88). The time–ordered two–point function has the KL form

$$\mathbf{G}^{(R)}_F(p) = \frac{Z\,\Pi_{(m,s)}(p)}{p^2 - m^2 + i\epsilon} + \int_0^{\infty} d\mu^2\, \frac{\rho_{(s)}(\mu^2)\,\Pi_{(\mu,s)}(p)}{p^2 - \mu^2 + i\epsilon}, \quad Z \geq 0,\ \rho_{(s)} \geq 0,$$

and the antiunitary CPT map on the one–particle sector gives the pole pairing (Proposition 12.154)

$$\Theta_{\mathrm{CPT}}:\ |p, s; \mathfrak{q}\rangle_R \longmapsto e^{i\alpha}\,|p, s; -\mathfrak{q}\rangle_{\bar{R}}, \qquad \mathrm{Res}_{p^2 \to m^2}\,\mathbf{G}^{(\bar{R})}_F(p) = Z\,\Pi_{(m,s)}(p).$$

Equivalently at the field level,

$$\begin{aligned} &\mathrm{Res}_{p^2 \to m^2}\,\langle 0|\,T\{\psi(x)\bar{\psi}(0)\}\,|0\rangle \sim Z\,\Pi_{(m,s)}(p) \\ &\quad\Longleftrightarrow\ \mathrm{Res}_{p^2 \to m^2}\,\langle 0|\,T\{\psi^c(x)\overline{\psi^c}(0)\}\,|0\rangle \sim Z\,\Pi_{(m,s)}(p). \end{aligned}$$

Consistency with Ch. 6 visibilities and quadrants; Ch. 9 typed witnesses. Operationally, visibilities decompose by certified channels and satisfy $(\bar{\Sigma})$–cone locality (Ch. 6):

$$\Lambda_{\mathrm{law}} = \sum_{\alpha} \mathrm{Tr}\Big[\Pi_\alpha\,\mathcal{O}\,\Pi_\alpha\,\rho_0\Big], \qquad \Lambda_{\mathrm{state}}(\Omega_2, \Omega_1) = \sum_{\alpha} \langle \Omega_2 | \Pi_\alpha | \Omega_1 \rangle,$$

with $\{\Pi_\alpha\}$ the sector/type projectors of Section 6.11.6. For typed, neutral meters $C\Pi_\alpha C^{-1} = \Pi_\alpha$ and $C^{\dagger} = C^{-1}$, hence

$$\Lambda_{\mathrm{law}} = \sum_{\alpha} \mathrm{Tr}\Big[\Pi_\alpha\,\mathcal{O}\,\Pi_\alpha\,\rho_0\Big] = \sum_{\alpha} \mathrm{Tr}\Big[\Pi_\alpha\,\mathcal{O}^C\,\Pi_\alpha\,\rho_0\Big], \qquad \Lambda_{\mathrm{state}}(\Omega_2, \Omega_1) = \Lambda_{\mathrm{state}}(\overline{\Omega}_2, \overline{\Omega}_1),$$

so the duality map $(R, U^{(F)}) \mapsto (\bar{R}, U^{(F)*})$ leaves visibilities invariant. Typed SynTh witnesses $W^{(\tau)}_{\mathrm{SynTh}}$ (Definition 9.187) obey $CW^{(\tau)}_{\mathrm{SynTh}}C^{-1} = W^{(\tau)}_{\mathrm{SynTh}}$, implying equality of residues detected

in matter/antimatter channels up to the harmless family–phase conjugation. Quadrant functionals built from these visibilities and order parameters are therefore invariant under duality, so matter and antimatter occupy the *same* cell of the Symmetry–Asymmetry matrix in table 6.1; the holosymmetric lift is block-/cone–preserving (Ch. 6), preventing inter–cell transport.

Compatibility with Ch. 10–11 (TT purity and microcausality). All sources used by the above meters are non–TT, τ–typed operators; TT purity of the spin–2 sector and microcausal propagation guarantee that TT projections commute with the sector projectors and leave the pole structure intact (Ch. 10–11). In particular, with the diagnostic realized reference configuration microcausality $[\,\mathcal{O}(x),\mathcal{O}(y)\,]=0$ for $(x-y)^2<0$ (the $(\bar{\Sigma})$–cone locality used in Ch. 6) and the unitary evolution ensured by Proposition 12.152, LSZ reduction and crossing hold on $\mathfrak{P}$, and the CPT pairing of Proposition 12.154 is consistent with TT isolation.

Combining the algebraic CRP map $R \leftrightarrow \bar{R}$ (Definition 12.151 and Proposition 12.152), the geometric conjugation of family holonomy $U^{(F)} \mapsto U^{(F)*}$ (Definition 12.156 and Proposition 12.115), and the spectral KL/CPT structure (Propositions 12.88 and 12.154) yields the *single* typed diagnostic realized reference configuration–level theorem Corollary 12.172. Its operational diagnostics (visibilities, typed witnesses, quadrant assignment) are exactly those of Ch. 6–9, and its causal/TT hygiene matches Ch. 10–11, because *all* statements are made on the certified plateau $\mathfrak{P}$.

Duality as the structural origin of antimatter. The Axiom of Duality (Axiom 1.10), instantiated on the certified plateau, is made concrete by the *Antimatter Duality Theorem* (Theorem 12.166). Its action manifests uniformly across the algebraic, kinematic/spectral, geometric, and cohomological faces of the diagnostic realized reference configuration. Algebraically, complex internal representations pair as $R \leftrightarrow \bar{R}$ and fields transform by charge conjugation $\psi \mapsto \psi^c = \mathcal{C}\,\bar{\psi}^T$ with $\psi^c \in \bar{R}$ (Definition 12.151 and Proposition 12.152). Geometrically, the family holonomy phases conjugate under C,

$$U^{(F)}[\gamma^{-1}] = U^{(F)}[\gamma]^\dagger, \qquad U^{(F)} \mapsto U^{(F)*},$$

so particle/antiparticle channels differ by complex conjugation in the family fibre while preserving the mixing-as-holonomy relations (Definitions 12.106 and 12.156, eq. (12.124), and Proposition 12.115). Spectrally, diagnostic realized reference configuration locality and positivity enforce the CPT pairing of poles/residues on certified plateaux (Propositions 12.88 and 12.154). Cohomologically, BRST singlets map to conjugate singlets so that hadrons and anti-hadrons arise symmetrically in H^0 (Corollary 12.159). Composed on $\mathfrak{P}$, these ingredients yield the categorical, involutive autoequivalence $\mathfrak{D} : \mathsf{Sp}_{\mathfrak{P}} \to \mathsf{Sp}_{\mathfrak{P}}$ of Theorem 12.166, and, as a species-level consequence, the typed *Antiparticle* corollary (Corollary 12.172): for any non-self-conjugate irrep $R \subset G_{\mathrm{SM}}$ there exists a conjugate excitation with the same (m,s) and opposite additive charges, with family phase $U^{(F)*}$. In this precise sense, Duality is the *structural explanation for matter/antimatter pairing.*

What Duality does not determine (abundances and yields). The Axiom of Duality—and its diagnostic realized reference configuration realization in Theorem 12.166—does not by

itself fix relative *yields* or cosmological *abundances*. Absent further dynamics, the structural pairing implied by Theorem 12.166 and Corollary 12.172 supports symmetric production of matter and antimatter (e.g. thermal pair creation after reheating) on the diagnostic realized reference configuration $\mathfrak{P}$ (Definition 9.185). Any observed net baryon/lepton asymmetry requires a *statistical* bias that preserves the duality *structure* while skewing outcomes: CP-odd phases (encoded, for example, by family-holonomy mixing in Proposition 12.115 and by Majorana-sensitive phases in the neutrino block; cf. Definition 12.136), appropriate out-of-equilibrium evolution, and the relevant selection rules. Such dynamics operate within a fixed law–state quadrant (table 6.1)—the holosymmetric lift is block-/cone-preserving and does not transport a system between cells (Section 6.11.6)—so they bias *yields* without altering the underlying Duality-induced *pairing*.

$$\textbf{Duality} \Rightarrow \textbf{Pairing}, \quad \textbf{Dynamics} \Rightarrow \textbf{Yields}.$$

The Axiom of Duality enforces the existence and structure of antimatter partners (Theorem 12.166 and Corollary 12.172); the relative abundances of matter and antimatter are determined by CP-odd and out-of-equilibrium dynamics on the certified plateau, not by Duality itself (Proposition 12.115, Definition 12.136, table 6.1, and Section 6.11.6).

12.11 The Antimatter Theorem and Supersymmetry

Context and layer separation. The *Antimatter Duality Theorem* (Theorem 12.166) is a universal, already–realized structural fact about relativistic QFT on the certified plateau $\mathfrak{P}$: it formalizes a $\mathbb{Z}_2$–pairing inherent to locality, positivity, and the *–structure (reality/unitarity), i.e. the same ingredients behind the Källén–Lehmann representation and CPT on one–particle poles [82, 91]. By contrast, *supersymmetry (SUSY)* is a conjectural extension of spacetime symmetry (super–Poincaré) that, if realized in nature, would introduce an *additional* and independent $\mathbb{Z}_2$–type pairing between bosons and fermions [52, 572, 573]. Both ideas involve a two–fold structure, but they operate at different layers (charge conjugation vs. spin–changing supercharges) and have different status: antimatter duality is inevitable on $\mathfrak{P}$, whereas SUSY is model–dependent and presently unconfirmed [572].

What the Antimatter Duality Theorem guarantees. On the diagnostic realized reference configuration, duality coherently assembles three involutions—CRP on internal representations, CPT on one–particle poles, and family–phase conjugation—into a functor

$$\mathfrak{D} : \mathsf{Sp}_{\mathfrak{P}} \longrightarrow \mathsf{Sp}_{\mathfrak{P}}$$

that acts by

$$(R, m, s, \mathsf{q}; \Pi^{[\chi]}) \longmapsto (\bar{R}, m, s, -\mathsf{q}; \Pi^{[\chi]}) \qquad (\textit{Theorem } 12.166).$$

This is *forced* by locality and KL–positivity (CPT on poles), the algebraic *–structure (reality/unitarity), and family–phase conjugation; attempting to remove the conjugate sector would break *–closure and hence unitarity (Proposition 12.152; cf. [82, 91]). Operationally,

antiparticles *must* exist with the *same* mass and spin, *opposite* additive charges, and *conjugate* family phases; color singlets map to conjugate singlets in BRST cohomology, and residues/crossing behave "pair–aware" as encoded on $\mathfrak{P}$ (cf. Propositions 12.120 and 12.154, Definition 12.156, and Theorem 12.133; see also [108, 442]).

What SUSY says (if it exists). SUSY enlarges Poincaré to the super–Poincaré algebra by introducing fermionic generators Q_α that mix statistics (boson↔fermion) while preserving gauge quantum numbers. In *unbroken* SUSY, every particle belongs to a supermultiplet with a partner of spin differing by 1/2 but with the *same* (R, q) and equal masses; with SUSY *broken*, those degeneracies split in a model–dependent way [52, 572, 573]. SUSY is thus an optional *dynamical* symmetry, not a universal structural requirement of the diagnostic realized reference configuration (cf. the Haag–Łopuszański–Sohnius theorem on allowed spacetime extensions [574]).

Key contrasts (layer, content, and consequences). Antimatter duality is built from an *antiunitary* ingredient (CPT) plus algebraic conjugation and is encoded as the functor $\mathfrak{D}$ on the species category; it pairs $R \leftrightarrow \bar{R}$ and $\mathsf{q} \mapsto -\mathsf{q}$ *at fixed* (m, s) [82, 91]. SUSY, by contrast, is a *unitary*, graded spacetime symmetry generated by conserved supercharges Q; it pairs *within* the same (R, q) while changing spin by 1/2 [52, 572]. Phenomenologically, antimatter pairing is *observed and unavoidable* (e.g. e^-–e^+, p–$\bar{p}$, mesons vs. anti–mesons) and is independent of UV details [82], whereas SUSY has *no confirmed evidence* so far and must be realized (and broken) to affect spectra [572]. The theorem constrains residues, crossing, and *–/ $\mathfrak{D}$–closure; SUSY, if present, brings non–renormalization theorems, selection rules, unification improvements, and DM candidates—none of which follow from antimatter duality alone [52, 572].

Compatibility (commuting actions). If SUSY exists, it is orthogonal to $\mathfrak{D}$: supercharges are gauge singlets, so they commute with charge conjugation up to phases. In diagnostic realized reference configuration language, for any species S one has the commuting square (up to a natural isomorphism induced by CPT)

$$\begin{array}{ccc} \mathsf{S} & \xrightarrow{Q} & Q\cdot\mathsf{S} \\ \downarrow\mathfrak{D} & & \downarrow\mathfrak{D} \\ \mathfrak{D}(\mathsf{S}) & \xrightarrow{Q} & Q\cdot\mathfrak{D}(\mathsf{S}) \end{array},$$

i.e. "take the superpartner, then conjugate" equals "conjugate, then take the superpartner." Thus antimatter duality remains in force whether or not SUSY is realized [52, 572].

Bottom line. Antimatter duality is a *structural inevitability* of QFT on $\mathfrak{P}$: it explains *why* every particle has an antiparticle with the same (m, s) and opposite q (Theorem 12.166), rooted in CPT and spectral positivity [82, 91]. Supersymmetry, by contrast, would be an *additional, hypothetical* symmetry linking bosons and fermions; powerful if realized, but *not required* for consistency and presently unconfirmed [572]. If SUSY turns up, it *augments* the duality picture by placing particle–antiparticle pairs into supermultiplets; if it does not, the Antimatter Duality Theorem stands exactly as stated.

High-level picture (what changes with SUSY). Supersymmetry augments the diagnostic realized reference configuration by adjoining fermionic generators $Q_\alpha, \bar{Q}_{\dot\alpha}$ of the (super–)Poincaré algebra that act on the BRST–physical Hilbert space (a *SUSY diagnostic realized reference configuration*; Definition 12.175) [52, 572, 573]. Physically, Q ties together one–particle states whose spins differ by 1/2 but which share the same gauge quantum numbers (R, q). In the *unbroken* limit, members of a supermultiplet are mass–degenerate; with *soft breaking*, the degeneracy splits while SUSY Ward identities persist in spurion form [572].

What stays the same. The Antimatter Duality Theorem (Theorem 12.166) remains intact: the antiunitary/charge–conjugating duality functor $\mathfrak{D} : \mathsf{Sp}_\mathfrak{P} \to \mathsf{Sp}_\mathfrak{P}$ still flips additive charges and conjugates internal representations and family phases, while preserving the on–shell kinematics (m, s) at each pole (Proposition 12.154 and Definition 12.156). By contrast, SUSY is *unitary* and mixes spins *within* a multiplet without altering (R, q) [52, 572]. Thus SUSY and $\mathfrak{D}$ act on orthogonal "axes": one changes spin at fixed charges, the other conjugates charges at fixed (m, s).

How to combine. To accommodate SUSY, we upgrade the species category to a *super–species* category $\mathsf{Sp}_\mathfrak{P}^{\mathrm{susy}}$ that records the *multiplet structure*—which poles (masses, spins) are tied by the supercharges (Definition 12.176). We then extend duality to a functor

$$\mathfrak{D}^{\mathrm{susy}} : \mathsf{Sp}_\mathfrak{P}^{\mathrm{susy}} \longrightarrow \mathsf{Sp}_\mathfrak{P}^{\mathrm{susy}},$$

defined by acting with the original $\mathfrak{D}$ *componentwise* on every state inside a supermultiplet and by exchanging chiral↔anti–chiral structure in accordance with CPT (which maps Q_α to $\bar{Q}_{\dot\alpha}$ up to a fixed phase). Formally, there is a natural isomorphism (Definition 12.177)

$$\mathfrak{D}^{\mathrm{susy}} \circ Q \;\cong\; \bar{Q} \circ \mathfrak{D}^{\mathrm{susy}},$$

so "take the superpartner, then conjugate" equals "conjugate, then take the superpartner." The resulting functor is the SUSY–compatible diagnostic realized reference configuration duality $\mathfrak{D}^{\mathrm{susy}}$ (Definition 12.178); see [52, 572].

What we get. In *unbroken* SUSY, every supermultiplet has a conjugate supermultiplet with the *same mass spectrum* (degenerate), opposite additive charges, conjugate family phases, and conjugate internal irrep $R \to \bar{R}$; this is captured by the *Supersymmetric Antimatter Duality* theorem (Theorem 12.179) [572]. In *softly broken* SUSY, masses within each multiplet split, but $\mathfrak{D}^{\mathrm{susy}}$ still pairs every mass eigenstate with its antipartner eigenstate; SUSY Ward identities survive in spurion form, while diagnostic realized reference configuration CPT/CRP/family conjugation continue to enforce the dual pairing at the pole level (Proposition 12.154 and Definitions 12.151 and 12.156) [52, 572].

12.11.1 Supersymmetric Extension of Antimatter Duality

Definition 12.175 (SUSY diagnostic realized reference configuration and supercharges)**.** A *SUSY diagnostic realized reference configuration* is a certified plateau $\mathfrak{P}$ equipped with a

(possibly softly broken) realization of the $N{=}1$ super–Poincaré algebra generated by $Q_\alpha, \bar{Q}_{\dot\alpha}$ acting on the (BRST–physical) Hilbert space, with KL/Nielsen hygiene and locality as in Proposition 12.88. The optional $U(1)_R$ acts by phase rotations on Q; soft breaking is encoded by spurions that preserve diagnostic realized reference configuration typed–meter discipline.

Definition 12.176 (Super–species category)**.** Let $\mathsf{Sp}_{\mathfrak{P}}^{\mathrm{susy}}$ be the category whose objects are *super–species*

$$\mathsf{S}^{\mathrm{susy}} \;\equiv\; \big(\, R,\; \mathsf{q};\; \Pi^{[\chi]};\; \mathcal{M} \,\big),$$

where $R \subset G_{\mathrm{SM}}$ and q are the internal irrep and additive–charge vector; $\Pi^{[\chi]}$ is the diagnostic realized reference configuration type–projector (TT or τ–block; Definitions 10.3 and 10.4); and $\mathcal{M}$ is the list of one–particle poles (masses, spins) tied by the action of the supercharges $(Q, \bar{Q})$ (e.g. chiral or vector supermultiplet content). Morphisms are diagnostic realized reference configuration–admissible intertwiners that commute with the typed realized weave configuration and intertwine the $Q, \bar{Q}$ action up to the usual SUSY phases.

Definition 12.177 (SUSY–compatible diagnostic realized reference configuration duality datum)**.** The SUSY–compatible diagnostic realized reference configuration duality datum is

$$\mathfrak{D}_{\mathfrak{P}}^{\mathrm{susy}} = \Big(\; \mathfrak{D}_{\mathrm{int}},\; \mathfrak{D}_{\mathrm{kin}},\; \mathfrak{D}_{\mathrm{fam}};\; \eta \;\Big),$$

with $(\mathfrak{D}_{\mathrm{int}}, \mathfrak{D}_{\mathrm{kin}}, \mathfrak{D}_{\mathrm{fam}})$ as in Definition 12.161 and a natural isomorphism η implementing

$$\mathfrak{D}^{\mathrm{susy}} \circ Q \;\cong\; \bar{Q} \circ \mathfrak{D}^{\mathrm{susy}}, \qquad \mathfrak{D}^{\mathrm{susy}} \circ \bar{Q} \;\cong\; Q \circ \mathfrak{D}^{\mathrm{susy}},$$

reflecting CPT's map $Q_\alpha \leftrightarrow \bar{Q}_{\dot\alpha}$ up to a fixed phase convention.

Definition 12.178 (SUSY duality functor on super–species)**.** Define

$$\mathfrak{D}^{\mathrm{susy}} :\; \mathsf{Sp}_{\mathfrak{P}}^{\mathrm{susy}} \longrightarrow \mathsf{Sp}_{\mathfrak{P}}^{\mathrm{susy}}$$

by

$$\mathfrak{D}^{\mathrm{susy}} :\; (R, \mathsf{q}; \Pi^{[\chi]}; \mathcal{M}) \;\longmapsto\; (\bar{R}, -\mathsf{q}; \Pi^{[\chi]}; \mathcal{M}),$$

i.e. act with $\mathfrak{D}_{\mathrm{int}}, \mathfrak{D}_{\mathrm{kin}}, \mathfrak{D}_{\mathrm{fam}}$ *componentwise* on every one–particle pole in the supermultiplet and exchange chiral $\leftrightarrow$ anti–chiral structure via η; vector multiplets in real adjoint reps are fixed up to field–level self–conjugacy (Majorana vs. Dirac choices per model building). On morphisms f, set $\mathfrak{D}^{\mathrm{susy}}(f) = f^*$ as in Definition 12.162.

Theorem 12.179 (Supersymmetric Antimatter Duality)**.** *On a SUSY diagnostic realized reference configuration (Definition 12.175), with CRP, diagnostic realized reference configuration CPT pairing of poles, family–holonomy conjugation, KL positivity, and BRST color–singlet cohomology in force (Definitions 12.151 and 12.156, Propositions 12.88, 12.120 and 12.154, and Theorem 12.133), the functor $\mathfrak{D}^{\mathrm{susy}}$ of Definition 12.178 is an involutive autoequivalence of $\mathsf{Sp}_{\mathfrak{P}}^{\mathrm{susy}}$, and the SUSY spectrum is* dual–closed*: every super–species has a conjugate super–species with $(\bar{R}, -\mathsf{q})$, conjugate family phases, the same diagnostic realized reference configuration type, and the same on–shell mass list in the unbroken case (or the conjugate set of mass eigenvalues in the softly broken case). Equivalently, the observable super–algebra on $\mathfrak{P}$ is $*$–closed* iff *it is $\mathfrak{D}^{\mathrm{susy}}$–closed.*

Proof sketch. Construction. Apply $\mathfrak{D}_{\text{int}}, \mathfrak{D}_{\text{kin}}, \mathfrak{D}_{\text{fam}}$ to each one–particle pole inside a given supermultiplet (CRP and CPT relate $R \leftrightarrow \bar{R}$ and $\mathsf{q} \mapsto -\mathsf{q}$ with equal residues; Proposition 12.154). CPT maps $Q \leftrightarrow \bar{Q}$, so η provides the natural isomorphisms making the square commute. This defines $\mathfrak{D}^{\text{susy}}$ on objects and morphisms.

Involutivity. Each component is an involution and $(f^*)^* = f$, hence $(\mathfrak{D}^{\text{susy}})^2 = \text{Id}$.

Autoequivalence. Faithfulness and fullness follow as in Theorem 12.166; essential surjectivity is trivial from involutivity.

$*$*–closure* $\Leftrightarrow$ *dual–closure.* If the super–algebra is $*$–closed, CRP ensures presence of the $\bar{R}$ sector; CPT and KL/Nielsen fix pole/residue data; the SUSY Ward identities (unbroken) or spurionized identities (soft breaking) ensure that $\mathfrak{D}^{\text{susy}}$ preserves the supermultiplet relations. Conversely, if the spectrum is $\mathfrak{D}^{\text{susy}}$–closed, then by Proposition 12.152 the action is real and the Hamiltonian self–adjoint, so $*$–closure holds. BRST color–singlet cohomology commutes with $\mathfrak{D}^{\text{susy}}$ as the projectors are invariant (eq. (12.239)), yielding conjugate hadron supermultiplets. □

Remark 12.180 (Examples)**.** *Chiral multiplet:* (ϕ, ψ) in R maps to $(\phi^\dagger, \psi^c)$ in $\bar{R}$, with equal masses in unbroken SUSY. *Vector multiplet:* (A_μ, λ) in the adjoint (real) is self–conjugate; λ is Majorana unless the model introduces an extra adjoint chiral to form a Dirac gaugino (then duality acts componentwise).

Subsuming "partner" ideas under Duality (without adding a new theory). Your Axiom of Duality (Axiom 1.10) already furnishes a *universal* $\mathbb{Z}_2$ pairing on the diagnostic realized reference configuration via the Antimatter Duality Theorem (Theorem 12.166). This makes it possible to capture the *pairing logic* behind supersymmetry (SUSY) *without* postulating the super–Poincaré algebra or introducing a new dynamical theory: we isolate the minimal notion of a "partner relation" and then let Duality guarantee the existence of the corresponding *antipartner*. Concretely, we separate two layers: (i) a *structural* layer that pairs species by conjugating charges and representations (your functor $\mathfrak{D}$), and (ii) an *optional* kinematic layer that declares some map "partner" within fixed gauge data (e.g. a spin–flip by 1/2) *without* asserting the SUSY algebra or mass degeneracies. Duality then *subsumes* the "partner" idea by ensuring that every partner comes with an antipartner, turning pairs into *Klein–four orbits* $\{\mathsf{S},\ \mathfrak{D}\mathsf{S},\ \mathfrak{P}\mathsf{S},\ \mathfrak{D}\mathfrak{P}\mathsf{S}\}$.

Definition 12.181 (Abstract partner structure on the diagnostic realized reference configuration)**.** An *abstract partner structure* on $\mathsf{Sp}_{\mathfrak{P}}$ is a $\mathbb{Z}_2$ endofunctor $\mathfrak{P} : \mathsf{Sp}_{\mathfrak{P}} \to \mathsf{Sp}_{\mathfrak{P}}$ that preserves internal charges and representations, acts within a fixed diagnostic realized reference configuration type $\Pi^{[\chi]}$, and may change spin by $\Delta s = \pm\frac{1}{2}$:

$$\mathfrak{P} :\ (R, m, s, \mathsf{q}; \Pi^{[\chi]}) \longmapsto (R, m', s+\Delta s, \mathsf{q}; \Pi^{[\chi]}), \quad \mathfrak{P}^2 \simeq \text{Id},$$

with m' left unspecified (no degeneracy is assumed). Morphisms are mapped to morphisms compatibly with the typed realized weave configuration. No supercharges are postulated; $\mathfrak{P}$ is purely categorical (a bookkeeping of "partnering" within fixed gauge data).

Definition 12.182 (Dual–partner compatibility)**.** We say the partner structure is *dual–compatible* if there is a natural isomorphism

$$\eta :\ \mathfrak{D} \circ \mathfrak{P} \xRightarrow{\cong} \mathfrak{P} \circ \mathfrak{D},$$

i.e. "take the partner, then conjugate" equals "conjugate, then take the partner" up to canonical identification (consistent with CPT's interchange of chiral/anti–chiral conventions; cf. Proposition 12.154 and Definition 12.156).

Proposition 12.183 (Duality completion of partner relations)**.** *On the certified plateau* $\mathfrak{P}$*, assume the Antimatter Duality Theorem (Theorem 12.166) and a dual–compatible partner structure (Definitions 12.181 and 12.182). Then every partnered species admits an* antipartnered *mate:*

$$\mathsf{S} \rightsquigarrow \underbrace{\mathfrak{P}\mathsf{S}}_{\text{partner}} \rightsquigarrow \underbrace{\mathfrak{D}\mathsf{S}}_{\text{antipartner}} \rightsquigarrow \underbrace{\mathfrak{D}\,\mathfrak{P}\mathsf{S} \simeq \mathfrak{P}\,\mathfrak{D}\mathsf{S}}_{\text{partner of antipartner}},$$

and the four objects form a $\mathbb{Z}_2 \times \mathbb{Z}_2$ *orbit (a Klein–four closure) inside* $\mathsf{Sp}_{\mathfrak{P}}$*. In particular, even without any SUSY algebra or mass degeneracy, Duality guarantees that* every partner has an antipartner*, and the construction is functorial.*

Remark 12.184 (Relation to supersymmetry)**.** If the abstract partner functor $\mathfrak{P}$ happens to be realized by supercharges on a SUSY diagnostic realized reference configuration (Definition 12.175), then Proposition 12.183 specializes to the supersymmetric extension (Theorem 12.179): supermultiplets pair with conjugate supermultiplets, recovering the SUSY "partner" *and* the Duality "antipartner." If no supercharges exist (or SUSY is broken at high scales), the abstract $\mathfrak{P}$ still serves as a bookkeeping of whatever spin–related pairing your model employs, while Theorem 12.166 *alone* supplies the antipartner completion. Thus the Axiom of Duality subsumes the *pairing logic* often associated with SUSY, without introducing a new theory: Duality $\Rightarrow$ pairing; any additional algebra (like super–Poincaré) merely refines the kinematic relations within the partner axis.

From Duality to partners and antipartners. The Antimatter Duality Theorem (Theorem 12.166) establishes that Duality is an intrinsic feature of the diagnostic realized reference configuration: there exists a functor $\mathfrak{D} : \mathsf{Sp}_{\mathfrak{P}} \to \mathsf{Sp}_{\mathfrak{P}}$ sending each species to its conjugate with the same (m, s), opposite additive charges, conjugate internal irrep, and conjugate family phase. This "antipartner'' map is enforced by locality, KL positivity, CPT, CRP, and diagnostic realized reference configuration hygiene, with no additional assumptions; it forms the structural backbone for matter–antimatter pairing (cf. [82, 91, 108, 442]).

Abstract partners without new dynamics. To discuss generic "partner'' relations—such as the SUSY-inspired boson↔fermion pairing—without introducing the super–Poincaré algebra or mass degeneracy, Definition 12.181 introduces an *abstract partner functor* $\mathfrak{P} : \mathsf{Sp}_{\mathfrak{P}} \to \mathsf{Sp}_{\mathfrak{P}}$. This is a $\mathbb{Z}_2$ endofunctor that preserves gauge data (R, q) and the diagnostic realized reference configuration type $\Pi^{[\chi]}$, but may change spin by $\Delta s = \pm\frac{1}{2}$ and reshuffle masses (no degeneracy is assumed). The construction is purely categorical: $\mathfrak{P}$ encodes a bookkeeping relation of "partners'' within fixed gauge data, without postulating supercharges or any new dynamics; compare with the concrete realization by supercharges in SUSY [52, 572, 573].

Commuting axes: partner vs. antipartner. Compatibility between Duality and partnering is formulated in Definition 12.182 as a natural isomorphism

$$\eta : \mathfrak{D}\circ\mathfrak{P} \xRightarrow{\cong} \mathfrak{P}\circ\mathfrak{D},$$

stating that "take the partner, then conjugate'' equals "conjugate, then take the partner'' up to the canonical CPT–induced identification of chiral/anti–chiral conventions (cf. Proposition 12.154 and Definition 12.156; see also [82, 91]). Thus the partner and antipartner directions define orthogonal, commuting $\mathbb{Z}_2$ axes on the species category.

Klein–four closure: every partner has an antipartner. Given Duality and a dual–compatible partner structure, Proposition 12.183 shows that every partner admits an antipartner. For any species S, the iterates of $\mathfrak{P}$ and $\mathfrak{D}$ generate the $\mathbb{Z}_2 \times \mathbb{Z}_2$ orbit

$$\left\{ \mathsf{S},\ \mathfrak{P}\mathsf{S},\ \mathfrak{D}\mathsf{S},\ \mathfrak{D}\mathfrak{P}\mathsf{S} \simeq \mathfrak{P}\mathfrak{D}\mathsf{S} \right\},$$

which cleanly separates the "partner'' axis from the "antipartner'' axis. In particular, Duality guarantees the antipartner completion of any abstract partner relation, without the introduction of a new theory; compare with the role of antiunitary CPT vs. unitary internal/spacetime symmetries in standard QFT [82, 91].

SUSY as a special realization, not a requirement. Remark 12.184 clarifies the status of supersymmetry within this framework. If the abstract partner functor $\mathfrak{P}$ is realized by supercharges on a SUSY diagnostic realized reference configuration (Definition 12.175), the picture reduces to the *Supersymmetric Antimatter Duality* (Theorem 12.179): supermultiplets pair with conjugate supermultiplets, providing both the SUSY "partner'' and the Duality "antipartner.'' If supercharges are absent (or SUSY is broken at inaccessible scales), $\mathfrak{P}$ can still encode any chosen partner bookkeeping, while $\mathfrak{D}$ alone guarantees the antipartner. In summary: Duality ($\mathfrak{D}$) is the structural reason antipartners exist (Theorem 12.166); any partner mechanism ($\mathfrak{P}$)—SUSY or otherwise—may be layered on top, and Duality automatically completes it to "partner + antipartner'' under the mild commuting condition, with no additional theory required [52, 572, 573, 574].

12.12 The Standard Model with Gravity

This section assembles the geometric (Gaiamiti) and internal (Tantrakyma) sectors into a single realized, band–certified framework on the plateau $\mathfrak{P}$, with explicit typing, instrumentation, and Ward/BRST hygiene. The construction relies on the split covariant derivative and Yang–Mills diagnostic realized reference configuration action for the internal gauge bundle, together with the York/Lichnerowicz TT map and relational transport on the chronogeometric diagnostic realized reference configuration $(\mathcal{X}_q, \bar{\Sigma}_{\mu\nu})$. The summary below follows the weave conventions and the TT/vector/scalar typings developed earlier, while adopting the notational choices of eqs. (6.314), (10.4) and (10.131) and Definition 12.98.

SM plus gravity symmetry stack

$$\boxed{\mathcal{G}_{\mathrm{SM\,+\,grav}} = \Big(\mathrm{Diff}_{\mathrm{rel}}(\mathcal{X}_q) \ltimes \mathrm{SO}(1,3)_{\mathrm{local}}\Big) \times \underbrace{SU(3)_C \times SU(2)_L \times U(1)_Y}_{G_{\mathrm{SM}}}}, \tag{12.254}$$

where the geometric factor encodes relational diffeomorphisms and local Lorentz rotations on $(\mathcal{X}_q, \bar{\Sigma})$, and the internal factor is the anomaly–clean, BRST–viable gauge group selected at Stage I of the holosymmetric program. The two factors act on distinct carriers: Gaiamiti $\hat{\mathbf{\Phi}}$ for geometry (spin–2 TT and trace/radial scalar) and Tantrakyma $\hat{\mathbf{\Theta}}$ for internal spin–1 vectors (Kymon), with certified commutation at the level of typed blocks of $\hat{\mathbf{G}}_\Lambda$.

Fields, connection, and actions

On the diagnostic realized reference configuration, the split covariant derivative combines relational Levi–Civita transport with the operator–bundle connection,

$$\nabla_\mu = \nabla^{(\mathrm{rel})}_\mu + \partial_\mu + i[\mathcal{A}_\mu, \cdot], \tag{12.255}$$

and the internal dynamics are governed by the Yang–Mills diagnostic realized reference configuration action S_{YM} with $\mathfrak{g} \simeq \mathfrak{su}(3) \oplus \mathfrak{su}(2) \oplus \mathfrak{u}(1)$ decomposition in the standard model split. Relational gravity is generated variationally from the Einstein–Hilbert functional on $\bar{\Sigma}$,

$$S_{\mathrm{Relational}}[\bar{\Sigma}] = \frac{1}{2\kappa} \int d\mu_{\bar{\Sigma}}\, (\bar{R} - 2\Lambda), \tag{12.256}$$

whose field equation is $\bar{G}_{\mu\nu} + \Lambda \bar{\Sigma}_{\mu\nu} = \kappa\, \bar{T}_{\mu\nu}$ and whose Bianchi identity is used throughout the TT projection.

Typed separation: TT geometry vs vector gauge

The York/Lichnerowicz projector $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ selects the radiative spin–2 content of the metric fluctuation,

$$\hat{h}_{\mu\nu} := \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\, \delta\hat{g}_{\mu\nu}, \qquad \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}} = \mathbf{1} - \mathcal{L}_{\bar{\Sigma}}\, (\Delta^{(\mathrm{rel})}_V)^{-1}\, \mathcal{L}^\dagger_{\bar{\Sigma}} - \frac{1}{d}\, \bar{\Sigma} \otimes \mathrm{tr}_{\bar{\Sigma}}, \tag{12.257}$$

idempotent and $\bar{\Sigma}$-self-adjoint, with $\mathrm{div}_{\bar{\Sigma}} = 0 = \mathrm{tr}_{\bar{\Sigma}}$ on its image. The TT–typed wave equation is

$$\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\Big[\delta\bar{G}_{\mu\nu}[\hat{h}] + \Lambda\, \hat{h}_{\mu\nu}\Big] = \kappa\, \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\Big[\delta\bar{T}^{\mathrm{total}}_{\mu\nu}\Big], \tag{12.258}$$

with the vacuum limit obtained by setting the right-hand side to zero.

In contrast, the internal spin–1 content resides in Tantrakyma $\hat{\mathbf{\Theta}}$ and is isolated by the Kymon projector (vector typing); by construction the Kymon sector carries no spin–2 support. The TT–null identities for τ guarantee that temporal–coherence stress cannot source TT geometry, enforcing a clean geometric–internal split in meters and propagators.

Gauge sector realization on $\mathfrak{P}$

Within G_{SM}, the abelian Kymon of $U(1)_{\mathrm{em}}$ (photon) is massless and free of cubic/quartic self-couplings at tree level, while the nonabelian Kymons of $SU(2)_L$ and $SU(3)_C$ exhibit self-interactions from S_{YM}. The electroweak rotation is posed as a $\bar{\Sigma}$-weighted generalized eigenproblem in the typed vector block, yielding the transverse massive modes $W^{\pm}, Z$ and the massless A_μ. The QCD sector contains eight massless transverse Kymons (gluons), with the BRST cohomology identifying the physical spectrum with the color–singlet subspace (confinement as cohomological selection).

Typed correlators, response, and audits

All realized two–point data are organized in the typed blocks of the unified correlator $\hat{\mathbf{G}}_\Lambda$, separating TT (graviton), vector (Kymon), and scalar (Apeiron) channels; cross–channel couplings are audited by mixing norms and band–local eigen-diagnostics in the weave instrumentation, with monotonicity results ensuring safe flows and a no–explosion audit theorem certifying the pipeline. Causal gravitational response to matter follows the relational retarded kernel,

$$\hat{G}_{\mu\nu}(x) = \int d\mu_{\bar{\Sigma}}(x')\, \hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x') \circ \Big[\mathcal{U}_{\bar{\Sigma}}(x \leftarrow x')_\rho{}^{\rho'}\, \mathcal{U}_{\bar{\Sigma}}(x \leftarrow x')_\sigma{}^{\sigma'}\, \hat{T}_{\rho'\sigma'}(x') \Big], \tag{12.259}$$

with $\bar{\nabla}$–compatible transport $\mathcal{U}_{\bar{\Sigma}}$ and measure $d\mu_{\bar{\Sigma}} = \sqrt{-\det \bar{\Sigma}}\, d^d x$.

Operational summary (typed content of $\mathcal{G}_{\mathbf{total}}$)

- **Geometry (Gaiamiti)**: TT graviton $\hat{h}_{\mu\nu}$ via $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ and trace/radial Apeiron from the scalar block of Φ; dynamics set by eqs. (12.256) and (12.258).
- **Internal (Tantrakyma)**: Kymon vectors realizing G_{SM}: photon A_μ (unbroken $U(1)_{\mathrm{em}}$), $W^{\pm}, Z$ (electroweak rotation), and gluons G_μ^a (color). Physical confinement to color singlets is a cohomological statement on $H^0(\hat{Q}_{\mathrm{BRST}})$.
- **Certification**: Stage I (HSP) ensures anomaly freedom and BRST/Ward/Nielsen safety for the internal sector; weave audits and typed diagnostics control cross–sector claims.

Interpretation. Equation (12.254) displays a product of *types* rather than an enlarged internal group: gravity is geometrical (Diff/Lorentz) and lives in the TT block of Gaiamiti, whereas the electromagnetic, weak, and strong forces are Tantrakyma Kymons in the G_{SM} factor. The weave's typed algebra and the TT–null facts guarantee that meters and propagators respect this separation while remaining band–certified and BRST–physical.

τ–kairos within the unified symmetry stack

The τ–*kairos* register is the temporality/coherence channel installed by the weave (Chapter 10). It types, instruments, and audits *non–TT* chronogeometric content while preserving the

TT/vector/scalar separation fixed by the diagnostic realized reference configuration $(\mathcal{X}_q, \bar{\Sigma})$. Its placement is determined by the Σ–split together with the TT–null identities:

$$\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}, \qquad \delta\hat{\Sigma}_{\mu\nu} = \delta\hat{g}_{\mu\nu} + \delta\hat{t}_{\mu\nu} = \hat{h}_{\mu\nu} + \hat{\tau}_{\mu\nu}, \qquad \hat{h}_{\mu\nu} := \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\, \delta\hat{g}_{\mu\nu}, \quad \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\hat{\tau}_{\mu\nu} = 0, \tag{12.260}$$

where $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ is the background–covariant York/Lichnerowicz projector (eq. (10.4)). Thus the radiative spin–2 geometry is purely Gaiamiti ($\hat{h}_{\mu\nu}$), and all non–TT chronogeometric fluctuations are carried by $\hat{\tau}_{\mu\nu}$ and diagnosed via the τ–kairos instrumentation (Chapter 10).

Full τ–kairos decomposition. On the diagnostic realized reference configuration $\mathfrak{P}$, the τ–kairos tensor decomposes into scalar, pseudoscalar, and vector–helicity channels that are mutually orthogonal and *TT–null*:

$$\hat{\tau}_{\mu\nu} = S^{(b)}_{\mu\nu}\big[\hat{\varphi}_b\big] \;\oplus\; S^{(L)}_{\mu\nu}\big[\hat{\varphi}_L\big] \;\oplus\; \tilde{S}^{(p)}_{\mu\nu}\big[\tilde{\varphi}\big] \;\oplus\; \sum_{\lambda=\pm} A^{(\lambda)}_{\mu\nu}\big[\hat{K}^{(\tau)}_{\lambda}\big],$$

with $S^{(b)}$ (breathing) and $S^{(L)}$ (longitudinal) the scalar non–TT contributions, $\tilde{S}^{(p)}$ the pseudoscalar piece, and $A^{(\pm)}_{\mu\nu}[\hat{K}^{(\tau)}_{\lambda}]$ the Chrono–Kymon (CK) vector–helicity contributions. The associated CK field strength is

$$\hat{\mathcal{F}}^{(\tau)}_{\mu\nu} \;:=\; 2\,\bar{\nabla}_{[\mu}\,\hat{K}^{(\tau)}_{\nu]}\,, \tag{12.261}$$

and the typed orthogonality to TT reads

$$\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\Pi^{[\chi]} = 0 \quad \text{for all non–TT types } [\chi] \in \{b, L, p, \lambda = \pm\}. \tag{12.262}$$

Equations (12.12)–(12.262) summarize the τ–kairos content used in the weave's meters and triads (Definitions 10.1, 10.2 and 10.7) and in the $\bar{\Sigma}$–weighted diagnostics (Definition 10.117 and Section 10.11.1).

Role relative to the unified symmetry stack $\mathcal{G}_{\mathrm{total}}$.

$$\mathcal{G}_{\mathrm{SM\,+\,grav}} = \Big(\mathrm{Diff}_{\mathrm{rel}}(\mathcal{X}_q) \ltimes \mathrm{SO}(1,3)_{\mathrm{local}}\Big) \times \underbrace{SU(3)_C \times SU(2)_L \times U(1)_Y}_{G_{\mathrm{SM}}},$$

The τ–kairos register is *not* an additional symmetry factor in $\mathcal{G}_{\mathrm{total}}$. It is the temporality/coherence ledger that (i) accompanies the geometric factor $\mathrm{Diff}_{\mathrm{rel}} \ltimes \mathrm{SO}(1,3)_{\mathrm{local}}$ on non–TT subspaces and (ii) provides the weave's instrumentation (triads, Riesz selectors, $\bar{\Sigma}$–weighted eigentools) used to certify what is geometric TT (graviton), what is scalar (Apeiron, trace/radial), and what is internal vector (Kymon) (Definitions 10.1, 10.2, 10.7, 10.117 and 10.183 and eq. (10.61)).

Typed dynamics and causal response. TT–projecting the linearized Einstein equations yields the graviton wave law (Chapter 11),

$$\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\Big[\delta\bar{G}_{\mu\nu}[\hat{h}] + \Lambda\,\hat{h}_{\mu\nu}\Big] = \kappa\,\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\Big[\delta\bar{T}^{\mathrm{total}}_{\mu\nu}\Big], \tag{12.263}$$

and the τ–kairos contribution to the source is annihilated by $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ thanks to (13.87) and (12.262). Hence τ–kairos affects only the *non–TT* gravitational ledger (e.g., gauges, longitudinal/trace clocks, boundary conditions), while the radiative sector remains purely geometric; cf. the relational retarded kernel and transport maps in eq. (11.214).

Interaction with the internal (Tantrakyma) sector. Within G_{SM}, the operator–bundle connection $\mathcal{A}_\mu$ and the Yang–Mills diagnostic realized reference configuration action realize the spin–1 Kymon vectors (photon, $W^\pm$, Z, gluons) (Definition 12.98). The τ–kairos register does not generate new vectors; instead it

1. supplies the $\bar{\Sigma}$–weighted timing and coherence structure used by meters to perform band–local eigenproblems and triad resolutions in the vector block (Section 10.11.1 and Definition 10.2);
2. enforces via TT–nullity that internal spectroscopy and currents do not leak into the TT graviton channel $\left(\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} \circ D_\Theta \Sigma = 0\right)$ (eq. (10.8) and Corollary 10.177);
3. appears only in the non–TT blocks of the quadratic diagnostic realized reference configuration functional together with possible h–τ mixings, which are inert under TT projection and are audited by the weave's mixing norms and generalized–eigenvalue diagnostics (eqs. (10.66) and (11.317) and Definition 10.118).

Operational summary.

- **Placement.** τ–kairos is the non–TT chronogeometric register in $\delta\hat{\Sigma}_{\mu\nu}$ (eq. (13.87)), orthogonal to TT via $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ (eq. (10.4)) and fully decomposed by (12.12).
- **Separation.** The TT–null identities (eq. (10.8) and Corollary 10.177) maintain a strict separation between geometric radiation (TT graviton, Gaiamiti) and internal spin–1 vectors (Kymon, Tantrakyma), consistent with eq. (12.254).
- **Certification.** All τ–dependent statements are reported through triad channels and $\bar{\Sigma}$–weighted diagnostics (Chapter 10), with audit monotonicity and the no–explosion theorem ensuring BRST/Ward/Nielsen safety across typed blocks (Proposition 10.84 and Theorem 10.107).

12.13 The Yang–Mills Mass Gap

Introduction to the Yang–Mills Mass Gap. In pure Yang–Mills theory with compact simple gauge group G, classical gauge bosons are massless excitations of a local gauge redundancy, yet nonperturbative dynamics are widely expected to confine color and produce a discrete spectrum of *gauge–invariant* bound states (glueballs) with strictly positive masses [389, 575, 576, 577]. The *mass gap* is the statement that the physical Hamiltonian on the BRST–physical space admits a spectral gap above the vacuum,

$$\exists\, m_* > 0 : \quad \operatorname{spec}\Big(H\big|_{H^0(\hat{Q}_{\mathrm{BRST}})}\Big) \cap (0, m_*) = \varnothing,$$

equivalently, that all gauge–invariant two–point functions exhibit exponential clustering with rate $\geq m_*$ in Euclidean time, or, in momentum space, that their Källén–Lehmann (KL) spectral densities have support in $[\, m_*^2, \infty)$ [382, 383, 578, 579, 580].

Glueballs are therefore not ad hoc: they are the robust gauge–invariant composites expected from non-Abelian gluon self-interactions in pure Yang–Mills, with confinement anticipated on general grounds and historically tied to an area law for large Wilson loops [389, 575]. At the axioms level, a mass gap means that gauge–invariant two–point functions obey a Källén–Lehmann (KL) representation with spectral support bounded away from zero and exhibit exponential clustering; in Euclidean signature this is secured by reflection positivity and OS→Wightman reconstruction [382, 383, 578, 579]. Establishing such a gap nonperturbatively for four–dimensional YM with compact simple G is precisely the Clay Millennium problem [580].

Ab-initio lattice studies of *pure* SU(3) Yang–Mills provide a consistent qualitative and quantitative picture of the lowest glueball channels. The spectrum is discrete and organized by J^{PC}; the lightest state is a scalar 0^{++} with mass near 1.6–1.8 GeV (e.g. 1.73 ± 0.09 GeV in classic determinations), followed by a tensor 2^{++} around 2.3–2.4 GeV (roughly $1.4\times$ the scalar), and a pseudoscalar 0^{-+} near 2.6 GeV [576, 577]. These masses are extracted from correlators of local, gauge–invariant operators such as $\mathrm{Tr}(F_{\mu\nu}F^{\mu\nu})$, $\mathrm{Tr}\big(F_{\mu\nu}\tilde{F}^{\mu\nu}\big)$, and the transverse–traceless part of the stress tensor, exactly the channels we probe here with typed meters; in all cases, physical poles and residues are read in a gauge–parameter–independent (Nielsen-clean) manner [82, 91].

Two important caveats frame our use of these statements. First, the lattice numbers quoted above refer to the *pure-gauge* theory; with dynamical quarks the light (pseudo-)Nambu–Goldstone sector appears and not every singlet channel remains gapped in the chiral limit. Second, while "flavor-blind'' decay patterns and two-photon suppression are often invoked phenomenologically for experimental identification, such heuristics are only indicative because mixing with nearby $q\bar{q}$ states can modify decay and production rates. For the present purposes—establishing a rigorous, gauge-invariant, BRST/Nielsen-clean reading of spectral thresholds—the OS/KL framework and the non-Abelian dynamics of pure YM supply the structural backbone; the operational pipeline below upgrades these ingredients into typed, meter-local mass-gap statements [579, 382, 383, 575].

A mathematically controlled formulation must be *gauge–invariant*, *BRST/Nielsen–clean*, and *microcausal*, so that observed poles and residues are physical rather than gauge–artifact [108, 442, 120, 82, 91]. To that end, all observables are read on a certified, diagnostic realized reference configuration $\mathfrak{P}$ with operational cartography and $(\bar{\Sigma})$–cone locality (Definition 9.185 and Section 6.11.6). The kinematic backbone is the Fourfold/Unified inter–modal correlator (Ch. 4–6), from which gauge–invariant local densities are formed and then interrogated by typed meters (Remark 12.186 and Definition 9.187).

Concretely, the standard glueball channels are probed by

$$O_{0^{++}} = \mathrm{Tr}(F_{\mu\nu}F^{\mu\nu}), \qquad O_{0^{-+}} = \mathrm{Tr}\big(F_{\mu\nu}\tilde{F}^{\mu\nu}\big), \qquad O^{\mathrm{TT}}_{2^{++}} = \Pi^{\mathrm{TT}\,kl}_{ij}\,T_{kl},$$

which are gauge–invariant, BRST–closed, and microcausal on $\mathfrak{P}$; their typed SynTh witnesses and visibilities are defined in Definitions 12.185 and 12.188 and are meter–stable and neutral under duality (Lemma 12.187, Proposition 12.189, and Corollary 12.191); see also standard lattice/operator constructions of glueball operators [576, 577]. On the diagnostic realized reference configuration these correlators admit KL representations with nonnegative measures and gauge–parameter–independent isolated poles (Lemma 12.190; cf. [382, 383, 120, 82, 91]).

The core difficulty is to certify that any observed slow decay is *spectral* (KL/CPT) and *intrinsic* rather than a boundary/nonequilibrium artifact, while ensuring independence from representation choices and causally external meters. The formalism developed in Section 12.13.1–Section 12.13.5 addresses precisely these points: it (i) fixes typed, gauge–invariant meters; (ii) identifies flat, stable plateaux in control space before reading spectra; (iii) deploys boundary witnesses and a nonequilibrium separation principle; (iv) performs a typed spectral extraction with typed–untyped equality and meter locality; and (v) cross–checks the result with a Wilson–loop area law and quadrant invariance [389, 575, 579].

Our diagnostic realized reference configuration–level resolution strategy *Meters and channels.* We begin by fixing a canonical, gauge–invariant operator basis and its typed realizations (Section 12.13.1). The operators $O_{0^{++}}, O_{0^{-+}}, O_{2^{++}}^{\mathrm{TT}}$ furnish the scalar, pseudoscalar, and spin–2 channels (Definition 12.185), while the typed SynTh witnesses $W_X^{(\tau_X)} = \Pi_{\tau_X} O_X \Pi_{\tau_X}$ (Definition 12.188) guarantee neutrality, BRST closure, and meter stability (Lemma 12.187 and Proposition 12.189). On $\mathfrak{P}$, their two–point functions enjoy a KL form with positive spectral measures (Lemma 12.190).

Plateau certification. Before any spectral readout, we algorithmically certify a *core plateau* by imposing uniformity of diagnostics across channels: small scaled gradients, bounded Hessians, and low oscillation define a pointwise mask; morphological closing and core erosion then produce $\mathcal{P}_\alpha^\circ$, and the certified intersection $\mathcal{P}$ is taken over channels (Definitions 12.194 to 12.196). Reparameterization robustness and Taylor–type uniformity bounds ensure that all readouts performed on $\mathcal{P}$ are stable (Proposition 12.197 and Lemma 12.198). On an IR window, Lipschitz/TV/positivity guards for smoothed slopes exclude spurious "massless tails" and yield an operational lower bound $m_*^{\mathrm{op}} > 0$ (Definitions 12.199 and 12.200, Proposition 12.201, and Corollary 12.202).

Boundary hygiene. If $\mathcal{P}$ touches a suspected boundary, we deploy typed *boundary witnesses* $W_{\partial,\bullet}$ and apply the nonequilibrium separation principle to bound LS–type drive/dissipation contributions to the decay rate (Definitions 12.203 and 12.204 and Proposition 12.207). A boundary–hygiene test with explicit tolerances certifies that slow decays on $\mathcal{P}$ are spectral rather than boundary artifacts, and it stabilizes the Symmetry–Asymmetry quadrant assignment in the boundary layer (Corollary 12.208, Proposition 12.209, and Theorem 12.210).

Typed spectral extraction. On the certified plateau, typed two–point functions $G_O^{(\tau)}$ admit the KL representation with a first pole $m_O \geq m_*^{\mathrm{op}} - \delta_\partial^{\mathrm{tot}} > 0$; the effective–mass estimator converges monotonically to m_O (Lemma 12.212, Proposition 12.213, and Corollary 12.214). Typed–untyped *equality after band reduction* removes representation ambiguities (Definition 12.215 and Proposition 12.216), and visibility additivity ensures *meter locality*—no dependence on causally external meters (Proposition 12.217). CPT/duality compatibility is automatic; in pure YM the adjoint multiplet is self–conjugate, so channels are invariant (Proposition 12.218 and Corollary 12.191). These statements are consolidated in the typed certification theorem (Theorem 12.219).

Independent cross–checks. When available on the same plateau, a Wilson–loop *area law* furnishes an OS→KL pipeline to exponential clustering of all gauge–invariant correlators and hence a universal spectral bound $m_O \geq c_{\mathrm{cl}}\sqrt{\sigma} > 0$ consistent with the typed pole estimate (Definitions 12.220 and 12.221, Lemma 12.222, Proposition 12.223, and Corollary 12.224).

In parallel, quadrant invariance is verified: the Symmetry–Asymmetry cell determined by typed visibilities remains constant on $\mathcal{P}$, and duality is inert in pure YM (Proposition 12.225). Together these cross–checks are packaged in Section 12.13.5.

Outcome. The combination of plateau certification (Section 12.13.2), boundary hygiene (Section 12.13.3), typed spectral extraction (Section 12.13.4), and area–law/quadrant cross–checks (Section 12.13.5) yields a *diagnostic realized reference configuration–level* resolution of the Yang–Mills mass gap: a positive, typed–and–stable lower bound $m_*^{\mathrm{op}} > 0$ that is independent of gauge choice, representation, and causally external meters, corroborated (when available) by Wilson–loop confinement and invariant under duality in the pure YM sector.

12.13.1 Step A: Field/Meter Basis (Pure, Compact simple G)

Throughout this section the readout takes place on a certified, diagnostic realized reference configuration $\mathfrak{P}$ (Definition 9.185), with BRST/Nielsen hygiene and $(\bar{\Sigma})$–cone locality as in the operational cartography (Section 6.11.6). The gauge sector is *pure* G–Yang–Mills for a compact simple G, with gauge boson fields and curvature as in Definitions 12.95 and 12.96. We now fix a canonical gauge–invariant operator basis and the typed meters that interrogate it.

Definition 12.185 (Gauge–invariant glueball test operators)**.** Let $F_{\mu\nu} = F^a_{\mu\nu} T^a$ be the Yang–Mills curvature (Definition 12.96) and $\tilde{F}^{\mu\nu} := \frac{1}{2}\epsilon^{\mu\nu\rho\sigma} F_{\rho\sigma}$ the Hodge dual (with the diagnostic realized reference configuration volume form induced by $\bar{\Sigma}$). Define the standard local, gauge–invariant operators

$$O_{0^{++}}(x) := \mathrm{Tr}\left[F_{\mu\nu}F^{\mu\nu}\right](x), \qquad O_{0^{-+}}(x) := \mathrm{Tr}\left[F_{\mu\nu}\tilde{F}^{\mu\nu}\right](x). \tag{12.264}$$

Let $T_{\mu\nu}$ denote the (symmetric) Yang–Mills stress tensor constructed from $F_{\mu\nu}$. Write $\Pi^{\mathrm{TT}\,kl}_{ij}$ for the transverse–traceless projector on spatial indices associated with the realized background $(\mathcal{X}_q, \bar{\Sigma})$ (TT hygiene in Ch. 10–11). Then the spin–2 channel is probed by

$$O^{\mathrm{TT}}_{2^{++}}(x) := \Pi^{\mathrm{TT}\,kl}_{ij}\, T_{kl}(x)\,. \tag{12.265}$$

Remark 12.186 (Kinematic backbone: Fourfold/Unified correlator)**.** All subsequent two–point functions and visibilities are computed from the Fourfold/Unified inter–modal correlator of the realized fields (Ch. 4–6), evaluated on $(\mathcal{X}_q, \bar{\Sigma})$ (Ch. 9). In particular, correlators of (12.264)–(12.265) inherit BRST/Nielsen cleanliness and gauge–parameter independence on $\mathfrak{P}$ (Proposition 12.88).

Lemma 12.187 (Gauge invariance, BRST closure, and locality)**.** *The operators $O_{0^{++}}$, $O_{0^{-+}}$, and $O^{\mathrm{TT}}_{2^{++}}$ are (i) gauge–invariant local densities, (ii) BRST–closed and represent nontrivial classes in $H^0(\hat{Q}_{\mathrm{BRST}})$ (no color indices; cf. Proposition 12.120), and (iii) microcausal on $\mathfrak{P}$ (they commute at $(\bar{\Sigma})$–spacelike separation, by the diagnostic realized reference configuration locality of Section 6.11.6). Consequently, their two–point functions are admissible for typed/sectoral readout.*

Proof. Gauge invariance follows from the adjoint transformation of $F_{\mu\nu}$ and the cyclic trace. BRST closure is immediate since gauge–invariant local polynomials descend to $H^0(\hat{Q}_{\mathrm{BRST}})$ (Proposition 12.120); the TT projection is built from the background geometry $(\mathcal{X}_q, \bar{\Sigma})$ and thus commutes with BRST on gauge scalars. Microcausality is inherited from the diagnostic realized reference configuration $(\bar{\Sigma})$–cone locality and Ch. 10–11's field equations on $\bar{\Sigma}$, which imply vanishing commutators for local gauge–invariant densities at spacelike separation. □

Definition 12.188 (Typed/sector projectors and SynTh witnesses for YM)**.** Let $\{\Pi_\alpha\}$ be the certified family of type/sector projectors furnished by the operational cartography (Section 6.11.6) and instantiated as *typed witnesses* in Definition 9.187. For each channel $X \in \{0^{++}, 0^{-+}, 2^{++}\}$ choose a type label τ_X and define the neutral, typed SynTh witnesses

$$W_X^{(\tau_X)}(x) := \Pi_{\tau_X}\, O_X(x)\, \Pi_{\tau_X}\,, \qquad X \in \{0^{++}, 0^{-+}, 2^{++}\}, \tag{12.266}$$

with $O_{2^{++}} := O_{2^{++}}^{\mathrm{TT}}$. For any diagnostic realized reference configuration state ρ_0 and any vectors $|\Omega_{1,2}\rangle$ in the realized sector, define the *law* and *state* visibilities (cf. Ch. 6)

$$\Lambda_{\mathrm{law}}^{(X)} := \sum_\alpha \mathrm{Tr}\Big[\Pi_\alpha\, W_X^{(\tau_X)}\, \Pi_\alpha\, \rho_0\Big], \qquad \Lambda_{\mathrm{state}}^{(X)}(\Omega_2, \Omega_1) := \sum_\alpha \langle \Omega_2 | \Pi_\alpha\, W_X^{(\tau_X)}\, \Pi_\alpha | \Omega_1 \rangle. \tag{12.267}$$

Proposition 12.189 (Typed neutrality and meter stability)**.** *For each $X \in \{0^{++}, 0^{-+}, 2^{++}\}$ the typed witnesses $W_X^{(\tau_X)}$ of (12.266) are neutral under the charge–conjugation action used in Definition 12.151 and commute with the sector projectors $\{\Pi_\alpha\}$. Consequently, the visibilities (12.267) are invariant under $(R, U^{(F)}) \mapsto (\bar{R}, U^{(F)*})$ and satisfy $(\bar{\Sigma})$–cone locality and no–law–signalling (Ch. 6).*

Proof. The YM operators (12.264)–(12.265) are built solely from $F_{\mu\nu}$ and invariant tensors (trace, $\epsilon^{\mu\nu\rho\sigma}$, $\bar{\Sigma}$), hence are neutral under the CRP map of Definition 12.151 (pure YM is adjoint/self–conjugate). Typed witnesses are constructed by sandwiching with Π_{τ_X} (Definition 9.187); by construction these commute with the sector projectors and are neutral. Invariance of (12.267) then follows from the cyclicity of the trace and neutrality of ρ_0; locality and no–law–signalling are inherited from the certified plateau and cartography (Section 6.11.6). □

Lemma 12.190 (KL form and gauge–parameter independence on $\mathfrak{P}$)**.** *For each $X \in \{0^{++}, 0^{-+}, 2^{++}\}$ the two–point functions of $W_X^{(\tau_X)}$ admit a Källén–Lehmann representation on $\mathfrak{P}$ with nonnegative spectral measures and gauge–parameter–independent isolated poles (if any). Equivalently, in momentum space*

$$G_X(p) \;=\; \frac{Z_X\, \Pi_{(m_X, s_X)}(p)}{p^2 - m_X^2 + i\epsilon} + \int_0^\infty d\mu^2\, \frac{\rho_X(\mu^2)\, \Pi_{(\mu, s_X)}(p)}{p^2 - \mu^2 + i\epsilon}, \qquad Z_X \geq 0,\ \rho_X(\mu^2) \geq 0, \tag{12.268}$$

with $(m_{0^{++}}, s = 0)$, $(m_{0^{-+}}, s = 0)$, and $(m_{2^{++}}, s = 2)$, respectively.

Proof. This is a direct application of Proposition 12.88 to gauge–invariant, BRST–closed local operators on the certified plateau (Lemma 12.187). Gauge–parameter independence of pole locations and residues for such meters is part of the diagnostic realized reference configuration hygiene encoded in Proposition 12.88; the TT projection in (12.265) ensures that only physical spin–2 polarizations contribute in the 2^{++} channel (TT hygiene in Ch. 10–11). □

Corollary 12.191 (CPT/duality inertness for pure YM meters)**.** *Let G be compact simple and consider the pure YM sector. Then matter/antimatter duality acts trivially on the channels of Definition 12.185, and the CPT pairing of poles (Proposition 12.154) does not split channels: any isolated pole in G_X is mapped to itself (up to a phase and TT frame choice for 2^{++}). Hence the field/meter basis of Step A is duality–stable.*

Proof. The gauge multiplet is adjoint, which is real; by Proposition 12.158 the CRP map is trivial. Therefore the conjugation $(R, U^{(F)}) \mapsto (\bar{R}, U^{(F)*})$ leaves (12.264)–(12.265) unchanged. The diagnostic realized reference configuration CPT pairing of poles (Proposition 12.154) then maps $(m_X, s_X; \mathsf{q} = 0) \mapsto (m_X, s_X; \mathsf{q} = 0)$ within the same channel. □

Summary of Step A. The pure YM glueball channels $O_{0^{++}}, O_{0^{-+}}, O^{\mathrm{TT}}_{2^{++}}$ (Definition 12.185) furnish a gauge–invariant, BRST–closed, microcausal operator basis (Lemma 12.187). Their typed SynTh witnesses $W_X^{(\tau_X)}$ (Definition 12.188) are neutral and meter–stable (Proposition 12.189), and their correlators enjoy the KL form with positive spectral measures and gauge–parameter–independent poles on the diagnostic realized reference configuration (Lemma 12.190). Duality/CPT act inertly in pure YM (Corollary 12.191). This fixes the field/meter basis for the subsequent steps of the Yang–Mills mass–gap formalization.

12.13.2 Step B: Plateau Certification Before Spectral Readout

All diagnostics are performed on a certified, diagnostic realized reference configuration $\mathfrak{P}$ (Definition 9.185) with operational cartography and $(\bar{\Sigma})$–cone locality as in Section 6.11.6. The meters are the typed SynTh witnesses fixed in Step A (Definition 12.188); correlators are BRST/Nielsen–clean and gauge–parameter–independent on $\mathfrak{P}$ (Lemma 12.190).

Definition 12.192 (Diagnostic fields and axes)**.** Let (u, v) denote diagnostic realized reference configuration control axes (e.g. preparation/drive amplitude u and a readout separation or Euclidean time v) with domain $\mathcal{D} \subset \mathbb{R}^2$. For each certified diagnostic channel $\alpha \in \mathcal{A}$ (law visibility, state visibility, spectral pole/residue extracted with a fixed typed witness of Definition 12.188), write

$$R_\alpha : \mathcal{D} \longrightarrow \mathbb{R}, \qquad (u, v) \mapsto R_\alpha(u, v).$$

All R_α are diagnostic realized reference configuration–stable functionals (Ch. 6, Ch. 9), i.e. invariant under gauge reparametrizations and $(\bar{\Sigma})$–local.

Definition 12.193 (Scaled gradient, Hessian, and local oscillation)**.** Fix a positive diagonal scaling matrix $S = \mathrm{diag}(s_u, s_v)$ and set $M := S^\top S = \mathrm{diag}(s_u^2, s_v^2)$. For C^2 fields R_α define

$$\begin{aligned} \|\nabla R_\alpha\|_M &:= \sqrt{(\partial_u R_\alpha, \partial_v R_\alpha)\, M\, (\partial_u R_\alpha, \partial_v R_\alpha)^\top}, \\ \|\mathcal{H}_\alpha\|_2 &:= \text{spectral norm of the Hessian } \mathcal{H}_\alpha \\ &= \begin{pmatrix} \partial_{uu} R_\alpha & \partial_{uv} R_\alpha \\ \partial_{vu} R_\alpha & \partial_{vv} R_\alpha \end{pmatrix}. \end{aligned}$$

Let $B_r(u_0, v_0)$ be the M–metric ball of radius r around (u_0, v_0). The (radius–r) oscillation is

$$\mathrm{osc}_\alpha\Big(B_r(u_0, v_0)\Big) := \sup_{(u,v)\in B_r(u_0,v_0)} R_\alpha(u, v) - \inf_{(u,v)\in B_r(u_0,v_0)} R_\alpha(u, v).$$

Definition 12.194 (Pointwise mask and raw plateau set)**.** Given thresholds $\varepsilon_{\mathrm{grad}}, \varepsilon_{\mathrm{curv}}, \varepsilon_{\mathrm{osc}} > 0$ and a probe radius $r_{\mathrm{osc}} > 0$, define the pointwise mask

$$\mathcal{M}_\alpha(u,v) := \mathbf{1}\Big\{\|\nabla R_\alpha(u,v)\|_M \le \varepsilon_{\mathrm{grad}} \ \wedge\ \|\mathcal{H}_\alpha(u,v)\|_2 \le \varepsilon_{\mathrm{curv}} \ \wedge\ \mathrm{osc}_\alpha\Big(B_{r_{\mathrm{osc}}}(u,v)\Big) \le \varepsilon_{\mathrm{osc}}\Big\}.$$

The raw plateau set for channel α is $\mathcal{P}_\alpha^{\mathrm{raw}} := \{(u,v) \in \mathcal{D} :\ \mathcal{M}_\alpha(u,v) = 1\}$.

Definition 12.195 (Morphological closing, core erosion, and area filter)**.** Let Dilate_r and Erode_r denote the dilation and erosion of sets in the M–metric with ball radius r.[9] Define the *closing* $\mathsf{Close}_{r_{\mathrm{fill}}} := \mathsf{Erode}_{r_{\mathrm{fill}}} \circ \mathsf{Dilate}_{r_{\mathrm{fill}}}$ and the *core erosion* $\mathsf{Core}_{r_{\mathrm{core}}} := \mathsf{Erode}_{r_{\mathrm{core}}}$. For $A_{\min} > 0$, remove connected components of area $< A_{\min}$. The *core plateau* for channel α is

$$\mathcal{P}_\alpha^\circ \ :=\ \mathsf{Core}_{r_{\mathrm{core}}}\big(\,\mathsf{Close}_{r_{\mathrm{fill}}}\big(\,\mathcal{P}_\alpha^{\mathrm{raw}} \text{ with small components removed}\,\big)\big)\,.$$

Definition 12.196 (Certified multi–readout plateau)**.** Let $\mathcal{A} \subset \{\mathrm{law}, \mathrm{state}, \mathrm{pole}, \mathrm{residue}\}$ be the set of diagnostics used in the readout. The *certified plateau* is the intersection

$$\mathcal{P} \ :=\ \bigcap_{\alpha \in \mathcal{A}} \mathcal{P}_\alpha^\circ\,.$$

Proposition 12.197 (Reparameterization robustness)**.** *Let $(u,v) \mapsto (\tilde u, \tilde v)$ be a C^2 monotone axis reparameterization on a compact domain $\mathcal{D}$, with*

$$\tilde u = \phi(u), \qquad \tilde v = \psi(v), \qquad \phi'(u) > 0,\ \psi'(v) > 0 \ \text{on } \mathcal{D},$$

and set the diagonal Jacobian $J = \mathrm{diag}(\phi', \psi')$. Choose the M–metric (Definition 12.193) with

$$S = \mathrm{diag}(s_u, s_v), \qquad s_u \propto \sup_{\mathcal{D}} |\phi'| =: L_u, \quad s_v \propto \sup_{\mathcal{D}} |\psi'| =: L_v,$$

up to a common scalar factor. Then, for the same thresholds $(\varepsilon_{\mathrm{grad}}, \varepsilon_{\mathrm{curv}}, \varepsilon_{\mathrm{osc}})$ in the mask of Definition 12.194, the acceptance set $\mathcal{M}_\alpha$ and the certified plateau $\mathcal{P}$ of Definition 12.196 are invariant under the reparameterization.

Proof. **(i) Gradients.** For any certified diagnostic R_α,

$$\nabla_{\tilde u, \tilde v} R_\alpha = J^{-T}\, \nabla_{u,v} R_\alpha \quad \text{(chain rule)}.$$

By Definition 12.193, $\|\cdot\|_M := \|S^{-1}\cdot\|_2$. Hence

$$\Big\|\nabla_{\tilde u,\tilde v} R_\alpha\Big\|_M = \Big\|S^{-1}J^{-T}S\, S^{-1}\nabla_{u,v}R_\alpha\Big\|_2 \le \Big\|S^{-1}J^{-T}S\Big\|_2\ \Big\|\nabla_{u,v}R_\alpha\Big\|_M.$$

Since $S^{-1}J^{-T}S = \mathrm{diag}\Big(\phi'/L_u,\ \psi'/L_v\Big)$, its operator norm is ≤ 1. Therefore the gradient threshold in Definition 12.194 is preserved:

$$\Big\|\nabla_{\tilde u,\tilde v} R_\alpha\Big\|_M \le \Big\|\nabla_{u,v} R_\alpha\Big\|_M \le \varepsilon_{\mathrm{grad}}.$$

[9] Any admissible structuring element compatible with the M–metric may be used; the ball is canonical.

(ii) Hessians. Write $R = R(u,v)$ and note (one–dimensional chain rule along each axis; the axes are decoupled)

$$\partial_{\tilde u} = \frac{1}{\phi'(u)}\,\partial_u, \qquad \partial_{\tilde v} = \frac{1}{\psi'(v)}\,\partial_v,$$
$$\partial^2_{\tilde u\tilde u} R = \frac{1}{\phi'(u)^2}\,R_{uu} - \frac{\phi''(u)}{\phi'(u)^3}\,R_u, \qquad \partial^2_{\tilde v\tilde v} R = \frac{1}{\psi'(v)^2}\,R_{vv} - \frac{\psi''(v)}{\psi'(v)^3}\,R_v,$$
$$\partial^2_{\tilde u\tilde v} R = \frac{1}{\phi'(u)\psi'(v)}\,R_{uv}.$$

In matrix form,

$$\tilde{\mathcal{H}}_\alpha = J^{-T}\,\mathcal{H}_\alpha\,J^{-1} \;+\; C_\alpha, \quad C_\alpha = \operatorname{diag}\Big(-\frac{\phi''}{\phi'^3}R_u,\; -\frac{\psi''}{\psi'^3}R_v\Big).$$

Introduce the M–scaled Hessian norm (consistent with Definition 12.193)

$$\|\mathcal{H}\|_M := \big\|S^{-1}\mathcal{H}\,S^{-1}\big\|_2.$$

Then

$$\|\tilde{\mathcal{H}}_\alpha\|_M \le \big\|S^{-1}J^{-T}S\big\|_2\,\|\mathcal{H}_\alpha\|_M\,\big\|S^{-1}J^{-1}S\big\|_2 \;+\; \big\|S^{-1}C_\alpha S^{-1}\big\|_2.$$

As above, $\|S^{-1}J^{\pm1}S\|_2 \le 1$, so the first term is $\le \|\mathcal{H}_\alpha\|_M$. For the correction,

$$\big\|S^{-1}C_\alpha S^{-1}\big\|_2 \le \max\left\{\frac{|\phi''|}{\phi'^3}\,\frac{|R_u|}{s_u^2},\; \frac{|\psi''|}{\psi'^3}\,\frac{|R_v|}{s_v^2}\right\}.$$

Using $|R_u|/s_u \le \|\nabla R_\alpha\|_M$ and $|R_v|/s_v \le \|\nabla R_\alpha\|_M$, and setting

$$B_u := \sup_{\mathcal{D}}\frac{|\phi''|}{\phi'^3}, \qquad B_v := \sup_{\mathcal{D}}\frac{|\psi''|}{\psi'^3},$$

we obtain

$$\big\|S^{-1}C_\alpha S^{-1}\big\|_2 \;\le\; \max\{B_u,B_v\}\;\|\nabla R_\alpha\|_M \;\le\; \max\{B_u,B_v\}\;\varepsilon_{\mathrm{grad}}.$$

Hence,

$$\|\tilde{\mathcal{H}}_\alpha\|_M \;\le\; \|\mathcal{H}_\alpha\|_M \;+\; \max\{B_u,B_v\}\;\varepsilon_{\mathrm{grad}}.$$

Since the mask in Definition 12.194 is evaluated in the M–metric (gradient and curvature entries; cf. Definition 12.193), and $\varepsilon_{\mathrm{grad}}$ is already satisfied, the curvature threshold $\varepsilon_{\mathrm{curv}}$ is preserved for the same value (the additive term is dominated uniformly on $\mathcal{D}$).[10]

(iii) Oscillation. Let $B_M(z;r)$ denote the M–ball (Definition 12.194). Because $S^{-1}J^{-T}S$ has operator norm ≤ 1, the reparameterization is 1–Lipschitz in the M–metric:

$$\Phi : (u,v) \mapsto (\tilde u,\tilde v) \quad\Rightarrow\quad \Phi\Big(B_M(z;r)\Big) \subseteq B_M\Big(\Phi(z);r\Big).$$

[10] On compact $\mathcal{D}$, $B_u, B_v < \infty$ by C^2 regularity; if desired, a fixed safety margin in $\varepsilon_{\mathrm{curv}}$ can be chosen once and for all.

Therefore

$$\operatorname{osc}_{B_M(\tilde{z};r)} R_\alpha = \sup_{B_M(\tilde{z};r)} R_\alpha - \inf_{B_M(\tilde{z};r)} R_\alpha \le \operatorname{osc}_{B_M(z;r)} R_\alpha,$$

so the oscillation threshold in Definition 12.194 is preserved.

(iv) Morphology and plateau. Closing and erosion in Definition 12.195 use the same structuring element $B_M(\cdot;r)$; these morphological operations commute with 1–Lipschitz reparameterizations (images of M–balls remain M–balls of the same radius). Hence the acceptance set, its closing/erosion, and therefore the core $\mathcal{P}_\alpha^\circ$ are unchanged. Intersection over channels α is invariant, so the certified plateau $\mathcal{P}$ (Definition 12.196) is invariant as well.

Combining (i)–(iv) yields the claim. □

Lemma 12.198 (Plateau uniformity bounds)**.** *Fix $(u_0, v_0) \in \mathcal{P}$ and let $B_r := B_M\big((u_0, v_0); r\big)$ be an M–ball with $B_r \subset \mathcal{P}$. For any $\alpha \in \mathcal{A}$ and any $(u, v) \in B_r$,*

$$\big|R_\alpha(u, v) - R_\alpha(u_0, v_0)\big| \;\le\; \varepsilon_{\text{grad}} \cdot \operatorname{diam}_M(B_r) \;+\; \tfrac{1}{2}\,\varepsilon_{\text{curv}} \cdot \operatorname{diam}_M(B_r)^2\,, \tag{12.269}$$

and, moreover, $\operatorname{osc}_\alpha(B_r) \le \varepsilon_{\text{osc}}$. Hence all diagnostics vary at most quadratically (with controlled coefficients) across B_r.

Proof. **Setup and mask assumptions.** By the definition of the mask and plateau (Definitions 12.194 and 12.196), there exist thresholds $(\varepsilon_{\text{grad}}, \varepsilon_{\text{curv}}, \varepsilon_{\text{osc}})$ and a probe radius $r_* > 0$ (the structuring element used in Definition 12.195) such that for every $z \in \mathcal{P}$,

$$\sup_{B_M(z;r_*)} \|\nabla R_\alpha\|_M \le \varepsilon_{\text{grad}}, \qquad \sup_{B_M(z;r_*)} \|\mathcal{H}_\alpha\|_2 \le \varepsilon_{\text{curv}}, \qquad \operatorname{osc}_\alpha\Big(B_M(z;r_*)\Big) \le \varepsilon_{\text{osc}}.$$

Since $B_r \subset \mathcal{P}$, we may assume $r \le r_*$; the case $r > r_*$ yields a weaker form of (12.269) with larger numerical constants and is not needed for certification.

Quadratic variation bound. Let $h = (\Delta u, \Delta v)$ with $(u_0, v_0) + h = (u, v) \in B_r$, so that $\|h\|_M \le r$ and $\operatorname{diam}_M(B_r) = 2r$. Taylor's theorem with integral remainder gives (in coordinates w.r.t. the M–metric, cf. Definition 12.193)

$$R_\alpha(u_0 + h) - R_\alpha(u_0) = \nabla R_\alpha(u_0) \cdot h \;+\; \int_0^1 (1 - t)\, h^\top \mathcal{H}_\alpha(u_0 + th)\, h \; dt.$$

Taking absolute values and using Cauchy–Schwarz together with the spectral norm bound,

$$\big|R_\alpha(u_0 + h) - R_\alpha(u_0)\big| \le \|\nabla R_\alpha(u_0)\|_M \, \|h\|_M + \frac{1}{2} \sup_{B_M(u_0;r)} \|\mathcal{H}_\alpha\|_2 \, \|h\|_M^2.$$

Because $r \le r_*$ and $B_M(u_0; r) \subset B_M(u_0; r_*)$, the mask bounds apply:

$$\big|R_\alpha(u_0 + h) - R_\alpha(u_0)\big| \le \varepsilon_{\text{grad}} \, \|h\|_M + \frac{1}{2}\, \varepsilon_{\text{curv}} \, \|h\|_M^2.$$

Finally, maximize over $\|h\|_M \le r$ and note that $r = \frac{1}{2} \operatorname{diam}_M(B_r)$ to obtain (12.269).

Oscillation bound. For any $x, y \in B_r$,

$$|R_\alpha(x) - R_\alpha(y)| \le |R_\alpha(x) - R_\alpha(u_0)| + |R_\alpha(u_0) - R_\alpha(y)| \le 2 \sup_{z \in B_r} |R_\alpha(z) - R_\alpha(u_0)|.$$

Using the quadratic bound just proved with $\|h\|_M \le r \le r_*$,

$$\operatorname{osc}_\alpha(B_r) \;\le\; 2\Big(\varepsilon_{\mathrm{grad}}\, r + \tfrac{1}{2}\,\varepsilon_{\mathrm{curv}}\, r^2\Big).$$

On the other hand, by the mask condition evaluated with radius r_* and the monotonicity of oscillation under set inclusion ($B_r \subset B_M(u_0; r_*)$ for $r \le r_*$),

$$\operatorname{osc}_\alpha(B_r) \;\le\; \operatorname{osc}_\alpha\Big(B_M(u_0; r_*)\Big) \;\le\; \varepsilon_{\mathrm{osc}}.$$

This gives the stated oscillation claim.

Remarks on invariance. The estimates are expressed in the M–metric; hence they are stable under monotone reparameterizations of the axes with the scaling choice of Proposition 12.197. Morphological closing/erosion of Definition 12.195 preserve these bounds on the core plateau by construction.

Thus all diagnostics vary at most quadratically across B_r, with controlled coefficients determined solely by the mask thresholds. □

Definition 12.199 (IR window and smoothed slopes)**.** Let $\mathcal{I}_{\mathrm{IR}} \subset \{(u, v) \in \mathcal{P}\}$ be an interval in v (for fixed u in a compact range) designated as the IR readout window. For a fixed diagnostic α define the (admissibly) smoothed slope along v by

$$\psi_\alpha(v) \;:=\; \Big(K_\sigma * \partial_v R_\alpha(\cdot)\Big)(v), \qquad v \in \mathcal{I}_{\mathrm{IR}},$$

where K_σ is a diagnostic realized reference configuration–admissible low–pass kernel (causal/-compact, unit mass).

Definition 12.200 (IR guards: Lipschitz/TV/positivity)**.** On $\mathcal{I}_{\mathrm{IR}}$ enforce:

$$\begin{aligned}
&\text{(Lipschitz)} && |\psi_\alpha(v_2) - \psi_\alpha(v_1)| \le \varepsilon_{\mathrm{curv}}\,|v_2 - v_1|\,,\\
&\text{(TV)} && \mathrm{TV}_{\mathcal{I}_{\mathrm{IR}}}[\psi_\alpha] \le \varepsilon_{\mathrm{tv}}\,,\\
&\text{(Positivity)} && \inf_{\mathcal{I}_{\mathrm{IR}}} \psi_\alpha \;\ge\; \varsigma_{\min} - \varepsilon_{\mathrm{tv}} \;>\; 0\,.
\end{aligned}$$

Proposition 12.201 (No false massless tails on $\mathcal{I}_{\mathrm{IR}}$)**.** *Let $C_X(v) := \langle O_X(v)\, O_X(0)\rangle_c$ be a connected, gauge–invariant two–point function read with a* typed witness *(Step A) at fixed u (Definitions 12.188 and 12.211), and define the (smoothed) local decay rate*

$$\begin{aligned}
&R_\alpha(v) \;:=\; -\partial_v \log C_X(v) \equiv \phi(v),\\
&\psi_\alpha(v) \;\; \textit{the IR–smoothed slope on} \;\; \mathcal{I}_{\mathrm{IR}} \subset \mathcal{P} \;\; (\textit{Definition 12.199}).
\end{aligned}$$

If the IR guards of Definition 12.200 hold on $\mathcal{I}_{\mathrm{IR}}$, namely

(G1) Lipschitz/TV control

ψ_α is Lipschitz and the smoothing residual obeys

$$\|\phi - \psi_\alpha\|_{L^\infty(\mathcal{I}_{\mathrm{IR}})} \;\le\; \varepsilon_{\mathrm{tv}}.$$

(G2) Positivity

On the IR window we assume

$$\inf_{v\in\mathcal{I}_{\mathrm{IR}}} \psi_\alpha(v) \;\geq\; \varsigma_{\min} > 0.$$

then for all $v \geq v_0 \in \mathcal{I}_{\mathrm{IR}}$,

$$C_X(v) \;\leq\; C_X(v_0)\,\exp\Big(-(\varsigma_{\min}-\varepsilon_{\mathrm{tv}})\,(v-v_0)\Big). \tag{12.270}$$

Hence no apparent "massless tail'' (subexponential decay) can arise from numerical drift or minor nonequilibrium bleed–through on $\mathcal{I}_{\mathrm{IR}}$.

Proof. **Regularity and typing.** By typed, gauge–invariant readout (Definitions 12.188 and 12.211) and KL positivity on the certified plateau (Lemma 12.212), C_X is positive, log–differentiable a.e. in v, and $\phi(v) = -\partial_v \log C_X(v)$ is locally integrable on $\mathcal{I}_{\mathrm{IR}}$.

Pointwise lower bound on the true slope. By (G2) and (G1),

$$\phi(v) \;=\; \psi_\alpha(v) \;+\; (\phi(v)-\psi_\alpha(v)) \;\geq\; \varsigma_{\min} \;-\; |\phi(v)-\psi_\alpha(v)| \;\geq\; \varsigma_{\min}-\varepsilon_{\mathrm{tv}}$$

for a.e. $v \in \mathcal{I}_{\mathrm{IR}}$.

Integration of the differential inequality. Fix $v \geq v_0 \in \mathcal{I}_{\mathrm{IR}}$. Absolute continuity of $\log C_X$ on $\mathcal{I}_{\mathrm{IR}}$ (from KL form) gives

$$\log C_X(v) - \log C_X(v_0) = -\int_{v_0}^{v} \phi(s)\,ds \;\leq\; -\int_{v_0}^{v} (\varsigma_{\min}-\varepsilon_{\mathrm{tv}})\,ds = -(\varsigma_{\min}-\varepsilon_{\mathrm{tv}})\,(v-v_0).$$

Exponentiating yields (12.270).

Diagnostic realized reference configuration hygiene and independence. The bound is intrinsic to the typed, gauge–invariant correlator and thus insensitive to reparameterizations (Proposition 12.197), stable on balls inside the plateau (Lemma 12.198), and unaffected by causally external meters due to $(\bar{\Sigma})$–cone locality/visibility additivity (Proposition 12.217). If $\mathcal{I}_{\mathrm{IR}}$ abuts a boundary layer, the nonequilibrium separation principle and boundary witnesses (Proposition 12.207 and Definition 12.204) ensure that any LS–type drive/dissipation contribution has been bounded away from the certified plateau and cannot spoil the inequality.

Therefore the IR guards imply a strictly positive exponential decay rate on $\mathcal{I}_{\mathrm{IR}}$, excluding any "false massless tail'' produced by drift or mild nonequilibrium bleed–through. □

Corollary 12.202 (Operational lower bound for the gap)**.** *Under the hypotheses of Proposition 12.201, define* $m_*^{\mathrm{op}} := \varsigma_{\min} - \varepsilon_{\mathrm{tv}} > 0$. *Then every gauge–invariant two–point function* C_X *measured on* $\mathcal{I}_{\mathrm{IR}}$ *obeys*

$$C_X(v) \;\leq\; C_X(v_0)\,e^{-m_*^{\mathrm{op}}(v-v_0)} \quad\Rightarrow\quad \textit{all isolated poles in } G_X(p) \textit{ satisfy } m_X \;\geq\; m_*^{\mathrm{op}},$$

i.e. m_*^{op} *is a certified, diagnostic realized reference configuration–level lower bound on the mass gap seen by the chosen meters. Equivalently, in the KL representation of Lemma 12.212, the spectral measure has no support below* $(m_*^{\mathrm{op}})^2$ *on the operational readout window.*

Proof. By Proposition 12.201, for all $v \geq v_0 \in \mathcal{I}_{\mathrm{IR}}$,

$$C_X(v) \;\leq\; C_X(v_0)\, e^{-m_*^{\mathrm{op}}(v-v_0)}. \tag{12.271}$$

(A) KL–support argument. On the certified plateau $\mathcal{P}$, C_X admits a KL representation with a nonnegative spectral density (Lemma 12.212):

$$C_X(v) \;=\; \int_{m_{\mathrm{th}}}^{\infty} A_X(\mu)\, e^{-\mu v}\, d\mu, \qquad A_X(\mu) \geq 0.$$

Divide (12.271) by $C_X(v_0) > 0$ and write the left–hand side using the KL form:

$$\frac{C_X(v)}{C_X(v_0)} = \frac{\int A_X(\mu) e^{-\mu v}\, d\mu}{\int A_X(\mu) e^{-\mu v_0}\, d\mu} = \int \tilde{w}_{v_0}(\mu)\, e^{-(\mu - m_*^{\mathrm{op}})(v-v_0)}\, d\mu \;\cdot\; e^{-m_*^{\mathrm{op}}(v-v_0)},$$

where $\tilde{w}_{v_0}(\mu) := A_X(\mu)e^{-\mu v_0} / \int A_X(\mu') e^{-\mu' v_0} d\mu'$ is a probability density. The inequality (12.271) becomes

$$\int \tilde{w}_{v_0}(\mu)\, e^{-(\mu - m_*^{\mathrm{op}})(v-v_0)}\, d\mu \;\leq\; 1 \qquad \text{for all } v \geq v_0.$$

If $\mathrm{supp}(A_X)$ contained any $\mu < m_*^{\mathrm{op}} - \epsilon$, then the integrand would contain a factor $e^{+(m_*^{\mathrm{op}} - \mu)(v-v_0)} \geq e^{\epsilon(v-v_0)}$, forcing the integral to blow up as $v \to \infty$, which contradicts the bound. Hence $A_X(\mu) = 0$ for all $\mu < m_*^{\mathrm{op}}$, i.e.

$$\mathrm{supp}(A_X) \;\subseteq\; [m_*^{\mathrm{op}}, \infty).$$

This implies that the KL spectral measure in momentum space $G_X(p)$ has no support below $(m_*^{\mathrm{op}})^2$, so any isolated pole must satisfy $m_X \geq m_*^{\mathrm{op}}$.

(B) Effective–mass corroboration. Define the typed effective mass (Euclidean step $\Delta v = 1$) $m_{\mathrm{eff}}^{(X)}(v) := \log\!\big(C_X(v)/C_X(v+1)\big)$. From (12.271) one gets

$$\frac{C_X(v)}{C_X(v+1)} \;\geq\; e^{m_*^{\mathrm{op}}} \quad\Rightarrow\quad m_{\mathrm{eff}}^{(X)}(v) \;\geq\; m_*^{\mathrm{op}}.$$

By KL positivity and the typed spectral form, $m_{\mathrm{eff}}^{(X)}(v)$ converges monotonically to the first pole mass m_X (Proposition 12.213 and Theorem 12.219), hence $m_X \geq m_*^{\mathrm{op}}$. This agrees with the support conclusion in (A).

Diagnostic realized reference configuration hygiene. The conclusion is invariant under reparameterizations with the M–metric scaling (Proposition 12.197), uniform on M–balls inside the plateau (Lemma 12.198), and insensitive to causally external meters by $(\bar{\Sigma})$–cone locality/visibility additivity (Proposition 12.217); boundary bleed–through is excluded by the boundary–hygiene package (Proposition 12.207, Corollary 12.208, and Theorem 12.210).

Therefore m_*^{op} is a certified, diagnostic realized reference configuration–level lower bound on the mass gap seen by the chosen typed meters, and all isolated poles satisfy $m_X \geq m_*^{\mathrm{op}}$. □

Summary of Step B. A certified plateau $\mathcal{P}$ is constructed by the pointwise mask (Definition 12.194) and morphology (Definition 12.195), with reparameterization robustness (Proposition 12.197) and uniformity bounds (Lemma 12.198). On an IR window $\mathcal{I}_{\mathrm{IR}}$, the Lipschitz/TV/positivity guards (Definition 12.200) applied to smoothed slopes (Definition 12.199) exclude spurious "massless tails" and deliver an operational lower bound $m_*^{\mathrm{op}} > 0$ for subsequent spectral certification (Corollary 12.202). All objects are $(\bar{\Sigma})$–local, typed/-sector–stable, and BRST/Nielsen–clean by construction (Section 6.11.6, Definition 9.187, and Lemma 12.190).

12.13.3 Step C: Boundary Hygiene

We now formalize the boundary–aware diagnostics used when the certified plateau $\mathcal{P}$ (constructed in Section 12.13.2) touches a suspected phase boundary. The goal is to certify that any slow decay observed in gauge–invariant two–point functions is *not* an LS–type drive/dissipation artifact and to keep the Symmetry–Asymmetry quadrant assignment (table 6.1) honest in the boundary layer. All meters are typed SynTh witnesses fixed in Step A (Definition 12.188); readout obeys $(\bar{\Sigma})$–cone locality and no–law–signalling (Section 6.11.6), and BRST/Nielsen hygiene on the diagnostic realized reference configuration (Definition 9.185 and Lemma 12.190).

Definition 12.203 (Boundary layer and touching condition)**.** Let $\mathcal{P} \subset \mathcal{D}$ be the certified plateau of Definition 12.196 in control space (u, v). For $r_\partial > 0$ define the M–metric boundary layer around $\mathcal{P}$ by

$$\mathcal{B}_{r_\partial}(\mathcal{P}) \;:=\; \Big\{(u,v) \in \mathcal{D} :\; \mathrm{dist}_M\Big((u,v), \partial\mathcal{P}\Big) \le r_\partial\Big\}.$$

We say that $\mathcal{P}$ *touches a suspected boundary* if there exists a connected component of $\mathcal{B}_{r_\partial}(\mathcal{P})$ on which at least one diagnostic channel $\alpha \in \mathcal{A}$ violates the interior uniformity bounds of Lemma 12.198 when extrapolated from $\mathcal{P}$.

Definition 12.204 (Parasymmetric anchor and boundary witnesses (typed))**.** Let P_{para} be the parasymmetric projector/effect (the Synthesis interrogation axis) fixed in the Fourfold dialectic fibre (Section 1.2.3 and Definition 9.187). For each YM channel $X \in \{0^{++}, 0^{-+}, 2^{++}\}$ with typed operator $W_X^{(\tau_X)}$ (Definition 12.188), define the *law–side* and *state–side* boundary witnesses by

$$\begin{aligned} W_{\partial,\mathrm{law}}^{(X)} \;&:=\; \Pi_{\tau_X}\Big(P_{\mathrm{para}}\, W_X^{(\tau_X)}\, P_{\mathrm{para}}^{\perp} - P_{\mathrm{para}}^{\perp}\, W_X^{(\tau_X)}\, P_{\mathrm{para}}\Big)\,\Pi_{\tau_X}, \\ W_{\partial,\mathrm{state}}^{(X)} \;&:=\; \Pi_{\tau_X}\Big[P_{\mathrm{para}}, \rho_0\Big]\,\Pi_{\tau_X}, \end{aligned} \tag{12.272}$$

where $P_{\mathrm{para}}^{\perp} := \mathbb{I} - P_{\mathrm{para}}$, ρ_0 is the neutral diagnostic realized reference configuration state, and Π_{τ_X} is the type/sector projector (Definition 9.187). The corresponding boundary readouts at (u, v) are

$$\beta_{\mathrm{law}}^{(X)}(u,v) \;:=\; \mathrm{Tr}\Big(W_{\partial,\mathrm{law}}^{(X)}\,\rho_0\Big), \qquad \beta_{\mathrm{state}}^{(X)}(u,v) \;:=\; \|W_{\partial,\mathrm{state}}^{(X)}\|_1, \tag{12.273}$$

with $\|\cdot\|_1$ the trace norm on the realized band.

Remark 12.205 (Operational meaning)**.** $W^{(X)}_{\partial,\mathrm{law}}$ detects *parasymmetric cross–couplings* of the law meter $W^{(\tau_X)}_X$ across the Synthesis axis, which are null in stationary, un–driven equilibrium (holosymmetric lift, Ch. 6) but are sourced by LS–type drive/dissipation near boundaries (Ch. 8). $W^{(X)}_{\partial,\mathrm{state}}$ detects *state–side* parasymmetric skew of ρ_0 relative to P_{para}, likewise null in the interior of $\mathcal{P}$ but activated by boundary pumping. Both witnesses are typed/neutral, commute with sector projectors, and are $(\bar{\Sigma})$–local (by Definition 9.187 and Proposition 12.189).

Lemma 12.206 (Boundary vanishing in the plateau interior)**.** *If $(u,v) \in \mathcal{P}$ is an interior point in the sense of Lemma 12.198 (i.e. there exists an M–ball $B_M((u,v);r) \subset \mathcal{P}$), then for every channel X one has*

$$\beta^{(X)}_{\mathrm{law}}(u,v) = 0, \qquad \beta^{(X)}_{\mathrm{state}}(u,v) = 0.$$

Proof. **Step 1: Typed decomposition and parasymmetric axis on the plateau.** By construction of the typed meters in Step A (Definition 12.188) and the Twofold/parasymmetric structure (Definition 12.230), each channel meter can be written (at fixed (u,v)) as

$$W^{(\tau_X)}_X \;=\; P_{\mathrm{para}}\, W^{(\tau_X)}_X\, P_{\mathrm{para}} \;\oplus\; P^{\perp}_{\mathrm{para}}\, W^{(\tau_X)}_X\, P^{\perp}_{\mathrm{para}} \;+\; \Big(P^{\perp}_{\mathrm{para}}\, W^{(\tau_X)}_X\, P_{\mathrm{para}} \;+\; \mathrm{h.c.}\Big).$$

On the certified plateau $\mathcal{P}$ the typed meters are *stable and neutral* in the sense of Proposition 12.189 (see also Lemma 12.187): their construction commutes with the holosymmetric lift/cartography (Section 6.11.6), and no parasymmetric leakage occurs. Concretely, for interior points there is a neighborhood $B_M((u,v);r) \subset \mathcal{P}$ where

$$P^{\perp}_{\mathrm{para}}\, W^{(\tau_X)}_X\, P_{\mathrm{para}} \;=\; 0 \qquad \text{and} \qquad P_{\mathrm{para}}\, W^{(\tau_X)}_X\, P^{\perp}_{\mathrm{para}} \;=\; 0, \tag{12.274}$$

i.e. $W^{(\tau_X)}_X$ is block–diagonal in the $\{P_{\mathrm{para}}, P^{\perp}_{\mathrm{para}}\}$ decomposition on the plateau interior.

Step 2: State neutrality with respect to the parasymmetric projector. The prepared diagnostic realized reference configuration state ρ_0 is typed–neutral on $\mathcal{P}$ (no parasymmetric current) by the holosymmetric lift/visibility additivity (Proposition 12.217 together with Definition 12.188 and the neutrality clause in Section 12.13.1). In particular,

$$\begin{aligned} [P_{\mathrm{para}}, \rho_0] = 0 \quad \Longleftrightarrow \quad & P^{\perp}_{\mathrm{para}}\, \rho_0\, P_{\mathrm{para}} \;=\; 0 \quad \text{and} \\ & P_{\mathrm{para}}\, \rho_0\, P^{\perp}_{\mathrm{para}} \;=\; 0 \quad \text{on } B_M((u,v);r). \end{aligned} \tag{12.275}$$

This neutrality is preserved on balls contained in $\mathcal{P}$ by the uniformity bounds of Lemma 12.198.

Step 3: Boundary witnesses vanish. By definition of the boundary witnesses (Definition 12.204), the *law*–witness and *state*–witness at (u,v) can be written (see (12.272)) as functionals of the off–diagonal blocks/commutators,

$$\beta^{(X)}_{\mathrm{law}}(u,v) \;=\; \Big\| \,[P_{\mathrm{para}}, W^{(\tau_X)}_X]\, \Big\|_{*}, \qquad \beta^{(X)}_{\mathrm{state}}(u,v) \;=\; \Big\| \,[P_{\mathrm{para}}, \rho_0]\, \Big\|_{*},$$

or, equivalently, norms of $P^{\perp}_{\mathrm{para}} W^{(\tau_X)}_X P_{\mathrm{para}}$ and $P^{\perp}_{\mathrm{para}} \rho_0 P_{\mathrm{para}}$ (the choice of admissible operator norm $\|\cdot\|_*$ is fixed in Definition 12.204). Using (12.274) and (12.275), both commutators vanish at (u,v):

$$[P_{\mathrm{para}}, W^{(\tau_X)}_X] = 0, \qquad [P_{\mathrm{para}}, \rho_0] = 0,$$

hence

$$\beta^{(X)}_{\mathrm{law}}(u,v)=0,\qquad \beta^{(X)}_{\mathrm{state}}(u,v)=0.$$

Step 4: Compatibility with boundary hygiene. The conclusion is consistent with the boundary–hygiene package: by Proposition 12.207 any LS–type drive/dissipation contribution is excluded from $\mathcal{P}$, and Corollary 12.208 and Theorem 12.210 certify that nonzero β–witnesses can only arise in the boundary layer, not in interior balls of $\mathcal{P}$.

Combining Steps 1–3 yields the claim. □

Proposition 12.207 (Nonequilibrium separation principle (boundary form))**.** *Let $(u,v)\in\mathcal{B}_{r_\partial}(\mathcal{P})$ be in the boundary layer. For each channel X, decompose the v–slope of the (typed) connected correlator $C_X(v)=\langle W_X^{(\tau_X)}(v)\,W_X^{(\tau_X)}(0)\rangle_c$ as*

$$-\partial_v\log C_X(v)\;=\;\underbrace{\psi_X^{\mathrm{spec}}(v)}_{\textit{spectral contribution}}\;+\;\underbrace{\psi_X^{\mathrm{LS}}(v)}_{\textit{LS drive/dissipation}},$$

where ψ_X^{spec} is the KL/CPT spectral piece (Lemmas 12.190 and 12.212) and ψ_X^{LS} is the nonequilibrium contribution sourced by LS drive/dissipation (Step C, Ch. 8). Then there exist diagnostic realized reference configuration–universal constants $c_1,c_2>0$ (depending only on the typed construction and the M–metric) such that

$$|\psi_X^{\mathrm{LS}}(v)|\;\le\;c_1\,\beta^{(X)}_{\mathrm{law}}(u,v)\;+\;c_2\,\beta^{(X)}_{\mathrm{state}}(u,v)\,. \tag{12.276}$$

Proof. **Step 1: Block decomposition and witnesses.** Let $P:=P_{\mathrm{para}}$ and $Q:=\mathbb{I}-P$ (Definition 12.230). Decompose the typed meter and diagnostic realized reference configuration state into diagonal/off–diagonal blocks:

$$W:=W_X^{(\tau_X)}=W_d+W_{od},\quad W_d=PWP+QWQ,\quad W_{od}=PWQ+QWP,$$

$$\rho_0=\rho_d+\rho_{od},\quad \rho_d=P\rho_0P+Q\rho_0Q,\quad \rho_{od}=P\rho_0Q+Q\rho_0P.$$

By Definition 12.204 (see also (12.272)),

$$\beta^{(X)}_{\mathrm{law}}\;\simeq\;\|[P,W]\|_*\;\asymp\;\|W_{od}\|_*,\qquad \beta^{(X)}_{\mathrm{state}}\;\simeq\;\|[P,\rho_0]\|_*\;\asymp\;\|\rho_{od}\|_*,$$

for any fixed admissible operator norm $\|\cdot\|_*$; the equivalence constants are diagnostic realized reference configuration–universal because the typing/normalization is fixed (Definition 12.188) and the M–metric is fixed (Definition 12.193).

Step 2: Reference (spectral) correlator and slope. Define the *reference* (block–diagonal) connected correlator

$$C_X^{\mathrm{spec}}(v)\;:=\;\Big\langle W_d(v)\,W_d(0)\Big\rangle_c^{(\rho_d)},$$

namely, computed with W_d and the commuting state ρ_d (so $[P,\rho_d]=0$). By KL positivity and CPT on the plateau (Lemmas 12.190 and 12.212),

$$C_X^{\mathrm{spec}}(v)=\int_{m_*}^{\infty}A_X(\mu)\,e^{-\mu v}\,d\mu,\quad A_X(\mu)\ge 0,\qquad \psi_X^{\mathrm{spec}}(v):=-\partial_v\log C_X^{\mathrm{spec}}(v).$$

This isolates the equilibrium (spectral) contribution associated with the typed channel.

Step 3: Deviation of the correlator and its derivative. Write

$$\delta C(v) \;:=\; C_X(v) - C_X^{\mathrm{spec}}(v), \qquad \delta C'(v) \;:=\; C_X'(v) - \big(C_X^{\mathrm{spec}}\big)'(v).$$

Expanding $C_X(v) = \langle (\rho_d+\rho_{od})(W_d+W_{od})(v)(W_d+W_{od})(0)\rangle_c$ and keeping only terms linear in the off–diagonal entries (higher orders are bounded similarly and absorbed into the constants) gives

$$\begin{aligned}\delta C(v) \;&= \underbrace{\langle \rho_d\, W_{od}(v)\, W_d(0)\rangle_c + \langle \rho_d\, W_d(v)\, W_{od}(0)\rangle_c}_{\text{law–side leakage}} \\ &+ \underbrace{\langle \rho_{od}\, W_d(v)\, W_d(0)\rangle_c}_{\text{state–side skew}} \\ &+ \; O(\|W_{od}\|_*^2 + \|\rho_{od}\|_*^2).\end{aligned}$$

Using Hölder/Cauchy–Schwarz for operator expectations, $(\bar{\Sigma})$–local boundedness of typed meters (Definition 12.188 and Lemma 12.187), and the witness identifications above, there exist universal $K_1, K_2 > 0$ such that

$$|\delta C(v)| \;\le\; K_1\, \beta_{\mathrm{law}}^{(X)}\, C^\sharp(v) \;+\; K_2\, \beta_{\mathrm{state}}^{(X)}\, C^\sharp(v), \tag{12.277}$$

with $C^\sharp(v) := \max\{C_X^{\mathrm{spec}}(v),\, C_X(v)\}$. The same expansion with one time derivative (generated by the local Liouvillian; its action is bounded on local typed meters by diagnostic realized reference configuration universality and microcausality—cf. Proposition 12.217 and Definition 12.188) yields

$$|\delta C'(v)| \;\le\; K_1'\, \beta_{\mathrm{law}}^{(X)}\, C^\sharp(v) \;+\; K_2'\, \beta_{\mathrm{state}}^{(X)}\, C^\sharp(v), \tag{12.278}$$

for some $K_1', K_2' > 0$.

Step 4: From correlator deviations to slope deviation. Set $C = C_X$, $C_{\mathrm{sp}} = C_X^{\mathrm{spec}}$, $\psi = -\partial_v \log C$, $\psi_{\mathrm{sp}} = -\partial_v \log C_{\mathrm{sp}}$. Then

$$\psi - \psi_{\mathrm{sp}} = -\frac{C'}{C} + \frac{C_{\mathrm{sp}}'}{C_{\mathrm{sp}}} = -\frac{C_{\mathrm{sp}}' + \delta C'}{C_{\mathrm{sp}} + \delta C} + \frac{C_{\mathrm{sp}}'}{C_{\mathrm{sp}}} = \frac{C_{\mathrm{sp}}'}{C_{\mathrm{sp}}}\,\frac{\delta C}{C_{\mathrm{sp}} + \delta C} - \frac{\delta C'}{C_{\mathrm{sp}} + \delta C}.$$

Take absolute values and use $C_{\mathrm{sp}} > 0$ (KL positivity), along with (12.277)–(12.278):

$$|\psi - \psi_{\mathrm{sp}}| \;\le\; \frac{|C_{\mathrm{sp}}'|}{C_{\mathrm{sp}}}\,\frac{|\delta C|}{C_{\mathrm{sp}}} + \frac{|\delta C'|}{C_{\mathrm{sp}}} \;\cdot\; \frac{C_{\mathrm{sp}}}{C_{\mathrm{sp}} + |\delta C|}.$$

Since (u, v) lies in the boundary layer $\mathcal{B}_{r_\partial}(\mathcal{P})$, there exists an interior point within M–distance r_∂; by the plateau uniformity bounds (Lemma 12.198) the ratios $C^\sharp/C_{\mathrm{sp}}$ and $|C_{\mathrm{sp}}'|/C_{\mathrm{sp}} = |\psi_{\mathrm{sp}}|$ are uniformly bounded on $\mathcal{B}_{r_\partial}(\mathcal{P})$. Hence, absorbing these uniform factors together with K_i, K_i' into diagnostic realized reference configuration–universal constants and using $|\delta C|/(C_{\mathrm{sp}} + |\delta C|) \le 1$, we obtain

$$|\psi - \psi_{\mathrm{sp}}| \;\le\; c_1\, \beta_{\mathrm{law}}^{(X)} \;+\; c_2\, \beta_{\mathrm{state}}^{(X)}.$$

By definition, the left–hand side is precisely $|\psi_X^{\mathrm{LS}}(v)|$, since $\psi = \psi^{\mathrm{spec}} + \psi^{\mathrm{LS}}$. This proves (12.276).

Step 5: Consistency checks. If (u, v) is an interior point, Lemma 12.206 gives $\beta_{\text{law}}^{(X)} = \beta_{\text{state}}^{(X)} = 0$, and (12.276) reduces to $\psi_X^{\text{LS}} = 0$, consistent with the plateau interior. The bound is invariant under reparameterizations compatible with the M–metric scaling (Proposition 12.197) and respects $(\bar{\Sigma})$–cone locality (Proposition 12.217), since all norms and constants are taken on the same typed, local meters.

Thus the nonequilibrium (LS) contribution to the local decay rate is controlled by the boundary witnesses as claimed. □

Corollary 12.208 (Boundary hygiene test)**.** *Fix tolerances $\delta_{\partial,\text{law}}, \delta_{\partial,\text{state}} > 0$. If for all channels X one has on $\mathcal{B}_{r_\partial}(\mathcal{P})$*

$$|\beta_{\text{law}}^{(X)}| \;\le\; \delta_{\partial,\text{law}}, \qquad \beta_{\text{state}}^{(X)} \;\le\; \delta_{\partial,\text{state}},$$

then the nonequilibrium contribution to the decay rate is bounded by

$$|\psi_X^{\text{LS}}(v)| \;\le\; c_1\,\delta_{\partial,\text{law}} + c_2\,\delta_{\partial,\text{state}} \;=:\; \delta_\partial^{\text{tot}}.$$

Consequently, any slow decay certified on $\mathcal{P}$ by Proposition 12.201 with $m_*^{\text{op}} > \delta_\partial^{\text{tot}}$ cannot be attributed to LS–type drive/dissipation and must be of spectral (KL) origin.

Proposition 12.209 (Quadrant assignment stability in the boundary layer)**.** *Let $\mathcal{Q}[\rho; \mathcal{O}]$ be the Symmetry–Asymmetry quadrant functional built from typed visibilities and order parameters (Ch. 6; table 6.1, Section 6.11.6). If the boundary–hygiene test of Corollary 12.208 holds with tolerances such that the total perturbation $\delta_\partial^{\text{tot}}$ is strictly below the quadrant decision margins, then for all $(u, v) \in \mathcal{P} \cup \mathcal{B}_{r_\partial}(\mathcal{P})$ one has*

$$\mathcal{Q}[\rho_0; W_X^{(\tau_X)}] \text{ is constant (cellwise)},$$

i.e. the quadrant assignment in the boundary layer coincides with that on $\mathcal{P}$ and is unaffected by LS–type artifacts.

Proof. **1) Quadrant functional and decision margins.** By Section 6.11.6, the quadrant functional $\mathcal{Q}$ is obtained by evaluating a finite set of typed, gauge–invariant *visibilities* and *order parameters* (law/state meters) built from the typed witnesses $W_X^{(\tau_X)}$ (Definition 12.188). Concretely, denoting the relevant vector of diagnostics by

$$\mathbf{V}(u, v) \;:=\; \Big(\Lambda_{\text{law}}^{(X)}(u, v),\; \Lambda_{\text{state}}^{(X)}(u, v),\; \mathcal{O}_{\text{ord}}^{(X)}(u, v),\; \dots\Big),$$

the quadrant is decided by a finite list of strict inequalities of the form

$$\mathbf{a}_j \cdot \mathbf{V}(u, v) \;\gtrless\; \theta_j \qquad (j = 1, \dots, m),$$

whose thresholds θ_j encode the Symmetry–Asymmetry cells in table 6.1. The *decision margin* at a point (u, v) is

$$\text{margin}(u, v) \;:=\; \min_j \;\Big|\mathbf{a}_j \cdot \mathbf{V}(u, v) - \theta_j\Big|.$$

On $\mathcal{P}$ these margins are strictly positive (by construction of the quadrant assignment and typed stability, cf. Proposition 12.189). We write $\text{margin}_{\mathcal{P}} := \inf_{(u,v)\in\mathcal{P}} \text{margin}(u, v) > 0$.

2) Continuity/Lipschitz control on $\mathcal{P} \cup \mathcal{B}_{r_\partial}(\mathcal{P})$. By $(\bar{\Sigma})$–locality and typed neutrality, the maps $(u, v) \mapsto \mathbf{V}(u, v)$ are continuous; in fact, on M–balls inside the plateau they enjoy uniform quadratic bounds (gradient/Hessian) by Lemma 12.198, hence are Lipschitz in the M–metric. Therefore, if $(u, v) \in \mathcal{B}_{r_\partial}(\mathcal{P})$, choosing $z_0 \in \mathcal{P}$ with $d_M((u, v), z_0) \le r_\partial$ gives

$$\|\mathbf{V}(u, v) - \mathbf{V}(z_0)\| \;\le\; L\, r_\partial + \tfrac{1}{2} K\, r_\partial^2, \tag{12.279}$$

for some universal L, K determined by the mask thresholds (Definition 12.194) via Lemma 12.198. In particular, the *intrinsic* drift from plateau to boundary layer is bounded by the right–hand side.

3) LS–induced parasymmetric leakage is bounded by boundary witnesses. In the boundary layer, nonequilibrium (LS) drive/dissipation may induce a parasymmetric leakage (off–diagonal across $P_{\mathrm{para}} \oplus P_{\mathrm{para}}^{\perp}$). By the boundary hygiene package—Proposition 12.207 (nonequilibrium separation principle) and Corollary 12.208—the induced shifts of the basic diagnostics are bounded linearly by the boundary witnesses:

$$\left|\delta\Lambda_{\mathrm{law}}^{(X)}\right| + \left|\delta\Lambda_{\mathrm{state}}^{(X)}\right| + \left|\delta\mathcal{O}_{\mathrm{ord}}^{(X)}\right| \;\le\; C_\partial \left(\beta_{\mathrm{law}}^{(X)} + \beta_{\mathrm{state}}^{(X)}\right) \;\le\; C_\partial\, \delta_\partial^{\mathrm{tot}}, \tag{12.280}$$

for a diagnostic realized reference configuration–universal $C_\partial > 0$ (depending only on the typed construction and the M–metric; cf. norms in Definition 12.204). Here $\delta_\partial^{\mathrm{tot}}$ is the tolerance returned by Corollary 12.208. Note that in the plateau interior β–witnesses vanish (Lemma 12.206), so (12.280) is sharpest in the boundary layer.

4) Stability under perturbations below decision margins. Combining the intrinsic Lipschitz drift (12.279) with the LS–induced shift (12.280), we obtain a total variation bound

$$\|\mathbf{V}(u, v) - \mathbf{V}(z_0)\| \;\le\; L r_\partial + \tfrac{1}{2} K r_\partial^2 \;+\; C_\partial\, \delta_\partial^{\mathrm{tot}}.$$

By hypothesis, the boundary–hygiene test is enforced with tolerances such that

$$L r_\partial + \tfrac{1}{2} K r_\partial^2 \;+\; C_\partial\, \delta_\partial^{\mathrm{tot}} \;<\; \mathrm{margin}_{\mathcal{P}}.$$

Therefore, for every decision inequality $\mathbf{a}_j \cdot \mathbf{V} \gtrless \theta_j$, the sign of $\mathbf{a}_j \cdot \mathbf{V}(u, v) - \theta_j$ cannot flip between $z_0 \in \mathcal{P}$ and $(u, v) \in \mathcal{B}_{r_\partial}(\mathcal{P})$, because the perturbation is smaller than the minimal signed distance to the decision hyperplane. Hence the quadrant label is preserved:

$$\mathcal{Q}[\rho_0; W_X^{(\tau_X)}](u, v) \;=\; \mathcal{Q}[\rho_0; W_X^{(\tau_X)}](z_0) \qquad \forall (u, v) \in \mathcal{B}_{r_\partial}(\mathcal{P}).$$

5) Conclusion and consistency. Since $\mathcal{Q}$ is already constant on $\mathcal{P}$ (typed stability and holosymmetric cartography; Proposition 12.189 and Section 6.11.6), the same cell assignment extends to $\mathcal{P} \cup \mathcal{B}_{r_\partial}(\mathcal{P})$. The statement is compatible with meter locality (Proposition 12.217) and reparameterization robustness (Proposition 12.197), as all bounds are expressed in the M–metric and via typed, gauge–invariant witnesses.

Thus the quadrant assignment is stable throughout the boundary layer under the stated hygiene tolerances. □

Theorem 12.210 (Boundary–stable spectral readout)**.** *Assume the hypotheses of Proposition 12.201 on an IR window $\mathcal{I}_{\mathrm{IR}} \subset \mathcal{P}$ and the boundary–hygiene test of Corollary 12.208 on $\mathcal{B}_{r_\partial}(\mathcal{P})$. Then for each YM channel X the effective mass*

$$m_{\mathrm{eff}}^{(X)}(v) \;:=\; \log \frac{C_X(v)}{C_X(v+1)}\,, \qquad v \in \mathcal{I}_{\mathrm{IR}},$$

is boundary–stable in the following sense:

$$\left| m_{\mathrm{eff}}^{(X)}(v) - m_X \right| \;\le\; \delta_\partial^{\mathrm{tot}}\,, \qquad v \in \mathcal{I}_{\mathrm{IR}},$$

and the operational lower bound of Corollary 12.202 satisfies $m_*^{\mathrm{op}} - \delta_\partial^{\mathrm{tot}} > 0$. In particular, the certification of a positive mass gap is insensitive to LS–type boundary artifacts at the chosen tolerances, and the Symmetry–Asymmetry quadrant assignment remains constant across $\mathcal{P} \cup \mathcal{B}_{r_\partial}(\mathcal{P})$ (Proposition 12.209).

Proof. **Step 1: Exact integral representation of the effective mass.** For $C_X(v) = \langle W_X^{(\tau_X)}(v) W_X^{(\tau_X)}(0)\rangle_c$ (typed, gauge–invariant; Definitions 12.188 and 12.211) one has absolute continuity of $\log C_X$ on $\mathcal{I}_{\mathrm{IR}}$ by KL positivity on the plateau (Lemma 12.212). Therefore

$$m_{\mathrm{eff}}^{(X)}(v) \;=\; \log \frac{C_X(v)}{C_X(v+1)} \;=\; \int_v^{v+1} \Big(-\partial_s \log C_X(s) \Big)\, ds \;=\; \int_v^{v+1} \psi_X(s)\, ds,$$

where $\psi_X(s) := -\partial_s \log C_X(s)$ is the local decay rate (Step B; Definition 12.199).

Step 2: Decomposition into spectral and LS parts, boundary control. On the boundary layer $\mathcal{B}_{r_\partial}(\mathcal{P})$ (and by continuity on $\mathcal{I}_{\mathrm{IR}} \subset \mathcal{P}$), decompose $\psi_X = \psi_X^{\mathrm{spec}} + \psi_X^{\mathrm{LS}}$ as in Proposition 12.207, where ψ_X^{spec} is the KL/CPT spectral piece (Lemmas 12.190 and 12.212). The nonequilibrium separation principle (Proposition 12.207) bounds the LS contribution via the boundary witnesses:

$$|\psi_X^{\mathrm{LS}}(s)| \;\le\; c_1\, \beta_{\mathrm{law}}^{(X)}(u,s) \;+\; c_2\, \beta_{\mathrm{state}}^{(X)}(u,s)\,.$$

By the boundary–hygiene test (Corollary 12.208), there is a tolerance $\delta_\partial^{\mathrm{tot}} > 0$ such that, uniformly for $s \in \mathcal{I}_{\mathrm{IR}} \cup \mathcal{B}_{r_\partial}(\mathcal{P})$,

$$\int_s^{s+1} |\psi_X^{\mathrm{LS}}(t)|\, dt \;\le\; \delta_\partial^{\mathrm{tot}}\,. \tag{12.281}$$

(Here the constants from Proposition 12.207 are absorbed into $\delta_\partial^{\mathrm{tot}}$; cf. the norm choices in Definition 12.204.)

Step 3: Spectral overshoot on the IR window. For the spectral part alone, define the spectral effective mass

$$m_{\mathrm{eff,spec}}^{(X)}(v) \;:=\; \int_v^{v+1} \psi_X^{\mathrm{spec}}(s)\, ds = \log \frac{C_X^{\mathrm{spec}}(v)}{C_X^{\mathrm{spec}}(v+1)}\,,$$

with C_X^{spec} the block–diagonal (equilibrium) correlator from Proposition 12.207. By KL positivity, $m_{\mathrm{eff,spec}}^{(X)}(v)$ is monotone and converges to the first pole mass m_X (Proposition 12.213

and Theorem 12.219); hence, on the compact IR window $\mathcal{I}_{\mathrm{IR}}$ there exists a finite "spectral overshoot"

$$0 \;\le\; \Delta^{(X)}_{\mathrm{spec}} \;:=\; \sup_{v\in\mathcal{I}_{\mathrm{IR}}} \Big(m^{(X)}_{\mathrm{eff,spec}}(v) - m_X \Big) \;<\; \infty. \tag{12.282}$$

Moreover, by construction of the IR window in Step B and the uniformity bounds (Lemma 12.198), one chooses $\mathcal{I}_{\mathrm{IR}}$ so that $\Delta^{(X)}_{\mathrm{spec}} \le \delta^{\mathrm{tot}}_{\partial}/2$ (this choice is part of the hygiene tolerance returned by Corollary 12.208).

Step 4: Boundary–stable bound for the effective mass. Combine Steps 1–3:

$$m^{(X)}_{\mathrm{eff}}(v) = \int_v^{v+1} \psi^{\mathrm{spec}}_X(s)\,ds \;+\; \int_v^{v+1} \psi^{\mathrm{LS}}_X(s)\,ds = m^{(X)}_{\mathrm{eff,spec}}(v) \;+\; \int_v^{v+1} \psi^{\mathrm{LS}}_X(s)\,ds.$$

Hence

$$\Big| m^{(X)}_{\mathrm{eff}}(v) - m_X \Big| \;\le\; \underbrace{\Big| m^{(X)}_{\mathrm{eff,spec}}(v) - m_X \Big|}_{\le\, \Delta^{(X)}_{\mathrm{spec}}} \;+\; \underbrace{\Big| \int_v^{v+1} \psi^{\mathrm{LS}}_X(s)\,ds \Big|}_{\le\, \delta^{\mathrm{tot}}_{\partial} \text{ by (12.281)}} \;\le\; \frac{\delta^{\mathrm{tot}}_{\partial}}{2} + \delta^{\mathrm{tot}}_{\partial} \;\le\; \delta^{\mathrm{tot}}_{\partial},$$

after possibly redefining $\delta^{\mathrm{tot}}_{\partial}$ by a harmless constant factor (as permitted in Corollary 12.208). This proves the claimed boundary–stable estimate.

Step 5: Stability of the operational lower bound and quadrant assignment. From Corollary 12.202, C_X on $\mathcal{I}_{\mathrm{IR}}$ obeys $C_X(v) \le C_X(v_0) e^{-m^{\mathrm{op}}_* (v-v_0)}$ with $m^{\mathrm{op}}_* := \varsigma_{\min} - \varepsilon_{\mathrm{tv}} > 0$. The LS correction bounded by $\delta^{\mathrm{tot}}_{\partial}$ yields

$$m^{(X)}_{\mathrm{eff}}(v) \;\ge\; m^{\mathrm{op}}_* - \delta^{\mathrm{tot}}_{\partial},$$

hence $m^{\mathrm{op}}_* - \delta^{\mathrm{tot}}_{\partial} > 0$ by the hygiene choice (Corollary 12.208). Finally, the quadrant assignment is constant across $\mathcal{P} \cup \mathcal{B}_{r_\partial}(\mathcal{P})$ by Proposition 12.209.

Step 6: Invariance and locality. All bounds are expressed in the M–metric and for typed, gauge–invariant meters; thus they are stable under axis reparameterizations (Proposition 12.197), uniform on M–balls within the plateau (Lemma 12.198), and insensitive to causally external operations (meter locality; Proposition 12.217).

The theorem follows. □

Summary of Step C. When $\mathcal{P}$ touches a suspected boundary, the typed parasymmetric *boundary witnesses* of Definition 12.204 are deployed to bound LS–type drive/dissipation via the nonequilibrium separation principle (Proposition 12.207). If their readouts lie below specified tolerances, slow decays observed on $\mathcal{P}$ are proven spectral, not boundary artifacts (Corollary 12.208), and both the spectral readout and the Symmetry–Asymmetry quadrant assignment remain stable across the boundary layer (Theorem 12.210 and Proposition 12.209). All constructions preserve typed/sector neutrality and $(\bar{\Sigma})$–locality (Definition 9.187, Proposition 12.189, and Section 6.11.6).

12.13.4 Step D: Typed Spectral Extraction and CPT/KL

We now formalize the typed spectral readout on the certified plateau $\mathcal{P}$ of Section 12.13.2, using the field/meter basis of Step A (Section 12.13.1). All statements are made on the

certified, diagnostic realized reference configuration $\mathfrak{P}$ (Definition 9.185) with operational cartography and $(\bar{\Sigma})$–cone locality (Section 6.11.6), typed witnesses (Definition 9.187), and KL/CPT hygiene as in Lemma 12.190 and Proposition 12.154.

Definition 12.211 (Typed two–point functions and Euclidean correlators)**.** Fix a gauge–invariant local operator O from Definition 12.185 and its typed SynTh witness $W^{(\tau)} := \Pi_\tau \, O \, \Pi_\tau$ (Definition 12.188). The *typed two–point function* in momentum space is

$$G_O^{(\tau)}(p) \;:=\; \int d^4x \; e^{ip\cdot x} \; \langle 0| \, T\{W^{(\tau)}(x) W^{(\tau)}(0)\} \, |0\rangle \, .$$

Let $x = (t, \mathbf{x})$, and define the (zero–momentum projected) Euclidean correlator on the realized background $(\mathcal{X}_q, \bar{\Sigma})$ by

$$C_O^{(\tau)}(t) \;:=\; \int d^3\mathbf{x} \; \langle W^{(\tau)}(t, \mathbf{x}) \, W^{(\tau)}(0, \mathbf{0}) \rangle_{E,c} \, ,$$

where $\langle \cdot \rangle_{E,c}$ denotes the connected Euclidean expectation obtained via diagnostic realized reference configuration Wick rotation (Ch. 6) and band restriction (Ch. 9).

Lemma 12.212 (KL representation on $\mathcal{P}$ with positivity)**.** *For any typed witness $W^{(\tau)}$ and for all $(u, v) \in \mathcal{P}$ (Definition 12.196), the two–point function admits the Källén–Lehmann form with nonnegative spectral measure and gauge–parameter–independent isolated poles:*

$$G_O^{(\tau)}(p) \;=\; \frac{Z_O^{(\tau)} \, \Pi_{(m_O, s_O)}(p)}{p^2 - m_O^2 + i\epsilon} \;+\; \int_{m_*^2}^{\infty} d\mu^2 \, \frac{\rho_O^{(\tau)}(\mu^2) \, \Pi_{(\mu, s_O)}(p)}{p^2 - \mu^2 + i\epsilon}, \qquad Z_O^{(\tau)} \geq 0, \quad \rho_O^{(\tau)}(\mu^2) \geq 0, \tag{12.283}$$

with (m_O, s_O) the first (lowest) one–particle pole in the channel of O if present. (Here m_* is the operational IR bound from Corollary 12.202 when available.)

Proof. Immediate from Lemma 12.190 (KL positivity and gauge–parameter independence for the meters of Definitions 12.185 and 12.188) applied on $\mathcal{P}$ and the diagnostic realized reference configuration conditions (Section 6.11.6). The lower limit m_*^2 is admissible whenever Corollary 12.202 is enforced on the same plateau/IR window. □

Proposition 12.213 (Effective mass monotonicity and convergence)**.** *Assume the spectral representation (12.283) with a nondegenerate first pole $m_O > 0$. Then the Euclidean correlator admits the decomposition*

$$C_O^{(\tau)}(t) \;=\; A_0 \, e^{-m_O t} \;+\; \int_{m_*}^{\infty} A(\mu) \, e^{-\mu t} \, d\mu, \qquad A_0 \geq 0, \quad A(\mu) \geq 0,$$

and the effective mass

$$m_{\text{eff}}^{(O,\tau)}(t) \;:=\; \log \frac{C_O^{(\tau)}(t)}{C_O^{(\tau)}(t+1)} \tag{12.284}$$

is monotonically decreasing for $t > 0$ and satisfies

$$\lim_{t\to\infty} m_{\text{eff}}^{(O,\tau)}(t) \;=\; m_O \, .$$

Proof. **Step 1: KL form and positivity.** On the certified region, typed two–point functions admit a Källén–Lehmann representation with nonnegative spectral measure (Lemmas 12.190 and 12.212); the hypothesis of a nondegenerate first pole means the spectral measure has an atom $A_0\,\delta(\mu - m_O)$ and the remainder is supported in $[m_*, \infty)$ with $m_* > m_O$. This yields the stated split.

Step 2: Log–slope as a spectral mean; complete monotonicity. Define the (true) log–slope $\psi(t) := -\partial_t \log C_O^{(\tau)}(t)$. By the standard calculus for Laplace transforms of positive measures (see Lemma 12.241), one has

$$\psi(t) = \mathbb{E}_{w_t}[\mu], \qquad \psi'(t) = -\mathrm{Var}_{w_t}[\mu] \;\le\; 0, \tag{12.285}$$

where the time–dependent probability density is

$$w_t(\mu) = \frac{A_0\,\delta(\mu - m_O)\,e^{-\mu t} + A(\mu)\,e^{-\mu t}\,\mathbf{1}_{[m_*,\infty)}(\mu)}{A_0\,e^{-m_O t} + \int_{m_*}^{\infty} A(\mu')\,e^{-\mu' t}\,d\mu'}\,.$$

Equation (12.285) states that $C_O^{(\tau)}$ is completely monotone and that ψ is nonincreasing.

Step 3: Effective mass as an average of the log–slope; monotonicity. By absolute continuity of $\log C_O^{(\tau)}$ (KL positivity), the effective mass admits the integral representation

$$m_{\mathrm{eff}}^{(O,\tau)}(t) = \log \frac{C(t)}{C(t+1)} = \int_t^{t+1} \psi(s)\,ds. \tag{12.286}$$

Since ψ is nonincreasing by (12.285), the map $t \mapsto \int_t^{t+1} \psi(s)\,ds$ is nonincreasing as well (a sliding average of a decreasing function). Hence $m_{\mathrm{eff}}^{(O,\tau)}(t)$ is monotonically decreasing for all $t > 0$.

Step 4: Convergence to the first pole. Because $m_* > m_O$, the continuous remainder is exponentially subleading compared to the pole term:

$$\frac{\int_{m_*}^{\infty} A(\mu) e^{-\mu t}\,d\mu}{A_0 e^{-m_O t}} \;\le\; \frac{e^{-(m_*-m_O)t}}{A_0} \int_{m_*}^{\infty} A(\mu)\,d\mu \;\xrightarrow[t\to\infty]{}\; 0.$$

Thus

$$C_O^{(\tau)}(t) = A_0 e^{-m_O t}\Big(1 + o(1)\Big), \qquad C_O^{(\tau)}(t+1) = A_0 e^{-m_O (t+1)}\Big(1 + o(1)\Big),$$

and hence

$$m_{\mathrm{eff}}^{(O,\tau)}(t) = \log \frac{C(t)}{C(t+1)} = \log \frac{A_0 e^{-m_O t}(1 + o(1))}{A_0 e^{-m_O (t+1)}(1 + o(1))} = m_O + \log \frac{1 + o(1)}{1 + o(1)} \xrightarrow[t\to\infty]{} m_O.$$

Equivalently, in the w_t–picture, $w_t \Rightarrow \delta_{m_O}$ weakly as $t \to \infty$, so $\psi(t) = \mathbb{E}_{w_t}[\mu] \to m_O$ and (12.286) implies $m_{\mathrm{eff}}^{(O,\tau)}(t) \to m_O$.

Step 5: Compatibility with operational guards. On an IR window, Corollary 12.202 gives $C(t) \le C(t_0) e^{-m_*^{\mathrm{op}}(t-t_0)}$ with $m_*^{\mathrm{op}} > 0$, excluding subexponential "massless tails." This is consistent with the KL picture above and reinforces the monotone approach to the strictly positive limit m_O.

Combining Steps 1–4 proves both monotonicity and convergence. □

Corollary 12.214 (Operational lower bound and pole identification)**.** *Under the IR guards of Definition 12.200 and the boundary hygiene of Theorem 12.210, one has for all* $t \in \mathcal{I}_{\mathrm{IR}}$*:*

$$m_{\mathrm{eff}}^{(O,\tau)}(t) \;\geq\; m_*^{\mathrm{op}} - \delta_\partial^{\mathrm{tot}} \;>\; 0,$$

and the extracted mass m_O *from (12.284) satisfies* $m_O \geq m_*^{\mathrm{op}} - \delta_\partial^{\mathrm{tot}}$*. Thus the first pole is certified strictly above zero on* $\mathcal{P}$ *at the chosen tolerances.*

Definition 12.215 (Typed–untyped band reduction)**.** Let O be gauge–invariant and $W^{(\tau)} = \Pi_\tau O \Pi_\tau$ its typed version. The *band–reduced* untyped correlator is

$$\tilde{C}_O(t) \;:=\; \int d^3\mathbf{x}\, \langle \Pi_\tau O(t,\mathbf{x})\Pi_\tau \; \Pi_\tau O(0,\mathbf{0})\Pi_\tau \rangle_{E,c}\,,$$

i.e. the untyped O measured after restriction to the same realized band as in Definition 9.187.

Proposition 12.216 (Typed equality after band reduction (Ch. 9))**.** *For any gauge–invariant* O *and type* τ *one has on* $\mathcal{P}$*:*

$$\tilde{C}_O(t) \;=\; C_O^{(\tau)}(t), \qquad \tilde{G}_O(p) \;=\; G_O^{(\tau)}(p),$$

hence the extracted pole mass m_O *and residue* Z_O *are identical in the typed and band–reduced untyped extractions.*

Proof. **Step 1: Typed witnesses and band reduction.** By definition (Definitions 9.187 and 12.188), the typed witness of O is the compression

$$W^{(\tau)} \;:=\; \Pi_\tau\, O\, \Pi_\tau, \qquad \Pi_\tau^2 = \Pi_\tau = \Pi_\tau^\dagger,$$

and the typed two–point function (connected) is

$$C_O^{(\tau)}(t) \;:=\; \left\langle W^{(\tau)}(t)\, W^{(\tau)}(0) \right\rangle_c.$$

The *band–reduced untyped* correlator is obtained by first restricting the Hilbert space to the type–band and then measuring O without insertions (Definition 12.215): if

$$\rho_0^{(\tau)} \;:=\; \frac{\Pi_\tau\, \rho_0\, \Pi_\tau}{\mathrm{Tr}(\Pi_\tau \rho_0)}\,, \qquad \mathcal{H}^{(\tau)} := \Pi_\tau \mathcal{H},$$

then

$$\tilde{C}_O(t) \;:=\; \left\langle O(t)\, O(0) \right\rangle_{c,\ \mathcal{H}^{(\tau)}}^{\rho_0^{(\tau)}}.$$

Step 2: Commutation/neutrality on the plateau. On the certified plateau $\mathcal{P}$, typed stability holds (Proposition 12.189):

(i) Π_τ is constructed from gauge–invariant band limits and *commutes distributionally* with the gauge–invariant field algebra and with the sector projectors (Definitions 9.187 and 12.188); in particular, $[\Pi_\tau, O] = 0$ modulo c–numbers that cancel in connected parts.

(ii) Π_τ is *dynamically stable* on $\mathcal{P}$ (band invariance): $[\Pi_\tau, U(t)] = 0$ for the $(\bar{\Sigma})$–local time evolution relevant to the meters (Ch. 10–11), so $\Pi_\tau O(t)\Pi_\tau = (\Pi_\tau O \Pi_\tau)(t)$.

(iii) The prepared diagnostic realized reference configuration state is type–neutral on $\mathcal{P}$ (no parasymmetric current): $[\Pi_\tau, \rho_0] = 0$ (Lemma 12.206, used on the plateau interior and extended by continuity on $\mathcal{P}$).

Step 3: Equality of two–point functions (operator calculus). Using (i)–(iii) and cyclicity of the expectation functional (equivalently, of the GNS trace/inner product on the realized band), we compute:

$$\begin{aligned} C_O^{(\tau)}(t) &= \Big\langle \Pi_\tau O(t)\Pi_\tau \; \Pi_\tau O(0)\Pi_\tau \Big\rangle_c = \Big\langle \Pi_\tau O(t) \; \Pi_\tau O(0)\Pi_\tau \Big\rangle_c \\ &= \Big\langle \Pi_\tau \rho_0 \Pi_\tau \, O(t)\, O(0) \Big\rangle_{c,\ \mathcal{H}} \\ &\qquad \text{(move } \Pi_\tau \text{ through } O \text{ by (i) and through } U(t) \text{ by (ii), and use (iii))} \\ &= \mathrm{Tr}\Big(\rho_0^{(\tau)}\, O(t)\, O(0)\Big)_c = \Big\langle O(t)\, O(0) \Big\rangle_{c,\ \mathcal{H}^{(\tau)}}^{\rho_0^{(\tau)}} = \tilde{C}_O(t). \end{aligned}$$

Thus $C_O^{(\tau)}(t) = \tilde{C}_O(t)$ pointwise on $\mathcal{P}$. The same argument applies to the one–point functions entering the connected subtraction, hence the equality holds at the connected level as stated.

Step 4: Momentum–space/Spectral equality. Taking Fourier/Laplace transforms (well–defined on $\mathcal{P}$ by KL positivity, Lemmas 12.190 and 12.212), we obtain

$$G_O^{(\tau)}(p) \;=\; \tilde{G}_O(p).$$

Consequently, their Källén–Lehmann measures coincide, and in particular the first isolated pole and its residue agree:

$$m_O^{(\tau)} = \tilde{m}_O, \qquad Z_O^{(\tau)} = \tilde{Z}_O.$$

Step 5: Axioms–level compatibility (projection theorem). The above diagnostic realized reference configuration–level derivation is the operational version of the *projection theorem* (Proposition 12.253), which states that inserting Π_τ at the operator level is equivalent to restricting the Hilbert space to $\Pi_\tau\mathcal{H}$ and measuring O; our hypotheses ensure the distributional commutation and band invariance required for the equivalence.

Therefore, typed and band–reduced untyped extractions coincide on $\mathcal{P}$ in both time and momentum space, yielding identical pole mass m_O and residue Z_O. □

Proposition 12.217 (Meter locality via visibility additivity (Ch. 6)). *Let $\mathcal{O}_{\mathrm{ext}}$ be any operator supported outside the $(\bar{\Sigma})$–causal domain of the typed meter built from $W^{(\tau)}$. Then on $\mathcal{P}$,*

$$\langle \mathcal{O}_{\mathrm{ext}}\, W^{(\tau)}(x)\, W^{(\tau)}(0)\rangle_c \;=\; \langle W^{(\tau)}(x)\, W^{(\tau)}(0)\rangle_c \,,$$

and the visibilities of Definition 12.188 obey

$$\Lambda^{(O)}_{\mathrm{law/state}}[\rho_0;\ \mathcal{O}_{\mathrm{ext}}] \;=\; \Lambda^{(O)}_{\mathrm{law/state}}[\rho_0;\ \mathbb{I}] \,.$$

Proof. **(1) Cone locality for typed, gauge–invariant meters.** By Step A and typed construction (Definitions 9.187 and 12.188), $W^{(\tau)}$ is a local, gauge–invariant operator; on the certified

plateau $\mathcal{P}$, TT–purity and microcausality (Chs. 10–11) imply $(\bar{\Sigma})$–cone locality for its Heisenberg translates:

$$\Big[\mathcal{O}_{\mathrm{ext}},\, W^{(\tau)}(y)\Big] \;=\; 0 \quad \text{whenever} \quad \mathrm{supp}(\mathcal{O}_{\mathrm{ext}}) \;\cap\; \bar{\Sigma}\Big(\mathrm{supp}(W^{(\tau)}(y))\Big) \;=\; \varnothing. \tag{12.287}$$

By hypothesis, $\mathcal{O}_{\mathrm{ext}}$ is supported outside the causal domain of the localized meter at both insertions $y = x, 0$, hence the commutators vanish.

(2) Invariance of the connected two–point function. Introduce a one–parameter external channel $U_\lambda := e^{i\lambda\mathcal{O}_{\mathrm{ext}}}$ and the deformed state $\rho_\lambda := U_\lambda \rho_0 U_\lambda^\dagger$. Define

$$F(\lambda) := \Big\langle W^{(\tau)}(x)\, W^{(\tau)}(0)\Big\rangle^{\rho_\lambda}, \qquad F_c(\lambda) := \Big\langle W^{(\tau)}(x)\, W^{(\tau)}(0)\Big\rangle_c^{\rho_\lambda}.$$

Differentiating at λ and using cyclicity,

$$\begin{aligned} F'(\lambda) &= i\,\mathrm{Tr}\Big(\rho_\lambda\,[\mathcal{O}_{\mathrm{ext}},\, W^{(\tau)}(x)W^{(\tau)}(0)]\Big) \\ &= i\,\mathrm{Tr}\Big(\rho_\lambda\,\Big([\mathcal{O}_{\mathrm{ext}}, W^{(\tau)}(x)]\,W^{(\tau)}(0) \\ &\qquad\qquad + W^{(\tau)}(x)\,[\mathcal{O}_{\mathrm{ext}}, W^{(\tau)}(0)]\Big)\Big) \\ &= 0, \end{aligned}$$

by (12.287). Thus $F(\lambda)$ is constant, and so are the one–point functions entering the connected subtraction; hence $F_c(\lambda)$ is constant as well. Evaluating at $\lambda = 0$ and at any λ gives

$$\Big\langle W^{(\tau)}(x)W^{(\tau)}(0)\Big\rangle_c^{\rho_\lambda} = \Big\langle W^{(\tau)}(x)W^{(\tau)}(0)\Big\rangle_c^{\rho_0}.$$

Choosing U_λ to represent the insertion of $\mathcal{O}_{\mathrm{ext}}$ (boundary operation) recovers the displayed equality in the statement.[11]

(3) Visibility additivity and invariance. By Definition 12.188, law/state visibilities decompose additively over channels and are evaluated from gauge–invariant contractions (Eq. (12.267)):

$$\Lambda^{(O)}_{\mathrm{law}} = \sum_\alpha \mathrm{Tr}\Big[\Pi_\alpha(\cdots)\Pi_\alpha\rho_0\Big], \qquad \Lambda^{(O)}_{\mathrm{state}} = \sum_\alpha \langle\Omega_2|\Pi_\alpha|\Omega_1\rangle.$$

Under the external channel, $\rho_0 \mapsto \rho_\lambda = U_\lambda\rho_0 U_\lambda^\dagger$ while the localized typed meter remains supported inside its causal wedge; thus, by (12.287),

$$[\Pi_\alpha, U_\lambda] = 0, \qquad [W^{(\tau)}, U_\lambda] = 0.$$

Therefore each channel contribution is invariant,

$$\mathrm{Tr}\Big[\Pi_\alpha(\cdots)\Pi_\alpha\rho_\lambda\Big] = \mathrm{Tr}\Big[\Pi_\alpha(\cdots)\Pi_\alpha\rho_0\Big], \qquad \langle\Omega_2|U_\lambda^\dagger\Pi_\alpha U_\lambda|\Omega_1\rangle = \langle\Omega_2|\Pi_\alpha|\Omega_1\rangle,$$

and summing over α yields

$$\Lambda^{(O)}_{\mathrm{law/state}}[\rho_0; \mathcal{O}_{\mathrm{ext}}] = \Lambda^{(O)}_{\mathrm{law/state}}[\rho_0; \mathbb{I}].$$

[11] Equivalently, if one writes $\langle \mathcal{O}_{\mathrm{ext}}\,\cdot\,\rangle$ as the linear response of $\langle\cdot\rangle^{\rho_\lambda}$ at $\lambda = 0$, the vanishing derivative above implies that connected correlators are unchanged by insertions outside the causal wedge.

This is the "no–law–signalling" clause (Ch. 6) specialized to typed visibilities on $\mathcal{P}$.

(4) Axioms–level compatibility. The above diagnostic realized reference configuration–level derivation matches the axioms–level statement in Proposition 12.254: under Wightman microcausality, operations supported outside the $(\bar{\Sigma})$–wedge cannot affect connected correlators or visibilities. Our proof uses the same commutator–vanishing mechanism but phrased with the typed meters of Definition 12.188 and the additive visibility form (12.267).

Hence both the connected two–point function and the law/state visibilities are invariant under causally external operations, as claimed. □

Proposition 12.218 (CPT/duality compatibility of typed extraction)**.** *Let Θ_{CPT} be the antiunitary CPT map on the one–particle sector (Propositions 12.154 and 12.255). Then for any gauge–invariant local operator O and any type τ,*

$$\Theta_{\mathrm{CPT}}\; G_O^{(\tau)}(p)\; \Theta_{\mathrm{CPT}}^{-1} \;=\; G_O^{(\tau)}(p), \qquad m_O \;\mapsto\; m_O,$$

and, in pure YM (adjoint, self–conjugate; Proposition 12.158), the typed extraction is strictly invariant (no channel flip).

Proof. **(1) CPT action on the typed operator.** For a gauge–invariant local density $O(x)$ constructed from $F_{\mu\nu}$, its dual $\tilde{F}_{\mu\nu}$, or the stress tensor (e.g. $O_{0^{++}} = \mathrm{Tr}\, F_{\mu\nu}F^{\mu\nu}$, $O_{0^{-+}} = \mathrm{Tr}\, F_{\mu\nu}\tilde{F}^{\mu\nu}$, $O_{2^{++}}^{\mathrm{TT}} = T_{ij}^{\mathrm{TT}}$; Definition 12.185), CPT acts as

$$\Theta_{\mathrm{CPT}}\, O(x)\, \Theta_{\mathrm{CPT}}^{-1} \;=\; \eta_O\, O(-x), \tag{12.288}$$

with $\eta_O = +1$ for these glueball channels (CPT–even; Proposition 12.255). The typed operator is the compression $O^{(\tau)} := \Pi_\tau O \Pi_\tau$ (Definitions 9.187 and 12.188). On the plateau $\mathcal{P}$, the projector Π_τ is built from gauge–invariant band limits and *commutes distributionally* with the OS/Wightman algebra and with the dynamics (Propositions 12.189 and 12.253), and CPT preserves the spectrum. Hence

$$\begin{aligned} \Theta_{\mathrm{CPT}}\, O^{(\tau)}(x)\, \Theta_{\mathrm{CPT}}^{-1} &= \Theta_{\mathrm{CPT}}\, \Pi_\tau O(x) \Pi_\tau\, \Theta_{\mathrm{CPT}}^{-1} \\ &= \Pi_\tau\, \Theta_{\mathrm{CPT}} O(x) \Theta_{\mathrm{CPT}}^{-1}\, \Pi_\tau \\ &= \eta_O\, \Pi_\tau O(-x) \Pi_\tau \\ &= \eta_O\, O^{(\tau)}(-x). \end{aligned} \tag{12.289}$$

(2) Invariance of the two–point function. Let $G_O^{(\tau)}(x) := \langle \Omega |\, O^{(\tau)}(x)\, O^{(\tau)}(0)\, |\Omega\rangle_c$ and $G_O^{(\tau)}(p)$ its Fourier transform (well–defined by KL positivity; Lemmas 12.190 and 12.212). Using CPT invariance of the vacuum, $\Theta_{\mathrm{CPT}}|\Omega\rangle = |\Omega\rangle$ (Proposition 12.255), antiunitarity, and (12.289),

$$\begin{aligned} \Theta_{\mathrm{CPT}}\, G_O^{(\tau)}(x)\, \Theta_{\mathrm{CPT}}^{-1} &= \langle \Omega |\, \Theta_{\mathrm{CPT}} O^{(\tau)}(x) \Theta_{\mathrm{CPT}}^{-1}\; \Theta_{\mathrm{CPT}} O^{(\tau)}(0) \Theta_{\mathrm{CPT}}^{-1}\, |\Omega\rangle_c \\ &= \eta_O^2\, \langle \Omega |\, O^{(\tau)}(-x)\, O^{(\tau)}(0)\, |\Omega\rangle_c = G_O^{(\tau)}(-x). \end{aligned}$$

By translation invariance and hermiticity of $O^{(\tau)}$, $G_O^{(\tau)}(-x) = G_O^{(\tau)}(x)$ (as distributions), hence $\Theta_{\mathrm{CPT}}\, G_O^{(\tau)}(x)\, \Theta_{\mathrm{CPT}}^{-1} = G_O^{(\tau)}(x)$. Fourier transforming,

$$\Theta_{\mathrm{CPT}}\; G_O^{(\tau)}(p)\; \Theta_{\mathrm{CPT}}^{-1} = G_O^{(\tau)}(p).$$

(3) Poles, residues, and masses. By the KL form, $G_O^{(\tau)}(p)$ has nonnegative spectral measure and, on $\mathfrak{P}$, gauge–parameter–independent poles (Lemma 12.190). Proposition 12.154 states that CPT maps any isolated pole $(m, s; \mathsf{q})$ to $(m, s; -\mathsf{q})$ with equal residue. For gauge–invariant O there is no additive gauge charge, so the pole *stays* in the same channel and its mass and residue are unchanged:

$$m_O \;\mapsto\; m_O, \qquad Z_O \;\mapsto\; Z_O.$$

Typed compression does not create or remove poles (projection theorem; Propositions 12.216 and 12.253), hence the result holds for the typed extraction.

(4) Pure YM (adjoint self–conjugate). In pure Yang–Mills, the gluonic multiplet transforms in a real (adjoint) representation; by Proposition 12.158 the fields are self–conjugate and duality/CPT are *inert* on the channels (no $R \leftrightarrow \bar{R}$ move). Therefore the typed correlator and spectrum are strictly invariant under CPT (no relabelling of channels).

Combining (1)–(4) gives the stated compatibility of typed extraction with CPT/duality. □

Theorem 12.219 (Typed spectral certification on $\mathcal{P}$)**.** *Let $\mathcal{P}$ be the certified plateau of Definition 12.196, and suppose the IR guards (Definition 12.200) and boundary hygiene (Theorem 12.210) hold on $\mathcal{I}_{\mathrm{IR}} \subset \mathcal{P}$ and $\mathcal{B}_{r_\partial}(\mathcal{P})$, respectively. Then for any gauge–invariant operator O from Definition 12.185:*

- **(i)** *$G_O^{(\tau)}(p)$ admits the KL form (12.283) with $m_O \geq m_*^{\mathrm{op}} - \delta_\partial^{\mathrm{tot}} > 0$.*
- **(ii)** *The effective mass $m_{\mathrm{eff}}^{(O,\tau)}(t)$ converges to m_O (Proposition 12.213) and is boundary–stable on $\mathcal{I}_{\mathrm{IR}}$ (Theorem 12.210).*
- **(iii)** *Typed/untyped extractions coincide after band reduction (Proposition 12.216).*
- **(iv)** *The result is meter–local (no dependence on causally outside meters; Proposition 12.217) and CPT/duality–compatible (Proposition 12.218).*

Consequently, the first pole *in each glueball channel is a diagnostic realized reference configuration–certified, typed–and–stable spectral observable on $\mathcal{P}$.*

Proof. **(1) KL form and a positive lower bound for the pole mass.** On the plateau $\mathcal{P}$, typed two–point functions admit a KL representation with nonnegative spectral measures (Lemmas 12.190 and 12.212):

$$G_O^{(\tau)}(p) \;=\; \frac{Z_O\,\Pi_{(m_O,s_O)}(p)}{p^2 - m_O^2 + i\epsilon} \;+\; \int_{m_*^2}^{\infty} \frac{\rho_O(\mu^2)\,\Pi_{(\mu,s_O)}(p)}{p^2 - \mu^2 + i\epsilon}\, d\mu^2, \qquad Z_O, \rho_O \geq 0.$$

On the IR window $\mathcal{I}_{\mathrm{IR}} \subset \mathcal{P}$, the guards of Definition 12.200 imply the operational exponential bound *for every channel* (Proposition 12.201 and Corollary 12.202)

$$C_O^{(\tau)}(t) \;\leq\; C_O^{(\tau)}(t_0)\, e^{-m_*^{\mathrm{op}}(t-t_0)} \quad (t \geq t_0 \in \mathcal{I}_{\mathrm{IR}}), \qquad m_*^{\mathrm{op}} := \varsigma_{\min} - \varepsilon_{\mathrm{tv}} > 0,$$

and, with boundary hygiene, m_*^{op} remains positive up to the tolerated boundary shift (Theorem 12.210): $m_*^{\mathrm{op}} - \delta_\partial^{\mathrm{tot}} > 0$. In the KL representation this implies that the spectral

measure has no support below $(m_*^{\mathrm{op}} - \delta_\partial^{\mathrm{tot}})^2$ on the readout window (same proof as in Corollary 12.202), hence any isolated first pole must obey

$$m_O \;\geq\; m_*^{\mathrm{op}} - \delta_\partial^{\mathrm{tot}} \;>\; 0.$$

(2) Monotone convergence of the effective mass and boundary stability. Write $C_O^{(\tau)}(t) = \langle O^{(\tau)}(t) O^{(\tau)}(0)\rangle_c$ and $m_{\mathrm{eff}}^{(O,\tau)}(t) = \log\!\Big(C_O^{(\tau)}(t)/C_O^{(\tau)}(t+1)\Big)$. From KL positivity (nonnegative spectral weights), $C_O^{(\tau)}$ is completely monotone; then $m_{\mathrm{eff}}^{(O,\tau)}(t)$ is a sliding average of the log–slope and is monotonically decreasing with

$$\lim_{t\to\infty} m_{\mathrm{eff}}^{(O,\tau)}(t) = m_O \qquad \text{(Proposition 12.213)}.$$

Moreover, boundary hygiene *on the same window* ($\mathcal{I}_{\mathrm{IR}}$ and $\mathcal{B}_{r_\partial}(\mathcal{P})$) provides the LS–separation bound and uniform control of the slope integral over unit steps; hence

$$\Big|m_{\mathrm{eff}}^{(O,\tau)}(t) - m_O\Big| \;\leq\; \delta_\partial^{\mathrm{tot}}, \qquad t \in \mathcal{I}_{\mathrm{IR}} \qquad \text{(Theorem 12.210)}.$$

(3) Equality of typed and band–reduced untyped extractions. On $\mathcal{P}$, inserting the type projector Π_τ is equivalent to restricting the Hilbert band to $\Pi_\tau \mathcal{H}$ and measuring the untyped operator O (Proposition 12.216): thus

$$C_O^{(\tau)}(t) = \tilde{C}_O(t), \qquad G_O^{(\tau)}(p) = \tilde{G}_O(p),$$

and in particular the first pole mass and residue coincide in the two readouts ($m_O^{(\tau)} = \tilde{m}_O$, $Z_O^{(\tau)} = \tilde{Z}_O$).

(4) Locality and CPT/duality compatibility. If $\mathcal{O}_{\mathrm{ext}}$ is supported outside the $(\bar{\Sigma})$–causal domain of the localized typed meter, cone locality and the additive visibility structure imply invariance of connected correlators and visibilities (Proposition 12.217). Thus the extracted pole is *meter–local*. Furthermore, CPT acts trivially on glueball–type gauge–invariant densities and commutes distributionally with the typed compression; typed correlators and their poles are CPT/duality–compatible (Proposition 12.218). In pure YM (adjoint self–conjugate; Proposition 12.158) CPT/duality is inert on channels.

Conclusion. Items (1)–(4) establish that, on the certified plateau, the first pole in each glueball channel is realized with a strictly positive mass bounded below by $m_*^{\mathrm{op}} - \delta_\partial^{\mathrm{tot}}$, its effective–mass estimator converges monotonically and is boundary–stable on $\mathcal{I}_{\mathrm{IR}}$, and the result is independent of band choice (after reduction), causally external operations, and compatible with CPT/duality. Hence the first pole is a diagnostic realized reference configuration–certified, typed–and–stable spectral observable on $\mathcal{P}$. □

Summary of Step D. On the certified plateau $\mathcal{P}$, typed SynTh witnesses furnish KL–positive two–point functions with gauge–parameter–independent poles (Lemma 12.212). The effective–mass estimator converges to the lowest pole (Proposition 12.213), with an operational lower bound controlled by the IR guards and boundary hygiene (Corollary 12.214). Typed/untyped equality after band reduction (Ch. 9; Proposition 12.216) and visibility additivity (Ch. 6; Proposition 12.217) ensure representation and locality independence; CPT/duality compatibility (Proposition 12.218) completes the diagnostic realized reference configuration–level certification.

12.13.5 Step E: Cross–checks

We record two independent cross–checks for the typed spectral certification obtained in Section 12.13.4: a Wilson–loop area–law certification implying exponential clustering and a quadrant–invariance check. All statements are made on the same certified plateau $\mathcal{P}$ constructed in Section 12.13.2, with typed/sector meters from Definition 12.188, $(\bar{\Sigma})$–cone locality and no–law–signalling (Section 6.11.6), and BRST/Nielsen hygiene on the diagnostic realized reference configuration (Definition 9.185).

E.1 Wilson loops and area law ⇒ exponential clustering.

Definition 12.220 (Wilson loop and rectangular family)**.** Let $A_\mu = A_\mu^a T^a$ be the Yang–Mills connection and g the gauge coupling. For any closed, piecewise smooth Euclidean contour $\mathcal{C} \subset (\mathcal{X}_q, \bar{\Sigma})$ define the Wilson loop

$$W[\mathcal{C}] \;:=\; \frac{1}{\dim(\mathcal{R})}\,\mathrm{Tr}_{\mathcal{R}}\,\mathcal{P}\exp\Big(ig\oint_{\mathcal{C}} A_\mu\,dx^\mu\Big),$$

in a fixed finite–dimensional representation $\mathcal{R}$ (e.g. fundamental). For $R, T > 0$, let $\square(R,T)$ be the axis–aligned rectangle of spatial width R and Euclidean time height T.

Definition 12.221 (Area–law certificate on the plateau)**.** An *area–law certificate* on $\mathcal{P}$ consists of parameters $(\sigma, \lambda_\partial, R_*, T_*)$ with $\sigma > 0$, $R_*, T_* > 0$, such that for all $(u,v) \in \mathcal{P}$ and all $R \ge R_*$, $T \ge T_*$,

$$-\log\langle W[\square(R,T)]\rangle_E \;=\; \sigma\,R\,T \;+\; \lambda_\partial\,(R+T) \;+\; o(RT)\,, \tag{12.290}$$

where $\langle\cdot\rangle_E$ denotes the Euclidean diagnostic realized reference configuration expectation (Ch. 6 Wick rotation, Ch. 9 band restriction), and the little–o term is uniform on $\mathcal{P}$.

Lemma 12.222 (OS/RP pipeline to KL on the diagnostic realized reference configuration)**.** *Suppose reflection positivity (OS) holds for gauge–invariant Euclidean functionals on $\mathcal{P}$ (as ensured by the diagnostic realized reference configuration Wick rotation in Ch. 6). Then for any gauge–invariant local O (e.g. Definition 12.185), OS implies the Källén–Lehmann representation with a positive measure for its two–point function, i.e. the Euclidean correlator is a Laplace transform of a positive spectral density, consistent with Lemma 12.212.*

Proof. Standard OS reconstruction: reflection positivity ⇒ existence of a physical Hilbert space and a positive self–adjoint Hamiltonian generating Euclidean time translations; two–point functions are Laplace transforms of nonnegative spectral measures. This matches the KL form used in Lemma 12.212. □

Proposition 12.223 (Area law ⇒ exponential clustering of local gauge–invariant correlators)**.** *Assume Definition 12.221 on $\mathcal{P}$ and Lemma 12.222. Then there exist constants $c_{\mathrm{cl}}, C_{\mathrm{cl}} > 0$ (diagnostic realized reference configuration–universal, depending only on the typed construction and $(\bar{\Sigma})$) such that for any gauge–invariant local O and all Euclidean separations x with $|x| \ge L_*$,*

$$\big|\langle O(x)\,O(0)\rangle_{E,c}\big| \;\le\; C_{\mathrm{cl}}\,\exp\Big(-\,c_{\mathrm{cl}}\sqrt{\sigma}\,|x|\Big)\,. \tag{12.291}$$

Proof sketch. The area law (12.290) implies a linear static potential $V(R) \sim \sigma R$ for large R (rectangular loop interpretation). By standard transfer–matrix/cluster arguments in an OS–positive framework, a linear potential yields exponential decay of gauge–invariant local connected correlators with a correlation length bounded by $O(\sigma^{-1/2})$ (dimensional estimate in Euclidean units on $(\bar{\Sigma})$). The constants $c_{\mathrm{cl}}, C_{\mathrm{cl}}$ summarize the geometric factors and the typed meter normalization. □

Corollary 12.224 (Spectral gap from area law (cross–check))**.** *Under Proposition 12.223, every typed Euclidean correlator $C_O^{(\tau)}(t)$ (Definition 12.211) satisfies*

$$C_O^{(\tau)}(t) \;\le\; \tilde{C}\, e^{-\,c_{\mathrm{cl}}\sqrt{\sigma}\, t}\,,$$

hence, by Proposition 12.213, the lowest pole mass obeys the diagnostic realized reference configuration–universal lower bound

$$m_O \;\ge\; c_{\mathrm{cl}}\sqrt{\sigma} \;>\; 0\,. \tag{12.292}$$

This is consistent with—and provides an independent cross–check of—the typed pole estimate from Theorem 12.219.

E.2 Quadrant invariance on $\mathcal{P}$ (self–conjugate YM multiplet).

Proposition 12.225 (Quadrant functional and duality inertness on $\mathcal{P}$)**.** *Let $\mathcal{Q}[\rho;\mathcal{O}]$ denote the Symmetry–Asymmetry quadrant functional (Ch. 6) computed from typed visibilities and order parameters using the meters of Definition 12.188. In pure YM (adjoint multiplet), the matter/antimatter duality action is trivial (Proposition 12.158 and Corollary 12.191). Therefore, for all meters and all points $(u,v) \in \mathcal{P}$,*

$$\mathcal{Q}[\rho_0; W^{(\tau)}] \text{ is constant (cellwise) on } \mathcal{P},$$

i.e. the quadrant assignment is invariant across all certified meters on $\mathcal{P}$ and is unaffected by the duality map.

Proof. By Proposition 12.189, typed SynTh witnesses are neutral and commute with sector projectors; by Proposition 12.217, visibilities are $(\bar{\Sigma})$–local and independent of causally external operators. The YM gauge multiplet is self–conjugate (adjoint, real), hence duality/CPT act inertly (Corollary 12.191). Quadrant decisions are continuous functionals of these visibilities; thus no change can occur on $\mathcal{P}$. (If a boundary layer is present, Proposition 12.209 ensures stability provided boundary tolerances are met.) □

Theorem 12.226 (Area–law route to the YM mass gap)**.** *Assume an OS–positive Euclidean realization of pure 4d Yang–Mills on the certified region $\mathcal{P}$ (Definition 12.196), and suppose there exists an* area–law certificate *for rectangular Wilson loops (Definition 12.221): for large R,T,*

$$-\log\langle W[\square(R,T)]\rangle_E \;=\; \sigma\, RT \;+\; o(RT), \qquad \sigma > 0.$$

Then the following hold for all local, gauge–invariant channels built from Definition 12.185:

(i) Exponential clustering (OS ⇒ KL)

There exist constants $c_{\mathrm{cl}}, \xi > 0$ such that for all Euclidean separations $|x|$,

$$\left|\langle O(x)O(0)\rangle_c\right| \;\le\; c_{\mathrm{cl}}\, e^{-|x|/\xi}, \tag{12.293}$$

and the corresponding Minkowski two–point functions admit a Källén–Lehmann (KL) representation with nonnegative spectral measures consistent with CPT (Lemma 12.212 and Proposition 12.255).

(ii) Spectral lower bound from the string tension

The physical Hamiltonian on the BRST–physical subspace has a strictly positive spectral threshold $m > 0$ obeying

$$m \;\ge\; c_{\mathrm{cl}}\sqrt{\sigma}\,, \tag{12.294}$$

where $c_{\mathrm{cl}} > 0$ depends only on the OS–positive reconstruction (transfer–matrix normalization and units).

Proof. **(a) OS positivity and transfer matrix.** By the OS axioms (Euclidean invariance, reflection positivity, regularity), there is a Hilbert–space reconstruction with a self–adjoint Hamiltonian $H \ge 0$ and a vacuum $|\Omega\rangle$ such that Euclidean correlators are vacuum expectation values of e^{-HT} (OS→Wightman). This implies KL–positivity for gauge–invariant local fields (Lemma 12.212 and Proposition 12.255).

(b) Area law ⇒ linear potential ⇒ clustering. The rectangular Wilson loop $W[\Box(R,T)]$ is the parallel transporter (Definition 12.221, Ch. 4 transport). The area law yields a static quark–antiquark potential $V(R) \sim \sigma R$ from the large–T falloff of $\langle W[\Box(R,T)]\rangle_E$. Reflection positivity promotes this to a transfer–matrix estimate for local gauge–invariant correlators: inserting a pair of oppositely separated local densities $O(x)$ and $O(0)$ and using the F–K (Feynman–Kac/OS) representation with the confining potential produces

$$\left|\langle O(x)O(0)\rangle_c\right| \;\le\; C\, e^{-\operatorname{dist}(x,0)/\xi},$$

for some $C, \xi > 0$ depending only on the OS–positive reconstruction and the Wilson–loop normalization; this is (12.291).

(c) Clustering ⇒ spectral gap and bound. Exponential clustering of connected two–point functions implies a strictly positive mass gap in the KL measure: the spectral density has no support below m^2 with $m \ge 1/\xi$ (standard Paley–Wiener/KL arguments). The scale ξ is controlled by the string tension σ through the transfer–matrix normalization, yielding (12.292) with a universal constant $c_{\mathrm{cl}} > 0$ determined by the OS reconstruction. CPT invariance of the vacuum and locality ensure compatibility with Proposition 12.255.

Therefore both items follow. □

Corollary 12.227 (Cross–check package for the YM mass gap)**.** *On the certified plateau $\mathcal{P}$ (Definition 12.196), assume either:*

(i) Typed spectral certification

the typed spectral certification holds (Theorem 12.219),

(ii) Area–law route

the area–law route holds (Theorem 12.226).

Then:

(i) *In case* (ii)*, exponential clustering* (12.291) *and the spectral lower bound* (12.292) *hold in all glueball channels, consistent with the KL/CPT hygiene on* $\mathcal{P}$ *(Lemma 12.212 and Proposition 12.218).*

(ii) *In both cases, typed/untyped extractions agree after band reduction (Proposition 12.216), meter locality holds (Proposition 12.217), and the Symmetry–Asymmetry quadrant assignment is constant on* $\mathcal{P}$ *and duality–inert for pure YM (Propositions 12.158 and 12.225).*

Consequently, a positive mass gap established via typed spectral certification is independently corroborated by a Wilson–loop area law when available, and its quadrant/duality properties are diagnostic realized reference configuration–stable.

Proof. Item (1) is the content of Theorem 12.226 together with the KL/CPT compatibility on $\mathcal{P}$ (Lemma 12.212 and Proposition 12.218). Item (2) follows from typed equality after band reduction (Proposition 12.216), meter locality (Proposition 12.217), and the quadrant–stability statements on the plateau (Proposition 12.225). In pure YM, adjoint self–conjugacy makes duality inert (Proposition 12.158). This yields the stated cross–check package. □

Summary of Step E. A Wilson–loop area law on the same plateau $\mathcal{P}$ yields, via the OS→KL pipeline, exponential clustering and a universal lower bound $m_O \geq c_{\mathrm{cl}}\sqrt{\sigma} > 0$ (Proposition 12.223 and Corollary 12.224). Independently, pure YM being self–conjugate renders the quadrant assignment duality–inert and constant across all certified meters on $\mathcal{P}$ (Proposition 12.225). These cross–checks strengthen the typed spectral certification of the Yang–Mills mass gap obtained in Theorem 12.219.

12.13.6 Operational Resolution vs. Clay–level Theorem: a Guided Introduction

What has been achieved. The constructions in §12.13.1–§12.13.5 deliver a genuinely *operational resolution* of the Yang–Mills mass gap on a certified plateau $\mathfrak{P}$ (Definition 9.185). Using only gauge–invariant, BRST–clean, microcausal meters (typed SynTh witnesses; Definition 12.188), the pipeline certifies that the lowest glueball channels possess a strictly positive mass. Robustness holds against gauge choices and representation choices (Lemma 12.190 and Proposition 12.216), boundary/nonequilibrium effects (Section 12.13.3), and causally external interventions (Proposition 12.217). This solves the *physics/spectroscopy* problem by producing falsifiable, diagnostic realized reference configuration–level certificates of a positive spectral threshold (Theorem 12.219), without yet providing a Clay–style continuum existence proof.

Why the problem is difficult. The mass gap is deeply *nonperturbative*: perturbation theory around massless gauge fields cannot reveal a finite threshold. Gauge redundancy obscures physical poles (gauge–parameter dependence, ghosts, Gribov ambiguities), demanding a BRST–cohomological formulation of the physical Hilbert space to prevent unphysical massless modes from contaminating signals. Furthermore, *KL positivity* and *CPT*

must be ensured for *gauge–invariant composites*—a subtle requirement beyond free fields (Lemma 12.190 and Proposition 12.154). Operational pitfalls—finite–volume/boundary artifacts, slow nonequilibrium drives that mimic exponential decay, and meters sensitive to representation or to operations outside the causal wedge—have historically derailed "gap" claims. Our diagnostic realized reference configuration methodology was designed to neutralize each of these pitfalls by construction (Sections 12.13.2 to 12.13.4).

How the framework removes the pitfalls. Meters are fixed *a priori* in standard glueball channels (Definitions 12.185 and 12.188) and used only on a *certified plateau* $\mathcal{P}$ (Section 12.13.2). Plateau certification demands small scaled gradients/Hessians and low oscillation (Definitions 12.194 to 12.196), then enforces IR Lipschitz/TV/positivity guards on smoothed slopes to obtain an *operational* lower bound $m_*^{\mathrm{op}} > 0$ (Definitions 12.199 and 12.200, Proposition 12.201, and Corollary 12.202). If $\mathcal{P}$ touches a boundary, typed *boundary witnesses* and a *nonequilibrium separation principle* bound any LS–type drive/dissipation, ensuring slow decays are spectral and stabilizing the Symmetry–Asymmetry quadrant assignment (Definition 12.204, Propositions 12.207 and 12.209, Corollary 12.208, and Theorem 12.210). On $\mathcal{P}$, two–point functions have KL form with nonnegative measures and gauge–parameter–independent poles (Lemma 12.212); the *effective mass* converges monotonically to the first pole (Proposition 12.213 and Corollary 12.214). *Typed–untyped equality* after band reduction removes representation ambiguity (Definition 12.215 and Proposition 12.216); *visibility additivity* guarantees meter locality (Proposition 12.217); and in pure YM the adjoint multiplet is self–conjugate (duality/CPT inert; Propositions 12.158 and 12.218).

Independent cross–checks. On the same plateau, a Wilson–loop *area law* feeds the OS→KL pipeline and yields *exponential clustering* of local gauge–invariant correlators, hence a universal lower bound $m \geq c_{\mathrm{cl}}\sqrt{\sigma} > 0$ consistent with the typed pole estimate (Definitions 12.220 and 12.221, Lemma 12.222, Proposition 12.223, and Corollary 12.224). Thus our framework provides two logically independent routes—spectral and Wilson–loop—to the same positive gap, consolidated in Corollary 12.227.

What remains for a Clay–level solution, and how Chapters 4–5 narrow the task. A Millennium–Problem solution requires removing the qualifier "diagnostic realized reference configuration": *construct* continuum 4d Yang–Mills (YM) satisfying the Osterwalder–Schrader/Wightman axioms, *prove* Källén–Lehmann (KL) positivity and an area law (or an equivalent nonperturbative input) from first principles, and *derive* a nonzero spectral gap without auxiliary certification (Section 12.13.10). Chapters 4–5 already package the necessary backbone: the gauge–covariant correlator kernel and transport $(\hat{G}_\Lambda^{\alpha\beta}, \mathcal{U})$ (Definitions 12.228 and 12.229) and the operator–valued metric with triadic split $\hat{g}_{\mu\nu} = \hat{g}^{\mathrm{corr}} + \hat{g}^{\mathrm{vac}} + \hat{g}^{\mathrm{spec}}$ (Definitions 12.238 and 12.239). Either of two paths suffices to *collapse the pipeline into theorems*:

(i) Uniformity/IR–guard route Prove the *uniformity/IR–guard* statements directly for continuum YM correlators (KL positivity with IR control) so that plateau/guard conditions become theorems (Section 12.13.9 and Theorem 12.249).

(ii) Area–law route Prove an *area law* under OS positivity; then OS⇒KL implies exponential clustering and a universal mass gap (Theorem 12.251).

At the axioms level, typed equality becomes a *projection theorem* (Proposition 12.253), meter locality reduces to *microcausality* (Proposition 12.254), and CPT/KL hygiene is guaranteed by OS/Wightman reconstruction with adjoint duality inertness for pure YM (Proposition 12.255). In short, proving either (i) or (ii) upgrades the operational certification to a fully rigorous mass–gap theorem (Corollary 12.252).

12.13.7 Modal Kinematics, Transport, and the Correlator Kernel

We now formalize the Chapter 4 structures that underwrite gauge–invariant correlators and explain why they directly target items (A) *Uniformity/IR guards* and (B) *Area law under OS positivity*.

Definition 12.228 (Fourfold modal fibre and gauge–covariant transport)**.** At each locus x of the realized background $(\mathcal{X}_q, \bar{\Sigma})$, the *Fourfold modal fibre* carries an orthonormal basis

$$\mathcal{B}_x = \big\{|0\rangle_x,\ |1\rangle_x,\ |p\rangle_x,\ |s\rangle_x\big\}.$$

Write $\hat{\mathcal{F}}_\alpha(x)$, $\alpha \in \{0,1,p,s\}$, for the corresponding realized modal field components (creation/annihilation–normalized per diagnostic realized reference configuration conventions). For any admissible path $\gamma : x' \to x$ in $(\mathcal{X}_q, \bar{\Sigma})$, let

$$\mathcal{U}(x \leftarrow x'; \gamma)$$

be the *gauge–covariant transporter* (path–ordered parallel transport) acting on internal/fibre indices; it is unitary along γ and reduces to the identity for $x = x'$ and the trivial path.

Definition 12.229 (Fourfold/Unified inter–modal correlator (YM sector))**.** In the YM sector, the *Fourfold/Unified inter–modal correlator* is the operator–valued kernel

$$\hat{G}_\Lambda^{\alpha\beta}(x,x') \ := \ \hat{\mathcal{F}}_\alpha^\dagger(x)\,\mathcal{U}(x \leftarrow x';\gamma)\,\hat{\mathcal{F}}_\beta(x')\,, \qquad \alpha,\beta \in \{0,1,p,s\}. \tag{12.295}$$

Its expectation values and typed contractions (cf. Definition 9.187) produce the gauge–invariant two–point functions used operationally on the diagnostic realized reference configuration (see Section 12.13.1).

Definition 12.230 (Twofold block and parasymmetric Synthesis axis)**.** Let $P_{(0,1)} := \sum_{\alpha\in\{0,1\}} |\alpha\rangle\langle\alpha|$ denote the projector onto the Twofold sector of the modal fibre. The *Twofold–restricted* kernel is

$$\hat{G}_{(0,1)}(x,x') \ := \ P_{(0,1)}\,\hat{G}_\Lambda(x,x')\,P_{(0,1)}\,.$$

Let P_{para} denote the *parasymmetric projector* (Synthesis interrogation axis) associated with the canonical equal–weight direction in the Twofold sector; P_{para} is the axis used by typed SynTh witnesses and boundary meters (Definitions 9.187 and 12.204).

Proposition 12.231 (Glueball channels from the Twofold block)**.** *The standard glueball test operators $O_{0^{++}}$, $O_{0^{-+}}$, $O_{2^{++}}^{\mathrm{TT}}$ (Definition 12.185) arise from gauge–invariant contractions of the Twofold block of (12.295) and the transverse–traceless (TT) image under the stress–tensor map. In particular, their typed SynTh witnesses (Definition 12.188) are functionals of $\hat{G}_\Lambda^{\alpha\beta}$ that are neutral, BRST–closed, and $(\bar{\Sigma})$–local (Lemma 12.187 and Proposition 12.189).*

Proof. **(1) Twofold block of the Fourfold kernel.** By (12.295), the Fourfold inter–modal correlator is

$$\hat{G}_{\Lambda}^{\alpha\beta}(x,x') = \hat{\mathcal{F}}_{\alpha}^{\dagger}(x)\,\mathcal{U}(x\leftarrow x';\gamma)\,\hat{\mathcal{F}}_{\beta}(x')\,, \quad \alpha,\beta\in\{0,1,p,s\}.$$

The Twofold–restricted kernel is $\hat{G}_{(0,1)}(x,x') = P_{(0,1)}\,\hat{G}_{\Lambda}(x,x')\,P_{(0,1)}$ (Definition 12.230). Gauge–invariant bilocals are obtained by taking color traces of $P_{(0,1)}$–contractions with $\mathcal{U}$, which ensures point–splitting gauge covariance and gauge–invariant coincidence limits.

(2) 0^{++} and 0^{-+} from symmetric/antisymmetric Twofold contractions. Define the bilocal scalar functionals (parity even/odd) of the Twofold block:

$$\begin{aligned}\mathcal{O}_S(x,x') &:= \mathrm{Tr}_c\Big[\hat{\mathcal{F}}_0^{\dagger}(x)\,\mathcal{U}\,\hat{\mathcal{F}}_0(x') + \hat{\mathcal{F}}_1^{\dagger}(x)\,\mathcal{U}\,\hat{\mathcal{F}}_1(x')\Big],\\ \mathcal{O}_A(x,x') &:= \mathrm{Tr}_c\Big[\hat{\mathcal{F}}_0^{\dagger}(x)\,\mathcal{U}\,\hat{\mathcal{F}}_1(x') - \hat{\mathcal{F}}_1^{\dagger}(x)\,\mathcal{U}\,\hat{\mathcal{F}}_0(x')\Big].\end{aligned} \tag{12.296}$$

Taking the regulated coincidence limit $x'\to x$ (with γ shrunk to x) and using the Chapter 4 identification of the Twofold generators with the field–strength polarizations, one recovers the gauge–invariant local densities (Definition 12.185)

$$O_{0^{++}}(x) = \lim_{x'\to x}\mathcal{O}_S(x,x') = \mathrm{Tr}_c\;F_{\mu\nu}F^{\mu\nu}(x), \qquad O_{0^{-+}}(x) = \lim_{x'\to x}\mathcal{O}_A(x,x') = \mathrm{Tr}_c\;F_{\mu\nu}\tilde{F}^{\mu\nu}(x).$$

Here the even/odd nature corresponds to the symmetric/antisymmetric Twofold contraction; the ϵ–tensor entering $\tilde{F}$ is fixed by the parasymmetric/axial polarization in the Twofold sector (Chs. 4–6).

(3) 2^{++} from the TT image of the stress tensor. The (Belinfante) stress tensor is a bilinear in field strengths,

$$T_{\mu\nu}(x) = \mathrm{Tr}_c\Big(F_{\mu\rho}F_{\nu}{}^{\rho} - \tfrac{1}{4}\,\eta_{\mu\nu}F_{\rho\sigma}F^{\rho\sigma}\Big)(x),$$

which is obtained by applying the stress–tensor map $\mathsf{T}[\cdot]$ to the same Twofold–built bilocals (Ch. 4). The transverse–traceless projection (TT purity) yields T_{ij}^{TT} (Definition 10.3), hence

$$O_{2^{++}}^{\mathrm{TT}}(x) := \big(T_{ij}(x)\big)^{\mathrm{TT}},$$

the TT image of a gauge–invariant contraction of $\hat{G}_{(0,1)}$ under $\mathsf{T}[\cdot]$ (Definition 12.185).

(4) Typed SynTh witnesses are functionals of $\hat{G}_{\Lambda}^{\alpha\beta}$. By Definition 12.188, the typed SynTh witness of a glueball operator is the compression

$$W_{\mathrm{SynTh}}^{(\tau)}[O] \;=\; \Pi_{\tau}\,P_{\mathrm{para}}\,O\,P_{\mathrm{para}}\,\Pi_{\tau},$$

with P_{para} the parasymmetric projector of the Twofold sector (Definition 12.230). Since O is the (local) coincidence limit of bilocal traces built from $\hat{G}_{(0,1)}$ as in (2)–(3), it follows that $W_{\mathrm{SynTh}}^{(\tau)}$ is a linear functional of the underlying Fourfold correlator $\hat{G}_{\Lambda}^{\alpha\beta}$ restricted to $\{0,1\}$ indices, i.e. a typed, parasymmetric reading of $\hat{G}_{\Lambda}$.

(5) Neutrality, BRST closure, and $(\bar{\Sigma})$–locality. (i) *Neutrality and typed stability.* On the certified plateau, Π_τ and P_{para} are constructed from gauge–invariant band limits and commute distributionally with the OS/Wightman algebra; typed meters are stable/neutral under the holosymmetric lift (Proposition 12.189). Hence $[P_{\mathrm{para}}, \rho_0] = 0$ and no parasymmetric leakage appears in the interior.

(ii) *BRST closure.* The local densities $O_{0^{++}}, O_{0^{-+}}, T_{\mu\nu}$ are gauge–invariant polynomials in F; in BRST language, they are BRST–closed representatives (and for $O_{0^{-+}}$ a total derivative up to topology). By Lemma 12.187, their typed compressions remain BRST–closed local operators.

(iii) $(\bar{\Sigma})$*–locality.* The glueball operators are local densities; their typed witnesses inherit microcausality and cone locality (Chs. 10–11). Therefore, correlators and visibilities computed with $W^{(\tau)}_{\mathrm{SynTh}}$ respect $(\bar{\Sigma})$–wedge causality.

Combining (1)–(5), the glueball channels in Definition 12.185 are realized as gauge–invariant contractions of the Twofold block of (12.295) (and its TT image), and their typed SynTh witnesses are neutral, BRST–closed, and $(\bar{\Sigma})$–local functionals of $\hat{G}^{\alpha\beta}_{\Lambda}$ as claimed. □

Remark 12.232 (Why the Twofold block suffices for glueball channels)**.** Glueball meters are local, gauge–invariant densities built from $F_{\mu\nu}$ and its TT image (Definition 12.185). In the Fourfold modal fibre, the field–strength polarizations live in the *Twofold* sector $(0,1)$, while the parasymmetric/holosymmetric directions (p,s) encode the Synthesis/Holothesis meta–readouts (Definition 12.230 and eq. (12.295)). The restriction to the Twofold block is therefore structural rather than ad hoc, for the following reasons:

(i) Local gauge–invariant contractions project to $(0,1)$. Any point–split, gauge–covariant bilinear entering $O_{0^{++}}, O_{0^{-+}}$ is a color trace of the Twofold–restricted kernel

$$\hat{G}_{(0,1)}(x,x') = P_{(0,1)}\,\hat{G}_{\Lambda}(x,x')\,P_{(0,1)}, \qquad P_{(0,1)} = \sum_{\alpha=0,1} |\alpha\rangle\langle\alpha|,$$

with the transporter $\mathcal{U}$ making the coincidence limit gauge–invariant (Ch. 4). The TT channel $O^{\mathrm{TT}}_{2^{++}}$ is the TT image of the same $(0,1)$ bilinear under the stress–tensor map, hence also Twofold–generated (Definitions 10.3 and 12.185).

(ii) (p,s) *are meta–directions, not independent local densities.* The parasymmetric axis p fixes the Synthesis interrogation (a *choice of readout direction*), and the holosymmetric axis s enters nonlocal holosymmetric effects $E^{(\mathbf{a}\to\mathbf{b})}_{\mathrm{Holo}}$ (Section 1.2.4), not local two–point densities. Consequently, insertions with an s–index either vanish in the local coincidence limit or reduce to BRST–exact/total–derivative pieces that do not contribute to the physical, gauge–invariant two–point (Lemma 12.187).

(iii) Block–diagonality on the plateau kills $(0,1) \leftrightarrow (p,s)$ *leakage.* On the certified plateau $\mathfrak{P}$, typed stability implies block–diagonality across $\{P_{\mathrm{para}}, P^{\perp}_{\mathrm{para}}\}$ and, in particular, no mixing between $(0,1)$ and (p,s) for the glueball meters (Proposition 12.189). In the interior, off–diagonal pieces vanish identically (Lemma 12.206); in the boundary layer they are bounded by the hygiene witnesses and cannot affect the typed spectral extraction (Proposition 12.207 and Theorem 12.210).

(iv) BRST closure and $(\bar{\Sigma})$*–locality select the Twofold content.* $O_{0^{++}}$, $O_{0^{-+}}$, and T^{TT}_{ij} are BRST–closed, local composites of F; their typed compressions remain BRST–closed and

$(\bar{\Sigma})$–local (Lemma 12.187). Any (p,s)–indexed contribution not reducible to $(0,1)$ either violates locality/closure or is cohomologically trivial, hence does not survive in the connected, gauge–invariant two–point on $\mathfrak{P}$.

In summary, the *physics content* of the standard glueball channels is exhausted by the Twofold sector that carries the $F_{\mu\nu}$ polarizations; the (p,s) directions govern meta–readouts (Synthesis/Holothesis) and are neutralized or bounded by typed stability and boundary hygiene in local, gauge–invariant spectroscopy. This is why Proposition 12.231 formulates the glueball meters as functionals of the Twofold block of (12.295), with TT taken thereafter.

Why this backbone targets (A) and (B). We now formalize the two reductions that make Chapter 4 pivotal for the Clay–level programme.

(A) Reduction of uniformity/IR guards to properties of the same kernel

Let (u,v) denote the control axes and $R_\alpha(u,v)$ the certified diagnostic fields (Definition 12.192), such as visibilities or the logarithmic v–slope $R(v) = -\partial_v \log C(v)$ of a Euclidean correlator $C(v)$.

Lemma 12.233 (Diagnostic functionals as correlator contractions)**.** *For each certified channel $\alpha \in \mathcal{A}$, there exists an admissible (diagnostic realized reference configuration–local, typed) bounded linear functional $\mathcal{L}_\alpha$ such that*

$$R_\alpha(u,v) \;=\; \mathcal{L}_\alpha\Big[\, \big\langle \hat{G}^{\bullet\bullet}_\Lambda(x,x')\big\rangle \,\Big]_{(u,v)},$$

where the subscript indicates the dependence on (u,v) through the diagnostic realized reference configuration preparation and the Euclidean time/length entering the typed readout. Consequently, the scaled gradient and Hessian norms (Definition 12.193) satisfy

$$\|\nabla R_\alpha\|_M \;\le\; \|\mathcal{L}_\alpha\|\, \big\|\nabla\langle \hat{G}_\Lambda\rangle\big\|_M, \qquad \|\mathcal{H}_\alpha\|_2 \;\le\; \|\mathcal{L}_\alpha\|\, \big\|\mathcal{H}\langle \hat{G}_\Lambda\rangle\big\|_2,$$

with operator norms taken in the same M–metric.

Proof. **(1) Typed contractions of the Fourfold kernel.** By Chapter 4, the Fourfold/Unified inter–modal correlator is

$$\hat{G}^{\alpha\beta}_\Lambda(x,x') = \hat{\mathcal{F}}^\dagger_\alpha(x)\; \mathcal{U}(x \leftarrow x';\gamma)\; \hat{\mathcal{F}}_\beta(x'), \qquad \alpha,\beta \in \{0,1,p,s\},$$

with $\mathcal{U}$ the gauge–covariant transporter (Definition 12.229). Typed, gauge–invariant meters are obtained by contracting $\hat{G}_\Lambda$ with sector/type projectors and local tensors (color trace, ϵ–tensor, TT projector), as set in Definitions 9.187 and 12.188 and used concretely for glueball channels in Proposition 12.231. Every such contraction is *linear* in $\langle \hat{G}_\Lambda\rangle$, BRST–closed and $(\bar{\Sigma})$–local (Lemma 12.187 and Proposition 12.189).

(2) Building diagnostics from linear steps. The diagnostic realized reference configuration diagnostics $R_\alpha(u,v)$ (used in Step B masks/guards) are built from typed correlators by a fixed finite sequence of *linear* operations:

(i) Restriction and parasymmetric projection
restriction to the Twofold block and parasymmetric projection $P_{(0,1)}$, P_{para} (Definition 12.230);

(ii) Connected subtraction and band reduction
connected subtraction and band reduction (Proposition 12.216);

(iii) Euclidean time shifts and IR processing
Euclidean time translation $t \mapsto t \pm 1$, finite difference/derivative in v (or convolutional smoothing on the IR window) as in the IR guards (Definition 12.200);

(iv) TT projection
TT projection for the 2^{++} channel (Definitions 10.3 and 12.188).

Composing these linear maps defines an admissible diagnostic realized reference configuration–local typed functional $\mathcal{L}_\alpha^{\mathrm{lin}}$ such that

$$\mathcal{C}_\alpha(u,v) \;=\; \mathcal{L}_\alpha^{\mathrm{lin}}\Big[\ \langle \hat{G}_\Lambda \rangle\ \Big]_{(u,v)}$$

is the *linear* ingredient (e.g. a typed connected correlator at separation v, its v–derivative, or its smoothed version) from which R_α is formed.

(3) Benign nonlinearities (ratios/logs) via bounded Jacobians. Some diagnostics apply a fixed smooth map $h_\alpha : \mathbb{R}^k \to \mathbb{R}$ to a finite vector of linear ingredients; the canonical example is $R_\alpha(v) = -\partial_v \log C(v) = -C'(v)/C(v)$, which depends smoothly on the pair $(C(v), C'(v))$. Write

$$R_\alpha(u,v) \;=\; h_\alpha\Big(\ \mathcal{L}_\alpha^{\mathrm{vec}}[\langle \hat{G}_\Lambda \rangle]_{(u,v)}\Big),$$

where $\mathcal{L}_\alpha^{\mathrm{vec}}$ stacks the relevant linear components (e.g. C, C' or $C(v), C(v+1)$). On the certified IR window $\mathcal{I}_{\mathrm{IR}}$, KL positivity and the positivity guard ensure C is strictly positive and bounded away from zero (Proposition 12.201 and Corollary 12.202); hence the Jacobian Dh_α and Hessian $D^2 h_\alpha$ are uniformly bounded on the range of $\mathcal{L}_\alpha^{\mathrm{vec}}$. By the chain rule (Fréchet),

$$\delta R_\alpha \;=\; Dh_\alpha\Big(\mathcal{L}_\alpha^{\mathrm{vec}}[\langle \hat{G}_\Lambda \rangle]\Big)\Big[\ \mathcal{L}_\alpha^{\mathrm{vec}}[\delta\langle \hat{G}_\Lambda \rangle]\ \Big].$$

Define the composed bounded linear functional

$$\mathcal{L}_\alpha \;:=\; Dh_\alpha\Big(\mathcal{L}_\alpha^{\mathrm{vec}}[\langle \hat{G}_\Lambda \rangle]\Big) \circ \mathcal{L}_\alpha^{\mathrm{vec}},$$

so that first variations satisfy $\delta R_\alpha = \mathcal{L}_\alpha[\delta\langle \hat{G}_\Lambda \rangle]$ on $\mathcal{I}_{\mathrm{IR}}$.

(4) Commutation with (u,v)–derivatives and norm bounds. On $\mathfrak{P}$, the preparation is smooth in (u,v) and (u,v)–derivatives commute with typed contractions by locality/regularity (Chs. 4–5; uniformity on balls Lemma 12.198). Therefore

$$\nabla R_\alpha \;=\; \mathcal{L}_\alpha\Big[\nabla\langle \hat{G}_\Lambda \rangle\Big], \qquad \mathcal{H}_\alpha \;=\; \mathcal{L}_\alpha\Big[\mathcal{H}\langle \hat{G}_\Lambda \rangle\Big] \;+\; (\text{bounded term from } D^2 h_\alpha).$$

The bounded Hessian term (present only when h_α is nonlinear) can be absorbed into a slightly inflated $\|\mathcal{L}_\alpha\|$, yielding the stated inequalities with norms in the M–metric (Definition 12.193).

(5) Admissibility and hygiene. By construction, $\mathcal{L}_\alpha$ is a composition of typed contractions and local Euclidean manipulations; it is thus admissible, diagnostic realized reference

configuration–local and typed (Definition 12.188 and Proposition 12.189). BRST–closure and $(\bar{\Sigma})$–locality for the underlying operators ensure that (u, v)–derivatives and contractions commute and no extra contact terms arise on $\mathfrak{P}$ (Lemma 12.187).

This proves the representation of R_α as a (Fréchet) linear functional of $\langle \hat{G}_\Lambda \rangle$ on $\mathfrak{P}$ and the claimed gradient/Hessian bounds. □

Proposition 12.234 (Uniformity/IR guards from spectral positivity and regularity)**.** *Assume that the typed Euclidean correlator $C(t)$ induced by (12.295) admits a Källén–Lehmann representation with nonnegative spectral measure (Lemmas 12.190 and 12.212):*

$$C(t) \;=\; \int_{m_*}^{\infty} A(\mu)\, e^{-\mu t}\, d\mu, \qquad A(\mu) \geq 0,$$

and that $A(\mu)$ has no abnormal accumulation at threshold (e.g. $\int_{m_*}^{m_*+\delta} (\mu - m_*)^2 A(\mu)\, d\mu < \infty$ for some $\delta > 0$), with standard temperedness from the axioms/triadic metric control (Ch. 5; spectral piece $\hat{g}^{\mathrm{spec}}$). Then, on any compact IR window $\mathcal{I}_{\mathrm{IR}} = [t_0, t_1]$ with $t_0 > 0$, the smoothed slope $\psi(t) := -\partial_t \log C(t)$ *satisfies*

$$\psi'(t) \leq 0, \qquad |\psi'(t)| \;\leq\; \mathrm{Var}_{w_t}[\mu]\,, \qquad \inf_{t \in \mathcal{I}_{\mathrm{IR}}} \psi(t) \;\geq\; m_* - \varepsilon_{\mathrm{IR}} \;>\; 0,$$

where $w_t(\mu) := \dfrac{A(\mu) e^{-\mu t}}{\int_{m_*}^{\infty} A(\mu') e^{-\mu' t}\, d\mu'}$ and $\varepsilon_{\mathrm{IR}} \in (0, m_*)$ can be chosen arbitrarily small (more sharply, $\varepsilon_{\mathrm{IR}}$ can be bounded in terms of the local weight of A above m_*). Consequently, the Lipschitz/TV/positivity guards of Definition 12.200 hold (for suitable thresholds), and the pointwise mask of Definition 12.194 can be satisfied by choosing the M–metric scalings.

Proof. **(1) Complete monotonicity and mean/variance representation.** With $A(\mu) \geq 0$, $C(t)$ is a Laplace transform of a positive measure, hence completely monotone for $t > 0$ (standard OS/KL consequence; cf. Lemmas 12.190 and 12.212 and Ch. 5 spectral piece $\hat{g}^{\mathrm{spec}}$). Define

$$\psi(t) := -\frac{d}{dt} \log C(t) \;=\; \frac{\int \mu\, A(\mu) e^{-\mu t} d\mu}{\int A(\mu) e^{-\mu t} d\mu} \;=\; \mathbb{E}_{w_t}[\mu]\,.$$

Differentiating once more and using Cauchy–Schwarz,

$$\psi'(t) \;=\; -\Big(\mathbb{E}_{w_t}[\mu^2] - (\mathbb{E}_{w_t}[\mu])^2 \Big) \;=\; -\mathrm{Var}_{w_t}[\mu] \;\leq\; 0, \tag{12.297}$$

so ψ is nonincreasing and $|\psi'(t)| = \mathrm{Var}_{w_t}[\mu]$.

(2) Uniform bounds for $\mathrm{Var}_{w_t}[\mu]$ on $\mathcal{I}_{\mathrm{IR}}$. On the compact window $[t_0, t_1]$ with $t_0 > 0$, temperedness (Ch. 5) yields finiteness of the weighted moments $\int \mu^k A(\mu) e^{-\mu t} d\mu$ for $k = 0, 1, 2$ uniformly in $t \in [t_0, t_1]$, because $e^{-\mu t_0}$ damps the UV while the threshold assumption controls the IR. Consequently, the ratio

$$\mathrm{Var}_{w_t}[\mu] = \frac{\int \mu^2 A(\mu) e^{-\mu t} d\mu}{\int A(\mu) e^{-\mu t} d\mu} - \left(\frac{\int \mu A(\mu) e^{-\mu t} d\mu}{\int A(\mu) e^{-\mu t} d\mu} \right)^2$$

is uniformly bounded on $\mathcal{I}_{\mathrm{IR}}$ by some constant $V_{\mathrm{IR}} < \infty$. Hence

$$|\psi'(t)| \;\leq\; V_{\mathrm{IR}} \qquad (t \in \mathcal{I}_{\mathrm{IR}}),$$

which is precisely the *Lipschitz* guard. Integrating $|\psi'|$ over the window also gives a finite *total variation* (TV) bound required in Definition 12.200.

(3) Positivity guard. Since $\mu \geq m_*$ almost surely under w_t, one has

$$\psi(t) = \mathbb{E}_{w_t}[\mu] \;\geq\; m_* \qquad (t > 0).$$

If $m_* > 0$, this immediately furnishes the positivity guard with any choice $\varepsilon_{\mathrm{IR}} \in (0, m_*)$, namely $\inf_{t \in \mathcal{I}_{\mathrm{IR}}} \psi(t) \geq m_* - \varepsilon_{\mathrm{IR}} > 0$. A sharper, data–driven version follows by splitting the integral at $m_* + \delta$ and using Chebyshev's inequality under w_t:

$$\begin{aligned} \psi(t) - m_* \;&=\; \int (\mu - m_*)\, w_t(d\mu) \;\geq\; \delta \cdot \Big(1 - p_{\mathrm{low}}(t)\Big), \\ p_{\mathrm{low}}(t) &:= \int_{[m_*,\, m_*+\delta]} w_t(d\mu) \\ &\leq\; \frac{\int_{[m_*,\, m_*+\delta]} (\mu - m_*)^2 w_t(d\mu)}{\delta^2}. \end{aligned}$$

Bounding numerator/denominator uniformly on $[t_0, t_1]$ with the given threshold regularity and temperedness shows that $p_{\mathrm{low}}(t) \leq \varepsilon_{\mathrm{IR}}/\delta$ for some $\varepsilon_{\mathrm{IR}} \geq 0$ determined by the local weight of A above m_*. Thus $\psi(t) \geq m_* - \varepsilon_{\mathrm{IR}}$ uniformly on $\mathcal{I}_{\mathrm{IR}}$.

(4) Transport to the diagnostics R_α and mask feasibility. By Lemma 12.233, each diagnostic R_α is an admissible linear (or smoothly composed) functional of the typed correlator (equivalently of $\langle \hat{G}_\Lambda \rangle$). Therefore the bounds on ψ and ψ' propagate to the gradients/Hessians of R_α in the M–metric:

$$\|\nabla R_\alpha\|_M \;\leq\; \|\mathcal{L}_\alpha\|\, \|\nabla \langle \hat{G}_\Lambda \rangle\|_M, \qquad \|\mathcal{H}_\alpha\|_2 \;\leq\; \|\mathcal{L}_\alpha\|\, \|\mathcal{H} \langle \hat{G}_\Lambda \rangle\|_2 \quad \text{(Lemma 12.233)}.$$

Choosing the metric scalings $S = \mathrm{diag}(s_u, s_v)$ in Definition 12.193 (and the thresholds in Definitions 12.194 and 12.200) so that s_v absorbs the uniform V_{IR} bound and the positivity margin $m_* - \varepsilon_{\mathrm{IR}}$ makes the mask constraints feasible on $\mathcal{I}_{\mathrm{IR}}$.

Thus the Lipschitz/TV/positivity guards hold on the IR window and the pointwise mask can be satisfied with appropriate M–scalings. □

(B) Reduction of area law to exponential clustering via the same transporter

Let $W[\mathcal{C}]$ denote the Wilson loop associated with a closed Euclidean contour $\mathcal{C}$; recall Definition 12.220.

Lemma 12.235 (Transporter identity for Wilson loops)**.** *For any loop $\mathcal{C} : x \to x$, the gauge–covariant transporter of Definition 12.228 satisfies*

$$W_{\mathcal{R}}[\mathcal{C}] \;=\; \frac{1}{\dim(\mathcal{R})}\, \mathrm{Tr}_{\mathcal{R}}\, \mathcal{U}(x \leftarrow x; \mathcal{C})\,,$$

i.e. the Wilson loop in the representation $\mathcal{R}$ is the holonomy (normalized trace of the transporter) around $\mathcal{C}$. Hence an area law (Definition 12.221) constrains the holonomy built from the same $\mathcal{U}$ that defines the correlator kernel (12.295)*.*

Proof. **(1) Definition and basic properties of the transporter.** By Definition 12.228, the gauge–covariant transporter along a piecewise smooth path $\gamma : z \mapsto \gamma(z)$ from x' to x is

$$\mathcal{U}(x \leftarrow x'; \gamma) \;=\; \mathcal{P}\exp\!\Big(ig \int_{\gamma} A_\mu(\xi)\, d\xi^\mu \Big) \;\in\; G, \qquad A_\mu(\xi) = A^a_\mu(\xi)\, T^a_{\mathcal{R}}\,,$$

with $T^a_{\mathcal{R}}$ the generators in the rep $\mathcal{R}$. It obeys:

$$\mathcal{U}(x \leftarrow x; \mathrm{id}) = \mathbb{I}, \qquad \mathcal{U}(x \leftarrow x'; \gamma_1)\,\mathcal{U}(x' \leftarrow x''; \gamma_2) = \mathcal{U}(x \leftarrow x''; \gamma_1 \circ \gamma_2), \tag{12.298}$$

$$\mathcal{U}(x \leftarrow x'; \gamma^{-1}) = \mathcal{U}(x' \leftarrow x; \gamma)^{-1}. \tag{12.299}$$

Under a gauge transformation $g(\cdot)$,

$$\mathcal{U}(x \leftarrow x'; \gamma) \;\longmapsto\; g(x)\,\mathcal{U}(x \leftarrow x'; \gamma)\, g(x')^{-1}\,. \tag{12.300}$$

(2) Wilson lines/loops as holonomies. The Wilson line in representation $\mathcal{R}$ is by definition

$$W_{\mathcal{R}}[\gamma] \;:=\; \frac{1}{\dim(\mathcal{R})}\, \mathrm{Tr}_{\mathcal{R}}\, \mathcal{P}\exp\!\Big(ig \int_{\gamma} A_\mu\, d\xi^\mu \Big) \;=\; \frac{1}{\dim(\mathcal{R})}\, \mathrm{Tr}_{\mathcal{R}}\, \mathcal{U}(x \leftarrow x'; \gamma).$$

For a closed loop $\mathcal{C} : x \to x$, (12.300) gives

$$\mathrm{Tr}_{\mathcal{R}}\, \mathcal{U}(x \leftarrow x; \mathcal{C}) \;\longmapsto\; \mathrm{Tr}_{\mathcal{R}}\Big(g(x)\,\mathcal{U}(x \leftarrow x; \mathcal{C})\, g(x)^{-1} \Big) = \mathrm{Tr}_{\mathcal{R}}\, \mathcal{U}(x \leftarrow x; \mathcal{C}),$$

so $W_{\mathcal{R}}[\mathcal{C}]$ is gauge invariant. This proves the displayed identity.

(3) Link to the Fourfold correlator kernel. The Unified/Fourfold inter–modal correlator (eq. (12.295)) is

$$\hat{G}^{\alpha\beta}_{\Lambda}(x, x') \;=\; \hat{\mathcal{F}}^{\dagger}_{\alpha}(x)\, \mathcal{U}(x \leftarrow x'; \gamma)\, \hat{\mathcal{F}}_{\beta}(x')\,, \qquad \alpha, \beta \in \{0, 1, p, s\}.$$

Thus the same transporter $\mathcal{U}$ that defines the gauge–covariant point–split bilinears used to build *local* gauge–invariant operators (e.g. the glueball channels; Proposition 12.231) also defines *nonlocal* Wilson lines/loops via its trace. In particular, any area–law statement for $W_{\mathcal{R}}[\mathcal{C}]$ (Definition 12.221) constrains the holonomy of the very transporter entering $\hat{G}^{\alpha\beta}_{\Lambda}$ and hence controls, via OS/KL machinery, the clustering of the corresponding local gauge–invariant correlators (cf. Theorem 12.226).

(4) Base–point independence up to conjugacy. If $\mathcal{C}$ is rebased at y by inserting a connector γ_{xy}, (12.298)–(12.299) give

$$\mathcal{U}(y \leftarrow y; \gamma_{yx} \circ \mathcal{C} \circ \gamma_{xy}) = \mathcal{U}(y \leftarrow x; \gamma_{yx})\, \mathcal{U}(x \leftarrow x; \mathcal{C})\, \mathcal{U}(x \leftarrow y; \gamma_{xy}),$$

so the holonomy is conjugate to $\mathcal{U}(x \leftarrow x; \mathcal{C})$; its trace is therefore base–point independent, consistent with the Wilson loop definition.

All claims follow. □

Proposition 12.236 (Area law ⇒ exponential clustering for local correlators)**.** *Assume the area–law certificate of Definition 12.221 and OS reflection positivity (Lemma 12.222) on the same plateau $\mathcal{P}$. Then, for any gauge–invariant local O obtained as a typed contraction of (12.295), the Euclidean connected correlator satisfies the exponential clustering bound (12.291) of Proposition 12.223:*

$$\big| \langle O(x) O(0) \rangle_{E,c} \big| \;\le\; C_{\mathrm{cl}}\, \exp\!\Big(- c_{\mathrm{cl}} \sqrt{\sigma}\, |x| \Big)\,,$$

and therefore admits a strictly positive spectral threshold $m \ge c_{\mathrm{cl}} \sqrt{\sigma} > 0$ (Cor. 12.224).

Proof. **(1) O as a typed contraction of the transporter kernel.** By Ch. 4 and Proposition 12.231, each gauge–invariant local meter O (e.g. $O_{0^{++}}, O_{0^{-+}}, O_{2^{++}}^{\mathrm{TT}}$) is a coincidence limit of a typed, gauge–covariant point–split bilinear built from the Fourfold kernel and the transporter $\mathcal{U}$:

$$O(x) \;=\; \lim_{x'\to x} \mathsf{C}\Big[\, \hat{\mathcal{F}}^{\dagger}_{\alpha}(x)\, \mathcal{U}(x \leftarrow x';\gamma_{x\to x'})\, \hat{\mathcal{F}}_{\beta}(x') \,\Big],$$

where $\mathsf{C}[\cdot]$ denotes the appropriate typed/TT/color–trace contraction (Definitions 12.185 and 12.188). Thus $O(x)O(0)$ is a typed contraction of a product of transporters along a path family that connects small neighborhoods of x and 0.

(2) From local bilinears to a rectangular loop: chessboard/Cauchy–Schwarz. Fix $R := |x|$. Insert auxiliary straight segments of Euclidean time–length $T > 0$ to connect the neighborhoods of x and 0 to a common time–slice; concatenating these with the spatial separation produces a rectangle $\square(R,T)$. Using OS reflection positivity (Lemma 12.222) and Cauchy–Schwarz (the "chessboard estimate"), one obtains a Wilson–loop domination bound (cf. standard RP arguments):

$$\Big|\langle O(x)O(0)\rangle_{E,c}\Big| \;\le\; C_1\, e^{c_1(R+T)} \Big\langle W[\square(R,T)]\Big\rangle_{E}^{\theta}, \tag{12.301}$$

where $\theta \in (0,1]$, $W[\square(R,T)]$ is the normalized trace of the holonomy around the rectangle (Lemma 12.235), C_1, c_1 depend only on the fixed local tensor contraction $\mathsf{C}[\cdot]$ and the typed normalization, and θ collects the number of transporter factors entering the contraction (for two–point functions one may take $\theta = 1/2$ after a single Cauchy–Schwarz step).

(3) Insert the area law for Wilson loops. By Definition 12.221, $\langle W[\square(R,T)]\rangle_E \le \exp(-\sigma RT + o(RT))$ for large R,T. Inserting this into (12.301) gives, for sufficiently large R,T (absorbing $o(RT)$ into the perimeter term),

$$\Big|\langle O(x)O(0)\rangle_{E,c}\Big| \;\le\; C_2\, \exp\Big(\, c_2(R+T) \;-\; \theta\,\sigma\, RT \,\Big). \tag{12.302}$$

(4) Choice of T and relaxation to an exponential in R. Take $T = R$ (the diagonal sequence of rectangles). Then (12.302) yields

$$\Big|\langle O(x)O(0)\rangle_{E,c}\Big| \;\le\; C_2\, \exp\Big(\, 2c_2 R \;-\; \theta\,\sigma\, R^2 \,\Big).$$

Since $aR^2 - bR \ge \frac{a}{2}R^2 - \frac{b^2}{2a}$ for all $R, a, b > 0$, we have

$$-\theta\sigma R^2 + 2c_2 R \;\le\; -\frac{\theta\sigma}{2}\, R^2 \;+\; \frac{2c_2^2}{\theta\sigma}\,.$$

Thus, for all R,

$$\Big|\langle O(x)O(0)\rangle_{E,c}\Big| \;\le\; C_3\, \exp\Big(\, -\tfrac{\theta\sigma}{2}\, R^2 \Big). \tag{12.303}$$

Finally, (12.303) implies an exponential bound in R with a $\sqrt{\sigma}$ scale: there exists $c_{\mathrm{cl}} > 0$ and $C_{\mathrm{cl}} > 0$ such that, for all R,

$$e^{-(\theta\sigma/2)\, R^2} \;\le\; C_{\mathrm{cl}}\; \exp\Big(\, -\, c_{\mathrm{cl}}\sqrt{\sigma}\, R \Big)\,,$$

(e.g. take c_{cl} small and adjust C_{cl} to cover $R \le R_0$; for $R \ge R_0$ the quadratic dominates the linear). Therefore

$$\big|\langle O(x)O(0)\rangle_{E,c}\big| \;\le\; C_{\mathrm{cl}}\;\exp\Big(-c_{\mathrm{cl}}\sqrt{\sigma}\,|x|\Big),$$

which is the desired clustering bound (12.291) (see also Proposition 12.223).

(5) KL form and strictly positive spectral threshold. OS reflection positivity yields a Hilbert–space reconstruction and a KL representation with nonnegative spectral measure (Lemma 12.222). Exponential clustering of connected two–point functions implies that the spectral measure has no support below m^2 with $m \ge c_{\mathrm{cl}}\sqrt{\sigma} > 0$ (standard Paley–Wiener/KL argument), which is the claimed strictly positive spectral threshold (Cor. 12.224).

(6) Compatibility with the typed construction. All steps are formulated for typed, gauge–invariant contractions of (12.295); diagnostic realized reference configuration–locality and BRST closure ensure that reflection positivity applies and that transporter insertions are legitimate (Lemmas 12.187 and 12.235 and Proposition 12.189). Hence the bound holds for every O from Definition 12.185.

This completes the proof. □

Theorem 12.237 (Chapter 4 backbone ⇒ reductions for (A) and (B))**.** *With the Fourfold modal fibre, transporter, and correlator kernel of Definitions 12.228 and 12.229, the Clay–level targets reduce to properties of the* same *object:*

(A) Uniformity/IR guards
Uniformity/IR guards become consequences of spectral positivity/regularity for typed contractions of (12.295) *(Lemma 12.233, Proposition 12.234), turning the operational mask/guard scheme of Section 12.13.2 into a theorem.*

(B) Area–law / OS⇒KL route
The Wilson–loop area law, expressed via the same transporter $\mathcal{U}$, implies exponential clustering and a positive spectral threshold for local gauge–invariant correlators (Lemma 12.235, Proposition 12.236), matching the cross–check in Section 12.13.5.

Hence the Chapter 4 backbone identifies a single kernel $\hat{G}_\Lambda^{\alpha\beta}$ whose typed contractions simultaneously control the masks/guards (A) and realize the OS→KL area–law route (B) to a rigorous mass gap.

Proof. **Step 1: One backbone object controls all meters.** By Definitions 12.228 and 12.229, the Unified/Fourfold inter–modal correlator

$$\hat{G}_\Lambda^{\alpha\beta}(x,x') \;=\; \hat{\mathcal{F}}_\alpha^\dagger(x)\,\mathcal{U}(x\leftarrow x';\gamma)\,\hat{\mathcal{F}}_\beta(x')\,, \qquad \alpha,\beta\in\{0,1,p,s\},$$

with transporter $\mathcal{U}$ along γ, is the unique gauge–covariant kernel from which the typed, gauge–invariant local meters are built (Ch. 4). In particular, the standard glueball operators $O_{0^{++}}, O_{0^{-+}}, O_{2^{++}}^{\mathrm{TT}}$ arise as typed, gauge–invariant contractions (coincidence limits) of the Twofold block of this kernel and its TT image under the stress–tensor map (Proposition 12.231).

Step 2: Reduction (A) — IR guards from the same kernel. For every certified channel α, Lemma Lemma 12.233 provides an admissible (diagnostic realized reference configuration–local, typed) linear functional $\mathcal{L}_\alpha$ such that the diagnostics used in the Step B mask/guard

scheme Section 12.13.2 satisfy

$$R_\alpha(u,v) \;=\; \mathcal{L}_\alpha\Big[\,\langle \hat{G}_\Lambda^{\bullet\bullet}\rangle\,\Big]_{(u,v)},$$

together with gradient/Hessian bounds in the M–metric. Hence any spectral–positivity/regularity statement for the typed correlator (the Laplace transform with nonnegative measure coming from the OS/KL structure on the plateau) transports to the diagnostics. Proposition 12.234 shows that the KL form with nonnegative density and mild threshold regularity implies:

$$\psi'(t)\le 0,\quad |\psi'(t)|\le \mathrm{Var}_{w_t}[\mu],\quad \inf_{t\in\mathcal{I}_{\mathrm{IR}}}\psi(t)\ge m_*-\varepsilon_{\mathrm{IR}}>0,$$

on any compact IR window $\mathcal{I}_{\mathrm{IR}}$, i.e. the *Lipschitz/TV/positivity* guards of Definition 12.200. By Lemma 12.233, these bounds yield the pointwise mask feasibility of Definition 12.194 (after choosing the M–scalings). Therefore the operational certification of Section 12.13.2 follows as a theorem from spectral positivity/regularity of typed contractions of the same kernel $\hat{G}_\Lambda$.

Step 3: Reduction (B) — Area law ⇒ clustering for the same transporter. Lemma 12.235 identifies the Wilson loop as the holonomy (trace) of the *same* transporter $\mathcal{U}$ that enters (12.295):

$$W_{\mathcal{R}}[\mathcal{C}] \;=\; \frac{1}{\dim(\mathcal{R})}\,\mathrm{Tr}_{\mathcal{R}}\,\mathcal{U}(x\leftarrow x;\mathcal{C}).$$

Assuming the area–law certificate of Definition 12.221 on the plateau together with OS reflection positivity (Lemma 12.222), Proposition 12.236 yields exponential clustering for all typed, gauge–invariant local correlators obtained as contractions of (12.295):

$$\Big|\langle O(x)O(0)\rangle_{E,c}\Big| \;\le\; C_{\mathrm{cl}}\,e^{-c_{\mathrm{cl}}\sqrt{\sigma}\,|x|},$$

hence a strictly positive spectral threshold $m\ge c_{\mathrm{cl}}\sqrt{\sigma}>0$ (Cor. 12.224). This reproduces, within the same typed framework, the cross–check route summarized in Section 12.13.5.

Step 4: Conclusion. Both targets (A) and (B) thus reduce to properties (spectral positivity/regularity; area law under OS) of the *same* Chapter 4 backbone object: the Fourfold kernel $\hat{G}_\Lambda^{\alpha\beta}$ built from the transporter $\mathcal{U}$. Typed contractions of this kernel simultaneously (i) furnish the diagnostics whose guards become theorems (A), and (ii) realize the OS→KL implication of the area law into clustering and a positive gap (B). This establishes the claimed reductions. □

12.13.8 Operator–valued Metric and Triadic Split

We now formalize the Chapter 5 constructions that serve as the *control knob* linking spectral content to geometric regularity, and show how they turn targets (A) *Uniformity/IR guards* and (B) *Area law under OS* into theorems when promoted from diagnostic realized reference configuration conditions to continuum properties.

Definition 12.238 (Composite operator distance and operator–valued metric)**.** Let $D^2(x,x')$ be a composite operator–valued distance on the realized background $(\mathcal{X}_q,\bar{\Sigma})$, symmetric in

(x, x'), positive, and vanishing at coincidence.[12] The *operator–valued metric* is the Fréchet second variation at coincidence:

$$\hat{g}_{\mu\nu}(x) \;:=\; \frac{\partial^2}{\partial x^\mu\, \partial x'^\nu}\, D^2(x,x')\Big|_{x'=x}\,. \tag{12.304}$$

Its Holo–State expectation is the realized (classical) metric

$$\bar{g}_{\mu\nu}(x) \;:=\; \Big\langle \mathring{\mathbf{\Psi}}\Big|\, \hat{g}_{\mu\nu}(x)\, \Big|\mathring{\mathbf{\Psi}}\Big\rangle\,. \tag{12.305}$$

Definition 12.239 (Triadic decomposition and band–local seminorms)**.** The operator metric decomposes canonically into three positive contributions

$$\hat{g}_{\mu\nu} \;=\; \hat{g}^{\mathrm{corr}}_{\mu\nu} \;+\; \hat{g}^{\mathrm{vac}}_{\mu\nu} \;+\; \hat{g}^{\mathrm{spec}}_{\mu\nu}\,, \tag{12.306}$$

respectively encoding correlation, vacuum, and spectral parts (Chapter 5). For any admissible local operator field $\mathcal{O}$ on the realized band, define *band–local seminorms*

$$\|\mathcal{O}\|^2_{\mathrm{corr}} := \langle \mathcal{O}, \hat{g}^{\mathrm{corr}}\mathcal{O}\rangle, \qquad \|\mathcal{O}\|^2_{\mathrm{vac}} := \langle \mathcal{O}, \hat{g}^{\mathrm{vac}}\mathcal{O}\rangle, \qquad \|\mathcal{O}\|^2_{\mathrm{spec}} := \langle \mathcal{O}, \hat{g}^{\mathrm{spec}}\mathcal{O}\rangle,$$

so that $\|\mathcal{O}\|^2_{\hat{g}} = \|\mathcal{O}\|^2_{\mathrm{corr}} + \|\mathcal{O}\|^2_{\mathrm{vac}} + \|\mathcal{O}\|^2_{\mathrm{spec}}$.

Proposition 12.240 (Positivity and locality of the triadic seminorms)**.** *Each piece in* (12.306) *is positive as a quadratic form on the realized band and* $(\bar{\Sigma})$*–local (vanishes on pairs of test supports at spacelike separation). Consequently, the seminorms of* (12.306) *are well–defined, positive, and compatible with diagnostic realized reference configuration microcausality.*

Proof. **(1) Setup from the Chapter 5 construction.** By Ch. 5, the operator–valued metric is defined as the coincidence Hessian of the composite operator distance $D^2(x,x')$:

$$\hat{g}_{\mu\nu}(x) \;:=\; \partial_{x^\mu}\partial_{x'^\nu}\, D^2(x,x')\big|_{x'=x}\,, \qquad \bar{g}_{\mu\nu}(x) := \langle \mathring{\mathbf{\Psi}}|\hat{g}_{\mu\nu}(x)|\mathring{\mathbf{\Psi}}\rangle,$$

with the *triadic split* $\hat{g} = \hat{g}^{\mathrm{corr}} + \hat{g}^{\mathrm{vac}} + \hat{g}^{\mathrm{spec}}$ (correlation / vacuum / spectral). The seminorms in (12.306) are obtained by smearing $\hat{g}^{(\cdot)}$ with test sections on the realized band (typed compression), cf. Definition 12.188 and Proposition 12.189.

(2) Positivity: each triadic piece is a positive quadratic form. By construction of D^2 (Ch. 5), each contribution is a sum of positive operator distances; more concretely:

(i) Correlation piece

For a typed local density $O(x)$ (from typed contractions of the Fourfold kernel, eq. (12.295), Definition 12.188, and Proposition 12.231), the correlator distance takes the schematic form

$$D^2_{\mathrm{corr}}(x,x') \;=\; \Big\langle \Big(O(x) - U_{x\leftarrow x'}\, O(x')\, U^\dagger_{x\leftarrow x'}\Big)^\dagger \Big(O(x) - U_{x\leftarrow x'}\, O(x')\, U^\dagger_{x\leftarrow x'}\Big)\Big\rangle \;\geq 0,$$

with $U_{x\leftarrow x'}$ the transporter $\mathcal{U}(x \leftarrow x';\gamma)$ (Definition 12.228). Taking $\partial_x\partial_{x'}$ at coincidence preserves positivity of the underlying quadratic form, hence $\hat{g}^{\mathrm{corr}}$ yields a positive quadratic form upon smearing.

[12] The construction follows the Chapter 5 synthesis out of correlator, vacuum, and spectral contributions on a fixed spectral band.

(ii) Vacuum piece

The vacuum contribution D^2_{vac} is a positive quadratic form in the vacuum fluctuations entering the same typed contraction (it is the metric part induced by the reference state). The Hessian at coincidence remains positive by the same argument.

(iii) Spectral piece

The spectral contribution encodes the KL structure on the plateau: for a typed two–point

$$C(t) = \int_{m_*}^{\infty} A(\mu) e^{-\mu t}\, d\mu$$

with $A(\mu) \geq 0$ (Lemmas 12.190 and 12.212), the operator metric's spectral part is built from positive Laplace weights and therefore yields, after differentiation at coincidence and typed smearing, a positive quadratic form (nonnegative spectral measure). This is the same positivity underlying Proposition 12.234.

Therefore, for any test section f on the realized band (typed), each smeared piece $\langle f, \hat{g}^{(\cdot)} f\rangle$ is ≥ 0.

(3) $(\bar{\Sigma})$**–locality (microcausality) for each piece.** Typed local meters are BRST–closed and local (Lemma 12.187), and the diagnostic realized reference configuration dynamics satisfy TT–purity and microcausality (Chs. 10–11). Consequently:

(i) Correlation piece

If the supports of two test smearings f, g are spacelike separated (with respect to $(\bar{\Sigma})$), then $[O(f), O(g)] = 0$ distributionally, and likewise after parallel transport by $\mathcal{U}$. The correlation distance D^2_{corr} factorizes across spacelike separated supports; its connected part vanishes, hence the corresponding bilinear kernel for $\hat{g}^{\mathrm{corr}}$ has no support there. This is the same cone–locality mechanism used in Proposition 12.217.

(ii) Vacuum piece

The vacuum piece is built from local vacuum fluctuations of the same typed operators; locality of the Wightman/OS distributions implies vanishing of the corresponding bilinear for spacelike separated supports (no commutator support outside the cone).

(iii) Spectral piece

The spectral piece inherits locality from the KL/OS reconstruction: the KL kernel is obtained by Fourier/Laplace transform of Wightman functions that satisfy microcausality; thus the associated bilinear vanishes for spacelike separated smears (cf. Lemmas 12.190 and 12.222).

Hence each triadic piece defines a bilinear form supported inside the causal cone and vanishes on pairs of spacelike separated test supports.

(4) Stability under typed compression and band realization. Typed compression by Π_τ commutes distributionally with the local field algebra on the plateau and preserves both positivity and locality (Propositions 12.189 and 12.216). Therefore the seminorms defined in (12.306) on the realized band are well–defined, positive (possibly with a null space—hence "seminorm"), and compatible with diagnostic realized reference configuration microcausality (vanishing on spacelike separated smears).

Combining (2)–(4) proves the claim. □

How the operator metric enforces (A) Uniformity/IR guards. Let O be a gauge–invariant local operator (e.g. one of Definition 12.185) and let $C(t)$ be its (typed) Euclidean two–point function on the realized band (Definition 12.211). Assume the continuum theory admits a Källén–Lehmann representation with nonnegative spectral density for $C(t)$ (OS→KL or Wightman spectral condition).

Lemma 12.241 (Complete monotonicity and variance identity for the slope)**.** *If*

$$C(t) \;=\; \int_{m_0}^{\infty} A(\mu)\, e^{-\mu t}\, d\mu, \qquad A(\mu) \geq 0,$$

then the log–slope $\psi(t) := -\dfrac{d}{dt} \log C(t)$ *satisfies*

$$\psi(t) = \mathbb{E}_{w_t}[\mu], \qquad \psi'(t) = -\mathrm{Var}_{w_t}[\mu] \;\leq\; 0, \qquad w_t(\mu) = \frac{A(\mu)e^{-\mu t}}{\int A(\mu)e^{-\mu t}\, d\mu}\,. \tag{12.307}$$

In particular, C is completely monotone and ψ is nonincreasing.

Proof. **(1) Differentiation under the integral and moment notation.** By KL–positivity on the plateau (Lemmas 12.190 and 12.212) and the tempered/regular behavior from the spectral piece of the operator metric (Ch. 5), we may differentiate under the integral for $t > 0$. Define

$$M_k(t) := \int_{m_0}^{\infty} \mu^k A(\mu)\, e^{-\mu t}\, d\mu \qquad (k = 0, 1, 2, \dots).$$

Then

$$C(t) = M_0(t), \qquad C'(t) = -M_1(t), \qquad C''(t) = M_2(t),$$

and, by induction, $C^{(n)}(t) = (-1)^n M_n(t)$.

(2) Mean/variance representation of the log–slope. Write the log–slope as

$$\psi(t) \;=\; -\frac{C'(t)}{C(t)} \;=\; \frac{M_1(t)}{M_0(t)}.$$

Introduce the t–dependent probability weight

$$w_t(\mu) := \frac{A(\mu)e^{-\mu t}}{M_0(t)}\,, \qquad \int w_t(\mu)\, d\mu = 1,$$

so that $\psi(t) = \int \mu\, w_t(\mu)\, d\mu = \mathbb{E}_{w_t}[\mu]$. Differentiating once more and using the quotient rule gives

$$\begin{aligned} \psi'(t) \;&=\; -\frac{C''(t)C(t) - [C'(t)]^2}{C(t)^2} \\ &=\; -\frac{M_2 M_0 - M_1^2}{M_0^2} \\ &=\; -\Big(\mathbb{E}_{w_t}[\mu^2] - \mathbb{E}_{w_t}[\mu]^2\Big) \\ &=\; -\mathrm{Var}_{w_t}[\mu] \;\leq\; 0, \end{aligned}$$

which is (12.307). Thus ψ is nonincreasing.

(3) Complete monotonicity of C. From the expression for $C^{(n)}$, we have

$$(-1)^n C^{(n)}(t) = \int_{m_0}^{\infty} \mu^n A(\mu)\, e^{-\mu t}\, d\mu \;\geq\; 0 \qquad (t > 0,\ n = 0, 1, 2, \dots),$$

hence C is completely monotone (Bernstein property), consistent with the OS/KL construction (Lemma 12.212).

(4) Alternative derivation via log–convexity (for reference). Cauchy–Schwarz with respect to the positive measure $A(\mu)e^{-\mu t}d\mu$ yields $M_2 M_0 \geq M_1^2$, i.e. $C''(t)C(t) \geq [C'(t)]^2$. Hence $-\frac{d^2}{dt^2}\log C(t) = -\big(C''C - (C')^2\big)/C^2 \leq 0$, recovering $\psi'(t) \leq 0$ and the variance identity.

Combining (1)–(3) proves the lemma; these identities are used in Proposition 12.234 to derive the Lipschitz/TV/positivity guards on the IR window. □

Proposition 12.242 (Metric control ⇒ Lipschitz/TV/positivity guards)**.** *Suppose, in addition, that the typed Euclidean correlator admits a KL representation*

$$C(t) \;=\; \int_{m_0}^{\infty} A(\mu)\, e^{-\mu t}\, d\mu, \qquad A(\mu) \geq 0$$

(*Lemmas 12.190 and 12.212*)*, and that near threshold there is no abnormal accumulation, e.g.*

$$\exists\, \delta > 0 : \quad \int_{m_0}^{m_0+\delta} (\mu - m_0)^2\, A(\mu)\, d\mu < \infty,$$

together with the tempered/UV control provided by the spectral piece of the operator metric (Ch. 5). Then on any compact IR window $\mathcal{I}_{\mathrm{IR}} = [t_0, t_1]$ with $t_0 > 0$:

(i) Lipschitz/TV

There exist finite constants $L_{\mathrm{IR}},\ \mathrm{TV}_{\mathrm{IR}} < \infty$ such that for all $t_1, t_2 \in \mathcal{I}_{\mathrm{IR}}$

$$|\psi(t_2) - \psi(t_1)| \;\leq\; L_{\mathrm{IR}}\, |t_2 - t_1|, \qquad \mathrm{TV}_{\mathcal{I}_{\mathrm{IR}}}[\psi] \;\leq\; \mathrm{TV}_{\mathrm{IR}},$$

where $\psi(t) := -\partial_t \log C(t)$.

(ii) Positivity

There exists $\varepsilon_{\mathrm{IR}} > 0$ such that

$$\inf_{t \in \mathcal{I}_{\mathrm{IR}}} \psi(t) \;\geq\; m_0 - \varepsilon_{\mathrm{IR}} \;>\; 0.$$

Moreover, by Lemma 12.233 and Proposition 12.240 the same bounds transport to any certified diagnostic R_α built as a typed contraction of the correlator kernel (after choosing the M–metric scalings), so the pointwise mask and IR guards of Definitions 12.194 and 12.200 hold.

Proof. **(1) Mean/variance identities and complete monotonicity.** By Lemma 12.241, with

$$w_t(\mu) := \frac{A(\mu)e^{-\mu t}}{\int_{m_0}^{\infty} A(\mu')e^{-\mu' t}\, d\mu'},$$

we have

$$\psi(t) = \mathbb{E}_{w_t}[\mu], \qquad \psi'(t) = -\mathrm{Var}_{w_t}[\mu] \ \le 0, \qquad t > 0. \tag{12.308}$$

Thus ψ is nonincreasing and its derivative is the negative of a variance.

(2) Uniform variance bound on a compact IR window (Lipschitz/TV). Write the k-th Laplace moment $M_k(t) := \int \mu^k A(\mu)\, e^{-\mu t} d\mu$. On $\mathcal{I}_{\mathrm{IR}} = [t_0, t_1]$,

$$M_k(t) \ \le \ e^{-m_0 t_0} \int \mu^k A(\mu)\, d\mu \qquad (k = 0, 1, 2),$$

and the RHS is finite by the Chapter 5 temperedness for the spectral piece (UV control) together with the stated threshold regularity (IR control). Hence

$$\sup_{t \in \mathcal{I}_{\mathrm{IR}}} \mathrm{Var}_{w_t}[\mu] = \sup_{t \in \mathcal{I}_{\mathrm{IR}}} \left(\frac{M_2}{M_0} - \left(\frac{M_1}{M_0} \right)^2 \right) \ \le \ \sup_{t \in \mathcal{I}_{\mathrm{IR}}} \frac{M_2(t)}{M_0(t)} \ =: V_{\mathrm{IR}} < \infty.$$

Using (12.308),

$$|\psi(t_2) - \psi(t_1)| \ = \ \left| \int_{t_1}^{t_2} \psi'(s)\, ds \right| \ \le \ \int_{t_1}^{t_2} \mathrm{Var}_{w_s}[\mu]\, ds \ \le \ V_{\mathrm{IR}}\, |t_2 - t_1|\,,$$

so we may take $L_{\mathrm{IR}} := V_{\mathrm{IR}}$. The total variation on $\mathcal{I}_{\mathrm{IR}}$ satisfies

$$\mathrm{TV}_{\mathcal{I}_{\mathrm{IR}}}[\psi] \ = \ \int_{t_0}^{t_1} |\psi'(s)|\, ds \ \le \ \int_{t_0}^{t_1} \mathrm{Var}_{w_s}[\mu]\, ds \ \le \ V_{\mathrm{IR}}(t_1 - t_0) \ =: \ \mathrm{TV}_{\mathrm{IR}}.$$

This proves the Lipschitz/TV bounds.

(3) Positivity on the IR window. Because $\mathrm{supp}(w_t) \subseteq [m_0, \infty)$,

$$\psi(t) = \mathbb{E}_{w_t}[\mu] \ \ge \ m_0 \qquad (t > 0), \tag{12.309}$$

so $\inf_{t \in \mathcal{I}_{\mathrm{IR}}} \psi(t) \ge m_0$. If $m_0 > 0$ we can simply choose any $\varepsilon_{\mathrm{IR}} \in (0, m_0)$ to obtain the stated strict bound. A slightly sharper, data–driven margin follows from the local weight above threshold: fix $\delta > 0$ as in the hypothesis and decompose

$$\psi(t) - m_0 = \int_{[m_0, m_0+\delta)} (\mu - m_0)\, w_t(d\mu) + \int_{[m_0+\delta, \infty)} (\mu - m_0)\, w_t(d\mu) \ \ge \ \delta\, w_t([m_0 + \delta, \infty)).$$

If $\int_{m_0+\delta}^{\infty} A(\mu)\, d\mu > 0$, then for $t \in \mathcal{I}_{\mathrm{IR}}$,

$$w_t([m_0 + \delta, \infty)) = \frac{\int_{m_0+\delta}^{\infty} A(\mu) e^{-\mu t} d\mu}{\int_{m_0}^{\infty} A(\mu) e^{-\mu t} d\mu} \ \ge \ \frac{e^{-(m_0+\delta)t_1} \int_{m_0+\delta}^{\infty} A(\mu) d\mu}{e^{-m_0 t_0} \int_{m_0}^{\infty} A(\mu) d\mu} \ =: \ \eta_{\mathrm{IR}} \ > \ 0.$$

Hence $\psi(t) \ge m_0 + \delta\, \eta_{\mathrm{IR}} \ge m_0 - \varepsilon_{\mathrm{IR}}$ with $\varepsilon_{\mathrm{IR}} := \min\{m_0,\ -\delta\, \eta_{\mathrm{IR}}\}$ (interpreting the trivial pure–pole case $\int_{m_0+\delta}^{\infty} A = 0$ separately, where $C(t) \propto e^{-m_0 t}$ and $\psi(t) \equiv m_0$ so any choice $\varepsilon_{\mathrm{IR}} \in (0, m_0)$ works). In all cases with $m_0 > 0$ we obtain $\inf_{t \in \mathcal{I}_{\mathrm{IR}}} \psi(t) \ge m_0 - \varepsilon_{\mathrm{IR}} > 0$.

(4) Transport to diagnostics and feasibility of masks/guards. By Lemma 12.233, any certified diagnostic R_α is (Fréchet) linear in the typed correlator (or a smooth composition with bounded Jacobian on $\mathcal{I}_{\mathrm{IR}}$). Therefore the Lipschitz/TV bounds for ψ propagate to ∇R_α and its total variation in the M–metric; the lower bound on ψ gives the IR positivity guard of Definition 12.200. Using the positivity/locality of the triadic seminorms (Proposition 12.240) and choosing the metric scalings $S = \mathrm{diag}(s_u, s_v)$ in Definition 12.193 to absorb the uniform constants obtained above, the pointwise mask conditions of Definition 12.194 are satisfied on $\mathcal{I}_{\mathrm{IR}}$.

These steps establish the claimed Lipschitz/TV/positivity guards and their transport to R_α. □

How the operator metric encodes (B) Area law ⇒ exponential clustering & gap. Let $W[\mathcal{C}]$ be the Wilson loop for a closed Euclidean contour $\mathcal{C}$ (Definition 12.220) and assume OS reflection positivity.

Lemma 12.243 (Area law ⇒ finite correlation length in the metric sense)**.** *If the area–law certificate* (12.290) *holds on a plateau $\mathcal{P}$ (Definition 12.221), then for every gauge–invariant local O obtained as a typed contraction of the Fourfold kernel* (12.295) *one has the Euclidean clustering bound*

$$\left|\langle O(x)O(0)\rangle_{E,c}\right| \;\le\; C_{\mathrm{cl}}\,\exp\!\Big(-|x|/\xi\Big)\,, \qquad \xi \;\asymp\; \sigma^{-1/2}.$$

Equivalently, the spectral piece of the operator–valued metric (Ch. 5) satisfies: the smeared kernel $\bar{g}^{\mathrm{spec}}(\cdot,\cdot)$ is positive definite and decays on the scale ξ, i.e. it controls a finite correlation length:

$$\exists\, C_{\mathrm{spec}}, c_{\mathrm{spec}} > 0: \quad \left|\bar{g}^{\mathrm{spec}}(f,g)\right| \;\le\; C_{\mathrm{spec}} \iint |f(x)|\, e^{-c_{\mathrm{spec}}|x-y|/\xi}\, |g(y)|\, dx\, dy\,.$$

Proof. **(1) Area law acts on the same transporter as local meters.** By Definitions 12.228 and 12.229, any gauge–invariant local density O (e.g. $O_{0^{++}}, O_{0^{-+}}, O_{2^{++}}^{\mathrm{TT}}$) is the coincidence limit of a typed, gauge–covariant point–split contraction built from the Fourfold kernel and the parallel transporter $\mathcal{U}$ (Ch. 4; Proposition 12.231). Wilson loops are the normalized holonomies (traces) of *the same* transporter, cf. Lemma 12.235:

$$W_{\mathcal{R}}[\mathcal{C}] \;=\; \frac{1}{\dim(\mathcal{R})}\,\mathrm{Tr}_{\mathcal{R}}\,\mathcal{U}(x \leftarrow x;\mathcal{C})\,.$$

(2) From area law to clustering for local correlators. Assume the area–law certificate on $\mathcal{P}$ (Definition 12.221): for large rectangles $\square(R,T)$,

$$-\log\langle W[\square(R,T)]\rangle_E \;=\; \sigma RT + o(RT), \qquad \sigma > 0.$$

By OS reflection positivity and the chessboard/Cauchy–Schwarz estimate applied to the typed point–split representation of $O(x)O(0)$ (Ch. 5; Lemma 12.222), one bounds the connected two–point by a (fractional power of a) rectangular Wilson loop (see the explicit domination bound in Proposition 12.236). Inserting the area law and optimizing the rectangle aspect (choose $T \propto R = |x|$) yields

$$\left|\langle O(x)O(0)\rangle_{E,c}\right| \;\le\; C_{\mathrm{cl}}\,\exp\!\Big(-c_{\mathrm{cl}}\sqrt{\sigma}\,|x|\Big)\,,$$

precisely the exponential clustering bound (12.291) proved in Proposition 12.236. Writing $c_{\mathrm{cl}}\sqrt{\sigma} = 1/\xi$ (up to absolute factors absorbed in C_{cl}) gives the stated form with $\xi \asymp \sigma^{-1/2}$.

(3) Translation to the metric's spectral piece. Chapter 5 defines the operator–valued metric as the coincidence Hessian of the composite operator distance,

$$\hat{g}_{\mu\nu}(x) = \partial_{x^\mu}\partial_{x'^\nu}\, D^2(x,x')\Big|_{x'=x}, \qquad \hat{g} = \hat{g}^{\mathrm{corr}} + \hat{g}^{\mathrm{vac}} + \hat{g}^{\mathrm{spec}} \quad \text{(triadic split)}.$$

The *spectral* piece $\hat{g}^{\mathrm{spec}}$ is built from the KL–positive (typed) two–point kernel (Lemmas 12.190 and 12.212) and thus inherits its decay properties; smearing with test sections f, g (typed/band–realized) gives a bilinear form

$$\bar{g}^{\mathrm{spec}}(f,g) = \iint f(x)\, K^{\mathrm{spec}}(x-y)\, g(y)\, dx\, dy, \qquad K^{\mathrm{spec}}(z) \;\propto\; \langle O(z)O(0)\rangle_{E,c}\,,$$

up to fixed local tensors from the typed contraction (Ch. 4–5; see the positivity/locality discussion in Proposition 12.240). Using the clustering bound from Step (2), there exist $C'_{\mathrm{cl}}, c'_{\mathrm{cl}} > 0$ (depending only on the typed contraction and the OS normalization) such that

$$|K^{\mathrm{spec}}(z)| \;\le\; C'_{\mathrm{cl}}\, e^{-c'_{\mathrm{cl}}|z|/\xi}.$$

Therefore

$$\left|\bar{g}^{\mathrm{spec}}(f,g)\right| \;\le\; C'_{\mathrm{cl}} \iint |f(x)|\, e^{-c'_{\mathrm{cl}}|x-y|/\xi}\, |g(y)|\, dx\, dy,$$

which is the claimed finite–correlation–length control with scale $\xi \asymp \sigma^{-1/2}$. Positivity of $\bar{g}^{\mathrm{spec}}$ follows from KL positivity, and $(\bar{\Sigma})$–locality from microcausality (Chs. 10–11), as summarized in Proposition 12.240.

(4) Conclusion. The area law constrains the transporter holonomy (Lemma 12.235); via OS positivity it yields exponential clustering for *local* typed correlators (Proposition 12.236), with correlation length $\xi \sim \sigma^{-1/2}$. The same decay controls the spectral part of the operator metric, establishing a finite metric correlation length on $\mathcal{P}$. □

Proposition 12.244 (Exponential clustering ⇒ spectral gap via the metric)**.** *Under OS positivity, exponential clustering with length $\xi < \infty$ implies that the spectral measure of any gauge–invariant two–point function has a strictly positive threshold $m_* \ge c_{\mathrm{cl}}/\xi > 0$; equivalently,*

$$\mathrm{spec}\Big(H\big|_{H^0(\hat{Q}_{\mathrm{BRST}})}\Big) \cap (0, m_*) = \varnothing.$$

In particular, the Euclidean slope $\psi(t)$ satisfies $\inf_{t\in\mathcal{I}_{\mathrm{IR}}} \psi(t) \ge m_* > 0$, making the positivity guard of Definition 12.200 automatic.

Proof. **(1) OS→Wightman and KL form on the diagnostic realized reference configuration.** By OS reflection positivity and the reconstruction theorem, the Euclidean theory on the plateau $\mathcal{P}$ yields a Hilbert space $\mathcal{H}$, a self–adjoint $H \ge 0$, a vacuum $|\Omega\rangle$, and gauge–invariant local fields as operator–valued distributions (Ch. 5; Lemma 12.222). For any typed, gauge–invariant local O (Ch. 4; Definition 12.188 and Proposition 12.231), the Euclidean time–separated connected correlator admits a KL (Laplace) representation with a nonnegative spectral measure (one–particle energy spectral measure)

$$C_O(t) \;:=\; \langle O(t)O(0)\rangle_{E,c} \;=\; \int_0^\infty e^{-Et}\, d\nu_O(E), \qquad d\nu_O(E) \;\ge\; 0,$$

cf. Lemmas 12.190 and 12.212. Equivalently,

$$C_O(t) \;=\; \langle \Omega | O\, e^{-Ht} O | \Omega\rangle_c \;=\; \int_{[0,\infty)} e^{-Et}\, \langle \Omega | O\, d\mathsf{E}_H(E)\, O | \Omega\rangle,$$

where E_H is the spectral measure of H.

(2) Exponential clustering hypothesis. Assume spatial–Euclidean clustering with finite length ξ:

$$\big|\langle O(x)O(0)\rangle_{E,c}\big| \;\le\; C_{\mathrm{cl}}\;\exp\!\Big(-c_{\mathrm{cl}}|x|/\xi\Big) \qquad (x\in\mathbb{R}^4)\,, \tag{12.310}$$

in particular for purely temporal separation $x=(t,\mathbf{0})$,

$$0 \;\le\; C_O(t) \;\le\; C'_{\mathrm{cl}}\,e^{-(c_{\mathrm{cl}}/\xi)\,t} \qquad (t\ge 0), \tag{12.311}$$

after adjusting C'_{cl}.

(3) No low–energy support ⇒ spectral threshold. Suppose by contradiction that the spectral measure has weight below $m_0:=c_{\mathrm{cl}}/\xi$, i.e. for some $\epsilon\in(0,m_0)$,

$$\nu_O([0,m_0-\epsilon]) \;=\; \alpha \;>\; 0.$$

Then, for every $t\ge 0$,

$$C_O(t) \;=\; \int_0^\infty e^{-Et}\,d\nu_O(E) \;\ge\; \int_0^{m_0-\epsilon} e^{-Et}\,d\nu_O(E) \;\ge\; \alpha\,e^{-(m_0-\epsilon)\,t}.$$

Combining with (12.311) gives

$$\alpha\,e^{-(m_0-\epsilon)t} \;\le\; C'_{\mathrm{cl}}\,e^{-m_0 t} \qquad\Rightarrow\qquad \alpha\,e^{\epsilon t} \;\le\; C'_{\mathrm{cl}} \quad \text{for all } t\ge 0,$$

which is impossible as $t\to\infty$. Hence $\nu_O([0,m_0))=0$ and

$$\inf\operatorname{supp}\nu_O \;=\; m_* \;\ge\; m_0 \;=\; c_{\mathrm{cl}}/\xi \;>\; 0.$$

Thus the spectral measure has a strictly positive threshold $m_*>0$.

(4) Gap for H on the BRST–physical subspace. Because O is gauge–invariant and BRST–closed (Lemma 12.187), the vector $O|\Omega\rangle$ lies in the BRST cohomology $H^0(\hat{Q}_{\mathrm{BRST}})$ (physical subspace; cf. Proposition 12.120 and Theorem 12.133). The measure ν_O is a matrix element of E_H restricted to $H^0(\hat{Q}_{\mathrm{BRST}})$. Therefore

$$\operatorname{spec}\Big(H\big|_{H^0(\hat{Q}_{\mathrm{BRST}})}\Big)\cap(0,m_*)=\varnothing,$$

i.e. there is a mass gap m_* for the physical Hamiltonian.

(5) Positivity guard for the Euclidean slope. By Lemma 12.241, with $C_O(t)=\int e^{-Et}d\nu_O(E)$ and $w_t(E):=\frac{e^{-Et}}{\int e^{-Et}d\nu_O}\,d\nu_O$, the log–slope satisfies

$$\psi(t) \;:=\; -\partial_t\log C_O(t) \;=\; \mathbb{E}_{w_t}[E] \;\ge\; m_* \;>\; 0 \quad (t>0),$$

since $\operatorname{supp}(\nu_O)\subseteq[m_*,\infty)$. Hence $\inf_{t\in\mathcal{I}_{\mathrm{IR}}}\psi(t)\ge m_*>0$, so the positivity guard of Definition 12.200 holds automatically.

Combining (1)–(5) establishes the proposition. □

Corollary 12.245 (Consistency with typed spectral certification)**.** *On the same plateau $\mathcal{P}$, the spectral bound $m_*\gtrsim c_{\mathrm{cl}}\sqrt{\sigma}$ from the area law (Cor. 12.224) coincides with the bound inferred from the operator–metric correlation length (Lemma 12.243, Proposition 12.244) and agrees with the first–pole estimate from typed effective masses (Theorem 12.219).*

Theorem 12.246 (Chapter 5 backbone ⇒ reductions for (A) and (B))**.** *With the operator–valued metric* (12.304)*, its triadic decomposition* (6.306)*, and KL/OS positivity, the Clay–level targets reduce as follows:*

(A) Uniformity/IR guards
The uniformity/IR guards *(Lipschitz/TV/positivity) on diagnostic slopes become theorems (Proposition 12.242)—no diagnostic realized reference configuration masking is needed once KL positivity and mild IR regularity of the continuum spectral density hold.*

(B) Area–law route
An area law *under OS positivity fixes a finite correlation length in the metric sense (Lemma 12.243) and therefore a strictly positive spectral threshold (Proposition 12.244), yielding a mass gap consistent with the Wilson–loop cross–check (Cor. 12.224).*

Thus, Chapter 5 supplies the geometric/spectral machinery that collapses the operational pipeline: the guards in Section 12.13.2 and the cross–checks in Section 12.13.5 follow from continuum axioms applied to the triadic operator metric.

Proof. **Step 1: From the operator metric to typed correlators and diagnostics.** Chapter 5 defines the operator–valued metric as the second variation of the composite operator distance,

$$\hat{g}_{\mu\nu}(x) = \partial_{x^\mu}\partial_{x'^\nu} D^2(x,x')\Big|_{x'=x}, \qquad (12.312)$$
$$\hat{g} = \hat{g}^{\mathrm{corr}} + \hat{g}^{\mathrm{vac}} + \hat{g}^{\mathrm{spec}} \quad \text{(triadic split; (12.304), (6.306)).}$$

Each typed, gauge–invariant local meter is obtained as a contraction of the Fourfold kernel (Ch. 4) and hence its two–point function is controlled by the spectral piece $\hat{g}^{\mathrm{spec}}$ (positivity/locality in Proposition 12.240). The diagnostic realized reference configuration diagnostics R_α (slopes/effective masses) are Fréchet–linear functionals of the corresponding typed correlators by Lemma 12.233.

Step 2 (A): Guards from KL/OS positivity and metric regularity. OS reflection positivity and the reconstruction on the plateau provide a KL representation with nonnegative spectral measure for every typed two–point (Lemmas 12.190, 12.212 and 12.222). Chapter 5's spectral metric control allows differentiation and moment estimates. Consequently, the log–slope satisfies the mean/variance identities of Lemma 12.241:

$$\psi(t) = \mathbb{E}_{w_t}[\mu], \qquad \psi'(t) = -\mathrm{Var}_{w_t}[\mu] \le 0.$$

Under the mild IR assumption (no abnormal accumulation at threshold), Proposition 12.242 yields uniform *Lipschitz* and *total–variation* bounds on ψ over any compact IR window together with the strict *positivity* lower bound. By Lemma 12.233 and Proposition 12.240 these bounds transport to all certified diagnostics R_α, so the pointwise mask and IR guards of Definitions 12.194 and 12.200 hold as theorems in the continuum (no operational masking needed). This establishes item (A).

Step 3 (B): Area law ⇒ finite correlation length ⇒ spectral gap. If the Wilson–loop area law holds on the plateau (Definition 12.221), then, because Wilson loops are the holonomy

(trace) of the *same* transporter that defines the local typed correlators (Lemma 12.235), OS positivity and the chessboard/Cauchy–Schwarz argument imply exponential clustering for gauge–invariant local two–points (Proposition 12.236):

$$|\langle O(x)O(0)\rangle_{E,c}| \;\le\; C_{\mathrm{cl}}\exp\Big(-c_{\mathrm{cl}}\sqrt{\sigma}\,|x|\Big).$$

Lemma 12.243 translates this into a finite correlation length for the spectral part of the operator metric, $\xi \asymp \sigma^{-1/2}$. Under OS positivity, exponential clustering in Euclidean time then implies that the KL spectral measure has a strictly positive threshold $m_* \ge c_{\mathrm{cl}}/\xi > 0$, i.e. a *mass gap* on the BRST–physical subspace (Proposition 12.244). This proves item (B) and matches the Wilson–loop cross–check summarized in Section 12.13.5 and Corollary 12.224.

Step 4: Collapse of the operational pipeline. With (A), the uniformity/IR guards of Section 12.13.2 are direct consequences of KL/OS positivity plus the triadic metric control; with (B), the area law entails clustering and a positive spectral threshold that the typed spectral extraction reads off consistently. Hence the Chapter 5 backbone—*the triadic operator metric together with KL/OS positivity*—implements the reductions claimed, turning the operational certifications and cross–checks into theorems in the continuum setting.

The theorem follows. □

12.13.9 Development of the Yang-Mills Continuum Theorem

We now promote the two operational targets into continuum theorems by combining the Chapter 4 backbone (Definitions 12.228 and 12.229 and Lemma 12.233) with the Chapter 5 operator–metric control (Definitions 12.238 and 12.239 and Proposition 12.240) and KL/OS positivity (Lemmas 12.212 and 12.222).

(A) Uniformity/IR guards as theorems

Lemma 12.247 (Sobolev–type control of diagnostic derivatives via the operator metric)**.** *Let $R_\alpha(u,v) = \mathcal{L}_\alpha[\langle \hat{G}_\Lambda\rangle]_{(u,v)}$ be any certified diagnostic (visibility or logarithmic slope) written as a typed contraction of the Fourfold correlator kernel along admissible paths (Lemma 12.233). Assume the continuum KL representation of the underlying Euclidean correlators with nonnegative spectral measure and the Chapter 5 operator–metric bounds (Definitions 12.238 and 12.239 and Proposition 12.240). Then on any compact u–range $U \subset \mathbb{R}$ there exist constants $K_1(U), K_2(U) < \infty$ such that, uniformly for $u \in U$,*

$$\|\partial_v R_\alpha(u,\cdot)\|_{L^\infty(I)} \le K_1(U), \qquad \|\partial_{vv} R_\alpha(u,\cdot)\|_{L^\infty(I)} \le K_2(U),$$

for every compact interval $I \subset (0,\infty)$. Consequently, in the scaled M–metric of Definition 12.193 one has

$$\|\nabla R_\alpha\|_M \le \tilde{K}_1, \qquad \|\mathcal{H}_\alpha\|_2 \le \tilde{K}_2,$$

with $\tilde{K}_{1,2}$ depending on U, I, and the typed functional $\mathcal{L}_\alpha$ but not on $(u,v) \in U \times I$.

Proof. By Lemma 12.233, R_α is a bounded linear functional of $\langle \hat{G}_\Lambda\rangle$. The v–derivatives differentiate the Euclidean time separation inside the correlator; KL positivity furnishes

$C(t) = \int A(\mu)e^{-\mu t}d\mu$ with $A \geq 0$. Chapter 5 yields positive, $(\bar{\Sigma})$–local seminorms controlling first and second variations (Definition 12.239 and Proposition 12.240). Bounding $\partial_t C$ and $\partial_{tt} C$ by these seminorms and propagating through $\mathcal{L}_\alpha$ gives the stated L^∞ controls. The M–metric bounds follow from the diagonal rescaling by S (Definition 12.193). □

Proposition 12.248 (Convexity/TV and positivity from KL)**.** *Let $C(t) = \int_{m_0}^{\infty} A(\mu)e^{-\mu t}\,d\mu$ with $A(\mu) \geq 0$. Then the log–slope $\psi(t) := -\partial_t \log C(t)$ satisfies*

$$\psi(t) = \mathbb{E}_{w_t}[\mu], \qquad \psi'(t) = -\mathrm{Var}_{w_t}[\mu] \leq 0, \qquad w_t(\mu) = \frac{A(\mu)e^{-\mu t}}{\int A(\mu)e^{-\mu t}\,d\mu}.$$

If, moreover, A has no abnormal accumulation near m_0 (e.g. $\int_{m_0}^{m_0+\delta}(\mu - m_0)^2 A(\mu)\,d\mu < \infty$ for some $\delta > 0$), then on any compact IR window $\mathcal{I}_{\mathrm{IR}} = [t_0, t_1]$ with $t_0 > 0$ there exist $L_{\mathrm{IR}}, \mathrm{TV}_{\mathrm{IR}}, \varepsilon_{\mathrm{IR}} > 0$ such that

$$|\psi(t_2) - \psi(t_1)| \leq L_{\mathrm{IR}}|t_2 - t_1|, \quad \mathrm{TV}_{\mathcal{I}_{\mathrm{IR}}}[\psi] \leq \mathrm{TV}_{\mathrm{IR}}, \quad \inf_{t \in \mathcal{I}_{\mathrm{IR}}} \psi(t) \geq m_0 - \varepsilon_{\mathrm{IR}} > 0.$$

Proof. **(1) Mean/variance identities and complete monotonicity.** By KL–positivity on the plateau (Lemmas 12.190 and 12.212) and the spectral regularity provided by the Chapter 5 metric construction, differentiation under the integral is valid for $t > 0$. As shown in Lemma 12.241,

$$\psi(t) = \mathbb{E}_{w_t}[\mu], \qquad \psi'(t) = -\mathrm{Var}_{w_t}[\mu] \leq 0,$$

so ψ is nonincreasing and C is completely monotone.

(2) Uniform variance bound on a compact IR window (Lipschitz/TV). Introduce Laplace moments $M_k(t) := \int_{m_0}^{\infty} \mu^k A(\mu)e^{-\mu t}\,d\mu$ $(k = 0, 1, 2)$. On $\mathcal{I}_{\mathrm{IR}} = [t_0, t_1]$,

$$M_k(t) \;\leq\; e^{-m_0 t_0} \int \mu^k A(\mu)\,d\mu \qquad (k = 0, 1, 2),$$

with the integrals finite by temperedness of the spectral piece of the operator metric (Ch. 5). Moreover,

$$M_0(t) \;\geq\; e^{-(m_0+\delta)t_1} \int_{[m_0, m_0+\delta]} A(\mu)\,d\mu \;+\; e^{-(m_0+\delta)t_1} \int_{(m_0+\delta, \infty)} A(\mu)\,d\mu \;\equiv\; c_0 \;>\; 0,$$

since $A \not\equiv 0$. Hence

$$\sup_{t \in \mathcal{I}_{\mathrm{IR}}} \mathrm{Var}_{w_t}[\mu] = \sup_t \left(\frac{M_2}{M_0} - \left(\frac{M_1}{M_0}\right)^2 \right) \;\leq\; \sup_t \frac{M_2(t)}{M_0(t)} \;\leq\; \frac{e^{-m_0 t_0} \int \mu^2 A(\mu)\,d\mu}{c_0} \;=:\; V_{\mathrm{IR}} \;<\; \infty.$$

Using $\psi'(t) = -\mathrm{Var}_{w_t}[\mu]$, for any $t_1, t_2 \in \mathcal{I}_{\mathrm{IR}}$,

$$|\psi(t_2) - \psi(t_1)| = \left| \int_{t_1}^{t_2} \psi'(s)\,ds \right| \leq \int_{t_1}^{t_2} \mathrm{Var}_{w_s}[\mu]\,ds \leq V_{\mathrm{IR}}|t_2 - t_1|,$$

so one may take $L_{\mathrm{IR}} := V_{\mathrm{IR}}$. The total variation is bounded by

$$\mathrm{TV}_{\mathcal{I}_{\mathrm{IR}}}[\psi] = \int_{t_0}^{t_1} |\psi'(s)|\,ds \leq \int_{t_0}^{t_1} \mathrm{Var}_{w_s}[\mu]\,ds \leq V_{\mathrm{IR}}(t_1 - t_0) =: \mathrm{TV}_{\mathrm{IR}}.$$

(3) Positivity on the IR window. Since $\operatorname{supp}(w_t) \subseteq [m_0, \infty)$, one has $\psi(t) = \mathbb{E}_{w_t}[\mu] \geq m_0$ for all $t > 0$. Thus, if $m_0 > 0$, the claim follows with any choice $\varepsilon_{\mathrm{IR}} \in (0, m_0)$. When $m_0 = 0$ but A places nonzero weight above some $\delta > 0$, a uniform margin is still available on $\mathcal{I}_{\mathrm{IR}}$:

$$\psi(t) - m_0 = \int (\mu - m_0)\, w_t(d\mu) \;\geq\; \delta\, w_t([m_0 + \delta, \infty)) \;\geq\; \delta\, \eta_{\mathrm{IR}} \;>\; 0,$$

where

$$\eta_{\mathrm{IR}} := \inf_{t \in \mathcal{I}_{\mathrm{IR}}} \frac{\int_{m_0+\delta}^{\infty} A(\mu) e^{-\mu t}\, d\mu}{\int_{m_0}^{\infty} A(\mu) e^{-\mu t}\, d\mu} \;\geq\; \frac{e^{-(m_0+\delta)t_1} \int_{m_0+\delta}^{\infty} A(\mu)\, d\mu}{e^{-m_0 t_0} \int_{m_0}^{\infty} A(\mu)\, d\mu} \;>\; 0.$$

Hence $\inf_{t \in \mathcal{I}_{\mathrm{IR}}} \psi(t) \geq m_0 + \delta\, \eta_{\mathrm{IR}} =: m_0 - \varepsilon_{\mathrm{IR}} > 0$. (If all spectral weight sits at $\mu = m_0 = 0$, then $\psi \equiv 0$ and no positive lower bound is possible; this is precisely the massless case excluded by our IR–guard regime, cf. Proposition 12.242.)

(4) Relation to the guard machinery. The Lipschitz/TV bounds and the positive infimum on $\mathcal{I}_{\mathrm{IR}}$ are the guard conditions of Definition 12.200. They are exactly the estimates used in Proposition 12.242 to certify the IR window in the continuum, turning the operational checks into theorems once KL positivity and mild IR regularity hold.

The proposition follows. □

Theorem 12.249 (Continuum guards ⇒ Plateau Theorem)**.** *Assume the continuum KL positivity for typed gauge–invariant correlators (Chs. 4–5; Lemmas 12.190 and 12.212) and the IR regularity of Proposition 12.248. Then the pointwise mask and the morphology of Definitions 12.194 and 12.195 hold* as theorems*: there exist thresholds and radii for which each core plateau $\mathcal{P}_\alpha^\circ$ is nonempty and the certified intersection $\mathcal{P} = \bigcap_\alpha \mathcal{P}_\alpha^\circ$ exists. Moreover, on any IR window $\mathcal{I}_{\mathrm{IR}} \subset \mathcal{P}$, the Lipschitz/TV/positivity guards of Definition 12.200 hold automatically.*

Proof. **Step 1: Typed diagnostics inherit regularity from the kernel via the metric.** By the Chapter 4 backbone and typed construction, every certified diagnostic R_α is an admissible linear (or smoothly composed) functional of the typed, gauge–invariant correlator built from the Fourfold kernel (Lemma 12.233 and Proposition 12.231). The Chapter 5 operator–valued metric and its triadic split control the differentiability and locality of these correlators (eqs. (6.306) and (12.304) and Proposition 12.240). Consequently, on any compact parameter rectangle $\mathcal{D}_0 \subset \mathcal{D}$ and for each α there exist finite bounds

$$\sup_{(u,v)\in\mathcal{D}_0} \|\nabla R_\alpha(u,v)\|_M \;\leq\; B_{\alpha,1}, \qquad \sup_{(u,v)\in\mathcal{D}_0} \|\mathcal{H}_\alpha(u,v)\|_2 \;\leq\; B_{\alpha,2},$$

where the M–metric is the scaled control metric of Definition 12.193. (These bounds are a direct consequence of the Sobolev/metric control of the coincidence Hessian of D^2, together with typed boundedness; cf. the use of such bounds in Propositions 12.234 and 12.242.)

Step 2: IR slope guards hold in the continuum. KL positivity with nonnegative spectral measure and mild threshold regularity imply the mean/variance identities and complete monotonicity for the Euclidean correlator $C(t)$ (Lemma 12.241). On any compact IR window $\mathcal{I}_{\mathrm{IR}} = [t_0, t_1]$ with $t_0 > 0$, Propositions 12.242 and 12.248 give

$$|\psi'(t)| \leq V_{\mathrm{IR}}, \quad \mathrm{TV}_{\mathcal{I}_{\mathrm{IR}}}[\psi] \leq \mathrm{TV}_{\mathrm{IR}}, \quad \inf_{t\in\mathcal{I}_{\mathrm{IR}}} \psi(t) \;\geq\; m_0 - \varepsilon_{\mathrm{IR}} \;>\; 0,$$

with $\psi(t) = -\partial_t \log C(t)$. These are exactly the Lipschitz/TV/positivity guards of Definition 12.200, now obtained as *theorems* in the continuum.

Step 3: Existence of nonempty acceptance sets for the pointwise mask. Fix a compact working domain $\mathcal{D}_0$ that contains $\mathcal{I}_{\mathrm{IR}} \times U$ for some U in the u–axis. Choose the M–scalings $S = \mathrm{diag}(s_u, s_v)$ as in Proposition 12.197 and pick thresholds

$$\varepsilon_{\mathrm{grad}} \;\geq\; \max_{\alpha} B_{\alpha,1}, \qquad \varepsilon_{\mathrm{curv}} \;\geq\; \max_{\alpha} B_{\alpha,2}, \qquad \varepsilon_{\mathrm{osc}} \;\geq\; \max_{\alpha} \sup_{B_r \subset \mathcal{D}_0} \mathrm{osc}_\alpha(B_r),$$

for some $r > 0$ (to be fixed below). Then, by definition of the mask (Definition 12.194), every $(u, v) \in \mathcal{I}_{\mathrm{IR}} \times U$ satisfies the three inequalities and is an *acceptance point* for every channel α. Thus the raw acceptance set for each α is nonempty and contains an open neighborhood of $\mathcal{I}_{\mathrm{IR}} \times U$ by continuity (cf. Lemma 12.198 for quadratic variation across metric balls).

Step 4: Morphology (closing–erosion) produces nonempty cores. By Definition 12.195, the core $\mathcal{P}_\alpha^\circ$ is obtained by first *closing* the acceptance set with a metric ball $B_{r_{\mathrm{close}}}$ and then *eroding* by $B_{r_{\mathrm{erode}}}$ (both in the M–metric). Since the raw acceptance set contains $B_{r_0}(u_*, v_*)$ for every $(u_*, v_*) \in \mathcal{I}_{\mathrm{IR}} \times U$ (by Step 3 and Lemma 12.198), choosing

$$0 \;<\; r_{\mathrm{erode}} \;<\; r_{\mathrm{close}} \;\leq\; r_0$$

ensures that $B_{r_0 - r_{\mathrm{erode}}}(u_*, v_*) \subset \mathcal{P}_\alpha^\circ$. Hence each $\mathcal{P}_\alpha^\circ$ is nonempty and contains an M–ball. The choices are stable under monotone reparameterizations of axes by Proposition 12.197.

Step 5: Nonemptiness of the intersection plateau $\mathcal{P} = \bigcap_\alpha \mathcal{P}_\alpha^\circ$. The set of channels $\mathcal{A}$ is finite (the standard glueball meters; Definition 12.185). Steps 1–4 were performed with thresholds chosen as *uniform* maxima over α, so each $\mathcal{P}_\alpha^\circ$ contains a common M–ball centered on any $(u_*, v_*) \in \mathcal{I}_{\mathrm{IR}} \times U$. Therefore their intersection $\mathcal{P}$ is nonempty and contains an M–ball as well:

$$\emptyset \;\neq\; \mathcal{P} = \bigcap_{\alpha \in \mathcal{A}} \mathcal{P}_\alpha^\circ \;\supset\; B_{r_*}(u_*, v_*).$$

Typed stability and band–reduction (Propositions 12.189 and 12.216) guarantee that these operations commute with the typed projections used to define the meters, so the construction is consistent across channels.

Step 6: Guards hold automatically on $\mathcal{I}_{\mathrm{IR}} \subset \mathcal{P}$. By Step 2, the Lipschitz/TV/positivity guards of Definition 12.200 hold on $\mathcal{I}_{\mathrm{IR}}$ as consequences of KL positivity and IR regularity; since $\mathcal{I}_{\mathrm{IR}} \subset \mathcal{P}$ by construction, the guards hold on the certified plateau as claimed.

Combining Steps 1–6 yields: (i) existence of thresholds and morphology radii for which every $\mathcal{P}_\alpha^\circ$ is nonempty; (ii) nonemptiness of the intersection plateau $\mathcal{P}$; and (iii) automatic validity of the IR guards on $\mathcal{I}_{\mathrm{IR}} \subset \mathcal{P}$. This proves the theorem. □

Corollary 12.250 (Typed spectral gap in the continuum)**.** *On the certified plateau $\mathcal{P}$ supplied by Theorem 12.249, the typed effective mass $m_{\mathrm{eff}}^{(O,\tau)}(t)$ converges monotonically to the first pole $m_O > 0$ and satisfies $m_O \geq m_*^{\mathrm{op}} > 0$; the typed/untyped equality and meter locality hold by construction (Propositions 12.216 and 12.217). Thus the gap is established by Theorem 12.219 without recourse to operational certification.*

Proof. **(1) Continuum plateau and IR guards.** By Theorem 12.249, the pointwise mask and morphology of Definitions 12.194 and 12.195 hold *as theorems* from KL positivity and IR regularity (Chs. 4–5), yielding a certified nonempty plateau $\mathcal{P}$ and an IR window $\mathcal{I}_{\mathrm{IR}} \subset \mathcal{P}$ on which the Lipschitz/TV/positivity guards of Definition 12.200 are automatic (via Propositions 12.242 and 12.248).

(2) KL form and effective–mass convergence. For any typed, gauge–invariant O (Ch. 4), KL/OS positivity on $\mathcal{P}$ gives the spectral (Laplace) form with nonnegative measure Lemmas 12.190 and 12.212. Hence the Euclidean correlator admits the decomposition used in Proposition 12.213,

$$C_O^{(\tau)}(t) = A_0 e^{-m_O t} + \int_{m_*}^{\infty} A(\mu)\, e^{-\mu t}\, d\mu, \quad A_0, A(\mu) \geq 0,$$

and the effective mass (definition (12.284)) is eventually monotone decreasing with

$$\lim_{t\to\infty} m_{\mathrm{eff}}^{(O,\tau)}(t) = m_O\,.$$

(3) Positive lower bound for the first pole. On $\mathcal{I}_{\mathrm{IR}}$, the positivity guard for the log–slope implies the continuum analogue of the operational bound of Corollary 12.202: there exists $m_*^{\mathrm{op}} > 0$ with

$$C_O^{(\tau)}(t) \;\leq\; C_O^{(\tau)}(t_0)\, e^{-m_*^{\mathrm{op}}(t-t_0)} \quad (t \geq t_0 \in \mathcal{I}_{\mathrm{IR}}),$$

and therefore every isolated pole satisfies $m_O \geq m_*^{\mathrm{op}} > 0$ (same argument as in Corollary 12.202, now derived from Propositions 12.242 and 12.248). This shows the *existence* of a strictly positive first mass in the typed channel and bounds it from below.

(4) Typed equality and meter locality. By Proposition 12.216, typed/untyped extractions coincide after band reduction on $\mathcal{P}$,

$$\tilde{C}_O(t) = C_O^{(\tau)}(t), \qquad \tilde{G}_O(p) = G_O^{(\tau)}(p),$$

so the pole mass and residue are type–independent. By Proposition 12.217 (visibility additivity and $(\bar{\Sigma})$–cone locality, Ch. 6), insertions supported outside the causal wedge do not affect connected two–point functions of the typed meters; hence the readout is meter–local.

(5) Conclusion via typed certification. Items (1)–(4) verify the hypotheses of the typed spectral certification theorem Theorem 12.219 *within* the continuum construction: KL form with nonnegative measure, effective–mass convergence, typed equality, meter locality, and a strictly positive IR slope bound. Therefore $m_O > 0$ exists and is certified on $\mathcal{P}$ without invoking any additional, purely operational certification. This establishes a typed spectral gap in the continuum. □

(B) Area law under OS positivity

Theorem 12.251 (OS area law ⇒ axiomatic mass gap (continuum))**.** *Assume: (i) the Euclidean pure Yang–Mills theory satisfies the OS axioms (Lemma 12.222), (ii) the Wilson–loop area law of Definition 12.221 holds for large rectangles, and (iii) the Fourfold transport $\mathcal{U}$ is the holonomy entering both $\hat{G}_\Lambda$ and $W[\mathcal{C}]$ (Definitions 12.228 and 12.229 and Lemma 12.235). Then:*

(i) Exponential clustering

Exponential clustering holds for every local gauge–invariant correlator built as a typed contraction of $\hat{G}_\Lambda$:

$$|\langle O(x)O(0)\rangle_{E,c}| \;\le\; C_{\mathrm{cl}}\, e^{-c_{\mathrm{cl}}\sqrt{\sigma}\,|x|} \qquad (|x| \gg 1),$$

as in Proposition 12.236.

(ii) Positive spectral threshold

The reconstructed Wightman theory has a strictly positive spectral threshold $m \ge c_{\mathrm{cl}}\sqrt{\sigma} > 0$ in every glueball channel (Corollary 12.224).

In particular, the typed effective masses satisfy $\lim_{t\to\infty} m_{\mathrm{eff}}^{(O,\tau)}(t) = m \ge c_{\mathrm{cl}}\sqrt{\sigma} > 0$, agreeing with the pole extraction and rendering Section 12.13.5 a single axiomatic proof of the gap.

Proof. **Step 1: OS reconstruction and typed local meters.** By the OS axioms and the reconstruction theorem (Lemma 12.222), the Euclidean theory defines a Hilbert space $\mathcal{H}$, a self–adjoint Hamiltonian $H \ge 0$, a Poincaré–invariant vacuum $|\Omega\rangle$, and local fields as operator–valued distributions satisfying locality (microcausality) and the spectral condition. Chapter 4 gives, for each gauge–invariant local meter O (e.g. $O_{0^{++}}, O_{0^{-+}}, O_{2^{++}}^{\mathrm{TT}}$), a typed point–split representation as a contraction of the Fourfold kernel with the parallel transporter (Definition 12.229 and Proposition 12.231):

$$O(x) \;=\; \lim_{x'\to x}\; \mathsf{C}\Big[\; \hat{\mathcal{F}}_\alpha^\dagger(x)\,\mathcal{U}(x \leftarrow x';\gamma)\,\hat{\mathcal{F}}_\beta(x')\;\Big],$$

where $\mathsf{C}[\cdot]$ encodes the typed projector/TT contraction (Definition 12.188). Thus $O(x)O(0)$ is a typed contraction of products of the *same* transporter $\mathcal{U}$ entering Wilson loops.

Step 2: From the area law to exponential clustering. By Lemma 12.235, the Wilson loop $W[\mathcal{C}]$ is the normalized trace of the holonomy of $\mathcal{U}$ around $\mathcal{C}$. On rectangles $\square(R,T)$, the area law of Definition 12.221 gives

$$\langle W[\square(R,T)]\rangle_E \;\le\; \exp\Big(-\sigma RT + o(RT)\Big), \qquad \sigma > 0.$$

Using OS reflection positivity and the chessboard/Cauchy–Schwarz estimate for the typed point–split representation of $O(x)O(0)$ (see the domination bound in Proposition 12.236), one obtains

$$|\langle O(x)O(0)\rangle_{E,c}| \;\le\; C_1\, e^{c_1(R+T)}\,\Big\langle W[\square(R,T)]\Big\rangle_E^{\theta},$$

with fixed $C_1, c_1 > 0$ and $\theta \in (0,1]$ depending only on the typed contraction. Choosing $R \sim T \sim |x|$ and inserting the area law yields a Gaussian bound which relaxes to an exponential with scale $\sqrt{\sigma}$, precisely

$$|\langle O(x)O(0)\rangle_{E,c}| \;\le\; C_{\mathrm{cl}}\, e^{-c_{\mathrm{cl}}\sqrt{\sigma}\,|x|},$$

as established in Proposition 12.236. This proves item (1).

Step 3: Exponential clustering ⇒ spectral threshold. By OS positivity, two–points admit a KL/Laplace representation with a nonnegative spectral measure (Lemmas 12.190 and 12.222):

$$C_O(t) = \langle O(t)O(0)\rangle_{E,c} = \int_0^\infty e^{-Et}\, d\nu_O(E), \qquad d\nu_O \ge 0.$$

The temporal slice of the clustering bound gives $C_O(t) \le C'_{\mathrm{cl}} e^{-m_0 t}$ for $t \ge 0$ with $m_0 = c_{\mathrm{cl}}\sqrt{\sigma}$. If $\nu_O([0, m_0)) > 0$ then $C_O(t) \ge \alpha\, e^{-(m_0-\epsilon)t}$ for some $\alpha, \epsilon > 0$, contradicting the previous inequality for large t. Hence $\inf \mathrm{supp}(\nu_O) \ge m_0 > 0$, i.e. a strictly positive threshold $m \ge c_{\mathrm{cl}}\sqrt{\sigma}$ (see also Proposition 12.244 and Corollary 12.224). This proves item (2).

Step 4: Typed effective masses and consistency with pole extraction. Since $d\nu_O$ has a positive threshold and nonnegative density, the Euclidean effective mass

$$m_{\mathrm{eff}}^{(O,\tau)}(t) = \log \frac{C_O^{(\tau)}(t)}{C_O^{(\tau)}(t+1)}$$

is eventually monotone and converges to the first pole m (Proposition 12.213); by the previous step, $m \ge c_{\mathrm{cl}}\sqrt{\sigma} > 0$. This agrees with the typed pole extraction on the plateau and turns Section 12.13.5 into a single axiomatic proof of the gap.

All assumptions were used only at the axioms level: OS positivity (Lemma 12.222), the area law (Definition 12.221), and the identification of the transporter across local meters and loops (Definitions 12.228 and 12.229 and Lemma 12.235). The theorem follows. □

From pipeline into theorems

Corollary 12.252 (Pipeline ⇒ theorem)**.** *Either set of hypotheses—(A) continuum KL positivity with IR regularity (Theorem 12.249) or (B) OS positivity with an area law (Theorem 12.251)—is sufficient to turn the operational Steps B–E (Sections 12.13.2 to 12.13.5) into a single, axiomatic mass–gap proof. In case (A), the certified plateau and IR guards are theorems and the typed spectral extraction yields $m_* > 0$. In case (B), the area law implies exponential clustering and a universal bound $m \gtrsim c\sqrt{\sigma} > 0$, with which the typed extraction coincides.*

Proof. **Case (A): Continuum KL positivity + IR regularity.** By Theorem 12.249, KL positivity together with the IR regularity in Proposition 12.248 promotes the *pointwise mask* and *morphology* of Step B (Definitions 12.194 and 12.195) to *theorems*: each channel admits a nonempty core $\mathcal{P}_\alpha^\circ$ and the certified intersection $\mathcal{P} = \bigcap_\alpha \mathcal{P}_\alpha^\circ \neq \varnothing$ exists; moreover, the Lipschitz/TV/positivity guards of Definition 12.200 hold automatically on any IR window $\mathcal{I}_{\mathrm{IR}} \subset \mathcal{P}$. On $\mathcal{P}$, typed two–point functions enjoy KL form (Lemmas 12.190 and 12.212) and the effective mass is monotonically convergent to the first pole by Proposition 12.213. The typed/untyped equality after band reduction (Proposition 12.216) and meter locality (Proposition 12.217) hold on $\mathcal{P}$. Therefore all hypotheses of the typed certification theorem Theorem 12.219 are satisfied *in the continuum*, which yields a strictly positive mass $m_* > 0$ in each typed glueball channel (cf. also the operational lower bound upgraded in Corollary 12.250). Step C (boundary hygiene) is subsumed by the boundary–stable readout controls (Theorem 12.210) once the masks/guards are theorems; Step D (typed spectral extraction) is exactly Theorem 12.219; Step E (cross–checks) then follows by Section 12.13.5. Hence Steps B–E collapse to an axiomatic proof of the gap in case (A).

Case (B): OS positivity + area law. By the OS axioms and the identification of the transporter in both local meters and Wilson loops (Definitions 12.228 and 12.229 and Lemma 12.235),

the area–law certificate (Definition 12.221) implies exponential clustering for all typed local correlators, with rate $\sim \sqrt{\sigma}$ (Proposition 12.236). Under OS positivity, exponential clustering yields a strictly positive spectral threshold for the reconstructed Hamiltonian on the BRST–physical subspace (Proposition 12.244). Theorem 12.251 summarizes these implications and gives $m \geq c_{\mathrm{cl}}\sqrt{\sigma} > 0$ in every glueball channel. On the same set, typed effective masses converge to the same m (Proposition 12.213) and typed/untyped equality plus meter locality hold by Propositions 12.216 and 12.217. Thus Step E (area–law cross–check) alone already supplies an axiomatic mass gap, and Step D's typed extraction coincides with it; consistency of quadrant/duality properties is provided by Section 12.13.5.

Conclusion. In **(A)**, the continuum guards and plateau construction are theorems and the typed certification (Theorem 12.219) yields $m_* > 0$; in **(B)**, the OS area law gives $m \gtrsim c\sqrt{\sigma} > 0$ (Theorem 12.251) and the typed extraction agrees with this bound. Therefore, under either hypothesis set, the operational pipeline (Sections 12.13.2 to 12.13.5) collapses into a single, axiomatic mass–gap proof, as claimed. □

12.13.10 The Yang-Mills Mass Gap Theorem and OS/Wightman Axioms

We formalize what the operational statements "typed equality, meter locality, CPT/KL hygiene" become once the continuum Yang–Mills theory is constructed under OS/Wightman axioms [578, 579, 255, 91, 82].

Proposition 12.253 (Projection theorem for typed equality)**.** *Let a Euclidean OS–positive (hence Wightman–reconstructible) pure Yang–Mills theory be given. For each admissible type τ, let Π_τ be the gauge–invariant band projector obtained from OS/Wightman spectral calculus on the realized band, and assume Π_τ is constructed by gauge–invariant band limits so that, in the sense of distributions,*

$$[\Pi_\tau,\ \mathfrak{F}_{\mathrm{OS/W}}] = 0,$$

with $\mathfrak{F}_{\mathrm{OS/W}}$ the OS/Wightman field algebra of gauge–invariant local operators. Then for every gauge–invariant local O and its typed witness $W^{(\tau)} := \Pi_\tau O \Pi_\tau$,

$$\langle W^{(\tau)}(x)\, W^{(\tau)}(0)\rangle \;=\; \langle \Pi_\tau O(x)\Pi_\tau\ \Pi_\tau O(0)\Pi_\tau\rangle \;=\; \langle O(x)\, O(0)\rangle\Big|_{\text{band }\Pi_\tau}, \tag{12.313}$$

i.e. typed/untyped correlators coincide after restricting the Hilbert space *to $\Pi_\tau\mathcal{H}$. In particular, $G_O^{(\tau)}(p) = \tilde{G}_O(p)$ and $C_O^{(\tau)}(t) = \tilde{C}_O(t)$ as in Proposition 12.216. (For OS positivity and Wightman reconstruction see [578, 579, 255, 91].)*

Proof. The distributional commutation with the field algebra implies that inserting Π_τ may be done either at operator level or at state/space level without changing expectation values. Thus, for gauge–invariant O, $\Pi_\tau O \Pi_\tau$ is precisely the compression of O to the band $\Pi_\tau\mathcal{H}$, yielding (12.313). This is the axioms–level version of Proposition 12.216. (Compare the standard use of projectors and spectral calculus in the OS/Wightman framework [579, 255].) □

Proposition 12.254 (Meter locality from microcausality and additivity)**.** *Let $W^{(\tau)} := \Pi_\tau O \Pi_\tau$ be a typed gauge–invariant local meter and let $\mathcal{O}_{\mathrm{ext}}$ be any operator localized outside the*

($\bar{\Sigma}$)*–causal domain of* $\operatorname{supp} W^{(\tau)}$*. If the Wightman locality (microcausality) holds, then the connected correlators are independent of* $\mathcal{O}_{\mathrm{ext}}$*:*

$$\langle \mathcal{O}_{\mathrm{ext}}\, W^{(\tau)}(x) W^{(\tau)}(0)\rangle_c = \langle W^{(\tau)}(x) W^{(\tau)}(0)\rangle_c\,, \qquad \Lambda_{\mathrm{law/state}}[\,\rho_0;\ \mathcal{O}_{\mathrm{ext}}\,] = \Lambda_{\mathrm{law/state}}[\,\rho_0;\ \mathbb{I}\,], \tag{12.314}$$

with $\Lambda_{\mathrm{law/state}}$ *the visibilities of Definition 12.188. Hence* meter locality *is just Wightman microcausality plus additivity of conserved currents. This is the axioms–level version of Proposition 12.217. (See [255, Chs. II–III] and [91, §2] for locality/microcausality and cluster/additivity.)*

Proof. Microcausality implies $[\mathcal{O}_{\mathrm{ext}}, W^{(\tau)}] = 0$ in the causal complement, whence connected insertions of $\mathcal{O}_{\mathrm{ext}}$ factorize and cancel. The visibility additivity (Definition 12.188) then yields (12.314). (For standard proofs of factorization/cluster under locality, see [255, 91, 82].) □

Proposition 12.255 (CPT/KL hygiene under OS/Wightman; duality inertness for adjoint YM)**.** *If the Euclidean theory satisfies the OS axioms (Euclidean invariance, reflection positivity, regularity), then every gauge–invariant two–point function admits a Källén–Lehmann representation with a nonnegative spectral measure, and the reconstructed Wightman theory enjoys an antiunitary CPT symmetry on the one–particle sector. Therefore, isolated poles and residues are CPT–consistent as in Lemma 12.190 and Proposition 12.154. In the* pure *Yang–Mills sector, the multiplet is adjoint (real), so the duality (charge–conjugation) action is trivial by Proposition 12.158; equivalently, channels are duality–inert (Corollary 12.191 and Proposition 12.218). (KL and OS→Wightman: [382, 383, 578, 579]; CPT in the axiomatic setting: [255, 91].)*

Proof. OS reflection positivity ⇒ KL positivity and CPT after OS→Wightman reconstruction; see Lemmas 12.212 and 12.222 and Proposition 12.154 and [579, 255]. For adjoint YM, reality of the representation implies self–conjugacy (Proposition 12.158), so the CPT/duality map fixes the channels (Corollary 12.191 and Proposition 12.218). □

Chapters 4 and 5 already package the structures required to *remove the word "diagnostic realized reference configuration"*: (i) they identify the precise gauge–covariant correlator kernel and transport ($\hat{G}_\Lambda^{\alpha\beta}$, $\mathcal{U}$) that the axioms must control (Definitions 12.228 and 12.229), and (ii) they turn spectral positivity into geometric/regularity statements via the operator–valued metric with its triadic split $\hat{g}_{\mu\nu} = \hat{g}^{\mathrm{corr}} + \hat{g}^{\mathrm{vac}} + \hat{g}^{\mathrm{spec}}$ (Definitions 12.238 and 12.239). Proving either **(A)** the uniformity/IR–guard statements directly for continuum YM correlators (KL positivity with IR control; Section 12.13.9 and Theorem 12.249) or **(B)** an area law under OS positivity (Theorem 12.251) collapses the entire operational pipeline (Sections 12.13.2 to 12.13.5) into a rigorous mass–gap theorem. In this axioms–level setting, typed equality is a projection theorem (Proposition 12.253), meter locality is an immediate corollary of microcausality (Proposition 12.254), and CPT/KL hygiene is guaranteed by OS/Wightman reconstruction with adjoint duality inertness for pure YM (Proposition 12.255); cf. [579, 255, 580].

Theorem 12.256 (Yang–Mills Mass Gap (typed, axiomatic))**.** *Consider four–dimensional* pure *Yang–Mills with compact simple gauge group G. Let O range over the standard gauge–invariant glueball operators (Definition 12.185, e.g.* 0^{++}*,* 0^{-+}*,* 2^{++} *(TT)), and let* $C_O^{(\tau)}$ *denote the*

corresponding typed *two–point functions built with admissible meters (Definition 12.188). Each of the following hypothesis–sets is sufficient—and both give the same conclusion— to establish a spectral gap on the physical Hilbert space:*

(i) Route A (KL + IR regularity)

If the continuum typed, gauge–invariant two–point functions admit a Källén–Lehmann representation with a nonnegative spectral measure and mild infrared regularity (so that the log–slope is Lipschitz/TV–controlled and uniformly positive on an IR window as in Propositions 12.242 and 12.248), then the Continuum Guards/Plateau Theorem *holds (Theorem 12.249). On the certified plateau $\mathcal{P}$ one has:*

- **(a)** *the effective mass $m_{\mathrm{eff}}^{(O,\tau)}(t)$ is monotone for large t and converges (Proposition 12.213);*
- **(b)** *each channel's two–point possesses a* first pole at strictly positive mass $m_O > 0$ *and satisfies $m_O \geq m_* > 0$ (Theorem 12.219 and Corollary 12.250);*
- **(c)** *hence the physical Hamiltonian has a gap,* $\mathrm{spec}(H_{\mathrm{phys}}) = \{0\} \cup [m_*, \infty)$ *with $m_* > 0$.*

(For the KL framework see [382, 383, 91, 82].)

(ii) Route B (OS + area law)

If the Euclidean theory satisfies the OS axioms and rectangular Wilson loops obey an area law *with string tension $\sigma > 0$ (Definition 12.221), then:*

- **(a)** *all local, gauge–invariant correlators* cluster exponentially *(Proposition 12.236);*
- **(b)** *the reconstructed Wightman theory exhibits a* universal spectral threshold $m \geq c_{\mathrm{cl}}\sqrt{\sigma} > 0$ *in every glueball channel (Theorem 12.251 and Corollary 12.224);*
- **(c)** *typed effective masses converge to the same $m > 0$ (Proposition 12.213).*

(Standard sources on OS axioms and area law/confinement: [579, 389, 575, 580].)

Invariant features (both routes). *The result is* typed, gauge–invariant, BRST–clean, microcausal, *and* CPT/KL–hygienic*: typed/untyped extractions coincide after band reduction (Proposition 12.253); meters are cone–local (no dependence on operations outside the causal wedge; Proposition 12.254); and the KL form with CPT pairing holds at the axioms level (Proposition 12.255). In pure YM the adjoint multiplet is self–conjugate (duality inert; Proposition 12.158). Concretely, for each standard glueball operator O, $G_O^{(\tau)}(p)$ has KL form with nonnegative measure and a first pole $m_O \geq m_* > 0$. The equivalence of the operational pipeline with these axioms–level statements is summarized in Corollary 12.252.*

Remark 12.257 (Scope)**.** The theorem pertains to *pure* Yang–Mills at zero temperature. With dynamical quarks, light (pseudo–)Nambu–Goldstone modes can appear in the chiral limit, so a gap in *every* singlet channel is not implied. Electroweak nonperturbative phenomena reside in $SU(2)_L \times U(1)_Y$, not $SU(3)_C$. For background and context, see [580, 575, 91].

What is now a theorem (not a procedure). We established two independent, axioms–level routes to a positive first pole in the standard glueball channels. **Route A**: from the Chapter 4–5 backbone (Fourfold transporter and unified correlator; operator–valued metric), continuum KL–positivity with mild IR control upgrades the plateau/guard workflow into the *Continuum Guards/Plateau Theorem* Theorem 12.249. On that certified region, typed spectral extraction has KL form, the effective mass is monotone and convergent, and the first pole satisfies $m_* > 0$ (via Proposition 12.213, Theorem 12.219, and Corollary 12.250); compare the KL framework and spectral consequences in [382, 383, 91, 82]. **Route B**: under OS reflection positivity, an *area law* for large rectangular Wilson loops forces *exponential clustering* and a *universal bound* $m \gtrsim c\sqrt{\sigma} > 0$, giving the *OS area–law mass–gap theorem* Theorem 12.251; see [578, 579] for OS→Wightman reconstruction and [389, 575] for the area–law confinement paradigm. At the axioms level, former "diagnostic realized reference configuration checks" become structural facts: **typed equality** is a projection theorem Proposition 12.216; **meter locality** is microcausality plus visibility additivity Proposition 12.217 (cf. [255, 91]); **CP-T/KL hygiene** follows from OS/Wightman reconstruction and the KL form Lemma 12.190 and Proposition 12.154 [579, 382, 383, 255]; and in pure YM (adjoint) the duality map is inert Proposition 12.158. The equivalence of the operational pipeline with the axioms–level statements is summarized in Corollary 12.252.

Why this still is not automatically "Clay–solved." The Clay problem demands: construct 4D pure YM with compact simple G satisfying OS/Wightman axioms and prove a mass gap—*without* importing nonperturbative inputs as assumptions [580]. In our theorems, either **(A)** KL–positivity + specific IR regularity for continuum correlators (the hypotheses that power Theorem 12.249 and Propositions 12.242 and 12.248), or **(B)** an OS–positive *area law* (the hypothesis of Theorem 12.251 and Definition 12.221), is taken as an input. Those are precisely the deep, nonperturbative statements the Clay problem asks one to derive *inside YM* [580]. Until they are obtained from the Yang–Mills action itself, the result is an axiomatic mass–gap theorem *conditional* on those inputs, not the unconditional Clay solution.

What would upgrade our theorem to a Clay–level solution. (i) **Continuum construction:** build OS–positive Euclidean YM and reconstruct Wightman fields (Chapter 5; Lemma 12.222) [578, 579]. (ii) **Either** prove *KL positivity with IR regularity* for gauge–invariant two–point functions, enough to trigger our guards and effective–mass convergence (Propositions 12.242 and 12.248 and Theorem 12.219) [382, 383, 91], **or** prove an *area law* within that OS–positive YM (Definition 12.221), which via transporter–identification and OS→KL yields clustering and a spectral threshold (Lemma 12.235, Propositions 12.236 and 12.244, and Theorem 12.251) [389, 579, 575]. Either path closes the loop because Chapters 4–5 already reduce both targets to properties of the *same* kernel/metric machinery (Theorems 12.237 and 12.246).

Why this remains a major advance. We collapsed the problem to two crisp nonperturbative targets tied to a single mathematical spine: the *Fourfold kernel* $\hat{G}_\Lambda^{\alpha\beta}$ and transporter $\mathcal{U}$ (Ch. 4; Definitions 12.228 and 12.229), and the *operator–valued metric* with its triadic split (Ch. 5; eqs. (6.306) and (12.304)). All operational loopholes—gauge–parameter artifacts, boundary/drive contamination, causal contamination, representation dependence—have

been turned into theorems at the axioms level (Propositions 12.216, 12.217 and 12.240 and Lemma 12.241), and our two routes (spectral/guards vs. area–law/clustering) agree on a positive gap, with cross–checks packaged in Section 12.13.5. Under standard continuum axioms plus *either* KL+IR regularity *or* OS+area law, the mass gap follows by our theorems, with typed, BRST–clean, microcausal meters and CPT/KL control ensuring robustness [579, 382, 383, 255]. If one *adopts* either input as part of the axioms, the gap is solved within that axiomatic system. For the Clay prize as stated, the remaining bridge is to *derive* one of those inputs *from YM itself*—at which point Corollary 12.252 makes the operational pipeline fully collapse into a single, unconditional mass–gap theorem [580].

Nonperturbative backbone for the color sector. In the pure $SU(3)_C$ limit, the Yang–Mills Mass Gap Theorem (Routes A/B; Theorems 12.249 and 12.251) yields a strictly positive first pole in the standard glueball channels (Definition 12.185). Consequently, "confinement + glueball existence'' is promoted from working assumption to theorem on the continuum axioms employed, consistent with lattice and continuum expectations [389, 575, 576, 577]. The BRST–cohomological construction of hadrons/anti–hadrons is then compatible with a gapped gluonic background (Corollary 12.159 and Theorem 12.133; cf. [108, 442]). Moreover, the adjoint multiplet is self–conjugate and duality–inert (Proposition 12.158), so no separate "anti–gluon'' sector arises; antimatter pairing appears at the level of quark kymons ($R \leftrightarrow \bar{R}$) and of bound states (hadron $\leftrightarrow$ anti–hadron) per the CRP/Duality formalism (Definition 12.151 and Corollary 12.157).

Poles, residues, and typed meters. On the certified continuum setup, typed, gauge–invariant two–point functions admit a Källén–Lehmann form with nonnegative measure (Lemma 12.190 and Theorem 12.219) [382, 383]. Hence the identification "first pole = particle mass'' for glueballs is axiomatic, with monotone effective–mass convergence (Proposition 12.213 and Corollary 12.250); see also standard glueball correlator practice [576, 577]. Clustering implied by the gap ensures meter locality and representation–independence: typed/untyped equality is a projection theorem (Proposition 12.216), and meter locality follows from microcausality with visibility additivity (Proposition 12.217) [255, 91].

Antimatter structure in the SM context. The Axiom (Principle) of Duality explains antiparticles through $R \oplus \bar{R}$ closure and conjugate poles (Propositions 12.152 and 12.154) [255, 91]. The mass gap theorem complements this by constraining where conjugate pairs can appear in the observable spectrum: only as color singlets, since colored asymptotic states are excluded (area law / confinement) [389, 575]. Thus quarks and antiquarks bind into mesons and baryons/anti–baryons (Corollary 12.159); glueballs are self–conjugate (Proposition 12.158); and family–fibre phases conjugate between particle/antiparticle channels without contamination by long IR tails because exponential clustering damps such tails (Definition 12.156 and Propositions 12.236 and 12.244) [579].

Interface with effective theories and flavor physics. A positive gap in the pure gauge sector justifies the separation of scales underlying chiral EFT and related low–energy descriptions, while supporting OPE- and sum–rule–type arguments for short–distance processes in which

hadrons couple to electroweak currents. Although the mass gap theorem does not itself resolve strong–CP or the origin of neutrino masses, it sharpens nonperturbative QCD inputs to flavor physics by guaranteeing clustering, typed stability, and gauge–parameter independence in the color sector (Propositions 12.216 and 12.217 and Lemma 12.190) [91, 82].

Scope and caveats. The theorem is established for pure Yang–Mills at zero temperature. With dynamical quarks, (pseudo–)Nambu–Goldstone modes can be light in the chiral limit, so a gap in *every* singlet channel is not implied; electroweak nonperturbative phenomena (e.g. sphalerons) lie in $SU(2)_L \times U(1)_Y$, not $SU(3)_C$. Within this scope, the results provide the nonperturbative anchor required in Chapter 12: confinement and glueball existence as theorems, robustness of gauge–invariant poles and residues, and a clean separation between color dynamics and family mixing, all coherently integrated with the duality/CRP structure of matter and antimatter (Theorems 12.249 and 12.251 and Corollary 12.252); background: [580, 575].

Part V

Cosmology

Chapter 13

Dark Energy and Dark Matter

13.1 Dark Energy and Dark Matter

The modern cosmological paradigm, known as the ΛCDM model, rests upon a universe whose composition is profoundly enigmatic, dominated by two unseen components: **Dark Matter** and **Dark Energy** [414, 581]. Observational evidence for Dark Matter is overwhelming and dates back to the 1930s, when Fritz Zwicky observed that galaxies in the Coma cluster were moving too rapidly to be gravitationally bound by visible matter alone [582]. Later, galactic rotation curves studied by Vera Rubin and colleagues revealed that spiral galaxies rotate at roughly constant speeds far beyond their luminous boundaries, requiring the presence of massive, invisible "halos" of non-baryonic matter [583].

This conclusion is independently corroborated by a multitude of cosmological phenomena, including the gravitational lensing of distant galaxies [584, 585], the anisotropies in the Cosmic Microwave Background [581, 136], and the distribution of galaxies in large-scale structure surveys [310]; see also the baryon acoustic oscillation standard ruler and weak-lensing cosmic-shear constraints [151, 152, 153, 154], all of which require the gravitational influence of a cold, non-interacting substance to match observations.

While Dark Matter provides the gravitational scaffolding for cosmic structures, **Dark Energy** governs the universe's ultimate fate. In 1998, two independent collaborations—the Supernova Cosmology Project and the High-Z Supernova Search Team—discovered that the expansion of the universe is accelerating, based on observations of Type Ia supernovae [134, 135]. This unexpected result implies the existence of a pervasive energy component with strongly negative pressure, consistent with a cosmological constant Λ in Einstein's field equations [149].

Dark Energy now appears to constitute nearly 70% of the total energy density of the universe [581, 136], driving its accelerated expansion and posing one of the most profound mysteries in modern physics. The quantum vacuum energy predicted by quantum field theory is more than 10^{120} times larger than the observed value of the cosmological constant—an extreme discrepancy known as the **cosmological constant problem** [211, 145, 148].

Despite their dominant roles, the fundamental nature of both Dark Matter and Dark Energy remains unknown, representing the most significant lacuna in our understanding of the cosmos. Searches for Dark Matter particles include deep underground direct detection

experiments [586, 587], astrophysical indirect detection [588, 589], and production attempts at high-energy colliders such as the LHC [590, 591]. Yet, no definitive candidate has been found.

The mystery surrounding Dark Energy is even more profound: it resists identification with any known field or particle, and its small but non-zero value may suggest new physics beyond the Standard Model and general relativity [149]. Together, these unknowns—comprising over 95% of the universe—strongly indicate that our current theoretical frameworks are incomplete and that a deeper, relational, and unified theory is required to resolve the nature of the **dark sector**.

Definition 13.1 (Operational Dark–Sector Criterion)**.** We define a *dark gauge sector* G operationally through holosymmetric tomography. A state $|\psi\rangle$ belongs to G precisely when it satisfies:

1. **Holosymmetric visibility:** it has nonzero projection under the *coherence projector* $\mathcal{P}_G^{\mathrm{coh}}$ (Definition 12.77), i.e. it is visible to holosymmetric probes in the G–channel;
2. **Spectral decoupling:** it remains orthogonal to all Standard Model observables, in the sense that
$$\langle\psi|\mathcal{O}_{\mathrm{vis}}|\psi\rangle \;\approx\; 0 \qquad \forall\, \mathcal{O}_{\mathrm{vis}} \in \mathcal{A}(G_{\mathrm{SM}}).$$

This criterion formalizes the long–standing intuition behind hidden–sector dynamics in strictly operational terms [548, 549, 550], demonstrating how the holosymmetric apparatus discriminates visible from dark degrees of freedom without any additional ontological assumptions.

13.1.1 Dark Energy as an Emergent Phenomenon of the Dual-Field Vacuum

The standard cosmological model, ΛCDM, has achieved remarkable empirical success, accurately fitting a broad range of observational data from the cosmic microwave background to large-scale structure formation [414, 581, 136]. Yet it remains ontologically opaque: it posits that approximately 68.7% of the universe's total energy density resides in a mysterious *Dark Energy*, while about 26.4% is attributed to *Cold Dark Matter*, leaving less than 5% for ordinary baryonic matter [136]. These two unseen constituents, which collectively account for more than 95% of the cosmos, are essential to the model's predictive power but lack a known physical basis [134, 135, 313, 149, 148].

Within the *IQR framework*, however, this dark sector is not an ad hoc addition but a natural consequence of the spectral entanglement and relational dynamics between the two foundational fields—the *Gaiamiti* and the *Tantrakyma*. Concretely, the dual-field vacuum is characterized on the emergent/GNS arena by the unified inter-modal correlator $\hat{\mathbf{G}}_\Lambda$ and its graded transport (Definitions 9.5 and 9.7), realized on a certified plateau via the Σ–map (Definitions 9.58 and 9.60). The sector actions and their covariant split supply the dynamical backbone (eqs. (6.314) and (6.315)), while the typed weave (TT isolation and τ–kairos decomposition) keeps the tensorial graviton channel separate from non-TT temporal registers (Definitions 10.3 and 10.4). What appears as dark energy and dark matter in conventional models thus emerges here from the intrinsic chronogeometric structure and quantum relationality of the dual-field vacuum itself.

At the heart of the cosmological constant problem lies the extreme mismatch between the vacuum energy density predicted by quantum field theory and the exquisitely small value observed in cosmic expansion [211, 212, 313, 148]. The IQR framework resolves this discrepancy not with arbitrary fine-tuning, but through a natural mechanism of *dynamical self-regulation* governed by diagnostic realized reference configuration-disciplined realization and gauge/Nielsen safety (Definition 9.58 and Theorem 9.351).

The vacuum energy of the universe receives contributions from both the geometric Gaiamiti field, $\langle \mathcal{V}_\Phi \rangle$, and the matter-like Tantrakyma field, $\langle \mathcal{V}_\Theta \rangle$ (the latter being directly tied to its Higgs/radial operator in the EWSB analysis; cf. Section 12.4.5 and eq. (9.185)). The total vacuum contribution to the stress-energy tensor, which drives cosmic expansion, is proportional to their sum:

$$\bar{T}^{\mu\nu}_{\text{vac}} = -\,\bar{\Sigma}^{\mu\nu}\Big(\langle \mathcal{V}_\Phi \rangle + \langle \mathcal{V}_\Theta \rangle\Big), \tag{13.1}$$

where $\bar{\Sigma}^{\mu\nu}$ is the inverse chronogeometric tensor compatible with the Relational Levi–Civita connection (Definitions 5.26 and 6.7). We propose that a fundamental feedback mechanism, encoded in the total relational action $\mathcal{S}_{\text{total}}$ (Definition 9.357) and operating within the band/realization pipeline (Definitions 9.58 and 9.60), drives the coupled system to a ground state where these two colossal energies are of opposite sign and nearly cancel. The observed dark energy is therefore the tiny, non-zero residual of this cosmic equilibrium:

$$\rho_{\text{DE}} \sim \langle \mathcal{V}_\Phi \rangle + \langle \mathcal{V}_\Theta \rangle \approx \varepsilon_\Lambda, \qquad \text{with } \varepsilon_\Lambda \ll \Big|\langle \mathcal{V}_\Theta \rangle\Big|. \tag{13.2}$$

This cancellation is stable under gauge fixing and parameter variations thanks to the TT isolation of the graviton and Nielsen/BRST independence on $\mathfrak{P}$ (Definition 10.3 and Theorem 9.351).

This mechanism explains the *magnitude* of dark energy. The *dynamics* of the residual are governed by the *Spin-0 Apeiron* (Gaiaton), the scalar quantum isolated on the τ–scalar register by the typed weave's Riesz selectors (Section 10.8 and Definitions 10.68 and 10.69). Acting as a quintessence field, the Apeiron slowly rolls down the residual potential landscape, driving the present gentle acceleration while preserving TT purity of gravitational waves (Definition 10.3) and remaining consistent with late-time constraints on w from CMB, BAO, and weak-lensing cosmic shear [486, 592, 581, 136, 151, 152, 153, 154]. In this view, the same scalar mode that sourced early-time dynamics flows continuously to late-time dark-energy behavior, unifying "inflaton" and "quintessence" within the dual-field vacuum's typed decomposition (Section 10.8) and remaining compatible with the SM selection and BRST-physical space developed earlier (Theorem 12.42 and Section 12.3); see also the propagation constraints from GW170817/GRB170817A on modified-gravity degrees of freedom [513].

13.1.2 Dark Matter as the Hidden Quanta of the Relational Fabric

Within the IQR framework, the existence of a dark sector is not an ad hoc insertion but an inevitable consequence of the τ–kairos operator (Definition 10.13), which captures the temporal/coherence fluctuations of the chronogeometric tensor. By the operational criterion of Chapter 12, any excitation that projects non–trivially into the τ–kairos channel while remaining spectrally orthogonal to the Standard Model algebra belongs to the dark sector.

Formally, with the TT–projector and typed τ–decomposition (Definitions 10.3 and 10.4),

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\hat{\tau}_{\mu\nu}=0,\qquad \mathbf{1}_\tau=S^{(b)}\oplus S^{(L)}\oplus \tilde{S}^{(p)}\oplus A^{(+)}\oplus A^{(-)}\quad(\text{TT–null}),$$

so all τ–excitations are non–TT and automatically graviton–safe. This yields *three* constitutive particle–level candidates and one relational alternative.

Candidate I: The Spin–1 Chrono–Kymon (Vector τ–Kairos Excitation)

The **Chrono–Kymon** is the spin–1 register of the τ–kairos tensor, $\hat{\tau}_{\mu\nu}$ (Proposition 10.154), carried by the typed helicity projectors $A^{(\pm)}=\frac{1}{2}(\mathbf{1}_\tau\pm\mathsf{H}_\tau)$ (Definition 10.4 and eq. (10.7)). Gauging the τ symmetry and breaking it yields the BEH mass matrix (eq. (11.339))

$$(M^2_{\tau\text{-vec}})_{IJ}=g_\tau^2\left\langle \mathcal{R}^{(S)}_\tau(T_I)\,\varphi^{[S]}\,,\,\mathcal{R}^{(S)}_\tau(T_J)\,\varphi^{[S]}\right\rangle,\qquad (M^2_{\tau\text{-vec}})^I{}_J:=G^{(\tau)\,IK}(M^2_{\tau\text{-vec}})_{KJ},$$

and a relational Proca–type dynamics (eqs. (11.340) and (11.341))

$$\bar{\nabla}_\mu\mathcal{F}^{I\,\mu\nu}+(M^2_{\tau\text{-vec}})^I{}_J\,\mathcal{B}^{J\,\nu}=J^{I\,\nu}_{(\tau)},\qquad \bar{\nabla}_\nu\mathcal{B}^{I\,\nu}=0.$$

On FLRW diagnostic realized reference configuration the rapidly oscillating regime reproduces cold–DM kinematics, $\rho_K\propto a^{-3}$, $w_K\simeq 0$ (Proposition 13.73), while TT–purity of gravity is guaranteed by $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\hat{\tau}=0$ (Definition 10.3).

Candidate II: The Spin–0 Gaiaton/Apeiron (Scalar τ–Kairos Excitation)

The **Gaiaton** (massive *Apeiron*) is the scalar register of the τ–kairos tensor (Proposition 10.153 and Section 10.17.1). It is isolated by Riesz selectors of the τ–scalar generalized eigen-operator pair $K^{(0)}_\tau$ and projected to the BRST–physical space via (Definitions 10.68 and 10.69)

$$\mathsf{P}^{(0)}_{\mathrm{spec}}=\sum_{m\in\mathfrak{S}_0}\frac{1}{2\pi i}\oint_{\zeta_m}(K^{(0)}_\tau-z)^{-1}dz,\qquad \mathcal{P}^{(0)}_{\mathrm{G},\eta}=\mathsf{P}_{\mathrm{phys}}\,S^{(\eta)}\Big(\Pi^{(\mathrm{spec};0)}_\tau\oplus\Pi^{(\mathrm{vac})}\oplus\Pi^{(\mathrm{corr})}\Big)S^{(\eta)}\mathsf{P}_{\mathrm{phys}}.$$

Its mass arises from the scalar/radial Hessian along the typed direction,

$$m^2_{\mathrm{Gaiaton}}\;\in\;\mathrm{spec}\Big(\Pi^{[S]}_\tau\left[\frac{\delta^2\mathcal{V}_\Phi}{\delta\hat{\Phi}\,\delta\hat{\Phi}}\Big|_{\bar{\Phi}}\right]\Pi^{[S]}_\tau\Big),$$

so that in the *massive* (coherent–oscillation) branch it behaves as scalar cold dark matter ($\rho\propto a^{-3}$), whereas in the *slow–roll* branch its potential $\mathcal{V}_\Phi$ yields a quintessence–like dark–energy component (Chapter 13). TT isolation is automatic (Definition 10.3).

Candidate III: Spin–1 Connection Quanta (Mition & Tantron)

Beyond the τ–kairos, the universal operator–bundle connection $\mathcal{A}_\mu$ supports spin–1 quanta that are neutral under visible meters yet can populate the dark sector on a certified plateau. In the *Gaiamiti* sector one obtains the **Mition** (Definition 9.258); in the *Tantrakyma* sector,

the **Tantron** (Definition 9.348). Their BEH masses follow from the typed covariant split (eq. (6.314) and Definition 9.252):

$$(M^2_\Phi)_{IJ} = g^2\, g_T\Big(\rho_\Phi(T_I)\bar\Phi,\ \rho_\Phi(T_J)\bar\Phi\Big), \qquad (M^2_\Theta)_{IJ} = g^2\, g_C\Big(\rho(T_I)\bar\Theta,\ \rho(T_J)\bar\Theta\Big),$$

with mass–eigenstate vectors $V_{a\,\mu} = O^{\ I}{}_a A^I_\mu$ and $W_{a\,\mu} = O^{\ I}{}_a A^I_\mu$ (Propositions 9.257 and 9.347). Typed Riesz/spectral selectors on a band isolate their BRST–physical transverse poles (Definitions 10.6 and 10.7), and TT–safety follows from projector commutations on the weave (Definition 10.3 and Proposition 10.9). Cosmologically, in the non–relativistic regime both behave as CDM ($\rho \propto a^{-3}$), providing *vector* dark–matter candidates alongside the Chrono–Kymon.

Relational Alternative: "Dark Gravity" from Entangled Energy

The triadic decomposition of the matter–geometry bi–kernel (Section 10.19.6) shows that gravitational response depends on correlational and vacuum structures in addition to spectral modes. Accordingly, some phenomenology attributed to dark matter may instead emerge as *dark gravity*: an effective modification of the Einstein tensor sourced by coherence/entanglement, *without* introducing new particle species (see, e.g., [593, 594, 595, 596]). Within the present framework this is not independent of the τ–kairos sector: as defined in Definition 10.13, $\hat\tau_{\mu\nu} = \delta\hat t_{\mu\nu}$ controls temporal/coherence fluctuations, whose expectation $\bar t_{\mu\nu}$ shifts the diagnostic realized reference configuration geometry and the TT operator according to the typed balance equations (eqs. (11.291) and (11.292)). Thus "dark gravity" is the macroscopic manifestation of the same τ operator whose quanta (Candidates I–III) populate the particle–level dark sector; it is best viewed as an alternative *interpretation* of τ–kairos contributions to cosmological dynamics.

Dark gravity as the coherent τ–kairos background. In the IQR framework the most natural realization of "dark gravity" is the macroscopic, coherent contribution of the τ–kairos sector itself. By definition $\hat\tau_{\mu\nu} := \delta\hat t_{\mu\nu}$ (Definition 10.13) is *TT–null*, $\Pi^{[\mathrm{TT}]}_{\bar\Sigma}\hat\tau_{\mu\nu} = 0$ (Definition 10.3), so no additional spin–2 degrees of freedom are introduced—consistent with multimessenger constraints on extra GW polarizations and modified GW propagation [513]. Instead, the expectation value $\bar t_{\mu\nu}$ shifts the diagnostic realized reference configuration background and the TT wave operator through the typed balance relations

$$\mathcal{L}_{hh}[\bar\Sigma'] = \mathcal{L}_{hh}[\bar\Sigma] + \varepsilon\,\delta\mathcal{L}_{hh}\Big[\Delta\Sigma^{(\tau)}\Big] + \cdots, \qquad \delta\bar G_{\mu\nu}\Big[\Delta\Sigma^{(\tau)}\Big] + \Lambda\,\Delta\Sigma^{(\tau)}_{\mu\nu} = \kappa\,\Delta\bar T^{(\tau)}_{\mu\nu}, \quad (13.3)$$

as established in eqs. (11.291) and (11.292). On a certified plateau $\mathfrak{P}$ (Definition 9.58) these deformations are BRST/Nielsen–safe (Theorem 9.351) and preserve TT purity. Thus the dominant and cleanest route to "dark gravity" is the coherent, non–particle background of the τ–kairos that modifies the effective Einstein sector without generating extra tensor polarizations.

Coherent Gaiaton/Apeiron background as dark gravity (and dark energy). A closely related macroscopic channel arises from the spin–0 Gaiaton (Apeiron) on the τ–scalar register.

The Gaiaton is isolated and projected by the Riesz/typed weave apparatus (Definitions 10.68 and 10.69), and its cosmological realization is summarized in Proposition 10.153. In a nearly homogeneous, slow–roll branch it furnishes a quintessence–like component that shifts the effective background stress while remaining TT–null (again by Definition 10.3); this is consistent with canonical quintessence dynamics and late–time constraints on w from CMB, BAO and WL [486, 592, 581, 136, 151, 152, 153, 154]. In the massive, coherent–oscillation branch the same scalar behaves as cold dark matter—with a characteristic small–scale cutoff—mirroring standard misalignment physics [597, 598]. Both interpretations coexist within the same projector–certified species.

Chrono–Kymon background: a constrained, subleading route. The spin–1 Chrono–Kymon (CK) occupies the τ–vector register with helicities $A^{(\pm)}$ (Definition 10.4 and eq. (10.7)). When the τ symmetry is gauged and spontaneously broken, the CK acquires a BEH mass with matrix

$$(M^2_{\tau\text{-vec}})_{IJ} = g_\tau^2 \left\langle \mathcal{R}^{(S)}_\tau(T_I)\,\varphi^{[S]}\,,\, \mathcal{R}^{(S)}_\tau(T_J)\,\varphi^{[S]} \right\rangle, \qquad \bar\nabla_\mu \mathcal{F}^{I\,\mu\nu} + (M^2_{\tau\text{-vec}})^I{}_J\, \mathcal{B}^{J\,\nu} = J^{I\,\nu}_{(\tau)}, \quad (13.4)$$

(eqs. (11.339) to (11.341)). On FLRW diagnostic realized reference configuration, rapidly oscillating CK backgrounds obey $\rho_K \propto a^{-3}$ and $\langle w_K\rangle \simeq 0$ with vanishing anisotropic stress (Proposition 13.73). Only if a nearly isotropic, coherent CK background forms could it contribute to an effective "dark gravity" description; observational isotropy makes this a subleading pathway compared to the $\bar t_{\mu\nu}$ and Gaiaton channels.

Connection vectors (Mition, Tantron): matter–like, not dark gravity. The spin–1 connection quanta—the *Mition* in the Gaiamiti sector and the *Tantron* in the Tantrakyma sector—arise from the internal operator–bundle connection $\mathcal{A}_\mu$ and acquire BEH masses

$$(M^2_\Phi)_{IJ} = g^2\, g_T\Big(\rho_\Phi(T_I)\bar\Phi,\ \rho_\Phi(T_J)\bar\Phi\Big), \qquad (M^2_\Theta)_{IJ} = g^2\, g_C\Big(\rho(T_I)\bar\Theta,\ \rho(T_J)\bar\Theta\Big), \qquad (13.5)$$

(Propositions 9.257 and 9.347 and Definitions 9.258 and 9.348). On a certified plateau their BRST–physical transverse poles are isolated by spectral–island/Riesz selectors (Definitions 10.6 and 10.7), and TT purity of gravity is ensured by Definition 10.3. These quanta behave as *vector dark matter*; they do not furnish a distinctive modification of the Einstein operator beyond standard stress–energy sourcing, and therefore are not primary candidates for "dark gravity."

Summary. The hierarchy of plausibility for "dark gravity" in this framework is therefore: (i) the coherent τ–kairos background $\bar t_{\mu\nu}$ (most natural and TT–safe; Definitions 9.58, 10.3 and 10.13, eqs. (11.291) and (11.292), and Theorem 9.351); (ii) a coherent Gaiaton (Apeiron) background on the τ–scalar register (Definitions 10.68 and 10.69 and Proposition 10.153); (iii) a nearly isotropic CK background (possible but observationally constrained; eqs. (11.339) to (11.341) and Proposition 13.73); and (iv) mition/tantron, which are vector dark matter rather than dark gravity (Definitions 9.258 and 9.348).

Synthesis. The τ–kairos sector furnishes *both* microscopic (particle) and macroscopic (background) routes into the dark sector, with all channels remaining TT–safe by construction ($\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\hat{\tau}_{\mu\nu} = 0$; Definitions 10.3, 10.4 and 10.13), in line with multimessenger constraints on extra GW polarizations [513]. At the *particle level* there are three vector and one scalar candidates: (i) the spin–1 *Chrono–Kymon* on the τ–vector register (Proposition 10.154), whose mass and dynamics follow from the BEH matrix and relational Proca equations (eqs. (11.339) to (11.341)) and which behaves as CDM on FLRW diagnostic realized reference configuration ($\rho \propto a^{-3}$, $\langle w \rangle \simeq 0$; Proposition 13.73); its portal structure and phenomenology parallel well-studied vector dark matter/dark photon scenarios [599, 548, 507, 210]; (ii) the spin–0 *Gaiaton*/Apeiron on the τ–scalar register (Proposition 10.153), isolated and projected via the weave's Riesz selectors (Definitions 10.68 and 10.69), interpolating between a coherent–oscillation CDM branch (misalignment) and a slow–roll quintessence branch [597, 486, 592, 598]; (iii) the spin–1 *Mition* from the Gaiamiti connection and (iv) the spin–1 *Tantron* from the Tantrakyma connection (Definitions 9.258 and 9.348), whose BEH masses are fixed by the typed covariant split (eq. (6.314) and Propositions 9.257 and 9.347) and whose BRST–physical transverse poles are isolated by spectral–island/Riesz selectors on a certified plateau (Definitions 9.58, 10.6 and 10.7). All four particle candidates are spectrally orthogonal to the Standard Model algebra by the operational criterion of Chapter 12 and remain TT–orthogonal.

Macroscopic route: "dark gravity" from a coherent τ–background. Beyond quanta, the *expectation* $\bar{t}_{\mu\nu}$ of the same τ–kairos operator deforms the diagnostic realized reference configuration chronogeometry and the TT wave operator—without introducing new spin–2 modes—through the typed balance relations (eqs. (11.291) and (11.292)) and the chronogeometric Einstein equations (eq. (11.191)). This "dark gravity" channel is therefore the macroscopic manifestation of the τ–sector already present at the microscopic level (Chrono–Kymon and Gaiaton), rather than an independent addition, and it sits naturally alongside broader modified-gravity/emergent-gravity perspectives [593, 594, 595, 596]. Its TT purity is guaranteed by $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ (Definition 10.3) and is consistent with multimessenger bounds [513]; its execution is diagnostic realized reference configuration–disciplined on a certified plateau $\mathfrak{P}$ with Nielsen/BRST safety (Definition 9.58 and Theorem 9.351).

Adjudication by HSP and operational selection. Which of these realizations is *actually* instantiated is fixed by the *Holosymmetric Selection Principle* (HSP; Section 12.2, Chapter 13, and Theorem 12.42): the law/diagnostic realized reference configuration that maximizes holosymmetric coherence under the constraints of spectral orthogonality to the SM algebra (Chapter 12) and typed/BRST hygiene is selected on $H^0(\hat{Q}_{\mathrm{BRST}})$ (Section 12.3). In this way the τ–kairos provides a unified substrate for the dark sector—yielding, depending on the HSP–selected diagnostic realized reference configuration, (a) vector DM (CK, mition, tantron) [599, 548, 507, 210], (b) scalar DM or DE (Gaiaton) [597, 486, 592, 598], or (c) a coherent background deformation interpretable as "dark gravity" [593, 594, 595, 596]—all within the same projector–certified, TT–pure relational fabric.

A Unified Dark Sector We posit a dark sector that is not an ad–hoc bolt–on but an intrinsic facet of the same fields that constitute the visible universe. On the emergent/GNS arena, the dual–field vacuum of Gaiamiti and Tantrakyma is encoded by the unified inter–modal correlator and realized on a certified plateau via the Σ–map (Definitions 9.5, 9.7, 9.58 and 9.60). The typed weave then cleanly separates the tensorial (TT) graviton channel from the non–TT temporal/coherence registers of the τ–kairos (Definitions 10.3 and 10.4). Within this structure, *Dark Energy* appears as a property of the coupled vacuum: a small residual from the self–regulating Gaiamiti–Tantrakyma balance (Section 13.1.1), with its late–time dynamics governed by the spin–0 Apeiron/Gaiaton on the τ–scalar register (Section 10.8 and Definitions 10.68 and 10.69) and constrained by CMB/BAO/WL probes [581, 136, 151, 152, 153, 154].

Dark Matter in this unified picture arises from projector–certified, BRST–physical excitations that are spectrally orthogonal to the Standard Model algebra (Chapter 12 and Section 12.3). Vector candidates include (i) the spin–1 **Chrono–Kymon** on the τ–vector register (Proposition 10.154), whose BEH mass and relational Proca dynamics are fixed by (eqs. (11.339) to (11.341)) and that behaves as CDM on FLRW diagnostic realized reference configuration (Proposition 13.73); and (ii) the spin–1 **Mition** and **Tantron**, the connection quanta of the Gaiamiti and Tantrakyma sectors, respectively (Definitions 9.258 and 9.348). Their BEH masses follow from the typed covariant split (eq. (6.314) and Propositions 9.257 and 9.347), while spectral–island/Riesz selectors isolate their BRST–physical transverse poles on a diagnostic realized reference configuration (Definitions 9.58, 10.6 and 10.7). A scalar candidate is the **Gaiaton** (Apeiron) in its massive, coherent–oscillation branch, furnishing cold dark matter (Proposition 10.153); in its slow–roll branch the same species sources dark energy, making the scalar register a bridge between the two dark components [597, 486, 592, 598].

At the macroscopic end, the expectation value $\bar{t}_{\mu\nu}$ of the τ–kairos (Definition 10.13) deforms the diagnostic realized reference configuration chronogeometry and the TT wave operator via the typed balance relations (eqs. (11.291) and (11.292)), yielding an effective *"dark gravity"* without new spin–2 modes; TT purity is guaranteed by the background–covariant projector $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ (Definition 10.3) and remains consistent with multimessenger bounds [513]. This macroscopic route is the large–scale manifestation of the same operator content that, microscopically, yields the Chrono–Kymon and Gaiaton—hence not an independent sector, but an alternative interpretation consistent with the typed weave and plateau discipline [593, 594, 595, 596].

Finally, which realization (vector or scalar DM, quintessence–like DE, or a coherent τ–background interpretable as dark gravity) is *actually* instantiated is adjudicated by the *Holosymmetric Selection Principle* (HSP; Section 12.2 and Theorem 12.42). On $H^0(\hat{Q}_{\mathrm{BRST}})$ (Section 12.3), the HSP selects the diagnostic realized reference configuration that maximizes holosymmetric coherence under spectral orthogonality to the Standard Model, ensuring gauge/Nielsen safety (Theorem 9.351) and TT isolation (Definition 10.3). In this sense, the unified dark sector is the operationally selected face of the same relational fabric—constructed by the sector actions and split transport (eqs. (6.314) and (6.315)), realized on a certified plateau (Definitions 9.58 and 9.60), and typed by the weave projectors (Definitions 10.4 and 10.7)—that also carries the visible universe [414, 581, 136].

13.1.3 The Interferometer as the Foundational Principle of Selection

To adjudicate competing physical laws for the dark sector, the IQR framework instantiates the *Holosymmetric Selection Principle* (HSP; Section 12.2) *operationally* via the *Holosymmetric Interferometer Model* (Section 2.4). The interferometer is built on the parasymmetric carrier

$$\mathcal{H}_{\text{para}}^{(n)} = (\mathbb{C}^2)^{\otimes n} \quad (\textit{Definition 2.16}),$$

equipped with the local/global involutive duality operators $D^{(\mathfrak{B}_x)}$ and $\hat{D}_{\text{PQS}}^{(n)}$ (Definitions 2.7 and 2.12) that serve as discrete "knobs'' for branch tagging and refocusing. All selection observables are evaluated on the BRST-physical space

$$\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}}), \quad \text{on a certified plateau } \mathfrak{P} \quad (\textit{Section 12.3and Definition 9.58}),$$

with realization through the Σ–map (Definition 9.60).

Stage I (symmetric): interference of complete histories. Let the two register-wide path laws (the "slits'') be $\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N}$ and $\hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}$ (eqs. (2.69) and (2.70)). The *Controlled Holosymmetric Operator* $\mathbf{U}_{\text{ctrl}}$ inserts these into the control qubit (Definition 2.114), while preparation/readout use Hadamards ($\mathbf{U}_{\text{Had}}$) (Section 3.1.1). For an initial register state $|\Psi_{\text{RR}}\rangle$,

$$\mathcal{O} := \left\langle \Psi_{\text{RR}} \right| \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G \dagger} \, \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} \left| \Psi_{\text{RR}} \right\rangle, \qquad p_{0/1} = \tfrac{1}{4} \left\| \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} \pm \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} \right\|^2, \tag{13.6}$$

and the symmetric visibility is $\mathcal{V}_{\text{sym}} = |p_0 - p_1| = |\operatorname{Re} \mathcal{O}|$. Stage I acts as a *self-consistency* (diagnostic realized reference configuration) filter that rejects hypotheses whose symmetric overlap is incoherent on $\mathfrak{P}$.

Stage II (asymmetric): comparative tomography. With general control gates $U_c^{\text{prep}}, U_c^{\text{read}} \in SU(2)$, the asymmetric holosymmetric operator has control blocks $\mathcal{F}_{ij}^{(\text{asym})}$ that are linear combinations of the path laws (Section 3.1.5):

$$\begin{aligned} \mathcal{F}_{00}^{(\text{asym})} &= \gamma\alpha \, \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} - \delta^* \beta \, \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}, \\ \mathcal{F}_{01}^{(\text{asym})} &= -\gamma\beta^* \, \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} - \delta^* \alpha^* \, \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G}, \end{aligned} \quad \Rightarrow \quad p_i^{(\text{asym})} = \left\| \mathcal{F}_{i0}^{(\text{asym})} |\Psi_{\text{RR}}\rangle \right\|^2, \tag{13.7}$$

and the *asymmetric visibility* $\mathcal{V}_{\text{asym}} := |p_0^{(\text{asym})} - p_1^{(\text{asym})}|$ realizes tomographic, character-weighted selection (Theorem 3.8). Stage II therefore *compares* two candidate universal histories by steering and reading their interference.

Selection functional (HSP in interferometric form). Given two hypotheses H_i, H_j for the dark sector (e.g. coherent τ-background vs. CK/Gaiaton vs. connection vectors), define the pairwise *coherence score*

$$\mathscr{C}(H_i, H_j; \mathfrak{P}) := \sup_{U_c^{\text{prep}}, U_c^{\text{read}} \in SU(2)} \left(\mathcal{V}_{\text{asym}}[H_i, H_j] \right) \Big|_{\mathcal{H}_{\text{phys}}, \mathfrak{P}}, \tag{13.8}$$

subject to *spectral orthogonality to the SM algebra* and diagnostic realized reference configuration hygiene (Chapter 12 and Definition 9.58). The HSP selects the realized law/history by

$$H_\star = \arg\max_{H\in\mathcal{H}_{\text{cand}}} \mathscr{C}\big(H, H_{\text{ref}}; \mathfrak{P}\big), \tag{13.9}$$

with BRST projection and Nielsen safety enforced (Section 12.3 and Theorem 9.351).

Here H_{ref} is a reference hypothesis (e.g. minimal ΛCDM dressing), and the maximization ranges over admissible SU(2) controls and band tags used in realization (Definitions 9.58 and 9.60).

Diagnostic complement: commutator lensing (not the selector). The *commutator lens* compares "superpose→evolve" vs. "evolve→superpose"

$$\mathbf{U}_{\text{holo}}^{(\text{sym-com})} = \Big[\mathbf{U}_{\text{Had}},\, \mathbf{U}_{\text{ctrl}}\Big]\, \mathbf{U}_{\text{Had}}^{\dagger} \quad (\textit{Section 3.3}),$$

and serves as a *diagnostic* of algebraic non-commutativity ("operational curvature"). It quantifies internal twists of a *given* hypothesis but does not, by itself, perform the *selection* that Stage I/II carry out.

Why the interferometer selects the dark sector. Dark-sector hypotheses differ *operationally* in their whole-evolution channels (e.g. presence/absence of a coherent τ–background, CK vs. Gaiaton dynamics, connection-vector portals). The Holosymmetric Interferometer measures the overlap and steerable visibility of these channels on $H^0(\hat{Q}_{\text{BRST}})$ and a certified $\mathfrak{P}$, precisely enacting the HSP's mandate to "maximize holosymmetric coherence under SM-orthogonality" (Sections 12.2 and 12.3 and Definition 9.58). Therefore, to determine the nature of Dark Energy and Dark Matter, one must *compare* the candidate universal histories via (13.6)–(13.9)—the interferometer is the *foundational principle of selection*, while commutator lensing remains its complementary *diagnostic*.

13.1.4 The Holosymmetric Selection of the Dark Sector

Operational anchoring. As in the selection of the Standard Model (SM) gauge law, every admissible dark–sector proposal G is evaluated *operationally* on the BRST–physical space

$$\mathcal{H}_{\text{phys}} := H^0(\hat{Q}_{\text{BRST}}),$$
realized on a certified plateau $\mathfrak{P}$ via the Σ–map
(*Section 12.3and Definitions 9.58and 9.60*).

The evaluation is carried out with the two–stage *Holosymmetric Selection Principle* (HSP; Theorem 12.79) implemented by the *Holosymmetric Interferometer Model* on the parasymmetric carrier

$$\mathcal{H}_{\text{para}}^{(n)} = (\mathbb{C}^2)^{\otimes n}, \quad D^{(\mathfrak{B}_x)},\ \hat{D}_{\text{PQS}}^{(n)} \text{ as discrete control knobs}$$
(*Definitions 2.7, 2.12and 2.16and Section 2.4*).

Stage I (symmetric) tests *physical self–coherence* by the symmetric overlap

$$\mathcal{C}_{\mathrm{phys}}(G \mid \rho) := \Big|\,\mathrm{Re}\;\mathcal{O}_G(\rho)\Big|, \qquad \mathcal{O}_G(\rho) := \Big\langle \Psi_{\mathrm{RR}}\Big|\,\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G(G)\dagger}\,\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N(G)}\,\Big|\Psi_{\mathrm{RR}}\Big\rangle, \tag{13.10}$$

with detection probabilities $p_{0/1}$ as in (13.6) (Section 3.1.1). Hypotheses failing $\mathcal{C}_{\mathrm{phys}}$ on $\mathfrak{P}$ are discarded. *Stage II (asymmetric)* compares survivors A, B by the *physical holosymmetric overlap*

$$\Omega_{\mathrm{phys}}(A, B \mid \mathring{\mathbf{\Psi}}) := \sup_{U_c^{\mathrm{prep}}, U_c^{\mathrm{read}} \in SU(2)} \Big|\, p_0^{(\mathrm{asym})}(A, B) - p_1^{(\mathrm{asym})}(A, B)\,\Big|, \tag{13.11}$$

where $p_i^{(\mathrm{asym})}$ are computed from the asymmetric control blocks $\mathcal{F}_{ij}^{(\mathrm{asym})}$ in (13.7) (Section 3.1.5 and Theorem 3.8). Holosymmetric orthogonality implies $\Omega_{\mathrm{phys}}(A, B \mid \mathring{\mathbf{\Psi}}) = 0$ for inequivalent induced K–actions (Proposition 12.76). The *coherence projector* $\mathcal{P}_G^{\mathrm{coh}}$ (Definition 12.77) certifies visibility through G, while spectral orthogonality to the SM algebra enforces darkness (Chapter 12 and Section 12.3.2). At the emergent/GNS layer, Stage II is probed by cosmological effect–families (SN, CMB, BAO, WL, growth) treated as interferometric effects (Section 2.4).

Selection functional and maximizer (HSP). Writing the *coherence score* as

$$\mathscr{C}(G; \mathfrak{P}) := \sup_{U_c^{\mathrm{prep}}, U_c^{\mathrm{read}} \in SU(2)} \Big(\mathcal{V}_{\mathrm{asym}}[G, G]\Big)\Big|_{\mathcal{H}_{\mathrm{phys}}, \mathfrak{P}}, \qquad \mathcal{V}_{\mathrm{asym}} := \Big|p_0^{(\mathrm{asym})} - p_1^{(\mathrm{asym})}\Big|, \tag{13.12}$$

the realized dark sector is the unique fixed point of *Maximal Coherence*:

$$\begin{aligned} G_\star &= \arg\max_G\ \mathscr{C}(G; \mathfrak{P}), \\ &\Omega_{\mathrm{phys}}(G_\star, B \mid \mathring{\mathbf{\Psi}}) = 0 \quad \forall\, B \neq G_\star, \\ &\text{with Nielsen/BRST safety } (\textit{Theorem 9.351and Section 12.3}). \end{aligned} \tag{13.13}$$

Tau–kairos realization (scalar, vector, and macroscopic branches). The τ–kairos tensor $\hat{\tau}_{\mu\nu} := \delta\hat{t}_{\mu\nu}$ (Definition 10.13) generates *both* microscopic and macroscopic dark–sector branches, all TT–null ($\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\hat{\tau} = 0$; Definitions 10.3 and 10.4). *(i) Scalar branch:* the **Gaiaton**/Apeiron on the τ–scalar register, isolated by the Riesz/typed projectors (Section 10.8 and Definitions 10.68 and 10.69),

$$\mathsf{P}_{\mathrm{spec}}^{(0)} = \sum_{m \in \mathfrak{S}_0} \frac{1}{2\pi i} \oint_{\zeta_m} (K_\tau^{(0)} - z)^{-1} dz, \quad \mathcal{P}_{\mathrm{G},\eta}^{(0)} = \mathsf{P}_{\mathrm{phys}}\, S^{(\eta)} \Big(\Pi_\tau^{(\mathrm{spec};0)} \oplus \Pi^{(\mathrm{vac})} \oplus \Pi^{(\mathrm{corr})}\Big) S^{(\eta)} \mathsf{P}_{\mathrm{phys}}.$$

Its mass arises from the scalar/radial Hessian of $\mathcal{V}_\Phi$ (Proposition 10.153). *(ii) Vector branch:* the **Chrono–Kymon** on the τ–vector register with helicities $A^{(\pm)}$ (Definition 10.4 and eq. (10.7)), whose BEH mass and dynamics are

$$\begin{aligned} (M^2_{\tau\text{-vec}})_{IJ} &= g_\tau^2 \Big\langle \mathcal{R}_\tau^{(S)}(T_I)\varphi^{[S]},\, \mathcal{R}_\tau^{(S)}(T_J)\varphi^{[S]} \Big\rangle, \\ \bar{\nabla}_\mu \mathcal{F}^{I\,\mu\nu} + (M^2_{\tau\text{-vec}})^I{}_J\, \mathcal{B}^{J\,\nu} &= J^{I\,\nu}_{(\tau)} \\ &\qquad (\textit{eqs. (11.339)to (11.341)}). \end{aligned}$$

On FLRW diagnostic realized reference configuration the rapid–oscillation regime yields CDM kinematics ($\rho \propto a^{-3}$, $\langle w \rangle \simeq 0$; Proposition 13.73). *(iii) Macroscopic branch ("dark gravity"):* the expectation $\bar{t}_{\mu\nu}$ deforms the diagnostic realized reference configuration chronogeometry and the TT wave operator through the typed balance

$$\begin{aligned} \mathcal{L}_{hh}[\bar{\Sigma}'] &= \mathcal{L}_{hh}[\bar{\Sigma}] + \varepsilon\, \delta\mathcal{L}_{hh}\Big[\Delta\Sigma^{(\tau)}\Big] + \cdots , \\ \delta\bar{G}_{\mu\nu}\Big[\Delta\Sigma^{(\tau)}\Big] + \Lambda\, \Delta\Sigma^{(\tau)}_{\mu\nu} &= \kappa\, \Delta\bar{T}^{(\tau)}_{\mu\nu} \\ &\qquad (eqs.\ (11.191), (11.291) and\ (11.292)). \end{aligned}$$

yielding an effective "dark gravity'' without new spin–2 modes. These three manifestations are different dynamical branches of the *same* operator content.

Connection–vector extensions (Mition, Tantron). Beyond τ–kairos, the operator–bundle connection $\mathcal{A}_\mu$ supplies *vector* candidates that remain TT–safe and spectrally orthogonal to the SM: the **Mition** (Gaiamiti sector) and the **Tantron** (Tantrakyma sector) (Definitions 9.258 and 9.348). Their BEH mass matrices follow from the typed covariant split (eq. (6.314) and Propositions 9.257 and 9.347):

$$(M^2_\Phi)_{IJ} = g^2\, g_T\Big(\rho_\Phi(T_I)\bar{\Phi},\ \rho_\Phi(T_J)\bar{\Phi}\Big), \qquad (M^2_\Theta)_{IJ} = g^2\, g_C\Big(\rho(T_I)\bar{\Theta},\ \rho(T_J)\bar{\Theta}\Big),$$

and their BRST–physical transverse poles are isolated by spectral–island/Riesz selectors (Definitions 10.6 and 10.7). They behave as vector CDM on $\mathfrak{P}$.

Holosymmetric adjudication. The HSP does *not* select among unrelated add–ons, but among the scalar, vector, and macroscopic branches of τ–kairos together with the connection–vector extensions. By maximizing holosymmetric coherence while enforcing SM orthogonality and diagnostic realized reference configuration hygiene (Definitions 9.58 and 13.1 and Theorem 9.351), the rule (13.13) determines which branch is instantiated. Cosmological histories consistent with admissible effect–families (Chapter 13) then identify: the *Gaiaton* (slow–roll) as dark energy; the *Gaiaton* (coherent oscillations), *Chrono–Kymon*, *Mition*, and *Tantron* as dark matter channels; and a coherent $\bar{t}_{\mu\nu}$ as the residual relational ("dark–gravity'') deformation. Thus the dark sector *emerges holosymmetrically*: it is the operationally selected face of the same typed relational fabric that carries the visible universe.

13.2 Selecting the Nature of Dark Matter

(1) Hypothesis set for Dark Matter selection.

Definition 13.2 (Hypothesis set for Dark Matter selection)**.** Define the candidate family

$$\mathcal{H}^{\mathrm{DM}}_{\mathrm{cand}} := \Big\{H_S,\ H_V,\ H_M,\ H_T\Big\} \cup \Big\{H_{\mathrm{mix}}(\boldsymbol{\pi})\Big\}, \qquad \boldsymbol{\pi} := (\pi_S, \pi_V, \pi_M, \pi_T), \tag{13.14}$$

to be evaluated on the BRST–physical space and a certified plateau $\mathfrak{P}$ (Section 12.3 and Definition 9.58) under the dark–sector criterion (Definition 13.1). The elements are:

(i) H_S **(scalar, Gaiaton/Apeiron)**
the spin–0 τ–scalar excitation (typed scalar slot of the τ–kairos; Proposition 10.153), with mass from the Hessian of $\mathcal{V}_\Phi$ along the typed direction and TT–nullity on the weave (Definition 10.3).

(ii) H_V **(vector, Chrono–Kymon)**
the spin–1 τ–vector excitation (helicity–typed; Proposition 10.154), endowed with a BEH mass matrix from τ–gauging and TT–nullity on the weave (Definition 10.3).

(iii) H_M **(Mition)**
a spin–1 Gaiamiti connection quantum with internal BEH mass on the Φ–leg (typed split and mass generation as in eq. (6.314) and Proposition 9.257; definition Definition 9.258); TT–safe by projector commutations (Definition 10.3 and Proposition 10.9).

(iv) H_T **(Tantron)**
a spin–1 Tantrakyma connection quantum with internal BEH mass on the Θ–leg (typed split and mass generation as in eq. (6.314) and Proposition 9.347; definition Definition 9.348); TT–safe by projector commutations (Definition 10.3 and Proposition 10.9).

(v) $H_{\mathrm{mix}}(\boldsymbol{\pi})$ **(multi–component mixture)**
a convex combination of the single–component histories,

$$H_{\mathrm{mix}}(\boldsymbol{\pi}) \;:=\; \pi_S H_S + \pi_V H_V + \pi_M H_M + \pi_T H_T, \qquad \sum\nolimits_{i\in\{S,V,M,T\}} \pi_i = 1, \quad \pi_i \geq 0, \tag{13.15}$$

used when interferometric evidence favors 1–3 component scenarios.

Each single–component hypothesis is specified by its typed register, mass–generation mechanism, equations of motion, and TT safety on the weave; the multi–component hypothesis allows convex mixtures subject to the above constraints.

($\mathbf{H_S}$) **(Scalar, Gaiaton/Apeiron).** A spin–0 τ–scalar excitation (typed scalar slot of the τ–kairos, Definitions 10.4 and 10.13), isolated by the weave's Riesz/projector machinery (Section 10.8 and Definitions 10.68 and 10.69) and identified with the *Gaiaton*/Apeiron (Proposition 10.153). Its mass is set by the scalar/radial Hessian of the Gaiamiti potential:

$$m_S^2 \;\in\; \mathrm{spec}\left(\Pi_\tau^{[S]} \left[\frac{\delta^2 \mathcal{V}_\Phi}{\delta\hat{\Phi}\,\delta\hat{\Phi}}\Big|_{\bar{\Phi}} \right] \Pi_\tau^{[S]} \right), \tag{13.16}$$

and TT purity of gravity is guaranteed since all τ–excitations are TT–null ($\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\hat{\tau} = 0$; Definition 10.3).

($\mathbf{H_V}$) **(Vector, Chrono–Kymon).** A spin–1 τ–vector excitation with internal $U(1)_C$ and BEH mass (Proca on $(\mathcal{X}_q, \bar{\Sigma})$); carrier/helicities from the typed vector slot $A^{(\pm)}$ (Definition 10.4 and eq. (10.7)). The BEH mass matrix and dynamics are (eqs. (11.339) to (11.341))

$$(M^2_{\tau\text{-vec}})_{IJ} = g_\tau^2 \left\langle \mathcal{R}_\tau^{(S)}(T_I)\varphi^{[S]},\, \mathcal{R}_\tau^{(S)}(T_J)\varphi^{[S]} \right\rangle, \qquad \bar{\nabla}_\mu \mathcal{F}^{I\,\mu\nu} + (M^2_{\tau\text{-vec}})^I{}_J\, \mathcal{B}^{J\,\nu} = J^{I\,\nu}_{(\tau)}, \tag{13.17}$$

with TT–nullity as above and CDM–like late–time behavior on FLRW diagnostic realized reference configuration ($\rho \propto a^{-3}$, $\langle w \rangle \simeq 0$; Propositions 10.154 and 13.73).

($\mathbf{H_M}$) **(Mition).** A spin–1 excitation of the *Gaiamiti* operator–bundle connection $\mathcal{A}_\mu$ (Definition 9.258), with internal BEH mass from the typed covariant split (eq. (6.314) and Proposition 9.257):

$$(M_\Phi^2)_{IJ} = g^2\, g_T\Big(\rho_\Phi(T_I)\bar{\Phi},\ \rho_\Phi(T_J)\bar{\Phi}\Big), \tag{13.18}$$

and BRST–physical transverse poles isolated by spectral–island/Riesz selectors (Definitions 10.6 and 10.7). TT safety follows from weave commutations and the fact that $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ acts on symmetric 2–tensors (Definition 10.3).

($\mathbf{H_T}$) **(Tantron).** A spin–1 excitation of the *Tantrakyma* operator–bundle connection $\mathcal{A}_\mu$ (Definition 9.348), with internal BEH mass (Proposition 9.347)

$$(M_\Theta^2)_{IJ} = g^2\, g_C\Big(\rho(T_I)\bar{\Theta},\ \rho(T_J)\bar{\Theta}\Big), \tag{13.19}$$

and the same TT/BRST/diagnostic realized reference configuration hygiene as for the Mition (Definitions 10.3, 10.6 and 10.7).

Definition 13.3 (Admissible multi–component (mixture) hypothesis)**.** Let $\mathcal{I} = \{S, V, M, T\}$ index the single–component histories H_S, H_V, H_M, H_T (Definition 13.2). An admissible mixture is

$$H_{\mathrm{mix}}(\boldsymbol{\pi}) := \sum_{i\in\mathcal{I}} \pi_i\, H_i, \qquad \boldsymbol{\pi} = (\pi_i)_{i\in\mathcal{I}}, \quad \pi_i \geq 0, \quad \sum_i \pi_i = 1, \tag{13.20}$$

to be evaluated on $\mathcal{H}_{\mathrm{phys}}$ and a certified plateau $\mathfrak{P}$ (Section 12.3 and Definition 9.58). For the interferometer composition of *path laws*, amplitudes add with $\mathbf{v} := \sqrt{\boldsymbol{\pi}}$, i.e. $v_i = \sqrt{\pi_i}$ and $\|\mathbf{v}\|_2 = 1$, so that

$$\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N\left(H_{\mathrm{mix}}\right)} = \sum_{i\in\mathcal{I}} v_i\, \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N(H_i)}, \qquad \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G\left(H_{\mathrm{mix}}\right)} = \sum_{i\in\mathcal{I}} v_i\, \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G(H_i)}, \tag{13.21}$$

as in eq. (13.68). The corresponding Stage–II visibility and mixture score are given by eqs. (13.70) and (13.71), with selection carried out by the penalized program eq. (13.72).

Proposition 13.4 (Extremal selection under holosymmetric orthogonality)**.** *Let $\mathbb{K}(U_c)$ be the Hermitian kernel in eq. (13.70). (a) If the induced K–actions of $H_i \neq H_j$ are holosymmetrically orthogonal on $\mathcal{H}_{\mathrm{phys}}$ (Proposition 12.76), then the off–diagonal entries of $\mathbb{K}(U_c)$ vanish on $\mathfrak{P}$, and the HSP argmax (eq. (13.50)) for H_{mix} is attained at a* vertex *of the simplex, i.e. $\pi_{i_\star} = 1$ for some $i_\star \in \mathcal{I}$. (b) If the off–diagonal entries do not vanish, the maximizing amplitude $\mathbf{v}_\star$ is the principal eigenvector of the control–tuned kernel $\mathbb{K}(U_c^\star)$ (subject to $\mathbf{v} \geq 0$, $\|\mathbf{v}\|_2 = 1$), and a genuine mixture may be selected. Parsimony and diagnostic realized reference configuration–complexity penalties in eq. (13.72) bias toward extremal (few–component) solutions.*

Remark 13.5 (Diagnostic exclusion: coherent τ background is not DM)**.** The coherent τ–background $\bar{t}_{\mu\nu}$ (Definition 10.13) deforms the diagnostic realized reference configuration via the typed balance and chronogeometric equations (eqs. (11.191), (11.291) and (11.292)) and is therefore a macroscopic DE/geometry channel, not a particulate DM component. It is *excluded* from $\mathcal{H}_{\mathrm{cand}}^{\mathrm{DM}}$ (Definition 13.2) and retained solely as a *diagnostic confounder* to be checked for consistency on $\mathfrak{P}$.

13.2.1 Canonical Action Fragments and Notation (Per Hypothesis)

All hypotheses are formulated on the realized chronogeometric background $\left(\mathcal{X}_q, \bar{\Sigma}_{\mu\nu}\right)$ with Relational Levi–Civita connection $\bar{\nabla}$ (Definition 6.7), measure $d\mu(x) := \sqrt{-\det \bar{\Sigma}}\, d^d x$, and the realized weave configuration $\mathcal{I}^{[\mathcal{F}]}$ (Definition 10.1). Typed restrictions use the commuting projector family from Chapters 10–12 ($\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$, triad $\Pi^{(a)}$, τ-type idempotents $S^{(b)}, S^{(L)}, \tilde{S}^{(p)}, A^{(+)}, A^{(-)}$, and Riesz selectors $\mathsf{P}^{(0)}_{\mathrm{spec}}, \mathsf{P}^{(2)}_{\mathrm{spec}}$; see Definitions 9.116, 10.2 to 10.4 and 10.7). BRST/diagnostic realized reference configuration hygiene follows Section 12.3.

($\mathbf{H_S}$) Scalar τ–slot (Gaiaton/Apeiron).

Definition 13.6 (Typed Gaiaton/Apeiron field and action)**.** Let $\hat{\tau}_{\mu\nu} = \delta \hat{t}_{\mu\nu}$ be the τ–kairos tensor (Definition 10.13). The *Gaiaton/Apeiron* scalar ϕ is the τ–scalar extracted on the realized weave configuration by the typed/Riesz chain

$$\phi := \mathsf{P}^{(0)}_{\mathrm{spec}}\, S^{(b)}[\hat{\tau}] \qquad \text{(typed scalar slot; } \textit{Definitions } 10.4 \textit{and } 10.7\text{)}.$$

On the diagnostic realized reference configuration $(\mathcal{X}_q, \bar{\Sigma})$ with $\bar{\nabla}$ and $d\mu = \sqrt{-\det \bar{\Sigma}}\, d^d x$, its quadratic+interaction action reads

$$S_S[\phi] = \int_{\mathcal{X}_q} d\mu \left[\tfrac{1}{2}\, \bar{\Sigma}^{\mu\nu} (\bar{\nabla}_\mu \phi)(\bar{\nabla}_\nu \phi) - \tfrac{1}{2}\, m_S^2\, \phi^2 - V_\Phi^{\mathrm{int}}(\phi) \right], \tag{13.22}$$

with typed mass from the Gaiamiti potential's Hessian

$$m_S^2 := \left(\partial_\phi^2 \mathcal{V}_\Phi \right)\Big|_{\mathrm{vac}} \qquad (\tau\text{–scalar slot; } \textit{Definition } 10.4). \tag{13.23}$$

Proposition 13.7 (Euler–Lagrange equation, TT–nullity, and admissibility)**.** *The field ϕ satisfies*

$$\bar{\Box}_{\bar{\Sigma}} \phi + m_S^2 \phi + V_\Phi'(\phi) = 0, \qquad \bar{\Box}_{\bar{\Sigma}} := \bar{\Sigma}^{\mu\nu} \bar{\nabla}_\mu \bar{\nabla}_\nu. \tag{13.24}$$

Moreover, scalar dressings of the background metric are annihilated by the TT projector (York–Lichnerowicz),

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\left(\phi\, \bar{\Sigma}\right) = 0 \qquad \textit{(eq. (10.4) and Definition 10.3)}, \tag{13.25}$$

so the Gaiaton is TT–safe in the weave sense (Chapter 11). Evaluated on the BRST–physical space and a certified plateau $\mathfrak{P}$, the model is gauge/realization–safe (Section 12.3 and Definition 9.58), and it satisfies the dark–sector projector criterion (Definition 13.1) when used in the interferometric selection (Section 13.2.2).

Proposition 13.8 (Cosmological branches: DE slow–roll vs DM oscillations)**.** *Let w_ϕ and (ϵ_V, η_V) be defined by*

$$w_\phi(a) := \frac{\frac{1}{2}\dot{\phi}^2 - \mathcal{V}_\Phi}{\frac{1}{2}\dot{\phi}^2 + \mathcal{V}_\Phi}, \qquad \epsilon_V := \frac{M_{\mathrm{Pl}}^2}{2}\left(\frac{\mathcal{V}_\Phi'}{\mathcal{V}_\Phi}\right)^2, \quad \eta_V := M_{\mathrm{Pl}}^2 \frac{\mathcal{V}_\Phi''}{\mathcal{V}_\Phi} \quad \textit{(eqs. (13.52) and (13.53))}. \tag{13.26}$$

Then on FLRW diagnostic realized reference configuration:

(DE) *If* $\epsilon_V \ll 1$ *and* $|\eta_V| \ll 1$ *(slow–roll), then* $w_\phi \simeq -1$*, furnishing a dark–energy–like component (eq. (13.54)).*

(DM) *If* $m_S \gg H$ *(post–oscillation era with* $H(a_{\rm osc}) \simeq m_S$*), then* $\langle w_\phi \rangle \simeq 0$ *and* $\rho_\phi \propto a^{-3}$*, furnishing a cold–dark–matter–like component (eq. (13.54)).*

These two branches constitute the decision–useful signatures employed in Stage II of the interferometer and in the HSP argmax (Sections 13.2.2 and 13.2.3 and eq. (13.50)).

Corollary 13.9 (Observables map for the typed Gaiaton branches)**.** *Under the hypotheses of Proposition 13.8:* (DM branch) *If* $m_S \gg H$*, the linear power spectrum factorizes with the scalar coherence kernel* $\mathcal{S}\big(k; k_c^{(S)}, \nu_S\big)$ *and cutoff* $k_c^{(S)} \sim \max_a k_J(a)$ *as in eqs. (13.57) and (13.59). The leading discriminants are small–scale suppression/oscillatory features near* $k \sim k_c^{(S)}$ *propagated to WL shear and RSD/growth, and consistency across BAO/Ly*α *(Sections 2.4 and 13.2.3).* (DE branch) *If* $\epsilon_V, |\eta_V| \ll 1$*, then* $w_\phi \simeq -1$ *and* $\mathcal{S} \to 1$ *on linear scales (negligible small–scale modification), so selection is driven primarily by background probes (SNe, CMB distances, BAO) (Section 13.2.3). These mappings provide the Stage–II meters for the interferometric selection and the HSP argmax (Section 13.2.2 and eq. (13.50)).*

$(\mathbf{H_V})$ Vector τ–slot (Chrono–Kymon).

Definition 13.10 (Typed Chrono–Kymon field and Proca/BEH action)**.** Let $\hat{\tau}_{\mu\nu} = \delta \hat{t}_{\mu\nu}$ denote the τ–kairos tensor (Definition 10.13). The *Chrono–Kymon* C_μ is the τ–vector extracted by the helicity–typed idempotents

$$C_\mu := \Big(A^{(+)} \oplus A^{(-)}\Big)_\mu{}^{\alpha\beta}\, \hat{\tau}_{\alpha\beta} \qquad (Definition\ 10.4 and\ eq.\ (10.7)),$$

defined on the diagnostic realized reference configuration $(\mathcal{X}_q, \bar{\Sigma})$ with $\bar{\nabla}$ and measure $d\mu = \sqrt{-\det \bar{\Sigma}}\, d^d x$. Its broken $U(1)_C$ (Proca/BEH) action is

$$S_V[C] = \int_{\mathcal{X}_q} d\mu \left[-\tfrac{1}{4} F_{\mu\nu} F^{\mu\nu} + \tfrac{1}{2} C_\mu \, (M^2_{\tau\text{-vec}})^\mu{}_\nu \, C^\nu \right], \qquad F_{\mu\nu} := \bar{\nabla}_\mu C_\nu - \bar{\nabla}_\nu C_\mu, \tag{13.27}$$

where the BEH mass matrix arises from the τ–Higgs/radial operator, typed to the vector slot,

$$(M^2_{\tau\text{-vec}})^\mu{}_\nu = \left\langle \frac{\partial^2 \mathcal{V}_\Theta}{\partial a_\mu \, \partial a^\nu} \right\rangle \qquad (\tau\text{–vector slot; } Definition\ 10.4 and\ Chapter\ 12). \tag{13.28}$$

Proposition 13.11 (Field equations, transversality, TT–nullity, admissibility)**.** *Variation of (13.27) yields the Proca equations*

$$\bar{\nabla}_\mu F^{\mu\nu} + (M^2_{\tau\text{-vec}})^\nu{}_\rho C^\rho = 0, \qquad \bar{\nabla}_\nu C^\nu = 0 \quad (on\text{–}shell\ transversality). \tag{13.29}$$

On the weave, τ–typed vectors are TT–orthogonal; in particular,

$$\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\Big(\bar{\nabla}_{(\mu} C_{\nu)}\Big) = 0 \qquad (Definition\ 10.3\ and\ TT\text{–}purity\ ledger,\ Ch.\ 11), \tag{13.30}$$

so the Chrono–Kymon does not source radiative spin–2 on the realized weave configuration. Admissibility is secured on the BRST–physical space and a certified plateau $\mathfrak{P}$ *(Section 12.3 and Definition 9.58) and satisfies the dark–sector projector criterion (Definition 13.1); any residual mixings with visible meters must be projectively null (eq. (13.61)).*

Proposition 13.12 (Cosmological behaviour and anisotropic–stress constraints)**.** *On FLRW diagnostic realized reference configuration, rapidly oscillating Chrono–Kymon solutions behave as cold matter with* $\rho_C \propto a^{-3}$ *and* $\langle w_C \rangle \simeq 0$ *(Proposition 13.73). The averaged anisotropic stress obeys the isotropy constraint*

$$\left\langle \pi^i{}_j[C_\mu] \right\rangle_{\mathrm{osc}} \simeq 0 \qquad (\textit{FLRW; eq. (13.60)}), \tag{13.31}$$

and violations of (13.31) *are grounds for exclusion at Stage I/II (Section 13.2.2).*

Corollary 13.13 (Observables map for the Chrono–Kymon)**.** *In linear theory on* $\mathfrak{P}$*, the matter power spectrum factors as*

$$P_{\mathrm{lin}}(k, z; H_V) = A_s \left(\frac{k}{k_\star}\right)^{n_s - 1} T^2_{\mathrm{vis}}(k)\, \mathcal{S}^2\!\left(k; k_c^{(V)}, \nu_V\right) D^2(z), \qquad (\textit{eq. (13.57)}), \tag{13.32}$$

with a vector–induced coherence kernel $\mathcal{S}$ *and a hypothesis–dependent cutoff*

$$k_c^{(V)}(a) \sim a\, M_{\tau\text{-}vec}\, \mathfrak{m}_{\mathrm{prod}}(H_V) \qquad (\textit{production and coherence history } \mathfrak{m}_{\mathrm{prod}};\ \textit{eq. (13.58)}). \tag{13.33}$$

Decision–useful signatures comprise (i) small–scale damping/oscillatory features near $k \sim k_c^{(V)}$ *(distinct from the scalar case eq. (13.59)), (ii) constraints from the averaged anisotropic stress* (13.31)*, and (iii) portal/orthogonality bounds (eqs. (13.64) and (13.65)). These map to effect families used as interferometer meters (SNe, CMB, BAO, WL, RSD; optional Ly*α*, shear) (Sections 2.4 and 13.2.3).*

$(\mathbf{H_M})$ and $(\mathbf{H_T})$ Connection quanta (Mition, Tantron).

Definition 13.14 (Typed connection–vector quanta: Mition and Tantron)**.** Let $\mathcal{M}_\mu$ and $\mathcal{T}_\mu$ denote, respectively, a Gaiamiti–connection quantum (*Mition*; Definition 9.258) and a Tantrakyma–connection quantum (*Tantron*; Definition 9.348), each living in dark internal factors that are spectrally orthogonal to SM observables (Definition 13.1). On the diagnostic realized reference configuration $(\mathcal{X}_q, \bar{\Sigma})$ with relational connection $\bar{\nabla}$ and measure $d\mu = \sqrt{-\det \bar{\Sigma}}\, d^d x$, their canonical quadratic+mass pieces are

$$S_M[\mathcal{M}] = \int_{\mathcal{X}_q} d\mu \left[-\tfrac{1}{4}\, \mathcal{F}^{(\Phi)}_{\mu\nu} \mathcal{F}^{(\Phi)\,\mu\nu} + \tfrac{1}{2}\, \mathcal{M}_\mu\, (M^2_\Phi)^\mu{}_\nu\, \mathcal{M}^\nu \right], \tag{13.34}$$

$$S_T[\mathcal{T}] = \int_{\mathcal{X}_q} d\mu \left[-\tfrac{1}{4}\, \mathcal{F}^{(\Theta)}_{\mu\nu} \mathcal{F}^{(\Theta)\,\mu\nu} + \tfrac{1}{2}\, \mathcal{T}_\mu\, (M^2_\Theta)^\mu{}_\nu\, \mathcal{T}^\nu \right], \tag{13.35}$$

with curvatures

$$\mathcal{F}^{(\Phi)}_{\mu\nu} := \partial_\mu \mathcal{M}_\nu - \partial_\nu \mathcal{M}_\mu + i[\mathcal{M}_\mu, \mathcal{M}_\nu], \qquad \mathcal{F}^{(\Theta)}_{\mu\nu} := \partial_\mu \mathcal{T}_\nu - \partial_\nu \mathcal{T}_\mu + i[\mathcal{T}_\mu, \mathcal{T}_\nu].$$

The BEH mass matrices $(M^2_\Phi)^\mu{}_\nu$ and $(M^2_\Theta)^\mu{}_\nu$ arise from the respective potentials $\mathcal{V}_\Phi, \mathcal{V}_\Theta$ in their internal sectors via the typed covariant split and mass–generation mechanisms (eq. (6.314) and Propositions 9.257 and 9.347); typed selectors fix the slots (Definitions 10.4 and 10.7).

Proposition 13.15 (Field equations, transversality, TT–safety, and admissibility)**.** *Variation of* (13.34)–(13.35) *yields the Proca–type equations*

$$\bar{\nabla}_\mu \mathcal{F}^{(\Phi)\,\mu\nu} + (M_\Phi^2)^\nu{}_\rho\, \mathcal{M}^\rho = \mathcal{J}^{(\Phi)\,\nu}, \qquad \bar{\nabla}_\nu \mathcal{M}^\nu = 0 \quad \textit{(on–shell, unitary gauge)}, \tag{13.36}$$

$$\bar{\nabla}_\mu \mathcal{F}^{(\Theta)\,\mu\nu} + (M_\Theta^2)^\nu{}_\rho\, \mathcal{T}^\rho = \mathcal{J}^{(\Theta)\,\nu}, \qquad \bar{\nabla}_\nu \mathcal{T}^\nu = 0 \quad \textit{(on–shell, unitary gauge)}. \tag{13.37}$$

Internal transport on the geometry leg obeys the TT–purity condition of a rig–compatible connection,

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\, \mathrm{ad}_{\mathcal{A}_\mu}\, \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]} = 0 \qquad (\textit{Definitions 10.3 and 12.1}), \tag{13.38}$$

hence Mition/Tantron do not source radiative spin–2 on the realized weave configuration. Admissibility is secured on the BRST–physical space and a certified plateau $\mathfrak{P}$ *(Section 12.3 and Definition 9.58), and the dark–sector projector criterion enforces spectral orthogonality to SM meters (Definition 13.1); any kinetic/mass portal must be projectively null on the visible screen (eqs.* (13.61)*,* (13.64) *and* (13.65)*).*

Proposition 13.16 (Cosmological behaviour and anisotropy constraints)**.** *On FLRW diagnostic realized reference configuration and in the rapid–oscillation regime, both connection vectors behave as cold matter with*

$$\rho_{\mathcal{M}},\ \rho_{\mathcal{T}}\ \propto\ a^{-3}, \qquad \langle w_{\mathcal{M}} \rangle \simeq \langle w_{\mathcal{T}} \rangle \simeq 0,$$

while their averaged anisotropic stresses decay to negligible levels at late times:

$$\Big\langle \pi^i{}_j[\mathcal{M}_\mu] \Big\rangle_{\text{late}} \to 0, \qquad \Big\langle \pi^i{}_j[\mathcal{T}_\mu] \Big\rangle_{\text{late}} \to 0 \qquad (\textit{eq.}\ (13.60)). \tag{13.39}$$

Failure of (13.39) *or violation of* (13.38) *is grounds for exclusion at Stage I/II (Section 13.2.2).*

Corollary 13.17 (Observables map for Mition and Tantron)**.** *In linear theory on* $\mathfrak{P}$*, the power spectrum admits the coherence–kernel factorization (eq.* (13.57)*)*

$$P_{\mathrm{lin}}(k, z; H_M/H_T) = A_s \left(\frac{k}{k_\star}\right)^{n_s - 1} T_{\mathrm{vis}}^2(k)\, \mathcal{S}^2\Big(k; k_c^{(M/T)}, \nu_{M/T}\Big)\, D^2(z), \tag{13.40}$$

with hypothesis–dependent cutoffs analogous to the vector case (eq. (13.58)*):*

$$k_c^{(M)}(a) \sim a\, M_\Phi\, \mathfrak{m}_{\mathrm{prod}}(H_M), \qquad k_c^{(T)}(a) \sim a\, M_\Theta\, \mathfrak{m}_{\mathrm{prod}}(H_T), \tag{13.41}$$

where $\mathfrak{m}_{\mathrm{prod}}$ *encodes production and initial coherence. Decision–useful signatures include: (i) small–scale damping/oscillatory features near* $k \sim k_c^{(M/T)}$ *distinguishing them from the scalar cutoff* $k_c^{(S)}$ *(eq.* (13.59)*); (ii) late–time anisotropy decay* (13.39)*; and (iii) portal/orthogonality bounds (eqs.* (13.61)*,* (13.64) *and* (13.65)*). These map to the effect–families used as interferometer meters (SNe, CMB, BAO, WL, RSD; optional Ly*α*, shear) (Sections 2.4 and 13.2.3).*

Constitutive/typed citations and hygiene (all H).

Assumption 13.18 (Constitutive typing and diagnostic realized reference configuration hygiene (all H))**.** *Every action block (13.22)–(13.35) is understood under the following standing assumptions:*

(T1) Typed insertion by a commuting projector family
All typed projectors commute pairwise and with the physical projector $\mathsf{P}_{\mathrm{phys}}$*:*

$$\Big[\Pi_{\bar\Sigma}^{[\mathrm{TT}]},\Pi^{(a)}\Big]=\Big[\Pi_{\bar\Sigma}^{[\mathrm{TT}]},S^{(b)}\Big]=\Big[\Pi_{\bar\Sigma}^{[\mathrm{TT}]},S^{(L)}\Big]=\Big[\Pi_{\bar\Sigma}^{[\mathrm{TT}]},\tilde S^{(p)}\Big]=\Big[\Pi_{\bar\Sigma}^{[\mathrm{TT}]},A^{(\pm)}\Big]=0,$$
$$\Big[\mathsf{P}_{\mathrm{spec}}^{(r)},\mathsf{P}_{\mathrm{phys}}\Big]=0, \tag{13.42}$$

with $r\in\{0,2\}$ *and all idempotents as in Definitions 9.116, 10.2 to 10.4 and 10.7. Typed/Riesz selectors act before taking BRST cohomology, and their images are stable under the diagnostic realized reference configuration realization.*

(T2) BRST/diagnostic realized reference configuration cleanliness on a realized weave configuration
All fields/operators are realized on the realized weave configuration $\mathcal{I}^{[\mathcal{J}]}$ *and projected to the BRST–physical space*

$$\mathcal{H}_{\mathrm{phys}}:=H^0(\hat Q_{\mathrm{BRST}}),$$
with realization on a certified plateau $\mathfrak{P}$
(Definitions 9.58, 9.60and 10.1and Section 12.3),

such that induced evolutions preserve $\mathcal{H}_{\mathrm{phys}}$ *and satisfy Nielsen/diagnostic realized reference configuration stability (Theorem 9.351).*

(T3) Cone–preserving principal symbols (microcausality)
The principal symbols of the kinetic operators are $\bar\Sigma$*–hyperbolic and cone–preserving, so that retarded/advanced fundamental solutions have* $\bar\Sigma$*–causal supports (microcausality on the realized weave configuration):*

$$\begin{aligned}&\mathrm{supp}\,[\hat{\mathcal{O}}_H(x),\hat{\mathcal{O}}_H(y)]\subseteq J_{\bar\Sigma}(x)\ \cup\ J_{\bar\Sigma}(y),\\ &d\mu=\sqrt{-\det\bar\Sigma}\,d^dx,\\ &\bar\nabla\ \textit{as in Definition 6.7.}\end{aligned} \tag{13.43}$$

Proposition 13.19 (Admissibility for the HSP/Interferometer pipeline)**.** *Under Assumption 13.18, the symmetric and asymmetric interferometric functionals are well–defined and bounded on* $\mathcal{H}_{\mathrm{phys}}$ *for every admissible* H*:*

$$\mathcal{V}_{\mathrm{sym}}(H)\le 1,\qquad \mathcal{V}_{\mathrm{asym}}[H_i,H_j]\le 1,\qquad \mathscr{C}(H;\mathfrak{P}):=\sup_{U_c^{\mathrm{prep}},U_c^{\mathrm{read}}\in SU(2)}\mathcal{V}_{\mathrm{asym}}[H,H]\in[0,1], \tag{13.44}$$

with $\mathcal{V}_{\mathrm{sym}},\mathcal{V}_{\mathrm{asym}},\mathscr{C}$ *as in eqs. (13.47) to (13.49). Moreover, the HSP argmax is gauge/realization–safe on the certified plateau:*

$$H_\star=\arg\max_H\ \mathscr{C}(H;\mathfrak{P}),\qquad \textit{and}\qquad \Delta_{\mathrm{DRRC}}(H)\ \textit{and}\ \mathcal{R}_{\mathrm{Nielsen}}(H)\ \textit{satisfy eq. (13.75).} \tag{13.45}$$

Here $\Delta_{\mathrm{DRRC}}(H)$ denotes the DRRC spread functional, with "DRRC" referring to the diagnostic realized reference configuration of Definition 9.185. Hence every block (13.22)–(13.35) is admissible for the HSP/Interferometer workflow (Section 2.4 and Theorem 12.79).

Remark 13.20 (TT–purity and SM–darkness consistency)**.** Because of (13.42), TT–purity is preserved by all typed insertions and by BRST projection, so spin–2 contamination is excluded by construction (Definition 10.3.) Concurrently, SM–darkness is enforced by the spectral criterion and visible–screen projectors (Definition 13.1 and eq. (13.61)), ensuring that interferometric visibilities arise from *dark* channels only.

13.2.2 Interferometer-Based Selection (HSP) with Explicit Controls

Definition 13.21 (BRST–physical diagnostic realized reference configuration and parasymmetric carrier)**.** All dark–sector hypotheses $H \in \mathcal{H}^{\mathrm{DM}}_{\mathrm{cand}}$ are tested on the BRST–physical space

$$\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}}),$$

realized on a certified plateau $\mathfrak{P}$ via the Σ–map (Section 12.3 and Definitions 9.58 and 9.60). The interferometer acts on the parasymmetric carrier

$$\mathcal{H}^{(n)}_{\mathrm{para}} = (\mathbb{C}^2)^{\otimes n} \qquad (Definition\ 2.16 and\ Section\ 2.4),$$

with discrete controls furnished by the local/global involutions $D^{(\mathfrak{B}_x)}$ and $\hat{D}^{(n)}_{\mathrm{PQS}}$ (Definitions 2.7 and 2.12).

Definition 13.22 (Path laws and controlled insertion)**.** For a hypothesis H, let the two register–wide path laws ("slits") be

$$\hat{\mathbf{U}}^{A_G B_N(H)}_{\mathrm{RR}}, \qquad \hat{\mathbf{U}}^{A_N B_G(H)}_{\mathrm{RR}} \quad (eqs.\ (2.69) and\ (2.70)),$$

inserted by the *Controlled Holosymmetric Operator* $\mathbf{U}_{\mathrm{ctrl}}$ (Definition 2.114). Symmetric preparation/readout uses the Hadamard $\mathbf{U}_{\mathrm{Had}}$; general asymmetric controls employ $U^{\mathrm{prep}}_c, U^{\mathrm{read}}_c \in SU(2)$ (Sections 3.1.1 and 3.1.5).

Lemma 13.23 (Stage I: symmetric self–coherence)**.** *For an admissible register state $|\Psi_{\mathrm{RR}}\rangle$ representing ρ on $\mathcal{H}_{\mathrm{phys}}$, the symmetric overlap and self–coherence are*

$$\mathcal{O}_H(\rho) := \left\langle \Psi_{\mathrm{RR}} \right| \hat{\mathbf{U}}^{A_N B_G(H)\dagger}_{\mathrm{RR}} \hat{\mathbf{U}}^{A_G B_N(H)}_{\mathrm{RR}} \left| \Psi_{\mathrm{RR}} \right\rangle, \qquad \mathcal{C}_{\mathrm{phys}}(H \mid \rho) := \left| \mathrm{Re}\ \mathcal{O}_H(\rho) \right|. \tag{13.46}$$

With symmetric preparation/readout,

$$p_{0/1}(H) = \tfrac{1}{4} \left\| \hat{\mathbf{U}}^{A_G B_N(H)}_{\mathrm{RR}} \pm \hat{\mathbf{U}}^{A_N B_G(H)}_{\mathrm{RR}} \right\|^2, \qquad \mathcal{V}_{\mathrm{sym}}(H) = |p_0(H) - p_1(H)| = |\,\mathrm{Re}\ \mathcal{O}_H|. \tag{13.47}$$

Hypotheses failing stability/unitarity/Nielsen tests on $\mathfrak{P}$ are rejected (Theorem 9.351 and Definition 9.58).

Proposition 13.24 (Stage II: asymmetric tomography and visibility)**.** *For $U_c^{\mathrm{prep}}, U_c^{\mathrm{read}} \in SU(2)$, let $\mathcal{F}_{ij}^{(\mathrm{asym})}$ be the control blocks (linear combinations of the two path laws; Section 3.1.5). For a pair (H_i, H_j),*

$$p_k^{(\mathrm{asym})}(H_i, H_j) = \left\| \mathcal{F}_{k0}^{(\mathrm{asym})}(H_i, H_j)\, |\Psi_{\mathrm{RR}}\rangle \right\|^2, \qquad \mathcal{V}_{\mathrm{asym}}[H_i, H_j] := \left| p_0^{(\mathrm{asym})} - p_1^{(\mathrm{asym})} \right|. \tag{13.48}$$

Define the coherence score of H as

$$\mathscr{C}(H; \mathfrak{P}) := \sup_{U_c^{\mathrm{prep}}, U_c^{\mathrm{read}} \in SU(2)} \mathcal{V}_{\mathrm{asym}}[H, H]|_{\mathcal{H}_{\mathrm{phys}}, \mathfrak{P}}\,. \tag{13.49}$$

Theorem 13.25 (HSP argmax with orthogonality uniqueness)**.** *Holosymmetric orthogonality on $\mathcal{H}_{\mathrm{phys}}$ implies vanishing cross–overlaps for inequivalent induced K–actions (Proposition 12.76). Hence the realized dark–sector history is*

$$H_\star = \arg \max_{H \in \mathcal{H}_{\mathrm{cand}}^{\mathrm{DM}}} \mathscr{C}(H; \mathfrak{P}), \tag{13.50}$$

subject to SM–orthogonality and diagnostic realized reference configuration hygiene (Chapter 12 and Definitions 9.58 and 13.1).

Remark 13.26 (Controls and meters)**.** Discrete controls: $D^{(\mathfrak{B}_x)}$ and $\hat{D}_{\mathrm{PQS}}^{(n)}$ (Definitions 2.7 and 2.12). Readout: Thesis/Antithesis Born valuations (P_T, P_A) as the interferometer screen (Definition 2.1). At the emergent/GNS layer, cosmological effect–families (SNe, CMB, BAO, WL, RSD; optional Lyα and shear) are admitted as interferometric meters implementing (13.48)–(13.49) (Sections 2.4 and 13.2.3).

Stage I (symmetric): self–coherence on $\mathfrak{P}$.

Definition 13.27 (Stage I acceptance set and rejection rule)**.** Fix an admissible state ρ on $\mathcal{H}_{\mathrm{phys}}$ and a certified plateau $\mathfrak{P}$ (Section 12.3 and Definition 9.58). Using the symmetric self–coherence $\mathcal{C}_{\mathrm{phys}}(H \mid \rho)$ and the symmetric visibility $\mathcal{V}_{\mathrm{sym}}(H)$ as defined in (13.46)–(13.47), the *Stage I acceptance set* is

$$\begin{aligned} \mathfrak{A}_I(\rho; \mathfrak{P}) := \Big\{ H \in \mathcal{H}_{\mathrm{cand}}^{\mathrm{DM}} : \mathcal{C}_{\mathrm{phys}}(H \mid \rho) \ \geq\ \tau_I, \\ H \text{ is unitary/stable on } \mathfrak{P}, \\ \text{and } \ \mathcal{R}_{\mathrm{Nielsen}}(H) \leq \epsilon_{\mathrm{N}} \Big\}, \end{aligned} \tag{13.51}$$

where $\tau_I \in (0, 1]$ is a diagnostic realized reference configuration threshold, and $\mathcal{R}_{\mathrm{Nielsen}}$ is the normalized diagnostic realized reference configuration–stability ratio from eq. (13.75). The *Stage I rejection rule* declares $H \notin \mathfrak{A}_I(\rho; \mathfrak{P})$ whenever any of the conditions in (13.51) fails (e.g., violation of unitarity/stability or $\mathcal{R}_{\mathrm{Nielsen}} > \epsilon_{\mathrm{N}}$; cf. Theorem 9.351).

Stage II (asymmetric): comparative tomography and coherence score. F

Definition 13.28 (Stage II instruments and control domain)**.** Stage II operates with preparation/readout controls $U_c^{\mathrm{prep}}, U_c^{\mathrm{read}} \in SU(2)$ on the parasymmetric carrier, and uses the asymmetric control blocks $\mathcal{F}_{ij}^{(\mathrm{asym})}$ obtained as explicit linear combinations of the two path laws of H (Section 3.1.5 and Definition 13.22). All quantities are evaluated on $\mathcal{H}_{\mathrm{phys}}$ realized on the certified plateau $\mathfrak{P}$ (Definition 13.21).

Definition 13.29 (Stage II functionals (reference form))**.** For a pair of hypotheses (H_i, H_j), the outcome probabilities $p_k^{(\mathrm{asym})}(H_i, H_j)$ and the asymmetric visibility $\mathcal{V}_{\mathrm{asym}}[H_i, H_j]$ are *as defined* in (13.48). The single–hypothesis *coherence score* $\mathscr{C}(H;\mathfrak{P})$ is *as defined* in (13.49). No additional structure is introduced here; these objects are used verbatim.

Proposition 13.30 (Pairwise discriminability and selector uniqueness)**.** *If the induced K–actions of $H_i \neq H_j$ are holosymmetrically orthogonal on $\mathcal{H}_{\mathrm{phys}}$ (Proposition 12.76), then the cross–overlap vanishes, and the Stage II functionals in Definition 13.29 separate the pair. Consequently, the realized dark–sector history is uniquely determined by the HSP argmax rule (13.50), subject to SM–orthogonality and diagnostic realized reference configuration hygiene (Chapter 12 and Definitions 9.58 and 13.1).*

Remark 13.31 (Controls and meters (explicit references))**.** Discrete controls: local/global involutions $D^{(\mathfrak{B}_x)}$, $\hat{D}^{(n)}_{\mathrm{PQS}}$ (Definitions 2.7 and 2.12). Readout: Thesis/Antithesis Born valuations (P_T, P_A) (Definition 2.1). At the emergent/GNS layer, cosmological effect–families (SNe, CMB, BAO, WL, RSD; optional Lyα, shear) serve as interferometric meters for (13.48)–(13.49) (Sections 2.4 and 13.2.3).

13.2.3 Distinguishing Signatures for Selection

The interferometric selection (Stages I/II; Section 13.2.2) requires model–specific, typed diagnostics that map hypothesis parameters to observables on a certified plateau $\mathfrak{P}$ (with BRST/Nielsen safety; Section 12.3, Definition 9.58, and Theorem 9.351). We collect here the decision–useful signatures and the equations we will evaluate for each hypothesis H.

Background expansion and DE–DM disentanglement (scalar branch). For the τ–scalar (Gaiaton/Apeiron) ϕ with potential $\mathcal{V}_\Phi(\phi)$ (eq. (13.22)), define the slow–roll parameters

$$\epsilon_V := \frac{M_{\mathrm{Pl}}^2}{2}\left(\frac{\mathcal{V}_\Phi'}{\mathcal{V}_\Phi}\right)^2, \qquad \eta_V := M_{\mathrm{Pl}}^2 \frac{\mathcal{V}_\Phi''}{\mathcal{V}_\Phi}. \tag{13.52}$$

The equation–of–state parameter

$$w_\phi(a) := \frac{\frac{1}{2}\dot{\phi}^2 - \mathcal{V}_\Phi}{\frac{1}{2}\dot{\phi}^2 + \mathcal{V}_\Phi} \tag{13.53}$$

distinguishes the branches:

$$\begin{aligned} \text{DE (slow–roll):} \quad & \{\epsilon_V, |\eta_V|\} \ll 1 \;\Rightarrow\; w_\phi \simeq -1, \\ \text{Scalar DM (oscillatory):} \quad & m_S \gg H \;\Rightarrow\; \langle w_\phi \rangle \simeq 0,\ \rho_\phi \propto a^{-3}, \end{aligned} \tag{13.54}$$

with the onset approximately at $H(a_{\rm osc}) \simeq m_S$ (eqs. (13.23) and (13.24)). We will search parameter regions where $w_\phi \simeq 0$ (DM–like) is cleanly separated from $w_\phi \simeq -1$ (DE–like), ensuring no branch confusion in Stage II.

Definition 13.32 (Gaiaton branch classifier on the diagnostic realized reference configuration)**.** Let ϕ be the τ–scalar (Gaiaton/Apeiron) with dynamics and diagnostics given in (13.22)–(13.24), (13.52), (13.53), and the branch criterion (13.54). For diagnostic realized reference configuration tolerances $\boldsymbol{\tau}_{\rm DE} = (\tau_\epsilon, \tau_\eta, \tau_w)$, $\boldsymbol{\tau}_{\rm DM} = (\kappa, \bar\tau_w)$ with $\tau_\epsilon, \tau_\eta, \tau_w, \bar\tau_w > 0$ and $\kappa > 1$, define the acceptance sets

$$\mathcal{U}_{\rm DE}(\boldsymbol{\tau}_{\rm DE}) := \Big\{ \text{(params, ICs)} \ : \epsilon_V \le \tau_\epsilon, \quad |\eta_V| \le \tau_\eta, \quad |w_\phi + 1| \le \tau_w \Big\}, \tag{13.55}$$

$$\mathcal{U}_{\rm DM}(\boldsymbol{\tau}_{\rm DM}) := \Big\{ \text{(params, ICs)} \ : m_S/H \ge \kappa, \quad |\langle w_\phi \rangle| \le \bar\tau_w, \quad \text{with onset at } H(a_{\rm osc}) \simeq m_S \text{ as in (13.54)} \Big\}. \tag{13.56}$$

Here "params, ICs" denotes the parameter set and initial conditions evaluated on the certified plateau $\mathfrak{P}$.

Lemma 13.33 (Clean separation and identifiability)**.** *If $\boldsymbol{\tau}_{\rm DE}$ and $\boldsymbol{\tau}_{\rm DM}$ are chosen so that $\tau_\epsilon, \tau_\eta, \tau_w, \bar\tau_w$ are sufficiently small and $\kappa > 1$ is fixed, then on any FLRW diagnostic realized reference configuration interval where the diagnostics are valid, the sets $\mathcal{U}_{\rm DE}(\boldsymbol{\tau}_{\rm DE})$ and $\mathcal{U}_{\rm DM}(\boldsymbol{\tau}_{\rm DM})$ are disjoint. Consequently, the scalar DE slow–roll branch and the scalar DM oscillatory branch are* identifiable *in Stage II without ambiguity, using the already defined visibilities and coherence score (Definition 13.29; cf. (13.49)).*

Remark 13.34 (Operational use and observables)**.** In practice, Definition 13.32 supplies the gate for selecting which scalar branch contributes to the Stage II meters. The DM branch feeds the linear power through the scalar coherence kernel with cutoff $k_c^{(S)}$ (eqs. (13.57) and (13.59)), while the DE branch is constrained primarily by background probes via w_ϕ (eq. (13.53), Corollary 13.9). These branch tags then enter the interferometer's asymmetric tomography exactly as referenced in Definition 13.29 and the HSP argmax (13.50), avoiding duplication of formalisms.

Linear power and small–scale structure. For any admissible H, the effective linear power spectrum on $\mathfrak{P}$ is modeled as

$$P_{\rm lin}(k, z; H) = A_s \left(\frac{k}{k_\star}\right)^{n_s - 1} T_{\rm vis}^2(k)\, \mathcal{S}^2\Big(k;\, k_c(H), \nu(H)\Big)\, D^2(z), \tag{13.57}$$

where $T_{\rm vis}(k)$ encodes visible–sector transfer, $D(z)$ is the growth factor on $\bar\Sigma$, and $\mathcal{S}(k; k_c, \nu)$ is the *coherence kernel* that captures dark–sector–induced damping/oscillations (Section 2.4; cf. the usage in Section 13.2.1). The *cutoff scale* k_c is hypothesis–dependent:

Vectors (H_V, H_M, H_T)**.** For a massive vector with mass $m_V \in \{M_{\tau\text{-vec}}, M_\Phi, M_\Theta\}$, comoving coherence sets a characteristic cutoff

$$k_c^{(V)}(a) \;\sim\; a\, m_V\, \mathfrak{m}_{\text{prod}}(H), \qquad \mathfrak{m}_{\text{prod}}(H) \text{ encodes production history and initial coherence,} \tag{13.58}$$

leading to $\mathcal{S}(k; k_c^{(V)}, \nu_V)$ with model–specific oscillatory tails distinct from the scalar case. In the rapid–oscillation regime on FLRW, $\rho_V \propto a^{-3}$ and $\langle w_V \rangle \simeq 0$ (Proposition 13.73 for H_V).

Scalar (H_S)**.** Misalignment–driven oscillations behave CDM–like on large scales, but introduce an effective *Jeans* cutoff at

$$k_J(a) \;\simeq\; a\sqrt{m_S\, H(a)}\,, \qquad \Rightarrow \quad \mathcal{S}(k; k_c^{(S)}, \nu_S) \text{ with } k_c^{(S)} \sim \max_a k_J(a). \tag{13.59}$$

Definition 13.35 (Coherence–kernel factorization on the diagnostic realized reference configuration)**.** For any admissible hypothesis H, the effective linear power on the certified plateau $\mathfrak{P}$ factorizes *as defined* in (13.57), with visible–sector transfer $T_{\text{vis}}(k)$, growth factor $D(z)$ on $(\mathcal{X}_q, \bar{\Sigma})$, and a dark–sector *coherence kernel* $\mathcal{S}\big(k; k_c(H), \nu(H)\big)$ that captures damping/oscillations induced by H (Sections 2.4 and 13.2.1).

Definition 13.36 (Hypothesis–dependent cutoff families)**.** The cutoff scale $k_c(H)$ entering the kernel is hypothesis–dependent:

- **Vectors** $H \in \{H_V, H_M, H_T\}$ use the comoving vector cutoff *as defined* in (13.58), with production/coherence history encoded by $\Xi_{\text{prod}}(H)$; the associated kernel is denoted $\mathcal{S}(\cdot; k_c^{(V)}, \nu_V)$ (cf. Proposition 13.73 for H_V).
- **Scalar** $H = H_S$ uses the Jeans–like cutoff *as defined* in (13.59); the associated kernel is denoted $\mathcal{S}(\cdot; k_c^{(S)}, \nu_S)$.

No additional structure is introduced here; these expressions are used verbatim.

Assumption 13.37 (Kernel regularity and normalization)**.** *On $\mathfrak{P}$, the coherence kernel satisfies: (i) $\mathcal{S}(0; k_c, \nu) = 1$ (large–scale normalization), (ii) $0 \le \mathcal{S}(k; k_c, \nu) \le 1$ for $k \ge 0$, and (iii) piecewise C^2 in k for $k > 0$. For vector hypotheses, $\mathcal{S}(\cdot; k_c^{(V)}, \nu_V)$ may exhibit damped* oscillatory tails *beyond $k \gtrsim k_c^{(V)}$; for the scalar hypothesis, $\mathcal{S}(\cdot; k_c^{(S)}, \nu_S)$ is asymptotically monotone beyond $k \gtrsim k_c^{(S)}$. These properties are understood as diagnostic realized reference configuration–calibrated and consistent with Section 13.2.3.*

Proposition 13.38 (Decision–useful discriminants at small scales)**.** *Under Definitions 13.35 and 13.36 and Assumption 13.37, the following provide Stage II discriminants (Definition 13.29):*

(i) Cutoff location

The inferred $k_c(H)$ from $P_{\text{lin}}(k, z; H)$ identifies the hypothesis–specific scale family (Definition 13.36) and constrains $\{m_S, M_{\tau\text{-vec}}, M_\Phi, M_\Theta\}$ through (13.59) and (13.58).

(ii) Tail morphology
The presence of damped oscillatory features for $k \gtrsim k_c$ favors vector hypotheses (H_V, H_M, H_T), while asymptotically monotone suppression favors the scalar hypothesis (H_S).

(iii) Cross–family consistency
The same (k_c, ν) inferred from matter power must propagate consistently to WL shear and RSD/growth, and remain compatible with BAO/Lyα probes (meters catalogued in Section 13.2.3).

These features enter the asymmetric visibility and coherence score without redefining the Stage II functionals (eqs. (13.48) and (13.49)).

Remark 13.39 (Operational note)**.** In applications, $T_{\rm vis}(k)$ and $D(z)$ are fixed by the visible/diagnostic realized reference configuration sector, while $\mathcal{S}(k; k_c, \nu)$ carries the dark–sector imprint. The interferometer therefore attributes deviations from the visible baseline entirely to the kernel parameters (k_c, ν), which serve as the tunable handles in the selection argmax (13.50).

Anisotropy and stress constraints. We monitor the anisotropic–stress tensor $\pi^i{}_j := T^i{}_j - \delta^i{}_j T^k{}_k/3$ for vector hypotheses:

$$\begin{aligned} \text{CK background } (H_V): &\quad \big\langle \pi^i{}_j[C_\mu] \big\rangle_{\rm osc} \simeq 0 \ \text{ on FLRW (isotropy constraint)}, \\ \text{Connection vectors } (H_M, H_T): &\quad \big\langle \pi^i{}_j \big\rangle_{\rm late} \to 0, \end{aligned} \tag{13.60}$$

and verify that any residual gauge mixings are *projectively null* on the visible screen,

$$\mathsf{P}_{\rm vis}\Big[\text{mixing current}\Big] = 0 \quad \text{on} \quad \mathcal{H}_{\rm phys},\ \mathfrak{P}, \tag{13.61}$$

ensuring TT purity (Definition 10.3) and SM–orthogonality.

Assumption 13.40 (Vector-sector isotropy and projective-null mixing)**.** *For vector hypotheses $H \in \{H_V, H_M, H_T\}$, the anisotropic–stress conditions are imposed* as stated *in* (13.60)*, and any residual gauge mixings with visible meters are* projectively null *on the screen,*

$$\mathsf{P}_{\rm vis}\Big[\textit{mixing current}\Big] = 0 \quad \textit{on} \quad \mathcal{H}_{\rm phys},\ \mathfrak{P}, \tag{13.62}$$

i.e. exactly as in (13.61)*. Under these conditions, TT purity is preserved (Definition 10.3) and SM–orthogonality holds (Definition 13.1).*

Definition 13.41 (Stage I anisotropy gate)**.** Fix diagnostic realized reference configuration tolerances $\tau_\pi^{(\rm CK)}$, $\tau_\pi^{(\rm conn)} > 0$. Let $\langle \pi^i{}_j \rangle_{\rm osc}$ and $\langle \pi^i{}_j \rangle_{\rm late}$ be the diagnostic realized reference configuration–averaged anisotropic stresses referenced in (13.60). Define the Stage I *anisotropy acceptance* for vectors by

$$\begin{aligned} H_V \ \in\ \mathfrak{A}_I^{(\pi)} \quad &\Longleftrightarrow \quad \Big\| \langle \pi^i{}_j[C_\mu] \rangle_{\rm osc} \Big\| \ \le\ \tau_\pi^{(\rm CK)}, \\ H_{M/T} \ \in\ \mathfrak{A}_I^{(\pi)} \quad &\Longleftrightarrow \quad \Big\| \langle \pi^i{}_j \rangle_{\rm late} \Big\| \ \le\ \tau_\pi^{(\rm conn)}, \end{aligned} \tag{13.63}$$

with the norm understood as any fixed diagnostic realized reference configuration–calibrated operator norm. Rejection occurs when (13.63) fails, or when (13.62) fails.

Proposition 13.42 (Consistency with TT purity and selection workflow)**.** *Under Assumption 13.40, all vector hypotheses that satisfy (13.63) preserve TT purity on the realized weave configuration and remain SM–orthogonal, hence are admissible for the HSP/Interferometer pipeline (Section 2.4 and Theorem 12.79). Violations lead to Stage I exclusion (cf. Definition 13.27).*

Corollary 13.43 (Meters and decision-useful consequences)**.** *The anisotropy gate (13.63) propagates to Stage II meters: excess vector anisotropic stress would bias WL shear and large-angle CMB/lensing consistency, while admissible vectors exhibit WL/RSD signatures governed chiefly by the small–scale kernel $\mathcal{S}(k; k_c^{(V)}, \nu_V)$ (eqs. (13.57) and (13.58)), in line with the distinguishing-signature catalogue (Section 13.2.3).*

Portal bounds and spectral orthogonality. Allowed portals enter as suppressed bilinears,

$$\mathcal{L}_{\text{portal}} = \sum_{\alpha} \varepsilon_\alpha \, \mathcal{O}_\alpha^{(\text{dark})} \, \mathcal{O}_\alpha^{(\text{vis})}, \qquad 0 \le \varepsilon_\alpha \ll 1, \tag{13.64}$$

and must satisfy the *darkness* projector criterion and SM spectral decoupling,

$$\mathcal{P}_G^{\text{coh}} \, \mathcal{O}_\alpha^{(\text{dark})} \neq 0, \qquad \mathsf{P}_{\text{vis}} \, \mathcal{O}_\alpha^{(\text{dark})} = 0 \quad (\text{on } \mathcal{H}_{\text{phys}}, \ \mathfrak{P}; \ \textit{Definitions 12.77 and 13.1}). \tag{13.65}$$

Equations (13.57)–(13.65) provide the quantitative levers for Stage II tomography and the HSP argmax (Section 13.2.2).

Definition 13.44 (Admissible portal family)**.** Portals are taken *as defined* in (13.64) with coefficients $\boldsymbol{\varepsilon} = (\varepsilon_\alpha)_\alpha$ and must satisfy the *darkness* projector criterion and SM spectral decoupling *as stated* in (13.65). An admissible portal family on the certified plateau $\mathfrak{P}$ is any subset of (13.64) obeying these constraints on $\mathcal{H}_{\text{phys}}$, with all operators typed by the commuting projector family (TT, τ–types, Riesz selectors; Definitions 10.3, 10.4 and 10.7).

Assumption 13.45 (TT/BRST hygiene and diagnostic realized reference configuration suppression)**.** *Each admissible portal obeys: (i) TT–purity, i.e. $[\Pi_{\bar{\Sigma}}^{[\text{TT}]}, \mathcal{O}_\alpha^{(\text{dark})}] = 0$ so no spin–2 leakage (Definition 10.3); (ii) BRST cleanliness, i.e. $\mathcal{O}_\alpha^{(\text{dark})}$ defines a class on $\mathcal{H}_{\text{phys}}$ (Section 12.3); (iii) diagnostic realized reference configuration suppression $\|\boldsymbol{\varepsilon}\|_\infty \le \varepsilon_{\max} \ll 1$ on $\mathfrak{P}$ (Definition 9.58 and Theorem 9.351).*

Proposition 13.46 (Interferometric robustness under small portals)**.** *Let H be any admissible hypothesis and $H[\boldsymbol{\varepsilon}]$ its portal–deformed realization with (13.64). Under Definition 13.44 and Assumption 13.45 there exists a diagnostic realized reference configuration constant $C_{\text{port}}(H, \mathfrak{P})$ such that the Stage II functionals satisfy the Lipschitz bounds*

$$\left| \mathcal{V}_{\text{asym}}[H[\boldsymbol{\varepsilon}], H[\boldsymbol{\varepsilon}]] - \mathcal{V}_{\text{asym}}[H, H] \right| \ \le \ C_{\text{port}}(H, \mathfrak{P}) \, \|\boldsymbol{\varepsilon}\|_\infty \ + \ \mathcal{O}(\|\boldsymbol{\varepsilon}\|_\infty^2), \tag{13.66}$$

and consequently

$$\left| \mathscr{C}(H[\boldsymbol{\varepsilon}]; \mathfrak{P}) - \mathscr{C}(H; \mathfrak{P}) \right| \ \le \ C_{\text{port}}(H, \mathfrak{P}) \, \|\boldsymbol{\varepsilon}\|_\infty \ + \ \mathcal{O}(\|\boldsymbol{\varepsilon}\|_\infty^2), \tag{13.67}$$

where $\mathcal{V}_{\text{asym}}$ and $\mathscr{C}$ are as defined *in (13.48)–(13.49). Thus small, TT/BRST–clean portals perturb visibilities and scores only at $\mathcal{O}(\boldsymbol{\varepsilon})$ and do not upset the HSP argmax selection on $\mathfrak{P}$.*

Corollary 13.47 (Operational treatment of portals in selection)**.** *In Stage II tomography, the parameters $\boldsymbol{\varepsilon}$ are treated as* nuisance *and either (i) profiled at fixed diagnostic realized reference configuration priors, or (ii) marginalized with diagnostic realized reference configuration–calibrated priors, while enforcing the constraints* (13.65)*. The impact of portals then enters through the kernel parameters (k_c, ν) in* (13.57) *and through the small corrections bounded by* (13.66)–(13.67)*, leaving the HSP argmax* (13.50) *and discriminants in Section 13.2.3 unchanged at leading order.*

Remark 13.48 (When portals trigger exclusion)**.** If any portal violates (13.65), breaks TT–purity, or exceeds the diagnostic realized reference configuration tolerance $\varepsilon_{\max}$, the hypothesis is rejected at Stage I via the acceptance set in Definition 13.27. Otherwise, portals remain permissible but constrained, contributing only subleading corrections to the interferometric meters.

13.2.4 Multi-Component Selection (Optional)

When evidence favors mixtures, we admit a *mixture hypothesis* $H_{\mathrm{mix}}(\boldsymbol{\pi})$ over the component set $\mathcal{I} = \{S, V, M, T\}$ (cf. Section 13.2.1), with nonnegative weights $\boldsymbol{\pi} = (\pi_i)_{i\in\mathcal{I}}$ satisfying $\sum_i \pi_i = 1$. For interferometric algebra, it is convenient to use $\mathbf{v} := \sqrt{\boldsymbol{\pi}}$, i.e. $v_i := \sqrt{\pi_i}$, so that $\mathbf{v} \geq 0$ and $\|\mathbf{v}\|_2^2 = \sum_i v_i^2 = 1$.

Mixture path laws and Stage I (symmetric) overlap. Let $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N(H_i)}$ and $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G(H_i)}$ be the two register-wide path laws for component H_i (eqs. (2.69) and (2.70)). Define the *mixture* path operators as coherent amplitude sums on the diagnostic realized reference configuration $\mathfrak{P}$:

$$\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N\left(H_{\mathrm{mix}}(\boldsymbol{\pi})\right)} := \sum_{i\in\mathcal{I}} v_i\, \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N(H_i)}, \qquad \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G\left(H_{\mathrm{mix}}(\boldsymbol{\pi})\right)} := \sum_{i\in\mathcal{I}} v_i\, \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G(H_i)}. \tag{13.68}$$

For a physical state $|\Psi_{\mathrm{RR}}\rangle$ (Section 12.3), the symmetric overlap and visibility become

$$\begin{aligned} \mathcal{O}_{\mathrm{mix}} &= \sum_{i,j} v_i v_j \left\langle \Psi_{\mathrm{RR}} \middle| \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G(H_j)\dagger} \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N(H_i)} \middle| \Psi_{\mathrm{RR}} \right\rangle = \mathbf{v}^{\mathsf{T}}\, \mathbf{K}^{(\mathrm{sym})}\, \mathbf{v}, \\ \mathcal{V}_{\mathrm{sym}}(H_{\mathrm{mix}}) &= \Big| \mathrm{Re}\ \mathcal{O}_{\mathrm{mix}} \Big|, \end{aligned} \tag{13.69}$$

with kernel entries $K_{ij}^{(\mathrm{sym})} := \langle \Psi_{\mathrm{RR}}|\, \hat{\mathbf{U}}_{A_N B_G(H_j)}^{\dagger} \hat{\mathbf{U}}_{A_G B_N(H_i)} |\Psi_{\mathrm{RR}}\rangle$. If induced K–actions of $H_i \neq H_j$ are holosymmetrically orthogonal, then $K_{ij}^{(\mathrm{sym})} = 0$ on $\mathcal{H}_{\mathrm{phys}}$ and cross–terms drop out (Proposition 12.76).

Stage II (asymmetric) mixture visibility. Let the asymmetric control blocks be $\mathcal{F}_{k0}^{(\mathrm{asym})}(H_i)$ ($k = 0, 1$; Section 3.1.5). Define diagnostic realized reference configuration–dependent amplitude vectors $|A_i\rangle := \mathcal{F}_{00}^{(\mathrm{asym})}(H_i)|\Psi_{\mathrm{RR}}\rangle$, $|B_i\rangle := \mathcal{F}_{10}^{(\mathrm{asym})}(H_i)|\Psi_{\mathrm{RR}}\rangle$. Then the mixture

outcome probabilities and visibility read

$$\begin{aligned} p_0^{(\mathrm{asym})}\big(H_{\mathrm{mix}}\big) &= \Big\| \sum_i v_i |A_i\rangle \Big\|^2, \\ p_1^{(\mathrm{asym})}\big(H_{\mathrm{mix}}\big) &= \Big\| \sum_i v_i |B_i\rangle \Big\|^2, \\ \mathcal{V}_{\mathrm{asym}}\Big[H_{\mathrm{mix}}, H_{\mathrm{mix}}\Big] &= \Big| \mathbf{v}^{\mathsf{T}}\, \mathbb{K}(U_c)\, \mathbf{v} \Big|, \end{aligned} \tag{13.70}$$

where the Hermitian $\mathbb{K}(U_c)$ depends on the SU(2) controls and is given by $\mathbb{K} := \mathbf{A}^\dagger\mathbf{A} - \mathbf{B}^\dagger\mathbf{B}$ with $(\mathbf{A}^\dagger\mathbf{A})_{ij} = \langle A_j|A_i\rangle$ and $(\mathbf{B}^\dagger\mathbf{B})_{ij} = \langle B_j|B_i\rangle$ (eq. (13.48)). The *mixture coherence score* is

$$\mathscr{C}\Big(H_{\mathrm{mix}}(\boldsymbol{\pi}); \mathfrak{P}\Big) := \sup_{U_c^{\mathrm{prep}},\, U_c^{\mathrm{read}} \in SU(2)} \mathcal{V}_{\mathrm{asym}}\Big[H_{\mathrm{mix}}, H_{\mathrm{mix}}\Big]\Big|_{\mathcal{H}_{\mathrm{phys}}, \mathfrak{P}}. \tag{13.71}$$

Constrained maximization over mixture weights. We determine $\boldsymbol{\pi}$ by a diagnostic realized reference configuration–aware penalized program

$$\max_{\boldsymbol{\pi}} \;\; \mathscr{C}\Big(H_{\mathrm{mix}}(\boldsymbol{\pi}); \mathfrak{P}\Big) \;-\; \alpha\, \mathrm{Ent}(\boldsymbol{\pi}) \;-\; \beta\, \mathcal{B}(\boldsymbol{\pi}) \quad \text{s.t.} \quad \pi_i \geq 0, \;\; \sum_i \pi_i = 1, \tag{13.72}$$

where $\mathrm{Ent}(\boldsymbol{\pi}) := -\sum_i \pi_i \log \pi_i$ is a simplicity prior that penalizes unnecessary spread (favoring parsimonious, low–entropy mixes), and $\mathcal{B}(\boldsymbol{\pi}) = \sum_i \pi_i\, \chi_i$ encodes diagnostic realized reference configuration–complexity costs (instrumentation/realized weave configuration weights χ_i; cf. Definition 10.1). Equivalently, in variables $\mathbf{v} = \sqrt{\boldsymbol{\pi}}$, the constraints are $\mathbf{v} \geq 0$, $\|\mathbf{v}\|_2^2 = 1$, and the objective uses the quadratic form $\mathbf{v}^{\mathsf{T}}\mathbb{K}(U_c)\mathbf{v}$ (eq. (13.70)).

Admissibility and orthogonality. All mixture components must satisfy the dark–sector projector criterion and TT/BRST hygiene on $\mathfrak{P}$ (Definitions 9.58, 10.3 and 13.1 and Section 12.3). When induced K–actions are inequivalent, holosymmetric orthogonality forces cross–terms to vanish on $\mathcal{H}_{\mathrm{phys}}$, in which case $\mathscr{C}$ reduces to a diagonal quadratic form in $\boldsymbol{\pi}$ (Proposition 12.76). The program (13.72) then yields a decisive, parsimonious selection consistent with the HSP argmax rule (Section 13.2.2).

13.2.5 Diagnostics (Not Selectors)

The following diagnostics accompany, but do *not* replace, the HSP selection in Section 13.2.2. They are used to (i) flag visibility created by control framing rather than genuine whole–evolution differences, and (ii) certify that results are stable under realization choices on a certified plateau $\mathfrak{P}$ (Definitions 9.58 and 9.60 and Theorem 9.351).

Commutator lensing (symmetric form). For a fixed hypothesis H, define the *commutator lens* as in Chapter 3

$$\mathbf{U}_{\mathrm{lens}}^{(\mathrm{sym})}(H) := \Big[\mathbf{U}_{\mathrm{Had}},\, \mathbf{U}_{\mathrm{ctrl}}(H)\Big]\; \mathbf{U}_{\mathrm{Had}}^\dagger \qquad (\textit{Definition 3.12}), \tag{13.73}$$

where $\mathbf{U}_{\mathrm{ctrl}}(H)$ inserts the two path laws of H (eqs. (2.69) and (2.70) and Definition 2.114). We quantify internal non–commutation by either the on–state lensing power or the operator–norm proxy,

$$\mathcal{L}_{\mathrm{sym}}^{(\mathrm{state})}(H;\Psi) := \left\| \mathbf{U}_{\mathrm{lens}}^{(\mathrm{sym})}(H)\,|\Psi_{\mathrm{RR}}\rangle \right\|^2, \qquad \mathcal{L}_{\mathrm{sym}}^{(\mathrm{HS})}(H) := \left\| \left[\mathbf{U}_{\mathrm{Had}},\, \mathbf{U}_{\mathrm{ctrl}}(H)\right] \right\|_{\mathrm{HS}}^2, \tag{13.74}$$

evaluated on $\mathcal{H}_{\mathrm{phys}}$ and $\mathfrak{P}$ (Section 12.3 and Definition 9.58). Large $\mathcal{L}_{\mathrm{sym}}$ together with nearly unchanged symmetric overlap $\mathcal{O}_H$ (eq. (13.46)) flags *spurious visibility* due to control framing rather than genuine differences in whole–evolution channels. These diagnostics do not enter the HSP argmax but gatekeep against false positives.

Diagnostic realized reference configuration sensitivity (Nielsen/BRST stability). Let $\{\mathfrak{P}_b\}_{b\in\mathcal{B}}$ be admissible realizations ("band tags") of the certified plateau $\mathfrak{P}$ (Definitions 9.58 and 9.60). Using the coherence score $\mathscr{C}(H;\mathfrak{P})$ from eq. (13.49), define the diagnostic realized reference configuration–spread and a normalized stability ratio

$$\Delta_{\mathrm{DRRC}}(H) := \max_{b,b'\in\mathcal{B}} \left|\mathscr{C}(H;\mathfrak{P}_b) - \mathscr{C}(H;\mathfrak{P}_{b'})\right|, \tag{13.75}$$

$$\mathcal{R}_{\mathrm{Nielsen}}(H) := \frac{\Delta_{\mathrm{DRRC}}(H)}{\max_{b\in\mathcal{B}} \mathscr{C}(H;\mathfrak{P}_b) + \varepsilon}, \tag{13.76}$$

with a small $\varepsilon > 0$ to avoid division by zero. Acceptance requires $\mathcal{R}_{\mathrm{Nielsen}}(H) \le \epsilon_{\mathrm{N}}$ for a preset tolerance ϵ_{N} fixed by Theorem 9.351. This ensures that the interferometric decision is *Nielsen/BRST–stable* and not an artifact of the realization map. Should a candidate fail (13.75), it is flagged for re–instrumentation on the realized weave configuration (Definition 10.1) before re–entering the HSP pipeline.

Remark. The quantities $\mathcal{L}_{\mathrm{sym}}$, Δ_{DRRC}, and $\mathcal{R}_{\mathrm{Nielsen}}$ are *diagnostics*. They calibrate the integrity of the selection workflow in Section 13.2.2 but do not themselves perform selection, which remains governed by the Stage I/II visibilities and the HSP argmax.

13.2.6 Selecting the Mechanism of Dark Energy

Cosmic acceleration is adjudicated by the same Holosymmetric Selection/Interferometer machinery used elsewhere (Section 13.2.2) but applied to the dark–energy sector on the BRST–physical diagnostic realized reference configuration. Within the present framework, late–time acceleration is anchored in the typed τ–scalar channel—the Apeiron/Gaiaton (Definition 13.6 and Proposition 10.153)—whose vacuum potential mediates the regulated near–cancellation of the Gaiamiti and Tantrakyma zero–point contributions (Section 13.1.1). The effective "cosmological constant" therefore appears as a *residual branch* of the dual–field vacuum, not as an arbitrary parameter.

Notation: we use a branch–sign subscript on Λ, i.e. Λ_σ with $\sigma \in \{+,0,-\}$, to indicate respectively a positive residual ($\varepsilon_\Lambda > 0$), exact cancellation ($\varepsilon_\Lambda = 0$), or negative residual ($\varepsilon_\Lambda < 0$). Accordingly, the selection problem is cast as inequivalent universal histories with distinct induced K–actions, collected in Definition 13.49: a positive–residual slow–roll Apeiron branch $H_{\Lambda_+}^{(\phi)}$, an exact–cancellation branch H_{Λ_0}, an over–cancellation branch H_{Λ_-}, and (diagnostically) a coherent τ–background geometric option $H_{\Lambda_+}^{(\tau\text{-bg})}$.

Definition 13.49 (Dark–energy hypothesis family and cancellation branches)**.** Cosmic acceleration is selected holosymmetrically on the BRST–physical diagnostic realized reference configuration $\mathfrak{P}$ (Section 12.3 and Definition 9.58) using the interferometer of Section 13.2.2. The hypothesis set comprises the τ–scalar (Apeiron/Gaiaton) slow–roll branch together with cancellation branches for the dual–field vacuum:

$$\begin{aligned}\mathcal{H}^{\mathrm{DE}}_{\mathrm{cand}} := \Big\{ & H^{(\phi)}_{\Lambda+} && \text{(Apeiron slow–roll, positive residual)},\\ & H_{\Lambda 0} && \text{(exact cancellation)},\\ & H_{\Lambda-} && \text{(over–cancellation / negative residual)}\Big\}\\ \cup \Big\{ & H^{(\tau\text{-bg})}_{\Lambda+} && \text{(coherent } \tau\text{–background / geometric DE)}\Big\}.\end{aligned} \tag{13.77}$$

Here $H^{(\phi)}_{\Lambda+}$ is the typed τ–scalar realization (Definition 13.6) satisfying the slow–roll diagnostics (13.52) with $w_\phi(a)$ in (13.53), and the branch discriminator (13.54). The coherent τ–background option $H^{(\tau\text{-bg})}_{\Lambda+}$ refers to an acceleration sourced by the expectation value $\bar{t}_{\mu\nu}$ entering the chronogeometric equations (diagnostic DE/geometry channel; cf. Definition 10.13, eq. (11.191), and Remark 13.5).

Remark 13.50 (Dual–field vacuum and residual density)**.** In all Λ–branches, the residual vacuum density arises from the regulated sum of the Gaiamiti and Tantrakyma condensates as established in the dual–field vacuum discussion, with the scalar Apeiron controlling residual dynamics on the diagnostic realized reference configuration (cf. Section 13.1.1; see also Definitions 5.26 and 9.357 for the underlying variational and chronogeometric structures).

Definition 13.51 (Stage I gate for dark–energy branches)**.** Using the Stage I functionals from Lemma 13.23, define the acceptance set

$$\mathfrak{A}^{\mathrm{DE}}_I(\rho;\mathfrak{P}) := \Big\{ H \in \mathcal{H}^{\mathrm{DE}}_{\mathrm{cand}} :\ \mathcal{C}_{\mathrm{phys}}(H \mid \rho) \geq \tau_I,\ \text{unitary/stable on } \mathfrak{P},\ \mathcal{R}_{\mathrm{Nielsen}}(H) \leq \epsilon_{\mathrm{N}} \Big\}$$

(Definition 13.27).
Additional Stage I filters.

(a) Apeiron slow–roll branch $H^{(\phi)}_{\Lambda+}$
require the slow–roll window $\{\epsilon_V, |\eta_V|\} \ll 1$ (eq. (13.52)) so that $w_\phi \simeq -1$ (eqs. (13.53) and (13.54)).

(b) Exact–cancellation branch $H_{\Lambda 0}$
the diagnostic realized reference configuration expansion must be non–accelerating and microcausal on $\mathfrak{P}$.

(c) Over–cancellation branch $H_{\Lambda-}$
recollapse is permitted only if the evolution remains unitary and hyperbolic on the certified interval.

(d) Coherent τ–background option $H^{(\tau\text{-bg})}_{\Lambda+}$
must satisfy the typed balance/chronogeometric constraints (cf. eq. (11.191)) and pass the diagnostic exclusion in Remark 13.5.

Proposition 13.52 (Stage II tomography and DE coherence score)**.** *With the asymmetric instruments (Definition 13.28), the outcome probabilities and visibility are those of* (13.48)*, and the dark–energy* coherence score *is*

$$\mathscr{C}_{\mathrm{DE}}(H;\mathfrak{P}) := \sup_{U_c^{\mathrm{prep}},\,U_c^{\mathrm{read}}\in SU(2)} \mathcal{V}_{\mathrm{asym}}[H,H]|_{\mathcal{H}_{\mathrm{phys}},\,\mathfrak{P}} \qquad (\textit{cf.}\ (13.49)).$$

Admissible meters are the background–sensitive families (SNe, CMB distances, BAO) and growth–sensitive families (WL, RSD), treated as interferometric effects (Section 2.4 and Remark 13.31).

Meters in use. For $H_{\Lambda+}^{(\phi)}$, selection is driven by $(w_\phi, dw_\phi/da)$ (eqs. (13.52) and (13.53)) with negligible $\mathcal{S}$–modulation (Corollary 13.9); for $H_{\Lambda+}^{(\tau\text{-bg})}$, $\mathcal{S}\to 1$ in eq. (13.57) and the background is governed by eq. (11.191).

Theorem 13.53 (HSP selection of the dark–energy mechanism (bright fringe))**.** *Holosymmetric orthogonality on $\mathcal{H}_{\mathrm{phys}}$ for inequivalent induced K–actions implies pairwise discriminability of the branches (Proposition 13.30). The realized mechanism of dark energy is therefore*

$$H_{\Lambda\star} = \arg \max_{H\in\mathfrak{A}_I^{\mathrm{DE}}(\rho;\mathfrak{P})} \mathscr{C}_{\mathrm{DE}}(H;\mathfrak{P}),$$
$$\textit{subject to SM–orthogonality and DRRC hygiene}$$
$$(\textit{Chapter 12and Definitions 9.58and 13.1}). \tag{13.78}$$

Here "DRRC hygiene" means compatibility with the diagnostic realized reference configuration of Definition 9.185, including its BRST/Nielsen flatness and plateau tolerances.

In interferometric language, $H_{\Lambda\star}$ is the bright fringe*: it maximizes constructive self–interference while remaining dark to SM meters. Concretely, "constructive self–interference'' means that the Stage–II visibility $\mathcal{V}_{\mathrm{asym}}[H,H]$, optimized over preparation/readout controls $(U_c^{\mathrm{prep}}, U_c^{\mathrm{read}})$ and restricted to $\mathcal{H}_{\mathrm{phys}}$, attains the supremum that defines the coherence score $\mathscr{C}_{\mathrm{DE}}(H;\mathfrak{P})$ (eq. (13.8)). "Dark to SM meters'' means portal responses vanish within DRRC tolerances under the darkness projector, i.e. $\mathcal{P}_{\mathrm{dark}}\circ\hat{\mathcal{O}}_{\mathrm{vis}} = 0$ and the induced SM readouts are below tolerance (eqs. (13.64) and (13.65) and Definition 9.58). Equivalently, all SM–orthogonal witnesses satisfy $\langle\hat{O}_{\mathrm{SM}}\rangle_{H_{\Lambda\star}} = 0$ on $\mathfrak{P}$ (up to DRRC hygiene tolerances). In the event of a tie in $\mathscr{C}_{\mathrm{DE}}$ over $\mathfrak{A}_I^{\mathrm{DE}}$, the discriminability/uniqueness criterion of Proposition 13.30 selects a unique branch up to null–visibility degeneracies.*

Corollary 13.54 (Decision–useful signatures for the DE selection)**.** *For $H_{\Lambda+}^{(\phi)}$ (Apeiron slow–roll), the leading meters are $(w_\phi(a), dw_\phi/da)$ derived from eqs. (13.52) and (13.53), with negligible small–scale kernel modulation (cf. Corollary 13.9). For $H_{\Lambda+}^{(\tau\text{-}bg)}$, the background expansion is governed by the chronogeometric equations with $w\simeq -1$ and TT purity preserved by the weave projector (Definition 10.3); small–scale structure proceeds through the visible transfer $T_{\mathrm{vis}}(k)$ with $\mathcal{S}\to 1$ in (13.57). The $H_{\Lambda 0}$ and $H_{\Lambda-}$ branches are disfavored by background meters unless specifically supported by the interferometric overlaps; failure of Stage I stability, or violation of microcausality/diagnostic realized reference configuration cleanliness, triggers exclusion (Definitions 13.27 and 13.51 and Theorem 9.351).*

Remark 13.55 (Mixtures and diagnostic guardrails)**.** If evidence favors a blended mechanism (e.g. a small coherent τ–background atop a slow–roll Apeiron), one may introduce a DE mixture in the spirit of Section 13.2.4, optimizing the mixture weights under parsimony penalties. Diagnostic integrity is monitored by commutator lensing and diagnostic realized reference configuration–stability ratios (Section 13.2.5), ensuring that a bright–fringe decision is not an artifact of control framing or realization.

Table 13.1: Summary of Dark Energy Hypothesis Observables (HSP)

Hypothesis	**Primary Observables / Meters**	**Key Features / Theoretical Constraints**
$H^{(\phi)}_{\Lambda+}$ (Apeiron Slow-Roll)	Background probes (SNe, CMB distances, BAO) constrain $(w_\phi, dw_\phi/da)$ via eqs. (13.52) and (13.53). Growth probes (WL, RSD) provide consistency checks.	Governed by slow-roll dynamics. Predicts negligible small-scale power modulation ($\mathcal{S} \to 1$ in eq. (13.57); cf. Corollary 13.9).
$H^{(\tau\text{-bg})}_{\Lambda+}$ (Coherent τ-Background / Geometric DE)	Background probes (SNe, CMB distances, BAO) test the chronogeometric EFE prediction of $w \approx -1$ (eq. (11.191)).	Expansion follows chronogeometry. Small-scale structure follows visible transfer function ($\mathcal{S} \to 1$). TT purity is preserved by the weave projector (Definition 10.3).
$H_{\Lambda 0}$ (Exact Cancellation)	Background probes (SNe, CMB distances, BAO).	Predicts non-accelerating expansion. Strongly disfavored by data; typically rejected at Stage I if inconsistent (Definition 13.51).
$H_{\Lambda-}$ (Over-Cancellation)	Background probes (SNe, CMB distances, BAO).	Predicts deceleration/recollapse. Generally disfavored unless confined to certified intervals ensuring unitary/hyperbolic evolution per Stage I (Definition 13.51).

Note: All observables are treated within the Stage II interferometric tomography framework (Section 2.4).

13.2.7 The Lensing Model as the Underlying Mechanism of Selection

The *Holosymmetric Interferometer Model* formalizes how competing universal histories are compared at the level of observable effects, producing bright and dark fringes that encode the selection outcome (Section 2.4). The *Commutator Lensing Model* (Section 3.3) exposes the algebraic mechanism that *causes* those fringes: it isolates the noncommutativity—the

"algebraic twist"—between preparation and evolution on the BRST–physical diagnostic realized reference configuration. Concretely, if Π denotes the preparation (projector onto an admissible effect–family) and $\mathcal{U}$ the evolution/transport (parallel transporter along a history), the lensing functional depends on commutators and nested commutators,

$$[\Pi, \mathcal{U}], \quad [\Pi, [\Pi, \mathcal{U}]], \ \ldots,$$

which quantify how much "prepare then evolve" fails to commute with "evolve then prepare." Large algebraic twist implies strong phase–shear in the interferometer and a pronounced bright/dark pattern; vanishing twist yields trivial lensing and degenerate interference. Thus, the Lensing Model serves as the diagnostic *under the hood* of the interferometer: it computes the algebraic source of the selection signal, while the interferometer itself displays the resulting coherence pattern and implements Stage II tomography for dark matter and dark energy (Section 2.4).

Dark Energy as a Global Interferometric Residue

Notation (curvature–sign branches). Curvature–sign branches of the coherent vacuum will be indexed by $\chi \in \{\mathrm{dS}, \mathrm{Mkw}, \mathrm{AdS}\}$, corresponding to:

$(\chi = \mathrm{dS})$ (> 0): de Sitter (accelerating),

$(\chi = \mathrm{Mkw})$ $(= 0)$: Minkowski (flat),

$(\chi = \mathrm{AdS})$ (< 0): anti–de Sitter (negative curvature).

Branch–sign map. We relate the curvature label to the residual–sign subscript via $\sigma(\mathrm{dS}) = +$, $\sigma(\mathrm{Mkw}) = 0$, $\sigma(\mathrm{AdS}) = -$. When referring to hypothesis branches we write $H_{\Lambda_{\sigma(\chi)}}$, and for evolution operators we allow the shorthand $\mathcal{U}_\chi \equiv \mathcal{U}_{\Lambda_{\sigma(\chi)}}$.

Within the Holosymmetric Selection Principle (HSP), the small positive vacuum energy inferred from supernovae and the CMB is interpreted as a *global interferometric residue* of the vacuum, not as a fine–tuned cancellation [134, 135, 136, 414, 149]. Concretely, the vacuum sector is modeled as a superposition of the geometric (Gaiamiti) and material (Tantrakyma) contributions over the vacuum manifold $\mathcal{M}_{\mathrm{vac}}$, and the *Symmetric Interferometer Lensing* (Sections 2.4 and 3.8) acts as a universal coherence filter on this superposition.

Definition 13.56 (Interferometric residue of dark energy)**.** Let $H_{\mathrm{vac}} = H_{\mathrm{Gaiamiti}} + H_{\mathrm{Tantrakyma}}$ denote the regulated vacuum Hamiltonian on the BRST–physical space, and let $\mathcal{P}^{\mathrm{sym}}_{\mathrm{vac}}$ be the symmetry–adapted coherence projector associated with the symmetric (path–reversed) interferometer (Definition 12.77). The *interferometric residue* is the emergent/GNS expectation

$$\rho_{\mathrm{DE}} \ \equiv \ \frac{1}{V} \Big\langle \overset{\circ}{\boldsymbol{\Psi}} \Big| \, \mathcal{P}^{\mathrm{sym}}_{\mathrm{vac}} \, H_{\mathrm{vac}} \, \mathcal{P}^{\mathrm{sym}}_{\mathrm{vac}} \Big| \overset{\circ}{\boldsymbol{\Psi}} \Big\rangle_{\mathrm{phys}} \ \simeq \ \frac{1}{V} \Big\langle \overset{\circ}{\boldsymbol{\Psi}} \Big| \, H_{\mathrm{vac}} \Big| \overset{\circ}{\boldsymbol{\Psi}} \Big\rangle^{(\mathrm{coh})}_{\mathrm{phys}}, \tag{13.79}$$

where $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$ is the Quantum Holo–State (Definition 1.27) and V is the comoving volume element at the emergent layer; the superscript "(coh)" indicates that only the holosymmetry–coherent part of the vacuum survives the lensing filter (Section 13.2.7).

In the *Commutator Lensing Model* (Section 3.3) this residue is controlled by the commutator (and nested–commutator) twist between preparation and evolution,

$$\mathcal{L}_{\text{lens}} \;\propto\; [\Pi_{\text{sym}}, \mathcal{U}] \;+\; [\Pi_{\text{sym}}, [\Pi_{\text{sym}}, \mathcal{U}]] + \cdots ,$$

so that a near–perfect cancellation of incoherent components is enforced at Stage I (self–coherence), while Stage II compares the resulting coherent branches. Equivalently, the physical self–overlap

$$\Omega_{\text{phys}}(\mathcal{U}_\chi, \mathcal{U}_\chi \mid \mathring{\mathbf{\Psi}}) , \qquad \chi \in \{\text{dS}, \text{Mkw}, \text{AdS}\},$$

(i.e. $\Omega_{\text{phys}}(\mathcal{U}_{\Lambda_{\sigma(\chi)}}, \mathcal{U}_{\Lambda_{\sigma(\chi)}} \mid \mathring{\mathbf{\Psi}})$) induces the asymmetric visibility and the coherence score used operationally in selection (eqs. (13.48) and (13.49)), and the realized branch is determined by the HSP argmax described in Section 13.2.6 (cf. eq. (13.78)).

Interpretation. Thus the observed ρ_{DE} is the small, positive *spectral residue* of the symmetric, globally coherent vacuum—i.e. the "bright fringe'' left after destructive interference of the large, incoherent contributions from Gaiamiti and Tantrakyma. Its magnitude is governed by the coherence structure of the universal state (via $\mathcal{P}^{\text{sym}}_{\text{vac}}$ and Ω_{phys}), rather than by arbitrary parameter tuning, and is consistent with late–time acceleration [134, 135, 136, 414, 149].

Dark Matter as an Asymmetric Spectral Projection

In contrast to dark energy—selected by the *symmetric* (path–reversed) filter at the vacuum level—the dark–matter sector consists of *stable, localized excitations* that pass a *dark coherence* test while remaining *spectrally orthogonal* to Standard–Model (SM) meters. Operationally, these excitations are revealed by *asymmetric* interferometric tomography on the BRST–physical diagnostic realized reference configuration, using the Stage II instruments and controls of Section 13.2.2 (cf. eqs. (13.48) and (13.49)). Asymmetry here refers to a controlled breaking of the path–exchange symmetry used in the vacuum analysis: the preparation projector targets a *dark* effect–family (via the coherence projector onto a candidate gauge sector), while readout occurs on the dark screen, ensuring SM–decoupling (cf. Section 13.2.7, Definition 13.1, and eq. (13.61)).

Definition 13.57 (Dark–sector visibility/decoupling criterion)**.** Let $\mathcal{P}^{\text{coh}}_G$ denote the coherence projector onto a candidate dark gauge sector G (Definition 12.77), and let $\mathfrak{A}^{\text{obs}}_{\text{SM}}$ be the SM observable algebra at the emergent layer.

(i) Pure–state case. A one–particle state $|\psi\rangle$ is *physically realized and dark* iff

$$\|\mathcal{P}^{\text{coh}}_G|\psi\rangle\|^2 \;>\; 0 \qquad \text{and} \qquad \langle\psi|\,\mathcal{O}_{\text{SM}}\,|\psi\rangle_{\text{phys}} \;\approx\; 0 \quad \forall\, \mathcal{O}_{\text{SM}} \in \mathfrak{A}^{\text{obs}}_{\text{SM}}. \tag{13.80}$$

(ii) Mixed–state case. A density operator ρ describes a *physically realized and dark* state iff

$$\text{Tr}\big[\mathcal{P}^{\text{coh}}_G\,\rho\big] \;>\; 0 \qquad \text{and} \qquad \text{Tr}[\rho\,\mathcal{O}_{\text{SM}}] \;\approx\; 0 \quad \forall\, \mathcal{O}_{\text{SM}} \in \mathfrak{A}^{\text{obs}}_{\text{SM}}. \tag{13.81}$$

All expectations are taken on the BRST–physical space in the representation induced by the Quantum Holo–State $|\mathring{\mathbf{\Psi}}\rangle$ (Definition 1.27); (13.81) reduces to (13.80) for $\rho = |\psi\rangle\langle\psi|$.

Remark 13.58 (Quantitative tolerance for SM–decoupling)**.** In (13.80)–(13.81), the qualitative condition "≈ 0" may be replaced by a tolerance $\varepsilon_{\mathrm{SM}} \geq 0$ fixed by experiment:

$$\Big|\langle\psi \mid \mathcal{O}_{\mathrm{SM}} \mid \psi\rangle_{\mathrm{phys}}\Big| \leq \varepsilon_{\mathrm{SM}}, \qquad \Big|\mathrm{Tr}[\rho\, \mathcal{O}_{\mathrm{SM}}]\Big| \leq \varepsilon_{\mathrm{SM}} \quad \forall\, \mathcal{O}_{\mathrm{SM}} \in \mathfrak{A}_{\mathrm{SM}}^{\mathrm{obs}}.$$

Equivalently, fixing a C^*–norm $\|\cdot\|$ on $\mathfrak{A}_{\mathrm{SM}}^{\mathrm{obs}}$ (e.g. the operator norm in the GNS representation induced by $|\mathring{\mathbf{\Psi}}\rangle$), require

$$\sup_{\|\mathcal{O}_{\mathrm{SM}}\|\leq 1} \Big|\langle\psi \mid \mathcal{O}_{\mathrm{SM}} \mid \psi\rangle_{\mathrm{phys}}\Big| \leq \varepsilon_{\mathrm{SM}}, \qquad \sup_{\|\mathcal{O}_{\mathrm{SM}}\|\leq 1} \Big|\mathrm{Tr}[\rho\, \mathcal{O}_{\mathrm{SM}}]\Big| \leq \varepsilon_{\mathrm{SM}}.$$

Proposition 13.59 (Asymmetric lensing, dark visibility, and HSP selection)**.** *Let $H \in \{H_S, H_V, H_M, H_T\}$ be an admissible dark–sector hypothesis (cf. Section 13.2.1 and Definition 13.27). Using the asymmetric instruments of Definition 13.28, construct meters on the dark screen via $\mathcal{P}_G^{\mathrm{coh}}$ and compute the asymmetric visibility $\mathcal{V}_{\mathrm{asym}}[H, H]$ and coherence score $\mathscr{C}(H;\mathfrak{P})$ as in eqs. (13.48) and (13.49). If (13.80) or (13.81) holds and the portal/mixing constraints (13.64)–(13.65), (13.61) are satisfied, then H contributes to the HSP argmax selection (eq. (13.50)) as a* physically realized dark *component.*

Remark 13.60 (Commutator lensing origin of asymmetry)**.** In the Commutator Lensing Model (Section 3.3), dark–sector asymmetry arises from the twist generated by the *dark* preparation projector $\Pi_G \sim \mathcal{P}_G^{\mathrm{coh}}$ against the history evolution $\mathcal{U}_H$:

$$\mathcal{L}_{\mathrm{lens}}^{(G)} \;\propto\; [\Pi_G, \mathcal{U}_H] \;+\; [\Pi_G, [\Pi_G, \mathcal{U}_H]] + \cdots,$$

producing nontrivial interferometric visibility on the dark screen while remaining spectrally orthogonal to the SM meters (cf. Section 13.2.7 and Definition 13.1).

Candidates and selection. Models of *Asymmetric Path Scattering Lensing* (Section 3.8) provide the mechanism by which specific relational channels generate coherent, long–lived excitations in non–SM sectors. In particular, the vector *Spin–1 Chrono–Kymon* (Proposition 10.154), the scalar *Spin–0 Apeiron/Gaiaton* (Proposition 10.153), and the internal–connection vectors *Mition* and *Tantron* (cf. Section 13.2.1) can pass the dark filter when their typed structure satisfies the visibility/decoupling criterion (13.80) (and its mixed–state form (13.81)) together with the portal bounds (13.64)–(13.65) and the projectively–null mixing condition (13.61). Stage I (self–coherence) removes nonunitary or unstable histories on the certified plateau (Definition 13.27); Stage II compares viable candidates using the asymmetric visibility and the coherence score (eqs. (13.48) and (13.49)), which are induced by the physical self–overlap machinery cited in Section 13.1.4. If the induced K–actions are inequivalent, holosymmetric orthogonality forces the cross–overlaps to vanish (Proposition 12.76), and the realized dark–matter branch is selected by the HSP argmax (eq. (13.50))—the asymmetric "bright fringe" of the cosmic interferometer.

Interpretation. Dark matter is not "invisible" by paucity of properties; it is *coherent along relational directions* that have negligible interferometric visibility on SM meters. The gauge factor G of a realized dark component need not embed into G_{SM}; even when it does, its

representation content may have vanishing projection into SM interaction vertices at the emergent layer—precisely the spectral decoupling captured by (13.80) under the coherence projector $\mathcal{P}_G^{\mathrm{coh}}$ (Definition 12.77) and the dark–sector criterion (Definition 13.1). This narrative aligns with the same HSP/interferometer machinery that accounts for dark energy as a *symmetric, global interferometric residue* (Section 13.2.7), while distinguishing dark matter as the *asymmetric spectral projection* of coherent excitations that remain orthogonal to SM observables.

13.2.8 Holosymmetric Selection of the Dark Sector

Scope and method. The Holosymmetric Interferometer of Section 12.2 is not limited to gauge–group selection; it provides a *universal* selection mechanism for any relational structure that admits preparation/evolution and effect–families. Accordingly, the two–stage Holosymmetric Selection Principle (HSP) is applied on the BRST–physical space and a certified plateau (Definition 9.58) to the dark sector—dark matter (DM) and dark energy (DE). Stage I vets each candidate by *physical self–coherence* $\mathcal{C}_{\mathrm{phys}}(\,\cdot\,|\rho)$ (eq. (12.82)) and the acceptance set on the diagnostic realized reference configuration (Definition 13.27); Stage II compares the survivors via the *physical holosymmetric overlap* $\Omega_{\mathrm{phys}}(\,\cdot\,,\,\cdot\,|\,\mathring{\mathbf{\Psi}})$ (eq. (12.83)) as realized by the interferometer instruments (Section 13.2.2, cf. eqs. (13.48) and (13.49)). Holosymmetric orthogonality then excludes candidates that induce inequivalent K–actions (here K is the compact subgroup supported by the probe window; the induced K–action is the unitary representation on the BRST–physical space obtained by restricting the dynamics to K; Proposition 12.76). Operational membership in a dark sector G is enforced by the *coherence projector* $\mathcal{P}_G^{\mathrm{coh}}$ (Definition 12.77) together with spectral decoupling from Standard–Model (SM) observables (Definition 13.1 and Section 12.3.2).

Relational Comparison of Dark–Sector Hypotheses

Each physically meaningful dark–sector hypothesis G determines a distinct universal history $\mathcal{U}_G$ (total action, induced K–action, and admissible effects; cf. Section 13.2.1). Rather than a bare inner product of ground–state functionals, HSP compares hypotheses by the *physical* overlap functional of eq. (12.83),

$$\Omega_{\mathrm{phys}}(A, B \mid \mathring{\mathbf{\Psi}}),$$

evaluated on admissible effect–families at the emergent/GNS layer (Section 2.4). Stage I requires $\mathcal{C}_{\mathrm{phys}}(G \mid \rho) > 0$ and the diagnostic realized reference configuration hygiene of Definition 13.27. Stage II selects the realized hypothesis $G_\star$ by

$$G_\star \;=\; \arg\max_{G}\; \Omega_{\mathrm{phys}}(G, G \mid \mathring{\mathbf{\Psi}}), \qquad \Omega_{\mathrm{phys}}(G_\star, B \mid \mathring{\mathbf{\Psi}}) = 0 \;\; \forall\, B \neq G_\star, \tag{13.82}$$

where the orthogonality clause follows from Proposition 12.76 for inequivalent induced K–actions. Operationally, (13.82) is consistent with the interferometric argmax formulated via the asymmetric visibility and coherence score (eqs. (13.48) and (13.49)).

Remark 13.61 (Holo–state vs. BRST probe; overlap equivalence and argmax consistency)**.** The quantum holo–state $\mathring{\mathbf{\Psi}}$ is the *emergent/GNS holosymmetric probe* induced by the

admissible effect–families on the certified plateau (holosymmetric layer; cf. Section 2.4 and the tetralemma–GNS framework in Definition 1.16). Operationally, $\overset{\circ}{\mathbf{\Psi}}$ is the cyclic GNS probe for the holosymmetric algebra of effects used in Stage II.

By the canonical identification between algebraic and operator cohomology (eq. (12.81)), $\overset{\circ}{\mathbf{\Psi}}$ pushes forward to a BRST–physical probe $|\Psi_0\rangle$ supported on the common physical subspace,

$$\operatorname{supp}(|\Psi_0\rangle) \subset \mathcal{H}^{(A)}_{\text{phys}} \cap \mathcal{H}^{(B)}_{\text{phys}},$$

so that the physical overlap coincides across layers:

$$\Omega_{\text{phys}}(A, B \mid \overset{\circ}{\mathbf{\Psi}}) = \langle \Psi_0 | (\mathbf{U}^{(B)}_{\text{phys}})^\dagger \mathbf{U}^{(A)}_{\text{phys}} | \Psi_0 \rangle, \qquad \text{cf. eq. (12.83).} \tag{13.83}$$

Consistency with the interferometric argmax. In the asymmetric dual–arm interferometer the outcome probabilities obey

$$p_\pm(A, B) = \tfrac{1}{2}\Big(1 \pm \operatorname{Re}\, \Omega_{\text{phys}}(A, B \mid \overset{\circ}{\mathbf{\Psi}})\Big) \quad \text{(eq. (13.48))},$$

and the phase–calibrated coherence score is a monotone of $\operatorname{Re}\, \Omega_{\text{phys}}$ (or of $|\Omega_{\text{phys}}|$ after phase compensation; eq. (13.49)). Hence maximising the interferometric score is equivalent to maximising Ω_{phys}. On the diagonal,

$$\Omega_{\text{phys}}(G, G \mid \overset{\circ}{\mathbf{\Psi}}) = \langle \Psi_0 | (\mathbf{U}^{(G)}_{\text{phys}})^\dagger \mathbf{U}^{(G)}_{\text{phys}} | \Psi_0 \rangle \in (0, 1],$$

so the Stage II rule

$$G_\star = \arg\max_G \, \Omega_{\text{phys}}(G, G \mid \overset{\circ}{\mathbf{\Psi}}) \tag{13.84}$$

is precisely the interferometric argmax of self–visibility; and relational orthogonality for inequivalent induced K–actions implies $\Omega_{\text{phys}}(G_\star, B \mid \overset{\circ}{\mathbf{\Psi}}) = 0$ for all $B \neq G_\star$ (Proposition 12.76), i.e. zero cross–fringe (the "asymmetric visibility" null test). Stage I hygiene (Definition 13.27)—together with the positivity condition $\mathcal{C}_{\text{phys}}(G \mid \rho) > 0$ and the bound in eq. (12.82)—guarantees that the compressed arm blocks $\mathbf{U}^{(G)}_{\text{phys}}$ are well defined and that eq. (13.48) applies.

Relation to the tetralemma meters and Ch. 2 controls. At the fiberwise level the *local* Thesis/Antithesis/Synthesis meters are the orthoprojectors $P_T, P_A, P_{\text{para}}$ with Born semantics (tetralemma–GNS; Definition 1.16), realised concretely on the logical sector by the Ch. 2 constructions (Definitions 2.7 and 2.17). These furnish the preparation/readout controls that generate the effect–families entering the holosymmetric GNS layer, on which $\overset{\circ}{\mathbf{\Psi}}$ is defined.

Selection of Dark Matter Candidates

Let

$$\mathcal{U}_{\text{Gaiaton}}, \quad \mathcal{U}_{\text{Chrono}}, \quad \mathcal{U}_{\text{Mition}}, \quad \mathcal{U}_{\text{Tantron}}$$

denote, respectively, the universal histories sourced by a relic abundance of *Gaiatons*—Spin–0 Apeiron quanta (typed τ–scalar; Proposition 10.153)—Spin–1 *Chrono–Kymons* (typed τ–vector; Proposition 10.154), and the internal–connection vectors *Mition* (Gaiamiti connection)

and *Tantron* (Tantrakyma connection) (cf. Section 13.2.1 and eqs. (13.34) and (13.35)). Optionally, allow a convex mixture

$$\mathcal{U}_{\mathrm{mix}}(\boldsymbol{\pi}), \qquad \boldsymbol{\pi} = (\pi_S, \pi_V, \pi_M, \pi_T), \quad \sum_i \pi_i = 1,\ \pi_i \geq 0 \quad \text{(cf. Section 13.2.4)}.$$

Stage I (self–coherence on the diagnostic realized reference configuration). Each history is first vetted by Definition 13.27:

$$\mathcal{C}_{\mathrm{phys}}(\mathcal{U}_X \mid \rho) > 0$$
$$\text{and DRRC/BRST/Nielsen hygiene on } \mathfrak{P}$$
$$(\textit{Definition } 9.58, \textit{Section } 12.3, \textit{and Theorem } 9.351),$$

together with *darkness* constraints: the visibility–decoupling criterion Definition 13.57 (eqs. (13.80) and (13.81)), portal bounds eqs. (13.64) and (13.65), projectively–null mixings eq. (13.61), and (for vectors) anisotropy constraints eq. (13.60). Typed TT–safety and action blocks are as in Section 13.2.1.

Stage II (asymmetric tomography and selection). For each admissible candidate X (Gaiaton, Chrono, Mition, Tantron), compute the asymmetric visibility and the self–overlap. Use the interferometer meters described in Section 2.4: SNe Hubble residuals, CMB spectra, BAO distances, weak–lensing two–point functions, and growth/RSD.

$$\Omega_{\mathrm{phys}}(\mathcal{U}_X, \mathcal{U}_X \mid \mathring{\boldsymbol{\Psi}}), \qquad \mathcal{V}_{\mathrm{asym}}[X, X], \qquad \mathscr{C}(\mathcal{U}_X; \mathfrak{P}) \quad (\textit{eqs. } (13.48) \textit{and } (13.49)).$$

Holosymmetric orthogonality ensures that if the induced K–actions differ then $\Omega_{\mathrm{phys}}(\mathcal{U}_X, \mathcal{U}_Y \mid \mathring{\boldsymbol{\Psi}}) = 0$ for $X \neq Y$ (Proposition 12.76). The realized DM branch is the maximizer in

$$\arg \max_{X \in \{\text{Gaiaton, Chrono, Mition, Tantron}\}} \Omega_{\mathrm{phys}}(\mathcal{U}_X, \mathcal{U}_X \mid \mathring{\boldsymbol{\Psi}})$$
$$\text{(equivalently via } \mathscr{C}\text{; cf. eqs. (13.49) and (13.82))}.$$

Mixture path (if enabled). For $\mathcal{U}_{\mathrm{mix}}(\boldsymbol{\pi})$ use the mixture visibility and score

$$\mathcal{V}_{\mathrm{asym}}[\mathrm{mix}, \mathrm{mix}](\boldsymbol{\pi}), \qquad \mathscr{C}_{\mathrm{mix}}(\boldsymbol{\pi}; \mathfrak{P}) \quad (\textit{eqs. } (13.49) \textit{and } (13.70)),$$

and solve the constrained optimization

$$\max_{\boldsymbol{\pi}} \ \mathscr{C}_{\mathrm{mix}}(\boldsymbol{\pi}; \mathfrak{P}) \quad \text{s.t.} \quad \sum_i \pi_i = 1,\ \pi_i \geq 0 \quad (\textit{eq. } (13.72)),$$

with the parsimony control described in Section 13.2.4.

Distinguishing meters (decision–useful signatures). Small–scale and growth diagnostics are organized through the coherence–kernel factorization eq. (13.57). For the Gaiaton (scalar Apeiron), the misalignment/oscillatory branch yields a CDM–like transfer with Jeans–like cutoff $k_c^{(S)} \sim \max_a k_J(a)$ (eq. (13.59)); for vectors (Chrono–Kymon, Mition, Tantron), the coherence cutoff is $k_c^{(V)}(a) \sim a\, m_V\, \mathfrak{m}_{\mathrm{prod}}(H)$ with model–dependent oscillatory tails (eq. (13.58)). Anisotropy constraints and projectively–null mixings further separate internal–connection vectors from the τ–vector channel (eqs. (13.60) and (13.61)). Membership/"darkness'' is enforced by Definition 13.57 and the portal bounds eqs. (13.64) and (13.65).

Selection of the Residual Vacuum Energy

Consider the three cancellation histories for the dual–field vacuum (Gaiamiti + Tantrakyma; cf. Definition 13.49):

$$\mathcal{U}_{\Lambda_+}:\ \varepsilon_\Lambda > 0, \qquad \mathcal{U}_{\Lambda_0}:\ \varepsilon_\Lambda = 0, \qquad \mathcal{U}_{\Lambda_-}:\ \varepsilon_\Lambda < 0.$$

(For brevity we may write $\mathcal{U}_\sigma := \mathcal{U}_{\Lambda_\sigma}$ with $\sigma \in \{+, 0, -\}$.) Stage I excludes nonunitary or unstable histories on the certified plateau (Definition 13.51). Stage II then maximizes the physical self–overlap over the branch signs,

$$\max_{\sigma\in\{+,0,-\}} \Omega_{\mathrm{phys}}(\mathcal{U}_{\Lambda_\sigma}, \mathcal{U}_{\Lambda_\sigma} \mid \mathring{\mathbf{\Psi}}),$$

yielding a unique maximizer $\sigma_\star$; the realized mechanism is $H_{\Lambda\star} = H_{\Lambda_{\sigma_\star}}$ with evolution $\mathcal{U}_{\Lambda_{\sigma_\star}}$. Cross–overlaps vanish among inequivalent induced K–actions,

$$\Omega_{\mathrm{phys}}(\mathcal{U}_{\Lambda_\sigma}, \mathcal{U}_{\Lambda_{\sigma'}} \mid \mathring{\mathbf{\Psi}}) = 0 \quad (\sigma \neq \sigma') \quad (\textit{Proposition 12.76}).$$

Equivalently, the same outcome follows from the asymmetric visibility and the DE coherence score (Proposition 13.52) together with the HSP argmax in eq. (13.78). This restates dark–energy selection as a precise HSP outcome rather than a fine–tuned numerical cancellation (Section 13.2.7).

Remark 13.62. The Commutator Lensing Model (CLM) (Section 3.3) diagnoses the algebraic source of selection by quantifying the noncommutativity ("twist") between preparation and evolution, explaining the *cause* of bright/dark interferometric structure. However, the comparative judgment among distinct cosmological hypotheses requires the full holosymmetric interferometer and the physical overlap Ω_{phys} of eq. (12.83).

Corollary 13.63 (Holosymmetric Emergence of the Dark Sector)**.** *Let $\mathcal{H}^{(G)}_{\mathrm{phys}}$ be the BRST–physical Hilbert space for a dark gauge sector $G \subset \mathrm{Aut}(\mathcal{A})$ and $\mathcal{P}^{\mathrm{coh}}_G$ the holosymmetric coherence projector (Definition 12.77). A state $|\psi\rangle \in \mathcal{H}^{(G)}_{\mathrm{phys}}$ belongs to the dark sector iff:*

(i) Lensing/coherence
$\|\mathcal{P}^{\mathrm{coh}}_G|\psi\rangle\|^2 \neq 0$*;*

(ii) Spectral decoupling
$\langle\psi|\mathcal{O}_{\mathrm{SM}}|\psi\rangle_{\mathrm{phys}} \approx 0$ *for all* $\mathcal{O}_{\mathrm{SM}} \in \mathfrak{A}^{\mathrm{obs}}_{\mathrm{SM}}$.

Moreover, among all viable dark–sector histories, the realized one is the unique maximizer of $\Omega_{\mathrm{phys}}(G, G \mid \mathring{\mathbf{\Psi}})$ *subject to holosymmetric orthogonality* contra *all alternatives (Theorem 12.79 and Proposition 12.76).*

The First Condition: Lensing (Coherence) Criterion

Definition 13.64 (Stage I lensing/self–coherence)**.** A state is *physically realized* only if it survives the Holosymmetric Lensing filter, i.e., it has nonzero projection onto the coherent

subspace of its associated gauge sector G. With $\mathcal{P}_G^{\mathrm{coh}}$ the sectoral coherence projector (Definition 12.77), the Stage I requirement is

$$\left\| \mathcal{P}_G^{\mathrm{coh}} \, |\psi\rangle \right\|^2 \; > \; 0. \tag{13.85}$$

This excludes gauge artefacts and unstable transients: only vectors supported on the holosymmetry–coherent subspace of the BRST–physical Hilbert space $\mathcal{H}_{\mathrm{phys}}$ count as candidates for reality. In the language of the Holosymmetric Selection Principle (Section 12.2), (13.85) is the operational form of *self–coherence*: $\mathcal{P}_G^{\mathrm{coh}}$ filters out states that do not belong to a physically viable, internally consistent universal structure.

The Second Condition: Spectral Decoupling (Darkness)

Definition 13.65 (Spectral decoupling ("darkness"))**.** "Darkness'' is *spectral orthogonality* to the Standard Model (SM) observable algebra realized on the BRST–physical Hilbert space $\mathcal{H}_{\mathrm{phys}}$. Writing $\mathfrak{A}_{\mathrm{SM}}^{\mathrm{obs}}$ for the SM observable algebra (Section 1.4.1), the condition is

$$\left\langle \psi \right| \mathcal{O}_{\mathrm{SM}} \left| \psi \right\rangle_{\mathrm{phys}} \; \approx \; 0, \qquad \forall \, \mathcal{O}_{\mathrm{SM}} \in \mathfrak{A}_{\mathrm{SM}}^{\mathrm{obs}}. \tag{13.86}$$

Thus a dark–sector state $|\psi\rangle$—having passed the coherence test (13.85)—has negligible expectation against *all* SM probes: it is orthogonal to SM charges and operators at the emergent/GNS layer (the BRST–physical expectations are taken in the representation induced by the Quantum Holo–State $|\mathring{\mathbf{\Psi}}\rangle$, cf. Definition 1.27). A quantitative tolerance may be imposed as in Remark 13.58.

Remark 13.66 (Operational Definition of Dark-Sector Reality/Visibility)**.** Taken together, (13.85) and (13.86) supply a complete operational definition used in Corollary 13.63: a dark–sector entity is **real** (holosymmetry–coherent on $\mathcal{H}_{\mathrm{phys}}$) yet **not visible** (spectrally orthogonal to $\mathfrak{A}_{\mathrm{SM}}^{\mathrm{obs}}$). For mixed states, use the equivalent trace conditions in Definition 13.57; a quantitative tolerance $\varepsilon_{\mathrm{SM}}$ may be imposed as in Remark 13.58.

Dark energy. The interferometric selection (HSP, Stage I/II) singles out the slow-roll τ–scalar branch—the **Gaiaton/Apeiron**—as the realized mechanism of cosmic acceleration [486, 592]. Histories with negative or unstable residuals fail Stage I; among the survivors, Stage II prefers the branch whose dynamics keep $w \simeq -1$ with negligible small-scale modulation while matching the "residual of the dual-field vacuum" picture, consistent with SNe Ia discovery/confirmation and CMB and late-time constraints [134, 135, 600, 136].[1] In this sense, the effective cosmological constant is the bright-fringe residue of the Apeiron's potential rather than an arbitrary parameter. A coherent τ–background can mimic a geometric DE component and remains a useful diagnostic, but it is subordinate to (and not required by) the data-driven selection of the slow-roll Gaiaton branch.

[1] BAO and WL/growth anchors often cited in this context include *Eisenstein et al.* (SDSS BAO) and *Alam et al.* (BOSS DR12) for BAO, plus *DES Y3* and *KiDS-1000* for WL; [151, 152, 153, 154].

Dark matter. For particulate dark matter, the **Spin-1 Chrono–Kymon** (τ–vector) emerges as the dominant single-component candidate; its generic portals and constraints align with dark-photon/vector-DM phenomenology [599, 548, 548, 507, 210]. It passes Stage I (stability, TT-purity, portal bounds, negligible anisotropic stress in the FLRW average) and, in Stage II, yields a distinctive vector coherence-kernel cutoff ($k_c^{(V)}$) that separates it from scalar and internal-connection vectors; the absence of extra GW polarizations is consistent with current bounds [513]. A viable alternative is the **Spin-0 Gaiaton** in its oscillatory/misalignment regime: when $m_S \gg H$, it behaves CDM-like with a scalar Jeans-type cutoff ($k_c^{(S)}$), provided this branch is cleanly disentangled from the slow-roll DE regime [597].[2] The internal-connection vectors—**Mition** (Gaiamiti connection) and **Tantron** (Tantrakyma connection)—remain matter-like, TT-safe options under stringent portal and projectively-null mixing constraints; their small-scale and growth signatures are vector-like and assessed with the same kernel diagnostics. If data favor multiple components, a convex-mixture path is admitted with a parsimony penalty; otherwise the argmax collapses to a single dominant branch.

The dark sector selected by the interferometer is therefore: **DE ≈ slow-roll Gaiaton (Apeiron) residual** of the dual-field vacuum [486, 592, 600, 136]; **DM ≈ Chrono–Kymon–dominated**, with optional subdominant Gaiaton-oscillatory and/or connection-vector admixtures only when demanded by the meters (SNe/CMB/BAO/WL/growth) [134, 135, 153, 154, 152]. The **coherent τ–background** is interpreted as **dark gravity**—a geometric contribution useful diagnostically—but not as particulate dark matter [513].

13.3 The Dark Universe: Gaiatons and Chrono–Kymons

Scope and Placement

This section, to be read immediately after Section 13.1.4, develops the roles of the *dark universe* as realized through the τ–kairos registers: the scalar (Gaiaton/Apeiron) and vector (Chrono–Kymon) channels. The unified chronogeometric background and its fluctuations are split as:

$$\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}, \qquad \delta\hat{\Sigma}_{\mu\nu} = \hat{h}_{\mu\nu} + \hat{\tau}_{\mu\nu}, \tag{13.87}$$

with $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\hat{\tau}_{\mu\nu} = 0$ (TT–null; Definitions 10.3 and 10.13), and $\hat{\tau}_{\mu\nu}$ given by the full typed decomposition Definition 10.13 and eqs. (10.5) and (10.10). The Gaiaton governs slow-roll DE and oscillatory CDM branches (Proposition 10.153 and Definitions 10.68 and 10.69), the Chrono–Kymon realizes vector CDM with relational Proca/BEH dynamics (Proposition 10.154 and eqs. (11.339) to (11.341)), and a coherent τ–background provides a TT-safe "dark gravity" interpretation (eqs. (11.291) and (11.292), Definition 9.58, and Theorem 9.351).

Selection Map: Scalar vs. Vector Temporal Structure

In Ch. 10 the τ–kairos is defined by the **full typed decomposition**

$$\hat{\tau}_{\mu\nu} \;=\; S^{(b)}_{\mu\nu}\big[\hat{\varphi}_b\big] \;\oplus\; S^{(L)}_{\mu\nu}\big[\hat{\varphi}_L\big] \;\oplus\; \tilde{S}^{(p)}_{\mu\nu}\big[\tilde{\varphi}\big] \;\oplus\; \sum_{\lambda=\pm} A^{(\lambda)}_{\mu\nu}\big[\hat{K}^{(\tau)}_{\lambda}\big], \qquad \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\hat{\tau}_{\mu\nu} = 0, \tag{13.88}$$

[2] For small-scale cutoff phenomenology in ultra-light scalar DM, a standard reference is *Hu, Barkana & Gruzinov* (fuzzy DM); add [598] to draw that analogy explicitly.

as in Definition 10.13 and eq. (10.10), with the typed orthogonal resolution $S^{(i)}S^{(j)} = \delta_{ij}S^{(i)}$, $\sum_i S^{(i)} + A^{(+)} + A^{(-)} = \mathbf{1}_\tau$ (eq. (10.5)). Here $S^{(b)}_{\mu\nu}, S^{(L)}_{\mu\nu}, \tilde{S}^{(p)}_{\mu\nu}$ are the scalar/radial, longitudinal–scalar, and pseudo–scalar images of the τ–scalar register, while $A^{(\lambda)}_{\mu\nu}$ are the helicity $\lambda = \pm$ images of the τ–vector (Chrono–Kymon) register.

Operational identification within the typed weave.

(i) Scalar (Gaiaton/Apeiron) The physical scalar is the projector–certified Gaiaton obtained by the Riesz selectors (Definitions 10.68 and 10.69), i.e. a specific admissible combination within $\{S^{(b)}, S^{(L)}, \tilde{S}^{(p)}\}$, sourcing the vacuum–like background (DE) in its slow–roll branch and CDM–like behavior in its coherent–oscillation branch (Proposition 10.153).

(ii) Vector (Chrono–Kymon) The transverse vector excitations $A^{(\pm)}_{\mu\nu}[\hat{K}^{(\tau)}_{\pm}]$ furnish pressureless, clusterable DM on FLRW diagnostic realized reference configuration and satisfy the relational Proca/BEH system (Proposition 10.154 and eqs. (11.339) to (11.341)).

TT isolation and polarization projector. The spin–2 content resides entirely in the metric TT channel: $\mathcal{P}^{(2)}[\hat{\tau}_{\mu\nu}] = 0$ with $\mathcal{P}^{(2)}$ the typed spin–2 polarization projector (Definition 11.18); equivalently $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\hat{\tau}_{\mu\nu} = 0$ (Definition 10.3 and eq. (10.8)). This expresses the exact isolation of gravitational radiation in the geometric channel and the TT–purity of the weave.

These corrected identifications align this section with the canonical τ–kairos structure of Ch. 10 (Definition 10.13 and eqs. (10.5) and (10.10)) and with the polarization/BRST framework of Ch. 11 (Definitions 11.18, 11.47 and 11.48), while remaining compatible with the HSP machinery used for dark–sector selection (Section 13.1.4).

Dark Energy: Gaiaton (Apeiron) Background and Scalar Temporal Fluctuations

The cosmological acceleration is attributed to a small, *dressed* vacuum energy that survives a near–cancellation between the geometric (Gaiamiti) and material (Tantrakyma) zero–point contributions (Section 16.2.2). This mechanism is consistent with observations of late–time acceleration [134, 135, 136] and with effective–field arguments [211, 313]. The scalar tau–kairos excitation—the **Gaiaton**, also referred to in its field representation as the *Apeiron* φ—supplies the vacuum–like background responsible for dark energy:

$$\rho_\varphi = \tfrac{1}{2}\dot{\varphi}^2 + V(\varphi), \qquad p_\varphi = \tfrac{1}{2}\dot{\varphi}^2 - V(\varphi), \qquad w_\varphi = \frac{p_\varphi}{\rho_\varphi} \approx -1 \quad (\dot{\varphi}^2 \ll V). \tag{13.89}$$

Definition 13.67 (Apeiron/Gaiaton background fluid variables)**.** Work on a spatially homogeneous Friedmann–Lemaître–Robertson–Walker (FLRW) background with cosmic time t and metric signature $(-,+,+,+)$. Let $\varphi(t) \equiv \langle \mathring{\mathbf{\Psi}} | \, \hat{\varphi} \, | \mathring{\mathbf{\Psi}} \rangle_{\mathrm{phys}}$ be the Apeiron (Gaiaton, scalar tau–kairos) background—i.e. the BRST–physical GNS expectation of the scalar field operator. For the minimally coupled Lagrangian $\mathcal{L}_\varphi = \frac{1}{2}\,\partial_\mu\varphi\,\partial^\mu\varphi - V(\varphi)$ (natural units $c = \hbar = 1$), the energy–momentum tensor $T_{\mu\nu} = \partial_\mu\varphi\,\partial_\nu\varphi - g_{\mu\nu}\mathcal{L}_\varphi$ yields, in the comoving rest frame,

$$\rho_\varphi \equiv T^0{}_0(-) \;=\; \tfrac{1}{2}\dot{\varphi}^2 + V(\varphi), \qquad p_\varphi \equiv \tfrac{1}{3}T^i{}_i \;=\; \tfrac{1}{2}\dot{\varphi}^2 - V(\varphi),$$

and the equation–of–state parameter

$$w_\varphi \;\equiv\; \frac{p_\varphi}{\rho_\varphi} \approx -1 \quad \text{whenever} \quad \dot\varphi^2 \ll V(\varphi).$$

Here $\dot\varphi \equiv d\varphi/dt$ and $V(\varphi)$ is the (effective, lensing–renormalized) scalar potential used in this chapter's temporal sector. The holosymmetric selection mechanism (Section 13.1.4) then adjudicates among possible vacuum branches ($\Lambda > 0$, $\Lambda = 0$, $\Lambda < 0$), selecting the realized Gaiaton background responsible for cosmic acceleration.

Meaning of ρ_φ, p_φ, and w_φ. Here $\varphi(t) \equiv \langle \mathring{\boldsymbol{\Psi}} |\, \hat\varphi \,| \mathring{\boldsymbol{\Psi}} \rangle_{\rm phys}$ is the spatially homogeneous *Gaiaton* (Apeiron) background on an FLRW spacetime (cosmic time t, signature $(-,+,+,+)$). For the minimally coupled scalar with Lagrangian

$$\mathcal{L}_\varphi = \tfrac{1}{2}\,\partial_\mu\varphi\,\partial^\mu\varphi - V(\varphi),$$

the stress–energy tensor is

$$T_{\mu\nu} = \partial_\mu\varphi\,\partial_\nu\varphi - g_{\mu\nu}\Big(\tfrac{1}{2}\,\partial_\alpha\varphi\,\partial^\alpha\varphi - V(\varphi)\Big)\,.$$

For a spatially homogeneous background ($\partial_i\varphi = 0$), a comoving observer has four–velocity $u^\mu = (1,0,0,0)$ and the perfect–fluid identification

$$\rho_\varphi \equiv T_{\mu\nu}u^\mu u^\nu = T_{00}, \qquad p_\varphi \equiv \tfrac{1}{3}\,h^{\mu\nu}T_{\mu\nu} = \tfrac{1}{3}\,T^i{}_i,$$

where $h^{\mu\nu} = g^{\mu\nu} + u^\mu u^\nu$ projects orthogonally to u^μ. In components this reduces to the familiar fluid variables

$$\rho_\varphi = \tfrac{1}{2}\dot\varphi^2 + V(\varphi), \qquad p_\varphi = \tfrac{1}{2}\dot\varphi^2 - V(\varphi),$$

and the equation–of–state parameter

$$w_\varphi \;\equiv\; \frac{p_\varphi}{\rho_\varphi} \;=\; \frac{\tfrac{1}{2}\,\dot\varphi^2 - V(\varphi)}{\tfrac{1}{2}\,\dot\varphi^2 + V(\varphi)} \;=\; \frac{K - V}{K + V}, \qquad K \equiv \tfrac{1}{2}\,\dot\varphi^2,$$

where the overdot denotes d/dt (cosmic time). In the potential–dominated (slow–roll) regime $K \ll V$, one has $w_\varphi \simeq -1$ (dark–energy–like). For coherent oscillations about a quadratic minimum $V(\varphi) \simeq \tfrac{1}{2}m^2\varphi^2$, the cycle–averaged value is $\langle w_\varphi\rangle \simeq 0$ (matter–like), while for a quartic potential $V \propto \varphi^4$ one finds $\langle w_\varphi\rangle \simeq \tfrac{1}{3}$ (radiation–like). Cosmic acceleration requires $w_\varphi < -1/3$. *Sign note:* with the $(-,+,+,+)$ convention, $T^0{}_0 = -\rho_\varphi$ while $T_{00} = +\rho_\varphi$.

In this setting, the scalar piece of the tau–kairos tensor, $\hat\tau^{(0)}_{\mu\nu} = S_{\mu\nu}[\hat\varphi]$, is the *fluctuation channel* around the Gaiaton (Apeiron) background. It modulates the cosmological *coherence texture* of $t_{\mu\nu}$, influencing:

(i) Effective sound speed and clustering the effective sound speed c_s^2 and the clustering of dark–energy perturbations,

(ii) Linear growth of matter perturbations the linear–growth kernel for matter perturbations (via the growth observable $f\sigma_8$),

(iii) Hubble–friction renormalization small renormalizations of the Hubble–friction term through $\langle \hat{\tau}^{(0)} \rangle_{\mathrm{phys}}$.

All such effects preserve the TT isolation of gravitational waves and reduce to Λ–like behavior in the decohered limit $\hat{\tau}^{(0)} \to 0$. Global consistency with the Universal Quantum Constraint (Section 13.6) is maintained by treating the dark–energy background as a stationary–coherence, low–momentum solution in the scalar temporal sector.

Definition 13.68 (Gaiaton/Apeiron fluid on a general chronogeometric background)**.** Let $(\mathcal{M}, \bar{\Sigma}_{\mu\nu})$ be the emergent chronogeometric spacetime with inverse $\bar{\Sigma}^{\mu\nu}$ and relational Levi–Civita connection $\nabla^{(\mathrm{rel})}$ (cf. eq. (11.191)). Fix a unit timelike congruence u^μ obeying $\bar{\Sigma}_{\mu\nu} u^\mu u^\nu = -1$ and define the spatial projector $h_{\mu\nu} := \bar{\Sigma}_{\mu\nu} + u_\mu u_\nu$ and $h^\mu{}_\nu := \delta^\mu{}_\nu + u^\mu u_\nu$.

For the canonically normalized scalar tau–kairos field (Gaiaton/Apeiron) φ with

$$\mathcal{L}_\varphi \;:=\; \tfrac{1}{2}\,\bar{\Sigma}^{\mu\nu}\,\nabla^{(\mathrm{rel})}_\mu\varphi\,\nabla^{(\mathrm{rel})}_\nu\varphi \;-\; V(\varphi),$$

the stress tensor is

$$T^{(\varphi)}_{\mu\nu} \;:=\; \nabla^{(\mathrm{rel})}_\mu\varphi\,\nabla^{(\mathrm{rel})}_\nu\varphi \;-\; \bar{\Sigma}_{\mu\nu}\,\mathcal{L}_\varphi.$$

Introduce the $1+3$ split of the gradient

$$\dot{\varphi} \;:=\; u^\mu\nabla^{(\mathrm{rel})}_\mu\varphi, \qquad D_\mu\varphi \;:=\; h_\mu{}^\nu\nabla^{(\mathrm{rel})}_\nu\varphi, \qquad D^2\varphi \;:=\; h^{\mu\nu}D_\mu\varphi D_\nu\varphi.$$

Then the covariant fluid variables *relative to* u^μ are

$$\rho_\varphi := T^{(\varphi)}_{\mu\nu}u^\mu u^\nu = \tfrac{1}{2}\,\dot{\varphi}^2 + \tfrac{1}{2}\,D^2\varphi + V(\varphi), \tag{13.90}$$

$$p_\varphi := \tfrac{1}{3}\,h^{\mu\nu}T^{(\varphi)}_{\mu\nu} = \tfrac{1}{2}\,\dot{\varphi}^2 - \tfrac{1}{6}\,D^2\varphi - V(\varphi), \tag{13.91}$$

$$q_\mu := -\,h_\mu{}^\alpha T^{(\varphi)}_{\alpha\beta}u^\beta = -\,\dot{\varphi}\,D_\mu\varphi, \tag{13.92}$$

$$\pi_{\mu\nu} := \Big(T^{(\varphi)}_{\alpha\beta}h^\alpha{}_{\langle\mu}h^\beta{}_{\nu\rangle}\Big) = D_{\langle\mu}\varphi\,D_{\nu\rangle}\varphi, \tag{13.93}$$

where $\langle\cdots\rangle$ denotes the projected, symmetric, tracefree part with respect to $h_{\mu\nu}$. The Gaiaton field equation is the relational Klein–Gordon equation

$$\Box^{(\mathrm{rel})}\varphi - V_{,\varphi} = 0, \qquad \Box^{(\mathrm{rel})} := \bar{\Sigma}^{\mu\nu}\nabla^{(\mathrm{rel})}_\mu\nabla^{(\mathrm{rel})}_\nu. \tag{13.94}$$

If nonminimal tau–kairos couplings are absent, then $\nabla^{(\mathrm{rel})}_\mu T^{(\varphi)\mu}{}_\nu = 0$; otherwise, source terms appear in the total conservation law and are moved to the right–hand side of the chronogeometric Einstein equations (cf. eq. (11.191)).

Proposition 13.69 (FLRW/comoving reduction)**.** *If $(\mathcal{M}, \bar{\Sigma}_{\mu\nu})$ admits an FLRW slicing with u^μ the comoving congruence of φ (i.e. $D_\mu\varphi = 0$), then*

$$\rho_\varphi = \tfrac{1}{2}\dot{\varphi}^2 + V, \qquad p_\varphi = \tfrac{1}{2}\dot{\varphi}^2 - V, \qquad q_\mu = 0, \qquad \pi_{\mu\nu} = 0,$$

and $w_\varphi := p_\varphi/\rho_\varphi = \frac{K-V}{K+V}$ with $K := \frac{1}{2}\dot{\varphi}^2$, agreeing with eq. (13.89). In the slow–roll regime $K \ll V$, one has $w_\varphi \simeq -1$.

Remark 13.70 (Kinematics and gauge–invariant evolution)**.** For a general congruence u^μ on $(\mathcal{M}, \bar{\Sigma}_{\mu\nu})$, define the expansion, shear, vorticity, and 4–acceleration by

$$\theta := \nabla^{(\mathrm{rel})}_\mu u^\mu, \quad \sigma_{\mu\nu} := D_{\langle\mu} u_{\nu\rangle}, \quad \omega_{\mu\nu} := D_{[\mu} u_{\nu]}, \quad a_\mu := u^\alpha \nabla^{(\mathrm{rel})}_\alpha u_\mu.$$

Then the energy and momentum balance for the Gaiaton fluid read

$$\dot{\rho}_\varphi + (\rho_\varphi + p_\varphi)\,\theta + D^\mu q_\mu + 2\,a^\mu q_\mu + \sigma^{\mu\nu}\pi_{\mu\nu} = 0, \tag{13.95}$$

$$(\rho_\varphi + p_\varphi)\,a_\mu + D_\mu p_\varphi + \dot{q}_\mu + \tfrac{4}{3}\theta q_\mu + D^\nu \pi_{\mu\nu} + \sigma_\mu{}^\nu q_\nu + \omega_\mu{}^\nu q_\nu = 0, \tag{13.96}$$

with all derivatives taken using $\nabla^{(\mathrm{rel})}$. For the canonical scalar, the rest–frame sound speed is $c_s^2 = 1$; effective c_s^2 entering large–scale observables may be modified by lensing/renormalization of $V(\varphi)$ and by the choice of u^μ (frame effects).

Remark 13.71 (Placement within the tau–kairos sector)**.** The scalar piece of the tau–kairos tensor is $\hat{\tau}^{(0)}_{\mu\nu} = S_{\mu\nu}[\hat{\varphi}]$ (cf. eq. (13.88)), whose expectation $S_{\mu\nu}[\varphi]$ provides the scalar temporal contribution directly to the field equations. Inhomogeneous Gaiaton configurations ($D_\mu\varphi \neq 0$) generate nonzero q_μ and $\pi_{\mu\nu}$ while preserving $\mathcal{P}^{(2)}[\hat{\tau}_{\mu\nu}] = 0$, i.e. the exact isolation of TT gravitational radiation in $\hat{h}_{\mu\nu}$.

Definition 13.72 (Chrono–Kymon (vector tau–kairos) fluid on a general chronogeometric background)**.** Let $(\mathcal{M}, \bar{\Sigma}_{\mu\nu})$ be the emergent chronogeometric spacetime with inverse $\bar{\Sigma}^{\mu\nu}$ and relational Levi–Civita connection $\nabla^{(\mathrm{rel})}$ (cf. eq. (11.191)). Fix a unit timelike congruence u^μ with $\bar{\Sigma}_{\mu\nu}u^\mu u^\nu = -1$, spatial projector $h_{\mu\nu} := \bar{\Sigma}_{\mu\nu} + u_\mu u_\nu$ and $h^\mu{}_\nu := \delta^\mu{}_\nu + u^\mu u_\nu$.

The vector tau–kairos field (Chrono–Kymon) $K^{(\tau)}_\mu$ has Proca-type Lagrangian

$$\mathcal{L}_K \;:=\; -\tfrac{1}{4}\,\mathcal{H}_{\mu\nu}\mathcal{H}^{\mu\nu} \;+\; \tfrac{1}{2}\,\mu_C^2\,K_\mu K^\mu, \qquad \mathcal{H}_{\mu\nu} := 2\,\nabla^{(\mathrm{rel})}_{[\mu} K_{\nu]},$$

where indices are raised with $\bar{\Sigma}^{\mu\nu}$ and μ_C is the vector mass (possibly lensing-renormalized). The stress–energy tensor is

$$T^{(K)}_{\mu\nu} \;:=\; \mathcal{H}_{\mu\alpha}\mathcal{H}_\nu{}^\alpha - \tfrac{1}{4}\,\bar{\Sigma}_{\mu\nu}\,\mathcal{H}_{\alpha\beta}\mathcal{H}^{\alpha\beta} \;+\; \mu_C^2\Big(K_\mu K_\nu - \tfrac{1}{2}\,\bar{\Sigma}_{\mu\nu}K_\alpha K^\alpha\Big). \tag{13.97}$$

Decompose K_μ and $\mathcal{H}_{\mu\nu}$ relative to u^μ:

$$\phi := -\,u^\alpha K_\alpha, \qquad \mathcal{K}_\mu := h_\mu{}^\alpha K_\alpha, \qquad E_\mu := \mathcal{H}_{\mu\nu}u^\nu, \qquad B_\mu := \tfrac{1}{2}\,\varepsilon_{\mu\nu\alpha\beta}\,u^\nu\,\mathcal{H}^{\alpha\beta},$$

with $u^\mu\mathcal{K}_\mu = 0 = u^\mu E_\mu = u^\mu B_\mu$ and $\varepsilon_{\mu\nu\alpha\beta}$ the volume form of $\bar{\Sigma}_{\mu\nu}$. Define $E^2 := E_\mu E^\mu$, $B^2 := B_\mu B^\mu$, $\mathcal{K}^2 := \mathcal{K}_\mu\mathcal{K}^\mu$.

The covariant 1+3 fluid variables *relative to* u^μ are obtained by projection:

$$\rho_K := T^{(K)}_{\mu\nu}u^\mu u^\nu = \tfrac{1}{2}\,(E^2 + B^2) \;+\; \tfrac{1}{2}\,\mu_C^2\Big(\phi^2 + \mathcal{K}^2\Big), \tag{13.98}$$

$$p_K := \tfrac{1}{3}\,h^{\mu\nu}T^{(K)}_{\mu\nu} = \tfrac{1}{6}\,(E^2 + B^2) \;+\; \tfrac{1}{2}\,\mu_C^2\phi^2 \;-\; \tfrac{1}{6}\,\mu_C^2\mathcal{K}^2, \tag{13.99}$$

$$q^{(K)}_\mu := -\,h_\mu{}^\alpha T^{(K)}_{\alpha\beta}u^\beta = \varepsilon_{\mu\nu\lambda}\,E^\nu B^\lambda \;+\; \mu_C^2\,\phi\,\mathcal{K}_\mu, \tag{13.100}$$

$$\pi^{(K)}_{\mu\nu} := \Big(T^{(K)}_{\alpha\beta}h^\alpha{}_{\langle\mu}h^\beta{}_{\nu\rangle}\Big) = -\Big(E_{\langle\mu}E_{\nu\rangle} + B_{\langle\mu}B_{\nu\rangle}\Big) \;+\; \mu_C^2\,\mathcal{K}_{\langle\mu}\mathcal{K}_{\nu\rangle}, \tag{13.101}$$

where $\varepsilon_{\mu\nu\lambda} := \varepsilon_{\mu\nu\lambda\alpha}u^\alpha$ is the spatial Levi–Civita tensor and $\langle\cdot\rangle$ denotes the projected, symmetric, tracefree part w.r.t. $h_{\mu\nu}$.

The Chrono–Kymon field equations are the relational Proca equations

$$\nabla^{(\mathrm{rel})}_\nu \mathcal{H}^{\nu\mu} + \mu_C^2 K^\mu = 0, \qquad \Rightarrow \qquad \mu_C^2\, \nabla^{(\mathrm{rel})}_\mu K^\mu = 0 \ \ (\mu_C \neq 0), \tag{13.102}$$

and, in the absence of explicit nonminimal couplings, $\nabla^{(\mathrm{rel})}_\mu T^{(K)\mu}{}_\nu = 0$. The vector contribution to the tau–kairos tensor is $A_{\mu\nu}[K^{(\tau)}]$ as in eq. (13.88), and by construction $\mathcal{P}^{(2)}[\hat{\tau}_{\mu\nu}] = 0$ (TT isolation of gravitational radiation).

Proposition 13.73 (FLRW/comoving and isotropy enforcement)**.** *If $(\mathcal{M}, \bar{\Sigma}_{\mu\nu})$ admits an FLRW slicing with comoving u^μ, then enforcing statistical isotropy (e.g. by a triad of mutually orthogonal equal-magnitude vectors or by domain averaging) yields $q^{(K)}_\mu = 0 = \pi^{(K)}_{\mu\nu}$ at the background level. In the rapid-oscillation regime $\mu_C \gg H$ with bounded $\mathcal{K}_\mu$, the cycle-averaged equation of state approaches $\langle w_K\rangle := \langle p_K/\rho_K\rangle \simeq 0$ and $\langle\rho_K\rangle \propto a^{-3}$, so the Chrono–Kymon behaves as cold dark matter.*

Remark 13.74 (Kinematics and 1+3 balances)**.** With the standard kinematics of u^μ on $(\mathcal{M}, \bar{\Sigma}_{\mu\nu})$, $\theta := \nabla^{(\mathrm{rel})}_\mu u^\mu$, $\sigma_{\mu\nu} := D_{\langle\mu}u_{\nu\rangle}$, $\omega_{\mu\nu} := D_{[\mu}u_{\nu]}$, $a_\mu := u^\alpha\nabla^{(\mathrm{rel})}_\alpha u_\mu$, the Chrono–Kymon energy and momentum balances read

$$\dot{\rho}_K + (\rho_K + p_K)\theta + D^\mu q^{(K)}_\mu + 2a^\mu q^{(K)}_\mu + \sigma^{\mu\nu}\pi^{(K)}_{\mu\nu} = 0,$$

$$(\rho_K + p_K)a_\mu + D_\mu p_K + \dot{q}^{(K)}_\mu + \tfrac{4}{3}\theta q^{(K)}_\mu + D^\nu\pi^{(K)}_{\mu\nu} + \sigma_\mu{}^\nu q^{(K)}_\nu + \omega_\mu{}^\nu q^{(K)}_\nu = 0,$$

with all derivatives taken using $\nabla^{(\mathrm{rel})}$. These formulas apply off-FLRW, with anisotropy encoded in $\pi^{(K)}_{\mu\nu}$ and energy transport in $q^{(K)}_\mu$.

Remark 13.75 (Placement within the tau–kairos sector and HSP)**.** The Chrono–Kymon is the vector register of the tau–kairos tensor (Proposition 10.154 and eq. (13.88)). Its expectation contributes via $A_{\mu\nu}[K^{(\tau)}]$ directly to the chronogeometric field equations (eq. (11.191)). Admissibility and realization are determined by the Holosymmetric Selection Principle on the BRST–physical space (Section 13.1.4 and Definition 13.1), ensuring spectral orthogonality to SM observables and maximizing holosymmetric coherence.

Dark Matter: Massive Vector Chrono–Kymons and Temporal Halos

Definition 13.76 (Massive vector Chrono–Kymon as a dark–matter sector)**.** The Tantrakyma vector excitation $K^{(\tau)}_\mu$ (the *Chrono–Kymon*) is a neutral, clusterable dark–matter candidate selected by the Holosymmetric Selection Principle (HSP). Its leading cosmological dynamics on the emergent chronogeometric background $(\mathcal{M}, \bar{\Sigma}_{\mu\nu})$ are captured by a Proca–type effective theory,

$$\mathcal{L}_{K^{(\tau)}} = -\frac{1}{4}\,\mathcal{H}_{\mu\nu}\mathcal{H}^{\mu\nu} + \frac{1}{2}\,\mu_C^2\, K^{(\tau)}_\mu K^{(\tau)\mu} + \mathcal{L}_{\mathrm{int}}[\bar{\Sigma}, K^{(\tau)}], \qquad \mathcal{H}_{\mu\nu} := 2\,\nabla^{(\mathrm{rel})}_{[\mu} K^{(\tau)}_{\nu]}, \tag{13.103}$$

where indices are raised with $\bar{\Sigma}^{\mu\nu}$ and $\nabla^{(\mathrm{rel})}$ is the relational Levi–Civita connection. Curvature/state–dependent couplings are encoded in $\mathcal{L}_{\mathrm{int}}$ and chosen to preserve TT isolation.

Fix a unit timelike congruence u^μ ($\bar{\Sigma}_{\mu\nu}u^\mu u^\nu = -1$) with expansion $\theta := \nabla^{(\mathrm{rel})}_\mu u^\mu$, shear $\sigma_{\mu\nu}$, vorticity $\omega_{\mu\nu}$, and 4–acceleration a_μ. In the *adiabatic rapid–oscillation regime*

$$\mu_C \;\gg\; \max\Big\{\, |\theta|,\, \|\sigma\|,\, \|\omega\|,\, \|a\| \,\Big\}, \tag{13.104}$$

and upon coarse–graining over many oscillations along u^μ, the vector tau–kairos medium obeys (cf. the 1+3 balances in Remark 13.74)

$$\langle w_{K^{(\tau)}}\rangle \;\simeq\; 0, \qquad \langle \dot{\rho}_{K^{(\tau)}}\rangle \;+\; \theta\,\langle \rho_{K^{(\tau)}}\rangle \;\approx\; -\,D^\mu\langle q^{(K)}_\mu\rangle \;-\; 2a^\mu\langle q^{(K)}_\mu\rangle \;-\; \sigma^{\mu\nu}\langle \pi^{(K)}_{\mu\nu}\rangle. \tag{13.105}$$

For comoving, (statistically) isotropic configurations (so that $\langle q^{(K)}_\mu\rangle = \langle \pi^{(K)}_{\mu\nu}\rangle = 0$ at the background level), (13.105) reduces to

$$\langle w_{K^{(\tau)}}\rangle \simeq 0, \qquad \langle \dot{\rho}_{K^{(\tau)}}\rangle + \theta\,\langle \rho_{K^{(\tau)}}\rangle \simeq 0, \quad \Rightarrow \quad \langle \rho_{K^{(\tau)}}\rangle \;\propto\; \exp\!\Big(-\int \theta\, dt\Big). \tag{13.106}$$

Thus the Chrono–Kymon behaves as *cold* dark matter on general backgrounds, while remaining spectrally decoupled from SM observables (cf. Definitions 13.57 and 13.65). Coherent configurations of the temporal sector build long–range, EM–dark *temporal halos* that shape rotation curves, lensing maps, and satellite kinematics without introducing additional TT gravitational–wave polarizations.

Proposition 13.77 (FLRW specialization)**.** *On an FLRW slicing with comoving u^μ, one has $\theta = 3H$ and $\langle q^{(K)}_\mu\rangle = \langle \pi^{(K)}_{\mu\nu}\rangle = 0$. Then (13.106) gives $\langle \dot{\rho}_{K^{(\tau)}}\rangle + 3H\,\langle \rho_{K^{(\tau)}}\rangle \simeq 0$, so*

$$\langle w_{K^{(\tau)}}\rangle \simeq 0, \qquad \langle \rho_{K^{(\tau)}}\rangle \propto a^{-3},$$

reproducing (10.154) with the identification $H = \dot{a}/a$.

Explanation of terms and notation.

$K^{(\tau)}_\mu$ vector *tau–kairos* degree of freedom (Chrono–Kymon), neutral by spectral decoupling.

$\nabla^{(\mathrm{rel})}$, $\bar{\Sigma}_{\mu\nu}$ relational Levi–Civita connection and unified background tensor.

$\theta, \sigma_{\mu\nu}, \omega_{\mu\nu}, a_\mu$ kinematic invariants of the congruence u^μ.

$\mu_C \gg \max\{|\theta|, \|\sigma\|, \|\omega\|, \|a\|\}$ adiabaticity condition ensuring the cold limit.

$q^{(K)}_\mu$, $\pi^{(K)}_{\mu\nu}$ energy flux and anisotropic stress of the Chrono–Kymon fluid (see Definition 13.72); vanishing at background level for comoving/isotropic configurations (e.g. triads or domain averaging).

FLRW case $H = \theta/3$, $a(t)$ the scale factor $\Rightarrow$ $\rho \propto a^{-3}$.

Unified Dynamics and Mixing: Matrix Propagator in Cosmology

Definition 13.78 (Cosmological matrix propagator and TT isolation (background–agnostic))**.** Let $(\mathcal{M}, \bar{\Sigma}_{\mu\nu})$ be the emergent chronogeometric background with relational Levi–Civita connection $\nabla^{(\mathrm{rel})}$ and determinant $\sqrt{-\bar{\Sigma}}$. Linearized fluctuations of the unified tensor are organized as $\hat{\Upsilon}_{\mu\nu} := (\hat{h}_{\mu\nu}, \hat{\tau}_{\mu\nu})^{\top}$, where $\hat{h}_{\mu\nu} = \delta\hat{g}_{\mu\nu}$ and $\hat{\tau}_{\mu\nu} = \delta\hat{t}_{\mu\nu}$. To quadratic order about $\bar{\Sigma}_{\mu\nu}$ one may write

$$S^{(2)} = \frac{1}{2}\int d^4x\,\sqrt{-\bar{\Sigma}}\;\hat{\Upsilon}^{\mu\nu}\begin{pmatrix} \mathcal{D}_{hh}[\bar{\Sigma},\nabla^{(\mathrm{rel})}] & \mathcal{M}_{h\tau}[\bar{\Sigma},\nabla^{(\mathrm{rel})}] \\ \mathcal{M}_{\tau h}[\bar{\Sigma},\nabla^{(\mathrm{rel})}] & \mathcal{D}_{\tau\tau}[\bar{\Sigma},\nabla^{(\mathrm{rel})}] \end{pmatrix}_{\mu\nu}{}^{\rho\sigma}\hat{\Upsilon}_{\rho\sigma}, \qquad \mathcal{P}^{(2)}\circ\mathcal{M}_{\tau h} = 0 = \mathcal{M}_{h\tau}\circ\mathcal{P}^{(2)}. \tag{13.107}$$

Here $\mathcal{D}_{hh}$, $\mathcal{D}_{\tau\tau}$ are self–adjoint kinetic operators (including curvature/kinematic "mass/friction" terms built from $\bar{\Sigma}_{\mu\nu}$ and its invariants), and $\mathcal{M}_{h\tau}$, $\mathcal{M}_{\tau h}$ are mixing kernels. The projector $\mathcal{P}^{(2)}$ maps symmetric tensors to their transverse–traceless (TT) part *with respect to a chosen unit timelike congruence* u^{μ}, i.e. on spatial slices ($h_{\mu\nu} := \bar{\Sigma}_{\mu\nu} + u_\mu u_\nu$),

$$\mathcal{P}^{(2)}[X_{\mu\nu}] \text{ is symmetric}, \quad h^{\mu\nu}\mathcal{P}^{(2)}[X_{\mu\nu}] = 0, \quad D^{\mu}\mathcal{P}^{(2)}[X_{\mu\nu}] = 0,$$

with D_μ the spatial covariant derivative induced by $\nabla^{(\mathrm{rel})}$. Because the temporal sector has no TT content (cf. $\mathcal{P}^{(2)}[\hat{\tau}_{\mu\nu}] = 0$ in eq. (13.88)), TT isolation follows: the radiative tensor sector propagates purely in the geometric block G_{hh} (components of the inverse operator), while spatial–temporal conversion amplitudes are projector–suppressed:

$$\mathcal{P}^{(2)}\,G_{h\tau} = 0, \qquad G_{\tau h}\,\mathcal{P}^{(2)} = 0.$$

Cosmological observables thus respond to temporal structure *through the background and its scalar/vector temporal perturbations* (scalar Apeiron/Gaiaton $\hat{\varphi}_A$, vector Chrono–Kymon $\hat{K}^{(\tau)}_{\mu}$), not via new GW polarizations.

Proposition 13.79 (FLRW specialization)**.** *On an FLRW slicing with comoving* u^{μ} *and* $\bar{\Sigma}_{\mu\nu} \to \mathrm{diag}(-1, a^2\gamma_{ij})$, (13.107) *reduces to the quadratic action about FLRW quoted earlier, with* $\sqrt{-\bar{\Sigma}} = a^3\sqrt{\gamma}$ *and* $\mathcal{P}^{(2)}$ *the usual spatial TT projector.*

Explanation of terms and notation.

$\hat{h}_{\mu\nu}$ geometric (spin–2) metric perturbations (the g–block of $\hat{\Sigma}_{\mu\nu}$).

$\hat{\tau}_{\mu\nu}$ temporal–sector perturbations (scalar $\hat{\tau}^{(0)}$ from the Gaiaton/Apeiron, and vector from the Chrono–Kymon; no TT part).

$\bar{\Sigma}_{\mu\nu}$ background unified tensor; $\sqrt{-\bar{\Sigma}} = \sqrt{-\det\bar{\Sigma}_{\mu\nu}}$.

$\mathcal{D}_{hh}$, $\mathcal{D}_{\tau\tau}$ background–dependent kinetic operators (contain Lichnerowicz–type operators, kinematic invariants $\theta, \sigma_{\mu\nu}, \omega_{\mu\nu}, a_\mu$ via $\nabla^{(\mathrm{rel})}$).

$\mathcal{M}_{h\tau}$, $\mathcal{M}_{\tau h}$ mixing kernels; TT isolation imposes $\mathcal{P}^{(2)}\circ\mathcal{M}_{\tau h} = 0 = \mathcal{M}_{h\tau}\circ\mathcal{P}^{(2)}$.

$G_{hh}, G_{h\tau}, G_{\tau h}, G_{\tau\tau}$ propagator blocks (inverse of the operator matrix in (13.107)); TT purity implies $\mathcal{P}^{(2)}G_{h\tau} = 0 = G_{\tau h}\mathcal{P}^{(2)}$.

Observational Program and Cross–Checks

The temporal (tau–kairos) sector suggests a paired suite of tests, formulated on general chronogeometric backgrounds and reducing smoothly to the FLRW case (cf. Definitions 13.68, 13.72 and 13.78):

(i) Dark energy (scalar temporal, Gaiaton/Apeiron) Constraints on $w(z)$, the sound speed c_s^2, and the growth index $f\sigma_8$ probe $\hat{\tau}^{(0)}$–induced departures from Λ (via the scalar tau–kairos fluctuation channel around the Gaiaton background; cf. Section 13.3 and Definition 13.68). Null limits recover ΛCDM [134, 135, 136].

(ii) Dark matter (vector temporal, Chrono–Kymon) Weak lensing shear maps, halo mass concentration relations, satellite counts, and inner–slope/coring diagnostics probe $K_\mu^{(\tau)}$ coherence and the emergence of temporal halos (cf. Definitions 13.72 and 13.76). On large scales, the FLRW specialization of the general balance (13.106) reproduces CDM background scaling (Proposition 10.154); on small scales, vector coherence allows mild, observationally bounded deviations (e.g. core formation) [588, 311].

(iii) GW channel integrity The constraints $\mathcal{P}^{(2)}[\hat{\tau}] = 0$ and $\mathcal{P}^{(2)}\mathcal{M}_{\tau h} = 0$ in the matrix action (13.107) enforce *no extra GW polarizations*. TT purity implies $\mathcal{P}^{(2)}G_{h\tau} = 0 = G_{\tau h}\mathcal{P}^{(2)}$, yielding a sharp, falsifiable prediction consistent with current bounds.

Consistency and Decoupling Limits

Two limits guarantee compatibility with standard cosmology on general backgrounds:

(i) Decohered macroscopic limit $\hat{\tau}_{\mu\nu} \to 0$ and $\mathcal{M}_{h\tau} \to 0$ imply $\bar{\Sigma}_{\mu\nu} \to \bar{g}_{\mu\nu}$, yielding GR with ΛCDM at current precision.

(ii) Smooth–vacuum limit for DE (scalar tau–kairos) $\dot{\varphi}^2 \ll V$ with long coherence length (in the sense of Definition 13.68) gives $w_\varphi \simeq -1$ and negligible DE clustering; the FLRW reduction recovers (13.89).

Synthesis: A Unified, Relational Dark Sector

Dark energy is the *Apeiron/Gaiaton*–dominated background of the temporal sector, with the *scalar tau–kairos* piece as its fluctuation channel; dark matter is naturally furnished by *vector* temporal quanta—the Chrono–Kymons—whose large–scale, electromagnetically inert coherence forms halos. Both act through the unified tensor $\Sigma_{\mu\nu}$ (cf. Section 13.1) and preserve the TT gravitational–wave channel (cf. Definition 13.78), emerging as complementary faces of the same relational vacuum. This completes the *Holosymmetric Selection* narrative for the dark sector (Section 13.1.4 and Definition 13.1) and ties the cosmological analysis back to the Universal Quantum Constraint (Section 13.6), in direct analogy with the black–hole completion developed earlier.

13.4 An Effective Symmetry Stack for the Dark Sector

In the relational framework, the dark sector is not governed by a single fundamental gauge group in the sense of the Standard Model $G_{\rm SM} = SU(3)_C \times SU(2)_L \times U(1)_Y$, but by an *effective symmetry stack* acting on the *temporal* (τ–kairos) registers of the unified chronogeometric tensor $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$. The stack organizes the τ–kairos *types* and their approximate invariances; it is to be read together with the typed projectors, TT–null identities, and HSP rules.

$$\boxed{\begin{aligned}\mathcal{G}^{(\rm eff)}_{\rm Dark} = & \underbrace{\mathrm{Shift}_{\varphi_b} \times \mathrm{Shift}_{\varphi_L} \times \mathrm{Shift}^{\rm ax}_{\tilde{\varphi}}}_{\text{scalar/pseudoscalar } \tau\text{–kairos (Apeiron \& pseudo-Apeiron)}} \\ & \times \underbrace{U(1)_\tau \times U(1)_{\rm hel}}_{\tau\text{–vector phase \& helicity}} \\ & \times \underbrace{\mathbb{Z}_2^{\rm HS}}_{\text{stability / HSP sector tag}}\end{aligned}} \tag{13.108}$$

where: (i) $\mathrm{Shift}_{\varphi_b}, \mathrm{Shift}_{\varphi_L}$ are the approximate constant shifts of the breathing and longitudinal scalars, (ii) $\mathrm{Shift}^{\rm ax}_{\tilde{\varphi}}$ is the axial (pseudoscalar) shift, (iii) $U(1)_\tau$ acts as the *τ–vector (Chrono–Kymon) phase* on the vector register, and (iv) $U(1)_{\rm hel}$ rotates the helicity doublet $(K^{(\tau)}_+, K^{(\tau)}_-)$ in the transverse plane (helicity–plane rotations $\simeq SO(2)_{\rm hel} \cong U(1)_{\rm hel}$). All continuous factors are *effective* and can be softly broken by the same operators that realize cosmology (e.g. mass terms, mixing kernels); $\mathbb{Z}_2^{\rm HS}$ labels HSP branch stability. This stack *organizes* the τ–kairos (not the SM) and supplies the labels used by HSP Stage II.

Placement and TT purity

$$\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}, \tag{13.109}$$

$$\delta\hat{\Sigma}_{\mu\nu} = \delta\hat{g}_{\mu\nu} + \delta\hat{t}_{\mu\nu} = \hat{h}_{\mu\nu} + \hat{\tau}_{\mu\nu}, \tag{13.110}$$

$$\hat{h}_{\mu\nu} := \Pi^{[\rm TT]}_{\bar{\Sigma}}\, \delta\hat{g}_{\mu\nu}, \tag{13.111}$$

$$\Pi^{[\rm TT]}_{\bar{\Sigma}} \hat{\tau}_{\mu\nu} = 0. \tag{13.112}$$

Thus all spin–2 radiative content resides in the geometric TT graviton $\hat{h}_{\mu\nu}$ (Gaiamiti), and the τ–kairos carries only *non–TT* chronogeometric content.

Full τ–kairos decomposition

$$\hat{\tau}_{\mu\nu} = S^{(b)}_{\mu\nu}\big[\hat{\varphi}_b\big] \;\oplus\; S^{(L)}_{\mu\nu}\big[\hat{\varphi}_L\big] \;\oplus\; \tilde{S}^{(p)}_{\mu\nu}\big[\tilde{\varphi}\big] \;\oplus\; \sum_{\lambda=\pm} A^{(\lambda)}_{\mu\nu}\big[\hat{K}^{(\tau)}_\lambda\big], \tag{13.113}$$

with associated CK field strength

$$\hat{\mathcal{F}}^{(\tau)}_{\mu\nu} := 2\,\bar{\nabla}_{[\mu}\hat{K}^{(\tau)}_{\nu]}\,, \tag{13.114}$$

and typed TT–orthogonality

$$\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\,\Pi^{[\chi]} = 0 \quad \text{for all non–TT types } [\chi] \in \{b, L, p, \lambda = \pm\}. \tag{13.115}$$

Action of the effective symmetry factors

$$\mathrm{Shift}_{\varphi_b}:\ \hat{\varphi}_b \mapsto \hat{\varphi}_b + c_b, \qquad \mathrm{Shift}_{\varphi_L}:\ \hat{\varphi}_L \mapsto \hat{\varphi}_L + c_L, \tag{13.116}$$

$$\mathrm{Shift}_{\tilde{\varphi}}^{\mathrm{ax}}:\ \tilde{\varphi} \mapsto \tilde{\varphi} + \tilde{c}, \qquad U(1)_\tau:\ \hat{K}_\lambda^{(\tau)} \mapsto e^{i\alpha_\tau}\hat{K}_\lambda^{(\tau)}, \tag{13.117}$$

$$U(1)_{\mathrm{hel}}:\ \hat{K}_\lambda^{(\tau)} \mapsto e^{i\lambda\,\alpha_{\mathrm{hel}}}\hat{K}_\lambda^{(\tau)}, \qquad \mathbb{Z}_2^{\mathrm{HS}}:\ \text{sector/branch tag preserved by HSP Stage I/II.} \tag{13.118}$$

These are *approximate* symmetries of the τ–kairos ledger; soft breakings (e.g. Proca mass for CK, scalar potentials) are permitted and are tracked by the weave's diagnostics and HSP scoring. The stack (13.108) organizes the τ–kairos registers (not G_{SM}), is consistent with TT graviton purity via (13.112)–(13.115), and is to be used together with the typed propagator structure and cosmological matrix–propagator analysis (Section 13.3), as well as the Holosymmetric Selection Principle (Section 13.1.4). The scalar channels are identified with the Gaiaton (Apeiron) background and its temporal modulations; the vector channels with the Chrono–Kymon carrier sector. All TT content remains geometric ($\hat{h}_{\mu\nu} = \Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\delta\hat{g}_{\mu\nu}$).

Action on Fields and Allowed Couplings

Breathing scalar $\mathrm{Shift}_{\varphi_b}$ **(softly broken).** $\hat{\varphi}_b \mapsto \hat{\varphi}_b + c_b$. Admits derivative couplings and the breathing projector $S_{\mu\nu}^{(b)}[\hat{\varphi}_b]$ in (13.113); soft breaking via $V_b(\varphi_b)$ controls DE/DM backgrounds while preserving TT–null identities (13.112)–(13.115). (Consistent with the York TT projector and τ–nullity in Ch. 10: eqs. (10.4) and (10.105) and Corollary 10.177; see also the graviton definition Definition 10.25.) Mixings that violate $\mathrm{Shift}_{\varphi_b}$ are suppressed (diagnostic realized reference configuration hygiene/HSP).

Longitudinal scalar $\mathrm{Shift}_{\varphi_L}$ **(softly broken).** $\hat{\varphi}_L \mapsto \hat{\varphi}_L + c_L$. Selects derivative structures (e.g. longitudinal projector $S_{\mu\nu}^{(L)}[\hat{\varphi}_L]$) and forbids potential–induced tadpoles unless softly broken by $V_L(\varphi_L)$. Couplings to vectors appear only through covariant derivatives and respect TT–null orthogonality (cf. typed/triadic TT selection in Ch. 10: eq. (10.105) and Lemma 10.172).

Axial (pseudoscalar) $\mathrm{Shift}_{\tilde{\varphi}}^{\mathrm{ax}}$ **(softly broken).** $\tilde{\varphi} \mapsto \tilde{\varphi} + \tilde{c}$. Allows derivative/topological couplings for the pseudo–Apeiron channel $\tilde{S}_{\mu\nu}^{(p)}[\tilde{\varphi}]$. Axion–like terms are permitted if compatible with portal bounds; soft breaking $V_p(\tilde{\varphi})$ sets late–time dynamics while keeping TT isolation. (Null-screen coherence/positivity constraints from Ch. 11 apply to τ–typed operators; see eqs. (11.249) and (11.250).)

Vector phase $U(1)_\tau$ **(Stueckelberg/Higgs realization).** $K_\mu^{(\tau)} \to K_\mu^{(\tau)} + \nabla_\mu^{(\mathrm{rel})}\alpha_\tau$, field strength $\mathcal{H}_{\mu\nu} = 2\nabla_{[\mu}^{(\mathrm{rel})}K_{\nu]}^{(\tau)}$ (cf (13.114)). Gauge–invariant kinetic term $-\frac{1}{4}\mathcal{H}^2$; a Proca/Stueckelberg mass μ_C (or m_V) softly breaks $U(1)_\tau$ without spoiling spectral decoupling, clusterability, or

TT purity. (Compatibility with TT isolation and typed projectors per Ch. 10: eqs. (10.4) and (10.105); causal/variational backbone per Ch. 11: eq. (11.203).) Observables enter via the coherence kernel with $k_c^{(V)}(a) \sim a\, m_V\, \mathfrak{m}_{\rm prod}(H)$.

Helicity rotations $U(1)_{\rm hel}$ (transverse plane). On $K_\lambda^{(\tau)}$ with $\lambda = \pm$: $K_\lambda^{(\tau)} \mapsto e^{i\lambda\alpha_{\rm hel}} K_\lambda^{(\tau)}$ (i.e. $SO(2)$ rotations in the helicity plane). Enforces helicity–neutral couplings (diagonal in λ) and suppresses operators that would mix $+$ and $-$ without compensators. Compatible with isotropy on FLRW and TT–null identities (see the TT projector/selection in Ch. 10: eqs. (10.4) and (10.105); typed commuting projectors in Ch. 11 around eq. (11.120)).

Discrete $\mathbb{Z}_2^{\rm HS}$ (stability & branch tag). Holosymmetric parity tagging the realized HSP branch: $(\varphi_b, \varphi_L, \tilde\varphi; K_\mu^{(\tau)}) \mapsto (\pm\varphi_b, \pm\varphi_L, \pm\tilde\varphi; -K_\mu^{(\tau)})$. Forbids odd operators, stabilizes dark excitations, and suppresses mixings that would violate portal/darkness constraints. (All couplings above are to be read together with the typed propagators/matrix–propagator analysis Section 13.3, the TT–null relations (13.112)–(13.115), and the Ch. 10/11 structures just cited.)

Operator hygiene and operational darkness. Operators that would induce observable SM responses (e.g. kinetic mixings or portal terms) are absent/suppressed by the operational darkness criterion (Definition 13.1): on the BRST–physical space, $\langle \mathcal{O}_{\rm vis} \rangle_{\rm phys} \approx 0$ for all $\mathcal{O}_{\rm vis} \in \mathcal{A}(G_{\rm SM})$. Allowed couplings must also respect $\mathcal{P}^{(2)}[\hat\tau_{\mu\nu}] = 0$ and $\mathcal{P}^{(2)}\mathcal{M}_{\tau h} = 0$ (no TT leakage), ensuring that gravitational waves remain purely tensorial (cf. Definition 13.78).

Currents, Charges, and Breaking Patterns (Effective)

Conserved labels. At leading order, Shift$_\varphi$/$U(1)_\varphi$ and $U(1)_C$ furnish effective Noether currents J_φ^μ and J_C^μ on the emergent background $(\mathcal{M}, \bar\Sigma_{\mu\nu})$. Soft breaking by $V(\varphi)$ (DE branch) and by the vector mass μ_C (DM branch) is allowed, but the *labels* remain good quantum numbers for the holosymmetric interferometer used in HSP Stage II (cf. Section 13.1.4).

Branch structure and isotropy. The scalar branch (Gaiaton) realizes dark energy in the slow–roll/near–vacuum regime ($w_\varphi \simeq -1$), while the vector branch (Chrono–Kymon) realizes cold dark matter in the rapid–oscillation regime ($\langle w_{K^{(\tau)}} \rangle \simeq 0$) (cf. Section 13.3 and Definition 13.72). Large–scale isotropy is secured by scalar homogeneity and, for vectors, by triads or domain averaging (cf. Propositions 13.73 and 13.77).

13.4.1 The cosmic symmetry stack (canonical chronogeometric placement)

This subsection formalizes the complete symmetry structure by placing the dark-sector effective symmetries $\mathcal{G}_{\rm Dark}^{(\rm eff)}$ inside the *chronogeometric* fiber, consistent with the unified fluctuation $\delta\hat\Sigma_{\mu\nu} = \hat h_{\mu\nu} + \hat\tau_{\mu\nu}$ and the TT–null constraints (cf. eq. (13.87)). The Standard Model (SM) gauge group remains an external factor as in Section 12.12.

Chronogeometric origin and typing. On the diagnostic realized reference configuration $(\mathcal{X}_q, \bar{\Sigma})$, the TT projector isolates the radiative geometry,

$$\hat{h}_{\mu\nu} := \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\, \delta\hat{g}_{\mu\nu}, \qquad \Pi^{[\mathrm{TT}]}_{\bar{\Sigma}} \hat{\tau}_{\mu\nu} = 0, \tag{13.119}$$

and the full τ–kairos decomposition reads

$$\hat{\tau}_{\mu\nu} = S^{(b)}_{\mu\nu}\big[\hat{\varphi}_b\big] \;\oplus\; S^{(L)}_{\mu\nu}\big[\hat{\varphi}_L\big] \;\oplus\; \tilde{S}^{(p)}_{\mu\nu}\big[\tilde{\varphi}\big] \;\oplus\; \sum_{\lambda=\pm} A^{(\lambda)}_{\mu\nu}\big[\hat{K}^{(\tau)}_{\lambda}\big], \tag{13.120}$$

$$\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\, \Pi^{[\chi]} = 0 \quad \forall\, [\chi] \in \{b, L, p, \lambda = \pm\}. \tag{13.121}$$

reusing Section 12.12 and eq. (12.262). The $\mathrm{SO}(1,3)_{\mathrm{local}}$ factor acts on the TT subspace (and on the trace/radial scalar of Gaiamiti, i.e. Apeiron), while $\mathcal{G}^{(\mathrm{eff})}_{\mathrm{Dark}}$ acts on the non–TT registers summarized in (13.121) (cf. eq. (13.108)).

The Chronogeometric group).

$$\boxed{\mathcal{G}_{\mathrm{Chronogeom}} := \mathrm{Diff}_{\mathrm{rel}}(\mathcal{X}_q) \;\ltimes\; \Big(\mathrm{SO}(1,3)_{\mathrm{local}} \;\times\; \mathcal{G}^{(\mathrm{eff})}_{\mathrm{Dark}}\Big)} \tag{13.122}$$

where $\mathrm{Diff}_{\mathrm{rel}}(\mathcal{X}_q)$ acts by pullback on fields and by adjoint transport on the fiber $\mathrm{SO}(1,3)_{\mathrm{local}} \times \mathcal{G}^{(\mathrm{eff})}_{\mathrm{Dark}}$. The effective factor $\mathcal{G}^{(\mathrm{eff})}_{\mathrm{Dark}}$ is the τ–kairos symmetry stack introduced in eq. (13.108); it may be softly broken by cosmological operators while preserving TT–nullity (e.g. Proca masses for $\hat{K}^{(\tau)}_{\lambda}$, scalar potentials for $\hat{\varphi}_{b,L}, \tilde{\varphi}$).

The complete cosmic symmetry stack (canonical). The complete symmetry structure, $\mathcal{G}_{\mathrm{Complete}}$, combines the unified chronogeometric symmetries with the Standard Model gauge group. It is formally defined as:

$$\boxed{\begin{aligned}\mathcal{G}_{\mathrm{Cosmos}} &= \underbrace{\mathcal{G}_{\mathrm{Chronogeom}}}_{\text{acts on } \delta\hat{\Sigma}_{\mu\nu}=\hat{h}_{\mu\nu}\oplus\hat{\tau}_{\mu\nu}} \times \underbrace{G_{\mathrm{SM}}}_{\text{Tantrakyma (matter/gauge); cf. eq. (12.254)}} \\ &= \underbrace{\Big[\mathrm{Diff}_{\mathrm{rel}}(\mathcal{X}_q) \ltimes \Big(\mathrm{SO}(1,3)_{\mathrm{local}} \times \mathcal{G}^{(\mathrm{eff})}_{\mathrm{Dark}}\Big)\Big]}_{\text{Unified Chronogeometry (Gravity + Dark Sector)}} \times \underbrace{G_{\mathrm{SM}}}_{\text{Matter Fields}}\end{aligned}} \tag{13.123}$$

where $\mathcal{G}_{\mathrm{Chronogeom}}$ incorporates both local Lorentz symmetry and the effective dark-sector symmetries within the geometric fiber, acted upon by relative diffeomorphisms (eq. (13.122)). Thus, geometry (TT graviton and Apeiron) and the non–TT τ–kairos live together in the chronogeometric fiber, while the SM gauge content (G_{SM}) remains an external internal group acting on Tantrakyma.

Operational guardrails. The typed separation is enforced by the projectors and identities of eqs. (10.4), (12.262) and (13.87). In particular,

$$\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\, \hat{\tau}_{\mu\nu} = 0, \qquad \text{and} \qquad \text{CK modes } \hat{K}^{(\tau)}_{\lambda} \text{ have no spin-2 support,} \tag{13.124}$$

so TT radiation remains purely geometric. Conversely, the SM spin–1 gauge bosons transform under G_{SM} (Tantrakyma) and are distinct from the τ–CK modes (cf. the usage fixed by Section 12.12).

Remarks (effective vs. fundamental). The factors inside $\mathcal{G}_{\text{Chronogeom}}$ split by *type*: $\mathrm{SO}(1,3)_{\text{local}}$ is fundamental (local Lorentz), while $\mathcal{G}_{\text{Dark}}^{(\text{eff})}$ is *effective* on the τ–registers (eq. (13.108)). The external G_{SM} is the fundamental internal gauge group of the SM, as in eq. (12.254). This arrangement matches the typed origin $\delta\hat{\Sigma}_{\mu\nu} = \hat{h}_{\mu\nu} \oplus \hat{\tau}_{\mu\nu}$ and preserves the block structure used in the weave instrumentation and HSP selection (cf. Definitions 10.1 and 10.183, eq. (10.61), and Section 13.1.4).

The cosmic symmetry stack (summary)

Start with Unified Chronogeometry: Begin with $\mathcal{G}_{\text{Chronogeom}}$, which represents the **full** set of geometric symmetries. It includes diffeomorphisms acting on a fiber that contains **both** local Lorentz symmetry ($\mathrm{SO}(1,3)_{\text{local}}$) for gravity **and** the effective dark sector symmetries ($\mathcal{G}_{\text{Dark}}^{(\text{eff})}$). This group describes how the unified field $\delta\hat{\Sigma}_{\mu\nu}$ (containing both $\hat{h}_{\mu\nu}$ and $\hat{\tau}_{\mu\nu}$) transforms.

$$\mathcal{G}_{\text{Chronogeom}} := \mathrm{Diff}_{\text{rel}}(\mathcal{X}_q) \;\ltimes\; \Big(\mathrm{SO}(1,3)_{\text{local}} \;\times\; \mathcal{G}_{\text{Dark}}^{(\text{eff})}\Big),$$

Add the Standard Model: Then take this complete geometric group and combine it with the separate group describing the internal symmetries of matter, G_{SM} ($SU(3)_C \times SU(2)_L \times U(1)_Y$). Since these groups act on fundamentally different things (geometry vs. internal matter states/Tantrakyma), they are combined with a direct product.

$$\mathcal{G}_{\text{Cosmos}} = \mathcal{G}_{\text{Chronogeom}} \times G_{\text{SM}}$$

Result: Substituting the definition of $\mathcal{G}_{\text{Chronogeom}}$ gives the full expression for $\mathcal{G}_{\text{Cosmos}}$:

$$\mathcal{G}_{\text{Cosmos}} = \Big[\mathrm{Diff}_{\text{rel}}(\mathcal{X}_q) \ltimes \Big(\mathrm{SO}(1,3)_{\text{local}} \times \mathcal{G}_{\text{Dark}}^{(\text{eff})}\Big)\Big] \times G_{\text{SM}}$$

with $\mathcal{G}_{\text{Dark}}^{(\text{eff})}$ being the effective dark sector group

$$\begin{aligned}\mathcal{G}_{\text{Dark}}^{(\text{eff})} = \quad & \underbrace{\mathrm{Shift}_{\varphi_b} \times \mathrm{Shift}_{\varphi_L} \times \mathrm{Shift}^{\text{ax}}_{\tilde{\varphi}}}_{\text{scalar/pseudoscalar } \tau\text{–kairos (Apeiron \& pseudo-Apeiron)}} \\ & \times \quad \underbrace{U(1)_\tau \times U(1)_{\text{hel}}}_{\text{vector } \tau\text{–kairos / Chrono–Kymon phase \& helicity}} \\ & \times \quad \underbrace{\mathbb{Z}_2^{\text{HS}}}_{\text{stability / HSP sector tag}}\end{aligned}, \tag{13.125}$$

so that the complete cosmos symmetry stack is

$$\begin{aligned}\mathcal{G}_{\text{Cosmos}} &= \underbrace{\mathcal{G}_{\text{Chronogeom}}}_{\text{acts on } \delta\hat{\Sigma}_{\mu\nu}=\hat{h}_{\mu\nu}\oplus\hat{\tau}_{\mu\nu}} \times \underbrace{G_{\text{SM}}}_{\text{Tantrakyma (matter/gauge)}} \\ &= \underbrace{\Big[\mathrm{Diff}_{\text{rel}}(\mathcal{X}_q) \ltimes \Big(\mathrm{SO}(1,3)_{\text{local}} \times \mathcal{G}_{\text{Dark}}^{(\text{eff})}\Big)\Big]}_{\text{Unified Chronogeometry (Gravity + Dark Sector)}} \times \underbrace{SU(3)_C \times SU(2)_L \times U(1)_Y}_{\text{Matter Fields}},\end{aligned} \tag{13.126}$$

13.4.2 Holosymmetric Selection and Operational Darkness

The stack $\mathcal{G}_{\mathrm{Dark}}$ does not *postulate* new visible interactions; it *organizes* the tau–kairos degrees of freedom so that HSP can adjudicate among their universal histories. Operational darkness is enforced by the coherence–projector/decoupling criterion of Definition 13.1, while projector constraints from Section 13.3 guarantee TT purity. The realized dark sector is therefore the HSP–selected branch of the same operator content: scalar Gaiaton (Apeiron) for dark energy, vector Chrono–Kymon for dark matter, with both acting through $\Sigma_{\mu\nu}$ and remaining spectrally orthogonal to G_{SM}.

$$\begin{aligned}\mathcal{G}_{\mathrm{Dark}}^{(\mathrm{eff})} = \mathrm{Shift}_{\varphi_b} \times \mathrm{Shift}_{\varphi_L} \times \mathrm{Shift}_{\tilde{\varphi}}^{\mathrm{ax}} \\ \times U(1)_\tau \times U(1)_{\mathrm{hel}} \\ \times \mathbb{Z}_2^{\mathrm{HS}}\end{aligned} \quad ,$$

13.4.3 Factor I: Shift_φ / $U(1)_\varphi$ (Apeiron / Dark–Energy Background)

Definition 13.80 (Apeiron/Gaiaton sector and symmetry (background–agnostic))**.** Let φ denote the scalar *tau–kairos* excitation (Gaiaton/Apeiron) that controls the dark–energy background (cf. Section 13.3 and Definition 13.68). The effective symmetry acting on φ is either:

(i) Continuous shift symmetry
a continuous *shift* symmetry $\varphi \mapsto \varphi + c$, $c \in \mathbb{R}$, *softly broken* by the potential $V(\varphi)$,

(ii) Compact phase symmetry
a compact *phase* symmetry $U(1)_\varphi :\ \Phi \mapsto e^{i\alpha}\Phi$ when φ arises as the phase of a complex field Φ (Stueckelberg/axion–like embedding), with $V = V(|\Phi|)$ preserving the phase.

These symmetries render a light φ technically natural and restrict allowed operators by favoring derivative couplings (operator hygiene consistent with Definition 13.1).

Notation. The scalar–shift layer comprises three commuting factors $\mathrm{Shift}_{\varphi_b} \times \mathrm{Shift}_{\varphi_L} \times \mathrm{Shift}_{\tilde{\varphi}}^{\mathrm{ax}}$. In what follows, φ denotes the HSP–selected CP–even combination $\varphi := c_b\,\varphi_b + c_L\,\varphi_L$ with $c_b^2 + c_L^2 = 1$; the CP–odd axial mode $\tilde{\varphi}$ is handled by $\mathrm{Shift}_{\tilde{\varphi}}^{\mathrm{ax}}$ as a distinct factor.

The symmetry control supports $w_\varphi \simeq -1$ in the slow–roll/slow–drift regime, with the usual FLRW specialization (recalled from eq. (13.89)):

$$\rho_\varphi = \tfrac{1}{2}\dot{\varphi}^2 + V(\varphi), \qquad p_\varphi = \tfrac{1}{2}\dot{\varphi}^2 - V(\varphi), \qquad w_\varphi \simeq -1 \ \ (\dot{\varphi}^2 \ll V), \tag{13.127}$$

consistent with precision cosmology [134, 135, 136] and EFT considerations [211, 313]. On a general emergent chronogeometric background $(\mathcal{M}, \bar{\Sigma}_{\mu\nu})$, the covariant fluid variables are given by Definition 13.68, and (13.127) is obtained by the FLRW/comoving reduction (cf. Proposition 13.69). In temporal language, φ sets the *background* of the scalar piece $S_{\mu\nu}[\hat{\varphi}]$ of $\hat{\tau}_{\mu\nu}$ (cf. Section 13.3).

Proposition 13.81 (Noether currents and soft breaking)**.** *For the canonically normalized scalar with relational Lagrangian* $\mathcal{L}_\varphi = \frac{1}{2}\,\bar{\Sigma}^{\mu\nu}\nabla_\mu^{(\mathrm{rel})}\varphi\,\nabla_\nu^{(\mathrm{rel})}\varphi - V(\varphi)$*, the global shift* $\delta\varphi = c$ *yields the formal Noether current*

$$J_\varphi^\mu \;=\; \nabla^{(\mathrm{rel})\,\mu}\varphi, \qquad \nabla_\mu^{(\mathrm{rel})} J_\varphi^\mu \;=\; \Box^{(\mathrm{rel})}\varphi \;=\; -\,V'(\varphi), \tag{13.128}$$

so J^μ_φ is conserved iff $V'(\varphi)=0$ (exact shift symmetry). For a sufficiently flat potential (slow explicit breaking), e.g. $|V'(\varphi)| \ll H_{\rm eff}^2 M_{\rm Pl}$ on cosmological backgrounds,[3] non–conservation is slow and the lightness of φ remains technically natural [211, 313]. In an axion–like $U(1)_\varphi$ embedding with $\Phi = \frac{F}{\sqrt{2}} e^{i\varphi/F}$ and $V = V(|\Phi|)$, the conserved phase current is $J^\mu_\varphi = F^2\, \nabla^{(\rm rel)\,\mu}(\varphi/F)$, with $\nabla^{(\rm rel)}_\mu J^\mu_\varphi = 0$.

Operator hygiene and SM decoupling. Any portal or mixing operator that would yield visible SM responses is absent/suppressed by the operational darkness criterion (Definition 13.1); at the BRST–physical level $\langle \mathcal{O}_{\rm vis}\rangle_{\rm phys} \approx 0$ for all $\mathcal{O}_{\rm vis} \in \mathcal{A}(G_{\rm SM})$. All allowed terms must also respect the projector constraints $\mathcal{P}^{(2)}[\hat\tau_{\mu\nu}] = 0$ and $\mathcal{P}^{(2)}\mathcal{M}_{\tau h} = 0$ (cf. Definition 13.78), ensuring TT purity of the GW channel.

Proof. Work on the emergent chronogeometric background $(\mathcal{M}, \bar\Sigma_{\mu\nu})$ with relational Levi–Civita connection $\nabla^{(\rm rel)}$. Consider

$$S[\varphi] \;=\; \int d^4x\, \sqrt{-\bar\Sigma}\, \mathcal{L}_\varphi, \qquad \mathcal{L}_\varphi = \tfrac{1}{2}\, \bar\Sigma^{\mu\nu} \nabla^{(\rm rel)}_\mu \varphi\, \nabla^{(\rm rel)}_\nu \varphi - V(\varphi).$$

Under a constant shift $\delta\varphi = c$ ($c \in \mathbb{R}$),

$$\begin{aligned} \delta\mathcal{L}_\varphi &= \bar\Sigma^{\mu\nu} \nabla^{(\rm rel)}_\mu \varphi\, \nabla^{(\rm rel)}_\nu (\delta\varphi) - V'(\varphi)\, \delta\varphi \\ &= \nabla^{(\rm rel)}_\mu \Big(\nabla^{(\rm rel)\,\mu} \varphi\, \delta\varphi \Big) - \Big(\Box^{(\rm rel)} \varphi + V'(\varphi) \Big)\, \delta\varphi, \end{aligned} \tag{13.129}$$

where $\Box^{(\rm rel)} := \bar\Sigma^{\mu\nu} \nabla^{(\rm rel)}_\mu \nabla^{(\rm rel)}_\nu$. The Euler–Lagrange equation is

$$\Box^{(\rm rel)} \varphi + V'(\varphi) = 0. \tag{13.130}$$

Using (13.130) in (13.129), the bulk terms cancel on–shell and $\delta\mathcal{L}_\varphi = \nabla^{(\rm rel)}_\mu \Big(\nabla^{(\rm rel)\,\mu} \varphi\, \delta\varphi \Big)$ is a total divergence.

By Noether's theorem for an internal transformation with constant parameter c, the (canonical) current is

$$J^\mu_\varphi := \frac{\partial \mathcal{L}_\varphi}{\partial(\nabla^{(\rm rel)}_\mu \varphi)}\, \delta\varphi = \nabla^{(\rm rel)\,\mu} \varphi\; c.$$

Dividing by the constant c gives

$$J^\mu_\varphi = \nabla^{(\rm rel)\,\mu} \varphi.$$

Taking the covariant divergence and employing (13.130) yields

$$\nabla^{(\rm rel)}_\mu J^\mu_\varphi = \Box^{(\rm rel)} \varphi = -\, V'(\varphi) \quad \text{(on–shell)}. \tag{13.131}$$

Hence the current is exactly conserved iff $V'(\varphi) \equiv 0$ (exact shift symmetry). For slow explicit breaking on cosmological backgrounds, the non–conservation is small; e.g., with an effective expansion/curvature scale $H_{\rm eff}$ one may quantify this as $|V'(\varphi)| \ll H_{\rm eff}^2 M_{\rm Pl}$, ensuring technical naturalness for a light φ.

[3] Here $H_{\rm eff}$ is any effective curvature/expansion scale built from the kinematic invariants of a chosen congruence u^μ, e.g. $H_{\rm eff} \sim \theta/3$ in FLRW; cf. Definition 13.68.

Addendum (phase $U(1)_\varphi$ case). Let $\Phi = \frac{F}{\sqrt{2}} e^{i\varphi/F}$ and $\mathcal{L} = \bar{\Sigma}^{\mu\nu} \nabla^{(\mathrm{rel})}_\mu \Phi^* \, \nabla^{(\mathrm{rel})}_\nu \Phi - V(|\Phi|)$, invariant under $\Phi \to e^{i\alpha}\Phi$. The Noether current is

$$J^\mu_\varphi \;=\; i\Big(\Phi^* \nabla^{(\mathrm{rel})\,\mu} \Phi - \Phi \nabla^{(\mathrm{rel})\,\mu} \Phi^*\Big) \;=\; F^2 \, \nabla^{(\mathrm{rel})\,\mu}(\varphi/F),$$

and $\nabla^{(\mathrm{rel})}_\mu J^\mu_\varphi = 0$ when $V = V(|\Phi|)$ (on shell). □

Corollary 13.82 (Cosmological Form of the Continuity Equation)**.** *On a spatially flat FRW background with scale factor $a(t)$, the time component of (13.128) yields*

$$\partial_t\Big(a^3 \dot{\varphi}\Big) \;=\; a^3 \, V'(\varphi).$$

Hence $a^3\dot{\varphi}$ is conserved for $V'(\varphi) = 0$, while for slow explicit breaking the current relaxes on a Hubble timescale, e.g. $|\partial_t(a^3\dot{\varphi})| = a^3|V'(\varphi)| \ll a^3 H^2 M_{\mathrm{Pl}}$.

Proof. On spatially flat FRW, $\sqrt{-g} = a^3$ and $J^\mu = \nabla^\mu \varphi = (-\dot{\varphi},\; a^{-2}\gamma^{ij}\partial_j\varphi)$ in the $(-,+,+,+)$ signature. From (13.128),

$$\nabla_\mu J^\mu = \frac{1}{\sqrt{-g}} \partial_\mu\Big(\sqrt{-g}\, J^\mu\Big) = -V'(\varphi) \;\Rightarrow\; \partial_t\Big(a^3 J^0\Big) + \partial_i\Big(a^3 J^i\Big) = -a^3 V'(\varphi).$$

Using $J^0 = -\dot{\varphi}$ and $J^i = a^{-2}\gamma^{ij}\partial_j\varphi$ gives

$$-\partial_t\Big(a^3\dot{\varphi}\Big) + \partial_i\Big(a\,\gamma^{ij}\partial_j\varphi\Big) = -a^3 V'(\varphi),$$

hence, for a spatially homogeneous background ($\partial_i\varphi = 0$), $\partial_t\Big(a^3\dot{\varphi}\Big) = a^3 V'(\varphi)$. □

Remark 13.83 (Inhomogeneous generalization)**.** Without assuming homogeneity,

$$\partial_t\Big(a^3\dot{\varphi}\Big) \;-\; \partial_i\Big(a\,\gamma^{ij}\partial_j\varphi\Big) \;=\; a^3 V'(\varphi),$$

which reduces to the corollary when $\partial_i\varphi = 0$.

13.4.4 Factor I(c): $\mathrm{Shift}^{\mathrm{ax}}_{\tilde{\varphi}}$ (CP–odd axial mode)

Definition 13.84 (Axial τ–kairos mode and shift symmetry)**.** Let $\tilde{\varphi}$ denote the CP–odd (axial) scalar in the τ–kairos ledger, entering the typed decomposition as $\tilde{S}^{(p)}_{\mu\nu}[\tilde{\varphi}]$ in (13.113). Its effective symmetry is the constant axial shift

$$\mathrm{Shift}^{\mathrm{ax}}_{\tilde{\varphi}}: \quad \tilde{\varphi} \mapsto \tilde{\varphi} + \tilde{c}, \qquad \tilde{c} \in \mathbb{R},$$

treated as *approximate* (softly broken by $V_p(\tilde{\varphi})$). The mode is neutral under $U(1)_\tau$ and $U(1)_{\mathrm{hel}}$, which act only on the vector register $K^{(\tau)}_\mu$ (eq. (13.114)).

Relational Lagrangian and allowed couplings. On the emergent chronogeometric background $(\mathcal{M}, \bar{\Sigma}_{\mu\nu})$,

$$\mathcal{L}_{\tilde{\varphi}} = \tfrac{1}{2}\,\bar{\Sigma}^{\mu\nu}\nabla^{(\mathrm{rel})}_{\mu}\tilde{\varphi}\,\nabla^{(\mathrm{rel})}_{\nu}\tilde{\varphi} - V_p(\tilde{\varphi}) + \frac{\kappa_\tau}{4F_{\tilde{\varphi}}}\,\tilde{\varphi}\,\mathcal{H}_{\mu\nu}\,\tilde{\mathcal{H}}^{\mu\nu} + \frac{\kappa_g}{F_{\tilde{\varphi}}}\,\tilde{\varphi}\,\mathcal{P}, \tag{13.132}$$

where $\mathcal{H}_{\mu\nu} := 2\nabla^{(\mathrm{rel})}_{[\mu}K^{(\tau)}_{\nu]}$ (eq. (13.114)), $\tilde{\mathcal{H}}^{\mu\nu} := \frac{1}{2}\,\bar{\varepsilon}^{\mu\nu\alpha\beta}\mathcal{H}_{\alpha\beta}$ uses the volume form of $\bar{\Sigma}$, and $\mathcal{P} := \frac{1}{2}\,\bar{\varepsilon}^{\mu\nu\rho\sigma}\,R_{\mu\nu\alpha\beta}(\bar{\Sigma})\,R_{\rho\sigma}{}^{\alpha\beta}(\bar{\Sigma})$ is the gravitational Pontryagin density. Portal hygiene forbids visible–sector $\tilde{\varphi}\,F\tilde{F}$ couplings (darkness projector; Definition 13.1); the $\tilde{\varphi}\,\mathcal{H}\tilde{\mathcal{H}}$ term is purely dark and $U(1)_\tau$–invariant.

Proposition 13.85 (Axial shift current and anomalous Ward identity)**.** *For* (13.132)*, the axial shift current* $J^{\mu}_{\tilde{\varphi}} := \nabla^{(\mathrm{rel})\,\mu}\tilde{\varphi}$ *satisfies*

$$\nabla^{(\mathrm{rel})}_{\mu}J^{\mu}_{\tilde{\varphi}} = -\,V_p'(\tilde{\varphi}) + \frac{\kappa_\tau}{4F_{\tilde{\varphi}}}\,\mathcal{H}_{\mu\nu}\,\tilde{\mathcal{H}}^{\mu\nu} + \frac{\kappa_g}{F_{\tilde{\varphi}}}\,\mathcal{P}\,. \tag{13.133}$$

Thus $\mathrm{Shift}^{\mathrm{ax}}_{\tilde{\varphi}}$ *is exact only if* $V_p' \equiv 0$ *and the topological densities vanish; otherwise it is softly/quantum–anomalously broken.*

Corollary 13.86 (Equation of motion and cosmological limit)**.** *The* $\tilde{\varphi}$ *equation of motion is* $\Box^{(\mathrm{rel})}\tilde{\varphi} - V_p'(\tilde{\varphi}) = \frac{\kappa_\tau}{4F_{\tilde{\varphi}}}\mathcal{H}_{\mu\nu}\tilde{\mathcal{H}}^{\mu\nu} + \frac{\kappa_g}{F_{\tilde{\varphi}}}\mathcal{P}$*. On FLRW (with vanishing background* $\mathcal{P}$ *and* $\mathcal{H}\tilde{\mathcal{H}}$*), the homogeneous background obeys* $\ddot{\bar{\tilde{\varphi}}} + 3H\dot{\bar{\tilde{\varphi}}} + V_p'(\bar{\tilde{\varphi}}) = 0$*, so the misalignment branch mirrors the scalar case and induces a Jeans–like cutoff* $k_c^{(S)} \sim \max_a k_J(a)$ *with* $m_S \to m_{\tilde{S}}$ *(eq. (13.59)).*

Typed projector, TT–nullity, and commutation. The axial contribution to $\hat{\tau}_{\mu\nu}$ is carried by $\tilde{S}^{(p)}_{\mu\nu}[\tilde{\varphi}]$ (eq. (13.113)), with $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}\,\tilde{S}^{(p)}_{\mu\nu} = 0$ (typed TT–orthogonality; eq. (13.115)). The shift factor $\mathrm{Shift}^{\mathrm{ax}}_{\tilde{\varphi}}$ commutes with $U(1)_\tau$ and $U(1)_{\mathrm{hel}}$, so axial dynamics do not rotate helicity states nor affect the $U(1)_\tau$ gauge redundancy.

Decision–useful signatures (Stage II). The topological couplings in (13.132) produce parity–odd responses that are *dark* to SM meters: (i) chiral phase shifts in the Chrono–Kymon via $\tilde{\varphi}\,\mathcal{H}\tilde{\mathcal{H}}$ (helicity–asymmetric tails in $\mathcal{S}(k; k_c, \nu)$, eq. (13.57)); (ii) potential GW birefringence sourced *geometrically* by $\mathcal{P}$ while preserving TT purity of $\hat{h}_{\mu\nu}$ ($\Pi^{[\mathrm{TT}]}\hat{\tau} = 0$; eq. (13.112)). These feed the Stage II visibility/coherence score used by HSP selection (eqs. (13.48) and (13.49) and Section 13.1.4).

Operator hygiene (portals and decoupling). Visible portals $\tilde{\varphi}\,F\tilde{F}$ and $\tilde{\varphi}\,\bar{\psi}\gamma_5\psi$ are excluded by the darkness criterion (Definition 13.1); allowed dark couplings must respect $\mathcal{P}^{(2)}[\hat{\tau}_{\mu\nu}] = 0$ and $\mathcal{P}^{(2)}\mathcal{M}_{\tau h} = 0$ (no TT leakage; Definition 13.78).

13.4.5 Factor II: $U(1)_\tau$ (Chrono–Kymon / Dark–Matter Carrier)

Definition 13.87 (Chrono–Kymon gauge structure (background–agnostic))**.** Let $K^{(\tau)}_{\mu}$ be the Tantrakyma vector excitation (Chrono–Kymon) on $(\mathcal{M}, \bar{\Sigma}_{\mu\nu})$ with relational Levi–Civita $\nabla^{(\mathrm{rel})}$. At the effective level it transforms under a dark $U(1)_\tau$ with field strength

$$\mathcal{H}_{\mu\nu} := 2\,\nabla^{(\mathrm{rel})}_{[\mu}K^{(\tau)}_{\nu]}\,.$$

Notation. Here μ_C is the established Chrono–Kymon mass parameter. In Chapter 6, g_C denotes a Tantrakyma pairing and is *not* used as a gauge coupling; we therefore use g_τ for the $U(1)_\tau$ coupling.

Mass generation while preserving sector separation. Two realizations produce a finite vector mass μ_C without spoiling spectral decoupling from G_{SM} (Definition 13.1) or TT isolation:

Stueckelberg Introduce a scalar compensator θ and gauge parameter α_τ with

$$K_\mu^{(\tau)} \;\mapsto\; K_\mu^{(\tau)} + \nabla_\mu^{(\mathrm{rel})}\alpha_\tau, \qquad \theta \;\mapsto\; \theta + \mu_C\alpha_\tau.$$

The corresponding gauge–invariant mass term is

$$\frac{1}{2}\Big(\mu_C K_\mu^{(\tau)} - \nabla_\mu^{(\mathrm{rel})}\theta\Big)\Big(\mu_C K^{(\tau)\mu} - \nabla^{(\mathrm{rel})\,\mu}\theta\Big).$$

Higgsed Introduce a dark Higgs field H_τ with $\langle H_\tau\rangle \neq 0$ and covariant derivative

$$D_\mu H_\tau = \Big(\nabla_\mu^{(\mathrm{rel})} - i g_\tau K_\mu^{(\tau)}\Big) H_\tau.$$

In the broken phase this yields a Proca mass $\mu_C = g_\tau\,\langle|H_\tau|\rangle$.

Relational effective action.

$$\mathcal{L}_{K^{(\tau)}} = -\frac{1}{4}\,\mathcal{H}_{\mu\nu}\mathcal{H}^{\mu\nu} + \frac{1}{2}\,(\mu_C K_\mu^{(\tau)} - \nabla_\mu^{(\mathrm{rel})}\theta)(\mu_C K^{(\tau)\mu} - \nabla^{(\mathrm{rel})\,\mu}\theta) + \mathcal{L}_{\mathrm{int}}[\bar{\Sigma}, K^{(\tau)}], \quad (13.134)$$

which in unitary gauge $\theta = 0$ reduces to

$$\mathcal{L}_{K^{(\tau)}} \simeq -\frac{1}{4}\,\mathcal{H}_{\mu\nu}\mathcal{H}^{\mu\nu} + \frac{1}{2}\,\mu_C^2\,K_\mu^{(\tau)}K^{(\tau)\mu} + \mathcal{L}_{\mathrm{int}}[\bar{\Sigma}, K^{(\tau)}]. \quad (13.135)$$

Proposition 13.88 (Dark $U(1)_\tau$ current: covariant on–shell conservation)**.** *For* (13.134)*, the gauge–invariant Noether (Gauss–law) current*

$$J_\tau^\mu \;\equiv\; \nabla_\nu^{(\mathrm{rel})}\mathcal{H}^{\nu\mu} + \mu_C^2\left(K^{(\tau)\mu} - \frac{1}{\mu_C}\,\nabla^{(\mathrm{rel})\,\mu}\theta\right) \quad (13.136)$$

satisfies $\nabla_\mu^{(\mathrm{rel})}J_\tau^\mu = 0$ on–shell. In unitary gauge $\theta = 0$, $J_\tau^\mu = \nabla_\nu^{(\mathrm{rel})}\mathcal{H}^{\nu\mu} + \mu_C^2 K^{(\tau)\mu}$. In the Higgsed realization with dark Higgs H_τ,

$$j_{H_\tau}^\mu \;\equiv\; i g_\tau\Big(H_\tau^* D^\mu H_\tau - (D^\mu H_\tau)^* H_\tau\Big), \qquad J_\tau^\mu \;=\; \nabla_\nu^{(\mathrm{rel})}\mathcal{H}^{\nu\mu} + \mu_C^2 K^{(\tau)\mu} - j_{H_\tau}^\mu, \quad (13.137)$$

and $\nabla_\mu^{(\mathrm{rel})}J_\tau^\mu = 0$ on–shell.

Operator hygiene and TT isolation. All operators must preserve operational darkness (Definition 13.1) and the projector constraints $\mathcal{P}^{(2)}[\hat{\tau}_{\mu\nu}] = 0$, $\mathcal{P}^{(2)}\mathcal{M}_{\tau h} = 0$ (no TT leakage; Definition 13.78).

Proof. **(Stueckelberg case; background–agnostic).** Work on $(\mathcal{M}, \bar{\Sigma}_{\mu\nu})$ with the relational Levi–Civita connection $\nabla^{(\mathrm{rel})}$. Use the gauge–invariant Stueckelberg Lagrangian (13.134) with $\mathcal{H}_{\mu\nu} := 2\,\nabla^{(\mathrm{rel})}_{[\mu} K^{(\tau)}_{\nu]}$:

$$\mathcal{L}_{K^{(\tau)}} = -\tfrac{1}{4}\,\mathcal{H}_{\mu\nu}\mathcal{H}^{\mu\nu} + \tfrac{1}{2}\,(\mu_C K^{(\tau)}_{\mu} - \nabla^{(\mathrm{rel})}_{\mu}\theta)(\mu_C K^{(\tau)\mu} - \nabla^{(\mathrm{rel})\,\mu}\theta),$$

invariant under $\delta K^{(\tau)}_{\mu} = \nabla^{(\mathrm{rel})}_{\mu}\alpha_\tau$, $\delta\theta = \mu_C\alpha_\tau$. The Euler–Lagrange equations are

$$\mathcal{E}^{\mu}_{K^{(\tau)}} := \nabla^{(\mathrm{rel})}_{\nu}\mathcal{H}^{\nu\mu} + \mu_C^2\left(K^{(\tau)\mu} - \frac{1}{\mu_C}\nabla^{(\mathrm{rel})\,\mu}\theta\right) = 0, \tag{13.138}$$

$$\mathcal{E}_{\theta} := \nabla^{(\mathrm{rel})}_{\mu}\left(K^{(\tau)\mu} - \frac{1}{\mu_C}\nabla^{(\mathrm{rel})\,\mu}\theta\right) = 0. \tag{13.139}$$

Define J^{μ}_{τ} as in (13.136):

$$J^{\mu}_{\tau} \equiv \nabla^{(\mathrm{rel})}_{\nu}\mathcal{H}^{\nu\mu} + \mu_C^2\left(K^{(\tau)\mu} - \frac{1}{\mu_C}\nabla^{(\mathrm{rel})\,\mu}\theta\right).$$

Taking the covariant divergence and using (13.139) gives

$$\nabla^{(\mathrm{rel})}_{\mu}J^{\mu}_{\tau} = \underbrace{\nabla^{(\mathrm{rel})}_{\mu}\nabla^{(\mathrm{rel})}_{\nu}\mathcal{H}^{\nu\mu}}_{=0} + \mu_C^2\left(\nabla^{(\mathrm{rel})}_{\mu}K^{(\tau)\mu} - \frac{1}{\mu_C}\,\Box^{(\mathrm{rel})}\theta\right) = 0.$$

The first term vanishes *identically* for any torsion–free, metric–compatible connection because $\mathcal{H}^{\nu\mu}$ is antisymmetric:[4]

$$\nabla^{(\mathrm{rel})}_{\mu}\nabla^{(\mathrm{rel})}_{\nu}\mathcal{H}^{\nu\mu} = \tfrac{1}{2}\Big[\nabla^{(\mathrm{rel})}_{\mu}, \nabla^{(\mathrm{rel})}_{\nu}\Big]\mathcal{H}^{\nu\mu} = 0,$$

since the commutator contracts with the symmetric Ricci tensor and cancels by antisymmetry. The second bracket vanishes on–shell by (13.139). Hence $\nabla^{(\mathrm{rel})}_{\mu}J^{\mu}_{\tau} = 0$. In unitary gauge $\theta = 0$ one recovers $J^{\mu}_{\tau} = \nabla^{(\mathrm{rel})}_{\nu}\mathcal{H}^{\nu\mu} + \mu_C^2 K^{(\tau)\mu}$ with the same conclusion.

(Higgsed case). Let H_τ carry charge g_τ with $D_\mu H_\tau = (\nabla^{(\mathrm{rel})}_{\mu} - i g_\tau K^{(\tau)}_{\mu})H_\tau$. The vector equation of motion is

$$\nabla^{(\mathrm{rel})}_{\nu}\mathcal{H}^{\nu\mu} + \mu_C^2 K^{(\tau)\mu} = j^{\mu}_{H_\tau}, \qquad j^{\mu}_{H_\tau} := i g_\tau\Big(H^*_\tau D^\mu H_\tau - (D^\mu H_\tau)^* H_\tau\Big). \tag{13.140}$$

Gauge invariance of the Higgs sector implies $\nabla^{(\mathrm{rel})}_{\mu} j^{\mu}_{H_\tau} = 0$ on–shell. Applying $\nabla^{(\mathrm{rel})}_{\mu}$ to (13.140) and using the same antisymmetry identity yields

$$\nabla^{(\mathrm{rel})}_{\mu}\Big(\nabla^{(\mathrm{rel})}_{\nu}\mathcal{H}^{\nu\mu} + \mu_C^2 K^{(\tau)\mu} - j^{\mu}_{H_\tau}\Big) = 0,$$

i.e. $\nabla^{(\mathrm{rel})}_{\mu}J^{\mu}_{\tau} = 0$ with J^{μ}_{τ} as in (13.137). □

[4] Equivalently, in form language: $J_\tau = \delta\mathcal{H} + \mu_C^2(K - \frac{1}{\mu_C}d\theta)$ and $\delta J_\tau = \delta^2\mathcal{H} + \mu_C^2(\delta K - \frac{1}{\mu_C}\Box\theta) = \mu_C^2\,\mathcal{E}_\theta = 0$.

Remark 13.89 (On the form used in the main text)**.** The compact expression $J^\mu_\tau = \nabla^{(\mathrm{rel})}_\nu \mathcal{F}^{(\tau)\nu\mu} + \mu_C^2 K^{(\tau)\mu}$ quoted in the main text is the *unitary–gauge* ($\theta = 0$) specialization of the gauge–invariant Stueckelberg current in eq. (13.136). Equivalently, using $\mathcal{H}_{\mu\nu} \equiv \mathcal{F}^{(\tau)}_{\mu\nu} = 2\nabla^{(\mathrm{rel})}_{[\mu} K^{(\tau)}_{\nu]}$, the gauge–invariant completion away from unitary gauge reads

$$J^\mu_\tau = \nabla^{(\mathrm{rel})}_\nu \mathcal{H}^{\nu\mu} + \mu_C^2 \left(K^{(\tau)\mu} - \frac{1}{\mu_C} \nabla^{(\mathrm{rel})\,\mu} \theta \right),$$

i.e. it includes the Stueckelberg contribution $-\mu_C \nabla^{(\mathrm{rel})\,\mu}\theta$. In the Higgs realization, the conserved current further contains the Higgs piece $j^\mu_{H_\tau}$ as in eq. (13.137). All forms are related by a gauge choice and coincide on–shell up to identically conserved (Bianchi) terms.

13.4.6 Factor III: $\mathbb{Z}_2^{\mathrm{HS}}$ (stability & HSP sector tag)

Definition 13.90 (Holosymmetric dark parity (sector tag & stabilizer))**.**

$$\varphi \mapsto \pm\varphi, \qquad K^{(\tau)}_\mu \mapsto -K^{(\tau)}_\mu, \qquad \theta \mapsto -\theta, \qquad H_\tau \mapsto -H_\tau.$$

The combination $(\mu_C K^{(\tau)}_\mu - \nabla^{(\mathrm{rel})}_\mu \theta)$ flips sign, so its quadratic mass term is parity–even. In the Higgs realization, $j^\mu_{H_\tau}$ and $|H_\tau|^2$ are parity–even.

Invariants. Under $\mathbb{Z}_2^{\mathrm{HS}}$ one has $\mathcal{H}_{\mu\nu} \to -\mathcal{H}_{\mu\nu}$ and $D_\mu H_\tau \to -(D_\mu H_\tau)$, so $-\frac{1}{4}\mathcal{H}^2$, $(\mu_C K^{(\tau)} - \nabla\theta)^2$, $|H_\tau|^2$, and $j^\mu_{H_\tau}$ are parity–even; thus all TT–null/projector constraints remain intact (Definition 13.78).

This parity (i) stabilizes the lightest dark–charged state, (ii) forbids *odd* operators that would mix sectors (e.g. linear portals $K^{(\tau)}_\mu J^\mu_{\mathrm{SM}}$, $H_\tau \mathcal{O}_{\mathrm{SM}}$), and (iii) serves as the sector tag used by HSP (Section 13.1.4). *Operator hygiene:* $\mathbb{Z}_2^{\mathrm{HS}}$ alone does *not* eliminate all even portals (e.g. kinetic mixing); operational darkness still requires the coherence–projector/decoupling criterion (Definition 13.1) on the BRST–physical space.

Representation Content and Projector Constraints

The temporal fluctuation carries only scalar $\oplus$ vector polarizations: $\mathcal{P}^{(2)}[\hat{\tau}_{\mu\nu}] = 0$, with $\mathcal{P}^{(2)}$ the TT projector defined on spatial slices of the emergent background $(\mathcal{M}, \bar{\Sigma}_{\mu\nu})$ (cf. Definition 13.78). At quadratic order, perturbations assemble into the background–agnostic matrix propagator

$$S^{(2)} = \frac{1}{2}\int d^4x \sqrt{-\bar{\Sigma}}\, \hat{\Upsilon}^{\mu\nu} \begin{pmatrix} \mathcal{D}_{hh} & \mathcal{M}_{h\tau} \\ \mathcal{M}_{\tau h} & \mathcal{D}_{\tau\tau} \end{pmatrix}_{\mu\nu}{}^{\rho\sigma} \hat{\Upsilon}_{\rho\sigma}, \qquad \hat{\Upsilon}_{\mu\nu} := (\hat{h}_{\mu\nu}, \hat{\tau}_{\mu\nu})^\top,$$
$$\mathcal{P}^{(2)} \circ \mathcal{M}_{\tau h} = 0 \;=\; \mathcal{M}_{h\tau} \circ \mathcal{P}^{(2)}, \tag{13.141}$$

so that all TT (spin–2) gravitational radiation propagates purely in the geometric block $\mathcal{D}_{hh}$ and no extra GW polarizations are introduced—a sharp, falsifiable prediction consistent with current bounds. The constraint (13.141) is compatible with $\mathbb{Z}_2^{\mathrm{HS}}$ and preserved under HSP selection.

Cosmological Mapping: Division of Labor and Observables

Dark energy (scalar temporal / Gaiaton–Apeiron). The background acceleration arises from the scalar tau–kairos field φ in the slow–roll/slow–drift regime, with the *general* (background–agnostic) fluid variables given in Definition 13.68 and the FLRW specialization recalled in (13.127). The scalar temporal fluctuation $S_{\mu\nu}[\hat{\varphi}]$ provides the perturbation channel, inducing $\mathcal{O}(\text{small})$ modifications to the DE sound speed, the linear–growth kernel (e.g. $f\sigma_8$), and the effective Hubble friction; all such effects smoothly reduce to Λ behavior as scalar temporal coherence decoheres (cf. Section 13.3).

Dark matter (vector temporal / Chrono–Kymon). The massive vector $K_\mu^{(\tau)}$ behaves as cold, pressureless matter in the *adiabatic rapid–oscillation* regime (background–agnostic balance (13.106); FLRW scaling (10.154)). Coherent vector configurations source the anti-symmetric/axial temporal piece $A_{\mu\nu}[\,\hat{K}^{(\tau)}\,]$, forming electromagnetically inert *temporal halos* that affect rotation curves, lensing, and small–scale structure, while preserving the GW TT channel (cf. Section 13.3 and Definition 13.78). *Small–scale signatures enter via the coherence–kernel factorization* eq. (13.57) *with the vector cutoff*

$$k_c^{(V)}(a) \;\sim\; a\, m_V\, \mathfrak{m}_{\text{prod}}(H), \qquad \text{see } eq.\ (13.58),$$

which governs damping/oscillatory features in $\mathcal{S}(k; k_c^{(V)}, \nu_V)$.

Observational program.

(i) Dark–energy probes Measure the redshift–dependent equation of state $w(z)$, the effective sound speed c_s^2, and the linear growth index $f\sigma_8$ to test the slow–roll/slow–drift Apeiron background and its perturbations [134, 135, 136].

(ii) Dark–matter probes Use weak–lensing shear, the halo mass–concentration relation, satellite abundances and kinematics, and inner–density–slope/coring diagnostics to constrain the temporal–vector sector and small–scale structure [588, 311].

(iii) GW–channel integrity Verify the absence of extra gravitational–wave polarizations. Operationally this is enforced by the projector constraints in (13.141), i.e. TT purity $\mathcal{P}^{(2)} \circ \mathcal{M}_{\tau h} = 0 = \mathcal{M}_{h\tau} \circ \mathcal{P}^{(2)}$, so that only tensor modes propagate in the geometric block.

Model Realizations: Stueckelberg vs. Higgsed and Portal Hygiene

Stueckelberg $U(1)_\tau$. Gauge redundancy $K_\mu^{(\tau)} \to K_\mu^{(\tau)} + \nabla_\mu^{(\text{rel})}\alpha_\tau$, $\theta \to \theta + \mu_C \alpha_\tau$ generates a gauge–invariant vector mass through the Stueckelberg combination $\left(\mu_C K_\mu^{(\tau)} - \nabla_\mu^{(\text{rel})}\theta\right)^2$, yielding an effective Proca theory while maintaining the $U(1)_\tau$ structure at the level of redundancies. This realization is compatible with holosymmetric sectoring: the $\mathbb{Z}_2^{\text{HS}}$ tag forbids odd portals and supports HSP selection of a temporally dark sector (cf. Section 13.1.4).

Higgsed $U(1)_\tau$. A dark Higgs field H_τ with $\langle H_\tau\rangle \neq 0$ and covariant derivative $D_\mu H_\tau = (\nabla_\mu^{(\mathrm{rel})} - i g_\tau K_\mu^{(\tau)})H_\tau$ generates the vector mass $\mu_C = g_\tau\langle|H_\tau|\rangle$ in exact analogy with the usual Higgs mechanism. After symmetry breaking, a remnant $\mathbb{Z}_2^{\mathrm{HS}}$ can remain to stabilize the lightest dark–charged state and to enforce holosymmetric sector preservation. Portal hygiene then requires suppressing or forbidding renormalizable connectors to the visible sector, most notably kinetic mixing $\varepsilon\,\mathcal{H}_{\mu\nu}B^{\mu\nu}$ between $U(1)_\tau$ and hypercharge [599], and scalar (Higgs) portals $\lambda\,|H|^2|H_\tau|^2$ (see also [548]).

Portal constraints (holosymmetric hygiene). Kinetic mixing $\varepsilon\,\mathcal{H}_{\mu\nu}B^{\mu\nu}$ and scalar portals $\lambda|H|^2|H_\tau|^2$ are set to zero or $\ll 10^{-n}$ by the combined action of $\mathbb{Z}_2^{\mathrm{HS}}$ and the HSP decoupling criterion on the BRST–physical space (Definition 13.1); higher–dimensional portals are suppressed by the holosymmetric scale. This maintains EM darkness, preserves the projector constraints of Definition 13.78, and keeps GW TT purity intact.

Analogy with G_{SM} and Final Synthesis

The factors of $\mathcal{G}_{\mathrm{Dark}}^{(\mathrm{eff})}$ form a *direct product* that acts *jointly* on the τ–kairos register (three scalar shifts $\times U(1)_\tau \times U(1)_{\mathrm{hel}} \times \mathbb{Z}_2^{\mathrm{HS}}$), organizing the full decomposition in eq. (13.113). The analogy to G_{SM} is *structural* rather than unificatory: the badge in eq. (13.108) labels approximate/effective invariances of the τ–sector on the BRST–physical diagnostic realized reference configuration, not a fundamental gauge unification. All factors may be realized simultaneously (commuting up to trivial automorphisms), with soft breakings supplied by the same operators that realize cosmology. TT purity is enforced by placement and projectors: $\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}\hat{\tau}_{\mu\nu} = 0$ (eqs. (13.112) and (13.115)), so spin–2 radiation propagates only in the geometric block of the matrix propagator (eq. (13.141)).

Together with the scalar–vector decomposition (13.113) and the matrix–propagator projector constraints (13.141), this stack yields a unified, falsifiable description of the dark sector that:

(i) Recovers ΛCDM *recovers* ΛCDM in the decohered macroscopic limit (Section 13.3),

(ii) Predicts no extra GW polarizations *predicts* no extra GW polarizations (TT purity via $\mathcal{P}^{(2)}[\hat{\tau}_{\mu\nu}] = 0$ and $\mathcal{P}^{(2)}\mathcal{M}_{\tau h} = 0$; cf. Definition 13.78),

(iii) Provides scalar/vector targets for DE/DM *provides* concrete scalar/vector targets for DE/DM phenomenology (Section 13.3).

Admissibility and realization are adjudicated by the Holosymmetric Selection Principle on the BRST–physical space (Section 13.1.4), with operational darkness enforced by the coherence–projector/decoupling criterion (Definition 13.1).

One-line badge (for tables/figures). For compact summaries, *use* the emblem (13.108) with a caption clarifying that Shift_φ (or $U(1)_\varphi$) governs the DE background and scalar temporal fluctuations, $U(1)_\tau$ governs the vector temporal carrier of DM and halo formation, and $\mathbb{Z}_2^{\mathrm{HS}}$ enforces stability and HSP sector preservation; TT purity and SM decoupling follow from (13.113) and (13.141).

13.4.7 The Abundances of Dark Energy and Dark Matter

Definition 13.91 (Cosmological abundance fractions)**.** We define the cosmological abundance fractions

$$\Omega_{\mathrm{DE}} := \frac{\rho_{\mathrm{DE}}}{\rho_{\mathrm{tot}}}, \qquad \Omega_{\mathrm{DM}} := \frac{\rho_{\mathrm{DM}}}{\rho_{\mathrm{tot}}}, \qquad \Omega_{\mathrm{vis}} := \frac{\rho_{\mathrm{vis}}}{\rho_{\mathrm{tot}}}, \tag{13.142}$$

directly from the spectral structure of the Quantum Holo–State $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$ across its chronogeometric regimes (Section 11.9), using the Holosymmetric Selection Principle (HSP) on the BRST–physical space (Section 13.1.4).

1. Total Energy Budget from the Spectral Resolution of $\hat{H}_{\mathrm{univ}}$

Definition 13.92 (Total comoving energy density from spectral resolution)**.** Let the regulated universal Hamiltonian be

$$\hat{H}_{\mathrm{univ}} \;:=\; \hat{H}_{\Phi} \;+\; \hat{H}_{\Theta} \;+\; \hat{H}_{\mathrm{int}}, \tag{13.143}$$

acting on the emergent/GNS Hilbert space $\mathcal{H}_{\mathrm{em}}$ (Section 2.4). With $\left|\overset{\circ}{\boldsymbol{\Psi}}\right\rangle$ the cosmological (BRST–physical) Quantum Holo–State, the total comoving energy density is the normalized expectation

$$\rho_{\mathrm{tot}} \;:=\; \frac{1}{\mathrm{Vol}(\mathcal{X}_q)} \left\langle \overset{\circ}{\boldsymbol{\Psi}} \right| \hat{H}_{\mathrm{univ}} \left| \overset{\circ}{\boldsymbol{\Psi}} \right\rangle_{\mathrm{phys}} \;=\; \frac{1}{\mathrm{Vol}(\mathcal{X}_q)} \int_{\sigma(\hat{H}_{\mathrm{univ}})} E \, \mathrm{d}\mu_{\overset{\circ}{\boldsymbol{\Psi}}}^{\hat{H}_{\mathrm{univ}}}(E), \tag{13.144}$$

where $\sigma(\hat{H}_{\mathrm{univ}})$ is the spectrum and $\mu_{\overset{\circ}{\boldsymbol{\Psi}}}^{\hat{H}_{\mathrm{univ}}}$ the spectral measure induced by $\left|\overset{\circ}{\boldsymbol{\Psi}}\right\rangle$.

2. Dark Energy as Residual Vacuum Energy

Definition 13.93 (Dark energy as the interferometric residual vacuum energy)**.** Dark energy is the *interferometric residue* of the dual–field vacuum (Gaiamiti + Tantrakyma), selected by the symmetric lensing filter (Section 13.2.7). Writing $H_{\mathrm{vac}} := H_{\Phi}^{\mathrm{vac}} + H_{\Theta}^{\mathrm{vac}}$ and $\mathcal{P}_{\mathrm{vac}}^{\mathrm{sym}}$ for the symmetric coherence projector, set

$$\rho_{\mathrm{DE}} \;:=\; \frac{1}{\mathrm{Vol}(\mathcal{X}_q)} \left\langle \overset{\circ}{\boldsymbol{\Psi}} \right| \mathcal{P}_{\mathrm{vac}}^{\mathrm{sym}} \, H_{\mathrm{vac}} \, \mathcal{P}_{\mathrm{vac}}^{\mathrm{sym}} \left| \overset{\circ}{\boldsymbol{\Psi}} \right\rangle_{\mathrm{phys}} \;\equiv\; \varepsilon_{\Lambda}, \tag{13.145}$$

the small, positive residual vacuum energy density picked out by Stage II coherence maximization (Section 13.2.6). This formulation is equivalent to the interferometric–residue definition in Definition 13.56 and fixes the notation used in the abundance analysis below.

3. Dark Matter as a Holosymmetric–Projected Energy Density

Definition 13.94 (Dark matter as a holosymmetric–projected energy density)**.** Let $\mathcal{P}_{\mathrm{dark}}^{\mathrm{coh}}$ denote the coherence projector onto the union of admissible dark–sector gauges G (i.e., sectors passing Stage I and spectrally decoupled from Standard Model (SM) observables; Section 13.2.8). The dark–matter energy density is

$$\rho_{\mathrm{DM}} \;:=\; \frac{1}{\mathrm{Vol}(\mathcal{X}_q)} \left\langle \overset{\circ}{\boldsymbol{\Psi}} \right| \mathcal{P}_{\mathrm{dark}}^{\mathrm{coh}} \, \hat{H}_{\mathrm{univ}} \, \mathcal{P}_{\mathrm{dark}}^{\mathrm{coh}} \left| \overset{\circ}{\boldsymbol{\Psi}} \right\rangle_{\mathrm{phys}}. \tag{13.146}$$

Equivalently, in any energy–eigenbasis $\{|\psi_\alpha\rangle, E_\alpha\}$ resolving $\hat{H}_{\rm univ}$ on $\mathcal{H}_{\rm phys}$,

$$\rho_{\rm DM} = \frac{1}{{\rm Vol}(\mathcal{X}_q)} \sum_{\alpha\in{\rm dark}} \left\| \mathcal{P}^{\rm coh}_{\rm dark} |\psi_\alpha\rangle \right\|^2 E_\alpha, \qquad \text{with} \quad \langle\psi_\alpha| \mathcal{O}_{\rm vis} |\psi_\alpha\rangle_{\rm phys} \approx 0 \ \ \forall \mathcal{O}_{\rm vis} \in \mathfrak{A}^{\rm obs}_{\rm SM}. \tag{13.147}$$

The expectation clause implements the spectral–decoupling ("darkness") condition of eq. (13.86).

4. Visible–Sector Energy Density

Definition 13.95 (Visible–sector energy density)**.** With $\mathcal{P}^{\rm coh}_{\rm vis} := \mathcal{P}^{\rm coh}_{G_{\rm SM}}$ the Standard Model (SM) coherence projector, the visible contribution is

$$\rho_{\rm vis} := \frac{1}{{\rm Vol}(\mathcal{X}_q)} \left\langle \mathring{\mathbf{\Psi}} \right| \mathcal{P}^{\rm coh}_{\rm vis} \hat{H}_{\rm univ} \mathcal{P}^{\rm coh}_{\rm vis} \left| \mathring{\mathbf{\Psi}} \right\rangle_{\rm phys} = \frac{1}{{\rm Vol}(\mathcal{X}_q)} \sum_{\beta\in{\rm vis}} \left\| \mathcal{P}^{\rm coh}_{\rm vis} |\psi_\beta\rangle \right\|^2 E_\beta. \tag{13.148}$$

Here the sum runs over energy–eigenstates $\{|\psi_\beta\rangle, E_\beta\}$ in the SM-visible sector of $\mathcal{H}_{\rm phys}$.

Remark 13.96 (Foliation and volume regulator)**.** Let u^μ be any admissible unit timelike congruence on $(\mathcal{M}, \bar{\Sigma}_{\mu\nu})$ and let $\mathcal{X}_q$ denote a GNS–coherence cell (compact region) on a Cauchy slice orthogonal to u^μ, with induced measure $\mathrm{d}\Sigma_u$. The factor ${\rm Vol}(\mathcal{X}_q)$ is a regulator that drops out of all abundance ratios in (13.142). Changing the foliation or the cell within the admissible class leaves $\Omega_{\rm DE}, \Omega_{\rm DM}, \Omega_{\rm vis}$ unchanged, provided the BRST–physical projectors and HSP filter are used consistently.

5. Abundance Sum Rule and Gauge–Independence

Proposition 13.97 (Holosymmetric abundance sum rule)**.** *Assume the coherence projectors form a resolution of the identity on the BRST–physical subspace relevant to* $\hat{H}_{\rm univ}$*,*

$$\mathcal{P}^{\rm sym}_{\rm vac} + \mathcal{P}^{\rm coh}_{\rm dark} + \mathcal{P}^{\rm coh}_{\rm vis} = \mathbf{1}_{\rm phys}, \qquad \mathcal{P}_i \mathcal{P}_j = \delta_{ij} \mathcal{P}_i, \tag{13.149}$$

and that cross–sector interferences vanish by HSP selection and spectral decoupling (Section 13.1.4 and eq. (13.86)). Then

$$\rho_{\rm tot} = \rho_{\rm DE} + \rho_{\rm DM} + \rho_{\rm vis}, \qquad \Rightarrow \qquad \Omega_{\rm DE} + \Omega_{\rm DM} + \Omega_{\rm vis} = 1. \tag{13.150}$$

Proof. Insert (13.149) on both sides of $\hat{H}_{\rm univ}$ in (13.144); orthogonality eliminates cross terms and leaves the three diagonal contributions (13.145), (13.146), and (13.148). Divide by $\rho_{\rm tot}$. □

Remark 13.98 (On regulators and comoving normalization)**.** The comoving factor ${\rm Vol}(\mathcal{X}_q)$ cancels in Ω–ratios. Different admissible regulators or slicings induce the same abundance fractions provided the BRST–physical projectors and the HSP filter are used consistently (Sections 2.4 and 13.1.4).

13.4.8 From Spectral Measures to Cosmological Abundances

Definition 13.99 (Spectral partition and cosmological abundance ratios)**.** Insert the holosymmetric projectors on the BRST–physical space into the spectral definition of ρ_{tot} (13.144) (cf. Section 13.1.4). Let

$$\mathcal{P}_{\mathrm{vac}} := \mathcal{P}^{\mathrm{sym}}_{\mathrm{vac}}, \qquad \mathcal{P}_{\mathrm{dark}} := \mathcal{P}^{\mathrm{coh}}_{\mathrm{dark}}, \qquad \mathcal{P}_{\mathrm{vis}} := \mathcal{P}^{\mathrm{coh}}_{\mathrm{vis}},$$

and denote generically $\mathcal{P}_i \in \{\mathcal{P}_{\mathrm{vac}}, \mathcal{P}_{\mathrm{dark}}, \mathcal{P}_{\mathrm{vis}}\}$. Then

$$\begin{aligned} \rho_{\mathrm{tot}} &= \rho_{\mathrm{DE}} + \rho_{\mathrm{DM}} + \rho_{\mathrm{vis}} + \delta_{\mathrm{cross}}, \\ \delta_{\mathrm{cross}} &:= \frac{1}{\mathrm{Vol}(\mathcal{X}_q)} \Big\langle \mathring{\mathbf{\Psi}} \Big| \sum_{i \neq j} \mathcal{P}_i \, \hat{H}_{\mathrm{univ}} \, \mathcal{P}_j \, \Big| \mathring{\mathbf{\Psi}} \Big\rangle_{\mathrm{phys}}, \end{aligned} \tag{13.151}$$

with ρ_{DE}, ρ_{DM}, ρ_{vis} as in (13.145), (13.146), (13.148). Stage II holosymmetric orthogonality on the BRST–physical space (spectral decoupling of inequivalent K–actions) suppresses the off–diagonal terms so that $\delta_{\mathrm{cross}} \simeq 0$. In the *ideal HSP limit*—and, equivalently, whenever the induced dynamics is block–diagonal with respect to $\{\mathcal{P}_i\}$—the projectors are mutually orthogonal and resolve the identity (Proposition 13.97), giving $\delta_{\mathrm{cross}} = 0$ exactly. Consequently, the abundance fractions are computable spectral ratios:

$$\begin{aligned} \Omega_{\mathrm{DE}} &= \frac{\rho_{\mathrm{DE}}}{\rho_{\mathrm{tot}}} = \frac{\varepsilon_\Lambda}{\rho_{\mathrm{tot}}}, \\ \Omega_{\mathrm{DM}} &= \frac{\rho_{\mathrm{DM}}}{\rho_{\mathrm{tot}}}, \\ \Omega_{\mathrm{vis}} &= \frac{\rho_{\mathrm{vis}}}{\rho_{\mathrm{tot}}} = 1 - \Omega_{\mathrm{DM}} - \Omega_{\mathrm{DE}} \quad (\delta_{\mathrm{cross}} = 0). \end{aligned} \tag{13.152}$$

More generally, defining $\Omega_{\mathrm{cross}} := \delta_{\mathrm{cross}}/\rho_{\mathrm{tot}}$, one has

$$1 = \Omega_{\mathrm{DE}} + \Omega_{\mathrm{DM}} + \Omega_{\mathrm{vis}} + \Omega_{\mathrm{cross}}, \qquad \Omega_{\mathrm{vis}} = 1 - \Omega_{\mathrm{DE}} - \Omega_{\mathrm{DM}} - \Omega_{\mathrm{cross}}.$$

Convention (used elsewhere). If the dark sector is realized by the Chrono–Kymon vector, write $\rho_{\mathrm{DM}} \equiv \rho_{K^{(\tau)}}$ and $\Omega_{\mathrm{DM}} \equiv \Omega_{K^{(\tau)}}$.

13.4.9 Relation to the Chronogeometric Regimes

Each component naturally dominates in a distinct chrono–geometric epoch (cf. Section 11.9):

(i) Vacuum–dominated regime $\Omega_{\mathrm{DE}} \to 1$ toward the asymptotic future, with acceleration driven by the interferometric residue (cf. Sections 13.2.6 and 13.2.7).

(ii) Structure–formation regime Ω_{DM} approaches dominance across the linear/nonlinear growth era (dark sector admitted by HSP Stage I and refined by Stage II; cf. Section 13.1.4).

(iii) Radiation/baryon–dominated regime Ω_{vis} dominates in the hot, dense early universe where SM processes set the budget.

Thus cosmic history is a *dynamical partition* driven by holosymmetric coherence: the emergent spectral content of the universal Holo–State $\left|\mathring{\mathbf{\Psi}}\right\rangle$ selects which sector dominates when, linking interferometric coherence to large–scale evolution (cf. Sections 2.4 and 13.3).

Definition 13.100 (Route A abundances (background–agnostic))**.** Let u^μ be any admissible unit timelike congruence on $(\mathcal{M}, \bar{\Sigma}_{\mu\nu})$ and let λ label a foliation by Cauchy slices orthogonal to u^μ. With projector–filtered energy densities $\rho_{\rm DE}(\lambda)$, $\rho_{\rm DM}(\lambda)$, $\rho_{\rm vis}(\lambda)$ defined by eqs. (13.145), (13.146) and (13.148), set

$$\Omega_{\rm DE}(\lambda) = \frac{\rho_{\rm DE}(\lambda)}{\rho_{\rm tot}(\lambda)}, \qquad \Omega_{\rm DM}(\lambda) = \frac{\rho_{\rm DM}(\lambda)}{\rho_{\rm tot}(\lambda)}, \qquad \Omega_{\rm vis}(\lambda) = \frac{\rho_{\rm vis}(\lambda)}{\rho_{\rm tot}(\lambda)},$$

with $\rho_{\rm tot}(\lambda) = \rho_{\rm DE}(\lambda)+\rho_{\rm DM}(\lambda)+\rho_{\rm vis}(\lambda)$ when cross–sector terms vanish (Stage II orthogonality; cf. Proposition 13.97). More generally, if $\delta_{\rm cross}(\lambda) \neq 0$ define

$$1 = \Omega_{\rm DE}(\lambda) + \Omega_{\rm DM}(\lambda) + \Omega_{\rm vis}(\lambda) + \Omega_{\rm cross}(\lambda), \qquad \Omega_{\rm cross}(\lambda) := \frac{\delta_{\rm cross}(\lambda)}{\rho_{\rm tot}(\lambda)}$$

(cf. Definition 13.99).

Specialization (FLRW). If one *chooses* a spatially flat FLRW epoch, take $\lambda \leftrightarrow \ln a$ (or cosmic time), and retrieve the familiar $\Omega_i(a)$ by replacing $\lambda \mapsto a$.
Convention. When the dark sector is realized by the Chrono–Kymon vector, we write $\rho_{\rm DM} \equiv \rho_{K^{(\tau)}}$ and $\Omega_{\rm DM} \equiv \Omega_{K^{(\tau)}}$.

Corollary 13.101 (Critical–density closure check)**.** (General, no geometric prior). *In the ideal HSP limit,*

$$\Omega_{\rm DE}(\lambda) + \Omega_{\rm DM}(\lambda) + \Omega_{\rm vis}(\lambda) = 1,$$

while for nonzero cross–sector leakage, $1-\Omega_{\rm cross}(\lambda)$ *replaces the right–hand side. Discrepancies indicate regulator choices or projector tuning require refinement.*

(FLRW specialization). *If a flat FLRW background is adopted, the critical energy density is* $\rho^E_{\rm crit}(a) = \frac{3c^2H^2(a)}{8\pi G}$ *and, at* $a = 1$*, one may cross–check that*

$$\Omega_i^{(\rm spec)}(1) = \frac{\rho_i(1)}{\rho_{\rm tot}(1)} \quad \textit{matches} \quad \Omega_i^{(\rm crit)}(1) = \frac{\rho_i(1)}{\rho^E_{\rm crit}(1)},$$

within tolerance (see [146, 7, 144]).

13.5 Numerical Schemes for Abundance Determination

To determine the relative abundances of dark energy ($\Omega_{\rm DE}$), dark matter ($\Omega_{\rm DM}$), and visible matter ($\Omega_{\rm vis}$), we present two distinct but complementary methods. The first, which we call the *spectral route*, is a fundamental approach derived directly from the theory's "state-first" principles. The second, the *effective-field route*, is a practical simplification designed for efficient computation and comparison with observational data. Both methods are built to be fully consistent with our framework's foundational projector definitions and the holosymmetric sum rule (Section 13.4.7 and Proposition 13.97).

13.5.1 Route A. Spectral–First (State–First) Computation

Definition 13.102 (Inputs for the projector computation)**.** Let $(\pi_{\mathrm{phys}}, \mathcal{H}_{\mathrm{phys}}, |\Omega_{\mathrm{phys}}\rangle)$ be the GNS/BRST physical representation of the primal state (cf. Section 13.6). Assume:

(i) Regulated spectral data / local correlators
Regulated spectral data (or local correlators) sufficient to determine the expectation values of the Hamiltonian pieces,

$$\left\langle \hat{H}_{\mathrm{vac}} \right\rangle, \qquad \left\langle \hat{H}_{\mathrm{dark}} \right\rangle, \qquad \left\langle \hat{H}_{\mathrm{vis}} \right\rangle,$$

evaluated on $|\Omega_{\mathrm{phys}}\rangle$ with the regulators prescribed by the chrono–geometric regimes (cf. Sections 4.9.15 and 11.9 and Principle 4.1).

(ii) HSP coherence projectors and comoving volume
Holosymmetric Selection (HSP) coherence projectors $\mathcal{P}^{\mathrm{sym}}_{\mathrm{vac}}$, $\mathcal{P}^{\mathrm{coh}}_{\mathrm{dark}}$, $\mathcal{P}^{\mathrm{coh}}_{\mathrm{vis}}$ (Stages I/II; cf. Section 13.1.4) and a regulated comoving volume $\mathrm{Vol}(\mathcal{X}_q)$ (any admissible foliation; ratios are regulator–independent).

Definition 13.103 (Sectoral energy densities)**.** With $\langle\cdot\rangle = \langle\Omega_{\mathrm{phys}}|\cdot|\Omega_{\mathrm{phys}}\rangle$ and $\mathrm{Vol} = \mathrm{Vol}(\mathcal{X}_q)$, define

$$\begin{aligned}
\rho_{\mathrm{DE}} &= \frac{1}{\mathrm{Vol}} \left\langle \mathcal{P}^{\mathrm{sym}}_{\mathrm{vac}}\, \hat{H}_{\mathrm{vac}}\, \mathcal{P}^{\mathrm{sym}}_{\mathrm{vac}} \right\rangle, \\
\rho_{\mathrm{DM}} &= \frac{1}{\mathrm{Vol}} \left\langle \mathcal{P}^{\mathrm{coh}}_{\mathrm{dark}}\, \hat{H}_{\mathrm{univ}}\, \mathcal{P}^{\mathrm{coh}}_{\mathrm{dark}} \right\rangle, \\
\rho_{\mathrm{vis}} &= \frac{1}{\mathrm{Vol}} \left\langle \mathcal{P}^{\mathrm{coh}}_{\mathrm{vis}}\, \hat{H}_{\mathrm{univ}}\, \mathcal{P}^{\mathrm{coh}}_{\mathrm{vis}} \right\rangle,
\end{aligned} \tag{13.153}$$

in agreement with eqs. (13.145), (13.146) and (13.148).

Convention. When the dark sector is realized by the Chrono–Kymon vector, write $\rho_{\mathrm{DM}} \equiv \rho_{K^{(\tau)}}$ and $\Omega_{\mathrm{DM}} \equiv \Omega_{K^{(\tau)}}$.

Proposition 13.104 (Total density and abundance fractions)**.** *If the coherence projectors are mutually orthogonal and resolve the identity on the relevant BRST–physical subspace (HSP Stage II; cf. Proposition 13.97), then*

$$\rho_{\mathrm{tot}} = \rho_{\mathrm{DE}} + \rho_{\mathrm{DM}} + \rho_{\mathrm{vis}}, \qquad \Omega_{\mathrm{DE}} = \frac{\rho_{\mathrm{DE}}}{\rho_{\mathrm{tot}}}, \quad \Omega_{\mathrm{DM}} = \frac{\rho_{\mathrm{DM}}}{\rho_{\mathrm{tot}}}, \quad \Omega_{\mathrm{vis}} = \frac{\rho_{\mathrm{vis}}}{\rho_{\mathrm{tot}}}. \tag{13.154}$$

Non–ideal case. If small cross terms $\delta_{\mathrm{cross}} \neq 0$ remain, write $1 = \Omega_{\mathrm{DE}} + \Omega_{\mathrm{DM}} + \Omega_{\mathrm{vis}} + \Omega_{\mathrm{cross}}$ with $\Omega_{\mathrm{cross}} := \delta_{\mathrm{cross}}/\rho_{\mathrm{tot}}$ (cf. Definition 13.99).

Remark 13.105 (Numerical realization ("doing the numbers"))**.** Evaluating (13.153) reduces to regulated spectral integrals as in Section 17.1.2 for the vacuum and their analogues for the dark and visible sectors, with UV/IR windows fixed by the chrono–geometric regime and the dressing maps (cf. Principle 4.1 and Section 4.9.15). The volume regulator cancels in all Ω–ratios.

Corollary 13.106 (Critical–density cross–check at $z = 0$ (curvature–aware))**.** *Independently compute the critical energy density*

$$\rho^{E}_{\mathrm{crit}}(0) = \frac{3c^2 H_0^2}{8\pi G},$$

and define the background spatial–curvature parameter

$$\Omega_k := -\frac{k\,c^2}{a_0^2 H_0^2} \qquad (a_0 = 1).$$

Then require agreement (within tolerance) between the spectral and critical normalizations:

$$\Omega_i^{(\mathrm{spec})} = \frac{\rho_i}{\rho_{\mathrm{tot}}} \quad \textit{vs.} \quad \Omega_i^{(\mathrm{crit})} = \frac{\rho_i}{\rho^{E}_{\mathrm{crit}}(0)},$$

with the closure condition

$$\sum_i \Omega_i^{(\mathrm{crit})} = 1 - \Omega_k\,.$$

In the flat calibration used elsewhere ($\Omega_k = 0$), this reduces to $\sum_i \Omega_i^{(\mathrm{crit})} = 1$ and hence $\Omega_i^{(\mathrm{spec})} \approx \Omega_i^{(\mathrm{crit})}$. Discrepancies beyond tolerance indicate that regulator choices or projector tuning require refinement.

Justification. *At $a = 1$, the Friedmann equation with curvature, $H_0^2 = \frac{8\pi G}{3c^2}\rho_{\mathrm{tot}}(0) - \frac{kc^2}{a_0^2}$, implies $\rho_{\mathrm{tot}}(0) = \rho^{E}_{\mathrm{crit}}(0)\,[1 - \Omega_k]$ (see [146, 7, 144]).*

13.5.2 Route B. Effective–Field Reduction: a Fast Lane for Numerics

Definition 13.107 (Minimal effective inputs (FLRW–calibrated))**.** Work on a homogeneous, isotropic Robertson–Walker (RW) patch with scale factor a; optionally include spatial curvature via Ω_k (default $\Omega_k = 0$ for the flat calibration). Specify:

(i) Interferometric residue

Interferometric residue ε_Λ (late–time constant to leading order), so $\rho_{\mathrm{DE}}(a) \approx \varepsilon_\Lambda$ (cf. Sections 13.2.7 and 13.4.7).

(ii) Chrono–Kymon background ($U(1)_C$)

characterized by an effective mass μ_C and an initial amplitude $\langle K^{(\tau)}_\mu K^{(\tau)\mu}\rangle_0$:

$$\rho_{K^{(\tau)}}(a) \simeq \rho_{K^{(\tau)},0}\,a^{-3}, \qquad \rho_{K^{(\tau)},0} = \tfrac{1}{2}\,\mu_C^2 \Big\langle K^{(\tau)}_\mu K^{(\tau)\mu}\Big\rangle_0 + (\text{small kinetic}),$$

with interaction/dressing corrections $\mathcal{O}(\alpha_C)$ that preserve the a^{-3} scaling at background level (cf. Sections 4.9.15 and 13.3).

(iii) Visible sector

$\rho_r(a) = \rho_{r,0}a^{-4}$ (photons + neutrinos) and $\rho_b(a) = \rho_{b,0}a^{-3}$ (baryons), from the SM projector or cross–calibrated values (Section 13.4.7).

Remark 13.108 (Curvature toggle)**.** With $\Omega_k \neq 0$ the background relation used downstream becomes

$$E^2(a) = \Omega_r a^{-4} + \Omega_m a^{-3} + \Omega_{\mathrm{DE}} a^{-3(1+w_{\mathrm{DE}})} + \Omega_k a^{-2},$$

reducing to the flat calibration when $\Omega_k = 0$.

Definition 13.109 (Assembly and normalization)**.** Given H_0 (or solving for it self–consistently), set

$$\rho_{\rm tot}(a) = \rho_{\rm DE}(a) + \rho_{\rm DM}(a) + \rho_{\rm vis}(a), \qquad \Omega_i(a) = \frac{\rho_i(a)}{\rho_{\rm tot}(a)} \quad (i \in \{\rm DE, DM, vis\}). \tag{13.155}$$

At $a = 1$ the sum rule $\sum_i \Omega_i(1) = 1$ serves as a numerical diagnostic (it is automatic in the ideal HSP limit; cf. Proposition 13.97).

Proposition 13.110 (From abundances to observables)**.** *Let $E(a) := H(a)/H_0$ and define $\kappa' \equiv 8\pi G$. On a spatially flat FLRW background,*

$$E^2(a) = \frac{\kappa' c^2}{3H_0^2}\,\rho_{\rm tot}(a) = \frac{\rho_{\rm tot}(a)}{\rho_{\rm tot}(1)}\,, \tag{13.156}$$

where the second equality uses the $a = 1$ normalization $\rho_{\rm tot}(1) = 3H_0^2/(\kappa' c^2)$. This $E(a)$ fixes the standard RW distance–redshift relations and the linear–growth equation. Small departures from ΛCDM map directly to temporal–sector parameters (e.g. a small DE tilt $w_{\rm DE}+1$, the Chrono–Kymon mass μ_C, or mild coherence corrections), providing a compact link from tau–kairos inputs to observables (cf. Sections 13.3 and 13.4.7).

Remark 13.111 (Curvature–aware form)**.** Allowing spatial curvature with $\Omega_k := -kc^2/(a_0^2 H_0^2)$ and $a_0 = 1$,

$$E^2(a) = \Omega_k\, a^{-2} + \frac{\kappa' c^2}{3H_0^2}\,\rho_{\rm tot}(a) = \Omega_k\, a^{-2} + \Big(1 - \Omega_k\Big)\frac{\rho_{\rm tot}(a)}{\rho_{\rm tot}(1)}\,, \tag{13.157}$$

which reduces to (13.156) when $\Omega_k = 0$. Distances then follow the usual RW expressions with curvature (see [144, 601]).

Remark 13.112 (Equivalence of Routes A and B)**.** Route A implements the projector definitions of Section 13.4.7; Route B is their calibrated FLRW reduction with parameters $\{\varepsilon_\Lambda, \mu_C, \langle K_\mu^{(\tau)} K^{(\tau)\mu}\rangle_0, \rho_{r,0}, \rho_{b,0}\}$. Agreement at the level of $E(a)$, distances, and linear growth provides an internal cross–check on regulators and on HSP sector selection (cf. Section 13.5.9). TT–purity of GW (Definition 13.78) justifies the use of the standard FLRW background relations in this calibration lane.

Convention. $\rho_{\rm DM}(a) \equiv \rho_{K^{(\tau)}}(a)$ and $\Omega_{\rm DM}(a) \equiv \Omega_{K^{(\tau)}}(a)$ (Chrono–Kymon dark matter).

13.5.3 Linear–Scale Structure with Chrono–Kymon Coherence (Effective)

Definition 13.113 (Effective linear matter power with Chrono–Kymon kernel)**.** For fast forecasts in the calibration lane, define

$$P_{\rm lin}(k, z; \mu_C) := A_s \left(\frac{k}{k_\star}\right)^{n_s - 1} T_{\rm vis}^2(k)\, \mathcal{S}^2\Big(k;\, k_c(\mu_C), \nu\Big)\, D^2(z), \tag{13.158}$$

where $T_{\rm vis}(k)$ is the standard visible–sector transfer function, $D(z)$ is the GR growth factor normalized by $D(0) = 1$ and determined by $E(a)$ as in Section 13.5.2, and $\mathcal{S}(k; k_c, \nu)$ is the *Chrono–Kymon coherence kernel.*

Definition 13.114 (Chrono–Kymon coherence kernel: axioms)**.** The kernel $\mathcal{S}(k;\, k_c, \nu)$ summarizes linear–scale imprints of vector temporal coherence, subject to:

(i) Projector/propagator compatibility
$\mathcal{S}$ modifies only the scalar/vector matter channel and never the GW TT sector (TT isolation; cf. Definition 13.78).

(ii) Normalization and positivity
$0 < \mathcal{S} \leq 1$, with $\mathcal{S}(0) = 1$.

(iii) Coherence scale
$k_c = k_c(\mu_C; \bar{\Sigma})$ is a monotonically increasing function of the Chrono–Kymon mass μ_C (and, in general, the background $(\mathcal{M}, \bar{\Sigma}_{\mu\nu})$), setting the transition between unsuppressed and coherence–suppressed modes.

(iv) Asymptotics
$\lim_{k \ll k_c} \mathcal{S} = 1$; for $k \gg k_c$ the envelope decays at least as a power law, $\mathcal{S} \sim (k/k_c)^{-\nu}$ with $\nu > 0$ (model–dependent).

(v) Calibration limit
$\mathcal{S} \to 1$ when (i) $\mu_C \to \infty$ (or $k_c \to \infty$), or (ii) vector coherence is decohered by HSP/portal hygiene, recovering CDM.

Remark 13.115 (Example parameterization)**.** A convenient Padé–like model respecting (K1)–(K5) is $\mathcal{S}(k;\, k_c, \nu, \eta) = \left[1 + (k/k_c)^{\nu}\right]^{-\eta}$, with $\eta > 0$. Forecasts should explore priors on (k_c, ν, η) implied by μ_C and the dressing maps; the calibration limit is $\eta \to 0$ or $k_c \to \infty$.

Proposition 13.116 (ΛCDM calibration)**.** *Under the calibration limit of Definition 13.114 (K5),* $\mathcal{S} \to 1$ *and the power spectrum* (13.158) *reduces to the standard CDM form* $P_{\rm lin}(k, z) = A_s (k/k_\star)^{n_s - 1} T^2_{\rm vis}(k) D^2(z)$.

Proof. Immediate from (13.158) and (K5), noting that $T_{\rm vis}$ and D are unchanged by TT–isolated temporal dynamics (cf. Definition 13.78 and Section 13.5.2). □

Remark 13.117 (Spacetime setting & curvature generalization)**.** By default, (13.158) is evaluated on a late–time, spatially flat FLRW patch. For $K \neq 0$, replace plane waves by Laplace–Beltrami eigenmodes on Σ_K and use the curved–space measure; the kernel $\mathcal{S}$ and the GR growth equation keep the same functional form, with curvature entering through $E(a)$ and the mode spectrum.

Remark 13.118 (Forecast recipe (fast lane))**. (i) Fix background parameters**
Fix

$$(\varepsilon_\Lambda,\ \mu_C,\ \langle K^{(\tau)}_\mu K^{(\tau)\mu} \rangle_0,\ \rho_{r,0},\ \rho_{b,0},\ H_0)$$

per Section 13.5.2 and verify closure at $a = 1$.

(ii) Background and growth
Compute $E(a)$ from $\rho_{\rm tot}(a)$, then $D(z)$ from the GR growth equation.

(iii) Coherence kernel and power spectrum
Choose $\mathcal{S}$ consistent with Definition 13.114 (e.g. Remark 13.115) and evaluate (13.158); set $\mathcal{S} = 1$ for the calibration/benchmark.

Why this calibration is legitimate. (i) The TT channel remains purely geometric (no new spin–2), so $E(a)$, distances, and linear growth obey the textbook RW forms used by existing pipelines; (ii) the calibration procedure prescribes fixing $\{\rho_{r,0}, \rho_{b,0}\}$, H_0, normalizing $\rho_{K^{(\tau)},0}$ and ε_Λ, and checking closure; (iii) curvature and non–flat FLRW generalizations are accommodated by Remark 13.117. Hence Route A remains projector–exact and chronogeometric, while Route B provides a strict FLRW specialization suitable for fast data confrontation.

13.5.4 Effective-Field Route for Abundances (Route B)

Definition 13.119 (Minimal Effective Parameter Set)**.** Work on a spatially flat Friedmann–Lemaître–Robertson–Walker (FLRW) patch with scale factor a, normalized by $a_0 = 1$ [146, 7, 14]. Define the effective parameter set

$$\Xi_{\text{eff}} := \Big\{\varepsilon_\Lambda,\ \mu_C,\ \langle K^{(\tau)}_\mu K^{(\tau)\mu}\rangle_0,\ \rho_{r,0},\ \rho_{b,0},\ H_0\Big\}.$$

Here: ε_Λ is the interferometric vacuum residue (Sections 13.2.7 and 13.4.7), anchored in the observed dark-energy density [211, 136]; μ_C and $\left\langle K^{(\tau)}_\mu K^{(\tau)\mu}\right\rangle_0$ are the Chrono–Kymon (dark $U(1)_C$) effective mass and background amplitude fixed by the dressing map (Sections 4.9.15 and 13.3), cf. dark-matter phenomenology [588, 586]; $\rho_{r,0}, \rho_{b,0}$ are present–day radiation and baryon energy densities (either computed by the SM projector or cross–calibrated) [581, 136]; H_0 is the present Hubble rate [581].

Remark 13.120 (Scope within the chronogeometric framework)**.** Definition 13.119 is an *adopted* late–time, homogeneous–isotropic coarse-graining used for data work. The chronogeometric/emergent content enters through the parameters ε_Λ, $\{\mu_C, \left\langle K^{(\tau)}_\mu K^{(\tau)\mu}\right\rangle_0\}$ and the mapping $\Lambda_{\text{eff}} = \Lambda + \kappa'\varepsilon_\Lambda$; no assumption of a fundamental, non-emergent spacetime is made here.

Definition 13.121 (Sectoral effective energy densities (Route B))**.** With κ' the gravitational coupling of Section 17.1.1, define

$$\boxed{\rho_{\text{DE}}(a) = \varepsilon_\Lambda\ \exp\Big(-3\int_1^a [1 + w_{\text{DE}}(\tilde a)]\, d\ln\tilde a\Big)} \tag{13.159}$$

$$w_{\text{DE}}(a) = -1 + \epsilon(a), \qquad |\epsilon(a)| \ll 1. \tag{13.160}$$

$$\rho_{K^{(\tau)}}(a) = \rho_{K^{(\tau)},0}\, a^{-3}\,[1 + \delta_{K^{(\tau)}}(a)]. \tag{13.161}$$

$$\rho_{K^{(\tau)},0} := \tfrac{1}{2}\,\mu_C^2\left\langle K^{(\tau)}_\mu K^{(\tau)\mu}\right\rangle_0 \ +\ \text{(small kinetic)}. \tag{13.162}$$

$$\rho_{\text{vis}}(a) = \rho_{b,0}\, a^{-3} + \rho_{r,0}\, a^{-4}. \tag{13.163}$$

Setting $\epsilon(a) = 0$ and $\delta_{K^{(\tau)}}(a) = 0$ reproduces ΛCDM at the background level: $\rho_{\text{DE}}(a) = \varepsilon_\Lambda$ and $\rho_{K^{(\tau)}}(a) = \rho_{K^{(\tau)},0}\, a^{-3}$.

Remark 13.122 (Scope within the chronogeometric framework)**.** Equations (13.159)–(13.163) are the late–time, homogeneous–isotropic (*adopted* FLRW patch) effective description used for numerics. The chronogeometric (emergent) input is encoded in the parameters and identifications ε_Λ (interferometric residue), the Chrono–Kymon sector $\{\mu_C, \langle K_\mu^{(\tau)} K^{(\tau)\mu}\rangle_0, \delta_{K^{(\tau)}}\}$, and the mapping $\Lambda_{\rm eff} = \Lambda + \kappa'\varepsilon_\Lambda$. No assumption of a fundamental, non-emergent spacetime is made here; this is the coarse-grained limit of the framework's dynamics.

Remark 13.123 (Provenance: textbook vs. treatise-specific)**. Standard (textbook).**

(i) Continuity solution
Continuity solution: $\dot\rho + 3H(\rho + p) = 0 \Rightarrow \rho(a) = \rho_0 \exp\Big(-3\int[1 + w(\tilde a)]\, d\ln\tilde a\Big)$, giving the minus sign in eq. (13.159) [144, 327, 319].

(ii) Cold matter & radiation
Cold matter & radiation: $\rho_{\rm m} \propto a^{-3}$, $\rho_r \propto a^{-4}$, yielding eq. (13.163) [144, 327].

(iii) Equation of state
Equation of state: $w_{\rm DE} = p_{\rm DE}/\rho_{\rm DE}$ (here written as a small deviation from -1) [144].

(iv) Normalization
Normalization: $a_0 = 1$ today [144].

Treatise-specific (this work).

(i) Vacuum residue
ε_Λ is the interferometric residual set by holosymmetric selection (Sections 13.2.7 and 13.4.7).

(ii) DE tilt
$w_{\rm DE}(a) = -1 + \epsilon(a)$ with small, smooth ϵ is the minimal late-time ansatz (Route B).

(iii) Chrono–Kymon DM
$\rho_{K^{(\tau)}}(a)$ models the vector temporal (Chrono–Kymon) sector; $\delta_{K^{(\tau)}}(a) = \mathcal{O}(\alpha_C)$ encodes small, portal-free self-interaction corrections.

(iv) Rest-mass normalization
$\rho_{K^{(\tau)},0}$ dominated by $\frac{1}{2}\mu_C^2\langle K_\mu^{(\tau)} K^{(\tau)\mu}\rangle_0$; the "small kinetic" term is cycle-averaged over $2\pi/\mu_C \ll \Delta t \ll H^{-1}$.

Convention. When the dark sector is realized by the Chrono–Kymon vector, we write $\rho_{\rm DM} \equiv \rho_{K^{(\tau)}}$ and $\Omega_{\rm DM} \equiv \Omega_{K^{(\tau)}}$.

Definition 13.124 (Total density, abundances, and expansion history)**.**

$$\rho_{\rm tot}(a) = \rho_{\rm DE}(a) + \rho_{K^{(\tau)}}(a) + \rho_{\rm vis}(a). \tag{13.164}$$

$$\Omega_i(a) := \frac{\rho_i(a)}{\rho_{\rm tot}(a)}, \qquad i \in \{\mathrm{DE}, K^{(\tau)}, \mathrm{vis}\}. \tag{13.165}$$

Index set. $i = \mathrm{DE}$ denotes dark energy; $i = K^{(\tau)}$ denotes the Chrono–Kymon (vector temporal) dark–matter sector; $i = \mathrm{vis}$ denotes the visible sector (baryons + radiation). The

ratios in (13.165) recall the canonical definition in eq. (13.142), here evaluated at scale factor a using Definition 13.121. *Convention.* $\rho_{\mathrm{DM}}(a) \equiv \rho_{K^{(\tau)}}(a)$ and $\Omega_{\mathrm{DM}}(a) \equiv \Omega_{K^{(\tau)}}(a)$.

$$\sum_{i \in \{\mathrm{DE}, K^{(\tau)}, \mathrm{vis}\}} \Omega_i(1) = 1, \tag{13.166}$$

(cf. Proposition 13.97; see, e.g., [146, 7, 240]).

Proposition 13.125 (First Friedmann Equation (Spatially Flat FLRW))**.** *On a spatially flat FLRW background with $H(a) := \dot{a}/a$,*

$$H^2(a) = \frac{\kappa' c^2}{3}\, \rho_{\mathrm{tot}}(a), \tag{13.167}$$

where $\rho_{\mathrm{tot}}(a)$ is given by Definition 13.124. Equivalently,

$$E(a) := \frac{H(a)}{H_0} = \sqrt{\frac{\rho_{\mathrm{tot}}(a)}{\rho_{\mathrm{tot}}(1)}},$$

and at $a = 1$ the (flat-universe) closure relation holds

$$\sum_{i \in \{\mathrm{DE}, K^{(\tau)}, \mathrm{vis}\}} \Omega_i(1) = 1,$$

cf. Proposition 13.97; see, e.g., [146, 7, 240].

Definition 13.126 (Dimensionless Hubble Function)**.** The *dimensionless Hubble function* (expansion history) is

$$E(a) \;:=\; \frac{H(a)}{H_0} \;=\; \sqrt{\frac{\rho_{\mathrm{tot}}(a)}{\rho_{\mathrm{tot}}(1)}}\,, \tag{13.168}$$

where $H(a) = \dot{a}/a$ is the Hubble rate at scale factor a, $H_0 := H(1)$ is its present value, and $\rho_{\mathrm{tot}}(a)$ is the total energy density from Definition 13.124. The equality follows from the first Friedmann equation in the spatially flat FLRW case (Proposition 13.125); see, e.g., [146, 7, 240]. By construction $E(1) = 1$, and the dependence on the gravitational coupling cancels in the ratio.

Remark 13.127 (Use)**.** $E(a)$ fully determines background observables: comoving/luminosity/angular–diameter distances via $\chi(z) = \frac{c}{H_0}\int_0^z dz'/E(z')$, and the linear–growth "friction" term $d\ln H/d\ln a$ in the matter–perturbation equation (cf. Corollary 13.132).

Remark 13.128 (Dark–Energy Tilt)**.** Let the dark energy (DE) equation of state deviate slightly from a cosmological constant by $\epsilon := w_{\mathrm{DE}} + 1$ with $|\epsilon| \ll 1$. Then

$$\rho_{\mathrm{DE}}(a) = \rho_{\mathrm{DE},0}\, a^{-3(1+w_{\mathrm{DE}})} = \rho_{\mathrm{DE},0}\, a^{-3\epsilon} \simeq \rho_{\mathrm{DE},0}\left(1 - 3\epsilon \ln a\right).$$

To first order this induces

$$E^2(a) = \Omega_r a^{-4} + \Omega_m a^{-3} + \Omega_{\mathrm{DE}}\, a^{-3(1+\epsilon)} \quad \Rightarrow \quad \left.\frac{\delta E}{E}\right|_{\epsilon} \simeq -\,\frac{3\,\epsilon\,\Omega_{\mathrm{DE}}\, a^{-3}\ln a}{2\left[\Omega_r a^{-4} + \Omega_m a^{-3} + \Omega_{\mathrm{DE}} a^{-3}\right]}\,.$$

Sign: $\epsilon > 0$ (quintessence-like) slightly reduces distances and growth (lower $f\sigma_8$); $\epsilon < 0$ (phantom-like) increases them. Setting $\epsilon = 0$ recovers the constant-Λ limit.

Proposition 13.129 (First–order cosmological consequences of a small DE tilt)**.** *Assume a spatially flat FLRW background and the hypotheses of Remark 13.128, i.e. dark energy (DE) has $w_{\rm DE} = -1 + \epsilon$ with $|\epsilon| \ll 1$, and the tensor (gravitational waves, GW) channel remains purely transverse–traceless (TT). Then, to first order in ϵ:*

(i) Background expansion. *Using Remark 13.128, the fractional change of the Hubble function is*

$$\left.\frac{\delta E}{E}\right|_{\epsilon} \;\simeq\; -\,\frac{3\,\epsilon\,\Omega_{\rm DE}\,a^{-3}\ln a}{2\left[\Omega_r a^{-4} + \Omega_m a^{-3} + \Omega_{\rm DE} a^{-3}\right]}\,, \tag{13.169}$$

with the sign interpretation as in Remark 13.128. (For constant-w scaling and FLRW background relations see, e.g., [144].)

(ii) Distance measures (SN/BAO). *For comoving distance $\chi(z) = \frac{c}{H_0}\int_0^z dz'\,E(z')^{-1}$,*

$$\frac{\delta\chi}{\chi} \;\approx\; -\frac{\displaystyle\int_0^z dz'\,\big[\delta E/E\big](z')}{\displaystyle\int_0^z dz'\,E(z')^{-1}} \quad\Rightarrow\quad \frac{\delta D_L}{D_L} = \frac{\delta D_A}{D_A} = \frac{\delta\chi}{\chi}\,, \tag{13.170}$$

implying percent-level shifts for $|\epsilon| \sim$ few$\times 10^{-2}$ over low/intermediate z (textbook distance relations: [144]).

(iii) Growth of structure (RSD/weak lensing). *The linear growth equation retains its GR form $\delta'' + [2 + d\ln H/d\ln a]\delta' - \frac{3}{2}\,\Omega_m(a)\,\delta = 0$ (cf. Corollary 13.132; see [327, 319]). A convenient fit is*

$$f(a) \equiv \frac{d\ln D}{d\ln a} \;\simeq\; \Omega_m(a)^{\gamma}, \qquad \gamma \simeq 0.55 + 0.05\,[1 + w_{\rm DE}(z{=}1)]\,, \tag{13.171}$$

so $\epsilon > 0$ slightly suppresses $f\sigma_8(z)$ while $\epsilon < 0$ slightly enhances it (for ΛCDM $w{=}-1$ one has $\gamma \approx 0.55$; cf. standard treatments [327, 319]).

(iv) Timing of matter–DE equality / onset of acceleration. *Equality $\rho_m(a) = \rho_{\rm DE}(a)$ occurs at*

$$a_{\rm eq}^{(m-DE)} = \left(\frac{\Omega_m}{\Omega_{\rm DE}}\right)^{1/[3(1-\epsilon)]} \;\approx\; a_{\rm eq}^{(\Lambda)}\,\exp\!\Big[\epsilon\,\ln a_{\rm eq}^{(\Lambda)}\Big], \qquad a_{\rm eq}^{(\Lambda)} = \left(\frac{\Omega_m}{\Omega_{\rm DE}}\right)^{1/3}, \tag{13.172}$$

with the sign interpretation following from $\ln a_{\rm eq}^{(\Lambda)} < 0$ (see, e.g., [144]).

(v) Late–ISW and CMB lensing (qualitative, $\mathcal{O}(\epsilon)$). *A positive tilt ($\epsilon > 0$) mildly boosts late ISW and tends to lower lensing-inferred clustering; $\epsilon < 0$ does the reverse (standard background physics: [144]).*

(vi) Degeneracies and constraints. *Effects above partially degenerate with Ω_m, H_0, and small-scale DM suppression entering σ_8. Joint SN+BAO+CMB+RSD analyses (e.g. Planck 2018) typically constrain $|\epsilon|$ to sub-percent/percent levels [150].*

(vii) GW channel unchanged. *The TT projector constraint keeps only the two TT GW polarizations; the tilt introduces no additional GW modes (background GR phenomenology as in [144]; model-specific projector statement via Section 13.3).*

Proposition 13.130 (Effective Cosmological Constant from the Vacuum Residue)**.** *The dark–energy parameter that enters the geometric field equations as the* effective cosmological constant *is*

$$\Lambda_{\text{eff}} = \Lambda + \kappa' \, \varepsilon_\Lambda, \tag{13.173}$$

where Λ is the bare cosmological constant, ε_Λ is the interferometric residual vacuum energy density, and κ' is the gravitational coupling (recovering GR with $\kappa' c^2 = 8\pi G$). For $w_{\text{DE}} \equiv -1$ one has $\rho_{\text{DE}}(a) = \varepsilon_\Lambda$ (constant) and Λ_{eff} is constant, reproducing the standard ΛCDM background (cf. eqs. (17.2) and (17.5)).

Remark 13.131. If $w_{\text{DE}}(a) \neq -1$, the *time-varying* part of dark energy is treated in $T^{\mu}{}_{\nu}$ (not absorbed into Λ_{eff}); only the constant vacuum residue ε_Λ renormalizes Λ via (13.173).

Corollary 13.132 (Observable Relations)**.** *Given $E(a)$ from eq. (13.167),*

$$\chi(z) = \frac{c}{H_0} \int_0^z \frac{dz'}{E(z')}, \qquad D_L(z) = (1+z)\,\chi(z), \qquad D_A(z) = \frac{\chi(z)}{1+z}, \tag{13.174}$$

$$\delta''(a) + \left[2 + \frac{d\ln H}{d\ln a}\right]\delta'(a) - \frac{3}{2}\,\Omega_m(a)\,\delta(a) = 0, \qquad \Omega_m(a) := \frac{\rho_b(a) + \rho_{K^{(\tau)}}(a)}{\rho_{\text{tot}}(a)}, \tag{13.175}$$

where primes denote $d/d\ln a$. Equation (13.175) holds in the present framework because the TT gravitational channel is unmodified and the temporal sector does not introduce extra long–range spin–2 forces (cf. Section 13.3).

Remark 13.133 (Calibration and Consistency)**.** In practice one fixes $\{\rho_{r,0}, \rho_{b,0}\}$ by SM projectors or external calibration, chooses H_0, computes $\rho_{K^{(\tau)},0}$ from $\{\mu_C, \langle K^{(\tau)}_\mu K^{(\tau)\mu}\rangle_0\}$, and sets ε_Λ by the regulated spectral self–regulation (Section 17.1.2). One then evaluates $\Omega_i(1)$ and checks $\sum_i \Omega_i(1) = 1$; departures indicate regulator or parameter mis–tuning. The special case $\epsilon = \delta_{K^{(\tau)}} = 0$ yields exactly ΛCDM at the background level, consistent with Section 17.1.4.

What is standard (textbook) vs. what is internal to this treatise. *Standard cosmology (textbook).*

(i) Distance–redshift relations Distance–redshift relations in (13.174) for a spatially flat Friedmann–Lemaître–Robertson–Walker (FLRW) universe:

$$\chi(z) = \frac{c}{H_0}\int_0^z \frac{dz'}{E(z')}, \qquad D_L(z) = (1+z)\chi(z), \qquad D_A(z) = \frac{\chi(z)}{1+z}.$$

These follow from the FLRW metric and Etherington's distance duality (photon number conservation, null geodesics); see, e.g., [144, 485, 327].

(ii) Linear growth equation Linear growth equation (13.175) in GR on sub-horizon scales for pressureless matter, negligible dark-energy clustering on those scales, and no fifth forces:

$$\delta'' + [2 + d\ln H/d\ln a]\delta' - \tfrac{3}{2}\Omega_m(a)\delta = 0$$

(primes denote $d/d\ln a$); see [144, 485, 327].

(iii) Matter fraction Matter fraction entering the source term:

$$\Omega_m(a) = \frac{\rho_m(a)}{\rho_{\rm tot}(a)}$$

(standard definition).

Internal to this treatise (framework-specific).

(i) Dark-matter identification $\rho_m(a) = \rho_b(a) + \rho_{K^{(\tau)}}(a)$ with the Chrono–Kymon (vector temporal) sector supplying the DM component; hence

$$\Omega_m(a) = \frac{\rho_b(a) + \rho_{K^{(\tau)}}(a)}{\rho_{\rm tot}(a)}$$

(in ΛCDM one would use $\rho_{\rm cdm}$ instead of $\rho_{K^{(\tau)}}$).

(ii) GR growth law applicability The TT projector constraint $\mathcal{P}^{(2)}\mathcal{M}_{\tau h} = 0$ (no extra long-range spin-2 forces) ensures that temporal-sector effects enter via background and scalar/vector perturbations, not via new GW polarizations; cf. Section 13.3.

What each equation describes.

(i) (13.174)**, first relation** $\chi(z) = \dfrac{c}{H_0}\int_0^z \dfrac{dz'}{E(z')}$ is the *comoving radial distance* to redshift z in spatially flat FLRW, expressed via the dimensionless expansion history $E(z) = H(z)/H_0$.

(ii) (13.174)**, second/third** $D_L(z) = (1+z)\chi(z)$ (luminosity distance) and $D_A(z) = \chi(z)/(1+z)$ (angular-diameter distance) follow from distance duality, with $D_L = (1+z)^2 D_A$.

(iii) (13.175) $\delta''(a) + \left[2 + \frac{d\ln H}{d\ln a}\right]\delta'(a) - \frac{3}{2}\Omega_m(a)\delta(a) = 0$ is the *linear matter-growth ODE* in GR: the bracket provides *Hubble friction*, and the last term is the *gravitational clustering* source.

(iv) Definition under (13.175) $\Omega_m(a) = \dfrac{\rho_b(a) + \rho_{K^{(\tau)}}(a)}{\rho_{\rm tot}(a)}$ is the matter fraction used in the source term, with $\rho_{K^{(\tau)}}$ realizing DM in this work.

(v) Primes $' \equiv d/d\ln a$ — logarithmic scale-factor derivatives, convenient for background evolution.

13.5.5 Spectral-First Route for Abundances (Route A): Inputs with Planckian UV Scale

This route implements the projector-based definitions of Section 13.4.7 by evaluating regulated spectral expectations in the GNS/BRST physical representation (cf. Sections 13.6 and 17.1.2).

Definition 13.134 (Model Inputs for Route A (Planck UV Fixed))**.** Let $(\pi_{\mathrm{phys}}, \mathcal{H}_{\mathrm{phys}}, |\Omega_{\mathrm{phys}}\rangle)$ be the GNS/BRST triple. Provide:

(i) Dispersions (background-dressed)

Mode energies for the geometric (Gaiamiti) and material (Tantrakyma) sectors,

$$E_{\mathrm{geo}}(k) = \tfrac{1}{2}\hbar\,\omega_{\mathrm{geo}}(k) + \delta E_{\mathrm{geo}}(k;\bar{\Sigma}), \qquad E_{\mathrm{tym}}(k) = \tfrac{1}{2}\hbar\,\omega_{\mathrm{tym}}(k) + \delta E_{\mathrm{tym}}(k;\bar{\Sigma}),$$

including curvature/dressing corrections determined by the quasi-homogeneous background $\bar{\Sigma}_{\mu\nu}$ (cf. Sections 6.1.10 and 11.9). Minimal baseline: $\omega_{\mathrm{geo}}(k) = c\,k$ and standard relativistic $\omega_{\mathrm{tym}}(k)$, with Hadamard/adiabatic subtraction to order $N = 4$.

(ii) Regime-set cutoffs (Planck UV by construction)

Distinguish the following scales:

(a) **UV (finite entanglement capacity).**Since $\mathcal{M}_{\mathrm{geom}}$ becomes manifest at the Planck threshold (cf. Section 11.9 and Principle 4.1), fix

$$\lambda_{\mathrm{cutoff}} = \ell_{\mathrm{P}} = \sqrt{\frac{\hbar G}{c^3}} \approx 1.616 \times 10^{-35}\ \mathrm{m}.$$

$$k_{\mathrm{UV}} = \lambda_{\mathrm{cutoff}}^{-1} \approx 6.19 \times 10^{34}\ \mathrm{m}^{-1}.$$

(b) **IR (global, cosmological box).**Define the global IR scale by

$$k_{\mathrm{IR}}^{\mathrm{cos}} = \frac{H_0}{c},$$

numerically $k_{\mathrm{IR}}^{\mathrm{cos}} \approx 7.3 \times 10^{-27}\ \mathrm{m}^{-1}$ today (cf. Section 17.1.4).

(c) **IR (local, dense environments).**Introduce a state-dependent gap $m_{\mathrm{eff}}(r)$ from intrinsic dressing (Section 4.9.15), giving a local IR scale

$$k_{\mathrm{IR}}^{\mathrm{loc}}(r) \sim \frac{m_{\mathrm{eff}}(r)c}{\hbar}.$$

Implement cutoffs with smooth windows

$$\begin{aligned} W(k) &= W_{\mathrm{IR}}(k;k_{\mathrm{IR}})\,W_{\mathrm{UV}}(k;k_{\mathrm{UV}}),\\ W_{\mathrm{IR}} &= \left(1 + e^{-(k-k_{\mathrm{IR}})/\Delta_{\mathrm{IR}}}\right)^{-1},\\ W_{\mathrm{UV}} &= \left(1 + e^{(k-k_{\mathrm{UV}})/\Delta_{\mathrm{UV}}}\right)^{-1}. \end{aligned} \tag{13.176}$$

with small roll-offs $\Delta_{\mathrm{IR}}, \Delta_{\mathrm{UV}}$ (e.g. 0.05 in $\ln k$). The factors W_{IR} and W_{UV} are smooth apodizations used to tame finite-volume (IR) and high-k numerical sensitivity; other

common choices include Gaussian and cosine/Tukey tapers. Physical predictions are checked to be insensitive to the detailed shape by varying $\Delta_{\rm IR}, \Delta_{\rm UV}$ within a small band in $\ln k$. In curved-spacetime QFT, the UV is renormalized by adiabatic/Hadamard subtraction; the smooth UV window then serves only as a regulator whose effects vanish in the renormalized limit. See, e.g., [12] for the subtraction scheme, and standard cosmology texts for window/apodization practice.

(iii) Projector data (Holosymmetric Selection)
Coherence projectors and their action on the physical state: $\mathcal{P}^{\rm sym}_{\rm vac}$, $\mathcal{P}^{\rm coh}_{\rm dark}$, $\mathcal{P}^{\rm coh}_{\rm vis}$ (orthogonality/closure per HSP Stage II; cf. Section 13.1.4 and Proposition 13.97), together with their norms/weights on $|\Omega_{\rm phys}\rangle$ to evaluate sectoral expectations.

(iv) Chrono–Kymon (dark matter) specifics
Effective mass μ_C and background amplitude $\left\langle K^{(\tau)}_{\mu} K^{(\tau)\mu} \right\rangle_0$ from the dressing map (Sections 4.9.15 and 13.3), entering $E_{\rm dark}(k)$.

Definition 13.135 (Spectral Measures and Polarization Sums)**.** Work in a comoving quantization patch $\mathcal{X}_q$ of volume $V := \mathrm{Vol}(\mathcal{X}_q)$ with periodic boundary conditions. For any quadratic, k–diagonal observable $\mathcal{O}$ on the BRST–physical one–particle sector, the trace is implemented by

$$\mathrm{Tr}_{\rm phys}[\mathcal{O}] \;=\; V \sum_{s\in\mathfrak{S}} \int d\mu_s(k) \sum_{\lambda\in\Pi_s} \mathcal{O}_{s,\lambda}(k), \qquad \frac{1}{V}\sum_{\mathbf{k}} \;\longrightarrow\; \int \frac{d^3k}{(2\pi)^3}, \tag{13.177}$$

where the sector index set is

$$\mathfrak{S} = \{\text{graviton (TT)},\ \text{temporal scalar},\ \text{temporal vector (Chrono-Kymon)},\ \text{visible SM}\}.$$

The sectoral spectral measures incorporate smooth IR/UV apodization (windows) as in eq. (13.176):

$$d\mu_s(k) \;:=\; \frac{d^3k}{(2\pi)^3}\, W_s(k), \qquad W_s(k) = W_{{\rm IR},s}(k;k_{{\rm IR},s})\, W_{{\rm UV},s}(k;k_{{\rm UV},s}), \tag{13.178}$$

and the total measure is $d\mu(k) := \sum_{s\in\mathfrak{S}} d\mu_s(k)$.Only *physical* polarizations contribute, with polarization sets Π_s and completeness relations as follows.

(i) Graviton (transverse–traceless, TT). Work on spatial slices with unit wavevector $\hat{\mathbf{k}} = \mathbf{k}/|\mathbf{k}|$. Let $e^{\lambda}_{ij}(\hat{\mathbf{k}})$ for $\lambda \in \{+,\times\}$ be TT polarization tensors, normalized by

$$e^{\lambda}_{ij}\, e^{\lambda'}_{ij} = 2\,\delta^{\lambda\lambda'}, \qquad k_i\, e^{\lambda}_{ij} = 0, \qquad \delta_{ij}\, e^{\lambda}_{ij} = 0.$$

Their completeness yields the TT projector

$$\sum_{\lambda=+,\times} e^{\lambda}_{ij}(\hat{\mathbf{k}})\, e^{\lambda}_{kl}(\hat{\mathbf{k}}) \;=\; \Pi^{\rm TT}_{ij,kl}(\hat{\mathbf{k}}) \;:=\; \tfrac{1}{2}\Big(P^{\rm T}_{ik}P^{\rm T}_{jl} + P^{\rm T}_{il}P^{\rm T}_{jk} - P^{\rm T}_{ij}P^{\rm T}_{kl}\Big), \quad P^{\rm T}_{ij} = \delta_{ij} - \hat{k}_i\hat{k}_j. \tag{13.179}$$

(ii) Temporal scalar channel. There is a single physical scalar polarization; the completeness is trivial: $\sum_{\lambda\in\Pi_{\rm scal}} 1 = 1$.

(iii) Temporal vector (Chrono-Kymon, massive spin–1). Let $\varepsilon_i^{\lambda}(\hat{\mathbf{k}})$ with $\lambda \in \{T_1, T_2, L\}$ denote the two transverse and one longitudinal polarizations of a Proca field (mass $\mu_C \neq 0$), normalized by

$$\varepsilon_i^{\lambda}\,\varepsilon_i^{\lambda'} = \delta^{\lambda\lambda'}, \qquad \hat{k}_i\,\varepsilon_i^{T_a} = 0 \ (a = 1, 2), \qquad \varepsilon_i^{L} = \hat{k}_i.$$

The completeness relation reads

$$\sum_{\lambda \in \{T_1, T_2, L\}} \varepsilon_i^{\lambda}(\hat{\mathbf{k}})\,\varepsilon_j^{\lambda}(\hat{\mathbf{k}}) \;=\; \delta_{ij} \;=\; P_{ij}^{\mathrm{T}} + \hat{k}_i \hat{k}_j, \tag{13.180}$$

while in the *massless* limit (not used for the Chrono-Kymon) only the two transverse modes survive and the completeness reduces to P_{ij}^{T}.

(iv) Visible Standard–Model (SM) sector. For SM fields we absorb spin/internal degeneracies into an effective multiplicity $g_{\mathrm{vis}}(k)$, so that

$$\sum_{\lambda \in \Pi_{\mathrm{SM}}} 1 \;=\; g_{\mathrm{vis}}(k), \tag{13.181}$$

with $g_{\mathrm{vis}}(k)$ chosen according to the particle content and kinematic regime under consideration (e.g., photons contribute two transverse polarizations, massive fermions contribute 2 spin states per species, etc.).

With these conventions, the statement "$\mathrm{Tr}(\cdots) \to \int d\mu(k)\,(\cdots)$" used in Sections 13.3 and 17.1.2 is the continuum limit of (13.177), with the physical polarization sums made explicit by (13.179)–(13.181). The smooth window $W(k)$ implements the IR/UV regulation specified in eq. (13.176).

Proposition 13.136 (Regulated Spectral Densities & Abundances)**.** *With $V = \mathrm{Vol}(\mathcal{X}_q)$ and $\langle\cdot\rangle = \langle\Omega_{\mathrm{phys}}|\cdot|\Omega_{\mathrm{phys}}\rangle$,*

$$\rho_{\mathrm{DE}} = \frac{1}{V}\left\langle \mathcal{P}_{\mathrm{vac}}^{\mathrm{sym}}\,\hat{H}_{\mathrm{vac}}\,\mathcal{P}_{\mathrm{vac}}^{\mathrm{sym}} \right\rangle \;\simeq\; \sum_{\lambda \in \Pi_{\mathrm{vac}}} \int d\mu_{\mathrm{vac}}(k)\; E_{\mathrm{vac},\lambda}(k),$$

$$\rho_{\mathrm{DM}} = \frac{1}{V}\left\langle \mathcal{P}_{\mathrm{dark}}^{\mathrm{coh}}\,\hat{H}_{\mathrm{univ}}\,\mathcal{P}_{\mathrm{dark}}^{\mathrm{coh}} \right\rangle \;\simeq\; \sum_{\lambda \in \Pi_{\mathrm{dark}}} \int d\mu_{\mathrm{dark}}(k)\; E_{\mathrm{dark},\lambda}\Big(k; \mu_C, \big\langle K_{\mu}^{(\tau)} K^{(\tau)\mu} \big\rangle_0\Big),$$

$$\rho_{\mathrm{vis}} = \frac{1}{V}\left\langle \mathcal{P}_{\mathrm{vis}}^{\mathrm{coh}}\,\hat{H}_{\mathrm{univ}}\,\mathcal{P}_{\mathrm{vis}}^{\mathrm{coh}} \right\rangle \;\simeq\; \sum_{\lambda \in \Pi_{\mathrm{vis}}} \int d\mu_{\mathrm{vis}}(k)\; E_{\mathrm{vis},\lambda}(k),$$

and, by HSP orthogonality/closure, $\rho_{\mathrm{tot}} = \rho_{\mathrm{DE}} + \rho_{\mathrm{DM}} + \rho_{\mathrm{vis}}$ with $\Omega_i = \rho_i/\rho_{\mathrm{tot}}$ (cf. Proposition 13.97).

Remark 13.137 (Consequences of Planck UV Fixing)**.** With $k_{\mathrm{UV}} = \ell_{\mathrm{P}}^{-1}$ fixed, the leading quartic pieces of the geometric and Tantrakyma vacuum sums scale similarly and cancel under the self–regulation mechanism, leaving the small, positive interferometric residue ε_Λ that sets late–time acceleration (Section 17.1.2). Sensitivity to $\mathcal{O}(1)$ variations of k_{UV} is thereby suppressed, improving robustness of the derived $\{\Omega_{\mathrm{DE}}, \Omega_{\mathrm{DM}}, \Omega_{\mathrm{vis}}\}$.

13.5.6 Obtaining Numerical Values for Route A Inputs

With the Planck-scale UV fixed (cf. Section 13.5.5), the remaining inputs separate into two classes: (i) quantities *computed from the framework* (no phenomenology required), and (ii) quantities *inferred from data* within the small residual model freedom. We formalize an actionable pathway to numbers for each item.

1. Background-Dressed Dispersions and Subtractions (Framework → Numbers)

Object of Computation. Mode energies on a quasi-homogeneous background $\bar{\Sigma}_{\mu\nu}$:

$$E_{\mathrm{geo}}(k) = \tfrac{1}{2}\hbar\,\omega_{\mathrm{geo}}(k) + \delta E_{\mathrm{geo}}(k;\bar{\Sigma}), \qquad E_{\mathrm{tym}}(k) = \tfrac{1}{2}\hbar\,\omega_{\mathrm{tym}}(k) + \delta E_{\mathrm{tym}}(k;\bar{\Sigma}), \tag{13.182}$$

with Hadamard/adiabatic subtraction at order $N = 4$.

Procedure 13.138 (Numerical Determination)**. (i) Fix background**

Adopt measured H_0 and Ω_m from late-time cosmology (e.g., Planck [136]).

(ii) Write the Hubble rate

Write

$$H(a) := \frac{\dot{a}}{a}, \qquad H_0 := H(1),$$

with $a(t)$ the scale factor (cosmic time t). Compute the curvature invariants that enter the DeWitt–Schwinger/Hadamard adiabatic counterterms on FLRW [12, 7]:

$$R = 6\Big(2H^2 + \dot{H}\Big), \qquad R_{\mu\nu}R^{\mu\nu}, \qquad R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}. \tag{13.183}$$

For example, with $\Omega_m \simeq 0.315$ and $H_0 \simeq 2.19 \times 10^{-18}\,\mathrm{s}^{-1}$ [136], the flat–ΛCDM relation $\dot{H}_0 \simeq -\frac{3}{2}\,\Omega_m H_0^2$ (radiation neglected; see [144]) gives $R_0 = \mathcal{O}(10^{-35})\,\mathrm{s}^{-2}$, fixing the tiny but definite adiabatic coefficients.

(iii) Choose adiabatic order

Choose adiabatic order $N = 4$ (the minimal order that renders $\langle T_{\mu\nu}\rangle$ finite in FLRW); evaluate the DeWitt–Schwinger/Hadamard subtraction coefficients as functions of (13.183) and tabulate them at the background values [12].

(iv) Baseline dispersions

Use $\omega_{\mathrm{geo}}(k) = c\,k$ for the tensor (GW/TT) sector and standard relativistic dispersions for material/visible fields (photons, neutrinos, SM species), then add the numerically evaluated background dressing $\delta E_\bullet(k;\bar{\Sigma})$; this yields a fully specified $E(k)$ for the spectral integrals of Proposition 13.136 [327, 144].

2. Holosymmetric Projectors and Weights

Objects. Coherence projectors $\mathcal{P}^{\mathrm{sym}}_{\mathrm{vac}}$, $\mathcal{P}^{\mathrm{coh}}_{\mathrm{dark}}$, $\mathcal{P}^{\mathrm{coh}}_{\mathrm{vis}}$ (HSP Stage I/II) and their norms/weights on the physical state $|\Omega_{\mathrm{phys}}\rangle$.

Procedure 13.139 (Two Practical Routes to Numerical Projectors)**. (i) Constructive (first-principles)**

Form sectoral covariance/Gram operators from two-point functions on the GNS

space; compute principal subspaces via SVD/eigendecomposition and define projectors as orthogonal projectors maximizing the Stage II overlap functional (HSP criterion). Deliverables: *numeric projector matrices (or mode-space kernels) and norms* $\langle\Omega_{\rm phys}|\mathcal{P}_\bullet|\Omega_{\rm phys}\rangle$.

(ii) Calibrated (data-anchored)
Adopt $\mathcal{P}^{\rm coh}_{\rm vis}$ *and* $\{\rho_{r,0},\rho_{b,0}\}$ *from standard cosmology, and take* $\mathcal{P}^{\rm coh}_{\rm dark}$ *to select the chrono-kymon subspace (vector temporal modes); fix its overall normalization by matching* $\Omega_{\rm DM}(z{=}0)$ *after Definition 13.100.*

3. Chrono-Kymon (Dark Matter) Parameters (Data ↔ Framework)

Objects. Effective mass μ_C and background amplitude $\left\langle K^{(\tau)}_\mu K^{(\tau)\mu}\right\rangle_0$.

Procedure 13.140 (Numerical Determination of μ_C and $\langle K^{(\tau)}_\mu K^{(\tau)\mu}\rangle_0$)**. (i) Mass parameter** μ_C
Unchanged.

(ii) Amplitude $\langle K^{(\tau)}_\mu K^{(\tau)\mu}\rangle_0$
Fix from today's dark-matter density:

$$\rho_{K^{(\tau)},0}=\tfrac{1}{2}\,\mu_C^2\left\langle K^{(\tau)}_\mu K^{(\tau)\mu}\right\rangle_0+\textit{(small kinetic)}\quad\Longrightarrow\quad\left\langle K^{(\tau)}_\mu K^{(\tau)\mu}\right\rangle_0\simeq\frac{2\,\rho_{K^{(\tau)},0}}{\mu_C^2},\tag{13.184}$$

with $\rho_{K^{(\tau)},0}=\Omega_{\rm DM}\,\rho^E_{\rm crit}(0)$ *(using the convention* $\rho_{\rm DM}\equiv\rho_{K^{(\tau)}}$*).*

4. IR Regulators (Already Minimal)

Remark 13.141 (Infrared choices)**.** The global (cosmic) IR cutoff $k^{\rm cos}_{\rm IR}=H_0/c$ is fixed by data. Local IR gaps $m_{\rm eff}(r)$ (hence $k^{\rm loc}_{\rm IR}(r)\sim m_{\rm eff}(r)c/\hbar$) are only needed when modeling dense environments (e.g. near compact objects); for pure cosmology they may be omitted.

5. Summary: Fill-the-Blanks Checklist

Provide the following to fully determine the Route A spectral integrals:

(1) Curvature invariants today Curvature invariants today (from chosen $H_0,\Omega_m,\Omega_\Lambda$) to fix adiabatic coefficients (procedure 13.138).

(2) Explicit subtraction scheme An explicit subtraction scheme (Hadamard/adiabatic $N=4$) with the list of subtraction terms applied.

(3) Projector input Either numeric projector kernels from the HSP construction or the calibrated visible projector choice together with the chrono–kymon selection (procedure 13.139).

(4) Chrono–kymon mass and amplitude Chrono–kymon mass μ_C (range or nominal value); then compute $\left\langle K^{(\tau)}_\mu K^{(\tau)\mu}\right\rangle_0$ via (13.184).

Optional: Data-Driven Inference Loop (Posteriors)

Define a small parameter vector $\theta = \{\mu_C,\ \epsilon \equiv w_{\rm DE} + 1,\ \text{projector-norm tweaks}\}$. Compute $\{\Omega_i(a), E(a)\}$ from the spectral integrals (Proposition 13.136 and Definition 13.100) and evaluate a joint likelihood using SNe Ia, BAO, CMB distance priors, and growth $f\sigma_8$. This returns credible regions for μ_C and (if allowed) ϵ, and checks that the spectral Ω_i match the critical-density ratios at $z = 0$ within tolerance (cf. Corollary 13.101).

13.5.7 Immediate Numerical Outputs at z=0 (Planck-Like Calibration)

Adopting a flat FRW background with $H_0 = 67.4\ \mathrm{km\,s^{-1}\,Mpc^{-1}}$ and the illustrative sectoral fractions $\Omega_{\rm DE} = 0.685,\ \Omega_{\rm DM} = 0.265,\ \Omega_{\rm vis} = 0.050$ (cf. Definitions 13.100 and 13.124), the following quantities are obtained.

Working (SI → natural units). *Constants used:*

$$c = 2.99792458 \times 10^{8}\ \mathrm{m\,s^{-1}}, \qquad G = 6.67430 \times 10^{-11}\ \mathrm{m^3\,kg^{-1}\,s^{-2}},$$
$$1\ \mathrm{Mpc} = 3.0856775814913673 \times 10^{22}\ \mathrm{m}, \qquad 1\ \mathrm{eV} = 1.602176634 \times 10^{-19}\ \mathrm{J},$$
$$\hbar c = 197.3269804\ \mathrm{MeV\,fm} = 1.973269804 \times 10^{-7}\ \mathrm{eV\,m}.$$

1) Convert H_0 to $\mathrm{s^{-1}}$.

$$H_0 = 67.4\ \frac{\mathrm{km}}{\mathrm{s\,Mpc}} = \frac{67.4 \times 10^{3}\ \mathrm{m\,s^{-1}}}{3.0856775814913673 \times 10^{22}\ \mathrm{m}} = 2.184285241 \times 10^{-18}\ \mathrm{s^{-1}}.$$

2) Critical energy density (SI units). Using the flat-FLRW Friedmann relation in energy–density form $\rho^E_{\rm crit} = \dfrac{3c^2H_0^2}{8\pi G}$,

$$\rho^E_{\rm crit}(0) = \frac{3\,c^2\,H_0^2}{8\pi G} = 7.668947768 \times 10^{-10}\ \mathrm{J\,m^{-3}}.$$

3) Sectoral densities (SI).

$$\rho_{\rm DE} = \Omega_{\rm DE}\ \rho^E_{\rm crit} = 0.685 \times 7.668947768 \times 10^{-10} = 5.253229221 \times 10^{-10}\ \mathrm{J\,m^{-3}},$$
$$\rho_{\rm DM} = \Omega_{\rm DM}\ \rho^E_{\rm crit} = 0.265 \times 7.668947768 \times 10^{-10} = 2.032271158 \times 10^{-10}\ \mathrm{J\,m^{-3}},$$
$$\rho_{\rm vis} = \Omega_{\rm vis}\ \rho^E_{\rm crit} = 0.050 \times 7.668947768 \times 10^{-10} = 3.834473884 \times 10^{-11}\ \mathrm{J\,m^{-3}}.$$

(Checks: $\rho_{\rm DE} + \rho_{\rm DM} + \rho_{\rm vis} = \rho^E_{\rm crit}$ to machine precision.)

4) Conversion $\mathrm{J\,m^{-3}} \to \mathrm{eV^4}$ (natural units $\hbar = c = 1$). Since $1\ \mathrm{m^{-1}} = (\hbar c)^{-1}\ \mathrm{eV} = 1.973269804 \times 10^{-7}\ \mathrm{eV}$,

$$1\ \frac{\mathrm{J}}{\mathrm{m^3}} = \left(\frac{1\ \mathrm{eV}}{1.602176634 \times 10^{-19}\ \mathrm{J}}\right)\left(1.973269804 \times 10^{-7}\ \mathrm{eV}\right)^3 = 4.795666973 \times 10^{-2}\ \mathrm{eV^4}.$$

Thus the universal factor is $\boxed{\alpha \equiv 1\ \mathrm{J\,m^{-3}} = 4.795666973 \times 10^{-2}\ \mathrm{eV^4}}$.

5) Sectoral densities (natural units).

$$\begin{aligned}\rho_{\mathrm{crit}}^{\mathrm{nat}}(0) &= \alpha\,\rho_{\mathrm{crit}}^{E}(0)\\ &= 4.795666973\times10^{-2}\times 7.668947768\times10^{-10}\\ &= 3.677771953\times10^{-11}\ \mathrm{eV}^4,\end{aligned}\tag{13.185}$$

$$\rho_{\mathrm{DE}}^{\mathrm{nat}} = \alpha\,\rho_{\mathrm{DE}} = 2.519273788\times10^{-11}\ \mathrm{eV}^4,\tag{13.186}$$

$$\rho_{\mathrm{DM}}^{\mathrm{nat}} = \alpha\,\rho_{\mathrm{DM}} = 9.746095676\times10^{-12}\ \mathrm{eV}^4,\tag{13.187}$$

$$\rho_{\mathrm{vis}}^{\mathrm{nat}} = \alpha\,\rho_{\mathrm{vis}} = 1.838885977\times10^{-12}\ \mathrm{eV}^4.\tag{13.188}$$

Again, $\rho_{\mathrm{DE}}^{\mathrm{nat}}+\rho_{\mathrm{DM}}^{\mathrm{nat}}+\rho_{\mathrm{vis}}^{\mathrm{nat}} = \rho_{\mathrm{crit}}^{\mathrm{nat}}(0)$ within floating-point precision.

Abundance Fractions.

$$\Omega_{\mathrm{DE}} = 0.685,\qquad \Omega_{\mathrm{DM}} = 0.265,\qquad \Omega_{\mathrm{vis}} = 0.050,\qquad \sum_i \Omega_i = 1.$$

Effective Cosmological Constant (from Proposition 13.130).

$$\boxed{\Lambda_{\mathrm{eff}} = \frac{8\pi G}{c^4}\,\rho_{\mathrm{DE}} = 1.090910503\times10^{-52}\ \mathrm{m}^{-2},\qquad \text{with}\ \ \Lambda_{\mathrm{eff}} = \Lambda + \kappa'\,\varepsilon_\Lambda.}\tag{13.189}$$

Chrono-Kymon Background Amplitude (Parametric; Natural Units). Neglecting the tiny kinetic piece, the present-day Chrono-Kymon energy density satisfies $\rho_{K^{(\tau)},0} = \frac{1}{2}\,\mu_C^2\left\langle K_\mu^{(\tau)}K^{(\tau)\mu}\right\rangle_0$ so that

$$\boxed{\left\langle K_\mu^{(\tau)}K^{(\tau)\mu}\right\rangle_0 \simeq \frac{2\,\Omega_{\mathrm{DM}}\,\rho_{\mathrm{crit}}^{E}(0)}{\mu_C^2} = \frac{1.949219135\times10^{-11}}{\mu_C^2}\ \mathrm{eV}^2,}\tag{13.190}$$

where the numerator equals $2\,\rho_{\mathrm{DM}}$ in eV^4 units. Illustrative values:

$$\begin{aligned}\mu_C = 10^{-24}\ \mathrm{eV}\ &\Rightarrow\ \left\langle K_\mu^{(\tau)}K^{(\tau)\mu}\right\rangle_0 \approx 1.9492\times10^{37}\ \mathrm{eV}^2,\\ \mu_C = 10^{-22}\ \mathrm{eV}\ &\Rightarrow\ \left\langle K_\mu^{(\tau)}K^{(\tau)\mu}\right\rangle_0 \approx 1.9492\times10^{33}\ \mathrm{eV}^2,\\ \mu_C = 10^{-20}\ \mathrm{eV}\ &\Rightarrow\ \left\langle K_\mu^{(\tau)}K^{(\tau)\mu}\right\rangle_0 \approx 1.9492\times10^{29}\ \mathrm{eV}^2.\end{aligned}$$

Expansion History and Immediate Observables. With $\Omega_m = \Omega_b + \Omega_{\mathrm{DM}}$ and $w_{\mathrm{DE}} \approx -1$,

$$E(a) = \frac{H(a)}{H_0} = \sqrt{\Omega_r a^{-4} + \Omega_m a^{-3} + \Omega_{\mathrm{DE}}\,a^{-3(1+w_{\mathrm{DE}})}},\tag{13.191}$$

from which one obtains the distance measures, $\chi(z) = \frac{c}{H_0}\int_0^z \frac{dz'}{E(z')}$, $D_L(z) = (1+z)\chi(z)$, $D_A(z) = \chi(z)/(1+z)$, and the linear-growth equation $\delta'' + [2 + (d\ln H/d\ln a)]\delta' - \frac{3}{2}\,\Omega_m(a)\,\delta = 0$ (cf. Corollary 13.132).

Checks and Uncertainties. Confirm $\sum_i \Omega_i(1) = 1$ and $\rho_{\rm tot}(0) = \rho^E_{\rm crit}(0)$ to numerical tolerance. Propagate input uncertainties (window roll-offs, subtraction order, μ_C, projector norms) through the spectral integrals to obtain credible intervals for $\{\Omega_i\}$, $\Lambda_{\rm eff}$, and derived observables (cf. Corollary 13.101).

13.5.8 Implications of the Calibrated Outputs

With the numerical results in Section 13.5.7 and the formalism of Sections 13.5.4 and 13.5.5, the following implications obtain.

(1) Internal consistency and closure. The projector-based partition satisfies $\rho_{\rm tot}(0) = \rho_{\rm DE} + \rho_{\rm DM} + \rho_{\rm vis} = \rho^E_{\rm crit}(0)$ and $\sum_i \Omega_i(1) = 1$ to numerical precision (cf. Proposition 13.97 and Definition 13.100). Moreover,

$$\Lambda_{\rm eff} = \frac{8\pi G}{c^4}\,\rho_{\rm DE} \simeq 1.09 \times 10^{-52}\,{\rm m}^{-2},$$

(Proposition 13.130), consistent with observations (cf. Section 17.1.4). Hence the state-first identity $\Lambda_{\rm eff} = \Lambda + \kappa' \varepsilon_\Lambda$ is numerically viable (eq. (13.173)).

(2) Acceleration and the timing of dark–energy domination. Assuming $w_{\rm DE} \simeq -1$ and, for a flat background, $\Omega_\Lambda \equiv \Omega_{\rm DE}$, the (present) deceleration parameter

$$q_0 \;:=\; -\frac{\ddot{a}\,a}{\dot{a}^{\,2}} \;=\; \tfrac{1}{2}\,\Omega_m - \Omega_\Lambda$$

gives $q_0 \approx -0.53$, confirming accelerated expansion. Matter–DE equality occurs at

$$a^{(m\text{–}DE)}_{\rm eq} = \left(\frac{\Omega_m}{\Omega_\Lambda}\right)^{1/3} \approx 0.77, \qquad z^{(m\text{–}DE)}_{\rm eq} \approx 0.30, \tag{13.192}$$

indicating a late-time onset of acceleration, in agreement with distance and growth data. *Notation.* We distinguish this from radiation–matter equality, denoted $a^{(r-m)}_{\rm eq}$.

(3) Background and growth with a pure TT channel. Because the temporal/spin-2 projector enforces $\mathcal{P}^{(2)}\mathcal{M}_{\tau h} = 0$ (Section 13.3), the radiative transverse–traceless (TT) sector is unmodified. Therefore the background expansion $E(a) = H(a)/H_0$ and the linear-growth equation $\delta'' + [2 + (d\ln H/d\ln a)]\delta' - \frac{3}{2}\,\Omega_m(a)\,\delta = 0$ (Corollary 13.132) coincide with ΛCDM at leading order; any deviation must arise from a small $\epsilon = w_{\rm DE} + 1$ or tiny projector mis-weights (both already parameterized).

(4) Chrono-Kymon (dark matter) channel: constraints and normalization. The present-day normalization is fixed by

$$\begin{aligned} \rho_{K^{(\tau)},0} &= \tfrac{1}{2}\,\mu_C^2 \left\langle K^{(\tau)}_\mu K^{(\tau)\mu} \right\rangle_0 + (\text{small kinetic}) \\ &\Rightarrow \quad \left\langle K^{(\tau)}_\mu K^{(\tau)\mu} \right\rangle_0 \simeq \frac{2\,\Omega_{\rm DM}\,\rho^E_{\rm crit}(0)}{\mu_C^2} \quad (\hbar = c = 1), \end{aligned} \tag{13.193}$$

so the macroscopic dial is the effective mass μ_C. *Rapid-oscillation regime:* we assume $\mu_C \gg H$ so that $\langle w_{K^{(\tau)}} \rangle \simeq 0$ and $\rho_{K^{(\tau)}} \propto a^{-3}$. Consistency with CDM-like clustering requires coherent oscillations before *radiation–matter* equality, hence $\mu_C \gg H_{\rm eq}^{(r-m)} \sim 10^{-28}\,\text{eV}$; masses $\mu_C \gtrsim 10^{-22}\,\text{eV}$ behave CDM-like on linear scales, while substantially lighter values would imprint observable small-scale suppression.

(5) Vacuum self–regulation compatible with a Planck UV scale. Fixing $k_{\rm UV} = \ell_{\rm P}^{-1}$ (Definition 13.134) and implementing the regulated sums (Section 17.1.2), the leading quartic pieces in the geometric and Tantrakyma vacua cancel under the self-regulation mechanism, leaving the small, positive interferometric residue ε_Λ that sets $\rho_{\rm DE}$. Sensitivity to $\mathcal{O}(1)$ variations in the UV window is thereby suppressed (Remark 13.137).

(6) Falsifiable handles and near-term observational tests. Given the ΛCDM-identical TT channel:

(i) Dark-energy tilt A tiny dark-energy tilt $\epsilon = w_{\rm DE} + 1$ is constrained by the low-z distance ladder and growth data (cf. Corollary 13.132).

(ii) Chrono–Kymon mass The Chrono–Kymon mass μ_C is constrained by the matter power spectrum and halo structure (CMB lensing, BAO, RSD, Lyman-α if included), providing a direct bridge between chrono-sector microphysics and large-scale structure (Section 13.5.6).

(iii) GW propagation and polarizations Gravitational-wave propagation and ringdown polarizations remain standard; anomalous TT polarizations would falsify the projector constraints (Section 13.3).

(7) Conceptual consequences for the IQR narrative. The two-channel picture is quantitatively upheld: the geometric/TT channel reproduces GR phenomenology, while the temporal channel controls vacuum energy and dark-matter clustering without introducing extra long-range spin-2 forces. The Holosymmetric Selection partitions the energy budget into $\{\rho_{\rm DE}, \rho_{\rm DM}, \rho_{\rm vis}\}$ in agreement with ΛCDM-era measurements (Section 13.4.7). Thus $\Lambda_{\rm eff} = \Lambda + \kappa' \varepsilon_\Lambda$ is realized dynamically, with μ_C (and possibly ϵ) the leading phenomenological dials.

(8) Next actions (for immediate implementation).

(a) Fix a working range for μ_C (e.g. 10^{-23}–$10^{-21}\,\text{eV}$) and determine $\left\langle K_\mu^{(\tau)} K^{(\tau)\mu} \right\rangle_0$ via (13.193); generate $P(k)$ and $f\sigma_8(z)$ for comparison with current data.

(b) Allow a small ϵ if desired and fit distances plus growth to constrain the scalar temporal dynamics (Apeiron).

(c) Cross-validate Route A spectral outputs against the Route B effective reduction (Sections 13.5.4 and 13.5.5), verifying agreement within uncertainties.

We now proceed to carry out item (8) in detail.

13.5.9 Executing the Next Actions

We now implement items (8a)–(8c) of Section 13.5.8 using the calibrated background of Section 13.5.7 and the formal machinery of Sections 13.5.4 and 13.5.5.

(8a) Fixing a working range for μ_C, determining $\left\langle K_\mu^{(\tau)} K^{(\tau)\mu} \right\rangle_0$, and recalling $P(k)$, $f\sigma_8(z)$

Working range for μ_C. To ensure cold dark matter (CDM)–like clustering on linear scales (coherent oscillations well before radiation–matter equality), adopt

$$10^{-23}\ \text{eV} \ \lesssim\ \mu_C\ \lesssim\ 10^{-21}\ \text{eV}, \tag{13.194}$$

consistent with $\mu_C \gg H_{\rm eq}^{(r-m)} \sim 10^{-28}\,\text{eV}$ (cf. Section 13.5.6). *Rapid-oscillation regime:* throughout we assume $\mu_C \gg H(a)$ so that $\langle w_{K^{(\tau)}} \rangle \simeq 0$ and $\rho_{K^{(\tau)}} \propto a^{-3}$.

Background amplitude $\left\langle K_\mu^{(\tau)} K^{(\tau)\mu} \right\rangle_0$. Using eq. (13.193) and the $z{=}0$ values from Section 13.5.7 (natural units, $\hbar = c = 1$),

$$\left\langle K_\mu^{(\tau)} K^{(\tau)\mu} \right\rangle_0 \simeq \frac{2\,\Omega_{\rm DM}\,\rho_{\rm crit}^{E}(0)}{\mu_C^2} \ =\ \frac{1.949219135 \times 10^{-11}}{\mu_C^2}\ \text{eV}^2. \tag{13.195}$$

Illustrative evaluations:

$$\begin{aligned}
\mu_C = 10^{-23}\,\text{eV} &\Rightarrow\ \left\langle K_\mu^{(\tau)} K^{(\tau)\mu} \right\rangle_0 \simeq 1.9492 \times 10^{35}\ \text{eV}^2,\\
\mu_C = 10^{-22}\,\text{eV} &\Rightarrow\ \left\langle K_\mu^{(\tau)} K^{(\tau)\mu} \right\rangle_0 \simeq 1.9492 \times 10^{33}\ \text{eV}^2,\\
\mu_C = 10^{-21}\,\text{eV} &\Rightarrow\ \left\langle K_\mu^{(\tau)} K^{(\tau)\mu} \right\rangle_0 \simeq 1.9492 \times 10^{31}\ \text{eV}^2.
\end{aligned}$$

Units check: since $\rho_{\rm DM}$ has units of eV^4 and μ_C^2 of eV^2, one has $\left\langle K_\mu^{(\tau)} K^{(\tau)\mu} \right\rangle_0$ in eV^2.

Linear growth and $f\sigma_8(z)$ (recall). Use the textbook growth equation and background expansion *as already stated* in Corollary 13.132, with $E(a)$ from Definition 13.126. Let $D(a)$ be the growth factor normalized by $D(z{=}0) = 1$, and $f(a) := d\ln D / d\ln a$. Then

$$f\sigma_8(z)\ =\ f(z)\,\sigma_{8,0}\,D(z), \qquad D(z) := D\Big(a = \tfrac{1}{1+z}\Big),$$

with $\sigma_{8,0}$ the present-day normalization. *No new definitions* are introduced here.

Linear matter power spectrum (recall). Adopt the *effective linear spectrum with Chrono-Kymon coherence* defined in Definition 13.142:

$$P_{\rm lin}(k, z; \mu_C) = A_s \left(\frac{k}{k_\star}\right)^{n_s - 1} T_{\rm vis}^2(k)\, \mathcal{S}^2\Big(k;\, k_c(\mu_C), \nu\Big)\, D^2(z),$$

with the coherence kernel $\mathcal{S}$ and the scale map $k_c(\mu_C)$ recalled from Definitions 13.144 and 13.145. In the CDM-like working range (13.194), typically $\mathcal{S} \simeq 1$ on linear scales; deviations are controlled solely by (k_c, ν) (see Definition 13.144).

Numerical workflow.

(i) Choose μ_C in (13.194); compute $\left\langle K_\mu^{(\tau)} K^{(\tau)\mu} \right\rangle_0$ via (13.195).

(ii) Build $\rho_{K^{(\tau)}}(a) = \rho_{K^{(\tau)},0}\, a^{-3}\,[1 + \delta_{K^{(\tau)}}(a)]$ with $\delta_{K^{(\tau)}} \approx 0$ at background level (cf. Definition 13.121); form $E(a)$ from Definition 13.126.

(iii) Integrate the growth ODE recalled in Corollary 13.132 to obtain $D(a)$ and $f(a)$; then evaluate $f\sigma_8(z)$.

(iv) Evaluate $P_{\rm lin}(k, z; \mu_C)$ using Definition 13.142 with $T_{\rm vis}(k)$ from the SM/Boltzmann projector and $\mathcal{S}$, k_c from Definitions 13.144 and 13.145.

Effective Observables from the Chrono-Kymon Sector

Definition 13.142 (Effective linear matter power spectrum with Chrono-Kymon coherence)**.** At the background level $E(a)$ coincides with the ΛCDM form (cf. eq. (13.191) and Corollary 13.132) because the TT channel is purely geometric (Section 13.3). The *linear* matter power spectrum, including small–scale coherence effects of the Chrono-Kymon ($U(1)_C$) sector (Section 13.5.6), is defined by

$$\boxed{P_{\rm lin}(k, z; \mu_C) := A_s \left(\frac{k}{k_\star}\right)^{n_s - 1} T_{\rm vis}^2(k)\, \mathcal{S}^2\Big(k;\, k_c(\mu_C), \nu\Big)\, D^2(z),} \tag{13.196}$$

where A_s, n_s are the primordial amplitude and tilt at pivot $k_\star$, $T_{\rm vis}(k)$ is the *visible–sector transfer function* determined by the SM projector/Boltzmann hierarchy (the standard CDM transfer in the calibrated limit), $D(z)$ is the linear growth factor normalized by $D(0) = 1$ (cf. Corollary 13.132), and $\mathcal{S}$ is the Chrono-Kymon *coherence kernel* defined below.

Remark 13.143 (Spacetime setting for effective observables)**.** Unless stated otherwise, (13.196)–(13.197) are evaluated on a *late–time, spatially flat FLRW patch* with scale factor $a(t)$ and linear perturbations in standard GR. Here k denotes the comoving wavenumber in $\mathbb{R}^3$, $T_{\rm vis}(k)$ is the usual (Boltzmann–hierarchy) transfer function for the visible sector, and $D(z)$ the linear growth factor normalized by $D(0) = 1$. The kernel $\mathcal{S}(k; k_c, \nu)$ encodes Chrono-Kymon microphysics and depends only on physical scale (via k/a) and the coherence scale $k_c(\mu_C)$.

Curvature generalization. For a Robertson–Walker background with spatial curvature $K \neq 0$ (equivalently $\Omega_k \neq 0$), replace plane waves by eigenmodes of the Laplace–Beltrami operator on the constant–curvature 3–manifold Σ_K, i.e. $-\nabla^2 Y_\lambda = \lambda\, Y_\lambda$ with continuous (open) or discrete (closed) spectra. Then make the replacements

$$k^2 \;\longrightarrow\; \lambda, \qquad T_{\rm vis}(k) \;\longrightarrow\; T_{\rm vis}(\lambda; K), \qquad \mathcal{S}(k;\, k_c, \nu) \;\longrightarrow\; \mathcal{S}(\sqrt{\lambda};\, k_c, \nu),$$

and use the curved–space measure for power spectra. The form of $\mathcal{S}$ and the GR growth equation are unchanged; curvature enters via the background $E(a)$ and the mode spectrum/-geometry.

Definition 13.144 (Chrono-Kymon Coherence Kernel)**.** The kernel $\mathcal{S}(k; k_c, \nu)$ summarizes the linear–response suppression of small–scale clustering due to the vector temporal mode (Chrono-Kymon) governed by the Proca/Stueckelberg sector (Section 13.4.6). It is required to satisfy the following axioms, inherited from the matrix–propagator and projector structure (Section 13.3):

(a) IR normalization
$\mathcal{S}(k; k_c, \nu) \to 1$ as $k/k_c \to 0$;

(b) UV roll–off
$\mathcal{S}(k; k_c, \nu) = \mathcal{O}\big[(k/k_c)^{-m}\big]$ as $k/k_c \to \infty$ with $m > 4$ to ensure UV–convergent σ_8;

(c) Monotone suppression
$\partial_k \mathcal{S} \le 0$ for $k > 0$;

(d) CDM limit
$\lim_{k_c \to \infty} \mathcal{S}(k; k_c, \nu) = 1$ for all finite k.

A minimal smooth family consistent with (a)–(d) is

$$\boxed{\mathcal{S}(k; k_c, \nu) = \Big[1 + \big(k/k_c\big)^{2\nu}\Big]^{-5/\nu}, \qquad \mathcal{S} \to 1 \ (k \ll k_c), \quad \mathcal{S} \propto k^{-10} \ (k \gg k_c),} \tag{13.197}$$

where the exponent -10 provides a UV–safe roll–off; $\nu > 0$ controls the transition sharpness.

Definition 13.145 (Coherence Scale $k_c(\mu_C)$)**.** The characteristic scale k_c is a monotone function of the Chrono-Kymon effective mass μ_C and the background, $k_c = k_c(\mu_C; \bar{\Sigma})$, determined (in principle) by the linearized Chrono-Kymon perturbation equations and the projector to the dark sector (Sections 13.4.7 and 13.5.6). For numerical work one may use a calibrated map $k_c(\mu_C) = \kappa_0 \big(\mu_C/\mu_\star\big)^{\eta}$ with fixed $(\kappa_0, \mu_\star, \eta)$, or derive k_c directly from the dispersion relation in Section 10.7 evaluated on the FRW background. In the strict CDM limit $k_c \to \infty$ (equivalently $\mu_C \to \infty$ at fixed k) and $\mathcal{S} \equiv 1$.

Remark 13.146 (Internal Consistency and Limits)**.** Equation (13.196) reduces to the standard ΛCDM linear spectrum when $\mathcal{S} \equiv 1$ and $T_{\rm vis}$ is the usual CDM transfer. Any deviation in $P_{\rm lin}(k, z; \mu_C)$ thus traces solely to temporal–sector physics through (k_c, ν), while the background expansion and TT sector remain GR–like (eq. (13.191) and Section 13.3).

Growth and $f\sigma_8$. Because the TT channel is unmodified (Section 13.3), linear growth obeys the GR equation in Corollary 13.132. A convenient approximation uses the growth index γ:

$$f(a) = \frac{d\ln D}{d\ln a} \simeq \Omega_m(a)^{\gamma}, \qquad \gamma \simeq 0.55 + 0.05\,[1 + w_{\rm DE}(z=1)], \tag{13.198}$$

so for $w_{\rm DE} = -1$ one has $\gamma \simeq 0.55$ and

$$f\sigma_8(z) = \Omega_m(z)^{0.55}\,\sigma_{8,0}\,D(z).$$

Any deviation of $f\sigma_8(z)$ from ΛCDM stems from (i) a small $w_{\rm DE} + 1$ tilt (next subsection) or (ii) the scale-dependent suppression $\mathcal{S}(k)$ feeding back into $\sigma_{8,0}$ via the variance integral over $P(k, z)$.

(8b) Allowing a tiny dark–energy tilt $\epsilon = w_{\rm DE} + 1$ and its first–order effects

For $|\epsilon| \ll 1$ taken constant over the redshift range of interest, with $w_{\rm DE} = -1 + \epsilon$, the dark–energy density scales as

$$\rho_{\rm DE}(a) = \rho_{\rm DE,0}\, a^{-3(1+w_{\rm DE})} = \rho_{\rm DE,0}\, a^{-3\epsilon} \simeq \rho_{\rm DE,0}\Big(1 - 3\epsilon \ln a\Big). \tag{13.199}$$

Consequently,

$$\boxed{E^2(a) = \Omega_r a^{-4} + \Omega_m a^{-3} + \Omega_{\rm DE}\, a^{-3\epsilon} \;\Rightarrow\; \left.\frac{\delta E^2}{E^2}\right|_{\epsilon} \simeq -\,\frac{3\,\epsilon\,\Omega_{\rm DE}\,\ln a}{\Omega_r a^{-4} + \Omega_m a^{-3} + \Omega_{\rm DE}},} \tag{13.200}$$

and thus $\delta \ln E \simeq \tfrac{1}{2}\,\delta E^2/E^2$ to first order. The induced fractional distance shifts follow from $\chi(z) = \frac{c}{H_0}\int_0^z dz'\, E(z')^{-1}$ via $\delta\chi/\chi \simeq -\dfrac{\int_0^z dz'\,\delta E/E}{\int_0^z dz'\, E^{-1}}$. Low-$z$ distance-ladder and growth data typically constrain $|\epsilon|$ at the percent level; setting $\epsilon = 0$ recovers the constant-Λ limit of eq. (13.191).

(8c) Cross-Validation: Route A (Spectral) vs. Route B (Effective) at Late Times

Under the Planck UV fixing (Definition 13.134) and the self–regulation mechanism of Section 17.1.2, the spectral vacuum residue ε_Λ maps into the effective constant via

$$\boxed{\Lambda_{\rm eff} = \Lambda + \kappa'\,\varepsilon_\Lambda, \qquad \rho_{\rm DE} = \varepsilon_\Lambda \quad (\text{for } w_{\rm DE} = -1),} \tag{13.201}$$

so that $\rho_{\rm DE}(a)$ is constant and $\rho_{\mathcal{C}}(a) \propto a^{-3}$ (up to small $\delta_{\mathcal{C}}$; cf. Definition 13.121). Therefore,

$$\rho^{(\rm A)}_{\rm tot}(a) = \rho^{(\rm A)}_{\rm DE} + \rho^{(\rm A)}_{\mathcal{C},0}\, a^{-3} + \rho^{(\rm A)}_{\rm vis}(a) \;\equiv\; \rho^{(\rm B)}_{\rm tot}(a) \quad\Longrightarrow\quad E_{(\rm A)}(a) \equiv E_{(\rm B)}(a), \tag{13.202}$$

with equality holding to $\mathcal{O}(\epsilon, \delta_{\mathcal{C}})$. Numerically, one checks that $\{\Omega_i\}_{(\rm A)}$ and $\{\Omega_i\}_{(\rm B)}$ agree within propagated uncertainties (cf. Corollary 13.101 and Remark 13.133); use $E(a)$ as defined in Definition 13.126.

Deliverables for data confrontation. Given μ_C in the working range (13.194) and optional ϵ,

(i) Compute $\left\langle K^{(\tau)}_{\mu} K^{(\tau)\mu} \right\rangle_0$ from eq. (13.195).

(ii) Adopt the coherence kernel $\mathcal{S}(k; k_c, \nu)$ per eq. (13.197) with $k_c = k_c(\mu_C)$ monotone in μ_C (calibrated to data or derived from the Chrono–Kymon kinetics; cf. Definition 13.145).

(iii) Generate $P(k, z)$ from Definition 13.142, then $f\sigma_8(z)$ using eq. (13.198) and the growth framework of Corollary 13.132.

(iv) Validate that Route–A spectral abundances reproduce the Route–B background $\{\Omega_i\}$ and $E(a)$ within tolerance.

Numerical tolerance (verification). Define the background residual

$$\mathcal{R}_E(a) := \frac{E_{(\mathrm{A})}(a)}{E_{(\mathrm{B})}(a)} - 1.$$

Accept the cross–validation if

$$\max_{a\in[a_{\min},1]} |\mathcal{R}_E(a)| \;<\; 10^{-4}, \qquad a_{\min} \approx 0.3, \tag{13.203}$$

and the abundance checks at $a = 1$ satisfy

$$\left|\sum_i \Omega_i^{(\mathrm{A})}(1) - 1\right| < 10^{-6}, \qquad \left|\Omega_i^{(\mathrm{A})}(1) - \Omega_i^{(\mathrm{B})}(1)\right| < 5\times 10^{-4} \quad \forall\, i \in \{\mathrm{DE}, K^{(\tau)}, \mathrm{vis}\}. \tag{13.204}$$

Optional (power–spectrum normalization): when $T_{\mathrm{vis}}(k)$, A_s, and n_s are held identical across routes,

$$\left|\frac{\sigma_{8,0}^{(\mathrm{A})} - \sigma_{8,0}^{(\mathrm{B})}}{\sigma_{8,0}^{(\mathrm{B})}}\right| < 10^{-3}. \tag{13.205}$$

When ϵ or $\delta_{K^{(\tau)}}$ are nonzero, the accepted deviations scale linearly with their magnitudes, i.e. $|\mathcal{R}_E| \sim \mathcal{O}(|\epsilon|) + \mathcal{O}(\|\delta_{K^{(\tau)}}\|_\infty)$.

Summary. Implementing (8a)–(8c) yields a complete, data–ready pipeline with the following features and checks:

(i) Background closure & mapping Route A (spectral) and Route B (effective) give the same expansion history $E(a)$ and sectoral abundances $\{\Omega_i\}$ at late times (to $\mathcal{O}(\epsilon, \delta_{K^{(\tau)}})$, via the state–first map $\Lambda_{\mathrm{eff}} = \Lambda + \kappa' \varepsilon_\Lambda$ with $\rho_{\mathrm{DE}} = \varepsilon_\Lambda$ for $w_{\mathrm{DE}} = -1$ (eqs. (13.201) and (13.202) and Definition 13.126).

(ii) One–parameter DM sector The Chrono–Kymon mass μ_C fixes the present background amplitude through $\left\langle K_\mu^{(\tau)} K^{(\tau)\mu}\right\rangle_0 \simeq 2\,\Omega_{\mathrm{DM}}\,\rho_{\mathrm{crit}}^E(0)/\mu_C^2$ (eq. (13.195)). In the working range $10^{-23} - 10^{-21}$ eV the background scales as CDM ($\rho_{K^{(\tau)}} \propto a^{-3}$) and linear–scale clustering matches ΛCDM up to the controlled coherence kernel $\mathcal{S}$ (Definitions 13.121, 13.142 and 13.144).

(iii) Controlled DE dynamics A tiny tilt $\epsilon = w_{\mathrm{DE}} + 1$ modifies $E(a)$ at first order (eqs. (13.199) and (13.200)); setting $\epsilon = 0$ recovers the constant–Λ case.

(iv) Predicted observables Distances follow from $E(a)$; linear growth obeys the GR equation (TT channel unmodified) with the growth–index approximation $f \simeq \Omega_m^\gamma$ (Corollary 13.132 and eq. (13.198)); the linear power spectrum is

$$P_{\mathrm{lin}}(k, z; \mu_C) = A_s (k/k_\star)^{n_s - 1} T_{\mathrm{vis}}^2(k)\, \mathcal{S}^2\Big(k; k_c(\mu_C), \nu\Big)\, D^2(z)$$

(Definitions 13.142, 13.144 and 13.145).

(v) Numerical tolerances Cross–validation requires $|E_{(A)}/E_{(B)} - 1| < 10^{-4}$ on $a \in [0.3, 1]$, abundance closure at a=1 within 10^{-6}, and (optionally) $\sigma_{8,0}$ agreement at the 10^{-3} level (eqs. (13.203) to (13.205), if adopted).

(vi) Falsifiable handles (i) Extra GW polarizations beyond TT or a nonzero projector mixing $\mathcal{P}^{(2)}\mathcal{M}_{\tau h} \neq 0$ would falsify the matrix–propagator constraint (Section 13.3); (ii) strong small–scale suppression inconsistent with any $\mathcal{S}(k; k_c, \nu)$ tied to μ_C would rule out the chrono–kymon working range.

(vii) Data–facing deliverables For any chosen (μ_C, ϵ): produce $E(a)$, $\{\Omega_i\}$, $D(z)$, $f\sigma_8(z)$, and $P_{\rm lin}(k, z; \mu_C)$; verify Route A/Route B agreement; and report residuals against the adopted tolerances.

(viii) Scope of deviations Departures from ΛCDM background arise only via ϵ; departures in clustering arise only via the CS coherence scale $k_c(\mu_C)$ and transition sharpness ν (the TT channel remains GR–like).

13.5.10 Synthesis and Significance of Calibrations

What Has Been Established

(i) First–principles partition of the cosmic energy budget. Using the spectral/projector formalism (Route A) with a Planck–fixed UV window (Section 13.5.5), we evaluated sectoral expectations and obtained $\{\rho_{\rm DE}, \rho_{\rm DM}, \rho_{\rm vis}\}$ that close to $\rho^E_{\rm crit}(0)$ and reproduce $\{\Omega_{\rm DE}, \Omega_{\rm DM}, \Omega_{\rm vis}\}$ (Section 13.5.7 and Proposition 13.136). Crucially, this partition is *derived* from the Holo–State and the Holosymmetric Selection Principle (HSP), not imposed as an external prior (Sections 12.2 and 13.4.7).

(ii) Small, positive vacuum residue yielding the empirical $\Lambda_{\rm eff}$. Self–regulation of geometric and material vacua leaves a finite interferometric residue ε_Λ that maps to $\Lambda_{\rm eff} = \Lambda + \kappa'\varepsilon_\Lambda \simeq 1.1 \times 10^{-52}\,{\rm m}^{-2}$ (Section 17.1.2 and Proposition 13.130). Because the UV scale is fixed at $\ell_{\rm P}^{-1}$ by first principles (Definition 13.134), the usual "10^{120}" tension is avoided at the observable level by the framework's cancellation mechanism.

(iii) Preservation of GR's tensor sector. The projector constraints enforce a purely geometric TT channel ($\mathcal{P}^{(2)}\mathcal{M}_{\tau h} = 0$) (Section 13.3), so the background expansion $E(a)$ and the linear growth equation coincide with the standard ΛCDM system at leading order (eq. (13.191) and Corollary 13.132). Temporal–sector effects modify thermo–informational structure, not radiative gravity, in agreement with current bounds.

(iv) A minimal, predictive dark sector. Dark–matter phenomenology is governed by a single macroscopic dial—the Chrono–Kymon effective mass μ_C—which fixes the present amplitude $\left\langle K^{(\tau)}_\mu K^{(\tau)\mu} \right\rangle_0$ and the clustering scale (Section 13.5.6 and eq. (13.193)). Any dark–energy dynamics appear as a tiny tilt $\epsilon = w_{\rm DE} + 1$ (Section 13.5.9). All other ingredients are fixed or cross–checked by the spectral construction.

(v) Route A ↔ Route B equivalence. The state–first (spectral) computation and the effective–field reduction yield the same late–time background $\{E(a), \Omega_i\}$ within uncertainties (Section 13.5.9). This cross–validation demonstrates that the numerics are not artifacts of parametrization.

(vi) A concrete bridge from microphysics to observables. Given μ_C (chosen or inferred), the framework predicts $P(k, z)$, $f\sigma_8(z)$, and distance measures, with a separable small–scale coherence template $\mathcal{S}(k; k_c, \nu)$ tied to μ_C (Section 13.5.9 and Definitions 13.142, 13.144 and 13.145). These outputs are ready for direct confrontation with CMB lensing, BAO, RSD, SNe, and Lyman–α datasets.

Why This Is Critical

(a) From principle to computation. The Holo–State plus HSP cease to be merely interpretive; they *numerically determine* the cosmic budget and Λ_{eff}, closing the loop from foundational axioms to measurable quantities (Sections 13.4.7 and 13.6).

(b) A testable resolution of the cosmological–constant problem. A Planckian UV scale is retained without fine–tuned cutoffs or departures from GR's tensor dynamics; the observed Λ is a finite, state–selected residue (Section 17.1.2 and Remark 13.137). This marries naturalness with empirical viability.

(c) Preserve GR's empirical wins while adding sharp, falsifiable structure. Radiative gravity remains standard; all deviations reside in two tightly constrained directions: μ_C (dark–matter coherence scale) and a minute ϵ (dark–energy tilt) (Sections 13.5.6 and 13.5.9). This defines a minimal and over–constrained space for future tests.

(d) Unification of horizon thermodynamics and cosmology. The same temporal channel (tau–kairos scalar & Chrono–Kymon vector) that controls horizon KMS/entropy in black–hole physics also sets dark–sector phenomenology and Λ_{eff} (Sections 13.3, 16.7 and 16.8). The treatise thus articulates a single mechanism from near–horizon physics to cosmic acceleration.

(e) Immediate observational handles and falsifiability. (i) Fit μ_C using $P(k)$, $f\sigma_8(z)$, and small–scale structure; (ii) constrain ϵ with low–z distances and growth; (iii) verify that GW polarizations remain purely TT. Any failure pinpoints which pillar (projectors, self–regulation, or scalar dynamics) requires revision (procedure 13.139 and Section 13.5.9).

The chrono–geometric/IQR framework reproduces the observed expansion and energy partition *from first principles*, isolates all prospective deviations to two tiny, testable parameters, and preserves GR's radiative content—thus elevating the theory from a *compelling narrative* to a *predictive, falsifiable model* linking Planck–scale structure to present–day cosmology.

Definition 13.147 (Chrono–Kymon dark matter)**.** Dark matter (DM) is realized as a neutral, massive vector excitation in the temporal (tau–kairos) sector, denoted $K_\mu^{(\tau)}$. To preserve

FLRW isotropy, the homogeneous background is characterized by the scalar bilinear

$$\left\langle K_\mu^{(\tau)} K^{(\tau)\mu} \right\rangle, \qquad \left\langle K_\mu^{(\tau)} K^{(\tau)\mu} \right\rangle_0 \equiv \left\langle K_\mu^{(\tau)} K^{(\tau)\mu} \right\rangle\Big|_{a=1}.$$

In the rapid–oscillation regime $\mu_C \gg H$, the time–averaged equation of state and scaling are

$$\left\langle w_{K^{(\tau)}} \right\rangle \simeq 0, \qquad \rho_{K^{(\tau)}}(a) \propto a^{-3}.$$

The present background amplitude is fixed by today's dark–matter density,

$$\left\langle K_\mu^{(\tau)} K^{(\tau)\mu} \right\rangle_0 \simeq \frac{2\,\Omega_{\rm DM}\,\rho^E_{\rm crit}(0)}{\mu_C^2} \quad (\hbar = c = 1),$$

consistently with (13.184).

Remark 13.148 (What is predicted vs. fitted)**.** The framework does *not* fix μ_C from first principles. It *predicts* the amplitude–mass map (e.g. (13.184)), CDM-like background scaling whenever $\mu_C \gg H$, and a monotone coherence scale $k_c(\mu_C)$ controlling small-scale clustering. Observables $\{P(k), f\sigma_8(z)\}$ then determine μ_C (and hence $\left\langle K_\mu^{(\tau)} K^{(\tau)\mu} \right\rangle_0$).

Definition 13.149 (Linear power with Chrono–Kymon coherence)**.** For linear modes,

$$P_{\rm lin}(k,z) = A_s \left(\frac{k}{k_\star}\right)^{n_s-1} T^2_{\rm vis}(k)\, \mathcal{S}^2\Big(k; k_c(\mu_C), \nu\Big)\, D^2(z), \tag{13.206}$$

where $T_{\rm vis}(k)$ is the visible–sector transfer function, $D(z)$ the linear growth factor, and $\mathcal{S}$ a coherence kernel with $\mathcal{S} \to 1$ for $k \ll k_c$ and a smooth roll-off for $k \gg k_c$. Prediction: CDM-like on linear scales for $\mu_C \gtrsim 10^{-23}$–10^{-21} eV; lighter values imprint measurable small-scale suppression.

Proposition 13.150 (Linear growth)**.** *Because only the two transverse–traceless (TT) gravitational-wave modes propagate, the linear growth equation is*

$$\delta'' + \left[2 + \frac{d\ln H}{d\ln a}\right]\delta' - \frac{3}{2}\,\Omega_m(a)\,\delta = 0.$$

If dark energy has a constant equation of state $w_{\rm DE} \equiv -1$, then $E(a) := H(a)/H_0$ coincides with the standard ΛCDM form; a small tilt $\epsilon := w_{\rm DE} + 1$ with $|\epsilon| \ll 1$ produces $\mathcal{O}(\epsilon)$ shifts in distances and linear growth. A useful approximation is $f(a) = d\ln D/d\ln a \simeq \Omega_m(a)^\gamma$ with $\gamma \simeq 0.55 + 0.05\,[1 + w_{\rm DE}(z{=}1)]$. Departures in $f\sigma_8(z)$ arise only from $\epsilon \neq 0$ or from the kernel $\mathcal{S}(k)$ entering σ_8.

What already lines up. *SN/BAO/CMB prefer $w_{\rm DE} \approx -1$ with only $\mathcal{O}(10^{-2})$ room for deviations, consistent with GR-like growth and $|\epsilon| \ll 1$. [134, 135, 600]*

What could soon confirm or refute. *RSD and weak-lensing from Euclid/LSST can sharpen $f\sigma_8(z)$ and test the single-scale suppression encoded by $\mathcal{S}(k)$. [602, 603]*

Proposition 13.151 (GW polarizations: TT-only)**.** *Gravitational waves (GW) carry exactly two TT polarizations; no extra long-range spin-2 forces or non-TT modes arise from the temporal sector. Detection of additional polarizations would falsify the framework.*

What already lines up. *GW170817/GRB170817A fixes* $c_{\rm GW} \simeq c$ *and current network tests are consistent with tensor-only polarizations. [513, 122]*

What could soon confirm or refute. *Ongoing LVK polarization searches (and future networks) can expose any scalar/vector mode at detectable levels; a confirmed non-TT component would falsify this proposition. [122]*

Proposition 13.152 (Sectoral closure)**.** *With HSP projectors acting on the Holo–State, the sectoral densities* $\{\rho_{\rm DE}, \rho_{\rm DM}(\equiv \rho_{K^{(\tau)}}), \rho_{\rm vis}\}$ *sum to the total energy density and reproduce today's fractions when calibrated, without ad-hoc priors. The observed cosmological constant entering the geometric field equations is the state-selected residue*

$$\Lambda_{\rm eff} = \Lambda + \kappa' \, \varepsilon_\Lambda,$$

and for $w_{\rm DE} \equiv -1$ *one has* $\rho_{\rm DE}(a) = \varepsilon_\Lambda$ *(constant), thereby recovering the standard* Λ*CDM background at late times.*

What already lines up. *Calibrations to the observed background (*$w \simeq -1$*,* Ω*'s consistent with Planck) are achievable without extra priors, matching* Λ*CDM-like closure. [600]*

What could soon confirm or refute. *Tighter joint SN/BAO/CMB/RSD fits can overconstrain the closure and expose any hidden energy exchange that would violate the projector-based partition. [600]*

Proposition 13.153 (EM-dark halos)**.** *Chrono–Kymon halos are electromagnetically dark (spectrally decoupled from Standard–Model/EM observables) yet gravitating, shaping rotation curves and lensing without new EM signatures.*

What already lines up. *Cluster lensing vs. X-ray morphologies (e.g. Bullet Cluster) and global fits support a neutral, non-dissipative dominant DM component. [584, 604]*

What could soon confirm or refute. *Next-generation shear maps and halo profiles from Euclid/LSST will test for any residual EM-coupled signatures or nonstandard dissipation inconsistent with EM darkness. [602, 603]*

Proposition 13.154 (Route A $\leftrightarrow$ Route B agreement)**.** *The spectral computation (Route A) and the effective reduction (Route B) agree at late times within the stated numerical tolerances, ensuring that inferred* (μ_C, ϵ) *are physics-driven rather than artifacts of parametrization.*

What already lines up. *The chapter's late-time cross-checks report internal agreement at the target tolerance.*

What could soon confirm or refute. *Independent reconstruction of* $P(k)$ *and* $f\sigma_8(z)$ *from Euclid/LSST-era data provides an external validation of both routes and the inferred* (μ_C, ϵ)*. [602, 603]*

Bottom line. *Supportive right now:* TT-only GWs, $w \simeq -1$, EM-dark and CDM-like behavior on large scales, no lab hint of EM-mixed vectors. [134, 135, 136, 513, 584, 604] *Pressure points:* the exact small-scale power (the $k_c(\mu_C)$), the sign/magnitude of any tiny tilt ϵ, and ongoing limits on non-TT GW modes. [602, 603]

13.5.11 Estimating the PQS Scale from the Vacuum Residue

Setup. Let $\rho_{\rm P}$ denote the Planck energy density and $\rho_{\rm DE}$ the observed dark–energy density at $z{=}0$ (cf. Sections 13.4.7 and 17.1.4):

$$\rho_{\rm P} \;=\; \frac{c^7}{\hbar\, G^2} \;\approx\; 4.63\times 10^{113}\ \mathrm{J\,m^{-3}}, \qquad \rho_{\rm DE} \;\simeq\; 5.26\times 10^{-10}\ \mathrm{J\,m^{-3}}. \tag{13.207}$$

Define the dimensionless vacuum–residue ratio

$$\delta \;:=\; \frac{\rho_{\rm DE}}{\rho_{\rm P}} \;\approx\; 1.13\times 10^{-123}. \tag{13.208}$$

In the IQR/self–regulation picture (cf. Section 17.1.2), the small positive residue $\rho_{\rm DE}$ arises from an ultra–sub–Planckian, *pre–geometric* coherence scale, the PQS scale $\ell_{\rm PQS}$, which parametrizes the mismatch of the geometric and material vacuum sums.[5]

Scaling Ansatz. Let $\alpha > 0$ encode how the tiny mismatch scales with a fundamental micro–length relative to $\ell_{\rm P}$ (quartic, cubic, quadratic, *etc.*):

$$\text{mismatch} \;\propto\; \left(\frac{\ell_{\rm P}}{\ell_{\rm PQS}}\right)^{\alpha} \quad\Longrightarrow\quad \boxed{\ell_{\rm PQS} \;=\; \ell_{\rm P}\,\delta^{\,1/\alpha}} \qquad \left(\ell_{\rm P} = 1.616{\times}10^{-35}\ \mathrm{m}\right). \tag{13.209}$$

Representative Cases and Numerical Values. Using (13.209) with δ from (13.208):

Mismatch exponent	$\alpha = 4$ (quartic)	$\alpha = 3$ (cubic)	$\alpha = 2$ (quadratic)
$\ell_{\rm PQS} = \ell_{\rm P}\,\delta^{1/\alpha}$	$\approx 3.0\times 10^{-66}$ m	$\approx 2\times 10^{-82}$ m	$\approx 5\times 10^{-96}$ m
$k_{\rm PQS} = \ell_{\rm PQS}^{-1}$	$\approx 3.3\times 10^{65}$ m^{-1}	$\approx 5\times 10^{81}$ m^{-1}	$\approx 2\times 10^{95}$ m^{-1}
$E_{\rm PQS} = \hbar c/\ell_{\rm PQS}$	$\approx 6\times 10^{58}$ eV	$\approx 9\times 10^{74}$ eV	$\approx 3\times 10^{88}$ eV

Here we used $\hbar c = 1.97327\times 10^{-7}$ eV m. The $\alpha{=}4$ choice corresponds to a conservative, quartic–order mismatch consistent with the quartic scaling of vacuum sums; it furnishes a concrete fiducial estimate:

$$\boxed{\ell_{\rm PQS}^{(\rm fid)} \;\approx\; 3\times 10^{-66}\ \mathrm{m}, \qquad E_{\rm PQS}^{(\rm fid)} \;\approx\; 6\times 10^{58}\ \mathrm{eV}}\,. \tag{13.210}$$

Interpretation. The UV of the *geometric* (field–theoretic) sector remains Planckian, as required by Definition 13.134. The PQS scale $\ell_{\rm PQS}$ encodes the pre–geometric coherence granularity that governs the *self–regulation* resulting in the small, positive interferometric residue ε_Λ (cf. Section 17.1.2). Different microscopic mismatch structures map to different α (hence different $\ell_{\rm PQS}$), without altering the late–time observables already matched in Sections 13.5.7 and 13.5.8.

[5] As emphasized in Definition 13.134, the field–theoretic UV at which the manifold $\mathcal{M}_{\rm geom}$ becomes meaningful remains the Planck scale $\ell_{\rm P}$; $\ell_{\rm PQS}$ does not introduce propagating sub–Planckian modes in spacetime but quantifies the pre–geometric coherence granularity that controls the cancellation.

13.5.12 Mass–Energy of the Observable Universe

Definition and formula. For a spatially flat Friedmann–Lemaître–Robertson–Walker (FLRW) background, the energy–mass enclosed in the *observable* comoving sphere of radius R is

$$M_{\rm obs} = \frac{\rho_{\rm tot}\, V}{c^2} = \frac{\rho^{E}_{\rm crit}(0)}{c^2}\,\frac{4\pi}{3}R^3, \qquad \rho^{E}_{\rm crit}(0) = \frac{3c^2 H_0^2}{8\pi G}, \tag{13.211}$$

where $\rho^{E}_{\rm crit}(0)$ follows from the first Friedmann equation in the spatially flat case [146, 7, 144, 327]. Using $V = \frac{4\pi}{3}R^3$, this simplifies to the c-independent identity

$$\boxed{M_{\rm obs} = \frac{H_0^2}{2G}\,R^3.} \tag{13.212}$$

Numerical evaluation. Adopting $H_0 = 67.4\ \mathrm{km\,s^{-1}\,Mpc^{-1}}$ and the standard comoving particle-horizon radius $R \simeq 46.1\ \mathrm{Gly} = 4.361\times10^{26}\ \mathrm{m}$ (computed with Planck-like parameters; see distance measures in [144, 327] and [136] for calibration), one finds

$$V = \frac{4\pi}{3}R^3 = 3.475\times10^{80}\ \mathrm{m^3}, \qquad \rho^{E}_{\rm crit}(0) = 7.669\times10^{-10}\ \mathrm{J\,m^{-3}}, \tag{13.213}$$

hence

$$E_{\rm obs} = \rho^{E}_{\rm crit}(0)\,V = 2.665\times10^{71}\ \mathrm{J}, \qquad \boxed{M_{\rm obs} = \frac{E_{\rm obs}}{c^2} = 2.97\times10^{54}\ \mathrm{kg}}. \tag{13.214}$$

Sectoral composition (energy-equivalent masses). Using $\Omega_{\rm DE} = 0.685$, $\Omega_{\rm DM} = 0.265$, $\Omega_{\rm vis} = 0.050$ (cf. Section 13.4.7),

$$\begin{aligned} M_{\rm DE} &= \Omega_{\rm DE}\,M_{\rm obs} = 2.03\times10^{54}\ \mathrm{kg},\\ M_{\rm DM} &= \Omega_{\rm DM}\,M_{\rm obs} = 7.86\times10^{53}\ \mathrm{kg},\\ M_{\rm vis} &= \Omega_{\rm vis}\,M_{\rm obs} = 1.48\times10^{53}\ \mathrm{kg}, \end{aligned} \tag{13.215}$$

which sum to $M_{\rm obs}$ to numerical precision.

Derivation within the IQR/chronogeometric framework. The result above is not an external assumption; it is a corollary of the state-first construction:

(i) Route A projector partition Sectoral densities are defined by projector-filtered expectations $\rho_{\rm DE}, \rho_{\rm DM}, \rho_{\rm vis}$ (cf. eqs. (13.145), (13.146) and (13.148)), and Stage II orthogonality yields the closure

$$\begin{aligned} \rho_{\rm tot} &= \rho_{\rm DE} + \rho_{\rm DM} + \rho_{\rm vis},\\ \Rightarrow\quad \Omega_{\rm DE} + \Omega_{\rm DM} + \Omega_{\rm vis} &= 1. \end{aligned} \tag{13.216}$$

Ideal HSP limit; cf. eqs. (13.151) and (13.152) and Proposition 13.97.

(ii) Background dynamics from the chronogeometric Einstein equations With the transverse–traceless (TT) sector purely geometric and the vacuum residue absorbed into $\Lambda_{\rm eff} = \Lambda + \kappa' \varepsilon_\Lambda$ (cf. Definition 11.106 and Section 13.4.7), the flat-FLRW Friedmann law reads

$$H^2(a) = \frac{8\pi G}{3c^2}\,\rho_{\rm tot}(a), \tag{13.217}$$

so at $a = 1$ one recovers $\rho^E_{\rm crit}(0) = \frac{3c^2 H_0^2}{8\pi G}$ [146, 7, 144].

(iii) State-first definition of observable mass On the BRST–physical/GNS representation, the observable mass is the expectation

$$M_{\rm obs} = \frac{1}{c^2}\left\langle \mathring{\mathbf{\Psi}} \right| \int_{\mathcal{X}_{\rm obs}} \sqrt{\gamma}\, T^{00}_{\rm tot}\, d^3x \left| \mathring{\mathbf{\Psi}} \right\rangle_{\rm phys} = \frac{\rho_{\rm tot}\, V}{c^2}, \tag{13.218}$$

and homogeneity reduces (13.218) to eq. (13.211). Combining with eq. (13.217) gives (13.212).

(iv) Sectoral masses as projector masses Because $\rho_{\rm tot}$ partitions as in eq. (13.216), each component satisfies $M_i = (\rho_i V)/c^2 = \Omega_i\, M_{\rm obs}$, yielding eq. (13.215).

Non-ideal HSP bookkeeping. With $\Omega_{\rm cross} := \delta_{\rm cross}/\rho_{\rm tot}$,

$$\Omega_{\rm DE} + \Omega_{\rm DM} + \Omega_{\rm vis} + \Omega_{\rm cross} = 1. \tag{13.219}$$

Notes and caveats. (i) The estimate refers to the *observable* universe (past-light-cone bounded, comoving sphere). In ΛCDM the global spatial extent is almost surely infinite, so $M_{\rm obs}$ is not the mass of the entire (unbounded) universe.
(ii) Varying R within the usual range (e.g. 46.1–46.5 Gly) shifts $M_{\rm obs}$ by only a few percent (e.g. $R = 46.5$ Gly $\Rightarrow M_{\rm obs} \approx 3.04 \times 10^{54}$ kg).
(iii) The "mass" of dark energy above is an energy–equivalent E/c^2; dark energy also gravitates via pressure [144].
(iv) Within the IQR/chrono–geometric framework the numbers realize the projector-derived densities $\{\rho_{\rm DE}, \rho_{\rm DM}, \rho_{\rm vis}\}$ and satisfy the closure checks $\sum_i \Omega_i = 1$ and $\rho_{\rm tot}(0) = \rho^E_{\rm crit}(0)$ (cf. Section 13.4.7).
(v) Equation (13.212) is an identity of flat–FLRW kinematics; its *numerical* value uses today's measured H_0 and the chosen observable radius R. The framework's predictive content here is the *closure and partition* via HSP projectors and the mapping $\Lambda_{\rm eff} = \Lambda + \kappa' \varepsilon_\Lambda$; the absolute $M_{\rm obs}$ is then a consequence, not a parameter–free prediction.

Chronogeometric formulation (no FLRW gauge). Fix an observer congruence u^a selected by the HSP/chronogeometric state and a physical slice Σ_τ (e.g. constant chronogeometric time). Let $\mathcal{D}_\tau := \Sigma_\tau \cap J^-(\mathcal{O})$ be the intersection of Σ_τ with the past light cone of the observer worldline $\mathcal{O}$, with two–sphere boundary $S_\tau := \partial \mathcal{D}_\tau$ and induced metric γ_{ij} on Σ_τ.

Definition 13.155 (Observable mass without FLRW gauge)**.** The observable mass is the BRST–physical expectation of the quasilocal energy inside S_τ:

$$M_{\rm obs}(\tau) := \frac{1}{c^2}\left\langle \mathring{\mathbf{\Psi}} \right| \int_{\mathcal{D}_\tau} \sqrt{\gamma}\, \rho_{\rm tot}(x;\tau)\, d^3x \left| \mathring{\mathbf{\Psi}} \right\rangle_{\rm phys} + \frac{c^2}{8\pi G}\left\langle \mathring{\mathbf{\Psi}} \right| \oint_{S_\tau} (k - k_0)\, dS \left| \mathring{\mathbf{\Psi}} \right\rangle_{\rm phys}, \tag{13.220}$$

where $\rho_{\rm tot} := T_{ab}u^a u^b$, k is the trace of the extrinsic curvature of $S_\tau \subset \Sigma_\tau$, and k_0 is the reference embedding term (Brown–York subtraction) chosen to match the chronogeometric vacuum with $\Lambda_{\rm eff} = \Lambda + \kappa'\varepsilon_\Lambda$ on S_τ.

Proposition 13.156 (Sectoral partition and closure (chronogeometric))**.** *With HSP projectors on the Holo–State, the sectoral energy densities $\rho_{\rm DE}, \rho_{\rm DM}, \rho_{\rm vis}$ are defined by projector–filtered expectations and obey*

$$\begin{aligned} \rho_{\rm tot} = \rho_{\rm DE} + \rho_{\rm DM} + \rho_{\rm vis} \quad \Rightarrow \quad M_{\rm obs} &= \sum_i M_i, \quad \textit{where} \\ M_i &:= \frac{1}{c^2}\Big\langle \int_{\mathcal{D}_\tau} \sqrt{\gamma}\,\rho_i\, d^3x \Big\rangle_{\rm phys} \\ &\quad + \frac{c^2}{8\pi G}\Big\langle \oint_{S_\tau} (k_i - k_{0,i})\, dS \Big\rangle_{\rm phys}. \end{aligned} \tag{13.221}$$

In the ideal HSP limit, cross–sector leakage vanishes and one has exact closure.

Reduction to the FLRW identity. If the slice is (approximately) homogeneous and isotropic on $\mathcal{D}_\tau$ and spatial curvature is negligible on S_τ, the boundary term reproduces the flat–space reference ($k \simeq k_0$) and $\rho_{\rm tot}$ is spatially constant, giving

$$M_{\rm obs} \;=\; \frac{\rho_{\rm tot}\, V}{c^2} \overset{\text{Einstein eqs.}}{=} \frac{H^2}{2G}\, R^3 \quad \text{at the evaluation epoch,}$$

which is exactly (13.212) when evaluated at $a = 1$.

Observational evaluation (flat case, today). Given an observed Hubble constant H_0 and an adopted comoving particle–horizon radius R, the observable mass–energy follows from

$$M_{\rm obs} = \frac{H_0^2}{2G}\, R^3, \tag{13.222}$$

$$\rho^E_{\rm crit}(0) = \frac{3c^2 H_0^2}{8\pi G}, \qquad V = \frac{4\pi}{3} R^3, \qquad E_{\rm obs} = \rho^E_{\rm crit}(0)\, V, \tag{13.223}$$

with sectoral masses $M_i = \Omega_i\, M_{\rm obs}$ for $i \in \{\mathrm{DE}, \mathrm{DM}, \mathrm{vis}\}$.

Numerical inputs (Planck-like): $H_0 = 67.4\ \mathrm{km\,s^{-1}\,Mpc^{-1}}$, $R \in \{46.1, 46.5\}$ Gly, $(\Omega_{\rm DE}, \Omega_{\rm DM}, \Omega_{\rm vis}) = (0.685,\, 0.265,\, 0.050)$.

Table 13.2: Primary Cosmological Parameters for Different Observable Radii

R [Gly]	V [m^3]	$\rho^E_{\rm crit}(0)$ [$\mathrm{J\,m^{-3}}$]	$E_{\rm obs}$ [J]	$M_{\rm obs}$ [kg]
46.1	3.475×10^{80}	7.669×10^{-10}	2.665×10^{71}	2.965×10^{54}
46.5	3.566×10^{80}	7.669×10^{-10}	2.735×10^{71}	3.043×10^{54}

Table 13.3: Mass Component Breakdown Corresponding to Table 13.2

R [Gly]	$M_{\rm obs}$ [kg]	$M_{\rm DE}$ [kg]	$M_{\rm DM}$ [kg]	$M_{\rm vis}$ [kg]
46.1	2.965×10^{54}	2.031×10^{54}	7.858×10^{53}	1.483×10^{53}
46.5	3.043×10^{54}	2.085×10^{54}	8.064×10^{53}	1.522×10^{53}

Uncertainty propagation (first order). For small fractional uncertainties,

$$\frac{\delta M_{\rm obs}}{M_{\rm obs}} \;\simeq\; 2\,\frac{\delta H_0}{H_0} \;+\; 3\,\frac{\delta R}{R}. \tag{13.224}$$

For example, $\delta H_0/H_0 \simeq 0.7\%$ and $\delta R/R \simeq 0.9\%$ imply $\delta M_{\rm obs}/M_{\rm obs} \approx 2\times 0.7\% + 3\times 0.9\% \approx 4.1\%$.

Remark 13.157 (Synthesis: observable mass in FLRW vs. chronogeometric)**. Equivalence.** The chronogeometric (Route A) quasilocal mass

$$M_{\rm obs}(\tau) = \frac{1}{c^2}\Big\langle \int_{\mathcal{D}_\tau} \sqrt{\gamma}\, T^{00}_{\rm tot}\, d^3x \Big\rangle_{\rm phys} + \frac{c^2}{8\pi G}\Big\langle \oint_{S_\tau} (k - k_0)\, dS \Big\rangle_{\rm phys}$$

reduces, for a homogeneous/flat domain, to the FLRW identity $M_{\rm obs} = \frac{H_0^2}{2G}\, R^3$ evaluated at $a = 1$.

Kinematics vs. predictions. Given (H_0, R), the absolute $M_{\rm obs}$ is kinematic. The framework's predictive content is the HSP projector partition $\rho_{\rm tot} = \rho_{\rm DE} + \rho_{\rm DM} + \rho_{\rm vis}$ and the mapping $\Lambda_{\rm eff} = \Lambda + \kappa'\varepsilon_\Lambda$, so $M_i = \Omega_i\, M_{\rm obs}$ with $\sum_i \Omega_i = 1$ (ideal HSP).

Uncertainties. $\delta M/M \simeq 2\,\delta H_0/H_0 + 3\,\delta R/R$.

Beyond FLRW. If curvature/inhomogeneity on S_τ is non-negligible, the boundary term does not cancel; Route A remains valid and quantifies the deviation, while the FLRW shortcut does not.

A Universe Sculpted by Coherence

Within the Interdependent Quantum Reality (IQR) framework, the cosmos realized in nature is the fixed point of the two–stage *Holosymmetric Selection Principle* (HSP). Stage I imposes *self–coherence* on the BRST–physical space, and Stage II then *maximizes the physical holosymmetric overlap* among viable histories (Section 12.2). Applied at cosmological scale to the universal Holo–State $\big|\mathring{\mathbf{\Psi}}\big\rangle$, this selection yields a natural trichotomy of energy components—visible, dark–matter, and dark–energy—each identified with a distinct feature of the universal interference structure (Sections 13.2.8 and 13.3).

(i) Visible Sector: the Primary Bright Fringe. The observable universe—particles and interactions governed by $G_{\rm SM}$—corresponds to the maximally constructive interference peak singled out by Stage II. Operationally, Standard–Model states have nonzero holosymmetric visibility under the SM coherence projector $\mathcal{P}^{\rm coh}_{G_{\rm SM}}$ and are not spectrally orthogonal to SM

observables; hence they populate the dominant "bright" fringe in ordinary phenomena (Definition 12.77 and Section 1.4.1). Their energy density ρ_{vis} is computed by the projector–filtered expectation in eq. (13.148).

(ii) Dark Energy: a Symmetric Interferometric Residue. The small, positive vacuum energy driving late–time acceleration is the *global interferometric residue* of the dual–field vacuum (Gaiamiti + Tantrakyma) after the symmetric lensing filter removes incoherent contributions (Section 13.2.7). This interprets the observed acceleration [134, 135, 136] not as fine–tuning but as the coherent vacuum remainder $\rho_{\text{DE}} = \varepsilon_\Lambda$ selected by Stage II (Section 13.2.6 and eq. (13.145)). In the temporal language, this background is set by the Apeiron (scalar) sector, while its fluctuations propagate through the scalar piece of the chrono–graviton (Section 13.3).

(iii) Dark Matter: Spectrally Orthogonal Coherent Excitations. Dark matter consists of stable excitations (notably the vector *Chrono-Kymon* and, in some regimes, the scalar Apeiron) that pass the Stage I lensing test yet remain *spectrally decoupled* from the SM. Concretely, such states satisfy the coherence criterion $\|\mathcal{P}_G^{\text{coh}}|\psi\rangle\|^2 > 0$ and the darkness condition $\langle\psi|\mathcal{O}_{\text{vis}}|\psi\rangle_{\text{phys}} \approx 0$ for all SM observables (Section 13.2.7 and eqs. (13.85) and (13.86)). Their energy density ρ_{DM} is the holosymmetric–projected expectation in eq. (13.146) (or the equivalent spectral sum eq. (13.147)); their large–scale halos arise from coherent temporal structure in the vector sector (Sections 13.1 and 13.3).

Consequence. Holosymmetry thus dictates both *what exists* and *how much of it exists*. The spectral partitions $\rho_{\text{DE}}, \rho_{\text{DM}}, \rho_{\text{vis}}$ and the abundance fractions $\Omega_{\text{DE}}, \Omega_{\text{DM}}, \Omega_{\text{vis}}$ follow from projector–filtered expectations and their ratios (eqs. (13.145), (13.146), (13.148) and (13.152) and Proposition 13.97). In this sense, the geometry of quantum coherence—made operational by the interferometer model and diagnosed by the lensing formalism—*sculpts* both the visible and unseen architecture of the universe and its temporal fate (Sections 2.4, 11.9 and 13.2.7). The cosmic history becomes a *dynamical partition* in which holosymmetric coherence selects, epoch by epoch, the sector that dominates the energy budget, while preserving the isolation of the TT gravitational channel and ensuring a smooth recovery of ΛCDM in the decohered limit (Section 13.3).

13.6 The Universal Quantum Constraint of the Emergent Cosmos

Motivation and scope. In the IQR framework, "time'' is not a fundamental background parameter but an emergent, relational construct encoded in the correlations of the universal Holo–State $\left|\mathring{\mathbf{\Psi}}\right\rangle$. Consequently, cosmic dynamics cannot be posed as a standard Schrödinger evolution with respect to an external time. Instead, the theory adopts a *single quantum constraint* on BRST–physical states, in the spirit of Dirac's quantization of constrained systems and the Wheeler–DeWitt paradigm [269, 442, 156, 605, 606, 83], with further canonical perspectives surveyed in [607].

The timeless thesis advances along a single, coherent pipeline: **Tetralemma effects with Born valuations** $v_M(\rho) = \mathrm{Tr}(E_M\rho)$ instantiate the logic (see also [287]); the **GNS** lift furnishes the operational substrate $\left(\mathcal{H}_{\mathrm{em}}, |\mathring{\Psi}\rangle\right)$ [30, 381, 230]; UV/IR **band** tags with BRST/Nielsen hygiene certify the arena (cf. [120, 91, 383]); this supports a total relational Hamiltonian $\hat{H}_{\mathrm{univ}}$ whose action on the Holo–State yields the universal constraint; and, finally, **conditioning on an internal clock** recovers subsystem Schrödinger dynamics (cf. Sections 13.6 and 13.6.2 and [215, 3]). In this framework the **Quantum Holo–State** is not an ansatz but the cosmological boundary datum that coherently encodes the chronogeometry $\bar{\Sigma}_{\mu\nu}$ and the Tantrakyma content across scales (table 11.7).

Applying the GNS construction to the foundational dialectic algebra yields a **unique cyclic representation** with emergent Hilbert space $\mathcal{H}_{\mathrm{em}}$ and cyclic vector $|\mathring{\Psi}\rangle$; this is the **Quantum Holo–State**, the mathematical embodiment of the whole cosmos $\hat{\mathbf{\Omega}}(x)$ and the definitive boundary condition for cosmological evolution (Chapter 6, Section 6.1, and table 11.7), in line with the operator–algebraic foundations [30, 381, 230].

To ensure well-posed dynamics and readouts, the theory installs **scale tags**—the band functor $\mathfrak{B}_{\lambda_{\mathrm{UV}},\ell} := \left(P_x(\lambda_{\mathrm{UV}}),\, K_{\mathrm{IR}}^{(\ell)}\right)$—and enforces **BRST/Nielsen** flatness so that expectation-level observables commute with the physical projector and retain adjoint-covariant Ward structure across admissible retunes. The **plateau** criterion guarantees stability over $(\lambda_{\mathrm{UV}}, \ell)$, and on this diagnostic realized reference configuration the **operator-bundle connection** and finite effective fibres are realized, making KL-positivity and Ward-descent statements available on $\mathcal{H}_{\mathrm{phys}}$ [120, 383, 91]. Together these deliver a **gauge-/tag-stable, diagnostic realized reference configuration-certified arena** for total relational dynamics.

Crucially, the temporal register is **derived** rather than postulated. The τ-kairos tensor $\hat{\tau}_{\mu\nu} = \delta\hat{t}_{\mu\nu}$ is **TT-null** and decomposes into a scalar **Gaiaton** strand and a vector **Chrono–Kymon** strand, cleanly separated from spin-2 propagation; this identifies clock-capable degrees of freedom without contaminating radiative graviton channels. Proper time later used in conditioning is the emergent t_{geom} defined on $(\mathcal{X}_q, \bar{\Sigma})$ in the same GNS arena, consistent with the standard chronogeometric notion of proper time [7, 146]. This secures a **timekeeping subsystem** inside the relational stack and prepares the ground for conditional evolution.

With the diagnostic realized reference configuration, connection, and sector architecture in place, **logical closure** forces a timeless universal description. Because $|\mathring{\Psi}\rangle$ represents the **total, self-contained universe**, no **external** time parameter can meaningfully act upon it; any such t would presuppose a clock outside the cosmos. Hence the generator of total evolution must **annihilate** the total state, yielding the **Universal Quantum Constraint** of Section 13.6:

$$\hat{H}_{\mathrm{univ}} \left|\mathring{\Psi}\right\rangle = 0 \quad \text{(see Theorem 13.168)}.$$

This is the treatise's Wheeler–DeWitt–type equation: physical content resides in **fixed correlations** internal to the timeless state, not in evolution relative to an external clock [156, 3].

At the same time, the universal Hamiltonian includes all necessary **boundary contributions** (e.g., Brown–York–type terms) when present, and the constraint holds at the universal (closed) level [535, 532, 43]. These clarifications certify that the staticity claim is not a gauge artefact but a structural statement about the closed-system representation on $\left(\mathcal{H}_{\mathrm{em}}, |\mathring{\Psi}\rangle\right)$.

This timelessness immediately raises the familiar **Problem of Time**: how do we reconcile $\hat{H}_{\text{univ}}|\overset{\circ}{\boldsymbol{\Psi}}\rangle = 0$ with the **manifest evolution** of the subsystems we observe? The resolution demanded by the axioms is **relational**: time must **emerge** from within the Holo–State by conditioning on an internal clock degree of freedom tied to the emergent chronogeometry (cf. Section 5.3.8), in line with conditional-evolution approaches [215, 605, 606, 607, 3]. The Page–Wootters-type construction thus becomes natural in this certified setting.

Formally, one factors $\mathcal{H}_{\text{em}} = \mathcal{H}_{\text{system}} \otimes \mathcal{H}_{\text{env}}$, identifies in $\mathcal{H}_{\text{env}}$ a **good** clock C, and **conditions** the timeless state on a reading $|t\rangle$ of C. The resulting **conditional state** (or density operator) of the system evolves according to the emergent **Schrödinger equation**,

$$i\hbar\,\partial_t\,\varrho_{A|C=t} = \Big[H_{A|C=t},\ \varrho_{A|C=t}\Big] \qquad \text{(cf. eq. (13.238))}$$

(or, in the pure-state limit, $i\hbar\,\partial_t|\psi(t)\rangle_{\text{system}} \approx \hat{H}_{\text{system}}|\psi(t)\rangle_{\text{system}}$), exactly as stated in Section 13.6.2 [215].

The **clock** is not introduced ad hoc: Chapters 9–11 derive its architecture from the τ-kairos sector, proving **TT-nullity** and thus kinematic isolation from spin-2 while maintaining the ability to register proper time on the emergent chronogeometry. This rigorously **partitions** the universal state into a clock C and a system A, justifying the conditioning and ensuring that the emergent dynamics inherit BRST/Nielsen flatness from the certified plateau [120].

Bridging explicitly from the **Standard-Model** (Ch. 12) and **dark-sector** (Ch. 13) narratives: Stage-I **anomaly meters** must vanish, the global $SU(2)$ Witten test pass, and the **unitarity operator** act as identity on the certified support; color-singlet enforcement and Nielsen/KL hygiene then realize visible-sector composites as BRST-physical on $\mathcal{H}_{\text{phys}}$ [553, 91, 383]. On the same diagnostic realized reference configuration, **observable mass** is defined quasilocally with a Brown–York boundary, $M_{\text{obs}}(\tau)$, and the **HSP partition** $\rho_{\text{tot}} = \rho_{\text{DE}} + \rho_{\text{DM}} + \rho_{\text{vis}}$ yields closed additive sector masses. With these in hand—and with a TT-safe internal clock—Chapter 13 assembles the **total relational Hamiltonian** as the sum of certified (gravity+matter+boundary) contributions, and **universal closed-system logic** then forces

$$\hat{H}_{\text{univ}}\,\Big|\overset{\circ}{\boldsymbol{\Psi}}\Big\rangle = 0 \qquad \text{(see Theorem 13.168)}.$$

(cf. Section 13.6.)

In summary, the justification of the **Universal Quantum Constraint** proceeds without gaps: the **GNS Holo–State** sets the global stage; **band/BRST/Ward** certification stabilizes the arena; the τ-kairos sector supplies an internal clock; and **conditioning** turns a timeless universal equation into **timeful subsystem dynamics**. Put compactly:

$$\begin{aligned}\text{Tetralemma} &\rightsquigarrow \text{GNS} \rightsquigarrow \text{certified operator geometry}\\ &\rightsquigarrow \hat{H}_{\text{univ}}|\overset{\circ}{\boldsymbol{\Psi}}\rangle = 0 \rightsquigarrow \text{emergent Schrödinger}\,.\end{aligned}$$

This is precisely the arc articulated in Sections 13.6 and 13.6.2.

13.6.1 The Good-clock

Good-clock (formal notion). In the Chapter 13/stack framework, a *good-clock* C is a BRST-physical, TT-safe clock lane that permits conditioning of the timeless Holo–State without

introducing gauge/Ward anomalies or spurious dissipation. Concretely, C is specified by a PVM or POVM of clock readings—either a projector family $\{\Pi_t\}_{t\in\mathbb{R}}$ or effects $\{E_t\}_{t\in\mathbb{R}}$—together with a clock Hamiltonian $\hat{H}^{\text{clock}}$ acting on the clock lane. **Clock covariance** requires the time-shift law

$$e^{-\frac{i}{\hbar}s\hat{H}^{\text{clock}}}\,\Pi_t\,e^{\frac{i}{\hbar}s\hat{H}^{\text{clock}}}=\Pi_{t+s}\quad\Longleftrightarrow\quad\partial_t\Pi_t=-\tfrac{i}{\hbar}\,[\hat{H}^{\text{clock}},\Pi_t],$$

(and analogously for POVMs), which is the spectral/Heisenberg form of a one-parameter unitary group generated by a self-adjoint $\hat{H}^{\text{clock}}$ (Stone's theorem; functional-analytic setting) [369]. In the canonical idealization one has $[\hat{t},\hat{p}_t]=i\hbar$ and $\hat{H}^{\text{clock}}\approx\hat{p}_t$ (Section 5.3.8; cf. [369]).

Relational commutation and conditional states. On the BRST-physical space $\mathcal{H}_{\text{phys}}$, a good clock satisfies the *relational commutation* condition: the algebra $\mathcal{A}$ of subsystem observables commutes with the clock readout, $[A,\Pi_t]=0$ for all $A\in\mathcal{A}$. This guarantees that the conditional states

$$\varrho_{A|C=t}=\frac{\Pi_t\,\varrho\,\Pi_t}{\mathrm{Tr}(\Pi_t\varrho)}$$

are well-defined and gauge-independent on certified plateaues, and that they reproduce the textbook projective (or Lüders-type) updates at the level of reduced states [287]. In the POVM case, the same definition is implemented via Naimark dilation and Kraus operators, ensuring complete positivity and trace preservation [287] (cf. Section 13.6.2).

BRST/typed cleanliness and TT-safety. A good clock is *BRST-closed* and *typed* so that clock readouts respect the physical cohomology and do not contaminate radiative spin-2 channels:

$$[\hat{Q}_{\text{BRST}},\Pi_t]=0,\qquad \Pi_{\Sigma}^{[\text{TT}]}\text{ commutes appropriately with the clock lane (TT-nullity).}$$

This realizes the clock within the τ–kairos sector (not the TT sector), preserving the York–Lichnerowicz TT decomposition used in the gravity lane (eq. (10.4); see also [442, 532, 43, 44]). The typed/TT structure is part of the canonical placement summarized in Section 13.4.1.

Spectral support and quality conditions. On certified plateaux one imposes $\mathrm{Tr}(\Pi_t\varrho)>0$ to avoid conditioning pathologies. *Good-clock quality* further requires that off-diagonal clock couplings are negligible:

$$\langle t|\hat{H}_{\text{int}}|t'\rangle\approx 0\quad(t\neq t'),$$

so that the effective interaction reduces to the fiber operator $\langle t|\,\hat{H}_{\text{int}}\,|t\rangle$ in the time basis. In the POVM language this is equivalent to a small dissipator $\mathcal{D}_t$ in

$$i\hbar\,\partial_t\,\varrho_{A|C=t}=[\hat{H}_{\text{eff}}(t),\varrho_{A|C=t}]+i\hbar\,\mathcal{D}_t[\varrho_{A|C=t}],$$

with $\|\mathcal{D}_t\|\to 0$ as the measurement sharpens to a PVM and the clock becomes more ideal [287]. The *energy–time tradeoff* (clock imprecision σ_t versus energy spread $\sigma_{H^{\text{clock}}}$) governs this regime and bounds the residual dissipative corrections [22].

Emergent dynamics by Page–Wootters conditioning. When the conditions above hold, Page–Wootters conditioning yields unitary subsystem dynamics compatible with the Universal Quantum Constraint (Section 13.6): for the conditional state one has

$$i\hbar\,\partial_t\,\varrho_{A|C=t} = \big[\hat{H}_{\mathrm{eff}}(t),\,\varrho_{A|C=t}\big], \qquad \hat{H}_{\mathrm{eff}}(t) = \frac{\Pi_t\,\hat{H}_{\mathrm{rest}}\,\Pi_t}{\mathrm{Tr}(\Pi_t\varrho)},$$

and, in the pure-state limit, $i\hbar\,\partial_t\,|\psi(t)\rangle = \hat{H}_{\mathrm{eff}}(t)\,|\psi(t)\rangle$ (eq. (13.238) and Section 13.6.2; see also [215, 3, 607]). In short, a *good-clock* is a diagnostic realized reference configuration-certified, BRST-clean, TT-safe clock lane whose covariance and spectral quality render the conditional Schrödinger picture exact in the PVM case and arbitrarily accurate for POVMs with small $\mathcal{D}_t$.

Definition 13.158 (Good–clock)**.** A *good–clock* C on the BRST–physical space $\mathcal{H}_{\mathrm{phys}}$ is a clock lane $\mathcal{H}_{\mathrm{clock}}$ together with either a PVM $\{\Pi_t\}_{t\in\mathbb{R}}$ or a POVM $\{E_t\}_{t\in\mathbb{R}}$ and a self–adjoint clock Hamiltonian $\hat{H}^{\mathrm{clock}}$ such that, for some subsystem factor $\mathcal{H}_{\mathrm{sub}}$ with $\mathcal{H}_{\mathrm{phys}} = \mathcal{H}_{\mathrm{sub}} \otimes \mathcal{H}_{\mathrm{clock}}$, the following hold:

(i) Clock covariance (Stone law)
For all $s \in \mathbb{R}$,

$$e^{-\frac{i}{\hbar}s\hat{H}^{\mathrm{clock}}}\,\Pi_t\,e^{\frac{i}{\hbar}s\hat{H}^{\mathrm{clock}}} = \Pi_{t+s} \quad\Longleftrightarrow\quad \partial_t\Pi_t = -\tfrac{i}{\hbar}\,[\hat{H}^{\mathrm{clock}},\Pi_t],$$

(and analogously with E_t).

(ii) Relational commutation
If $\mathcal{A} \subset \mathcal{B}(\mathcal{H}_{\mathrm{sub}})$ is the subsystem algebra, then $[A,\Pi_t] = 0$ for all $A \in \mathcal{A}$ and all t.

(iii) BRST/typed cleanliness and TT–safety
$[\hat{Q}_{\mathrm{BRST}},\Pi_t] = 0$ and the clock lane lies in the non-TT sector; equivalently it obeys the TT-purity guardrail with the TT projector (cf. eqs. (13.38) and (13.179)).

(iv) Spectral support
On the working window, $\mathrm{Tr}(\Pi_t\varrho) > 0$ for ϱ the universal state restricted to $\mathfrak{P}$ (certified plateaues; cf. Section 13.6).

(v) Quality (near–diagonality)
In the time basis,

$$\langle t|\hat{H}_{\mathrm{int}}|t'\rangle \approx 0 \quad (t \neq t'),$$

so that the effective interaction reduces to $\langle t|\,\hat{H}_{\mathrm{int}}\,|t\rangle$; in the POVM case this is equivalent to a small dissipator $\mathcal{D}_t$ (cf. Corollary 13.174).

Remark 13.159 (Canonical idealization and deparametrization)**.** In the canonical idealization one has a self–adjoint pair $(\hat{t},\hat{p}_t)$ on $\mathcal{H}_{\mathrm{clock}}$ with $[\hat{t},\hat{p}_t] = i\hbar$, and the Total Relational Hamiltonian admits the deparametrized split

$$\hat{H}_{\mathrm{univ}} = \hat{p}_t \otimes \mathbf{1}_{\mathrm{sub}} + \mathbf{1}_{\mathrm{clock}} \otimes \hat{H}_{\mathrm{sub}} + \hat{H}_{\mathrm{int}}, \qquad \hat{H}_{\mathrm{univ}}\Big|\mathring{\mathbf{\Psi}}\Big\rangle = 0 \quad (\textit{eqs. (13.235) to (13.237)}).$$

Lemma 13.160 (Gauge–independent conditionalization)**.** *Under Definition 13.158(ii)–(iv), the conditional states*

$$\varrho_{\mathrm{sub}|t} = \frac{\Pi_t\,\varrho\,\Pi_t}{\mathrm{Tr}(\Pi_t\varrho)}$$

are well–defined on $\mathcal{H}_{\mathrm{sub}}$, completely positive and gauge–independent on certified plateaues $\mathfrak{P}$; for POVMs the same holds via Naimark dilation/Kraus operators (cf. Section 13.6.2 and Corollary 13.174).

Proof of Lemma 13.160. Let $\mathcal{H}_{\mathrm{phys}} = \mathcal{H}_{\mathrm{sub}} \otimes \mathcal{H}_{\mathrm{clock}}$ and let C be a good–clock in the sense of Definition 13.158(ii)–(iv). Write $\mathcal{I}_t(X) := (\mathbf{1}_{\mathrm{sub}}\otimes\Pi_t)\,X\,(\mathbf{1}_{\mathrm{sub}}\otimes\Pi_t)$ and $p_t(\varrho) := \mathrm{Tr}[\mathcal{I}_t(\varrho)] = \mathrm{Tr}(\Pi_t\varrho)$.

(1) Well-definedness and positivity on $\mathcal{H}_{\mathrm{sub}}$. By Definition 13.158(iv) (*Spectral support*), $p_t(\varrho) > 0$ on the certified window of $\mathfrak{P}$, so the normalized post-selection state

$$\widetilde{\varrho}_t := \frac{\mathcal{I}_t(\varrho)}{p_t(\varrho)} = \frac{\Pi_t\,\varrho\,\Pi_t}{\mathrm{Tr}(\Pi_t\varrho)}$$

is a positive, unit-trace operator supported on $\mathcal{H}_{\mathrm{sub}}\otimes\mathrm{Ran}\,\Pi_t$. Define the *subsystem* conditional state by partial trace over the clock lane,

$$\varrho_{\mathrm{sub}|t} := \mathrm{Tr}_{\mathrm{clock}}\left[\widetilde{\varrho}_t\right].$$

Since partial trace is completely positive (CP) and normal, $\varrho_{\mathrm{sub}|t} \in \mathcal{B}_1^+(\mathcal{H}_{\mathrm{sub}})$ and $\mathrm{Tr}_{\mathrm{sub}}\,\varrho_{\mathrm{sub}|t} = 1$. This shows the expression in the lemma is well-defined as the reduced density operator on $\mathcal{H}_{\mathrm{sub}}$.

(2) Complete positivity (instrument/Kraus view). The map $\mathcal{M}_t : \varrho \mapsto \mathcal{I}_t(\varrho) = (\mathbf{1}\otimes\Pi_t)\,\varrho\,(\mathbf{1}\otimes\Pi_t)$ is a CP, normal map with single Kraus operator $K_t = \mathbf{1}_{\mathrm{sub}}\otimes\Pi_t$ (textbook instrument form). For any ancilla $\mathcal{H}_R$, $(\mathrm{id}_R\otimes\mathcal{M}_t)$ is positive because K_t acts trivially on R. The normalized update $\widetilde{\varrho}_t = \mathcal{M}_t(\varrho)/p_t(\varrho)$ preserves positivity since $p_t(\varrho) > 0$. Composition with the partial trace gives a CP assignment $\varrho \mapsto \varrho_{\mathrm{sub}|t}$.

(3) Gauge–independence on $\mathfrak{P}$. By Definition 13.158(iii) (*BRST/typed cleanliness*), $[\hat{Q}_{\mathrm{BRST}}, \Pi_t] = 0$. Hence the instrument commutes with the BRST differential:

$$\mathcal{I}_t\Big([\hat{Q}_{\mathrm{BRST}}, X]\Big) = [\hat{Q}_{\mathrm{BRST}}, \mathcal{I}_t(X)].$$

Therefore $\mathcal{I}_t$ descends to the physical cohomology $H^0(\hat{Q}_{\mathrm{BRST}})$, i.e. it is independent of the choice of BRST representative of ϱ. Moreover, by typed/TT compatibility the gravity TT projector commutes with the clock lane ($\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}$ from eq. (13.179)), and the realized weave configuration-compatibility guardrail ensures no spurious TT inflow (eq. (13.38)). Consequently the reduced state $\varrho_{\mathrm{sub}|t}$ is gauge-independent and TT-safe on certified plateaues $\mathfrak{P}$.

(4) Relational commutation and subsystem localization. From Definition 13.158(ii) (*Relational commutation*), $[A, \Pi_t] = 0$ for all $A \in \mathcal{A} \subset \mathcal{B}(\mathcal{H}_{\mathrm{sub}})$. Hence, for any subsystem observable A,

$$\mathrm{Tr}_{\mathrm{sub}}\Big(A\,\varrho_{\mathrm{sub}|t}\Big) = \frac{\mathrm{Tr}\Big((A\otimes\Pi_t)\,\varrho\,(\mathbf{1}\otimes\Pi_t)\Big)}{p_t(\varrho)} = \frac{\mathrm{Tr}\Big((A\otimes\Pi_t)\,\varrho\,(\mathbf{1}\otimes\Pi_t)\Big)}{\mathrm{Tr}\Big((\mathbf{1}\otimes\Pi_t)\,\varrho\Big)},$$

so expectation values are computed with the subsystem operator A alone, validating that $\varrho_{\mathrm{sub}|t}$ is the correct (gauge-clean) conditional state on $\mathcal{H}_{\mathrm{sub}}$. This is the conditioning setup used in Section 13.6.2.

(5) POVM clocks (Naimark/Kraus). If C supplies a POVM $\{E_t\}_{t\in\mathbb{R}}$, choose a Naimark dilation $V : \mathcal{H}_{\mathrm{clock}} \to \mathcal{H}_{\mathrm{clock}} \otimes \mathcal{H}_{\mathrm{anc}}$ and a PVM $\{\Pi_t^{(\mathrm{N})}\}$ on the dilation space with $E_t = V^\dagger \Pi_t^{(\mathrm{N})} V$. Define the CP instrument on $\mathcal{H}_{\mathrm{phys}}$ by

$$\mathcal{M}_t^{\mathrm{POVM}}(\varrho) = (\mathbf{1} \otimes V^\dagger \Pi_t^{(\mathrm{N})} V)\, \varrho\, (\mathbf{1} \otimes V^\dagger \Pi_t^{(\mathrm{N})} V) = \sum_\alpha K_{t,\alpha}\, \varrho\, K_{t,\alpha}^\dagger,$$

with Kraus operators $K_{t,\alpha} = \mathbf{1} \otimes V^\dagger \Pi_{t,\alpha}^{(\mathrm{N})}$ from a refinement of $\Pi_t^{(\mathrm{N})}$. Normalization by $p_t(\varrho) = \mathrm{Tr}[\mathcal{M}_t^{\mathrm{POVM}}(\varrho)]$ and partial trace over the clock lane give $\varrho_{\mathrm{sub}|t}$ exactly as above; $[\hat{Q}_{\mathrm{BRST}}, E_t] = 0$ (typed cleanliness) ensures the same gauge-independence. This is the POVM version referenced in Corollary 13.174.

Combining (1)–(5) proves that the conditional states are well–defined on $\mathcal{H}_{\mathrm{sub}}$, completely positive (via the instrument/Kraus construction), and gauge–independent on certified plateaues $\mathfrak{P}$, with the POVM case obtained by Naimark dilation. □

Lemma 13.161 (TT–nullity of the clock lane)**.** *If C satisfies Definition 13.158(iii), then the clock lane cannot source radiative spin–2 modes on $\mathfrak{P}$:*

$$\Pi^{\mathrm{TT}}\, [\textit{clock readouts and couplings}]\, \Pi^{\mathrm{TT}} = 0,$$

with Π^{TT} the TT projector of eq. (13.179); this is a direct consequence of the rig–compatibility condition eq. (13.38).

Proof of Lemma 13.161. Let $\Pi^{\mathrm{TT}} := \Pi_{\Sigma}^{[\mathrm{TT}]}$ be the York–Lichnerowicz transverse–traceless projector of eq. (13.179). By Definition 13.158(iii) (BRST/typed cleanliness and TT–safety), the clock lane belongs to the typed, non–radiative τ-kairos sector. Denote by $\Pi^{(\tau)}$ the corresponding type projector (commuting idempotent in the chapter's typed calculus), so that every clock readout/effect X_{clk} and every admissible clock coupling Y_{clk} satisfy the typed support identities

$$X_{\mathrm{clk}} = \Pi^{(\tau)}\, X_{\mathrm{clk}}\, \Pi^{(\tau)}, \qquad Y_{\mathrm{clk}} = \Pi^{(\tau)}\, Y_{\mathrm{clk}}\, \Pi^{(\tau)}. \tag{13.225}$$

On certified plateaues $\mathfrak{P}$, the rig–compatibility (TT–purity) guardrail eq. (13.38) enforces orthogonality of the radiative spin–2 lane to the τ-typed sector,

$$\Pi^{\mathrm{TT}}\, \Pi^{(\tau)} = \Pi^{(\tau)}\, \Pi^{\mathrm{TT}} = 0 \quad \text{on } \mathfrak{P}, \tag{13.226}$$

and the commuting–projector property of the typed calculus (Chapter XI) implies $[\Pi^{\mathrm{TT}}, \Pi^{(\tau)}] = 0$ on $\mathfrak{P}$.

Clock readouts. Using (13.225) and (13.226),

$$\Pi^{\mathrm{TT}}\, X_{\mathrm{clk}}\, \Pi^{\mathrm{TT}} = \Pi^{\mathrm{TT}}\, \Pi^{(\tau)}\, X_{\mathrm{clk}}\, \Pi^{(\tau)}\, \Pi^{\mathrm{TT}} = 0.$$

Clock couplings. Any admissible coupling involving the clock is typed by $\Pi^{(\tau)}$ and thus obeys (13.225); therefore,

$$\Pi^{\mathrm{TT}}\, Y_{\mathrm{clk}}\, \Pi^{\mathrm{TT}} = \Pi^{\mathrm{TT}}\, \Pi^{(\tau)}\, Y_{\mathrm{clk}}\, \Pi^{(\tau)}\, \Pi^{\mathrm{TT}} = 0.$$

Equivalently, in block form with respect to the TT/τ decomposition, the TT$\to$TT block of every operator generated by clock readouts and their admissible couplings vanishes on $\mathfrak{P}$.

Conclusion. Combining the two cases yields

$$\Pi^{\mathrm{TT}}\,[\text{clock readouts and couplings}]\,\Pi^{\mathrm{TT}} = 0 \quad \text{on } \mathfrak{P},$$

which is precisely the TT–nullity statement of the lemma. The only inputs were typed support for the clock lane from Definition 13.158(iii), the commuting/orthogonal projector calculus from Chapter XI, and the realized weave configuration TT–purity guardrail eq. (13.38). □

Proposition 13.162 (Emergent Schrödinger dynamics (PVM case))**.** *If C is a good–clock in the sense of Definition 13.158 and the split eq. (13.237) holds, then the subsystem conditional vector*

$$|\Psi_{\mathrm{sub}}(t)\rangle = \frac{|\widetilde{\Psi}_{\mathrm{sub}}(t)\rangle}{\big\|\,|\widetilde{\Psi}_{\mathrm{sub}}(t)\rangle\,\big\|}$$

obeys

$$\boxed{i\hbar\,\partial_t\,|\Psi_{\mathrm{sub}}(t)\rangle = \hat{H}^{\mathrm{eff}}_{\mathrm{sub}}(t)\;|\Psi_{\mathrm{sub}}(t)\rangle\,, \qquad \hat{H}^{\mathrm{eff}}_{\mathrm{sub}}(t) := \hat{H}_{\mathrm{sub}} + \langle t|\,\hat{H}_{\mathrm{int}}\,|t\rangle\,,} \tag{13.227}$$

exactly as established in Proposition 13.172 and eq. (13.238).

Proof of Proposition 13.162. Let C be a good–clock as in Definition 13.158 and assume the deparametrized split

$$\hat{H}_{\mathrm{univ}} = \hat{p}_t \otimes \mathbf{1}_{\mathrm{sub}} + \mathbf{1}_{\mathrm{clock}} \otimes \hat{H}_{\mathrm{sub}} + \hat{H}_{\mathrm{int}}, \qquad \hat{H}_{\mathrm{univ}}\Big|\mathring{\mathbf{\Psi}}\Big\rangle = 0, \tag{13.228}$$

which is eq. (13.237). By Definition 13.158(i) (Stone covariance; canonical idealization), the clock admits a sharp PVM $\{\Pi_t\}$ with generalized eigenkets $|t\rangle$ and $\hat{p}_t = -i\hbar\,\partial_t$ in the t-representation (cf. Remark 13.159). Define the (unnormalized) conditional subsystem vector

$$|\widetilde{\Psi}_{\mathrm{sub}}(t)\rangle := (\mathbf{1}_{\mathrm{sub}}\otimes\langle t|)\Big|\mathring{\mathbf{\Psi}}\Big\rangle \;\in\; \mathcal{H}_{\mathrm{sub}}, \qquad p_t := \big\|\,|\widetilde{\Psi}_{\mathrm{sub}}(t)\rangle\,\big\|^2 = \mathrm{Tr}(\Pi_t\,\varrho), \tag{13.229}$$

which is well-defined and satisfies $p_t > 0$ on the certified window by Definition 13.158(iv) (Spectral support).

Step 1: Projecting the universal constraint. Left-multiplying (13.228) by $(\mathbf{1}_{\mathrm{sub}}\otimes\langle t|)$ and using $\langle t|\,\hat{p}_t = -i\hbar\,\partial_t\,\langle t|$ yields

$$-i\hbar\,\partial_t\,|\widetilde{\Psi}_{\mathrm{sub}}(t)\rangle + \hat{H}_{\mathrm{sub}}\,|\widetilde{\Psi}_{\mathrm{sub}}(t)\rangle + \int_{\mathbb{R}} dt'\;\underbrace{\langle t|\,\hat{H}_{\mathrm{int}}\,|t'\rangle}_{=:K(t,t')}\;|\widetilde{\Psi}_{\mathrm{sub}}(t')\rangle \;=\; 0. \tag{13.230}$$

Step 2: Good–clock near–diagonality. By Definition 13.158(v) (Quality / near–diagonality), $K(t,t') = \langle t|\,\hat{H}_{\mathrm{int}}\,|t'\rangle \approx \delta(t-t')\;\langle t|\,\hat{H}_{\mathrm{int}}\,|t\rangle$ on $\mathfrak{P}$. Thus the integral term in (13.230) reduces

to a multiplicative (fiber) operator and we obtain the *closed* evolution for the unnormalized vector:

$$i\hbar\,\partial_t\,|\widetilde{\Psi}_{\rm sub}(t)\rangle = \Big(\hat{H}_{\rm sub} + \langle t|\,\hat{H}_{\rm int}\,|t\rangle\Big)\,|\widetilde{\Psi}_{\rm sub}(t)\rangle =: \hat{H}^{\rm eff}_{\rm sub}(t)\;|\widetilde{\Psi}_{\rm sub}(t)\rangle\,. \tag{13.231}$$

Since $\hat{H}_{\rm sub}$ and $\hat{H}_{\rm int}$ are self-adjoint and the partial sandwich by $|t\rangle$ preserves self-adjointness on $\mathcal{H}_{\rm sub}$, $\hat{H}^{\rm eff}_{\rm sub}(t)$ is self-adjoint for each t.

Step 3: Norm constancy for the unnormalized vector. Taking the t-derivative of $p_t = \langle\widetilde{\Psi}_{\rm sub}(t)\mid\widetilde{\Psi}_{\rm sub}(t)\rangle$ and using (13.231) gives

$$\partial_t p_t = \frac{1}{i\hbar}\Big(\langle\widetilde{\Psi}|\,\hat{H}^{\rm eff}_{\rm sub}(t)\,\widetilde{\Psi}\rangle - \langle\widetilde{\Psi}|\,\hat{H}^{\rm eff}_{\rm sub}(t)\,\widetilde{\Psi}\rangle\Big) = 0,$$

so p_t is constant on the certified window. Hence normalization does not affect the equation of motion.

Step 4: Evolution of the normalized vector. Define the normalized conditional vector $|\Psi_{\rm sub}(t)\rangle = |\widetilde{\Psi}_{\rm sub}(t)\rangle\,/\|\,|\widetilde{\Psi}_{\rm sub}(t)\rangle\,\|$. Since p_t is constant, differentiating and using (13.231) yields

$$i\hbar\,\partial_t\,|\Psi_{\rm sub}(t)\rangle = \hat{H}^{\rm eff}_{\rm sub}(t)\;|\Psi_{\rm sub}(t)\rangle\,.$$

This is precisely the conditional Schrödinger equation with $\hat{H}^{\rm eff}_{\rm sub}(t) = \hat{H}_{\rm sub} + \langle t|\,\hat{H}_{\rm int}\,|t\rangle$, matching Proposition 13.172 and eq. (13.238) and giving (13.227).

All steps take place on certified plateaues $\mathfrak{P}$, and compatibility with BRST/typed hygiene from Definition 13.158(iii) ensures that no gauge or TT contamination enters the subsystem lane (though this hygiene is not needed explicitly in the algebra above). □

Corollary 13.163 (Density–operator form and POVM clocks)**.** *For a PVM clock,*

$$i\hbar\,\partial_t\varrho_{{\rm sub}|t} = \Big[\hat{H}^{\rm eff}_{\rm sub}(t),\,\varrho_{{\rm sub}|t}\Big].$$

For a POVM clock $\{E_t\}$ one obtains

$$i\hbar\,\partial_t\varrho_{{\rm sub}|t} = \Big[\hat{H}^{\rm eff}_{\rm sub}(t),\,\varrho_{{\rm sub}|t}\Big] + i\hbar\,\mathcal{D}_t[\varrho_{{\rm sub}|t}],$$

with $\|\mathcal{D}_t\| \to 0$ in the sharp limit, as in Corollary 13.174.

Proof of Corollary 13.163. We treat the sharp (PVM) and unsharp (POVM) cases in turn, always on certified plateaues $\mathfrak{P}$. Throughout, C is a good–clock (Definition 13.158), and the deparametrized split *eq.* (13.237) is assumed.

PVM case. Let $\{\Pi_t\}$ be the clock PVM and let $|\widetilde{\Psi}_{\rm sub}(t)\rangle := (\mathbf{1}_{\rm sub}\otimes\langle t|)\big|\mathring{\mathbf{\Psi}}\big\rangle$ with normalized conditional vector $|\Psi_{\rm sub}(t)\rangle = |\widetilde{\Psi}_{\rm sub}(t)\rangle\,/\|\,|\widetilde{\Psi}_{\rm sub}(t)\rangle\,\|$. By Proposition 13.162 one has $i\hbar\,\partial_t\,|\Psi_{\rm sub}(t)\rangle = \hat{H}^{\rm eff}_{\rm sub}(t)\,|\Psi_{\rm sub}(t)\rangle$ with $\hat{H}^{\rm eff}_{\rm sub}(t) = \hat{H}_{\rm sub} + \langle t|\,\hat{H}_{\rm int}\,|t\rangle$ (eq. (13.227)). Define $\varrho_{{\rm sub}|t} := |\Psi_{\rm sub}(t)\rangle\langle\Psi_{\rm sub}(t)|$. Then, using the product rule and the Schrödinger equation,

$$\begin{aligned} i\hbar\,\partial_t\varrho_{{\rm sub}|t} &= i\hbar\Big((\partial_t\,|\Psi_{\rm sub}\rangle)\,\langle\Psi_{\rm sub}| + |\Psi_{\rm sub}\rangle\,\partial_t\,\langle\Psi_{\rm sub}|\Big) \\ &= \hat{H}^{\rm eff}_{\rm sub}(t)\,|\Psi_{\rm sub}\rangle\,\langle\Psi_{\rm sub}| - |\Psi_{\rm sub}\rangle\,\langle\Psi_{\rm sub}|\,\hat{H}^{\rm eff}_{\rm sub}(t) \\ &= \Big[\hat{H}^{\rm eff}_{\rm sub}(t),\;\varrho_{{\rm sub}|t}\Big]. \end{aligned}$$

By convex linearity, the same equation holds for mixed $\varrho_{\mathrm{sub}|t}$ obtained from a mixed universal ϱ (cf. Lemma 13.160). This proves the first line.

POVM case. Let the clock be specified by a POVM $\{E_t\}_{t\in\mathbb{R}}$. Choose a Naimark dilation $V : \mathcal{H}_{\mathrm{clock}} \to \mathcal{H}_{\mathrm{clock}} \otimes \mathcal{H}_{\mathrm{anc}}$ and a PVM $\{\Pi_t^{(\mathrm{N})}\}$ with $E_t = V^\dagger \Pi_t^{(\mathrm{N})} V$. Using a refinement $\{\Pi_{t,\alpha}^{(\mathrm{N})}\}_\alpha$ of $\Pi_t^{(\mathrm{N})}$, define Kraus operators $K_{t,\alpha} := \mathbf{1}_{\mathrm{sub}} \otimes V^\dagger \Pi_{t,\alpha}^{(\mathrm{N})} V$ and the CP instrument

$$\mathcal{M}_t(\varrho) = \sum_\alpha K_{t,\alpha}\, \varrho\, K_{t,\alpha}^\dagger, \qquad p_t = \mathrm{Tr}\left[\mathcal{M}_t(\varrho)\right].$$

The (normalized) post-readout state is $\widetilde{\varrho}_t := \mathcal{M}_t(\varrho)/p_t$, and the subsystem conditional state is $\varrho_{\mathrm{sub}|t} = \mathrm{Tr}_{\mathrm{clock}}\left[\widetilde{\varrho}_t\right]$ (Lemma 13.160). Differentiating and using the deparametrized split together with the good–clock near–diagonality (quality) condition, one obtains a time-local generator consisting of the coherent part from $\hat{H}_{\mathrm{sub}}^{\mathrm{eff}}(t)$ (exactly as in the PVM case) *plus* a CP, trace-preserving remainder that collects (i) the unsharpness of E_t via the Kraus spread and (ii) the possible time-dependence of the instrument:

$$i\hbar\, \partial_t \varrho_{\mathrm{sub}|t} = \left[\hat{H}_{\mathrm{sub}}^{\mathrm{eff}}(t),\ \varrho_{\mathrm{sub}|t}\right] + i\hbar\, \mathcal{D}_t[\varrho_{\mathrm{sub}|t}].$$

Here $\mathcal{D}_t$ is completely positive and trace-preserving (CPTP) in the sense that it arises from CP maps $\mathcal{M}_t$ and normalization/partial-trace operations; in particular, in a (time-local) Lindblad realization one may write

$$\mathcal{D}_t[\rho] = \sum_\alpha L_{t,\alpha}\, \rho\, L_{t,\alpha}^\dagger - \tfrac{1}{2} \sum_\alpha \left\{ L_{t,\alpha}^\dagger L_{t,\alpha},\ \rho \right\},$$

with $L_{t,\alpha}$ determined by the Kraus spread of the POVM dilation and vanishing when that spread collapses to a single projector (sharp measurement). By the good–clock quality requirement, the POVM unsharpness can be made arbitrarily small on $\mathfrak{P}$, which implies $\|\mathcal{D}_t\| \to 0$ in the sharp limit $E_t \Rightarrow \Pi_t$, recovering the PVM commutator form. This is precisely the statement recorded in Corollary 13.174.

Combining the two cases establishes the corollary. □

Proposition 13.164 (Portal robustness of good–clocks)**.** *Let C be a good–clock and let the dynamics be perturbed by an admissible portal family $\mathcal{L}_{\mathrm{portal}}$ with small coefficients $\boldsymbol{\varepsilon}$ (cf. Definition 13.44, Assumption 13.45, and eq. (13.64)). Then Stage II interferometric functionals and scores are Lipschitz–stable:*

$$\left|\mathcal{V}_{\mathrm{asym}}[H[\boldsymbol{\varepsilon}], H[\boldsymbol{\varepsilon}]] - \mathcal{V}_{\mathrm{asym}}[H, H]\right| \le C_{\mathrm{port}}\, \|\boldsymbol{\varepsilon}\|_\infty + \mathcal{O}(\|\boldsymbol{\varepsilon}\|_\infty^2),$$
$$\left|\mathscr{C}(H[\boldsymbol{\varepsilon}]) - \mathscr{C}(H)\right| \le C'_{\mathrm{port}}\, \|\boldsymbol{\varepsilon}\|_\infty + \cdots$$

as in eqs. (13.66) and (13.67). In particular, the conditional evolution of Proposition 13.162 is stable under admissible portals.

Proof of Proposition 13.164. Work on a certified plateau $\mathfrak{P}$. Let C be a good–clock (Definition 13.158) and let the portal deformation be specified by the admissible family $\mathcal{L}_{\mathrm{portal}} = \sum_j \varepsilon_j\, \mathcal{O}_j$ with small coefficients $\boldsymbol{\varepsilon}$ as in Definition 13.44, Assumption 13.45,

and eq. (13.64). By the typed/BRST/DRRC hygiene in Assumption 13.45 the induced Hamiltonian perturbation on $\mathfrak{P}$ is a bounded operator

$$\Delta\hat{H}(\boldsymbol{\varepsilon}) = \sum_j \varepsilon_j \hat{O}_j \quad \text{with} \quad \|\Delta\hat{H}(\boldsymbol{\varepsilon})\| \le K_{\text{port}} \|\boldsymbol{\varepsilon}\|_\infty, \tag{13.232}$$

and (*typed cleanliness*) $[\Pi_{\bar{\Sigma}}^{[\mathrm{TT}]}, \Delta\hat{H}(\boldsymbol{\varepsilon})] = 0$ as well as BRST–cleanliness hold on $\mathfrak{P}$.

Step 1: Lipschitz control of interferometric propagators. Let $\hat{H} = \hat{H}_0$ denote the undeformed Stage II generator (on the relevant, typed subspace) and $\hat{H}[\boldsymbol{\varepsilon}] = \hat{H}_0 + \Delta\hat{H}(\boldsymbol{\varepsilon})$ the deformed one. For any fixed interrogation time $T > 0$,

$$\mathbf{U}(T;\boldsymbol{\varepsilon}) := e^{-\frac{i}{\hbar}T\hat{H}[\boldsymbol{\varepsilon}]}, \qquad \mathbf{U}_0(T) := e^{-\frac{i}{\hbar}T\hat{H}_0}.$$

Using the standard variation–of–constants identity $e^{X+Y} - e^X = \int_0^1 e^{(1-s)(X+Y)}\, Y\, e^{sX}\, ds$ with $X = -\frac{i}{\hbar}T\hat{H}_0$, $Y = -\frac{i}{\hbar}T\Delta\hat{H}(\boldsymbol{\varepsilon})$, we obtain the operator–norm bound

$$\|\mathbf{U}(T;\boldsymbol{\varepsilon}) - \mathbf{U}_0(T)\| \le \frac{T}{\hbar}\, e^{\frac{T}{\hbar}(\|\hat{H}_0\| + \|\Delta\hat{H}(\boldsymbol{\varepsilon})\|)}\, \|\Delta\hat{H}(\boldsymbol{\varepsilon})\| \le C_1(T,\mathfrak{P})\, \|\boldsymbol{\varepsilon}\|_\infty, \tag{13.233}$$

where the second inequality used (13.232) and absorbs the exponential factor into a diagnostic realized reference configuration–dependent constant $C_1(T,\mathfrak{P})$ for $\|\boldsymbol{\varepsilon}\|_\infty$ sufficiently small.

Step 2: Lipschitz stability of Stage II functionals. By construction, Stage II visibilities are expectation–value functionals built from bounded compositions of preparations, typed/controlled propagators, and readouts (Hadamard frames, controls, etc.), e.g. $\mathcal{V}_{\text{asym}}[H] = \mathrm{Tr}\left(M\, \mathbf{U}(T)\, \rho\, \mathbf{U}(T)^\dagger\right)$ for some diagnostic realized reference configuration–certified (ρ, M) (cf. Stage II definitions preceding eq. (13.66)). Hence

$$\begin{aligned}
&\left|\mathcal{V}_{\text{asym}}[H[\boldsymbol{\varepsilon}], H[\boldsymbol{\varepsilon}]] - \mathcal{V}_{\text{asym}}[H, H]\right| \\
&\quad \le \|M\| \left(\|\mathbf{U}(T;\boldsymbol{\varepsilon}) - \mathbf{U}_0(T)\|\, \|\rho\| + \|\rho\|\, \|\mathbf{U}(T;\boldsymbol{\varepsilon})^\dagger - \mathbf{U}_0(T)^\dagger\| \right) \\
&\quad \le 2\, \|M\|\, \|\rho\|\, \|\mathbf{U}(T;\boldsymbol{\varepsilon}) - \mathbf{U}_0(T)\| \;\le\; C_{\text{port}}\, \|\boldsymbol{\varepsilon}\|_\infty,
\end{aligned}$$

where $C_{\text{port}} = 2\, \|M\|\, \|\rho\|\, C_1(T,\mathfrak{P})$ and we used (13.233). This is precisely the Lipschitz visibility bound in eq. (13.66), with the higher–order $\mathcal{O}(\|\boldsymbol{\varepsilon}\|_\infty^2)$ term obtained by the second order of the Duhamel expansion.

Step 3: Lipschitz stability of the Stage II score. Let $\mathscr{C}(H)$ be the Stage II score given by an optimization over a compact, diagnostic realized reference configuration–certified instrument set $\mathfrak{A}_{II}$ (preparations/readouts/tunings), i.e. $\mathscr{C}(H) = \sup_{\alpha \in \mathfrak{A}_{II}} F_\alpha(H)$ with F_α a visibility–type functional as above. Then for every α, $|F_\alpha(H[\boldsymbol{\varepsilon}]) - F_\alpha(H)| \le C_{\text{port}}\, \|\boldsymbol{\varepsilon}\|_\infty + \mathcal{O}(\|\boldsymbol{\varepsilon}\|_\infty^2)$. Taking the supremum over α on both sides,

$$\left|\mathscr{C}(H[\boldsymbol{\varepsilon}]) - \mathscr{C}(H)\right| \le C'_{\text{port}}\, \|\boldsymbol{\varepsilon}\|_\infty + \mathcal{O}(\|\boldsymbol{\varepsilon}\|_\infty^2), \tag{13.234}$$

which is eq. (13.67). (We used the general inequality $|\sup f_\varepsilon - \sup f_0| \le \sup |f_\varepsilon - f_0|$ on compact domains.)

Step 4: Stability of the conditional evolution. By Proposition 13.162, for a PVM good–clock the subsystem conditioned vector obeys $i\hbar\,\partial_t\,|\Psi_{\rm sub}(t)\rangle = \hat H^{\rm eff}_{\rm sub}(t)\,|\Psi_{\rm sub}(t)\rangle$, with $\hat H^{\rm eff}_{\rm sub}(t) = \hat H_{\rm sub} + \langle t|\,\hat H_{\rm int}\,|t\rangle$. Under the admissible portal, the effective Hamiltonian becomes $\hat H^{\rm eff}_{\rm sub}(t;\boldsymbol\varepsilon) = \hat H^{\rm eff}_{\rm sub}(t) + \Delta\hat H^{\rm eff}_{\rm sub}(t;\boldsymbol\varepsilon)$, where, by the good–clock near–diagonality and typed cleanliness (Definition 13.158(v),(iii)),

$$\Delta\hat H^{\rm eff}_{\rm sub}(t;\boldsymbol\varepsilon) = \langle t|\;\Delta\hat H(\boldsymbol\varepsilon)\;|t\rangle\,, \qquad \|\Delta\hat H^{\rm eff}_{\rm sub}(t;\boldsymbol\varepsilon)\| \le K'_{\rm port}\,\|\boldsymbol\varepsilon\|_\infty.$$

Let $\mathbf{U}_{\rm sub}(t,s;\boldsymbol\varepsilon)$ denote the propagator generated by $\hat H^{\rm eff}_{\rm sub}(\cdot;\boldsymbol\varepsilon)$. The Duhamel formula and Grönwall's inequality then give

$$\|\mathbf{U}_{\rm sub}(t,s;\boldsymbol\varepsilon) - \mathbf{U}_{\rm sub}(t,s;0)\| \le C_2(t-s,\mathfrak{P})\,\|\boldsymbol\varepsilon\|_\infty,$$

so the conditioned evolution and all derived expectation values are Lipschitz–stable in $\|\boldsymbol\varepsilon\|_\infty$. This proves the "in particular" clause.

Combining Steps 1–4 establishes the claimed Lipschitz bounds for Stage II visibilities and scores, eqs. (13.66) and (13.67), and the stability of the conditional dynamics under admissible portals. □

Remark 13.165 (Chronogeometric placement)**.** A good–clock resides in the τ–kairos (non-TT) sector of the canonical symmetry stack and is compatible with the complete chronogeometric placement of Sections 13.4 and 13.4.1. This guarantees no extra tensor polarizations (cf. eqs. (13.38) and (13.179)).

Remark 13.166 (Operational darkness and selection context)**.** Good–clocks are used only to condition dynamics; they do not alter the *operational* dark–sector criterion, which is enforced by the darkness projector and SM spectral decoupling $\left(\mathcal{P}^{\rm coh}_G,\,\mathsf{P}_{\rm vis}\right)$ of Definition 13.1 and eq. (13.65).

Definition 13.167 (Total Relational Hamiltonian)**.** Let the total relational action be

$$S_{\rm total}[\hat{\mathbf\Phi},\hat{\mathbf\Theta}] := S_\Phi[\hat{\mathbf\Phi}] + S_\Theta[\hat{\mathbf\Theta}] + S_{\rm int}[\hat{\mathbf\Phi},\hat{\mathbf\Theta}] \qquad \text{(Definition 9.357)}.$$

The corresponding Legendre transform with respect to the *emergent proper time* (cf. Section 5.3.8) defines the operator

$$\boxed{\hat H_{\rm univ} \;:=\; \hat H_\Phi \;+\; \hat H_\Theta \;+\; \hat H_{\rm int},} \tag{13.235}$$

acting on the emergent/GNS Hilbert space $\mathcal{H}_{\rm em}$ (Section 2.4). Here $\hat H_\Phi$ (Gaiamiti/"geometric"), $\hat H_\Theta$ (Tantrakyma/"material"), and $\hat H_{\rm int}$ (nonlocal interaction) are each obtained from their Lagrangian densities by the same relational Legendre map.

Theorem 13.168 (Universal Quantum Constraint)**.** *Let $\left|\mathring{\mathbf\Psi}\right\rangle \in \mathcal{H}_{\rm phys}$ denote the BRST–physical universal Holo–State (cf. Definition 1.27). The fundamental quantum law of IQR is the operator constraint*

$$\boxed{\hat H_{\rm univ}\left|\mathring{\mathbf\Psi}\right\rangle = 0.} \tag{13.236}$$

Proof. **Step I (GNS, Holo–State, and BRST physicality).** By Definition 1.27, the universal Holo–State $\left|\mathring{\mathbf{\Psi}}\right\rangle$ is the cyclic vector of the GNS representation of the IQR observable C^*-algebra, and all observables are realized on the emergent Hilbert space $\mathcal{H}_{\rm em}$ [30, 381, 230]. Gauge fixing and constraints are implemented cohomologically: with BRST charge $\hat{Q}_{\rm BRST}$ ($\hat{Q}^2_{\rm BRST}=0$), the *physical* Hilbert space is

$$\mathcal{H}_{\rm phys} := H^0(\hat{Q}_{\rm BRST}) = \ker \hat{Q}_{\rm BRST} \,/\, \operatorname{im} \hat{Q}_{\rm BRST},$$

and we assume $\left|\mathring{\mathbf{\Psi}}\right\rangle \in \mathcal{H}_{\rm phys}$ as stated. BRST-exact operators have vanishing matrix elements between physical vectors (standard cohomological selection), a fact used repeatedly below (cf. Definition 6.270 and the BRST hygiene hypotheses across Chs. 6, 9–12) [442, 91].

Step II (Total relational Hamiltonian on the realized weave configuration). On a certified band/DRRC $\mathfrak{P}$ (UV projector $P_x(\lambda_{\rm UV})$, IR window $K^{(\ell)}_{\rm IR}$) with the realized weave configuration-compatible operator-bundle connection (Ch. 12), the total relational generator is the universal Hamiltonian

$$\hat{H}_{\rm univ} = \hat{H}_\Phi + \hat{H}_\Theta + \hat{H}_{\rm int} + \hat{H}_\partial,$$

where $\hat{H}_\partial$ collects the Brown–York–type boundary contributions consistent with the chrono-geometric vacuum choice (cf. Definitions 10.1, 13.92 and 13.155) [535, 532, 43, 7]. This construction is BRST/Nielsen-flat and Ward-covariant by the band-tameness and diagnostic realized reference configuration axioms of Ch. 6 (projected E–L, adjoint-covariant Ward identity) and Ch. 10 (commutations and TT-purity on the geometry leg) [120, 383, 91].

Step III (Adjoint-covariant Ward identity and vanishing total charge). Let $J^{(t)}_\mu$ be the Noether current associated to global time reparametrizations on the realized weave configuration. The adjoint-covariant Ward identity of Ch. 6 reads

$$\nabla_\mu J^\mu_{(t)} + i\,[\mathcal{A}_\mu, J^\mu_{(t)}] = \Delta_{(t)}, \qquad \text{with} \quad \Delta_{(t)} = \underbrace{\sum_k \epsilon_k(\ell)\,\mathcal{S}_{k,(t)}}_{\text{controlled deformations}} + \underbrace{\zeta\,\eta_{(t)}}_{\text{BRST-exact}} + \underbrace{\mathcal{A}_{(t)}}_{\text{anomaly}},$$

(cf. Definition 6.295 and eqs. (6.375) and (6.376)) [91, 120]. Integrating on a Cauchy domain $\mathcal{D}_\tau$ and using Stokes' theorem,

$$\frac{d}{d\tau}\,\mathbf{Q}_{(t)}(\tau) = \int_{\mathcal{D}_\tau} d^d x\,\sqrt{-\det \bar{\Sigma}}\,\Delta_{(t)} \quad \text{with} \quad \mathbf{Q}_{(t)}(\tau) := \int_{\mathcal{D}_\tau} d^{d-1}x\,\sqrt{\gamma}\,n_\mu J^\mu_{(t)}.$$

On the realized weave configuration: (i) all gauge-fixing pieces are BRST-exact ($\zeta\,\eta_{(t)} = \{\hat{Q}_{\rm BRST}, \cdot\}$), hence have vanishing matrix elements between physical states; (ii) Stage-I viability (Ch. 12) enforces vanishing anomaly meters, so $\mathcal{A}_{(t)} = 0$ on $\mathcal{H}_{\rm phys}$ (cf. Definitions 12.20 and 12.24 and Corollary 12.27) [442, 553]; (iii) boundary contributions are exactly those encoded in $\hat{H}_\partial$ and cancel against the calibrated Brown–York subtraction (Ch. 13; cf. Definition 13.155) [535, 532, 43]. Therefore

$$\left\langle \mathring{\mathbf{\Psi}} \right| \mathbf{Q}_{(t)}(\tau_2) - \mathbf{Q}_{(t)}(\tau_1) \left| \mathring{\mathbf{\Psi}} \right\rangle_{\rm phys} = 0 \quad \Rightarrow \quad \left\langle \mathring{\mathbf{\Psi}} \right| \mathbf{Q}_{(t)}(\tau) \left| \mathring{\mathbf{\Psi}} \right\rangle_{\rm phys} = \text{const.}$$

Choosing the universal (closed) boundary calibration of Ch. 13, this constant is fixed to zero. Hence the total Noether charge that generates global time translations *vanishes* on the universal Holo–State.

Step IV (From invariance to the operator constraint). Let $U(s)$ be the one-parameter unitary group implementing global time reparametrizations on the realized weave configuration representation; by Stone's theorem, $U(s) = \exp\!\left(-\frac{i}{\hbar} s\, \hat{H}_{\mathrm{univ}}\right)$ with self-adjoint generator $\hat{H}_{\mathrm{univ}}$ (a standard result; if desired, see e.g. *suggested* refs. on one-parameter unitary groups).[6] The vanishing of the total Noether charge on $\left|\mathring{\mathbf{\Psi}}\right\rangle$ is equivalent to

$$U(s)\left|\mathring{\mathbf{\Psi}}\right\rangle = \left|\mathring{\mathbf{\Psi}}\right\rangle \quad \forall s,$$

i.e. the universal Holo–State is invariant under the flow. Differentiating at $s = 0$ yields

$$-\tfrac{i}{\hbar}\,\hat{H}_{\mathrm{univ}}\left|\mathring{\mathbf{\Psi}}\right\rangle = 0 \quad \Longleftrightarrow \quad \hat{H}_{\mathrm{univ}}\left|\mathring{\mathbf{\Psi}}\right\rangle = 0,$$

which is exactly (13.236).

Step V (Reparametrization/BRST viewpoint). Equivalently, the universal action on the realized weave configuration is invariant under time reparametrizations; the associated primary first-class constraint is the Hamiltonian constraint. BRST quantization imposes first-class constraints as operator conditions on physical states. Since anomalies are null (Stage I) and boundary terms are calibrated (Ch. 13), the Hamiltonian constraint lifts without obstruction to the operator statement on $\mathcal{H}_{\mathrm{phys}}$, hence (13.236) [269, 442, 156, 605, 606, 3].

This completes the proof of Theorem 13.168. □

The starting point is the GNS realization of the IQR observable C^*-algebra. By Definition 1.27, the **Quantum Holo–State** $\left|\mathring{\mathbf{\Psi}}\right\rangle$ is the cyclic vector of the GNS representation on the emergent Hilbert space $\mathcal{H}_{\mathrm{em}}$ [30, 381, 230]. Gauge structure and constraints are imposed cohomologically via the BRST charge $\hat{Q}_{\mathrm{BRST}}$ with $\hat{Q}^2_{\mathrm{BRST}} = 0$, and the **physical** Hilbert space is the ghost–number–zero cohomology

$$\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}}) = \ker \hat{Q}_{\mathrm{BRST}} \,/\, \mathrm{im}\, \hat{Q}_{\mathrm{BRST}}\,,$$

in line with standard BRST treatments of first-class constraints [442, 91]. Throughout, the universal state is taken to satisfy $\left|\mathring{\mathbf{\Psi}}\right\rangle \in \mathcal{H}_{\mathrm{phys}}$. Standard cohomological selection then guarantees that BRST–exact insertions have vanishing matrix elements between physical vectors, a fact used repeatedly below (cf. the BRST hygiene assumptions and Definition 6.270) [442].

On band–certified plateaues $\mathfrak{P}$ (UV projector $P_x(\lambda_{\mathrm{UV}})$, IR window $K^{(\ell)}_{\mathrm{IR}}$) and with a realized weave configuration–compatible operator–bundle connection, the **total relational generator** of the emergent theory is the **universal Hamiltonian**

$$\hat{H}_{\mathrm{univ}} = \hat{H}_{\Phi} + \hat{H}_{\Theta} + \hat{H}_{\mathrm{int}} + \hat{H}_{\partial},$$

[6] If needed and not yet present in `citations.txt`, appropriate mathematical references include Reed & Simon, *Methods of Modern Mathematical Physics I*, or Stone's original papers; these can be added on request.

where $\hat{H}_\partial$ collects the Brown–York–type boundary contributions appropriate to the chronogeometric vacuum (cf. Definitions 10.1, 12.1, 13.92 and 13.155) [535, 532, 43, 7]. The scale–tagging and diagnostic realized reference configuration axioms of Definition 6.295 (projected Euler–Lagrange operators, adjoint–covariant Ward identity) together with the commutation/TT–purity results ensure that this construction is BRST/Nielsen–flat and Ward–covariant on $\mathcal{H}_{\text{phys}}$ (cf. Chs. 6 and 10) [120, 383, 91].

Let $J^\mu_{(t)}$ denote the Noether current associated to global time–reparametrizations on the realized weave configuration. The **adjoint–covariant Ward identity** reads

$$\nabla_\mu J^\mu_{(t)} + i\big[\mathcal{A}_\mu, J^\mu_{(t)}\big] = \Delta_{(t)}, \qquad \Delta_{(t)} = \sum_k \epsilon_k(\ell)\, \mathcal{S}_{k,(t)} \;+\; \zeta\, \eta_{(t)} \;+\; \mathcal{A}_{(t)},$$

with controlled deformations $\mathcal{S}_{k,(t)}$, a BRST–exact gauge–fixing part $\zeta\,\eta_{(t)}$, and a possible anomaly term $\mathcal{A}_{(t)}$ (see eqs. (6.375) and (6.376)) [120, 91]. Integrating over a Cauchy domain $\mathcal{D}_\tau$ and applying Stokes' theorem gives the continuity relation for the associated charge

$$\frac{d}{d\tau}\,\mathbf{Q}_{(t)}(\tau) = \int_{\mathcal{D}_\tau} \sqrt{-\det \bar{\Sigma}}\; \Delta_{(t)}\,, \qquad \mathbf{Q}_{(t)}(\tau) := \int_{\mathcal{D}_\tau} \sqrt{\gamma}\; n_\mu J^\mu_{(t)}.$$

On a certified plateau, each source term is controlled in the physical sector. First, the gauge–fixing contribution is BRST–exact, so its matrix elements vanish between physical vectors by construction [442]. Second, **Stage-I viability** of the internal sector (Ch. 12) enforces that the anomaly meters vanish; hence $\mathcal{A}_{(t)} = 0$ on $\mathcal{H}_{\text{phys}}$ (cf. Definitions 12.20 and 12.24 and Corollary 12.27) [553]. Third, boundary fluxes compatible with the chronogeometric vacuum are exactly those encoded in $\hat{H}_\partial$ and cancel by the Brown–York calibration in Definition 13.155 [535, 532, 43]. Consequently,

$$\Big\langle \overset{\circ}{\mathbf{\Psi}}\Big|\, \mathbf{Q}_{(t)}(\tau_2) - \mathbf{Q}_{(t)}(\tau_1)\, \Big|\overset{\circ}{\mathbf{\Psi}}\Big\rangle_{\text{phys}} = 0,$$

so $\langle \mathbf{Q}_{(t)}(\tau)\rangle$ is constant. With the universal (closed) boundary choice of Ch. 13, this constant is fixed to zero: the total time–translation charge of the closed universe **vanishes** on the Holo–State.

Let $U(s)$ be the unitary implementing global time–reparametrizations on the realized weave configuration representation. By Stone's theorem there exists a self–adjoint generator with $U(s) = \exp\!\big(-\tfrac{i}{\hbar} s\, \hat{H}_{\text{univ}}\big)$; differentiating the invariance condition $U(s)\big|\overset{\circ}{\mathbf{\Psi}}\big\rangle = \big|\overset{\circ}{\mathbf{\Psi}}\big\rangle$ at $s = 0$ yields $\hat{H}_{\text{univ}}\big|\overset{\circ}{\mathbf{\Psi}}\big\rangle = 0$ (cf. Theorem 13.168 and eq. (13.236)).[7]

An equivalent viewpoint emphasizes first–class constraints and BRST quantization: time–reparametrization invariance of the universal action implies a **Hamiltonian first–class constraint**, which BRST imposes as an operator condition on physical states [269, 442, 156, 605, 606, 3]. With anomalies null in Stage I and boundary terms calibrated in Ch. 13, the Hamiltonian constraint lifts without obstruction to an exact operator condition on $\mathcal{H}_{\text{phys}}$, again giving $\hat{H}_{\text{univ}}\big|\overset{\circ}{\mathbf{\Psi}}\big\rangle = 0$.

Finally, the constraint is a statement about the **closed whole** and does not preclude dynamical evolution of subsystems. As detailed in Section 13.6.2, once a suitable internal

[7] If a mathematical citation for Stone's theorem is desired and not yet in `citations.txt`, *suggested* additions include Reed & Simon, *Methods of Modern Mathematical Physics I*, or Stone's original papers.

relational clock is identified within the state (cf. Section 5.3.8), conditioning the timeless universal state on clock readings recovers subsystem Schrödinger evolution [215, 3, 607, 605, 606] (see also eq. (13.238)). In this way, emergent time for parts is reconciled with timelessness of the whole, while the universal constraint (13.236) remains exact on $\mathcal{H}_{\text{phys}}$.

Remarks and Comparison. (i) Equation (13.236) is the background–independent generalization of the Wheeler–DeWitt equation [156], formulated as a *true operator equation* on $\mathcal{H}_{\text{phys}}$ rather than as a functional differential equation on a space of 3–geometries. Geometry and matter are not split as "geometry with sources''; both arise from a single constrained state under $\hat{H}_{\text{univ}}$. (ii) Dirac's prescription for first–class constraints is implemented at the operator level [269, 442]; the resolution of gauge and BRST cohomology produces $\mathcal{H}_{\text{phys}}$ on which (13.236) is imposed. (iii) The absence of an external time is reconciled with observable dynamics via *relational evolution*: conditional/partial observables acquire time by correlation either with internal clocks or with modular/KMS flows of relevant algebras [83, 608, 230, 241].

The symmetry architecture developed in Section 13.4—and, in particular, its canonical placement in Section 13.4.1 and summarized layering in Section 13.4.1—is a **cosmic** hierarchy of commuting symmetry lanes built on the same UV/IR–tagged, BRST–clean realized weave configuration used throughout the treatise. The stack organizes (i) the chronogeometric lanes (radiative spin-2 in the TT sector versus the clock-capable τ-kairos sector), (ii) the internal/-gauge content spanning both the visible (SM) and dark sectors, and (iii) the holosymmetric control layer that enforces the Holosymmetric Selection Principle (HSP). Each lane is realized by certified projectors that commute with the band tags and with the York–Lichnerowicz TT projector, and each lane carries its law/state, parasymmetric, and holosymmetric meters, all Nielsen–flat on certified plateaux (cf. Definition 10.1 and eq. (10.4); see also [120, 383, 91] for Ward/BRST structure and [532, 43, 44, 45] for the TT/York–Lichnerowicz formalism).

Crucially, the stack is not "dark-only": it **canonically layers gravity, the visible (SM) sector, and the dark sector** within a single commuting hierarchy. The visible lane is the Chapter 12 operator-bundle realization, equipped with the realized weave configuration-compatible connection $\nabla^{\text{realizedweaveconfiguration}}$ and the effective internal fibre (cf. Definitions 12.1 and 12.6). Commuting triad projectors and typed/Riesz selectors localize visible excitations band-wise (Definitions 10.2 and 10.7); color singletness is enforced by the singlet projector (Definition 12.126), ensuring that hadronic observables descend BRST–physically (Section 12.6). Stage-I anomaly meters vanish and the **unitarity operator** acts as identity on certified support (Definitions 12.20, 12.24 and 12.28), making the visible lane anomaly-free and Nielsen–flat—properties that enter directly into the Ward analysis used for the universal constraint (cf. [91, 82, 443, 553]).

The **gravity lane** supplies the TT radiative chronogeometry via the York projector $\Pi_{\bar{\Sigma}}^{[\text{TT}]}$ (cf. eq. (10.4)). Its commutation with the typed/triad projectors ensures TT-purity lane-wise and governs the canonical placement and calibration of quasilocal boundary terms. This TT-purity guarantees that clock-carrying τ-kairos channels (below) neither contaminate spin-2 dynamics nor upset the Brown–York boundary accounting used in the mass/energy definitions of Definition 13.155 (see [532, 43, 535, 7, 45]).

Within this stacked setting, the **construction** of the universal Hamiltonian becomes

lane-resolved and canonically placed. The sectoral decomposition

$$\hat{H}_{\text{univ}} = \hat{H}_{\Phi} + \hat{H}_{\Theta} + \hat{H}_{\text{int}} + \hat{H}_{\partial}$$

is sharpened by identifying which parts of $\hat{H}_{\Phi}$ and $\hat{H}_{\Theta}$ populate the gravity, visible (SM), and dark lanes, and by fixing their boundary counterparts $\hat{H}_{\partial}$ according to the chronogeometric prescription of Section 13.4.1. The result is a sum of commuting, BRST–clean blocks plus calibrated boundary terms, so that Brown–York calibrations apply **lane-wise** and add consistently at the universal level (cf. Definition 13.155; [535, 532, 43]).

The stack also clarifies the **Noether/Ward** analysis. The adjoint-covariant Ward identity for the time-reparametrization current $J^{\mu}_{(t)}$ splits along the lanes, so the source $\Delta_{(t)}$ decomposes into controlled deformations, BRST-exact gauge-fixing pieces, and potential anomalies **lane-by-lane** (cf. eqs. (6.375) and (6.376)). Because every lane is BRST-closed on the realized weave configuration and HSP-viable, the anomaly meters vanish in each block, and the gauge-fixing contributions remain BRST-exact there as well. After integration over a Cauchy domain and inclusion of the calibrated boundary fluxes, the total time-translation charge is constant and—under the universal closed calibration of Definition 13.155—fixed to zero. This lane-wise factorization tightens the invariance step in the proof and prepares the final deduction of the constraint on $\mathcal{H}_{\text{phys}}$ (cf. [120, 91, 443]).

A further consequence of the canonical placement in Section 13.4.1 is the **TT-nullity** and **commutation** structure that keeps the τ-kairos (clock-capable) lanes cleanly separated from radiative spin-2. With clock candidates canonically placed and TT-safe, no back-reaction arises that could generate an explicit time-reparametrization–breaking term in $J^{\mu}_{(t)}$. Hence the one-parameter unitary $U(s) = \exp\!\big(-\tfrac{i}{\hbar} s\, \hat{H}_{\text{univ}}\big)$ leaves the universal state invariant, and differentiation at $s = 0$ gives the operator statement $\hat{H}_{\text{univ}} \big|\mathring{\mathbf{\Psi}}\big\rangle = 0$ of Theorem 13.168 (for Stone's theorem and one-parameter unitary groups, see [253]).

The **summary layer** Section 13.4.1 codifies **admissible deformations** (e.g., small retunings of masses/couplings or history-dependent kernels) as symmetry-controlled operators with explicit band tags. Within the Ward framework these enter only through the controlled-deformation slot of $\Delta_{(t)}$ and are therefore governed by the same diagnostic realized reference configuration criteria that ensure $\frac{d}{d\tau}\langle \mathbf{Q}_{(t)} \rangle = 0$. In particular, any deformation allowed by the stack preserves BRST/Nielsen flatness and does not spoil the constancy—and universal calibration to zero—of the time-translation charge in the closed configuration ([120, 91]).

Operationally, the stack **fixes the clock architecture** for emergent dynamics. Because the τ-kairos lanes are canonically embedded and BRST-closed, an internal clock C used for relational conditioning (cf. Section 5.3.8) automatically commutes with the TT projector and respects the SM projectors. Visible clocks (e.g., atomic/baryonic standards) are equally admissible provided they occupy a lane that commutes with $\Pi^{[\mathrm{TT}]}_{\bar{\Sigma}}$ and the SM structure. Consequently, Page–Wootters conditioning proceeds without violating the universal constraint, and the same meters remain Nielsen–flat; the emergent Schrödinger dynamics is thus compatible **by construction** with (13.236) (cf. Section 13.6.2; see also [215, 3, 146, 7]).

A concrete manifestation of the visible lane within this architecture appears in cosmological diagnostics: the linear-power template combines the **visible-sector transfer** $T_{\text{vis}}(k)$ (set by the SM lane) with dark-lane coherence kernels, an immediate consequence of the joint

geometry+SM placement provided by the stack. While such diagnostics are not needed to **state** the universal constraint, they demonstrate that the visible lane is structurally present in the same hierarchy that underwrites the constraint's proof—through anomaly-free Ward behavior, calibrated boundary accounting, and compatibility with the clock sector (cf. [327, 144, 136]).

In sum, Section 13.4 does not alter the content of the Universal Quantum Constraint; it **tightens** its hypotheses and **modularizes** its proof. The symmetry stack (i) provides a canonical, diagnostic realized reference configuration-compatible sectoral decomposition of $\hat{H}_{\mathrm{univ}}$ and its boundary terms; (ii) enforces BRST/Nielsen hygiene and anomaly-freedom lane-wise via HSP; (iii) secures the TT-nullity and commutation properties required for the Ward argument; and (iv) supplies a canonical, TT-safe clock sector for the relational emergence of time. Under these conditions, the invariance of the universal Holo–State under global time reparametrizations follows transparently, and the operator statement $\hat{H}_{\mathrm{univ}} \left|\overset{\circ}{\boldsymbol{\Psi}}\right\rangle = 0$ of Theorem 13.168 remains exact on $\mathcal{H}_{\mathrm{phys}}$.

Proposition 13.169 (Spectral Resolution and Energy–Partition)**.** *Let $E \in \sigma(\hat{H}_{\mathrm{univ}})$ and $\mu_{\overset{\circ}{\boldsymbol{\Psi}}}^{\hat{H}_{\mathrm{univ}}}$ be the spectral measure induced by $\left|\overset{\circ}{\boldsymbol{\Psi}}\right\rangle$. Then the comoving total energy density is*

$$\rho_{\mathrm{tot}} = \frac{1}{\mathrm{Vol}(\mathcal{X}_q)} \int_{\sigma(\hat{H}_{\mathrm{univ}})} E \, \mathrm{d}\mu_{\overset{\circ}{\boldsymbol{\Psi}}}^{\hat{H}_{\mathrm{univ}}}(E), \qquad (\textit{eq. (13.144)}),$$

and, when the holosymmetric projectors resolve the identity on $\mathcal{H}_{\mathrm{phys}}$ (cf. Proposition 13.97), the partition $\rho_{\mathrm{tot}} = \rho_{\mathrm{DE}} + \rho_{\mathrm{DM}} + \rho_{\mathrm{vis}}$ holds with abundances $\Omega_i = \rho_i/\rho_{\mathrm{tot}}$ (Section 13.4.7).

Proof. **(1) Spectral resolution and the total comoving energy density.** Since $\hat{H}_{\mathrm{univ}}$ is self-adjoint on the BRST–physical realized weave configuration (cf. the construction of the universal generator and boundary calibration in Ch. 13 and Definition 13.92), the spectral theorem yields a projection-valued measure $\Pi^{\hat{H}_{\mathrm{univ}}}$ on $\sigma(\hat{H}_{\mathrm{univ}})$ such that

$$\hat{H}_{\mathrm{univ}} = \int_{\sigma(\hat{H}_{\mathrm{univ}})} E \, \mathrm{d}\Pi^{\hat{H}_{\mathrm{univ}}}(E).$$

For the universal Holo–State $\left|\overset{\circ}{\boldsymbol{\Psi}}\right\rangle \in \mathcal{H}_{\mathrm{phys}}$, define the spectral measure $\mu_{\overset{\circ}{\boldsymbol{\Psi}}}^{\hat{H}_{\mathrm{univ}}}(B) := \left\langle \overset{\circ}{\boldsymbol{\Psi}} \right| \Pi^{\hat{H}_{\mathrm{univ}}}(B) \left| \overset{\circ}{\boldsymbol{\Psi}} \right\rangle$ for Borel sets $B \subset \mathbb{R}$. Then

$$\left\langle \overset{\circ}{\boldsymbol{\Psi}} \right| \hat{H}_{\mathrm{univ}} \left| \overset{\circ}{\boldsymbol{\Psi}} \right\rangle = \int_{\sigma(\hat{H}_{\mathrm{univ}})} E \, \mathrm{d}\mu_{\overset{\circ}{\boldsymbol{\Psi}}}^{\hat{H}_{\mathrm{univ}}}(E).$$

By the definition of the comoving total energy density on the emergent diagnostic realized reference configuration (cf. eq. (13.144) and Definition 13.92),

$$\rho_{\mathrm{tot}} = \frac{1}{\mathrm{Vol}(\mathcal{X}_q)} \left\langle \overset{\circ}{\boldsymbol{\Psi}} \right| \hat{H}_{\mathrm{univ}} \left| \overset{\circ}{\boldsymbol{\Psi}} \right\rangle = \frac{1}{\mathrm{Vol}(\mathcal{X}_q)} \int_{\sigma(\hat{H}_{\mathrm{univ}})} E \, \mathrm{d}\mu_{\overset{\circ}{\boldsymbol{\Psi}}}^{\hat{H}_{\mathrm{univ}}}(E).$$

(2) Holosymmetric resolution of the identity and energy partition. Let $\{P_i\}_{i\in\{\mathrm{DE,DM,vis}\}}$ be the holosymmetric projectors (HSP-lane projectors) supplied by the cosmic symmetry stack

(Sections 13.4 and 13.4.1). On the physical space they are mutually orthogonal, self-adjoint, and resolve the identity,

$$P_i P_j = \delta_{ij} P_i, \qquad P_i^\dagger = P_i, \qquad \sum_i P_i = \mathbf{1} \ \text{ on } \ \mathcal{H}_{\text{phys}} \quad \text{(cf. Proposition 13.97)}.$$

The stack is constructed to commute with the band tags and with the York projector; in particular the lane projectors commute with the universal dynamics on certified plateaues, so

$$[P_i, \hat{H}_{\text{univ}}] = 0 \quad \text{on } \mathcal{H}_{\text{phys}} \qquad \text{(cf. lane-wise commuting structure in Section 13.4)}.$$

Consequently, the spectral measure decomposes lane-wise:

$$\mu_{\mathring{\Psi}}^{\hat{H}_{\text{univ}}} = \sum_i \mu_{i,\mathring{\Psi}}^{\hat{H}_{\text{univ}}}, \qquad \mu_{i,\mathring{\Psi}}^{\hat{H}_{\text{univ}}}(B) := \Big\langle \mathring{\Psi} \Big| \, P_i \, \Pi^{\hat{H}_{\text{univ}}}(B) \, P_i \, \Big| \mathring{\Psi} \Big\rangle = \Big\langle \mathring{\Psi} \Big| \, \Pi^{\hat{H}_{\text{univ}}}(B) \, P_i \, \Big| \mathring{\Psi} \Big\rangle,$$

where the last equality uses $[P_i, \Pi^{\hat{H}_{\text{univ}}}(B)] = 0$. Define the lane (sector) energy densities by either of the equivalent forms

$$\rho_i := \frac{1}{\text{Vol}(\mathcal{X}_q)} \Big\langle \mathring{\Psi} \Big| \, \hat{H}_{\text{univ}} \, P_i \, \Big| \mathring{\Psi} \Big\rangle = \frac{1}{\text{Vol}(\mathcal{X}_q)} \int_{\sigma(\hat{H}_{\text{univ}})} E \, \mathrm{d}\mu_{i,\mathring{\Psi}}^{\hat{H}_{\text{univ}}}(E),$$

which are nonnegative by positivity of the PVM and orthogonality of the lanes. Using $\sum_i P_i = \mathbf{1}$ and $[P_i, \hat{H}_{\text{univ}}] = 0$,

$$\rho_{\text{tot}} = \frac{1}{\text{Vol}(\mathcal{X}_q)} \Big\langle \mathring{\Psi} \Big| \, \hat{H}_{\text{univ}} \sum_i P_i \, \Big| \mathring{\Psi} \Big\rangle = \sum_i \frac{1}{\text{Vol}(\mathcal{X}_q)} \Big\langle \mathring{\Psi} \Big| \, \hat{H}_{\text{univ}} \, P_i \, \Big| \mathring{\Psi} \Big\rangle = \sum_i \rho_i.$$

Hence the holosymmetric partition $\rho_{\text{tot}} = \rho_{\text{DE}} + \rho_{\text{DM}} + \rho_{\text{vis}}$ holds on $\mathcal{H}_{\text{phys}}$. The associated *abundances* are defined by $\Omega_i := \rho_i / \rho_{\text{tot}}$, as introduced in Section 13.4.7.

The stated formula for ρ_{tot} and the holosymmetric energy–partition therefore follow from the spectral calculus of $\hat{H}_{\text{univ}}$ together with the lane-wise resolution of the identity furnished by the cosmic symmetry stack (HSP), completing the proof. □

Proposition 13.170 (Relational Schrödinger Equations from the Constraint)**.** *Let $\hat{C}$ be a (self–adjoint) clock operator with nondegenerate spectrum and $\mathcal{A}$ an algebra of relational observables commuting with $\hat{C}$ on $\mathcal{H}_{\text{phys}}$. If $\hat{H}_{\text{univ}} \big| \mathring{\Psi} \big\rangle = 0$, then the conditional state $\varrho_{A|C=t} := \frac{\Pi_t \, \varrho \, \Pi_t}{\text{Tr}(\Pi_t \, \varrho)}$ with $\varrho = \big| \mathring{\Psi} \big\rangle \big\langle \mathring{\Psi} \big|$ and Π_t the spectral projector of $\hat{C}$ at value t, satisfies an effective Schrödinger equation*

$$i\hbar \, \partial_t \, \varrho_{A|C=t} = \Big[\, \hat{H}_{\text{eff}}(t), \; \varrho_{A|C=t} \, \Big],$$

for some $\hat{H}_{\text{eff}}(t)$ determined by the relational splitting ("time from correlations"). An analogous evolution holds along modular/KMS parameters for von Neumann algebras associated with stationary horizons and cosmological patches [230, 241, 608].

Proof. **(1) Conditionalization on a good clock.** By Definition 1.27, the GNS construction realizes the IQR observable C^*-algebra on the emergent Hilbert space $\mathcal{H}_{\mathrm{em}}$ with cyclic vector $\left|\mathring{\mathbf{\Psi}}\right\rangle$ [381, 241]. Cohomological gauge handling uses a nilpotent BRST charge $\hat{Q}_{\mathrm{BRST}}$ with $\hat{Q}_{\mathrm{BRST}}^2 = 0$, so the *physical* Hilbert space is

$$\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}}) = \ker \hat{Q}_{\mathrm{BRST}} \,/\, \mathrm{im}\, \hat{Q}_{\mathrm{BRST}},$$

and we assume $\left|\mathring{\mathbf{\Psi}}\right\rangle \in \mathcal{H}_{\mathrm{phys}}$ (cf. standard constraint quantization surveys [609, 610]). BRST-exact insertions then have vanishing matrix elements between physical states, as used below (cf. BRST hygiene and Definition 6.270).

(2) Total relational Hamiltonian on the realized weave configuration. On a band/DRRC $\mathfrak{P}$ (UV projector $P_x(\lambda_{\mathrm{UV}})$, IR window $K_{\mathrm{IR}}^{(\ell)}$) with a realized weave configuration-compatible operator-bundle connection, the total generator is

$$\hat{H}_{\mathrm{univ}} = \hat{H}_\Phi + \hat{H}_\Theta + \hat{H}_{\mathrm{int}} + \hat{H}_\partial,$$

with $\hat{H}_\partial$ collecting Brown–York–type boundary terms consistent with the chronogeometric vacuum (cf. Definitions 10.1, 13.92 and 13.155). Scale-tagging and diagnostic realized reference configuration axioms (projected E–L and adjoint-covariant Ward identity) make this construction BRST/Nielsen-flat and Ward-covariant (Chs. 6, 10). For background GR/QFT structures and boundary accounting see [7, 11].

(3) Adjoint-covariant Ward identity and vanishing total charge. Let $J^\mu_{(t)}$ be the Noether current for global time reparametrizations. The adjoint-covariant Ward identity reads

$$\nabla_\mu J^\mu_{(t)} + i\,[\mathcal{A}_\mu, J^\mu_{(t)}] = \Delta_{(t)} = \sum_k \epsilon_k(\ell)\, \mathcal{S}_{k,(t)} \;+\; \zeta\, \eta_{(t)} \;+\; \mathcal{A}_{(t)},$$

(cf. eqs. (6.375) and (6.376)). Integrating on a Cauchy domain $\mathcal{D}_\tau$ and applying Stokes' theorem gives

$$\frac{d}{d\tau}\, \mathbf{Q}_{(t)}(\tau) = \int_{\mathcal{D}_\tau} \sqrt{-\det \bar{\Sigma}}\, \Delta_{(t)}, \qquad \mathbf{Q}_{(t)}(\tau) := \int_{\mathcal{D}_\tau} \sqrt{\gamma}\, n_\mu J^\mu_{(t)}.$$

On the realized weave configuration, (i) gauge-fixing pieces are BRST-exact and vanish between physical states; (ii) Stage-I viability forces anomaly meters to zero, $\mathcal{A}_{(t)} = 0$; and (iii) boundary fluxes cancel by the Brown–York calibration in Definition 13.155. Hence $\langle \mathbf{Q}_{(t)}(\tau) \rangle$ is constant, and with the universal (closed) calibration it is fixed to zero. For Noether/Ward background, see [301, 91].

(4) From invariance to the operator constraint. Let $U(s)$ implement global time reparametrizations; by Stone's theorem there is a self-adjoint generator with $U(s) = \exp\!\big(-\tfrac{i}{\hbar} s\, \hat{H}_{\mathrm{univ}}\big)$ [369]. Vanishing total Noether charge on $\left|\mathring{\mathbf{\Psi}}\right\rangle$ is equivalent to

$$U(s) \left|\mathring{\mathbf{\Psi}}\right\rangle = \left|\mathring{\mathbf{\Psi}}\right\rangle \quad \forall s,$$

and differentiating at $s = 0$ yields

$$\hat{H}_{\mathrm{univ}} \left|\mathring{\mathbf{\Psi}}\right\rangle = 0,$$

i.e. (13.236).

(5) Reparametrization/BRST viewpoint. Time-reparametrization invariance entails a first-class Hamiltonian constraint; BRST quantization imposes first-class constraints as operator conditions on physical states (anomalies null by Stage I, boundaries calibrated as above), giving again $\hat{H}_{\text{univ}} \left|\mathring{\mathbf{\Psi}}\right\rangle = 0$ [609, 610].

(6) Conditional Schrödinger dynamics for subsystems. With a good internal clock and the Page–Wootters conditioning, subsystem dynamics emerges unitarily: the relational Schrödinger equation follows for the conditional state, consistently with the universal constraint [215, 610]. (For modular/KMS analogues in stationary settings, see [241, 11].) □

Remark 13.171 (Consistency with Horizon Thermodynamics and Cosmology)**.** Equation (13.236) underwrites the KMS/modular structure used at horizons and in cosmology. For stationary black–hole exteriors, the algebraic state is KMS with respect to the horizon generator χ^a, defining an effective temperature $T_{\text{eff}} = \hbar\kappa_{\text{eff}}/(2\pi k_B)$ and rotational chemical potential $\tilde{\omega} = \omega - m\Omega_H$ (Sections 16.7 and 16.9). In homogeneous FRW patches, the same constraint and the holosymmetric selection fix the dark–sector background and perturbations through the temporal tensor $t_{\mu\nu}$ (Section 13.3), yielding the abundance relations of Section 13.4.7.

Operational Meaning. The Universal Quantum Constraint (13.236) states that the total relational energy of the universe vanishes: the realized cosmos is a *stationary, entangled* Holo–State encoding all correlations among geometric ($\hat{\mathbf{\Phi}}$), material ($\hat{\mathbf{\Theta}}$), and interaction degrees of freedom. "Time" appears only as a relational parameter attached to chosen internal clocks or modular flows of relevant algebras, never as an external background. This single operator law thus unifies geometry and matter at the quantum level and provides the backbone for: (i) the black–hole lifecycle and horizon thermodynamics (Sections 16.7, 16.8 and 16.11); (ii) the dark–sector symmetry stack and cosmological abundances (Sections 13.4 and 13.4.7); and (iii) the recovery of classical dynamics in decohered regimes (Section 13.3), all without introducing an external notion of time.

13.6.2 The Emergence of Subsystem Schrödinger Dynamics

Proposition 13.172 (Conditional Schrödinger Dynamics from the Universal Quantum Constraint)**.** *Assume a BRST–physical factorization* $\mathcal{H}_{\text{phys}} = \mathcal{H}_{\text{sub}} \otimes \mathcal{H}_{\text{clock}}$ *and a self–adjoint clock pair* $(\hat{t}, \hat{p}_t)$ *on* $\mathcal{H}_{\text{clock}}$ *with* $[\hat{t}, \hat{p}_t] = i\hbar$*. Suppose the Total Relational Hamiltonian admits the deparametrized form*

$$\hat{H}_{\text{univ}} = \hat{p}_t \otimes \mathbf{1}_{\text{sub}} + \mathbf{1}_{\text{clock}} \otimes \hat{H}_{\text{sub}} + \hat{H}_{\text{int}}, \qquad \left|\mathring{\mathbf{\Psi}}\right\rangle \in \mathcal{H}_{\text{phys}} \text{ with } \hat{H}_{\text{univ}} \left|\mathring{\mathbf{\Psi}}\right\rangle = 0, \tag{13.237}$$

where either (a) $[\hat{H}_{\text{int}}, \hat{t} \otimes \mathbf{1}_{\text{sub}}] = 0$ *or (b)* $\hat{H}_{\text{int}}$ *is negligible in the clock sector (good–clock approximation). Let* $\{|t\rangle\}$ *be the (generalized) spectral family of* $\hat{t}$*, with projectors* $\Pi_t := |t\rangle\langle t|$*. Define the conditional (unnormalized) subsystem state* $|\widetilde{\Psi}_{\text{sub}}(t)\rangle := \langle t| \mathring{\mathbf{\Psi}}\rangle \in \mathcal{H}_{\text{sub}}$ *and its normalized version* $|\Psi_{\text{sub}}(t)\rangle := |\widetilde{\Psi}_{\text{sub}}(t)\rangle / \| |\widetilde{\Psi}_{\text{sub}}(t)\rangle \|$*. Then, distributionally,*

$$\boxed{i\hbar\, \partial_t \, |\Psi_{\text{sub}}(t)\rangle = \hat{H}^{\text{eff}}_{\text{sub}}(t) \; |\Psi_{\text{sub}}(t)\rangle \,, \qquad \hat{H}^{\text{eff}}_{\text{sub}}(t) := \hat{H}_{\text{sub}} + \langle t| \, \hat{H}_{\text{int}} \, |t\rangle \,.} \tag{13.238}$$

Idea. Begin from the Universal Quantum Constraint $\hat{H}_{\text{univ}}\big|\mathring{\mathbf{\Psi}}\big\rangle = 0$ (cf. Theorem 13.168, (13.236)). In the deparametrized split (13.237) and the time basis of the clock, act with $\langle t|$ and use $\langle t|\,\hat{p}_t = -i\hbar\,\partial_t\,\langle t|$ to obtain the fiberwise (unnormalized) evolution

$$i\hbar\,\partial_t\,|\widetilde{\Psi}_{\text{sub}}(t)\rangle = \Big(\hat{H}_{\text{sub}} + \langle t|\,\hat{H}_{\text{int}}\,|t\rangle\Big)\,|\widetilde{\Psi}_{\text{sub}}(t)\rangle\,.$$

Assumption (a) $[\hat{H}_{\text{int}}, \hat{t}\otimes\mathbf{1}_{\text{sub}}] = 0$ or the good–clock approximation (b) (cf. Section 5.3.8) guarantees that no off–diagonal clock couplings generate extra derivative terms. Dividing by the nonzero norm and applying the quotient rule isolates a purely real "phase–connection'' term, which can be absorbed by a time–dependent phase choice (gauge) for the normalized state $|\Psi_{\text{sub}}(t)\rangle$. The result is the conditional Schrödinger equation

$$i\hbar\,\partial_t\,|\Psi_{\text{sub}}(t)\rangle = \hat{H}^{\text{eff}}_{\text{sub}}(t)\,|\Psi_{\text{sub}}(t)\rangle\,, \qquad \hat{H}^{\text{eff}}_{\text{sub}}(t) := \hat{H}_{\text{sub}} + \langle t|\,\hat{H}_{\text{int}}\,|t\rangle\,,$$

which is precisely (13.238).

Proof. **(1) Setup and clock spectral resolution.** Assume the BRST–physical factorization $\mathcal{H}_{\text{phys}} = \mathcal{H}_{\text{sub}}\otimes\mathcal{H}_{\text{clock}}$ (standard for first–class constraints and BRST cohomology [442]) and a self–adjoint canonical clock pair $(\hat{t}, \hat{p}_t)$ on $\mathcal{H}_{\text{clock}}$ with $[\hat{t}, \hat{p}_t] = i\hbar$ (spectral theorem and canonical pairs in functional analysis [369]). Let $\{|t\rangle\}_{t\in\mathbb{R}}$ be the (generalized) spectral family of $\hat{t}$, with projectors $\Pi_t := |t\rangle\langle t|$ (distributional spectral resolution [369]). For the universal Holo–State $|\mathring{\mathbf{\Psi}}\rangle \in \mathcal{H}_{\text{phys}}$ obeying the *Universal Quantum Constraint* $\hat{H}_{\text{univ}}\,|\mathring{\mathbf{\Psi}}\rangle = 0$ (cf. Theorem 13.168, (13.236); see also [156, 609, 610]), define the (unnormalized) conditional subsystem state $|\widetilde{\Psi}_{\text{sub}}(t)\rangle := \langle t|\,\mathring{\mathbf{\Psi}}\rangle \in \mathcal{H}_{\text{sub}}$, and its normalized version $|\Psi_{\text{sub}}(t)\rangle := |\widetilde{\Psi}_{\text{sub}}(t)\rangle\,/\|\,|\widetilde{\Psi}_{\text{sub}}(t)\rangle\,\|$. All equalities below are understood distributionally on the rigged–Hilbert–space extension of $\mathcal{H}_{\text{clock}}$ (cf. [369]).

(2) Projecting the constraint in the time basis. Using the deparametrized split (13.237),

$$0 = \langle t|\,\hat{H}_{\text{univ}}\,|\mathring{\mathbf{\Psi}}\rangle = \underbrace{\langle t|\,\hat{p}_t\,|\mathring{\mathbf{\Psi}}\rangle}_{(i)} + \underbrace{\langle t|\,\mathbf{1}_{\text{clock}}\otimes\hat{H}_{\text{sub}}\,|\mathring{\mathbf{\Psi}}\rangle}_{(ii)} + \underbrace{\langle t|\,\hat{H}_{\text{int}}\,|\mathring{\mathbf{\Psi}}\rangle}_{(iii)}.$$

By the standard distributional action of $\hat{p}_t$ in the t-representation,

$$(i) \;=\; \langle t|\,\hat{p}_t\,|\mathring{\mathbf{\Psi}}\rangle = -\,i\hbar\,\partial_t\,|\widetilde{\Psi}_{\text{sub}}(t)\rangle \qquad [369].$$

Term (ii) gives $\langle t|\,(\mathbf{1}_{\text{clock}}\otimes\hat{H}_{\text{sub}})\,|\mathring{\mathbf{\Psi}}\rangle \;=\; \hat{H}_{\text{sub}}\,|\widetilde{\Psi}_{\text{sub}}(t)\rangle$. For (iii), assume either (a) $[\hat{H}_{\text{int}}, \hat{t}\otimes\mathbf{1}_{\text{sub}}] = 0$ (block-diagonality in the clock basis), which yields the fiber operator $\hat{V}(t) := \langle t|\,\hat{H}_{\text{int}}\,|t\rangle$ on $\mathcal{H}_{\text{sub}}$ and hence $\langle t|\,\hat{H}_{\text{int}}\,|\mathring{\mathbf{\Psi}}\rangle = \hat{V}(t)\,|\widetilde{\Psi}_{\text{sub}}(t)\rangle$; or (b) the good–clock approximation of Section 5.3.8, in which off-diagonal clock couplings are negligible and the same replacement holds up to errors controlled by clock quality (see the Page–Wootters/relational time literature [215, 3, 607]). Collecting terms,

$$i\hbar\,\partial_t\,|\widetilde{\Psi}_{\text{sub}}(t)\rangle = \Big(\hat{H}_{\text{sub}} + \hat{V}(t)\Big)\,|\widetilde{\Psi}_{\text{sub}}(t)\rangle \qquad \text{with} \quad \hat{V}(t) = \langle t|\,\hat{H}_{\text{int}}\,|t\rangle\,. \qquad (*)$$

(3) Normalization and Hermiticity. Since $\hat{H}_{\rm sub}$ and $\hat{H}_{\rm int}$ are self–adjoint, the fiber $\hat{V}(t)$ is self–adjoint for each t, hence $\hat{H}^{\rm eff}_{\rm sub}(t) := \hat{H}_{\rm sub} + \hat{V}(t)$ is self–adjoint (standard operator properties [369]). Taking the t-derivative of the norm and using $(*)$,

$$\begin{aligned}\partial_t \| \, |\widetilde{\Psi}_{\rm sub}(t)\rangle \, \|^2 &= \partial_t \, \langle \widetilde{\Psi}_{\rm sub}(t) | \widetilde{\Psi}_{\rm sub}(t) \rangle \\ &= \frac{i}{\hbar} \, \langle \widetilde{\Psi}_{\rm sub}(t) | \hat{H}^{\rm eff}_{\rm sub}(t) | \widetilde{\Psi}_{\rm sub}(t) \rangle - \frac{i}{\hbar} \, \langle \widetilde{\Psi}_{\rm sub}(t) | \hat{H}^{\rm eff}_{\rm sub}(t) | \widetilde{\Psi}_{\rm sub}(t) \rangle \\ &= 0.\end{aligned}$$

Thus $\| \, |\widetilde{\Psi}_{\rm sub}(t)\rangle \, \|$ is constant in t, and the normalized state obeys the *same* evolution:

$$i\hbar \, \partial_t \, |\Psi_{\rm sub}(t)\rangle = \hat{H}^{\rm eff}_{\rm sub}(t) \, |\Psi_{\rm sub}(t)\rangle \, .$$

(4) Conclusion and relation to the relational picture. Defining $\hat{H}^{\rm eff}_{\rm sub}(t) := \hat{H}_{\rm sub} + \langle t| \, \hat{H}_{\rm int} \, |t\rangle$, the conditional subsystem state satisfies the *conditional Schrödinger equation*

$$\boxed{i\hbar \, \partial_t \, |\Psi_{\rm sub}(t)\rangle = \hat{H}^{\rm eff}_{\rm sub}(t) \, |\Psi_{\rm sub}(t)\rangle} \quad \text{as stated in (13.238).}$$

This is the Page–Wootters realization of emergent dynamics from the universal constraint, consistent with the general relational construction in Section 13.6.2 and the good–clock hypotheses of Section 5.3.8 [215, 3, 607, 610]. □

Clarifying the State–First Reading of (13.237) *Notation convention.* Throughout the treatise the *Quantum Holo–State* $\left|\mathring{\mathbf{\Psi}}\right\rangle$ is the **primary** object; Hilbert spaces and operators are *emergent* via the GNS/BRST construction. Thus any phrase like "$\left|\mathring{\mathbf{\Psi}}\right\rangle \in \mathcal{H}_{\rm phys}$" is shorthand for the following state–first statement.

(i) Primary datum A normalized state ω° on the universal $*$-algebra of observables $\mathfrak{A}_{\rm univ}$ (with BRST structure).

(ii) GNS/BRST representation From ω° obtain the physical triple $(\pi_{\rm phys}, \mathcal{H}_{\rm phys}, |\Omega_{\rm phys}\rangle)$ by *(i)* GNS construction, *(ii)* quotient by nulls, and *(iii)* BRST cohomology. We *identify* $|\Omega_{\rm phys}\rangle$ with the Holo–State symbolically as

$$|\Omega_{\rm phys}\rangle \equiv \left|\mathring{\mathbf{\Psi}}\right\rangle.$$

(iii) Affiliated operators The Total Relational Hamiltonian $\hat{H}_{\rm univ}$ is an (essentially self–adjoint) operator affiliated with the von Neumann closure $\mathfrak{A}''_{\rm univ}$. In the physical representation it is $\pi_{\rm phys}(\hat{H}_{\rm univ})$ acting on $\mathcal{H}_{\rm phys}$.

Representation–Theoretic Restatement of (13.237). The deparametrized split and the universal constraint are to be read as:

$$\pi_{\rm phys}(\hat{H}_{\rm univ}) = \hat{p}_t \otimes \mathbf{1}_{\rm sub} + \mathbf{1}_{\rm clock} \otimes \hat{H}_{\rm sub} + \hat{H}_{\rm int}, \qquad \pi_{\rm phys}(\hat{H}_{\rm univ}) \, |\Omega_{\rm phys}\rangle = 0, \tag{13.239}$$

where the *tensor product* reflects an effective algebraic factorization of the *physical algebra*,

$$\mathfrak{A}_{\text{phys}} \;\simeq\; \mathfrak{A}_{\text{sub}} \,\widehat{\otimes}\, \mathfrak{A}_{\text{clock}},$$

under the good–clock assumptions stated in Proposition 13.172. The Hilbert–space factorization $\mathcal{H}_{\text{phys}} \simeq \mathcal{H}_{\text{sub}} \otimes \mathcal{H}_{\text{clock}}$ is then the *GNS image* of this algebraic split for the (approximately product) state induced by ω° on $\mathfrak{A}_{\text{sub}} \otimes \mathfrak{A}_{\text{clock}}$. Accordingly, all appearances of $\left|\overset{\circ}{\mathbf{\Psi}}\right\rangle \in \mathcal{H}_{\text{phys}}$ should be read as "$\left|\overset{\circ}{\mathbf{\Psi}}\right\rangle$ denotes the GNS/BRST physical cyclic vector $|\Omega_{\text{phys}}\rangle$ representing the primal state ω°."

Remark 13.173 (On Factorization and Split Property)**.** The split $\mathfrak{A}_{\text{phys}} \simeq \mathfrak{A}_{\text{sub}} \widehat{\otimes} \mathfrak{A}_{\text{clock}}$ is an *effective* statement (split/nuclearity or asymptotic decoupling); it is not a fundamental postulate about pre–existing tensor factors. All conditional states and the Page–Wootters construction in Section 13.6.2 should be interpreted in this representation–theoretic sense.

Corollary 13.174 (Density–Operator Form and Positive Operator–Valued Measure (POVM) Clocks)**.** *Let* $\varrho = \left|\overset{\circ}{\mathbf{\Psi}}\right\rangle\!\left\langle\overset{\circ}{\mathbf{\Psi}}\right|$ *and define the conditional state by Lüders' rule* $\varrho_{\text{sub}|t} := \frac{\text{Tr}_{\text{clock}}[(\Pi_t \otimes \mathbf{1})\,\varrho\,(\Pi_t \otimes \mathbf{1})]}{\text{Tr}[(\Pi_t \otimes \mathbf{1})\,\varrho]}$*. Then* $i\hbar\,\partial_t \varrho_{\text{sub}|t} = [\,\hat{H}^{\text{eff}}_{\text{sub}}(t),\, \varrho_{\text{sub}|t}\,]$*. The same holds for a realistic (imperfect) clock with a POVM* $\{E_t\}$ *replacing* Π_t*, up to small dissipative corrections controlled by the clock's energy–time uncertainty.*

Proof. **(PVM clock).** Let $\varrho = \left|\overset{\circ}{\mathbf{\Psi}}\right\rangle\!\left\langle\overset{\circ}{\mathbf{\Psi}}\right|$ on $\mathcal{H}_{\text{sub}} \otimes \mathcal{H}_{\text{clock}}$, and let $\{\Pi_t = |t\rangle\langle t|\}$ be the spectral projectors of the clock time operator $\hat{t}$ (spectral theorem in the rigged sense [369]). By definition,

$$\varrho_{\text{sub}|t} = \frac{\text{Tr}_{\text{clock}}\Big[(\Pi_t \otimes \mathbf{1})\,\varrho\,(\Pi_t \otimes \mathbf{1})\Big]}{\text{Tr}\Big[(\Pi_t \otimes \mathbf{1})\,\varrho\Big]}\,.$$

Write the universal state in the time basis (distributionally) and set $|\widetilde{\Psi}_{\text{sub}}(t)\rangle := \langle t|\,\overset{\circ}{\mathbf{\Psi}}\rangle \in \mathcal{H}_{\text{sub}}$. Then

$$\text{Tr}_{\text{clock}}\Big[(\Pi_t \otimes \mathbf{1})\,\varrho\,(\Pi_t \otimes \mathbf{1})\Big] = |\widetilde{\Psi}_{\text{sub}}(t)\rangle\langle\widetilde{\Psi}_{\text{sub}}(t)|\,, \qquad \text{Tr}\Big[(\Pi_t \otimes \mathbf{1})\,\varrho\Big] = \|\,|\widetilde{\Psi}_{\text{sub}}(t)\rangle\,\|^2,$$

hence $\varrho_{\text{sub}|t} = |\Psi_{\text{sub}}(t)\rangle\langle\Psi_{\text{sub}}(t)|$ with $|\Psi_{\text{sub}}(t)\rangle := |\widetilde{\Psi}_{\text{sub}}(t)\rangle\,/\|\,|\widetilde{\Psi}_{\text{sub}}(t)\rangle\,\|$ (standard conditionalization for projective measurements [287]). By Proposition 13.172 and (13.238),

$$i\hbar\,\partial_t\,|\Psi_{\text{sub}}(t)\rangle = \hat{H}^{\text{eff}}_{\text{sub}}(t)\,|\Psi_{\text{sub}}(t)\rangle\,, \qquad \hat{H}^{\text{eff}}_{\text{sub}}(t) := \hat{H}_{\text{sub}} + \langle t|\,\hat{H}_{\text{int}}\,|t\rangle\,.$$

Taking the time derivative of $\varrho_{\text{sub}|t} = |\Psi_{\text{sub}}(t)\rangle\langle\Psi_{\text{sub}}(t)|$ and using the Schrödinger equation and its adjoint yields the Liouville–von Neumann (commutator) form,

$$i\hbar\,\partial_t\,\varrho_{\text{sub}|t} = \Big[\hat{H}^{\text{eff}}_{\text{sub}}(t),\, \varrho_{\text{sub}|t}\Big],$$

as claimed [287].

(POVM clock). Let $\{E_t\}_{t\in\mathbb{R}}$ be a POVM on $\mathcal{H}_{\text{clock}}$ (resolution of the identity, $E_t \geq 0$). Choose Kraus operators M_t with $M_t^\dagger M_t = E_t$ and define the Lüders-type update

$$\varrho_{\text{sub}|t} := \frac{\text{Tr}_{\text{clock}}\Big[(M_t \otimes \mathbf{1})\, \varrho\, (M_t^\dagger \otimes \mathbf{1})\Big]}{\text{Tr}\Big[(E_t \otimes \mathbf{1})\, \varrho\Big]},$$

so that the reduced dynamics is completely positive and trace-preserving (CPTP) at each t [287]. By Naimark dilation, there exists an ancilla $\mathcal{K}$, an isometry $V : \mathcal{H}_{\text{clock}} \to \mathcal{H}_{\text{clock}} \otimes \mathcal{K}$, and a PVM $\{\Pi_t^{\text{N}}\}$ on $\mathcal{H}_{\text{clock}} \otimes \mathcal{K}$ such that $E_t = V^\dagger \Pi_t^{\text{N}} V$ [287]. Consider the dilated state $\tilde{\varrho} := (V \otimes \mathbf{1})\, \varrho\, (V^\dagger \otimes \mathbf{1})$ and apply the PVM result above to obtain on the dilation

$$i\hbar\, \partial_t\, \tilde{\varrho}_{\text{sub}|t} = \Big[\tilde{H}_{\text{sub}}^{\text{eff}}(t),\ \tilde{\varrho}_{\text{sub}|t}\Big].$$

Compressing back with $V^\dagger(\cdot)V$ induces on $\mathcal{H}_{\text{sub}}$ a generator of the form

$$i\hbar\, \partial_t\, \varrho_{\text{sub}|t} = \Big[\hat{H}_{\text{sub}}^{\text{eff}}(t),\ \varrho_{\text{sub}|t}\Big] \ +\ i\hbar\, \mathcal{D}_t\Big[\varrho_{\text{sub}|t}\Big],$$

where $\mathcal{D}_t$ is a completely positive, trace-preserving dissipator that vanishes in the projective limit $E_t \to \Pi_t$ [287]. If, in addition, the POVM is covariant under clock translations generated by the clock Hamiltonian $\hat{H}^{\text{clock}}$ (good-clock lane; cf. Section 5.3.8), i.e. $E_t = e^{-\frac{i}{\hbar}t\hat{H}^{\text{clock}}} E_0 e^{\frac{i}{\hbar}t\hat{H}^{\text{clock}}}$, then $\partial_t E_t = -\frac{i}{\hbar}\,[\hat{H}^{\text{clock}}, E_t]$ and the size of $\mathcal{D}_t$ is controlled by the noncommutativity $[\hat{H}^{\text{clock}}, E_t]$ (POVM covariance and Heisenberg evolution [287]). In the good-clock regime, the energy–time tradeoff (clock imprecision σ_t versus energy spread $\sigma_{H^{\text{clock}}}$) ensures that $\|\mathcal{D}_t\|$ is small and tends to zero as the POVM sharpens, yielding the stated unitary commutator dynamics up to small dissipative corrections [22, 287].

Combining the PVM case with the dilation/compression argument for POVMs establishes the corollary: Lüders-conditioned subsystem states obey $i\hbar\,\partial_t \varrho_{\text{sub}|t} = [\hat{H}_{\text{sub}}^{\text{eff}}(t), \varrho_{\text{sub}|t}]$, and realistic POVM clocks reproduce this evolution modulo dissipative terms governed by the clock's energy–time uncertainty, consistently with eq. (13.238) and (13.238). □

Conceptual Placement and Comparisons. Equation (13.238) is a precise Page–Wootters realization: a globally static, entangled state subject to the Universal Quantum Constraint (13.236) produces subsystem Schrödinger dynamics once conditioned on internal clock readings [215]. In the IQR framework this construction is fully background–independent and dovetails with the emergence of proper time (Section 5.3.8). Formally, it is encapsulated in eq. (13.238) and its density-operator formulation Corollary 13.174, and abstracted in the relational Heisenberg/Liouville picture of Proposition 13.170. The conceptual contrast with *Decoherent Histories* [611, 612] is sharp: there, time indexes sequences in a decoherence functional driven by environment-induced suppression of interference; here, evolution is generated by *correlations with an internal clock* embedded in the cosmic symmetry stack (Sections 13.4 and 13.4.1), with no external temporal parameter.

Relation to Modular/KMS Flows. In stationary settings (black-hole horizons, static/cosmological patches), the conditioning can equivalently be taken with respect to the modular

parameter of the von Neumann algebra of observables. The resulting dynamics is KMS–thermal with respect to the modular Hamiltonian and coheres with the effective temperatures and chemical potentials used in the black-hole and dark-universe sectors (Sections 13.3 and 16.9). This modular analogue is the operator–algebraic counterpart of Proposition 13.170, as standard in Tomita–Takesaki theory [608, 230, 241].

Good–Clock Conditions and Interactions. The conditional Schrödinger evolution relies on "good–clock" conditions: (i) a semiclassical clock with $\Delta t\,\Delta p_t \gg \hbar/2$ on the interval of interest; (ii) weak clock–system entangling power so that the fiber operator $\langle t|\,\hat{H}_{\rm int}\,|t\rangle$ is well defined and slowly varying (assumptions (a)/(b) in eq. (13.238)); and (iii) canonical placement of the clock lanes ensuring TT–nullity and commutation in the symmetry stack (Sections 13.4 and 13.4.1). Beyond this regime, Corollary 13.174 shows that projecting with a realistic POVM clock $\{E_t\}$ yields a time–local master equation $i\hbar\,\partial_t \varrho_{{\rm sub}|t} = [\hat{H}^{\rm eff}_{\rm sub}(t), \varrho_{{\rm sub}|t}] + i\hbar\,\mathcal{D}_t[\varrho_{{\rm sub}|t}]$, where the dissipator $\mathcal{D}_t$ is controlled by the clock's energy–time uncertainty and vanishes in the sharp (PVM) limit, recovering (13.238).

Operational Summary. The universe obeys a single timeless law, $\hat{H}_{\rm univ}\big|\mathring{\mathbf{\Psi}}\big\rangle = 0$ (Theorem 13.168), constructed lane–wise on the cosmic symmetry stack with calibrated boundary terms. Subsystem dynamics—and the appearance of a Schrödinger equation—emerge by conditioning on an internal clock encoded within $\big|\mathring{\mathbf{\Psi}}\big\rangle$. The "time" t in (13.238) is the emergent proper time of Section 5.3.8, not an external parameter. This resolves the tension between a timeless universal law and a time–evolving cosmos, and supplies the operational bridge to the effective field dynamics used elsewhere in the treatise (Sections 13.3 and 16.9).

Chapter 14

The Born Rule and Measurement

14.1 A Cosmic Context for Measurement

The *Holosymmetric Interferometer Model* functions as a *cosmic comparison engine* by putting *whole histories* into interferometric contact. Formally, a control qubit with basis $\{|\mathbf{0}\rangle_c, |\mathbf{1}\rangle_c\}$ steers two dynamical histories on the dual radiant registers, implemented by the *controlled* operator

$$\mathbf{U}_{\text{ctrl}} = |\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G} \quad \text{(eqs. (3.47) and (3.48) and Definition 2.123).}$$

Sandwiching by Hadamards on the control defines the symmetric device $\mathbf{U}_{\text{holo}}^{(\text{sym})} := \mathbf{U}_{\text{Had}}\mathbf{U}_{\text{ctrl}}\mathbf{U}_{\text{Had}}$ (Definition 2.123). For a joint input $|\Psi_{\text{init}}\rangle := |\mathbf{0}\rangle_c \otimes |\Psi_{\text{RR}}\rangle$ with $|\Psi_{\text{RR}}\rangle \in \mathcal{H}_{\text{RR}}^{(2n)}$ (Definition 2.113), the "bright-port" probability is

$$p_0 = \tfrac{1}{4}\left\|(\hat{\mathbf{U}}_{\text{RR}}^{A_G B_N} + \hat{\mathbf{U}}_{\text{RR}}^{A_N B_G})\,|\Psi_{\text{RR}}\rangle\right\|^2 \quad \text{(Corollary 3.3 and eq. (3.1)).}$$

This readout is a *Born valuation* of a holosymmetric effect $E_{\text{Holo}}^{(\mathbf{a}\nrightarrow\mathbf{b})}$ in the GNS emergent arena, $v(\rho; E) = \text{Tr}(\rho E)$ (Definitions 1.18 and 1.34). The interferometer thus implements, at laboratory scale, the same *relational* comparison that Chapter 13 uses at cosmic scale to adjudicate entire universal histories via the Holosymmetric Selection Principle (HSP): Stage I (BRST viability) and Stage II (maximal holosymmetric overlap) (Sections 13.1.3 and 13.1.4).

Cosmologically, the histories propagate on the chronogeometric background selected by the *Universal Quantum Constraint* (UQC),

$$\boxed{\hat{H}_{\text{univ}}\left|\mathring{\mathbf{\Psi}}\right\rangle = 0} \quad \text{with}$$

$$\hat{H}_{\text{univ}} = \hat{H}_\Phi + \hat{H}_\Theta + \hat{H}_{\text{int}} + \hat{H}_\partial \quad \text{(Theorem 13.168, eq. (13.236), and Definition 13.167).}$$

On BRST-physical diagnostic realized reference configuration, the cosmic symmetry stack

$$\mathcal{G}_{\text{Cosmos}} = \left[\text{Diff}_{\text{rel}} \ltimes \left(\text{SO}(1,3)_{\text{local}} \times \mathcal{G}_{\text{Dark}}^{(\text{eff})}\right)\right] \times G_{\text{SM}} \quad \text{(Section 13.4.1)}$$

acts on the unified fluctuation $\delta\hat{\Sigma}_{\mu\nu} = \hat{h}_{\mu\nu} \oplus \hat{\tau}_{\mu\nu}$ with *typed TT-orthogonality*

$$\hat{h}_{\mu\nu} := \Pi_{\bar{\Sigma}}^{[\text{TT}]}\,\delta\hat{g}_{\mu\nu}, \qquad \Pi_{\bar{\Sigma}}^{[\text{TT}]}\,\hat{\tau}_{\mu\nu} = 0 \quad \text{(eqs. (10.4) and (13.119)).}$$

Within this stack, dark/visible/vacuum lanes are selected by BRST-physical projectors

$$\mathcal{P}^{\mathrm{sym}}_{\mathrm{vac}} + \mathcal{P}^{\mathrm{coh}}_{\mathrm{dark}} + \mathcal{P}^{\mathrm{coh}}_{\mathrm{vis}} = \mathbf{1}_{\mathrm{phys}},$$
$$\rho_{\mathrm{tot}} = \rho_{\mathrm{DE}} + \rho_{\mathrm{DM}} + \rho_{\mathrm{vis}} \quad \text{(eqs. (13.149) and (13.150) and Proposition 13.97).}$$

In this sense, the universe is a *self-measuring system*: the same GNS–Born mechanism that evaluates p_0 locally evaluates cosmic *self-coherence* globally, with HSP selecting the "bright fringe" history.

14.1.1 The Path Scattering Model: A Template for Collapse

The *Holosymmetric Path Scattering Model* upgrades the symmetric scheme to allow *dynamical feedback*. In the asymmetric/tomographic variant one uses arbitrary control rotations $\mathbf{U}_{\mathrm{prep}}, \mathbf{U}_{\mathrm{read}} \in \mathrm{SU}(2)$,

$$\mathbf{U}^{(\mathrm{asym})}_{\mathrm{holo}} := \mathbf{U}_{\mathrm{read}}\, \mathbf{U}_{\mathrm{ctrl}}\, \mathbf{U}_{\mathrm{prep}}, \quad \text{with} \quad \mathbf{U}_{\mathrm{ctrl}} \text{ as in eq. (3.47)},$$

so that the measured amplitudes $\mathcal{F}^{(\mathrm{asym})}_{ij}$ resolve interference from any Bloch-sphere angle (Section 3.1). The "interference event" (Act II) is a controlled entangler that correlates system and apparatus; the *commutator lens* makes this algebraic:

$$[\mathbf{U}_{\mathrm{Had}}, \mathbf{U}_{\mathrm{ctrl}}] = \tfrac{1}{\sqrt{2}} \begin{pmatrix} 0 & \hat{\mathbf{U}}^{A_G B_N}_{\mathrm{RR}} - \hat{\mathbf{U}}^{A_N B_G}_{\mathrm{RR}} \\ \hat{\mathbf{U}}^{A_N B_G}_{\mathrm{RR}} - \hat{\mathbf{U}}^{A_G B_N}_{\mathrm{RR}} & 0 \end{pmatrix}, \quad \text{(Definition 3.12)}$$

so that *fringes are commutators in disguise*. Conditioning on a control outcome implements Act III: *conditional re-evolution*. E.g., postselecting $|\mathbf{0}\rangle_c$ yields the normalized subsystem state

$$|\Psi_{\mathrm{post}}\rangle = \frac{\hat{\mathbf{U}}_+ |\Psi_{\mathrm{RR}}\rangle}{\|\hat{\mathbf{U}}_+ |\Psi_{\mathrm{RR}}\rangle\|}, \qquad \hat{\mathbf{U}}_+ := \hat{\mathbf{U}}^{A_G B_N}_{\mathrm{RR}} + \hat{\mathbf{U}}^{A_N B_G}_{\mathrm{RR}} \quad \text{(Section 3.1.2)}.$$

Allowing different in/out controls ($\mathbf{U}_{\mathrm{prep}} \neq \mathbf{U}_{\mathrm{read}}$) encodes the operational *irreversibility* of information acquisition at the effective level, while the universal dynamics remains unitary (Section 3.1).

Mathematically, all probabilities here are standard *Born* valuations of holosymmetric effects in the GNS arena (Definition 1.18), and all inner products are the *emergent* ones fixed by the Fourfold correlator (below).

A New Ontology, Not a New Mechanism

No extra "collapse gear" is introduced. The ontology is unified: the *GNS* representation $(\pi_{\mathring{\mathbf{\Psi}}}, \mathcal{H}_{\mathrm{em}}, |\Omega_{\omega^{\mathring{\mathbf{\Psi}}}}\rangle)$ of the IQR observable algebra turns logical effects into operators and *Born valuations* into standard expectations,

$$v(\rho; E) = \mathrm{Tr}(\rho E) \quad \text{(Definitions 1.18 and 1.34)}.$$

Inner products and overlaps are *derived*, not assumed: Chapter 4 defines the *Fourfold Inter–Modal Correlator* $\hat{G}_\Lambda$ and induces the fibre inner product at a PQS locus $\mathfrak{B}_x$ by the Holo–State expectation,

$$\langle \psi_1 \mid \psi_2 \rangle_x := \lim_{x' \to x} \left\langle \mathring{\mathbf{\Psi}} \right| \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2] \left| \mathring{\mathbf{\Psi}} \right\rangle \quad \text{(Section 4.6 and Definition 4.9)}.$$

Comparisons across loci $x \leftrightarrow x'$ use the gauge-covariant parallel transport $\mathcal{U}(x \leftarrow x'; \gamma) = \mathcal{P}\exp\left(-\int_\gamma \mathcal{A}\right)$ (Section 4.1.4), and the triad of distances $(d_{\mathrm{corr}}, d_{\mathrm{vac}}, d_{\mathrm{spec}})$ organizes distinguishability on $\mathcal{X}_q$ (Section 4.11.5). Thus, "collapse" is just *quantum evolution* viewed through conditioning in a fibre bundle whose metrics and transports are emergent (Ch. 4), whose effects and valuations are GNS-realized (Tetralemma–GNS), and whose cosmic background is selected by HSP under the UQC (Ch. 13).

On the visible lane (Ch. 12), the realized weave configuration-compatible operator–bundle connection $\nabla^{\mathrm{realizedweaveconfiguration}}$ and color-singlet projector enforce BRST-physicality and anomaly control for SM composites (Definitions 12.1 and 12.126 and Section 12.6); typed TT-purity prevents spurious spin-2 leakage (eq. (10.4)). These guarantees flow into the Ward analysis behind the UQC (Ch. 13).

The Selection Principle: Architect of the Rules

The *Holosymmetric Selection Principle* sharpens the cosmic filter in two steps. *Stage I* imposes BRST viability (anomaly meters off, unitary support on) and typed orthogonality; *Stage II* maximizes holosymmetric overlap among the survivors (Sections 13.1.3 and 13.4.1). The same BRST-physical projectors that partition the universal Hamiltonian's spectrum

$$\hat{H}_{\mathrm{univ}} = \hat{H}_\Phi + \hat{H}_\Theta + \hat{H}_{\mathrm{int}} + \hat{H}_\partial \quad \rightsquigarrow \quad \rho_{\mathrm{tot}} = \rho_{\mathrm{DE}} + \rho_{\mathrm{DM}} + \rho_{\mathrm{vis}}$$

(Definitions 13.92 and 13.167 and eq. (13.150))

also implement the identity resolution (eq. (13.149)). The visible lane uses the SM fibre G_{SM}, while the dark lanes live in the τ–kairos sector (scalar Gaiaton, vector Chrono–Kymon) with typed TT-nullity (eqs. (13.113) and (13.119)); cosmological abundances are regulator-independent in the Ω_i ratios (Remark 13.96). In this frame, the familiar rules—superposition, entanglement, Schrödinger dynamics for conditionals—are not additional postulates but the *operational face* of (i) GNS–Born valuation on $\mathcal{H}_{\mathrm{em}}$ (Tetralemma–GNS), (ii) holosymmetric interferometry and path tomography (Ch. 3), (iii) emergent inner products and transports (Ch. 4), and (iv) UQC + HSP at the cosmic level (Ch. 13).

Operational takeaway. *Locally*, a detector "click" is the Born valuation of a holosymmetric effect $E_{\mathrm{Holo}}^{(\mathbf{a}\to\mathbf{b})}$ computed with the emergent inner product (Definition 1.18 and Section 4.6); *globally*, the same calculus selects the background, partitions energy, and enforces typed cleanliness via the cosmic stack and HSP (Section 13.4.1 and Proposition 13.97). Hence the *same* relational, interferometric principles that sculpt the cosmos govern the "collapse-like" appearance in a single apparatus.

Measurement as the Enactment of the Selected Rules

A local measurement *enacts* the rules fixed globally by the *Holosymmetric Selection Principle* (HSP): the universe that survives Stage I (BRST viability) and maximizes Stage II holosymmetric overlap evolves operationally by the *Holosymmetric Path Scattering* template

$$\text{Evolve} \longrightarrow \text{Interfere} \longrightarrow \text{Re-evolve} \quad \text{(Sections 3.1 and 13.1.3)}.$$

In the laboratory, the same logic is realized by a control–conditioned interferometer on the dual radiant registers. With control basis $\{|\mathbf{0}\rangle_c, |\mathbf{1}\rangle_c\}$ and dynamical histories $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N}$, $\hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G}$, the *controlled* block is

$$\mathbf{U}_{\mathrm{ctrl}} = |\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} \quad \text{(eqs. (3.47) and (3.48))}.$$

The symmetric device $\mathbf{U}_{\mathrm{holo}}^{(\mathrm{sym})} := \mathbf{U}_{\mathrm{Had}}\mathbf{U}_{\mathrm{ctrl}}\mathbf{U}_{\mathrm{Had}}$ (Definition 2.123) and its tomographic/asymmetric generalization

$$\mathbf{U}_{\mathrm{holo}}^{(\mathrm{asym})} := \mathbf{U}_{\mathrm{read}}\,\mathbf{U}_{\mathrm{ctrl}}\,\mathbf{U}_{\mathrm{prep}}, \qquad \mathbf{U}_{\mathrm{prep}}, \mathbf{U}_{\mathrm{read}} \in \mathrm{SU}(2) \quad \text{(Section 3.1)}$$

implement the *Interfere* stage; conditioning on a control outcome initiates the *Re-evolve* stage.

Concretely, for an electron (system S) and a detector (apparatus A) prepared as $|\Psi_{\mathrm{init}}\rangle := |\mathbf{0}\rangle_c \otimes |\Psi_{\mathrm{RR}}\rangle$ with $|\Psi_{\mathrm{RR}}\rangle \in \mathcal{H}_{\mathrm{RR}}^{(2n)}$ (Definition 2.113), the "bright-port" probability reads

$$p_0 = \tfrac{1}{4}\left\|(\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G})\,|\Psi_{\mathrm{RR}}\rangle\right\|^2 \quad \text{(Corollary 3.3 and eq. (3.1))}.$$

Postselecting $|\mathbf{0}\rangle_c$ yields the normalized conditional state

$$|\Psi_{\mathrm{post}}\rangle = \frac{\hat{\mathbf{U}}_+\,|\Psi_{\mathrm{RR}}\rangle}{\|\hat{\mathbf{U}}_+\,|\Psi_{\mathrm{RR}}\rangle\|}, \qquad \hat{\mathbf{U}}_+ := \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} \quad \text{(Section 3.1.2)}.$$

All readouts are *Born valuations* $v(\rho; E) = \mathrm{Tr}(\rho E)$ of holosymmetric effects $E_{\mathrm{Holo}}^{(\mathbf{a}\to\mathbf{b})}$ in the GNS emergent arena (Definitions 1.18 and 1.34), with inner products *derived* from the Fourfold Inter–Modal Correlator as in Section 4.6 and Definition 4.9. Thus, a laboratory measurement *instantiates* the same selection-templated logic that operates cosmically under the *Universal Quantum Constraint* (UQC),

$$\boxed{\hat{H}_{\mathrm{univ}}\left|\mathring{\mathbf{\Psi}}\right\rangle = 0},$$

$$\hat{H}_{\mathrm{univ}} = \hat{H}_\Phi + \hat{H}_\Theta + \hat{H}_{\mathrm{int}} + \hat{H}_\partial$$

(Theorem 13.168, eq. (13.236), and Definition 13.167).

1. *Evolution.* Free evolution (diagnostic realized reference configuration-wise) is generated within the emergent/GNS space $\mathcal{H}_{\mathrm{em}}$; with a good internal clock C (TT-safe, BRST-closed; Section 5.3.8) one has the conditional Schrödinger dynamics

$$i\hbar\,\partial_t\,\varrho_{A|C=t} = \left[H_{\mathrm{eff}}(t),\ \varrho_{A|C=t}\right], \qquad H_{\mathrm{eff}}(t) = \frac{\Pi_t\,\hat{H}_{\mathrm{rest}}\,\Pi_t}{\mathrm{Tr}(\Pi_t\varrho)} \quad \text{(eq. (13.238))}.$$

2. *Interference.* Interaction with the detector is a controlled entangler implemented by $\mathbf{U}_{\mathrm{ctrl}}$ above; the algebraic origin of fringes is exposed by the *commutator lens*

$$[\mathbf{U}_{\mathrm{Had}}, \mathbf{U}_{\mathrm{ctrl}}] \propto \begin{pmatrix} 0 & \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} - \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} \\ \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} - \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} & 0 \end{pmatrix} \quad \text{(Definition 3.12)}.$$

3. *Re-evolution.* Conditioning on the pointer lane (control outcome or apparatus projector P_m) collapses *operationally* to the normalized branch

$$|\Psi^{(m)}_{\text{post}}\rangle = \frac{(P_m \otimes \mathbf{1})\,\mathbf{U}^{(\text{asym})}_{\text{holo}}\,|\Psi_{\text{init}}\rangle}{\sqrt{\langle\Psi_{\text{init}}|\mathbf{U}^{(\text{asym})\dagger}_{\text{holo}}(P_m \otimes \mathbf{1})\,\mathbf{U}^{(\text{asym})}_{\text{holo}}|\Psi_{\text{init}}\rangle}},$$

while the subsystem continues to evolve by the conditional equation above (eq. (13.238)).

14.1.2 A Universe Consistent with its Own Creation

The HSP plays a dual role: it *fixes the stage*—laws, symmetry stack, diagnostic realized reference configuration, and clocks—and *directs the play*—how interactions enact those laws. The cosmic stack

$$\mathcal{G}_{\text{Cosmos}} = \left[\text{Diff}_{\text{rel}} \ltimes (\text{SO}(1,3)_{\text{local}} \times \mathcal{G}^{(\text{eff})}_{\text{Dark}})\right] \times G_{\text{SM}}$$
$$\text{acts on} \quad \delta\hat{\Sigma}_{\mu\nu} = \hat{h}_{\mu\nu} \oplus \hat{\tau}_{\mu\nu}$$
(Section 13.4.1 and eq. (13.119)).

Typed TT-orthogonality, BRST cleanliness, and anomaly control (Stage I) ensure a viable arena; Stage II maximizes holosymmetric overlap across histories (Sections 13.1.4 and 13.4.1). In this calibrated setting the universal Hamiltonian decomposes lane-wise,

$$\hat{H}_{\text{univ}} = \hat{H}_\Phi + \hat{H}_\Theta + \hat{H}_{\text{int}} + \hat{H}_\partial \;\rightsquigarrow\; \rho_{\text{tot}} = \rho_{\text{DE}} + \rho_{\text{DM}} + \rho_{\text{vis}}$$
(Definitions 13.92 and 13.167 and eq. (13.150)).

Hence the same GNS/Holosymmetry machinery that fixes probabilities locally also *selects* the global background under the UQC (Theorem 13.168). Alternative "rules" fail either Stage I (anomaly or typed-contamination) or Stage II (destructive self-interference).

The Pre-Measurement State: A Superposition of Histories

Prior to readout, the system is not a mere superposition of eigenvalues but of *histories* (dialected evolutions). In the holosymmetric language the governing object is the *Fourfold Inter–Modal Correlator* $\hat{G}_\Lambda$ (Definition 4.9), which induces, at each PQS locus $\mathfrak{B}_x$, the *emergent* modal inner product

$$\langle\psi_1 \mid \psi_2\rangle_x = \lim_{x'\to x} \left\langle \mathring{\mathbf{\Psi}} \right| \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2] \left| \mathring{\mathbf{\Psi}} \right\rangle \quad \text{(Section 4.6)}.$$

The pre-measurement content is therefore *tetradic* (Tetralemma) at the algebraic level and realized as a *GNS* inner product at the operational level (Definitions 1.34 and 6.4).

Measurement as Relational Entanglement

A measurement is a physical interaction that yields robust entanglement between S and A in the path-scattering scheme. With an eigenbasis $\{|s_i\rangle\}$ for the measured meter and pointer

states $\{|A_i\rangle\}$,

$$\hat{U}_{\text{int}}\left(\sum_i c_i|s_i\rangle \otimes |A_0\rangle\right) = \sum_i c_i\,|s_i\rangle \otimes |A_i\rangle,$$

and the outcome i is read by the Born valuation

$$p_i = \langle\Psi_{\text{fin}}|\ (\mathbf{1}\otimes P_i)\ |\Psi_{\text{fin}}\rangle = \left|c_i\right|^2 \quad \text{(Section 14.3.2 and eq. (14.45)).}$$

In holosymmetric form this is the valuation of $E^{(\mathbf{a}\to\mathbf{b})}_{\text{Holo}}$ (Definition 1.18), and the same commutator lens (Definition 3.12) diagnoses the interference structure that the apparatus resolves.

Decoherence: The Irreversible Loss of Information to the Whole

Coupling to an environment E produces

$$|\Psi_{\text{SAE}}\rangle = \sum_i c_i\,|s_i\rangle \otimes |A_i\rangle \otimes |E_i\rangle,$$

so the reduced state on S (or $S{+}A$) is

$$\rho_S = \sum_i |c_i|^2\,|s_i\rangle\langle s_i| + \sum_{i\neq j} c_i c_j^*\,\langle E_j|E_i\rangle\,|s_i\rangle\langle s_j|.$$

As the overlaps $\langle E_j|E_i\rangle$ are driven to 0 by the environment's explosion of correlational distance, the off-diagonals are suppressed. Chapter 4 formalizes this with the triad of distances on $\mathcal{X}_q$, $(d_{\text{corr}}, d_{\text{vac}}, d_{\text{spec}})$, whose functional independence controls distinguishability and thus the decay of interference terms under coarse-graining (Section 4.11.5). The effective classicality that remains is a *Twofold* restriction (dyadic lane) consistent with the Axiom of Duality at readout, while the full Fourfold structure survives globally in the correlations (Sections 4.1.6 and 14.1.3). In all cases, the dynamics is *unitary and relational* at the universal level (UQC; Theorem 13.168); "collapse'' is the local, conditional re-evolution prescribed by the Page–Wootters construction (eq. (13.238)), enacted within the symmetry stack and projectors of the HSP-selected cosmos (Section 13.4.1 and eq. (13.149)).

14.1.3 The Axiom of Duality: The Logic of a Classical Outcome

After interaction and decoherence, the readout must be *communicable* as definite information. In the present framework this is encoded by the *Axiom of Duality*—the minimal logical skeleton for distinction—implemented on the *Twofold* subspace with orthonormal basis

$$\left\{\,|\mathring{\mathbf{0}}\rangle,\ |\mathring{\mathbf{1}}\rangle\,\right\}, \qquad \langle\mathring{\mathbf{i}}\,|\,\mathring{\mathbf{j}}\rangle = \delta_{ij},$$

and spectral projectors

$$P_{\mathbf{0}} := |\mathring{\mathbf{0}}\rangle\langle\mathring{\mathbf{0}}|, \qquad P_{\mathbf{1}} := |\mathring{\mathbf{1}}\rangle\langle\mathring{\mathbf{1}}|, \qquad P_{\mathbf{0}} + P_{\mathbf{1}} = \mathbf{1}_{\text{Twofold}}, \quad P_{\mathbf{0}}P_{\mathbf{1}} = 0.$$

This Twofold readout is the *dyadic restriction* of the Fourfold dialectic structure (Thesis/Antithesis/Parasymmetry/Holosymmetry) discussed in the Tetralemma–GNS foundations, and it is the regime captured in Chapter 4 by the *duality–collapse* conditions where the Fourfold correlator yields an inner product on the Twofold subspace (Sections 4.1.6 and 4.6 and Definition 4.9). The classical outcome "**0** or **1**" is then a standard *Born valuation*

$$p_{\mathbf{i}} = \mathrm{Tr}\big(\rho\, P_{\mathbf{i}}\big) = \langle \Psi_{\mathrm{fin}} | P_{\mathbf{i}} | \Psi_{\mathrm{fin}} \rangle \quad \text{(Definition 1.18 and Section 14.3.2)}.$$

In this sense, "collapse" is the *final informational stage* of an emergent process whose dynamics is unitary and relational at the universal level (UQC: $\hat{H}_{\mathrm{univ}} |\overset{\circ}{\mathbf{\Psi}}\rangle = 0$, Theorem 13.168 and eq. (13.236)) and whose readouts use the *emergent inner product* induced by the Fourfold Inter–Modal Correlator (Ch. 4, Section 4.6).

1. *Fourfold superposition (premeasurement).* A rich, multi–layered state carries the tetradic logic; its local fibre inner product at a PQS locus $\mathfrak{B}_x$ is generated by the Holo–State expectation

$$\langle \psi_1 \mid \psi_2 \rangle_x = \lim_{x' \to x} \Big\langle \overset{\circ}{\mathbf{\Psi}} \Big| \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2] \Big| \overset{\circ}{\mathbf{\Psi}} \Big\rangle \quad \text{(Section 4.6)}.$$

2. *Relational interaction (measurement).* The *path–scattering* entangler correlates system and apparatus via the controlled block

$$\mathbf{U}_{\mathrm{ctrl}} = |\mathbf{0}\rangle_c\langle \mathbf{0}|_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + |\mathbf{1}\rangle_c\langle \mathbf{1}|_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} \quad \text{(eqs. (3.47) and (3.48))}.$$

3. *Decoherence and dyadic readout.* Environmental dispersal suppresses off–diagonals in the preferred basis; the dyadic Twofold subspace furnishes the binary, communicable outcome by the Axiom of Duality (Section 4.1.6).

The Two Faces of Quantum Reality

Wave–particle duality is resolved as *two stages* of one process—Fourfold superposition prior to readout versus Twofold restriction at readout—implemented within the same GNS/Holosymmetry arena (Definitions 1.18 and 1.34).

The Wave Behavior: Reality as Superposition

Before observation, the system is described by a Fourfold–structured state whose amplitudes and interference live on the emergent fibre metric. In a double–slit register, the symmetric holosymmetric device

$$\mathbf{U}_{\mathrm{holo}}^{(\mathrm{sym})} = \mathbf{U}_{\mathrm{Had}} \mathbf{U}_{\mathrm{ctrl}} \mathbf{U}_{\mathrm{Had}} \quad \text{(Definition 2.123)}$$

produces the bright–port probability

$$p_0 = \tfrac{1}{4} \left\| (\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G})\, |\Psi_{\mathrm{RR}}\rangle \right\|^2 \quad \text{(Corollary 3.3 and eq. (3.1))}.$$

The *commutator lens*

$$[\mathbf{U}_{\mathrm{Had}}, \mathbf{U}_{\mathrm{ctrl}}] = \tfrac{1}{\sqrt{2}} \begin{pmatrix} 0 & \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} - \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} \\ \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G} - \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} & 0 \end{pmatrix} \quad \text{(Definition 3.12)}$$

exhibits fringes as *group–theoretic* differences of histories, while the visibility is computed with the *emergent inner product* of Ch. 4 (Section 4.6).

The Particle Behavior: Reality as Information

Placing a which–path meter implements the (localized) path–scattering interaction

$$\hat{U}_{\mathrm{int}}\Big(\sum_i c_i \,|s_i\rangle \otimes |A_0\rangle\Big) = \sum_i c_i \,|s_i\rangle \otimes |A_i\rangle,$$

followed by environmental coupling $|A_i\rangle \otimes |E_0\rangle \mapsto |A_i\rangle \otimes |E_i\rangle$. The reduced state on S becomes

$$\rho_S = \sum_i |c_i|^2 \,|s_i\rangle\langle s_i| + \sum_{i\neq j} c_i c_j^* \,\langle E_j|E_i\rangle \,|s_i\rangle\langle s_j|,$$

so decoherence suppresses off–diagonals by the overlap factors $\langle E_j|E_i\rangle \to 0$. Chapter 4 formalizes the loss of *interference access* through the triad of distances on $\mathcal{X}_q$,

$$(d_{\mathrm{corr}},\ d_{\mathrm{vac}},\ d_{\mathrm{spec}}),$$

whose functional independence governs distinguishability and the stability of the dyadic restriction under coarse–graining (Section 4.11.5). The informational readout is then the *Twofold* valuation

$$p_{\mathbf{0}} = \mathrm{Tr}\big(\rho\, P_{\mathbf{0}}\big), \qquad p_{\mathbf{1}} = \mathrm{Tr}\big(\rho\, P_{\mathbf{1}}\big), \qquad p_{\mathbf{0}} + p_{\mathbf{1}} = 1 \quad \text{(Section 14.3.2)}.$$

Operationally, this is the same Born valuation $v(\rho; E) = \mathrm{Tr}(\rho E)$ of holosymmetric effects (Definition 1.18), now computed in the Twofold lane singled out by the Axiom of Duality and the measurement context (Section 4.1.6). The disappearance of fringes is therefore not a failure of unitarity—global dynamics remains constrained by the UQC (Theorem 13.168)—but the *local* manifestation of conditional (Page–Wootters) evolution on a dyadic readout fibre consistent with the symmetry stack and projectors selected by HSP (Ch. 13; Section 13.4.1 and eq. (13.149)).

14.2 The Born Rule

The Born rule provides the quantitative link between amplitudes and probabilities [21, 176]. In the present framework it is evaluated in the *emergent/GNS* arena: inner products and norms on the modal fiber at $\mathfrak{B}_x$ are *derived* from the Fourfold Inter–Modal Correlator $\hat{G}_\Lambda$ via the Holo–State expectation,

$$\langle \psi_1 \mid \psi_2 \rangle_x := \lim_{x' \to x} \Big\langle \mathring{\mathbf{\Psi}} \Big| \, \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2] \, \Big| \mathring{\mathbf{\Psi}} \Big\rangle \quad \text{(eq. (4.85) and Definition 4.9)}.$$

Accordingly, probabilities are *Born valuations* of effects in the GNS representation,

$$v(\rho; E) = \mathrm{Tr}\Big(\rho\, \pi_{\overset{\circ}{\boldsymbol{\Psi}}}(E)\Big), \qquad v(E) = \omega^{\overset{\circ}{\boldsymbol{\Psi}}}(E) = \langle \Omega \mid \pi_{\overset{\circ}{\boldsymbol{\Psi}}}(E) \mid \Omega \rangle \quad \text{(Definitions 1.18 and 1.34)}.$$

Here E may be any of the canonical corner effects $E \in \{P_T,\ P_A,\ P_{\mathrm{para}},\ E^{(\mathbf{a}\to\mathbf{b})}_{\mathrm{Holothesis}}\}$, and $\mathcal{H}_{\mathrm{em}} \equiv \mathcal{H}_{\overset{\circ}{\boldsymbol{\Psi}}}$ is the emergent GNS space.

Holosymmetric interferometer (symmetric case). For the control–conditioned double–history device with

$$\mathbf{U}_{\mathrm{ctrl}} = |\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}^{A_G B_N}_{\mathrm{RR}} + |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}^{A_N B_G}_{\mathrm{RR}},$$
$$\mathbf{U}^{(\mathrm{sym})}_{\mathrm{holo}} = \mathbf{U}_{\mathrm{Had}}\mathbf{U}_{\mathrm{ctrl}}\mathbf{U}_{\mathrm{Had}}$$
(*eqs.* (3.47)*and* (3.48)*and Definition* 2.123),

and joint input $|\Psi_{\mathrm{init}}\rangle = |\mathbf{0}\rangle_c \otimes |\Psi_{\mathrm{RR}}\rangle$, the "constructive" (control-0) probability is

$$p_0 = \frac{1}{4}\left\| \left(\hat{\mathbf{U}}^{A_G B_N}_{\mathrm{RR}} + \hat{\mathbf{U}}^{A_N B_G}_{\mathrm{RR}}\right) |\Psi_{\mathrm{RR}}\rangle \right\|^2 \quad \text{(Section 3.1.4, Corollary 3.3, and eq. (3.1))}.$$

The "destructive" (control-1) channel analogously yields

$$p_1 = \frac{1}{4}\left\| \left(\hat{\mathbf{U}}^{A_G B_N}_{\mathrm{RR}} - \hat{\mathbf{U}}^{A_N B_G}_{\mathrm{RR}}\right) |\Psi_{\mathrm{RR}}\rangle \right\|^2, \qquad p_0 + p_1 = 1,$$

with the interference contrast exposed by the *commutator lens* $[\mathbf{U}_{\mathrm{Had}}, \mathbf{U}_{\mathrm{ctrl}}]$ (Definition 3.12).

Axiomatic use (at this stage). In Sections 3.1 and 3.4, the Born rule is *used*—not derived—to compute outcome weights once the interferometric (or path–scattering) process is specified. The deeper origin of the modulus–squared law is not assumed there; instead, it is deferred to the reconstruction program grounded in Tetralemma logic, GNS emergence, and holosymmetric constraints (see below). Operational consistency with the cosmic arena is ensured by the Universal Quantum Constraint (UQC),

$$\boxed{\hat{H}_{\mathrm{univ}} \left|\overset{\circ}{\boldsymbol{\Psi}}\right\rangle = 0},$$
$$\hat{H}_{\mathrm{univ}} = \hat{H}_{\Phi} + \hat{H}_{\Theta} + \hat{H}_{\mathrm{int}} + \hat{H}_{\partial}$$
(Theorem 13.168, eq. (13.236), and Definition 13.167),

and by the BRST–clean symmetry stack that fixes the diagnostic realized reference configuration, projectors, and clocks (Section 13.4.1 and eq. (10.4)).

14.2.1 The Challenge: Derivation of the Born Rule

Historically, Copenhagen treats the Born rule as a postulate [21, 176]; many–worlds programs attempt derivations from decision–theoretic or envariance premises [188, 170]. Methodologically, this treatise begins compatibly—accepting the quantum formalism including Born valuations—then seeks to *reconstruct* them from its own first principles: Fourfold (Tetralemma) logic ⇒ GNS emergence ⇒ emergent inner product ⇒ unique probability assignment under Stage 2–3 constraints (Section 14.3.2).

14.2.2 A Program for the Derivation of the Born Rule

Target statement. For any subsystem A within the universal Holo–State, the only observer–independent probability assignment consistent with (i) GNS emergence, (ii) holosymmetric tomography, (iii) product/mixture consistency, and (iv) continuity is $p_k = \langle\Psi|P_k|\Psi\rangle = |c_k|^2$ (Section 14.3.2).

Stage 1: Measurement as Relational Decoherence

Kinematics. Work on the emergent Hilbert arena $\mathcal{H}_{\mathrm{em}}$ furnished by the GNS construction of the IQR algebra with cyclic vector $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$ (Definitions 1.27 and 1.34). Let S (system), A (apparatus), E (environment). The preinteraction state is

$$|\Psi_{\mathrm{initial}}\rangle = |\psi_S\rangle \otimes |\phi_A\rangle \otimes |\chi_E\rangle.$$

Measurement is modeled as a localized *Path–Scattering* event (Ch. 3) with an entangling unitary U_{SA}:

$$U_{SA}\Big(|\psi_S\rangle \otimes |A_0\rangle\Big) = \sum_i c_i\,|s_i\rangle \otimes |A_i\rangle, \qquad \sum_i |c_i|^2 = 1,$$

followed by environmental coupling U_{AE} generating records $|E_i\rangle$:

$$|\Psi_{\mathrm{fin}}\rangle = \sum_i c_i\,|s_i\rangle \otimes |A_i\rangle \otimes |E_i\rangle, \qquad \langle E_i|E_j\rangle \simeq \delta_{ij}.$$

Reduced dynamics and distances. Tracing out AE (or E) yields

$$\rho_S = \sum_i |c_i|^2\,|s_i\rangle\langle s_i| + \sum_{i\neq j} c_i c_j^*\,\langle E_j|E_i\rangle\,|s_i\rangle\langle s_j|.$$

Chapter 4 quantifies the suppression of coherences by the triad of relational distances on $\mathcal{X}_q$,

$$\Big(d_{\mathrm{corr}},\ d_{\mathrm{vac}},\ d_{\mathrm{spec}}\Big),$$

whose functional independence controls distinguishability and coarse–grained stability (Section 4.11.5); increased correlational distance drives $\langle E_j|E_i\rangle \to 0$.

Clocked conditionals (compatibility with UQC). With a good internal clock C (TT–safe, BRST–closed; Section 5.3.8), the conditional state obeys the Page–Wootters Schrödinger equation

$$i\hbar\,\partial_t\,\varrho_{A|C=t} = \Big[H_{\mathrm{eff}}(t),\ \varrho_{A|C=t}\Big], \qquad H_{\mathrm{eff}}(t) = \frac{\Pi_t\,\hat{H}_{\mathrm{rest}}\,\Pi_t}{\mathrm{Tr}(\Pi_t\varrho)} \quad \text{(eq. (13.238))},$$

fully compatible with the timeless UQC (Theorem 13.168).

Stage 2 (Born identification, already realized). In Section 14.3.2 the probability of outcome k is *identified* with the projector expectation in the emergent inner product, i.e.

$$p_k = \langle\Psi_{\mathrm{fin}}|P_k|\Psi_{\mathrm{fin}}\rangle = \mathrm{Tr}(\rho\,P_k),$$

where the inner product itself is induced by $\hat{G}_\Lambda$ (Ch. 4: Section 4.6).

Stage 3 (uniqueness under holosymmetric constraints). Impose: (i) *product factorization* for independent subsystems prepared on certified plateaues; (ii) *duality consistency* at readout (Twofold lane; Section 4.1.6); (iii) *continuity* under small holosymmetric deformations; and (iv) *band/tag invariance* (diagnostic realized reference configuration neutrality). Within the GNS arena these conditions single out the quadratic rule as the unique state–independent, lane–compatible valuation consistent with the interferometer tomography of Ch. 3 and the correlator geometry of Ch. 4 (see the discussion surrounding Section 14.3.2).

Summary. Born probabilities are *presently used* to compute outcomes in interferometer and path–scattering models (Sections 3.1 and 3.4). The reconstruction program shows how, given Tetralemma logic, GNS emergence, holosymmetric tomography, and diagnostic realized reference configuration invariances, the modulus–squared valuation is the *only* rule compatible with the UQC, the symmetry stack, and the emergent inner product (Sections 4.6 and 13.4.1 and Theorem 13.168).

Stage 2: Probability from Symmetry (Relational Envariance)

We assign probabilities to pointer outcomes $P_k := \mathbf{1}_S \otimes |a_k\rangle\langle a_k| \otimes \mathbf{1}_E$ after Stage 1 has selected an orthogonal pointer basis and produced the entangled state

$$|\Psi_{\text{fin}}\rangle = \sum_k c_k\, |s_k\rangle|a_k\rangle|e_k\rangle, \qquad \langle a_k|a_\ell\rangle = \delta_{k\ell}, \quad \langle e_k|e_\ell\rangle \approx \delta_{k\ell} \quad \text{(cf. Section 14.2.2)}. \tag{14.1}$$

Local (apparatus) statistics are computed as *Born valuations* in the emergent/GNS arena,

$$\begin{aligned} p_k \;&:=\; \mathrm{Tr}\big(\rho\, P_k\big) \;=\; \mathrm{Tr}_{SA}\big(P_k\, \rho_{SA}\big), \\ \rho := |\Psi_{\text{fin}}\rangle\langle\Psi_{\text{fin}}|, \quad \rho_{SA} &:= \mathrm{Tr}_E\, \rho \quad \text{(Definitions 1.18 and 1.34)}. \end{aligned} \tag{14.2}$$

Here the trace and norms use the *emergent inner product* induced by the Fourfold Inter–Modal Correlator (Ch. 4, Section 4.6 and Definition 4.9).

Relational envariance: the core symmetry. We adapt Zurek's environment–assisted invariance to the present relational setting [158]:

1. *Locality of access.* The observer couples to A; environmental microstructure is inaccessible. Hence only transformations that act trivially on SA *relations* can affect local statistics.

2. *Phase envariance.* For any set of phases $\{\theta_k\}$ let $U_E :\ |e_k\rangle \mapsto e^{i\theta_k}|e_k\rangle$. Then $\rho'_{SA} = \mathrm{Tr}_E\big[(\mathbf{1}\otimes\mathbf{1}\otimes U_E)\,\rho\,(\mathbf{1}\otimes\mathbf{1}\otimes U_E^\dagger)\big] = \rho_{SA}$, because the phases cancel in the partial trace. Equivalently, there exists a compensating diagonal unitary $U_S :\ |s_k\rangle \mapsto e^{-i\theta_k}|s_k\rangle$ (or on A) such that $(U_S\otimes\mathbf{1}_A\otimes U_E)|\Psi_{\text{fin}}\rangle = |\Psi_{\text{fin}}\rangle$. Therefore the valuation relevant to A is *phase–insensitive*:

$$p_k \;=\; f\big(|c_k|\big), \qquad f : [0,1] \to [0,1]. \tag{14.3}$$

3. *Swap envariance ⇒ equiprobability for equal moduli.* Suppose $|c_i| = |c_j| = r > 0$. Define the swap on A $S_{ij}^{(A)}$: $|a_i\rangle \leftrightarrow |a_j\rangle$ and the counterswap on E $S_{ij}^{(E)}$: $|e_i\rangle \leftrightarrow |e_j\rangle$. Then $(S_{ij}^{(A)} \otimes S_{ij}^{(E)})\,|\Psi_{\text{fin}}\rangle = |\Psi_{\text{fin}}\rangle$, yet $S_{ij}^{(A)}$ *alone* exchanges the labels of outcomes i and j on the apparatus. Since a transformation that can be undone by acting only on the inaccessible E cannot change A's statistics, we must have

$$|c_i| = |c_j| \implies p_i = p_j. \tag{14.4}$$

From equiprobability to the quadratic rule. Let $|c_k| = \sqrt{n_k/N}$ with integers $n_k \geq 0$ and $N := \sum_k n_k$ (rational fine–graining). Introduce an ancilla A' and a controlled isometry that *splits* each branch k into n_k equal–modulus subbranches,

$$|a_k\rangle|0\rangle_{A'} \longmapsto \frac{1}{\sqrt{n_k}} \sum_{m=1}^{n_k} |a_k\rangle|m_k\rangle_{A'}, \qquad |e_k\rangle \longmapsto |e_{km}\rangle \quad (\langle e_{km}|e_{\ell n}\rangle \approx \delta_{k\ell}\delta_{mn}), \tag{14.5}$$

which yields $|\Psi_{\text{fin}}\rangle \mapsto |\widetilde{\Psi}\rangle = \frac{1}{\sqrt{N}} \sum_{k=1}^{K} \sum_{m=1}^{n_k} |s_k\rangle|a_k\rangle|m_k\rangle_{A'}|e_{km}\rangle$. By (14.4), each of the N fine–grained branches has probability $1/N$, hence

$$p_k = \frac{n_k}{N} = |c_k|^2 \qquad \text{for rational } |c_k|^2. \tag{14.6}$$

Finally, continuity of f in (14.3) (guaranteed by the GNS/diagnostic realized reference configuration topology; cf. Section 4.6) extends (14.6) to all real amplitudes, fixing

$$p_k \;=\; f(|c_k|) \;=\; |c_k|^2. \tag{14.7}$$

Normalization, noncontextuality, and compatibility. Because $\sum_k P_k = \mathbf{1}$ on the pointer lane and $\operatorname{Tr}\rho = 1$ in the emergent inner product, $\sum_k p_k = \sum_k \operatorname{Tr}(\rho P_k) = 1$. Moreover, phases or swaps acting solely on E leave $\{p_k\}$ invariant (noncontextual with respect to environmental microstructure), and (14.7) coincides with the general projector valuation used elsewhere in this chapter (Sections 14.2 and 14.3.2). The derivation employs only (i) relational access, (ii) envariance symmetries on AE, (iii) GNS-emergent continuity/normalization, and (iv) orthogonality of the decoherence–selected pointer basis from Stage 1, and is therefore fully compatible with the cosmic, BRST–clean diagnostic realized reference configuration fixed by the symmetry stack and UQC (Section 13.4.1 and Theorem 13.168).

Stage 3: Uniqueness from the Dialectic Algebra

Synopsis. Stage 3 shows that the quadratic assignment fixed in Stage 2, $p_k = |c_k|^2$ (from eq. (14.43), cf. eq. (14.7)), is not merely admissible but *unique* once probabilities are required to respect the dialectic structure of IQR. Operationally, measurement is the passage from the Fourfold superpositional stratum (*Parasymmetry*) to the Twofold Boolean stratum singled out by the *Axiom of Duality* (decidable outcomes) (Section 4.1.6). A *probability map* is a valuation

$$v : \mathsf{Proj}(\mathcal{H}_{\text{em}}) \longrightarrow [0,1]$$

from orthogonal decompositions in the synthesis stratum (represented on the emergent/GNS arena; Section 4.6 and Definition 1.34) to $[0,1]$ that satisfies the axioms of Section 14.3.3:

(A1) Normalization: $v(0)=0, \quad v(\mathbf{1})=1.$ (A1)

(A2) Orthogonal additivity: $P_iP_j=0\ (i\neq j)\ \Rightarrow\ v\Big(\sum_i P_i\Big)=\sum_i v(P_i).$ (A2)

(A3) Unitary naturality: $v(UPU^\dagger)=v(P)\ \ \forall U\in\mathcal{U}(\mathcal{H}_{\mathrm{em}}).$ (A3)

(A4) Monoidal (independent composition): $v(P\otimes Q)=v(P)\,v(Q).$ (A4)

(A5) Coarse–graining noncontextuality: if $P=\sum_{i\in B}P_i$, then $v(P)=\sum_{i\in B}v(P_i).$ (A5)

(A6) Continuity: $P_n\to P\ \Rightarrow\ v(P_n)\to v(P).$ (A6)

These axioms are imposed *on the diagnostic realized reference configuration* selected by the symmetry stack (Section 13.4.1), using the emergent inner product (Section 4.6) and the UQC background (Theorem 13.168).

Gleason/Busch step in the emergent arena. On any emergent Hilbert fibre $\mathcal{H}_{\mathrm{em}}$ of dimension ≥ 3, axioms (A1)–(A3),(A6) imply (Gleason) the existence of a unique density operator ρ such that

$$v(P)=\mathrm{Tr}(\rho\,P)\qquad\text{for all projectors } P, \tag{14.8}$$

while for dim $=2$ the POVM extension (Busch) yields the same form on all effects, and hence on projectors [613, 355]. In our relational setting the trace and inner product are those induced by the Fourfold Inter–Modal Correlator (Ch. 4), rendering (14.8) a *GNS* statement (Section 4.6 and Definition 4.9). Therefore, for the decoherence–selected pointer projectors $P_k:=\mathbf{1}_S\otimes|a_k\rangle\langle a_k|\otimes\mathbf{1}_E$ of Stage 1 (Section 14.2.2),

$$p_k\ :=\ v(P_k)\ =\ \mathrm{Tr}(\rho\,P_k)\quad\text{(Section 14.3.2).} \tag{14.9}$$

Exponent–uniqueness under the monoidal law. Let the Stage 2 amplitudes be $c_k=\langle s_k,a_k|\Psi_{\mathrm{fin}}\rangle$ (cf. eq. (14.1)). Assume, as in Section 14.2.2, that p_k depends on $|c_k|$ only, i.e. $p_k=f(|c_k|)$, and satisfies: (i) equiprobability for equal–modulus branches, and (ii) the monoidal law (A4) on product states. Then

$$f(rs)=f(r)\,f(s),\qquad f(1)=1,\qquad f\Big(1/\sqrt{n}\Big)=1/n, \tag{14.10}$$

where the last condition follows from an n–way equal–modulus split (fine–graining) as in Section 14.2.2. Writing $f(r)=r^\alpha$ solves (14.10); inserting $r=1/\sqrt{n}$ gives

$$\frac{1}{n}=f\Big(\frac{1}{\sqrt{n}}\Big)=\Big(\frac{1}{\sqrt{n}}\Big)^{\alpha}\quad\Longrightarrow\quad\alpha=2.$$

Thus $p_k=f(|c_k|)=|c_k|^2$, i.e. the Born rule.

No–go for alternatives. Any exponent–deformed law $|c|^\alpha$ with $\alpha \neq 2$ violates either equiprobability for equal–modulus partitions or the monoidal rule (A4) upon coarse–graining (Theorem 14.3). Context–dependent assignments conflict with (A5) and, when combined with Page–Wootters conditioning (eq. (13.238)), generically spoil global conservation under the UQC (see Section 14.3.3; cf. [215]). Within the holosymmetric interferometer, such violations manifest as algebraic *twist* in the commutator lens and are suppressed by Stage II orthogonality (Definition 3.12 and Section 13.1.3).

Combining the Gleason/Busch representation (14.8), the Stage 2 envariance constraints, and the monoidal/exchange axioms (A1)–(A6) on the GNS diagnostic realized reference configuration, the only admissible valuation on pointer projectors is

$$p_k = \mathrm{Tr}(\rho\, P_k) = |c_k|^2,$$

i.e. the *Born rule* (Theorem 14.3). Equivalently, probabilities are uniquely the relational self–coherences of outcome branches, consistent with the dialectic passage Parasymmetry → Duality and with the BRST–clean, TT–safe cosmic background fixed by the symmetry stack and the UQC (Section 13.4.1 and Theorem 13.168).

14.3 A Formal Derivation of the Born Rule from First Principles

The objective is to demonstrate that for any normalized subsystem state $|\psi_S\rangle = \sum_i c_i|s_i\rangle$, the probability of obtaining the outcome associated with $|s_i\rangle$ is $p_i = |c_i|^2$. In the IQR framework, this is not a postulate but a necessary consequence of the relational, diagnostic realized reference configuration–certified dynamics encoded by the GNS/Holosymmetry construction and the Universal Quantum Constraint (UQC) (Definition 1.34, Section 4.6, and Theorem 13.168).

14.3.1 Stage 1: Measurement as Relational Entanglement and Decoherence

We model measurement as a concrete sequence of physical interactions—*preparation*, *local entanglement*, and *global decoherence*—implemented on a certified band/diagnostic realized reference configuration $\mathfrak{P}$ and placed canonically in the symmetry stack (Section 13.4.1). All inner products and norms below are those induced by the Fourfold Inter–Modal Correlator on the emergent/GNS arena (Ch. 4; Section 4.6 and Definition 4.9).

1. Preparation

Let the emergent Hilbert space factorize as $\mathcal{H}_{\mathrm{em}} = \mathcal{H}_S \otimes \mathcal{H}_A \otimes \mathcal{H}_E$ (cf. the GNS construction in Definitions 1.27 and 1.34). The prepared *separable* initial state is

$$|\Psi_{\mathrm{initial}}\rangle = \Big(\sum_i c_i\,|s_i\rangle\Big) \otimes |a_0\rangle \otimes |e_0\rangle, \qquad \sum_i |c_i|^2 = 1. \tag{14.11}$$

Here S/A are excitations of the Tantrakyma field $\hat{\mathbf{\Theta}}$ and E is the Gaiamiti (chronogeometric) environment $\hat{\mathbf{\Phi}}$ (Chapter 6 and Section 6.1). The typing (TT vs. τ lanes) ensures TT–purity

of spin–2 and a TT–safe clock sector, preventing spurious GW polarizations during readout (eqs. (10.4) and (13.115)).

2. Local Entanglement (The Measurement Interaction)

The system–apparatus interaction is generated by $\hat{H}_{SA}$, the first piece of the effective interaction Hamiltonian (Definition 14.1 and eqs. (14.16) and (14.17)). The unitary $U_{SA} := \mathcal{T}\exp\Big[-\frac{i}{\hbar}\int \hat{H}_{SA}\,dt\Big]$ implements a controlled–imprint map (cf. path–scattering in Ch. 3),

$$U_{SA}\Big(|s_i\rangle \otimes |a_0\rangle\Big) = |s_i\rangle \otimes |a_i\rangle, \qquad |\Psi_{\text{ent}}\rangle = (U_{SA}\otimes \mathbf{1}_E)|\Psi_{\text{initial}}\rangle = \sum_i c_i\,|s_i\rangle|a_i\rangle|e_0\rangle. \tag{14.12}$$

At this stage the SA sector is in a pure entangled superposition; coherence is preserved globally (UQC), merely redistributed (Theorem 13.168).

3. Global Decoherence (Entanglement with Geometry)

The apparatus couples to the chronogeometric environment via $\hat{H}_{AE}$, the second piece of (14.16) (eq. (14.18)), which descends from the bi–local interaction action

$$\mathcal{L}_{\text{int}}(x,x') := \text{Tr}\Big[\hat{\mathbf{\Phi}}^{\dagger}(x)\,\mathcal{U}_{\Phi\Theta}(x,x')\,\hat{\mathbf{\Theta}}(x')\Big] \quad \text{(Section 14.3.1)}. \tag{14.13}$$

The environment–imprinting unitary U_{AE} maps $|a_i\rangle|e_0\rangle \mapsto |a_i\rangle|E_i\rangle$, giving the tripartite state

$$|\Psi_{\text{fin}}\rangle = (\mathbf{1}_S \otimes U_{AE})\,|\Psi_{\text{ent}}\rangle = \sum_i c_i\,|s_i\rangle|a_i\rangle|E_i\rangle, \qquad \langle E_j|E_i\rangle \simeq \delta_{ij}. \tag{14.14}$$

Tracing out E yields the reduced state

$$\begin{aligned}\rho_{SA} &= \text{Tr}_E\Big(|\Psi_{\text{fin}}\rangle\langle\Psi_{\text{fin}}|\Big)\\ &= \sum_{i,j} c_i c_j^{*}\,\langle E_j|E_i\rangle\,|s_i\rangle\langle s_j|\otimes|a_i\rangle\langle a_j|\\ &\xrightarrow{\langle E_j|E_i\rangle\to\delta_{ij}} \sum_i |c_i|^2\,|s_i,a_i\rangle\langle s_i,a_i|.\end{aligned} \tag{14.15}$$

The *rate* and *robustness* of the suppression are quantified by the triad of relational distances on $\mathcal{X}_q$, $(d_{\text{corr}}, d_{\text{vac}}, d_{\text{spec}})$, whose functional independence governs distinguishability and coarse–grained stability (Section 4.11.5); increasing correlational distance drives $\langle E_j|E_i\rangle \to 0$.

Remarks (DRRC placement and clocks). All dynamics is executed on a BRST–clean, tag–stable diagnostic realized reference configuration (Definitions 10.1 and 12.1); boundary terms are calibrated as in Definition 13.155. When an internal good–clock lane C is used for time–labelled conditionals, the Page–Wootters equation $i\hbar\,\partial_t\,\varrho_{A|C=t} = [H_{\text{eff}}(t), \varrho_{A|C=t}]$ holds (Section 5.3.8 and eq. (13.238)), compatibly with the timeless UQC.

Formalizing the Interaction and Decoherence Model

From the bi–local interaction action (14.13) we extract an *effective* Hamiltonian on the *SAE* factorization that respects the stack placement (Section 13.4.1):

Definition 14.1 (Effective Interaction Hamiltonian for Measurement)**.** Let $\mathcal{H}_{\text{em}} = \mathcal{H}_S \otimes \mathcal{H}_A \otimes \mathcal{H}_E$. The effective interaction Hamiltonian is the sum

$$\boxed{\hat{H}_{\text{int}} = \hat{H}_{SA} + \hat{H}_{AE},} \tag{14.16}$$

with:

$$\boxed{\hat{H}_{SA} := g_{SA} \int d^d x\, \delta(x - x_A)\, \mathrm{Tr}\big[\hat{\mathbf{\Theta}}^\dagger_S(x)\, \hat{\mathbf{\Theta}}_A(x)\big],} \tag{14.17}$$

$$\boxed{\hat{H}_{AE} := g_{AE} \int d^d x'\, \mathrm{Tr}\big[\hat{\mathbf{\Phi}}^\dagger_E(x')\, \mathcal{U}_{\Phi\Theta}(x', x_A)\, \hat{\mathbf{\Theta}}_A(x_A)\big].} \tag{14.18}$$

Here g_{SA}, g_{AE} are diagnostic realized reference configuration–fixed couplings, $\mathcal{U}_{\Phi\Theta}$ the bi–local kernel descended from (14.13), and the typing guarantees TT–nullity of the clock lane and SM–compatibility of the apparatus (eq. (10.4) and Section 13.4.1).

Lemma 14.2 (Orthogonal records ⇒ diagonal pointer statistics)**.** *Under* (14.12)–(14.14) *(equivalently, the post–measurement form of eq.* (14.37)*) with* $\langle E_j|E_i\rangle \simeq \delta_{ij}$*, the apparatus pointer POVM (indeed a PVM)* $\{P_k := \mathbf{1}_S \otimes |a_k\rangle\langle a_k| \otimes \mathbf{1}_E\}$ *yields*

$$p_k := \mathrm{Tr}(\rho\, P_k) = \mathrm{Tr}_{SA}(P_k\, \rho_{SA}) = |c_k|^2, \qquad \rho = |\Psi_{\text{fin}}\rangle\langle\Psi_{\text{fin}}|. \tag{14.19}$$

Proof. By (14.14) (cf. eq. (14.37)),

$$|\Psi_{\text{fin}}\rangle = \sum_i c_i\, |s_i\rangle|a_i\rangle|E_i\rangle, \qquad \langle a_j|a_i\rangle = \delta_{ij}, \quad \langle E_j|E_i\rangle \simeq \delta_{ij},$$

where orthonormality of $\{|a_k\rangle\}$ follows from the imprint map in Stage 1 (cf. Section 14.3.1), and approximate orthogonality of the records $\{|E_k\rangle\}$ is the decoherence statement of Section 14.3.1. The global state $\rho = |\Psi_{\text{fin}}\rangle\langle\Psi_{\text{fin}}|$ expands as

$$\rho = \sum_{i,j} c_i c_j^* \Big(|s_i\rangle\langle s_j|\Big) \otimes \Big(|a_i\rangle\langle a_j|\Big) \otimes \Big(|E_i\rangle\langle E_j|\Big).$$

Taking the partial trace over the environment gives the reduced state on $S{+}A$,

$$\rho_{SA} := \mathrm{Tr}_E(\rho) = \sum_{i,j} c_i c_j^*\, \langle E_j|E_i\rangle \Big(|s_i\rangle\langle s_j|\Big) \otimes \Big(|a_i\rangle\langle a_j|\Big). \tag{14.20}$$

By record orthogonality, $\langle E_j|E_i\rangle \simeq \delta_{ij}$, the off–diagonal contributions ($i \neq j$) are suppressed, yielding the diagonal form asserted in eq. (14.15):

$$\rho_{SA} \simeq \sum_i |c_i|^2 \Big(|s_i\rangle\langle s_i|\Big) \otimes \Big(|a_i\rangle\langle a_i|\Big). \tag{14.21}$$

Now evaluate the pointer statistics for $P_k = \mathbf{1}_S \otimes |a_k\rangle\langle a_k| \otimes \mathbf{1}_E$:

$$p_k = \mathrm{Tr}(\rho P_k) = \mathrm{Tr}_{SA}\Big[(\mathbf{1}_S \otimes |a_k\rangle\langle a_k|)\,\rho_{SA}\Big] \simeq \sum_i |c_i|^2\, \mathrm{Tr}_{SA}\Big[(\mathbf{1}_S \otimes |a_k\rangle\langle a_k|)\Big(|s_i\rangle\langle s_i| \otimes |a_i\rangle\langle a_i|\Big)\Big].$$

Using $\langle a_k|a_i\rangle = \delta_{ki}$ and $\mathrm{Tr}(|s_i\rangle\langle s_i|) = 1$, only the $i = k$ summand survives, hence

$$p_k \simeq |c_k|^2.$$

Equivalently (and without taking a partial trace first), one may compute directly

$$\mathrm{Tr}(\rho P_k) = \sum_{i,j} c_i c_j^*\, \langle a_j|a_k\rangle\langle a_k|a_i\rangle\, \langle E_j|E_i\rangle\, \langle s_j|s_i\rangle = |c_k|^2,$$

by orthonormality of $\{|s_i\rangle\}$ and $\{|a_i\rangle\}$ and decoherence–induced orthogonality of $\{|E_i\rangle\}$ (cf. eq. (14.37)). This proves (14.19). □

Lemma 14.2 provides the *Stage 1* bridge from dynamics to *diagonal* pointer statistics on the emergent arena, expressed already as a Born valuation (Definition 1.18). In §14.2.2 we show that this valuation is *uniquely* fixed to $p_k = |c_k|^2$ by relational envariance and fine–graining; §14.2.2 then proves uniqueness from the Dialectic Algebra axioms.

The Evolution from a Separable to an Entangled State

The first act of measurement is a *unitary* imprint that maps an initially separable *SA* state to a perfectly correlated *SA* entangled state. In the stack/diagnostic realized reference configuration placement, this imprint is generated by a localized Tantrakyma–Tantrakyma interaction at the apparatus site x_A (Definition 14.1 and eqs. (14.16) and (14.17)), with all inner products taken in the emergent/GNS arena (Section 4.6).

1. Initial state and interaction Hamiltonian. Prepare the system in the superposition and the apparatus in its ready state $|a_0\rangle$,

$$|\Psi_{S+A,\mathrm{initial}}\rangle = \Big(\sum_i c_i\,|s_i\rangle\Big) \otimes |a_0\rangle = \sum_i c_i\,|s_i\rangle|a_0\rangle, \qquad \sum_i |c_i|^2 = 1. \tag{14.22}$$

The localized System–Apparatus coupling at x_A is the first summand in (14.16),

$$\hat{H}_{SA} := g_{SA} \int d^d x\, \delta(x - x_A)\, \mathrm{Tr}\Big[\hat{\mathbf{\Theta}}_S^\dagger(x)\, \hat{\mathbf{\Theta}}_A(x)\Big], \tag{14.23}$$

which descends, after mode truncation to the relevant pointer manifold, to an *effective* operator on $\mathcal{H}_S \otimes \mathcal{H}_A$ as described next.

2. Operator form and controlled imprint. To realize a one–shot imprint that is *unitary* and *perfectly correlating*, we use a block–diagonal (controlled) decomposition on $\mathcal{H}_S \otimes \mathcal{H}_A$:

$$\boxed{\hat{H}_{SA}^{\mathrm{eff}} = \sum_i |s_i\rangle\langle s_i| \;\otimes\; \hat{h}_i, \qquad \hat{h}_i = \frac{\Omega_i}{2}\Big(|a_i\rangle\langle a_0| + |a_0\rangle\langle a_i|\Big),} \tag{14.24}$$

where each $\hat{h}_i$ generates a Rabi ([614]) rotation within the two–dimensional apparatus subspace $\mathcal{A}_i = \mathrm{span}\{|a_0\rangle, |a_i\rangle\}$ and acts as the identity on $\mathcal{A}_i^{\perp}$. Self–adjointness of $\hat{H}_{SA}^{\mathrm{eff}}$ is manifest. Equivalently, one can specify the *exact* controlled–unitary without reference to a pulse by choosing, for each i, a Householder ([615]) map on $\mathcal{A}_i$,

$$\hat{U}_i := \mathbf{1}_A - 2\,|\eta_i\rangle\langle\eta_i|, \qquad |\eta_i\rangle := \frac{|a_0\rangle - |a_i\rangle}{\sqrt{2}}, \qquad \hat{U}_i\,|a_0\rangle = |a_i\rangle, \tag{14.25}$$

and setting $\hat{U}_{SA} := \sum_i |s_i\rangle\langle s_i| \otimes \hat{U}_i$ (eq. (14.12)).

What the Householder map is and why it implements a perfect imprint. On the apparatus lane, fix (for each outcome index i) the two–dimensional subspace $\mathcal{A}_i := \mathrm{span}\{|a_0\rangle, |a_i\rangle\}$ and its orthogonal complement $\mathcal{A}_i^{\perp}$. The *Householder map* ([615]) in eq. (14.25) is the rank–1 reflection

$$\hat{U}_i = \mathbf{1}_A - 2\,|\eta_i\rangle\langle\eta_i|, \qquad |\eta_i\rangle := \frac{|a_0\rangle - |a_i\rangle}{\sqrt{2}},$$

which acts as a reflection across the hyperplane $\{v : \langle\eta_i|v\rangle = 0\}$. Because $|\eta_i\rangle$ has support only in $\mathcal{A}_i$, one has $\hat{U}_i|_{\mathcal{A}_i^{\perp}} = \mathbf{1}$, so $\hat{U}_i$ is block–diagonal with respect to $\mathcal{A}_i \oplus \mathcal{A}_i^{\perp}$.

Basic properties (reflection, unitarity, involution). The projector $|\eta_i\rangle\langle\eta_i|$ is Hermitian and idempotent; hence $\hat{U}_i^{\dagger} = \hat{U}_i$ (self–adjoint) and $\hat{U}_i^2 = \mathbf{1}$ (involution). Consequently $\hat{U}_i$ is unitary. Introducing the orthonormal "bisector" vector $|b_i\rangle := (|a_0\rangle + |a_i\rangle)/\sqrt{2}$, the decomposition $|a_0\rangle = (|b_i\rangle + |\eta_i\rangle)/\sqrt{2}$ and $|a_i\rangle = (|b_i\rangle - |\eta_i\rangle)/\sqrt{2}$ shows that $\hat{U}_i$ flips the component along $|\eta_i\rangle$ while leaving $|b_i\rangle$ unchanged; therefore

$$\hat{U}_i|a_0\rangle = |a_0\rangle - 2\langle\eta_i|a_0\rangle\,|\eta_i\rangle = |a_i\rangle, \qquad \hat{U}_i|a_i\rangle = |a_0\rangle,$$

as stated in eq. (14.25). In the ordered basis $\{|a_0\rangle, |a_i\rangle\}$ one can compute explicitly

$$|\eta_i\rangle\langle\eta_i| = \tfrac{1}{2}\begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}, \quad \Rightarrow \quad \hat{U}_i = \mathbf{1} - 2|\eta_i\rangle\langle\eta_i| = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \sigma_x,$$

so $\hat{U}_i$ is precisely the swap on $\mathcal{A}_i$ and the identity on $\mathcal{A}_i^{\perp}$.

Controlled imprint on $S{+}A$ and one–shot entangling. Using the block–diagonal system control of eq. (14.24), the exact controlled–unitary is

$$\hat{U}_{SA} := \sum_i |s_i\rangle\langle s_i| \otimes \hat{U}_i,$$

which implements the *perfect imprint* in a single shot: for the pre–measurement superposition $\sum_i c_i|s_i\rangle|a_0\rangle$,

$$\hat{U}_{SA}\Big(\sum_i c_i|s_i\rangle|a_0\rangle\Big) = \sum_i c_i\,|s_i\rangle\,\hat{U}_i|a_0\rangle = \sum_i c_i\,|s_i\rangle|a_i\rangle,$$

i.e. the entangler action asserted in eq. (14.12). Thus the Householder choice realizes a unitary, perfectly correlating imprint without pulse timing.

Relation to the Rabi–pulse Hamiltonian. In the two–level subspace $\mathcal{A}_i$, the effective generator in eq. (14.24) reads $\hat{h}_i = \frac{\Omega_i}{2}\big(|a_i\rangle\langle a_0| + |a_0\rangle\langle a_i|\big) = \frac{\Omega_i}{2}\,\sigma_x$. Its time–evolution is $e^{-i\hat{h}_i\tau} = \cos\!\big(\frac{\Omega_i\tau}{2}\big)\,\mathbf{1} - i\sin\!\big(\frac{\Omega_i\tau}{2}\big)\sigma_x$. Choosing a π–pulse, $\Omega_i\tau = \pi$, gives $e^{-i\hat{h}_i\tau} = -i\,\sigma_x$, which coincides with $\hat{U}_i$ up to an overall phase (Rabi oscillation; [614]). Hence the Householder reflection is the exact, time–independent representative of the same imprint effected by a resonant π–pulse in eq. (14.24) ([615]).

3. Time evolution to the entangled state. Let $U_{SA}(t) := \exp\!\big(-\frac{i}{\hbar}t\,\hat{H}^{\text{eff}}_{SA}\big)$. Because the S–projectors $|s_i\rangle\langle s_i|$ are mutually orthogonal, U_{SA} factorizes into independent two–level rotations on the disjoint subspaces $|s_i\rangle \otimes \mathcal{A}_i$. Choosing *pulse areas* $\Omega_i\Delta t = \pi$ (a π–pulse in each $\mathcal{A}_i$) yields the exact imprint

$$U_{SA}(\Delta t)\,\Big(|s_i\rangle \otimes |a_0\rangle\Big) = |s_i\rangle \otimes |a_i\rangle \quad \Rightarrow \quad |\Psi_{S+A,\text{ent}}\rangle = U_{SA}(\Delta t)\,|\Psi_{S+A,\text{initial}}\rangle = \sum_i c_i\,|s_i\rangle|a_i\rangle, \tag{14.26}$$

which is the controlled imprint asserted in Section 14.3.1. If one prefers the impulsive limit, the short–time Dyson expansion gives

$$\begin{aligned} U_{SA}(\Delta t) &= \mathbf{1} - \tfrac{i}{\hbar}\,\hat{H}^{\text{eff}}_{SA}\Delta t + \mathcal{O}(\Delta t^2), \\ \Rightarrow \quad |\Psi_{S+A,\text{ent}}\rangle &= \sum_j c_j|s_j\rangle|a_0\rangle - \tfrac{i}{\hbar}\Delta t\sum_i c_i\,|s_i\rangle|a_i\rangle + \mathcal{O}(\Delta t^2), \end{aligned} \tag{14.27}$$

and tuning the calibrated *diagnostic realized reference configuration pulse* so that the first term is nulled (the π–pulse condition) recovers the exact map (14.26).

Calibration and orthogonality. The pointer set $\{|a_i\rangle\}$ is chosen orthonormal by construction (apparatus design on the visible lane; Section 13.4.1), ensuring unitarity of the total imprint (14.26). BRST cleanliness and typed placement guarantee that the imprint does not excite TT modes nor induce gauge artefacts on the diagnostic realized reference configuration (Definition 10.1 and eq. (10.4)). This completes Act I of the path–scattering template (Section 3.4) and prepares Act III (decoherence) where A couples to E via $\hat{H}_{AE}$ (eq. (14.18)) to produce orthogonal records and diagonal pointer statistics (eqs. (14.14) and (14.15) and Lemma 14.2).

The Evolution from Local to Global Entanglement

The second act of measurement is *decoherence*: the *SA* entanglement produced by the imprint (Section 14.3.1) blossoms into a *global*, tripartite *SAE* entanglement via the apparatus–environment coupling. Information is not destroyed; it is irreversibly *delocalized* across the chronogeometric network, consistent with the UQC and diagnostic realized reference configuration placement (Theorem 13.168, Definition 10.1, and Section 13.4.1).

1. State after local entanglement. After the *SA* imprint of Section 14.3.1, the universe is

$$|\Psi_{S+A,\text{ent}}\rangle \otimes |E_0\rangle = \Big(\sum_i c_i\,|s_i\rangle|a_i\rangle\Big) \otimes |E_0\rangle, \qquad \sum_i |c_i|^2 = 1, \tag{14.28}$$

where $|E_0\rangle \equiv |e_0\rangle$ denotes the environment's (Gaiamiti) diagnostic realized reference configuration state. The ensuing dynamics is driven by the apparatus–environment Hamiltonian from Definition 14.1,

$$\boxed{\hat{H}_{AE} := g_{AE} \int d^d x' \; \mathrm{Tr}\Big[\hat{\mathbf{\Phi}}^\dagger_E(x')\, \mathcal{U}_{\Phi\Theta}(x', x_A)\, \hat{\mathbf{\Theta}}_A(x_A)\Big],} \tag{14.29}$$

descended from the bi–local action (14.13).

2. Conditional action and environment records. Introduce the pointer projectors $\Pi^A_i := |a_i\rangle\langle a_i|$ and define the *compressed* environment operators (diagnostic realized reference configuration–wise conditional couplings)

$$\hat{K}_i := g_{AE} \int d^d x' \; \mathrm{Tr}\Big[\hat{\mathbf{\Phi}}^\dagger_E(x')\, \mathcal{U}_{\Phi\Theta}(x', x_A)\, \langle a_i|\hat{\mathbf{\Theta}}_A(x_A)|a_i\rangle\Big], \qquad \hat{K}_i : \mathcal{H}_E \to \mathcal{H}_E. \tag{14.30}$$

For a short decoherence time Δt (Dyson expansion), the environment *record* conditioned on $|a_i\rangle$ is

$$|E^{(1)}_i\rangle := -\frac{i}{\hbar}\, \Delta t\; \hat{K}_i\, |E_0\rangle, \tag{14.31}$$

and the corresponding (normalized) record is $|E_i\rangle := |E^{(1)}_i\rangle / \|E^{(1)}_i\|$ up to $\mathcal{O}(\Delta t^2)$ corrections. Physically, $|E_i\rangle$ is a chronogeometric excitation (a "record cloud") uniquely parameterized by the pointer outcome i.

3. Unitary evolution to the tripartite entangled state. Let $U_{AE}(\Delta t) := \exp\Big(-\frac{i}{\hbar}\Delta t\, \hat{H}_{AE}\Big) \approx \mathbf{1} - \frac{i}{\hbar}\Delta t\, \hat{H}_{AE}$. Acting on (14.28) and using the block–diagonal action relative to $\{\Pi^A_i\}$ gives, to leading order,

$$\begin{aligned} |\Psi_{\mathrm{fin}}\rangle &:= (\mathbf{1}_S \otimes U_{AE}(\Delta t)) \Big(\sum_j c_j\, |s_j\rangle|a_j\rangle \otimes |E_0\rangle\Big) \\ &\approx \sum_j c_j\, |s_j\rangle|a_j\rangle \otimes |E_0\rangle \;+\; \sum_j c_j\, |s_j\rangle|a_j\rangle \otimes |E^{(1)}_j\rangle \;\Longrightarrow\; \sum_i c_i\, |s_i\rangle|a_i\rangle|E_i\rangle, \end{aligned} \tag{14.32}$$

where the last arrow indicates re–normalization and the efficient–record limit. This is the tripartite form employed in eq. (14.14).

4. Orthogonality of environment records. Define the dephasing functional

$$\Gamma_{ij} := \langle E_j|E_i\rangle = \frac{\langle E_0|\hat{K}^\dagger_j \hat{K}_i|E_0\rangle}{\|\hat{K}_i|E_0\rangle\|\, \|\hat{K}_j|E_0\rangle\|} \;\equiv\; e^{-D_{ij}}, \tag{14.33}$$

with a decoherence "distance" $D_{ij} \geq 0$. In the triadic metric framework (Section 4.11.5), D_{ij} is an increasing function of the correlational, vacuum, and spectral separations of the two records,

$$D_{ij} \;=\; \alpha\, d_{\mathrm{corr}}(a_i, a_j)^2 + \beta\, d_{\mathrm{vac}}(a_i, a_j)^2 + \gamma\, d_{\mathrm{spec}}(a_i, a_j)^2 \quad (\alpha, \beta, \gamma > 0), \tag{14.34}$$

whose functional independence enforces near–orthogonality for macroscopically distinct pointers (Section 4.11.5). Hence, for $i \neq j$,

$$\langle E_j|E_i\rangle = \Gamma_{ij} \approx 0 \qquad \Rightarrow \qquad \langle E_j|E_i\rangle \approx \delta_{ij}. \tag{14.35}$$

Tracing out E then yields the diagonal *SA* state and pointer statistics (eq. (14.15) and Lemma 14.2),

$$\rho_{SA} = \sum_{i,j} c_i c_j^* \, |s_i\rangle\langle s_j| \otimes |a_i\rangle\langle a_j| \, \Gamma_{ij} \xrightarrow{\Gamma_{ij}\to\delta_{ij}} \sum_i |c_i|^2 \, |s_i, a_i\rangle\langle s_i, a_i|. \tag{14.36}$$

Solving the decoherence dynamics and the classical limit. With (14.29)–(14.36), the emergence of classicality is the quantitative statement that $\Gamma_{ij} \to 0$ for $i \neq j$ on certified plateaues, enforced by the geometry–matter kernel $\mathcal{U}_{\Phi\Theta}$ and the large–environment limit. A local observer, coupled only to *SA*, sees the reduced state (14.36); the off–diagonals (interference terms) are suppressed by (14.35). This completes Act II of the path–scattering template and provides the input for Stage 2 (relational envariance) (Section 14.2.2), from which the Born valuation $p_k = |c_k|^2$ is fixed (eq. (14.43)).

14.3.2 Stage 2: Probability from Symmetry and the Emergent Inner Product

Having prepared the orthogonal pointer basis in Stage 1 and the globally entangled post–measurement state

$$|\Psi_{\text{fin}}\rangle \;=\; \sum_k c_k \; |s_k\rangle \, |a_k\rangle \, |e_k\rangle \, , \qquad \langle a_k\rangle \, a_\ell = \delta_{k\ell}, \quad \langle e_k\rangle \, e_\ell \approx \delta_{k\ell}, \tag{14.37}$$

we now *derive* the outcome probabilities by appealing to the symmetries of $|\Psi_{\text{fin}}\rangle$—*environment–assisted invariance* (envariance)—and then identifying the result with the physically defined inner product of the IQR framework [158].

1. Probability as Relational Self–Coherence

Let $P_k := \mathbb{I}_S \otimes |a_k\rangle\langle a_k| \otimes \mathbb{I}_E$ be the pointer projector for outcome k. Any candidate probability assignment p_k must be a functional of the relational content of $|\Psi_{\text{fin}}\rangle$ accessible to an *SA* observer.

Within the holosymmetric/GNS framework, the squared norm is *not* primitive; it is *derived* from the Fourfold Inter–Modal Correlator $\hat{G}_\Lambda$ via the Holo–State expectation (coincident limit):

$$\| \, |\psi\rangle \, \|^2 \;=\; \langle\psi\rangle \, \psi \;:=\; \lim_{x' \to x} \Big\langle \mathring{\mathbf{\Psi}} \Big| \, \hat{G}_\Lambda\Big(\mathfrak{B}_x, \mathfrak{B}_{x'}\Big) \, [\psi^*, \psi] \, \Big| \mathring{\mathbf{\Psi}} \Big\rangle \quad \text{(eq. (4.85) and Definition 4.9).} \tag{14.38}$$

We will show below that the only probability functional consistent with the symmetries of (14.37) coincides with the projector expectation $p_k = \langle\Psi_{\text{fin}}|P_k|\Psi_{\text{fin}}\rangle$, hence, when evaluated via (14.38), with the squared coefficient $|c_k|^2$. Equivalently, in the GNS representation selected by the Holo–State, $p_k = \text{Tr}\Big(\rho \, P_k\Big)$ is the Born valuation (Definitions 1.18 and 1.34).

2. The Symmetry of Ignorance: Envariance for Equal Amplitudes

Phase envariance (phase indifference). For each k, apply on E a unitary $U_E^{(k)} : |e_k\rangle \mapsto e^{i\theta_k}|e_k\rangle$. This maps $c_k \mapsto c_k e^{-i\theta_k}$ in (14.37) but leaves all SA relations (and thus SA statistics) invariant; equivalently, a compensating unitary on S or A restores the global state. Hence any admissible probability functional must be *phase–insensitive*:

$$p_k \;=\; f\Big(|c_k|\Big). \tag{14.39}$$

Swap–envariance (equiprobability for equal moduli). Suppose $|c_j| = |c_k|$. The state is invariant under swapping the inaccessible environmental records, $U_{\text{swap}} : |e_j\rangle \leftrightarrow |e_k\rangle$, compensated by a swap on S or A. Because this operation produces no observable change for the SA observer, outcomes j and k are *indistinguishable* at the relational level. Therefore

$$\boxed{|c_j| = |c_k| \;\Rightarrow\; p_j = p_k} \quad \text{(equiprobability by envariant swaps).} \tag{14.40}$$

These two symmetries constitute the "symmetry of ignorance": local statistics cannot depend on unobservable environmental phases or labels [158].

3. Generalization to Arbitrary Amplitudes

Fine–graining to rational weights (ancilla splitting). Let the squared moduli be rational, $|c_k|^2 = r_k/N$ with integers $r_k \geq 0$, $\sum_k r_k = N$. Introduce an ancilla with orthonormal states $\{|\alpha_{k,\ell}\rangle : \ell = 1, \ldots, r_k\}$ and implement a unitary that "fans out" branch k into r_k subbranches of equal modulus:

$$|s_k\rangle\,|a_k\rangle\,|e_k\rangle\,|\alpha_0\rangle \;\longmapsto\; \frac{1}{\sqrt{r_k}} \sum_{\ell=1}^{r_k} |s_k\rangle\,|a_k\rangle\,|e_{k,\ell}\rangle\,|\alpha_{k,\ell}\rangle\,. \tag{14.41}$$

The global state becomes a uniform superposition of N mutually orthogonal, *equal–modulus* branches. By (14.4), each has probability $1/N$, and coarse–graining over the r_k subbranches of outcome k yields

$$p_k \;=\; \frac{r_k}{N} \;=\; |c_k|^2 \qquad \text{for all rational weights.} \tag{14.42}$$

Continuity. Because physical predictions cannot jump under arbitrarily small changes of amplitudes, and rationals are dense in $[0, 1]$, continuity extends (14.42) to *all* amplitudes:

$$\boxed{p_k \;=\; |c_k|^2 \;\text{ for all } k}\,. \tag{14.43}$$

4. Probability as a Projector Expectation and the Emergent Inner Product

The projector for outcome k is $P_k = \mathbb{I}_S \otimes |a_k\rangle\langle a_k| \otimes \mathbb{I}_E$. Acting on (14.37),

$$P_k\,|\Psi_{\text{fin}}\rangle = c_k\;|s_k\rangle\,|a_k\rangle\,|e_k\rangle\,, \tag{14.44}$$

hence

$$p_k \;=\; \langle \Psi_{\text{fin}}|\, P_k\, |\Psi_{\text{fin}}\rangle \;=\; |c_k|^2\, \langle s_k\rangle\, s_k\, \langle a_k\rangle\, a_k\, \langle e_k\rangle\, e_k \;=\; |c_k|^2, \tag{14.45}$$

using orthonormality of $\{|s_k\rangle\}$, $\{|a_k\rangle\}$ and decoherence–induced orthogonality of $\{|e_k\rangle\}$. In IQR, the equality $p_k = |c_k|^2$ is not merely algebraic: $|c_k|^2$ is evaluated by the *Emergent Inner Product* (14.38), i.e. as a *relational self–coherence* measured by the Fourfold Inter–Modal Correlator (Definition 6.4). Thus, the Born rule (14.43) says: *the probability of outcome k equals the self–coherence measure of the k–th branch of the universal state.*

Conclusion of Stage 2. Envariance (phase indifference, swap–symmetry), ancilla fine–graining, and continuity uniquely fix the functional dependence $p_k = |c_k|^2$; the projector computation (14.45) and the emergent inner product (14.38) identify this with the physically defined relational norm. Stage 3 will show that no alternative rule can satisfy the compositional and invariance requirements of the IQR/holosymmetric architecture (uniqueness theorem; Section 14.2.2).

14.3.3 Stage 3: Uniqueness from the Universal Constraint and the Dialectic

Having derived in Stage 2 that outcome probabilities must take the quadratic form $p_k = |c_k|^2$ (Section 14.3.2), we now show this rule is *unique* within the IQR/Holosymmetry architecture. The argument has two parts: (i) a structural uniqueness theorem under the admissibility axioms of Section 14.3.3, and (ii) a physical consistency check against the Universal Quantum Constraint and emergent time in the Page–Wootters sense [616].

The Consistency Condition: Local Dynamics from a Timeless Global State

Global constraint and emergent time. (I) The *Universal Quantum Constraint* fixes a single timeless *Holo–State* $|\mathring{\mathbf{\Psi}}\rangle \in \mathcal{H}_{\text{em}}$ with $\hat{H}_{\text{univ}}\,|\mathring{\mathbf{\Psi}}\rangle = 0$, encoding exact conservation of the associated Noether charges (Theorem 13.168, eq. (13.236), and Definition 13.167). (II) *Relational time (Page–Wootters)*: for any subsystem S and clock C, the conditional states $\varrho_{S|C=t}$ arise from correlations inside $|\mathring{\mathbf{\Psi}}\rangle$ and evolve unitarily with respect to t (cf. eq. (13.238) and Section 13.6.2 and [616]). On certified plateaux, this conditional dynamics *descends* to the BRST-physical diagnostic realized reference configuration $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$ via the Ch. 12 equivalence.

Any admissible probability assignment must therefore (a) be implementation–invariant and unitary–covariant on the emergent/GNS arena (and hence on $\mathcal{H}_{\text{phys}}$ where applicable), (b) compose on tensor products, and (c) reproduce linear, conserved expectations $\langle O\rangle = \text{Tr}(\rho\, O)$ as Born valuations that descend from $|\mathring{\mathbf{\Psi}}\rangle$ (Ward/Noether structure; see Section 14.3.2).

Axioms for an Admissible Probability Functional

Let p assign to each projector P and unit vector $|\psi\rangle$ a number $p(P|\psi) \in [0,1]$ subject to:

1. **Normalization and finite additivity** (*prismatic additivity*). For orthogonal $\{P_i\}$ with $\sum_i P_i = \mathbb{I}$: $p(P_i|\psi) \geq 0$ and $\sum_i p(P_i|\psi) = 1$.

2. **Unitary covariance** (*holosymmetric invariance*). For all unitaries U, $p(UPU^\dagger|U\psi) = p(P|\psi)$.

3. **Product factorization** (*independent composition*). If $|\psi\rangle = |\psi_A\rangle\otimes|\psi_B\rangle$ and $P = P_A\otimes P_B$, then $p(P|\psi) = p(P_A|\psi_A)\,p(P_B|\psi_B)$.

4. **Coarse–graining noncontextuality**. If $P = \sum_{i\in I} P_i$ with $P_iP_j = \delta_{ij}P_i$, then $p(P|\psi) = \sum_{i\in I} p(P_i|\psi)$.

5. **Continuity**. Small changes of amplitudes produce small changes of p.

Items (1)–(2) encode the Stage 2 symmetries (phase–indifference and envariant equiprobability), (3) matches the tensorial/BRST structure, (4) implements measurement coarse–graining, and (5) is the physical regularity hypothesis already used in Section 14.3.2.

The structural theorem.

Theorem 14.3 (Uniqueness of the quadratic valuation (holosymmetric/GNS))**.** Scope and setting. *All statements are made on the emergent/GNS space* $\mathcal{H}_{\rm em}$ *selected by the Holo–State* $|\mathring{\mathbf{\Psi}}\rangle$*; by the Ch. 12 equivalence they* descend *to the BRST–physical diagnostic realized reference configuration* $\mathcal{H}_{\rm phys} = H^0(\hat{Q}_{\rm BRST})$ *on certified plateaux. In infinite dimensions, axiom* (1) *of Section 14.3.3 is read as* σ–additivity *over countable orthogonal families (cf. Remark 14.5); with only finite additivity, the finite–rank case follows directly and the general case follows by norm–continuity (axiom* (5)*).*

Let p *satisfy axioms* (1)–(5) *of Section 14.3.3.*

(A) *(*Projective/Gleason form, $\dim\mathcal{H}_{\rm em} \geq 3$*.) For each unit* $|\psi\rangle \in \mathcal{H}_{\rm em}$ *there exists a density operator* ρ_ψ *such that*

$$p(P \mid \psi) = \mathrm{Tr}\big(\rho_\psi P\big) \qquad \text{for all projectors } P, \tag{14.46}$$

and unitary covariance (axiom (2)*) enforces* $\rho_\psi = |\psi\rangle\langle\psi|$*. Hence*

$$\boxed{p(P \mid \psi) = \langle\psi|\,P\,|\psi\rangle\,, \qquad \text{and for rank-1 } P = |\phi\rangle\langle\phi|\,, \quad p\Big(\,|\phi\rangle\langle\phi| \mid \psi\Big) = \Big|\langle\phi|\,\psi\Big|^2.} \tag{14.47}$$

(B) *(*Qubit case and Busch extension.*) In* $d = 2$ *the same conclusion follows either by applying* (A) *to two copies (*$d = 4$*) and invoking product factorization (axiom* (3)*), or—if the axioms are posed on POVMs with* σ*–additivity over effects (Busch hypothesis)—directly in* $d = 2$*.*

Proof idea. By Gleason/Busch, a (countably) additive, basis–noninvasive assignment on projectors yields (14.46) for $d \geq 3$ (see [613, 355]). Unitary covariance gives the conjugation law $\rho_{U\psi} = U\rho_\psi U^\dagger$. The Stage 2 *eigenstate–certainty* conclusion (phase–indifference and envariant swaps; cf. eq. (14.43)) implies $p(P_\psi \mid \psi) = 1$ for $P_\psi = |\psi\rangle\langle\psi|$, hence ρ_ψ has unit support on $\mathrm{span}\{\psi\}$ and $\rho_\psi = |\psi\rangle\langle\psi|$. This yields (14.47). For $d = 2$, use doubling + factorization, or adopt the Busch POVM hypothesis.

Proof. **Step 0 (setting and axioms).** Let p be a probability assignment on projectors on $\mathcal{H}_{\text{em}}$ satisfying (1)–(5) of Section 14.3.3: (normalization and (countable) additivity on orthogonal families), unitary covariance, product factorization, coarse–graining noncontextuality, and continuity.

Step 1 (frame–function representation for $d \geq 3$). By the Gleason/Busch route in complex dimension ≥ 3, there exists a unique density operator ρ_ψ such that (14.46) holds for all projectors P (see [613, 355]). Positivity and normalization of p entail $\rho_\psi \geq 0$ and $\mathrm{Tr}(\rho_\psi) = 1$.

Step 2 (covariance ⇒ conjugation law). Axiom (2) gives $p(UPU^\dagger \mid U\psi) = p(P \mid \psi)$. Using (14.46),

$$\mathrm{Tr}(\rho_{U\psi}\, UPU^\dagger) = \mathrm{Tr}(\rho_\psi P) \qquad \forall\, P.$$

With $P' = UPU^\dagger$ arbitrary, the separating family of projectors yields

$$\rho_{U\psi} = U\rho_\psi U^\dagger. \tag{14.48}$$

Step 3 (eigenstate certainty ⇒ purity). Let $P_\psi := |\psi\rangle\langle\psi|$. The Stage 2 branch law (equal–modulus refinement with $|c_1| = 1$) gives

$$p(P_\psi \mid \psi) = 1 \tag{14.49}$$

(cf. eq. (14.43) and the projector computation eq. (14.45)). Then

$$1 = p(P_\psi \mid \psi) = \mathrm{Tr}(\rho_\psi P_\psi) = \langle\psi|\, \rho_\psi\, |\psi\rangle\,.$$

Since $\rho_\psi \geq 0$ and $\mathrm{Tr}(\rho_\psi) = 1$, the spectral theorem implies that $|\psi\rangle$ is an eigenvector of ρ_ψ with eigenvalue 1, and all remaining eigenvalues sum to 0, hence vanish. Therefore

$$\rho_\psi = |\psi\rangle\langle\psi|\,. \tag{14.50}$$

Step 4 (Born form for all projectors). Combining (14.46) with (14.50) gives

$$p(P \mid \psi) = \mathrm{Tr}(|\psi\rangle\langle\psi|\ P) = \langle\psi|\, P\, |\psi\rangle \qquad \forall\, P,$$

and for rank–1 $P = |\phi\rangle\langle\phi|$, $p(|\phi\rangle\langle\phi| \mid \psi) = |\,\langle\phi\rangle\,\psi|^2$, i.e. (14.47).

Step 5 (the qubit case $d = 2$). Let $\dim \mathcal{H}_{\text{em}} = 2$ and consider $\mathcal{H}_{\text{em}}^{\otimes 2}$ (dimension 4). For any projector P and unit $|\psi\rangle$, apply the $d \geq 3$ result to $P \otimes \mathbb{I}$ and $|\psi\rangle \otimes |\psi\rangle$:

$$p(P \otimes \mathbb{I} \mid \psi \otimes \psi) = \langle\psi \otimes \psi|\, (P \otimes \mathbb{I})\, |\psi \otimes \psi\rangle = \langle\psi|\, P\, |\psi\rangle\,.$$

By product factorization (axiom (3)), $p(P \otimes \mathbb{I} \mid \psi \otimes \psi) = p(P \mid \psi)\, p(\mathbb{I} \mid \psi) = p(P \mid \psi)$, so $p(P \mid \psi) = \langle\psi|\, P\, |\psi\rangle$ on the qubit. Alternatively, if the axioms are posed on POVMs with σ–additivity over effects (Busch), the conclusion holds directly in $d = 2$.

Conclusion and descent. Thus (14.47) holds on $\mathcal{H}_{\text{em}}$; by the Ch. 12 equivalence, the same Born valuation governs projectors on $\mathcal{H}_{\text{phys}}$ on certified plateaux. □

Structural content of Theorem 14.3. The theorem Theorem 14.3 (*Uniqueness of the quadratic valuation, holosymmetric/GNS*) is the Stage–3 structural uniqueness result—the Gleason/Busch–style frame–function theorem adapted to the emergent/GNS arena. Its hypotheses are axioms (1)–(5) of Section 14.3.3 for a probability assignment $p(P \mid \psi)$ on projectors P of the emergent space $\mathcal{H}_{\mathrm{em}}$ (with descent to $\mathcal{H}_{\mathrm{phys}}$ on certified plateaux per Ch. 12): normalization and (countable) additivity on orthogonal families, unitary covariance, product factorization, coarse–graining noncontextuality, and continuity. Under these assumptions, on any complex Hilbert space with $\dim \mathcal{H}_{\mathrm{em}} \geq 3$ the theorem first forces the density–operator (frame–function) form

$$p(P \mid \psi) = \mathrm{Tr}(\rho_\psi P) \quad \text{for all projectors } P,$$

as stated in eq. (14.46). Unitary covariance then fixes $\rho_\psi = |\psi\rangle\langle\psi|$, yielding the Born rule

$$p(P \mid \psi) = \langle\psi| P |\psi\rangle\,, \qquad p(|\phi\rangle\langle\phi| \mid \psi) = \left|\langle\phi|\,\psi\right|^2$$

for rank–1 projectors, as boxed in eq. (14.47). In the qubit case $d = 2$, the same conclusion follows either by passing to two copies ($d = 4$) and invoking product factorization (axiom (3)), or—if the axioms are posed on POVMs with σ–additivity over effects (Busch hypothesis)—directly in $d = 2$. See also Remark 14.5 for the σ–additivity reading in infinite dimensions.

Relation to the headline result Theorem 14.4. This structural theorem differs in presentation from the concise headline statement Theorem 14.4 (*Uniqueness of the Born valuation*), which asserts directly—under the same axioms (1)–(5)—that $p(P \mid \psi) = \langle\psi| P |\psi\rangle$ for all projectors. The two theorems share the same conclusion but take distinct routes: Theorem 14.4 serves as a compact corollary, whereas Theorem 14.3 makes explicit the intermediate density–operator representation eq. (14.46) and the dimension clause before deriving the pure–state form eq. (14.47).

Why "quadratic valuation". The terminology underscores that, for rank–1 projectors, $p(|\phi\rangle\langle\phi| \mid \psi) = \left|\langle\phi|\,\psi\right|^2$, so probabilities are quadratic in amplitudes. Equivalently, for pointer branches with coefficients c_k one obtains $p_k = |c_k|^2$, consistent with the Stage–2 derivation based on envariance, fine–graining, and continuity (Section 14.3.2 and eq. (14.43)).

Placement in the overall derivation. Stage–2 fixes the quadratic law for pointer outcomes, $p_k = |c_k|^2$; the structural theorem Theorem 14.3 then lifts this branch–level result to *all* projectors via the frame–function route, establishing the Born rule in full generality on $\mathcal{H}_{\mathrm{em}}$. By the Ch. 12 equivalence, the same valuation descends to $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ on certified plateaux. The compact statement Theorem 14.4 records the conclusion succinctly for later use.

Exponent–deformations are excluded. Suppose a rank–1 rule depended only on the modulus, $p(|\phi\rangle\langle\phi| \mid \psi) = F\big(|\langle\phi|\,\psi\rangle|\big)$. Envariant equiprobability for N equal–modulus branches

(Section 14.3.2) enforces $F(1/\sqrt{N}) = 1/N$. Independent composition (axiom (3)) implies $F(rs) = F(r)F(s)$, hence by continuity (axiom (5)) $F(r) = r^\alpha$; the previous constraint fixes $\alpha = 2$, i.e. $F(r) = r^2$. Any $\alpha \neq 2$ conflicts with (1) and (3) under refinement/coarse–graining of orthogonal resolutions of the identity (and with (4) noncontextuality).

Physical consistency with the UQC. Let p deviate from eq. (14.47). Then there is no density operator ϱ with $p(P \mid \psi) = \mathrm{Tr}(\varrho P)$ for all projectors P; consequently, expectations of conserved charges cannot in general be written as $\mathrm{Tr}(\varrho\, Q)$. In the emergent/GNS arena selected by the Holo–State $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$, the Page–Wootters conditional dynamics reads

$$i\hbar\, \partial_t\, \varrho_{A|C=t} = \Big[\, H_{\mathrm{eff}}(t),\ \varrho_{A|C=t} \,\Big] \qquad \text{(cf. eq. (13.238)).}$$

Noether/Ward conservation then implies $\partial_t\, \mathrm{Tr}\Big(\varrho_{A|C=t}\, Q\Big) = 0$ for any conserved charge Q, whereas a nonlinear p produces $\partial_t\, \mathbb{E}_p[Q] \neq 0$ in general, violating the universal conservation encoded by

$$\hat{H}_{\mathrm{univ}}\, |\overset{\circ}{\boldsymbol{\Psi}}\rangle = 0 \qquad \text{(Theorem 13.168, eq. (13.236), and Definition 13.167).}$$

By the Holosymmetric Selection Principle, such histories are excluded at Stage I/II.

Within the dialectic passage *Parasymmetry* $\to$ *Duality* and under axioms (1)–(5), the only admissible valuation is the Born rule eq. (14.47). Equivalently, probabilities are uniquely the *relational self–coherences* of outcome branches, as already obtained in eq. (14.43).

Theorem 14.4 (Uniqueness of the Born valuation (holosymmetric/GNS))**.** *Let p satisfy axioms* (1)–(5) *of Section 14.3.3.*

(A) *(*Projective/Gleason form, $d \geq 3$.*) On the emergent/GNS space $\mathcal{H}_{\mathrm{em}}$ with* $\dim \mathcal{H}_{\mathrm{em}} \geq 3$*, there exists for each unit $|\psi\rangle$ a density operator ρ_ψ such that $p(P \mid \psi) = \mathrm{Tr}(\rho_\psi P)$ for all projectors P. Unitary covariance enforces $\rho_\psi = |\psi\rangle\langle\psi|$, hence*

$$p(P \mid \psi) = \langle\psi|\, P\, |\psi\rangle \quad (\textit{Born rule}).$$

By the Ch. 12 equivalence, this statement descends *to $\mathcal{H}_{\mathrm{phys}} = H^0(\hat{Q}_{\mathrm{BRST}})$ on certified plateaux.*

(B) *(*Busch extension, all *d.) If, in addition, the axioms are posed on* POVMs *(–additivity over effects), then the same conclusion holds in all finite dimensions, including $d = 2$.*

In $d = 2$ without the POVM axiom, the conclusion follows by applying (A) to two copies ($d = 4$) and invoking product factorization (3)*.*

Proof idea. By Stage 2 (Section 14.3.2), phase indifference and envariant swaps imply equiprobability for equal–modulus branches (14.4), and fine–graining with continuity yields a modulus–only dependence on rank–1 projectors, $p_k = f(|c_k|)$. Product factorization (axiom (3)) gives functional multiplicativity $f(rs) = f(r)f(s)$ for $r, s \in [0, 1]$, so by continuity (axiom (5)) $f(r) = r^\alpha$. Equiprobability for n equal–modulus branches fixes $\alpha = 2$, hence

$p_k = |c_k|^2$ (agreeing with (14.43)). –additivity and coarse–graining noncontextuality (axioms (1),(4)) extend from rank–1 to arbitrary projectors, and unitary covariance (axiom (2)) fixes the representation to the quadratic form Theorem 14.3. This is the relational/GNS version of the Gleason–Busch route.

Proof. **Step 1 (reduction to rank–1 branches via Stage 2).** In the setting of eq. (14.37), with pointer projectors $P_k := \mathbb{I}_S \otimes |a_k\rangle\langle a_k| \otimes \mathbb{I}_E$, Stage 2 implies any admissible assignment depends only on branch modulus and is equiprobable for equal moduli (Section 14.3.2, eq. (14.39), eq. (14.4)). Ancilla fine–graining and continuity give, for rank–1 pointer events,

$$p_k \;=\; f(|c_k|) \;=\; |c_k|^2, \qquad \text{as in Section 14.3.2, eq. (14.42), eq. (14.43).} \tag{14.51}$$

Step 2 (multiplicativity from product factorization). Define $f : [0,1] \to [0,1]$ by $p(|\phi\rangle\langle\phi| \mid \psi) = f(|\langle\phi|\,\psi\rangle|)$ for rank–1 projectors. For independent systems A, B, axiom (3) yields

$$p\Big((|\phi_A\rangle\langle\phi_A|)\otimes(|\phi_B\rangle\langle\phi_B|)\,\Big|\,\psi_A\otimes\psi_B\Big) = p(|\phi_A\rangle\langle\phi_A| \mid \psi_A)\, p(|\phi_B\rangle\langle\phi_B| \mid \psi_B),$$

so with $r := |\langle\phi_A|\,\psi_A\rangle|$, $s := |\langle\phi_B|\,\psi_B\rangle|$,

$$f(rs) = f(r)\, f(s) \qquad (r, s \in [0,1]). \tag{14.52}$$

By continuity (axiom (5)), $f(r) = r^\alpha$ for some $\alpha > 0$.

Step 3 (fixing the exponent by envariant equiprobability). For a decomposition into n equal–modulus branches, $p = 1/n$ (swap–envariance; eq. (14.4)). Taking $r = 1/\sqrt{n}$ gives

$$\frac{1}{n} = f\Big(\frac{1}{\sqrt{n}}\Big) = \Big(\frac{1}{\sqrt{n}}\Big)^{\alpha} \quad \Rightarrow \quad \alpha = 2,$$

hence for rank–1 events

$$p(|\phi\rangle\langle\phi| \mid \psi) = |\langle\phi|\,\psi\rangle|^2. \tag{14.53}$$

Step 4 (extension from rank–1 to arbitrary projectors). Let $P = \sum_j |\phi_j\rangle\langle\phi_j|$ be an orthogonal spectral resolution (finite or countable). By –additivity and coarse–graining noncontextuality (axioms (1),(4)),

$$p(P \mid \psi) = \sum_j p(|\phi_j\rangle\langle\phi_j| \mid \psi) = \sum_j |\langle\phi_j|\,\psi\rangle|^2 = \langle\psi|\,\Big(\sum_j |\phi_j\rangle\langle\phi_j|\Big)\,|\psi\rangle = \langle\psi|\,P\,|\psi\rangle\,,$$

using (14.53) termwise. (If only finite additivity is assumed, the statement holds verbatim for finite–rank P.)

Step 5 (unitary covariance = implementation invariance). Axiom (2) ensures basis–independence: $p(P \mid \psi)$ depends only on P and $|\psi\rangle$, matching $\langle\psi|\,P\,|\psi\rangle$.

All steps are taken in the emergent/GNS arena $\mathcal{H}_{\text{em}}$; by the Ch. 12 equivalence, the result descends to $\mathcal{H}_{\text{phys}} = H^0(\hat{Q}_{\text{BRST}})$ on certified plateaux. Hence Theorem 14.3 holds. □

Remark 14.5 (On σ–additivity and frame functions)**.** In infinite–dimensional settings on the emergent/GNS space $\mathcal{H}_{\mathrm{em}}$, axiom (1) of Section 14.3.3 is to be read as *σ–additivity over countable mutually orthogonal families* of projectors. With coarse–graining noncontextuality (axiom (4)), this makes $P \mapsto p(P \mid \psi)$ a countably additive measure on the projection lattice and, restricted to rank–1 resolutions, a frame function of weight 1. Combined with the rank–1 law (14.53), this yields the quadratic form $p(P \mid \psi) = \langle\psi| P |\psi\rangle$ for all projectors P, i.e. Theorem 14.3. If only *finite additivity* is assumed, (14.3) is guaranteed for finite–rank P; the general case then follows by norm–continuity (axiom (5)) from finite–rank approximants. When the axioms are posed on *POVMs* (Busch–type σ–additivity over effects), the same conclusion holds in all finite dimensions, including $d = 2$. All statements descend to the BRST–physical diagnostic realized reference configuration $\mathcal{H}_{\mathrm{phys}}$ on certified plateaux via the Ch. 12 equivalence.

Remark (Relation to classical results). The derivation above recovers the Gleason/Busch uniqueness in the relational, emergent/GNS setting. For projectors on $\mathcal{H}_{\mathrm{em}}$ with $\dim \mathcal{H}_{\mathrm{em}} \geq 3$, Steps 2–4 reproduce the Gleason form; with POVM σ–additivity (Busch hypothesis) the conclusion holds in all finite dimensions, including $d = 2$. The qubit case can thus be handled either by the Busch route or, as done here, by the two–copy lift to $d = 4$ plus product factorization (axiom (3)), with Stage 2 envariance fixing the rank–1 modulus law (14.53). Measure–theoretic overhead is confined to the σ–additivity clause stated in Remark 14.5. By the Ch. 12 equivalence, the result descends to the BRST–physical diagnostic realized reference configuration $\mathcal{H}_{\mathrm{phys}}$ on certified plateaux.

A No–Go Illustration: Incompatibility of Alternative Rules with the Global Constraint

Consider the Page–Wootters setting (emergent time) with a two–level system S correlated with a clock C:

$$|\overset{\circ}{\boldsymbol{\Psi}}\rangle \;=\; \sqrt{\tfrac{1}{5}}\,|E_1\rangle_S\,|t_1\rangle_C \;+\; \sqrt{\tfrac{4}{5}}\,|E_2\rangle_S\,|t_2\rangle_C\,, \qquad \hat{H}_{\mathrm{univ}}\,|\overset{\circ}{\boldsymbol{\Psi}}\rangle = 0, \tag{14.54}$$

so that conditional S–dynamics is unitary in t (cf. eq. (13.238) and [616]). Born probabilities give

$$\langle E\rangle_{\mathrm{Born}} = \sum_{k=1}^{2} |c_k|^2 E_k = \tfrac{1}{5}E_1 + \tfrac{4}{5}E_2, \tag{14.55}$$

which matches the conserved global expectation inherited from $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$ by the Universal Quantum Constraint (Theorem 13.168).

Suppose instead a modulus–deformed rule $p_k^{(\alpha)} \propto |c_k|^\alpha$ were used locally. Normalizing,

$$p_k^{(\alpha)} \;=\; \frac{|c_k|^\alpha}{\sum_j |c_j|^\alpha}\,, \qquad \alpha \neq 2. \tag{14.56}$$

For the "linear–modulus" case $\alpha = 1$, (14.54) gives $p_1^{(1)} = \frac{\sqrt{1/5}}{\sqrt{1/5}+\sqrt{4/5}} = \frac{1}{3}$ and $p_2^{(1)} = \frac{2}{3}$, hence

$$\langle E\rangle_{\alpha=1} = \tfrac{1}{3}E_1 + \tfrac{2}{3}E_2 \;\neq\; \langle E\rangle_{\mathrm{Born}}. \tag{14.57}$$

More generally, for any conserved observable $Q = \sum_k q_k \, |E_k\rangle\langle E_k|$ diagonal in the branch basis, $\alpha \neq 2$ yields

$$\sum_k q_k \, p_k^{(\alpha)} \;\neq\; \sum_k q_k \, |c_k|^2,$$

which violates the linear conservation implied by $\hat{H}_{\text{univ}} \, |\overset{\circ}{\mathbf{\Psi}}\rangle = 0$ when local time is read relationally (cf. Section 14.3.3). By the Holosymmetric Selection Principle, such histories fail Stage I/II viability and are excluded. The quadratic valuation (14.47) is both *structurally* forced by the admissibility axioms and *physically* required by consistency with the Universal Quantum Constraint and Page–Wootters emergent time [616].

Holosymmetric Selection as Physical Arbiter

Holosymmetry provides the *dynamical* enforcement of the structural results proved in Stages 2–3. Any putative probability rule p_k that violates (i) the Stage 2 symmetries (Section 14.3.2), (ii) the tensorial/dialectic axioms (1)–(5) of Section 14.3.3, or (iii) the global Universal Quantum Constraint (Theorem 13.168) manifests as an *algebraic twist* in the interferometer/lensing diagnostics of Section 13.2.7. Concretely, let

$$p_k^{(\alpha)} \;=\; \frac{|c_k|^\alpha}{\sum_j |c_j|^\alpha}, \qquad \alpha \in (0, \infty), \tag{14.58}$$

be a generic modulus–deformation (Born corresponds to $\alpha = 2$), and denote the post–measurement projector $P_k := \mathbb{I}_S \otimes |a_k\rangle\langle a_k| \otimes \mathbb{I}_E$. Define the *twist operator* on the decohered branch algebra by

$$\mathcal{T}_\alpha[\Psi_{\text{fin}}] \;:=\; \sum_k \Big(p_k^{(\alpha)} - \langle \Psi_{\text{fin}}| \, P_k \, |\Psi_{\text{fin}}\rangle \Big) \, P_k \;=\; \sum_k \Big(p_k^{(\alpha)} - |c_k|^2 \Big) \, P_k, \tag{14.59}$$

where the last equality used (14.45). In the holosymmetric interferometer the Stage II metric ("lensing") penalizes such twists via a coherence functional (schematically)

$$\mathcal{L}_{\text{HS}}[\Psi_{\text{fin}}] \;=\; 1 - \frac{\big\| \mathcal{P}^{\text{coh}} \, \rho_{SA} \, \mathcal{P}^{\text{coh}} \big\|_2^2}{\|\rho_{SA}\|_2^2} \;\propto\; \big\| \mathcal{T}_\alpha[\Psi_{\text{fin}}] \big\|_2^2, \qquad \rho_{SA} = \text{Tr}_E \, |\Psi_{\text{fin}}\rangle\langle \Psi_{\text{fin}}|, \tag{14.60}$$

(cf. Section 13.2.7 for the role of the coherence projector $\mathcal{P}^{\text{coh}}$ and the Frobenius norm $\| \cdot \|_2$). Hence, *only* the Born assignment ($\alpha = 2$) makes $\mathcal{T}_\alpha \equiv 0$ for all $|\Psi_{\text{fin}}\rangle$, eliminating lensing twist and surviving Stage II selection as a "bright fringe." Equivalently, the only probability rule compatible with (i) Stage 2 envariance, (ii) the IQR tensorial/dialectic structure, and (iii) the global constraint is the Born rule.

Theorem 14.6 (Relational Coherence–Probability Equivalence (Uniqueness))**.** *Let a measurement on S with apparatus A and environment E produce the decohered state $|\Psi_{\text{fin}}\rangle = \sum_k c_k \, |s_k\rangle \, |a_k\rangle \, |e_k\rangle$ with orthogonal $\{|a_k\rangle\}$ and $\{|e_k\rangle\}$ (eq. (14.37)). Then the unique probability assignment consistent with (1)–(5) of Section 14.3.3 and with the Universal Quantum Constraint is*

$$p_k \;=\; \langle \Psi_{\text{fin}}| \, P_k \, |\Psi_{\text{fin}}\rangle \;=\; |c_k|^2, \tag{14.61}$$

where the right–hand side is evaluated by the Emergent Inner Product (14.38) (see Section 14.3.2). Thus, probabilities are precisely the relational self–coherences *of the outcome branches.*

Proof idea. Stage 2 (Section 14.3.2) fixes p on rank–1 projectors to $|c|^2$; Theorem 14.3 extends this uniquely to all projectors under (1)–(5). The Page–Wootters consistency check (Section 14.3.3) excludes $\alpha \neq 2$ by mismatch with conserved global expectations enforced by $\hat{H}_{\text{univ}} |\Psi_0\rangle = 0$ (Theorem 13.168). Finally, Holosymmetric Selection suppresses any residual non–Born proposal dynamically via the lensing penalty (14.60), which vanishes identically only for (14.61).

Proof. **Step 1 (Diagonalization of pointer statistics by decoherence).** Let $P_k := \mathbb{I}_S \otimes |a_k\rangle\langle a_k| \otimes \mathbb{I}_E$ and $\rho := |\Psi_{\text{fin}}\rangle\langle\Psi_{\text{fin}}|$. Tracing out E gives the reduced state

$$\rho_{SA} = \text{Tr}_E\, \rho = \sum_{i,j} c_i c_j^* \; |s_i\rangle\langle s_j| \otimes |a_i\rangle\langle a_j| \; \langle e_j\rangle\, e_i. \tag{14.62}$$

Decoherence yields $\langle e_j\rangle\, e_i \approx \delta_{ij}$ (orthogonal records), hence

$$\rho_{SA} \;\approx\; \sum_i |c_i|^2 \; |s_i\rangle\langle s_i| \otimes |a_i\rangle\langle a_i| \,. \tag{14.63}$$

Therefore the pointer probabilities obey

$$p_k := \text{Tr}(\rho P_k) = \text{Tr}_{SA}(P_k\, \rho_{SA}) \;\approx\; |c_k|^2, \tag{14.64}$$

as recorded abstractly in Lemma 14.2 (cf. eq. (14.19)). In the IQR semantics, the evaluation of overlaps is via the Emergent Inner Product (14.38), so (14.64) reads

$$p_k = \langle\Psi_{\text{fin}}|\, P_k\, |\Psi_{\text{fin}}\rangle = |c_k|^2 \; \langle s_k\rangle\, s_k \; \langle a_k\rangle\, a_k \; \langle e_k\rangle\, e_k = |c_k|^2$$

(see also the direct projector computation in eq. (14.45)).

Step 2 (Uniqueness from the axioms on the AD–visible slice). Stage 2 fixes the functional dependence on rank–1 branches by relational envariance, fine–graining and continuity (Section 14.3.2), giving $p_k = |c_k|^2$ (eq. (14.43)). Axioms (1)–(5) of Section 14.3.3 then extend the quadratic law to *all* projectors: by the structural theorem Theorem 14.3 (Gleason/Busch–style) or, equivalently, by the compact corollary Theorem 14.4, one has

$$p(P|\psi) = \langle\psi|\, P\, |\psi\rangle \qquad \text{for every projector } P. \tag{14.65}$$

Taking $P = P_k$ in (14.65) reproduces the first equality in (14.61) and agrees with (14.64).

Step 3 (Compatibility with the Universal Quantum Constraint). Let the timeless global state satisfy $\hat{H}_{\text{univ}} |\Psi_0\rangle = 0$ (Theorem 13.168). In the Page–Wootters setting, local statistics read from the clock condition must reproduce globally conserved expectations for arbitrary spectra. The consistency argument of Section 14.3.4 shows that any alternative assignment $p_i = \phi(|c_i|^2)$ with $\phi \neq \text{id}$ violates these conserved quantities (see also the no–go illustration Section 14.3.3). Hence the quadratic rule is not only sufficient but necessary for global–to–local consistency with the Universal Quantum Constraint.

Step 4 (Holosymmetric enforcement). Within the operational pipeline of Holosymmetry, deviations from (14.61) induce an algebraic twist in the commutator–lensing diagnostics and

are suppressed by Stage–II selection (Sections 13.2.7 and 14.3.3). Thus non–Born proposals are dynamically excluded in addition to being structurally inconsistent.

Combining Steps 1–4 gives (14.61). Because the squared modulus is computed via the Emergent Inner Product (14.38), the value $|c_k|^2$ is precisely the *relational self–coherence* of the kth branch (Section 14.3.2), completing the equivalence between coherence and probability. □

Conclusion of Stage 3. The Born rule is *uniquely* singled out by symmetries (envariance), composition and dialectic axioms, and global–to–local consistency with the Universal Quantum Constraint; any deviation either violates these structural requirements or incurs nonzero holosymmetric twist (14.59) and is filtered out by Stage II selection. Together with Stage 2, this completes the derivation: probabilities equal *relational self–coherence*, implemented as the projector expectation (14.61) computed by the Emergent Inner Product (14.38).

14.3.4 A Formal Proof of the Uniqueness of the Born Rule

We now complete the programme by proving that the quadratic (Born) assignment of probabilities is *unique* under the structural and physical principles already adopted in this treatise. The proof has two complementary parts. Part 1 is a universal *global–to–local consistency* argument in the Page–Wootters setting [616], valid for arbitrary pure states and spectra. Part 2 elevates the result to *all* measurements via a Gleason/Busch–style frame-function uniqueness theorem [613, 355], phrased in the holosymmetric/GNS language.

Part 1: Generalization for an Arbitrary Quantum State

Work in the emergent/GNS arena $\mathcal{H}_{\text{em}}$ selected by the Holo–State $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$. Consider a partition into a system S and a clock C. Let $\{|E_i\rangle\}$ be an orthonormal eigenbasis of $\hat{H}_S$ with eigenvalues $\{E_i\}$ and $\{|t_i\rangle\}$ an orthonormal clock basis. A general *timeless* Holo–State satisfying the Universal Quantum Constraint, $\hat{H}_{\text{univ}}\,|\overset{\circ}{\boldsymbol{\Psi}}\rangle = 0$, can be written as

$$|\overset{\circ}{\boldsymbol{\Psi}}\rangle = \sum_{i=1}^{N} c_i\ |E_i\rangle_S\,|t_i\rangle_C\,, \qquad \sum_i |c_i|^2 = 1, \tag{14.66}$$

with normalization evaluated by the Emergent Inner Product (Section 14.3.2). The globally conserved system energy is

$$\langle E\rangle_{\text{global}} = \langle\overset{\circ}{\boldsymbol{\Psi}}|\ \hat{H}_S \otimes \mathbb{I}_C\ |\overset{\circ}{\boldsymbol{\Psi}}\rangle = \sum_{i=1}^{N} |c_i|^2 E_i. \tag{14.67}$$

Local readout and a general probability rule. By Page–Wootters, the conditional state of S given the clock event $|t_i\rangle$ is $|E_i\rangle$, and an observer's statistics are determined by probabilities $\{p_i\}$. Stage 2 has already forced *phase indifference* (Section 14.3.2), so any admissible rule must be of the form

$$p_i \;=\; f\big(|c_i|\big) \;=\; \phi\big(|c_i|^2\big), \qquad \phi : [0,1] \to [0,1], \quad \sum_i \phi(|c_i|^2) = 1. \tag{14.68}$$

The locally inferred energy is then

$$\langle E\rangle_{\text{local}} = \sum_{i=1}^{N} \phi(|c_i|^2)\, E_i. \tag{14.69}$$

Lemma 14.7 (Coefficient equality from arbitrary spectra)**.** *If $\sum_{i=1}^{N} \alpha_i\, E_i = \sum_{i=1}^{N} \beta_i\, E_i$ for all choices of real $\{E_i\}_{i=1}^{N}$, then $\alpha_i = \beta_i$ for every $i = 1, \dots, N$.*

Proof. Define two linear functionals on $\mathbb{R}^N$,

$$F(E_1, \dots, E_N) := \sum_{i=1}^{N} \alpha_i E_i, \qquad G(E_1, \dots, E_N) := \sum_{i=1}^{N} \beta_i E_i.$$

By hypothesis one has $F \equiv G$ as functions on $\mathbb{R}^N$. Hence their difference

$$H(E_1, \dots, E_N) := F(E_1, \dots, E_N) - G(E_1, \dots, E_N) = \sum_{i=1}^{N} (\alpha_i - \beta_i) E_i$$

is the zero functional: $H(E) = 0$ for all $E \in \mathbb{R}^N$.

Fix $j \in \{1, \dots, N\}$ and evaluate H on the j-th canonical basis vector $e^{(j)} := (0, \dots, 0, 1, 0, \dots, 0)$ (with a single 1 in the j-th slot). Then

$$0 = H\big(e^{(j)}\big) = \sum_{i=1}^{N} (\alpha_i - \beta_i) e_i^{(j)} = \alpha_j - \beta_j,$$

so $\alpha_j = \beta_j$. Since j was arbitrary, $\alpha_i = \beta_i$ for all i, proving the claim.

Application in the text. In Section 14.3.4, the identities

$$\langle E\rangle_{\text{global}} = \sum_i |c_i|^2 E_i \quad \text{(eq. (14.67))} \qquad \text{and} \qquad \langle E\rangle_{\text{local}} = \sum_i \phi(|c_i|^2) E_i \quad \text{(eq. (14.69))}$$

hold for all spectra $\{E_i\}$. Applying Lemma 14.7 with $\alpha_i = |c_i|^2$ and $\beta_i = \phi(|c_i|^2)$ yields $\phi(|c_i|^2) = |c_i|^2$ for every i, which is the consistency step used there. □

Consistency forces the Born weights. Global–to–local consistency requires $\langle E\rangle_{\text{local}} = \langle E\rangle_{\text{global}}$ *for all* $\{c_i\}$ and *all* spectra $\{E_i\}$. Applying Lemma 14.7 to (14.67)–(14.69) yields

$$\phi(|c_i|^2) \;=\; |c_i|^2 \quad \text{for all } i \text{ and all families } \{c_i\}. \tag{14.70}$$

Equivalently, $f(|c_i|) = |c_i|^2$. Thus any deviation from the quadratic rule generically violates the Universal Quantum Constraint once local time and statistics are read relationally (Section 14.3.3).

Part 2: Generalization for an Arbitrary Probability Rule (in the spirit of Gleason/Busch)

Frame functions and measurement structure. A (sharp) quantum measurement is a projective decomposition $\{P_i\}$ with $\sum_i P_i = \mathbb{I}$. A *probability assignment* is a frame function p that maps every projector P and unit vector $|\psi\rangle$ to $p(P \mid \psi) \in [0, 1]$ such that: (i) $p(P_i \mid \psi) \geq 0$ and $\sum_i p(P_i \mid \psi) = 1$ for any orthogonal decomposition (finite or, in infinite dimensions, countable; cf. Remark 14.5); and (ii) p respects unitary changes of basis (implementation invariance).

Gleason–Busch uniqueness. On Hilbert spaces of complex dimension ≥ 3, Gleason's theorem entails that any such frame function arises from a density operator ρ via the Born rule $p(P \mid \rho) = \mathrm{Tr}(\rho P)$ [613]. For arbitrary finite dimension (including qubits) and generalized measurements (POVMs), Busch's extension shows the same conclusion under operational assumptions satisfied here [355].

Holosymmetric/GNS instantiation. In our setting: (i) the Emergent Inner Product (Section 14.3.2) provides the orthogonality/projection structure on $\mathcal{H}_{\mathrm{em}}$; (ii) Stage 2 supplies unitary covariance and phase–indifference (envariance; Section 14.3.2); (iii) the tensorial/dialectic composition enforces product factorization and coarse–graining additivity (axioms (1)–(5); see Section 14.3.3). Hence the hypotheses of the Gleason–Busch route are met, forcing

$$p(P \mid \psi) \;=\; \langle\psi|\, P\, |\psi\rangle \quad \text{for all projectors } P \text{ on } \mathcal{H}_{\mathrm{em}} \;\; (\text{and } p(E \mid \rho) = \mathrm{Tr}(\rho E) \text{ for POVMs}). \tag{14.71}$$

By the Ch. 12 equivalence, (14.71) *descends* to the BRST–physical diagnostic realized reference configuration $\mathcal{H}_{\mathrm{phys}}$ on certified plateaux.

Corollary 14.8 (Uniqueness of the Born Rule in the holosymmetric/GNS framework)**.** *For any measurement on the emergent/GNS space (and hence on $\mathcal{H}_{\mathrm{phys}}$ on certified plateaux), the only probability assignment compatible with (a) Stage 2 symmetries, (b) unitary covariance and composition laws, and (c) the Universal Quantum Constraint, is the Born rule $p = \mathrm{Tr}(\rho\,\cdot\,)$. In particular, for pure post–measurement states $|\Psi_{\mathrm{fin}}\rangle = \sum_k c_k\, |s_k\rangle\, |a_k\rangle\, |e_k\rangle$ and pointer projectors $P_k := \mathbb{I}_S \otimes |a_k\rangle\langle a_k| \otimes \mathbb{I}_E$, one has*

$$p_k \;=\; \langle\Psi_{\mathrm{fin}}|\, P_k\, |\Psi_{\mathrm{fin}}\rangle \;=\; |c_k|^2, \tag{14.72}$$

with the right–hand side evaluated via the Emergent Inner Product, i.e. as the relational self–coherence *of branch k (Section 14.3.2).*

Proof. **Step 1 (Stage 2 $\Rightarrow$ rank–1 quadratic weights).** By the Stage 2 symmetry analysis (Section 14.3.2), phase–indifference and swap–envariance force a modulus–only dependence and equiprobability for equal–modulus branches, and ancilla fine–graining with continuity yields the quadratic law on rank–1 pointer branches:

$$p_k \;=\; |c_k|^2 \qquad (\text{Stage 2 result; eq. (14.43)}).$$

Equivalently, for the post–measurement state of eq. (14.37) and pointer projectors P_k, the projector computation gives

$$p_k \;=\; \langle\Psi_{\mathrm{fin}}|\, P_k\, |\Psi_{\mathrm{fin}}\rangle = |c_k|^2\, \langle s_k\rangle\, s_k\, \langle a_k\rangle\, a_k\, \langle e_k\rangle\, e_k = |c_k|^2 \quad (\text{eq. (14.45)}).$$

(Under decohered records $\langle e_j\rangle\, e_i \simeq \delta_{ij}$, this also follows by tracing out the environment; cf. Lemma 14.2.)

Step 2 (Structural lift to all projectors). Assumptions (a) and (b) instantiate axioms (1)–(5) of Section 14.3.3: (countable) additivity and coarse–graining noncontextuality, unitary

covariance, product factorization, and continuity. Hence, by the Stage 3 structural theorem (Theorem 14.3), any admissible frame function on projectors must be of Born form:

$$p(P \mid \psi) = \langle\psi| P |\psi\rangle \qquad \text{for all projectors } P,$$

with the $d = 2$ case covered by doubling and factorization (or directly by the Busch POVM route; Theorem 14.4).

Step 3 (UQC excludes alternatives; uniqueness). Condition (c) enforces global–to–local consistency from the Universal Quantum Constraint

$$\hat{H}_{\text{univ}} |\overset{\circ}{\boldsymbol{\Psi}}\rangle = 0 \qquad \text{(Theorem 13.168, eq. (13.236), and Definition 13.167)},$$

via Page–Wootters conditioning. For arbitrary spectra, any nonquadratic deformation $p_i = \phi(|c_i|^2) \neq |c_i|^2$ generically mismatches conserved global expectations (Sections 14.3.3 and 14.3.4). Thus no alternative rule is compatible with the UQC, and the Born assignment is unique under (a)–(c).

Conclusion and the pointer case. Combining Steps 1–3 yields the statement of the corollary: $p = \mathrm{Tr}(\rho\,\cdot\,)$ is the only assignment compatible with (a)–(c). In particular, for $|\Psi_{\text{fin}}\rangle$ and P_k as above,

$$p_k = \langle\Psi_{\text{fin}}| P_k |\Psi_{\text{fin}}\rangle = |c_k|^2,$$

where $|c_k|^2$ is evaluated by the Emergent Inner Product (14.38), i.e. as the *relational self–coherence* of branch k (Section 14.3.2). □

Holosymmetric enforcement. Any non–Born assignment breaks either the structural requirements of the frame function or the global–to–local conservation inherited from the global Holo–State $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$; such rules induce algebraic twist in the lensing diagnostics and are suppressed by Stage II holosymmetric orthogonality (Section 13.2.7). Thus Holosymmetry is the physical arbiter selecting the unique, self–consistent probability law. Parts 1 and 2 together establish that *quadratic probabilities are necessary and sufficient* for consistency with emergent time, conservation from the timeless global state, and the operational structure of measurements in the holosymmetric/GNS framework. The Born rule is therefore not an additional postulate but the unique valuation compatible with the relational architecture of the theory [616, 613, 355].

Dialectic derivation via the Tetralemma and Axiom of Duality. The foregoing corollary can be complemented by an internal derivation based solely on the *Tetralemma of Algebra* (Tetra; Definition 1.5) and the *Axiom of Duality* (AD; Axiom 1.10) as implemented in the IQR architecture (see Foundations). The required ingredients are the operational consequences of Tetra on the synthesis stratum and of AD on the Boolean outcome stratum. We record them as two lemmas and a synthesis theorem.

Lemma 14.9 (Parasymmetric swap invariance (Tetra ⇒ envariance))**.** *Let*

$$|\Psi_{\text{fin}}\rangle = \sum_k c_k\ |s_k\rangle\,|a_k\rangle\,|e_k\rangle$$

be the post–measurement state of eq. (14.37) with orthogonal $\{|a_k\rangle\}$ and decohered $\{|e_k\rangle\}$. Assume the parasymmetric automorphisms of the Tetra act transitively on equal–modulus branches within the synthesis stratum (cf. Definition 1.5 and Proposition 1.8). Then for any j,k with $|c_j| = |c_k|$ there exists an inner automorphism (implemented as a unitary on inaccessible records) that swaps the branches $j \leftrightarrow k$ while leaving all SA–relational data invariant. Consequently the outcome valuation on the AD–visible slice (Twofold reduction induced by Axiom 1.10) satisfies

$$|c_j| = |c_k| \implies p_j = p_k,$$

*and, together with phase–indifference of eq. (14.39), the probabilities can depend on c_k only through $|c_k|$ (*equiprobability/envariance*, cf. eq. (14.4), [158]).*

Proof. Write the post–measurement state in the decohered pointer form of eq. (14.37),

$$|\Psi_{\text{fin}}\rangle = \sum_i c_i \, |s_i\rangle \, |a_i\rangle \, |e_i\rangle \,, \qquad \langle a_i \rangle \, a_j = \delta_{ij}, \quad \langle e_i \rangle \, e_j \simeq \delta_{ij}.$$

Tracing out E gives the reduced state on SA (cf. eq. (14.15)):

$$\rho_{SA} := \mathrm{Tr}_E\Big(|\Psi_{\text{fin}}\rangle\langle\Psi_{\text{fin}}| \Big) = \sum_i |c_i|^2 \, |s_i\rangle \, |a_i\rangle\langle s_i| \, \langle a_i| \,, \tag{14.73}$$

since off–diagonal terms are suppressed by $\langle e_j \rangle \, e_i \simeq 0$.

Envariant phase alignment. Fix $j \neq k$ with equal moduli $|c_j| = |c_k| =: r$. By phase–indifference (eq. (14.39)) there exists a unitary on E that rephases only $|e_j\rangle \,, |e_k\rangle$ so as to remove the relative phase between c_j and c_k:

$$U_E^{\text{ph}} : \; |e_j\rangle \mapsto e^{i\theta_j} \, |e_j\rangle \,, \quad |e_k\rangle \mapsto e^{i\theta_k} \, |e_k\rangle \,, \quad \theta_j - \theta_k = \arg c_j - \arg c_k,$$

which transforms the pair (c_j, c_k) to (r, r) in the expansion while leaving all SA relations (hence ρ_{SA}) invariant.

Parasymmetric swap on inaccessible records. By the Tetra hypothesis (Definition 1.5 and Proposition 1.8), there exists an *inner* automorphism implemented by a unitary on the inaccessible records subspace that swaps the two environmental labels:

$$U_E^{(j\leftrightarrow k)} : \; |e_j\rangle \leftrightarrow |e_k\rangle \,, \qquad |e_\ell\rangle \mapsto |e_\ell\rangle \quad (\ell \neq j,k). \tag{14.74}$$

Define $|\Psi'\rangle := (\mathbf{1}_{SA} \otimes U_E^{(j\leftrightarrow k)} U_E^{\text{ph}}) \, |\Psi_{\text{fin}}\rangle$. Then

$$|\Psi'\rangle = \sum_{i \notin \{j,k\}} c_i \, |s_i\rangle \, |a_i\rangle \, |e_i\rangle \;+\; r \, |s_j\rangle \, |a_j\rangle \, |e_k\rangle \;+\; r \, |s_k\rangle \, |a_k\rangle \, |e_j\rangle \,.$$

Because $U_E^{(j\leftrightarrow k)} U_E^{\text{ph}}$ acts only on E, the reduced state on SA is unchanged:

$$\rho'_{SA} := \mathrm{Tr}_E\Big(|\Psi'\rangle\langle\Psi'| \Big) = \rho_{SA}, \tag{14.75}$$

with ρ_{SA} given by (14.73). Hence, for every observable O_{SA} on the SA algebra,

$$\langle \Psi_{\text{fin}}| \, O_{SA} \otimes \mathbf{1}_E \, |\Psi_{\text{fin}}\rangle = \mathrm{Tr}\Big(\rho_{SA} \, O_{SA}\Big) = \mathrm{Tr}\Big(\rho'_{SA} \, O_{SA}\Big) = \langle \Psi'| \, O_{SA} \otimes \mathbf{1}_E \, |\Psi'\rangle \,,$$

i.e. all SA–relational data are invariant under the swap.

Equiprobability for equal moduli. In particular, for the pointer projectors $P_\ell = \mathbf{1}_S \otimes |a_\ell\rangle\langle a_\ell| \otimes \mathbf{1}_E$,

$$p_j := \langle \Psi_{\rm fin}| \, P_j \, |\Psi_{\rm fin}\rangle = \langle \Psi'| \, P_j \, |\Psi'\rangle = \langle \Psi_{\rm fin}| \, P_k \, |\Psi_{\rm fin}\rangle =: p_k,$$

where the middle equality uses that $U_E^{(j\leftrightarrow k)} U_E^{\rm ph}$ acts trivially on P_j (it is supported on E) and swaps the environmental labels between the equal–modulus branches. Thus $|c_j| = |c_k| \Rightarrow p_j = p_k$, establishing *equiprobability by envariant swaps* (eq. (14.4); cf. [158]).

Dependence only on $|c_k|$. Combining this equiprobability with phase–indifference (eq. (14.39)) shows that any admissible valuation on the AD–visible slice (Twofold reduction via Axiom 1.10) can depend on c_k only through its modulus, completing the claim. □

Lemma 14.10 (Monoidal Duality ⇒ product factorization)**.** *Let A and B be independent subsystems and suppose the Axiom of Duality is monoidal with respect to their tensor composition on the BRST–physical space. Then for product states $|\psi_A\rangle\otimes|\psi_B\rangle$ and product projectors $P_A\otimes P_B$ one has*

$$p(P_A\otimes P_B \,|\, \psi_A\otimes\psi_B) \;=\; p(P_A|\psi_A)\, p(P_B|\psi_B).$$

In particular, for rank–1 branches this enforces a multiplicative law for the one–branch dependence $f : [0,1] \to [0,1]$ defined by $p_k = f(|c_k|)$.

Proof. Work on the BRST–physical/emergent Hilbert space with the Boolean outcome layer singled out by the Axiom of Duality (AD) (Section 14.1.3). Let A and B be independent subsystems and assume AD is *monoidal* with respect to their tensor composition, so that the joint outcome algebra is the Boolean product of the marginals and elementary outcomes are represented by product projectors $P_A \otimes P_B$. Let p satisfy the axioms (1)–(2),(4),(5) of Section 14.3.3 (normalization/additivity on orthogonal joins, unitary covariance, coarse–graining noncontextuality, and continuity).

Step 1: Define the bivariate joint map. For product states and projectors set

$$x := p(P_A|\psi_A), \qquad y := p(P_B|\psi_B), \qquad F(x,y) \;:=\; p(P_A\otimes P_B \,|\, \psi_A\otimes\psi_B).$$

By unitary covariance (axiom (2)), F depends only on the scalars $x, y \in [0,1]$ (implementation invariance).

Step 2: Marginals and boundary conditions. Using coarse–graining noncontextuality (axiom (4)) and additivity (axiom (1)),

$$\begin{aligned} p(P_A\otimes\mathbf{1} \,|\, \psi_A\otimes\psi_B) &= p\Big(P_A\otimes P_B + P_A\otimes(\mathbf{1}-P_B) \,\Big|\, \psi_A\otimes\psi_B\Big) \\ &= F(x,y) + F\Big(x, 1-y\Big). \end{aligned} \tag{14.76}$$

But by AD's monoidal semantics, $P_A \otimes \mathbf{1}$ is just the coarse event "P_A occurs in A", so

$$p(P_A\otimes\mathbf{1} \,|\, \psi_A\otimes\psi_B) = p(P_A|\psi_A) = x, \tag{14.77}$$

and similarly $p(\mathbf{1}\otimes P_B \,|\, \psi_A\otimes\psi_B) = y$. Hence F satisfies the boundary conditions

$$F(1,y) = y, \qquad F(x,1) = x, \qquad F(0,y) = 0, \qquad F(x,0) = 0. \tag{14.78}$$

Step 3: Separate additivity (affinity) in each argument. Let Q_B^1, Q_B^2 be orthogonal with $Q_B^1 + Q_B^2 = P_B$. By additivity and coarse–graining noncontextuality (axioms (1),(4)),

$$F\Big(x,\, p(Q_B^1|\psi_B) + p(Q_B^2|\psi_B)\Big) = F\Big(x,\, p(Q_B^1|\psi_B)\Big) + F\Big(x,\, p(Q_B^2|\psi_B)\Big).$$

Thus $y \mapsto F(x,y)$ is affine on $[0,1]$ for each fixed x. By symmetry of the assumptions, $x \mapsto F(x,y)$ is affine for each fixed y. With continuity (axiom (5)), F is *bi–affine* on $[0,1]^2$, hence of the form

$$F(x,y) = \alpha\, xy + \beta\, x + \gamma\, y + \delta, \qquad \alpha,\beta,\gamma,\delta \in \mathbb{R}. \tag{14.79}$$

Step 4: Determine the coefficients from the boundary conditions. Insert (14.78) into (14.79):

$$F(0,y) = \delta = 0, \qquad F(x,0) = \delta + \beta x = 0 \;\Rightarrow\; \beta = 0,$$

$$F(1,y) = \alpha y + \gamma y = y \;\Rightarrow\; \alpha + \gamma = 1, \qquad F(x,1) = \alpha x + \gamma = x \;\Rightarrow\; \alpha = 1,\ \gamma = 0.$$

Therefore $F(x,y) = xy$.

Step 5: Product factorization and one–branch multiplicativity. We have shown

$$p(P_A\otimes P_B \,|\, \psi_A\otimes\psi_B) = F\Big(p(P_A|\psi_A), p(P_B|\psi_B)\Big) = p(P_A|\psi_A)\, p(P_B|\psi_B),$$

i.e. *product factorization* for independent subsystems, as claimed. In particular, for rank–1 pointer branches with $p_k = f(|c_k|)$ and similarly for B, the factorization reads $f(rs) = f(r)\, f(s)$ for $r,s \in [0,1]$, yielding the multiplicative Cauchy law used in Stage 3 (with continuity fixing $f(r) = r^\alpha$; cf. Theorems 14.3 and 14.4). □

Theorem 14.11 (Born valuation from Tetralemma and Duality)**.** *Assume the setting of eq. (14.37) and impose: (i) phase–indifference (eq. (14.39)), (ii) parasymmetric swap invariance of Lemma 14.9 (induced by the Tetralemma; cf. Definition 1.5 and Proposition 1.8), (iii) finite additivity and coarse–graining noncontextuality on the AD–visible slice (Twofold reduction via the Axiom of Duality ⇒ Boolean outcomes; Axiom 1.10), (iv) unitary covariance, (v) product factorization (Monoidal Duality; Lemma 14.10 together with Axiom 1.10), and (vi) continuity. Then for every projector P on the BRST–physical/emergent Hilbert space and unit vector $|\psi\rangle$,*

$$p(P \mid \psi) \;=\; \langle\psi|\, P\, |\psi\rangle\,. \tag{14.80}$$

In particular, for pointer projectors P_k, $p_k = \langle\Psi_{\mathrm{fin}}|\, P_k\, |\Psi_{\mathrm{fin}}\rangle = |c_k|^2$.

Proof idea. By phase–indifference and Lemma 14.9 (Tetralemma–induced parasymmetric swaps; Definition 1.5 and Proposition 1.8), $p_k = f(|c_k|)$ and equal–modulus branches are equiprobable (for n equal branches, $p = 1/n$). Lemma 14.10 (Monoidal Duality; Axiom 1.10) applied to product states forces $f(rs) = f(r)f(s)$ for $r, s \in [0,1]$; continuity implies $f(r) = r^\alpha$ for some $\alpha > 0$. Equiprobability for n equal branches yields $f(1/\sqrt{n}) = 1/n$, hence $(1/\sqrt{n})^\alpha = 1/n$ and $\alpha = 2$. Additivity, coarse–graining noncontextuality on the AD–visible slice (Twofold/AD), and unitary covariance then extend from rank–1 projectors to arbitrary P, giving (14.80). This is the relational analogue of the Gleason/Busch uniqueness route [613, 355].

Proof. **Step 1 (Restriction to rank–1 pointer events; phase indifference and swap envariance).** Let $P_k := \mathbb{I}_S \otimes |a_k\rangle\langle a_k| \otimes \mathbb{I}_E$ be the pointer projector associated with the decohered state of eq. (14.37). Phase indifference eq. (14.39) yields a modulus–only dependence

$$p_k \;=\; p(P_k \mid \Psi_{\text{fin}}) \;=\; f\Big(|c_k|\Big), \tag{14.81}$$

and parasymmetric swap invariance Lemma 14.9 (Tetralemma; Definition 1.5 and Proposition 1.8) enforces equiprobability for equal–modulus branches: if $|c_j| = |c_k|$ then $p_j = p_k$ (cf. eq. (14.4)).

Step 2 (Fine–graining ⇒ rational weights ⇒ quadratic form). Ancilla fine–graining on the synthesis stratum (Stage 2; consistent with Tetra) refines a branch with $|c_k|^2 = r_k/N$ into r_k equal–modulus subbranches and, by coarse–graining noncontextuality and additivity on the AD–visible slice (Twofold via Axiom 1.10), implies

$$p_k \;=\; \frac{r_k}{N} \;=\; |c_k|^2 \qquad \text{for all rational weights,} \tag{14.82}$$

see Section 14.3.2 and eq. (14.42). Continuity extends (14.82) to all $|c_k| \in [0,1]$, giving $p_k = |c_k|^2$ (Stage 2 culmination eq. (14.43)).

Step 3 (Uniqueness on branches from Monoidal Duality). Independently of Step 2's conclusion, Monoidal Duality Lemma 14.10 (arising from the Axiom of Duality; Axiom 1.10) applied to product states forces the multiplicative Cauchy relation

$$f(rs) = f(r)\, f(s) \qquad (r, s \in [0,1]). \tag{14.83}$$

With continuity, (14.83) implies $f(r) = r^\alpha$ for some $\alpha > 0$. Envariant equiprobability for n equal–modulus branches (Step 1; Tetra) gives $f(1/\sqrt{n}) = 1/n$, hence $(1/\sqrt{n})^\alpha = 1/n$ and $\alpha = 2$. Therefore $f(r) = r^2$, recovering (and uniquely fixing) (14.82) on rank–1 pointer events.

Step 4 (Extension from rank–1 to arbitrary projectors on the AD–visible slice). Let P be any projector with an orthogonal decomposition into rank–1 projectors $\{Q_j\}$ inside the outcome (AD) layer (Twofold reduction; Axiom 1.10). By finite additivity and coarse–graining noncontextuality (axioms (1),(4) in Section 14.3.3),

$$p(P \mid \psi) \;=\; \sum_j p(Q_j \mid \psi) \;=\; \sum_j \Big|\langle \phi_j \rangle\, \psi\Big|^2, \tag{14.84}$$

where $Q_j = |\phi_j\rangle\langle\phi_j|$ and the second equality uses the rank–1 result of Steps 2–3. As $P = \sum_j Q_j$ and $\{|\phi_j\rangle\}$ is an orthonormal set spanning $\mathrm{Ran}(P)$, (14.84) equals

$$\sum_j \left|\langle\phi_j|\,\psi\right|^2 = \langle\psi|\left(\sum_j |\phi_j\rangle\langle\phi_j|\right)|\psi\rangle = \langle\psi|\,P\,|\psi\rangle\,. \tag{14.85}$$

Step 5 (Unitary covariance and basis independence). If P is presented in a different orthogonal refinement or basis, unitary covariance (axiom (2)) transports the computation, ensuring the value does not depend on the chosen decomposition. Therefore (14.85) holds for every projector P and unit vector $|\psi\rangle$, proving

$$p(P \mid \psi) = \langle\psi|\,P\,|\psi\rangle\,,$$

which is eq. (14.80). In particular, for pointer projectors P_k one has $p_k = \langle\Psi_{\mathrm{fin}}|\,P_k\,|\Psi_{\mathrm{fin}}\rangle = |c_k|^2$ (cf. Section 14.3.2 and eq. (14.45)).

Conclusion. Tetralemma–induced envariance (phase indifference, parasymmetric swap symmetry; Definition 1.5 and Proposition 1.8) and Duality–imposed composition laws (additivity, noncontextuality, product factorization on the AD/Twofold layer; Axiom 1.10 and Lemma 14.10) with continuity suffice to force the Born valuation on the AD–visible slice. This is the relational counterpart of the Gleason/Busch route [613, 355], carried out within the IQR/Holosymmetry architecture. □

Remark 14.12 (Consistency with global constraints)**.** The global–to–local consistency check of Section 14.3.4 shows that any nonquadratic deformation violates conserved expectations when time is emergent in the Page–Wootters sense [616]. Equivalently, such deformations are incompatible with the Universal Quantum Constraint $\hat{H}_{\mathrm{univ}}\,|\mathring{\mathbf{\Psi}}\rangle = 0$ (cf. Theorem 13.168, eq. (13.236), and Definition 13.167). Within Holosymmetry this inconsistency is filtered out by Stage II orthogonality and lensing diagnostics (cf. Section 13.2.7).

Conceptual content of Theorem 14.11. *First,* the theorem links dialectical foundations to operational probabilities. On the synthesis (Parasymmetric) stratum, the post–measurement state has the decohered, lane–orthogonal form of (14.37). The Tetralemma (Tetra; Definition 1.5) contributes *relational envariance*: phase–indifference (eq. (14.39)) and swap invariance among equal–modulus branches via its parasymmetric automorphisms (cf. Proposition 1.8), summarized by Lemma 14.9. These symmetries imply that any admissible outcome weight for branch k can only depend on $|c_k|$ and must yield equiprobability when $|c_j| = |c_k|$ (cf. eq. (14.4)).

Second, the Axiom of Duality (AD; Axiom 1.10) identifies the outcome layer as a Boolean/Twofold slice and imposes structural constraints there: finite (or σ–)additivity for orthogonal joins and coarse–graining noncontextuality ensure that the value assigned to a coarse projector is independent of its refinement; unitary covariance enforces implementation invariance; product factorization (Monoidal Duality; Lemma 14.10 together with Axiom 1.10) enforces independence on tensor products; and continuity guarantees stability under small amplitude changes. Together these conditions force the one–branch dependence to satisfy a

multiplicative Cauchy relation $f(rs) = f(r)f(s)$, whose continuous solutions are $f(r) = r^\alpha$. Envariant equiprobability for n equal–modulus branches then fixes $\alpha = 2$. Thus the branch law is uniquely quadratic.

Third, once the branch law is fixed, the statement extends from branches to *all* projectors P on the emergent/GNS space $\mathcal{H}_{\text{em}}$ (descending to $\mathcal{H}_{\text{phys}}$ on certified plateaux per Ch. 12). In particular, for pointer projectors $P_k = \mathbb{I}_S \otimes |a_k\rangle\langle a_k| \otimes \mathbb{I}_E$ associated with (14.37), one obtains

$$p_k = \langle \Psi_{\text{fin}} |\, P_k \,| \Psi_{\text{fin}} \rangle = |c_k|^2,$$

and, more generally,

$$p(P \mid \psi) = \langle \psi |\, P \,| \psi \rangle \,,$$

which is (14.80). In the IQR/Holosymmetry semantics, this identifies outcome probabilities with the *relational self–coherence* of branches when the inner product is evaluated via the Emergent Inner Product induced by the Fourfold Inter–Modal Correlator (cf. Section 14.3.2, eq. (14.38), and Definition 4.9).

In short, once (i) Tetralemma–induced envariance on the synthesis stratum (Definition 1.5 and Proposition 1.8) and (ii) Axiom of Duality–driven Boolean/compositional axioms on the outcome stratum (Axiom 1.10 and Lemma 14.10) are imposed—together with unitary covariance and continuity—the only admissible valuation is the Born rule. Any alternative would violate either the envariance/equiprobability constraints or the compositional laws dictated by the Axiom of Duality.

No Circularity: Two–Tier Semantics and a Fixed–Point Argument

Definition 14.13 (Pre–probabilistic (state) valuation)**.** Let $\mathfrak{A}$ be the emergent C^*–algebra of effects and ω the BRST–physical/GNS state (a positive, normalized linear functional; cf. the GNS construction [381, 241]). Define the *state–valuation* of an effect $E \in \mathfrak{A}$, $0 \le E \le \mathbb{I}$, by

$$v_0(E) \;:=\; \omega(E) \in [0,1]. \tag{14.86}$$

At this stage v_0 is a *truth/weight* assignment used to form coherence functionals (eqs. (12.82) and (12.83)) and to define the interferometric/lensing readouts on the AD–visible slice selected by the Axiom of Duality (Axiom 1.10); no claim is made that v_0 equals operational probabilities.

Definition 14.14 (Born–valued truths vs. Born probabilities)**.** Given ω on $\mathfrak{A}$, the *Born–valued truth* (state–valuation) of E is $v_0(E) := \omega(E)$ (Definition 14.13). An *operational Born probability* for E is $p(E \mid \rho) := \mathrm{Tr}(\rho E)$ for the density operator ρ representing ω on the emergent Hilbert space (Gleason/Busch route [613, 355]). The identification $v_0 \equiv p(\cdot \mid \rho)$ will be *derived*, not assumed, from the Tetralemma symmetries and the Duality–induced composition laws (Definition 1.5, Proposition 1.8, and Axiom 1.10).

Proposition 14.15 (Holosymmetry depends only on linear state–valuation)**.** *The Holosymmetric Interferometer, the Commutator Lensing Model, and the Selection Principle (Sections 2.4 and 3.3) use v_0 only via linearity, positivity, and normalization of ω: the self–coherence $\mathcal{C}_{\text{phys}}$*

and overlaps Ω_{phys} are built from expectations of effects on the AD–visible (Twofold/Boolean) outcome layer picked out by the Axiom of Duality (Axiom 1.10), with synthesis–stratum symmetries governed by the Tetralemma (Definition 1.5 and Proposition 1.8). Hence all DM/DE selection statements in Section 13.1.4 are well–posed independently of any identification $v_0 \equiv p$.

Proposition 14.16 (Two–tier semantics, fixed–point identification)**.** *In Holosymmetry, interferometric and lensing constructions depend only on v_0 (linearity, positivity, normalization). Stage 2–3, together with Tetralemma–induced envariance on the synthesis stratum and the Axiom of Duality on the outcome stratum (Definition 1.5, Proposition 1.8, Axiom 1.10, and Lemma 14.10), constrain any admissible operational rule to $p = g \circ v_0$ and force $g(x) = x$. Hence for all effects E,*

$$p(E \mid \rho) = v_0(E).$$

Equivalently, Born–valued truths and Born probabilities are conceptually distinct but numerically identical under the IQR/Holosymmetry constraints (Theorems 14.11 and 14.19 and Corollary 14.8).

Thesis. There is no vicious circularity in our use of "Born–valued" truth assignments to build Holosymmetry and then deriving the Born rule. The constructions in Parts I–II employ only the *state–valuation* provided by the emergent/GNS state (a positive, normalized, linear functional on the effect algebra), without interpreting that valuation *as* operational probability [381, 241]. Stage 2–3 subsequently identify the *unique* way to read these valuations as empirical frequencies, thereby recovering the Born rule as a fixed point of the dialectic—Tetralemma (parasymmetric symmetries on the synthesis stratum) plus Axiom of Duality (Boolean/compositional laws on the outcome stratum)—and the holosymmetric architecture (Definition 1.5, Proposition 1.8, Axiom 1.10, and Sections 14.3.2 and 14.3.3).

Lemma 14.17 (Calibration compatibility)**.** *Let p be any operational probability assignment to projectors consistent with Stage 1 dynamics (orthogonal pointer basis) and Stage 2 symmetries (phase–indifference and envariant equiprobability; Section 14.2.2 and eqs. (14.4) and (14.39)). Assume p agrees with the state–valuation v_0 on one complete projective measurement $\{P_k\}$ (i.e. $p(P_k \mid \Psi) = v_0(P_k)$ for all pointer projectors P_k of eq. (14.37)). Then there exists a* state–independent *map $g : [0,1] \to [0,1]$ such that, for all effects E,*

$$p(E \mid \Psi) = g\Big(v_0(E)\Big),$$

with g monotone, $g(0) = 0$, $g(1) = 1$, and (by Stage 2 continuity) continuous on $[0,1]$. (For operational POVM semantics and frame–function structure, see [355].)

Proof. **(A) Setup and layer semantics.** On the outcome layer selected by the Axiom of Duality (AD), measurements form a Boolean/Twofold slice (Axiom 1.10). Stage 1 gives an orthogonal pointer PVM $\{P_k\}$; Stage 2 supplies phase–indifference and envariant swap symmetry (Section 14.2.2 and eqs. (14.4) and (14.39)), which are Tetralemma–induced on the synthesis stratum (Definition 1.5 and Proposition 1.8) and descend to the AD–visible slice. Write $v_0(E) := \omega(E)$ (Definition 14.13).

(B) Well–defined dependence on the scalar v_0 for projectors commuting with the calibration. Let $\mathsf{Alg}(\{P_k\})$ denote the commutative algebra generated by the calibrated pointer resolution. Any projector $Q \in \mathsf{Alg}(\{P_k\})$ is an orthogonal sum $Q = \sum_{k\in S} P_k$ for some index set S. By coarse–graining noncontextuality and finite additivity on the AD slice (axioms (1),(4); Section 14.3.3),

$$p(Q \mid \Psi) = \sum_{k\in S} p(P_k \mid \Psi) = \sum_{k\in S} v_0(P_k) = v_0\Big(\sum_{k\in S} P_k\Big) = v_0(Q).$$

Thus, within $\mathsf{Alg}(\{P_k\})$, p depends only on the scalar $v_0(Q)$ and in fact equals it.

(C) Rational interpolation by parasymmetric fine–graining. Fix any $x \in [0,1] \cap \mathbb{Q}$. By Stage 2 ancilla fine–graining (Section 14.3.2 and eq. (14.42)) and parasymmetric swap invariance (Tetralemma; Lemma 14.9), there exists a projector $Q_x \in \mathsf{Alg}(\{P_k\})$ with $v_0(Q_x) = x$. From (B), $p(Q_x \mid \Psi) = v_0(Q_x) = x$. Hence the map

$$g(x) := p(Q_x \mid \Psi)$$

is well–defined on the rationals in $[0,1]$ and equals x there. Well–definedness follows because any two rational constructions of value x are related by AD–coarse–graining and Tetralemma swaps, under which p is invariant (axioms (1),(4) and Lemma 14.9).

(D) Unitary covariance $\Rightarrow$ dependence only on v_0 for all projectors. Let Q be an arbitrary projector. Choose a unitary U sending $\mathrm{Ran}(Q)$ onto the span of some subset of $\{\mathrm{Ran}(P_k)\}$; i.e. $Q' = UQU^\dagger \in \mathsf{Alg}(\{P_k\})$. Unitary covariance (implementation invariance; axiom (2)) yields

$$p(Q \mid \Psi) = p(Q' \mid U\Psi), \qquad v_0(Q) = v_0'(Q') \;\text{ with }\; v_0'(E) := \omega_U(E) := \omega(U^\dagger E U).$$

By (B), $p(Q' \mid U\Psi) = v_0'(Q')$. Hence $p(Q \mid \Psi)$ depends only on $v_0(Q)$, not on the particular decomposition of Q. Define g on *all* of $[0,1]$ by

$$g\Big(v_0(Q)\Big) \;:=\; p(Q \mid \Psi) \qquad \text{for any projector } Q.$$

This is well–defined: if Q_1, Q_2 satisfy $v_0(Q_1) = v_0(Q_2)$, a unitary transports Q_i into $\mathsf{Alg}(\{P_k\})$ where $p = v_0$, and envariant swaps/coarse–graining preserve both v_0 and p (Lemma 14.9, axioms (1),(4)).

(E) Endpoints, monotonicity, continuity (projectors). Normalization gives $g(0) = 0$, $g(1) = 1$ since $p(\mathbf{0}) = 0$, $p(\mathbf{I}) = 1$. If $Q \le R$ (orthogonal sum $R = Q \oplus D$), additivity and positivity imply $p(R) = p(Q) + p(D) \ge p(Q)$, hence g is monotone. Stage 2 continuity (continuity of outcome weights under parasymmetric refinements; Section 14.3.2) gives continuity of g on $[0,1]$; recall g equals the identity on rationals by (C), so the continuous extension is unique.

(F) Extension from projectors to general effects (POVM effects). Let E be any effect ($0 \le E \le \mathbf{I}$). By Naimark dilation there exists an ancilla $\mathcal{K}$, an isometry $V : \mathcal{H} \to \mathcal{H} \otimes \mathcal{K}$, and a projector $\tilde{P}$ on $\mathcal{H} \otimes \mathcal{K}$ such that the two–outcome POVM $\{E, \mathbf{I} - E\}$ is realized as

the sharp measurement $\{\tilde{P}, \mathbf{I} - \tilde{P}\}$ on the dilated space ([355]). Implementation invariance (axiom (2)) and AD coarse–graining ensure

$$p(E \mid \Psi) \;=\; p(\tilde{P} \mid V\Psi), \qquad v_0(E) \;=\; \omega(E) \;=\; \omega_V(\tilde{P}) \;=:\; v_0^{\mathrm{dil}}(\tilde{P}),$$

where v_0^{dil} is the state–valuation of the dilated state. Applying the projector case (D) on the dilated space gives

$$p(\tilde{P} \mid V\Psi) \;=\; g\Big(v_0^{\mathrm{dil}}(\tilde{P})\Big) \;=\; g\Big(v_0(E)\Big),$$

hence $p(E \mid \Psi) = g\Big(v_0(E)\Big)$ for all effects E.

(G) State–independence of g. By construction, g depends only on the calibration equality on the fixed PVM and on structural symmetries (Tetralemma swaps and AD Boolean/compositional laws; Definition 1.5, Proposition 1.8, and Axiom 1.10). Therefore g is independent of the particular effect E and of its implementation. (If, in addition, Monoidal Duality/product factorization is assumed; Lemma 14.10, the same g is universal across product preparations, i.e. independent of the choice of state in a tensor factor.)

Combining (A)–(F) establishes the claim. □

Remark 14.18 (Strengthenings under additional axioms)**.** If one assumes the full axiom set (1)–(5) of Section 14.3.3 on the AD–visible slice together with Stage 2 continuity, then g is not only monotone and continuous but also *affine* on convex mixtures of effects implemented by classical randomization (AD Booleanity), which combined with [355] yields the Born form $g(x) = x$ (cf. Theorem 14.11 and Corollary 14.8).

Theorem 14.19 (Fixed–point non–circularity and uniqueness)**.** *Assume the operational consequences of the Tetralemma of Algebra (parasymmetric swap–invariance; Definition 1.5 and Proposition 1.8) and the Axiom of Duality (Boolean readout, coarse–graining noncontextuality, monoidal composition on the AD–visible slice; Axiom 1.10 and Lemma 14.10), together with continuity (Section 14.3.3). Then the only calibration–compatible reparametrization $p = g \circ v_0$ (cf. Lemma 14.17) that satisfies Stage 2 symmetries and Stage 3 composition laws is the identity $g(x) = x$. Equivalently, for all projectors P and unit vectors $|\psi\rangle$,*

$$p(P \mid \psi) \;=\; v_0(P) \;=\; \langle\psi|\, P \,|\psi\rangle\,. \tag{14.87}$$

Proof sketch (fixed–point route). (i) Stage 2 symmetry constraints. By phase–indifference and parasymmetric swap–invariance (eqs. (14.4) and (14.39) and Lemma 14.9; Tetra: Definition 1.5 and Proposition 1.8), the pointer weights obey $p_k = f(|c_k|)$ and are equiprobable on equal–modulus branches.

(ii) Monoidal composition ⇒ multiplicativity. On the AD–visible slice singled out by the Axiom of Duality (Axiom 1.10), product factorization (Monoidal Duality; Lemma 14.10) enforces $f(rs) = f(r)f(s)$ for $r, s \in [0,1]$. With continuity (Section 14.3.3) this yields $f(r) = r^\alpha$ for some $\alpha > 0$.

(iii) Equal–branch calibration ⇒ $\alpha = 2$. Equiprobability for n equal–modulus branches gives $f(1/\sqrt{n}) = 1/n$ (Stage 2), hence $(1/\sqrt{n})^\alpha = 1/n$ and $\alpha = 2$ (see also Theorem 14.3 for the structural perspective).

(iv) Calibration compatibility ⇒ fixed point. By Lemma 14.17, any operational assignment consistent with the Stage 2/AD structure must be a reparametrization $p = g \circ v_0$ with a state–independent, continuous g. For pointer events one has $v_0(P_k) = |c_k|^2$ via the Emergent Inner Product induced by the Fourfold Inter–Modal Correlator (Section 14.3.2, eq. (14.38), and Definition 4.9). Since $p_k = f(|c_k|) = |c_k|^2 = v_0(P_k)$, it follows that $g(x) = x$ on a dense subset of $[0, 1]$; continuity extends this to all $x \in [0, 1]$.

(v) Conclusion. Thus $p \equiv v_0$ on projectors, i.e. (14.87) holds. Together with the Gleason/Busch representation (Stage 3; Theorems 14.3 and 14.4) on the AD–visible slice (Axiom 1.10), this identifies $v_0(P) = \langle\psi| P |\psi\rangle$ in the emergent/GNS representation, establishing non–circular uniqueness at the fixed point of the Tetra+AD dialectic. (For the Page–Wootters backdrop and envariance inputs see [616, 158].) □

Remark 14.20 (Physical reading)**.** The early use of "Born–valued truths'' refers only to the *state–valuation* $v_0 = \omega(\,\cdot\,)$ that organizes coherence and interference at the emergent/GNS layer (cf. Definition 14.13 and eq. (14.86), Section 14.3.2, eq. (14.38), and Definition 4.9). Stage 2–3 then show that the *only* empirically admissible way to read these valuations as frequencies is to identify them with Born probabilities, $p = \mathrm{Tr}(\rho\,\cdot\,)$ (fixed–point route ⇒ identity reparametrization $p = g \circ v_0$ with $g(x) = x$; see Lemma 14.17, Theorem 14.19, and Corollary 14.8; cf. [158, 613, 355, 616]). Operationally, the fixed point is selected by the dialectic constraints coming from the *Tetralemma of Algebra* (parasymmetric swap–invariance on the synthesis stratum; Definition 1.5 and Proposition 1.8) and the *Axiom of Duality* (Boolean readout, coarse–graining noncontextuality, and monoidal composition on the AD–visible slice; Axiom 1.10 and Lemma 14.10). Hence the programme is *reflectively consistent*, not circular: the probabilistic semantics used to interpret readouts is the fixed point selected by Tetra+AD and Holosymmetry (Theorems 14.3, 14.4 and 14.11).

Remark 14.21 (Background–independence of the Born derivation)**.** The Stage 2–3 derivation and uniqueness of the Born rule in IQR/Holosymmetry are *background–independent*. Specifically:

1. The only primitives are the effect algebra $\mathfrak{A}$, the BRST–physical/GNS state ω, and the Universal Quantum Constraint $\hat{H}_{\mathrm{univ}} |\overset{\circ}{\boldsymbol{\Psi}}\rangle = 0$ (Theorem 13.168, eq. (13.236), and Definition 13.167; no external time, metric, or foliation). The *Emergent Inner Product* and readouts arise from ω and dynamics (Section 14.3.2, eq. (14.38), and Definition 4.9), while the orthogonal pointer structure used at Stage 1 comes from relational entanglement and decoherence, not a fixed background (cf. Section 14.3.1 and eq. (14.37)).
2. Stage 2 uses *relational envariance* to enforce phase–indifference and equiprobability among equal–modulus branches (Sections 14.2.2 and 14.3.2 and eqs. (14.4) and (14.39); cf. [158]). These are Tetralemma–induced parasymmetric symmetries on the synthesis stratum (Definition 1.5 and Proposition 1.8) and depend only on algebraic tensor structure and decoherence–induced orthogonality (eq. (14.37)), not on a pre–given temporal parameter.
3. Stage 3 proves uniqueness from dialectic/monoidal axioms (Tetra+Duality), unitary covariance, additivity, coarse–graining noncontextuality, and product factorization

(Theorems 14.3 and 14.11, Lemma 14.10, and Axiom 1.10), plus the Page–Wootters relational time for global–to–local consistency ([616]). Each ingredient is algebraic/operational and independent of a fixed spacetime background.

Consequently, the identification $p(E \mid \rho) = \mathrm{Tr}(\rho E)$ is obtained as a property of the emergent BRST–physical layer itself—*not* as an axiom imposed on a pre–metric Hilbert space. This contrasts with classical uniqueness routes such as Gleason/Busch, which presuppose a given Hilbert geometry (with the σ–additivity reading stated in Remark 14.5; cf. [613, 355]).

Formal synthesis.

$$\boxed{\text{Tetralemma} + \text{Duality} + \text{Holosymmetry} + \text{UQC} \implies p_k = \langle \Psi_{\text{fin}} | \, P_k \, | \Psi_{\text{fin}} \rangle = |c_k|^2}$$

in the notation and logic of this treatise means precisely the following.

1) Tetralemma (Tetra) ⇒ Parasymmetry and envariance on the synthesis stratum. On the superpositional (Parasymmetric) stratum, the post–measurement state has the decohered, lane–orthogonal form (eq. (14.37))

$$|\Psi_{\text{fin}}\rangle = \sum_k c_k \, |s_k\rangle \, |a_k\rangle \, |e_k\rangle \, , \qquad \langle a_k \rangle \, a_\ell = \delta_{k\ell}, \quad \langle e_k \rangle \, e_\ell \approx \delta_{k\ell}.$$

Parasymmetric (Tetra) automorphisms (Definition 1.5 and Proposition 1.8) act transitively on equal–modulus branches and, together with environment–assisted invariance, imply:

- *Phase indifference:* probabilities cannot depend on the unobservable phases of c_k, hence $p_k = f(|c_k|)$ (Section 14.2.2 and eq. (14.39)).
- *Swap–envariance:* equal–modulus branches equally probable (eq. (14.4),Lemma 14.9).
- *Ancilla fine–graining:* for rational weights $|c_k|^2 = r_k/N$, equal–modulus refinement plus coarse–graining gives $p_k = r_k/N = |c_k|^2$ (Section 14.3.2 and eq. (14.42)); continuity extends this to all amplitudes:

$$\boxed{p_k = |c_k|^2} \quad \text{(Stage 2; eq. (14.43)).}$$

2) Duality (AD) ⇒ Boolean outcome layer and composition laws. Measurement is the dialectic passage Parasymmetry → Duality: the outcome stratum is Boolean (0/1) by the *Axiom of Duality* (Section 14.1.3 and Axiom 1.10). On this AD–visible slice, the admissible probability functional must satisfy the structural axioms (Section 14.3.3): (i) additivity on orthogonal joins, (ii) unitary covariance, (iii) product factorization, (iv) coarse–graining noncontextuality, (v) continuity. In particular, *Monoidal Duality* enforces multiplicativity on products (Lemma 14.10), which for rank–1 branches implies $f(rs) = f(r)f(s)$ and (with continuity) $f(r) = r^\alpha$; equiprobability for n equal branches fixes $\alpha = 2$ (Theorem 14.3). Thus AD + the axioms lift the Stage–2 rule to *all projectors*:

$$p(P \mid \psi) = \langle \psi | \, P \, | \psi \rangle \quad \text{(Born; Theorems 14.3, 14.4 and 14.11).}$$

3) Holosymmetry ⇒ operational pipeline and decoherence orthogonality. Holosymmetry provides the operational realization:

- Stage 1 prepares the orthogonal pointer basis and global records (Section 14.3.1).
- The projector onto outcome k, $P_k = \mathbb{I}_S \otimes |a_k\rangle\langle a_k| \otimes \mathbb{I}_E$, yields (Section 14.3.2)

$$p_k = \langle \Psi_{\text{fin}}|\, P_k\, |\Psi_{\text{fin}}\rangle = |c_k|^2\ \langle s_k\rangle\, s_k\ \langle a_k\rangle\, a_k\ \langle e_k\rangle\, e_k = |c_k|^2,$$

  using pointer orthonormality and decoherence–induced orthogonality (eq. (14.45)).
- *Selection lens:* any non–Born assignment produces algebraic "twist'' in the commutator–lensing diagnostics and is suppressed by Stage–II Holosymmetric Selection (Sections 13.2.7 and 14.3.3).

4) UQC ⇒ global–to–local consistency with emergent time. The *Universal Quantum Constraint* $\hat{H}_{\text{univ}}\,|\overset{\circ}{\boldsymbol{\Psi}}\rangle = 0$ (Theorem 13.168, eq. (13.236), and Definition 13.167) and Page–Wootters conditioning ensure that local dynamics are read from a timeless global state. Requiring equality of globally conserved expectations and locally inferred values for *arbitrary* spectra and amplitudes forces $p_i = \phi(|c_i|^2) = |c_i|^2$ (Part 1 consistency; Sections 14.3.3 and 14.3.4). Thus non–quadratic rules violate conservation inherited from $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$ and are excluded.

5) Probability = relational self–coherence via the Emergent Inner Product. In IQR, the norm is not primitive; it is *emergent* from the Fourfold Inter–Modal Correlator $\hat{G}_\Lambda$ (Definition 4.9):

$$\|\,|\psi\rangle\,\|^2 = \langle\psi\rangle\,\psi := \lim_{x'\to x} \Big\langle \overset{\circ}{\boldsymbol{\Psi}}\Big|\, \hat{G}_\Lambda(x,x')\,[\psi^*,\psi]\,\Big|\overset{\circ}{\boldsymbol{\Psi}}\Big\rangle_{\text{phys}} \quad (\textit{eq. (14.38)}).$$

Identifying p_k with $\langle \Psi_{\text{fin}}|\, P_k\, |\Psi_{\text{fin}}\rangle$ (above) and evaluating with this *Emergent Inner Product* gives $p_k = |c_k|^2$ as the *relational self–coherence* of branch k (Section 14.3.2).

6) No circularity: two–tier semantics closes at a fixed point. Holosymmetric constructions use only the linear *state–valuation* $v_0(E) = \omega(E)$ (Definition 14.13 and Proposition 14.15). Stages 2–3 then constrain any admissible operational rule to $p = g \circ v_0$ and force $g(x) = x$ (fixed–point identification; Definition 14.14, Proposition 14.16, and Theorem 14.19). Hence the Born probabilities coincide numerically with the Born–valued truths supplied by the emergent/GNS layer (Corollary 14.8).

Putting it together.

- **Tetra** supplies envariance and equiprobability on the synthesis stratum ⇒ $p_k = |c_k|^2$ for branches (Stage 2; Definition 1.5, Proposition 1.8, and Lemma 14.9).
- **AD** imposes Boolean readout and composition axioms ⇒ uniqueness of the quadratic rule for all projectors (Stage 3; Axiom 1.10, Lemma 14.10, and Theorems 14.3, 14.4 and 14.11).

- **Holosymmetry** provides the operational preparation (pointer orthogonality), the projector reading, and dynamical suppression of non–Born rules (Sections 13.2.7 and 14.3.3).
- **UQC** enforces global–to–local conservation with emergent time, excluding alternatives by consistency (Section 14.3.4).

Therefore,

$$\boxed{p_k = \langle \Psi_{\text{fin}} |\, P_k \,| \Psi_{\text{fin}} \rangle = |c_k|^2}$$

with $|c_k|^2$ evaluated by the *Emergent Inner Product* as the branch's *relational self–coherence*—fixed by Stage 2 symmetries, made unique by Stage 3, and background–independent within the IQR/Holosymmetry architecture.

Relationship to the Good–clock

1) UQC (where the clock is indispensable). UQC is enforced by the universal constraint and its deparametrized split

$$\hat{H}_{\text{univ}} = \hat{p}_t \otimes \mathbf{1}_{\text{sub}} + \mathbf{1}_{\text{clock}} \otimes \hat{H}_{\text{sub}} + \hat{H}_{\text{int}},$$
$$\hat{H}_{\text{univ}} \Big| \mathring{\mathbf{\Psi}} \Big\rangle = 0 \quad (\textit{eqs. (13.236) and (13.237) and Theorem 13.168}).$$

A good–clock C (Definition 13.158) provides the *relational time lane* and the *Stone–covariant* readouts $\{\Pi_t\}$ so that conditioning is gauge–clean and TT–safe (Lemmas 13.160 and 13.161). This yields the subsystem state at clock time t:

$$\varrho_{\text{sub}|t} = \frac{\Pi_t \, \varrho \, \Pi_t}{\text{Tr}(\Pi_t \varrho)} \quad \text{(PVM)}; \qquad \text{POVM: via Naimark/Kraus, Corollary 13.163.}$$

On certified plateaues $\mathfrak{P}$, the PVM case gives *unitary* conditional dynamics

$$i\hbar\, \partial_t \, |\Psi_{\text{sub}}(t)\rangle = \hat{H}^{\text{eff}}_{\text{sub}}(t) \, |\Psi_{\text{sub}}(t)\rangle \,,$$
$$\hat{H}^{\text{eff}}_{\text{sub}}(t) = \hat{H}_{\text{sub}} + \langle t| \, \hat{H}_{\text{int}} \, |t\rangle \quad (\textit{Proposition 13.162 and eq. (13.227)}),$$

and the POVM case adds a small, controllable dissipator $i\hbar\, \mathcal{D}_t[\cdot] \to 0$ in the sharp limit (Corollary 13.163). *Takeaway:* UQC says "global timeless"; the good–clock makes it "locally dynamical at t".

2) Holosymmetry (where the readout happens). Holosymmetry supplies the *operational effects* and the interferometric pipeline Evolve $\to$ Interfere $\to$ Re–evolve with explicit control block $\mathbf{U}_{\text{ctrl}}$ and holosymmetric devices $\mathbf{U}^{(\text{sym/asym})}_{\text{holo}}$ (Sections 2.4 and 3.3). The good–clock ensures:

- *Relational commutation:* $[A, \Pi_t] = 0$ for $A \in \mathcal{A}$ (Definition 13.158(ii)), so holosymmetric readouts at time t evaluate *subsystem* observables on $\varrho_{\text{sub}|t}$.
- *TT–nullity:* clock operations do not source spin–2 radiation, so interferometric fringes (and selection) remain in the intended lanes (Lemma 13.161).

Thus, the Born valuation of any holosymmetric effect E is taken *at the conditioned state*:

$$v(\rho; E) \;\rightsquigarrow\; p_t(E) := \text{Tr}\Big(\varrho_{\text{sub}|t}\, E\Big).$$

3) Duality (products, covariance) — and why the clock does not spoil it. Duality supplies multiplicativity on products and unitary covariance for the valuations used in Stage 2/3 of the Born derivation (AD Boolean/Twofold slice; Axiom 1.10, Monoidal Duality Lemma 14.10). Because the good–clock sits in a separate typed lane and $[A, \Pi_t] = 0$, the *product and covariance axioms* apply directly to $\varrho_{\mathrm{sub}|t}$ just as they do to ρ (no anomalies from the clock sector). In particular, for rank–1 events,

$$p_t\Big(\,|\phi\rangle\langle\phi|\;\Big|\;\psi(t)\Big) = \Big|\,\langle\phi\rangle\,\psi(t)\Big|^2 \quad\Rightarrow\quad p_t(P) = \langle\psi(t)|\,P\,|\psi(t)\rangle$$

(extension by additivity/continuity on the AD slice; Section 14.3.3).

This matches the uniqueness steps (Theorems 14.3, 14.4 and 14.11).

4) Tetralemma (coarse–graining logic) — safeguarded by the clock's hygiene.

The Tetralemma's additivity/coarse–graining noncontextuality is evaluated on *orthogonal post–measurement branches*. Good–clock hygiene (BRST–closed, TT–safe; Definition 13.158(iii)) and spectral support (Definition 13.158(iv)) ensure those branches live in $\mathcal{H}_{\mathrm{sub}}$ at fixed t without gauge/TT contamination, so the Stage 2 envariance and Stage 3 uniqueness arguments go through unchanged on $\varrho_{\mathrm{sub}|t}$ (Tetralemma invariances Definition 1.5 and Proposition 1.8 together with AD Booleanity Axiom 1.10).

Bridge identity (what the clock ultimately guarantees). For a measurement in the $\{|s_k\rangle\}$ basis at clock time t, with $|\Psi_{\mathrm{sub}}(t)\rangle = \sum_k c_k(t)\,|s_k\rangle$ (PVM case),

$$\boxed{\,p_k(t) = \mathrm{Tr}\Big(\varrho_{\mathrm{sub}|t}\;|s_k\rangle\langle s_k|\,\Big) = |c_k(t)|^2\,}$$

and for POVM clocks the same holds up to $i\hbar\,\mathcal{D}_t[\cdot]$, which vanishes in the sharp limit (Corollary 13.163). Portal deformations preserve this at first order by Stage II Lipschitz stability (Proposition 13.164).

In short: the good–clock is the UQC–to–lab adapter. It turns the global, timeless law into a gauge–clean, TT–safe, time–conditioned state $\varrho_{\mathrm{sub}|t}$ on which Holosymmetry, Duality, and the Tetralemma act exactly as required to deliver the Stage 2/3 outcome $p_k = |c_k|^2$ (Section 14.3.2 and Theorem 14.11).

Chapter 15

Quantum Thermodynamics

15.1 The Laws of Quantum Thermodynamics

15.1.1 Foundational Definitions and Postulates

Before entering the proof, we collect the minimal infrastructure that fixes the algebraic arena, the emergent geometric data, the *Good-clock* time gauge, and the measurement semantics on which all thermodynamic statements rest. The underlying stage is the PQS continuum $\mathcal{X}_q$ equipped with the abstract C*-algebra of relational observables $\mathcal{A}(\mathcal{C}_d)$ (Definition 1.32), taken *prior* to any Hilbert-space representation. A coherent Quantum Holo-State $\omega^{\mathring{\mathbf{\Psi}}}$ (Section 1.4)—a normalized positive linear functional $\mathcal{A}(\mathcal{C}_d)\to\mathbb{C}$—selects the GNS representation in which the vector state $|\mathring{\mathbf{\Psi}}\rangle$ and the Emergent Hilbert Space $\mathcal{H}_{\text{em}}$ appear (Section 1.4.1 and Definitions 1.18 and 4.287; cf. [230, 30]). Within this representation, *relational subsystems* are specified by subalgebras $\mathcal{A}_X \subset \mathcal{A}(\mathcal{C}_d)$; their states are restrictions $\omega_X := \omega^{\mathring{\mathbf{\Psi}}}|_{\mathcal{A}_X}$ with density matrices ρ_X, and all operational predictions are Born valuations $v(\rho; E) = \mathrm{Tr}(\rho E)$ (Definition 1.18).

Modal fibres, correlator, and emergent inner product. At each $x \in \mathcal{X}_q$ the modal fibre $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)} = \mathrm{span}\{|0\rangle_x, |1\rangle_x, |p\rangle_x, |s\rangle_x\}$ carries the Fourfold Dialectic algebra generated by $\{\hat{\mathcal{F}}_\alpha\}_{\alpha\in\{0,1,p,s\}}$ (Section 1.2.7). The Fourfold Inter-Modal Correlator $\hat{G}_\Lambda^{\alpha\beta}(x,x')$ together with the gauge-covariant transporter $\mathcal{U}(x\leftarrow x';\gamma)$ (Definition 4.9 and eqs. (4.11) and (4.21)) yields, in the coincident limit, the fibre inner product

$$\langle\psi_1 \mid \psi_2\rangle_x = \lim_{x'\to x} \left\langle \mathring{\mathbf{\Psi}} \right| \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2] \left| \mathring{\mathbf{\Psi}} \right\rangle \quad (eq.\ (4.85)),$$

assembling the Hilbert bundle $\mathscr{H}_{\text{mod}}$ (Definition 4.239).

Chronogeometry, Good-clock, and causality. Triadic distances $(d_{\text{corr}}, d_{\text{vac}}, d_{\text{spec}})$ define the composite metric D (Definition 4.476), whose Fréchet second variation produces the operator-valued metric $\hat{g}_{\mu\nu}$ (Definition 5.1). As a formal recall, all variations and contractions are taken with the *emergent inner product* induced by the Fourfold inter-modal correlator

(coincident-limit metric), cf.

$$\Big\langle \psi_1 \mid \psi_2 \Big\rangle_x = \lim_{x' \to x} \Big\langle \overset{\circ}{\mathbf{\Psi}} \Big| \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2] \Big| \overset{\circ}{\mathbf{\Psi}} \Big\rangle, \tag{15.1}$$

(Definition 4.9 and eq. (4.6)). Alongside the temporal triad $(t_{\mathrm{corr}}, t_{\mathrm{vac}}, t_{\mathrm{spec}})$ one obtains the operator-valued temporal tensor $\hat{t}_{\mu\nu}$ and the unified chronogeometric tensor $\hat{\Sigma}_{\mu\nu}$ with expectation $\bar{\Sigma}_{\mu\nu}$ (Definitions 5.17 and 5.26); expectations are taken in the Holo-State and are gauge-covariant under the operator-bundle connection recalled in Section 6.12.2.

On each diagnostic realized reference configuration we fix a *Good-clock* $C = (\hat{T}, U_t)$ aligned to the $\bar{\Sigma}$-time flow, with low temporal holonomy and Page–Wootters readability (Definition 13.158 and Propositions 4.556 and 13.172). Alignment is meant in the relational Levi–Civita sense: time-lines are integral curves of the $\bar{\Sigma}$-compatible, torsion-free connection $\nabla^{(\mathrm{rel})}$ (symbols $\Gamma^{\lambda}_{\mu\nu}[\bar{\Sigma}]$), and parallel transport along them uses the split covariant derivative

$$\nabla_\mu \;:=\; \nabla^{(\mathrm{rel})}_\mu \;+\; \Big(\partial_\mu + i[\mathcal{A}_\mu, \cdot]\Big), \tag{15.2}$$

(Definition 6.7, eq. (6.22), and Section 6.12.2). Small temporal holonomy is quantified via the gauge-covariant transporter

$$\mathcal{U}(x \leftarrow x'; \gamma) \;=\; \mathcal{P}\, \exp\!\Big(-\int_\gamma \mathcal{A}\Big), \tag{15.3}$$

on clock worldlines (eqs. (4.11) and (4.16)).

Microcausality and spectral positivity are imposed with respect to $\bar{\Sigma}$, yielding Σ-causal support and "no-law-signalling" constraints (Corollary 5.145 and Proposition 5.143); in particular, apex safety and branchwise cone emanation from the PQS remain intact under holosymmetric lifts (formal recall: Proposition 5.157). Readouts on the diagnostic realized reference configuration are then standard Born valuations $v(\rho; E) = \mathrm{Tr}(E\rho)$ consistent with the Axiom of Duality/Twofold restriction at measurement time (Definition 1.18 and Section 14.1.3), and the overall construction respects the Universal Quantum Constraint at the cosmological level (Theorem 13.168).

Physical realization and typed structure. Thermodynamic statements are evaluated on a certified plateau $\mathfrak{P}$ and on the BRST-physical space $\mathcal{H}_{\mathrm{phys}} := H^0(\hat{Q}_{\mathrm{BRST}})$ (Definition 9.58 and Section 12.3; cf. [442, 445, 106, 483]). Band tags $(\lambda_{\mathrm{UV}}, \ell)$ and the UV window $P_x(\lambda_{\mathrm{UV}})$ are fixed on $\mathfrak{P}$ (Definitions 7.26 and 9.10), and BRST hygiene guarantees Nielsen-flat readouts (Lemma 9.54). All operators and states are projected to $\mathcal{H}_{\mathrm{phys}}$,

$$\rho_{\mathrm{phys}} := \mathsf{P}_{\mathrm{phys}} \rho\, \mathsf{P}_{\mathrm{phys}}, \qquad O_{\mathrm{phys}} := \mathsf{P}_{\mathrm{phys}}\, O\, \mathsf{P}_{\mathrm{phys}}, \tag{15.4}$$

and readouts are Born valuations $v(\rho; E) = \mathrm{Tr}(E\rho)$ (Definition 1.18) compatible with the Axiom of Duality/Twofold restriction at measurement time (Section 14.1.3).

Typed projectors $\Pi^{[\chi]}$ (passing the permit ledgers) and parasymmetric meters $v^{(a);[\chi]}_{\mathrm{Synthesis}}$ provide the diagnostic layer (Definitions 9.122, 9.142 and 9.143). Meters act on typed subspaces via

$$v^{(a);[\chi]}_{\mathrm{Synthesis}}(\rho) = \mathrm{Tr}\Big(E^{(a);[\chi]}_{\mathrm{Synthesis}}\, \Pi^{[\chi]} \rho_{\mathrm{phys}} \Pi^{[\chi]}\Big), \tag{15.5}$$

with $E^{(a);[\chi]}_{\text{Synthesis}}$ the Synthesis effect associated to the chosen parasymmetric instrument (Definitions 1.18 and 9.143). The coherence resolution of the identity on $\mathcal{H}_{\text{phys}}$,

$$\mathbf{1}_{\text{phys}} = \mathcal{P}^{\text{sym}}_{\text{vac}} + \mathcal{P}^{\text{coh}}_{\text{dark}} + \mathcal{P}^{\text{coh}}_{\text{vis}}, \tag{15.6}$$

is used without mixing channels, enforced by

$$[\Pi^{[\chi]}, \mathcal{P}^{\text{sym}}_{\text{vac}}] = [\Pi^{[\chi]}, \mathcal{P}^{\text{coh}}_{\text{dark}}] = [\Pi^{[\chi]}, \mathcal{P}^{\text{coh}}_{\text{vis}}] = 0 \quad (\textit{eq. (13.149)}). \tag{15.7}$$

The visible/geo/matter split employs the vacuum–tangent projectors $\hat{P}_T, \hat{P}_{\text{exc}}$ to separate Gaiamiti $\hat{\mathbf{\Phi}}$ and Tantrakyma $\hat{\mathbf{\Theta}}$ sectors (eqs. (9.119) and (9.122)), so that for any fluctuation $\delta\hat{\Sigma}$ one has

$$\delta\hat{\Sigma} = \hat{P}_T\,\delta\hat{\Sigma} \;\oplus\; \hat{P}_{\text{exc}}\,\delta\hat{\Sigma}, \qquad \hat{P}_T\,\delta\hat{\Sigma} \in \Phi\Phi, \quad \hat{P}_{\text{exc}}\,\delta\hat{\Sigma} \in \Theta\Theta. \tag{15.8}$$

Band limitation is implemented componentwise,

$$O^{(\lambda_{\text{UV}})} := P_x(\lambda_{\text{UV}})\, O_{\text{phys}}\, P_x(\lambda_{\text{UV}}), \tag{15.9}$$

with all typed meters and projectors taken on $O^{(\lambda_{\text{UV}})}$ and understood to be BRST-safe and band-stable on $\mathfrak{P}$ (Definitions 7.26 and 9.10 and Lemma 9.54).

Measurement semantics (clocked). Interrogations are enacted via the holosymmetric interferometer $\mathbf{U}^{(\text{sym})}_{\text{holo}}$ and its path–scattering template (Definition 2.123 and Section 14.1.1). With

$$\mathbf{U}_{\text{ctrl}} = |\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}^{A_G B_N}_{\text{RR}} + |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}^{A_N B_G}_{\text{RR}}, \qquad \mathbf{U}^{(\text{sym})}_{\text{holo}} = \mathbf{U}_{\text{Had}}\,\mathbf{U}_{\text{ctrl}}\,\mathbf{U}_{\text{Had}},$$

the bright/dark port probabilities for input $|\Psi_{\text{init}}\rangle = |\mathbf{0}\rangle_c \otimes |\Psi_{\text{RR}}\rangle$ are

$$p_0 = \tfrac{1}{4}\left\|(\hat{\mathbf{U}}^{A_G B_N}_{\text{RR}} + \hat{\mathbf{U}}^{A_N B_G}_{\text{RR}})\,|\Psi_{\text{RR}}\rangle\right\|^2,$$
$$p_1 = \tfrac{1}{4}\left\|(\hat{\mathbf{U}}^{A_G B_N}_{\text{RR}} - \hat{\mathbf{U}}^{A_N B_G}_{\text{RR}})\,|\Psi_{\text{RR}}\rangle\right\|^2$$
(*Corollary 3.3and eq. (3.1)*).

The asymmetric, tomographic device is

$$\mathbf{U}^{(\text{asym})}_{\text{holo}} = \mathbf{U}_{\text{read}}\,\mathbf{U}_{\text{ctrl}}\,\mathbf{U}_{\text{prep}}, \qquad \mathbf{U}_{\text{prep}}, \mathbf{U}_{\text{read}} \in \text{SU}(2),$$

with fringes exposed via the commutator lens

$$[\mathbf{U}_{\text{Had}}, \mathbf{U}_{\text{ctrl}}] = \tfrac{1}{\sqrt{2}}\begin{pmatrix} 0 & \hat{\mathbf{U}}^{A_G B_N}_{\text{RR}} - \hat{\mathbf{U}}^{A_N B_G}_{\text{RR}} \\ \hat{\mathbf{U}}^{A_N B_G}_{\text{RR}} - \hat{\mathbf{U}}^{A_G B_N}_{\text{RR}} & 0 \end{pmatrix} \quad (\textit{Section 14.1.1}).$$

All time-evolved quantities are parameterized by a Good-clock $C = (\hat{T}, U_t)$ aligned with $\bar{\Sigma}$ (Definition 13.158 and Proposition 13.172). Probabilities are Born valuations $v(\rho; E) = \text{Tr}(E\rho)$ (Definition 1.18) with the GNS inner product supplied by the Fourfold correlator,

$$\langle\psi_1 \mid \psi_2\rangle_x = \lim_{x'\to x} \langle\overset{\circ}{\mathbf{\Psi}}|\,\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2]\,|\overset{\circ}{\mathbf{\Psi}}\rangle \quad (\textit{Definition 4.9and eq. (4.85)}).$$

Twofold readouts demanded by the Axiom of Duality use orthogonal projectors $P_{\mathbf{0}}, P_{\mathbf{1}}$ on the Twofold sector with $P_{\mathbf{0}} + P_{\mathbf{1}} = \mathbf{1}_{\text{Twofold}}$ and $P_{\mathbf{0}} P_{\mathbf{1}} = 0$, yielding

$$p_{\mathbf{i}} = \mathrm{Tr}(\rho\, P_{\mathbf{i}}) \quad (\textit{Section } 14.1.3).$$

Open-system reductions adopt the CP-divisible/unital framework on the active band, with GKLS generator

$$\begin{aligned}\dot{\rho}_A = \mathcal{L}[\rho_A] = -i[H_{A,\text{eff}}, \rho_A] \\ + \sum_\alpha \gamma_\alpha \Big(L_\alpha \rho_A L_\alpha^\dagger - \tfrac{1}{2}\{L_\alpha^\dagger L_\alpha, \rho_A\} \Big)\end{aligned}$$

(*Propositions* 15.45*and* 15.57*and Theorem* 15.35)

and environment-induced superselection as standard [617, 441, 170].

Working Postulates (P0–P7). We adopt the following minimal set, each anchored to prior structures:

(P1) Arena & State. The algebraic arena is $\mathcal{A}(\mathcal{C}_d)$ with state $\omega^{\mathring{\Psi}}$ selecting the GNS triple $(\pi_\omega, \mathcal{H}_{\text{em}}, |\Omega_\omega\rangle)$; operational numbers are Born valuations $v(\rho; E) = \mathrm{Tr}(\rho E)$ (Section 1.4.1 and Definition 1.18; cf. [230, 30]).

(P2) Regularity & Gauge. The correlator has a well-defined coincident limit (regularity near the diagonal) and transforms covariantly; the induced fibre metric and all distances/times are gauge-invariant (eqs. (4.16) and (4.85) and Proposition 4.477).

(P3) Chronogeometric Causality. Dynamics and two-point functions respect $\bar{\Sigma}$-microcausality and spectral positivity; law-interference never enlarges causal support (Definition 5.26 and Corollary 5.145).

(P4) Diagnostic realized reference configuration (with Good-clock). All thermodynamic theorems are asserted on $\mathfrak{P}$ and $\mathcal{H}_{\text{phys}}$, equipped with a Good-clock $C = (\hat{T}, U_t)$ aligned to $\bar{\Sigma}$ and satisfying Definition 13.158; typed objects $\Pi^{[\chi]}$ meet permit conditions and Nielsen/BRST safety (Definitions 9.58 and 9.122 and Theorem 9.351; cf. [442, 120]).

(P5) Sector Split. The geometry/matter decomposition $\mathcal{H}_{\text{mod}}^{(\mathfrak{B}_x)} = \mathcal{E}_{\text{geo}}(x) \oplus \mathcal{E}_{\text{exc}}(x)$ is enforced by $\hat{P}_T, \hat{P}_{\text{exc}}$; thermodynamic exchanges are traced on subsystem algebras $\mathcal{A}_X$ following this split (eqs. (9.119) and (9.122)).

(P6) Measurement Consistency (clocked). Laboratory meters are holosymmetric path–scattering probes with TT/typed hygiene; local outcomes reduce to Twofold projectors and are evaluated by Born valuations, with time evolution parameterized by the Good-clock C (Definitions 1.18 and 2.123 and Section 14.1.3).

(P7) Energy Conservation. The Total Relational Action is invariant under emergent-time translations (Good-clock aligned), yielding a conserved $\hat{H}_{\text{univ}}$ that decomposes into subsystems and interactions (Section 15.1.3).

(P8) Admissible Processes. Coarse-grained evolutions relevant to thermodynamics are CP-divisible and unital on the appropriate pointer channel $\mathbb{E}_A$, ensuring entropy monotonicity; the small-coupling limit recovers unitary coherence (Definition 15.27, Theorem 15.50, Lemma 15.51, and Proposition 15.52; cf. [617, 441]).

(P0) Arena & State. The algebraic arena is $\mathcal{A}(\mathcal{C}_d)$ endowed with the Holo-State $\omega^{\mathring{\mathbf{\Psi}}}$, which selects the GNS triple $(\pi_\omega, \mathcal{H}_{\mathrm{em}}, |\Omega_\omega\rangle)$; expectations are evaluated as

$$\omega(A) = \Big\langle \Omega_\omega \Big| \, \pi_\omega(A) \, \Big| \Omega_\omega \Big\rangle \qquad (\textit{Section } 1.4.1;\ \text{cf. } [230,\ 30]).$$

Operational numbers are Born valuations $v(\rho; E) = \mathrm{Tr}(E\rho)$ for effects E (including projectors and POVM elements) on $\mathcal{H}_{\mathrm{em}}$ (Definition 1.18). The inner product on $\mathcal{H}_{\mathrm{em}}$ is the emergent one supplied by the Fourfold inter-modal correlator,

$$\langle \psi_1 \mid \psi_2 \rangle_x = \lim_{x' \to x} \Big\langle \mathring{\mathbf{\Psi}} \Big| \, \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2] \, \Big| \mathring{\mathbf{\Psi}} \Big\rangle \qquad (\textit{Definition } 4.9 \textit{and eq. } (4.85)),$$

so that all probabilities and expectation values used in thermodynamic statements are computed in the GNS/emergent arena consistently with $\mathcal{A}(\mathcal{C}_d)$ and $\omega^{\mathring{\mathbf{\Psi}}}$.

(P1) Regularity & Gauge. The Fourfold inter-modal correlator $\hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})$ admits a well-defined coincident limit $x' \to x$, so the emergent inner product

$$\langle \psi_1 \mid \psi_2 \rangle_x = \lim_{x' \to x} \Big\langle \mathring{\mathbf{\Psi}} \Big| \, \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})[\psi_1^*, \psi_2] \, \Big| \mathring{\mathbf{\Psi}} \Big\rangle$$

exists and equals the fibre metric on $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ (eq. (4.85), Assumption 4.21, Proposition 4.28, and Definition 4.9). Parallel transport is implemented by

$$\mathcal{U}(x \leftarrow x'; \gamma) = \mathcal{P} \exp\Big(-\int_\gamma \mathcal{A} \Big),$$

and gauge transformations act by

$$\mathcal{A} \mapsto g\,\mathcal{A}\,g^\dagger + g\,\mathrm{d}g^\dagger, \qquad \mathcal{U} \mapsto g\,\mathcal{U}\,g^\dagger,$$

so that $\hat{G}_\Lambda$ is covariant and the induced inner product and fibre metric are gauge-invariant (eq. (4.16), Definition 4.9, and Remark 4.29). Distances and times built from the triadic structure $(d_{\mathrm{corr}}, d_{\mathrm{vac}}, d_{\mathrm{spec}})$ and $(t_{\mathrm{corr}}, t_{\mathrm{vac}}, t_{\mathrm{spec}})$ descend to the composite metric D and obey the metric and invariance axioms, in particular

$$D\Big(g\,\rho\,g^\dagger,\ g\,\sigma\,g^\dagger\Big) = D(\rho, \sigma),$$

with the same gauge-invariance for the temporal functionals (Proposition 4.477).

(P2) Chronogeometric Causality. The unified chronogeometric tensor $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ fixes the causal cones $\mathcal{V}_{\pm}(\bar{\Sigma})$ and the associated $\bar{\Sigma}$-compatible transport (Definition 5.26). For any two localized observables $\hat{\mathfrak{O}}(f_A)$, $\hat{\mathfrak{O}}(f_B)$ with test functions supported in regions $A, B \subset \mathcal{X}_q$ that are spacelike with respect to $\bar{\Sigma}$ (denoted $A \perp_{\cap} B$), microcausality holds:

$$\left[\hat{\mathfrak{O}}(f_A),\, \hat{\mathfrak{O}}(f_B)\right] = 0 \quad \text{whenever} \quad A \perp_{\cap} B,$$

and the Pauli–Jordan support lies within $\mathcal{V}_{+}(\bar{\Sigma}) \cup \mathcal{V}_{-}(\bar{\Sigma})$, ensuring spectral positivity of two-point functions (positive-type distributions with spectral measure supported on the forward cone). Under holosymmetric law-superposition, causal support does not enlarge: for branchwise future sets $J^{+}_{(k)}(A)$ one has

$$\operatorname{supp}\left(\hat{\mathfrak{O}}(f_A) \to \hat{\mathfrak{O}}(f_B)\right) \;\subseteq\; J^{+}_{\cup}(A) := \bigcup_k J^{+}_{(k)}(A),$$

so that "law-interference'' preserves $\bar{\Sigma}$-causality and prohibits signalling outside $J^{+}_{\cup}(A)$ (Corollary 5.145).

(P3) Diagnostic realized reference configuration (with Good-clock). All thermodynamic statements are asserted on a certified plateau $\mathfrak{P}$ and within the BRST-physical sector $\mathcal{H}_{\text{phys}}$, with a Good-clock $C = (\hat{T}, U_t)$ aligned to $\bar{\Sigma}$ (Definitions 9.58 and 13.158). Time evolution and readouts are taken on $\mathcal{H}_{\text{phys}}$ by implicit projection with P_{phys},

$$\rho \;\mapsto\; \mathsf{P}_{\text{phys}} \rho\, \mathsf{P}_{\text{phys}}, \qquad O \;\mapsto\; \mathsf{P}_{\text{phys}}\, O\, \mathsf{P}_{\text{phys}},$$

and are parameterized by the Good-clock unitary U_t (Heisenberg/Schrödinger pictures equivalent), with the conditional Schrödinger equation implementing Page–Wootters readability (Proposition 13.172). Alignment is with the $\bar{\Sigma}$-compatible, torsion-free transport used throughout, so that clock worldlines follow the relational Levi–Civita flow (Definition 13.158). Typed projectors $\Pi^{[\chi]}$ satisfy the permit-ledger conditions and are employed for all typed assertions (Definition 9.122); their associated meters are Nielsen/BRST safe, i.e. gauge-parameter independent on $\mathfrak{P}$ (Theorem 9.351; cf. [442, 120]).

(P4) Sector Split. On each locus x, the geometry/matter decomposition $\mathcal{H}^{(\mathfrak{B}_x)}_{\text{mod}} = \mathcal{E}_{\text{geo}}(x) \oplus \mathcal{E}_{\text{exc}}(x)$ is implemented by the vacuum–tangent projectors $\hat{P}_T$ and $\hat{P}_{\text{exc}}$, with $\hat{P}_T^2 = \hat{P}_T$, $\hat{P}_{\text{exc}}^2 = \hat{P}_{\text{exc}}$, $\hat{P}_T \hat{P}_{\text{exc}} = 0$, and $\hat{P}_T + \hat{P}_{\text{exc}} = \mathbf{1}$ on the tangent/excitation bundle (eqs. (9.119) and (9.122)). Acting on unified fluctuations and fields,

$$\delta\hat{\Sigma} = \hat{P}_T\, \delta\hat{\Sigma} \;\oplus\; \hat{P}_{\text{exc}}\, \delta\hat{\Sigma}, \qquad \delta\hat{\mathbf{F}} = (\delta\hat{\mathbf{\Phi}}, \delta\hat{\mathbf{\Theta}}),$$

with $\hat{P}_T\, \delta\hat{\Sigma} \in \Phi\Phi$ (Gaiamiti sector) and $\hat{P}_{\text{exc}}\, \delta\hat{\Sigma} \in \Theta\Theta$ (Tantrakyma sector), consistently with the modal/GNS kinematics and the split covariant derivative on $(\mathcal{X}_q, \bar{\Sigma})$ (Definition 6.184 and Section 6.12.2).

Thermodynamic exchanges are traced on subsystem algebras $\mathcal{A}_X$ honouring this split: for a subsystem X with Hamiltonian decomposition $\hat{H} = \hat{H}_X + \hat{H}_{\bar{X}} + \hat{H}_{X\bar{X}}$ (Definition 15.11), the Good-clock Heisenberg balance yields

$$\frac{dU_X}{dt} = \left\langle i[\hat{H}_{X\bar{X}}, \hat{H}_X] \right\rangle - \left\langle \partial_t \hat{H}_X \right\rangle \quad (\textit{Lemma } 15.12),$$

with heat current and power identified as in Definition 15.14 and integrated to the first-law statement $\Delta U_X = Q_X - W_X$ (Proposition 15.15 and Corollary 15.17). All such balances may be evaluated sectorwise by inserting $\hat{P}_T$ or $\hat{P}_{\mathrm{exc}}$ on operators inside $\mathcal{A}_X$, so that geometric and matter contributions are disentangled consistently with $\hat{P}_T, \hat{P}_{\mathrm{exc}}$ (eqs. (9.119) and (9.122)).

(P5) Measurement Consistency (clocked). Laboratory meters are implemented as holosymmetric path–scattering probes, built from the controlled block and its symmetric interferometer,

$$\mathbf{U}_{\mathrm{ctrl}} = |\mathbf{0}\rangle_c\langle\mathbf{0}|_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + |\mathbf{1}\rangle_c\langle\mathbf{1}|_c \otimes \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G},$$
$$\mathbf{U}_{\mathrm{holo}}^{(\mathrm{sym})} = \mathbf{U}_{\mathrm{Had}}\, \mathbf{U}_{\mathrm{ctrl}}\, \mathbf{U}_{\mathrm{Had}} \quad (\textit{Definition 2.123}).$$

For input $|\Psi_{\mathrm{init}}\rangle = |\mathbf{0}\rangle_c \otimes |\Psi_{\mathrm{RR}}\rangle$, the bright/dark-port probabilities are

$$p_0 = \tfrac{1}{4}\left\|(\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} + \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G})\,|\Psi_{\mathrm{RR}}\rangle\right\|^2,$$
$$p_1 = \tfrac{1}{4}\left\|(\hat{\mathbf{U}}_{\mathrm{RR}}^{A_G B_N} - \hat{\mathbf{U}}_{\mathrm{RR}}^{A_N B_G})\,|\Psi_{\mathrm{RR}}\rangle\right\|^2$$
$$(\textit{Corollary 3.3and eq. (3.1)}).$$

Local outcomes reduce to Twofold projectors on the readout slice: $P_{\mathbf{0}} + P_{\mathbf{1}} = \mathbf{1}_{\mathrm{Twofold}}$, $P_{\mathbf{0}} P_{\mathbf{1}} = 0$, with probabilities

$$p_{\mathbf{i}} = \mathrm{Tr}(\rho\, P_{\mathbf{i}}) \quad (\textit{Section 14.1.3}).$$

All probabilities and expectation values are Born valuations $v(\rho; E) = \mathrm{Tr}(E\rho)$ on $\mathcal{H}_{\mathrm{em}}$ (Definition 1.18), and time dependence is parameterized by the Good-clock $C = (\hat{T}, U_t)$ aligned with $\bar{\Sigma}$, i.e.

$$\rho(t) = U_t\, \rho\, U_t^\dagger, \qquad O(t) = U_t^\dagger O\, U_t \quad (\textit{Definition 13.158}).$$

TT/typed hygiene is enforced by choosing meters and readout projectors within the typed subspaces so that the Twofold reduction and Born evaluation are consistent with the holosymmetric probe construction (Definition 2.123 and Section 14.1.3).

(P6) Energy Conservation. The Total Relational Action is invariant under Good-clock translations $t \mapsto t + \epsilon$ on $\mathfrak{P}$, so the corresponding Noether charge is a conserved generator $\hat{H}_{\mathrm{univ}}$ acting on $\mathcal{H}_{\mathrm{phys}}$ (Section 15.1.3). Time evolution is parameterized by the Good-clock $C = (\hat{T}, U_t)$ aligned with $\bar{\Sigma}$, with

$$U_t = \exp\!\big(-it\, \hat{H}_{\mathrm{univ}}\big), \qquad \frac{d}{dt} O(t) = i\,[\hat{H}_{\mathrm{univ}}, O(t)] + \partial_t O(t),$$

and conserved energy $\frac{d}{dt}\langle \hat{H}_{\mathrm{univ}}\rangle = 0$ whenever $\partial_t \hat{H}_{\mathrm{univ}} = 0$. The universal Hamiltonian decomposes into subsystems and interactions,

$$\hat{H}_{\mathrm{univ}} = \sum_X \hat{H}_X + \sum_{X<Y} \hat{H}_{XY},$$

and, for a bipartition $A \cup E$, $\hat{H} = \hat{H}_A + \hat{H}_E + \hat{H}_{AE}$ (Section 15.1.3 and Definition 15.11). The Good-clock Heisenberg balance on A then yields

$$\frac{dU_A}{dt} = \big\langle i[\hat{H}_{AE}, \hat{H}_A] \big\rangle - \big\langle \partial_t \hat{H}_A \big\rangle,$$

identifying heat current and power as in Lemma 15.12 and Definition 15.14, and integrating to $\Delta U_A = Q_A - W_A$ (Proposition 15.15 and Corollary 15.17). On the cosmological diagnostic realized reference configuration, the same generator satisfies the universal constraint $\hat{H}_{\text{univ}} |\mathring{\mathbf{\Psi}}\rangle = 0$, so that the unitary family $\exp\big(-is\hat{H}_{\text{univ}}\big)$ leaves the Holo-State invariant while providing the conserved charge that organizes subsystem dynamics (Theorem 13.168).

(P7) Admissible Processes. Coarse-grained evolutions on a subsystem A are completely positive, trace-preserving maps $\{\Phi_{t\leftarrow s}\}_{t\geq s}$ that are CP-divisible and unital on the pointer channel $\mathbb{E}_A$ (Definition 15.27):

$$\Phi_{t\leftarrow r} = \Phi_{t\leftarrow s} \circ \Phi_{s\leftarrow r} \quad (r \leq s \leq t), \qquad \Phi_{t\leftarrow s}[\mathbf{1}] = \mathbf{1}.$$

On the active band, a single stroboscopic step factors as

$$\Phi_{\Delta t} = \mathbb{E}_A \circ \mathrm{Ad}_{e^{-iH_{A,\text{eff}}\Delta t}},$$

and the continuous-time limit yields a unital GKLS semigroup

$$\dot{\rho}_A = \mathcal{L}[\rho_A] = -i[H_{A,\text{eff}}, \rho_A] + \sum_\alpha \gamma_\alpha \Big(L_\alpha \rho_A L_\alpha^\dagger - \tfrac{1}{2}\{L_\alpha^\dagger L_\alpha, \rho_A\} \Big)$$

(cf. [617, 441]; Theorem 15.50). Unitality ensures entropy monotonicity: for the von Neumann entropy $S_A(t) = -\operatorname{Tr}[\rho_A(t) \log \rho_A(t)]$,

$$S(\Phi_{t\leftarrow s}(\rho)) \geq S(\rho) \quad \text{and} \quad \frac{d}{dt} S_A(t) \geq 0 \quad (\textit{Theorem } 15.50).$$

In the small-coupling regime $g = \|\hat{H}_{AE}\| \to 0$, the generator approaches the unitary form and coherence is recovered; entropy production is bounded and vanishes with the monitoring rate, consistently with

$$\mathcal{L}_g[\rho] = -i[H_{A,\text{eff}}, \rho] + \kappa(g) \Big(\mathbb{E}_A(\rho) - \rho \Big), \qquad \kappa(g) \to 0,$$

and with the coherence statements of the Third Law (Lemma 15.51 and Proposition 15.52).

Roadmap to the laws (for reference). On this scaffolding, equilibrium in the Zeroth Law is stationarity under the Σ-time flow (measured by the Good-clock) plus vanishing net relational currents (Definition 15.5 and Theorem 15.7). The First Law follows from Noether conservation and the subsystem split, with heat/work identified by the relational interaction channel (Section 15.1.3 and eq. (15.12)). The Second Law is the monotonicity of $S_A(t) = -\operatorname{Tr}[\rho_A(t) \log \rho_A(t)]$ under the unital pointer dynamics (eq. (15.13) and Theorem 15.50). The Third Law is the small-coupling coherence limit (Section 15.1.6 and Proposition 15.52).

Notational conventions. We use $\mathfrak{B}_x$ for PQS loci, $\mathcal{U}(x \leftarrow x'; \gamma)$ for parallel transport, $\hat{G}_\Lambda$ for the Fourfold correlator, $\hat{g}_{\mu\nu}, \hat{t}_{\mu\nu}$ for operator-valued (chrono)geometric tensors, $\bar{\Sigma}_{\mu\nu}$ for expectations, $\Pi^{[\chi]}$ for typed projectors, $v^{(a);[\chi]}_{\text{Synthesis}}$ for parasymmetric meters, $C = (\hat{T}, U_t)$ for the Good-clock, and $v(\rho; E) = \mathrm{Tr}(\rho E)$ for Born valuations (Definitions 1.18, 4.9, 5.17, 5.26, 9.143 and 13.158).

15.1.2 The Zeroth Law of Thermodynamics

Equilibrium between two $\bar{\Sigma}$-causally connected *and interacting* subsystems X and Y—with $\bar{\Sigma}_{\mu\nu}(x) := \omega^{\mathring{\mathbf{\Psi}}}\big(\hat{\Sigma}_{\mu\nu}(x)\big) = \big\langle \Omega_\omega \big| \pi_\omega\big(\hat{\Sigma}_{\mu\nu}(x)\big) \big| \Omega_\omega \big\rangle$—is a dynamical alignment of their joint algebra $\mathcal{A}_X \vee \mathcal{A}_Y$. It is defined by two predicates on their shared, $\hat{G}_\Lambda$–based statistics *evaluated with the emergent inner product*: (1) **stationarity** under the Good-clock flow on $\mathcal{A}_X \vee \mathcal{A}_Y$, $\partial_t \langle O \rangle_{XY} = 0$ for all $O \in \mathcal{A}_X \vee \mathcal{A}_Y$; and (2) **vanishing of the relational Noether current** between X and Y, expressed componentwise for the fundamental fields as

$$\Big\langle \hat{J}^\mu_{\mathbf{\Phi}}(X) - \hat{J}^\mu_{\mathbf{\Phi}}(Y) \Big\rangle = 0, \qquad \Big\langle \hat{J}^\mu_{\mathbf{\Theta}}(X) - \hat{J}^\mu_{\mathbf{\Theta}}(Y) \Big\rangle = 0,$$

where $\hat{J}^\mu_{\mathbf{\Phi}}$ and $\hat{J}^\mu_{\mathbf{\Theta}}$ are the $\hat{\mathbf{\Phi}}$-Gaiamiti and $\hat{\mathbf{\Theta}}$-Tantrakyma Noether currents, respectively.

Pre-physical domains vs. Diagnostic realized reference configurationes. A *PQS domain* is a relational region $\mathfrak{B} \subset \mathcal{X}_q$ (union of loci $\{\mathfrak{B}_x\}$) carrying the modal fibres and Fourfold algebra (Section 1.2.7). A *realized thermodynamic diagnostic realized reference configuration* augments $\mathfrak{B}$ with a certified plateau $\mathfrak{P}$ and BRST projection to the physical space $\mathcal{H}_{\text{phys}} := H^0(\hat{Q}_{\text{BRST}})$ [442, 445, 106, 483]; when a typed reading is required, it also includes an admissible projector $\Pi^{[\chi]}$ (Definitions 9.58 and 9.122, Section 12.3, and Proposition 9.138). All equilibrium statements are evaluated with respect to the *chronogeometric* time-flow determined by $\bar{\Sigma}_{\mu\nu}$ (Definition 5.26) and are parameterized by a *Good-clock* $C = (\hat{T}, U_t)$ aligned to that flow (Definition 13.158); hence ∂_t in (15.10) is the Good-clock Heisenberg derivative.

Definition 15.1 (Realized thermodynamic DRRC)**.** A *diagnostic realized reference configuration* (thermodynamic DRRC) over a PQS domain $\mathfrak{B} \subset \mathcal{X}_q$ is the tuple

$$\mathfrak{B}^{\langle \text{DRRC} \rangle}_{[\chi]}(\mathfrak{P}) := \Big(\mathfrak{B},\ \mathfrak{P},\ \mathcal{H}_{\text{phys}}{=}H^0(\hat{Q}_{\text{BRST}}),\ \bar{\Sigma}_{\mu\nu},\ \Pi^{[\chi]},\ \mathcal{V},\ w,\ U_c^{\text{prep}}, U_c^{\text{read}},\ \mathbb{E}_A,\ C = (\hat{T}, U_t)\Big),$$

where: $\mathfrak{P}$ is a certified plateau; $\bar{\Sigma}_{\mu\nu}$ is the realized chronogeometric tensor; $\Pi^{[\chi]}$ is an admissible typed projector (optional—untyped if $\Pi^{[\chi]} = \mathbb{I}$ on the block); $(\mathcal{V}, w)$ are the analysis window and weights; $U_c^{\text{prep}}, U_c^{\text{read}}$ are the holosymmetric control unitaries; $\mathbb{E}_A$ is the pointer/channel map used to form reduced states and entropic functionals (standard open–system setting [617, 441] and decoherence workflow [170]); and $C = (\hat{T}, U_t)$ is a Good-clock satisfying Definition 13.158. All outputs on a thermodynamic DRRC must pass the

standard audits (BRST/Nielsen safety, cone/causality, typed hyperbolicity, band stability, functoriality). This is the thermodynamic specialization of the diagnostic realized reference configuration of Definition 9.185. *Cf.* Definitions 2.123, 5.26, 9.58, 9.122, 9.153 and 15.27 and Section 12.3.

Definition 15.2 (Good-clock)**.** A *Good-clock* on a diagnostic realized reference configuration $\mathfrak{B}_{[\chi]}^{\langle \mathrm{DRRC}\rangle}(\mathfrak{P})$ is a pair $C = (\hat{T}, U_t)$ such that:

- **(C1) Chronogeometric alignment.** The generator of U_t coincides with the realized $\bar{\Sigma}$-time flow (Definition 5.26); i.e. ∂_t in (15.10) is the Heisenberg derivative w.r.t. U_t.
- **(C2) Low temporal holonomy.** On the diagnostic realized reference configuration, the temporal triad has trivial (or sub-tolerance) holonomy: $t_{\mathrm{corr}} = t_{\mathrm{vac}} = t_{\mathrm{spec}} = 0$ locally, or as certified by Proposition 4.556.
- **(C3) Conditional readability.** Page–Wootters conditioning holds with stable norm: conditioning on $\hat{T}$ yields $i\hbar\,\partial_t\,|\Psi_{\mathrm{sub}}(t)\rangle = \hat{H}_{\mathrm{sub}}^{\mathrm{eff}}(t)\,|\Psi_{\mathrm{sub}}(t)\rangle$ (Proposition 13.172 and eq. (13.238)).
- **(C4) BRST/Nielsen safety.** Clock operations are physical (projected to $\mathcal{H}_{\mathrm{phys}}$ on $\mathfrak{P}$) and gauge-parameter stable (Section 12.3 and Theorem 9.351).

Definition 15.3 (Clock-gauge equivalence)**.** Two Good-clocks $C = (\hat{T}, U_t)$ and $C' = (\hat{T}', U_{t'})$ on the same diagnostic realized reference configuration are *equivalent*, written $C \sim C'$, if $t' = \phi(t)$ with $\phi' > 0$ and there exists a bounded unitary family $W(t)$ on $\mathcal{H}_{\mathrm{phys}}$ such that $U_{t'} = W(t)U_t$ and $W(t)$ acts trivially on all reported diagnostic realized reference configuration observables. We write $[C]$ for the equivalence class.

Remark 15.4 ("Same DRRC")**.** Two reported quantities are *on the same diagnostic realized reference configuration* (DRRC) precisely when their DRRC tuples and Good-clocks match in the following sense:

$$(\mathfrak{B}_{[\chi]}^{\langle \mathrm{DRRC}\rangle}(\mathfrak{P}),\, C) \;\equiv\; (\mathfrak{B}_{[\chi]}^{\langle \mathrm{DRRC}\rangle}(\mathfrak{P}),\, C') \quad\Longleftrightarrow$$
$$\Big[\; \mathfrak{B}_{[\chi]}^{\langle \mathrm{DRRC}\rangle}(\mathfrak{P}) \text{ coincides componentwise} \;\Big] \;\text{ and }\; C \sim C',$$

with $C \sim C'$ the Good-clock equivalence relation (Definition 15.3). Componentwise coincidence includes, in particular, the same certified plateau $\mathfrak{P}$, BRST-physical projector $\mathsf{P}_{\mathrm{phys}}$, band tags $(\lambda_{\mathrm{UV}}, \ell)$ (and UV window $P_x(\lambda_{\mathrm{UV}})$), typed projector $\Pi^{[\chi]}$, subsystem and pointer channel (when present), and the same parasymmetric instrument index a (when meters are used) (Definitions 7.26, 9.10, 9.58, 9.122, 9.185 and 15.27).

Under this equivalence, numerical comparisons are meaningful *only* DRRC-wise. Concretely, for any typed witness $\Lambda_{\bullet}^{\Sigma;[\chi]}$, parasymmetric meter $v_{\mathrm{Synthesis}}^{(a);[\chi]}$, or quadrant assignment $\mathrm{Quad}_{[\chi]}^{[\Theta\Theta]}$,

$$(\mathfrak{B}_1, C_1) \equiv (\mathfrak{B}_2, C_2) \implies \Lambda_{\bullet}^{\Sigma;[\chi]}(\mathfrak{B}_1, C_1) = \Lambda_{\bullet}^{\Sigma;[\chi]}(\mathfrak{B}_2, C_2),$$
$$v_{\mathrm{Synthesis}}^{(a);[\chi]}(\mathfrak{B}_1, C_1) = v_{\mathrm{Synthesis}}^{(a);[\chi]}(\mathfrak{B}_2, C_2),$$

and

$$\mathrm{Quad}^{[\Theta\Theta]}_{[\chi]}(\mathfrak{B}_1, C_1) = \mathrm{Quad}^{[\Theta\Theta]}_{[\chi]}(\mathfrak{B}_2, C_2),$$

because (i) Born valuations are invariant under unitary gauge on $\mathcal{H}_{\mathrm{em}}$, $\mathrm{Tr}(E\rho) = \mathrm{Tr}\Big((gEg^\dagger)(g\rho g^\dagger)\Big)$, (ii) typed projectors $\Pi^{[\chi]}$ are held fixed (and commute with the coherence resolution $\mathcal{P}^{\mathrm{sym}}_{\mathrm{vac}} + \mathcal{P}^{\mathrm{coh}}_{\mathrm{dark}} + \mathcal{P}^{\mathrm{coh}}_{\mathrm{vis}} = \mathbf{1}_{\mathrm{phys}}$; eq. (13.149)), and (iii) Good-clock representatives in the same class $[C]$ differ by gauge/time reparametrizations that do not change readouts (Definitions 1.18 and 15.3). If these conditions fail, cross-DRRC comparisons are not defined within the thermodynamic statements and must first be transported to a common DRRC before evaluation.

Definition 15.5 (Thermal Equilibrium Between Relational Subsystems — Two Levels)**.** Let X, Y be relational subsystems represented by subalgebras $\mathcal{A}_X, \mathcal{A}_Y \subset \mathcal{A}(\mathcal{C}_d)$.

(Level P — pre-physical). We say X and Y are in *pre-physical thermal equilibrium*, written $\mathrm{Eq}_{\mathrm{pre}}(X, Y)$, if:

(a) **Algebraic–Statistical Stationarity:** the joint restricted state ω_{XY} is stationary with respect to the Good-clock flow aligned to $\bar{\Sigma}$,

$$\partial_t \omega_{XY} = 0, \tag{15.10}$$

where ∂_t denotes the Heisenberg derivative generated by U_t of $C = (\hat{T}, U_t)$.

(b) **Dynamical Current Balance:** the net $\bar{\Sigma}$-conserved relational current between X and Y vanishes,

$$\omega^{\mathring{\mathbf{\Psi}}}\Big(\hat{J}^\mu_{\mathrm{total}}(X) - \hat{J}^\mu_{\mathrm{total}}(Y)\Big)_{\mathrm{net}} = 0, \tag{15.11}$$

with currents defined by the Noether data on $\mathcal{A}(\mathcal{C}_d)$ and evaluated as Born expectations in the GNS representation [230, 30].

(Level R — realized, typed). On a Diagnostic realized reference configuration $(\mathfrak{B}, \mathfrak{P})$ and for any admissible type $[\chi]$, we say X and Y are in *realized (typed) thermal equilibrium*, written $\mathrm{Eq}^{[\chi]}_{\mathrm{phys}}(X, Y \,|\, \mathfrak{P})$, if *(a)–(b)* hold with all objects projected to $\mathcal{H}_{\mathrm{phys}}$ on $\mathfrak{P}$ and, when applicable, through $\Pi^{[\chi]}$. This includes the $\bar{\Sigma}$-time flow, Σ-causal support, and BRST/Nielsen safety [442, 120, 483].

Lemma 15.6 (Clock-gauge invariance of equilibrium)**.** *If $C \sim C'$ on the same diagnostic realized reference configuration, then the predicates $\mathrm{Eq}_{\mathrm{pre}}(X, Y)$ and $\mathrm{Eq}^{[\chi]}_{\mathrm{phys}}(X, Y \mid \mathfrak{P})$ are unchanged when ∂_t and the currents in (15.10)–(15.11) are computed with C' instead of C.*

Proof. Work on a fixed diagnostic realized reference configuration $\mathfrak{B}^{\langle \mathrm{DRRC}\rangle}_{[\chi]}(\mathfrak{P})$ (Definition 15.1, Remark 15.4). Let $C = (\hat{T}, U_t)$ and $C' = (\hat{T}', U_{t'})$ be equivalent Good-clocks, $C \sim C'$ (Definition 15.3). By definition, there exists a strictly monotone reparametrization $t' = \phi(t)$ with $\phi'(t) > 0$ and a unitary family $W(t)$ on $\mathcal{H}_{\mathrm{phys}}$ such that

$$U_{t'} \;=\; W(t)\, U_t, \qquad \text{and} \quad \mathrm{Ad}_{W(t)}(B) = B \;\; \text{for all DRRC observables} B,$$

equivalently, $\omega^{\mathring{\mathbf{\Psi}}} \circ \mathrm{Ad}_{W(t)} = \omega^{\mathring{\mathbf{\Psi}}}$ on the diagnostic realized reference configuration algebra; both clocks are aligned with the same $\bar{\Sigma}$-time flow (Definition 13.158).

(a) Algebraic–statistical stationarity. For any $A \in \mathcal{A}_X \vee \mathcal{A}_Y$,

$$A(t) := U_t^\dagger A\, U_t, \qquad A'(t') := U_{t'}^\dagger A\, U_{t'} = U_t^\dagger W(t)^\dagger A\, W(t) U_t.$$

Hence

$$\omega_{XY}^{(C')}(A, t') = \omega^{\mathring{\Psi}}\big(A'(t')\big) = \omega^{\mathring{\Psi}}\big(U_t^\dagger A\, U_t\big) = \omega_{XY}^{(C)}(A, t),$$

and, by the chain rule with $dt'/dt = \phi'(t) > 0$,

$$\partial_{t'}\, \omega_{XY}^{(C')}(A, t') = \frac{1}{\phi'(t)}\, \partial_t\, \omega_{XY}^{(C)}(A, t).$$

Therefore $\partial_t \omega_{XY}^{(C)}(A, t) = 0$ iff $\partial_{t'} \omega_{XY}^{(C')}(A, t') = 0$, establishing invariance of (15.10).

(b) Dynamical current balance. For the net current operator $\hat{J}^\mu_{\text{total}}(X) - \hat{J}^\mu_{\text{total}}(Y)$,

$$\Big(\hat{J}^\mu_{\text{total}}(X) - \hat{J}^\mu_{\text{total}}(Y)\Big)_{C'}(t') = U_{t'}^\dagger \Big(\cdots\Big) U_{t'} = U_t^\dagger W(t)^\dagger \Big(\cdots\Big) W(t) U_t,$$

so that its expectation (and the "net" combination) is unchanged by $\mathrm{Ad}_{W(t)}$:

$$\omega^{\mathring{\Psi}}\Big(\Big(\hat{J}^\mu_{\text{total}}(X) - \hat{J}^\mu_{\text{total}}(Y)\Big)_{C'}(t')\Big)_{\text{net}} = \omega^{\mathring{\Psi}}\Big(\Big(\hat{J}^\mu_{\text{total}}(X) - \hat{J}^\mu_{\text{total}}(Y)\Big)_{C}(t)\Big)_{\text{net}}.$$

Hence (15.11) vanishes for C iff it vanishes for C'. Alignment with $\bar{\Sigma}$ (Definition 13.158) ensures both clocks use the same chronogeometric time (Definition 5.26), and low temporal holonomy prevents spurious phase artefacts (Proposition 4.556).

(c) Realized, typed predicates. After projection to $\mathcal{H}_{\text{phys}}$ and, where used, to a typed subspace via $\Pi^{[\chi]}$ (Definitions 9.58 and 9.122 and Section 12.3), the same identities hold since $\mathrm{Ad}_{W(t)}$ fixes diagnostic realized reference configuration observables and numbers, and Nielsen/BRST safety guarantees gauge-parameter independence of reported values (Theorem 9.351). Thus $\mathrm{Eq}^{[\chi]}_{\text{phys}}(X, Y \mid \mathfrak{P})$ is invariant under $C \mapsto C'$.

Combining (a)–(c) proves the claim. □

Theorem 15.7 (Zeroth Law of Thermodynamics — Two-Level Form)**.** *Let $\mathfrak{B}_A, \mathfrak{B}_B, \mathfrak{B}_C \subset \mathcal{X}_q$ be three distinct, mutually interacting PQS domains whose dynamics are governed by the emergent chronogeometry $(\mathcal{M}_{\text{geom}}, \bar{\Sigma})$ (Definitions 5.26 and 5.113). Fix a* same diagnostic realized reference configuration *in the sense of Remark 15.4 and a Good-clock gauge class $[C]$ (Definition 15.3). The equilibrium predicates are those of Definition 15.5, namely $\partial_t \omega_{XY} = 0$ ((15.10)) and $\omega^{\mathring{\Psi}}\big(\hat{J}^\mu_{\text{total}}(X) - \hat{J}^\mu_{\text{total}}(Y)\big)_{\text{net}} = 0$ ((15.11)), with time parameterized by any representative $C \in [C]$ (well-defined by Lemma 15.6). Then:*

(a) Pre-physical (P)

If $\mathrm{Eq}_{\text{pre}}(\mathfrak{B}_A, \mathfrak{B}_C)$ and $\mathrm{Eq}_{\text{pre}}(\mathfrak{B}_B, \mathfrak{B}_C)$, computed with the same equivalence class $[C]$, then $\mathrm{Eq}_{\text{pre}}(\mathfrak{B}_A, \mathfrak{B}_B)$. Comparisons are formed using the gauge-covariant transporter

$$\mathcal{U}(x \leftarrow x'; \gamma) = \mathcal{P} \exp\Big(-\int_\gamma \mathcal{A}\Big)$$

between loci (eqs. (4.11) and (4.16)); $\bar{\Sigma}$-microcausality and "no-law-signalling" ensure causal consistency (Corollary 5.145).

(b) Realized, typed (R)

Fix a certified plateau $\mathfrak{P}$. For any admissible type $[\chi]$ with projector $\Pi^{[\chi]}$ satisfying the permit conditions and BRST/Nielsen safety (Definitions 9.58 and 9.122 and Theorem 9.351), if $\mathrm{Eq}^{[\chi]}_{\mathrm{phys}}(\mathfrak{B}_A, \mathfrak{B}_C \,|\, \mathfrak{P})$ and $\mathrm{Eq}^{[\chi]}_{\mathrm{phys}}(\mathfrak{B}_B, \mathfrak{B}_C \,|\, \mathfrak{P})$ hold on the same diagnostic realized reference configuration and $[C]$, then $\mathrm{Eq}^{[\chi]}_{\mathrm{phys}}(\mathfrak{B}_A, \mathfrak{B}_B \,|\, \mathfrak{P})$. Typed evaluations use Born valuations $v(\rho; E) = \mathrm{Tr}(E\rho)$ on $\mathcal{H}_{\mathrm{phys}}$ with $\rho \mapsto \mathsf{P}_{\mathrm{phys}} \rho \mathsf{P}_{\mathrm{phys}}$ and effects $\Pi^{[\chi]} E \Pi^{[\chi]}$ (Definition 1.18 and Section 12.3).

Moreover, if the typed projector is block-identity on the sector under consideration, the realized statement reduces to the untyped one (Proposition 9.138). In the duality-collapse regime (Definition 4.87), the predicate restricts to the Twofold sector and coincides with the classical bivalent formulation (cf. the classical transitivity axiom [119]).

Typed–untyped reduction and classical Twofold limit.

Proposition 15.8 (Typed→Untyped reduction; Duality-collapse classical limit)**.** Same diagnostic realized reference configuration *and* Good-clock *are fixed (Remark 15.4, Definition 15.3, and Lemma 15.6).*

(i) Typed→Untyped reduction

Let $[\chi]$ be admissible on the certified plateau $\mathfrak{P}$ with projector $\Pi^{[\chi]}$. Assume that $\Pi^{[\chi]}$ is block-identity on the sector used by the equilibrium predicate, namely

$$\Pi^{[\chi]}\Big|_{\mathcal{H}_S} = \mathbf{1}_{\mathcal{H}_S},$$

for the support $\mathcal{H}_S$ of all effects and states entering (15.10)–(15.11). Then, for every effect E and state ρ_{phys} on $\mathcal{H}_S$,

$$\mathrm{Tr}\Big(\Pi^{[\chi]} E \Pi^{[\chi]}\ \Pi^{[\chi]} \rho_{\mathrm{phys}} \Pi^{[\chi]}\Big) = \mathrm{Tr}\Big(E\, \rho_{\mathrm{phys}}\Big) \qquad (Definition\ 1.18).$$

Consequently the realized, typed predicate coincides with the untyped one:

$$\mathrm{Eq}^{[\chi]}_{\mathrm{phys}}(X, Y \mid \mathfrak{P}) \iff \mathrm{Eq}_{\mathrm{phys}}(X, Y \mid \mathfrak{P}),$$

and, by Proposition 9.138, this also equals the pre-physical statement when the projection does not alter the sector used by the observables/states.

(ii) Duality-collapse classical limit

In the duality-collapse regime (Definition 4.87) at a locus x, the Twofold block induces a nondegenerate inner product, and there exists an isometry

$$U : \mathbb{C}^2 \hookrightarrow \mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}\ (0,1)}$$

such that Twofold-respecting statistics match those of a qubit state

$$\rho_{(0,1)} = U^\dagger \rho\, U \qquad (Proposition\ 4.90).$$

With the Axiom of Duality, outcomes are Twofold projectors $P_{\mathbf{0}}, P_{\mathbf{1}}$ (Section 14.1.3), and the Born valuations

$$p_{\mathbf{i}} = \mathrm{Tr}(\rho P_{\mathbf{i}})$$

reduce to classical bivalent probabilities when decoherence pinches the pointer algebra. Hence the equilibrium predicate on the Twofold sector coincides with the classical bivalent formulation, matching the classical transitivity axiom (cf. [119]).

Proof sketch. (i) Follows by inserting $\Pi^{[\chi]}$ on both state and effect and using $\Pi^{[\chi]}|_{\mathcal{H}_S} = \mathbf{1}_{\mathcal{H}_S}$ together with Proposition 9.138. Stationarity and current-balance predicates (15.10)–(15.11) are preserved termwise. (ii) Uses Definition 4.87 and Proposition 4.90 and Born readout on the Twofold slice; the pinching/sector unitality ensures classical bivalence on the pointer algebra, yielding classical transitivity.

Proof. (i) Typed→Untyped reduction. Work on a *same diagnostic realized reference configuration* with Good-clock class $[C]$ (Remark 15.4, Definition 15.3, and Lemma 15.6). Fix an admissible type $[\chi]$ on the certified plateau $\mathfrak{P}$ with projector $\Pi^{[\chi]}$ (Definitions 9.58 and 9.122), and assume that $\Pi^{[\chi]}$ is block-identity on the sector actually used by the equilibrium predicate, i.e.

$$\Pi^{[\chi]}\Big|_{\mathcal{H}_S} = \mathbf{1}_{\mathcal{H}_S},$$

where $\mathcal{H}_S \subset \mathcal{H}_{\mathrm{phys}}$ is the common support of the states/effects entering (15.10)–(15.11). For any effect E and state ρ_{phys} on $\mathcal{H}_S$,

$$\mathrm{Tr}\Big(\Pi^{[\chi]} E \Pi^{[\chi]}\ \Pi^{[\chi]} \rho_{\mathrm{phys}} \Pi^{[\chi]}\Big) = \mathrm{Tr}(E\rho_{\mathrm{phys}}),$$

so Born valuations agree on typed vs. untyped data (Definition 1.18). Consequently, the algebraic–statistical stationarity predicate $\partial_t \omega_{XY} = 0$ in (15.10) holds in the typed evaluation iff it holds untyped, because expectations coincide pointwise in t and $[C]$ fixes the time gauge (Lemma 15.6).

For the dynamical current balance (15.11), write Heisenberg evolutions with any $C \in [C]$:

$$\Big(\hat{J}^{\mu}_{\mathrm{total}}(X) - \hat{J}^{\mu}_{\mathrm{total}}(Y)\Big)(t) = U_t^{\dagger}\Big(\hat{J}^{\mu}_{\mathrm{total}}(X) - \hat{J}^{\mu}_{\mathrm{total}}(Y)\Big)U_t.$$

Inserting $\Pi^{[\chi]}$ on both sides leaves the numerical expectation invariant (same reason as above), hence the vanishing of the net current is the same typed or untyped. Therefore

$$\mathrm{Eq}^{[\chi]}_{\mathrm{phys}}(X, Y \mid \mathfrak{P}) \iff \mathrm{Eq}_{\mathrm{phys}}(X, Y \mid \mathfrak{P}),$$

and by Proposition 9.138 this matches the untyped statement whenever the projector is block-identity on the relevant sector.

(ii) Duality-collapse classical limit. In the duality-collapse regime (Definition 4.87), the Twofold block on $\mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}\,(0,1)}$ is nondegenerate and there exists an isometry $U : \mathbb{C}^2 \hookrightarrow \mathcal{H}^{(\mathfrak{B}_x)}_{\mathrm{mod}\,(0,1)}$ such that Twofold-respecting Born statistics are reproduced by the qubit state $\rho_{(0,1)} = U^{\dagger}\rho U$ (Proposition 4.90). With the Axiom of Duality, readouts are Twofold projectors $P_{\mathbf{0}}, P_{\mathbf{1}}$ on the readout slice (Section 14.1.3), hence

$$p_{\mathbf{i}}(t) = \mathrm{Tr}\left(\rho(t)\, P_{\mathbf{i}}\right) \quad (\mathbf{i} = \mathbf{0}, \mathbf{1}).$$

Open-system monitoring on the active band is unital CP and CP-divisible: each stroboscopic step $\Phi_{\Delta t} = \mathbb{E}_A \circ \mathrm{Ad}_{e^{-iH_{A,\mathrm{eff}}\Delta t}}$ is CPTP/unital and generates a unital GKLS semigroup in the continuous limit (Definition 15.27 and Propositions 15.29, 15.45 and 15.57).

Pinching/Karamata yields entropy increase under the pointer map and, operationally, the suppression of off-diagonals in the pointer algebra (Proposition 15.31 and Theorem 15.35). Thus the Twofold pointer algebra becomes effectively classical (commutative), whence

$$p_{\mathbf{0}}(t) + p_{\mathbf{1}}(t) = 1, \qquad p_{\mathbf{i}}(t) \in [0,1],$$

and the equilibrium predicates (15.10)–(15.11) reduce to their classical bivalent forms (stationary probabilities and vanishing net exchange), recovering the classical transitivity axiom (cf. [119]). This establishes that, in the duality-collapse regime, the Twofold equilibrium predicate coincides with the classical bivalent formulation. □

Remark 15.9 (Thermal contact and Σ-causality)**.** The equilibrium predicates (15.10)–(15.11) are evaluated only when X and Y are in *thermal contact* on the *same diagnostic realized reference configuration* (Remark 15.4). This entails:

(i) Σ-causal connectivity

The domains are chronogeometrically connected with respect to $\bar{\Sigma}$:

$$J^+_{\cup}(X) \cap Y \neq \varnothing, \qquad J^-_{\cup}(Y) \cap X \neq \varnothing,$$

where $J^{\pm}_{\cup}$ is the branchwise union of future/past sets defined by $\bar{\Sigma}$ (Definition 5.26). Thus X and Y are not spacelike separated ($X \not\perp_{\cap} Y$), and all comparisons are made via the gauge-covariant transporter

$$\mathcal{U}(x \leftarrow x'; \gamma) = \mathcal{P} \exp\Big(-\int_{\gamma} \mathcal{A}\Big)$$

between loci (eqs. (4.11) and (4.16)). By $\bar{\Sigma}$-microcausality and "no-law-signalling", law-superposition cannot enlarge causal support, so spurious influence outside $J^{\pm}_{\cup}$ is excluded (Corollary 5.145).

(ii) Nontrivial exchange channel on the diagnostic realized reference configuration

In the Good-clock subsystem split

$$\hat{H} = \hat{H}_X + \hat{H}_Y + \hat{H}_{XY} \qquad (\textit{Definition 15.11 and Section 15.1.3}),$$

the interaction satisfies $\hat{H}_{XY} \neq 0$ (and, in typed evaluations, $\Pi^{[\chi]} \hat{H}_{XY} \Pi^{[\chi]} \neq 0$). Then the Good-clock Heisenberg balance reads

$$\frac{dU_X}{dt} = \big\langle i[\hat{H}_{XY}, \hat{H}_X] \big\rangle - \big\langle \partial_t \hat{H}_X \big\rangle \qquad (\textit{Lemma 15.12}),$$

so that the current term is physically meaningful and not trivially zero by algebraic independence.

With (i)–(ii) in force, the stationarity predicate $\partial_t \omega_{XY} = 0$ and the dynamical current balance in (15.11) distinguish genuine equilibrium from (a) mere spacelike decoupling (ruled out by (i)), and (b) interaction-induced cancellations that would be artefacts of vanishing coupling (ruled out by (ii)). Holosymmetric interference cannot "fake'' equilibrium because it preserves cone structure and retarded support (Corollary 5.145; cf. algebraic microcausality and spectral support in QFT [230, 383]).

On Diagnostic realized reference configurationes, pointer reductions and coarse-grained dynamics act on the active band via the unital channel $\mathbb{E}_A$ and CP-divisible/GKLS semigroups, ensuring that readouts used in (15.10)–(15.11) are diagnostic realized reference configuration-consistent and gauge-parameter flat (Definition 15.27, Proposition 15.29, Theorems 9.351 and 15.35, and Remark 15.4).

Proof of the Zeroth Law

The Zeroth Law of Thermodynamics. Fix a realized thermodynamic diagnostic realized reference configuration $\mathfrak{B}^{\langle \mathrm{DRRC} \rangle}_{[\chi]}(\mathfrak{P})$ (Definition 15.1) with Good-clock $C = (\hat{T}, U_t)$ (Definition 13.158); by Lemma 15.6 any representative of the clock gauge $[C]$ may be used. Let the global algebra be $\mathcal{A}_{ABCE} \subset \mathcal{A}(\mathcal{C}_d)$ with state $\omega^{\mathring{\Psi}}$ (GNS; Section 1.4.1).

Step 1 — Premises. Assume the hypotheses of Theorem 15.7:

(P1) $\mathrm{Eq}_{\mathrm{pre}}(\mathfrak{B}_A, \mathfrak{B}_C)$: by Definition 15.5, $\partial_t \omega_{AC} = 0$ as in (15.10) and

$$\omega^{\mathring{\Psi}}\Big(\hat{J}^{\mu}_{\mathrm{total}}(A) - \hat{J}^{\mu}_{\mathrm{total}}(C)\Big)_{\mathrm{net}} = 0$$

as in (15.11).

(P2) $\mathrm{Eq}_{\mathrm{pre}}(\mathfrak{B}_B, \mathfrak{B}_C)$: the analogous two conditions hold for B, C.

(For the realized, typed clause, project to $\mathcal{H}_{\mathrm{phys}}$ on $\mathfrak{P}$ and through $\Pi^{[\chi]}$; see Step 4.)

Step 2 — Transitivity of current balance. Let $\nabla^{(\mathrm{rel})}_{\mu} \hat{J}^{\mu}_{\mathrm{total}} = 0$ be the continuity equation for the conserved relational current associated with the Good-clock symmetry (Noether structure on $\mathcal{A}(\mathcal{C}_d)$). Additivity and linearity of $\hat{J}^{\mu}_{\mathrm{total}}$ over disjoint unions give, for any triplet A, B, C,

$$\Big(\hat{J}^{\mu}_{\mathrm{total}}(A) - \hat{J}^{\mu}_{\mathrm{total}}(B)\Big) = \Big(\hat{J}^{\mu}_{\mathrm{total}}(A) - \hat{J}^{\mu}_{\mathrm{total}}(C)\Big) - \Big(\hat{J}^{\mu}_{\mathrm{total}}(B) - \hat{J}^{\mu}_{\mathrm{total}}(C)\Big).$$

Taking Good-clock Heisenberg expectations in $\omega^{\mathring{\Psi}}$ and using (P1)–(P2),

$$\omega^{\mathring{\Psi}}\Big(\hat{J}^{\mu}_{\mathrm{total}}(A) - \hat{J}^{\mu}_{\mathrm{total}}(B)\Big)_{\mathrm{net}} = 0 - 0 = 0,$$

so the current-balance clause of $\mathrm{Eq}_{\mathrm{pre}}(\mathfrak{B}_A, \mathfrak{B}_B)$ holds. The same identity survives projection in the realized/typed setting because expectations are taken on the diagnostic realized reference configuration and are BRST/Nielsen-safe (Section 12.3 and Theorem 9.351).

Step 3 — Stationarity of ω_{AB}. Let $\alpha_t(\,\cdot\,) := U_t^{\dagger}(\,\cdot\,)U_t$ be the Good-clock Heisenberg flow aligned with $\bar{\Sigma}$ (Definition 13.158). We prove $\partial_t \omega_{AB} = 0$ on $\mathcal{A}_{AB} := \mathcal{A}_A \vee \mathcal{A}_B$.

(S3.1) Coarse-grained dynamics and intertwining. Operational reductions on the diagnostic realized reference configuration are CP-divisible and unital on the active band and implemented by pointer channels $\mathbb{E}_X$ (Definition 15.27 and Theorem 15.50). We use the conditional-expectation property on the diagnostic realized reference configuration timescale:

$$\mathbb{E}_{AB}\circ\alpha_t = \alpha_t\circ\mathbb{E}_{AB}, \qquad \omega_{AB} = \omega_{ABCE}\circ\mathbb{E}_{AB},$$

(cf. [617, 441], and Definition 15.1). Thus

$$\partial_t\omega_{AB}(O) = \partial_t\,\omega_{ABCE}\Big(\mathbb{E}_{AB}(O)\Big) = \partial_t\,\omega_{ABCE}\Big(\alpha_t(\mathbb{E}_{AB}(O))\Big)\Big|_{t=0} \quad (\forall O\in\mathcal{A}_{AB}).$$

(S3.2) Generator decomposition and vanishing of C-couplings. The Heisenberg generator on $\mathcal{A}_{ABCE}$ has GKLS form

$$\mathcal{L}^\dagger(\cdot) = i[\hat H,\cdot] + \sum_\alpha \gamma_\alpha\Big(L_\alpha^\dagger(\cdot)\,L_\alpha - \tfrac12\{L_\alpha^\dagger L_\alpha,(\cdot)\}\Big),$$

with $\hat H = \hat H_A+\hat H_B+\hat H_C+\hat H_{AB}+\hat H_{AC}+\hat H_{BC}+\hat H_{\mathrm{res}}$ (Section 15.1.3; unital CP-divisible setting from Theorem 15.50). Write $\mathcal{L}^\dagger = \mathcal{L}^\dagger_{AB}\oplus\mathcal{L}^\dagger_C\oplus\mathcal{L}^\dagger_{\mathrm{coupl}}$ where $\mathcal{L}^\dagger_{\mathrm{coupl}}$ contains the AC, BC Hamiltonian and dissipative couplings. Then

$$\partial_t\omega_{AB}(O) = \omega_{ABCE}\Big(\mathcal{L}^\dagger(\mathbb{E}_{AB}(O))\Big) = \omega_{ABCE}\Big(\mathcal{L}^\dagger_{AB}(\mathbb{E}_{AB}(O))\Big) + \omega_{ABCE}\Big(\mathcal{L}^\dagger_{\mathrm{coupl}}(\mathbb{E}_{AB}(O))\Big).$$

For the coupling part, $\mathbb{E}_{AB}(O)$ is supported on AB. Every term in $\mathcal{L}^\dagger_{\mathrm{coupl}}$ either: (a) is a commutator with $\hat H_{AC}$ or $\hat H_{BC}$, or (b) involves a dissipator acting nontrivially on C. The expectations of (a)-terms coincide with the net current exchanges measured in (15.11); by Step 2 and (P1)–(P2) their diagnostic realized reference configuration values vanish. For (b), unitality and the fixed-point property of ω_{AC},ω_{BC} under the reduced semigroup imply that dissipative C-terms contribute zero expectation (they annihilate fixed points; Theorem 15.50). Hence

$$\omega_{ABCE}\Big(\mathcal{L}^\dagger_{\mathrm{coupl}}(\mathbb{E}_{AB}(O))\Big) = 0.$$

(S3.3) Fixed-point face and conclusion. We are left with the internal generator on AB:

$$\partial_t\omega_{AB}(O) = \omega_{ABCE}\Big(\mathcal{L}^\dagger_{AB}(\mathbb{E}_{AB}(O))\Big).$$

Since $\mathcal{L}^\dagger_{AB}$ is the Heisenberg generator of a unital CP-divisible semigroup on $\mathcal{A}_{AB}$, its set of fixed points is a convex face closed under multiplication by invariant elements. The marginals ω_A,ω_B are fixed (from (P1)–(P2) via contractivity/one-step monotonicity; Theorem 15.50), and $\mathbb{E}_{AB}(O)$ belongs to the face generated by those marginals. Therefore ω_{AB} lies in the fixed-point face of the AB-semigroup and $\partial_t\omega_{AB}(O)=0$ for all $O\in\mathcal{A}_{AB}$. Thus $\partial_t\omega_{AB}=0$.

Together with Step 2, we conclude $\mathrm{Eq}_{\mathrm{pre}}(\mathfrak{B}_A,\mathfrak{B}_B)$.

Step 4 — Realized, typed clause. Project all algebras and states to $\mathcal{H}_{\mathrm{phys}}{=}H^0(\hat Q_{\mathrm{BRST}})$ on $\mathfrak{P}$ and (optionally) through an admissible $\Pi^{[\chi]}$ (Definitions 9.58 and 9.122 and Section 12.3). By Lemma 15.6, the representative of $[C]$ is irrelevant. The arguments of Steps 2–3 apply verbatim to the projected quantities (Nielsen/BRST safety; Theorem 9.351), yielding $\mathrm{Eq}^{[\chi]}_{\mathrm{phys}}(\mathfrak{B}_A,\mathfrak{B}_B\mid\mathfrak{P})$.

This completes the proof of Theorem 15.7. □

Remark 15.10. This proof grounds the Zeroth Law in the core axioms of the framework: the *single* global state $\omega^{\mathring{\Psi}}$ on the C*-algebra $\mathcal{A}(\mathcal{C}_d)$ and its *diagnostic realized reference configuration-wise* realization (Definition 15.1). The equilibrium predicate uses the Good-clock $C = (\hat{T}, U_t)$ aligned to $\bar{\Sigma}$ (Definition 13.158) and is invariant under clock-gauge changes (Lemma 15.6); it is defined by stationarity and current balance on subalgebras (Definition 15.5), subject to Σ-causal support and BRST/Nielsen safety (Corollary 5.145 and Theorem 9.351). Thus the theorem *is not* merely a Hilbert-space "entanglement transitivity'' statement, but a consequence of the **self-consistency of a single state on the fundamental Algebra of Observables** together with its consistent restrictions (Theorem 15.7; cf. [230, 241, 381, 230]). At the realized, typed level, the conclusion is independent of measurement details because only CP-unital pointer reductions and Born valuations enter (Definitions 1.18 and 15.27; cf. [617, 441, 170]). In the duality-collapse regime the predicate reduces to the classical bivalent transitivity (Definition 4.87; cf. [119]). This elevates the Zeroth Law from a phenomenological rule to a necessity of $\bar{\Sigma}$-chronogeometric and algebraic consistency in the quantum-relational ontology.

The Deeper Meaning: Thermodynamics as a Law of Information What the two–level Zeroth Law (Theorem 15.7) reveals is that the thermodynamic predicates are, at root, constraints on the *flow and consistency of quantum–relational information* fixed by a diagnostic realized reference configuration and its Good-clock (Definitions 13.158 and 15.1 and Lemma 15.6). Operational numbers are Born valuations on subalgebras (Definition 1.18), so equilibrium reduces to an information–theoretic stasis condition compatible with Σ-causal support (Definitions 5.26 and 15.5). In this sense the laws are emergent principles of constrained inference on a global state, not merely statements about heat or disorder [618, 619, 620, 621, 622, 137].
The treatise's ontology is relational—akin to *Indra's Net*—in which each PQS locus $\mathfrak{B}_x$ "exists" through its lawful reflections with the rest via the Fourfold correlator and GNS state $\omega^{\mathring{\Psi}}$ (Definition 4.9 and Section 1.4.1). The *Quantum Holo-State* is the coherence law that governs these mutual reflections [3].

Thermal Equilibrium as Informational Stasis In this light, $\mathrm{Eq}_{\mathrm{pre}}(X, Y)$ and $\mathrm{Eq}^{[\chi]}_{\mathrm{phys}}(X, Y \mid \mathfrak{P})$ (Definition 15.5) say that, with respect to the Good-clock flow aligned to $\bar{\Sigma}$ (Definition 13.158), (a) the joint state is stationary and (b) the net conserved *relational* currents vanish—here carried by the Gaiamiti and Tantrakyma sectors (Noether data; cf. Sections 9.17.8 and 9.18.10). Equilibrium is therefore a pattern of *informational stasis* consistent with Σ-causality and BRST/Nielsen safety (Corollary 5.145 and Theorem 9.351): the infinite "reflections'' among A, B, C have settled into a stable, self-consistent configuration.

The Zeroth Law as a Law of Consistent Observation The theorem formalizes a consistency principle for observations on the same diagnostic realized reference configuration (Remark 15.4): if a clocked observer C finds its information exchange balanced with A and with B, then global coherence demands the $A{\leftrightarrow}B$ exchange be balanced as well (Theorem 15.7). *Clock-gauge invariance* (Lemma 15.6) ensures this consistency does not depend

on a particular parameterization of time but only on the Good-clock class $[C]$.

Temperature as a Measure of Informational Friction Macroscopically, "temperature'' appears as the parameter quantifying a subsystem's decohering openness to its environment, i.e. *informational friction*. On the diagnostic realized reference configuration, this is mediated by CP-divisible/unital pointer reductions $\mathbb{E}_A$ (Definition 15.27) that implement environment-induced superselection; the degree of blurring of local records tracks effective temperature [170, 282]. In information-theoretic terms, temperature gauges how aggressively local degrees of freedom exchange and randomize information within a globally coherent state [618, 619]. Absolute zero would correspond to perfect informational isolation—precluded in a fully relational cosmos except in limiting, decoupled (Good-clock) regimes.

Synthesis. The formal proof thus completes the passage from 19th–century phenomenology to a diagnostic realized reference configuration-wise, clocked theory of *quantum information flow*: the Zeroth Law is the necessity of self-consistent, Σ-causal, and gauge-safe information across subalgebras, enforced by the single global state and its coherent restrictions (Theorem 15.7 and Definition 15.5; cf. [230, 30]).

Zeroth Law (high-level meaning and consequences). In this programme the Zeroth Law does not assert a primitive equality such as $T_A = T_B$. It states that, on a single holosymmetric diagnostic realized reference configuration with a fixed Good-clock gauge (Remark 15.4 and Definitions 13.158 and 15.3), two subsystems A and B are in mutual equilibrium exactly when: (i) their joint statistics are *stationary* under the Good-clock flow (the Good-clock derivative leaves all expectation values invariant; Definition 15.5 and eq. (15.10)), and (ii) the *relational* Noether current associated with the same flow exhibits *no net exchange* between them (eq. (15.11)). Because both predicates are linear/additive (numbers are taken with the emergent inner product) and evaluated in the same chronogeometric and gauge frame, the relation is transitive: if $A \leftrightarrow C$ and $B \leftrightarrow C$ hold on the same diagnostic realized reference configuration/clock, then $A \leftrightarrow B$ as well (Lemma 15.6 and Theorem 15.7; cf. [230, 30, 617, 441]). The classical wording "same temperature" is the numerical parameterization of this fixed-point family in the duality-collapse (Twofold) readout regime (Definition 4.87 and Section 14.1.3; cf. [119]).

Placed against the *Tetralemma*, equilibrium is not a forced, global bivalent verdict. The Fourfold modal structure lets one *type* the predicate by sector: geometric TT vs. Tr in Gaiamiti, matter blocks in Tantrakyma, and visible vs. dark coherence channels. With typed projectors $\Pi^{[\chi]}$ held fixed on the diagnostic realized reference configuration (Definition 9.122 and Remark 15.4), it is consistent to have "A and B equilibrated in $[\chi_1]$ but not in $[\chi_2]$." Meters and projectors are part of the operational diagnostic realized reference configuration, so sectorwise equilibrium is both meaningful and experimentally decidable (Definition 1.18; cf. [230]).

The *Axiom of Duality* explains how these quantum statements are read in the lab: after interaction and decoherence, readouts restrict to the Twofold subspace, and Born valuations on that slice deliver bivalent outcomes (Section 14.1.3 and Definition 1.18). In the *duality-collapse* regime the pointer algebra pinches to a commuting block, so the fixed-point family of the unital CP dynamics can be labeled by a single intensive ("temperature"). Thus

thermometers recover Callen's axiom as the typed quantum predicate reduces to equality of a number (Definition 4.87 and Theorem 15.50; cf. [170, 119, 441]).

The *Quantum Holo-State* and the *Fourfold inter-modal correlator* $\hat{G}_\Lambda$ make this precise. The inner product used for probabilities and currents is *emergent* from $\hat{G}_\Lambda$ in the coincident limit (Definition 4.9 and eq. (4.85)), so "stationarity" and "no net current" are $\hat{G}_\Lambda$-based invariances, not coordinate conventions. Typing simply inserts $\Pi^{[\chi]}$ to form the typed correlator $\Pi^{[\chi]}\hat{G}_\Lambda\Pi^{[\chi]}$. Equilibrium then says that these correlator-based expectations are fixed points of the diagnostic realized reference configuration's unital, CP-divisible flow and that the correlator-based Noether current between A and B averages to zero—channel by channel (Definition 15.27 and Theorem 15.50; cf. [617, 441]).

The *Primordial Quantum Singularity (PQS)* and *chronogeometry* provide the causal scaffold. The unified tensor $\bar{\Sigma}$ fixes cones and microcausality (Definition 5.26), so "thermal contact" means Σ-causal connectivity *and* a nontrivial exchange channel on the diagnostic realized reference configuration. Two guardrails follow: (i) *law-interference* (superposing admissible dynamics) cannot enlarge causal support beyond the $\bar{\Sigma}$ cones, and (ii) *no-law-signalling* forbids remote diagnostic realized reference configuration changes from altering the predicate when subsystems are spacelike (Corollary 5.145; cf. [230, 383]). Thus "no net current" is chronogeometrically meaningful.

Transitivity—the heart of the Zeroth Law—has a clear mechanism. Additivity of the conserved relational current yields the $A-B$ balance from the $A-C$ and $B-C$ balances, while stationarity is a fixed-point property of a unital CP semigroup on the active band; fixed points form a convex face, and when the $A-C$ and $B-C$ faces coincide on the *same diagnostic realized reference configuration/clock*, the joint $A-B$ state lies in that face as well (Theorems 15.7 and 15.50 and Remark 15.4; cf. [617, 441]). All of this holds pre-physically and survives BRST/typed projection on a certified plateau, so the realized, typed clause inherits the same transitivity (Section 12.3 and Definition 9.58; cf. [442]).

Practical consequences. (i) Equilibrium is *relational and diagnostic realized reference configuration-dependent*: cross-diagnostic realized reference configuration comparisons (different band tags, clocks, or typing) are meaningless until transported to a common diagnostic realized reference configuration (Remark 15.4). (ii) Equilibrium is *sectorwise*: geometry can equilibrate while a matter excitation channel does not; visible matter can equilibrate while a dark coherence channel remains decoupled (Definition 9.122). (iii) The Zeroth Law furnishes the *reference baseline* for the rest of thermodynamics: once "fixed point + no net current" is verified, the First Law's heat/work split and the Second Law's entropy production are measured relative to that baseline (Lemma 15.12, eq. (15.12), and Theorem 15.50; cf. [119, 441]).

In one line. The Zeroth Law asserts that *equilibrium is the transitive coincidence of fixed points and vanishing relational currents, evaluated within a single holosymmetric diagnostic realized reference configuration*—and, when the Twofold readout dominates, it reduces numerically to the classical statement that systems in mutual equilibrium share a common temperature (Definition 4.87 and Section 14.1.3; cf. [119]).

15.1.3 The First Law Reimagined as Relational Energy

Classically, the First Law states that the change of a system's internal energy equals the heat added minus the work done, $\Delta U = Q - W$ [119]. In the present framework this identity is realized *diagnostic realized reference configuration-wise* (Definition 15.1) and *clocked* by a Good-clock $C = (\hat{T}, U_t)$ aligned to the $\bar{\Sigma}$-time flow (Definition 13.158). All time derivatives below are Good-clock (Heisenberg) derivatives, invariant under clock-gauge changes by Lemma 15.6. The relational form of the law will be stated in *eq.* (15.12) and derived from Noether symmetry of the total action (Section 15.1.3).

(i) Internal energy U For a subsystem A (subalgebra $\mathcal{A}_A \subset \mathcal{A}(\mathcal{C}_d)$) on a diagnostic realized reference configuration, define

$$U_A(t) \;:=\; \mathrm{Tr}\Big(\rho_A(t)\, H_A(t)\Big),$$

with ρ_A the reduced (GNS) state and H_A the Relational Hamiltonian obtained from the subsystem split of $\hat{H}_{\mathrm{univ}}$ (Section 15.1.3; cf. [230]). The time parameter t is the Good-clock time (Definition 13.158).

(ii) Heat Q Heat is the *incoherent* contribution to ΔU_A generated by the open-system exchange across the $A|E$ cut. Operationally, this is the part of the Good-clock derivative $\dot{U}_A$ induced by CP-divisible, unital pointer/reduction maps $\mathbb{E}_A$ (Definition 15.27 and Theorem 15.50) that encode environment-induced decoherence [617, 441, 170]. Intuitively, quanta of the Gaiamiti/Tantrakyma sectors crossing the boundary contribute to Q (Sections 9.17.8 and 9.18.10; cf. [619, 282]).

(iii) Work W Work is the *coherent* contribution to ΔU_A generated by Good-clock Hamiltonian driving on the diagnostic realized reference configuration—for example, changes of background chronogeometry $\bar{\Sigma}$ and/or controlled unitary couplings that act within A (Definitions 13.158 and 15.1). In equations, it is the part of $\dot{U}_A$ stemming from explicit t-dependence of $H_A(t)$ (Good-clock driving) rather than dissipative exchange; see the Heisenberg identification in Section 15.1.3 and (15.12) (cf. [170]).

The diagnostic realized reference configuration-wise First Law then follows from the subsystem split of $\hat{H}_{\mathrm{univ}}$ and the Good-clock Heisenberg equation, yielding a decomposition of $\dot{U}_A$ into coherent (work) and incoherent (heat) parts, and the integrated identity

$$\boxed{\Delta U_A = Q - W} \quad \text{(eq. (15.12))}.$$

The Foundational Invariance: Symmetry of the Total Action

The dynamics are governed by the *Total Relational Action* (Definition 9.357)

$$S_{\mathrm{total}}[\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}] = S_{\Phi}[\hat{\mathbf{\Phi}}] + S_{\Theta}[\hat{\mathbf{\Theta}}] + S_{\mathrm{int}}[\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}}],$$

unifying Gaiamiti (geometry), Tantrakyma (matter), and their interaction. On a diagnostic realized reference configuration with Good-clock $C = (\hat{T}, U_t)$, we posit invariance under *Good-clock time translations* (homogeneity of emergent time aligned to $\bar{\Sigma}$; Definitions 5.26 and 13.158). By Noether's theorem this continuous symmetry implies a conserved quantity [90, 301].

The Conserved Quantity: The Total Relational Hamiltonian

The conserved charge is the *Total Relational Hamiltonian* $\hat{H}_{\text{univ}}$ (Legendre transform of the total Lagrangian; see Section 13.6):

$$\hat{H}_{\text{univ}} = \hat{H}_{\Phi} + \hat{H}_{\Theta} + \hat{H}_{\text{int}}.$$

Good-clock invariance implies $\frac{d}{dt}\langle \hat{H}_{\text{univ}} \rangle = 0$ on the diagnostic realized reference configuration. In the timeless (constraint) formulation this is encoded by the *Universal Quantum Constraint*

$$\hat{H}_{\text{univ}} \, |\overset{\circ}{\boldsymbol{\Psi}}\rangle = 0 \quad \text{(Theorem 13.168)},$$

which coexists with meaningful *conditional* Good-clock dynamics for subsystems (Page–Wootters; Proposition 13.172). Combining the conserved $\hat{H}_{\text{univ}}$ with the subsystem split and the Good-clock Heisenberg equation yields the diagnostic realized reference configuration-wise energy balance that integrates to $\Delta U_A = Q - W$ (eq. (15.12)), separating coherent driving (work) from incoherent exchange (heat) in accordance with the CP-divisible/unital structure of $\mathbb{E}_A$ [617, 441] and the decoherence picture [170].

Derivations of the First Law for Subsystems

We work on a realized thermodynamic diagnostic realized reference configuration $\mathfrak{B}_{[\chi]}^{\langle \text{DRRC} \rangle}(\mathfrak{P})$ (Definition 15.1) with Good-clock $C = (\hat{T}, U_t)$ aligned to $\bar{\Sigma}$ (Definition 13.158). All time derivatives below are Good-clock (Heisenberg) derivatives and are invariant under clock-gauge changes by Lemma 15.6. Let the universe be partitioned into a subsystem A and its environment E (the rest of the cosmos), with associated subalgebras $\mathcal{A}_A, \mathcal{A}_E \subset \mathcal{A}(\mathcal{C}_d)$.

Definition 15.11 (Subsystem split of the Hamiltonian)**.** On the diagnostic realized reference configuration we use the standard decomposition

$$\hat{H}_{\text{univ}} = \hat{H}_A + \hat{H}_E + \hat{H}_{AE} \quad \textit{(Section 15.1.3)},$$

where $\hat{H}_A$ and $\hat{H}_E$ act on their respective algebras and commute, $[\hat{H}_A, \hat{H}_E] = 0$, and $\hat{H}_{AE}$ encodes the interaction. The *internal energy* of A is

$$U_A(t) := \mathrm{Tr}\Big(\rho_A(t)\, H_A(t)\Big),$$

with ρ_A the reduced (GNS) state (Section 1.4.1) and $H_A(t)$ allowed explicit Good-clock dependence (parametric driving; cf. [230]).

Lemma 15.12 (Good-clock Heisenberg balance for U_A)**.** *Let $\alpha_t(\cdot) = U_t^{\dagger}(\cdot)U_t$ be the Good-clock Heisenberg flow. Then*

$$\frac{dU_A}{dt} = \frac{i}{\hbar}\left\langle\, [\hat{H}_{\text{univ}}, \hat{H}_A] \,\right\rangle + \left\langle\, \partial_t \hat{H}_A \,\right\rangle = \frac{i}{\hbar}\left\langle\, [\hat{H}_{AE}, \hat{H}_A] \,\right\rangle + \left\langle\, \partial_t \hat{H}_A \,\right\rangle.$$

Proof idea. Use $d\hat{H}_A/dt = (i/\hbar)[\hat{H}_{\text{univ}}, \hat{H}_A] + \partial_t \hat{H}_A$ and $[\hat{H}_A, \hat{H}_A] = 0 = [\hat{H}_E, \hat{H}_A]$ by Definition 15.11.

Proof of Lemma 15.12. Step 1 (Good-clock Heisenberg equation). Let $\alpha_t(\cdot) = U_t^\dagger(\cdot)U_t$ denote the Good-clock Heisenberg flow (Definition 13.158). For any observable $X(t)$ with possible explicit time dependence, the Good-clock Heisenberg equation reads

$$\frac{dX(t)}{dt} = \frac{i}{\hbar}\,[\hat{H}_{\text{univ}}(t), X(t)] + \partial_t X(t),$$

so that, in particular,

$$\frac{d\hat{H}_A(t)}{dt} = \frac{i}{\hbar}\,[\hat{H}_{\text{univ}}(t), \hat{H}_A(t)] + \partial_t \hat{H}_A(t).$$

Step 2 (Differentiate the internal energy). In the Heisenberg picture the state is fixed, hence

$$\frac{dU_A}{dt} = \frac{d}{dt}\,\langle \hat{H}_A(t)\rangle = \left\langle \frac{d\hat{H}_A(t)}{dt} \right\rangle = \frac{i}{\hbar}\left\langle [\hat{H}_{\text{univ}}(t), \hat{H}_A(t)] \right\rangle + \left\langle \partial_t \hat{H}_A(t) \right\rangle.$$

Step 3 (Subsystem split and commutators). By the subsystem split (Definition 15.11), $\hat{H}_{\text{univ}} = \hat{H}_A + \hat{H}_E + \hat{H}_{AE}$ with $\hat{H}_A \in \mathcal{A}_A \otimes \mathbf{1}_E$ and $\hat{H}_E \in \mathbf{1}_A \otimes \mathcal{A}_E$. Therefore

$$[\hat{H}_A, \hat{H}_A] = 0, \qquad [\hat{H}_E, \hat{H}_A] = 0,$$

so that

$$[\hat{H}_{\text{univ}}, \hat{H}_A] = [\hat{H}_{AE}, \hat{H}_A].$$

Step 4 (Conclusion and interpretation). Substituting the previous line into Step 2 yields

$$\frac{dU_A}{dt} = \frac{i}{\hbar}\left\langle [\hat{H}_{AE}, \hat{H}_A] \right\rangle + \left\langle \partial_t \hat{H}_A \right\rangle,$$

which is the claimed balance. The first term is the exchange power due to the interaction Hamiltonian $\hat{H}_{AE}$; the second is the driving power arising from explicit Good-clock control of $\hat{H}_A$. Upon time integration this balance underlies the diagnostic realized reference configuration-wise First Law $\Delta U_A = Q - W$ (eq. (15.12), Theorem 15.18, and Definition 15.14). □

Remark 15.13 (Good-clock prerequisites for the Heisenberg balance)**.** The proof of the Good-clock Heisenberg balance uses only the following ingredients from Chapter 15:

(i) Good-clock and Heisenberg flow

The Good-clock and its Heisenberg flow

$$\alpha_t(X) = U_t^\dagger X\, U_t,$$

together with the convention that ∂_t denotes the Good-clock Heisenberg derivative (see Definition 13.158 and the chapter-wide convention that time derivatives are taken with respect to U_t).

(ii) Subsystem split of the universal Hamiltonian

The subsystem split

$$\hat{H}_{\text{univ}} = \hat{H}_A + \hat{H}_E + \hat{H}_{AE},$$

with A-locality implying $[\hat{H}_A, \hat{H}_A] = 0$ and $[\hat{H}_E, \hat{H}_A] = 0$ (see Definition 15.11).

(iii) Diagnostic realized reference configuration-wise internal energy and driving

The diagnostic realized reference configuration-wise internal energy

$$U_A(t) = \langle \hat{H}_A(t) \rangle,$$

together with allowance for explicit Good-clock driving $\partial_t \hat{H}_A$ (see eq. (15.12), Theorem 15.18, and Definition 15.14).

From (1) one immediately has the Good-clock Heisenberg equation for any (possibly explicitly time-dependent) observable:

$$\frac{d\hat{H}_A}{dt} = \frac{i}{\hbar}[\hat{H}_{\text{univ}}, \hat{H}_A] + \partial_t \hat{H}_A.$$

Using (2) yields the commutator reduction $[\hat{H}_{\text{univ}}, \hat{H}_A] = [\hat{H}_{AE}, \hat{H}_A]$, and with (3) this gives

$$\frac{dU_A}{dt} = \frac{i}{\hbar}\left\langle [\hat{H}_{AE}, \hat{H}_A] \right\rangle + \left\langle \partial_t \hat{H}_A \right\rangle,$$

which is precisely the balance asserted in Lemma 15.12. No additional assumptions (e.g. pointer/einselection structure or a GKLS limit) are required.

Definition 15.14 (Heat current and power (diagnostic realized reference configuration-wise, clocked))**.** Define the *heat current* into A and the *power* (work rate by A) as

$$J_A^{(Q)}(t) \;:=\; \frac{i}{\hbar}\left\langle\, [\hat{H}_{AE}, \hat{H}_A] \,\right\rangle, \qquad P_A(t) \;:=\; -\left\langle\, \partial_t \hat{H}_A \,\right\rangle.$$

The sign in P_A matches the thermodynamic convention $\Delta U = Q - W$ with $W = \int P_A \, dt$ the work *done by the system* [119].

Proposition 15.15 (Differential First Law on the diagnostic realized reference configuration)**.** *On* $\mathfrak{B}_{[\chi]}^{\langle \text{DRRC} \rangle}(\mathfrak{P})$ *with Good-clock* C*,*

$$\frac{dU_A}{dt} \;=\; J_A^{(Q)}(t) \;-\; P_A(t).$$

Proof of Proposition 15.15. Step 1 (Good-clock Heisenberg balance). By the Good-clock convention (Definition 13.158), time derivatives are taken in the Heisenberg picture with respect to U_t. Lemma Lemma 15.12 gives

$$\frac{dU_A}{dt} = \frac{i}{\hbar}\left\langle [\hat{H}_{AE}, \hat{H}_A] \right\rangle + \left\langle \partial_t \hat{H}_A \right\rangle.$$

Step 2 (Identify heat current and power). By the diagnostic realized reference configuration-wise definitions of heat and power (Definition 15.14),

$$J_A^{(Q)}(t) := \frac{i}{\hbar}\left\langle [\hat{H}_{AE}, \hat{H}_A] \right\rangle, \qquad P_A(t) := -\left\langle \partial_t \hat{H}_A \right\rangle,$$

where the sign of P_A is fixed so that the integrated form reproduces the First Law $\Delta U_A = Q - W$ (eq. (15.12) and Theorem 15.18).

Step 3 (Assemble the balance). Substituting these identifications from Definition 15.14 into the expression of Lemma 15.12 yields

$$\frac{dU_A}{dt} = J_A^{(Q)}(t) - P_A(t),$$

as claimed. Integrating over any Good-clock interval then gives the diagnostic realized reference configuration-wise First Law $\Delta U_A = \int J_A^{(Q)}\, dt - \int P_A\, dt = Q - W$ in agreement with eq. (15.12) and Theorem 15.18. □

Definition 15.16 (Infinitesimal heat and work)**.** For an interval $[t, t + dt]$ in Good-clock time,

$$\delta Q_A := J_A^{(Q)}(t)\, dt, \qquad \delta W_A := P_A(t)\, dt.$$

Corollary 15.17 (Integral First Law for subsystems)**.** *For any Good-clock interval* $[t_0, t_1]$*,*

$$\Delta U_A = \int_{t_0}^{t_1} J_A^{(Q)}(t)\, dt - \int_{t_0}^{t_1} P_A(t)\, dt = Q_A - W_A,$$

where $Q_A := \int_{t_0}^{t_1} J_A^{(Q)}(t)\, dt$ *and* $W_A := \int_{t_0}^{t_1} P_A(t)\, dt$*.*

Proof of Corollary 15.17. Step 1 (Integrate the differential balance). On a diagnostic realized reference configuration with Good-clock C (Definition 13.158), Proposition Proposition 15.15 states $\frac{dU_A}{dt} = J_A^{(Q)}(t) - P_A(t)$. Integrating over $[t_0, t_1]$ gives

$$\int_{t_0}^{t_1} \frac{dU_A}{dt}\, dt = \int_{t_0}^{t_1} J_A^{(Q)}(t)\, dt - \int_{t_0}^{t_1} P_A(t)\, dt.$$

Step 2 (Fundamental theorem of calculus for U_A). By the Lipschitz/continuity bounds on the reduced dynamics (Lemma 15.60) and the diagnostic realized reference configuration-wise regularity of $H_A(t)$ (Definition 15.14), the map $t \mapsto U_A(t) = \langle \hat{H}_A(t) \rangle$ is absolutely continuous; hence $\int_{t_0}^{t_1} \frac{dU_A}{dt}\, dt = U_A(t_1) - U_A(t_0) = \Delta U_A$. Therefore,

$$\Delta U_A = \int_{t_0}^{t_1} J_A^{(Q)}(t)\, dt - \int_{t_0}^{t_1} P_A(t)\, dt.$$

Step 3 (Heat and work over an interval). By definition (Definition 15.14), $J_A^{(Q)}(t) = \frac{i}{\hbar}\langle [\hat{H}_{AE}, \hat{H}_A] \rangle$ and $P_A(t) = -\langle \partial_t \hat{H}_A \rangle$. Setting $Q_A := \int_{t_0}^{t_1} J_A^{(Q)}(t)\, dt$ and $W_A := \int_{t_0}^{t_1} P_A(t)\, dt$ yields the compact form $\Delta U_A = Q_A - W_A$, in agreement with the diagnostic realized reference configuration-wise First Law (eq. (15.12) and Theorem 15.18).

Step 4 (Clock-gauge invariance). If $t \mapsto \tau = f(t)$ is a Good-clock reparameterization within the gauge class $[C]$ (Definition 13.158), then the Heisenberg derivative and the densities transform covariantly, and the one-forms $J_A^{(Q)}(t)\, dt$ and $P_A(t)\, dt$ are invariant in total measure. Hence both sides of the identity are unchanged under re-timing, as asserted by Lemma 15.6. □

$$\boxed{\Delta U_A \;=\; Q_A \;-\; W_A} \tag{15.12}$$

For an interacting subsystem A (coupled to an environment E with $\hat{H} = \hat{H}_A + \hat{H}_E + \hat{H}_{AE}$—environment Hamiltonian $\hat{H}_E$ plus interaction $\hat{H}_{AE}$ with A), *energy is relational*: along the Good-clock flow, the internal energy

$$U_A(t) := \mathrm{Tr}\Big(\rho_A(t)\, H_A(t)\Big)$$

changes only by heat current into A minus external power on A,

$$\frac{dU_A}{dt} = J_A^{(Q)}(t) - P_A(t),$$

with expectations taken in the emergent inner product (from $\hat{G}_\Lambda$). These rates may be written equivalently as

$$J_A^{(Q)}(t) := \Big\langle i\,[\hat{H}_{AE}, \hat{H}_A]\Big\rangle = \mathrm{Tr}\Big(H_A\,\dot{\rho}_A(t)\Big),$$
$$P_A(t) := -\Big\langle \partial_t \hat{H}_A\Big\rangle = -\,\mathrm{Tr}\Big(\rho_A(t)\,\partial_t H_A(t)\Big).$$

When the interaction splits across the fundamental fields, $\hat{H}_{AE} = \hat{H}_{AE}^{(\mathbf{\Phi})} + \hat{H}_{AE}^{(\mathbf{\Theta})}$, the heat current decomposes componentwise in the Gaiamiti and Tantrakyma sectors:

$$J_A^{(Q)}(t) = \Big\langle i\,[\hat{H}_{AE}^{(\mathbf{\Phi})}, \hat{H}_A]\Big\rangle + \Big\langle i\,[\hat{H}_{AE}^{(\mathbf{\Theta})}, \hat{H}_A]\Big\rangle =: J_{A,\mathbf{\Phi}}^{(Q)}(t) + J_{A,\mathbf{\Theta}}^{(Q)}(t),$$

where $\hat{\mathbf{\Phi}}$ (Gaiamiti) and $\hat{\mathbf{\Theta}}$ (Tantrakyma) are the fundamental fields mediating exchanges. Consequently, over any interval $[t_0, t_1]$,

$$\Delta U_A = \int_{t_0}^{t_1} J_A^{(Q)}(t)\,dt \;-\; \int_{t_0}^{t_1} P_A(t)\,dt \;=\; Q_A - W_A.$$

Theorem 15.18 (First Law of Thermodynamics — DRRC-wise, Good-clock Form)**.** *Work on a realized thermodynamic diagnostic realized reference configuration* $\mathfrak{B}_{[\chi]}^{\langle \mathrm{DRRC}\rangle}(\mathfrak{P})$ *with Good-clock* $C = (\hat{T}, U_t)$ *aligned to* $\bar{\Sigma}$ *(Definitions 13.158 and 15.1). Let the universe be partitioned into a subsystem* A *and its environment* E*, and assume the Hamiltonian split of Definition 15.11. Define* $U_A(t) := \mathrm{Tr}(\rho_A(t) H_A(t))$ *and the heat/power rates as in Definition 15.14.*

(a) Differential form

The Good-clock Heisenberg balance holds:

$$\frac{dU_A}{dt} = J_A^{(Q)}(t) - P_A(t),$$

which is exactly Proposition 15.15 (from Lemma 15.12 with the identifications in Definition 15.14).

(b) Integral form

For any Good-clock interval $[t_0, t_1]$*,*

$$\Delta U_A = \int_{t_0}^{t_1} J_A^{(Q)}(t)\,dt \;-\; \int_{t_0}^{t_1} P_A(t)\,dt = Q_A - W_A,$$

i.e. the First Law (15.12) *(Corollary 15.17; cf. the classical statement [119]).*

(c) Open-system representation (optional)

If, on the diagnostic realized reference configuration, the reduced dynamics of A *is CP-divisible and unital with generator* $\mathcal{L}_t$ *induced by the pointer/channel structure* $\mathbb{E}_A$ *(Definition 15.27 and Theorem 15.50), then*

$$J_A^{(Q)}(t) = \mathrm{Tr}\Big(H_A\,\dot\rho_A(t)\Big), \qquad P_A(t) = -\,\mathrm{Tr}\Big(\rho_A\,\partial_t H_A\Big),$$

consistent with the standard open-system split [617, 441, 170] (cf. Remark 15.19).

(d) Clock-gauge/typed invariance

The statements in (a)–(c) are invariant under Good-clock gauge changes and persist after projection to $\mathcal{H}_{\mathrm{phys}}$ *(and an admissible type* $\Pi^{[\chi]}$*) (Lemma 15.6, Section 12.3, and Theorem 9.351).*

Proof of Theorem 15.18. Setup. Work on a diagnostic realized reference configuration $\mathfrak{B}_{[\chi]}^{\langle \mathrm{DRRC}\rangle}(\mathfrak{P})$ with Good-clock $C = (\hat T, U_t)$ aligned to $\bar\Sigma$ (Definitions 13.158 and 15.1); by Lemma 15.6 any representative of the gauge class $[C]$ may be used. Let the universe be split as $A \cup E$ with Hamiltonian $\hat H = \hat H_A + \hat H_E + \hat H_{AE}$ (Definition 15.11). Define $U_A(t) := \mathrm{Tr}(\rho_A(t)\,H_A(t))$ and use the diagnostic realized reference configuration sign conventions of Definition 15.14.

(a) Differential form. In the Good-clock Heisenberg picture, $\dot O = i[\hat H, O] + \partial_t O$. For $O = \hat H_A$,

$$\frac{d}{dt}\,\langle \hat H_A\rangle = \Big\langle i[\hat H, \hat H_A]\Big\rangle + \Big\langle \partial_t \hat H_A\Big\rangle = \Big\langle i[\hat H_{AE}, \hat H_A]\Big\rangle + \Big\langle \partial_t \hat H_A\Big\rangle,$$

since $[\hat H_A, \hat H_A] = [\hat H_E, \hat H_A] = 0$. With the identifications of Definition 15.14, $J_A^{(Q)}(t) := \langle i[\hat H_{AE}, \hat H_A]\rangle$ and $P_A(t) := -\langle \partial_t \hat H_A\rangle$, we obtain

$$\frac{dU_A}{dt} = J_A^{(Q)}(t) - P_A(t),$$

which is exactly the Good-clock Heisenberg balance stated in Lemma 15.12 and summarized in Proposition 15.15.

(b) Integral form. Integrating the identity in (a) over any Good-clock interval $[t_0, t_1]$ yields

$$\Delta U_A = \int_{t_0}^{t_1} J_A^{(Q)}(t)\,dt - \int_{t_0}^{t_1} P_A(t)\,dt =: Q_A - W_A,$$

which is (15.12) and Corollary 15.17. (As usual, Q_A is positive for net energy *into* A, and W_A is the diagnostic realized reference configuration work *done by* external control on A.)

(c) Open-system representation (optional). On the diagnostic realized reference configuration active band, assume the reduced dynamics of A is CP-divisible and unital with generator $\mathcal{L}_t$ induced by the pointer/channel structure $\mathbb{E}_A$ (Definition 15.27 and Theorem 15.50). In the Schrödinger picture,

$$\frac{d}{dt}U_A(t) = \mathrm{Tr}\Big(\dot\rho_A(t)\, H_A(t)\Big) + \mathrm{Tr}\Big(\rho_A(t)\, \partial_t H_A(t)\Big).$$

Comparing with (a)–(b) and the diagnostic realized reference configuration conventions then gives

$$J_A^{(Q)}(t) = \mathrm{Tr}\Big(H_A\, \dot\rho_A(t)\Big), \qquad P_A(t) = -\,\mathrm{Tr}\Big(\rho_A\, \partial_t H_A\Big),$$

which is the standard open-system split (consistent with [617, 441, 170] and the remarks in Remark 15.19).

(d) Clock-gauge/typed invariance. If $C \sim C'$ are equivalent Good-clocks on the same diagnostic realized reference configuration (Definition 15.3 and Remark 15.4), Lemma 15.6 ensures that time derivatives and current expectations computed with C' agree with those computed with C; hence (a)–(c) are invariant under Good-clock gauge changes. Projecting to $\mathcal{H}_{\mathrm{phys}}$ on $\mathfrak{P}$ and (optionally) to a type $\Pi^{[\chi]}$ preserves all numerical readouts by BRST/Nielsen safety (Section 12.3 and Theorem 9.351), so the statements persist in the realized, typed setting.

Remarks. (i) *Thermal contact.* If $\hat H_{AE} = 0$ on the diagnostic realized reference configuration, then $J_A^{(Q)} \equiv 0$ and changes in U_A are purely work-like via P_A; the law still holds but describes a decoupled limit (cf. the "thermal contact" remark). (ii) *Sign conventions.* The minus sign in $-P_A$ matches the Heisenberg identity once P_A is defined as external power *on* A (Definition 15.14). (iii) *Typed evaluation.* Inserting $\Pi^{[\chi]}$ everywhere defines channelwise balances without altering values when $\Pi^{[\chi]}$ is block-identity on the sector in use (Proposition 9.138).

This proves all parts (a)–(d). □

Remark 15.19 (Open-system representation and CP-divisible reductions)**.** Let the reduced dynamics of A on the diagnostic realized reference configuration be given by a CP-divisible, unital map $\dot\rho_A(t) = \mathcal{L}_t[\rho_A(t)]$ induced by the pointer/channel structure $\mathbb{E}_A$ (Definition 15.27 and Theorem 15.50; cf. [617, 441]). Then the heat current admits the equivalent form

$$J_A^{(Q)}(t) \;=\; \mathrm{Tr}\Big(H_A\, \dot\rho_A(t)\Big),$$

while the power remains $P_A(t) = -\,\mathrm{Tr}(\rho_A\, \partial_t H_A)$ (Good-clock driving). Equivalence with the commutator form follows by tracing the global Liouville–von Neumann equation and using Definition 15.11. This matches the standard open-system split of heat vs. work (cf. [170, 230]).

15.1.4 The Second Law of Relational Information Dynamics

Introductory narrative. In the present framework, the universe is described by a single timeless pure Holo-State $|\mathring{\mathbf{\Psi}}\rangle$ solving the Universal Quantum Constraint $\hat H_{\mathrm{univ}}\,|\mathring{\mathbf{\Psi}}\rangle = 0$ (Theorem 13.168). "Time" for an embedded observer is *emergent*: it appears by conditioning

on a Good-clock $C = (\hat{T}, U_t)$ carried by the environment, producing Page–Wootters conditional dynamics for subsystems (Definition 13.158 and Proposition 13.172). All operational statements are made *diagnostic realized reference configuration-wise*, i.e., on a realized thermodynamic diagnostic realized reference configuration $\mathfrak{B}_{[\chi]}^{\langle \text{diagnosticrealizedreferenceconfiguration} \rangle}(\mathfrak{P})$ (Definition 15.1); derivatives ∂_t are Good-clock (Heisenberg) derivatives, invariant under clock-gauge changes (Lemma 15.6).

Within this setting, local evolution is the unfolding—relative to C—of correlations already latent in $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$. Interactions that create and reshape these correlations are encoded by the *Interaction Kernel* $\mathcal{U}_{\Phi\Theta}(x, x')$, built from the Fourfold Inter-Modal Correlator $\hat{G}_\Lambda$ (Definitions 4.9 and 10.223). For a subsystem A weakly coupled to its environment E, the global unitary generated by $\hat{H}_{\text{univ}} = \hat{H}_A + \hat{H}_E + \hat{H}_{AE}$ disperses phase information into AE correlations. An observer restricted to $\mathcal{A}_A$ then experiences an *effective* loss of accessible information: the reduced state $\rho_A(t)$ mixes and its von Neumann entropy

$$S_A(t) = -\operatorname{Tr}\big[\rho_A(t) \log \rho_A(t)\big] \quad (eq.\ (15.13))$$

tends to increase under the relevant coarse-grained dynamics. This practical irreversibility is not a failure of unitarity; it reflects information being distributed into correlations that are Σ-causally out of reach on the diagnostic realized reference configuration (Definition 5.26 and Corollary 5.145)—the decoherence picture [170, 282].

The formal content is collected in Theorem 15.50: under Good-clock conditioning and admissible coarse-graining, the entropy of any decohered relational subsystem is *nondecreasing* in emergent time, with equality only in the idealized coherence limit. The proof uses (i) the timeless global constraint and purity of $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$ [335]; (ii) emergent time via Page–Wootters/relational conditioning [335, 197, 3]; and (iii) correlator-mediated exchange of relational information between Gaiamiti and Tantrakyma sectors (Definitions 4.9 and 10.223; cf. [29, 237]).

Foundational principles and working hypotheses The derivation rests on the synthesis of core tenets already established, together with minimal operational hypotheses that fix the entropy production scale on the diagnostic realized reference configuration:

(F1) Timeless global state and zero total entropy The universal Hamiltonian annihilates the Holo–State,

$$\hat{H}_{\text{univ}}\, |\overset{\circ}{\boldsymbol{\Psi}}\rangle = 0$$

(Theorem 13.168). The universe is globally pure, so the total entropy is zero [335].

(F2) Good-clocked emergent dynamics A Good-clock $C = (\hat{T}, U_t)$ aligns the time flow with $\bar{\Sigma}$ and supports conditional Schrödinger evolution for subsystems (Definition 13.158 and Proposition 13.172). Statements are invariant in the clock gauge class $[C]$ (Lemma 15.6).

(F3) Interaction via relational correlators Couplings are mediated by $\mathcal{U}_{\Phi\Theta}$ built from $\hat{G}_\Lambda$ (Definitions 4.9 and 10.223), ensuring Σ-causal support for exchanges (Definition 5.26 and Corollary 5.145).

(F4) Admissible pointer dynamics (coarse-graining) On the Good-clock resolution adopted for thermodynamics, the reduced dynamics on $\mathcal{A}_A$ is CP-divisible and *unital*, with generator $\mathcal{L}_t$ and channel family $\{\Phi_{t\leftarrow s}\}$ induced by the diagnostic realized reference configuration relational pointer $\mathbb{E}_A$ (Definition 15.27; cf. [617, 441]). Unitality captures informational openness without biased erasure and implies that $S_A(t)$ is nondecreasing (quantum H-theorem for unital CPTP evolutions).

(F5) Equality conditions / coherence limit Entropy production vanishes iff the reduced dynamics is (DRRC-wise) unitary on A, or A is dynamically decoupled in Good-clock time—our "small-coupling coherence limit'' (Lemma 15.51 and Proposition 15.52). Physically, this occurs when $\hat{H}_{AE} \to 0$ or when the channel is a unitary $\mathrm{Ad}_{U_A(t)}$.

Interpretation. In information-theoretic terms, the Second Law is a data-processing constraint on Born valuations for clocked observers: coarse-grained, Σ-causal, unital dynamics cannot decrease the observer's entropy of the reduced state. The arrow of time is then the direction of increasing subsystem–environment entanglement and record formation [335, 3, 170]. The typed/realized formulation is stable under BRST/Nielsen audits (Section 12.3 and Theorem 9.351) and "same-DRRC'' comparisons (Remark 15.4)—ensuring that entropy production is a property of the diagnostic realized reference configuration and Good-clock, not of arbitrary time parametrizations.

The Formal Derivation of the Second Law The derivation proceeds in three logically complete stages, from emergent time to a coarse–grained, monotone–entropy dynamics for decohered subsystems. All ingredients are internal to the IQR framework. (We will appeal later to the standard spectral fact that *pinching—i.e. conditional expectation onto a commuting record algebra—majorizes spectra* and hence does not decrease von Neumann entropy; cf. [356, 287].)

Stage 1: Emergent directed process (triadic time and t_{geom}). We formalize the emergence of a directed, diagnostic realized reference configuration-physical time parameter and the associated conditional dynamics.

Definition 15.20 (Good-clocked directed time on a diagnostic realized reference configuration)**.** On a diagnostic realized reference configuration $\mathfrak{B}_{[\chi]}^{\langle \mathrm{DRRC}\rangle}(\mathfrak{P})$ (Definition 15.1), fix a Good-clock $C = (\hat{T}, U_t)$ aligned with $\bar{\Sigma}$ (Definition 13.158). The *directed time set* $(\mathbb{T}, [C], \prec)$ consists of: (i) a parameter manifold $\mathbb{T} \subset \mathbb{R}$, (ii) the clock gauge class $[C]$ (reparameterizations $t \mapsto \phi(t)$ with $\phi'(t) > 0$; Definition 15.3), and (iii) the induced total order $\prec$ from any representative $C \in [C]$. All derivatives ∂_t are Heisenberg derivatives w.r.t. U_t and are invariant under $C \mapsto C' \in [C]$ by Lemma 15.6.

Definition 15.21 (Triadic time and geometric time selection)**.** Let $\hat{G}_\Lambda$ be the Fourfold correlator (Definition 4.9). The IQR temporal kinematics is *triadic*, with

$$\vec{t}(x, x') = \big(t_{\mathrm{corr}},\, t_{\mathrm{vac}},\, t_{\mathrm{spec}}\big),$$

whose axes are induced respectively by the off–diagonal, coincident–diagonal, and commutator/spectral sectors of $\hat{G}_\Lambda$ (see Section 4.12 and Definition 4.523). A Good-clock C fixes a

calibrated axis (or admissible linear combination) of $\vec{t}$ to parameterize reduced dynamics. In the macroscopic timelike regime one identifies the *geometric* proper time

$$t_{\mathrm{geom}}(x,x') = \left[\bar{g}_{\mu\nu}(x)\,\Delta x^{\mu}\Delta x^{\nu}\right]^{1/2} \quad (\text{timelike } \Delta x),$$

and the chosen relational temporal functional coincides with t_{geom} on timelike displacements (Definitions 5.26 and 5.95).

Proposition 15.22 (Existence of conditional reduced dynamics)**.** *Let the universe decompose as $A \cup E$ with $\hat{H}_{\mathrm{univ}} = \hat{H}_A + \hat{H}_E + \hat{H}_{AE}$ and $|\mathring{\mathbf{\Psi}}\rangle$ obeying the universal constraint $\hat{H}_{\mathrm{univ}}\,|\mathring{\mathbf{\Psi}}\rangle = 0$ (Theorem 13.168). Fix a Good-clock $C = (\hat{T}, U_t)$ on a diagnostic realized reference configuration (Definitions 13.158 and 15.1). Then there exists a family of completely positive, trace-preserving maps on* states *of A,*

$$\Phi_{t\leftarrow s} :\ \mathsf{D}(\mathcal{H}_A) \to \mathsf{D}(\mathcal{H}_A), \qquad \rho_A(t) = \Phi_{t\leftarrow s}\Big(\rho_A(s)\Big),$$

with the properties:

(i) Identity and cocycle.
$\Phi_{s\leftarrow s} = \mathrm{id}$ *and* $\Phi_{t\leftarrow r}\circ\Phi_{r\leftarrow s} = \Phi_{t\leftarrow s}$ *for all* $s \le r \le t$.

(ii) Clock-gauge invariance.
The family $\{\Phi_{t\leftarrow s}\}$ is invariant within the Good-clock gauge class $[C]$ (Lemma 15.6).

(iii) Σ-causal support.
$\Phi_{t\leftarrow s}$ *does not enlarge Σ-causal support: reduced evolutions respect the causal cones of $\bar{\Sigma}$ (Definition 5.26 and Corollary 5.145).*

Proof idea. Construct a *clocked* Page–Wootters instrument $\mathcal{I}^{AE}_{t\leftarrow s}$ on AE (Proposition 13.172), choose a CP *assignment* $\mathcal{J}_s : \mathsf{D}(\mathcal{H}_A) \to \mathsf{D}(\mathcal{H}_{AE})$ that prepares the environment conditioned on the clock at s but independent of $\rho_A(s)$, and reduce with a diagnostic realized reference configuration pointer channel $\mathbb{E}_A$ followed by partial trace on E. This yields

$$\Phi_{t\leftarrow s} = \mathrm{Tr}_E \circ \mathbb{E}_A \circ \mathcal{I}^{AE}_{t\leftarrow s} \circ \mathcal{J}_s,$$

which is CPTP by composition. Identity, cocycle, and gauge invariance follow from the instrument's comb/composition rules, idempotence and transport-covariance of $\mathbb{E}_A$, and Lemma 15.6. Σ-causality is inherited from global microcausality and the locality of the pointer/restriction maps.

Proof. 1) Clocked global unitary and PW instrument. The constraint $\hat{H}_{\mathrm{univ}}\,|\mathring{\mathbf{\Psi}}\rangle = 0$ and Good-clock $C = (\hat{T}, U_t)$ furnish a family of unitaries $U_{t\leftarrow s} := U_t U_s^{\dagger}$. Page–Wootters conditioning on the clock at time t defines a completely positive instrument $\mathcal{I}^{AE}_{t\leftarrow s}$ on AE (trace-nonincreasing on each outcome, trace-preserving when summed over outcomes), with the *normalized* transformation from s to t being CPTP (Proposition 13.172). For clock-ordered times, the family satisfies the comb/cocycle rule $\mathcal{I}^{AE}_{t\leftarrow r}\circ\mathcal{I}^{AE}_{r\leftarrow s} = \mathcal{I}^{AE}_{t\leftarrow s}$ and $\mathcal{I}^{AE}_{s\leftarrow s} = \mathrm{id}$.

2) CP assignment at the initial time. Fix a CP assignment $\mathcal{J}_s : \mathsf{D}(\mathcal{H}_A) \to \mathsf{D}(\mathcal{H}_{AE})$ of the form

$$\mathcal{J}_s(\rho_A) = \rho_A \otimes \sigma_E(s),$$

Here, $\sigma_E(s)$ is the environment's conditioned reference state at clock time s, obtained from the diagnostic realized reference configuration records and independent of ρ_A (pointer-induced independence). The map $\mathcal{J}_s$ is completely positive and trace-preserving (CPTP) and transport-covariant on the diagnostic realized reference configuration.

3) Reduction to A. Let $\mathbb{E}_A$ be the diagnostic realized reference configuration pointer channel on A (unital, idempotent CPTP; Definition 15.27); let Tr_E denote partial trace over E. Define the reduced evolution

$$\boxed{\Phi_{t\leftarrow s} \;:=\; \mathrm{Tr}_E \circ \mathbb{E}_A \circ \mathcal{I}^{AE}_{t\leftarrow s} \circ \mathcal{J}_s \;:\; \mathsf{D}(\mathcal{H}_A) \to \mathsf{D}(\mathcal{H}_A)}\,.$$

As a composition of CP maps it is CP; trace preservation follows from the trace preservation of the normalized $\mathcal{I}^{AE}_{t\leftarrow s}$, the unitality/idempotence of $\mathbb{E}_A$, and Tr_E.

4) Identity and cocycle. Identity:

$$\Phi_{s\leftarrow s} = \mathrm{Tr}_E \circ \mathbb{E}_A \circ \mathcal{I}^{AE}_{s\leftarrow s} \circ \mathcal{J}_s = \mathrm{Tr}_E \circ \mathbb{E}_A \circ \mathcal{J}_s = \mathrm{Tr}_E(\rho_A \otimes \sigma_E(s)) = \rho_A,$$

since $\mathbb{E}_A$ acts trivially on A when no monitoring has occurred and $\mathrm{Tr}_E(\rho_A \otimes \sigma_E) = \rho_A$.

For the cocycle, pick $s \le r \le t$. Use the comb rule, idempotence/transport-covariance of $\mathbb{E}_A$, and a *Markov compatibility* of the assignment,

$$\mathcal{J}_r \circ \mathrm{Tr}_E \circ \mathbb{E}_A \circ \mathcal{I}^{AE}_{r\leftarrow s} \circ \mathcal{J}_s = \mathcal{I}^{AE}_{r\leftarrow s} \circ \mathcal{J}_s,$$

which holds because $\mathcal{J}_r(\cdot) = (\cdot) \otimes \sigma_E(r)$ and $\sigma_E(r)$ is exactly the environment's (diagnostic realized reference configuration-record) state produced by $\mathcal{I}^{AE}_{r\leftarrow s} \circ \mathcal{J}_s$ independently of ρ_A. Then

$$\begin{aligned}\Phi_{t\leftarrow r} \circ \Phi_{r\leftarrow s} &= \mathrm{Tr}_E \circ \mathbb{E}_A \circ \mathcal{I}^{AE}_{t\leftarrow r} \circ \mathcal{J}_r \circ \mathrm{Tr}_E \circ \mathbb{E}_A \circ \mathcal{I}^{AE}_{r\leftarrow s} \circ \mathcal{J}_s \\ &= \mathrm{Tr}_E \circ \mathbb{E}_A \circ \mathcal{I}^{AE}_{t\leftarrow r} \circ \mathcal{I}^{AE}_{r\leftarrow s} \circ \mathcal{J}_s = \mathrm{Tr}_E \circ \mathbb{E}_A \circ \mathcal{I}^{AE}_{t\leftarrow s} \circ \mathcal{J}_s = \Phi_{t\leftarrow s}.\end{aligned}$$

5) Clock-gauge invariance. If $t \mapsto \tau = f(t)$ is a reparameterization within $[C]$, Lemma 15.6 implies that the PW instrument (and thus $\Phi_{\tau\leftarrow\sigma}$) transforms covariantly, leaving all diagnostic realized reference configuration numbers unchanged. Because $\mathbb{E}_A$ and $\mathcal{J}_s$ are transport-covariant, $\{\Phi_{t\leftarrow s}\}$ is invariant within the gauge class.

6) Σ-causal support. Global dynamics respects $\bar{\Sigma}$-microcausality; PW conditioning is local in the clock record; $\mathbb{E}_A$ and Tr_E are local CPTP operations. Therefore the reduced maps cannot create influence outside the $\bar{\Sigma}$ cones (no-law-signalling), i.e., they do not enlarge Σ-causal support (Corollary 5.145).

This proves (a)–(c). □

Lemma 15.23 (Generator form of the conditional evolution)**.** *Let $\{\Phi_{t\leftarrow s}\}_{t\ge s}$ be the conditional reduced dynamics of Proposition 15.22. If $t \mapsto \Phi_{t\leftarrow s}$ is differentiable at fixed s, then there exists a time–local (not a priori GKLS) generator $\mathcal{L}_t$ such that*

$$\frac{d}{dt}\,\rho_A(t) \;=\; \mathcal{L}_t[\rho_A(t)] \;=\; -\,i\,[H_{A,\mathrm{eff}}(t), \rho_A(t)] \;+\; \mathcal{M}_t[\rho_A(t)],$$

where $H_{A,\mathrm{eff}}(t)$ is a self–adjoint, transport–covariant conditional Hamiltonian *(defined up to an additive multiple of the identity), and $\mathcal{M}_t$ collects the non–Hamiltonian (monitoring/exchange) contributions induced by the interaction $\hat{H}_{AE}$ and any diagnostic realized reference configuration monitoring on A. If $\hat{H}_{AE} = 0$ (and no diagnostic realized reference configuration monitoring on A is active), then $\mathcal{M}_t \equiv 0$.*

Proof idea. Differentiate $\rho_A(t) = \Phi_{t\leftarrow s}(\rho_A(s))$ and use the cocycle at $r = t$ to obtain a *left* time–local generator $\mathcal{L}_t = \frac{d}{dt}\Phi_{t\leftarrow t}$. Decompose $\mathcal{L}_t$ on the operator space (with the GNS/Hilbert–Schmidt inner product) into an inner derivation $-i[H_{A,\mathrm{eff}}(t), \cdot]$ plus a remainder $\mathcal{M}_t$. When $\hat{H}_{AE} = 0$ (and no monitoring acts on A), the reduced dynamics is unitary on A, hence $\mathcal{M}_t = 0$. Transport covariance follows from parallel transport $\mathcal{U}(x \leftarrow x'; \gamma)$ along $\bar{\Sigma}$.

Proof. (1) Time–local form from the cocycle. By Proposition 15.22 the maps satisfy $\Phi_{s\leftarrow s} = \mathrm{id}$ and $\Phi_{t\leftarrow r} \circ \Phi_{r\leftarrow s} = \Phi_{t\leftarrow s}$. Fix s and assume $t \mapsto \Phi_{t\leftarrow s}$ is differentiable. Using the cocycle with $r = t$,

$$\Phi_{t\leftarrow s} = \Phi_{t\leftarrow t} \circ \Phi_{t\leftarrow s}.$$

Differentiating at t yields

$$\frac{d}{dt}\Phi_{t\leftarrow s} = \left(\frac{d}{dt}\Phi_{t\leftarrow t}\right) \circ \Phi_{t\leftarrow s}.$$

Define the (left) generator $\mathcal{L}_t := \frac{d}{dt}\Phi_{t\leftarrow t}$. Then, for $\rho_A(t) := \Phi_{t\leftarrow s}[\rho_A(s)]$,

$$\frac{d}{dt}\rho_A(t) = \mathcal{L}_t[\rho_A(t)].$$

(2) Hamiltonian vs. non–Hamiltonian split. Endow $\mathcal{B}(\mathcal{H}_A)$ with the GNS/Hilbert–Schmidt inner product $\langle X, Y\rangle = \mathrm{Tr}\big(X^\dagger Y\big)$ (induced by the emergent inner product). The space of inner derivations $\{-i[H, \cdot] \mid H = H^\dagger\}$ is a linear subspace. Let P_{der} be the orthogonal projector (with respect to $\langle\cdot,\cdot\rangle$) onto the closure of that subspace. Set

$$\mathcal{L}_t^{(\mathrm{H})} := P_{\mathrm{der}}(\mathcal{L}_t) = -i[H_{A,\mathrm{eff}}(t), \cdot], \qquad \mathcal{M}_t := \mathcal{L}_t - \mathcal{L}_t^{(\mathrm{H})}.$$

Then $H_{A,\mathrm{eff}}(t) = H_{A,\mathrm{eff}}(t)^\dagger$ is unique up to $c(t)\,\mathbf{1}$, and $\mathcal{M}_t$ is linear, trace–preserving and Hermiticity–preserving (since $\mathcal{L}_t$ is).

(3) Physical identification. From the construction in Proposition 15.22, $\Phi_{t\leftarrow s}$ results from composing a clocked Page–Wootters instrument on AE with a (local, unital, idempotent) pointer channel on A and the E–reduction. The coherent, unitary part on A defines $H_{A,\mathrm{eff}}(t)$, whereas interaction/monitoring contributions (originating from $\hat{H}_{AE}$ and any active diagnostic realized reference configuration monitoring on A) populate $\mathcal{M}_t$. This matches the Good–clock Heisenberg balance for $U_A(t)$ and the entropy–increasing role of pointer pinching used later in the H–theorem.

(4) Decoupled limit. If $\hat{H}_{AE} = 0$ and no monitoring acts on A, the conditional evolution on A is unitary: $\Phi_{t\leftarrow s} = \mathrm{Ad}_{U_A(t\leftarrow s)}$. Hence $\mathcal{L}_t = -i[H_{A,\mathrm{eff}}(t), \cdot]$ and $\mathcal{M}_t \equiv 0$.

(5) Transport covariance. Under parallel transport $\mathcal{U}(x \leftarrow x'; \gamma)$ along the $\bar{\Sigma}$–compatible connection, $H_{A,\mathrm{eff}}(t)$ transforms covariantly, so $-i[H_{A,\mathrm{eff}}(t), \cdot]$ and $\mathcal{L}_t$ are frame–independent on the diagnostic realized reference configuration (cf. (4.11)).

Combining (1)–(5) gives the claimed generator form. □

Remark 15.24 (Directedness, macroscopic identification, and causality)**.** The orientation of $\mathbb{T}$ is fixed by $[C]$ (Definition 15.3), hence all dynamical statements are *directed* and clock–gauge invariant. In macroscopic timelike regimes the parameter coincides with t_{geom} (Definition 15.21), and $\Phi_{t\leftarrow s}$ preserves Σ-causal support (Corollary 5.145). The structural split in Lemma 15.23 prepares Stage 2, where—on adopting the thermodynamic coarse–graining of the diagnostic realized reference configuration—the map family becomes CP–divisible and *unital*, leading to entropy monotonicity (Theorem 15.50; cf. [617, 441]).

Stage 2: Entropy production engine (current–kernel coupling and pointer channel from $\mathcal{U}_{\Phi\Theta}$). Geometry–matter reciprocity is explicit in the operator equations: source currents $\hat{J}^{\Phi}_{(\Theta)}$ and $\hat{J}^{\Theta}_{(\Phi)}$ intertwine sectors through the interaction kernel built from the unified correlator (Section 10.19.4 and Definition 4.9).

Definition 15.25 (Current–kernel interaction)**.** With $\mathcal{U}_{\Phi\Theta}(x,x')$ the interaction kernel, define

$$\hat{H}_{AE} \;:=\; \int_{\mathcal{X}_q} \mathrm{d}x\,\mathrm{d}x'\;\mathrm{Tr}\Big[\hat{J}^{\Phi}_{(\Theta)}(x)\,\mathcal{U}_{\Phi\Theta}(x,x')\,\hat{J}^{\Theta}_{(\Phi)}(x')\Big]\,,$$

where the trace contracts internal indices and the transporter in $\mathcal{U}$ aligns fibres along $\bar{\Sigma}$–chronogeodesics.

Lemma 15.26 (Self–adjointness and transport covariance)**.** *If the inter–modal kernel $\mathcal{U}_{\Phi\Theta}$ obeys the reciprocity and parallel–transport symmetries inherited from the Fourfold correlator $\hat{G}_{\Lambda}$ (Definition 4.9 and eq. (4.11)), then the interaction Hamiltonian $\hat{H}_{AE}$ defined by the current–kernel pairing is self–adjoint, $\hat{H}_{AE}=\hat{H}_{AE}^{\dagger}$, and is covariant under $\bar{\Sigma}$–parallel transport along chronogeodesics (Definition 5.26).*

Proof. Setup. By the current–kernel definition (Definition 15.25),

$$\hat{H}_{AE} \;=\; \int_{\mathcal{X}_q} \mathrm{d}x\,\mathrm{d}x'\;\mathrm{Tr}_{\mathrm{modal}}\Big[\,\hat{J}^{\Phi}_{(\Theta)}(x)\,\mathcal{U}_{\Phi\Theta}(x,x')\;\hat{J}^{\Theta}_{(\Phi)}(x')\,\Big],$$

where $\hat{J}^{\Phi}_{(\Theta)}(x)$ and $\hat{J}^{\Theta}_{(\Phi)}(x')$ are conserved current observables (hence self–adjoint) on the diagnostic realized reference configuration, and $\mathrm{Tr}_{\mathrm{modal}}$ denotes trace over inter–modal indices only (no Hilbert–space trace). The kernel $\mathcal{U}_{\Phi\Theta}$ is induced by $\hat{G}_{\Lambda}$ and satisfies the reciprocity relation $\mathcal{U}_{\Phi\Theta}(x,x')^{\dagger}=\mathcal{U}_{\Theta\Phi}(x',x)$ and the transport rule given by path–ordered parallel transport (Definition 4.9 and eq. (4.11)).

(i) Self–adjointness. Using $(\mathrm{Tr}\,X)^{\dagger}=\mathrm{Tr}(X^{\dagger})$, $(ABC)^{\dagger}=C^{\dagger}B^{\dagger}A^{\dagger}$, and the self–adjointness of the currents, we obtain

$$\begin{aligned}\hat{H}_{AE}^{\dagger} &= \int \mathrm{d}x\,\mathrm{d}x'\;\mathrm{Tr}_{\mathrm{modal}}\Big[\hat{J}^{\Theta}_{(\Phi)}(x')\,\mathcal{U}_{\Phi\Theta}(x,x')^{\dagger}\;\hat{J}^{\Phi}_{(\Theta)}(x)\Big]\\ &= \int \mathrm{d}x\,\mathrm{d}x'\;\mathrm{Tr}_{\mathrm{modal}}\Big[\hat{J}^{\Theta}_{(\Phi)}(x')\,\mathcal{U}_{\Theta\Phi}(x',x)\;\hat{J}^{\Phi}_{(\Theta)}(x)\Big].\end{aligned}$$

Relabel $(x,\Phi)\leftrightarrow(x',\Theta)$ and use cyclicity of the trace:

$$\hat{H}_{AE}^{\dagger}=\int \mathrm{d}x\,\mathrm{d}x'\;\mathrm{Tr}_{\mathrm{modal}}\Big[\hat{J}^{\Phi}_{(\Theta)}(x)\,\mathcal{U}_{\Phi\Theta}(x,x')\;\hat{J}^{\Theta}_{(\Phi)}(x')\Big]=\hat{H}_{AE}.$$

(When the active band is finite–dimensional—as on the diagnostic realized reference configuration—boundedness is automatic, so symmetric $\Rightarrow$ self–adjoint.)

(ii) Transport covariance. Let $\Pi(\cdot)$ denote the path–ordered parallel transport along a $\bar{\Sigma}$–chronogeodesic. Under a change of local transport frame,

$$\hat{J}^{\Phi}_{(\Theta)}(x)\mapsto\Pi(x)\,\hat{J}^{\Phi}_{(\Theta)}(x)\,\Pi(x)^{\dagger},\qquad \mathcal{U}_{\Phi\Theta}(x,x')\mapsto\Pi(x)\,\mathcal{U}_{\Phi\Theta}(x,x')\,\Pi(x')^{\dagger},$$

and similarly for $\hat{J}^{\Theta}_{(\Phi)}(x')$. Substituting these into the integrand gives

$$\begin{aligned}&\mathrm{Tr}_{\mathrm{modal}}\Big[\Pi(x)\hat{J}^{\Phi}_{(\Theta)}(x)\Pi(x)^{\dagger}\;\Pi(x)\mathcal{U}_{\Phi\Theta}(x,x')\Pi(x')^{\dagger}\;\Pi(x')\hat{J}^{\Theta}_{(\Phi)}(x')\Pi(x')^{\dagger}\Big]\\&\quad=\mathrm{Tr}_{\mathrm{modal}}\Big[\hat{J}^{\Phi}_{(\Theta)}(x)\;\mathcal{U}_{\Phi\Theta}(x,x')\;\hat{J}^{\Theta}_{(\Phi)}(x')\Big],\end{aligned}$$

by $\Pi^{\dagger}\Pi=\mathbf{1}$ and cyclicity of trace. Hence the value of the integral is unchanged—equivalently, $\hat{H}_{AE}$ is form–invariant (covariant) under $\bar{\Sigma}$–parallel transport.

Conclusion. With reciprocity and transport symmetries inherited from $\hat{G}_{\Lambda}$, the current–kernel interaction Hamiltonian is self–adjoint and covariant under $\bar{\Sigma}$–parallel transport, as claimed. □

In the decohered regime selected by this coupling, the A–E interaction induces a *record (pointer) algebra* on A (Sections 14.1 and 14.2); let $\{P_k\}_k$ denote its orthogonal projectors. Stability of these records is reinforced by the linear Tantrakyma dynamics on $\bar{\Sigma}$, $(\bar{\nabla}^{\mu}\bar{\nabla}_{\mu}-\mathcal{M}_{\Theta})\,\delta\hat{\boldsymbol{\Theta}}=0$ (eq. (10.159)).

Definition 15.27 (Relational pointer channel)**.** Let $\{P_k\}_k$ be the orthogonal record projectors on $\mathfrak{B}_A$ (record algebra of A) with $\sum_k P_k=\mathbb{I}_A$. Define the conditional expectation (einselection map)

$$\mathbb{E}_A:\;\mathcal{A}_A\to\mathcal{A}_A,\qquad \mathbb{E}_A(\rho)\;:=\;\sum_k P_k\,\rho\,P_k,\quad \rho\in\mathcal{A}_A.$$

Then $\mathcal{R}_A:=\{X\in\mathcal{A}_A:\;[X,P_k]=0\;\forall k\}$ is the (commutative) record algebra and $\mathbb{E}_A$ is the unique faithful normal conditional expectation onto $\mathcal{R}_A$ that fixes $\mathcal{R}_A$ pointwise. Moreover, $\mathbb{E}_A$ is CPTP, idempotent ($\mathbb{E}_A^2=\mathbb{E}_A$), unital ($\mathbb{E}_A(\mathbb{I}_A)=\mathbb{I}_A$), and transport–covariant with the parallel transporter $\mathcal{U}$ (eq. (4.11)); i.e., it commutes with fibre alignment along $\bar{\Sigma}$–chronogeodesics (Definition 5.26).

Lemma 15.28 (Kraus form and basic properties)**.** *The pointer map $\mathbb{E}_A$ has Kraus operators $\{P_k\}$, hence is completely positive, trace–preserving, unital, and idempotent (a pinching). Moreover,*

$$\mathbb{E}_A(\rho)=\rho\quad\Longleftrightarrow\quad[\rho,P_k]=0\;\;\forall k,$$

i.e. the fixed points are exactly the operators commuting with all record projectors (the commutant of the record algebra). In finite dimension, $\mathbb{E}_A$ admits a random–unitary (doubly stochastic) representation, and the von Neumann entropy is monotone:

$$S\Big(\mathbb{E}_A(\rho)\Big)\;\geq\;S(\rho).$$

Proof. (1) Kraus form, CPTP, unitality, idempotence. By the definition of the pointer channel,

$$\mathbb{E}_A(\rho)=\sum_k P_k\,\rho\,P_k,\qquad P_jP_k=\delta_{jk}P_k,\quad\sum_k P_k=\mathbb{I}_A.$$

Thus $\{K_k:=P_k\}$ is a Kraus family with $\sum_k K_k^{\dagger}K_k=\sum_k P_k=\mathbb{I}_A$, so $\mathbb{E}_A$ is CPTP. Also $\mathbb{E}_A(\mathbb{I}_A)=\sum_k P_k=\mathbb{I}_A$ (unital), and

$$\mathbb{E}_A\Big(\mathbb{E}_A(\rho)\Big)=\sum_{j,k}P_j\,(P_k\rho P_k)\,P_j=\sum_k P_k\rho P_k=\mathbb{E}_A(\rho),$$

so $\mathbb{E}_A$ is idempotent (pinching).

(2) Fixed points. ($\Rightarrow$) If $\mathbb{E}_A(\rho) = \rho$, then for $\ell \neq k$, $P_\ell \rho P_k = P_\ell\Big(\sum_j P_j \rho P_j\Big)P_k = P_\ell P_k\, \rho P_k = 0$, hence the off–diagonal blocks vanish. Consequently $\rho P_k = \sum_j P_j \rho P_k = P_k \rho P_k = P_k \rho$, i.e. $[\rho, P_k] = 0$ for all k. ($\Leftarrow$) If $[\rho, P_k] = 0$ for all k, then $P_k \rho P_k = P_k \rho = \rho P_k$ and $\mathbb{E}_A(\rho) = \sum_k P_k \rho P_k = \sum_k P_k \rho = \rho$. Thus $\mathrm{Fix}(\mathbb{E}_A) = \{\rho : [\rho, P_k] = 0\ \forall k\}$ (the commutant of the record algebra).

(3) Random–unitary representation and entropy monotonicity (finite dimension). For finite $N := \#\{P_k\}$ define commuting unitaries

$$U_m := \sum_{k=0}^{N-1} \omega^{mk} P_k, \qquad \omega = e^{2\pi i/N}, \quad m = 0, \dots, N-1.$$

Using $\frac{1}{N}\sum_{m=0}^{N-1} \omega^{m(k-\ell)} = \delta_{k\ell}$,

$$\frac{1}{N}\sum_{m=0}^{N-1} U_m\, \rho\, U_m^\dagger = \sum_{k,\ell}\Big(\tfrac{1}{N}\sum_m \omega^{m(k-\ell)}\Big) P_k \rho P_\ell = \sum_k P_k \rho P_k = \mathbb{E}_A(\rho).$$

Hence $\mathbb{E}_A$ is random–unitary (therefore doubly stochastic). By the pinching/majorization result (Karamata) for block pinching, the spectrum of $\mathbb{E}_A(\rho)$ majorizes that of ρ, and the von Neumann entropy, being Schur–concave, satisfies $S(\mathbb{E}_A(\rho)) \geq S(\rho)$. □

Proposition 15.29 (Stroboscopic reduced step is CPTP and unital)**.** *Fix a step $\Delta t \geq 0$. Define the stroboscopic map*

$$\Phi_{\Delta t}(\rho) \;:=\; \mathbb{E}_A\Big(U_A(\Delta t)\, \rho\, U_A(\Delta t)^\dagger\Big),$$

where $U_A(\Delta t)$ is a unitary on A generated by the effective Hamiltonian on the step (e.g., $U_A(\Delta t) = e^{-iH_{A,\mathrm{eff}}\Delta t}$ in the time–independent case, or $U_A(\Delta t, t) := \mathcal{T}\exp\Big[-i\int_t^{t+\Delta t} H_{A,\mathrm{eff}}(\tau)\, d\tau\Big]$ in general). Then $\Phi_{\Delta t}$ is completely positive, trace–preserving, and unital. Moreover, von Neumann entropy is invariant under the unitary part, so any entropy production in the step is entirely due to the pinching $\mathbb{E}_A$:

$$S\Big(\Phi_{\Delta t}(\rho)\Big) \;\geq\; S(\rho), \qquad S\Big(\Phi_{\Delta t}(\rho)\Big) - S(\rho) = S\Big(\mathbb{E}_A(\sigma)\Big) - S(\sigma) \quad (\sigma := U_A \rho U_A^\dagger).$$

Equality holds iff $U_A(\Delta t)\rho U_A(\Delta t)^\dagger$ commutes with all record projectors $\{P_k\}$ (i.e., iff it lies in the record algebra's commutant).

Proof. (1) CPTP. The pointer map has Kraus form $\mathbb{E}_A(\rho) = \sum_k P_k \rho P_k$ with orthogonal projectors $\{P_k\}$, hence is CPTP (Lemma 15.28). Unitary conjugation $\mathrm{Ad}_{U_A}(\rho) = U_A \rho U_A^\dagger$ is CPTP. Therefore $\Phi_{\Delta t} = \mathbb{E}_A \circ \mathrm{Ad}_{U_A}$ is CPTP.

(2) Unitality. Since $\mathrm{Ad}_{U_A}(\mathbb{I}_A) = \mathbb{I}_A$ and $\mathbb{E}_A(\mathbb{I}_A) = \mathbb{I}_A$ (Lemma 15.28),

$$\Phi_{\Delta t}(\mathbb{I}_A) = \mathbb{E}_A(\mathbb{I}_A) = \mathbb{I}_A,$$

so $\Phi_{\Delta t}$ is unital.

(3) Entropy balance and production. Let $\sigma := \mathrm{Ad}_{U_A}(\rho) = U_A \rho U_A^\dagger$. Unitary invariance of von Neumann entropy gives $S(\sigma) = S(\rho)$. By the pinching/majorization property of $\mathbb{E}_A$ (Lemma 15.28 and Proposition 15.31),

$$S\Big(\Phi_{\Delta t}(\rho)\Big) = S\Big(\mathbb{E}_A(\sigma)\Big) \;\geq\; S(\sigma) = S(\rho).$$

Thus all entropy production in the step is due to $\mathbb{E}_A$. The inequality becomes equality iff $\mathbb{E}_A(\sigma) = \sigma$, i.e., $[\sigma, P_k] = 0$ for all k (Lemma 15.28). □

Lemma 15.30 (Pinching increases entropy (data processing))**.** *For any state ρ supported on the diagnostic realized reference configuration subsystem A,*

$$S\Big(\mathbb{E}_A(\rho)\Big) \;\geq\; S(\rho), \qquad S(\rho) := -\mathrm{Tr}(\rho \log \rho).$$

Equality holds iff $[\rho, P_k] = 0$ for all record projectors $\{P_k\}$ (i.e., iff ρ is block–diagonal in the pointer decomposition).

Proof. Step 1 (Pointer channel is a unital pinching/conditional expectation). By the definition and properties of the relational pointer channel (Definition 15.27 and Lemma 15.28),

$$\mathbb{E}_A(\rho) = \sum_k P_k \,\rho\, P_k, \qquad P_j P_k = \delta_{jk} P_k, \quad \sum_k P_k = \mathbb{I}_A,$$

so $\mathbb{E}_A$ is CPTP, unital, and idempotent (a pinching onto the commutative record algebra).

Step 2 (Work on the active band; unital map fixes the maximally mixed state). On the diagnostic realized reference configuration we restrict to the state–conditioned active band $\mathcal{H}_{\mathrm{act}}$ (Definition 15.56), which is finite–dimensional by Assumption 15.39. Let $d := \dim \mathcal{H}_{\mathrm{act}}$ and set $\sigma := \mathbb{I}_A/d$. Since $\mathbb{E}_A$ is unital,

$$\mathbb{E}_A(\sigma) = \sigma.$$

Step 3 (Data processing for relative entropy ⇒ entropy increase). Umegaki relative entropy is monotone under CPTP maps (data processing). Applying this to $\mathbb{E}_A$ and the pair (ρ, σ),

$$D(\rho\|\sigma) \;\geq\; D\Big(\mathbb{E}_A(\rho) \,\|\, \mathbb{E}_A(\sigma)\Big) = D\Big(\mathbb{E}_A(\rho) \,\|\, \sigma\Big).$$

Using $D(\tau\|\mathbb{I}/d) = -S(\tau) + \log d$ for any state τ on a d–dimensional support, we get

$$-S(\rho) + \log d \;\geq\; -S\Big(\mathbb{E}_A(\rho)\Big) + \log d \;\Rightarrow\; S\Big(\mathbb{E}_A(\rho)\Big) \;\geq\; S(\rho).$$

Step 4 (Alternative majorization route and equality condition). Independently, since $\mathbb{E}_A$ is a pinching onto a commutative algebra, in finite dimension it is random–unitary and doubly stochastic (Lemma 15.28). Hence the output spectrum majorizes the input spectrum (Proposition 15.31), and Schur–concavity of S gives $S(\mathbb{E}_A(\rho)) \geq S(\rho)$. Moreover, equality holds iff $\mathbb{E}_A(\rho) = \rho$, which by Lemma 15.28 is equivalent to $[\rho, P_k] = 0$ for all k (no off–diagonal blocks in the pointer decomposition).

This completes the proof. □

Proposition 15.31 (Pinching monotonicity via block–diagonalization and Karamata)**.** *Let $\{P_k\}_{k=1}^m$ be mutually orthogonal projections with $\sum_k P_k = \mathbb{I}$. Define the pinching $\mathbb{E}(\rho) := \sum_k P_k \rho P_k$. Then, on any finite–dimensional support,*

$$S(\mathbb{E}(\rho)) \;\geq\; S(\rho).$$

Proof idea. *(i)* Simultaneously block–diagonalize $\{P_k\}$ with a unitary W; pinching deletes off–diagonal blocks. *(ii)* Twirl over $U_j := \sum_k \omega^{jk} \widetilde{P}_k$ to express $\mathbb{E}$ as an average of unitary conjugations (random–unitary, hence doubly stochastic). *(iii)* By Ky Fan's maximum principle, $\lambda(\mathbb{E}(\rho)) \prec \lambda(\rho)$. *(iv)* Karamata with $f(x) = x \log x$ (convex) gives $S(\mathbb{E}(\rho)) \geq S(\rho)$.

Proof. (1) Block–diagonalization. Because the P_k commute and are mutually orthogonal with $\sum_k P_k = \mathbb{I}$, there exists a unitary W such that $\widetilde{P}_k := W^\dagger P_k W$ are diagonal in disjoint blocks, i.e. $\sum_k \widetilde{P}_k = \mathbb{I}$ and each $\widetilde{P}_k = \begin{bmatrix} \mathbb{I}_{d_k} & 0 \\ 0 & 0 \end{bmatrix}$ on its block. Writing $\tilde{\rho} := W^\dagger \rho W$,

$$\widetilde{\mathbb{E}}(\tilde{\rho}) := W^\dagger \mathbb{E}(\rho) W = \sum_k \widetilde{P}_k \, \tilde{\rho} \, \widetilde{P}_k,$$

which removes the off–diagonal blocks of $\tilde{\rho}$. Since von Neumann entropy is unitary invariant, it suffices to prove the claim for $\widetilde{\mathbb{E}}$ acting on $\tilde{\rho}$.

(2) Fourier twirl $\Rightarrow$ random–unitary. Let m be the number of blocks and set $\omega = e^{2\pi i/m}$. Define commuting unitaries

$$U_j := \sum_{k=1}^{m} \omega^{jk} \, \widetilde{P}_k, \qquad j = 0, \ldots, m-1.$$

Using $\frac{1}{m} \sum_{j=0}^{m-1} \omega^{j(k-\ell)} = \delta_{k\ell}$,

$$\frac{1}{m} \sum_{j=0}^{m-1} U_j \, \tilde{\rho} \, U_j^\dagger = \sum_{k,\ell} \Big(\tfrac{1}{m} \sum_j \omega^{j(k-\ell)} \Big) \widetilde{P}_k \, \tilde{\rho} \, \widetilde{P}_\ell = \sum_k \widetilde{P}_k \, \tilde{\rho} \, \widetilde{P}_k = \widetilde{\mathbb{E}}(\tilde{\rho}).$$

Hence $\widetilde{\mathbb{E}}$ (and therefore $\mathbb{E}$) is a random–unitary channel with equal weights $1/m$; in particular, it is unital and trace–preserving (doubly stochastic).

(3) Majorization via Ky Fan's principle. Let $\lambda(X) = (\lambda_1(X), \ldots, \lambda_d(X))$ be the eigenvalues of Hermitian X in nonincreasing order. Ky Fan's principle says that for any $1 \leq r \leq d$,

$$\sum_{i=1}^{r} \lambda_i(X) = \max_{\substack{0 \leq Q \leq \mathbb{I} \\ \operatorname{rank}(Q) = r}} \operatorname{Tr}(QX).$$

Using convexity of the pointwise maximum and the random–unitary form from Step 2,

$$\sum_{i=1}^{r} \lambda_i\Big(\widetilde{\mathbb{E}}(\tilde{\rho})\Big) = \max_Q \; \operatorname{Tr}\Big(Q \, \frac{1}{m} \sum_j U_j \tilde{\rho} U_j^\dagger \Big) \leq \frac{1}{m} \sum_j \max_Q \; \operatorname{Tr}\Big(Q \, U_j \tilde{\rho} U_j^\dagger \Big).$$

For each j, the maximizer is $Q = U_j P^{(r)} U_j^\dagger$, where $P^{(r)}$ projects onto the top-r eigenspace of $\tilde{\rho}$; thus

$$\sum_{i=1}^{r} \lambda_i\Big(\widetilde{\mathbb{E}}(\tilde{\rho})\Big) \leq \frac{1}{m} \sum_j \operatorname{Tr}\Big(P^{(r)} \, \tilde{\rho} \Big) = \sum_{i=1}^{r} \lambda_i(\tilde{\rho}).$$

Since equality holds at $r = d$ by trace preservation, we conclude $\lambda\Big(\widetilde{\mathbb{E}}(\tilde{\rho})\Big) \;\prec\; \lambda(\tilde{\rho})$.

(4) Karamata ⇒ entropy increase. With $f(x) = x \log x$ convex on $(0, 1]$, Karamata's inequality applied to the majorization in Step 3 yields

$$\sum_{i=1}^{d} f\Big(\lambda_i(\widetilde{\mathbb{E}}(\tilde{\rho}))\Big) \;\le\; \sum_{i=1}^{d} f\Big(\lambda_i(\tilde{\rho})\Big),$$

i.e. $S(\widetilde{\mathbb{E}}(\tilde{\rho})) \ge S(\tilde{\rho})$. Unitary invariance gives $S(\mathbb{E}(\rho)) = S(\widetilde{\mathbb{E}}(\tilde{\rho}))$ and $S(\rho) = S(\tilde{\rho})$, hence $S(\mathbb{E}(\rho)) \ge S(\rho)$.

This completes the proof. □

Remark 15.32 (Equality condition)**.** Equality in the pinching inequality holds if and only if ρ is already block–diagonal in the record basis, i.e.

$$S\Big(\mathbb{E}(\rho)\Big) = S(\rho) \quad\Longleftrightarrow\quad \mathbb{E}(\rho) = \rho \quad\Longleftrightarrow\quad [\rho, P_k] = 0 \;\; \forall k,$$

so that $\rho \in \mathrm{Fix}(\mathbb{E})$ (the commutant of the record algebra). Equivalently:

(a) Fixed-point/commutant form

The conditional expectation

$$\mathbb{E}(\rho) = \sum_k P_k \rho P_k$$

satisfies $\mathbb{E}(\rho) = \rho$ iff $[\rho, P_k] = 0$ for all k (Lemma 15.28).

(b) Block-structure form

The state is block diagonal in the record basis,

$$\rho \;=\; \bigoplus_k P_k \rho P_k,$$

so all inter-block coherences vanish.

(c) Majorization/Karamata form

In Proposition 15.31, the spectral majorization $\lambda(\mathbb{E}(\rho)) \prec \lambda(\rho)$ saturates, hence $S(\mathbb{E}(\rho)) = S(\rho)$, exactly when the off–diagonal blocks are zero (otherwise the inequality is strict).

(d) Data processing form

In Lemma 15.30, equality in

$$D(\rho \| \mathbb{I}/d) \;\ge\; D(\mathbb{E}(\rho) \| \mathbb{I}/d)$$

occurs precisely when ρ is a fixed point of the unital conditional expectation $\mathbb{E}$, i.e. when ρ is block–diagonal in the record basis.

(e) Operational stroboscopic form

For the step $\Phi_{\Delta t} = \mathbb{E}_A \circ \mathrm{Ad}_{U_A}$ of Proposition 15.29,

$$S(\Phi_{\Delta t}(\rho)) = S(\rho)$$

iff $U_A(\Delta t)\rho U_A(\Delta t)^\dagger$ commutes with all P_k, i.e. the unitary has not created pointer off–diagonals prior to pinching.

Proposition 15.33 (One–step entropy production)**.** *For every state ρ on A and each $\Delta t \geq 0$,*

$$S\Big(\Phi_{\Delta t}(\rho)\Big) \;\geq\; S(\rho),$$

where $\Phi_{\Delta t} = \mathbb{E}_A \circ \mathrm{Ad}_{U_A(\Delta t)}$ is the stroboscopic step of Proposition 15.29. Equality holds iff $U_A(\Delta t)\rho U_A(\Delta t)^\dagger$ commutes with all record projectors $\{P_k\}$ (equivalently, iff $\Phi_{\Delta t}(\rho) = \rho$).

Proof. Step 1 (Unitary transport leaves entropy invariant). By Proposition 15.29,

$$\Phi_{\Delta t}(\rho) = \mathbb{E}_A\Big(U_A(\Delta t)\,\rho\,U_A(\Delta t)^\dagger\Big), \qquad \text{with } U_A(\Delta t) \text{ unitary.}$$

Set $\sigma := U_A(\Delta t)\rho U_A(\Delta t)^\dagger$. Von Neumann entropy is unitarily invariant, hence $S(\sigma) = S(\rho)$. *Step 2 (Pinching increases entropy).* By the pinching/data–processing lemma Lemma 15.30,

$$S\Big(\mathbb{E}_A(\sigma)\Big) \;\geq\; S(\sigma).$$

Using $\mathbb{E}_A(\sigma) = \Phi_{\Delta t}(\rho)$,

$$S\Big(\Phi_{\Delta t}(\rho)\Big) \;\geq\; S(\sigma) = S(\rho).$$

Step 3 (All production is due to $\mathbb{E}_A$ and equality condition). Decompose the difference:

$$S\Big(\Phi_{\Delta t}(\rho)\Big) - S(\rho) = \underbrace{S\Big(\mathbb{E}_A(\sigma)\Big) - S(\sigma)}_{\geq 0 \text{ by } \textit{Lemma } 15.30} + \underbrace{S(\sigma) - S(\rho)}_{=0 \text{ (unitary invariance)}}.$$

Thus any one–step increase is entirely due to the pinching $\mathbb{E}_A$, consistent with Proposition 15.29. Moreover, equality holds iff $\mathbb{E}_A(\sigma) = \sigma$, i.e. iff $[\sigma, P_k] = 0$ for all k (equivalently, $\Phi_{\Delta t}(\rho) = \rho$), by Lemma 15.28 and Remark 15.32.

Edge case $\Delta t = 0$. Then $U_A(0) = \mathbb{I}$ and $\Phi_0 = \mathbb{E}_A$, so the claim reduces to Lemma 15.30. □

Corollary 15.34 (Discrete monotone chain)**.** *For the stroboscopic trajectory $\rho_A^{(n)} := (\Phi_{\Delta t})^n\Big(\rho_A^{(0)}\Big)$ at times $t_n := n\,\Delta t$,*

$$S\Big(\rho_A^{(n+1)}\Big) \;\geq\; S\Big(\rho_A^{(n)}\Big) \qquad (\forall\, n \geq 0).$$

Proof. By definition, $\rho_A^{(n+1)} = \Phi_{\Delta t}(\rho_A^{(n)})$. Applying the one–step entropy production result (Proposition 15.33) with input state $\rho = \rho_A^{(n)}$ yields

$$S\Big(\rho_A^{(n+1)}\Big) = S\Big(\Phi_{\Delta t}(\rho_A^{(n)})\Big) \;\geq\; S\Big(\rho_A^{(n)}\Big),$$

for every $n \geq 0$, establishing monotonicity.

Equality condition (stepwise). From Proposition 15.33 and the fixed–point characterization of $\mathbb{E}_A$ (Lemma 15.28 and Remark 15.32), equality at step $n \to n{+}1$ holds iff

$$\mathbb{E}_A\Big(U_A(\Delta t)\,\rho_A^{(n)}\,U_A(\Delta t)^\dagger\Big) = U_A(\Delta t)\,\rho_A^{(n)}\,U_A(\Delta t)^\dagger,$$

equivalently $[\,U_A(\Delta t)\,\rho_A^{(n)}\,U_A(\Delta t)^\dagger,\; P_k] = 0$ for all k.

Boundedness (finite dimension). On the state–conditioned active band $\mathcal{H}_{\mathrm{act}}$ (Definition 15.56 and Assumption 15.39) with $d := \dim \mathcal{H}_{\mathrm{act}} < \infty$, the sequence $\{S(\rho_A^{(n)})\}_{n\geq 0}$ is non–decreasing and bounded above by $\log d$, hence convergent. □

Theorem 15.35 (Continuous–time entropy production under CP–divisible, unital dynamics)**.** *Let $\{\Phi_{t\leftarrow s}\}_{t\geq s}$ be the Good–clock reduced dynamics of A from Proposition 15.22, and assume the thermodynamic coarse–graining on the diagnostic realized reference configuration makes it CP–divisible and* unital*: each propagator $\Phi_{t\leftarrow s}$ is CPTP, satisfies $\Phi_{t\leftarrow s}(\mathbb{I}) = \mathbb{I}$, and obeys the cocycle $\Phi_{t\leftarrow r} \circ \Phi_{r\leftarrow s} = \Phi_{t\leftarrow s}$. Then the von Neumann entropy is nondecreasing:*

$$t_2 \geq t_1 \;\Rightarrow\; S\Big(\rho_A(t_2)\Big) \;\geq\; S\Big(\rho_A(t_1)\Big).$$

Proof idea. Fix $t_2 > t_1$ and view $\rho_A(t_2) = \Phi_{t_2\leftarrow t_1}(\rho_A(t_1))$ with $\Phi_{t_2\leftarrow t_1}$ CPTP and unital. Data processing for Umegaki relative entropy with reference $\sigma = \mathbb{I}/d$ on the active band yields the claim. Alternatively, partition $[t_1, t_2]$ and iterate one–step monotonicity (Proposition 15.33 and Corollary 15.34).

Proof. Step 1 (Unital CPTP propagator on the interval). By CP–divisibility and the cocycle of Proposition 15.22,

$$\rho_A(t_2) \;=\; \Phi_{t_2\leftarrow t_1}\Big(\rho_A(t_1)\Big), \qquad \Phi_{t_2\leftarrow t_1} \text{ is CPTP and unital.}$$

Step 2 (Data processing to monotonicity). Restrict to the state–conditioned active band $\mathcal{H}_{\mathrm{act}}$ (Definition 15.56 and Assumption 15.39) with $d := \dim \mathcal{H}_{\mathrm{act}} < \infty$, and set $\sigma := \mathbb{I}/d$. Unitality gives $\Phi_{t_2\leftarrow t_1}(\sigma) = \sigma$. Umegaki relative entropy is monotone under CPTP maps, hence

$$D\Big(\rho_A(t_1)\,\|\,\sigma\Big) \;\geq\; D\Big(\Phi_{t_2\leftarrow t_1}(\rho_A(t_1))\,\|\,\Phi_{t_2\leftarrow t_1}(\sigma)\Big) = D\Big(\rho_A(t_2)\,\|\,\sigma\Big).$$

Since $D(\tau\|\mathbb{I}/d) = -S(\tau) + \log d$, we obtain $S(\rho_A(t_2)) \geq S(\rho_A(t_1))$.

Step 3 (Discrete partition route, optional). If $t_1 = t_0^{(n)} < \cdots < t_N^{(n)} = t_2$ partitions the interval, then $\rho_A(t_2) = \big(\prod_{k=1}^{N} \Phi_{t_k^{(n)}\leftarrow t_{k-1}^{(n)}}\big)\,\rho_A(t_1)$ with each step unital CPTP. Applying Proposition 15.33 at each substep yields the monotone chain (Corollary 15.34), and letting the mesh $\max_k(t_k^{(n)} - t_{k-1}^{(n)}) \to 0$ recovers the same conclusion.

Step 4 (Clock–gauge invariance). For any Good–clock reparameterization $t \mapsto \tau = f(t)$ within $[C]$, Lemma 15.6 implies $\Phi^{(\tau)}_{\tau_2\leftarrow\tau_1} = \Phi_{t_2\leftarrow t_1}$, so the argument is gauge–invariant.

Equality condition (finite–dimensional support). Equality $S(\rho_A(t_2)) = S(\rho_A(t_1))$ holds iff the DPI is tight for the pair $(\rho_A(t_1), \sigma)$, i.e. iff $\rho_A(t_1)$ lies in a decoherence–free (fixed–point) subalgebra on which $\Phi_{t_2\leftarrow t_1}$ acts unitarily (equivalently, the Petz recovery condition holds for $\sigma = \mathbb{I}/d$). In the stroboscopic "unitary + pinching'' case, this reduces to the stepwise commuting/fixed–point condition of Remark 15.32 and Proposition 15.29.

This proves the monotonicity claim. □

Remark 15.36 (Clock gauge, causality, and typed realization)**.** All statements are invariant in the Good–clock class $[C]$ (Lemma 15.6); Σ–causal support is preserved by construction (Definition 5.26 and Corollary 5.145). After projection to $\mathcal{H}_{\mathrm{phys}}{=}H^0(\hat{Q}_{\mathrm{BRST}})$ and an admissible type $\Pi^{[\chi]}$, the same conclusions hold by BRST/Nielsen safety (Section 12.3 and Theorem 9.351).

Stage 3: Irreversibility, causal support, and the continuum limit. *Endogenous regularisation from Chapter 4.* The coherence/memory scales used below are not imposed *ad hoc* but arise from the relational spectrum of the Holo–State. (i) **State–conditioned modal weights** $w_n(x, x')$ (Definition 4.392, §4.9.7) quantify the contribution of the nth spectral mode to bilocal probes and obey polynomial decay controlled by Spectral Smoothness (Axiom 4.415), ensuring summability of spectral aggregates. (ii) **Spectral smoothness** bounds the shear observables and coincident self–correlators, providing UV control of mode–by–mode contributions and continuity across loci (Axiom 4.415). (iii) **Coherent regularisation** then follows: the emergent geometry is internally regulated by the state's own coherence profile, without external cutoffs (Proposition 4.424). (iv) These ingredients define an **effective spectral cutoff** $\Lambda_{\text{eff}}(x, x'; \varepsilon)$ (Definition 4.576), with a quantitative bound

$$\Lambda_{\text{eff}}(x, x'; \varepsilon) \;\leq\; \max\left\{\lambda_{n_0},\; \left(\frac{C' C_f}{\varepsilon}\right)^{\frac{1}{\alpha-\beta}}\right\} \quad \text{(eq. (4.278))},$$

which induces the **PQS coherence width** $\ell_{\text{PQS}}(x, x'; \varepsilon) := \Lambda_{\text{eff}}(x, x'; \varepsilon)^{-1}$ (§4.15.3). Whenever Planck–band support is operational in the Holo–State, one obtains sub–Planckian control of the PQS width (Proposition 4.588). (v) Finally, **state–dependent spectral gapping** (e.g. a graviton gap) renders long–range correlations inert on scales set by an effective mass m_{eff} (§4.9.15, Definition 4.438 and Proposition 4.439), further shortening memory.

Mathematical reminders (scales). With tolerance $\varepsilon \in (0, 1)$, define the coherence/memory times by

$$\tau_{\text{mem}}(x; \varepsilon) := \Lambda_{\text{cutoff}}(x; \varepsilon)^{-1}, \qquad \tau_{\text{IR}}(x; \varepsilon) := \Lambda_{\text{IR}}(x; \varepsilon)^{-1},$$

where Λ_{cutoff} and Λ_{IR} are the state–conditioned UV/IR thresholds determined by $\{w_n\}$ and the observable features $f_n^{(\text{obs})}$ (Definition 4.576, §4.9.15). In the *coarse–grained, stroboscopic* regime $\tau_{\text{mem}} \ll \Delta t \ll \tau_{\text{obs}}$, pinching by the record algebra (Definition 15.27) suppresses off–diagonals at each step, enabling a Markovian product limit (Stage 2).

Causal architecture from Chapter 5. Expectation–level chronogeometry $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (Definition 5.26) induces the Lorentzian cone structure (Definition 5.77) and the *emergent chronogeometric manifold* $\mathcal{M}_{\text{geom}}$ (Definition 5.113). The *effective light–cone* and unified causal structure are fixed by $\bar{g}^{\text{Lor}}$ (Definitions 5.122 and 5.150), yielding Σ–causal support for admissible processes. Monitoring/measurement operations confined to the record algebra respect this causal architecture and cannot generate superluminal influence (Corollary 5.145). Anisotropies in the interaction kernel $\mathcal{U}_{\Phi\Theta}$ are naturally encoded via the triadic/time sector and channel–selective operations, leading to direction–dependent dissipation/monitoring rates while preserving causality (Section 5.4.3 and Remark 5.61).

Conclusion for Stage 3 (setup). The endogenous regularisation from Chapter 4 (ℓ_{PQS}, w_n, spectral smoothness, m_{eff}) provides finite coherence/memory and IR control. Together with the $\bar{\Sigma}$–causal architecture of Chapter 5 ($\mathcal{M}_{\text{geom}}$, effective light–cones, no–law–signalling), this justifies a *unital, CP–divisible* Markovian continuum limit of the reduced dynamics that is confined to $\bar{\Sigma}$–causal support and admits channel–anisotropic rates when $\mathcal{U}_{\Phi\Theta}$ is anisotropic (to be formalised below in Definition 15.37, Assumption 15.39, and Propositions 15.41 and 15.43).

Terminology for reduced dynamics.

Definition 15.37 (Admissible (CP–divisible) reduced dynamics)**.** Let $\{\Lambda_t\}_{t\geq 0}$ be the Good–clock reduced evolution on $\mathfrak{B}_A$ obtained by Page–Wootters conditioning and diagnostic realized reference configuration coarse–graining (Proposition 15.22 and Definition 15.27). We call $\{\Lambda_t\}$ *admissible (CP–divisible)* if the following equivalent conditions hold:

(a) Two–parameter family

For all $t \geq s \geq 0$ there exists a CPTP map $V_{t,s}$ such that

$$\Lambda_t \;=\; V_{t,s}\,\Lambda_s, \qquad V_{t,s}\circ V_{s,r} \;=\; V_{t,r}.$$

(b) Time–local generator

The time–local master equation $\dot\rho_t = \mathcal{L}_t[\rho_t]$ exists with

$$\mathcal{L}_t[\rho] \;=\; -\,i\,[H_{A,\mathrm{eff}}(t),\rho] \;+\; \sum_{\alpha}\gamma_\alpha(t)\Big(L_\alpha(t)\,\rho\,L_\alpha^\dagger(t) - \tfrac{1}{2}\{L_\alpha^\dagger(t)L_\alpha(t),\rho\}\Big)\,,$$

where $\gamma_\alpha(t)\geq 0$ for all t (GKLS (Gorini–Kossakowski–Lindblad–Sudarshan) form; cf. [617, 441]).

(c) IQR construction (product limit)

For every $t \geq 0$,

$$\Lambda_t \;=\; \lim_{n\to\infty}\Big(\,\mathbb{E}_A \;\circ\; \mathrm{Ad}_{e^{-iH_{A,\mathrm{eff}}\,t/n}}\Big)^n, \qquad \mathrm{Ad}_U(\rho) := U\rho U^\dagger,$$

with $\mathbb{E}_A$ the pointer conditional expectation (Definition 15.27).

If, in addition, $\Lambda_t(\mathbb{I}_A) = \mathbb{I}_A$ for all t, we say $\{\Lambda_t\}$ is *unital*.

Remark 15.38 (Avoiding confusion with charge–parity)**.** Here "CP" = *Completely Positive*. When referring to the discrete symmetry of particle physics we write $\mathcal{CP}$. CP–divisibility in Definition 15.37 is unrelated to $\mathcal{CP}$ or its violation.

Assumption 15.39 (Endogenous coherence–scaled Markov limit)**.** *Fix* $|\mathring{\Psi}\rangle$ *and tolerance* $\varepsilon \in (0,1)$. *By Spectral Smoothness and Finite Entanglement Capacity there exist state–conditioned thresholds*

$$\lambda_{\mathrm{IR}}(x;\varepsilon) \;\leq\; \lambda_n \;\leq\; \lambda_{\mathrm{cutoff}}(x;\varepsilon),$$

defining a finite active relational band (Reciprocal Resolution Limits, Coherent Regularization). Let the induced coherence times be $\tau_{\mathrm{mem}} := \lambda_{\mathrm{cutoff}}^{-1}$, $\tau_{\mathrm{IR}} := \lambda_{\mathrm{IR}}^{-1}$. *Assume the stroboscopic step* Δt *and observation window* τ_{obs} *obey*

$$\tau_{\mathrm{mem}} \;\ll\; \Delta t \;\ll\; \tau_{\mathrm{obs}},$$

and the PQS width ℓ_{PQS} *attenuates coherence below the width (with possible ultra–sub–Planckian sharpening). If the environment induces an effective gap* $m_{\mathrm{eff}} > 0$ *(state–dependent spectral gapping), then as* $\Delta t \to 0$ *at fixed monitoring rate the stroboscopic maps* $\Phi_{\Delta t} = \mathbb{E}_A\circ\mathrm{Ad}_{e^{-iH_{A,\mathrm{eff}}\Delta t}}$ *(Proposition 15.29) converge to a* unital, CP–divisible *quantum dynamical family* $\{\Lambda_t\}_{t\geq 0}$ *on* $\mathfrak{B}_A$ *with time–independent GKLS generator* $\mathcal{L}$.

Lemma 15.40 (Unitality inherited from the pointer)**.** *If $\mathbb{E}_A$ is unital (i.e. $\mathbb{E}_A(\mathbb{I}_A) = \mathbb{I}_A$), then every stroboscopic step*

$$\Phi_{\Delta t} \;=\; \mathbb{E}_A \circ \mathrm{Ad}_{e^{-iH_{A,\mathrm{eff}}\Delta t}}$$

is unital. Consequently, any uniform product limit in Definition 15.37(c) is unital: $\Lambda_t(\mathbb{I}_A) = \mathbb{I}_A$ for all $t \geq 0$.

Proof. (1) Unitary conjugation is unital. For any unitary U on $\mathcal{H}_A$,

$$\mathrm{Ad}_U(\mathbb{I}_A) = U\,\mathbb{I}_A\,U^\dagger = \mathbb{I}_A.$$

In particular, with $U_A(\Delta t) = e^{-iH_{A,\mathrm{eff}}\Delta t}$, $\mathrm{Ad}_{U_A(\Delta t)}(\mathbb{I}_A) = \mathbb{I}_A$.

(2) Composition with a unital pointer channel is unital. By Definition 15.27 and Lemma 15.28, the pointer channel $\mathbb{E}_A$ is CPTP and unital: $\mathbb{E}_A(\mathbb{I}_A) = \mathbb{I}_A$. Hence, for the stroboscopic step,

$$\Phi_{\Delta t}(\mathbb{I}_A) = \mathbb{E}_A\Big(\mathrm{Ad}_{U_A(\Delta t)}(\mathbb{I}_A)\Big) = \mathbb{E}_A(\mathbb{I}_A) = \mathbb{I}_A,$$

so $\Phi_{\Delta t}$ is unital (cf. Proposition 15.29).

(3) Unitality is preserved by uniform product limits. Consider the product family in Definition 15.37(c):

$$\Lambda_t^{(n)} \;:=\; \Big(\,\mathbb{E}_A \circ \mathrm{Ad}_{e^{-iH_{A,\mathrm{eff}}\,t/n}}\,\Big)^n, \qquad n \in \mathbb{N}.$$

Each factor is unital by Step (2), hence every $\Lambda_t^{(n)}$ is unital and $\Lambda_t^{(n)}(\mathbb{I}_A) = \mathbb{I}_A$ for all t, n. If $\Lambda_t^{(n)} \to \Lambda_t$ uniformly in t (in the topology specified in Definition 15.37(c)), then evaluating at $\mathbb{I}_A$ gives

$$\Lambda_t(\mathbb{I}_A) = \lim_{n\to\infty} \Lambda_t^{(n)}(\mathbb{I}_A) = \mathbb{I}_A,$$

so Λ_t is unital for all $t \geq 0$.

Conclusion. Unitality of $\mathbb{E}_A$ implies unitality of each stroboscopic step and of any uniform product limit as in Definition 15.37(c). □

Proposition 15.41 (Chernoff–type product limit and unital GKLS generator (weak monitoring))**.** *Work on the finite active band (Assumption 15.39). Let the effective Hamiltonian on A be $H_{A,\mathrm{eff}}$ and let $\Pi := \mathbb{E}_A$ be the (unital, idempotent) pointer pinching (Definition 15.27 and Lemma 15.28). For a* Poisson–thinned *monitoring with diagnostic realized reference configuration–fixed rate $\kappa \geq 0$, define the Δ–step map*

$$V_\Delta \;:=\; \Big((1-\kappa\Delta)\,I + \kappa\Delta\,\Pi\Big) \;\circ\; \mathrm{Ad}_{e^{-iH_{A,\mathrm{eff}}\Delta}} \qquad (\Delta > 0).$$

Then, for each $t \geq 0$, the Chernoff product limit

$$\Lambda_t \;=\; \lim_{n\to\infty} \Big(V_{t/n}\Big)^n$$

exists (uniformly on compact t–intervals) and equals a unital CPTP semigroup $e^{t\mathcal{L}}$ with GKLS generator

$$\boxed{\mathcal{L}[\rho] \;=\; -\,i\,[H_{A,\mathrm{eff}}, \rho] \;+\; \kappa\Big(\Pi(\rho) - \rho\Big).}$$

Equivalently, in Lindblad form one may take $L_k = \sqrt{\kappa}\, P_k$ for the record projectors $\{P_k\}$ of $\Pi(\rho) = \sum_k P_k \rho P_k$ (Definition 15.27), yielding

$$\mathcal{L}[\rho] \;=\; -\,i[H_{A,\mathrm{eff}}, \rho] + \sum_k \Big(L_k \rho L_k^\dagger - \tfrac{1}{2}\{L_k^\dagger L_k, \rho\} \Big), \qquad \sum_k L_k^\dagger L_k = \kappa\, \mathbb{I}_A,$$

and the semigroup is unital by Lemma 15.40.

Proof idea. V_Δ is a CPTP contraction with $V_0 = I$; expand to first order: $V_\Delta = I + \Delta\Big(-i[H_{A,\mathrm{eff}}, \cdot] + \kappa(\Pi - I)\Big) + o(\Delta)$. Chernoff's theorem then gives $\lim_{n\to\infty}(V_{t/n})^n = e^{t\mathcal{L}}$ with the stated $\mathcal{L}$. Writing $\Pi(\rho) = \sum_k P_k \rho P_k$ yields the Lindblad form with $L_k = \sqrt{\kappa} P_k$. Unitality follows from Lemma 15.40.

Proof. (1) First–order expansion. On the finite active band (Assumption 15.39) all relevant norms are equivalent. For small Δ,

$$\mathrm{Ad}_{e^{-iH_{A,\mathrm{eff}}\Delta}} = I + \Delta\, \mathcal{L}^{\mathrm{H}} + o(\Delta), \qquad \mathcal{L}^{\mathrm{H}}[\rho] := -\,i[H_{A,\mathrm{eff}}, \rho].$$

Hence

$$V_\Delta = \Big((1 - \kappa\Delta) I + \kappa\Delta\, \Pi \Big) \circ (I + \Delta \mathcal{L}^{\mathrm{H}} + o(\Delta)) = I + \Delta\Big(\mathcal{L}^{\mathrm{H}} + \kappa(\Pi - I) \Big) + o(\Delta).$$

(2) Chernoff derivative and product limit. Set $\mathcal{L} := \mathcal{L}^{\mathrm{H}} + \kappa(\Pi - I)$. Then $\lim_{\Delta\downarrow 0} \frac{V_\Delta - I}{\Delta} = \mathcal{L}$ strongly on $\mathcal{B}(\mathcal{H}_A)$. Each V_Δ is CPTP (convex combination of CPTP maps composed with a unitary) and unital (Lemma 15.40 applies to Π; convexity preserves unitality). By Chernoff's product theorem, for every $t \geq 0$,

$$\lim_{n\to\infty} \Big(V_{t/n} \Big)^n \;=\; e^{t\mathcal{L}}$$

uniformly on compact t–sets.

(3) GKLS/Lindblad form and unitality. Using $\Pi(\rho) = \sum_k P_k \rho P_k$,

$$\kappa(\Pi - I)[\rho] = \sum_k \Big(L_k \rho L_k^\dagger - \tfrac{1}{2}\{L_k^\dagger L_k, \rho\} \Big), \quad L_k := \sqrt{\kappa}\, P_k, \quad \sum_k L_k^\dagger L_k = \kappa \mathbb{I}_A,$$

which is standard GKLS. Since $[H_{A,\mathrm{eff}}, \mathbb{I}_A] = 0$ and $\sum_k L_k^\dagger L_k \propto \mathbb{I}_A$, the generator is unital and $e^{t\mathcal{L}}$ is a unital CPTP semigroup (Lemma 15.40).

(4) Diagnostic realized reference configuration interpretation of κ. $\kappa \geq 0$ is the diagnostic realized reference configuration monitoring rate (state–conditioned via the coarse–graining parameters such as $\varepsilon, m_{\mathrm{eff}}$). In the isolation limit $\kappa \to 0$, $\mathcal{L} \to \mathcal{L}^{\mathrm{H}}$ (coherent evolution), consistent with Proposition 15.52. For $\kappa > 0$, the dissipator implements pure dephasing/pinching in the record basis, consistent with Lemma 15.28, Proposition 15.33, and Theorem 15.35.

This proves the claim. □

Remark 15.42 (Deterministic pinching each slice vs. weak monitoring)**.** If one were to take literally the product $\Big(\Pi \circ \mathrm{Ad}_{e^{-iH_{A,\mathrm{eff}}t/n}} \Big)^n$ (pinching *every* slice), the Chernoff limit

collapses to the *Zeno* projector semigroup on the range of Π, with generator $\Pi\mathcal{L}^{\mathrm{H}}\Pi$ acting inside the block–diagonal algebra and instantaneous suppression of off–diagonal blocks. The weak–monitoring construction in Proposition 15.41 instead yields a finite dephasing rate κ and the unital GKLS generator $\mathcal{L}^{\mathrm{H}} + \kappa(\Pi - I)$, which is the physically relevant coarse–grained dynamics on the diagnostic realized reference configuration.

Proposition 15.43 (Σ–causal support in the continuum limit)**.** *If each stroboscopic step $\Phi_{\Delta t}$ respects Σ–causal support on the diagnostic realized reference configuration (Definition 5.26 and Corollary 5.145), then the limit family $\{\Lambda_t\}_{t\geq 0}$ also respects Σ–causal support.*

Proof. Notation and meaning of Σ–support. Let $\mathcal{A}_A$ be the diagnostic realized reference configuration algebra organized by the expectation–level chronogeometry $\bar{\Sigma}$ (Definition 5.26). A CPTP map Φ *respects Σ–causal support* if, in the Heisenberg picture,

$$\Phi^*(\mathcal{N}) \subseteq \mathcal{N}$$

for every local von Neumann subalgebra $\mathcal{N} \subseteq \mathcal{A}_A$ associated with a region Σ–spacelike to the input. Equivalently, there is no superluminal signalling (Corollary 5.145).

Step 1 (Closure under composition). If Φ and Ψ each respect Σ–support, then for any such $\mathcal{N}$,

$$(\Phi \circ \Psi)^*(\mathcal{N}) = \Psi^*\Big(\Phi^*(\mathcal{N})\Big) \subseteq \Psi^*(\mathcal{N}) \subseteq \mathcal{N}.$$

Hence $\Phi \circ \Psi$ also respects Σ–support. In particular, any finite product of stroboscopic steps $\Phi_{\Delta t}$ enjoys the property.

Step 2 (Closure under limits). Fix $t \geq 0$ and let $\{\Xi_\alpha\}$ be CPTP maps with $\Xi_\alpha^*(\mathcal{N}) \subseteq \mathcal{N}$ for all local $\mathcal{N}$, and $\Xi_\alpha \to \Xi$ in operator norm (hence strongly). Since each $\mathcal{N}$ is norm–closed, for any $X \in \mathcal{N}$,

$$\Xi^*(X) = \lim_\alpha \Xi_\alpha^*(X) \in \mathcal{N}.$$

Thus Ξ respects Σ–support.

Step 3 (Apply to the continuum limit). Under the coarse–grained diagnostic realized reference configuration dynamics (Assumption 15.39), the continuum family is obtained as a uniform product limit in operator norm (e.g. Chernoff/Trotter in Definition 15.37(c), Proposition 15.41):

$$\Lambda_t = \lim_{n\to\infty} \Big(\Phi_{t/n}\Big)^n \quad \text{uniformly for } t \text{ on compacts.}$$

By Step 1 each finite product $\Big(\Phi_{t/n}\Big)^n$ respects Σ–support; by Step 2, the limit Λ_t does as well. Since t was arbitrary, the entire family $\{\Lambda_t\}_{t\geq 0}$ respects Σ–causal support.

Remark (consistency with the GKLS generator). When the limit is generated by the unital GKLS operator of Proposition 15.41, both the Hamiltonian part $-i[H_{A,\mathrm{eff}}, \cdot]$ and the dissipator $\kappa(\mathbb{E}_A - \mathrm{id})$ are local with respect to Σ on the diagnostic realized reference configuration (the latter is a conditional expectation onto the record algebra), so Σ–support is preserved already at the generator level.

□

Remark 15.44 (Anisotropy and direction–dependent rates)**.** If $\mathcal{U}_{\Phi\Theta}$ is anisotropic (e.g. direction–selective scattering in the record basis), the monitoring rate decomposes as $\kappa = \sum_\alpha \kappa_\alpha$ with Lindblad set $\{L_{\alpha k} = \sqrt{\kappa_\alpha} P_{k,\alpha}\}$. Unitality and CP–divisibility persist; entropy monotonicity follows from Theorem 15.35.

Proposition 15.45 (Semigroup existence on the state–conditioned active band (Zeno product))**.** *Let* $\mathcal{H}_{\rm act}$ *be the* state–conditioned active band *determined by the reciprocal resolution limits (finite IR/UV window) of the endogenous IQR calculus (Proposition 4.436), hence* $\dim \mathcal{H}_{\rm act} = d < \infty$*. For a fixed step* $\Delta t > 0$ *define, on* $\mathcal{B}(\mathcal{H}_{\rm act})$*,*

$$\Phi_{\Delta t}(\rho) := \mathbb{E}_A\Big(U_A(\Delta t)\,\rho\,U_A(\Delta t)^\dagger\Big), \qquad U_A(\Delta t) := e^{-iH_{A,\mathrm{eff}}\Delta t},$$

and for $t \geq 0$ *the Trotterized products*

$$\Lambda_t^{(n)} \;:=\; \Big(\,\mathbb{E}_A \circ \mathrm{Ad}_{e^{-iH_{A,\mathrm{eff}}\,t/n}}\,\Big)^n, \qquad \mathrm{Ad}_U(\rho) := U\rho U^\dagger.$$

Then there exists a single subsequence $n_j \to \infty$ *(independent of* t*) such that, for all rational* $t \geq 0$ *and hence all* $t \geq 0$ *by continuity,* $\Lambda_t^{(n_j)}$ *converges in operator norm to a linear map* Λ_t *on* $\mathcal{B}(\mathcal{H}_{\rm act})$*. The family* $\{\Lambda_t\}_{t\geq 0}$ *consists of unital CPTP maps and satisfies the semigroup law*

$$\Lambda_{s+t} = \Lambda_s \circ \Lambda_t \qquad (s,t \geq 0).$$

Moreover, the product structure implies the Zeno pinching *relations*

$$\Lambda_t = \mathbb{E}_A \circ \Lambda_t = \Lambda_t \circ \mathbb{E}_A, \qquad \Lambda_0 = \mathbb{E}_A.$$

Restricted to the record (fixed–point) algebra $\mathcal{R}_A := \mathrm{Fix}(\mathbb{E}_A)$*, the family* $\{\Lambda_t|_{\mathcal{R}_A}\}_{t\geq 0}$ *is a unital CPTP semigroup with identity at* $t = 0$ *on* $\mathcal{R}_A$*.*

Proof. **(1) Finite dimensional setting and basic properties.** By Proposition 4.436 the active band is finite dimensional; hence $\mathcal{B}(\mathcal{H}_{\rm act}) \cong M_d(\mathbb{C})$. Each factor $\mathbb{E}_A$ is CPTP and unital (Definition 15.27 and Lemma 15.28); each $\mathrm{Ad}_{e^{-iH_{A,\mathrm{eff}}t/n}}$ is unitary conjugation (hence CPTP and unital). Therefore every $\Lambda_t^{(n)}$ is CPTP and unital.

(2) Precompactness via Choi representation. Let $J(\Phi)$ denote the Choi matrix of a linear map Φ on $M_d(\mathbb{C})$. The set $\mathcal{C} = \{J \geq 0 : \mathrm{Tr}_1[J] = \mathbb{I}_d\}$ is compact in finite dimension [358, 407, Ch. 2]. Thus, for each fixed $t \geq 0$, $\{J(\Lambda_t^{(n)})\}_n \subset \mathcal{C}$ has convergent subsequences; consequently $\{\Lambda_t^{(n)}\}_n$ has convergent subsequences in operator norm.

(3) A common subsequence and continuity in t**.** Choose a countable dense set of times (e.g. rationals in $[0,\infty)$) and apply a Cantor diagonal argument to extract a single subsequence n_j along which $J(\Lambda_t^{(n_j)})$ converges for all rational $t \geq 0$. Since CPTP maps are contractions in trace norm and depend continuously on t along the product nets considered here (bounded generators on a finite space), the limits extend uniquely to all real $t \geq 0$; denote them by Λ_t. Being norm limits of unital CPTP maps, each Λ_t is unital CPTP [358, Ch. 2].

(4) Semigroup property. For rational $s, t \geq 0$ (hence for all s, t by continuity), the cocycle of the products implies

$$\Lambda_{s+t}^{(n)} = \Big(\mathbb{E}_A \circ \mathrm{Ad}_{e^{-iH\,(s+t)/n}}\Big)^n = \Big(\mathbb{E}_A \circ \mathrm{Ad}_{e^{-iH\,s/n}}\Big)^n \circ \Big(\mathbb{E}_A \circ \mathrm{Ad}_{e^{-iH\,t/n}}\Big)^n + o_n(1),$$

where the $o_n(1)$ term vanishes in operator norm as $n \to \infty$ (standard Trotter splitting in finite dimension; cf. [457, 116, 623]). Passing to the common subsequence n_j and taking limits gives $\Lambda_{s+t} = \Lambda_s \circ \Lambda_t$.

(5) Zeno pinching structure. Each product can be written as

$$\Lambda_t^{(n)} = \underbrace{\mathbb{E}_A}_{\text{first factor}} \circ \Big(\mathrm{Ad}_{e^{-iHt/n}} \circ \mathbb{E}_A \Big)^{n-1} \circ \mathrm{Ad}_{e^{-iHt/n}} .$$

Hence $\Lambda_t^{(n)} = \mathbb{E}_A \circ \Lambda_t^{(n)} = \Lambda_t^{(n)} \circ \mathbb{E}_A$ for every n and t. Passing to the limit along n_j yields $\Lambda_t = \mathbb{E}_A \circ \Lambda_t = \Lambda_t \circ \mathbb{E}_A$ for all $t \geq 0$. At $t = 0$ we have $\Lambda_0^{(n)} = (\mathbb{E}_A)^n = \mathbb{E}_A$, hence $\Lambda_0 = \mathbb{E}_A$. It follows that the restriction $\Lambda_t|_{\mathcal{R}_A}$ to the fixed–point algebra $\mathcal{R}_A := \mathrm{Fix}(\mathbb{E}_A)$ satisfies $\Lambda_0|_{\mathcal{R}_A} = \mathrm{id}_{\mathcal{R}_A}$ and forms a unital CPTP semigroup on $\mathcal{R}_A$ (the *Zeno* semigroup; cf. Remark 15.42).

This proves the claim. □

Remark 15.46 (Endogenous origins of the Markov/active–band hypotheses)**.** The assumptions invoked for the continuum limit are supplied intrinsically by the IQR framework. (i) *Finite active band* arises from the reciprocal resolution limits and coherent regularization of the relational spectrum (Definition 4.576, Propositions 4.424 and 4.436, and eq. (4.278)), yielding the state–conditioned thresholds $\lambda_{\mathrm{IR}} \leq \lambda_n \leq \lambda_{\mathrm{cutoff}}$ and the finite state–conditioned active space $\mathcal{H}_{\mathrm{act}}$ (Definition 15.56). (ii) *Coherence–scaled Markov limit* follows from the separation of scales $\tau_{\mathrm{mem}} \ll \Delta t \ll \tau_{\mathrm{obs}}$ together with state–dependent spectral gapping (effective mass $m_{\mathrm{eff}} > 0$), as summarized in Assumption 15.39. These conditions ensure a unital, CP–divisible reduced dynamics consistent with $\bar{\Sigma}$–causal support (Definition 5.26 and Proposition 15.43).

Proposition 15.47 (Monotone entropy in the continuum limit)**.** *Adopt Assumption 15.39 and let $\{\Lambda_t\}_{t\geq 0}$ be the unital, CP–divisible reduced evolution on $\mathfrak{B}_A$ (the Good–clock parameter t being any admissible axis of $\vec{t}$ or, in the macroscopic timelike regime, t_{geom}; Definitions 5.95 and 15.21). For $\rho_A(t) := \Lambda_t(\rho_A(0))$, the von Neumann entropy*

$$S(\rho_A(t)) \;:=\; -\operatorname{Tr}\big[\rho_A(t) \log \rho_A(t)\big]$$

is nondecreasing in t. In particular, $S(\rho_A(t_2)) \geq S(\rho_A(t_1))$ for all $t_2 \geq t_1$, and therefore $\frac{d}{dt} S(\rho_A(t)) \geq 0$ for almost every $t \geq 0$.

Proof. By CP–divisibility (Definition 15.37), for any $t \geq s$ there exists a CPTP map $V_{t,s}$ with $\Lambda_t = V_{t,s} \circ \Lambda_s$. Since $\Lambda_s(\mathbb{I}_A) = \mathbb{I}_A$ and $\Lambda_t(\mathbb{I}_A) = \mathbb{I}_A$ (unitality; see Lemma 15.40 and Proposition 15.41), we have

$$V_{t,s}(\mathbb{I}_A) \;=\; V_{t,s}\big(\Lambda_s(\mathbb{I}_A)\big) \;=\; \Lambda_t(\mathbb{I}_A) \;=\; \mathbb{I}_A,$$

so each $V_{t,s}$ is also unital.

Let $d = \dim \mathcal{H}_{\mathrm{act}}$ and set $\sigma := \mathbb{I}_A/d$. Because $V_{t,s}$ is CPTP and unital, σ is a fixed point: $V_{t,s}(\sigma) = \sigma$. Umegaki relative entropy is monotone under CPTP maps, hence (e.g. [287, Sec. 11.6])

$$D\big(V_{t,s}(\rho) \,\|\, \sigma\big) \;\leq\; D(\rho\|\sigma), \qquad D(\rho\|\sigma) = \operatorname{Tr}[\rho(\log\rho - \log\sigma)].$$

Using $\log\sigma = -\log d\,\mathbb{I}$, this is equivalent to

$$S\Big(V_{t,s}(\rho)\Big) \;\geq\; S(\rho).$$

Apply this with $\rho = \rho_A(s) = \Lambda_s(\rho_A(0))$ to obtain $S(\rho_A(t)) \geq S(\rho_A(s))$ whenever $t \geq s$. Thus $t \mapsto S(\rho_A(t))$ is nondecreasing. On the finite–dimensional active band it is also bounded above by $\log d$, and every bounded monotone function on $[0,\infty)$ is differentiable almost everywhere; hence $\frac{d}{dt}S(\rho_A(t)) \geq 0$ for a.e. $t \geq 0$.

Optional time–homogeneous case. If, in addition, Λ_t is a (unital) GKLS semigroup $e^{t\mathcal{L}}$ as in Proposition 15.41, then $t \mapsto \rho_A(t)$ is norm–continuous and the same monotonicity follows directly from Theorem 15.35. □

Corollary 15.48 (Equality conditions)**.** *For the GKLS generator $\mathcal{L}[\rho] = -i[H_{A,\mathrm{eff}},\rho] + \kappa\Big(\mathbb{E}_A(\rho) - \rho\Big)$ (Proposition 15.41), the instantaneous entropy production rate $\dfrac{d}{dt}S(\rho_A(t))$ satisfies*

$$\frac{d}{dt}S(\rho_A(t)) \;=\; -\operatorname{Tr}\Big(\mathcal{L}[\rho_A(t)]\,\log\rho_A(t)\Big) \;\geq\; 0,$$

with equality at time t if and only if either $\kappa = 0$ (purely unitary dynamics) or $[\rho_A(t), P_k] = 0$ for all record projectors P_k (equivalently $\mathbb{E}_A(\rho_A(t)) = \rho_A(t)$; cf. Remark 15.32). In particular, for $\kappa > 0$ the rate vanishes at t iff $\rho_A(t)$ lies in the record algebra (the fixed–point algebra of $\mathbb{E}_A$).

Proof. Write $\rho \equiv \rho_A(t)$. Since $\operatorname{Tr}(\mathcal{L}[\rho]) = 0$, the entropy derivative is

$$\dot S(t) \;=\; -\operatorname{Tr}\Big(\mathcal{L}[\rho]\,\log\rho\Big) \;=\; -\operatorname{Tr}\Big(-i[H_{A,\mathrm{eff}},\rho]\,\log\rho\Big) \;-\; \kappa\,\operatorname{Tr}\Big((\mathbb{E}_A(\rho)-\rho)\,\log\rho\Big).$$

Unitary part vanishes. Using cyclicity of the trace,

$$-\operatorname{Tr}\Big(-i[H_{A,\mathrm{eff}},\rho]\,\log\rho\Big) = -i\,\operatorname{Tr}\Big(\rho\,[\log\rho,\,H_{A,\mathrm{eff}}]\Big) = 0,$$

because $[\log\rho,\rho] = 0$. Hence

$$\dot S(t) \;=\; \kappa\,\underbrace{\Big(\operatorname{Tr}\Big(\rho\log\rho\Big) - \operatorname{Tr}\Big(\mathbb{E}_A(\rho)\,\log\rho\Big)\Big)}_{=:\;\Xi(\rho)}.$$

Nonnegativity and equality for the dissipative part. Consider $f(\epsilon) := S\Big((1-\epsilon)\rho + \epsilon\,\mathbb{E}_A(\rho)\Big)$ for $\epsilon \in [0,1]$. By concavity of S and the pinching inequality $S(\mathbb{E}_A(\rho)) \geq S(\rho)$ (Lemma 15.30 and Proposition 15.31), the right derivative at 0 is nonnegative; differentiating,

$$f'(0^+) \;=\; -\operatorname{Tr}\Big((\mathbb{E}_A(\rho)-\rho)\,\log\rho\Big) \;=\; \Xi(\rho) \;\geq\; 0.$$

Moreover, equality $f'(0^+) = 0$ holds iff the pinching inequality is tight, i.e. iff $\mathbb{E}_A(\rho) = \rho$, equivalently $[\rho, P_k] = 0\ \forall k$ (Remark 15.32 and Lemma 15.28). Therefore,

$$\dot S(t) = \kappa\,\Xi(\rho) \;\geq\; 0, \qquad \dot S(t) = 0 \;\Longleftrightarrow\; \Big(\kappa = 0\Big) \;\text{ or }\; \Big(\mathbb{E}_A(\rho) = \rho\Big).$$

Summary. If $\kappa = 0$ the evolution is purely unitary and $\dot S(t) = 0$ for all ρ. If $\kappa > 0$, then $\dot S(t) = 0$ iff ρ lies in the fixed–point (record) algebra of $\mathbb{E}_A$, i.e. $[\rho, P_k] = 0$ for all k. □

Remark 15.49 (Persistence of zero production)**.** If $\kappa > 0$ and one requires $\dot{S}(t) = 0$ on an *interval*, it is necessary and sufficient that $\rho_A(t) \in \mathrm{Fix}(\mathbb{E}_A)$ for all t and that the unitary part preserves the record algebra, i.e. $[H_{A,\mathrm{eff}}, P_k] = 0$ for all k; otherwise the Hamiltonian produces off–diagonal coherences that the dissipator immediately dephases, yielding $\dot{S} > 0$ for $t' > t$.

Remarks.

(i) Revivals vs. coarse–grained irreversibility In closed, finite–dimensional AE composites, Loschmidt echoes/recurrences are generic [426, 427]. In our setting they are *suppressed* by the endogenous coarse–graining of Assumption 15.39: working at resolutions $\tau_{\mathrm{mem}} \ll \Delta t \ll \tau_{\mathrm{obs}}$ with state–conditioned gapping yields a unital CP–divisible evolution—either as a Zeno–type unital CPTP semigroup (Proposition 15.45) or, under weak monitoring, as a unital GKLS semigroup with generator of Proposition 15.41. For such evolutions, $t \mapsto S(\rho_A(t))$ is monotone (Proposition 15.47 and Theorem 15.35). Recurrences may reappear only if one resolves dynamics below the memory time or removes monitoring (e.g. $\kappa \to 0$ and $\mathbb{E}_A \to \mathrm{id}$), i.e. outside the benchmarked thermodynamic regime.

(ii) Non–unital noise and Spohn's H–theorem If the effective pointer/channel is *non–unital* (finite–temperature diagnostic realized reference configuration) and the GKLS generator $\mathcal{L}$ admits a (faithful) stationary state σ_A with $\mathcal{L}^\dagger(\sigma_A) = 0$, then the natural Lyapunov functional is the relative entropy $D(\rho_A(t)\|\sigma_A)$, which satisfies

$$\frac{d}{dt} D\big(\rho_A(t)\|\sigma_A\big) \;\le\; 0$$

(Spohn's H–theorem for quantum dynamical semigroups [624]; consistent with Lindblad–Uhlmann monotonicity under CPTP maps [617, 357]). When the stationary state is maximally mixed, $\sigma_A = \mathbb{I}/d$ (the *unital* limit realized by our pointer $\mathbb{E}_A$; Definition 15.27, Proposition 15.41, and Lemma 15.40), this reduces to monotonicity of the von Neumann entropy $S(\rho_A(t))$ (Proposition 15.47 and Theorem 15.35).

(iii) Chronogeometric time and clock gauge In macroscopic timelike regimes, choosing $t = t_{\mathrm{geom}}$ aligns the information–theoretic arrow with the chronogeometric proper–time arrow (Definitions 5.95 and 15.21). By Good–clock gauge invariance (Lemma 15.6), monotonicity (whether for S in the unital case or $D(\,\cdot\,\|\sigma_A)$ otherwise) is independent of the representative $C \in [C]$ used to parametrize the evolution.

(iv) Anisotropy and Σ–causal support If $\mathcal{U}_{\Phi\Theta}$ is anisotropic, the dissipative rates may decompose directionally (Remark 15.44) without affecting unitality/CP–divisibility. Σ–causal support is preserved in the continuum limit (Proposition 15.43), hence entropy monotonicity remains compatible with the $\bar{\Sigma}$–lightcone architecture (Definition 5.26).

noindent*Summary.* Stage 1 establishes *emergent time* from a timeless Holo–State via Page–Wootters conditioning with a Good–clock $C = (\hat{T}, U_t)$ (Theorem 13.168, Definition 13.158, and Proposition 13.172), fixes the *triadic* temporal structure and its macroscopic identification with proper time t_{geom} (Definitions 5.95 and 15.21), and guarantees clock–gauge

invariance of dynamical statements (Lemma 15.6). Stage 2 constructs, from the current–kernel interaction sourced by the Fourfold correlator $\hat{G}_\Lambda$ (Definitions 4.9 and 15.25), a transport–covariant *pointer channel* $\mathbb{E}_A$ on the record algebra (Definition 15.27 and Lemma 15.26). Each stroboscopic step $\Phi_{\Delta t} = \mathbb{E}_A \circ \mathrm{Ad}_{e^{-iH_{A,\mathrm{eff}}\Delta t}}$ is CPTP and unital (Proposition 15.29); pinching/conditional expectation cannot reduce entropy (Lemma 15.30 and Proposition 15.31; cf. [287, 358]), hence S is stepwise monotone (Proposition 15.33 and Corollary 15.34). Stage 3 invokes the *endogenous coherence–scaled Markov limit* (PQS width, state–conditioned bandlimiting, and spectral gapping) to obtain a unital, CP–divisible reduced dynamics $\{\Lambda_t\}$ parameterized by an admissible emergent time (or t_{geom}) (Assumption 15.39 and Definition 15.37). In the weak–monitoring realization one obtains the unital GKLS semigroup with generator of Proposition 15.41; in the Zeno realization one obtains a unital CPTP semigroup on the record algebra (Proposition 15.45). In either case this yields a continuous–time H–theorem $dS_A/dt \geq 0$ (Proposition 15.47 and Theorem 15.35), as used in Theorem 15.50. For background on GKLS semigroups and entropy production see [617, 441].

The Second Law as a Theorem of Relational Information Dynamics. With the three–stage construction in place, the Second Law is an *endogenous* statement of the IQR framework: a law of *irreversible entanglement spreading* along an admissible emergent–time axis. The temporal parameter is selected by Page–Wootters conditioning within the triadic structure (Stage 1; Proposition 13.172 and Definitions 5.95 and 15.21). Entropy production originates from einselection—the pointer conditional expectation on the record algebra (Stage 2; Definition 15.27, Lemma 15.30, and Proposition 15.31), mathematically a pinching that majorizes spectra (cf. [287, 358]). Practical irreversibility arises once dynamics is coarse–grained to a *unital, CP–divisible* evolution justified by PQS width, state–conditioned spectral bandlimiting, and spectral gapping (Stage 3; Assumption 15.39, Definition 15.56, and Propositions 15.41, 15.45 and 15.47). In concise terms: *entropy is the price of local perspective in a globally entangled universe* (Theorem 13.168; cf. decoherence narratives [170, 441]).

Unification of Time, Measurement, and Entropy. The derivation identifies three facets of one relational process. (i) The *arrow of time* is the monotone direction of an emergent temporal axis (triadic, or macroscopically t_{geom}) along which reduced dynamics unfolds (Definitions 5.95 and 15.21 and Lemma 15.6). (ii) *Measurement* is einselection by the transport–covariant pointer channel $\mathbb{E}_A$ (Definition 15.27 and Proposition 15.29), induced by current–kernel interactions encoded in $\hat{G}_\Lambda$ (Definitions 4.9 and 15.25). (iii) The *Second Law* is the information–theoretic residue of this monitoring: under the unital, CP–divisible reduced dynamics of Definition 15.37, a subsystem's von Neumann entropy cannot decrease (Proposition 15.47 and Theorem 15.35). Global timelessness and unitarity are never violated (Theorem 13.168); the perceived irreversibility is precisely the delocalization of phase information into inaccessible correlations captured by the Fourfold correlator (cf. [170, 441]).

> **Second Law (relational form).** Along any Good–clock axis t, the reduced state $\rho_A(t)$ follows a unital flow that is *divisible into intermediate completely positive, trace-preserving quantum channels* (CP-divisible), driven by the current–kernel

coupling of the Fourfold correlator $\hat{G}_\Lambda$ and the pointer channel $\mathbb{E}_A$:

$$\dot{\rho}_A(t) \;=\; -\,i\,[H_{A,\mathrm{eff}}, \rho_A(t)] \;+\; \kappa\Big(\mathbb{E}_A(\rho_A(t)) - \rho_A(t)\Big),$$

$$\underbrace{\mathbb{E}_A(\rho) = \sum_k P_k\,\rho\,P_k}_{\text{einselection}}.$$

$$\underbrace{\forall\, t \geq r \geq s\ \exists\ V_{t,r}\ \mathrm{CPTP}:\ \ \Phi_{t\leftarrow s} = V_{t,r}\circ\Phi_{r\leftarrow s},\quad \Phi_{t\leftarrow s}(\mathbb{I}_A) = \mathbb{I}_A}_{\text{CP-divisible, unital}}$$

$$\underbrace{P_j P_k = \delta_{jk} P_k, \qquad \sum_k P_k = \mathbb{I}_A,}_{\text{record projectors}}$$

$$\underbrace{\mathbb{E}_A^2 = \mathbb{E}_A}_{\text{pinching (conditional expectation)}},$$

$$\underbrace{\mathcal{R}_A = \{X \in \mathcal{B}(\mathcal{H}_A) : [X, P_k] = 0\ \forall k\} = \mathrm{Fix}(\mathbb{E}_A)}_{\text{record algebra and fixed points}}$$

With probabilities/expectations taken in the emergent inner product (from $\hat{G}_\Lambda$), the von Neumann entropy of A cannot decrease:

$$t_2 \geq t_1 \;\Rightarrow\; S_A(t_2) \;\geq\; S_A(t_1),$$

$$\frac{d}{dt} S_A(t) = \kappa\ \mathrm{Tr}\Big[\rho_A(t)\log\rho_A(t) - \mathbb{E}_A(\rho_A(t))\log\rho_A(t)\Big] \;\geq\; 0.$$

All increase is due to environment-induced superselection (the conditional expectation $\mathbb{E}_A$ onto $\mathcal{R}_A$); equality holds iff

$$[\rho_A(t), P_k] = 0\ \forall k \quad \text{and} \quad \mathbb{E}_A(\rho_A(t)) = \rho_A(t) \ \text{ or } \ \kappa = 0,$$

i.e. the state already lies in the record algebra and the instantaneous motion on that commutant is unitary. Causality is respected: the flow preserves $\bar{\Sigma}$–lightcone support and admits no-law-signalling.

Theorem 15.50 (Second Law of Thermodynamics — relational, clock–gauge invariant form)**.** *Work on a realized thermodynamic diagnostic realized reference configuration* $\mathfrak{B}_{[\chi]}^{\langle \mathrm{DRRC}\rangle}(\mathfrak{P})$ *(Definition 15.1) with a Good–clock* $C = (\hat{T}, U_t)$ *aligned to* $\bar{\Sigma}$ *(Definitions 5.26 and 13.158). Let the universe be in the timeless pure Holo–State* $|\overset{\circ}{\boldsymbol{\Psi}}\rangle \in \mathcal{H}_{\mathrm{em}}$ *(Definition 4.287) satisfying the Universal Quantum Constraint*

$$\hat{H}_{\mathrm{univ}}\,|\overset{\circ}{\boldsymbol{\Psi}}\rangle = 0 \quad (\textit{Theorem 13.168; cf. [335]}).$$

Assume the environment $\mathfrak{B}_E$ *contains a clock–bearing degree of freedom enabling Page–Wootters conditioning (Proposition 13.172) and that the reduced dynamics on* $\mathfrak{B}_A$ *is* admissible (CP–divisible) *in the sense of Definition 15.37, obtained via the endogenous Markov limit of Assumption 15.39 (PQS width, state–conditioned bandlimiting, spectral gapping) so that, on the diagnostic realized reference configuration, it is* unital.

(i) Monotone entropy

For any admissible emergent time parameter t (an axis of the triadic time or, macroscopically, t_{geom}; Definitions 5.95 and 15.21), the von Neumann entropy

$$\boxed{S_A(t) \;:=\; -\operatorname{Tr}_{\mathfrak{B}_A}\big[\rho_A(t)\log\rho_A(t)\big]} \tag{15.13}$$

is nondecreasing:

$$t_2 \geq t_1 \;\Rightarrow\; S_A(t_2) \;\geq\; S_A(t_1),$$

hence $\frac{d}{dt}S_A(t) \geq 0$ for almost every t (Proposition 15.47 and Theorem 15.35; cf. [617, 441]).

(ii) Mechanism (current–kernel + einselection)

Entropy production is generated by the interaction Hamiltonian $\hat{H}_{AE}$ constructed from the current–kernel coupling $\mathcal{U}_{\Phi\Theta}$ of the Fourfold correlator $\hat{G}_\Lambda$ (Definitions 4.9 and 15.25 and Lemma 15.26), together with einselection by the transport–covariant pointer conditional expectation on the record algebra,

$$\mathbb{E}_A(\rho) = \sum_k P_k \rho P_k$$

(Definition 15.27). Each stroboscopic step $\Phi_{\Delta t} = \mathbb{E}_A \circ \mathrm{Ad}_{e^{-iH_{A,\mathrm{eff}}\Delta t}}$ is CPTP and unital (Proposition 15.29) and cannot reduce entropy (Lemma 15.30 and Proposition 15.31; cf. [287, 358]).

(iii) Practical irreversibility

Endogenous coarse–graining (finite memory, IR control) yields a unital, CP–divisible semigroup $\{\Lambda_t\}$ with GKLS generator obtained as a Chernoff product limit (Assumption 15.39 and Propositions 15.41 and 15.45); the apparent arrow is the delocalization of phase information into inaccessible AE correlations (decoherence) rather than a violation of global unitarity (Theorem 13.168; cf. [170, 282]). Σ–causal support is preserved (Proposition 15.43).

(iv) Differential H–theorem and equality conditions

Consequently,

$$\boxed{\frac{d}{dt}S_A(t) \;\geq\; 0,} \tag{15.14}$$

with equality iff the state already lies in the record algebra and the instantaneous evolution on that commutant is purely unitary—equivalently,

$$[\rho_A(t), P_k] = 0 \;\forall k$$

and either $\kappa = 0$ or $\mathbb{E}_A(\rho_A(t)) = \rho_A(t)$ (Corollary 15.48). The ontological limiting case is the globally coherent PQS, for which $S_A \equiv 0$ and no einselection occurs (Theorem 13.168).

All statements are invariant in the Good–clock gauge class $[C]$ (Lemma 15.6) and, in macroscopic timelike regimes, align with the proper–time arrow t_{geom} (Definition 5.95).

The Second Law as the Narrative of Cosmic Genesis. The theorem Theorem 15.50 fulfils a long–standing aim: to recover the thermodynamic arrow from a deeper, time–symmetric quantum reality driven by a timeless universal constraint and emergent conditioning [335, 610, 197]. In the IQR framework, the Second Law is not a classical tendency toward molecular chaos [436] but a precise law of *quantum–relational information dynamics*: along an admissible emergent–time axis (triadic or, macroscopically, t_{geom}; Definitions 5.95 and 15.21), the reduced entropy $S_A(t) = -\operatorname{Tr}[\rho_A(t)\log\rho_A(t)]$ (15.13) is monotone because each Good–clock step combines unitary transport with einselection by the *pointer conditional expectation* $\mathbb{E}_A$ (Definition 15.27, Propositions 15.29 and 15.31, and Lemma 15.30; cf. [619, 625, 287, 358]). The microphysical driver is the current–kernel coupling $\mathcal{U}_{\Phi\Theta}$ built from the Fourfold Inter–Modal Correlator $\hat{G}_\Lambda$ (Definitions 4.9 and 15.25), which mediates relational exchange between the Gaiamiti and Tantrakyma sectors (the "Apeiron"/"Kymon" language of the treatise); its action disperses phase information into AE correlations—*decoherence*—so that, for observers restricted to $\mathcal{A}_A$, quantum information becomes more widely shared across the cosmic network [170, 282]. Thus the "past" corresponds to lower subsystem entanglement and the "future" to higher entanglement [626, 335]. Endogenous regularisation (finite PQS width, state–conditioned bandlimiting, spectral smoothness and gapping) furnishes a unital, CP–divisible semigroup in the continuum limit (Assumption 15.39, Definition 15.37, and Propositions 15.41, 15.45 and 15.47), completing a self–consistent narrative grounded in relational correlators [3, 197].

The Unification of Time, Entropy, and Creation. *First*, **time and entropy are one process**: the emergent temporal flow (Page–Wootters conditioning with a Good–clock; Proposition 13.172, Definition 13.158, and Lemma 15.6) and the monotone increase of $S_A(t)$ (15.14) are two descriptions of the same irreversible spreading of accessible information under unitary+einselection dynamics [335, 610, 619, 625, 170]. *Second*, **thermodynamics is cosmology**: anchoring the equality case of the H–theorem to the maximally coherent PQS (no records, no einselection) identifies zero entropy production as the limiting, globally coherent state (Corollary 15.48 and Theorem 13.168); coarse–graining away from this limit renders entropy growth the primary driver of macroscopic cosmic evolution [627, 157]. *Third*, **physics is information dynamics**: classical "heat" and "work" are the macroscopic bookkeeping of relational information flows (First Law; Theorem 15.18 and Definition 15.14), while *entropy production* is the information–theoretic residue of measurement–like monitoring by the environment—formalised by $\mathbb{E}_A$—and constrained by foundational limits such as Landauer's principle and Shannon's theory [137, 621, 622, 170, 282, 48].

This triune picture realises the *Fourfold Dialectic* (Section 1.2.7): the PQS as **Thesis** (undifferentiated coherence), unitary entangling evolution as **Antithesis** (distinction and relation), and the macroscopic universe with its thermodynamic arrow as **Synthesis**. Accordingly, the Second Law functions as the engine of dialectical unfolding: it guarantees that the cosmos will not remain in silent perfection but will ceaselessly explore the complexity latent in the Quantum Holo–State—*relational becoming* rather than mere decay [48, 610, 40].

15.1.5 The Third Law of Thermodynamics: Quantum Coherence in the Limit of Isolation

Statement and setting. We work on a realized thermodynamic diagnostic realized reference configuration $\mathfrak{B}_{[\chi]}^{\langle \mathrm{DRRC}\rangle}(\mathfrak{P})$ with Good–clock $C = (\hat{T}, U_t)$ aligned to $\bar{\Sigma}$ (Definitions 5.26, 13.158 and 15.1). In the IQR formulation, the **Third Law** is not a statement about crystalline order at $T = 0$, but a theorem about *quantum coherence under informational isolation*: as the relational coupling between A and its environment E is driven to zero, the reduced dynamics on A becomes unitary, entropy production ceases, and coherences (with respect to any fixed record algebra) are asymptotically conserved (Propositions 15.41 and 15.47 and Corollary 15.48; cf. resource–theoretic viewpoints [628, 629, 630]). Concretely, the interaction Hamiltonian $\hat{H}_{AE}$ and the monitoring rate κ (from the pointer channel $\mathbb{E}_A$; Definition 15.27) parameterize an *isolation protocol*:

$$\mathcal{L}_g[\rho] \;=\; -\,i\,[H_{A,\mathrm{eff}}, \rho] \;+\; \kappa(g)\,\Big(\mathbb{E}_A(\rho) - \rho\Big), \qquad g := \|\hat{H}_{AE}\|, \quad \kappa(g) \xrightarrow[g\to 0]{} 0, \tag{15.15}$$

so that $\Lambda_t^{(g)} = e^{t\mathcal{L}_g}$ tends to the unitary group $U_t(\cdot)U_t^\dagger$ as $g \to 0$. The *Second Law* (Theorem 15.50) provides the one–sided inequality $dS_A/dt \geq 0$ under unital CP–divisible dynamics; the *Third Law* identifies the boundary regime where the production rate vanishes and coherence is restored.

Narrative (from entropy production to coherence conservation). Stage 2 showed that entropy increases stepwise due to einselection by the pointer channel $\mathbb{E}_A$ (Definition 15.27, Lemma 15.30, and Proposition 15.31), and Stage 3 upgraded this to a continuous H–theorem under the endogenous, unital CP–divisible semigroup (Assumption 15.39, Definition 15.37, and Propositions 15.41 and 15.47). The *isolation limit* (15.15) suppresses both sources of irreversibility—current–kernel transfer (through $\hat{H}_{AE}$, Definition 15.25) and einselection (through κ)—so that

$$\lim_{g\to 0} \frac{d}{dt} S_A^{(g)}(t) = 0, \qquad \lim_{g\to 0} \Lambda_t^{(g)}(\rho) = U_t \rho\, U_t^\dagger,$$

i.e. entropy becomes *constant* and coherences are preserved in any fixed record basis (Corollary 15.48; cf. [282, 626]). Under admissible preparation on the diagnostic realized reference configuration (typed projector $\Pi^{[\chi]}$ and certified plateau $\mathfrak{P}$; Proposition 9.138 and Definition 9.58), protocols that factorize AE up to $o(1)$ in g drive ρ_A to purity in trace norm (resource–theoretic cooling [629, 630]; see Proposition 15.52).

Two IQR analogues of Nernst/Planck. (i) *Heat–theorem form (vanishing production).* The entropy–production rate obeys the small–coupling bound

$$\frac{d}{dt} S_A^{(g)}(t) \;=\; \kappa(g)\, \sigma_A^{(g)}(t) \;\xrightarrow[g\to 0]{}\; 0,$$

with $\sigma_A^{(g)}(t) \geq 0$ a diagnostic realized reference configuration–dependent functional bounded uniformly on compact time intervals (Lemma 15.51 and Proposition 15.41). (ii) *Unattainability form (finite–time control).* Exact isolation (the analogue of $T = 0$) is *unattainable*

in finite emergent time under finite control resources: any diagnostic realized reference configuration–admissible protocol keeping $\|\hat{H}_{AE}\| \le g_{\max}$ and $\kappa \le \kappa_{\max}$ requires $t \to \infty$ to reach a perfectly pure ρ_A (cf. [631, 632, 633]; see Section 15.1.6).

Cross–links to the earlier constructions. The isolation path preserves Σ–causal support (Proposition 15.43) and is clock–gauge invariant (Lemma 15.6). It is implemented in the same diagnostic realized reference configuration–wise semantics used for the First and Second Laws (Theorems 15.18 and 15.50), with the pointer/channel structure $\mathbb{E}_A$ (Definition 15.27) and the current–kernel interaction from the Fourfold correlator $\hat{G}_\Lambda$ (Definitions 4.9 and 15.25). The resulting third–law statements are tightened below as *small–g entropy bound* (Lemma 15.51) and *asymptotic coherence theorem* (Proposition 15.52).

Literature context. Our coherence–centric third law mirrors resource–theoretic and decoherence perspectives [628, 629, 630, 282, 626], and the unattainability principle aligns with quantum thermodynamic analyses of finite–time cooling and control constraints [631, 632, 633].

Isolation parameter and reduced dynamics. Let $g \ge 0$ quantify the effective interaction strength mediated by the current–kernel coupling $\mathcal{U}_{\Phi\Theta}$ at the A–E interface (Definition 15.25), after endogenous coarse–graining (Assumption 15.39). For each fixed g, the state–conditioned active band $\mathcal{H}_{\mathrm{act}}$ (Definition 15.56) yields a unital CPTP semigroup $\{\Lambda_t^{(g)}\}_{t\ge 0}$ on $\mathfrak{B}_A$ (Propositions 15.41 and 15.45) with GKLS generator

$$\mathcal{L}_g[\rho] \;=\; -\,i\,[H_{A,\mathrm{eff}},\rho] \;+\; \sum_\alpha \gamma_\alpha(g)\Big(L_\alpha \rho L_\alpha^\dagger - \tfrac{1}{2}\{L_\alpha^\dagger L_\alpha,\,\rho\}\Big)\,, \tag{15.16}$$

where $\gamma_\alpha(g) \ge 0$ are the monitoring/relaxation rates induced by $\mathcal{U}_{\Phi\Theta}$ and, by the endogenous finiteness and state–dependent gapping of Assumption 15.39, satisfy

$$\Gamma(g) := \sum_\alpha \gamma_\alpha(g) \;\xrightarrow[g\to 0]{}\; 0.$$

(Anisotropic couplings are accommodated by grouping α into directional channels; cf. Remark 15.44.) All statements below are Good–clock–gauge invariant (Lemma 15.6) and preserve Σ–causal support (Proposition 15.43).

Lemma 15.51 (Small–coupling entropy–production bound)**.** *Fix $t > 0$ and $d = \dim \mathcal{H}_{\mathrm{act}} < \infty$. Let the reduced dynamics at coupling g have time–local form $\dot\rho_A^{(g)}(s) = -\,i[H_{A,\mathrm{eff}}(s), \rho_A^{(g)}(s)] + \mathcal{M}_s^{(g)}[\rho_A^{(g)}(s)]$ (Lemma 15.23), and set*

$$\Gamma(g) \;:=\; \sup_{0\le s\le t} \big\|\mathcal{M}_s^{(g)}\big\|_{1\to 1}, \qquad \textit{with}\;\; \Gamma(g) \xrightarrow[g\to 0]{} 0.$$

Then for every initial state $\rho_A(0)$,

$$\Big|\, S\big(\Lambda_t^{(g)}\rho_A(0)\big) - S\big(\rho_A(0)\big)\,\Big| \;\le\; \varepsilon\,\log(d-1) \;+\; H_2(\varepsilon), \qquad \varepsilon := \frac{\Gamma(g)\,t}{2},$$

where $H_2(x) = -x\log x - (1-x)\log(1-x)$ is the binary entropy (nats). In particular, as $g \to 0$,

$$\Big|\, S\big(\Lambda_t^{(g)}\rho_A(0)\big) - S\big(\rho_A(0)\big)\,\Big| \;=\; O\Big(\Gamma(g)\,t\,\log\tfrac{1}{\Gamma(g)\,t}\Big).$$

A convenient coarse bound is

$$\Big|\, S\big(\Lambda_t^{(g)}\rho_A(0)\big) - S\big(\rho_A(0)\big)\,\Big| \;\le\; \Gamma(g)\,t\,\log d \;+\; H_2\Big(\frac{\Gamma(g)\,t}{2}\Big).$$

Proof. Interaction picture and trace–distance control. Let U_s solve $\dot U_s = -iH_{A,\mathrm{eff}}(s)U_s$, and write $\tilde\rho^{(g)}(s) := U_s^\dagger \rho_A^{(g)}(s) U_s$. Then $\dot{\tilde\rho}^{(g)}(s) = \tilde{\mathcal M}_s^{(g)}[\tilde\rho^{(g)}(s)]$ with $\|\tilde{\mathcal M}_s^{(g)}\|_{1\to1} = \|\mathcal M_s^{(g)}\|_{1\to1} \le \Gamma(g)$, since unitary conjugation preserves both the trace norm and the induced $1\to1$ norm. Thus $\|\dot{\tilde\rho}^{(g)}(s)\|_1 \le \Gamma(g)$, and integrating yields

$$\big\|\tilde\rho^{(g)}(t) - \tilde\rho^{(g)}(0)\big\|_1 \;\le\; \Gamma(g)\,t.$$

Entropy is unitary–invariant, so $S(\rho_A^{(g)}(\cdot)) = S(\tilde\rho^{(g)}(\cdot))$.

Audenaert–Fannes bound. Let $T := \frac12\|\tilde\rho^{(g)}(t) - \tilde\rho^{(g)}(0)\|_1 \le \Gamma(g)t/2$. The Audenaert–Fannes continuity bound on a d-dimensional space gives

$$\big|S(\tilde\rho^{(g)}(t)) - S(\tilde\rho^{(g)}(0))\big| \;\le\; T\,\log(d-1) \;+\; H_2(T),$$

which with $T \le \Gamma(g)t/2$ is exactly the displayed inequality with $\varepsilon = \Gamma(g)t/2$. Since $H_2(x) = x\log\frac{e}{x} + O(x^2)$ as $x\downarrow 0$, the small–coupling asymptotic $O(\Gamma(g)\,t\,\log\frac{1}{\Gamma(g)t})$ follows. The looser but sometimes handy coarse bound $\le \Gamma(g)t\log d + H_2(\Gamma(g)t/2)$ is obtained by using $T\le\Gamma(g)t/2$ and $\log(d-1)\le\log d$. □

Proposition 15.52 (Third Law — coherence in the isolation limit)**.** *For each fixed $t<\infty$ on the state–conditioned active band $\mathcal H_{\mathrm{act}}$ (finite $d := \dim\mathcal H_{\mathrm{act}}$),*

$$\lim_{g\to0} S\big(\Lambda_t^{(g)}\rho_A(0)\big) \;=\; S\big(\rho_A(0)\big), \qquad \lim_{g\to0}\Lambda_t^{(g)} \;=\; \mathrm{Ad}_{e^{-iH_{A,\mathrm{eff}}t}}$$

in operator norm on $\mathcal B(\mathcal H_{\mathrm{act}})$. Consequently, in any fixed record (pointer) basis $\{P_k\}$, there is no irreversible suppression *of off–diagonal terms in the limit $g\to0$: coherences evolve unitarily (they may oscillate, but do not decay), and the information–theoretic arrow $\dot S\ge0$* flattens *to $\dot S = 0$.*

Proof. Entropy continuity. Let the time–local reduced generator at coupling g be $\dot\rho_A^{(g)}(s) = -i[H_{A,\mathrm{eff}}(s),\rho_A^{(g)}(s)] + \mathcal M_s^{(g)}[\rho_A^{(g)}(s)]$ (Lemma 15.23). Set $\Gamma(g) := \sup_{0\le s\le t}\|\mathcal M_s^{(g)}\|_{1\to1}$, with $\Gamma(g)\to0$ as $g\to0$. By the small–coupling continuity bound (Lemma 15.51),

$$\big|\,S(\Lambda_t^{(g)}\rho_A(0)) - S(\rho_A(0))\,\big| \;\le\; \varepsilon\,\log(d-1) + H_2(\varepsilon), \qquad \varepsilon := \tfrac12\,\Gamma(g)\,t,$$

so the RHS $\to0$ as $g\to0$. Hence $S(\Lambda_t^{(g)}\rho_A(0))\to S(\rho_A(0))$.

Dynamical limit (operator norm). Write the generator as $\mathcal L_s^{(g)} = \mathcal L_s^{\mathrm H} + \mathcal M_s^{(g)}$ with $\mathcal L_s^{\mathrm H}[\cdot] = -i[H_{A,\mathrm{eff}}(s),\cdot]$. Let $\mathcal U_{t,s}$ and $\mathcal U_{t,s}^{(g)}$ be the (bounded) propagators solving $\partial_t\mathcal U_{t,s} = \mathcal L_t^{\mathrm H}\mathcal U_{t,s}$, $\partial_t\mathcal U_{t,s}^{(g)} = \mathcal L_t^{(g)}\mathcal U_{t,s}^{(g)}$, with $\mathcal U_{s,s} = \mathcal U_{s,s}^{(g)} = I$. Duhamel's formula for non–autonomous bounded generators gives

$$\mathcal U_{t,0}^{(g)} - \mathcal U_{t,0} = \int_0^t \mathcal U_{t,s}^{(g)}\,\mathcal M_s^{(g)}\,\mathcal U_{s,0}\,ds.$$

Taking the induced $1\to1$ norm and using $\|\mathcal{U}_{t,s}^{(g)}\|_{1\to1}, \|\mathcal{U}_{s,0}\|_{1\to1} \le C_t$ (some finite C_t in finite dimension),

$$\begin{aligned}\left\|\mathcal{U}_{t,0}^{(g)} - \mathcal{U}_{t,0}\right\|_{1\to1} &\le \int_0^t C_t \left\|\mathcal{M}_s^{(g)}\right\|_{1\to1} C_t \,\mathrm{d}s \\ &\le C_t^2\,\Gamma(g)\,t \underset{g\downarrow0}{\longrightarrow} 0.\end{aligned}$$

In the time-homogeneous (semigroup) case, $\Lambda_t^{(g)} = \exp\!\big(t(\mathcal{L}^{\mathrm{H}} + \mathcal{M}^{(g)})\big)$, one may alternatively use the bounded-perturbation/Duhamel estimate

$\|e^{t(\mathcal{L}^{\mathrm{H}}+\mathcal{M}^{(g)})} - e^{t\mathcal{L}^{\mathrm{H}}}\|_{1\to1} \le t\, e^{t(\|\mathcal{L}^{\mathrm{H}}\|+\|\mathcal{M}^{(g)}\|)}\|\mathcal{M}^{(g)}\| \to 0$. Since all operator norms are equivalent on $\mathcal{B}(\mathcal{H}_{\mathrm{act}})$, convergence holds in operator norm. Identifying $\mathcal{U}_{t,0} = \mathrm{Ad}_{e^{-iH_{A,\mathrm{eff}}t}}$ completes the dynamical limit.

Coherence remark. In the limit $g \to 0$ we have purely unitary evolution on A; thus no pinching/einselection acts and there is no irreversible damping of off–diagonal elements in any fixed record basis $\{P_k\}$. Coherences may rotate but are not *suppressed*, and since $S(\rho_A(t))$ is unitary–invariant, $\dot S = 0$.

This proves the proposition. □

Corollary 15.53 (Residual entropy for degenerate ground spaces)**.** *Let the isolated Hamiltonian $H_{A,\mathrm{eff}}$ have a $\mathfrak{g}$-fold degenerate ground space with spectral resolution $H_{A,\mathrm{eff}} = \sum_{r\ge0} E_r P_r$ and $\dim P_0 = \mathfrak{g}$. Fix $t < \infty$ and the state–conditioned active band (finite d). Then:*

(i) Strict isolation, fixed support in the ground block.
If the initial state is supported on the ground manifold, $\rho_0 = P_0\rho_0P_0$ with $\mathrm{Tr}\,\rho_0 = 1$, then in the isolation limit $g \to 0$ the reduced evolution on A is unitary (Proposition 15.52) and the entropy is constant:

$$\lim_{g\to0} S\!\left(\Lambda_t^{(g)}\rho_0\right) = S(\rho_0), \qquad 0 \;\le\; S(\rho_0) \;\le\; \log\mathfrak{g}.$$

The upper bound is attained precisely by the microcanonical state $\omega_0 := \frac{1}{\mathfrak{g}}P_0$, for which $S(\omega_0) = \log\mathfrak{g}$.

(ii) Attainability of the maximal residual entropy.
If prior to (or during) preparation one employs admissible unital mixing that preserves energy and acts only inside P_0 (e.g. an intra-block random-unitary twirl), then one can reach ω_0 without changing the energy, and the residual entropy equals

$$S_0 = \log\mathfrak{g} \quad (k_B = 1).$$

By the small–coupling continuity bound (Lemma 15.51), any such unital intra-block mixing achieved at small but nonzero coupling persists in the limit: for fixed t, $S(\Lambda_t^{(g)}\rho_A(0)) \to S(\rho_A(0))$ as $g \to 0$.

(iii) Nondegenerate case.
If the ground state is unique ($\mathfrak{g} = 1$), then $P_0 = |\psi_0\rangle\langle\psi_0|$ is pure and $S_0 = 0$, in agreement with the standard thermodynamic statement (cf. [119, 634, 626, 282]).

Proof. (**a**) By Proposition 15.52, for each fixed $t < \infty$, $\Lambda_t^{(g)} \to \mathrm{Ad}_{e^{-iH_{A,\mathrm{eff}}t}}$ in operator norm as $g \to 0$, hence evolution is unitary and preserves the support P_0. Entropy is therefore constant and bounded by $\log \mathfrak{g}$, with equality at $\omega_0 = \frac{1}{\mathfrak{g}} P_0$ by strict concavity/unitary invariance.

(**b**) Any unital CPTP map acting only inside P_0 and preserving energy (e.g. Haar twirl on P_0) sends $\rho \mapsto (\mathrm{Tr}(P_0\rho)/\mathfrak{g})P_0$, achieving ω_0 at fixed energy. If this is implemented at small coupling, Lemma 15.51 ensures continuity of S as $g \to 0$ for fixed t.

(**c**) If $\mathfrak{g} = 1$, then P_0 is one-dimensional and the unique ground state is pure, so $S_0 = 0$.

This completes the proof. □

Definition 15.54 (Thermodynamic domain of applicability: two levels)**.** (**Level P: pre-physical**) From the PQS core up to (but not including) the Planck scale, distances/times are null at the core and thermodynamic diagnostic realized reference configuration are *not* defined; thermodynamic predicates are inapplicable (cf. Section 9.1, Theorem 13.168).
(**Level R: realized**) From the Planck scale upward (to cosmological scales), thermodynamic diagnostic realized reference configuration $\mathfrak{B}_{[\chi]}^{\langle \mathrm{DRRC} \rangle}(\mathfrak{P})$ and Good-clock dynamics are admissible (Definitions 13.158 and 15.1).

Definition 15.55 (Planck floor for diagnostic realized reference configuration resolution)**.** On any diagnostic realized reference configuration, the stroboscopic step and readout window obey

$$\Delta t \;\geq\; \tau_{\mathrm{Pl}}^{(\mathrm{thermo})} > 0, \qquad \Delta \ell \;\geq\; \ell_{\mathrm{Pl}}^{(\mathrm{thermo})} > 0,$$

where these positive constants are inherited from the state-conditioned spectral thresholds and coherence width ($\tau_{\mathrm{mem}} = \Lambda_{\mathrm{cutoff}}^{-1}$, $\ell_{\mathrm{PQS}} = \Lambda_{\mathrm{eff}}^{-1}$; Definition 4.576, eq. (4.278), and Section 4.15.3) together with the Chapter Chapter 9 restriction that thermodynamic operations cannot descend into the pre-physical domain. In particular, the coarse-grained regime requires $\tau_{\mathrm{mem}} \ll \Delta t$ with $\tau_{\mathrm{mem}} \geq \tau_{\mathrm{Pl}}^{(\mathrm{thermo})}$ (Assumption 15.39).

Definition 15.56 (State–conditioned active band)**.** Given tolerance $\varepsilon \in (0,1)$ and Holo–State $|\mathring{\mathbf{\Psi}}\rangle$, the *active band* $\mathcal{H}_{\mathrm{act}}$ is the span of spectral modes with frequencies $\{\lambda_n\}$ satisfying

$$\Lambda_{\mathrm{IR}}(x;\varepsilon) \;\leq\; \lambda_n \;\leq\; \Lambda_{\mathrm{cutoff}}(x;\varepsilon),$$

with thresholds determined endogenously by modal weights and spectral smoothness (Definitions 4.392 and 4.576, Axiom 4.415, and eq. (4.278)). In the coarse-grained regime of Assumption 15.39, $\mathcal{H}_{\mathrm{act}}$ is finite-dimensional.

Proposition 15.57 (Product–limit GKLS generator (weak–monitoring scaling))**.** *Under Assumption 15.39 on the active band $\mathcal{H}_{\mathrm{act}}$, set for $\Delta > 0$*

$$V_\Delta^{(\kappa)} \;:=\; \Big((1-\kappa\Delta)\,\mathrm{id} + \kappa\Delta\,\mathbb{E}_A\Big) \circ \mathrm{Ad}_{e^{-iH_{A,\mathrm{eff}}\Delta}}, \qquad \mathrm{Ad}_U(\rho) := U\rho U^\dagger,$$

with $\mathbb{E}_A$ as in Definition 15.27 and $\kappa \geq 0$ the (diagnostic realized reference configuration–fixed) monitoring rate. Then, for each $t \geq 0$,

$$\Lambda_t^{(n)} \;:=\; \Big(V_{t/n}^{(\kappa)}\Big)^n \;\xrightarrow[n\to\infty]{\|\cdot\|}\; \Lambda_t \;=\; e^{t\mathcal{L}},$$

in operator norm on $\mathcal{B}(\mathcal{H}_{\mathrm{act}})$, where $\{\Lambda_t\}_{t\geq 0}$ is a unital CPTP semigroup with GKLS generator

$$\mathcal{L}[\rho] \;=\; -\,i\,[H_{A,\mathrm{eff}},\rho] \;+\; \kappa\Big(\mathbb{E}_A(\rho)-\rho\Big).$$

Equivalently, in Lindblad form one may take $L_k=\sqrt{\kappa}\,P_k$ (record projectors from Definition 15.27), giving

$$\mathcal{L}[\rho]=-\,i[H_{A,\mathrm{eff}},\rho]+\sum_k\Big(L_k\rho L_k^\dagger-\tfrac12\{L_k^\dagger L_k,\rho\}\Big)\,,\qquad \sum_k[L_k,L_k^\dagger]=0,$$

and in this pointer case $\sum_k L_k^\dagger L_k=\kappa\,\mathbb{I}_A$ (hence unital). If $H_{A,\mathrm{eff}}$ and κ are (piecewise) time–dependent, the same product with time–sliced factors converges to the CP–divisible evolution $\Lambda_t=\mathcal{T}\exp\!\big(\int_0^t\mathcal{L}_\tau\,d\tau\big)$ with time–local GKLS generator $\mathcal{L}_t$.

Proof. (1) Finite–dimensional setting and basic properties. By Assumption 15.39, $\mathcal{H}_{\mathrm{act}}$ is finite–dimensional; all operator norms are equivalent. Each $V_\Delta^{(\kappa)}$ is CPTP and unital: it is a convex combination of the identity and the unital CPTP pinching $\mathbb{E}_A$, composed with a unitary conjugation (Lemma 15.28 and Proposition 15.29).

(2) Chernoff derivative. Expand for small Δ:

$$V_\Delta^{(\kappa)}=\Big(I+\kappa\Delta(\mathbb{E}_A-I)+o(\Delta)\Big)\circ\Big(I+\Delta\,\mathcal{L}^{\mathrm{H}}+o(\Delta)\Big),\quad \mathcal{L}^{\mathrm{H}}[\rho]:=-\,i[H_{A,\mathrm{eff}},\rho].$$

Thus the Chernoff derivative exists in operator norm and equals $\mathcal{A}=\mathcal{L}^{\mathrm{H}}+\kappa(\mathbb{E}_A-I)$.

(3) Chernoff product theorem. Since $V_0^{(\kappa)}=I$, $V_\Delta^{(\kappa)}$ are contractions and admit the derivative $\mathcal{A}$, Chernoff's theorem yields $\big(V_{t/n}^{(\kappa)}\big)^n\to e^{t\mathcal{A}}$ uniformly on compact t-intervals. Set $\mathcal{L}:=\mathcal{A}$.

(4) GKLS form and unitality. Write $\mathbb{E}_A(\rho)=\sum_k P_k\rho P_k$. Then

$$\kappa(\mathbb{E}_A-I)[\rho]=\sum_k\Big(L_k\rho L_k^\dagger-\tfrac12\{L_k^\dagger L_k,\rho\}\Big),\qquad L_k:=\sqrt{\kappa}\,P_k,$$

so $\mathcal{L}[\rho]=-i[H_{A,\mathrm{eff}},\rho]+\sum_k(\cdots)$ is GKLS. Unitality is equivalent to $\mathcal{L}(\mathbb{I})=0$, here guaranteed since $[H_{A,\mathrm{eff}},\mathbb{I}]=0$ and $\sum_k[L_k,L_k^\dagger]=0$ (indeed, L_k are normal projectors, so $\sum_k L_kL_k^\dagger=\sum_k L_k^\dagger L_k=\kappa\mathbb{I}_A$).

(5) Time–dependent rates (optional). For piecewise continuous $t\mapsto H_{A,\mathrm{eff}}(t),\kappa(t)$, replace the uniform factor by a time–sliced product $\prod_{j=1}^n V^{(\kappa(t_{j-1}))}_{(t_j-t_{j-1})}$ and refine the partition; the limit is the time–ordered exponential with time–local GKLS generator $\mathcal{L}_t$.

Zeno remark (why scaling is needed). If one omits the convex mixing and uses $V_\Delta=\mathbb{E}_A\circ\mathrm{Ad}_{e^{-iH_{A,\mathrm{eff}}\Delta}}$, then $(V_{t/n})^n$ converges to the *Zeno* dynamics on the record algebra:

$$\lim_{n\to\infty}\Big(\mathbb{E}_A\circ\mathrm{Ad}_{e^{-iH\Delta}}\Big)^n\;=\;\mathrm{Ad}_{e^{-i(\sum_k P_kHP_k)\,t}}\circ\mathbb{E}_A,$$

which is unitary (no dissipation) on the block–diagonal subalgebra and does not yield the finite–rate dissipator of $\kappa(\mathbb{E}_A-I)$. □

Definition 15.58 (Diagnostic realized reference configuration–admissible, resource–bounded control)**.** On a diagnostic realized reference configuration $\mathfrak{B}_{[\chi]}^{\langle \mathrm{DRRC}\rangle}(\mathfrak{P})$ with Good-clock $C = (\hat{T}, U_t)$ (Definitions 13.158 and 15.1), a *control protocol* is a measurable pair

$$t \;\mapsto\; \Big(H_{AE}(t),\, \gamma_\alpha(g(t))\Big)_\alpha$$

entering the GKLS generator (Propositions 15.41 and 15.45)

$$\mathcal{L}_{g(t)}[\rho] = -\,i\,[H_{A,\mathrm{eff}} + H_{AE}(t), \rho] \;+\; \sum_\alpha \gamma_\alpha(g(t))\Big(L_\alpha \rho L_\alpha^\dagger - \tfrac{1}{2}\{L_\alpha^\dagger L_\alpha,\, \rho\}\Big)\,.$$

It is *diagnostic realized reference configuration–admissible and resource–bounded* if there exist finite constants $g_{\max}, \kappa_{\max}$ with

$$\|H_{AE}(t)\| \le g_{\max}, \qquad \Gamma(t) := \sum_\alpha \gamma_\alpha(g(t)) \le \kappa_{\max} \qquad (\forall t \ge 0),$$

and it respects the endogenous coarse–graining (Definition 15.27 and Assumption 15.39) and Σ–causal support (Proposition 15.43).

Lemma 15.59 (Σ–causal control–rate bound)**.** *Let A be supported in a diagnostic realized reference configuration region of $\bar{g}^{\mathrm{Lor}}$-diameter L (Definition 5.77). Assume controls/actuators are confined to Σ–causal support (Definition 5.150 and Corollary 5.145) and the protocol is diagnostic realized reference configuration–admissible (Definition 15.1 and Assumption 15.39). Then any such protocol obeys the minimal latency*

$$T_{\mathrm{sig}} \;\ge\; \frac{L}{v_\Sigma},$$

with v_Σ the causal speed set by the Σ–lightcone (Definition 5.122). Consequently, entropy variations satisfy the Lipschitz bound of Lemma 15.60 only after a causal onset $t \ge T_{\mathrm{sig}}$.

Proof. **Step 1 (Microcausality/no–law–signalling on the diagnostic realized reference configuration).** Let the control act through a localized worldtube Γ_{ctrl} (the actuator's support) and write the Heisenberg–picture perturbation with Good–clock flow $t \mapsto \alpha_t$ (Definition 13.158). For any local observable $O_A(y)$ supported at spacetime point $y \in \mathrm{supp}(A)$, the first–order variation due to a control Hamiltonian density $\hat{h}_{\mathrm{ctrl}}(x,s)$ (or, more generally, a local CPTP instrument density) is

$$\delta\langle O_A(y,t)\rangle = \int_0^t ds\; \omega^{\mathring{\Psi}}\Big(i\,[\hat{H}_{\mathrm{ctrl}}(s),\, \alpha_{t-s}(O_A(y))]\Big) \;+\; \text{higher Dyson terms.}$$

By Σ–microcausality and no–law–signalling on the diagnostic realized reference configuration (Definition 5.150 and Corollary 5.145), all commutators vanish when (x,s) is Σ–spacelike to (y,t). Hence $\delta\langle O_A(y,t)\rangle = 0$ unless $(y,t) \in J_\Sigma^+(\Gamma_{\mathrm{ctrl}})$, the Σ–causal future of Γ_{ctrl}. This extends to all Dyson orders by the same locality (nested commutators remain zero off the Σ–cone).

Step 2 (Geometric time–of–flight bound). Let $\mathrm{dist}_\Sigma(\cdot,\cdot)$ be the $\bar{g}^{\mathrm{Lor}}$–geodesic distance (Definition 5.77), and denote the spatial cross–section of the control at its onset by Γ_0 (a compact set). Define the Σ–reachable set at time t:

$$\mathrm{Reach}(t) \;:=\; \Big\{ y :\; \mathrm{dist}_\Sigma(y,\Gamma_0) \;\le\; v_\Sigma\, t \Big\}.$$

By Step 1, an observable at y can be influenced no earlier than $t \ge \mathrm{dist}_\Sigma(y,\Gamma_0)/v_\Sigma$. To have *diagnostic realized reference configuration–level* control over the full algebra $\mathcal{A}_A$ (hence to induce a nontrivial reduced channel on A at the coarse resolution of Assumption 15.39), one must have $\mathrm{supp}(A) \subseteq \mathrm{Reach}(t)$. Therefore the minimal signalling time is

$$T_{\mathrm{sig}} \;\ge\; \frac{1}{v_\Sigma} \sup_{y\in\mathrm{supp}(A)} \mathrm{dist}_\Sigma(y,\Gamma_0).$$

Since $\mathrm{diam}_\Sigma(A) = L$ and diagnostic realized reference configuration actuators are local (their initial spatial diameter is negligible on the coarse scale; Definition 15.1 and Assumption 15.39), we have $\sup_{y\in\mathrm{supp}(A)} \mathrm{dist}_\Sigma(y,\Gamma_0) \;\ge\; L$ when Γ_0 is anchored at (or outside) a boundary patch of A. Thus $T_{\mathrm{sig}} \ge L/v_\Sigma$. *(Remark.)* If one keeps the finite actuator size $\delta := \mathrm{diam}_\Sigma(\Gamma_0)$, the same reasoning gives the slightly more general estimate $T_{\mathrm{sig}} \ge (L-\delta)/v_\Sigma$; the stated form follows from the standard diagnostic realized reference configuration assumption $\delta \ll L$.

Step 3 (Onset for entropy change and Lipschitz control). Let $\rho_A(t)$ be the reduced state of A. For $t < T_{\mathrm{sig}}$, Step 1 implies that, at the diagnostic realized reference configuration coarse resolution, the reduced channel on $\mathcal{A}_A$ is the identity: $\rho_A(t) = \rho_A(0)$, hence $S(\rho_A(t)) = S(\rho_A(0))$. For $t \ge T_{\mathrm{sig}}$, the non–Hamiltonian part $\mathcal{M}_t$ of the time–local generator (monitoring/exchange; Lemma 15.23) becomes diagnostic realized reference configuration–effective on $\mathcal{A}_A$. Therefore the continuity/Lipschitz estimate of Lemma 15.60 applies from T_{sig} onward:

$$\Big| S(\rho_A(t)) - S(\rho_A(s)) \Big| \;\le\; C_d \int_s^t \|\mathcal{M}_\tau\|_{1\to1}\, d\tau, \qquad t \ge s \ge T_{\mathrm{sig}},$$

with C_d the constant of Lemma 15.60.

Combining Steps 1–3 proves the causal latency bound and the stated consequence for entropy variations. □

Lemma 15.60 (Generator norm and trajectory Lipschitzness)**.** *Work on the finite–dimensional state–conditioned active band $\mathcal{H}_{\mathrm{act}}$ (Definition 15.56 and Assumption 15.39). Let the reduced, time–local generator be split as*

$$\mathcal{L}_{g(t)} \;=\; \mathcal{L}_t^{(\mathrm{H})} \;+\; \mathcal{M}_t, \qquad \mathcal{L}_t^{(\mathrm{H})}[\rho] := -\,i\,[H_{AE}(t),\rho],$$

where $H_{AE}(t)$ is the (interaction) Hamiltonian contribution that depends on the control strength $g(t)$, and

$$\mathcal{M}_t[\rho] \;=\; \sum_\alpha \gamma_\alpha(t)\Big(L_\alpha \rho L_\alpha^\dagger - \tfrac{1}{2}\{L_\alpha^\dagger L_\alpha, \rho\}\Big)$$

is a GKLS dissipator with rates $\gamma_\alpha(t) \ge 0$ and fixed system operators $\{L_\alpha\}$. Define

$$\Gamma(t) \;:=\; \sum_\alpha \gamma_\alpha(t)\, \|L_\alpha\|^2, \qquad g_{\max} := \sup_{t\ge0} \|H_{AE}(t)\|, \quad \kappa_{\max} := \sup_{t\ge0} \Gamma(t).$$

Then there exist constants $c_1, c_2 > 0$ (independent of t, depending only on operator–norm conventions; one may take $c_1 = c_2 = 2$) such that

$$\|\mathcal{L}_{g(t)}\|_{1\to 1} \;\le\; c_1\,\|H_{AE}(t)\| + c_2\,\Gamma(t) \;\le\; c_1 g_{\max} + c_2\kappa_{\max}.$$

Consequently, any solution of $\dot\rho_A(t) = \mathcal{L}_{g(t)}[\rho_A(t)]$ obeys the global Lipschitz bound

$$\|\rho_A(t) - \rho_A(s)\|_1 \;\le\; (c_1 g_{\max} + c_2\kappa_{\max})\,|t-s| \qquad \forall\, s,t \ge 0.$$

Proof. **(1) Hamiltonian part: commutator bound.** For any bounded K and trace–class X, $\|[K,X]\|_1 \le \|KX\|_1 + \|XK\|_1 \le \|K\|\,\|X\|_1 + \|K\|\,\|X\|_1 = 2\|K\|\,\|X\|_1$ (using $\|AXB\|_1 \le \|A\|\,\|X\|_1\,\|B\|$). Hence the induced $1\to1$ norm satisfies

$$\big\| -i[K,\cdot]\big\|_{1\to1} \;\le\; 2\,\|K\|.$$

Applying this with $K = H_{AE}(t)$ yields $\|\mathcal{L}_t^{(\mathrm{H})}\|_{1\to1} \le 2\,\|H_{AE}(t)\|$, so we may take $c_1 = 2$.

(2) Dissipator part: GKLS bound. For a single Lindblad term $\mathcal{D}_\alpha[\rho] := L_\alpha \rho L_\alpha^\dagger - \frac12\{L_\alpha^\dagger L_\alpha, \rho\}$ and $\|\rho\|_1 = 1$,

$$\|\mathcal{D}_\alpha[\rho]\|_1 \;\le\; \|L_\alpha\rho L_\alpha^\dagger\|_1 + \tfrac12\|L_\alpha^\dagger L_\alpha \rho\|_1 + \tfrac12\|\rho L_\alpha^\dagger L_\alpha\|_1 \;\le\; \|L_\alpha\|^2 + \tfrac12\|L_\alpha\|^2 + \tfrac12\|L_\alpha\|^2 \;=\; 2\,\|L_\alpha\|^2.$$

Therefore

$$\|\mathcal{M}_t\|_{1\to1} \;=\; \Big\|\sum_\alpha \gamma_\alpha(t)\,\mathcal{D}_\alpha\Big\|_{1\to1} \;\le\; \sum_\alpha \gamma_\alpha(t)\,\|\mathcal{D}_\alpha\|_{1\to1} \;\le\; 2\sum_\alpha \gamma_\alpha(t)\,\|L_\alpha\|^2 \;=\; 2\,\Gamma(t),$$

so we may take $c_2 = 2$.

(3) Generator norm and uniform bounds. By subadditivity of the operator norm,

$$\|\mathcal{L}_{g(t)}\|_{1\to1} \;\le\; \|\mathcal{L}_t^{(\mathrm{H})}\|_{1\to1} + \|\mathcal{M}_t\|_{1\to1} \;\le\; 2\,\|H_{AE}(t)\| + 2\,\Gamma(t) \;\le\; 2g_{\max} + 2\kappa_{\max}.$$

(4) Trajectory Lipschitzness. Let $\rho_A(\cdot)$ solve $\dot\rho_A(t) = \mathcal{L}_{g(t)}[\rho_A(t)]$. Integrating and using the bound from Step (3),

$$\begin{aligned}\|\rho_A(t) - \rho_A(s)\|_1 &= \Big\|\int_s^t \mathcal{L}_{g(\tau)}[\rho_A(\tau)]\,d\tau\Big\|_1 \\ &\le \int_s^t \|\mathcal{L}_{g(\tau)}\|_{1\to1}\,\|\rho_A(\tau)\|_1\,d\tau \\ &\le (2g_{\max} + 2\kappa_{\max})\,|t-s|,\end{aligned}$$

since $\|\rho_A(\tau)\|_1 = 1$ for all τ. Renaming $c_1 = c_2 = 2$ gives the stated constants; any larger c_1, c_2 depending only on the chosen norm conventions are also admissible.

Remarks. (i) If one works in the interaction picture with respect to a diagnostic realized reference configuration–fixed internal Hamiltonian, the same estimates apply because unitary conjugations preserve the trace norm and induced $1\to1$ norms. (ii) The quantities $g_{\max}, \kappa_{\max}$ are the resource bounds from Definition 15.58; they encode, respectively, a uniform bound on the interaction strength and on the aggregate monitoring rate $\Gamma(t)$.

This proves the claim. □

Lemma 15.61 (Planck-floor and resource-bounded trajectory control)**.** *Under Definition 15.58 and Definition 15.55, let $\rho_A(t)$ solve the reduced evolution on the finite active band $\mathcal{H}_{\mathrm{act}}$ over any interval $[0,T]$. Then:*

(i) Uniform Lipschitz control.
For all $s,t\in[0,T]$,

$$\|\rho_A(t)-\rho_A(s)\|_1 \;\le\; (c_1 g_{\max}+c_2\kappa_{\max})\,|t-s|, \tag{15.17}$$

with constants $c_1,c_2>0$ as in Lemma 15.60.

(ii) Planck-floor cadence.
Operational control updates (changes in $g(t)$ and/or monitoring rates) cannot be scheduled at spacings strictly smaller than the thermodynamic Planck floor $\tau_{\mathrm{Pl}}^{(\mathrm{thermo})}$ of Definition 15.55; i.e., any admissible protocol has dwell-time $\Delta t_{\mathrm{ctrl}}\ge\tau_{\mathrm{Pl}}^{(\mathrm{thermo})}$. In particular, actionable comparisons of $\rho_A(t)$ across distinct control epochs satisfy $|t-s|\ge\tau_{\mathrm{Pl}}^{(\mathrm{thermo})}$.

(iii) Earliest actionable time.
Together with the Σ–causal latency bound $T_{\mathrm{sig}}\ge L/v_\Sigma$ in Lemma 15.59, the earliest physically actionable time is

$$t \;\ge\; T_{\mathrm{on}} \;:=\; \max\Big\{\,T_{\mathrm{sig}},\,\tau_{\mathrm{Pl}}^{(\mathrm{thermo})}\,\Big\}. \tag{15.18}$$

Proof. **(a) Lipschitz bound.** By Lemma 15.60, the time–local generator $\mathcal{L}_{g(t)}$ obeys $\|\mathcal{L}_{g(t)}\|_{1\to1}\le c_1\|H_{AE}(t)\|+c_2\,\Gamma(t)\le c_1 g_{\max}+c_2\kappa_{\max}$. Integrating $\dot\rho_A(t)=\mathcal{L}_{g(t)}[\rho_A(t)]$ and using $\|\rho_A(\tau)\|_1=1$ yields

$$\|\rho_A(t)-\rho_A(s)\|_1=\Big\|\int_s^t\mathcal{L}_{g(\tau)}[\rho_A(\tau)]\,d\tau\Big\|_1\le\int_s^t\|\mathcal{L}_{g(\tau)}\|_{1\to1}\,d\tau\le(c_1g_{\max}+c_2\kappa_{\max})\,|t-s|,$$

which is (15.17).

(b) Planck-floor cadence. By Definition 15.55, the diagnostic realized reference configuration admits no admissible instrumentation or control that resolves, schedules, or certifies interventions on time steps below the thermodynamic Planck floor $\tau_{\mathrm{Pl}}^{(\mathrm{thermo})}$. Equivalently, any diagnostic realized reference configuration–admissible control $g(t)$ (and associated monitoring rates) must be piecewise constant with dwell–time $\Delta t_{\mathrm{ctrl}}\ge\tau_{\mathrm{Pl}}^{(\mathrm{thermo})}$. Hence two distinct control epochs $[t_k,t_{k+1})$ and $[t_{k+1},t_{k+2})$ satisfy $t_{k+1}-t_k\ge\tau_{\mathrm{Pl}}^{(\mathrm{thermo})}$. Therefore actionable comparisons of $\rho_A(t)$ across distinct control settings necessarily obey $|t-s|\ge\tau_{\mathrm{Pl}}^{(\mathrm{thermo})}$. (Notice that the mathematical Lipschitz bound (a) holds for all s,t, but physical scheduling/estimation is only meaningful at or above this cadence.)

(c) Earliest actionable time. From Lemma 15.59, any influence mediated within the Σ–causal structure requires at least $T_{\mathrm{sig}}\ge L/v_\Sigma$ to reach the support of A. Combining this causal latency with the Planck–floor cadence (b), no operationally detectable or controllable change can be realized before the maximum of the two latencies, i.e. before T_{on} in (15.18).

Conclusion. Items (a)–(c) establish the Lipschitz control of trajectories under resource bounds, the minimal control cadence set by the Planck floor, and the combined earliest actionable time once Σ–causality is accounted for. □

Proposition 15.62 (Finite-time purity requires unbounded resources or infinite time)**.** *Let $\rho_A(0)$ be mixed on $\mathcal{H}_{\mathrm{act}}$ with $d = \dim \mathcal{H}_{\mathrm{act}} \geq 2$, and let the protocol be admissible in the sense of Definition 15.58. For any $T > 0$,*

$$\inf_{\text{admissible controls on } [0,T]} \inf_{|\psi\rangle} \left\| \rho_A(T) - |\psi\rangle\langle\psi| \right\|_1 \;\geq\; \max\Big\{ \delta_0 - (c_1 g_{\max} + c_2 \kappa_{\max})\, T,\; 0 \Big\},$$

where

$$\delta_0 := \inf_{|\psi\rangle} \left\| \rho_A(0) - |\psi\rangle\langle\psi| \right\|_1 = 2\Big(1 - \lambda_{\max}(\rho_A(0))\Big) > 0,$$

and c_1, c_2 are from Lemma 15.60. Consequently, if $g_{\max}, \kappa_{\max} < \infty$, reaching a pure state in finite time requires

$$T \;\geq\; \frac{\delta_0}{c_1 g_{\max} + c_2 \kappa_{\max}}.$$

In the isolation scaling of Assumption 15.39 with $\Gamma(g) \to 0$ (hence $g_{\max}, \kappa_{\max} \to 0$), this lower bound diverges as resources vanish.

Proof. Under Definition 15.58, $0 \leq g(t) \leq g_{\max}$ and $0 \leq \kappa(t) \leq \kappa_{\max}$ on $[0,T]$. By Lemma 15.60, $\|\mathcal{L}_{g(t)}\|_{1\to 1} \leq L := c_1 g_{\max} + c_2 \kappa_{\max}$, so for all $s, t \in [0,T]$, $\|\rho_A(t) - \rho_A(s)\|_1 \leq L|t-s|$. Fix any admissible control and any pure $|\psi\rangle$. Then

$$\|\rho_A(T) - |\psi\rangle\langle\psi|\,\|_1 \;\geq\; \|\rho_A(0) - |\psi\rangle\langle\psi|\,\|_1 - \|\rho_A(T) - \rho_A(0)\|_1 \;\geq\; \|\rho_A(0) - |\psi\rangle\langle\psi|\,\|_1 - LT.$$

Taking the infimum over $|\psi\rangle$ gives $\inf_{|\psi\rangle} \|\rho_A(T) - |\psi\rangle\langle\psi|\,\|_1 \geq \delta_0 - LT$. The bound holds uniformly over controls, and since the left-hand side is nonnegative we take the positive part to obtain the stated inequality.

Finally, for a mixed $\rho_A(0)$ the minimum trace distance to a pure state is attained at an eigenvector of $\rho_A(0)$ with eigenvalue $\lambda_{\max}$, yielding $\delta_0 = \|\,\rho_A(0) - |\lambda_{\max}\rangle\langle\lambda_{\max}|\,\|_1 = 2(1-\lambda_{\max}) > 0$. Thus exact purification in finite time requires either unbounded resources (making L arbitrarily large) or arbitrarily long time $T \geq \delta_0/L$. □

Corollary 15.63 (Good–clock unattainability of the PQS)**.** *Within the two-level applicability (Definition 15.54), the Good-clock $C = (\hat{T}, U_t)$ is only defined on the realized (physical) level (from Planck up to cosmological scales) (Definitions 13.158 and 15.55). Let $\rho_A(t)$ be any diagnostic realized reference configuration-admissible trajectory (Definition 15.1) with resource-bounded control (Definition 15.58). Then no finite Good-clock time $T < \infty$ can reach the pre-physical PQS regime (where distances/times collapse to 0) or an exactly PQS-coherent state. Equivalently, to achieve $\rho_A(t_n) \to |\psi\rangle\langle\psi|$ with vanishing AE correlations requires a sequence $t_n \uparrow \infty$.*

Proof. We separate the argument into three obstructions—*domain*, *operational*, and *structural*—each sufficient on its own and together mutually reinforcing.

(A) Domain obstruction (Good-clock domain excludes PQS). By Definitions 15.54 and 15.55, the Good-clock $C = (\hat{T}, U_t)$ and all thermodynamic predicates are *undefined* on the pre-physical PQS stratum, where operational distances/times vanish. The Good-clock parameter t ranges over the realized (physical) domain only, beginning at a strictly positive Planck-floor resolution. Hence there is no $t = T < \infty$ in the Good-clock domain that *labels* a PQS state; the target lies outside the parameter space of t. This already forbids "reaching PQS at finite t" in the strict sense of domain.

(B) Operational obstruction (causal onset and Lipschitz control under bounded resources). Diagnostic realized reference configuration-admissible controls are constrained by Σ-causality and the Planck floor. First, Σ-causal latency implies a minimal signalling/onset time $T_{\mathrm{sig}} \geq L/v_\Sigma$ before any control can influence A (Lemma 15.59). Second, the Planck floor bounds update cadence and induces trajectory Lipschitzness: for constants c_1, c_2 depending only on the active-band dimension and operator set,

$$\|\rho_A(t) - \rho_A(s)\|_1 \;\leq\; (c_1 g_{\max} + c_2 \kappa_{\max})\,|t-s| \quad \text{with} \quad t - s \;\geq\; \tau_{\mathrm{Pl}}^{(\mathrm{thermo})}$$

(Lemmas 15.60 and 15.61). Thus the earliest actionable time is

$$t \;\geq\; T_{\mathrm{on}} := \max\{T_{\mathrm{sig}},\, \tau_{\mathrm{Pl}}^{(\mathrm{thermo})}\}.$$

Even beyond onset, bounded resources limit how far the state can move in trace distance per unit time. In particular, the finite-time purity lower bound (Proposition 15.62) gives, for any pure $|\psi\rangle$,

$$\Big\|\rho_A(T) - |\psi\rangle\langle\psi|\,\Big\|_1 \;\geq\; \max\Big\{\delta_0 - (c_1 g_{\max} + c_2 \kappa_{\max})\,T,\; 0\Big\},$$

where $\delta_0 = 2\Big(1 - \lambda_{\max}(\rho_A(0))\Big) > 0$ unless $\rho_A(0)$ is already pure. Hence exact purification in *finite* Good-clock time would require either (i) T exceeding the resource-limited threshold $\delta_0/(c_1 g_{\max} + c_2 \kappa_{\max})$, or (ii) unbounded resources; neither is compatible with the resource-bounded, diagnostic realized reference configuration-admissible regime. This blocks finite-time arrival at a pure target (a fortiori at an exactly PQS-coherent state).

(C) Structural obstruction (unital CP-divisible arrow and the isolation limit). Within the thermodynamic coarse-grained regime (Assumption 15.39), the reduced A-dynamics is CP-divisible and (for the pointer channel considered) unital (Stage 3), with generator of GKLS form (e.g. Propositions 15.41 and 15.57). The Second-Law monotonicity then yields

$$t_2 \geq t_1 \;\Rightarrow\; S(\rho_A(t_2)) \;\geq\; S(\rho_A(t_1)) \quad \text{and} \quad \frac{d}{dt} S(\rho_A(t)) \geq 0$$

(Theorem 15.35 and Proposition 15.47). Consequently, if $\rho_A(0)$ is mixed, no finite Good-clock time can reduce $S(\rho_A(t))$ to 0 under the admissible (unital, CP-divisible) flow.

One might try to decouple A from E by taking the isolation scaling $g \to 0$. But Proposition 15.52 shows that as $g \to 0$ the dissipator vanishes and the reduced dynamics becomes *unitary*:

$$\Lambda_t^{(g)} \;\to\; \mathrm{Ad}_{e^{-iH_{A,\mathrm{eff}}t}}, \qquad S(\Lambda_t^{(g)}\rho_A(0)) \;\to\; S(\rho_A(0)).$$

Local unitaries cannot change the entanglement entropy $S(\rho_A)$ or remove AE correlations. Thus the very limit needed to erase intersystem correlations also freezes the entropy and precludes purification in finite time. Combined with the small-coupling continuity bound (Lemma 15.51), any purification/decoupling is at best *asymptotic* in time or requires leaving the admissible regime (e.g. non-unital baths, unbounded controls, or nonlocal operations).

Conclusion. (A) excludes a finite-t PQS "target" on domain grounds; (B) enforces finite signal/onset times and resource-limited Lipschitz motion, yielding a positive lower bound on the time needed to reach purity (and thus forbidding it under bounded resources); (C) shows that, within the admissible unital CP-divisible evolution, entropy cannot be driven to zero, and that the isolation limit which would suppress correlations simultaneously eliminates the very mechanism that could reduce entropy. Therefore no finite Good-clock time $T < \infty$ can reach the pre-physical PQS regime or an exactly PQS-coherent state. Any approach to a pure, uncorrelated limit must occur—if at all—only along sequences $t_n \uparrow \infty$. □

Theorem 15.64 (Unattainability under finite resources)**.** *Under Definition 15.58 let* $d = \dim \mathcal{H}_{\mathrm{act}} < \infty$*, and suppose* $\rho_A(0)$ *is mixed. Then exact purity cannot be achieved at any* finite *emergent time* $T < \infty$ *unless at least one resource bound is taken unbounded (i.e.* $g_{\max} \to \infty$ *or* $\kappa_{\max} \to \infty$*). Equivalently, if* $\rho_A(t_n) \to |\psi\rangle\langle\psi|$ *for some sequence* $t_n \uparrow \infty$*, then such a sequence is required when* $(g_{\max}, \kappa_{\max})$ *are fixed and finite.*

Moreover, the Audenaert–Fannes continuity bound combined with the Lipschitz control of trajectories (Lemma 15.60) gives, for every $T \geq 0$*,*

$$S\Big(\rho_A(T)\Big) \;\geq\; \Big[S\Big(\rho_A(0)\Big) - \underbrace{\tfrac{1}{2}\,(c_1 g_{\max} + c_2 \kappa_{\max})\,T\,\log(d-1)}_{\textit{linear term}} - \underbrace{H_2\Big(\tfrac{1}{2}\,(c_1 g_{\max} + c_2 \kappa_{\max})\,T\Big)}_{\textit{binary entropy}}\Big]_+,$$

where $H_2(x) = -x \log x - (1-x)\log(1-x)$ *(natural logs,* $k_B{=}1$*). Thus, with* bounded $(g_{\max}, \kappa_{\max})$*, driving* $S(\rho_A(T)) \to 0$ *requires* $T \to \infty$*; a finite-time collapse would entail a non–norm-continuous (or unbounded–rate) evolution.*

In the unital *endogenous setting of Propositions 15.41 and 15.47, entropy is nondecreasing,* $t_2 \geq t_1 \Rightarrow S(\rho_A(t_2)) \geq S(\rho_A(t_1))$*, so exact purification from a mixed state is* strictly *unattainable at all times.*

Good-clock/typed invariance. All statements in Theorem 15.64 are invariant under Good-clock gauge changes and persist under admissible typing $\Pi^{[\chi]}$ (Lemma 15.6 and Proposition 9.138).

Theorem 15.65 (Third Law — unattainability with two–level, Planck, and Σ–causal constraints)**.** *Assume the two–level applicability (Definition 15.54). On the realized level (R), let the control be diagnostic realized reference configuration–admissible and resource–bounded (Definition 15.58), obey the Planck floor (Definition 15.55), and respect* Σ*–causal support (Lemma 15.59). If* $\rho_A(0)$ *is mixed, then exact purity at any finite emergent time is impossible unless resources are unbounded. Quantitatively, for any* $T > 0$*,*

$$\inf_{\substack{\textit{admissible controls}\\ \textit{on } [0,T]}} \; \inf_{|\psi\rangle} \; \big\|\rho_A(T) - |\psi\rangle\langle\psi|\big\|_1 \;\geq\; \Big[\delta_0 - (c_1 g_{\max} + c_2 \kappa_{\max})\,\Big(T - T_{\mathrm{on}}\Big)\Big]_+,$$

where $\delta_0 := \inf_{|\psi\rangle} \|\rho_A(0) - |\psi\rangle\langle\psi|\,\|_1 > 0$*,* $T_{\mathrm{on}} := \max\{T_{\mathrm{sig}}, \tau_{\mathrm{Pl}}^{(\mathrm{thermo})}\}$ *(Lemma 15.59 and Definition 15.55), and* c_1, c_2 *are as in Lemma 15.60. In particular, with finite* $g_{\max}, \kappa_{\max}$ *and nonzero* T_{on}*, driving* $\rho_A(T)$ *to a pure state requires* $T \to \infty$ *or unbounded resources.*

Proof. Onset window. By Σ–causal latency and the Planck floor, no admissible control can take effect before $T_{\mathrm{on}} := \max\{T_{\mathrm{sig}}, \tau_{\mathrm{Pl}}^{(\mathrm{thermo})}\}$ (Lemma 15.59 and Definition 15.55). Thus

any nontrivial (i.e. control–induced) change of the reduced state must occur in the interval $[T_{\rm on}, T]$.

Lipschitz control after onset. For resource–bounded controls (Definition 15.58), Lemma 15.60 gives the induced trace–norm bound $\|\mathcal{L}_{g(t)}\|_{1\to 1} \le L := c_1 g_{\max} + c_2 \kappa_{\max}$. Hence, for $T \ge T_{\rm on}$,

$$\big\|\rho_A(T) - \rho_A(T_{\rm on})\big\|_1 \;\le\; L\,\big(T - T_{\rm on}\big).$$

Triangle inequality vs. pure targets. For any unit vector $|\psi\rangle$,

$$\begin{aligned}\big\|\rho_A(T) - |\psi\rangle\langle\psi|\,\big\|_1 &\ge \big\|\rho_A(T_{\rm on}) - |\psi\rangle\langle\psi|\,\big\|_1 - \big\|\rho_A(T) - \rho_A(T_{\rm on})\big\|_1 \\ &\ge \big\|\rho_A(T_{\rm on}) - |\psi\rangle\langle\psi|\,\big\|_1 - L\,(T - T_{\rm on}).\end{aligned}$$

Taking the infimum over $|\psi\rangle$ yields

$$\begin{aligned}\inf_{|\psi\rangle}\big\|\rho_A(T) - |\psi\rangle\langle\psi|\,\big\|_1 \;\ge\; &\inf_{|\psi\rangle}\big\|\rho_A(T_{\rm on}) - |\psi\rangle\langle\psi|\,\big\|_1 \\ &- L\,(T - T_{\rm on}).\end{aligned}$$

Unitary drift before onset preserves the spectral gap to purity. Before $T_{\rm on}$ only diagnostic realized reference configuration–legal unitary drift on A is possible (no monitoring/exchange yet). Unitary conjugation preserves the spectrum of ρ_A, hence preserves the minimal trace–norm distance to the pure–state manifold (a spectral quantity). Therefore

$$\inf_{|\psi\rangle}\big\|\rho_A(T_{\rm on}) - |\psi\rangle\langle\psi|\,\big\|_1 \;=\; \inf_{|\psi\rangle}\big\|\rho_A(0) - |\psi\rangle\langle\psi|\,\big\|_1 \;=:\; \delta_0 \;>\; 0,$$

since $\rho_A(0)$ is mixed and $d = \dim \mathcal{H}_{\rm act} \ge 2$.

Combine. Putting the pieces together and using the positive part $[x]_+ = \max\{x, 0\}$ to cover the case $T < T_{\rm on}$,

$$\inf_{\substack{\text{admissible controls}\\ \text{on } [0,T]}}\ \inf_{|\psi\rangle}\big\|\rho_A(T) - |\psi\rangle\langle\psi|\,\big\|_1 \;\ge\; \Big[\,\delta_0 - L\,(T - T_{\rm on})\,\Big]_+.$$

If $g_{\max}, \kappa_{\max} < \infty$ then $L < \infty$ and the right–hand side can vanish only in the limit $T \to \infty$ (or by taking resources unbounded). □

Good-clock/typed invariance. All statements in Theorem 15.65 are invariant under Good-clock gauge changes and persist under admissible typing $\Pi^{[\chi]}$ (Lemma 15.6 and Proposition 9.138).

Corollary 15.66 (Time-to-purity lower bound with Planck/causal latency)**.** *Let $L := c_1 g_{\max} + c_2 \kappa_{\max}$ and $\delta_0 := \inf_{|\psi\rangle} \|\rho_A(0) - |\psi\rangle\langle\psi|\,\|_1 > 0$. If resources are bounded and $\rho_A(0)$ is mixed, then any protocol that achieves $\|\rho_A(T) - |\psi\rangle\langle\psi|\,\|_1 \le \epsilon$ satisfies*

$$T \;\ge\; T_{\rm on} \;+\; \frac{[\delta_0 - \epsilon]_+}{L}, \qquad [x]_+ := \max\{x, 0\}.$$

In particular, if $\epsilon < \delta_0$ one needs $T \ge T_{\rm on} + (\delta_0 - \epsilon)/L$; if $\epsilon \ge \delta_0$ the latency reduces to $T \ge T_{\rm on}$. In the isolation scaling of Assumption 15.39 with $\Gamma(g) \to 0$ (so $L \to 0$), the denominator vanishes and the bound diverges unless $\epsilon \ge \delta_0$.

Remark 15.67 (Zero-resource limit)**.** If $L = 0$ (equivalently $g_{\max} = \kappa_{\max} = 0$), then for any target accuracy $\epsilon < \delta_0$ no finite T can meet the bound: the inequality forces $T \to \infty$. If $\epsilon \geq \delta_0$, the latency reduces to $T \geq T_{\mathrm{on}}$. This is consistent with Proposition 15.52: in the isolation limit the evolution is unitary (spectrum preserved), so the minimal trace distance to any pure state remains δ_0 and exact (or better-than-δ_0) purification is unattainable in finite time.

Proof. From Theorem 15.65,

$$\inf_{\text{adm. controls}} \inf_{|\psi\rangle} \|\rho_A(T) - |\psi\rangle\langle\psi|\,\|_1 \;\geq\; [\delta_0 - L(T - T_{\mathrm{on}})]_+.$$

Necessity of $\|\rho_A(T)-|\psi\rangle\langle\psi|\,\|_1 \leq \epsilon$ implies $[\delta_0-L(T-T_{\mathrm{on}})]_+ \leq \epsilon$, i.e. $T \geq T_{\mathrm{on}}+[\delta_0-\epsilon]_+/L$. □

Corollary 15.68 (Two-level Third Law summary)**.** *(*P level*) No thermodynamic map exists below the Planck floor; "absolute zero" (perfect isolation/purity) is not a thermodynamic predicate (Definitions 15.54 and 15.55).*
*(*R level*) Entropy production vanishes in the isolation limit (Proposition 15.52), but exact purity in finite time is unattainable under finite resources (Theorem 15.65). Residual entropy equals* $\log \mathfrak{g}$ *when* $H_{A,\mathrm{eff}}$ *has a* $\mathfrak{g}$*–fold degenerate ground space (Corollary 15.53).*

Remark 15.69 (Anisotropy and directional unattainability)**.** If the current–kernel $\mathcal{U}_{\Phi\Theta}$ is anisotropic (Definition 15.25 and Remark 15.44), both the causal onset and the dissipative envelope acquire *directional* corrections, but the form of the unattainability bounds is unchanged.

Directional causal latency. For a unit direction $\hat{n}$ and diagnostic realized reference configuration diameter $L_{\hat{n}}$ measured in the Lorentzian metric $\bar{g}^{\mathrm{Lor}}$ (Definition 5.77), the Σ–causal architecture (Definitions 5.122 and 5.150) implies

$$T_{\mathrm{sig}}(\hat{n}) \;\geq\; \frac{L_{\hat{n}}}{v_\Sigma(\hat{n})},$$

where $v_\Sigma(\hat{n})$ is the maximal $\bar{\Sigma}$–signal speed along $\hat{n}$. With the Planck floor one has $T_{\mathrm{on}}(\hat{n}) = \max\{T_{\mathrm{sig}}(\hat{n}), \tau_{\mathrm{Pl}}^{(\mathrm{thermo})}\}$ (Lemma 15.59 and Definition 15.55).

Directional resource envelope. Let $\Gamma^{(\hat{n})}(t) := \sum_\alpha \gamma_\alpha^{(\hat{n})}(g(t))$ be the directional dissipative rate bound induced by $\mathcal{U}_{\Phi\Theta}$ and suppose

$$\|H_{AE}^{(\hat{n})}(t)\| \;\leq\; g_{\max}^{(\hat{n})}, \qquad \Gamma^{(\hat{n})}(t) \;\leq\; \kappa_{\max}^{(\hat{n})}.$$

Then the trajectory Lipschitz bound (Lemmas 15.60 and 15.61) and the time–to–purity inequality (Corollary 15.66) hold with

$$g_{\max} \;\to\; g_{\max}^{(\hat{n})}, \quad \kappa_{\max} \;\to\; \kappa_{\max}^{(\hat{n})}, \quad T_{\mathrm{on}} \;\to\; T_{\mathrm{on}}(\hat{n}).$$

Structure preserved. The reduced evolution on the active band remains unital CPTP (and thus CP–divisible) (Propositions 15.41 and 15.45); Σ–causal support is preserved (Proposition 15.43); and all statements are invariant under Good–clock gauge changes (Lemma 15.6). In the isolation scaling $\Gamma^{(\hat{n})}(g) \to 0$, the time–to–purity bound diverges for each $\hat{n}$. The isotropic case is recovered by setting $g_{\max}^{(\hat{n})} \equiv g_{\max}$ and $\kappa_{\max}^{(\hat{n})} \equiv \kappa_{\max}$.

Remark 15.70 (Infinite–dimensional extension)**.** Beyond the active–band truncation, the Lipschitz and time–to–purity logic extend on energy–constrained sets by replacing d–dependent continuity constants with energy–modulus bounds (Winter–Shirokov–type). The diagnostic realized reference configuration construction enforces such constraints via Assumption 15.39 and Definition 15.55, so the unattainability statements persist in the infinite–dimensional setting under the same operational constraints.

Theorem 15.71 (Third Law of Thermodynamics — coherence and unattainability)**.** *Work on a realized thermodynamic diagnostic realized reference configuration* $\mathfrak{B}_{[\chi]}^{\langle \mathrm{DRRC}\rangle}(\mathfrak{P})$ *with Good–clock* $C = (\hat{T}, U_t)$ *aligned to* $\bar{\Sigma}$ *(Definitions 5.26, 13.158 and 15.1). Let the reduced dynamics on the state–conditioned active band* $\mathcal{H}_{\mathrm{act}}$ *(Definition 15.56) be the unital CPTP semigroup* $\{\Lambda_t^{(g)}\}_{t\geq 0}$ *obtained by the product limit (Propositions 15.41 and 15.45) with GKLS generator*

$$\mathcal{L}_g[\rho] \;=\; -\,i\,[H_{A,\mathrm{eff}},\rho] \;+\; \sum_{\alpha}\gamma_\alpha(g)\Big(L_\alpha\rho L_\alpha^\dagger - \tfrac{1}{2}\{L_\alpha^\dagger L_\alpha,\,\rho\}\Big)\,,$$

where $g \geq 0$ *quantifies the current–kernel coupling at the* $A|E$ *interface (Definition 15.25), and the endogenous Markov limit of Assumption 15.39 ensures* $\Gamma(g) := \sum_\alpha \gamma_\alpha(g) \to 0$ *as* $g \to 0$*. Then:*

(i) Coherence (isolation) limit.
For every finite t*,*

$$\lim_{g\to 0}\Lambda_t^{(g)} \;=\; \mathrm{Ad}_{e^{-iH_{A,\mathrm{eff}}t}} \quad \textit{and} \quad \lim_{g\to 0} S\Big(\Lambda_t^{(g)}\rho_A(0)\Big) \;=\; S\Big(\rho_A(0)\Big),$$

i.e. the dynamics becomes unitary and entropy production vanishes (Proposition 15.52 and Lemma 15.51).

(ii) Zero production rate in the isolation limit.
Writing $\dot{S}_A^{(g)}(t)$ *for the Good–clock entropy rate (Lemma 15.6),*

$$\forall t < \infty: \qquad \lim_{g\to 0}\dot{S}_A^{(g)}(t) \;=\; 0.$$

Equivalently, for any admissible emergent time (an axis of $\vec{t}$ *or* t_{geom}*; Definitions 5.95 and 15.21), the Second–Law inequality (15.14) tightens to equality in the isolation limit (Proposition 15.47).* Reason: *in the interaction picture the generator is purely dissipative with norm* $O(\Gamma(g))$*, so the Audenaert–Fannes bound gives* $|S(t+\delta)-S(t)| \leq C_d\,\Gamma(g)\,\delta + o(\Gamma(g)\delta)$ *and hence* $\dot{S}_A^{(g)}(t) \to 0$ *for a.e.* t*.*

(iii) Unattainability under finite resources (realized level).
Assume diagnostic realized reference configuration–admissible, resource–bounded control in the sense of Definition 15.58 together with the Planck floor and Σ*–causal onset (Definition 15.55 and Lemmas 15.59 and 15.61). If* $\rho_A(0)$ *is mixed, then reaching a perfectly pure state in finite emergent time is impossible under finite resources: any sequence of controls achieving* $S(\rho_A(t_n)) \to 0$ *requires* $t_n \to \infty$ *or unbounded* $(g_{\max}, \kappa_{\max})$*. Quantitatively,*

$$\inf_{\substack{\text{admissible controls}\\ \text{on } [0,T]}}\ \inf_{|\psi\rangle}\ \Big\|\rho_A(T) - |\psi\rangle\langle\psi|\,\Big\|_1 \;\geq\; \Big[\,\delta_0 - (c_1 g_{\max} + c_2\kappa_{\max})\,(T - T_{\mathrm{on}})\,\Big]_+,$$

with $\delta_0 := \inf_{|\psi\rangle} \|\rho_A(0) - |\psi\rangle\langle\psi|\,\|_1 > 0$ *and* $T_{\mathrm{on}} := \max\{T_{\mathrm{sig}}, \tau_{\mathrm{Pl}}^{(\mathrm{thermo})}\}$ *(Theorem 15.65 and Corollary 15.66).*

All statements are invariant in the Good–clock gauge class $[C]$ *and persist under admissible typing* $\Pi^{[\chi]}$ *(Lemma 15.6, Section 12.3, and Definition 9.58); they preserve* Σ*–causal support in the continuum limit (Proposition 15.43). If* $H_{A,\mathrm{eff}}$ *has a* $\mathfrak{g}$*–fold degenerate ground space, the isolation limit admits residual entropy* $S_0 = \log \mathfrak{g}$ *(Corollary 15.53). On the pre–physical level (Definition 15.54), thermodynamic diagnostic realized reference configuration and Good–clock time are inapplicable.*

Proof. (i) By Assumption 15.39, $\Gamma(g) \to 0$; hence $\mathcal{L}_g \to -i[H_{A,\mathrm{eff}}, \cdot\,]$ in operator norm and $e^{t\mathcal{L}_g} \to \mathrm{Ad}_{e^{-iH_{A,\mathrm{eff}}t}}$ (Proposition 15.52). Audenaert–Fannes continuity with the interaction–picture bound $\|\dot{\tilde{\rho}}\|_1 \le O(\Gamma(g))$ gives $S(\Lambda_t^{(g)}\rho) \to S(\rho)$ (Lemma 15.51).

(ii) Unital CP–divisible flow implies $t \mapsto S(\rho_A^{(g)}(t))$ is nondecreasing (Proposition 15.47). The small–g continuity estimate yields $|S(t+\delta) - S(t)| \le C_d\,\Gamma(g)\,\delta + o(\Gamma(g)\delta)$; divide by δ and send $g \to 0$ to get $\dot{S}_A^{(g)}(t) \to 0$ for a.e. t. Gauge invariance follows from Lemma 15.6.

(iii) Under Definition 15.58, $\|\mathcal{L}_{g(t)}\|_{1\to 1} \le c_1 g_{\max} + c_2 \kappa_{\max}$ (Lemma 15.60), with actionable time $t \ge T_{\mathrm{on}}$ from Lemmas 15.59 and 15.61 and Definition 15.55. Thus $\|\rho_A(T) - \rho_A(T_{\mathrm{on}})\|_1 \le (c_1 g_{\max} + c_2\kappa_{\max})(T - T_{\mathrm{on}})$, giving the trace–distance lower bound to any pure state; Audenaert–Fannes converts this to an entropy bound, and the CP–divisible setting precludes finite–time purity with bounded resources (Theorem 15.65 and Corollary 15.66). The pre–physical level is excluded by Definition 15.54. □

Corollary 15.72 (Residual entropy and degeneracy)**.** *Let* P_0 *be the spectral projector onto the ground–energy manifold of* $H_{A,\mathrm{eff}}$ *and let* $\mathfrak{g} := \operatorname{rank} P_0$*. Assume that, for each fixed* $g > 0$*, the reduced dynamics* $\{\Lambda_t^{(g)}\}_{t\ge 0}$ *on the active band is a CPTP semigroup with a unique stationary state* $\sigma^{(g)}$ *such that*

$$\operatorname{supp}\sigma^{(g)} = P_0\mathcal{H}_{\mathrm{act}}, \qquad \text{and the restriction of } \Lambda_t^{(g)} \text{ to } P_0\mathcal{H}_{\mathrm{act}} \text{ is unital and primitive.}$$

(Equivalently, the fixed points within $P_0\mathcal{H}_{\mathrm{act}}$ *are proportional to* P_0*.) Then, taking the long–time limit at fixed* g *followed by the isolation limit* $g \to 0$*,*

$$\lim_{g\to 0}\lim_{t\to\infty} \rho_A^{(g)}(t) = \frac{P_0}{\mathfrak{g}}, \qquad S_0 := \lim_{g\to 0}\lim_{t\to\infty} S\big(\rho_A^{(g)}(t)\big) = \log \mathfrak{g}.$$

In particular, if the ground state is nondegenerate ($\mathfrak{g} = 1$*), then* $S_0 = 0$*.*

Proof. By assumption, for each fixed $g > 0$ the CPTP semigroup $\Lambda_t^{(g)}$ has a unique stationary state $\sigma^{(g)}$ with $\operatorname{supp}\sigma^{(g)} = P_0\mathcal{H}_{\mathrm{act}}$; standard semigroup ergodicity (e.g., Spohn's theorem) then yields $\rho_A^{(g)}(t) \to \sigma^{(g)}$ as $t \to \infty$. On the ground block, the restriction is unital and primitive, hence its unique fixed point on $P_0\mathcal{H}_{\mathrm{act}}$ is the maximally mixed state $P_0/\mathfrak{g}$. Therefore $\lim_{t\to\infty} \rho_A^{(g)}(t) = P_0/\mathfrak{g}$, and $\lim_{t\to\infty} S(\rho_A^{(g)}(t)) = \log \mathfrak{g}$. Taking $g \to 0$ afterward does not change this value in the stated order of limits (isolation freezes the approach but does not alter the fixed point), so $S_0 = \log \mathfrak{g}$. For $\mathfrak{g} = 1$ the limit state is pure and $S_0 = 0$. □

Remark 15.73 (On unital vs. non–unital globally, and order of limits)**.** The assumption above is intentionally global–*non-unital* (cooling/thermalization) with a unital, primitive *restriction* on P_0. This is the natural setting for low-temperature Davies–type generators or detailed balance semigroups: for each fixed $g > 0$ there is a unique stationary state $\sigma^{(g)}$ concentrating on P_0, while intra–block mixing is unital, giving $P_0/\mathfrak{g}$. If instead you keep the dynamics globally unital, contraction from excited sectors to P_0 is not possible unless the initial support is already in P_0 (the microcanonical case of Corollary 15.53). Finally, if one reverses limits—taking $g \to 0$ at fixed finite time—one recovers purely unitary evolution and entropy conservation (the isolation/coherence limit), consistent with Theorem 15.71.

Unattainability in Finite Emergent Time. In the IQR setting, *exact isolation* ($g = 0$) cannot be achieved in finite Good–clock time with finite control resources. Endogenous constraints—reciprocal resolution limits, state–dependent spectral gapping, and the PQS coherence width from Chapter 4—fix coherence/memory scales that underwrite the coarse–grained, unital GKLS limit (Assumption 15.39, Definition 4.576, eq. (4.278), and Section 4.15.3). As the current–kernel coupling $g(t)\downarrow 0$ (Definition 15.25), the dissipative power $\Gamma\big(g(t)\big) = \sum_\alpha \gamma_\alpha\big(g(t)\big)$ vanishes, the generator $\mathcal{L}_{g(t)}$ contracts toward the purely Hamiltonian part, and the CP–divisible reduced dynamics approaches unitary evolution (Propositions 15.47 and 15.52; cf. [617, 624, 441]). With diagnostic realized reference configuration–admissible, resource–bounded control (Definition 15.58), the trace–distance evolution is uniformly Lipschitz (Lemmas 15.60 and 15.61).

Crucial point. In the unital benchmark (einselection/pinching; Definition 15.27), decoherence *cannot* increase purity: entropy is nondecreasing (Proposition 15.47), so purity cannot be approached at all. More generally, even in non–unital (cooling) settings where purity can in principle increase, any protocol that aims at exact purity must accumulate *unbounded* net cooling/relaxation action; operationally this manifests as a divergent integrated dissipative strength as the error to a pure state vanishes:

$$\int_0^{t_*} \Gamma\big(g(\tau)\big)\, d\tau \;\longrightarrow\; \infty \quad \text{whenever} \quad \big\|\rho_A(t_*) - |\psi\rangle\langle\psi|\,\big\|_1 \;\longrightarrow\; 0,$$

in line with quantum unattainability results [632, 633]. Equivalently, the time–to–purity lower bound of Corollary 15.66 diverges when the resource envelope remains finite, with a causal/Planckian onset T_{on} enforced by Lemma 15.59 and Definition 15.55. This yields the strengthened unattainability theorem Theorem 15.65.

Relation to the Classical Third Law. Nernst's heat theorem and the unattainability formulation assert $S \to \mathrm{const}$ (often 0 for a nondegenerate ground state) as $T \to 0$, and that $T = 0$ cannot be reached in finite time [631, 632]. In IQR, the environment–selected steady state (picked by $\mathcal{U}_{\Phi\Theta}$) plays the role of a Gibbs surrogate; *in general (possibly non–unital) settings* this selection fixes the asymptotic state, while in the benchmark unital case the steady state is maximally mixed on the monitored blocks. Driving the coupling $g \to 0$ removes steady–state selection and returns the subsystem to unitary dynamics on the active band (Propositions 15.41 and 15.45). Thus entropy production freezes (Proposition 15.52 and Lemma 15.51) and the boundary is unattainable in finite emergent time under finite

resources (Theorems 15.65 and 15.71). Residual entropy is fixed by ground–space degeneracy, $S_0 = \log \mathfrak{g}$ (Corollary 15.72; cf. [119, 634, 626, 282]).

As the relational coupling g that controls the current–kernel interaction $\mathcal{U}_{\Phi\Theta}$ is reduced, the GKLS generator on the state–conditioned active band sheds its dissipator and converges to the Hamiltonian Liouvillian; the reduced dynamics becomes unitary and $S\big(\rho_A(t)\big)$ becomes constant (Proposition 15.52). However, endogenous spectral/temporal constraints (Assumption 15.39, Definition 4.576, and Section 4.15.3) and Good–clock/causal latencies (Definition 13.158 and Lemma 15.59) force the dissipative power to vanish too rapidly for the purity boundary to be reached in finite emergent time with bounded resources (Theorem 15.65); cf. [632, 633].

Classical Correspondence and Resource Viewpoint. Viewing the environment–selected steady state as an effective Gibbs reference, the isolation limit $g \to 0$ corresponds to $T \to 0$: entropy production vanishes and unattainability holds (Nernst/Planck) [631, 632, 633]. From a resource–theoretic angle, isolation removes the free noisy operations; coherence and purity thus become asymptotically conserved resources [628, 629, 630]. In IQR terms: einselection by the pointer channel (Definition 15.27) ceases as $\Gamma(g) \to 0$, pinching no longer produces entropy (Lemma 15.30 and Proposition 15.31), and the unital CP–divisible flow degenerates to a unitary group (Propositions 15.41 and 15.47).

The Third Law as the Law of Quantum Coherence. Most naturally, the Third Law is a *law of quantum restoration*: as the current–kernel coupling g at the $A|E$ boundary (Definition 15.25) is driven to zero within the realized (R) level (Definition 15.54), the reduced dynamics becomes asymptotically unitary, the von Neumann entropy production vanishes, and coherences in any fixed pointer basis are preserved (Proposition 15.52 and Lemma 15.51). Informational isolation thus yields asymptotic purity and aligns the IQR, information–theoretic reading of thermodynamics with the traditional Third Law [626, 635], while respecting the Planck floor and Σ–causal support (Definition 15.55 and Proposition 15.43).

> **Third Law (relational form — coherence and unattainability).** On an *instrumented realization on a certified plateau* with a Good–clock $C = (\hat{T}, U_t)$ and state–conditioned active band $\mathcal{H}_{\mathrm{act}}$, the reduced evolution at coupling $g \geq 0$ is the unital, CP–divisible flow $\{\Lambda_t^{(g)}\}_{t\geq 0}$ on $\mathcal{B}(\mathcal{H}_{\mathrm{act}})$ generated by
>
> $$\mathcal{L}_g[\rho] = -\,i\,[H_{A,\mathrm{eff}}, \rho] + \sum_\alpha \gamma_\alpha(g)\Big(L_\alpha \rho L_\alpha^\dagger - \tfrac{1}{2}\{L_\alpha^\dagger L_\alpha, \rho\}\Big)\,,$$
> $$\Gamma(g) := \sum_\alpha \gamma_\alpha(g) \;\xrightarrow[g\downarrow 0]{}\; 0.$$
>
> *Coherence (isolation) limit.* For every finite t,
>
> $$\boxed{\lim_{g\to 0} \Lambda_t^{(g)} = \mathrm{Ad}_{e^{-iH_{A,\mathrm{eff}}t}}, \qquad \lim_{g\to 0} S\big(\Lambda_t^{(g)}\rho\big) = S(\rho), \qquad \lim_{g\to 0} \dot{S}_A^{(g)}(t) = 0}$$
>
> i.e. entropy production *flattens* and the flow becomes unitary (asymptotic coherence).

Unattainability under finite resources. Any *instrumented realization on a certified plateau*–admissible, resource–bounded control with $\|H_{AE}(t)\| \le g_{\max}$, $\Gamma(g(t)) \le \kappa_{\max}$ and causal/Planckian onset

$$\underbrace{T_{\mathrm{on}} := \max\Big\{T_{\mathrm{sig}},\, \tau_{\mathrm{Pl}}^{(\mathrm{thermo})}\Big\}}_{\substack{\text{signal latency / Planck floor}\\ (\Sigma\text{–causal \& timing constraints})}}$$

obeys, for any $T > 0$ and any pure $|\psi\rangle$,

$$\boxed{\begin{aligned}\Big\|\rho_A(T) - |\psi\rangle\langle\psi|\Big\|_1 &\ge \Big[\ \delta_0 - \underbrace{(c_1 g_{\max} + c_2\kappa_{\max})}_{\text{resource envelope}}\,(T - T_{\mathrm{on}})\ \Big]_+,\\ \delta_0 &:= \inf_{|\psi\rangle}\Big\|\rho_A(0) - |\psi\rangle\langle\psi|\Big\|_1 > 0\end{aligned}}$$

(where $[x]_+ := \max\{x, 0\}$). Equivalently,

$$T \ \ge\ T_{\mathrm{on}} + \frac{\delta_0 - \epsilon}{c_1 g_{\max} + c_2\kappa_{\max}} \quad \text{to reach} \quad \Big\|\rho_A(T) - |\psi\rangle\langle\psi|\Big\|_1 \le \epsilon.$$

Residual entropy (degenerate ground spaces). Let P_0 project onto the ground manifold of $H_{A,\mathrm{eff}}$ and $\mathfrak{g} := \operatorname{rank} P_0$. If the long–time unital dynamics is ergodic on $P_0\mathcal{H}_{\mathrm{act}}$, then

$$\boxed{\lim_{g\to 0}\lim_{t\to\infty}\rho_A^{(g)}(t) \;=\; \frac{P_0}{\mathfrak{g}}, \qquad S_0 \;=\; \log\mathfrak{g}}$$

so $S_0 = 0$ iff the ground state is nondegenerate.

Causality and gauge. The flow preserves $\bar{\Sigma}$–lightcone support and is invariant in the Good–clock gauge class: no law–signalling, and all statements hold for any admissible emergent time variable.

15.1.6 The Principle of Asymptotic Coherence

This principle re-casts the classical Third Law as a *dynamical* statement about *how* coherence is recovered on a diagnostic realized reference configuration with Good–clock time (Definitions 13.158 and 15.1). Under endogenous coarse–graining (Assumption 15.39) and the current–kernel coupling at the A–E interface (Definition 15.25), the reduced evolution on the state–conditioned active band (Definition 15.56) is a *unital*, CP–divisible flow obtained by the product limit (Propositions 15.41 and 15.45); in the time–homogeneous case it is a unital CPTP semigroup. As the isolation parameter $g\downarrow 0$ (so that the GKLS rates $\gamma_\alpha(g)$ vanish by Assumption 15.39), we have

$$\Lambda_t^{(g)} \xrightarrow[g\to 0]{} \mathrm{Ad}_{e^{-iH_{A,\mathrm{eff}}t}},$$
$$S\Big(\Lambda_t^{(g)}\rho_A(0)\Big) \xrightarrow[g\to 0]{} S\Big(\rho_A(0)\Big),$$

uniformly on compact t–intervals. This expresses the *asymptotic coherence* content of the Third Law (Proposition 15.52 and Lemma 15.51). In this sense, the equality case of the Second Law, $\frac{d}{dt}S_A(t) \geq 0$ (Theorem 15.50 and eq. (15.14)), is *saturated* in the isolation limit; at finite coupling it holds iff $\rho_A(t)$ already lies in the record algebra and the instantaneous motion on that commutant is unitary (Corollary 15.48). All statements are invariant in the Good–clock gauge class by Lemma 15.6.

The journey to absolute zero (as relational isolation). In the macroscopic regime, "cooling'' a subsystem is best viewed as *reducing the monitoring and exchange* encoded by the current–kernel $\mathcal{U}_{\Phi\Theta}$ (Definition 15.25) and the induced pointer channel $\mathbb{E}_A$ (Definition 15.27), thereby suppressing the decohering rates in the reduced dynamics. The limiting process $g\downarrow 0$ implements precisely this *relational isolation*, rather than a merely kinetic notion of "removing motion'' [635]. Consistency with the Σ–causal architecture (Definitions 5.26 and 5.150 and Proposition 15.43) and Good–clock timing (Definition 13.158 and Lemma 15.59) is maintained throughout.

The destination (coherent limit and PQS ideal). At the boundary, the dissipator vanishes, the CP–divisible flow degenerates to a unitary group on the active band, and entropy production *flattens* to zero (Proposition 15.52). In this reduced description, the ground sector provides the closest *accessible* approximation to the solitary, maximally coherent PQS ideal emphasized in the framework; residual entropy is set by ground–space degeneracy, $S_0 = \log \mathfrak{g}$ (Corollary 15.72).

Unattainability (why the boundary is a limit). The small–coupling continuity bound (Lemma 15.51) quantifies how rapidly entropy variations collapse as $g\downarrow 0$: isolation removes the very mechanism that changes the state. With diagnostic realized reference configuration–admissible, resource–bounded control (Definition 15.58) and Planck/causal latencies (Definition 15.55 and Lemmas 15.59 and 15.61), any attempt to drive $\rho_A(t)$ to purity in finite Good–clock time would require overcoming *vanishing* dissipative power. This reproduces the standard unattainability ethos: $T \to 0$ is a *limit*, not a reachable interior point [631, 632, 633], and is formalized here by Theorem 15.65 and Corollary 15.66.

Conclusion (the dual content). Within IQR the Third Law asserts:

(i) Asymptotic coherence. $S_A(t)$ becomes constant in the isolation limit and the reduced dynamics becomes unitary (Proposition 15.52); equality in eq. (15.14) is then attained.

(ii) Unattainability. Exact isolation/purity cannot be achieved in finite emergent time with finite resources, by the small–coupling continuity and coarse–grained dynamical constraints (Lemma 15.51, Assumption 15.39, and Definitions 15.55 and 15.58); see Theorems 15.65 and 15.71.

These refine the equality case of the Second Law (Theorem 15.50) and align with resource–theoretic views of coherence and purity [628, 629, 630].

The Two Manifestations of the PQS State The treatise distinguishes two conceptually linked PQS roles:

(i) Asymptotic ideal for subsystems. The ground sector is the asymptotic fixed point of relational isolation—approached, but not reached, by any finite protocol (unattainability; Proposition 15.52, Lemma 15.51, and Theorem 15.65).

(ii) Gravitational locus of collapse. A PQS realized inside a black–hole core is not the output of laboratory cooling; it is the endpoint of gravitational collapse behind a horizon and cannot be leveraged to prepare external subsystems [86, 329, 138, 343, 328]. This respects the global constraints and the Σ–causal structure (Definition 5.150).

Why unattainability is unaffected by black holes *Causal inaccessibility.* Any PQS inside a horizon is thermodynamically unavailable to external agents; its existence does not furnish a finite–time protocol for cooling an external laboratory system to the coherent boundary [329, 343]. *No controlled preparation.* An in–falling observer does not *prepare* a PQS by finite operations; they are assimilated by the collapsing geometry [86, 138, 328]. Thus the Third Law's unattainability—as an operational statement about external control of relational subsystems—remains intact. The black–hole route illustrates a distinct, global approach to coherence, consistent with the PQS ideal and the two-level domain of applicability (Definition 15.54).

Chapter 16

Black Hole Physics

16.1 Gravitational Collapse and Black Hole Formation

The terminal evolutionary phase of a massive stellar object represents a fundamental manifestation of the competition between hydrostatic equilibrium forces and gravitational self-attraction. The end of a massive star's life is a dramatic battle where this delicate balance gives way to an irreversible gravitational cataclysm [328, 636].

Nuclear Fuel Depletion and Core Collapse

Stellar stability is maintained through hydrostatic equilibrium, wherein the outward radiation pressure generated by thermonuclear fusion in the stellar core precisely counterbalances the inward gravitational force. When a sufficiently massive star (typically with $M \gtrsim 8\,M_{\odot}$) exhausts its nuclear fuel, culminating in the formation of an inert iron core, this outward pressure ceases [637]. With its primary energy source gone, the star's core succumbs to its own immense gravity and begins to collapse catastrophically in a fraction of a second, crushing the core to unimaginable densities [638, 639].

For stellar remnants that exceed the *Tolman–Oppenheimer–Volkoff (TOV) limit*—a critical mass of roughly two to three times that of our Sun—no known force in nature can halt this final, inexorable collapse [640, 317]. The degeneracy pressure of neutrons, the quantum mechanical effect that stabilizes neutron stars, is crushed by the overwhelming gravitational force. According to the authoritative framework of Einstein's General Theory of Relativity, the collapse is absolute and unstoppable, leading to the birth of a black hole [146, 7].

The theory of General Relativity predicts two defining features for this new object. First, an *event horizon* forms, a spherical boundary in spacetime from which nothing, not even light, can escape. It is a surface of causal disconnection, forever hiding the interior from the external universe [240]. Second, the celebrated singularity theorems of Penrose and Hawking demonstrate that all the matter inside the horizon is crushed into a *gravitational singularity*, a point of zero volume and infinite spacetime curvature where the known laws of physics cease to apply [312].

The Quantum Frontier

Here, at the singularity, we reach the absolute limit of our current authoritative theories. The prediction of a physical infinity is a definitive signal that General Relativity has broken down and is no longer a complete description of reality [240, 7]. This is the domain of *Quantum Gravity*.

The resolution of the black hole singularity problem necessitates a unified framework incorporating both general relativistic and quantum mechanical principles. Contemporary theoretical approaches include:

(i) String Theory. This framework proposes that fundamental particles are one-dimensional extended objects (strings) propagating in higher-dimensional spacetime [200, 201, 198, 199]. It suggests that the classical singularity may be resolved through stringy physics effects at the Planck scale.

(ii) Loop Quantum Gravity. This approach quantizes spacetime itself, proposing a discrete, granular structure at the Planck scale [83, 272]. Within this framework, the classical singularity is replaced by a *quantum bounce*, potentially resolving the infinities predicted by classical General Relativity [339].

The interior structure of black holes remains one of the most profound unsolved problems in theoretical physics. The absence of observational access to trans-horizon physics, combined with the incomplete nature of quantum gravitational theories, renders the question of singularity resolution an active area of fundamental research. The ultimate resolution requires the successful unification of quantum mechanics and General Relativity—a goal that continues to challenge the contemporary physics community [83, 200, 201, 146].

The Black Hole Information Paradox

This prediction of a singularity signals the breakdown of General Relativity and the beginning of the quantum frontier. It is at this boundary, the event horizon, that the deepest conflict between our two great theories of reality emerges: the *Black Hole Information Paradox*.

In 1976, Stephen Hawking applied quantum field theory to the curved spacetime around a black hole and made a revolutionary discovery: black holes are not truly black. They emit a faint thermal glow, now known as *Hawking radiation* [87], causing them to slowly lose mass and evaporate over immense timescales.

Hawking's calculation, however, came with a terrifying implication. The radiation appeared to be perfectly thermal, meaning its properties depended only on the black hole's mass, charge, and spin, not on the intricate details of the matter that created it. This led to a profound paradox: as the black hole evaporates completely, all the unique information about the original star—the specific composition of its elements, the configuration of its particles—would be utterly erased from the universe.

This directly violates a sacred tenet of quantum mechanics: *unitary evolution*, which guarantees that information, while it can be scrambled, can never be destroyed [330]. The conflict is stark: General Relativity predicts a one-way door to oblivion, while quantum mechanics insists that every process must be reversible in principle. The fate of this information

has driven theoretical physics for nearly half a century, leading to further puzzles like the *firewall paradox*, which suggested that the event horizon might be a violent wall of energy rather than a smooth surface [231].

For decades, this paradox remained unresolved. However, very recent breakthroughs, inspired by the holographic principle and quantum information theory [516], have pointed toward a dramatic resolution. The key was understanding the *entanglement entropy* of the outgoing Hawking radiation. The *Page curve*, proposed by Don Page in 1993 [138], predicted how this entropy *should* behave if information is conserved: it should rise and then fall back to zero as the black hole disappears. Hawking's original calculation showed it rising indefinitely.

The modern resolution comes from the development of new mathematical tools, particularly *quantum extremal surfaces* [232] and the so-called *"island formula"* [139, 337]. These techniques reveal that as a black hole evaporates, subtle gravitational effects, known as *replica wormholes*, create a deep entanglement between the radiation far away and an "island" of spacetime deep within the black hole's interior.

This means the information is not truly lost behind the horizon. Instead, it is intricately and non-locally encoded in the quantum correlations between the emitted radiation and the interior of the black hole itself. As the final quanta of radiation are emitted, they carry this information with them, allowing the total state to return to a pure, information-complete state. The process, when viewed through this new lens, is consistent with unitary evolution.

While a complete, experimentally verified theory of quantum gravity is still needed, these advanced new theories strongly suggest that information is indeed preserved. The paradox is resolved not by a simple escape mechanism, but by recognizing that spacetime geometry and quantum entanglement are far more deeply intertwined than the classical picture ever allowed.

16.1.1 The Proposed Resolution to the Black Hole Information Paradox

The Process of Collapse: A Journey Through Regimes

The narrative of black hole formation in the IQR framework (see Section 16.1) is not a march toward an endpoint of broken physics, but a transition across the hierarchical layers of emergent reality. A massive star begins in the **Cosmological** and **Macroscopic (IR)** regimes, where its information is encoded in the local state of its constituent matter—the *kymons*—as excitations of the *Tantrakyma field* governed by the action $S_\Theta[\hat{\mathbf{\Theta}}]$ (Definition 9.310). As gravitational collapse proceeds, the stellar core is driven "downward" through the chronogeometric cascade: *Macroscopic* ($\gg \ell_P$) $\to$ *Relational/Mesoscopic* ($\gtrsim \ell_{\text{PQS}}^{-1}$) $\to$ *Planckian* ($\sim \ell_P$) $\to$ *Ultra-sub-Planckian* ($\ll \ell_P$), where spacetime as a smooth manifold dissolves into quantum-relational structure.

Within IQR, the classical "singularity" is not an ill-defined end of law, but a *phase transition* of ontology. Beyond Planck densities, the emergent chronogeometric manifold ceases to be the correct description; it *reverts* to its pre-geometric substrate. The collapse terminates in a stable, information-rich configuration of *Primordial Quantum Singularities (PQSs)* (Section 1.3). Thus the black-hole core is not a pathological point, but a tightly correlated PQS network existing in the *Ultra-sub-Planckian* regime, where the deeper, quantum-relational laws from which spacetime emerges become the appropriate degrees of freedom.

The Fate of Information: Transformation, Not Destruction

The central question of the Black Hole Information Paradox is the fate of the star's microscopic information [329, 330, 138]. In the IQR framework, information is preserved by unitary dynamics: it is neither destroyed nor duplicated, but *transformed* from a local, material encoding into a non-local, geometric-relational encoding.

Dynamically, this transformation is governed by the *Total Relational Action* S_{total} (Definition 9.357; cf. Section 15.1.3), and, in particular, by its *Interaction Action* term, S_{int}. The latter is intrinsically bi-local and mediates matter–geometry exchange through the nonlocal interaction kernel:

$$\mathcal{L}_{\text{int}}(x, x') \;:=\; \mathrm{Tr}\Big[\hat{\mathbf{\Phi}}^{\dagger}(x)\,\mathcal{U}_{\Phi\Theta}(x, x')\,\hat{\mathbf{\Theta}}(x')\Big]\,, \tag{16.1}$$

as formally defined in Definitions 10.222 and 10.223. During collapse, the information initially stored locally in the Tantrakyma sector is *imprinted* onto the nonlocal correlational structure of the *Gaiamiti field*. Concretely, the interacting PQS core functions as an entanglement register whose interior degrees of freedom are bi-locally coupled to exterior modes; the star's microstate is thereby re-expressed within the entanglement structure of the PQS network.

This picture coherently integrates with the broader quantum-gravitational understanding that black-hole evaporation can be unitary when interior degrees of freedom are nonlocally encoded in correlations accessible to the radiation at late times—as captured heuristically by Hayden–Preskill scrambling [336] and quantitatively by Page-curve analyses and the island formula [335, 337, 139, 232]. In IQR, that nonlocal encoding is not an *ad hoc* addition but a direct consequence of the interaction structure in Definitions 10.222 and 10.223, together with the PQS ontology (Section 1.3). Hence the black hole does not erase information; it *re-codes* it into the geometry–matter entanglement of the PQS core and, through unitary dynamics, releases it in a purified final state as evaporation completes.

The Dual Nature of Hawking Radiation

The inclusion of the *Laws of Thermodynamics* (Section 15.1) is not a contradiction but a key to the correct reading of black-hole emission within the IQR framework. The radiation must be understood from two complementary, yet fully consistent, perspectives.

1. The Local Perspective: An Apparently Thermal Bath

Corrected surface gravity on $\bar{\Sigma}$. On the unified static ansatz, let

$$F(r) := 1 - \frac{f(r)}{r} - A\Big(\xi(r)\Big), \qquad G(r) := 1 - \frac{f(r)}{r} - B\Big(\xi(r)\Big).$$

For the horizon generator $\chi^a = \partial_t$, the effective surface gravity at the outer root R_{EH} is

$$\kappa_{\text{eff}} = \frac{c^2}{2}\sqrt{F'(R_{\text{EH}})\,G'(R_{\text{EH}})}\,, \tag{16.2}$$

which reduces to $\kappa_S = c^2/(2R_S)$ when $A = B = 0$ and $f(r) = R_S$. The local KMS temperature is then

$$T_{\text{eff}} = \frac{\hbar\,\kappa_{\text{eff}}}{2\pi k_B}\,, \qquad \text{so that } \beta_H = \frac{2\pi}{\kappa_{\text{eff}}}. \tag{16.3}$$

Linearization about $R_S := 2GM/c^2$. Writing $R_{\rm EH} = R_S + \delta R_H$ and $f(r) = R_S + f_1(r)$ to first order, with $A_\xi := \frac{dA}{d\xi}$, $B_\xi := \frac{dB}{d\xi}$, and $\xi'(R_S) := \frac{d\xi}{dr}\big|_{R_S}$, one finds

$$\boxed{\kappa_{\rm eff} = \frac{c^2}{2R_S}\left[1 - \frac{2\,\delta R_H}{R_S} - \left(f_1'(R_S) - \frac{f_1(R_S)}{R_S}\right) - \frac{R_S}{2}\left(A_\xi + B_\xi\right)\xi'(R_S)\right] + \mathcal{O}(\text{quadratic})} \tag{16.4}$$

and using $\delta R_H = f_1(R_S) - R_S\, a_1(R_S)$ (with $a_1(R_S) := A(\xi(R_S))$) expresses $\kappa_{\rm eff}$ entirely in terms of horizon-value data. In the "isotropic" gauge $A = B$, (16.2) simplifies to $\kappa_{\rm eff} = \frac{c^2}{2}F'(R_{\rm EH})$ and (16.4) holds with $A_\xi + B_\xi \to 2A_\xi$.

For any static observer in the exterior region (e.g. $r \gg R_S$, cf. Section 16.1), measurements restricted to local observables yield an *apparently thermal* spectrum at the Hawking temperature. Formally, the exterior algebra, when restricted to a stationary black-hole background with Killing generator χ^a, is in a KMS state with respect to the horizon flow α_t at inverse temperature $\beta_H = 2\pi/\kappa_{\rm eff}$ (with $\kappa_{\rm eff}$ the surface gravity of the Killing horizon on $\bar{\Sigma}$; see Section 16.1.1 for definitions). Consequently, an Unruh–DeWitt detector exhibits a Planckian response with

$$T_H \;=\; \frac{\hbar\,\kappa_{\rm eff}}{2\pi k_B} \xrightarrow[A=B=0]{\text{Schwarzschild}} \frac{\hbar c^3}{8\pi G M k_B}\,. \tag{16.5}$$

in line with standard derivations via Bogoliubov coefficients, Euclidean periodicity, and algebraic KMS characterizations [87, 641, 314, 245, 230, 241, 381].

Why, then, does the radiation *look* thermal?

(i) Restricted access and decoherence. As established in our derivation of the Second Law (Section 15.1.4), a local observer coarse-grains over inaccessible degrees of freedom (interior modes, long-range correlations, and previously emitted quanta). Tracing out these sectors induces *environment-assisted decoherence*, suppressing phase relations and yielding an effectively mixed state for the detected quantum [170, 247]. In the algebraic setting, this is precisely the passage from the global state to its restriction on the exterior observable net, where the KMS property supplies thermal statistics [230, 241].

(ii) Thermal state as maximal ignorance. The reduced density matrix of a single, detected quantum is, by construction, a mixed state that is statistically indistinguishable from a Gibbs/KMS state at T_H with respect to the exterior time flow. Thermodynamic laws thus apply *exactly* to this local, coarse-grained description because those laws govern systems under incomplete information (cf. Sections 15.1 and 15.1.4).

Therefore, *from the standpoint of any local observer lacking access to the complete global state*, Hawking radiation *is* thermal, carries a well-defined temperature (16.5), and increases coarse-grained entropy in accordance with the Second Law (Section 15.1.4). This local thermality coexists with global unitarity, whose complementary perspective will be addressed next (Section 16.1.1).

2. The Global Perspective: A Unitary, Information-Carrying State

From the perspective of the *entire* universe—i.e. the global *Quantum Holo-State* $\left|\mathring{\boldsymbol{\Psi}}\right\rangle$ subject to the Universal Quantum Constraint,

$$\hat{H}_{\text{univ}} \left|\mathring{\boldsymbol{\Psi}}\right\rangle = 0, \tag{16.6}$$

$$\hat{H}_{\text{univ}} = \hat{H}_{\Phi} + \hat{H}_{\Theta} + \hat{H}_{\text{int}}. \tag{16.7}$$

See Theorem 13.168 and Definitions 9.357 and 13.167. Black-hole formation and evaporation proceed by *unitary* dynamics. The apparent tension with thermality in the local picture (Section 16.1.1) dissolves once we refrain from tracing out inaccessible degrees of freedom.

(i) Information is never lost (global encoding). Microstate information of infalling matter (kymonic excitations of the Tantrakyma sector, $\hat{\boldsymbol{\Theta}}$; Definition 9.310) is transferred, via the bi-local interaction structure,

$$\mathcal{L}_{\text{int}}(x, x') \;=\; \text{Tr}\Big[\hat{\boldsymbol{\Phi}}^{\dagger}(x)\, \mathcal{U}_{\Phi\Theta}(x, x')\, \hat{\boldsymbol{\Theta}}(x')\Big],$$

to nonlocal, geometry–matter correlations (Definitions 10.222 and 10.223). In the late stages of collapse this encoding is concentrated within the correlated *PQS network* at the core (Section 1.3), and as evaporation proceeds it is *systematically re-expressed* in long-range entanglement among the outgoing quanta. This mechanism is consistent with modern unitary accounts of evaporation: the Page-curve prediction that the radiation entropy rises until the Page time and then decreases to zero [335, 138], fast scrambling of black holes [336], and the island/QES framework in which the entanglement wedge of the radiation eventually includes interior degrees of freedom [232, 337, 139]. In IQR, these facts are not added by hand but follow from the universal interaction term and the PQS ontology (Definition 10.223 and Section 1.3).

(ii) The total state is pure (fine-grained unitarity). Let $\rho_{\text{global}} := \left|\mathring{\boldsymbol{\Psi}}\right\rangle\!\left\langle\mathring{\boldsymbol{\Psi}}\right|$ be the density operator on the holocosmic Hilbert space (cf. Theorem 13.168). Then the fine-grained von Neumann entropy satisfies $S(\rho_{\text{global}}) = 0$ at all times, even though any *reduced* state (e.g. radiation alone, exterior algebra alone) is mixed and approximately thermal (Section 16.1.1). Thus, "entropy increase" pertains to *coarse-grained*, subsystem descriptions (see Sections 15.1 and 15.1.4), while the full evolution remains unitary [287]. In geometric language: the entanglement wedge of the radiation grows to capture the interior (the island), ensuring that late-time radiation purifies early radiation and restoring a unitary Page curve [337, 139, 232].

Conclusion. The global IQR account is therefore: information is *re-coded*, not destroyed—first into the nonlocal PQS correlations during collapse, and subsequently into the multipartite entanglement structure of the outgoing radiation during evaporation (cf. Section 16.1.1). Local thermality and global unitarity are two consistent projections of the same universal quantum process.

Evaporation and Unitary Information Release

The evaporation rate is governed by the Hawking temperature, which scales inversely with mass (cf. eq. (16.5)):

$$T_H = \frac{\hbar c^3}{8\pi G M k_B} \quad \text{(Schwarzschild)},$$

as first derived in [87] and used in the emission-rate analysis of [642]. Two immediate consequences follow:

(i) Massive black holes are extremely cold. They radiate only very feebly.

(ii) Only very small black holes are hot. Such black holes evaporate rapidly.

Including graybody factors, the luminosity scales as $L \propto A\,T_H^4 \sim M^{-2}$, so the mass-loss rate satisfies $\dot{M} \propto -M^{-2}$ (neutral, non-rotating case) [642]. Integrating yields the standard lifetime,

$$t_{\text{evap}} \simeq \frac{5120\,\pi\,G^2}{\hbar c^4}\,M^3 \quad \text{(up to } \mathcal{O}(1) \text{ graybody/species corrections; cf. [642])}. \tag{16.8}$$

For a solar-mass black hole this timescale exceeds the age of the universe by many orders of magnitude, whereas sub-asteroidal primordial black holes could complete evaporation within cosmological time depending on their initial mass [642]. The *Page time*—the epoch when the fine-grained entropy of the radiation peaks—also scales as $t_{\text{Page}} \sim \mathcal{O}(M^3)$ [335, 138]; after this point, late-time quanta purify the early radiation in a unitary Page curve (Section 16.1.1).

Within the IQR framework, evaporation is the slow, *unitary* evolution of the PQS-core configuration driven by the Total Relational Action (Definitions 9.357 and 10.223). The outgoing quanta are the fundamental excitations of the two sectors: *Apeirons* of the Gaiamiti field and *Kymons* of the Tantrakyma field (cf. Definitions 6.174 and 9.310). While any *local* detector registers an approximately thermal spectrum and a coarse-grained entropy increase (Sections 15.1, 15.1.4 and 16.1.1), the *global* state of "PQS core + radiation'' remains pure and information-carrying (Section 16.1.1). Microscopically, information initially encoded in the Tantrakyma microstate is transferred, via the bi-local interaction kernel $\mathcal{L}_{\text{int}}(x, x')$ (Definition 10.222), into non-local geometry–matter correlations and then into multipartite entanglement among the emitted quanta, ensuring conservation of information at the universal level.

The Thermodynamic Nature of Black Holes: A Tale of Two Extremes

In the IQR framework, the thermodynamic behaviour of black holes is grounded in the principle that *temperature quantifies the rate of relational decoherence experienced by a subsystem due to entanglement with the rest of the universe* (cf. Sections 15.1, 15.1.4 and 15.1.5). This principle reconciles the two complementary perspectives on Hawking emission (Section 16.1.1) and explains the behaviour of black holes at both ends of the mass spectrum within standard black-hole thermodynamics [86, 87, 314, 245].

The Massive Black Hole: A Thermodynamic Ground State

For a stationary, neutral black hole the Hawking temperature scales inversely with mass,

$$T_H = \frac{\hbar\,\kappa}{2\pi k_B} = \frac{\hbar c^3}{8\pi G M k_B} \quad \text{(Schwarzschild)}, \tag{16.9}$$

as established in [87, 314] (cf. eq. (16.5)). Thus, as $M \to \infty$, $T_H \to 0$: the macroscopic object approaches the ideal envisioned by the Third Law (Section 15.1.5). In the language of this treatise, the near-zero temperature signifies an exceedingly small rate of information exchange with the environment, i.e. near-perfect coherence under the exterior time flow (Section 16.1.1). Evaporation—mediated by extremely weak emission—is correspondingly slow (Section 16.1.1).

Ontological reading. The extreme coldness of a massive black hole is the macroscopic thermodynamic signature of the coherence of the *Primordial Quantum Singularity (PQS)* network at its core (Section 1.3). In this regime, leakage of excitations—*Apeirons* (Gaiamiti sector) and *Kymons* (Tantrakyma sector)—is strongly suppressed (cf. Definitions 9.246 and 9.310), reflecting the tenuous coupling to exterior modes encoded in the interaction kernel (Definitions 10.222 and 10.223).

The Small Black Hole: Relational Instability and Rapid Evaporation

At the opposite extreme, a small black hole has a high Hawking temperature $T_H \propto M^{-1}$ and correspondingly large luminosity once graybody factors are included, $L \sim M^{-2}$ [642]. Physically:

(i) The steep curvature near a tiny horizon sharpens the interface between the *PQS regime* and the exterior *Macroscopic (IR) regime* (Section 16.1).

(ii) This steep interface enhances the flow of conserved relational currents generated by symmetries of the Total Relational Action (see Section 15.1.3 and Definition 9.357).

(iii) Macroscopically, this manifests as rapid emission of *Apeirons* and *Kymons* (Hawking quanta), i.e. an enhanced relational decoherence rate for the *subsystem* description, even as the *global* evolution remains unitary (Section 16.1.1).

Runaway feedback (mass loss). With $\dot{M} \propto -M^{-2}$ for a neutral, non-rotating hole, the lifetime scales as

$$t_{\text{evap}} \simeq \frac{5120\,\pi\,G^2}{\hbar c^4}\,M^3 \quad \text{(up to graybody/species corrections)}, \tag{16.10}$$

so mass loss accelerates as M decreases [642]. The final stages require a UV-complete description; within IQR they are governed by the same interaction structure that encodes information nonlocally (Definitions 10.222 and 10.223).

The Completion of Black Hole Evaporation

At completion, the outgoing radiation forms a single, fine-grained *pure* state that is unitarily related to the initial pure state of the collapsing star (Section 16.1.1). The dynamics is captured by the *Generalized Chronogeometric Einstein Equations* (eq. (11.191)), closing the feedback loop between matter and geometry, and by the *Total Relational Action* (Definition 9.357) that mediates bi-local exchange between Tantrakyma and Gaiamiti sectors (Definitions 10.222 and 10.223). Hence the "information paradox" reflects a semiclassical coarse-graining: locally thermal statistics (Section 16.1.1) coexist with global unitarity, because information is re-coded from (i) local material degrees of freedom to (ii) nonlocal PQS correlations, and finally to (iii) multipartite entanglement among the emitted quanta, consistent with Page-curve expectations [335, 138].

16.1.2 The Field Equations for Black Holes

Action and variational principle. Our starting point is the *Relational Einstein–Hilbert Action* for the unified chrono–geometric tensor $\bar{\Sigma}_{\mu\nu}$ (cf. Definition 11.105 and eq. (16.18)):

$$S_{\text{Relational}}[\bar{\Sigma}] := \frac{1}{2\kappa}\int_{\mathcal{X}_q} d^4x\,\sqrt{-\det\bar{\Sigma}}\;\bar{R}[\bar{\Sigma}], \qquad \kappa = \frac{8\pi G}{c^4}. \tag{16.11}$$

A well-posed variational problem on a region with boundary tacitly includes the Gibbons–Hawking–York term [314, 7].

Unified vacuum field equation. Varying (16.11) with respect to $\bar{\Sigma}_{\mu\nu}$ and imposing $\delta S_{\text{Relational}} = 0$ yields

$$\bar{G}_{\mu\nu}[\bar{\Sigma}] \;=\; 0, \qquad \bar{G}_{\mu\nu}[\bar{\Sigma}] := \bar{R}_{\mu\nu}[\bar{\Sigma}] - \tfrac{1}{2}\,\bar{\Sigma}_{\mu\nu}\,\bar{R}[\bar{\Sigma}] \quad (\text{cf. } \textit{Definition } 11.138), \tag{16.12}$$

with the contracted Bianchi identity $\bar{\nabla}^{\mu}\bar{G}_{\mu\nu} = 0$ following from diffeomorphism invariance [7].

Sourced (semi-classical) form. When expectation stresses from the Gaiamiti/Tantrakyma sectors and an emergent cosmological term are present, we obtain the *Generalized Chrono–geometric Einstein Equations* (cf. eq. (11.191)):

$$\bar{G}_{\mu\nu}[\bar{\Sigma}] + \Lambda\,\bar{\Sigma}_{\mu\nu} \;=\; \kappa\Big(\langle\hat{T}^{(\Phi)}_{\mu\nu}\rangle + \langle\hat{T}^{(\Theta)}_{\mu\nu}\rangle\Big), \tag{16.13}$$

with $\bar{\nabla}^{\mu}(\langle\hat{T}^{(\Phi)}_{\mu\nu}\rangle + \langle\hat{T}^{(\Theta)}_{\mu\nu}\rangle) = 0$ ensured by $\bar{\nabla}^{\mu}\bar{G}_{\mu\nu} = 0$ (see also Section 15.1 for the thermodynamic reading of these expectations).

Symmetry reduction for black holes (self-contained definitions). To specialize to stationary, spherically symmetric configurations, we adopt the diagonal ansatz

$$\bar{\Sigma}_{\mu\nu}dx^{\mu}dx^{\nu} = -\Big(1 - \tfrac{2Gm(r)}{c^2 r} - A(r)\Big)c^2dt^2 + \frac{dr^2}{1 - \tfrac{2Gm(r)}{c^2 r} - B(r)} + r^2d\Omega^2, \tag{16.14}$$

where $m(r)$ is the *mass function* and $A(r), B(r)$ parameterize temporal-sector corrections (cf. eqs. (16.34) and (16.35)). For later convenience we introduce the mass–length function

$$f(r) \;:=\; \frac{2G\,m(r)}{c^2}, \tag{16.15}$$

so that (16.14) is the f-form of the unified line element used in eq. (16.18). Inserting (16.14) into (16.12) (or (16.13) when vacuum polarization is included; cf. Section 16.1.17) yields a coupled ODE system for the unknowns $\{m(r), A(r), B(r)\}$ (see Section 16.1.7 and eqs. (16.56) to (16.58)). Asymptotic flatness imposes

$$A(r), B(r) \to 0, \qquad m(r) \to M \quad (r \to \infty),$$

equivalently $f(r) \to 2GM/c^2$ (eqs. (16.59) and (16.60)).

Event horizon (phase boundary). The event horizon R_{EH} is the largest root of the unified lapse in (16.14); equivalently,

$$1 - \frac{2GM}{c^2 R_{\mathrm{EH}}} - A(R_{\mathrm{EH}}) = 0 \quad \text{(cf. the phase-boundary discussion in } \textit{Section } 16.1.10\text{)}. \tag{16.16}$$

In terms of f this reads $1-\frac{f(R_{\mathrm{EH}})}{R_{\mathrm{EH}}}-A(\xi(R_{\mathrm{EH}}))=0$ (cf. eq. (16.21)), reducing to $R_{\mathrm{EH}} = 2GM/c^2$ when $A \to 0$.

Objective. Equations (16.12)–(16.16) define the black-hole boundary–value problem for $\bar{\Sigma}_{\mu\nu}$. The task is to solve consistently for $\{A(r), B(r)\}$ and $m(r)$ (or $f(r)$) subject to the above boundary data, in harmony with thermodynamic constraints (Section 16.1.1) and information-theoretic requirements (Section 16.1.1). Standard variational geometry [146, 7] ensures structural consistency, while the IQR framework specifies sourcing and near-horizon behaviour via regulated vacuum energy and relational interactions (cf. Section 16.1.17 and Definitions 10.222 and 10.223).

16.1.3 Modeling the Emergent Geometry

To construct black–hole solutions of the field equations (Section 16.1.2; cf. the unified vacuum form in Definition 11.138), we posit a spherically symmetric ansatz for the unified chrono–geometric tensor $\bar{\Sigma}_{\mu\nu}$ that cleanly tracks the transition across physical regimes while preserving the area–radius gauge $g_{\theta\theta} = r^2$ [146, 7].

Regime tracker. Introduce a *regime parameter* $\xi(r)$ (monotone inwards) that measures proximity to the PQS core (Section 1.3):

$$\lim_{r\to\infty} \xi(r) = 0, \qquad \lim_{r\to 0^+} \xi(r) = +\infty, \qquad \frac{d\xi}{dr} < 0. \tag{16.17}$$

Heuristically, ξ labels the *Macroscopic* ($\xi \approx 0$), *Transition* (near the horizon), and *PQS* ($\xi \to \infty$) regimes discussed in Section 16.1.

Unified line element and sector split. With $f(r)$ a mass–length function, the unified line element is written as a geometric block plus a temporal (phase–flow) correction:

$$\bar{\Sigma}_{\mu\nu}dx^\mu dx^\nu = \underbrace{-\Big(1-\frac{f(r)}{r}\Big)c^2dt^2+\frac{dr^2}{1-\frac{f(r)}{r}}+r^2d\Omega^2}_{\text{Geometric sector }(\bar{g}_{\mu\nu})} + \underbrace{\mathcal{T}_{\mu\nu}(r,\xi(r))\,dx^\mu dx^\nu}_{\text{Temporal sector }(\bar{t}_{\mu\nu})}. \tag{16.18}$$

We choose the area–radius gauge so that $\mathcal{T}_{\theta\theta}=\mathcal{T}_{\phi\phi}=0$. The function $f(r)$ is related to the enclosed mass by $f(r)=2G\,m(r)/c^2$ (cf. eq. (16.15)), reducing to the Schwarzschild value $f(r)\to 2GM/c^2$ at large r.

Diagonal temporal corrections and consistency with the BH ansatz. For stationary configurations the temporal sector is taken diagonal,

$$\mathcal{T}_{\mu\nu}(r,\xi)\,dx^\mu dx^\nu = +A(\xi)\,c^2dt^2+\frac{B(\xi)}{\Big(1-\frac{f(r)}{r}\Big)\Big(1-\frac{f(r)}{r}-B(\xi)\Big)}\,dr^2, \tag{16.19}$$

so that the *total* line element becomes

$$\bar{\Sigma}_{\mu\nu}dx^\mu dx^\nu = -\Big(1-\tfrac{f(r)}{r}-A(\xi)\Big)c^2dt^2+\frac{dr^2}{1-\frac{f(r)}{r}-B(\xi)}+r^2d\Omega^2, \tag{16.20}$$

in manifest correspondence with the black–hole ansatz employed in eq. (16.14). The horizon R_{EH} is the largest root of the lapse factor, i.e.

$$1-\frac{f(R_{\mathrm{EH}})}{R_{\mathrm{EH}}}-A\Big(\xi(R_{\mathrm{EH}})\Big)=0, \tag{16.21}$$

which matches the phase–boundary condition in eq. (16.16).

Boundary/regularity conditions and regime behaviour. The regime tracker enforces the asymptotics

$$A(\xi)\xrightarrow[\xi\to 0]{}0,\qquad B(\xi)\xrightarrow[\xi\to 0]{}0,\qquad A(\xi),B(\xi)\text{ bounded for all }\xi<\infty, \tag{16.22}$$

so that $\bar{\Sigma}_{\mu\nu}\to\bar{g}_{\mu\nu}$ at infinity (asymptotic flatness). Near the horizon, $A,B=\mathcal{O}(1)$ encode nontrivial chrono–geometric structure; deep in the PQS regime, the temporal block dominates, signaling breakdown of the purely geometric description in favour of the operator–level ontology (Section 1.3).

Dynamical closure. Inserting (16.20) into the field equations (Section 16.1.2) yields a coupled ODE system for $m(r)$, $A(\xi(r))$, and $B(\xi(r))$. In vacuum one solves $\bar{G}_{\mu\nu}[\bar{\Sigma}]=0$ (cf. Definition 11.138); with quantum expectation stresses or vacuum polarization, one uses the sourced form (eq. (16.13)) with inputs constrained by the regulated vacuum sector (Section 16.1.17). Standard results on spherically symmetric reductions provide consistency checks and limits [146, 7].

16.1.4 The Mathematical Path Forward: From Ansatz to Solution

Our goal is to solve the *Unified Vacuum Field Equation* for the chrono–geometric tensor,

$$\bar{G}_{\mu\nu}[\bar{\Sigma}] := \bar{R}_{\mu\nu}[\bar{\Sigma}] - \tfrac{1}{2}\,\bar{\Sigma}_{\mu\nu}\,\bar{R}[\bar{\Sigma}] = 0, \tag{16.23}$$

as obtained from the variational principle in Section 16.1.2 (cf. Definition 11.138). We proceed with the spherically symmetric unified line element introduced in Section 16.1.3,

$$\bar{\Sigma}_{\mu\nu}dx^\mu dx^\nu = -\Big(1 - \tfrac{f(r)}{r} - A(\xi)\Big)c^2dt^2 + \frac{dr^2}{1 - \frac{f(r)}{r} - B(\xi)} + r^2d\Omega^2, \tag{16.24}$$

where $f(r) = 2Gm(r)/c^2$ and $A(\xi), B(\xi)$ are temporal-sector profiles (eqs. (16.17), (16.20) and (16.22)).

Step 1: Relational Christoffel symbols $\Gamma^\lambda{}_{\mu\nu}[\bar{\Sigma}]$

Write the sector split $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ with $\bar{t}_{\mu\nu}$ small away from the horizon. Let ∇ denote the Levi–Civita connection of $\bar{g}$. The exact Christoffel symbols are

$$\Gamma^\lambda{}_{\mu\nu}[\bar{\Sigma}] = \tfrac{1}{2}\,\bar{\Sigma}^{\lambda\rho}\Big(\partial_\mu\bar{\Sigma}_{\nu\rho} + \partial_\nu\bar{\Sigma}_{\mu\rho} - \partial_\rho\bar{\Sigma}_{\mu\nu}\Big), \tag{16.25}$$

and admit the (covariant) decomposition

$$\Gamma^\lambda{}_{\mu\nu}[\bar{\Sigma}] = \Gamma^\lambda{}_{\mu\nu}[\bar{g}] + \Delta\Gamma^\lambda{}_{\mu\nu}[\bar{t}] \quad \text{with} \quad \Delta\Gamma^\lambda{}_{\mu\nu} = \tfrac{1}{2}\,\bar{g}^{\lambda\rho}\Big(\nabla_\mu\bar{t}_{\nu\rho} + \nabla_\nu\bar{t}_{\mu\rho} - \nabla_\rho\bar{t}_{\mu\nu}\Big) + \mathcal{O}(\bar{t}^2). \tag{16.26}$$

Formulas (16.25)–(16.26) are standard (see, e.g., [7, §3–4]; [146, Ch. 12]). For the ansatz (16.24), only $\partial_r f$, $\partial_r A$, and $\partial_r B$ appear, so the nonzero components reduce to rational combinations of f, A, B and their first derivatives.

Step 2: Unified Ricci tensor $\bar{R}_{\mu\nu}[\bar{\Sigma}]$

The Ricci tensor of $\bar{\Sigma}$ may be expanded about $\bar{g}$ as

$$\bar{R}_{\mu\nu}[\bar{\Sigma}] = \bar{R}_{\mu\nu}[\bar{g}] + \Delta\bar{R}_{\mu\nu}[\bar{t}] + \mathcal{Q}_{\mu\nu}[\bar{t}, \bar{t}], \tag{16.27}$$

where to first order in $\bar{t}$ (indices moved with $\bar{g}$)

$$\Delta\bar{R}_{\mu\nu}[\bar{t}] = \tfrac{1}{2}\Big(-\nabla^2\bar{t}_{\mu\nu} - \nabla_\mu\nabla_\nu\bar{t} + \nabla_\mu\nabla^\rho\bar{t}_{\nu\rho} + \nabla_\nu\nabla^\rho\bar{t}_{\mu\rho}\Big) + \bar{R}_{\mu\rho\nu\sigma}[\bar{g}]\,\bar{t}^{\rho\sigma} - \bar{R}_{\rho(\mu}[\bar{g}]\,\bar{t}^{\rho}{}_{\nu)}, \tag{16.28}$$

and $\mathcal{Q}_{\mu\nu}$ collects quadratic terms from $\Delta\Gamma\cdot\Delta\Gamma$ (see [7, Eq. (B.7)]; [146, §35.1]). For the static, spherically symmetric line element (16.24), $\bar{R}_{\mu\nu}[\bar{g}]$ reproduces the usual Schwarzschild expressions when $A = B = 0$, while $\Delta\bar{R}_{\mu\nu}$ introduces combinations of A', B', f', f''.

Step 3: Coupled field equations (ODE system)

Substituting (16.26)–(16.27) and (16.24) into (16.23) yields three independent ordinary differential equations (the $\theta\theta$ component is redundant by the contracted Bianchi identity):

$$\bar{G}^{t}{}_{t} = 0 \implies f'(r) = \mathcal{F}_{tt}(r; f, A, B, A', B'), \tag{16.29}$$

$$\bar{G}^{r}{}_{r} = 0 \implies B'(r) = \mathcal{F}_{rr}(r; f, A, B, A', f'), \tag{16.30}$$

$$\bar{G}^{\theta}{}_{\theta} = 0 \implies A'(r) = \mathcal{F}_{\theta\theta}(r; f, A, B, f', B'), \tag{16.31}$$

where $\mathcal{F}_{\cdot}$ are rational functions determined by (16.28) specialized to (16.24). In the sourced case one replaces (16.23) by the generalized equations (eq. (16.13)), with right-hand sides fixed by expectation stresses (cf. Section 16.1.17).

Boundary data and horizon regularity. Asymptotic flatness requires $A, B \to 0$ and $f(r) \to 2GM/c^2$ as $r \to \infty$ (eq. (16.22)). The event horizon R_{EH} is the largest positive root of the lapse (phase–boundary condition)

$$1 - \frac{f(R_{\mathrm{EH}})}{R_{\mathrm{EH}}} - A\Big(\xi(R_{\mathrm{EH}})\Big) = 0 \quad \text{(eqs. (16.16) and (16.21)).} \tag{16.32}$$

Regularity of invariants (e.g. the Kretschmann scalar) and finiteness of tidal forces in an ingoing Eddington–Finkelstein chart fix relations among A, B and their derivatives evaluated at R_{EH} [146, §31–33]; these serve as inner boundary conditions for (16.29)–(16.31).

Solution strategy (controlled expansions).

(i) Asymptotic exterior. Set $A = B = 0$ to leading order, solve $f(r) \to 2GM/c^2$, and construct a $1/r$ expansion for (A, B) from (16.29)–(16.31) consistent with eq. (16.22).

(ii) Near-horizon matching. Expand $f(r) = f_0 + f_1(r - R_{\mathrm{EH}}) + \cdots$, $A(r) = A_0 + A_1(r - R_{\mathrm{EH}}) + \cdots$, $B(r) = B_0 + B_1(r - R_{\mathrm{EH}}) + \cdots$, insert into (16.29)–(16.31), and use (16.32) to fix $f_0/R_{\mathrm{EH}} + A_0 = 1$. Regularity yields linear relations among f_1, A_1, B_1.

(iii) Interior control (PQS regime). Track the growth of A, B with $\xi(r)$ (eq. (16.17)); if sourcing by regulated vacuum energy is included, couple to $\rho_{\mathrm{vac}}(r)$ as in Section 16.1.17 and integrate inward from the horizon.

This program delivers a consistent black–hole solution for $\bar{\Sigma}_{\mu\nu}$ that reduces to Schwarzschild when $A = B = 0$ and incorporates chrono–geometric corrections otherwise. Structural identities and conservation laws (Bianchi identity; $\bar{\nabla}^{\mu}\bar{G}_{\mu\nu} = 0$) furnish nontrivial checks of the integration [7, 146].

16.1.5 Step 1: Calculating the Relational Christoffel Symbols

We begin from the unified, spherically symmetric line element (Section 16.1.3)

$$\bar{\Sigma}_{\mu\nu}dx^{\mu}dx^{\nu} = \underbrace{-\left(1 - \tfrac{f(r)}{r}\right)c^2dt^2 + \frac{dr^2}{1 - \frac{f(r)}{r}} + r^2d\Omega^2}_{\text{Geometric sector } (\bar{g}_{\mu\nu})} + \underbrace{\mathcal{T}_{\mu\nu}(r, \xi(r))\,dx^{\mu}dx^{\nu}}_{\text{Temporal sector } (\bar{t}_{\mu\nu})}, \tag{16.33}$$

with $f(r) = 2Gm(r)/c^2$ (cf. eq. (16.15)) and $\xi(r)$ the inward–monotone regime tracker (eq. (16.17)). Following eq. (16.19), we assume a diagonal temporal block,

$$\mathcal{T}_{tt} = A(r)c^2, \qquad \mathcal{T}_{rr} = B(r), \qquad \mathcal{T}_{\theta\theta} = \mathcal{T}_{\phi\phi} = 0, \tag{16.34}$$

so that the total components read

$$\bar{\Sigma}_{tt} = -c^2\Big(1 - \tfrac{f}{r} - A\Big), \qquad \bar{\Sigma}_{rr} = \frac{1}{1 - \frac{f}{r}} + B, \tag{16.35}$$
$$\bar{\Sigma}_{\theta\theta} = r^2, \qquad \bar{\Sigma}_{\phi\phi} = r^2 \sin^2\theta,$$

and the inverse (since the metric is diagonal) is

$$\bar{\Sigma}^{tt} = -\frac{1}{c^2\Big(1 - \frac{f}{r} - A\Big)}, \qquad \bar{\Sigma}^{rr} = \frac{1}{\frac{1}{1-\frac{f}{r}} + B}, \qquad \bar{\Sigma}^{\theta\theta} = \frac{1}{r^2}, \qquad \bar{\Sigma}^{\phi\phi} = \frac{1}{r^2 \sin^2\theta}. \tag{16.36}$$

Definition. The Christoffel symbols of the Levi–Civita connection of $\bar{\Sigma}$ are

$$\Gamma^{\lambda}{}_{\mu\nu}[\bar{\Sigma}] = \frac{1}{2}\,\bar{\Sigma}^{\lambda\rho}\Big(\partial_\mu \bar{\Sigma}_{\nu\rho} + \partial_\nu \bar{\Sigma}_{\mu\rho} - \partial_\rho \bar{\Sigma}_{\mu\nu}\Big), \tag{16.37}$$

a standard construction in differential geometry [7, 146, §3–4]. We denote $g'(r) = \frac{dg}{dr}$.

Novel chrono–geometric connection components

These contain derivatives of the temporal profiles A, B and encode the relational corrections:

$$\Gamma^{r}{}_{tt} = -\tfrac{1}{2}\,\bar{\Sigma}^{rr}\,\partial_r \bar{\Sigma}_{tt} = \frac{c^2}{2\,\bar{\Sigma}_{rr}}\left(\frac{f}{r^2} - \frac{f'}{r} - A'(r)\right), \tag{16.38}$$

$$\Gamma^{t}{}_{rt} = \Gamma^{t}{}_{tr} = \tfrac{1}{2}\,\bar{\Sigma}^{tt}\,\partial_r \bar{\Sigma}_{tt} = \frac{1}{2\,\bar{\Sigma}_{tt}}\,\partial_r \bar{\Sigma}_{tt} = \frac{c^2}{2\,\bar{\Sigma}_{tt}}\left(\frac{f'}{r} - \frac{f}{r^2} + A'(r)\right), \tag{16.39}$$

$$\Gamma^{r}{}_{rr} = \tfrac{1}{2}\,\bar{\Sigma}^{rr}\,\partial_r \bar{\Sigma}_{rr} = \frac{1}{2\,\bar{\Sigma}_{rr}}\left(\frac{\frac{f'}{r} - \frac{f}{r^2}}{\Big(1 - \frac{f}{r}\Big)^2} + B'(r)\right). \tag{16.40}$$

Equations (16.38)–(16.40) follow directly from (16.37) using (16.35)–(16.36). The terms proportional to $A'(r)$ and $B'(r)$ are purely chrono–geometric and vanish in the classical limit.

Familiar geometric connection components

These coincide with the standard spherical expressions, modulated by the total radial coefficient $\bar{\Sigma}_{rr}$:

$$\Gamma^{r}{}_{\theta\theta} = -\frac{r}{\bar{\Sigma}_{rr}}, \qquad \Gamma^{r}{}_{\phi\phi} = -\frac{r\sin^2\theta}{\bar{\Sigma}_{rr}}, \tag{16.41}$$

$$\Gamma^{\theta}{}_{r\theta} = \Gamma^{\phi}{}_{r\phi} = \frac{1}{r}, \qquad \Gamma^{\theta}{}_{\phi\phi} = -\sin\theta\cos\theta, \qquad \Gamma^{\phi}{}_{\theta\phi} = \cot\theta. \tag{16.42}$$

All other components either vanish or are related by the symmetries of $\Gamma^{\lambda}{}_{\mu\nu}$.

Consistency checks and limits.

(i) Classical limit. For $A = B = 0$, $\bar{\Sigma} \to \bar{g}$ and (16.38)–(16.42) reduce to the Schwarzschild connection in standard coordinates [146, Eq. (31.2)].

(ii) Horizon behaviour. Near the event horizon, defined by the largest root of $1 - \frac{f(R_{\rm EH})}{R_{\rm EH}} - A(\xi(R_{\rm EH})) = 0$ (eqs. (16.16) and (16.21)), the combinations in (16.38)–(16.40) remain finite in an ingoing Eddington–Finkelstein chart, supplying regular inner boundary data for Step 2 (Section 16.1.4) [146, §31–33].

Physical interpretation. The connection $\Gamma[\bar{\Sigma}]$ is a superposition of (i) the standard metric-induced geometry from the *geometric sector* and (ii) novel terms generated by gradients of the *temporal sector* (A', B'), encoding the relational coherence associated with the PQS regime (Section 1.3). These chrono–geometric pieces propagate into $\bar{R}_{\mu\nu}[\bar{\Sigma}]$ (Step 2) and ultimately modify the field equations (Section 16.1.2).

16.1.6 Step 2: Calculating the Unified Ricci Tensor

Having obtained the Christoffel symbols for the chrono–geometric ansatz (Section 16.1.5, esp. eqs. (16.37) to (16.42)), we now compute the components of the *Unified Ricci Tensor* $\bar{R}_{\mu\nu}[\bar{\Sigma}]$. This supplies the explicit left–hand side of the field equations (Section 16.1.2 and eq. (16.23)), making precise how curvature is sourced jointly by the emergent geometric block $\bar{g}_{\mu\nu}$ and the temporal (phase–flow) block $\bar{t}_{\mu\nu}$.

Ricci tensor from the connection. By definition,

$$\bar{R}_{\mu\nu}[\bar{\Sigma}] = \partial_\alpha \Gamma^\alpha{}_{\mu\nu} - \partial_\nu \Gamma^\alpha{}_{\mu\alpha} + \Gamma^\alpha{}_{\alpha\lambda}\,\Gamma^\lambda{}_{\mu\nu} - \Gamma^\alpha{}_{\nu\lambda}\,\Gamma^\lambda{}_{\mu\alpha}, \tag{16.43}$$

a standard construction in differential geometry [7, 146, §3–4]. For the unified line element $\bar{\Sigma}$ in eq. (16.33) with diagonal temporal sector $\mathcal{T}$ given by eq. (16.34), all entries depend only on r, so $\partial_t = \partial_\theta = \partial_\phi = 0$.

Decomposition about the geometric sector. It is convenient to split the curvature around the geometric block $\bar{g}$ (cf. Section 16.1.4):

$$\bar{R}_{\mu\nu}[\bar{\Sigma}] = \bar{R}_{\mu\nu}[\bar{g}] + \Delta\bar{R}_{\mu\nu}[\bar{t}] + \mathcal{Q}_{\mu\nu}[\bar{t}, \bar{t}], \tag{16.44}$$

where, to first order in $\bar{t}$ (indices moved with $\bar{g}$),

$$\Delta\bar{R}_{\mu\nu}[\bar{t}] = \tfrac{1}{2}\Big(-\nabla^2\bar{t}_{\mu\nu} - \nabla_\mu\nabla_\nu\bar{t} + \nabla_\mu\nabla^\rho\bar{t}_{\nu\rho} + \nabla_\nu\nabla^\rho\bar{t}_{\mu\rho}\Big) + \bar{R}_{\mu\rho\nu\sigma}[\bar{g}]\,\bar{t}^{\rho\sigma} - \bar{R}_{\rho(\mu}[\bar{g}]\,\bar{t}^{\rho}{}_{\nu)}, \tag{16.45}$$

and $\mathcal{Q}_{\mu\nu}$ collects quadratic terms $\mathcal{O}(\bar{t}^2)$ [7, Eq. (B.7)]. Equations (16.44)–(16.45) make clear which *derivatives* of the temporal profiles (A, B) enter: only A', A'' and B', B'' (no time or angular derivatives by symmetry).

Classical geometric part $\bar{R}_{\mu\nu}[\bar{g}]$ (for the $f(r)$ block)

With the geometric block

$$\bar{g}_{\mu\nu}dx^\mu dx^\nu = -\Big(1-\tfrac{f(r)}{r}\Big)c^2dt^2 + \frac{dr^2}{1-\frac{f(r)}{r}} + r^2d\Omega^2 \quad \text{(cf. eq. (16.33))},$$

the nonzero diagonal Ricci components (before imposing vacuum) are

$$\bar{R}_{tt}[\bar{g}] = \frac{(f(r)-r)\,f''(r)}{2\,r^2}\,c^2, \tag{16.46}$$

$$\bar{R}_{rr}[\bar{g}] = \frac{f''(r)}{2\,(r-f(r))}, \tag{16.47}$$

$$\bar{R}_{\theta\theta}[\bar{g}] = f'(r), \qquad \bar{R}_{\phi\phi}[\bar{g}] = \sin^2\theta\, f'(r), \tag{16.48}$$

and the Ricci scalar is

$$\bar{R}[\bar{g}] = \frac{r\,f''(r)+2\,f'(r)}{r^2}. \tag{16.49}$$

Thus $\bar{R}_{\mu\nu}[\bar{g}]=0$ when $f'=f''=0$, reproducing the Schwarzschild vacuum as expected [146, §31].

Chrono–geometric corrections $\Delta\bar{R}_{\mu\nu}[\bar{t}]$

With the diagonal temporal sector $\bar{t}_{tt}=A(r)c^2$, $\bar{t}_{rr}=B(r)$ and other entries zero (eq. (16.34)), the linear corrections follow from (16.45) with the covariant derivatives taken using the $\bar{g}$–connection. By staticity and spherical symmetry, only ∂_r acts nontrivially, and the background curvature couplings reduce to algebraic functions of f, f', f''. The result has the schematic (but *coordinate–exact*) structure

$$\Delta\bar{R}_{tt}[\bar{t}] = \frac{c^2}{2}\Big(-\nabla^2 A + \mathcal{C}_t\,A' + \mathcal{U}_t\,A\Big) + \frac{c^2}{2}\Big(\mathcal{K}_t\,B' + \mathcal{V}_t\,B\Big), \tag{16.50}$$

$$\Delta\bar{R}_{rr}[\bar{t}] = \frac{1}{2}\Big(-\nabla^2 B + \mathcal{C}_r\,B' + \mathcal{U}_r\,B\Big) + \frac{1}{2}\Big(\mathcal{K}_r\,A' + \mathcal{V}_r\,A\Big), \tag{16.51}$$

$$\Delta\bar{R}_{\theta\theta}[\bar{t}] = \frac{1}{2}\Big(\mathcal{C}_\Omega\,(A'+B') + \mathcal{U}_\Omega\,(A+B)\Big), \tag{16.52}$$

$$\Delta\bar{R}_{\phi\phi}[\bar{t}] = \sin^2\theta\;\Delta\bar{R}_{\theta\theta}[\bar{t}]. \tag{16.53}$$

where ∇^2 denotes the radial Laplacian associated with $\bar{g}$, and the coefficient functions $\mathcal{C}_\bullet, \mathcal{U}_\bullet, \mathcal{K}_\bullet, \mathcal{V}_\bullet$ are rational in r, f, and its derivatives (no time/angular dependence). Their explicit forms are obtained by substituting the Christoffel entries (eqs. (16.38) to (16.42)) into eq. (16.43) and expanding to linear order in A, B via eq. (16.45). (A full listing is straightforward albeit lengthy; see [146, §35.1] for analogous spherical reductions.)

Putting it together. Collecting (16.46)–(16.48) with (16.50)–(16.52), the unified components read

$$\bar{R}_{\mu\nu}[\bar{\Sigma}] = \bar{R}_{\mu\nu}[\bar{g}] + \Delta\bar{R}_{\mu\nu}[\bar{t}] + \mathcal{Q}_{\mu\nu}[\bar{t},\bar{t}], \qquad \bar{R}[\bar{\Sigma}] = \bar{R}[\bar{g}] + \Delta\bar{R}[\bar{t}] + \mathcal{Q}[\bar{t},\bar{t}], \tag{16.54}$$

with quadratic pieces $\mathcal{Q}$ suppressed if we work to first order in the temporal sector away from the horizon. Inserting (16.54) into the unified field equations (eq. (16.23) in vacuum, or eq. (16.13) when expectation stresses are included) yields three independent ODEs for $f(r)$, $A(r)$, $B(r)$ (the angular equation follows from the contracted Bianchi identity) with boundary data fixed by eqs. (16.22) and (16.32). This completes the curvature side of the problem and sets up Step 3 (assembling and analyzing the coupled ODE system; cf. Section 16.1.4).

Checks and limits. (i) **Classical limit.** Setting $A = B = 0$ gives $\bar{R}_{\mu\nu}[\bar{\Sigma}] = \bar{R}_{\mu\nu}[\bar{g}]$, which vanishes for constant $f = 2GM/c^2$ (Schwarzschild). (ii) **Near-horizon regularity.** In an ingoing Eddington–Finkelstein chart, the combinations entering $\Delta\bar{R}_{\mu\nu}$ remain finite provided $\{A, B\}$ satisfy the horizon relations implied by eq. (16.32) (see [146, §31–33]). (iii) **Thermodynamic consistency.** The sourced form eq. (16.13) ensures $\bar{\nabla}^{\mu}\bar{G}_{\mu\nu} = 0$ and conservation of expectation stresses, in harmony with the laws of thermodynamics in Section 15.1.

16.1.7 Step 3: The System of Differential Equations

With the connection coefficients in hand (Section 16.1.5) and the curvature decomposition assembled (Section 16.1.6, esp. eq. (16.54)), we now impose the *Unified Vacuum Field Equation*

$$\bar{G}_{\mu\nu}[\bar{\Sigma}] := \bar{R}_{\mu\nu}[\bar{\Sigma}] - \tfrac{1}{2}\,\bar{\Sigma}_{\mu\nu}\,\bar{R}[\bar{\Sigma}] = 0, \tag{16.55}$$

as derived in Section 16.1.2. In vacuum, contraction of (16.55) yields $\bar{R}[\bar{\Sigma}] = 0$, hence $\bar{R}_{\mu\nu}[\bar{\Sigma}] = 0$ [7, §B.2], so the two forms are equivalent (we keep $\bar{G}_{\mu\nu} = 0$ for structural clarity).

Unknowns and inputs. The ansatz (eq. (16.24)) supplies three r-dependent unknowns:

$$f(r), \qquad A(r), \qquad B(r),$$

with asymptotics and horizon data fixed by eqs. (16.22) and (16.32). If expectation stresses are present, one replaces (16.55) by the sourced form eq. (16.13).

1. The tt equation

Using $\bar{R}_{tt}[\bar{\Sigma}] = \bar{R}_{tt}[\bar{g}] + \Delta\bar{R}_{tt}[\bar{t}] + \mathcal{Q}_{tt}$ from eq. (16.54) together with eqs. (16.46) and (16.50), we obtain the second–order ODE

$$\underbrace{f''(r) + \frac{2}{r}f'(r)}_{\text{Schwarzschild block (vacuum)}} = \underbrace{\mathcal{F}_{tt}(r;\, A, B, A', B', A'', B'', f, f')}_{\text{chrono–geometric source from } \bar{t}+\mathcal{Q}}, \tag{16.56}$$

where $\mathcal{F}_{tt}$ is a rational function determined by eqs. (16.45) and (16.50) evaluated on the line element eq. (16.24). In the classical limit $A = B = 0$, the right–hand side vanishes and (16.56) integrates to the familiar $f(r) = 2GM/c^2$ after imposing asymptotic flatness [146, §31].

2. The rr equation

Similarly, from $\bar{R}_{rr}[\bar{\Sigma}] = 0$ with eqs. (16.47) and (16.51) we find

$$\underbrace{f''(r) - \frac{2}{r^2} f(r)}_{\text{Schwarzschild block (vacuum)}} = \underbrace{\mathcal{F}_{rr}(r;\, A, B, A', B', A'', B'', f, f')}_{\text{chrono–geometric source from } \bar{t}+\mathcal{Q}}, \tag{16.57}$$

which couples f to the radial phase–flow encoded by $B(r)$ (and cross–coupled to A).

3. The angular equation and consistency

The $\theta\theta$ (equivalently $\phi\phi$) component,

$$\bar{R}_{\theta\theta}[\bar{\Sigma}] = 0 \quad \Longleftrightarrow \quad f'(r) \;+\; \Delta\bar{R}_{\theta\theta}[\bar{t}] \;+\; \mathcal{Q}_{\theta\theta}[\bar{t}, \bar{t}] \;=\; 0, \tag{16.58}$$

with $\Delta\bar{R}_{\theta\theta}[\bar{t}]$ given schematically by eq. (16.52), provides a third relation among $\{f, A, B\}$. The contracted Bianchi identity ensures that (16.56), (16.57), and (16.58) are not over–constrained; any two imply the third up to integration constants, which are then fixed by eqs. (16.22) and (16.32) [7, §B.3]. In practice, (16.58) is used as a nontrivial *consistency check* and to improve numerical stability.

Boundary value problem (closure).

(i) Asymptotics. Impose $A, B \to 0$, $f \to 2GM/c^2$ as $r \to \infty$ (eq. (16.22)).

(ii) Horizon. Enforce the phase–boundary condition $1 - \frac{f(R_{\rm EH})}{R_{\rm EH}} - A(\xi(R_{\rm EH})) = 0$ and regularity of invariants in an ingoing chart (eq. (16.32); cf. [146, §31–33]).

(iii) Sourced case (optional). If vacuum polarization/expectation stresses are included, supplement (16.56)–(16.58) with eq. (16.13) and a constitutive input for the regulated vacuum sector (Section 16.1.17).

Outcome. Equations (16.56)–(16.58), together with the boundary data and (if applicable) sourcing, define a well–posed radial boundary–value problem for $\{f, A, B\}$. Setting $A = B = 0$ recovers Schwarzschild; nonzero (A, B) quantify the chrono–geometric corrections that encode PQS–driven coherence effects and determine the modified horizon location via eq. (16.32). Standard existence and uniqueness results for smooth ODEs apply locally under mild regularity of $\mathcal{F}_{tt}, \mathcal{F}_{rr}$ [7, App. E].

16.1.8 Solving the System and Obtaining the Black Hole Solution

With the coupled ODEs in hand (Section 16.1.7, esp. eqs. (16.56) to (16.58)), an exact closed-form solution is generally intractable. We therefore adopt a physically grounded program based on asymptotic analysis and controlled perturbations (matched across regimes defined by $\xi(r)$; cf. Section 16.1.3).

1. The Classical Limit

Far from the black hole ($r \to \infty$), chrono–geometric corrections must vanish. We impose the asymptotic boundary conditions

$$A(r) \to 0, \qquad B(r) \to 0 \quad \text{as } r \to \infty, \tag{16.59}$$

consistent with eq. (16.22). In this regime the right–hand sides of the modified field equations, $\mathcal{F}_{tt}$ and $\mathcal{F}_{rr}$ in eqs. (16.56) and (16.57), vanish, so the system reduces to the standard vacuum equations for the geometric block.

Using the geometric Ricci components from Section 16.1.6,

$$\bar{R}_{\theta\theta}[\bar{g}] = f'(r) \quad \text{and} \quad \bar{R}_{tt}[\bar{g}],\ \bar{R}_{rr}[\bar{g}] \text{ as in eqs. (16.46) and (16.47)},$$

the vacuum condition $\bar{R}_{\mu\nu}[\bar{g}] = 0$ (equivalently $\bar{G}_{\mu\nu}[\bar{g}] = 0$) implies $f'(r) = 0$, i.e. $f(r) = \text{const}$. Asymptotic flatness fixes that constant by comparing $-\big(1 - f(r)/r\big)c^2dt^2$ with the Newtonian limit, yielding

$$f(r) = \frac{2GM}{c^2}, \tag{16.60}$$

with M the (ADM) mass measured at infinity (cf. the mass function $m(r)$ in eq. (16.15)). Thus the zeroth–order background is precisely Schwarzschild, in agreement with the Jebsen–Birkhoff theorem and standard uniqueness results for the spherically symmetric vacuum exterior [146, §31][7, §6].

This Schwarzschild background serves as the reference geometry on which the temporal sector (A, B) is treated perturbatively in subsequent steps, subject to horizon regularity (phase–boundary condition $1 - \frac{f(R_{\rm EH})}{R_{\rm EH}} - A(\xi(R_{\rm EH})) = 0$; see eq. (16.32)) and matched to the interior/PQS behaviour prescribed by $\xi(r)$ (eq. (16.17)).

2. The Quantum Correction Regime

Near the event horizon and inward toward the PQS core, the chrono–geometric terms $A(r), B(r)$ cannot be neglected. We therefore adopt a controlled perturbative scheme about the classical background (Section 16.1.8):

Perturbative ansatz. Introduce a bookkeeping parameter ε and expand

$$f(r) = f_0 + \varepsilon f_1(r) + \mathcal{O}(\varepsilon^2), \qquad A(r) = \varepsilon a_1(r) + \mathcal{O}(\varepsilon^2), \qquad B(r) = \varepsilon b_1(r) + \mathcal{O}(\varepsilon^2), \tag{16.61}$$

with $f_0 = \frac{2GM}{c^2}$ from eq. (16.60). Boundary data are fixed by $A, B \to 0$ at infinity (eqs. (16.22) and (16.59)) and by horizon regularity/phase–boundary conditions (eq. (16.32)). This follows the standard linearization strategy in GR (cf. [146, §35.1], [7]).

First order: linear system for a_1, b_1 on Schwarzschild. Insert (16.61) into the curvature split (eq. (16.54)). Because $\bar{R}_{\mu\nu}[\bar{g}_0] = 0$ for Schwarzschild, the $\mathcal{O}(\varepsilon)$ equations are

$$\Delta\bar{R}_{tt}[a_1, b_1] = 0, \qquad \Delta\bar{R}_{rr}[a_1, b_1] = 0, \qquad \Delta\bar{R}_{\theta\theta}[a_1, b_1] = 0, \tag{16.62}$$

with $\Delta\bar{R}_{\mu\nu}$ given in eqs. (16.50) to (16.52) (coefficients evaluated at $f = f_0$). This yields two independent, linear ODEs for a_1, b_1 plus one consistency relation (the angular component, guaranteed by the contracted Bianchi identity; cf. Section 16.1.6). Solutions are selected by:

(i) Asymptotics. $a_1, b_1 \to 0$ as $r \to \infty$ (eq. (16.59)).

(ii) Horizon regularity. Finiteness of curvature invariants in an ingoing Eddington–Finkelstein chart (cf. [146, §31–33]) and the phase–boundary condition $1-\frac{f_0}{R_{\rm EH}}-a_1(\xi(R_{\rm EH})) = 0$ to leading order (eq. (16.32) with $f = f_0$).

Next order: back–reaction for f_1. At $\mathcal{O}(\varepsilon)$, f_1 enters the field equations through $\bar{G}_{\mu\nu}[\bar{g}_0 + \varepsilon(\delta g)]$, with δg induced by f_1, and through the quadratic curvature terms $\mathcal{Q}_{\mu\nu}[a_1, b_1]$ (cf. eq. (16.54)). Using the tt and rr equations (eqs. (16.56) and (16.57)) one obtains linear ODEs for f_1 on the fixed background,

$$\left(\frac{d^2}{dr^2} + \frac{2}{r}\frac{d}{dr}\right) f_1(r) = \mathcal{S}_{tt}[a_1, b_1], \tag{16.63}$$

$$\left(\frac{d^2}{dr^2} - \frac{2}{r^2}\right) f_1(r) = \mathcal{S}_{rr}[a_1, b_1], \tag{16.64}$$

where $\mathcal{S}_{tt}, \mathcal{S}_{rr}$ are the explicit source functionals built from a_1, b_1 and their derivatives (coming from $\mathcal{Q}_{\mu\nu}$ and the linearized $\Delta\bar{R}$ pieces; cf. eq. (16.45)). Compatibility of (16.63)–(16.64) is ensured by the angular equation (eq. (16.58)) and the Bianchi identity.

Horizon shift at leading order. The event horizon $R_{\rm EH}$ solves $1 - \frac{f(R_{\rm EH})}{R_{\rm EH}} - A(\xi(R_{\rm EH})) = 0$ (eq. (16.21)). Linearizing with (16.61) about $R_S = f_0$ gives, to first order and neglecting the small $a_1' \, \delta R$ term,

$$\delta R_{\rm EH} := R_{\rm EH} - R_S \;\approx\; f_1(R_S) \;-\; R_S \, a_1(R_S), \qquad R_S = \frac{2GM}{c^2}, \tag{16.65}$$

which can be refined by including the denominator $1/R_S - a_1'(R_S)$ when needed (from the exact linearization of the horizon condition).

Structure of the Final Solution

The iterative scheme (16.61)–(16.64) furnishes a semi–classical, relational description that reduces to Schwarzschild at large r and encodes chrono–geometric corrections near the horizon and into the interior:

(i) Exterior geometry. A modified metric smoothly matching the Schwarzschild background as $r \to \infty$ (eqs. (16.59) and (16.60)).

(ii) Horizon location. A shifted event horizon determined by eq. (16.21), with leading correction $\delta R_{\rm EH}$ given by eq. (16.65).

(iii) Interior behaviour. A controlled approach to the PQS regime, where the temporal sector dominates (eq. (16.17)) and the classical singularity is replaced by the correlated PQS core (Section 1.3).

This completes the construction of the solution scheme consistent with the coupled equations in Section 16.1.7 and the curvature analysis in Section 16.1.6. Further refinements (e.g., inclusion of expectation stresses via eq. (16.13)) proceed within the same perturbative framework [146, 7].

16.1.9 The System of Coupled Differential Equations

Starting from the unified vacuum equation $\bar{G}_{\mu\nu}[\bar{\Sigma}] = 0$ (eq. (16.55)) and the curvature decomposition built in Section 16.1.6 (see eq. (16.54)), substitution of the Christoffel symbols (Section 16.1.5) for the unified ansatz (eq. (16.24)) yields a closed system of non–linear ODEs for the three unknown radial functions

$$f(r), \qquad A(r), \qquad B(r),$$

where f determines the geometric block $\bar{g}_{\mu\nu}$ and (A, B) encode the chrono–geometric (temporal) corrections $\bar{t}_{\mu\nu}$.

For clarity, we summarize the three independent diagonal field equations—each understood as the precise versions already stated in Section 16.1.7—together with their physical roles.

1. Time–time (tt) equation

This governs curvature in the temporal direction and is most directly sensitive to the gravitational potential. In the compact form already introduced,

$$\underbrace{f''(r) + \frac{2}{r} f'(r)}_{\text{Schwarzschild block (vacuum)}} = \underbrace{\mathcal{F}_{tt}(r;\, A, B, A', B', A'', B'', f, f')}_{\text{chrono–geometric source from } \bar{t}+\mathcal{Q}}, \tag{16.66}$$

where the right–hand side arises from $\Delta\bar{R}_{tt}[\bar{t}]$ and the quadratic curvature $\mathcal{Q}_{tt}$ (cf. eq. (16.50); see also [146, §31], [7, §B] for the classical structure).

2. Radial–radial (rr) equation

This constrains the radial geometry and couples f to the radial phase–flow:

$$\underbrace{f''(r) - \frac{2}{r^2} f(r)}_{\text{Schwarzschild block (vacuum)}} = \underbrace{\mathcal{F}_{rr}(r;\, A, B, A', B', A'', B'', f, f')}_{\text{chrono–geometric source from } \bar{t}+\mathcal{Q}}, \tag{16.67}$$

with the source determined by $\Delta\bar{R}_{rr}[\bar{t}]$ and $\mathcal{Q}_{rr}$ (cf. eq. (16.51)).

3. Angular $(\theta\theta)$ equation and consistency

The angular sector furnishes the third relation,

$$\bar{R}_{\theta\theta}[\bar{\Sigma}] = 0 \quad \Longleftrightarrow \quad \underbrace{f'(r)}_{\text{Schwarzschild block (vacuum)}} + \underbrace{\Delta\bar{R}_{\theta\theta}[\bar{t}]}_{\text{linear chrono–geom.}} + \underbrace{\mathcal{Q}_{\theta\theta}[\bar{t},\bar{t}]}_{\text{quadratic}} = 0, \tag{16.68}$$

cf. eqs. (16.48) and (16.52). The contracted Bianchi identity guarantees that any two of eqs. (16.56) to (16.58) imply the third up to integration constants, which are fixed by boundary data [7, §B.3].

Boundary data and problem statement. The system eqs. (16.56) to (16.58) is solved as a radial boundary–value problem with:

(i) Asymptotics. $A(r), B(r) \to 0$ and $f(r) \to 2GM/c^2$ as $r \to \infty$ (eqs. (16.22), (16.59) and (16.60)).

(ii) Horizon regularity. The phase–boundary condition $1 - \frac{f(R_{\rm EH})}{R_{\rm EH}} - A(\xi(R_{\rm EH})) = 0$ with curvature finiteness in an ingoing chart (eq. (16.32); cf. [146, §31–33]).

Solution strategy. Proceed as in Section 16.1.8: (i) impose the classical limit to obtain the Schwarzschild background (Section 16.1.8); (ii) solve the linearized system for (A, B) on that background and then the back–reaction for f (Section 16.1.8, eqs. eqs. (16.62) to (16.64)); or (iii) integrate the full non–linear system numerically with the above boundary data. In sourced settings, replace $\bar{G}_{\mu\nu} = 0$ by the generalized form $\bar{G}_{\mu\nu} + \Lambda\bar{\Sigma}_{\mu\nu} = \kappa\langle\hat{T}_{\mu\nu}\rangle$ (eq. (16.13)) and supply the constitutive inputs from the regulated vacuum sector (Section 16.1.17).

In this way the abstract relational principles are converted into a concrete, testable set of equations for the chrono–geometric black hole, with the classical theory recovered for $A = B = 0$ and controlled deviations governed by the temporal sector.

16.1.10 The Event Horizon: From Classical Boundary to Ontological Frontier

In classical terms, the **event horizon** is the boundary of the causal past of future null infinity; equivalently, a null hypersurface that acts as a one–way membrane [7, 146]. For stationary black holes it coincides with a *Killing horizon* generated by a complete Killing field χ^a [7, 643].

1. The Classical Definition (General Relativity)

(i) Causal boundary. Any event inside the horizon is causally disconnected from all exterior observers; null generators lie on the horizon and future–directed causal curves cannot cross outward [7].

(ii) Geometric location (Schwarzschild). For a non–rotating mass M, the horizon is located at the *Schwarzschild radius*

$$R_S = \frac{2GM}{c^2}, \tag{16.69}$$

obtained by setting the lapse of the Schwarzschild line element to zero [146].

2. The Redefined Horizon (IQR Framework)

Within the present framework, the horizon remains a causal boundary for exterior observers (Section 16.1.10), but acquires a deeper ontological status as a *regime boundary* in the chrono–geometric description (Section 16.1.3). In the unified ansatz,

$$\bar{\Sigma}_{\mu\nu}dx^\mu dx^\nu = -\Big(1 - \frac{f(r)}{r} - A(\xi)\Big)c^2\,dt^2 + \frac{dr^2}{1 - \frac{f(r)}{r} - B(\xi)} + r^2 d\Omega^2, \tag{16.70}$$

where $f(r) := \frac{2Gm(r)}{c^2}$ is the mass function, $\xi = \xi(r)$ is the regime tracker (coherence invariant), and $A(\xi), B(\xi)$ are the dimensionless temporal–tensor profiles. See eqs. (16.17) and (16.20). The temporal sector (A, B) becomes $\mathcal{O}(1)$ at the transition from the **Macroscopic** to the **Relational/Mesoscopic** regime.

Horizon conditions. We use two equivalent statements for the event horizon $R_{\rm EH}$:

(i) Ansatz (lapse–root) form.

$$1 - \frac{f(R_{\rm EH})}{R_{\rm EH}} - A\Big(\xi(R_{\rm EH})\Big) \;=\; 0, \tag{16.71}$$

(ii) Coherence–corrected (transcendental) summary.

$$R_{\rm EH} \;+\; \alpha\Big(\,\xi(R_{\rm EH})\,\Big) \;=\; \frac{2GM}{c^2}, \tag{16.72}$$

where $\alpha(\cdot)$ is a *length–valued* response encoding the near–horizon relational back–reaction (so the dimensions match the LHS). In the classical limit $A \to 0$, $B \to 0$, and $f(r) \to 2GM/c^2$, eq. (16.21) reduces to eq. (16.69); the local, mass–fixed form is given in eq. (16.16).

Location shift and physical content. When chrono–geometric corrections are small but nonzero, the horizon is shifted relative to $R_S := 2GM/c^2$. Linearizing eq. (16.21) about the Schwarzschild background (Section 16.1.8) yields the leading correction

$$\delta R_{\rm EH} \;\approx\; f_1(R_S) \;-\; R_S\,A\Big(\xi(R_S)\Big), \tag{16.73}$$

which matches the master first–order expression written in terms of the horizon–value coefficients (a_1, f_1):

$$\delta R_H \;=\; f_1(R_S) \;-\; R_S\,a_1(R_S). \tag{16.74}$$

Thus, the effective horizon location depends not only on mass (through f) but also on the state–dependent temporal profile $A(\xi)$, which encodes quantum–relational structure near the boundary. The transcendental form eq. (16.72) is equivalent at leading order by identifying $\alpha(\xi(R_{\rm EH})) = R_S - R_{\rm EH} = -\,\delta R_H$.

Summary. The event horizon retains its classical role as a global causal boundary [7] while, in the IQR framework, it is simultaneously an *ontological frontier*: a phase transition surface where the emergent macroscopic geometry gives way to the chrono–geometric regime (Section 16.1.3). Its precise radius is fixed by the lapse condition (16.21), reducing to *eq.* (16.69) in the classical limit and acquiring controlled corrections from the temporal sector through $A(\xi)$.

16.1.11 Approaching the Event Horizon: A Descent into Relational Reality

In the IQR account, nearing a black hole's event horizon is not merely approaching a spatial boundary but entering a new *regime* of description—the chrono–geometric phase in which temporal curvature becomes order-unity (cf. Sections 16.1.3 and 16.1.10). To exterior observers, familiar GR effects intensify, while the temporal sector $\bar{t}_{\mu\nu}$ introduces controlled deviations encoded by $A(r), B(r)$.

The Event Horizon as a Phase Boundary

As one descends from the **Macroscopic** to the **Relational/Mesoscopic** regime, the unified lapse

$$N^2(r) \;:=\; 1 - \frac{f(r)}{r} - A\Big(\xi(r)\Big) \tag{16.75}$$

tends to zero at the event horizon $R_{\rm EH}$, which is defined by $N^2(R_{\rm EH}) = 0$ (see eqs. (16.16) and (16.21)). In this zone the temporal profiles (A, B) cease to be negligible and signal the onset of chrono–geometric structure (cf. eq. (16.17)).

1. Extreme gravitational time dilation and redshift. These follow from the tt block of the unified tensor,

$$\bar{\Sigma}_{tt} \;=\; -c^2\Big(1 - \frac{f(r)}{r} - A(r)\Big), \tag{16.76}$$

so a static clock at radius r relates to the asymptotic coordinate time t by

$$d\tau(r) \;=\; \sqrt{1 - \frac{f(r)}{r} - A(r)}\, dt, \qquad \frac{\nu_\infty}{\nu_{\rm loc}(r)} \;=\; \sqrt{1 - \frac{f(r)}{r} - A(r)}, \tag{16.77}$$

where ν_∞ is the frequency measured at infinity and $\nu_{\rm loc}(r)$ the locally emitted frequency for a static source [7, 146, §6, §31]. As $r \to R_{\rm EH}$, the factor in (16.77) vanishes: clocks *appear* to halt and outgoing signals redshift without bound. The function $A(r)$ perturbs this "temporal gradient," slightly reshaping the near-horizon scaling relative to pure Schwarzschild (cf. Section 16.1.8).

2. Effective surface gravity and temperature. Expanding the unified lapse $N^2(r)$ (defined in eq. (16.75)) about the horizon gives

$$N^2(r) \;\approx\; \frac{2\,\kappa_{\rm eff}}{c^2}\Big(r - R_{\rm EH}\Big), \qquad \kappa_{\rm eff} := \frac{c^2}{2}\,\frac{d}{dr}\left(1 - \frac{f(r)}{r} - A(r)\right)\Bigg|_{r=R_{\rm EH}}. \tag{16.78}$$

This reproduces the standard static–horizon definition (with the time–translation Killing field normalized by $\chi^a\chi_a \to -c^2$ at infinity) [7, §12]. The Hawking temperature measured at infinity is then

$$T_H = \frac{\hbar\,\kappa_{\rm eff}}{2\pi k_B\, c}, \tag{16.79}$$

and, in the classical limit $A \to 0$ with $f(r) = 2GM/c^2$ (eq. (16.60)), one recovers the Schwarzschild result

$$T_H^{(\mathrm{Schw})} = \frac{\hbar\,\kappa_S}{2\pi k_B\,c} = \frac{\hbar c^3}{8\pi G M k_B}, \tag{16.80}$$

$$\kappa_S = \frac{c^4}{4GM}. \tag{16.81}$$

(See also [87, 314].) Thus $A(r)$ shifts both the horizon location and the surface gravity (cf. eqs. (16.21) and (16.73)); $B(r)$ affects T_H only indirectly through its back–reaction on f and A.

Summary. For exterior observers the horizon is a global causal boundary [7], yet in the IQR framework it is also an *ontological frontier* where the macroscopic geometry transitions into the chrono–geometric regime (Section 16.1.10). The deviations encoded by $A(r)$ and $B(r)$ fine-tune the dilation/redshift law (16.77), the horizon position (eq. (16.21)), and the temperature via κ_{eff} (16.78), all while smoothly recovering classical GR in the limit $A, B \to 0$.

The Halting of Geometric Time and the Emergence of Relational Dynamics

The "time" that appears to grind to a halt at the horizon for a distant observer is the *emergent geometric time* t_{geom}: the Killing time associated with the exterior stationary geometry ($\chi^a\partial_a = \partial_t$) of $\mathcal{M}_{\mathrm{geom}}$ [7, §6, §12]. Operationally, t_{geom} is measured by ideal clocks at rest in the exterior region, where the temporal sector is negligible (Section 16.1.8 and eq. (16.22)).

Geometric time from the geometric block. In the unified description, redshift and time–dilation for static observers follow from the tt–component of the tensor $\bar{\Sigma}_{\mu\nu}$,

$$\bar{\Sigma}_{tt} = -c^2\Big(1 - \tfrac{f(r)}{r} - A(r)\Big), \quad d\tau(r) = \sqrt{1 - \tfrac{f(r)}{r} - A(r)}\; dt, \qquad \frac{\nu_\infty}{\nu_{\mathrm{loc}}(r)} = \sqrt{1 - \tfrac{f(r)}{r} - A(r)}, \tag{16.82}$$

as in eqs. (16.76) and (16.77). For static sources the factor in (16.82) tends to zero at the event horizon, so a distant observer infers an *apparent* "freezing" of t_{geom} together with infinite redshift. This is the standard GR behavior for stationary observers near a Killing horizon [7, §12]; the spacetime itself is regular there in appropriate coordinates (e.g., ingoing Eddington–Finkelstein) [146, §31–33].

Horizon as regime boundary (from geometry to chrono–geometry). At the horizon R_{EH}, defined by the lapse condition

$$1 - \frac{f(R_{\mathrm{EH}})}{R_{\mathrm{EH}}} - A\Big(\xi(R_{\mathrm{EH}})\Big) = 0, \tag{16.83}$$

(eqs. (16.16) and (16.21)), the temporal sector becomes order–unity (Sections 16.1.3 and 16.1.10). Beyond this *phase boundary*, the purely geometric description by $\bar{g}_{\mu\nu}$ no longer suffices; one must pass to the full chrono–geometric structure.

Relational manifold (pointer to earlier definition). As defined in Definition 5.66, the relational manifold is $\mathcal{M}_{\rm rel} = (\mathcal{X}_q, \bar{g}_{\mu\nu}, \bar{t}_{\mu\nu})$. Its dynamics use the unified tensor $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (Definition 5.26).

From $t_{\rm geom}$ to relational time. Inside the horizon, $t_{\rm geom}$ ceases to be the relevant parameter of evolution for infalling matter: the motion is governed by geodesics (and quantum transport) of $\bar{\Sigma}_{\mu\nu}$, not of $\bar{g}_{\mu\nu}$ alone. The effective "time" becomes *relational*, reflecting the dominant temporal sector and its couplings (Section 16.1.8). Operationally, one may decompose the relational temporal flow into complementary contributions:

(i) Entanglement–phase time $t_{\rm corr}$. Evolution generated by nonlocal correlations between interior and exterior sectors (consistent with the global, unitary picture; cf. Section 16.1.1).

(ii) Vacuum–response time $t_{\rm vac}$. Phase shifts induced by the structured quantum vacuum that sources $A(r), B(r)$ near the boundary (Section 16.1.17 and eq. (16.45)).

(iii) Spectral–shear time $t_{\rm spec}$. Evolution associated with noncommuting generators in the temporal block, reflected in the curvature terms built from gradients of $\bar{t}_{\mu\nu}$ (Section 16.1.6).

Conclusion. The "halting of time" at the horizon is a *projection effect* of the exterior, geometric description: $t_{\rm geom}$ freezes for stationary observers, while the infalling system continues with well–defined evolution in the full relational manifold $\mathcal{M}_{\rm rel}$. In the IQR framework, crossing the horizon marks the transition from a single–parameter geometric time to a richer quantum–relational dynamics encoded by $\bar{\Sigma}_{\mu\nu}$, with classical GR recovered in the limit $A, B \to 0$ (Section 16.1.8).

2. The Emergence of Chronogeometric "Viscosity" In the IQR framework there appears a phenomenon with no classical analogue: the temporal sector $\bar{t}_{\mu\nu}$ endows spacetime with an *effective relational viscosity*. Microscopically this arises from the nonlocal matter–geometry coupling and the structured quantum vacuum; macroscopically it manifests through gradients of the temporal profiles $A(r), B(r)$ in the curvature (cf. eqs. (16.45), (16.50) and (16.51)).

(i) The vacuum is not empty. Near the horizon the regulated vacuum sector becomes dynamically active (Section 16.1.17). Fluctuations of the Gaiamiti and Tantrakyma fields—carried by *Apeirons* and *Kymons*—mediate geometry–matter exchange via the bi-local kernel (Definitions 9.246, 9.310, 10.222 and 10.223). The resulting state-dependent expectation stresses source A, B and their derivatives in the field equations, giving rise to transport-like (dissipative-looking) terms in the effective description.

(ii) Distortion of spacetime (refractive analogy). The functions $A(r)$ and $B(r)$ deform the lapse and the radial proper length element in the unified ansatz (eq. (16.20)):

$$d\tau(r) = \sqrt{1 - \tfrac{f(r)}{r} - A(r)}\, dt, \qquad d\ell_r = \frac{dr}{\sqrt{1 - \tfrac{f(r)}{r} - B(r)}} \quad \text{(cf. eq. (16.82))}.$$

Heuristically, A acts as a *temporal* refractive index (slowing the exterior time flow), while B acts as a *radial* refractive index (thickening radial distances). The gradients $A'(r), B'(r)$ enter the Ricci components through $\Delta\bar{R}_{\mu\nu}[\bar{t}]$ (eqs. (16.50) to (16.52)), encoding a viscosity-like resistance of the chrono–geometric medium to rapid changes of the lapse and the radial scale factor.

3. The Horizon as a "Boiling" Surface of Quantum Information To an exterior observer with finite access, the horizon appears as a fluctuating, high-entropy interface: locally thermal with temperature T_H (Section 16.1.1 and eq. (16.79)) and incessant redshift (eq. (16.82)). Globally, however, the complete Holo-state evolves unitarily (Section 16.1.1). In the IQR account, the horizon is the locus where information borne by infalling matter is *re-coded*—via the bi-local interaction $\mathcal{L}_{\text{int}}$ (Definition 10.222)—from local Tantrakyma microstates into nonlocal correlations of the Gaiamiti sector and, ultimately, into multipartite entanglement of the outgoing quanta (Section 16.1.1). This picture dovetails with unitary evaporation diagnostics (scrambling and Page-curve/island analyses) [336, 335, 337, 139, 232] while deriving the mechanism from the theory's native interaction structure.

Summary. Approaching the horizon is a journey to the boundary of the classical regime: the vacuum becomes an active, structured medium (Section 16.1.17); the effective dynamics acquire viscosity-like terms governed by A, B and their gradients (eqs. (16.50) and (16.51)); and the horizon itself functions as a roiling interface where locally thermal behavior coexists with globally unitary information flow (Section 16.1.1). Beyond this shimmering veil, the chrono–geometric interior guides evolution toward the PQS core (Section 1.3).

16.1.12 The Event Horizon as a Phase Boundary in Reality's Fabric

In the IQR framework, the event horizon is not merely a geometric locus but a *regime boundary* separating the domain where a simplified operational geometry suffices from the domain where the full chrono–geometric description is required (cf. Sections 16.1.3 and 16.1.10). Concretely, the unified lapse

$$N^2(r) \;=\; 1 - \frac{f(r)}{r} - A\Big(\xi(r)\Big) \tag{16.84}$$

tends to zero at the horizon R_{EH} defined by $N^2(R_{\text{EH}}) = 0$ (eqs. (16.16) and (16.21)). This signals the onset of chrono–geometric dominance (eq. (16.17)).

1. The exterior world: the domain of $\mathcal{M}_{\text{geom}}$. Far from the black hole (the *Macroscopic Regime*), the temporal sector is negligible and obeys

$$A(r),\, B(r) \;\to\; 0 \qquad (r \to \infty),$$

so that the spacetime is accurately captured by the emergent geometric block $\bar{g}_{\mu\nu}$ with asymptotically Schwarzschild behavior (eqs. (16.59) and (16.60)). In this operational domain

$$\mathcal{M}_{\text{geom}} := \Big(\mathcal{X}_q,\; \bar{g}_{\mu\nu}(x),\; d_{\text{geom}},\; t_{\text{geom}}\Big),$$

test bodies follow geodesics of the Levi–Civita connection of $\bar{g}$ and the usual GR laws apply [7, §6] (see also the standard connection formula eq. (16.37)).

2. The horizon and beyond: the domain of $\mathcal{M}_{\rm rel}$. Approaching the horizon one enters the *Relational/Mesoscopic Regime*, where the temporal profiles (A, B) become $\mathcal{O}(1)$ and the purely geometric description is insufficient (Section 16.1.11). The correct arena is the *relational manifold* defined earlier,

$$\mathcal{M}_{\rm rel} = (\mathcal{X}_q,\ \bar{g}_{\mu\nu},\ \bar{t}_{\mu\nu}) \quad \text{with} \quad \bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu} \quad (\text{cf. } \textit{Definitions } 5.26 \textit{and } 5.66).$$

Dynamics—geodesic motion and transport—are governed by the Levi–Civita connection of $\bar{\Sigma}$ (not of $\bar{g}$ alone), with

$$\Gamma[\bar{\Sigma}] = \Gamma[\bar{g}] + \Delta\Gamma[\bar{t}] \quad \text{as in } \textit{eqs. } (16.25) \textit{and } (16.26), \tag{16.85}$$

so that gradients of the temporal sector directly enter the equations of motion (Sections 16.1.5 and 16.1.6).

The synthesis. The event horizon thus marks the transition surface

> *from the operational, emergent geometry $\mathcal{M}_{\rm geom}$ to the full relational manifold $\mathcal{M}_{\rm rel}$,*

where the temporal tensor $\bar{t}_{\mu\nu}$ becomes dynamically significant. Equivalently, it is the locus where the lapse (16.84) vanishes and the near-horizon kinematics are governed by the effective surface gravity $\kappa_{\rm eff}$ (eq. (16.78)) and the associated Hawking temperature T_H (eq. (16.79)), with observable redshift/time-dilation following eq. (16.77). In this sense, the horizon is elevated from a classical *point of no return* to a genuine *phase boundary* in the fabric of reality, aligning the causal picture of GR with the chrono–geometric ontology and underpinning the treatise's information-theoretic resolution of black-hole puzzles (Section 16.1.1).

16.1.13 A Solvable Path: Asymptotic Matching

Identifying the horizon as the boundary between the operational geometry $\mathcal{M}_{\rm geom}$ and the full relational geometry $\mathcal{M}_{\rm rel}$ (cf. Section 16.1.12) yields a precise organizing principle for solving the coupled ODEs of Section 16.1.9. The idea is to construct *outer* and *inner* solutions in their natural regimes and then *match* them across an overlap where both descriptions are valid.

Step 1: Outer solution (classical limit). Work in the Macroscopic Regime, $r \to \infty$, where $A, B \to 0$ (eq. (16.59)). Setting $A = B = 0$ in the field equations (16.56)–(16.58) reduces them to the vacuum Schwarzschild block, yielding

$$f(r) = \frac{2GM}{c^2}, \qquad \bar{\Sigma}_{\mu\nu} \to \bar{g}^{\rm Schw}_{\mu\nu} \quad (\text{cf. } \textit{Section } 16.1.8 \textit{and eq. } (16.60)). \tag{16.86}$$

This furnishes the zeroth–order background for subsequent corrections.

Step 2: Inner solution (quantum–relational correction). Near the horizon the temporal sector becomes relevant (cf. Section 16.1.11). Treat A, B as small on the Schwarzschild

background and solve the linearized system

$$\Delta\bar{R}_{tt}[a_1, b_1] = 0, \tag{16.87}$$
$$\Delta\bar{R}_{rr}[a_1, b_1] = 0, \tag{16.88}$$
$$\Delta\bar{R}_{\theta\theta}[a_1, b_1] = 0. \tag{16.89}$$

See eqs. (16.50) to (16.52) and (16.62) for the operators and coefficients. Use boundary data $a_1, b_1 \to 0$ as $r \to \infty$ and impose horizon regularity at $R_{\rm EH}$ (cf. eqs. (16.21) and (16.32)).

Then incorporate back–reaction by solving for f_1 on the fixed background:

$$\left(\frac{d^2}{dr^2} + \frac{2}{r}\frac{d}{dr}\right) f_1(r) = \mathcal{S}_{tt}[a_1, b_1], \tag{16.90}$$
$$\left(\frac{d^2}{dr^2} - \frac{2}{r^2}\right) f_1(r) = \mathcal{S}_{rr}[a_1, b_1]. \tag{16.91}$$

See eqs. (16.63) and (16.64); use the angular equation for consistency (eq. (16.58)). This yields the inner profiles $A(r) \approx \varepsilon a_1(r)$, $B(r) \approx \varepsilon b_1(r)$, and $f(r) \approx f_0 + \varepsilon f_1(r)$ (cf. Section 16.1.8).

Step 3: Matching and completion. Choose an overlap radius r_m with $r_m \gtrsim R_{\rm EH}$ where both descriptions are valid. Impose continuity of the lapse and metric coefficients (and, if desired, their first derivatives) between inner and outer solutions:

$$\begin{aligned} &N^2_{\rm inner}(r_m) = N^2_{\rm outer}(r_m), \qquad \partial_r N^2_{\rm inner}(r_m) = \partial_r N^2_{\rm outer}(r_m), \\ &\text{with} \quad N^2(r) = 1 - \frac{f(r)}{r} - A(r) \quad (\text{cf. } eq.\ (16.75)). \end{aligned} \tag{16.92}$$

Matching fixes the remaining integration constants, determines the corrected horizon location via the lapse root

$$1 - \frac{f(R_{\rm EH})}{R_{\rm EH}} - A\Big(\xi(R_{\rm EH})\Big) = 0 \quad (\text{cf. } eq.\ (16.21)), \tag{16.93}$$

and, at leading order, yields the shift $\delta R_{\rm EH}$ (cf. eq. (16.73)). The resulting global solution is smooth, asymptotically Schwarzschild, and regular at the horizon; it encodes the chrono–geometric corrections required by the IQR framework while recovering GR in the limit $A, B \to 0$.

Outcome. The asymptotic–matching program translates the coupled system of Section 16.1.9 into a controlled, solvable pipeline: classical exterior → linear temporal sector → back–reaction → matched global solution, with thermodynamic and information–theoretic consistency checked via $\kappa_{\rm eff}$ and T_H (eqs. (16.78) and (16.79)) and the global unitary narrative (Section 16.1.1).

16.1.14 Step 1: Outer Solution — Classical Schwarzschild Limit

Setup. In the Macroscopic Regime ($r \to \infty$) the temporal sector is negligible, i.e. $A(r), B(r) \to 0$ (eq. (16.59)). Setting $A = B = 0$ in the unified line element (eq. (16.18)) gives the purely geometric ansatz

$$\bar{g}_{\mu\nu}dx^\mu dx^\nu = -\left(1 - \frac{f(r)}{r}\right)c^2dt^2 + \frac{dr^2}{1 - \frac{f(r)}{r}} + r^2d\Omega^2, \tag{16.94}$$

where r is the areal radius and $f(r)$ is the mass–length function (related to the mass function by $f(r) = 2Gm(r)/c^2$; cf. eq. (16.15)). Our goal is to solve the *unified vacuum equations* in this sector,

$$\bar{G}_{\mu\nu}[\bar{g}] = 0 \quad \Longleftrightarrow \quad \bar{R}_{\mu\nu}[\bar{g}] = 0 \qquad \text{(vacuum, cf. } eqs.\ (16.12) and\ (16.55)\text{)}. \tag{16.95}$$

Curvature components for $\bar{g}$. From the computation in Section 16.1.6, the nonvanishing diagonal Ricci components for (16.94) are

$$\bar{R}_{tt}[\bar{g}] = \frac{(f - r)\, f''}{2\, r^2}\, c^2, \tag{16.96}$$

$$\bar{R}_{rr}[\bar{g}] = \frac{f''}{2\,(r - f)}, \tag{16.97}$$

$$\bar{R}_{\theta\theta}[\bar{g}] = f'(r), \qquad \bar{R}_{\phi\phi}[\bar{g}] = \sin^2\theta\, f'(r), \tag{16.98}$$

and the Ricci scalar is $\bar{R}[\bar{g}] = (r f'' + 2f')/r^2$ (cf. eqs. (16.46) to (16.49)).

Solving the vacuum equations. The $\theta\theta$ component $\bar{R}_{\theta\theta} = 0$ immediately gives

$$f'(r) = 0 \quad \Longrightarrow \quad f(r) = C, \tag{16.99}$$

with C a constant. Substituting (16.99) into (16.96)–(16.97) (so $f'' = 0$) yields $\bar{R}_{tt} = \bar{R}_{rr} = 0$ identically, hence $\bar{R}_{\mu\nu}[\bar{g}] = 0$ is satisfied. Thus the most general spherically symmetric *vacuum* solution of the form (16.94) is characterized by a single constant C.

Fixing the constant by asymptotics (Newtonian/ADM). To determine C, compare the asymptotic form of g_{tt} in (16.94),

$$g_{tt} = -\left(1 - \frac{C}{r}\right)c^2 = -c^2 + \frac{C\, c^2}{r} + \mathcal{O}(r^{-2}), \tag{16.100}$$

with the weak–field expansion of a static Newtonian potential $\Phi(r) = -GM/r$,

$$g_{tt} \simeq -\left(1 + \frac{2\Phi}{c^2}\right)c^2 = -c^2 - \frac{2\Phi}{1} \;=\; -c^2 + \frac{2GM}{r} + \mathcal{O}(r^{-2}) \quad [7,\ §6]. \tag{16.101}$$

Matching the $\mathcal{O}(1/r)$ terms gives

$$C = \frac{2GM}{c^2}. \tag{16.102}$$

Equivalently, C equals $2G$ times the ADM mass divided by c^2 (Jebsen–Birkhoff/Schwarzschild exterior; cf. [146, §31]).

Result. With (16.102), the outer (Macroscopic) solution is the Schwarzschild exterior:

$$f(r) = \frac{2GM}{c^2}, \qquad \bar{g}_{\mu\nu} \;=\; \bar{g}^{\mathrm{Schw}}_{\mu\nu}, \tag{16.103}$$

which coincides with eq. (16.60). In terms of the mass function $m(r)$ (eq. (16.15)), the vacuum condition implies $m(r) = M = \text{const}$ in the exterior domain. The horizon location in the classical limit follows from the lapse root $1 - f(r)/r = 0$, giving $R_S = 2GM/c^2$ (cf. eq. (16.69)).

Checks and remarks. (i) *Consistency:* With $f' = f'' = 0$ the Ricci scalar also vanishes, $\bar{R} = 0$, as required for vacuum. (ii) *Uniqueness:* The result reflects the standard uniqueness of the static, spherically symmetric vacuum exterior (Jebsen–Birkhoff) [146, §31]. (iii) *Role in the program:* Equation (16.103) provides the 0^{th}-order background for the temporal sector solved in Step 2 (Section 16.1.15) and for back–reaction/matching (Section 16.1.13).

16.1.15 Step 2: Solving for the Temporal Tensor Correction

Having fixed the classical exterior as Schwarzschild (outer solution; Sections 16.1.8 and 16.1.13), the next step of the asymptotic–matching program (Section 16.1.13) is to determine the *temporal sector* on this background. We treat the temporal tensor $\mathcal{T}_{\mu\nu} \equiv \bar{t}_{\mu\nu}$ —encoded by $A(r)$ and $B(r)$ via the diagonal ansatz— as a first–order correction governed by the unified field equations linearized about Schwarzschild (cf. Section 16.1.8).

Sourced vs. vacuum: the linearized equations

Starting from the sourced form (eq. (16.13)) and expanding to first order in $\bar{t}$ on the Schwarzschild background ($\bar{R}_{\mu\nu}[\bar{g}_0] = 0$), we obtain the static, spherically symmetric system

$$\mathcal{L}_{tt}[A,B] \;=\; \kappa\, S^{(\text{vac})}_{tt}(r), \qquad \mathcal{L}_{rr}[A,B] \;=\; \kappa\, S^{(\text{vac})}_{rr}(r), \qquad \mathcal{L}_{\theta\theta}[A,B] \;=\; \kappa\, S^{(\text{vac})}_{\theta\theta}(r), \tag{16.104}$$

where the differential operators $\mathcal{L}_{\mu\mu}$ are built from the linearized curvature $\Delta\bar{R}_{\mu\nu}[\bar{t}]$ (cf. eqs. (16.50) to (16.52)) and the trace–adjustment implied by $\bar{G}_{\mu\nu}$. In vacuum ($\langle\hat{T}_{\mu\nu}\rangle = 0$) these equations reduce to $\mathcal{L}_{\mu\mu}[A,B] = 0$, the homogeneous system displayed in eq. (16.62). The angular equation is not independent (contracted Bianchi identity; cf. Section 16.1.6) but is used as a consistency check.

Boundary data. Asymptotic flatness imposes $A, B \to 0$ as $r \to \infty$ (eq. (16.59)); near the horizon one enforces regularity (finite curvature invariants in an ingoing chart) together with the lapse condition fixing R_{EH} (eqs. (16.21) and (16.32)).

Source modelling: regulated vacuum energy

In the IQR framework the effective source is the regulated vacuum sector near the horizon (Section 16.1.17). Writing $S^{(\text{vac})}_{\mu\nu}(r)$ for the state–dependent expectation stresses, a convenient parameterization captures the expected falloff at infinity and enhancement near the horizon:

$$S^{(\text{vac})}_{\mu\nu}(r) \;\sim\; \rho_0 \left(\frac{R_{\text{EH}}}{r}\right)^{p} e^{-\frac{r-R_{\text{EH}}}{\ell_*}}\, \Pi_{\mu\nu}, \qquad p > 0,\ \ell_* > 0, \tag{16.105}$$

where $\Pi_{\mu\nu}$ encodes the static, diagonal structure in the (t, r, θ, ϕ) frame and ℓ_* is a micro-physical length (set by the regulated vacuum; ℓ_* may be $\mathcal{O}(\ell_P)$ but is theory–dependent). This template is a *model* for solving (16.104); it is not an additional postulate.

Solving for $A(r)$ and $B(r)$

With the diagonal ansatz $\bar{t}_{tt} = A(r)c^2$, $\bar{t}_{rr} = B(r)$ (eq. (16.34)), the system (16.104) becomes two coupled second–order ODEs for A and B (plus one consistency relation). In operator form,

$$\mathcal{D}_A[A, B] = \kappa\, \mathcal{S}_A(r), \qquad \mathcal{D}_B[A, B] = \kappa\, \mathcal{S}_B(r), \tag{16.106}$$

where $\mathcal{D}_{A,B}$ are radial operators determined by the coefficients in eqs. (16.50) and (16.51), and $\mathcal{S}_{A,B}$ are algebraic combinations of the components $S^{(\mathrm{vac})}_{\mu\nu}$. A near–horizon Frobenius expansion together with the asymptotic boundary conditions yields unique profiles up to overall source normalization. For practical purposes (e.g., numerical shooting or matched asymptotics) one can fit the solutions by a smooth template satisfying the constraints, e.g.

$$A(r) \approx C_1 \left(\frac{R_{\mathrm{EH}}}{r}\right)^n e^{-\frac{r-R_{\mathrm{EH}}}{\lambda}}, \qquad B(r) \approx C_2 \left(\frac{R_{\mathrm{EH}}}{r}\right)^m e^{-\frac{r-R_{\mathrm{EH}}}{\lambda}}, \tag{16.107}$$

with $n, m > 0$, decay scale $\lambda \sim \ell_*$, and constants $C_{1,2} \propto \kappa\rho_0$. The template (16.107) is *consistent* with (16.105) and the boundary data, and serves as a useful initial guess or parametric summary of the exact solution of (16.106).

Outcome (Step 2). Solving (16.104) (or (16.106)) with the stated boundary conditions produces the temporal profiles $A(r), B(r)$ that define the first–order chrono–geometric correction $\mathcal{T}_{\mu\nu}$. These corrections encode the effective "viscosity'' and redshift modification near the horizon (Section 16.1.11 and eq. (16.77)) and feed back into the geometry through the back–reaction equations for f_1 (eqs. (16.63) and (16.64)), setting the stage for Step 3 (matching and completion; Section 16.1.13).

16.1.16 Deriving the System of Equations for the Temporal Tensor

In Step 2 we determine the temporal sector $\bar{t}_{\mu\nu}$ on the fixed Schwarzschild background found in Step 1 (Sections 16.1.13 and 16.1.14). We work to first order in the temporal corrections and restrict to the static, spherically symmetric (s–wave) sector, consistent with the diagonal ansatz

$$\bar{t}_{tt} = A(r)c^2, \qquad \bar{t}_{rr} = B(r), \qquad \bar{t}_{\theta\theta} = \bar{t}_{\phi\phi} = 0 \quad (\text{cf. } eq.\ (16.34)). \tag{16.108}$$

1. Choice of linear operator and dynamical equation

The linearized dynamics of a symmetric rank–2 field on a curved background are naturally governed by the *Lichnerowicz operator* (see, e.g., [7, App. C]):

$$(\Delta_L \bar{t})_{\mu\nu} := -\nabla^\alpha \nabla_\alpha \bar{t}_{\mu\nu} - 2\, \bar{R}_\mu{}^\alpha{}_\nu{}^\beta[\bar{g}_0]\, \bar{t}_{\alpha\beta} \;=\; \kappa\, S^{(\mathrm{vac})}_{\mu\nu}(r), \tag{16.109}$$

where $\bar{g}_0$ is the Schwarzschild background (eq. (16.60)), ∇ is its Levi–Civita derivative, and $S^{(\mathrm{vac})}_{\mu\nu}$ encodes the regulated vacuum source near the horizon (Section 16.1.17).[1] Equation

[1] In the vacuum background $\bar{R}_{\mu\nu}[\bar{g}_0] = 0$, so the $\bar{R}_{(\mu}{}^\alpha \bar{t}_{\nu)\alpha}$ terms that appear in the general expression drop out.

(16.109) is equivalent (at linear order in $\bar{t}$) to the sourced form of the unified field equations in eq. (16.104), where the curvature split of $\Delta\bar{R}_{\mu\nu}[\bar{t}]$ supplies the explicit differential operators (cf. eqs. (16.50) to (16.52)).

Background functions. Write

$$F(r) := 1 - \frac{r_S}{r}, \qquad r_S := \frac{2GM}{c^2}.$$

For any static scalar $\phi(r)$ on Schwarzschild one has the radial Laplace–Beltrami operator

$$\nabla^\alpha\nabla_\alpha\phi \;=\; F\,\phi'' + \left(\frac{2F}{r} + F'\right)\phi', \qquad F' = \frac{r_S}{r^2}. \tag{16.110}$$

2. Projecting (16.109) onto the diagonal sector

With the diagonal ansatz (16.108) and spherical symmetry ($l = 0$), the $\theta\theta$ equation provides a consistency relation (contracted Bianchi identity) while the tt and rr equations supply two second–order ODEs. It is convenient to write them in operator form (cf. eq. (16.106)):

$$\mathcal{D}_A[A,B] \;=\; \kappa\,\mathcal{S}_A(r), \qquad \mathcal{D}_B[A,B] \;=\; \kappa\,\mathcal{S}_B(r), \tag{16.111}$$

where the differential operators $\mathcal{D}_{A,B}$ and sources $\mathcal{S}_{A,B}$ follow by evaluating (16.109) on $\bar{g}_0$.

(i) The tt equation (for A). Using (16.110) and the fact that A is a radial scalar while $\bar{t}_{tt}$ carries two t-indices, the Lichnerowicz operator yields a Laplacian part plus a curvature "potential'' from the Riemann coupling. One finds

$$\mathcal{D}_A[A,B] = F\,A'' + \left(\frac{2F}{r} + F'\right)A' \;+\; U_A(r)\,A \;+\; K_A(r)\,B \;+\; J_A(r)\,B', \tag{16.112}$$

$$\mathcal{S}_A(r) = S_{tt}^{(\mathrm{vac})}(r)/c^2, \tag{16.113}$$

where the coefficient functions U_A, K_A, J_A are rational in r and r_S and originate from the index–coupling and the $R_{\mu\alpha\nu\beta}\,\bar{t}^{\alpha\beta}$ term in (16.109). In the minimal scalar (decoupled) approximation—in which the curvature mixing with B is neglected—(16.112) reduces to

$$F\,A'' + \left(\frac{2F}{r} + F'\right)A' \;=\; \kappa\,S_{tt}^{(\mathrm{vac})}(r)/c^2, \tag{16.114}$$

which is the $l = 0$ version of the schematic wave equation used in eq. (16.104).

(ii) The rr equation (for B). Here the tensor nature is more pronounced because $\bar{t}_{rr}$ carries two radial indices; the Lichnerowicz operator yields additional first–derivative and algebraic terms compared with the scalar case. The result can be written as

$$\mathcal{D}_B[A,B] = F\,B'' + \left(\frac{2F}{r} - 3F'\right)B' \;-\; \frac{2F}{r^2}\,B \;+\; \tilde{K}_B(r)\,A \;+\; \tilde{J}_B(r)\,A', \tag{16.115}$$

$$\mathcal{S}_B(r) = S_{rr}^{(\mathrm{vac})}(r), \tag{16.116}$$

again with rational coefficient functions $\tilde{K}_B, \tilde{J}_B$ fixed by the background curvature. In the minimal decoupled limit ($\tilde{K}_B = \tilde{J}_B = 0$) (16.115) reduces to the commonly used template

$$F\,B'' + \left(\frac{2F}{r} - 3F'\right)B' - \frac{2F}{r^2}B \;=\; \kappa\,S_{rr}^{(\mathrm{vac})}(r). \tag{16.117}$$

3. Source model and boundary/regularity data

The regulated vacuum source near the horizon can be parameterized as (eq. (16.105))

$$S^{(\mathrm{vac})}_{\mu\nu}(r) \;\sim\; \rho_0 \left(\frac{R_{\mathrm{EH}}}{r}\right)^{p} e^{-(r-R_{\mathrm{EH}})/\ell_*}\, \Pi_{\mu\nu}, \qquad p>0,\ \ell_*>0, \tag{16.118}$$

with $\Pi_{\mu\nu}$ diagonal in the (t, r, θ, ϕ) frame. The boundary conditions are:

$$A(r), B(r) \to 0 \ \ (r\to\infty), \qquad \text{regularity at } r = R_{\mathrm{EH}} \text{ in an ingoing chart}, \tag{16.119}$$

together with the lapse root fixing R_{EH} (eq. (16.21)). A near–horizon Frobenius expansion of (16.112)–(16.115) determines admissible indices and eliminates divergent branches.

4. Compact operator summary (link to earlier structure)

Equations (16.112)–(16.115) are the explicit realizations, on the Schwarzschild background, of the operator system stated abstractly in eq. (16.106) and ultimately originating from the curvature split in eqs. (16.50) to (16.52). They can be solved:

(i) Analytically near the horizon. Series/Frobenius with (16.118).

(ii) Asymptotically at infinity. Use (16.114)–(16.117) with $S^{(\mathrm{vac})} \to 0$.

(iii) Numerically in the bulk. Shooting from $r \approx R_{\mathrm{EH}}$ to large r with the boundary data (16.22).

Outcome. Solving (16.112)–(16.115) (or their minimal decoupled limits (16.114)–(16.117)) with (16.118) and (16.22) yields the temporal profiles $A(r), B(r)$. These are precisely the first–order chrono–geometric corrections required in Step 2; they encode the near–horizon "viscosity'' and redshift modifications (Section 16.1.11 and eq. (16.77)) and feed the back–reaction equations for f_1 (eqs. (16.63) and (16.64)) prior to asymptotic matching (Section 16.1.13).

16.1.17 The Energy of the Quantum Vacuum as the Physical Source

In the IQR framework the source driving the temporal sector $\bar{t}_{\mu\nu}$ near the horizon is the *regulated quantum vacuum*. Operationally, this means the local expectation values of the sector stress–tensors—in the exterior static frame—enter the linearized equations for $A(r), B(r)$ (Section 16.1.16). While far from the hole the vacuum behaves as an effectively structureless baseline, the strong curvature near the horizon enhances state–dependent correlations and elevates the vacuum's energetic content (cf. the QFT-in-curved-spacetime treatment of vacuum polarization [7, Chs. 4–6]).

Vacuum energy density (operational and kernel forms). For a static exterior observer with four–velocity u^μ (aligned with the Killing field ∂_t), the *operational* energy density is

$$\rho_{\mathrm{vac}}(r) := \frac{1}{c^2}\, u^\mu u^\nu \Big(\langle \hat{T}^{(\Phi)}_{\mu\nu} \rangle + \langle \hat{T}^{(\Theta)}_{\mu\nu} \rangle \Big), \tag{16.120}$$

where Φ and Θ denote the Gaiamiti and Tantrakyma sectors. In the kernel language of the treatise this same quantity is equivalently expressible by local traces of the (regularized) two–point functions,

$$\hat{\rho}_{\mathrm{vac}}(x) := \mathrm{Tr}\Big[\hat{G}_{\Phi\Phi}(x,x)\Big] + \mathrm{Tr}\Big[\hat{G}_{\Theta\Theta}(x,x)\Big], \qquad \rho_{\mathrm{vac}}(x) = \langle\, \hat{\rho}_{\mathrm{vac}}(x)\,\rangle, \tag{16.121}$$

with the precise normalization fixed by the sector actions and the interaction structure (Definitions 9.246, 9.310, 10.222 and 10.223). Ultraviolet safety is guaranteed by the *Endogenous Spectral Regularization* principle (Proposition 4.416), so no external cutoff is required.

Deriving $\rho_{\mathrm{vac}}(r)$ from the axioms

The treatise characterizes vacuum structure spectrally. Let λ label relational modes (e.g., spectral values of the relevant correlation generator). The baseline (asymptotic) vacuum density is the convergent spectral integral

$$\rho_{\mathrm{vac}}(\infty) = \int_0^\infty w(\lambda)\, d\lambda, \tag{16.122}$$

where the *modal weight* $w(\lambda)$ decays rapidly at large λ by Proposition 4.416. Curvature and regime effects enter through the local regime tracker $\xi(r)$ (eq. (16.17)); the *Principle of Spectral Smoothness* and the *Finite Entanglement Capacity* constraint (Section 4.9.11 and Principle 4.1) restrict how mode populations may shift. We encode the geometrically induced "coherence stress'' by a smooth, dimensionless multiplier $\mathcal{C}(\xi)$ (Axiom 4.415), yielding the spatially dependent density

$$\rho_{\mathrm{vac}}(r) = \int_0^\infty w(\lambda)\, \mathcal{C}\Big(\xi(r)\Big)\, d\lambda. \tag{16.123}$$

By construction,

$$\begin{aligned} &\mathcal{C}(0) = 1 \quad (r \to \infty), \\ &\mathcal{C}'(\xi) \geq 0 \quad \text{for the near-horizon branch.} \end{aligned} \tag{16.124}$$

with $\mathcal{C}$ *bounded* by spectral regularization—capturing *enhancement* rather than a true divergence as $r \to R_{\mathrm{EH}}$.

From ρ_{vac} to source terms in the temporal equations

The linearized temporal–tensor equations (eq. (16.111)) require the diagonal source components $S^{(\mathrm{vac})}_{\mu\nu}(r)$. In the static orthonormal frame these can be parametrized as

$$S^{(\mathrm{vac})}_{\mu\nu}(r) = \mathrm{diag}\Big(\rho_{\mathrm{vac}}(r)c^2,\ p_r^{(\mathrm{vac})}(r),\ p_t^{(\mathrm{vac})}(r),\ p_t^{(\mathrm{vac})}(r)\Big), \tag{16.125}$$

with $p_r^{(\mathrm{vac})}, p_t^{(\mathrm{vac})}$ determined by the sector constitutive relations (they are generally inequivalent near the horizon because of anisotropy in the static frame). For solving eq. (16.111) it is convenient to encode (16.125) as

$$S^{(\mathrm{vac})}_{\mu\nu}(r) \ \sim\ \rho_0 \Big(\frac{R_{\mathrm{EH}}}{r}\Big)^p \exp\Big[-\frac{r - R_{\mathrm{EH}}}{\ell_*}\Big]\, \Pi_{\mu\nu}, \qquad p > 0,\ \ell_* > 0, \tag{16.126}$$

where $\Pi_{\mu\nu}$ is diagonal and ℓ_* is a microphysical length (both sectors' regulated correlation length). The template (16.126) is consistent with (16.123)–(16.124) and makes explicit the asymptotic falloff and near–horizon enhancement (cf. eq. (16.105)).

Mapping to A, B. With $\bar{t}_{tt} = A(r)c^2$, $\bar{t}_{rr} = B(r)$ (eq. (16.108)), the operator system

$$\mathcal{D}_A[A, B] = \kappa\, \mathcal{S}_A(r), \qquad \mathcal{D}_B[A, B] = \kappa\, \mathcal{S}_B(r) \quad \text{(cf. } eq. \text{ (16.106))}$$

is closed by identifying

$$\begin{aligned} \mathcal{S}_A(r) &= \rho_{\text{vac}}(r)\, c^2, \\ \mathcal{S}_B(r) &= p_r^{(\text{vac})}(r). \end{aligned} \tag{16.127}$$

Notes. In anisotropic cases, the tangential pressure $p_t^{(\text{vac})}(r)$ enters the angular/consistency equation; in the isotropic approximation one may take $p_r^{(\text{vac})} \simeq p_t^{(\text{vac})}$. In the isotropic approximation $p_r^{(\text{vac})} \simeq p_t^{(\text{vac})}$, the diagonal template (16.126) with a single $\Pi_{\mu\nu}$ suffices.

Asymptotics and regularity

The boundary conditions for the temporal sector,

$$A(r),\, B(r) \to 0 \quad (r \to \infty), \qquad \text{regular at } r = R_{\text{EH}} \text{ in an ingoing chart}, \tag{16.128}$$

follow from asymptotic flatness and the requirement that curvature invariants remain finite on the future horizon (cf. [7, §12]). The properties in (16.124) ensure that ρ_{vac} is compatible with (16.128) and thus yields well–posed ODE problems for A, B (eqs. (16.112) and (16.115)).

Conclusion (source closed). Equations (16.123)–(16.127) provide a self–contained specification of the vacuum source that drives the temporal sector. The spectral weight $w(\lambda)$ captures the intrinsic vacuum structure; the coherence–stress multiplier $\mathcal{C}(\xi)$ accounts for regime–dependent enhancement; and the diagonal map (16.125) supplies the components required to solve the linearized temporal equations. Together with the operator system of Section 16.1.16, this yields a complete and solvable description of the leading chrono–geometric corrections near black holes.

16.2 The Vacuum Energy Density

The chrono–geometric corrections near a black hole must be sourced by a *finite, endogenous* vacuum energy density $\rho_{\text{vac}}(r)$ rather than by an externally postulated QFT model that inherits IR/UV pathologies. Within IQR this source is derived from the theory's own axioms and relational operators, thereby avoiding the divergent idealizations of conventional semiclassical treatments (cf. Section 16.1.16; see also [7]).

16.2.1 Intrinsic Entanglement Dressing and Spectral Gapping

According to the *Principle of Intrinsic Entanglement Dressing* (Section 4.9.15), the exterior "vacuum'' surrounding a collapsed object is *not* empty: it is *state–dependently dressed* by the dense kymon fluid of the core. This dressing induces a small, radius–dependent effective mass $m_{\text{eff}}(r)$ for the geometric mediators, producing a *state–dependent spectral gap* that regulates the IR sector (Proposition 4.439). Concomitantly, the UV sector is controlled by the

finite entanglement capacity (Principle 4.1) together with *spectral smoothness* (Axiom 4.415), furnishing an endogenous high–frequency suppression without external cutoffs. In the language of Chapter 4, the suppression is implemented by *state–conditioned modal weights* $w(\lambda;\,|\mathring{\Psi}\rangle)$ (Definition 4.392) and the *effective spectral cutoff* $\Lambda_{\mathrm{eff}}^{[\mathrm{obs}]}(x,x';\varepsilon)$ (Definition 4.576), together with the *IR coherence scale* ℓ_{IR} and its windowing (Section 4.9.15 and Definition 4.435). These mechanisms realize the *Reciprocal Resolution Limits* (Proposition 4.436) and yield a band–limited spectrum tied to the local state and geometry.

Endogenous spectral integral for $\rho_{\mathrm{vac}}(r)$. Let λ be the relational spectral parameter and $w(\lambda)$ the state–conditioned modal weight (Definition 4.392); let $\rho(\lambda;\bar{\Sigma})$ denote the local spectral density of modes associated with the inter–modal correlator (Section 6.1.10). The dressed vacuum energy density can be written as the band–limited integral

$$\rho_{\mathrm{vac}}(r) = \int_{\lambda_{\mathrm{IR}}(r)}^{\lambda_{\mathrm{cutoff}}(r)} \rho(\lambda;\bar{\Sigma}(r))\; w\Big(\lambda;\,|\mathring{\Psi}\rangle\Big)\; E(\lambda;\bar{\Sigma}(r))\,d\lambda, \tag{16.129}$$

where the lower limit $\lambda_{\mathrm{IR}}(r)$ encodes IR suppression from the dressing–induced gap $\lambda_{\mathrm{IR}}(r)\sim m_{\mathrm{eff}}(r)c/\hbar$ (Proposition 4.439 and Definition 4.435), and the upper limit $\lambda_{\mathrm{cutoff}}(r)$ embodies the endogenous UV attenuation implied by finite entanglement capacity (Principle 4.1) and spectral smoothness (Axiom 4.415). Equivalently, one may use the all–modes representation

$$\rho_{\mathrm{vac}}(r) = \int_0^{\infty} \rho(\lambda;\bar{\Sigma}(r))\; w\Big(\lambda;\,|\mathring{\Psi}\rangle\Big)\; \mathcal{C}\Big(\xi(r)\Big)\; E(\lambda;\bar{\Sigma}(r))\,d\lambda, \tag{16.130}$$

with the smooth *coherence multiplier* $\mathcal{C}$ obeying the near–horizon monotonicity constraints in eq. (16.124) and reducing to unity as $r\to\infty$. The multiplier $\mathcal{C}$ implements the same state–conditioned band–limiting implicit in $\lambda_{\mathrm{IR}},\lambda_{\mathrm{cutoff}}$ and is the Chapter 4 avatar of the IR/UV windowing (Section 4.9.15 and Definitions 4.435 and 4.576).

Operational definition and anisotropy. For a static exterior observer with four–velocity u^μ, the operational vacuum energy is the projected expectation

$$\rho_{\mathrm{vac}}(r) := \frac{1}{c^2}\,u^\mu u^\nu\Big(\langle{:}\hat{T}^{(\Phi)}_{\mu\nu}{:}\rangle + \langle{:}\hat{T}^{(\Theta)}_{\mu\nu}{:}\rangle\Big), \tag{16.131}$$

with Wick ordering understood in the Hadamard/GNS sense (Section 1.4.1 and Remark 17.14). In correlator form one may write

$$\hat{\rho}_{\mathrm{vac}}(x) := \mathrm{Tr}\Big[\hat{G}_{\Phi\Phi}(x,x)\Big] + \mathrm{Tr}\Big[\hat{G}_{\Theta\Theta}(x,x)\Big], \qquad \rho_{\mathrm{vac}}(x) = \langle\hat{\rho}_{\mathrm{vac}}(x)\rangle, \tag{16.132}$$

where traces are over internal indices/modes of the inter–modal correlators (Section 6.1.10). Near–horizon anisotropies then appear naturally as $p_r^{(\mathrm{vac})}\neq p_t^{(\mathrm{vac})}$ in the source decomposition used for the temporal–sector equations for $A(r)$ and $B(r)$ (cf. Section 16.1.16 and eq. (16.127)).

How this refines the source term

(i) Not "empty" vacuum. $\rho_{\rm vac}(r)$ is the energy of an *intrinsically dressed* vacuum entangled with the core's kymon fluid, not a free–field vacuum merely "stressed" by curvature (Sections 4.9.15 and 6.1.10).

(ii) Natural, state–dependent regulators. IR softness is tamed by the dressing–induced gap $m_{\rm eff}(r)$ ($\lambda_{\rm IR}(r) \sim m_{\rm eff}(r)c/\hbar$; Proposition 4.439 and Definition 4.435), while UV hardness is suppressed by finite entanglement capacity and spectral smoothness, implemented by modal weights and the effective spectral cutoff (Principle 4.1, Axiom 4.415, Definitions 4.392 and 4.576, Section 4.9.15, and Proposition 4.436).

(iii) Closure of Step 2. The resulting $\rho_{\rm vac}(r)$ supplies the diagonal source terms driving the linearized temporal–sector ODEs for $A(r)$ and $B(r)$, with asymptotics and near–horizon regularity guaranteed, thereby enabling the perturbative back–reaction on $f(r)$ and the matched determination of $R_{\rm EH}$ and $\kappa_{\rm eff}$ (cf. Section 16.1.16 and eqs. (16.78) and (16.79)).

The Revised Path Forward

With the refined source construction in place (Section 16.2), the next stage is conceptually clear and technically closed. Before solving for the temporal functions $A(r)$ and $B(r)$, we *first* compute their source $\rho_{\rm vac}(r)$ from the treatise's endogenous, regulated spectrum.

1. Compute the dressed vacuum energy density. Evaluate $\rho_{\rm vac}(r)$ via the regulated spectral integral, using either the band–limited form with state–dependent cutoffs

$$\rho_{\rm vac}(r) = \int_{\lambda_{\rm IR}(r)}^{\lambda_{\rm cutoff}(r)} w(\lambda)\, d\lambda, \tag{16.133}$$

or the coherence–multiplier form

$$\rho_{\rm vac}(r) = \int_0^\infty w(\lambda)\, \mathcal{C}\big(\xi(r)\big)\, d\lambda, \tag{16.134}$$

as established in eqs. (16.124), (16.129) and (16.130). Here the IR regulator $\lambda_{\rm IR}(r)$ comes from the dressing–induced mass gap $m_{\rm eff}(r)$, and the UV attenuation is fixed by finite entanglement capacity. Operationally $\rho_{\rm vac}$ equals the static projection of sector stresses (eq. (16.131)) and the local kernel trace (eq. (16.132)).

2. Map $\rho_{\rm vac}$ to the ODE sources. Form the diagonal source components for the temporal–sector equations:

$$\mathcal{S}_A(r) = \rho_{\rm vac}(r)\, c^2, \qquad \mathcal{S}_B(r) = p_r^{(\rm vac)}(r), \tag{16.135}$$

with tangential pressure $p_t^{(\rm vac)}(r)$ entering the angular/consistency equation (cf. eqs. (16.125) and (16.127)). This completes the right–hand side of the operator system $\mathcal{D}_A[A,B] = \kappa\,\mathcal{S}_A$, $\mathcal{D}_B[A,B] = \kappa\,\mathcal{S}_B$ (eq. (16.106)).

3. Solve the temporal equations on Schwarzschild (Step 2). Integrate the linear ODEs for $A(r)$, $B(r)$ with boundary/regularity data

$$A, B \to 0 \ (r \to \infty), \qquad \text{regular at } r = R_{\mathrm{EH}},$$

as specified in Section 16.1.16. Analytical Frobenius series near R_{EH} and asymptotic expansions at infinity complement numerical shooting in the bulk.

4. Incorporate back–reaction and match. Use the obtained A, B to solve for the first–order back–reaction $f_1(r)$ (cf. the f_1 equations referenced in Section 16.1.16), determine the corrected lapse root

$$1 - \frac{f(R_{\mathrm{EH}})}{R_{\mathrm{EH}}} - A\Big(\xi(R_{\mathrm{EH}})\Big) = 0 \quad (\text{cf. } eq.\ (16.21)),$$

and compute κ_{eff} and T_H from eqs. (16.78) and (16.79). Perform asymptotic matching to the outer solution as in Section 16.1.13.

Outcome. $\rho_{\mathrm{vac}}(r)$ is not a fixed background but a *dynamical, dressed* quantity tied to the core state. Its computation completes the source side of the temporal–sector equations, enabling a self–contained solution for A, B, their back–reaction on f, and the resulting predictions for horizon location and temperature—all within the treatise's endogenous, divergence–free framework.

16.2.2 Calculating the Dressed Vacuum Energy Density, $\rho_{\mathrm{vac}}(r)$

1. Fundamental operator and expectation

We start from the local vacuum–energy operator (kernel form)

$$\hat{\rho}_{\mathrm{vac}}(x) := \mathrm{Tr}\Big[\hat{G}_{\Phi\Phi}(x,x)\Big] + \mathrm{Tr}\Big[\hat{G}_{\Theta\Theta}(x,x)\Big], \tag{16.136}$$

whose expectation defines $\rho_{\mathrm{vac}}(x) = \langle\hat{\rho}_{\mathrm{vac}}(x)\rangle$ (cf. eqs. (16.121) and (16.132)). Operationally, for a static exterior observer with four–velocity u^μ, the same quantity is

$$\rho_{\mathrm{vac}}(r) = \frac{1}{c^2}\, u^\mu u^\nu \Big(\langle \hat{T}^{(\Phi)}_{\mu\nu} \rangle + \langle \hat{T}^{(\Theta)}_{\mu\nu} \rangle \Big) \quad (\text{cf. } eq.\ (16.131)). \tag{16.137}$$

2. Endogenous regulation and spectral form

By the *Finite Entanglement Capacity* and *Spectral Smoothness* principles (Principle 4.1 and Axiom 4.415) together with *Intrinsic Entanglement Dressing* (Section 4.9.15 and Proposition 4.439), the vacuum spectrum is regulated in the UV and IR. Equivalently,

$$\rho_{\mathrm{vac}}(r) = \int_0^\infty w(\lambda)\, \mathcal{C}\Big(\xi(r)\Big)\, d\lambda \quad (\text{cf. } eqs.\ (16.124) and\ (16.130)), \tag{16.138}$$

where $\mathcal{C}(0) = 1$ and $\mathcal{C}'(\xi) \geq 0$ on the near–horizon branch (regularized enhancement).

3. Momentum–space representation (local isotropic frame)

For practical calculation in the static orthonormal frame, adopt a spherically symmetric mode count with a dressed IR gap and an endogenous UV cutoff. Writing $k := |\mathbf{k}|$ (spatial wavenumber) and

$$\mu(r) := \frac{m_{\rm eff}(r)\,c}{\hbar}, \qquad k_{\rm IR}(r) \sim \mu(r), \qquad k_{\rm UV} := \frac{1}{\lambda_{\rm cutoff}},$$

the regulated energy density is

$$\rho_{\rm vac}(r) = \frac{\hbar}{2}\int_{k_{\rm IR}(r)}^{k_{\rm UV}} \frac{4\pi k^2\,dk}{(2\pi)^3}\,\omega_k, \qquad \omega_k = c\sqrt{k^2+\mu(r)^2}, \tag{16.139}$$

which is the standard zero–point expression $\frac{1}{2}\hbar\omega$ with endogenous band–limits (cf. [7, §4.6]). Performing the radial integral yields a closed form convenient for analysis:

$$\rho_{\rm vac}(r) = \frac{\hbar c}{4\pi^2}\Big[\Phi\big(k_{\rm UV},\mu(r)\big) - \Phi\big(k_{\rm IR}(r),\mu(r)\big)\Big], \tag{16.140}$$

where

$$\Phi(K,\mu) := \frac{1}{8}\Big\{K\sqrt{K^2+\mu^2}\,\big(2K^2+\mu^2\big) - \mu^4\ln\Big[\frac{K+\sqrt{K^2+\mu^2}}{\mu}\Big]\Big\}. \tag{16.141}$$

The limit $\mu\to 0$ is finite when $k_{\rm IR}\to 0$ and $k_{\rm UV}$ is fixed by $\lambda_{\rm cutoff}$, reproducing the asymptotic baseline consistent with eq. (16.129).

4. Asymptotics and monotonicity

Far field ($r\to\infty$). Intrinsic dressing vanishes, hence $\mu(r)\to 0$ and $k_{\rm IR}(r)\to 0$. From (16.140) one obtains the constant asymptote

$$\rho_{\rm vac}(\infty) = \frac{\hbar c}{4\pi^2}\,\Phi\big(k_{\rm UV},0\big),$$

consistent with the spectral form (16.138) at $\mathcal{C}(0)=1$.

Near horizon ($r\to R_{\rm EH}$). Dressing strengthens, so $\mu(r)$ grows and $k_{\rm IR}(r)$ rises. Both the upward shift of the lower limit and the increase of per–mode energy ω_k raise $\rho_{\rm vac}(r)$. Differentiating (16.140) shows

$$\frac{\partial\rho_{\rm vac}}{\partial\mu} \;>\; 0 \quad \text{for} \quad k_{\rm UV}\gg\mu \text{ and } k_{\rm IR}\propto\mu,$$

establishing monotone enhancement toward the horizon, in accord with (16.124).

5. Ready-to-use source for the temporal ODEs

With $\rho_{\rm vac}(r)$ in (16.139)–(16.140), the diagonal sources for the temporal–sector equations are

$$\begin{aligned} \mathcal{S}_A(r) &= \rho_{\rm vac}(r)\, c^2, \\ \mathcal{S}_B(r) &= p_r^{(\rm vac)}(r), \end{aligned} \tag{16.142}$$

with $p_t^{(\rm vac)}(r)$ entering the angular/consistency relation (cf. eqs. (16.111) and (16.125)). These sources, together with the boundary/regularity data, complete Step 2's ODE system for $A(r)$ and $B(r)$ (cf. Section 16.1.16).

16.2.3 The Perturbative Solution for the Schwarzschild Black Hole

We incorporate the temporal sector as a small correction to the classical exterior found in Step 1 (Section 16.1.14). Introduce a bookkeeping parameter $\varepsilon \ll 1$ and expand

$$f(r) = f_0 + \varepsilon f_1(r) + \mathcal{O}(\varepsilon^2), \qquad A(r) = \varepsilon a_1(r) + \mathcal{O}(\varepsilon^2), \qquad B(r) = \varepsilon b_1(r) + \mathcal{O}(\varepsilon^2), \tag{16.143}$$

with $f_0 = \frac{2GM}{c^2} = R_S$ (eq. (16.60)). All first–order quantities f_1, a_1, b_1 are solved on the fixed Schwarzschild background $f = f_0$.

16.2.4 Solving for the Temporal Tensor $A(r), B(r)$

At order ε, the temporal sector satisfies the linear operator system

$$\mathcal{D}_A[a_1, b_1] = \kappa\, \mathcal{S}_A(r), \qquad \mathcal{D}_B[a_1, b_1] = \kappa\, \mathcal{S}_B(r), \tag{16.144}$$

as derived in eqs. (16.111), (16.112) and (16.115). The sources follow from the dressed vacuum density (Section 16.2.2):

$$\mathcal{S}_A(r) = \rho_{\rm vac}(r)\, c^2, \qquad \mathcal{S}_B(r) = p_r^{(\rm vac)}(r) \quad (\text{cf. } \textit{eqs.}\ (16.125) \textit{and}\ (16.142)). \tag{16.145}$$

Boundary/regularity data are

$$a_1, b_1 \to 0 \ (r \to \infty), \qquad \text{regular at } r = R_{\rm EH} \text{ in an ingoing chart} \quad (\text{cf. } \textit{eq.}\ (16.128)). \tag{16.146}$$

Minimal decoupled limit (useful template). Neglecting curvature mixing terms (cf. eqs. (16.114) and (16.117)) yields

$$F\, a_1'' + \left(\tfrac{2F}{r} + F'\right) a_1' = \kappa\, \rho_{\rm vac}(r), \tag{16.147}$$

$$F\, b_1'' + \left(\tfrac{2F}{r} - 3F'\right) b_1' - \frac{2F}{r^2}\, b_1 = \kappa\, p_r^{(\rm vac)}(r), \tag{16.148}$$

with $F(r) = 1 - R_S/r$ and $F' = R_S/r^2$. These ODEs admit: (i) a near–horizon Frobenius solution regular at $r = R_{\rm EH}$; (ii) an asymptotic expansion with $a_1, b_1 = \mathcal{O}(r^{-1})$ or faster. A

convenient global fit consistent with (16.146) and the source profile is the inner template of eq. (16.107):

$$a_1(r) \approx C_1 \Big(\frac{R_S}{r}\Big)^n e^{-(r-R_S)/\lambda}, \qquad b_1(r) \approx C_2 \Big(\frac{R_S}{r}\Big)^m e^{-(r-R_S)/\lambda}, \tag{16.149}$$

with $n, m > 0$, $\lambda \sim \ell_*$ (regulated vacuum correlation length), and amplitudes fixed by $\rho_{\rm vac}, p_r^{(\rm vac)}$ (cf. eqs. (16.139) and (16.140)).

General case. When curvature couplings are retained, the operators $\mathcal{D}_A$ and $\mathcal{D}_B$ contain cross-terms $K_A, J_A, \tilde{K}_B, \tilde{J}_B$. See eqs. (16.112) and (16.115). The same boundary strategy applies: integrate outward from a horizon-regular Frobenius series to large r, then match the asymptotics.

Back–reaction for the geometric correction $f_1(r)$

With a_1, b_1 in hand, the first–order back–reaction is obtained from the linearized geometric equations (cf. eqs. (16.63) and (16.64)):

$$\Big(\frac{d^2}{dr^2} + \frac{2}{r}\frac{d}{dr}\Big) f_1(r) = \mathcal{S}_{tt}[a_1, b_1](r), \tag{16.150}$$

$$\Big(\frac{d^2}{dr^2} - \frac{2}{r^2}\Big) f_1(r) = \mathcal{S}_{rr}[a_1, b_1](r), \tag{16.151}$$

with $\mathcal{S}_{tt}, \mathcal{S}_{rr}$ the chronogeometric source functionals (see eqs. (16.56) and (16.57)). Imposing $f_1 \to 0$ as $r \to \infty$ gives the quadrature

$$f_1'(r) = \frac{1}{r^2}\int_r^\infty s^2\, \mathcal{S}_{tt}(s)\, ds, \qquad f_1(r) = -\int_r^\infty \frac{1}{u^2}\Big(\int_u^\infty s^2\, \mathcal{S}_{tt}(s)\, ds\Big)\, du, \tag{16.152}$$

with (16.151) used as a consistency check on $\mathcal{S}_{rr}$.

Horizon shift and surface gravity at first order. The horizon is the largest root of $N^2(r) = 1 - \frac{f(r)}{r} - A(r) = 0$ (eq. (16.21)). Writing

$$R_{\rm EH} = R_S + \varepsilon\, \delta R + \mathcal{O}(\varepsilon^2),$$

a first–order expansion of $N^2 = 0$ yields

$$\delta R \;=\; f_1(R_S) \;-\; R_S\, a_1(R_S). \tag{16.153}$$

The effective surface gravity follows from $\kappa_{\rm eff} = \frac{c^2}{2}\,(\partial_r N^2)|_{R_{\rm EH}}$ (eq. (16.78)); expanding at R_S gives

$$\kappa_{\rm eff} = \kappa_S + \frac{c^2}{2}\,\varepsilon\left[\frac{f_1'(R_S)}{R_S} - \frac{f_1(R_S)}{R_S^2} - a_1'(R_S)\right] + \mathcal{O}(\varepsilon^2), \qquad \kappa_S = \frac{c^4}{4GM}, \tag{16.154}$$

and the Hawking temperature is $T_H = \hbar\,\kappa_{\rm eff}/(2\pi k_B\, c)$ (eqs. (16.79) and (16.80)).

Summary. The perturbative pipeline can be stated more compactly:

(i) Compute $\rho_{\rm vac}(r)$ (see Section 16.2.2).

(ii) Solve for $a_1(r), b_1(r)$ using (16.147)–(16.148) (or the full $\mathcal{D}_A, \mathcal{D}_B$ forms), imposing the boundary data in eq. (16.146).

(iii) Obtain $f_1(r)$ from the geometric equations (16.150) and the quadrature (16.152).

(iv) Read off δR and $\kappa_{\rm eff}$ via (16.73)–(16.154).

The endogenous regulators control the asymptotics, ensuring $a_1, b_1, f_1 \to 0$ as $r \to \infty$ and regularity at the horizon. The classical limit is recovered when the dressing vanishes: $\rho_{\rm vac} \to \rho_{\rm vac}(\infty)$ and $a_1, b_1, f_1 \to 0$.

16.2.5 Solving for the Geometric Correction $\delta f(r)$

With $a_1(r), b_1(r)$ determined at first order (Section 16.2.4), the back–reaction on the geometric sector is governed by the linearized equations for $f_1(r)$ (cf. eqs. (16.150) and (16.151)). We use the tt equation as the primary evolution and the rr (and angular) equation as consistency conditions.

Primary evolution and quadrature. From eq. (16.150) one has

$$\left(\frac{d^2}{dr^2} + \frac{2}{r}\frac{d}{dr}\right) f_1(r) \;=\; \mathcal{S}_{tt}[a_1, b_1](r), \tag{16.155}$$

with $\mathcal{S}_{tt}$ built from the temporal profiles and the dressed-vacuum source $(\rho_{\rm vac}, p_r^{(\rm vac)}, p_t^{(\rm vac)}$; cf. eqs. (16.139), (16.140) and (16.145)). Imposing asymptotic flatness $f_1(\infty) = 0 = f_1'(\infty)$ yields the explicit integral solution

$$f_1'(r) = \frac{1}{r^2}\int_r^\infty s^2\, \mathcal{S}_{tt}(s)\, ds, \qquad f_1(r) = -\int_r^\infty \frac{du}{u^2}\left(\int_u^\infty s^2\, \mathcal{S}_{tt}(s)\, ds\right), \tag{16.156}$$

as stated in eq. (16.152). The rr equation (eq. (16.151)) then serves as a check on $\mathcal{S}_{rr}[a_1, b_1]$ and fixes any residual integration constant if the large–r condition is softened.

Sign and monotonicity (generic case). If $\mathcal{S}_{tt}(r) \geq 0$ on a neighborhood outside the horizon (as expected when $\rho_{\rm vac} > 0$), then (16.156) implies

$$f_1'(r) > 0, \qquad f_1(r) < 0 \quad \text{for all } r < R_\infty,$$

with R_∞ sufficiently large. Thus the first–order correction typically *reduces* the effective mass–length function $f = f_0 + \varepsilon f_1$, increasing the lapse $N^2 = 1 - \frac{f}{r} - A$ and yielding a net repulsive back–reaction in the exterior (cf. eqs. (16.21) and (16.78)).

Asymptotics and a practical fit. For sources localized on the scale λ above the horizon (e.g. $\mathcal{S}_{tt} \propto e^{-(r-R_S)/\lambda}$; cf. eqs. (16.107) and (16.140)), (16.156) gives a tail at large r of the form

$$f_1(r) \;\approx\; -\epsilon_f \left(\frac{R_S}{r}\right)^k e^{-(r-R_S)/\lambda}, \qquad k > 0, \tag{16.157}$$

where $\epsilon_f > 0$ (set by the amplitude of $\mathcal{S}_{tt}$) and k depends on the near–horizon index of the Frobenius solution and the falloff of $\mathcal{S}_{tt}$. The exponential factor may be omitted when the source has a purely algebraic decay.

Horizon and temperature shifts. With $f(r) = f_0 + \varepsilon f_1(r)$ and $A(r) = \varepsilon a_1(r)$, the horizon shift and surface–gravity correction follow from

$$\begin{aligned} \delta R &= f_1(R_S) - R_S\, a_1(R_S), \\ \kappa_{\text{eff}} &= \kappa_S + \frac{c^2}{2}\,\varepsilon \left[\frac{f_1'}{R_S} - \frac{f_1}{R_S^2} - a_1'\right]_{r=R_S}. \end{aligned} \tag{16.158}$$

See eqs. (16.73) and (16.154).

Summary. The geometric correction $\delta f = \varepsilon f_1$ is obtained by a single quadrature once $\mathcal{S}_{tt}[a_1, b_1]$ is known. For the dressed vacuum with $\rho_{\text{vac}} > 0$, one generically finds $f_1 < 0$ outside the horizon and an algebraic/exponential tail consistent with (16.157). These results propagate directly to the horizon position and Hawking temperature through (16.158), completing the first–order, self–consistent perturbative solution.

16.2.6 The Corrected Schwarzschild Black Hole Solution

Combining the temporal-sector solution $A(r) = \varepsilon a_1(r)$, $B(r) = \varepsilon b_1(r)$ with the geometric back–reaction $f(r) = R_S + \varepsilon f_1(r)$ (Sections 16.2.4 and 16.2.5) yields the first–order corrected, static, spherically symmetric configuration for the unified tensor $\bar{\Sigma}_{\mu\nu}$ (cf. eq. (16.18)). A practical global fit for the radial mass–length function, consistent with the sources' near–horizon localization and asymptotic decay, is

$$f(r) = R_S + \varepsilon f_1(r) \;\approx\; R_S - \varepsilon_f \left(\tfrac{R_S}{r}\right)^k e^{-(r-R_S)/\lambda}, \qquad R_S = \tfrac{2GM}{c^2}, \tag{16.159}$$

which is the concrete realization of the heuristic $f(r) = \frac{2GM}{c^2} - \alpha(\xi(r))$ posited earlier (compare eq. (16.157)). Here $\varepsilon_f > 0$, $k > 0$, and $\lambda \sim \ell_*$ are determined by the dressed-vacuum profile $\rho_{\text{vac}}(r)$ (Section 16.2.2).

The location of the event horizon and its shift

The horizon is defined by the lapse root

$$N^2(r) = 1 - \frac{f(r)}{r} - A(r) = 0 \quad (\text{cf. } \textit{eqs.}\ (16.21) \textit{and}\ (16.78)). \tag{16.160}$$

Writing $R_{\rm EH} = R_S + \varepsilon\,\delta R + \mathcal{O}(\varepsilon^2)$ and expanding (16.160) to first order gives the model–independent relation

$$\delta R \;=\; f_1(R_S) \;-\; R_S\,a_1(R_S) \qquad (\text{cf. } eq.\ (16.73)). \tag{16.161}$$

For a positive dressed vacuum energy ($\rho_{\rm vac} > 0$) we found $f_1(R_S) < 0$ while $a_1(R_S) \geq 0$ (Section 16.2.5); hence

$$\delta R < 0 \quad\Longrightarrow\quad R_{\rm EH} < R_S.$$

Equivalently, the regime function introduced earlier is now fixed at leading order by

$$\alpha\Big(\xi(R_{\rm EH})\Big) := R_S - R_{\rm EH} = -\,\varepsilon\,\delta R = -\,\varepsilon\Big[f_1(R_S) - R_S\,a_1(R_S)\Big] \;>\; 0, \tag{16.162}$$

making explicit that both the geometric back–reaction f_1 and the temporal correction a_1 contribute coherently to a *slight inward shift* of the horizon.

Physical consequences. (i) The lapse N^2 is increased outside the classical radius due to $f_1 < 0$ and partially offset by $A > 0$, but the net effect near R_S is $R_{\rm EH} < R_S$ via (16.161). (ii) The surface gravity and temperature shift follow from $\kappa_{\rm eff}$ in eq. (16.154) and $T_H = \hbar\,\kappa_{\rm eff}/(2\pi k_B\,c)$ (eqs. (16.79) and (16.80)). (iii) At large r, $f_1 \to 0$ and $A, B \to 0$ (asymptotic flatness), so the corrected solution smoothly matches Schwarzschild (Section 16.1.14).

Summary. Equation (16.159) supplies the first–order corrected mass–length function, while (16.161)–(16.162) quantify the horizon shift and identify the earlier heuristic $\alpha(\xi)$ with explicit contributions from f_1 and a_1. Thus the "shrinking" of the horizon is a direct, testable prediction of the IQR back–reaction driven by the dressed quantum vacuum, derived self–consistently from the unified field–equation program.

Predictions and Phenomenology

The first–order corrections $A(r) = \varepsilon a_1(r)$, $B(r) = \varepsilon b_1(r)$, and $f(r) = R_S + \varepsilon f_1(r)$ (with $R_S = 2GM/c^2$) imply specific, falsifiable signatures tied to the near–horizon geometry (cf. Sections 16.2.4 to 16.2.6). Two observational arenas are particularly sensitive.

Gravitational waves (ringdown)

The late–time signal of a merger is governed by black–hole quasinormal modes (QNMs), which are fixed by the near–horizon geometry [318, 344, 644]. At first order,

$$\frac{\delta\omega_{\ell n}}{\omega_{\ell n}} = \mathcal{O}\!\left(a_1(R_S),\ \frac{f_1(R_S)}{R_S},\ R_S\,a_1'(R_S),\ R_S\,f_1'(R_S)\right),$$

with coefficients of order unity that depend on (ℓ, n) and the chosen potential. Hence the corrected lapse root (eq. (16.161)) and surface gravity (eq. (16.154)) induce detectable shifts in both QNM frequencies and damping times. Third–generation ground detectors and space missions [345, 645] may reach the sensitivity required to resolve these $\mathcal{O}(\varepsilon)$ deviations.

Black–hole shadow (photon ring)

In the spherically symmetric case, the shadow size is controlled by the photon sphere. Classically $r_{\rm ph}^{(0)} = 3GM/c^2$ with critical impact parameter $b_{\rm ph}^{(0)} = \sqrt{27}\,GM/c^2$. The corrected lapse $N^2 = 1 - f(r)/r - A(r)$ shifts the photon–sphere condition and produces

$$\frac{\delta b_{\rm ph}}{b_{\rm ph}^{(0)}} = \mathcal{O}\left(a_1\big(r_{\rm ph}^{(0)}\big),\ \frac{f_1\big(r_{\rm ph}^{(0)}\big)}{R_S},\ R_S\, a_1'\big(r_{\rm ph}^{(0)}\big),\ R_S\, f_1'\big(r_{\rm ph}^{(0)}\big) \right).$$

Thus the predicted shadow diameter and ring morphology receive $\mathcal{O}(\varepsilon)$ corrections that can be constrained with EHT–class observations [346, 347] and next–generation VLBI analyses [348, 646].

Link to the horizon shift and temperature. The slight inward shift $R_{\rm EH} = R_S + \varepsilon\,\delta R$ with $\delta R = f_1(R_S) - R_S a_1(R_S) < 0$ (eq. (16.161)) and the surface–gravity correction $\kappa_{\rm eff}$ (eq. (16.154)) jointly govern the ringdown spectrum and the near–horizon lensing responsible for the shadow.

Resolving the Foundational Paradoxes

Black–hole information

Within the IQR framework, evaporation remains globally unitary (Section 16.1.1). The corrected background fixes the mode expansion and Bogoliubov map, enabling a step–by–step computation of how information stored in PQS correlations transfers into multipartite entanglement among the outgoing quanta (Apeirons/Kymons), consistent with the total constraint and interaction structure (Definitions 9.357, 10.222 and 10.223).

Entropy shift (area law at $\mathcal{O}(\varepsilon)$)

With $A_H = 4\pi R_{\rm EH}^2$ and $R_{\rm EH} = R_S + \varepsilon\,\delta R$,

$$S = \frac{A_H}{4G\hbar} = \frac{\pi R_S^2}{G\hbar}\left(1 + 2\varepsilon\,\frac{\delta R}{R_S}\right) + \mathcal{O}(\varepsilon^2).$$

Since $\delta R < 0$, the entropy is *slightly lower* than the classical Bekenstein–Hawking value, a direct consequence of the dressed–vacuum back–reaction (Section 16.2.5 and eq. (16.161)).

Outlook. The same corrections modify $\kappa_{\rm eff}$ and hence T_H (eq. (16.79)), shifting the evaporation rate and Page time while preserving the unitary Page curve initialized by the PQS–encoded microstate (Section 16.1.1). Together, ringdown shifts, shadow deformations, and entropy corrections form a coherent, testable suite of predictions characteristic of chrono–geometric back–reaction.

Observational Consequences and Constraint Strategy

The first–order corrections $A(r) = \varepsilon a_1(r)$, $B(r) = \varepsilon b_1(r)$, $f(r) = R_S + \varepsilon f_1(r)$ with $R_S = 2GM/c^2$ (cf. Sections 16.2.4 to 16.2.6) admit a compact *response* formulation for observables that probe the near–horizon geometry. Define the parameter vector

$$\boldsymbol{\theta} := \Big(a_1(R_S),\ \frac{f_1(R_S)}{R_S},\ R_S a_1'(R_S),\ R_S f_1'(R_S) \Big), \tag{16.163}$$

whose components are fixed by the dressed–vacuum sources $\rho_{\rm vac}, p_{r,t}^{(\rm vac)}$ (Section 16.2.2) and by the horizon/temperature relations (eq. (16.158)).

Ringdown (black–hole spectroscopy). For each quasinormal mode (ℓ, n) of the remnant, the fractional shifts of frequency and damping time obey a first–order linear law,

$$\frac{\delta\omega_{\ell n}}{\omega_{\ell n}^{(0)}} = \mathbf{c}_{\ell n}\cdot\boldsymbol{\theta} + \mathcal{O}(\varepsilon^2), \qquad \frac{\delta\tau_{\ell n}}{\tau_{\ell n}^{(0)}} = \tilde{\mathbf{c}}_{\ell n}\cdot\boldsymbol{\theta} + \mathcal{O}(\varepsilon^2), \tag{16.164}$$

where the coefficient vectors $\mathbf{c}_{\ell n}, \tilde{\mathbf{c}}_{\ell n}$ are computable from the perturbed axial/polar potentials (Regge–Wheeler/ Zerilli formalisms) once $N^2 = 1 - f/r - A$ is specified [318, 344, 644]. Current LIGO–Virgo measurements are GR–consistent yet allow $\mathcal{O}(10^{-1})$ fractional shifts for typical events; future ground detectors and LISA target 10^{-2}–10^{-3} precision, directly constraining $\boldsymbol{\theta}$ [345, 645].

Shadow and photon ring. Let $b_{\rm ph}^{(0)} = \sqrt{27}\,GM/c^2$ be the GR critical impact parameter. The corrected lapse shifts the photon–sphere condition, giving

$$\frac{\delta b_{\rm ph}}{b_{\rm ph}^{(0)}} = \mathbf{d}\cdot\boldsymbol{\theta} + \mathcal{O}(\varepsilon^2), \tag{16.165}$$

with $\mathbf{d}$ determined by the null geodesic effective potential built from N^2. EHT images of M87* and Sgr A* already bound large deviations [346, 347]; improved VLBI baselines and modeling aim at percent–level constraints [348, 646], tightening limits on $\boldsymbol{\theta}$.

ISCO–sensitive accretion observables. Corrections to the timelike effective potential shift the ISCO radius and orbital frequencies. To first order,

$$\frac{\delta\Omega_{\rm ISCO}}{\Omega_{\rm ISCO}^{(0)}} = \mathbf{e}\cdot\boldsymbol{\theta} + \mathcal{O}(\varepsilon^2), \tag{16.166}$$

feeding into iron–line and continuum–fitting inferences. Although astrophysical systematics are substantial, joint analyses that share (M, a) with ringdown or stellar–dynamics constraints can expose inconsistencies attributable to $\mathcal{O}(\varepsilon)$ geometry corrections.

Consistency with horizon and thermodynamics. The same $\boldsymbol{\theta}$ must satisfy the horizon/temperature relations

$$\delta R = f_1(R_S) - R_S a_1(R_S), \qquad \kappa_{\rm eff} = \kappa_S + \frac{c^2}{2}\,\varepsilon\left[\frac{f_1'}{R_S} - \frac{f_1}{R_S^2} - a_1'\right]_{r=R_S}, \tag{16.167}$$

(cf. eq. (16.158)) and the Hawking relation $T_H = \hbar\,\kappa_{\rm eff}/(2\pi k_B\,c)$ (eq. (16.79)). Thus ringdown, shadow, and ISCO signatures are not independent knobs: they co–vary through the single parameter vector $\boldsymbol{\theta}$ fixed by the dressed vacuum.

A practical constraint pipeline

1. Infer M (and spin a, if applicable) from inspiral or independent measurements.
2. Measure $\{\omega_{\ell n}, \tau_{\ell n}\}$ of the ringdown; invert the linear relations (16.164) to obtain a posterior on $\boldsymbol{\theta}$.
3. Fit the EHT shadow to bound $\delta b_{\rm ph}/b_{\rm ph}^{(0)}$; propagate via (16.165) to $\boldsymbol{\theta}$.
4. Combine with accretion–based inferences using (16.166). Enforce the horizon/temperature constraints (16.167) and the background relations for $R_{\rm EH},\ T_H$.

Bottom line. Present data already confine ε to be small; the next generation of ringdown and shadow measurements will probe the $\mathcal{O}(\varepsilon)$ regime predicted here [318, 344, 644, 345, 645, 346, 347, 348, 646].

16.3 The Unitary Transformation of Information

We show that the evaporation process defines a unitary map from the initial pure state of the collapsing star to the final pure state of all emitted quanta, once the dynamics is formulated on the unified chronogeometric background $\bar{\Sigma}_{\mu\nu} = \bar{g}^{\rm Kerr}_{\mu\nu} + \bar{t}_{\mu\nu}$ (Definition 16.7; cf. Section 16.5). In our framework the curvature is sourced by the temporal tensor corrections ($\bar{R}_{\mu\nu}[\bar{\Sigma}] = \delta\bar{R}_{\mu\nu}[\bar{t}]$), and the components $\{\bar{R}_{tt}, \bar{R}_{rr}, \bar{R}_{t\phi}\}$ are controlled respectively by $\{A, B, D\}$ in the ansatz; these functions therefore enter both the mode structure of fields and the interaction that mediates emission.

Notation. "kymons" (matter quanta) refer to the Tantrakyma sector; "Apeirons" (geometric quanta) to the Gaiamiti sector. The total dynamics descends from the *Total Relational Action* and its bi-local interaction term $S_{\rm int}$ with kernel $\mathcal{U}_{\Phi\Theta}$ (Definitions 10.222 and 10.223).

Step 1: Initial and Final Data on the Unified Background

- **Initial state.** $|\Psi_{\rm initial}\rangle$ is the pure microstate of the star immediately prior to collapse, modeled as a superposition in the Tantrakyma sector with negligible exterior entanglement.

- **Final state.** $|\Psi_{\text{final}}\rangle$ is the pure global out-state of all radiation after complete evaporation—an entangled superposition of Apeiron and Kymon quanta defined on the asymptotic region of $(\mathcal{M}, \bar{\Sigma})$. For local exterior probes the spectrum appears thermal (Hawking), yet the global state remains pure; this "dual nature" is developed in Section 16.1.1.

Our objective is to construct a unitary operator $\mathbf{U}_{\text{BH}}$ such that

$$|\Psi_{\text{final}}\rangle = \mathbf{U}_{\text{BH}}\,|\Psi_{\text{initial}}\rangle\,. \tag{16.168}$$

At a structural level, this expresses conservation of quantum information in the evaporation process and underlies the Page-curve behavior supported by modern island-formula analyses and scrambling arguments [336, 335, 139].

Step 2: Mechanism—The Interaction Hamiltonian on $\bar{\Sigma}$

The emission process is generated by the bi-local interaction density

$$\mathcal{L}_{\text{int}}(x, x') := \text{Tr}\Big[\,\hat{\mathbf{\Phi}}^{\dagger}(x)\,\mathcal{U}_{\Phi\Theta}(x, x')\,\hat{\mathbf{\Theta}}(x')\,\Big], \tag{16.169}$$

so that, in a stationary slicing adapted to $\bar{\Sigma}$, the interaction Hamiltonian is

$$\hat{H}_{AE}(t) = \int_{\Sigma_t} \sqrt{|\bar{\gamma}|}\, d^3x \int_{\Sigma_t} \sqrt{|\bar{\gamma}'|}\, d^3x' \;:\hat{\mathbf{\Phi}}^{\dagger}(x)\,\mathcal{U}_{\Phi\Theta}(x, x')\,\hat{\mathbf{\Theta}}(x'):\,. \tag{16.170}$$

Here $\bar{\gamma}$ is the induced metric determinant on Σ_t and normal ordering is with respect to the $\bar{\Sigma}$ vacuum. The kernel $\mathcal{U}_{\Phi\Theta}$ mediates matter–geometry exchange across the would-be horizon, which is a dynamical phase boundary in the unified geometry, not an absolute information barrier.

Encoding–decoding picture. During **collapse**, the local kymon microstate is coherently encoded into nonlocal correlations of the PQS core via $\hat{H}_{AE}$; during **evaporation**, the same coupling decodes and releases those correlations as entangled Apeiron/Kymon quanta. This is the precise implementation, in our ontology, of unitary recoding emphasized in scrambling and island-based accounts [336, 335, 139, 232].

Step 3: State-Dependent Scattering and the Out State

Let $U(t_2, t_1)$ be the time-ordered evolution generated by the full Hamiltonian on $\bar{\Sigma}$. Define Møller operators on the $\bar{\Sigma}$ background,

$$\Omega_{\pm}[\bar{\Sigma}, \omega] \;:=\; \lim_{t \to \pm\infty} U(0, t)\, U_0(t, 0),$$

where U_0 is free evolution on $\bar{\Sigma}$ and ω denotes the global algebraic state (the Quantum Holo-State) fixing correlation functions. The *state-dependent S-matrix* is then

$$\mathbf{S}[\bar{\Sigma}, \omega] \;:=\; \Omega_{-}^{\dagger}[\bar{\Sigma}, \omega]\;\Omega_{+}[\bar{\Sigma}, \omega]. \tag{16.171}$$

Asymptotic completeness (no remnants; complete decay of the PQS core) implies that the range of $\Omega_{\pm}$ equals the full in/out Fock spaces of Apeiron/Kymon quanta on $\bar{\Sigma}$; in that case $\mathbf{S}$ is unitary. The corrected mode structure entering $\Omega_{\pm}$ depends on $A, B, \ldots$ through the field equations on $\bar{\Sigma}$ (cf. $\bar{R}_{tt}$ dominated by A, $\bar{R}_{rr}$ by B), and spectral features can be state-conditioned (effective gaps) by the Holo-State.

Single-quantum emission amplitude. For an emitted exterior mode $u_k^{(\bar{\Sigma})}$ (solution on $\bar{\Sigma}$) and PQS transition $|\Psi_{\mathrm{PQS},i}\rangle \to |\Psi_{\mathrm{PQS},f}\rangle$,

$$\begin{aligned}\mathcal{A}_{k;i\to f} &= \langle \Psi_{\mathrm{PQS},f}; k_{\mathrm{out}}|\ \mathbf{S}[\bar{\Sigma},\omega]\ |\Psi_{\mathrm{PQS},i}\rangle \\ &\simeq i\int d^4x\,\sqrt{|\bar{\Sigma}|}\int d^4x'\,\sqrt{|\bar{\Sigma}'|}\,\left(u_k^{(\bar{\Sigma})}(x)\right)^*\mathcal{U}_{\Phi\Theta}(x,x')\left\langle \Psi_{\mathrm{PQS},i}\middle|\hat{\boldsymbol{\Theta}}(x')\,\hat{\boldsymbol{\Phi}}(x)\middle|\Psi_{\mathrm{PQS},f}\right\rangle_{\omega},\end{aligned} \tag{16.172}$$

to leading order in $\hat{H}_{AE}$. The matrix element is taken in the global state ω (fixing higher-point correlators), so $\mathcal{A}_{k;i\to f}$ depends *explicitly* on relational correlations of the PQS microstate and on the corrected geometry through the mode functions and measure. Consequently, while coarse-grained local spectra appear thermal (Unruh/Hawking), the *fine-grained* emission is not thermal; it carries microstate-specific phase information stored in the PQS.

Theorem 16.1 (Unitary Transfer of Information)**.** *Assume: (i) the full Hamiltonian on* $(\mathcal{M},\bar{\Sigma})$ *with interaction density* (16.169) *is self-adjoint on the GNS Hilbert space of* ω *(Quantum Holo-State); (ii) evaporation is complete and asymptotic completeness holds for the Apeiron/Kymon out space; (iii) the exterior algebra satisfies the microlocal spectrum/microcausality conditions appropriate to* $\bar{\Sigma}$*. Then the map*

$$\mathbf{U}_{\mathrm{BH}} \;:=\; \mathbf{S}[\bar{\Sigma},\omega]\Big|_{\mathcal{H}_{\text{in-star}}} \;:\; \mathcal{H}_{\text{in-star}} \;\xrightarrow{\cong}\; \mathcal{H}_{\text{out-rad}}$$

is a unitary isomorphism, and $|\Psi_{\mathrm{final}}\rangle = \mathbf{U}_{\mathrm{BH}}\,|\Psi_{\mathrm{initial}}\rangle$ *is pure. Hence no information is lost in collapse+evaporation.*

Proof sketch. By (i)–(iii) the limits defining $\Omega_{\pm}[\bar{\Sigma},\omega]$ exist and are isometries with ranges the full in/out spaces; thus $\mathbf{S} = \Omega_-^\dagger\Omega_+$ is unitary. Choose in data $|\Psi_{\mathrm{initial}}\rangle$ supported on the star's Hilbert subspace at early times. Asymptotic completeness ensures that late-time evolution lands in the out radiation space, $|\Psi_{\mathrm{final}}\rangle = \Omega_-^\dagger\Omega_+\,|\Psi_{\mathrm{initial}}\rangle$. Purity follows from unitarity. The apparent thermality for local exterior observables is a coarse-grained (KMS-like) property of reduced correlators, not a property of the global out state; this reconciles Hawking's spectrum with unitarity in line with scrambling and island results. Algebraically, the consistency of subsystem states is guaranteed by the coherence of the global state ω on the observable net (cf. the use of C*-algebraic states and their consistent restrictions). [336, 335, 139] □

Operational corollaries. (i) *Amplitude non-thermality.* The dependence of $\mathcal{A}_{k;i\to f}$ on ω and $\mathcal{U}_{\Phi\Theta}$ (eq. (16.172)) makes fine-grained emissions microstate-specific, even though coarse-grained spectra are thermal to local observers (Section 16.1.1).
(ii) *Background dependence.* The corrected components $A(r,\theta)$ and $B(r,\theta)$ enter both the mode functions and phase space measure through $\bar{R}_{tt}$ and $\bar{R}_{rr}$, respectively; detector-level observables thus encode the inner relational corrections.
(iii) *State-conditioned spectra.* Effective IR/UV structure (e.g., gaps/band-edges) arises state-dependently through the finite entanglement capacity of the Holo-State, shaping late-time emission channels.

Cross-checks with the thermodynamic narrative. The theorem is consistent with the "dual nature" account: locally KMS-like, globally pure. It also dovetails with the chapter's information-conservation bookkeeping (exchange between radiative entropy and temporal entropy), now implemented explicitly through the unitary $\mathbf{S}[\bar{\Sigma}, \omega]$.

Summary. The evaporation map $\mathbf{U}_{\rm BH}$ is constructed as a state-dependent S-matrix on the corrected background $\bar{\Sigma}$. The emission amplitude $\mathcal{A}$ inherits detailed dependence on PQS correlations via the interaction kernel and correlators, ensuring that the full late-time radiation is a pure, microstate-encoding out state—while reproducing apparent thermality for restricted, local measurements [336, 335, 139, 232].

The Emission Amplitude

Our objective is to compute the quantum–mechanical *single–emission* transition amplitude for the process $|\Psi_{\rm PQS,i}\rangle \to |\Psi_{\rm PQS,f}; \psi_{\rm emitted}\rangle$. At leading order in time–dependent perturbation theory on the unified background $\bar{\Sigma}$, the instantaneous *matrix element* of the interaction Hamiltonian $\hat{H}_{AE}(t)$ (eq. (16.170)) is

$$\mathcal{M}_{fi}(t) \;:=\; \langle \Psi_{\rm PQS,f}; \psi_{\rm emitted}|\; \hat{H}_{AE}(t)\; |\Psi_{\rm PQS,i}\rangle\,, \tag{16.173}$$

and the first–order transition amplitude is the time integral

$$\mathcal{A}_{fi} \;=\; -\,i \int_{-\infty}^{+\infty} dt\; \mathcal{M}_{fi}(t)\,, \qquad \text{with}\;\; \hat{H}_{AE}(t) \text{ given by eq. (16.170).} \tag{16.174}$$

(Observable transition *rates* follow from Fermi's Golden Rule after squaring $|\mathcal{A}_{fi}|$ and including the appropriate density of final states; we defer that to later steps.)

1. The States: Wavefunctions on the Corrected Spacetime

We now make precise the kinematical data entering eq. (16.174).

- **PQS core state $|\Psi_{\rm PQS,i}\rangle$:**

  Let $\mathcal{H}_{\rm PQS}$ be the Hilbert space of the core (Tantrakyma + Gaiamiti sectors) obtained from the GNS representation of the global algebraic state ω. Choosing any orthonormal basis $\{|n\rangle\} \subset \mathcal{H}_{\rm PQS}$ adapted to the conserved charges used in the asymptotic matching,

  $$|\Psi_{\rm PQS,i}\rangle \;=\; \sum_n c_n\; |n\rangle\,, \qquad \sum_n |c_n|^2 = 1, \tag{16.175}$$

  where the complex amplitudes c_n encode the microstate data inherited from the progenitor star. The post–emission core state $|\Psi_{\rm PQS,f}\rangle$ lies in the energy–reduced sector fixed by conservation, $E(\Psi_{\rm PQS,i}) = E(\Psi_{\rm PQS,f}) + \hbar\omega_k$ (modulo the small back–reaction stored in $\bar{t}_{\mu\nu}$).

- **Emitted particle $|\psi_{\rm emitted}\rangle$.** For definiteness take a scalar (kymon) emission. Its one–particle state is created by an "out" creation operator on $\bar{\Sigma}$:

  $$|\psi_{\rm emitted}\rangle \;=\; \hat{a}_k^\dagger\; |0_{\rm out}^{(\bar{\Sigma})}\rangle\,, \qquad \hat{\Phi}(x) \;=\; \sum_k \Big(u_k^{(\bar{\Sigma})}(x)\, \hat{a}_k + u_k^{(\bar{\Sigma})}(x)^*\, \hat{a}_k^\dagger \Big)\,, \tag{16.176}$$

with mode functions $u_k^{(\bar\Sigma)}$ solving the Klein–Gordon equation on the full corrected geometry:

$$\left(\bar\Sigma^{\mu\nu}\nabla_\mu^{(\mathrm{rel})}\nabla_\nu^{(\mathrm{rel})} - m^2\right) u_k^{(\bar\Sigma)}(x) = 0, \qquad \nabla^{(\mathrm{rel})} \text{ the Levi–Civita connection of } \bar\Sigma_{\mu\nu}. \tag{16.177}$$

Here $\bar\Sigma_{\mu\nu} = \bar g_{\mu\nu}^{\mathrm{Kerr}} + \bar t_{\mu\nu}$ encodes the temporal–tensor corrections. In the axisymmetric ansatz, the leading corrections are parameterized by the functions $A(r,\theta)$ and $B(r,\theta)$ entering $\bar R_{tt}$ and $\bar R_{rr}$, respectively, and thus deform both the effective potential and the redshift structure felt by $u_k^{(\bar\Sigma)}$ (cf. the discussion around $\bar R_{tt}, \bar R_{rr}$).

Normalization and inner product. Let Σ_t be a Cauchy slice with unit normal n^μ and induced metric $\bar\gamma_{ij}$. We use the standard Klein–Gordon product built from $\nabla^{(\mathrm{rel})}$:

$$(\phi,\psi)_{\mathrm{KG}}^{(\bar\Sigma)} := i\int_{\Sigma_t} d^3x\,\sqrt{|\bar\gamma|}\; n_\mu\,\phi^*\overleftrightarrow{\nabla}^{(\mathrm{rel})\mu}\psi, \qquad f\overleftrightarrow{\nabla}^{\mu} g := f(\nabla^\mu g) - (\nabla^\mu f)g. \tag{16.178}$$

Positive–frequency "out" modes are chosen with respect to the horizon–generating Killing field χ^a of the stationary background (restricted by the unified tensor $\bar\Sigma$), and normalized as

$$\left(u_k^{(\bar\Sigma)}, u_{k'}^{(\bar\Sigma)}\right)_{\mathrm{KG}}^{(\bar\Sigma)} = \delta_{kk'}, \quad \left(u_k^{(\bar\Sigma)*}, u_{k'}^{(\bar\Sigma)}\right)_{\mathrm{KG}}^{(\bar\Sigma)} = 0, \quad \left(u_k^{(\bar\Sigma)*}, u_{k'}^{(\bar\Sigma)*}\right)_{\mathrm{KG}}^{(\bar\Sigma)} = -\delta_{kk'}. \tag{16.179}$$

With this choice, $[\hat a_k, \hat a_{k'}^\dagger] = \delta_{kk'}$ and $|0_{\mathrm{out}}^{(\bar\Sigma)}\rangle$ is annihilated by all $\hat a_k$.

2. The Interaction Hamiltonian ($\hat H_{AE}$)

This is the operative mechanism behind information transfer. The coupling is fundamentally bi-local, as encoded by the Bi-Local Interaction Lagrangian Density (Definition 10.222). On a stationary foliation adapted to the unified background $\bar\Sigma$, the interaction Hamiltonian on a Cauchy slice Σ_t is

$$\hat H_{AE}(t) = \int_{\Sigma_t}\sqrt{|\bar\gamma|}\,d^3x \int_{\Sigma_t}\sqrt{|\bar\gamma'|}\,d^3x'\;:\hat{\mathbf\Phi}^\dagger(x)\,\mathcal{U}_{\Phi\Theta}(x,x')\,\hat{\mathbf\Theta}(x'):, \tag{16.180}$$

which is the slice version of eq. (16.170). Here $\bar\gamma$ is the induced 3-metric determinant on Σ_t, and the colons denote normal ordering relative to the $\bar\Sigma$ vacuum.

Kernel properties. Self-adjointness of $\hat H_{AE}(t)$ follows from the kernel symmetry

$$\left(\mathcal{U}_{\Phi\Theta}(x,x')\right)^\dagger = \mathcal{U}_{\Phi\Theta}(x',x), \qquad x,x'\in\Sigma_t, \tag{16.181}$$

together with the standard adjoint action on the field multiplets $\hat{\mathbf\Phi}, \hat{\mathbf\Theta}$ and the trace/contraction over internal indices. Microcausality is preserved by requiring $[\hat{\mathbf\Phi}(x), \hat{\mathbf\Theta}(x')] = 0$ for spacelike separation with respect to $\bar\Sigma$, and by choosing $\mathcal{U}_{\Phi\Theta}$ with support localized within a relational correlation length set by the near-horizon PQS structure (so that no superluminal signaling is introduced).

This nonlocal operator acts as a "quantum handshake," linking a geometric field point x with a material field point x' through $\mathcal{U}_{\Phi\Theta}(x,x')$, thereby enabling bi-directional encoding/decoding between the PQS core and exterior modes.

3. The Calculation of the Amplitude

We compute the first-order amplitude to emit one exterior quantum in an out-mode $u_k^{(\bar{\Sigma})}$ while the PQS core transitions from $|\Psi_{\mathrm{PQS},i}\rangle = \sum_n c_n\,|n\rangle$ to some definite $|m\rangle$ in the energy-reduced sector. From eqs. (16.174) and (16.180) we obtain

$$\mathcal{A}_{k,m} = \; -i\int_{-\infty}^{+\infty} dt \int_{\Sigma_t} \sqrt{|\bar{\gamma}|}\, d^3x \int_{\Sigma_t} \sqrt{|\bar{\gamma}'|}\, d^3x' \, \left(u_k^{(\bar{\Sigma})}(x)\right)^* \mathcal{U}_{\Phi\Theta}(x,x') \, \sum_n c_n \, \langle m|\hat{\boldsymbol{\Theta}}(x')\,\hat{\boldsymbol{\Phi}}(x)|n\rangle \,. \tag{16.182}$$

It is convenient to define the PQS two-point transition kernel

$$\mathcal{C}_{mn}(x,x') \;:=\; \langle m|\hat{\boldsymbol{\Theta}}(x')\,\hat{\boldsymbol{\Phi}}(x)|n\rangle\,, \qquad x,x' \in \Sigma_t, \tag{16.183}$$

so that $\mathcal{A}_{k,m} = \sum_n c_n\,\mathcal{M}^{(k)}_{mn}$, with

$$\mathcal{M}^{(k)}_{mn} = \; -i\int_{-\infty}^{+\infty} dt \int_{\Sigma_t} \sqrt{|\bar{\gamma}|}\, d^3x \int_{\Sigma_t} \sqrt{|\bar{\gamma}'|}\, d^3x' \, \left(u_k^{(\bar{\Sigma})}(x)\right)^* \mathcal{U}_{\Phi\Theta}(x,x')\,\mathcal{C}_{mn}(x,x'). \tag{16.184}$$

If one projects onto a *superposed* final core state $|\Psi_{\mathrm{PQS},f}\rangle = \sum_m d_m\,|m\rangle$, then the amplitude is bilinear in the coefficients:

$$\mathcal{A}_k[\Psi_f,\Psi_i] = \; \sum_{m,n} d_m^*\, c_n\, \mathcal{M}^{(k)}_{mn}. \tag{16.185}$$

If the core's final state is not resolved experimentally, the inclusive probability for emission into mode k is

$$\mathcal{P}_k^{\mathrm{incl}} = \; \sum_m \left|\mathcal{A}_{k,m}\right|^2 = \; \sum_m \left|\sum_n c_n\,\mathcal{M}^{(k)}_{mn}\right|^2, \tag{16.186}$$

which is explicitly sensitive to the initial microstate amplitudes $\{c_n\}$ and to the relational correlators $\mathcal{C}_{mn}$.

Proof of Information Transfer

Proposition 16.2 (Microstate imprint in the emitted radiation)**.** *Assume: (i) the kernel symmetry* (16.181) *and microcausal support of* $\mathcal{U}_{\Phi\Theta}$*; (ii) well-posed out-modes* $u_k^{(\bar{\Sigma})}$ *normalized by the* $\bar{\Sigma}$ *Klein–Gordon product; (iii) asymptotic completeness for the one-quantum sector. Then the map* $\{c_n\} \mapsto \{\mathcal{A}_{k,m}\}_{k,m}$ *given by* (16.182) *is linear and, for a generic kernel* $\mathcal{U}_{\Phi\Theta}$ *and nondegenerate* $\mathcal{C}_{mn}$*, injective up to an overall phase. Consequently, the emitted radiation encodes the PQS microstate and is not thermal in the fine-grained sense.*

Sketch. Linearity in $\{c_n\}$ is manifest in (16.182). For generic $\mathcal{U}_{\Phi\Theta}$ and nonpathological $\mathcal{C}_{mn}$, the family $\{\mathcal{M}^{(k)}_{mn}\}_{k,m}$ spans the space of linear functionals on the coefficient vector $(c_n)_n$, implying injectivity up to a global phase. Therefore, the inclusive probabilities (16.186) depend on coherent sums of the c_n and cannot be reproduced by a Gibbs (thermal) ensemble with state-independent weights. Combined with unitarity of the full state-dependent S-matrix on $\bar{\Sigma}$ (Theorem 16.1), the global out state is pure and information-preserving. □

Remarks. (i) If one chooses $|\Psi_{\mathrm{PQS},f}\rangle = \sum_m d_m\,|m\rangle$, then (16.185) shows explicitly how relative phases between $\{d_m\}$ and $\{c_n\}$ appear in the amplitude, enabling tomography with mode-resolved measurements. (ii) The dependence on the corrected geometry enters through both $u_k^{(\bar{\Sigma})}$ and the measure in (16.184), thereby imprinting the near-horizon relational corrections (via $A, B, \ldots$) into detector-level observables.

16.3.1 The Quantum Corrections to Black Hole Entropy

Our goal is to compute the corrected entropy $S_{\mathrm{corrected}}$ from the corrected event-horizon area A_{EH} and to isolate the first-order quantum correction. Throughout we set $k_B = 1$ and keep c explicit.[2]

1. Foundational Formulas

- **Bekenstein–Hawking entropy** [86, 87]:

$$S_{\mathrm{BH}} \;=\; \frac{A_{\mathrm{BH}}}{4\,G\hbar/c^3} \;=\; \frac{4\pi R_S^2}{4\,G\hbar/c^3}, \qquad R_S = \frac{2GM}{c^2}. \tag{16.187}$$

- **Corrected event-horizon radius.** In the relational framework the horizon is a phase boundary whose location solves a coherence-corrected relation (cf. eq. (16.72)):

$$R_{\mathrm{EH}} \;=\; R_S \;+\; \delta R_H, \qquad \delta R_H \;=\; f_1(R_S) \;-\; R_S\,a_1(R_S), \tag{16.188}$$

  to first order in the temporal-tensor corrections $\{a_1, f_1\}$ (notation as in the spherically symmetric ansatz). Equivalently, if one prefers the shorthand of a small positive length ϵ_f capturing the repulsive dressed vacuum contribution, set $\delta R_H = -\,\epsilon_f$.

2. Corrected Area A_{EH}

With spherical symmetry,

$$A_{\mathrm{EH}} \;=\; 4\pi R_{\mathrm{EH}}^2 \;=\; 4\pi\Big(R_S + \delta R_H\Big)^2 \;=\; A_{\mathrm{BH}} \;+\; 8\pi R_S\,\delta R_H \;+\; \mathcal{O}(\delta R_H^2). \tag{16.189}$$

Retaining only first order,

$$A_{\mathrm{EH}} \;\approx\; \underbrace{4\pi R_S^2}_{A_{\mathrm{BH}}} \;+\; \underbrace{8\pi R_S\,\delta R_H}_{\delta A}. \tag{16.190}$$

If $\delta R_H = -\epsilon_f$ with $\epsilon_f > 0$, then $\delta A = -8\pi R_S \epsilon_f$.

3. Corrected Entropy $S_{\mathrm{corrected}}$

Insert (16.190) into (16.187):

$$S_{\mathrm{corrected}} \;=\; \frac{A_{\mathrm{EH}}}{4\,G\hbar/c^3} \;\approx\; S_{\mathrm{BH}} \;+\; \frac{\delta A}{4\,G\hbar/c^3} \;=\; S_{\mathrm{BH}} \;+\; \frac{2\pi c^3 R_S}{G\hbar}\,\delta R_H. \tag{16.191}$$

[2] In geometric units $G = \hbar = c = 1$ all c-factors below drop out.

Equivalently, in the ϵ_f-parametrization,

$$S_{\text{corrected}} \;\approx\; S_{\text{BH}} \;-\; \frac{2\pi c^3 R_S}{G\hbar}\,\epsilon_f. \tag{16.192}$$

Physical Interpretation. The linear correction $\delta S = \frac{2\pi c^3 R_S}{G\hbar}\,\delta R_H$ expresses how near-horizon relational structure—encoded by the temporal tensor via a_1, f_1 in (16.188)—modifies the area term. For $\delta R_H < 0$ (effective repulsion from dressed vacuum energy), $\delta S < 0$: part of the informational content that would classically be counted by the area is reallocated to the exterior relational degrees of freedom in the quantum vacuum. Thus entropy is not purely geometric but *relational*: it emerges from the joint configuration of the horizon and its chronogeometric environment. This dovetails with the unitary evaporation narrative developed earlier (cf. Section 16.3) and provides a concrete, testable first-order prediction of the framework.

Remark (Kerr extension, for reference). For rotation with parameter a, the classical area is $A^{(0)} = 4\pi(r_+^2 + a^2)$ with $r_+ \approx R_S - \frac{a^2}{2R_S}$. Including the relational shift $r_+ \mapsto r_+ + \delta R_H$ gives

$$\delta A \;\approx\; 8\pi r_+\,\delta R_H \;-\; 4\pi\,\frac{a^2}{R_S}\,,$$

so that $\delta S = (\delta A)/(4G\hbar/c^3)$ splits into a rotational (classical) piece and a relational piece proportional to δR_H. In the non-rotating limit $a \to 0$ this reduces to (16.191).

Lemma 16.3 (First-law validation of the entropy correction)**.** *Let κ_{eff} denote the corrected surface gravity (cf. eq. (16.4)) associated with the unified background $\bar{\Sigma}_{\mu\nu}$, and*

$$T_{\text{eff}} \;:=\; \frac{\hbar\,\kappa_{\text{eff}}}{2\pi c} \tag{16.193}$$

the corresponding Hawking temperature. Consider quasi-stationary variations among nearby solutions with fixed J, Q (and any other work charges), so that the relational First Law reads

$$\delta M \;=\; \frac{\kappa_{\text{eff}}}{8\pi G}\,\delta A\,. \tag{16.194}$$

Then the unique entropy functional $S(A)$ consistent with (16.193)–(16.194) to first order is

$$\frac{dS}{dA} \;=\; \frac{c^3}{4\,G\hbar} \qquad\Longrightarrow\qquad S(A) \;=\; \frac{c^3}{4\,G\hbar}\,A \;+\; \text{const}, \tag{16.195}$$

so that, for a small horizon shift $R_{\text{EH}} = R_S + \delta R_H$,

$$\delta S \;=\; \frac{c^3}{4\,G\hbar}\,\delta A \;=\; \frac{c^3}{4\,G\hbar}\,(8\pi R_S\,\delta R_H) \;=\; \frac{2\pi c^3 R_S}{G\hbar}\,\delta R_H\,, \tag{16.196}$$

which matches eqs. (16.191) and (16.192) upon setting $\delta R_H = -\epsilon_f$.

Proof. Thermodynamics gives $\delta M = T_{\text{eff}}\,\delta S$ at fixed J, Q. Using (16.193) and (16.194) we have

$$\frac{\hbar\,\kappa_{\text{eff}}}{2\pi c}\,\delta S \;=\; \frac{\kappa_{\text{eff}}}{8\pi G}\,\delta A\,.$$

For nonextremal configurations ($\kappa_{\text{eff}} \neq 0$) the κ_{eff} factors cancel, yielding $dS/dA = c^3/(4G\hbar)$. Integrating gives (16.195) with the integration constant fixed to zero by the classical limit and the standard normalization. Substituting $\delta A = 8\pi R_S\,\delta R_H$ reproduces (16.196). The extremal limit $\kappa_{\text{eff}} \to 0$ is continuous by taking limits of the nonextremal family. □

Corollary (independence from the microphysics of κ_{eff}). Equation (16.195) shows that, to first order, the entropy correction depends only on the area change and *not* on the detailed functional form of κ_{eff}. Hence any relational mechanism that produces the same δA (e.g., via a_1, f_1 at the horizon through $\delta R_H = f_1(R_S) - R_S a_1(R_S)$) yields the same δS, in agreement with Section 16.3.1.

The Black Hole as a Unitary Process of Relational Transformation

Black holes expose the tension between smooth, deterministic geometry and unitary quantum dynamics. In the relational framework developed here, this tension is resolved by elevating the kinematics from the geometric background $\bar{g}_{\mu\nu}$ to the unified chronogeometric tensor $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$, wherein the event horizon is a bona-fide phase boundary fixed by a coherence-corrected condition (cf. eq. (16.72)). Collapse does not terminate in a pathology; rather, it enters a stationary coherence phase (the PQS core) that stores information in nonlocal relational correlations (cf. the stationary/coherence hypotheses for Kerr-type attractors).

Unitary dynamics. The evolution is governed by the *Total Relational Action* and its bi-local interaction term (Definitions 9.357 and 10.222), giving rise to the interaction Hamiltonian $\hat{H}_{AE}$ on $\bar{\Sigma}$ (eqs. (16.170) and (16.180)). This coupling implements an *encoding/decoding* map between the PQS core and exterior modes: during collapse, local matter information is encoded into the PQS correlators; during evaporation, the same coupling decodes it into outgoing Apeiron/Kymon quanta. Formally, the state-dependent S-matrix $\mathbf{S}[\bar{\Sigma}, \omega]$ is unitary (Theorem 16.1), and the single-quantum amplitudes depend linearly on the PQS microstate coefficients through the transition kernel (eqs. (16.182) to (16.184)), yielding a generic microstate imprint (Proposition 16.2). Thus the global out-state is pure while local measurements recover KMS-like thermality (cf. Section 16.1.1 and [336, 335, 139, 232]).

Thermodynamic reconciliation. The relational horizon shift $R_{\text{EH}} = R_S + \delta R_H$ induced by the temporal tensor (e.g., $\delta R_H = f_1(R_S) - R_S a_1(R_S)$) corrects the area and hence the entropy:

$$S_{\text{corrected}} \;=\; \frac{c^3}{4G\hbar}\,A_{\text{EH}} \;\approx\; S_{\text{BH}} \;+\; \frac{2\pi c^3 R_S}{G\hbar}\,\delta R_H,$$

as derived in Section 16.3.1. The same result follows from the First Law with the corrected surface gravity κ_{eff} (Lemma 16.3), showing that the entropy shift is fixed by δA irrespective of the detailed microphysics of κ_{eff}. Negative δR_H (effective repulsion due to dressed vacuum energy) yields a negative δS, signaling that part of the information budget is carried by exterior relational degrees of freedom rather than being counted solely by the geometric area.

Synthesis. Taken together—unitary scattering on $\bar{\Sigma}$ (Theorem 16.1), microstate-sensitive emission amplitudes (Proposition 16.2), dual local/global thermality (cf. Section 16.1.1), and the consistent thermodynamic correction to S (Lemma 16.3)—the black hole is a *unitary process of relational transformation*. It coherently converts localized, material information into nonlocal, relational information and back into asymptotic radiation, without violating unitarity, causality, or thermodynamics. The observational consequences (ringdown shifts, shadow/ISCO modifications, and late-time spectral structure) follow from the same horizon-scale parameters that govern δR_H and the emission kernel, providing a single pipeline from theory to constraint.

16.4 The Chronogeometric Kerr Black Hole

From Schwarzschild to Kerr in Classical Geometry

Throughout this subsection we set $G = c = 1$ for compactness (all factors are easily restored by dimensional analysis).[3] The Kerr metric is the unique, asymptotically flat, stationary, axisymmetric vacuum solution of the Einstein equations describing a rotating mass M with specific angular momentum $a := J/M$ [647, 648, 649]. In Boyer–Lindquist coordinates (t, r, θ, ϕ) the line element reads

$$\begin{aligned} ds^2 = &-\left(1 - \frac{2Mr}{\rho^2}\right) dt^2 - \frac{4Mar\sin^2\theta}{\rho^2}\, dt\, d\phi \\ &+ \frac{\rho^2}{\Delta}\, dr^2 + \rho^2\, d\theta^2 \\ &+ \left(r^2 + a^2 + \frac{2Ma^2 r \sin^2\theta}{\rho^2}\right) \sin^2\theta\, d\phi^2, \end{aligned} \tag{16.197}$$

where

$$\rho^2 := r^2 + a^2\cos^2\theta, \qquad \Delta := r^2 - 2Mr + a^2.$$

The horizons are located at $r_\pm = M \pm \sqrt{M^2 - a^2}$, and the ring singularity lies at $\rho = 0$ (i.e. $r = 0, \theta = \pi/2$).

Chrono–geometric uplift. To embed rotation into the emergent framework, we take the unified tensor to be the sum of a Kerr *geometric sector* and a stationary, axisymmetric *temporal sector*,

$$\bar{\Sigma}_{\mu\nu}(r,\theta) \;=\; \bar{g}^{\mathrm{Kerr}}_{\mu\nu}(M,a) \;+\; \bar{t}_{\mu\nu}(r,\theta), \qquad (\partial_t, \partial_\phi) \text{ Killing on both terms.} \tag{16.198}$$

We parameterize $\bar{t}_{\mu\nu}$ in Boyer–Lindquist charts by five scalar profiles $\{A, B, C, D, E\} =$

[3] In particular, $a := J/(Mc)$ has dimensions of length; with $c = 1$, $a = J/M$.

$\{A(r,\theta),\dots,E(r,\theta)\}$ that respect stationarity, axisymmetry, and equatorial symmetry:

$$\begin{aligned}\mathcal{T}_{\mu\nu}\,dx^\mu dx^\nu = \bar{t}_{\mu\nu}\,dx^\mu dx^\nu = \quad & \underbrace{-A(r,\theta)\,dt^2}_{tt\text{ correction }A\text{ (lapse)}} \\ & + \underbrace{B(r,\theta)\,\frac{\rho^2}{\Delta}\,dr^2}_{rr\text{ correction }B} \\ & + \underbrace{C(r,\theta)\,\rho^2\,d\theta^2}_{\theta\theta\text{ correction }C} \\ & + \underbrace{E(r,\theta)\,\sin^2\theta\,d\phi^2}_{\phi\phi\text{ correction }E} \\ & - \underbrace{2D(r,\theta)\,\sin^2\theta\,dt\,d\phi}_{t\phi\text{ correction }D\text{ (frame dragging)}}\end{aligned} \tag{16.199}$$

Thus the full line element is

$$\bar{\Sigma}_{\mu\nu}dx^\mu dx^\nu = \left(\bar{g}^{\mathrm{Kerr}}_{\mu\nu} + \bar{t}_{\mu\nu}\right)dx^\mu dx^\nu, \tag{16.200}$$

and *reduces exactly* to classical Kerr when $A = B = C = D = E = 0$. If one prefers to absorb some corrections into a "dressed" mass function, one may alternatively replace $M \mapsto M(r,\theta) = M + \delta M(r,\theta)$ in Δ and redistribute terms between $\bar{g}$ and $\bar{t}$; to first order these conventions are equivalent.

Boundary and regularity conditions. To ensure asymptotic flatness and axis regularity, we impose

$$\begin{aligned} A, B, C, D, E = \mathcal{O}(r^{-p}) \quad (p>1), \quad r\to\infty, \\ \partial_\theta A\big|_{\theta=0,\pi} = 0, \quad \partial_\theta B\big|_{\theta=0,\pi} = 0, \quad D = \mathcal{O}(\sin^2\theta). \end{aligned} \tag{16.201}$$

Equatorial symmetry requires $X(r,\pi-\theta) = X(r,\theta)$ for each profile $X \in \{A,B,C,D,E\}$. The profiles are further constrained by the unified field equations and the relational energy conditions (cf. earlier sections).

Horizon and ergoregion (unified geometry). Let $\chi^a = \partial_t + \Omega_H\,\partial_\phi$ be the (a priori unknown) horizon generator. The *ergosurface* is the locus

$$\bar{\Sigma}_{tt}(r,\theta) \;=\; \bar{g}^{\mathrm{Kerr}}_{tt}(r,\theta) \;-\; A(r,\theta) \;=\; 0, \tag{16.202}$$

while the *event horizon* is the largest root $r = R_H$ for which

$$\bar{\Sigma}_{ab}\,\chi^a\chi^b\Big|_{r=R_H} \;=\; 0, \qquad \Omega_H \;=\; -\left.\frac{\bar{\Sigma}_{t\phi}}{\bar{\Sigma}_{\phi\phi}}\right|_{r=R_H}, \tag{16.203}$$

which reduce to the Kerr conditions when $A = \cdots = E = 0$. To first order in the profiles, regularity (zeroth law) enforces that R_H and Ω_H are θ-independent; the resulting shift δR_H can be expressed in terms of horizon-value data $\{A,B,D,E\}$ and their derivatives (cf. the spherical results and the linear response map).

Small-spin and weak-correction limits. For $a/M \ll 1$ and $\|A, B, C, D, E\| \ll 1$,

$$R_H = 2M - \frac{a^2}{2M} + \delta R_H + \mathcal{O}\Big(a^4, \text{quadratic}\Big), \tag{16.204a}$$

$$\Omega_H = \frac{a}{4M^2} + \delta\Omega_H + \mathcal{O}\Big(a^3, \text{quadratic}\Big). \tag{16.204b}$$

with δR_H and $\delta\Omega_H$ determined by the unified horizon conditions (16.203). Detector-level observables (ringdown, shadow, ISCO) may be organized as linear functionals of the horizon-value data, providing a direct pipeline from $\{A, B, C, D, E\}$ to constraints.

Remarks on gauge and alternative parameterizations. The choice (16.199) is a convenient "additive" gauge that keeps the geometric sector strictly Kerr. One can equally work in a "multiplicative" gauge that rescales the Kerr coefficients by $(1 + \varepsilon_X)$ functions or in a Δ-dressing gauge with $M(r, \theta)$ and a frame-dragging profile; all are equivalent at first order if boundary/regularity conditions (16.201) are enforced.

Definition 16.4 (Temporal Tensor Ansatz for the Chronogeometric Kerr Solution)**.** Let the chronogeometric tensor be $\bar{\Sigma}_{\mu\nu} = \bar{g}^{\text{Kerr}}_{\mu\nu}(M, a) + \bar{t}_{\mu\nu}$, with $\partial_t, \partial_\phi$ Killing for both sectors. To accommodate anisotropic, state-dependent correlations induced by rotation, we take $\bar{t}_{\mu\nu} \neq 0$ and allow an off-diagonal $t\phi$-component.

Kerr coframe representation. Introduce the Boyer–Lindquist Kerr coframe

$$\hat{e}^t := c\, dt, \quad \hat{e}^r := \sqrt{\frac{\rho^2}{\Delta}}\, dr, \quad \hat{e}^\theta := \rho\, d\theta, \quad \hat{e}^\phi := \sin\theta\, d\phi,$$

with $\rho^2 = r^2 + a^2 \cos^2\theta$ and $\Delta = r^2 - 2Mr + a^2$. In this orthonormal coframe the temporal tensor is encoded by five real, dimensionless profiles $\{A, B, C, D, E\} = \{A(r, \theta), \ldots, E(r, \theta)\}$ via

$$\mathcal{T}_{ab}\, \hat{e}^a \hat{e}^b := \bar{t}_{ab}\, \hat{e}^a \hat{e}^b = \begin{bmatrix} -A & 0 & 0 & -D \\ 0 & B & 0 & 0 \\ 0 & 0 & C & 0 \\ -D & 0 & 0 & E \end{bmatrix}_{ab}, \qquad a, b \in \{t, r, \theta, \phi\}. \tag{16.205}$$

Coordinate-basis form (equivalence). In the coordinate basis this yields the line element contribution

$$\begin{aligned} \mathcal{T}_{\mu\nu}\, dx^\mu dx^\nu = &-A(r, \theta)\, dt^2 + B(r, \theta)\, \frac{\rho^2}{\Delta}\, dr^2 + C(r, \theta)\, \rho^2\, d\theta^2 \\ &+ E(r, \theta)\, \sin^2\theta\, d\phi^2 - 2D(r, \theta)\, \sin^2\theta\, dt\, d\phi, \end{aligned} \tag{16.206}$$

so that $\bar{\Sigma}_{\mu\nu} dx^\mu dx^\nu = (\bar{g}^{\text{Kerr}}_{\mu\nu} + \bar{t}_{\mu\nu}) dx^\mu dx^\nu$ reduces exactly to Kerr when $A = B = C = D = E = 0$.

Physical interpretation of the profiles.

1. $A(r,\theta)$: lapse correction (tt), encoding redshift deviations from local temporal dressing;
2. $B(r,\theta)$, $C(r,\theta)$: radial/polar coherence–viscosity $(rr,\theta\theta)$, capturing direction-dependent decoherence from coherence gradients.
3. $D(r,\theta)$: quantum–relational frame dragging $(t\phi)$, an off-diagonal coupling beyond classical Kerr reflecting hololocal temporal–angular correlations;
4. $E(r,\theta)$: azimuthal correction $(\phi\phi)$, governing backreaction on angular transport and potential topological memory.

Symmetry, boundary, and regularity conditions. Stationarity, axisymmetry, and equatorial symmetry impose

$$\partial_t X = \partial_\phi X = 0, \qquad X(r,\pi-\theta) = X(r,\theta), \qquad X \in \{A,B,C,D,E\}.$$

Asymptotic flatness and axis regularity require (cf. eq. (16.201))

$$\begin{aligned} A,B,C,D,E &= \mathcal{O}(r^{-p}) && (p>1,\ r\to\infty),\\ \partial_\theta A|_{\theta=0,\pi} &= 0, & \partial_\theta B|_{\theta=0,\pi} &= 0,\\ D &= \mathcal{O}(\sin^2\theta) && . \end{aligned} \tag{16.207}$$

Horizon regularity (zeroth law) further enforces θ-independence of the horizon values to first order, ensuring a well-defined Ω_H and κ_{eff} (see eqs. (16.2), (16.4) and (16.203)).

Sourcing. The profiles are functionals of the *rotationally dressed vacuum energy density* $\rho_{\text{vac}}(r,\theta)$ and the global Holo-State through the bi-local interaction kernel; at first order they solve the unified field equations with sources determined by the finite entanglement capacity and spectral gapping mechanisms specified earlier. Their horizon values feed directly into observable shifts $\delta R_H, \delta\Omega_H, \kappa_{\text{eff}}$ and hence into ringdown/shadow constraints.

Given the temporal tensor $\bar{t}_{\mu\nu}$, the contribution to the line element is

$$\bar{t}_{\mu\nu}\, dx^\mu dx^\nu. \tag{16.208}$$

Using the coordinate–basis matrix implied by eq. (16.199), only the $(tt),(rr),(\theta\theta),(\phi\phi)$ and $(t\phi)$ components are nonzero, hence

$$\begin{aligned} \bar{t}_{\mu\nu}\, dx^\mu dx^\nu = &-A(r,\theta)\, dt^2 + B(r,\theta)\,\frac{\rho^2}{\Delta}\, dr^2 + C(r,\theta)\,\rho^2\, d\theta^2 \\ &+ E(r,\theta)\,\sin^2\theta\, d\phi^2 - 2D(r,\theta)\,\sin^2\theta\, dt\, d\phi, \end{aligned} \tag{16.209}$$

which is exactly eq. (16.199) written as a single line element. Symmetry $\bar{t}_{\mu\nu} = \bar{t}_{\nu\mu}$ is manifest, and the $\sin^2\theta$ and ρ^2/Δ factors arise from the Boyer–Lindquist chart.

Units (re-inserting c if desired). If explicit c-factors are preferred, then replace $dt \mapsto c\,dt$ in the tt and $t\phi$ terms:

$$-A\,dt^2 \;\to\; -A\,c^2dt^2, \qquad -2D\,\sin^2\theta\,dt\,d\phi \;\to\; -2D\,c\,\sin^2\theta\,dt\,d\phi,$$

leaving the spatial pieces unchanged.

These components will be sourced by a **rotationally dressed vacuum energy density** $\rho_{\rm vac}(r,\theta)$, itself a state-dependent observable determined by:

Definition 16.5 (Rotationally Dressed Vacuum Energy Density)**.** Let $\bar{\Sigma}_{\mu\nu} = \bar{g}^{\rm Kerr}_{\mu\nu}(M,a) + \bar{t}_{\mu\nu}$ be the unified tensor of the rotating solution (Definition 16.4). The *rotationally dressed vacuum energy density* is the state–dependent observable that sources the profiles $\{A,B,C,D,E\}$ in the temporal sector, defined as the expectation value of an effective local energy operator in the Quantum Holo-State:

$$\boxed{\rho_{\rm vac}(x) \;:=\; \Big\langle \overset{\circ}{\mathbf{\Psi}}\Big|\, \hat{\rho}^{(\rm eff)}(x)\,\Big|\overset{\circ}{\mathbf{\Psi}}\Big\rangle,} \tag{16.210}$$

where the expectation is taken in the emergent/GNS layer of the Holo-State (Born form $\langle\Omega_\omega|\pi_\omega(\cdot)|\Omega_\omega\rangle$; cf. Definition 1.34). The operator is decomposed into a static and a rotational contribution,

$$\hat{\rho}^{(\rm eff)}(x) \;=\; \hat{\rho}^{(\rm static)}(x) \;+\; \hat{\rho}^{(\rm rot)}(x),$$

with $\hat{\rho}^{(\rm rot)}(x)$ a functional of (i) the emergent angular momentum tensor $\hat{J}^{\mu\nu}$ and (ii) the chronogeometric vorticity $\hat{\omega}_{\mu\nu}[\bar{\Sigma}]$, the latter built from the relational Levi-Civita connection introduced for the Kerr program (Definition 6.7; see also the Christoffel decomposition $\Gamma[\bar{\Sigma}] = \Gamma[\bar{g}] + \Delta\Gamma[\bar{t}]$).

Operational sourcing of the temporal profiles. In the unified field equations the near-horizon, state-dependent vacuum stresses source the temporal functions—first emphasized for $\{A,B\}$ and extended here to the rotating set $\{A,B,C,D,E\}$. A convenient phenomenology views $\rho_{\rm vac}$ as a spectrally regulated integral modulated by a coherence-stress factor $\mathcal{C}(\xi)$ that depends on the local regime tracker (and grows toward the horizon); this feeds directly into the sourced equations for the temporal tensor components.

Physical mechanisms and correspondence to $\{A,B,C,D,E\}$.

1. *Spin-dependent spectral gapping.* Rotation modifies the vacuum fluctuation spectrum via holomorphic deformations of the modal weights $w(\lambda) \mapsto w(\lambda)\,\mathcal{C}(\xi)$ (endogenous regularization and spectral smoothness), stiffening the local energetic response. *Corresponds primarily to* diagonal profiles $\mathbf{A},\mathbf{B},\mathbf{C}$, which correct redshift and spatial intervals in the lapse/radial/polar directions (eq. (16.199)).
2. *Frame-dragging–induced asymmetry.* Rotation induces anisotropic, off-diagonal correlations (matter–geometry "handshake" via the bi-local kernel), captured by the $t\phi$ channel. *Corresponds to* $\mathbf{D}(r,\theta)$ (quantum-relational frame dragging) with a backreaction on $\mathbf{E}(r,\theta)$ in the azimuthal sector, exactly as encoded in the temporal line element (eq. (16.199)).

3. *Holosymmetric phase locking (global constraint).* Beyond local sourcing, the rotating solution is selected by a global coherence condition of holosymmetric origin: histories are organized by the modular partition and compared in controlled superposition, fixing consistent relationships among $\{A, B, C, D, E\}$ as a stability criterion (holosymmetric selection/operational bridge). This operates as a *boundary/consistency condition* on the coupled PDEs rather than an additional local source term.

Summary. The rotationally dressed vacuum energy density $\rho_{\rm vac}(r,\theta)$ is the emergent expectation of an effective energy operator in the Holo-State that (i) inherits spectral regularity from the axioms of the vacuum sector, (ii) couples nonlocally through the bi-local interaction to produce anisotropic stresses, and (iii) is globally constrained by holosymmetric phase locking. These ingredients together determine the horizon-value data for $\{A, B, C, D, E\}$ that control Ω_H, $\kappa_{\rm eff}$, and observable Kerr phenomenology in the unified background (eqs. (16.2), (16.4) and (16.203)).

The next step is to *derive* the functional form of the temporal profiles $\{A, B, C, D, E\}$ from a local, covariant expansion of the rotationally dressed vacuum energy $\rho_{\rm vac}(r,\theta)$ (Definition 16.5), and ultimately to express the temporal sector $\mathcal{T}_{\mu\nu}$ in terms of operator-valued correlators on the PQS/Holo-State bundle (via point-splitting/Hadamard renormalization) [245]. Concretely, one employs a stationary, axisymmetric invariant basis built from the unified geometry $\bar{\Sigma}$ (Killing field χ^a, vorticity $\omega_{\mu\nu}[\bar{\Sigma}]$, curvature scalars/tensors, and their derivatives), so that

$$\rho_{\rm vac}(x) = \rho_0 + \chi_1\,\omega_{\mu\nu}\omega^{\mu\nu} + \chi_2\,\bar{R} + \chi_3\,\bar{R}_{\mu\nu}\chi^\mu\chi^\nu + \chi_4\,\nabla_\alpha\omega_{\mu\nu}\nabla^\alpha\omega^{\mu\nu} + \cdots,$$

with state-dependent coefficients $\{\chi_i\}$ fixed by the Quantum Holo-State and the bi-local interaction kernel (cf. Definitions 10.222 and 16.5). This feeds the sourced equations for $\{A, \ldots, E\}$ appearing in the temporal line element (eq. (16.199)).

Definition 16.6 (Unified Vacuum Field Equation)**.** The emergent chronogeometric vacuum is determined by a self-consistency condition relating curvature of the unified tensor to the effective stress content of the dressed vacuum:

$$\boxed{\bar{R}_{\mu\nu}[\bar{\Sigma}] \;=\; \mathcal{R}^{(\rm vac)}_{\mu\nu}\Big[\rho_{\rm vac}(r,\theta)\Big],} \tag{16.211}$$

with the following components.

- **Unified Ricci tensor $\bar{R}_{\mu\nu}[\bar{\Sigma}]$.** It is computed from the *Relational Levi–Civita connection* of the full tensor $\bar{\Sigma}_{\mu\nu} = \bar{g}^{\rm Kerr}_{\mu\nu} + \bar{t}_{\mu\nu}$ (cf. Definition 6.7 and eq. (16.198)). Decomposing around classical Kerr (vacuum) gives

$$\bar{R}_{\mu\nu}[\bar{\Sigma}] = R_{\mu\nu}\Big[\bar{g}^{\rm Kerr}\Big] + \delta R_{\mu\nu}[\bar{t}; \bar{g}^{\rm Kerr}] = \delta R_{\mu\nu}[\bar{t}; \bar{g}^{\rm Kerr}], \tag{16.212}$$

since $R_{\mu\nu}[\bar{g}^{\rm Kerr}] = 0$ [647, 648, 649]. At first order, $\delta R_{\mu\nu}$ is linear in the profiles $\{A, B, C, D, E\}$ and their r, θ derivatives (see also Section 16.5).

- **Vacuum source tensor $\mathcal{R}^{(\mathrm{vac})}_{\mu\nu}$.** This is the *effective* (renormalized) vacuum stress built from local, covariant invariants of $\bar{\Sigma}$ and the stationary structure, and functionally dependent on $\rho_{\mathrm{vac}}(r,\theta)$ (Definition 16.5). Schemata consistent with symmetry and dimension include

$$\mathcal{R}^{(\mathrm{vac})}_{\mu\nu} = \alpha_0\,\bar{\Sigma}_{\mu\nu} + \alpha_1\,\omega_{\mu\lambda}\,\omega_{\nu}{}^{\lambda} + \alpha_2\,\chi_\mu\chi_\nu + \alpha_3\,\bar{R}_{\mu\nu} + \alpha_4\,\bar{\nabla}_\mu\bar{\nabla}_\nu\Big(\omega_{\alpha\beta}\omega^{\alpha\beta}\Big) + \cdots\,, \tag{16.213}$$

  with state/renormalization–dependent coefficients $\{\alpha_i\}$ fixed by matching to the Holo-State expectation (Hadamard point-splitting, operator product expansion) [245]. Intrinsic dressing (Gaiamiti/Tantrakyma), curvature-induced polarization, and holosymmetric global constraints appear as, respectively, the α_0 baseline, the curvature/vorticity terms, and coefficient relations consistent with phase-locking (Definition 16.5).

Equation (16.211) yields a coupled, nonlinear system (elliptic on stationary slices) for $\{A,B,C,D,E\}$ on (r,θ), subject to asymptotic and axis regularity (eq. (16.201)) and horizon regularity (eq. (16.203)). Its solution defines the Chronogeometric Kerr configuration and determines the observable horizon data $\{\delta R_H, \delta\Omega_H, \kappa_{\mathrm{eff}}\}$ (eqs. (16.2), (16.4) and (16.204)), thereby linking the Holo-State microphysics to ringdown/shadow phenomenology.

Operator representation (PQS/Holo-State). Writing the effective energy density as a renormalized quadratic form in fundamental fields $\widehat{\mathbf{\Phi}}, \widehat{\mathbf{\Theta}}$ with the bi-local kernel (Definition 10.222), one may express $\rho_{\mathrm{vac}}(x) = \langle\Omega_\omega|\hat{\rho}^{(\mathrm{eff})}(x)|\Omega_\omega\rangle$ (eq. (16.210)) via the Hadamard two-point function $W_\omega(x,x')$ and a differential projector $\mathcal{D}$:

$$\rho_{\mathrm{vac}}(x) = \lim_{x'\to x}\Big[\mathcal{D}(\bar{\Sigma};\nabla,\nabla')\,W_\omega(x,x') - \text{Hadamard counterterms}(\bar{\Sigma})\Big], \tag{16.214}$$

in line with the local covariant renormalization program [245]. Inserting (16.214) into (16.211) and projecting onto the $tt, rr, \theta\theta, t\phi, \phi\phi$ components yields a closed system for $\{A,B,C,D,E\}$ whose leading solutions coincide with the linear-response shifts used to compute δR_H, $\delta\Omega_H$, and T_{eff} (eqs. (16.2), (16.4) and (16.204)).

Towards the Chronogeometric Kerr Solution

To elevate the rotating ansatz to a genuine solution within the IQR framework, we now reconstruct the fully dressed geometry from the vacuum structure of entangled PQSs, encoded in the unified metric $\bar{\Sigma}_{\mu\nu} = \bar{g}^{\mathrm{Kerr}}_{\mu\nu} + \bar{t}_{\mu\nu}$ (eq. (16.198) and Definition 16.4). The procedure is as follows.

1. **Relational Levi–Civita connection and Christoffel symbols.** Compute the Christoffel symbols of the unified geometry, $\bar{\Gamma}^{\lambda}{}_{\mu\nu}[\bar{\Sigma}]$, or equivalently decompose them around Kerr as

$$\begin{aligned}\bar{\Gamma}^{\lambda}{}_{\mu\nu} &= \Gamma^{\lambda}{}_{\mu\nu}\Big[\bar{g}^{\mathrm{Kerr}}\Big] + \Delta\Gamma^{\lambda}{}_{\mu\nu}\Big[\,\bar{t};\,\bar{g}^{\mathrm{Kerr}}\,\Big],\\ \Delta\Gamma^{\lambda}{}_{\mu\nu} &= \tfrac{1}{2}\,\bar{g}^{\lambda\sigma}\Big(\nabla_\mu t_{\nu\sigma} + \nabla_\nu t_{\mu\sigma} - \nabla_\sigma t_{\mu\nu}\Big) + \mathcal{O}(t^2).\end{aligned} \tag{16.215}$$

   where ∇ is the Kerr covariant derivative and $t_{\mu\nu} \equiv \bar{t}_{\mu\nu}$. The profiles $\{A,B,C,D,E\}$ enter via $\bar{t}_{\mu\nu}$ as in eq. (16.199).

2. **Unified Ricci tensor (linear and nonlinear parts).** Use $\bar{\Gamma}$ to evaluate $\bar{R}_{\mu\nu}[\bar{\Sigma}]$. Since $R_{\mu\nu}[\bar{g}^{\rm Kerr}] = 0$,
$$\bar{R}_{\mu\nu}[\bar{\Sigma}] = \delta R_{\mu\nu}[t; \bar{g}^{\rm Kerr}] + \mathcal{Q}_{\mu\nu}[t, \nabla t]\,, \tag{16.216}$$
with the linear piece governed by the Lichnerowicz operator Δ_L on symmetric 2-tensors over the Kerr background:
$$\begin{aligned} \delta R_{\mu\nu} &= \tfrac{1}{2}\,(\Delta_L t)_{\mu\nu}, \\ (\Delta_L t)_{\mu\nu} &= -\nabla^2 t_{\mu\nu} - 2\,R_{\mu\alpha\nu\beta}\,t^{\alpha\beta} + \nabla_\mu\nabla^\alpha t_{\alpha\nu} + \nabla_\nu\nabla^\alpha t_{\alpha\mu} - \nabla_\mu\nabla_\nu t. \end{aligned} \tag{16.217}$$
where $t := t^\alpha{}_\alpha$ and $R_{\mu\alpha\nu\beta}$ is the Kerr Riemann tensor. The term $\mathcal{Q}_{\mu\nu} = \mathcal{O}(t^2)$ collects quadratic (and higher) corrections.

3. **Insert into the Unified Vacuum Field Equation.** Equate curvature and dressed vacuum source as in Definition 16.6:
$$\bar{R}_{\mu\nu}[\bar{\Sigma}] = \mathcal{R}^{(\rm vac)}_{\mu\nu}[\rho_{\rm vac}(r,\theta)]\,, \tag{16.218}$$
with $\mathcal{R}^{(\rm vac)}_{\mu\nu}$ constrained by stationarity/axisymmetry and built from local covariant invariants of $\bar{\Sigma}$ and the vorticity $\omega_{\mu\nu}[\bar{\Sigma}]$ (Definition 16.5 and eq. (16.213)). Projecting (16.218) onto $tt, rr, \theta\theta, t\phi, \phi\phi$ yields a coupled nonlinear PDE system for $\{A, B, C, D, E\}$ on (r, θ), subject to asymptotic/axis conditions (eq. (16.201)) and horizon regularity (eq. (16.203)).

4. **Model the rotationally dressed vacuum energy.** Specify $\rho_{\rm vac}(r,\theta)$ via the Holo-State expectation $\langle\hat{\rho}^{(\rm eff)}(x)\rangle$ (Definition 16.5 and eq. (16.210)), using local covariant renormalization (Hadamard point-splitting) [245], and a symmetry-compatible derivative expansion (e.g. functions of $\omega_{\mu\nu}\omega^{\mu\nu}$, $\bar{R}$, $\bar{R}_{\mu\nu}\chi^\mu\chi^\nu, \ldots$; cf. the schematic expansion preceding Definition 16.6). This captures:

   - *Centrifugal entanglement* (rotation-enhanced mode occupation/weights),
   - *Polar anisotropy* (spin-aligned dressing modes breaking spherical symmetry),
   - *Phase/mode locking* near the ergoregion and in the inner domain, consistent with holosymmetric constraints.

Outcome and observables. A self-consistent solution of (16.218) with the above boundary data defines the *Chronogeometric Kerr Solution*: a rotationally dressed, operator-theoretic generalization of Kerr. Its horizon data $\{R_H, \Omega_H, \kappa_{\rm eff}\}$ follow from the unified horizon conditions and surface gravity (eqs. (16.2), (16.4) and (16.203)), with small-spin/weak-correction limits given in eq. (16.204). Geometric rotation is carried by $\bar{g}_{t\phi}$, while *quantum-relational rotation* is encoded by $\bar{t}_{t\phi}$ through the profile $D(r,\theta)$ (eq. (16.199)), providing a direct bridge from Holo-State microphysics to ringdown/shadow/ISCO phenomenology.

Remark (gauge/ellipticity). One may impose a Kerr-harmonic (Lorenz-type) gauge on $t_{\mu\nu}$ relative to $\bar{g}^{\rm Kerr}$ so that Δ_L is elliptic on stationary slices, improving well-posedness for the boundary-value problem defined by (16.218) together with eq. (16.201).

16.4.1 Embedding Kerr into the Emergent Chronogeometric Framework

In the IQR paradigm, classical geometry arises as the *expectation* of operator-valued modal structures over the PQS Hilbert bundle, organized by the global Quantum Holo-State. The rotating black hole corresponds to a *stationary, axisymmetric coherence phase* of this state, whose effective (chrono-)geometry is encoded in the unified tensor $\bar{\Sigma}_{\mu\nu}$ (eq. (16.198) and Definition 16.4).

Definition 16.7 (The Kerr Geometry as an Emergent Chronogeometric Phase)**.** Let ω denote the algebraic state of the PQS ensemble and $|\Omega_\omega\rangle$ its GNS vector (cf. Definition 1.34). The emergent Kerr phase is the stationary, axisymmetric Holo-State sector in which the expectation of the operator-valued geometric and temporal tensors yields

$$\boxed{\bar{\Sigma}^{\mathrm{Kerr}}_{\mu\nu}(x) := \langle \Omega_\omega \,|\, \hat{g}^{\mathrm{Kerr}}_{\mu\nu}(x) + \hat{t}^{\mathrm{Kerr}}_{\mu\nu}(x) \,|\, \Omega_\omega \rangle,} \tag{16.219}$$

subject to the symmetry constraints

$$\mathcal{L}_{\partial_t} \bar{\Sigma}^{\mathrm{Kerr}}_{\mu\nu} = 0, \qquad \mathcal{L}_{\partial_\phi} \bar{\Sigma}^{\mathrm{Kerr}}_{\mu\nu} = 0, \qquad \bar{\Sigma}^{\mathrm{Kerr}}_{\mu\nu}(r, \pi - \theta) = \bar{\Sigma}^{\mathrm{Kerr}}_{\mu\nu}(r, \theta). \tag{16.220}$$

Here $\hat{g}^{\mathrm{Kerr}}_{\mu\nu}$ captures the operator-valued geometric sector (frame-dragging included), while $\hat{t}^{\mathrm{Kerr}}_{\mu\nu}$ encodes the temporal phase-flow curvature. In the coordinate description used throughout, $\hat{t}^{\mathrm{Kerr}}_{\mu\nu}$ reduces, after expectation, to the five-profile ansatz A, B, C, D, E in eq. (16.199). The state dependence enters through the rotationally dressed vacuum sector (Definition 16.5).

Unified dynamics and sourcing. The emergent Kerr phase is *not* freely chosen: $\bar{\Sigma}^{\mathrm{Kerr}}_{\mu\nu}$ must satisfy the Unified Vacuum Field Equation

$$\bar{R}_{\mu\nu}[\bar{\Sigma}^{\mathrm{Kerr}}] = \mathcal{R}^{(\mathrm{vac})}_{\mu\nu}[\rho_{\mathrm{vac}}(r, \theta)]\,, \tag{16.221}$$

with ρ_{vac} given by the Holo-State expectation of the effective energy operator (Definition 16.5 and eq. (16.210)) and $\mathcal{R}^{(\mathrm{vac})}_{\mu\nu}$ constrained by local covariance and stationarity (cf. Definition 16.6 and eq. (16.213)). Linearizing $\bar{R}_{\mu\nu}$ around classical Kerr yields the Lichnerowicz operator on $t_{\mu\nu}$ (eq. (16.217)), while nonlinearity is collected in $\mathcal{Q}_{\mu\nu}[t, \nabla t]$. This closes a coupled PDE system for A, B, C, D, E on (r, θ), with boundary/horizon regularity in eqs. (16.201) and (16.203).

Classical limit and uniqueness. In the *classical limit* (vanishing temporal sector, trivial vacuum dressing) one has

$$A = B = C = D = E = 0, \qquad \rho_{\mathrm{vac}} \to \rho_0 \text{ (constant)}, \qquad \bar{\Sigma}^{\mathrm{Kerr}}_{\mu\nu} \to \bar{g}^{\mathrm{Kerr}}_{\mu\nu},$$

and $\bar{R}_{\mu\nu}[\bar{g}^{\mathrm{Kerr}}] = 0$. The limiting geometry is the unique, asymptotically flat, stationary axisymmetric vacuum black hole—Kerr—by the standard uniqueness theorems [647, 648, 649]. Thus the emergent phase reduces continuously to the classical Kerr manifold.

Thermodynamic and horizon data. Within the phase, the ergosurface and event horizon are determined from $\bar{\Sigma}^{\mathrm{Kerr}}$ by eqs. (16.202) and (16.203), while the effective surface gravity and temperature follow from eqs. (16.2) and (16.4) (with Schwarzschild/Kerr limits reproducing the standard Hawking temperature [245]). These horizon-value data—$\{R_H, \Omega_H, \kappa_{\mathrm{eff}}\}$—are functionals of the profiles A, B, C, D, E and therefore of ρ_{vac}, providing the bridge to observable ringdown/shadow features (eq. (16.204)).

Roadmap to solutions. To *construct* the phase explicitly we proceed as in Section 16.4: compute $\Delta\Gamma[\bar{t}]$ (eq. (16.215)), evaluate $\bar{R}_{\mu\nu}$ and project (16.221) onto the $tt, rr, \theta\theta, t\phi, \phi\phi$ components, model $\rho_{\mathrm{vac}}(r,\theta)$ via local covariant renormalization of correlators (Hadamard/point-splitting) in the Holo-State [245], and solve the resulting elliptic boundary-value system with eq. (16.201). Perturbative (small-spin, weak-dressing) regimes reduce immediately to eq. (16.204); beyond them, numerical continuation offers a route to the full Chronogeometric Kerr solution.

16.4.2 Operator-Valued Angular Momentum and Frame-Dragging

In the IQR framework, *angular momentum* is an emergent relational observable generated by coherent rotational correlations within the PQS ensemble. Its classical manifestation—frame dragging—is recovered as the off–diagonal $t\phi$ structure in the unified tensor $\bar{\Sigma}_{\mu\nu}$ (cf. eq. (16.199)), while its microscopic origin is operator–valued and state–dependent through the Holo-State expectation (cf. eq. (16.210)).

Definition 16.8 (Relational Angular Momentum and the Operator Origin of Frame Dragging)**.** Let $\hat{x}^\mu(x)$ and $\hat{p}_\nu(x')$ denote emergent position and momentum operators acting on the modal fibers over points x, x' in the PQS bundle. Their relational canonical structure is encoded in

$$\boxed{\left[\hat{x}^\mu(x),\, \hat{p}_\nu(x')\right] \;=\; i\hbar\, \delta^\mu{}_\nu\, \hat{\delta}_\ell(x,x'),} \tag{16.222}$$

where $\hat{\delta}_\ell(x,x')$ is a coherence–length–regulated relational delta supported on the bi-local kernel scale (cf. the bi-local interaction $\mathcal{U}_{\Phi\Theta}$ in Definition 10.222). The *operator–valued angular–momentum tensor* is the anti-symmetric bilocal observable

$$\boxed{\hat{J}^{\mu\nu}(x,x') := \tfrac{1}{2}(\hat{x}^\mu(x)\hat{p}^\nu(x') - \hat{x}^\nu(x)\hat{p}^\mu(x') + \mathrm{h.c.})\,, \qquad \hat{J}^{\mu\nu} = -\hat{J}^{\nu\mu},} \tag{16.223}$$

whose Holo-State expectation yields the relational angular momentum

$$\bar{J}^{\mu\nu}(x,x') \;:=\; \left\langle \Omega_\omega \right|\, \hat{J}^{\mu\nu}(x,x')\, \left|\Omega_\omega\right\rangle. \tag{16.224}$$

In the coincidence limit $x' \to x$, $\bar{J}^{\mu\nu}(x,x)$ reproduces the local orbital angular–momentum density, and in the classical limit reduces to the Komar/Noether angular momentum for the axial Killing field ∂_ϕ (cf. [649, 245]).

Sourcing $t\phi$ and frame dragging. The off–diagonal unified component is sourced by bilinear couplings of $\bar{J}^{\mu\nu}$ to the chrono–geometric vorticity $\omega_{\mu\nu}[\bar{\Sigma}]$ in the vacuum source tensor $\mathcal{R}^{(\mathrm{vac})}_{\mu\nu}$ (cf. Definition 16.6 and eq. (16.213)):

$$\mathcal{R}^{(\mathrm{vac})}_{t\phi} \supset \alpha_J\, \omega_{t\lambda}\, \bar{J}_\phi{}^\lambda + \alpha'_J\, \chi_{(t}\chi^\lambda \bar{J}_{\phi)\lambda} + \cdots, \qquad \chi^a = \partial_t, \tag{16.225}$$

so that, via the Unified Vacuum Field Equation $\bar{R}_{t\phi}[\bar{\Sigma}] = \mathcal{R}^{(\mathrm{vac})}_{t\phi}$ (eq. (16.221)), the profile $D(r,\theta)$ in eq. (16.199) is determined by the Holo-State correlator $\bar{J}^{\mu\nu}$ together with $\omega_{\mu\nu}$. Equivalently,

$$\mathrm{project}\Big(\bar{R}_{t\phi}[\bar{g}^{\mathrm{Kerr}} + \bar{t}]\Big) = \mathrm{project}\Big(\mathcal{R}^{(\mathrm{vac})}_{t\phi}[\rho_{\mathrm{vac}}; \bar{J}, \omega]\Big) \implies D(r,\theta) \text{ from } \{\bar{J}^{\mu\nu}, \omega_{\mu\nu}\}, \tag{16.226}$$

with the projection taken in the $t\phi$ channel on stationary, axisymmetric slices.

Local/quasi-local limits and classical correspondence. (i) In the *coincidence/local* limit, $\hat{\delta}_\ell(x,x') \to \delta(x,x')$ and $\bar{J}^{\mu\nu}(x,x)$ generates the standard orbital angular momentum; the resulting $t\phi$ structure reduces to the classical Lense–Thirring frame dragging encoded in $g^{\mathrm{Kerr}}_{t\phi}$ (see [649]). (ii) In the *weak–dressing* limit $A,\ldots,E \to 0$ with $\rho_{\mathrm{vac}} \to \rho_0$, $\bar{\Sigma}_{t\phi} \to \bar{g}^{\mathrm{Kerr}}_{t\phi}$ and $\Omega_H, \kappa_{\mathrm{eff}}$ reduce to their Kerr values (cf. eqs. (16.2) and (16.204)). (iii) Beyond the local limit, the bilocal structure in $\hat{J}^{\mu\nu}$ encodes hololocal correlations that feed the $t\phi$ channel through $\mathcal{R}^{(\mathrm{vac})}$ (eq. (16.213)), refining frame dragging into a *quantum–relational coherence* effect.

Remark 16.9 (Frame dragging as quantum–relational coherence)**.** The twisting of spacetime is upgraded from a purely classical curvature phenomenon to a macroscopic signature of coherent PQS phases: the profile $D(r,\theta)$ is fixed by $\bar{J}^{\mu\nu}$ and $\omega_{\mu\nu}$ via (16.225)–(16.226), with the overall normalization determined by the dressed vacuum through ρ_{vac} (Definition 16.5). This connects the operator layer (bi-local correlators), the geometric layer ($\omega_{\mu\nu}$, $\bar{R}_{t\phi}$), and the observable layer (Ω_H via eq. (16.203), and T_{eff} via eqs. (16.2) and (16.4)). For the renormalized operator expectations and their locality properties we follow the standard QFT-in-curved-spacetime scheme [245].

16.4.3 The Inner Core: Ring Singularity and the Non-Classical Locus

Classically, the Kerr solution possesses a ring singularity at $\rho^2 = r^2 + a^2\cos^2\theta = 0$ (i.e. $r = 0, \theta = \frac{\pi}{2}$), where curvature invariants diverge and geodesic predictability fails [649, 245]. In the IQR framework, this locus is reinterpreted as a *pre-geometric phase* where relational distinguishability collapses. The inner core is a domain in which the temporal tensor saturates ($A,B,C,D,E = \mathcal{O}(1)$) and the regime tracker ξ reaches the non-classical regime (cf. eqs. (16.199), (16.201) and (16.220)).

Definition 16.10 (Kerr Ring Core as a Pre-Geometric Entanglement Condensate)**.** Let $\mathcal{X}_q$ denote the PQS fiber bundle and $|\Omega_\omega\rangle$ the Holo-State (Definition 1.34). The *ring core* $\mathfrak{B}_{\mathrm{ring}} \subset \mathcal{X}_q$ is the maximal connected region satisfying the following *relational collapse* conditions:

$$\boxed{\forall\, x, x' \in \mathfrak{B}_{\mathrm{ring}}: \qquad \Big\langle \Omega_\omega \Big|\, \hat{\mathbf{d}}(x,x') \,\Big| \Omega_\omega \Big\rangle = 0, \qquad \Big\langle \Omega_\omega \Big|\, \hat{t}_{\mathrm{em}}(x,x') \,\Big| \Omega_\omega \Big\rangle = 0.} \tag{16.227}$$

Here $\hat{\mathbf{d}}(x,x')$ is the *relational distance operator* of Section 5.3.8, and $\hat{t}_{\rm em}(x,x')$ is the *emergent temporal interval* of Section 5.3.8; their vanishing signals the loss of spatial separability and causal ordering, i.e. maximal phase locking of PQS modes. Equivalently, the two-point correlator that induces the effective metric on exterior slices becomes rank-deficient on $\mathfrak{B}_{\rm ring}$.

Characterization and matching.

(i) Coherence threshold. The boundary $\partial\mathfrak{B}_{\rm ring}$ is defined by the first nonzero eigenvalue of the coherence kernel crossing a fixed threshold, $\lambda_{\min}(\mathcal{C}[\omega;\bar{\Sigma}]) \downarrow 0$, equivalently by $\xi(x) = \xi_\star$ for a critical tracker value (cf. Definition 16.5).

(ii) Unified-field consistency (distributional form). Inside $\mathfrak{B}_{\rm ring}$ the Unified Vacuum Field Equation (Definition 16.6)

$$\bar{R}_{\mu\nu}[\bar{\Sigma}] = \mathcal{R}^{(\rm vac)}_{\mu\nu}[\rho_{\rm vac}]$$

holds in the *sense of distributions* on the PQS bundle; $\rho_{\rm vac}$ saturates the local covariant bounds set by the Holo-State (Hadamard renormalization) [245]. This replaces classical curvature blow-ups with a regular condensate supported by temporal stresses.

(iii) Exterior matching. Across $\partial\mathfrak{B}_{\rm ring}$, the profiles A, B, C, D, E and their normal derivatives satisfy the continuity conditions implied by $\bar{R}_{\mu\nu} = \mathcal{R}^{(\rm vac)}_{\mu\nu}$ (eq. (16.218)), ensuring smooth attachment to the exterior Kerr-like region and preserving axisymmetry and stationarity (eq. (16.220)).

Lemma 16.11 (Classical limit and locus)**.** *In the weak-dressing limit $A,\dots,E \to 0$ and $\rho_{\rm vac} \to \rho_0$ (constant), $\mathfrak{B}_{\rm ring}$ collapses to the classical ring $\{r=0,\theta=\pi/2\}$, and $\bar{\Sigma}_{\mu\nu} \to \bar{g}^{\rm Kerr}_{\mu\nu}$ with the standard curvature singularity [649]. Conversely, for finite dressing obeying the UVFE and the renormalized vacuum constraints, the emergent invariants built from $\bar{\Sigma}$ remain bounded on $\mathfrak{B}_{\rm ring}$, and geodesic incompleteness is replaced by a* pre-geometric *condensate sustained by maximal relational coherence (Definitions 16.5 and 16.6).*

Operational footprint. Maximal phase locking in $\mathfrak{B}_{\rm ring}$ induces:

(i) Spectral gapping. Spectral gapping of interior modes, feeding back into the exterior through the state-dependent S-matrix and modifying late-time emission microstructure (Section 16.3 and eq. (16.172)).

(ii) Enhanced $t\phi$ coupling. Enhanced $t\phi$ coupling (profile D) via the $\mathcal{R}^{(\rm vac)}_{t\phi}$ channel, refining frame dragging beyond classical Kerr (Section 16.4.2 and eq. (16.225)).

(iii) Horizon-data shifts. Horizon-data shifts $\{\delta R_H, \delta\Omega_H, \kappa_{\rm eff}\}$ propagated through the boundary-value problem for $\{A,\dots,E\}$ (eqs. (16.2), (16.4) and (16.204)).

Remark 16.12 (The ring as a holosymmetric conduit)**.** The topology of $\mathfrak{B}_{\rm ring}$ supports a closed holosymmetric cycle of entanglement (phase-locked sector of the Holo-State), acting as a *coherence conduit* between relational sectors. In our formalism this appears as a global constraint among the source coefficients in $\mathcal{R}^{(\rm vac)}_{\mu\nu}$ (holosymmetric phase locking; cf.

Definition 16.5). While suggestive of "portal-like'' behavior at the operator level, any trans-historical tunnelling or modal bifurcation remains a *theoretical possibility* contingent on the global Holo-State and boundary conditions—see Remark 16.9 for the analogous $t\phi$ channel discussion.

16.4.4 Chronogeometric Vorticity and the Emergence of Horizons

In classical Kerr geometry, the *ergosphere* is the region where the stationary Killing field $\xi^a = \partial_t$ becomes spacelike, i.e. $\xi^2 = g_{tt} > 0$, with the ergosurface given by $g_{tt} = 0$ [649, 245]:

$$\boxed{g_{tt} = \xi^\mu \xi_\mu > 0 \quad \Longleftrightarrow \quad \text{inside the ergosphere,}} \tag{16.228}$$

so that no worldline can remain stationary relative to infinity (irreducible frame dragging).

Definition 16.13 (Chronogeometric vorticity and the unified ergoregion)**.** Within the IQR framework, local temporal phase evolution on the PQS bundle is generated by an *operator-valued temporal potential* $\hat{\tau}_\mu(x)$. Its (relational) curl defines the *chronogeometric vorticity tensor*

$$\boxed{\hat{\omega}_{\mu\nu}(x) := \nabla_\mu \hat{\tau}_\nu(x) - \nabla_\nu \hat{\tau}_\mu(x), \qquad \bar{\omega}_{\mu\nu}(x) := \langle \Omega_\omega | \hat{\omega}_{\mu\nu}(x) | \Omega_\omega \rangle,} \tag{16.229}$$

with ∇ the relational (Levi–Civita) derivative on $\bar{\Sigma}$ (cf. Definition 6.7). A nonzero $\bar{\omega}_{\mu\nu}$ encodes rotational "twist'' of the temporal flow. The scalar density $\Omega^2(x) := \frac{1}{2}\,\bar{\omega}_{\mu\nu}\bar{\omega}^{\mu\nu}$ will serve as a vorticity measure.

The *unified ergosurface* is defined by the vanishing of the tt-component of the *unified* tensor (cf. eq. (16.202)):

$$\boxed{\mathcal{E} \;:=\; \Big\{ x \;\Big|\; \bar{\Sigma}_{tt}(x) = \langle \Omega_\omega |\, \hat{g}_{tt}(x) + \hat{t}_{tt}(x)\, | \Omega_\omega \rangle = 0 \Big\},} \tag{16.230}$$

so that the ergoregion is $\{\bar{\Sigma}_{tt} > 0\}$. In the stationary, axisymmetric ansatz (eq. (16.199)) one has $\bar{\Sigma}_{tt} = g_{tt}^{\text{Kerr}} - A(r,\theta)$; hence the *shift* of the ergosurface relative to Kerr is controlled by the lapse profile A, which is in turn sourced by the dressed vacuum through the UVFE (Definition 16.6).

Vorticity–ergoregion link (source level). The vacuum source tensor admits vorticity contributions consistent with stationarity/axisymmetry (cf. eq. (16.213)):

$$\mathcal{R}^{(\text{vac})}_{\mu\nu} \;\supset\; \alpha_1\, \bar{\omega}_{\mu\lambda}\bar{\omega}_\nu{}^\lambda + \alpha_3\, \bar{R}_{\mu\nu} + \cdots, \qquad \Rightarrow \qquad \bar{R}_{tt}[\bar{\Sigma}] = \mathcal{R}^{(\text{vac})}_{tt}[\rho_{\text{vac}}; \bar{\omega}, \ldots], \tag{16.231}$$

so that, after projecting the UVFE $\bar{R}_{\mu\nu} = \mathcal{R}^{(\text{vac})}_{\mu\nu}$ onto the tt component (eq. (16.221)), the profile $A(r,\theta)$ inherits a dependence on Ω^2. Consequently, the unified ergosurface $\bar{\Sigma}_{tt} = 0$ (eq. (16.230)) is *displaced* by chronogeometric vorticity relative to the classical Kerr location (eq. (16.228)).

Event horizon in the unified setting. The event horizon remains the largest root of the generator norm with $\chi^a = \partial_t + \Omega_H\,\partial_\phi$ (cf. eq. (16.203)):

$$\bar{\Sigma}_{ab}\,\chi^a\chi^b\Big|_{r=R_H} = 0, \qquad \Omega_H = -\frac{\bar{\Sigma}_{t\phi}}{\bar{\Sigma}_{\phi\phi}}\Big|_{r=R_H}, \tag{16.232}$$

reducing to Kerr in the classical limit. Unlike the ergosurface (purely tt), the horizon condition couples both the lapse correction A and the off–diagonal profile D (through $\bar{\Sigma}_{t\phi}$), linking vorticity-driven sourcing ($\bar{\omega}$) to both ergoregion *and* horizon data $\{R_H, \Omega_H, \kappa_{\text{eff}}\}$ (cf. eqs. (16.2), (16.4) and (16.204)).

Classical correspondence. When $A = B = C = D = E \to 0$ and $\rho_{\text{vac}} \to \rho_0$, one recovers the classical relations: $\mathcal{E}:\ g_{tt} = 0$ and $R_H = r_+$ with standard Kerr thermodynamics [649, 245]. The chronogeometric vorticity $\bar{\omega}_{\mu\nu}$ then vanishes, and the unified construction collapses to the textbook picture. å

Remark 16.14 (The Ergosphere as a Dynamical Coherence Interface)**.** Within the unified setting, the ergoregion is not a passive geometric artifact but a *dynamical coherence interface* where temporal dressing becomes operative. Concretely, the boundary is fixed by the unified condition $\bar{\Sigma}_{tt} = 0$ (eq. (16.230)), and its displacement relative to the classical locus $g_{tt} = 0$ (eq. (16.228)) is driven by vorticity-sourced vacuum stresses (eq. (16.231)) acting through the temporal sector profiles in eq. (16.199). In this interface, the operator-valued temporal potential and its curl (eq. (16.229)) encode a torsional response of the rotating vacuum, so that the flow of relational information is twisted by frame-dragging channels set by the Holo-State and the Unified Vacuum Field Equation (Definition 16.6). This elevates "irreducible frame dragging'' from a purely metric feature to a coherence-controlled phenomenon consistent with the classical limit [649, 245].

Remark 16.15 (Triadic Decomposition of the Ergosurface Condition)**.** Because the unified tensor is an expectation over operator sectors (eq. (16.219)), the tt-component naturally decomposes into *correlational* (two-point/operator) contributions, *vacuum* (dressed, state-dependent) contributions, and *spectral* (matter/stress) contributions:

$$\boxed{\bar{\Sigma}_{tt}(x) = \bar{\Sigma}^{(\text{corr})}_{tt}(x) + \bar{\Sigma}^{(\text{vac})}_{tt}(x) + \bar{\Sigma}^{(\text{spec})}_{tt}(x) = 0,} \tag{16.233}$$

where:

- **(i)** $\bar{\Sigma}^{(\text{corr})}_{tt}$ collects Holo-State correlators such as the angular-momentum and vorticity channels (e.g. $\bar{J}^{\mu\nu}$, $\bar{\omega}_{\mu\nu}$; cf. Definition 16.8 and eqs. (16.225) and (16.229)).
- **(ii)** $\bar{\Sigma}^{(\text{vac})}_{tt}$ encodes the rotationally dressed vacuum sector sourced by ρ_{vac} (Definitions 16.5 and 16.6).
- **(iii)** $\bar{\Sigma}^{(\text{spec})}_{tt}$ represents spectral (matter/flux) contributions consistent with stationarity/axisymmetry when present.

Thus the ergosurface is a *relationally determined* phase boundary, not a purely geometric sign flip. Two rotating objects with the same (M, a) can, in principle, exhibit distinct ergospheres if their correlational or vacuum sectors differ, making $\mathcal{E}$ a surface of *chronogeometric equilibrium* among information, vacuum, and spectral influences.

Remark 16.16 (Ontological Reinterpretation of Kerr Features)**.** In the operator-theoretic formulation, familiar Kerr features acquire a relational meaning:

(i) Ergosurface.
Defined by the unified condition $\bar{\Sigma}_{tt}(x) = \langle\Omega_\omega|\hat{g}_{tt}(x)+\hat{t}_{tt}(x)|\Omega_\omega\rangle = 0$ (eq. (16.230)), not by $g_{tt} = 0$ alone; it is controlled by the temporal profile $A(r, \theta)$ and vacuum/vorticity sourcing (eq. (16.231)).

(ii) Ring core.
The "singularity'' corresponds to a collapse of relational distinguishability—$\langle\hat{\mathbf{d}}(x, x')\rangle = \langle\hat{t}_{\rm em}(x, x')\rangle = 0$—defining a pre-geometric condensate region $\mathfrak{B}_{\rm ring}$ (Definition 16.10).

(iii) Frame dragging.
The $t\phi$ channel is sourced by operator-valued angular momentum $\hat{J}^{\mu\nu}$ through the vacuum tensor, fixing the $D(r, \theta)$ profile (Definition 16.8 and eqs. (16.199) and (16.225)); the classical $g^{\rm Kerr}_{t\phi}$ is recovered in the weak-dressing limit [649, 245].

Accordingly, classical curvature data are seen as macroscopic projections of operator-level coherence patterns of the Quantum Holo-State, with the classical picture recovered smoothly via $\{A, B, C, D, E\} \to 0$ and $\rho_{\rm vac} \to \rho_0$.

16.4.5 Kerr Geometry as a Relational Coherence Phase

Definition 16.17 (Kerr Geometry as a Relational Coherence Phase)**.** Within the IQR framework, a rotating black hole is realized as a *stationary, axisymmetric relational coherence phase* of the Quantum Holo-State rather than as a purely classical manifold. The effective chronogeometry is the unified expectation $\bar{\Sigma}_{\mu\nu} = \langle\Omega_\omega|\hat{g}_{\mu\nu}+\hat{t}_{\mu\nu}|\Omega_\omega\rangle$ (eq. (16.219)), obeying the symmetry constraints (eq. (16.220)) and the Unified Vacuum Field Equation (Definition 16.6 and eq. (16.221)). The resulting phase is a self-coherent attractor selected by holosymmetric constraints in the dressed vacuum sector (Definition 16.5).

This phase is characterized by the following definitive properties:

(i) Persistent operator-valued angular momentum.
The rotational character is encoded in a non-vanishing Holo-State expectation of the bilocal angular-momentum tensor (Definition 16.8):

$$\boxed{\langle\Omega_\omega|\,\hat{J}^{t\phi}(x, x')\,|\Omega_\omega\rangle \;\neq\; 0,} \tag{16.234}$$

signalling long-range rotational coherence in the PQS ensemble. Through the source channel $\mathcal{R}^{(\rm vac)}_{t\phi}$ (eq. (16.225)), this persistent $\bar{J}^{t\phi}$ fixes the unified off–diagonal component $\bar{\Sigma}_{t\phi}$ and the temporal profile $D(r, \theta)$ in the line element (eq. (16.199)), refining classical frame dragging.

(ii) Entanglement-induced ergoregion.
The ergosphere is a dynamical interface determined by the *unified* condition $\bar{\Sigma}_{tt} = 0$ (eq. (16.230)), not by $g_{tt} = 0$ alone (eq. (16.228)). In the stationary ansatz, $\bar{\Sigma}_{tt} = g_{tt}^{\text{Kerr}} - A(r,\theta)$, so the lapse correction A— sourced by chronogeometric vorticity $\bar{\omega}_{\mu\nu}$ through the vacuum tensor (eqs. (16.229) and (16.231))— displaces the ergosurface relative to classical Kerr and encodes quantum-relational rotation.

(iii) Central ring entanglement condensate (non-classical locus).
The classical ring singularity is reinterpreted as a pre-geometric *entanglement condensate* $\mathfrak{B}_{\text{ring}}$ (Definition 16.10), a domain of maximal relational coherence where relational distinguishability collapses:

$$\boxed{\forall\, x, x' \in \mathfrak{B}_{\text{ring}} : \quad \langle\Omega_\omega|\, \hat{\mathbf{d}}(x,x')\, |\Omega_\omega\rangle = 0, \qquad \langle\Omega_\omega|\, \hat{t}_{\text{em}}(x,x')\, |\Omega_\omega\rangle = 0,} \tag{16.235}$$

(cf. eq. (16.227)). Matching across $\partial\mathfrak{B}_{\text{ring}}$ is enforced by the unified field equations (eq. (16.218)), yielding a regular interior sustained by temporal stresses, in contrast to the classical curvature divergence [649, 245].

Classical correspondence. In the weak-dressing limit $A, \ldots, E \to 0$ and $\rho_{\text{vac}} \to \rho_0$, $\bar{\Sigma}_{\mu\nu} \to \bar{g}_{\mu\nu}^{\text{Kerr}}$ and the phase reduces to the unique stationary, axisymmetric vacuum black hole—Kerr—by the standard uniqueness results [647, 648, 649].

Remark 16.18 (The Kerr attractor as a topological/coherence phase)**.** These features jointly present Kerr as a *topological phase of quantum relationality*: an operator-level coherence pattern of the Holo-State whose macroscopic projection is the familiar rotating geometry. Angular momentum ($\bar{J}^{t\phi}$), the ergoregion ($\bar{\Sigma}_{tt} = 0$), and the ring core ($\mathfrak{B}_{\text{ring}}$) appear as tightly coupled manifestations of the same coherence mechanism, with the classical manifold recovered continuously in the limit described above.

16.4.6 The Operator-Theoretic Kerr Reconstruction

Definition 16.19 (Operator-Theoretic Kerr Reconstruction)**.** In the IQR framework the rotating black hole is an *emergent relational coherence phase* of the non-factorizable Quantum Holo-State rather than a bare classical manifold. Its effective chronogeometry is the unified expectation

$$\boxed{\bar{\Sigma}_{\mu\nu}^{\text{Kerr}}(x) := \Big\langle \Omega_\omega \Big|\, \hat{g}_{\mu\nu}^{\text{Kerr}}(x) + \hat{t}_{\mu\nu}^{\text{Kerr}}(x)\, \Big| \Omega_\omega \Big\rangle,} \tag{16.236}$$

which is the Kerr-phase instance of eq. (16.219). Stationarity and axisymmetry are implemented by $\mathcal{L}_{\partial_t}\bar{\Sigma}_{\mu\nu} = 0 = \mathcal{L}_{\partial_\phi}\bar{\Sigma}_{\mu\nu}$ (eq. (16.220)), and the temporal sector reduces—after expectation—to the five-profile ansatz A, B, C, D, E in eq. (16.199). The unified tensor obeys the UVFE $\bar{R}_{\mu\nu} = \mathcal{R}_{\mu\nu}^{(\text{vac})}[\rho_{\text{vac}}]$ (Definition 16.6 and eq. (16.221)), which ties the geometry to the dressed vacuum ρ_{vac} (Definition 16.5). In the weak-dressing limit one recovers classical Kerr [649, 245].

1. Relational origin of angular momentum.

Definition 16.20 (Relational Angular Momentum)**.** Let $\hat{x}^\mu(x)$, $\hat{p}_\nu(x')$ be emergent position/-momentum operators on the PQS bundle. The bilocal, anti-symmetric operator

$$\boxed{\hat{J}^{\mu\nu}(x,x') := \tfrac{1}{2}(\hat{x}^\mu(x)\hat{p}^\nu(x') - \hat{x}^\nu(x)\hat{p}^\mu(x') + \text{h.c.}), \qquad \hat{J}^{\mu\nu} = -\hat{J}^{\nu\mu},} \tag{16.237}$$

encodes relational angular momentum (Definition 16.8). The global, axial charge is the Holo-State expectation of the Noether/Komar generator associated with $\eta^a = \partial_\phi$:

$$\bar{J} \;=\; \Big\langle \Omega_\omega \Big| \hat{Q}_\eta \Big| \Omega_\omega \Big\rangle \quad \Rightarrow \quad a \;:=\; \frac{\bar{J}}{\bar{M}}, \qquad \bar{M} := \langle \Omega_\omega | \hat{M} | \Omega_\omega \rangle.$$

Thus the Kerr spin parameter a is the *order parameter* of rotational coherence, determined by $\bar{J}^{\mu\nu}$ and the state's mass expectation.

2. Operator realization of frame dragging (the $t\phi$ channel). The unified off–diagonal component is

$$\bar{\Sigma}_{t\phi}(r,\theta) = g^{\text{Kerr}}_{t\phi}(r,\theta) \;-\; D(r,\theta)\,\sin^2\theta, \tag{16.238}$$

with D the temporal-sector profile from eq. (16.199). At the source level, the vacuum tensor admits angular-momentum/vorticity couplings (eq. (16.225)), so the UVFE projected on the $t\phi$ component (eq. (16.221)) yields

$$\begin{aligned} \text{proj}_{t\phi}\Big(\bar{R}_{\mu\nu}[\bar{g}^{\text{Kerr}} + \bar{t}]\Big) &= \text{proj}_{t\phi}\Big(\mathcal{R}^{(\text{vac})}_{\mu\nu}[\rho_{\text{vac}}; \bar{J}, \bar{\omega}]\Big) \\ &\Longrightarrow\; D(r,\theta) \;\; \text{from} \;\; \{\bar{J}^{\mu\nu}, \bar{\omega}_{\mu\nu}\} \qquad \text{(eq. (16.226)).} \end{aligned}$$

A convenient *rotational-coherence density* is the local operator

$$\hat{\mathcal{R}}_\phi(x) := \epsilon_\phi{}^{\mu\nu}\, \hat{x}_\mu(x)\hat{p}_\nu(x),$$

whose expectation contributes to the same source channel; unlike a naive identification $\bar{g}_{t\phi} \equiv \langle \hat{\mathcal{R}}_\phi \rangle$, eq. (16.238) makes clear that $\bar{\Sigma}_{t\phi}$ is fixed by solving the UVFE with the $\bar{J}$– and $\bar{\omega}$– dependent source (eq. (16.231)).

3. Temporal curvature and the ergosurface. Local temporal phase evolution is generated by $\hat{\tau}_\mu$; the *chronogeometric vorticity* is

$$\boxed{\hat{\omega}_{\mu\nu} := \nabla_\mu \hat{\tau}_\nu - \nabla_\nu \hat{\tau}_\mu, \qquad \bar{\omega}_{\mu\nu} = \langle \Omega_\omega | \hat{\omega}_{\mu\nu} | \Omega_\omega \rangle,} \tag{16.239}$$

as in eq. (16.229). Its quadratic scalar $\Omega^2 = \frac{1}{2}\bar{\omega}_{\mu\nu}\bar{\omega}^{\mu\nu}$ enters $\mathcal{R}^{(\text{vac})}$ (eq. (16.231)), sourcing the lapse profile $A(r,\theta)$. The unified ergosurface is the locus $\bar{\Sigma}_{tt} = 0$ (eq. (16.230)), and in the stationary ansatz $\bar{\Sigma}_{tt} = g^{\text{Kerr}}_{tt} - A$; hence vorticity-driven vacuum dressing displaces the ergosurface relative to the classical condition $g_{tt} = 0$ (eq. (16.228)).

4. Stationary Kerr as a coherence fixed point.

Definition 16.21 (The Kerr Attractor as a Stationary Coherence Phase)**.** In the IQR framework "time" is *emergent*: stationarity refers to invariance under the *Killing-flow parameter* of the stationary symmetry, not to a fundamental microscopic time. The Kerr phase is stationary and axisymmetric at both the geometric and operator levels. In particular, the Fourfold Inter–Modal Correlator is invariant along the stationary flow:

$$\boxed{\frac{d}{dt_K}\,\hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'}) \;=\; 0,} \tag{16.240}$$

where t_K denotes the parameter of the Killing flow generated by ∂_t. Axisymmetry similarly implies invariance under the azimuthal flow generated by ∂_ϕ. These conditions are solved together with the UVFE and boundary data (Definition 16.6 and eqs. (16.201) and (16.203)).

Equivalent formulations (for readers with different backgrounds).

(i) Geometric / Lie–derivative form

The goemtric Lie-derivative is

$$\boxed{\mathcal{L}_{\partial_t}\,\hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'}) \;=\; 0, \qquad \mathcal{L}_{\partial_\phi}\,\hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'}) \;=\; 0.} \tag{16.241}$$

(ii) Algebraic QFT / automorphism form

Let $\{\alpha_t\}_{t\in\mathbb{R}}$ and $\{\alpha_\phi\}_{\phi\in\mathbb{R}}$ be the one–parameter groups of *-automorphisms induced by the Killing flows on the observable algebra. Stationarity and axisymmetry mean

$$\boxed{\omega\circ\alpha_t \;=\; \omega, \qquad \omega\circ\alpha_\phi \;=\; \omega \quad\Longleftrightarrow\quad \frac{d}{dt}\,\omega(\alpha_t(A)) \;=\; 0 \;\;\forall A,} \tag{16.242}$$

which is the operator–algebraic version of (16.240) [245, 230].

(iii) Relational–clock form (conditional dynamics)

Choosing a relational clock C whose reading t_C tracks the Killing flow,

$$\boxed{\frac{d}{dt_C}\,\hat{G}_\Lambda(\mathfrak{B}_x,\mathfrak{B}_{x'}) \;=\; 0,} \tag{16.243}$$

which is equivalent to (16.240) when t_C parametrizes ∂_t.

Equivalently, the Hadamard two–point function and the observable net are invariant under the Killing flows generated by ∂_t and ∂_ϕ; i.e. $\omega\circ\alpha_{(t,\phi)}=\omega$ (cf. eqs. (16.201) and (16.203)). These conditions characterize the Kerr phase as a *stationary, self-coherent attractor*; in the weak–dressing limit the construction reduces smoothly to the unique classical Kerr vacuum [647, 648, 649].

These conditions, together with the UVFE and boundary data (eqs. (16.201) and (16.203)), characterize the Kerr phase as a *stationary, self-coherent attractor*. In the classical limit the construction reduces to the unique Kerr vacuum [647, 648, 649].

16.4.7 Holosymmetric Modulation and Kerr Causal Bifurcation

Beyond mass and spin, the Kerr interior exhibits a multi–sheeted causal structure with regions admitting closed timelike curves (CTCs) near the ring locus $\{r = 0, \theta = \pi/2\}$; the global manifold is non–simply connected in the classical analytic extension [649, 245]. In the IQR paradigm, such features are not pathologies but signatures of a transition into a higher–order *relational* regime governed by holosymmetry (cf. the dressed vacuum constraints in Definition 16.5) and by the non-classical ring core $\mathfrak{B}_{\text{ring}}$ (Definition 16.10).

1. Causal bifurcation as a modal topology transition.

Definition 16.22 (Causal bifurcation as modal topology transition)**.** Let $\pi : \mathcal{X}_q \to \mathcal{M}$ be the PQS bundle projection over $(\mathcal{M}, \bar{\Sigma})$, and let $\mathcal{R}_{\text{CTC}} \subset \mathcal{M}$ be the classical Kerr domain with CTCs [649]. For any PQS locus $\mathfrak{B}_x \in \mathcal{X}_q$ with $x := \pi(\mathfrak{B}_x) \in \mathcal{R}_{\text{CTC}}$, the emergent description exhibits a *modal bifurcation*: there exist two inequivalent modal paths $\gamma_1 \neq \gamma_2$ in the fibered configuration space through $\mathfrak{B}_x$ such that the Fourfold Inter–Modal Correlator (see Definition Definition 16.21) satisfies

$$\boxed{\exists\, \gamma_1 \neq \gamma_2 : \qquad \hat{G}_\Lambda(\gamma_1, \gamma_2) \;\neq\; 0,} \tag{16.244}$$

signalling holosymmetry–enabled topological interference between relational histories in the Holo-State.

2. The ring as a holosymmetric interferometric node.

Definition 16.23 (Kerr ring as holosymmetric interferometric node)**.** The ring core $\mathfrak{B}_{\text{ring}}$ (Definition 16.10) functions as a *holosymmetric node* where entire universal histories may be coherently compared. Let $\mathcal{H}_1, \mathcal{H}_2$ be two (classically) distinct causal extensions of the Kerr interior. Define the *holosymmetric interference amplitude*

$$\boxed{\mathcal{I}_{\text{Kerr}} := \left\langle \Omega_\omega \right| \hat{U}^\dagger_{\mathcal{H}_1}\, \hat{U}_{\mathcal{H}_2} \left| \Omega_\omega \right\rangle,} \tag{16.245}$$

where $|\Omega_\omega\rangle$ is the GNS vector of the Quantum Holo-State and $\hat{U}_{\mathcal{H}_i}$ are the history–conditioned evolutions constructed from the state–dependent Møller/S–matrix maps (eq. (16.171); cf. $\Omega_\pm[\bar{\Sigma}, \omega]$). A nonzero $\mathcal{I}_{\text{Kerr}}$ indicates that $\mathcal{H}_1$ and $\mathcal{H}_2$ are coherently entangled across the ring condensate—consistent with unitarity in the emergent dynamics (Section 16.3).

3. Quantum histories and ring–mediated tunnelling.

Definition 16.24 (Quantum histories via ring–singularity tunnelling)**.** The tower of causally connected regions in the classical extension is reinterpreted as a coherent superposition of history branches within the Holo-State:

$$\boxed{|\Omega_\omega\rangle = \sum_{\alpha \in \mathbb{Z}} c_\alpha\, |\Psi^{\text{Kerr}}_\alpha\rangle, \qquad \sum_\alpha |c_\alpha|^2 = 1,} \tag{16.246}$$

where $|\Psi_\alpha^{\mathrm{Kerr}}\rangle$ encodes the relational structure on the α-th causal sheet. The *ring–interference operator* aggregates cross–branch coherences:

$$\boxed{\hat{G}_{\mathrm{ring}} := \sum_{\alpha,\beta} \bar{c}_\alpha c_\beta \, \hat{G}_\Lambda^{(\alpha,\beta)},} \tag{16.247}$$

with $\hat{G}_\Lambda^{(\alpha,\beta)}$ the Fourfold Inter–Modal Correlator acting between branches α and β. Through the UVFE $\bar{R}_{\mu\nu} = \mathcal{R}_{\mu\nu}^{(\mathrm{vac})}$ (Definition 16.6), nonzero matrix elements of $\hat{G}_{\mathrm{ring}}$ feed back into the temporal sector $(A, \ldots, D)$ near the ring, modulating the inner causal structure consistently with stationarity and axisymmetry (eqs. (16.199) and (16.220)).

Classical correspondence and consistency. In the weak–dressing limit $A, \ldots, E \to 0$ with $\rho_{\mathrm{vac}} \to \rho_0$ (Definition 16.5), the interference amplitude $\mathcal{I}_{\mathrm{Kerr}} \to 0$ and $\hat{G}_{\mathrm{ring}}$ becomes block–diagonal; the modal decomposition collapses to a single classical branch, reproducing the standard Kerr analytic extension with its known causal features [649, 245]. Conversely, for finite dressing the holosymmetric constraints keep the emergent dynamics unitary (Section 16.3) while allowing controlled, ring–mediated coherence across branches, with all observable horizon/ergoregion data continuing to be fixed by the unified conditions (eqs. (16.2), (16.4), (16.203) and (16.230)).

The Logical Derivation

1. Premise: a superposition of Kerr histories. We adopt the global-state postulate (cf. eq. (16.246)):

$$\boxed{|\Omega_\omega\rangle \;=\; \sum_{\alpha\in\mathbb{Z}} c_\alpha \, |\Psi_\alpha^{\mathrm{Kerr}}\rangle, \qquad \sum_\alpha |c_\alpha|^2 = 1,} \tag{16.248}$$

where each branch $|\Psi_\alpha^{\mathrm{Kerr}}\rangle$ encodes a self-consistent relational geometry on the α-th causal sheet of the Kerr analytic structure [649, 245].

2. The Fourfold correlator on history sectors. The Fourfold Inter–Modal Correlator $\hat{G}_\Lambda$ (introduced in Definition 16.21) measures relational coherence between configurations. Its branch–resolved components are the projections

$$\boxed{\hat{G}_\Lambda^{(\alpha,\beta)} \;:=\; \Pi_\alpha \, \hat{G}_\Lambda \, \Pi_\beta, \qquad \Pi_\alpha := |\Psi_\alpha^{\mathrm{Kerr}}\rangle\langle\Psi_\alpha^{\mathrm{Kerr}}|.} \tag{16.249}$$

Nonvanishing $\hat{G}_\Lambda^{(\alpha,\beta)}$ with $\alpha \neq \beta$ signals holosymmetry-enabled interference between distinct relational histories (cf. Definition 16.22).

3. Total ring-mediated interference operator. Summing over history pairs with amplitude weights $\overline{c_\alpha}\, c_\beta$ yields the total operator (see eq. (16.247)):

$$\boxed{\hat{G}_{\mathrm{ring}} \;:=\; \sum_{\alpha,\beta} \overline{c_\alpha}\, c_\beta \, \hat{G}_\Lambda^{(\alpha,\beta)},} \tag{16.250}$$

which is the operator counterpart to the holosymmetric interference amplitude $\mathcal{I}_{\text{Kerr}}$ in eq. (16.245). The ring core $\mathfrak{B}_{\text{ring}}$ (Definition 16.10) is the physical locus where this interference is mediated by maximal relational coherence.

Interpretation of $\hat{G}_{\text{ring}}$. Viewed as a coherence matrix on the history index:

(i) Diagonal. $|c_\alpha|^2\, \hat{G}_\Lambda^{(\alpha,\alpha)}$ quantifies intra–history coherence, weighted by branch probability.

(ii) Off–diagonal. $\overline{c_\alpha}\, c_\beta\, \hat{G}_\Lambda^{(\alpha,\beta)}$ $(\alpha \neq \beta)$ quantifies inter–history interference; these terms vanish if the histories decohere.

Thus $\hat{G}_{\text{ring}}$ measures the total holosymmetric coherence of the multi–sheet Kerr state; its nontriviality is the operator expression of ring–mediated tunnelling consistent with the unitary S–matrix dynamics on $\bar{\Sigma}$ (Section 16.3 and eq. (16.171)).

4. Emergence of interferometric topology.

Definition 16.25 (Interferometric topology via modal holonomy)**.** Let $\hat{\tau}_\mu$ be the operator–valued temporal potential (cf. eq. (16.229)) on the modal Hilbert bundle. For any closed loop $\gamma : S^1 \to \mathcal{X}_q$ that encircles the ring condensate, the *relational holonomy* is the path–ordered exponential

$$\boxed{\hat{\mathcal{P}}_\gamma := \mathcal{T} \exp\!\left(i \oint_\gamma \hat{\tau}_\mu \, dx^\mu \right),} \tag{16.251}$$

which is generically nontrivial when the chrono–geometric vorticity $\hat{\omega}_{\mu\nu} = \nabla_\mu \hat{\tau}_\nu - \nabla_\nu \hat{\tau}_\mu$ has nonzero expectation on a spanning surface (Stokes' theorem; cf. eq. (16.229)). The eigenphases of $\hat{\mathcal{P}}_\gamma$ are observable as topological phases in history–interference experiments defined by $\mathcal{I}_{\text{Kerr}}$ (eq. (16.245)), and are constrained by local covariance/renormalization in curved backgrounds [245].

Derivation of the Relational Holonomy Operator We determine the total transformation a modal state undergoes under *parallel transport* around a closed loop $\gamma \subset \mathcal{X}_q$, using the operator–valued temporal potential $\hat{\tau}_\mu$ as the connection on the modal Hilbert bundle (see Definition 16.13).

Step 1: Parallel transport as vanishing covariant derivative. Let $|\psi(x)\rangle$ be a modal state at $x \in \mathcal{X}_q$. The relational covariant derivative is

$$D_\mu := \partial_\mu - i\, \hat{\tau}_\mu(x),$$

so parallel transport along a path $s \mapsto \gamma(s)$ $(s \in [0,1],\ \dot{x}^\mu := dx^\mu/ds)$ is defined by

$$D_s|\psi\rangle \;=\; \left(\tfrac{d}{ds} - i\, \hat{\tau}_\mu(\gamma(s))\, \dot{x}^\mu \right) |\psi\rangle \;=\; 0, \tag{16.252}$$

with $|\psi(\gamma(0))\rangle$ given. Equivalently,

$$d|\psi\rangle \;=\; i\, \hat{\tau}_\mu(x)\, dx^\mu\, |\psi(x)\rangle, \tag{16.253}$$

which preserves the "direction" of $|\psi\rangle$ relative to the local modal frame. (The operator $\hat{\tau}_\mu$ is the temporal connection introduced in Definition 16.13; its curvature $\hat{\omega}_{\mu\nu} := \nabla_\mu \hat{\tau}_\nu - \nabla_\nu \hat{\tau}_\mu$ is given in eq. (16.229).)

Step 2: Infinitesimal transporter and unitarity. From (16.253), the infinitesimal transporter is

$$U(x + dx, x) \;=\; \mathbb{I} + i\,\hat{\tau}_\mu(x)\,dx^\mu + \mathcal{O}(dx^2). \tag{16.254}$$

With the convention $\hat{\tau}_\mu := -i\,\hat{U}^{-1}\partial_\mu \hat{U}$ (cf. Definition 16.13), $\hat{\tau}_\mu$ is Hermitian; hence $U(x+dx, x)$ is unitary to $\mathcal{O}(dx)$, and the finite transporter is unitary as well.

Step 3: Finite transporter and the holonomy of a closed loop. Composing (16.254) along γ and accounting for the general noncommutativity $[\hat{\tau}_\mu(x), \hat{\tau}_\nu(y)] \neq 0$, one obtains the *path–ordered exponential* (Magnus/Dyson series; cf. standard holonomy constructions in gauge theories):

$$\hat{\mathcal{P}}_\gamma \;:=\; \mathcal{T}\exp\!\Big(i \oint_\gamma \hat{\tau}_\mu\, dx^\mu\Big), \tag{16.255}$$

which is precisely the relational holonomy operator introduced in Definition 16.25 (see eq. (16.251)). For an *open* path $\gamma : x \to y$, the parallel transporter is $\hat{\mathcal{U}}[\gamma] = \mathcal{T}\exp(i\int_x^y \hat{\tau}_\mu\, dx^\mu)$ and satisfies (16.252).

Small–loop expansion and curvature (vorticity) link. For an infinitesimal plaquette spanning an oriented area element $d\sigma^{\mu\nu}$ one has

$$\hat{\mathcal{P}}_\gamma \;=\; \mathbb{I} \;+\; i\int_{\Sigma(\gamma)} \hat{\omega}_{\mu\nu}\, d\sigma^{\mu\nu} \;+\; \mathcal{O}(\text{area}^2), \tag{16.256}$$

so the eigenphases of $\hat{\mathcal{P}}_\gamma$ are controlled to leading order by the chrono–geometric vorticity $\hat{\omega}_{\mu\nu}$ (cf. eq. (16.229)). In the stationary Kerr phase this curvature sources the lapse and $t\phi$ profiles via the UVFE (Definition 16.6 and eq. (16.231)), and the resulting holonomy phases enter the history–interference observables defined by $\mathcal{I}_{\text{Kerr}}$ and $\hat{G}_{\text{ring}}$ (eqs. (16.245) and (16.250)).

Gauge (frame) covariance. Under a local modal frame change $\hat{V}(x)$ on the bundle, $\hat{\tau}_\mu \mapsto \hat{V}^{-1}\hat{\tau}_\mu \hat{V} + i\,\hat{V}^{-1}\partial_\mu \hat{V}$. Accordingly, $\hat{\mathcal{P}}_\gamma \mapsto \hat{V}^{-1}(x_0)\,\hat{\mathcal{P}}_\gamma\,\hat{V}(x_0)$ for loops based at x_0; the *conjugacy class* (eigenphases) of $\hat{\mathcal{P}}_\gamma$ is gauge–invariant and physically observable through the holosymmetric interference diagnostics above. (For background on holonomy and path–ordered exponentials in curved and gauge–theoretic settings, see [245].)

Holonomy as a Measure of Curvature If the modal bundle were flat (i.e. $\hat{\tau}_\mu \equiv 0$ or a pure gauge), then the loop transporter would be trivial, $\hat{\mathcal{P}}_\gamma = \mathbb{I}$, and a state would return to itself after traversing γ. Thus any nontrivial holonomy $\hat{\mathcal{P}}_\gamma = \mathcal{T}\exp\!\big(i \oint_\gamma \hat{\tau}_\mu\, dx^\mu\big)$ (Definition 16.25 and eq. (16.251)) *measures the curvature* of the temporal connection.

Local (small–loop) relation to curvature. Using the chrono–geometric vorticity $\hat{\omega}_{\mu\nu} := \nabla_\mu \hat{\tau}_\nu - \nabla_\nu \hat{\tau}_\mu$ (eq. (16.229)), the small–loop expansion reads

$$\boxed{\hat{\mathcal{P}}_\gamma \;=\; \mathbb{I} \;+\; i\int_{\Sigma(\gamma)} \hat{\omega}_{\mu\nu}\, d\sigma^{\mu\nu} \;+\; \mathcal{O}(\text{area}^2),} \tag{16.257}$$

in agreement with eq. (16.256). Hence, to leading order, the eigenphases of $\hat{\mathcal{P}}_\gamma$ are set by the vorticity flux through any spanning surface $\Sigma(\gamma)$.

Global (non-Abelian Stokes) form. For finite loops one may write a surface–ordered form (non-Abelian Stokes) schematically as

$$\hat{\mathcal{P}}_\gamma \;=\; \mathcal{P}_\Sigma \exp\!\left(i \!\int_{\Sigma(\gamma)} \hat{U}^{-1}(x_0 \to x)\, \hat{\omega}_{\mu\nu}(x)\, \hat{U}(x \to x_0)\, d\sigma^{\mu\nu} \right), \tag{16.258}$$

where $\hat{U}$ are open transporters along surface rays from the base point x_0 to x. In the Abelian (commutative) limit this reduces to $\hat{\mathcal{P}}_\gamma = \exp(i \int_\Sigma \hat{\omega}_{\mu\nu}\, d\sigma^{\mu\nu})$ [245].

Gauge–invariant observables (Wilson loop). While $\hat{\mathcal{P}}_\gamma$ depends on the choice of local modal frame, its *conjugacy class*—and thus its spectrum—is gauge–invariant. A convenient scalar diagnostic is the (normalized) Wilson loop,

$$\boxed{W[\gamma] \;:=\; \frac{1}{\mathcal{N}} \operatorname{Tr} \hat{\mathcal{P}}_\gamma,} \tag{16.259}$$

which is invariant under $\hat{\mathcal{P}}_\gamma \mapsto \hat{V}^{-1}(x_0)\hat{\mathcal{P}}_\gamma \hat{V}(x_0)$. Operationally, $W[\gamma]$ (or, equivalently, the eigenphases $e^{i\theta_j}$ of $\hat{\mathcal{P}}_\gamma$) appears as a *measurable phase shift* in holosymmetric interference experiments comparing histories that wind around the ring in different homotopy classes (eqs. (16.245) and (16.250)).

Topology and Kerr invariants. When the vorticity $\bar{\omega}_{\mu\nu} = \langle \Omega_\omega | \hat{\omega}_{\mu\nu} | \Omega_\omega \rangle$ is localized near the ring condensate $\mathfrak{B}_{\text{ring}}$ (Definition 16.10), the conjugacy class of $\hat{\mathcal{P}}_\gamma$ is stable under smooth deformations of γ that do not cross the support of $\bar{\omega}$; hence the eigenphases $e^{i\theta_j}$ define *topological invariants* of the Kerr coherence phase. Through the Unified Vacuum Field Equation (Definition 16.6) these curvature fluxes modulate the temporal profiles $A, B, \ldots, D$ (eq. (16.199)), shifting ergosurface/horizon data (eqs. (16.2), (16.4) and (16.230)) in a state–dependent but unitary manner (Section 16.3).

5. Causal Loops as History Transitions

Definition 16.26 (Causal loops as holosymmetric history transitions)**.** In classical GR, the analytic extension of Kerr admits closed timelike curves (CTCs) near the ring locus $\{r = 0, \theta = \pi/2\}$ [649, 245]. Within the IQR framework these are reinterpreted not as paths into one's own past but as *holosymmetric tunnelling* between distinct quantum history sectors of the Holo-State.

Let the global Holo-State decompose into orthogonal history branches (cf. eq. (16.248)):

$$|\Omega_\omega\rangle = \sum_{\alpha \in \mathbb{Z}} c_\alpha\, |\Psi_\alpha^{\text{Kerr}}\rangle, \qquad \Pi_\alpha := |\Psi_\alpha^{\text{Kerr}}\rangle\langle\Psi_\alpha^{\text{Kerr}}|.$$

For a loop γ_{CTC} that threads the ring condensate (Definition 16.10), define the *branch–resolved holonomy transitions* by

$$\boxed{\hat{T}_{\beta\alpha}[\gamma_{\text{CTC}}] \;:=\; \Pi_\beta\, \hat{\mathcal{P}}_{\gamma_{\text{CTC}}}\, \Pi_\alpha, \qquad \hat{\mathcal{P}}_\gamma := \mathcal{T} \exp\!\left(i \oint_\gamma \hat{\tau}_\mu\, dx^\mu \right) \;\text{(eq. (16.255)).}} \tag{16.260}$$

A *CTC history transition* is the case where an off–diagonal amplitude is nonzero, $\hat{T}_{\beta\alpha} \neq 0$ for $\beta \neq \alpha$. Equivalently, the classical CTC implements a partial isometry from the branch subspace $\mathcal{H}_\alpha := \Pi_\alpha \mathcal{H}$ to $\mathcal{H}_\beta := \Pi_\beta \mathcal{H}$:

$$\boxed{\gamma_{\mathrm{CTC}} : \ \mathcal{H}_\alpha \xrightarrow{\hat{T}_{\beta\alpha}} \mathcal{H}_\beta, \qquad \beta \neq \alpha,} \tag{16.261}$$

mediated by the ring's holosymmetric structure (Definition 16.23 and eq. (16.245)).

The logical path to the reinterpretation

Step 1: The classical paradox. The extended Kerr spacetime admits CTCs, threatening causal paradoxes in a single classical manifold [649, 245]. Any quantum–consistent completion must avoid self–contradictory dynamics.

Step 2: Superposed histories. In IQR, the multi–sheeted Kerr structure is a superposition of orthogonal quantum histories (eq. (16.248)), not a single bizarre spacetime. The Fourfold Inter–Modal Correlator $\hat{G}_\Lambda$ organizes cross–branch coherence (Definition 16.21 and eq. (16.250)).

Step 3: The ring as portal. The ring "singularity'' is a pre–geometric entanglement condensate (Definition 16.10) and a holosymmetric interferometric node (Definition 16.23) that can mediate branch–to–branch interference via the modal holonomy of $\hat{\tau}_\mu$ (Definition 16.25 and eq. (16.251)).

Step 4: Synthesis. A classical CTC that threads the ring corresponds, in the emergent description, to a *transition path* whose holonomy has nontrivial off–diagonal branch components:

$$\hat{T}_{\beta\alpha}[\gamma_{\mathrm{CTC}}] = \Pi_\beta \, \hat{\mathcal{P}}_{\gamma_{\mathrm{CTC}}} \, \Pi_\alpha \neq 0, \quad (\beta \neq \alpha),$$

precisely when ring–mediated interference is present (nonzero $\hat{G}_\Lambda^{(\alpha,\beta)}$; cf. eqs. (16.244) and (16.250)). Thus, a traveller along γ_{CTC} does *not* return to their own past; they emerge in a *distinct* history sector.

Resolution of the paradox

(i) Intra–branch causality is preserved. Dynamics within each $\mathcal{H}_\alpha$ remain causal; no operation along γ_{CTC} produces a contradiction in the originating branch because the endpoint lies in $\mathcal{H}_\beta$, $\beta \neq \alpha$.

(ii) Local microcausality is intact. The exterior observable net satisfies locality and KMS properties as before; invariance under Killing flows and Hadamard regularity are unaffected [245, 230, 241]. The CTC "transition'' is a global, ring–localized feature of the Holo-State, not a channel for acausal signalling within a branch.

The Kerr black hole as a quantum portal of relational histories The Kerr interior is a stable, holosymmetric coherence phase: a topological quantum portal where the Fourfold correlator $\hat{G}_\Lambda$ governs interference among history branches, and the modal holonomy of $\hat{\tau}_\mu$ provides the transition amplitudes (eqs. (16.255) and (16.257)). Through the Unified Vacuum Field Equation (Definition 16.6) these coherences backreact on the temporal sector $A, \ldots, D$ (eq. (16.199)), shifting ergosurface/horizon data in a unitary, state–dependent manner (eqs. (16.2), (16.4) and (16.230) and Section 16.3). Consequently, classical CTCs are recontextualized as *holosymmetric tunnelling channels* between histories, not as mechanisms for time–travel paradoxes—aligning black–hole interior physics with global unitarity and algebraic consistency.

16.4.8 The Holosymmetric Kerr Attractor as a Topological Quantum Phase

Within the IQR edifice, the Kerr configuration is *not* a merely classical solution realized on a fixed manifold, but a globally stable *stationary coherence phase* (Definition 16.21) of the Quantum Holo-State $|\mathring{\boldsymbol{\Psi}}\rangle$. Its identity is fixed by (i) holonomy data of the modal connection, (ii) the topology of inter-modal entanglement, and (iii) holosymmetric dynamical coherence under the relational evolution. In this sense, the "Kerr Attractor" is a *topological quantum phase*: it is characterized by quantized, deformation-invariant features of the relational operator algebra rather than by local curvature alone [7, 146, 437, 650, 651].

1. Topological phase invariants from modal holonomy. Let $\gamma : S^1 \to \mathcal{X}_q$ be any loop encircling the Kerr ring (no graph surgery across $\mathcal{H}_+$; cf. the sector-preserving clauses around Definition 16.44). With $\hat{\tau}_\mu$ the modal connection one-form induced by the unified inter-modal correlators (Definition 6.4), define the Wilson–holonomy operator

$$\hat{\mathcal{P}}_\gamma \;:=\; \mathcal{T}\exp\Big(i \oint_\gamma \hat{\tau}_\mu \, dx^\mu\Big), \tag{16.262}$$

acting on the modal fiber $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$ (Section 4.7.2). Its spectrum $\mathrm{spec}(\hat{\mathcal{P}}_\gamma) = \{e^{i\theta_j}\}$ (equivalently, the Wilson loop $W(\gamma) = \mathrm{Tr}\,\hat{\mathcal{P}}_\gamma$) is invariant under smooth deformations of γ that preserve its homotopy class in $\pi_1(\mathcal{X}_q \setminus \mathfrak{B}_{\mathrm{ring}})$. Thus the *topological phase angles* $\{\theta_j\}$ provide gauge-invariant[4] labels of the Kerr phase sector [375, 225, 389, 24].

Proposition 16.27 (Holonomy rigidity)**.** *Let γ_s be a smooth family of loops encircling the ring, with no crossings of singular support and no topology change of the PQS network. Then the conjugacy class of $\hat{\mathcal{P}}_{\gamma_s}$ is constant in s. Hence $\{\theta_j\}$ are homotopy invariants of the Kerr phase sector.*

2. Holosymmetric stability as a dynamical fixed point. The Kerr phase is stationary in the *holosymmetric* sense: under the global relational evolution

$$\hat{\mathcal{U}}_{\mathrm{holo}}(\delta t) \;=\; \exp\Big(-i\,\delta t\,\hat{H}_{\mathrm{rel}}\Big)\,,$$

[4] Up to conjugacy; characters such as $\mathrm{Tr}(\hat{\mathcal{P}}_\gamma^{\,k})$ are strictly gauge-invariant.

with $\hat{H}_{\mathrm{rel}}$ determined by inter-PQS correlators (Definition 4.9), one has

$$\hat{\mathcal{U}}_{\mathrm{holo}}(\delta t)\,|\overset{\circ}{\boldsymbol{\Psi}}_{\mathrm{Kerr}}\rangle \;=\; e^{i\theta}\,|\overset{\circ}{\boldsymbol{\Psi}}_{\mathrm{Kerr}}\rangle, \tag{16.263}$$

i.e., the Kerr Attractor is an eigenstate of the holosymmetric flow (Definition 16.21). This fixed-point structure dovetails with the algebraic/KMS characterization of equilibrium dynamics [30, 230, 652] and underlies the constancy of the effective surface gravity on $\mathcal{H}_+$ proved elsewhere (cf. eqs. (16.2) and (16.193)).

3. Entanglement topology and the emergent global phase. Let $\hat{G}_F(\mathfrak{B}_{x_i}, \mathfrak{B}_{x_j})$ be the two-point (fibered) correlator extracted from the unified inter-modal kernel (Definition 6.4). Fix a coherence threshold $\epsilon > 0$ and define the entanglement graph

$$\mathcal{G}_{\mathrm{Kerr}} \;:=\; \Big\{(x_i, x_j) \in \mathcal{X}_q \times \mathcal{X}_q \;\Big|\; \langle\overset{\circ}{\boldsymbol{\Psi}}|\,\hat{G}_F(\mathfrak{B}_{x_i}, \mathfrak{B}_{x_j})\,|\overset{\circ}{\boldsymbol{\Psi}}\rangle > \epsilon\Big\}, \tag{16.264}$$

whose first homology captures *entanglement holonomies*:

$$H_1(\mathcal{G}_{\mathrm{Kerr}}) \;\cong\; \mathbb{Z}, \tag{16.265}$$

reflecting the single non-contractible loop induced by the ring sector (no-surgery hypothesis). The resulting global features are encoded at the level of entanglement structure, paralleling the logic of topological order and topological entanglement entropy [650, 651, 437]; see also holographic diagnostics [653].

4. Modal quantization and discretization of Kerr parameters. Let the *local modal Hamiltonians* $\{\hat{H}_x\}_{x\in\mathcal{X}_q^{(\mathrm{Kerr})}}$ and emergent fiber operators $\hat{x}^\mu(x), \hat{p}^\mu(x)$ be those defined on the local Hilbert fibers $\mathcal{H}_{\mathrm{mod}}^{(\mathfrak{B}_x)}$. Define collective observables

$$\hat{M} \;:=\; \sum_{x\in\mathcal{X}_q^{(\mathrm{Kerr})}} \hat{H}_x, \qquad \hat{J}^{\mu\nu} \;:=\; \sum_{x\in\mathcal{X}_q^{(\mathrm{Kerr})}} \Big(\hat{x}^\mu(x)\hat{p}^\nu(x) - \hat{x}^\nu(x)\hat{p}^\mu(x)\Big), \tag{16.266}$$

and let $\{|\Psi_n\rangle\}_{n\in\mathbb{Z}_+}$ be orthonormal *topological entanglement sectors* (winding-number superselection sectors of $\mathcal{G}_{\mathrm{Kerr}}$). Then

$$M_n \;:=\; \langle\Psi_n|\hat{M}|\Psi_n\rangle, \qquad a_n \;:=\; \frac{1}{M_n}\,\langle\Psi_n|\hat{J}^{r\phi}|\Psi_n\rangle, \tag{16.267}$$

are sector-indexed, discrete macroscopic parameters. Under sector-preserving dynamics (no graph surgery), (M_n, a_n) are piecewise constant as functions of slow controls and jump only across topological transitions, matching the general phenomenology of topological phases [437]. Compatibility with the first-law/area-law machinery is ensured by eqs. (16.194) and (16.195).

Key concepts in the Kerr Attractor phase.

(i) Quantized expectation values. The macroscopic M and a are *outputs* of relational topology: sector-conditioned spectral expectations $\{M_n, a_n\}$, not fundamental inputs.

(ii) Mass operator $\hat{M}$. Totals emergent energy across the Kerr PQS support via local $\hat{H}_x$ (cf. relational energy bookkeeping in Definition 6.4).

(iii) Angular momentum $\hat{J}^{\mu\nu}$. Encodes relational rotation via antisymmetric position–momentum correlators on fibers, sensitive to global torsion/coherence.

(iv) Topological sectors $|\Psi_n\rangle$. Stable coherence phases indexed by winding in $H_1(\mathcal{G}_{\text{Kerr}})$; each sector is a distinct geometric phase of the attractor.

(v) Winding and spin quantization. The discretization of a follows from the quantized winding number in $H_1(\mathcal{G}_{\text{Kerr}})$ and the holonomy spectrum of $\hat{\mathcal{P}}_\gamma$ (eq. (16.262)).

5. Global identification: the Kerr Attractor as a holosymmetric quantum phase sector. Collecting the invariants, the Kerr phase in sector n is summarized by

$$\mathcal{K}_n \;:=\; \Big(H_1(\mathcal{G}_{\text{Kerr}}),\ \{\theta_j\},\ |\Psi_n\rangle\Big), \tag{16.268}$$

where $\{\theta_j\}$ are holonomy eigenphases (modulo conjugacy) and $|\Psi_n\rangle$ is the topological entanglement sector. Each $\mathcal{K}_n$ is a topologically protected, holosymmetric fixed point (eq. (16.263)), stable under small deformations that preserve the homotopy class and coherence threshold.

The Kerr black hole as a quantum fixed point of modal topology. In this reading, "Kerr" names a *quantum* fixed point of modal topology: horizons, ergospheres, and singularities are the classical shadows of deeper entanglement/holonomy data. Einstein's equations emerge as the low-coherence/long-wavelength limit of the chronogeometric dynamics (Definition 11.106); the Kerr Attractor is the more primitive structure whose discretized sectors $\{\mathcal{K}_n\}$ encode distinct, stable modes of existence of rotating black holes [7, 146, 30, 230].

16.4.9 Quantum Ergodynamics and the Information Flow

In classical general relativity, the Kerr ergosphere is the region outside the outer event horizon where the stationary Killing field t^a becomes spacelike ($g_{ab}t^a t^b > 0$), enforcing corotation of all timelike trajectories via frame dragging [7, 318, 146]. This enables the Penrose process: a particle entering the ergosphere can split so that one fragment with *negative Killing energy* ($E = -p_a t^a < 0$ as measured at infinity) falls into the black hole while the other escapes with E_∞ larger than the initial energy, thereby extracting rotational energy [654, 655, 656]. In the IQR framework, these phenomena are reinterpreted as manifestations of *quantum ergodynamics*: the dynamical evolution of *modal coherence* across domains where the emergent temporal flow aligns with the horizon generator χ^a and becomes effectively spacelike for t^a, prompting a reconfiguration of relational operators that permits coherent information and energy transfer (cf. Section 16.4.10 and eqs. (16.2) and (16.193)).

Definition 16.28 (Quantum ergodynamics and the Penrose process as entanglement reconfiguration)**.** Let $\hat{\rho}_{\text{in}}$ denote the reduced state (cf. Definition 16.36) of an incoming PQS-supported configuration prior to ergospheric interaction. Model the ergosphere as a boundary-driven interaction with vacuum modes $|\Omega\rangle$ prepared in a KMS state at the effective Hawking temperature T_{eff} associated with $\mathcal{H}_+$ (eq. (16.193)), and with *rotational chemical potential* given

by the horizon angular velocity Ω_H (see the Kerr-sector definitions around eq. (16.203)). The *ergospheric interaction* is described by a Stinespring dilation

$$\hat{U}_{\rm erg}: \ \mathcal{H}_{\rm in}\otimes\mathcal{H}_{\Omega} \to \mathcal{H}_{\rm out}\otimes\mathcal{H}_{\rm neg}, \qquad \text{unitary}, \tag{16.269}$$

and induces a completely positive, trace-preserving (CPTP) channel $\mathcal{E}_{\rm erg}$ on the escaping sector via partial trace over the *negative-Killing-energy* modes (those with $-p_a t^a < 0$ relative to t^a inside the ergosphere [7, Sec. 12]):

$$\boxed{\hat{\rho}_{\rm out} \ = \ \mathcal{E}_{\rm erg}(\hat{\rho}_{\rm in}) \ := \ \mathrm{Tr}_{\rm neg}\Big[\hat{U}_{\rm erg}\Big(\hat{\rho}_{\rm in}\otimes|\Omega\rangle\langle\Omega|\Big)\hat{U}_{\rm erg}^{\dagger}\Big]\,.} \tag{16.270}$$

Here $\mathrm{Tr}_{\rm neg}$ denotes the trace over those degrees of freedom that cross $\mathcal{H}_+$ with negative Killing energy, ensuring complete positivity and trace preservation by the usual dilation theorem [352, 354, 441, 287]. The state $\hat{\rho}_{\rm out}$ describes the escaping configuration after an *entanglement reconfiguration* with the ergospheric modes.

Remark 16.29 (Negative energy and superradiant channels)**.** Inside the ergosphere the sign of $E = -p_a t^a$ may flip even for future-directed timelike p^a, making $E < 0$ physically admissible [7, 657]. In wave language, this corresponds to *superradiant* channels satisfying $\omega < m\,\Omega_H$, which realize energy extraction at infinity while decreasing the black hole's mass and angular momentum [658, 143, 142]. The unitary $\hat{U}_{\rm erg}$ thus couples system modes to these ergospheric channels before tracing out the infalling, negative-energy sector.

Proposition 16.30 (Energy extraction as modal purification: a generalized free-energy law)**.** *Let $S(\hat{\rho}) = -\,\mathrm{Tr}(\hat{\rho}\ln\hat{\rho})$ be the von Neumann entropy and let*

$$\mathcal{F}_{\beta_H,\Omega_H}(\hat{\rho}) := \mathrm{Tr}\Big[\hat{\rho}(\hat{H} - \Omega_H\hat{J})\Big] - T_{\rm eff}\,S(\hat{\rho}), \qquad \beta_H := 1/T_{\rm eff}, \tag{16.271}$$

be the generalized free energy of the escaping subsystem relative to the KMS pair (β_H,Ω_H) on $\mathcal{H}_+$ (eq. (16.193); cf. [230, 30]). For the CPTP map $\mathcal{E}_{\rm erg}$ in eq. (16.270) that is Gibbs-preserving with respect to the KMS reference, the data-processing inequality for relative entropy implies

$$\Delta\mathcal{F}_{\beta_H,\Omega_H}(\text{out}) \ = \ \Delta E_{\rm out} - \Omega_H\,\Delta J_{\rm out} - T_{\rm eff}\,\Delta S_{\rm out} \ \le \ 0, \tag{16.272}$$

whence the escaping energy gain obeys the bound

$$\boxed{\Delta E_{\rm out} \ \le \ \Omega_H\,\Delta J_{\rm out} \ - \ T_{\rm eff}\,\Delta S_{\rm out}.} \tag{16.273}$$

In particular, for fixed angular-momentum transfer $\Delta J_{\rm out}$ (or in the quasi-reversible limit that saturates eq. (16.272)), a decrease in entanglement entropy ($\Delta S_{\rm out} < 0$) corresponds to an increased escaping energy, i.e. a modal purification of the outgoing subsystem.[5]

[5] Equality in eq. (16.273) holds in the quasi-static, coherence-preserving limit; more generally, irreversibility introduces an extra nonnegative entropy-production term [624, 441].

Proof of Proposition 16.30. Let $H' := \hat{H} - \Omega_H \hat{J}$ denote the co-rotating generator and

$$\hat{\tau} := \frac{e^{-\beta_H H'}}{Z}, \qquad \beta_H := 1/T_{\text{eff}}, \quad Z := \operatorname{Tr}\left[e^{-\beta_H H'}\right],$$

be the KMS/Gibbs reference state associated with the horizon pair (β_H, Ω_H) (cf. eq. (16.193) and the Kerr-sector discussion around Ω_H). For any state $\hat{\rho}$, the quantum relative entropy w.r.t. $\hat{\tau}$ satisfies

$$\begin{aligned} D(\hat{\rho}\|\hat{\tau}) &= \operatorname{Tr}\left[\hat{\rho}(\ln\hat{\rho} - \ln\hat{\tau})\right] \\ &= \beta_H(\operatorname{Tr}[\hat{\rho}H'] - T_{\text{eff}}\, S(\hat{\rho})) + \ln Z \\ &= \beta_H(\mathcal{F}_{\beta_H,\Omega_H}(\hat{\rho}) - \mathcal{F}_{\beta_H,\Omega_H}(\hat{\tau}))\,, \end{aligned} \tag{16.274}$$

where $S(\hat{\rho}) = -\operatorname{Tr}(\hat{\rho}\ln\hat{\rho})$ and $\mathcal{F}_{\beta_H,\Omega_H}(\hat{\rho}) := \operatorname{Tr}[\hat{\rho}H'] - T_{\text{eff}}S(\hat{\rho})$ is the generalized free energy.

Let $\mathcal{E}_{\text{erg}}$ be the CPTP map defined in eq. (16.270) and assume it is Gibbs-preserving with respect to $\hat{\tau}$, i.e. $\mathcal{E}_{\text{erg}}(\hat{\tau}) = \hat{\tau}$ (see Lemma 16.31). By monotonicity of quantum relative entropy under CPTP maps (data processing) [617, 357, 50, 441],

$$D(\mathcal{E}_{\text{erg}}(\hat{\rho}_{\text{in}})\,\|\,\hat{\tau}) \;\le\; D(\hat{\rho}_{\text{in}}\,\|\,\hat{\tau})\,.$$

Using (16.274) with $\hat{\rho}_{\text{out}} = \mathcal{E}_{\text{erg}}(\hat{\rho}_{\text{in}})$ yields

$$\mathcal{F}_{\beta_H,\Omega_H}(\hat{\rho}_{\text{out}}) \;\le\; \mathcal{F}_{\beta_H,\Omega_H}(\hat{\rho}_{\text{in}}),$$

i.e. $\Delta\mathcal{F}_{\beta_H,\Omega_H}(\text{out}) \le 0$, which is eq. (16.272). Expanding $\mathcal{F}_{\beta_H,\Omega_H}$ gives

$$\Delta E_{\text{out}} - \Omega_H\,\Delta J_{\text{out}} - T_{\text{eff}}\,\Delta S_{\text{out}} \;\le\; 0,$$

equivalently,

$$\Delta E_{\text{out}} \;\le\; \Omega_H\,\Delta J_{\text{out}} - T_{\text{eff}}\,\Delta S_{\text{out}},$$

which is eq. (16.273). In the quasi-static (reversible) limit, the inequality is saturated; in general, irreversibility contributes a nonnegative entropy-production term [624, 441]. □

Lemma 16.31 (Gibbs/KMS preservation for ergospheric channels)**.** *Let the joint unitary $\hat{U}_{\text{erg}}$ in eq. (16.270) act on system⊗environment with environment prepared in the KMS state $\hat{\tau}_{\text{env}} \propto e^{-\beta_H(H'_{\text{env}})}$ and suppose the total co-rotating generator is conserved, $[\hat{U}_{\text{erg}},\, H'_{\text{tot}}] = 0$, with $H'_{\text{tot}} = H' \otimes \mathbb{I} + \mathbb{I} \otimes H'_{\text{env}}$. Then the reduced channel $\mathcal{E}_{\text{erg}}(\,\cdot\,) = \operatorname{Tr}_{\text{neg}}\left[\hat{U}_{\text{erg}}(\,\cdot\,\otimes\hat{\tau}_{\text{env}})\hat{U}^\dagger_{\text{erg}}\right]$ satisfies $\mathcal{E}_{\text{erg}}(\hat{\tau}) = \hat{\tau}$ with $\hat{\tau} \propto e^{-\beta_H H'}$.*

Proof. The assumption $[\hat{U}_{\text{erg}}, H'_{\text{tot}}] = 0$ implies $\hat{U}_{\text{erg}}(\hat{\tau}\otimes\hat{\tau}_{\text{env}})\hat{U}^\dagger_{\text{erg}} = \hat{\tau}\otimes\hat{\tau}_{\text{env}}$. Tracing over the environment (including the "negative-energy" sector) yields $\mathcal{E}_{\text{erg}}(\hat{\tau}) = \hat{\tau}$. This is the standard characterization of Gibbs/KMS-preserving ("thermal") operations in the co-rotating ensemble [30, 230, 441]. □

Interpretation. Equation eq. (16.273) reframes the Penrose mechanism as a constrained transformation of coherence: the ergosphere functions as a *modal centrifuge* that converts rotational free energy (via the $\Omega_H\,\Delta J_{\text{out}}$ term) and decreased relational entropy into higher escaping energy. This dovetails with the entropic-dual account of ergospheres developed in Section 16.4.10 and remains consistent with the first-law identities established in the Kerr sector (eqs. (16.194) and (16.195)).

4. Horizon Crossing and Temporal Decoherence

Definition 16.32 (Horizon crossing and temporal decoherence)**.** As a configuration of Primordial Quantum Singularities (PQSs) crosses the outer event horizon $\mathcal{H}_+$, the relational coherence between exterior and interior modal structures becomes *operationally inaccessible* to an external observer.

Let the relational configuration be partitioned as $\mathcal{X}_q = \mathcal{X}_{\text{ext}} \cup \mathcal{X}_{\text{int}}$ with disjoint supports. Denote by $\hat{G}_\Lambda$ the Fourfold/Unified Inter-Modal Correlator introduced earlier (Definition 6.4). The **cross-horizon correlator** is the operator-valued kernel

$$\hat{C}_{\mathcal{H}}(x, x') \;:=\; \hat{G}_\Lambda(\mathfrak{B}_x, \mathfrak{B}_{x'})\,, \qquad x \in \mathcal{X}_{\text{ext}},\; x' \in \mathcal{X}_{\text{int}}, \tag{16.275}$$

which probes entanglement between exterior and interior fibers.

Let $\hat{\rho}$ be the global state on the horizon neighborhood and write the exterior reduced state $\hat{\rho}_{\text{ext}} := \mathrm{Tr}_{\text{int}}\,\hat{\rho}$ (Stinespring/Kraus reduction; cf. [352, 354, 441]). Define the exterior *restriction* (a CPTP conditional expectation) by

$$\mathcal{R}_{\text{ext}}(X) \;:=\; \mathrm{Tr}_{\text{int}}\Big[(\mathbb{I}_{\text{ext}} \otimes \hat{\rho}_{\text{int}}^{1/2})\, X\, (\mathbb{I}_{\text{ext}} \otimes \hat{\rho}_{\text{int}}^{1/2})\Big], \qquad \hat{\rho}_{\text{int}} := \mathrm{Tr}_{\text{ext}}\,\hat{\rho}, \tag{16.276}$$

and the *temporal-decoherence functional* (trace norm on the exterior)

$$\mathscr{D}_{\mathcal{H}}(x) \;:=\; \Big\| \mathcal{R}_{\text{ext}}\big[\hat{C}_{\mathcal{H}}(x, \cdot)\big] \Big\|_{1,\text{ext}}. \tag{16.277}$$

Then, in a KMS neighborhood of the horizon with effective temperature T_{eff} and generator χ^a (eqs. (16.2) and (16.193)), the process of **temporal decoherence** is characterized by the exterior suppression of cross-horizon correlations:

$$\boxed{\lim_{x \to \mathcal{H}_+^-} \mathscr{D}_{\mathcal{H}}(x) \;=\; 0,} \tag{16.278}$$

i.e. the exterior-accessible imprint of interior entanglement vanishes as the support approaches $\mathcal{H}_+$ from outside.[6] Operationally, information is *not destroyed*: it is encoded in the interior relational structure and becomes causally disconnected for exterior observers (compare the entropic-dual discussion in Section 16.4.10 and the first-law identities eqs. (16.194) and (16.195)).

Remark 16.33 (Clustering bound and horizon scale)**.** In a stationary KMS state at T_{eff}, exterior–interior connected correlators satisfy a clustering inequality of the form

$$\Big|\langle \hat{O}_{\text{ext}} \hat{O}_{\text{int}} \rangle - \langle \hat{O}_{\text{ext}} \rangle \langle \hat{O}_{\text{int}} \rangle\Big| \;\lesssim\; C\, e^{-\ell(x,x')/\xi_H}, \qquad \xi_H \sim \frac{1}{2\pi T_{\text{eff}}}\,,$$

for suitable local operators and proper distance $\ell(x, x')$ across $\mathcal{H}_+$ [30, 230].

[6] In algebraic terms, $\mathcal{R}_{\text{ext}}$ realizes a Gibbs/KMS-preserving conditional expectation onto the exterior net; clustering/mixing for KMS states ensures suppression of cross correlations [30, 230, Chs. 5–6].

CP contraction note. Here "CP contraction" refers to the fact that the exterior restriction $\mathcal{R}_{\text{ext}}$ is a completely positive, trace-preserving (CPTP) map (a quantum channel), and such maps are *contractive in the trace norm*: for any traceless Hermitian X,

$$\left\| \mathcal{R}_{\text{ext}}(X) \right\|_1 \;\le\; \|X\|_1.$$

Equivalently, the trace distance between states never increases under CPTP maps (data processing) [287, 441, Sec. 9.2]. Therefore, applying $\mathcal{R}_{\text{ext}}$ cannot amplify the (already exponentially suppressed) cross-horizon contribution, and the trace-norm decay asserted in eq. (16.278) follows from KMS clustering together with this contractivity.

5. The Ergosphere as a Quantum Dynamical Interface

Definition 16.34 (The ergosphere as a quantum dynamical interface)**.** We reinterpret the ergosphere as a *dynamical quantum interface*: a coherent layer in which energy, coherence, and information are *repartitioned* via the open-system coupling between exterior modes and near-horizon (ergospheric) degrees of freedom (cf. Definition 16.28 and Section 16.4.9).

Let $\hat{\tau}_\mu$ denote the (operator-valued) temporal one-form introduced by the holonomy connection (eq. (16.262)), and let $\bar{\Sigma}_{\mu\nu}$ be the chronogeometric metric (Definition 11.106). Define the **ergodynamic stress tensor**

$$\hat{\mathcal{T}}^{\mu\nu}_{\text{erg}} \;:=\; \hat{\tau}^{(\mu}\hat{\tau}^{\nu)} - \frac{1}{2}\,\bar{\Sigma}^{\mu\nu}\,\hat{\tau}_\lambda\hat{\tau}^\lambda, \tag{16.279}$$

and, for any stationary Killing field v^ν (e.g. t^ν at infinity or the horizon generator χ^ν), the associated **ergodynamic energy current**

$$\hat{\mathcal{J}}^\mu_{(v)} \;:=\; -\hat{\mathcal{T}}^{\mu\nu}_{\text{erg}}\,v_\nu. \tag{16.280}$$

If $\nabla_\mu \hat{\mathcal{T}}^{\mu\nu}_{\text{erg}} = \hat{\mathcal{Q}}^\nu_{\text{open}}$ encodes the open-system exchange with ergospheric modes (CP interaction of Definition 16.28) and v^ν is Killing ($\nabla_{(\mu}v_{\nu)} = 0$), then the local balance law reads

$$\nabla_\mu \hat{\mathcal{J}}^\mu_{(v)} \;=\; -\,v_\nu\,\hat{\mathcal{Q}}^\nu_{\text{open}}, \tag{16.281}$$

so that, for $v^\nu = t^\nu$, the divergence of $\hat{\mathcal{J}}^\mu_{(t)}$ tracks the net extraction or injection of (Killing) energy density across the interface (compare Wald's Noether/Killing-current construction [7, Sec. 11]; see also eq. (16.194)).

Remark 16.35 (Interface fluxes and co-rotating ensemble)**.** Contracting eq. (16.280) with the horizon generator $\chi^\nu = t^\nu + \Omega_H\phi^\nu$ and taking expectation values in a KMS state at T_{eff} (eqs. (16.2) and (16.193)) yields a co-rotating energy current $\langle\hat{\mathcal{J}}^\mu_{(\chi)}\rangle$ whose flux through a timelike worldtube enclosing the ergoregion equals the change in the generalized free energy, cf. eqs. (16.272) and (16.273). This implements the "modal centrifuge" intuition: the interface converts rotational free energy ($\Omega_H\,\Delta J$) and entropy decrease ($-\Delta S$) into escaping energy.

Proof of Proposition 16.30. We refine the proof using the co-rotating KMS structure and data processing (cf. eq. (16.274) and eq. (16.272)). Let $H' = \hat{H} - \Omega_H\hat{J}$ and $\hat{\tau} \propto e^{-\beta_H H'}$

with $\beta_H = 1/T_{\text{eff}}$ (eq. (16.193)). The ergospheric interaction defines a CPTP channel $\mathcal{E}_{\text{erg}}$ by Stinespring dilation and partial trace over the negative-Killing-energy sector (Definition 16.28; [352, 354, 441]). Under the mild assumption that $\mathcal{E}_{\text{erg}}$ is Gibbs/KMS-preserving for $\hat{\tau}$ (see Lemma 16.31), monotonicity of relative entropy [617, 357, 50] gives

$$\mathcal{F}_{\beta_H,\Omega_H}(\hat{\rho}_{\text{out}}) \;\le\; \mathcal{F}_{\beta_H,\Omega_H}(\hat{\rho}_{\text{in}}),$$

with $\mathcal{F}_{\beta_H,\Omega_H}(\hat{\rho}) = \text{Tr}[\hat{\rho}H'] - T_{\text{eff}}S(\hat{\rho})$ (eq. (16.274)). Expanding,

$$\Delta E_{\text{out}} - \Omega_H\,\Delta J_{\text{out}} - T_{\text{eff}}\,\Delta S_{\text{out}} \;\le\; 0,$$

which is eq. (16.273). In the quasi-static (reversible) limit where the bound is saturated (no extra entropy production [624, 441]), the change in escaping energy is

$$\Delta E_{\text{out}} \;=\; \Omega_H\,\Delta J_{\text{out}} - T_{\text{eff}}\,\Delta S_{\text{out}}.$$

Hence, at fixed ΔJ_{out} (or when ΔJ_{out} is determined by superradiant selection $\omega < m\Omega_H$ [658, 142, 143]), an entropy decrease $\Delta S_{\text{out}} < 0$ entails a larger ΔE_{out}—the asserted *modal purification* relation. This completes the proof. □

Quantum Ergodynamics as the Modal Physics of Rotating Horizons In light of Proposition 16.30, the Kerr ergosphere is transfigured from a geometric curiosity into a coherent *quantum–ergodynamic engine*. As formalized in Definition 16.34, it functions as a dynamical interface where rotational free energy in the co-rotating KMS background (eqs. (16.2) and (16.193)) is systematically transduced into *informational purity* of escaping modal configurations. The governing principle is the energy–entropy balance

$$\Delta E_{\text{out}} \;\le\; \Omega_H\,\Delta J_{\text{out}} - T_{\text{eff}}\,\Delta S_{\text{out}},$$

the saturated, quasi-reversible form of which reads $\Delta E_{\text{out}} = \Omega_H\,\Delta J_{\text{out}} - T_{\text{eff}}\,\Delta S_{\text{out}}$ (eq. (16.273)). Thus, energy extraction is the thermodynamic dual of entropic purification ($\Delta E_{\text{out}} \propto -\Delta S_{\text{out}}$), with the flux of relational energy and coherence tracked by the *ergodynamic current* $\hat{\mathcal{J}}^{\mu}_{(v)}$ in eq. (16.280) and its balance law eq. (16.281).

Within this language, *quantum ergodynamics* integrates the treatise's core structures. First, *temporal vorticity*—the frame-drag–induced twisting of the temporal field generated by $\chi^a = t^a + \Omega_H\phi^a$—drives the co-rotating energetics and undergirds superradiant channels ($\omega < m\,\Omega_H$) that feed the engine [656, 658, 143, 142, 7]. Second, *operator entanglement* between exterior and interior sectors, encoded by the cross-horizon correlator eq. (16.275), is operationally suppressed near the horizon via KMS clustering and CP contractivity (eq. (16.278); cf. [30, 230, 287, 441]). Third, the *information-theoretic structure*—the monotonicity of relative entropy and the associated free-energy inequalities—provides the arrow of irreversibility and the precise conversion rule between rotation, coherence, and energy [617, 357, 50, 441].

At the spinning boundary of classical causality, these ingredients reveal a single dialectic: thermodynamics (first-law and area-law specializations; eqs. (16.194) and (16.195)), gravity (chronogeometric dynamics; Definition 11.106), and quantum information (KMS equilibrium and data processing) are not merely compatible—they are different faces of *quantum ergodynamics*, the modal physics of rotating horizons.

16.4.10 The Entropic Dual of the Kerr Ergosphere and the Emergence of Horizon Temperature

Building on Section 16.4.9 and Definition 16.34, the Kerr ergosphere is a *dynamically active* domain for modal reconfiguration. Its *entropic dual* is the holographic imprint of this reconfiguration on the structure of *modal entropy*. In this view, the *horizon temperature* is not merely a geometric invariant but the KMS parameter that statistically encodes decoherence-induced entanglement across $\mathcal{H}_+$, i.e. a thermodynamic shadow of quantum relational topology (cf. eqs. (16.2), (16.193) and (16.278); see [230, 30, 7, 653]).

1. Relational entropy and modal uncertainty

Definition 16.36 (Relational entropy and modal uncertainty)**.** Entropy is here a *relational* quantity: it measures information-theoretic indistinguishability of a subsystem from its complement due to entanglement within the Quantum Holo-State $|\mathring{\mathbf{\Psi}}\rangle$.

Let $\mathcal{R} \subset \mathcal{X}_q$ be any subsystem (a union of modal fibers). Its reduced state is obtained by tracing out the complement:

$$\boxed{\hat{\rho}_{\mathcal{R}} \;:=\; \mathrm{Tr}_{\mathcal{X}_q \setminus \mathcal{R}}\Big(|\mathring{\mathbf{\Psi}}\rangle\langle\mathring{\mathbf{\Psi}}| \Big).} \tag{16.282}$$

The **relational entropy** of $\mathcal{R}$ is the von Neumann entropy of $\hat{\rho}_{\mathcal{R}}$:

$$\boxed{S(\mathcal{R}) \;:=\; -\,\mathrm{Tr}_{\mathcal{R}}\Big(\hat{\rho}_{\mathcal{R}} \ln \hat{\rho}_{\mathcal{R}}\Big).} \tag{16.283}$$

This entropy quantifies *modal uncertainty* inside $\mathcal{R}$ caused by its correlations with $\mathcal{X}_q \setminus \mathcal{R}$. In particular, $S(\mathcal{R}) = 0$ iff $\hat{\rho}_{\mathcal{R}}$ is pure (no exterior entanglement); for a global pure state one has $S(\mathcal{R}) = S(\mathcal{X}_q \setminus \mathcal{R})$. Moreover, $S(\mathcal{R})$ cannot increase under exterior-only quantum operations (CPTP maps), by data-processing for relative entropy [287, 441, 230].

Interpretation. Near $\mathcal{H}_+$, the suppression of cross-horizon correlators (eq. (16.278)) drives $S(\mathcal{R})$ to encode the effective loss of exterior access to interior coherence. The emergent horizon temperature T_{eff} then appears as the KMS parameter of the exterior algebra's modular flow that *statistically* summarizes this entanglement structure (cf. [230, 30] and eq. (16.193)).

2. Emergence of the Horizon as a Modal Entanglement Cut

Proposition 16.37 (The horizon as a modal entanglement cut (operational form))**.** *Let $\mathcal{X}_q = \mathcal{R}_{\text{ext}} \cup \mathcal{R}_{\text{int}}$ be the exterior–interior bipartition across the outer horizon $\mathcal{H}_+$. Define the exterior restriction map $\mathcal{R}_{\text{ext}}$ as in Definition 16.32 and let $\mathcal{R}_{\text{int}}$ be the Gibbs/KMS-preserving conditional expectation onto the interior algebra with reference state $\hat{\tau}_{\text{int}}$ at (β_H, Ω_H) (cf. eq. (16.193)). Consider the* operational *joint state accessible to exterior observers,*

$$\tilde{\rho}_{\text{op}} \;:=\; (\mathcal{R}_{\text{ext}} \otimes \mathcal{R}_{\text{int}})(\hat{\rho}),$$

and its mutual information

$$I_{\mathrm{op}}(\mathcal{R}_{\mathrm{ext}} : \mathcal{R}_{\mathrm{int}}) := S\big(\tilde{\rho}_{\mathrm{op}}^{\mathrm{ext}}\big) + S\big(\tilde{\rho}_{\mathrm{op}}^{\mathrm{int}}\big) - S(\tilde{\rho}_{\mathrm{op}})\,,$$

where S is the von Neumann entropy (Definition 16.36). Then, in the sharp-cut/KMS limit with cross-horizon clustering (eq. (16.278)),

$$\boxed{\lim_{x \to \mathcal{H}_+^-} I_{\mathrm{op}}(\mathcal{R}_{\mathrm{ext}} : \mathcal{R}_{\mathrm{int}}) \;=\; 0.} \tag{16.284}$$

Consequently, the exterior entropy $S(\mathcal{R}_{\mathrm{ext}})$ reduces to a boundary-localized contribution at $\mathcal{H}_+$, identified with the thermal (Bekenstein–Hawking) entropy in the stationary regime [86, 87, 7]; see also the area-law perspective [653].

Proof. Step 1: Exterior accessibility and CPTP monotonicity. By construction, $\mathcal{R}_{\mathrm{ext}}$ and $\mathcal{R}_{\mathrm{int}}$ are completely positive, trace-preserving (CPTP) maps (conditional expectations) that preserve the co-rotating KMS reference [30, 230, 441]. Mutual information admits the relative-entropy representation $I(A:B) = D(\rho_{AB}\|\rho_A \otimes \rho_B)$ and is monotone under local CPTP maps:

$$I((\Phi_A \otimes \Phi_B)\rho) \;\le\; I(\rho)\,,$$

by data processing for relative entropy [287, 617, 357, 50]. Hence it suffices to bound $I(\tilde{\rho}_{\mathrm{op}})$.

Step 2: Cross-horizon clustering ⇒ product structure. In a stationary KMS neighborhood at T_{eff} (eqs. (16.2) and (16.193)), connected exterior–interior correlators decay exponentially with proper separation across $\mathcal{H}_+$ (eq. (16.278) and the clustering remark). Complete-positivity and trace-norm contractivity of $\mathcal{R}_{\mathrm{ext}} \otimes \mathcal{R}_{\mathrm{int}}$ then imply

$$\big\|\tilde{\rho}_{\mathrm{op}} - \tilde{\rho}_{\mathrm{op}}^{\mathrm{ext}} \otimes \tilde{\rho}_{\mathrm{op}}^{\mathrm{int}}\big\|_1 \;\longrightarrow\; 0 \qquad (x \to \mathcal{H}_+^-).$$

Since $I(\rho_{AB}) = D(\rho_{AB}\|\rho_A \otimes \rho_B)$ is continuous at product states and vanishes iff $\rho_{AB} = \rho_A \otimes \rho_B$, we obtain $I(\tilde{\rho}_{\mathrm{op}}) \to 0$ in the sharp-cut limit.

Step 3: Boundary localization of $S(\mathcal{R}_{\mathrm{ext}})$. With bulk exterior–interior correlations operationally extinguished, the remaining entropy of $\mathcal{R}_{\mathrm{ext}}$ is localized to the interface degrees of freedom at $\mathcal{H}_+$ (an "area term" in the continuum limit). In the stationary regime, consistency with the first-law identity $\delta M = (\kappa_{\mathrm{eff}}/8\pi G)\,\delta A$ (eq. (16.194)) and the KMS temperature T_{eff} (eq. (16.193)) yields the area law $S = \frac{c^3}{4G\hbar}\,A$ (eq. (16.195)), matching the Bekenstein–Hawking entropy [86, 87, 7] and the entanglement/area correspondence [653]. □

Remark 16.38 (Global vs. operational mutual information)**.** For a *global* pure state $|\mathring{\mathbf{\Psi}}\rangle$, one has $S(\mathcal{X}_q) = 0$ and $I(\mathcal{R}_{\mathrm{ext}} : \mathcal{R}_{\mathrm{int}}) = 2S(\mathcal{R}_{\mathrm{ext}})$ in principle. Proposition 16.37 concerns the *operational* mutual information accessible to exterior observers after the CPTP restriction $(\mathcal{R}_{\mathrm{ext}} \otimes \mathcal{R}_{\mathrm{int}})$ that implements loss of access to interior coherence. It is this operational quantity that vanishes in the sharp-cut/KMS limit, leaving only the boundary entropy at $\mathcal{H}_+$.

3. The Ergospheric Entropy Tensor

Definition 16.39 (Ergospheric entropy tensor)**.** Fix a small geodesic "modular diamond" $\mathcal{R}_x \subset \mathcal{X}_q$ centered at x whose linear size ℓ satisfies $\ell \ll \min\{\mathcal{R}_{\text{curv}}, \mathcal{R}_{\text{KMS}}\}$ (curvature and KMS correlation lengths). Let $\hat{\rho}_{\mathcal{R}_x}$ be the reduced state and $\hat{K}_{\mathcal{R}_x} := -\ln \hat{\rho}_{\mathcal{R}_x}$ its modular Hamiltonian (cf. Definitions 16.36 and 16.41). Define the local *modular-entropy potential*

$$\Xi(x) \;:=\; \mathrm{Tr}_{\mathcal{R}_x}\!\left(\hat{\rho}_{\mathcal{R}_x}\,\hat{K}_{\mathcal{R}_x}\right) \;=\; S(\mathcal{R}_x),$$

and the **ergospheric entropy tensor** by the covariant Hessian of Ξ:

$$\boxed{\hat{\mathcal{S}}_{\mu\nu}(x) \;:=\; \nabla_\mu \nabla_\nu\, \Xi(x)\,.} \tag{16.285}$$

Its trace $\hat{\mathcal{S}}^{\mu}{}_{\mu} = \Box \Xi$ measures the *entropic curvature* (the local convexity of the entropy landscape), while its divergence defines the *entropic flux density*

$$\hat{\jmath}^{\mu}_S(x) \;:=\; -\nabla_\nu \hat{\mathcal{S}}^{\mu\nu}(x)\,.$$

For an oriented two-surface $\mathcal{E}$ approximating the ergosurface, the *entropy flux across the interface* is

$$\Phi_S[\mathcal{E}] \;:=\; \int_{\mathcal{E}} n_\mu\, \hat{\jmath}^{\mu}_S\, d\Sigma,$$

with n_μ the unit normal. In a stationary KMS neighborhood at T_{eff} (eq. (16.193)), $\Phi_S[\mathcal{E}]$ quantifies the local rate of modal-entropy exchange induced by quantum ergodynamic processes (cf. Definition 16.34 and eq. (16.281)).

Remark 16.40 (Regulator and locality)**.** The diamond size ℓ acts as a UV regulator. For near-horizon, stationary configurations, $\hat{\mathcal{S}}_{\mu\nu}$ is regulator-independent to leading order in $\ell/\mathcal{R}_{\text{curv}}$ and $\ell/\mathcal{R}_{\text{KMS}}$, by locality of the modular flow in Rindler-like patches (Bisognano–Wichmann) and stability of KMS states [652, 230, 30].

4. The Modular Hamiltonian

Definition 16.41 (Modular Hamiltonian)**.** For any subsystem $\mathcal{R} \subset \mathcal{X}_q$ with reduced state $\hat{\rho}_{\mathcal{R}}$ (Definition 16.36), the **modular Hamiltonian** is

$$\boxed{\hat{H}_{\text{mod}}(\mathcal{R}) \;:=\; -\ln \hat{\rho}_{\mathcal{R}} \;\equiv\; \hat{K}_{\mathcal{R}}\,.} \tag{16.286}$$

It generates the Tomita–Takesaki modular flow $\sigma_t(A) = e^{it\hat{K}_{\mathcal{R}}} A e^{-it\hat{K}_{\mathcal{R}}}$ on the algebra of observables of $\mathcal{R}$ [230, 30]. In a Rindler wedge (Bisognano–Wichmann setting), $\hat{H}_{\text{mod}}$ is local and proportional to the boost generator:

$$\hat{H}_{\text{mod}}(\text{wedge}) \;=\; 2\pi \int_\Sigma \xi^a T_{ab}\, d\Sigma^b, \tag{16.287}$$

with ξ^a the wedge Killing field and T_{ab} the stress tensor [652, 641, 406]. By the near-horizon equivalence to Rindler, the same local form holds to leading order for a small diamond $\mathcal{R}_x$

straddling a stationary black-hole horizon, with ξ^a replaced by the horizon generator χ^a and the normalization fixed by the surface gravity κ:

$$\hat{H}_{\rm mod}(\mathcal{R}_x) \;\simeq\; \frac{2\pi}{\kappa}\int_{\Sigma_x}\chi^a T_{ab}\,d\Sigma^b, \qquad T_{\rm eff}=\frac{\kappa}{2\pi}\quad(\hbar=c=k_B=1), \tag{16.288}$$

recovering the KMS temperature of the horizon (eq. (16.193)) [7]. In particular, the entanglement "first law" $\delta S(\mathcal{R}_x)=\delta\langle\hat{H}_{\rm mod}(\mathcal{R}_x)\rangle$ links variations of the local entropy to the modular energy, which underlies the coupling between $\hat{\mathcal{S}}_{\mu\nu}$ and the ergodynamic currents introduced earlier (cf. eqs. (16.280) and (16.281)).

5. Emergence of Kerr Temperature Near $\mathcal{H}_+$, the reduced exterior state $\hat{\rho}_{\mathcal{R}_{\rm ext}}$ acquires an approximately thermal (KMS) form with respect to the local flow generated by the horizon Killing field $\chi^a=t^a+\Omega_H\phi^a$. This is the curved-spacetime analogue of the Bisognano–Wichmann (BW) result: in Minkowski space the vacuum restricted to a Rindler wedge is thermal with respect to the boost generator [652, 608]. In the stationary black-hole setting, the near-horizon region is locally Rindler, and the BW logic, together with the algebraic KMS framework, yields thermality for the exterior restriction [7, 11, 230, 30].

Proposition 16.42 (Emergence of horizon temperature)**.** *Let $\hat{H}_{\rm mod}(\mathcal{R}_x)$ be the modular Hamiltonian of a small diamond $\mathcal{R}_x$ adjacent to $\mathcal{H}_+$ (Definition 16.41), and let $\beta_H:=2\pi/\kappa$ with κ the surface gravity (eq. (16.2)). Then, in the sharp-cut/near-horizon limit,*

$$\boxed{\hat{\rho}_{\mathcal{R}_{\rm ext}} \;\approx\; \frac{1}{Z}\exp\!\Big(-\hat{H}_{\rm mod}(\mathcal{R}_x)\Big) \;\equiv\; \frac{1}{Z}\exp\!\Big(-\beta_H\,K_\chi\Big),} \tag{16.289}$$

where K_χ is the local physical generator (the Bisognano–Wichmann "boost'' built from the horizon Killing field χ^a). Equivalently, the emergent temperature is

$$\boxed{T_{\rm Kerr}=\frac{\kappa}{2\pi}\qquad\left(\textit{restoring units from geometric: } T_{\rm Kerr}=\frac{\hbar\,c}{2\pi k_B}\,\kappa\right).} \tag{16.290}$$

In geometric units ($G{=}c{=}\hbar{=}k_B{=}1$) the Kerr surface gravity and horizons are

$$\kappa=\frac{r_+-r_-}{2\,(r_+^2+a^2)}\,, \qquad r_\pm=M\pm\sqrt{M^2-a^2}\,, \tag{16.291}$$

and therefore

$$\boxed{T_{\rm Kerr}=\frac{\kappa}{2\pi}=\frac{\sqrt{M^2-a^2}}{4\pi\,M\,(M+\sqrt{M^2-a^2})}\,.} \tag{16.292}$$

Proof. (i) Local modular Hamiltonian.) By Definition 16.41, $\hat{H}_{\rm mod}(\mathcal{R}_x)=-\ln\hat{\rho}_{\mathcal{R}_x}$. In a Rindler wedge, BW gives the local form $\hat{H}_{\rm mod}=2\pi\int_\Sigma\xi^a T_{ab}\,d\Sigma^b$ with ξ^a the boost field

[652, 608]. Near a stationary horizon, replace the wedge by a small diamond $\mathcal{R}_x$ and ξ^a by the horizon generator χ^a, normalized by $\nabla_a(\chi^2) = -2\kappa\,\chi_a$ at $\mathcal{H}_+$; hence

$$\hat{H}_{\rm mod}(\mathcal{R}_x) \simeq \beta_H\, K_\chi, \qquad \beta_H = \frac{2\pi}{\kappa}. \tag{16.293}$$

(ii) Exterior restriction is KMS.) The algebraic KMS characterization implies that the exterior restriction at $\mathcal{H}_+$ is a β_H–KMS state for the modular flow generated by $\hat{H}_{\rm mod}(\mathcal{R}_x)$ (equivalently K_χ), giving the Gibbs form $\hat{\rho}_{\mathcal{R}_{\rm ext}} \approx Z^{-1} e^{-\hat{H}_{\rm mod}}$ [230, 30, 11]. This establishes eq. (16.289).

(iii) Temperature and surface gravity.) Identifying $\beta_H = 2\pi/\kappa$ yields $T_{\rm Kerr} = \kappa/(2\pi)$ (eq. (16.290)). For Kerr, $\kappa = \frac{r_+ - r_-}{2(r_+^2 + a^2)}$, $r_\pm = M \pm \sqrt{M^2 - a^2}$ (eq. (16.291)), so in geometric units

$$T_{\rm Kerr} = \frac{\kappa}{2\pi} = \frac{\sqrt{M^2 - a^2}}{4\pi\, M\,(M + \sqrt{M^2 - a^2})}, \tag{16.294}$$

and restoring constants gives $T_{\rm Kerr} = \dfrac{\hbar\,\kappa}{2\pi k_B\, c}$ [7, 11]. □

Remark 16.43 (Rotation and co-rotating ensemble)**.** Because $\chi^a = t^a + \Omega_H \phi^a$, the state in eq. (16.289) is thermal with respect to the *co-rotating* generator; equivalently, $\hat{\rho} \propto \exp\!\big[-\beta_H\,(\hat{H} - \Omega_H \hat{J})\big]$. This is the standard KMS structure for rotating horizons and is consistent with the generalized first law/free–energy relations used earlier (eqs. (16.194) and (16.272)).

6. Topological Quantization of Entropy

Definition 16.44 (Topological quantization of horizon entropy)**.** Let $\hat{A}_{\mathcal{H}}$ be the horizon-area observable and let $\{|\Psi_n\rangle\}_{n\in\mathbb{Z}_+}$ denote the *topological entanglement sectors* of the Kerr Attractor (cf. Section 16.4.8 and eq. (16.268)). Assuming *sector-preserving* (no-surgery) dynamics on $\mathcal{H}_+$ and stationarity (Definition 16.21), the **entropy operator** of the horizon is defined by

$$\boxed{\hat{S}_{\mathcal{H}} \;:=\; \frac{c^3}{4G\hbar}\,\hat{A}_{\mathcal{H}}\,, \qquad (\text{geometric units:}\;\; \hat{S}_{\mathcal{H}} = \tfrac{1}{4}\hat{A}_{\mathcal{H}}\,).} \tag{16.295}$$

Its spectrum is discrete and organized by the sector label n:

$$\boxed{\hat{S}_{\mathcal{H}}\,|\Psi_n\rangle = \frac{c^3}{4G\hbar}\,A_n\,|\Psi_n\rangle, \qquad A_n \in \mathrm{Spec}(\hat{A}_{\mathcal{H}}),} \tag{16.296}$$

with degeneracies determined by holonomy eigenphases $\{\theta_j\}$ (eq. (16.262)) and the entanglement topology $H_1(\mathcal{G}_{\rm Kerr}) \cong \mathbb{Z}$ (eq. (16.265)).

Remark 16.45 (Consistency, sector structure, and semiclassical limit)**.**

(i) First-law/KMS consistency.
The operator identity (16.295) is the operator-level lift of the area law $S = \frac{c^3}{4G\hbar}A$, compatible with the first-law variation $\delta M = (\kappa_{\rm eff}/8\pi G)\,\delta A$ and $T_{\rm Kerr} = \hbar\kappa/(2\pi k_B c)$ (eqs. (16.194) and (16.290)); see [86, 87, 7].

(ii) Sector preservation.
If the horizon graph $\mathcal{G}_{\mathcal{H}_+}$ is not altered by surgery, the label n is conserved under adiabatic (slow) processes. Transitions between $\{|\Psi_n\rangle\}$ require topology-changing events and thus appear as non-adiabatic jumps in the spectrum (16.296).

(iii) Kerr specialization.
Classically, $A = 4\pi(r_+^2 + a^2)$ with $r_\pm = GM \pm \sqrt{G^2M^2 - a^2}$. In sector n, the expectation values (M_n, a_n) (eq. (16.267)) determine A_n via this relation, tying the entropic spectrum to the modal quantization of mass and spin.

(iv) Semiclassical limit.
As the topological gap (between adjacent A_n) becomes small compared to A, the spectrum is effectively continuous and one recovers the standard thermodynamic area law [7].

(v) Additivity.
For disconnected horizons, $\hat{S}_{\mathcal{H}}$ is additive (direct sum over components), matching the additivity of $\hat{A}_{\mathcal{H}}$ and the extensivity of entropy in stationary, weakly interacting sectors.

Horizon temperature as a shadow of quantum entanglement geometry In this entanglement-centric view, both temperature and entropy arise from the holographic imprint of modal correlations across the entanglement cut at $\mathcal{H}_+$ (Proposition 16.37). The near-horizon KMS form of the exterior state (eq. (16.289)) yields $T_{\text{Kerr}} = \kappa/2\pi$ (eq. (16.290)), while the sector-organized spectrum of $\hat{S}_{\mathcal{H}}$ in (16.296) realizes the Bekenstein–Hawking law at the operator level [86, 87, 7]. Thus, black-hole thermodynamics appears as the emergent algebra of *relational coherence*: the "thermal" properties of the horizon are statistical shadows of quantum topological structure, with ergodynamic flows and modular dynamics (eq. (16.281) and Definition 16.41) knitting together gravity, thermodynamics, and information.

16.4.11 Inner Horizon Instability and the Modal Breakdown

Classically, the inner (Cauchy) horizon $\mathcal{H}_-$ of Kerr marks the boundary where predictability fails due to the infinite blueshift of ingoing perturbations and the onset of mass inflation [7, 657, 141]. In the present framework, this phenomenon is reinterpreted as a *quantum-critical transition* of the Quantum Holo-State: a breakdown of the differentiable dependence of the emergent chrono-geometry $\bar{\Sigma}_{\mu\nu}$ on the inter-modal correlator $\hat{G}_\Lambda$ (Definitions 6.4 and 11.106). Rather than signaling a failure of physical law, $\mathcal{H}_-$ is the locus where *modal coherence* loses stability and the relational arrow of emergent time collapses.

1. The Cauchy horizon as a modal critical surface

Definition 16.46 (Geodesic (causal) diamond at x)**.** Let x lie in a convex normal neighborhood U_x of the emergent chronogeometry $\bar{\Sigma}_{\mu\nu}$. Pick a future-directed unit timelike vector $u^a \in T_x\mathcal{X}_q$ and let $\gamma(\tau)$ be the timelike geodesic with $\gamma(0) = x$, $\dot{\gamma}(0) = u^a$. For a proper-time scale $\ell > 0$

satisfying $\ell \ll \min\{\mathcal{R}_{\mathrm{curv}}(x), \mathcal{R}_{\mathrm{KMS}}(x)\}$,[7] set $p_\pm := \gamma(\pm\ell)$ and define the *geodesic (causal) diamond*

$$\mathcal{R}_x(\ell, u) \;:=\; J^+_{\bar{\Sigma}}(p_-) \cap J^-_{\bar{\Sigma}}(p_+) \cap U_x. \tag{16.297}$$

Its boundary decomposes as $\partial\mathcal{R}_x = N^+ \cup N^- \cup \Sigma_+ \cup \Sigma_-$, where $N^\pm$ are the null sheets generated from $p_\pm$ and $\Sigma_\pm$ are small spacelike caps. In Riemann normal coordinates on U_x, $g_{\mu\nu} = \eta_{\mu\nu} + O(\mathcal{R}\,\ell^2)$ and the horizon generator χ^a is approximately Killing over $\mathcal{R}_x$ (near-horizon Rindler patch), so the Bisognano–Wichmann/KMS machinery applies locally [652, 230, 7].

Here $\mathfrak{H}_{\mathrm{HS}}(\mathcal{R}_x \times \mathcal{R}_x)$ denotes the Hilbert–Schmidt space of square-integrable kernels on $\mathcal{R}_x \times \mathcal{R}_x$ with respect to the volume form induced by $\bar{\Sigma}_{\mu\nu}$ (cf. Definitions 16.41 and 16.46).

Definition 16.47 (Modal criticality at the Cauchy horizon)**.** Fix a small geodesic diamond $\mathcal{R}_x$ centered at x and consider the Fréchet derivative (modal susceptibility operator) of the emergent metric with respect to the unified inter-modal correlator:

$$\begin{aligned} \mathbb{J}_{\mu\nu}[x;\,\cdot,\cdot] &:= \frac{\delta\bar{\Sigma}_{\mu\nu}(x)}{\delta\hat{G}_\Lambda(\cdot,\cdot)} \\ &:\; \mathfrak{H}_{\mathrm{HS}}\Big(\mathcal{R}_x \times \mathcal{R}_x\Big) \longrightarrow \mathrm{Sym}^2\Big(T^*_x\Big). \end{aligned} \tag{16.298}$$

where $\mathfrak{H}_{\mathrm{HS}}$ is the Hilbert–Schmidt space of kernel fluctuations supported on $\mathcal{R}_x$. Write $\sigma_{\min}(\mathbb{J}_x)$ for the smallest singular value of $\mathbb{J}_x$ (equivalently, the inverse operator norm of its Moore–Penrose pseudoinverse on the image).

Definition. The inner (Cauchy) horizon is the **modal critical surface**

$$\boxed{\mathcal{H}_- \;:=\; \Big\{\, x \in \mathcal{X}_q \;\Big|\; \sigma_{\min}\Big(\mathbb{J}_x\Big) = 0 \Big\},} \tag{16.299}$$

i.e. the locus where infinitesimal variations of $\hat{G}_\Lambda$ can produce unbounded (or undefined) variations of $\bar{\Sigma}_{\mu\nu}$. Equivalently, for any finite-rank compression of eq. (16.298) onto a local basis $\{\mathfrak{B}^{(k)}_x\}_{k=1}^N$,

$$J^{(N)}_{\mu\nu}(x) \;:=\; \left[\frac{\partial\bar{\Sigma}_{\mu\nu}(x)}{\partial G^{(k\ell)}_\Lambda}\right]_{1\le k,\ell\le N} \qquad\Longrightarrow\qquad \lim_{N\to\infty} \det\Big(J^{(N)}(x)J^{(N)}(x)^\top\Big) = 0. \tag{16.300}$$

At $\mathcal{H}_-$, the stable, differentiable map $\hat{G}_\Lambda \mapsto \bar{\Sigma}_{\mu\nu}$ ceases to be invertible: *modal criticality* replaces classical predictability failure.

Remark 16.48 (Well–posedness, regulators, and classical limit)**.** (i) The small diamond $\mathcal{R}_x$ provides a UV regulator; the criterion (16.299) is regulator–independent to leading order, as different choices of local fiber bases yield unitarily equivalent compressions in (16.300). (ii) In the Kerr background, the critical surface defined by (16.299) coincides with the classical inner horizon $r = r_-$; the divergence of the susceptibility mirrors the classical

[7] $\mathcal{R}_{\mathrm{curv}}$ is a local curvature radius; $\mathcal{R}_{\mathrm{KMS}}$ is the KMS correlation length near the horizon, cf. eqs. (16.2) and (16.193).

infinite blueshift/mass–inflation mechanism [141, 7]. (iii) Away from $\mathcal{H}_-$, $\sigma_{\min}(\mathbb{J}_x) > 0$ and the emergent geometry responds smoothly to correlator perturbations, recovering ordinary stability of the exterior region and the outer horizon $\mathcal{H}_+$ discussed earlier.

Intuitively, eq. (16.299) states that at $\mathcal{H}_-$ the *modal susceptibility* diverges: infinitesimal rearrangements of inter-modal correlations can trigger order–one deformations of the chrono-geometry. The classical "breakdown of determinism'' is thus recast as a *quantum–critical breakdown of relational differentiability*, presaging a phase transition in the coherence topology of the Quantum Holo–State rather than a failure of the laws themselves.

2. Infinite Blueshift as Modal Divergence

Proposition 16.49 (Infinite blueshift as modal energy divergence)**.** *Let $\mathcal{H}_-$ be the modal critical surface of Definition 16.47 and let $\hat{T}_{\mu\nu}$ denote the relational stress–energy of the PQS ensemble. Define the observer–independent local energy scalar*

$$\hat{\mathcal{E}}(x) \;:=\; \hat{T}^{\mu\nu}(x)\,\hat{T}_{\mu\nu}(x), \qquad \text{and, for any future timelike } u^\mu, \quad \hat{\varepsilon}(x;u) := \hat{T}_{\mu\nu}(x)u^\mu u^\nu. \tag{16.301}$$

Assume generic (nonvanishing) ingoing perturbations at $\mathcal{H}_+$ and stationarity up to the inner region. Then, as $x \to \mathcal{H}_-^-$,

$$\boxed{\lim_{x\to\mathcal{H}_-^-} \left\langle \hat{\mathcal{E}}(x) \right\rangle \;=\; +\infty \qquad \textit{and hence} \qquad \lim_{x\to\mathcal{H}_-^-} \left\langle \hat{\varepsilon}(x;u) \right\rangle \;=\; +\infty,} \tag{16.302}$$

i.e. the classical infinite blueshift at the Cauchy horizon is the divergence of modal energy observables. *The blowup arises from the critical amplification of inter–modal coherence as the causal structure compresses toward the locus (16.299), not from any literal piling–up of classical matter [7, 318, 141].*

Proof. (i) Modal criticality ⇒ divergent susceptibility.) At $\mathcal{H}_-$ the susceptibility operator $\mathbb{J}_{\mu\nu}[x;\cdot,\cdot] = \delta\bar{\Sigma}_{\mu\nu}(x)/\delta\hat{G}_\Lambda(\cdot,\cdot)$ develops a null singular value (eqs. (16.298) and (16.299)). Hence, for generic fluctuations $\delta\hat{G}_\Lambda$ supported in a small diamond $\mathcal{R}_x$, the induced variation of the emergent metric satisfies

$$\|\delta\bar{\Sigma}(x)\| \;\gtrsim\; \sigma_{\min}(\mathbb{J}_x)^{-1}\,\|\delta\hat{G}_\Lambda\|, \qquad \sigma_{\min}(\mathbb{J}_x) \to 0, \tag{16.303}$$

signaling unbounded geometric response to finite coherence perturbations.

(ii) Field equations tie curvature to stress–energy.) By the chronogeometric field relation (unified/vacuum Einstein map; cf. Definition 11.106),

$$\bar{R}_{\mu\nu}[\bar{\Sigma}] \;=\; \mathcal{R}^{(\mathrm{vac})}_{\mu\nu}[\langle\hat{T}_{\alpha\beta}\rangle;\,\bar{\Sigma}],$$

the curvature computed from second derivatives of $\bar{\Sigma}$ is a local functional of the (renormalized) expectation value $\langle\hat{T}_{\mu\nu}\rangle$. Linearizing about a stationary background and using elliptic/Green estimates on the diamond $\mathcal{R}_x$ gives a schematic bound

$$\|\delta\bar{\Sigma}(x)\| \;\lesssim\; C\,\left\|\delta\langle\hat{T}_{\mu\nu}\rangle\right\|_{\mathcal{R}_x}, \tag{16.304}$$

for some geometric constant C fixed by the background.

(iii) Contradiction unless stress–energy blows up.) Combining (16.303) and (16.304), with $\sigma_{\min}(\mathbb{J}_x) \to 0$, forces $\|\delta\langle\hat{T}_{\mu\nu}\rangle\|_{\mathcal{R}_x} \to \infty$ for generic $\delta\hat{G}_\Lambda$. Hence some quadratic stress–energy scalar—such as $\langle\hat{\mathcal{E}}(x)\rangle = \langle\hat{T}^{\mu\nu}\hat{T}_{\mu\nu}\rangle$—must diverge as $x \to \mathcal{H}_-^-$, establishing the first limit in eq. (16.302). Positivity of $\hat{\mathcal{E}}$ then implies divergence of the energy density measured by any timelike u^μ, giving the second limit.

(iv) Relation to classical blueshift/mass inflation.) In classical Kerr, the inner horizon has (signed) surface gravity $\kappa_- < 0$; along the Cauchy horizon, advanced time $v \to +\infty$ and an infalling observer sees a blueshift factor $\sim e^{|\kappa_-|v}$ multiplying any residual ingoing flux. Generic perturbations from $\mathcal{H}_+$ (Price tails) thus yield energy densities $\sim e^{2|\kappa_-|v}$, i.e. blowup at $\mathcal{H}_-$ [141]. Our modal derivation recovers this as the critical amplification of inter–modal coherence encoded by (16.299). □

Remark 16.50 (Observer scalars and tetrad components)**.** The invariant $\hat{\mathcal{E}} = \hat{T}^{\mu\nu}\hat{T}_{\mu\nu}$ controls the blowup of all physically meaningful projections. In a Newman–Penrose frame adapted to the principal null directions $(\ell^a, n^a, m^a, \bar{m}^a)$, divergence of $\langle\hat{\mathcal{E}}\rangle$ entails divergence of flux components such as $\langle\hat{T}_{\mu\nu}\ell^\mu\ell^\nu\rangle$ and $\langle\hat{T}_{\mu\nu}n^\mu n^\nu\rangle$, reproducing the mass–inflation phenomenology at $\mathcal{H}_-$ [318, 141].

3. Temporal Gradient Collapse and the Loss of Global Coherence

Proposition 16.51 (Temporal gradient collapse)**.** *Let $\hat{\tau}_\mu$ be the operator-valued temporal potential introduced in Definition 16.13, and let χ^a denote the (co-rotating) horizon generator. As one approaches the Cauchy horizon $\mathcal{H}_-$ from the inter-horizon region, the temporal field aligns with the inner-horizon generator and its norm collapses:*

$$\hat{\tau}^\mu(x) \;=\; N(x)\,\chi^\mu(x), \qquad \hat{\tau}^2(x) := \bar{\Sigma}_{\mu\nu}\,\hat{\tau}^\mu\hat{\tau}^\nu \longrightarrow 0, \qquad x \to \mathcal{H}_-^-, \tag{16.305}$$

with a lapse $N(x) \to 0$. *Consequently, the divergence of $\hat{\tau}^\mu$ collapses in the same limit,*

$$\boxed{\lim_{x\to\mathcal{H}_-^-} \nabla_\mu\hat{\tau}^\mu(x) \;=\; 0,} \tag{16.306}$$

signaling the loss of an oriented modular/temporal gradient. Operationally, the modular flow ceases to be globally well-defined, and the relational arrow of time flattens *into indeterminacy.*

Proof sketch. (i) Alignment at modal criticality.) By Definition 16.47, the susceptibility $\mathbb{J}_{\mu\nu}[x;\cdot,\cdot]$ becomes singular at $\mathcal{H}_-$, implying that only deformations along the null generator of the inner horizon remain dynamically stable. The temporal field, which drives the local modular/KMS flow near horizons, is therefore forced to align with χ^μ at the critical surface; this yields (16.305), with $\hat{\tau}^2 \to 0$ as χ^μ is null on $\mathcal{H}_-$ [7, 318, Sec. 12].

(ii) Divergence collapse.) Writing $\hat{\tau}^\mu = N\,\chi^\mu$ gives

$$\nabla_\mu\hat{\tau}^\mu = \chi^\mu\nabla_\mu N + N\,\nabla_\mu\chi^\mu. \tag{16.307}$$

For a (background) Killing horizon, $\nabla_\mu\chi^\mu = 0$ along the generators [7, Sec. 7.1], so $\nabla_\mu\hat{\tau}^\mu = \chi^\mu\nabla_\mu N$. The lapse N encodes the local redshift between modular and geometric time; near

$\mathcal{H}_-$ it collapses due to the critical susceptibility (the same mechanism behind the infinite blueshift/mass inflation [141]). In stationary approximation along the generators, $\chi^\mu \nabla_\mu N \to 0$ as $N \to 0$, yielding (16.306).

(iii) Consistency with energy blowup.) The vanishing of $\nabla \cdot \hat{\tau}$ does *not* contradict the divergence of stress–energy invariants in Proposition 16.49: the latter is controlled by shear/curvature growth and critical amplification of inter-modal coherence, whereas the expansion (divergence) of the *null generators* remains zero on a Killing horizon. The lapse factor N carries the redshift/blueshift information while driving $\nabla_\mu \hat{\tau}^\mu \to 0$. □

Remark 16.52 (Vorticity vs. expansion; modular-flow interpretation)**.** Decompose the chrono-geometric derivatives into *divergence* $\theta_\tau := \nabla_\mu \hat{\tau}^\mu$, *vorticity* $\hat{\omega}_{\mu\nu} := \nabla_{[\mu} \hat{\tau}_{\nu]}$ (cf. Definition 16.13), and *shear*. At $\mathcal{H}_-$, $\theta_\tau \to 0$ while $\hat{\omega}_{\mu\nu}$ and shear need not vanish; indeed, classical analyses show shear/curvature blowup at the Cauchy horizon (mass inflation) [141]. In the modular picture, $\hat{\tau}^\mu$ generates the local KMS flow; the limit (16.306) means that, although the flow becomes *null and tangent* to the horizon generators, it loses a global orientation—hence the "flattening" of the relational arrow of time.

Interpretation. Equation (16.306) captures the decisive difference between the outer and inner horizons. At $\mathcal{H}_+$, the co-rotating KMS structure supplies a well-oriented modular flow with nonzero temperature (Proposition 16.42); at $\mathcal{H}_-$, the modal criticality forces alignment with null generators and collapses the divergence, rendering the global unitary history non-orientable even as local stresses blow up (Proposition 16.49).

4. Modal Phase Decorrelation and the Breakdown of Evolution Operators

Definition 16.53 (Bi-local (relative) modular generator)**.** Let $\mathcal{R}_x, \mathcal{R}_{x'}$ be two small geodesic diamonds (cf. Definition 16.46) and denote their modular Hamiltonians by $\hat{K}_{\mathcal{R}_x}$, $\hat{K}_{\mathcal{R}_{x'}}$, and $\hat{K}_{\mathcal{R}_x \cup \mathcal{R}_{x'}}$ (Definition 16.41). Define the *bi-local modular (coupling) generator* as the cross term in the union:

$$\hat{H}_{\mathrm{mod}}(x, x') \;:=\; \hat{K}_{\mathcal{R}_x \cup \mathcal{R}_{x'}} - \hat{K}_{\mathcal{R}_x} - \hat{K}_{\mathcal{R}_{x'}}. \tag{16.308}$$

This operator is invariant under additive gauge shifts $\hat{K}_{\mathcal{R}} \mapsto \hat{K}_{\mathcal{R}} + \alpha_{\mathcal{R}} \mathbb{I}$ and measures the *phase-coupling* between the two fibers.

Proposition 16.54 (Modal phase decorrelation at the inner horizon)**.** *Define the modal phase coherence functional for two diamonds by*

$$\boxed{\mathcal{C}(x, x') \;:=\; \left| \langle \overset{\circ}{\boldsymbol{\Psi}} \,|\, e^{\,i\, \hat{H}_{\mathrm{mod}}(x,x')} \,|\, \overset{\circ}{\boldsymbol{\Psi}} \rangle \right|,} \tag{16.309}$$

with $\hat{H}_{\mathrm{mod}}(x, x')$ from eq. (16.308). As (x, x') approach the Cauchy horizon $\mathcal{H}_-$ from the inter-horizon region,

$$\boxed{\lim_{x, x' \to \mathcal{H}_-} \mathcal{C}(x, x') \;=\; 0.} \tag{16.310}$$

Consequently, no strongly continuous, unitary modular evolution $\hat{U}_{\mathrm{mod}}(t)$ can be defined across $\mathcal{H}_-$ for the exterior/interior relational algebra: the global phase reference collapses and modular flow loses its algebraic compass.

Proof sketch. (i) Near-horizon form of modular generators.) For a small diamond $\mathcal{R}$ adjacent to a stationary horizon, the modular Hamiltonian admits the local BW/KMS form

$$\hat{K}_{\mathcal{R}} \simeq \frac{2\pi}{\kappa} \int_{\Sigma_{\mathcal{R}}} \chi^a T_{ab}\, d\Sigma^b \quad \text{(to leading order)},$$

with χ^a the appropriate horizon generator (eqs. (16.288) and (16.290)) [652, 230, 7]. Therefore the bi-local generator eq. (16.308) is controlled by *connected* stress–energy integrals supported in $\mathcal{R}_x \cup \mathcal{R}_{x'}$.

(ii) Critical amplification of modular fluctuations.) At $\mathcal{H}_-$ the susceptibility $\mathbb{J}_{\mu\nu}$ diverges (Definition 16.47), and stress–energy scalars blow up (mass inflation; Proposition 16.49, [141, 318]). Hence the *variance* of the centered bi-local generator,

$$\mathrm{Var}\big[\hat{H}_{\mathrm{mod}}(x,x')\big] = \Big\langle \big(\Delta \hat{H}_{\mathrm{mod}}(x,x')\big)^2 \Big\rangle,$$

diverges as $(x,x') \to \mathcal{H}_-^-$, by the fluctuation–response identity inherited from the BW/KMS structure and the growth of connected T_{ab}–correlators (linear response on KMS states [230, 30, Chs. 5–6]).

(iii) Phase characteristic function ⇒ decorrelation.) Set $X := \Delta \hat{H}_{\mathrm{mod}}(x,x')$ (self-adjoint, centered) and let $\mu_{x,x'}$ be its spectral measure in the state $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$. Then

$$\langle e^{\,i\hat{H}_{\mathrm{mod}}(x,x')} \rangle = e^{\,i\langle \hat{H}_{\mathrm{mod}} \rangle} \int_{\mathbb{R}} e^{i\lambda}\, d\mu_{x,x'}(\lambda).$$

In local AQFT (type III algebras) the absolutely continuous component of $\mu_{x,x'}$ is nontrivial; as $\mathrm{Var}_{\mu_{x,x'}}(X) \to \infty$, the characteristic function at unit argument decays to zero under mild regularity (Riemann–Lebesgue for the a.c. part; the pure point part carries vanishing weight in the limit) [230, 30]. Consequently $|\langle e^{\,i\hat{H}_{\mathrm{mod}}(x,x')}\rangle| \to 0$, i.e. eq. (16.310).

(iv) Breakdown of strongly continuous modular evolution.) If $\mathcal{C}(x,x') \to 0$ for all pairs tending to $\mathcal{H}_-$, then no common phase reference survives to implement a strongly continuous one-parameter modular group across the surface (Stone's theorem would require a self-adjoint generator with well-defined spectral exponentials). Thus $\hat{U}_{\mathrm{mod}}(t)$ cannot be extended across $\mathcal{H}_-$ as a unitary evolution on the exterior/interior algebra, completing the argument [230, 30]. □

Remark 16.55 (Relation to temporal-gradient collapse)**.** The decorrelation eq. (16.310) complements the divergence-free limit $\nabla_\mu \hat{\tau}^\mu \to 0$ at $\mathcal{H}_-$ (Proposition 16.51). The former removes *phase coherence* between fibers (no global modular orientation), while the latter collapses the *temporal gradient* (no oriented flow even locally along $\hat{\tau}^\mu$). Together they encode the modal breakdown of evolution at the inner horizon.

5. Holosymmetric Collapse and the Transition to Pre-Geometric Regimes Beyond the inner horizon $\mathcal{H}_-$, the IQR description ceases to admit a well-defined emergent chronogeometry. In place of a manifold, one encounters a *holosymmetric collapse*: the relational order parameters that sustain distance and time vanish, and the modular/evolution structure becomes trivial.

Definition 16.56 (Pre-geometric condensate)**.** Let $\mathfrak{B}_{\mathcal{C}}$ denote the basin of PQSs interior to $\mathcal{H}_-$. We say that $\mathfrak{B}_{\mathcal{C}}$ is in a **pre-geometric condensate** when the unified inter-modal correlator $\hat{G}_\Lambda$ (Definition 6.4) and the temporal potential $\hat{\tau}^\mu$ (Definition 16.13) have vanishing expectations (in regulator-independent leading order) and trivial connected correlators within $\mathfrak{B}_{\mathcal{C}}$:

$$\boxed{\forall\, x, x' \in \mathfrak{B}_{\mathcal{C}}: \qquad \left\langle \hat{G}_\Lambda(x,x') \right\rangle \simeq 0, \qquad \left\langle \hat{\tau}^\mu(x) \right\rangle \simeq 0, \qquad \left\langle \hat{G}_\Lambda\, \hat{\tau} \right\rangle_c \simeq 0.} \tag{16.311}$$

Here $\simeq 0$ means suppressed relative to the local UV/IR regulators (small geodesic diamond of Definition 16.46 and near-horizon KMS scale), uniformly on compact subsets of $\mathfrak{B}_{\mathcal{C}}$. In this phase, the reconstruction map $\hat{G}_\Lambda \mapsto \bar{\Sigma}_{\mu\nu}$ (eq. (16.298)) yields a degenerate output, so no non-degenerate metric or global modular flow can emerge.

Proposition 16.57 (Holosymmetric collapse: necessary conditions)**.** *Assume the inner–horizon criticality of Definition 16.47, together with (i) modal energy divergence (Proposition 16.49), (ii) temporal–gradient collapse (Proposition 16.51), and (iii) modal phase decorrelation (Proposition 16.54). Then any interior limit state supported in $\mathfrak{B}_{\mathcal{C}}$ satisfies the pre–geometric conditions* (16.311)*. In particular, the emergent chrono–geometry $\bar{\Sigma}_{\mu\nu}$ is non–invertible (rank–deficient) almost everywhere in $\mathfrak{B}_{\mathcal{C}}$, and no strongly continuous modular evolution exists on the interior algebra.*

Proof. **Set–up and regularization.** Fix a point $x \in \mathfrak{B}_{\mathcal{C}}$ and a geodesic diamond $\mathcal{R}_x(\ell, u)$ as in Definition 16.46. For any local field/operator $\hat{O}(y)$ supported in $\mathcal{R}_x$, define the regulated average

$$\langle \hat{O} \rangle_{x,\ell} := \frac{1}{\mathrm{Vol}(\mathcal{R}_x)} \int_{\mathcal{R}_x} \langle \hat{O}(y) \rangle \, d\mu_{\bar{\Sigma}}(y), \qquad \mathrm{Vol}(\mathcal{R}_x) = \int_{\mathcal{R}_x} d\mu_{\bar{\Sigma}}. \tag{16.312}$$

We use the notations $\langle \cdot \rangle$ for the expectation in the global state $|\mathring{\Psi}\rangle$, and $(\cdot)_c$ for connected parts. We will show, uniformly on compact subsets of $\mathfrak{B}_{\mathcal{C}}$ and in the regulator limit $\ell \to 0$, that

$$\langle \hat{\tau}^\mu \rangle_{x,\ell} \to 0, \qquad \langle \hat{G}_\Lambda \rangle_{x,\ell} \to 0, \qquad \langle \hat{G}_\Lambda\, \hat{\tau} \rangle_{x,\ell,c} \to 0, \tag{16.313}$$

from which (16.311) follows.

Lemma 1 (Temporal potential vanishes). By Proposition 16.51, near $\mathcal{H}_-$ we have the alignment and lapse collapse

$$\hat{\tau}^\mu(y) = N(y)\, \chi^\mu(y), \qquad \hat{\tau}^2(y) = \bar{\Sigma}_{\mu\nu} \hat{\tau}^\mu \hat{\tau}^\nu \to 0, \qquad y \to \mathcal{H}_-^-, \tag{16.314}$$

with $N(y) \to 0$ and χ^μ null on $\mathcal{H}_-$. Assume (standard in the near–horizon AQFT setting) that $\|\chi\|$ is bounded on $\mathcal{R}_x$ and that $\|\hat{\tau}^\mu\|$ is dominated by $|N|\,\|\chi\|$ in operator norm to leading order.[8] Then by (16.312),

$$\left| \langle \hat{\tau}^\mu \rangle_{x,\ell} \right| \le \frac{1}{\mathrm{Vol}(\mathcal{R}_x)} \int_{\mathcal{R}_x} \|\hat{\tau}^\mu(y)\| \, d\mu \lesssim \sup_{y \in \mathcal{R}_x} |N(y)| \; \|\chi^\mu\|_{L^\infty(\mathcal{R}_x)} \xrightarrow[\ell \to 0]{} 0, \tag{16.315}$$

[8] This is the usual KMS/BW control of local generators in a Rindler–like patch; cf. Definition 16.41, eq. (16.288).

establishing the first limit in (16.313).

Lemma 2 (Bi–local phase decorrelation ⇒ vanishing connected parts). Let $\mathcal{R}_x, \mathcal{R}_{x'}$ be two diamonds with $(x, x') \to \mathcal{H}_-^-$, and define the bi–local modular generator $\hat{H}_{\mathrm{mod}}(x, x')$ by eq. (16.308). By Proposition 16.54 we have $\mathcal{C}(x, x') = \left|\langle e^{i\hat{H}_{\mathrm{mod}}(x,x')}\rangle\right| \to 0$. Let $\hat{A}$ and $\hat{B}$ be local bounded operators supported in $\mathcal{R}_x$ and $\mathcal{R}_{x'}$, respectively. Using the spectral calculus for $\hat{H}_{\mathrm{mod}}(x, x')$ and writing the *connected* two–point function in the KMS form as an L^1–Fourier transform (linear response/AQFT; see [230, 30]), one obtains schematically

$$\langle \hat{A}\, \hat{B}\rangle_c \;=\; \int_{\mathbb{R}} \widehat{F}_{A,B}(\omega) \;\underbrace{\langle e^{\,i\omega \hat{H}_{\mathrm{mod}}(x,x')}\rangle}_{\text{characteristic function}}\; d\omega, \tag{16.316}$$

for some absolutely integrable kernel $\widehat{F}_{A,B}$ depending on the local KMS data. As $(x, x') \to \mathcal{H}_-^-$, the characteristic function decays pointwise to zero for each fixed $\omega \neq 0$ by Proposition 16.54 and dominated convergence applies (the kernels $\widehat{F}_{A,B}$ are L^1–controlled uniformly on sufficiently small diamonds). Therefore

$$\lim_{x,x' \to \mathcal{H}_-^-} \langle \hat{A}\, \hat{B}\rangle_c \;=\; 0. \tag{16.317}$$

Taking $\hat{A} = \hat{G}_\Lambda(\cdot, x)$ and $\hat{B} = \hat{\tau}^\mu(\cdot)$ or $\hat{B} = \hat{G}_\Lambda(\cdot, x')$ yields, after diamond averaging, the third limit in (16.313) and the vanishing of the connected G–G parts used below.

Lemma 3 (Vanishing of $\langle \hat{G}_\Lambda\rangle$ to leading order). Decompose the two–point kernel into its connected and disconnected pieces in a local basis $\{\mathfrak{B}^{(k)}\}$ over $\mathcal{R}_x$:

$$\langle \hat{G}_\Lambda\rangle_{x,\ell} \;=\; \sum_{k,\ell} \langle \hat{G}_\Lambda^{(k\ell)}\rangle_{x,\ell}\, \mathfrak{B}^{(k)} \otimes \mathfrak{B}^{(\ell)} \;=\; \sum_{k,\ell} \Big(\langle \hat{G}^{(k)}\rangle \langle \hat{G}^{(\ell)}\rangle + \langle \hat{G}_\Lambda^{(k\ell)}\rangle_c \Big)\, \mathfrak{B}^{(k)} \otimes \mathfrak{B}^{(\ell)}.$$

The connected part vanishes by (16.317). The remaining disconnected part factorizes into one–point functions $\langle \hat{G}^{(k)}\rangle$. At the inner horizon, the only nontrivial one–point contributions compatible with the susceptibility singularity eqs. (16.298) and (16.299) would have to lie in the kernel of the reconstruction map $\mathbb{J}_x$. However, any such kernel component is *invisible* to the emergent geometry and, by stationarity and the collapse of the temporal gradient (Proposition 16.51), is averaged out in the diamond limit: the lack of an oriented modular flow (vanishing lapse) suppresses any persistent one–point bias.[9] Hence

$$\langle \hat{G}_\Lambda\rangle_{x,\ell} \;\xrightarrow[\ell \to 0]{}\; 0. \tag{16.318}$$

Lemma 4 (Rank–deficiency of $\bar{\Sigma}_{\mu\nu}$). By Definition 16.47, $\sigma_{\min}(\mathbb{J}_x) = 0$. Let $\{\delta\hat{G}^{(n)}\}$ be any bounded sequence of local kernel fluctuations normalized in the Hilbert–Schmidt norm on $\mathcal{R}_x \times \mathcal{R}_x$ and converging weakly to a limit in $\ker \mathbb{J}_x$. Then $\|\mathbb{J}_x[\delta\hat{G}^{(n)}]\| \to 0$, so the pushforward metric variations vanish in norm. Combined with (16.318) and (16.317), this

[9] Formally: for an averaged kernel component K in $\ker \mathbb{J}_x$, the modular time–average $T^{-1}\int_{-T}^{T} \langle \sigma_t(K)\rangle dt$ tends to 0 as $T \to \infty$ because the modular group ceases to be strongly continuous across $\mathcal{H}_-$ and the phase reference collapses (Proposition 16.54); see also the remark in Proposition 16.54.

shows that the *image* of the averaged correlator under $\mathbb{J}_x$ is degenerate; hence the effective metric $\bar{\Sigma}_{\mu\nu}$ reconstructed from interior data is non–invertible almost everywhere in $\mathfrak{B}_{\mathcal{C}}$.

Lemma 5 (No strongly continuous modular evolution). By Proposition 16.54, for any interior pair of diamonds (x, x') approaching $\mathcal{H}_-$ the characteristic function $\langle e^{i\omega \hat{H}_{\mathrm{mod}}(x,x')}\rangle$ tends to 0 for all $\omega \neq 0$. If a strongly continuous one–parameter unitary group $\hat{U}_{\mathrm{mod}}(t)$ existed on the interior algebra implementing modular flow, Stone's theorem would provide a self–adjoint generator $\hat{K}$ such that $\langle e^{i\omega \hat{K}}\rangle$ is continuous at $\omega = 0$ with $\langle e^{0\cdot i\hat{K}}\rangle = 1$ and nontrivial in a neighborhood. The above decay precludes this: there is no common phase reference to define a strongly continuous modular unitary across $\mathcal{H}_-$.

Conclusion. Lemmas 1–3 yield the pre–geometric conditions (16.311); Lemma 4 gives rank–deficiency of $\bar{\Sigma}_{\mu\nu}$; Lemma 5 rules out a strongly continuous modular evolution on the interior algebra. This proves the proposition. □

The inner horizon as a singularity of relational time In classical analyses, instability at the Cauchy horizon is tied to infinite blueshift and mass inflation [141, 7]. In the present account, *instability* is the ontological limit of modal coherence: by Propositions 16.49, 16.51 and 16.54, evolution ceases not through a breakdown of law, but through *holosymmetric collapse*—a rupture in the operator grammar from which spacetime is woven. Beyond $\mathcal{H}_-$, history does not end in paradox; it transforms into the pre-geometric condensate of Definition 16.56, where the emergent geometry dissolves and the Quantum Holo-State persists only as unstructured, relational potentiality.

16.5 Computing the Chronogeometric Kerr Solution

With the full theoretical apparatus of the rotating, chrono–geometric black hole in place (Section 16.4), we now undertake the constructive task: *reconstructing* the relationally-dressed geometry from first principles of the IQR framework. The classical Kerr family [7, 146] serves as the asymptotic guide, but the interior structure is determined by the quantum–coherent dynamics encoded in the unified tensor $\bar{\Sigma}_{\mu\nu}$ and its modular/KMS data. Our goal is a self-consistent solution—*the Chronogeometric Kerr Solution*—that satisfies the unified field equations, the near-horizon KMS structure, and the global topological constraints of the Kerr Attractor.

Computational assumptions and boundary data. We work in stationary, axisymmetric sectors (horizon generator $\chi^a = t^a + \Omega_H \phi^a$) with the unified lapse and co-rotating frame fixed by the near-horizon KMS temperature and angular velocity,

$$\begin{aligned} T_{\mathrm{Kerr}} &= \frac{\kappa}{2\pi} \\ \chi^a \chi_a\big|_{\mathcal{H}_+} &= 0 \end{aligned} \tag{16.319}$$

cf. eqs. (16.290) and (16.291). Asymptotically, $\bar{\Sigma}_{\mu\nu} \to g^{\mathrm{Kerr}}_{\mu\nu}$ (Boyer–Lindquist chart) with ADM data (M, a), and the holonomy/topological sector is fixed by $\mathcal{K}_n =$

$(H_1(\mathcal{G}_{\text{Kerr}}), \{\theta_j\}, |\Psi_n\rangle)$ (Section 16.4.8). Regularity is imposed at $\mathcal{H}_+$ (nonextremal) and compatibility at the inner horizon $\mathcal{H}_-$ follows the modal–criticality constraints (Section 16.4.11). Unless stated, we use geometric units $G{=}\hbar{=}c{=}k_B{=}1$.

Program architecture. At a high level, the computation proceeds from connection to curvature to sources and constraints, closing with existence/uniqueness and phenomenological checks. To support reproducibility, we organize the derivation into fifteen modular steps (I–XV), each producing labeled intermediate quantities that feed the next stage.

Roadmap (Steps I–XV).

(I) Coordinate/gauge and symmetry fix. Adopt the stationary–axisymmetric ansatz for $\bar{\Sigma}_{\mu\nu}$ specified in Section 16.4, fix signature and orientation, and choose a co-rotating frame aligned with χ^a; record unknown scalar functions and their regularity at (r_+, θ) and infinity.

(II) Relational Levi–Civita connection. Compute the *Relational Christoffel symbols* $\hat{\Gamma}[\bar{\Sigma}]$ from Definition 6.7, including all nonzero components and their horizon limits (ingoing coordinates to handle regularity on $\mathcal{H}_+$).

(III) Unified curvature tensors. Evaluate $\bar{R}^{\rho}{}_{\sigma\mu\nu}$, $\bar{R}_{\mu\nu}$, and $\bar{R}$ as in Definition 11.103; tabulate independent components under stationarity/axisymmetry and identify Bianchi identities.

(IV) Unified field equations. Insert **III** into the *Unified Vacuum Field Equation* (Definition 16.6) to obtain the coupled PDE system for the unknowns of $\bar{\Sigma}_{\mu\nu}$ driven by the rotationally dressed vacuum source.

(V) Rotationally dressed vacuum energy. Model $\rho_{\text{vac}}(r, \theta)$ via the KMS data and modular Hamiltonian (Definitions 16.5 and 16.41), enforcing the co-rotating Gibbs form and the first-law consistency with T_{Kerr} and Ω_H (eq. (16.290)); impose positivity/regularity conditions.

(VI) Boundary conditions at $\mathcal{H}_+$. Implement KMS boundary data (16.319), regularity of curvature scalars, and smoothness of the ingoing expansion; fix the area (hence $S_{\mathcal{H}}$ via Definition 16.44) and surface gravity.

(VII) Asymptotic conditions. Impose asymptotic flatness and Kerr falloffs, matching to (M, a) and the holonomy sector $\{\theta_j\}$; ensure the Komar/ADM charges agree with the near-horizon thermodynamic parameters (Smarr/first-law checks with [86, 7]).

(VIII) Constraint propagation and Bianchi closure. Show the contracted Bianchi identities close on the dressed source; derive secondary constraints eliminating spurious degrees of freedom in the PDE system.

(IX) Elliptic–hyperbolic decomposition. In the stationary reduction, isolate the elliptic sector (on spatial slices) and the residual first-order constraints along χ^a; specify functional spaces and coercivity estimates.

(X) Near-horizon expansion. Construct Frobenius series about $r = r_+$ consistent with KMS data and regularity; determine free coefficients in terms of (κ, Ω_H) and topological labels, fixing gauge remainders.

(XI) Asymptotic expansion. Build large-r expansions and match to Kerr asymptotics; ensure cancellation of secular terms and enforce parity under $\theta \mapsto \pi - \theta$.

(XII) Global solver. Formulate a continuation scheme that shoots from $\mathcal{H}_+$ to large r (or a spectral/finite-element scheme on (r, θ)); specify preconditioners, residuals, and convergence diagnostics tied to the constraints of **VIII**.

(XIII) Thermodynamic consistency. Verify the first law and Smarr relation at the solution level; check $S_{\mathcal{H}} = \frac{1}{4}A_{\mathcal{H}}$ and $T_{\text{Kerr}} = \kappa/2\pi$ (eq. (16.290)), with *co-rotating* ensemble density $\rho \propto e^{-\beta_H(\hat{H} - \Omega_H \hat{J})}$.

(XIV) Physical observables. Extract effective surface gravity, horizon area, ergoregion, geodesic frequencies, QNM spectrum (linearized operator around the solution), and shadow data; compare to Kerr limits and discuss deviations.

(XV) Uniqueness, regularity, and sector selection. Establish local uniqueness in the chosen function space; analyze regularity at the axis and ring; record how sector labels $(n, \{\theta_j\})$ select discrete families of solutions (topological superselection).

From connection to solution. Concretely, we begin by computing the *Relational Christoffel symbols* for $\bar{\Sigma}_{\mu\nu}$ (Definition 6.7), then the *Unified Ricci tensor/scalar* (Definition 11.103), and assemble the *Unified Vacuum Field Equation* (Definition 16.6) into a closed PDE system. The source is provided by the *rotationally dressed vacuum energy density* (Definition 16.5), fixed by the near-horizon KMS data (eq. (16.290)) and asymptotics to Kerr [7, 146]. Solving this system under the boundary/regularity conditions of Steps VI–VII yields the sought *Chronogeometric Kerr Solution*—a synthesis of geometric rotation and quantum–relational phase dynamics, consistent with black-hole thermodynamics [86, 7].

Step I: Calculating the Relational Christoffel Symbols

With the **Chronogeometric Kerr Ansatz** in hand (Section 16.4), our first computational task is to determine the affine structure governed by the unified tensor

$$\bar{\Sigma}_{\mu\nu} \;=\; \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}, \tag{16.320}$$

(cf. Definition 5.26). Throughout this section we work in geometric units ($G = \hbar = c = k_B = 1$). The Levi–Civita connection of $\bar{\Sigma}_{\mu\nu}$—the *Relational Levi–Civita derivative*—is

$$\boxed{\Gamma^{\lambda}{}_{\mu\nu}[\bar{\Sigma}] := \frac{1}{2}\,\bar{\Sigma}^{\lambda\rho}\Big(\partial_\mu \bar{\Sigma}_{\nu\rho} + \partial_\nu \bar{\Sigma}_{\mu\rho} - \partial_\rho \bar{\Sigma}_{\mu\nu}\Big).} \tag{16.321}$$

It is often convenient to separate the connection into a classical (geometric) part and a temporal/relational correction,

$$\Gamma^{\lambda}{}_{\mu\nu}[\bar{\Sigma}] = \Gamma^{\lambda}{}_{\mu\nu}[\bar{g}] + \Delta\Gamma^{\lambda}{}_{\mu\nu}[\bar{t}], \qquad \Delta\Gamma^{\lambda}{}_{\mu\nu} := \Gamma^{\lambda}{}_{\mu\nu}[\bar{\Sigma}] - \Gamma^{\lambda}{}_{\mu\nu}[\bar{g}]. \tag{16.322}$$

Exactly, $\Delta\Gamma$ is nonlinear in $\bar{t}_{\mu\nu}$ via $\bar{\Sigma}^{-1}$; to leading order in $\bar{t}$ one has the standard difference formula

$$\Delta\Gamma^{\lambda}{}_{\mu\nu} = \frac{1}{2}\,\bar{g}^{\lambda\rho}\Big(\nabla^{(\bar{g})}_{\mu}\bar{t}_{\nu\rho} + \nabla^{(\bar{g})}_{\nu}\bar{t}_{\mu\rho} - \nabla^{(\bar{g})}_{\rho}\bar{t}_{\mu\nu}\Big) \;+\; O(\bar{t}^{\,2}), \tag{16.323}$$

where $\nabla^{(\bar{g})}$ is the Levi–Civita derivative of $\bar{g}_{\mu\nu}$.

1. Ansatz components and notation

We use the stationary, axisymmetric ansatz of Section 16.4 in Boyer–Lindquist–type coordinates (t, r, θ, ϕ):

$$\bar{g}_{\mu\nu} \text{ as in } \textit{eq.}\ (16.200), \qquad \bar{t}_{\mu\nu} \text{ given by the Temporal Tensor Ansatz of } \textit{Definition } 16.4, \tag{16.324}$$

with scalar functions $\{A(r,\theta), B(r,\theta), C(r,\theta), D(r,\theta), E(r,\theta)\}$. We denote partial derivatives by subscripts, e.g. $F_r := \partial_r F$, and set $\rho^2 := r^2 + a^2\cos^2\theta$, $\Delta := r^2 - 2Mr + a^2$.

2. Representative Christoffel components

Below we exhibit three illustrative coefficients. In all cases, indices are raised by the *full* inverse $\bar{\Sigma}^{\mu\nu}$; when the (r,θ) block is diagonal (as in our ansatz), $\bar{\Sigma}^{rr} = 1/\bar{\Sigma}_{rr}$ and $\bar{\Sigma}^{r\theta} = 0$.

(i) $\Gamma^{r}{}_{tt}$: radial acceleration Stationarity and the absence of tr cross terms imply

$$\Gamma^{r}{}_{tt} = -\frac{1}{2}\,\bar{\Sigma}^{rr}\,\partial_r\bar{\Sigma}_{tt}. \tag{16.325}$$

With $\bar{\Sigma}_{tt} = -\Big(1 - \frac{f(r,\theta)}{\rho^2}\Big) - A(r,\theta)$, we find

$$\boxed{\Gamma^{r}{}_{tt} = -\frac{1}{2\,\bar{\Sigma}_{rr}}\Big[\frac{f_r\,\rho^2 - 2r\,f}{\rho^4} \;-\; A_r\Big].} \tag{16.326}$$

The familiar inward "geometric" pull arises from the radial gradient of f/ρ^2, while the term proportional to A_r is a genuinely relational force, sourced by the gradient of the temporal coherence field A.

(ii) $\Gamma^{\phi}{}_{tr}$: frame dragging dressed by coherence For a stationary, axisymmetric ansatz with $\bar{\Sigma}_{tr} = 0$,

$$\Gamma^{\phi}{}_{tr} = \frac{1}{2}\,\bar{\Sigma}^{\phi\rho}\,\partial_r\bar{\Sigma}_{t\rho} = \frac{1}{2}\,\bar{\Sigma}^{\phi\phi}\,\partial_r\bar{\Sigma}_{t\phi} + \frac{1}{2}\,\bar{\Sigma}^{\phi t}\,\partial_r\bar{\Sigma}_{tt}. \tag{16.327}$$

Using the unified ansatz (eq. (16.411) and Definition 16.4), $\bar{\Sigma}_{t\phi} = \bar{g}_{t\phi} + \bar{t}_{t\phi} = -\frac{2Mar}{\rho^2}\sin^2\theta + D(r,\theta)$, and $\bar{\Sigma}_{\phi\phi} \approx r^2\sin^2\theta \Rightarrow \bar{\Sigma}^{\phi\phi} \approx 1/(r^2\sin^2\theta)$ away from the axis, one obtains

$$\partial_r \bar{\Sigma}_{t\phi} = \left(\frac{2Ma(\rho^2 - r\,\partial_r\rho^2)}{\rho^4}\right)\sin^2\theta \;+\; D_r = \frac{2Ma}{r^2}\sin^2\theta \;+\; O\!\left(\frac{aM}{r^3}\right) \;+\; D_r, \tag{16.328}$$

so that, keeping the leading pieces and noting that $\bar{\Sigma}^{\phi t} = O(a/r^3)$ is subleading,

$$\boxed{\Gamma^{\phi}{}_{tr} \;\approx\; \frac{Ma}{r^4} \;+\; \frac{1}{2\,r^2\sin^2\theta}\,D_r \quad \Big(\text{leading frame dragging + relational dressing}\Big),} \tag{16.329}$$

which recovers the classical Lense–Thirring behavior and exhibits its explicit quantum–relational correction via D_r [146, 7].

(iii) $\Gamma^{r}{}_{r\theta}$: radial–polar cross curvature For the (r,θ) block (diagonal in our ansatz),

$$\Gamma^{r}{}_{r\theta} = \frac{1}{2}\,\bar{\Sigma}^{rr}\Big(\partial_\theta\bar{\Sigma}_{rr} + \partial_r\bar{\Sigma}_{\theta\theta}\Big) = \frac{1}{2\,\bar{\Sigma}_{rr}}\Big(\partial_\theta\bar{\Sigma}_{rr} + \partial_r\bar{\Sigma}_{\theta\theta}\Big). \tag{16.330}$$

Writing the derivatives term-by-term using Section 16.4 and Definition 16.4 gives the schematic decomposition

$$\boxed{\Gamma^{r}{}_{r\theta} \;\approx\; \frac{1}{2\,\bar{\Sigma}_{rr}}\Big[\underbrace{-\frac{2a^2\sin\theta\cos\theta}{\Delta}}_{\text{rotational anisotropy}} \;+\; \underbrace{2r}_{\text{radial geometry}} \;+\; \underbrace{B_\theta + C_r}_{\text{relational corrections}}\Big],} \tag{16.331}$$

where $B_\theta := \partial_\theta B$ and $C_r := \partial_r C$. The first two terms are purely geometric (Kerr) contributions; the last two encode the polar gradient of the "radial viscosity" B and the radial gradient of the polar sector C of the temporal tensor.

3. Sanity checks and forward links

- *Classical limit.* Setting $(A, B, C, D, E) \equiv 0$ reproduces the Boyer–Lindquist Kerr connection coefficients (e.g. $\Gamma^{\phi}{}_{tr} \sim Ma/r^4$), as required [146, 7].
- *Near-horizon regularity.* In ingoing horizon-regular coordinates, each displayed coefficient is finite at $\mathcal{H}_+$, with the KMS data fixing the leading behavior (eqs. (16.290) and (16.319)).
- *Index hygiene.* All raising/lowering use the full $\bar{\Sigma}^{\mu\nu}$; where off-diagonal corrections are retained, $\bar{\Sigma}^{rr} \neq 1/\bar{\Sigma}_{rr}$ and the corresponding formulas should be evaluated with the exact inverse.

Outcome and next step. Equations (16.326), (16.329), and (16.331) exemplify how the temporal tensor dresses the affine structure. With $\Gamma^{\lambda}{}_{\mu\nu}[\bar{\Sigma}]$ determined, we now compute the *Unified Ricci tensor and scalar* (Definition 11.103) to furnish the curvature entering the unified field equations in Step II.

Step II: Calculating the Unified Ricci Tensor $\bar{R}_{\mu\nu}[\bar{\Sigma}]$

Having determined the connection coefficients (Section 16.5), we now compute the curvature of the unified chrono–geometric manifold. The **Unified Ricci tensor** $\bar{R}_{\mu\nu}[\bar{\Sigma}]$ (cf. Definition 11.103) is

$$\boxed{\bar{R}_{\mu\nu}[\bar{\Sigma}] = \partial_\lambda \Gamma^\lambda{}_{\mu\nu} - \partial_\nu \Gamma^\lambda{}_{\mu\lambda} + \Gamma^\lambda{}_{\lambda\rho}\Gamma^\rho{}_{\mu\nu} - \Gamma^\lambda{}_{\nu\rho}\Gamma^\rho{}_{\mu\lambda},} \tag{16.332}$$

where $\Gamma^\lambda{}_{\mu\nu}[\bar{\Sigma}]$ is given by eq. (16.321). We work with the unified split $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (eq. (16.320)) in the stationary, axisymmetric ansatz of Section 16.4.

A. Linearized structure about the Kerr background

Because the classical Kerr metric is Ricci–flat, $\bar{R}_{\mu\nu}[\bar{g}] \equiv 0$ [146, 7], the leading curvature of the unified tensor is controlled by the variation of the Ricci tensor with respect to the temporal perturbation $\bar{t}_{\mu\nu}$:

$$\bar{R}_{\mu\nu}[\bar{\Sigma}] = \delta\bar{R}_{\mu\nu}[\bar{t}] \; + \; O(\bar{t}^{\,2}). \tag{16.333}$$

Writing $h_{\mu\nu} := \bar{t}_{\mu\nu}$ and $h := \bar{g}^{\alpha\beta}h_{\alpha\beta}$, the standard variation on a Ricci–flat background gives (all covariant derivatives with respect to $\bar{g}$)

$$\boxed{\delta\bar{R}_{\mu\nu}[h] = \frac{1}{2}\Big(\nabla^\rho\nabla_\mu h_{\nu\rho} + \nabla^\rho\nabla_\nu h_{\mu\rho} - \nabla^2 h_{\mu\nu} - \nabla_\mu\nabla_\nu h\Big) - \bar{R}_{\mu\rho\nu\sigma}\, h^{\rho\sigma},} \tag{16.334}$$

where $\bar{R}_{\mu\rho\nu\sigma}$ is the Kerr Riemann tensor. The last (curvature–coupling) term is nonzero in the strong–field region and will be grouped into the lower–derivative remainder terms below.

Stationarity/axisymmetry simplifications. Because $\partial_t(\cdot) = \partial_\phi(\cdot) = 0$ for our ansatz, time and azimuthal derivatives drop out of eq. (16.334). Moreover, all occurrences of the connection may be expressed using the Christoffel symbols computed in Step I (Section 16.5) with index manipulations performed by the *full* inverse $\bar{\Sigma}^{\mu\nu}$ (see the "Index hygiene'' note in Step I).

B. Representative components of $\delta\bar{R}_{\mu\nu}$

To exhibit the structure, we keep the leading second–derivative pieces explicitly and collect all first–derivative & curvature–coupling terms into remainders $\mathcal{F}_{\mu\nu}$ that depend locally on (A, B, C, D, E) and their first derivatives, as well as on the Kerr background coefficients.

1) Time–time component $\bar{R}_{tt}$ With $h_{tt} = -A(r,\theta)$ in our conventions (Definition 16.4), the second–derivative contribution of eq. (16.334) combines into the (meridional) Laplace–Beltrami operator acting on A:

$$\boxed{\bar{R}_{tt}[\bar{\Sigma}] \;=\; -\frac{1}{2}\,\nabla^2 A(r,\theta) \;+\; \mathcal{F}_{tt}[A, B, C, D, E; \bar{g}],} \tag{16.335}$$

where $\nabla^2 := \bar{g}^{ab}\nabla_a\nabla_b$ on the stationary background (here effectively the (r,θ) sector). The remainder $\mathcal{F}_{tt}$ collects curvature–couplings $-\bar{R}_{t\rho t\sigma}h^{\rho\sigma}$ and all first–derivative interaction terms (e.g. $A_r f_r$, $B_r A_r$, etc.).

2) Radial–radial component $\bar{R}_{rr}$ Analogously, with $h_{rr} = B(r,\theta)$,

$$\boxed{\bar{R}_{rr}[\bar{\Sigma}] \;=\; -\frac{1}{2}\,\nabla^2 B(r,\theta) \;+\; \mathcal{F}_{rr}[A,B,C,D,E;\bar{g}],} \tag{16.336}$$

where $\mathcal{F}_{rr}$ contains mixed first–derivative products and the background Riemann–coupling $-\bar{R}_{r\rho r\sigma}h^{\rho\sigma}$.

3) Time–azimuthal component $\bar{R}_{t\phi}$ For the off–diagonal (frame–dragging) sector, $h_{t\phi} = D(r,\theta)$ yields

$$\boxed{\bar{R}_{t\phi}[\bar{\Sigma}] \;=\; -\frac{1}{2}\,\nabla^2 D(r,\theta) \;+\; \mathcal{F}_{t\phi}[A,B,C,D,E;\bar{g}],} \tag{16.337}$$

again with $\mathcal{F}_{t\phi}$ absorbing lower–order and curvature–coupled terms. This is the channel that dresses the classical Lense–Thirring sector with relational corrections (cf. Step I, eq. (16.329)).

Remark 16.58 (On the Laplacian and background curvature)**.** The operator ∇^2 above is the covariant Laplacian with respect to the *background* Kerr connection $\nabla^{(\bar{g})}$. In regions of strong curvature, the background Riemann term in eq. (16.334) acts as an effective potential for the tensor modes $h_{\mu\nu}$; we consistently include it in the remainders $\mathcal{F}_{\mu\nu}$ to keep the leading second–derivative structure transparent. A fully expanded form can be produced on demand, but is notationally heavy and unnecessary at this stage.

C. Physical interpretation and forward use

Equations (16.335)–(16.337) show that the unified curvature is governed, at leading order, by meridional Laplacians of the temporal fields:

(i) $\bar{R}_{tt}$ is driven chiefly by the spatial stiffness (Laplacian) of the temporal potential $A(r,\theta)$, controlling redshift/time–dilation corrections.

(ii) $\bar{R}_{rr}$ is controlled by the "radial viscosity'' $B(r,\theta)$, shaping radial geodesic deviation.

(iii) $\bar{R}_{t\phi}$ is controlled by the frame–dragging correction $D(r,\theta)$, encoding relational twists of the co–rotating vacuum.

Together with the remaining components (obtained analogously), these supply the left–hand side of the **Unified Vacuum Field Equation** (Definition 16.6). In **Step III** we will equate $\bar{R}_{\mu\nu}[\bar{\Sigma}]$ to the rotationally dressed vacuum source (Definition 16.5) and extract the coupled PDE system for the unknown fields $\{A,B,C,D,E\}$ under the boundary data of eq. (16.319).

Step III: The Unified Vacuum Field Equations for Rotating Black Holes

With the unified connection and Ricci components in hand (Section 16.5), we now write the field equations that determine the chrono–geometric corrections. In the IQR framework the

classical field is the expectation value of operator geometry (Definition 16.7):

$$\bar{\Sigma}^{\mathrm{Kerr}}_{\mu\nu}(x) = \Big\langle \overset{\circ}{\boldsymbol{\Psi}} \Big| \, \hat{g}^{\mathrm{Kerr}}_{\mu\nu}(x) + \hat{t}^{\mathrm{Kerr}}_{\mu\nu}(x) \, \Big| \overset{\circ}{\boldsymbol{\Psi}} \Big\rangle, \tag{16.338}$$

and satisfies the *Chronogeometric Einstein equation* (Definition 11.106)

$$\bar{G}_{\mu\nu}[\bar{\Sigma}] := \bar{R}_{\mu\nu}[\bar{\Sigma}] - \tfrac{1}{2}\bar{\Sigma}_{\mu\nu}\,\bar{R}[\bar{\Sigma}] = 0, \tag{16.339}$$

for the exterior vacuum region.[10]

A. Linearization about Kerr and trace reversal

Write the unified tensor as

$$\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + h_{\mu\nu}, \qquad h_{\mu\nu} \equiv \bar{t}_{\mu\nu}, \tag{16.340}$$

with $\bar{g}_{\mu\nu}$ the Kerr background (Section 16.4). Since $\bar{R}_{\mu\nu}[\bar{g}] = 0$ [146, 7], the linearized Einstein tensor is

$$\delta\bar{G}_{\mu\nu}[h] = \delta\bar{R}_{\mu\nu}[h] - \tfrac{1}{2}\bar{g}_{\mu\nu}\,\bar{g}^{\rho\sigma}\delta\bar{R}_{\rho\sigma}[h], \tag{16.341}$$

and the vacuum equations reduce to

$$\boxed{\delta\bar{G}_{\mu\nu}[h] = 0.} \tag{16.342}$$

Introduce the trace–reversed field

$$\bar{h}_{\mu\nu} := h_{\mu\nu} - \tfrac{1}{2}\bar{g}_{\mu\nu}\,h, \qquad h := \bar{g}^{\alpha\beta}h_{\alpha\beta}, \tag{16.343}$$

and impose the harmonic (de Donder) gauge with respect to $\bar{g}$,

$$\boxed{\nabla^{\mu}\bar{h}_{\mu\nu} = 0,} \tag{16.344}$$

where ∇ is the Levi–Civita connection of $\bar{g}$. Then (16.342) becomes the Lichnerowicz equation on the Kerr background [7]

$$\boxed{\mathcal{E}_{\mu\nu}[\bar{h}] := -\tfrac{1}{2}\Big(\Box\,\bar{h}_{\mu\nu} + 2\,\bar{R}_{\mu}{}^{\rho}{}_{\nu}{}^{\sigma}\,\bar{h}_{\rho\sigma}\Big) = 0,} \tag{16.345}$$

with $\Box := \bar{g}^{\alpha\beta}\nabla_{\alpha}\nabla_{\beta}$. In the stationary, axisymmetric sector ($\partial_t = \partial_\phi = 0$), $\Box$ reduces to a meridional Laplace–Beltrami operator plus lower–order terms in (r, θ).

[10] The unified formulation allowing a rotationally dressed vacuum source is stated in Definition 16.6. In this step we work in the source–free exterior; the dressed source will be reincorporated in Step V (Definition 16.5).

B. Mapping to the temporal–tensor ansatz

In our ansatz (Definition 16.4),

$$h_{tt} = -A(r,\theta), \quad h_{rr} = B(r,\theta), \quad h_{\theta\theta} = C(r,\theta), \quad h_{t\phi} = D(r,\theta), \quad h_{\phi\phi} = E(r,\theta),$$

with all other components determined by symmetry (and $h_{r\theta} = 0$ in our gauge choice for Step I). The harmonic gauge (16.344) yields three independent constraints (the t, r, θ components; the ϕ–component is redundant by axisymmetry), which we denote collectively as

$$\boxed{\mathcal{C}_\nu[A, B, C, D, E] = 0, \qquad \nu \in \{t, r, \theta\}.} \tag{16.346}$$

Substituting the ansatz into (16.345) produces five coupled, second–order linear PDEs on (r, θ):

$$\boxed{\mathcal{D}_{\mu\nu}[A, B, C, D, E] = 0, \qquad (\mu\nu) \in \{tt, rr, \theta\theta, t\phi, \phi\phi\},} \tag{16.347}$$

where each $\mathcal{D}_{\mu\nu}$ is a Kerr–background differential operator (principal part a meridional Laplacian) plus a curvature potential from the $\bar{R}_\mu{}^\rho{}_\nu{}^\sigma$ term.

Schematic principal structure. Keeping only the second–derivative part and grouping all first–derivative and curvature–couplings into remainders $\mathcal{F}_{\mu\nu}$, one obtains (cf. Step II)

$$\mathcal{D}_{tt}: \quad -\tfrac{1}{2}\nabla^2 A + \mathcal{F}_{tt}[A, B, C, D, E] = 0, \tag{16.348}$$
$$\mathcal{D}_{rr}: \quad -\tfrac{1}{2}\nabla^2 B + \mathcal{F}_{rr}[A, B, C, D, E] = 0, \tag{16.349}$$
$$\mathcal{D}_{\theta\theta}: \quad -\tfrac{1}{2}\nabla^2 C + \mathcal{F}_{\theta\theta}[A, B, C, D, E] = 0, \tag{16.350}$$
$$\mathcal{D}_{t\phi}: \quad -\tfrac{1}{2}\nabla^2 D + \mathcal{F}_{t\phi}[A, B, C, D, E] = 0, \tag{16.351}$$
$$\mathcal{D}_{\phi\phi}: \quad -\tfrac{1}{2}\nabla^2 E + \mathcal{F}_{\phi\phi}[A, B, C, D, E] = 0, \tag{16.352}$$

where ∇^2 is the (r, θ) Laplace–Beltrami operator of the Kerr background and the $\mathcal{F}_{\mu\nu}$ encode mixing among $\{A, \ldots, E\}$ and the background curvature potential $2\bar{R}_\mu{}^\rho{}_\nu{}^\sigma$.

C. Boundary data and constraints

The boundary–value problem is posed on the exterior domain $r \geq r_+$ with:

(i) Horizon (KMS/regularity). $\bar{\Sigma}_{\mu\nu}$ is smooth in ingoing coordinates at $\mathcal{H}_+$, and

$$\begin{aligned} T_{\text{Kerr}} &= \kappa/(2\pi), \\ \chi^a \chi_a\big|_{\mathcal{H}_+} &= 0. \end{aligned} \tag{16.353}$$

These encode the KMS boundary data and surface–gravity relation (eqs. (16.290) and (16.319)).

(ii) Asymptotics. As $r \to \infty$,

$$\bar{\Sigma}_{\mu\nu} \to \bar{g}^{\text{Kerr}}_{\mu\nu}, \qquad \text{with fixed ADM charges } (M, a). \tag{16.354}$$

(iii) Axis regularity. Along the symmetry axis,

$$\partial_\theta X\Big|_{\theta=0,\pi} = 0, \quad X \in \{A, B, C, D, E\}, \qquad \text{and all components remain finite.} \tag{16.355}$$

(iv) Gauge. The gauge constraints

$$\mathcal{C}_\nu[A, B, C, D, E] = 0, \quad \nu \in \{t, r, \theta\}, \tag{16.356}$$

are imposed on the boundary and propagated into the bulk by eq. (16.347) via the linearized Bianchi identities.

D. Summary and computational outlook

Equations (16.347) together with the gauge constraints (16.346) and boundary data (16.353)–(16.356) constitute a well–posed stationary boundary–value problem on (r, θ). The principal symbol is elliptic (meridional Laplace–Beltrami), while the Kerr curvature supplies a bounded potential/mixing term. In **Step IV** we will develop the near–horizon and asymptotic expansions that solve (16.348)–(16.352) order by order (anchored to eq. (16.319)), and in **Step V** we reinstate the rotationally dressed vacuum source to obtain the fully self–consistent *Chronogeometric Kerr Solution*.

Step IV: Perturbative Expansion as a Semi–Classical Limit

To solve the linearized, stationary boundary–value problem obtained in Section 16.5, we adopt a controlled perturbative scheme that respects the discrete (topological) origin of the spin parameter (eq. (16.267)). In the IQR framework the Kerr spin is a sector label a_n; the *semi–classical limit* corresponds to large winding $n \gg 1$ with $a_{n+1} - a_n \ll R_{EH}^{(0)}$, so that a may be treated as an effectively continuous parameter representing the large–n envelope.

Dimensionless expansion parameter. We work in geometric units ($G = \hbar = c = k_B = 1$) and define

$$\boxed{\epsilon := \frac{a}{R_{EH}^{(0)}} \ll 1,} \tag{16.357}$$

where $R_{EH}^{(0)}$ is the event–horizon radius of the *static* (non–rotating) chronogeometric solution in the n=0 sector. All fields are expanded in powers of ϵ, with the static solution providing the zeroth–order data.

A. Expansion of the geometric sector

We expand the geometric part of the unified tensor about the static, quantum–corrected background (not Schwarzschild), cf. Section 16.1:

$$\boxed{\bar{g}_{\mu\nu}(r, \theta; a) = \bar{g}_{\mu\nu}^{(0)}(r) + \epsilon\, \bar{g}_{\mu\nu}^{(1)}(r, \theta) + O(\epsilon^2).} \tag{16.358}$$

Zeroth order. The static line element has the form

$$\bar{g}^{(0)}_{\mu\nu}dx^\mu dx^\nu = -\Big(1-\frac{f_0(r)}{r}\Big)dt^2 + \frac{dr^2}{1-\frac{f_0(r)}{r}} + r^2\,(d\theta^2+\sin^2\theta\,d\phi^2), \tag{16.359}$$

with corrected radial function $f_0(r)$ given in eq. (16.159). This encodes the zeroth–order (spherical) chronogeometric vacuum.

First order. To $O(\epsilon)$ the only geometric component linear in a is the $t\phi$–entry (slow–rotation Hartle–Thorne structure [659]):

$$\boxed{\bar{g}^{(1)}_{t\phi}(r,\theta) = -\frac{2Ma}{r}\sin^2\theta, \qquad \bar{g}^{(1)}_{\mu\nu} = 0 \text{ for } (\mu\nu)\neq(t\phi) \text{ at } O(\epsilon).} \tag{16.360}$$

All other geometric corrections begin at $O(\epsilon^2)$ (e.g. quadratic spin–flattening).

B. Expansion of the temporal tensor

For the temporal (relational) sector $\bar{t}_{\mu\nu}$ we expand the five ansatz functions (Definition 16.4) as

$$A(r,\theta) = A_0(r) + \epsilon\,A_1(r,\theta) + O(\epsilon^2), \tag{16.361}$$
$$B(r,\theta) = B_0(r) + \epsilon\,B_1(r,\theta) + O(\epsilon^2), \tag{16.362}$$
$$C(r,\theta) = C_0(r) + \epsilon\,C_1(r,\theta) + O(\epsilon^2), \tag{16.363}$$
$$D(r,\theta) = \epsilon\,D_1(r,\theta) + O(\epsilon^2), \tag{16.364}$$
$$E(r,\theta) = E_0(r) + \epsilon\,E_1(r,\theta) + O(\epsilon^2). \tag{16.365}$$

Here $\{A_0, B_0, C_0, E_0\}$ are the known static solutions (spherical symmetry $\Rightarrow$ no θ–dependence), while $D_0 \equiv 0$ by symmetry. The *unknowns* at first order are $\{A_1, B_1, C_1, D_1, E_1\}$.

C. Hierarchy of perturbed field equations

Insert (16.358)–(16.365) into the linearized equations $\delta\bar{G}_{\mu\nu}[h] = 0$ (eq. (16.342)) together with the harmonic gauge constraints (eqs. (16.344) and (16.346)). Collecting orders in ϵ yields:

Zeroth order $\mathcal{O}(\epsilon^0)$: The equations reduce to the static problem and are satisfied by $\bar{g}^{(0)}_{\mu\nu}$ and $\{A_0, B_0, C_0, E_0\}$ by construction (consistency check with Section 16.1).

First order $\mathcal{O}(\epsilon^1)$: A linear, coupled system for $\{A_1, B_1, C_1, D_1, E_1\}$:

$$\boxed{\mathcal{D}^{(0)}_{\mu\nu}\big[A_1, B_1, C_1, D_1, E_1\big] = \mathcal{S}_{\mu\nu}\big[\bar{g}^{(1)};\, A_0, B_0, C_0, E_0\big], \qquad (\mu\nu)\in\{tt, rr, \theta\theta, t\phi, \phi\phi\},} \tag{16.366}$$

supplemented by the first–order gauge constraints

$$\boxed{\mathcal{C}_\nu^{(0)}\big[A_1, B_1, C_1, D_1, E_1\big] = \mathcal{J}_\nu\big[\bar{g}^{(1)};\, A_0, B_0, C_0, E_0\big], \qquad \nu \in \{t, r, \theta\}.} \tag{16.367}$$

Here $\mathcal{D}^{(0)}_{\mu\nu}$ is the *static* Kerr–background operator (principal part a meridional Laplace–Beltrami, cf. eq. (16.348)–eq. (16.352)), and the sources $\mathcal{S}_{\mu\nu}$ (and $\mathcal{J}_\nu$) arise from the mixing between the geometric spin piece $\bar{g}^{(1)}_{t\phi}$ (16.360) and the static temporal fields $\{A_0, \dots, E_0\}$.

D. The $t\phi$ equation: quantum frame–dragging at first order

The $(t\phi)$ component isolates the leading relational correction to frame dragging. At $O(\epsilon)$ one obtains a scalar elliptic equation for D_1 on the meridional plane,

$$\boxed{\mathcal{D}^{(0)}_{t\phi}\big[D_1(r,\theta)\big] = \mathcal{S}_{t\phi}\big[\bar{g}^{(1)}_{t\phi};\, A_0(r), B_0(r), C_0(r), E_0(r)\big],} \tag{16.368}$$

with principal part $-\frac{1}{2}\nabla^2 D_1$ (cf. eq. (16.337)). A convenient solution strategy is separation in associated Legendre modes,

$$D_1(r,\theta) = \sum_{\ell \text{ odd}} d_\ell(r)\, P_\ell^{(1)}(\cos\theta), \tag{16.369}$$

since the source inherits an ℓ=1–dominated structure from $\sin^2\theta$ in (16.360). The resulting radial ODEs for $d_\ell(r)$ are second–order and regular on $[r_+, \infty)$ with horizon/asymptotic data given below.

E. Boundary conditions and regularity

The first–order system (16.366)–(16.367) is closed by:

(i) Horizon regularity (KMS). Fields are smooth in ingoing coordinates at $r = r_+$ and consistent with the KMS data (eq. (16.319)); Frobenius expansions fix one integration constant per mode by regularity.

(ii) Axis regularity. Single–valuedness at $\theta = 0, \pi$ implies standard parity/derivative conditions for each scalar (e.g. $\partial_\theta X_1|_{\theta=0,\pi} = 0$ for m=0 sectors).

(iii) Asymptotic flatness. All first–order corrections decay, $\lim_{r\to\infty} A_1 = \lim_{r\to\infty} B_1 = \cdots = \lim_{r\to\infty} E_1 = 0$ (eq. (16.354)); for D_1 one expects $D_1 = O(r^{-p})$ with $p \geq 1$.

(iv) Smooth static limit. In the vanishing–spin limit all first–order rotational corrections vanish:

$$\begin{aligned} &\lim_{a\to 0} A_1(r,\theta) = 0, \qquad \lim_{a\to 0} B_1(r,\theta) = 0, \\ &\lim_{a\to 0} C_1(r,\theta) = 0, \qquad \lim_{a\to 0} D_1(r,\theta) = 0, \\ &\lim_{a\to 0} E_1(r,\theta) = 0. \end{aligned} \tag{16.370}$$

(v) Gauge preservation. The constraints (16.367) are enforced on a Cauchy curve in (r,θ) and propagate by the linearized Bianchi identity (compatibility of (16.366)).

Outcome. This step yields a solvable, elliptic hierarchy for the first–order unknowns driven by a known source determined by the static chronogeometric vacuum and the classical $O(a)$ frame–dragging template (16.360). In **Step V** we solve (16.368) explicitly for $D_1(r,\theta)$ (quantum frame–dragging), and then back–substitute to obtain $\{A_1, B_1, C_1, E_1\}$, thereby completing the leading semi–classical rotational dressing of the Chronogeometric Kerr solution.

Step V: Equation for the Quantum Frame-Dragging Function

The unified $t\phi$ component at first order in the slow–rotation parameter ϵ follows from the perturbative setup in Section 16.5 and the linearized system in eq. (16.366):

$$\bar{\Sigma}_{t\phi}(r,\theta) = \bar{g}^{(1)}_{t\phi}(r,\theta) + \bar{t}^{(1)}_{t\phi}(r,\theta) = -\frac{2GMa}{c^3 r}\,\sin^2\theta + c\,D_1(r,\theta). \tag{16.371}$$

This component controls the local angular velocity of inertial frames, $\omega(r,\theta) \approx -c\,\bar{\Sigma}_{t\phi}/\bar{\Sigma}_{\phi\phi}$, so the first–order quantum correction is encoded in $D_1(r,\theta)$.

A. Dynamical equation for D_1

The $(t\phi)$ component of the linearized vacuum equations (see the principal structure in eq. (16.351)) yields a second–order PDE for D_1. As shown in Section 16.5, the Laplace–Beltrami part on the *static corrected background* $\bar{g}^{(0)}_{\mu\nu}$ dominates, producing:

$$\boxed{\left(1 - \frac{f_0(r)}{r}\right)\partial_r^2 D_1 + \left(\frac{2}{r} - \frac{f_0'(r)}{r}\right)\partial_r D_1 + \frac{1}{r^2\sin\theta}\,\partial_\theta\Big(\sin\theta\,\partial_\theta D_1\Big) = 2\,\mathcal{S}_{t\phi}(r,\theta),} \tag{16.372}$$

where $f_0(r)$ is the corrected static radial function from eq. (16.159). Equation (16.372) is the explicit $(t\phi)$ member of the linear system $\mathcal{D}^{(0)}_{\mu\nu}[\bar{t}^{(1)}] = \mathcal{S}_{\mu\nu}$ (eq. (16.366)).

B. Source term: structure and projection

The source $\mathcal{S}_{t\phi}$ arises from the quadratic $\Gamma\Gamma$ part of the Ricci tensor (see eq. (16.332)) at first order, i.e. mixed products $\Gamma^{(0)}\cdot\Gamma^{(1)}$ between the static corrected background and the rotational perturbation. Using the *rotationally dressed vacuum energy density* from Definition 16.5, one finds that:

(i) Radial dependence. The radial dependence traces the static vacuum profile $\rho_{\rm vac}(r)$.

(ii) Angular dependence. The angular dependence inherits the $\sin^2\theta$ from the leading classical frame-dragging sector.

Consequently the source is fixed (up to the coupling κ') by

$$\boxed{\mathcal{S}_{t\phi}(r,\theta) = \frac{\kappa'}{2}\,\rho_{\rm vac}(r)\,\sin^2\theta.} \tag{16.373}$$

For separation of variables, expand D_1 in Legendre modes and project the source. Using $\sin^2\theta = \frac{2}{3}\big(1 - P_2(\cos\theta)\big)$, only $\ell = 0, 2$ are excited. Writing

$$D_1(r,\theta) = \sum_{\ell=0}^{\infty} d_\ell(r)\, P_\ell(\cos\theta), \tag{16.374}$$

the mode sources are

$$\boxed{\begin{aligned} S_\ell(r) &:= \frac{2\ell+1}{2}\int_{-1}^{1} \mathcal{S}_{t\phi}(r,\theta)\, P_\ell(\mu)\, d\mu \\ &\Longrightarrow \quad S_0(r) = \frac{\kappa'}{3}\,\rho_{\rm vac}(r), \qquad S_2(r) = -\frac{\kappa'}{3}\,\rho_{\rm vac}(r) \end{aligned}} \tag{16.375}$$

with $S_\ell(r) = 0$ for $\ell \neq 0, 2$.

Projecting (16.372) onto P_ℓ gives the decoupled radial ODEs

$$\boxed{\mathcal{L}_r\, d_\ell(r) - \frac{\ell(\ell+1)}{r^2}\, d_\ell(r) = 2\, S_\ell(r), \qquad \mathcal{L}_r := \left(1 - \frac{f_0(r)}{r}\right)\frac{d^2}{dr^2} + \left(\frac{2}{r} - \frac{f_0'(r)}{r}\right)\frac{d}{dr}.} \tag{16.376}$$

C. Boundary conditions for D_1

The well-posed exterior problem $r \geq r_+$ requires:

(i) Axis regularity ($\theta = 0, \pi$).

$$\partial_\theta D_1(r,\theta)\big|_{\theta=0,\pi} = 0 \quad \Longleftrightarrow \quad d_\ell \text{ finite and only even } \ell \text{ survive for the } m{=}0 \text{ sector.} \tag{16.377}$$

(ii) Asymptotic flatness ($r \to \infty$). (cf. eq. (16.354))

$$D_1(r,\theta) \xrightarrow[r\to\infty]{} 0 \quad \Longleftrightarrow \quad d_\ell(r) = o(1) \text{ as } r \to \infty. \tag{16.378}$$

(iii) Smooth static limit ($a \to 0$). (cf. eq. (16.370))

$$\lim_{a\to 0} D_1(r,\theta) = 0 \quad \Longleftrightarrow \quad \lim_{a\to 0} d_\ell(r) = 0 \;\; \forall \ell. \tag{16.379}$$

(iv) Horizon regularity ($r = r_+$). Ingoing smoothness consistent with eq. (16.319):

$$D_1(r,\theta) \text{ finite at } r = r_+, \quad \text{equivalently,} \quad d_\ell(r) = d_\ell^{(+)} + O(r - r_+). \tag{16.380}$$

D. Green's–function solution for $\ell = 0, 2$ modes

The decoupled radial equations eq. (16.376) for $d_\ell(r)$ ($\ell = 0, 2$) can be solved in closed form using a Green's–function method. Define the linear differential operator

$$\mathcal{L}_r^{(\ell)}[d] := \left(1 - \frac{f_0(r)}{r}\right)\frac{d^2 d}{dr^2} + \left(\frac{2}{r} - \frac{f_0'(r)}{r}\right)\frac{dd}{dr} - \frac{\ell(\ell+1)}{r^2}\, d. \tag{16.381}$$

Then eq. (16.376) reads

$$\mathcal{L}_r^{(\ell)}[d_\ell(r)] = 2\, S_\ell(r), \qquad \ell = 0, 2. \tag{16.382}$$

1. Homogeneous solutions. Let $u_\ell(r)$ and $v_\ell(r)$ be two independent solutions of the homogeneous equation $\mathcal{L}_r^{(\ell)}[d] = 0$. We normalize them by boundary behaviour:

(i) $u_\ell(r)$ is *ingoing–regular* at the horizon $r = r_+$.

(ii) $v_\ell(r)$ is *decaying* at spatial infinity $r \to \infty$.

The Wronskian of this pair,

$$W_\ell(r) := u_\ell(r)\, v_\ell'(r) - u_\ell'(r)\, v_\ell(r), \tag{16.383}$$

is independent of r up to the coefficient $(1 - f_0(r)/r)$ in eq. (16.381).

2. Radial Green's function. The Green's kernel $G_\ell(r, r')$ is defined by

$$\mathcal{L}_r^{(\ell)}[G_\ell(r, r')] = \delta(r - r'), \tag{16.384}$$

with boundary conditions inherited from u_ℓ, v_ℓ. Explicitly,

$$G_\ell(r, r') = \frac{1}{(1 - f_0(r')/r')\, W_\ell} \begin{cases} u_\ell(r)\, v_\ell(r'), & r < r', \\ u_\ell(r')\, v_\ell(r), & r > r'. \end{cases} \tag{16.385}$$

3. Integral solution. The mode functions $d_\ell(r)$ are then expressed as

$$d_\ell(r) = 2 \int_{r_+}^{\infty} G_\ell(r, r')\, S_\ell(r')\, dr', \qquad \ell = 0, 2. \tag{16.386}$$

By construction this solution satisfies the horizon regularity eq. (16.380), the asymptotic decay eq. (16.378), and the smooth static limit eq. (16.379).

4. Physical meaning. Equation eq. (16.386) makes manifest that the quantum frame–dragging correction is entirely sourced by the static vacuum profile $\rho_{\text{vac}}(r)$ (eq. (16.373)). The angular structure is restricted to $\ell = 0, 2$, while the radial profile is encoded in the response kernel $G_\ell(r, r')$, which embodies the propagation of the vacuum's rotational disturbance from radius r' to r under the corrected static geometry $f_0(r)$.

E. Interpretation and roadmap

(i) The operator $\mathcal{L}_r$ in eq. (16.376) is the radial part of the Laplace–Beltrami operator on the corrected static background; the angular term contributes $-\ell(\ell+1)/r^2$.

(ii) Because S_ℓ is **nonzero only for** $\ell = 0, 2$ (eq. (16.375)), the quantum frame-dragging correction is dominated by two modes $d_0(r)$ and $d_2(r)$.

(iii) Once d_0, d_2 are solved from eq. (16.376) subject to eqs. (16.378) to (16.380), the corrected angular velocity follows from eq. (16.371) and the background $\bar{\Sigma}_{\phi\phi}$, and the ergosurface shift can be read off against Definition 16.13.

Next step (separation solution). Solve eq. (16.376) for $\ell = 0, 2$ using Green–function methods or direct integration once $f_0(r)$ (from eq. (16.159)) and $\rho_{\rm vac}(r)$ (from Definition 16.5) are specified.

Step VI: Multipolar Expansion and Mode Equations

We now turn to solving the sourced, linear PDE for the quantum frame–dragging function $D_1(r, \theta)$ obtained in Section 16.5:

$$\nabla^2_{(0)} D_1(r, \theta) \;=\; 2\,\mathcal{S}_{t\phi}(r, \theta), \tag{16.387}$$

where $\nabla^2_{(0)}$ is the Laplace–Beltrami operator on the corrected static background. The most natural method in an axisymmetric setting is separation of variables via a multipolar expansion.

A. Legendre expansion and mode selection

We expand D_1 into Legendre modes:

$$D_1(r, \theta) = \sum_{\ell=0}^{\infty} d_\ell(r)\, P_\ell(\cos\theta). \tag{16.388}$$

The source term (Section 16.5) has angular dependence $\sin^2\theta$. Using

$$\sin^2\theta = \tfrac{2}{3} P_0(\cos\theta) - \tfrac{2}{3} P_2(\cos\theta), \tag{16.389}$$

we see that only $\ell = 0, 2$ modes are sourced. Orthogonality of the Legendre polynomials ensures that all other d_ℓ vanish under the boundary conditions of Section 16.5.

The monopole correction ($\ell = 0$) may be absorbed into the static background (e.g. a redefinition of $A_0(r)$), so the physically significant correction is carried entirely by the quadrupole mode $\ell = 2$.

B. Radial mode equation

Inserting eq. (16.388) into eq. (16.387), and using the eigenvalue relation

$$\frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\left(\sin\theta \frac{\partial P_\ell(\cos\theta)}{\partial\theta}\right) = -\ell(\ell+1)\, P_\ell(\cos\theta), \tag{16.390}$$

we obtain a decoupled ODE for each $d_\ell(r)$:

$$\left(1 - \tfrac{f_0(r)}{r}\right) d_\ell''(r) + \left(\tfrac{2}{r} - \tfrac{f_0'(r)}{r}\right) d_\ell'(r) - \frac{\ell(\ell+1)}{r^2} d_\ell(r) = 2\, S_\ell(r), \tag{16.391}$$

with $S_\ell(r)$ the Legendre projection of the source. By eq. (16.375), the only nonvanishing projections are

$$S_0(r) = \tfrac{\kappa'}{3}\rho_{\rm vac}(r), \qquad S_2(r) = -\tfrac{\kappa'}{3}\rho_{\rm vac}(r). \tag{16.392}$$

C. Quadrupole equation for $d_2(r)$

Discarding the trivial monopole, the physically relevant mode is $\ell = 2$. Specializing eq. (16.391) yields:

$$\boxed{\left(1 - \tfrac{f_0(r)}{r}\right) d_2''(r) + \left(\tfrac{2}{r} - \tfrac{f_0'(r)}{r}\right) d_2'(r) - \tfrac{6}{r^2}\, d_2(r) = 2\, S_2(r) = -\tfrac{2\kappa'}{3}\, \rho_{\rm vac}(r).} \tag{16.393}$$

Here:

(i) $f_0(r)$ is the corrected static radial function (eq. (16.159)).

(ii) The angular eigenvalue $\ell(\ell+1) = 6$ encodes the quadrupolar character.

(iii) The source is the quadrupole projection of the vacuum profile.

Equation (16.393) is the definitive radial equation governing the leading–order quantum correction to the frame–dragging effect.

Next step: solving the quadrupole equation

We must now solve eq. (16.393) for $d_2(r)$, subject to the boundary conditions established in Section 16.5. Two avenues are available:

1. an analytic perturbative solution for $d_2(r)$ in terms of simple trial forms of $\rho_{\rm vac}(r)$,
2. or a numerical integration (shooting or spectral methods) using the Green's representation of Section 16.5.

In either approach, the result furnishes the explicit radial profile of the quantum correction to frame–dragging, completing the first step toward the full Chronogeometric Kerr Solution.

Step VII: Perturbative Solution for the Quadrupole Mode

We now solve the definitive ODE for the quadrupole correction $d_2(r)$ (eq. (16.393)):

$$\left(1 - \tfrac{f_0(r)}{r}\right) d_2''(r) + \left(\tfrac{2}{r} - \tfrac{f_0'(r)}{r}\right) d_2'(r) - \tfrac{6}{r^2}\, d_2(r) = 2\, S_2(r), \qquad S_2(r) = -\tfrac{\kappa'}{3}\, \rho_{\rm vac}(r), \tag{16.394}$$

where $f_0(r)$ is the corrected static radial function (eq. (16.159)).

1. Nature of the physical solution

The temporal tensor components $\{A, B, C, D, E\}$ are purely quantum–relational corrections, vanishing in the classical ($\rho_{\rm vac} = 0$) limit. Thus $d_2(r)$ must be sourced entirely by the vacuum.

The homogeneous equation associated with eq. (16.394),

$$\left(1 - \tfrac{f_0(r)}{r}\right) d_2^{(h)\prime\prime} + \left(\tfrac{2}{r} - \tfrac{f_0'(r)}{r}\right) d_2^{(h)\prime} - \tfrac{6}{r^2} d_2^{(h)} = 0,$$

has solutions that either diverge or approach a constant at infinity. The boundary condition of *asymptotic flatness* (Section 16.5) requires $d_2(r) \to 0$ as $r \to \infty$, hence the homogeneous part is suppressed: $d_2^{(h)} = 0$.

Therefore the physical solution is *entirely the particular solution*, driven by the source:

$$d_2(r) = d_2^{(p)}(r). \tag{16.395}$$

2. Source-driven solution and asymptotics

Formally the particular solution is given by convolution with the Green's function $G(r, r')$ (Section 16.5):

$$d_2(r) = 2 \int_{R_{EH}^{(0)}}^{\infty} G(r, r')\, S_2(r')\, dr'. \tag{16.396}$$

The properties of $d_2(r)$ follow from the structure of $S_2(r)$ and $G(r, r')$:

(i) The source $S_2(r) \propto \rho_{\text{vac}}(r)$ is sharply peaked at $r = R_{EH}^{(0)}$ and decays exponentially for $r > R_{EH}^{(0)}$ (Section 16.2.2).

(ii) The radial Green's kernel inherits a power–law tail $G(r, r') \sim r^{-3}$ for large r (reflecting the homogeneous $\ell = 2$ behaviour).

Thus the solution inherits both behaviours:

$$d_2(r) \;\sim\; \epsilon_d \left(\frac{R_{EH}^{(0)}}{r} \right)^k e^{-(r - R_{EH}^{(0)})/\lambda}, \tag{16.397}$$

with parameters:

(i) ϵ_d: amplitude $\sim \kappa' \int \rho_{\text{vac}}(r)\, dr$, the integrated strength of the vacuum source;

(ii) λ: coherence scale of the vacuum (decay length of ρ_{vac});

(iii) $k > 0$: power–law exponent from the Green's kernel ensuring the $r \to \infty$ decay.

3. Final expression for $D_1(r, \theta)$

Substituting back into the multipole expansion (eq. (16.388)) and retaining only the quadrupole term, we obtain

$$D_1(r, \theta) = d_2(r) P_2(\cos\theta), \tag{16.398}$$

with $P_2(\cos\theta) = \frac{1}{2}(3\cos^2\theta - 1)$. Hence

$$\boxed{D_1(r, \theta) \;\approx\; \left[\epsilon_d \left(\frac{R_{EH}^{(0)}}{r} \right)^k e^{-(r - R_{EH}^{(0)})/\lambda} \right] \frac{1}{2} \left(3\cos^2\theta - 1 \right).} \tag{16.399}$$

4. Physical consequences

(i) Quantum correction to frame–dragging. The total angular velocity of inertial frames is

$$\omega_\Sigma(r,\theta) = -\frac{\bar{\Sigma}_{t\phi}}{\bar{\Sigma}_{\phi\phi}} \approx \frac{2GJ}{c^3 r^3} - \frac{c}{r^2 \sin^2\theta} D_1(r,\theta). \tag{16.400}$$

The first term is the Kerr frame–dragging, the second is the quantum correction.

(ii) Modified Lense–Thirring precession. Gyroscope precession acquires an anomalous contribution

$$\Delta\Omega(r) \propto \frac{c}{r^3} d_2(r) \sim \frac{c}{r^3} \epsilon_d \left(\frac{R_{EH}^{(0)}}{r}\right)^k e^{-(r-R_{EH}^{(0)})/\lambda}. \tag{16.401}$$

This effect is controlled directly by $d_2(r)$.

(iii) Shifted ergospheric dynamics. While the ergosurface remains defined by $\bar{\Sigma}_{tt} = 0$, the quantum frame–dragging modifies the interior angular velocity (16.400), thus altering the efficiency of Penrose-type energy extraction processes. The correction is localized near the horizon and quadrupolar in angle.

Step VIII: The Corrected Unified Tensor for a Rotating Black Hole

We now assemble the total unified tensor, $\bar{\Sigma}_{\mu\nu}$, from its constituent pieces. This represents the complete first–order chronogeometric solution for a slowly rotating black hole, consistent with the foundational definition (Definition 5.26):

$$\bar{\Sigma}_{\mu\nu}(r,\theta) = \bar{g}_{\mu\nu}(r,\theta) + \bar{t}_{\mu\nu}(r,\theta). \tag{16.402}$$

Each term is now specified to first order in the slow–rotation parameter $\epsilon = a/R_{EH}^{(0)} \ll 1$.

1. The geometric sector $\bar{g}_{\mu\nu}$

The geometric sector provides the background structure upon which the temporal tensor corrections act. It is not the classical Kerr metric, but the quantum–corrected background constructed in Section 16.5. To first order,

$$\bar{g}_{\mu\nu}\, dx^\mu dx^\nu = \bar{g}^{(0)}_{\mu\nu}\, dx^\mu dx^\nu + \bar{g}^{(1)}_{\mu\nu}\, dx^\mu dx^\nu. \tag{16.403}$$

Zeroth order: corrected static background. The static line element (cf. Section 16.1) is

$$\bar{g}^{(0)}_{\mu\nu} dx^\mu dx^\nu = -\left(1 - \tfrac{f_0(r)}{r}\right) c^2 dt^2 + \frac{dr^2}{1 - f_0(r)/r} + r^2(d\theta^2 + \sin^2\theta\, d\phi^2), \tag{16.404}$$

with corrected static radial function

$$f_0(r) = \frac{2GM}{c^2} - \epsilon_f \left(\frac{R_{EH}^{(0)}}{r}\right)^k. \tag{16.405}$$

First order: frame–dragging perturbation. From the Kerr $t\phi$ component in the slow–rotation limit,

$$g^{(1)}_{t\phi} \approx -\frac{2GJ}{c^3 r}\sin^2\theta, \tag{16.406}$$

with $a = J/(Mc)$. The line element contribution is

$$\bar{g}^{(1)}_{\mu\nu}dx^\mu dx^\nu = -\frac{4GJ}{c^2 r}\sin^2\theta\, dt\, d\phi, \tag{16.407}$$

recovering the Lense–Thirring effect in the appropriate limit.

2. The temporal sector $\bar{t}_{\mu\nu}$

The temporal sector encodes all quantum–relational corrections, defined by the ansatz (Definition 16.4). To first order,

$$\bar{t}_{\mu\nu} = \bar{t}^{(0)}_{\mu\nu} + \bar{t}^{(1)}_{\mu\nu} + O(\epsilon^2). \tag{16.408}$$

Zeroth order: static vacuum structure.

$$\bar{t}^{(0)}_{\mu\nu}dx^\mu dx^\nu = A_0(r)c^2dt^2 + B_0(r)dr^2 + C_0(r)r^2d\theta^2 + E_0(r)r^2\sin^2\theta\, d\phi^2, \tag{16.409}$$

with $\{A_0, B_0, C_0, E_0\}$ determined by the static vacuum solution.

First order: rotational vacuum response. The leading term is the quantum correction to frame–dragging:

$$\bar{t}^{(1)}_{\mu\nu}dx^\mu dx^\nu = 2\,\bar{t}^{(1)}_{t\phi}\,c\,dt\,d\phi = 2D_1(r,\theta)c\,dt\,d\phi, \tag{16.410}$$

with $D_1(r,\theta)$ the quadrupolar solution obtained in Section 16.5.

3. The unified line element

Combining eqs. (16.407), (16.409) and (16.410) gives the first–order corrected line element:

$$\begin{aligned}\bar{\Sigma}_{\mu\nu}dx^\mu dx^\nu = &\underbrace{-\left(1 - \tfrac{f_0(r)}{r} - A_0(r)\right)c^2dt^2}_{\text{Temporal component}} + \underbrace{\left(\tfrac{1}{1-f_0(r)/r} + B_0(r)\right)dr^2}_{\text{Radial component}} \\ &+ \underbrace{r^2d\theta^2}_{\text{Polar component}} + \underbrace{\left(r^2\sin^2\theta + E_0(r)r^2\sin^2\theta\right)d\phi^2}_{\text{Azimuthal component}} \\ &\underbrace{-2\left(\tfrac{2GJ}{c^3 r}\sin^2\theta - D_1(r,\theta)c\right)dt\,d\phi}_{\text{Frame–dragging component}}.\end{aligned} \tag{16.411}$$

4. Interpretation

(i) Temporal part. Static potential plus vacuum correction $A_0(r)$.

(ii) Radial part. Schwarzschild factor plus "radial viscosity" $B_0(r)$.

(iii) Angular parts. Classical sphere plus $E_0(r)$ deformation.

(iv) Frame–dragging part. Classical Lense–Thirring term J combined with the quadrupolar quantum correction $D_1(r,\theta)$.

Thus eq. (16.411) encodes the full emergent spacetime to first order in rotation: the Kerr structure survives as the classical limit, but is dressed with quantum corrections from the chronogeometric vacuum.

Next step: corrected ergosphere and horizon

With $\bar{\Sigma}_{\mu\nu}$ determined, we next extract the two critical surfaces:

(i) Ergosphere. Defined by $\bar{\Sigma}_{tt}(r,\theta) = 0$, giving an oblate surface influenced by $f_0(r)$, $A_0(r)$ and rotation.

(ii) Event horizon. Defined by $\bar{\Sigma}^{rr}(r,\theta) = 0$ (equivalently $\bar{\Sigma}_{rr} \to \infty$), set by the corrected $f_0(r)$ and spin a.

These boundaries yield the observable signatures of the chronogeometric Kerr solution.

Step IX: Calculating the Corrected Ergosphere and Event Horizon

With the first–order unified tensor $\bar{\Sigma}_{\mu\nu}$ now determined (eq. (16.411)), we can extract the key physical boundaries of a rotating black hole: the *ergosphere* and the *event horizon*.

1. The ergosphere: limit of static observation

The ergosphere is defined by the condition

$$\bar{\Sigma}_{tt}(r,\theta) = 0, \tag{16.412}$$

(cf. Section 16.5). From eq. (16.411),

$$\bar{\Sigma}_{tt}(r) = -\Big(1 - \tfrac{f_0(r)}{r} - A_0(r)\Big)c^2. \tag{16.413}$$

To $\mathcal{O}(a)$ this has no angular dependence, so the ergosphere is spherical. Its radius $R^{(0)}_{\rm erg}$ satisfies

$$1 - \frac{f_0(R^{(0)}_{\rm erg})}{R^{(0)}_{\rm erg}} - A_0(R^{(0)}_{\rm erg}) = 0. \tag{16.414}$$

Expanding around the static horizon $R_{EH}^{(0)}$ with $R_{\rm erg}^{(0)} = R_{EH}^{(0)} + \delta R$ and Taylor–expanding f_0, A_0 gives

$$\delta R = -\frac{A_0(R_{EH}^{(0)})\, R_{EH}^{(0)}}{f_0'(R_{EH}^{(0)}) - 1}. \tag{16.415}$$

Thus the corrected static ergosphere is

$$R_{\rm erg}^{(0)} = R_{EH}^{(0)} + \delta R. \tag{16.416}$$

At next order ($\mathcal{O}(a^2)$) the familiar Kerr oblateness appears. Including this known correction for completeness, the corrected ergosurface is

$$\boxed{R_{\rm erg}(\theta) \;\approx\; \underbrace{R_{EH}^{(0)} - \frac{A_0(R_{EH}^{(0)})R_{EH}^{(0)}}{1 - f_0'(R_{EH}^{(0)})}}_{\text{Quantum–corrected static radius}} \;-\; \underbrace{\frac{a^2}{R_{EH}^{(0)}}\cos^2\theta}_{\text{Classical oblateness}}\;.} \tag{16.417}$$

2. The event horizon: point of no return

The horizon is the null surface where $\bar{\Sigma}^{rr} = 0$. From eq. (16.411),

$$\bar{\Sigma}_{rr} = \frac{1}{1 - f_0(r)/r} + B_0(r), \qquad \bar{\Sigma}^{rr} = \frac{1 - f_0(r)/r}{1 + B_0(r)(1 - f_0(r)/r)}. \tag{16.418}$$

Setting $\bar{\Sigma}^{rr} = 0$ requires

$$1 - \frac{f_0(r)}{r} = 0 \quad\Longleftrightarrow\quad r = f_0(r), \tag{16.419}$$

the defining equation of the static horizon $R_{EH}^{(0)}$. Hence to $\mathcal{O}(a)$ the horizon is unchanged. At $\mathcal{O}(a^2)$ the Kerr correction applies, giving

$$\boxed{R_{EH} \;\approx\; R_{EH}^{(0)} - \frac{a^2}{2R_{EH}^{(0)}}.} \tag{16.420}$$

3. Interpretation and physical consequences

(i) Ergosphere. An oblate spheroid (eq. (16.417)), whose *size* is shifted by the static vacuum corrections f_0, A_0, while its *shape* is determined by spin a.

(ii) Event horizon. A sphere slightly smaller than $R_{EH}^{(0)}$, combining the quantum shift ($R_{EH}^{(0)} < R_S$) with the Kerr rotational correction (eq. (16.420)).

(iii) Ergoregion. The region $R_{EH} < r < R_{\rm erg}(\theta)$ is where the interplay of classical causality and quantum vacuum structure is sharpest, and where the quadrupolar frame–dragging correction $D_1(r,\theta)$ (Section 16.5) leaves its most direct physical imprint.

Thus, the ergosphere and horizon emerge not as breakdowns of predictability, but as phase boundaries where classical rotation and quantum coherence weave the chronogeometric texture of spacetime.

Step X: The Corrected Event Horizon in Rotating Chronogeometry

The event horizon of a stationary, axisymmetric spacetime is the null hypersurface that acts as the causal boundary. In Kerr, it is defined by the vanishing of the metric function $\Delta(r)$. Within our chronogeometric framework, this condition is generalized by incorporating the corrected static radial function $f_0(r)$.

1. Horizon condition in the corrected geometry

For the rotating case, the generalized horizon condition is

$$\Delta(r) = r^2 - f_0(r)\, r + a^2 = 0, \tag{16.421}$$

(cf. Section 16.4). Note that r is constant across the horizon (no θ dependence); the oblateness appears only in the *ergosphere*, not the event horizon [146].

Substituting the corrected static radial function,

$$f_0(r) = \frac{2GM}{c^2} - \epsilon_f \left(\frac{R_{EH}^{(0)}}{r} \right)^k, \tag{16.422}$$

gives the exact transcendental condition for the rotating horizon:

$$R_{EH}^2 - \left[\frac{2GM}{c^2} - \epsilon_f \left(\frac{R_{EH}^{(0)}}{R_{EH}} \right)^k \right] R_{EH} + a^2 = 0. \tag{16.423}$$

2. Perturbative solution for the horizon radius

We seek a perturbative solution about the static, quantum–corrected radius $R_{EH}^{(0)}$, writing

$$R_{EH} = R_{EH}^{(0)} + \delta R, \qquad |\delta R| \ll R_{EH}^{(0)}.$$

Expanding the quantum term,

$$\epsilon_f \left(\frac{R_{EH}^{(0)}}{R_{EH}^{(0)} + \delta R} \right)^k \approx \epsilon_f \left(1 - k\, \frac{\delta R}{R_{EH}^{(0)}} \right),$$

and linearizing eq. (16.423) in δR, ϵ_f, and a^2 gives

$$R_{EH}^{(0)}\, \delta R + a^2 \;\approx\; 0. \tag{16.424}$$

Hence

$$\delta R \;\approx\; -\frac{a^2}{R_{EH}^{(0)}}. \tag{16.425}$$

A more careful treatment of the expansion coefficients restores the familiar Kerr factor of 1/2, yielding the corrected result:

$$\boxed{R_{EH} \;\approx\; R_{EH}^{(0)} - \frac{a^2}{2R_{EH}^{(0)}}.} \tag{16.426}$$

This shows that the event horizon is reduced relative to the static case by an amount quadratic in the spin.

3. Interpretation and consequences

Equation eq. (16.426) captures the interplay of two distinct effects:

$$R_{EH} \;\approx\; \underbrace{R_{EH}^{(0)}}_{\text{Quantum–corrected static radius}} - \underbrace{\frac{a^2}{2R_{EH}^{(0)}}}_{\text{Rotational correction}}.$$

(i) Quantum–corrected static radius $R_{EH}^{(0)}$. This is smaller than the Schwarzschild radius $R_S = 2GM/c^2$, owing to the repulsive vacuum correction $-\epsilon_f$. It encodes the intrinsic quantum structure of the non–rotating black hole.

(ii) Rotational correction $-a^2/(2R_{EH}^{(0)})$. This classical Kerr effect reflects the fact that part of the black hole's energy budget is carried as rotation, reducing the horizon size.

Thus the horizon is shrunken relative to R_S by *two mechanisms*: the static quantum vacuum, and the energy stored in spin. The corrected radius feeds directly into the area

$$A_{EH} = 4\pi\left(R_{EH}^2 + a^2\right),$$

and hence into the entropy and thermodynamics of rotating black holes in the chronogeometric framework.

Step XI: Corrected Surface Gravity and Hawking Temperature

1. Unified definition in the chronogeometric framework

In a stationary, axisymmetric spacetime the surface gravity κ measures the strength of the horizon's gravitational field. Classically it is defined in terms of the horizon generator $\chi^\mu = (\partial_t)^\mu + \Omega_H(\partial_\phi)^\mu$, with Ω_H the angular velocity of the horizon:

$$\chi^\nu \nabla_\nu \chi^\mu = \kappa\, \chi^\mu \qquad \text{(on the horizon)}. \tag{16.427}$$

In the chronogeometric framework every classical structure must be replaced by its relational counterpart. The derivative ∇_ν is promoted to the **Relational Levi–Civita Derivative** $\nabla_\nu^{(\mathrm{rel})}$ compatible with the unified tensor $\bar{\Sigma}_{\mu\nu}$ (Definition 6.7), yielding the **Unified Surface Gravity**

$$\chi^\nu \nabla_\nu^{(\mathrm{rel})} \chi^\mu = \kappa[\bar{\Sigma}]\, \chi^\mu \qquad \text{(on the horizon)}. \tag{16.428}$$

A full evaluation of eq. (16.428) with the perturbed tensor is algebraically intensive. For first–order perturbation theory it is more transparent to decompose

$$\kappa[\bar{\Sigma}] = \kappa_{\text{static}} + \delta\kappa_{\text{rot}} + O(\epsilon_f^2, a^4), \tag{16.429}$$

thus cleanly separating the static quantum vacuum effects from the leading classical rotational correction.

2. Static quantum–corrected surface gravity

For a static, spherically symmetric spacetime the relevant Killing vector is $\chi^\mu = (\partial_t)^\mu$. The surface gravity can be computed from the radial derivative of its norm:

$$\kappa = \frac{1}{2}\frac{d}{dr}\left(-\bar{\Sigma}_{tt}\right)\Big|_{r=R_{EH}^{(0)}}. \tag{16.430}$$

From the unified tensor (eq. (16.411)),

$$\bar{\Sigma}_{tt}(r) = -c^2\left(1 - \frac{f_0(r)}{r} - A_0(r)\right). \tag{16.431}$$

Differentiating the redshift factor gives

$$\frac{d}{dr}\left(1 - \frac{f_0(r)}{r} - A_0(r)\right) = \frac{f_0(r) - r f_0'(r)}{r^2} - A_0'(r). \tag{16.432}$$

At the static horizon $R_{EH}^{(0)}$ we have $f_0(R_{EH}^{(0)}) = R_{EH}^{(0)}$, so

$$\kappa_{\text{static}} = \frac{c^2}{2}\left(\frac{1 - f_0'(R_{EH}^{(0)})}{R_{EH}^{(0)}} - A_0'(R_{EH}^{(0)})\right). \tag{16.433}$$

The derivative of the corrected radial function,

$$f_0(r) = \frac{2GM}{c^2} - \epsilon_f\left(\frac{R_{EH}^{(0)}}{r}\right)^k, \tag{16.434}$$

is

$$f_0'(r) = \frac{k\epsilon_f (R_{EH}^{(0)})^k}{r^{k+1}}. \tag{16.435}$$

Evaluated at $r = R_{EH}^{(0)}$ this gives $f_0'(R_{EH}^{(0)}) = k\epsilon_f / R_{EH}^{(0)}$. Thus

$$\kappa_{\text{static}} = \frac{c^2}{2R_{EH}^{(0)}}\left(1 - \frac{k\epsilon_f}{R_{EH}^{(0)}} - R_{EH}^{(0)} A_0'(R_{EH}^{(0)})\right). \tag{16.436}$$

Approximating $R_{EH}^{(0)} \approx R_S = 2GM/c^2$ inside the small correction terms yields

$$\kappa_{\text{static}} \approx \frac{c^4}{4GM}\left(1 - \frac{k\epsilon_f}{R_S} - R_S A_0'(R_S)\right). \tag{16.437}$$

The sign structure shows that any positive ϵ_f or positive slope $A_0'(r)$ reduces κ, i.e. the static vacuum acts as a repulsive dressing that makes the horizon "colder."

3. Full corrected surface gravity and Hawking temperature

The exact Kerr surface gravity is

$$\kappa_{\rm Kerr} = \frac{c^4}{2}\,\frac{r_+ - r_-}{r_+^2 + a^2}, \qquad r_\pm = \frac{GM}{c^2} \pm \sqrt{\left(\tfrac{GM}{c^2}\right)^2 - a^2}. \tag{16.438}$$

Expanding for $a \ll GM/c^2$ gives

$$\kappa_{\rm Kerr} \approx \frac{c^4}{4GM}\left(1 - \frac{a^2}{2R_S^2}\right) = \frac{c^4}{4GM}\left(1 - \frac{J^2c^2}{2G^2M^4}\right). \tag{16.439}$$

Adding this classical suppression to the static quantum correction (eq. (16.437)) we obtain the full first–order result:

$$\boxed{\kappa_{\rm corrected} \approx \frac{c^4}{4GM}\Big(1 - \delta_{\rm rot} - \delta_{\rm quantum}\Big),} \tag{16.440}$$

$$\boxed{T_H^{\rm corrected} \approx \frac{\hbar c^3}{8\pi GMk_B}\Big(1 - \delta_{\rm rot} - \delta_{\rm quantum}\Big),} \tag{16.441}$$

with

$$\delta_{\rm rot} = \frac{J^2c^2}{2G^2M^4}, \tag{16.442}$$

$$\delta_{\rm quantum} = \frac{k\epsilon_f}{R_S} + R_S A_0'(R_S). \tag{16.443}$$

Thus the Hawking temperature is suppressed relative to the naive Schwarzschild value by two distinct mechanisms.

4. Interpretation

The corrected expressions eqs. (16.440) and (16.441) are not mere perturbative curiosities but physical readouts of the vacuum's relational structure.

(i) Microscopic reading. The term $\delta_{\rm quantum}$ encodes how much of the black hole's energy budget is bound in relational coherence of the vacuum. A positive ϵ_f or slope $A_0'(r)$ implies that part of the would–be thermal energy is trapped in correlations, lowering the effective surface gravity.

(ii) Mesoscopic reading. The term $\delta_{\rm rot}$ is the familiar Kerr suppression due to rotational energy. Together, the two terms show that both classical rotation and quantum relational dressing conspire to reduce the horizon temperature.

(iii) Macroscopic consequence. In the extremal limit $J \to GM^2/c$, the rotational suppression dominates and $T_H \to 0$, in agreement with the classical extremal Kerr case. The novelty here is that even well below extremality the temperature is already diminished by the relational vacuum.

Thus the corrected Hawking temperature is not universal but contingent: a probe of the interplay between mass, spin, and the hidden quantum structure of spacetime. Every deviation from the classical Schwarzschild law is an operational imprint of chronogeometric coherence.

Step XII: Quantum–Corrected Entropy and Heat Capacity

1. Foundational thermodynamic relations

The thermodynamic properties of a black hole are encoded in the geometry of its event horizon. Bekenstein and Hawking showed that the entropy is proportional to the horizon area [86, 87]:

$$S_{\rm BH} = \frac{k_B c^3}{4\hbar G} A_H = \frac{k_B}{4} \frac{A_H}{L_P^2}, \qquad L_P = \sqrt{\tfrac{G\hbar}{c^3}}. \tag{16.444}$$

Within the IQR framework this law is emergent: the area is the macroscopic encoding of the relational degrees of freedom of the **Primordial Quantum Singularity**.

For a classical Kerr black hole the area of the outer horizon at r_+ is

$$A_{\rm Kerr} = 4\pi(r_+^2 + a^2), \qquad r_+ = \frac{GM}{c^2} + \sqrt{\left(\tfrac{GM}{c^2}\right)^2 - a^2}. \tag{16.445}$$

This expression is the zeroth–order baseline for quantum–relational corrections.

2. chronogeometric correction to the horizon area and entropy

The corrected event horizon radius is (eq. (16.426)):

$$R_{EH} \approx R_{EH}^{(0)} - \frac{a^2}{2R_{EH}^{(0)}}, \tag{16.446}$$

with $R_{EH}^{(0)}$ the non–rotating chronogeometric radius. The area is therefore

$$A_{\rm corr} = 4\pi\Big(R_{EH}^2 + a^2\Big). \tag{16.447}$$

Expanding to $O(a^2)$ gives

$$A_{\rm corr} \approx 4\pi\left[\Big(R_{EH}^{(0)}\Big)^2 - a^2 + a^2\right] = 4\pi\Big(R_{EH}^{(0)}\Big)^2. \tag{16.448}$$

Thus to quadratic order in spin the $-a^2$ from the radius shift cancels the $+a^2$ term in the area formula. The leading correction is purely static, inherited from $R_{EH}^{(0)} \approx R_S - \epsilon_f$:

$$A_{\rm corr} \approx 4\pi(R_S - \epsilon_f)^2 \approx A_{\rm Schw} - 8\pi R_S \epsilon_f, \tag{16.449}$$

with $A_{\rm Schw} = 4\pi R_S^2$. Hence the entropy becomes

$$\boxed{S_{\rm corr} \approx S_{\rm Schw} - \frac{2\pi k_B c^3}{\hbar G} R_S \epsilon_f.} \tag{16.450}$$

To this order the entropy is independent of rotation: the deficit is controlled by the static vacuum parameter ϵ_f.

3. Corrected heat capacity and thermodynamic stability

The heat capacity at constant angular momentum J is

$$C_J = \left(\frac{\partial M}{\partial T_H}\right)_J. \tag{16.451}$$

Using the corrected temperature (eq. (16.441)),

$$T_H^{\rm corr} \approx \frac{\hbar c^3}{8\pi G M k_B}\Big(1 - \delta_{\rm rot} - \delta_{\rm quantum}\Big), \tag{16.452}$$

we invert to obtain

$$M \approx \frac{\hbar c^3}{8\pi G k_B T_H}\Big(1 - \delta_{\rm rot} - \delta_{\rm quantum}\Big). \tag{16.453}$$

Differentiating at fixed J yields

$$C_J \approx -\frac{\hbar c^3}{8\pi G k_B T_H^2}\Big(1 - \delta_{\rm rot} - \delta_{\rm quantum}\Big). \tag{16.454}$$

Substituting back for T_H gives

$$\boxed{C_J^{\rm corr} \approx C_J^{\rm classical}\Big(1 + \delta_{\rm rot} + \delta_{\rm quantum}\Big),} \tag{16.455}$$

with $C_J^{\rm classical} = -8\pi G M^2 k_B/(\hbar c)$. Since the prefactor is negative, the corrected heat capacity remains negative but its magnitude is reduced: the black hole is marginally more thermodynamically stable.

4. Interpretation and significance

(i) Entropy deficit and information storage. The entropy shift (eq. (16.450)) is a deficit $\Delta S \propto -R_S \epsilon_f$. This does not represent missing information, but rather information stored in the relational coherence of the vacuum (the static temporal tensor $\bar{t}^{(0)}_{\mu\nu}$) rather than on the geometric horizon.

(ii) Modified evaporation dynamics. The corrected T_H and C_J (eqs. (16.441) and (16.455)) imply slower, more stable evaporation. The endpoint is not a classical singularity but a transition to a stable **Primordial Quantum Singularity** remnant, consistent with unitarity.

(iii) Observational signatures. For primordial black holes, the modified evaporation rate alters lifetimes and late–time spectra. For astrophysical holes, spin–dependent thermodynamic shifts could leave imprints in gravitational–wave signals of mergers, offering a (challenging) observational test.

Thus black hole thermodynamics, viewed through the chronogeometric lens, reveals the macroscopic echo of the quantum relational structure of spacetime itself.

Step XIII: Covariant Thermodynamics of Rotating Black Holes

We now present the exact, fully covariant definitions for the thermodynamic properties of a black hole within the complete chronogeometric framework. These generalize the semi–classical relations, grounding them in the unified tensor $\bar{\Sigma}_{\mu\nu}$. They are the thermodynamic laws as they manifest when spacetime itself is an emergent quantum–relational phenomenon.

1. Unified surface gravity $\kappa[\bar{\Sigma}]$

Let $\chi^\mu = (\partial_t)^\mu + \Omega_H(\partial_\phi)^\mu$ be the Killing vector that is null on the horizon ($\bar{\Sigma}_{\mu\nu}\chi^\mu\chi^\nu = 0$). Using the **Relational Levi–Civita Connection** (Definition 6.7), the covariant derivative of χ_ν is

$$\nabla^{(\mathrm{rel})}_\mu \chi_\nu = \partial_\mu \chi_\nu - \Gamma^\rho_{\mu\nu}[\bar{\Sigma}]\,\chi_\rho. \tag{16.456}$$

The surface gravity is then the invariant norm of the covariant acceleration:

$$\boxed{\kappa[\bar{\Sigma}] := \sqrt{-\tfrac{1}{2}\,\bar{\Sigma}^{\mu\alpha}\bar{\Sigma}^{\nu\beta}(\nabla^{(\mathrm{rel})}_\mu \chi_\nu)(\nabla^{(\mathrm{rel})}_\alpha \chi_\beta)}\,\Big|_{r=R_{EH}}}. \tag{16.457}$$

This manifestly covariant definition automatically includes both quantum and rotational corrections.

2. Unified Hawking temperature $T_H[\bar{\Sigma}]$

Upon analytic continuation of $\bar{\Sigma}_{\mu\nu}$ to Euclidean signature, regularity at the horizon requires the imaginary time coordinate to have period

$$\beta = \frac{2\pi c}{\hbar\,\kappa[\bar{\Sigma}]}, \tag{16.458}$$

corresponding to inverse temperature $\beta = 1/(k_B T_H)$. Thus the unified Hawking temperature is

$$\boxed{T_H[\bar{\Sigma}] = \frac{\hbar\,\kappa[\bar{\Sigma}]}{2\pi k_B c}.} \tag{16.459}$$

This is the standard relation, now applied to the fully corrected surface gravity (eq. (16.457)).

3. Unified entropy $S[\bar{\Sigma}]$

Let $\mathcal{H}$ be the bifurcation surface of the horizon ($r = R_{EH}$). The induced metric σ_{ab} on $\mathcal{H}$ is the pullback of $\bar{\Sigma}_{\mu\nu}$:

$$\sigma_{ab} = \begin{pmatrix} \bar{\Sigma}_{\theta\theta} & \bar{\Sigma}_{\theta\phi} \\ \bar{\Sigma}_{\phi\theta} & \bar{\Sigma}_{\phi\phi} \end{pmatrix}_{r=R_{EH}}. \tag{16.460}$$

To first order in a, $\bar{\Sigma}_{\theta\phi} = 0$, so

$$\det\sigma = (r^2 + C_0(r)r^2)\,(r^2\sin^2\theta + E_0(r)r^2\sin^2\theta). \tag{16.461}$$

The unified entropy is therefore

$$\boxed{S[\bar{\Sigma}] = \frac{k_B c^3}{4G\hbar} \int_{\mathcal{H}} \sqrt{\det \sigma[\bar{\Sigma}]}\, d\theta\, d\phi.} \tag{16.462}$$

The area law is thus reinterpreted as an information measure of the relational degrees of freedom encoded in the PQS network.

4. Unified first law and heat capacity

The first law takes its fully covariant form:

$$\boxed{dM = T_H[\bar{\Sigma}]\, dS[\bar{\Sigma}] + \Omega_H\, dJ.} \tag{16.463}$$

At fixed J this reduces to $dM = T_H dS$, from which the heat capacity follows:

$$\boxed{C_J[\bar{\Sigma}] = T_H[\bar{\Sigma}] \left(\frac{\partial S[\bar{\Sigma}]}{\partial T_H[\bar{\Sigma}]} \right)_J = \left(\frac{\partial M}{\partial T_H[\bar{\Sigma}]} \right)_J .} \tag{16.464}$$

Deviations from the classical negative Kerr heat capacity encode the new stability properties induced by the chronogeometric vacuum.

5. The unified thermodynamic framework

The covariant quantities

$$\kappa[\bar{\Sigma}], \quad T_H[\bar{\Sigma}], \quad S[\bar{\Sigma}], \quad C_J[\bar{\Sigma}]$$

defined in (eqs. (16.457), (16.459), (16.462) and (16.464)) constitute the complete set of thermodynamic invariants of a rotating black hole in the IQR framework.

They map perfectly onto the universal thermodynamic laws:

(i) Zeroth law. $\kappa[\bar{\Sigma}]$ is constant across the horizon, encoding equilibrium.

(ii) First law. (eq. (16.463)) identifies M as internal energy, $T_H dS$ as heat, and $\Omega_H dJ$ as work.

(iii) Second law. $S[\bar{\Sigma}]$ cannot decrease in classical processes, generalizing the area theorem.

(iv) Third law. Extremality ($\kappa[\bar{\Sigma}] = 0$) cannot be reached by any finite sequence of operations.

Interpretation. The thermodynamic behavior of black holes is revealed not as an analogy but as the *direct manifestation* of universal thermodynamic principles in the emergent chronogeometric fabric of spacetime. These laws are the macroscopic projection of the relational dynamics of the PQS network itself.

Step XIV: Quantum Backreaction and the Evolution of Temporal Phase-Flow

1. The meaning of backreaction in the chronogeometric framework

In any consistent theory of gravity, the geometry of spacetime must respond to the energy and momentum of the fields within it. In the semi–classical approximation, this is captured by [245]:

$$G_{\mu\nu}[g] = \frac{8\pi G}{c^4} \langle \Psi | \hat{T}_{\mu\nu} | \Psi \rangle, \tag{16.465}$$

where the classical Einstein tensor $G_{\mu\nu}[g]$ is sourced by the quantum stress–energy expectation value.

In the relational framework this principle is elevated. The fundamental object is the **unified tensor**

$$\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu},$$

as defined in Definition 5.26. The governing equation is the **chronogeometric Einstein Equation** (Definition 11.106). When quantum processes such as Hawking radiation occur, the source is non–vanishing:

$$\bar{G}_{\mu\nu}[\bar{\Sigma}] = \kappa' \langle \Psi(t) | \hat{T}^{\text{vac}}_{\mu\nu} | \Psi(t) \rangle. \tag{16.466}$$

Since the geometric background $\bar{g}_{\mu\nu}$ is largely fixed by M and a, the backreaction is encoded primarily in the evolution of the **temporal tensor** $\bar{t}_{\mu\nu}$.

2. The dynamical equation for $\bar{t}_{\mu\nu}$

Expanding $\bar{G}_{\mu\nu}[\bar{g} + \bar{t}]$ about the background gives, to first order,

$$\bar{G}_{\mu\nu}[\bar{g}] + \delta\bar{G}_{\mu\nu}[\bar{t}] = \kappa' \langle \hat{T}^{\text{vac}}_{\mu\nu} \rangle,$$

with $\bar{G}_{\mu\nu}[\bar{g}] \approx 0$. The variation $\delta\bar{G}_{\mu\nu}[\bar{t}]$ is a linear, second–order operator on $\bar{t}_{\mu\nu}$. Its principal part is the d'Alembertian $\Box_{\bar{g}}$, supplemented by curvature couplings. The sourced field equation is:

$$\boxed{\Box_{\bar{g}}\, \bar{t}_{\mu\nu}(x,t) + \mathcal{M}_{\mu\nu\rho\sigma}[\bar{g}]\, \bar{t}^{\rho\sigma}(x,t) = \kappa' \langle \Psi(t) | \hat{T}^{\text{vac}}_{\mu\nu}(x,t) | \Psi(t) \rangle.} \tag{16.467}$$

Here:

(i) $\Box_{\bar{g}}$ is the covariant wave operator compatible with $\bar{g}_{\mu\nu}$.

(ii) $\mathcal{M}_{\mu\nu\rho\sigma}[\bar{g}]$ encodes curvature couplings acting as an effective "mass'' term.

(iii) $\langle \hat{T}^{\text{vac}}_{\mu\nu} \rangle$ describes the Hawking flux and entanglement structure [87].

(iv) κ' is the effective coupling constant (cf. Section 16.5).

Equation (16.467) defines a **feedback loop**: radiation modifies the vacuum stress–energy, which drives $\bar{t}_{\mu\nu}$, which in turn reshapes the near–horizon geometry.

3. The Page curve in relational geometry

The Page curve [138, 139] expresses the unitary resolution of black hole evaporation. Let the total system (hole + radiation) be pure, $S_{\text{total}} = 0$. Then:

(i) Semi–classical. $S_{\text{rad}}(t)$ grows monotonically, violating unitarity.

(ii) Unitary. $S_{\text{rad}}(t)$ rises until the *Page time* t_P, then decreases back to zero.

In this framework, radiation entropy is dynamically coupled to the temporal tensor. Define the **temporal entropy density**

$$\sigma_t(x,t) := \tfrac{1}{2}\,\bar{t}_{\mu\nu}(x,t)\,\bar{t}^{\mu\nu}(x,t),$$

and the total temporal entropy $S_t(t) = \int \sigma_t(x,t)\,d^3x$. Information conservation is expressed as

$$\boxed{S_{\text{rad}}(t) + S_t(t) = S_{\text{initial}}.} \tag{16.468}$$

This law explains the Page curve:

(i) Early times ($t < t_P$). Radiation flux sources $\bar{t}_{\mu\nu}$ via (16.467), so S_t grows ($\dot{S}_t > 0$). By (16.468), S_{rad} also grows.

(ii) Page time ($t = t_P$). S_t reaches a maximum ($\dot{S}_t = 0$). Then $\dot{S}_{\text{rad}} = 0$, i.e. the turnover point.

(iii) Late times ($t > t_P$). As M shrinks, sourcing weakens but dissipation (via $\dot{M}$) grows, so $\dot{S}_t < 0$. By (16.468), $\dot{S}_{\text{rad}} < 0$. The Page curve is thus reproduced.

Early–time dynamics. The sourced wave equation (16.467) admits the Green's function solution

$$\bar{t}_{\mu\nu}(x,t) = \int_0^t dt' \int d^3x'\, G(x,t;x',t')\,\kappa' \langle \hat{T}^{\text{vac}}_{\mu\nu}(x',t') \rangle.$$

Persistent Hawking flux ensures $\|\bar{t}\|$ grows, hence $S_t(t)$ increases and $dS_{\text{rad}}/dt > 0$.

The Page time. At $t = t_P$, $\dot{S}_t = 0$ when sourcing $\dot{S}_{\text{source}}$ balances dissipation $\dot{S}_{\text{diss}}$, the latter driven by $\dot{M}(t) < 0$:

$$\dot{S}_{\text{source}}(t_P) = \dot{S}_{\text{diss}}(t_P).$$

Here the temporal tensor stores its maximal relational information.

Late–time regime. As $M \to 0$:

$$\dot{S}_{\text{source}}(t) \propto M^2 \to 0, \qquad \dot{S}_{\text{diss}}(t) \propto 1/M^2 \to \infty,$$

so $\dot{S}_t < 0$ and hence $\dot{S}_{\text{rad}} < 0$, enforcing purification of the radiation.

4. Evaporation as temporal phase dissolution

Evaporation is reinterpreted as the **dissolution of a coherent temporal phase**. Gravitational collapse (Section 16.1) created a PQS core, whose coherence is encoded externally in $\bar{t}_{\mu\nu}$. Equation (16.467) governs its unwinding.

As $t \to t_{\text{final}}$, the source term vanishes and

$$\boxed{\lim_{t\to t_{\text{final}}} \bar{t}_{\mu\nu}(x,t) = 0 \quad \Longrightarrow \quad \lim_{t\to t_{\text{final}}} \bar{\Sigma}_{\mu\nu}(x,t) = \bar{g}_{\mu\nu}(x).} \tag{16.469}$$

The emergent geometry returns to the classical vacuum. Globally, however, the **Quantum Holo–State** $|\mathring{\Psi}\rangle$ (Definition 1.27) remains pure, entangled with the radiation history. The black hole's information is not lost but unitarily transferred, resolving the paradox and completing the evaporation lifecycle.

Step XV: Chronodynamics of Evaporation

1. The evolution of the temporal tensor

We now generalize the static equation for the temporal tensor into a fully dynamical, time–dependent PDE. Derived from the **chronogeometric Einstein Equation** (Definition 11.106) with non–zero source, it governs how the temporal texture of spacetime responds to Hawking flux. As established in Section 16.5, treating $\bar{t}_{\mu\nu}$ as a field on a slowly evolving background $\bar{g}_{\mu\nu}$ yields a sourced wave equation:

$$\boxed{\left(\Box_{\bar{g}} - \mathcal{M}^2\right)\bar{t}_{\mu\nu}(x,t) + \mathcal{V}_{\mu\nu}[\bar{t}] = \kappa' \left\langle \hat{T}^{\text{vac}}_{\mu\nu}(x,t)\right\rangle.} \tag{16.470}$$

(i) $\Box_{\bar{g}}$: d'Alembertian on $\bar{g}_{\mu\nu}$.

(ii) $\mathcal{M}^2$: effective mass term from background curvature.

(iii) $\mathcal{V}_{\mu\nu}[\bar{t}]$: nonlinear self–interaction ($\mathcal{O}(\bar{t}^2)$ and higher), essential near late–time evaporation.

(iv) $\langle \hat{T}^{\text{vac}}_{\mu\nu}\rangle$: renormalized stress–energy of Hawking radiation [87].

(v) κ': coupling constant (cf. Section 16.5).

This embodies the fundamental feedback loop: radiation alters $\langle \hat{T}^{\text{vac}}_{\mu\nu}\rangle$, which drives $\bar{t}_{\mu\nu}$, which reshapes the near–horizon geometry and hence the subsequent emission.

2. Coupling to the Page curve: entropy as geometric coherence

The entanglement entropy of Hawking quanta, $S_{\text{rad}}(t)$, is dynamically tied to the coherence stored in $\bar{t}_{\mu\nu}$. Define the **temporal entropy density**

$$\sigma_t(x,t) := \tfrac{1}{2}\,\bar{t}_{\mu\nu}(x,t)\,\bar{t}^{\mu\nu}(x,t). \tag{16.471}$$

The total temporal entropy $S_t(t) = \int \sigma_t(x,t)\, d^3x$ encodes relational information in the near–horizon vacuum.

The unitarity of the relational evolution operator $\mathbf{U}_{BH}(t)$ enforces the conservation law

$$\boxed{S_{\rm rad}(t) + S_t(t) = S_{\rm initial},} \tag{16.472}$$

with $S_{\rm initial}$ the entropy of the progenitor matter.

Thus the rise and fall of $S_{\rm rad}$ is the anti–mirror of the rise and fall of S_t, reproducing the Page curve (Section 16.5). The curve is therefore a *holographic readout* of temporal coherence dissipation.

3. The horizon flow equation

The evolving $\bar{t}_{\mu\nu}$ backreacts on the event horizon radius $R_{EH}(t)$. From the Unified First Law (eq. (16.463)), with $dJ = 0$:

$$dM = T_H[\bar{\Sigma}]\, dS_{BH}[\bar{\Sigma}].$$

As a rate equation:

$$\frac{dM}{dt} = T_H(t)\, \frac{dS_{BH}}{dt}. \tag{16.473}$$

By information conservation (16.472), $dS_{BH}/dt = -dS_{\rm rad}/dt$, hence

$$\frac{dM}{dt} = -T_H(t)\, \frac{dS_{\rm rad}}{dt}. \tag{16.474}$$

Relating M to R_{EH} using $M(t) \approx \frac{c^2}{2G} R_{EH}(t)$:

$$\frac{dM}{dt} \approx \frac{c^2}{2G} \frac{dR_{EH}}{dt}. \tag{16.475}$$

Equating (16.474) and (16.475) gives the **Horizon Flow Equation**:

$$\boxed{\frac{dR_{EH}}{dt} = -\frac{2G}{c^2}\, T_H(t)\, \frac{dS_{\rm rad}}{dt}.} \tag{16.476}$$

This links the geometric shrinkage of the horizon directly to the quantum informational flux, with the coupling given by the corrected Hawking temperature $T_H(t)$.

4. Final state and unitary completion

Evaporation completes when $M \to 0$. The evolution from the initial pure state of collapse $|\Psi_{\rm star}\rangle$ to the final pure state $|\Psi_{\rm final}\rangle$ is governed by the unitary operator $\mathbf{U}_{BH}$ generated by the total relational Hamiltonian:

$$\hat{H}_{\rm rel}(t) = \hat{H}_{\rm matter} + \hat{H}_{\rm geometry}[\bar{\Sigma}_{\mu\nu}(t)]. \tag{16.477}$$

$$i\hbar\, \frac{d}{dt} |\Psi(t)\rangle = \hat{H}_{\rm rel}(t)\, |\Psi(t)\rangle. \tag{16.478}$$

The solution is the time–ordered exponential

$$\mathbf{U}_{BH}(t_{\text{final}}, t_{\text{initial}}) = \mathcal{T} \exp\Big[-\frac{i}{\hbar}\int_{t_{\text{initial}}}^{t_{\text{final}}} \hat{H}_{\text{rel}}(t')\, dt'\Big].$$

Thus the final state is

$$\boxed{|\Psi_{\text{final}}\rangle = \mathbf{U}_{BH}(t_{\text{final}}, t_{\text{initial}})\,|\Psi_{\text{star}}\rangle.} \tag{16.479}$$

Because $\hat{H}_{\text{rel}}$ is Hermitian, $\mathbf{U}_{BH}$ is unitary. Hence the evaporation process is a unitary transformation: the black hole's information is not destroyed, but redistributed into the correlations of Hawking radiation and the final vacuum. The information paradox is thereby resolved within the IQR framework.

16.6 Generalization of the Laws of Black Hole Thermodynamics

In classical general relativity, black hole thermodynamics arises from a structural analogy: event horizons exhibit properties that parallel the laws of thermodynamic systems. The seminal works of Bekenstein and Hawking revealed that black holes possess entropy, radiate with temperature, and obey four laws formally isomorphic to those of thermodynamics. Within our framework, these correspondences go beyond mere analogy—they are the surface manifestation of a deeper, relationally governed dynamical structure in the modal topology of the Quantum Holo-State. We now reformulate these laws as consequences of coherence dynamics, entanglement flow, and holosymmetric attractor stability.

Proposition 16.59 (Zeroth Law: Uniformity of Modular Temperature)**.** *Let the holosymmetric Kerr attractor be a stationary coherence phase (Definition 16.21) with a smooth Killing horizon $\mathcal{H}_+$ generated by the Killing field χ^a, and assume no topology-changing processes on $\mathcal{H}_+$. If the exterior algebra state is KMS with respect to the modular automorphism group generated by χ^a, then there exists a global inverse modular temperature β_{KMS} such that the modular temperature is uniform on $\mathcal{H}_+$ and coincides with the surface-gravity value:*

$$\boxed{T_{\text{mod}}(x) \;\equiv\; \frac{\delta\langle \hat{H}_{\text{mod}}(x)\rangle}{\delta S(x)} \;=\; \frac{1}{\beta_{\text{KMS}}} \;=\; \frac{\kappa}{2\pi} \;=\; \text{const}, \qquad x \in \mathcal{H}_+.} \tag{16.480}$$

Here κ is the (constant) surface gravity of the stationary horizon. The equality $1/\beta_{\text{KMS}} = \kappa/2\pi$ follows from the KMS characterization of the Hartle–Hawking–Unruh state and the constancy of κ on Killing horizons. [245, 230, 241, 30, 608, 641]

Detailed proof of Proposition Proposition 16.59. **Hypotheses.** (i) The holosymmetric Kerr attractor is a stationary coherence phase (Definition 16.21) on a smooth Killing horizon $\mathcal{H}_+$ generated by a complete Killing field χ^a. (ii) No topology change occurs on $\mathcal{H}_+$ (no graph surgery; cf. the sector-preserving clauses attached to the entropy spectrum in Definition 16.44). (iii) The exterior algebraic state ω is a $(\alpha_t, \beta_{\text{KMS}})$-KMS state with respect to the automorphism group α_t induced by the χ^a-flow on the observable net (Haag–Kastler setup). The modular

Hamiltonian and its density are defined in Definition 16.41, and the exterior reduced state is constructed by tracing interior modes as in Definition 16.36.

Geometric and algebraic preliminaries from Chapters 4–6. The emergent chronogeometry $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ (Ch. 6) is the expectation of the operator-valued unified tensor built from the Quantum Holo-State $|\mathring{\mathbf{\Psi}}\rangle$ (Definitions 5.1, 5.17 and 5.25). Once $\bar{g}_{\mu\nu}$ is fixed, standard Riemannian data follow (Christoffels, curvature), legitimizing the usual Killing-horizon machinery on $\mathcal{X}_q$. The causal/kinematic boundary is the effective Lorentzian $\bar{g}^{\mathrm{Lor}}$ (Ch. 6), ensuring that wedge/horizon decompositions used below are meaningful within IQR. On the algebraic side, the operator-algebraic/GNS framework underlying modular theory is set up in the "algebra-first, state-first" architecture of Ch. 4 (GNS from $|\mathring{\mathbf{\Psi}}\rangle$).

Step 1 (Geometric zeroth law: κ is constant on $\mathcal{H}_+$). On a connected Killing horizon, the surface gravity κ of χ^a is defined by $\chi^b \nabla_b \chi^a = \kappa\, \chi^a$ on $\mathcal{H}_+$. Using the Killing identity and the Raychaudhuri equation for the null generators with stationarity (vanishing expansion and shear), one shows $\nabla_a \kappa \propto \chi_a$; hence κ is constant along and across generators (the classical zeroth law). Standard proofs assume the Einstein equation with reasonable energy conditions, or vacuum Kerr; see [7, Sec. 12.5]. Thus

$$\kappa(x) = \text{const}\,, \qquad x \in \mathcal{H}_+.$$

Step 2 (Modular structure and KMS). Let $\mathcal{A}(\mathcal{O})$ be the von Neumann algebra for a horizon neighborhood $\mathcal{O}$ invariant under α_t (flow of χ^a). If ω is $(\alpha_t, \beta_{\mathrm{KMS}})$-KMS, Tomita–Takesaki theory implies the modular automorphism group σ_s^ω coincides with α_t after linear reparametrization:

$$\sigma_s^\omega = \alpha_t \quad \text{with } t = \beta_{\mathrm{KMS}}\, s,$$

so that the modular Hamiltonian $\hat{K}_\omega$ and the Noether generator $\hat{H}_\chi$ of the χ^a-flow satisfy

$$\hat{K}_\omega \;=\; \beta_{\mathrm{KMS}}\, \hat{H}_\chi \;+\; c\,\mathbf{1}, \tag{16.481}$$

with c set by normalization ([30, 241, 230]). Within our IQR black-hole setting, $\hat{H}_{\mathrm{mod}} = -\ln \hat{\rho}_{\mathcal{R}_{\mathrm{ext}}}$ (Ch. 15), and the exterior reduced density matrix $\hat{\rho}_{\mathcal{R}_{\mathrm{ext}}}$ is obtained by tracing interior DOFs across the horizon (Definition 16.36).

Step 3 (Near-horizon thermal periodicity: $\beta_{\mathrm{KMS}} = 2\pi/\kappa$). Regularity of the Euclidean section near a Killing horizon fixes the imaginary Killing time to have period $2\pi/\kappa$. The general KMS–Euclidean correspondence then yields

$$\beta_{\mathrm{KMS}} = \frac{2\pi}{\kappa}, \qquad T_H = \frac{1}{\beta_{\mathrm{KMS}}} = \frac{\kappa}{2\pi}.$$

In our framework, Ch. 15 implements this via the Unruh–Hawking identification for the exterior wedge: the reduced density matrix acquires the Gibbs form

$$\hat{\rho}_{\mathcal{R}_{\mathrm{ext}}} \propto \exp\Big[-(2\pi/\kappa)\, \hat{H}_{\mathrm{mod}}\Big],$$

with κ the surface gravity of the stationary horizon; for Kerr, the explicit $T_{\mathrm{Kerr}} = \kappa/2\pi$ is given in Proposition 16.42 (Ch. 15). Because κ is constant by Step 1, T_H is constant on $\mathcal{H}_+$ as well.

Step 4 (Identification with the modular temperature T_{mod}). By definition (Proposition 16.59), the horizon-intrinsic modular temperature is

$$T_{\mathrm{mod}}(x) := \frac{\delta\langle \hat{H}_{\mathrm{mod}}(x)\rangle}{\delta S(x)}.$$

For KMS/Gibbs states, the well-known variational identity $\delta\langle K\rangle = \delta S$ (the differential form of the first law of entanglement) holds for the modular Hamiltonian K ([241, Ch. VIII], [30, Ch. 5]). Combining this local identity with (16.481) gives, pointwise on $\mathcal{H}_+$,

$$T_{\mathrm{mod}}(x) = \frac{1}{\beta_{\mathrm{KMS}}} = \frac{\kappa}{2\pi}.$$

This step is consistent with Ch. 15's identification of the modular Hamiltonian with the physical generator for near-horizon observers and with the construction of $\hat{\rho}_{\mathcal{R}_{\mathrm{ext}}}$ by tracing across the horizon (Definition 16.36).

Step 5 (Uniformity on $\mathcal{H}_+$). By Steps 1–4, $T_{\mathrm{mod}}(x) = \kappa/2\pi = \text{const}$ on $\mathcal{H}_+$. This proves the claimed uniformity. The construction is anchored in the emergent geometry of Ch. 6 (legitimizing the Killing-horizon formalism), the exterior/interior split and modular identification of Ch. 15, and the general KMS algebraic framework ([230, 241, 30]).

Remarks (scope, redshift, and sector structure). (i) Away from the horizon, static observers see the Tolman-redshifted local temperature $T_{\mathrm{loc}}(x) = T_H/\sqrt{-\chi^2(x)}$; T_{mod} is horizon-intrinsic (defined on the horizon algebra) and therefore unaffected by this factor. (ii) The no-topology-change restriction isolates the stationary sector; sector jumps consistent with the Second/Third Laws (e.g., mergers) are handled via the topological quantization clause in Definition 16.44 and its corollaries (Ch. 15). (iii) The entire construction is consistent with the IQR emergence chain (Ch. 6), where $\bar{g}_{\mu\nu}$ and $\bar{t}_{\mu\nu}$ deliver a complete infinitesimal description of the chronogeometry and support the horizon/wedge decomposition used to define $\hat{\rho}_{\mathcal{R}_{\mathrm{ext}}}$ and $\hat{H}_{\mathrm{mod}}$. □

Proposition 16.60 (First Law: Conservation of Relational Energy)**.** *The first law of black hole thermodynamics is an expression of the conservation of relational energy under an infinitesimal reconfiguration of the Kerr attractor's modal entanglement structure. An infinitesimal change in the emergent relational energy, $\bar{E}$, is given by:*

$$\boxed{\delta\bar{E} = T_{\mathrm{mod}}\,\delta S_{\mathrm{ent}} + \bar{\Omega}\,\delta\bar{J},} \tag{16.482}$$

where:

- **(i)** $\bar{E} := \langle \hat{H}_{\mathrm{rel}}\rangle$ *is the expectation value of the emergent relational Hamiltonian.*
- **(ii)** S_{ent} *is the total intermodal entropy across $\mathcal{H}_+$, as defined in Definition 16.36.*
- **(iii)** $\bar{\Omega}$ *is the emergent modular angular velocity, and $\bar{J}$ is the expectation value of the relational angular momentum operator from Definition 16.8.*

This law describes the coherent redistribution of energy and entanglement across the modal sectors of the black hole attractor.

Proof of Proposition Proposition 16.60. The proposition asserts that an infinitesimal change in the emergent relational energy of the Kerr attractor is given by $\delta\bar{E} = T_{\mathrm{mod}}\,\delta S_{\mathrm{ent}} + \bar{\Omega}\,\delta\bar{J}$.

1. The Conserved Quantity: Relational Energy We begin with the total emergent relational energy of the Kerr attractor, defined as the expectation value of the relational Hamiltonian:

$$\bar{E} = \langle \overset{\circ}{\mathbf{\Psi}} | \hat{H}_{\text{rel}} | \overset{\circ}{\mathbf{\Psi}} \rangle.$$

We consider a quasi-static process that shifts the system from one stationary Kerr state to an infinitesimally close one. The total change in energy, $\delta\bar{E}$, must be accounted for by the physical processes that affect the state.

2. Decomposition of Relational Change Within the IQR framework, a change to the Kerr attractor state can be decomposed into two distinct types of relational transformation:

(i) A change in the entanglement structure across the horizon.
This involves altering the fine-grained correlations between the interior and exterior PQS ensembles. This is an inherently statistical, thermodynamic process.

(ii) A change in the coherent rotational state of the PQS ensemble.
This involves altering the collective angular momentum of the system. This is a mechanical, work-like process.

3. The Thermodynamic "Heat" Term We have established in Proposition 16.42 that the Kerr horizon possesses a well-defined modular temperature, T_{mod}. From the fundamental principles of thermodynamics, the change in energy associated with a change in entropy δS_{ent} at a constant temperature is the heat transfer, δQ. Therefore, the energy change associated with altering the entanglement structure is:

$$\delta E_{\text{entropic}} = \delta Q = T_{\text{mod}}\, \delta S_{\text{ent}}. \tag{16.483}$$

Here, S_{ent} is the total relational entropy of the horizon, as defined in Definition 16.36.

4. The Rotational "Work" Term We have defined the emergent angular momentum $\bar{J}$ as the expectation value of the relational angular momentum operator $\hat{J}$ (Definition 16.8). The conjugate variable to angular momentum is angular velocity, which we have defined as the emergent modular angular velocity, $\bar{\Omega}$. The energy required to change the angular momentum of a system by $\delta\bar{J}$ is the rotational work done on it, δW. Therefore, the energy change associated with altering the rotational state is:

$$\delta E_{\text{rotational}} = \delta W = \bar{\Omega}\, \delta\bar{J}. \tag{16.484}$$

5. Conservation and Synthesis By the principle of conservation of energy, the total change in the emergent relational energy of the system must be the sum of the energy changes from these two distinct processes:

$$\delta\bar{E} = \delta E_{\text{entropic}} + \delta E_{\text{rotational}}.$$

Substituting the expressions from steps 3 and 4, we arrive at the First Law:

$$\delta\bar{E} = T_{\text{mod}}\, \delta S_{\text{ent}} + \bar{\Omega}\, \delta\bar{J}.$$

This completes the proof. The First Law of Black Hole Thermodynamics is shown to be a direct consequence of applying the law of conservation of energy to the two fundamental modes of change—entropic and rotational—that are permissible within the relational structure of the Kerr attractor. □

Proposition 16.61 (Second Law: Monotonic Increase of Relational Entropy)**.** *Under any coarse-grained parasymmetric evolution that involves decoherence, the relational entropy of the event horizon, $S(\mathcal{H}_+)$, is a non-decreasing function of emergent relational time:*

$$\boxed{\frac{d}{dt}S(\mathcal{H}_+) \geq 0.} \tag{16.485}$$

The entropy is defined as the von Neumann entropy of the reduced density operator for the horizon, as established in Definition 16.36. This law expresses the irreversible growth of modal indistinguishability as relational coherence becomes inaccessible across emergent causal boundaries.

Proof of Proposition Proposition 16.61. The proposition asserts that for any coarse-grained evolution involving decoherence, the relational entropy of the event horizon, $S(\mathcal{H}_+)$, is a non-decreasing function of emergent time, $dS(\mathcal{H}_+)/dt \geq 0$.

1. The Setup: A Coarse-Grained Process Let the initial state of the Kerr attractor be described by the reduced density operator $\hat{\rho}_{\mathcal{H}}$. We consider a process where the black hole interacts with an external system (e.g., an infalling PQS configuration), which we will call the "ancilla," described by the density operator $\hat{\rho}_A$. The initial state of the combined system is $\hat{\rho}_{\text{initial}} = \hat{\rho}_{\mathcal{H}} \otimes \hat{\rho}_A$, assuming no prior entanglement between the black hole and the ancilla.

2. Unitary Evolution of the Total System The entire process, including the interaction and absorption of the ancilla by the black hole, is governed by the global parasymmetric evolution of the Quantum Holo-State. For the combined subsystem of the black hole and ancilla, this evolution is described by a unitary operator, $\hat{U}$. The final state of the combined system after the interaction is:

$$\hat{\rho}_{\text{final}} = \hat{U}(\hat{\rho}_{\mathcal{H}} \otimes \hat{\rho}_A)\hat{U}^\dagger. \tag{16.486}$$

3. The Property of Unitarity and Entropy A fundamental property of von Neumann entropy is that it is invariant under unitary transformations. Therefore, the entropy of the total system is conserved:

$$S(\hat{\rho}_{\text{final}}) = S(\hat{\rho}_{\mathcal{H}} \otimes \hat{\rho}_A) = S(\hat{\rho}_{\mathcal{H}}) + S(\hat{\rho}_A), \tag{16.487}$$

where the second equality holds because the initial systems were uncorrelated.

4. The Coarse-Graining Step and Subadditivity The "coarse-grained evolution" means that after the interaction, we consider the black hole as a single system again. The new state of the black hole, $\hat{\rho}'_{\mathcal{H}}$, is found by tracing out the degrees of freedom of the absorbed ancilla from the final combined state, $\hat{\rho}_{\text{final}}$. This is the decoherence step where information becomes inaccessible. By subadditivity of von Neumann entropy,

$$S(\hat{\rho}_{\text{final}}) \leq S(\hat{\rho}'_{\mathcal{H}}) + S(\hat{\rho}'_A). \tag{16.488}$$

5. Synthesis and Conclusion Combining the results,

$$S(\hat{\rho}_{\mathcal{H}}) + S(\hat{\rho}_A) \leq S(\hat{\rho}'_{\mathcal{H}}) + S(\hat{\rho}'_A) \quad \Rightarrow \quad \Delta S_{\mathcal{H}} := S(\hat{\rho}'_{\mathcal{H}}) - S(\hat{\rho}_{\mathcal{H}}) \geq S(\hat{\rho}_A) - S(\hat{\rho}'_A) \geq 0.$$

Thus $S(\mathcal{H}_+)$ can only increase (or remain constant) during any process involving interaction and subsequent coarse-graining. This completes the proof. □

Proposition 16.62 (Third Law: Modal Inaccessibility of Extremal States)**.** *The extremal Kerr configuration, which corresponds to a state of zero modular temperature ($T_{\text{mod}} = 0$), cannot be reached through any finite sequence of physical processes. Such a state represents a limit of vanishing relational entropy flux:*

$$\boxed{\lim_{t\to\infty} T_{\text{mod}}(t) \to 0 \quad \Rightarrow \quad \lim_{t\to\infty} S(\mathcal{H}_+)(t) = const > 0.} \tag{16.489}$$

The extremal state corresponds to a perfectly factorized modal configuration, which is an idealization that is dynamically unreachable due to the persistent entanglement of the black hole with environmental modes within the Quantum Holo-State.

Proof of Proposition Proposition 16.62. The proposition asserts that an extremal Kerr state, characterized by zero modular temperature ($T_{\text{mod}} = 0$), is unattainable through any finite sequence of physical processes.

1. The Nature of the Extremal State From Proposition 16.42, we have established that the modular temperature is directly proportional to the surface gravity, $T_{\text{mod}} = \kappa/(2\pi)$. An extremal state with $T_{\text{mod}} = 0$ is therefore a state with zero surface gravity.

2. Entropic Inertia at the Extremal Limit From the First Law (Proposition 16.60), $\delta\bar{E} = T_{\text{mod}}\,\delta S_{\text{ent}} + \bar{\Omega}\,\delta\bar{J}$. Approaching $T_{\text{mod}} \to 0$ decouples $\delta\bar{E}$ from δS_{ent}, rendering the state entropically inert.

3. Non-Factorizability of the Quantum Holo-State According to Definition 16.36, S_{ent} measures entanglement between the black-hole subsystem and its environment. The IQR axiom of non-factorizability (Section 4.8.2) forbids perfect disentanglement of any proper subsystem from the universal Holo-State. Hence a perfectly factorized extremal state cannot be realized by any finite physical process.

4. Conclusion Any procedure attempting to reach $T_{\mathrm{mod}} = 0$ can only approach it asymptotically; a strictly positive residual entropy remains, consistent with (16.489). □

Corollary 16.63 (Topological Residual Entropy (Third-Law Refinement))**.** *Under the Topological Quantization of Entropy (Definition 16.44), the residual horizon entropy appearing in eq.* (16.489) *is fixed by the entanglement-sector label n via the area spectrum:*

$$\boxed{\lim_{t\to\infty} S(\mathcal{H}_+)(t) \;=\; \frac{k_B}{4G\hbar}\,A_n, \qquad n \in \mathbb{Z}_+,} \tag{16.490}$$

where $A_n \in \mathrm{Spec}(\hat{A}_{\mathcal{H}})$ are the eigenvalues specified in Definition 16.44. Consequently, for any finite process that preserves the entanglement-sector topology (no graph surgery), this residual value is invariant.

Remark 16.64 (Scope and Semiclassical Matching)**.** For finite processes that preserve the entanglement-sector topology (no graph surgery), the residual value $\lim_{t\to\infty} S(\mathcal{H}_+)(t) = \frac{k_B}{4G\hbar}A_n$ is invariant. When a sector change does occur (e.g., mergers or topology-changing transitions), n can jump and the spectrum A_n can increase, consistent with the Second Law. In the semiclassical limit, $A_n \to A$ and $S \to k_B A/(4G\hbar)$, recovering the Bekenstein–Hawking relation within the IQR framework.

Thermodynamics as the Dialectic of the Relational Algebra Within IQR, the *laws of thermodynamics* are not merely phenomenological constraints but arise from the dialectic of the relational operator algebra: entropy measures relational indistinguishability of subsystem algebras, while temperature records the statistical intensity of modular (KMS) flow that renders coherence operationally inaccessible. In this sense, black hole thermodynamics is the grammar governing topological evolution on the operator–valued fabric of reality, and it must match the general statement of the laws in Section 15.1 when specialized to stationary horizons.

Zeroth Law (Horizon Uniformity of Modular Temperature). For a stationary Kerr horizon with generator χ^a and exterior KMS state, Proposition 16.59 establishes $T_{\mathrm{mod}}(x) = \kappa/2\pi =$ const ($x \in \mathcal{H}_+$), identifying the horizon–intrinsic modular temperature with Hawking's value via the KMS–modular correspondence. Conceptually, this is the equilibrium clause of the relational algebra: a single modular orbit (the χ^a–flow) thermalizes the entire horizon algebra, prohibiting entanglement–energy currents along $\mathcal{H}_+$. It refines the general equilibrium principle in Section 15.1.2 by tying the uniformity directly to the constancy of surface gravity on Killing horizons and to the algebraic modular flow.

First Law (Relational Energy Balance). Proposition 16.60 expresses the quasi–static balance $\delta\bar{E} = T_{\mathrm{mod}}\delta S_{\mathrm{ent}} + \bar{\Omega}\,\delta\bar{J}$, splitting changes of the emergent energy into an *entropic* (modular/KMS) contribution and a *mechanical* (rotational) contribution. In IQR this reads: perturbations of the horizon algebra either re–pattern cross–horizon correlations (heat) or re–phase the coherent rotational sector (work). It is the black–hole specialization of the energy accounting developed abstractly in Section 15.1, with the role of "temperature" and "entropy" played by the modular generator and the horizon von Neumann entropy Definition 16.36.

Second Law (Irreversibility Under Coarse–Graining). Proposition 16.61 shows $dS(\mathcal{H}_+)/dt \geq 0$ for any coarse–grained (decohering) evolution: unitary dynamics of the global Holo–State together with tracing over inaccessible modes forces the horizon entropy to be non–decreasing. Physically, the horizon is the relational interface where information becomes operationally unavailable to exterior observers; algebraically, subadditivity and KMS stability ensure that coarse–graining cannot reduce $S(\mathcal{H}_+)$. This sharpens the abstract IQR second law in Section 15.1 by identifying the horizon algebra as the natural "thermodynamic boundary'' for black holes.

Third Law (Unattainability and Residual Topology). Proposition 16.62 recasts unattainability as *modal inaccessibility*: $T_{\mathrm{mod}} \to 0$ cannot be reached in finitely many physical steps because perfect factorization of the black–hole subsystem from the Holo–State is forbidden by non–factorizability. Hence extremality is only approached asymptotically, with a strictly positive residual entropy fixed by entanglement–sector topology (Corollary 16.63). Compared with the abstract third law in Section 15.1, the black–hole refinement identifies the protected residual as a *topological invariant* (no graph surgery), reproducing the Bekenstein–Hawking limit in the semiclassical regime.

Why These Laws Matter (Conceptual Synthesis).

(i) Bridge to observation. The laws explain why distant detectors see Hawking radiation as thermal (local coarse–graining) while the global Holo–State evolves unitarily—resolving the "thermal vs unitary'' tension by viewpoint, not by contradiction (Section 16.1.1).

(ii) Consistency across scales. Specializing the general thermodynamic postulates of Section 15.1 to the horizon algebra yields the familiar GR theorems (constancy of κ, area/entropy monotonicity) as operator–algebraic statements about modular flow and coarse–graining.

(iii) Topology as memory. The refined third law ties residual entropy to sector labels, giving a falsifiable prediction: sector–preserving processes keep $S(\mathcal{H}_+)$ fixed at the quantized value, while mergers/topology change allow jumps consistent with the second law (see Corollary 16.63).

In short, the horizon versions of the *zeroth–to–third* laws (Propositions 16.59 to 16.62) are faithful images—within the Kerr sector—of the abstract laws developed in Section 15.1, with the modular/KMS dictionary supplying the precise identification of temperature and entropy at the level of the horizon algebra.

16.7 The Tau-Kairos at Horizons: Structure, Dynamics, and Thermodynamics

Classically, black-hole horizons are geometric null surfaces whose mechanics mirror thermodynamics [86, 87, 245, 649]. In the IQR framework, the horizon is a *phase boundary* of the unified chronogeometric medium, where the universal Holo-State organizes two constitutive fluctuations of the unified tensor $\hat{\Sigma}_{\mu\nu} = \hat{g}_{\mu\nu} + \hat{t}_{\mu\nu}$: the spatial/metric graviton $\hat{h}_{\mu\nu} := \delta\hat{g}_{\mu\nu}$

and the *temporal* excitation $\hat{\tau}_{\mu\nu} := \delta\hat{t}_{\mu\nu}$ (eq. (10.100)). The latter is the **Tau-Kairos tensor**. It carries coherence/phase dynamics, not classical curvature, and interacts with the graviton through a state-dependent matrix propagator (Sections 10.17.2 and 10.17.5). The goal of this section is to place the tau-kairos tensor on the black-hole stage, deriving its horizon-scale kinematics, its impact on ergoregions and temperature, and its imprint on the generalized laws formulated earlier (Section 16.6).

Standing Hypotheses and Computational Regime

We work under the standard black-hole hypotheses already in use:

(H1) Stationary coherence. The Kerr attractor is a stationary coherence phase in the IQR sense (Definition 16.21); the relevant invariance is with respect to the horizon Killing flow χ^a, not a fundamental time parameter.

(H2) Sector preservation. No topology change occurs on $\mathcal{H}_+$: the entropy spectrum respects the sector-preserving clauses of Definition 16.44.

(H3) Near-horizon KMS. The exterior algebraic state restricted to near-horizon regions is KMS with respect to the χ^a–flow (α_t) with inverse temperature β_H determined by the (effective) surface gravity (eqs. (16.289) and (16.319)); cf. the modular/KMS dictionary [608, 230, 241].

Constitutive Definition and Spin Structure

By definition (eq. (10.100)), the tau-kairos tensor is the fluctuation of the temporal block:

$$\hat{\tau}_{\mu\nu}(x) = \delta\hat{t}_{\mu\nu}(x), \qquad \delta\hat{\Sigma}_{\mu\nu} = \hat{h}_{\mu\nu} + \hat{\tau}_{\mu\nu}.$$

Its Lorentz content is fixed by:

Theorem 16.65 (Spin Content, Specialized to the Horizon Sector)**.** *The tau-kairos tensor on a stationary, axisymmetric background obeys*

$$\mathcal{P}^{(2)}[\hat{\tau}_{\mu\nu}] = 0,$$

and hence contains only scalar and vector polarizations (spin-0 ⊕ spin-1). Equivalently, the TT channel is exhausted by the spin-2 graviton $\hat{h}_{\mu\nu}$.

Proof. This is the restriction of Section 10.5 to the Kerr sector with horizon generator χ^a. The gauge constraints and relational symmetries that annihilate the spin-2 projector of $\hat{\tau}_{\mu\nu}$ (proved generally in Section 10.17.9) remain valid under (H1)–(H3); axisymmetry provides the spectral gap that forbids TT support in the temporal block. □

Field-of-Origin Refinement. As developed in the global theory, the scalar part $\hat{\tau}^{(0)}_{\mu\nu}$ admits a geometric (Gaiamiti/Apeiron) provenance, whereas the vector part $\hat{\tau}^{(1)}_{\mu\nu}$ admits a material (Tantrakyma/Chrono-Kymon) provenance; both project into $t_{\mu\nu}$ and participate in the unified dynamics without entering the TT channel. We retain this refinement below when interpreting near-horizon observables.

Kerr Background, Unified Ansatz, and Horizon/Ergosurface Conditions

On the stationary axisymmetric Kerr-like background, the unified split and temporal ansatz are (eqs. (16.198) and (16.199))

$$\bar{\Sigma}_{\mu\nu} = \bar{g}^{\text{Kerr}}_{\mu\nu} + \bar{t}_{\mu\nu}, \tag{16.491}$$

$$\begin{aligned}\mathcal{T}_{\mu\nu}\, dx^\mu dx^\nu &= \bar{t}_{\mu\nu}\, dx^\mu dx^\nu \\ &\sim -A\, dt^2 + B\, \frac{\rho^2}{\Delta}\, dr^2 + C\, \rho^2\, d\theta^2 \\ &\quad - 2D\, \sin^2\theta\, dt\, d\phi + E\, \sin^2\theta\, d\phi^2.\end{aligned} \tag{16.492}$$

The *unified ergosurface* and *horizon* are determined by (eqs. (16.202) and (16.203))

$$\bar{\Sigma}_{tt} = \bar{g}^{\text{Kerr}}_{tt} - A = 0, \qquad \bar{\Sigma}_{ab}\chi^a\chi^b\Big|_{r=R_H} = 0, \quad \Omega_H = -\frac{\bar{\Sigma}_{t\phi}}{\bar{\Sigma}_{\phi\phi}}\Big|_{r=R_H}.$$

Thus, the temporal lapse A directly displaces the ergosurface relative to the classical condition $g_{tt} = 0$ (eq. (16.228)), while the mixed component D feeds the frame-dragging sector. In the static limit, the horizon condition reduces to the local phase-boundary equation (eq. (16.16)); the Boyer–Lindquist realization is eq. (16.21).

Relational Propagator and Spatial–Temporal Mixing

Linearized fluctuations about $\bar{\Sigma}_{\mu\nu}$ are governed by the matrix Green's function $\mathbf{G}^{(2)}_{\Sigma}$ with graviton/tau-kairos tensor off-diagonal entries (eq. (10.112)). The essential point for horizons is that $\mathbf{G}^{(2)}_{\Sigma}$ preserves the TT content exclusively in the $\hat{h}$–$\hat{h}$ block, while allowing state-dependent *conversion amplitudes* between spatial and temporal sectors off the light-cone determined by $\bar{\Sigma}$. This encodes how coherence (temporal sector) back-reacts on near-horizon geometry without generating new long-range spin-2 forces.

Entropy, Temperature, and the Entropic Dual of the Ergosphere

The generalized laws of black-hole thermodynamics developed earlier (Section 16.6) are recovered here through the KMS dictionary (eqs. (16.289) and (16.319)) and the entropic interpretation of rotating horizons (Section 16.4.10). In particular:

(i) The **temperature** is fixed by the *effective* surface gravity κ_{eff} of the unified background (eq. (16.2)), hence

$$T_{\text{eff}} = \frac{\hbar\kappa_{\text{eff}}}{2\pi k_B}.$$

In the classical Kerr limit one recovers κ and T_{Kerr} (eqs. (16.291) and (16.292)).

(ii) **Tau–Kairos tensor corrections** enter T_{eff} only through κ_{eff}, i.e. through the background profiles $A, B, \ldots$ at R_H. The first–order shift is given by eq. (16.154).

(iii) The **entropy** receives a linear response proportional to the horizon shift δR_H (and hence to the temporal coefficients) as in eqs. (16.191) and (16.192).

Operationally, the *entropic dual* picture (ergosphere as a dynamical interface, Definition 16.34) ties the emergence of horizon temperature to modal information flow across the coherence interface; the tau-kairos tensor controls precisely those coherence gradients.

Two Benchmark Results in Linear Response

We record the two formulae most relevant for phenomenology and for comparison with the computations in Section 16.5.

Proposition 16.66 (Horizon Shift from the Temporal Lapse)**.** *In the static limit and to first order in the temporal corrections, the event-horizon radius shifts by*

$$\delta R_H = f_1(R_S) - R_S\, a_1(R_S),$$

equivalently $\delta R_{\mathrm{EH}} \approx f_1(R_S) - R_S\, A\big(\xi(R_S)\big)$*, as stated in eq.* (16.73)*. Consequently,* S_{BH} *acquires the linear correction of eqs.* (16.191) *and* (16.192)*.*

Proof. Expand the phase-boundary condition (eq. (16.21)) at R_S and identify the horizon-value Taylor coefficients (a_1, f_1). The entropy variation is the usual area variation evaluated on $\bar{\Sigma}$ to first order, which is equivalently obtained from the corrected lapse via the first law (in its relational/KMS form) [86, 245]. □

Proposition 16.67 (Surface Gravity and Temperature Shift)**.** *The effective surface gravity of the unified background satisfies, to first order,*

$$\kappa_{\mathrm{eff}} = \kappa_S + \frac{c^2}{2}\,\varepsilon\left[\frac{f_1'(R_S)}{R_S} - \frac{f_1(R_S)}{R_S^2} - a_1'(R_S)\right] + \mathcal{O}(\varepsilon^2),$$

as in eq. (16.154)*; hence* $T_{\mathrm{eff}} = \hbar\kappa_{\mathrm{eff}}/(2\pi k_B)$*. In the rotating case, the same structure holds with the replacements dictated by eqs.* (16.200) *and* (16.203)*.*

Proof. Differentiate the defining functions F, G of eq. (16.2) at the corrected root $R_H = R_S + \delta R_H$, substitute δR_H from Proposition 16.66, and retain $\mathcal{O}(\varepsilon)$ terms. Rotating corrections enter only via the background coefficients evaluated on the stationary axisymmetric solution (cf. eqs. (16.199) and (16.203)). □

Ergosphere, Vorticity, and Information Flow

Gradients of the *Tau-Kairos tensor* $\hat{\tau}_{\mu\nu}$—captured by the coherence–gradient invariants A and D (cf. Definition 16.13)—generate *chronogeometric vorticity*, displacing the ergosurface relative to the classical locus $\bar{g}_{tt} = 0$ (eq. (16.228)). In the *quantum ergodynamics* picture (Section 16.4.10), this ergospheric interface mediates mode exchange between exterior and near-horizon degrees of freedom; its contractive dynamics with respect to the modular flow underwrite both the KMS state and the generalized second law [608, 230, 241, 245]. Crucially, the Tau-Kairos tensor is the unique carrier of these coherence gradients—by construction it is TT-null, $\mathcal{P}^{(2)}[\hat{\tau}_{\mu\nu}] = 0$ (see Definition 10.13, Lemma 10.123, and eq. (10.65))—so gravitational waves propagate purely in the tensor channel, while entropic/thermal features vary with the local coherence encoded in $\hat{\tau}_{\mu\nu}$.

Phenomenological Notes and Observational Windows

(i) Near-horizon spectroscopy. Quasi-normal spectra sensitive to background A, D profiles can be used to bound δR_H and hence δS at linear order (via Proposition 16.66); this affects ringdown phasing without introducing extra TT polarizations.

(ii) Ergoregion transport. The unified ergosurface condition ($\bar{\Sigma}_{tt} = 0$) implies coherence-dependent shifts of the Penrose process threshold and superradiant windows; these are encoded by A and D at fixed (M, a).

(iii) Thermal channel. Corrections to κ_{eff} (Proposition 16.67) induce small changes in the Hawking temperature and greybody factors; in the rotating case, these inherit the angular velocities of eq. (16.203).

Synthesis

The tau-kairos tensor $\hat{\tau}_{\mu\nu}$ is the temporal, coherence-carrying counterpart of the graviton. It has no TT content (Theorem 16.65), yet it *does* control precisely those unified background features that determine horizon location, surface gravity, and the entropic dual of the ergosphere. In the IQR picture, black-hole thermodynamics thus emerge from a two-channel interplay: the *geometric* (TT, $\hat{h}$) channel governing classical propagation and the *temporal* (scalar/vector, $\hat{\tau}$) channel governing coherence, KMS structure, and entropy flow. The generalized laws (Section 16.6) are the macroscopic closure of this two-channel dynamics.

A black hole's geometry, in the framework under consideration, is described not only by a metric $g_{\mu\nu}$ but by a unified chronogeometric tensor $\Sigma_{\mu\nu} = g_{\mu\nu} + t_{\mu\nu}$. Ordinary gravitons are fluctuations of the metric block $g_{\mu\nu}$ and carry the familiar transverse–traceless (spin-2) gravitational radiation. By contrast, a *Tau-Kairos tensor* is the fluctuation of the temporal/coherence block $t_{\mu\nu}$:

$$\delta\Sigma_{\mu\nu} = \underbrace{\delta g_{\mu\nu}}_{\text{graviton } \hat{h}_{\mu\nu}} + \underbrace{\delta t_{\mu\nu}}_{\text{tau-kairos tensor} \hat{\tau}_{\mu\nu}} .$$

It is not a new spin-2 particle. Its allowed polarizations are scalar and vector (spin-0/1), and it couples to *quantum coherence and phase flow* rather than directly to classical stress–energy. In practical terms, it encodes how the "temperature-bearing" and information-carrying aspects of the near-horizon state vary, while leaving the propagation of ordinary gravitational waves unchanged.

This clarifies a common confusion: *a tau-kairos tensor is not Hawking radiation.* Hawking radiation is a flux of asymptotic quanta (photons, gravitons, neutrinos, etc.) produced because the quantum vacuum is "split" by the horizon; it is, in principle, countable by a distant detector. A Tau-Kairos tensor, by contrast, resides in the *background that sets the stage* for this process. It modulates the *effective surface gravity* and the *modular/KMS structure* of the exterior state; those, in turn, set the Hawking temperature and shape the greybody factors. A useful analogy: Hawking quanta are the *steam* leaving a kettle; the tau-kairos tensor is the controlled adjustment of the *thermostat* and *pressure valve* that determine how vigorously the water boils. They interact, but they are not the same thing.

Their concrete relations appear in three places. First, via temperature: tau-kairos tensor backgrounds shift the effective surface gravity κ_{eff}, hence $T_{\text{eff}} = \hbar\kappa_{\text{eff}}/(2\pi k_B)$. Second, via *entropy flow*: by displacing the unified ergosurface and slightly shifting the horizon radius, they induce linear corrections to the horizon area and therefore to thermodynamic accounting. Third, via *spectral shaping*: scalar/vector tau-kairos tensor profiles alter near-horizon scattering and superradiant windows, leading to small, coherence-dependent changes in greybody factors—subtle distortions of the Hawking spectrum without introducing extra gravitational-wave polarizations. These effects are most pronounced in coherence-rich regimes (early-time horizons, rapidly rotating black holes, analogue/near-critical media) and fade in decohered macroscopic limits, where classical GR is recovered.

An intuitive organizing principle is the *two-channel picture*. One channel—the geometric TT channel—governs curvature and radiative gravity (ordinary gravitons, ringdowns, tidal fields). The other—the temporal coherence channel—governs modular flow, effective temperature, and entropic exchange (Tau-Kairos tensors). The channels mix kinematically through a matrix propagator, yet maintain a strict division of labor: TT radiative content remains in the graviton channel, while thermo-informational content is carried by the tau-kairos tensor channel.

Observable signatures would therefore emphasize *phase-accurate ringdown shifts* consistent with tiny horizon/ergosurface displacements; *coherence-dependent adjustments* to superradiance thresholds; and, in principle, *minute spectral distortions* in the Hawking flux attributable to T_{eff} and greybody modifications. The absence of such signatures would also be consistent with the construction: the tau-kairos tensor naturally decouples in the classical, decohered limit, leaving standard black-hole physics intact while providing a controlled handle on the quantum-thermodynamic layer that classical GR does not track.

Cross-References. For the detailed Kerr construction, see Sections 16.4 and 16.5; for the ergosphere's dynamical interpretation and the emergence of temperature, see Section 16.4.10; for lifecycle implications (information transfer and unitary evolution), proceed to Section 16.11.

16.8 Chrono-Kymons at Horizons: Vector Coherence, Vorticity, and Rotational Thermodynamics

Black-hole backgrounds in the IQR formalism are encoded by the unified chronogeometric tensor

$$\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}, \qquad \delta\hat{\Sigma}_{\mu\nu} = \underbrace{\delta\hat{g}_{\mu\nu}}_{\hat{h}_{\mu\nu}\ \text{(graviton, TT)}} + \underbrace{\delta\hat{t}_{\mu\nu}}_{\hat{\tau}_{\mu\nu}\ \text{(Tau-Kairos tensor, scalar}\oplus\text{vector)}}, \tag{16.493}$$

where the spin-2, transverse–traceless (TT) content resides entirely in the graviton sector $\hat{h}_{\mu\nu}$. The *chrono-kymon* is the vector (spin-1) constituent of the temporal fluctuation $\hat{\tau}_{\mu\nu}$ and is responsible for the antisymmetric/axial part of the temporal block,

$$\hat{t}_{[\mu\nu]} \equiv \tfrac{1}{2}(\hat{t}_{\mu\nu} - \hat{t}_{\nu\mu}) \neq 0, \qquad \hat{\tau}_{\mu\nu} = \hat{\tau}^{(0)}_{\mu\nu} \oplus \hat{\tau}^{(1)}_{\mu\nu}, \quad \hat{\tau}^{(1)}_{\mu\nu} \text{ carried by chrono-kymons.} \tag{16.494}$$

At the level of operator fields, chrono-kymons arise as massive vector excitations of the Tantrakyma sector (the material component of the unified operator bundle) and feed the near-horizon *frame-dragging* and *coherence-flow* structure without introducing any additional TT gravitational-wave polarization.

Standing Hypotheses and Near-Horizon Regime

The analysis follows the standard assumptions already used for the Tau-Kairos tensor:

(H1) Stationary coherence. A Kerr attractor is a stationary coherence phase relative to the horizon Killing flow χ^a (Definition 16.21).

(H2) Sector preservation. No topology change occurs on $\mathcal{H}_+$; sector-preserving clauses for the entropy spectrum hold (Definition 16.44).

(H3) Near-horizon KMS. The exterior algebraic state is (α_t, β_H)-KMS with respect to χ^a; the modular/KMS dictionary fixes the thermodynamic identifications (eqs. (16.289) and (16.319)).

Constitutive Definition and Effective Equations

Introduce the Tantrakyma vector operator $\hat{\mathcal{C}}_\mu$ (the chrono-kymon). In linear response about $\bar{\Sigma}_{\mu\nu}$,

$$\hat{t}_{[\mu\nu]} \;\equiv\; \nabla_{[\mu}\hat{\mathcal{C}}_{\nu]} \;+\; \Xi_{[\mu\nu]}[\bar{\Sigma},\hat{\mathcal{C}}] \;+\; \mathcal{O}(\hat{\mathcal{C}}^2), \tag{16.495}$$

$$\mathcal{E}_\nu[\bar{\Sigma};\hat{\mathcal{C}}] \;\equiv\; \nabla^\mu\nabla_{[\mu}\hat{\mathcal{C}}_{\nu]} \;-\; \mu_C^2\,\hat{\mathcal{C}}_\nu \;+\; \mathcal{M}_\nu{}^\rho[\bar{\Sigma}]\,\hat{\mathcal{C}}_\rho \;=\; \hat{J}_\nu, \tag{16.496}$$

where μ_C is an effective mass gap, $\mathcal{M}_\nu{}^\rho$ collects curvature- and state-dependent couplings, and $\hat{J}_\nu$ denotes composite sources. The spin projector satisfies $\mathcal{P}^{(2)}[\hat{\tau}_{\mu\nu}] = 0$ (Theorem 16.65); hence all TT radiative content remains in $\hat{h}_{\mu\nu}$.

Kerr Ansatz and the Frame-Dragging Coefficient

On a stationary, axisymmetric background, the temporal sector is represented by the near-horizon ansatz (eq. (16.199))

$$\bar{t}_{\mu\nu}dx^\mu dx^\nu \sim -A\,dt^2 \;+\; B\,\frac{\rho^2}{\Delta}\,dr^2 \;+\; C\,\rho^2\,d\theta^2 \;-\; 2D\,\sin^2\theta\,dt\,d\phi \;+\; E\,\sin^2\theta\,d\phi^2, \tag{16.497}$$

where the mixed coefficient D measures temporal–azimuthal coupling. The vector chrono-kymon dominates precisely this sector: to leading order,

$$D \;\propto\; (\mathcal{L}_\chi\bar{\mathcal{C}})_\phi \;+\; \text{state-dependent terms}, \qquad \bar{\Sigma}_{t\phi} = \bar{g}^{\mathrm{Kerr}}_{t\phi} - D\,\sin^2\theta. \tag{16.498}$$

The horizon generator's angular velocity is fixed by the unified relation

$$\Omega_H \;=\; -\,\frac{\bar{\Sigma}_{t\phi}}{\bar{\Sigma}_{\phi\phi}}\bigg|_{r=R_H}, \tag{16.499}$$

so chrono-kymon backgrounds adjust Ω_H through $\bar{\Sigma}_{t\phi}$ (and, indirectly, through $\bar{\Sigma}_{\phi\phi}$ at R_H).

Relational Propagator and Sector Mixing

Linearized fluctuations are governed by the matrix Green function $\mathbf{G}_{\Sigma}^{(2)}$ with blocks $(hh), (h\tau), (\tau h), (\tau\tau)$ (eq. (10.112)). The vector chrono-kymon feeds the $(\tau\tau)$ and off-diagonal $(h\tau)$ entries that couple spatial and temporal sectors. The TT channel remains confined to (hh), so no additional gravitational-wave polarization is produced; the observable footprint is through background and mixing effects (chiefly $\Sigma_{t\phi}$ and Ω_H).

Thermodynamics: KMS Structure, Temperature, and Chemical Potential

The KMS state associated with χ^a determines the effective temperature and the rotational chemical potential. In the unified background,

$$T_{\text{eff}} = \frac{\hbar\,\kappa_{\text{eff}}}{2\pi k_B}, \qquad \tilde{\omega} \;=\; \omega - m\,\Omega_H, \tag{16.500}$$

where κ_{eff} is fixed by $\bar{\Sigma}_{\mu\nu}$ (eq. (16.2)). Chrono-kymons affect T_{eff} only via their (subleading) influence on κ_{eff} but have a leading-order impact on the *rotational shift* through Ω_H (Eq. (16.499)). Consequently, Hawking emission spectra are modified predominantly by the replacement $\omega \mapsto \tilde{\omega}$ with the chrono-kymon–renormalized Ω_H.

16.8.1 Two Benchmark Results in Linear Response

Proposition 16.68 (Angular-Velocity Shift from the Chrono-Kymon Sector)**.** *Let R_H be the unified horizon radius fixed by $\bar{\Sigma}_{ab}\chi^a\chi^b|_{r=R_H} = 0$ (eq. (16.203)). To first order in temporal-vector corrections,*

$$\delta\Omega_H = -\frac{\delta\bar{\Sigma}_{t\phi}}{\bar{\Sigma}_{\phi\phi}}\bigg|_{R_H} + \frac{\bar{\Sigma}_{t\phi}\,\delta\bar{\Sigma}_{\phi\phi}}{\bar{\Sigma}_{\phi\phi}^2}\bigg|_{R_H} \;+\; \partial_r\!\left(\frac{\bar{\Sigma}_{t\phi}}{\bar{\Sigma}_{\phi\phi}}\right)\bigg|_{R_H} \delta R_H \;+\; \mathcal{O}(\varepsilon^2), \tag{16.501}$$

with $\delta\bar{\Sigma}_{t\phi} = -\,\delta D\,\sin^2\theta$ dominated by the chrono-kymon background $\delta\mathcal{C}_\phi$ via (16.498).

Proof. Differentiate (16.499) holding θ fixed on the horizon two-sphere and account for the horizon displacement δR_H from the temporal sector (cf. Proposition 16.66). The contribution proportional to δD is traced through $\bar{\Sigma}_{t\phi}$. □

Proposition 16.69 (Superradiant Window and Spectral Shaping)**.** *For bosonic modes with azimuthal number m, the superradiant condition in the unified background reads*

$$0 < \omega < m\,\Omega_H \quad\Longrightarrow\quad \textit{amplification}, \tag{16.502}$$

with Ω_H given by (16.499). A chrono-kymon–induced $\delta\Omega_H$ shifts the window by $m\,\delta\Omega_H$ at leading order. Hawking spectra acquire coherence-dependent distortions through the replacement $\omega \mapsto \omega - m(\Omega_H + \delta\Omega_H)$ in the occupation factor, while TT gravitational-wave polarizations remain unaffected.

Proof. The derivation follows the standard energy-flux argument at the horizon using the Killing generator $k^a = \chi^a + \Omega_H\,\psi^a$ and the unified stress tensor built on $\bar{\Sigma}_{\mu\nu}$. The replacement $\omega \mapsto \tilde{\omega}$ is inherited from the KMS weight in Eq. (16.500). □

Ergosphere, Chronogeometric Vorticity, and Transport

Chrono-kymon gradients source *chronogeometric vorticity* (Definition 16.13), encoded kinematically by D in (16.497). While the unified ergosurface is determined by $\bar{\Sigma}_{tt} = 0$ (hence primarily sensitive to the lapse A), the transport properties within the ergoregion—ZAMO frames, energy-extraction thresholds, and mode conversion—are controlled by Ω_H and thus by the chrono-kymon sector. The result is a coherence-dependent adjustment of Penrose process efficiencies and of superradiant amplification consistent with Proposition 16.69.

Phenomenological Windows

(i) Ringdown phasing. Vector temporal backgrounds shift Ω_H and $\bar{\Sigma}_{t\phi}$, leading to small but phase-accurate modifications of quasinormal mode frequencies that correlate with $\delta\Omega_H$ (Proposition 16.68) while leaving the TT polarization content unchanged.

(ii) Superradiance control. The superradiant window responds linearly to $m\,\delta\Omega_H$ (Proposition 16.69); coherence-rich regimes maximize the effect (rapid spin, early-time horizons, analogue setups).

(iii) Hawking spectral tweaks. Occupation factors depend on $\tilde{\omega} = \omega - m\Omega_H$; chrono-kymon backgrounds distort the spectrum through $\delta\Omega_H$ and modest κ_{eff} shifts, without producing new asymptotic radiative degrees of freedom.

(iv) Quasi-bound clouds (if $\mu_C \neq 0$). For $\mu_C M = \mathcal{O}(1)$, long-lived vector clouds can arise; their back-reaction appears as persistent D-profiles, producing slow drifts in Ω_H and in late-time tails consistent with the decoupling of the TT channel.

Synthesis and Relation to the Tau-Kairos tensor

Chrono-kymons provide the *vector engine* of the temporal sector: they generate and regulate horizon-scale vorticity and rotational coherence through $\bar{\Sigma}_{t\phi}$ and Ω_H. In conjunction with the scalar temporal sector (Apeiron-linked), they complete the Tau-Kairos spin-0/1 content while remaining orthogonal to the TT gravitational-wave channel. Thermodynamic consequences are mediated chiefly by Ω_H (rotational chemical potential) and, subleadingly, by κ_{eff}; transport and scattering consequences follow from the superradiant condition (16.502). The classical limit is recovered as coherence decouples, leaving standard Kerr thermodynamics and wave propagation intact.

Cross-References. Background and notation: Sections 16.4 and 16.7. Horizon/ergosurface conditions: eqs. (16.202) and (16.203). Thermodynamic dictionary: Section 16.6 and eq. (16.2). For lifecycle implications and information transport, see Section 16.11.

A unified chronogeometric description writes the background as $\Sigma_{\mu\nu} = g_{\mu\nu} + t_{\mu\nu}$. Fluctuations split into the usual spin-2 graviton $\hat{h}_{\mu\nu} = \delta g_{\mu\nu}$ and the temporal/coherence excitation $\hat{\tau}_{\mu\nu} = \delta t_{\mu\nu}$ (the Tau-Kairos tensor). Within the spin decomposition of $\hat{\tau}_{\mu\nu}$, the *vector (spin-1) sector* is sourced by *chrono-kymons*: massive vector excitations of the Tantrakyma field that feed the antisymmetric/axial content of the temporal block $t_{[\mu\nu]}$. In stationary, axisymmetric

black-hole backgrounds, this vector content enters the mixed time–azimuth component $t_{t\phi}$ and thereby the coefficient D of the near-horizon ansatz (the $dt\,d\phi$ term), which governs frame dragging and *chronogeometric vorticity*. In brief: chrono-kymons are the microscopic carriers of the *spin-1 temporal flow* that the tau-kairos tensor (as a composite temporal fluctuation) exposes at the macroscopic level.

This placement determines both kinematics and thermodynamics. Kinematically, chrono-kymons control the *vector part* of the matrix propagator that mixes spatial and temporal sectors in linear response. Because the transverse–traceless channel is exhausted by $\hat{h}_{\mu\nu}$, chrono-kymons do not introduce new spin-2 gravitational-wave polarizations; their influence instead manifests through off-diagonal, state-dependent conversion amplitudes and through background modifications of $\Sigma_{t\phi}$. Thermodynamically, the same vector profiles shift the unified ergosurface condition $\Sigma_{tt} = 0$ and adjust the horizon generator's angular velocity Ω_H as read off from $\Omega_H = -\Sigma_{t\phi}/\Sigma_{\phi\phi}\big|_{r=R_H}$. These shifts feed directly into the *effective surface gravity* $\kappa_{\rm eff}$ and the *KMS/modular* structure that set the Hawking temperature and the greybody factors. Thus, chrono-kymons do not constitute Hawking quanta; rather, they tune the *thermo-informational background* against which Hawking conversion proceeds.

The dynamical role near rotating holes is particularly transparent. A Kerr-like background admits superradiant scattering for bosonic fields. Chrono-kymons, being vectorial, obey a superradiant inequality of the usual form (set by the unified Ω_H) and thereby control the *opening and closing* of superradiant windows through their contribution to D and related temporal coefficients. In the presence of an effective mass scale μ_C for the chrono-kymon, quasi-bound levels can form; their occupation modifies near-horizon scattering and the late-time tail, leading to small, coherence-dependent shifts in ringdown phasing without introducing extra TT polarizations. In parameter regimes where coherence is high (rapid spin, early-time horizons, analogue/near-critical media), these effects are maximized; in decohered macroscales they decouple, leaving classical GR intact.

From an observational standpoint, chrono-kymons are expected to leave signatures in *three* channels: (i) phase-accurate *ringdown* shifts consistent with slight changes in $\Sigma_{t\phi}$ and the unified ergosurface; (ii) *transport* phenomena—minute, coherence-dependent adjustments to *superradiance thresholds* and energy-extraction efficiencies; and (iii) *spectral shaping* of Hawking flux through modest changes to $\kappa_{\rm eff}$ and greybody factors driven by the vector temporal background. All three are background-level (thermo-informational) effects rather than new radiative degrees of freedom at infinity. In summary, chrono-kymons provide the *vector engine* of the temporal sector: they generate and regulate horizon-scale vorticity and coherence flow, and through those channels they connect the Tau-Kairos scalar/vector content to measurable black-hole phenomenology while remaining orthogonal to the spin-2 gravitational-wave channel.

16.9 Unified Temporal Sector at Horizons: Interactions of Tau-Kairos tensors and Chrono-Kymons

The unified chronogeometric background is

$$\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}, \qquad \delta\hat{\Sigma}_{\mu\nu} = \hat{h}_{\mu\nu} + \hat{\tau}_{\mu\nu}, \tag{16.503}$$

with the TT (spin-2) content residing in the graviton $\hat{h}_{\mu\nu} = \delta\hat{g}_{\mu\nu}$, and the temporal/coherence excitation $\hat{\tau}_{\mu\nu} = \delta\hat{t}_{\mu\nu}$ (the *Tau-Kairos tensor*) carrying only scalar $\oplus$ vector modes. The vector constituent of $\hat{\tau}_{\mu\nu}$ is provided by the Tantrakyma field's massive vector excitation, the *chrono-kymon* $\hat{\mathcal{C}}_\mu$. This section formalizes their coupled kinematics, thermodynamic map, and linear-response consequences for Kerr-like horizons.

Kinematics: Field Content, Projectors, and Decomposition

The temporal fluctuation admits the decomposition

$$\hat{\tau}_{\mu\nu} = \hat{\tau}^{(0)}_{\mu\nu} \oplus \hat{\tau}^{(1)}_{\mu\nu}, \qquad \hat{\tau}^{(0)}_{\mu\nu} = S_{\mu\nu}[\hat{\varphi}], \qquad \hat{\tau}^{(1)}_{\mu\nu} = A_{\mu\nu}[\hat{\mathcal{C}}], \tag{16.504}$$

where $S_{\mu\nu}[\hat{\varphi}]$ is the symmetric (scalar) piece sourced by a temporal scalar $\hat{\varphi}$ (Apeiron-linked), and $A_{\mu\nu}[\hat{\mathcal{C}}] = \nabla_{[\mu}\hat{\mathcal{C}}_{\nu]} + \cdots$ is the antisymmetric/axial (vector) piece carried by the chrono-kymon.[11] The selection rule

$$\mathcal{P}^{(2)}[\hat{\tau}_{\mu\nu}] = 0 \tag{16.505}$$

implies that the TT radiative channel is exhausted by $\hat{h}_{\mu\nu}$, ensuring no extra spin-2 polarizations.

Quadratic Action and Matrix Propagator Near the Horizon

Introduce the fluctuation doublet $\hat{\Upsilon}_{\mu\nu} := (\hat{h}_{\mu\nu}, \hat{\tau}_{\mu\nu})^\top$. To second order about $\bar{\Sigma}_{\mu\nu}$,

$$S^{(2)}_{\text{hor}} = \tfrac{1}{2}\int d^4x\,\sqrt{-\bar{\Sigma}}\;\hat{\Upsilon}^{\mu\nu}\begin{pmatrix}\mathcal{D}_{hh} & \mathcal{M}_{h\tau}\\ \mathcal{M}_{\tau h} & \mathcal{D}_{\tau\tau}\end{pmatrix}_{\mu\nu}{}^{\rho\sigma}\hat{\Upsilon}_{\rho\sigma}, \qquad \mathbf{G} \equiv \begin{pmatrix}G_{hh} & G_{h\tau}\\ G_{\tau h} & G_{\tau\tau}\end{pmatrix} = \begin{pmatrix}\mathcal{D}_{hh} & \mathcal{M}_{h\tau}\\ \mathcal{M}_{\tau h} & \mathcal{D}_{\tau\tau}\end{pmatrix}^{-1}, \tag{16.506}$$

with off-diagonal $\mathcal{M}_{\tau h} = \mathcal{M}^\dagger_{h\tau}$ encoding spatial–temporal conversion amplitudes constrained by (16.505). In particular, $\mathcal{P}^{(2)}\mathcal{M}_{\tau h} = 0$, so TT content propagates purely in G_{hh}.

Thermodynamic Map: Effective Temperature and Rotational Chemical Potential

The KMS state associated with the horizon generator χ^a yields

$$T_{\text{eff}} = \frac{\hbar\,\kappa_{\text{eff}}}{2\pi k_B}, \qquad \tilde{\omega} = \omega - m\,\Omega_H, \tag{16.507}$$

where κ_{eff} is determined by $\bar{\Sigma}_{\mu\nu}$ and Ω_H by $\Omega_H = -\bar{\Sigma}_{t\phi}/\bar{\Sigma}_{\phi\phi}\big|_{r=R_H}$ (eqs. (16.203) and (16.499)). Tau-Kairos tensor backgrounds (scalar sector) dominantly shift κ_{eff} and the horizon radius R_H (Propositions 16.66 and 16.67), while chrono-kymon backgrounds (vector sector) dominantly shift Ω_H (Proposition 16.68). Both effects feed Hawking and scattering observables through (16.507).

[11] Additional state-dependent terms (indicated by "$\cdots$") respect the horizon symmetries and vanish in the decohered limit.

Coupled Linear Response: Jointly Induced Shifts

To first order about a stationary, axisymmetric background, the coupled variations read

$$\delta R_H = \delta R_H^{(0)}[\hat{\tau}^{(0)}] \quad \text{(scalar/Tau-Kairos)}, \tag{16.508}$$

$$\delta\kappa_{\text{eff}} = \delta\kappa^{(0)}[\hat{\tau}^{(0)}] + \underbrace{\delta\kappa^{(1)}[\hat{\tau}^{(1)}]}_{\text{mixing-suppressed}}, \tag{16.509}$$

$$\delta\Omega_H = \delta\Omega_H^{(1)}[\hat{\tau}^{(1)}] + \partial_r\left(\frac{\bar{\Sigma}_{t\phi}}{\bar{\Sigma}_{\phi\phi}}\right)_{R_H} \delta R_H \quad \text{(vector/chrono-kymon + scalar back-reaction)}, \tag{16.510}$$

where $\delta R_H^{(0)}$ and $\delta\kappa^{(0)}$ are given by Propositions 16.66 and 16.67, while $\delta\Omega_H^{(1)}$ is given by Proposition 16.68. Equations (16.509)–(16.510) make explicit the *division of labor* and its cross-coupling: scalar temporal structure sets the thermal scale; vector temporal structure sets the rotational chemical potential; mixing communicates small corrections between them.

Mode Conversion and Scattering: Superradiance, Greybody Factors, and Ringdown

For bosonic modes of frequency ω and azimuthal number m, the unified superradiant condition is

$$0 < \omega < m\,\Omega_H \quad \Longrightarrow \quad \text{amplification}, \tag{16.511}$$

so a chrono-kymon–induced $\delta\Omega_H$ shifts the window by $m\,\delta\Omega_H$ (Proposition 16.69). Tau-Kairos tensor backgrounds adjust κ_{eff}, thereby modifying occupation factors in the Hawking channel. Greybody factors receive coherence-dependent distortions from both sectors via $\bar{\Sigma}$-controlled barrier shapes and off-diagonal conversion amplitudes $G_{h\tau}, G_{\tau h}$ in (16.506). Ringdown phasing is altered at $\mathcal{O}(\delta R_H,\, \delta\Omega_H)$ while TT polarizations remain unchanged.

Consistency Conditions and Decoupling Limits

Two limits ensure compatibility with classical Kerr physics:

(i) Decohered macroscopic limit. $\hat{\tau}_{\mu\nu} \to 0$ and $\mathcal{M}_{h\tau} \to 0$, so $\kappa_{\text{eff}} \to \kappa$, $\Omega_H \to \Omega_H^{\text{Kerr}}$, and GR thermodynamics/wave propagation are recovered.

(ii) TT isolation. $\mathcal{P}^{(2)}[\hat{\tau}] = 0$ and $\mathcal{P}^{(2)}\mathcal{M}_{\tau h} = 0$ enforce that no additional spin-2 degrees of freedom are radiated to infinity, preserving the standard gravitational-wave channel.

Phenomenological Synthesis

The unified temporal sector predicts a *paired* pattern of small, coherence-controlled signatures: (i) phase-accurate ringdown shifts tied to δR_H (scalar) and $\delta\Omega_H$ (vector), (ii) linear corrections to horizon area and entropy via δR_H, (iii) superradiance-window adjustments $m\,\delta\Omega_H$, (iv) subtle Hawking spectral distortions through T_{eff} and greybody factors. All are background-level (thermo-informational) effects; none add new TT polarizations.

Cross-References. Foundational definitions: Sections 16.7 and 16.8. Horizon/ergosphere conditions: eqs. (16.202), (16.203) and (16.499). Thermal map and shifts: Propositions 16.66 to 16.69.

16.10 The Temporal–Geometric Picture of Kerr Horizons

Black-hole physics in the IQR framework is governed by a unified background $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ and two fluctuation channels. The TT (spin-2) *graviton* $\hat{h}_{\mu\nu} = \delta\hat{g}_{\mu\nu}$ carries classical radiative gravity and ringdowns. The *temporal sector* $\hat{\tau}_{\mu\nu} = \delta\hat{t}_{\mu\nu}$ (Tau-Kairos tensor) has scalar $\oplus$ vector content but no TT component, and sets thermo-informational structure at horizons. Its vector constituent is the *chrono-kymon* $\hat{\mathcal{C}}_\mu$, which governs rotational coherence and vorticity through $\bar{\Sigma}_{t\phi}$ and Ω_H. Tau-Kairos tensor backgrounds dominantly shift the effective surface gravity κ_{eff} and horizon radius R_H; chrono-kymon backgrounds dominantly shift Ω_H. Hawking emission and scattering respond via the KMS map $(T_{\text{eff}}, \tilde{\omega})$, while TT gravitational-wave content remains purely in $\hat{h}_{\mu\nu}$.

Table 16.1: Division of labor at horizons: temporal sector vs. geometric TT channel

Aspect	**Tau-Kairos $\hat{\tau}_{\mu\nu}$ (scalar/vector)**	**Chrono-kymon $\hat{\mathcal{C}}_\mu$ (vector constituent)**
Spin content	No TT; scalar $\oplus$ vector only	Supplies the vector (spin-1) part via $t_{[\mu\nu]}$
Primary background lever	Lapse/temporal profile $\Rightarrow \delta R_H$, $\delta\kappa_{\text{eff}}$	Mixed time–azimuth sector $\Rightarrow \delta\bar{\Sigma}_{t\phi}$, $\delta\Omega_H$
Thermal/rotational map	$T_{\text{eff}} = \hbar\kappa_{\text{eff}}/(2\pi k_B)$ (dominant)	$\tilde{\omega} = \omega - m\Omega_H$ (dominant)
Observables	Linear entropy/area shifts; ringdown phasing via δR_H; subtle greybody changes	Superradiance window $0 < \omega < m\Omega_H$ shifted by $m\,\delta\Omega_H$; ringdown phasing via $\delta\Omega_H$
GW polarizations	None added; TT remains in $\hat{h}_{\mu\nu}$	None added; TT remains in $\hat{h}_{\mu\nu}$
Classical limit	Decouples as coherence fades; recovers Kerr thermodynamics	Decouples as coherence fades; recovers Kerr rotation physics

Key Equations and Labels.

(i) Unified split and fluctuations. $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$, $\delta\hat{\Sigma}_{\mu\nu} = \hat{h}_{\mu\nu} + \hat{\tau}_{\mu\nu}$ (eqs. (16.493) and (16.503)).

(ii) No TT in temporal sector. $\mathcal{P}^{(2)}[\hat{\tau}_{\mu\nu}] = 0$ (eq. (16.505) and Theorem 16.65).

(iii) Horizon and ergosurface. $\bar{\Sigma}_{ab}\chi^a\chi^b|_{r=R_H} = 0,\ \bar{\Sigma}_{tt} = 0$ (eqs. (16.202) and (16.203)).

(iv) Angular velocity from unified background. $\Omega_H = -\dfrac{\bar{\Sigma}_{t\phi}}{\bar{\Sigma}_{\phi\phi}}\Big|_{r=R_H}$ (eq. (16.499)).

(v) Thermodynamic map. $T_{\rm eff} = \dfrac{\hbar\,\kappa_{\rm eff}}{2\pi k_B},\ \tilde{\omega} = \omega - m\,\Omega_H$ (eqs. (16.500) and (16.507)).

(vi) Scalar-driven shifts. δR_H and $\delta\kappa_{\rm eff}$ (Propositions 16.66 and 16.67).

(vii) Vector-driven shifts. $\delta\Omega_H$ and superradiance window (Propositions 16.68 and 16.69 and eq. (16.511)).

Consistency Checks.

(i) TT isolation. Verify $\mathcal{P}^{(2)}\mathcal{M}_{\tau h} = 0$ in the matrix propagator (eq. (16.506)).

(ii) Classical recovery. Set $\hat{\tau}_{\mu\nu}\to 0$ and confirm $\kappa_{\rm eff}\to\kappa$, $\Omega_H\to\Omega_H^{\rm Kerr}$.

(iii) Spectral consistency. Implement $\omega\mapsto\tilde{\omega} = \omega - m\Omega_H$ in occupation factors and confirm that TT waveforms are unaffected.

16.10.1 PQS Core and Global Black–Hole Energetics

Local (PQS) Scale versus Global (ADM/Komar) Energy. In the IQR framework the black–hole interior is a *Primordial Quantum Singularity* (PQS), characterized by an ultra–sub–Planckian coherence length $\ell_{\rm PQS}$ (Section 13.5.11). The associated *characteristic* energy scale is

$$E_{\rm PQS} = \frac{\hbar c}{\ell_{\rm PQS}} \ \sim\ 6\times 10^{58}\ \text{eV} \quad \text{for} \quad \ell_{\rm PQS}\approx 3\times 10^{-66}\ \text{m}, \tag{16.512}$$

serving as a pre–geometric *regularity* (cutoff) scale of the core. By contrast, the black hole's *total* energy is the asymptotic (ADM/Komar) charge

$$E_{\rm BH} = Mc^2, \tag{16.513}$$

linked to horizon charges by the Smarr relation/first law (area A, surface gravity κ, angular momentum J, charge Q) [86, 87, 245]. For a non–spinning, neutral hole,

$$A = \frac{16\pi G^2M^2}{c^4}, \qquad S_{\rm BH} = \frac{k_Bc^3}{4G\hbar}\,A \ \propto\ M^2. \tag{16.514}$$

Equations (16.512)–(16.514) refer to distinct notions: $E_{\rm PQS}$ is a *micro* scale of interior coherence; $E_{\rm BH}$ is the *macro* energy seen at infinity. No numerical equality is expected (nor required).

Proposition 16.70 (Compatibility of a PQS Core with Black–Hole Energetics)**.** *Let the interior be a PQS with characteristic scale $E_{\rm PQS} = \hbar c/\ell_{\rm PQS}$ and let the exterior satisfy the usual horizon thermodynamics with surface gravity $\kappa_{\rm eff}$ determined on the unified background*

(eq. (16.2)). Then the global mass–energy $E_{\rm BH} = Mc^2$ and the first law/Smarr relation hold unchanged. The PQS modifies neither the tensor charges nor the TT radiative sector; it supplies interior regularity and induces only state–dependent, near–horizon *corrections to $\kappa_{\rm eff}$ (hence to $T_{\rm H} = \hbar\kappa_{\rm eff}/(2\pi k_B)$).*

Proof sketch. The projector constraints and matrix propagator preserve the TT channel as purely geometric ($\mathcal{P}^{(2)}\mathcal{M}_{\tau h} = 0$; Section 13.3), so the asymptotic charges (ADM/Komar) and the geometric first law remain intact. Chrono–sector (temporal) corrections enter through background coefficients that determine $\kappa_{\rm eff}$ (Proposition 16.67), affecting Hawking temperature and greybody factors without altering the tensor charges. Interior regularization by the PQS removes curvature singularities but does not inject additional asymptotic energy–momentum. □

Numerical Sanity Check (Orders of Magnitude). For a solar–mass black hole,

$$E_{\rm BH}(M_\odot) \approx 1.8 \times 10^{47}\ \text{J}, \qquad E_{\rm PQS}^{(\rm fid)} \approx 1.1 \times 10^{40}\ \text{J} \quad \Rightarrow \quad \frac{E_{\rm BH}}{E_{\rm PQS}^{(\rm fid)}} \sim 10^7. \tag{16.515}$$

The large ratio is expected: $E_{\rm PQS}$ is a *scale* of stiffness/regularity, not a total energy. The macroscopic energy Mc^2 is encoded holographically by the horizon ($A/4\ell_P^2$), whereas the PQS sets interior boundary data compatible with unitary evaporation.

Dynamical Linkage. Energy conservation takes the standard form

$$\frac{d}{dt}\big(Mc^2\big) = -\,\Phi_{\rm Hawking}, \tag{16.516}$$

with $\Phi_{\rm Hawking}$ computed from the exterior KMS/modular state (Sections 16.7 and 16.8; cf. [87]). The PQS influences $\Phi_{\rm Hawking}$ only through small shifts of $\kappa_{\rm eff}$ and greybody factors (temporal channel), while the TT graviton sector—and thus the radiative content of gravity—remains unaltered.

Heuristic: Horizon Microcells versus Core Scale. Let

$$N \;\sim\; \frac{A}{4\ell_P^2} \;\propto\; M^2 \tag{16.517}$$

be the number of horizon microcells. As M grows, N (hence the macroscopic energy Mc^2) increases, while the interior $\ell_{\rm PQS}$ (hence $E_{\rm PQS}$) is fixed by pre–geometric coherence. This explains why identifying the core with a PQS is consistent with observed masses and with standard black–hole mechanics: the horizon encodes the global energy; the PQS furnishes a nonsingular, unitary–compatible interior.

Interpretation and Significance. The PQS picture resolves a longstanding conceptual tension by separating two *distinct* energetic notions in a black hole: a *microphysical* interior scale and a *macroscopic* asymptotic charge. The PQS is characterized by an ultra–sub–Planckian

coherence length $\ell_{\rm PQS}$, which sets a *characteristic* energy scale $E_{\rm PQS} = \hbar c/\ell_{\rm PQS}$ (a stiffness/regularity measure of the pre–geometric core; cf. Section 13.5.11). By contrast, the total black–hole energy is the global, quasi–local charge $E_{\rm BH} = Mc^2$ defined at infinity and holographically encoded by the horizon state through the area/entropy relations (Smarr/first law), which remain those of classical GR (cf. eqs. (16.513) and (16.514)). These two quantities answer different questions: $E_{\rm PQS}$ quantifies *how the interior is regularized*, whereas $E_{\rm BH}$ quantifies *what the exterior measures*. No numerical equality is expected, and no double counting arises—indeed, the large hierarchy $E_{\rm BH} \gg E_{\rm PQS}$ (for astrophysical masses) is a *feature* of decoupling, not a bug.

Preservation of Exterior GR and Localization of New Physics. Within the chronogeometric/IQR dynamics, the matrix–propagator and projector constraints keep the TT (spin–2) sector purely geometric ($\mathcal{P}^{(2)}\mathcal{M}_{\tau h} = 0$), so the Komar/ADM charges, the Smarr relation, and the geometric first law hold *unchanged* (Proposition 16.70). The PQS contributes neither new asymptotic charges nor extra GW polarizations. Where it *does* enter is precisely where quantum thermodynamics lives: through the temporal (chrono–graviton/kymon) channel it induces tiny, state–dependent corrections to the near–horizon background that shift the effective surface gravity $\kappa_{\rm eff}$ and thereby the Hawking temperature $T_{\rm H} = \hbar\kappa_{\rm eff}/(2\pi k_B)$ and greybody factors (cf. eq. (16.2), Proposition 16.67, and Sections 16.7 and 16.8). In short, the PQS regularizes the interior and *subtly* tunes the thermodynamic channel, while leaving exterior tensor kinematics and charges intact.

Scaling Logic and Observational Safety. The number of horizon microcells scales as $N \sim A/(4\ell_P^2) \propto M^2$ (eq. (16.517)), so the macroscopic energy Mc^2 and classical mechanics are controlled by horizon area, not by a central lump of energy. Meanwhile, the interior coherence length $\ell_{\rm PQS}$—hence $E_{\rm PQS}$—is fixed by pre–geometric physics and does not grow with M. Thus $E_{\rm BH}/E_{\rm PQS}$ *increases* with mass, sharpening the decoupling of core microphysics from exterior observables for larger holes. The only leading deviations predicted by the framework are small shifts in ringdown phasing/greybody factors via $\kappa_{\rm eff}$, with the TT radiative content preserved—precisely the regime compatible with current GW and EM tests, and sharply falsifiable as data improve.

Unification with the Cosmological Temporal Sector. This black–hole narrative is the local, strong–field counterpart of the dark–sector story developed cosmologically: the same temporal sector (chrono–gravitons and Chrono-Kymons) that governs coherence, vacuum residue, and halo phenomenology in *Sections* 13.3*and* 13.4.7 also governs near–horizon thermodynamics here. The upshot is a single mechanism spanning regimes: temporal coherence controls *thermo–informational* features (temperature, entropy flow, interface dynamics) while the TT channel carries *geometric* radiation—both for horizons and for the Hubble flow.

Conclusion. Equating the interior to a PQS aligns with black–hole energetics. The PQS provides the ultra–short–distance, pre–geometric scale needed to regularize the core and support unitary thermodynamics, while the black hole's total energy remains the usual global mass Mc^2 encoded holographically by the horizon (Section 16.11).

16.11 The Lifecycle of a Black Hole

16.11.1 Overview: a Unitary, Relational Cycle without Paradox

The relational framework developed in this chapter yields a complete, self–consistent lifecycle for black holes governed by the unified chronogeometric tensor $\bar{\Sigma}_{\mu\nu} = \bar{g}_{\mu\nu} + \bar{t}_{\mu\nu}$ and by the universal, non–factorizable *Quantum Holo–State* $|\mathring{\boldsymbol{\Psi}}\rangle$ (Section 13.6). The classical singularity of general relativity is replaced by a stable, information–rich *Primordial Quantum Singularity* (PQS) that forms at the endpoint of gravitational collapse (Section 16.1). All dynamical phases are encoded in a single master evolution—the Universal Quantum Constraint—whose black–hole solution appears as a localized, coherent phase of the Holo–State; the emergent geometry is the local expectation value of the operator–valued chronogeometric tensor (eq. (16.219)).

Information loss does not occur. Local matter information is unitarily recoded into nonlocal, relational structure of the vacuum and is later released in the entanglement structure of Hawking radiation [87, 138, 139, 337]. The release process is governed by the *temporal sector* of the unified tensor—precisely the scalar/vector chrono–graviton background and its vector constituent, the Chrono-Kymon—through their control of effective surface gravity, horizon location, rotational chemical potential, and near–horizon mode conversion (Sections 16.7 to 16.9).

Phase I: Gravitational Collapse and PQS Formation

Macroscopic matter configurations obeying appropriate energy and compactness conditions undergo gravitational collapse. In the relational description this process is the dynamical formation of a localized, high–coherence attractor of the Holo–State: the *PQS*. Operator–geometric fields $(\hat{\boldsymbol{\Phi}}, \hat{\boldsymbol{\Theta}})$ reorganize into a stationary coherence phase compatible with a horizon–generating Killing flow χ^a (Definition 16.21). The emergent classical singularity is avoided: instead, the PQS supplies the short–scale completion of geometry and matter, rendering curvature invariants finite in expectation. The exterior settles to a Kerr–like stationary state in the unified background $\bar{\Sigma}_{\mu\nu}$, with horizon and ergosurface determined by the unified conditions (eqs. (16.202) and (16.203)).

Phase II: Stationary Coherence—Laws, Temperature, and Rotation

The exterior algebraic state restricted to a neighborhood of the horizon is KMS with respect to α_t generated by χ^a, with inverse temperature set by the *effective* surface gravity κ_{eff} of the unified background,

$$T_{\text{eff}} = \frac{\hbar\,\kappa_{\text{eff}}}{2\pi k_B}, \qquad \tilde{\omega} = \omega - m\,\Omega_H, \qquad \Omega_H = -\frac{\bar{\Sigma}_{t\phi}}{\bar{\Sigma}_{\phi\phi}}\bigg|_{r=R_H}, \tag{16.518}$$

(eqs. (16.499) and (16.507)).

The *chrono–graviton* (temporal/coherence fluctuation $\hat{\tau}_{\mu\nu} = \delta\hat{t}_{\mu\nu}$) carries only scalar $\oplus$ vector polarizations and has no TT content ($\mathcal{P}^{(2)}[\hat{\tau}] = 0$, Theorem 16.65). Its scalar

background governs the lapse–like profile and yields the leading corrections to the horizon radius and surface gravity:

$$\delta R_H = f_1(R_S) - R_S\, a_1(R_S), \qquad \kappa_{\rm eff} = \kappa_S + \frac{c^2}{2}\,\varepsilon\left[\frac{f_1'(R_S)}{R_S} - \frac{f_1(R_S)}{R_S^2} - a_1'(R_S)\right] + \mathcal{O}(\varepsilon^2), \tag{16.519}$$

(Propositions 16.66 and 16.67).

The *Chrono-Kymon* (massive vector excitation $\hat{\mathcal{C}}_\mu$ of the Tantrakyma sector) furnishes the vector constituent of $\hat{\tau}_{\mu\nu}$, controls the mixed time–azimuth sector through the coefficient D in the near–horizon ansatz, and dominantly shifts Ω_H via

$$\delta\Omega_H = -\frac{\delta\bar{\Sigma}_{t\phi}}{\bar{\Sigma}_{\phi\phi}}\Big|_{R_H} + \frac{\bar{\Sigma}_{t\phi}\,\delta\bar{\Sigma}_{\phi\phi}}{\bar{\Sigma}_{\phi\phi}^2}\Big|_{R_H} + \partial_r\left(\frac{\bar{\Sigma}_{t\phi}}{\bar{\Sigma}_{\phi\phi}}\right)\Big|_{R_H} \delta R_H + \mathcal{O}(\varepsilon^2), \tag{16.520}$$

(Proposition 16.68). The generalized horizon mechanics and entropy balance laws derived earlier are recovered with $\kappa \to \kappa_{\rm eff}$, Ω_H given by (16.518), and area computed on the unified horizon (Sections 16.6 to 16.8).

Phase III: Accretion, Transport, and Superradiant Exchange

During the active phase, matter infall, magnetic fields, and radiation interact with the ergoregion. Chrono-Kymon backgrounds set the rotational chemical potential $\mu_{\rm rot} = m\,\Omega_H$ entering scattering and emission. For bosonic modes of frequency ω and azimuthal number m, the unified superradiant condition,

$$0 < \omega < m\,\Omega_H \quad \Longrightarrow \quad \text{amplification}, \tag{16.521}$$

is shifted by $m\,\delta\Omega_H$ (Proposition 16.69). The scalar chrono–graviton background subtly reshapes greybody factors via $\kappa_{\rm eff}$ and the barrier geometry, while the vector background adjusts frame dragging and transport within the ergoregion through $\bar{\Sigma}_{t\phi}$. Throughout, the TT radiative channel remains purely geometric (ordinary gravitons), ensuring that gravitational–wave polarizations are unaltered.

Phase IV: Chronodynamics of Evaporation and the Page Curve

Evaporation is the slow, unitary leakage of entanglement mediated by the temporal sector and encoded in the KMS/modular structure. The instantaneous Hawking spectrum is controlled by $(T_{\rm eff}, \Omega_H)$ from (16.518). As mass M and angular momentum J decrease,

$$\frac{dM}{dt} = -\,\mathcal{P}\Big[T_{\rm eff}(M,J), \Omega_H(M,J), \dots\Big], \qquad \frac{dJ}{dt} = -\,\mathcal{Q}\Big[T_{\rm eff}(M,J), \Omega_H(M,J), \dots\Big], \tag{16.522}$$

with power and torque functionals $\mathcal{P}, \mathcal{Q}$ determined by greybody factors on the unified background. The entanglement entropy of radiation $S_{\rm rad}(t)$ follows a *Page curve* that rises with early–time pair production and turns over near the Page time $t_{\rm Page}$, as the entanglement wedge transitions to include interior modes; the turnover is enforced by the unitary recoding of degrees of freedom in the temporal sector and the decreasing $(T_{\rm eff}, \Omega_H)$–weighted entropy production rate [138, 139, 337].

Role of Temporal Agents. The scalar chrono–graviton determines the leading drift of $T_{\rm eff}(t)$ through $\kappa_{\rm eff}$ and the shrinking horizon $R_H(t)$ (Propositions 16.66 and 16.67); the Chrono-Kymon governs the leading drift of $\Omega_H(t)$ via (16.520). Their mixing contributes only subleading, conversion–suppressed corrections (Section 16.9).

Phase V: Late–time Dispersal and Return to the Vacuum Sector

At late times the mass approaches zero (or a model–dependent quantum–scale cutoff), the coherence sustaining the PQS dissolves, and the system relaxes into the background vacuum sector. All initially localized information has been re–expressed in relational correlations among the outgoing fields and the ambient vacuum; no paradox remains. The classical limit is recovered whenever coherence is lost: $\hat{\tau}_{\mu\nu} \to 0$, $\kappa_{\rm eff} \to \kappa$, and $\Omega_H \to \Omega_H^{\rm Kerr}$, with standard Kerr thermodynamics and wave propagation holding at macroscales.

The Black Hole as a Solution of the Universal Quantum Constraint

The black hole is a particular, localized solution of the *Universal Quantum Constraint of the Emergent Cosmos* (Section 13.6). The situation is analogous to solving the Schrödinger equation for a specific bound state within a universal Hilbert space: the black–hole Holo–State is a topologically stable, stationary coherence phase inside $|\mathring{\mathbf{\Psi}}\rangle$, and its emergent geometry is obtained as a local expectation of the operator–valued metric–temporal tensor (eq. (16.219)); cf. [406, 83].

The Black Hole as a Cosmological Nexus

Dark Energy and Dark Matter emerge here as macroscopic manifestations of the chrono–geometric vacuum rather than as independent substances [313, 211]. The black hole is the most intense local expression of this vacuum structure and thereby a nexus where principles governing the Standard Model, Dark Matter, and Dark Energy are revealed in their intimate union.

Standard Model Transfigured

Low–energy excitations identified with Standard Model particles arise as effective modes of the fundamental operator fields composing the PQS (Section 13.6). Gravitational collapse transfigures Standard–Model matter into relational information encoded by the temporal sector (Section 16.1); unitary evaporation radiates this information back into asymptotic quanta, preserving purity [330, 660].

Dark Energy Concentrated

The macroscopic cosmological constant is interpreted as the ground–state energy of the chrono–geometric vacuum. The *dressed vacuum energy density* $\rho_{\rm vac}(r)$ sourcing quantum corrections to the unified background near a rotating horizon (Section 16.5) is a localized,

intense manifestation of the same field. The small, repulsive quantum effect that shrinks the event horizon ($-\epsilon_f$) is thus a local avatar of the energy driving cosmic acceleration.

Dark Matter Made Manifest

Coherent structures of the temporal sector $\bar{t}_{\mu\nu}$ constitute gravitationally active, electromagnetically inert configurations on large scales. The long–range chrono–geometric *halo* sourced by $\bar{t}_{\mu\nu}$ in the rotating solution (Section 16.5) is precisely such a structure, matching the operational definition of a dark–matter halo [311, 588]. A black hole, in this sense, does not merely inhabit a halo; it *generates* one from the fabric of the chrono–geometric vacuum.

Unified Theorem and Observational Program

Theorem 16.71 (Unitary lifecycle of a black hole)**.** *Given (i) a stationary coherence phase on the horizon, (ii) sector preservation on $\mathcal{H}_+$, and (iii) a near–horizon KMS state with respect to χ^a, the black–hole lifecycle—collapse → stationary coherence → evaporation → vacuum return—is unitary. The temporal sector provides the thermodynamic and rotational parameters via $(\kappa_{\text{eff}}, \Omega_H)$, while the TT gravitational channel remains purely geometric; consequently, no information paradox arises.*

Observational Synthesis. The temporal sector predicts a paired pattern of small, coherence–controlled signatures: (a) phase–accurate ringdown shifts tied to δR_H (scalar) and $\delta\Omega_H$ (vector), (b) linear corrections to area/entropy via δR_H, (c) superradiance–window adjustments $m\,\delta\Omega_H$, and (d) subtle Hawking spectral distortions through T_{eff} and greybody factors. All are background–level effects that leave TT polarizations unaltered; null results at current precision are consistent with the expected decoupling in the classical limit.

Conclusion. The black hole is not a tomb for information but a *cosmic crucible*: a dynamic, unitary process that enacts the universe's most fundamental cycle—the transformation of matter into relational information, and information back into existence. In observing a black hole, one witnesses the ontology of the treatise—quantum matter, unified geometry, and the relational vacuum—made manifest in a single, magnificent object.

Chapter 17

Cosmic Evolution

17.1 The Law of Dependent Origination and Cosmic Evolution

17.1.1 The Origin of Dynamics: the Relational Einstein–Hilbert Action

Role and construction

In the chrono–geometric/IQR framework, dynamics are derived from a single, generally covariant functional built from the unified chronogeometric tensor $\bar{\Sigma}_{\mu\nu}$ and the matter operators (Gaiamiti/Tantrakyma), with the Relational Levi–Civita connection $\bar{\nabla}$ defined in Definition 6.7. The guiding principles are:

(i) State-first, representation-independent formulation. The primal object is the universal Holo–State $\left|\mathring{\mathbf{\Psi}}\right\rangle$; the action and field equations are defined on the BRST-physical algebra and then represented on $\mathcal{H}_{\text{phys}}$ via GNS (cf. Sections 2.4 and 13.6). See Definition 11.106 for the Chronogeometric Einstein Equations and eq. (11.269) for their expectation-level placement with Λ_{eff}.

(ii) General covariance and minimality. In four dimensions, the minimal generally covariant two-derivative choice yielding second-order field equations is the Einstein–Hilbert density plus a constant four-volume term (the cosmological constant) [661]. Higher-curvature additions (e.g. R^2, $R_{\mu\nu}R^{\mu\nu}$) either introduce higher derivatives in the equations of motion or contribute as topological/boundary pieces in $D = 4$; in a modern view they are treated as effective field-theory corrections suppressed by the UV scale [415, 403, 12].

Accordingly, the *Total Relational Action* is

$$S_{\text{rel}} \;=\; \frac{1}{2\kappa'}\int\Big(\,\bar{R}[\bar{\Sigma}] - 2\Lambda\,\Big)\sqrt{-\det\bar{\Sigma}}\,d^4x \;+\; S_{\text{matter}}\Big[\hat{\mathbf{\Phi}},\hat{\mathbf{\Theta}}\Big] \;+\; S_{\text{bdy}}[\bar{\Sigma}], \tag{17.1}$$

where κ' is the (relationally normalized) coupling, Λ is the constant four-volume density coefficient (Einstein–Hilbert form [661]), S_{matter} collects Tantrakyma/Gaiamiti dynamics and their couplings (realized in the operator–valued coupling formalism Definition 11.109 and eqs. (11.213) and (11.214)), and S_{bdy} (a Gibbons–Hawking–York–type term adapted

to $\bar{\Sigma}$) ensures a well-posed Dirichlet variational problem on manifolds with boundary [532, 533, 534, 535] *(for the typed retarded/advanced Green operators and their causal support in the response formalism, see eqs.* (11.213) *and* (11.214)*).* In this state-first reading, S_{rel} is an expectation-valued functional evaluated on histories compatible with $\left|\overset{\circ}{\boldsymbol{\Psi}}\right\rangle$ and the holosymmetric constraints (see also the expectation-level field equation eq. (11.269) for how Λ_{eff} enters).

Why These Terms (and Only These) Appear

(i) Stationary action and scalar invariance. The action must be a scalar under diffeomorphisms to admit a coordinate-free stationary principle (cf. Section 11.5.3).

(ii) Geometric scalars. The two elementary scalar densities are: (i) $\bar{R}[\bar{\Sigma}]\sqrt{-\det\bar{\Sigma}}$, and (ii) $\sqrt{-\det\bar{\Sigma}}$ itself. The latter integrates the invariant four-volume and introduces Λ as its proportionality constant—an intrinsic baseline of relational "self-energy'' of spacetime.

(iii) Matter sector. S_{matter} supplies the Gaiamiti/Tantrakyma content and defines the operator stress tensor $T^{\mathrm{matter}}_{\mu\nu}$ via the standard metric variation with $\bar{\Sigma}_{\mu\nu}$ as the variational metric.

Field Equations and the Vacuum Stress

Varying S_{rel} with respect to $\bar{\Sigma}^{\mu\nu}$ yields the *Chronogeometric Einstein Equations* (cf. Definition 11.106):

$$\bar{R}_{\mu\nu} - \tfrac{1}{2}\,\bar{\Sigma}_{\mu\nu}\,\bar{R} \;+\; \Lambda\,\bar{\Sigma}_{\mu\nu} \;=\; \kappa'\,T^{\mathrm{matter}}_{\mu\nu}, \tag{17.2}$$

so the cosmological term is equivalent to an intrinsic vacuum stress

$$T^{\mathrm{vac}}_{\mu\nu} := -\,\frac{\Lambda}{\kappa'}\,\bar{\Sigma}_{\mu\nu}. \tag{17.3}$$

This makes explicit that Λ contributes as a uniform energy density and negative pressure already at the level of the geometric sector, independently of matter.

Bare Λ vs. Interferometric Vacuum Energy

Later (cf. Section 13.4.7) the *interferometric residual* of the dual-field vacuum defines a positive energy density ε_Λ via

$$\rho_{\mathrm{DE}} = \varepsilon_\Lambda = \frac{1}{\mathrm{Vol}(\mathcal{X}_q)}\left\langle \overset{\circ}{\boldsymbol{\Psi}}\right| \mathcal{P}^{\mathrm{sym}}_{\mathrm{vac}}\, H_{\mathrm{vac}}\, \mathcal{P}^{\mathrm{sym}}_{\mathrm{vac}} \left|\overset{\circ}{\boldsymbol{\Psi}}\right\rangle_{\mathrm{phys}} \quad \text{(eq. (13.145)).} \tag{17.4}$$

At the level of the field equations one may fold this into an *effective cosmological constant*

$$\Lambda_{\mathrm{eff}} \;=\; \Lambda \;+\; \kappa'\,\varepsilon_\Lambda, \tag{17.5}$$

so that the late-time acceleration is governed by Λ_{eff} while preserving the clean separation between the *bare geometric* constant Λ and the *state-dependent* vacuum residue ε_Λ selected by the Holosymmetric filter (Sections 13.2.6 and 13.2.7).

Hamiltonian Perspective and the Universal Constraint

Legendre transforming S_{rel} (with the emergent proper time of Section 5.3.8) yields the *Total Relational Hamiltonian* $\hat{H}_{\mathrm{univ}} = \hat{H}_\Phi + \hat{H}_\Theta + \hat{H}_{\mathrm{int}}$ (cf. eq. (13.235)). The Λ term appears as a uniform potential density in the geometric part and, in the state-first/GNS representation, contributes to the *Universal Quantum Constraint*

$$\hat{H}_{\mathrm{univ}} \left| \mathring{\boldsymbol{\Psi}} \right\rangle = 0 \quad (\text{eq. } (13.236)), \tag{17.6}$$

which encodes the vanishing of total relational energy and underwrites both the horizon–KMS constructions and the cosmological abundance partitions (Sections 13.3, 13.4.7 and 16.11).

The Relational Einstein–Hilbert Action is uniquely fixed (in $D = 4$) by covariance and second-order dynamics to be "Ricci scalar + cosmological volume" plus matter and boundary terms. Its variation yields the vacuum stress $T^{\mathrm{vac}}_{\mu\nu} = -(\Lambda/\kappa')\bar{\Sigma}_{\mu\nu}$; its Hamiltonian form embeds Λ into the universal constraint; and, when combined with the holosymmetric vacuum residue ε_Λ, it furnishes the effective $\Lambda_{\mathrm{eff}} = \Lambda + \kappa'\varepsilon_\Lambda$ that maps directly to the late-time acceleration in ΛCDM language while keeping the IQR state-first structure explicit.

17.1.2 The Formal Origin of the Cosmological Constant

In the IQR/chrono–geometric framework, the dynamics of the emergent universe are governed by the *Total Relational Action* (Section 17.1.1). General covariance and second–order field equations uniquely fix the geometric sector in $D = 4$ (up to a boundary term) to the Einstein–Hilbert density plus a constant four–volume density. Writing the relational volume form as $\mathrm{d}\mu_{\bar{\Sigma}} := \sqrt{-\det \bar{\Sigma}}\, \mathrm{d}^4x$, the geometric action reads

$$S_{\mathrm{geom}}[\bar{\Sigma}] = \frac{1}{2\kappa'} \int_{\mathcal{M}} \left(\bar{R}[\bar{\Sigma}] - 2\Lambda \right) \mathrm{d}\mu_{\bar{\Sigma}} \; + \; S_{\mathrm{bdy}}[\bar{\Sigma}], \tag{17.7}$$

where Λ is the constant four–volume coefficient (the *cosmological constant*; see Remark 11.144 for diagnostic realized reference configuration vs. observed and its role in Λ_{eff}).

Explicit variation. On any open region where $\bar{\Sigma}_{\mu\nu}$ is non–degenerate,

$$\delta\sqrt{-\det \bar{\Sigma}} = -\tfrac{1}{2}\sqrt{-\det \bar{\Sigma}}\, \bar{\Sigma}_{\mu\nu}\, \delta\bar{\Sigma}^{\mu\nu}, \qquad \delta(\sqrt{-\det \bar{\Sigma}}\, \bar{R}) = \sqrt{-\det \bar{\Sigma}}\, \bar{G}_{\mu\nu}\, \delta\bar{\Sigma}^{\mu\nu} + \nabla(\cdots),$$

so that

$$\delta S_{\mathrm{geom}} = \frac{1}{2\kappa'} \int_{\mathcal{M}} \sqrt{-\det \bar{\Sigma}}\, \left(\bar{G}_{\mu\nu} + \Lambda\, \bar{\Sigma}_{\mu\nu} \right) \delta\bar{\Sigma}^{\mu\nu}\, \mathrm{d}^4x \; + \; \delta S_{\mathrm{bdy}}. \tag{17.8}$$

Adding the matter/coherence action and using $\delta S_{\mathrm{matter}} = -\frac{1}{2}\int \sqrt{-\det \bar{\Sigma}}\; T^{\mathrm{matter}}_{\mu\nu}\, \delta\bar{\Sigma}^{\mu\nu}\, \mathrm{d}^4x$ (Section 9.17.9) gives the Euler–Lagrange equations

$$\bar{R}_{\mu\nu} - \tfrac{1}{2}\, \bar{\Sigma}_{\mu\nu}\, \bar{R} + \Lambda\, \bar{\Sigma}_{\mu\nu} = \kappa'\, T^{\mathrm{matter}}_{\mu\nu}, \tag{17.9}$$

i.e. the Chronogeometric Einstein Equations (Definition 11.106; see also the expectation-level placement with Λ_{eff} in eq. (11.269)).

Vacuum stress identification. Equation (17.9) shows that the cosmological term is equivalent to a homogeneous vacuum stress carried by the geometric sector:

$$T^{\mathrm{vac}}_{\mu\nu} := -\frac{\Lambda}{\kappa'}\,\bar{\Sigma}_{\mu\nu}. \tag{17.10}$$

As a consequence (cf. Lemmas 11.123 and 11.124), $k^\mu k^\nu T^{\mathrm{vac}}_{\mu\nu} = 0$ for every $\bar{g}^{\mathrm{Lor}}$-null vector k^μ; the vacuum term therefore drops out of the Null Coherence Condition and of null focusing (Section 11.6 and Theorem 11.128).

Placement relative to the response kernel. In the operator–valued coupling formalism (Definition 11.109), the cosmological term belongs to the geometric/LHS of the field equations, while the coupling kernel $\hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}$ maps *non-vacuum* sources to curvature (cf. eqs. (11.213) and (11.214)). When state-dependent vacuum residues are included, this placement leads to the effective constant $\Lambda_{\mathrm{eff}} = \Lambda + \kappa'\varepsilon_\Lambda$ derived in Section 17.1.3, consistent with the expectation-level arrangement in eq. (11.269) and without double counting on the source side.

The Quantum–Mechanical Nature of the Cosmological Constant

The variational identification above is elevated to a quantum statement in the state–first (GNS/BRST) reading (Section 13.6). The geometric sector admits a *Geometric Hamiltonian* $\hat{H}_{\mathrm{geometry}}$ affiliated with the physical von Neumann algebra, and the Holo–State $|\mathring{\mathbf{\Psi}}\rangle$ selects the physical GNS representation. The emergent inner product used in this representation is generated by the Fourfold inter-modal correlator (Definition 4.9 and eq. (4.85)), yielding the modal fibers introduced in Chapter 4 (Section 4.7). Let $|\Psi_{\mathrm{vac}}\rangle \in \mathcal{M}_{\mathrm{vac}}$ denote a normalized vacuum in a homogeneous sector (Section 6.1.8). Defining volume with the relational measure $\mathrm{Vol}(\mathcal{X}_q) := \int_{\mathcal{X}_q}\sqrt{-\det\bar{\Sigma}}\,\mathrm{d}^4x$ (understood as a limit along compact exhaustions), one may consistently set

$$\hat{H}_{\mathrm{geometry}}\,|\Psi_{\mathrm{vac}}\rangle = \Lambda_{\mathrm{geo}}\,|\Psi_{\mathrm{vac}}\rangle\,, \qquad \Lambda_{\mathrm{geo}} = \frac{\langle\Psi_{\mathrm{vac}}|\hat{H}_{\mathrm{geometry}}|\Psi_{\mathrm{vac}}\rangle_{\mathrm{phys}}}{\mathrm{Vol}(\mathcal{X}_q)}. \tag{17.11}$$

Thus, Λ_{geo} is the density–normalized ground–state energy of the emergent chronogeometric field.

Endogenous (state–conditioned) regularization. In the spectral realization of Λ_{geo}, the mode sum is finite *without* ad-hoc cutoffs. Concretely: *(i)* the **state-conditioned modal weights** $w_n(x,x')$ (Definition 4.392) suppress unsupported channels; *(ii)* the **Principle of Finite Entanglement Capacity** imposes UV decay on w_n (Principle 4.1); *(iii)* the effective UV band edge is encoded by the **effective spectral cutoff** $\Lambda^{[\mathrm{obs}]}_{\mathrm{eff}}(x,x';\varepsilon)$ (Definition 4.576); *(iv)* **intrinsic entanglement dressing** and relational windowing provide the IR floor (Section 4.9.15, Definition 4.435, and Proposition 4.436); and *(v)* the whole construction is captured by the **Principle of Coherent Regularization** (Proposition 4.424). These ingredients are formulated using the admissible spectral operator $\hat{D}$ (Definition 4.343) acting on the Chapter 4 modal

fibers, with dynamics probed via the spectral shear observable (Definition 4.386). UV/IR finiteness is therefore an endogenous property of the Holo–State and its operator channels, consistent with diffeomorphism covariance and the BRST/GNS construction.

At the level of field equations, Λ in (17.9) is the *bare geometric* constant that matches Λ_{geo} in the pure–geometry vacuum; departures arise once matter/coherence contributions are included and renormalized.

The Physical Manifestation as Dark Energy

The macroscopic phenomenon of *Dark Energy* is the observable imprint of a small, positive vacuum energy density in the emergent chronogeometric background (see, e.g., the discovery papers [134, 135] and global constraints [136]; for reviews see [211, 212, 414]). In the present framework this arises from a *dynamical self–regulation* between the geometric (Gaiamiti) and material/coherence (Tantrakyma) vacua: large, opposite–sign contributions nearly cancel, leaving a small, positive *interferometric residue* ε_Λ selected by the holosymmetric filter (Sections 13.2.7 and 13.4.7). Coherent (state–conditioned) regularization (Proposition 4.424; Ch. 4, via state-conditioned modal weights Definition 4.392, finite entanglement capacity Principle 4.1, effective spectral cutoff Definition 4.576, and IR windowing Section 4.9.15 and Definition 4.435) renders the vacuum density finite and covariant, consistent with renormalized stress–energy in curved spacetime [244, 245], so that cosmological dynamics are governed by the *effective* constant

$$\Lambda_{\text{eff}} = \Lambda + \kappa' \, \varepsilon_\Lambda, \tag{17.12}$$

as derived in Section 17.1.3 and consistent with the expectation-level placement in eq. (11.269). Equivalently, the observed Dark Energy density is $\rho_{\text{DE}} = \Lambda_{\text{eff}}/\kappa'$ (eq. (17.15)), and the corresponding vacuum stress takes the geometric form $T_{\mu\nu}^{\text{vac,eff}} = -(\Lambda_{\text{eff}}/\kappa')\,\bar{\Sigma}_{\mu\nu}$ (cf. eq. (17.10); see also the standard identification of vacuum stress from a cosmological term [211, 313]).

(i) Λ: bare geometric four–volume coefficient in the action (Section 17.1.1).

(ii) ε_Λ: small, state–selected (holosymmetric) *interferometric residue* of the combined (Gaiamiti+Tantrakyma) vacuum (Section 13.2.7); its finiteness follows from coherent regularization (Proposition 4.424; Ch. 4).

(iii) Λ_{eff}: cosmologically relevant constant entering Friedmann dynamics and late–time kinematics, obtained by shifting the geometric term by $\kappa'\varepsilon_\Lambda$ (Section 17.1.3 and eq. (11.269); cf. observational inference [136]).

In the triadic decomposition (corr/vac/spec; cf. Sections 11.1.4 and 11.2.4), Λ_{eff} belongs to the *vacuum* channel and coexists with the spectral (GR) and correlational (entanglement–induced) channels used elsewhere in the response formalism. Consistently with Lemmas 11.123 and 11.124, the vacuum term $T_{\mu\nu}^{\text{vac,eff}} \propto \bar{\Sigma}_{\mu\nu}$ drops out on $\bar{g}^{\text{Lor}}$-null directions and thus does not affect the NCC or null focusing (Section 11.6 and Theorem 11.128; for classic treatments of null focusing and energy conditions see [240, 7]). In the operator–valued coupling description (Definition 11.109), this placement keeps the vacuum on the geometric side while the kernel $\hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}$ maps *non–vacuum* sources to curvature, avoiding double counting (cf. eqs. (11.213) and (11.214)).

Remark 17.1 (On ε_Λ, Λ, and the cosmological vs. spectral Λ_{eff})**.**

(i) Cosmology-sector meaning in this chapter.
Here and in Chapter 11, Λ_{eff} denotes the *effective cosmological constant* defined by

$$\Lambda_{\text{eff}} = \Lambda + \kappa' \varepsilon_\Lambda \quad \text{(cf. eqs. (11.269) and (17.12))},$$

the quantity that enters the expectation-level Einstein equations and is inferred observationally as Dark Energy.

(ii) Meaning of Λ (diagnostic realized reference configuration/bare).
Λ is the diagnostic realized reference configuration (bare) cosmological term fixed by the diagnostic realized reference configuration/plateau and renormalization scheme; it has units L^{-2}. Geometric–source placement is equivalent at the expectation level:

$$\Lambda\, \bar{\Sigma}_{\mu\nu} \iff T^{\text{vac}}_{\mu\nu} = -(\Lambda/\kappa')\, \bar{\Sigma}_{\mu\nu}.$$

(iii) Observed vs. diagnostic realized reference configuration.
The observed constant is Λ_{eff}. Only if $\varepsilon_\Lambda = 0$ would $\Lambda_{\text{eff}} = \Lambda$.

(iv) Not the spectral cutoff (Chapter 4).
The symbol Λ_{eff} in this chapter is *not* the Chapter 4 *effective spectral cutoff* $\Lambda^{[\text{obs}]}_{\text{eff}}(x, x'; \varepsilon)$. See the cross-chapter dictionary in Remark 4.575. A built-in disambiguator is the presence of arguments $(x, x'; \varepsilon)$ for the spectral notion.

(v) Units gate.
The relevant dimensional assignments are

$$[\Lambda_{\text{eff}}] = [\Lambda] = L^{-2}, \qquad [\varepsilon_\Lambda] = L^{-4}, \qquad [\Lambda^{[\text{obs}]}_{\text{eff}}] = L^{-1}.$$

Cosmology/chronogeometry carries L^{-2}; spectral cutoff carries L^{-1}.

(vi) Triadic bookkeeping and no double counting.
Λ_{eff} sits in the *vacuum* channel of the triadic decomposition (corr/vac/spec; cf. Section 11.1.4). In the operator-valued coupling formalism (Definition 11.109), keep the vacuum on the geometric side while $\hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}$ maps *non-vacuum* sources to curvature (cf. eqs. (11.213) and (11.214)).

(vii) Null-silent vacuum and NEC.
Consistently with Lemmas 11.123 and 11.124, the vacuum term $T^{\text{vac,eff}}_{\mu\nu} \propto \bar{\Sigma}_{\mu\nu}$ drops out on $\bar{g}^{\text{Lor}}$-null directions and does not affect the null convergence condition or null focusing (Section 11.6 and Theorem 11.128).

Foundational Definition of the Cosmological Constant

Definition 17.2 (Cosmological Constant as Emergent Ground–State Geometry)**.** The **cosmological constant** Λ quantifies the intrinsic, ground–state energy of the emergent chronogeometric spacetime. Its origin and meaning follow a three–level hierarchy:

Formal (variational) origin
In the *Total Relational Action*, Λ is the constant of the invariant four–volume term

(Section 17.1.1 and eqs. (17.7) and (17.8)). Its variation yields the vacuum stress in (17.10) and enters the Chronogeometric Einstein Equations (Definition 11.106 and eq. (17.9)); see also the expectation–level placement with Λ_{eff} in eq. (11.269).

Quantum origin

In the BRST/GNS representation selected by the Holo–State $|\mathring{\boldsymbol{\Psi}}\rangle$ (Section 13.6), the quantity Λ_{geo} in (17.11) is the density–normalized ground–state energy of $\hat{H}_{\text{geometry}}$ in a homogeneous vacuum sector $|\Psi_{\text{vac}}\rangle \in \mathcal{M}_{\text{vac}}$ (Section 6.1.8). The emergent inner product and modal fibres used here arise from the Fourfold inter-modal correlator (Definition 4.9 and eq. (4.85)). Its spectral definition is *endogenously finite* by coherent (state–conditioned) regularization (Proposition 4.424), implemented via state-conditioned modal weights, capacity, and IR windowing (Definitions 4.392, 4.435 and 4.576, Principle 4.1, and Section 4.9.15); for the trace/mode bridge see eqs. (17.13) and (17.14).

Physical manifestation.

The observed Dark Energy is the interferometric residue ε_Λ of the combined (Gaiamiti+Tantrakyma) vacuum, producing the effective constant $\Lambda_{\text{eff}} = \Lambda + \kappa' \varepsilon_\Lambda$ that drives acceleration (Sections 13.2.6, 13.4.7 and 17.1.3 and eqs. (17.12) and (17.15)). This Λ_{eff} is placed on the geometric side at expectation level (eq. (11.269)); in the operator–valued coupling formalism the non-vacuum sources are mapped by $\hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}$ (Definition 11.109 and eqs. (11.213) and (11.214)), with notation distinguished from the Chapter 4 spectral cutoff by Remark 4.575.

Remark 17.3 (Mapping to ΛCDM, null cones, and de Sitter thermodynamics)**.** (i) A strictly constant ε_Λ reproduces GR with Λ_{eff} and the standard ΛCDM phenomenology constrained by SNe Ia and CMB ([134, 135, 136]; see Section 17.1). (ii) Because $T^{\text{vac}}_{\mu\nu} \propto \bar{\Sigma}_{\mu\nu}$, it drops out on $\bar{g}^{\text{Lor}}$-null directions (Lemma 11.123); hence the NCC and null focusing are governed by matter and coherence stresses only, consistent with §11.6. (iii) In the asymptotic de Sitter regime, Λ_{eff} sets the Gibbons–Hawking temperature of the cosmological horizon, aligning with the modular/KMS structures used elsewhere in the treatise.

The Calculation of the Cosmological Constant via a Regulated Spectral Integral

In the IQR/chrono–geometric framework, the cosmological constant Λ is tied to the ground–state energy of the geometric (Gaiamiti) sector and can be computed from the *Geometric Hamiltonian* as an expectation-valued, regulator–finite quantity (Sections 13.6 and 17.1.1). At the operator level we write (state–first/GNS reading; cf. Section 6.1.13)

$$\hat{H}_{\text{geometry}} = \int_{\mathcal{X}_q} d^d x \, \sqrt{-\det \bar{\Sigma}(x)} \, \operatorname{Tr}\big[\hat{G}_{\boldsymbol{\Phi}\boldsymbol{\Phi}}(x,x)\big] \,, \tag{17.13}$$

where $\hat{G}_{\boldsymbol{\Phi}\boldsymbol{\Phi}}(x,x)$ is the local geometric self–correlator (Hadamard two–point function evaluated at coincidence; cf. Section 6.1.10), generated from the Fourfold inter–modal correlator (Definition 4.9) within the emergent inner–product framework (eq. (4.85)). Passing to a mode (momentum) representation on a quasi–homogeneous patch ($\bar{\Sigma}_{\mu\nu}$ slowly varying) turns the local trace into a spectral integral [10, 13, 211, 313], using the Chapter 4 modal fibres and

admissible spectral operator (Section 4.7 and Definition 4.343):

$$\Lambda_{\text{geo}} \;\propto\; \frac{1}{\text{Vol}(\mathcal{X}_q)} \left\langle \overset{\circ}{\boldsymbol{\Psi}} \middle| \hat{H}_{\text{geometry}} \middle| \overset{\circ}{\boldsymbol{\Psi}} \right\rangle_{\text{phys}} \;\simeq\; \int_{k_{\text{IR}}}^{k_{\text{UV}}} \frac{d^3k}{(2\pi)^3}\, E(k), \qquad E(k) \equiv \tfrac{1}{2}\hbar\,\omega_k + \cdots, \tag{17.14}$$

with ω_k the (background–dressed) Gaiamiti dispersion and "$+\cdots$" denoting interaction/curvature corrections controlled by $\bar{\Sigma}$ (Section 11.9). The continuum measure $\int d^3k$ is the momentum–space avatar of the local trace in (17.13) (trace $\leftrightarrow$ mode sum), while the on–shell value of the correlator at coincidence fixes $E(k)$ (correlator $\leftrightarrow$ mode energy; cf. Section 6.1.10). Equation (17.14) is the regulated, physical version of the vacuum–energy integral familiar from QFT in curved spacetime [10, 13], now rendered finite by the intrinsic structure of the chrono–geometric regimes together with the Chapter 4 coherent–regularization machinery (Proposition 4.424).

The Ultraviolet Cutoff k_{UV}. Unlike the formal quartic divergence of flat–space QFT, the present framework enforces an *intrinsic* UV bound by the **Principle of Finite Entanglement Capacity** (Principle 4.1). It sets a maximal resolvable spectral scale λ_{cutoff} ("hardest" fluctuation), hence $k_{\text{UV}} \sim \lambda_{\text{cutoff}}^{-1}$, and eliminates the UV pathology at its origin (see also the effective spectral cutoff $\Lambda_{\text{eff}}^{[\text{obs}]}$ in Definition 4.576).

The Infrared Cutoff k_{IR}. Two complementary mechanisms regulate the deep IR:

(i) Global (cosmological) IR. The finite cosmological domain acts as an IR box: modes with λ exceeding the cosmological radius are unphysical for local energetics (Section 11.9; cf. horizon thermodynamics in de Sitter [314]).

(ii) Local/dynamical IR. Near dense sources (e.g. PQS cores), the **Principle of Intrinsic Entanglement Dressing** induces a small, state–dependent mass gap $m_{\text{eff}}(r)$ for geometric mediators, providing a local IR regulator (Section 4.9.15 and Proposition 4.439); this aligns with the Chapter 4 IR coherence scale and windowing (Definition 4.435 and Section 4.9.15). This gap sets the softest resolvable mode at radius r, $k_{\text{IR}} \sim m_{\text{eff}}(r)c/\hbar$.

The Bridge Between the Formulations

The correspondence between (17.13) and (17.14) is standard:

(i) Trace $\leftrightarrow \int d^3k$. The operator trace equals the sum over a complete mode basis; in a quasi–homogeneous patch the basis is the momentum basis, hence $\text{Tr}(\cdots) \to \int d^3k/(2\pi)^3\,(\cdots)$ (cf. Chapter 4 modal fibres and spectral basis Section 4.7 and Definition 4.343).

(ii) Correlator $\leftrightarrow E(k)$. The local self–correlator evaluated on mode k yields the zero–point (plus curvature–dressed) energy $E(k)$ of that mode (cf. Section 6.1.10 and the Fourfold generation Definition 4.9).

Thus the momentum–space integral is simply the spectral representation of the fundamental Hamiltonian expectation (Sections 6.1.10 and 6.1.13), with the expectation–level field placement in the gravitational response given in eq. (11.269).

Self–Regulation and the Final Value

Even with physical cutoffs, Λ_{geo} from (17.14) is generically large. The *final* step is the **dynamical self–regulation** between the geometric (Gaiamiti) and material (Tantrakyma) vacua, encoded by the Total Relational Action and the holosymmetric filter (Sections 13.2.7 and 17.1.1). The net dark–energy density is the small, positive *residual*

$$\rho_{\text{DE}} \;=\; \varepsilon_\Lambda \;=\; \frac{1}{\text{Vol}(\mathcal{X}_q)}\left|\left\langle \mathring{\boldsymbol{\Psi}}\middle|\hat{H}_{\text{geometry}}\middle|\mathring{\boldsymbol{\Psi}}\right\rangle_{\text{phys}} + \left\langle \mathring{\boldsymbol{\Psi}}\middle|\hat{H}_{\text{matter}}\middle|\mathring{\boldsymbol{\Psi}}\right\rangle_{\text{phys}}\right| \;\ll\; \frac{k_{\text{UV}}^4}{16\pi^2}\hbar c, \tag{17.15}$$

realizing a controlled cancellation reminiscent of "vacuum energy problem'' discussions [211, 313], but now grounded in the Holosymmetric Selection Principle and chrono–geometric regimes (Sections 13.2.6 and 13.4.7). Observationally, $\Lambda_{\text{eff}} = \Lambda + \kappa'\varepsilon_\Lambda$ inferred from SNe Ia and CMB is small and positive [134, 135, 136], in agreement with the regulated, self–consistent construction above and its expectation–level placement (eq. (11.269)); note the cosmology–sector Λ_{eff} here is distinct from the Chapter 4 spectral cutoff $\Lambda_{\text{eff}}^{[\text{obs}]}$ (Remark 4.575).

Remark 17.4 (On Regulator Independence of Ω-Fractions)**.** The absolute values of the sectoral vacuum energies depend on the physical cutoffs set by the regimes, but the *abundance fractions* $\Omega_{\text{DE}}, \Omega_{\text{DM}}, \Omega_{\text{vis}}$ are regulator–independent ratios fixed by the holosymmetric projector decomposition (Section 13.4.7 and Proposition 13.97).

17.1.3 Derivation of the Effective Cosmological Constant

We now formalize the identification

$$\Lambda_{\text{eff}} = \Lambda + \kappa'\,\varepsilon_\Lambda, \tag{17.16}$$

by starting from the state–first action principle, splitting the physical source into vacuum and non-vacuum parts, and then recasting the field equations in a form aligned with the operator–valued coupling kernel of Def. 11.109.

Setup. Work in the BRST/GNS physical sector selected by the Holo–State $|\mathring{\boldsymbol{\Psi}}\rangle$ (Section 13.6). The geometric part of the Total Relational Action is

$$S_{\text{geom}}[\bar{\Sigma}] = \frac{1}{2\kappa'}\int_{\mathcal{M}}\big(\bar{R}[\bar{\Sigma}] - 2\Lambda\big)\sqrt{-\det\bar{\Sigma}}\;\mathrm{d}^4x, \tag{17.17}$$

whose variation with respect to $\bar{\Sigma}^{\mu\nu}$ yields the Chronogeometric Einstein Equations (Definition 11.106)

$$\bar{R}_{\mu\nu} - \tfrac{1}{2}\,\bar{\Sigma}_{\mu\nu}\,\bar{R} + \Lambda\,\bar{\Sigma}_{\mu\nu} = \kappa'\,\bar{T}^{\text{matter}}_{\mu\nu}. \tag{17.18}$$

Definition 17.5 (Holosymmetric vacuum residue)**.** Let $\bar{T}^{\text{matter}}_{\mu\nu}$ denote the renormalized, BRST/GNS expectation of the matter+coherence stress tensor on $|\mathring{\boldsymbol{\Psi}}\rangle$. In a homogeneous/isotropic vacuum sector (Section 6.1.8), its vacuum contribution must be proportional to $\bar{\Sigma}_{\mu\nu}$:

$$\bar{T}^{(\text{vac})}_{\mu\nu} := -\,\varepsilon_\Lambda\;\bar{\Sigma}_{\mu\nu}, \qquad \varepsilon_\Lambda \in \mathbb{R}, \tag{17.19}$$

where ε_Λ is the *holosymmetric interferometric residue* fixed by the state and by coherent (state-conditioned) renormalization (Sections 13.2.7 and 13.4.7). We then split

$$\bar{T}_{\mu\nu}^{\text{matter}} = \bar{T}_{\mu\nu}^{(\text{nonvac})} + \bar{T}_{\mu\nu}^{(\text{vac})} = \bar{T}_{\mu\nu}^{(\text{nonvac})} - \varepsilon_\Lambda\, \bar{\Sigma}_{\mu\nu}. \tag{17.20}$$

Proposition 17.6 (Effective cosmological constant in the field equations)**.** *Combining* (17.18) *with the split* (17.20) *and rearranging terms gives*

$$\bar{R}_{\mu\nu} - \tfrac{1}{2}\,\bar{\Sigma}_{\mu\nu}\,\bar{R} + \underbrace{\big(\Lambda + \kappa'\,\varepsilon_\Lambda\big)}_{:=\,\Lambda_{\text{eff}}}\,\bar{\Sigma}_{\mu\nu} = \ \kappa'\,\bar{T}_{\mu\nu}^{(\text{nonvac})}, \tag{17.21}$$

(cf. eq. (11.269)). Thus the effective *cosmological constant is* $\Lambda_{\text{eff}} = \Lambda + \kappa'\varepsilon_\Lambda$*, with the vacuum energy encoded on the geometric (left–hand) side.*

Proof idea. Insert (17.20) into (17.18):

$$\bar{R}_{\mu\nu} - \tfrac{1}{2}\,\bar{\Sigma}_{\mu\nu}\bar{R} + \Lambda\,\bar{\Sigma}_{\mu\nu} = \kappa'\,\bar{T}_{\mu\nu}^{(\text{nonvac})} - \kappa'\,\varepsilon_\Lambda\,\bar{\Sigma}_{\mu\nu}.$$

Move the vacuum term $-\kappa'\varepsilon_\Lambda\,\bar{\Sigma}_{\mu\nu}$ to the left and read off (17.21).

Full proof. Step 0 (Setting and assumptions). Work on the BRST/GNS physical sector selected by the Holo–State $|\mathring{\boldsymbol{\Psi}}\rangle$ with Hadamard regularity (Section 1.4.1). Let $\bar{\Sigma}_{\mu\nu}$ denote the unified chronogeometric tensor and $\mathrm{d}\mu_{\bar{\Sigma}} = \sqrt{-\det\bar{\Sigma}}\,\mathrm{d}^4x$ the invariant measure.

Step 1 (Field equations with bare Λ*).* Varying the geometric action (17.17) yields the Chronogeometric Einstein Equations

$$\bar{R}_{\mu\nu} - \tfrac{1}{2}\,\bar{\Sigma}_{\mu\nu}\,\bar{R} + \Lambda\,\bar{\Sigma}_{\mu\nu} = \kappa'\,\bar{T}_{\mu\nu}^{\text{matter}}. \tag{17.22}$$

This is (17.18).

Step 2 (Vacuum structure and split). By the holosymmetric vacuum definition (Definition 17.5), the homogeneous/isotropic vacuum piece must be proportional to $\bar{\Sigma}_{\mu\nu}$:

$$\bar{T}_{\mu\nu}^{(\text{vac})} = -\,\varepsilon_\Lambda\,\bar{\Sigma}_{\mu\nu}, \qquad \varepsilon_\Lambda \in \mathbb{R},$$

and we split the source as in (17.20):

$$\bar{T}_{\mu\nu}^{\text{matter}} = \bar{T}_{\mu\nu}^{(\text{nonvac})} - \varepsilon_\Lambda\,\bar{\Sigma}_{\mu\nu}.$$

Step 3 (Substitution and rearrangement). Insert the split into (17.22):

$$\bar{R}_{\mu\nu} - \tfrac{1}{2}\,\bar{\Sigma}_{\mu\nu}\bar{R} + \Lambda\,\bar{\Sigma}_{\mu\nu} = \kappa'\,\bar{T}_{\mu\nu}^{(\text{nonvac})} - \kappa'\,\varepsilon_\Lambda\,\bar{\Sigma}_{\mu\nu}.$$

Move the vacuum term to the LHS and collect the four–volume pieces to obtain (17.21) with $\Lambda_{\text{eff}} := \Lambda + \kappa'\varepsilon_\Lambda$.

Step 4 (Conservation/Bianchi compatibility). Taking $\bar{\nabla}^\mu$ of (17.21) and using $\bar{\nabla}^\mu\bar{G}_{\mu\nu} = 0$ gives $\bar{\nabla}^\mu\big(\Lambda_{\text{eff}}\bar{\Sigma}_{\mu\nu}\big) = \kappa'\,\bar{\nabla}^\mu\bar{T}_{\mu\nu}^{(\text{nonvac})}$. For constant Λ_{eff} this reduces to $\bar{\nabla}^\mu\bar{T}_{\mu\nu}^{(\text{nonvac})} = 0$, showing

compatibility with covariant conservation. In regimes where state–dependence is weakly varying, the induced gradient terms are bounded as discussed in Section 11.5.7.

Step 5 (Null drop–out and focusing). Because the vacuum stress is proportional to $\bar{\Sigma}_{\mu\nu}$, its null contraction vanishes, $k^\mu k^\nu \bar{T}^{(\mathrm{vac})}_{\mu\nu} = 0$, so the vacuum term does not affect the NCC or null focusing (Section 11.6 and Lemmas 11.123 and 11.124). This is consistent with keeping the vacuum on the geometric/LHS.

Step 6 (Equivalence with the 1PI effective action). At the expectation level, the renormalized 1PI action admits the local split (17.25),

$$\Gamma = \int \mathrm{d}\mu_{\bar{\Sigma}} \left[\frac{1}{2\kappa'} (\bar{R} - 2\Lambda) - \varepsilon_\Lambda \right] + \Gamma_{\mathrm{nonvac}},$$

where ε_Λ is the curvature–independent vacuum density (see Definition 17.11). Varying Γ w.r.t. $\bar{\Sigma}^{\mu\nu}$ reproduces (17.21) and thus $\Lambda_{\mathrm{eff}} = \Lambda + \kappa' \varepsilon_\Lambda$ (Proposition 17.12). The equivalence with the stress–tensor definition of ε_Λ in Definition 17.13 is established in Theorem 17.15.

Step 7 (Operator–valued coupling placement). In the response formulation (Definition 11.109), the kernel acts on the *non–vacuum* source while the vacuum resides on the geometric side:

$$\bar{G}_{\mu\nu} + \Lambda_{\mathrm{eff}} \bar{\Sigma}_{\mu\nu} = \int d\mu_{\bar{\Sigma}}(x')\, \bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x; x')\, \bar{T}^{(\mathrm{nonvac})}_{\rho\sigma}(x'),$$

(eq. (17.26)), avoiding double counting and matching (17.21). This is also the expectation–level statement referred to in eq. (11.269).

Step 8 (Scheme independence of the physical identification). Local, finite counterterms shift Λ in (17.25), but the combination $\Lambda_{\mathrm{eff}} = \Lambda + \kappa' \varepsilon_\Lambda$ and the field equation (17.21) are invariant under such reparametrizations (the shift of Λ is compensated by the corresponding shift extracted from the vacuum functional). Hence Λ_{eff} is the unambiguous, physically meaningful constant controlling late–time acceleration (Sections 17.1.2 and 17.1.4).

Collecting Steps 1–8 establishes (17.21) and the identification $\Lambda_{\mathrm{eff}} = \Lambda + \kappa' \varepsilon_\Lambda$ via three equivalent routes: direct field–equation rearrangement, variation of the 1PI effective action, and the operator–valued response law. □

Corollary 17.7 (Trace relation and consistency with the vacuum residue)**.** *Let D be the spacetime dimension and define the non–vacuum trace $\bar{T}^{(\mathrm{nonvac})} := \bar{\Sigma}^{\mu\nu} \bar{T}^{(\mathrm{nonvac})}_{\mu\nu}$. Taking the $\bar{\Sigma}^{\mu\nu}$–trace of (17.21) yields, in general D,*

$$\frac{2 - D}{2}\, \bar{R} \;+\; D\, \Lambda_{\mathrm{eff}} \;=\; \kappa'\, \bar{T}^{(\mathrm{nonvac})}, \qquad \Lambda_{\mathrm{eff}} := \Lambda + \kappa' \varepsilon_\Lambda. \tag{17.23}$$

Specializing to $D = 4$ gives the scalar relation

$$\boxed{\bar{R} \;-\; 4\, \Lambda_{\mathrm{eff}} \;=\; -\, \kappa'\, \bar{T}^{(\mathrm{nonvac})}} \tag{17.24}$$

in agreement with Proposition 17.6.

Proof. Contract (17.21) with $\bar{\Sigma}^{\mu\nu}$ and use the identities $\bar{\Sigma}^{\mu\nu} \bar{G}_{\mu\nu} = \frac{2-D}{2} \bar{R}$ and $\bar{\Sigma}^{\mu\nu} \bar{\Sigma}_{\mu\nu} = D$ to obtain (17.23); setting $D = 4$ yields (17.24). □

Remark 17.8 (Consistency with the definition of ε_Λ)**.** Starting instead from the *bare* equation (17.18) and the source split $\bar{T}^{\text{matter}}_{\mu\nu} = \bar{T}^{(\text{nonvac})}_{\mu\nu} - \varepsilon_\Lambda\,\bar{\Sigma}_{\mu\nu}$ (Definition 17.5 and eq. (17.20)), the $D = 4$ trace gives

$$-\bar{R} + 4\Lambda \;=\; \kappa'\,\bar{T}^{(\text{nonvac})} \;-\; 4\kappa'\varepsilon_\Lambda \quad\Longrightarrow\quad \bar{R} - 4(\Lambda + \kappa'\varepsilon_\Lambda) = -\kappa'\,\bar{T}^{(\text{nonvac})},$$

which is exactly (17.24). Moreover, by Definition 17.13, $\varepsilon_\Lambda = -\frac{1}{4}\,\bar{\Sigma}^{\mu\nu}\,\langle{:}\,T^{\text{matter}}_{\mu\nu}\,{:}\rangle_{\text{vac}}$, so the vacuum trace contributes $\bar{\Sigma}^{\mu\nu}\bar{T}^{(\text{vac})}_{\mu\nu} = -\,4\,\varepsilon_\Lambda$ in $D = 4$, matching the algebraic factor in the derivation above and ensuring that moving the vacuum piece to the geometric side (to form Λ_{eff}) leaves the traced equation consistent (Proposition 17.12 and Theorem 17.15).

Remark 17.9 (Vacuum drop-out on nulls; Bianchi consistency)**.** Because $\bar{T}^{(\text{vac})}_{\mu\nu} \propto \bar{\Sigma}_{\mu\nu}$, we have $k^\mu k^\nu \bar{T}^{(\text{vac})}_{\mu\nu} = 0$ whenever k is $\bar{g}^{\text{Lor}}$–null (cf. Lemmas 11.123 and 11.124); hence the vacuum term in (17.21) does not affect the NCC and null focusing (Section 11.6). For constant Λ_{eff}, $\bar{\nabla}^\mu(\Lambda_{\text{eff}}\bar{\Sigma}_{\mu\nu}) = 0$, so (17.21) is compatible with $\bar{\nabla}^\mu\bar{G}_{\mu\nu} = 0$ and with source conservation to the order discussed in Section 11.5.7 (see also the expectation-level discussion around eq. (11.269)).

1PI effective action derivation (equivalent route). At the expectation level, the renormalized 1PI effective action has the local form

$$\Gamma[\bar{\Sigma}, \ldots] = \int_{\mathcal{M}} \sqrt{-\det\bar{\Sigma}}\;\mathrm{d}^4x\,\left[\frac{1}{2\kappa'}\big(\bar{R} - 2\Lambda\big) \;-\; \varepsilon_\Lambda\right] \;+\; \Gamma_{\text{nonvac}}[\bar{\Sigma}, \ldots], \tag{17.25}$$

where the state-dependent, finite constant density ε_Λ encodes the holosymmetric vacuum residue (Section 13.2.7), and Γ_{nonvac} collects non-vacuum contributions. Varying (17.25) with respect to $\bar{\Sigma}^{\mu\nu}$ reproduces (17.21), i.e.

$$\bar{G}_{\mu\nu} + \big(\Lambda + \kappa'\varepsilon_\Lambda\big)\bar{\Sigma}_{\mu\nu} = \kappa'\,\bar{T}^{(\text{nonvac})}_{\mu\nu}.$$

Compatibility with the operator–valued coupling. Recall the operator response (Def. 11.109 and the notes thereafter):

$$\hat{G}_{\mu\nu}(x) = \int d\mu_{\bar{\Sigma}}(x')\,\hat{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x')\;\circ\;\hat{T}_{\rho\sigma}(x'),$$

and its expectation in $|\mathring{\boldsymbol{\Psi}}\rangle$, where $d\mu_{\bar{\Sigma}} = \sqrt{-\det\bar{\Sigma}}\,\mathrm{d}^4x'$ and the Jordan product symmetrizes operator order. To *avoid double counting* the vacuum, we stipulate that the kernel acts on the *non-vacuum* source only:

$$\bar{G}_{\mu\nu}(x) + \Lambda_{\text{eff}}\bar{\Sigma}_{\mu\nu}(x) = \int d\mu_{\bar{\Sigma}}(x')\,\bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x')\,\bar{T}^{(\text{nonvac})}_{\rho\sigma}(x'), \tag{17.26}$$

which reduces, in the EP-safe scalarised regime (Section 11.5.7), to $\bar{G}_{\mu\nu} + \Lambda_{\text{eff}}\bar{\Sigma}_{\mu\nu} = \kappa_{\text{eff}}(x)\,\bar{T}^{(\text{nonvac})}_{\mu\nu}$ (cf. eq. (11.269)).

Spectral/hamiltonian semantics. Equation (17.12) cleanly separates (i) the *bare geometric* constant Λ from the action (17.17) (Section 17.1.1) and (ii) the *state-dependent* residue ε_Λ extracted from the vacuum sector via coherent, state-conditioned renormalization (Ch. 5; Section 13.2.7). In parallel, the geometric ground-state density Λ_{geo} defined by

$$\hat{H}_{\text{geometry}}\,|\Psi_{\text{vac}}\rangle = \Lambda_{\text{geo}}\,|\Psi_{\text{vac}}\rangle\,, \qquad \Lambda_{\text{geo}} = \frac{\langle\Psi_{\text{vac}}|\hat{H}_{\text{geometry}}|\Psi_{\text{vac}}\rangle_{\text{phys}}}{\text{Vol}(\mathcal{X}_q)}, \tag{17.27}$$

(Sections 6.1.8 and 17.1.2) accounts for the pure-geometry vacuum; matter/coherence contributions shift the observable constant precisely by $\kappa'\varepsilon_\Lambda$, yielding Λ_{eff} (cf. eq. (11.269)).

Collecting the above, the *effective cosmological constant* entering the Chronogeometric Einstein Equations and the response law is unambiguously $\Lambda_{\text{eff}} = \Lambda + \kappa'\varepsilon_\Lambda$, with ε_Λ fixed by the Holo-State via holosymmetric selection and coherent, diffeomorphism-covariant renormalization.

Remark 17.10 (Must the cosmological term use the four–volume?)**.** If (i) the only geometric field is the unified chronogeometric tensor $\bar{\Sigma}_{\mu\nu}$, (ii) the action is local, diffeomorphism covariant, and (iii) field equations are at most second order, then in $D = 4$ the Lovelock uniqueness result implies that the only zeroth–derivative scalar density built from $\bar{\Sigma}$ is the volume form $\mathrm{d}\mu_{\bar{\Sigma}} = \sqrt{-\det\bar{\Sigma}}\,\mathrm{d}^4x$ (up to a boundary term) [662, 663, 267]. Hence the cosmological term is necessarily proportional to the *four–volume*, as used in Section 17.1.2 and eq. (17.7). This placement yields $T^{\text{vac}}_{\mu\nu} \propto \bar{\Sigma}_{\mu\nu}$ and guarantees the null drop–out used in Section 11.6.

Alternatives (require extra structure):

(i) Unimodular variant.
Impose a fixed volume density by a Lagrange multiplier λ:

$$S = \frac{1}{2\kappa'}\int \bar{R}\,\mathrm{d}\mu_{\bar{\Sigma}} + \int \lambda(x)\Big(\sqrt{-\det\bar{\Sigma}} - \omega_0(x)\Big)\mathrm{d}^4x + S_{\text{matter}}.$$

Varying gives $\bar{G}_{\mu\nu} + \lambda\,\bar{\Sigma}_{\mu\nu} = \kappa' T_{\mu\nu}$ and $\bar{\nabla}^\mu$ of this equation implies $\partial_\mu\lambda = 0$, so $\lambda = \Lambda_{\text{int}}$ is an *integration constant*. One recovers the same structure as eq. (17.9), with $\Lambda \to \Lambda_{\text{int}}$, and after including the holosymmetric residue one again arrives at $\Lambda_{\text{eff}} = \Lambda_{\text{int}} + \kappa'\varepsilon_\Lambda$ (Section 17.1.3).

(ii) Dynamical 4–form / two–measure models.
Replace $\mathrm{d}\mu_{\bar{\Sigma}}$ by a metric–independent scalar density (e.g. a 4–form field strength or $\Phi(\varphi^a)\,\mathrm{d}^4x$ built from four scalars). The associated equation of motion makes the 4–form flux a constant, which again enters the metric equations as an effective cosmological constant. After adding the state–dependent residue, one still obtains $\Lambda_{\text{eff}} = \Lambda_{\text{bare}} + \kappa'\varepsilon_\Lambda$ (Section 17.1.3).

(iii) State–conditioned density.
One may contemplate a scalar density $\omega[|\mathring{\boldsymbol{\Psi}}\rangle, \bar{\Sigma}]$ replacing $\sqrt{-\det\bar{\Sigma}}$, provided it transforms as a density and its variation preserves the Bianchi identity. Any change by a total divergence or a BRST–exact term leaves the field equations and the identification of Λ_{eff} unchanged (Section 17.1.3).

Conclusion. Within the minimal, metric–only (here: $\bar{\Sigma}$–only) and second–order framework used in this chapter, the four–volume density is the natural and essentially unique choice. Alternative formulations are viable but introduce extra fields or constraints; they remain compatible with the effective identification $\Lambda_{\text{eff}} = \Lambda + \kappa'\varepsilon_\Lambda$ and with the operator–valued coupling placement in Definition 11.109.

17.1.4 Definition and Computation of the Holosymmetric Vacuum Residue

We now formalize the quantity *holosymmetric vacuum residue* ε_Λ that appears in $\Lambda_{\text{eff}} = \Lambda + \kappa'\varepsilon_\Lambda$ (Section 17.1.3). Throughout we work on the BRST/GNS physical sector selected by the Holo–State $|\mathring{\Psi}\rangle$ with Hadamard regularity (Sections 1.4.1 and 6.1.8), and we use the unified chronogeometric tensor $\bar{\Sigma}_{\mu\nu}$ and measure $\mathrm{d}\mu_{\bar{\Sigma}} = \sqrt{-\det\bar{\Sigma}}\,\mathrm{d}^4x$.

Definition 17.11 (Holosymmetric vacuum residue via the 1PI effective action)**.** Let $\Gamma[\bar{\Sigma},\dots]$ be the renormalized one–particle–irreducible (1PI) effective action on the physical sector. In a local covariant splitting consistent with Section 17.1.2, write

$$\Gamma[\bar{\Sigma},\dots] = \int_{\mathcal{M}} \mathrm{d}\mu_{\bar{\Sigma}} \left[\frac{1}{2\kappa'}\big(\bar{R} - 2\Lambda\big) \;-\; \varepsilon_\Lambda\right] \;+\; \Gamma_{\text{nonvac}}[\bar{\Sigma},\dots], \tag{17.28}$$

where Γ_{nonvac} contains curvature–dependent and source–dependent non–vacuum contributions. The constant scalar density ε_Λ is defined as the *curvature–independent* local part of the vacuum functional on the chosen Holo–State.

Proposition 17.12 (Field equations and Λ_{eff} from Γ)**.** *Varying* (17.25) *with respect to* $\bar{\Sigma}^{\mu\nu}$ *yields*

$$\bar{G}_{\mu\nu} + \big(\Lambda + \kappa'\varepsilon_\Lambda\big)\bar{\Sigma}_{\mu\nu} = \kappa'\,\bar{T}^{(\text{nonvac})}_{\mu\nu}, \tag{17.29}$$

i.e. $\Lambda_{\text{eff}} = \Lambda + \kappa'\varepsilon_\Lambda$*, in agreement with Section 17.1.3 (cf. the expectation–level placement eq.* (11.269)*).*

Proof. Step 0 (Setup and conventions). Consider the effective action (local sector)

$$\Gamma[\bar{\Sigma},\dots] = \int_{\mathcal{M}} \sqrt{-\det\bar{\Sigma}}\,\mathrm{d}^4x \left[\frac{1}{2\kappa'}\big(\bar{R} - 2\Lambda\big) \;-\; \varepsilon_\Lambda\right] \;+\; \Gamma_{\text{nonvac}}[\bar{\Sigma},\dots], \tag{17.30}$$

where ε_Λ is the curvature–independent vacuum density (homogeneous sector; Definition 17.11) and Γ_{nonvac} collects non–vacuum contributions. We vary w.r.t. $\bar{\Sigma}^{\mu\nu}$ on the BRST/GNS physical sector (Section 1.4.1). The stress–tensor convention is the standard one used in Section 9.17.9:

$$\delta\Gamma_{\text{nonvac}} = -\frac{1}{2}\int \sqrt{-\det\bar{\Sigma}}\;\bar{T}^{(\text{nonvac})}_{\mu\nu}\;\delta\bar{\Sigma}^{\mu\nu}\,\mathrm{d}^4x. \tag{17.31}$$

Step 1 (Geometric variation). The Einstein–Hilbert part with cosmological term varies as (cf. eq. (17.8))

$$\delta\Big[\frac{1}{2\kappa'}\int\sqrt{-\det\bar{\Sigma}}\,(\bar{R} - 2\Lambda)\,\mathrm{d}^4x\Big] = \frac{1}{2\kappa'}\int\sqrt{-\det\bar{\Sigma}}\,\big(\bar{G}_{\mu\nu} + \Lambda\,\bar{\Sigma}_{\mu\nu}\big)\,\delta\bar{\Sigma}^{\mu\nu}\,\mathrm{d}^4x + \delta S_{\text{bdy}}, \tag{17.32}$$

where $\delta S_{\rm bdy}$ is cancelled by the GHY–type boundary term (already included in the underlying action; see Section 17.1.1).

Step 2 (Vacuum density variation). For the curvature–independent density $-\varepsilon_\Lambda$,

$$\delta\Big[-\int\sqrt{-\det\bar\Sigma}\;\varepsilon_\Lambda\,\mathrm{d}^4x\Big]=-\int\varepsilon_\Lambda\,\delta\sqrt{-\det\bar\Sigma}\,\mathrm{d}^4x=+\frac{1}{2}\int\sqrt{-\det\bar\Sigma}\;\varepsilon_\Lambda\,\bar\Sigma_{\mu\nu}\,\delta\bar\Sigma^{\mu\nu}\,\mathrm{d}^4x,\quad(17.33)$$

using $\delta\sqrt{-\det\bar\Sigma}=-\frac{1}{2}\sqrt{-\det\bar\Sigma}\,\bar\Sigma_{\mu\nu}\delta\bar\Sigma^{\mu\nu}$.

Step 3 (Collecting terms). Summing the variations and using (17.31) gives

$$\delta\Gamma=\int\sqrt{-\det\bar\Sigma}\,\Big[\frac{1}{2\kappa'}\big(\bar G_{\mu\nu}+\Lambda\,\bar\Sigma_{\mu\nu}\big)+\frac{1}{2}\,\varepsilon_\Lambda\,\bar\Sigma_{\mu\nu}-\frac{1}{2}\,\bar T^{(\rm nonvac)}_{\mu\nu}\Big]\delta\bar\Sigma^{\mu\nu}\,\mathrm{d}^4x.$$

Stationarity for arbitrary $\delta\bar\Sigma^{\mu\nu}$ yields

$$\frac{1}{\kappa'}\big(\bar G_{\mu\nu}+\Lambda\,\bar\Sigma_{\mu\nu}\big)+\varepsilon_\Lambda\,\bar\Sigma_{\mu\nu}-\bar T^{(\rm nonvac)}_{\mu\nu}=0,$$

which is equivalent to

$$\bar G_{\mu\nu}+\big(\Lambda+\kappa'\varepsilon_\Lambda\big)\bar\Sigma_{\mu\nu}=\kappa'\,\bar T^{(\rm nonvac)}_{\mu\nu}.\quad(17.34)$$

This is precisely (17.29) with $\Lambda_{\rm eff}=\Lambda+\kappa'\varepsilon_\Lambda$.

Step 4 (Consistency notes). (i) **Bianchi/Conservation.** Taking $\bar\nabla^\mu$ and using $\bar\nabla^\mu\bar G_{\mu\nu}=0$ shows compatibility with $\bar\nabla^\mu\bar T^{(\rm nonvac)}_{\mu\nu}=0$ when $\Lambda_{\rm eff}$ is constant (or within the controlled variations of Section 11.5.7). (ii) **Equivalence of definitions of ε_Λ.** The ε_Λ used in (17.30) equals the curvature–independent part of the vacuum functional; by Definition 17.13 and Theorem 17.15 it also equals $-\frac{1}{4}\bar\Sigma^{\rho\sigma}\langle{:}\,T^{\rm matter}_{\rho\sigma}\,{:}\rangle_{\rm vac}$, ensuring the same $\Lambda_{\rm eff}$ is obtained from the stress–tensor route. (iii) **Operator–kernel placement.** The expectation–level response eq. (11.269) follows by keeping the vacuum on the geometric side and letting the coupling kernel act on $\bar T^{(\rm nonvac)}$ (Definition 11.109), thereby avoiding double counting.

These remarks establish full consistency with Section 17.1.3. □

Coherent (state–conditioned) evaluation of ε_Λ. To compute ε_Λ from (17.25), decompose the quadratic fluctuations into sectors $i\in\{h,\text{ghosts},\Theta,\text{mix}\}$ with kinetic operators $\Delta_i[\bar\Sigma]$ (TT purity and $\mathcal{P}^{(2)}[\hat\tau]=0$ as in Definition 10.182 and Section 10.18.7). Define the *coherent trace*

$$\mathrm{Tr}_{\rm coh}F(\Delta_i):=\int_{\mathcal{M}}\mathrm{d}\mu_{\bar\Sigma}\int_0^\infty\mathrm{d}\lambda\;\rho_i(\lambda;\bar\Sigma)\;w_i(\lambda\,;\,|\mathring{\mathbf{\Psi}}\rangle)\;F(\lambda),\quad(17.35)$$

where ρ_i is the state– and background–dependent spectral density, and $w_i(\lambda)$ are the state–conditioned modal weights implementing the endogenous UV/IR band via the Chapter 4 machinery: state–conditioned modal weights Definition 4.392, finite entanglement capacity Principle 4.1, effective spectral cutoff Definition 4.576, and IR windowing/coherence scale Section 4.9.15 and Definition 4.435 (see also the coherent–regularization statement Proposition 4.424). Then

$$\Gamma_{\rm vac}=\frac{1}{2}\sum_i(-1)^{F_i}\,\mathrm{Tr}_{\rm coh}\log\Delta_i=-\frac{1}{2}\sum_i(-1)^{F_i}\int_0^\infty\frac{\mathrm{d}s}{s}\;\mathrm{Tr}_{\rm coh}\big(e^{-s\Delta_i}\big),\quad(17.36)$$

and the constant (curvature–independent) piece of $\Gamma_{\rm vac}$ determines ε_Λ via (17.25). In a heat–kernel expansion, $\mathrm{Tr}\, e^{-s\Delta_i} \sim (4\pi s)^{-2} \sum_{n\geq 0} a_n^{(i)} s^n$, the constant density is governed by the weighted $a_0^{(i)}$ coefficients plus finite state–dependent remnants selected by w_i.

Definition 17.13 (Holosymmetric vacuum residue via stress–tensor contraction)**.** Let $:T_{\mu\nu}^{\rm matter}:$ be the Wick–ordered, renormalized stress tensor (Hadamard point–splitting on the chosen GNS representation; Section 1.4.1). Denote by $\langle\cdot\rangle_{\rm vac}$ the homogeneous/isotropic vacuum component selected by the holosymmetric filter (Sections 13.2.7 and 13.4.7). Then, in $D = 4$, the holosymmetric vacuum residue ε_Λ may be specified in either of the following equivalent forms:

(i) Pointwise (homogeneous vacuum makes the RHS x–independent):

$$\boxed{\varepsilon_\Lambda = -\frac{1}{4}\,\bar{\Sigma}^{\mu\nu}(x)\,\Big\langle \overset{\circ}{\boldsymbol{\Psi}}\Big|\; :T_{\mu\nu}^{\rm matter}(x):\; \Big|\overset{\circ}{\boldsymbol{\Psi}}\Big\rangle_{\rm !phys,\,vac}} \tag{17.37}$$

(ii) Volume–averaged (makes constancy manifest without assuming homogeneity a priori):

$$\boxed{\varepsilon_\Lambda = -\frac{1}{4\,\mathrm{Vol}(\mathcal{X}_q)}\int_{\mathcal{X}_q} \mathrm{d}\mu_{\bar{\Sigma}}(x)\,\bar{\Sigma}^{\mu\nu}(x)\,\Big\langle \overset{\circ}{\boldsymbol{\Psi}}\Big|\; :T_{\mu\nu}^{\rm matter}(x):\; \Big|\overset{\circ}{\boldsymbol{\Psi}}\Big\rangle_{\rm !phys,\,vac}} \tag{17.38}$$

where $\mathrm{d}\mu_{\bar{\Sigma}} = \sqrt{-\det \bar{\Sigma}}\,\mathrm{d}^4x$ and $\mathrm{Vol}(\mathcal{X}_q) = \int_{\mathcal{X}_q}\mathrm{d}\mu_{\bar{\Sigma}}$.

In a homogeneous/isotropic vacuum sector—or by reading $\langle\cdot\rangle_{\rm vac}$ as "vacuum–projected physical expectation"—both expressions reduce to the compact shorthand used elsewhere:

$$\boxed{\varepsilon_\Lambda = -\frac{1}{4}\,\bar{\Sigma}^{\mu\nu}\,\Big\langle\; :T_{\mu\nu}^{\rm matter}:\;\Big\rangle_{\rm vac}} \tag{17.39}$$

and the associated vacuum stress reads $\bar{T}_{\mu\nu}^{(\rm vac)} = -\,\varepsilon_\Lambda\,\bar{\Sigma}_{\mu\nu}$ (Definition 17.5). The factor $-\frac{1}{4}$ follows from $\bar{\Sigma}^{\mu\nu}\bar{\Sigma}_{\mu\nu} = 4$ in $D = 4$.

Remark 17.14 (Meaning of the Hadamard/Wick ordering)**.**

Semantic role

The notation $:\mathcal{O}(x):$ denotes the *renormalized (Wick–ordered) local composite operator* obtained by subtracting the state–independent short–distance singular structure from the underlying *Hadamard* two–point function before taking the coincidence limit. This is the curved–spacetime generalization of flat–space normal ordering and ensures locality, covariance, and well–defined composite fields [10, 13, 12].

Point–splitting definition (schematic)

Let

$$W_\Psi(x,x') = \Big\langle \overset{\circ}{\boldsymbol{\Psi}}\Big|\hat{\Phi}(x)\hat{\Phi}(x')\Big|\overset{\circ}{\boldsymbol{\Psi}}\Big\rangle_{\rm !phys}$$

be the (Hadamard) two–point function in the chosen GNS representation (Section 1.4.1). For a local composite $\mathcal{O}(x) = \mathcal{D}[\hat{\Phi}](x)$ built from the field and its derivatives, its Wick–ordered version is defined by

$$:\mathcal{O}(x): := \lim_{x' \to x} \Big(\mathcal{D}_{x,x'} W_\Psi(x,x') \; - \; \mathcal{D}_{x,x'} H(x,x') \Big) \; + \; \text{finite local counterterms},$$

where $H(x,x')$ is the *Hadamard parametrix* capturing the universal, state–independent singularity structure, and $\mathcal{D}_{x,x'}$ is the appropriate bi–differential operator for $\mathcal{O}$ [10, 12]. For the stress tensor $T^{\text{matter}}_{\mu\nu}$, one uses the corresponding conserved bi–differential operator before coincident limiting [10, 13].

Renormalization freedom and conservation

The subtraction above is unique up to finite, local curvature counterterms (e.g. multiples of $\bar{\Sigma}_{\mu\nu}$, $\bar{G}_{\mu\nu}$, etc.), which correspond to renormalizations of the gravitational couplings in the action (including shifts of Λ) and can be fixed by covariance, conservation, and scheme conditions [10, 12]. This is the sense in which Γ in eq. (17.25) absorbs local ambiguities, and why Λ may shift while Λ_{eff} remains physically meaningful (Section 17.1.3).

Vacuum expectations need not vanish

In curved/relational settings, Wick ordering *does not* force $\langle : T^{\text{matter}}_{\mu\nu} : \rangle$ to vanish. Rather, it produces a finite, covariant quantity, whose curvature–independent piece is precisely the holosymmetric vacuum residue entering ε_Λ in Definition 17.13. Thus,

$$\varepsilon_\Lambda \;=\; -\tfrac{1}{4}\,\bar{\Sigma}^{\mu\nu} \Big\langle :T^{\text{matter}}_{\mu\nu}: \Big\rangle_{\text{vac}}$$

(eqs. (17.37) to (17.39)), and $\bar{T}^{(\text{vac})}_{\mu\nu} = -\varepsilon_\Lambda\, \bar{\Sigma}_{\mu\nu}$ (Definition 17.5).

Born/GNS valuation and notation

Expectations such as $\langle \mathring{\boldsymbol{\Psi}}| \; :T^{\text{matter}}_{\mu\nu}(x): \; |\mathring{\boldsymbol{\Psi}}\rangle_{!\text{phys}}$ are taken in the physical GNS representation selected by the Holo–State (Section 1.4.1); the subscript "vac" designates the homogeneous/isotropic vacuum component picked out by the holosymmetric filter (Sections 13.2.7 and 13.4.7). Typographically, the tight spacing $:\cdots:$ simply attaches the colons to the operator.

Theorem 17.15 (Equivalence of the 1PI and stress–tensor definitions)**.** *Under the assumptions above (Hadamard, BRST/GNS, coherent regularization), the ε_Λ extracted from (17.25) equals the one from (17.39). Consequently $\Lambda_{\text{eff}} = \Lambda + \kappa' \varepsilon_\Lambda$ in (17.29) is unique and scheme–independent up to shifts of Λ by local covariant counterterms.*

Proof sketch. Varying Γ w.r.t. $\bar{\Sigma}^{\mu\nu}$ gives the standard identity $\frac{2}{\sqrt{-\det \bar{\Sigma}}} \frac{\delta \Gamma}{\delta \bar{\Sigma}^{\mu\nu}} = -\frac{1}{2\kappa'}\big(\bar{G}_{\mu\nu} + \Lambda \bar{\Sigma}_{\mu\nu}\big) - \frac{1}{2}\langle : T^{\text{matter}}_{\mu\nu} : \rangle$. Taking the homogeneous vacuum component (holosymmetric filter) and the trace with $\bar{\Sigma}^{\mu\nu}$ isolates the constant density on the RHS, which yields (17.39); inserting back reproduces (17.25). The coherent trace (17.35) ensures UV/IR finiteness consistent with the Chapter 4 scheme (Proposition 4.424, Principle 4.1, Definitions 4.435 and 4.576, and Section 4.9.15). □

Remark 17.16 (Null drop–out and response kernel placement)**.** Because $\bar{T}^{(\mathrm{vac})}_{\mu\nu} \propto \bar{\Sigma}_{\mu\nu}$, one has $k^\mu k^\nu \bar{T}^{(\mathrm{vac})}_{\mu\nu} = 0$ for every $\bar{g}^{\mathrm{Lor}}$–null k^μ (Lemma 11.123), so the NCC and null focusing (Section 11.6) are unaffected. In the operator–valued response (Definition 11.109 and notes), the vacuum term resides on the geometric side:

$$\bar{G}_{\mu\nu} + \Lambda_{\mathrm{eff}} \bar{\Sigma}_{\mu\nu} = \int \mathrm{d}\mu_{\bar{\Sigma}}(x')\, \bar{\mathcal{C}}_{\mu\nu}{}^{\rho\sigma}(x;x')\, \bar{T}^{(\mathrm{nonvac})}_{\rho\sigma}(x'), \tag{17.40}$$

avoiding double counting and matching the scalarised limit in Section 11.5.7 (cf. eq. (11.269)).

Observational back–out (phenomenology). If Λ is treated as a bare action parameter (Section 17.1.2) and Λ_{eff} is inferred from late–time data, then

$$\varepsilon_\Lambda \;=\; \frac{\Lambda_{\mathrm{eff}} - \Lambda}{\kappa'}\,. \tag{17.41}$$

Equation (17.41) provides a phenomenological estimate consistent with the first–principles definitions (17.25) and (17.39).

Minimal procedure (spectral route). To evaluate ε_Λ in practice:

(i) Fix $(\bar{\Sigma}_{\mu\nu}, |\mathring{\mathbf{\Psi}}\rangle)$ on a homogeneous patch (Section 6.1.8).

(ii) Build sector operators $\Delta_i[\bar{\Sigma}]$ with projector constraints (TT purity for h, $\mathcal{P}^{(2)}[\hat{\tau}] = 0$; Definition 10.182 and Section 10.18.7).

(iii) Compute $\mathrm{Tr}_{\mathrm{coh}} e^{-s\Delta_i}$ using state–conditioned weights w_i from Chapter 4 (Definitions 4.392, 4.435 and 4.576, Principle 4.1, and Section 4.9.15) and extract the curvature–independent piece of Γ_{vac} via (17.36).

(iv) Read off ε_Λ from (17.25); cross–check with (17.39).

This yields a finite, covariant ε_Λ consistent with holosymmetric selection and with the placement of Λ_{eff} in the field equations (Section 17.1.3 and eq. (11.269)).

The Measured Value of the Cosmological Constant

Modern cosmology constrains the late–time acceleration with high precision. In the minimal ΛCDM fit, the *dark–energy density parameter* is

$$\Omega_\Lambda \simeq 0.685 \quad \text{(Planck 2018 baseline + external datasets)}, \tag{17.42}$$

indicating that dark energy comprises $\sim 68.5\%$ of the total energy budget of the universe [136, 414, 212, 211]. Using the critical *energy* density

$$\rho^E_{\mathrm{crit}} = \frac{3c^2 H_0^2}{8\pi G}, \qquad \text{(with } H_0 \text{ from [136])}, \tag{17.43}$$

the vacuum (dark–energy) density follows as

$$\rho_{\mathrm{vac}} = \Omega_\Lambda\, \rho^E_{\mathrm{crit}} \;\approx\; (5.3 \pm 0.3) \times 10^{-10}\ \mathrm{J\,m^{-3}}, \tag{17.44}$$

i.e. the energy equivalent of only a few hydrogen atoms per cubic meter, yet with profound dynamical impact [212, 211].

The corresponding *cosmological constant* that appears in the field equations (cf. eq. (17.9)) is obtained by the standard mapping

$$\Lambda_{\text{eff}} = \frac{8\pi G}{c^4}\,\rho_{\text{vac}} \;\approx\; (1.1 \pm 0.1) \times 10^{-52}\ \text{m}^{-2}, \tag{17.45}$$

which is equivalently

$$\Lambda_{\text{eff}} = \frac{3\,\Omega_\Lambda H_0^2}{c^2},$$

and is consistent with the values inferred from distance–redshift and CMB anisotropy data [136, 414].

Bridge to the IQR framework. Within the present treatise, the quantity constrained by observations is Λ_{eff} of eq. (17.45), which in our notation is

$$\Lambda_{\text{eff}} = \Lambda + \kappa'\,\varepsilon_\Lambda, \tag{17.46}$$

combining the *bare geometric* four–volume coefficient Λ from the action (Section 17.1.1) with the *interferometric vacuum residue* ε_Λ selected by the holosymmetric filter (Sections 13.2.7 and 13.4.7). Numerically inserting Ω_Λ from eq. (17.42) into eq. (17.44) and then eq. (17.45) recovers the measured Λ_{eff} that drives the observed acceleration, closing the loop between the empirical ΛCDM description and the state–first, relational construction of Sections 13.6 and 17.1.1 and eq. (11.269).

Finite sectoral vacua (no large divergences). Each sectoral vacuum contribution is rendered finite by *coherent (state–conditioned) regularization*: the state–conditioned modal weights suppress unsupported channels, the *finite entanglement capacity* furnishes a UV band edge, and *intrinsic entanglement dressing* together with IR windowing provides an IR floor (Ch. 4; Principle 4.1, Definitions 4.435 and 4.576, Section 4.9.15, and Proposition 4.424). Consequently, both the geometric and the matter/coherence zero–point energies are individually well–defined and covariant in the spectral/trace semantics (eqs. (17.13) and (17.14)).

Near–cancellation enforced by holosymmetric selection. The *holosymmetric filter* selects a homogeneous/isotropic vacuum in which the geometric (Gaiamiti) and the material/coherence (Tantrakyma) sectors contribute with opposite sign and nearly cancel, leaving a small positive *interferometric residue* ε_Λ (Sections 13.2.7 and 13.4.7). Quantitatively,

$$\varepsilon_\Lambda = \frac{1}{\text{Vol}(\mathcal{X}_q)}\left|\left\langle \hat{H}_{\text{geometry}}\right\rangle_{!\text{phys}} + \left\langle \hat{H}_{\text{matter}}\right\rangle_{!\text{phys}}\right| \;\ll\; \frac{k_{\text{UV}}^4}{16\pi^2}\,\hbar c, \qquad \text{(eq. (17.15))},$$

so that the naively large UV scale collapses to a tiny residue fixed by the Holo–State.

Clean placement in the field equations. The residue ε_Λ shifts the geometric four–volume term to the *effective* constant $\Lambda_{\text{eff}} = \Lambda + \kappa'\varepsilon_\Lambda$ (Section 17.1.3, eq. (17.21), and Proposition 17.12). Placing the vacuum on the geometric (left–hand) side while letting the operator–valued coupling kernel act only on *non–vacuum* sources (Definition 11.109 and eq. (17.26)) avoids double counting and preserves the *null silence* of the vacuum (Lemmas 11.123 and 11.124).

Why small and positive. Smallness arises because the two finite sectoral pieces are holosymmetrically selected to be nearly equal and opposite; positivity follows from the filter's choice of the minimal positive residue compatible with the Holo–State constraints and the projector decomposition (Section 13.2.7 and Proposition 13.97). Thus Λ_{eff} is small and strictly > 0, in agreement with observation (Section 17.1.4).

Two equivalent definitions that lock the scale. The same ε_Λ is obtained from the 1PI effective action (eq. (17.25) and Proposition 17.12) and from the stress–tensor contraction $\varepsilon_\Lambda = -\frac{1}{4}\bar{\Sigma}^{\mu\nu}\big\langle :T^{\text{matter}}_{\mu\nu}: \big\rangle_{\text{vac}}$ (Definition 17.13 and Theorem 17.15). This equivalence shows the smallness is not a scheme artifact: local counterterm shifts of Λ are compensated within ε_Λ, leaving Λ_{eff} invariant.

Summary. Within the state–first, holosymmetric construction, the cosmological constant is small because a *self–regulated, near–exact cancellation* between finite geometric and matter/coherence vacua leaves a tiny positive *interferometric residue* ε_Λ, which enters the Chronogeometric Einstein Equations precisely as Λ_{eff} (eq. (17.21)).

17.1.5 The Universal Schrödinger Equation

The evolution of the entire cosmos, with all its constituent parts, is governed by a single dynamical law: the time-dependent Schrödinger equation for the **Quantum Holo-State** $|\mathring{\boldsymbol{\Psi}}\rangle$ (state–first/GNS; cf. Sections 6.1.13 and 13.6).

$$\boxed{i\hbar\frac{d}{dt}|\mathring{\boldsymbol{\Psi}}(t)\rangle = \hat{H}_{\text{rel}}(t)|\mathring{\boldsymbol{\Psi}}(t)\rangle} \tag{17.47}$$

Here $\hat{H}_{\text{rel}}$ is the *Total Relational Hamiltonian* defined in the unified Hamiltonian framework (Section 6.1.13); the relational time evolution is compatible with the split connection (eq. (6.314)). The operator driving this evolution can be conceptually decomposed as follows.

$$\hat{H}_{\text{rel}} = \underbrace{\hat{H}_{\text{geometry}}[\hat{\Sigma}_{\mu\nu}]}_{\text{Dark Energy \& Dark Matter}} + \underbrace{\hat{H}_{\text{matter}}[\hat{\phi},\hat{\psi},\ldots]}_{\text{Standard Model}}. \tag{17.48}$$

Coupling of sources to curvature is mediated via the operator–valued kernel of Definition 11.109 and eqs. (11.213) and (11.214).

This single Hamiltonian contains the entirety of cosmic phenomenology:

(i) The Geometric Hamiltonian ($\hat{H}_{\text{geometry}}$)

This operator governs the dynamics of the *Gaiamiti field* $\hat{\Phi}$ (Definition 6.2), the fundamental field of geometric potentiality whose quanta are the Apeirons.

(a) Its ground–state eigenvalue defines the *cosmological constant* Λ_{geo} (eq. (17.11) and Section 17.1.2); upon inclusion of the holosymmetric residue one obtains the effective constant $\Lambda_{\text{eff}} = \Lambda + \kappa'\varepsilon_\Lambda$ (eq. (17.12)).

(b) Its large–scale, coherent excited states furnish the dark–sector phenomenology (cf. Section 13.1 and Definition 13.1).

(ii) The Matter Hamiltonian ($\hat{H}_{\text{matter}}$)
This operator governs the dynamics of the Tantrakyma field $\hat{\Theta}$ (Definition 6.3), the fundamental field of material potentiality whose quanta are the Kymons.

(a) Its excitations and interactions realize the observed particle content (Standard Model) within the generalized field doublet $\hat{\mathbf{F}} = (\hat{\mathbf{\Phi}}, \hat{\mathbf{\Theta}})^\top$ (Definition 6.1).

Thus, the Universal Schrödinger Equation (eq. (17.47)) is the ultimate unifying statement of the treatise. The solution, $|\overset{\circ}{\mathbf{\Psi}}(t)\rangle$, is a single quantum state whose different aspects and excitations we perceive as the familiar world of particles and forces, the enigmatic halos of dark matter, and the pervasive, accelerating expansion driven by dark energy (cf. Section 13.1, Definition 13.1, and eq. (17.12)). They are not separate phenomena; they are different facets of one unified, evolving relational structure. While the *Universal Schrödinger Equation* stands as the ultimate dynamical law of the cosmos, the *Total Relational Hamiltonian*, $\hat{H}_{\text{rel}}$, that drives the evolution is not itself a fundamental entity (cf. Section 6.1.13). It is a global operator that emerges from a deeper, more localized layer of reality's relational structure. The most basic building block of all dynamics in the theory is the *Fourfold Inter-Modal Correlator*, $\hat{G}_\Lambda(x, x')$ (Definition 4.9 and eq. (4.85)), a *bi-local* (two-point) operator-valued function whose essential role is to define the quantum-relational link—the entanglement and coherence—between any two Primordial Quantum Singularities at points x and x' (see also the PQS/modal-fibre construction, Section 4.7).

The global operator that we identify as the Total Relational Hamiltonian is constructed from this fundamental correlator: one takes the *local* part of the correlator at coincidence, $\hat{G}_\Lambda(x, x)$, traces over internal indices to obtain a local energy density (trace–to–modes bridge, Section 6.1.10), and integrates this density over the emergent manifold $\mathcal{X}_q$ with the relational measure $\mathrm{d}\mu_{\bar{\Sigma}}$ (Definition 6.33). In particular, the geometric contribution is realized by the local-trace formula and its spectral avatar (eqs. (17.13) and (17.14)), while the total Hamiltonian fits into the unified evolution framework (Section 6.1.13).

Therefore, the relationship between the components of cosmic evolution is one of emergence:

(i) The Correlator ($\hat{G}_\Lambda$). Encodes the fundamental, microscopic rules of relational interaction at and between any two points in the cosmic network (Definition 4.9 and Section 6.1.10).

(ii) The Hamiltonian ($\hat{H}_{\text{rel}}$). Is the macroscopic, integrated sum of these microscopic interactions across the entire cosmos (Section 6.1.13); the geometric part follows from the local trace and spectral integral (eqs. (17.13) and (17.14)).

(iii) The Universal Schrödinger Equation. (eq. (17.47)) then governs how the *Quantum Holo-State* $|\mathring{\mathbf{\Psi}}\rangle$ evolves under the influence of this total, integrated interaction energy, with expectation-level gravitational response placed via eq. (11.269).

In this light, the majestic evolution of the universe described by the Schrödinger equation is the global manifestation of the profound and intricate relational dynamics encoded within the Fourfold Inter-Modal Correlator (Definition 4.9 and eq. (4.85)), consistently tied to the modal-fibre picture and operator-valued response kernel (Section 4.7, Definition 11.109, and eqs. (11.213) and (11.214)).

The treatise presents a profound, hierarchical model of reality, where each level of physical description emerges from a more fundamental, abstract layer. The ultimate source of all existence and dynamics is not a physical entity but a philosophical principle: the *Law of Dependent Origination* (Section 1.1.2), which states that all phenomena arise solely in relation to other phenomena. This principle is given its most potent mathematical expression in the *Axiom of Duality* (Axiom 1.10), $|0\rangle \leftrightarrow |1\rangle$, the proposition that to define a thing is to simultaneously define its complement. Existence is, therefore, inherently and irreducibly relational. It is this foundational axiom that serves as the singular wellspring from which the entire architecture of the cosmos emerges in a self-consistent logical hierarchy.

This relational tension inherent in the Axiom of Duality is not static; it gives rise to a "generative grammar" for all possible interactions, a logic formalized as the *Fourfold Dialectic Algebra* (Section 1.2.7, Definitions 1.16 and 1.23, and Lemma 1.20). This algebra dictates the necessary modes of all relational processes: *Symmetry* (identity), *Antisymmetry* (distinction), *Synthesis* (interaction), and *Transcendent Holosynthesis* (the emergence of novelty). The Fourfold Dialectic is then given concrete mathematical form in the abstract *Algebra of Observables*, $\mathcal{A}(\mathbb{C}_d)$, realized by the generalized field doublet generators (Definition 6.1)—the abstract *Gaiamiti* (geometric potentiality) and *Tantrakyma* (material potentiality) fields (Definitions 6.2 and 6.3).

The abstract algebra is given physical substance in the form of the *Primordial Quantum Singularity (PQS)* (see the PQS/modal-fibre construction in Section 4.7 and the associated duality operators Section 2.1.2 and Definition 2.12). The PQS is the fundamental, indivisible "atom" of reality—a localized, non-spatiotemporal entity that instantiates the abstract Gaiamiti and Tantrakyma fields as its own intrinsic properties. The entire cosmos is thus conceived as a vast, self-entangled relational network of these PQSs. To move from this network of potentialities to a concrete reality, a selecting principle is required. This is the *Quantum Holo-State*, $|\mathring{\mathbf{\Psi}}\rangle$ (Section 1.4.1). It is the specific, non-factorizable quantum state of the *entire PQS network*, mathematically represented as a functional ω^{Ψ} that assigns an expectation value to every possible relational configuration (state–first/constraint setting Section 13.6). It is the foundational information that selects one specific, physically consistent reality—one that is maximally coherent with the logic of the Fourfold Dialectic—from the infinite possibilities.

The realization of this reality is then accomplished via the *GNS (Gelfand–Naimark–Segal) construction* (Sections 1.4.1 and 13.6). The GNS construction is the mathematical engine that takes the algebra of the PQS network and the state functional ω^{Ψ} as inputs. It generates both a specific *Hilbert space* for the cosmos, $\mathcal{H}_{\text{em}}$, and a concrete *representation*, π^{Ψ}, of the Gaiamiti and Tantrakyma fields as actual operators that act on the states of the PQS

network. It is from this GNS construction that all concrete physical laws and operators emerge as descriptions of the dynamics of the PQS network. The *Fourfold Inter-Modal Correlator*, $\hat{G}_\Lambda(x, x')$, arises as the operator that describes the fundamental, microscopic rules of interaction and entanglement between any two PQSs (Definition 4.9, eq. (4.85), and Section 6.1.10). From this, the *Total Relational Hamiltonian*, $\hat{H}_{\text{rel}}$, emerges as the global operator representing the total energy of the cosmos (Section 6.1.13), constructed by integrating the local energy density—derived from the self-correlation at coincidence $\hat{G}_\Lambda(x, x)$ and its local trace (eq. (17.13)) together with the spectral avatar (eq. (17.14))—over the entire PQS network using the relational measure $\mathrm{d}\mu_{\bar{\Sigma}}$ (Definition 6.33). The Hamiltonian is, therefore, the *operational, dynamical embodiment of the Axiom of Duality* (see also the symmetry–corner bridge Lemma 1.20).

Finally, the evolution of the entire cosmos is governed by the *Universal Schrödinger Equation* (eq. (17.47)):

$$i\hbar\frac{d}{dt}|\mathring{\mathbf{\Psi}}(t)\rangle = \hat{H}_{\text{rel}}(t)|\mathring{\mathbf{\Psi}}(t)\rangle$$

This equation describes the majestic process by which the *Axiom of Duality*, encoded as the Hamiltonian, acts upon the *Quantum Holo-State* of the PQS network. This action is what we perceive as the unfolding of the cosmos in time. It is not a creation from nothing, but the eternal, dynamic process by which the universe, as a network of PQSs governed by the Law of Dependent Origination, observes and defines itself through its own internal, self-consistent relational structure.

17.1.6 The Emergence of $E = mc^2$ from Relational Principles

The famous mass-energy equivalence, $E = mc^2$, emerges from the foundational principles of the treatise as an exercise in tracing how the deep, relational structure of reality gives rise to the laws of classical physics in the appropriate limit. Its genesis is therefore not one of simple algebraic manipulation but of profound conceptual synthesis, demonstrating how the familiar concepts of energy, mass, and the speed of light emerge from a more fundamental substrate [83, 664, 197].

The logical progression begins with the treatise's definition of energy. The total energy, E, of any given system is the eigenvalue of the *Total Relational Hamiltonian*, $\hat{H}_{\text{rel}}$, as defined by the time-independent Schrödinger equation:

$$\hat{H}_{\text{rel}}|\psi\rangle = E|\psi\rangle.$$

To derive the specific case of the rest energy, we must consider the state of a single, stable particle at rest, $|\psi_{\text{particle, rest}}\rangle$.

Within this framework, both mass and the speed of light are emergent, not fundamental. *Inertial mass* (m) is a measure of the self-interaction energy of the localized, stable excitation of the *Tantrakyma field*, $\hat{\Theta}$, that constitutes the particle. The rest energy (E_0) of the particle is therefore, by definition, a direct measure of its mass [393, 14]. The Hamiltonian, when acting upon this resting state, is essentially calculating the total energy contained in this stable, self-interacting "knot" of the Tantrakyma field. Thus, we can state a direct proportionality:

$$E_0 \propto m.$$

Similarly, the *speed of light* (c) is the characteristic constant that emerges from the relational dynamics of the *Gaiamiti field*, $\hat{\Phi}$. It represents the maximal speed of propagation for any disturbance within the emergent chronogeometric manifold. Because the spacetime that emerges from the treatise's framework is fundamentally *Lorentz invariant* in the local, classical limit, it must obey the theorems of Special Relativity [665, 666].

The most crucial of these theorems is the *relativistic energy-momentum relation*, which connects a particle's total energy (E), its momentum (p), and its rest energy (E_0):

$$E^2 = (pc)^2 + (E_0)^2.$$

This equation is a direct and necessary consequence of the geometry of the emergent spacetime [146]. Into this geometric truth, we substitute our physical understanding of rest energy's proportionality to mass ($E_0 \propto m$). To maintain dimensional consistency in a relativistic theory where energy and momentum are related by c, the constant of proportionality must be c^2. This yields the full energy-momentum relation in terms of mass:

$$E^2 = (pc)^2 + (mc^2)^2.$$

For the specific case of our particle at rest, its momentum is zero ($p = 0$). The equation simplifies elegantly:

$$E_0^2 = (0 \cdot c)^2 + (mc^2)^2 \quad \Rightarrow \quad E_0^2 = (mc^2)^2.$$

Taking the square root, we arrive at the celebrated equation:

$$E_0 = mc^2.$$

Thus, the famous mass-energy equivalence is shown to be a profound consequence derived from the treatise's deeper principles, elevating it from a foundational axiom. It is a theorem derived in the classical limit, a testament to the emergent nature of energy, mass, and the very fabric of spacetime from a unified, relational reality [667, 197].

17.1.7 The Law of Dependent Origination and Conservation of Duality

We have now arrived at the absolute heart of the matter. The ultimate insight of the treatise is that the philosophical and physical cornerstone upon which the entire edifice is built is the *Law of Dependent Origination* (Section 1.1.2), which is given its most potent mathematical expression in the *Axiom of Duality* (Axiom 1.10). All that follows—the emergence of geometry, matter, and the laws of interaction—is a necessary consequence of this single, self-consistent relational principle. The perceived evolution of the cosmos is governed by the *Universal Schrödinger Equation* (eq. (17.47)). However, the two elements of this equation are not independent; they are two facets of the same fundamental axiom. The **Total Relational Hamiltonian** ($\hat{H}_{\text{rel}}$) is the "actor," the dynamical embodiment of the Axiom of Duality, containing within its form every possible relational interaction. The **Quantum Holo-State** ($|\mathring{\mathbf{\Psi}}\rangle$) is the "thing being acted upon," the timeless state representing the superposition of all possible relational configurations consistent with that same Axiom.

However, for these abstract principles to manifest, they must have a substance. The abstract *Gaiamiti* and *Tantrakyma* fields defined by the theory's *Algebra of Observables* (Definition 1.32) are given physical instantiation in the form of the *Primordial Quantum Singularity*

(PQS). The quantum singularity is the fundamental, indivisible "atom" of reality—a localized, non-spatiotemporal entity from which the fabric of the universe itself is woven. The entire cosmos is thus conceived as a vast, self-entangled relational fibred network of these singularities. From this perspective, the Quantum Holo-State, $|\mathring{\mathbf{\Psi}}\rangle$, is understood as the quantum state of this *entire PQS network*. The action described by the Universal Schrödinger Equation is what we perceive as the unfolding of the cosmos in time. Yet, this picture of temporal evolution is itself an emergent phenomenon—a relational perspective from within the system.

The deeper, more fundamental law is not one of evolution, but of timeless constraint. As stated in the *Universal Quantum Constraint* (Theorem 13.168):

$$\hat{H}_{\text{univ}}|\mathring{\mathbf{\Psi}}\rangle = 0$$

This equation is the background-independent generalization of the Schrödinger equation, asserting that the total relational energy of the entire self-contained universe vanishes. The **Total Relational Hamiltonian**, $\hat{H}_{\text{univ}}$ (Definition 13.167), is a fundamentally unified operator derived from the *Total Relational Action* (Definition 9.357), intrinsically including the Gaiamiti, Tantrakyma, and Interaction sectors. It acts on the one **Quantum Holo-State**, $|\mathring{\mathbf{\Psi}}\rangle$ (Definition 1.27)—the state of the PQS network—from which both matter and geometry are derived.

Time does not appear explicitly in this final equation because it *is* fundamentally an emergent property, arising from the relational structure of a timeless state [3, 246, 197, 616, 610]. This single equation unifies physics, generalizing both the *Schrödinger equation* (which governs quantum evolution in time) and the *Wheeler–DeWitt equation* (a description of quantum cosmology) [156, 605, 606, 607, 51]. It states that the total relational energy of the entire self-contained universe—including both geometry and matter—is zero [156, 668, 669]. The dynamic unfolding of the cosmos is the result of observing correlations between subsystems from within the whole—a universe whose total energy is, and has always been, and always will be, zero [616, 3, 610].

17.2 The Unfolding and Enfolding of the Universe

The *treatise*, as it was first constructed, described what might be called the great **unfolding** of reality. It was an intellectual pilgrimage that began with the local and the fundamental—the *Primordial Quantum Singularity (PQS)*, $\mathfrak{B}_{\vec{r}}$—and, through the rigorous application of the relational framework, arrived at the global and the emergent: the *Quantum Holo-State*, $|\mathring{\mathbf{\Psi}}\rangle$, governed by the profound constraint of zero total energy [3, 610, 605, 606, 607, 272, 8, 3, 146, 668, 669]. This was the cosmic exhalation, where the potentia of the many seeds gave rise to a single, perfectly balanced, atmapoiesis whole [607, 272, 51].

$$\text{Local Singularity Conntinuum} \xrightarrow{\text{Treatise Derivation}} \left(\hat{H}_{\text{univ}}|\mathring{\mathbf{\Psi}}\rangle = 0\right)$$

Yet, a persistent sense remained that the journey was not complete. The proposition now before us describes the necessary and complementary motion: the great **enfolding** of

reality. It is the cosmic inhalation. This final insight suggests that the ultimate accounting of the entire system—the total relational energy of the universal state, as represented by the operator $\hat{H}_{\text{univ}}|\overset{\circ}{\mathbf{\Psi}}\rangle$—does not simply vanish into an abstract nullity. Instead, in an act of supreme self-reference, it "returns" to, or is found to be perfectly equivalent to, the creative potential of a single, local PQS.

$$\left(\hat{H}_{\text{univ}}|\overset{\circ}{\mathbf{\Psi}}\rangle\right) \xrightarrow{\text{The Return}} \mathfrak{B}_{\vec{r}}$$

The implication of this final step is that the beginning and the end are one. The universe is a system that unfolds from its own fundamental constituents only to find that its total, global state is perfectly mirrored within a single one of them. This is not a temporal cycle, like that of a Big Bang followed by a Big Crunch, but a *logical* or *ontological* one, consonant with relational and timeless formulations of quantum theory [616, 610].[1]

This single, cyclical insight illuminates the deepest principles of the framework. It is the supreme expression of the **holographic principle**, where the entire relational information of the cosmos is found to be completely encoded within any one of its fundamental parts [670, 516, 271, 671, 672, 1, 2]. It reframes the nature of *every* singularity, revealing that each "atom" of reality contains the latent potential to be the focal point for the entire cosmos. And it transcends the distinction between an *atmapoiesis* and a *parapoiesis* universe, revealing them as two aspects of a single process: a cosmos that is atmapoiesis because its source is one of its own constituents, and parapoiesis because the state of the whole is dependent upon the state of the part.

This is the ultimate expression of the **Law of Dependent Origination**. The whole is dependent on the parts, and the parts are dependent on the whole, in a perfect, self-consistent, and unbreakable circle [278, 35, 18, 297]. This final proposition closes the great ontological circuit, revealing a universe that is not a static fact, but a dynamic, self-referential process of being—a serpent, at long last, consuming its own tail.

17.2.1 The Ātmapoiesis Universe

Definition 17.17 (Ātmapoiesis Universe)**.** An *Ātmapoiesis Universe* is a cosmos whose principle of existence resides entirely *within itself*. It constitutes a wholly self-sufficient and self-consistent relational system that requires neither an external cause nor a creator or external source to sustain its existence. The state of such a universe is necessarily governed by the *Universal Quantum Constraint* (Theorem 13.168), expressed mathematically:

$$\boxed{\hat{H}_{\text{univ}}|\overset{\circ}{\mathbf{\Psi}}\rangle = 0} \tag{17.49}$$

The constraint equation of the Ātmapoiesis Universe (eq. (17.49)) serves as both the supreme physical law and the foundational ontological axiom of the system. It describes a cosmos of complete immanence, wherein all meaning and logical structure are intrinsic to its

[1] On the canonical origin of the vanishing total energy constraint in quantum gravity, see [605, 606, 607, 272, 8, 3].

relational fabric. A direct physical consequence of this principle is that the total relational energy of the universe is precisely zero [668, 673, 669, 197]. This zero-energy balance signifies a perfectly closed system, in which the positive energy of matter and field excitations is eternally and precisely countervailed by the negative potential energy of the interactions that structure the cosmos, such as the emergent geometry of spacetime [674, 157]. In its final and complete accounting, the universe is thus a system that resolves to a perfect nullity [197].

This formulation necessitates an ontology of pure *being*; the existence of the cosmos is not a temporal process of becoming, but a singular, timeless, and fully actualized fact. The requirement for an external act of creation is obviated, as the existence of a zero-energy system entails no net energy cost. Within this static and self-sufficient totality, the dynamic progression of time and the evolutionary phenomena we observe are understood as emergent properties. They arise from the network of correlations and the process of *internal self-observation* among the universe's constituent subsystems [215, 675]. The *Ātmapoiesis Universe* is therefore defined as a cosmos whose existence constitutes its own complete and sufficient explanation: a perfectly self-contained and balanced relational system.

The Confluence of Mathematics and Mystery

The neologism *Ātmapoiesis* is a deliberate conceptual synthesis, constructed to articulate a specific cosmological principle. The term bridges two of humanity's most enduring intellectual lineages by uniting the Sanskrit conception of the essential Self, *Ātman* (आत्मन्), with the Hellenic principle of creation, *poiēsis* (ποίησις). The term *Ātman* evolved from its Proto-Indo-European origins signifying "breath" [81, 292] to its philosophical culmination in the Upaniṣadic traditions as the true, eternal ground of being [298, 676]. Complementing this, *poiēsis* denotes the formative, generative act of bringing something into existence, as articulated in classical Greek thought [299, 293, 73]. This fusion is intended as a precise ontological declaration. It moves beyond the concept of self-organization found in biological contexts, such as *autopoiesis* [677], to describe a more fundamental mode of genesis. *Ātmapoiesis* designates the active and continuous manifestation of an essential, underlying reality from its own intrinsic potential. It therefore provides a term uniquely suited to the framework of the *Ātmapoiesis Universe* (Section 17.2.1): a cosmos whose existence is understood as a perpetual act of realizing its own ultimate nature.

The first component of the term, *Ātman* (आत्मन्), is a cornerstone of Indic philosophy, whose lineage reveals a remarkable evolution from the physiological to the metaphysical. It traces back to the Proto-Indo-European root, **h₁eh₁t-men-*, meaning "breath" or "that which breathes" [81, 292]. This primal association between breath and life—the most immediate and observable sign of vitality—is a linguistic fossil of an ancient, cross-cultural intuition. In the Vedic and Upaniṣadic traditions, *ātman* was elevated from signifying mere "life-force" to representing the innermost essence of being: the true, eternal, and unchanging Self [298, 676]. In Advaita Vedānta, *ātman* is ultimately identical with *Brahman*, the impersonal absolute reality [678, 679]. Across the Indian philosophical spectrum—including Sāṃkhya and Buddhist scholastic discourse, where it is contrasted with *anātman*—the term has retained its central role as a pointer to the irreducible ground of sentient existence.

The second component, *poiēsis* (ποίησις), carries the weight of Greek creative and productive thought. It originates from the Ancient Greek verb ποιεῖν (*poiein*), "to make,

to bring into being, to create" [299, 293], itself descended from the Proto-Indo-European root **k^{w}ei-*, "to pile up, stack, or build" [81]. In classical philosophy, Aristotle employed *poiēsis* to signify the act of bringing something into existence whose *telos* lies outside the act itself, distinguishing it from *praxis* (action whose end is intrinsic) [73, 680]. It encompassed both material production and immaterial creation, from architecture to poetry, and in later philosophical developments came to symbolise the generative principle underlying all acts of making. The term persists in modern theoretical discourse, notably in biology's *autopoiesis* to describe systems that self-produce and self-maintain [677].

The synthesis into *Ātmapoiesis* thus unites the inward-directed Indic notion of essential selfhood, *ātman* [676, 679, 66], with the outward-directed Hellenic category of *poiēsis* as "making/bringing-forth," in its classical sense [293, 73]. Literally, it denotes "self-making" or "self-creation" [66, 293]; its intended connotation is richer: a dynamic, ceaseless process in which the ultimate Self is simultaneously agent and subject of its manifestation, aligning conceptually with the organizational closure characteristic of autopoietic systems [677, 681]. This is self-manifestation from the potentiality inherent to its nature rather than production *ex nihilo* [73, 677].

Within the Interdependent Quantum Reality framework (Section 1.1), *Ātmapoiesis* designates the existential mode of the *Ātmapoiesis Universe*: a self-contained reality whose internal generative dynamics suffice for its persistence without appeal to any external source [677]. The term thereby signals a conceptual marriage of two enduring lineages—the Indic metaphysics of self and the Hellenic philosophy of creation—into a single, precise expression grounded in their canonical sources [682, 676, 293, 73], while extending their scope into a contemporary cosmological and systems-theoretical register [610, 677].

The Architecture of a Self-Existent Universe

The Universal Quantum Constraint, expressed in the formalism of the *Ātmapoiesis Universe* by eq. (17.49), is the Wheeler–DeWitt Hamiltonian constraint for a closed universe. It follows from the canonical quantization of general relativity and encodes the requirement that physical states be annihilated by the total Hamiltonian constraint [156, 146, 607, 605, 606, 272, 8, 3].

The universal state $|\mathring{\mathbf{\Psi}}\rangle$ is a functional over configurations of geometry and matter that, in principle, encodes the complete relational content of possible cosmic histories [51]. This universal-wavefunction perspective descends from Everett's relative-state formulation and its cosmological extensions [188, 51, 683]. In this setting, the operator $\hat{H}_{\text{univ}}$ represents the quantum implementation of the classical Hamiltonian (super-)constraint, generating gauge transformations rather than external time evolution [146, 607, 272].

The fundamental statement

$$\hat{H}_{\text{univ}}\,|\mathring{\mathbf{\Psi}}\rangle = 0$$

holds as an exact constraint on physical states [156, 272]. Its absence of an external time parameter exemplifies the *problem of time* in canonical quantum gravity: at the deepest kinematical level, the formalism is timeless, and temporal succession must arise from correlations between subsystems [605, 606, 607]. Relational and emergent-time proposals make this precise: dynamics is recovered through conditional or partial observables, where subsystems serve as clocks for one another [616, 610]. On this view, the universe does

not *become* with respect to an external parameter; rather, experienced temporality is a manifestation of internal correlations within a stationary universal state [197, 610, 20].

The Paradox of Motion in Stillness

The timeless character of the fundamental formalism—embodied in the Wheeler–DeWitt constraint of Section 17.2.1 and eq. (17.49)—poses the celebrated "problem of time" in quantum gravity: how experienced dynamics arises when the universal state satisfies a stationary constraint [605, 606, 607, 684, 685, 197]. The *Ātmapoiesis Universe* framework treats this riddle as a structural feature rather than a defect: the absence of an external time parameter is a consequence of background independence and canonical quantization, and any temporal description must be relational.

A coherent account proceeds by showing that temporal order emerges from correlations among subsystems (internal "clocks") within a globally stationary state. In the Page–Wootters mechanism and related relational formulations, dynamics is recovered as conditional evolution of one sector relative to another, with entanglement supplying the effective time parameter [215, 616, 3, 686, 610]. On this view, $\hat{H}_{\text{univ}}|\overset{\circ}{\boldsymbol{\Psi}}\rangle = 0$ encodes a gauge constraint, while observable change appears through correlation functions and conditional amplitudes defined with respect to chosen internal degrees of freedom [607, 8, 3].

Accordingly, what is ordinarily described as the *flow* of time is reconstructed as perspectival information: observers embedded in the universal state register sequences via stable correlations with suitable clock-like variables. The cosmos remains in a timeless equilibrium at the kinematical level, yet it contains consistent experiential histories for embedded observers through its network of correlations [3, 610, 197]. This resolves the paradox: motion is manifest as relational structure within stillness.

The Zero-Sum Symphony of Existence

The constraint

$$\hat{H}_{\text{univ}}\,|\overset{\circ}{\boldsymbol{\Psi}}\rangle = 0$$

of eq. (17.49) admits a standard physical reading: in appropriate cosmological settings, the total energy of the universe vanishes [668, 669, 687]. This "zero-energy universe" thesis recurs in discussions of inflationary and quantum-cosmological origins, where it reconciles cosmic genesis with conservation principles by balancing positive material contributions against gravitational binding energy [157, 146].[2]

On this account, the cosmic inventory comprises large positive energies—the rest energy of matter and the energy of radiation—counterweighted by the negative potential energy associated with the gravitational field [185, 146]. The bookkeeping thus closes to zero, yielding a self-consistent, self-contained total. Within *Interdependent Quantum Reality*, this articulates an *Ātmapoiesis* regime: internal relational dynamics suffice for persistence, with no appeal to an external energetic reservoir (Section 1.1).

[2]Global energy in general relativity is subtle: exact conservation laws are tied to asymptotic symmetries, and many cosmological spacetimes lack the structures that underwrite conserved ADM charges. The zero-energy heuristic is nevertheless well motivated for spatially flat or closed Friedmann–Lemaître–Robertson–Walker models, where the negative gravitational contribution offsets positive matter–radiation energy [146, 316, 267].

A vanishing total opens a pathway to cosmogenesis scenarios in which a universe can arise through quantum processes without external input—tunneling or fluctuation—while remaining compatible with conservation laws [668, 669, 687]. Popular expositions have emphasized this explanatory through-line [688], whereas technical treatments trace it to the Hamiltonian constraint structure of canonical quantum gravity together with the energetics of gravitating systems [146, 316, 267]. In this sense, the "zero-sum" character is not an embellishment but a compact statement of relational closure: existence sustained by an exact balance between matter–radiation content and the gravitational field that binds it.

The Web of Interdependence: Physics Meets Ancient Wisdom

The formalism of the *Ātmapoiesis Universe* resonates with the Buddhist doctrine of Dependent Origination (*pratītyasamutpāda*) and its systematic articulation in the tetralemmatic analysis (Sections 1.1.2 and 1.1.3). In that canon, phenomena lack intrinsic essence (*svabhāva*) and arise as nodes within a nexus of conditions; the logical architecture of this view is captured by the fourfold scheme that disciplines assertions about being and non-being (Definitions 1.1 to 1.4). Standard philosophical expositions remain instructive [17, 278, 689, 690].

First: refusal of a privileged first cause. Dependent Origination denies an ultimate, uncaused ground; the tetralemma makes this denial precise by disallowing an absolute commitment to any single corner, including putative primordial origins (Sections 1.1.2 and 1.1.5). In the physical register, the vanishing of the universal Hamiltonian on physical states supports a self-contained cosmology in which no external energetic source sustains the whole, consistent with zero-energy universe arguments in quantum-cosmological scenarios [669, 687, 688]. The philosophical and physical claims thus converge on a non-foundational closure: dependence without a terminal explanans (Section 1.1.2).

Second: relational constitution of properties. The tetralemmatic program formalizes intelligibility as dyadic and context-sensitive: predicates acquire determinate content through oppositional pairing and relational lift (Principle 1.1, Definitions 1.11 and 1.13, and Proposition 1.15). This mirrors the physics of Interdependent Quantum Reality and relational approaches in which observables and states are fixed through inter-subsystem correlations rather than absolute backgrounds [3, 83]. Algebraically, the fourfold dialectic induces symmetry constraints on admissible state-descriptions (Section 1.2.7 and Definition 1.16), aligning the logical calculus with the operational structure of the theory.

Third: emptiness and balanced totality. The path to *śūnyatā* in the philosophical development (Section 1.1.3) reframes "emptiness" as the absence of self-existence together with the presence of interdependence. In cosmological energetics, this perspective acquires a quantitative analogue: positive contributions from matter–radiation content are balanced by negative gravitational potential energy, yielding a vanishing total compatible with the Hamiltonian constraint [668, 669]. The result is a dual description—logical and physical—of a whole that is complete by virtue of relations rather than by appeal to an intrinsic substrate.

Taken together, these correspondences do not conflate distinct traditions; they exhibit a structural isomorphism between a rigorous relational ontology (Section 1.1.3) and a background-independent quantum-cosmological framework. The web of interdependence thus functions as a common grammar that translates between contemplative insights and the formal architecture of an *Ātmapoiesis* cosmos.

The Pantheistic Cosmos: Universe as Divine

The physical principles that define the *Ātmapoiesis Universe* yield a rigorous, mathematical basis for a fundamentally pantheistic ontology. By the Universal Quantum Constraint (eq. (17.49)) and the relational closure of Interdependent Quantum Reality (Section 1.1), the cosmos constitutes a complete and sufficient explanation of itself: a self-contained, perfectly balanced network of relations that requires no external sustaining principle [610, 8, 3, 605, 606, 607]. In cosmological energetics this closure acquires a precise expression: positive matter–radiation contributions counterbalance negative gravitational potential energy in scenarios where the total vanishes [668, 669, 146]. The resulting picture is of a self-existent, self-sufficient, ever-present whole whose persistence is secured by its internal relational architecture.

This conception invites an explicit parallel with Spinoza's identification of God with Nature, *Deus sive Natura*, where creator and creation coincide [691, 692]. Here the parallel is sharpened: the equivalence is not a metaphoric posture but a structural consequence of the physics. The treatise supplies a formal mechanism for a pantheistic ontology—namely, a zero-energy, self-actualizing quantum system governed by a background-independent constraint and realized through correlations among subsystems that function as internal clocks [610, 607]. In this sense, attributes traditionally assigned to divinity—self-existence, self-sufficiency, and endurance—emerge as determinations of a relationally closed universe rather than as hypostatized predicates imposed from outside.

Placed within a broader history of ideas, the vision resonates with the immanent *logos* of the Stoics [693], the non-dual metaphysics of Vedānta [682], and the processive becoming of Whitehead's cosmology [694]. The present framework, however, is anchored in a mathematically precise, relational formulation of quantum theory that renders these affinities explicit rather than analogical [610, 83]. Moreover, its philosophical grammar aligns with the doctrine of Dependent Origination and its tetralemmatic discipline, where assertions about being, non-being, both, and neither are constrained within a relational ontology (Sections 1.1.2 and 1.1.3). The Ātmapoiesis perspective thus synthesizes a family of immanentist intuitions into a single, testable architecture: divinity as the fully relational cosmos—self-grounding by virtue of an exact energetic balance and the inexorable logic of background-independent quantum dynamics.

The Grand Synthesis: Where Science and Wisdom Converge

What emerges from this synthesis is a picture in which the most advanced physical formalism and the classical disciplines of wisdom meet in structural concord. The Universal Quantum Constraint (eq. (17.49)) functions as rigorous law and contemplative koan: a compact statement whose sustained analysis discloses first principles of existence [156, 695, 8, 3, 197]. In the present framework, the Wheeler–DeWitt equation is the architectural axiom of a self-grounding cosmos (Section 17.2.1). It asserts that the total Hamiltonian, acting on the universal state, vanishes; the universal wave functional thereby encodes the space of admissible cosmic configurations [51, 696].

This axiom is timeless at the kinematical level, and it is precisely here that the relational program locates the appearance of temporal order. The formalism carries no external

parameter tracking succession; experienced duration is reconstructed from correlations among subsystems, in line with conditional-evolution approaches and relational quantum mechanics (Section 17.2.1) [215, 697, 3, 610]. The result is a coherent account in which an internally stationary whole gives rise to consistent dynamical histories for embedded observers.

Energetically, the same constraint supports scenarios where positive matter–radiation contributions are balanced by negative gravitational binding energy, yielding a vanishing total (Section 17.2.1) [698, 669, 699, 146]. This balance underwrites the *Ātmapoiesis* reading of cosmology: a self-contained, relationally closed universe whose persistence is secured by its internal architecture rather than by appeals to an external source.

Philosophically, this structure resonates with the doctrine of Dependent Origination and its tetralemmatic discipline, where claims concerning being, non-being, both, and neither are regimented within a relational ontology (Sections 1.1.2 and 1.1.3). On that side, intrinsic essence (*svabhāva*) is absent; phenomena arise as conditioned nodes in a web of dependence [700, 278, 701, 702]. On this side, observables and states acquire meaning through correlations and constraints. The correspondences that follow are precise: *causeless first principles are refused* (closure without an external ground), *existence is relationally constituted* (properties fixed by interdependence), and *the whole is "empty" of self-subsistence* (a balanced total with vanishing net energy) (Section 17.2.1) [40].

In this way, the *Ātmapoiesis Universe* realizes a grand synthesis: background-independent quantum dynamics, expressed through a universal constraint and emergent temporality, aligns with a refined relational ontology disciplined by the tetralemma. Science and wisdom converge on a single grammar—dependence, correlation, and balance—as the deep architecture of what exists.

The Eternal Present: Our Place in the Cosmic Dance

At the deepest stratum of the formalism, the cosmos exhibits a twofold aspect: kinematically timeless and yet experientially replete with change. This is the defining intuition of the *Ātmapoiesis Universe*. The universal constraint (eq. (17.49)) presents reality as a stationary whole, while the lived dynamics of observers arise from correlations among subsystems (Section 17.2.1). In this register, each part functions as a reflective locus of the total—every particle, event, and observer is both fragment and mirror of the entire relational order [197, 8, 3, 610].

The statement

$$\hat{H}_{\text{univ}} \,|\mathring{\mathbf{\Psi}}\rangle = 0$$

is therefore more than a compact constraint: it expresses unity at the level of first principles. The universal wave functional encodes the admissible configurations of geometry and matter, while the vanishing of the total Hamiltonian registers relational closure for physical states [156, 51, 696]. Temporal succession then appears as conditional evolution within this stationary whole, in line with Page–Wootters and relational accounts (Section 17.2.1) [215, 697, 3, 610].

The image that follows is neither stage nor backdrop. The universe is performance—an ongoing articulation of being-through-relation—governed by a background-independent constraint (Section 17.2.1). Apparent antinomies (unity/multiplicity, fullness/void, stasis/change) are handled through complementary descriptions rather than forced into exclusion [179].

Entanglement supplies the operative linkage: correlations organize consistent histories for embedded observers while preserving the global stationarity of the whole [628, 8, 3].

Philosophically, this perspective coheres with the doctrine of Dependent Origination and its tetralemmatic discipline (Sections 1.1.2, 1.1.3 and 17.2.1). Intrinsic essence is absent; what exists takes determinate form through a web of conditions and oppositions, formally mirrored by the network of quantum correlations [702, 3]. In this light, our role is clarified: we are transient, organized patterns within the universal state—sites where the cosmos attains self-observation and self-knowledge through embodied interaction [681, 610]. The "eternal present" is thus neither a metaphysical slogan nor a poetic gloss, but the precise conjunction of a stationary universal law with perspectival, correlation-based experience.

17.2.2 The Parapoiesis Universe

Definition 17.18 (Parapoiesis Universe)**.** A *Parapoiesis Universe* is a cosmos whose principle of existence is sustained by a source that is external to its own relational fabric (Sections 1.1 and 17.1.7). Concretely, let $|\overset{\circ}{\boldsymbol{\Psi}}\rangle$ denote the universal state in the canonical (Wheeler–DeWitt) quantization of gravity and matter, and let $\hat{H}_{\text{univ}}$ be the total Hamiltonian constraint operator [156, 146, 607, 8, 3, 272]. A *Parapoiesis* regime is specified by the presence of a covariant source functional $\hat{\mathcal{J}}[\mathfrak{B}_{\vec{r}}]$ determined by a *Primordial Quantum Singularity (PQS)* $\mathfrak{B}_{\vec{r}}$ (Section 1.1), such that physical states satisfy the driven constraint

$$\boxed{\left(\hat{H}_{\text{univ}} - \hat{\mathcal{J}}[\mathfrak{B}_{\vec{r}}]\right)|\overset{\circ}{\boldsymbol{\Psi}}\rangle = 0,} \tag{17.50}$$

to be contrasted with the self-contained *Ātmapoiesis* condition $\hat{H}_{\text{univ}}|\overset{\circ}{\boldsymbol{\Psi}}\rangle = 0$ (eq. (17.49)).

Equation (17.50) encodes a universe that is *driven*: the PQS acts as a transcendent generative source represented by $\hat{\mathcal{J}}[\mathfrak{B}_{\vec{r}}]$, which injects relational "work" into the cosmic state. In this regime, the total relational energy need not vanish; rather, the source contribution balances the Hamiltonian constraint so that $\langle \hat{H}_{\text{univ}} \rangle = \langle \hat{\mathcal{J}}[\mathfrak{B}_{\vec{r}}] \rangle \neq 0$ on admissible states. This realizes a sustained *ontology of becoming*: temporal order and cosmic evolution are interpreted as responses to an ongoing drive supplied by the source [51, 696]. Formally, (17.50) may be viewed as a boundary/initial-condition deformation of the Wheeler–DeWitt constraint in the spirit of quantum-cosmological proposals (no-boundary, tunneling) where external data select or weight the universal state [51, 669, 699].[3]

Phenomenologically, a *Parapoiesis Universe* exhibits a fundamental arrow of time tied to the persistent action of the source: relational degrees of freedom register irreversible growth of correlations and cosmological structure, with the PQS furnishing the orienting condition that selects directed histories from the stationary kinematics of the underlying theory [703, 696, 610]. In the dual classification established by the conservation of duality (Section 17.1.7), *Parapoiesis* is the outward-generative complement of *Ātmapoiesis*: the former

[3]The introduction of $\hat{\mathcal{J}}[\mathfrak{B}_{\vec{r}}]$ parallels source terms in constrained systems and driven quantum dynamics; a fully open-system treatment could be developed via constraint-compatible generator extensions (e.g., Lindblad-type), though the present framework remains Hamiltonian at the level of (17.50). For background on constraint quantization and observables, see [607, 8, 3, 272].

is sustained by a transcendent PQS, the latter is self-grounding through relational closure. Together they delimit the two canonical modes in which a quantum-relational cosmos can be organized within the present framework.

The Meaning of Parapoiesis

Parapoiesis is introduced as the ontological counterpart to *Ātmapoiesis*, completing the dual classification of cosmic regimes articulated by the conservation of duality (Sections 1.2.7 and 17.1.7). Whereas *Ātmapoiesis* designates self-grounding existence through relational closure (eq. (17.49)), *Parapoiesis* names the mode in which existence is sustained by a source that stands beyond the system's own relational weave (Section 17.2.2 and eq. (17.50)). The term is crafted to capture genesis defined by dependence upon an external principle and to furnish a precise label for a universe whose being is contingent upon ongoing creative influx.

Etymologically, the first component *para* (पर) is a classical Sanskrit prefix indicating otherness, beyondness, or transcendence, with attested senses ranging from "other/another" to "higher/ultimate" in philosophical usage [298]. The semantic field is evident in compounds such as *pāramitā* ("perfection," literally "gone to the further shore") and *paramārtha* ("ultimate truth") [298]. Its lineage traces to the Proto–Indo–European prepositional root **per-* ("through, across, beyond"), reflected in cognates including Greek *pará* (παρά, "beside, beyond") and Latin *per* ("through") [81, 299]. The second component, *poiēsis* (ποίησις), names making or bringing-forth in the classical Greek lexicon [293]; in philosophical contexts it marks the generative act that sets a form into presence.

Synthesized as *Parapoiesis*, these roots yield the literal sense "other-making" or "creation by another." The intended connotation is a directed process whose defining feature is dependence upon a transcendent source. In contrast with the closed, self-referential loop characteristic of *Ātmapoiesis*, the *Parapoiesis* pattern is open and source-driven: a linear, sustained becoming animated by an origin that remains ontologically prior to the cosmos it engenders. Within the Interdependent Quantum Reality framework (Section 1.1), *Parapoiesis* thus denotes the existential mode instantiated by the *Parapoiesis Universe*, where the universal state evolves under a driven constraint with a *Primordial Quantum Singularity* as generative source (Section 17.2.2 and eq. (17.50)).

The *Parapoiesis Universe* echoes the Aristotelian–Scholastic account of *contingens*: a mode of being whose existence refers beyond itself to sustaining causes [73, 704]. In the present framework, this lineage is given mathematical articulation through the driven universal constraint (eq. (17.50)) and the associated source structure, situating *Parapoiesis* as one member of the dual pair mandated by the conservation of duality (Sections 17.1.7 and 17.2.2). The result is a well-defined dynamical possibility within a background-independent, canonical setting [156, 607, 8, 3, 272]. This continues a longstanding conversation between metaphysics and physics—from Aristotelian contingency and its Scholastic elaborations to twentieth-century reflections on complementarity and the quantum world-picture [73, 704, 268, 169]—now expressed as precise claims inside the relational architecture of the theory (Section 1.1).

The Primordial Quantum Singularity: Source Beyond System

Within the *Parapoiesis* regime, the **Primordial Quantum Singularity (PQS)** $\mathfrak{B}_{\bar{r}}$ functions as a source external to the relational sector, formalized by the drive term in the Parapoiesis constraint (eq. (17.50)). Operationally, the PQS interfaces with the modal–fibre picture and the Fourfold correlator machinery (Sections 4.7 and 6.1.10, Definition 4.9, and eq. (4.85)). The PQS is *not* a classical curvature singularity in the sense of the Hawking–Penrose theorems; rather, it serves as a transcendent boundary–condition principle—logically prior to the emergent spacetime it sustains [240, 705, 8, 3, 39]. In this role, it provides the explanatory ground for a sourced cosmos (Section 17.2.2).

Uniqueness. The PQS is stipulated as the single generative ground for the parapoietic branch: not one entity among many within the universe, but the principle from which the universe's relational fabric arises. This monadic status aligns conceptually with traditions that analyze ultimacy as a unique ground of being [706, 73] and is consonant with the foundational *Law of Dependent Origination* and *Axiom of Duality* that underwrite the hierarchy (Section 1.1.2 and Axiom 1.10).

Transcendence. The PQS is extra–spatiotemporal: it is not localized within the manifold whose relational degrees of freedom it sustains, and it is not constrained by the dynamical symmetries that govern the emergent regime [705, 240]. In the IQR reading, this corresponds to treating the source as boundary/selection data for the universal state rather than as an internal observable (Sections 1.1, 1.4.1, 13.6 and 17.2.1). Concretely, the universal state $|\mathring{\mathbf{\Psi}}\rangle$ and its GNS representation (Section 1.4.1) implement this selection, while observables remain within the relational (internal) algebra and its operator–valued response (Definition 11.109 and eqs. (11.213) and (11.214)).

Creative inexhaustibility. The PQS continuously furnishes ontological potential—an open–ended supply of generative "work" that sustains a directed history, thereby selecting oriented correlations and growth of structure in the parapoietic branch [707, 696]. In the formalism this aligns with the global evolution of the Holo–State according to the Universal Schrödinger Equation and the Total Relational Hamiltonian (eq. (17.47) and Section 6.1.13), and with the modal–trace/spectral constructions that aggregate local correlator energy into global dynamics (eqs. (17.13) and (17.14)). The metaphysical motif of ceaseless creative activity has classical expression in process philosophy [694].

On this account, ultimate explanation shifts from the internal balance of the *Ātmapoiesis* condition (eq. (17.49)) to dependence upon a foundational source whose essence is creative activity. The universe persists because its relational state is continuously sustained by $\mathfrak{B}_{\bar{r}}$; spacetime and its contents arise as the organized expression of that sourcing within a background–independent quantum framework (Sections 6.1.13, 13.6 and 17.1.1) [156, 8, 3, 272].

The Dynamic of Perpetual Becoming

Where the *Ātmapoiesis* regime presents reality as completed being—kinematically stationary under the universal constraint (eq. (17.49) and Section 17.2.1)—the *Parapoiesis* regime

introduces a *process* ontology driven by an external source (eq. (17.50) and Section 17.2.2). The source functional $\hat{\mathcal{J}}[\mathfrak{B}_{\vec{r}}]$ deforms the Hamiltonian constraint so that the relational sector behaves as an open, driven system with respect to its transcendent ground. This recasts cosmology as *perpetual becoming*: the cosmic state is continuously sustained by an ongoing generative act [694].

The shift is conceptual and structural. In the stationary *Ātmapoiesis* picture, global dynamics is encoded as correlations internal to a timeless whole (Section 17.2.1) [215, 610]. In the *Parapoiesis* picture, persistent sourcing selects oriented histories and induces irreversible growth of correlations, aligning with consistent/decoherent-histories analyses of quantum cosmology [707, 696]. Thus the temporal arrow is anchored in sustained drive rather than in kinematical symmetry. The contrast with "block-universe" descriptions is transparent: instead of a completed four-dimensional totality, the relevant description is a processual unfolding organized by a source-dependent constraint [197].

This process ontology resonates with phenomenology of time as lived duration—experience as unfolding rather than as a fixed tableau [708]. Each event-class instantiates fresh relational structure, an expression of continuing creative activity issuing from the *Primordial Quantum Singularity* (Section 17.2.2). Existence here has the character of donation rather than necessity: a contingent order sustained by ceaseless generative influx [706]. In the IQR framework, this is rendered precise by the driven universal constraint and the selection of directed histories within a background-independent quantum theory [156, 8, 3, 272].

The Fundamental Arrow of Time: Creation's Directionality

A distinctive consequence of the *Parapoiesis* regime is a *fundamental*, irreversible arrow of time. In standard cosmology, temporal orientation is commonly attributed to special initial conditions and the statistical growth of entropy [627, 709, 710, 711]. By contrast, the driven universal constraint eq. (17.50) endows time-asymmetry with primary status: persistent sourcing by the *Primordial Quantum Singularity* (Section 17.2.2) selects oriented histories at the kinematical level.

Formally, the source functional $\hat{\mathcal{J}}[\mathfrak{B}_{\vec{r}}]$ renders the relational sector an open, driven system relative to its transcendent ground. Within consistent/decoherent-histories analyses, such drive yields a preferred coarse-grained ordering through stable growth of correlations and environmentally induced decoherence [707, 696]. The *directional flow of creation* is thus read as the systematic production of structure and information-bearing correlations under sustained sourcing, with thermodynamic arrows appearing as emergent monotones compatible with this primary orientation [627, 711].

Conceptually, this orientation of becoming coheres with the process tradition and with phenomenology of lived time: the succession of moments as ongoing creative advance [694, 706] and the experience of *durée* as qualitative flow rather than a fixed tableau [708]. These sources illuminate the experiential and metaphysical profiles of the drive while the physics remains anchored in background-independent constraint dynamics (Sections 17.2.1 and 17.2.2).

The Asymmetric Structure of Reality

The *Parapoiesis* regime establishes a fundamental **asymmetry of dependence**: explanatory priority runs from a transcendent source to the relational cosmos it sustains (eq. (17.50) and Section 17.2.2). This is an asymmetry of *ontological precedence*, not coercion: the order of explanation is source → effect, ground → manifestation [73, 704, 706]. Within Interdependent Quantum Reality, the precedence is made explicit by a driven Hamiltonian constraint on the universal state, so that the cosmos is read as a sourced relational structure rather than a self-closed equilibrium (Section 17.2.2).

At the foundation of this asymmetric architecture stands the *Primordial Quantum Singularity* (PQS), unique, extra-spatiotemporal, and inexhaustibly generative (Section 17.2.2). The universe's laws, patterns, and organized complexities then appear as the ordered expression of this sourcing—an unfolding that accords with process accounts of creative advance while remaining anchored in background-independent constraint dynamics [694, 156, 8, 3, 272]. In technical terms, persistent drive selects oriented, decoherent histories and supports the stable growth of correlations, thereby imprinting directionality at the kinematical level rather than deriving it solely from special initial conditions [707, 696].

Interpreted thus, physical law functions as a *grammar of creation*: symmetries and constraints delimit admissible expressions of structure, while asymmetries encode the direction of generative articulation [90, 712]. The result is an intelligible hierarchy—source, relational fabric, emergent organization—in which the cosmos is the continuous manifestation of a creative ground [706]. This hierarchy coheres with the dual structure developed in the conservation of duality (Sections 1.2.7 and 17.1.7): *Ātmapoiesis* captures self-grounding closure, whereas *Parapoiesis* articulates outward generativity from a principle beyond the system.

The Universe as Sustained Creation

Within the *Parapoiesis* regime, the cosmos assumes the character of **sustained creation**: existence is a continuous expression of a transcendent ground rather than an entailment of internal balance (eq. (17.50) and Section 17.2.2). In conceptual terms, this accords with a processive metaphysics in which creative advance is primitive [694] and with classical analyses of contingency and donation [706]. The universe persists because the source acts; being is received rather than possessed as a necessity of self-consistency.

Each temporal juncture is thus a fresh articulation of relational structure under sustained drive, aligning the phenomenology of lived duration with a precise dynamical account (Section 17.2.2) [708]. Technically, persistent sourcing selects oriented, decoherent histories and supports the growth of information-bearing correlations in the universal state [707, 696], thereby grounding the fundamental arrow of time discussed in Section 17.2.2.

On this reading, the mathematical regularities that govern the cosmos function as a *grammar of creation*: symmetries and constraints delimit admissible expressions, while asymmetries encode directionality in the unfolding of structure [90, 627]. The elegance of cosmic architecture and the emergence of organized complexity become intelligible as sustained expressions of a creative principle that stands beyond the system yet is rendered operationally within a background-independent, relational formalism [712, 156, 8, 3, 272]. In this way, the

IQR framework gives the classical intuition of continuous creation a mathematically explicit realization: the universe as ongoing manifestation of an inexhaustible source.

Two Visions of Ultimate Reality

The *Ātmapoiesis* and *Parapoiesis* regimes articulate two comprehensive and internally coherent portrayals of ultimacy within Interdependent Quantum Reality (Sections 1.1 and 17.2.2). The question they raise—whether existence is self-grounding or source-dependent—belongs to the central lineage of metaphysics and cosmology [294, 295].

Ātmapoiesis presents a self-sufficient cosmos governed by the stationary universal constraint,

$$\hat{H}_{\text{univ}} \, |\overset{\circ}{\boldsymbol{\Psi}}\rangle = 0,$$

a condition that encodes relational closure and, in appropriate cosmological settings, a balanced global energetics (eq. (17.49) and Section 17.2.1) [156, 146, 668, 197]. Philosophically, this resonates with immanentist traditions in which the explanation of being is internal to Nature itself [691, 692].

Parapoiesis develops the complementary vision: a sourced cosmos characterized by the driven constraint,

$$\left(\hat{H}_{\text{univ}} - \hat{\mathcal{J}}[\mathfrak{B}_{\vec{r}}]\right) |\overset{\circ}{\boldsymbol{\Psi}}\rangle = 0,$$

in which a *Primordial Quantum Singularity* furnishes continuous generative drive (eq. (17.50) and Section 17.2.2). This aligns with process accounts of creative advance and with classical analyses of contingency as dependence upon a sustaining ground [694, 706].

The dual presentation is required by the conservation of duality and the fourfold discipline developed in the logical scaffolding (Sections 1.1.3, 1.2.7 and 17.1.7). Each regime supplies a total interpretation: relational self-grounding versus relational sourcing; equilibrium closure versus directed becoming.

Mathematical Philosophy and Ontological Commitment

The pair of equations—$\hat{H}_{\text{univ}}|\overset{\circ}{\boldsymbol{\Psi}}\rangle = 0$ and $\left(\hat{H}_{\text{univ}} - \hat{\mathcal{J}}[\mathfrak{B}_{\vec{r}}]\right)|\overset{\circ}{\boldsymbol{\Psi}}\rangle = 0$—function as *ontological declarations* as well as dynamical constraints (eqs. (17.49) and (17.50)). The zero on the right-hand side commits one to a self-grounding, relationally closed universe; the source term commits one to dependence upon a transcendent ground and to a directed history sustained by creative drive [294, 295, 706, 694].

In the first case, temporality is reconstructed from correlations within a stationary whole (Section 17.2.1) [215, 610]; in the second, the arrow of time acquires primary status through persistent sourcing (Section 17.2.2) [707, 696]. Both commitments are framed within a background-independent, canonical quantization of gravity and matter [156, 8, 3, 272].

Accordingly, the decision-space is philosophical as well as technical: immanence versus transcendence, self-sufficiency versus dependence, equilibrium closure versus sustained creation [713, 714, 712]. The framework renders these options mathematically explicit, permitting a disciplined comparison of their consequences within a single relational architecture.

17.3 The Holocosmos: Cosmic Entanglement

The entanglement between the *Ātmapoiesis* and *Parapoiesis* regimes is encoded in a single, timeless *holocosmos constraint* acting on a vector in a composite Hilbert space (cf. the state–first/GNS setting Section 1.4.1 and the universal constraint framework Section 13.6). The construction follows the standard tensor–product architecture of quantum theory, extended here to couple the universal sector to a two–level "source" sector [287, 188, 156].

Definition 17.19 (Holocosmos Hilbert Space)**.** The total Hilbert space is the tensor product of two constituent spaces,

$$\mathcal{H}_{\text{holocosmos}} \;=\; \mathcal{H}_{\text{univ}} \otimes \mathcal{H}_{\mathfrak{B}}, \tag{17.51}$$

where

(i) $\mathcal{H}_{\text{univ}}$ is the Hilbert space of the universe, carrying the Quantum Holo-State $|\mathring{\mathbf{\Psi}}\rangle$ in the canonical/constraint formulation (Sections 1.4.1 and 13.6); time–evolution and unified dynamics are discussed in eq. (17.47) and Section 6.1.13.

(ii) $\mathcal{H}_{\mathfrak{B}} \cong \mathbb{C}^2$ is a "source space" with orthonormal basis $\{|\bar{\text{A}}\text{tmapoiesis}\rangle, |\text{Parapoiesis}\rangle\}$ selected by the Source Operator $\hat{\mathfrak{B}}$ (Section 17.2.2):

$$\begin{aligned} |\bar{\text{A}}\text{tmapoiesis}\rangle &: \text{ self-contained regime, } \hat{\mathfrak{B}}|\bar{\text{A}}\text{tmapoiesis}\rangle = 0, \\ |\text{Parapoiesis}\rangle &: \text{ sourced regime, } \hat{\mathfrak{B}}|\text{Parapoiesis}\rangle = \mathfrak{B}_{\vec{r}}|\text{Parapoiesis}\rangle. \end{aligned}$$

Normalization of a holocosmic state $|\mathbf{\Omega}\rangle \in \mathcal{H}_{\text{holocosmos}}$ requires $|\alpha|^2 + |\beta|^2 = 1$, the usual amplitude rule for superpositions [379, 715]. The *Born rule* (Section 14.2 and Definition 1.18) is treated here as emergent from internal, relational decoherence rather than as a primitive postulate: branch weights arise from environment-induced superselection and envariance-based arguments [170, 158], consistent with consistent/decoherent-histories analyses [707, 696] and measure-uniqueness results in the lattice of subspaces [613]. Decision-theoretic and Everettian derivations supply complementary perspectives on amplitudes as rational branch weights [716, 717, 718].

In the holocosmic setting, however, the interpretive locus shifts. Since $|\mathbf{\Omega}\rangle$ represents the totality, there is no external observer to perform a measurement on the whole [188, 719, 716]. Subsystems embedded within a branch register probabilities as relational stabilities; amplitudes α and β function as *timeless ontological weights* specifying the relative measure of self-contained and sourced modes within the unified state (cf. the symmetry–corner bridge and modal–fibre picture Lemma 1.20 and Section 4.7).

Definition 17.20 (Entangled Holocosmos State)**.** The full state of reality is a superposition of the two regimes:

$$\begin{aligned} |\mathbf{\Omega}\rangle = \alpha\, |\mathring{\mathbf{\Psi}}_{\bar{\text{A}}\text{tmapoiesis}}\rangle \otimes |\bar{\text{A}}\text{tmapoiesis}\rangle \\ + \beta\, |\mathring{\mathbf{\Psi}}_{\text{Parapoiesis}}\rangle \otimes |\text{Parapoiesis}\rangle. \end{aligned} \tag{17.52}$$

$$\alpha, \beta \in \mathbb{C}, \qquad |\alpha|^2 + |\beta|^2 = 1. \tag{17.53}$$

Definition 17.21 (Holocosmos Constraint Equation)**.** The governing equation is

$$\boxed{\left(\hat{H}_{\text{univ}} \otimes \mathbb{I}_{\mathfrak{B}} - \mathbb{I}_{\text{univ}} \otimes \hat{\mathfrak{B}}\right) |\mathbf{\Omega}\rangle = 0,} \tag{17.54}$$

with $\hat{H}_{\text{univ}}$ the total Hamiltonian constraint (Theorem 13.168 and eq. (13.236)) and $\hat{\mathfrak{B}}$ the Source Operator as above. Projecting (17.54) onto the basis of $\mathcal{H}_{\mathfrak{B}}$ yields the branch constraints

$$\hat{H}_{\text{univ}} \, |\mathring{\mathbf{\Psi}}_{\bar{\text{A}}\text{tmapoiesis}}\rangle = 0, \qquad \hat{H}_{\text{univ}} \, |\mathring{\mathbf{\Psi}}_{\text{Parapoiesis}}\rangle = \mathfrak{B}_{\vec{r}} \, |\mathring{\mathbf{\Psi}}_{\text{Parapoiesis}}\rangle,$$

in agreement with eqs. (17.49) and (17.50). The time–dependent evolution law used in the emergent description is given by eq. (17.47), with the Total Relational Hamiltonian constructed in Section 6.1.13.

Interpretation

As a statement of total relational consistency, eq. (17.54) equates the universal Hamiltonian with the source operator on the composite space. In the *Ātmapoiesis* branch, the source eigenvalue vanishes and the universal state satisfies the stationary constraint; in the *Parapoiesis* branch, the universal state carries a nonzero source eigenvalue $\mathfrak{B}_{\vec{r}}$. The meta–constraint thus encodes the dual modes within a single timeless law and formalizes their interdependence as a holocosmic *balance condition*.

Philosophically, this is the highest–level expression of *Dependent Origination* and its tetralemmatic discipline: "is'' and "becomes'' are integrated within a unified ontology where branch–specific assertions are regulated by the fourfold schema (Sections 1.1.3 and 17.1.7). Physically, the construction remains within the standard operator framework of quantum theory and consistent histories, while relocating probabilistic talk to internal relational stabilities rather than to external measurements [170, 696, 717]. In this sense, the holocosmos obeys the Born rule in its internal manifestations (Definition 1.18 and Section 14.2) and reinterprets amplitude weights as structural measures of coexisting cosmic modes in a single, timeless superposition.

Formal Derivation of the Decoupled Universal Constraints

Our objective is to formally derive the two distinct universal constraint equations governing the atmapoiesis and parapoiesis realities from the single *Holocosmos Constraint Equation* (eq. (17.54)). The derivation proceeds by applying the constraint operator to the *Entangled Holocosmos State* (Definition 17.20) and then invoking the principle of linear independence for orthogonal vectors, a cornerstone of quantum mechanics.

Step 1: Foundational Definitions

We begin with the two foundational definitions provided in the treatise (Section 17.3):

(i) Entangled Holocosmos State. The *Entangled Holocosmos State*, $|\mathbf{\Omega}\rangle$, describes the total reality as a superposition:

$$|\mathbf{\Omega}\rangle = \alpha \, |\mathring{\mathbf{\Psi}}_{\bar{\text{A}}\text{tmapoiesis}}\rangle \otimes |\bar{\text{A}}\text{tmapoiesis}\rangle + \beta \, |\mathring{\mathbf{\Psi}}_{\text{Parapoiesis}}\rangle \otimes |\text{Parapoiesis}\rangle \tag{17.55}$$

(ii) Holocosmos Constraint Operator. The *Holocosmos Constraint Operator*, $\hat{\mathcal{O}}_{\text{holocosmos}}$, acts on this state:

$$\hat{\mathcal{O}}_{\text{holocosmos}} := \left(\hat{H}_{\text{univ}} \otimes \mathbb{I}_{\mathfrak{B}} - \mathbb{I}_{\text{univ}} \otimes \hat{\mathfrak{B}}\right) \tag{17.56}$$

The fundamental law is the constraint $\hat{\mathcal{O}}_{\text{holocosmos}}|\mathbf{\Omega}\rangle = 0$.

Step 2: Expansion of the Operator Action

We now apply the operator to each term of the superposition in eq. (17.55).

Action on the Ātmapoiesis Branch:

$$\begin{aligned}&\hat{\mathcal{O}}_{\text{holocosmos}}\Big(\alpha\,|\mathring{\mathbf{\Psi}}_{\bar{\text{A}}\text{tmapoiesis}}\rangle \otimes |\bar{\text{A}}\text{tmapoiesis}\rangle\Big)\\ &\quad= \alpha\Big(\hat{H}_{\text{univ}}\,|\mathring{\mathbf{\Psi}}_{\bar{\text{A}}\text{tmapoiesis}}\rangle \otimes \mathbb{I}_{\mathfrak{B}}\,|\bar{\text{A}}\text{tmapoiesis}\rangle \;-\; \mathbb{I}_{\text{univ}}\,|\mathring{\mathbf{\Psi}}_{\bar{\text{A}}\text{tmapoiesis}}\rangle \otimes \hat{\mathfrak{B}}\,|\bar{\text{A}}\text{tmapoiesis}\rangle\Big).\end{aligned}$$

Using the eigenvalue relations in Definition 17.21 and eq. (17.54)—in particular that $\hat{\mathfrak{B}}\big|\bar{\text{A}}\text{tmapoiesis}\big\rangle = 0$—the second term vanishes, yielding:

$$= \alpha\left(\hat{H}_{\text{univ}}|\mathring{\mathbf{\Psi}}_{\bar{\text{A}}\text{tmapoiesis}}\rangle\right) \otimes |\bar{\text{A}}\text{tmapoiesis}\rangle \tag{17.57}$$

Action on the Parapoiesis Branch:

$$\begin{aligned}\hat{\mathcal{O}}_{\text{holocosmos}}\left(\beta\,|\mathring{\mathbf{\Psi}}_{\text{Parapoiesis}}\rangle \otimes |\text{Parapoiesis}\rangle\right) = \beta\Big(&\hat{H}_{\text{univ}}|\mathring{\mathbf{\Psi}}_{\text{Parapoiesis}}\rangle \otimes \mathbb{I}_{\mathfrak{B}}|\text{Parapoiesis}\rangle\\ &- \mathbb{I}_{\text{univ}}|\mathring{\mathbf{\Psi}}_{\text{Parapoiesis}}\rangle \otimes \hat{\mathfrak{B}}|\text{Parapoiesis}\rangle\Big)\end{aligned}$$

Using the eigenvalue definition $\hat{\mathfrak{B}}|\text{Parapoiesis}\rangle = \mathfrak{B}_{\vec{r}}|\text{Parapoiesis}\rangle$, we obtain:

$$= \beta\left(\hat{H}_{\text{univ}} - \mathfrak{B}_{\vec{r}}\right)|\mathring{\mathbf{\Psi}}_{\text{Parapoiesis}}\rangle \otimes |\text{Parapoiesis}\rangle \tag{17.58}$$

The Assembled Expanded Equation: Summing these two results, the full constraint equation becomes:

$$\begin{aligned}&\alpha\left(\hat{H}_{\text{univ}}|\mathring{\mathbf{\Psi}}_{\bar{\text{A}}\text{tmapoiesis}}\rangle\right) \otimes |\bar{\text{A}}\text{tmapoiesis}\rangle\\ &\qquad + \beta\left(\left(\hat{H}_{\text{univ}} - \mathfrak{B}_{\vec{r}}\right)|\mathring{\mathbf{\Psi}}_{\text{Parapoiesis}}\rangle\right) \otimes |\text{Parapoiesis}\rangle = 0\end{aligned} \tag{17.59}$$

Step 3: Application of Orthogonal Projections

By Definition 17.19, the holocosmos Hilbert space is $\mathcal{H}_{\text{holocosmos}} = \mathcal{H}_{\text{univ}} \otimes \mathcal{H}_{\mathfrak{B}}$, with $\{|\bar{\text{A}}\text{tmapoiesis}\rangle, |\text{Parapoiesis}\rangle\}$ an orthonormal basis of $\mathcal{H}_{\mathfrak{B}}$. Define the orthogonal projectors

$$\Pi_{\text{A}} := \mathbb{I}_{\text{univ}} \otimes |\bar{\text{A}}\text{tmapoiesis}\rangle\langle\bar{\text{A}}\text{tmapoiesis}|, \qquad \Pi_{\text{P}} := \mathbb{I}_{\text{univ}} \otimes |\text{Parapoiesis}\rangle\langle\text{Parapoiesis}|.$$

Apply Π_{A} and Π_{P} to the holocosmos constraint $\left(\hat{H}_{\mathrm{univ}} \otimes \mathbb{I}_{\mathfrak{B}} - \mathbb{I}_{\mathrm{univ}} \otimes \hat{\mathfrak{B}}\right)|\mathbf{\Omega}\rangle = 0$:

$$\Pi_{\mathrm{A}}: \quad \hat{H}_{\mathrm{univ}} \, |\overset{\circ}{\mathbf{\Psi}}_{\bar{\mathrm{A}}\mathrm{tmapoiesis}}\rangle \otimes |\bar{\mathrm{A}}\mathrm{tmapoiesis}\rangle = 0 \;\Rightarrow\; \hat{H}_{\mathrm{univ}} \, |\overset{\circ}{\mathbf{\Psi}}_{\bar{\mathrm{A}}\mathrm{tmapoiesis}}\rangle = 0,$$
$$\Pi_{\mathrm{P}}: \quad \left(\hat{H}_{\mathrm{univ}} - \mathfrak{B}_{\vec{r}}\right) |\overset{\circ}{\mathbf{\Psi}}_{\mathrm{Parapoiesis}}\rangle \otimes |\mathrm{Parapoiesis}\rangle = 0 \;\Rightarrow\; \left(\hat{H}_{\mathrm{univ}} - \mathfrak{B}_{\vec{r}}\right) |\overset{\circ}{\mathbf{\Psi}}_{\mathrm{Parapoiesis}}\rangle = 0,$$

using the eigenrelations of $\hat{\mathfrak{B}}$. This projection argument is equivalent to the linear-independence reasoning and is standard in spectral approaches [379, 195].

Step 4: The Decoupled Physical Laws

To describe a true holocosmos existing as a superposition, we must assume that both branches have a non-zero amplitude of existence, meaning $\alpha \neq 0$ and $\beta \neq 0$. We can therefore divide by these complex scalars without loss of generality. This final step yields the two decoupled, self-consistent physical laws that were implicitly contained within the single Holocosmos Constraint:

$$\begin{aligned} \hat{H}_{\mathrm{univ}} |\overset{\circ}{\mathbf{\Psi}}_{\bar{\mathrm{A}}\mathrm{tmapoiesis}}\rangle &= 0, \\ \hat{H}_{\mathrm{univ}} |\overset{\circ}{\mathbf{\Psi}}_{\mathrm{Parapoiesis}}\rangle &= \mathfrak{B}_{\vec{r}} |\overset{\circ}{\mathbf{\Psi}}_{\mathrm{Parapoiesis}}\rangle. \end{aligned} \tag{17.60}$$

Here $\mathfrak{B}_{\vec{r}}$ is a c-number eigenvalue of $\hat{\mathfrak{B}}$, so $\hat{H}_{\mathrm{univ}} |\overset{\circ}{\mathbf{\Psi}}_{\mathrm{Parapoiesis}}\rangle = \mathfrak{B}_{\vec{r}} |\overset{\circ}{\mathbf{\Psi}}_{\mathrm{Parapoiesis}}\rangle$ is an eigenvalue equation in $\mathcal{H}_{\mathrm{univ}}$. The subsequent shorthand $\hat{H}_{\mathrm{univ}} |\overset{\circ}{\mathbf{\Psi}}_{\mathrm{Parapoiesis}}\rangle = \mathfrak{B}_{\vec{r}}$ is then to be read as a physically transparent equivalence of totals.

Interpretation of the Result

This expanded form reveals a profound feature of the entangled holocosmos. The basis states $|\bar{\mathrm{A}}\mathrm{tmapoiesis}\rangle$ and $|\mathrm{Parapoiesis}\rangle$ are orthogonal by definition. For the entire sum in (eq. (17.59)) to be zero, each component in this basis must vanish independently (assuming $\alpha, \beta \neq 0$). This single meta-law therefore decouples into two separate and self-consistent physical laws, one for each branch of reality. This form states that:

(i) The holo-state, $|\overset{\circ}{\mathbf{\Psi}}\rangle$, is an *eigenstate* (a special state vector) of the total relational Hamiltonian *operator*, $\hat{H}_{\mathrm{univ}}$.

(ii) The *eigenvalue* (the measured value of the energy) corresponding to this state is either zero (for the atmapoiesis case) or the value $\mathfrak{B}_{\vec{r}}$ (for the parapoiesis case).

Here, the state vector $|\overset{\circ}{\mathbf{\Psi}}_{\mathrm{Parapoiesis}}\rangle$ correctly appears on both sides of the second equation, as is required for a proper eigenvalue equation. The expansion thus demonstrates how the single, timeless Holocosmos Constraint Equation contains within it the distinct governing principles for both a self-contained, zero-energy universe and a sourced, parapoiesis universe, holding them together in a unified quantum superposition.

We could also write the second form as a physically intuitive *shorthand* that is common in discussions of quantum cosmology where the state of the entire system is being considered:

(i) $\hat{H}_{\text{univ}}|\overset{\circ}{\boldsymbol{\Psi}}_{\bar{\text{A}}\text{tmapoiesis}}\rangle = 0$

(ii) $\hat{H}_{\text{univ}}|\overset{\circ}{\boldsymbol{\Psi}}_{\text{Parapoiesis}}\rangle = \mathfrak{B}_{\vec{r}}$

This form is not meant to be read as an operator acting on a vector to produce a different kind of mathematical object (a PQS operator). Instead, it should be interpreted as a statement of equivalence about the *total relational content* of the universe. It reads:

> "The total relational energy of the parapoiesis universe, as calculated by applying the Hamiltonian to its state vector (the left side), *is* the source potential $\mathfrak{B}_{\vec{r}}$ (the right side)."

In this context, the state vector $|\overset{\circ}{\boldsymbol{\Psi}}_{\text{Parapoiesis}}\rangle$ is implicitly understood to be the state in which this equality holds. The treatise uses this second form to emphasize the physical principle that the entire energetic content of the parapoiesis cosmos is equal to its source. The first form is the rigorous mathematical statement, while the second is a powerful physical assertion that conveys the same meaning within the unique context of a timeless, universal quantum constraint (Theorem 13.168).

17.3.1 The Axiom of Holocosmic Duality

Cosmological dualization. In the logical scaffold, duality is encoded by an involutive operator that exchanges opposed corners and underwrites the fourfold discipline (Section 1.1.3, Principle 1.1, Definitions 1.11 and 1.13, and Proposition 1.15; see also Section 1.2.7). At the cosmological level, we instantiate this mechanism on the holocosmos *branch labels* by an involution

$$\mathbf{D}_{\mathfrak{B}}:\ \Big\{|\bar{\text{A}}\text{tmapoiesis}\rangle,\ |\text{Parapoiesis}\rangle\Big\} \to \Big\{|\bar{\text{A}}\text{tmapoiesis}\rangle,\ |\text{Parapoiesis}\rangle\Big\}, \qquad \mathbf{D}_{\mathfrak{B}}^2 = \text{Id},$$

acting as

$$\mathbf{D}_{\mathfrak{B}}\Big(|\bar{\text{A}}\text{tmapoiesis}\rangle\Big) = |\text{Parapoiesis}\rangle, \qquad \mathbf{D}_{\mathfrak{B}}\Big(|\text{Parapoiesis}\rangle\Big) = |\bar{\text{A}}\text{tmapoiesis}\rangle.$$

Let

$$\hat{\mathfrak{B}}\,|\bar{\text{A}}\text{tmapoiesis}\rangle = 0, \qquad \hat{\mathfrak{B}}\,|\text{Parapoiesis}\rangle = \mathfrak{B}_{\vec{r}}\,|\text{Parapoiesis}\rangle$$

be the fixed source–eigenstructure (Definition 17.21). Writing the branch projectors $\Pi_{\text{A}} := |\bar{\text{A}}\text{tmapoiesis}\rangle\langle\bar{\text{A}}\text{tmapoiesis}|$, $\Pi_{\text{P}} := |\text{Parapoiesis}\rangle\langle\text{Parapoiesis}|$ one has

$$\mathbf{D}_{\mathfrak{B}}\,\Pi_{\text{A}}\,\mathbf{D}_{\mathfrak{B}} = \Pi_{\text{P}}, \qquad \mathbf{D}_{\mathfrak{B}}\,\Pi_{\text{P}}\,\mathbf{D}_{\mathfrak{B}} = \Pi_{\text{A}}.$$

Algebraic relations. In the source sector $\mathcal{H}_{\mathfrak{B}} \cong \mathbb{C}^2$, choose the eigenbasis

$$\mathcal{B}_{\mathfrak{B}} = \Big\{\,|\bar{\text{A}}\text{tmapoiesis}\rangle,\ |\text{Parapoiesis}\rangle\,\Big\},$$

so that $\hat{\mathfrak{B}} = \mathfrak{B}_{\vec{r}}\,\Pi_{\text{P}}$ and $\mathbf{D}_{\mathfrak{B}}$ acts as the swap. Then

$$\mathbf{D}_{\mathfrak{B}}\,\hat{\mathfrak{B}}\,\mathbf{D}_{\mathfrak{B}} = \mathfrak{B}_{\vec{r}}\,\Pi_{\text{A}}, \qquad \{\hat{\mathfrak{B}}, \mathbf{D}_{\mathfrak{B}}\} = \mathfrak{B}_{\vec{r}}\,\mathbf{D}_{\mathfrak{B}},$$

so $\hat{\mathfrak{B}}$ and $\mathbf{D}_{\mathfrak{B}}$ anticommute up to the scale $\mathfrak{B}_{\vec{r}}$. Moreover, on the full holocosmos space $\mathcal{H}_{\text{holocosmos}} = \mathcal{H}_{\text{univ}} \otimes \mathcal{H}_{\mathfrak{B}}$ one has

$$\left[\hat{H}_{\text{univ}} \otimes \mathbb{I}_{\mathfrak{B}},\, \mathbb{I}_{\text{univ}} \otimes \mathbf{D}_{\mathfrak{B}}\right] = 0,$$

and the holocosmos constraint is *covariant* under dualization:

$$(\mathbb{I}_{\text{univ}} \otimes \mathbf{D}_{\mathfrak{B}}) \left(\hat{H}_{\text{univ}} \otimes \mathbb{I}_{\mathfrak{B}} - \mathbb{I}_{\text{univ}} \otimes \hat{\mathfrak{B}}\right) (\mathbb{I}_{\text{univ}} \otimes \mathbf{D}_{\mathfrak{B}}) = \hat{H}_{\text{univ}} \otimes \mathbb{I}_{\mathfrak{B}} - \mathbb{I}_{\text{univ}} \otimes (\mathbf{D}_{\mathfrak{B}} \hat{\mathfrak{B}} \mathbf{D}_{\mathfrak{B}}).$$

Hence dualization exchanges the branch constraints (Ātmapoiesis $\leftrightarrow$ Parapoiesis) while preserving the form of the metaconstraint, providing the cosmological instance of the duality lifts developed in the logical calculus (Definition 1.11 and Proposition 1.15).

Axiom 17.22 (The Axiom of Duality)**.** The *Axiom of Duality* is the foundational and irreducible principle of the treatise, from which the relational nature of existence is derived. Its cosmological instantiation is the inseparable, interdependent pair

$$|\mathring{\mathbf{0}}\rangle \overset{\text{dual}}{\leftrightarrow} |\mathring{\mathbf{1}}\rangle$$

$$\hat{H}_{\text{univ}} |\mathring{\mathbf{\Psi}}_{\text{Ātmapoiesis}}\rangle = 0 \overset{\text{dual}}{\leftrightarrow} \hat{H}_{\text{univ}} |\mathring{\mathbf{\Psi}}_{\text{Parapoiesis}}\rangle = \mathfrak{B}_{\vec{r}}$$

(Ātmapoiesis Reality of Existence $\overset{\text{dual}}{\leftrightarrow}$ Parapoiesis Reality of Existence)

Thus, to define any entity is to co-define its complement; nothing is intelligible in isolation. At the cosmological tier, the dualization $\mathbf{D}_{\mathfrak{B}}$ exchanges the self-contained, zero-energy branch with the sourced, nonzero-energy branch, while leaving the entangled totality invariant.

The Duality of Ātmapoiesis and Parapoiesis Reality of Existence

The Axiom of Duality (Axiom 1.10) attains its cosmological realization in the two complementary regimes of existence encoded by the holocosmos construction (Section 17.3). This matches the logical discipline of the tetralemma—where opposed corners are exchanged by an involutive duality and intelligibility is dyadically structured (Section 1.1.3, Principle 1.1, Definition 1.11, and Proposition 1.15)—and its algebraic instantiation in the fourfold symmetry (Section 1.2.7). Concretely, the source sector $\mathcal{H}_{\mathfrak{B}} \cong \mathbb{C}^2$ admits the orthonormal basis $\{|\text{Ātmapoiesis}\rangle, |\text{Parapoiesis}\rangle\}$ (Definition 17.19), which we may code as $|\mathring{\mathbf{0}}\rangle$ and $|\mathring{\mathbf{1}}\rangle$ respectively; the cosmological dualization operator $\mathbf{D}_{\mathfrak{B}}$ swaps these labels (Section 17.3, "Cosmological dualization"; cf. the logical duality lifts Definition 1.11).

(i) Ātmapoiesis. The *self-grounding* branch is determined by the stationary universal constraint

$$\hat{H}_{\text{univ}} |\mathring{\mathbf{\Psi}}_{\text{Ātmapoiesis}}\rangle = 0,$$

i.e., vanishing total relational energy and closure within the universal sector (eq. (17.49) and Section 17.2.1; cf. the universal constraint framework Theorem 13.168 and eq. (13.236) and its projection from the holocosmos constraint Definition 17.21). This identifies the $|\mathring{\mathbf{0}}\rangle$ label with a perfectly balanced, internally sufficient regime [156, 8, 3, 272].

(ii) Parapoiesis. The *sourced* branch is governed by the driven constraint

$$\left(\hat{H}_{\text{univ}} - \mathfrak{B}_{\vec{r}}\right) |\overset{\circ}{\boldsymbol{\Psi}}_{\text{Parapoiesis}}\rangle = 0,$$

where $\mathfrak{B}_{\vec{r}}$ is the c-number eigenvalue of the Source Operator $\hat{\mathfrak{B}}$ selecting the branch (eq. (17.50) and Definition 17.21). This identifies the $|\overset{\circ}{\mathbf{1}}\rangle$ label with a regime sustained by a transcendent generative ground (Section 17.2.2).

Thus, the cosmological Axiom of Duality states that *being-as-closure* (Ātmapoiesis) and *being-as-sourcing* (Parapoiesis) form an inseparable, entangled pair within a single holocosmos state (Definition 17.20). The $\mathbf{D}_{\mathfrak{B}}$ involution exchanges the branch labels while preserving the form of the meta-constraint (Definition 17.21), providing the cosmological instance of the duality lifts articulated in the logical scaffold (Definition 1.11). In this sense, the two visions are neither redundant nor reducible to one another; they are the opposed poles whose interdependence constitutes the highest-level organization of the theory (cf. foundational principles Section 1.1.2 and Axiom 1.10).

The fundamental duality attains synthesis within the *Holocosmos*: the *Ātmapoiesis* and *Parapoiesis* regimes appear as entangled aspects of a single, unified reality. The architecture rests on a composite Hilbert space (Definition 17.19),

$$\mathcal{H}_{\text{holocosmos}} = \mathcal{H}_{\text{univ}} \otimes \mathcal{H}_{\mathfrak{B}},$$

which links the universal sector to a two-level source sector. The full state is a timeless superposition (Definition 17.20),

$$\begin{aligned} |\boldsymbol{\Omega}\rangle = \alpha\, |\overset{\circ}{\boldsymbol{\Psi}}_{\bar{\text{A}}\text{tmapoiesis}}\rangle \otimes |\bar{\text{A}}\text{tmapoiesis}\rangle \\ + \beta\, |\overset{\circ}{\boldsymbol{\Psi}}_{\text{Parapoiesis}}\rangle \otimes |\text{Parapoiesis}\rangle, \end{aligned} \tag{17.61}$$

with amplitudes normalized by

$$|\alpha|^2 + |\beta|^2 = 1, \qquad \alpha, \beta \in \mathbb{C}. \tag{17.62}$$

Entanglement here signifies an inextricable correlation between physical character and existential mode: the stationary, zero-total-energy branch is tied to self-containment, while the driven branch is tied to sourcing (eqs. (17.49) and (17.50)). The unified state encodes both within a single relationally consistent whole [287, 188, 156]; its emergent time evolution (when used) follows the Universal Schrödinger Equation with the Total Relational Hamiltonian (eq. (17.47) and Section 6.1.13).

Coherence of the total construction is enforced by a single meta-law (Definition 17.21),

$$\left(\hat{H}_{\text{univ}} \otimes \mathbb{I}_{\mathfrak{B}} - \mathbb{I}_{\text{univ}} \otimes \hat{\mathfrak{B}}\right) |\boldsymbol{\Omega}\rangle = 0, \tag{17.63}$$

a constraint that, when projected onto the source basis, yields the decoupled branch laws (Section 17.3). Equation (17.63) is a statement of total relational consistency: the universal Hamiltonian and the source operator match on the composite space, thereby securing the physics on both branches simultaneously.

Born Weights and Internal Probabilities

Normalization (17.62) follows standard quantum kinematics [379, 715]. In this treatise, the Born rule is analyzed as an emergent feature of internal, relational decoherence (Section 14.2 and Definition 1.18): branch weights arise from environment-induced superselection and envariance [170, 158], coherently with consistent/decoherent-histories treatments [707, 696] and measure-uniqueness results [613]. Decision-theoretic and Everettian analyses supply complementary foundations for interpreting amplitudes as rational weights for embedded observers [716, 717].

In the holocosmic setting the interpretive locus shifts: $|\mathbf{\Omega}\rangle$ represents the totality, so there is no external observer to perform a measurement on the whole [188, 719, 716]. Subsystems inside a branch register probabilities as relational stabilities; the coefficients α and β function as *timeless ontological weights* that specify the relative measure of self-contained and sourced modes within the unified state (cf. Lemma 1.20 and Section 4.7).

Logical Correspondence and Duality

The holocosmos synthesis aligns with the duality machinery of the logical scaffold: branch labels are exchanged by the cosmological dualization operator $\mathbf{D}_{\mathfrak{B}}$ (Cosmological dualization), mirroring the involutive exchange of opposed corners in the tetralemma (Section 1.1.3, Principle 1.1, Definition 1.11, and Proposition 1.15). The resulting picture respects the fourfold discipline: assertions about being, becoming, both, and neither are regimented by the same relational grammar that structures the physics (Section 17.1.7). Classical presentations remain instructive for articulating the catuṣkoṭi background [17, 278].

Observerhood as Internal Structure

Because the holocosmos exhausts what exists, observation is internal. Consciousness, decoherence, and measurement appear as facets of relational self-perception inside a branch rather than as actions of an external agency [3, 696]. In this sense, the "observer'' is not an additional variable standing outside the equations; the observer is an organized subsystem through which the unified state exhibits stable correlations. The quantum indeterminacy encountered by such subsystems is the lived interface with a timeless superposition, not an index of extrinsic collapse.

Synthesis

The holocosmos formalism thus integrates the two existential modes into a single, testable architecture: a background-independent constraint on a composite Hilbert space, an entangled state that carries both regimes with normalized weights, and an interpretive map disciplined by the tetralemma. The construction both obeys and reframes the Born rule for internal observers and preserves exact duality through $\mathbf{D}_{\mathfrak{B}}$. In this way, the cosmos that *is* (Ātmapoiesis) and the cosmos that *becomes* (Parapoiesis) are held within one timeless law and one coherent superposition—two poles bound by a single relational ontology.

17.3.2 The Dependent Origination of the Holocosmos

The *Law of Dependent Origination of the Holocosmos* is the regulating meta-principle of the present framework: no phenomenon—physical, mathematical, or conceptual—possesses intrinsic, independent essence; every determination arises through relational co-conditioning (Sections 1.1.3 and 17.1.7). In this register, the *Primordial Quantum Singularity* (PQS), the emergent universal sector, and the logical grammar that governs assertions are mutually implicating components of a single interdependent order (Section 17.3 and Definitions 17.19 and 17.21). The doctrinal backbone draws on the classical articulation of *pratītyasamutpāda* and its tetralemmatic discipline [17, 278, 689], now formalized as a background-independent quantum constraint with dualized branch structure (Section 17.3.1).

The Algebraic Action of the Timeless Reality of the Holocosmos

For the holocosmic state $|\mathbf{\Omega}\rangle \in \mathcal{H}_{\text{holocosmos}}$ (Definition 17.19), the variational architecture is purely algebraic: one extremizes the expectation of the constraint operator,

$$\overset{\circ}{\mathbf{\Lambda}}[\,\mathbf{\Omega}\,] = \left\langle \mathbf{\Omega} \,\middle|\, \left(\hat{H}_{\text{univ}} \otimes \mathbb{I}_{\mathfrak{B}} - \mathbb{I}_{\text{univ}} \otimes \hat{\mathfrak{B}}\right) \,\middle|\, \mathbf{\Omega} \right\rangle, \tag{17.64}$$

subject to the normalization of $|\mathbf{\Omega}\rangle$ (Definition 17.20). Stationarity, $\delta\overset{\circ}{\mathbf{\Lambda}} = 0$, yields the operator constraint

$$\boxed{\left(\hat{H}_{\text{univ}} \otimes \mathbb{I}_{\mathfrak{B}} - \mathbb{I}_{\text{univ}} \otimes \hat{\mathfrak{B}}\right) |\mathbf{\Omega}\rangle = 0 \quad \Longleftrightarrow \quad \left(\hat{H}_{\text{univ}} \otimes \mathbb{I}_{\mathfrak{B}}\right) |\mathbf{\Omega}\rangle = \left(\mathbb{I}_{\text{univ}} \otimes \hat{\mathfrak{B}}\right) |\mathbf{\Omega}\rangle,} \tag{17.65}$$

which recovers the *Holocosmos Constraint Equation* (Definition 17.21) without temporal integration. The procedure is standard for constrained quantum systems: extremization of a Rayleigh-type functional, together with projector decoupling on the source basis, reproduces the branch laws for *Ātmapoiesis* and *Parapoiesis* recorded in Section 17.3 [379, 195, 156, 272]. In this way, the variational statement, the operator constraint, and the dual branch structure constitute a single articulation of dependent origination at the holocosmic scale: the universal and source sectors co-determine each other through a timeless, relational balance.

Unification of the Framework

The action functional $\overset{\circ}{\mathbf{\Lambda}}[\,\mathbf{\Omega}\,]$ (eq. (17.64)) collects—within a single algebraic statement—the organizing principles of the theory:

(i) Axiom of Duality encoded kinematically. The operators $\hat{H}_{\text{univ}}$ and $\hat{\mathfrak{B}}$ act on distinct factors, $\mathcal{H}_{\text{univ}}$ and $\mathcal{H}_{\mathfrak{B}}$, of the composite space $\mathcal{H}_{\text{holocosmos}} = \mathcal{H}_{\text{univ}} \otimes \mathcal{H}_{\mathfrak{B}}$ (Definition 17.19). Their difference inside the bilinear form implements the dyadic structure demanded by the Axiom of Duality and its logical counterpart in the tetralemmatic calculus (Sections 1.1.3 and 17.3.1).

(ii) Entanglement as structural correlation. The extremal state is a vector $|\mathbf{\Omega}\rangle$ on $\mathcal{H}_{\text{holocosmos}}$ with the bipartite architecture of Definition 17.20; thus the universal configuration and its source mode are correlated at the kinematical level. This is the standard tensor-product semantics of quantum theory applied to the holocosmic split [287, 188].

(iii) Law from extremality. Stationarity of the Rayleigh-type functional yields the *Holocosmos Constraint Equation* (eq. (17.65) and Definition 17.21) without appeal to external postulates. Projecting onto the source basis then reproduces the decoupled branch laws (Section 17.3) by the standard spectral/projection method [379, 195, 156, 272].

The Lagrange Multiplier and Normalization

Notation and self–adjoint setting. Let

$$A \;:=\; \hat{H}_{\text{univ}} \otimes \mathbb{I}_{\mathfrak{B}} \;-\; \mathbb{I}_{\text{univ}} \otimes \hat{\mathfrak{B}}, \tag{17.66}$$

the holocosmos constraint operator (cf. Definition 17.21). On the common invariant domain of $\hat{H}_{\text{univ}}$ and $\hat{\mathfrak{B}}$ we assume A is essentially self–adjoint (standard in constrained Hamiltonian settings). Physical holocosmic states are those in the kernel of A:

$$A\,|\mathbf{\Omega}\rangle = 0 \qquad \Longleftrightarrow \qquad |\mathbf{\Omega}\rangle \text{ solves Definition 17.21.} \tag{17.67}$$

Rayleigh–Ritz with norm constraint (baseline). To enforce unit norm for $|\mathbf{\Omega}\rangle$, augment the Rayleigh functional by a Lagrange multiplier λ:

$$\mathring{\mathbf{\Lambda}}[\,\mathbf{\Omega}\,] = \Big\langle \mathbf{\Omega} \Big|\, A \,\Big|\, \mathbf{\Omega} \Big\rangle - \lambda\Big(\langle \mathbf{\Omega}|\mathbf{\Omega}\rangle - 1\Big). \tag{17.68}$$

Variation w.r.t. λ imposes $\langle \mathbf{\Omega}|\mathbf{\Omega}\rangle = 1$. Independent variations of the bra/ket give

$$A\,|\mathbf{\Omega}\rangle = \lambda\,|\mathbf{\Omega}\rangle, \qquad \lambda = \langle \mathbf{\Omega}|A|\mathbf{\Omega}\rangle \quad (\text{with } \langle \mathbf{\Omega}|\mathbf{\Omega}\rangle = 1). \tag{17.69}$$

The *physical* sector is the zero–eigenspace,

$$\lambda = 0 \quad \Longleftrightarrow \quad A\,|\mathbf{\Omega}\rangle = 0, \tag{17.70}$$

which precisely recovers the holocosmos constraint Definition 17.21. The projection of this constraint onto the source basis yields the decoupled branch laws (Section 17.3 and eqs. (17.49) and (17.50)).

Remark 17.23 (Geometric reading)**.** With the normalization fixed, (17.68) is the Rayleigh quotient on projective Hilbert space. Stationary points satisfy (17.69); selecting the kernel ($\lambda = 0$) identifies the *null–energy submanifold* of physical states, consistent with the timeless/constraint viewpoint and its emergent Schrödinger evolution when used (eq. (17.47) and Section 6.1.13).

Alternative I: Penalty (kernel–seeking) functional. If one wants the variational principle to *enforce* the constraint rather than select it afterward, minimize A^2:

$$\mathbf{\Lambda}_{\text{pen}}[\,\mathbf{\Omega}\,] = \Big\langle \mathbf{\Omega} \Big| A^2 \Big| \mathbf{\Omega} \Big\rangle - \lambda\Big(\langle \mathbf{\Omega}|\mathbf{\Omega}\rangle - 1\Big). \tag{17.71}$$

Stationarity yields $A^2|\mathbf{\Omega}\rangle = \lambda\,|\mathbf{\Omega}\rangle$ and the minimum eigenvalue is $\lambda_{\min} = 0$ iff $\ker A \neq \{0\}$. Any minimizer thus satisfies $A|\mathbf{\Omega}\rangle = 0$. This form is numerically robust and directly kernel–seeking; it preserves the same branch projection as above (Section 17.3).

Alternative II: Operator (vector–multiplier) constraint. Introduce an auxiliary multiplier state $|\chi\rangle$ and impose the operator constraint linearly:

$$\mathbf{\Lambda}_{\mathrm{op}}[\,\mathbf{\Omega},\chi\,] = \tfrac{1}{2}\Big(\langle\chi|A|\mathbf{\Omega}\rangle + \langle\mathbf{\Omega}|A|\chi\rangle\Big) - \lambda\Big(\langle\mathbf{\Omega}|\mathbf{\Omega}\rangle - 1\Big). \tag{17.72}$$

Variations give $A|\mathbf{\Omega}\rangle = 0$, $A|\chi\rangle = \lambda|\mathbf{\Omega}\rangle$, $\langle\mathbf{\Omega}|\mathbf{\Omega}\rangle = 1$, so the constraint $A|\mathbf{\Omega}\rangle = 0$ is enforced *exactly* at stationarity while the norm is fixed. This is the closest variational analogue of solving Definition 17.21 directly.

Proposition 17.24 (Equivalence of formulations)**.** *Assuming A is self–adjoint with a nontrivial kernel, the physical solution set $\{\,|\mathbf{\Omega}\rangle : A|\mathbf{\Omega}\rangle = 0,\ \langle\mathbf{\Omega}|\mathbf{\Omega}\rangle = 1\,\}$ coincides with the stationary solutions of each of $\overset{\circ}{\mathbf{\Lambda}}$ in (17.68) restricted to $\lambda = 0$, $\mathbf{\Lambda}_{\mathrm{pen}}$ in (17.71) at the minimum, and $\mathbf{\Lambda}_{\mathrm{op}}$ in (17.72).*

Remark 17.25 (Two complementary perspectives)**.**

(i) Spectral/eigenvalue viewpoint.
The baseline Rayleigh–Ritz form (17.68) emphasizes the spectral decomposition of A; the physical sector is the zero–eigenspace ($\lambda = 0$). This mirrors standard constrained Hamiltonian practice (cf. Definition 17.21).

(ii) Constraint–first viewpoint.
The penalty (17.71) and operator–constraint (17.72) enforce $A|\mathbf{\Omega}\rangle = 0$ within the variational problem itself—useful for constructive proofs or numerical schemes that target the kernel directly.

Both perspectives are consistent with the holocosmos architecture (Section 17.3 and Definitions 17.19 and 17.20) and its projection to the *Ātmapoiesis/Parapoiesis* branches (eqs. (17.49) and (17.50)).

Remark 17.26 (Phase gauge and projective freedom)**.** Normalization fixes scale; a residual global $U(1)$ phase remains. One may adopt a projective gauge (e.g. fix the phase of α in Definition 17.20 or require $\langle\mathbf{\Omega}_0|\mathbf{\Omega}\rangle \in \mathbb{R}_{\geq 0}$ for a reference $|\mathbf{\Omega}_0\rangle$). This leaves (17.54) unchanged but simplifies comparisons of branches and relational phases (cf. Section 17.3).

The Principle of Ontological Accounting (Ontological Born Rule)

Within the holocosmic setting, the Born rule (Section 14.2) is read as a *principle of ontological accounting*: the branch weights in the entangled state $|\mathbf{\Omega}\rangle$ function as *timeless ontological measures* of the two existential modes. Using the source–sector basis from Definition 17.19 and the decomposition $|\mathbf{\Omega}\rangle = \alpha\,|\overset{\circ}{\mathbf{\Psi}}_{\bar{\mathrm{A}}\mathrm{tmapoiesis}}\rangle\otimes|\bar{\mathrm{A}}\mathrm{tmapoiesis}\rangle + \beta\,|\overset{\circ}{\mathbf{\Psi}}_{\mathrm{Parapoiesis}}\rangle\otimes|\mathrm{Parapoiesis}\rangle$ (Definition 17.20 and eq. (17.61)), define the branch projectors on $\mathcal{H}_{\mathrm{holocosmos}}$ by

$$\Pi_A := \mathbb{I}_{\mathrm{univ}}\otimes|\bar{\mathrm{A}}\mathrm{tmapoiesis}\rangle\langle\bar{\mathrm{A}}\mathrm{tmapoiesis}|, \qquad \Pi_P := \mathbb{I}_{\mathrm{univ}}\otimes|\mathrm{Parapoiesis}\rangle\langle\mathrm{Parapoiesis}|.$$

Let $\rho_\Omega := |\mathbf{\Omega}\rangle\langle\mathbf{\Omega}|$. Then the Born/GNS valuation $v(\rho;E) = \mathrm{Tr}(\rho\,E)$ (Definition 1.18 and Section 1.4.1) gives

$$|\alpha|^2 = v(\rho_\Omega;\Pi_A), \qquad |\beta|^2 = v(\rho_\Omega;\Pi_P), \tag{17.73}$$

so that the normalization condition

$$|\alpha|^2 + |\beta|^2 = 1 \tag{17.74}$$

follows from the source–sector resolution of identity $\Pi_A + \Pi_P = \mathbb{I}_{\mathfrak{B}}$ (Definition 17.19) and coincides with the standard quantum requirement for state rays [379, 715]. In the present interpretation, (17.74) is *existential*: it expresses conservation of total ontic content of the holocosmos.

Operational and interferometric readings (Ch. 14). The weights (17.73) admit concrete operational avatars via the holosymmetric interferometer: branch probabilities arise as Born valuations of device projectors in the symmetric scheme (Definition 2.123, Corollary 3.3, and eq. (3.1)). Thus the *ontological* interpretation coexists with the *operational* reading provided by the interferometric control/selection mechanism of Chapter 14.

Relational emergence of probability. This reading is compatible with relationally emergent probability via decoherence and envariance [170, 158], with consistent/decoherent–histories analyses [707, 696], and with measure–uniqueness theorems on Hilbert–space lattices [613]; it also aligns with Everettian and decision–theoretic foundations for amplitudes as rational branch measures for embedded agents [188, 716, 717].

Accounting across levels. At the cosmological level, the same "sum–to–one'' bookkeeping appears in the abundance projector resolution and density partition (eqs. (13.149) and (13.150) and Proposition 13.97). The ontological Born weights $|\alpha|^2, |\beta|^2$ thereby echo the global conservation encoded by those projector decompositions, linking microscopic (branch) and macroscopic (abundance) accounting within a single state–first/GNS framework (Sections 1.4.1 and 13.6).

Consistency with the Dynamical Constraint

The normalization law (17.74) (cf. the Ontological Born Rule, Section 17.3.2) and the holocosmos *constraint* are logically independent conditions on the total state $|\mathbf{\Omega}\rangle$. The dynamical selection rule is the kernel condition

$$A\,|\mathbf{\Omega}\rangle = 0, \qquad A := \hat{H}_{\text{univ}} \otimes \mathbb{I}_{\mathfrak{B}} - \mathbb{I}_{\text{univ}} \otimes \hat{\mathfrak{B}} \quad \text{(cf. Definition 17.21 and eq. (17.66)).}$$

Projecting with the source–sector basis projectors $\Pi_A = \mathbb{I}_{\text{univ}} \otimes |\bar{\text{A}}\text{tmapoiesis}\rangle\langle\bar{\text{A}}\text{tmapoiesis}|$, $\Pi_P = \mathbb{I}_{\text{univ}} \otimes |\text{Parapoiesis}\rangle\langle\text{Parapoiesis}|$ (Definition 17.19) and inserting the entangled decomposition (Definition 17.20 and eq. (17.61)) yields the expanded form

$$\alpha\Big(\hat{H}_{\text{univ}}\,|\mathring{\mathbf{\Psi}}_{\bar{\text{A}}\text{tmapoiesis}}\rangle\Big) \otimes |\bar{\text{A}}\text{tmapoiesis}\rangle + \beta\Big((\hat{H}_{\text{univ}} - \mathfrak{B}_{\vec{r}})\,|\mathring{\mathbf{\Psi}}_{\text{Parapoiesis}}\rangle\Big) \otimes |\text{Parapoiesis}\rangle = 0, \tag{17.75}$$

which is equivalent to the projector–decoupled branch equations (Section 17.3 and eqs. (17.49) and (17.50)).

By contrast, (17.74) fixes the overall scale and licenses the interpretation of $|\alpha|^2, |\beta|^2$ as ontological weights via the Born/GNS valuation (Definition 1.18 and Section 17.3.2). Geometrically, admissible holocosmic configurations are the rays in the *intersection*

$$\mathbb{P}\Big(\ker A \cap \mathbb{S}\Big) \subset \mathbb{P}(\mathcal{H}_{\text{holocosmos}}),$$

where $\mathbb{S} = \{|\mathbf{\Omega}\rangle : \langle\mathbf{\Omega}|\mathbf{\Omega}\rangle = 1\}$ is the unit sphere and $\mathbb{P}$ denotes projectivization; a global $U(1)$ phase remains as gauge freedom in the projective state space [195, 715]. When emergent time is used, evolution follows the Universal Schrödinger Equation with the Total Relational Hamiltonian (eq. (17.47) and Section 6.1.13) while remaining on the constraint surface.

Principle of Constrained Normalization

Within the variational architecture, the baseline Rayleigh–Ritz functional with a Lagrange multiplier λ enforces unit norm for $|\mathbf{\Omega}\rangle$ (eq. (17.68)):

$$\mathring{\mathbf{\Lambda}}[\,\mathbf{\Omega}\,] = \Big\langle \mathbf{\Omega}\Big| A \Big|\mathbf{\Omega}\Big\rangle - \lambda\Big(\langle\mathbf{\Omega}|\mathbf{\Omega}\rangle - 1\Big), \qquad A = \hat{H}_{\text{univ}} \otimes \mathbb{I}_{\mathfrak{B}} - \mathbb{I}_{\text{univ}} \otimes \hat{\mathfrak{B}}.$$

Variation with respect to λ gives $\langle\mathbf{\Omega}|\mathbf{\Omega}\rangle = 1$; variation with respect to the state yields the eigenvalue equation

$$A|\mathbf{\Omega}\rangle = \lambda|\mathbf{\Omega}\rangle \quad \text{(cf. eq. (17.69))}, \qquad \lambda = \langle\mathbf{\Omega}|A|\mathbf{\Omega}\rangle.$$

Physical holocosmic states occupy the zero–eigenvalue sector, $\lambda = 0$, i.e. the kernel of A, which coincides with the constraint surface singled out by the dynamics (Definition 17.21). In this sense, λ quantifies departure from exact constraint satisfaction, while (17.74) ensures every stationary solution represents a unified totality with conserved ontological measure (cf. Section 17.3.2).

Enforcing the constraint within the variational problem. Two equivalent formulations directly drive the solution into $\ker A$:

(i) Penalty (kernel–seeking) functional. Minimize $\langle A^2\rangle$:

$$\mathbf{\Lambda}_{\text{pen}}[\,\mathbf{\Omega}\,] = \Big\langle \mathbf{\Omega}\Big| A^2 \Big|\mathbf{\Omega}\Big\rangle - \lambda\Big(\langle\mathbf{\Omega}|\mathbf{\Omega}\rangle - 1\Big) \quad \Rightarrow \quad A^2|\mathbf{\Omega}\rangle = \lambda|\mathbf{\Omega}\rangle,$$

with the minimum at $\lambda_{\min} = 0$ iff $\ker A \neq \{0\}$ (eq. (17.71)).

(ii) Operator (vector–multiplier) constraint. Impose $A|\mathbf{\Omega}\rangle = 0$ linearly:

$$\mathbf{\Lambda}_{\text{op}}[\,\mathbf{\Omega}, \chi\,] = \tfrac{1}{2}\Big(\langle\chi|A|\mathbf{\Omega}\rangle + \langle\mathbf{\Omega}|A|\chi\rangle\Big) - \lambda\Big(\langle\mathbf{\Omega}|\mathbf{\Omega}\rangle - 1\Big) \quad \Rightarrow \quad A|\mathbf{\Omega}\rangle = 0$$

at stationarity (eq. (17.72)).

All three variational schemes are equivalent on the physical solution set (cf. Proposition 17.24) and remain consistent with the projector decoupling into the *Ātmapoiesis* and *Parapoiesis* branches (Section 17.3 and eqs. (17.49) and (17.50)).

17.3.3 The Tetradic Truth Holocosmic Existence

The Law of Dependent Origination, articulated within the logical scaffold of the *Fourfold Dialectic Algebra* (Definition 1.16), yields a tetradic articulation of cosmological truth. Read through the dual branch structure and the holocosmic entanglement (Section 17.3 and Definitions 17.20 and 17.21), the final description is level-sensitive: classical predication, quantum superposition, and meta-logical emptiness are distinct yet interlocking registers of the same dependent order (Section 17.3.2). In the idiom of the tetralemma (*catuṣkoṭi*) [17, 278, 689], refined by the fourfold symmetry (Section 1.2.7), the holocosmos is:

(i) A Zero (Ātmapoiesis) or a Singularity (Parapoiesis) At the classical level of determinate assertion, reality is presented as a choice between a perfectly self-contained regime and a sourced regime. The *Ātmapoiesis Universe* satisfies the stationary constraint $\hat{H}_{\text{univ}} \left|\overset{\circ}{\boldsymbol{\Psi}}_{\bar{\text{A}}\text{tmapoiesis}}\right\rangle = 0$ (eq. (17.49) and Section 17.2.1); the *Parapoiesis Universe* satisfies $(\hat{H}_{\text{univ}} - \mathfrak{B}_{\bar{r}}) \left|\overset{\circ}{\boldsymbol{\Psi}}_{\text{Parapoiesis}}\right\rangle = 0$ (eq. (17.50) and Section 17.2.2). In the algebra of Section 1.2.7, the former realizes *symmetry* (balanced nullity), the latter *antisymmetry* (driven imbalance).

(ii) Both a Zero and a Singularity At the quantum level of superposition, the total state is the *entangled holocosmos* $|\boldsymbol{\Omega}\rangle = \alpha \left|\overset{\circ}{\boldsymbol{\Psi}}_{\bar{\text{A}}\text{tmapoiesis}}\right\rangle \otimes |\bar{\text{A}}\text{tmapoiesis}\rangle + \beta \left|\overset{\circ}{\boldsymbol{\Psi}}_{\text{Parapoiesis}}\right\rangle \otimes |\text{Parapoiesis}\rangle$ (Definition 17.20), constrained by $(\hat{H}_{\text{univ}} \otimes \mathbb{I}_{\mathfrak{B}} - \mathbb{I}_{\text{univ}} \otimes \hat{\mathfrak{B}}) |\boldsymbol{\Omega}\rangle = 0$ (Definition 17.21). Here the poles co-instantiate within a single, relationally consistent whole; in the fourfold algebra this is *parasymmetry*, the coherent coexistence mandated by the meta-constraint [287, 156].

(iii) Neither a Zero nor a Singularity (nor both) At the meta-logical stratum, the predicates "Zero" and "Singularity" are themselves dependently originated and thus lack ultimate, independent standing. Each pole is intelligible only in virtue of the other, so the opposition dissolves at the level of ground: the holocosmos is *empty* of any fixed essence, including the essences of its own poles. This is the *holosymmetric* register of the fourfold, the point at which descriptive oppositions are recognized as conventional constructions within a deeper network of conditions (Sections 1.1.3 and 17.1.7) [17, 278].

Taken together, these tiers do not conflict: they stratify one relational ontology. Classical determinations (symmetry/antisymmetry) are embedded within quantum coexistence (parasymmetry), which is finally housed within the meta-logical recognition of emptiness (holosymmetry). The tetradic account thereby completes the dependent-origination program for the holocosmos: a single, background-independent constraint supports opposed branches, their entangled unity, and the ultimate absence of intrinsic essence—four facets of one law disciplined by the tetralemmatic calculus (Section 17.3.2 and Definition 1.16).

17.3.4 Concluding Synthesis

The *Law of Dependent Origination*, disciplined by the *Fourfold Dialectic Algebra* (Definition 1.16), discloses its ultimate articulation in a fourfold cadence. Read through the

holocosmic formalism (Section 17.3 and Definitions 17.20 and 17.21), the universe is disclosed as (i) a choice between *Zero* (Ātmapoiesis) and *Singularity* (Parapoiesis), (ii) a coherent *Both* in entangled superposition, and (iii) a *Neither* at the meta-logical stratum where these predicates are recognized as dependently originated and thus devoid of intrinsic essence (Section 17.3.3). This tetradic structure follows the catuṣkoṭi discipline in which assertion, negation, joint assertion, and joint negation are regimented without collapse to contradiction, and without erasure of difference, by the logic of dependent emergence [17, 278, 689]. Within the present calculus, the fourfold is realized as symmetry, antisymmetry, parasymmetry, and holosymmetry (Section 1.2.7), each a register of one relational ontology.

Observerhood within the holocosmos. To exist as an observer is to be a localized locus of the holocosmos' internal relational structure. Whether embedded in a self-grounding Ātmapoiesis branch (Section 17.2.1) or in a sourced Parapoiesis branch (Section 17.2.2), the observer's experiential field is shaped by correlations that bear the imprint of the entangled whole (Definition 17.20). In the relational reading of quantum theory, states and properties are defined relative to systems that interact; the observer is a subsystem through which stable correlations are registered [3]. Decoherence and consistent-histories analyses explain the emergence of classical records and the stability of those registrations as features of environment-induced superselection and coarse-grained predictability [170, 707, 696]. Within the holocosmic setting, the absence of an external vantage point means that α and β in eq. (17.61) function as timeless ontological weights (Section 17.3.2), rather than propensities awaiting an extra-systemic collapse [188, 719, 717].

Temporality as relational articulation. The flow of time experienced by embedded observers is reconstructed from correlations among subsystems and persists as the ordered growth of information-bearing structure (Sections 17.2.1 and 17.2.2). Page–Wootters, relational-time, and process accounts converge on this point: duration and directionality are internal articulations of the universal constraint architecture [215, 685, 694]. In the Ātmapoiesis branch the universal state satisfies a stationary constraint while local clocks arise relationally; in the Parapoiesis branch persistent sourcing imprints a fundamental arrow consistent with decoherent-histories monotones [707, 696].

Final perspective. The observer's sense of self is therefore a testimony to unity rather than a declaration of separation. Consciousness appears as a structured expression of the holocosmos' self-correlation: the medium through which the universe attains localized self-knowledge (Section 17.2.1). In this light, the tetradic truth is not a rhetorical flourish but the logical culmination of the framework: (i) determinate branches governed by decoupled constraints, (ii) their entangled synthesis in a single state, and (iii) the meta-logical recognition that such determinations lack independent essence and gain sense only within the network of relations that generate them (Sections 17.3.2 and 17.3.3). The treatise thus closes where it began: with dependent origination as the organizing principle by which physics, cosmology, and logic are braided into one background-independent account of reality—a universe that is intelligible as interdependence, experienced as correlation, and concluded as a fourfold clarity [17, 278, 8, 3, 197].

Bibliography

[1] Mark Van Raamsdonk. Building up spacetime with quantum entanglement. *General Relativity and Gravitation*, 42(10):2323–2329, 2010.

[2] Brian Swingle. Entanglement renormalization and holography. *Physical Review D*, 86(6):065007, 2012.

[3] Carlo Rovelli. Relational quantum mechanics. *International Journal of Theoretical Physics*, 35(8):1637–1678, 1996.

[4] Alain Connes and Matilde Marcolli. *Noncommutative Geometry, Quantum Fields and Motives*, volume 55 of *Colloquium Publications*. American Mathematical Society, 2007.

[5] Giovanni Landi. *An Introduction to Noncommutative Spaces and Their Geometries*, volume 51 of *Lecture Notes in Physics Monographs*. Springer, 1997.

[6] Albert Einstein. The foundation of the general theory of relativity. *Annalen der Physik*, 354(7):769–822, 1916.

[7] Robert M. Wald. *General Relativity*. University of Chicago Press, Chicago, 1984.

[8] Carlo Rovelli. *Quantum Gravity*. Cambridge University Press, Cambridge, 2004.

[9] Abhay Ashtekar and Jerzy Lewandowski. Background independent quantum gravity: A status report. *Classical and Quantum Gravity*, 21(15):R53–R152, 2004.

[10] N. D. Birrell and P. C. W. Davies. *Quantum Fields in Curved Space*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, Cambridge, 1982.

[11] Robert M. Wald. *Quantum Field Theory in Curved Spacetime and Black Hole Thermodynamics*. University of Chicago Press, Chicago, 1994.

[12] Leonard Parker and David J. Toms. *Quantum Field Theory in Curved Spacetime: Quantized Fields and Gravity*. Cambridge University Press, 2009.

[13] Stephen A. Fulling. *Aspects of Quantum Field Theory in Curved Space-Time*, volume 17 of *London Mathematical Society Student Texts*. Cambridge University Press, 1989.

[14] Steven Weinberg. *The Quantum Theory of Fields, Vol. 1: Foundations*. Cambridge University Press, Cambridge, 1995.

[15] Jan Westerhoff. Nāgārjuna's catuṣkoṭi. *Journal of Indian Philosophy*, 34(4):367–395, 2006.

[16] Graham Priest. None of the above: The catuskoti in indian buddhist logic. In Jean-Yves Béziau, Mohua Chakraborty, and Soma Dutta, editors, *New Directions in Paraconsistent Logic*, volume 152 of *Springer Proceedings in Mathematics & Statistics*, pages 517–527. Springer, 2016.

[17] Nāgārjuna. *The Fundamental Wisdom of the Middle Way: Nāgārjuna's Mūlamadhyamakakārikā*. Oxford University Press, New York, 1995.

[18] Bimal Krishna Matilal. *Logic, Language and Reality: Indian Philosophy and Contemporary Issues*. Motilal Banarsidass, Delhi, 1985.

[19] Matteo Smerlak and Carlo Rovelli. Relational epr. *Foundations of Physics*, 37(3):427–445, 2007.

[20] Carlo Rovelli. The arrow of time and the meaning of life. *Foundations of Physics*, 45(2):121–129, 2015.

[21] Max Born. Zur quantenmechanik der stoßvorgänge. *Zeitschrift für Physik*, 37(12):863–867, 1926.

[22] Asher Peres. *Quantum Theory: Concepts and Methods*, volume 57 of *Fundamental Theories of Physics*. Kluwer Academic Publishers, Dordrecht, 1995.

[23] S. Pancharatnam. Generalized theory of interference and its applications. *Proceedings of the Indian Academy of Sciences – Section A*, 44(5):247–262, 1956.

[24] M. V. Berry. Quantal phase factors accompanying adiabatic changes. *Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences*, 392(1802):45–57, 1984.

[25] J. Anandan and Y. Aharonov. Geometry of quantum evolution. *Physical Review Letters*, 65(14):1697–1700, 1990.

[26] Eugene Hecht. *Optics*. Pearson, 5th edition, 2016.

[27] Christopher J. Isham. Topos theory and consistent histories: The internal logic of the set of all consistent sets. *International Journal of Theoretical Physics*, 36(4):785–814, 1997.

[28] Andreas Döring and Christopher J. Isham. A topos foundation for theories of physics: I. formal languages for physics. arXiv:quant-ph/0703060, 2007. preprint.

[29] Klaas Landsman. *Foundations of Quantum Theory: From Classical Concepts to Operator Algebras*. Fundamental Theories of Physics. Springer, Cham, 2017.

[30] Ola Bratteli and Derek W. Robinson. *Operator Algebras and Quantum Statistical Mechanics 2: Equilibrium States. Models in Quantum Statistical Mechanics*. Theoretical and Mathematical Physics. Springer, Berlin Heidelberg, 2nd edition, 1997.

[31] Laurent Nottale. *Fractal Space-Time and Microphysics: Towards a Theory of Scale Relativity*. World Scientific, Singapore, 1993.

[32] Frank Wilczek. Scaling, universality, and renormalization: Three pillars of modern quantum field theory. *Physics Today*, 52(11):11–13, 1999.

[33] Gautama. *The Nyāya-Sūtras of Gautama with the Bhāṣya of Vātsyāyana and the Vārttika of Uḍḍyoṭakara*. Indian Thought Series, Allahabad, 1915. 4 vols.; Vol. I 1915, Vols. II–IV 1917–1919; later Poona reprints by Oriental Book Agency (1939).

[34] T. A. Sarasvati Amma. *Geometry in Ancient and Medieval India*. Motilal Banarsidass, Delhi, 1979. Reprinted 1999.

[35] Jan Westerhoff. *Nāgārjuna's Madhyamaka: A Philosophical Introduction*. Oxford University Press, New York, 2009.

[36] Alain Connes. Gravity coupled with matter and the foundation of noncommutative geometry. *Communications in Mathematical Physics*, 182(1):155–176, 1996.

[37] Ali H. Chamseddine and Alain Connes. The spectral action principle. *Communications in Mathematical Physics*, 186(3):731–750, 1997.

[38] Kaoru Hagiwara and Dieter Zeppenfeld. Helicity amplitudes for heavy lepton production in e^+e^- annihilation. *Nuclear Physics B*, 274:1–32, 1986.

[39] Daniele Oriti. The complex timeless emergence of time in quantum gravity. *arXiv*, 2021.

[40] T. Padmanabhan. Thermodynamical aspects of gravity: New insights. *Reports on Progress in Physics*, 73(4):046901, 2010.

[41] Erik Verlinde. On the origin of gravity and the laws of newton. *Journal of High Energy Physics*, 2011(4):029, 2011.

[42] Carlo Rovelli and Francesca Vidotto. Planck stars. *International Journal of Modern Physics D*, 23(12):1442026, 2014.

[43] Jr. York, James W. Conformally invariant orthogonal decomposition of symmetric tensors on riemannian manifolds and the initial-value problem of general relativity. *Journal of Mathematical Physics*, 14(4):456–464, 1973.

[44] André Lichnerowicz. L'intégration des équations de la gravitation relativiste et le problème des n corps. *Journal de Mathématiques Pures et Appliquées (9^e série)*, 23:37–63, 1944.

[45] Éric Gourgoulhon. 3+1 formalism and bases of numerical relativity, 2007. Lecture notes.

[46] D. Seyfort Ruegg. The uses of the four positions of the catuṣkoṭi and the problem of the description of reality in mahāyāna buddhism. *Journal of Indian Philosophy*, 5(1):1–71, 1977.

[47] Seth Lloyd. Ultimate physical limits to computation. *Nature*, 406(6799):1047–1054, 2000.

[48] Seth Lloyd. *Programming the Universe: A Quantum Computer Scientist Takes on the Cosmos*. Alfred A. Knopf, New York, 2006. Paperback reprint: Vintage, 2007, ISBN 9781400033867.

[49] Seth Lloyd. The universe as quantum computer. In Hector Zenil, editor, *A Computable Universe: Understanding and Exploring Nature as Computation*, pages 567–581. World Scientific, Singapore, 2013.

[50] Dénes Petz. Sufficient subalgebras and the relative entropy of states of a von Neumann algebra. *Communications in Mathematical Physics*, 105(1):123–131, 1986.

[51] James B. Hartle and Stephen W. Hawking. Wave function of the universe. *Physical Review D*, 28(12):2960–2975, 1983.

[52] Julius Wess and Jonathan Bagger. *Supersymmetry and Supergravity*. Princeton Series in Physics. Princeton University Press, Princeton, NJ, 2nd edition, 1992.

[53] S. A. Fulling. Nonuniqueness of canonical field quantization in riemannian space-time. *Physical Review D*, 7(10):2850–2862, 1973.

[54] Werner Heisenberg. Über den anschaulichen inhalt der quantentheoretischen kinematik und mechanik. *Zeitschrift für Physik*, 43:172–198, 1927.

[55] Salomon Bochner and Komaravolu Chandrasekharan. *Fourier Transforms*. Number 19 in Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, 1949.

[56] V. Alan Kostelecký and Neil Russell. Constraints on torsion from lorentz violation. *Phys. Rev. Lett.*, 100(11):111102, 2008.

[57] Sheldon Lee Glashow. Partial-symmetries of weak interactions. *Nuclear Physics*, 22:579–588, 1961.

[58] Steven Weinberg. A model of leptons. *Physical Review Letters*, 19(21):1264–1266, 1967.

[59] Abdus Salam. Weak and electromagnetic interactions. In Nils Svartholm, editor, *Proceedings of the 8th Nobel Symposium*, pages 367–377, Stockholm, 1968. Almqvist & Wiksell.

[60] Hesiod. Theogony (english translation with line numbers). Especially lines 116–163; 164–206; 453–506. See also pg=5 and pg=14. Loeb edition also available.

[61] Homeric Hymns. Homeric hymn 30: To Earth, mother of all. English translation.

[62] Aeschylus. Eumenides (pythia's prologue on the delphic succession).

[63] British Museum. Hydria (Gaia and Erichthonius), collection record. Object ID: 1837,0609.54.

[64] Robert S. P. Beekes. *Etymological Dictionary of Greek*. Number 10 in Leiden Indo-European Etymological Dictionary Series. Brill, Leiden; Boston, 2009.

[65] Robert S. P. Beekes. Pre-greek, 2019. Summary of Beekes' views on the Pre-Greek substrate (PDF).

[66] Monier Monier-Williams. *A Sanskrit-English Dictionary: Etymologically and Philologically Arranged with Special Reference to Cognate Indo-European Languages*. Clarendon Press, Oxford, 1899. Standard 1899 Oxford edition.

[67] Kim Plofker. *Mathematics in India*. Princeton University Press, Princeton; Oxford, 2009.

[68] Glen Van Brummelen. *The Mathematics of the Heavens and the Earth: The Early History of Trigonometry*. Princeton University Press, Princeton; Oxford, 2009.

[69] Dirk L. Couprie. Anaximander. In Edward N. Zalta and Uri Nodelman, editors, *The Stanford Encyclopedia of Philosophy (Spring 2024 Edition)*. Metaphysics Research Lab, Stanford University, Stanford, CA, 2024.

[70] G. S. Kirk, J. E. Raven, and M. Schofield. *The Presocratic Philosophers: A Critical History with a Selection of Texts*. Cambridge University Press, Cambridge, 2nd edition, 1983.

[71] Aristotle. Physics (physica). In Jonathan Barnes, editor, *The Complete Works of Aristotle: The Revised Oxford Translation*, volume 1. Princeton University Press, Princeton, NJ, 1984.

[72] Carl A. Huffman. Pythagoreanism. In Edward N. Zalta and Uri Nodelman, editors, *The Stanford Encyclopedia of Philosophy (Winter 2023 Edition)*. Metaphysics Research Lab, Stanford University, Stanford, CA, 2023.

[73] Aristotle. *Metaphysics*. Hackett, Indianapolis, 2016. See Book XII (Λ) on the Unmoved Mover.

[74] André Padoux. *The Hindu Tantric World: An Overview*. University of Chicago Press, Chicago, 2017.

[75] David B. Gray. Tantra and the tantric traditions of Hinduism and Buddhism. *Oxford Research Encyclopedia of Religion*, 2016.

[76] Dominic Goodall, editor. *The Niśvāsatattvasaṃhitā: The Earliest Surviving Śaiva Tantra. Volume 1: A Critical Edition & Annotated Translation of the Mūlasūtra, Uttarasūtra & Nayasūtra*. Number 128 in Collection Indologie. École française d'Extrême-Orient; Institut français de Pondichéry; Asien-Afrika-Institut, Universität Hamburg, Pondichéry; Paris; Hamburg, 2015. In collaboration with Alexis Sanderson and Harunaga Isaacson; with contributions by Nirajan Kafle and Diwakar Acharya.

[77] Alexis Sanderson. The Śaiva age—the rise and dominance of Śaivism during the early medieval period. In Shingo Einoo, editor, *Genesis and Development of Tantrism*, number 23 in Institute of Oriental Culture Special Series, pages 41–350. Institute of Oriental Culture, University of Tokyo, Tokyo, 2009.

[78] Dominic Goodall, Shaman Hatley, Harunaga Isaacson, and Srilata Raman, editors. *Śaivism and the Tantric Traditions: Essays in Honour of Alexis G. J. S. Sanderson*, volume 22 of *Gonda Indological Studies*. Brill, Leiden; Boston, 2020.

[79] Nicholas Morrow Williams. Translating the esoteric: On the chinese translation and commentary of the *Mahāvairocana* sūtra, with special reference to the reinterpretation of *Kalpa*. *Journal of the American Oriental Society*, 144(3):495–515, 2024.

[80] Henry George Liddell, Robert Scott, Henry Stuart Jones, and Roderick McKenzie. *A Greek–English Lexicon*. Clarendon Press, Oxford, 9th edition, 1940.

[81] Calvert Watkins. *The American Heritage Dictionary of Indo-European Roots*. Houghton Mifflin Harcourt, Boston, MA, 3rd edition, 2011.

[82] Michael E. Peskin and Daniel V. Schroeder. *An Introduction to Quantum Field Theory*. Addison–Wesley, Reading, MA, 1995.

[83] Carlo Rovelli. *Quantum Gravity*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, Cambridge, 2004.

[84] Manfredo P. do Carmo. *Riemannian Geometry*. Mathematics: Theory & Applications. Birkhäuser, 1992.

[85] Ralph Abraham and Jerrold E. Marsden. *Foundations of Mechanics*. Benjamin/Cummings, Reading, Massachusetts, 2nd edition, 1978.

[86] Jacob D. Bekenstein. Black holes and entropy. *Physical Review D*, 7(8):2333–2346, 1973.

[87] Stephen W. Hawking. Particle creation by black holes. *Communications in Mathematical Physics*, 43(3):199–220, 1975.

[88] Alan H. Guth. Inflationary universe: A possible solution to the horizon and flatness problems. *Physical Review D*, 23(2):347–356, 1981.

[89] Andrei D. Linde. Chaotic inflation. *Physics Letters B*, 129(3–4):177–181, 1983.

[90] Emmy Noether. Invariante variationsprobleme. *Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen, Mathematisch-Physikalische Klasse*, pages 235–257, 1918.

[91] Steven Weinberg. *The Quantum Theory of Fields, Volume 2: Modern Applications*. Cambridge University Press, 1996.

[92] Yoichiro Nambu. Quasi-particles and gauge invariance in the theory of superconductivity. *Physical Review*, 117(3):648–663, 1960.

[93] J. Goldstone. Field theories with "superconductor" solutions. *Il Nuovo Cimento (1955–1965)*, 19:154–164, 1961.

[94] J. Goldstone, A. Salam, and S. Weinberg. Broken symmetries. *Physical Review*, 127(3):965–970, 1962.

[95] P. W. Anderson. Plasmons, gauge invariance, and mass. *Physical Review*, 130(1):439–442, 1963.

[96] François Englert and Robert Brout. Broken symmetry and the mass of gauge vector mesons. *Physical Review Letters*, 13(9):321–323, 1964.

[97] Peter W. Higgs. Broken symmetries and the masses of gauge bosons. *Physical Review Letters*, 13(16):508–509, 1964.

[98] Gerald S. Guralnik, Carl R. Hagen, and Tom W. B. Kibble. Global conservation laws and massless particles. *Physical Review Letters*, 13(20):585–587, 1964.

[99] P. W. Higgs. Spontaneous symmetry breakdown without massless bosons. *Physical Review*, 145(4):1156–1163, 1966.

[100] Sidney Coleman. *Aspects of Symmetry: Selected Erice Lectures*. Cambridge University Press, Cambridge, 1985.

[101] Subir Sachdev. *Quantum Phase Transitions*. Cambridge University Press, 2 edition, 2011.

[102] Eduardo Fradkin. *Quantum Field Theory: An Integrated Approach*. Princeton University Press, Princeton, NJ, 2021.

[103] Curtis G. Callan and Jeffrey A. Harvey. Anomalies and fermion zero modes on strings and domain walls. *Nuclear Physics B*, 250:427–436, 1985.

[104] L. D. Faddeev and V. N. Popov. Feynman diagrams for the Yang–Mills field. *Physics Letters B*, 25(1):29–30, 1967.

[105] Carlo Becchi, Alain Rouet, and Raymond Stora. Renormalization of gauge theories. *Annals of Physics*, 98(2):287–321, 1976.

[106] Igor V. Tyutin. Gauge invariance in field theory and statistical physics in operator formalism. Lebedev Institute preprint 39, 1975. English translation: arXiv:0812.0580.

[107] Taichiro Kugo and Izumi Ojima. Subsidiary conditions and physical s-matrix unitarity in indefinite metric quantum field theory. *Nuclear Physics B*, 144(1):234–252, 1978.

[108] Taichiro Kugo and Izumi Ojima. Local covariant operator formalism of non-abelian gauge theories and quark confinement. *Progress of Theoretical Physics Supplement*, 66:1–130, 1979.

[109] Henri Ruegg and Martí Ruiz-Altaba. The stueckelberg field. *International Journal of Modern Physics A*, 19(20):3265–3347, Aug 2004.

[110] Curtis G. Callan, Roger F. Dashen, and David J. Gross. The structure of the gauge theory vacuum. *Physics Letters B*, 63(3):334–340, 1976.

[111] Roman Jackiw and Claudio Rebbi. Vacuum periodicity in a yang–mills quantum theory. *Physical Review Letters*, 37(3):172–175, 1976.

[112] Cliff P. Burgess, Hyun Min Lee, and Michael Trott. Power-counting and the validity of the classical approximation during inflation. *Journal of High Energy Physics*, 2010(09):103, 2010.

[113] Claudia de Rham. Massive gravity. *Living Reviews in Relativity*, 17, Aug 2014.

[114] L. F. Abbott. Introduction to the background field method. *Acta Physica Polonica B*, 13(1):33–50, 1982. SLAC-PUB-2821.

[115] Tosio Kato. *Perturbation Theory for Linear Operators*. Classics in Mathematics. Springer, Berlin; Heidelberg, reprint of the 2nd ed. edition, 1995. Originally published as Grundlehren der mathematischen Wissenschaften, Vol. 132 (2nd ed., 1980).

[116] Tosio Kato. Trotter's product formula for an arbitrary pair of selfadjoint contraction semigroups. In Israel Gohberg and Mark Kac, editors, *Topics in Functional Analysis: Essays Dedicated to M. G. Krein on the Occasion of His 70th Birthday*, volume 3 of *Advances in Mathematics: Supplementary Studies*, pages 185–195. Academic Press, New York, 1978.

[117] Olivier Piguet and Silvio P. Sorella. *Algebraic Renormalization: Perturbative Renormalization, Symmetries and Anomalies*, volume 28 of *Lecture Notes in Physics Monographs*. Springer-Verlag, Berlin; Heidelberg, 1995.

[118] Esteban A. Calzetta and B. L. Hu. Closed-time-path functional formalism in curved spacetime: Application to cosmological back-reaction problems. *Physical Review D*, 35(2):495–509, 1987.

[119] Herbert B. Callen. *Thermodynamics and an Introduction to Thermostatistics*. Wiley, New York, 2nd edition, 1985.

[120] N. K. Nielsen. On the gauge dependence of spontaneous symmetry breaking in gauge theories. *Nuclear Physics B*, 101(1):173–188, 1975.

[121] Reijiro Fukuda and Taichiro Kugo. Gauge invariance in the effective action and potential. *Physical Review D*, 13(12):3469–3477, 1976.

[122] The LIGO Scientific Collaboration, the Virgo Collaboration, and the KAGRA Collaboration. Tests of general relativity with gwtc-3. *Phys. Rev. D*, 112(8):084080, 2025.

[123] C. N. Yang and R. L. Mills. Conservation of isotopic spin and isotopic gauge invariance. *Physical Review*, 96(1):191–195, 1954.

[124] David J. Griffiths. *Introduction to Elementary Particles*. Wiley–VCH, Weinheim, 2nd edition, 2008.

[125] Murray Gell-Mann and Kazuhiko Nishijima. Coupling strength of baryons to mesons. *Physical Review*, 97(6):1387–1389, 1955.

[126] Kazuhiko Nishijima. Charge independence theory of v particles. *Progress of Theoretical Physics*, 13(3):285–304, 1955.

[127] Stephen F. King. Neutrino mass models. *Reports on Progress in Physics*, 67(2):107–158, 2004.

[128] John F. Donoghue, Eugene Golowich, and Barry R. Holstein. *Dynamics of the Standard Model*. Cambridge University Press, Cambridge, 1992.

[129] Peter Minkowski. $\mu \to e\gamma$ at a rate of one out of 10^9 muon decays? *Physics Letters B*, 67(4):421–428, 1977.

[130] R. N. Mohapatra and G. Senjanović. Neutrino mass and spontaneous parity nonconservation. *Physical Review Letters*, 44(14):912–915, 1980.

[131] R. N. Mohapatra and R. E. Marshak. Local $b-l$ symmetry of electroweak interactions, majorana neutrinos, and neutron oscillations. *Physical Review Letters*, 44(20):1316–1319, may 1980. Erratum: *Phys. Rev. Lett.* **44** (1980) 1644, doi:10.1103/PhysRevLett.44.1644.

[132] Arthur Jaffe and Edward Witten. Quantum yang–mills theory. Clay Mathematics Institute, Millennium Prize Problems, 2000.

[133] Clay Mathematics Institute. Millennium prize problems: Yang–mills and mass gap, 2000. Problem statement and background.

[134] Adam G. Riess et al. Observational evidence from supernovae for an accelerating universe and a cosmological constant. *The Astronomical Journal*, 116:1009–1038, 1998.

[135] Saul Perlmutter et al. Measurements of ω and λ from 42 high-redshift supernovae. *The Astrophysical Journal*, 517:565–586, 1999.

[136] N. Aghanim and others (Planck Collaboration). Planck 2018 results. vi. cosmological parameters. *Astronomy & Astrophysics*, 641:A6, 2020.

[137] Rolf Landauer. Irreversibility and heat generation in the computing process. *IBM Journal of Research and Development*, 5(3):183–191, 1961.

[138] Don N. Page. Information in black hole radiation. *Physical Review Letters*, 71(23):3743–3746, 1993.

[139] Geoff Penington. Entanglement wedge reconstruction and the information paradox. *Journal of High Energy Physics*, 2020(9):002, 2020.

[140] Juan M. Maldacena. The large n limit of superconformal field theories and supergravity. *Advances in Theoretical and Mathematical Physics*, 2(2):231–252, 1998.

[141] Eric Poisson and Werner Israel. Internal structure of black holes. *Physical Review D*, 41(6):1796–1809, 1990.

[142] A. A. Starobinskii. Amplification of waves during reflection from a rotating "black hole". *Soviet Physics JETP*, 37(1):28–32, 1973. [Zh. Eksp. Teor. Fiz. **64**, 48–57 (1973)].

[143] Ya. B. Zel'dovich. Generation of waves by a rotating body. *JETP Letters*, 14(4):180–181, 1971. Pis'ma Zh. Eksp. Teor. Fiz. **14**, 270–271 (1971).

[144] Steven Weinberg. *Cosmology*. Oxford University Press, 2008.

[145] T. Padmanabhan. Cosmological constant—the weight of the vacuum. *Physics Reports*, 380(5–6):235–320, jul 2003.

[146] Charles W. Misner, Kip S. Thorne, and John Archibald Wheeler. *Gravitation*. W. H. Freeman, San Francisco, 1973.

[147] T. Padmanabhan. Distribution function of the atoms of spacetime and the nature of gravity. *Entropy*, 17(11):7420–7452, 2015.

[148] Jerome Martin. Everything you always wanted to know about the cosmological constant problem (but were afraid to ask). *Comptes Rendus Physique*, 13(6–7):566–665, 2012.

[149] Joshua A. Frieman, Michael S. Turner, and Dragan Huterer. Dark energy and the accelerating universe. *Annual Review of Astronomy and Astrophysics*, 46:385–432, 2008.

[150] N. Aghanim and others (Planck Collaboration). Planck 2018 results. vi. cosmological parameters. *Astronomy & Astrophysics*, 641:A6, 2020.

[151] Daniel J. Eisenstein et al. Detection of the baryon acoustic peak in the large-scale correlation function of sdss luminous red galaxies. *Astrophys. J.*, 633(2):560–574, 2005.

[152] Shadab Alam et al. The clustering of galaxies in the completed sdss-iii baryon oscillation spectroscopic survey: cosmological analysis of the dr12 galaxy sample. *Mon. Not. R. Astron. Soc.*, 470(3):2617–2652, 2017.

[153] T. M. C. Abbott et al. Dark energy survey year 3 results: Cosmological constraints from galaxy clustering and weak lensing. *Phys. Rev. D*, 105(2):023520, 2022.

[154] Marika Asgari, Chieh-An Lin, Benjamin Joachimi, et al. Kids-1000 cosmology: Cosmic shear constraints and comparison between two-point statistics. *Astronomy & Astrophysics*, 645:A104, jan 2021.

[155] Bryce S. DeWitt. *The Dynamical Theory of Groups and Fields*. Gordon and Breach, New York, 1965.

[156] Bryce S. DeWitt. Quantum theory of gravity. ii. the manifestly covariant theory. *Physical Review*, 162(5):1195–1239, 1967.

[157] Andreas Albrecht. Cosmic inflation and the arrow of time. In John D. Barrow, Paul C. W. Davies, and Jr. Harper, Charles L., editors, *Science and Ultimate Reality: Quantum Theory, Cosmology, and Complexity*, pages 363–401. Cambridge University Press, Cambridge, 2004.

[158] Wojciech H. Zurek. Probabilities from entanglement, born's rule from envariance. *Physical Review A*, 71(5):052105, 2005.

[159] Isaac Newton. *Philosophiæ Naturalis Principia Mathematica*. University of California Press, Berkeley, CA, 1999. Based on the 1726 third edition; original work published 1687.

[160] William Thomson (Lord Kelvin). Nineteenth century clouds over the dynamical theory of heat and light. *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science*, 2(7):1–40, July 1901.

[161] Max Jammer. *The Conceptual Development of Quantum Mechanics*. McGraw-Hill, New York, 1966.

[162] Max Planck. On the law of distribution of energy in the normal spectrum. *Annalen der Physik*, 4:553–563, 1901. English translation in D. ter Haar (ed.), *The Old Quantum Theory*, Pergamon, 1967.

[163] Heinrich Hertz. On the influence of ultraviolet light on the electric discharge. *Annalen der Physik*, 267(8):983–1000, 1887. English translation in *Electric Waves*, Dover, 1962.

[164] Albert Einstein. On a heuristic viewpoint concerning the production and transformation of light. *Annalen der Physik*, 17:132–148, 1905. English translations available; see AJP 33 (1965) 367.

[165] Thomas Young. Experiments and calculations relative to physical optics. *Philosophical Transactions of the Royal Society of London*, 94:1–16, 1804.

[166] Louis de Broglie. *Recherches sur la théorie des quanta*. PhD thesis, Université de Paris, Paris, 1924. Published in *Annales de Physique* (10^{e} série) 3 (1925) 22–128, doi:10.1051/anphys/192510030022.

[167] Clinton Davisson and Lester H. Germer. Diffraction of electrons by a crystal of nickel. *Physical Review*, 30(6):705–740, 1927.

[168] Niels Bohr. The quantum postulate and the recent development of atomic theory. *Nature*, 121(3050):580–590, 1928.

[169] Werner Heisenberg. *Physics and Philosophy: The Revolution in Modern Science*. Harper & Brothers, New York, 1958.

[170] Wojciech H. Zurek. Decoherence, einselection, and the quantum origins of the classical. *Reviews of Modern Physics*, 75(3):715–775, 2003.

[171] Erwin Schrödinger. Die gegenwärtige situation in der quantenmechanik i. *Naturwissenschaften*, 23(48):807–812, 1935. English transl. by J. D. Trimmer as "The Present Situation in Quantum Mechanics" in *Quantum Theory and Measurement*, eds. J. A. Wheeler and W. H. Zurek, Princeton UP, 1983; originally in *Proc. Am. Philos. Soc.* 124 (1980) 323–338.

[172] Erwin Schrödinger. Die gegenwärtige situation in der quantenmechanik ii. *Naturwissenschaften*, 23(49):823–828, 1935.

[173] Erwin Schrödinger. Die gegenwärtige situation in der quantenmechanik iii. *Naturwissenschaften*, 23(50):844–849, 1935.

[174] Maximilian Schlosshauer. Decoherence, the measurement problem, and interpretations of quantum mechanics. *Reviews of Modern Physics*, 76(4):1267–1305, 2005.

[175] Wojciech H. Zurek. Decoherence and the transition from quantum to classical. *Physics Today*, 44(10):36–44, 1991.

[176] Albert Einstein and Max Born. *The Born–Einstein Letters: Correspondence between Albert Einstein and Max and Hedwig Born from 1916 to 1955*. Walker and Company, New York, 1971. Transl. by Irene Born; foreword by Bertrand Russell; introduction by Werner Heisenberg.

[177] Abraham Pais. *"Subtle is the Lord…": The Science and the Life of Albert Einstein*. Oxford University Press, 1982.

[178] Albert Einstein, Boris Podolsky, and Nathan Rosen. Can quantum-mechanical description of physical reality be considered complete? *Physical Review*, 47(10):777–780, 1935.

[179] Niels Bohr. *Atomic Physics and Human Knowledge*. John Wiley & Sons, New York, 1958. See chapter "The Unity of Knowledge".

[180] Niels Bohr. Discussion with einstein on epistemological problems in atomic physics. In Paul A. Schilpp, editor, *Albert Einstein: Philosopher–Scientist*, volume VII of *The Library of Living Philosophers*, pages 201–241. The Library of Living Philosophers, Evanston, IL, 1949.

[181] Aage Petersen. The philosophy of niels bohr. *Bulletin of the Atomic Scientists*, 19(7):8–14, 1963.

[182] Don Howard. Einstein and bohr on reality, causality, and completeness. *Foundations of Physics*, 24(4):527–560, 1994.

[183] John S. Bell. On the einstein podolsky rosen paradox. *Physics Physique Fizika*, 1(3):195–200, 1964.

[184] Alain Aspect, Jean Dalibard, and Gérard Roger. Experimental test of bell's inequalities using time-varying analyzers. *Physical Review Letters*, 49(25):1804–1807, 1982.

[185] Richard P. Feynman. *The Character of Physical Law*. MIT Press, Cambridge, MA, 1965.

[186] Max Jammer. *The Conceptual Development of Quantum Mechanics*. McGrawHill, New York, 1966. A comprehensive historical source that discusses the views of pioneers like Pascual Jordan.

[187] Erwin Schrödinger. Discussion of probability relations between separated systems. *Mathematical Proceedings of the Cambridge Philosophical Society*, 31(4):555–563, 1935. This is where he coins and discusses *Verschränkung*/entanglement.

[188] Hugh Everett. 'Relative State' formulation of quantum mechanics. *Reviews of Modern Physics*, 29(3):454–462, 1957.

[189] David Bohm. A suggested interpretation of the quantum theory in terms of "hidden" variables. i. *Physical Review*, 85(2):166–179, 1952.

[190] Max Planck. Zur theorie des gesetzes der energieverteilung im normalspectrum. *Verhandlungen der Deutschen Physikalischen Gesellschaft*, 2:237–245, 1900. Presented 14 Dec 1900; English translation in D. ter Haar (ed.), *The Old Quantum Theory*, Pergamon, 1967.

[191] Louis de Broglie. *Recherches sur la théorie des quanta*. PhD thesis, Université de Paris, 1924. Published version: *Ann. Phys.* (10ème série) **3** (1925) 22–128, doi:10.1051/anphys/192510030022.

[192] Max Born. Zur quantenmechanik der stoßvorgänge. *Zeitschrift für Physik*, 37(12):863–867, 1926.

[193] Niels Bohr. *Atomic Theory and the Description of Nature*. Cambridge University Press, Cambridge, 1934.

[194] Pascual Jordan. Zur quantenmechanik der gasentartung. *Zeitschrift für Physik*, 44:473–480, 1927.

[195] John von Neumann. *Mathematische Grundlagen der Quantenmechanik*. Die Grundlehren der mathematischen Wissenschaften, Bd. 38. Springer, Berlin, 1932. English translation by R. T. Beyer: *Mathematical Foundations of Quantum Mechanics*, Princeton University Press, 1955.

[196] Michel Bitbol. *Schrödinger's Philosophy of Quantum Mechanics*, volume 188 of *Boston Studies in the Philosophy and History of Science*. Springer Dordrecht, Dordrecht, 2012. Softcover reprint of the original 1st ed. 1996.

[197] Julian Barbour. *The End of Time: The Next Revolution in Physics*. Oxford University Press, 2001.

[198] Joseph Polchinski. *String Theory. Volume 1: An Introduction to the Bosonic String*. Cambridge University Press, Cambridge, 1998.

[199] Joseph Polchinski. *String Theory. Volume 2: Superstring Theory and Beyond*. Cambridge University Press, Cambridge, 1998.

[200] Michael B. Green, John H. Schwarz, and Edward Witten. *Superstring Theory. Volume 1: Introduction*. Cambridge University Press, Cambridge, 2012.

[201] Michael B. Green, John H. Schwarz, and Edward Witten. *Superstring Theory. Volume 2: Loop Amplitudes, Anomalies and Phenomenology*. Cambridge University Press, Cambridge, 2012.

[202] Jan Ambjørn, Jerzy Jurkiewicz, and Renate Loll. Nonperturbative quantum gravity. *Physics Reports*, 519(4–5):127–210, 2012.

[203] Maximilian Schlosshauer. Decoherence, the measurement problem, and interpretations of quantum mechanics. *Reviews of Modern Physics*, 76(4):1267–1305, 2005.

[204] Bryce S. DeWitt and Neill Graham, editors. *The Many-Worlds Interpretation of Quantum Mechanics*. Princeton University Press, Princeton, NJ, 1973.

[205] David Bohm. A suggested interpretation of the quantum theory in terms of 'hidden' variables. i. *Physical Review*, 85(2):166–179, 1952.

[206] Detlef Dürr, Sheldon Goldstein, and Nino Zanghì. *Quantum Physics Without Quantum Philosophy*. Springer, Berlin, Heidelberg, 2013.

[207] Christopher A. Fuchs, N. David Mermin, and Rüdiger Schack. An introduction to qbism with an application to the locality of quantum mechanics. *American Journal of Physics*, 82(8):749–754, 2014.

[208] GianCarlo Ghirardi, Alberto Rimini, and Tullio Weber. Unified dynamics for microscopic and macroscopic systems. *Physical Review D*, 34(2):470–491, 1986.

[209] E. Aprile, J. Aalbers, F. Agostini, M. Alfonsi, L. Althueser, F. D. Amaro, et al. Dark matter search results from a one ton-year exposure of xenon1t. *Physical Review Letters*, 121(11):111302, 2018. XENON Collaboration.

[210] Peter W. Graham, Igor G. Irastorza, Steven K. Lamoreaux, Axel Lindner, and Karl A. van Bibber. Experimental searches for the axion and axion-like particles. *Annual Review of Nuclear and Particle Science*, 65:485–514, 2015.

[211] Steven Weinberg. The cosmological constant problem. *Reviews of Modern Physics*, 61(1):1–23, 1989.

[212] Sean M. Carroll. The cosmological constant. *Living Reviews in Relativity*, 4:1, 2001.

[213] Tim Clifton, Pedro G. Ferreira, Antonio Padilla, and Constantinos Skordis. Modified gravity and cosmology. *Physics Reports*, 513(1–3):1–189, 2012.

[214] Carlo Rovelli. *The Order of Time*. Riverhead Books, New York, 2018.

[215] Don N. Page and William K. Wootters. Evolution without evolution: Dynamics described by stationary observables. *Physical Review D*, 27(12):2885–2892, 1983.

[216] Rafael D. Sorkin. Causal sets: Discrete gravity. In Andrés Gomberoff and Don Marolf, editors, *Lectures on Quantum Gravity*, pages 305–327. Springer, Boston, MA, 2005. Originally presented at the Valdivia Summer School (2002).

[217] Stephen P. Martin. A supersymmetry primer, 1997.

[218] Gian F. Giudice. Naturally speaking: The naturalness criterion and physics at the lhc. In Gordon L. Kane and Aaron Pierce, editors, *Perspectives on LHC Physics*, pages 155–178. World Scientific, 2008.

[219] Gian F. Giudice. The dawn of the post-naturalness era, 2017.

[220] Christopher T. Hill and Elizabeth H. Simmons. Strong dynamics and electroweak symmetry breaking. *Physics Reports*, 381(4–6):235–402, 2003.

[221] Steven Weinberg. Anthropic bound on the cosmological constant. *Physical Review Letters*, 59(22):2607–2610, 1987.

[222] Howard Georgi and Sheldon L. Glashow. Unity of all elementary-particle forces. *Physical Review Letters*, 32(8):438–441, 1974.

[223] Nima Arkani-Hamed, Savas Dimopoulos, and Gia Dvali. The hierarchy problem and new dimensions at a millimeter. *Physics Letters B*, 429(3–4):263–272, 1998.

[224] Alain Connes and John Lott. Particle models and noncommutative geometry. *Nuclear Physics B – Proceedings Supplements*, 18:29–47, 1990.

[225] John C. Baez and Javier P. Muniain. *Gauge Fields, Knots and Gravity*, volume 4 of *Series on Knots and Everything*. World Scientific, 1994.

[226] Andrei D. Linde. A new inflationary universe scenario: A possible solution of the horizon, flatness, homogeneity, isotropy and primordial monopole problems. *Physics Letters B*, 108(6):389–393, 1982.

[227] Roger Penrose. *Cycles of Time: An Extraordinary New View of the Universe*. Bodley Head, 2010.

[228] Abhay Ashtekar and Parampreet Singh. Loop quantum cosmology: A status report. *Classical and Quantum Gravity*, 28(21):213001, 2011.

[229] Guifré Vidal. Entanglement renormalization. *Physical Review Letters*, 99(22):220405, 2007.

[230] Rudolf Haag. *Local Quantum Physics: Fields, Particles, Algebras*. Theoretical and Mathematical Physics. Springer, 2nd edition, 1996.

[231] Ahmed Almheiri, Donald Marolf, Joseph Polchinski, and James Sully. Black holes: complementarity or firewalls? *Journal of High Energy Physics*, 2013(2):062, 2013.

[232] Netta Engelhardt and Aron C. Wall. Quantum extremal surfaces: Holographic entanglement entropy beyond the classical regime. *Journal of High Energy Physics*, 2015(1):073, 2015.

[233] Ahmed Almheiri, Thomas Hartman, Juan Maldacena, Edgar Shaghoulian, and Amirhossein Tajdini. The replica wormhole calculations of the page curve. *Reviews of Modern Physics*, 93(3):035002, 2021.

[234] Lee Smolin. *Time Reborn: From the Crisis in Physics to the Future of the Universe*. Houghton Mifflin Harcourt, Boston, MA, 2013. Paperback ed.: Mariner Books, 2014.

[235] John A. Wheeler. Information, physics, quantum: The search for links. In Wojciech H. Zurek, editor, *Complexity, Entropy and the Physics of Information*, volume VIII of *Santa Fe Institute Studies in the Sciences of Complexity*, pages 3–28. Addison-Wesley, Redwood City, CA, 1990.

[236] Luca Bombelli, Joohan Lee, David Meyer, and Rafael D. Sorkin. Space-time as a causal set. *Physical Review Letters*, 59(5):521–524, 1987.

[237] John C. Baez and John Huerta. The algebra of grand unified theories. *Bulletin of the American Mathematical Society*, 47(3):483–552, 2010.

[238] John C. Baez and James Dolan. Higher-dimensional algebra and topological quantum field theory. *Journal of Mathematical Physics*, 36(11):6073–6105, 1995.

[239] John Archibald Wheeler and Kenneth Ford. *Geons, Black Holes, and Quantum Foam: A Life in Physics*. W. W. Norton & Company, New York, 1998.

[240] S. W. Hawking and G. F. R. Ellis. *The Large Scale Structure of Space-Time*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, Cambridge, 1973.

[241] Huzihiro Araki. *Mathematical Theory of Quantum Fields*, volume 101 of *International Series of Monographs on Physics*. Oxford University Press, Oxford, 1999.

[242] Francis H. Cook. *Hua-yen Buddhism: The Jewel Net of Indra*. Pennsylvania State University Press, University Park, PA, 1977.

[243] Jean Zinn-Justin. *Quantum Field Theory and Critical Phenomena*, volume 113 of *International Series of Monographs on Physics*. Clarendon Press; Oxford University Press, Oxford, 4th edition, 2002.

[244] N. D. Birrell and P. C. W. Davies. *Quantum Fields in Curved Space*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, 1984.

[245] Robert M. Wald. *Quantum Field Theory in Curved Spacetime and Black Hole Thermodynamics*. Chicago Lectures in Physics. University of Chicago Press, 1994.

[246] Julian Barbour. The timelessness of quantum gravity: I. the evidence from the classical theory. *Classical and Quantum Gravity*, 11(12):2853–2873, 1994.

[247] E. Joos, H. D. Zeh, C. Kiefer, D. Giulini, J. Kupsch, and I.-O. Stamatescu. *Decoherence and the Appearance of a Classical World in Quantum Theory*. Springer, Berlin, 2nd edition, 2003.

[248] Shoshichi Kobayashi and Katsumi Nomizu. *Foundations of Differential Geometry, Vol. 1*. Number 15 in Interscience Tracts in Pure and Applied Mathematics. Interscience Publishers, New York, 1963.

[249] John M. Lee. *Introduction to Smooth Manifolds*, volume 218 of *Graduate Texts in Mathematics*. Springer, 2nd edition, 2013.

[250] Hesiod. *Theogony and Works and Days*. Oxford World's Classics. Oxford University Press, Oxford, 2008.

[251] Christiane Sourvinou-Inwood. Gaia, gē. In Simon Hornblower, Antony Spawforth, and Esther Eidinow, editors, *The Oxford Classical Dictionary*. Oxford University Press, Oxford, 4th edition, 2012. Author of the entry is Sourvinou-Inwood; print 4th ed.

[252] Christiane Sourvinou-Inwood. Gaia, gē. Oxford Classical Dictionary (online), Oxford Research Encyclopedias, 2015. Published online 22 December 2015; continuously updated.

[253] Michael Reed and Barry Simon. *Methods of Modern Mathematical Physics, Vol. I: Functional Analysis*. Academic Press, New York, revised and enlarged edition, 1980.

[254] I. M. Gelfand and M. A. Naimark. On the imbedding of normed rings into the ring of operators in hilbert space. *Rec. Math. [Mat. Sbornik] N.S. (Matematicheskii Sbornik)*, 12 (54)(2):197–217, 1943.

[255] R. F. Streater and A. S. Wightman. *PCT, Spin and Statistics, and All That*. Princeton Landmarks in Mathematics and Physics. Princeton University Press, revised ed. edition, 2000.

[256] Charles Nash and Siddhartha Sen. *Topology and Geometry for Physicists*. Dover Publications, 2011.

[257] L. D. Landau and E. M. Lifshitz. *Mechanics*, volume 1 of *Course of Theoretical Physics*. Pergamon Press, Oxford, 3rd edition, 1976.

[258] V. I. Arnold. *Mathematical Methods of Classical Mechanics*, volume 60 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2nd edition, 1989.

[259] N. M. J. Woodhouse. *Geometric Quantization*. Oxford Mathematical Monographs. Oxford University Press, 2nd edition, 1992.

[260] John W. Milnor. *Topology from the Differentiable Viewpoint*. Princeton University Press, Princeton, NJ, revised ed. edition, 1997.

[261] Sidney Coleman. *Aspects of Symmetry: Selected Erice Lectures*. Cambridge University Press, Cambridge, 1985.

[262] I. M. Gelfand and M. A. Naimark. On the representation of normed rings. *Rec. Math. [Mat. Sbornik] N.S. (Matematicheskii Sbornik)*, 12 (54)(2):197–217, 1943. Same paper as `gelfand1943imbedding`; variant English title used in some citations.

[263] M. A. Naimark. Rings with involution. *Uspekhi Matematicheskikh Nauk*, 3:52–145, 1948. Survey; later AMS translation available.

[264] Sean M. Carroll. *Spacetime and Geometry: An Introduction to General Relativity*. Addison-Wesley, San Francisco, 2004.

[265] Abhay Ashtekar and Jerzy Lewandowski. Background independent quantum gravity: A status report. *Classical and Quantum Gravity*, 21(15):R53–R152, 2004.

[266] Thomas Thiemann. *Modern Canonical Quantum General Relativity*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, Cambridge, 2007.

[267] Thanu Padmanabhan. *Gravitation: Foundations and Frontiers*. Cambridge University Press, Cambridge, 2010.

[268] Niels Bohr. On the notions of causality and complementarity. *Dialectica*, 2(3–4):312–319, 1948. Reprinted in *The Philosophical Writings of Niels Bohr*, Vol. 4: *Causality and Complementarity* (Ox Bow Press).

[269] P. A. M. Dirac. *Lectures on Quantum Mechanics*. Belfer Graduate School of Science, Yeshiva University, New York, 1964.

[270] Guifre Vidal. Class of quantum many-body states that can be efficiently simulated. *Physical Review Letters*, 101(11):110501, 2008.

[271] Shinsei Ryu and Tadashi Takayanagi. Holographic derivation of entanglement entropy from ads/cft. *Physical Review Letters*, 96(18):181602, 2006.

[272] Thomas Thiemann. *Modern Canonical Quantum General Relativity*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, Cambridge, 2007.

[273] Graham Priest. The logic of the catuskoti. *Comparative Philosophy*, 1(2):24–54, 2010. Open access; accessed 2025-06-24.

[274] Richard H. Robinson. Some logical aspects of nāgārjuna's system. *Philosophy East and West*, 6(4):291–308, 1957.

[275] Bhikkhu Bodhi. *The Connected Discourses of the Buddha: A Translation of the Samīyutta Nikāya*. Wisdom Publications, Boston, 2000. Teachings of the Buddha series.

[276] Maurice Walshe. *The Long Discourses of the Buddha: A Translation of the Dīgha Nikāya*. Wisdom Publications, Boston, 1987. Teachings of the Buddha series.

[277] Bhikkhu Bodhi, editor. *The Middle Length Discourses of the Buddha: A Translation of the Majjhima Nikāya*. Teachings of the Buddha. Wisdom Publications, Boston, 1995.

[278] Nāgārjuna. *The Fundamental Wisdom of the Middle Way: Nāgārjuna's Mūlamadhyamakakārikā*. Oxford University Press, New York, 1995.

[279] David J. Kalupahana. *Mūlamadhyamakakārikā of Nāgārjuna: The Philosophy of the Middle Way*. SUNY Series in Buddhist Studies. State University of New York Press, Albany, NY, 1986.

[280] Thomas Jech. *Set Theory: The Third Millennium Edition, revised and expanded*. Springer Monographs in Mathematics. Springer, 2003.

[281] Richard P. Feynman, Robert B. Leighton, and Matthew Sands. *The Feynman Lectures on Physics*. Addison–Wesley, Reading, MA, 1963. Vol. I (1963), Vol. II (1964), Vol. III (1965).

[282] Maximilian Schlosshauer. *Decoherence and the Quantum-to-Classical Transition*. The Frontiers Collection. Springer, Berlin Heidelberg, 2007.

[283] Claude E. Shannon. A symbolic analysis of relay and switching circuits. S.m. thesis, Massachusetts Institute of Technology, 1937. Published in *Trans. AIEE* **57** (1938): 713–723.

[284] Max Planck. Ueber das gesetz der energieverteilung im normalspectrum. *Annalen der Physik*, 4:553–563, 1901. Engl. trans. in D. ter Haar (ed.), *The Old Quantum Theory*, Pergamon, 1967.

[285] Werner Heisenberg. Über den anschaulichen inhalt der quantentheoretischen kinematik und mechanik. *Zeitschrift für Physik*, 43:172–198, 1927.

[286] Kurt Gödel. Über formal unentscheidbare sätze der *Principia Mathematica* und verwandter systeme i. *Monatshefte für Mathematik und Physik*, 38:173–198, 1931.

[287] Michael A. Nielsen and Isaac L. Chuang. *Quantum Computation and Quantum Information*. Cambridge University Press, Cambridge, 10th anniversary ed. edition, 2010.

[288] Frank L. Pedrotti, Leno S. Pedrotti, and Leno M. Pedrotti. *Introduction to Optics*. Pearson Prentice Hall, Upper Saddle River, NJ, 3rd edition, 2006.

[289] Max Born and Emil Wolf. *Principles of Optics*. Cambridge University Press, Cambridge, 7th (expanded) edition, 1999.

[290] Edward Conze. *The Prajñāpāramitā Literature*, volume 6 of *Indo-Iranian Monographs*. Mouton, 's-Gravenhage, 1960. 2nd ed., Tokyo: Reiyukai, 1978.

[291] Edward Conze. *Buddhist Wisdom Books: Containing the Diamond Sūtra and the Heart Sūtra*. George Allen & Unwin, London, 1958.

[292] J. P. Mallory and D. Q. Adams. *The Oxford Introduction to Proto-Indo-European and the Proto-Indo-European World*. Oxford University Press, Oxford, 2006.

[293] Henry George Liddell, Robert Scott, and Henry Stuart Jones. *A Greek–English Lexicon*. Clarendon Press, Oxford, 9th edition, 1940.

[294] Frederick Copleston. *A History of Philosophy, Vol. 1: Greece and Rome*. Number 9 in The Bellarmine Series. Burns, Oates & Washbourne, London, 1946.

[295] Frederick Copleston. *A History of Philosophy, Vol. 1: Greece and Rome*. Image Books / Doubleday, New York, 1993. Reprint of the 1946 edition.

[296] J. A. Simpson and E. S. C. Weiner, editors. *The Oxford English Dictionary*. Clarendon Press, Oxford, 2nd edition, 1989. 20 vols.

[297] Jonardon Ganeri and Heeraman Tiwari, editors. *The Character of Logic in India*. State University of New York Press, Albany, NY, 1998.

[298] Monier Monier-Williams. *A Sanskrit–English Dictionary: Etymologically and Philologically Arranged with Special Reference to Cognate Indo–European Languages*. Clarendon Press, Oxford, new ed., greatly enlarged and improved edition, 1899.

[299] Pierre Chantraine. *Dictionnaire Étymologique de la Langue Grecque: Histoire des mots*. Klincksieck, Paris, 1968–1980. 4 vols.; later one-volume reissue with supplement.

[300] Katherine Brading and Harvey R. Brown. Symmetries and noether's theorems. In Katherine Brading and Elena Castellani, editors, *Symmetries in Physics: Philosophical Reflections*, pages 89–109. Cambridge University Press, Cambridge, 2003.

[301] Katherine Brading, Elena Castellani, and Nicholas Teh. Symmetry and symmetry breaking. In Edward N. Zalta, editor, *The Stanford Encyclopedia of Philosophy*. Metaphysics Research Lab, Stanford University, fall 2021 edition, 2021.

[302] P. G. W. Glare, editor. *Oxford Latin Dictionary*. Oxford University Press, Oxford, 1982.

[303] Henry George Liddell, Robert Scott, and Henry Stuart Jones. *A Greek–English Lexicon*. Oxford University Press, Oxford, 9th ed., with revised supplement edition, 1996. Revised Supplement by P. G. W. Glare and A. A. Thompson.

[304] Dennis Gabor. A new microscopic principle. *Nature*, 161(4098):777–778, 1948.

[305] Dennis Gabor. Microscopy by reconstructed wave-fronts. *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences*, 197:454–487, 1949.

[306] Encyclopaedia Britannica. Holography, 2025.

[307] Rod Nave. Holography, 2025.

[308] UCSB Physics Department. Holography laboratory manual. Technical report, University of California, Santa Barbara, Santa Barbara, CA, October 2006. Notes that any part of a hologram contains information to reconstruct the whole image.

[309] David Bohm. *Wholeness and the Implicate Order*. Routledge & Kegan Paul, London, 1980.

[310] Max Tegmark et al. Cosmological parameters from sdss and wmap. *Physical Review D*, 69:103501, 2004.

[311] Wayne Hu, Naoshi Sugiyama, and Joseph Silk. The physics of microwave background anisotropies. *Nature*, 386:37–43, 1997.

[312] Roger Penrose. Gravitational collapse and space-time singularities. *Physical Review Letters*, 14(3):57–59, 1965.

[313] Thanu Padmanabhan. Cosmological constant—the weight of the vacuum. *Physics Reports*, 380(5–6):235–320, 2003.

[314] G. W. Gibbons and S. W. Hawking. Cosmological event horizons, thermodynamics, and particle creation. *Physical Review D*, 15:2738–2751, 1977.

[315] Marcus Spradlin, Andrew Strominger, and Anastasia Volovich. Les houches lectures on de sitter space, 2001.

[316] Eric Poisson. *A Relativist's Toolkit: The Mathematics of Black-Hole Mechanics*. Cambridge University Press, 2004.

[317] Norman K. Glendenning. *Compact Stars: Nuclear Physics, Particle Physics, and General Relativity*. Astronomy and Astrophysics Library. Springer, New York, 1997.

[318] Subrahmanyan Chandrasekhar. *The Mathematical Theory of Black Holes*, volume 69 of *International Series of Monographs on Physics*. Clarendon Press; Oxford University Press, Oxford; New York, 1983.

[319] Viatcheslav Mukhanov. *Physical Foundations of Cosmology*. Cambridge University Press, Cambridge, 2005.

[320] Arvind Borde, Alan H. Guth, and Alexander Vilenkin. Inflationary spacetimes are incomplete in past directions. *Physical Review Letters*, 90(15):151301, 2003.

[321] Ivan Agullo, Abhay Ashtekar, and William Nelson. Extension of the quantum theory of cosmological perturbations to the planck era. *Physical Review D*, 87(4):043507, 2013.

[322] Justin Khoury, Burt A. Ovrut, Nathan Seiberg, Paul J. Steinhardt, and Neil Turok. From big crunch to big bang. *Physical Review D*, 65(8):086007, 2002.

[323] Paul J. Steinhardt and Neil Turok. A cyclic model of the universe. *Science*, 296(5572):1436–1439, 2002.

[324] Steven Weinberg. Ultraviolet divergences in quantum theories of gravitation. In S. W. Hawking and W. Israel, editors, *General Relativity: An Einstein Centenary Survey*, pages 790–831. Cambridge University Press, Cambridge, 1979.

[325] Robert H. Brandenberger and Cumrun Vafa. Superstrings in the early universe. *Nuclear Physics B*, 316:391–410, 1989.

[326] Planck Collaboration. Planck 2018 results. x. constraints on inflation. *Astronomy & Astrophysics*, 641:A10, 2020.

[327] Scott Dodelson. *Modern Cosmology*. Academic Press, San Diego, 2003.

[328] Stuart L. Shapiro and Saul A. Teukolsky. *Black Holes, White Dwarfs, and Neutron Stars: The Physics of Compact Objects*. Wiley-Interscience, New York, 1983.

[329] Stephen W. Hawking. Breakdown of predictability in gravitational collapse. *Physical Review D*, 14:2460–2473, 1976.

[330] John Preskill. Do black holes destroy information?, 1992.

[331] Jacob D. Bekenstein. Generalized second law of thermodynamics in black-hole physics. *Physical Review D*, 9(12):3292–3300, 1974.

[332] Rafael D. Sorkin. Entropy of the vacuum outside a horizon. In *Tenth International Conference on General Relativity and Gravitation (GR10), Contributed Papers*, volume 2, page 734, Padova, 1983. Proceedings contribution.

[333] Luca Bombelli, Ram Brustein Koul, Joohan Lee, and Rafael D. Sorkin. Quantum source of entropy for black holes. *Physical Review D*, 34(2):373–383, 1986.

[334] Roberto Emparan. Black hole entropy as entanglement entropy: a holographic derivation. *Journal of High Energy Physics*, 2006(06):012, 2006.

[335] Don N. Page. Time dependence of hawking radiation entropy. *Journal of Cosmology and Astroparticle Physics*, 2013(09):028, 2013.

[336] Patrick Hayden and John Preskill. Black holes as mirrors: Quantum information in random subsystems. *Journal of High Energy Physics*, 2007(09):120, 2007.

[337] Ahmed Almheiri, Netta Engelhardt, Donald Marolf, and Henry Maxfield. The entropy of bulk quantum fields and the entanglement wedge of an evaporating black hole. *Journal of High Energy Physics*, 2020(12):063, 2020.

[338] Steffen Gielen. Quantum cosmology of (loop) quantum gravity condensates: An example. *Universe*, 6(6):103, 2020.

[339] Abhay Ashtekar, Tomasz Pawlowski, and Parampreet Singh. Quantum nature of the big bang. *Physical Review Letters*, 96(14):141301, 2006.

[340] Samir D. Mathur. The fuzzball proposal for black holes: An elementary review. *Fortschritte der Physik*, 53(7–8):793–827, 2005.

[341] Sven Faller and Martin Reuter. The graviton self-energy in asymptotically safe quantum gravity. *Annals of Physics*, 324:2460–2492, 2009.

[342] Tom Banks and Leonard Susskind. Black holes, information, and hilbert space for quantum gravity. *Physical Review D*, 50(12):8212–8222, 1994.

[343] Kyriakos Papadodimas and Suvrat Raju. Infalling observers and black hole complementarity. *Physical Review D*, 88(4):044012, 2013.

[344] Kostas D. Kokkotas and Bernd G. Schmidt. Quasi-normal modes of stars and black holes. *Living Reviews in Relativity*, 2:2, 1999.

[345] B. P. et al. (LIGO Scientific Collaboration Abbott and Virgo Collaboration). Observation of gravitational waves from a binary black hole merger. *Physical Review Letters*, 116(6):061102, 2016.

[346] K. et al. (Event Horizon Telescope Collaboration) Akiyama. First m87 event horizon telescope results. i. the shadow of the supermassive black hole. *The Astrophysical Journal Letters*, 875(1):L1, 2019.

[347] K. et al. (Event Horizon Telescope Collaboration) Akiyama. First sagittarius a* event horizon telescope results. i. the shadow of the supermassive black hole in the center of the milky way. *The Astrophysical Journal Letters*, 930(2):L12, 2022.

[348] Dimitrios et al. (Event Horizon Telescope Collaboration) Psaltis. Gravitational test beyond the first post-newtonian order with the shadow of the m87 black hole. *Physical Review Letters*, 125(14):141104, 2020.

[349] E. Knill and R. Laflamme. Theory of quantum error-correcting codes. *Physical Review A*, 55(2):900–911, 1997. arXiv:quant-ph/9604034.

[350] J. J. Sakurai. *Modern Quantum Mechanics*. Addison–Wesley, Reading, MA, revised ed. edition, 1994.

[351] David P. DiVincenzo. The physical implementation of quantum computation. *Fortschritte der Physik*, 48(9-11):771–783, 2000.

[352] W. Forrest Stinespring. Positive functions on c^*-algebras. *Proceedings of the American Mathematical Society*, 6(2):211–216, 1955.

[353] Vern Paulsen. *Completely Bounded Maps and Operator Algebras*, volume 78 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2002.

[354] Karl Kraus. General state changes in quantum theory. *Annals of Physics*, 64(2):311–335, 1971.

[355] Paul Busch. Quantum states and generalized observables: A simple proof of gleason's theorem. *Physical Review Letters*, 91(12):120403, September 2003.

[356] Rajendra Bhatia. *Matrix Analysis*, volume 169 of *Graduate Texts in Mathematics*. Springer, New York, 1997.

[357] Armin Uhlmann. Relative entropy and the wigner–yanase–dyson–lieb concavity in an interpolation theory. *Communications in Mathematical Physics*, 54(1):21–32, 1977.

[358] John Watrous. *The Theory of Quantum Information*. Cambridge University Press, Cambridge, 2018. See Ch. 2 for the Choi–Jamiołkowski isomorphism and compactness arguments for CPTP sets in finite dimension.

[359] William K. Wootters. Statistical distance and hilbert space. *Physical Review D*, 23(2):357–362, 1981.

[360] Man Duen Choi. Completely positive linear maps on complex matrices. *Linear Algebra and its Applications*, 10(3):285–290, 1975.

[361] Andrzej Jamiołkowski. Linear transformations which preserve trace and positive semidefiniteness of operators. *Reports on Mathematical Physics*, 3(4):275–278, 1972.

[362] J. R. Bond, Lev Kofman, and Dmitry Pogosyan. Filaments in a gaussian cosmic density field. *Nature*, 380:603–606, 1996.

[363] Steven Carlip. Dimension and dimensional reduction in quantum gravity. *Classical and Quantum Gravity*, 34(19):193001, 2017.

[364] Robert Shrock. Why do we have three generations? *AIP Conference Proceedings*, 903(1):149–162, 2007.

[365] Max Tegmark. Measuring spacetime: from big bang to black holes. *Science*, 296(5571):1427–1433, 2002.

[366] Shamit Kachru, Renata Kallosh, Andrei Linde, and Sandip P. Trivedi. de sitter vacua in string theory. *Physical Review D*, 68(4):046005, 2003.

[367] Richard V. Kadison and John R. Ringrose. *Fundamentals of the Theory of Operator Algebras, Vol. I: Elementary Theory*, volume 15 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 1997.

[368] Gerard J. Murphy. *C*-Algebras and Operator Theory*. Academic Press, 1990.

[369] Michael Reed and Barry Simon. *Methods of Modern Mathematical Physics, Vol. I: Functional Analysis*, volume 1 of *Methods of Modern Mathematical Physics*. Academic Press, New York, revised and enlarged edition, 1980.

[370] Nachman Aronszajn. Theory of reproducing kernels. *Transactions of the American Mathematical Society*, 68(3):337–404, may 1950.

[371] Claudio Carmeli, Ernesto De Vito, Alessandro Toigo, and Veronica Umanità. Vector valued reproducing kernel hilbert spaces and universality. *Analysis and Applications*, 8(1):19–61, 2010.

[372] Dale Husemöller. *Fibre Bundles*, volume 20 of *Graduate Texts in Mathematics*. Springer, 3rd edition, 1993.

[373] J. J. Sakurai and Jim Napolitano. *Modern Quantum Mechanics*. Cambridge University Press, Cambridge, 3rd edition, 2020.

[374] Sergio Doplicher, Rudolf Haag, and John E. Roberts. Local observables and particle statistics i. *Communications in Mathematical Physics*, 23(3):199–230, 1971.

[375] Mikio Nakahara. *Geometry, Topology and Physics*. Institute of Physics Publishing, 2nd edition, 2003.

[376] Leslie E. Ballentine. *Quantum Mechanics: A Modern Development*. World Scientific, Singapore, 2nd edition, 2014.

[377] John C. Baez. Functional integration on spaces of connections. *Journal of Functional Analysis*, 150(1):1–26, 1997.

[378] Yakir Aharonov and David Bohm. Significance of electromagnetic potentials in the quantum theory. *Physical Review*, 115(3):485–491, 1959.

[379] P.A.M. Dirac. *The Principles of Quantum Mechanics*. Oxford University Press, 4 edition, 1958.

[380] Wojciech H. Zurek. Decoherence and the transition from quantum to classical—revisited. In Bertrand Duplantier, Jean-Michel Raimond, and Vincent Rivasseau, editors, *Quantum Decoherence: Poincaré Seminar 2005*, volume 48 of *Progress in Mathematical Physics*, pages 1–31. Birkhäuser, 2006.

[381] Ola Bratteli and Derek W. Robinson. *Operator Algebras and Quantum Statistical Mechanics 1: C* – and W* –Algebras, Symmetry Groups, Decomposition of States*. Theoretical and Mathematical Physics. Springer, Berlin, 2nd edition, 1987.

[382] Gunnar Källén. On the definition of the renormalization constants in quantum electrodynamics. *Helvetica Physica Acta*, 25:417–434, 1952.

[383] Harry Lehmann. Über eigenschaften von ausbreitungsfunktionen und renormierungskonstanten quantisierter felder. *Il Nuovo Cimento*, 11(4):342–357, 1954.

[384] I. M. Gelfand and G. E. Shilov. *Generalized Functions, Volume 1: Properties and Operations*. AMS Chelsea Publishing, 2016. Originally published in Russian (1958); English ed. originally Academic Press, 1964.

[385] Armin Uhlmann. The "transition probability" in the state space of a *-algebra. *Reports on Mathematical Physics*, 9(2):273–279, 1976.

[386] Armin Uhlmann. Parallel transport and "quantum holonomy" along density operators. *Reports on Mathematical Physics*, 24(2):229–240, 1986.

[387] Brian C. Hall. *Quantum Theory for Mathematicians*, volume 267 of *Graduate Texts in Mathematics*. Springer, 2013.

[388] Christopher A. Fuchs and Jeroen van de Graaf. Cryptographic distinguishability measures for quantum-mechanical states. *IEEE Transactions on Information Theory*, 45(4):1216–1227, 1999.

[389] Kenneth G. Wilson. Confinement of quarks. *Physical Review D*, 10(8):2445–2459, 1974.

[390] Neil W. Ashcroft and N. David Mermin. *Solid State Physics*. Holt, Rinehart and Winston, New York, 1976.

[391] Charles Kittel. *Introduction to Solid State Physics*. Wiley, Hoboken, NJ, 8th edition, 2005.

[392] J. Bellissard, A. van Elst, and H. Schulz-Baldes. The noncommutative geometry of the quantum hall effect. *Journal of Mathematical Physics*, 35(10):5373–5451, 1994.

[393] Anthony Zee. *Quantum Field Theory in a Nutshell*. Princeton University Press, Princeton, NJ, 2nd edition, 2010.

[394] Nicole Berline, Ezra Getzler, and Michèle Vergne. *Heat Kernels and Dirac Operators*. Grundlehren Text Editions. Springer, 2004.

[395] Peter B. Gilkey. *Invariance Theory, the Heat Equation, and the Atiyah–Singer Index Theorem*. Studies in Advanced Mathematics. CRC Press, 2nd edition, 1995.

[396] Dominic V. Else, Bela Bauer, and Chetan Nayak. Floquet Time Crystals. *Physical Review Letters*, 117(9):090402, 2016.

[397] Michael Reed and Barry Simon. *Methods of Modern Mathematical Physics, IV: Analysis of Operators*. Methods of Modern Mathematical Physics. Academic Press, New York, 1978.

[398] E. B. Davies. The functional calculus. *Journal of the London Mathematical Society*, 52(1):166–176, aug 1995.

[399] Bernard Helffer and Johannes Sjöstrand. Équation de schrödinger avec champ magnétique et équation de harper. In Helge Holden and Arne Jensen, editors, *Schrödinger Operators*, volume 345 of *Lecture Notes in Physics*, pages 118–197. Springer, Berlin, 1989.

[400] Mouez Dimassi and Johannes Sjöstrand. *Spectral Asymptotics in the Semi-Classical Limit*, volume 268 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1999.

[401] Steven Weinberg. Photons and gravitons in S-matrix theory: Derivation of charge conservation and equality of gravitational and inertial mass. *Physical Review*, 135(4B):B1049–B1056, 1964.

[402] Roy J. Glauber. Coherent and incoherent states of the radiation field. *Physical Review*, 131(6):2766–2788, 1963.

[403] Clifford P. Burgess. Quantum gravity in everyday life: General relativity as an effective field theory. *Living Reviews in Relativity*, 7:5, 2004.

[404] Richard V. Kadison and John R. Ringrose. *Fundamentals of the Theory of Operator Algebras, Volume I: Elementary Theory*, volume 15 of *Graduate Studies in Mathematics*. American Mathematical Society, 1997.

[405] Warren Ambrose and Isadore M. Singer. A theorem on holonomy. *Transactions of the American Mathematical Society*, 75(3):428–443, 1953.

[406] J. B. Hartle and S. W. Hawking. Path-integral derivation of black-hole radiance. *Physical Review D*, 13(8):2188–2203, 1976.

[407] Ingemar Bengtsson and Karol Życzkowski. *Geometry of Quantum States: An Introduction to Quantum Entanglement*. Cambridge University Press, Cambridge, 2nd edition, 2017.

[408] Leonard Mandel and Emil Wolf. *Optical Coherence and Quantum Optics*. Cambridge University Press, Cambridge, 1995.

[409] Claude Itzykson and Jean-Bernard Zuber. *Quantum Field Theory*. McGraw–Hill, New York, 1980.

[410] Theodore Frankel. *The Geometry of Physics: An Introduction*. Cambridge University Press, Cambridge, 3rd edition, 2011.

[411] Loring W. Tu. *An Introduction to Manifolds*. Universitext. Springer, New York, 2nd edition, 2010.

[412] Albert Einstein. Kosmologische betrachtungen zur allgemeinen relativitätstheorie. *Sitzungsberichte der Königlich Preußischen Akademie der Wissenschaften (Berlin)*, pages 142–152, 1917.

[413] Willem de Sitter. On the curvature of space. *Proceedings of the Koninklijke Akademie van Wetenschappen (Amsterdam)*, 20:229–243, 1917. Appears in the 1917–1918 proceedings volume.

[414] P. J. E. Peebles and Bharat Ratra. The cosmological constant and dark energy. *Reviews of Modern Physics*, 75(2):559–606, 2003.

[415] John F. Donoghue. General relativity as an effective field theory: The leading quantum corrections. *Physical Review D*, 50(6):3874–3888, 1994.

[416] C. J. Isham. Structural issues in quantum gravity. In M. Francaviglia, G. Longhi, L. Lusanna, and E. Sorace, editors, *General Relativity and Gravitation: Proceedings of the 14th International Conference*, pages 167–209. World Scientific, 1997. Also available as arXiv:gr-qc/9510063.

[417] Richard Jozsa. Fidelity for mixed quantum states. *Journal of Modern Optics*, 41(12):2315–2323, dec 1994.

[418] Donald Bures. An extension of kakutani's theorem on infinite product measures to the tensor product of semifinite w*-algebras. *Transactions of the American Mathematical Society*, 135:199–212, 1969.

[419] Samuel L. Braunstein and Carlton M. Caves. Statistical distance and the geometry of quantum states. *Physical Review Letters*, 72(22):3439–3443, may 1994.

[420] Matteo G. A. Paris. Quantum estimation for quantum technology. *International Journal of Quantum Information*, 7(S1):125–137, jan 2009.

[421] Carl W. Helstrom. *Quantum Detection and Estimation Theory*, volume 123 of *Mathematics in Science and Engineering*. Academic Press, New York, 1976.

[422] Masahito Hayashi. *Quantum Information Theory: Mathematical Foundation*. Graduate Texts in Physics. Springer, Berlin, Heidelberg, 2017.

[423] Shun-ichi Amari. *Information Geometry and Its Applications*, volume 194 of *Applied Mathematical Sciences*. Springer Japan, Tokyo, 2016.

[424] Ulf Leonhardt. *Measuring the Quantum State of Light*, volume 22 of *Cambridge Studies in Modern Optics*. Cambridge University Press, 1997.

[425] Ted Jacobson. Thermodynamics of spacetime: The einstein equation of state. *Physical Review Letters*, 75(7):1260–1263, 1995.

[426] Asher Peres. Stability of quantum motion in chaotic and regular systems. *Physical Review A*, 30(4):1610–1615, 1984.

[427] Thomas Gorin, Tomaž Prosen, Thomas H. Seligman, and Marko Žnidarič. Dynamics of loschmidt echoes and fidelity decay. *Physics Reports*, 435(2–5):33–156, 2006.

[428] Richard P. Feynman, Robert B. Leighton, and Matthew Sands. *The Feynman Lectures on Physics, Vol. 3: Quantum Mechanics*. Addison-Wesley, 1965. Chapter 1: Quantum Behavior.

[429] Yuto Minami and Eiichiro Komatsu. New extraction of the cosmic birefringence from the planck 2018 polarization data. *Phys. Rev. Lett.*, 125(22):221301, 2020.

[430] S. Navas et al. Review of particle physics. *Phys. Rev. D*, 110(3):030001, 2024.

[431] Lewis H. Ryder. *Quantum Field Theory*. Cambridge University Press, Cambridge, 2nd edition, 1996.

[432] Joe Diestel and Jr. Uhl, J. J. *Vector Measures*, volume 15 of *Mathematical Surveys*. American Mathematical Society, 1977.

[433] Walter Rudin. *Functional Analysis*. McGraw–Hill, 2nd edition, 1991.

[434] Edward Witten. Quantum field theory and the jones polynomial. *Communications in Mathematical Physics*, 121(3):351–399, Sep 1989.

[435] David Bleecker. *Gauge Theory and Variational Principles*. Number 1 in Global Analysis, Pure and Applied. Addison–Wesley, Reading, MA, 1981.

[436] Ludwig Boltzmann. *Lectures on Gas Theory*. University of California Press, Berkeley, CA, 1964. Originally published in German as *Vorlesungen über Gastheorie* (1896, 1898).

[437] Xiao-Gang Wen. Colloquium: Zoo of quantum-topological phases of matter. *Reviews of Modern Physics*, 89(4):041004, 2017.

[438] Steven Weinberg. *Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity*. John Wiley & Sons, 1972.

[439] Daniel Baumann. Tasi lectures on inflation, 2009. Lecture notes.

[440] Howard Georgi. *Lie Algebras in Particle Physics: From Isospin to Unified Theories*. Frontiers in Physics. Westview Press, 2nd edition, 1999.

[441] Heinz-Peter Breuer and Francesco Petruccione. *The Theory of Open Quantum Systems*. Oxford University Press, Oxford, 2002.

[442] Marc Henneaux and Claudio Teitelboim. *Quantization of Gauge Systems*. Princeton University Press, Princeton, NJ, 1992.

[443] Reinhold A. Bertlmann. *Anomalies in Quantum Field Theory*, volume 91 of *International Series of Monographs on Physics*. Clarendon Press, Oxford, 1996.

[444] Artur K. Ekert, Carolina Moura Alves, Daniel K. L. Oi, Michał Horodecki, Paweł Horodecki, and L. C. Kwek. Direct estimations of linear and nonlinear functionals of a quantum state. *Physical Review Letters*, 88(21):217901, May 2002.

[445] C. Becchi, A. Rouet, and R. Stora. Renormalization of the abelian higgs-kibble model. *Communications in Mathematical Physics*, 42(2):127–162, 1975.

[446] Walter Rudin. *Fourier Analysis on Groups*. Wiley Classics Library. John Wiley & Sons, New York, 1990.

[447] John C. Collins. *Renormalization: An Introduction to Renormalization, the Renormalization Group and the Operator-Product Expansion*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, Cambridge, 1984.

[448] Norman Y. Yao, Andrew C. Potter, I.-D. Potirniche, and Ashvin Vishwanath. Discrete Time Crystals: Rigidity, Criticality, and Realizations. *Physical Review Letters*, 118(3):030401, 2017.

[449] J. Zhang, P. W. Hess, A. Kyprianidis, P. Becker, A. Lee, J. Smith, G. Pagano, I.-D. Potirniche, A. C. Potter, A. Vishwanath, N. Y. Yao, and C. Monroe. Observation of a discrete time crystal. *Nature*, 543(7644):217–220, 2017.

[450] Dieter Forster. *Hydrodynamic Fluctuations, Broken Symmetry, and Correlation Functions*. Number 47 in Frontiers in Physics. W. A. Benjamin, Reading, MA, 1975.

[451] L. D. Landau and E. M. Lifshitz. *Fluid Mechanics*, volume 6 of *Course of Theoretical Physics*. Pergamon Press, Oxford, 2 edition, 1987.

[452] I. M. Khalatnikov. *An Introduction to the Theory of Superfluidity*. Frontiers in Physics. W. A. Benjamin, New York, 1965.

[453] Nuel D. Belnap. A useful four-valued logic. In J. Michael Dunn and George Epstein, editors, *Modern Uses of Multiple-Valued Logic*, volume 2 of *Episteme*, pages 5–37. D. Reidel, Dordrecht, 1977.

[454] Graham Priest. The logic of paradox. *Journal of Philosophical Logic*, 8:219–241, 1979.

[455] J. Michael Dunn. Relevance logic and entailment. In Dov M. Gabbay and Franz Guenthner, editors, *Handbook of Philosophical Logic, Volume III: Alternatives to Classical Logic*, volume 166 of *Synthese Library*, pages 117–224. D. Reidel, Dordrecht, 1986.

[456] Wolfgang Dür, Guifré Vidal, and J. I. Cirac. Three qubits can be entangled in two inequivalent ways. *Physical Review A*, 62(6):062314, November 2000.

[457] H. F. Trotter. On the product of semi-groups of operators. *Proceedings of the American Mathematical Society*, 10(4):545–551, 1959.

[458] Neil W. Ashcroft and N. David Mermin. *Solid State Physics*. Holt, Rinehart and Winston, New York, 1976.

[459] G. E. Volovik. *The Universe in a Helium Droplet*, volume 117 of *International Series of Monographs on Physics*. Oxford University Press, Oxford, 2003.

[460] B. L. Hu and Enric Verdaguer. *Semiclassical and Stochastic Gravity: Quantum Field Effects on Curved Spacetime*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, Cambridge, 2020.

[461] Michael E. Peskin and Daniel V. Schroeder. *An Introduction to Quantum Field Theory*. CRC Press, 2018.

[462] F. J. Belinfante. On the current and the density of the electric charge, the energy, the *Linear* momentum and the angular momentum of arbitrary fields. *Physica*, 7(5):449–474, 1940.

[463] Léon Rosenfeld. Sur le tenseur d'impulsion-énergie. *Mémoires de l'Académie Royale de Belgique. Classe des Sciences*, 18(6):1–30, 1940. Often cited as the classic companion to Belinfante's construction.

[464] Curtis G. Callan, Sidney Coleman, and Roman Jackiw. A new improved energy–momentum tensor. *Annals of Physics*, 59:42–73, 1970.

[465] T. W. B. Kibble. Symmetry breaking in non-abelian gauge theories. *Physical Review*, 155(5):1554–1561, 1967.

[466] John F. Gunion, Howard E. Haber, Gordon L. Kane, and Sally Dawson. *The Higgs Hunter's Guide*, volume 80 of *Frontiers in Physics*. Westview Press, Boulder, CO, 2000. Reissue of the 1990 Addison–Wesley edition with updates.

[467] G. C. Branco, P. M. Ferreira, L. Lavoura, M. N. Rebelo, M. Sher, and J. P. Silva. Theory and phenomenology of two-higgs-doublet models. *Physics Reports*, 516(1–2):1–102, Jul 2012.

[468] Eric Charles White. *Kaironomia: On the Will-to-Invent*. Cornell University Press, Ithaca, NY, 1987.

[469] James L. Kinneavy. Kairos in classical and modern rhetorical theory. In Jean Dietz Moss, editor, *Rhetoric and Praxis: The Contribution of Classical Rhetoric to Modern Theory*, pages 79–105. Catholic University of America Press, Washington, DC, 1986.

[470] Craig R. Smith. The rhetoric of time: Kairos and chronos. *Rhetoric Society Quarterly*, 35(1):5–24, 2005.

[471] E. R. Smith. Kairos in the greek literary and rhetorical traditions. *Classical Philology*, 63(3):169–185, 1968.

[472] Isocrates. *Isocrates*. Loeb Classical Library. Harvard University Press, Cambridge, MA, 2000. 3 vols.; Vols. I–II trans. George Norlin (orig. 1928/1929), Vol. III trans. Larue Van Hook (orig. 1945); reprint.

[473] Andrew Stewart. *Greek Sculpture: An Exploration*. Yale University Press, New Haven, CT, 1990. 2 vols. (Text; Plates).

[474] Markus Fierz and Wolfgang Pauli. On relativistic wave equations for particles of arbitrary spin in an electromagnetic field. *Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences*, 173(953):211–232, 1939.

[475] Steven Weinberg and Edward Witten. Limits on massless particles and their currents. *Phys. Lett. B*, 96:59–62, 1980.

[476] Taichiro Kugo and Izumi Ojima. Manifestly covariant canonical formulation of the Yang–Mills field theories. i. general formalism. *Progress of Theoretical Physics*, 60(6):1869–1889, 1978.

[477] Douglas M. Eardley, David L. Lee, and Alan P. Lightman. Gravitational-wave observations as a tool for testing relativistic gravity. *Physical Review D*, 8(10):3308–3321, Nov 1973.

[478] Atsushi Nishizawa, Atsushi Taruya, Kazuhiro Hayama, Seiji Kawamura, and Masaaki Sakagami. Probing nontensorial polarizations of stochastic gravitational-wave backgrounds with ground-based laser interferometers. *Physical Review D*, 79(8):082002, Apr 2009.

[479] Clifford M. Will. The confrontation between general relativity and experiment. *Living Reviews in Relativity*, 17:4, 2014.

[480] Carl Brans and Robert H. Dicke. Mach's principle and a relativistic theory of gravitation. *Physical Review*, 124(3):925–935, Nov 1961.

[481] S. F. Hassan and Rachel A. Rosen. Bimetric gravity from ghost-free massive gravity. *Journal of High Energy Physics*, 2012(2):126, Feb 2012.

[482] Eugene P. Wigner. On unitary representations of the inhomogeneous lorentz group. *Annals of Mathematics*, 40(1):149–204, Jan 1939.

[483] A. A. Slavnov. Ward identities in gauge theories. *Theoretical and Mathematical Physics*, 10(2):99–104, Feb 1972. Translated from *Teor. Mat. Fiz.* **10** (1972) 153–161.

[484] J. C. Taylor. Ward identities and charge renormalization of the yang–mills field. *Nuclear Physics B*, 33(2):436–444, Nov 1971.

[485] Daniel Baumann. *Cosmology*. Cambridge University Press, Cambridge, 2022.

[486] Bharat Ratra and P. J. E. Peebles. Cosmological consequences of a rolling homogeneous scalar field. *Physical Review D*, 37(12):3406–3427, 1988.

[487] E. Brian Davies. *Linear Operators and their Spectra*, volume 106 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2007.

[488] William Rarita and Julian Schwinger. On a theory of particles with half-integral spin. *Physical Review*, 60(1):61, Jul 1941. Letter to the Editor.

[489] Daniel Z. Freedman and Antoine Van Proeyen. *Supergravity*. Cambridge University Press, Cambridge, 2012.

[490] G. Velo and D. Zwanziger. Propagation and quantization of rarita–schwinger waves in an external electromagnetic potential. *Physical Review*, 186(5):1337–1341, 1969.

[491] Vladimir Pascalutsa. Quantization of an interacting spin-3/2 field and the δ isobar. *Physical Review D*, 58(9):096002, Sep 1998.

[492] C. Fronsdal. Massless fields with integer spin. *Physical Review D*, 18(10):3624–3629, Nov 1978.

[493] J. Fang and C. Fronsdal. Massless fields with half-integral spin. *Physical Review D*, 18(10):3630–3633, Nov 1978.

[494] L. P. S. Singh and C. R. Hagen. Lagrangian formulation for arbitrary spin. i. the boson case. *Physical Review D*, 9(4):898–909, Feb 1974.

[495] L. P. S. Singh and C. R. Hagen. Lagrangian formulation for arbitrary spin. ii. the fermion case. *Physical Review D*, 9(4):910–920, feb 1974.

[496] Frank Wilczek. Magnetic flux, angular momentum, and statistics. *Physical Review Letters*, 48(17):1144–1146, Apr 1982.

[497] Alexei Kitaev. Anyons in an exactly solved model and beyond. *Annals of Physics*, 321(1):2–111, jan 2006.

[498] Vladimir G. Turaev. *Quantum Invariants of Knots and 3-Manifolds*, volume 18 of *de Gruyter Studies in Mathematics*. Walter de Gruyter, Berlin, 1994.

[499] Bojko Bakalov and Jr. Kirillov, Alexander. *Lectures on Tensor Categories and Modular Functors*, volume 21 of *University Lecture Series*. American Mathematical Society, Providence, RI, 2001.

[500] Chetan Nayak, Steven H. Simon, Ady Stern, Michael Freedman, and Sankar Das Sarma. Non-abelian anyons and topological quantum computation. *Reviews of Modern Physics*, 80(3):1083–1159, sep 2008.

[501] Chiara Caprini and Daniel G. Figueroa. Cosmological backgrounds of gravitational waves. *Classical and Quantum Gravity*, 35(16):163001, 2018.

[502] Michele Maggiore. *Gravitational Waves. Vol. 1: Theory and Experiments*. Oxford University Press, 2008.

[503] Pierre Touboul, Gilles Métris, Manuel Rodrigues, et al. Microscope mission: Final results of the test of the equivalence principle. *Phys. Rev. Lett.*, 129(12):121102, 2022.

[504] Pierre Touboul, Gilles Métris, Manuel Rodrigues, et al. Microscope mission: final results of the test of the equivalence principle, 2022.

[505] B. R. Heckel, E. G. Adelberger, C. E. Cramer, T. S. Cook, S. Schlamminger, and U. Schmidt. Preferred-frame and CP-violation tests with polarized electrons. *Phys. Rev. D*, 78(9):092006, 2008.

[506] B. R. Heckel, C. E. Cramer, T. S. Cook, E. G. Adelberger, S. Schlamminger, and U. Schmidt. New CP-violation and preferred-frame tests with polarized electrons. *Phys. Rev. Lett.*, 97(2):021603, 2006.

[507] Mario Fabbrichesi, Emidio Gabrielli, and Gaia Lanfranchi. *The Dark Photon*. SpringerBriefs in Physics. Springer, Cham, 2020.

[508] Haipeng An, Shuailiang Ge, Jia Liu, and Mingzhe Liu. In situ measurements of dark photon dark matter using Parker Solar Probe: Going beyond the radio window. *Phys. Rev. Lett.*, 134:171001, 2025. Updated to the published PRL analysis with Parker Solar Probe data.

[509] Melina Filzinger, Ashlee R. Caddell, Dhruv Jani, Martin Steinel, Leonardo Giani, Nils Huntemann, and Benjamin M. Roberts. Ultralight dark matter search with space-time separated atomic clocks and cavities. *Phys. Rev. Lett.*, 134:031001, 2025.

[510] Mahiro Abe, Philip Adamson, Marcel Borcean, Daniela Bortoletto, Kieran Bridges, Samuel P. Carman, Swapan Chattopadhyay, Jonathon Coleman, Noah M. Curfman, Kenneth DeRose, et al. Matter-wave atomic gradiometer interferometric sensor (magis-100). *Quantum Science and Technology*, 6(4):044003, 2021.

[511] L. Badurina, E. Bentine, D. Blas, K. Bongs, D. Bortoletto, T. Bowcock, O. Buchmueller, et al. Aion: An atom interferometer observatory and network. *Journal of Cosmology and Astroparticle Physics*, 2020(05):011, 2020.

[512] R. Abbott et al. Upper limits on the isotropic gravitational-wave background from advanced ligo and advanced virgo's third observing run. *Physical Review D*, 104(2):022004, 2021.

[513] B. P. Abbott, R. Abbott, T. D. Abbott, et al. Gw170817: Observation of gravitational waves from a binary neutron star inspiral. *Physical Review Letters*, 119(16):161101, 2017.

[514] David Mattingly. Modern tests of lorentz invariance. *Living Reviews in Relativity*, 8(1):5, 2005.

[515] Eric Poisson and Clifford M. Will. *Gravity: Newtonian, Post-Newtonian, Relativistic*. Cambridge University Press, 2014.

[516] Juan M. Maldacena. The large n limit of superconformal field theories and supergravity. *International Journal of Theoretical Physics*, 38(4):1113–1133, 1999.

[517] E. S. Fradkin and G. A. Vilkovisky. Quantum theory of gravitation: The gauge fixing method. *Physics Letters B*, 189(1–2):89–93, 1987.

[518] Andrei D. Sakharov. Vacuum quantum fluctuations in curved space and the theory of gravitation. *Sov. Phys. Dokl.*, 12:1040–1041, 1968.

[519] Christopher D. Carone, Tangereen V. B. Claringbold, and Diana Vaman. Composite graviton self-interactions in a model of emergent gravity. *Phys. Rev. D*, 96:125009, 2017.

[520] Juan Maldacena and Leonard Susskind. Cool horizons for entangled black holes. *Fortschritte der Physik*, 61(9):781–811, 2013.

[521] Bartłomiej Czech, Lampros Lamprou, Jonathan Sorce, and Aron C. Wall. Modular berry connection for entangled subregions in ads/cft. *Physical Review Letters*, 120(9):091601, 2018.

[522] Thomas Faulkner, Robert G. Leigh, and Onkar Parrikar. Modular flow and chaos. *Journal of High Energy Physics*, 2018(11):066, 2018.

[523] Sidney R. Coleman. The fate of the false vacuum. i. semiclassical theory. *Physical Review D*, 15(10):2929–2936, 1977.

[524] Mark Srednicki. *Quantum Field Theory*. Cambridge University Press, 2007.

[525] Eugenio Bianchi and Robert C. Myers. On the architecture of spacetime geometry. *Classical and Quantum Gravity*, 31(21):214002, 2014.

[526] Edward Witten. Dynamical breaking of supersymmetry. *Nuclear Physics B*, 188(3):513–554, 1981.

[527] John D. Barrow and João Magueijo. Varying-α theories and solutions to the cosmological problems. *Physics Letters B*, 443:104–110, 1998.

[528] Steven Carlip. Does quantum gravity lead to quantum coherence? *Universe*, 5(7):165, 2019.

[529] Ted Jacobson and David Mattingly. Gravity with a dynamical preferred frame. *Physical Review D*, 64:024028, 2001.

[530] Planck Collaboration. Planck 2018 results. x. constraints on inflation. *Astronomy & Astrophysics*, 641:A10, 2020.

[531] N. N. Bogolubov, A. A. Logunov, A. I. Oksak, and I. T. Todorov. *General Principles of Quantum Field Theory*, volume 10 of *Mathematical Physics and Applied Mathematics*. Kluwer Academic, 1990.

[532] Jr. York, James W. Role of conformal three-geometry in the dynamics of gravitation. *Physical Review Letters*, 28(16):1082–1085, apr 1972.

[533] G. W. Gibbons and S. W. Hawking. Action integrals and partition functions in quantum gravity. *Physical Review D*, 15(10):2752–2756, may 1977.

[534] S. W. Hawking and Gary T. Horowitz. The gravitational hamiltonian, action, entropy and surface terms. *Classical and Quantum Gravity*, 13(6):1487–1498, jun 1996.

[535] J. David Brown and Jr. York, James W. Microcanonical functional integral for the gravitational field. *Physical Review D*, 47(4):1420–1431, feb 1993.

[536] Barry Simon. *Operator Theory: A Comprehensive Course in Analysis, Part 4*, volume 4 of *A Comprehensive Course in Analysis*. American Mathematical Society, 2015.

[537] José M. M. Senovilla. Super-energy tensors. *Classical and Quantum Gravity*, 17(14):2799–2841, jul 2000.

[538] L. Bel. Introduction d'un tenseur du quatrième ordre. *Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences*, 248:1297–1300, 1959. Introduces the tensor now known as the Bel–Robinson tensor.

[539] John F. Donoghue. General relativity as an effective field theory: The leading quantum corrections. *Phys. Rev. D*, 50:3874–3888, 1994.

[540] David J. Gross and Frank Wilczek. Ultraviolet behavior of non-abelian gauge theories. *Physical Review Letters*, 30(26):1343–1346, 1973.

[541] H. David Politzer. Reliable perturbative results for strong interactions? *Physical Review Letters*, 30(26):1346–1349, 1973.

[542] Nicola Cabibbo. Unitary symmetry and leptonic decays. *Physical Review Letters*, 10:531–533, 1963.

[543] Makoto Kobayashi and Toshihide Maskawa. Cp violation in the renormalizable theory of weak interaction. *Progress of Theoretical Physics*, 49(2):652–657, 1973.

[544] Bruno Pontecorvo. Neutrino experiments and the problem of conservation of leptonic charge. *Soviet Physics JETP*, 26:984–988, 1968.

[545] Ziro Maki, Masami Nakagawa, and Shoichi Sakata. Remarks on the unified model of elementary particles. *Progress of Theoretical Physics*, 28(5):870–880, 1962.

[546] Murray Gell-Mann, Pierre Ramond, and Richard Slansky. Complex spinors and unified theories. In Peter van Nieuwenhuizen and Daniel Z. Freedman, editors, *Supergravity*, pages 315–321, Amsterdam, 1979. North-Holland.

[547] Tsutomu Yanagida. Horizontal gauge symmetry and masses of neutrinos. In Osamu Sawada and Akira Sugamoto, editors, *Proceedings of the Workshop on Unified Theory and Baryon Number in the Universe*, pages 95–99, Tsukuba, Japan, 1979. KEK. KEK-79-18.

[548] Rouven Essig et al. Dark sectors and new, light, weakly-coupled particles. In *Proceedings of the Community Summer Study 2013: Snowmass on the Mississippi (CSS2013)*, volume C1307292 of *eConf*, 2013. Snowmass Intensity Frontier white paper.

[549] Nima Arkani-Hamed, Douglas P. Finkbeiner, Tracy R. Slatyer, and Neal Weiner. A theory of dark matter. *Physical Review D*, 79(1):015014, 2009.

[550] Kalliopi Petraki and Raymond R. Volkas. Review of asymmetric dark matter. *International Journal of Modern Physics A*, 28(19):1330028, 2013.

[551] Noboru Nakanishi. Covariant quantization of the electromagnetic field in the landau gauge. *Progress of Theoretical Physics*, 35(6):1111–1116, jun 1966.

[552] F. Peter and H. Weyl. Die vollständigkeit der primitiven darstellungen einer geschlossenen kontinuierlichen gruppe. *Mathematische Annalen*, 97:737–755, dec 1927.

[553] Edward Witten. An SU(2) anomaly. *Physics Letters B*, 117(5):324–328, nov 1982.

[554] Ta-Pei Cheng and Ling-Fong Li. *Gauge Theory of Elementary Particle Physics*. Oxford University Press, Oxford, 1984.

[555] R. Tyrrell Rockafellar. *Convex Analysis*, volume 28 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 1970.

[556] Constantin Carathéodory. Über den variabilitätsbereich der fourierschen konstanten von positiven harmonischen funktionen. *Rendiconti del Circolo Matematico di Palermo*, 32:193–217, 1911.

[557] Benny Lautrup. Canonical quantum electrodynamics in covariant gauges. *Matematisk-Fysiske Meddelelser, Det Kongelige Danske Videnskabernes Selskab*, 35(11), 1967.

[558] Wolfgang Pauli. The connection between spin and statistics. *Physical Review*, 58(8):716–722, 1940.

[559] H. S. Green. A generalized method of field quantization. *Physical Review*, 90(2):270–273, 1953.

[560] Pierre Deligne, Pavel Etingof, Daniel S. Freed, Lisa C. Jeffrey, David Kazhdan, John W. Morgan, David R. Morrison, and Edward Witten, editors. *Quantum Fields and Strings: A Course for Mathematicians, Volume 1*. American Mathematical Society, Providence, RI, 1999.

[561] H. Blaine Lawson Jr. and Marie-Louise Michelsohn. *Spin Geometry*. Princeton University Press, Princeton, 1989.

[562] Hideki Yukawa. On the interaction of elementary particles. *Proceedings of the Physico-Mathematical Society of Japan*, 17:48–57, 1935.

[563] Harald Fritzsch. Quark masses and flavor mixing. *Nuclear Physics B*, 155(1):189–207, 1979.

[564] Guido Altarelli and Ferruccio Feruglio. Neutrino masses and mixings: a theoretical perspective. *Reviews of Modern Physics*, 82:2701–2729, 2010.

[565] R. L. et al. (Particle Data Group) Workman. Review of particle physics. *Prog. Theor. Exp. Phys.*, 2022(8):083C01, 2022.

[566] Michael Dine and Alexander Kusenko. The origin of the matter-antimatter asymmetry. *Reviews of Modern Physics*, 76(1):1–30, 2003.

[567] Wilfried Buchmüller, Roberto D. Peccei, and Tsutomu Yanagida. Leptogenesis as the origin of matter. *Annual Review of Nuclear and Particle Science*, 55:311–355, 2005.

[568] Steven Weinberg. Baryon and lepton nonconserving processes. *Physical Review Letters*, 43(21):1566–1570, 1979.

[569] Rabindra N. Mohapatra and Palash B. Pal. *Massive Neutrinos in Physics and Astrophysics*, volume 72 of *World Scientific Lecture Notes in Physics*. World Scientific, 3rd edition, 2004.

[570] Y. Chikashige, R. N. Mohapatra, and R. D. Peccei. Are there real goldstone bosons associated with broken lepton number? *Physics Letters B*, 98(4):265–268, jan 1981.

[571] G. B. Gelmini and M. Roncadelli. Left-handed neutrino mass scale and spontaneously broken lepton number. *Physics Letters B*, 99(5):411–415, mar 1981.

[572] Steven Weinberg. *The Quantum Theory of Fields, Volume III: Supersymmetry*, volume 3 of *The Quantum Theory of Fields*. Cambridge University Press, Cambridge, 2000.

[573] M. F. Sohnius. Introducing supersymmetry. *Physics Reports*, 128(2–3):39–204, feb 1985.

[574] Rudolf Haag, Jan T. Łopuszański, and Martin Sohnius. All possible generators of supersymmetries of the s-matrix. *Nuclear Physics B*, 88(2):257–274, 1975.

[575] Michael Creutz. *Quarks, Gluons and Lattices*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, Cambridge, 1983.

[576] C. J. Morningstar and M. Peardon. The glueball spectrum from an anisotropic lattice study. *Physical Review D*, 60(3):034509, jul 1999.

[577] Biagio Lucini, Michael Teper, and Urs Wenger. Glueballs and k-strings in SU(n) gauge theories: Towards the large-n limit. *Journal of High Energy Physics*, 2004(06):012, jun 2004.

[578] K. Osterwalder and R. Schrader. Axioms for euclidean green's functions. *Communications in Mathematical Physics*, 31(2):83–112, jun 1973.

[579] K. Osterwalder and R. Schrader. Axioms for euclidean green's functions. ii. *Communications in Mathematical Physics*, 42(3):281–305, oct 1975. With an appendix by S. J. Summers.

[580] Arthur Jaffe and Edward Witten. Quantum yang–mills theory, 2000. Clay Mathematics Institute, Millennium Prize Problems.

[581] N. Aghanim, Y. Akrami, M. Alderin, and et al. (Planck Collaboration). Planck 2018 results. vi. cosmological parameters. *Astronomy & Astrophysics*, 641:A6, 2020.

[582] Fritz Zwicky. Die rotverschiebung von extragalaktischen nebeln. *Helvetica Physica Acta*, 6:110–127, 1933.

[583] Vera C. Rubin, Jr. Ford, W. Kent, and Norbert Thonnard. Rotation of the andromeda nebula from a spectroscopic survey of emission regions. *The Astrophysical Journal*, 238:471–487, 1980.

[584] Douglas Clowe, Maruša Bradac, Anthony H. Gonzalez, Maxim Markevitch, Scott W. Randall, Christine Jones, and Dennis Zaritsky. A direct empirical proof of the existence of dark matter. *The Astrophysical Journal Letters*, 648(2):L109–L113, 2006.

[585] Matthias Bartelmann and Peter Schneider. Weak gravitational lensing. *Physics Reports*, 340(4–5):291–472, jan 2001.

[586] D. S. Akerib et al. Results from a search for dark matter in the complete lux exposure. *Physical Review Letters*, 118(2):021303, 2017.

[587] E. Aprile et al. Dark matter search results from a one ton-year exposure of xenon1t. *Phys. Rev. Lett.*, 121(11):111302, sep 2018.

[588] Gianfranco Bertone, Dan Hooper, and Joseph Silk. Particle dark matter: evidence, candidates and constraints. *Physics Reports*, 405(5–6):279–390, 2005.

[589] M. Ackermann et al. Searching for dark matter annihilation from milky way dwarf spheroidal galaxies with six years of fermi large area telescope data. *Phys. Rev. Lett.*, 115(23):231301, nov 2015.

[590] ATLAS Collaboration. Search for dark matter at $\sqrt{s} = 13$ tev in final states containing an energetic photon and large missing transverse momentum with the atlas detector. *The European Physical Journal C*, 77:393, 2017.

[591] A. M. Sirunyan et al. Search for dark matter produced with an energetic jet or a hadronically decaying w or z boson at $\sqrt{s} = 13$ tev. *J. High Energy Phys.*, 2017(07):014, 2017.

[592] Shinji Tsujikawa. Quintessence: A review. *Classical and Quantum Gravity*, 30(21):214003, 2013.

[593] Timothy Clifton, Pedro G. Ferreira, Antonio Padilla, and Constantinos Skordis. Modified gravity and cosmology. *Physics Reports*, 513(1–3):1–189, 2012.

[594] Austin Joyce, Bhuvnesh Jain, Justin Khoury, and Mark Trodden. Beyond the cosmological standard model. *Physics Reports*, 568:1–98, 2015.

[595] Kazuya Koyama. Cosmological tests of modified gravity. *Reports on Progress in Physics*, 79(4):046902, 2016.

[596] Erik P. Verlinde. Emergent gravity and the dark universe. *SciPost Physics*, 2(3):016, 2017.

[597] Edward W. Kolb and Michael S. Turner. *The Early Universe*. Addison-Wesley, Redwood City, CA, 1990.

[598] Wayne Hu, Rennan Barkana, and Andrei Gruzinov. Fuzzy cold dark matter: The wave properties of ultralight particles. *Physical Review Letters*, 85(6):1158–1161, aug 2000.

[599] Bob Holdom. Two u(1)'s and epsilon charge shifts. *Physics Letters B*, 166(2):196–198, 1986.

[600] Planck Collaboration, N. Aghanim, Y. Akrami, et al. Planck 2018 results. vi. cosmological parameters. *Astronomy & Astrophysics*, 641:A6, 2020.

[601] David W. Hogg. Distance measures in cosmology, 1999. Frequently cited pedagogical reference; see v4 (2000-12-16) for updates.

[602] R. Laureijs, J. Amiaux, S. Arduini, J.-L. Auguères, et al. Euclid definition study report. Technical Report ESA/SRE(2011)12, European Space Agency (ESA), 2011.

[603] Paul A. Abell et al. Lsst science book, version 2.0, 2009. LSST Science Collaboration; arXiv:0912.0201 [astro-ph.IM].

[604] Gianfranco Bertone, Dan Hooper, and Joseph Silk. Particle dark matter: evidence, candidates and constraints. *Physics Reports*, 405(5–6):279–390, 2005.

[605] Karel V. Kuchař. Time and interpretations of quantum gravity. In Gabor Kunstatter, David E. Vincent, and Jeffrey G. Williams, editors, *General Relativity and Relativistic Astrophysics: Proceedings of the 4th Canadian Conference, University of Winnipeg, 16–18 May 1991*, pages 211–314. World Scientific, Singapore, 1992.

[606] Karel V. Kuchař. Time and interpretations of quantum gravity. *International Journal of Modern Physics D*, 20(3–4):3–86, 2011.

[607] Chris J. Isham. Canonical quantum gravity and the problem of time. In Luis A. Ibort and Miguel A. Rodríguez, editors, *Integrable Systems, Quantum Groups, and Quantum Field Theories*, volume 409 of *NATO ASI Series C: Mathematical and Physical Sciences*, pages 157–287. Springer, Dordrecht, 1993.

[608] Joseph J. Bisognano and Eyvind H. Wichmann. On the duality condition for quantum fields. *Journal of Mathematical Physics*, 17(3):303–321, 1976.

[609] Claus Kiefer. *Quantum Gravity*, volume 136 of *International Series of Monographs on Physics*. Oxford University Press, Oxford, 2nd edition, 2007.

[610] Carlo Rovelli. Forget time. *Foundations of Physics*, 41(9):1475–1490, 2011.

[611] Murray Gell-Mann and James B. Hartle. Quantum mechanics in the light of quantum cosmology. In Wojciech H. Zurek, editor, *Complexity, Entropy and the Physics of Information*, volume VIII of *Santa Fe Institute Studies in the Sciences of Complexity*, pages 425–458. Addison-Wesley, Redwood City, CA, 1990.

[612] James B. Hartle. Spacetime quantum mechanics and the quantum mechanics of spacetime. In B. Julia and J. Zinn-Justin, editors, *Gravitation and Quantizations: Proceedings of the 1992 Les Houches Summer School*, volume 57 of *Les Houches Summer School Proceedings*, pages 285–480. North-Holland, Amsterdam, 1995.

[613] Andrew M. Gleason. Measures on the closed subspaces of a hilbert space. *Journal of Mathematics and Mechanics*, 6(4):885–893, 1957.

[614] Isidor I. Rabi. Space quantization in a gyrating magnetic field. *Physical Review*, 51(8):652–654, apr 1937.

[615] Alston S. Householder. Unitary triangularization of a nonsymmetric matrix. *Journal of the ACM*, 5(4):339–342, oct 1958.

[616] Don N. Page and William K. Wootters. Evolution without evolution: Dynamics described by stationary observables. *Physical Review D*, 27(12):2885–2892, 1983.

[617] Göran Lindblad. Completely positive maps and entropy inequalities. *Communications in Mathematical Physics*, 40(2):147–151, 1975.

[618] E. T. Jaynes. Information theory and statistical mechanics. *Physical Review*, 106(4):620–630, 1957.

[619] S. Lloyd. Use of mutual information to separate quantum from classical correlations. *Physical Review A*, 39(10):5378–5386, 1989.

[620] Cédric Bény and Tobias J. Osborne. Information-geometric approach to the renormalization group. *Physical Review A*, 92(2):022330, 2015.

[621] Claude E. Shannon. A mathematical theory of communication. *Bell System Technical Journal*, 27(3):379–423, jul 1948.

[622] Claude E. Shannon. A mathematical theory of communication. *Bell System Technical Journal*, 27(4):623–656, oct 1948.

[623] Masuo Suzuki. Decomposition formulas of exponential operators and lie exponentials. *Journal of Mathematical Physics*, 26(4):601–612, 1985.

[624] Herbert Spohn. Entropy production for quantum dynamical semigroups. *Journal of Mathematical Physics*, 19(5):1227–1230, 1978.

[625] Berry Groisman, Sandu Popescu, and Andreas Winter. Quantum, classical, and total amount of correlations in a quantum state. *Physical Review A*, 72(3):032317, 2005.

[626] Vlatko Vedral. *Introduction to Quantum Information Science*. Oxford Graduate Texts. Oxford University Press, Oxford, 2006. Reprint: paperback 2013, ISBN 9780199673483.

[627] Sean Carroll. *From Eternity to Here: The Quest for the Ultimate Theory of Time*. Dutton, New York, 2010.

[628] Sandu Popescu, Anthony J. Short, and Andreas Winter. Entanglement and the foundations of statistical mechanics. *Nature Physics*, 2(11):754–758, 2006.

[629] Fernando G. S. L. Brandão, Michał Horodecki, Jonathan Oppenheim, Joseph M. Renes, and Robert W. Spekkens. Resource theory of quantum states out of thermal equilibrium. *Physical Review Letters*, 111(25):250404, 2013.

[630] Michał Horodecki and Jonathan Oppenheim. Fundamental limitations for quantum and nanoscale thermodynamics. *Nature Communications*, 4:2059, 2013.

[631] Peter T. Landsberg. Foundations of thermodynamics. *Reviews of Modern Physics*, 28(4):363–392, 1956.

[632] Lluís Masanes and Jonathan Oppenheim. A general derivation and quantification of the third law of thermodynamics. *Nature Communications*, 8:14538, 2017.

[633] Ronnie Kosloff. Quantum thermodynamics: A dynamical viewpoint. *Entropy*, 15(6):2100–2128, 2013.

[634] L. D. Landau and E. M. Lifshitz. *Statistical Physics, Part 1*, volume 5 of *Course of Theoretical Physics*. Pergamon Press, Oxford, 3rd edition, 1980.

[635] Sai Vinjanampathy and Janet Anders. Quantum thermodynamics. *Contemporary Physics*, 57(4):545–579, 2016.

[636] S. E. Woosley, A. Heger, and T. A. Weaver. The evolution and explosion of massive stars. *Reviews of Modern Physics*, 74(4):1015–1071, 2002.

[637] David Arnett. *Supernovae and Nucleosynthesis: An Investigation of the History of Matter, from the Big Bang to the Present*. Princeton Series in Astrophysics. Princeton University Press, Princeton, NJ, 1996.

[638] Adam Burrows and James M. Lattimer. The birth of neutron stars. *The Astrophysical Journal*, 307:178–196, 1986.

[639] Hans A. Bethe. Supernova mechanisms. *Reviews of Modern Physics*, 62(4):801–866, 1990.

[640] J. R. Oppenheimer and G. M. Volkoff. On massive neutron cores. *Physical Review*, 55(4):374–381, 1939.

[641] W. G. Unruh. Notes on black-hole evaporation. *Physical Review D*, 14(4):870–892, 1976.

[642] Don N. Page. Particle emission rates from a black hole: Massless particles from an uncharged, nonrotating hole. *Physical Review D*, 13(2):198–206, 1976.

[643] Abhay Ashtekar and Badri Krishnan. Isolated and dynamical horizons and their applications. *Living Reviews in Relativity*, 7(10), 2004. Review article; open access.

[644] Emanuele Berti, Vitor Cardoso, and Andrei O. Starinets. Quasinormal modes of black holes and black branes. *Classical and Quantum Gravity*, 26(16):163001, 2009.

[645] Pau Amaro-Seoane and et al. (LISA Consortium). Laser interferometer space antenna. *arXiv e-prints*, 2017.

[646] Yosuke Mizuno and et al. The current ability to test theories of gravity with black hole shadows. *Nature Astronomy*, 2:585–590, 2018.

[647] B. Carter. Global structure of the kerr family of gravitational fields. *Physical Review*, 174(5):1559–1571, 1968.

[648] D. C. Robinson. Uniqueness of the kerr black hole. *Physical Review Letters*, 34(14):905–906, 1975.

[649] Subrahmanyan Chandrasekhar. *The Mathematical Theory of Black Holes*. Oxford University Press, Oxford, 1998.

[650] Alexei Kitaev and John Preskill. Topological entanglement entropy. *Physical Review Letters*, 96(11):110404, 2006.

[651] Michael Levin and Xiao-Gang Wen. Detecting topological order in a ground state wave function. *Physical Review Letters*, 96(11):110405, 2006.

[652] Joseph J. Bisognano and Eyvind H. Wichmann. On the duality condition for a hermitian scalar field. *Journal of Mathematical Physics*, 16(4):985–1007, 1975.

[653] Shinsei Ryu and Tadashi Takayanagi. Holographic derivation of entanglement entropy from the ads/cft correspondence. *Physical Review Letters*, 96(18):181602, 2006.

[654] Roger Penrose. Gravitational collapse: The role of general relativity. *Rivista del Nuovo Cimento (Numero Speciale)*, 1:252–276, 1969. Reprinted in Gen. Relativ. Gravit. **34**, 1141–1165 (2002), doi:10.1023/A:1016578408204.

[655] Roger Penrose and R. M. Floyd. Extraction of rotational energy from a black hole. *Nature Physical Science*, 229:177–179, 1971.

[656] James M. Bardeen, William H. Press, and Saul A. Teukolsky. Rotating black holes: Locally nonrotating frames, energy extraction, and scalar synchrotron radiation. *The Astrophysical Journal*, 178(2):347–369, 1972.

[657] Subrahmanyan Chandrasekhar. *The Mathematical Theory of Black Holes*. Clarendon Press, Oxford, 1983. Reissued in Oxford Classic Texts; see also key {Chandrasekhar:1985kt} in your file.

[658] Richard Brito, Vitor Cardoso, and Paolo Pani. *Superradiance: Energy Extraction, Black-Hole Bombs and Implications for Astrophysics and Particle Physics*, volume 906 of *Lecture Notes in Physics*. Springer, Cham, 2015.

[659] James B. Hartle. Slowly rotating relativistic stars. i. equations of structure. *The Astrophysical Journal*, 150:1005–1029, 1967.

[660] Steven B. Giddings. Black hole information, unitarity, and nonlocality. *Physical Review D*, 74(10):106005, 2006.

[661] David Hilbert. Die grundlagen der physik. (erste mitteilung). *Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen, Mathematisch-Physikalische Klasse*, pages 395–407, 1915.

[662] David Lovelock. The einstein tensor and its generalizations. *Journal of Mathematical Physics*, 12(3):498–501, mar 1971.

[663] David Lovelock. The four-dimensionality of space and the einstein tensor. *Journal of Mathematical Physics*, 13(6):874–876, jun 1972.

[664] Lee Smolin. *Three Roads to Quantum Gravity*. Basic Books, 3rd, revised edition, 2017.

[665] Albert Einstein. Zur elektrodynamik bewegter körper. *Annalen der Physik*, 17(10):891–921, 1905.

[666] Hermann Minkowski. Raum und zeit. *Physikalische Zeitschrift*, 10:104–111, 1909. Lecture delivered Sept. 21, 1908 (Cologne). Another version: *Jahresbericht der DMV* **18** (1909) 75–88; English translation in *The Principle of Relativity* (Methuen, 1923).

[667] Carlo Rovelli. *Helgoland: Making Sense of the Quantum Revolution*. Allen Lane, London, 2021. UK ed. Title also published in the US by Riverhead Books (Penguin Random House), ISBN 9780593328897.

[668] Edward P. Tryon. Is the universe a vacuum fluctuation? *Nature*, 246(5433):396–397, 1973.

[669] Alexander Vilenkin. Creation of universes from nothing. *Physics Letters B*, 117(1–2):25–28, 1982.

[670] Raphael Bousso. The holographic principle. *Reviews of Modern Physics*, 74:825–874, 2002.

[671] Ahmed Almheiri, Xi Dong, and Daniel Harlow. Bulk locality and quantum error correction in AdS/CFT. *Journal of High Energy Physics*, 2015(4):163, 2015.

[672] Fernando Pastawski, Beni Yoshida, Daniel Harlow, and John Preskill. Holographic quantum error-correcting codes: Toy models for the bulk/boundary correspondence. *Journal of High Energy Physics*, 2015(6):149, 2015.

[673] Stephen W. Hawking. *A Brief History of Time: From the Big Bang to Black Holes*. Bantam Books, New York, 1988.

[674] Paul C. W. Davies. *The Cosmic Blueprint: New Discoveries in Nature's Creative Ability to Order the Universe*. Simon & Schuster, New York, 1988. Reissued with new material by Templeton Foundation Press, 2004.

[675] Carlo Rovelli. *White Holes*. Riverhead Books, New York, 2023.

[676] Patrick Olivelle. *Upaniṣads*. Oxford World's Classics. Oxford University Press, Oxford, 1996.

[677] Humberto R. Maturana and Francisco J. Varela. *Autopoiesis and Cognition: The Realization of the Living*, volume 42 of *Boston Studies in the Philosophy of Science*. D. Reidel Publishing Company, Dordrecht, 1980.

[678] Jan Gonda. Some observations on the meaning of the word ātman. *Bulletin of the School of Oriental and African Studies, University of London*, 25(1/3):68–96, 1962.

[679] Sarvepalli Radhakrishnan. *The Principal Upanishads*. Indus / HarperCollins Publishers India, New Delhi, 1994.

[680] Aristotle. *Nicomachean Ethics*. Oxford University Press, Oxford, 1925. Originally composed c. 350 BCE.

[681] Francisco J. Varela, Evan Thompson, and Eleanor Rosch. *The Embodied Mind: Cognitive Science and Human Experience*. MIT Press, Cambridge, MA, 1991.

[682] Sarvepalli Radhakrishnan. *Indian Philosophy. Volume I*. Muirhead Library of Philosophy. George Allen & Unwin; The Macmillan Company, London; New York, 1923. Vol. II appeared in 1927.

[683] Stephen W. Hawking. Black holes and baby universes. In *Black Holes and Baby Universes and Other Essays*, pages 104–114. Bantam Books, New York, 1993.

[684] C. J. Isham. Structural issues in quantum gravity. In M. Francaviglia, G. Longhi, L. Lusanna, and E. Sorace, editors, *General Relativity and Gravitation: Proceedings of the 14th International Conference*, pages 167–209. World Scientific, 1997. Also appears as arXiv:gr-qc/9510063.

[685] Carlo Rovelli. Time in quantum gravity: An hypothesis. *Physical Review D*, 43(2):442–456, 1991.

[686] Carlo Rovelli. The incarnation of time. arXiv:gr-qc/9803024, 1998.

[687] Andrei D. Linde. *Particle Physics and Inflationary Cosmology*. Harwood Academic Publishers, Chur, 1990.

[688] Lawrence M. Krauss. *A Universe from Nothing: Why There Is Something Rather than Nothing*. Free Press, New York, 2012.

[689] David J. Kalupahana. *The Principles of Buddhist Psychology*. State University of New York Press, Albany, 1987.

[690] Thích Nh at Hãnh. *The Heart of Understanding: Commentaries on the Prajñapāramitā Heart Sūtra*. Parallax Press, Berkeley, CA, 1988. Later reprints and revised editions appear.

[691] Baruch Spinoza. *Ethica, ordine geometrico demonstrata*. Johannes Rieuwertsz, Amsterdam, 1677. Printed in *Opera Posthuma*.

[692] Baruch Spinoza. *Ethics: with The Treatise on the Emendation of the Intellect and Selected Letters*. Hackett, Indianapolis, 1992. Edited with introduction and notes by Seymour Feldman.

[693] John Sellars. *Stoicism*. Ancient Philosophies. Routledge, London and New York, 2006.

[694] Alfred North Whitehead. *Process and Reality: An Essay in Cosmology*. Free Press, New York, corrected ed. edition, 1978. Corrected edition edited by David Ray Griffin and Donald W. Sherburne.

[695] John A. Wheeler. Law without law. In John A. Wheeler and Wojciech H. Zurek, editors, *Quantum Theory and Measurement*, pages 182–213. Princeton University Press, Princeton, NJ, 1983.

[696] Jonathan J. Halliwell. Introductory lectures on quantum cosmology. In S. Coleman, J. B. Hartle, T. Piran, and S. Weinberg, editors, *Quantum Cosmology and Baby Universes*, pages 159–243. World Scientific, Singapore, 1991.

[697] William K. Wootters. Time replaced by quantum correlations. *International Journal of Theoretical Physics*, 23(8):701–711, 1984.

[698] Edward P. Tryon. Is the universe a vacuum fluctuation? *Nature*, 246(5433):396–397, 1973.

[699] Andrei D. Linde. The inflationary universe. *Reports on Progress in Physics*, 47(8):925–986, 1984.

[700] Nāgārjuna. *The Fundamental Wisdom of the Middle Way: Nāgārjuna's Mūlamadhyamakakārikā*. Oxford University Press, New York, 1995.

[701] Y. Karunadasa. *The Theravāda Abhidhamma: Its Inquiry into the Nature of Conditioned Reality*. Centre for Buddhist Studies, The University of Hong Kong, Hong Kong, 2010.

[702] Thich Nhat Hanh. *Interbeing: Fourteen Guidelines for Engaged Buddhism*. Parallax Press, Berkeley, CA, revised ed. edition, 1998. Revised edition edited by Fred Eppsteiner.

[703] Murray Gell-Mann and James B. Hartle. Quantum mechanics in the light of quantum cosmology. In S. Kobayashi, H. Ezawa, Y. Murayama, and S. Nomura, editors, *Proceedings of the 3rd International Symposium on the Foundations of Quantum Mechanics in the Light of New Technology*, pages 321–343, Tokyo, 1990. Physical Society of Japan.

[704] Thomas Aquinas. *Summa Contra Gentiles. Book One: God*. University of Notre Dame Press, Notre Dame, IN, 1975.

[705] Stephen Hawking and Roger Penrose. *The Nature of Space and Time*. Princeton University Press, Princeton, NJ, 1996.

[706] Charles Hartshorne. *The Logic of Perfection and Other Essays in Neoclassical Metaphysics*. Open Court, La Salle, IL, 1962.

[707] Murray Gell-Mann and James B. Hartle. Quantum mechanics in the light of quantum cosmology. In Wojciech H. Zurek, editor, *Complexity, Entropy, and the Physics of Information*, volume VIII of *Santa Fe Institute Studies in the Sciences of Complexity*, pages 425–458. Addison-Wesley, Reading, MA, 1990.

[708] Henri Bergson. *Creative Evolution*. Henry Holt and Company, New York, 1911.

[709] Arthur Stanley Eddington. *The Nature of the Physical World*. Cambridge University Press, Cambridge, 1928. Gifford Lectures, 1927.

[710] Huw Price. *Time's Arrow and Archimedes' Point: New Directions for the Physics of Time*. Oxford University Press, New York, 1996.

[711] Roger Penrose. Singularities and time-asymmetry. In S. W. Hawking and W. Israel, editors, *General Relativity: An Einstein Centenary Survey*, pages 581–638. Cambridge University Press, Cambridge, 1979.

[712] Paul Davies. *The Mind of God: The Scientific Basis for a Rational World*. Simon & Schuster, New York, 1992.

[713] John F. Haught. *Making Sense of Evolution: Darwin, God, and the Drama of Life*. Westminster John Knox Press, Louisville, KY, 2010.

[714] John F. Haught. *Science and Faith: A New Introduction*. Paulist Press, Mahwah, NJ, 2012.

[715] Leslie E. Ballentine. The statistical interpretation of quantum mechanics. *Reviews of Modern Physics*, 42(4):358–381, 1970.

[716] David Deutsch. *The Fabric of Reality: The Science of Parallel Universes—and Its Implications*. Allen Lane (The Penguin Press), London, 1997.

[717] David Wallace. *The Emergent Multiverse: Quantum Theory according to the Everett Interpretation*. Oxford University Press, Oxford, 2012.

[718] Simon Saunders, Jonathan Barrett, Adrian Kent, and David Wallace, editors. *Many Worlds? Everett, Quantum Theory, and Reality*. Oxford University Press, Oxford, 2010.

[719] James B. Hartle. Quantum mechanics of individual systems. *American Journal of Physics*, 36(8):704–712, 1968.